



(86) Date de dépôt PCT/PCT Filing Date: 2015/09/07
(87) Date publication PCT/PCT Publication Date: 2016/03/10
(85) Entrée phase nationale/National Entry: 2017/03/02
(86) N° demande PCT/PCT Application No.: IB 2015/056837
(87) N° publication PCT/PCT Publication No.: 2016/035059
(30) Priorité/Priority: 2014/09/05 (BE2014/0668)

(51) Cl.Int./Int.Cl. *A23J 3/18* (2006.01),
A23J 3/22 (2006.01)
(71) Demandeur/Applicant:
SYRAL BELGIUM NV, BE
(72) Inventeurs/Inventors:
REDL, ANDREAS, BE;
FENEUIL, AURELIEN, BE;
VOGEL, FABRICE, FR
(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : INCLUSION CONTENANT UN ANALOGUE DE VIANDE PROTEINIQUE AYANT UNE TEXTURE AMELIOREE
ET UNE DUREE DE CONSERVATION PROLONGEE
(54) Title: AN INCLUSION CONTAINING PROTEINACEOUS MEAT ANALOGUE HAVING AN IMPROVED TEXTURE
AND AN EXTENDED SHELF-LIFE




Fig. 1

(57) **Abrégé/Abstract:**

The invention concerns an extended shelf-life proteinaceous meat analogue wherein said meat analogue comprises at least one inclusion included in a matrix which comprises vegetable proteins among which at least wheat gluten and a plasticizer said

(57) **Abrégé(suite)/Abstract(continued):**

proteinaceous meat analogue having a water content of less than 20% (w/w) and a torque value at 10mn of at least 6 gram meter (g*m), as determined by a test A. The invention is further about a method for obtaining an extended shelf- life proteinaceous meat analogue comprising the step of (i) mixing vegetable proteins among which at least vital wheat gluten and a plasticizer for obtaining a dough, (ii) shaping said dough by at least cutting the dough in dough pieces and agglomerating said dough pieces with at least one inclusion to obtain an agglomerate, and (iii) heating said agglomerate wherein said extended shelf-life proteinaceous meat analogue has a water content of less than 20% (w/w). The invention also concerns a foodstuff comprising said extended shelf-life proteinaceous meat analogue and the method for obtaining said foodstuff.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(43) International Publication Date
10 March 2016 (10.03.2016)(10) International Publication Number
WO 2016/035059 A3

- (51) International Patent Classification:
A23J 3/18 (2006.01) A23J 3/22 (2006.01)
- (21) International Application Number:
PCT/IB2015/056837
- (22) International Filing Date:
7 September 2015 (07.09.2015)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
2014/0668 5 September 2014 (05.09.2014) BE
- (71) Applicant: SYRAL BELGIUM NV [BE/BE]; Burcht-
straat 10, B-9300 Aalst (BE).
- (72) Inventors: REDL, Andreas; Burchtstraat 10, B-9300
Aalst (BE). FENEUIL, Aurelien; Burchtstraat 10, B-9300
Aalst (BE).
- (74) Agents: CHIELENS, Kristof et al.; President Kennedy-
park 31 C, B-8500 Kortrijk (BE).
- (81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,

KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))


Published:

— with international search report (Art. 21(3))

— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))

- (88) Date of publication of the international search report:
28 April 2016

(54) Title: AN INCLUSION CONTAINING PROTEINACEOUS MEAT ANALOGUE HAVING AN IMPROVED TEXTURE
AND AN EXTENDED SHELF-LIFE

**Fig. 1**

(57) **Abstract:** The invention concerns an extended shelf-life proteinaceous meat analogue wherein said meat analogue comprises at least one inclusion included in a matrix which comprises vegetable proteins among which at least wheat gluten and a plasticizer said proteinaceous meat analogue having a water content of less than 20% (w/w) and a torque value at 10mn of at least 6 gram meter (g*m), as determined by a test A. The invention is further about a method for obtaining an extended shelf- life proteinaceous meat analogue comprising the step of (i) mixing vegetable proteins among which at least vital wheat gluten and a plasticizer for obtaining a dough, (ii) shaping said dough by at least cutting the dough in dough pieces and agglomerating said dough pieces with at least one inclusion to obtain an agglomerate, and (iii) heating said agglomerate wherein said extended shelf-life proteinaceous meat analogue has a water content of less than 20% (w/w). The invention also concerns a foodstuff comprising said extended shelf-life proteinaceous meat analogue and the method for obtaining said foodstuff.



WO 2016/035059 A3

An inclusion containing proteinaceous meat analogue having an improved texture and an extended shelf-life

FIELD OF THE INVENTION

5 This invention concerns an extended shelf-life proteinaceous meat analogue and the method for obtaining such extended shelf-life proteinaceous meat analogue. The invention also concerns a ready to eat proteinaceous meat analogue and the method for obtaining such ready to eat proteinaceous meat analogue. The invention further concerns a foodstuff comprising said extended shelf-life proteinaceous meat
10 analogue or ready to eat proteinaceous meat analogue and the method for obtaining said foodstuff.

BACKGROUND OF THE INVENTION

 Several types of wheat gluten-based meat analogues or wheat gluten based intermediates for preparing meat analogues have already been described in the
15 state of the art.

 The best known wheat gluten-based meat analogue is "SEITAN". This is in fact a wheat gluten dough which is seasoned. This dough is then cooked to provide the "wheat meat". Before final processing, SEITAN must be stored cold, or in frozen conditions, because of its high moisture content (>50% moisture).

20 Apart from SEITAN, also other wheat protein based meat replacers have been described in the prior art. However, it has been observed that the products of the prior art, if used as meat analogues or meat extenders, do show a number of shortcomings with regard to texture, storage stability and/or ease of processing. This might explain the relatively limited success of such products in the market.

25 Among those products from the prior art texturized meat analogues can be mentioned, such products are obtained by an extrusion step at high temperature (more than 120°C), those products are usually named high moisture texturized proteins (HMTP) or low moisture texturized proteins (LMTP). The methods for obtaining LMTP are for example disclosed in US5922392, GB1288193, US3769029
30 or WO2012/008994 and provide products having spongy texture and which cannot be shaped easily without using binders such as eggs. The HMTP are difficult to manufacture and may have a mushy consistency if expansion directly after the

extrusion at high temperature. Usually, a cooling die is used at the end of the extrusion in order to avoid any expansion of the product. However, such cooling die is regularly clogged, thereby varying pressure within the barrel of the extruder and resulting in an unpredictable efflux of the product from the die as mentioned in US 20100136201. Moreover, an incorrect orientation of fibers in the product may take place inducing additional treatment of the product to re-orientate the fibers. Such treatment leads to complex and costly installations. A simple and cheap process providing a proteinaceous meat analogue having a good texture is needed.

Besides, the applicant has developed a low moisture plasticized wheat gluten composition (WO2007006431). However, this composition is only suitable for use in chewy pet food applications, because of its high breaking force.

Prior art also described meat analogues obtained by agglomerating protein filaments or protein fibers by using a binder such as albumin notably by making an emulsion then coagulating the emulsion obtained, such process is disclosed in GB977238. Such protein filaments or protein fibers are obtained by spinning process, which consists in the precipitation of a protein solution to form stretched fibers. However, spinning produces large waste water streams. In addition, the necessity for low pH, high salt concentrations and chemical additives makes the process very complex to implement (Manski et al., 2007).

Therefore the goal of this invention is to provide proteinaceous meat analogues showing improved meat-like properties with regard to texture, chewiness and bite, compared to already existing products, but also to provide proteinaceous meat analogues which can be shaped easily without adding any binder so as to form big pieces of meat analogues with irregular shapes such as natural pieces of meat but also so as to include easily pieces food (vegetables..) to provide a foodstuff having a good nutritional intake. At the same time it is the purpose of this invention to provide storage stable intermediates, which after hydration, provide easily ready to eat meat analogues showing a feeling in mouth which is as close as possible to meat without any additional treatment. Further, the invention provides processes for preparing these proteinaceous meat analogues and their storage-stable intermediates.

DESCRIPTION OF THE INVENTION

Extended shelf-life proteinaceous meat analogue and the method for obtaining such extended shelf-life proteinaceous meat analogue

The invention is about a **method for obtaining an extended shelf-life**
5 **proteinaceous meat analogue** comprising the step of:

- mixing at a temperature of less than 100°C i) vegetable proteins among which at least **vital wheat gluten** (VWG) ii) a **plasticizer** and optionally iii) **fibers** for obtaining a dough,
- shaping said dough by at least cutting the dough in dough pieces and
10 agglomerating said dough pieces with at least one inclusion to obtain an agglomerate, and
- heating preferably **static heating** between 120 to 160°C during 1 minute to 1 hour of said agglomerate for obtaining an extended shelf-life proteinaceous meat analogue

15 wherein said extended shelf-life proteinaceous meat analogue has a **water content** of less than 20% (w/w) and preferably, wherein said plasticizer is selected among a polyhydroxy alcohol, a starch hydrolysate, a carboxylic acid and mixture thereof.

Typically, the **mixing step** may be implemented by any mixing device capable of handling high viscosity materials (such as z blenders, ribbon mixers,
20 planetary mixers or co rotating intermeshing extruders). Advantageously, the mixing step is carried out in a batch or a continuous mixing unit, preferably at a temperature of less than 100°C (such as between 20 to 90°C, preferably, 30 to 65°C more preferably 40 to 50°C) for a period sufficient to obtain a dough-like composition. Typically, the mixing step is an extrusion step.

25 According to the invention, an “**extrusion**” refers to a process in which a material is pushed under compressive stresses through a deformation control element such as a die to form an elongated product. Continuous extrusion refers to an extrusion process where such deformation is carried out on a product of unlimited length. Advantageously, the extrusion according to the invention is
30 implemented at a temperature of less than 100°C, such as between 20 to 90°C, preferably, 30 to 65°C more preferably 40 to 50°C. Indeed, the inventors have

shown that a mixing step at a temperature above 100°C notably by extrusion at temperature above 100°C provides a product having spongy texture which is not observed for the product of the invention.

Advantageously, a food product, a food additive and/or a processed food
5 may be added during the mixing step. Such food product, food additive and/or processed food may be in a dried form or a hydrated form, typically by having a water content of more than 50%.

As used herein "**processed food**" refers to a food which is significantly modified from its natural state, as by mechanical alteration (such as grinding or
10 chopping), combination with other food products or additives, and/or cooking. As used herein, "processed food" excludes foods which substantially maintain their natural state after processing. For example, fresh produce may be washed, sorted, coated or treated, and packaged, but remain substantially in its natural state after processing, and would not be considered a "processed food" for the purpose of this
15 disclosure. A "processed food" also refers to an extract of food. The "*extract*" refers to the resultant solid or liquid material from an extraction. Indeed, an extract obtained from animal origins (such as meat or fish) or vegetal origins, such extract may be soluble or insoluble carbohydrates, *proteins*, *fibers*, *fat*, or combinations thereof. The processed food may comprise oil or fat particles, and particularly
20 preferred is vegetable oil or fat, especially that used in the form of sunflower oil. The processed food may also comprise particles of an inorganic salt. Calcium or magnesium salts are preferred. The processed food may comprise an insoluble material, for example an insoluble organic or inorganic salt.

As used herein "**food products**" refers to pieces of meat or fish or
25 vegetables such as for example nuts, cereals, carrots, leguminous.

As used herein "**food additives**" preferably includes simulated meat flavorings such as pork flavor, pepperoni flavor, smoke powder, chicken flavor, beef flavor, seafood flavor, savory flavorings (e.g., onion, garlic), vitamins (such as vitamins B12) and mixtures thereof.

30 According to the invention, said dough obtained by mixing of vegetable proteins and at least one plasticizer can further be shaped before the static heating.

The **shaping step** according to the invention comprises at least one or a combination of cutting, molding, sheeting and agglomerating steps. Preferably, the shaping step comprises at least a cutting step and a molding or a sheeting step, more preferably a cutting step, a sheeting step and a molding step.

5 According to an embodiment, the shaping step comprises a cutting step, an agglomerating step and optionally a molding and/or a sheeting step.

The cutting step may be done by using any means having at least one blade, for example a meat mincer (mincing machine), or a meat cutter.

10 Typically, the shaping step comprises a cutting step, an agglomerating step and optionally a molding and/or a sheeting step.

Advantageously, the agglomeration is carried out by i) hydrating the dough pieces and/or said at least one inclusion and ii) mixing to obtain an agglomerate, typically having a water content of between 5%-30%, preferably 7-25%, more preferably 10-20%.

15 According to one embodiment, the agglomeration may be carried out by

- i) hydrating the dough pieces, preferably by spraying 1-10% water by weight of dough pieces, ii) adding the at least one inclusion, and iii) mixing to obtain an agglomerated mixture typically having a water content of between 5%-30%, , preferably 7-25%, more preferably 10-20% or
- 20 • i) hydrating the mixture of dough pieces and the at least one inclusion, preferably by spraying 1-10% water by weight of dough pieces, and ii) mixing the hydrated mixture to obtained an agglomerated mixture typically having a water content of between 5%-30%, preferably 7-25%, more preferably 10-20% or
- 25 • i) hydrating at least one inclusion preferably by spraying 1-10% water by weight of dough pieces, ii) adding the dough pieces, and iii) mixing to obtain an agglomerated mixture typically having a water content of between 5%-30%, preferably 7-25%, more preferably 10-20%.
- 30

As used herein, the term "**inclusion**" refers to a coherent mass which is a food product a food additive and/or a processed food. The inclusion is not a dough piece according to the invention. The inclusion should be of a size that is visible, as a discrete piece or pieces, in the meat analogue prepared. Preferably the inclusion
5 has a maximum linear dimension of at least 1 mm, preferably of at least 3 mm, preferably between 1 and 8mm. Unless stated otherwise, the size of the inclusion is quoted as the maximum linear dimension of the inclusion, i.e. the maximum length in any dimension. Typically, the inclusion may be a dried or an hydrated product or may contained naturally water.

10 According to the invention, the dough piece has a mass of at least 0.03g preferably 0.04 to 300g more preferably 0.05 to 30g, 0.05g to 3g typically 0.05g to 0.2 g. Advantageously, said dough piece may have a long or a circular shape.

The **sheeting step** may be implemented by compressing dough pieces between cylindrical rolls

15 The **molding step** may be implemented by compressing dough pieces into molds of any desired shape.

The method of the invention is particularly advantageous in that without any added binder, the product develops a very good agglomerating and molding properties providing big pieces of meat analogues having very high volume.
20 Moreover, the texture of the product obtained is dense enough to provide a uniform piece of meat analogue. Examples of size and texture which can be provided is the one of an escalope or a wiener schnitzel. Such pieces cannot be obtained by using the meat analogues of the prior art without adding a binder such as eggs or xanthan or starch providing a product which cannot be eaten by vegan people or having poor
25 texture.

Moreover, the meat analogue according to the invention is particularly easy to mold and provides structure having irregular forms close to natural meat steaks or beefsteaks. Besides, the shaping steps such as the cutting, agglomerating, sheeting and molding steps can be repeated in order to obtain a product having a fibrillar
30 structure which can be close to the myofril structure of skeletal muscles. Typically, a very nice fibrillar structure can be obtained when the dough pieces are of longitudinal shape.

As used herein the term “**static heating**” refers to a heating step without any stirring or shearing of the dough to be heated. An example of static heating may be oil frying, microwaving, or by using an oven, or a hot plate. Typically, the **heating step** is carried out at a temperature between 120 to 160°C during 1 minute to 1 hour, typically, 10min to 30mn. The duration of the heating step can be adapted by the man skilled in the art depending on the volume of mixed proteins to be heated. The inventors have shown that boiling (100°C) or an heating at 110°C of the mix of vegetable proteins and plasticizer does not provide the meat analogue of the invention. Moreover, an heating step above 160°C for example a classical frying step (at 185°C) provides a product which has a very soft and spongy texture and a dark color..

It should be emphasized that meat analogues in accordance with the present invention may be used as meat replacements to provide a meatless foodstuff or as meat enhancers/extenders to replace a portion of the meat that would normally be present in a foodstuff

As used herein **an extended shelf-life product** is a product which is shelf-stable at ambient temperatures for an extended period of time for e.g. up to and above 60 days. Such extended shelf-life product is microbiologically stable because of its reduced water content. Such an extended shelf-life product may be eaten after hydration and cooking. Such extended shelf life product may be a petfood, typically designated to cats and dogs.

As used herein the term “**vital wheat gluten**” refers to those forms of dried wheat gluten that have been subjected to only minimal or no heat denaturation during drying. Upon reconstitution with water, vital wheat gluten shows physical properties (e.g., elasticity, gumminess, etc.) similar to those of freshly prepared wet wheat gluten.

Preferably, the **vegetable proteins** are powdered vegetable proteins (as measured according to AOAC 979.09; Kjeldahl method with a conversion factor N*6,25). Advantageously, said vegetable protein powder has a water content of between 1 to 15% preferably 2 to 12%, typically, 3 to 7%. Said vegetable proteins may be vital wheat gluten (100% of vital wheat gluten) advantageously a vital wheat gluten powder. Typically, said vegetable proteins are a mixture of vital wheat gluten and vegetable proteins from at least one other origin. According to one embodiment,

vegetable proteins comprise more than 50% of vital wheat gluten, preferably, more than 60, 70, 80, 85, 95% of vital wheat gluten. Preferably, the "vegetable proteins from at least one other origin" or "vegetable protein other than VWG" may be selected from a group consisting of potato, lupine, soya, pea, chick pea plants, alfalfa, faba bean, lentil, bean, rapeseed, sunflower and cereals such as corn, barley, malt and oats. Said vegetable proteins are typically in the form of flour, concentrate or isolate.

Vital wheat gluten or **VWG** refers to the dried, insoluble gluten portion of wheat flour from which the starch and soluble components have been removed by a washing process. Typically, vital wheat gluten is then dried to a fine powdered state. Vital wheat gluten powder typically has a percent protein on a dry basis of about 80% or greater (as measured according to AOAC 979.09; Kjeldahl method with a conversion factor N*6,25). Vital wheat gluten is typically not denatured as determined by the test procedure in "Approved Methods of the American Association of Cereal Chemist", Method 38 entitled "Vital Wheat Gluten" (December 1962). Useful vital wheat gluten powder is commercially available under the trade designation AMYGLUTEN®.

As used herein, a "**plasticizer**" refers to a compound that increases the plasticity or fluidity of the material to which it is added. Typically, the plasticizer of the invention is a "food grade plasticizer" which is a plasticizer approved to be used in foods. Advantageously, the plasticizer is a non-aqueous plasticizer, typically said plasticizer has a **water content** of less than 20% (w/w). Advantageously the plasticizer may be a polyhydroxy alcohol (such as glycerol, sorbitol, ethylene glycol, polyethylene glycol propyleneglycol, butanediol, polyethylenglycol and mixture thereof), a starch hydrolysate (such as a glucose syrup), a carboxylic acid and mixture thereof.

As used herein, "**fibers**" may be insoluble fibers, preferably from cereal, tuber, seed or leguminosae.

The term "**water content**" refers to the content of water based upon the Loss on Drying method as described in *Pharmacopeial Forum*, Vol. 24, No. 1, page 5438 (January-February 1998). The calculation of water content is based upon the percent of weight that is lost by drying.

According to one embodiment, said proteinaceous meat analogue comprises:

- 5
 - 30 to 90%, of vegetable proteins on dry mass, preferably 40 to 85%, more preferably 45 to 80% (w/w), even more preferably, 50% to 78% of vegetable proteins typically VWG notably VWG powder, advantageously said vegetable proteins comprises 40 to 90% of VWG powder and 0 to 40% of vegetable proteins other than VWG
- 10
 - 10 to 40% (w/w), preferably, 15 to 30%, typically 20 to 27% of plasticizer on dry mass
 - 0 to 8% (w/w) of fibers on dry mass

Advantageously said method for obtaining an extended shelf-life proteinaceous meat analogue comprises mixing at a temperature of less than 100°C of i) 30 to 90%, preferably 40 to 85% of VWG powder, ii) 10 to 40% of a plasticizer, 15 iii) 0 to 40% preferably 3 to 30%, typically 5 to 20% of a vegetable protein other than VWG and iv) 0 to 8%, preferably, 1 to 5% of fibers.

The invention is also about an **extended shelf-life proteinaceous meat analogue directly obtained by the method** according to the invention.

The invention is further about an **extended shelf-life proteinaceous meat analogue** comprising at least one inclusion included in a matrix which comprises:

- 20
 - 30 to 90% (w/w), preferably 40 to 85%, more preferably 45 to 80% (w/w), even more preferably, 50% to 78% of vegetable proteins among which at least wheat gluten, preferably at least 50% of wheat gluten,
 - 25
 - 10 to 40% (w/w), preferably, 15 to 30%, typically 20 to 27% of plasticizer,
 - 0 to 8% (w/w) of fibers
 - less than 20% (w/w) of water content.

Preferably said proteinaceous meat analogue has a water content of less than 20% (w/w) and a torque value at 10mn of at least 6 gram meter (g*m)

(preferably between 6 to 20, more preferably between 8 to 15), or a ratio torque value at 30sec/ torque value at 10mn of about 1 to 1.7, as determined by a test A.

Indeed, the aim of the invention is to provide a meat analogue, meaning a product which may be a good meat replacer. The inventors have shown that the characteristic which is important to be a good meat analogue is to give the feeling of meat during mastication. The inventors have shown that to the opposite of the meat analogues of the market such as for example, the low moisture texturized proteins, the meat analogue of the invention gives a feeling which is very close to meat due to its physicochemical characteristics. The feeling during the first bite (30 sec) than after several mastications (10min) can be evaluated by the ratio torque value at 30sec/ torque value at 10min.

The invention is further about an **extended shelf-life proteinaceous meat analogue** comprising at least one inclusion included in a matrix comprising vegetable proteins among which at least **wheat gluten**, notably VWG such as for example VWG powder and a **plasticizer** said proteinaceous meat analogue having a water content of less than 20% (w/w) and a torque value at 10mn of at least 6 gram meter (g*m) (preferably between 6 to 20, more preferably between 8 to 15), or a ratio torque value at 30sec/ torque value at 10mn of about 1 to 1.7, as determined by a test A, preferably said proteinaceous meat analogue also comprises fibers.

As used herein the “**torque value at 30sec**” is the value measured in the first 30 sec of the Brabender Plastograph assay (Test A or B). As used herein the “**torque value at 10mn**” is the value measured after 10 min of the Brabender Plastograph assay. The torque values are measured in gram meter (g*m).

As used herein the **torque value at 30sec** and the **torque value at 10mn** of the extended shelf-life proteinaceous meat analogue are evaluated according to a **Test A** by using a Brabender Plastograph, type EC equipped with a type 50 measuring head and a mixing chamber comprising two counter-rotating identical sigma shaped mixing blades (references of the blades: Sigma (S)), the samples are hydrated to their equilibrium (their water holding capacity), meaning between 55 to 70% of water content by for example a boiling step of 20mn or by hydrating the samples in excess of water during 48h at room temperature, 70g of cube shaped samples (15*15mm) are then introduced in a mixing chamber thermo-stated at 37°C

and measurement is obtained by counter-rotating the mixing blades at differential speeds of 34rpm and 22.67rpm ($34 \times 2/3$).

Preferably, the extended shelf-life proteinaceous meat analogue has a **ratio torque value at 30sec/ torque value at 10mn** of about 1 to 1.7, typically, 1 to 1,6%, more preferably 1 to 1.5, preferably 1 to 1.4, more preferably 1 to 1.3, of about 1.1 to 1.25.

The invention, further concerns **a foodstuff comprising the extended shelf-life proteinaceous meat analogue** of the invention. Typically, said foodstuff is an extended shelf-life foodstuff advantageously having a water content of less than 20%. Said, foodstuff may be obtained by simply adding or mixing the extended shelf-life proteinaceous meat analogue with a food products, food additives and/or processed food.

The term “**foodstuff**” means any material, substance, additive, that can be used as food, feed or that may be added to food or feed. Typically the foodstuff is any composition which an animal, preferably a mammal such as a human, a dog or a cat may consume as part of its diet.

Said foodstuff may be for example dried food such as dry pet food, pet food treats but also food aiming at long transportation like food for soldiers, astronauts. Typically dried burger, sausage, schnitzel, “meat” ball, wok piece, filet.

A ready to eat proteinaceous meat analogue and the method for obtaining such ready to eat proteinaceous meat analogue

The invention concerns a method **for obtaining a ready-to-eat proteinaceous meat analogue** wherein said method comprises step of

- obtaining an extended shelf-life proteinaceous meat analogue according to the method of the invention,
- hydrating said extended shelf-life proteinaceous meat analogue preferably, until a water content of more than 50% is reached preferably, 60 to 70% preferably 62 to 68% more preferably 65% water content and
- optionally, cooking said hydrated proteinaceous meat analogue
- obtaining a ready-to-eat proteinaceous meat analogue

According to the invention, the **cooking** step is implemented at a temperature of more than 100°C, typically, at a temperature between 30 to 200°C during 1 minute to 1 hour. Said cooking step may be for example a sterilization step.

5 The invention further concerns the ready-to-eat proteinaceous meat analogue directly obtained by the method of the invention.

The invention is further about a **ready-to-eat proteinaceous meat analogue** wherein, said ready-to-eat meat analogue comprises vegetable proteins among which at least **wheat gluten**, a **plasticizer** and preferably fibers and has a water content of more than 50%, preferably, between 60 to 70%, more preferably 62
10 to 68%, even more preferably 65% water content, a torque value at 10mn of at least 6 gram meter meter (preferably between 6 to 20, more preferably between 8 to 15) or a **ratio torque value at 30sec/ torque value at 10mn** of about 1 to 1.7, as determined by a Test B.

As used herein the **torque value at 30sec** and the **torque value at 10mn**
15 of the extended shelf-life proteinaceous meat analogue comprised an extended shelf-life foodstuff are evaluated according to a **Test A** by using a Brabender Plastograph, type EC equipped with a type 50 measuring head and a mixing chamber comprising two counter-rotating identical sigma shaped mixing blades (references of the blades: Sigma (S)), hydrated to their equilibrium (their water
20 holding capacity), meaning between 55 to 70% of water content by for example a boiling step of 20mn or by hydrating the samples in excess of water during 48h at room temperature, 70g of cube shaped samples (15*15mm) are then introduced in a mixing chamber thermo-stated at 37°C and measurement is obtained by counter-rotating the mixing blades at differential speeds of 34rpm and 22.67rpm (34*2/3).

25 Preferably, the proteinaceous meat analogue has a **ratio torque value at 30sec/ torque value at 10mn** of about 1 to 1.5, preferably 1 to 1.4, more preferably 1 to 1.3, of about 1.1 to 1.25.

As used herein "**ready-to-eat proteinaceous meat analogue**" refers to a proteinaceous meat analogue which can be eaten as is. A **ready-to-eat foodstuff**
30 refers to a hydrated (having more than 50% (w/w) water content, between 60 to 70%, preferably 62 to 68%, more preferably 65% of water content) and optionally cooked foodstuff (with the same procedure such as meat e.g. at a temperature between 30 to 200°C during 1 minute to 1 hour) or a foodstuff comprising a hydrated

and optionally cooked (with the same procedure such as meat e.g. at a temperature between 30 to 200°C during 1 minute to 1 hour) proteinaceous meat analogue.

As used herein the torque value at 30sec and the torque value at 10mn of
5 the proteinaceous meat analogue comprised an ready-to-eat foodstuff are evaluated according to a **Test B** by using a Brabender Plastograph, type EC equipped with a type 50 measuring head and a mixing chamber comprising two counter-rotating identical sigma shaped mixing blades (references of the blades: Sigma (S)), 70g of cube shaped samples (15*15mm) are introduced in a mixing chamber thermo-stated
10 at 37°C and measurement is obtained by counter-rotating the mixing blades at differential speeds of 34rpm and 22.67rpm (34*2/3). If needed, the samples may be washed before analysis in order to remove any residual sauce, for example by several washing steps with water and if needed hot water.

According to the invention, said proteinaceous meat analogue is a cohesive
15 material. Typically, said proteinaceous meat analogue has a Fi value of more than 30%, preferably between 50% and 80%.

As used herein, the "**Fi value**" refers to the percentage of protein insoluble in a SDS buffer. The Fi value reflects the level of crosslinking of the gluten proteins. Indeed, it has been demonstrated that a Fi value of vital gluten is less than 10%
20 (*Redl A, Morel MH, Bonicel J, Guilbert S, Vergnes B. Rheol Acta 1999;38(4):311–20*). A Fi value of more than 30% is observed for example when the gluten has undergone a thermal treatment.

The Fi value may be measured according to a method well known from the man skilled in the art, and is described for example in *Redl A, Morel MH, Bonicel J, Guilbert S, Vergnes B. Rheol Acta 1999;38(4):311–20*. The Fi value of the
25 proteinaceous meat analogue is provided by the static heating step of the method of the invention.

The invention is also about a **ready-to-eat proteinaceous meat analogue comprising:**

- 30
- 60 to 70%, preferably 62 to 68%, more preferably 65% water content

- 15-25% proteins (preferably 17 -20%) among which more than 60% wheat gluten (preferably more than 70%, more preferably between 75 and 80%, or typically 100% of wheat gluten)
- 2 – 5% fat (3 to 4%)
- 5 • 8 – 15% (w/w) carbohydrates, typically between, 10 to 12% (w/w) among which less than 4% (w/w) of plasticizer, preferably 1 to 3.5% (w/w), more preferably 1.5 to 3%(w/w),
- 2 to 8% fibres, typically, between 3 to 6%, more preferably 3,2 to 5,8%.

10 As used herein, the term “**fat**” refers to triacylglycerides or triglycerides formed by the esterification reaction of long chain-, medium chain- or short chain-fatty acids with glycerol, a trihydroxy alcohol, or a mixture thereof, in any of solid, liquid or suspension forms, regardless of whether they are obtained from animal, fowl, fish or plants sources or are made synthetically, so long as they are safe for

15 consumption by mammals, particularly humans.

As used herein, “Carbohydrate” refers to at least a source of carbohydrates such as, but not limited to, monosaccharides, disaccharides, oligosaccharides, polysaccharides or derivatives thereof.

Preferably, said ready to eat proteinaceous meat analogue has a water

20 content of more than 50%, preferably, between 60 to 70%, more preferably 62 to 68%, even more preferably 65% water content, and/or a torque value at 10mn of at least 6 gram meter meter (preferably between 6 to 20, more preferably between 8 to 15) and/or a **ratio torque value at 30sec/ torque value at 10mn** of about 1 to 1.7, as determined by a Test B.

25 The invention is further about **a foodstuff comprising the ready to eat proteinaceous meat analogue** according to the invention. Typically, said foodstuff is an extended shelf-life foodstuff advantageously having a water content of more than 50%, preferably, between 60 to 70%, more preferably 62 to 68%, even more preferably 65% water content. Said, foodstuff may be obtained by simply adding or

30 mixing the ready to eat proteinaceous meat analogue with a food products, food additives and/or processed food.

Said foodstuff may be a ready to eat foodstuff, for example a burger, a sausage, a schnitzel, a “meat” ball, a wok piece, a filet.

Although having distinct meanings, the terms “comprising”, “having”,
5 “containing” and “consisting of” may be replaced with one another throughout the above description of the invention.

The invention will be further evaluated in view of the following examples and figures.

10 **BRIEF DESCRIPTION OF THE FIGURES**

Figure 1. Comparison of torque values between commercial products and product according to the invention (P3 6% fibers)

EXAMPLES

Example 1: Preparation of wheat gluten based products of the
15 invention

Vital wheat gluten powder (AMYGLUTEN®) (75 parts), glycerol (100%)(25 parts) and wheat fibre (VITACELL® 400) (0, 3 or 6 parts) are continuously mixed together in a twin screw extruder (Werner & Pfleiderer ZSK25) at a temperature of 65°C, and shaped and cut to obtain small particles with a ZGF30 pelletizer (Werner
20 & Pfleiderer). These particles can then be stored at room temperature, without risk of spoilage. Three different kinds of products (A1, A2, and A3) are obtained, having a different content of fibres (0, 3 and 6%).

Agglomerates of these particles are prepared by spraying 3 parts of water over the particles, and manually pressing the hydrated particles together into a
25 mould having the shape of a pork chop. The agglomerates obtained are then fried in cooking oil at 150°C for 5 minutes, thereby providing products P1, P2 and P3.

Example 2: Chewing and bite properties of meat analogues, and meat products:

The products prepared in example 1 were hydrated and then cooked in
30 boiling water for 20 minutes, resulting in products containing 60-65% water (35-40%

dry substance). Products differed in fibre content whereby P1 contained 0% fibre, P2 contained 3% fibre and P3 contained 6% fibre on dry substance.

The texture analysis was made in a Brabender Plastograph, type EC equipped with a type 50 measuring head and a mixing chamber comprising two counter-rotating identical sigma shaped mixing blades (references of the blades: Sigma (S)). 70g of cube shaped samples (15*15mm) are introduced in the mixing chamber thermo-stated at 37°C and measurement is obtained by counter-rotating the mixing blades at differential speeds of 34rpm and 22.67rpm (34*2/3).

The torque values were recorded and compared with cooked meat products such as cooked beef, cooked chicken and cooked schnitzel. Also torque values for meat analogues available in commerce, such as SEITAN, and commercial products such as "VIANA" cowgirl veggie steak, and "TOPAZ" Wheaty Vegankebab Döner were compared with the products of the invention. All these products were also boiled in water for 20 minutes.

In the following table, torque values are given after 30 seconds, 10 minutes and 14 minutes for all the above cited products.

<i>Torque (g*m) after</i>	<i>30 sec</i>	<i>10 min</i>	<i>14 min</i>
Cooked beef	70	35	32
Cooked chicken	30	20	18
Cooked schnitzel	13	13	11
Comm. Product 1	13	7	7
Comm. Product 2	15	8	7
Seitan	10	5	5
P1	13	12	10
P2	14	13	13
P3	17	14	13

Table 1: *Comparative analysis of torque values of meat pieces (beef, chicken, schnitzel) and meat analogues of the market (SEITAN, TOPAZ, VIANA) and of the invention at 30sec, 10min and 14min.*

A sensorial evaluation of the cooked pieces has been done in order to evaluate their textural properties compared with the measured values:

- Cooked beef is experienced as quite hard in bite (as expected for a beef cooked for 20 min in water). This observation is confirmed by the torque results which are very high at 30 sec, a slight reduction is observed at 10 and 14mn,
- 5 • Cooked chicken is experienced softer in bite than beef but still on the hard side. The torque values show quite constant results confirming the sensory evaluation.
- Cooked schnitzel is experienced as having a nice texture in a well-balanced equilibrium between hardness and softness of the texture
- 10 • The commercial products 1 and 2 were good in the first bite but lost structure very soon after the first bite and got too soft, the commercial products 1 and 2 shows a ratio torque value at 30 sec / torque value at 10mn of 1.9 which indicate a quick loss of texture in the mouth.
- Commercial SEITAN product was too soft from the very first bite, all
15 the torque values of SEITAN are below 10g*m, which seems to be too soft for a tasty meat analogue.

The examples of the invention (P1, P2 and P3) were considered as very nice in the first bite and also maintained a good texture during chewing.

20 The results clearly show that measured torque values correspond well to the experience of eating the test pieces. Indeed, it is clear that the products of the invention do show much better chewing and bite properties than actually commercialised wheat gluten based meat analogues.

At the same time these values illustrate that the products of the invention approach chewing and bite properties of natural meat products.

25 **Example 3:**

Vital wheat gluten powder (AMYTEX®)(50 parts), glycerol (100%)(33 parts), chick pea flour (13 parts) and wheat fibre (VITACEL® WF200) (4 parts) are continuously mixed together in a twin screw extruder (Werner & Pfleiderer ZSK25) at a temperature of 65°C, and shaped and cut to obtain small particles with a ZGF30
30 pelletizer (Werner & Pfleiderer), thereby providing product A4. These particles can then be stored at room temperature, without risk of spoilage.

Agglomerates of these particles are prepared by spraying 11 parts of water over the particles, and manually pressing the hydrated particles together into a mould of 15 mm thickness, thereby providing product W4. The product is then fried in cooking oil at 150°C for 10 minutes, thereby providing the product P4, and
 5 hydrated for 48 hours in water, resulting in products containing 60-65% water (35-40% dry substance). This product is prepared and analyzed with the Plastograph as described in example 2.

Torque (g*m) after	30 sec	10 min	14 min
P4	16	10	10

Table 2: Analysis of torque value of the product of the invention at 30 sec, 10min
 10 and 14min.

Example 4 : Different cooking conditions

The product W4 from example 3 is cut in pieces of 20 x 20 mm and cooked in several conditions.

The pieces are fried for 15 min either at 110°C or at 150°C or at 185°C. The pieces
 15 are cooked in an oven at 150°C either for 15 min or for 30 min. The pieces are cooked in boiling water (100°C) for 15 min. All these cooked products are then hydrated for 48 hours in water and tasted after cooking in a pan for 5 minutes.

The products cooked at 150°C (in the oven or in the fryer) were good in texture and have a nice yellow to light brown color.

20 The product cooked at 185°C is very dark (brownish, nearly black). Moreover, this product was found more spongy and softer in texture than products cooked at 150°C.

The products fried at 110°C or boiled at 100°C are found very white in appearance, so not attractive for a consumer and their texture was much softer than products
 25 cooked at 150°C.

Example 5 :

Vital wheat gluten powder (AMYGLUTEN®)(46 parts), chick pea flour (21 parts), wheat fibre (VITACEL® WF200) (4 parts) and glycerol (100%)(29 parts) are

continuously mixed together in a twin screw extruder (Werner & Pfleiderer ZSK25) at a temperature of 65°C, and shaped and cut to obtain small particles with a ZGF30 pelletizer (Werner & Pfleiderer), thereby providing product A5. These particles can then be stored at room temperature, without risk of spoilage.

- 5 Agglomerates of these particles are prepared by adding 1 part of salt, 13 parts of water and 25 parts of hydrated chick pea over the particles in a Meissner bowl cutter (RS 35) and cutting for 1 min (speed 3 for the bowl and speed 2 for the blade). Then, the agglomerates are grinded continuously through a Hobart meat grinder attachment (Hobart A200N, plate of #12 1/8"), thereby providing multiples strands.

10 **Example 6: Vegan Wiener schnitzel**

The continuous product of example 5 is cut in portions of 100 mm length. Each portion is passed through a manual dough sheeter set with a thickness between the steel cylinders of 1 to 3 mm in order to form a schnitzel having dimensions of up to about 300 mm in diameter and of 4 to 7 mm in thickness.

- 15 The product is then fried in cooking oil at 150°C for 15 minutes and hydrated for 48 hours in a vegetable broth (one vegetable bouillon cube of 11 g for 500 mL of water).

This product is then rolled in a mix of flour, yeast flakes, salt and water. Then, it is rolled in bread crumbs and fried at 170°C for 5 minutes or cooked in a pan for 5

- 20 minutes in order to prepare a vegan Wiener schnitzel.

The obtained Wiener Schnitzel is very nice in taste, texture and appearance. It has a crusty crumb with a firm but still tender bite.

Example 7 :

- 25 The product of example 5 is passed continuously through a manual dough sheeter set with a thickness between the steel cylinders of 5 mm, thereby providing a continuous strip of 15 mm thickness and 150 mm width. Pieces of 15 mm x 30 mm are cut from this strip.

- 30 These pieces are then fried in cooking oil at 150°C for 15 minutes, thereby providing the product P5, and hydrated for 48 hours, resulting in products containing 60-65% water (35-40% dry substance). This product is prepared and analyzed with the Plastograph as described in example 2.

Torque after	(g*m)	30 sec	10 min	14 min
P5		13	8	8

Table 3: Analysis of torque value of the product of the invention at 30 sec, 10min and 14min.

The nutritional content of the meat analogue of the invention was calculated and then compared to the ones of the meat analogue of the market (see table 4)

By comparing the products of the invention to the ones of the market based on wheat, pea or soy proteins, it can be seen that the product of the invention has a very low content in lipids but a high content in fibers, proteins and carbohydrates giving to this product a very nice nutritional profile.

Overall composition		Moisture	Proteins	Fibres	Fat	Salt	Carboh ydrates	Sugars	Gross Energy
		g/100g	g/100g	g/100g	g/100g	g/100g	g/100g	g/100g	KJ/100g
Ready-to-eat product of invention	P5 (analyzed)	64,2	17,6	4,1	3,1	0,3	10,5	0,3	619,6
	P5 (calculated)	65,0	18,8	5,3	2,3	0,7	6,5	0,5	644,0
	P4 (calculated)	65,0	22,4	3,8	2,0	0,2	6,5	0,7	594,0
Wheat protein based products	Seitanmedaillons	55,5	31,4		6,6	2,1	4,1	0,3	869,0
	DAS Schnitzel	51,6	25,1		9,4	2,5	9,2	2,2	969,0
	Frankenberger	53,1	25,6		12,0	1,4	6,2	1,7	981,0
	Weenies	49,0	30,6		14,6	1,7	3,7	0,4	1123,0
	Kebab gyros	53,7	26,4		12,4	2,5	3,5	1,5	993,0
Soy or pea protein products	Beefy crumble	63,6	23,6	1,8	9,1	1,2	1,8	1,8	837,4
	Feisty crumble	63,6	23,6	1,8	9,1	1,2	1,8	1,8	837,4
	crumble								
	grilled strips	68,2	23,5	1,2	5,9	0,8	1,2	1,2	591,1
	lightly seasoned strips	65,9	23,5	2,4	3,5	0,8	4,7	0,0	591,1
	Beast Burger	59,3	20,4	3,5	14,2	0,8	2,7	0,0	963,3
	Tofu Natur Bjorg	70,6	16,9	1,5	9,9	0,0	0,4	0,7	684,0

Table 4: Nutritional analysis of the soy, pea, or wheat based meat analogues of the market and the one of the invention

Example 8 : Product of invention in a complete dish

The product P5 from example 7 is hydrated for 48 hours in a vegetable broth (one vegetable bouillon cube of 11 g for 500 mL of water).

The hydrated pieces (1280 g) were cooked with:

- 80 g of Chinese dried black mushrooms (hydrated in excess of warm tap water)
- 1,6L of Coconut milk
- 0,8L of Coconut cream
- 5 - 150 g of Ginger (minced in SEB Optimo cutter)
- 100g of Lemon Grass (cut in pieces)
- 50 g green curry
- 50 g Basil (cut in fine pieces)
- 50 g of Soy sauce
- 10 - 2 g dried lime grass
- 100 mL of sunflower oil

The whole recipe was cooked on gentle heat and stirred for 30 min.

Upon stirring and heating, pieces remain entire without any disintegration. The whole dish is very attractive in appearance and taste.

15 **Example 9:** Sausages containing carrots, parsnips and leek

600 g of product A4 from example 3 are mixed with 150 g of leek, 125 g of carrots and 125g of parsnips in a Stephan cutter (UM 12 V Type). The mixture is then formed into sausages with a Bomann meat grinder (FW 443 CB Type) equipped with a sausage filling funnel and cellulose casings (Viscofan, composed of
20 65% regenerated cellulose, 18% glycerine, 15 % eau and 2% oil). After removing the casings, the sausages are cooked in a Miele steam oven at 121°C for 15 min, hydrated for 48 h and grilled in a pan. The obtained sausages have an attractive texture and color.

Example 10: Vegan meat balls

25 740 g of product A5 from example 5 are mixed with 150 g cashew nuts, 100 g of water and 10g of salt in a Stephan cutter (UM 12 V Type). The mixture is then formed through a Bomann meat grinder (FW 443 CB Type) and a very cohesive vegetable minced meat is obtained. The minced meat is formed by hand into "meat" balls The so formed meat balls are fried in oil at 150°C for 15 min, hydrated thereafter in excess water and cooked in a
30 pan. The obtained meat balls have an attractive taste, texture and appearance.

Example 11: Vegan hamburgers

690 g of product A5 from example 5 are mixed with 300 g of carrots and 10g of salt in a Stephan cutter (UM 12 V Type). The mixture is then formed into hamburgers using a Bomann meat grinder (FW 443 CB Type) and a hamburger shape. The so formed
5 hamburgers are fried in oil at 150°C for 15 min, hydrated thereafter in excess water and cooked in a pan. The obtained hamburgers have an attractive taste, texture and appearance. The carrot pieces can still be distinguished in the product and are very strongly embedded in the vegetable protein matrix.

CLAIMS

1. **Method for obtaining an extended shelf-life proteinaceous meat analogue**
5 comprising the step of:
- mixing at a temperature of less than 100°C i) vegetable proteins among which at least **vital wheat gluten** (VWG) ii) a **plasticizer** and optionally iii) **fibers** for obtaining a dough,
 - 10 • shaping said dough by at least cutting the dough in dough pieces and agglomerating said dough pieces with at least one inclusion to obtain an agglomerate, and
 - heating preferably **static heating** between 120 to 160°C during 1 minute to 1 hour of said agglomerate for obtaining an extended shelf-life proteinaceous meat analogue
- 15 wherein said extended shelf-life proteinaceous meat analogue has a **water content** of less than 20% (w/w) and preferably, wherein said plasticizer is selected among a polyhydroxy alcohol, a starch hydrolysate, a carboxylic acid and mixture thereof.
- 20 2. The method according to claim 1 wherein the proteinaceous meat analogue comprises:
- 30 to 90% (w/w) of vegetable proteins, preferably powdered vegetable proteins,
 - 10 to 40% (w/w) of plasticizer,
 - 25 • 0 to 8% (w/w) of fibers.
3. The method according to one of claim 1 or 2 wherein the **vegetable proteins** are a mixture of vital wheat gluten and vegetable proteins from at least one origin selected from a group consisting of potato, lupine, soya, pea, chick pea

plants, alfalfa, faba bean, lentil, bean, rapeseed, sunflower and cereals such as corn, barley, malt and oats.

4. The method according to any one of claims 1 to 3 wherein the mixing
5 step at a temperature of less than 100°C comprises mixing of 30 to 90% of VWG powder, 10 to 40% of a **plasticizer**, 0 to 40% of a vegetable protein other than VWG and 0 to 8% of **fibers**.
5. The method according to any one of claims 1 to 4 wherein the **fibers** are insoluble fibers, preferably from cereal, tuber, seed or leguminosae.
- 10 6. The method according to any one of claims 1 to 5 wherein the agglomeration is carried out by i) hydrating the dough pieces and/or said at least one inclusion and ii) mixing to obtain an agglomerate, typically having a water content of between 5%-30%, preferably 7-25%, more preferably 10-20%.
- 15 7. The method according to any one of claims 1 to 6 wherein a food product, a food additive and/or a processed food is added during the mixing step.
8. **An extended shelf-life proteinaceous meat analogue** directly obtained by the method according to any one of claims 1 to 7.
- 20 9. **An extended shelf-life proteinaceous meat analogue** comprising at least one inclusion included in a matrix comprising vegetable proteins among which at least **wheat gluten** and a **plasticizer** said proteinaceous meat analogue having a water content of less than 20% (w/w) and a torque value at 10mn of at least 6 gram meter or a **ratio torque value at 30sec/ torque value**
25 **at 10mn** of about 1 to 1.7, as determined by a test A, preferably said proteinaceous meat analogue also comprises fibers.
10. **An extended shelf-life proteinaceous meat analogue** comprising at least one inclusion included in a matrix which comprises:
 - 30 to 90% (w/w) of vegetable proteins among which at least wheat
30 gluten, preferably at least 50% of wheat gluten,
 - 10 to 40% (w/w) of plasticizer,

- 0 to 8% (w/w) of fibers
 - less than 20% (w/w) of water content.
11. **A foodstuff** comprising the extended shelf-life proteinaceous meat analogue according to any one of claims 8 to 10.
- 5 12. **Method for obtaining a ready-to-eat proteinaceous meat analogue** wherein said method comprises the step of:
- obtaining an extended shelf-life proteinaceous meat analogue according to the method of claims 1 to 7,
 - hydrating said extended shelf-life proteinaceous meat analogue preferably, until a water content of more than 50% is reached and
 - optionally, cooking the hydrated proteinaceous meat analogue
 - obtaining a ready-to-eat proteinaceous meat analogue.
- 10
13. **A ready-to-eat proteinaceous meat analogue directly obtained by the method** according to claim 12.
- 15
14. **A ready-to-eat proteinaceous meat analogue** wherein said meat analogue comprises at least one inclusion included in a matrix which comprises vegetable proteins among which at least **wheat gluten**, a **plasticizer** and preferably fibers and has a water content of more than 50%, a torque value at 10mn of at least 6 gram meter or a **ratio torque value at 30sec/ torque value at 10mn** of about 1 to 1.7, as determined by a Test B.
- 20
15. **A ready-to-eat proteinaceous meat analogue** comprising at least one inclusion included in a matrix said ready to eat **proteinaceous meat analogue** comprising
- 60 to 70%, preferably 62 to 68%, more preferably 65% water content
 - 15-25% proteins among which more than 60% wheat gluten
 - 2 – 5% fat,
- 25

26

- 8 – 15% carbohydrates among which less than 4% (w/w) of plasticizer, preferably 1 to 3.5%, more preferably 1.5 to 3%
- 2 to 8% fibres.

5 **16 A foodstuff comprising the ready to eat proteinaceous meat analogue**
according to any one of claims of claim 12 to 15.

10




Fig. 1




Fig. 1