



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2649977 C 2016/05/03

(11)(21) **2 649 977**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) **Date de dépôt PCT/PCT Filing Date:** 2007/06/14
(87) **Date publication PCT/PCT Publication Date:** 2007/12/21
(45) **Date de délivrance/Issue Date:** 2016/05/03
(85) **Entrée phase nationale/National Entry:** 2008/12/12
(86) **N° demande PCT/PCT Application No.:** AU 2007/000832
(87) **N° publication PCT/PCT Publication No.:** 2007/143788
(30) **Priorité/Priority:** 2006/06/14 (AU2006903203)

(51) **Cl.Int./Int.Cl. A01N 57/20** (2006.01),
A01N 25/06 (2006.01), **A01N 25/30** (2006.01),
A01P 13/00 (2006.01)
(72) **Inventeurs/Inventors:**
PENTLAND, PHILIP EDWARD, AU;
FLYNN, ANTHONY GERARD, AU
(73) **Propriétaire/Owner:**
EUREKA AGRESEARCH PTY LTD., AU
(74) **Agent:** BLAKE, CASSELS & GRAYDON LLP

(54) **Titre : COMPOSITION HERBICIDE ET PROCEDE D'ELIMINATION DE FEUILLAGE INDESIRABLE**
(54) **Title: HERBICIDAL COMPOSITION AND METHOD FOR REMOVING UNWANTED FOLIAGE**

(57) **Abrégé/Abstract:**

This invention relates to a method of preparing a spray tank mix of glyphosate comprising (a) providing glyphosate acid concentrate; (b) providing an alkaline composition; and (c) forming a mixture of glyphosate acid concentrate and alkaline composition in the spray tank by addition of a composition comprising the glyphosate acid concentrate to a diluted aqueous mixture of the alkaline composition.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 December 2007 (21.12.2007)

PCT

(10) International Publication Number
WO 2007/143788 A1

(51) International Patent Classification:

A01N 57/02 (2006.01) A01N 25/30 (2006.01)

(21) International Application Number:

PCT/AU2007/000832

(22) International Filing Date: 14 June 2007 (14.06.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

2006903203 14 June 2006 (14.06.2006) AU

(71) Applicant (for all designated States except US): **EUREKA AGRESEARCH PTY LTD** [AU/AU]; Dairy Road, Werribee, Victoria 3030 (AU).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **PENTLAND, Philip, Edward** [AU/AU]; 520 Racecourse Road, Flemington, Victoria 3031 (AU). **FLYNN, Anthony, Gerard** [AU/AU]; 24 James Cook Drive, Wandana Heights, Victoria 3216 (AU).

(74) Agent: **PHILLIPS ORMONDE & FITZPATRICK**; Level 21 & 22, 367 Collins Street, Melbourne, Victoria 3000 (AU).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- with amended claims

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HERBICIDAL COMPOSITION AND METHOD FOR REMOVING UNWANTED FOLIAGE

(57) Abstract: This invention relates to a method of preparing a spray tank mix of glyphosate comprising (a) providing glyphosate acid concentrate; (b) providing an alkaline composition; and (c) forming a mixture of glyphosate acid concentrate and alkaline composition in the spray tank by addition of a composition comprising the glyphosate acid concentrate to a diluted aqueous mixture of the alkaline composition.

WO 2007/143788 A1

HERBICIDAL COMPOSITION AND METHOD FOR REMOVING UNWANTED FOLIAGE

Field

- 5 This patent relates to herbicidal formulations and to their use in removing unwanted foliage. In particular the invention relates to a composition and method of use of the herbicide glyphosate.

10 In current practice the weed-killer glyphosate is transported to the ware-house of the distributor, and from thence to the farm or other site of application most commonly as a concentrate liquid aqueous formulation containing glyphosate in the form of a salt such as the isopropylamine or ammonium salts. In the manufacture of glyphosate the acid form is precipitated from the reaction mixture and the resulting wet cake is reacted with the appropriate base to form a water
15 soluble salt. The intermediate glyphosate technical grade acid is poorly water soluble and not generally useful as an herbicide. The herbicidal composition of salts of glyphosate is then prepared from the salt as an aqueous solution concentrate and may contain adjuvants such as a surfactant that reinforces the herbicidal effect of glyphosate.

20

The preparation of glyphosate as an aqueous concentrate of glyphosate salts necessarily entails disadvantages, such as the need to transport larger than necessary weights or volumes of prepared glyphosate salt liquor, the need to dispose of contaminated, rigid packaging materials, the danger that spilling the
25 prepared glyphosate salt liquor prior to its addition to the spray tank will create environmental damage, and the cost penalties associated with using robust packaging materials, and with the processing step of salt-formation.

30 US 6475954 (Hamroll et al. filed October 2, 2001) teaches the application of glyphosate acid (as distinct from glyphosate salt) in a composition which further comprises a carbonate or bicarbonate, organic acid chosen from citric, oxalic and

adipic acids, hydrosoluble cellulose, a glyphosate-synergising surfactant, and a bonding agent such as ammonium or sodium sulphate or urea. The formulations are generally presented as a tablet for addition to water, for convenient dispensing by small-scale users. Carbon dioxide is generated as a result of the acid-carbonate reaction to facilitate rapid disintegration of the tablet. Submerging the composition in water causes the glyphosate acid to convert into a soluble salt and solubilises the glyphosate by the formation of the salt. The teachings of this patent are not useful when weed control is required over a large area because the deposition of a large number of tablets into a large spray tank and the resultant foaming is not practical for broad acre use.

US Patent 5118338 describes a composition containing glyphosate acid solid powder or granules which are formulated using a very specific surfactant having a chain length of 16 to 18 carbon atoms and 25 units of ethylene oxide and an HLB of about 16. The surfactant is said to be very important to the success of the formulation.

US 6746988 (Hopkinson et al; filed September 5, 2002) teaches an agricultural composition comprising at least one agriculturally active compound (which may be glyphosate in salt form). This composition also comprises at least one alkyl polyglycoside, at least one anionic surfactant (selected from the set polyarylphenolpolyalkoxyether sulphate and polyarylphenolpolyalkoxyether phosphate) and at least one basic compound wherein the at least one anionic surfactant is neutralised to the inflection point in the titration curve with the at least one basic compound. Preferred basic compounds are selected from the set tallowamine ethoxylate, cocoamine alkoxylate, oleylamine alkoxylate, stearylamine alkoxylate, linoleic diethanolamide. The basic compounds are preferably present in an amount of 0.1 to 8% by weight, more preferably in an amount of 0.5 to 4% by weight. US 6746988 teaches a preference that anionic surfactants and basic compounds are in a ratio of about 1:1. The preferred range for anionic surfactants in the composition is 0.1 to 8% by weight, more

preferably 1-4%. The above formulation is a complex, multi-component formulation and is designed to provide robust in-tank performance for many different active ingredients. These formulations are intended for use as a one-pack formulation which includes the agriculturally active compound.

5

Patent US 6207617 (Gillespie J, filed April 1999) assigned to Monsanto teaches a concentrate composition of a plant treatment compound in acid form, wherein the plant treatment compound has a solubility in deionised water at 25 deg C of less than about 50 g/l and is present predominantly in the acid form. The composition further comprises an ethoxylated aliphatic hydrocarbon surfactant wherein the hydrocarbon has about 24 to 60 carbon atoms and the number of ethoxy moieties is about 5 to about 100. The plant treatment compound can be glyphosate. The weight ratio of surfactant to plant treatment compound, expressed as acid equivalent, is about 1:2 to 1:30. This invention is limited to one-pack formulations wherein all components of the plant treatment compound are contained in one pack. The surfactant is not a standard surfactant for herbicidal formulations.

Two pack formulations of glyphosate have also been proposed. Nufarm of Pipe Road Laverton, Victoria, Australia have sold "Credit plus Bonus", wherein one pack (Credit) comprises glyphosate salt (in fact a dual salt, more particularly isopropylamine and mono-ammonium salts) and the other pack (Bonus) comprises adjuvant described as a proprietary blend of surfactants, acidifiers and water conditioners. The two packs are added to the spray tank in a 1:1 ratio.

25

The UGA Cotton Newsletter of April 26, 2000 provides a table of glyphosate products together with adjuvant recommendations (provided as a separate component to the spray tank). Products in this table include products by BASF, Nufarm, Cheminova, Dow, Griffin LLC, Helena, Monsanto, Dupont and Zeneca. The products invariably comprise a glyphosate salt (generally isopropylamine salt

30

but occasionally ammonium or trimethylsulfonium or other salts) and the common adjuvants are ammonium sulphate (AMS) and/or non-ionic surfactant.

5 The discussion of documents, acts, materials, devices, articles and the like is included in this specification solely for the purpose of providing a context for the present invention. It is not suggested or represented that any or all of these matters formed part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed before the priority date of each claim of this application.

10

There is a need for improved glyphosate formulation systems and methods which allow the active to be transported in a highly concentrated form without the requirement or cost of specialist surfactants and additives.

15 **Summary**

A method of preparing a spray tank mix of glyphosate comprising

- (a) providing a glyphosate acid concentrate;
 - (b) providing an alkaline composition; and
 - (c) forming a dilute mixture of glyphosate acid concentrate and alkaline
- 20 composition in the spray tank by addition of a composition comprising the glyphosate acid concentrate to a diluted aqueous mixture of the alkaline composition. The composition may be directly applied to plants from the spray tank.

25 The glyphosate acid concentrate is preferably a solid particulate material which typically contains at least 50% by weight glyphosate acid (preferably at least 50% and most preferably at least 95% glyphosate acid). This significantly reduces the expense and problems of transport and handling.

30 An aqueous dispersion of the glyphosate acid concentrate is generally mixed with an aqueous mixture containing the alkaline composition.

In a further embodiment the invention provides a glyphosate formulation system for preparing a glyphosate spray tank mixture the system comprising a pack comprising a glyphosate acid concentrate composition for addition to a spray tank and a pack comprising a water soluble alkaline for reaction with the glyphosate acid in the spray tank to form a water soluble salt thereof *in situ*.

It is particularly preferred to add the glyphosate acid as a dispersion formed from the glyphosate acid concentrate to the alkaline composition which has been dispersed in water in the spray tank.

Throughout the description and the claims of this specification the word "comprise" and variations of the word, such as "comprising" and "comprises" is not intended to exclude other additives, components, integers or steps.

15

Detailed Description

One aspect of the invention provides a method of preparing a spray tank mix of glyphosate comprising

- (a) providing a glyphosate acid concentrate;
- 20 (b) providing an alkaline composition; and
- (c) forming a mixture of glyphosate acid concentrate and alkaline composition in the spray tank by addition of a composition comprising the glyphosate acid concentrate to a diluted aqueous mixture of the alkaline composition.

25 The invention also provides a method of controlling vegetation comprising the additional step of applying the resulting mixture to plants from the spray tank. The process generally involves dilution of at least one of the components with water to form a dilute composition suitable for application to plants.

30 In one embodiment the invention provides a method of preparing a spray tank mix of glyphosate comprising:

- (a) providing a glyphosate acid concentrate;
- (b) providing an alkaline composition;
- (c) diluting the alkaline composition with water in the spray tank to provide a diluted alkaline mixture;
- 5 (d) adding a composition comprising the glyphosate acid concentrate, optionally diluted with water, to the dilute alkaline mixture; and
- (e) optionally adding further water to the spray tank to provide the desired volume.

10 **Glyphosate Acid Concentrate**

The invention utilises glyphosate acid which term is used herein to mean N-phosphonomethyl glycine in the form of the free acid as distinguished from the highly water soluble salts such as the isopropyl amine, ammonium and potassium salts which are presently used in glyphosate compositions.

15

The invention is concerned with preparation of a spray tank mix. Spray tank mixes are prepared in the industry to provide dilution of a herbicidal concentrate, which is sold for farm use, to provide an aqueous mixture having the concentration of pesticide required for application at the site of use by spraying.

20 The concentrate is combined with a diluent, generally water, shortly before use and may be mixed with adjuvants to enhance performance. In contrast to conventional glyphosate concentrates which are based on a salt of glyphosate the present invention utilises a concentrate in the form of glyphosate acid. The concentrate preferably contains glyphosate in the form of particles or granules.

25 The particles may be in the form of an aqueous dispersion or may be adapted to be agitated to provide an aqueous dispersion. Alternatively and more preferably the particulate glyphosate is in the form of dry solid particles.

While a wide range of particle sizes may be used the glyphosate acid
30 concentrate is preferably in the form of a finely divided solid comprising particles

of size in the range of from 50 microns to 5 mm, preferably 0.01 mm to 2 mm and most preferably 0.05 mm to 1.5 mm.

The glyphosate acid concentrate may comprise fillers, additives or adjuvants if desired. It is preferred that the glyphosate concentrate is a particulate material comprising at least 50% by weight glyphosate acid, more preferably the glyphosate acid concentrate comprises at least 80% by weight glyphosate acid, still more preferably at least 95% glyphosate acid and most preferably the glyphosate acid concentrate consists essentially of glyphosate acid.

The glyphosate acid concentrate is preferably dry although it will be understood that dry particles may contain some residual water. Accordingly the glyphosate acid concentrate is typically in the form of particles of than 25% preferably less than 20% more preferably less than 15% still more preferably less than 10% more preferably less than 8% by weight water and most preferably less than 5% water.

The invention utilises an alkaline component which preferably forms a water soluble salt with glyphosate on mixing with the glyphosate acid.


The alkaline composition may contain one or more of a wide range of alkaline materials. It is particularly preferred that the alkaline composition contain at least one material selected from the group of alkaline materials registered for use with glyphosate, for example alkali metal bases such as sodium hydroxide and potassium hydroxide, alkananolamines such as monoethanolamine, diethanolamine, triethanolamine and monoisopropanolamine, alkylamine alkoxylates particularly fatty amine ethoxylates such as tallowamine ethoxylate and amines such as alkyl amines including isopropylamines. The most preferred alkaline materials are selected from mono and dialkanolamines and alkyl alkanolamines. Specific examples of preferred alkaline materials include

monoethanolamine, isopropanolamine, diethanolamine, triethanolamine and mixtures thereof.

The alkaline composition may, and preferably will contain an adjuvant particularly a surfactant. The preferred surfactants are those which enhance spray performance and or the activity of glyphosate. The surfactant may be of a type registered for use with glyphosate. It is preferred that the alkaline composition comprises a surfactant chosen from the group consisting of amidoaliphaticbetaines, alkylamine alkoxyates particularly fatty amine ethoxyates such as tallowamine ethoxyates, cocobetaine, cocoamidopropyl betaines, alkoxyated aliphaticamines, alkylpolyglucosides, alkylglycosides, aliphaticamines quaternary amines, aliphatic acetates, ethylene diamine ethoxyates, amphoteric surfactants such as polyoxyethylene aliphaticamine oxides and polyoxyethylene aliphatic ether salts, amine oxides and amphotoacetates. The term aliphatic is used to include saturated as well as unsaturated hydrocarbon chains, and includes linear and branched chains. Glyphosate-synergising surfactants typically contain at least one aliphatic group containing 8 to 22, more frequently 12 to 18 carbon atoms. Discussions of glyphosate-synergising surfactants are provided in US 6207617, WO 95/16351, US Pub 20030158042, US Pub 20050170965 and US Pub 20040224846.

The preferred surfactants are selected from aliphatic, betaines amidoaliphaticbetaines and tallow amine ethoxyates.


Betaine surfactants are generally of formula I:



Wherein: R is the aliphatic group of from 1 to 30 carbon atoms or aliphatic amidoalkyl and preferably containing more than 13 carbon atoms; R^1 and R^2 are each, independently, hydrogen C_1 to C_6 alkyl or C_1 to C_6 hydroxyalkyl and

preferably is an ethyl, hydroxyethyl or most preferably a methyl group; m is 1 or 2; X is a phosphono, sulphono, or, preferably, carboxy group; and n is 0 or, preferably 1.

- 5 R preferably has from 14 to 25 carbon atoms and is desirably a straight chain alkyl or alkyl amido alkylene group, especially a group of the formula $\text{CH}_3(\text{CH}_2)_a\text{CONH}(\text{CH}_2)_b$ where (a+b) is from 12 to 23 and b is preferably 2 or most preferably 3.
- 10 In one embodiment, the betaine surfactant is a compound according to formula II preferably of formula IIa



wherein:


- 15 R is alkyl, alkenyl, or alkylamidoalkyl, and R^1 and R^2 are each independently (C_1 to C_6)alkyl or hydroxy(C_1 to C_6)alkyl.

As used herein, "alkyl" means a saturated straight or branched chain hydrocarbon radical, typically a (C_1 - C_{30}) saturated straight or branched chain hydrocarbon radical, such as for example selected from the group consisting of methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, t-butyl, pentyl, n-hexyl, cyclohexyl, octyl, decyl, dodecyl, tetradecyl, hexadecyl, octadecyl, eicosyl, behenyl and tricosyl.

- 25 As used herein, the term "alkenyl" means an unsaturated straight chain, branched chain, or cyclic hydrocarbon radical having at least one carbon-carbon

double bond per radical, preferably having from C₂ to C₃₀ alkenyl such as for example, propenyl, butenyl, octadecenyl.

As used herein, "alkylamidoalkyl" means a group according to formula III:



wherein R⁴ is alkyl or alkenyl, typically (C₁ to C₃₀)alkyl, and R⁵ is an alkylenyl radical, typically (C₁ to C₃₀)alkylenyl and includes, for example, dodecylamidopropyl, cocoamidopropyl and tetradecylamidoethyl.

10 As used herein, "hydroxy(C₁ to C₆)alkyl" means a hydroxyalkyl group having from 1 to 6 carbon atoms per group, such as for example hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, hydroxypentyl, and hydroxyhexyl. Suitable betaine surfactants include for example, (C₁₂ to C₁₈) alkyldimethyl betaine, cocoamidopropyl betaine, and mixtures thereof.

15 In one embodiment, the betaine surfactant comprises a (C₁₂ to C₁₆) alkyldimethylbetaine.


20 A particularly preferred surfactant component for use in the alkaline composition comprises an alkylbetaine surfactant, preferably cocobetaine.

Other surfactants which may be used include ethoxylated alkyl amines and alkyl amines.

25 The alkyl amines (also referred to as fatty amines) or ethoxylated alkyl/fatty amines can comprise at least one hydrocarbon group containing 2 to 24 carbon atoms, optionally polyalkoxylated.

The fatty amines or alkoxylated fatty amines can more particularly be selected from amines comprising at least one linear or branched, saturated or unsaturated group containing 2 to 24 carbon atoms, preferably 8 to 18 carbon atoms, optionally comprising 2 to 30 oxyethylene groups and or polypropylene groups,
 5 or a mixture of a plurality thereof. Examples include ethoxylated tallow amines.

The fatty amines or ethoxylated fatty amines can be selected from ethoxylated fatty amines comprising at least one linear or branched, saturated or unsaturated groups containing 6 to 24 carbon atoms, preferably 8 to 20 carbon atoms,
 10 comprising 2 to 30 oxyethylene groups, or a mixture of a plurality thereof. Examples include the compounds having the following formula IV:



wherein R^6 represents a linear or branched, saturated or unsaturated hydrocarbon group containing 6 to 24 carbon atoms, preferably 8 to 20 carbon
 15 atoms; OA represents an oxypropylene group; and q, q^1 , which may or may not be identical, represent a mean number in the range 1 to 30. Examples of such amines that can be cited are amines derived from copra and containing 5 oxyethylene (OE) motifs, oleic amines containing 5 OE, amines derived from tallow containing 5-20 OE, for example 10, compounds corresponding to the
 20 above formula, in which R^6 is an alkyl group containing 12 to 15 carbon atoms, the number of OE motifs being in the range 20 to 30.

Examples of alkyl amine and ethoxylated alkyl amine surfactants include amine oxide surfactants such as:

- 25
- an amine oxide having formula $\text{R}^7\text{R}^8\text{R}^8\text{N} \rightarrow \text{O}$ (V),
 - an amine oxide having formula $\text{R}^7\text{-CO-NH-R}^9\text{-R}^8\text{R}^8\text{N} \rightarrow \text{O}$ (IV), or
 - a mixture or association thereof,

wherein:

R^7 which is identical or different, is a linear or branched hydrocarbon group, preferably an alkyl group containing 3 to 30 carbon atoms, preferably 3 to 20 carbon atoms,

5 R^8 , which is identical or different, is a C_1 - C_3 alkyl group, preferably a methyl group, and

R^9 , which is identical or different, is divalent linear or branched hydrocarbon group containing 1 to 6 carbon atoms, optionally substituted with a hydroxyl group, preferably a group of formula $-CH_2-CH_2-CH_2-$ or $-CH_2-CHOH-CH_2-$.

10 The alkaline composition preferably has a pH (1% solution in water) in the range of from 8 to 13.5.

The alkaline composition may if desired contain other components such as
15 adjuvants (for example crop oils, wetters, stickers, foaming agents and marking agents) and other active agents. Examples of complementary active agents which may be used include at least one selected from the group consisting of fungicides, insecticides and herbicides. Specific examples of additional active agents which may be present in the alkaline composition may comprise at least
20 one selected from 2,4D, amitrole, ammonium sulphate, atrazine, carfentrazone, chlorsulfuron, dicamba, diuron, MCPA, metolachlor, metribuzin, metsulfuron, oryzalin, oxyflourfen, pendimethalin, picloram, simazine, terbutryne, triallate, triasulfuron, triclopyr and trifluralin.

25 The method of the invention involves addition of glyphosate acid to a dilute alkaline composition in a spray tank. Generally the alkaline composition will be diluted with water in the spray tank before the addition of glyphosate acid to the diluted alkaline composition. The glyphosate acid may be added as a concentrate or alternatively the glyphosate acid concentrate is dispersed in water
30 prior to mixing with the alkaline composition in the spray tank.

In this embodiment the aqueous dispersion of the glyphosate acid may be formed from the glyphosate acid concentrate in a separate vessel and then added to the tank or it may be formed in or during addition to the spray tank. In general the glyphosate will be mixed with water and this may be achieved by stirring or agitating the glyphosate acid concentrate in water or by the impact of a stream of water on solid glyphosate acid concentrate.

The aqueous dispersion of glyphosate need not be a particularly stable dispersion as it may be added to the dilute alkaline composition within the spray tank within a short period. For example, the dispersion may be added immediately after it is formed to the diluted alkaline composition. A dispersion in water may therefore be suitable if it would otherwise settle within a short period of say 1 minute. Indeed the dispersion may be maintained entirely by the turbulence or mixing prior to addition to the spray tank.

An aqueous dispersion may be achieved by adding the glyphosate acid concentrate to a bucket of water and stirring with a stick, paddle or agitator, or by adding the glyphosate acid concentrate to a mixing vessel for vigorous agitation with water. The mixing vessel may provide mixing using a vortex of the diluent (typically water). A suitable mixing vessel for vortex mixing is available from Hardi Industries under the trade mark "Granni Pot" Chemical Indicator or from Gold Acres of St Arnaud, Victoria, Australia under the name of a Chemical Induction Hopper. The aqueous dispersion may also be achieved by adding solid particulate glyphosate acid concentrate to a perforated container such as a filter container located on or over the spray tank, and flushing the glyphosate through the perforations with a charge of water directed into the perforated container. In a particularly preferred embodiment the method of the invention involves placing solid particulate glyphosate acid concentrate in a container having perforations and dispersing the glyphosate acid in water by impacting the concentrate with a stream of water and draining the water from the perforated container into the spray tank. It is even more preferred in this embodiment that the water drains

from the perforated container into the spray tank wherein the spray tank contains the alkaline composition optionally diluted with water.

The alkaline composition is preferably diluted with water in the spray tank.

5

For example the spray tank may be charged with an initial charge of water and alkaline composition and an aqueous dispersion of the glyphosate acid then added.

10 Accordingly in a particularly preferred embodiment of the invention we provide a method of preparing a spray tank composition of glyphosate comprising:

- (a) providing an initial charge of water to the spray tank, which initial charge of water may, for example, provide $\frac{1}{4}$ to $\frac{3}{4}$ of the total spray water volume and preferably approximately $\frac{1}{2}$ of the total spray water volume;
- 15 (b) adding the alkali composition to the initial charge of water to provide a diluted alkaline composition;
- (c) dispersing solid particulate glyphosate acid concentrate in water;
- (d) mixing the aqueous dispersion of glyphosate acid with the diluted alkaline composition; and
- 20 (e) optionally adding more water to provide the desired volume.

The aqueous dispersion of glyphosate acid is preferably added to the alkaline composition (which is preferably diluted with water) having a pH of in the range of 7.5 to 10.5.

25

The volume of diluted herbicidal composition formed in the spray tank in accordance with the method of the invention is preferably in the range of from 50 to 20,000 litres, more preferably from 100 to 10,000 litres and most preferably from 400 to 5,000 litres.

30

The pH of the spray liquor after the combination of the compositions and dilution with water is preferably in the range 3.5 to 7 and more preferably from 4 to 6.0.

5 In the case of some amines including alkyl amines such as fatty amines (eg tallowamines) there is an advantage in working at lower pH values of for example from 2 to 5.5 and most preferably at about 2.

10 The process of the invention may further comprise the steps of preparing the glyphosate acid concentrate composition, transporting the glyphosate acid concentrate composition and forming a spray tank composition as hereinabove described. The spray tank composition will typically be prepared within several days of use and typically within a day of use whereas the glyphosate acid concentrate is stable for many months of storage and transport and allows diluents and solvents to be significantly reduced when compared with
15 conventional glyphosate (salt type) concentrates.

The concentration of glyphosate (acid equivalent) in the final spray tank composition is preferably in the range 0.01% to 9.0% by weight, more preferably in the range 0.4 to 4% by weight and most preferably in the range 0.4 to 3% by
20 weight.

The two pack system in accordance with the invention comprises a first pack containing the alkaline composition and a second pack containing the glyphosate concentrate as hereinbefore described. The two packs contain amounts of the
25 respective glyphosate and alkaline component to provide at least 80% (based on glyphosate acid) of the glyphosate as water soluble glyphosate salt.

The two pack system may comprise instructions for mixing the packs of the system in accordance with the above described method steps.

In one preference, the ratio of glyphosate-synergising surfactant (in the first pack) to glyphosate (acid equivalent) in the second pack is in the range of from 1:1 to 1:20 for the active component of the adjuvant and more preferably 1:1 to 1:14. Typically the active surfactant component of the adjuvant is present in a diluent
5 so that the ratio of formulated glyphosate-synergising surfactant to glyphosate is typically from 1:7 to 1:1 and preferably 1:3 to 1:4.

In one preference the first pack comprises alkaline material and glyphosate synergising surfactant.
10

Preferably the range of water in the first pack is less than 70% and preferably less than 50%.

In one preference the glyphosate synergising surfactant has a low aquatic animal
15 toxicity.

In another preference the glyphosate synergising surfactant is an ethoxylated tallow amine.

20 Preferably the pH of liquor in the first pack is less than 13 and more preferably less than 12.

In one preference the alkaline material is monoethanolamine.

25 The contents of the two packs are typically combined in accordance with the above-described method.

Typically the molar ratio of glyphosate acid to alkaline component of the alkaline composition is in the range of from 6:1:1:2, preferably 0.8:1 to 1:1.2 and more
30 preferably from 1:1.05 to 1:1.2.

In the case of alkyl amines including fatty amines such as tallowamine we have surprisingly found that a ratio as low as from 6:1 to 1:1.2, (such as from 6:1 to 1:1 or 1:1 to 1.2:1) is sufficient to solubilise the glyphosate and provide a useful product of high activity.

5

The invention will now be described with reference to a number of examples which are provided for the purpose of further understanding embodiments of the invention but are not intended to limit the scope or applicability of the invention to the specific examples.

10

Examples

Compositions and methods used in the examples, in particular the results of efficacy trials are described with reference to the attached drawings.

15 In the drawings:

Figure 1 is a graph comparing the efficacy on ryegrass of a two pack glyphosate system of the invention of Example 1 with a commercial glyphosate herbicide formulation at various application rates.

20

Figure 2 is a graph comparing the efficacy on oats of a two pack glyphosate system of the invention of Example 2 with a commercial glyphosate herbicidal formulation at various application rates.

25

Figure 3 is a graph comparing the field efficacy on ryegrass of a two pack glyphosate system of the invention with a commercial glyphosate formulation at various application rates as described in Example 3.

Example 1

30

This example compares the bioefficacy of a 2 Pack glyphosate herbicide of the invention with a commercial glyphosate standard.

METHOD

Plant propagation

Annual ryegrass (*Lolium rigidum*) seeds (5/pot) were sown to a depth of 10mm in 10cm diameter pots filled with potting mix (Australian Standard 3743) that had been amended with macro and micronutrients for optimal growth.

One week after seedling emergence, seedlings were thinned for uniform size to one seedling per pot. Plants were grown in a temperature-controlled greenhouse (14°C – 25°C) for 14 days then outdoors for 20 days prior to spray application to more closely simulate field conditions. After the application of herbicides the pots were returned to the greenhouse for an additional 14 days before plants were assessed for fresh weight.

Preparation of 2 Packs Herbicide

Approximately 13L of water was added to each of five 20 litre ("L") plastic buckets (Table 1). Pack 1 was made by taking 37.5 parts monoethanolamine ("MEA") and 25.3 parts of a cocobetaine surfactant called Empigen™ BB from Huntsman Chemical Company Australia. Pack 1 was also called the Alkaline – Surfactant Blend in this and later examples. Pack 1 was added to the water at the rates given in Table 1.

Glyphosate acid technical material with a purity of 95% and a particle size smaller than 1 mm (Pack 2) was then added to the buckets at the rates given in Table 1 and stirred with a plastic paddle until all the acid disappeared. This took 3-5 minutes at the higher concentrations.

1.3L of each mix was then transferred to a 6L spray canister and pressurized using compressed air.

Table 1. Mixing ratios of Alkali Surfactant Blend (Pack 1), glyphosate acid (Pack 2) and water and equivalent rate of glyphosate applied per hectare. NB ai = active ingredient measured as the pure acid.

Glyphosate Rate g ai/ha	Glyphosate concentration g ai/L	Pack 1 – Alkali-Surfactant blend ml/bucket	Pack 2- Glyphosate acid g/bucket	Water L/bucket
45	0.70	5.79	9.62	12.99
90	1.41	11.58	19.24	12.99
180	2.81	23.17	38.49	12.98
360	5.63	46.34	76.97	12.95
450	7.03	57.92	96.22	12.94

5 Commercial Herbicide

A commercial glyphosate formulation Roundup Biactive™ was used as the commercial standard. Roundup Biactive™ herbicide is based on an isopropylamine (IPA) salt and contains a surfactant.

- 10 The commercial standard herbicide was added directly to the water in the spray canister at the required rate (Table 2) to match the concentration of pure glyphosate acid in the 2 Pack system described previously. The commercial standard was not mixed in a bucket.

15 Table 2. Mixing ratios of Roundup Biactive and water.

Glyphosate Rate g ai/ha	Glyphosate concentration g ai/L	Roundup Biactive™ concentration g/L	Roundup Biactive™ ml/canister	Water L/canister
45	0.70	360	2.54	1297.5
90	1.41	360	5.08	1294.9
180	2.81	360	10.16	1289.8
360	5.63	360	20.32	1279.7
450	7.03	360	25.39	1274.6

Herbicide application

The annual ryegrass was at the early tillering stage (2 tillers) when sprayed with the herbicide treatments.

- 5 Herbicide formulations were applied using an enclosed laboratory track-sprayer fitted with three 110° flat fan nozzles ("Teejet"® XR11001-VS) spaced at 50cm intervals across the boom. The boom moved along a fixed track at 6 km h⁻¹, sprayed at a water volume of 64 L ha⁻¹ with a pressure of 200 kPa.
- 10 There were eight replicates for each treatment.

Assessment

Seedlings were harvested 14 days after spray application by cutting foliage off at the base immediately prior to weighing on an "AND FX" 300 electronic balance (range 0-300 g).

15 **Statistical analysis**

Data was analysed using a factorial design with two factors, glyphosate formulation (Formulation) and spray application rate (Rate). 95% least significant differences (LSD) were calculated for the mean of each treatment.

20 **RESULTS**

- Averaged across all rates there was no significant difference in the fresh weight of ryegrass treated with either of the formulations indicating that both formulations were of equal efficacy (Table 3). As expected there was a highly significant effect of application rate. There was no interaction between Rate and
- 25 Formulation.

Table 3. Statistic analysis data for fresh weight of annual ryegrass plants at harvest. NS = not significant (P<0.05). LSD = least significant difference (P=0.05)

FACTOR	Prob (F)	LSD
Formulation	0.71	NS
Rate	0.0001	0.43 g/plant
Formulation x Rate	0.10	NS

Raw data and means for fresh weight of annual ryegrass at harvest are tabled (Table 4) and graphed (Figure 1).

- 5 **Table 4.** Fresh weight (g/plant) of plants harvested after treatment with two glyphosate formulations. Mean Rate data in the same column or Mean Formulation data in the same row that are followed by the same letter are not significantly different ($P < 0.05$).

Glyphosate application rate g ai/ha	2 Pack System	Roundup Biactive™	Rate Mean
0	14.1	14.1	
45	12.9	12.4	12.6 e
90	9.77	10.1	9.94 d
180	5.97	6.36	6.17 c
360	2.55	2.00	2.28 b
450	1.76	1.71	1.74 a
Formulation Mean (not including the 0g ai/ha rate)	6.58 a	6.53 a	

10

Example 2

This example compares the herbicidal activity of a 2 Pack glyphosate system in accordance with the invention with commercial glyphosate standards on canola and oats.

15 METHOD

Plant propagation

Oat (*Avena sativa*) and canola (*Brassica napus*) seeds (3/pot) were sown to a depth of 15mm for the oats and 3mm for the canola in 10cm diameter pots filled

with potting mix (Australian Standard 3743) that had been amended with macro and micronutrients for optimal growth.

One week after seedling emergence, seedlings were thinned for uniform size to one seedling per pot. Plants were grown in a temperature-controlled greenhouse (14°C – 25°C) for 14 days then outdoors for 10 days prior to spray application to more closely simulate field conditions. After the application of herbicides the pots were returned to the greenhouse for an additional 14 days before plants were assessed for fresh weight.

2 Pack herbicide preparation

Approximately 13L of water was added to seven 20L plastic buckets (Table 5). A blend of 37.5 parts MEA and 25.3 parts of the cocobetain surfactant Empigen BB was made to form the Alkali Surfactant Blend (Pack 1). Pack 1 was then added to the water at the rates given in Table 5. Glyphosate acid (Pack 2) with a purity of 95% was then added to the buckets at the rates given in Table 5 and stirred with a plastic paddle until all the acid disappeared. This took 3-5 minutes at the higher concentrations.

1.3L of each mix was then transferred to a 6L spray canister and pressurized using compressed air.

Table 5. Mixing ratios of Alkali Surfactant Blend (Pack 1), glyphosate acid (Pack 2) and water at an equivalent rate of glyphosate applied per hectare. NB ai = active ingredient measured as the pure acid.

Glyphosate Rate g ai/ha	Glyphosate concentration g ai/L	Pack 1 – Alkali-Surfactant Blend ml/bucket	Pack 2- Glyphosate acid g/bucket	Water L/bucket
30	0.47	3.86	6.41	13.00
60	0.94	7.72	12.83	12.99
90	1.41	11.58	19.24	12.99
120	1.88	15.45	25.66	12.98
180	2.81	23.17	38.49	12.98
240	3.75	30.89	51.32	12.97
450	7.03	57.92	96.22	12.94

Commercial Herbicides

- 5 Two commercial glyphosate formulations Roundup Biactive™ brand and Macphersons® Glyphosate 450 SL brand were used as commercial standards. Macphersons® Glyphosate 450SL is based on the MEA salt. Roundup Biactive™ is based on an IPA salt.
- 10 The commercial standard herbicides were added directly to the water in the spray canister at the required rate to match the concentration of pure glyphosate acid in the 2 Pack system described above (Table 6 and 7). The commercial standards were not mixed in a bucket.

15 “Roundup Biactive”™

Table 6. Mixing ratios of “Roundup Biactive”™ and water.

Glyphosate Rate g ai/ha	Glyphosate concentration g ai/L	Roundup Biactive™ concentration g/L	Roundup Biactive™ ml/canister	Water L/canister
30	0.47	360	1.69	1298.3
60	0.94	360	3.39	1296.6
90	1.41	360	5.08	1294.9
120	1.88	360	6.77	1293.2
180	2.81	360	10.16	1289.8
240	3.75	360	13.54	1286.5
450	7.03	360	25.39	1274.6

"Macphersons"[®] Glyphosate 450 SL**Table 7.** Mixing ratios of "Macphersons"[®] Glyphosate 450SL and water.

Glyphosate Rate g ai/ha	Glyphosate concentration g ai/L	Macphersons[®] Glyphosate 450SL concentration g/L	Macphersons[®] Glyphosate 450SL ml/canister	Water L/canister
30	0.47	450	1.35	1298.6
60	0.94	450	2.71	1297.3
90	1.41	450	4.06	1295.9
120	1.88	450	5.42	1294.6
180	2.81	450	8.13	1291.9
240	3.75	360	13.54	1286.5
450	7.03	360	25.39	1274.6

5 Herbicide application

Both the oats and the canola were at the three leaf stage when sprayed with the herbicide treatments.

Herbicide formulations were applied using an enclosed laboratory track-sprayer fitted with three 110° flat fan nozzles ("Teejet"[®] XR11001-VS brand) spaced at 50cm intervals across the boom. The boom moved along a fixed track at 7.7 km h⁻¹, sprayed at a water volume of 50 L ha⁻¹ with a pressure of 200 kPa.

There were 8 replicates for each treatments.

15 Assessment

Seedlings were harvested 14 days after spray application by cutting foliage off at base immediately prior to weighing on an "AND FX" 300 electronic balance (range 0-300 g).

20 Statistical analysis

Data was analysed using a factorial design with two factors, glyphosate formulation (Formulation) and spray application rate (Rate). 95% least significant differences (LSD) were calculated for the mean of each treatment.

5 RESULTS

Oats

When the fresh weight of the oats was averaged across all rates there was no significant difference between the formulations indicating that the three formulations had equal efficacy (Table 8). As expected there was a highly significant effect of application rate. There was no interaction between Rate and Formulation.

Table 8. Statistic analysis data for fresh weight of oat plants at harvest. NS = not significant ($P < 0.05$).

FACTOR	Prob (F)	LSD
Formulation	0.39	NS
Rate	0.0001	0.39 g/plant
Formulation x Rate	0.99	NS

Raw data for fresh weight and means for oats at harvest are tabled (Table 9) and graphed in (Figure 2).

Table 9. Fresh weight (g/plant) of oat plants harvested after treatment with various glyphosate formulations. Mean Rate data in the same column or Mean Formulation data in the same row that are followed by the same letter are not significantly different ($P < 0.05$).

Glyphosate application rate g ai/ha	2 Pack System	MacPhersons 450SL	Roundup Biactive™	Rate Mean
0	6.11	6.11	6.11	
30	5.61	5.43	5.52	5.52 e
60	4.26	4.17	4.48	4.30 d
90	2.60	2.66	2.94	2.73 c
120	2.02	1.76	1.95	1.91 b
180	1.19	1.14	0.90	1.08 a
240	1.08	0.79	0.94	0.94 a
450	0.73	0.61	0.95	0.76 a
Formulation Mean (not including the 0g ai/ha rate)	2.50 a	2.36 a	2.53 a	

Canola

When the fresh weight of the canola was averaged across all rates there was no significant difference between the formulations indicating that the three formulations had equal efficacy (Table 10). As seen with the oats there was a highly significant effect of application rate. There was no interaction between Rate and Formulation.

Table 10. Statistic analysis data for fresh weight of canola plants at harvest. NS = not significant ($P < 0.05$).

FACTOR	Prob (F)	LSD
Formulation	0.16	NS
Rate	0.0001	0.29 g/plant
Formulation x Rate	0.88	NS

Raw data for fresh weight and means for canola at harvest are tabled (Table 11).

5 **Table 11.** Fresh weight (g/plant) of oat plants harvested after treatment with various glyphosate formulations. Mean Rate data in the same column or Mean Formulation data in the same row that are followed by the same letter are not significantly different ($P < 0.05$).

Glyphosate application rate g ai/ha	2 Pack System	MacPhersons 450SL	Roundup Biactive™	Rate Mean
0	7.22	7.22	7.22	
30	5.09	4.81	4.99	4.96 e
60	3.13	2.74	2.69	2.86 d
90	1.43	1.17	1.37	1.32 c
120	1.15	0.92	0.67	0.91 b
180	0.53	0.68	0.57	0.59 a
240	0.54	0.52	0.48	0.51 a
450	0.47	0.46	0.42	0.45 a
Formulation Mean (not including the 0g ai/ha rate)	1.76 a	1.61 a	1.60 a	

Example 3

10 This example compares the field bioefficacy of a 2 Pack glyphosate herbicide made according to the invention with a commercial glyphosate standard on annual ryegrass.

METHOD

Site Details

15 A field trial site was selected for uniformity of annual ryegrass (*Lolium rigidum*) near Werribee, Victoria, Australia. The weeds were at the late tillering stage (Zadocks GS 25-29) and were growing vigorously. Soil moisture conditions were optimum for plant growth.

The site was sprayed on the 6th September 2005, when wind conditions were slight (<5kph).

Plots were 2m x 15m with a 0.5m buffer between each plot.

5

Mixing Tank Details

The mixing tank was a 200L, wide mouth (mouth was 20cm diameter) plastic tank with a diameter of approximately 65cm and height 60cm.

10

2 Pack herbicide preparation

70L of water was added to the 200L tank. A blend was made of 37 parts MEA and 25 parts of the cocobetaine surfactant Empigen BB to form the Alkali Surfactant Blend (Pack 1). The Alkali Surfactant Blend was then added to the water at one of the rates given in Table 12. A plastic mesh container insert (with
15 holes approximately 4mm diameter) was then placed into the mouth of the container and the corresponding amount of glyphosate acid (Pack 2) for the previously added Alkali Surfactant Blend (Table 12) was then added onto the mesh and washed into the container using a steady stream of water from a hose
20 unto the total volume in the spray tank was 130L. The tank had no agitation other than that caused by the action of the water stream leaving the hose while the additional 60L of water was added.

A small plot experiment was done using a 1.3L sample of the solution taken from the 200L tank above. This sample was poured into a 6L pressurised spray
25 canister. After the 1.3L of solution was transferred to the 6L spray canister the canister was pressurized using compressed air.

This process was repeated for each of the spray application rates given in Table 12.

Table 12. Mixing ratios of Alkali Surfactant Blend (Pack 1), glyphosate acid (Pack 2) and water and equivalent rate of glyphosate applied per hectare. NB ai = active ingredient measured as the pure acid.

5

Glyphosate Rate g ai/ha	Glyphosate concentration g ai/L	Pack 1 – Alkali-Surfactant Blend ml/spray tank	Pack 2- Glyphosate acid g/spray tank	Water L/spray tank
45	0.70	57.9	96.2	129.9
90	1.41	115.8	192.4	129.9
180	2.81	231.7	384.9	129.8
360	5.63	463.4	769.7	129.5
450	7.03	579.2	962.2	129.4

Commercial Herbicide

A commercial glyphosate formulation Roundup Biactive™ was used as the commercial standard.

- 10 The commercial standard herbicide was added directly to the water in the spray canister at the required rate to match the concentration of pure glyphosate acid in the 2 Pack system described above. The commercial standard was not pre-mixed in a larger container (Table 13).

15 “Roundup Biactive”™

Table 13. Mixing ratios of “Roundup Biactive”™ and water.

Glyphosate Rate g ai/ha	Glyphosate concentration g ai/L	Roundup Biactive™ concentration g/L	Roundup Biactive™ ml/canister	Water L/canister
45	0.71	360	2.54	1297.5
90	1.41	360	5.08	1294.9
180	2.81	360	10.16	1289.8
360	5.62	360	20.32	1279.7
450	7.03	360	25.39	1274.6

Herbicide application

Herbicide formulations were applied using a hand held boom fitted with three 110° flat fan nozzles ("Teejet"® XR11001-VS) spaced at 50cm intervals across the boom. The boom was moved along the plots at 6 km h⁻¹ and sprayed at a water volume of 64 L ha⁻¹ with a pressure of 200 kPa.

There were five replicates for each treatment.

Assessment

Weed counts were taken using 10 x 0.1m² quadrants per replicate, 28 days after treatment.

Statistical analysis

Data was analysed using a factorial design with two factors, glyphosate formulation (Formulation) and spray application rate (Rate). 95% least significant differences (LSD) were calculated for the mean of each treatment.

RESULTS

Averaged across all rates there was no significant difference in the density of annual ryegrass between the formulations indicating that the two formulations had equal efficacy (Table 14). There was a highly significant effect of application rate. There was no interaction between Rate and Formulation.

Table 14. Statistic analysis data for the density of annual ryegrass plants. NS = not significant (P<0.05).

FACTOR	Prob (F)	LSD
Formulation	0.52	NS
Rate	0.0001	5.8 plants /m2
Formulation x Rate	0.68	NS

Raw data for the weed counts is tabled (Table 15) and graphed in (Figure 3).

Table 15. Annual ryegrass density (plant/m²) after treatment with two glyphosate formulations. Mean Rate data in the same column or Mean Formulation data in the same row that are followed by the same letter are not significantly different (P<0.05).

Glyphosate application rate g ai/ha	2 Pack System	Roundup Biactive™	Rate Mean
0	86.2	86.2	
45	77.4	71.6	74.5 d
90	24.2	26.8	25.5 c
180	9.6	8.8	9.2 b
360	5.8	3.8	4.8 ab
450	0.6	0.8	0.7 a
Formulation Mean (not including 0g ai/ha)	23.5 a	22.4 a	

5

Example 4

This example trials the mixing of a two pack system of the invention in a 1000L commercial spray tank without plastic mesh insert.

10 **METHOD**

Spray Tank Details

A commercial "Silvan" Trailing Boom Spray fitted with a 1000L tank was used for this experiment. The boom was 10m wide.

15 **Glyphosate Technical Material**

The glyphosate material used was glyphosate acid (purity 95%) with a particle size where 95% can pass through a 1.2mm sieve.

Addition

20 The spray tank was filled with 900L of water. The recirculation or agitation system was driven by a power take off ("PTO") from the tractor. The recirculation system consisted of a wet boom, submerged at the base of the tank with outlets

to one side of the tank's midline. This resulted in approximately 75% of the base of the tank on the same side as the jet outlets being swept by water. This is acceptable agitation by commercial spray boom standards.

- 5 With the PTO operating, a blend made from 0.979kg of the surfactant Empigen BB and 1.57kg of the monoethanolamine was poured into the tank and allowed to disperse for one minute.

- 10 4.26kg of glyphosate acid was then tipped into the water. The glyphosate acid could be seen stuck to the base and sides of the tank and was not removed by the agitation of the water.

Five minutes after the addition of the glyphosate there was still some glyphosate acid adhering to surfaces but almost all disappeared over the next 5 minutes.

15

Although the time taken to completely dissolve was commercially feasible a product would ideally have a shorter dissolution time, ideally 3-5 minutes.

Example 5

- 20 This example describes a process of mixing a two pack glyphosate in a system in accordance with the invention, where one pack is a premilled glyphosate acid (particle size less than 0.5mm), in a 1000L commercial spray tank without plastic mesh insert.

25 METHOD

Spray Tank Details

A commercial "Silvan" brand Trailing Boom Spray fitted with a 1000L tank was use for this experiment. The boom was 10m wide.

30

Glyphosate Technical Material

Glyphosate acid (purity 95%) with a particle size where 95% can pass through a 0.5mm sieve.

Addition

- 5 The spray tank was filled with 900L of water. The recirculation or agitation system was driven by a PTO from the tractor. The recirculation system consisted of a wet boom at the base of the tank with outlets to one side of the tank's midline. This resulted in approximately 75% of the base of the tank on the same side as the jet outlets being swept by water. This is adequate agitation by
10 commercial spray boom standards.

With the PTO operating, a blend made from 0.979kg of the surfactant Empigen BB and 1.57kg of the monoethanolamine was poured into the tank and allowed to disperse for one minute.

15

4.26kg of the milled glyphosate acid was tipped into the water. As with the unmilled glyphosate acid the milled acid could be seen stuck to the base and sides of the tank and was dissolved over time by the agitation of the water.

- 20 Five minutes after addition there was still some glyphosate acid adhering to surfaces but by 10 minutes from addition all the acid had disappeared.

Example 6

- This example describes mixing a two pack glyphosate system of the invention
25 using a plastic mesh insert in a 1000L commercial spray tank. The first pack (Alkali Surfactant Blend) contains a cocobetaine surfactant.

METHOD

30 Spray Tank Details

A commercial "Silvan" Trailing Boom Spray fitted with a 1000L tank was use for this experiment. The boom was 10m wide.

Glyphosate Technical Material

Glyphosate acid (purity 95%) with a particle size where 95% can pass through a 1.2mm sieve but not through a 0.5mm sieve.

5

Addition

The spray tank was filled with 900L of water. The recirculation or agitation system was driven by a PTO from the tractor. The recirculation system consisted of a wet boom at the base of the tank with outlets to one side of the tank's
10 midline. This resulted in approximately 75% of the base of the tank on the same side as the jet outlets being swept by water. This is adequate agitation by commercial spray boom standards.

A blend made from 0.979kg of the surfactant Empigen BB, a cocobetaine
15 surfactant, and 1.57kg of the monoethanolamine (MEA) was poured into the tank and allowed to disperse for one minute. A sample of water was collected and analysed for pH.

A plastic mesh insert with holes of 3-4mm diameter was fitted into the mouth of
20 the tank and 4.26kg of glyphosate acid was tipped into the insert. A low pressure hose was used to wash the acid through the mesh over a period of 1-2 minutes. Immediately after the mesh was empty it was removed and the water was inspected for evidence that the acid was not dissolved.

25 RESULTS

There was no evidence of undissolved acid in the tank.

A sample of water was then taken and analysed for pH. The pH of the water taken from the tap and used to fill the tank was 7.3 and this increased to 9.9 with
30 the addition of the alkali. The addition of the acid reduced this to 4.8, within the desirable range for glyphosate.

Table 16. pH of water at various stages of addition of ingredients.

	pH
Water from tap	7.3
Water after addition of Alkali Surfactant Blend	9.9
Water after addition of Alkali Surfactant Blend and Glyphosate	4.8

Example 7

- 5 This example compares field bioefficacy of a 2 Pack glyphosate herbicide in accordance with the invention with a commercial glyphosate standard on annual ryegrass, capeweed and barley grass.

METHOD**10 Site Details**

A field trial site was selected for uniformity of annual ryegrass (*Lolium rigidum*), capeweed (*Arctotheca calendula*) and barley grass (*Hordeum leporinum*) near Ballarat, Victoria, Australia. The capeweed was flowering and the barley grass and annual ryegrass were just coming out in head and all were growing
15 vigorously. Soil moisture conditions were optimum for plant growth.

The site was sprayed on the 16th December 2005 when wind conditions were slight (<5kph).

- 20 Plots were 10m x 100m with a 10m buffer between each plot. There were four replicates for each treatment.

Spray Solution

- 25 The spray solution in this example was that described in Example 6. The commercial standard was Roundup CT[®] glyphosate herbicide composition 450g glyphosate ai/L which was added directly to the tank to give a 4.47g ai/L concentration identical to that of the 2 Pack system.

The herbicide solutions were sprayed at 100L/ha to deliver 447g ai/ha.

Assessment

Plots were assessed visually for brown out seven days and 28 days after spraying. Brown out is an estimation of the proportion of plants that are dead or the proportion of an individual plant that has turned yellow or brown in response to the damage caused by the glyphosate herbicide.

RESULTS

Annual Ryegrass

The two formulations were identical in their rate of brown out or the rate at which they killed annual ryegrass (Table 17).

Table 17. Annual ryegrass mean percent brown out 7 and 28 days after spraying with 450g glyphosate acid /ha. Data in the same column and followed by the same letter and are not significantly different ($P < 0.05$).

Treatment	Brown out (%) 7 Days	Brown Out (%) 28 days
Control	0	0
2 Pack	42 a	99 a
Roundup CT	37 a	98 a
LSD ($P=0.05$)	8	1.2

Barley grass

The two formulations were identical in their rate of brown out or the rate at which they killed barley grass (Table 18).

Table 18. Barley grass mean percent brown out 7 and 28 days after spraying with 450g glyphosate acid /ha. Data in the same column and followed by the same letter are not significantly different ($P < 0.05$).

Treatment	Brown out (%) 7 Days	Brown Out (%) 28 days
Control	0	0
2 Pack	62 a	99 a
Roundup CT	57 a	99 a
LSD ($P=0.05$)	11	1.1

Capeweed

The two formulations were statistically similar in their rate of brown out or the rate at which they killed capeweed (Table 19).

Table 19. Capeweed mean percent brown out 7 and 28 days after spraying with 450g glyphosate acid /ha. Data in the same column and followed by the same letter are not significantly different ($P < 0.05$).

Treatment	Brown out (%) 7 Days	Brown Out (%) 28 days
Control	0	0
2 Pack	32 a	91 a
Roundup CT	35 a	89 a
LSD ($P=0.05$)	6	3.8

Example 8

In this example four Alkali Surfactant Blends were formulated using the proportion of ingredients given in Table 20. The surfactants in this example included a cocobetaine, an alkyl polyglycoside/EDA alkoxyate blend, an alkyl polyglycoside and a tallow amine ethoxyate. Each of these formulations was allowed to settle at room temperature for 24 hours and was then observed for

phase separation. None of these four formulations showed any sign of separation (Table 21).

Each of these Alkali Surfactant Blends was added to 800ml of water in a 1L glass container. Sufficient glyphosate acid was then added to give a typical field concentration of 4.5g/L. The ratio of Alkali Surfactant Blend to glyphosate acid was one part Alkali Surfactant Blend to 1.66 parts glyphosate acid. Tank mixes were tested for pH of resultant solution and time taken to dissolve completely (rate of dissolution). All formulations passed both of these tests, with the acid dissolving in less than 3 minutes and the pH being less than 5 (Table 21).

Table 20. Compositions of four 2 Pack glyphosate formulations (Parts on a weight basis)

	EUR25-1	EUR25-2	EUR25-3	EUR25-4
Glyphosate acid	100	100	100	100
Alkali				
MEA	37.5	-	-	37.5
KOH	-	35.6	35.6	-
Surfactant				
Empigen BB	25.3	-	-	-
Terwet 1215	-	25.3	-	-
Terwet 3780	-	-	-	25.3
Terwet 3001	-	-	25.3	-
Water	-	17.5	10.0	-

Ingredients

- Empigen BB is a cocobetaine
 Terwet 1215 is an alkyl polyglycoside/EDA alkoxyate blend
 Terwet 3001 is an alkyl polyglycoside
 Terwet 3780 is a tallow amine ethoxyate

Table 21. Performance tests on four Alkali Surfactant Blends.

	EUR25-1	EUR25-2	EUR25-3	EUR25-4
Pack 1 stability	Stable, no separation	Stable, no separation	Stable, no separation	Stable, no separation
Tank mix				
pH	4.5	4.8	4.8	4.7
Rate of dissolution	<2 minutes	< 2 minutes	< 2 minutes	<2 minutes

Example 9

This example describes a bioefficacy comparison of a 2 Pack glyphosate system of the invention, made using an alternative betaine type surfactant, with commercial glyphosate standards on annual ryegrass and wild radish

INTRODUCTION

This example relates to a system of the invention containing a betaine (specifically a cocamidopropyl betaine) in the Alkali Surfactant Blend component.

METHOD**Plant propagation**

Wild radish (*Raphanus raphanistrum*) seeds (3/pot) were sown to 2mm depth and annual ryegrass (*Lolium rigidum*) seeds (3/pot) were sown to a depth of 5mm in 10cm diameter pots filled with potting mix (Australian Standard 3743) that had been amended with macro and micronutrients for optimal growth.

One week after seedling emergence, seedlings were thinned for uniform size to one seedling per pot. Plants were grown in a temperature-controlled greenhouse (14°C – 25°C) for 18 days for annual ryegrass and 14 days for wild radish then outdoors for 22 days for annual ryegrass and 16 days for wild radish prior to spray application to more closely simulate field conditions. After the application of herbicides the pots were returned to the greenhouse for an additional 14 days before plants were assessed for fresh weight.

2 Pack herbicide preparation

In this example the monoethanolamine and cocamidopropyl betaine (also known as alkyl amidopropyl betaine) surfactant and sold as Huntsman's Empigen BS/FA surfactant blend was made using 37.5 parts monoethanolamine to 25.3 parts of cocamidopropyl betaine (aqueous concentration 30%). This blend is called the Alkali Surfactant Blend.

Approximately 13L of water was added to seven 20L plastic buckets (Table 22). The Alkali Surfactant Blend of monoethanolamine and the cocamidopropyl betaine (Pack 1) was then added to the water at the rates given in the Table. Glyphosate acid (Pack 2) with a purity of 95% was then added to the buckets at the corresponding rates (Table 22) and stirred with a plastic paddle until all the acid disappeared. This took approximately 3 minutes at the higher concentrations.

1.3L of each mix was then transferred to a 6L spray canister and pressurized using compressed air.

Table 22. Mixing ratios of Alkali Surfactant Blend (Pack 1), glyphosate acid (Pack 2) and water and equivalent rate of glyphosate applied per hectare. NB ai = active ingredient measured as the pure acid. BS/FA = Huntsman's Empigen BS/FA

Glyphosate Rate g ai/ha	Glyphosate concentration g ai/L	Pack 1 – Alkali-Surfactant blend (MEA-BS/FA) ml/bucket	Pack 2- Glyphosate acid g/bucket	Water L/bucket
30	0.47	3.86	6.41	13.00
60	0.94	7.72	12.83	12.99
90	1.41	11.58	19.24	12.99
120	1.88	15.45	25.66	12.98
180	2.81	23.17	38.49	12.98
240	3.75	30.89	51.32	12.97
450	7.03	57.92	96.22	12.94

Commercial Herbicides

Two commercial glyphosate formulations Roundup Biactive (containing an isopropylamine salt of glyphosate) and Macphersons Glyphosate 450 SL (containing a monoethanolamine salt of glyphosate) were used as commercial standards.

The commercial standard herbicides were added directly to the water in the spray canister at the required rate to match the concentration of pure glyphosate acid in the 2 Pack system described above. See Table 23 for Roundup Biactive

and Table 24 for Macphersons Glyphosate 450SL mixing ratios with water in the spray canister. The commercial standards were not mixed in a bucket.

Roundup Biactive

5 **Table 23.** Mixing ratios of Roundup Biactive and water.

Glyphosate Rate g ai/ha	Glyphosate concentration g ai/L	Roundup Biactive concentration g/L	Roundup Biactive g/canister	Water L/canister
30	0.469	360	1.69	1298.3
60	0.938	360	3.39	1296.6
90	1.41	360	5.08	1294.9
120	1.875	360	6.77	1293.2
180	2.81	360	10.16	1289.8
240	3.75	360	13.54	1286.5
450	7.03	360	25.39	1274.6

Macphersons Glyphosate 450 SL

Table 24. Mixing ratios of Macphersons Glyphosate 450SL and water.

Glyphosate Rate g ai/ha	Glyphosate concentration g ai/L	Macphersons Glyphosate 450SL concentration g/L	Macphersons Glyphosate 450SL g/canister	Water L/canister
30	0.469	450	1.35	1298.6
60	0.938	450	2.71	1297.3
90	1.41	450	4.06	1295.9
120	1.875	450	5.42	1294.6
180	2.81	450	8.13	1291.9
240	3.75	360	13.54	1286.5
450	7.03	360	25.39	1274.6

10

Herbicide application

The annual ryegrass was at early tillering stage and the wild radish at the early rosette stage when sprayed with the herbicide treatments.

Herbicide formulations were applied using an enclosed laboratory track-sprayer fitted with three 110° flat fan nozzles (Teejet XR11001-VS) spaced at 50cm intervals across the boom. The boom moved along a fixed track at 7.7 km h⁻¹, sprayed at a water volume of 50 L ha⁻¹ with a pressure of 200 kPa.

5

There were 8 replicates for each treatments.

Assessment

Seedlings were harvested 14 days after spray application by cutting foliage off at base immediately prior to weighing on an AND FX 300 electronic balance (range 0-300 g).

10

Statistical analysis

Data was analysed using a factorial design with two factors, glyphosate formulation (Formulation) and spray application rate (Rate). 95% least significant differences (LSD) were calculated for the mean of each treatment.

15

RESULTS

Annual Ryegrass

20

Averaged across all rates there was a significant difference between the formulations (Table 25). As expected there was a highly significant effect of application rate. There was a significant interaction between Formulation and Rate.

25

Table 25. Statistic analysis data for fresh weight of annual ryegrass plants at harvest.

FACTOR	Prob (F)	LSD
Formulation	<0.01	0.09
Rate	<0.01	0.08
Formulation x Rate	<0.01	0.26

Mean data (8 replicates) for fresh weight at harvest is tabled (Table 26).

When the fresh weight of the annual ryegrass was averaged across all rates there was no significant difference between the 2 Pack system (containing a cocamidopropyl betaine surfactant) and Macphersons 450SL. However, in this experiment both of these formulations were more efficacious than Roundup Biactive (Table 26). As expected there was a highly significant effect of application rate.

- 10 There was a significant interaction between Formulation and Rate due largely to higher than expected fresh weight for plants treated with Roundup Biactive in the mid rate range of 90 to 180g ai/ha.

15 **Table 26.** Mean fresh weight (g/plant) of annual ryegrass plants harvested after treatment with a 2 Pack System (which contains a cocamidopropyl betaine as the surfactant component) compared with commercial standards Macphersons 450SL and Roundup Biactive. Mean data followed by the same letter are not significantly different ($P < 0.05$).

Glyphosate application rate g ai/ha	2 Pack System	MacPhersons 450SL	Roundup Biactive
0	6.21	6.21	6.21
30	5.52	5.91	5.60
60	4.75	5.08	5.04
90	2.71	2.73	3.28
120	2.11	2.05	2.85
180	1.30	1.03	1.47
240	0.61	0.48	0.76
450	0.30	0.19	0.23
Formulation Mean (not including 0g ai/ha)	2.47 a	2.50 a	2.75 b

Wild Radish

Averaged across all rates there was a significant difference between the formulations (Table 27). As expected there was a highly significant effect of application rate. There was a significant interaction between Formulation and Rate.

5 **Table 27.** Statistic analysis data for fresh weight of wild radish plants at harvest.

FACTOR	Prob (F)	LSD
Formulation	<0.01	0.08
Rate	<0.01	0.17
Formulation x Rate	<0.01	0.21

Mean data (eight replicates) for fresh weight at harvest is tabled (Table 25).

10 When the fresh weight of the wild radish was averaged across all rates there was no significant difference between the 2 Pack system (containing a cocamidobetaine surfactant) and Macphersons 450SL. However, as with the ryegrass, both of these formulations were more efficacious than Roundup Biactive (Table 28). As expected there was a highly significant effect of
15 application rate.

There was a significant interaction between Formulation and Rate once again due largely to higher than expected harvested weights for wild radish plants treated with Roundup Biactive in the mid rate range of 90 - 180g ai/ha.

20

Table 28. Mean fresh weight (g/plant) of wild radish plants harvested after treatment with a 2 Pack System (which contains a cocamidopropyl betaine as the surfactant component) compared with commercial standards Macphersons 450SL and Roundup Biactive. Mean data followed by the same letter are not
25 significantly different ($P < 0.05$).

Glyphosate application rate g ai/ha	2 Pack System	MacPhersons 450SL	Roundup Biactive
0	7.25	7.25	7.25
30	5.90	6.40	6.36
60	4.94	5.33	5.48
90	4.14	4.21	4.48
120	3.35	3.39	3.68
180	2.38	2.20	3.07
240	1.41	1.17	1.46
450	0.27	0.22	0.26
Average	3.20 a	3.27 a	3.54 b

Experiment 10

5 Bioefficacy comparison on annual ryegrass of a 2 Pack glyphosate system using a cocamidopropyl betaine (also know as alkyl amidopropyl betaine) and triethanolamine glyphosate salt with commercial glyphosate standards.

INTRODUCTION

10 Triethanolamine is an alternative alkali to monethanolamine. Cocamidopropyl betaine is a surfactant option that has been used in Example 9.

METHOD

Plant propagation

15 Annual ryegrass (*Lolium rigidum*) seeds (3/pot) were sown to a depth of 5mm in 10cm diameter pots filled with potting mix (Australian Standard 3743) that had been amended with macro and micronutrients for optimal growth.

20 One week after seedling emergence, seedlings were thinned for uniform size to one seedling per pot. Plants were grown in a temperature-controlled greenhouse

(14°C – 25°C) for 18 days then outdoors for 22 days prior to spray application to more closely simulate field conditions. After the application of herbicides the pots were returned to the greenhouse for an additional 14 days before plants were assessed by harvesting all foliage for fresh weight.

5 2 Pack herbicide preparation

In this example the triethanolamine - cocamidopropyl betaine surfactant blend was made using 91.6 parts triethanolamine to 25.3 parts cocamidopropyl betaine (aqueous concentration 30%) and is called the Alkali Surfactant Blend.

- 10 Approximately 13L of water was added to seven 20L plastic buckets. A blend of triethanolamine and the cocamidopropyl betaine (also know as alkyl amidopropyl betaine) surfactant sold as Huntsman's Empigen BS/FA (Pack 1) was then added to the water at the rates given in the Table 29. Glyphosate acid (Pack 2) with a purity of 95% was then added to the buckets at the corresponding rate
- 15 (Table 29) and stirred with a plastic paddle until all the acid disappeared.

1.3L of each mix was then transferred to a 6L spray canister and pressurized using compressed air.

- 20 **Table 29.** Mixing ratios of the Alkali Surfactant Blend (Pack 1), glyphosate acid (Pack 2) and water and equivalent rate of glyphosate applied per hectare. NB ai = active ingredient measured as the pure acid. BS/FA = Huntsman's Empigen BS/FA

Glyphosate Rate g ai/ha	Glyphosate concentration g ai/L	Pack 1 – Alkali-Surfactant blend (triethanolamine - BS/FA) ml/bucket	Pack 2- Glyphosate acid g/bucket	Water L/bucket
30	0.47	3.86	6.41	13.00
60	0.94	7.72	12.83	12.99
90	1.41	11.58	19.24	12.99
120	1.88	15.45	25.66	12.98
180	2.81	23.17	38.49	12.98
240	3.75	30.89	51.32	12.97
450	7.03	57.92	96.22	12.94

Commercial Herbicides

Two commercial glyphosate formulations Roundup Biactive and Macphersons Glyphosate 450 SL were used as commercial standards.

- 5 The commercial standard herbicides were added directly to the water in the spray canister at the required rate to match the concentration of pure glyphosate acid in the 2 pack system described above. See Table 30 for Roundup Biactive and Table 31 for Macphersons Glyphosate 450SL mixing ratios with water in the spray canister. The commercial standards were not mixed in a bucket.

10

Roundup Biactive

Table 30. Mixing ratios of Roundup Biactive and water.

Glyphosate Rate g ai/ha	Glyphosate concentration g ai/L	Roundup Biactive concentration g/L	Roundup Biactive g/canister	Water L/canister
30	0.469	360	1.69	1298.3
60	0.938	360	3.39	1296.6
90	1.41	360	5.08	1294.9
120	1.875	360	6.77	1293.2
180	2.81	360	10.16	1289.8
240	3.75	360	13.54	1286.5
450	7.03	360	25.39	1274.6

Macphersons Glyphosate 450 SL

Table31. Mixing ratios of Macphersons Glyphosate 450SL and water.

Glyphosate Rate g ai/ha	Glyphosate concentration g ai/L	Macphersons Glyphosate 450SL concentration g/L	Macphersons Glyphosate 450SL g/canister	Water L/canister
30	0.469	450	1.35	1298.6
60	0.938	450	2.71	1297.3
90	1.41	450	4.06	1295.9
120	1.875	450	5.42	1294.6
180	2.81	450	8.13	1291.9
240	3.75	360	13.54	1286.5
450	7.03	360	25.39	1274.6

15

Herbicide application

The annual ryegrass was at early tillering stage when sprayed with the herbicide treatments.

- 5 Herbicide formulations were applied using an enclosed laboratory track-sprayer fitted with three 110° flat fan nozzles (Teejet XR11001-VS) spaced at 50cm intervals across the boom. The boom moved along a fixed track at 7.7 km h⁻¹, sprayed at a water volume of 50 L ha⁻¹ with a pressure of 200 kPa.
- 10 There were 8 replicates for each treatments.

Assessment

Seedlings were harvested 14 days after spray application by cutting foliage off at base immediately prior to weighing on an AND FX 300 electronic balance (range 0-300 g).

15

Statistical analysis

Data was analysed using a factorial design with two factors, glyphosate formulation (Formulation) and spray application rate (Rate). 95% least significant differences (LSD) were calculated for the mean of each treatment.

20

RESULTS

- Averaged across all rates there was a no significant difference between the formulations proving that the efficacy of the 2 Pack system containing a triethanolamine base and cocamidopropyl betaine surfactant and the commercial standards was similar (Table 32). As expected there was a highly significant effect of application rate.
- 25

Table 32. Statistic analysis data for fresh weight of annual ryegrass plants at harvest. NS = Not significantly different ($P < 0.05$).

FACTOR	Prob (F)	LSD
Formulation	NS	
Rate	<0.01	0.17
Formulation x Rate	NS	

5 Mean data (8 replicates) for fresh weight at harvest is tabled (Table 33).

10 **Table 33.** Mean Fresh weight (g/plant) of annual ryegrass plants harvested after treatment with a 2 Pack system (which contains a cocamidopropyl betaine as the surfactant component and triethanolamine as the alkali) with commercial standards Macphersons 450SL and Roundup Biactive. Mean data followed by the same letter are not significantly different ($P < 0.05$).


Glyphosate application rate g ai/ha	2 Pack System	MacPhersons 450SL	Roundup Biactive
0	6.41	6.41	6.41
30	5.79	5.73	5.67
60	4.83	4.87	4.82
90	2.63	2.55	2.59
120	2.17	2.29	2.23
180	1.32	1.34	1.33
240	0.64	0.64	0.63
450	0.30	0.31	0.30
Means	2.53 a	2.53 a	2.51 a

The scope of the claims should not be limited by the preferred embodiments set forth in the examples but should be given the broadest interpretation consistent with the description as a whole.


CLAIMS:

1. A method of preparing a spray tank mix of glyphosate for application to plants from a spray tank, the method comprising the steps of:
 - (a) providing a two pack formulation system comprising (i) a pack of a glyphosate acid concentrate in the form of solid particulate material of particle size in the range 0.1mm to 5mm said glyphosate acid concentrate comprising at least 80% by weight glyphosate acid and (ii) a pack of a water soluble alkaline composition comprising an alkanolamine and a surfactant;
 - (b) providing a spray tank for spray application to plants;
 - (c) diluting the alkaline composition with water in the spray tank to provide a diluted alkaline mixture comprising alkanolamine and surfactant;
 - (d) forming an aqueous dispersion of the particulate glyphosate acid concentrate with mixing;
 - (e) adding the mixed aqueous dispersion of particulate glyphosate acid to the dilute alkaline mixture to thereby react therewith and form an alkanolamine salt of glyphosate in situ in the spray tank; and
 - (f) optionally adding further water to the spray tank to provide the spray tank mix of alkanolamine salt of glyphosate for application to plants from the spray tank wherein the spray tank mix comprises in the range of from 0.1 to 9.0% by weight glyphosate salt based on glyphosate acid equivalent.
2. A method according to claim 1 wherein the concentrate comprising glyphosate acid is a solid particulate material of particle size in the range of from 0.1 mm to 2 mm.
3. A method according to claim 2 wherein the glyphosate acid concentrate is in the form of particles comprising less than 10% by weight water.
4. A method according to any one of claims 1-3 wherein the alkaline composition has pH (1% aqueous solution) in the range of from 8 to 13.5.
5. A method according to claim 1 wherein the alkanolamine is monoethanolamine, diethanolamine, triethanolamine or alkyl alkanolamine.

6. A method according to any one of claims 1-5 wherein the alkaline composition further comprises one or more surfactants wherein the surfactant is alkylamine ethoxylates, cocobetaine, alkoxyated aliphaticamine, alkylpolyglucoside, alkylglycoside, aliphatic amine quaternary amines, aliphaticamine acetate, ethylene diamine ethoxylate, polyoxyethylene aliphaticamine oxides and polyoxyethylene alkyl ethers salts, amine oxides, amphotoacetates, cocobetaine or cocamidopropyl betaines.
7. A method according to any one of claims 1-6 wherein the alkaline composition comprises at least one surfactant wherein the surfactant is:
- (a) Betaine surfactant of formula IIa



wherein R is C₁₂ to C₁₈ alkyl, C₁₂ to C₁₈ alkenyl, or alkylamidoalkyl, and R¹ and R² are each independently (C₁ to C₆) alkyl or hydroxyl (C₁ to C₆) alkyl;
wherein "alkylamidoalkyl" means a group according to formula III:



wherein R⁴ is C₁ to C₃₀ alkyl or C₁ to C₃₀ alkenyl and R⁵ is a C₁ to C₆ alkylenyl;

- (b) aliphatic amines and ethoxylated aliphatic or ethoxylated alkyl/fatty amines comprising at least one hydrocarbon group containing 12 to 18 carbon atoms, optionally polyalkoxylated; or
- (c) amine oxide surfactant:
- an amine oxide having formula R⁷R⁸R⁸N→O (V),
 - an amine oxide having formula R⁷-CO-NH-R⁹-R⁸R⁸N→O (VI) or
 - a mixture or association thereof,

wherein:

R⁷ which is identical or different, is a linear or branched hydrocarbon group,

R⁸, which is identical or different, is a C₁-C₃ alkyl group, and

R⁹, which is identical or different, is divalent linear or branched hydrocarbon group containing 1 to 6 carbon atoms.


8. A method according to claim 5 wherein the glyphosate acid and alkaline composition are mixed in a molar ratio of glyphosate acid:alkanolamine in the range of 0.8 to 1 to 1 to 1.2.
9. A method according to any one of claims 1-8 wherein the final volume of spray tank mix is in the range of from 50 to 20,000 litres.
10. A method according to any one of claims 1-9 wherein the glyphosate acid concentrate and alkaline composition are mixed to provide a pH in the spray tank mix of alkanolamine salt of glyphosate in the range of from 3.5 to 6.0.
11. A method according to claim 1 wherein the aqueous dispersion of the particulate glyphosate acid concentrate is prepared in a mixing vessel providing a vortex of water.
12. A method of preparing a spray tank mix of glyphosate for application to plants from a spray tank, the method comprising the steps of:
 - (a) providing a two pack formulation system comprising
 - (i) a pack of glyphosate acid concentrate in the form of solid particulate material having a particle size in the range 0.1 to 2 mm and comprising less than 10% by weight water, said concentrate composition comprising at least 80% by weight glyphosate acid, and
 - (ii) a pack of water soluble alkaline composition comprising ethanolamine and surfactant;
 - (b) providing a spray tank for spray application to plants;
 - (c) diluting the alkaline composition with water in the spray tank to provide a dilute alkaline mixture comprising ethanolamine and surfactant;
 - (d) forming an aqueous dispersion of the particulate glyphosate acid concentrate with mixing;

- (e) adding the mixed aqueous dispersion of particulate glyphosate acid to the dilute alkaline mixture to react with the alkaline mixture to form the ethanolamine salt of glyphosate in situ in the spray tank; and
 - (f) optionally adding further water to the spray tank;
- to provide the spray tank mix of glyphosate for application to plants from a spray tank wherein the final volume of spray tank mix is in the range of from 50 to 20,000 litres and comprises in the range of from 0.1 to 9.0% by weight glyphosate salt based on glyphosate acid equivalent.
13. A method according to claim 1 wherein the alkanolamine is ethanolamine.
14. A method of preparing a spray tank mix of glyphosate for application to plants from a spray tank, the method comprising the steps of:
- (a) providing a two pack formulation system comprising (i) a pack of a glyphosate acid concentrate in the form of solid particulate material of particle size in the range 0.1mm to 5mm said glyphosate acid concentrate comprising at least 80% by weight glyphosate acid and (ii) a pack of a water soluble alkaline composition comprising an alkanolamine and a betaine surfactant;
 - (b) providing a spray tank for spray application to plants;
 - (c) diluting the alkaline composition with water in the spray tank to provide a diluted alkaline mixture comprising alkanolamine and betaine surfactant;
 - (d) forming an aqueous dispersion of the particulate glyphosate acid concentrate with mixing;
 - (e) adding the mixed aqueous dispersion of particulate glyphosate acid to the dilute alkaline mixture to react therewith and form the alkanolamine salt of glyphosate in situ in the spray tank; and
 - (f) optionally adding further water to the spray tank to provide the spray tank mix of alkanolamine salt of glyphosate for application to plants from the spray tank wherein the spray tank mix comprises in the range of from 0.1 to 9.0% by weight glyphosate salt based on glyphosate acid equivalent.
15. A method according to claim 14 where the betaine surfactant is an amidoaliphaticbetaine or cocobetaine surfactant.

16. A method according to claim 14, further comprising preparing a kit comprising a first pack of the glyphosate acid concentrate by forming a composition comprising at least 80% glyphosate acid into granules;
and a second pack comprising the alkaline composition; and transporting the kit to the site of use.
17. The method according to claim 7, wherein R^7 is an alkyl group containing 3 to 30 carbon atoms.
18. The method according to claim 7, wherein R^7 is an alkyl group containing 3 to 20 carbon atoms.
19. The method according to claim 7, wherein R^8 is a methyl group.
20. The method according to claim 7, wherein R^9 is further substituted with a hydroxyl group.


1/3

Figure 1



2/3

Figure 2



3/3

Figure 3

