

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 July 2009 (02.07.2009)

PCT

(10) International Publication Number
WO 2009/079768 A1

(51) International Patent Classification:

A61K 39/395 (2006.01) A61P 35/04 (2006.01)
A61K 31/7105 (2006.01) C12Q 1/48 (2006.01)
A61K 45/00 (2006.01) C12Q 1/68 (2006.01)
A61K 48/00 (2006.01) G01N 33/574 (2006.01)
A61P 35/00 (2006.01) G01N 33/68 (2006.01)

(21) International Application Number:

PCT/CA2008/002228

(22) International Filing Date:

19 December 2008 (19.12.2008)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/008,649 21 December 2007 (21.12.2007) US

(71) Applicant (for all designated States except US): **UNIVERSITY HEALTH NETWORK** [CA/CA]; 190 Elizabeth Street, Toronto, Ontario M5G 2C4 (CA).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **PAN, Guohua** [CA/CA]; 182 Trelawn Avenue, Oakville, Ontario L6J 4R2 (CA).

(74) Agents: **ERRATT, Judy, A.** et al.; Gowling Lafleur Henderson LLP, 160 Elgin Street, suite 2600, Ottawa, Ontario K1P 1C3 (CA).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

(54) Title: METHODS OF INHIBITING TUMOR GROWTH USING TTK ANTAGONISTS

(57) Abstract: The present invention relates to methods for treating TTK positive breast cancers or soft-tissue sarcomas in a mammalian subject by administering a therapeutically effective amount of a TTK antagonist. The invention also provides compositions comprising a TTK antagonist and a HER-2 antagonist, as well as methods of diagnosing a basal-like breast cancer and methods of determining the prognosis of a subject having a cancer by assessing expression of TTK in a tumor sample from a subject.

WO 2009/079768 A1

METHODS OF INHIBITING TUMOR GROWTH USING TTK ANTAGONISTS

RELATED APPLICATION

This application claims the benefit of U.S. Provisional Application No. 61/008,649, filed on December 21, 2007.

- 5 The entire teachings of the above application are incorporated herein by reference.

BACKGROUND OF THE INVENTION

- Cancer is the second leading cause of death after heart disease and is the primary cause of death in women between the ages of 35 and 74 in the United
10 States. Based on estimates of the National Institutes of Health, overall costs for cancer in the year 2000 were 180.2 billion U.S. dollars. Cancer-related costs account for about 10 percent of the total amount spent on disease treatment in the United States. Thus, cancer is a major national burden.

- Breast cancer is the most common cancer among American women, except
15 for skin cancers. In addition, breast cancer is the second leading cause of cancer death in women, exceeded only by lung cancer. The chance of developing invasive breast cancer at some time in a woman's life is about 1 in 8 (12%). Women living in North America have the highest rate of breast cancer in the world and it is estimated that about 178,480 new cases of invasive breast cancer will be diagnosed among
20 women in the United States in 2007.

- Recent studies have led to the identification of classes, or sub-types, of breast cancer that are defined by gene expression profiles and molecular features (Lønning *et al.*, *Endocrine-Related Cancer* 8:259-263 (2001)). These include Luminal A, Luminal B, HER2-type, normal breast-like and basal-like breast cancers. Basal-like
25 breast cancers are high-grade cancers that are associated with rapid growth and poor prognosis, including high risk for metastasis, recurrence, and death. There is no definitive test or validated clinical assay at this time that accurately identifies basal-like breast cancers. Accordingly, there is an urgent need to identify new markers

that can be used to recognize basal-like breast cancer more readily and to develop targeted treatments for this disease.

Human TTK protein kinase (TTK), also known as tyrosine threonine kinase, dual specificity protein kinase TTK, Monopolar Spindle 1 (Mps1) and

5 Phosphotyrosine-Picked Threonine Kinase (PYT), is a conserved multispecific kinase that is capable of phosphorylating serine, threonine and tyrosine residues when expressed in *E. coli* (Mills *et al.*, *J. Biol. Chem.* 22(5): 16000-16006 (1992)). TTK mRNA is not expressed in the majority of physiologically normal tissues in human (*Id*). TTK mRNA is expressed in some rapidly proliferating tissues, such as
10 testis and thymus, as well as in some tumors (TTK mRNA was not expressed in renal cell carcinoma, was expressed in 50% of breast cancer samples, was expressed in testicular tumors and ovarian cancer samples) (*Id*). TTK is expressed in some cancer cell lines and tumors relative to normal counterparts(*Id.*; see also WO 02/068444 A1).

15 The yeast homolog of human TTK, Mps1, is required for normal cell cycle progression and has been shown to have a role in centrosome/spindle body duplication, the mitotic spindle checkpoint and cytokinesis (Winey and Huneycutt, *Oncogene* 21: 6161-6169 (2002)). Studies in *Drosophila* and vertebrates have confirmed that several of these functions are conserved in higher eukaryotes,
20 including humans (Fisk and Winey, *Curr. Biol.* 14: R1058-1060 (2004); Fischer *et al.*, *Curr. Biol.* 14: 2019-2024 (2004); Fisk *et al.*, *PNAS* 100(25): 14875-14880 (2003)). The specific biochemical role of TTK in cellular physiology is not currently understood.

The requirement of normal TTK kinase activity for controlled cell cycle
25 progression in yeast, *Drosophila*, and vertebrates, including humans, combined with the detection of TTK overexpression in some human cancer samples and cancer cell lines indicates that TTK should be investigated further as a potential target for therapeutic anti-cancer agents. However, the role of TTK in the pathology of breast cancer and particular subtypes of breast cancer is not understood.

30 There is a need to determine whether TTK expression and/or activity is associated with particular subtypes of breast cancer. In addition, there is a need to identify and develop therapeutic agents that target TTK expression and/or activity to treat breast cancer subtypes that are characterized by TTK overexpression.

SUMMARY OF THE INVENTION

The present invention relates to a method for treating TTK positive non-luminal breast cancer (*e.g.*, a basal-like breast cancer, a HER-2 positive breast cancer) in a mammalian subject. In the method of the invention, a therapeutically effective amount of a TTK antagonist is administered to the mammalian subject. The TTK positive non-luminal breast cancer can be a basal-like breast cancer or a HER-2 positive breast cancer, such as a HER-2 positive, estrogen receptor (ER) negative breast cancer. The TTK antagonist inhibits the expression and/or activity of a TTK gene or gene product and can be, for example, an antibody, an antigen-binding fragment of an antibody, a small interfering ribonucleic acid (siRNA), a peptide, a peptidomimetic, an antisense oligonucleotide, or a small molecule. The TTK antagonist can be administered with one or more other therapeutic agents, including, but not limited to, a HER-2 antagonist, such as trastuzumab.

The invention also relates to a method for treating a TTK positive soft tissue sarcoma in a mammalian subject. The method comprises administering a therapeutically effective amount of a TTK antagonist to the mammalian subject. Examples of TTK positive soft-tissue sarcomas that can be treated include TTK positive fibrosarcomas, leiomyosarcomas, dedifferentiated liposarcomas, pleiomorphic liposarcomas, malignant fibrous histiocytomas, round cell sarcomas and synovial sarcomas. The TTK antagonist can be, for example, an antibody, an antigen-binding fragment of an antibody, a small interfering ribonucleic acid (siRNA), a peptide, a peptidomimetic, an antisense oligonucleotide, or a small molecule, and can be administered with one or more other therapeutic agents.

The invention also relates to a method of treating a TTK positive basal-like breast cancer tumor in a mammalian subject comprising administering a therapeutically effective amount of a TTK antagonist to the mammalian subject. The TTK antagonist can be, for example, an antibody, an antigen-binding fragment of an antibody, a small interfering ribonucleic acid (siRNA), a peptide, a peptidomimetic, an antisense oligonucleotide, or a small molecule, and can be administered with one or more other therapeutic agents. Examples of TTK positive basal-like breast cancer tumors that can be treated include TTK positive metastatic tumors and carcinomas.

The invention also provides a method of treating a TTK positive breast cancer that is also HER-2 positive and ER negative in a mammalian subject. The method comprises administering a therapeutically effective amount of a TTK antagonist to the mammalian subject. The TTK antagonist can be, for example, an antibody, an antigen-binding fragment of an antibody, a small interfering ribonucleic acid (siRNA), a peptide, a peptidomimetic, an antisense oligonucleotide, or a small molecule, and can be administered with one or more other therapeutic agents, including, but not limited to, a HER-2 antagonist, such as trastuzumab.

The invention further relates to a composition comprising a TTK antagonist, a HER-2 antagonist and a physiologically acceptable carrier. The HER-2 antagonist can be, for example, trastuzumab.

The invention further relates to a method for identifying a candidate for an anti-cancer therapy using a TTK antagonist. The method comprises providing a suitable tumor sample obtained from a subject and assessing expression of TTK in the tumor sample. According to the invention, expression of TTK by the tumor, or increased expression of TTK by the tumor relative to a suitable control, indicates that the subject is a candidate for an anti-cancer therapy using a TTK antagonist.

The invention also encompasses a method of diagnosing a non-luminal breast cancer in a subject. The method comprises providing a suitable tumor sample from breast tissue of the subject and assessing expression of TTK in said tumor sample. Expression of TTK by the tumor, or increased expression of TTK by the tumor relative to a suitable control, indicates that the subject has a basal-like breast cancer.

The invention also relates to a method of determining the prognosis of a subject who has a cancer. The method comprises providing a suitable tumor sample from the subject and assessing expression of TTK in the tumor sample. According to the invention, expression of TTK by the tumor, or increased expression of TTK by the tumor relative to a suitable control, indicates reduced patient survival, increased risk of metastases or increased risk of relapse.

The invention further provides a method for screening a breast cancer patient as an aid for selecting aggressive cancer therapy. The method comprises providing a suitable sample from a patient and determining TTK expression in the sample. According to the invention, increased TTK expression in the sample from the patient

as compared with a suitable control indicates the patient is a candidate for aggressive cancer therapy.

BRIEF DESCRIPTION OF THE DRAWINGS

The patent or application file contains at least one drawing executed in color.

5 Copies of this patent or patent application publication with color drawings will be provided by the Office upon request and payment of the necessary fee.

FIG. 1 is a microarray gene expression profile of 36 normal tissues illustrating that only certain physiologically normal tissues in human express TTK RNA.

10 FIG. 2A is a graph depicting TTK RNA expression in several normal breast cell lines and breast cancer cell lines, as determined by quantitative polymerase chain reaction (QPCR). TTK RNA is overexpressed in several breast cancer cell lines.

FIG. 2B is a picture of a Western blot depicting increased expression of TTK
15 protein in breast cancer cell lines relative to normal breast cell lines. Hs578T, MDA-MB-231, MDA-MB-435, and MDA-MB-468 are non-luminal breast cancer cell lines and BT-474, MDA-MB-330, MDA-MB-453 and HCC1419 are luminal breast cancer cell lines. β -actin levels serve as a loading control.

FIG. 3 is a graph depicting microarray expression data illustrating that TTK
20 RNA is overexpressed more frequently and to a greater extent in non-luminal breast cancer cell lines than in luminal, non-malignant breast cancer cell lines, relative to normal breast cell lines or non-malignant breast cells.

FIG. 4 is a microarray gene expression profile illustrating TTK RNA
25 overexpression in non-basal-like, BRCA1-associated and basal-like cancer cell lines relative to normal breast cell lines. TTK RNA expression is shown to be highest in basal-like breast cancer cell lines.

FIG. 5 is a graph illustrating TTK RNA expression levels in different normal
soft tissue (normal, GIST) and soft tissue sarcoma samples (fibrosarcoma,
leiomyosarcoma, Lipo dediff, Lipo pleo, MFH, Round cell, Synovial sarcoma).
30 Increased TTK RNA levels were observed in sarcoma samples relative to samples from physiologically normal tissues.

FIG. 6 is a graph illustrating a strong association ($p=3.7E-08$) between high TTK expression in breast and reduced patient survival in three data sets (NEJM, Perou, PNAS).

FIG. 7 is a graph illustrating an association ($p=0.014$) between high TTK expression in breast and increased risk of developing metastases.

FIGS. 8A and 8B are graphs illustrating the association between TTK expression in breast and risk of relapse in two data sets (NEJM295, Perou). FIG. 8A illustrates an association ($p=5.9E-4$) between high TTK expression and increased risk of breast cancer relapse.

FIG. 9A is a graph illustrating siRNA-mediated knockdown of TTK RNA expression levels in MD-MB-468 breast cancer cells treated with one of four different siRNAs that target human TTK (siRNA #1, siRNA #2, siRNA #3, siRNA #4), as determined by quantitative polymerase chain reaction (QPCR). siCONTROL is a non-targeting siRNA that targets firefly luciferase. siRNA POOL is a combination of siRNA #1, siRNA #2, siRNA #3 and siRNA #4. siRNA #2 and siRNA #3 display a TTK knockdown efficiency of greater than 70% (see downward pointing arrows).

FIG. 9B is a picture of a Western blot depicting siRNA-mediated knockdown of TTK protein expression in lysates of MD-MB-468 breast cancer cells treated with one of four different siRNAs that target human TTK (siRNA #1, siRNA #2, siRNA #3, siRNA #4). siCONTROL is a non-targeting siRNA that targets firefly luciferase. siRNA POOL is a combination of siRNA #1, siRNA #2, siRNA #3 and siRNA #4. β -actin levels serve as a loading control.

FIG. 10A-10C is a series of fluorescence histograms illustrating flow cytometry analysis of MCF7 breast cancer cells treated with one of two TTK-targeting siRNAs (TTK siRNA #2 (FIG. 10B), TTK siRNA #3 (FIG. 10C)) or non-targeting control siRNA that targets firefly luciferase (siCONTROL) (FIG. 10A). Treatment of the cells with TTK siRNA #2 or TTK siRNA #3, but not siCONTROL siRNA, caused apoptosis of the cancer cells, indicated by increased G2/M and <G1 populations.

FIG. 11A is a graph illustrating a reduction in the number of colonies formed in soft agar by T-47D breast cancer cells treated with TTK-targeting siRNAs (siRNA #2 (FIG. 11C), siRNA #3 (FIG. 11D), siRNA POOL (FIG. 11E)), relative to

T-47D breast cancer cells treated with non-targeting control siRNA that targets firefly luciferase (siCONTROL) (FIG. 11B). Results are based on two independent experiments.

FIG. 11B-11E is a series of photographs depicting a reduction in the number and size of colonies formed in soft agar by T-47D breast cancer cells treated with TTK-targeting siRNAs (siRNA #2, siRNA #3, siRNA POOL), relative to T-47D breast cancer cells treated with non-targeting control siRNA that targets firefly luciferase (siCONTROL). A 40 nM concentration of siRNA was used in each transfection.

FIG. 12 is a graph illustrating reduced survival of breast cancer cells from several different breast cancer cell lines following treatment with SP600125, a small molecule antagonist of TTK.

FIG. 13A-13C depicts the chemical structures of SP6000125 (FIG. 13A) and particular substituted derivatives of SP600125 (FIG. 13B-13C).

FIG. 14A-14C depicts the chemical structures of cincreasin (FIG. 14A) and particular cincreasin derivatives (FIG. 14B-14C).

FIGS. 15A and 15B show the human TTK cDNA sequence (Genbank Accession No. M86699).

FIG. 16 shows the human TTK protein sequence (Genbank Accession No. NP003309).

FIG. 17A is a graph depicting normalized TTK transcript levels in MDA-MB-468 cells infected with constructs expressing either control shRNA (pSIREN-shLUC) or TTK shRNA (pSIREN-shTTK). TTK transcript levels in MDA-MB-468 cells expressing TTK shRNA were reduced by about 50% relative to controls.

FIG. 17B is a graph depicting significant suppression of tumor growth in mice that were injected with MDA-MB-468 cells infected with a construct expressing TTK shRNA (pSIREN-shTTK) relative to mice injected with MDA-MB-468 cells infected with a construct expressing control shRNA (pSIREN-shLUC); 2.5×10^6 pSIREN-shLUC (control) and pSIREN-shTTK infected MDA-MB-468 cells were injected subcutaneously into the left and right hindlimb, respectively of nude mice (n=5).

DETAILED DESCRIPTION OF THE INVENTION

Definitions

As used herein, "TTK protein," "TTK kinase," or "TTK" refers to a naturally occurring or endogenous TTK (*e.g.*, mammalian, human) protein, and to proteins having an amino acid sequence that is the same as that of naturally occurring or endogenous TTK protein (*e.g.*, recombinant proteins, synthetic proteins).

Accordingly "TTK protein," "TTK kinase," or "TTK" includes polymorphic or allelic variants and other isoforms of a TTK protein (*e.g.*, mammalian, human) produced by, *e.g.*, alternative splicing or other cellular processes, that occur naturally in mammals (*e.g.*, humans, non-human primates). Preferably, the TTK protein is a human protein that has the amino acid sequence of SEQ ID NO: 2. (See, Genbank Accession No. NP003309 and FIG. 16).

As defined herein, a "TTK antagonist" is an agent (*e.g.*, nucleic acid, protein, peptide, peptidomimetic, antibody, small molecule), which specifically and, preferably, selectively binds a TTK protein and inhibits (*e.g.*, reduces, prevents) one or more activities of a TTK protein; or an agent that inhibits (*e.g.*, reduces, prevents) the expression of a TTK gene and/or protein. A TTK antagonist can inhibit the activity of a TTK protein by, for example, inhibiting the binding of ATP, blocking the active site of the protein's kinase domain (*e.g.*, amino acids 525-797 of SEQ ID NO:2) and/or blocking the association of TTK with one or more of its substrates (*e.g.*, BLM helicase, CHK2/hCds1 protein kinase, Smad2, Smad3) in a cell. A TTK antagonist that inhibits the expression and/or activity of a TTK can be, for example, a natural or synthetic nucleic acid or nucleic acid analog, antisense molecule, small interfering RNA (siRNA), protein, peptide, antibody, small molecule, chemical compound or the like.

As used herein, the term "peptide", refers to a compound consisting of from about 2 to about 100 amino acid residues wherein the amino group of one amino acid is linked to the carboxyl group of another amino acid by a peptide bond. Such peptides are typically less than about 100 amino acid residues in length and preferably are about 10, about 20, about 30, about 40 or about 50 residues.

As used herein, the term "peptidomimetic", refers to molecules which are not polypeptides, but which mimic aspects of their structures. Peptidomimetic antagonists can be prepared by conventional chemical methods (see *e.g.*, Damewood J.R. "Peptide Mimetic Design with the Aid of Computational Chemistry" in *Reviews*

in *Computational Biology*, 2007, Vol. 9, pp.1-80, John Wiley and Sons, Inc., New York, 1996; Kazmierski W.K., "*Methods of Molecular Medicine: Peptidomimetic Protocols*," Humana Press, New Jersey, 1999).

As defined herein, "therapy" is the administration of a particular therapeutic or prophylactic agent to a subject (*e.g.*, a mammal, a human).

As defined herein a "treatment regimen" is a regimen in which one or more therapeutic or prophylactic agents are administered to a mammalian subject at a particular dose (*e.g.*, level, amount, quantity) and on a particular schedule or at particular intervals (*e.g.*, minutes, days, weeks, months).

As defined herein, "direct inhibition of tumor growth" refers to inhibited tumor growth (*e.g.*, reduced tumor cell proliferation, tumor cell death) caused by the interaction of a therapeutic agent with a target in or on a tumor cell. Thus, a TTK antagonist can directly inhibit tumor growth by binding a TTK protein expressed by the cells of the tumor and inhibiting the activity of the TTK protein, for example. In addition, a TTK antagonist can directly inhibit tumor growth by inhibiting expression (*e.g.*, decreasing nucleic acid (*e.g.*, RNA) and/or protein) of a TTK gene product or protein in the cells of the tumor.

As defined herein, a "therapeutically effective amount" is an amount sufficient to achieve the desired therapeutic or prophylactic effect under the conditions of administration, such as an amount sufficient to inhibit (*i.e.*, reduce, prevent) tumor cell growth (proliferation, size) and/or tumor progression (invasion, metastasis) for a particular cancer. The effectiveness of a therapy (*e.g.*, the reduction/elimination of a tumor and/or prevention tumor growth) can be determined by suitable methods (*e.g.*, *in situ* immunohistochemistry, imaging (MRI, NMR), ³H-thymidine incorporation).

As defined herein, an "anti-tumor effective amount" is an amount sufficient to directly inhibit tumor cell growth (*e.g.*, proliferation) or survival.

As defined herein, an "anti-angiogenic effective amount" is an amount sufficient to inhibit angiogenesis.

As used herein, "aggressive therapy" is the administration of a therapeutic agent or agents at higher doses, more frequent doses, or a combination thereof, than is normally administered in a typical therapeutic regime. Aggressive therapy can also be the administration of a combination of therapeutic agents that are not

typically administered in the same therapeutic regime. Aggressive therapy is often at or near the limit of tolerance for a subject receiving such therapy. For example, aggressive chemotherapy is sufficiently toxic that the subject's bone marrow is likely to fail (*e.g.*, the bone marrow will no longer be able to produce hematopoietic cells after aggressive therapy). To get around this anticipated side effect of aggressive therapy, the subject may receive an autologous bone marrow transplant, or receive a tissue-type matched bone marrow transplant.

As described herein, TTK expression or overexpression is associated with certain types of cancers, in particular non-luminal breast cancer subtypes, as well as soft-tissue sarcomas. In particular, the inventor have determined that TTK gene products (*e.g.*, mRNA, protein) are expressed at higher levels in particular breast cancer subtypes, such as basal-like breast cancer and HER-2 positive/estrogen receptor negative breast cancer, relative to normal breast cells or tissues. In addition, the inventor have shown that antagonists of TTK can directly inhibit proliferation (*e.g.*, by inducing apoptosis, by inducing cell cycle arrest, by inhibiting anchorage-independent cell growth) of cancer cells that express TTK. Thus, antagonists of TTK can be used to treat cancers, for example, by inhibiting tumor growth and/or progression (*e.g.*, in cancer patients). Accordingly, the invention provides a method for the targeted treatment of a cancer (*e.g.*, a non-luminal breast cancer, a basal-like breast cancer tumor, a HER-2 positive and ER-negative breast cancer tumor, a soft tissue sarcoma) that expresses TTK and, further, provides for a pharmaceutical composition comprising a TTK antagonist. The invention also provides a method of treating a non-luminal subtype breast cancer by administering a TTK antagonist alone or in combination with one or more other therapeutic agents (*e.g.*, a HER-2 antagonist).

Inhibition of the expression or activity of a TTK protein provides an effective and selective mechanism by which to treat cancers (*e.g.*, tumors) that express TTK. Thus, one aspect of the present invention relates to a method for treating cancer in a mammalian subject comprising administering to the subject a therapeutically effective amount of a TTK antagonist.

TTK antagonists

The TTK antagonist can be an antibody or antigen-binding fragment thereof that selectively binds a TTK protein. The term "antibody" is intended to encompass all types of polyclonal and monoclonal antibodies (*e.g.*, human, chimeric, humanized, primatized, veneered, single chain, domain antibodies (dAbs)) and
5 antigen-binding fragments of antibodies (*e.g.*, Fv, Fc, Fd, Fab, Fab', F(ab'), dAb). (See *e.g.*, Harlow *et al.*, *Antibodies A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988). In a particular embodiment, the TTK-specific antibody is a human antibody or humanized antibody. TTK-specific antibodies can also be directly or indirectly linked to a cytotoxic agent.

10 Several antibodies that selectively bind TTK have been produced and are commercially available (*e.g.*, from Abgent, Abnova Corporation, ABR-Affinity BioReagents, Bethyl laboratories, Cell Signaling Technology, GeneTex, Lifespan Biosciences, Novus Biologicals, Proteintech Group, Inc., Santa Cruz Biotechnology, Inc.). In addition, antibodies that selectively bind to and inhibit the activity of a
15 TTK protein include monoclonal antibodies N1, N2 and C1 described in Stucke *et al.*, *EMBO J.* 21(7):1723-1732 (2002).

Other antibodies or antibody fragments which selectively bind to and inhibit the activity of a TTK protein can also be produced, constructed, engineered and/or isolated by conventional methods or other suitable techniques. For example,
20 antibodies which are specific for a TTK protein can be raised against an appropriate immunogen, such as a recombinant mammalian (*e.g.*, human) TTK protein or portion thereof (including synthetic molecules, *e.g.*, synthetic peptides). A variety of methods have been described (see *e.g.*, Kohler *et al.*, *Nature*, 256: 495-497 (1975) and *Eur. J. Immunol.* 6: 511-519 (1976); Milstein *et al.*, *Nature* 266: 550-552 (1977); Koprowski *et al.*, U.S. Patent No. 4,172,124; Harlow, E. and D.
25 Lane, 1988, *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory: Cold Spring Harbor, NY); *Current Protocols In Molecular Biology*, Vol. 2 (Supplement 27, Summer '94), Ausubel, F.M. *et al.*, Eds., (John Wiley & Sons: New York, NY), Chapter 11, (1991)). Antibodies can also be raised by immunizing
30 a suitable host (*e.g.*, mouse) with cells that express TTK (*e.g.*, cancer cells/cell lines) or cells engineered to express TTK (*e.g.*, transfected cells). (See *e.g.*, Chuntharapai *et al.*, *J. Immunol.*, 152:1783-1789 (1994); Chuntharapai *et al.* U.S. Patent No. 5,440, 021). For the production of monoclonal antibodies, a hybridoma can be

produced by fusing a suitable immortal cell line (*e.g.*, a myeloma cell line such as SP2/0 or P3X63Ag8.653) with antibody producing cells. The antibody producing cells can be obtained from the peripheral blood, or preferably, the spleen or lymph nodes, of humans or other suitable animals immunized with the antigen of interest.

- 5 The fused cells (hybridomas) can be isolated using selective culture conditions, and cloned by limited dilution. Cells which produce antibodies with the desired specificity can be selected by a suitable assay (*e.g.*, ELISA).

Antibody fragments can be produced by enzymatic cleavage or by recombinant techniques. For example, papain or pepsin cleavage can generate Fab or F(ab')₂ fragments, respectively. Other proteases with the requisite substrate specificity can also be used to generate Fab or F(ab')₂ fragments. Antibodies can also be produced in a variety of truncated forms using antibody genes in which one or more stop codons has been introduced upstream of the natural stop site. For example, a chimeric gene encoding a F(ab')₂ heavy chain portion can be designed to include DNA sequences encoding the CH₁ domain and hinge region of the heavy chain. Single chain antibodies, and human, chimeric, humanized or primatized (CDR-grafted), or veneered antibodies, as well as chimeric, CDR-grafted or veneered single chain antibodies, comprising portions derived from different species, and the like are also encompassed by the present invention and the term "antibody". The various portions of these antibodies can be joined together chemically by conventional techniques, or can be prepared as a contiguous protein using genetic engineering techniques. For example, nucleic acids encoding a chimeric or humanized chain can be expressed to produce a contiguous protein. See, *e.g.*, Cabilly *et al.*, U.S. Patent No. 4,816,567; Cabilly *et al.*, European Patent No. 0,125,023 B1; Boss *et al.*, U.S. Patent No. 4,816,397; Boss *et al.*, European Patent No. 0,120,694 B1; Neuberger, M.S. *et al.*, WO 86/01533; Neuberger, M.S. *et al.*, European Patent No. 0,194,276 B1; Winter, U.S. Patent No. 5,225,539; Winter, European Patent No. 0,239,400 B1; Queen *et al.*, European Patent No. 0 451 216 B1; and Padlan, E.A. *et al.*, EP 0 519 596 A1. See also, Newman, R. *et al.*, *BioTechnology*, 10: 1455-1460 (1992), regarding primatized antibody, and Ladner *et al.*, U.S. Patent No. 4,946,778 and Bird, R.E. *et al.*, *Science*, 242: 423-426 (1988)) regarding single chain antibodies.

Humanized antibodies can be produced using synthetic or recombinant DNA technology using standard methods or other suitable techniques. Nucleic acid (*e.g.*, cDNA) sequences coding for humanized variable regions can also be constructed using PCR mutagenesis methods to alter DNA sequences encoding a human or
5 humanized chain, such as a DNA template from a previously humanized variable region (see *e.g.*, Kamman, M., *et al.*, *Nucl. Acids Res.*, 17: 5404 (1989)); Sato, K., *et al.*, *Cancer Research*, 53: 851-856 (1993); Daugherty, B.L. *et al.*, *Nucleic Acids Res.*, 19(9): 2471-2476 (1991); and Lewis, A.P. and J.S. Crowe, *Gene*, 101: 297-302 (1991)). Using these or other suitable methods, variants can also be readily
10 produced. In one embodiment, cloned variable regions (*e.g.*, dAbs) can be mutated, and sequences encoding variants with the desired specificity can be selected (*e.g.*, from a phage library; see *e.g.*, Krebber *et al.*, U.S. 5,514,548; Hoogenboom *et al.*, WO 93/06213, published April 1, 1993).

Other suitable methods of producing or isolating antibodies of the requisite
15 specificity can be used, including, for example, methods which select a recombinant antibody or antibody-binding fragment (*e.g.*, dAbs) from a library (*e.g.*, a phage display library), or which rely upon immunization of transgenic animals (*e.g.*, mice). Transgenic animals capable of producing a repertoire of human antibodies are well-known in the art (*e.g.*, Xenomouse[®] (Abgenix, Fremont, CA)) and can be produced
20 using suitable methods (see *e.g.*, Jakobovits *et al.*, *Proc. Natl. Acad. Sci. USA*, 90: 2551-2555 (1993); Jakobovits *et al.*, *Nature*, 362: 255-258 (1993); Lonberg *et al.*, U.S. Patent No. 5,545,806; Surani *et al.*, U.S. Patent No. 5,545,807; Lonberg *et al.*, WO 97/13852).

A TTK antagonist can be a peptide (*e.g.*, synthetic, recombinant, fusion or
25 derivatized) that specifically binds to, and inhibits (reduces, prevents, decreases) the activity of, the TTK protein. The peptide can be linear, branched or cyclic, *e.g.*, a peptide having a heteroatom ring structure that includes several amide bonds. In a particular embodiment, the peptide is a cyclic peptide.

Peptides, including cyclic peptides, that are selective for binding to a
30 particular domain (*e.g.*, unique domain) of a TTK protein can be produced. A peptide can be, for example, derived or removed from a native protein by enzymatic or chemical cleavage, or can be synthesized by suitable methods, for example, solid phase peptide synthesis (*e.g.*, Merrifield-type synthesis) (see, *e.g.*, Bodanszky *et al.*

"*Peptide Synthesis*," John Wiley & Sons, Second Edition, 1976). Peptides that are TTK antagonists can also be produced, for example, using recombinant DNA methodologies or other suitable methods (see, *e.g.*, Sambrook J. and Russell D.W., *Molecular Cloning: A Laboratory Manual*, 3rd Edition, Cold Spring Harbor

- 5 Laboratory Press, Cold Spring Harbor, New York, 2001). TTK antagonists can also be fusion peptides fused, for example to a carrier protein (*e.g.*, myc, his, glutathione sulfhydryl transferase) and/or tagged (*e.g.*, radiolabeled, fluorescently labeled).

A peptide can comprise any suitable L-and/or D-amino acid, for example, common α -amino acids (*e.g.*, alanine, glycine, valine), non- α -amino acids (*e.g.*, β -alanine, 4-aminobutyric acid, 6-aminocaproic acid, sarcosine, statine), and unusual
10 amino acids (*e.g.*, citrulline, homocitrulline, homoserine, norleucine, norvaline, ornithine). The amino, carboxyl and/or other functional groups on a peptide can be free (*e.g.*, unmodified) or protected with a suitable protecting group. Suitable protecting groups for amino and carboxyl groups, and methods for adding or
15 removing protecting groups are known in the art and are disclosed in, for example, Green and Wuts, "*Protecting Groups in Organic Synthesis*", John Wiley and Sons, 1991. The functional groups of a peptide can also be derivatized (*e.g.*, alkylated) using art-known methods.

Peptides can be synthesized and assembled into libraries comprising a few to
20 many discrete molecular species. Such libraries can be prepared using methods of combinatorial chemistry, and can be screened using any suitable method to determine if the library comprises peptides with a desired biological activity. Such peptide antagonists can then be isolated using suitable methods.

The peptide can comprise modifications (*e.g.*, amino acid linkers, acylation, acetylation, amidation, methylation, terminal modifiers (*e.g.*, cyclizing
25 modifications)), if desired. The peptide can also contain chemical modifications (*e.g.*, N-methyl- α -amino group substitution). In addition, the peptide antagonist can be an analog of a known and/or naturally-occurring peptide, for example, a peptide analog having conservative amino acid residue substitution(s). These modifications
30 can improve various properties of the peptide (*e.g.*, solubility, binding), including its TTK antagonist activity.

Peptidomimetics can be prepared that are TTK antagonists. For example, polysaccharides can be prepared that have the same functional groups as peptides.

Peptidomimetics can be designed, for example, by establishing the three dimensional structure of a peptide agent in the environment in which it is bound or will bind to a target molecule. The peptidomimetic comprises at least two components, the binding moiety or moieties and the backbone or supporting structure.

The binding moieties are the chemical atoms or groups which will react or form a complex (*e.g.*, through hydrophobic or ionic interactions) with a target molecule, for example, with the amino acid(s) at or near the active site of the kinase. For example, the binding moieties in a peptidomimetic can be the same as those in a peptide or protein antagonist. The binding moieties can be an atom or chemical group which reacts with the receptor in the same or similar manner as the binding moiety in the peptide antagonist. For example, computational chemistry can be used to design peptide mimetics of the ATP-binding site and/or active site of the TTK protein, for instance. Examples of binding moieties suitable for use in designing a peptidomimetic for a basic amino acid in a peptide include nitrogen containing groups, such as amines, ammoniums, guanidines and amides or phosphoniums. Examples of binding moieties suitable for use in designing a peptidomimetic for an acidic amino acid include, for example, carboxyl, lower alkyl carboxylic acid ester, sulfonic acid, a lower alkyl sulfonic acid ester or a phosphorous acid or ester thereof.

The supporting structure is the chemical entity that, when bound to the binding moiety or moieties, provides the three dimensional configuration of the peptidomimetic. The supporting structure can be organic or inorganic. Examples of organic supporting structures include polysaccharides, polymers or oligomers of organic synthetic polymers (such as, polyvinyl alcohol or polylactide). It is preferred that the supporting structure possess substantially the same size and dimensions as the peptide backbone or supporting structure. This can be determined by calculating or measuring the size of the atoms and bonds of the peptide and peptidomimetic. In one embodiment, the nitrogen of the peptide bond can be substituted with oxygen or sulfur, for example, forming a polyester backbone. In another embodiment, the carbonyl can be substituted with a sulfonyl group or sulfinyl group, thereby forming a polyamide (*e.g.*, a polysulfonamide). Reverse amides of the peptide can be made (*e.g.*, substituting one or more-CONH-groups for

a-NHCO-group). In yet another embodiment, the peptide backbone can be substituted with a polysilane backbone.

These compounds can be manufactured by known methods. For example, a polyester peptidomimetic can be prepared by substituting a hydroxyl group for the corresponding α -amino group on amino acids, thereby preparing a hydroxyacid and sequentially esterifying the hydroxyacids, optionally blocking the basic and acidic side chains to minimize side reactions. Determining an appropriate chemical synthesis route can generally be readily identified upon determining the chemical structure.

Peptidomimetics can be synthesized and assembled into libraries comprising a few to many discrete molecular species. Such libraries can be prepared using well-known methods of combinatorial chemistry, and can be screened to determine if the library comprises one or more peptidomimetics which have the desired activity. Such peptidomimetic antagonists can then be isolated by suitable methods.

TTK antagonists can also be small molecules. Examples of small molecules include organic compounds, organometallic compounds, inorganic compounds, and salts of organic, organometallic or inorganic compounds. Atoms in a small molecule are typically linked together via covalent and/or ionic bonds. The arrangement of atoms in a small organic molecule may represent a chain (*e.g.* a carbon-carbon chain or a carbon-heteroatom chain), or may represent a ring containing carbon atoms, *e.g.* benzene or a polycyclic system, or a combination of carbon and heteroatoms, *i.e.*, heterocycles such as a pyrimidine or quinazoline. Although small molecules can have any molecular weight, they generally include molecules that are less than about 5,000 daltons. For example, such small molecules can be less than about 1000 daltons and, preferably, are less than about 750 daltons or, more preferably, are less than about 500 daltons. Small molecules and other non-peptidic TTK antagonists can be found in nature (*e.g.*, identified, isolated, purified) and/or produced synthetically (*e.g.*, by traditional organic synthesis, bio-mediated synthesis, or a combination thereof). See *e.g.* Ganesan, Drug Discov. Today 7(1): 47-55 (January 2002); Lou, Drug Discov. Today, 6(24): 1288-1294 (December 2001). Examples of naturally occurring small molecules include, but are not limited to, hormones, neurotransmitters, nucleotides, amino acids, sugars, lipids, and their derivatives.

A small molecule TTK antagonist according to the present invention, and physiologically acceptable salts thereof, can inhibit the kinase activity of TTK (*e.g.*, by directly competing with ATP for binding to the ATP-binding pocket of TTK). Suitable small molecule TTK antagonists for use in the methods of the invention include, for example, anthrapyrazolone compounds and physiologically acceptable salts thereof. Examples of specific anthrapyrazolone compounds useful in the methods of the invention include anthra(1,9-cd)pyrazol-6(2H)-one 1,9-pyrazoloanthrone (SP600125) (FIG. 13A), which is a reversible ATP-competitive inhibitor TTK and Jun N-terminal family kinases (JNK), and particular substituted derivatives of SP600125 (see, for example, FIG. 13B-13C) (Bennett *et al.*, *PNAS* 98(24):13681-13686 (2001); Schmidt *et al.*, *EMBO Reports* 6(9): 866-872 (2005)). SP600125 is available commercially (A.G. Scientific, Inc.; Sigma-Aldrich; Biomol International, LP; SuperArray Bioscience Corporation). Other examples of small molecule TTK antagonists include cincreasin (FIG. 14A) and related compounds, such as isocyanate and halogen-substituted derivatives (see, for example, FIG. 14B-14C) of cincreasin (Dorer *et al.*, *Curr. Biol.* 15:1070-1076 (2005)).

Agents having TTK binding specificity, including small molecules, can be identified in a screen, for example, a high-throughput screen of chemical compounds and/or libraries (*e.g.*, chemical, peptide, nucleic acid libraries). Compounds or small molecules can be identified from numerous available libraries of chemical compounds from, for example, the Chemical Repository of the National Cancer Institute and the Molecular Libraries Small Molecules Repository (PubChem), as well as libraries of the Institute of Chemistry and Cell Biology at Harvard University and other libraries that are available from commercial sources (*e.g.*, Chembridge, Peakdale, CEREP, MayBridge, Bionet). Such libraries or collections of molecules can also be prepared using well-known chemical methods, such as well-known methods of combinatorial chemistry. The libraries can be screened to identify compounds that bind and inhibit TTK. Identified compounds can serve as lead compounds for further diversification using well-known methods of medicinal chemistry. For example, a collection of compounds that are structural variants of the lead can be prepared and screened for TTK binding and/or inhibitory activity. This can result in the development of a structure activity relationship that links the

structure of the compounds to biological activity. Compounds that have suitable binding and inhibitory activity can be developed further for *in vivo* use.

Agents that bind TTK can be evaluated further for TTK antagonist activity. For example, a composition comprising a TTK protein can be used in a screen or
5 binding assay to detect and/or identify agents that bind and antagonize the TTK protein. Compositions suitable for use include, for example, cells that naturally express a TTK protein (*e.g.*, one or more non-luminal breast cancer cell line described herein), extracts of such cells, and recombinant TTK protein.

An agent that binds a TTK protein can be identified in a competitive binding
10 assay, for example, in which the ability of a test agent to inhibit the binding of TTK to a reference agent is assessed. The reference agent can be a substrate, such as ATP, an endogenous substrate (*e.g.*, BLM helicase, CHK2/hCds1 protein kinase, Smad2, Smad3), or an exogenous substrate (*e.g.*, myelin basic protein) that is phosphorylated by TTK. Suitable endogenous substrates for TTK include BLM
15 helicase, CHK2/hCds1 protein kinase, Smad2 and Smad3. The reference agent can be labeled with a suitable label (*e.g.*, radioisotope, epitope label, affinity label (*e.g.*, biotin and avidin or streptavidin), spin label, enzyme, fluorescent group, chemiluminescent group, dye, metal (*e.g.*, gold, silver), magnetic bead) and the amount of labeled reference agent required to saturate the TTK protein in the assay
20 can be determined. The specificity of the formation of the complex between the TTK protein and the test agent can be determined using a suitable control (*e.g.*, unlabeled agent, label alone).

The capacity of a test agent to inhibit formation of a complex between the reference agent and a TTK protein can be determined as the concentration of test
25 agent required for 50% inhibition (IC_{50} value) of specific binding of labeled reference agent. Specific binding is preferably defined as the total binding (*e.g.*, total label in complex) minus the non-specific binding. Non-specific binding is preferably defined as the amount of label still detected in complexes formed in the presence of excess unlabeled reference agent. Reference agents suitable for use in
30 the method include molecules and compounds which specifically bind to TTK, *e.g.*, an endogenous or exogenous substrate or an antibody specific for TTK (*e.g.*, sc-540, Santa Cruz Biotechnology).

An agent that antagonizes a TTK protein can be identified by screening for agents that have an ability to antagonize (reduce, prevent, inhibit) one or more activities of TTK, such as, for example, a binding activity (*e.g.*, the binding of TTK to a substrate) or a kinase activity (*e.g.*, the phosphorylation of a substrate by TTK).

5 Such activities can be assessed using an appropriate *in vitro* or *in vivo* assay.

Exemplary *in vitro* kinase assays for TTK activity have been described previously (Stucke *et al.*, *EMBO J.* 21(7):1723-1732 (2002); Wei *et al.*, *J. Biol. Chem.* 280(9): 7748-7757 (2005)). For example, TTK protein (*e.g.*, endogenous TTK protein that is present in, or isolated from, a human cell extract, recombinant TTK protein) is
10 washed three times in TTK kinase buffer [50 mM Tris-HCl pH 7.5, 10 mM MgCl₂, 0.5 mM dithiothreitol (DTT), 10 mM β -glycerophosphate, 100 μ M Na₃VO₄].

Kinase reactions are carried out for 30 min at 30°C in TTK kinase buffer supplemented with 10 μ M ATP, 2 μ Ci of [γ ³²P]ATP (Amersham) and 0.5 mg/ml of myelin basic protein (MBP) as a substrate. Reactions are stopped by addition of gel

15 sample buffer and heating at 95°C for 5 min. Proteins are then resolved by SDS-PAGE and ³²P incorporation is visualized by autoradiography. A particularly suitable high-throughput ELISA based assay has also been described (CycLex[®] Human Mps1/TTK kinase Assay/Inhibitor Screening Kit, Catalog # CY-1179, CycLex Co., Ltd., Nagano, Japan). Exemplary assays for assessing TTK kinase
20 activity *in vivo* also have been described (See, for example, Leng *et al.*, *PNAS* 103(31):11485-11490 (2006); Zhu *et al.*, *J. Biol. Chem.* 282(25): 18327-18338 (2007)).

The ability of an agent that binds a TTK protein to antagonize one or more activities of TTK can also be assessed by measuring one or more of the TTK-
25 mediated activities. Such TTK-mediated activities include inhibition of cell death of cancer cells, inhibition of cell cycle arrest of cancer cells and anchorage-dependent cell growth *in vitro*. Thus, assays for detecting these TTK-mediated activities can be used to evaluate the antagonist activity of a test agent (*e.g.*, the ability of a test agent to inhibit one or more activities of TTK). Suitable assays include standard cell
30 proliferation assays (*e.g.*, BrdU incorporation, ³H-thymidine incorporation; See *e.g.*, Friedlander M. *et al.* *Science* 270:1500-1502, 1995; Klemke R.L., *J. Cell. Biol.* 131:791-805, 1995; Kerr J.S. *et al.*, *Anticancer Res.* 19:959-968, 1999), as well as the SRB cell viability assay, the flow cytometry assay and the colony formation

assay described herein (See Exemplification under *Materials and Methods* and Example 3).

Once a TTK antagonist is identified, the ability of the TTK antagonist to interfere with (*e.g.*, reduce, inhibit, prevent) one or more biological functions or properties associated with TTK kinase activity in a cell can be assessed, for example, using a cell-based assay designed to measure a particular biological function or property associated with TTK. Such biological functions and properties that are known to be associated with TTK expression and/or activity include, but are not limited to, centrosome duplication, mitotic checkpoint arrest, and localization to kinetocohores (Winey and Huneycutt, *Oncogene* 21: 6161-6169 (2002)); Stucke *et al.*, *EMBO J.* 21(7): 1723-1732 (2002); Fisk *et al.*, *PNAS* 100(25): 14875-14880 (2003)).

TTK antagonists are also agents that inhibit (reduce, decrease, prevent) the expression of a TTK protein. Agents (small molecules, peptides, nucleic acids, oligonucleotides) that inhibit TTK gene expression (*e.g.*, transcription, mRNA processing, translation) are effective TTK antagonists. For example, small interfering ribonucleic acids (siRNAs) and, similarly, short hairpin ribonucleic acids (shRNAs) which are processed into short siRNA-like molecules in a cell, can prevent the expression (translation) of the TTK protein. siRNA molecules can be polynucleotides that are generally about 20 to about 25 nucleotides long and are designed to bind specific RNA sequence (*e.g.*, TTK mRNA). siRNAs silence gene expression in a sequence-specific manner, binding to a target RNA (*e.g.*, an RNA having the complementary sequence) and causing the RNA to be degraded by endoribonucleases. siRNA molecules able to inhibit the expression of the TTK gene product can be produced by suitable methods. There are several algorithms that can be used to design siRNA molecules that bind the sequence of a gene of interest (see *e.g.*, Mateeva O. *et al.* *Nucleic Acids Res.* 35(8):Epub, 2007; Huesken D. *et al.*, *Nat. Biotechnol.* 23:995-1001; Jagla B. *et al.*, *RNA* 11:864-872, 2005; Shabalinea S.A. *BMC Bioinformatics* 7:65, 2005; Vert J.P. *et al.* *BMC Bioinformatics* 7:520, 2006). Expression vectors that can stably express siRNA or shRNA are available. (See *e.g.*, Brummelkamp, T.R., *Science* 296: 550-553, 2002, Lee, NS, *et al.*, *Nature Biotechnol.* 20:500-505, 2002; Miyagishi, M., and Taira, K. *Nature Biotechnol.* 20:497-500, 2002; Paddison, P.J., *et al.*, *Genes & Dev.* 16:948-958, 2002; Paul,

C.P., *et al.*, *Nature Biotechnol.* 20:505-508; 2002; Sui, G., *et al.*, *Proc. Natl. Acad. Sci. USA* 99(6):5515-5520, 2002; Yu, J-Y, *et al.*, *Proc. Natl. Acad. Sci. USA* 99(9):6047-6052, 2002; Elbashir, SM, *et al.*, *Nature* 411:494-498, 2001.). Stable expression of siRNA/shRNA molecules is advantageous in the treatment of cancer
5 as it enables long-term expression of the molecules, potentially reducing and/or eliminating the need for repeated treatments.

Antisense oligonucleotides (*e.g.*, DNA, riboprobes) can also be used as TTK antagonists to inhibit TTK expression. Antisense oligonucleotides are generally short (~13 to ~25 nucleotides) single-stranded nucleic acids which specifically
10 hybridize to a target nucleic acid sequence (*e.g.*, mRNA) and induce the degradation of the target nucleic acid (*e.g.*, degradation of the RNA through RNase H-dependent mechanisms) or sterically hinder the progression of splicing or translational machinery. (See *e.g.*, Dias N. and Stein C.A., *Mol. Can. Ther.* 1:347-355, 2002). There are a number of different types of antisense oligonucleotides that can be used
15 as TTK antagonists including methylphosphonate oligonucleotides, phosphorothioate oligonucleotides, oligonucleotides having a hydrogen at the 2'-position of ribose replaced by an *O*-alkyl group (*e.g.*, a methyl), polyamide nucleic acid (PNA), phosphorodiamidate morpholino oligomers (deoxyribose moiety is replaced by a morpholine ring), PN (N3'→P5' replacement of the oxygen at the 3'
20 position on ribose by an amine group) and chimeric oligonucleotides (*e.g.*, 2'-*O*-Methyl/phosphorothioate). Antisense oligonucleotides can be designed to be specific for a protein using predictive algorithms. (See *e.g.*, Ding, Y., and Lawrence, C. E., *Nucleic Acids Res.*, 29:1034-1046, 2001; Sczakiel, G., *Front. Biosci.*, 5:D194-D201, 2000; Scherr, M., *et al.*, *Nucleic Acids Res.*, 28:2455-2461,
25 2000; Patzel, V., *et al.* *Nucleic Acids Res.*, 27:4328-4334, 1999; Chiang, M.Y., *et al.*, *J. Biol. Chem.*, 266:18162-18171, 1991; Stull, R. A., *et al.*, *Nucleic Acids Res.*, 20:3501-3508, 1992; Ding, Y., and Lawrence, C. E., *Comput. Chem.*, 23:387-400, 1999; Lloyd, B. H., *et al.*, *Nucleic Acids Res.*, 29:3664-3673, 2001; Mir, K. U., and Southern, E. M., *Nat. Biotechnol.*, 17:788-792, 1999; Sohail, M., *et al.*, *Nucleic
30 Acids Res.*, 29:2041-2051, 2001; Altman, R. K., *et al.*, *J. Comb. Chem.*, 1:493-508, 1999). The antisense oligonucleotides can be produced by suitable methods; for example, nucleic acid (*e.g.*, DNA, RNA, PNA) synthesis using an automated nucleic acid synthesizer (from, *e.g.*, Applied Biosystems) (see also Martin, P., *Helv. Chim.*

Acta 78:486-504, 1995). Antisense oligonucleotides can also be stably expressed in a cell containing an appropriate expression vector.

Antisense oligonucleotides can be taken up by target cells (*e.g.*, tumor cells) via the process of adsorptive endocytosis. Thus, in the treatment of a subject (*e.g.*, mammalian), antisense TTK oligonucleotides can be delivered to target cells (*e.g.*, tumor cells) by, for example, injection or infusion. For instance, purified oligonucleotides or siRNA/shRNA, can be administered alone or in a formulation with a suitable drug delivery vehicle (*e.g.*, liposomes, cationic polymers, (*e.g.*, poly-L-lysine; PAMAM dendrimers, polyalkylcyanoacrylate nanoparticles and polyethyleneimine) or coupled to a suitable carrier peptide (*e.g.*, homeotic transcription factor, the Antennapedia peptide, Tat protein of HIV-1, E5CA peptide).

Ribozymes can also be used as TTK antagonists to inhibit TTK expression. Ribozymes are RNA molecules possessing enzymatic activity. One class of ribozymes is capable of repeatedly cleaving other separate RNA molecules into two or more pieces in a nucleotide base sequence specific manner. See Kim *et al.*, *Proc Natl Acad Sci USA*, 84:8788 (1987); Haseloff & Gerlach, *Nature*, 334:585 (1988); and Jefferies *et al.*, *Nucleic Acid Res*, 17:1371 (1989). Such ribozymes typically have two functional domains: a catalytic domain and a binding sequence that guides the binding of ribozymes to a target RNA through complementary base-pairing. Once a specifically-designed ribozyme is bound to a target mRNA, it enzymatically cleaves the target mRNA, typically reducing its stability and destroying its ability to directly translate an encoded protein. After a ribozyme has cleaved its RNA target, it is released from that target RNA and thereafter can bind and cleave another target. That is, a single ribozyme molecule can repeatedly bind and cleave new targets.

In accordance with the present invention, a ribozyme may target any portion of the mRNA encoding TTK. Methods for selecting a ribozyme target sequence and designing and making ribozymes are generally known in the art. See *e.g.*, U.S. Pat. Nos. 4,987,071; 5,496,698; 5,525,468; 5,631,359; 5,646,020; 5,672,511; and 6,140,491, each of which is incorporated herein by reference in its entirety. For example, suitable ribozymes may be designed in various configurations such as hammerhead motifs, hairpin motifs, hepatitis delta virus motifs, group I intron motifs, or RNase P RNA motifs. See *e.g.*, U.S. Pat. Nos. 4,987,071; 5,496,698; 5,525,468; 5,631,359; 5,646,020; 5,672,511; and 6,140,491; Rossi *et al.*, *AIDS Res*

Human Retroviruses 8:183 (1992); Hampel & Tritz, *Biochemistry* 28:4929 (1989); Hampel *et al.*, *Nucleic Acids Res*, 18:299 (1990); Perrotta & Been, *Biochemistry* 31:16 (1992); and Guerrier-Takada *et al.*, *Cell*, 35:849 (1983).

Ribozymes can be synthesized by the same methods used for normal RNA
5 synthesis. For example, suitable methods are disclosed in Usman *et al.*, *J Am Chem Soc*, 109:7845-7854 (1987) and Scaringe *et al.*, *Nucleic Acids Res*, 18:5433-5441 (1990). Modified ribozymes may be synthesized by the methods disclosed in, *e.g.*, U.S. Pat. No. 5,652,094; International Publication Nos. WO 91/03162; WO 92/07065 and WO 93/15187; European Patent Application No. 92110298.4; Perrault
10 *et al.*, *Nature*, 344:565 (1990); Pieken *et al.*, *Science*, 253:314 (1991); and Usman & Cedergren, *Trends Biochem Sci*, 17:334 (1992).

Methods of Therapy

Using the methods of the invention, cancer can be treated by inhibiting tumor growth (*e.g.*, directly inhibited) using a TTK antagonist (*e.g.*, antibodies, siRNA
15 molecules, small organic molecules, antisense oligonucleotides, chemical compounds, peptides, peptide mimetics).

Accordingly, one aspect of the invention relates to a method for treating a TTK positive non-luminal breast cancer in a mammalian subject comprising administering to the subject a therapeutically effective amount of a TTK antagonist.
20 The non-luminal breast cancer (*e.g.*, non-luminal breast cancer tumor) can be any estrogen-receptor (ER)-negative breast cancer (*e.g.*, a HER-2 positive breast cancer, a basal-like breast cancer) that expresses TTK. In a particular embodiment, the non-luminal breast cancer treated is a HER-2 positive non-luminal breast cancer. In a more particular embodiment, a TTK positive basal-like breast cancer (*e.g.*, a TTK
25 positive basal-like breast cancer tumor) is treated by administering a TTK antagonist.

In another aspect, the invention relates to a method for treating a TTK positive soft tissue sarcoma (*e.g.*, malignant tumor) in a mammalian subject comprising administering to the subject a therapeutically effective amount of a TTK
30 antagonist. The soft tissue sarcoma can be derived from any soft tissue, such as a soft tissue that connects, supports, or surrounds various structures and organs of the body, including, but not limited to, smooth muscle, skeletal muscle, tendons, fibrous

tissues, fatty tissue, blood and lymph vessels, perivascular tissue, nerves, mesenchymal cells and synovial tissues. In a particular embodiment, the soft tissue sarcoma is a fibrosarcoma, a gastrointestinal sarcoma, a leiomyosarcoma, a dedifferentiated liposarcoma, a pleomorphic liposarcoma, a malignant fibrous histiocytoma, a round cell sarcoma, or a synovial sarcoma.

In a particular aspect of the method, a TTK antagonist inhibits tumor growth directly by inducing the death (*e.g.*, apoptosis) of the cells of the tumor or by inhibiting the growth (*e.g.*, proliferation) of the cells of the tumor.

A therapeutically effective amount of the TTK antagonist is administered in the methods of the invention. In one aspect, an "anti-tumor effective amount" of a TTK antagonist is administered to a patient in need thereof. For example, agents which directly inhibit tumor growth (*e.g.*, chemotherapeutic agents) are conventionally administered at a particular dosing schedule and level to achieve the most effective therapy (*e.g.*, to best kill tumor cells). Generally, about the maximum tolerated dose is administered during a relatively short treatment period (*e.g.*, one to several days), which is followed by an off-therapy period. In a particular example, the chemotherapeutic cyclophosphamide is administered at a maximum tolerated dose of 150 mg/kg every other day for three doses, with a second cycle given 21 days after the first cycle. (Browder *et al. Can Res* 60:1878-1886, 2000). Similarly, the anti-HER-2 monoclonal antibody, trastuzumab, is administered to HER-2 positive breast cancer patients in one larger initial dose (4 mg/kg) given over period of about 90 minutes, followed by smaller weekly maintenance doses (2 mg/kg) that are administered over a shorter period of time, about 30 minutes. When administered in conjunction with other adjuvant cancer therapies (*e.g.*, chemotherapy, hormone therapy), the anti-HER-2 monoclonal antibody is administered on the same or similar cycles as the other cancer therapy.

An anti-tumor effective amount of TTK antagonist that directly inhibits the expression or activity of a TTK protein in a tumor cell (*e.g.*, inhibitory small molecules, neutralizing antibodies, inhibitory nucleic acids (*e.g.*, siRNA, antisense nucleotides)) can be administered, for example, in a first cycle in which about the maximum tolerated dose of the antagonist is administered in one interval/dose, or in several closely spaced intervals (minutes, hours, days) with another/second cycle administered after a suitable off-therapy period (*e.g.*, one or more weeks). Suitable

dosing schedules and amounts for a TTK antagonist can be readily determined by a clinician of ordinary skill. Decreased toxicity of a particular TTK antagonist as compared to chemotherapeutic agents can allow for the time between administration cycles to be shorter. When used as an adjuvant therapy (to, *e.g.*, surgery, radiation therapy, other primary therapies), an anti-tumor effective amount of a
5 TTK antagonist is preferably administered on a dosing schedule that is similar to that of the other cancer therapy (*e.g.*, chemotherapeutics), or on a dosing schedule determined by the skilled clinician to be more/most effective at inhibiting (reducing, preventing) tumor growth. A treatment regimen for an anti-tumor effective amount
10 of a TTK antagonist, for example, a small molecule, can be 0.001 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 0.01 mg/kg to about 1 mg/kg, every 1 to 7 days over a period of about 4 to about 6 months. In addition, treatment regimen for an anti-tumor effective amount of a TTK antagonist, for example, an antibody, can be from
15 about 0.01 mg/kg to about 300 mg/kg body weight per treatment and preferably from about 0.01 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 1 mg/kg to about 10 mg/kg every 1 to 7 days over a period of about 4 to about 6 months.

Accordingly, one aspect of the invention also relates to a method for directly
20 inhibiting the growth of a tumor that expresses a TTK protein comprising administering to a patient with the tumor a therapeutically effective amount (*e.g.*, an anti-tumor effective amount) of a TTK antagonist. In one embodiment, the TTK antagonist directly inhibits the growth of the tumor by inducing the apoptosis of the tumor cells or by inhibiting the proliferation of the tumor cells. The
25 TTK antagonist can inhibit the expression (*e.g.*, siRNA, antisense oligonucleotides) or activity (*e.g.*, antibody, small molecule (*e.g.*, SP600125, cincreasin), peptide, peptide mimetic) of a TTK protein, thereby directly inhibiting the growth of the cells of the tumor.

In another aspect, a TTK antagonist can be administered in a metronomic
30 dosing regime, whereby a lower dose is administered more frequently relative to maximum tolerated dosing. A number of preclinical studies have demonstrated superior anti-tumor efficacy, potent antiangiogenic effects, and reduced toxicity and

side effects (e.g., myelosuppression) of metronomic regimes compared to maximum tolerated dose (MTD) counterparts (Bocci, et al., *Cancer Res*, 62:6938-6943, (2002); Bocci, et al., *Proc. Natl. Acad. Sci.*, 100(22):12917-12922, (2003); and Bertolini, et al., *Cancer Res*, 63(15):4342-4346, (2003)). Metronomic chemotherapy appears to
5 be effective in overcoming some of the shortcomings associated with chemotherapy.

A TTK antagonist can be administered in a metronomic dosing regime to inhibit (reduce, prevent) angiogenesis in a patient in need thereof as part of an anti-angiogenic therapy. Such anti-angiogenic therapy may indirectly affect (inhibit, reduce) tumor growth by blocking the formation of new blood vessels that supply
10 tumors with nutrients needed to sustain tumor growth and enable tumors to metastasize. Starving the tumor of nutrients and blood supply in this manner can eventually cause the cells of the tumor to die by necrosis and/or apoptosis. Previous work has indicated that the clinical outcomes (inhibition of endothelial cell-mediated tumor angiogenesis and tumor growth) of cancer therapies that involve the blocking
15 of angiogenic factors (e.g., VEGF, bFGF, TGF- α , IL-8, PDGF) or their signaling have been more efficacious when lower dosage levels are administered more frequently, providing a continuous blood level of the antiangiogenic agent. (See Browder et al. *Can. Res.* 60:1878-1886, 2000; Folkman J., *Sem. Can. Biol.* 13:159-167, 2003). An anti-angiogenic treatment regimen has been used with a targeted
20 inhibitor of angiogenesis (thrombospondin1 and platelet growth factor-4 (TNP-470)) and the chemotherapeutic agent cyclophosphamide. Every 6 days, TNP-470 was administered at a dose lower than the maximum tolerated dose and cyclophosphamide was administered at a dose of 170 mg/kg. *Id.* This treatment regimen resulted in complete regression of the tumors. *Id.* In fact, anti-angiogenic
25 treatments are most effective when administered in concert with other anti-cancer therapeutic agents, for example, those agents that directly inhibit tumor growth (e.g., chemotherapeutic agents). *Id.*

Other therapies

A therapeutically effective amount of a TTK antagonist can be administered
30 alone, as an adjuvant therapy to a primary cancer therapy (surgery, radiation), with anti-angiogenic therapies (e.g., avastatin, endostatin, tumstatin, angiostatin) or as a primary therapy with other adjuvant therapies (e.g., chemotherapeutic, hormone).

The effectiveness of a therapy (*e.g.*, the reduction or elimination of a tumor, the prevention or inhibition of tumor growth, the treatment or prevention of an angiogenesis disorder, and/or the prevention or treatment of cancer metastasis) can be determined by any suitable method (*e.g.*, *in situ* immunohistochemistry, imaging
5 (MRI, NMR), ³H-thymidine incorporation).

The methods described herein comprise administering a TTK antagonist. The TTK antagonist may be administered to the individual in need thereof as a primary therapy (*e.g.*, as the principal therapeutic agent in a therapy or treatment regimen); as an adjunct therapy (*e.g.*, as a therapeutic agent used together with
10 another therapeutic agent in a therapy or treatment regime, wherein the combination of therapeutic agents provides the desired treatment; “adjunct therapy” is also referred to as “adjunctive therapy”); in combination with an adjunct therapy; as an adjuvant therapy (*e.g.*, as a therapeutic agent that is given to the subject in need thereof after the principal therapeutic agent in a therapy or treatment regimen has
15 been given); or in combination with an adjuvant therapy (*e.g.*, chemotherapy (*e.g.*, dacarbazine (DTIC), Cis-platinum, cimetidine, tamoxifen, cyclophosphamide), radiation therapy, hormone therapy (*e.g.*, anti-estrogen therapy, androgen deprivation therapy (ADT), luteinizing hormone-releasing hormone (LH-RH) agonists, aromatase inhibitors (AIs, such as anastrozole, exemestane, letrozole),
20 estrogen receptor modulators (*e.g.*, tamoxifen, raloxifene, toremifene)), or biological therapy). Numerous other therapies can also be administered during a cancer treatment regime to mitigate the effects of the disease and/or side effects of the cancer treatment including therapies to manage pain (narcotics, acupuncture), gastric discomfort (antacids), dizziness (anti-vertigo medications), nausea (anti-nausea
25 medications), infection (*e.g.*, medications to increase red/white blood cell counts) and the like, all of which are readily appreciated by the person skilled in the art.

Thus, a TTK antagonist can be administered as an adjuvant therapy (*e.g.*, with another primary cancer therapy or treatment). As an adjuvant therapy, the TTK antagonist can be administered before, after or concurrently with a primary therapy
30 like radiation and/or the surgical removal of a tumor(s). In some embodiments, the method comprises administering a therapeutically effective amount of a TTK antagonist and one or more other therapies (*e.g.*, adjuvant therapies, other targeted therapies). An adjuvant therapy (*e.g.*, a chemotherapeutic agent) and/or the one or

more other targeted therapies (*e.g.*, a PLK1 antagonist) and the TTK antagonist can be co-administered simultaneously (*e.g.*, concurrently) as either separate formulations or as a joint formulation. Alternatively, the therapies can be administered sequentially, as separate compositions, within an appropriate time frame (*e.g.*, a cancer treatment session/interval such as 1.5 to 5 hours) as determined by the skilled clinician (*e.g.*, a time sufficient to allow an overlap of the pharmaceutical effects of the therapies). The adjuvant therapy and/or one or more other targeted therapies (*e.g.*, a PLK1 antagonist) and the TTK antagonist can be administered in a single dose or multiple doses in an order and on a schedule suitable to achieve a desired therapeutic effect (*e.g.*, inhibition of tumor growth, inhibition of angiogenesis, and/or inhibition of cancer metastasis).

One or more agents that are a TTK antagonist can be administered in single or multiple doses. Suitable dosing and regimens of administration can be determined by a clinician and are dependent on the agent(s) chosen, pharmaceutical formulation and route of administration, various patient factors and other considerations. With respect to the administration of a TTK antagonist with one or more other therapies or treatments (adjuvant, targeted, cancer treatment-associated, and the like) the TTK antagonist is typically administered as a single dose (by *e.g.*, injection, infusion, orally), followed by repeated doses at particular intervals (*e.g.*, one or more hours) if desired or indicated.

The amount of the TTK antagonist to be administered (*e.g.*, a therapeutically effective amount, an anti-tumor effective amount, an anti-angiogenesis effective amount, an anti-metastasis effective amount) can be determined by a clinician using the guidance provided herein and other methods known in the art and is dependent on several factors including, for example, the particular agent chosen, the subject's age, sensitivity, tolerance to drugs and overall well-being. For example, suitable dosages for a small molecule can be from about 0.001 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 0.01 mg/kg to about 1 mg/kg body weight per treatment. Suitable dosages for antibodies can be from about 0.01 mg/kg to about 300 mg/kg body weight per treatment and preferably from about 0.01 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 1 mg/kg to about 10 mg/kg body weight per treatment. Where the TTK antagonist is a polypeptide

(linear, cyclic, mimetic), the preferred dosage will result in a plasma concentration of the peptide from about 0.1 µg/mL to about 200 µg/mL. Determining the dosage for a particular agent, patient and cancer is well within the abilities of one of skill in the art. Preferably, the dosage does not cause or produces minimal adverse side effects (e.g., immunogenic response, nausea, dizziness, gastric upset, hyperviscosity syndromes, congestive heart failure, stroke, pulmonary edema).

HER-2 positive tumor therapy

Breast cancers that are positive for HER-2 expression represent a breast cancer subtype that is characterized by rapid, aggressive tumor growth and progression, a high-grade appearance under the microscope and is associated with a poor clinical outcome and poor prognosis. Despite the negative attributes associated with HER-2 positive cancers, a number of therapeutic agents have been developed which have some efficacy in the treatment of such cancers. For example, in HER-2 positive breast cancers, a therapeutic agent that inhibited HER-2 activity (e.g., an anti-HER-2 antibody), when used in conjunction with adjuvant therapies (e.g., chemotherapy, hormone therapy) reduced the risk of cancer recurrence or death by about half (Romond E.H. *et al.* *N. Engl. J Med.* 353(16):1673-1684, 2005).

Therapeutic agents that target HER-2 expression or activity include, for example, monoclonal antibodies (e.g., trastuzumab (Herceptin®, Genetech, Inc.)), small molecule compounds and antisense HER-2 oligonucleotides. An indication for a treatment involving administration of an anti-HER-2 agent can be confirmed using any suitable methods such as fluorescence *in situ* hybridization (FISH) which can be used to detect the presence of excess copies of the HER-2 gene in the tumor cells, or immunohistochemistry. However, as shown herein, there is a strong association between TTK expression in breast tumors and HER-2 positive breast cancer (see, e.g., Example 3).

Accordingly, in one embodiment of the method for treating cancer in a mammalian subject by administering to the subject a therapeutically effective amount of a TTK antagonist further comprises administering a HER-2 antagonist, such as trastuzumab.

Another aspect of the invention provides for a method of treating a TTK positive, HER-2 positive tumor comprising administering to a mammalian subject a

therapeutically effective amount of a TTK antagonist and a therapeutically effective amount of a HER-2 antagonist, such as trastuzumab. As defined herein, HER-2 positive tumors are tumors overexpressing the HER-2 protein, generally due to the presence of extra copies of the HER-2 gene in the tumor cells. Overexpression of the HER-2 protein, which can be determined immunohistochemically in cultured, biopsied or surgical tumor tissue samples (or using other suitable methods). In a particular embodiment, the tumor is a HER-2 positive breast cancer tumor that also expresses TTK. Several types of breast cancer tumors (*e.g.*, carcinoma) can be treated with a combination of anti-HER-2 and anti-TTK antagonists, including tumors associated with ductal breast cancer, lobular breast cancer and nipple breast cancer.

Another aspect of the invention provides for a method of treating a TTK positive, HER-2 positive, estrogen receptor negative tumor comprising administering to a mammalian subject a therapeutically effective amount of a TTK antagonist and a therapeutically effective amount of a HER-2 antagonist, such as trastuzumab. In a particular embodiment, the tumor is a HER-2 positive breast cancer tumor that also expresses a TTK. Several types of breast cancer tumors (*e.g.*, carcinoma) can be treated with a combination of anti-HER-2 and anti-TTK antagonists, including ductal breast cancer, lobular breast cancer and nipple breast cancer.

In yet another aspect, the invention provides for a composition comprising a TTK antagonist and a HER-2 antagonist. In one embodiment, the composition comprises a TTK antagonist, a HER-2 antagonist and a physiologically or pharmaceutically acceptable carrier. In another embodiment, the composition comprises trastuzumab, a TTK antagonist and a physiologically or pharmaceutically acceptable carrier. In another embodiment, the composition comprises a small molecule which selectively binds a TTK protein, a HER-2 antagonist and a physiologically or pharmaceutically acceptable carrier. In yet another embodiment, the composition comprises trastuzumab, a small molecule which selectively binds a TTK protein and a physiologically or pharmaceutically acceptable carrier. Pharmaceutical compositions and formulations are discussed herein.

Basal-like breast cancer tumor therapy

Basal-like breast cancer represents another breast cancer subtype that is characterized by rapid, aggressive tumor growth and progression, a high-grade appearance under the microscope and is associated with a poor clinical outcome and poor prognosis. In general, basal-like breast cancers are estrogen receptor negative and express normal amounts of HER-2 protein, making these cancers particularly difficult to treat and diagnose. Thus, basal subtype breast cancers cannot be treated with ER-targeted or HER2 targeted therapies. As a result, basal subtype breast cancer is primarily treated with chemotherapy. Carey, L.A., *Breast Cancer Research*, 9(Suppl 1):S13 (2007). Basal-like breast cancer is common in women who have a BRCA gene mutation, in particular a BRCA1 gene mutation. Basal-like breast cancers typically express genes that are characteristic of basal breast cells, including basal cytokeratins (e.g., cytokeratin 5/6, cytokeratin 17), laminin, B4, HER1 and c-KIT, among others (Nielsen *et al.*, *Clinical Cancer Research* 10:5367-5374 (2004)). Expression of genes that are characteristic of basal breast cells can be determined immunohistochemically in cultured, biopsied or surgical tumor tissue samples (or using other suitable methods). As shown herein, there is a strong association between TTK expression and basal-like breast cancer tumor samples and cell lines (see, e.g., Example 3).

Accordingly, in one aspect, the invention provides for a method of treating a TTK positive basal-like breast cancer tumor comprising administering to a mammalian subject a therapeutically effective amount of a TTK antagonist. The method can further comprise administering one or more other therapeutic agents used for the treatment of basal-like breast cancer.

Methods for Administration

According to the methods of the invention, a therapeutically effective amount (anti-tumor effective amount, anti-angiogenesis effective amount) is administered to a mammalian subject to treat cancer. The term "mammalian subject" is defined herein to include mammals such as primates (e.g., humans) cows, sheep, goats, horses, dogs cats, rabbits, guinea pigs, rats, mice or other bovine, ovine, equine, canine feline, rodent or murine species.

TTK antagonists can be administered in single or multiple doses. Suitable dosing and regimens of administration can be determined by a practitioner and are

dependent on the agent(s) chosen, pharmaceutical formulation and route of administration, various patient factors and other considerations. With respect to the administration of a TTK antagonist with one or more other therapeutic agents (adjuvant, targeted, cancer treatment-associated), the antagonist is typically administered as a single dose (by *e.g.*, injection, infusion), followed by repeated doses at particular intervals (*e.g.*, one or more hours) if indicated or desired.

The amount of the TTK antagonist to be administered (*e.g.*, therapeutically effective amount, anti-tumor effective amount, anti-angiogenesis effective amount) can be determined by a clinician using the guidance provided herein and other methods known in the art and is dependent on several factors including, for example, the particular agent chosen, the subject's age, sensitivity, tolerance to drugs and overall well-being. For example, suitable dosages for a small molecule can be from about 0.001 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 0.01 mg/kg to about 1 mg/kg body weight per treatment. Suitable dosages for antibodies can be from about 0.1 mg/kg to about 300 mg/kg body weight per treatment and preferably from about 0.01 mg/kg to about 100 mg/kg body weight per treatment. Preferably, the dosage does not cause, or produces minimal, adverse side effects (*e.g.*, immunogenic response, nausea, dizziness, gastric upset, hyperviscosity syndromes, congestive heart failure, stroke, pulmonary edema). Where the TTK antagonist is a polypeptide (linear, cyclic, mimetic), the preferred dosage will result in a plasma concentration of the peptide from about 0.1 µg/mL to about 200 µg/mL. Determining the dosage for a particular agent, patient and cancer is well within the abilities of one of skill in the art.

A variety of routes of administration can be used including, for example, oral, dietary, topical, transdermal, rectal, parenteral (*e.g.*, intravenous, intraarterial, intramuscular, subcutaneous injection, intradermal injection), intravenous infusion and inhalation (*e.g.*, intrabronchial, intranasal or oral inhalation, intranasal drops) routes of administration, depending on the agent and the particular cancer to be treated. Administration can be local or systemic as indicated. The preferred mode of administration can vary depending on the particular agent chosen; however, oral or parenteral administration is generally preferred (*e.g.*, to administer small molecule TTK antagonists).

The TTK antagonist can be administered to a mammalian subject as part of a pharmaceutical or physiological composition, for example, as part of a pharmaceutical composition comprising an antagonist of TTK and a pharmaceutically acceptable carrier. Formulations or compositions comprising a TTK antagonist or compositions comprising a TTK antagonist and one or more other targeted therapies (*e.g.*, a HER-2 antagonist) will vary according to the route of administration selected (*e.g.*, solution, emulsion or capsule). Suitable pharmaceutical carriers can contain inert ingredients which do not interact with the TTK antagonist. Standard pharmaceutical formulation techniques can be employed, such as those described in Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA. Suitable pharmaceutical carriers for parenteral administration include, for example, sterile water, physiological saline, bacteriostatic saline (saline containing about 0.9% mg/ml benzyl alcohol), phosphate-buffered saline, Hank's solution, Ringer's lactate and the like. Formulations can also include small amounts of substances that enhance the effectiveness of the active ingredient (*e.g.*, emulsifying, solubilizing, pH buffering, wetting agents). Methods of encapsulation compositions (such as in a coating of hard gelatin or cyclodextran) are known in the art. For inhalation, the agent can be solubilized and loaded into a suitable dispenser for administration (*e.g.*, an atomizer or nebulizer or pressurized aerosol dispenser).

For example, nucleic acid-based TTK antagonists (*e.g.*, siRNAs, antisense oligonucleotides, natural or synthetic nucleic acids, nucleic acid analogs) can be introduced into a mammalian subject of interest in a number of ways. For instance, nucleic acids may be expressed endogenously from expression vectors or PCR products in host cells or packaged into synthetic or engineered compositions (*e.g.*, liposomes, polymers, nanoparticles) that can then be introduced directly into the bloodstream of a mammalian subject (by, *e.g.*, injection, infusion). Anti-TTK nucleic acids or nucleic acid expression vectors (*e.g.*, retroviral, adenoviral, adeno-associated and herpes simplex viral vectors, engineered vectors, non-viral-mediated vectors) can also be introduced into a mammalian subject directly using established gene therapy strategies and protocols (see *e.g.*, Tochilin V.P. *Annu Rev Biomed Eng* 8:343-375, 2006; Recombinant DNA and Gene Transfer, Office of Biotechnology Activities, National Institutes of Health Guidelines).

Similarly, where the agent is a protein or polypeptide, the agent can be administered via *in vivo* expression of recombinant protein. *In vivo* expression can be accomplished by somatic cell expression according to suitable methods (see, *e.g.*, U.S. Patent No. 5,399,346). Further, a nucleic acid encoding the polypeptide can
5 also be incorporated into retroviral, adenoviral or other suitable vectors (preferably, a replication deficient infectious vector) for delivery, or can be introduced into a transfected or transformed host cell capable of expressing the polypeptide for delivery. In the latter embodiment, the cells can be implanted (alone or in a barrier device), injected or otherwise introduced in an amount effective to express the
10 polypeptide in a therapeutically effective amount.

Diagnostic and Prognostic Methods

The present invention encompasses diagnostic and prognostic methods that comprise assessing expression of TTK_Δ in a suitable tumor sample from a subject. For diagnostic methods of the invention, expression of TTK_Δ by the tumor, or
15 increased expression of TTK_Δ by the tumor relative to a suitable control, indicates that the subject has a non-luminal breast cancer and/or that the subject is a candidate for an anti-cancer therapy using a antagonist and/or the subject is a candidate for an aggressive cancer therapy.

For prognostic methods of the invention, expression of TTK_Δ by the tumor, or
20 increased expression of TTK_Δ by the tumor relative to a suitable control, indicates a poor prognosis. The prognosis can be a prognosis for patient survival, a prognosis for risk of metastases and/or a prognosis for risk of relapse. As described herein, a strong association exists between high TTK expression in breast cancer samples and reduced patient survival, increased risk of metastases and increased risk of relapse
25 (see, *e.g.*, Example 2).

Suitable tumor samples for these methods include a tissue sample, a biological fluid sample, a cell(s) (*e.g.*, a tumor cell) sample, and the like. Any means of sampling from a subject, for example, by blood draw, spinal tap, tissue smear or scrape, or tissue biopsy can be used to obtain a sample. Thus, the sample
30 can be a biopsy specimen (*e.g.*, tumor, polyp, mass (solid, cell)), aspirate, smear or blood sample. The sample can be from a tissue that has a tumor (*e.g.*, cancerous growth) and/or tumor cells, or is suspecting of having a tumor and/or tumor cells.

For example, a tumor biopsy can be obtained in an open biopsy, a procedure in which an entire (excisional biopsy) or partial (incisional biopsy) mass is removed from a target area. Alternatively, a tumor sample can be obtained through a percutaneous biopsy, a procedure performed with a needle-like instrument through a small incision or puncture (with or without the aid of a imaging device) to obtain individual cells or clusters of cells (*e.g.*, a fine needle aspiration (FNA)) or a core or fragment of tissues (core biopsy). The biopsy samples can be examined cytologically (*e.g.*, smear), histologically (*e.g.*, frozen or paraffin section) or using any other suitable method (*e.g.*, molecular diagnostic methods). A tumor sample can also be obtained by *in vitro* harvest of cultured human cells derived from an individual's tissue. Tumor samples can, if desired, be stored before analysis by suitable storage means that preserve a sample's protein and/or nucleic acid in an analyzable condition, such as quick freezing, or a controlled freezing regime. If desired, freezing can be performed in the presence of a cryoprotectant, for example, dimethyl sulfoxide (DMSO), glycerol, or propanediol-sucrose. Tumor samples can be pooled, as appropriate, before or after storage for purposes of analysis. The tumor sample can be from a patient who has a TTK positive cancer, for example, a TTK positive non-luminal breast cancer (*e.g.*, a basal-like cancer, a HER-2 positive cancer).

Suitable assays can be used to assess the presence or amount of a TTK in a sample (*e.g.*, biological sample). Methods to detect a TTK protein or peptide can include immunological and immunochemical methods like flow cytometry (*e.g.*, FACS analysis), enzyme-linked immunosorbent assays (ELISA), including chemiluminescence assays, radioimmunoassay, immunoblot (*e.g.*, Western blot), and immunohistology, or other suitable methods such as mass spectroscopy. For example, antibodies to TTK can be used to determine the presence and/or expression level of TTK in a sample directly or indirectly using, for instance, immunohistology. For instance, paraffin sections can be taken from a biopsy, fixed to a slide and combined with one or more antibodies by suitable methods.

Methods to detect a TTK gene or expression thereof (*e.g.*, DNA, mRNA) include TTK nucleic acid amplification and/or visualization. To detect a TTK gene or expression thereof, nucleic acid can be isolated from an individual by suitable methods which are routine in the art (see, *e.g.*, Sambrook *et al.*, 1989). Isolated

nucleic acid can then be amplified (by *e.g.*, polymerase chain reaction (PCR) (*e.g.*, direct PCR, quantitative real time PCR, reverse transcriptase PCR), ligase chain reaction, self sustained sequence replication, transcriptional amplification system, Q-Beta Replicase, or the like) and visualized (by *e.g.*, labeling of the nucleic acid
5 during amplification, exposure to intercalating compounds/dyes, probes). TTK gene or expression thereof can also be detected using a nucleic acid probe, for example, a labeled nucleic acid probe (*e.g.*, fluorescence *in situ* hybridization (FISH)) directly in a paraffin section of a tissue sample taken from, *e.g.*, a tumor biopsy, or using other suitable methods. TTK gene or expression thereof can also be assessed by
10 Southern blot or in solution (*e.g.*, dyes, probes). Further, a gene chip, microarray, probe (*e.g.*, quantum dots) or other such device (*e.g.*, sensor, nanosensor/detector) can be used to detect expression and/or differential expression of a TTK gene.

In one embodiment, a TTK positive non-luminal breast cancer can be diagnosed by detecting expression of a TTK gene product (*e.g.*, TTK mRNA, TTK
15 protein) in a sample from a patient. Thus, the method does not require that TTK expression in the sample from the patient be compared to the expression of TTK in a control. The presence or absence of TTK can be ascertained by the methods described herein or other suitable assays. In another embodiment, an increase in expression of TTK can be determined by comparison of TTK expression in the
20 sample to that of a suitable control. Suitable controls include, for instance, a non-neoplastic tissue sample from the individual, non-cancerous cells, non-metastatic cancer cells, non-malignant (benign) cells or the like, or a suitable known or determined standard. The control can be a known or determined typical, normal or normalized range or level of expression of a TTK protein or gene (*e.g.*, an
25 expression standard). Thus, the method does not require that expression of the gene/protein be assessed in a suitable control. TTK expression can be compared to a known or determined standard.

The present invention will now be illustrated by the following Examples, which are not intended to be limiting in any way.

EXEMPLIFICATION

Material and Methods

siRNAs and reagents: The following siRNAs targeting the human TTK protein kinase gene (SEQ ID NO:1) (Genbank Accession No. M86699, see also

5 FIGS. 13A-13B) were used:

siRNA1: 5'-GCAAUACCUUGGAUGAUUA-3' (SEQ ID NO:3);

siRNA2: 5'-GGUAUUAACUGCCCAAGAA-3' (SEQ ID NO:4);

siRNA3: 5'-GCACGUGACUACUUUCAA-3' (SEQ ID NO:5);

siRNA4: 5'-GAUAAGAUCAUCCGACUUU-3' (SEQ ID NO:6).

10 Non-targeting control siRNA (siCONTROL), which targets a firefly luciferase gene sequence, was purchased from Dharmacon. Transfection reagent Lipofectamine 2000 was purchased from Invitrogen Canada, Burlington, ON, Canada. The reagents for the Sulforhodamine B (SRB) cell viability assay were from Sigma Canada, Oakville, ON, Canada. TTK antibody (C-19) was purchased from Santa Cruz
15 Biotechnology, Inc., and β -actin antibody (A2066) was purchased from Sigma-Aldrich Co.

TTK QPCR: Total RNA (tRNA) was prepared using Trizol Reagent (Invitrogen Corporation) according to manufacturer's instructions. tRNA was DNase treated at room temperature for 15 min. Primers were designed to amplify
20 short segments of target cDNA (amplicon) with a size range of 100-150 bp. Quantitative real time PCR (QRT-PCR) reactions were performed using One Step SYBR QPCR Kit (Invitrogen Corporation, Cat #117146-100) according to manufacturer's instructions. Direct detection of the PCR product was monitored by measuring the increase in fluorescence caused by the binding of SYBR Green I dye
25 to the double-stranded (ds) DNA product using the RealPlex4 System (Eppendorf). The conditions for the assay were 53°C for 20 min, 95°C for 5 min, followed by 42 cycles of 95°C for 15 sec, 60°C for 30 sec and 72°C for 30 sec. All PCR efficiencies were above 95%. Relative quantification of gene expression was performed using the standard curve method comprising six serial dilution points
30 (ranging from 0.32 ng to 1000 ng). Each tRNA sample was analyzed by calculating the average C_t values from triplicate PCR reactions. β -Actin gene expression was used for data normalization. Data were expressed as a mean $\pm$ SD. Amplification primers for TTK are 5'-TGATGGCAAACAACCCAGAGG-3' (forward) (SEQ ID

NO:7) and 5'-TTGCTTGACTGTAACGACCAAT-3' (reverse) (SEQ ID NO:8), and β -Actin are 5'-GGCACTCTTCCAGCCTTCCTT-3' (forward) (SEQ ID NO:9) and 5'-TCTCCTTCTGCATCCTGTCTG-3' (reverse) (SEQ ID NO:10).

5 *TTK Western*: Whole cell lysates (40 μ g) were resolved by SDS-PAGE and transferred to nitrocellulose membranes (Invitrogen Corporation) for immunoblot analysis using TTK (1/500) and β -actin (1 μ g/ml) rabbit polyclonal antibodies. For the detection, peroxidase-Goat Anti-Rabbit IgG (1/5000; Amersham Biosciences) conjugated antisera was used. The proteins were visualized using ECL (Pierce).

10 *TTK Microarray expression profiling*: All gene expression analyses were done in three individual datasets and in a combined dataset. The datasets were combined after normalizing expression values, e.g. logratios, of each gene by mean and standard deviation in each dataset. Regression models in the combined dataset contained a dataset term in addition to the logratio term. To verify that results did not depend on normality assumption, another combined dataset was assembled using
15 logratio ranks, computed separately in each dataset and scaled to vary from 0 to 1. The same analyses were performed and general agreement with the results from the normalized log ratio dataset was verified.

Transfection of siRNAs into normal breast and breast cancer cell lines: Different normal breast and breast cancer cell lines were used for siRNA
20 transfection. Cells were seeded at various concentrations, ranging from 1500 to 6000 per well according to cell growth rate, into 96 well plates. 40nM individual siRNAs or 40nM siRNA pool including four individual siRNAs at 10nM each were transfected into cells using Lipofectamine2000 24 hrs after cell seeding. Cells were then incubated at 37°C for five days before cell viability assay were conducted.

25 *Sulforhodamine B (SRB) assay*: SRB assay was performed to assess cell survival. SRB is a water-soluble dye that binds to the basic amino acids of the cellular proteins. Thus, colorimetric measurement of the bound dye provides an estimate of the total protein mass and is related to the cell number. The cells were fixed *in situ* by gently aspirating off the culture media and adding 50 μ l ice cold
30 10% Tri-chloroacetic Acid (TCA) per well and incubate at 4°C for 30-60 min. The plates were washed with tap water five times and allowed to air dry for 5 min. 50 μ l of 0.4%(w/v) Sulforhodamine B solution in 1% (v/v) acetic acid was added per well and incubated for 30min at RT for staining. Following staining, plates were washed

four times with 1% acetic acid to remove any unbound dye and then allowed to air dry for 5 min. The stain was solubilized with 100 μ l of 10 mM Tris pH 10.5 per well. Absorbance was read at 570 nm. The cell survival percentage after each siRNA(s) knock down was calculated over the non-silencing control siCTRL or siPOOL as well as siTOX as a transfection efficiency control.

Colony formation in soft agar: Colony formation in soft agar: Cells were transfected with 40 nM individual siRNAs, 40 nM siRNA pool including four individual siRNAs at 10 nM each or 40 nM non-silencing control siCONTROL using Lipofectamine 2000 Transfection Reagent (Invitrogen Corporation), mixed with culture medium containing 0.7% agar in 6-well plates and cultured at 37°C. After 2 weeks, the top layer of the culture was stained with 0.2% p-iodonitrotetrazolium violet and colonies were counted using a Sorcerer Colony Counter (Optomax).

SP600125 experiment: Compound SP600125 was purchased from Biomol International (PA, USA). 10mM stock was prepared in 100% DMSO. Cells were seeded at various numbers, ranging from 1500 to 4000 per 80 μ l in each well according to cell growth rate, into 96 well plates, 24 hours before compound overlay. The 10mM SP600125 compounds stock in 100% DMSO was diluted with Opti-MEM I Reduced-Serum Medium (Invitrogen, Burlington, ON, Canada) to concentrations range from 50 nM to 250 μ M. 20 μ l from each concentration was overlay to cells to make the final concentrations range from 10 nM to 50 μ M. The cells were cultured for 5 days before Sulforhodamine B assay. SRB assay is performed to assess cell survival. SRB is a water-soluble dye that binds to the basic amino acids of the cellular proteins. Thus, colorimetric measurement of the bound dye provides an estimate of the total protein mass that is related to the cell number. the cells are fixed in situ by gently aspirating off the culture media and adding 50 μ l ice cold 10% Tri-chloroacetic Acid (TCA) per well and incubate at 4°C for 30-60 min, The plates are washed with tap water five times and allowed to air dry for 5 min. Add 50 μ l 0.4%(w/v) Sulforhodamine B solution in 1% (v/v) acetic acid per well and incubate for 30min at RT for staining. Following staining, plates are washed four times with 1% acetic acid to remove any unbound dye and then allowed to air dry for 5 min. Stain is solubilized with 100ul of 10mM Tris pH 10.5 per well. Absorbance is read at 570 nm. The cell survival percentage in each compound

concentration was calculated over the cell DMSO control, which was growing in the presence of 0.1% DMSO (0.5% in the case of 50 μ M compounds). The GI_{50} were calculated using GraphPad PRISM software.

EXAMPLE 1: TTK expression in normal and cancer cells

5 Microarray expression analysis revealed that expression of TTK RNA in physiologically normal human tissues is restricted to fetal liver, placenta, testis, thymus and uterus (FIG. 1). Little detectable expression was observed in normal breast tissue. However, TTK RNA (FIG. 2A) and protein (FIG. 2B) was found to be expressed in several different breast cancer cell lines, with non-luminal breast cancer
10 cell lines displaying the highest levels of TTK protein (FIG. 2B).

These results indicate that TTK is a target for breast cancer therapy.

EXAMPLE 2: High TTK expression is associated with poor prognosis in breast cancer patients

 The association between TTK expression and survival in patients with breast
15 cancer was evaluated using three data sets (NEJM 295 (Hu, *et al.*, The molecular portraits of breast tumors are conserved across microarray platforms. *BMC Genomics* 7:96 (2006)), Perou (van di Vijver, M.J., *et al.*, A gene-expression signature as a predictor of survival in breast cancer. *N Engl. J. Med.* 347(25):1999-2009 (2002)), and PNAS (Miller, L.D., *et al.* An expression signature for p53 status
20 in human breast cancer predicts mutation status, transcriptional effects, and patient survival. *Proc. Natl. Acad. Sci. U S A* 102(38):13550-5) (2005)). A strong association between high TTK expression and reduced survival was discovered in each of the three data sets (FIG. 6). The Cox regression p value for the combined data set was 3.7e-08.

25 In addition, the association between TTK expression and risk of developing metastases was also evaluated in patients with breast cancer. High TTK expression was associated with an increased risk of metastasis (FIG. 7). The Cox regression p value for the data set was 0.014.

 The association between TTK expression and risk of relapse was also
30 evaluated using two data sets (NEJM295, Perou) (FIGS. 8A-8B). High TTK expression was associated with an increased risk of breast cancer relapse in the

NEJM295 data set (FIG. 8A). The Cox regression p value for this data set was 0.00059.

Survival analysis was performed using Cox regression of survival times vs. gene expression log ratios, followed by the estimation of false discovery rates by Benjamini and Hochberg's method. NEJM 295 dataset provided data on recurrence and survival; Perou dataset on recurrence, metastasis, and survival; PNAS dataset on survival only; the combined dataset could be analyzed for survival, i.e., risk of death, only. High expression of TTK was associated with higher risks of recurrence, metastasis, and death in all datasets where such data were available, as well as in the combined dataset ($P < 0.05$). These associations were observed in all cancers combined and in ER-positive cancers as a subgroup. A strong association of TTK expression with poor survival was also observed in p53⁻ cancers in the PNAS dataset ($P < 0.05$).

Tumor gene expression profiles were classified into subtypes defined by Hu et al. (2006). The classification of the NEJM 295 dataset was provided by its authors; Perou and PNAS datasets were classified into Luminal A, Luminal B, Basal-like and HER2+/ER- subtypes using the intrinsic gene set defined by Hu et al. Different subtypes were compared to each other and, where available, to normal controls.

NEJM 295 dataset was the only one that contained a normal control group. All cancers as a group and individual subtypes were compared to the control group using t tests. Significantly higher than normal expression of TTK, with p values < 0.05 and false discovery rates < 0.1 , was detected in all cancers as a group and in Basal-like, Luminal B, and HER2+/ER- subtypes. The over-expression was the highest and most significant in the basal-like subtype, where it was, on average, 8-fold higher than normal.

An additional analysis was performed on the NEJM 295 dataset to determine if a fraction of cancers expressed a gene significantly outside of the normal range. This analysis was designed to detect over-expression in a subset of a potentially heterogeneous population of cancers. The normal range was defined as the mean $\pm$ 3 standard deviations of the normal control group. The fractions of tumor samples that fell above and below this range were recorded as percentages of the total

number of tumors, in all cancers as a group and in individual subtypes. TTK was found to be over-expressed in 38% of all tumors and in 85% of basal-like tumors.

To verify over-expression of TTK in cancers compared to normal controls, we performed similar analyses in an additional dataset (Richardson *et al.*, X
5 chromosomal abnormalities in basal-like human breast cancer. *Cancer Cell* 9: 121-132 (2006)) that contained a normal control group. In the Richardson dataset, TTK appeared to be over-expressed, on average, 20-fold in all cancers and 37-fold in basal-like cancers; 100% of basal-like cancers expressed TTK above the normal range.

10 Different tumor subtypes were compared to each other using one way ANOVA and pair-wise t tests in all three datasets. There were statistically significant differences in TTK expression between different tumor subtypes in all 3 datasets, with estimated false discovery rates in 10^{-19} - 10^{-31} range for the ANOVA p values. Expression was highest in basal-like tumors and lowest in normal controls
15 in NEJM 295 dataset. Therefore, higher TTK expression was associated with more aggressive subtypes of breast cancer.

EXAMPLE 3: Validation of TTK as a target for breast cancer therapy

To validate TTK as a target for breast cancer therapy, the biological effects of TTK antagonists were investigated.

20 A. siRNA antagonists

TTK expression was inhibited using RNA interference. MD-MB-468 breast cancer cells were transfected with a 40 nM concentration of one of four individual siRNAs that target different TTK mRNA sequences (siRNA #1, siRNA #2, siRNA #3, siRNA #4), a pool of the four TTK-targeting siRNAs (siRNA POOL), or a non-
25 targeting control siRNA (siCONTROL) that targets firefly luciferase mRNA. Total RNA was isolated 48 hours post-transfection and TTK mRNA levels were determined by quantitative reverse-transcriptase polymerase chain reaction. TTK mRNA levels were normalized over beta-actin mRNA levels. Of the four TTK-targeting siRNAs, siRNAs #2 and #3 showed greater than 70% knockdown
30 efficiency (FIG. 9A).

In addition, Western blot analysis of TTK protein levels in lysates of MD-MB-468 breast cancer cells that had been transfected with siRNA revealed that

transfection with any of the four TTK-targeting siRNAs, as well as the siRNA POOL, but not the siCONTROL siRNA, resulted in significantly reduced levels of TTK protein (FIG. 9B).

To assess the effects of TTK knockdown on the viability of breast cancer
5 cells, various breast cancer cell lines were transfected with each of the four TTK-
targeting siRNAs individually, the pooled siRNA containing all four of the TTK-
targeting siRNAs, or the siCONTROL siRNA, all at a concentration of 40 nM. At
day 5 post-transfection, the viability of the cells was determined using the SRB
assay described herein. TTK expression levels post-transfection with the targeting
10 siRNA were normalized over TTK levels following transfection with the non-
silencing control. siRNA-mediated depletion of TTK levels substantially inhibited
the viability of breast cancer cells in 7 out of 12 different breast cancer cell lines,
including 3 basal-like breast cancer cell lines (Table 1).

Table 1. TTK knockdown inhibits viability of several different breast cancer cell lines (% survival).

	MDA-MB-468	MCF-7	MDA-MB-435	SKBr-3	CAMA-1	Hs578T	MDA-MB-231	T-47D	HCC 1419	MDA-MB-453	BT 474	MDA-MB-330
siRNA #1	0.69	<i>0.27</i>	<i>0.78</i>	<i>0.32</i>	<i>0.46</i>	<i>1.02</i>	<i>0.21</i>	<i>0.62</i>	<i>0.91</i>	<i>0.33</i>	<i>0.64</i>	<i>0.31</i>
siRNA #2	<i>0.00</i>	<i>0.27</i>	<i>0.25</i>	<i>0.34</i>	<i>0.30</i>	<i>1.25</i>	<i>-0.09</i>	<i>0.44</i>	<i>0.75</i>	<i>-0.04</i>	<i>0.66</i>	<i>0.38</i>
siRNA #3	<i>0.27</i>	<i>0.23</i>	<i>0.60</i>	<i>0.49</i>	<i>0.52</i>	<i>0.53</i>	<i>0.23</i>	<i>0.74</i>	<i>0.91</i>	<i>0.18</i>	<i>0.83</i>	<i>0.63</i>
siRNA #4	<i>1.46</i>	<i>0.63</i>	<i>0.93</i>	<i>0.91</i>	<i>0.77</i>	<i>1.16</i>	<i>0.58</i>	<i>1.19</i>	<i>1.05</i>	<i>0.56</i>	<i>0.90</i>	<i>0.72</i>
siRNA POOL	<i>0.34</i>	<i>0.28</i>	<i>0.57</i>	<i>0.48</i>	<i>0.34</i>	<i>0.65</i>	<i>0.24</i>	<i>0.78</i>	<i>0.97</i>	<i>0.66</i>	<i>0.98</i>	<i>0.55</i>

TTK expression levels were normalized over siCONTROL. Basal cell lines are MDA-MB-469, MDA-MB-435, Hs578T, and MDA-MB-231.

Numbers in italics signify inhibition of greater than 50%.

To assess the effects of TTK knockdown on the physiology of normal breast and breast cancer cell lines, luminal (MCF7) or basal-like (MDA-MB-435, MDA-MB-468) breast cancer cell lines, a lung cancer cell line (A549) and a normal breast cell line (184A1) each were transfected with 40 nM of TTK-targeting siRNA #2, siRNA #3, the pooled siRNA containing all four of the TTK-targeting siRNAs, or the siCONTROL siRNA. At day 3 post-transfection, the cell cycle profile of the transfected cells was analyzed by flow cytometry to determine the effects of TTK knockdown on cell cycle arrest and cell death. The results are presented in Table 2 and FIG. 10. Depletion of TTK in cells transfected with TTK-targeting siRNAs caused significant cell cycle arrest and cell death.

To determine the effects of siRNA-mediated knockdown of TTK gene expression on anchorage-independent cell growth in breast cancer cells, T-47D breast cancer cell line was transfected with either the siRNA#2 or siRNA#3 TTK-targeting siRNA, the siRNA POOL containing 4 TTK-targeting siRNAs or the siCONTROL non-targeting siRNA. Following transfection colony formation in soft agar was assessed. Transfection of T-47D cells with siRNA#2, siRNA#3 or the siRNA POOL resulted in a reduction in both colony size and number, relative to the siCONTROL-transfected cells, in two independent experiments (FIGS. 11A-11B).

Table 2. TTK knockdown induces cell cycle arrest and cell death in some cancer cell lines, but not in normal cell lines.

<u>CELL LINE</u>	<u>siCONTROL</u> (Neg. Control)		<u>TTK siRNA #2</u>		<u>TTK siRNA #3</u>	
	G2/M	<G1	G2/M	<G1	G2/M	<G1
MCF-7 (Luminal)	16.17	2.11	25.18	12.03	65.23	5.61
MDA- MB-435 (Basal)	18.77	0.85	19.89	4.69	40.08	2.66
MDA- MB-468 (Basal)	20.59	4.09	15.72	15.10	28.41	11.12
184A1 (Normal)	10.35	1.03	27.75	5.35	17.49	3.92
A549 (Lung)	10.57	0.84	20.58	6.03	24.87	11.38

B. Small molecule antagonist

To assess the effects of inhibition of TTK activity on the viability of breast cancer cell lines, varying concentrations of SP600125, a small molecule TTK antagonist (Schmidt *et al.*, *EMBO Reports* 6(9): 866-872 (2005)), in 0.1% DMSO, were added to the culture media of several different breast cancer cell lines 24 hr after the cells were seeded. Cell viability was assessed using the SRB assay described herein, which was performed 5 days after the addition of SP600125 to the culture media. Prism software was used to calculate GI₅₀ (concentration of the compound required to achieve 50% growth inhibition) (Table 3). Treatment of multiple cancer cell lines with compound SP600125 inhibited proliferation of cancer cells in a dose dependent manner with a GI₅₀ at low μ M levels, suggesting TTK antagonists have anti-cancer activity.

Table 3. Concentrations of compound SP600125 required to achieve 50% growth inhibition (GI₅₀) for various cell lines.

Cell line	CAMA-1	Hs578T	MCF-7	MDA-MB-231	MDA-MB-435	MDA-MB-468	SKBR-3
GI ₅₀ (μ M)	N/A	1.839	0.6715	0.8938	0.8225	0.4834	0.8119
Cell line	T47D	HCC1419	MDA-MB-453	MDA-MB-330	BT474	184A	
GI ₅₀ (μ M)	2.516	10.04	2.299	7.016	8.585	1.59	

EXAMPLE 4: TTK overexpression in basal-like breast cancer cells

The inventor have discovered an association between TTK overexpression and basal-like breast cancers and soft-tissue sarcomas. In particular, overexpression of TTK was detected in non-luminal (*e.g.*, basal-like, non-basal-like, HER-2 positive/ER-negative) breast cancer cells more frequently than in luminal (*e.g.*, luminal A, luminal B) or non-malignant breast cells (FIG. 3). Of the non-luminal breast cancer cell lines evaluated, TTK overexpression was detected most frequently in basal-like breast cancer cells (Table 4), which also displayed the highest levels of TTK protein expression (FIG. 4).

Table 4. TTK is overexpressed in different breast cancer subtypes

	% ↑ OVERALL	% ↑ BASAL	% ↑ HER2/ER-	% ↑ LUMINAL A	% ↑ LUMINAL B
TTK <i>6q14.1</i>	38	85	38	6	18

Perou Dataset: % increase in tumors vs. normal samples.

Microarray expression analysis revealed elevated TTK gene expression in several soft-tissue sarcoma samples derived from various tissues, as compared to TTK gene expression in physiologically normal tissues (FIG. 5).

These results indicate that TTK is a target for cancer therapy, particularly basal-like breast cancer therapy and soft tissue sarcoma therapy.

EXAMPLE 5: RNAi silencing of TTK suppresses tumor growth *in vivo*

10

A double-stranded oligonucleotide encoding a human TTK gene-specific shRNA (sense insert sequence 5'-GCAGUCAUGCCCAUUUGGAA-3' (SEQ ID NO:11)) was ligated into the RNAi-Ready-pSIREN-RetroQ-ZsGreen retroviral vector (Clontech). Amphotropic-Phoenix packaging cells (ATCC, Manassas, VA) were transiently transfected with either control RNAi-Ready-pSIREN-RetroQ-ZsGreen-shLUC (Clontech) or RNAi-Ready-pSIREN-RetroQ-ZsGreen-shTTK using FuGENE 6 transfection reagent (Roche Diagnostics, Indianapolis, IN). Culture supernatants were collected 2 days after transfection and filtered through 0.45-μm pore-size filters. MDA-MB-468 breast cancer cells (ATCC, Manassas, VA) were infected with retroviruses by culturing the cells for 24 hours in 1:1 Phoenix conditioned media (Dulbecco's Modified Eagle's Media, 10% FCS, supplemented with 8 μg/ml Polybrene; Sigma-Aldrich). This transfection process was repeated three times to increase the transfection efficiency. One day after the final infection, the RNAi-Ready-pSIREN-RetroQ-ZsGreen-shLUC and RNAi-Ready-pSIREN-RetroQ-ZsGreen-shTTK infected cancer cells were trypsinized, counted and injected subcutaneously into the left and right hindlimb, respectively, of nude mice at concentrations of 2.5×10^6 cells (5 mice per group). The infected cells were also analyzed for TTK expression by RT-PCR using TTK-specific primers. The tumors were measured and viable tumor area was calculated twice weekly for approximately 10 weeks.

Infection of MDA-MB-468 cells with the TTK shRNA-encoding construct led to the reduction of TTK expression in these cells by approximately 50% compared to that of the control cells (FIG. 17A). Furthermore, reduction of TTK mRNA levels resulted in a significant suppression of tumor growth in a mouse xenograft model (FIG. 17B). Taken together, these results suggest that inhibition of TTK activity in cancer cells inhibits tumor growth in tumors that express TTK.

The teachings of all patents, published applications and references cited herein are incorporated by reference in their entirety.

While this invention has been particularly shown and described with references to example embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

CLAIMS

What is claimed is:

1. A method for treating TTK positive non-luminal breast cancer in a mammalian subject, comprising administering to the subject a therapeutically effective amount of a TTK antagonist.
5
2. The method of Claim 1, wherein said cancer is selected from the group consisting of basal-like breast cancer and HER-2 positive breast cancer.
3. The method of Claim 1, wherein said TTK antagonist is selected from the group consisting of an antibody, an antigen-binding fragment of an antibody,
10 a small interfering ribonucleic acid (siRNA), a peptide, a peptidomimetic, an antisense oligonucleotide, and a small molecule.
4. The method of Claim 1, further comprising administering one or more other therapeutic agents.
5. The method of Claim 4, wherein said non-luminal breast cancer is a HER-2
15 positive breast cancer and a HER-2 antagonist is also administered.
6. The method of Claim 5, wherein said HER-2 antagonist is trastuzumab.
7. The method of Claim 5, wherein said HER-2 positive breast cancer is estrogen receptor (ER) negative.
8. The method of Claim 1, wherein said mammalian subject has a tumor.
- 20 9. The method of Claim 8, wherein said tumor is a metastatic tumor.
10. The method of Claim 8, wherein said tumor is a HER-2 positive tumor.
11. The method of Claim 10, wherein said HER-2 positive tumor is estrogen receptor (ER) negative.

12. A method for treating a TTK positive soft tissue sarcoma in a mammalian subject, comprising administering to the subject a therapeutically effective amount of a TTK antagonist.
13. The method of Claim 12, wherein said sarcoma is selected from the group consisting of a fibrosarcoma, a leiomyosarcoma, a dedifferentiated liposarcoma, a pleiomorphic liposarcoma, a malignant fibrous histiocytoma, a round cell sarcoma and a synovial sarcoma.
14. The method of Claim 12, wherein said TTK antagonist is selected from the group consisting of an antibody, an antigen-binding fragment of an antibody, a small interfering ribonucleic acid (siRNA), a peptide, a peptidomimetic, an antisense oligonucleotide, and a small molecule.
15. The method of Claim 12, further comprising administering one or more other therapeutic agents.
16. A method of treating a TTK positive basal-like breast cancer tumor in a mammalian subject comprising administering to the subject a therapeutically effective amount of a TTK antagonist.
17. The method of Claim 16, wherein the TTK antagonist is selected from the group consisting of an antibody, an antigen-binding fragment of an antibody, a small interfering ribonucleic acid (siRNA), a peptide, a peptidomimetic, an antisense oligonucleotide, and a small molecule.
18. The method of Claim 16, wherein said tumor is a metastatic tumor.
19. The method of Claim 16, wherein said tumor is a carcinoma.
20. The method of Claim 16, further comprising administering one or more other therapeutic agents.
21. A method of treating a TTK positive breast cancer in a mammalian subject, wherein said breast cancer is HER-2 positive and ER negative, comprising

administering to the subject a therapeutically effective amount of a TTK antagonist.

22. The method of Claim 21, wherein the TTK antagonist is selected from the group consisting of an antibody, an antigen-binding fragment of an antibody,
5 a small interfering ribonucleic acid (siRNA), a peptide, a peptidomimetic, an antisense oligonucleotide, and a small molecule.
23. The method of Claim 21, further comprising administering a HER-2 antagonist.
24. The method of Claim 23, wherein the HER-2 antagonist is trastuzumab.
- 10 25. A composition comprising a TTK antagonist, a HER-2 antagonist and a physiologically acceptable carrier.
26. The composition of Claim 25, wherein said HER-2 antagonist is trastuzumab.
- 15 27. A method for identifying a candidate for an anti-cancer therapy using a TTK antagonist comprising:
a) providing a suitable sample obtained from a subject with a tumor;
and
b) assessing expression of TTK in said sample,
wherein expression of TTK in said sample, or increased expression of
20 TTK in said sample relative to a suitable control, indicates that the subject is a candidate for an anti-cancer therapy using a TTK antagonist.
28. The method of Claim 27, wherein the sample is a tumor sample.
29. A method of diagnosing a non-luminal breast cancer in a subject, comprising:
25 a) providing a suitable sample from a subject with a breast tumor; and
b) assessing expression of TTK in said sample,

wherein expression of TTK in said sample, or increased expression of TTK in said sample relative to a suitable control, indicates that the subject has a non-luminal breast cancer.

30. The method of Claim 29, wherein the sample is a tumor sample.
- 5 31. A method of determining the prognosis of a subject having a cancer, comprising:
- a) providing a suitable sample from a subject with a tumor; and
- b) assessing expression of TTK in said sample,
- 10 wherein expression of TTK in said sample, or increased expression of TTK in said sample relative to a suitable control, indicates a prognosis selected from the group consisting of reduced patient survival, increased risk of metastases and increased risk of relapse.
32. The method of Claim 31, wherein the sample is a tumor sample.
33. A method for screening a breast cancer patient as an aid for selecting
- 15 aggressive cancer therapy of said patient, comprising:
- (a) providing a suitable sample from said patient; and
- (b) determining TTK expression in said sample;
- wherein increased TTK expression in said sample as compared with a suitable control indicates the patient is a candidate for aggressive cancer
- 20 therapy of said patient.
34. The method of Claim 33, wherein the sample is a tumor sample.






FIG. 1






FIG. 3




FIG. 4




FIG. 5

FIG. 6




FIG. 7




FIG. 8A







FIG. 8B





siControl




FIG. 10A

TTK siRNA #2




FIG. 10B

TTK siRNA #3





FIG. 10C






FIG. 12




FIG. 13A




FIG. 13B




FIG. 13C




FIG. 14A




FIG. 14B




FIG. 14C

15/19

```

1   ggaattcctt tttttttttt tttgagatgg agtttcactc ttgttgGCCa ggctggagtG
61  caatggcaca atctcagctt actgcaacct cgcctcccGg ggttcaagcg attctcctgc
121 ctcagcctct caagtagctg ggattacagg catgtgccac caccctggc taactaattt
181 cttttctatt tagtagagat ggggtttcac catgttggtc aggtgggtct tgaactcctg
241 acctcaggtg atccacttgc cttggcctcc caaagtgcta ggattacagc cgtgaaactg
301 tgccctggctg attctttttt tgttggttga tttttgaaac agggctctccc ttggtcgcgc
361 aggtctggagt gcagtgggtg gatcttggct cactataacc tccacctcct ggtttcaagt
421 gatcctccca ctttagcctc ctgagtagct gtgattacag gcgtgcacca ccacaccggg
481 ctaatttttg tatttttatt agagacaggg tttcaccatg ttggccaggc tgttctcaaa
541 ctccctggact caagggatcc gcctgcctcc acttcccaaa gtcccagatg tacaggtgtg
601 agtcaccatg cctgacctta taattcttaa gtcatttttt ctggtccatt tcttccttag
661 ggctctcaca acaaactctg attaggcggg acaataatcc ttaacttcat gattcacaaa
721 aggaagatga agtgattcat gatttagaaa ggggaagtag taagccact gcacactcct
781 ggatgatgat cctaaatcca gatacagtaa aaatggggtg tgggaaggta gaatacaaaa
841 tttgggttaa attaattatc taaatatcta aaaacatttt tggatacatt gttgatgtga
901 atgtaagact gtacagactt cctagaaaac agtttgggtt ccatcttttc atttcccag
961 tgcagttttc tgtagaaatg gaatccgagg atttaagtgg cagagaattg acaattgatt
1021 ccataatgaa caaagtgaga gacattaaaa ataagttaa aaatgaagac cttactgatg
1081 aactaagctt gaataaaatt tctgctgata ctacagataa ctcggaact gttaacaaaa
1141 ttatgatgat ggcaacaac ccagaggact ggttgagttt gttgctcaaa ctagagaaaa
1201 acagtgttcc gctaagtgat gctcttttaa ataaattgat tggctgttac agtcaagcaa
1261 ttgaagcgct tccccagat aaatatggcc aaaatgagag ttttgctaga attcaagtga
1321 gatttgctga attaaaagct attcaagagc cagatgatgc acgtgactac tttcaaatgg
1381 ccagagcaaa ctgcaagaaa tttgcttttg ttcatatatc ttttgcaaa tttgaactgt
1441 cacaaggtaa tgtcaaaaaa agtaaacaac ttcttcaaaa agctgtagaa cgtggagcag
1501 taccactaga aatgctggaa attgccctgc ggaatttaaa cctccaaaaa aagcagctgc
1561 tttcagagga ggaaaagaag aatttatcag catctacggt attaactgcc caagaatcat
1621 tttccggttc acttgggcat ttacagaata ggaacaacag ttgtgattcc agaggacaga
1681 ctactaaagc cagggttttta tatggagaga acatgccacc acaagatgca gaaataggtt
1741 accggaattc attgagacaa actaacaaaa ctaaacagtc atgcccattt ggaagagtcc
1801 cagttaacct tctaaatagc ccagattgtg atgtgaagac agatgattca gttgtacctt
1861 gttttatgaa aagacaaaacc tctagatcag aatgccgaga tttggttggt cctggatcta
1921 aaccaagtgg aaatgattcc tgtgaattaa gaaatttaaa gtctgttcaa aatagtcatt
1981 tcaaggaacc tctggtgtca gatgaaaaga gttctgaact tattattact gattcaataa
2041 ccctgaagaa taaaacggaa tcaagtcttc tagctaaatt agaagaaact aaagagtatc
2101 aagaaccaga ggttccagag agtaaccaga aacagtggca agctaagaga aagtcagagt
2161 gtattaacca gaatcctgct gcattctcaa atcactggca gattccggag ttagcccgaa
2221 aagttaatac agagcagaaa cataccactt ttgagcaacc tgtcttttca gtttcaaaac
2281 agtcaccacc aatatcaaca tctaaatggg ttgacccaaa atctatttgt aagacaccaa
2341 gcagcaatac cttggatgat tacatgagct gttttagaac tccagttgta aagaatgact
2401 ttccacctgc ttgtcagttg tcaacacctt atggccaacc tgccgtttc cagcagcaac
2461 agcatcaaat acttgccact ccacttcaaa atttacaggt tttagcatct tcttcagcaa
2521 atgaatgcat ttcggttaaa ggaagaattt attccatatt aaagcagata ggaagtggag
2581 gttcaagcaa ggtatttcag gtgttaaag aaaagaaaca gatatatgct ataaaatatg
2641 tgaacttaga agaagcagat aaccaaactc ttgatagtta ccggaacgaa atagcttatt
2701 tgaataaact acaacaacac agtgataaga tcatccgact ttatgattat gaaatcacgg
2761 accagtacat ctacatggta atggagtgtg gaaatattga tcttaatagt tggcttaaaa
2821 agaaaaaatc cattgatcca tgggaacgca agagttactg gaaaaatatg ttagaggcag
2881 ttcacacaat ccatcaacat ggcattgttc acagtgatct taaaccagct aactttctga
2941 tagttgatgg aatgctaaag ctaattgatt ttgggattgc aaaccaaatg caaccagata
3001 caacaagtgt tgttaaagat tctcaggttg gcacagttaa ttatatgcca ccagaagcaa
3061 tcaaaagatat gtcttcctcc agagagaatg ggaaatctaa gtcaaagata agccccaaaa
3121 gtgatgtttg gtccttagga tgtattttgt actatatgac ttacgggaaa acaccatttc
3181 agcagataat taatcagatt tctaaattac atgccataat tgatcctaatt catgaaattg
3241 aatttcccga tattccagag aaagatcttc aagatgtgtt aaagtgtgtt taaaaaggg

```

FIG. 15A

16/19

```
3301 acccaaaaca gaggatatcc attcctgagc tcctgggtca tccatatggtt caaattcaaa
3361 ctcatccagt taaccaaatg gccaagggaa ccactgaaga aatgaaatat gttctgggcc
3421 aacttggttg tctgaattct cctaactcca ttttgaaagc tgctaaaact ttatatgaac
3481 actatagtgg tggtgaaagt cataattctt catcctccaa gacttttgaa aaaaaaggg
3541 gaaaaaaatg atttgcagtt attcgtaatg tcagatagga ggtataaaat atattggact
3601 gttatactct tgaatccctg tggaaatcta catttgaaga caacatcact ctgaagtgtt
3661 atcagcaaaa aaaattcagt gagattatct ttaaaagaaa actgtaaaaa tagcaaccac
3721 ttatggcact gtatatattg tagacttggtt ttctctgttt tatgctcttg tgtaatctac
3781 ttgacatcat ttactcttg gaatagtggg tggatagcaa gtatattcta aaaaactttg
3841 taaataaagt tttgtggcta aaatga (SEQ ID NO: 1)
```

FIG. 15B

17/19

1 mesedlsgre ltidsimnkvr dikanfkne dltdeislnk isadttndsg tvnqimman
61 npedwlslll kleknsvppls dallnkligr ysqaiealpp dkygqnesfa riqvrfaelk
121 aigepddard yfqmaranck kfafvhisfa qfelsqgnvk kskqllqkav ergavpleml
181 eialrnlnlq kkqllseeek knlsastvlt aqesfsgslg hlqnrnnsd srgqttkarf
241 lygenmppqd aeigyrsnr qtnktkqscp fgrvpvnlln spdcdvktdd svvpfcfmrq
301 tsrsecrdlv vpgskpsgnd scelrnkxsv qnshfkeplv sdeksseli tdsitlknkt
361 essllaklee tkeyqepevp esnqkqwqsk rksecinqnp aassnhwqip elarkvnteq
421 khttfeqpvf svskqsppis tskwfdpxsi cktpssntld dymscfrtpv vkndfppacq
481 lstpygqpac fqqqqhqla tplqnlqvla sssanecisv kgriysilkq igsggsskvf
541 qvlnekkqiy aikyvnllea dnqtlidsyrn eiaylnklq hsdkiirlyd yeitdqyiy
601 vmecgnidln swlkkkksid pwerksywkn mlevhtihq hgivhsdlkp anflivdgml
661 klidfgianq mqpdttsvvk dsqvgtnym ppeaikdmss srengksksk ispksdvws
721 gcilyymtyg ktpfqqiinq isklhaiidp nheiefpdip ekdlqdvkcl clkrdpkqri
781 sipellahpy vqiqtphvng makgtteemk yvlgqlvgln spnsilkaak tlyehysgge
841 shnssssktf ekkrgkk (SEQ ID NO: 2)

FIG. 16




FIG. 17A

19/19




FIG. 17B

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2008/002228

A. CLASSIFICATION OF SUBJECT MATTER

IPC: **A61K 39/395** (2006.01) , **A61K 31/7105** (2006.01) , **A61K 45/00** (2006.01) , **A61K 48/00** (2006.01) ,
A61P 35/00 (2006.01), **A61P 35/04** (2006.01) (additional International Patent Classifications found on extra sheet)
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K*(2006.01), A61P*(2006.01), C12Q*(2006.01), G01N*(2006.01)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic database(s) consulted during the international search (name of database(s) and, where practicable, search terms used)

Databases: Canadian Patent Database, Delphion, QPAT, SCOPUS, Pubmed

Keywords: tyrosine threonine kinase, TTK, MPS1, PYT, breast cancer, basal-like, non-luminal, soft tissue sarcoma, TTK antagonist, HER-2, trastuzumab.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02/068444 A1 (REINHARD et al.) 6 September 2002 (06-09-2002)	1-4, 8- 11, 16-22, 27-34
Y	*whole document*	5-7, 23-26
X	YUAN et al. Increased expression of mitotic checkpoint genes in breast cancer cells with chromosomal instability. Clinical Cancer Research. January 15, 2006. Vol. 12, No. 2, pages 405-410. ISSN: 1078-0432	29-34
	whole document	
Y	SLAMON et al. Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. The New England Journal of Medicine. March 15, 2001. Vol. 344, No. 11, pages 783-792. ISSN: 0028-4793	5-7, 23-26
	Abstract	

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier application or patent but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

11 February 2009 (11-02-2009)

Date of mailing of the international search report

23 March 2009 (23-03-2009)

Name and mailing address of the ISA/CA
Canadian Intellectual Property Office
Place du Portage I, C114 - 1st Floor, Box PCT
50 Victoria Street
Gatineau, Quebec K1A 0C9
Facsimile No.: 001-819-953-2476

Authorized officer

Scott Gurd 819- 994-4157

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2008/002228

Classification of subject matter continued:

C12Q 1/48 (2006.01) , *C12Q 1/68* (2006.01) , *G01N 33/574* (2006.01) , *G01N 33/68* (2006.01)

INTERNATIONAL SEARCH REPORTInternational application No.
PCT/CA2008/002228**Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of the first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons :

1. ☒ Claim Nos. : 1-24, 27-34

because they relate to subject matter not required to be searched by this Authority, namely :

Claims 1-24 and 27-34 are directed to methods of medical treatment and diagnostic methods practised on the human or animal body which the International Search Authority is not required to search. However, this Authority has carried out a search based on the alleged effects or purposes/uses of the product defined in claims 1-24 and 27-34.

2. ☐ Claim Nos. :

because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically :

3. ☐ Claim Nos. :

because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows :

See extra sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claim Nos. :
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim Nos. :

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.

☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.

☐ No protest accompanied the payment of additional search fees.

Continuation of Box III:

Group 1 : Claims 1-11, 16-26, 29, 30, 33, 34 (completely) and claims 27, 28, 31, 32 (partially) are directed methods and compositions for treating tyrosine threonine kinase (TTK) positive non-luminal breast cancer and methods for the diagnosis and prognosis of non-luminal breast cancer comprising the assessment of TTK expression levels in cancer cells or tissues.

Group 2: Claims 12-15 (completely) and 27, 28, 31, 32 (partially) are directed to methods for treating TTK positive soft tissue sarcomas and methods for the diagnosis and prognosis of cancer comprising the assessment of TTK expression levels in cancer cells or tissues.

The above groups of inventions all involve TTK in the diagnosis of cancer and as a therapeutic target in cancer. However, diagnostic, prognostic, and therapeutic methods which take advantage of the differential expression of TTK in mammalian cancers have been disclosed in PCT application WO 02/068444 (D1). Since the use of TTK for the diagnosis, prognosis, and treatment of cancer is known in the art, the methods involving non-luminal breast cancer and soft tissue sarcomas do not possess unity *a posteriori* and are thus treated as separate inventions.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/CA2008/002228

Patent Document Cited in Search Report	Publication Date	Patent Family Member(s)	Publication Date
WO 02/068444 A1	06-09-2002	US20050063974 A1	24-03-2005
		US20050059630 A1	17-03-2005
		US20050058627 A1	17-03-2005
		NZ0527421 A	27-05-2005
		JP2004526716 T2	02-09-2004
		EP1377596 A1	07-01-2004
		EP1377596 A4	13-07-2005
		CN1492875 A	28-04-2004
		CA2438092 AA	06-09-2002
		AU2244112 BB	10-05-2007