



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/00		A2	(11) International Publication Number: WO 99/34785 (43) International Publication Date: 15 July 1999 (15.07.99)
(21) International Application Number: PCT/IL99/00003			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 5 January 1999 (05.01.99)			
(30) Priority Data: 122883 9 January 1998 (09.01.98) IL 127102 17 November 1998 (17.11.98) IL			
(71) Applicant (for all designated States except US): MOR-RESEARCH APPLICATIONS LTD. [IL/IL]; Hasivim Street 23, P.O. Box 7590, Kiryat Matalon, 49170 Petach Tikva (IL).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): MELAMED, Eldad [IL/IL]; Tagore Street 44, Ramat Aviv, 69341 Tel Aviv (IL). DJALDETTI, Ruth [IL/IL]; Nathan Street 7, 52450 Ramat Gan (IL). ZIV, Ilan [IL/IL]; Shezaf Street 5, 44418 Kfar-Sava (IL).			
(74) Agent: REINHOLD COHN AND PARTNERS; P.O Box 4060, 61040 Tel Aviv (IL).			

(54) Title: TREATMENT OF DYSKINESIAS

(57) Abstract

Levodopa-induced dyskinesia and tardative dyskinesia are ameliorated by administration of riluzole to needing subjects.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TREATMENT OF DYSKINESIAS

FIELD OF THE INVENTION

The present invention concerns pharmaceutical compositions for the treatment of dyskinesias, particularly levodopa-induced dyskinesia and tardative dyskinesia.

5

BACKGROUND OF THE INVENTION

Parkinson's disease is an age related, progressive neurodegenerative disorder. The prevalence rate is approximately 0.5% in the population aged 50-59, 1% in ages 60-69, 2% in the 70-79 age group and rises to over 3% in 10 those who are 80 and older. Prevalence rates are similar in Europe.

Parkinson's disease is characterized by a relatively selective degeneration of dopaminergic neurons in the *substantia nigra pars compacta* with loss of striatal dopamine. The pathology shows depigmentation of the *substantia nigra* and intracellular inclusions (Lewy bodies). The cardinal 15 features of the disease include resting tremor, rigidity, bradykinesia and postural instability. Current treatment of the motor signs of Parkinson's disease is based on dopamine replacement. This involves the administration of levodopa, usually combined with a decarboxylase inhibitor. Exogenous levodopa is converted in the striatum to dopamine and replenishes the reduced 20 dopaminergic concentrations in the basal ganglia. Dopamine agonists may be helpful as well. During the first years of treatment the patients enjoy a smooth and stable response to this treatment. However, after 2-5 years of chronic treatment with dopaminergic preparations, 75% of patients develop disabling

and incapacitating motor complications. One of the most common side effects is the levodopa-induced dyskinesias (choreiform involuntary movements). They occur in the majority (80-100%) of the patients as their illness progresses. Dyskinesias may be initially mild but they can become more and more progressive, complex, generalized, violent, and may severely interfere with motor function, speech, coordination and postural stability. Patients and families are often shameful by the unaesthetic and bizarre movements which can affect facial muscles, eyelids, mouth, cheeks, lips and tongue, upper and lower limbs and even trunk and respiratory muscles. This is one of the major 5 reasons for social decline of afflicted patients.

10

At the beginning, when they first emerge, dyskinesias are mainly the peak-dose type, i.e., they are most prominent when levodopa plasma levels are high. When patients later develop response fluctuations after chronic levodopa treatment, dyskinesias may also appear at the beginning and again at the 15 termination of an individual levodopa dose beneficial effect. When disease is more advanced, dyskinesias predominate in an "all or none" fashion, i.e., they are present throughout the duration of an "on" period, induced by a successful single oral dose of levodopa. Such levodopa-induced dyskinesias also represent a major limiting factor in the pharmacological treatment of Parkinson's disease.

20 When their illness progresses, the patients need increases in their daily levodopa dosage and the addition of other dopaminergic agents, e.g., dopamine agonists and MAO-B inhibitors. This is invariably associated with a rapid and intolerable increase of the frequency, distribution and severity of dyskinesias necessitating reduction of drugs. Dykinesias are probably and primarily caused 25 by the action of excessive exogenous dopamine on denervation-supersensitive post-synaptic dopaminergic receptors. Normally, the dopamine formed from levodopa is stored in vesicles within the dopaminergic nerve-endings for regulated release into the synapse. As the disease progresses, more nigral

dopaminergic neurons degenerate and there is more severe loss of their nerve-terminals in the basal ganglia (caudate and putamen nuclei).

It is believed that the decarboxylation of exogenous levodopa therefore shifts to non-dopaminergic striatal compartments. Since the generated 5 dopamine molecules are not stored, they immediately interact and hyperactivate the postsynaptic dopaminergic receptors (mainly of the D2 subtype) resulting in the involuntary movements. There is no satisfactory treatment for dyskinesias. Discontinuation or reduction of levodopa and other dopaminergic drugs or addition of neuroleptic drugs that block dopaminergic 10 receptors, can abolish the abnormal movements, but at the expense of severe aggravation of the Parkinsonian symptoms. The control-release levodopa preparations have been proven unhelpful. This despairing state of affairs suggested that it would be difficult if not impossible to dissociate the beneficial anti-Parkinsonian from the bad dyskinetic producing effects of levodopa.

15 Behavioral and psychiatric disorders are usually treated by the administration of by various anti-psychotics, also termed: "*neuroleptic drugs*" the majority act by blockage of dopamine D₂ receptor. Prolonged administration of anti-psychotic drugs often results in development of involuntary movements, termed: "*tardative dyskinesia*" .

20 Riluzole (2-amino-6-trifluoromethoxy benzothiazole) has recently emerged as a pharmacological agent potentially useful to slow down the evolution of neurodegenerative diseases, such as amyotrophic lateral sclerosis. (Ben Simon *et al.*, *New Engl. J. Med.*, **330**:585-91 (1994)). In addition, this molecule has been shown to display anticonvulsant, anti-ischemic, and 25 neuroprotective properties under various experimental conditions. A clear understanding of the site and mechanism of action of this molecule is still lacking.

There has been recently a report (Palti *et al.* *Exper. Neuro* 1.146 13J-141 (1997)) that riluzole reduces various abnormal motor movement in baboons

which were induced by 3-nitroproponic acid, serving as a model for progressive striatal degeneration, which is a model for Huntington's Disease.

SUMMARY OF THE INVENTION

5 The present invention provides, by one of its aspects, a pharmaceutical composition for the amelioration of levodopa-induced dyskinesia and tardative dyskinesia, comprising as an active ingredient, a pharmaceutically effective amount of riluzole.

10 The present invention provides, by another of its aspects, use of riluzole for the preparation of a pharmaceutical composition for the amelioration of levodopa-induced dyskinesia and tardative dyskinesia.

15 The term "*amelioration*" refers to a decrease in the abnormal involuntary movements characterizing these two types of dyskinesia, as can be determined for example, by using the Abnormal Involuntary Movement Scale (AIMS) as will be specified hereinbelow.

The term "*levodopa-induced dyskinesia*" refers to dyskinesia, i.e. involuntary choreiform movements, brought about by the chronic administration of levodopa, for example in patients suffering from Parkinson's Disease.

20 The term "*tardative dyskinesia*" refers to dyskinesia brought about by the chronic administration of neuroleptic, anti-psychotic drugs of the Dopaminergic-receptor blocker type.

The term "*riluzole*" refers to 2-amino-6 trifluoromethoxy-benzothiazole.

25 The term "*effective amount*" refers to an amount that brings about a reduction in the AIMS of the patients without causing severe side effects.

The dosage of the active ingredient should be tested empirically for each specific indication, and depends on various factors, such as the patient's weight, the length of time of administration of the levodopa or the neuroleptic pharmaceutical composition, age, etc. Generally speaking, the dosage should be

of about 25 to about 200 mg per day, preferably of about 50 to about 200 mg per day, most preferably of about 50 to about 100 mg per day.

The pharmaceutical composition of the invention, may comprise solely riluzole and a pharmaceutically acceptable carrier.

5 Alternatively, it is possible to include in one dosage form, both the dyskinesia causing agent such as the neuroleptic drug (in the case of tardative dyskinesia), or levodopa (in the case of levodopa-induced dyskinesia) together with the riluzole.

10 The present invention further concerns a method for ameliorating levodopa-induced dyskinesia or tardative dyskinesia by administering to a subject in need of such treatment, a therapeutically effective amount of riluzole.

15 According to the method of the present invention, the riluzole may be administrated separately, i.e. not simultaneously with the dyskinesia-causing agent (such as the neuroleptic drug or the levodopa), or alternatively may be administered together with the dyskinesia-causing agents either by administration of the two medicaments simultaneously or by forming both medicaments in a single dosage form.

20 The invention will now be described with reference to some non-limiting examples.

DETAILED DESCRIPTION OF THE INVENTION

I. Clinical Procedures

25 Twelve patients suffering from advanced Parkinson's Disease featuring dyskinesia, and twelve patients having appropriate criteria for tardative dyskinesia are participating in an open experiment for assessing the influence of riluzole on involuntary movement.

The Parkinson patients are balanced by optimal dopaminergic treatment in the three months prior to the clinical trial. The patients with tardative dyskinesia, which are already balanced by neuroleptic treatment, do not reduce

the dosage of the neuroleptic drug, and do not cease other treatments, which they receive.

The clinical assessment of the Parkinson patient is carried out by using the Unified Parkinson's Disease Rating Scale (UPDRS) and the assessment of 5 involuntary movement will be carried out by the Abnormal Involuntary Movement Scale (AIMS).

Assessment of patients with tardative dyskinesia is carried out utilizing AIMS. The trial is carried out for six weeks. Prior to the beginning of the trial, patients undergo blood and urine tests, a chest X-ray, an ECG, as well as 10 general physical and neurological evaluations. During the clinical trial, the patients are treated with riluzole having an initial dosage of 25 mg. twice a day, and after a week, the dosage will be increased to 50 mg. twice a day. Patients are monitored once every two weeks in order to carry out a blood count and SMA (biochemical blood tests). In addition, a clinical 15 assessment-involuntary-movement is carried out separately by two independent physicians. During the clinical trial, patients are asked to fill a detailed diary which will grade the severity of the dyskinesia and will assess daily function according to the following scale:

20 **THE INTENSITY OF THE DYSKINESIA:**

- 0-without dyskinesia
- 1-mild dyskinesia
- 2-medium dyskinesia,
- 3-severe dyskinesia

25

RATING OF DAILY FUNCTION:

- 1 - An improvement of daily function as compared to the basic condition
- 2 - No improvement in daily function
- 3 - Deterioration of daily function as compared to the basic condition

At the end of the clinical trial, patients undergo blood tests, ECG 's, as well as neurological and psychological assessment.

In addition, patients are invited for routine check-ups two weeks from the end of the trial.

5

II. Clinical trials

Six patients suffering from advanced parkinson's disease with severe levodopa-induced dyskinesias participated in an open-label pilot study to assess safety, tolerability and efficacy of riluzole in attenuating the involuntary movements.

10

The patients were given optimal anti-Parkinsonian drug treatment during the three months prior to the clinical trial. Duration of study was six weeks. First two weeks served to accumulate baseline data. Patients filled up dyskinesias diaries in which they marked at every waking hour whether involuntary movements were present and their severity (mild/moderate and severe). Each patient was administered with half a tablet of riluzole (25 mg) once, in the morning, for four days. The dose was then increased to 25 mg, b.i.d. for additional four days (once in the morning and once in the early afternoon). The dose was then further increased to two 50 mg tablets (total of 20 100 mg. daily) which the patients took for three additional weeks. They continued to fill up their dyskinesias diaries throughout the trial period.

Results

Treatment with riluzole was found to be effective in attenuating the dyskinesias. Mean daily waking hours spent with dyskinesias decreased by about 24% from 6.92 ± 3.67 hours before treatment to 5.26 ± 4.23 hours during treatment ($P < 0.01$; paired t-test). Mean daily waking hours spent in severe dyskinesias reduced by about 30% from 2.76 ± 1.77 hours before treatment to 1.94 ± 2.40 hours during treatment ($0.01 < p < 0.05$; paired t-test).

25

There was no worsening of the Parkinsonian signs and symptoms when patients took riluzole. Likewise, there was no decrease in the efficacy of levodopa and other anti-Parkinsonian drugs and in the total daily time patients spent in "on" periods.

5 Riluzole was well-tolerated and there was no report of adverse-effects. This preliminary open-label study indicates that administration of riluzole (50 mg. b.i.d) can attenuate levodopa-induced dyskinesias in patients with Parkinson's disease without causing deterioration of the Parkinsonian signs and without suppression of levodopa efficacy.

CLAIMS:

1. A pharmaceutical composition for the amelioration of levodopa-induced dyskinesia and tardative dyskinesia, comprising as an active ingredient, a pharmaceutically effective amount of riluzole.
2. A pharmaceutical composition according to Claim 1 in a dosage form for the administration of about 25 to about 200 mg of riluzole per day.
3. A pharmaceutical composition according to Claim 2 in a dosage form for the administration of about 50 to about 100 mg of riluzole per day.
4. A method for ameliorating levodopa-induced dyskinesia or tardative dyskinesia comprising administering to a subject in need of such treatment, a therapeutically effective amount of riluzole.
5. A method according to Claim 4, wherein the riluzole is together with levodopa or with an anti-psychotic drug.
6. A method according to Claim 4, wherein the riluzole is administered in an amount of about 25 to about 200 mg per day.
7. A method according to Claim 6, wherein the riluzole is administered in an amount of about 50 to about 100 mg per day.
8. Use of riluzole for the preparation of a pharmaceutical composition, for the amelioration of levodopa-induced dyskinesia and tardative dyskinesia.
9. Use according to Claim 8, wherein the prepared pharmaceutical composition is in a dosage form for administration of 25 to 200 mg of riluzole per day.
10. Use according to Claim 9, wherein the prepared pharmaceutical composition is in a dosage form for administration of 50 to 200 mg of riluzole per day.
11. Use according to Claim 10, wherein the prepared pharmaceutical composition is in a dosage form for administration of 50 to 100 mg of riluzole per day.