



(86) Date de dépôt PCT/PCT Filing Date: 2009/09/25  
 (87) Date publication PCT/PCT Publication Date: 2010/04/01  
 (45) Date de délivrance/Issue Date: 2012/09/04  
 (85) Entrée phase nationale/National Entry: 2011/03/11  
 (86) N° demande PCT/PCT Application No.: US 2009/058463  
 (87) N° publication PCT/PCT Publication No.: 2010/036947  
 (30) Priorité/Priority: 2008/09/27 (US61/100,714)

(51) Cl.Int./Int.Cl. *A61K 47/44* (2006.01),  
*A61K 31/506* (2006.01), *A61K 31/58* (2006.01),  
*A61P 17/14* (2006.01)

(72) Inventeurs/Inventors:  
 ALI, SHOUKATH M., US;  
 AHAMD, ATEEQ, US;  
 AHMAD, MOGHIS U., US;  
 SHEIKH, SAIFUDDIN, US;  
 AHMAD, IMRAN, US

(73) Propriétaire/Owner:  
 JINA PHARMACEUTICALS, INC., US

(74) Agent: SMART & BIGGAR

(54) Titre : PREPARATIONS PHARMACEUTIQUES A BASE DE LIPIDE(S) A USAGE ORAL ET TOPIQUE, LEURS  
 COMPOSITIONS, PROCEDES ET UTILISATIONS  
 (54) Title: LIPID BASED PHARMACEUTICAL PREPARATIONS FOR ORAL AND TOPICAL APPLICATION; THEIR  
 COMPOSITIONS, METHODS, AND USES THEREOF

(57) **Abrégé/Abstract:**

The present invention relates to the topical and oral delivery of a composition comprising one or more active agents for treating a disease or symptoms in a subject. In some embodiments, the present invention comprises a composition comprising at least one active compound, e.g., finasteride or minoxidil, and one or more lipids. In some embodiments, the present invention relates to composition and method of preparation for treating androgenic alopecia (AGA), prevention of hair loss and female hirsutism. In some embodiment, the present invention comprises finasteride and at least one lipid component for the treatment of benign prostatic hyperplasia. In some embodiment, the present invention comprises tacrolimus or amphotericin B and at least one lipid component for the treatment of skin or eye related diseases. The present invention provides a method of preparation of a composition comprising at least one active compound and at least one lipid and administering the composition to a subject by oral or topical delivery. In certain embodiments the subject is a mammal. In certain preferred embodiment, the subject is human.



## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
1 April 2010 (01.04.2010)(10) International Publication Number  
**WO 2010/036947 A3**

## (51) International Patent Classification:

A61K 47/44 (2006.01) A61K 31/58 (2006.01)  
A61P 17/14 (2006.01) A61K 31/506 (2006.01)

## (21) International Application Number:

PCT/US2009/058463

## (22) International Filing Date:

25 September 2009 (25.09.2009)

## (25) Filing Language:

English

## (26) Publication Language:

English

## (30) Priority Data:

61/100,714 27 September 2008 (27.09.2008) US

(71) Applicant (for all designated States except US): **JINA PHARMACEUTICALS, INC.** [US/US]; 28100 N. Ashley Circle, Suite 103, Libertyville, Illinois 60048 (US).

## (72) Inventors; and

(75) Inventors/Applicants (for US only): **ALI, Shoukath M.** [US/US]; 29681 North Waukegan Road, #204, Waukegan, Illinois 60048 (US). **AHAMD, Ateeq** [US/US]; 4795 West Pebble Beach Drive, Wadsworth, Illinois 60083 (US). **AHMAD, Moghis U.** [US/US]; 3050 North Forrest Hills Court, Wadsworth, Illinois 60083 (US). **SHEIKH, Saifuddin** [US/US]; 1059 South West Avenue, Waukegan, Illinois 60085 (US). **AHMAD, Imran** [US/US]; 4731 West Pebble Beach Drive, Wadsworth, Illinois 60083 (US).(74) Agent: **BROW, Mary Ann D.**; Casimir Jones, S.C., 440 Science Drive, Suite 203, Madison, Wisconsin 53711 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

## Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

## (88) Date of publication of the international search report:

10 June 2010

(54) Title: LIPID BASED PHARMACEUTICAL PREPARATIONS FOR ORAL AND TOPICAL APPLICATION; THEIR COMPOSITIONS, METHODS, AND USES THEREOF

(57) Abstract: The present invention relates to the topical and oral delivery of a composition comprising one or more active agents for treating a disease or symptoms in a subject. In some embodiments, the present invention comprises a composition comprising at least one active compound, e.g., finasteride or minoxidil, and one or more lipids. In some embodiments, the present invention relates to composition and method of preparation for treating androgenic alopecia (AGA), prevention of hair loss and female hirsutism. In some embodiment, the present invention comprises finasteride and at least one lipid component for the treatment of benign prostatic hyperplasia. In some embodiment, the present invention comprises tacrolimus or amphotericin B and at least one lipid component for the treatment of skin or eye related diseases. The present invention provides a method of preparation of a composition comprising at least one active compound and at least one lipid and administering the composition to a subject by oral or topical delivery. In certain embodiments the subject is a mammal. In certain preferred embodiment, the subject is human.



WO 2010/036947 A3



**LIPID BASED PHARMACEUTICAL PREPARATIONS FOR ORAL AND  
TOPICAL APPLICATION; THEIR COMPOSITIONS, METHODS, AND  
USES THEREOF**

**FIELD OF THE INVENTION**

5

The invention relates to compositions comprising active components or compounds, *e.g.*, pharmaceutical compounds, and lipids, including, *e.g.*, solutions, complexes, micelles, emulsions, liposomes or lipidic particle, and mixture of micelles and vesicles. The invention further relates to their methods of preparation, and uses in the treatment of diseases. The invention also relates to a method for the topical delivery of active agents, and compositions used therein. In some embodiments, the invention relates to compositions comprising, hair loss preventing or hair growth drugs such as finasteride and minoxidil. In some embodiments, the invention relates to composition comprising active compounds for use in treating skin or eye infections, or diseases related to skin or eyes, *e.g.*, in mammals. In some embodiments, the invention also related to a method for the oral delivery of active agents. Methods according to the present invention are suitable for practice on an industrial manufacturing scale, and may be practiced, *e.g.*, as a continuous process.

10  
15  
20 

**BACKGROUND OF THE INVENTION**

[0001] Androgenic alopecia (AGA) is the most common form of hair loss, affecting approximately 30 to 40 percent of adult men and women. The incidence is generally considered to be greater in males than females. In men, the condition is also called male-pattern baldness. Commonly treatments for androgenic alopecia include hair follicle transplants, topical therapies, and orally prescribed anti-androgens (J.L. Roberts, 1997). The hyperandrogenic stimulation that causes alopecia also produces other undesirable physiological symptoms including acne vulgaris, benign prostatic hyperplasia, female hirsutism, and seborrhea.

25  
30 [0002] Early attempts to provide a chemotherapeutic agent to counter the undesirable results of hyperandrogenicity resulted in the discovery of several steroidal antiandrogens having undesirable hormonal activities of their own. The

principal mediator of androgenic activity in some target organs, e.g. the prostate, is 5 $\alpha$ -dihydrotestosterone (DHT), formed locally in the target organ by the action of testosterone-5 $\alpha$ -reductase. Inhibitors of testosterone-5 $\alpha$ -reductase will serve to prevent or lessen symptoms of hyperandrogenic stimulation in these organs (Gormley *et al.* US Pat. No. 5,981,543, 1999 and Rasmusson *et al.* US Pat. No. 4,377,584, 1983).

10 [0003] Minoxidil (6-(1-piperidinyl)-2,4-pyrimidinediamide 3 oxide) is a drug known for the treatment of AGA. Minoxidil is effective when delivered topically at a concentration of about 0.01% to about 5%. The topical solution of Minoxidil is currently marketed as "Rogaine" having 2% minoxidil concentration in a solution of (60% v/v) propylene glycol, and water. Disadvantageously, clinical trials have shown that the topical application of a 2% minoxidil solution to patients experiencing hair loss results stimulated dense hair regrowth in only less than about 5% of the patients and moderate hair regrowth in only about 30% of the patients (E.A. Olsen, *et al.*, 1985; J. Roberts, 1987), S. Niemiec *et al.* US Patent No. 6,419,913 B1, 2002).

20

[0004] Finasteride is a synthetic androgen inhibitor currently marketed as PROSCAR<sup>®</sup> for the treatment of benign prostatic hyperplasia. It has also been marketed orally in low dosage form as PROPECIA<sup>®</sup> for the treatment of androgenic alopecia (AGA) (hair loss). Chemically, finasteride is (5 $\alpha$ , 17 $\beta$ )-N-(1, 1-dimethylethyl)-3-oxo-4-azaandrost-1-ene-17-carboxamide and is practically insoluble in water, and is soluble only in organic solvents such as ethanol and methanol. Finasteride is known to inhibit 5 $\alpha$ -reductase type 2.

25 [0005] Although, finasteride is currently administered orally, it is not available topically as approved product for the treatment of AGA. The side effects as a result of the oral administration for finasteride include erectile dysfunction, impotence, low libido, gynecomestica, and facial hair growth.

30



5 [0006] Development of new lipid - based formulation is needed to improve oral and or transdermal delivery and reduce the toxicity associated with compositions such as finasteride and minoxidil.

## SUMMARY OF THE INVENTION

10 [0007] The present invention relates to a new methods of preparing lipid formulations of active compounds, and methods of using the formulation in treating a disease or symptom in a subject. In some embodiments, the present invention comprises a composition comprising at least one active agent, such as finasteride or minoxidil. In some embodiments, the present invention comprises a method  
15 comprising preparing a composition comprising at least one active compound, *e.g.*, finasteride, and one or more lipids and administering the composition to a subject. In certain embodiments the subject is a mammal. In certain preferred embodiments, the subject is human.

20 [0008] Another objective of the present invention is to provide lipid formulations comprising at least one active component and at least one lipid, *e.g.*, a phospholipid, or neutral lipid formed in a suitable vehicle. The suitable vehicle in the present invention comprises a non-toxic solvent or an aqueous medium.

25 [0009] The amount of phospholipid included in a composition according to the present invention is not limited to any particular amount or percentage (*e.g.*, by weight) of the final composition. In some embodiments, the proportion of at least one phospholipid is between about 0.5% to about 98% of total lipids (*e.g.*, a commercially usable form) by weight. In some preferred embodiments, the amount  
30 of the at least one phospholipid is in between 0.5% to 90% of the lipid complex by weight.

[0010] A lipid formulation of the present invention is not limited to any particular use or application. For example, a lipid formulation of an active  
35 component according to the present invention comprising a pharmaceutically active ingredient can be used for different pharmaceutical applications. An aqueous

5 system or a non-toxic solvent system of the present invention can also be used in the formation of unloaded lipid complexes (*e.g.*, without any encapsulated active ingredient) for use, *e.g.*, as controls for complexes comprising active components.

[0011] As for example, it is possible to encapsulate or entrap finasteride, in the inventive liposome system. Such a pharmaceutical product is particularly suitable for oral or topical application. Furthermore, the known active ingredients are for the treatment of benign prostatic hyperplasia, androgenic alopecia (male and female baldness), and prevention of hair loss, female hirsutism, acne vulgaris and seborrhea.

[0012] In some embodiments, the one or more lipids of a composition according to the present invention comprise one or more cholesterol, cholesteryl sulfate and its salts (*e.g.*, sodium salt), cholesteryl hemisuccinate, cholesteryl succinate, cholesteryl oleate, polyethylene glycol derivatives of cholesterol (cholesterol-PEG), coprostanol, cholestanol, cholestane, cholic acid, cortisol, corticosterone, hydrocortisone, or calciferol, while in some embodiments, the one or more lipids comprises a sterol. In certain embodiments, the sterol is  $\beta$ -sitosterol, stigmasterol, stigmastanol, lanosterol,  $\alpha$ -spinasterol, lathosterol, campesterol or a mixture thereof. In some embodiments, the sterol is guggulsterol, derivatives of guggulsterol (*e.g.*, guggulsulfate and its salt such as sodium salt, guggulmyristate, guggullaureate, gugguloleate, or short chain fatty acid derivatives of guggulsterol or mixture thereof.

[0013] In some embodiments, the one or more lipids of a composition according to the present invention comprises one or more of fatty acids having a chain length of about C<sub>4</sub>-C<sub>34</sub>. In some embodiments, one or more fatty acid chains are unsaturated, while in some embodiments, one or more of the fatty acid chains are saturated. In some embodiments, one or more of the fatty acids are in salt form, while in some embodiments; one or more of the fatty acids are in acidic form. In some embodiments, one or more fatty acids are in the form of an ester.

[0014] In some embodiments, the one or more lipids of a composition according to the present invention comprises one or more of long alkyl chain alcohols having a



chain length of about C<sub>4</sub>-C<sub>34</sub>. In some embodiments, one or more alkyl chains are unsaturated, while in some embodiments, one or more of the alkyl chains are saturated.

[0015] In some embodiments, one or more lipids of a composition according to the present invention comprise a phospholipid. In some preferred embodiments, one or more of the lipids of the composition comprises a phosphatidylcholine or phosphatidylglycerol, while in some preferred embodiments; one or more of the lipids of the composition comprises a phosphatidylethanolamine, phosphatidylserine, phosphatidylinositol, or phosphatidic acid. In some preferred embodiments, one or more lipids of the present invention comprise a soybean phospholipid. In some particularly preferred embodiments, a soybean phospholipid used in the methods and compositions of the present invention comprises a large concentration of phosphatidylcholine. In still more particularly preferred embodiments, a soybean phospholipid used in the methods and compositions of the present invention contains at least 90% by weight phosphatidylcholine. In some embodiments, one or more phospholipids are pegylated (PEG) derivatives of phospholipids. In certain embodiments, one or more of the lipids of the composition comprise a pegylated derivative of a distearoylphosphatidylglycerol, a dimyristoylphosphatidylglycerol, or a dioleoylphosphatidylglycerol phospholipid.

[0016] In some embodiments, one or more lipids of a composition according to the present invention comprise a monoglyceride, a diglyceride, or a triglyceride lipid.

[0017] The method of composition, wherein said fatty acids of mono-, di-, and triglycerides are selected from a group of saturated and unsaturated fatty acids having short chain or long chain.

[0018] In some embodiments, a composition according to the present invention further comprises polyethylene glycol (PEG). In some embodiments, the PEG has an average molecular weight ranging from 200-20,000, while in certain preferred embodiments, the average molecular weight of the PEG is in between 500-2000.

[0019] In some embodiments, a composition according to the present invention comprises active compound (for example finasteride), cholesterol or cholesterol derivatives and one or more phospholipids. In some embodiments in which the composition comprises a cholesterol derivative, the cholesterol derivative is cholesteryl sulfate. In some embodiments wherein the phospholipid comprises soy phosphatidylcholine or hydrogenated phosphatidylcholine. In some preferred embodiments, the mole ratio of active compound (for example, finasteride) and cholesterol or cholesterol derivative is in the range of about 1:0.5 and 1:50, while in certain particularly preferred embodiments, the mole ratio of active compound (for example, finasteride) and cholesterol or cholesterol derivative is in between about 1:1 and 1:10.

[0020] In some embodiments, one or more lipids of a composition according to the present invention comprise hydrogenated soy phosphatidylcholine, wherein the mole ratio of active compound (for example, finasteride) and hydrogenated soy phosphatidylcholine is in between about 1:1 and 1:80. In certain preferred embodiments, the mole ratio of active compound (for example, finasteride) and hydrogenated soy phosphatidylcholine is in between about 1:1 and 1:60.

[0021] In some embodiments, one or more lipids of a composition according to the present invention comprise soy phosphatidylcholine, wherein the mole ratio of active compound (for example, finasteride) and soy phosphatidylcholine is in between about 1:1 and 1:80. In certain preferred embodiments, the mole ratio of active compound (for example, finasteride) and soy phosphatidylcholine is in between about 1:1 and 1:60.

[0022] In some embodiments, a composition according to the present invention comprises active compound (for example, finasteride,) at a concentration of from about 0.5 mg/mL to about 200 mg/mL while in some preferred embodiments, the active compound (for example, finasteride) of the composition is at a concentration of from about 1 mg/mL to about 10 mg/mL. In some particularly preferred embodiments, the composition of the invention comprises active compound (for example, finasteride) is at a concentration of about 1 mg/mL to about 5mg/mL.



5 [0023] In some embodiments, a composition according to the present invention comprises active compound (for example, finasteride,) at a concentration of from about 0.5mg/g to about 200mg/g while in some preferred embodiments, the active compound (for example, finasteride) of the composition is at a concentration of from about 1 mg/g to about 50 mg/g. In some particularly preferred embodiments, the composition of the invention comprises active compound (for example, finasteride) is at a concentration of about 1 mg/g to about 10 mg/g.

10

[0024] In some embodiments, a composition according to the present invention comprises a total lipid concentration or proportion of from about 0.5% by weight to about 95% by weight, while in some preferred embodiments; the composition comprises a total lipid concentration of from about 0.5% by weight to about 80% by weight. In certain particularly preferred embodiments, the composition comprises a total lipid concentration of from about 10% by weight to about 90% by weight.

15

[0025] In some embodiments, a composition according to the present invention comprises active compound (for example, finasteride) and total lipid(s) having a weight-to-weight ratio ranging from about 1:1 to about 1:100, while in certain preferred embodiments, the ratio is in between 1:1 to about 1:60.

20

[0026] In some embodiments, a composition according to the present invention comprises a composition selected from the group consisting of a micelle and an emulsion. In certain preferred embodiments, the composition comprises a plurality of micelles, wherein said micelles are in the form of monomeric, dimeric, polymeric or mixed micelles.

25

[0027] The present invention is not limited to any particular form of composition of the invention. For example, in some embodiments, a composition according to the present invention is in a lyophilized form. In some, embodiments, the composition further comprise a cryoprotectant. In certain, preferred embodiments, the cryoprotectant comprise one or more sugars, while in particularly preferred embodiments; the one or more sugars comprise trehalose, maltose, lactose, sucrose, glucose, and/or dextran.

35

5 [0028] In some embodiments, a composition according to the present invention comprises complexes, liposomes, micelles, vesicles that have a diameter of about 20 microns or less, while in some embodiments, the complexes, liposomes, micelles, vesicles that have a diameter of about 10 microns or less. In some embodiments, the complexes, liposomes, micelles, vesicles that have a diameter of about 5 microns or less, while in some embodiments, the complexes, liposomes, micelles, vesicles that have a diameter of about 1 micron or less. In some embodiments, the complexes, liposomes, micelles, vesicles that have a diameter of about 500 nm or less, while in some embodiments, the complexes, liposomes, micelles, vesicles that have a diameter of about 200 nm or less. In some preferred embodiments, the complexes, liposomes, micelles, vesicles that have a diameter of about 100 nm or less.

15

[0029] In some embodiments, a composition according to the present invention is in a powder form, while in some embodiments, the complex is in a solution form. In some embodiments, the composition is in a suspension form, while in other embodiments, the composition is in an emulsion form, while in still other 20 embodiments, the composition is in a micelle form or mixed micellar form or in a liposome form. In some embodiments, the composition is in a lyophilized or gel form, while in some embodiments, the composition is in a paste form. In some embodiments, the composition is a mixture of mixed micelles, liposomes or vesicles form.

25

[0030] The composition produced according to the present inventive method can be filled directly in corresponding ampoules, vials, bottles, tubes in a condition ready to use. The product can be lyophilized, if desired to provide the composition in powder form, which can be mixed in a suitable vehicle for topical application. 30 Examples of suitable vehicles include but not limited to oil, gel, paste, lotion, shampoo, soap, and the like. In some embodiments, the lyophilized powder can be used for oral delivery of therapeutic agent (for example, finasteride).

35 [0031] It is contemplated that the methods and compositions of the present invention are used to treat androgenic alopecia such as male and female baldness,



prevention of hair loss, female hirsutism, acne vulgaris, and seborrhea. In some  
embodiments, the present invention comprises use of the compositions described  
5 above in the preparation of a medicament for treatment of a subject for, *e.g.*,  
androgenic alopecia such as male and female baldness, prevention of hair loss,  
female hirsutism, acne vulgaris, and seborrhea.

[0032] It is contemplated that in some embodiments, the exposing of a cell in a  
10 subject comprises topical delivery of the composition to the subject. In some  
preferred embodiments, the subject is a mammal. In some particularly preferred  
embodiments, the mammal is human.

## DEFINITIONS

15 [0033] The term "lipid composition" as used herein refers to amphoteric  
compounds which are capable of liposome formation, vesicle formation, micelle  
formation, emulsion formation, and are substantially non-toxic when administered  
at the necessary concentrations as liposomes. The lipid composition may include  
without limitation egg phosphatidylcholine (EPC), egg phosphatidylglycerol (EPG),  
20 soy phosphatidylcholine (SPC), hydrogenated soy phosphatidylcholine (HSPC),  
dimyristoylphosphatidylcholine (DMPC), dimyristoylphosphatidylglycerol  
(DMPG), Dipalmitoylphosphatidylcholine (DPPC), distearylphosphatidylglycerol  
(DSPG), dipalmitoylphosphatidylglycerol (DMPG), cholesterol (Chol), cholesterol  
sulfate and its salts (CS), cholesterol hemisuccinate and its salts (Chems),  
25 cholesterol phosphate and its salts (CP), cholesterylphospholine and other  
hydroxycholesterol or amino cholesterol derivatives, guggulsterol or guggulsterol  
derivatives, *e.g.*, guggulsulfate and its salts, guggullaurate, guggulmyristate or other  
saturated or unsaturated fatty acid derivatives of guggulsterol.

30 [0034] As used herein, the term "aqueous" as used in reference to a solvent,  
fluid, or system, refers to a water-based solvent, fluid or system that does not  
contain any organic solvents. The aqueous system may further contain buffer(s).  
Examples of base or buffer includes but not limited to sodium succinate dibasic,  
sodium acetate, sodium phosphate monobasic, sodium phosphate dibasic, sodium  
35 phosphate tribasic, sodium citrate, sodium hydroxide, and the like.

5 [0035] As used herein, the term "non-toxic solvent" refers to alcohols, and oils. Examples of alcohols include but not limited to ethanol, 1-propanol, isopropyl alcohol, propylene glycol, butanol, and t-butanol. Suitable examples of oils include but not limited to olive oil, soybean oil, coconut oil, almond oil, sunflower oil, sun flower oil, vegetable oil, canola oil, coconut oil, sesame seed oil, avocado oil, cod oil and the like.

10 [0036] The term "encapsulating amount" or "entrapping amount" refers to the amount of lipid necessary to encapsulate or entrap the poorly soluble compound and form liposome or lipidic particles of appropriate mean particle size less than 5,000nm in diameter, preferably between 30-1000nm. The encapsulating amount will depend on the pharmaceutically active compounds and process conditions  
15 selected, but in general range in between from 2:1 to about 1:100 compound: lipid ratio; preferably about 1:1 to about 1:50.

[0037] The term "lipidic particle" as used herein refers to particles of undefined structure which consist of a suitable lipid and an encapsulated or complexed  
20 pharmaceutically active compound. Lipidic particles may have a lamellar structure but are not required to exhibit any defined structure.

[0038] As used herein, the term "effective amount" refers to the amount of an active composition (*e.g.*, a pharmaceutical compound or composition provided as a  
25 component in a lipid formulation) sufficient to effect beneficial or desired results. An effective amount can be administered in one or more administrations, applications or dosages and is not intended to be limited to a particular formulation or administration route.

30 [0039] As used herein, the terms "active" or "pharmaceutically active" as used in reference to an agent, composition, or compound, refers to an agent that, upon administration or application, causes a beneficial, desired, or expected result. The administration may be in one or more administrations, applications or dosages and is not intended to be limited to a particular formulation or administration route. The  
35 term is not limited to any particular level of activity. For example, a lipid formulation of an active agent need not have the same level of activity as a different



5 formulation of an active agent, so long as the active agent in the lipid formulation is sufficiently active that an effective amount of the active agent can be administered by administration of the lipid formulation of the agent.

10 [0040] The terms "agent" and "compound" are used herein interchangeably to refer to any atom, molecule, mixture, or more complex composition having an attributed feature. For example, an "active agent" or "active compound" refers to any atom, molecule, preparation, mixture, *etc.*, that, upon administration or application, causes a beneficial, desired, or expected result.

15 [0041] As used herein, the term "administration" refers to the act of giving a drug, prodrug, or other active agent, or therapeutic treatment (*e.g.*, compositions of the present invention) to a subject. Exemplary routes of administration to the human body can be through mouth and through skin. Administration may be in one or more administrations, applications or dosages, and is not intended to be limited to a particular administration route.

20 [0042] As used herein, the term "co-administration" refers to the administration of at least two agent(s) (*e.g.*, two separate lipid compositions, containing different active compounds) or therapies to a subject. In some embodiments, the co-administration of two or more agents or therapies is concurrent. In other embodiments, a first agent/therapy is administered prior to a second agent/therapy. 25 Those of skill in the art understand that the formulations and/or routes of administration of the various agents or therapies used may vary. The appropriate dosage for co-administration can be readily determined by one skilled in the art. In some embodiments, when agents or therapies are co-administered, the respective agents or therapies are administered at lower dosages than appropriate for their administration alone. Thus, co-administration is especially desirable in 30 embodiments where the co-administration of the agents or therapies lowers the requisite dosage of a potentially harmful (*e.g.*, toxic) agent(s).

35 [0043] As used herein, the term "toxic" refers to any detrimental or harmful effects on a subject, a cell, or a tissue as compared to the same cell or tissue prior to the administration of the toxicant.

5 [0044] As used herein, the term "pharmaceutical composition" refers to the combination of an active agent (*e.g.*, an active pharmaceutical compound) with a carrier, inert or active (*e.g.*, a phospholipid), making the composition especially suitable for diagnostic or topical therapeutic use.

10 [0045] The terms "pharmaceutically acceptable" or "pharmacologically acceptable," as used herein, refer to compositions that do not substantially produce adverse reactions, *e.g.*, toxic, allergic, or immunological reactions, when administered to a subject.

15 [0046] As used herein, the term "topically" refers to application of the compositions of the present invention to the surface of the skin and mucosal cells and tissues (*e.g.*, alveolar, buccal, lingual, masticatory, or nasal mucosa, and other tissues and cells which line hollow organs or body cavities).

20 [0047] As used herein, the term "pharmaceutically acceptable carrier" refers to any of the standard pharmaceutical carriers including, but not limited to non-toxic solvent, phosphate buffered saline solution, water, emulsions (*e.g.*, such as an oil/water or water/oil emulsions), and various types of wetting agents, any and all solvents, dispersion media, coatings, sodium lauryl sulfate, isotonic and absorption delaying agents, disintegrants (*e.g.*, potato starch or sodium starch glycolate), and the like.. The compositions also can include stabilizers and preservatives. For  
25 examples of carriers, stabilizers, and adjuvants. (See *e.g.*, Martin, Remington's Pharmaceutical Sciences, 15th Ed., Mack Publ. Co., Easton, Pa. (1975), incorporated herein by reference). Moreover, in certain embodiments, the compositions of the present invention may be formulated for horticultural or  
30 agricultural use. Such formulations include dips, sprays, seed dressings, stem injections, sprays, and mists.

35 [0048] As used herein, the term "pharmaceutically acceptable salt" refers to any salt (*e.g.*, obtained by reaction with an acid or a base) of a compound of the present invention that is physiologically tolerated in the target subject (*e.g.*, a mammalian subject). "Salts" of the compounds of the present invention may be derived from



inorganic or organic acids and bases. Examples of acids include, but are not limited to, hydrochloric, hydrobromic, sulfuric, nitric, perchloric, fumaric, maleic, phosphoric, glycolic, lactic, salicylic, succinic, toluene-p-sulfonic, tartaric, acetic, citric, methanesulfonic, ethanesulfonic, formic, benzoic, malonic, sulfonic, naphthalene-2-sulfonic, benzenesulfonic acid, and the like. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts.

**[0049]** Examples of bases include, but are not limited to, alkali metal (*e.g.*, sodium) hydroxides, alkaline earth metal (*e.g.*, magnesium) hydroxides, ammonia, and compounds of formula  $NW_4^+$ , wherein W is  $C_{1-4}$  alkyl, and the like.

**[0050]** Examples of salts include, but are not limited to: acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, flucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, chloride, bromide, iodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, palmoate, pectinate, persulfate, phenyl propionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, undecanoate, and the like. Other examples of salts include anions of the compounds of the present invention compounded with a suitable cation such as  $Na^+$ ,  $NH_4^+$ , and  $NW_4^+$  (wherein W is a  $C_{1-4}$  alkyl group), and the like. For therapeutic use, salts of the compounds of the present invention are contemplated as being pharmaceutically acceptable. However, salts of acids and bases that are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound.

**[0051]** For therapeutic use, salts of the compounds of the present invention are contemplated as being pharmaceutically acceptable. However, salts of acids and bases that are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound.

5 [0052] The term “Polyethylene glycol (PEG)” includes polymers of lower alkylene oxide, in particular ethylene oxide (polyethylene glycols) having an esterifiable hydroxyl group at least at one end of the polymer molecule, as well as derivatives of such polymers having esterifiable carboxyl groups. Polyethylene glycols of an average molecular weight ranging from 200-20,000 are preferred; those having an average molecular weight ranging from 500-2000 are particularly preferred.

10

[0053] The use of terms “a” and “an” and “the” and similar referents in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. The terms “comprising”, “including”, “having”, and “containing” are to be construed as open-ended terms (i.e. meaning “including but not limited to”) unless otherwise noted. The use of any and all examples, or exemplary language (e.g., “such as”) provided herein, is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention unless otherwise claimed. No language in the specifications should be constructed as indicating any non-claimed element as essential to the practice of the invention.

15

20

#### **DETAILED DESCRIPTION OF THE INVENTION**

25 [0054] This invention relates to the preparation of a solution, suspension, liposomes, lipid complex, or micelles in an aqueous or a non-toxic solvent system. The inventive preparation involves at least one phospholipid such as Soy phosphatidylcholine in aqueous media or in a non-toxic solvent with therapeutically active insoluble or poorly soluble compounds.

30 [0055] Particular embodiments of the invention are described in the Summary, and in this Detailed Description of the Invention. Although the invention has been described in connection with specific embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. For example, the compositions and methods of the present invention are described in connection with particular hair growth agent, such as finasteride or minoxidil. It

35



should be understood that the present invention is not limited to methods or compositions using or comprising finasteride or minoxidil.

5

[0056] In some embodiments, the present invention comprises a lipid composition with active compound (for example, finasteride) in which the composition contains lipid or a mixture of lipids. In some embodiments, the composition is in the form of solutions, micelles, emulsions or mixture of micelles and vesicles, gels, paste. The micelles of the present invention can be in the form, *e.g.*, of monomeric, dimeric, polymeric or mixed micelles. In some embodiments, the composition including micelles, emulsions or mixture of micelles and vesicles are predominately in the size range of 50nm-20 micron, while in some preferred embodiments, the micelles and emulsions are in the size range of 50nm-5 micron.

15

[0057] In some embodiments, active agent-lipid composition (for example, finasteride-lipid composition contain cholesterol or cholesterol derivatives. Examples of cholesterol derivatives that find use in the present invention include but are not limited to cholesteryl sulfate, cholesteryl hemisuccinate, cholesteryl succinate, cholesteryl oleate, cholesteryl linoleate, cholesteryl eicosapentenoate, cholesteryl linolenate, cholesteryl arachidonate, cholesteryl palmitate, cholesteryl stearate, cholesteryl myristate, polyethylene glycol derivatives of cholesterol (cholesterol-PEG), water soluble cholesterol (for example, cholesterol methyl- $\beta$ -cyclodextrin), coprostanol, cholestanol, or cholestane, cholic acid, cortisol, corticosterone or hydrocortisone and 7-dehydrocholesterol.

20  
25

[0058] In some preferred embodiments, the compositions also include  $\alpha$ -,  $\beta$ -,  $\gamma$ -tocopherols, vitamin E, calciferol, organic acid derivatives of  $\alpha$ -,  $\beta$ -,  $\gamma$ -tocopherols, such as  $\alpha$ -tocopherol hemisuccinate (THS),  $\alpha$ -tocopherol succinate, or mixtures thereof.

30

[0059] In some preferred embodiments, active agent-lipid compositions (for example, finsteride-lipid composition) contain sterols. Examples of sterols that find use in the present invention include  $\beta$ -sitosterol, stigmasterol, stigmastanol, lanosterol,  $\alpha$ -spinasterol, lathosterol, campesterol and/or mixtures thereof.

35

[0060] In some preferred embodiments, active agent-lipid composition ( for example, finasteride-lipid composition) contain guggulsterol and guggulsterol derivatives such as, guggul sulfate and its salt (preferably sodium salt), guggullaurate and guggulsterol derivatives containing long chain or short chain, saturated or unsaturated fatty acids (such as gugguloleate).

[0061] Compositions of the present invention also include active compounds (for example, finasteride) with free and/or salts or esters of fatty acid. In some preferred embodiments, fatty acids range from carbon chain lengths of about C<sub>2</sub> to C<sub>34</sub>, preferably between about C<sub>4</sub> and about C<sub>24</sub>, and include tetraoic acid (C<sub>4:0</sub>), pentanoic acid (C<sub>5:0</sub>), hexanoic acid (C<sub>6:0</sub>), heptanoic acid (C<sub>7:0</sub>), octanoic acid (C<sub>8:0</sub>), nonanoic acid (C<sub>9:0</sub>), decanoic acid (C<sub>10:0</sub>), undecanoic acid (C<sub>11:0</sub>), dodecanoic acid (C<sub>12:0</sub>), tridecanoic acid (C<sub>13:0</sub>), tetradecanoic (myristic) acid (C<sub>14:0</sub>), pentadecanoic acid (C<sub>15:0</sub>), hexadecanoic (palmatic) acid (C<sub>16:0</sub>), heptadecanoic acid (C<sub>17:0</sub>), octadecanoic (stearic) acid (C<sub>18:0</sub>), nonadecanoic acid (C<sub>19:0</sub>), eicosanoic (arachidic) acid (C<sub>20:0</sub>), heneicosanoic acid (C<sub>21:0</sub>), docosanoic (behenic) acid (C<sub>22:0</sub>), tricosanoic acid (C<sub>23:0</sub>), tetracosanoic acid (C<sub>24:0</sub>), 10-undecenoic acid (C<sub>11:1</sub>), 11-dodecenoic acid (C<sub>12:1</sub>), 12-tridecenoic acid (C<sub>13:1</sub>), myristoleic acid (C<sub>14:1</sub>), 10-pentadecenoic acid (C<sub>15:1</sub>), palmitoleic acid (C<sub>16:1</sub>), oleic acid (C<sub>18:1</sub>), linoleic acid (C<sub>18:2</sub>), linolenic acid (C<sub>18:3</sub>), eicosenoic acid (C<sub>20:1</sub>), eicosdienoic acid (C<sub>20:2</sub>), eicosatrienoic acid (C<sub>20:3</sub>), arachidonic acid (*cis*-5,8,11,14-eicosatetraenoic acid), and *cis*-5,8,11,14,17-eicosapentaenoic acid, among others. Other fatty acid chains also can be employed in the compositions. Examples of such include saturated fatty acids such as ethanoic (or acetic) acid, propanoic (or propionic) acid, butanoic (or butyric) acid, hexacosanoic (or cerotic) acid, octacosanoic (or montanic) acid, triacontanoic (or melissic) acid, dotriacontanoic (or lacceroic) acid, tetratriacontanoic (or gheddic) acid, pentatriacontanoic (or ceroplastic) acid, and the like; monoethenoic unsaturated fatty acids such as *trans*-2-butenic (or crotonic) acid, *cis*-2-butenic (or isocrotonic) acid, 2-hexenoic (or isohydrosorbic) acid, 4-decanoic (or obtusilic) acid, 9-decanoic (or caproleic) acid, 4-dodecenoic (or linderic) acid, 5-dodecenoic (or denticetic) acid, 9-dodecenoic (or lauroleic) acid, 4-tetradecenoic (or tsuzuic) acid, 5-tetradecenoic (or physeteric) acid, 6-octadecenoic (or petroselenic) acid, *trans*-9-octadecenoic ( or elaidic) acid, *trans*-11-octadecenoic ( or vaccinic) acid, 9-eicosenoic ( or gadoleic) acid, 11-eicosenoic ( or gondoic)



acid, 11-docosenoic (or cetoleic) acid, 13-decosenoic ( or erucic) acid, 15-tetracosenoic (or nervonic) acid, 17-hexacosenoic ( or ximenic) acid, 21-  
5 triacontenoic ( or lumequeic) acid, and the like; dienoic unsaturated fatty acids such as 2,4-pentadienoic (or  $\beta$ -vinylacrylic) acid, 2,4-hexadienoic (or sorbic) acid, 2,4-decadienoic (or stillingic) acid, 2,4-dodecadienoic acid, 9,12-hexadecadienoic acid, *cis*-9, *cis*-12-octadecadienoic (or  $\alpha$ -linoleic) acid, *trans*-9, *trans*-12-octadecadienoic (or linlolelaidic) acid, *trans*-10,*trans*-12-octadecadienoic acid, 11,14-eicosadienoic  
10 acid, 13,16-docosadienoic acid, 17,20-hexacosadienoic acid and the like; trienoic unsaturated fatty acids such as 6,10,14-hexadecatrienoic (or hiragonic) acid, 7,10,13-hexadecatrienoic acid, *cis*-6, *cis*-9- *cis*-12-octadecatrienoic (or  $\gamma$ -linoleic) acid, *trans*-8, *trans*-10- *trans*-12-octadecatrienoic (or  $\beta$ -calendic) acid, *cis*-8, *trans*-10- *cis*-12-octadecatrienoic acid, *cis*-9, *cis*-12- *cis*-15-octadecatrienoic (or  $\alpha$ -  
15 linolenic) acid, *trans*-9, *trans*-12- *trans*-15-octadecatrienoic (or  $\alpha$ -linolenelaidic) acid, *cis*-9, *trans*-11- *trans*-13-octadecatrienoic (or  $\alpha$ -eleostearic) acid, *trans*-9, *trans*-11- *trans*-13-octadecatrienoic (or  $\beta$ -eleostearic) acid, *cis*-9, *trans*-11- *cis*-13-octadecatrienoic (or puniic) acid, 5,8,11-eicosatrienoic acid, 8,11,14-eicosatrienoic acid and the like; tetraenoic unsaturated fatty acids such as 4,8,11,14-  
20 hexadecatetraenoic acid, 6,9,12,15- hexadecatetraenoic acid, 4,8,12,15-octadecatetraenoic (or moroctic) acid, 6,9,12,15- octadecatetraenoic acid, 9,11,13,15- octadecatetraenoic ( or  $\alpha$  -or  $\beta$ -parinaric) acid, 9,12,15,18-octadecatetraenoic acid, 4,8,12,16-eicosatetraenoic acid, 6,10,14,18-eicosatetraenoic acid, 4,7,10,13-docasatetraenoic acid, 7,10,13,16-docosatetraenoic acid, 8,12,16,19-  
25 docosatetraenoic acid and the like; penta- and hexa-enoic unsaturated fatty acids such as 4,8,12,15,18-eicosapentaenoic (or timnodonic) acid, 4,7,10,13,16-docosapentaenoic acid, 4,8,12,15,19-docosapentaenoic (or clupanodonic) acid, 7,10,13,16,19-docosapentaenoic, 4,7,10, 13,16,19-docosahexaenoic acid, 4,8,12,15,18,21-tetracosahexaenoic (or nisinic) acid and the like; branched-chain  
30 fatty acids such as 3-methylbutanoic (or isovaleric) acid, 8-methyldodecanoic acid, 10-methylundecanoic (or isolauric) acid, 11-methyldodecanoic (or isoundecylic) acid, 12-methyltridecanoic (or isomyristic) acid, 13-methyltetradecanoic (or isopentadecylic) acid, 14-methylpentadecanoic (or isopalmitic) acid, 15-methylhexadecanoic, 10-methylheptadecanoic acid, 16-methylheptadecanoic (or  
35 isostearic) acid, 18-methylnonadecanoic (or isoarachidic) acid, 20-methylheneicosanoic (or isobehenic) acid, 22-methyltricosanoic (or isolignoceric)

acid, 24-methylpentacosanoic (or isocerotic) acid, 26-methylheptacosanoic (or isomonatonic) acid, 2,4,6-trimethyloctacosanoic (or mycoceranic or mycoserosic) acid, 2-methyl-*cis*-2-butenoic(angelic)acid, 2-methyl-*trans*-2-butenoic (or tiglic) acid, 4-methyl-3-pentenoic (or pyroterebic) acid and the like.

**[0062]** Compositions of the present invention also include active compounds (for example, finasteride) with alcohols having long alkyl chain. In some preferred embodiments, alcohols range from carbon chain lengths of about C<sub>5</sub> to C<sub>34</sub>, preferably between about C<sub>4</sub> and about C<sub>24</sub>. The alkyl chain in the present invention may be saturated or unsaturated.

**[0063]** In certain preferred embodiments, active compounds (for example, finasteride) comprise phospholipids. Any suitable phospholipids can be used. For example, phospholipids can be obtained from natural sources or chemically synthesized. Examples of phospholipids that find use in the present invention include phosphatidylethanolamine (PE), phosphatidylglycerol (PG), phosphatidylserine (PS), phosphatidylcholine (PC), phosphatidylinositol (PI), phosphatidic acid (PA), sphingomyelin and the like, either used separately or in combination. Phosphatidylglycerols may be having short chain or long chain, saturated or unsaturated such as dimyristoylphosphatidylglycerol, dioleoylphosphatidylglycerol, distearoylphosphatidylglycerol, dipalmitoylphosphatidylglycerol, diarachidonoylphosphatidylglycerol, short chain phosphatidylglycerol (C<sub>6</sub>-C<sub>8</sub>), and mixtures thereof. Examples of phosphatidylcholines includes dimyristoylphosphatidylcholine, distearoylphosphatidylcholine, dipalmitoylphosphatidylcholine, dioleoylphosphatidylcholine, diarachidonoylphosphatidylcholine, egg phosphatidylcholine, soy phosphatidylcholine or hydrogenated soy phosphatidylcholine can be used, as can mixtures thereof.

**[0064]** In some embodiments, the present invention provides compositions comprising at least one active compound (for example, finasteride) and derivatives of mono-, di- and tri-glycerides. Examples of the glycerides that find use in the present invention include but are not limited to 1-oleoyl glycerol (monoolein) and 1, 2-dioctanoyl-*sn*-glycerol.



5 [0065] Another aspect of the invention is to provide a composition having at least one active compound (for example, finasteride) with at least one carbohydrate-based lipid. Examples of carbohydrate-based lipids that find use in the present invention include but are not limited to galactolipids, mannosylphosphatidylcholine, galactolecithin and the like.

10 [0066] Another aspect of the invention is to provide a composition comprising at least one active compound (for example, finsteride) with derivatives of phospholipids such as pegylated phospholipids. Examples include but not limited to the polyethylene glycol (Pegylated, PEG) derivatives of distearoylphosphatidylglycerol, dimyristoylphosphatidylglycerol, 15 dioleoylphosphatidylglycerol and the like.

[0067] Another further aspect of the present invention provides compositions comprising at least one active compound (for example, finasteride) and polyethylene glycol (PEG) and one or more lipids.

20

[0068] According to another aspect, the present invention provides compositions comprising at least one active compound (for example, finasteride) and one or more lipids. Example includes compositions comprising finasteride, cholesterol or cholesterol derivatives and one or more phospholipids. Other 25 examples of compositions according to the invention include finasteride,  $\beta$ -sitosterol, and one or more phospholipids. In some preferred embodiments, the composition of the present invention finasteride, cholesteryl sulfate and hydrogenated soy phosphatidylcholine or soy phosphatidylcholine.

30 [0069] According to another aspect, the present invention provides compositions comprising at least one active compound (for example, tacrolimus) and one or more lipids. Examples include compositions comprising tacrolimus, cholesterol or cholesterol derivatives and one or more phospholipids. Other 35 examples of compositions according to the invention include finasteride,  $\beta$ -sitosterol, and one or more phospholipids. In some preferred embodiments, the

composition of the present invention tacrolimus, cholesteryl sulfate and hydrogenated soy phosphatidylcholine or soy phosphatidylcholine.

5

[0070] According to another aspect, the present invention provides compositions comprising at least one active compound such as amphotericin B, and one or more lipids. Examples include compositions comprising amphotericin B, cholesterol or cholesterol derivatives and one or more phospholipids. Other examples of compositions according to the invention include amphotericin B,  $\beta$ -sitosterol, and one or more phospholipids. In some preferred embodiments, the composition of the present invention include amphotericin B, cholesteryl sulfate and hydrogenated soy phosphatidylcholine or soy phosphatidylcholine.

[0071] The composition of the present invention can be made by dissolving an active compound, for example, finasteride) in 70% isopropyl alcohol at a concentration of about 0.5 mg/mL to about 100 mg/mL. In some embodiments, the active compound is dissolved at a concentration between 1 mg/mL and about 50 mg/mL. In certain preferred embodiments, the active compound is dissolved at a concentration of between 1 mg/mL and 20 mg/mL. In particularly preferred embodiments, the active compound is dissolved at a concentration of between 1 mg/mL and 10 mg/mL.

[0072] In some embodiments, compositions of the present invention contain about 2.5% to about 95% by weight of total lipid, preferably about 10% to about 90% by weight of total lipid or more, preferably about 20% to about 90% by weight of total lipid.

[0073] In some embodiments, compositions of the present invention contain at least one active compound (for example, finasteride) and lipid(s) in mole ratio between 1:1 to 1:100, *e.g.*, in between 1:1 and 1:20 molar ratio or in between 1:1 and 1:30 molar ratio or in between 1:1 and 1:40 molar ratio or in between 1:1 and 1:50 molar ratio, in between 1:1 and 1:60 molar ratio, in between 1:1 and 1:70 molar ratios, and in between 1:1 and 1:80 molar ratios. As used herein, the term "in between" is inclusive of the limits of a recited range. For example, a mole ratio "in between" 1:1 and 1:20 molar ratio includes ratios of 1:1 and 1:20.



5 [0074] In certain preferred embodiments, the mole ratio of active compound (for example, finasteride) and cholesteryl sulfate in a composition containing active compound (for example, finasteride), cholesteryl sulfate and hydrogenated soy phosphatidylcholine is in between 1:0.25 and 1:20, such as in between 1:0.5 and 1:10, or in between 1:1 and 1:5 or 1:1 and 1:2. In particularly preferred  
10 embodiments, the mole ratio of active compound (for example, finasteride) and cholesteryl sulfate is in between 1:0.5 and 1:5.

15 [0075] In certain preferred embodiments, the mole ratio of active compound (for example, finasteride) and guggul sulfate in a composition containing active compound (for example, finasteride), guggul sulfate and hydrogenated soy phosphatidylcholine is in between 1:0.25 and 1:20, such as in between 1:0.5 and 1:10, or in between 1:1 and 1:5 or 1:1 and 1:2. In particularly preferred  
embodiments, the mole ratio of active compound (for example, finasteride) and guggul sulfate is in between 1:0.5 and 1:5

20 [0076] In certain preferred embodiments, the mole ratio of active compound (for example, finasteride) and hydrogenated soy phosphatidylcholine in a composition containing active compound (for example, finasteride), cholesteryl sulfate and hydrogenated soy phosphatidylcholine is in between about 1:1 and 1:90, *e.g.*, in between 1:1 and 1:70 or 1:1 and 1:60 or 1:1 and 1:50 or 1:1 and 1:40 and 1:1 and  
25 1:30. In particularly preferred embodiments, the mole ratio of active compound (for example, finasteride) and hydrogenated soy phosphatidylcholine is in between 1:5 and 1:60.

30 [0077] In certain preferred embodiments, the mole ratio of active compound (for example, finasteride) and soy phosphatidylcholine in a composition containing active compound (for example, finasteride), cholesteryl sulfate and soy phosphatidylcholine is in between 1:1 and 1:90, *e.g.*, in between 1:1 and 1:70 or 1:1 and 1:60 or 1:1 and 1:50 or 1:1 and 1:40 and 1:1 and 1:30. In particularly preferred  
35 embodiments, the mole ratio of active compound (for example, finasteride) and soy phosphatidylcholine is in between 1:5 and 1:60.

5 [0078] In certain preferred embodiments, the mole ratio of active compound (for example, finasteride) and soy phosphatidylcholine in a composition containing active compound (for example, finasteride), guggul sulfate and soy phosphatidylcholine is in between 1:1 and 1:90, *e.g.*, in between 1:1 and 1:70 or 1:1 and 1:60 or 1:1 and 1:50 or 1:1 and 1:40 and 1:1 and 1:30. In particularly preferred embodiments, the mole ratio of active compound (for example, finasteride) and soy phosphatidylcholine is in between 1:5 and 1:60.

10

[0079] In some embodiments, compositions of the present invention contain active compound (for example, finasteride) and total lipids having weight-to-weight ratio between 1:1 to 1:100 ratio such as in between 1:1 and 1:20 ratio or in between 1:1 and 1:30 ratio or in between 1:1 and 1:40 ratio or in between 1:1 and 1:50 ratio, 15 or in between 1:1 and 1:60 ratio, or in between 1:1 and 1:70 ratio, and in between 1:1 and 1:80 ratio, or in between 1:1 and 1:90 ratio.

[0080] In some embodiments, compositions of the present invention contain active compound (for example, finasteride) and soy phosphatidylcholine having weight-to-weight ratio between 1:1 to 1:100 ratio such as in between 1:1 and 1:20 ratio or in between 1:1 and 1:30 ratio or in between 1:1 and 1:40 ratio or in between 1:1 and 1:50 ratio, or in between 1:1 and 1:60 ratio, or in between 1:1 and 1:70 ratio, and in between 1:1 and 1:80 ratio, or in between 1:1 and 1:90 ratio.

25 [0081] In some embodiments, compositions of the present invention contain two active compounds (for example, finasteride and minoxidil) and soy phosphatidylcholine having weight-to-weight ratio between 1:1 to 1:100 ratio such as in between 1:1 and 1:20 ratio or in between 1:1 and 1:30 ratio or in between 1:1 and 1:40 ratio or in between 1:1 and 1:50 ratio, or in between 1:1 and 1:60 ratio, or 30 in between 1:1 and 1:70 ratio, and in between 1:1 and 1:80 ratio, or in between 1:1 and 1:90 ratio.

[0082] In some embodiments, the methods of the present invention involve solubilizing active compound(s) for example, finasteride and lipid(s), in water or suitable buffer. The active compound-lipid complex solution can be filtered 35 through suitable filters to control the size distribution of the formed complexes.



[0083] In some embodiments, the methods of the present invention involve  
5 dissolving active compound, *e.g.*, finasteride, in non-toxic solvent and mixing the  
dissolved active compound and the lipid(s) together. The non-toxic solvent in the  
present invention include but not limited to ethanol, 1-propanol, isopropanol,  
propylene glycol, and 1-butanol. The percentage of the non-toxic solvent in the  
present invention ranges between 100% and 10% such as in between 80% and 20%,  
10 such as in between 60% and 40%. In preferred embodiment the percentage of non-  
toxic solvent ranges between 80% and 50%.

[0084] In some embodiments, the methods of the present invention involve  
dissolving active compound, *e.g.*, finasteride, in non-toxic solvent-water system and  
15 mixing the dissolved active compound and the lipid(s) together. The non-toxic  
solvent in non-toxic solvent-water system in the present invention includes but not  
limited to ethanol, 1-propanol, isopropanol, propylene glycol, and 1-butanol. The  
percentage of the non-toxic solvent in the present invention ranges between 100%  
and 10% such as in between 80% and 20%, such as in between 60% and 40%. In  
20 preferred embodiment the percentage of non-toxic solvent ranges between 80% and  
50%.

[0085] In some embodiment, the methods of the present invention involve  
dissolving active compound(s) *e.g.* finasteride and/or minoxidil and lipid(s) together  
25 in more than one non-toxic solvents and water. For example, finasteride and or  
monoxidil can be dissolved in isopropyl alcohol-propylene glycol-water system.  
The percentage of the total non-toxic solvent in the present invention ranges  
between 90% and 10% such as in between 80% and 20%, such as in between 60%  
and 40%. In preferred embodiment the percentage of non-toxic solvent ranges  
30 between 80% and 50%.

[0086] In some embodiments, the methods of the present invention involve  
mixing active compound and lipid (s) for example, finasteride and soy  
phosphatidylcholine in isopropyl alcohol-water solution until the solution is  
35 completely clear. The percentage of isopropyl alcohol in isopropyl-water solution

ranges between 10 and 90 percent such as in between 40 and 80 percent, such as in between 50 and 70 percent.

5

[0087] In some embodiments, the methods of the present invention involve mixing active compound and lipid (s) for example, finasteride and soy phosphatidylcholine in ethanol-water solution until the solution is completely clear. The percentage of ethanol in isopropyl-water solution ranges between 10 and 90 percent such as in between 40 and 80 percent, such as in between 50 and 70 percent

10

[0088] In some embodiments, the compositions of present invention involve more than one active compound and one more lipids. For example, the composition contains finasteride as one active agent and minoxidil as second active agent. The mole ratio of one active agent to another active agent in the present invention ranges in between 1:0.5 to 1:100, *e.g.*, in between 1:1 and 1:20 molar ratio or in between 1:1 and 1:30 molar ratio or in between 1:1 and 1:40 molar ratio or in between 1:1 and 1:50 molar ratio, in between 1:1 and 1:60 molar ratio, in between 1:1 and 1:70 molar ratios, and in between 1:1 and 1:80 molar ratios. As used herein, the term "in between" is inclusive of the limits of a recited range. For example, a mole ratio "in between" 1:1 and 1:20 molar ratio includes ratios of 1:1 and 1:20.

15

20

[0089] In some embodiments, the methods of the present invention involve mixing active compound(s) (for example, finasteride and minoxidil) and lipid(s) for example, soy phosphatidylcholine in isopropyl alcohol-water solution until the solution is completely clear. The percentage of isopropyl alcohol in isopropyl-water solution ranges between 30 and 90 percent such as in between 40 and 80 percent, such as in between 50 and 70 percent.

25

[0090] In some embodiments, the methods of the present invention involve mixing active compound(s) (for example, finasteride and / or minoxidil) and lipid(s) for example, soy phosphatidylcholine in ethanol-water solution until the solution is completely clear. The percentage ethanol in ethanol-water solution ranges between 30 and 90 percent such as in between 40 and 80 percent, such as in between 50 and 70 percent.

30

35



[0091] In some embodiments, the methods of the present invention involve mixing active compound(s) (for example, finasteride and / or minoxidil) and lipid(s) for example, soy phosphatidylcholine in ethanol-water solution containing polysorbate 80 until the solution is completely clear. The percentage of ethanol in ethanol -water solution ranges between 30 and 90 percent such as in between 40 and 80 percent, such as in between 50 and 70%. The percentage of polysorbate 80 in ethanol-water mixture ranges between 10 and 60 percent such as in between 20 and 50 percent such as in between 25 and 40 percent.

[0092] In some embodiments, the methods of the present invention involve mixing active compound(s) (for example, finasteride and / or minoxidil) and lipid(s) for example, soy phosphatidylcholine in isopropyl alcohol-water solution containing polysorbate 80 until the solution is completely clear. The percentage of isopropyl alcohol in isopropyl-water solution ranges between 30 and 90 percent such as in between 40 and 80 percent, such as in between 50 and 70 percent. The percentage of polysorbate 80 in ethanol-water mixture ranges between 10 and 60 percent such as in between 20 and 50 percent such as in between 25 and 40 percent

[0093] In some embodiments, the methods of the present invention involve mixing active compound(s) (for example, finasteride and / or minoxidil) and lipid(s) for example, soy phosphatidylcholine in oleyl alcohol until the solution is clear.

[0094] In some embodiment, the methods of present invention involve mixing active compound(s), (for example, finasteride and / or minoxidil) and lipid(s) for example, soy phosphatidylcholine in propylene glycol and homogenize or sonicate until the solution is completely clear.

[0095] In some embodiments, the methods of the present invention involve mixing active compound(s) (for example, finasteride and / or minoxidil) and lipid(s) for example, soy phosphatidylcholine in propylene glycol-isopropyl alcohol-water solution until the solution is completely clear. The percentage of isopropyl alcohol in propylene glycol-isopropyl alcohol-water solution ranges between 10 and 90 percent such as in between 30 and 80 percent, such as in between 40 and 70 percent. The percentage of propylene glycol in propylene glycol-isopropyl alcohol-water

solution ranges between 10 and 80 percent such as in between 20 and 60 percent such as in between 25 and 50 percent.

5

[0096] In some embodiments, the methods of the present invention involve mixing active compound(s) (for example, finasteride and / or minoxidil) and lipid(s) for example, soy phosphatidylcholine in propylene glycol-ethanol-water solution until the solution is completely clear. The percentage of ethanol in propylene glycol-ethanol-water solution ranges between 10 and 90 percent such as in between 30 and 80 percent, such as in between 40 and 70 percent. The percentage of propylene glycol in propylene glycol-ethanol-water solution ranges between 10 and 80 percent such as in between 20 and 60 percent such as in between 25 and 50 percent.

10

[0097] In some embodiments, the methods of the present invention involve mixing active compound(s) (for example, finasteride and / or minoxidil) and lipid(s) for example, soy phosphatidylcholine in suitable oil and homogenize or sonicate until it is completely clear. Examples of suitable oil includes but not limited to vegetable oil, such as olive oil, castor oil, saff flower oil, sunflower oil, canola oil, soybean oil, pea nut oil, coconut oil, sesame seed oil, almond oil.

15

[0098] In some embodiments, compositions of the present invention having more than one active agent, (for example, finasteride and minoxidil), the weight-to-weight ratio of one active agent and the second active agent is in between 1:0.5 to 1:100 ratio such as in between 1:1 and 1:20 ratio or in between 1:1 and 1:30 ratio or in between 1:1 and 1:40 ratio or in between 1:1 and 1:50 ratio, or in between 1:1 and 1:60 ratio, or in between 1:1 and 1:70 ratio, and in between 1:1 and 1:80 ratio, or in between 1:1 and 1:90 ratio.

20

25

[0099] In some embodiments, the method of the present invention involves mixing lipid(s) and active compound(s) together in non-toxic solvent. The active compound-lipid composition solution can be filtered through suitable filters to control the size distribution of the formed complexes.

30

35



[00100] In some embodiments, the method of preparation of present invention comprising mixing active compound (for example, finasteride), cholesteryl derivative (for example, cholesteryl sulfate) and phosphatidylcholine such as soy phosphatidylcholine or hydrogenated soy phosphatidylcholine in water or buffer. The resulting suspension can be homogenized or sonicated at any desired temperature ranging from 20-60°C. Examples of base or buffer includes but not limited to sodium succinate dibasic, sodium acetate, sodium phosphate monobasic, sodium phosphate dibasic, sodium phosphate tribasic, sodium citrate, sodium hydroxide, and the like. The resulting suspension can be homogenized or sonicated to reduce the particle size. In some embodiments, the hydrated suspension is filtered through suitable filters to control the size distribution of the formed complexes. In some composition, the hydrated suspension can be lyophilized to obtain the composition in powder form. In some embodiments, the hydrated composition can be autoclaved. In some embodiments, a cryoprotectant such as sugars may be added to facilitate lyophilization. Examples of sugars include but not limited to trehalose, dextrose, sucrose, lactose, maltose, and the like. The percentage of sugar may range from 5% to about 25%.

20

[00101] In some embodiments, a method of preparation of the present invention comprising mixing one or more active compounds (*e.g.*, tacrolimus or amphotericin B), a cholesterol derivative (*e.g.*, cholesteryl sulfate) and a phosphatidyl choline such as soy phosphatidylcholine or hydrogenated soy phosphatidylcholine in ethanol or isopropanol. The resulting solution or suspension can be mixed with suitable base used commonly used is ointments and sprays.

25

[00102] In some embodiments, the composition of the present invention can be sterilized by filtering through 0.22  $\mu\text{m}$  or 0.45  $\mu\text{m}$  filter under aseptic conditions. In other embodiments, the composition of the present invention can be sterilized by autoclaving in the range of 110°C-130°C for duration of 15-20 minutes.

30

[00103] Pharmaceutical preparations that find use in the present invention include but are not limited to solutions, suspensions, emulsions, ointments; suppositories, gels can be suitable pharmaceutical preparations. In some embodiments, *e.g.*, for topical application and suppositories, active compound-lipid

35

composition (for example, finasteride-lipid composition) is provided in the form of solutions, gels, oils, and emulsions, such as are known by the addition of suitable water-soluble or water-insoluble excipients, for example polyethylene glycols, certain fats, and esters, compounds having a higher content of polyunsaturated fatty acids and derivatives thereof. Derivatives include but are not limited to mono-, di-, and triglycerides and their aliphatic esters (for example, fish oils, vegetable oils etc.) or mixtures of these substances. In some embodiments, excipients that find use in conjunction with the compositions of the present invention comprise those in which the drug complexes are sufficiently stable to allow for therapeutic use. In some embodiments, the oral mode of administration, the composition of present invention is used in the form of tablets, capsules, lozenges, powders, syrups, aqueous solutions, suspensions, and the like.

15

**[00104]** In some embodiments, the composition of the present invention can be admixed with other carrier substances known in the art. Examples of carrier substances include but not limited to glycerine, oils, mineral oil, propylene glycol, alcohol, aloe vera gel, allantoin, vitamin A and E, PPG2 myristyl propionate, and the like. The said mixture of composition and carrier substance can be administered topically in the form of solutions, creams, gels, lotions, shampoo, paste (See, e.g, Rasmusson *et al.*, EP 0285 382, 1988).

20

**[00105]** Preparations of active compound-lipid composition (for example, finasteride-lipid composition) of the present invention can comprise composition of varying size, or can comprise composition of substantially uniform size. For example, in some embodiments the composition have a size range of about 1 mm or less, while in preferred embodiments, the composition are in the micron or sub-micron range. In some embodiments, the composition have a diameter of about 5  $\mu\text{m}$  or less, such as 0.2  $\mu\text{m}$  or less, or even 01.  $\mu\text{m}$  or less.

25  
30

**[00106]** Active compound-lipid composition (for example, finasteride-lipid composition) of the present invention may comprise or consist essentially of micelles, mixed micelles, liposomes and vesicles of different shape and sizes.

35



5 [00107] As noted above, the technology outlined in the present invention for the preparation of finasteride composition is also suitable for use with any other water-insoluble drugs.

10 [00108] The composition of present invention can be employed (for example, finasteride-lipid composition or finasteride-minoxidil-lipid composition) to treat androgenic alopecia and promoting hair growth in males. In some embodiments, the composition of present invention can also be employed in preventing hair loss in males. In some embodiments, the composition of present invention can also be employed to treat hirsutism in females. In some embodiments, the composition of present invention can also be used to treat acne vulgaris and seborrhea.

15 [00109] The composition of present invention (for example, finasteride-lipid composition) can also be employed to treat benign prostatic hyperplasia in males.

20 [00110] The examples of the present invention are illustrated below but the invention is not limited to the following examples and modifications can be made without departing from the purports described in this application.

#### EXAMPLE I

25 [00111] Finasteride (100 mg) and Soya Phosphatidylcholine (4 g) was blended together in sodium citrate, monobasic (100 mL) and then subjected to high pressure homogenization. The particle size was determined using Nicomp Particle Sizer 380. The mean volume diameter amounted to less than 200 nm.

#### EXAMPLE II

30 [00112] Finasteride (100 mg), sodium cholesteryl sulfate (66 mg) and Soya Phosphatidylcholine (3.94 g) was blended together in sodium citrate, monobasic (100 mL) and then subjected to high pressure homogenization. The particle size was determined using Nicomp Particle Sizer 380. The mean volume diameter amounted to less than 200 nm.

35

**EXAMPLE III**

5 [00113] Finasteride (10 mg) and Soya Phosphatidylcholine (25 mg) was mixed together in soybean oil (2 mL) and sonicated for 40 minutes.

**EXAMPLE IV**

10 [00114] Finasteride (200 mg) and Soy phosphatidylcholine (1 g) was mixed together in 70% isopropyl alcohol (40 mL) and sonicated until a solution is formed.

**EXAMPLE V**

[00115] Finasteride (10 mg), soy phosphatidylcholine (50 mg), and polysorbate 80 (0.5 mL) was solubilized in ethanol (0.5 mL) and water (1 mL) was added and stirred vigorously.

15

**EXAMPLE VI**

[00116] Finasteride (10 mg), soy phosphatidylcholine (50 mg), and polysorbate 80 (0.5 mL) was solubilized in ethanol (0.5 mL) and water (1 mL) was added and stirred vigorously.

**EXAMPLE VII**

20 [00117] Finasteride (10 mg) and soy phosphatidylcholine (50 mg) was solubilized in isopropyl alcohol (0.5 mL) and water (0.5 mL) was added and stirred vigorously.

**EXAMPLE VIII**

25 [00118] Finasteride (10 mg) and soy phosphatidylcholine (50 mg) was solubilized in isopropyl alcohol (2.5 mL) and water (2.5 mL) was added and stirred vigorously.

**EXAMPLE VIII**

30 [00119] Finasteride ( 2 mg ) and soy phosphatidylcholine (10 mg) was solubilized in 70% isopropyl alcohol in water (1 mL) and added to a suspension of Minoxidil (50 mg) in 60% propylene glycol in water (1 mL). The suspension was homogenized or sonicated until the solution is clear.

**EXAMPLE IX**

35 [00120] Finasteride ( 2 mg ) and soy phosphatidylcholine (20 mg) was solubilized in 70% isopropyl alcohol in water (1 mL) and added to a suspension of



Minoxidil (80 mg) in 60% propylene glycol in water (1 mL). The suspension was homogenized or sonicated until the solution is clear.

5

**EXAMPLE X**

[00121] Finasteride (2 mg), Minoxidil (100 mg) and soy phosphatidylcholine (20 mg) was solubilized in propylene glycol (2 mL).

10

**EXAMPLE XI**

[00122] Finasteride (2 mg), Minoxidil (100 mg) and soy phosphatidylcholine (20 mg) was solubilized in a solution containing isopropyl alcohol (0.7 mL), propylene glycol (0.6 mL) and water (0.6 mL).

[00123]

15

**EXAMPLE XII**

[00124] Finasteride (2 mg), Minoxidil (100 mg) and soy phosphatidylcholine (20 mg) was solubilized in a solution containing isopropyl alcohol (0.0.32mL), propylene glycol (1.2 mL) and water (0.48 mL).

20

**EXAMPLE XIII**

[00125] Finasteride (2 mg), Minoxidil (100 mg) and soy phosphatidylcholine (20 mg) was solubilized in a solution containing isopropyl alcohol (1.2 mL), propylene glycol (0.4 mL) and water (0.4 mL).

25

**EXAMPLE XIV**

[00126] Finasteride (2 mg), Minoxidil (100 mg) and soy phosphatidylcholine (20 mg) was solubilized in a solution containing isopropyl alcohol (0.8 mL), propylene glycol (0.6 mL) and water (0.6 mL).

30

**EXAMPLE XIV**

[00127] Finasteride (2 mg), Minoxidil (100 mg) and soy phosphatidylcholine (20 mg) was solubilized in a solution containing isopropyl alcohol (0.8 mL), propylene glycol (0.6 mL) and water (0.6 mL).

35

**EXAMPLE XV**

5 [00128] Finasteride (20 mg), Minoxidil (1 g ) and soy phosphatidylcholine (20 mg) was solubilized in a solution containing isopropyl alcohol (0.4 mL), propylene glycol (1.2 mL) and water (0.4 mL).

**EXAMPLE XVI**

10 [00129] Finasteride (20 mg) and Minoxidil (1 g) and soy phosphatidylcholine (200 mg ) was taken in propylene glycol (20 mL) and homogenized or sonicated until all ingredients are completely soluble.

**EXAMPLE XVII**

15 [00130] Finasteride (20 mg) and Minoxidil ( 1g ) and soy phosphatidylcholine (200 mg) was taken in propylene glycol (20 mL) and homogenized or sonicated until the solution is completely clear.

**EXAMPLE XVIII**

20 [00131] Finasteride (40 mg) and Minoxidil ( 2 g ) and soy phosphatidylcholine (200 mg) was taken in isopropanol:propylene glycol:water (50:30:20, 40 mL) and homogenized or sonicated until the solution is completely clear.

**EXAMPLE XIX**

25 [00132] Finasteride (40 mg) and Minoxidil ( 2 g ) and soy phosphatidylcholine (400 mg) was taken in isopropanol:propylene glycol:water (50:30:20, 40 mL) and homogenized or sonicated until the solution is completely clear.

**EXAMPLE XX**

30 [00133] Finasteride (40 mg) and Minoxidil ( 2 g ) and soy phosphatidylcholine (200 mg) was taken in isopropanol:propylene glycol:water (50:40:10, 40 mL) and homogenized or sonicated until the solution is completely clear.



**EXAMPLE XXI**

5 [00134] Finasteride (40 mg) and Minoxidil ( 2 g ) and soy phosphatidylcholine (400 mg) was taken in isopropanol:propylene glycol:water (50:40:10, 40 mL) and homogenized or sonicated until the solution is completely clear.

**EXAMPLE XXII**

10 [00135] Finasteride (40 mg) and Minoxidil ( 2 g ) and soy phosphatidylcholine (200 mg) was taken in isopropanol:propylene glycol:water (40:50:10, 40 mL) and homogenized or sonicated until the solution is completely clear.

**EXAMPLE XXIII**

15 [00136] Finasteride (40 mg) and Minoxidil ( 2 g ) and soy phosphatidylcholine (400 mg) was taken in isopropanol:propylene glycol:water (40:50:10, 40 mL) and homogenized or sonicated until the solution is completely clear.

**EXAMPLE XXIV**

20 [00137] Tacrolimus (20 mg) and soy phosphatidylcholine (200 mg) was taken in isopropanol and homogenized or sonicated until the solution is completely clear.

**REFERENCES**

1. Deray G.; Mercadal, L.; Bagnis, C. *Nephrologie*, **2002**, 23, 119-122.
- 25 2. Gormley, G.J., Kauffmann, K.D., Stoner, E., Waldstreicher, J. US Patent No 5,981,543, 1999.
3. Hammarstrom, L.; and Smith, C.I.E. *Acta Patho. Microbial. Scand.* **1977**, 85, 277-283.
4. Roberts, J. *Dermatology Nursing*, **1977**, 9, 379-386
- 30 5. Roberts, J. *J. Amer. Acad. Derm.*, **1987**, 16, 705-710.
6. Medoff, G. and Kobayashi, G.S. *J. Am. Med. Assoc.* **1975**, 232, 619-620.
7. Niemiec, S.M., Wang, J.C.T., Wisniewski, S.J., Stenn, K.S., Lu, G.W. US Patent No 6,419,913 B1, 2002
8. Olsen, E.A., Weiner, M.S., DeLong, E.R., Pinnell, S. *J. Amer. Acad. Derm.*, **1985**,  
35 13, 185-192.
9. Rasmusson, G.H., Johnston, D.B.R., Arth, G.E. US Patent No. 4,377,584, 1983.

10. Rasmusson, G.H., Reynolds, G.F. EP Patent No. 0285382B1, 1988

11. Valeriote, F.; Lynch, R.; Medoff, G.; and Kumar, B.V. *J. Natl. Cancer. Inst.* 1976,  
5 56, 557-559.

[00138]

10

[00139] Preferred embodiments of this invention are described, including the best  
mode known to the inventors for carrying out the invention.

15

Accordingly, this invention includes all modifications  
and equivalents of the subject matter recited in the claims appended hereto as  
permitted by applicable law. Moreover, any combination of the above-described  
20 elements in all possible variations thereof is encompassed by the invention unless  
otherwise indicated herein or otherwise clearly contradicted by context. Indeed, any  
modifications of the described modes for carrying out the invention that are obvious  
to those skilled in the relevant fields are intended to be within the scope of the  
25 following claims.



**Claims:**

1. A composition for use as a topical or transdermal treatment of a human subject for: androgenic alopecia; or, to prevent or reduce hair loss; wherein the composition comprises:
  - an effective amount of finasteride and minoxidil; and,
  - a phosphatidylcholine;
  - in an aqueous formulation.
  
2. The composition of claim 1, further comprising an active compound selected from the group consisting of Duasteride, Amphotericin B, and Tacrolimus.
  
3. The composition of claim 1 or 2, wherein said phosphatidylcholine is egg phosphatidylcholine (EPC), soy phosphatidylcholine (SPC), hydrogenated soy phosphatidylcholine (HSPC), dimyristoylphosphatidylcholine (DMPC), or dipalmitoylphosphatidylcholine (DPPC).
  
4. The composition of any one of claims 1 to 3, further comprising one or more of: egg phosphatidylglycerol (EPG), dimyristoylphosphatidylglycerol (DMPG), disteroylphosphatidylglycerol (DSPG), dipalmitoylphosphatidylglycerol (DMPG), phosphatidylethanolamines, phosphatidylserine, phosphatidylinositol, phosphatidic acid, cholesterol (Chol), cholesterol sulfate and its salts (CS), cholesterol hemisuccinate and its salts (Chems), cholesterol phosphate and its salts (CP), cholesterylphosphocholine and other hydroxycholesterol or amino cholesterol derivatives, cholesteryl succinate, cholesteryl oleate, polyethylene glycol derivatives of cholesterol (cholesterol-PEG), coprostanol, cholestanol, cholestane, cholic acid, cortisol, corticosterone, hydrocortisone, and calciferol; guggulsterol-and guggulsterol derivatives containing long chain or short chain, saturated or unsaturated fatty acids, guggul sulfate and its salts; carbohydrate-based lipids selected from a group consisting of galactolipid, mannosylphosphatidylcholine, galactolecithin; fatty acids selected from a group consisting of saturated or unsaturated fatty acids having a chain length of C<sub>4</sub>-C<sub>34</sub>; long chain alcohols selected from a group consisting of saturated or unsaturated alcohols having a chain length of C<sub>5</sub>-C<sub>34</sub>;

pegylated derivative of a phospholipid selected from the group consisting of pegylated derivatives of disteoylphosphatidylglycerol, dipalmitoylphosphatidylglycerol, dimyristoylphosphatidylglycerol, or dioleoylphosphatidylglycerol.

5. The composition of any one of claims 1-4, wherein the composition further comprises polyethylene glycol (PEG) having an average molecular weight ranging from about 200-20,000.
6. The composition of any one of claims 1-5, wherein the composition further comprises one or more of isopropanol, propylene glycol, ethanol, propyl alcohol, or butanol.
7. The composition of any one of claims 1-6, wherein the composition further comprises one or more of: soybean oil, vegetable oil, olive oil, coconut oil, sunflower oil, almond oil, canola oil, sesame seed oil, peanut oil, corn oil, or cod oil.
8. The composition of any one of claims 1-7, wherein said formulation is a solution, suspension, emulsion, micelle, liposomal formulation, lipidic particle formulation, gel, shampoo, paste, or cream form.
9. The composition of any one of claims 1 to 8, wherein said composition comprises soy phosphatidylcholine and 70% isopropyl alcohol in water.
10. The composition of any one of claims 1 to 8, wherein said composition comprises tacrolimus, cholesteryl sulfate, and soy phosphatidylcholine in isopropanol or ethanol.
11. The composition of any one of claims 1 to 8, wherein said composition comprises amphotericin B, cholesteryl sulfate, and soy phosphatidylcholine in isopropanol or ethanol.
12. Use of a composition for topical or transdermal treatment of a human subject for: androgenic alopecia; or, to prevent or reduce hair loss; wherein the composition comprises: an effective amount of finasteride and minoxidil; and,



a phosphatidylcholine;  
in an aqueous formulation.

13. The use according to claim 12, wherein the composition further comprises an active compound selected from the group consisting of Duasteride, Amphotericin B, and Tacrolimus.

14. The use according to claim 12 or 13, wherein said phosphatidylcholine is egg phosphatidylcholine (EPC), soy phosphatidylcholine (SPC), hydrogenated soy phosphatidylcholine (HSPC), dimyristoylphosphatidylcholine (DMPC), or dipalmitoylphosphatidylcholine (DPPC).

15. The use according to any one of claims 12 to 14, wherein the composition further comprises one or more of: egg phosphatidylglycerol (EPG), dimyristoylphosphatidylglycerol (DMPG), distearylphosphatidylglycerol (DSPG), dipalmitoylphosphatidylglycerol (DMPG), phosphatidylethanolamines, phosphatidylserine, phosphatidylinositol, phosphatidic acid, cholesterol (Chol), cholesterol sulfate and its salts (CS), cholesterol hemisuccinate and its salts (Chems), cholesterol phosphate and its salts (CP), cholesterylphosphocholine and other hydroxycholesterol or amino cholesterol derivatives, cholesteryl succinate, cholesteryl oleate, polyethylene glycol derivatives of cholesterol (cholesterol-PEG), coprostanol, cholestanol, cholestane, cholic acid, cortisol, corticosterone, hydrocortisone, and calciferol; guggulsterol and other guggulsterol derivatives containing long chain or short chain, saturated or unsaturated fatty acids, guggul sulfate and its salts; carbohydrate-based lipids selected from a group consisting of galactolipid, mannanolipid, galactolecithin; fatty acids selected from a group consisting of saturated or unsaturated fatty acids having a chain length of C<sub>4</sub>-C<sub>34</sub>; long chain alcohols selected from a group consisting of saturated or unsaturated alcohols having a chain length of C<sub>5</sub>-C<sub>34</sub>; pegylated derivative of a phospholipid selected from the group consisting of pegylated derivatives of distearylphosphatidylglycerol, dipalmitoylphosphatidylglycerol, dimyristoylphosphatidylglycerol, or dioleoylphosphatidylglycerol.

16. The use according to any one of claims 12 to 15, wherein the composition further comprises polyethylene glycol (PEG) having an average molecular weight ranging from about 200-20,000.
17. The use according to any one of claims 12 to 16, wherein the composition further comprises one or more of isopropanol, propylene glycol, ethanol, propyl alcohol, or butanol.
18. The use according to any one of claims 12 to 17, wherein the composition further comprises one or more of: soybean oil, vegetable oil, olive oil, coconut oil, sunflower oil, almond oil, canola oil, sesame seed oil, peanut oil, corn oil, or cod oil.
19. The use according to any one of claims 12 to 18, wherein said formulation is a solution, suspension, emulsion, micelle, liposomal formulation, lipidic particle formulation, gel, shampoo, paste, or cream form.
20. The use according to any one of claims 12 to 19, wherein said composition comprises soy phosphatidylcholine and 70% isopropyl alcohol in water.
21. The use according to any one of claims 12 to 19, wherein said composition comprises tacrolimus, cholesteryl sulfate, and soy phosphatidylcholine in isopropanol or ethanol.
22. The use according to any one of claims 12 to 19, wherein said composition comprises amphotericin B, cholesteryl sulfate, and soy phosphatidylcholine in isopropanol or ethanol.
23. The composition of any one of claims 1 to 11, or the use according to any one of claims 12 to 22, wherein the human subject is a male.