

FASCICULE DEBREVET D'INVENTION

21 Numéro de dépôt: 1201300105
(PCT/US11/029441)

22 Date de dépôt : 22/03/2011

30 Priorité(s) :
US n° 12/885,917 du 20/09/2010
US n° 13/050,820 du 17/03/2011

24 Délivré le : 31/03/2014

45 Publié le : 11.05.2015

73 Titulaire(s) :

GILEAD SCIENCES, INC.,
333 Lakeside Drive,
FOSTER CITY, CA 94404 (US)

72 Inventeur(s) :


CHO, Aesop (US)
KIM, Choung (US)
RAY, Adrian (US)

74 Mandataire : SCP AKKUM, AKKUM & Associates,
Quartier Mballa II, Dragages,
B.P. 4966, YAOUNDE (CM).

54 Titre : 2'-fluoro substituted carba-nucleoside analogs for antiviral treatment.

57 Abrégé :

Provided are select imidazo[1,2-f][1,2,4] triazinyl nucleosides, nucleoside phosphates and prodrugs thereof, wherein the 2' position of the nucleoside sugar is substituted with halogen and carbon substituents. The compounds, compositions, and methods provided are useful for the treatment of Flaviviridae virus infections, particularly hepatitis C infections caused by both wild type and mutant strains of HCV.



5

2'-FLUORO SUBSTITUTED CARBA-NUCLEOSIDE ANALOGS FOR
ANTIVIRAL TREATMENT

10

FIELD OF THE INVENTION

The invention relates generally to compounds with antiviral activity, more particularly nucleosides active against *Flaviviridae* infections and most particularly to inhibitors of hepatitis C virus RNA-dependent RNA polymerase.

15

BACKGROUND OF THE INVENTION

Viruses comprising the *Flaviviridae* family comprise at least three distinguishable genera including *pestiviruses*, *flaviviruses*, and *hepaciviruses* (Calisher, *et al.*, J. Gen. Virol., 1993, 70, 37-43). While *pestiviruses* cause many economically important animal diseases such as bovine viral diarrhea virus (BVDV), classical swine fever virus (CSFV, hog cholera) and border disease of sheep (BDV), their importance in human disease is less well characterized (Moennig, V., *et al.*, Adv. Vir. Res. 1992, 48, 53-98). *Flaviviruses* are responsible for important human diseases such as dengue fever and yellow fever while *hepaciviruses* cause hepatitis C virus infections in humans. Other important viral infections caused by the *Flaviviridae* family include West Nile virus (WNV) Japanese encephalitis virus (JEV), tick-borne encephalitis virus, Junjin virus, Murray Valley encephalitis, St Louis encephalitis, Omsk hemorrhagic fever virus and Zika virus. Combined, infections from the *Flaviviridae* virus family cause significant mortality, morbidity and economic losses throughout the world. Therefore, there is a need to develop effective treatments for *Flaviviridae* virus infections.



5 The hepatitis C virus (HCV) is the leading cause of chronic liver disease worldwide (Boyer, N. et al. *J Hepatol.* 32:98-112, 2000) so a significant focus of current antiviral research is directed toward the development of improved methods of treatment of chronic HCV infections in humans (Di Besceglie, A.M. and Bacon, B. R., *Scientific American*, Oct.: 80-85, (1999); Gordon, C. P., et al., *J. Med. Chem.* 10 **2005**, 48, 1-20; Maradpour, D.; et al., *Nat. Rev. Micro.* **2007**, 5(6), 453-463). A number of HCV treatments are reviewed by Bymock et al. in *Antiviral Chemistry & Chemotherapy*, 11:2; 79-95 (2000).

 RNA-dependent RNA polymerase (RdRp) is one of the best studied targets for the development of novel HCV therapeutic agents. The NS5B polymerase is a target 15 for inhibitors in early human clinical trials (Sommadosi, J., WO 01/90121 A2, US 2004/0006002 A1). These enzymes have been extensively characterized at the biochemical and structural level, with screening assays for identifying selective inhibitors (De Clercq, E. (2001) *J. Pharmacol. Exp. Ther.* 297:1-10; De Clercq, E. (2001) *J. Clin. Virol.* 22:73-89). Biochemical targets such as NS5B are important in 20 developing HCV therapies since HCV does not replicate in the laboratory and there are difficulties in developing cell-based assays and preclinical animal systems.

 Currently, there are primarily two antiviral compounds, ribavirin, a nucleoside analog, and interferon-alpha (α) (IFN), that are used for the treatment of chronic HCV infections in humans. Ribavirin alone is not effective in reducing viral RNA levels, 25 has significant toxicity, and is known to induce anemia. The combination of IFN and ribavirin has been reported to be effective in the management of chronic hepatitis C (Scott, L. J., et al. *Drugs* **2002**, 62, 507-556) but less than half the patients infected with some genotypes show a persistent benefit when given this treatment. Other patent applications disclosing the use of nucleoside analogs to treat hepatitis C virus 30 include WO 01/32153, WO 01/60315, WO 02/057425, WO 02/057287, WO 02/032920, WO 02/18404, WO 04/046331, WO2008/089105 and WO2008/141079 but additional treatments for HCV infections have not yet become available for patients.

X

5 Virologic cures of patients with chronic HCV infection are difficult to achieve because of the prodigious amount of daily virus production in chronically infected patients and the high spontaneous mutability of HCV virus (Neumann, et al., *Science* 1998, 282, 103-7; Fukimoto, et al., *Hepatology*, 1996, 24, 1351-4; Domingo, et al., *Gene*, 1985, 40, 1-8; Martell, et al., *J. Virol.* 1992, 66, 3225-9. Experimental anti-
10 viral nucleoside analogs have been shown to induce viable mutations in the HCV virus both *in vivo* and *in vitro* (Migliaccio, et al., *J. Biol. Chem.* 2003, 926; Carroll, et al., *Antimicrobial Agents Chemotherapy* 2009, 926; Brown, A. B., *Expert Opin. Investig. Drugs* 2009, 18, 709-725). Therefore, drugs having improved antiviral properties, particularly enhanced activity against resistant strains of virus; improved
15 oral bioavailability; fewer undesirable side effects and extended effective half-life *in vivo* (De Francesco, R. et al. (2003) *Antiviral Research* 58:1-16) are urgently needed.

Certain ribosides of the nucleobases pyrrolo[1,2-f][1,2,4]triazine, imidazo[1,5-f][1,2,4]triazine, imidazo[1,2-f][1,2,4]triazine, and [1,2,4]triazolo[4,3-f][1,2,4]triazine have been disclosed in *Carbohydrate Research* 2001, 331(1), 77-82; *Nucleosides &*
20 *Nucleotides* (1996), 15(1-3), 793-807; *Tetrahedron Letters* (1994), 35(30), 5339-42; *Heterocycles* (1992), 34(3), 569-74; *J. Chem. Soc. Perkin Trans. 1* 1985, 3, 621-30; *J. Chem. Soc. Perkin Trans. 1* 1984, 2, 229-38; WO 2000056734; *Organic Letters* (2001), 3(6), 839-842; *J. Chem. Soc. Perkin Trans. 1* 1999, 20, 2929-2936; and *J. Med. Chem.* 1986, 29(11), 2231-5. However, these compounds have not been
25 disclosed as useful for the treatment of HCV.

Ribosides of pyrrolo[1,2-f][1,2,4]triazinyl, imidazo[1,5-f][1,2,4]triazinyl, imidazo[1,2-f][1,2,4]triazinyl, and [1,2,4]triazolo[4,3-f][1,2,4]triazinyl nucleobases with antiviral, anti-HCV, and anti-RdRp activity have been disclosed by Babu, Y. S., WO2008/089105 and WO2008/141079; Cho, et al., WO2009/132123 and Francom,
30 et al. WO2010/002877. Butler, et al., WO2009/132135, has disclosed anti-viral pyrrolo[1,2-f][1,2,4]triazinyl, imidazo[1,5-f][1,2,4]triazinyl, imidazo[1,2-f][1,2,4]triazinyl, and [1,2,4]triazolo[4,3-f][1,2,4]triazinyl nucleosides wherein the 1' position of the nucleoside sugar is substituted.




5

SUMMARY OF THE INVENTION

Provided are compounds that inhibit viruses of the *Flaviviridae* family. The invention also comprises compounds of Formula I or Formula IV-VI that inhibit viral nucleic acid polymerases, particularly HCV RNA-dependent RNA polymerase (RdRp), rather than cellular nucleic acid polymerases. The compounds of Formula I or Formula IV-VI have been discovered to be efficacious against both wild type and S282T mutant strains of HCV virus. Therefore, a compound of Formula I or Formula IV-VI are useful for treating *Flaviviridae* infections in humans and other animals.

In one embodiment, provided are compounds of Formula I:



Formula I

or a pharmaceutically acceptable salt, thereof;

wherein:

R¹ is (C₁-C₈)alkyl, (C₄-C₈)carbocyclalkyl, (C₁-C₈)substituted alkyl, (C₂-C₈)alkenyl, (C₂-C₈)substituted alkenyl, (C₂-C₈)alkynyl, (C₂-C₈)substituted alkynyl, or aryl(C₁-C₈)alkyl;

R² is halogen;

each R³, R⁴, or R⁵ is independently H, OR^a, N(R^a)₂, N₃, CN, NO₂, S(O)_nR^a, halogen, (C₁-C₈)alkyl, (C₄-C₈)carbocyclalkyl, (C₁-C₈)substituted alkyl, (C₂-C₈)alkenyl, (C₂-C₈)substituted alkenyl, (C₂-C₈)alkynyl, (C₂-C₈)substituted alkynyl, or aryl(C₁-C₈)alkyl;

h

- 5 or any two of R^3 , R^4 or R^5 on adjacent carbon atoms when taken together are $-O(CO)O-$ or when taken together with the ring carbon atoms to which they are attached form a double bond;


R^6 is H, OR^a , $N(R^a)_2$, N_3 , CN, NO_2 , $S(O)_nR^a$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$,

- 10 $-SO_2NR^{11}R^{12}$, halogen, (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl (C_1-C_8) alkyl;

each n is independently 0, 1, or 2;


- 15 each R^a is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, or $-SO_2NR^{11}R^{12}$;

R^7 is H, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, $-SO_2NR^{11}R^{12}$, or



- 20 each Y or Y^1 is, independently, O, S, NR, $^+N(O)(R)$, $N(OR)$, $^+N(O)(OR)$, or $N-NR_2$;

W^1 and W^2 , when taken together, are $-Y^3(C(R^y)_2)_3Y^3-$; or one of W^1 or W^2 together with either R^3 or R^4 is $-Y^3-$ and the other of W^1 or W^2 is Formula Ia; or W^1 and W^2 are each, independently, a group of the Formula Ia:



a

5

Formula Ia

wherein:


each Y^2 is independently a bond, O, CR_2 , NR, $^+N(O)(R)$, $N(OR)$, $^+N(O)(OR)$, $N-NR_2$, S, S-S, S(O), or S(O)₂;

each Y^3 is independently O, S, or NR;

10

M2 is 0, 1 or 2;

each R^x is independently R^y or the formula:



wherein:

each M1a, M1c, and M1d is independently 0 or 1;

15

M12c is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

each R^y is independently H, F, Cl, Br, I, OH, R, $-C(=Y^1)R$, $-C(=Y^1)OR$, $-C(=Y^1)N(R)_2$, $-N(R)_2$, $^+N(R)_3$, $-SR$, $-S(O)R$, $-S(O)_2R$, $-S(O)(OR)$, $-S(O)_2(OR)$, $-OC(=Y^1)R$, $-OC(=Y^1)OR$, $-OC(=Y^1)N(R)_2$, $-SC(=Y^1)R$, $-SC(=Y^1)OR$, $-SC(=Y^1)N(R)_2$, $-N(R)C(=Y^1)R$, $-N(R)C(=Y^1)OR$, $-N(R)C(=Y^1)N(R)_2$, $-SO_2NR_2$, $-CN$, $-N_3$, $-NO_2$, $-OR$, or W^3 ; or when taken together, two R^y on the same carbon atom form a carbocyclic ring of 3 to 7 carbon atoms;

20

each R is independently H, (C₁-C₈) alkyl, (C₁-C₈) substituted alkyl, (C₂-C₈) alkenyl, (C₂-C₈) substituted alkenyl, (C₂-C₈) alkynyl, (C₂-C₈) substituted alkynyl, C₆-C₂₀ aryl, C₆-C₂₀ substituted aryl, C₂-C₂₀ heterocyclyl, C₂-C₂₀ substituted heterocyclyl, arylalkyl or substituted arylalkyl;

25

W^3 is W^4 or W^5 ; W^4 is R, $-C(Y^1)R^y$, $-C(Y^1)W^5$, $-SO_2R^y$, or $-SO_2W^5$; and W^5 is a carbocycle or a heterocycle wherein W^5 is independently substituted with 0 to 3 R^y groups;

each X^1 or X^2 is independently C-R¹⁰ or N;

1

5 each R^8 is halogen, $NR^{11}R^{12}$, $N(R^{11})OR^{11}$, $NR^{11}NR^{11}R^{12}$, N_3 , NO , NO_2 , CHO , CN , $-CH(=NR^{11})$, $-CH=NNHR^{11}$, $-CH=N(OR^{11})$, $-CH(OR^{11})_2$, $-C(=O)NR^{11}R^{12}$, $-C(=S)NR^{11}R^{12}$, $-C(=O)OR^{11}$, $(C_1-C_8)alkyl$, $(C_2-C_8)alkenyl$, $(C_2-C_8)alkynyl$, $(C_4-C_8)carbocyclylalkyl$, optionally substituted aryl, optionally substituted heteroaryl, $-C(=O)(C_1-C_8)alkyl$, $-S(O)_n(C_1-C_8)alkyl$, $aryl(C_1-C_8)alkyl$, OR^{11} or SR^{11} ;

10 each R^9 or R^{10} is independently H , halogen, $NR^{11}R^{12}$, $N(R^{11})OR^{11}$, $NR^{11}NR^{11}R^{12}$, N_3 , NO , NO_2 , CHO , CN , $-CH(=NR^{11})$, $-CH=NNHR^{11}$, $-CH=N(OR^{11})$, $-CH(OR^{11})_2$, $-C(=O)NR^{11}R^{12}$, $-C(=S)NR^{11}R^{12}$, $-C(=O)OR^{11}$, R^{11} , OR^{11} or SR^{11} ;

each R^{11} or R^{12} is independently H , $(C_1-C_8)alkyl$, $(C_2-C_8)alkenyl$, $(C_2-C_8)alkynyl$, $(C_4-C_8)carbocyclylalkyl$, optionally substituted aryl, optionally substituted heteroaryl, $-C(=O)(C_1-C_8)alkyl$, $-S(O)_n(C_1-C_8)alkyl$ or $aryl(C_1-C_8)alkyl$; or R^{11} and R^{12} taken together with a nitrogen to which they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with $-O-$, $-S-$ or $-NR^a-$;

20 wherein each $(C_1-C_8)alkyl$, $(C_2-C_8)alkenyl$, $(C_2-C_8)alkynyl$ or $aryl(C_1-C_8)alkyl$ of each R^1 , R^3 , R^4 , R^5 , R^6 , R^{11} or R^{12} is, independently, optionally substituted with one or more halo, hydroxy, CN , N_3 , $N(R^a)_2$ or OR^a ; and wherein one or more of the non-terminal carbon atoms of each said $(C_1-C_8)alkyl$ may be optionally replaced with $-O-$, $-S-$ or $-NR^a-$.

In another embodiment, provided are compounds of Formula I or Formula IV-
25 VI and pharmaceutically acceptable salts thereof and all racemates, enantiomers, diastereomers, tautomers, polymorphs, pseudopolymorphs and amorphous forms thereof.

In another embodiment, provided are novel compounds of Formula I or Formula IV-VI with activity against infectious *Flaviviridae* viruses. Without wishing
30 to be bound by theory, the compounds of the invention may inhibit viral RNA-dependent RNA polymerase and thus inhibit the replication of the virus. They are useful for treating human patients infected with a human virus such as hepatitis C.



5 In another embodiment, provided are pharmaceutical compositions comprising an effective amount of a Formula I or a Formula IV-VI compound, or a pharmaceutically acceptable salt thereof, in combination with a pharmaceutically acceptable diluent or carrier.

10 In another embodiment, the present application provides for combination pharmaceutical agent comprising:

- a) a first pharmaceutical composition comprising a compound of Formula I or Formula IV-VI; or a pharmaceutically acceptable salt, solvate, or ester thereof; and
- b) a second pharmaceutical composition comprising at least one
15 additional therapeutic agent selected from the group consisting of interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, other anti-fibrotic agents, endothelin antagonists, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-
20 nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers and other drugs for treating HCV; or mixtures thereof.

25 In another embodiment, the present application provides for a method of inhibiting HCV polymerase, comprising contacting a cell infected with HCV with an effective amount of a compound of Formula I or Formula IV-VI; or a pharmaceutically acceptable salts, solvate, and/or ester thereof.

30 In another embodiment, the present application provides for a method of inhibiting HCV polymerase, comprising contacting a cell infected with HCV with an effective amount of a compound of Formula I or Formula IV-VI; or a pharmaceutically acceptable salts, solvate, and/or ester thereof; and at least one additional therapeutic agent.

 In another embodiment, the present application provides for a method of treating and/or preventing a disease caused by a viral infection wherein the viral infection is caused by a virus selected from the group consisting of dengue virus,

2

- 5 yellow fever virus, West Nile virus, Japanese encephalitis virus, tick-borne encephalitis virus, Junjin virus, Murray Valley encephalitis virus, St Louis encephalitis virus, Omsk hemorrhagic fever virus, bovine viral diarrhea virus, Zika virus and Hepatitis C virus; by administering to a subject in need thereof a therapeutically effective amount of a compound of Formula I or Formula IV-VI, or a
10 pharmaceutically acceptable salt thereof.

In another embodiment, the present application provides for a method of treating HCV in a patient, comprising administering to said patient a therapeutically effective amount of a compound of Formula I or Formula IV-VI; or a pharmaceutically acceptable salt, solvate, and/or ester thereof.

- 15 In another embodiment, the present application provides for a method of treating HCV in a patient, comprising administering to said patient a therapeutically effective amount of a compound of Formula I or Formula IV-VI; or a pharmaceutically acceptable salt, solvate, and/or ester thereof; and at least one additional therapeutic agent.

- 20 Another aspect of the invention provides a method for the treatment or prevention of the symptoms or effects of an HCV infection in an infected animal which comprises administering to, i.e. treating, said animal with a pharmaceutical combination composition or formulation comprising an effective amount of a Formula I compound or Formula IV-VI, and a second compound having anti-HCV properties.

- 25 In another aspect, the invention also provides a method of inhibiting HCV, comprising administering to a mammal infected with HCV an amount of a Formula I or Formula IV-VI compound, effective to inhibit the replication of HCV in infected cells in said mammal.

- In another aspect, provided is the use of a compound of Formula I or Formula
30 IV-VI for the manufacture of a medicament for the treatment of *Flaviviridae* viral infections. In another aspect, provided is a compound of Formula I or Formula IV-VI for use in treating a *Flaviviridae* viral infection. In one embodiment, the *Flaviviridae* viral infection is acute or chronic HCV infection. In one embodiment of each aspect
- 2

5 of use and compound, the treatment results in the reduction of one or more of the viral loads or clearance of RNA in the patient.

In another aspect, the invention also provides processes and novel intermediates disclosed herein which are useful for preparing Formula I or Formula IV-VI compounds of the invention.


10 In other aspects, novel methods for synthesis, analysis, separation, isolation, purification, characterization, and testing of the compounds of this invention are provided.

15

DETAILED DESCRIPTION OF EXEMPLARY EMBODIMENTS

Reference will now be made in detail to certain embodiments of the invention, examples of which are illustrated in the accompanying description, structures and
20 formulas. While the invention will be described in conjunction with the enumerated embodiments, it will be understood that they are not intended to limit the invention to those embodiments. On the contrary, the invention is intended to cover all alternatives, modifications, and equivalents, which may be included within the scope of the present invention.

25 In another aspect, compounds of Formula I are represented by Formula II:



Handwritten mark resembling a stylized 'd' or 'a'.

5

Formula II

or a pharmaceutically acceptable salt, thereof;

wherein:


- R^1 is (C_1-C_8) alkyl, (C_4-C_8) carbocyclylalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl (C_1-C_8) alkyl;
- each R^3 , R^4 , or R^5 is independently H, OR^a , $N(R^a)_2$, N_3 , CN, NO_2 , $S(O)_nR^a$, halogen, (C_1-C_8) alkyl, (C_4-C_8) carbocyclylalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl (C_1-C_8) alkyl;
- or any two of R^3 , R^4 or R^5 on adjacent carbon atoms when taken together are $-O(CO)O-$ or when taken together with the ring carbon atoms to which they are attached form a double bond;

- R^6 is H, OR^a , $N(R^a)_2$, N_3 , CN, NO_2 , $S(O)_nR^a$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, $-SO_2NR^{11}R^{12}$, halogen, (C_1-C_8) alkyl, (C_4-C_8) carbocyclylalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl (C_1-C_8) alkyl;

each n is independently 0, 1, or 2;

- each R^a is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl (C_1-C_8) alkyl, (C_4-C_8) carbocyclylalkyl, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, or $-SO_2NR^{11}R^{12}$;


R^7 is H, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, $-SO_2NR^{11}R^{12}$, or



- each Y or Y^1 is, independently, O, S, NR, $^+N(O)(R)$, $N(OR)$, $^+N(O)(OR)$, or $N-NR_2$;


✓

- 5 W^1 and W^2 , when taken together, are $-Y^3(C(R^y)_2)_3Y^3-$; or one of W^1 or W^2 together with either R^3 or R^4 is $-Y^3-$ and the other of W^1 or W^2 is Formula Ia; or W^1 and W^2 are each, independently, a group of the Formula Ia:



Formula Ia

- 10 wherein:
 each Y^2 is independently a bond, O, CR_2 , NR, $^+N(O)(R)$, $N(OR)$, $^+N(O)(OR)$, $N-NR_2$, S, S-S, S(O), or S(O)₂;
 each Y^3 is independently O, S, or NR;
 M2 is 0, 1 or 2;
- 15 each R^x is independently R^y or the formula:



- wherein:
 each M1a, M1c, and M1d is independently 0 or 1;
 M12c is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;
- 20 each R^y is independently H, F, Cl, Br, I, OH, R, $-C(=Y^1)R$, $-C(=Y^1)OR$, $-C(=Y^1)N(R)_2$, $-N(R)_2$, $^+N(R)_3$, $-SR$, $-S(O)R$, $-S(O)_2R$, $-S(O)(OR)$, $-S(O)_2(OR)$, $-OC(=Y^1)R$, $-OC(=Y^1)OR$, $-OC(=Y^1)N(R)_2$, $-SC(=Y^1)R$, $-SC(=Y^1)OR$, $-SC(=Y^1)N(R)_2$, $-N(R)C(=Y^1)R$, $-N(R)C(=Y^1)OR$, $-N(R)C(=Y^1)N(R)_2$, $-SO_2NR_2$,

- 5 -CN, -N₃, -NO₂, -OR, or W³; or when taken together, two R^y on the same carbon atom form a carbocyclic ring of 3 to 7 carbon atoms;

each R is independently H, (C₁-C₈) alkyl, (C₁-C₈) substituted alkyl, (C₂-C₈) alkenyl, (C₂-C₈) substituted alkenyl, (C₂-C₈) alkynyl, (C₂-C₈) substituted alkynyl, C₆-C₂₀ aryl, C₆-C₂₀ substituted aryl, C₂-C₂₀ heterocyclyl, C₂-C₂₀ substituted

- 10 heterocyclyl, arylalkyl or substituted arylalkyl;

W³ is W⁴ or W⁵; W⁴ is R, -C(Y¹)R^y, -C(Y¹)W⁵, -SO₂R^y, or -SO₂W⁵; and W⁵ is a carbocycle or a heterocycle wherein W⁵ is independently substituted with 0 to 3 R^y groups;

each X¹ or X² is independently C-R¹⁰ or N;

- 15 each R⁸ is halogen, NR¹¹R¹², N(R¹¹)OR¹¹, NR¹¹NR¹¹R¹², N₃, NO, NO₂, CHO, CN, -CH(=NR¹¹), -CH=NNHR¹¹, -CH=N(OR¹¹), -CH(OR¹¹)₂, -C(=O)NR¹¹R¹², -C(=S)NR¹¹R¹², -C(=O)OR¹¹, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₄-C₈)carbocyclalkyl, optionally substituted aryl, optionally substituted heteroaryl, -C(=O)(C₁-C₈)alkyl, -S(O)_n(C₁-C₈)alkyl, aryl(C₁-C₈)alkyl, OR¹¹ or SR¹¹;

- 20 each R⁹ or R¹⁰ is independently H, halogen, NR¹¹R¹², N(R¹¹)OR¹¹, NR¹¹NR¹¹R¹², N₃, NO, NO₂, CHO, CN, -CH(=NR¹¹), -CH=NNHR¹¹, -CH=N(OR¹¹), -CH(OR¹¹)₂, -C(=O)NR¹¹R¹², -C(=S)NR¹¹R¹², -C(=O)OR¹¹, R¹¹, OR¹¹ or SR¹¹;

each R¹¹ or R¹² is independently H, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₄-C₈)carbocyclalkyl, optionally substituted aryl, optionally

- 25 substituted heteroaryl, -C(=O)(C₁-C₈)alkyl, -S(O)_n(C₁-C₈)alkyl or aryl(C₁-C₈)alkyl; or R¹¹ and R¹² taken together with a nitrogen to which they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with -O-, -S- or -NR^a;

- 30 wherein each (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl or aryl(C₁-C₈)alkyl of each R¹, R³, R⁴, R⁵, R⁶, R¹¹ or R¹² is, independently, optionally substituted with one or more halo, hydroxy, CN, N₃, N(R^a)₂ or OR^a; and wherein one or more of the non-terminal carbon atoms of each said (C₁-C₈)alkyl may be optionally replaced with -O-, -S- or -NR^a.

L

5 In one embodiment of the invention of Formula II, R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment, R^1 is methyl. In another aspect of this embodiment, R^1 is (C_1-C_8) alkyl and R^6 is H. In another aspect of this embodiment, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 .

In one embodiment of Formula II, R^3 is H, OR^a , $N(R^a)_2$, N_3 , CN, SR^a , halogen, (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In one aspect of this embodiment, R^3 is H. In another aspect of this embodiment, R^3 is H and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^3 is H and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^3 is H and R^1 methyl, CH_2F , or ethynyl. In another aspect of this embodiment, R^3 is H and R^1 is methyl. In another aspect of this embodiment, R^3 is H, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^3 is H, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^3 is H, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^3 is H, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^3 is H, R^1 is methyl and R^6 is H.

In one embodiment of Formula II, R^4 is H, OR^a , $N(R^a)_2$, N_3 , CN, SR^a , halogen, (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this

α

- embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.
- 15 In one embodiment of Formula II, R^5 is H, OR^a , $N(R^a)_2$, N_3 , CN, SR^a , halogen, (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6



5 is CN, OH, or CH₃. In another aspect of this embodiment, R⁴ is OH, R¹ is methyl and R⁶ is H. In another aspect of this embodiment, R⁵ is N₃.

In another embodiment of Formula II, R⁵ is H. In another aspect of this embodiment, R⁴ is H or OR^a. In another aspect of this embodiment, R⁴ is OR^a. In another aspect of this embodiment, R⁴ is OR^a and R¹ is (C₁-C₈)alkyl, (C₂-C₈) alkenyl or (C₂-C₈)alkynyl. In another aspect of this embodiment, R⁴ is OR^a and R¹ is (C₁-C₈)alkyl, (C₂-C₈) alkenyl or (C₂-C₈)alkynyl. In another aspect of this embodiment, R⁴ is OR^a and R¹ is (C₁-C₈)alkyl. In another aspect of this embodiment, R⁴ is OR^a and R¹ is methyl. In another aspect of this embodiment, R⁴ is OR^a, R¹ is (C₁-C₈)alkyl and at least one of X¹ or X² is N. In another aspect of this embodiment, R⁴ is OR^a, R¹ is methyl and at least one of X¹ or X² is N. In another aspect of this embodiment, R⁴ is OR^a, R¹ is (C₁-C₈)alkyl and R⁶ is CN, OH, or CH₃. In another aspect of this embodiment, R⁴ is OR^a, R¹ is methyl and R⁶ is CN, OH, or CH₃. In another aspect of this embodiment, R⁴ is OR^a, R¹ is methyl and R⁶ is H. In another aspect of this embodiment, R⁴ is OH and R¹ is methyl. In another aspect of this embodiment, R⁴ is OH, R¹ is (C₁-C₈)alkyl and at least one of X¹ or X² is N. In another aspect of this embodiment, R⁴ is OH, R¹ is methyl and at least one of X¹ or X² is N. In another aspect of this embodiment, R⁴ is OH, R¹ is (C₁-C₈)alkyl and R⁶ is CN, OH, or CH₃. In another aspect of this embodiment, R⁴ is OH, R¹ is methyl and R⁶ is CN, OH, or CH₃. In another aspect of this embodiment, R⁴ is OH, R¹ is methyl and R⁶ is H.

In another embodiment of Formula II, R⁶ is H, CN, OR^a or CH₃. In another aspect of this embodiment R⁶ is H. In another aspect of this embodiment R⁶ is CN. In another aspect of this embodiment R⁶ is OR^a. In another aspect of this embodiment R⁶ is OH. In another aspect of this embodiment R⁶ is CH₃. In another aspect of this embodiment, R⁴ is H or OR^a. In another aspect of this embodiment, R⁴ is OR^a. In another aspect of this embodiment, R⁴ is OR^a and R¹ is (C₁-C₈)alkyl, (C₂-C₈) alkenyl or (C₂-C₈)alkynyl. In another aspect of this embodiment, R⁴ is OR^a and R¹ is (C₁-C₈)alkyl, (C₂-C₈) alkenyl or (C₂-C₈)alkynyl. In another aspect of this



embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.


In another embodiment of Formula II, R^6 is CN, OR^a or CH_3 . In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of




- 5 X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 .

In one embodiment of Formula II, R^7 is H, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, -

- 10 $C(=O)SR^{11}$ or



. In a aspect of this embodiment, R^7 is H. In another aspect of this embodiment, R^7 is $-C(=O)R^{11}$. In another aspect of this embodiment, R^7 is $-C(=O)R^{11}$ wherein R^{11} is (C_1-C_8) alkyl. In another aspect of this embodiment, R^7 is



- . In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In

L

- 5 another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.
- 10 In one embodiment of Formula II, X^1 is N or $C-R^{10}$. In another aspect of this embodiment, X^1 is N. In another aspect of this embodiment, X^1 is $C-R^{10}$. In another aspect of this embodiment, X^2 is C-H. In another aspect of this embodiment, X^1 is N and X^2 is C-H. In another aspect of this embodiment, X^1 is $C-R^{10}$ and X^2 is CH. In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is
- 15 CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a
- 20 and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this
- 25 embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 .
- 30 In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

In another embodiment of Formula II, each R^8 is independently halogen, $NR^{11}R^{12}$, $N(R^{11})OR^{11}$, $NR^{11}NR^{11}R^{12}$, OR^{11} or SR^{11} . In another aspect of this embodiment, R^1 is methyl, CH_2F or ethynyl. In another aspect of this embodiment,

- 5 R^1 is methyl. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$ and R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$ and R^1 is methyl. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen and
- 10 R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen and R^1 is methyl. In another aspect of this embodiment, R^8 and R^9 are each NH_2 . In another aspect of this embodiment, R^8 and R^9 are each NH_2 and R^1 is methyl. In another aspect of this embodiment, R^8 and R^9 are each NH_2 and R^1 is methyl, CH_2F or ethynyl. In another aspect of this embodiment, R^8 is OH and R^9 is
- 15 NH_2 . In another aspect of this embodiment, R^8 is OH, R^9 is NH_2 and R^1 is methyl. In another aspect of this embodiment, R^8 is OH, R^9 is NH_2 and R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 .
- 20 In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is
- 25 (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another
- 30 aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6


5 is CN, OH, or CH₃. In another aspect of this embodiment, R⁴ is OH, R¹ is methyl and R⁶ is H.

In another embodiment of Formula II, each R¹⁰ is, independently, H, halogen, CN or optionally substituted heteroaryl. In another aspect of this embodiment, R¹ is methyl. In another aspect of this embodiment, R¹ is methyl, CH₂F or ethynyl. In
 10 another aspect of this embodiment, R⁹ is H, halogen, or NR¹¹R¹². In another aspect of this embodiment, R⁹ is H, halogen, or NR¹¹R¹² and R¹ is methyl. In another aspect of this embodiment, R⁹ is H, halogen, or NR¹¹R¹² and R¹ is methyl, CH₂F, or ethynyl. In another aspect of this embodiment, R⁸ is NH₂ and R⁹ is H or halogen. In another aspect of this embodiment, R⁸ is NH₂ and R⁹ is H or halogen and R¹ is methyl. In
 15 another aspect of this embodiment, R⁸ is NH₂ and R⁹ is H or halogen and R¹ is methyl, CH₂F, or ethynyl. In another aspect of this embodiment, R⁸ and R⁹ are each NH₂. In another aspect of this embodiment, R⁸ and R⁹ are each NH₂ and R¹ is methyl. In another aspect of this embodiment, R⁸ and R⁹ are each NH₂ and R¹ is methyl, CH₂F or ethynyl. In another aspect of this embodiment, R⁸ is OH and R⁹ is NH₂. In another
 20 aspect of this embodiment, R⁸ is OH, R⁹ is NH₂ and R¹ is methyl. In another aspect of this embodiment, R⁸ is OH, R⁹ is NH₂ and R¹ is methyl, CH₂F, or ethynyl. In another aspect of this embodiment R⁶ is H. In another aspect of this embodiment R⁶ is CN. In another aspect of this embodiment R⁶ is OR^a. In another aspect of this embodiment R⁶ is OH. In another aspect of this embodiment R⁶ is CH₃. In another aspect of this
 25 embodiment, R⁴ is H or OR^a. In another aspect of this embodiment, R⁴ is OR^a. In another aspect of this embodiment, R⁴ is OR^a and R¹ is (C₁–C₈)alkyl, (C₂–C₈) alkenyl or (C₂–C₈)alkynyl. In another aspect of this embodiment, R⁴ is OR^a and R¹ is (C₁–C₈)alkyl. In another aspect of this embodiment, R⁴ is OR^a and R¹ is methyl. In another aspect of this embodiment, R⁴ is OR^a, R¹ is (C₁–C₈)alkyl and at least one of
 30 X¹ or X² is N. In another aspect of this embodiment, R⁴ is OR^a, R¹ is methyl and at least one of X¹ or X² is N. In another aspect of this embodiment, R⁴ is OR^a, R¹ is (C₁–C₈)alkyl and R⁶ is CN, OH, or CH₃. In another aspect of this embodiment, R⁴ is OR^a, R¹ is methyl and R⁶ is CN, OH, or CH₃. In another aspect of this embodiment, R⁴ is OR^a, R¹ is methyl and R⁶ is H. In another aspect of this embodiment, R⁴ is OH

✓

- 5 and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this
- 10 this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

In another embodiment, compounds of Formula I or Formula II are represented by Formula III:



Formula III

- 15 or a pharmaceutically acceptable salt, thereof;
wherein:

R^1 is CH_3 , CH_2F , or ethynyl and all remaining variables are defined as for Formula I.

- In one embodiment of Formula III, R^4 is H, OR^a , $N(R^a)_2$, N_3 , CN, SR^a , halogen, (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this
- 20 embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is CH_3 , CH_2F , or ethynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another
- 25 aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this

Handwritten signature or mark.


5 embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.


In another embodiment of Formula III, R^6 is H, CN, OR^a or CH_3 . In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In
 10 another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another
 15 aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In
 20 another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

In another embodiment of Formula III, R^6 is CN, OR^a or CH_3 . In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of
 25 this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one
 30 of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 .

In one embodiment of Formula III, R^7 is H, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)SR^{11}$ or



- 5  . In a aspect of this embodiment, R^7 is H. In another aspect of this embodiment, R^7 is $-C(=O)R^{11}$. In another aspect of this embodiment, R^7 is $-C(=O)R^{11}$ wherein R^{11} is (C_1-C_8) alkyl. In another aspect of this embodiment, R^7 is

-  . In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In
10 another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of
15 this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

- 20 In one embodiment of Formula III, X^1 is N or $C-R^{10}$. In another aspect of this embodiment, X^1 is N. In another aspect of this embodiment, X^1 is $C-R^{10}$. In another aspect of this embodiment, X^2 is C-H. In another aspect of this embodiment, X^1 is N and X^2 is C-H. In another aspect of this embodiment, X^1 is $C-R^{10}$ and X^2 is CH. In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is
25 CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N.



- 5 In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.
- 10 In another embodiment of Formula III, each R^8 is independently halogen, $NR^{11}R^{12}$, $N(R^{11})OR^{11}$, $NR^{11}NR^{11}R^{12}$, OR^{11} or SR^{11} . In another aspect of this embodiment, R^1 is methyl, CH_2F or ethynyl. In another aspect of this embodiment, R^1 is methyl. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$ and R^1 is methyl,
- 15 CH_2F , or ethynyl. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$ and R^1 is methyl. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen and R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen and R^1 is methyl. In another aspect of this embodiment, R^8 and R^9 are each NH_2 . In another aspect of this embodiment, R^8 and R^9 are each NH_2 and R^1 is methyl, CH_2F or ethynyl. In another aspect of this embodiment, R^8 and R^9 are each NH_2 and R^1 is methyl. In another aspect of this embodiment, R^8 is OH and R^9 is NH_2 . In another aspect of this embodiment, R^8 is OH, R^9 is NH_2 and R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment, R^8 is OH, R^9 is NH_2 and R^1 is methyl.
- 25 In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is
- 30




- 5 methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

In another embodiment of Formula III, each R^{10} is, independently, H, halogen, CN or optionally substituted heteroaryl. In another aspect of this embodiment, R^1 is methyl. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$. In

- 10 another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$ and R^1 is methyl. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen and R^1 is methyl. In another aspect of this embodiment, R^8 and R^9 are each NH_2 . In another aspect of this embodiment, R^8 and R^9 are each NH_2 and R^1 is methyl. In another aspect of this
- 15 embodiment, R^8 is OH and R^9 is NH_2 . In another aspect of this embodiment, R^8 is OH, R^9 is NH_2 and R^1 is methyl. In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In
- 20 another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this
- 25 embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

In another embodiment, provided are compounds of Formula IV:



Formula IV


or a pharmaceutically acceptable salt, thereof;

wherein:

- R^1 is (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl,
 10 (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted
 alkynyl, or aryl (C_1-C_8) alkyl;
 R^2 is halogen;
 R^3 , R^4 , and R^5 are each independently H, halogen, OR^a , $N(R^a)_2$, N_3 , CN, NO_2 ,
 $S(O)_nR^a$, (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl,
 15 (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted
 alkynyl, or aryl (C_1-C_8) alkyl;
 or any two of R^3 , R^4 or R^5 on adjacent carbon atoms when taken together are –
 $O(CO)O$ - or when taken together with the ring carbon atoms to which they are
 attached form a double bond;
 20 each n is independently 0, 1, or 2;
 each R^a is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl,
 aryl (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$,
 $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, or $-SO_2NR^{11}R^{12}$;
 R^7 is H, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$,
 25 $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, $-SO_2NR^{11}R^{12}$, or

L

28



5


Y is O, S, NR, $^+N(O)(R)$, N(OR), $^+N(O)(OR)$, or N-NR₂;

W¹ and W², when taken together, are $-Y^3(C(R^y)_2)_3Y^3-$; or

one of W¹ or W² together with either R³ or R⁴ is $-Y^3-$ and the other of W¹ or

10 W² is Formula Ia; or

W¹ and W² are each, independently, a group of Formula IVa:



Formula IVa

wherein:

15 each Y¹ is, independently, O, S, NR, $^+N(O)(R)$, N(OR), $^+N(O)(OR)$, or N-NR₂;


each Y² is independently a bond, O, CR₂, NR, $^+N(O)(R)$, N(OR), $^+N(O)(OR)$, N-NR₂, S, S-S, S(O), or S(O)₂;

each Y³ is independently O, S, or NR;

20 M2 is 0, 1 or 2;

each R^x is a group of Formula IVb:

✂



Formula IVb

wherein:

each M1a, M1c, and M1d is independently 0 or 1;

10 M12c is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

each R^y is independently H, F, Cl, Br, I, OH, -C(=Y¹)R, -C(=Y¹)R¹³, -C(=Y¹)OR, -C(=Y¹)N(R)₂, -N(R)₂, -⁺N(R)₃, -SR, -S(O)R, -S(O)₂R, -S(O)₂R¹³, -S(O)(OR), -S(O)₂(OR), -OC(=Y¹)R, -OC(=Y¹)OR, -OC(=Y¹)(N(R)₂), -SC(=Y¹)R, -SC(=Y¹)OR, -SC(=Y¹)(N(R)₂), -N(R)C(=Y¹)R, -N(R)C(=Y¹)OR, -N(R)C(=Y¹)N(R)₂,
 15 -SO₂NR₂, -CN, -N₃, -NO₂, -OR, (C₁-C₈) alkyl, (C₂-C₈) alkenyl, (C₂-C₈) alkynyl, C₆-C₂₀ aryl, C₃-C₂₀ carbocyclyl, C₂-C₂₀ heterocyclyl, arylalkyl, heteroarylalkyl;

wherein each (C₁-C₈) alkyl, (C₂-C₈) alkenyl, (C₂-C₈) alkynyl, C₆-C₂₀ aryl,

C₃-C₂₀ carbocyclyl, C₂-C₂₀ heterocyclyl, arylalkyl, or heteroarylalkyl

is optionally substituted with 1-3 R²⁰ groups;

20 or when taken together, two R^y on the same carbon atom form a carbocyclic ring of 3 to 7 carbon atoms;

each R is independently H, (C₁-C₈) alkyl, (C₂-C₈) alkenyl, (C₂-C₈) alkynyl,

C₆-C₂₀ aryl, C₃-C₂₀ carbocyclyl, C₂-C₂₀ heterocyclyl, or arylalkyl;

each R⁸ is halogen, NR¹¹R¹², N(R¹¹)OR¹¹, NR¹¹NR¹¹R¹², N₃, NO, NO₂, OR¹¹
 25 or S(O)_nR¹¹;

each R⁹ is independently H, halogen, NR¹¹R¹², N(R¹¹)OR¹¹, NR¹¹NR¹¹R¹², N₃, NO, NO₂, CHO, CN, -CH(=NR¹¹), -CH=NHNR¹¹, -CH=N(OR¹¹), -CH(OR¹¹)₂, -C(=O)NR¹¹R¹², -C(=S)NR¹¹R¹², -C(=O)OR¹¹, R¹¹, OR¹¹ or S(O)_nR¹¹;

each R¹¹ or R¹² is independently H, (C₁-C₈) alkyl, (C₂-C₈) alkenyl, (C₂-C₈) alkynyl, (C₄-C₈) carbocyclalkyl, optionally substituted aryl, optionally
 30

5 substituted heteroaryl, $-C(=O)(C_1-C_8)alkyl$, $-S(O)_n(C_1-C_8)alkyl$ or $aryl(C_1-C_8)alkyl$; or R^{11} and R^{12} taken together with a nitrogen to which they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with $-O-$, $-S-$ or $-NR^b$;

10 each R^{13} is independently a carbocycle or heterocycle optionally substituted with 1-3 R^{20} groups;

each R^{20} is independently, halogen, CN , N_3 , $N(R)_2$, OR , $-SR$, $-S(O)R$, $-S(O)_2R$, $-S(O)(OR)$, $-S(O)_2(OR)$, $-C(=Y^1)R$, $-C(=Y^1)OR$, or $C(=Y^1)N(R)_2$;

15 wherein each $(C_1-C_8)alkyl$, $(C_2-C_8)alkenyl$, $(C_2-C_8)alkynyl$ or $aryl(C_1-C_8)alkyl$ of each R^1 , R^3 , R^4 , R^5 , R^6 , R^{11} or R^{12} is, independently, optionally substituted with one or more halo, hydroxy, CN , N_3 , $N(R^b)_2$ or OR^b ; and wherein one or more of the non-terminal carbon atoms of each said $(C_1-C_8)alkyl$ may be optionally replaced with $-O-$, $-S-$ or $-NR^b$;

20 each R^b is independently H , $(C_1-C_8)alkyl$, $(C_2-C_8)alkenyl$, $(C_2-C_8)alkynyl$, $aryl(C_1-C_8)alkyl$, $(C_4-C_8)carbocyclalkyl$, $-C(=O)R^{21}$, $-C(=O)OR^{21}$, $-C(=O)NR^{21}R^{22}$, $-C(=O)SR^{21}$, $-S(O)R^{21}$, $-S(O)_2R^{21}$, $-S(O)(OR^{21})$, $-S(O)_2(OR^{21})$, or $-SO_2NR^{21}R^{22}$;


each R^{21} or R^{22} is independently H , $(C_1-C_8)alkyl$, $(C_2-C_8)alkenyl$, $(C_2-C_8)alkynyl$, $(C_4-C_8)carbocyclalkyl$, $-C(=O)(C_1-C_8)alkyl$, $-S(O)_n(C_1-C_8)alkyl$ or $aryl(C_1-C_8)alkyl$;

25 with the optional proviso that compounds 1, 1d, 1e, 2, TP-1, A-1, 8, and 21 are excluded.

In another aspect of this embodiment Y and Y^1 is O . In another aspect of this embodiment R^8 is halogen, $NR^{11}R^{12}$, $N(R^{11})OR^{11}$, $NR^{11}NR^{11}R^{12}$, OR^{11} or $S(O)_nR^{11}$. In another aspect of this embodiment R^9 is H , halogen, $S(O)_nR^{11}$ or $NR^{11}R^{12}$. In another aspect of this embodiment R^4 is OR^a . In another aspect of this embodiment R^1 is CH_3 . In another aspect of this embodiment R^2 is F . In another aspect of this embodiment R^7 is




31



5

wherein Y is -O-; W¹ is Formula Ia and W² together with R⁴ is -O-. In another embodiment, compounds of Formula IV are represented by Formula V:




10

Formula V

wherein R¹ is methyl or ethynyl, and R⁴ is OR^a. In another aspect of this embodiment R⁷ is H or

15







In another aspect of this embodiment, compound of Formula V are represented the following structures:


20

32

5




10



In another embodiment, provided are compounds of Formula VI:

33




Formula VI

or a pharmaceutically acceptable salt, thereof;

wherein:


- 10 R^4 is OR^a ;
- each n is independently 0, 1, or 2;
- each R^a is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, or $-SO_2NR^{11}R^{12}$;
- 15 R^7 is H, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, $-SO_2NR^{11}R^{12}$, or



- 20 Y is O, S, NR, $^+N(O)(R)$, $N(OR)$, $^+N(O)(OR)$, or $N-NR_2$;
- W^1 and W^2 , when taken together, are $-Y^3(C(R^y)_2)_3Y^3-$; or
- one of W^1 or W^2 together with R^4 is $-Y^3-$ and the other of W^1 or W^2 is Formula Ia; or
- W^1 and W^2 are each, independently, a group of Formula VIa:

Handwritten signature or mark.

34



Formula VIa

wherein:


each Y^1 is, independently, O, S, NR, $^+\text{N}(\text{O})(\text{R})$, N(OR), $^+\text{N}(\text{O})(\text{OR})$, or N-NR₂;

each Y^2 is independently a bond, O, CR₂, NR, $^+\text{N}(\text{O})(\text{R})$, N(OR), $^+\text{N}(\text{O})(\text{OR})$, N-NR₂, S, S-S, S(O), or S(O)₂;

each Y^3 is independently O, S, or NR;

M2 is 0, 1 or 2;

each R^x is a group of Formula VIb:



Formula VIb

wherein:

each M1a, M1c, and M1d is independently 0 or 1;

M12c is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

each R^y is independently H, F, Cl, Br, I, OH, $-\text{C}(=\text{Y}^1)\text{R}$, $-\text{C}(=\text{Y}^1)\text{R}^{13}$, $-\text{C}(=\text{Y}^1)\text{OR}$, $-\text{C}(=\text{Y}^1)\text{N}(\text{R})_2$, $-\text{N}(\text{R})_2$, $^+\text{N}(\text{R})_3$, $-\text{SR}$, $-\text{S}(\text{O})\text{R}$, $-\text{S}(\text{O})_2\text{R}$, $-\text{S}(\text{O})_2\text{R}^{13}$, $-\text{S}(\text{O})(\text{OR})$, $-\text{S}(\text{O})_2(\text{OR})$, $-\text{OC}(=\text{Y}^1)\text{R}$, $-\text{OC}(=\text{Y}^1)\text{OR}$, $-\text{OC}(=\text{Y}^1)(\text{N}(\text{R})_2)$, $-\text{SC}(=\text{Y}^1)\text{R}$, $-\text{SC}(=\text{Y}^1)\text{OR}$, $-\text{SC}(=\text{Y}^1)(\text{N}(\text{R})_2)$, $-\text{N}(\text{R})\text{C}(=\text{Y}^1)\text{R}$, $-\text{N}(\text{R})\text{C}(=\text{Y}^1)\text{OR}$, $-\text{N}(\text{R})\text{C}(=\text{Y}^1)\text{N}(\text{R})_2$,


L

- 5 $-\text{SO}_2\text{NR}_2$, $-\text{CN}$, $-\text{N}_3$, $-\text{NO}_2$, $-\text{OR}$, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, C_6-C_{20} aryl, C_3-C_{20} carbocyclyl, C_2-C_{20} heterocyclyl, arylalkyl, heteroarylalkyl; wherein each (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, C_6-C_{20} aryl, C_3-C_{20} carbocyclyl, C_2-C_{20} heterocyclyl, arylalkyl, or heteroarylalkyl is optionally substituted with 1-3 R^{20} groups;
- 10 each R is independently H , (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, C_6-C_{20} aryl, C_3-C_{20} carbocyclyl, C_2-C_{20} heterocyclyl, or arylalkyl; each R^{11} or R^{12} is independently H , (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, (C_4-C_8) carbocyclylalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-\text{C}(=\text{O})(\text{C}_1-\text{C}_8)$ alkyl, $-\text{S}(\text{O})_n(\text{C}_1-\text{C}_8)$ alkyl or aryl (C_1-C_8) alkyl;
- 15 each R^{13} is independently a carbocycle or heterocycle optionally substituted with 1-3 R^{20} groups; each R^{20} is independently, halogen, CN , N_3 , $\text{N}(\text{R})_2$, OR , $-\text{SR}$, $-\text{S}(\text{O})\text{R}$, $-\text{S}(\text{O})_2\text{R}$, $-\text{S}(\text{O})(\text{OR})$, $-\text{S}(\text{O})_2(\text{OR})$, $-\text{C}(=\text{Y}^1)\text{R}$, $-\text{C}(=\text{Y}^1)\text{OR}$, or $\text{C}(=\text{Y}^1)\text{N}(\text{R})_2$; wherein each (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl or aryl (C_1-C_8) alkyl
- 20 of each R^4 , R^{11} or R^{12} is, independently, optionally substituted with one or more halo, hydroxy, CN , N_3 , $\text{N}(\text{R}^b)_2$ or OR^b ; and wherein one or more of the non-terminal carbon atoms of each said (C_1-C_8) alkyl may be optionally replaced with $-\text{O}-$, $-\text{S}-$ or $-\text{NR}^b$; each R^b is independently H , (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl (C_1-C_8) alkyl, (C_4-C_8) carbocyclylalkyl, $-\text{C}(=\text{O})\text{R}^{21}$, $-\text{C}(=\text{O})\text{OR}^{21}$, $-\text{C}(=\text{O})\text{NR}^{21}\text{R}^{22}$,
- 25 $-\text{C}(=\text{O})\text{SR}^{21}$, $-\text{S}(\text{O})\text{R}^{21}$, $-\text{S}(\text{O})_2\text{R}^{21}$, $-\text{S}(\text{O})(\text{OR}^{21})$, $-\text{S}(\text{O})_2(\text{OR}^{21})$, or $-\text{SO}_2\text{NR}^{21}\text{R}^{22}$; each R^{21} or R^{22} is independently H , (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, (C_4-C_8) carbocyclylalkyl, $-\text{C}(=\text{O})(\text{C}_1-\text{C}_8)$ alkyl, $-\text{S}(\text{O})_n(\text{C}_1-\text{C}_8)$ alkyl or aryl (C_1-C_8) alkyl; and
- with the optional proviso that compounds 1, 1c, 1d, 1e, 2, TP-1, A-1, 8, and 21
- 30 are excluded.
- In another aspect of this embodiment R^a is H , (C_1-C_8) alkyl, or $-\text{C}(=\text{O})(\text{C}_1-\text{C}_6)$ alkyl; R^7 or R^7 together with R^4 is



36

5



wherein

10

a is the point of attachment to R^7 ;

b is the point of attachment to R^4 ;

Ar is phenyl or naphthyl, wherein the phenyl and naphthyl are optionally substituted with 1-3 R^{20} groups;


15

each R^y is independently (C_1-C_8) alkyl or C_5-C_6 carbocyclyl, wherein the alkyl and carbocyclyl are optionally substituted with 1-3 R^{20} groups;

each R is independently H, (C_1-C_6) alkyl, or arylalkyl; and

each R^{20} is independently halogen, CN, $N(R)_2$, OR, -SR, -S(O)R, -S(O)₂R, -S(O)(OR), -S(O)₂(OR), -C(=O)R, -C(=O)OR, or C(=O) $N(R)_2$.


In another embodiment, compounds of Formula IV-VI are represented by compounds having a structure:




[Handwritten signature]

37

5













10




X


38



5




, or



;

or a pharmaceutically acceptable salt, thereof.

- 10 In one embodiment of Formulas I-III and Formulas IV-VI, R^{11} or R^{12} is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, (C_4-C_8) carbocyclalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-C(=O)(C_1-C_8)$ alkyl, $-S(O)_n(C_1-C_8)$ alkyl or aryl (C_1-C_8) alkyl. In another embodiment, R^{11} and R^{12} taken together with a nitrogen to which they are both attached, form a 3 to
- 15 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with -O-, -S- or $-NR^a$ -. Therefore, by way of example and not limitation, the moiety $-NR^{11}R^{12}$ can be represented by the heterocycles:




and the like.

- 20 In another embodiment of Formulas I-III and Formulas IV-VI, each R^3 , R^4 , R^5 , R^6 , R^{11} or R^{12} is, independently, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl or aryl (C_1-C_8) alkyl, wherein said (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl or aryl (C_1-C_8) alkyl are, independently, optionally substituted with one or more halo, hydroxy, CN, N_3 , $N(R^a)_2$ or OR^a . Therefore, by way of example and not limitation,
- 25 R^3 , R^4 , R^5 , R^6 , R^{11} or R^{12} could represent moieties such as $-CH(NH_2)CH_3$, -

- 5 $\text{CH(OH)CH}_2\text{CH}_3$, $-\text{CH(NH}_2\text{)CH(CH}_3\text{)}_2$, $-\text{CH}_2\text{CF}_3$, $-(\text{CH}_2)_2\text{CH(N}_3\text{)CH}_3$, $-(\text{CH}_2)_6\text{NH}_2$
and the like.

In another embodiment of Formula I-III and Formula IV-VI, R^3 , R^4 , R^5 , R^6 , R^{11} or R^{12} is $(\text{C}_1\text{-C}_8)$ alkyl wherein one or more of the non-terminal carbon atoms of each said $(\text{C}_1\text{-C}_8)$ alkyl may be optionally replaced with $-\text{O}-$, $-\text{S}-$ or $-\text{NR}^a$ -. Therefore,
10 by way of example and not limitation, R^3 , R^4 , R^5 , R^6 , R^{11} or R^{12} could represent moieties such as $-\text{CH}_2\text{OCH}_3$, $-\text{CH}_2\text{OCH}_2\text{CH}_3$, $-\text{CH}_2\text{OCH(CH}_3\text{)}_2$, $-\text{CH}_2\text{SCH}_3$, $-(\text{CH}_2)_6\text{OCH}_3$, $-(\text{CH}_2)_6\text{N(CH}_3\text{)}_2$ and the like.

In another embodiment, Formulas I-III is a compound selected from the group consisting of



9

3

2

1

2

2

2

1

2


2

3




42


5








5



44



5




10



46

5





and

or a pharmaceutically acceptable salt or ester thereof.

In another embodiment, provided is a compound useful for the synthesis of the compounds of Formula I selected from the group consisting of

10





DEFINITIONS

10 Unless stated otherwise, the following terms and phrases as used herein are intended to have the following meanings:

When trade names are used herein, applicants intend to independently include the trade name product and the active pharmaceutical ingredient(s) of the trade name product.

15 As used herein, "a compound of the invention" or "a compound of Formula I" means a compound of Formula I or a pharmaceutically acceptable salt, thereof. Similarly, with respect to isolatable intermediates, the phrase "a compound of Formula (number)" means a compound of that formula and pharmaceutically acceptable salts, thereof.

20 "Alkyl" is hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms. For example, an alkyl group can have 1 to 20 carbon atoms (*i.e.*, C₁-C₂₀ alkyl), 1 to 8 carbon atoms (*i.e.*, C₁-C₈ alkyl), or 1 to 6 carbon atoms (*i.e.*, C₁-C₆ alkyl). Examples of suitable alkyl groups include, but are not limited to, methyl (Me, -CH₃), ethyl (Et, -CH₂CH₃), 1-propyl (*n*-Pr, *n*-propyl, -CH₂CH₂CH₃), 2-propyl (*i*-Pr,

- 5 *i*-propyl, $-\text{CH}(\text{CH}_3)_2$, 1-butyl (*n*-Bu, *n*-butyl, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2-methyl-1-propyl (*i*-Bu, *i*-butyl, $-\text{CH}_2\text{CH}(\text{CH}_3)_2$), 2-butyl (*s*-Bu, *s*-butyl, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$), 2-methyl-2-propyl (*t*-Bu, *t*-butyl, $-\text{C}(\text{CH}_3)_3$), 1-pentyl (*n*-pentyl, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2-pentyl ($-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$), 3-pentyl ($-\text{CH}(\text{CH}_2\text{CH}_3)_2$), 2-methyl-2-butyl ($-\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_3$), 3-methyl-2-butyl ($-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)_2$), 3-methyl-1-butyl ($-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$), 2-methyl-1-butyl ($-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$), 1-hexyl ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2-hexyl ($-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 3-hexyl ($-\text{CH}(\text{CH}_2\text{CH}_3)(\text{CH}_2\text{CH}_2\text{CH}_3)$), 2-methyl-2-pentyl ($-\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_2\text{CH}_3$), 3-methyl-2-pentyl ($-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$), 4-methyl-2-pentyl ($-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}(\text{CH}_3)_2$), 3-methyl-3-pentyl ($-\text{C}(\text{CH}_3)(\text{CH}_2\text{CH}_3)_2$), 2-methyl-3-pentyl ($-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}(\text{CH}_3)_2$), 2,3-dimethyl-2-butyl ($-\text{C}(\text{CH}_3)_2\text{CH}(\text{CH}_3)_2$), 3,3-dimethyl-2-butyl ($-\text{CH}(\text{CH}_3)\text{C}(\text{CH}_3)_3$), and octyl ($-(\text{CH}_2)_7\text{CH}_3$).

“Alkoxy” means a group having the formula $-\text{O-alkyl}$, in which an alkyl group, as defined above, is attached to the parent molecule via an oxygen atom. The alkyl portion of an alkoxy group can have 1 to 20 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_{20}$ alkoxy), 1 to 12 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_{12}$ alkoxy), or 1 to 6 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_6$ alkoxy). Examples of suitable alkoxy groups include, but are not limited to, methoxy ($-\text{O-CH}_3$ or $-\text{OMe}$), ethoxy ($-\text{OCH}_2\text{CH}_3$ or $-\text{OEt}$), *t*-butoxy ($-\text{O-C}(\text{CH}_3)_3$ or $-\text{OtBu}$) and the like.

“Haloalkyl” is an alkyl group, as defined above, in which one or more hydrogen atoms of the alkyl group is replaced with a halogen atom. The alkyl portion of a haloalkyl group can have 1 to 20 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_{20}$ haloalkyl), 1 to 12 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_{12}$ haloalkyl), or 1 to 6 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_6$ alkyl). Examples of suitable haloalkyl groups include, but are not limited to, $-\text{CF}_3$, $-\text{CHF}_2$, $-\text{CFH}_2$, $-\text{CH}_2\text{CF}_3$, and the like.

“Alkenyl” is a hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms with at least one site of unsaturation, *i.e.* a carbon-carbon, sp^2 double bond. For example, an alkenyl group can have 2 to 20 carbon atoms (*i.e.*, $\text{C}_2\text{-C}_{20}$ alkenyl), 2 to 8 carbon atoms (*i.e.*, $\text{C}_2\text{-C}_8$ alkenyl), or 2 to 6 carbon atoms (*i.e.*, $\text{C}_2\text{-C}_6$ alkenyl). Examples of suitable alkenyl groups include, but are not limited to, ethylene

- 5 or vinyl ($-\text{CH}=\text{CH}_2$), allyl ($-\text{CH}_2\text{CH}=\text{CH}_2$), cyclopentenyl ($-\text{C}_5\text{H}_7$), and 5-hexenyl ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$).

“Alkynyl” is a hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms with at least one site of unsaturation, *i.e.* a carbon-carbon, *sp* triple bond. For example, an alkynyl group can have 2 to 20 carbon atoms (*i.e.*, $\text{C}_2\text{-C}_{20}$ alkynyl), 2 to 8 carbon atoms (*i.e.*, $\text{C}_2\text{-C}_8$ alkyne), or 2 to 6 carbon atoms (*i.e.*, $\text{C}_2\text{-C}_6$ alkynyl). Examples of suitable alkynyl groups include, but are not limited to, acetylenic ($-\text{C}\equiv\text{CH}$), propargyl ($-\text{CH}_2\text{C}\equiv\text{CH}$), and the like.

“Alkylene” refers to a saturated, branched or straight chain or cyclic hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkane. For example, an alkylene group can have 1 to 20 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon atoms. Typical alkylene radicals include, but are not limited to, methylene ($-\text{CH}_2-$), 1,1-ethyl ($-\text{CH}(\text{CH}_3)-$), 1,2-ethyl ($-\text{CH}_2\text{CH}_2-$), 1,1-propyl ($-\text{CH}(\text{CH}_2\text{CH}_3)-$), 1,2-propyl ($-\text{CH}_2\text{CH}(\text{CH}_3)-$), 1,3-propyl ($-\text{CH}_2\text{CH}_2\text{CH}_2-$), 1,4-butyl ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$), and the like.

“Alkenylene” refers to an unsaturated, branched or straight chain or cyclic hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkene. For example, an alkenylene group can have 1 to 20 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon atoms. Typical alkenylene radicals include, but are not limited to, 1,2-ethylene ($-\text{CH}=\text{CH}-$).

“Alkynylene” refers to an unsaturated, branched or straight chain or cyclic hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkyne. For example, an alkynylene group can have 1 to 20 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon atoms. Typical alkynylene radicals include, but are not limited to, acetylene ($-\text{C}\equiv\text{C}-$), propargyl ($-\text{CH}_2\text{C}\equiv\text{C}-$), and 4-pentynyl ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{C}-$).

“Amino” refers generally to a nitrogen radical which can be considered a derivative of ammonia, having the formula $-\text{N}(\text{X})_2$, where each “X” is independently H,



- 5 substituted or unsubstituted alkyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, etc. The hybridization of the nitrogen is approximately sp^3 . Nonlimiting types of amino include $-NH_2$, $-N(alkyl)_2$, $-NH(alkyl)$, $-N(carbocyclyl)_2$, $-NH(carbocyclyl)$, $-N(heterocyclyl)_2$, $-NH(heterocyclyl)$, $-N(aryl)_2$, $-NH(aryl)$, $-N(alkyl)(aryl)$, $-N(alkyl)(heterocyclyl)$, $-N(carbocyclyl)(heterocyclyl)$, $-N(aryl)(heteroaryl)$, $-N(alkyl)(heteroaryl)$, etc. The term "alkylamino" refers to an amino group substituted with at least one alkyl group. Nonlimiting examples of amino groups include $-NH_2$, $-NH(CH_3)$, $-N(CH_3)_2$, $-NH(CH_2CH_3)$, $-N(CH_2CH_3)_2$, $-NH(phenyl)$, $-N(phenyl)_2$, $-NH(benzyl)$, $-N(benzyl)_2$, etc. Substituted alkylamino refers generally to alkylamino groups, as defined above, in which at least one substituted alkyl, as defined herein, is attached to the amino nitrogen atom. Non-limiting examples of substituted alkylamino includes $-NH(alkylene-C(O)-OH)$, $-NH(alkylene-C(O)-O-alkyl)$, $-N(alkylene-C(O)-OH)_2$, $-N(alkylene-C(O)-O-alkyl)_2$, etc.

"Aryl" means an aromatic hydrocarbon radical derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic ring system. For example, an aryl group can have 6 to 20 carbon atoms, 6 to 14 carbon atoms, or 6 to 10 carbon atoms. Typical aryl groups include, but are not limited to, radicals derived from benzene (e.g., phenyl), substituted benzene, naphthalene, anthracene, biphenyl, and the like.

"Arylalkyl" refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with an aryl radical. Typical arylalkyl groups include, but are not limited to, benzyl, 2-phenylethan-1-yl, naphthylmethyl, 2-naphthylethan-1-yl, naphthobenzyl, 2-naphthophenylethan-1-yl and the like. The arylalkyl group can comprise 7 to 20 carbon atoms, e.g., the alkyl moiety is 1 to 6 carbon atoms and the aryl moiety is 6 to 14 carbon atoms.

"Arylalkenyl" refers to an acyclic alkenyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, but also an sp^2 carbon atom, is replaced with an aryl radical. The aryl portion of the arylalkenyl can include, for example, any of the aryl groups disclosed herein, and the

L

5 alkenyl portion of the arylalkenyl can include, for example, any of the alkenyl groups disclosed herein. The arylalkenyl group can comprise 8 to 20 carbon atoms, *e.g.*, the alkenyl moiety is 2 to 6 carbon atoms and the aryl moiety is 6 to 14 carbon atoms.

“Arylalkynyl” refers to an acyclic alkynyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, but
 10 also an sp carbon atom, is replaced with an aryl radical. The aryl portion of the arylalkynyl can include, for example, any of the aryl groups disclosed herein, and the alkynyl portion of the arylalkynyl can include, for example, any of the alkynyl groups disclosed herein. The arylalkynyl group can comprise 8 to 20 carbon atoms, *e.g.*, the alkynyl moiety is 2 to 6 carbon atoms and the aryl moiety is 6 to 14 carbon atoms.

15 The term “substituted” in reference to alkyl, alkylene, aryl, arylalkyl, alkoxy, heterocyclyl, heteroaryl, carbocyclyl, etc., for example, “substituted alkyl”, “substituted alkylene”, “substituted aryl”, “substituted arylalkyl”, “substituted heterocyclyl”, and “substituted carbocyclyl”, unless otherwise indicated, means alkyl, alkylene, aryl, arylalkyl, heterocyclyl, carbocyclyl respectively, in which one or more
 20 hydrogen atoms are each independently replaced with a non-hydrogen substituent. Typical substituents include, but are not limited to, $-X$, $-R^b$, $-O^-$, $=O$, $-OR^b$, $-SR^b$, $-S^-$, $-NR^b_2$, $-N^+R^b_3$, $=NR^b$, $-CX_3$, $-CN$, $-OCN$, $-SCN$, $-N=C=O$, $-NCS$, $-NO$, $-NO_2$, $=N_2$, $-N_3$, $-NHC(=O)R^b$, $-OC(=O)R^b$, $-NHC(=O)NR^b_2$, $-S(=O)_2-$, $-S(=O)_2OH$, $-S(=O)_2R^b$, $-OS(=O)_2OR^b$, $-S(=O)_2NR^b_2$, $-S(=O)R^b$, $-OP(=O)(OR^b)_2$, $-P(=O)(OR^b)_2$, $-P(=O)(O^-)_2$,
 25 $-P(=O)(OH)_2$, $-P(O)(OR^b)(O^-)$, $-C(=O)R^b$, $-C(=O)X$, $-C(S)R^b$, $-C(O)OR^b$, $-C(O)O^-$, $-C(S)OR^b$, $-C(OSR^b)$, $-C(S)SR^b$, $-C(O)NR^b_2$, $-C(S)NR^b_2$, $-C(=NR^b)NR^b_2$, where each X is independently a halogen: F, Cl, Br, or I; and each R^b is independently H, alkyl, aryl, arylalkyl, a heterocycle, or a protecting group or prodrug moiety. Alkylene, alkenylene, and alkynylene groups may also be similarly substituted. Unless otherwise
 30 indicated, when the term “substituted” is used in conjunction with groups such as arylalkyl, which have two or more moieties capable of substitution, the substituents can be attached to the aryl moiety, the alkyl moiety, or both.

The term “prodrug” as used herein refers to any compound that when administered to a biological system generates the drug substance, *i.e.*, active ingredient,




- 5 as a result of spontaneous chemical reaction(s), enzyme catalyzed chemical reaction(s), photolysis, and/or metabolic chemical reaction(s). A prodrug is thus a covalently modified analog or latent form of a therapeutically active compound.

One skilled in the art will recognize that substituents and other moieties of the compounds of Formula I-III and Formula IV-VI should be selected in order to provide a
10 compound which is sufficiently stable to provide a pharmaceutically useful compound which can be formulated into an acceptably stable pharmaceutical composition. The definitions and substituents for various genus and subgenus of the present compounds are described and illustrated herein. It should be understood by one skilled in the art that any combination of the definitions and substituents described above should not
15 result in an inoperable species or compound. "Inoperable species or compounds" means compound structures that violates relevant scientific principles (such as, for example, a carbon atom connecting to more than four covalent bonds) or compounds too unstable to permit isolation and formulation into pharmaceutically acceptable dosage forms.

20 "Heteroalkyl" refers to an alkyl group where one or more carbon atoms have been replaced with a heteroatom, such as, O, N, or S. For example, if the carbon atom of the alkyl group which is attached to the parent molecule is replaced with a heteroatom (e.g., O, N, or S) the resulting heteroalkyl groups are, respectively, an alkoxy group (e.g., -OCH₃, etc.), an amine (e.g., -NHCH₃, -N(CH₃)₂, etc.), or a thioalkyl group (e.g.,
25 -SCH₃). If a non-terminal carbon atom of the alkyl group which is not attached to the parent molecule is replaced with a heteroatom (e.g., O, N, or S) the resulting heteroalkyl groups are, respectively, an alkyl ether (e.g., -CH₂CH₂-O-CH₃, etc.), an alkyl amine (e.g., -CH₂NHCH₃, -CH₂N(CH₃)₂, etc.), or a thioalkyl ether (e.g., -CH₂-S-CH₃). If a terminal carbon atom of the alkyl group is replaced with a heteroatom (e.g., O, N, or S),
30 the resulting heteroalkyl groups are, respectively, a hydroxyalkyl group (e.g., -CH₂CH₂-OH), an aminoalkyl group (e.g., -CH₂NH₂), or an alkyl thiol group (e.g., -CH₂CH₂-SH). A heteroalkyl group can have, for example, 1 to 20 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon atoms. A C₁-C₆ heteroalkyl group means a heteroalkyl group having 1 to 6 carbon atoms.




- 5 "Heterocycle" or "heterocyclyl" as used herein includes by way of example and not limitation those heterocycles described in Paquette, Leo A.; Principles of Modern Heterocyclic Chemistry (W.A. Benjamin, New York, 1968), particularly Chapters 1, 3, 4, 6, 7, and 9; The Chemistry of Heterocyclic Compounds, A Series of Monographs" (John Wiley & Sons, New York, 1950 to present), in particular
- 10 Volumes 13, 14, 16, 19, and 28; and *J. Am. Chem. Soc.* (1960) 82:5566. In one specific embodiment of the invention "heterocycle" includes a "carbocycle" as defined herein, wherein one or more (e.g. 1, 2, 3, or 4) carbon atoms have been replaced with a heteroatom (e.g. O, N, or S). The terms "heterocycle" or
- "heterocyclyl" includes saturated rings, partially unsaturated rings, and aromatic rings
- 15 (i.e., heteroaromatic rings). Substituted heterocyclyls include, for example, heterocyclic rings substituted with any of the substituents disclosed herein including carbonyl groups. A non-limiting example of a carbonyl substituted heterocyclyl is:



- Examples of heterocycles include by way of example and not limitation
- 20 pyridyl, dihydropyridyl, tetrahydropyridyl (piperidyl), thiazolyl, tetrahydrothiophenyl, sulfur oxidized tetrahydrothiophenyl, pyrimidinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, imidazolyl, tetrazolyl, benzofuranyl, thianaphthalenyl, indolyl, indolenyl, quinolinyl, isoquinolinyl, benzimidazolyl, piperidynyl, 4-piperidonyl, pyrrolidinyl, 2-pyrrolidonyl, pyrrolinyl, tetrahydrofuranyl,
- 25 tetrahydroquinolinyl, tetrahydroisoquinolinyl, decahydroquinolinyl, octahydroisoquinolinyl, azocinyl, triazinyl, 6H-1,2,5-thiadiazinyl, 2H,6H-1,5,2-dithiazinyl, thienyl, thianthrenyl, pyranyl, isobenzofuranyl, chromenyl, xanthenyl, phenoxathinyl, 2H-pyrrolyl, isothiazolyl, isoxazolyl, pyrazinyl, pyridazinyl, indoliziny, isoindolyl, 3H-indolyl, 1H-indazolyl, purinyl, 4H-quinoliziny,
- 30 phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, pteridinyl, 4aH-carbazolyl, carbazolyl, β -carbolinyl, phenanthridinyl, acridinyl, pyrimidinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, furazanyl, phenoxazinyl, isochromanyl,

- 5 chromanyl, imidazolidinyl, imidazoliny, pyrazolidinyl, pyrazoliny, piperazinyl, indoliny, isoindoliny, quinuclidiny, morpholiny, oxazolidinyl, benzotriazolyl, benzisoxazolyl, oxindolyl, benzoxazoliny, isatinoyl, and bis-tetrahydrofurany:



- By way of example and not limitation, carbon bonded heterocycles are bonded
- 10 at position 2, 3, 4, 5, or 6 of a pyridine, position 3, 4, 5, or 6 of a pyridazine, position 2, 4, 5, or 6 of a pyrimidine, position 2, 3, 5, or 6 of a pyrazine, position 2, 3, 4, or 5 of a furan, tetrahydrofuran, thiofuran, thiophene, pyrrole or tetrahydropyrrole, position 2, 4, or 5 of an oxazole, imidazole or thiazole, position 3, 4, or 5 of an isoxazole, pyrazole, or isothiazole, position 2 or 3 of an aziridine, position 2, 3, or 4 of
- 15 an azetidine, position 2, 3, 4, 5, 6, 7, or 8 of a quinoline or position 1, 3, 4, 5, 6, 7, or 8 of an isoquinoline. Still more typically, carbon bonded heterocycles include 2-pyridyl, 3-pyridyl, 4-pyridyl, 5-pyridyl, 6-pyridyl, 3-pyridazinyl, 4-pyridazinyl, 5-pyridazinyl, 6-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 6-pyrimidinyl, 2-pyrazinyl, 3-pyrazinyl, 5-pyrazinyl, 6-pyrazinyl, 2-thiazolyl, 4-
- 20 thiazolyl, or 5-thiazolyl.

- By way of example and not limitation, nitrogen bonded heterocycles are bonded at position 1 of an aziridine, azetidine, pyrrole, pyrrolidine, 2-pyrroline, 3-pyrroline, imidazole, imidazolidine, 2-imidazoline, 3-imidazoline, pyrazole, pyrazoline, 2-pyrazoline, 3-pyrazoline, piperidine, piperazine, indole, indoline, 1H-
- 25 indazole, position 2 of a isoindole, or isoindoline, position 4 of a morpholine, and position 9 of a carbazole, or β -carboline. Still more typically, nitrogen bonded heterocycles include 1-aziridyl, 1-azetetyl, 1-pyrrolyl, 1-imidazolyl, 1-pyrazolyl, and 1-piperidinyl.

- "Heterocyclalkyl" refers to an acyclic alkyl radical in which one of the
- 30 hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with a heterocyclalkyl radical (*i.e.*, a heterocyclalkylene- moiety). Typical heterocyclalkyl groups include, but are not limited to heterocyclalk- CH_2 -, 2-

5 (heterocyclyl)ethan-1-yl, and the like, wherein the "heterocyclyl" portion includes any of the heterocyclyl groups described above, including those described in Principles of Modern Heterocyclic Chemistry. One skilled in the art will also understand that the heterocyclyl group can be attached to the alkyl portion of the heterocyclyl alkyl by means of a carbon-carbon bond or a carbon-heteroatom bond, with the proviso that
10 the resulting group is chemically stable. The heterocyclyl alkyl group comprises 3 to 20 carbon atoms, *e.g.*, the alkyl portion of the arylalkyl group is 1 to 6 carbon atoms and the heterocyclyl moiety is 2 to 14 carbon atoms. Examples of heterocyclylalkyls include by way of example and not limitation 5-membered sulfur, oxygen, and/or nitrogen containing heterocycles such as thiazolylmethyl, 2-thiazolyethan-1-yl,
15 imidazolylmethyl, oxazolylmethyl, thiadiazolylmethyl, etc., 6-membered sulfur, oxygen, and/or nitrogen containing heterocycles such as piperidinylmethyl, piperazinylmethyl, morpholinylmethyl, pyridinylmethyl, pyridizylmethyl, pyrimidylmethyl, pyrazinylmethyl, etc.

"Heterocyclylalkenyl" refers to an acyclic alkenyl radical in which one of the
20 hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, but also a sp^2 carbon atom, is replaced with a heterocyclyl radical (*i.e.*, a heterocyclyl-alkenylene- moiety). The heterocyclyl portion of the heterocyclyl alkenyl group includes any of the heterocyclyl groups described herein, including those described in Principles of Modern Heterocyclic Chemistry, and the alkenyl portion of the
25 heterocyclyl alkenyl group includes any of the alkenyl groups disclosed herein. One skilled in the art will also understand that the heterocyclyl group can be attached to the alkenyl portion of the heterocyclyl alkenyl by means of a carbon-carbon bond or a carbon-heteroatom bond, with the proviso that the resulting group is chemically stable. The heterocyclyl alkenyl group comprises 4 to 20 carbon atoms, *e.g.*, the
30 alkenyl portion of the heterocyclyl alkenyl group is 2 to 6 carbon atoms and the heterocyclyl moiety is 2 to 14 carbon atoms.

"Heterocyclylalkynyl" refers to an acyclic alkynyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, but also an sp carbon atom, is replaced with a heterocyclyl radical (*i.e.*, a heterocyclyl-



5 alkynylene- moiety). The heterocyclyl portion of the heterocyclyl alkynyl group includes any of the heterocyclyl groups described herein, including those described in Principles of Modern Heterocyclic Chemistry, and the alkynyl portion of the heterocyclyl alkynyl group includes any of the alkynyl groups disclosed herein. One skilled in the art will also understand that the heterocyclyl group can be attached to
10 the alkynyl portion of the heterocyclyl alkynyl by means of a carbon-carbon bond or a carbon-heteroatom bond, with the proviso that the resulting group is chemically stable. The heterocyclyl alkynyl group comprises 4 to 20 carbon atoms, e.g., the alkynyl portion of the heterocyclyl alkynyl group is 2 to 6 carbon atoms and the heterocyclyl moiety is 2 to 14 carbon atoms.

15 "Heteroaryl" refers to an aromatic heterocyclyl having at least one heteroatom in the ring. Non-limiting examples of suitable heteroatoms which can be included in the aromatic ring include oxygen, sulfur, and nitrogen. Non-limiting examples of heteroaryl rings include all of those aromatic rings listed in the definition of "heterocyclyl", including pyridinyl, pyrrolyl, oxazolyl, indolyl, isoindolyl, purinyl,
20 furanyl, thienyl, benzofuranyl, benzothiophenyl, carbazolyl, imidazolyl, thiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, quinolyl, isoquinolyl, pyridazyl, pyrimidyl, pyrazyl, etc.

"Carbocycle" or "carbocyclyl" refers to a saturated (i.e., cycloalkyl), partially unsaturated (e.g., cycloalkenyl, cycloalkadienyl, etc.) or aromatic ring having 3 to 7
25 carbon atoms as a monocycle, 7 to 12 carbon atoms as a bicycle, and up to about 20 carbon atoms as a polycycle. Monocyclic carbocycles have 3 to 7 ring atoms, still more typically 5 or 6 ring atoms. Bicyclic carbocycles have 7 to 12 ring atoms, e.g., arranged as a bicyclo [4,5], [5,5], [5,6] or [6,6] system, or 9 or 10 ring atoms arranged as a bicyclo [5,6] or [6,6] system, or spiro-fused rings. Non-limiting examples of
30 monocyclic carbocycles include cyclopropyl, cyclobutyl, cyclopentyl, 1-cyclopent-1-enyl, 1-cyclopent-2-enyl, 1-cyclopent-3-enyl, cyclohexyl, 1-cyclohex-1-enyl, 1-cyclohex-2-enyl, 1-cyclohex-3-enyl, and phenyl. Non-limiting examples of bicyclo carbocycles includes naphthyl, tetrahydronaphthalene, and decaline.

"Carbocyclalkyl" refers to an acyclic alkyl radical in which one of the



- 5 hydrogen atoms bonded to a carbon atom is replaced with a carbocyclyl radical as described herein. Typical, but non-limiting, examples of carbocyclylalkyl groups include cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

- “Arylheteroalkyl” refers to a heteroalkyl as defined herein, in which a
- 10 hydrogen atom (which may be attached either to a carbon atom or a heteroatom) has been replaced with an aryl group as defined herein. The aryl groups may be bonded to a carbon atom of the heteroalkyl group, or to a heteroatom of the heteroalkyl group, provided that the resulting arylheteroalkyl group provides a chemically stable moiety. For example, an arylheteroalkyl group can have the general formulae -alkylene-
- 15 O-aryl, -alkylene-O-alkylene-aryl, -alkylene-NH-aryl, -alkylene-NH-alkylene-aryl, -alkylene-S-aryl, -alkylene-S-alkylene-aryl, etc. In addition, any of the alkylene moieties in the general formulae above can be further substituted with any of the substituents defined or exemplified herein.

- “Heteroarylalkyl” refers to an alkyl group, as defined herein, in which a
- 20 hydrogen atom has been replaced with a heteroaryl group as defined herein. Non-limiting examples of heteroaryl alkyl include -CH₂-pyridinyl, -CH₂-pyrrolyl, -CH₂-oxazolyl, -CH₂-indolyl, -CH₂-isoindolyl, -CH₂-purinyl, -CH₂-furanlyl, -CH₂-thienyl, -CH₂-benzofuranyl, -CH₂-benzothiophenyl, -CH₂-carbazolyl, -CH₂-imidazolyl, -CH₂-thiazolyl, -CH₂-isoxazolyl, -CH₂-pyrazolyl, -CH₂-isothiazolyl,
- 25 -CH₂-quinolyl, -CH₂-isoquinolyl, -CH₂-pyridazyl, -CH₂-pyrimidyl, -CH₂-pyrazyl, -CH(CH₃)-pyridinyl, -CH(CH₃)-pyrrolyl, -CH(CH₃)-oxazolyl, -CH(CH₃)-indolyl, -CH(CH₃)-isoindolyl, -CH(CH₃)-purinyl, -CH(CH₃)-furanlyl, -CH(CH₃)-thienyl, -CH(CH₃)-benzofuranyl, -CH(CH₃)-benzothiophenyl, -CH(CH₃)-carbazolyl, -CH(CH₃)-imidazolyl, -CH(CH₃)-thiazolyl, -CH(CH₃)-isoxazolyl,
- 30 -CH(CH₃)-pyrazolyl, -CH(CH₃)-isothiazolyl, -CH(CH₃)-quinolyl, -CH(CH₃)-isoquinolyl, -CH(CH₃)-pyridazyl, -CH(CH₃)-pyrimidyl, -CH(CH₃)-pyrazyl, etc.

The term “optionally substituted” in reference to a particular moiety of the compound of Formula I-III and Formula IV-VI (e.g., an optionally substituted aryl





- 5 group) refers to a moiety wherein all substituents are hydrogen or wherein one or more of the hydrogens of the moiety may be replaced by substituents such as those listed under the definition of "substituted" or as otherwise indicated.

The term "optionally replaced" in reference to a particular moiety of the compound of Formula I-III and Formula IV-VI (e.g., the carbon atoms of said (C₁-
10 C₈)alkyl may be optionally replaced by -O-, -S-, or -NR^a-) means that one or more of the methylene groups of the (C₁-C₈)alkyl may be replaced by 0, 1, 2, or more of the groups specified (e.g., -O-, -S-, or -NR^a-).

The term "non-terminal carbon atom(s)" in reference to an alkyl, alkenyl, alkynyl, alkylene, alkenylene, or alkynylene moiety refers to the carbon atoms in the
15 moiety that intervene between the first carbon atom of the moiety and the last carbon atom in the moiety. Therefore, by way of example and not limitation, in the alkyl moiety -CH₂(C^{*})H₂(C^{*})H₂CH₃ or alkylene moiety -CH₂(C^{*})H₂(C^{*})H₂CH₂- the C^{*} atoms would be considered to be the non-terminal carbon atoms.

Certain Y and Y¹ alternatives are nitrogen oxides such as ⁺N(O)(R) or
20 ⁺N(O)(OR). These nitrogen oxides, as shown here attached to a carbon atom, can also

be represented by charge separated groups such as  or ,
respectively, and are intended to be equivalent to the aforementioned representations for the purposes of describing this invention.


"Linker" or "link" means a chemical moiety comprising a covalent bond or a
25 chain of atoms. Linkers include repeating units of alkyloxy (e.g. polyethyleneoxy, PEG, polymethyleneoxy) and alkylamino (e.g. polyethyleneamino, JeffamineTM); and diacid ester and amides including succinate, succinamide, diglycolate, malonate, and caproamide.

The terms such as "oxygen-linked", "nitrogen-linked", "carbon-linked",
30 "sulfur-linked", or "phosphorous-linked" mean that if a bond between two moieties can be formed by using more than one type of atom in a moiety, then the bond formed between the moieties is through the atom specified. For example, a nitrogen-linked




- 5 amino acid would be bonded through a nitrogen atom of the amino acid rather than through an oxygen or carbon atom of the amino acid.

Unless otherwise specified, the carbon atoms of the compounds of Formula I-III and Formula IV-VI are intended to have a valence of four. In some chemical structure representations where carbon atoms do not have a sufficient number of variables attached to produce a valence of four, the remaining carbon substituents
10 needed to provide a valence of four should be assumed to be hydrogen. For example,



has the same meaning as



- “Protecting group” refers to a moiety of a compound that masks or alters the
15 properties of a functional group or the properties of the compound as a whole. The chemical substructure of a protecting group varies widely. One function of a protecting group is to serve as an intermediate in the synthesis of the parental drug substance. Chemical protecting groups and strategies for protection/deprotection are well known in the art. See: “Protective Groups in Organic Chemistry”, Theodora W.
20 Greene (John Wiley & Sons, Inc., New York, 1991. Protecting groups are often utilized to mask the reactivity of certain functional groups, to assist in the efficiency

Handwritten signature

5 of desired chemical reactions, e.g. making and breaking chemical bonds in an ordered and planned fashion. Protection of functional groups of a compound alters other physical properties besides the reactivity of the protected functional group, such as the polarity, lipophilicity (hydrophobicity), and other properties which can be measured by common analytical tools. Chemically protected intermediates may themselves be
10 biologically active or inactive.

Protected compounds may also exhibit altered, and in some cases, optimized properties *in vitro* and *in vivo*, such as passage through cellular membranes and resistance to enzymatic degradation or sequestration. In this role, protected compounds with intended therapeutic effects may be referred to as prodrugs. Another
15 function of a protecting group is to convert the parental drug into a prodrug, whereby the parental drug is released upon conversion of the prodrug *in vivo*. Because active prodrugs may be absorbed more effectively than the parental drug, prodrugs may possess greater potency *in vivo* than the parental drug. Protecting groups are removed either *in vitro*, in the instance of chemical intermediates, or *in vivo*, in the case of
20 prodrugs. With chemical intermediates, it is not particularly important that the resulting products after deprotection, e.g. alcohols, be physiologically acceptable, although in general it is more desirable if the products are pharmacologically innocuous.

“Prodrug moiety” means a labile functional group which separates from the
25 active inhibitory compound during metabolism, systemically, inside a cell, by hydrolysis, enzymatic cleavage, or by some other process (Bundgaard, Hans, “Design and Application of Prodrugs” in Textbook of Drug Design and Development (1991), P. Krogsgaard-Larsen and H. Bundgaard, Eds. Harwood Academic Publishers, pp. 113-191). Enzymes which are capable of an enzymatic activation mechanism with the
30 phosphonate prodrug compounds of the invention include, but are not limited to, amidases, esterases, microbial enzymes, phospholipases, cholinesterases, and phosphatases. Prodrug moieties can serve to enhance solubility, absorption and lipophilicity to optimize drug delivery, bioavailability and efficacy.

A prodrug moiety may include an active metabolite or drug itself.

✓

5 Exemplary prodrug moieties include the hydrolytically sensitive or labile
acyloxymethyl esters $-\text{CH}_2\text{OC}(=\text{O})\text{R}^{30}$ and acyloxymethyl carbonates
 $-\text{CH}_2\text{OC}(=\text{O})\text{OR}^{30}$ where R^{30} is C_1-C_6 alkyl, C_1-C_6 substituted alkyl, C_6-C_{20} aryl or
 C_6-C_{20} substituted aryl. The acyloxyalkyl ester was used as a prodrug strategy for
carboxylic acids and then applied to phosphates and phosphonates by Farquhar et al
10 (1983) *J. Pharm. Sci.* 72: 324; also US Patent Nos. 4816570, 4968788, 5663159 and
5792756. In certain compounds of the invention, a prodrug moiety is part of a
phosphate group. The acyloxyalkyl ester may be used to deliver phosphoric acids
across cell membranes and to enhance oral bioavailability. A close variant of the
acyloxyalkyl ester, the alkoxycarbonyloxyalkyl ester (carbonate), may also enhance
15 oral bioavailability as a prodrug moiety in the compounds of the combinations of the
invention. An exemplary acyloxymethyl ester is pivaloyloxymethoxy, (POM)
 $-\text{CH}_2\text{OC}(=\text{O})\text{C}(\text{CH}_3)_3$. An exemplary acyloxymethyl carbonate prodrug moiety is
pivaloyloxymethylcarbonate (POC) $-\text{CH}_2\text{OC}(=\text{O})\text{OC}(\text{CH}_3)_3$.

The phosphate group may be a phosphate prodrug moiety. The prodrug
20 moiety may be sensitive to hydrolysis, such as, but not limited to those comprising a
pivaloyloxymethyl carbonate (POC) or POM group. Alternatively, the prodrug
moiety may be sensitive to enzymatic potentiated cleavage, such as a lactate ester or a
phosphoramidate-ester group.

Aryl esters of phosphorus groups, especially phenyl esters, are reported to
25 enhance oral bioavailability (DeLambert et al (1994) *J. Med. Chem.* 37: 498). Phenyl
esters containing a carboxylic ester ortho to the phosphate have also been described
(Khamnei and Torrence, (1996) *J. Med. Chem.* 39:4109-4115). Benzyl esters are
reported to generate the parent phosphonic acid. In some cases, substituents at the
ortho-or *para*-position may accelerate the hydrolysis. Benzyl analogs with an
30 acylated phenol or an alkylated phenol may generate the phenolic compound through
the action of enzymes, e.g. esterases, oxidases, etc., which in turn undergoes cleavage
at the benzylic C-O bond to generate the phosphoric acid and the quinone methide
intermediate. Examples of this class of prodrugs are described by Mitchell et al
(1992) *J. Chem. Soc. Perkin Trans. I* 2345; Brook et al WO 91/19721. Still other



- 5 benzylic prodrugs have been described containing a carboxylic ester-containing group attached to the benzylic methylene (Glazier et al WO 91/19721). Thio-containing prodrugs are reported to be useful for the intracellular delivery of phosphonate drugs. These proesters contain an ethylthio group in which the thiol group is either esterified with an acyl group or combined with another thiol group to form a disulfide.
- 10 Deesterification or reduction of the disulfide generates the free thio intermediate which subsequently breaks down to the phosphoric acid and episulfide (Puech et al (1993) *Antiviral Res.*, 22: 155-174; Benzaria et al (1996) *J. Med. Chem.* 39: 4958). Cyclic phosphonate esters have also been described as prodrugs of phosphorus-containing compounds (Erion et al, US Patent No. 6312662).
- 15 It is to be noted that all enantiomers, diastereomers, and racemic mixtures, tautomers, polymorphs, pseudopolymorphs of compounds within the scope of Formula I, Formula II, Formula III, Formula IV, Formula V, or Formula VI and pharmaceutically acceptable salts thereof are embraced by the present invention. All mixtures of such enantiomers and diastereomers are within the scope of the present
- 20 invention.

A compound of Formula I-III and Formula IV-VI and its pharmaceutically acceptable salts may exist as different polymorphs or pseudopolymorphs. As used herein, crystalline polymorphism means the ability of a crystalline compound to exist in different crystal structures. The crystalline polymorphism may result from

25 differences in crystal packing (packing polymorphism) or differences in packing between different conformers of the same molecule (conformational polymorphism). As used herein, crystalline pseudopolymorphism means the ability of a hydrate or solvate of a compound to exist in different crystal structures. The pseudopolymorphs of the instant invention may exist due to differences in crystal packing (packing

30 pseudopolymorphism) or due to differences in packing between different conformers of the same molecule (conformational pseudopolymorphism). The instant invention comprises all polymorphs and pseudopolymorphs of the compounds of Formula I-III and Formula IV-VI and their pharmaceutically acceptable salts.



5 A compound of Formula I-III and Formula IV-VI and its pharmaceutically acceptable salts may also exist as an amorphous solid. As used herein, an amorphous solid is a solid in which there is no long-range order of the positions of the atoms in the solid. This definition applies as well when the crystal size is two nanometers or less. Additives, including solvents, may be used to create the amorphous forms of the
10 instant invention. The instant invention comprises all amorphous forms of the compounds of Formula I-III and Formula IV-VI and their pharmaceutically acceptable salts.

 Selected substituents comprising the compounds of Formula I-III and Formula IV-VI are present to a recursive degree. In this context, "recursive substituent" means
15 that a substituent may recite another instance of itself. Because of the recursive nature of such substituents, theoretically, a large number of compounds may be present in any given embodiment. For example, R^x comprises a R^y substituent. R^y can be R. R can be W^3 . W^3 can be W^4 and W^4 can be R or comprise substituents comprising R^y . One of ordinary skill in the art of medicinal chemistry understands
20 that the total number of such substituents is reasonably limited by the desired properties of the compound intended. Such properties include, by way of example and not limitation, physical properties such as molecular weight, solubility or log P, application properties such as activity against the intended target, and practical properties such as ease of synthesis.


25 By way of example and not limitation, W^3 and R^y are recursive substituents in certain embodiments. Typically, each recursive substituent can independently occur 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, or 0, times in a given embodiment. More typically, each recursive substituent can independently occur 12 or fewer times in a given embodiment. Even more typically, each recursive
30 substituent can independently occur 3 or fewer times in a given embodiment. For example, W^3 will occur 0 to 8 times, R^y will occur 0 to 6 times in a given embodiment. Even more typically, W^3 will occur 0 to 6 times and R^y will occur 0 to 4 times in a given embodiment.



5 Recursive substituents are an intended aspect of the invention. One of ordinary skill in the art of medicinal chemistry understands the versatility of such substituents. To the degree that recursive substituents are present in an embodiment of the invention, the total number will be determined as set forth above.

10 The modifier "about" used in connection with a quantity is inclusive of the stated value and has the meaning dictated by the context (e.g., includes the degree of error associated with measurement of the particular quantity).

The compounds of the Formula I-III and Formula IV-VI may comprise a




phosphate group as R^7 , which may be a prodrug moiety

wherein

each Y or Y^1 is, independently, O, S, NR, $^+N(O)(R)$, N(OR), $^+N(O)(OR)$, or N-NR₂;

15 W^1 and W^2 , when taken together, are $-Y^3(C(R^y)_2)_3Y^3-$; or one of W^1 or W^2 together with either R^3 or R^4 is $-Y^3-$ and the other of W^1 or W^2 is Formula Ia; or W^1 and W^2 are each, independently, a group of Formula Ia:



wherein:

20 each Y^2 is independently a bond, O, CR₂, NR, $^+N(O)(R)$, N(OR), $^+N(O)(OR)$, N-NR₂, S, S-S, S(O), or S(O)₂;

each Y^3 is independently O, S, or NR;


M2 is 0, 1 or 2;

25 each R^y is independently H, F, Cl, Br, I, OH, R, $-C(=Y^1)R$, $-C(=Y^1)OR$, $-C(=Y^1)N(R)_2$, $-N(R)_2$, $-^+N(R)_3$, $-SR$, $-S(O)R$, $-S(O)_2R$, $-S(O)(OR)$, $-S(O)_2(OR)$, $-OC(=Y^1)R$, $-OC(=Y^1)OR$, $-OC(=Y^1)(N(R)_2)$, $-SC(=Y^1)R$, $-SC(=Y^1)OR$, -

2

- 5 $\text{SC}(=\text{Y}^1)(\text{N}(\text{R})_2)$, $-\text{N}(\text{R})\text{C}(=\text{Y}^1)\text{R}$, $-\text{N}(\text{R})\text{C}(=\text{Y}^1)\text{OR}$, or $-\text{N}(\text{R})\text{C}(=\text{Y}^1)\text{N}(\text{R})_2$, $-\text{SO}_2\text{NR}_2$, $-\text{CN}$, $-\text{N}_3$, $-\text{NO}_2$, $-\text{OR}$, a protecting group or W^3 ; or when taken together, two R^y on the same carbon atom form a carbocyclic ring of 3 to 7 carbon atoms;

each R^x is independently R^y , a protecting group, or the formula:



- 10 wherein:

M1a , M1c , and M1d are independently 0 or 1;

M12c is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

each R is H , halogen, $(\text{C}_1\text{--C}_8)$ alkyl, $(\text{C}_1\text{--C}_8)$ substituted alkyl, $(\text{C}_2\text{--C}_8)$ alkenyl, $(\text{C}_2\text{--C}_8)$ substituted alkenyl, $(\text{C}_2\text{--C}_8)$ alkynyl, $(\text{C}_2\text{--C}_8)$ substituted alkynyl, $\text{C}_6\text{--C}_{20}$ aryl, $\text{C}_6\text{--C}_{20}$ substituted aryl, $\text{C}_2\text{--C}_{20}$ heterocycle, $\text{C}_2\text{--C}_{20}$ substituted heterocyclyl, arylalkyl, substituted arylalkyl or a protecting group;

W^3 is W^4 or W^5 ; W^4 is R , $-\text{C}(\text{Y}^1)\text{R}^y$, $-\text{C}(\text{Y}^1)\text{W}^5$, $-\text{SO}_2\text{R}^y$, or $-\text{SO}_2\text{W}^5$; and W^5 is a carbocycle or a heterocycle wherein W^5 is independently substituted with 0 to 3 R^y groups.


- 20 W^5 carbocycles and W^5 heterocycles may be independently substituted with 0 to 3 R^y groups. W^5 may be a saturated, unsaturated or aromatic ring comprising a mono- or bicyclic carbocycle or heterocycle. W^5 may have 3 to 10 ring atoms, e.g., 3 to 7 ring atoms. The W^5 rings are saturated when containing 3 ring atoms, saturated or mono-unsaturated when containing 4 ring atoms, saturated, or mono- or di-unsaturated when containing 5 ring atoms, and saturated, mono- or di-unsaturated, or aromatic when containing 6 ring atoms.

A W^5 heterocycle may be a monocycle having 3 to 7 ring members (2 to 6 carbon atoms and 1 to 3 heteroatoms selected from N , O , P , and S) or a bicycle having 7 to 10 ring members (4 to 9 carbon atoms and 1 to 3 heteroatoms selected from N , O ,

K



- 5 P, and S). W^5 heterocyclic monocycles may have 3 to 6 ring atoms (2 to 5 carbon atoms and 1 to 2 heteroatoms selected from N, O, and S); or 5 or 6 ring atoms (3 to 5 carbon atoms and 1 to 2 heteroatoms selected from N and S). W^5 heterocyclic bicycles have 7 to 10 ring atoms (6 to 9 carbon atoms and 1 to 2 heteroatoms selected from N, O, and S) arranged as a bicyclo [4,5], [5,5], [5,6], or [6,6] system; or 9 to 10 ring atoms (8 to 9 carbon atoms and 1 to 2 hetero atoms selected from N and S) arranged as a bicyclo [5,6] or [6,6] system. The W^5 heterocycle may be bonded to Y^2 through a carbon, nitrogen, sulfur or other atom by a stable covalent bond.

- 15 W^5 heterocycles include for example, pyridyl, dihydropyridyl isomers, piperidine, pyridazinyl, pyrimidinyl, pyrazinyl, s-triazinyl, oxazolyl, imidazolyl, thiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, furanyl, thiofuranyl, thienyl, and pyrrolyl. W^5 also includes, but is not limited to, examples such as:





- 20 W^5 carbocycles and heterocycles may be independently substituted with 0 to 3 R groups, as defined above. For example, substituted W^5 carbocycles include:


67



5




Examples of substituted phenyl carbocycles include:



10

5 Embodiments of R^7 or R^7 together with R^4 include the structures



wherein

a is the point of attachment to R^7 ;

b is the point of attachment to R^4 ;


Ar is phenyl or naphthyl, wherein the phenyl and naphthyl are optionally

15 substituted with 1-3 R^{20} groups;

each R^y is independently (C_1-C_8) alkyl or C_5-C_6 carbocyclyl, wherein the alkyl and carbocyclyl are optionally substituted with 1-3 R^{20} groups;

each R is independently H, (C_1-C_6) alkyl, or arylalkyl; and

20 each R^{20} is independently halogen, CN, $N(R)_2$, OR, -SR, -S(O)R, -S(O)₂R, -S(O)(OR), -S(O)₂(OR), -C(=O)R, -C(=O)OR, or C(=O) $N(R)_2$.




Embodiments of

of Formula I-III and Formula IV-VI

compounds include substructures such as:


[Handwritten signature]

69




5

wherein each Y^{2b} is, independently, O or N(R). In another aspect of this embodiment, each Y^{2b} is O and each R^x is independently:




wherein M12c is 1, 2 or 3 and each Y^2 is independently a bond, O, CR_2 , or S.

- 10 In another aspect of this embodiment, one $Y^{2b}-R^x$ is $NH(R)$ and the other $Y^{2b}-R^x$ is $O-R^x$ wherein R^x is:




wherein M12c is 2. In another aspect of this embodiment, each Y^{2b} is O and each R^x is independently:




15

wherein M12c is 2. In another aspect of this embodiment, each Y^{2b} is O and each R^x is independently:


70




wherein M12c is 1 and Y² is a bond, O, or CR₂.




Other embodiments of of Formulas I-III and Formulas IV-VI compounds include substructures such as:



wherein each Y³ is, independently, O or N(R). In another aspect of this embodiment, each Y³ is O. In another aspect of this embodiment, the substructure is:




wherein R^y is W⁵ as defined herein.



Another embodiment of of Formula I-III and Formula IV-VI includes the substructures:


2

71



5



wherein each Y^{2c} is, independently, O, $N(R^y)$ or S.





Another embodiment of of Formula I-III and Formula IV-VI

compounds includes the substructures wherein one of W^1 or W^2 together with either R^3 or R^4 is $-Y^3-$ and the other of W^1 or W^2 is Formula Ia. Such an embodiment is

10 represented by a compound of Formula Ib selected from:






Formula Ib

In another aspect of the embodiment of Formula Ib, each Y and Y³ is O. In another aspect of the embodiment of Formula Ib, W¹ or W² is Y^{2b}-R^x; each Y, Y³ and Y^{2b} is O and R^x is:


10



wherein M12c is 1, 2 or 3 and each Y² is independently a bond, O, CR₂, or S. In another aspect of the embodiment of Formula Ib, W¹ or W² is Y^{2b}-R^x; each Y, Y³ and Y^{2b} is O and R^x is:


15

73




M12c

wherein M12c is 2. In another aspect of the embodiment of Formula Ib, W^1 or W^2 is $\text{Y}^{2b}-\text{R}^x$; each Y, Y^3 and Y^{2b} is O and R^x is:




M12c

wherein M12c is 1 and Y^2 is a bond, O, or CR_2 .




Another embodiment of of Formula I-III and Formula IV-VI compounds includes a substructure:




wherein W^5 is a carbocycle such as phenyl or substituted phenyl. In another aspect of this embodiment, the substructure is:

74




5

wherein Y^{2b} is O or N(R) and the phenyl carbocycle is substituted with 0 to 3 R groups. In another aspect of this embodiment of the substructure, R^x is:



wherein M12c is 1, 2 or 3 and each Y^2 is independently a bond, O, CR_2 , or S.


10



Another embodiment of

of Formula I-III and Formula IV-VI


includes substructure:




The chiral carbon of the amino acid and lactate moieties may be either the *R* or

15 *S* configuration or the racemic mixture.


75



5 Another embodiment of of Formula I-III and Formula IV-VI is substructure




wherein each Y^2 is, independently, $-O-$ or $-NH-$. In another aspect of this embodiment, R^Y is (C_1-C_8) alkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl or (C_2-C_8) substituted alkynyl. In another aspect of this embodiment, R^Y is (C_1-C_8) alkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl or (C_2-C_8) substituted alkynyl; and R is CH_3 . In another aspect of this embodiment, R^Y is (C_1-C_8) alkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl or (C_2-C_8) substituted alkynyl; R is CH_3 ; and each Y^2 is $-NH-$. In a aspect of this embodiment, W^1 and W^2 are, independently, nitrogen-linked, naturally occurring amino acids or naturally occurring amino acid esters. In another aspect of this embodiment, W^1 and W^2 are, independently, naturally-occurring 2-hydroxy carboxylic acids or naturally-occurring 2-hydroxy carboxylic acid esters wherein the acid or ester is linked to P through the 2-hydroxy group.



Another embodiment of of Formula I, Formula II, Formula III, Formula IV, Formula V, or Formula VI is substructure:


76



5




In one aspect of this embodiment, each R^x is, independently, (C_1-C_8) alkyl. In another aspect of this embodiment, each R^x is, independently, C_6-C_{20} aryl or C_6-C_{20} substituted aryl.

In a preferred embodiment,




10


is selected from



or




- 5 Another embodiment of of Formulas I-III and Formula IV-VI is substructure

























- wherein W^1 and W^2 are independently selected from one of the formulas in Tables 20.1-20.37 and Table 30.1 below. The variables used in Tables 20.1-20.37 (e.g., W^{23} , R^{21} , etc.) pertain only to Tables 20.1-20.37, unless otherwise indicated.
- The variables used in Tables 20.1 to 20.37 have the following definitions:
- each R^{21} is independently H or (C_1-C_8) alkyl;
 - each R^{22} is independently H, R^{21} , R^{23} or R^{24} wherein each R^{24} is independently substituted with 0 to 3 R^{23} ;
 - each R^{23} is independently R^{23a} , R^{23b} , R^{23c} or R^{23d} , provided that when R^{23} is bound to a heteroatom, then R^{23} is R^{23c} or R^{23d} ;
 - each R^{23a} is independently F, Cl, Br, I, -CN, N_3 or -NO₂;
 - each R^{23b} is independently Y^{21} ;
 - each R^{23c} is independently $-R^{2x}$, $-N(R^{2x})(R^{2x})$, $-SR^{2x}$, $-S(O)R^{2x}$, $-S(O)_2R^{2x}$, $-S(O)(OR^{2x})$, $-S(O)_2(OR^{2x})$, $-OC(=Y^{21})R^{2x}$, $-OC(=Y^{21})OR^{2x}$, $-OC(=Y^{21})(N(R^{2x})(R^{2x}))$, $-SC(=Y^{21})R^{2x}$, $-SC(=Y^{21})OR^{2x}$, $-SC(=Y^{21})(N(R^{2x})(R^{2x}))$, $-N(R^{2x})C(=Y^{21})R^{2x}$, $-N(R^{2x})C(=Y^{21})OR^{2x}$, or $-N(R^{2x})C(=Y^{21})(N(R^{2x})(R^{2x}))$;
 - each R^{23d} is independently $-C(=Y^{21})R^{2x}$, $-C(=Y^{21})OR^{2x}$ or $-C(=Y^{21})(N(R^{2x})(R^{2x}))$;
 - each R^{2x} is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl, heteroaryl; or two R^{2x} taken together with a nitrogen to which they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with -O-, -S- or $-NR^{21}-$; and wherein

- 5 one or more of the non-terminal carbon atoms of each said (C₁-C₈)alkyl may be optionally replaced with -O-, -S- or -NR²¹-;
 each R²⁴ is independently (C₁-C₈)alkyl, (C₂-C₈)alkenyl, or (C₂-C₈)alkynyl;
 each R²⁵ is independently R²⁴ wherein each R²⁴ is substituted with 0 to 3 R²³ groups;
- 10 each R^{25a} is independently (C₁-C₈)alkylene, (C₂-C₈)alkenylene, or (C₂-C₈)alkynylene any one of which said (C₁-C₈)alkylene, (C₂-C₈)alkenylene, or (C₂-C₈)alkynylene is substituted with 0-3 R²³ groups;
 each W²³ is independently W²⁴ or W²⁵;
 each W²⁴ is independently R²⁵, -C(=Y²¹)R²⁵, -C(=Y²¹)W²⁵, -SO₂R²⁵, or -
- 15 SO₂W²⁵;
 each W²⁵ is independently carbocycle or heterocycle wherein W²⁵ is independently substituted with 0 to 3 R²² groups; and
 each Y²¹ is independently O or S.


20 Table 20.1





5 Table 20.2Table 20.3

5 Table 20.4Table 20.5


81

5 Table 20.6


31




32




33

Table 20.7


34




35




36




37




38





39







40














41

5 Table 20.8Table 20.9

5 Table 20.10Table 20.11

5 Table 20.12Table 20.13

85

5 Table 20.14Table 20.15

5 Table 20.16










Table 20.17




5 Table 20.18Table 20.19

10



Table 20.20

5 Table 20.21Table 20.22


10


Table 20.23

15


5 Table 20.24Table 20.25

10


Table 20.26

5 Table 20.27


172




173




174




175




176



177



178




179


10

Table 20.28


91




180




181




182



183




184




185

5


Table 20.29




186




187




188




189




190



191




192




193


5

Table 20.30


194




195




196



197











198









199


Table 20.31




5

Table 20.32




5 Table 20.33


214




215




216




217




218



219



220



221

Table 20.34
























Table 20.35





5 Table 20.36Table 20.37

5 Table 30.1



5

Embodiments of R^x include esters, carbamates, carbonates, thioesters, amides, thioamides, and urea groups:



10

Any reference to the compounds of the invention described herein also includes a reference to a physiologically acceptable salt thereof. Examples of physiologically acceptable salts of the compounds of the invention include salts derived from an appropriate base, such as an alkali metal or an alkaline earth (for example, Na^+ , Li^+ , K^+ , Ca^{+2} and Mg^{+2}), ammonium and NR_4^+ (wherein R is defined herein). Physiologically acceptable salts of a nitrogen atom or an amino group include (a) acid addition salts formed with inorganic acids, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, sulfamic acids, phosphoric acid, nitric acid and the like; (b) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, isethionic acid, lactobionic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, benzenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, malonic acid, sulfosalicylic acid, glycolic acid, 2-hydroxy-3-naphthoate, pamoate, salicylic acid, stearic acid, phthalic acid, mandelic acid, lactic acid, ethanesulfonic acid, lysine, arginine, glutamic acid, glycine, serine, threonine, alanine, isoleucine, leucine and the like; and (c) salts formed from elemental anions for example, chlorine, bromine, and iodine. Physiologically acceptable salts of a compound of a hydroxy group include the anion of said compound in combination with a suitable cation such as Na^+ and NR_4^+ .

30

For therapeutic use, salts of active ingredients of the compounds of the invention will be physiologically acceptable, i.e. they will be salts derived from a

5 physiologically acceptable acid or base. However, salts of acids or bases which are not physiologically acceptable may also find use, for example, in the preparation or purification of a physiologically acceptable compound. All salts, whether or not derived from a physiologically acceptable acid or base, are within the scope of the present invention.

10 Finally, it is to be understood that the compositions herein comprise compounds of the invention in their un-ionized, as well as zwitterionic form, and combinations with stoichiometric amounts of water as in hydrates.

The compounds of the invention, exemplified by Formula I-III and Formula IV-VI may have chiral centers, e.g. chiral carbon or phosphorus atoms. The
15 compounds of the invention thus include racemic mixtures of all stereoisomers, including enantiomers, diastereomers, and atropisomers. In addition, the compounds of the invention include enriched or resolved optical isomers at any or all asymmetric, chiral atoms. In other words, the chiral centers apparent from the depictions are provided as the chiral isomers or racemic mixtures. Both racemic and diastereomeric
20 mixtures, as well as the individual optical isomers isolated or synthesized, substantially free of their enantiomeric or diastereomeric partners, are all within the scope of the invention. The racemic mixtures are separated into their individual, substantially optically pure isomers through well-known techniques such as, for example, the separation of diastereomeric salts formed with optically active adjuncts,
25 e.g., acids or bases followed by conversion back to the optically active substances. In most instances, the desired optical isomer is synthesized by means of stereospecific reactions, beginning with the appropriate stereoisomer of the desired starting material.

The term "chiral" refers to molecules which have the property of non-superimposability of the mirror image partner, while the term "achiral" refers to
30 molecules which are superimposable on their mirror image partner.

The term "stereoisomers" refers to compounds which have identical chemical constitution, but differ with regard to the arrangement of the atoms or groups in space.

"Diastereomer" refers to a stereoisomer with two or more centers of chirality and whose molecules are not mirror images of one another. Diastereomers have



5 different physical properties, e.g. melting points, boiling points, spectral properties, and reactivities. Mixtures of diastereomers may separate under high resolution analytical procedures such as electrophoresis and chromatography.

"Enantiomers" refer to two stereoisomers of a compound which are non-superimposable mirror images of one another.

10 Stereochemical definitions and conventions used herein generally follow S. P. Parker, Ed., McGraw-Hill Dictionary of Chemical Terms (1984) McGraw-Hill Book Company, New York; and Eliel, E. and Wilen, S., Stereochemistry of Organic Compounds (1994) John Wiley & Sons, Inc., New York. Many organic compounds exist in optically active forms, i.e., they have the ability to rotate the plane-

15 polarized light. In describing an optically active compound, the prefixes D and L or R and S are used to denote the absolute configuration of the molecule about its chiral center(s). The prefixes d and l, D and L, or (+) and (-) are employed to designate the sign of rotation of plane-polarized light by the compound, with S, (-), or l meaning that the compound is levorotatory while a compound prefixed with R, (+), or d is

20 dextrorotatory. For a given chemical structure, these stereoisomers are identical except that they are mirror images of one another. A specific stereoisomer may also be referred to as an enantiomer, and a mixture of such isomers is often called an enantiomeric mixture. A 50:50 mixture of enantiomers is referred to as a racemic mixture or a racemate, which may occur where there has been no stereoselection or

25 stereospecificity in a chemical reaction or process. The terms "racemic mixture" and "racemate" refer to an equimolar mixture of two enantiomeric species, devoid of optical activity.

Whenever a compound described herein is substituted with more than one of the same designated group, e.g., "R" or "R¹", then it will be understood that the


30 groups may be the same or different, i.e., each group is independently selected. Wavy lines, ~~~~~, indicate the site of covalent bond attachments to the adjoining substructures, groups, moieties, or atoms.

The compounds of the invention can also exist as tautomeric isomers in certain cases. Although only one delocalized resonance structure may be depicted, all such

2

5 forms are contemplated within the scope of the invention. For example, ene-amine tautomers can exist for purine, pyrimidine, imidazole, guanidine, amidine, and tetrazole systems and all their possible tautomeric forms are within the scope of the invention.

One skilled in the art will recognize that the pyrrolo[1,2-f][1,2,4]triazine,
 10 imidazo[1,5-f][1,2,4]triazine, imidazo[1,2-f][1,2,4]triazine, and [1,2,4]triazolo[4,3-f][1,2,4]triazine nucleosides can exist in tautomeric forms. For example, but not by way of limitation, structures (a) and (b) can have equivalent tautomeric forms as shown below:



15

All possible tautomeric forms of the heterocycles in all of the embodiments disclosed herein are within the scope of the invention.

Methods of Inhibition of HCV polymerase

Another aspect of the invention relates to methods of inhibiting the activity of
 20 HCV polymerase comprising the step of treating a sample suspected of containing

- 5 HCV with a composition of the invention.

Compositions of the invention may act as inhibitors of HCV polymerase , as intermediates for such inhibitors or have other utilities as described below. The inhibitors will bind to locations on the surface or in a cavity of HCV polymerase having a geometry unique to HCV polymerase . Compositions binding HCV

- 10 polymerase may bind with varying degrees of reversibility. Those compounds binding substantially irreversibly are ideal candidates for use in this method of the invention. Once labeled, the substantially irreversibly binding compositions are useful as probes for the detection of HCV polymerase . Accordingly, the invention relates to methods of detecting HCV polymerase in a sample suspected of containing
- 15 HCV polymerase comprising the steps of: treating a sample suspected of containing HCV polymerase with a composition comprising a compound of the invention bound to a label; and observing the effect of the sample on the activity of the label. Suitable labels are well known in the diagnostics field and include stable free radicals, fluorophores, radioisotopes, enzymes, chemiluminescent groups and chromogens.
- 20 The compounds herein are labeled in conventional fashion using functional groups such as hydroxyl, carboxyl, sulfhydryl or amino.

- Within the context of the invention, samples suspected of containing HCV polymerase include natural or man-made materials such as living organisms; tissue or cell cultures; biological samples such as biological material samples (blood, serum,
- 25 urine, cerebrospinal fluid, tears, sputum, saliva, tissue samples, and the like); laboratory samples; food, water, or air samples; bioproduct samples such as extracts of cells, particularly recombinant cells synthesizing a desired glycoprotein; and the like. Typically the sample will be suspected of containing an organism which produces HCV polymerase , frequently a pathogenic organism such as HCV.
- 30 Samples can be contained in any medium including water and organic solvent\water mixtures. Samples include living organisms such as humans, and man made materials such as cell cultures.

The treating step of the invention comprises adding the composition of the invention to the sample or it comprises adding a precursor of the composition to the



- 5 sample. The addition step comprises any method of administration as described above.

If desired, the activity of HCV polymerase after application of the composition can be observed by any method including direct and indirect methods of detecting HCV polymerase activity. Quantitative, qualitative, and semiquantitative methods of
10 determining HCV polymerase activity are all contemplated. Typically one of the screening methods described above are applied, however, any other method such as observation of the physiological properties of a living organism are also applicable.

Organisms that contain HCV polymerase include the HCV virus. The compounds of this invention are useful in the treatment or prophylaxis of HCV
15 infections in animals or in man.

However, in screening compounds capable of inhibiting human immunodeficiency viruses, it should be kept in mind that the results of enzyme assays may not correlate with cell culture assays. Thus, a cell based assay should be the primary screening tool.

20 Screens for HCV polymerase Inhibitors.

Compositions of the invention are screened for inhibitory activity against HCV polymerase by any of the conventional techniques for evaluating enzyme activity. Within the context of the invention, typically compositions are first screened for inhibition of HCV polymerase *in vitro* and compositions showing inhibitory
25 activity are then screened for activity *in vivo*. Compositions having *in vitro* K_i (inhibitory constants) of less than about 5×10^{-6} M, typically less than about 1×10^{-7} M and preferably less than about 5×10^{-8} M are preferred for *in vivo* use.

Useful *in vitro* screens have been described in detail and will not be elaborated here. However, the examples describe suitable *in vitro* assays.

30 Pharmaceutical Formulations

The compounds of this invention are formulated with conventional carriers and excipients, which will be selected in accord with ordinary practice. Tablets will contain excipients, glidants, fillers, binders and the like. Aqueous formulations are



- 5 prepared in sterile form, and when intended for delivery by other than oral administration generally will be isotonic. All formulations will optionally contain excipients such as those set forth in the "Handbook of Pharmaceutical Excipients" (1986). Excipients include ascorbic acid and other antioxidants, chelating agents such as EDTA, carbohydrates such as dextran, hydroxyalkylcellulose, hydroxyalkylmethylcellulose, stearic acid and the like. The pH of the formulations ranges from about 3 to about 11, but is ordinarily about 7 to 10.

While it is possible for the active ingredients to be administered alone it may be preferable to present them as pharmaceutical formulations. The formulations, both for veterinary and for human use, of the invention comprise at least one active ingredient, as above defined, together with one or more acceptable carriers therefore and optionally other therapeutic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and physiologically innocuous to the recipient thereof.

The formulations include those suitable for the foregoing administration routes. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. Techniques and formulations generally are found in Remington's Pharmaceutical Sciences (Mack Publishing Co., Easton, PA). Such methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be administered as a bolus, electuary or paste.

A tablet is made by compression or molding, optionally with one or more



5 accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered active ingredient moistened with an inert liquid
10 diluent. The tablets may optionally be coated or scored and optionally are formulated so as to provide slow or controlled release of the active ingredient therefrom.

For infections of the eye or other external tissues e.g. mouth and skin, the formulations are preferably applied as a topical ointment or cream containing the active ingredient(s) in an amount of, for example, 0.075 to 20% w/w (including active
15 ingredient(s) in a range between 0.1% and 20% in increments of 0.1% w/w such as 0.6% w/w, 0.7% w/w, etc.), preferably 0.2 to 15% w/w and most preferably 0.5 to 10% w/w. When formulated in an ointment, the active ingredients may be employed with either a paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base.

20 If desired, the aqueous phase of the cream base may include, for example, at least 30% w/w of a polyhydric alcohol, i.e. an alcohol having two or more hydroxyl groups such as propylene glycol, butane 1,3-diol, mannitol, sorbitol, glycerol and polyethylene glycol (including PEG 400) and mixtures thereof. The topical formulations may desirably include a compound which enhances absorption or
25 penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethyl sulphoxide and related analogs.

The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner. While the phase may comprise merely an
30 emulsifier (otherwise known as an emulgent), it desirably comprises a mixture of at least one emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make up the so-called emulsifying wax, and

L

5 the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations.

Emulgents and emulsion stabilizers suitable for use in the formulation of the invention include Tween® 60, Span® 80, cetostearyl alcohol, benzyl alcohol, myristyl alcohol, glyceryl mono-stearate and sodium lauryl sulfate.

10 The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties. The cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids,
15 isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters known as Crodamol CAP may be used, the last three being preferred esters. These may be used alone or in combination depending on the properties required. Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils are used.

20 Pharmaceutical formulations according to the present invention comprise a combination according to the invention together with one or more pharmaceutically acceptable carriers or excipients and optionally other therapeutic agents. Pharmaceutical formulations containing the active ingredient may be in any form suitable for the intended method of administration. When used for oral use for
25 example, tablets, troches, lozenges, aqueous or oil suspensions, dispersible powders or granules, emulsions, hard or soft capsules, syrups or elixirs may be prepared. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents including sweetening agents, flavoring agents,
30 coloring agents and preserving agents, in order to provide a palatable preparation. Tablets containing the active ingredient in admixture with non-toxic pharmaceutically acceptable excipient which are suitable for manufacture of tablets are acceptable. These excipients may be, for example, inert diluents, such as calcium or sodium carbonate, lactose, calcium or sodium phosphate; granulating and disintegrating

2

5 agents, such as maize starch, or alginic acid; binding agents, such as starch, gelatin or acacia; and lubricating agents, such as magnesium stearate, stearic acid or talc.

Tablets may be uncoated or may be coated by known techniques including microencapsulation to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time
10 delay material such as glyceryl monostearate or glyceryl distearate alone or with a wax may be employed.

Formulations for oral use may be also presented as hard gelatin capsules where the active ingredient is mixed with an inert solid diluent, for example calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed
15 with water or an oil medium, such as peanut oil, liquid paraffin or olive oil.

Aqueous suspensions of the invention contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients include a suspending agent, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropyl methylcellulose, sodium alginate,
20 polyvinylpyrrolidone, gum tragacanth and gum acacia, and dispersing or wetting agents such as a naturally-occurring phosphatide (e.g., lecithin), a condensation product of an alkylene oxide with a fatty acid (e.g., polyoxyethylene stearate), a condensation product of ethylene oxide with a long chain aliphatic alcohol (e.g., heptadecaethyleneoxycetanol), a condensation product of ethylene oxide with a partial
25 ester derived from a fatty acid and a hexitol anhydride (e.g., polyoxyethylene sorbitan monooleate). The aqueous suspension may also contain one or more preservatives such as ethyl or n-propyl p-hydroxy-benzoate, one or more coloring agents, one or more flavoring agents and one or more sweetening agents, such as sucrose or saccharin.

30 Oil suspensions may be formulated by suspending the active ingredient in a vegetable oil, such as arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oral suspensions may contain a thickening agent, such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation.



- 5 These compositions may be preserved by the addition of an antioxidant such as ascorbic acid.

Dispersible powders and granules of the invention suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent, and one or more
10 preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those disclosed above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or
15 arachis oil, a mineral oil, such as liquid paraffin, or a mixture of these. Suitable emulsifying agents include naturally-occurring gums, such as gum acacia and gum tragacanth, naturally-occurring phosphatides, such as soybean lecithin, esters or partial esters derived from fatty acids and hexitol anhydrides, such as sorbitan monooleate, and condensation products of these partial esters with ethylene oxide,
20 such as polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening and flavoring agents. Syrups and elixirs may be formulated with sweetening agents, such as glycerol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, a flavoring or a coloring agent.

The pharmaceutical compositions of the invention may be in the form of a
25 sterile injectable preparation, such as a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, such
30 as a solution in 1,3-butane-diol or prepared as a lyophilized powder. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile fixed oils may conventionally be employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In

L

- 5 addition, fatty acids such as oleic acid may likewise be used in the preparation of injectables.

The amount of active ingredient that may be combined with the carrier material to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a time-release formulation
10 intended for oral administration to humans may contain approximately 1 to 1000 mg of active material compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95% of the total compositions (weight:weight). The pharmaceutical composition can be prepared to provide easily measurable amounts for administration. For example, an aqueous solution intended
15 for intravenous infusion may contain from about 3 to 500 μg of the active ingredient per milliliter of solution in order that infusion of a suitable volume at a rate of about 30 mL/hr can occur.

Formulations suitable for topical administration to the eye also include eye drops wherein the active ingredient is dissolved or suspended in a suitable carrier,
20 especially an aqueous solvent for the active ingredient. The active ingredient is preferably present in such formulations in a concentration of 0.5 to 20%, advantageously 0.5 to 10%, and particularly about 1.5% w/w.

Formulations suitable for topical administration in the mouth include lozenges comprising the active ingredient in a flavored basis, usually sucrose and acacia or
25 tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

Formulations for rectal administration may be presented as a suppository with a suitable base comprising for example cocoa butter or a salicylate.

30 Formulations suitable for intrapulmonary or nasal administration have a particle size for example in the range of 0.1 to 500 microns, such as 0.5, 1, 30, 35 etc., which is administered by rapid inhalation through the nasal passage or by inhalation through the mouth so as to reach the alveolar sacs. Suitable formulations include aqueous or oily solutions of the active ingredient. Formulations suitable for aerosol or



5 dry powder administration may be prepared according to conventional methods and may be delivered with other therapeutic agents such as compounds heretofore used in the treatment or prophylaxis of HCV infections as described below.

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in
10 addition to the active ingredient such carriers as are known in the art to be appropriate.

Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the
15 intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents.

The formulations are presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for
20 injection, immediately prior to use. Extemporaneous injection solutions and suspensions are prepared from sterile powders, granules and tablets of the kind previously described. Preferred unit dosage formulations are those containing a daily dose or unit daily sub-dose, as herein above recited, or an appropriate fraction thereof, of the active ingredient.

25 It should be understood that in addition to the ingredients particularly mentioned above the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavoring agents.

The invention further provides veterinary compositions comprising at least
30 one active ingredient as above defined together with a veterinary carrier therefore.

Veterinary carriers are materials useful for the purpose of administering the composition and may be solid, liquid or gaseous materials which are otherwise inert or acceptable in the veterinary art and are compatible with the active ingredient. These veterinary compositions may be administered orally, parenterally or by any



5 other desired route.

Compounds of the invention are used to provide controlled release pharmaceutical formulations containing as active ingredient one or more compounds of the invention ("controlled release formulations") in which the release of the active ingredient are controlled and regulated to allow less frequency dosing or to improve
10 the pharmacokinetic or toxicity profile of a given active ingredient.

Effective dose of active ingredient depends at least on the nature of the condition being treated, toxicity, whether the compound is being used prophylactically (lower doses) or against an active viral infection, the method of delivery, and the pharmaceutical formulation, and will be determined by the clinician
15 using conventional dose escalation studies. It can be expected to be from about 0.0001 to about 100 mg/kg body weight per day; typically, from about 0.01 to about 10 mg/kg body weight per day; more typically, from about .01 to about 5 mg/kg body weight per day; most typically, from about .05 to about 0.5 mg/kg body weight per day. For example, the daily candidate dose for an adult human of approximately 70
20 kg body weight will range from 1 mg to 1000 mg, preferably between 5 mg and 500 mg, and may take the form of single or multiple doses.

Routes of Administration

One or more compounds of the invention (herein referred to as the active ingredients) are administered by any route appropriate to the condition to be treated.
25 Suitable routes include oral, rectal, nasal, topical (including buccal and sublingual), vaginal and parenteral (including subcutaneous, intramuscular, intravenous, intradermal, intrathecal and epidural), and the like. It will be appreciated that the preferred route may vary with for example the condition of the recipient. An advantage of the compounds of this invention is that they are orally bioavailable and
30 can be dosed orally.

Combination Therapy

Combinations of the compounds of Formula I-III and Formula IV-VI are typically selected based on the condition to be treated, cross-reactivities of ingredients



5 and pharmaco-properties of the combination. For example, when treating an infection (e.g., HCV), the compositions of the invention are combined with other active therapeutic agents (such as those described herein).

Compositions of the invention are also used in combination with one or more other active ingredients. Preferably, the other active therapeutic ingredients or agents
10 are interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, other anti-fibrotic agents, endothelin antagonists, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A
15 inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers or other drugs for treating HCV; or mixtures thereof.

More specifically, one or more compounds of the present invention may be combined with one or more compounds selected from the group consisting of

1) interferons, *e.g.*, pegylated rIFN-alpha 2b (PEG-Intron), pegylated rIFN-alpha 2a (Pegasys), rIFN-alpha 2b (Intron A), rIFN-alpha 2a (Roferon-A), interferon alpha (MOR-22, OPC-18, Alfaferone, Alfanative, Multiferon, subalin), interferon alfacon-1 (Infergen), interferon alpha-n1 (Wellferon), interferon alpha-n3 (Alferon), interferon-beta (Avonex, DL-8234), interferon-omega (omega DUROS, Biomed 510), albinterferon alpha-2b (Albuferon), IFN alpha XL, BLX-883 (Locteron), DA-3021,
25 glycosylated interferon alpha-2b (AVI-005), PEG-Infergen, PEGylated interferon lambda (PEGylated IL-29), and belerofon,

2) ribavirin and its analogs, *e.g.*, ribavirin (Rebetol, Copegus), and taribavirin (Viramidine),

3) HCV NS3 protease inhibitors, *e.g.*, boceprevir (SCH-503034, SCH-7),
30 telaprevir (VX-950), VX-813, TMC-435 (TMC435350), ABT-450, BI-201335, BI-1230, MK-7009, SCH-900518, VBY-376, VX-500, GS-9256, GS-9451, BMS-790052, BMS-605339, PHX-1766, AS-101, YH-5258, YH5530, YH5531, and ITMN-191 (R-7227),



- 5 4) alpha-glucosidase 1 inhibitors, *e.g.*, celgosivir (MX-3253), Miglitol, and UT-231B,
- 5) hepatoprotectants, *e.g.*, emericasan (IDN-6556), ME-3738, GS-9450 (LB-84451), silibinin, and MitoQ,
- 6) nucleoside or nucleotide inhibitors of HCV NS5B polymerase, *e.g.*, R1626,
- 10 R7128 (R4048), IDX184, IDX-102, PSI-7851, BCX-4678, valopicitabine (NM-283), and MK-0608,
- 7) non-nucleoside inhibitors of HCV NS5B polymerase, *e.g.*, filibuvir (PF-868554), ABT-333, ABT-072, BI-207127, VCH-759, VCH-916, JTK-652, MK-3281, VBY-708, VCH-222, A848837, ANA-598, GL60667, GL59728, A-63890, A-48773,
- 15 A-48547, BC-2329, VCH-796 (nesbuvir), GSK625433, BILN-1941, XTL-2125, and GS-9190,
- 8) HCV NS5A inhibitors, *e.g.*, AZD-2836 (A-831), AZD-7295 (A-689), and BMS-790052,
- 9) TLR-7 agonists, *e.g.*, imiquimod, 852A, GS-9524, ANA-773, ANA-975,
- 20 AZD-8848 (DSP-3025), PF-04878691, and SM-360320,
- 10) cyclophilin inhibitors, *e.g.*, DEBIO-025, SCY-635, and NIM811,
- 11) HCV IRES inhibitors, *e.g.*, MCI-067,
- 12) pharmacokinetic enhancers, *e.g.*, BAS-100, SPI-452, PF-4194477, TMC-41629, GS-9350, GS-9585, and roxythromycin,
- 25 13) other drugs for treating HCV, *e.g.*, thymosin alpha 1 (Zadaxin), nitazoxanide (Alinea, NTZ), BIVN-401 (virostat), PYN-17 (altirex), KPE02003002, actilon (CPG-10101), GS-9525, KRN-7000, civacir, GI-5005, XTL-6865, BIT225, PTX-111, ITX2865, TT-033i, ANA 971, NOV-205, tarvacin, EHC-18, VGX-410C, EMZ-702, AVI 4065, BMS-650032, BMS-791325, Bavituximab, MDX-1106 (ONO-
- 30 4538), Oglufanide, FK-788, and VX-497 (merimepodib)
- 14) mevalonate decarboxylase antagonists, *e.g.*, statins, HMGCoA synthase inhibitors (*e.g.*, hymeglusin), squalene synthesis inhibitors (*e.g.*, zaragozic acid);
- 15) angiotensin II receptor antagonists, *e.g.*, losartan, irbesartan, olmesartan, candesartan, valsartan, telmisartan, eprosartan;



5 16) angiotensin-converting enzyme inhibitors, e.g., captopril, zofenopril, enalapril, ramipril, quinapril, perindopril, lisinopril, benazepril, fosinopril;

17) other anti-fibrotic agents, e.g., amiloride and

18) endothelin antagonists, e.g. bosentan and ambrisentan.

10 In yet another embodiment, the present application discloses pharmaceutical compositions comprising a compound of the present invention, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, in combination with at least one additional therapeutic agent, and a pharmaceutically acceptable carrier or excipient.

15 According to the present invention, the therapeutic agent used in combination with the compound or composition of the present invention can be any agent having a therapeutic effect when used in combination with the compound of the present invention. For example, the therapeutic agent used in combination with the compound or composition of the present invention can be interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of
20 the renin-angiotensin system, other anti-fibrotic agents, endothelin antagonists, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophillin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers or other drugs for treating HCV; or mixtures thereof.

25 More specifically, compositions of one or more compounds of the present invention may be combined with one or more compounds selected from the group consisting of

1) interferons, e.g., pegylated rIFN-alpha 2b (PEG-Intron), pegylated rIFN-alpha 2a (Pegasys), rIFN-alpha 2b (Intron A), rIFN-alpha 2a (Roferon-A), interferon
30 alpha (MOR-22, OPC-18, Alfaferone, Alfanaive, Multiferon, subalin), interferon alfacon-1 (Infergen), interferon alpha-n1 (Wellferon), interferon alpha-n3 (Alferon), interferon-beta (Avonex, DL-8234), interferon-omega (omega DUROS, Biomed 510), albinterferon alpha-2b (Albuferon), IFN alpha XL, BLX-883 (Locteron), DA-3021,



- 5 glycosylated interferon alpha-2b (AVI-005), PEG-Infergen, PEGylated interferon
lambda (PEGylated IL-29), and belerofon,
- 2) ribavirin and its analogs, *e.g.*, ribavirin (Rebetol, Copegus), and taribavirin
(Viramidine),
- 3) HCV NS3 protease inhibitors, *e.g.*, boceprevir (SCH-503034 , SCH-7),
10 telaprevir (VX-950), VX-813, TMC-435 (TMC435350), ABT-450, BI-201335, BI-
1230, MK-7009, SCH-900518, VBY-376, VX-500, GS-9256, GS-9451, BMS-
790052, BMS-605339, PHX-1766, AS-101, YH-5258, YH5530, YH5531, and
ITMN-191 (R-7227),
- 4) alpha-glucosidase 1 inhibitors, *e.g.*, celgosivir (MX-3253), Miglitol, and
15 UT-231B,
- 5) hepatoprotectants, *e.g.*, emericasan (IDN-6556), ME-3738, GS-9450 (LB-
84451), silibinin, and MitoQ,
- 6) nucleoside or nucleotide inhibitors of HCV NS5B polymerase, *e.g.*, R1626,
R7128 (R4048), IDX184, IDX-102, PSI-7851, BCX-4678, valopicitabine (NM-283),
20 and MK-0608,
- 7) non-nucleoside inhibitors of HCV NS5B polymerase, *e.g.*, filibuvir (PF-
868554), ABT-333, ABT-072, BI-207127, VCH-759, VCH-916, JTK-652, MK-3281,
VBY-708, VCH-222, A848837, ANA-598, GL60667, GL59728, A-63890, A-48773,
A-48547, BC-2329, VCH-796 (nesbuvir), GSK625433, BILN-1941, XTL-2125, and
25 GS-9190,
- 8) HCV NS5A inhibitors, *e.g.*, AZD-2836 (A-831), AZD-7295 (A-689), and
BMS-790052,
- 9) TLR-7 agonists, *e.g.*, imiquimod, 852A, GS-9524, ANA-773, ANA-975,
AZD-8848 (DSP-3025), PF-04878691, and SM-360320,
- 30 10) cyclophilin inhibitors, *e.g.*, DEBIO-025, SCY-635, and NIM811,
11) HCV IRES inhibitors, *e.g.*, MCI-067,
12) pharmacokinetic enhancers, *e.g.*, BAS-100, SPI-452, PF-4194477, TMC-
41629, GS-9350, GS-9585, and roxythromycin,



- 5 13) other drugs for treating HCV, *e.g.*, thymosin alpha 1 (Zadaxin),
nitazoxanide (Alinea, NTZ), BIVN-401 (virostat), PYN-17 (altirex), KPE02003002,
actilon (CPG-10101), GS-9525, KRN-7000, civacir, GI-5005, XTL-6865, BIT225,
PTX-111, ITX2865, TT-033i, ANA 971, NOV-205, tarvacin, EHC-18, VGX-410C,
EMZ-702, AVI 4065, BMS-650032, BMS-791325, Bavituximab, MDX-1106 (ONO-
10 4538), Oglufanide, FK-788, and VX-497 (merimepodib)
- 14) mevalonate decarboxylase antagonists, *e.g.*, statins, HMGCoA synthase
inhibitors (*e.g.*, hymeglusin), squalene synthesis inhibitors (*e.g.*, zaragozic acid);
- 15) angiotensin II receptor antagonists, *e.g.*, losartan, irbesartan, olmesartan,
candesartan, valsartan, telmisartan, eprosartan;
- 15 16) angiotensin-converting enzyme inhibitors, *e.g.*, captopril, zofenopril,
enalapril, ramipril, quinapril, perindopril, lisinopril, benazepril, fosinopril;
- 17) other anti-fibrotic agents, *e.g.*, amiloride and
- 18) endothelin antagonists, *e.g.* bosentan and ambrisentan.

- In yet another embodiment, the present application provides a combination
20 pharmaceutical agent comprising:
- a) a first pharmaceutical composition comprising a compound of the
present invention, or a pharmaceutically acceptable salt, solvate, or ester thereof; and
- b) a second pharmaceutical composition comprising at least one
additional therapeutic agent selected from the group consisting of HIV protease
25 inhibiting compounds, HIV non-nucleoside inhibitors of reverse transcriptase, HIV
nucleoside inhibitors of reverse transcriptase, HIV nucleotide inhibitors of reverse
transcriptase, HIV integrase inhibitors, gp41 inhibitors, CXCR4 inhibitors, gp120
inhibitors, CCR5 inhibitors, interferons, ribavirin analogs, NS3 protease inhibitors,
NS5a inhibitors, alpha-glucosidase 1 inhibitors, cyclophilin inhibitors,
30 hepatoprotectants, non-nucleoside inhibitors of HCV, and other drugs for treating
HCV, and combinations thereof.

Combinations of the compounds of Formula I-III and Formula IV-VI and
additional active therapeutic agents may be selected to treat patients infected with
HCV and other conditions such as HIV infections. Accordingly, the compounds of



5 Formula I-III and Formula IV-VI may be combined with one or more compounds
useful in treating HIV, for example HIV protease inhibiting compounds, HIV non-
nucleoside inhibitors of reverse transcriptase, HIV nucleoside inhibitors of reverse
transcriptase, HIV nucleotide inhibitors of reverse transcriptase, HIV integrase
inhibitors, gp41 inhibitors, CXCR4 inhibitors, gp120 inhibitors, CCR5 inhibitors,
10 interferons, ribavirin analogs, NS3 protease inhibitors, NS5a inhibitors, alpha-
glucosidase 1 inhibitors, cyclophilin inhibitors, hepatoprotectants, non-nucleoside
inhibitors of HCV, and other drugs for treating HCV.

More specifically, one or more compounds of the present invention may be
combined with one or more compounds selected from the group consisting of 1) HIV
15 protease inhibitors, *e.g.*, amprenavir, atazanavir, fosamprenavir, indinavir, lopinavir,
ritonavir, lopinavir + ritonavir, nelfinavir, saquinavir, tipranavir, brecanavir,
darunavir, TMC-126, TMC-114, mozenavir (DMP-450), JE-2147 (AG1776),
AG1859, DG35, L-756423, RO0334649, KNI-272, DPC-681, DPC-684, and
GW640385X, DG17, PPL-100, 2) a HIV non-nucleoside inhibitor of reverse
20 transcriptase, *e.g.*, capravirine, emvirine, delaviridine, efavirenz, nevirapine, (+)
calanolide A, etravirine, GW5634, DPC-083, DPC-961, DPC-963, MIV-150, and
TMC-120, TMC-278 (rilpivirine), efavirenz, BILR 355 BS, VRX 840773, UK-
453,061, RDEA806, 3) a HIV nucleoside inhibitor of reverse transcriptase, *e.g.*,
zidovudine, emtricitabine, didanosine, stavudine, zalcitabine, lamivudine, abacavir,
25 amdoxovir, elvucitabine, alovudine, MIV-210, racivir ($\pm$ -FTC), D-d4FC,
emtricitabine, phosphazide, fozivudine tidoxil, fosaltivudine tidoxil, apricitibine
(AVX754), amdoxovir, KP-1461, abacavir + lamivudine, abacavir + lamivudine +
zidovudine, zidovudine + lamivudine, 4) a HIV nucleotide inhibitor of reverse
transcriptase, *e.g.*, tenofovir, tenofovir disoproxil fumarate + emtricitabine, tenofovir
30 disoproxil fumarate + emtricitabine + efavirenz, and adefovir, 5) a HIV integrase
inhibitor, *e.g.*, curcumin, derivatives of curcumin, chicoric acid, derivatives of
chicoric acid, 3,5-dicaffeoylquinic acid, derivatives of 3,5-dicaffeoylquinic acid,
aurintricarboxylic acid, derivatives of aurintricarboxylic acid, caffeic acid phenethyl
ester, derivatives of caffeic acid phenethyl ester, tyrphostin, derivatives of tyrphostin,



- 5 quercetin, derivatives of quercetin, S-1360, zintevir (AR-177), L-870812, and L-870810, MK-0518 (raltegravir), BMS-707035, MK-2048, BA-011, BMS-538158, GSK364735C, 6) a gp41 inhibitor, *e.g.*, enfuvirtide, sifuvirtide, FB006M, TRI-1144, SPC3, DES6, Locus gp41, CovX, and REP 9, 7) a CXCR4 inhibitor, *e.g.*, AMD-070, 8) an entry inhibitor, *e.g.*, SP01A, TNX-355, 9) a gp120 inhibitor, *e.g.*, BMS-488043 and BlockAide/CR, 10) a G6PD and NADH-oxidase inhibitor, *e.g.*, immunitin, 10) a CCR5 inhibitor, *e.g.*, aplaviroc, vicriviroc, INCB9471, PRO-140, INCB15050, PF-232798, CCR5mAb004, and maraviroc, 11) an interferon, *e.g.*, pegylated rIFN-alpha 2b, pegylated rIFN-alpha 2a, rIFN-alpha 2b, IFN alpha-2b XL, rIFN-alpha 2a, consensus IFN alpha, infergen, rebif, locteron, AVI-005, PEG-infergen, pegylated
- 15 IFN-beta, oral interferon alpha, feron, reaferon, intermax alpha, r-IFN-beta, infergen + actimmune, IFN-omega with DUROS, and albuferon, 12) ribavirin analogs, *e.g.*, rebetol, copegus, VX-497, and viramidine (taribavirin) 13) NS5a inhibitors, *e.g.*, A-831, A-689 and BMS-790052, 14) NS5b polymerase inhibitors, *e.g.*, NM-283, valopicitabine, R1626, PSI-6130 (R1656), IDX184, PSI-7851, HCV-796, BILB 1941, 20 MK-0608, NM-107, R7128, VCH-759, PF-868554, GSK625433, and XTL-2125, 15) NS3 protease inhibitors, *e.g.*, SCH-503034 (SCH-7), VX-950 (Telaprevir), ITMN-191, and BILN-2065, 16) alpha-glucosidase 1 inhibitors, *e.g.*, MX-3253 (celgosivir) and UT-231B, 17) hepatoprotectants, *e.g.*, IDN-6556, ME 3738, MitoQ, and LB-84451, 18) non-nucleoside inhibitors of HCV, *e.g.*, benzimidazole derivatives, benzo-
- 25 1,2,4-thiadiazine derivatives, and phenylalanine derivatives, 19) other drugs for treating HCV, *e.g.*, zadaxin, nitazoxanide (alinea), BIVN-401 (virostat), DEBIO-025, VGX-410C, EMZ-702, AVI 4065, bavituximab, oglufanide, PYN-17, KPE02003002, actilon (CPG-10101), KRN-7000, civacir, GI-5005, ANA-975, XTL-6865, ANA 971, NOV-205, tarvacin, EHC-18, and NIM811, 19) pharmacokinetic enhancers, *e.g.*,
- 30 BAS-100 and SPI452, 20) RNAse H inhibitors, *e.g.*, ODN-93 and ODN-112, 21) other anti-HIV agents, *e.g.*, VGV-1, PA-457 (bevirimat), ampligen, HRG214, cytolin, polymun, VGX-410, KD247, AMZ 0026, CYT 99007, A-221 HIV, BAY 50-4798, MDX010 (iplimumab), PBS119, ALG889, and PA-1050040.

5 It is also possible to combine any compound of the invention with one or more other active therapeutic agents in a unitary dosage form for simultaneous or sequential administration to a patient. The combination therapy may be administered as a simultaneous or sequential regimen. When administered sequentially, the combination may be administered in two or more administrations.

10 Co-administration of a compound of the invention with one or more other active therapeutic agents generally refers to simultaneous or sequential administration of a compound of the invention and one or more other active therapeutic agents, such that therapeutically effective amounts of the compound of the invention and one or more other active therapeutic agents are both present in the body of the patient.

15 Co-administration includes administration of unit dosages of the compounds of the invention before or after administration of unit dosages of one or more other active therapeutic agents, for example, administration of the compounds of the invention within seconds, minutes, or hours of the administration of one or more other active therapeutic agents. For example, a unit dose of a compound of the invention
20 can be administered first, followed within seconds or minutes by administration of a unit dose of one or more other active therapeutic agents. Alternatively, a unit dose of one or more other therapeutic agents can be administered first, followed by administration of a unit dose of a compound of the invention within seconds or minutes. In some cases, it may be desirable to administer a unit dose of a compound
25 of the invention first, followed, after a period of hours (e.g., 1-12 hours), by administration of a unit dose of one or more other active therapeutic agents. In other cases, it may be desirable to administer a unit dose of one or more other active therapeutic agents first, followed, after a period of hours (e.g., 1-12 hours), by administration of a unit dose of a compound of the invention.

30 The combination therapy may provide "synergy" and "synergistic", i.e. the effect achieved when the active ingredients used together is greater than the sum of the effects that results from using the compounds separately. A synergistic effect may be attained when the active ingredients are: (1) co-formulated and administered or delivered simultaneously in a combined formulation; (2) delivered by alternation or in



5 parallel as separate formulations; or (3) by some other regimen. When delivered in alternation therapy, a synergistic effect may be attained when the compounds are administered or delivered sequentially, e.g. in separate tablets, pills or capsules, or by different injections in separate syringes. In general, during alternation therapy, an effective dosage of each active ingredient is administered sequentially, i.e. serially,
10 whereas in combination therapy, effective dosages of two or more active ingredients are administered together. A synergistic anti-viral effect denotes an antiviral effect which is greater than the predicted purely additive effects of the individual compounds of the combination.

In still yet another embodiment, the present application provides for methods
15 of inhibiting HCV polymerase in a cell, comprising: contacting a cell infected with HCV with an effective amount of a compound of Formula I-III and Formula IV-VI, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, whereby HCV polymerase is inhibited.

In still yet another embodiment, the present application provides for methods
20 of inhibiting HCV polymerase in a cell, comprising: contacting a cell infected with HCV with an effective amount of a compound of Formula I-III and Formula IV-VI, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and at least one additional active therapeutic agent, whereby HCV polymerase is inhibited.

In still yet another embodiment, the present application provides for methods
25 of inhibiting HCV polymerase in a cell, comprising: contacting a cell infected with HCV with an effective amount of a compound of Formula I-III and Formula IV-VI, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and at least one additional active therapeutic agent selected from the group consisting of one or more interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors,
30 alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, other anti-fibrotic agents, endothelin antagonists, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7



- 5 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers and other drugs for treating HCV; or mixtures thereof.

In still yet another embodiment, the present application provides for methods of treating HCV in a patient, comprising: administering to the patient a therapeutically effective amount of a compound of Formula I-III and Formula IV-VI, or a
10 pharmaceutically acceptable salt, solvate, and/or ester thereof.

In still yet another embodiment, the present application provides for methods of treating HCV in a patient, comprising: administering to the patient a therapeutically effective amount of a compound of Formula I-III and Formula IV-VI, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and at least one
15 additional active therapeutic agent, whereby HCV polymerase is inhibited.

In still yet another embodiment, the present application provides for methods of treating HCV in a patient, comprising: administering to the patient a therapeutically effective amount of a compound of Formula I-III and Formula IV-VI, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and at least one
20 additional active therapeutic agent selected from the group consisting of one or more interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, other anti-fibrotic agents, endothelin antagonists, nucleoside or nucleotide inhibitors of HCV NS5B polymerase,
25 non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers and other drugs for treating HCV; or mixtures thereof.

In still yet another embodiment, the present application provides for the use of a compound of the present invention, or a pharmaceutically acceptable salt, solvate,
30 and/or ester thereof, for the preparation of a medicament for treating an HCV infection in a patient.

Metabolites of the Compounds of the Invention

Also falling within the scope of this invention are the *in vivo* metabolic products of the compounds described herein, to the extent such products are novel and



5 unobvious over the prior art. Such products may result for example from the
oxidation, reduction, hydrolysis, amidation, esterification and the like of the
administered compound, primarily due to enzymatic processes. Accordingly, the
invention includes novel and unobvious compounds produced by a process
10 comprising contacting a compound of this invention with a mammal for a period of
time sufficient to yield a metabolic product thereof. Such products typically are
identified by preparing a radiolabelled (e.g. ^{14}C or ^3H) compound of the invention,
administering it parenterally in a detectable dose (e.g. greater than about 0.5 mg/kg) to
an animal such as rat, mouse, guinea pig, monkey, or to man, allowing sufficient time
15 for metabolism to occur (typically about 30 seconds to 30 hours) and isolating its
conversion products from the urine, blood or other biological samples. These
products are easily isolated since they are labeled (others are isolated by the use of
antibodies capable of binding epitopes surviving in the metabolite). The metabolite
structures are determined in conventional fashion, e.g. by MS or NMR analysis. In
20 general, analysis of metabolites is done in the same way as conventional drug
metabolism studies well-known to those skilled in the art. The conversion products,
so long as they are not otherwise found *in vivo*, are useful in diagnostic assays for
therapeutic dosing of the compounds of the invention even if they possess no HCV
polymerase inhibitory activity of their own.

Recipes and methods for determining stability of compounds in surrogate
25 gastrointestinal secretions are known. Compounds are defined herein as stable in the
gastrointestinal tract where less than about 50 mole percent of the protected groups
are deprotected in surrogate intestinal or gastric juice upon incubation for 1 hour at
37°C. Simply because the compounds are stable to the gastrointestinal tract does not
mean that they cannot be hydrolyzed *in vivo*. The prodrugs of the invention typically
30 will be stable in the digestive system but may be substantially hydrolyzed to the
parental drug in the digestive lumen, liver or other metabolic organ, or within cells in
general.

Examples

35



- 5 Certain abbreviations and acronyms are used in describing the experimental details. Although most of these would be understood by one skilled in the art, Table 1 contains a list of many of these abbreviations and acronyms.

Table 1. List of abbreviations and acronyms.

Abbreviation	Meaning
Ac ₂ O	acetic anhydride
AIBN	2,2'-azobis(2-methylpropionitrile)
Bn	benzyl
BnBr	benzylbromide
BSA	bis(trimethylsilyl)acetamide
BzCl	benzoyl chloride
CDI	carbonyl diimidazole
DABCO	1,4-diazabicyclo[2.2.2]octane
DBN	1,5-diazabicyclo[4.3.0]non-5-ene
DDQ	2,3-dichloro-5,6-dicyano-1,4-benzoquinone
DBU	1,5-diazabicyclo[5.4.0]undec-5-ene
DCA	dichloroacetamide
DCC	dicyclohexylcarbodiimide
DCM	dichloromethane
DMAP	4-dimethylaminopyridine
DME	1,2-dimethoxyethane
DMTCI	dimethoxytrityl chloride
DMSO	dimethylsulfoxide
DMTr	4, 4'-dimethoxytrityl
DMF	dimethylformamide
EtOAc	ethyl acetate
ESI	electrospray ionization
HMDS	hexamethyldisilazane

HPLC	High pressure liquid chromatography
LDA	lithium diisopropylamide
LRMS	low resolution mass spectrum
MCPBA	meta-chloroperbenzoic acid
MeCN	acetonitrile
MeOH	methanol
MMTC	mono methoxytrityl chloride
m/z or m/e	mass to charge ratio
MH ⁺	mass plus 1
MH ⁻	mass minus 1
MsOH	methanesulfonic acid
MS or ms	mass spectrum
NBS	N-bromosuccinimide
Ph	phenyl
rt or r.t.	room temperature
TBAF	tetrabutylammonium fluoride
TMSCl	chlorotrimethylsilane
TMSBr	bromotrimethylsilane
TMSI	iodotrimethylsilane
TMSOTf	(trimethylsilyl)trifluoromethylsulfonate
TEA	triethylamine
TBA	tributylamine
TBAP	tributylammonium pyrophosphate
TBSCl	t-butyltrimethylsilyl chloride
TEAB	triethylammonium bicarbonate
TFA	trifluoroacetic acid
TLC or tlc	thin layer chromatography
Tr	triphenylmethyl
Tol	4-methylbenzoyl




Turbo Grignard	1:1 mixture of isopropylmagnesium chloride and lithium chloride
δ	parts per million down field from tetramethylsilane


5

Preparation of Compounds


Compound 1



- 10 To a suspension of 7-bromo-2,4-bis-methylsulfanyl-imidazo[2,1-f][1,2,4]triazine (prepared according to WO2008116064, 500 mg, 1.72 mmol) in anhydrous THF (5 mL) was dropwise added BuLi (1.6 M in hexanes, 1.61 mL, 2.41 mmol) at -78°C . The suspension became red brown solution after 5 min, and then a mixture of **1a** (prepared according to WO 200631725, 675 mg, 1.81 mmol) and boron trifluoride etherate (2.40 mL, 1.89 mmol) in THF (5 mL) was added dropwise to the
- 15 mixture. After stirring for 2 h at -78°C , saturated NH_4Cl was added to quench the reaction. The mixture was diluted with ethyl acetate; the organic layer was washed with brine and concentrated *in vacuo*. The residue was purified by silica gel column chromatography (EtOAc / hexanes), affording **1b** as a rich yellow foam (650 mg,
- 20 67%). ^1H NMR (400 MHz, CDCl_3): δ 8.13 (d, 2H), 8.03 (d, 2H), 7.81 (d, 1H), 7.59 (t, 1H), 7.45 (m, 3H), 7.36 (t, 2H), 6.40 (brs, 1H), 6.01 (dd, 1H), 4.78 (m, 2H), 4.60 (dd, 1H), 2.68 (s, 3H), 2.45 (s, 3H), 1.62 (d, 3H). ^{19}F NMR (376 MHz, CDCl_3): δ -167.5. MS = 585.1 ($\text{M} + \text{H}^+$).




To a solution of **1b** (820 mg, 1.40 mmol) in dichloromethane (20 mL) were added boron trifluoride etherate (2 mL) and triethylsilane (2 mL), and stirred at room temperature for 16 h. Additional boron trifluoride etherate (1 mL) and triethylsilane (1 mL) were added, and stirred for 7 d. The mixture was diluted with dichloromethane and saturated sodium bicarbonate. The organic layer was washed sequentially with water, saturated ammonium chloride and brine, dried over magnesium sulfate, and concentrated. The residue was purified by silica gel column chromatography (EtOAc / hexanes), affording **1c** (605 mg, 76%). ^1H NMR (400 MHz, CDCl_3): δ 8.10 (d, $J = 7.2\text{ Hz}$, 2H), 8.00 (d, $J = 7.2\text{ Hz}$, 2H), 7.66 (s, 1H), 7.61 (t, $J = 7.2\text{ Hz}$, 1H), 7.53 (t, $J = 7.2\text{ Hz}$, 1H), 7.46 (t, $J = 7.2\text{ Hz}$, 2H), 7.38 (t, $J = 7.2\text{ Hz}$, 2H), 5.78 (m, 2H), 4.80 (dd, 1H), 4.68 (m, 1H), 4.60 (dd, 1H), 2.68 (s, 3H), 2.65 (s, 3H), 1.32 (d, 3H). ^{19}F NMR (376 MHz, CDCl_3): δ -149.9. MS = 569.1 ($\text{M} + \text{H}^+$).



Compound **1c** (635 mg, 1.12 mmol) was placed in a steel bomb reactor. Liquid ammonia (~30 mL) was charged and the bomb reactor was tightly sealed. The mixture was stirred at 50°C for 16 h. After cooling to room temperature, ammonia was evaporated and the solid residue was dissolved in THF (10 mL) and MeOH (10 mL). Sodium ethoxide (25% wt. 0.63 mL) was added and stirred at 60°C for 40 min.


- 5 The mixture was neutralized with AcOH and concentrated. The residue was purified by RP HPLC, affording the product **1d** (175 mg, 48%). ^1H NMR (400 MHz, DMSO- d_6): δ 8.21 (brs, 2H), 7.60 (s, 1H), 5.45 (brs, 1H), 5.43 (d, 1H), 4.91 (t, 1H), 3.92 (m, 1H), 3.76 (m, 2H), 3.57 (m, 1H), 2.44 (s, 3H), 1.09 (d, 3H). ^{19}F NMR (376 MHz, DMSO- d_6): δ -153.5. MS = 330.1 ($\text{M} + \text{H}^+$).

10



To a solution of **1d** (175 mg, 0.53 mmol) in dichloromethane (11 mL) was added MCPBA (370 mg, ~ 1.5 mmol) and stirred at room temperature for 16 h. The mixture was concentrated, affording crude **1e** which was used for the next reaction without purification. MS = 362.0 ($\text{M} + \text{H}^+$).

15




- Compound **1e** (obtained from the previous reaction) was placed in a steel bomb reactor. Liquid ammonia (~ 30 mL) was charged, and the bomb reactor was tightly sealed. The mixture was stirred at 115°C for 3 d. After cooling to room temperature, ammonia was evaporated. The solid residue was purified by RP HPLC, affording compound **1** (105 mg, 66% in two steps). ^1H NMR (400 MHz, D_2O): δ 7.31 (s, 1H), 5.43 (d, $J = 25.2$ Hz, 1H), 4.07 (dd, $J = 9.6, 23.2$, 1H), 3.89 (m, 1H), 3.83 (dd,

20


- 5 $J = 2.4, 12.8$ Hz, 1H), 3.67 (dd, $J = 4.8, 12.8$ Hz, 1H), 1.05 (d, $J = 22.8$ Hz, 3H). ^{19}F NMR (376 MHz, D_2O): δ -153.5. MS = 299.2 ($\text{M} + \text{H}^+$).

Compound 2



- To a solution of compound **1** (82 mg, 0.28 mmol) in water (340 mL) was added adenosine deaminase (A5168 bovine spleen type IX from Sigma-Aldrich, 0.125 Unit per mL of water) and stirred at 37°C for 4 h. The mixture was concentrated and purified by RP HPLC, affording compound **2** (56 mg, 68%). ^1H NMR (400 MHz, D_2O): δ 7.35 (s, 1H), 5.46 (d, $J = 25.2$ Hz, 1H), 4.08 (dd, $J = 9.6, 22.6$, 1H), 3.93 (m, 1H), 3.87 (dd, $J = 2.4, 12.8$ Hz, 1H), 3.71 (dd, $J = 4.8, 12.8$ Hz, 1H), 1.12 (d, $J = 23.2$ Hz, 3H). ^{19}F NMR (376 MHz, D_2O): δ -153.4. MS = 300.2 ($\text{M} + \text{H}^+$).
- 15

Compound 3




To a suspension of 7-bromo-pyrrolo[2,1-f][1,2,4]triazin-4-ylamine (prepared according to WO2007056170, 2.13 g, 10 mmol) in THF (20 mL) was added TMSCl (2.66 mL, 21 mmol) and stirred at room temperature for 16 h under argon. After cooling to -78°C, a solution of BuLi (1.6 M, 21 mL, 33 mmol) in hexanes was added

5 dropwise. The mixture was stirred for 1 h at the same temperature. A solution of **1a**
 (prepared according to WO 200631725, 4.46 g, 12 mmol) in THF (10 mL) was then
 added. After stirring for 2 h at -78°C , saturated ammonium chloride was added to
 quench the reaction. The mixture was extracted with ethyl acetate. The organic
 extract was concentrated *in vacuo*. The residue was purified by silica gel
 10 chromatography (ethyl acetate / hexanes), affording **3b** as a yellow solid (1.6 g, 32%).
 MS = 507.1 ($\text{M} + \text{H}^+$).


**Alternative procedure for Compound 3b using 1,2-bis-
 [(chlorodimethyl)silanyl]ethane instead of chlorotrimethylsilane**

15 To a suspension of 7-bromo-pyrrolo[2,1-f][1,2,4]triazin-4-ylamine (500 mg,
 2.35 mmol) in THF (6.5 mL) was added BuLi (1.6 M in hexanes, 1.6 mL) at -78°C .
 After 30 min., a solution of 1,2-bis-[(chlorodimethyl)silanyl]ethane (538 mg, 2.4
 mmol) in THF (1.2 mL) was added. After 45 min., BuLi (1.6 mL) was added. After
 20 an additional 30 min., BuLi (1.5 mL) was added. After 30 min., a solution of **1a** (610
 mg, 1.64 mmol) in THF (2 mL) was then added dropwise. The resulting mixture was
 stirred at -78°C for 2 h under argon. Acetic acid (0.7 mL) was added dropwise to
 quench the reaction, followed by addition of saturated ammonium chloride. The
 mixture was extracted with ethyl acetate. The organic extract was concentrated *in*
 25 *vacuo*. The residue was purified by silica gel chromatography (ethyl acetate /
 hexanes), affording **3b** (320 mg, 40%). The starting **1a** was also recovered (350 mg)
 from the chromatography.



K


- 5 To a solution of compound **3b** (50 mg, 0.1 mmol) and TMSCN (67 μ L, 0.5 mmol) in acetonitrile (2.0 mL) at 0°C was added TMSOTf (91 μ L, 0.5 mmol). The reaction mixture was stirred at room temperature for 1 h, then at 65°C for 3 d. The reaction was quenched with saturated NaHCO₃ at room temperature, and diluted with CH₃CO₂Et. The organic phase was separated, washed with brine, dried over Na₂SO₄,
 10 filtered and concentrated. The residue was purified by RP-HPLC (acetonitrile / water), to give the desired compound **3c** (28 mg, 54%). MS = 516.1 (M + H⁺).



- To a solution of **3c** (56 mg, 0.11 mmol) in methanol (1.2 mL) was added
 15 ammonium hydroxide (28% in water, 0.8 mL) and stirred at room temperature for 16 h. The mixture was concentrated and the residue was purified by RP HPLC (water / acetonitrile), affording compound **3** (20 mg, 60%). ¹H NMR (500 MHz, D₂O): δ 7.88 (s, 1H), 7.07 (d, 1H), 6.92 (d, 1H), 4.17 (m, 2H), 4.04 (dd, 1H), 3.87 (dd, 1H), 1.15 (d, 3H). MS = 308.1 (M + H⁺).

20

Compound 4




Handwritten signature

5 To a solution of compound **3b** (60 mg, 0.12 mmol) in methanol (0.5 mL) was added ammonium hydroxide (28% in water, 0.5 mL) and stirred at room temperature for 16 h. The mixture was concentrated and the residue was purified by RP HPLC (water / acetonitrile), affording compound **4** (25 mg, 70%). MS = 299.1 ($M + H^+$).

10

Compound 5




Compound **3b** was converted to compound **5a** by a procedure similar to conversion of **1b** to **1c**. Compound **5a** was then converted to compound **5** by a procedure similar to conversion of **3c** to **3**. ^1H NMR (300 MHz, D_2O): δ 7.68 (s, 1H), 6.75 (d, $J = 4.5$ Hz, 1H), 6.65 (d, $J = 4.5$ Hz, 1H), 5.65 (d, $J = 25.2$ Hz, 1H), 3.95 (m, 3H), 3.74 (dd, 1H), 0.98 (d, $J = 22.8$ Hz, 3H). ^{19}F NMR (282 MHz, D_2O): δ -154.2. MS = 283.2 ($M + H^+$).

20

General procedure for preparation of a nucleoside triphosphate:

- 5 To a pear-shaped flask (5-15 mL) is charged with a nucleoside (~20 mg). Trimethyl phosphate (0.5-1.0 mL) is added. The solution is cooled with ice-water bath. POCl_3 (40-45 mg) is added and stirred at 0°C until the reaction is complete (1 to 4 h; the reaction progress is monitored by ion-exchange HPLC; analytical samples are prepared by taking ~3 μL of the reaction mixture and diluting it with 1.0 M
- 10 $\text{Et}_3\text{NH}_2\text{CO}_3$ (30-50 μL)). A solution of pyrophosphate- Bu_3N (250 mg) and Bu_3N (90-105 mg) in acetonitrile or DMF (1-1.5 mL) is then added. The mixture is stirred at 0°C for 0.3 to 2.5 h, and then the reaction is quenched with 1.0 M $\text{Et}_3\text{NH}_2\text{CO}_3$ (~5 mL). The resulting mixture is stirred for additional 0.5-1 h while warming up to room temperature. The mixture is concentrated to dryness, re-dissolved in water (4 mL),
- 15 and purified by ion exchange HPLC. The fractions containing the desired product is concentrated to dryness, dissolved in water (~5 mL), concentrated to dryness, and again dissolved in water (~5 mL). NaHCO_3 (30-50 mg) is added and concentrated to dryness. The residue is dissolved in water and concentrated to dryness again. This process is repeated 2-5 times. The residue is then subjected to C-18 HPLC
- 20 purification, affording the desired product as a sodium or salt. Alternatively, the crude reaction mixture is subjected to C-18 HPLC first and then ion exchange HPLC purification to afford the desired product as a triethylammonium salt.


Compound TP-1



25 **Compound TP-1** was prepared by the general method using **Compound 2** as starting material. ^1H NMR (300 MHz, D_2O): δ 7.44 (s, 1H), 5.45 (d, $J = 25.5$ Hz, 1H), 4.0-4.4 (m, 4H), 3.05 (m, NCH_2CH_3), 1.10 (m, NCH_2CH_3 and $2'\text{-C-CH}_3$). ^{31}P

- 5 NMR (121.4 MHz, D₂O): δ -9.5 (d, J = 22.1 Hz), -11.0 (d, J = 19.9 Hz), -23.2 (t, J = 23.0 Hz). ¹⁹F NMR (282 MHz, D₂O): δ -153.9.

Compound TP-2




10

Compound TP-2 was prepared by the general method using **Compound 3** as starting material. ¹H NMR (300 MHz, D₂O): δ 7.82 (s, 1H), 7.03 (d, 1H), 6.90 (d, 1H), 4.1-4.4 (m, 4H), 3.05 (m, NCH₂CH₃), 1.10 (m, NCH₂CH₃ and 2'-C-CH₃). ³¹P NMR (121.4 MHz, D₂O): δ -10.7 (d, J = 19.5 Hz), -11.3 (d, J = 19.8 Hz), -23.1 (t, J = 19.8 Hz).

15

Compound TP-3




20

Compound TP-3 was prepared by the general method using **Compound 5** as starting material. ¹H NMR (300 MHz, D₂O): δ 7.73 (s, 1H), 6.87 (d, 1H), 6.82 (d, 1H), 5.71 (d, J = 24.6 Hz, 1H), 4.0-4.4 (m, 4H), 3.05 (m, NCH₂CH₃), 1.14 (m, NCH₂CH₃), 1.00 (d, J = 22.8 Hz, 3H, 2'-C-CH₃). ³¹P NMR (121.4 MHz, D₂O): δ -8.1 (d, J = 22.1 Hz), -11.1 (d, J = 19.9 Hz), -22.7 (t, J = 23.0 Hz). ¹⁹F NMR (282 MHz, D₂O): δ -155.6. MS = 520.9 (M - H⁺).

25


5 Compound TP-8a



TP-8a

Compound TP-8a was prepared by the general method using Compound 8 as starting material. ^1H NMR (300 MHz, D_2O): δ 7.95 (s, 1H), 7.68 (s, 1H), 5.63 (d, J = 25.5 Hz, 1H), 4.0-4.4 (m, 4H), 3.05 (m, NCH_2CH_3), 1.10 (m, NCH_2CH_3 and 2'-C- CH_3). ^{31}P NMR (121.4 MHz, D_2O): δ -9.20 (d, J = 22.1 Hz), -11.07 (d, J = 19.9 Hz), -23.82 (t, J = 23.0 Hz). ^{19}F NMR (282 MHz, D_2O): δ -155.9. MS = 521.6 ($\text{M} - \text{H}^+$).


General procedure for preparation of a nucleoside prodrug (Method A):



To a solution of a nucleoside (0.1 mmol) in trimethylphosphite (1.0 mL) are added 1H-tetrazole (42 mg, 0.6 mmol) followed by addition of 2,2-dimethyl-thiopropionic acid S-(2-{diisopropylamino-[2-(2,2-dimethyl-propionyl)sulfanyl)]-)

- 5 ethoxy]-phosphanyloxy}-ethyl) ester (prepared according to *J. Med. Chem.*, **1985**, 38, 3941, 90 mg, 0.2 mmol) at 0°C. After stirring for 2 h, 30% hydrogen peroxide in H₂O (140 µL) was added to the mixture. The mixture was then allowed to warm up to room temperature. After 30 min stirring, 1 M Na₂S₂O₃ in H₂O (5 mL) was added to quench the reaction. The organic layer was washed with saturated aqueous Na₂CO₃ (10 mL x 2), brine, concentrated *in vacuo*. The residue was purified by RP-HPLC (MeCN-H₂O gradient) to afford a prodrug **A**.

Compound A-1



A-1


- 15 Compound **A-1** was prepared by Method A using compound **1** as starting material. ¹H NMR (400 MHz, CDCl₃): δ 7.42 (s, 1H), 5.47 (d, *J* = 26.4 Hz, 1H), 4.95 (brs, 2H), 4.59 (m, 2H), 4.35 (m, 1H, 4'-H), 4.18 (m, 2H, 5'-H), 4.10 (m, 4H), 3.13 (m, 4H), 1.24 (d, 3H), 1.22 (s, 9H), 1.19 (d, 9H). ³¹P NMR (161.9 MHz, CDCl₃): δ -
- 20 1.26. MS = 667.1 (M + H⁺).

General procedure for preparation of a nucleoside prodrug (Method B):

- Non-limiting examples of mono-phosphoramidate prodrugs comprising the
- 25 instant invention may be prepared according to general Scheme 1.

Scheme 1


136




The general procedure comprises the reaction of an amino acid ester salt **19b**, e.g., HCl salt, with an aryl dichlorophosphate **19a** in the presence of about two to ten equivalents of a suitable base to give the phosphoramidate **19c**. Suitable bases include, but are not limited to, imidazoles, pyridines such as lutidine and DMAP, tertiary amines such as triethylamine and DABCO, and substituted amidines such as DBN and DBU. Tertiary amines are particularly preferred. Preferably, the product of each step is used directly in the subsequent steps without recrystallization or chromatography. Specific, but non-limiting, examples of **19a**, **19b**, and **19c** can be found in WO 2006/121820 that is hereby incorporated by reference in its entirety. A nucleoside base **19d** reacts with the phosphoramidate **19c** in the presence of a suitable base. Suitable bases include, but are not limited to, imidazoles, pyridines such as lutidine and DMAP, tertiary amines such as triethylamine and DABCO, and substituted amidines such as DBN and DBU. The product **B** may be isolated by recrystallization and/or chromatography.

Compound B-1

137

**B-1**


Phenyl ethoxyalaninyl phosphorochloridate (124 mg, 0.42 mmol; prepared according to McGuigan et al, *J. Med. Chem.* **1993**, 36, 1048–1052) was added to a mixture of **Compound 3** (20 mg, 0.065 mmol) and *N*-methylimidazole (42 μ L, 0.52 mmol) in anhydrous trimethyl phosphate (0.8 mL). The reaction mixture is stirred for 3 h at room temperature, and then methanol was added to quench the reaction. The methanol solvent is removed under reduced pressure. The residue was purified by reverse-phase HPLC and then by silica gel column chromatography (100% ethyl acetate), affording compound **B-1** (10 mg, 27%). ^{31}P NMR (121.4 MHz, CDCl_3): δ - 3.42, 3.77. MS = 563.0 ($\text{M} + \text{H}^+$), 561.0 ($\text{M} - \text{H}^+$).

Compound B-2**B-2**

About 3.1 mmol of 4-chlorophenyl 2-propoxyalaninyl phosphorochloridate (prepared according to McGuigan et al, *J. Med. Chem.* **1993**, 36, 1048–1052) is added to a mixture of about 0.5 mmol of **Compound 3** and about 3.8 mmol of *N*-methylimidazole in about 3 mL anhydrous trimethyl phosphate. The reaction mixture is stirred for about one hour to 24 hours at room temperature and methanol is added to

- 5 quench the reaction. The methanol solvent is removed under reduced pressure. The residue is purified by reverse-phase HPLC to give compound **B-2**.

Compound B-3




10

B-3

Compound B-3 was obtained by a similar procedure used for compound **B-1**.

^{31}P NMR (121.4 MHz, CDCl_3): δ -3.50, 3.76. MS = 577.2 ($\text{M} + \text{H}^+$).

15 **Compound B-4**




B-4

Compound B-4 was obtained by a similar procedure used for compound **B-1**.

20 ^{31}P NMR (162 MHz, CD_3OD): δ 2.2. MS = 633.4 ($\text{M} + \text{H}^+$).

Compound B-5



(Handwritten signature)

139


5

B-5

Compound B-5 was obtained by a similar procedure used for compound B-1.

^{31}P NMR (162 MHz, CDCl_3): δ 4.15, 4.27. MS = 549.3 ($\text{M} + \text{H}^+$).

Compound B-6



10


B-6

Compound B-6 was obtained by a similar procedure used for compound B-1.

^{31}P NMR (162 MHz, CDCl_3): δ 3.50, 4.07. MS = 613.1 ($\text{M} + \text{H}^+$).

15

Compound B-7




B-7

20

Compound B-7 was obtained by a similar procedure used for compound B-1, using compound 5 as parent nucleoside. ^{31}P NMR (162 MHz, CDCl_3): δ 3.37, 3.97. MS = 538.1 ($\text{M} + \text{H}^+$).

Compound B-8


140

**B-8**

Compound B-8 was obtained by a similar procedure used for compound **B-1**, using compound **5** as parent nucleoside. ^{31}P NMR (162 MHz, CDCl_3): δ 3.69, 4.39.

MS = 588.1 ($\text{M} + \text{H}^+$).

Alternative procedure for preparation of a nucleoside prodrug (Method C):

**C-1a**

Into a flask containing ethyl *L*-valine hydrochloride (2.5 g, 13.8 mmol, 1 equiv.) was added CH_2Cl_2 (46 mL, 0.3 M) and phenyl dichlorophosphate (2.1 mL, 13.8 mmol, 1 equiv.) before being cooled to -10°C . After 10 minutes, TEA (3.8 mL, 13.8 mmol, 1 equiv) was added slowly to the reaction mixture over five minutes. The reaction was allowed to proceed for an hour before *p*-nitrophenol (1.9 g, 13.8 mmol,


1 equiv.) was added to the reaction mixture followed by addition of more TEA (3.8 mL, 13.8 mmol, 1 equiv.) over five minutes. The reaction was allowed to warm up and proceed for another two hours. The reaction was concentrated *in vacuo* and taken up in diethyl ether (200 mL). The insoluble salts were filtered off and the filtrate concentrated *in vacuo*. Flash column chromatography was carried out using 4/1 Hex / EtOAc to furnish a clear oil as **C-1a**.

^1H NMR (400 MHz, CDCl_3): δ 8.21 (s, 2 H), 7.41 – 7.20 (m, 7 H), 4.22 – 4.05 (m, 3 H), 2.46 (s, 2 H), 1.99 (dd, $J = 23.0, 20.1$ Hz, 2 H), 1.68 (s, 1 H), 1.20 – 1.05 (m, 8 H).

^{31}P NMR (162 MHz, CDCl_3): δ -2.79 (dd, $J = 28.0, 4.2$ Hz).

LC MS m/z 422.99 [$\text{M} + \text{H}^+$].

Compound C-1



C-1


Into a flask containing **compound 3** (70 mg, 0.23 mmol, 1 equiv.) was added THF (1 mL, 0.2 M) and NMP (1 mL, 0.2 M) before cooling to 0°C. *t*-BuMgCl (560 μL , 2.5 equiv., 1M THF) was added slowly and allowed to stir for 5 minutes before the above phenolate **C-1a** (207 mg, 0.46 mmol, 2 equiv. dissolved in 500 μL of THF) was added. The reaction mixture was warmed to 50°C. The reaction was monitored by LCMS. Once the reaction was complete, the mixture was then concentrated *in vacuo*, and the residue was purified by HPLC, affording **Compound C-1**.

^1H NMR (400 MHz, CDCl_3) δ 7.87 (s, 1 H), 7.24 – 7.10 (m, 4 H), 7.03 (t, $J = 7.2$ Hz, 1H), 6.81 (d, $J = 4.6$ Hz, 1 H), 6.52 (d, $J = 4.7$ Hz, 1 H), 5.61 (s, 2H), 4.46 (dd, $J = 24.0, 11.4$ Hz, 2 H), 4.33 – 4.14 (m, 2 H), 4.06 (dt, $J = 7.2, 4.2$ Hz, 2 H), 3.82 – 3.70 (m, 1 H), 3.63 (t, $J = 10.6$ Hz, 2 H), 1.98 (s, 1 H), 1.17 (dd, $J = 14.8, 7.6$ Hz, 3 H),

0.82 (dd, $J = 22.8, 6.8$ Hz, 6 H).

α

- 5 ^{31}P NMR (162 MHz, CDCl_3): d 5.11.
 ^{19}F NMR (376 MHz, CDCl_3): d -152.28.
 LC MS m/z 591.21 $[\text{M} + \text{H}^+]$.



C-2a


Compound C-2a was obtained in a procedure similar to that exemplified for **Compound C-1a** but using the methionine ester.

^1H NMR (400 MHz, CDCl_3) d 8.19 (s, 2 H), 7.44 – 7.03 (m, 7 H), 4.11 (s, 2 H), 3.81 (d, $J = 44.5$ Hz, 1H), 2.04 (s, 3 H), 1.61 (s, 2 H), 1.21 (d, $J = 6.1$ Hz, 2 H), 1.01 – 0.65 (m, 4 H).

^{31}P NMR (162 MHz, CDCl_3) d -2.00 (d, $J = 12.9$ Hz).

LC MS m/z 455.03 $[\text{M} + \text{H}^+]$.


Compound C-2



C-2

Compound C-2 was obtained in a procedure similar to that exemplified for **Compound C-1** using **Compound 3** and **C-2a**.


- 5 ^1H NMR (400 MHz, CDCl_3) δ 7.96 (d, $J = 15.8$ Hz, 1H), 7.40 – 7.06 (m, 13H), 6.93 (d, $J = 6.7$ Hz, 1H), 6.70 (s, 1H), 5.98 (s, 1H), 4.54 (dd, $J = 21.6, 11.7$ Hz, 2H), 4.32 (d, $J = 12.0$ Hz, 2H), 4.14 (dt, $J = 13.0, 6.4$ Hz, 4H), 2.44 (d, $J = 7.5$ Hz, 2H), 2.00 (d, $J = 16.2$ Hz, 5H), 1.89 (s, 2H), 1.35 – 1.13 (m, 7H).
- ^{31}P NMR (162 MHz, CDCl_3) δ 4.12, 3.58.
- 10 ^{19}F NMR (376 MHz, CDCl_3) δ -152.28 (s).
- LC MS m/z 623.27 $[\text{M} + \text{H}^+]$.

**C-3a**

- 15 **Compound C-3a** was obtained in a procedure similar to that exemplified for **Compound C-1a** but using a tryptophan ester.
- ^1H NMR (400 MHz, CDCl_3) δ 8.18 – 8.03 (m, 3 H), 7.29 – 7.08 (m, 8 H), 7.36 – 6.98 (m, 3 H), 4.41 – 4.11 (m, 1 H), 4.15 – 3.95 (m, 2 H), 3.68 – 3.80 (m, 1 H), 3.33 – 3.04 (m, 2 H), 1.06 – 1.17 (m, 3 H).
- 20 ^{31}P NMR (162 MHz, CDCl_3) δ -2.87, -2.99.
- LC MS m/z 510.03 $[\text{M} + \text{H}^+]$.

Compound C-3

144

**C-3**


Compound C-3 was obtained in a procedure similar to that exemplified for **Compound C-1** using **Compound 3** and **C-3a**.

¹H NMR (400 MHz, CDCl₃) δ 8.27 (s, 1H), 7.84 (s, 1H), 7.47 (s, 1H), 7.36 – 6.77 (m, 11 H), 6.57 (s, 1 H), 4.40 – 3.96 (m, 6 H), 3.20 (s, 4 H), 2.60 (s, 1H), 1.30 – 1.04 (m, 6 H).

³¹P NMR (162 MHz, CDCl₃) δ 4.02, 3.75

¹⁹F NMR (376 MHz, CDCl₃) δ -152.13.

LC MS m/z 678.32 [M + H⁺].

**C-4a**

Compound C-4a was obtained in a procedure similar to that exemplified for **Compound C-1a** by substituting the phenylalanine ester.


¹H NMR (400 MHz, CDCl₃) δ 8.15 (t, *J* = 8.7 Hz, 2H), 7.43 – 7.11 (m, 10 H), 7.04 (ddd, *J* = 11.4, 6.7, 2.9 Hz, 2 H), 4.32 (ddd, *J* = 15.3, 11.3, 6.1 Hz, 4 H), 4.15 – 3.99 (m, 7 H), 3.74 (td, *J* = 11.0, 5.0 Hz, 8 H), 3.01 (d, *J* = 5.7 Hz, 2 H), 1.17 (td, *J* = 7.1, 5.2 Hz, 2 H).

145

5 ^{31}P NMR (162 MHz, CDCl_3) δ -2.97, -2.99.

LC MS m/z 471.03 $[\text{M} + \text{H}^+]$.

Compound C-4



C-4

Compound C-4 was obtained in a procedure similar to that exemplified for Compound C-1 using Compound 3 and C-4a.


15 ^1H NMR (400 MHz, CDCl_3) δ 7.92 (d, $J = 13.2$ Hz, 1H), 7.46 – 6.97 (m, 17H), 6.91 (s, 1H), 6.75 (s, 1H), 4.10 (dd, $J = 29.6, 19.2$ Hz, 8H), 2.97 (s, 3H), 1.32 – 1.05 (m, 7H).

^{31}P NMR (162 MHz, CDCl_3) δ 5.11.

^{19}F NMR (376 MHz, CDCl_3) δ -152.34 (s).

LC MS m/z 639.24 $[\text{M} + \text{H}^+]$.

20



C-5a

Compound C-5a was obtained in a procedure similar to that exemplified for Compound C-1a but using the proline ester.

2


- 5 ^1H NMR (400 MHz, CDCl_3) d 8.20 (d, $J = 7.8$ Hz, 2 H), 7.45 – 7.08 (m, 7 H), 4.37 (td, $J = 8.0, 3.8$ Hz, 2 H), 4.17 – 3.98 (m, 2 H), 3.61 – 3.34 (m, 2 H), 2.21 – 1.77 (m, 3 H), 1.19 (td, $J = 7.1, 3.8$ Hz, 3 H).

^{31}P NMR (162 MHz, CDCl_3) d -3.92, -3.96.

LC MS m/z 420.98 $[\text{M} + \text{H}^+]$.

10

Compound C-5



C-5

- 15 **Compound C-5** was obtained in a procedure similar to that exemplified for **Compound C-1** using **Compound 3** and **C-5a**.

^1H NMR (400 MHz, CDCl_3) d 7.95 (d, $J = 4.5$ Hz, 1 H), 7.39 – 7.10 (m, 4 H), 6.92 (dd, $J = 16.0, 4.6$ Hz, 1 H), 6.69 (s, 1H), 6.03 (bs, 2 H), 4.46 – 4.36 (m, 1 H), 4.36 – 3.96 (m, 4 H), 3.37 (d, $J = 58.9$ Hz, 2 H), 2.26 – 1.66 (m, 4 H), 1.39 – 1.12 (m, 8 H).


- 20 ^{31}P NMR (162 MHz, CDCl_3) d 3.47, 2.75.

^{19}F NMR (376 MHz, CDCl_3) d -152.36.

LC MS m/z 589.14 $[\text{M} + \text{H}^+]$.

Handwritten signature

147



5

Compound C-6

Compound C-6 was obtained in a procedure similar to that exemplified for **Compound C-1** using **Compound 3** and the sulphone analog of **C-1a**.

¹H NMR (400 MHz, CDCl₃) δ 7.93 (s, 1 H), 7.89 (s, 1 H), 7.35 – 7.01 (m, 5 H), 6.93 (d, *J* = 2.8 Hz, 1 H), 6.58 (d, *J* = 2.8 Hz, 1 H), 5.79 (bs, 2 H), 4.30 (s, 6 H), 4.11 (d, *J* = 7.0 Hz, 6 H), 3.10 – 2.84 (m, 3 H), 2.75 (s, 3 H), 2.54 (s, 6 H), 1.31 – 1.15 (m, 6 H).

³¹P NMR (162 MHz, CDCl₃) δ 3.39, 3.33.


¹⁹F NMR (376 MHz, CDCl₃) δ -152.40

LC MS *m/z* 655.24 [M + H⁺].

15

Compound PD-A-8b

α



5

To a solution of **Compound 8** (200 mg, 0.71 mmol) in THF (1 mL) and NMP (1 mL) under an atmosphere of argon at 0°C was added tert-butyl magnesium chloride (1.0 M in THF, 1.06 mL, 1.06 mmol). After 15 minutes, compound **30d-1** (280 mg, 0.71 mmol) was added as a solution in THF. After 5 minutes, the reaction mixture was allowed to warm to room temperature and was stirred for 2 hours. The reaction mixture was cooled to 0°C, quenched with MeOH, and concentrated. The reaction was purified by silica gel chromatography and then RP HPLC, affording **PD-A-8b** (225 mg, 59%). ¹H NMR (400 MHz, CDCl₃): δ 8.09 (two s, 1H), 7.54 (two s, 1H), 7.31-7.12 (m, 5H), 5.66 (dd, 1H), 4.52-4.45 (m, 2H), 4.19-4.03 (m, 4H), 3.87-3.69 (m, 1H), 1.35-1.15 (m, 9H). ³¹P NMR (161 MHz, CDCl₃): δ 4.14 (s), 3.55 (s). LC/MS = 539 (M + H⁺).

Retention time: 1.94 min

LC: Thermo Electron Surveyor HPLC


MS: Finnigan LCQ Advantage MAX Mass Spectrometer

20 Column: Phenomenex Polar RP 30 mm X 4.6 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.


5 Preparation of 30d-1



Compound **30d-1** was prepared from **30a** in a matter similar to that of **30d-2** substituting alanine ethyl ester hydrochloride for alanine isopropyl ester hydrochloride.


10

Compound (S)-PD-A-8c



- 5 Compound (**S**)-**PD-A-8c** was prepared in a matter similar to that of **PD-A-8b** substituting (**S**)-**30d-2** for **30d-1**. ^1H NMR (400 MHz, CDCl_3): δ 8.14 (s, 1H), 7.60 (s, 1H), 7.1-7.3 (m, 5H), 5.66 (dd, 1H), 5.02 (m, 1H), 4.50 (m, 1H), 4.40 (m, 1H), 4.1-4.3 (m, 2H), 3.98 (m, 1H), 3.78 (m, 1H), 3.18 (brs, 1H), 1.15-1.4 (m, 12H). ^{31}P NMR (161 MHz, CDCl_3): δ 3.70 (s).
- 10 LC/MS = 553 ($\text{M} + \text{H}^+$).

Preparation of (**S**)-**30d-2**




- 15 Alanine isopropyl ester hydrochloride (7.95 g, 47.4 mmol) was suspended in dichloromethane (100 mL). Compound **31a** (10 g, 47.4 mmol) was added. Triethylamine (13.2 mL, 95 mmol) was then dropwise added over a period of 15 min. (internal reaction temperature; $-10^\circ\text{C} \sim -3^\circ\text{C}$). When the reaction was almost complete (by phosphorous NMR), *p*-nitrophenol (6.29 g, 45.0 mmol) was added as a
- 20 solid in one portion. To the resulting slurry was added triethylamine (6.28 mL, 45 mmol) over a period of 15 min. The mixture was then warmed up to room temperature. When the reaction was complete, MTBE (100 mL) was added. The white precipitate was removed by filtration. The filter cake was washed with MTBE (3 x 50 mL). The filtrate and washings were combined and concentrated. The residue
- 25 was purified by silica gel column chromatography (0 to 50% ethyl acetate / hexanes), affording compound **30d-2** as a 1:1 ratio of diastereomeric mixture (14.1 g, 77%). ^1H NMR (300 MHz, CDCl_3): δ 8.22 (2d, 2H), 7.2-7.4 (m, 7H), 5.0 (m, 1H), 4.09 (m,

2

- 5 ^1H), 3.96 (m, 1H), 1.39 (2d, 3H), 1.22 (m, 6H). MS = 409.0 ($\text{M} + \text{H}^+$), 407.2 ($\text{M} - \text{H}^+$).

Separation of two diastereomers of compound 30d-2



10

The two diastereomers were separated by chiral column chromatography under the following conditions;

Column: Chiralpak IC, 2 x 25 cm

Solvent system: 70% heptane and 30% isopropanol (IPA)

15 Flow rate: 6 mL/min.

Loading volume per run: 1.0 mL

Concentration of loading sample: 150 mg / mL in 70% heptane and 30% IPA

(S)-compound 30d-2: retention time 43 min. ^{31}P NMR (162.1 MHz, CDCl_3): δ -2.99 (s).



20 (R)-compound 30d-2: retention time 62 min. ^{31}P NMR (162.1 MHz, CDCl_3): δ -3.02 (s).

Alternatively, the two diastereomers were separated by crystallization under the following procedure;

- 5 Compound **30d-2** was dissolved in diethyl ether (~10 mL/ gram). While stirring, hexanes was then added until the solution became turbid. Seed crystals (~10 mg / gram of compound **30d-2**) were added to promote crystallization. The resulting suspension was gently stirred for 16 h, cooled to ~ 0 °C, stirred for an additional 2 h, and filtered to collect the crystalline material (recovery yield of the crystalline
- 10 material 35%-35%). The crystalline material contains ~95% of (S)-compound **30d-2** and ~5% of (R)-compound **30d-2**. Re-crystallization afforded 99% diastereomerically pure (S)-isomer.

The following **PD-A** compounds as examples are made by the general procedures:


15

Compound PD-A-8d20 **Compound PD-A-8e****Compound PD-A-8f**



25

153

5

**Compound PD-A-8g**


10

**Compound PD-A-8h****General procedure for preparation of a nucleoside prodrug (Method D):**

15

Non-limiting examples of 3'-O-acylated mono-phosphoramidate prodrugs comprising the instant invention may be prepared according to general Scheme 2.


Scheme 2



The general procedure comprises the reaction of **PD-A** ($\text{R}^4 = \text{OH}$) with a carboxylic acid or an activated carboxylate such as an acyl chloride or an acid anhydride, which is generally known to those skilled in the art (*Journal of Medicinal Chemistry*, **2006**, 49, 6614 and *Organic Letters*, **2003**, 6, 807). When $\text{R}^8 = \text{NH}_2$, protection of the amino group may be necessary. Briefly, to a solution of compound **PD-A** in acetonitrile (2 mL) is added *N,N*-dimethylformamide dimethyl acetal (~ 1.1 eq.) and stirred at room temperature for 1 h. After the protection of 6-amino group is complete, the mixture is then concentrated to dryness. To the residue are added a dehydrating agent such as DCC (~ 4 eq.), acetonitrile and a carboxylic acid (~ 2 eq.). The mixture is stirred at room temperature for 24 -48 h. Water (0.2 mL) and trifluoroacetic acid (0.1 mL) are added at 0°C and stirred at room temperature for 64 h. Sodium bicarbonate was added at 0°C . The mixture is stirred at room temperature for 0.5 h and filtered. The filtrate is concentrated and the residue was purified by silica gel column chromatography to afford compound **PD-B**. If an acyl chloride or an acid anhydride is used, a suitable base, such as triethylamine, is added instead of a dehydrating agent.

Compound PD-B-8i

Handwritten signature or mark.



To a solution of **PD-A-8b** (100 mg, 0.19 mmol) in DCM (1.0 mL) under an atmosphere of argon at room temperature was added *N,N*-dimethylformamide-dimethylacetal (25 μ L, 0.19 mmol). After 30 minutes, the reaction mixture was concentrated. The reaction was taken up in DCM and concentrated. This process was repeated twice. The resulting residue was taken up in THF (1.0 mL) and cooled to 0°C under an atmosphere of argon. To the solution was added triethylamine (79 μ L, 0.57 mmol) and DMAP (5mg, 0.04 mmol). After 5 minutes, isobutyryl chloride (60 μ L, 0.57 mmol) was added. After 10 minutes, the reaction was allowed to warm to room temperature and was stirred for 3 hours. The mixture was cooled to 0°C, quenched with a 5% TFA solution in water, and then allowed to stir at room temperature for 4 hours. The resulting mixture was extracted with ethyl acetate (3x). The combined organic layers were dried with sodium sulfate, filtered and concentrated. The residue was purified by RP HPLC (acetonitrile / water), affording **PD-B-8i** (71 mg, 61%). ¹H NMR (400 MHz, CDCl₃): δ 8.17 (two s, 1H), 7.66 (two s, 1H), 7.34-7.14 (m, 5H), 5.69 (dd, 1H), 5.56-5.43 (m, 1H), 4.55-4.01 (m, 5H), 3.79-3.69 (m, 1H), 2.70-2.64 (m, 1H), 1.37-1.17 (m, 15H). ³¹P NMR (161 MHz, CDCl₃): δ 2.99 (s), 2.88 (s).

LC/MS = 609 ($M + H^+$).

Retention time: 2.21 min

LC: Thermo Electron Surveyor HPLC

MS: Finnigan LCQ Advantage MAX Mass Spectrometer


Column: Phenomenex Polar RP 30 mm X 4.6 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid


- 5 Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

The following **PD-B** compounds as examples are made by the general procedures:


10 **Compound PD-B-8j**



Compound PD-B-8k




Compound PD-B-8l




2

5 Compound PD-B-8m




Compound PD-B-8n

10



Compound PD-B-8o

15




General procedure for preparation of a nucleoside prodrug (Method E):

Handwritten signature or mark.

- 5 Non-limiting examples of 3',5'-cyclic mono-phosphoramidate prodrugs comprising the instant invention may be prepared according to general Scheme 3.

Scheme 3



10


- 15 Scheme 3 illustrates chemical processes that may be useful for preparation of compound **PD-C**. Accordingly, **PD-A1** is converted to **PD-C** in the presence of a base when Ar is substituted with an electron withdrawing group such as *p*-nitro or *p*-chloro group (*European Journal of Medicinal Chemistry*, **2009**, 44, 3769).

Alternatively, compound **40** is converted to **Compound 41** according to *Bioorganic and Medicinal Chemistry Letters*, **2007**, 17, 2452, which is then coupled with an amino acid ester salt to form **PD-C**.

20 Compound PD-C-8q

2

159




5

A solution of **PD-A-8p** in DMSO is treated at room temperature with potassium *t*-butoxide (~1 eq.) and the resulting mixture is stirred for about 10 min. to about 2 h. The mixture is then cooled to 0°C and neutralized with 1N HCl to ~ pH 6. The mixture is purified by HPLC to afford compound **PD-C-8q**.

10


Additionally, the following **PD-C** compounds as examples are made by the general procedures:

Compound **PD-C-8r**





15

Compound **PD-C-8s**




, and

5 **Compound PD-C-8t****Compound PD-D-8u**

10

Compound **8** is dissolved in PO(OMe)₃ (0.1 - 0.5 M solution) and cooled to 0°C under argon. To this stirring solution is added POCl₃ (1.0 - 5.0 eq.) dropwise, and the reaction mixture is allowed to warm to room temperature for about 2 -16 h. The resulting solution is added dropwise to a rapidly stirring solution of acetonitrile and 0.05 - 0.5 M aqueous KOH. When addition is complete, the solvents are removed under reduced pressure. The resulting residue is dissolved in water and purified by HPLC to give **Compound 41-1**.

15



20


5

A solution of **Compound 41-1** in DCM and $\text{PO}(\text{OMe})_3$ is prepared and cooled to 0°C . To this solution is added oxalyl chloride (1.0 - 5.0 eq.) followed by a catalytic amount of DMF. The mixture is allowed to stir for about 10 min. to about 1h. When activation is complete, a large volume of 2-propanol is added to the reaction mixture and allowed to stir and warm to room temperature. The solvents are removed under reduced pressure, and the resulting crude material is purified by preparative HPLC to give **Compound PD-D-8u**.

10


Compound PD-E-8v

15




Compound PD-E-8v is prepared from **Compound 41-1** in a matter similar to that of **Compound PD-D-8u** substituting 2-aminopropane for 2-propanol.

20

Compound PD-F-8w**Compound PD-F-8w**

- 5 **Compound PD-F-8w** is prepared in a matter similar to that of **Compound 20** substituting **Compound 8** for **Compound 18**.


Compound PD-G-8x



Compound PD-G-8x

- About 90 mM **Compound 8** in THF is cooled to about -78°C and about 2.2 to
 10 about 5 equivalents of *t*-butylmagnesium chloride (about 1 M in THF) is added. The mixture is warmed to about 0°C for about 30 min and is again cooled to about -78°C . A solution of (2*S*)-2-{{[chloro(1-phenoxy)phosphoryl]amino}ethyl isobutyrate (WO2008085508) (1 M in THF, about 2 equivalents) is added dropwise. The cooling is removed and the reaction is stirred for about one to about 24 hours. The reaction is
 15 quenched with water and the mixture is extracted with ethyl acetate. The extracts are dried and evaporated and the residue purified by chromatography to give **Compound PD-G-8x**.

Compound 6




20

- Compound 4** (about 0.04 mmol) and anhydrous MeOH (about 5 mL) is treated with acetic acid (about 5 mL) and the reaction is stirred overnight at room temperature. Saturated NaHCO_3 is added to neutralize the reaction mixture and the
 25 crude material is purified using a HPLC system (acetonitrile- H_2O) to give **6**.

5


Compound 7



To a dry, argon purged round bottom flask (50 mL) is added compound **3b** (about 0.39 mmol) and anhydrous dichloromethane (about 10 mL). The flask is placed into a dry ice/acetone bath ($\sim -78^\circ\text{C}$) and the solution is stirred for about 10 min. $\text{BF}_3\cdot\text{Et}_2\text{O}$ (about 0.10 mL) is added dropwise and the reaction is stirred for about 10 min. AlMe_3 (about 1.16 mmol, 2.0 M in toluene) is then added. After a few minutes, the dry ice/acetone bath is removed and the reaction mixture is stirred at room temperature to about 45°C over about 4 h to about 4 d. A solution of pyridine (about 2 mL) in MeOH (about 10 mL) is added and the solvent is removed under reduced pressure. The crude material is purified by chromatography and is treated with ammonium hydroxide in methanol for about 16 h at about room temperature. The mixture is concentrated and the residue is purified by HPLC to give **7**.

20

Compound 8



To a suspension of 7-bromoimidazo[1,2-f][1,2,4]triazin-4-amine (obtained according to *ACS Medicinal Chemistry Letters*, 2010, 1, 286; 375 mg, 1.75 mmol) in

- 5 THF (4.0 mL) under an atmosphere of argon was added 1,2-bis-
 [(chlorodimethyl)silanyl]ethane (452 mg, 2.10 mmol). After 60 min, the reaction was
 cooled to -78°C and BuLi (1.6 M in THF, 3.8 mL, 6.10 mmol) was added. After 10
 min at -78°C , a solution of **1a** (obtained according to WO 200631725, 782 mg, 2.10
 mmol) in THF (1.0 mL) was added dropwise. The resulting mixture was stirred at -
 10 78°C for 1 hour. Saturated aqueous ammonium chloride was added and allowed to
 warm to 0°C . Water was added until all solids became soluble. The mixture was
 extracted with ethyl acetate. The organic extract was dried with sodium sulfate,
 filtered and concentrated in vacuo. The residue was purified by silica gel
 chromatography (ethyl acetate / hexanes), affording **8b** (606 mg, 59%) as a yellow
 15 solid.

LC/MS = 508 ($\text{M} + \text{H}^+$)

Retention time: 2.17-2.26 min


LC: Thermo Electron Surveyor HPLC

MS: Finnigan LCQ Advantage MAX Mass Spectrometer

- 20 Column: Phenomenex Polar RP 30 mm X 4.6 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid


Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5
 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.



25

To a solution of compound **8b** (510 mg, 1.39 mmol) in dichloroethane (10.0
 mL) at 0°C under an atmosphere of argon, was added triethyl silane (1.77 mL, 11.09
 mmol) and then $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (1.41 mL, 11.09 mmol). The reaction mixture was stirred
 at 55°C for 16h. The reaction was cooled to 0°C and quenched with saturated
 30 NaHCO_3 (aq). The reaction was extracted with DCM and then EtOAc. The

- 5 combined organic phase was dried over sodium sulfate, filtered and concentrated. The residue was purified by silica gel chromatography (ethyl acetate/hexanes), affording **8c** (453 mg, 64%). ¹H NMR (400 MHz, CDCl₃): δ 8.10-7.94 (m, 5H), 7.6-7.33 (m, 7H), 5.91 (dd, 1H), 5.78 (d, *J*=24.6 Hz, 1H), 4.87 (dd, 1H), 4.70 (m, 1H), 4.58 (dd, 1H), 1.31 (d, *J*=22.4 Hz, 3H).
- 10 LC/MS = 491 (M⁺).
Retention time: 2.36 min.
LC: Thermo Electron Surveyor HPLC
MS: Finnigan LCQ Advantage MAX Mass Spectrometer
Column: Phenomenex Polar RP 30 mm X 4.6 mm
- 15 Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid
Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.




- 20 To a solution of **8c** (500 mg, 01.02 mmol) in THF (5.0 mL) was added lithium hydroxide (122 mg, 5.09 mmol) as a solution in H₂O (5.0 mL) and was stirred at room temperature for 1h. The reaction was cooled to 0°C and was neutralized with 1N HCl in water (5.1 mL). The mixture was concentrated and the residue was purified by RP HPLC (water / acetonitrile), affording **Compound 8** (185 mg, 64%). ¹H NMR (400
- 25 MHz, CD₃OD): δ 7.97 (s, 1H), 7.63 (s, 1H), 5.54 (d, *J*=24.8 Hz, 1H), 4.03 (dd, 1H), 3.88 (m, 1H), 3.71 (dd, 1H), 1.80 (d, *J*=22.1 Hz, 3H).
LC/MS = 284 (M + H⁺).
Retention time: 1.06 min.
LC: Thermo Electron Surveyor HPLC
- 30 MS: Finnigan LCQ Advantage MAX Mass Spectrometer

5 Column: Phenomenex Polar RP 30 mm X 4.6 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid


Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

10 Alternative procedure for **Compound 8**



Compound **1e** (crude obtained from the previous reaction step) was dissolved in EtOH. Excess sodium borohydride was added in portions until the reaction was nearly complete. The mixture was neutralized with acetic acid. The mixture was concentrated and the solid residue was purified by silica gel column chromatography (0-10% MeOH / dichloromethane), affording compound **27** (210 mg, 50% in two steps).


20 Additional alternative procedure for **Compound 8**



Raney Ni (about 500 mg) was neutralized by washing with H₂O, and added to a solution of **1d** (about 100 mg) in ethanol (about 10 mL). The mixture was then heated to 80°C until the reaction is complete. The catalyst was removed by filtration

- 5 and the solution was concentrated *in vacuo*. The mixture was concentrated and the residue was purified by HPLC to give **8**.

Compound 9




10

9

- Into a flask containing **Compound 3** (120 mg, 0.39 mmol, 1 equiv.) was added $\text{PO}(\text{OMe})_3$ (1.5 mL, 0.25 M) and cooled to 0°C before adding POCl_3 (125 μL , 1.37 mmol, 3.5 equiv.). The reaction mixture was allowed to stir for 5 hr before the reaction was quenched with water. It was directly purified by HPLC to furnish the
- 15 monophosphate **Compound 9**.
LC MS m/z 387.95 $[\text{M} + \text{H}^+]$.

Compound 10




20

10

- Into a flask containing **Compound 9** (30 mg, 0.078 mmol, 1 equiv.) was added NMP (0.8 mL, 0.1 M) followed by addition of TEA (43 μL , 0.31 mmol, 4 equiv.), tetrabutylammonium bromide (25 mg, 0.078 mmol, 1 equiv.) before adding chloromethylisopropyl carbonate (60 μL , 0.38 mmol, 5 equiv.). The reaction
- 25 mixture was heated to 50°C and allowed to stir overnight. It was purified directly by HPLC, affording **Compound 10**.


- 5 ^1H NMR (400 MHz, CDCl_3) δ 7.98 (s, 1 H), 7.01 (d, $J = 4.7$ Hz, 1 H), 6.72 (d, $J = 4.7$ Hz, 1 H), 6.04 (bs, 2 H), 5.74 – 5.61 (m, 4 H), 4.91 (ddt, $J = 12.6, 9.4, 6.3$ Hz, 2 H), 4.64 – 4.28 (m, 4 H), 1.37 – 1.19 (m, 15 H).
- ^{31}P NMR (162 MHz, CDCl_3) δ -4.06.
- ^{19}F NMR (376 MHz, CDCl_3) δ -76.58, -151.95 TFA salt.
- 10 LC MS m/z 620.03 $[\text{M} + \text{H}^+]$.

Compound 11**11**

- 15 A solution of **Compound B-2** in DMSO is treated with about 3 mole equivalents of potassium *t*-butoxide for about 15 min to 24 hours. The reaction is quenched with 1N HCl and **Compound 11** is isolated by reverse-phase HPLC.

Compound 12


169



Compound 1b (about 1 mmol) is placed in a steel bomb reactor. The reactor is charged with liquid ammonia (about 30 mL) and the mixture is stirred at about 0°C to 50°C for about 16 h. The ammonia is evaporated and the residue is purified to give **12a**. A solution of **12a** (about 100 mg) in ethanol (about 10 mL) is treated with Raney Ni (about 500 mg) that is neutralized by washing with H₂O. The mixture is then heated to about 35 to about 80°C until the reaction is complete. The catalyst is removed by filtration and the solution is concentrated *in vacuo*. The mixture is concentrated and the residue is purified by HPLC to give **12b**. To a solution of compound **12b** (about 50 mg) and TMSCN (about 0.5 mmol) in acetonitrile (about 2.0 mL) at about 0°C is added TMSOTf (about 0.5 mmol). The reaction mixture is stirred at room temperature for about 1 h, then at 65°C for about 3 d. The reaction is quenched with saturated NaHCO₃ at room temperature, and diluted with CH₃CO₂Et. The organic phase was separated, washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue is purified by RP-HPLC then dissolved in methanol (about 1 mL). Ammonium hydroxide (28% in water, about 0.8 mL) is added and the

- 5 mixture is stirred at about room temperature for 16 h. The mixture is concentrated and the residue is purified by RP HPLC to give 12.


Compound 13



13

10 **Compound 13** is prepared in the same manner as **Compound 9** using **Compound 12** as a starting material.


Compound 14



14

15 **Compound 14** is prepared by treating **Compound 13** with about one to about five equivalents of DCC in pyridine and heating the reaction to reflux for about one to about 24 hours. **Compound 14** is isolated by conventional ion exchange and reverse-phase HPLC.


Compound 15




15

5 A solution of about 0.4 mmol of **Compound 14** in about 10 mL of DMF is treated with about 0.8 mmol of DIPEA and about 0.8 mmol of chloromethyl isopropyl carbonate (WO2007/027248). The reaction is heated to about 25 to about 80°C for about 15 min to about 24 hours. The solvent is removed under vacuum and the residue is purified by HPLC to give **Compound 15**.


10

Compound 16**16**


Compound **3** (about 0.22 mmol) is dissolved in anhydrous pyridine (about 2 mL) and chlorotrimethylsilane (about 0.17 mL) is added. The mixture is stirred at about 0°C to about 25°C for about one to about 24 hours. Additional chlorotrimethylsilane (about 0.1 mL) is added and the reaction is stirred for about one to about 24 hours. 4,4'-Dimethoxytrityl chloride (about 0.66 mmol) and DMAP (about 0.11 to about 0.22 mmol) is sequentially added. The mixture is stirred for about one to about 24 hours. A solution of TBAF (1.0 M, about 0.22 mL) in THF is added and the reaction is stirred for about one to about 24 hours. The mixture is partitioned between ethyl acetate and water. The ethyl acetate layer is dried and concentrated. The residue is purified chromatography to afford **Compound 16**.

25 **Compound 17****17**

5 A mixture of about 1.25 mmol of **Compound 16** and about 1.9 mmol of triethylammonium 2-(2,2-dimethyl-3-(trityloxy)propanoylthio)ethyl phosphinate (WO2008082601) is dissolved in anhydrous pyridine (about 19 mL). Pivaloyl chloride (about 2.5 mmol) is added dropwise at about -30 to about 0 °C and the solution is stirred at for about 30 min to about 24 hours. The reaction is diluted with
10 methylene chloride and is neutralized with aqueous ammonium chloride (about 0.5M). The methylene chloride phase is evaporated and the residue is dried and is purified by chromatography to give **Compound 17**.


Compound 18**18**

To a solution of about 0.49 mmol of **Compound 17** in anhydrous carbon tetrachloride (about 5 mL) is added dropwise benzylamine (about 2.45 mmol). The reaction mixture is stirred for about one to about 24 hours. The solvent is evaporated
20 and the residue is purified by chromatography to give **Compound 18**.

Compound 20**20**


- 5 A solution of about 2 mmol of **Compound 18** in methylene chloride (about 10 mL) is treated with an aqueous solution of trifluoroacetic acid (90%, about 10 mL). The reaction mixture is stirred at about 25 to about 60 °C for about one to about 24 hours. The reaction mixture is diluted with ethanol, the volatiles are evaporated and the residue is purified by chromatography to give **Compound 20**.

10

Compound 21**21**

- 15 About 90 mM **Compound 2** in THF is cooled to about -78 °C and about 2.2 to about 5 equivalents of *t*-butylmagnesium chloride (about 1 M in THF) is added. The mixture is warmed to about 0 °C for about 30 min and is again cooled to about -78 °C.

A solution of (2*S*)-2-{{chloro(1-phenoxy)phosphoryl}amino}propyl pivalate (WO2008085508) (1 M in THF, about 2 equivalents) is added dropwise. The cooling is removed and the reaction is stirred for about one to about 24 hours. The reaction is quenched with water and the mixture is extracted with ethyl acetate. The extracts are dried and evaporated and the residue purified by chromatography to give **Compound 21**.

Compound 22

174

5

22a


Compound 22a was obtained in a procedure similar to that for preparation of

C-1a.

^1H NMR (400 MHz, CDCl_3) δ 8.11 (d, $J = 9.0$ Hz, 2 H), 8.02 (s, 1 H), 7.48 (t, $J = 7.5$ Hz, 2 H), 7.42 – 7.25 (m, 4 H), 7.21 (dt, $J = 14.9, 5.5$ Hz, 2 H), 7.08 (t, $J = 7.3$ Hz, 2 H), 5.17 – 5.03 (m, 2 H), 4.99 (dd, $J = 16.5, 9.7$ Hz, 2 H), 3.44 (s, 1H), 3.35 – 3.21 (m, 2 H), 3.19 (d, $J = 9.2$ Hz, 1H), 3.00 – 2.80 (m, 2 H).

^{31}P NMR (162 MHz, CDCl_3) δ 4.27.

LC MS m/z 452.09 $[\text{M} + \text{H}^+]$.



22b


Compound 22b was obtained in a procedure similar to that for preparation of

C-1 using Compound 3 and 22a.

^1H NMR (400 MHz, CD_3OD) δ 7.76 (d, $J = 6.3$ Hz, 1H), 7.38 (t, $J = 8.2$ Hz, 1 H), 7.27 – 7.12 (m, 4 H), 7.06 – 6.81 (m, 3 H), 6.74 (dd, $J = 4.6, 3.5$ Hz, 1 H), 4.95 – 4.79 (m, 1 H), 4.35 – 3.90 (m, 4 H), 3.23 (dt, $J = 3.2, 1.6$ Hz, 3H), 3.18 – 3.05 (m, 2 H), 2.82 (dt, $J = 14.7, 7.3$ Hz, 2 H), 1.15 (d, $J = 22.4$ Hz, 3 H).

^{31}P NMR (162 MHz, CD_3OD) δ 10.76, 10.71.

LC MS m/z 620.05 $[\text{M} + \text{H}^+]$.



2

25

5

Compound 22

Into a flask containing the **22b** (50 mg, 0.08 mmol, 1 equiv.) was added ethanol (4 mL) followed by $\text{Pd}(\text{OH})_2$ (56 mg, 0.08 mmol, 1 equiv.) and ammonium formate (42 mg, 0.64 mmol, 8 equiv.). The reaction was heated to 80°C for about an hour. The solid was filtered off and the material purified by HPLC.

^1H NMR (400 MHz, DMSO-d_6) δ 10.72 (s, 1H), 7.91 (s, 1H), 7.95 – 7.89 (bs, 2H), 7.41 (d, $J = 7.7$ Hz, 1H), 7.26 (d, $J = 8.1$ Hz, 1H), 7.19 – 6.66 (m, 3H), 4.20 – 3.75 (m, 3H), 2.99 (dd, $J = 16.5, 9.6$ Hz, 2H), 2.89 – 2.70 (m, 2H), 2.48 – 2.58 (m, 8H), 1.10 (d, $J = 22.3$ Hz, 3H).


^{31}P NMR (162 MHz, DMSO-d_6) δ 7.49.

^{19}F NMR (376 MHz, DMSO-d_6) δ -154.89.

LC MS m/z 530.21 $[\text{M} + \text{H}^+]$.

20

Compound 23



Compound 3

Compound 23

Compound **3** (250mg, 0.82mmol) was dissolved in $\text{PO}(\text{OMe})_3$ (5 mL, 0.16M) and cooled to 0°C under argon. To this stirring solution was added POCl_3 (0.32 mL, 4.1 mmol) slowly dropwise, and the reaction mixture allowed to warm to room temperature for 16 h. The resulting solution was added dropwise to a rapidly stirring solution of acetonitrile (400 mL) and 0.08M aqueous KOH (300 mL). When addition was complete, the reaction progress was checked by LCMS. When the reaction was

5 complete, solvents were removed under reduced pressure. The resulting solid residue was dissolved in water and purified by HPLC to give 140mg of **Compound 23** (yield; 47%).

$^1\text{H-NMR}$ (400MHz; CD_3OD) : d 8.15 (s, 1H), 7.40 (d, 1H; $J = 4.8\text{Hz}$), 7.09 (d, 1H; $J = 4.8\text{Hz}$), 4.64 (dd, 1H; $J = 24\text{Hz}$, 7.2Hz), 4.50-4.36 (m, 3H), 1.32 (d, 3H; $J=22\text{Hz}$).


10 $^{19}\text{F-NMR}$ (376MHz; CD_3OD) : d -153.11.

$^{31}\text{P-NMR}$ (162MHz; CD_3OD) : d -2.20.

MS $[\text{M} + \text{H}^+] = 370.2$.

Compound 24

15



A solution of **Compound 23** (7mg, 0.02 mmol) in DCM (2 mL) and $\text{PO}(\text{OMe})_3$ (1 mL) was prepared and cooled to 0°C . To this solution was added oxalyl-Cl (10 μL) followed by DMF (2 μL). The mixture was allowed to stir for 1 min before an aliquot was taken out and quenched in MeOH and then checked by LCMS for activation. Successive amounts of oxalyl-Cl (10 μL) and DMF (2 μL) were added until activation was complete. At this point, a large volume of 2-propanol (5 mL) was added to the reaction mixture and allowed to stir and warm to room temperature. Once the reaction was complete, the solvents were removed under reduced pressure, and the resulting crude material was purified by preparative HPLC to give 5.5 mg of **Compound 24** (yield 70%).

$^1\text{H-NMR}$ (400MHz; DMSO-d_6) : d 8.26 (br, 1H), 8.15 (br, 1H), 7.97 (s, 1H), 7.00 (d, 1H; $J = 4.4\text{Hz}$), 6.88 (d, 1H; $J = 4.4\text{Hz}$), 4.59-4.51 (m, 2H), 4.37-4.25 (m, 2H), 1.23 (d, 3H; $J=22.8\text{Hz}$).

2


5 ^{19}F -NMR (376MHz; CD_3OD) : d -151.72.

^{31}P -NMR (162MHz; CD_3OD) : d -5.69.

MS $[\text{M} + \text{H}^+] = 412.0$.

Compound 25

10



Compound 25 was prepared from **Compound 23** in a matter similar to that of **Compound 24** substituting the heptyl ester of alanine for 2-propanol (yield 5.3%).


15 ^1H -NMR (400MHz; CD_3OD) : d 7.91 (s, 1H), 6.98 (d, 1H; $J = 4.8\text{Hz}$), 6.92 (d, 1H; $J = 4.8\text{Hz}$), 5.29 (dd, 1H; $J = 24.4\text{Hz}$, 8.8Hz), 4.66-4.60 (m, 2H), 4.48-4.40 (m, 1H), 4.15-4.11(m, 3H), 3.92(dd, 1H; $J = 9.6\text{Hz}$, 7.2Hz), 1.67-1.64 (m, 3H), 1.40-1.27 (m, 15H), 0.91-0.87(m, 6H).

^{19}F -NMR (376MHz; CD_3OD) : d -151.46.

20 ^{31}P -NMR (162MHz; CD_3OD) : d 7.36.

MS $[\text{M} + \text{H}^+] = 539.4$.

Compound 26



Handwritten signature

5

Compound 26

Compound 26 is prepared from compound **22** in a matter similar to that for preparation of compound **10**.

10 Antiviral Activity

Another aspect of the invention relates to methods of inhibiting viral infections, comprising the step of treating a sample or subject suspected of needing such inhibition with a composition of the invention.

- 15 Within the context of the invention samples suspected of containing a virus include natural or man-made materials such as living organisms; tissue or cell cultures; biological samples such as biological material samples (blood, serum, urine, cerebrospinal fluid, tears, sputum, saliva, tissue samples, and the like); laboratory samples; food, water, or air samples; bioproduct samples such as extracts of cells, particularly recombinant cells synthesizing a desired glycoprotein; and the like.
- 20 Typically the sample will be suspected of containing an organism which induces a viral infection, frequently a pathogenic organism such as a tumor virus. Samples can be contained in any medium including water and organic solvent\water mixtures. Samples include living organisms such as humans, and man made materials such as cell cultures.

- 25 If desired, the anti-virus activity of a compound of the invention after application of the composition can be observed by any method including direct and indirect methods of detecting such activity. Quantitative, qualitative, and semiquantitative methods of determining such activity are all contemplated. Typically one of the screening methods described above are applied, however, any
- 30 other method such as observation of the physiological properties of a living organism are also applicable.

The antiviral activity of a compound of the invention can be measured using standard screening protocols that are known. For example, the antiviral activity of a



- 5 compound can be measured using the following general protocols.

Cell-based Flavivirus Immunodetection assay

BHK21 or A549 cells are trypsinized, counted and diluted to 2×10^5 cells/mL in Hams F-12 media (A549 cells) or RPMI-1640 media (BHK21 cells) supplemented with 2% fetal bovine serum (FBS) and 1% penicillin/streptomycin. 2×10^4 cells are
10 dispensed in a clear 96-well tissue culture plates per well and placed at 37°C , 5% CO_2 overnight. On the next day, the cells are infected with viruses at multiplicity of infection (MOI) of 0.3 in the presence of varied concentrations of test compounds for 1 hour at 37°C and 5% CO_2 for another 48 hours. The cells are washed once with PBS and fixed with cold methanol for 10 min. After washing twice with PBS, the
15 fixed cells are blocked with PBS containing 1% FBS and 0.05% Tween-20 for 1 hour at room temperature. The primary antibody solution (4G2) is then added at a concentration of 1:20 to 1:100 in PBS containing 1% FBS and 0.05% Tween-20 for 3 hours. The cells are then washed three times with PBS followed by one hour incubation with horseradish peroxidase(HRP)-conjugated anti-mouse IgG (Sigma,
20 1:2000 dilution). After washing three times with PBS, 50 microliters of 3,3',5,5'-tetramethylbenzidine (TMB) substrate solution (Sigma) is added to each well for two minutes. The reaction is stopped by addition of 0.5 M sulfuric acid. The plates are read at 450 nm absorbance for viral load quantification. After measurement, the cells are washed three times with PBS followed by incubation with propidium iodide for 5
25 min. The plate is read in a Tecan SafireTM reader (excitation 537 nm, emission 617 nm) for cell number quantification. Dose response curves are plotted from the mean absorbance versus the log of the concentration of test compounds. The EC_{50} is calculated by non-linear regression analysis. A positive control such as N-nonyl-deoxynojirimycin may be used.

30 Cell-based Flavivirus cytopathic effect assay

For testing against West Nile virus or Japanese encephalitis virus, BHK21 cells are trypsinized and diluted to a concentration of 4×10^5 cells/mL in RPMI-1640



5 media supplemented with 2% FBS and 1% penicillin/streptomycin. For testing against dengue virus, Huh7 cells are trypsinized and diluted to a concentration of 4×10^5 cells/mL in DMEM media supplemented with 5% FBS and 1% penicillin/streptomycin. A 50 microliter of cell suspension (2×10^4 cells) is dispensed per well in a 96-well optical bottom PIT polymer-based plates (Nunc). Cells are
10 grown overnight in culture medium at 37°C, 5% CO₂, and then infected with West Nile virus (e.g. B956 strain) or Japanese encephalitis virus (e.g. Nakayama strain) at MOI = 0.3, or with dengue virus (e.g. DEN-2 NGC strain) at MOI = 1, in the presence of different concentrations of test compounds. The plates containing the virus and the compounds are further incubated at 37°C, 5% CO₂ for 72 hours. At the end of
15 incubation, 100 microliters of CellTiter-Glo™ reagent is added into each well. Contents are mixed for 2 minutes on an orbital shaker to induce cell lysis. The plates are incubated at room temperature for 10 minutes to stabilize luminescent signal. Luminescence reading is recorded using a plate reader. A positive control such as N-nonyl-deoxynojirimycin may be used.

20 Antiviral Activity in a Mouse Model of Dengue Infection.

Compounds are tested *in vivo* in a mouse model of dengue virus infection (Schul *et al.* J. Infectious Dis. 2007; 195:665-74). Six to ten week old AG129 mice (B&K Universal Ltd, Hill, UK) are housed in individually ventilated cages. Mice are injected intraperitoneally with 0.4 mL TSV01 dengue virus 2 suspension. Blood
25 samples are taken by retro orbital puncture under isoflurane anesthesia. Blood samples are collected in tubes containing sodium citrate to a final concentration of 0.4%, and immediately centrifuged for 3 minutes at 6000g to obtain plasma. Plasma (20 microliters) is diluted in 780 microliters RPMI-1640 medium and snap frozen in liquid nitrogen for plaque assay analysis. The remaining plasma is reserved for
30 cytokine and NS1 protein level determination. Mice develop dengue viremia rising over several days, peaking on day 3 post-infection.

For testing of antiviral activity, a compound of the invention is dissolved in



- 5 vehicle fluid, e.g. 10% ethanol, 30% PEG 300 and 60% D5W (5% dextrose in water; or 6N HCl (1.5 eq):1N NaOH (pH adjusted to 3.5): 100 mM citrate buffer pH 3.5 (0.9% v/v:2.5% v/v: 96.6% v/v). Thirty six 6-10 week old AG129 mice are divided into six groups of six mice each. All mice are infected with dengue virus as described above (day 0). Group 1 is dosed by oral gavage of 200 μ L/mouse with 0.2 mg/kg of
10 a compound of the invention twice a day (once early in the morning and once late in the afternoon) for three consecutive days starting on day 0 (first dose just before dengue infection). Groups 2, 3 and 4 are dosed the same way with 1 mg/kg, 5 mg/kg and 25 mg/kg of the compound, respectively. A positive control may be used, such as
15 (2R,3R,4R,5R)-2-(2-amino-6-hydroxy-purin-9-yl)-5-hydroxymethyl-3-methyl-tetrahydro-furan-3,4-diol, dosed by oral gavage of 200 microliters/mouse the same way as the previous groups. A further group is treated with only vehicle fluid.

On day 3 post-infection approximately 100 microliter blood samples (anti-coagulated with sodium citrate) are taken from the mice by retro-orbital puncture under isoflurane anesthesia. Plasma is obtained from each blood sample by
20 centrifugation and snap frozen in liquid nitrogen for plague assay analysis. The collected plasma samples are analyzed by plague assay as described in Schul *et al.* Cytokines are also analyzed as described by Schul. NS1 protein levels are analyzed using a PlateliaTM kit (BioRad Laboratories). An anti-viral effect is indicated by a reduction in cytokine levels and/or NS1 protein levels.

25 Typically, reductions in viremia of about 5-100 fold, more typically 10-60 fold, most typically 20-30 fold, are obtained with 5-50 mg/kg bid dosages of the compounds of the invention.

HCV IC₅₀ Determination

30 Assay Protocol: Either wild type or S282T (Migliaccio, et al, *J. Biol. Chem.* **2003**, 49164-49170; Klumpp, et al., *J. Biol. Chem.* **2006**, 3793-3799) mutant polymerase enzyme was used in this assay. NS5b polymerase assay (40 μ L) was







L

5 assembled by adding 28 μ L polymerase mixture (final concentration: 50 mM Tris-HCl at pH 7.5, 10 mM KCL, 5 mM $MgCl_2$, 1 mM DTT, 10 mM EDTA, 4 ng/ μ L of RNA template, and 75 nM HCV Δ 21 NS5b polymerase) to assay plates followed by 4 μ L of compound dilution. The polymerase and compound were pre-incubated at 35°C for 10 minute before the addition of 8 μ L of nucleotide substrate mixture (33P-
10 α -labeled competing nucleotide at K_M and 0.5 mM of the remaining three nucleotides). The assay plates were covered and incubated at 35°C for 90 min. Reactions were then filtered through 96-well DEAE-81 filter plates via vacuum. The filter plates were then washed under vacuum with multiple volumes of 0.125 M $NaHPO_4$, water, and ethanol to remove unincorporated label. Plates were then
15 counted on TopCount to assess the level of product synthesis over background controls. The IC₅₀ value is determined using Prism fitting program.

Preferably, compounds described herein inhibited NS5b polymerase with an IC₅₀'s below 1000 μ M, more preferably below 100 μ M, and most preferably below 10 μ M. For example, compound **TP-1** has an IC₅₀ of 0.15 μ M against both wild type
20 HCV polymerase and the S282T mutant enzyme. Table II below shows the activity of **TP-1** and **TP-2** against both wild type and the S282T mutant enzyme compared to the activities obtained with the triphosphate of 2'-methyl guanidine and the triphosphate of (2R,3R,4R,5R)-2-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-3,4-dihydroxy-5-(hydroxymethyl)-3-methyl-tetrahydrofuran-2-carbonitrile. This
25 demonstrates that replacing the 2' OH of the pyrrolo[1,2-f][1,2,4]triazin-7-yl nucleosides with a 2' F unexpectedly confers activity against resistant S282T HCV mutant strains of virus.

Table II

Triphosphate	WT IC ₅₀ (μ M)	S282T IC ₅₀ (μ M)	Note
--------------	-----------------------------------	--------------------------------------	------

 2'-C-MeGTP	0.1	20	from <i>J. Bio. Chem.</i> , 2003, 278, 49164 (200 fold shift)
 TP-1	0.15	0.15	(1 fold shift)
 TP-2	0.525	111	WO/2009/132135 (242 fold shift)
 TP-3	0.24	1.60	(7 fold shift)
 TP-8a	0.034		
 TP-8a	0.30	1.6	(5.3 fold shift)



5 HCV EC₅₀ Determination

Replicon cells were seeded in 96-well plates at a density of 8×10^3 cells per well in 100 μ L of culture medium, excluding Geneticin. Compound was serially diluted in 100% DMSO and then added to the cells at a 1:200 dilution, achieving a final concentration of 0.5% DMSO and a total volume of 200 μ L. Plates were incubated at 37°C for 3 days, after which culture medium was removed and cells were lysed in lysis buffer provided by Promega's luciferase assay system. Following the manufacturer's instruction, 100 μ L of luciferase substrate was added to the lysed cells and luciferase activity was measured in a TopCount luminometer. Preferably, compounds described herein have EC₅₀'s below 1000 μ M, more preferably below 100 μ M, and most preferably below 10 μ M. The activities of representative compounds of Formula I are shown in the Table III below.

Table III

Compound No.	EC ₅₀ , μ M
A-1	23
B-1	1.4-4.3
B-3	16-28
B-4	8.4-19
B-5	1.93-25.5
B-6	3.75-11.1
B-7	63-73
B-8	35-60
C-1	67-70
C-2	3.9-12
C-3	43-84
C-4	9.8-31
C-5	24-28



C-6	11
10	6.5-8
22	31-45
23	39.4-40.3
24	40.3-70.5
25	9.7-10
PD-A-8b	0.68

5

The cytotoxicity of a compound of the invention can be determined using the following general protocol.

Metabolism Studies:

10 Applicants have observed that monophosphate prodrugs of nucleoside analogs with a nitrogen at the X¹ position can have enhanced activity over their counterparts with a carbon at the X¹ position. This difference in activity correlates to the amount of the active triphosphate analogs of the compounds in cells. This can be quantified by a metabolism study which quantifies the intracellular concentration of the
15 triphosphate analogs. The higher intracellular concentration of the triphosphate metabolite correlates to the prodrug with enhanced activity.

For example, comparison of the prodrug compound **B-7** with prodrug compound **PD-A-8b** shows increased activity when the X¹ position is nitrogen. This can be observed in Table III, where the HCV EC₅₀ for the compound where the X¹
20 position is nitrogen (compound **PD-A-8b**) is 0.68 μ M compared to 63-73 μ M for compound **B-7**. The activation of prodrug analog **PD-A-8b** (to its triphosphate analog **TP-8a**) was found to be more than two orders of magnitude more efficient than that observed for its prodrug counterpart where the X¹ position is carbon, **B-7** (to its triphosphate analog **TP-3**), as seen in Table IV.

25

Experimental:



5 Huh-luc/neo replicon cells containing HCV genotype 1b subgenomic replicons were maintained in Dulbecco's modified eagle medium containing glutamax supplemented with 10% heat inactivated fetal bovine serum, penicillin-streptomycin, and G418 disulphate salt solution. Cells were transferred to twelve well tissue culture plates by trypsonization and grown to confluency (0.88×10^6 cells/well). Cells were
 10 treated for 24 hours with 10 μ M nucleoside, or 10 μ M prodrug. After 24 hours, cells were washed 2 times with 2.0 mL ice cold 0.9% sodium chloride saline. Cells were then scraped into 0.5 mL 70% methanol (MeOH) and frozen overnight to facilitate the extraction of nucleotide metabolites. Extracted cell material in 70% MeOH was transferred into tubes and dried. After drying, samples were resuspended in 1mM
 15 Ammonium phosphate pH 8.5 containing internal standard (100 nM ClATP). Intracellular levels of the nucleoside triphosphates were quantified based on authentic standard curves by liquid chromatography coupled to tandem mass spectrometry.

Results

20

Table IV: Intracellular triphosphate analog concentrations formed in Huh-luc/neo replicon cells following 24 hour incubations with 10 μ M **PD-A-8b** and **B-7**.

Prodrug	Triphosphate	Intracellular Triphosphate Analog Concentration (pmol/million)
B-7	TP-3	< 0.11 ^a
PD-A-8b	TP-8a	20.5 ^b
^a Intracellular concentrations were below the lower limit of quantification of the assay.		
^b Value is the average of results from 2 separate wells.		

25 Cytotoxicity Cell Culture Assay (Determination of CC50):

The assay is based on the evaluation of cytotoxic effect of tested compounds using a metabolic substrate.

Assay protocol for determination of CC50:

1. Maintain MT-2 cells in RPMI-1640 medium supplemented with 5% fetal bovine
 30 serum and antibiotics.



- 5 2. Distribute the cells into a 96-well plate (20,000 cell in 100 μ L media per well) and add various concentrations of the tested compound in triplicate (100 μ L/well). Include untreated control.
3. Incubate the cells for 5 days at 37°C.
4. Prepare XTT solution (6 ml per assay plate) in dark at a concentration of 2mg/ml
10 in a phosphate-buffered saline pH 7.4. Heat the solution in a water-bath at 55°C for 5 min. Add 50 μ L of N-methylphenazonium methasulfate (5 μ g/mL) per 6 mL of XTT solution.
5. Remove 100 μ L media from each well on the assay plate and add 100 μ L of the XTT substrate solution per well. Incubate at 37°C for 45 to 60 min in a CO₂
15 incubator.
6. Add 20 μ L of 2% Triton X-100 per well to stop the metabolic conversion of XTT.
7. Read the absorbance at 450 nm with subtracting off the background at 650 nm.
8. Plot the percentage absorbance relative to untreated control and estimate the CC50 value as drug concentration resulting in a 50% inhibition of the cell growth.
- 20 Consider the absorbance being directly proportional to the cell growth.


All publications, patents, and patent documents cited herein above are incorporated by reference herein, as though individually incorporated by reference. The invention has been described with reference to various specific and preferred
25 embodiments and techniques. However, one skilled in the art will understand that many variations and modifications may be made while remaining within the spirit and scope of the invention.



15/03/2013

5 What is claimed is

1. A compound of Formula IV:



Formula IV

10 or a pharmaceutically acceptable salt, thereof;

wherein:

R^1 is (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl (C_1-C_8) alkyl;

15 R^2 is halogen;


R^3 , R^4 , and R^5 are each independently H, halogen, OR^a , $N(R^a)_2$, N_3 , CN, NO_2 , $S(O)_nR^a$, (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl (C_1-C_8) alkyl;

20 or any two of R^3 , R^4 or R^5 on adjacent carbon atoms when taken together are $-O(CO)O-$ or when taken together with the ring carbon atoms to which they are attached form a double bond;

each n is independently 0, 1, or 2;


25 each R^a is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, or $-SO_2NR^{11}R^{12}$;

- 5 R^7 is H, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, $-SO_2NR^{11}R^{12}$, or



- 10 Y is O, S, NR, $^+N(O)(R)$, N(OR), $^+N(O)(OR)$, or N-NR₂;
 W^1 and W^2 , when taken together, are $-Y^3(C(R^y)_2)_3Y^3-$; or
 one of W^1 or W^2 together with either R^3 or R^4 is $-Y^3-$ and the other of W^1 or W^2 is Formula Ia; or

W^1 and W^2 are each, independently, a group of Formula IVa:



15

Formula IVa

wherein:

each Y^1 is, independently, O, S, NR, $^+N(O)(R)$, N(OR), $^+N(O)(OR)$, or N-NR₂;

- 20 each Y^2 is independently a bond, O, CR₂, NR, $^+N(O)(R)$, N(OR), $^+N(O)(OR)$, N-NR₂, S, S-S, S(O), or S(O)₂;


each Y^3 is independently O, S, or NR;

M2 is 0, 1 or 2;

each R^x is a group of Formula IVb:

Handwritten signature or mark.

190



Formula IVb

wherein:

each M1a, M1c, and M1d is independently 0 or 1;

M12c is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

each R^y is independently H, F, Cl, Br, I, OH, $-C(=Y^1)R$, $-C(=Y^1)R^{13}$, $-C(=Y^1)OR$, $-C(=Y^1)N(R)_2$, $-N(R)_2$, $-N(R)_3$, $-SR$, $-S(O)R$, $-S(O)_2R$, $-S(O)_2R^{13}$, $-S(O)(OR)$, $-S(O)_2(OR)$, $-OC(=Y^1)R$, $-OC(=Y^1)OR$, $-OC(=Y^1)(N(R)_2)$, $-SC(=Y^1)R$, $-SC(=Y^1)OR$, $-SC(=Y^1)(N(R)_2)$, $-N(R)C(=Y^1)R$, $-N(R)C(=Y^1)OR$, $-N(R)C(=Y^1)N(R)_2$, $-SO_2NR_2$, $-CN$, $-N_3$, $-NO_2$, $-OR$, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, C_6-C_{20} aryl, C_3-C_{20} carbocyclyl, C_2-C_{20} heterocyclyl, arylalkyl, heteroarylalkyl;

wherein each (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, C_6-C_{20} aryl,

C_3-C_{20} carbocyclyl, C_2-C_{20} heterocyclyl, arylalkyl, or heteroarylalkyl

is optionally substituted with 1-3 R^{20} groups;

or when taken together, two R^y on the same carbon atom form a carbocyclic ring of 3 to 7 carbon atoms;

each R is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, C_6-C_{20} aryl, C_3-C_{20} carbocyclyl, C_2-C_{20} heterocyclyl, or arylalkyl;

each R^8 is halogen, $NR^{11}R^{12}$, $N(R^{11})OR^{11}$, $NR^{11}NR^{11}R^{12}$, N_3 , NO, NO_2 , OR^{11} or $S(O)_nR^{11}$;

each R^9 is independently H, halogen, $NR^{11}R^{12}$, $N(R^{11})OR^{11}$, $NR^{11}NR^{11}R^{12}$, N_3 , NO, NO_2 , CHO, CN, $-CH(=NR^{11})$, $-CH=NHNR^{11}$, $-CH=N(OR^{11})$, $-CH(OR^{11})_2$, $-C(=O)NR^{11}R^{12}$, $-C(=S)NR^{11}R^{12}$, $-C(=O)OR^{11}$, R^{11} , OR^{11} or $S(O)_nR^{11}$;

each R^{11} or R^{12} is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, (C_4-C_8) carbocyclylalkyl, optionally substituted aryl, optionally

5 substituted heteroaryl, $-C(=O)(C_1-C_8)alkyl$, $-S(O)_n(C_1-C_8)alkyl$ or $aryl(C_1-C_8)alkyl$; or R^{11} and R^{12} taken together with a nitrogen to which they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with $-O-$, $-S-$ or $-NR^b$;

each R^{13} is independently a carbocycle or heterocycle optionally substituted
10 with 1-3 R^{20} groups;

each R^{20} is independently, halogen, CN , N_3 , $N(R)_2$, OR , $-SR$, $-S(O)R$, $-S(O)_2R$, $-S(O)(OR)$, $-S(O)_2(OR)$, $-C(=Y^1)R$, $-C(=Y^1)OR$, or $C(=Y^1)N(R)_2$;

wherein each $(C_1-C_8)alkyl$, $(C_2-C_8)alkenyl$, $(C_2-C_8)alkynyl$ or $aryl(C_1-C_8)alkyl$ of each R^1 , R^3 , R^4 , R^5 , R^6 , R^{11} or R^{12} is, independently, optionally substituted with one
15 or more halo, hydroxy, CN , N_3 , $N(R^b)_2$ or OR^b ; and wherein one or more of the non-terminal carbon atoms of each said $(C_1-C_8)alkyl$ may be optionally replaced with $-O-$, $-S-$ or $-NR^b$;

each R^b is independently H , $(C_1-C_8)alkyl$, $(C_2-C_8)alkenyl$, $(C_2-C_8)alkynyl$, $aryl(C_1-C_8)alkyl$, $(C_4-C_8)carbocyclalkyl$, $-C(=O)R^{21}$, $-C(=O)OR^{21}$, $-C(=O)NR^{21}R^{22}$,
20 $-C(=O)SR^{21}$, $-S(O)R^{21}$, $-S(O)_2R^{21}$, $-S(O)(OR^{21})$, $-S(O)_2(OR^{21})$, or $-SO_2NR^{21}R^{22}$; and

each R^{21} or R^{22} is independently H , $(C_1-C_8)alkyl$, $(C_2-C_8)alkenyl$, $(C_2-C_8)alkynyl$, $(C_4-C_8)carbocyclalkyl$, $-C(=O)(C_1-C_8)alkyl$, $-S(O)_n(C_1-C_8)alkyl$ or $aryl(C_1-C_8)alkyl$.

25 2. The compound according to claim 1 wherein each Y and Y^1 is O .

3. The compound according to claim 2 wherein R^8 is halogen, $NR^{11}R^{12}$, $N(R^{11})OR^{11}$, $NR^{11}NR^{11}R^{12}$, OR^{11} or $S(O)_nR^{11}$.


30 4. The compound according to claim 3 wherein R^9 is H , halogen, $S(O)_nR^{11}$ or $NR^{11}R^{12}$.

5. The compound according to claim 4 wherein R^4 is OR^a .




- 5 6. The compound according to claim 5 wherein R^1 is CH_3 .
7. The compound according to claim 6 wherein R^2 is F.
8. The compound according to claim 7 wherein R^7 is

10



wherein Y is $-O-$; W^1 is Formula Ia and W^2 together with R^4 is $-O-$.

- 15 9. The compound according to claim 1 represented by Formula V:




Formula V

- 20 wherein R^1 is methyl or ethynyl, and R^4 is OR^a .

10. The compound according to claim 9 wherein R^7 is H or






193




5

11. The compound according to claim 1 with the following structures:

10




194



5

12. A compound of Formula VI:



10

Formula VI

or a pharmaceutically acceptable salt, thereof;

wherein:

15 R^4 is OR^a ;

each n is independently 0, 1, or 2;

each R^a is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl (C_1-C_8) alkyl, (C_4-C_8) carbocyclylalkyl, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, or $-SO_2NR^{11}R^{12}$;

20 R^7 is H, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, $-SO_2NR^{11}R^{12}$, or

α



wherein:

each M1a, M1c, and M1d is independently 0 or 1;

M12c is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;


- 10 each R^y is independently H, F, Cl, Br, I, OH, -C(=Y¹)R, -C(=Y¹)R¹³, -C(=Y¹)OR, -C(=Y¹)N(R)₂, -N(R)₂, -⁺N(R)₃, -SR, -S(O)R, -S(O)₂R, -S(O)₂R¹³, -S(O)(OR), -S(O)₂(OR), -OC(=Y¹)R, -OC(=Y¹)OR, -OC(=Y¹)(N(R)₂), -SC(=Y¹)R, -SC(=Y¹)OR, -SC(=Y¹)(N(R)₂), -N(R)C(=Y¹)R, -N(R)C(=Y¹)OR, -N(R)C(=Y¹)N(R)₂, -SO₂NR₂, -CN, -N₃, -NO₂, -OR, (C₁-C₈) alkyl, (C₂-C₈) alkenyl, (C₂-C₈) alkynyl,
- 15 C₆-C₂₀ aryl, C₃-C₂₀ carbocyclyl, C₂-C₂₀ heterocyclyl, arylalkyl, heteroarylalkyl; wherein each (C₁-C₈) alkyl, (C₂-C₈) alkenyl, (C₂-C₈) alkynyl, C₆-C₂₀ aryl, C₃-C₂₀ carbocyclyl, C₂-C₂₀ heterocyclyl, arylalkyl, or heteroarylalkyl is optionally substituted with 1-3 R²⁰ groups;
- each R is independently H, (C₁-C₈) alkyl, (C₂-C₈) alkenyl, (C₂-C₈) alkynyl,
- 20 C₆-C₂₀ aryl, C₃-C₂₀ carbocyclyl, C₂-C₂₀ heterocyclyl, or arylalkyl; each R¹¹ or R¹² is independently H, (C₁-C₈) alkyl, (C₂-C₈) alkenyl, (C₂-C₈) alkynyl, (C₄-C₈) carbocyclylalkyl, optionally substituted aryl, optionally substituted heteroaryl, -C(=O)(C₁-C₈) alkyl, -S(O)_n(C₁-C₈) alkyl or aryl(C₁-C₈) alkyl; each R¹³ is independently a carbocycle or heterocycle optionally substituted
- 25 with 1-3 R²⁰ groups; each R²⁰ is independently, halogen, CN, N₃, N(R)₂, OR, -SR, -S(O)R, -S(O)₂R, -S(O)(OR), -S(O)₂(OR), -C(=Y¹)R, -C(=Y¹)OR, or C(=Y¹)N(R)₂;
- wherein each (C₁-C₈) alkyl, (C₂-C₈) alkenyl, (C₂-C₈) alkynyl or aryl(C₁-C₈) alkyl of each R⁴, R¹¹ or R¹² is, independently, optionally substituted with one or more halo, hydroxy, CN, N₃, N(R^b)₂ or OR^b; and wherein one or more of the non-terminal carbon
- 30 atoms of each said (C₁-C₈) alkyl may be optionally replaced with -O-, -S- or -NR^b; each R^b is independently H, (C₁-C₈) alkyl, (C₂-C₈) alkenyl, (C₂-C₈) alkynyl, aryl(C₁-C₈) alkyl, (C₄-C₈) carbocyclylalkyl, -C(=O)R²¹, -C(=O)OR²¹, -C(=O)NR²¹R²², -C(=O)SR²¹, -S(O)R²¹, -S(O)₂R²¹, -S(O)(OR²¹), -S(O)₂(OR²¹), or -SO₂NR²¹R²², and

- 5 each R^{21} or R^{22} is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, (C_4-C_8) carbocyclalkyl, $-C(=O)(C_1-C_8)$ alkyl, $-S(O)_n(C_1-C_8)$ alkyl or aryl (C_1-C_8) alkyl.

13. The compound according to claim 12 wherein:

10 R^a is H, (C_1-C_8) alkyl, or $-C(=O)(C_1-C_6)$ alkyl;

R^7 or R^7 together with R^4 is



wherein

a is the point of attachment to R^7 ;

b is the point of attachment to R^4 ;

20 Ar is phenyl or naphthyl, wherein the phenyl and naphthyl are optionally substituted with 1-3 R^{20} groups;


each R^y is independently (C_1-C_8) alkyl or C_5-C_6 carbocycl, wherein the alkyl and carbocycl are optionally substituted with 1-3 R^{20} groups;

each R is independently H, (C_1-C_6) alkyl, or arylalkyl; and

L



- 5 each R^{20} is independently halogen, CN, $N(R)_2$, OR, -SR, -S(O)R, -S(O) $_2$ R, -S(O)(OR), -S(O) $_2$ (OR), -C(=O)R, -C(=O)OR, or C(=O) $N(R)_2$.

14. A compound having a structure:






199



5





10



5



10



or a pharmaceutically acceptable salt, thereof.

15

15. A pharmaceutical composition comprising a therapeutically effective amount of a compound of claim 1 and a pharmaceutically acceptable carrier.

L

16. The pharmaceutical composition of claim 15 further comprising at least one additional therapeutic agent selected from the group consisting of interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, endothelin antagonists, other anti-fibrotic agents, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers and other drugs for treating HCV; or mixtures thereof.

17. Use of a compound of claim 1 in the manufacture of a pharmaceutical composition for treating a *Flaviviridae* virus infection in a mammal in need thereof, wherein said composition is for administration of a therapeutically effective amount of said compound of claim 1.

18. The use of claim 17 wherein the viral infection is a Hepatitis C virus infection.

19. The use of claim 18 wherein the viral infection is caused by a S282T mutant of Hepatitis C virus.

20. The use of claim 17 further comprising administration of at least one additional therapeutic agent selected from the group consisting of interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, endothelin antagonists, other anti-fibrotic agents, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers and other drugs for treating HCV; or mixtures thereof.

