



(86) Date de dépôt PCT/PCT Filing Date: 2011/06/17
(87) Date publication PCT/PCT Publication Date: 2011/12/22
(85) Entrée phase nationale/National Entry: 2012/12/11
(86) N° demande PCT/PCT Application No.: EP 2011/002993
(87) N° publication PCT/PCT Publication No.: 2011/157430
(30) Priorité/Priority: 2010/06/18 (US61/356,292)

(51) Cl.Int./Int.Cl. *C12N 9/12* (2006.01)
(71) Demandeur/Applicant:
F. HOFFMAN-LA ROCHE AG, CH
(72) Inventeurs/Inventors:
REICHERT, FRED, US;
BAUER, KEITH, US;
MYERS, THOMAS W., US
(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : ADN POLYMERASES PRESENTANT UNE DISCRIMINATION ACCRUE DES MESAPPARIEMENTS A L'EXTREMITÉ 3'
(54) Title: DNA POLYMERASES WITH INCREASED 3'-MISMATCH DISCRIMINATION

Figure 1

★

Z05	EEEVFRLAGH P FN L N S R D Q L E R V L F D E LRLPALGKT	(SEQ ID NO:12)
Taq	EAEVFRLAGH P FN L N S R D Q L E R V L F D E LGLPAIGKT	(SEQ ID NO:13)
Tfi	EAEVHRLAGH P FN L N S R D Q L E R V L F D E LGLPPIGKT	(SEQ ID NO:14)
Tfl	EEEVFRLAGH P FN L N S R D Q L E R V L F D E LGLPAIGKT	(SEQ ID NO:15)
Sps17	EAEVHRLAGH P FN L N S R D Q L E R V L F D E LGLPPIGKT	(SEQ ID NO:16)
Tth	EEEVFRLAGH P FN L N S R D Q L E R V L F D E LRLPALGKT	(SEQ ID NO:17)
Tca	EEEVFRLAGH P FN L N S R D Q L E R V L F D E LRLPALGKT	(SEQ ID NO:18)
Tma	AEEIYRIAGE P FN I N S P K Q V S R I L F E K LGIKPRGKT	(SEQ ID NO:19)
Tne	AEKIYQIAGE P FN I N S P K Q V S N I L F E K LGIKPRGKT	(SEQ ID NO:20)
Taf	KEKVFEIAGE T FN L N S S T Q V A Y I L F E K LNIAPY-KK	(SEQ ID NO:21)
Dra	ESQIHEYAGE E F H I R S P K Q L E T V L Y D K LELASSKKT	(SEQ ID NO:23)
Bst	ERRIYELAGQ E FN I N S P K Q L G T V L F D K LQLPVLKKT	(SEQ ID NO:24)
Bca	EQRIYELAGQ E FN I N S P K Q L G V I L F E K LQLPVLKKS	(SEQ ID NO:25)
	-----AGX ₁ X ₂ FX ₃ X ₄ X ₅ X ₆ X ₇ X ₈ QX ₉ X ₁₀ X ₁₁ X ₁₂ LX ₁₃ X ₁₄ X ₁₅ L-----	(SEQ ID NO:26)

(57) Abrégé/Abstract:

Disclosed are mutant DNA polymerases having increased 3'-mismatch discrimination relative to a corresponding, unmodified polymerase. The mutant polymerases are useful in a variety of disclosed primer extension methods. Also disclosed are related compositions, including recombinant nucleic acids, vectors, and host cells, which are useful, e.g., for production of the mutant DNA polymerases.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
22 December 2011 (22.12.2011)



(10) International Publication Number
WO 2011/157430 A1

(51) International Patent Classification:
C12N 9/12 (2006.01)

(21) International Application Number:
PCT/EP2011/002993

(22) International Filing Date:
17 June 2011 (17.06.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/356,292 18 June 2010 (18.06.2010) US

(71) Applicant (for DE only): **ROCHE DIAGNOSTICS GMBH** [DE/DE]; Sandhoferstrasse 116, 68305 Mannheim (DE).

(71) Applicant (for all designated States except DE, US): **F. HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basel (CH).

(72) Inventors: **REICHERT, Fred**; 675 Haas Avenue, San Leandro, CA 94577-2929 (US). **BAUER, Keith**; 62 W. Crescent Drive, San Rafael, CA 94901 (US). **MYERS, Thomas, W.**; 600 Loyola Drive, Los Altos, CA 94204 (US).

(74) Agent: **ROCHE DIAGNOSTICS GMBH**; -Patent Department-, 68298 Mannheim (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

(54) Title: DNA POLYMERASES WITH INCREASED 3'-MISMATCH DISCRIMINATION

Figure 1

*

Z05	EEEVFRLAGH P FN L N S R D Q L E R V L F D E LRLPALGKT (SEQ ID NO:12)
Taq	EA EVFRLAGH P FN L N S R D Q L E R V L F D E LGLPAIGKT (SEQ ID NO:13)
Tfi	EA EVHRLAGH P FN L N S R D Q L E R V L F D E LGLPPIGKT (SEQ ID NO:14)
Tfl	EEEVFRLAGH P FN L N S R D Q L E R V L F D E LGLPAIGKT (SEQ ID NO:15)
Sps17	EA EVHRLAGH P FN L N S R D Q L E R V L F D E LGLPPIGKT (SEQ ID NO:16)
Tth	EEEVFRLAGH P FN L N S R D Q L E R V L F D E LRLPALGKT (SEQ ID NO:17)
Tca	EEEVFRLAGH P FN L N S R D Q L E R V L F D E LRLPALGKT (SEQ ID NO:18)
Tma	AEIYRIAGE P FN I N S P K Q V S R I L F E K LGIKPRGKT (SEQ ID NO:19)
Tne	AEKIYQIAGE P FN I N S P K Q V S N I L F E K LGIKPRGKT (SEQ ID NO:20)
Taf	KEKVFEIAGE T FN L N S S T Q V A Y I L F E K LNIAPY-KK (SEQ ID NO:21)
Dra	ESQIHEYAGE E FH I R S P K Q L E T V L Y D K LELASSKKT (SEQ ID NO:23)
Bst	ERRIYELAGQ E FN I N S P K Q L G T V L F D K LQLPVLKKT (SEQ ID NO:24)
Bca	EQRIYELAGQ E FN I N S P K Q L G V I L F E K LQLPVLKKS (SEQ ID NO:25)
	-----AGX ₁ X ₂ FX ₃ X ₄ X ₅ X ₆ X ₇ X ₈ QX ₉ X ₁₀ X ₁₁ X ₁₂ LX ₁₃ X ₁₄ X ₁₅ L----- (SEQ ID NO:26)

(57) Abstract: Disclosed are mutant DNA polymerases having increased 3'-mismatch discrimination relative to a corresponding, unmodified polymerase. The mutant polymerases are useful in a variety of disclosed primer extension methods. Also disclosed are related compositions, including recombinant nucleic acids, vectors, and host cells, which are useful, e.g., for production of the mutant DNA polymerases.

DNA POLYMERASES WITH INCREASED 3'-MISMATCH DISCRIMINATION

FIELD OF THE INVENTION

5 The present invention provides DNA polymerases with increased 3'-mismatch discrimination and their use in various applications, including nucleic acid polynucleotide extension and amplification.

BACKGROUND OF THE INVENTION

DNA polymerases are responsible for the replication and maintenance of the genome, a role
10 that is central to accurately transmitting genetic information from generation to generation. DNA polymerases function in cells as the enzymes responsible for the synthesis of DNA. They polymerize deoxyribonucleoside triphosphates in the presence of a metal activator, such as Mg^{2+} , in an order dictated by the DNA template or polynucleotide template that is copied. *In vivo*, DNA polymerases participate in a spectrum of DNA synthetic processes including
15 DNA replication, DNA repair, recombination, and gene amplification. During each DNA synthetic process, the DNA template is copied once or at most a few times to produce identical replicas. In contrast, *in vitro*, DNA replication can be repeated many times such as, for example, during polymerase chain reaction (*see, e.g.*, U.S. Patent No. 4,683,202).

In the initial studies with polymerase chain reaction (PCR), the DNA polymerase was added at
20 the start of each round of DNA replication (*see* U.S. Patent No. 4,683,202, *supra*). Subsequently, it was determined that thermostable DNA polymerases could be obtained from bacteria that grow at elevated temperatures, and that these enzymes need to be added only once (*see* U.S. Patent No. 4,889,818 and U.S. Patent No. 4,965,188). At the elevated temperatures used during PCR, these enzymes are not irreversibly inactivated. As a result, one
25 can carry out repetitive cycles of polymerase chain reactions without adding fresh enzymes at the start of each synthetic addition process. DNA polymerases, particularly thermostable polymerases, are the key to a large number of techniques in recombinant DNA studies and in medical diagnosis of disease. For diagnostic applications in particular, a target nucleic acid

sequence may be only a small portion of the DNA or RNA in question, so it may be difficult to detect the presence of a target nucleic acid sequence without amplification.

The overall folding pattern of DNA polymerases resembles the human right hand and contains three distinct subdomains of palm, fingers, and thumb. (See Beese *et al.*, *Science* 260:352-355, 1993); Patel *et al.*, *Biochemistry* 34:5351-5363, 1995). While the structure of the fingers and thumb subdomains vary greatly between polymerases that differ in size and in cellular functions, the catalytic palm subdomains are all superimposable. For example, motif A, which interacts with the incoming dNTP and stabilizes the transition state during chemical catalysis, is superimposable with a mean deviation of about one Å amongst mammalian pol α and prokaryotic pol I family DNA polymerases (Wang *et al.*, *Cell* 89:1087-1099, 1997). Motif A begins structurally at an antiparallel β-strand containing predominantly hydrophobic residues and continues to an α-helix. The primary amino acid sequence of DNA polymerase active sites is exceptionally conserved. In the case of motif A, for example, the sequence DYSQIELR (SEQ ID NO:28) is retained in polymerases from organisms separated by many millions years of evolution, including, *e.g.*, *Thermus aquaticus*, *Chlamydia trachomatis*, and *Escherichia coli*.

In addition to being well-conserved, the active site of DNA polymerases has also been shown to be relatively mutable, capable of accommodating certain amino acid substitutions without reducing DNA polymerase activity significantly. (See, *e.g.*, U.S. Patent No. 6,602,695) Such mutant DNA polymerases can offer various selective advantages in, *e.g.*, diagnostic and research applications comprising nucleic acid synthesis reactions. Thus, there is a need in the art for identification of amino acid positions amenable to mutation to yield improved polymerase activities. The present invention, as set forth herein, meets these and other needs.

25

BRIEF SUMMARY OF THE INVENTION

Provided herein are DNA polymerases having increased 3'-mismatch discrimination relative to a corresponding, unmodified control polymerase, and methods of making and using such DNA polymerases. In some embodiments, the polymerase is a thermostable DNA polymerase. In some embodiments, the DNA polymerase is a thermoactive DNA polymerase. In some

embodiments, the DNA polymerase is derived from a *Thermus* species. In some embodiments, the DNA polymerase is derived from a *Thermotoga* species.

As described herein, according to the invention the amino acid of a DNA polymerase corresponding to position 497 of SEQ ID NO:1 can be mutated from the native amino acid at
5 that position to produce an enzyme having increased 3'-mismatch discrimination relative to a corresponding, unmodified control polymerase. In DNA polymerases derived from a *Thermus* species, the native amino acid corresponding to position 497 of SEQ ID NO:1 is F. In DNA polymerases derived from a *Thermotoga* species, the native amino acid corresponding to position 497 of SEQ ID NO:1 is F. In DNA polymerases derived from other non-*Thermus*,
10 non-*Thermotoga* species, the amino acid corresponding to position 497 of SEQ ID NO:1 can be F or an amino acid other than F, for example, Y. See, Figure 1.

In some embodiments, the amino acid of the DNA polymerase corresponding to position 497 of SEQ ID NO:1 is any amino acid other than F or Y, and the control DNA polymerase has the same amino acid sequence as the DNA polymerase except that the amino acid of the control
15 DNA polymerase corresponding to position 497 of SEQ ID NO:1 is F or Y. For example, in some embodiments, the amino acid at the position corresponding to position 497 of SEQ ID NO:1 is selected from G, A, V, L, I, M, W, P, S, T, C, N, Q, D, E, K, R or H. In some embodiments, the amino acid at the position corresponding to position 497 of SEQ ID NO:1 is selected from A, G, S, T, D or K. In some embodiments, the DNA polymerase of the invention
20 is derived from a *Thermus* species, and the amino acid of the DNA polymerase corresponding to position 497 of SEQ ID NO:1 is any amino acid other than F, and the control DNA polymerase has the same amino acid sequence as the DNA polymerase except that the amino acid of the control DNA polymerase corresponding to position 497 of SEQ ID NO:1 is F. For example, where the DNA polymerase is derived from a *Thermus* species, the amino acid at the
25 position corresponding to position 497 of SEQ ID NO:1 can be selected from G, A, V, L, I, M, W, P, S, T, C, N, Q, D, E, K, R, H, or Y. In some embodiments the amino acid at the position corresponding to position 497 of SEQ ID NO:1 is selected from A, G, S, T, Y, D or K. In some embodiments the amino acid at the position corresponding to position 497 of SEQ ID NO:1 is selected from A, G, S, T, D or K.

In some embodiments, the DNA polymerase of the invention is derived from a *Thermus* species, and the amino acid corresponding to position 497 of SEQ ID NO:1 is an amino acid having a polar, uncharged side-chain (e.g., N, Q, H, S, T, or Y), a nonpolar, uncharged side-chain (other than F, e.g., G, A, L, M, W, P, C, V, or I), a polar, positively charged side-chain (e.g., R or K), or polar, negatively charged side-chain (e.g., D or E) at neutral pH. In some embodiments, the amino acid corresponding to position 497 of SEQ ID NO:1 having a polar, uncharged side-chain is S, T, or Y. In some embodiments, the amino acid corresponding to position 497 of SEQ ID NO:1 having a nonpolar, uncharged side-chain is A or G. In some embodiments, the amino acid corresponding to position 497 of SEQ ID NO:1 having a polar, positively charged side-chain is K. In some embodiments, the amino acid corresponding to position 497 of SEQ ID NO:1 having a polar, negatively charged side-chain is D.

In some embodiments, the DNA polymerase having increased 3'-mismatch discrimination comprises a motif in the polymerase domain comprising

A-G-X₁-X₂-F-X₃-X₄-X₅-S-X₇-X₈-Q-X₉-X₁₀-X₁₁-X₁₂-L-X₁₃-X₁₄-X₁₅-L, wherein:

X₁ is H, E or Q;

X₂ is P, T or E;

X₃ is N or H;

X₄ is L or I;

X₅ is N or R;

X₇ is R, P, or S;

X₈ is D, K or T;

X₉ is L or V;

X₁₀ is E, S, A or G;

X₁₁ is R, N, Y, T or V;

X₁₂ is V or I;

X₁₃ is any amino acid other than F or Y;

X₁₄ is D or E; and

X₁₅ is E or K (SEQ ID NO:8).

In some embodiments, X₁₃ is selected from G, A, L, M, W, P, S, T, R, K, C, N, Q, D, E, V, I or H (SEQ ID NO:49).

In some embodiments, the DNA polymerase having increased 3'-mismatch discrimination comprises a motif in the polymerase domain comprising

A-G-X₁-P-F-N-X₄-N-S-X₇-X₈-Q-X₉-X₁₀-X₁₁-X₁₂-L-X₁₃-X₁₄-X₁₅-L, wherein:

X₁ is H or E;

5 X₄ is L or I;

X₇ is R or P;

X₈ is D or K;

X₉ is L or V;

X₁₀ is E or S;

10 X₁₁ is R or N;

X₁₂ is V or I;

X₁₃ is any amino acid other than F;

X₁₄ is D or E; and

X₁₅ is E or K (SEQ ID NO:9).

15 In some embodiments, the DNA polymerase having increased 3'-mismatch discrimination comprises a motif in the polymerase domain comprising

A-G-H-P-F-N-L-N-S-R-D-Q-L-E-R-V-L-X₁₃-D-E-L, wherein:

X₁₃ is any amino acid other than F (SEQ ID NO:10).

20 In some embodiments, X₁₃ is an amino acid having a polar, uncharged side-chain (e.g., N, Q, H, S, T, or Y), a nonpolar, uncharged side-chain (other than F, e.g., G, A, L, M, W, P, C, V, or I), a polar, positively charged side-chain (e.g., R or K), or polar, negatively charged side-chain (e.g., D or E).

In some embodiments, X₁₃ is A, G, S, T, Y, D, or K (SEQ ID NO:11). In some embodiments, X₁₃ is A, G, S, T, D, or K (SEQ ID NO:48).

25 In some embodiments, the DNA polymerase further comprises a mutation at one or more amino acids corresponding to a position selected from 488 and/or 493 of SEQ ID NO:1. For example, in some embodiments, the amino acid of the DNA polymerase corresponding to position 493 of SEQ ID NO:1 is any amino acid other than E, S, A, or G. In some
30 ID NO:1 is selected from Q, R, V, L, I, M, F, W, P, T, C, N, D, Y, K, or H. In some

embodiments, the amino acid of the DNA polymerase corresponding to position 493 of SEQ ID NO:1 is selected from K or R. In some embodiments, the DNA polymerase of the invention is derived from a *Thermus* species, and the amino acid of the DNA polymerase corresponding to position 493 of SEQ ID NO:1 is any amino acid other than E. For example, where the DNA polymerase is derived from a *Thermus* species, the DNA polymerase can further comprise an amino acid selected from S, A, G, V, L, I, M, F, W, P, T, C, Y, N, Q, D, K, R or H at the position corresponding to position 493 of SEQ ID NO:1. In some embodiments the amino acid at the position corresponding to position 493 of SEQ ID NO:1 is selected from S, A, Q, G, K, or R. In some embodiments the amino acid at the position corresponding to position 493 of SEQ ID NO:1 is selected from A, G, K, or R. In some embodiments the amino acid at the position corresponding to position 493 of SEQ ID NO:1 is K.

In some embodiments, the DNA polymerase can further comprise any amino acid other than S at the position corresponding to position 488 of SEQ ID NO:1. For example, in some embodiments, the amino acid at the position corresponding to position 488 of SEQ ID NO:1 is selected from G, A, V, L, I, M, F, W, P, T, C, Y, N, Q, D, E, K, R or H. In some embodiments, the amino acid at the position corresponding to position 488 of SEQ ID NO:1 is selected from G, A, D, F, K, C, T, or Y.

In some embodiments, the amino acid of the DNA polymerase corresponding to position 580 of SEQ ID NO:1 is any amino acid other than D or E. In some embodiments, the amino acid of the DNA polymerase corresponding to position 580 of SEQ ID NO:1 is any amino acid other than D. In some embodiments, the amino acid of the DNA polymerase corresponding to position 580 of SEQ ID NO:1 is selected from the group consisting of L, G, T, Q, A, S, N, R and K. In some embodiments, the amino acid of the DNA polymerase corresponding to position 580 of SEQ ID NO:1 is G.

Various DNA polymerases are amenable to mutation according to the present invention. Particularly suitable are thermostable polymerases, including wild-type or naturally occurring thermostable polymerases from various species of thermophilic bacteria, as well as synthetic thermostable polymerases derived from such wild-type or naturally occurring enzymes by amino acid substitution, insertion, or deletion, or other modification. Exemplary unmodified forms of polymerase include, *e.g.*, CS5 (SEQ ID NO:29), CS6 (SEQ ID NO:30) or Z05 DNA

polymerase (SEQ ID NO:1), or a functional DNA polymerase having at least 80%, preferably at least 90%, more preferably at least 95% sequence identity thereto. Other unmodified polymerases include, *e.g.*, DNA polymerases from any of the following species of thermophilic bacteria (or a functional DNA polymerase having at least 80%, preferably at least 90%, more preferably at least 95% sequence identity to such a polymerase): *Thermotoga*
 5 *maritima* (SEQ ID NO:38); *Thermus aquaticus* (SEQ ID NO:2); *Thermus thermophilus* (SEQ ID NO:6); *Thermus flavus* (SEQ ID NO:4); *Thermus filiformis* (SEQ ID NO:3); *Thermus sp.* Sps17 (SEQ ID NO:5); *Thermus sp.* Z05 (SEQ ID NO:1); *Thermotoga neopolitana* (SEQ ID NO:39); *Thermosipho africanus* (SEQ ID NO:37); *Thermus caldophilus* (SEQ ID NO:7),
 10 *Deinococcus radiodurans* (SEQ ID NO:36), *Bacillus stearothermophilus* (SEQ ID NO:40) or *Bacillus caldotenax* (SEQ ID NO:41). Suitable polymerases also include those having reverse transcriptase (RT) activity and/or the ability to incorporate unconventional nucleotides, such as ribonucleotides or other 2'-modified nucleotides.

While thermostable DNA polymerases possessing efficient 3'-mismatch discrimination
 15 activity are particularly suited for performing PCR, thermoactive, but not thermostable DNA polymerases possessing efficient 3'-mismatch discrimination activity also are amenable to mutation according to the present invention.

In some embodiments, the DNA polymerase is a *Thermus* DNA polymerase. For example, in some embodiments, the DNA polymerase has at least 80%, preferably at least 90%, more
 20 preferably at least 95% sequence identity to a polymerase selected from the group consisting of:

- (a) a *Thermus sp.* Z05 DNA polymerase (Z05) (SEQ ID NO:1);
- (b) a *Thermus aquaticus* DNA polymerase (Taq) (SEQ ID NO:2);
- (c) a *Thermus filiformis* DNA polymerase (Tfi) (SEQ ID NO:3);
- 25 (d) a *Thermus flavus* DNA polymerase (Tfl) (SEQ ID NO:4);
- (e) a *Thermus sp.* Sps17 DNA polymerase (Sps17) (SEQ ID NO:5);
- (f) a *Thermus thermophilus* DNA polymerase (Tth) (SEQ ID NO:6); and
- (g) a *Thermus caldophilus* DNA polymerase (Tca) (SEQ ID NO:7).

In some embodiments, the DNA polymerase is a *Thermotoga* DNA polymerase. For example,
 30 in some embodiments, the DNA polymerase has at least 80%, preferably at least 90%, more

preferably at least 95% sequence identity to a polymerase selected from the group consisting of:

- (a) a *Thermotoga maritima* DNA polymerase (Tma) (SEQ ID NO:38);
- (b) a *Thermotoga neopolitana* DNA polymerase (Tne) (SEQ ID NO:39);

- 5 In certain embodiments, the DNA polymerase has at least 80%, preferably at least 90%, preferably at least 95% sequence identity to SEQ ID NO:1. In some embodiments, the DNA polymerase is a *Thermus sp.* Z05 DNA polymerase (Z05) (i.e., SEQ ID NO:1), except that the amino acid at position 497 is any amino acid other than F. For example, in some
- 10 Y, N, Q, D, E, K, R or H. In some embodiments, the DNA polymerase is a Z05 DNA polymerase, and the amino acid at position 497 is A, G, S, T, Y, D, or K. In some embodiments, the DNA polymerase is derived from other non-*Thermus*, non-*Thermotoga* species, and the amino acid at position 497 is any amino acid other than F or Y, for example, A, G, S, T, D, or K. In some embodiments, the DNA polymerase is a Z05 DNA polymerase
- 15 further comprising a substitution at position 580, and the amino acid at position 580 is any amino acid other than D or E. In some embodiments, the DNA polymerase is a Z05 DNA polymerase, and the amino acid at position 580 is any amino acid other than D. In some embodiments, the DNA polymerase is a Z05 DNA polymerase, and the amino acid at position 580 is selected from the group consisting of L, G, T, Q, A, S, N, R and K. In some
- 20 20 embodiments, the DNA polymerase is a Z05 DNA polymerase, and the amino acid at position 580 is G.

The mutant or improved polymerase can include other, non-substitutional modifications. One such modification is a thermally reversible covalent modification that inactivates the enzyme, but which is reversed to activate the enzyme upon incubation at an elevated temperature, such

25 as a temperature typically used for polynucleotide extension. Exemplary reagents for such thermally reversible modifications are described in U.S. Patent Nos. 5,773, 258 and 5,677,152 to Birch *et al.*.

In some embodiments, the 3'-mismatch activity is determined using a mutant BRAF V600R target polynucleotide having the nucleic acid sequence of SEQ ID NO:35 (wild type BRAF =

30 SEQ ID NO:34) in the presence of a forward primer that is perfectly matched to the mutant

sequence and has a single 3' A:C mismatch to the wild type sequence in one or more reaction mixtures having a predetermined number of copies of wild-type BRAF V600 target polynucleotide and a predetermined number of copies of mutant BRAF V600R target polynucleotide equal in number or fewer than the number of copies of wild-type target (*e.g.*, 5 10,000 or fewer copies). Two or more reaction mixtures can have titrated numbers of copies of mutant BRAF V600R target polynucleotide (*e.g.*, 1:5 titrations, 1:10 titrations, *e.g.*, 10,000 copies, 1000 copies, 100 copies, 10 copies, 1 copy, 0 copies in several reaction mixtures). The 3'-mismatch discrimination ability of a polymerase of the invention can be compared to the 3'-mismatch discrimination ability of a reference polymerase (*e.g.*, a naturally occurring or 10 unmodified polymerase), over a preselected unit of time, as described herein. Polymerases with increased 3'-mismatch discrimination ability will not amplify the wild-type sequence when contacted with a primer that is perfectly matched to a mutant allele, or will require a greater number of PCR cycles to amplify the wild-type sequence using the mutant allele-specific primer (*i.e.*, exhibit a higher C_p value), in comparison to a naturally occurring or 15 unmodified polymerase.

In various other aspects, the present invention provides a recombinant nucleic acid encoding a mutant or improved DNA polymerase as described herein, a vector comprising the recombinant nucleic acid, and/or a host cell transformed with the vector. In certain 20 embodiments, the vector is an expression vector. Host cells comprising such expression vectors are useful in methods of the invention for producing the mutant or improved polymerase by culturing the host cells under conditions suitable for expression of the recombinant nucleic acid. The polymerases of the invention may be contained in reaction mixtures and/or kits. The embodiments of the recombinant nucleic acids, host cells, vectors, expression vectors, reaction mixtures and kits are as described above and herein.

25 In yet another aspect, a method for conducting polynucleotide extension is provided. The method generally includes contacting a DNA polymerase having increased 3'-mismatch discrimination as described herein with a primer, a polynucleotide template, and nucleoside triphosphates under conditions suitable for extension of the primer, thereby producing an extended primer. The polynucleotide template can be, for example, an RNA or DNA template. 30 The nucleoside triphosphates can include unconventional nucleotides such as, *e.g.*, ribonucleotides and/or labeled nucleotides. Further, the primer and/or template can include

one or more nucleotide analogs. In some variations, the polynucleotide extension method is a method for polynucleotide amplification that includes contacting the mutant or improved DNA polymerase with a primer pair, the polynucleotide template, and the nucleoside triphosphates under conditions suitable for amplification of the polynucleotide. The
5 polynucleotide extension reaction can be, *e.g.*, PCR, isothermal extension, or sequencing (*e.g.*, 454 sequencing reaction).

In some embodiments, the primer extension method is a method for conducting polymerase chain reaction (PCR).

The present invention also provides a kit useful in such a polynucleotide extension method.
10 Generally, the kit includes at least one container providing a mutant or improved DNA polymerase as described herein. In certain embodiments, the kit further includes one or more additional containers providing one or more additional reagents. For example, in specific variations, the one or more additional containers provide nucleoside triphosphates; a buffer suitable for polynucleotide extension; and/or a primer hybridizable, under polynucleotide
15 extension conditions, to a predetermined polynucleotide template.

Further provided are reaction mixtures comprising the polymerases of the invention. The reactions mixtures can also contain a template nucleic acid (DNA and/or RNA), one or more primer or probe polynucleotides, nucleoside triphosphates (including, *e.g.*, deoxyribonucleotides, ribonucleotides, labeled nucleotides, unconventional nucleotides),
20 buffers, salts, labels (*e.g.*, fluorophores).

Further embodiments of the invention are described herein.

DEFINITIONS

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains.
25 Although essentially any methods and materials similar to those described herein can be used in the practice or testing of the present invention, only exemplary methods and materials are described. For purposes of the present invention, the following terms are defined below.

The terms “a,” “an,” and “the” include plural referents, unless the context clearly indicates otherwise.

An “amino acid” refers to any monomer unit that can be incorporated into a peptide, polypeptide, or protein. As used herein, the term “amino acid” includes the following twenty
5 natural or genetically encoded alpha-amino acids: alanine (Ala or A), arginine (Arg or R), asparagine (Asn or N), aspartic acid (Asp or D), cysteine (Cys or C), glutamine (Gln or Q), glutamic acid (Glu or E), glycine (Gly or G), histidine (His or H), isoleucine (Ile or I), leucine (Leu or L), lysine (Lys or K), methionine (Met or M), phenylalanine (Phe or F), proline (Pro or P), serine (Ser or S), threonine (Thr or T), tryptophan (Trp or W), tyrosine (Tyr or Y), and
10 valine (Val or V). In cases where “X” residues are undefined, these should be defined as “any amino acid.” The structures of these twenty natural amino acids are shown in, e.g., Stryer et al., Biochemistry, 5th ed., Freeman and Company (2002). Additional amino acids, such as selenocysteine and pyrrolysine, can also be genetically coded for (Stadtman (1996) “Selenocysteine,” Annu Rev Biochem. 65:83-100 and Ibba et al. (2002) “Genetic code:
15 introducing pyrrolysine,” Curr Biol. 12(13):R464-R466). The term “amino acid” also includes unnatural amino acids, modified amino acids (e.g., having modified side chains and/or backbones), and amino acid analogs. See, e.g., Zhang et al. (2004) “Selective incorporation of 5-hydroxytryptophan into proteins in mammalian cells,” Proc. Natl. Acad. Sci. U.S.A. 101(24):8882-8887, Anderson et al. (2004) “An expanded genetic code with a functional
20 quadruplet codon” Proc. Natl. Acad. Sci. U.S.A. 101(20):7566-7571, Ikeda et al. (2003) “Synthesis of a novel histidine analogue and its efficient incorporation into a protein in vivo,” Protein Eng. Des. Sel. 16(9):699-706, Chin et al. (2003) “An Expanded Eukaryotic Genetic Code,” Science 301(5635):964-967, James et al. (2001) “Kinetic characterization of ribonuclease S mutants containing photoisomerizable phenylazophenylalanine residues,”
25 Protein Eng. Des. Sel. 14(12):983-991, Kohrer et al. (2001) “Import of amber and ochre suppressor tRNAs into mammalian cells: A general approach to site-specific insertion of amino acid analogues into proteins,” Proc. Natl. Acad. Sci. U.S.A. 98(25):14310-14315, Bacher et al. (2001) “Selection and Characterization of Escherichia coli Variants Capable of Growth on an Otherwise Toxic Tryptophan Analogue,” J. Bacteriol. 183(18):5414-5425,
30 Hamano-Takaku et al. (2000) “A Mutant Escherichia coli Tyrosyl-tRNA Synthetase Utilizes the Unnatural Amino Acid Azatyrosine More Efficiently than Tyrosine,” J. Biol. Chem.

275(51):40324-40328, and Budisa et al. (2001) "Proteins with {beta}-(thienopyrrolyl)alanines as alternative chromophores and pharmaceutically active amino acids," Protein Sci. 10(7):1281-1292.

To further illustrate, an amino acid is typically an organic acid that includes a substituted or unsubstituted amino group, a substituted or unsubstituted carboxy group, and one or more side chains or groups, or analogs of any of these groups. Exemplary side chains include, e.g., thiol, seleno, sulfonyl, alkyl, aryl, acyl, keto, azido, hydroxyl, hydrazine, cyano, halo, hydrazide, alkenyl, alkynyl, ether, borate, boronate, phospho, phosphono, phosphine, heterocyclic, enone, imine, aldehyde, ester, thioacid, hydroxylamine, or any combination of these groups. Other representative amino acids include, but are not limited to, amino acids comprising photoactivatable cross-linkers, metal binding amino acids, spin-labeled amino acids, fluorescent amino acids, metal-containing amino acids, amino acids with novel functional groups, amino acids that covalently or noncovalently interact with other molecules, photocaged and/or photoisomerizable amino acids, radioactive amino acids, amino acids comprising biotin or a biotin analog, glycosylated amino acids, other carbohydrate modified amino acids, amino acids comprising polyethylene glycol or polyether, heavy atom substituted amino acids, chemically cleavable and/or photocleavable amino acids, carbon-linked sugar-containing amino acids, redox-active amino acids, amino thioacid containing amino acids, and amino acids comprising one or more toxic moieties.

The term "aptamer" refers to a single-stranded DNA that recognizes and binds to DNA polymerase, and efficiently inhibits the polymerase activity as described in U.S. Pat. No. 5,693,502.

The term "mutant," in the context of DNA polymerases of the present invention, means a polypeptide, typically recombinant, that comprises one or more amino acid substitutions relative to a corresponding, naturally-occurring or unmodified DNA polymerase.

The term "unmodified form," in the context of a mutant polymerase, is a term used herein for purposes of defining a mutant DNA polymerase of the present invention: the term "unmodified form" refers to a functional DNA polymerase that has the amino acid sequence of the mutant polymerase except at one or more amino acid position(s) specified as characterizing the mutant polymerase. Thus, reference to a mutant DNA polymerase in terms

of (a) its unmodified form and (b) one or more specified amino acid substitutions means that, with the exception of the specified amino acid substitution(s), the mutant polymerase otherwise has an amino acid sequence identical to the unmodified form in the specified motif. The “unmodified polymerase” (and therefore also the modified form having increased 3'-mismatch discrimination) may contain additional mutations to provide desired functionality, e.g., improved incorporation of dideoxyribonucleotides, ribonucleotides, ribonucleotide analogs, dye-labeled nucleotides, modulating 5'-nuclease activity, modulating 3'-nuclease (or proofreading) activity, or the like. Accordingly, in carrying out the present invention as described herein, the unmodified form of a DNA polymerase is predetermined. The unmodified form of a DNA polymerase can be, for example, a wild-type and/or a naturally occurring DNA polymerase, or a DNA polymerase that has already been intentionally modified. An unmodified form of the polymerase is preferably a thermostable DNA polymerases, such as DNA polymerases from various thermophilic bacteria, as well as functional variants thereof having substantial sequence identity to a wild-type or naturally occurring thermostable polymerase. Such variants can include, for example, chimeric DNA polymerases such as, for example, the chimeric DNA polymerases described in U.S. Patent No. 6,228,628 and U.S. Application Publication No. 2004/0005599. In certain embodiments, the unmodified form of a polymerase has reverse transcriptase (RT) activity.

The term “thermostable polymerase,” refers to an enzyme that is stable to heat, is heat resistant, and retains sufficient activity to effect subsequent polynucleotide extension reactions and does not become irreversibly denatured (inactivated) when subjected to the elevated temperatures for the time necessary to effect denaturation of double-stranded nucleic acids. The heating conditions necessary for nucleic acid denaturation are well known in the art and are exemplified in, e.g., U.S. Patent Nos. 4,683,202, 4,683,195, and 4,965,188. As used herein, a thermostable polymerase is suitable for use in a temperature cycling reaction such as the polymerase chain reaction (“PCR”). Irreversible denaturation for purposes herein refers to permanent and complete loss of enzymatic activity. For a thermostable polymerase, enzymatic activity refers to the catalysis of the combination of the nucleotides in the proper manner to form polynucleotide extension products that are complementary to a template nucleic acid strand. Thermostable DNA polymerases from thermophilic bacteria include, e.g., DNA polymerases from *Thermotoga maritima*, *Thermus aquaticus*, *Thermus thermophilus*, *Thermus*

flavus, *Thermus filiformis*, *Thermus* species Sps17, *Thermus* species Z05, *Thermus caldophilus*, *Bacillus caldotenax*, *Thermotoga neopolitana*, and *Thermosipho africanus*.

The term "thermoactive" refers to an enzyme that maintains catalytic properties at temperatures commonly used for reverse transcription or anneal/extension steps in RT-PCR and/or PCR reactions (i.e., 45-80 °C). Thermostable enzymes are those which are not irreversibly inactivated or denatured when subjected to elevated temperatures necessary for nucleic acid denaturation. Thermoactive enzymes may or may not be thermostable.

Thermoactive DNA polymerases can be DNA or RNA dependent from thermophilic species or from mesophilic species including, but not limited to, *Escherichia coli*, *Moloney murine leukemia viruses*, and *Avian myoblastosis virus*.

As used herein, a "chimeric" protein refers to a protein whose amino acid sequence represents a fusion product of subsequences of the amino acid sequences from at least two distinct proteins. A chimeric protein typically is not produced by direct manipulation of amino acid sequences, but, rather, is expressed from a "chimeric" gene that encodes the chimeric amino acid sequence. In certain embodiments, for example, an unmodified form of a mutant DNA polymerase of the present invention is a chimeric protein that consists of an amino-terminal (N-terminal) region derived from a *Thermus* species DNA polymerase and a carboxy-terminal (C-terminal) region derived from Tma DNA polymerase. The N-terminal region refers to a region extending from the N-terminus (amino acid position 1) to an internal amino acid. Similarly, the C-terminal region refers to a region extending from an internal amino acid to the C-terminus.

In the context of DNA polymerases, "correspondence" to another sequence (e.g., regions, fragments, nucleotide or amino acid positions, or the like) is based on the convention of numbering according to nucleotide or amino acid position number and then aligning the sequences in a manner that maximizes the percentage of sequence identity. Because not all positions within a given "corresponding region" need be identical, non-matching positions within a corresponding region may be regarded as "corresponding positions." Accordingly, as used herein, referral to an "amino acid position corresponding to amino acid position [X]" of a specified DNA polymerase refers to equivalent positions, based on alignment, in other DNA polymerases and structural homologues and families. In some embodiments of the present

invention, "correspondence" of amino acid positions are determined with respect to a region of the polymerase comprising one or more motifs of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 36, 37, 38, 39, 40, or 41. When a polymerase polypeptide sequence differs from SEQ ID NOS:1, 2, 3, 4, 5, 6, 7, 36, 37, 38, 39, 40, or 41 (*e.g.*, by changes in amino acids or addition or deletion of amino acids), it may be that a particular mutation associated with improved activity as discussed herein will not be in the same position number as it is in SEQ ID NOS:1, 2, 3, 4, 5, 6, 7, 36, 37, 38, 39, 40, or 41. This is illustrated, for example, in Table 1.

"Recombinant," as used herein, refers to an amino acid sequence or a nucleotide sequence that has been intentionally modified by recombinant methods. By the term "recombinant nucleic acid" herein is meant a nucleic acid, originally formed *in vitro*, in general, by the manipulation of a nucleic acid by endonucleases, in a form not normally found in nature. Thus an isolated, mutant DNA polymerase nucleic acid, in a linear form, or an expression vector formed *in vitro* by ligating DNA molecules that are not normally joined, are both considered recombinant for the purposes of this invention. It is understood that once a recombinant nucleic acid is made and reintroduced into a host cell, it will replicate non-recombinantly, *i.e.*, using the *in vivo* cellular machinery of the host cell rather than *in vitro* manipulations; however, such nucleic acids, once produced recombinantly, although subsequently replicated non-recombinantly, are still considered recombinant for the purposes of the invention. A "recombinant protein" is a protein made using recombinant techniques, *i.e.*, through the expression of a recombinant nucleic acid as depicted above.

A nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation.

The term "host cell" refers to both single-cellular prokaryote and eukaryote organisms (*e.g.*, bacteria, yeast, and actinomycetes) and single cells from higher order plants or animals when being grown in cell culture.

The term "vector" refers to a piece of DNA, typically double-stranded, which may have inserted into it a piece of foreign DNA. The vector or may be, for example, of plasmid origin. Vectors contain "replicon" polynucleotide sequences that facilitate the autonomous replication

of the vector in a host cell. Foreign DNA is defined as heterologous DNA, which is DNA not naturally found in the host cell, which, for example, replicates the vector molecule, encodes a selectable or screenable marker, or encodes a transgene. The vector is used to transport the foreign or heterologous DNA into a suitable host cell. Once in the host cell, the vector can replicate independently of or coincidental with the host chromosomal DNA, and several copies of the vector and its inserted DNA can be generated. In addition, the vector can also contain the necessary elements that permit transcription of the inserted DNA into an mRNA molecule or otherwise cause replication of the inserted DNA into multiple copies of RNA. Some expression vectors additionally contain sequence elements adjacent to the inserted DNA that increase the half-life of the expressed mRNA and/or allow translation of the mRNA into a protein molecule. Many molecules of mRNA and polypeptide encoded by the inserted DNA can thus be rapidly synthesized.

The term "nucleotide," in addition to referring to the naturally occurring ribonucleotide or deoxyribonucleotide monomers, shall herein be understood to refer to related structural variants thereof, including derivatives and analogs, that are functionally equivalent with respect to the particular context in which the nucleotide is being used (*e.g.*, hybridization to a complementary base), unless the context clearly indicates otherwise.

The term "nucleic acid" or "polynucleotide" refers to a polymer that can be corresponded to a ribose nucleic acid (RNA) or deoxyribose nucleic acid (DNA) polymer, or an analog thereof. This includes polymers of nucleotides such as RNA and DNA, as well as synthetic forms, modified (*e.g.*, chemically or biochemically modified) forms thereof, and mixed polymers (*e.g.*, including both RNA and DNA subunits). Exemplary modifications include methylation, substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as uncharged linkages (*e.g.*, methyl phosphonates, phosphotriesters, phosphoamidates, carbamates, and the like), pendent moieties (*e.g.*, polypeptides), intercalators (*e.g.*, acridine, psoralen, and the like), chelators, alkylators, and modified linkages (*e.g.*, alpha anomeric nucleic acids and the like). Also included are synthetic molecules that mimic polynucleotides in their ability to bind to a designated sequence via hydrogen bonding and other chemical interactions. Typically, the nucleotide monomers are linked via phosphodiester bonds, although synthetic forms of nucleic acids can comprise other linkages (*e.g.*, peptide nucleic acids as described in Nielsen *et al.* (*Science* 254:1497-1500,

1991). A nucleic acid can be or can include, e.g., a chromosome or chromosomal segment, a vector (e.g., an expression vector), an expression cassette, a naked DNA or RNA polymer, the product of a polymerase chain reaction (PCR), an oligonucleotide, a probe, and a primer. A nucleic acid can be, e.g., single-stranded, double-stranded, or triple-stranded and is not limited to any particular length. Unless otherwise indicated, a particular nucleic acid sequence comprises or encodes complementary sequences, in addition to any sequence explicitly indicated.

The term "oligonucleotide" refers to a nucleic acid that includes at least two nucleic acid monomer units (e.g., nucleotides). An oligonucleotide typically includes from about six to about 175 nucleic acid monomer units, more typically from about eight to about 100 nucleic acid monomer units, and still more typically from about 10 to about 50 nucleic acid monomer units (e.g., about 15, about 20, about 25, about 30, about 35, or more nucleic acid monomer units). The exact size of an oligonucleotide will depend on many factors, including the ultimate function or use of the oligonucleotide. Oligonucleotides are optionally prepared by any suitable method, including, but not limited to, isolation of an existing or natural sequence, DNA replication or amplification, reverse transcription, cloning and restriction digestion of appropriate sequences, or direct chemical synthesis by a method such as the phosphotriester method of Narang *et al.* (*Meth. Enzymol.* 68:90-99, 1979); the phosphodiester method of Brown *et al.* (*Meth. Enzymol.* 68:109-151, 1979); the diethylphosphoramidite method of Beaucage *et al.* (*Tetrahedron Lett.* 22:1859-1862, 1981); the triester method of Matteucci *et al.* (*J. Am. Chem. Soc.* 103:3185-3191, 1981); automated synthesis methods; or the solid support method of U.S. Pat. No. 4,458,066, or other methods known to those skilled in the art.

The term "primer" as used herein refers to a polynucleotide capable of acting as a point of initiation of template-directed nucleic acid synthesis when placed under conditions in which polynucleotide extension is initiated (e.g., under conditions comprising the presence of requisite nucleoside triphosphates (as dictated by the template that is copied) and a polymerase in an appropriate buffer and at a suitable temperature or cycle(s) of temperatures (e.g., as in a polymerase chain reaction)). To further illustrate, primers can also be used in a variety of other oligonucleotide-mediated synthesis processes, including as initiators of *de novo* RNA synthesis and *in vitro* transcription-related processes (e.g., nucleic acid sequence-based amplification (NASBA), transcription mediated amplification (TMA), etc.). A primer is typically a single-

stranded oligonucleotide (*e.g.*, oligodeoxyribonucleotide). The appropriate length of a primer depends on the intended use of the primer but typically ranges from 6 to 40 nucleotides, more typically from 15 to 35 nucleotides. Short primer molecules generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. A primer need
5 not reflect the exact sequence of the template but must be sufficiently complementary to hybridize with a template for primer elongation to occur. In certain embodiments, the term "primer pair" means a set of primers including a 5' sense primer (sometimes called "forward") that hybridizes with the complement of the 5' end of the nucleic acid sequence to be amplified and a 3' antisense primer (sometimes called "reverse") that hybridizes with the 3' end of the
10 sequence to be amplified (*e.g.*, if the target sequence is expressed as RNA or is an RNA). A primer can be labeled, if desired, by incorporating a label detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include ^{32}P , fluorescent dyes, electron-dense reagents, enzymes (as commonly used in ELISA assays), biotin, or haptens and proteins for which antisera or monoclonal antibodies are
15 available.

The term "5'-nuclease probe" refers to an oligonucleotide that comprises at least one light emitting labeling moiety and that is used in a 5'-nuclease reaction to effect target nucleic acid detection. In some embodiments, for example, a 5'-nuclease probe includes only a single light emitting moiety (*e.g.*, a fluorescent dye, etc.). In certain embodiments, 5'-nuclease probes
20 include regions of self-complementarity such that the probes are capable of forming hairpin structures under selected conditions. To further illustrate, in some embodiments a 5'-nuclease probe comprises at least two labeling moieties and emits radiation of increased intensity after one of the two labels is cleaved or otherwise separated from the oligonucleotide. In certain embodiments, a 5'-nuclease probe is labeled with two different fluorescent dyes, *e.g.*, a 5'
25 terminus reporter dye and the 3' terminus quencher dye or moiety. In some embodiments, 5'-nuclease probes are labeled at one or more positions other than, or in addition to, terminal positions. When the probe is intact, energy transfer typically occurs between the two fluorophores such that fluorescent emission from the reporter dye is quenched at least in part. During an extension step of a polymerase chain reaction, for example, a 5'-nuclease probe
30 bound to a template nucleic acid is cleaved by the 5' to 3' nuclease activity of, *e.g.*, a *Taq* polymerase or another polymerase having this activity such that the fluorescent emission of

the reporter dye is no longer quenched. Exemplary 5'-nuclease probes are also described in, e.g., U.S. Pat. No. 5,210,015; U.S. Pat. No. 5,994,056; and U.S. Pat. No. 6,171,785. In other embodiments, a 5' nuclease probe may be labeled with two or more different reporter dyes and a 3' terminus quencher dye or moiety.

5 The term "FRET" or "fluorescent resonance energy transfer" or "Foerster resonance energy transfer" refers to a transfer of energy between at least two chromophores, a donor chromophore and an acceptor chromophore (referred to as a quencher). The donor typically transfers the energy to the acceptor when the donor is excited by light radiation with a suitable wavelength. The acceptor typically re-emits the transferred energy in the form of light
10 radiation with a different wavelength. When the acceptor is a "dark" quencher, it dissipates the transferred energy in a form other than light. Whether a particular fluorophore acts as a donor or an acceptor depends on the properties of the other member of the FRET pair. Commonly used donor-acceptor pairs include the FAM-TAMRA pair. Commonly used quenchers are DABCYL and TAMRA. Commonly used dark quenchers include BlackHole Quenchers™
15 (BHQ), (Biosearch Technologies, Inc., Novato, Cal.), Iowa Black™ (Integrated DNA Tech., Inc., Coralville, Iowa), and BlackBerry™ Quencher 650 (BBQ-650) (Berry & Assoc., Dexter, Mich.).

The term "conventional" or "natural" when referring to nucleic acid bases, nucleoside triphosphates, or nucleotides refers to those which occur naturally in the polynucleotide being
20 described (*i.e.*, for DNA these are dATP, dGTP, dCTP and dTTP). Additionally, dITP, and 7-deaza-dGTP are frequently utilized in place of dGTP and 7-deaza-dATP can be utilized in place of dATP in *in vitro* DNA synthesis reactions, such as sequencing. Collectively, these may be referred to as dNTPs.

The term "unconventional" or "modified" when referring to a nucleic acid base, nucleoside, or
25 nucleotide includes modification, derivations, or analogues of conventional bases, nucleosides, or nucleotides that naturally occur in a particular polynucleotide. Certain unconventional nucleotides are modified at the 2' position of the ribose sugar in comparison to conventional dNTPs. Thus, although for RNA the naturally occurring nucleotides are ribonucleotides (*i.e.*, ATP, GTP, CTP, UTP, collectively rNTPs), because these nucleotides have a hydroxyl group
30 at the 2' position of the sugar, which, by comparison is absent in dNTPs, as used herein,

ribonucleotides are unconventional nucleotides as substrates for DNA polymerases. As used herein, unconventional nucleotides include, but are not limited to, compounds used as terminators for nucleic acid sequencing. Exemplary terminator compounds include but are not limited to those compounds that have a 2',3' dideoxy structure and are referred to as

5 dideoxynucleoside triphosphates. The dideoxynucleoside triphosphates ddATP, ddTTP, ddCTP and ddGTP are referred to collectively as ddNTPs. Additional examples of terminator compounds include 2'-PO₄ analogs of ribonucleotides (*see, e.g.*, U.S. Application Publication Nos. 2005/0037991 and 2005/0037398). Other unconventional nucleotides include phosphorothioate dNTPs ([α]-S)dNTPs), 5'-[α]-borano-dNTPs, [α]-methyl-phosphonate

10 dNTPs, and ribonucleoside triphosphates (rNTPs). Unconventional bases may be labeled with radioactive isotopes such as ³²P, ³³P, or ³⁵S; fluorescent labels; chemiluminescent labels; bioluminescent labels; hapten labels such as biotin; or enzyme labels such as streptavidin or avidin. Fluorescent labels may include dyes that are negatively charged, such as dyes of the fluorescein family, or dyes that are neutral in charge, such as dyes of the rhodamine family, or

15 dyes that are positively charged, such as dyes of the cyanine family. Dyes of the fluorescein family include, *e.g.*, FAM, HEX, TET, JOE, NAN and ZOE. Dyes of the rhodamine family include Texas Red, ROX, R110, R6G, and TAMRA. Various dyes or nucleotides labeled with FAM, HEX, TET, JOE, NAN, ZOE, ROX, R110, R6G, Texas Red and TAMRA are marketed by Perkin-Elmer (Boston, MA), Applied Biosystems (Foster City, CA), or

20 Invitrogen/Molecular Probes (Eugene, OR). Dyes of the cyanine family include Cy2, Cy3, Cy5, and Cy7 and are marketed by GE Healthcare UK Limited (Amersham Place, Little Chalfont, Buckinghamshire, England).

As used herein, "percentage of sequence identity" is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the sequence in the

25 comparison window can comprise additions or deletions (*i.e.*, gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total

30 number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

The terms "identical" or "identity," in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same. Sequences are "substantially identical" to each other if they have a specified percentage of nucleotides or amino acid residues that are the same (*e.g.*, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% identity over a specified region), when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. These definitions also refer to the complement of a test sequence. Optionally, the identity exists over a region that is at least about 50 nucleotides in length, or more typically over a region that is 100 to 500 or 1000 or more nucleotides in length.

The terms "similarity" or "percent similarity," in the context of two or more polypeptide sequences, refer to two or more sequences or subsequences that have a specified percentage of amino acid residues that are either the same or similar as defined by a conservative amino acid substitutions (*e.g.*, 60% similarity, optionally 65%, 70%, 75%, 80%, 85%, 90%, or 95% similar over a specified region), when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. Sequences are "substantially similar" to each other if they are at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, or at least 55% similar to each other. Optionally, this similarity exists over a region that is at least about 50 amino acids in length, or more typically over a region that is at least about 100 to 500 or 1000 or more amino acids in length.

For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters are commonly used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities or similarities for the test sequences relative to the reference sequence, based on the program parameters.

A "comparison window," as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of from 20 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well known in the art. Optimal alignment of sequences for comparison can be conducted, for example, by the local homology algorithm of Smith and Waterman (*Adv. Appl. Math.* 2:482, 1970), by the homology alignment algorithm of Needleman and Wunsch (*J. Mol. Biol.* 48:443, 1970), by the search for similarity method of Pearson and Lipman (*Proc. Natl. Acad. Sci. USA* 85:2444, 1988), by computerized implementations of these algorithms (e.g., GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by manual alignment and visual inspection (see, e.g., Ausubel *et al.*, *Current Protocols in Molecular Biology* (1995 supplement)).

Algorithms suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul *et al.* (*Nuc. Acids Res.* 25:3389-402, 1997), and Altschul *et al.* (*J. Mol. Biol.* 215:403-10, 1990), respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm

parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3, and expectation (E) of 10, and the
5 BLOSUM62 scoring matrix (*see* Henikoff and Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915, 1989) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (*see, e.g.*, Karlin and Altschul, *Proc. Natl. Acad. Sci. USA* 90:5873-87, 1993). One
10 measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, typically less than
15 about 0.01, and more typically less than about 0.001.

The term “mismatch discrimination” refers to the ability of a biocatalyst (e.g., an enzyme, such as a polymerase, ligase, or the like) to distinguish a fully complementary sequence from a mismatch-containing sequence when extending a nucleic acid (e.g., a primer or other oligonucleotide) in a template-dependent manner by attaching (e.g., covalently) one or more
20 nucleotides to the nucleic acid. The term “3’-mismatch discrimination” refers to the ability of a biocatalyst to distinguish a fully complementary sequence from a mismatch-containing (nearly complementary) sequence where the nucleic acid to be extended (e.g., a primer or other oligonucleotide) has a mismatch at the nucleic acid’s 3’ terminus compared to the template to which the nucleic acid hybridizes. In some embodiments, the nucleic acid to be
25 extended comprises a mismatch at the 3’ end relative to the fully complementary sequence. In some embodiments, the nucleic acid to be extended comprises a mismatch at the penultimate (N-1) 3’ position and/or at the N-2 position relative to the fully complementary sequence.

The term “Cp value” or “crossing point” value refers to a value that allows quantification of input target nucleic acids. The Cp value can be determined according to the second-derivative
30 maximum method (Van Luu-The, et al., “Improved real-time RT-PCR method for high-

throughput measurements using second derivative calculation and double correction,” BioTechniques, Vol. 38, No. 2, February 2005, pp. 287–293). In the second derivative method, a C_p corresponds to the first peak of a second derivative curve. This peak corresponds to the beginning of a log-linear phase. The second derivative method calculates a second
 5 derivative value of the real-time fluorescence intensity curve, and only one value is obtained. The original C_p method is based on a locally defined, differentiable approximation of the intensity values, *e.g.*, by a polynomial function. Then the third derivative is computed. The C_p value is the smallest root of the third derivative. The C_p can also be determined using the fit point method, in which the C_p is determined by the intersection of a parallel to the threshold
 10 line in the log-linear region (Van Luu-The, et al., BioTechniques, Vol. 38, No. 2, February 2005, pp. 287–293). These computations are easily carried out by any person skilled in the art.

The term “PCR efficiency” refers to an indication of cycle to cycle amplification efficiency for the perfectly matched primer template. PCR efficiency is calculated for each condition using the equation: % PCR efficiency = $(10^{(\text{slope})} - 1) \times 100$, wherein the slope was calculated by
 15 linear regression with the log copy number plotted on the y-axis and C_p plotted on the x-axis.

The term “multiplex” refers to amplification with more than one set of primers, or the amplification of more than one polymorphism site in a single reaction.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts an amino acid sequence alignment of a region from the polymerase domain
 20 of exemplary DNA polymerases from various species of bacteria: *Thermus* species Z05 (Z05) (SEQ ID NO:12), *Thermus aquaticus* (Taq) (SEQ ID NO:13), *Thermus filiformis* (Tfi) (SEQ ID NO:14), *Thermus flavus* (Tfl) (SEQ ID NO:15), *Thermus* species Sps17 (Sps17) (SEQ ID NO:16), *Thermus thermophilus* (Tth) (SEQ ID NO:17), *Thermus caldophilus* (Tca) (SEQ ID NO:18), *Thermotoga maritima* (Tma) (SEQ ID NO:19), *Thermotoga neopolitana* (Tne) (SEQ
 25 ID NO:20), *Thermosipho africanus* (Taf) (SEQ ID NO:21), *Deinococcus radiodurans* (Dra) (SEQ ID NO:23), *Bacillus stearothermophilus* (Bst) (SEQ ID NO:24), and *Bacillus caldotenax* (Bca) (SEQ ID NO:25). In addition, the polypeptide regions shown comprise the amino acid motif A-G-X₁-X₂-F-X₃-X₄-X₅-X₆-X₇-X₈-Q-X₉-X₁₀-X₁₁-X₁₂-L-X₁₃-X₁₄-X₁₅-L (SEQ ID NO:26), the variable positions of which are further defined herein. This motif is

highlighted in bold type for each polymerase sequence. Amino acid positions amenable to mutation in accordance with the present invention are indicated with an asterisk (*).

Figure 2 provides sequence identities among the following DNA Polymerase I enzymes:

Thermus sp. Z05 DNA polymerase (Z05); *Thermus aquaticus* DNA polymerase (Taq);

- 5 *Thermus filiformis* DNA polymerase (Tfi); *Thermus flavus* DNA polymerase (Tfl); *Thermus sp.* Sps17 DNA polymerase (Sps17); *Thermus thermophilus* DNA polymerase (Tth); *Thermus caldophilus* DNA polymerase (Tca); *Deinococcus radiodurans* DNA polymerase (Dra); *Thermotoga maritima* DNA polymerase (Tma); *Thermotoga neopolitana* DNA polymerase (Tne); *Thermosipho africanus* DNA polymerase (Taf); *Bacillus stearothermophilus* DNA
10 polymerase (Bst); and *Bacillus caldotenax* DNA polymerase (Bca). (A) sequence identities over the entire polymerase I enzyme (corresponding to amino acids 1-834 of Z05); and (B) sequence identities over the polymerase sub domain corresponding to amino acids 420-834 of Z05.

DETAILED DESCRIPTION

- 15 The present invention provides improved DNA polymerases in which one or more amino acids in the polymerase domain have been identified as improving one or more polymerase activity or characteristics. The DNA polymerases of the invention are active enzymes having increased 3'-mismatch discrimination activity (i.e., the inventive polymerases described herein are less likely to extend primers that are mismatched to template at or near the 3' end of the
20 primer) relative to the unmodified form of the polymerase otherwise identical except for the amino acid difference noted herein. The DNA polymerases are useful in a variety of applications involving polynucleotide extension or amplification of polynucleotide templates, including, for example, applications in recombinant DNA studies and medical diagnosis of disease.

25 *Polymerases of the Invention*

DNA polymerases of the invention can be characterized by having the following motif:

Ala-Gly-X₁-X₂-Phe-X₃-X₄-X₅-Ser-X₇-X₈-Gln-X₉-X₁₀-X₁₁-X₁₂-Leu-X₁₃-X₁₄-
X₁₅-Leu (also referred to herein in the one-letter code as A-G-X₁-X₂-F-X₃-X₄-
X₅-S-X₇-X₈-Q-X₉-X₁₀-X₁₁-X₁₂-L-X₁₃-X₁₄-X₁₅-L); wherein:

X₁ is His (H), Glu (E) or Gln (Q);

X₂ is Pro (P), Thr (T) or Glu (E);

X₃ is Asn (N) or His (H);

X₄ is Leu (L) or Ile (I);

5 X₅ is Asn (N) or Arg (R);

X₇ is Arg (R), Pro (P), or Ser (S);

X₈ is Asp (D), Lys (K) or Thr (T);

X₉ is Leu (L) or Val (V);

X₁₀ is Glu (E), Ser (S), Ala (A) or Gly (G);

10 X₁₁ is Arg (R), Asn (N), Tyr (Y), Thr (T) or Val (V);

X₁₂ is Val (V) or Ile (I);

X₁₃ is any amino acid other than Phe (F) or Tyr (Y);

X₁₄ is Asp (D) or Glu (E); and

X₁₅ is Glu (E) or Lys (K) (SEQ ID NO:8).

15 In some embodiments, X₁₃ is selected from G, A, V, L, I, M, W, P, S, T, C, N, Q, D, E, K, R or H (SEQ ID NO:49).

In some embodiments, DNA polymerases of the invention can be characterized by having the following motif (corresponding to *Thermus* and *Thermotoga*):

Ala-Gly-X₁-Pro-Phe-Asn-X₄-Asn-Ser-X₇-X₈-Gln-X₉-X₁₀-X₁₁-X₁₂-Leu-X₁₃-X₁₄-

20 X₁₅-Leu (also referred to herein in the one-letter code as A-G-X₁-P-F-N-X₄-N-S-X₇-X₈-Q-X₉-X₁₀-X₁₁-X₁₂-L-X₁₃-X₁₄-X₁₅-L); wherein:

X₁ is His (H) or Glu (E);

X₄ is Leu (L) or Ile (I);

X₇ is Arg (R) or Pro (P);

25 X₈ is Asp (D) or Lys (K);

X₉ is Leu (L) or Val (V);

X₁₀ is Glu (E) or Ser (S);

X₁₁ is Arg (R) or Asn (N);

X₁₂ is Val (V) or Ile (I);

30 X₁₃ is any amino acid other than Phe (F);

X₁₄ is Asp (D) or Glu (E); and

X₁₅ is Glu (E) or Lys (K) (SEQ ID NO:9).

In some embodiments, DNA polymerases of the invention can be characterized by having the following motif:

5 Ala-Gly-His-Pro-Phe-Asn-Leu-Asn-Ser-Arg-Asp-Gln-Leu-Glu-Arg-Val-Leu-
X₁₃-Asp-Glu-Leu (also referred to herein in the one-letter code as A-G-H-P-F-
N-L-N-S-R-D-Q-L-E-R-V-L-X₁₃-D-E-L); wherein:
X₁₃ is any amino acid other than Phe (F) (SEQ ID NO:10).

10 In some embodiments, DNA polymerases of the invention can be characterized by having the following motif:

 Ala-Gly-His-Pro-Phe-Asn-Leu-Asn-Ser-Arg-Asp-Gln-Leu-Glu-Arg-Val-Leu-
X₁₃-Asp-Glu-Leu (also referred to herein in the one-letter code as A-G-H-P-F-
N-L-N-S-R-D-Q-L-E-R-V-L-X₁₃-D-E-L); wherein:
X₁₃ is Ala (A), Gly (G), Ser (S), Thr (T), Tyr (Y), Asp (D) or Lys (K) (SEQ ID
15 NO:11).

In some embodiments, DNA polymerases of the invention can be characterized by having the following motif:

 Ala-Gly-His-Pro-Phe-Asn-Leu-Asn-Ser-Arg-Asp-Gln-Leu-Glu-Arg-Val-Leu-
X₁₃-Asp-Glu-Leu (also referred to herein in the one-letter code as A-G-H-P-F-
20 N-L-N-S-R-D-Q-L-E-R-V-L-X₁₃-D-E-L); wherein:
X₁₃ is Ala (A), Gly (G), Ser (S), Thr (T), Asp (D) or Lys (K) (SEQ ID NO:48).

Further, in some embodiments, the DNA polymerases of the invention can comprise additional amino acid substitutions, for example, at positions X₆ and X₁₀ of the native motif (SEQ ID
25 NO:26). The additional substitutions at positions X₆ and X₁₀ can also result in increased 3' mismatch discrimination. Thus, in some embodiments, DNA polymerases of the invention can be characterized by having the following motif:

Ala-Gly-X₁-X₂-Phe-X₃-X₄-X₅-X₆-X₇-X₈-Gln-X₉-X₁₀-X₁₁-
X₁₂-Leu-X₁₃-X₁₄-X₁₅-Leu (also referred to herein in the one-

letter code as A-G-X₁-X₂-F-X₃-X₄-X₅-X₆-X₇-X₈-Q-X₉-X₁₀-

X₁₁-X₁₂-L-X₁₃-X₁₄-X₁₅-L); wherein

X₁ is His (H), Glu (E) or Gln (Q);

X₂ is Pro (P), Thr (T) or Glu (E);

5 X₃ is Asn (N) or His (H);

X₄ is Leu (L) or Ile (I);

X₅ is Asn (N) or Arg (R);

X₆ is any amino acid;

X₇ is Arg (R), Pro (P), or Ser (S);

10 X₈ is Asp (D), Lys (K) or Thr (T);

X₉ is Leu (L) or Val (V);

X₁₀ is any amino acid;

X₁₁ is Arg (R), Asn (N), Tyr (Y), Thr (T) or Val (V);

X₁₂ is Val (V) or Ile (I);

15 X₁₃ is any amino acid other than Phe (F) or Tyr (Y);

X₁₄ is Asp (D) or Glu (E); and

X₁₅ is Glu (E) or Lys (K) (SEQ ID NO:42)

In some embodiments, DNA polymerases of the invention can be characterized by having the following motif:

20 Ala-Gly-X₁-Pro-Phe-Asn-X₄-Asn-X₆-X₇-X₈-Gln-X₉-X₁₀-

X₁₁-X₁₂-Leu-X₁₃-X₁₄-X₁₅-Leu (also referred to herein in the

one-letter code as A-G-X₁-P-F-N-X₄-N-X₆-X₇-X₈-Q-X₉-

X₁₀-X₁₁-X₁₂-L-X₁₃-X₁₄-X₁₅-L); wherein

X₁ is His (H) or Glu (E);

25 X₄ is Leu (L) or Ile (I);

X₆ is any amino acid;

X₇ is Arg (R) or Pro (P);

X₈ is Asp (D) or Lys (K);

X₉ is Leu (L) or Val (V);

30 X₁₀ is any amino acid;

X₁₁ is Arg (R) or Asn (N);

X_{12} is Val (V) or Ile (I);
 X_{13} is any amino acid other than Phe (F);
 X_{14} is Asp (D) or Glu (E); and
 X_{15} is Glu (E) or Lys (K) (SEQ ID NO:43)

- 5 In some embodiments, DNA polymerases of the invention can be characterized by having the following motif (corresponding to DNA polymerases derived from *Thermus* and *Thermotoga*):

Ala-Gly- X_1 -Pro-Phe-Asn- X_4 -Asn-Ser- X_7 - X_8 -Gln- X_9 - X_{10} -
 Arg- X_{12} -Leu- X_{13} - X_{14} - X_{15} -Leu (also referred to herein in the
 one-letter code as A-G- X_1 -P-F-N- X_4 -N-S- X_7 - X_8 -Q- X_9 - X_{10} -
 10 R- X_{12} -L- X_{13} - X_{14} - X_{15} -L); wherein:

X_1 is His (H) or Glu (E);

X_4 is Leu (L) or Ile (I);

X_7 is Arg (R) or Pro (P);

X_8 is Asp (D) or Lys (K);

15 X_9 is Leu (L) or Val (V);

X_{10} is Glu (E) or Ser (S);

X_{12} is Val (V) or Ile (I);

X_{13} is any amino acid other than Phe (F);

X_{14} is Asp (D) or Glu (E); and

20 X_{15} is Glu (E) or Lys (K) (SEQ ID NO:44).

In some embodiments, DNA polymerases of the invention can be characterized by having the following motif:

Ala-Gly-His-Pro-Phe-Asn-Leu-Asn- X_6 -Arg-Asp-Gln-Leu-
 X_{10} -Arg-Val-Leu- X_{13} -Asp-Glu-Leu (also referred to herein
 25 in the one-letter code as A-G-H-P-F-N-L-N- X_6 -R-D-Q-L-
 X_{10} -R-V-L- X_{13} -D-E-L); wherein

X_6 is any amino acid;

X_{10} is any amino acid; and

X_{13} is any amino acid other than Phe (F) (SEQ ID NO:45).

In some embodiments, DNA polymerases of the invention can be characterized by having the following motif:

Ala-Gly-His-Pro-Phe-Asn-Leu-Asn-X₆-Arg-Asp-Gln-Leu-
 X₁₀-Arg-Val-Leu-X₁₃-Asp-Glu-Leu (also referred to herein
 5 in the one-letter code as A-G-H-P-F-N-L-N-X₆-R-D-Q-L-
 X₁₀-R-V-L-X₁₃-D-E-L); wherein
 X₆ is any amino acid other than Ser (S);
 X₁₀ is any amino acid other than Glu (E); and
 X₁₃ is any amino acid other than Phe (F) (SEQ ID NO:46).

10 In some embodiments, DNA polymerases of the invention can be characterized by having the following motif:

Ala-Gly-His-Pro-Phe-Asn-Leu-Asn-X₆-Arg-Asp-Gln-Leu-X₁₀-Arg-
 Val-Leu-X₁₃-Asp-Glu-Leu (also referred to herein in the one-letter
 code as A-G-H-P-F-N-L-N-X₆-R-D-Q-L-X₁₀-R-V-L-X₁₃-D-E-L);
 15 wherein
 X₆ is Ser (S), Gly (G), Ala (A), Asp (D), Phe (F), Lys (K), Cys (C),
 Thr (T) or Tyr (Y);
 X₁₀ is Glu (E), Ser (S), Gln (Q), Ala (A), Gly (G), Lys (K) or Arg
 (R); and
 20 X₁₃ is Ala (A), Gly (G), Ser (S), Thr (T), Tyr (Y), Asp (D) or Lys
 (K) (SEQ ID NO:47).

This motif is present within the “thumb” domain of many Family A type DNA-dependent DNA polymerases, particularly thermostable DNA polymerases from thermophilic bacteria (Li et al., *EMBO J.* 17:7514-7525, 1998). For example, Figure 1 shows an amino acid
 25 sequence alignment comprising the native sequence corresponding to the motif above in DNA polymerases from several species of bacteria: *Escherichia coli*, *Bacillus caldotenax*, *Bacillus stearothermophilus*, *Deinococcus radiodurans*, *Thermosipho africanus*, *Thermotoga maritima*, *Thermotoga neopolitana*, *Thermus aquaticus*, *Thermus caldophilus*, *Thermus filiformis*, *Thermus flavus*, *Thermus* sp. Sps17, *Thermus* sp. Z05, and *Thermus thermophilus*. As shown,
 30 the motif of SEQ ID NO:8 (except where X13 is F) is present in each of these polymerases,

indicating a conserved function for this region of the polymerase. Figure 2 provides sequence identities among these DNA polymerases.

Accordingly, in some embodiments, the invention provides for a polymerase comprising SEQ ID NO:8, 9, 10, or 11 (e.g., where X₃ is selected, as appropriate in the consensus sequence, from G, A, V, L, I, M, W, P, S, T, C, N, Q, D, E, K, R, H, or Y), having the improved activity and/or characteristics described herein, and wherein the DNA polymerase is otherwise a wild-type or a naturally occurring DNA polymerase, such as, for example, a polymerase from any of the species of thermophilic bacteria listed above, or is substantially identical to such a wild-type or a naturally occurring DNA polymerase. For example, in some embodiments, the polymerase of the invention comprises SEQ ID NO:8, 9, 10, or 11 and is at least 80%, 85%, 90%, or 95% identical to SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 36, 37, 38, 39, 40, or 41. In one variation, the unmodified form of the polymerase is from a species of the genus *Thermus*. In some embodiments of the invention, the unmodified polymerase is from a thermophilic species other than *Thermus*, e.g., *Thermotoga*. The full nucleic acid and amino acid sequence for numerous thermostable DNA polymerases are available. The sequences each of *Thermus aquaticus* (Taq) (SEQ ID NO:2), *Thermus thermophilus* (Tth) (SEQ ID NO:6), *Thermus* species Z05 (SEQ ID NO:1), *Thermus* species Sps17 (SEQ ID NO:5), *Thermotoga maritima* (Tma) (SEQ ID NO:38), and *Thermosipho africanus* (Taf) (SEQ ID NO:37) polymerase have been published in PCT International Patent Publication No. WO 92/06200. The sequence for the DNA polymerase from *Thermus flavus* (SEQ ID NO:4) has been published in Akhmetzjanov and Vakhitov (*Nucleic Acids Research* 20:5839, 1992). The sequence of the thermostable DNA polymerase from *Thermus caldophilus* (SEQ ID NO:7) is found in EMBL/GenBank Accession No. U62584. The sequence of the thermostable DNA polymerase from *Thermus filiformis* can be recovered from ATCC Deposit No. 42380 using, e.g., the methods provided in U.S. Pat. No. 4,889,818, as well as the sequence information provided in Table 1. The sequence of the *Thermotoga neapolitana* DNA polymerase (SEQ ID NO:39) is from GeneSeq Patent Data Base Accession No. R98144 and PCT WO 97/09451. The sequence of the thermostable DNA polymerase from *Bacillus caldotenax* (SEQ ID NO:41) is described in, e.g., Uemori et al. (*J Biochem (Tokyo)* 113(3):401-410, 1993; *see also*, Swiss-Prot database Accession No. Q04957 and GenBank Accession Nos. D12982 and BAA02361). Examples of unmodified forms of DNA polymerases that can be modified as described herein

are also described in, e.g., U.S. Pat. Nos. 6,228,628; 6,346,379; 7,030,220; 6,881,559; 6,794,177; 6,468,775; and U.S. Pat. Appl. Nos. 20040005599; 20020012970; 20060078928; 20040115639. Representative full length polymerase sequences are also provided in the sequence listing.

- 5 In some embodiments, the polymerase of the invention, as well as having a polymerase domain comprising SEQ ID NOS:8, 9, 10, or 11, also comprises a nuclease domain (e.g., corresponding to positions 1 to 291 of Z05).

In some embodiments, a polymerase of the invention is a chimeric polymerase, i.e., comprising polypeptide regions from two or more enzymes. Examples of such chimeric DNA
10 polymerases are described in, e.g., U.S. Patent No. 6,228,628. Particularly suitable are chimeric CS-family DNA polymerases, which include the CS5 (SEQ ID NO:29) and CS6 (SEQ ID NO:30) polymerases and variants thereof having substantial sequence identity or similarity to SEQ ID NO:29 or SEQ ID NO:30 (typically at least 80% sequence identity and more typically at least 90%, most typically at least 95% sequence identity) and can thus be
15 modified to contain SEQ ID NO:8. The CS5 and CS6 DNA polymerases are chimeric enzymes derived from *Thermus* sp. Z05 and *Thermotoga maritima* (*Tma*) DNA polymerases. They comprise the N-terminal 5'-nuclease domain of the *Thermus* enzyme and the C-terminal 3'-5' exonuclease and the polymerase domains of the *Tma* enzyme. These enzymes have efficient reverse transcriptase activity, can extend nucleotide analog-containing primers, and
20 can utilize alpha-phosphorothioate dNTPs, dUTP, dITP, and also fluorescein- and cyanine-dye family labeled dNTPs. The CS5 and CS6 polymerases are also efficient Mg^{2+} -activated PCR enzymes. The CS5 and CS6 chimeric polymerases are further described in, e.g., U.S. Pat. Application Publication No. 2004/0005599.

In some embodiments, the polymerase of the invention comprises SEQ ID NO:8, 9, 10, or 11
25 and further comprises one or more additional amino acid changes (e.g., by amino acid substitution, addition, or deletion) compared to a native polymerase. In some embodiments, such polymerases retain the amino acid motif of SEQ ID NO:8 (or a motif of SEQ ID NO:9, 10 or 11), and further comprise the amino acid motif of SEQ ID NO:27 (corresponding to the D580X mutation of Z05 (SEQ ID NO:1)) as follows:

T-G-R-L-S-S-X₇-X₈-P-N-L-Q-N; wherein

X₇ is Ser (S) or Thr (T); and

X₈ is any amino acid other than D or E (SEQ ID NO:27)

5 The mutation characterized by SEQ ID NO:27 is discussed in more detail in, e.g., US Patent Publication No. 2009/0148891. In some embodiments, such functional variant polymerases typically will have substantial sequence identity or similarity to the wild-type or naturally occurring polymerase (e.g., SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 39, 40, 41, 42, 43, or 44), typically at least 80% sequence identity and more typically at least 90%, most typically at least 95% sequence identity.

10 In some embodiments, the amino acid at position X₁₃ is substituted with an amino acid as set forth in SEQ ID NO:8, 9, 10 or 11, and the amino acid at position X₈ is substituted with an amino acid as set forth in SEQ ID NO:27. Thus, in some embodiments, the amino acid at position X₁₃ is any amino acid other than Phe (F) and the amino acid at position X₈ is any amino acid other than Asp (D) or Glu (E). In some embodiments, amino acid substitutions
15 include Leucine (L), Glycine (G), Threonine (T), Glutamine (Q), Alanine (A), Serine (S), Asparagine (N), Arginine (R), and Lysine (K) at position X₈ of SEQ ID NO:27. In certain embodiments, amino acid substitutions independently include Alanine (A), Glycine (G), Serine (S), Threonine (T), Tyrosine (Y), Aspartic Acid (D), or Lysine (K) at position X₁₃, and Glycine (G) at position X₈. Other suitable amino acid substitution(s) at one or more of the
20 identified sites can be determined using, e.g., known methods of site-directed mutagenesis and determination of polynucleotide extension performance in assays described further herein or otherwise known to persons of skill in the art.

Because the precise length of DNA polymerases vary, the precise amino acid positions corresponding to each of X₁₃ and X₈ can vary depending on the particular polymerase used.
25 Amino acid and nucleic acid sequence alignment programs are readily available (*see, e.g.*, those referred to *supra*) and, given the particular motifs identified herein, serve to assist in the identification of the exact amino acids (and corresponding codons) for modification in accordance with the present invention. The positions corresponding to each of X₁₃ and X₈ are shown in Table 1 for representative chimeric thermostable DNA polymerases and
30 thermostable DNA polymerases from exemplary thermophilic species.

Table 1. Amino Acid Positions Corresponding to Motif Positions X₆, X₁₀, X₁₃ (e.g., of SEQ ID NOS:8, 9, 10, and 11) and X₈ (of SEQ ID NO:27) in Exemplary Polymerases.

<u>Organism or Chimeric Sequence</u>	<u>Amino Acid Position</u>			
Consensus (SEQ ID NO:)	X ₆	X ₁₀	X ₁₃	X ₈ (of SEQ ID NO:27)
<i>T. thermophilus</i> (6)	488	493	497	580
<i>T. caldophilus</i> (7)	488	493	497	580
<i>T. sp. Z05</i> (1)	488	493	497	580
<i>T. aquaticus</i> (2)	486	491	495	578
<i>T. flavus</i> (4)	485	490	494	577
<i>T. filiformis</i> (3)	484	489	493	576
<i>T. sp. Sps17</i> (5)	484	489	493	576
<i>T. maritima</i> (38)	548	553	557	640
<i>T. neapolitana</i> (39)	548	553	557	640
<i>T. africanus</i> (37)	548	553	557	639
<i>B. caldotenax</i> (41)	530	535	539	621
<i>B. stearothermophilus</i> (40)	530	535	539	620
CS5 (29)	548	553	557	640
CS6 (30)	548	553	557	640

- 5 In some embodiments, the DNA polymerase of the present invention is derived from *Thermus*
sp. Z05 DNA polymerase (SEQ ID NO:1) or a variant thereof (e.g., carrying the D580G
mutation or the like). As referred to above, in *Thermus sp. Z05* DNA polymerase, position X₆
corresponds to Serine (S) at position 488; position X₁₀ corresponds to Glutamic acid (E) at
position 493; and position X₁₃ corresponds to Phenylalanine (F) at position 497; position X₈
10 corresponds to Aspartate (D) at position 580. Thus, in certain variations of the invention, the
mutant polymerase comprises at least one amino acid substitution, relative to a *Thermus sp.*
Z05 DNA polymerase, at S488, E493, F497, and D580. Thus, typically, the amino acid at


position S488 is not S. In some embodiments, the amino acid at position 488 is selected from G, A, V, L, I, M, F, W, P, T, C, Y, N, Q, D, E, K, R or H. In certain embodiments, amino acid residue at position S488 is G, A, D, F, K, C, T or Y. Typically, the amino acid at position E493, if substituted, is not E. In some embodiments, the amino acid at position 493 is selected from G, A, V, L, I, M, F, W, P, T, C, Y, N, Q, D, S, K, R or H. In certain embodiments, the amino acid residue at position E493 can be substituted or unsubstituted, and is E, S, A, Q, G, K or R. Typically, the amino acid at position F497, if substituted, is not F. In some embodiments, the amino acid at position 497 is selected from G, A, V, L, I, M, S, W, P, T, C, Y, N, Q, D, E, K, R or H. In certain embodiments, amino acid residue at position F497 can be substituted or unsubstituted, and is F, A, S, G, T, Y, D or K. In certain embodiments, amino acid residues at position D580 can be selected from Leucine (L), Glycine (G), Threonine (T), Glutamine (Q), Alanine (A), Serine (S), Asparagine (N), Arginine (R), and Lysine (K). Exemplary *Thermus sp.* Z05 DNA polymerase mutants include those comprising the amino acid substitution(s) F497A, F497G, F497S, F497T, F497Y, F497D, or F497K and D580G.

15 In some embodiments, the DNA polymerase of the invention is derived from a *Thermus* species, and the amino acid corresponding to position 497 of SEQ ID NO:1 is an amino acid having a polar, uncharged side-chain (e.g., N, Q, H, S, T, or Y), a nonpolar, uncharged side-chain (other than F, e.g., G, A, L, M, W, P, C, V, or I), a polar, positively charged side-chain (e.g., R or K), or polar, negatively charged side-chain (e.g., D or E) at neutral pH (e.g., about pH 7.4). In some embodiments, the amino acid corresponding to position 497 of SEQ ID NO:1 having a polar, uncharged side-chain is S, T, or Y. In some embodiments, the amino acid corresponding to position 497 of SEQ ID NO:1 having a nonpolar, uncharged side-chain is A or G. In some embodiments, the amino acid corresponding to position 497 of SEQ ID NO:1 having a polar, positively charged side-chain is K. In some embodiments, the amino acid corresponding to position 497 of SEQ ID NO:1 having a polar, negatively charged side-chain is D.


In some embodiments, the DNA polymerases of the present invention can also include other, non-substitutional modification(s). Such modifications can include, for example, covalent modifications known in the art to confer an additional advantage in applications comprising polynucleotide extension. For example, in certain embodiments, the mutant DNA polymerase further includes a thermally reversible covalent modification. DNA polymerases comprising

such thermally reversible modifications are particularly suitable for hot-start applications, such as, *e.g.*, various hot-start PCR techniques. Thermally reversible modifier reagents amenable to use in accordance with the mutant DNA polymerases of the present invention are described in, for example, U.S. Patent No. 5,773,258 to Birch *et al.*.

- 5 For example, particularly suitable polymerases comprising a thermally reversible covalent modification are produced by a reaction, carried out at alkaline pH at a temperature which is less than about 25 °C, of a mixture of a thermostable enzyme and a dicarboxylic acid anhydride having a general formula as set forth in the following formula I:



- 10 where R₁ and R₂ are hydrogen or organic radicals, which may be linked; or having the following formula II:



where R₁ and R₂ are organic radicals, which may linked, and the hydrogens are cis, essentially as described in Birch *et al, supra*.

- 15 The DNA polymerases of the present invention can be constructed by mutating the DNA sequences that encode the corresponding unmodified polymerase (*e.g.*, a wild-type polymerase or a corresponding variant from which the polymerase of the invention is derived), such as by using techniques commonly referred to as site-directed mutagenesis. Nucleic acid molecules encoding the unmodified form of the polymerase can be mutated by a variety of polymerase
20 chain reaction (PCR) techniques well-known to one of ordinary skill in the art. (*See, e.g., PCR Strategies* (M. A. Innis, D. H. Gelfand, and J. J. Sninsky eds., 1995, Academic Press, San Diego, CA) at Chapter 14; *PCR Protocols : A Guide to Methods and Applications* (M. A. Innis, D. H. Gelfand, J. J. Sninsky, and T. J. White eds., Academic Press, NY, 1990).

By way of non-limiting example, the two primer system, utilized in the Transformer Site-Directed Mutagenesis kit from Clontech, may be employed for introducing site-directed mutants into a polynucleotide encoding an unmodified form of the polymerase. Following denaturation of the target plasmid in this system, two primers are simultaneously annealed to the plasmid; one of these primers contains the desired site-directed mutation, the other contains a mutation at another point in the plasmid resulting in elimination of a restriction site. Second strand synthesis is then carried out, tightly linking these two mutations, and the resulting plasmids are transformed into a mutS strain of *E. coli*. Plasmid DNA is isolated from the transformed bacteria, restricted with the relevant restriction enzyme (thereby linearizing the unmutated plasmids), and then retransformed into *E. coli*. This system allows for generation of mutations directly in an expression plasmid, without the necessity of subcloning or generation of single-stranded phagemids. The tight linkage of the two mutations and the subsequent linearization of unmutated plasmids result in high mutation efficiency and allow minimal screening. Following synthesis of the initial restriction site primer, this method requires the use of only one new primer type per mutation site. Rather than prepare each positional mutant separately, a set of "designed degenerate" oligonucleotide primers can be synthesized in order to introduce all of the desired mutations at a given site simultaneously. Transformants can be screened by sequencing the plasmid DNA through the mutagenized region to identify and sort mutant clones. Each mutant DNA can then be restricted and analyzed by electrophoresis, such as for example, on a Mutation Detection Enhancement gel (Mallinckrodt Baker, Inc., Phillipsburg, NJ) to confirm that no other alterations in the sequence have occurred (by band shift comparison to the unmutagenized control). Alternatively, the entire DNA region can be sequenced to confirm that no additional mutational events have occurred outside of the targeted region.

Verified mutant duplexes in pET (or other) overexpression vectors can be employed to transform *E. coli* such as, *e.g.*, strain *E. coli* BL21 (DE3) pLysS, for high level production of the mutant protein, and purification by standard protocols. The method of FAB-MS mapping, for example, can be employed to rapidly check the fidelity of mutant expression. This technique provides for sequencing segments throughout the whole protein and provides the necessary confidence in the sequence assignment. In a mapping experiment of this type, protein is digested with a protease (the choice will depend on the specific region to be

modified since this segment is of prime interest and the remaining map should be identical to the map of unmutagenized protein). The set of cleavage fragments is fractionated by, for example, microbore HPLC (reversed phase or ion exchange, again depending on the specific region to be modified) to provide several peptides in each fraction, and the molecular weights of the peptides are determined by standard methods, such as FAB-MS. The determined mass of each fragment are then compared to the molecular weights of peptides expected from the digestion of the predicted sequence, and the correctness of the sequence quickly ascertained. Since this mutagenesis approach to protein modification is directed, sequencing of the altered peptide should not be necessary if the MS data agrees with prediction. If necessary to verify a changed residue, CAD-tandem MS/MS can be employed to sequence the peptides of the mixture in question, or the target peptide can be purified for subtractive Edman degradation or carboxypeptidase Y digestion depending on the location of the modification.

Mutant DNA polymerases with more than one amino acid substituted can be generated in various ways. In the case of amino acids located close together in the polypeptide chain, they may be mutated simultaneously using one oligonucleotide that codes for all of the desired amino acid substitutions. If however, the amino acids are located some distance from each other (separated by more than ten amino acids, for example) it is more difficult to generate a single oligonucleotide that encodes all of the desired changes. Instead, one of two alternative methods may be employed. In the first method, a separate oligonucleotide is generated for each amino acid to be substituted. The oligonucleotides are then annealed to the single-stranded template DNA simultaneously, and the second strand of DNA that is synthesized from the template will encode all of the desired amino acid substitutions. An alternative method involves two or more rounds of mutagenesis to produce the desired mutant. The first round is as described for the single mutants: DNA encoding the unmodified polymerase is used for the template, an oligonucleotide encoding the first desired amino acid substitution(s) is annealed to this template, and the heteroduplex DNA molecule is then generated. The second round of mutagenesis utilizes the mutated DNA produced in the first round of mutagenesis as the template. Thus, this template already contains one or more mutations. The oligonucleotide encoding the additional desired amino acid substitution(s) is then annealed to this template, and the resulting strand of DNA now encodes mutations from both the first and second rounds of mutagenesis. This resultant DNA can be used as a template in a third round

of mutagenesis, and so on. Alternatively, the multi-site mutagenesis method of Seyfang & Jin (*Anal. Biochem.* 324:285-291. 2004) may be utilized.

Accordingly, also provided are recombinant nucleic acids encoding any of the DNA polymerases of the present invention (e.g., polymerases comprising any of SEQ ID NOS:8, 9, 10, or 11). Using a nucleic acid of the present invention, encoding a DNA polymerase of the invention, a variety of vectors can be made. Any vector containing replicon and control sequences that are derived from a species compatible with the host cell can be used in the practice of the invention. Generally, expression vectors include transcriptional and translational regulatory nucleic acid regions operably linked to the nucleic acid encoding the mutant DNA polymerase. The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. In addition, the vector may contain a Positive Retroregulatory Element (PRE) to enhance the half-life of the transcribed mRNA (*see* Gelfand *et al.* U.S. Patent No. 4,666,848). The transcriptional and translational regulatory nucleic acid regions will generally be appropriate to the host cell used to express the polymerase. Numerous types of appropriate expression vectors, and suitable regulatory sequences are known in the art for a variety of host cells. In general, the transcriptional and translational regulatory sequences may include, e.g., promoter sequences, ribosomal binding sites, transcriptional start and stop sequences, translational start and stop sequences, and enhancer or activator sequences. In typical embodiments, the regulatory sequences include a promoter and transcriptional start and stop sequences. Vectors also typically include a polylinker region containing several restriction sites for insertion of foreign DNA. In certain embodiments, "fusion flags" are used to facilitate purification and, if desired, subsequent removal of tag/flag sequence, e.g., "His-Tag". However, these are generally unnecessary when purifying an thermoactive and/or thermostable protein from a mesophilic host (e.g., *E. coli*) where a "heat-step" may be employed. The construction of suitable vectors containing DNA encoding replication sequences, regulatory sequences, phenotypic selection genes, and the mutant polymerase of interest are prepared using standard recombinant DNA procedures. Isolated plasmids, viral vectors, and DNA fragments are cleaved, tailored, and ligated together in a specific order to generate the desired vectors, as is well-known in the art (*see, e.g.,*

Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, New York, NY, 2nd ed. 1989)).

In certain embodiments, the expression vector contains a selectable marker gene to allow the selection of transformed host cells. Selection genes are well known in the art and will vary
5 with the host cell used. Suitable selection genes can include, for example, genes coding for ampicillin and/or tetracycline resistance, which enables cells transformed with these vectors to grow in the presence of these antibiotics.

In one aspect of the present invention, a nucleic acid encoding a DNA polymerase of the invention is introduced into a cell, either alone or in combination with a vector. By
10 "introduced into" or grammatical equivalents herein is meant that the nucleic acids enter the cells in a manner suitable for subsequent integration, amplification, and/or expression of the nucleic acid. The method of introduction is largely dictated by the targeted cell type. Exemplary methods include CaPO₄ precipitation, liposome fusion, LIPOFECTIN®, electroporation, viral infection, and the like.

15 In some embodiments, prokaryotes are used as host cells for the initial cloning steps of the present invention. They are particularly useful for rapid production of large amounts of DNA, for production of single-stranded DNA templates used for site-directed mutagenesis, for screening many mutants simultaneously, and for DNA sequencing of the mutants generated. Suitable prokaryotic host cells include *E. coli* K12 strain 94 (ATCC No. 31,446), *E. coli* strain
20 W3110 (ATCC No. 27,325), *E. coli* K12 strain DG116 (ATCC No. 53,606), *E. coli* X1776 (ATCC No. 31,537), and *E. coli* B; however many other strains of *E. coli*, such as HB101, JM101, NM522, NM538, NM539, and many other species and genera of prokaryotes including bacilli such as *Bacillus subtilis*, other enterobacteriaceae such as *Salmonella typhimurium* or *Serratia marcesans*, and various *Pseudomonas* species can all be used as
25 hosts. Prokaryotic host cells or other host cells with rigid cell walls are typically transformed using the calcium chloride method as described in section 1.82 of Sambrook *et al.*, *supra*. Alternatively, electroporation can be used for transformation of these cells. Prokaryote transformation techniques are set forth in, for example Dower, in *Genetic Engineering, Principles and Methods* 12:275-296 (Plenum Publishing Corp., 1990); Hanahan *et al.*, *Meth.*
30 *Enzymol.*, 204:63, 1991. Plasmids typically used for transformation of *E. coli* include pBR322,

pUCI8, pUCI9, pUCI18, pUCI19, and Bluescript M13, all of which are described in sections 1.12-1.20 of Sambrook *et al.*, *supra*. However, many other suitable vectors are available as well.

In some embodiments, the DNA polymerases of the present invention are produced by
5 culturing a host cell transformed with an expression vector containing a nucleic acid encoding the DNA polymerase, under the appropriate conditions to induce or cause expression of the DNA polymerase. Methods of culturing transformed host cells under conditions suitable for protein expression are well-known in the art (*see, e.g.*, Sambrook *et al.*, *supra*). Suitable host
10 cells for production of the polymerases from lambda pL promotor-containing plasmid vectors include *E. coli* strain DG116 (ATCC No. 53606) (*see* US Pat. No. 5,079,352 and Lawyer, F.C. *et al.*, *PCR Methods and Applications* 2:275-87, 1993). Following expression, the polymerase can be harvested and isolated. Methods for purifying the thermostable DNA polymerase are described in, for example, Lawyer *et al.*, *supra*.

Once purified, a DNA polymerase's 3' mismatch discrimination can be assayed. For example,
15 in some embodiments, 3' mismatch discrimination activity is determined by comparing the amplification of a target sequence perfectly matched to the primer to amplification of a target that has a single base mismatch at the 3' end of the primer. Amplification can be detected, for example, in real time by use of TaqMan™ probes. Ability of a polymerase to distinguish between the two target sequences can be estimated by comparing the Cps of the two reactions.
20 Optionally, simultaneous amplification of a second target gene in each well can be performed and detected in a second optical channel as a control. "Delta Cp values" refer to the difference in value between the Cp associated with the mismatched template minus the Cp of the matched target (*see, e.g.*, the Examples). In some embodiments, the improved polymerases of the invention have a delta Cp value of at least 1, 2, 3, 4, 5, or more compared to an otherwise
25 identical control polymerase having a native amino acid (*e.g.*, N) at position X₃ of SEQ ID NO:8. In some embodiments, this determination is made with the precise materials and conditions set forth in the Examples.

Methods of the invention

The improved DNA polymerases of the present invention may be used for any purpose in
30 which such enzyme activity is necessary or desired. The improved DNA polymerase can be a

thermoactive or thermostable DNA polymerase, as described herein. Accordingly, in one aspect of the invention, methods of polynucleotide extension, including PCR, using the polymerases of the invention are provided. In some embodiments, the invention provides a thermoactive DNA polymerase that is useful to extend an RNA or DNA template when
5 amplification of the template nucleic acid is not required, for example, when it is desired to immediately detect the presence of a target nucleic acid. In some embodiments, the invention provides a thermostable DNA polymerase that is useful when it is desired to extend and/or amplify a target nucleic acid. Conditions suitable for polynucleotide extension are known in the art. (See, e.g., Sambrook *et al.*, *supra*. See also Ausubel *et al.*, *Short Protocols in*
10 *Molecular Biology* (4th ed., John Wiley & Sons 1999). Generally, a primer is annealed, *i.e.*, hybridized, to a target nucleic acid to form a primer-template complex. The primer-template complex is contacted with the mutant DNA polymerase and nucleoside triphosphates in a suitable environment to permit the addition of one or more nucleotides to the 3' end of the primer, thereby producing an extended primer complementary to the target nucleic acid. The
15 primer can include, *e.g.*, one or more nucleotide analog(s). In addition, the nucleoside triphosphates can be conventional nucleotides, unconventional nucleotides (*e.g.*, ribonucleotides or labeled nucleotides), or a mixture thereof. In some variations, the polynucleotide extension reaction comprises amplification of a target nucleic acid. Conditions suitable for nucleic acid amplification using a DNA polymerase and a primer pair are also
20 known in the art (*e.g.*, PCR amplification methods). (See, *e.g.*, Sambrook *et al.*, *supra*; Ausubel *et al.*, *supra*; *PCR Applications: Protocols for Functional Genomics* (Innis *et al.* eds., Academic Press 1999).

In some embodiments, use of the present polymerases, which provide increased 3' mismatch discrimination, allow for, *e.g.*, rare allele detection. For example, the fidelity of 3' mismatch
25 discrimination of a particular polymerase sets its sensitivity (ability to accurately detect small quantities of a target sequence in the presence of larger quantities of a different but related non-target sequence). Thus, increased 3'- mismatch discrimination results in greater sensitivity for detection of rare alleles. Rare allele detection is useful, for example, when screening biopsies or other samples for rare genetic changes, *e.g.*, a cell carrying a cancer allele in a
30 mass of normal cells.

In some embodiments, the improved polymerases are used for polynucleotide extension in the context of allele specific PCR or single nucleotide polymorphism (SNP) detection. Exemplary SNP detection methods are described in Chen et al., "Single nucleotide polymorphism genotyping: biochemistry, protocol, cost and throughput" *Pharmacogenomics J.* 3(2):77-96 (2003); Kwok et al., "Detection of single nucleotide polymorphisms" *Curr. Issues Mol. Biol.* 5(2):43-60 (April 2003); Shi, "Technologies for individual genotyping: detection of genetic polymorphisms in drug targets and disease genes" *Am. J. Pharmacogenomics* 2(3):197-205 (2002); and Kwok, "Methods for genotyping single nucleotide polymorphisms" *Annu. Rev. Genomics Hum. Genet.* 2:235-58 (2001). Exemplary techniques for high-throughput SNP detection are described in Marnellos, "High-throughput SNP analysis for genetic association studies" *Curr. Opin. Drug Discov. Devel.* 6(3):317-21 (May 2003). Common SNP detection methods include, but are not limited to, TaqMan assays, molecular beacon assays, nucleic acid arrays, allele-specific primer extension, allele-specific PCR, arrayed primer extension, homogeneous primer extension assays, primer extension with detection by mass spectrometry, pyrosequencing, multiplex primer extension sorted on genetic arrays, ligation with rolling circle amplification, homogeneous ligation, OLA (U.S. Pat. No. 4,988,167), multiplex ligation reaction sorted on genetic arrays, restriction-fragment length polymorphism, single base extension-tag assays, and the Invader assay. Such methods may be used in combination with detection mechanisms such as, for example, luminescence or chemiluminescence detection, fluorescence detection, time-resolved fluorescence detection, fluorescence resonance energy transfer, fluorescence polarization, mass spectrometry, and electrical detection.

Detection of multiple different alleles can also be accomplished using multiplex reactions, which allow the detection of multiple different alleles in a single reaction. In multiplex reactions, two or more allele-specific primers are used to extend and amplify SNPs or multiple nucleotide polymorphisms or alleles. Exemplary methods for multiplex detection of single and multiple nucleotide polymorphisms are described in U.S. Patent Publication No. 2006/0172324.

Other methods for detecting extension products or amplification products using the improved polymerases described herein include the use of fluorescent double-stranded nucleotide binding dyes or fluorescent double-stranded nucleotide intercalating dyes. Examples of fluorescent double-stranded DNA binding dyes include SYBR-green (Molecular Probes).

Examples of fluorescent double-stranded intercalating dyes include ethidium bromide. The double stranded DNA binding dyes can be used in conjunction with melting curve analysis to measure primer extension products and/or amplification products. The melting curve analysis can be performed on a real-time PCR instrument, such as the ABI 5700/7000 (96 well format) or ABI 7900 (384 well format) instrument with onboard software (SDS 2.1). Alternatively, the melting curve analysis can be performed as an end point analysis. Exemplary methods of melting point analysis are described in U.S. Patent Publication No. 2006/0172324.

In yet other embodiments, the polymerases of the invention are used for primer extension in the context of DNA sequencing, DNA labeling, or labeling of primer extension products. For example, DNA sequencing by the Sanger dideoxynucleotide method (Sanger *et al.*, *Proc. Natl. Acad. Sci. USA* 74: 5463, 1977) is improved by the present invention for polymerases capable of incorporating unconventional, chain-terminating nucleotides. Advances in the basic Sanger *et al.* method have provided novel vectors (Yanisch-Perron *et al.*, *Gene* 33:103-119, 1985) and base analogues (Mills *et al.*, *Proc. Natl. Acad. Sci. USA* 76:2232-2235, 1979; and Barr *et al.*, *Biotechniques* 4:428-432, 1986). In general, DNA sequencing requires template-dependent primer extension in the presence of chain-terminating base analogs, resulting in a distribution of partial fragments that are subsequently separated by size. The basic dideoxy sequencing procedure involves (i) annealing an oligonucleotide primer, optionally labeled, to a template; (ii) extending the primer with DNA polymerase in four separate reactions, each containing a mixture of unlabeled dNTPs and a limiting amount of one chain terminating agent such as a ddNTP, optionally labeled; and (iii) resolving the four sets of reaction products on a high-resolution denaturing polyacrylamide/urea gel. The reaction products can be detected in the gel by autoradiography or by fluorescence detection, depending on the label used, and the image can be examined to infer the nucleotide sequence. These methods utilize DNA polymerase such as the Klenow fragment of *E. coli* Pol I or a modified T7 DNA polymerase.

The availability of thermostable polymerases, such as Taq DNA polymerase, resulted in improved methods for sequencing with thermostable DNA polymerase (*see* Innis *et al.*, *Proc. Natl. Acad. Sci. USA* 85:9436, 1988) and modifications thereof referred to as "cycle sequencing" (Murray, *Nuc Acids Res.* 17:8889, 1989). Accordingly, thermostable polymerases of the present invention can be used in conjunction with such methods. As an alternative to basic dideoxy sequencing, cycle sequencing is a linear, asymmetric amplification of target

sequences complementary to the template sequence in the presence of chain terminators. A single cycle produces a family of extension products of all possible lengths. Following denaturation of the extension reaction product from the DNA template, multiple cycles of primer annealing and primer extension occur in the presence of terminators such as ddNTPs.

- 5 Cycle sequencing requires less template DNA than conventional chain-termination sequencing. Thermostable DNA polymerases have several advantages in cycle sequencing; they tolerate the stringent annealing temperatures which are required for specific hybridization of primer to nucleic acid targets as well as tolerating the multiple cycles of high temperature denaturation which occur in each cycle, *e.g.*, 90-95 °C. For this reason, AMPLITAQ® DNA
10 Polymerase and its derivatives and descendants, *e.g.*, AmpliTaq CS DNA Polymerase and AmpliTaq FS DNA Polymerase have been included in Taq cycle sequencing kits commercialized by companies such as Perkin-Elmer (Norwalk, CT) and Applied Biosystems (Foster City, CA).

- The improved polymerases find use in 454 sequencing (Roche) (Margulies, M et al. 2005,
15 Nature, 437, 376-380). 454 sequencing involves two steps. In the first step, DNA is sheared into fragments of approximately 300-800 base pairs, and the fragments are blunt ended. Oligonucleotide adaptors are then ligated to the ends of the fragments. The adaptors serve as primers for amplification and sequencing of the fragments. The fragments can be attached to DNA capture beads, *e.g.*, streptavidin-coated beads using, *e.g.*, Adaptor B, which contains 5'-
20 biotin tag. The fragments attached to the beads are PCR amplified within droplets of an oil-water emulsion. The result is multiple copies of clonally amplified DNA fragments on each bead. In the second step, the beads are captured in wells (pico-liter sized). Pyrosequencing is performed on each DNA fragment in parallel. Addition of one or more nucleotides generates a light signal that is recorded by a CCD camera in a sequencing instrument. The signal strength
25 is proportional to the number of nucleotides incorporated.

Pyrosequencing makes use of pyrophosphate (PPi) which is released upon nucleotide addition. PPi is converted to ATP by ATP sulfurylase in the presence of adenosine 5' phosphosulfate. Luciferase uses ATP to convert luciferin to oxyluciferin, and this reaction generates light that is detected and analyzed.

Variations of chain termination sequencing methods include dye-primer sequencing and dye-terminator sequencing. In dye-primer sequencing, the ddNTP terminators are unlabeled, and a labeled primer is utilized to detect extension products (Smith *et al.*, *Nature* 32:674-679, 1986). In dye-terminator DNA sequencing, a DNA polymerase is used to incorporate dNTPs and
5 fluorescently labeled ddNTPs onto the end of a DNA primer (Lee *et al.*, *Nuc. Acids. Res.* 20:2471, 1992). This process offers the advantage of not having to synthesize dye labeled primers. Furthermore, dye-terminator reactions are more convenient in that all four reactions can be performed in the same tube.

Both dye-primer and dye-terminator methods may be automated using an automated
10 sequencing instrument produced by Applied Biosystems (Foster City, CA) (U.S. Pat. No. 5,171,534). When using the instrument, the completed sequencing reaction mixture is fractionated on a denaturing polyacrylamide gel or capillaries mounted in the instrument. A laser at the bottom of the instrument detects the fluorescent products as they are electrophoretically separated according to size through the gel.

15 Two types of fluorescent dyes are commonly used to label the terminators used for dye-terminator sequencing-negatively charged and zwitterionic fluorescent dyes. Negatively charged fluorescent dyes include those of the fluorescein and BODIPY families. BODIPY dyes (4,4-difluoro-4-bora-3a,4a-diaza-s-indacene) are described in International Patent Publication WO 97/00967. Zwitterionic fluorescent dyes include those of the rhodamine
20 family. Commercially available cycle sequencing kits use terminators labeled with rhodamine derivatives. However, the rhodamine-labeled terminators are rather costly and the product must be separated from unincorporated dye-ddNTPs before loading on the gel since they co-migrate with the sequencing products. Rhodamine dye family terminators seem to stabilize hairpin structures in GC-rich regions, which causes the products to migrate anomalously. This
25 can involve the use of dITP, which relaxes the secondary structure but also affects the efficiency of incorporation of terminator.

In contrast, fluorescein-labeled terminators eliminate the separation step prior to gel loading since they have a greater net negative charge and migrate faster than the sequencing products. In addition, fluorescein-labeled sequencing products have better electrophoretic migration than
30 sequencing products labeled with rhodamine. Although wild-type Taq DNA polymerase does

not efficiently incorporate terminators labeled with fluorescein family dyes, this can now be accomplished efficiently by use of the modified enzymes as described in U.S. Patent Application Publication No. 2002/0142333. Accordingly, modifications as described in US 2002/0142333 can be used in the context of the present invention to produce fluorescein-
5 family-dye-incorporating thermostable polymerases having improved primer extension rates. For example, in certain embodiments, the unmodified DNA polymerase in accordance with the present invention is a modified thermostable polymerase as described in US 2002/0142333 and having the motif set forth in SEQ ID NO:8 (or a motif of SEQ ID NO:9, 10 or 11), and optionally the motif of SEQ ID NO:27.

- 10 Other exemplary nucleic acid sequencing formats in which the mutant DNA polymerases of the invention can be used include those involving terminator compounds that include 2'-PO₄ analogs of ribonucleotides (*see, e.g.*, U.S. Application Publication Nos. 2005/0037991 and 2005/0037398, and US Patent Application No. 12/174,488).

Kits

- 15 In another aspect of the present invention, kits are provided for use in primer extension methods described herein. In some embodiments, the kit is compartmentalized for ease of use and contains at least one container providing a DNA polymerase of the invention having increased 3' mismatch discrimination in accordance with the present invention. One or more additional containers providing additional reagent(s) can also be included. Such additional
20 containers can include any reagents or other elements recognized by the skilled artisan for use in primer extension procedures in accordance with the methods described above, including reagents for use in, *e.g.*, nucleic acid amplification procedures (*e.g.*, PCR, RT-PCR), DNA sequencing procedures, or DNA labeling procedures. For example, in certain embodiments, the kit further includes a container providing a 5' sense primer hybridizable, under primer
25 extension conditions, to a predetermined polynucleotide template, or a primer pair comprising the 5' sense primer and a corresponding 3' antisense primer. In some embodiments, the kit includes one or more containers containing one or more primers that are fully complementary to single nucleotide polymorphisms or multiple nucleotide polymorphisms, wherein the primers are useful for multiplex reactions, as described above. In other, non-mutually
30 exclusive variations, the kit includes one or more containers providing nucleoside

triphosphates (conventional and/or unconventional). In specific embodiments, the kit includes alpha-phosphorothioate dNTPs, dUTP, dITP, and/or labeled dNTPs such as, *e.g.*, fluorescein- or cyanin-dye family dNTPs. In still other, non-mutually exclusive embodiments, the kit includes one or more containers providing a buffer suitable for a primer extension reaction. In
5 some embodiments, the kit includes one or more labeled or unlabeled probes. Examples of probes include dual-labeled FRET (fluorescence resonance energy transfer) probes and molecular beacon probes. In another embodiment, the kit contains an aptamer, *e.g.*, for hot start PCR assays.

Reaction Mixtures

10 In another aspect of the present invention, reaction mixtures are provided comprising the polymerases with increased 3'-mismatch discrimination activity, as described herein. The reaction mixtures can further comprise reagents for use in, *e.g.*, nucleic acid amplification procedures (*e.g.*, PCR, RT-PCR), DNA sequencing procedures, or DNA labeling procedures. For example, in certain embodiments, the reaction mixtures comprise a buffer suitable for a
15 primer extension reaction. The reaction mixtures can also contain a template nucleic acid (DNA and/or RNA), one or more primer or probe polynucleotides, nucleoside triphosphates (including, *e.g.*, deoxyribonucleotides, ribonucleotides, labeled nucleotides, unconventional nucleotides), salts (*e.g.*, Mn^{2+} , Mg^{2+}), and labels (*e.g.*, fluorophores). In some embodiments, the reaction mixture further comprises double stranded DNA binding dyes, such as SYBR
20 green, or double stranded DNA intercalating dyes, such as ethidium bromide. In some embodiments, the reaction mixtures contain a 5'-sense primer hybridizable, under primer extension conditions, to a predetermined polynucleotide template, or a primer pair comprising the 5'-sense primer and a corresponding 3' antisense primer. In certain embodiments, the reaction mixture further comprises a fluorogenic FRET hydrolysis probe for detection of
25 amplified template nucleic acids, for example a Taqman® probe. In some embodiments, the reaction mixture contains two or more primers that are fully complementary to single nucleotide polymorphisms or multiple nucleotide polymorphisms. In some embodiments, the reaction mixtures contain alpha-phosphorothioate dNTPs, dUTP, dITP, and/or labeled dNTPs such as, *e.g.*, fluorescein- or cyanin-dye family dNTPs.

EXAMPLES

The following examples are offered to illustrate, but not to limit the claimed invention.

Example 1: Identification of mutant DNA polymerases with increased 3'-mismatch discrimination

Mutations in Z05 D580G polymerase were identified that provide a reduced ability to extend an oligonucleotide primer with a 3'-mismatch to a template. In brief, the steps in this screening process included library generation, expression and partial purification of the mutant enzymes, screening of the enzymes for the desired property, DNA sequencing, clonal purification, and further characterization of selected candidate mutants. Each of these steps is described further below.

Clonal Library generation: A nucleic acid encoding the polymerase domain of Z05 D580G DNA polymerase was subjected to error-prone (mutagenic) PCR between Bln I and Bgl II restriction sites of a plasmid including this nucleic acid sequence. The amplified sequence is provided as SEQ ID NO:33. The primers used for this are given below:

Forward Primer: 5'- CTACCTCCTGGACCCCTCCAA-3' (SEQ ID NO:31); and,

Reverse Primer: 5'- ATAACCAACTGGTAGTGGCGTGTA-3' (SEQ ID NO:32).

PCR was performed using a range of Mg^{2+} concentrations from 1.8-3.6 mM, in order to generate libraries with a range of mutation rates. Buffer conditions were 50 mM Bicine pH 8.2, 115 mM KOAc, 8% w/v glycerol, and 0.2 mM each dNTPs. A GeneAmp® AccuRT Hot Start PCR enzyme was used at 0.15 U/ μ L. Starting with 5×10^5 copies of linearized Z05 D580G plasmid DNA per reaction volume of 50 μ L, reactions were denatured using a temperature of 94 °C for 60 seconds, then 30 cycles of amplification were performed, using a denaturation temperature of 94 °C for 15 seconds, an annealing temperature of 60 °C for 15 seconds, an extension temperature of 72 °C for 120 seconds, and followed by a final extension at a temperature of 72 °C for 5 minutes.

The resulting amplicon was purified with a QIAquick PCR Purification Kit (Qiagen, Inc., Valencia, CA, USA) and cut with Bln I and Bgl II, and then re-purified with a QIAquick PCR

Purification Kit. A Z05 D580G vector plasmid was prepared by cutting with the same two restriction enzymes and treating with alkaline phosphatase, recombinant (RAS, cat# 03359123001) and purified with a QIAquick PCR Purification Kit. The cut vector and the mutated insert were mixed at a 1:3 ratio and treated with T4 DNA ligase for 5 minutes at room temperature (NEB Quick Ligation™ Kit). The ligations were purified with a QIAquick PCR Purification Kit and transformed into an *E.coli* host strain by electroporation.

Aliquots of the expressed cultures were plated on ampicillin-selective medium in order to determine the number of unique transformants in each transformation. Transformations were stored at -70 °C to -80 °C in the presence of glycerol as a cryo-protectant.

Each library was then spread on large format ampicillin-selective agar plates. Individual colonies were transferred to 384-well plates containing 2X Luria broth with ampicillin and 10% w/v glycerol using an automated colony picker (QPix2, Genetix Ltd). These plates were incubated overnight at 30 °C to allow the cultures to grow and then stored at -70 °C to -80 °C. The glycerol added to the 2X Luria broth was low enough to permit culture growth and yet high enough to provide cryo-protection. Several thousand colonies at several mutagenesis (Mg^{2+}) levels were prepared in this way for later use.

Extract library preparation Part 1—Fermentation: From the clonal libraries described above, a corresponding library of partially purified extracts suitable for screening purposes was prepared. The first step of this process was to make small-scale expression cultures of each clone. These cultures were grown in 96-well format; therefore there were 4 expression culture plates for each 384-well library plate. 0.5 µL was transferred from each well of the clonal library plate to a well of a 96 well seed plate, containing 150 µL of Medium A (see Table 3 below). This seed plate was shaken overnight at 1150 rpm at 30 °C, in an iEMS plate incubator/shaker (ThermoElectron). These seed cultures were then used to inoculate the same medium, this time inoculating 20 µL into 250 µL Medium A in large format 96 well plates (Nunc # 267334). These plates were incubated overnight at 37 °C with shaking. The expression plasmid contained transcriptional control elements, which allow for expression at 37 °C but not at 30 °C. After overnight incubation, the cultures expressed the clone protein at typically 1-10% of total cell protein. The cells from these cultures were harvested by

centrifugation. These cells were either frozen (-20 °C) or processed immediately, as described below.

Table 2. Medium A (Filter-sterilized prior to use)

Component	Concentration
MgSO ₄ ·7H ₂ O	0.2 g/L
Citric acid·H ₂ O	2 g/L
K ₂ HPO ₄	10 g/L
NaNH ₄ PO ₄ ·4H ₂ O	3.5 g/L
MgSO ₄	2 mM
Casamino acids	2.5 g/L
Glucose	2 g/L
Thiamine·HCl	10 mg/L
Ampicillin	100 mg/L

- 5 ***Extract library preparation Part 2—Extraction:*** Cell pellets from the fermentation step were resuspended in 25 µL Lysis buffer (Table 3 below) and transferred to 384-well thermocycler plates and sealed. Note that the buffer contained lysozyme to assist in cell lysis, and DNase to remove DNA from the extract. To lyse the cells the plates were incubated at 37 °C for 15 minutes, frozen overnight at -20 °C, and incubated again at 37 °C for 15 minutes. Ammonium sulfate was added (1.5 µL of a 2 M solution) and the plates incubated at 75 °C for 15 minutes in order to precipitate and inactivate contaminating proteins, including the exogenously added nucleases. The plates were centrifuged at 3000 x g for 15 minutes at 4 °C and the supernatants transferred to a fresh 384-well thermocycler plate. These extract plates were frozen at -20 °C for later use in screens. Each well contained about 0.5-3 µM of the mutant library polymerase enzyme.
- 10
- 15

Table 3. Lysis Buffer

Component	Concentration or Percentage
Tris pH 7.5	50 mM
EDTA	1 mM
MgCl ₂	6 mM
Tween 20	0.5% v/v
Lysozyme (from powder)	1 mg/mL
DNase I	0.05 Units/ μ L

Screening extract libraries for reduced 3' primer mismatch extension rate: The extract library was screened by comparing the extension rate of a primer perfectly matched to an oligonucleotide template vs. the extension rate of a primer with a 3' G:T mismatch.

The enzyme extracts above were diluted 10-fold for primer extension reactions by combining 2.5 μ l extract with 22.5 μ l of a buffer containing 20 mM Tris-HCl, pH 8, 100 mM KCl, 0.1 mM EDTA, and 0.2% Tween-20 in a 384-well thermocycler plate, covering and heating for 10 minutes at 90 °C. Control reactions with perfect match primer combined 0.5 μ l of the diluted extract with 15 μ l master mix in 384-well PCR plates. Extension of the primed template was monitored every 10 seconds in a modified kinetic thermal cycler using a CCD camera (see, Watson, *supra*). Master mix contained 50 nM primed primer template, 25 mM Tricine, pH 8.3, 100 mM KOAc, 0.6X SYBR Green I, 200 μ M each dNTP, 100 nM Aptamer, and 2.5 mM Magnesium Acetate. In order to distinguish extension-derived fluorescence from background fluorescence, parallel wells were included in the experiment in which primer strand extension was prevented by leaving out the nucleotides from the reaction master mix. Reactions with the 3'-mismatched primer were performed as above except 1.5 μ l the diluted extract was added to each reaction and 1.5 mM Manganese Acetate was substituted for the Magnesium Acetate. Increasing the amount of extract three fold and using Manganese as the metal activator both make mismatch extension more likely and therefore improve the selectivity of the screen for those enzymes with the greatest ability to discriminate against 3'-mismatch extension.

Approximately 5000 mutant extracts were screened using the above protocol. Approximately 7% of the original pool was chosen for rescreening based on a perfect match primer extension

value above an arbitrary cutoff and low mismatch to perfect match extension ratio. Culture wells corresponding to the top extracts were sampled to fresh growth medium and re-grown to produce a new culture plates containing the best mutants, as well as a number of parental cultures to be used for comparison. These culture plates were then used to make fresh extracts

5 which were rescreened to confirm the original screen phenotype. The primer extension rates for the reactions with the perfect 3'-matched and the 3'-mismatched primers were calculated as the slope of the rise in fluorescence over time for the linear portion of the curve. The ratio of mismatched extension slope divided by the perfect matched extension slope was used to rank and select the best candidates. Selected clones from the rescreening, plus for comparison

10 the parental clone Z05 D580G, with their respective genotypes and phenotypes are included in the table below.

Table 4.

Enzyme	Perfect Match Slope	Mismatch Slope	MM Slope / PM Slope
Z05 D580G	8.29	8.04	0.97
Z05 D580G F497S	5.01	0.11	0.02

Various Amino Acid Substitutions at the Z05 F497 position: The effect of various

15 substitutions at the F497 position of Z05 DNA polymerase on mismatch discrimination in allele-specific PCR was examined. These substitutions were created in both Z05 DNA polymerase and Z05 D580G DNA polymerase by cloning synthetic gene fragments into vectors for one or both enzymes and the expressed mutant enzymes were purified and quantified. Z05 F497 mutants A (Alanine), G (Glycine), S (Serine), T (Threonine), and Y

20 (Tyrosine); and Z05 D580G mutants D (Aspartic Acid), and K (Lysine) were compared to their respective parental enzyme in an allele-specific PCR assay.

The control DNA polymerases of this example are a *Thermus sp.* Z05 DNA polymerase of SEQ ID NO:1 or a *Thermus sp.* Z05 DNA polymerase of SEQ ID NO:1 except that the amino acid at position 580 is Glycine (e.g., a D580G substitution) (hereinafter Z05 D580G

25 polymerase).

Primers were used that amplify a region of the human BRAF gene and are perfectly matched to the target when said target carries a mutation in codon 600 of BRAF, V600K. Against wild-

type BRAF target, present in human genomic DNA, the allele selective primer results in a single A:C mismatch at the 3' end. The common primer is perfectly matched to the BRAF gene, as is the probe sequence, which allows for real-time, TaqMan detection of amplification. Each reaction had 10,000 copies (33 ng) of wild-type Human Genomic cell line DNA, or
5 either 10,000 or 100 copies of a linearized plasmid containing the BRAF V600R mutant sequence in a final volume of 16 μ l. To allow for the different salt optima of the enzymes, amplifications were performed using a range of KCl concentrations from 25 to 130 mM. Buffer conditions were 50 mM Tris-HCl pH 8.0, 2.5 mM MgCl₂, 0.2 mM each dNTP, 0.02 U/ μ l UNG, and 200 nM Aptamer. Forward and Reverse primers were at 100 nM and the probe
10 was at 25 nM. All DNA polymerases were assayed at 20 nM and add 2 % (v/v) enzyme storage buffer (50% v/v glycerol, 100 mM KCl, 20 mM Tris pH 8.0, 0.1 mM EDTA, 1 mM DTT, 0.5% Tween 20) to the reactions. The reactions were performed in a Roche LightCycler 480 thermal cycler and denatured using a temperature of 95 °C for 60 seconds, then 99 cycles of amplification were performed, using a denaturation temperature of 92 °C for 10 seconds and
15 an annealing temperature of 62 °C for 30 seconds.

Reactions were in duplicate, crossing points ("Cps") were calculated by the Abs Quant/2nd derivative Max method and the Cps were averaged. PCR efficiency was calculated from the slope of the 100 and 10,000 copy perfect match plasmid reactions at the KCl concentration which resulted in the earliest 10,000 copy perfect match plasmid Cp. High Copy delta Cp is
20 equal to the difference between the Cp values of the reactions with 10,000 copy of 3'-mismatched wild-type genomic target and the Cp values of the reactions with 10,000 copy of perfect match plasmid target.

Table 5 below contains the averaged Cp values at the KCl concentration for each enzyme which resulted in the earliest high copy plasmid Cp and the calculated PCR efficiency and
25 high copy delta Cp. Z05 D580G F497S DNA polymerase which was indentified in the initial mutant screen as described above is included for reference. The data indicate that several amino acid substitutions at position F497 of Z05 DNA polymerase result in improved discrimination of primer mismatches in allele-selective PCR.

Table 5. Cps of Amplification of BRAF V600K mutant plasmid vs. Human genomic DNA

Enzyme	100 copies mutant plasmid	10,000 copies mutant plasmid	10,000 copies human gDNA	KCl Optimum	PCR Efficiency	High Copy delta Cp (gDNA- plasmid)
Z05	31.3	24.5	26.8	100	96.1	2.3
Z05 F497A	39.1	30.1	43.2	25	67.7	13.1
Z05 F497G	39.6	30.6	43.8	25	66.5	13.3
Z05 F497S	39.5	30.7	44.3	25	69.1	13.6
Z05 F497T	40.7	31.4	46.7	25	64.0	15.3
Z05 F497Y	31.0	24.3	26.6	100	98.1	2.4
Z05 D580G*	31.2	24.3	26.3	115	96.3	2.0
Z05 D580G F497D	37.0	28.9	40.8	25	76.9	12.0
Z05 D580G F497K	35.9	27.9	40.7	55	76.9	12.8
Z05 D580G F497S	34.3	26.8	34.7	55	84.1	7.9

* Average of 4 experiments

This example demonstrates that the F497A, F497G, F497S, F497T, F497Y, F497D, F497K mutant enzymes have improved rare allele detection relative to the control parental enzymes, Z05 and Z05 D580G.

Example 2: Identification of additional mutant polymerases with increased 3'-mismatch discrimination

This example demonstrates that various substitutions at the E493 position of Z05 DNA polymerase result in increased 3'-mismatch discrimination. These substitutions were created as described in Example 1. Z05 E493 mutants G (Glycine), K (Lysine), and R (Arginine); and Z05 D580G mutants A (Alanine), G (Glycine), K (Lysine), and R (Arginine) were compared to their respective parental enzyme in an allele-specific PCR assay.

Control enzymes and reaction conditions were as described above in Example 1. Table 6 shows that several amino acid substitutions at position E493 of Z05 DNA polymerase result in

improved discrimination of primer mismatches in allele-selective PCR when compared to the parental enzymes, Z05 and Z05 D580G.

Table 6. Cps of Amplification of BRAF V600K mutant plasmid vs. Human genomic DNA using E493 mutant Z05 enzymes

Enzyme	100 copies mutant plasmid	10,000 copies mutant plasmid	10,000 copies gDNA	KCl Optimum	PCR Efficiency	High Copy delta Cp (gDNA- plasmid)
Z05	31.3	24.5	26.8	100	96.1	2.3
Z05 E493G	31.1	24.5	27.7	85	100.8	3.2
Z05 E493K	33.4	26.3	30.3	100	90.9	4.0
Z05 E493R	31.7	24.7	29.2	85	94.1	4.5
Z05 D580G*	31.2	24.3	26.3	115	96.3	2.0
Z05 D580G E493A	31.0	24.1	26.2	115	93.3	2.1
Z05 D580G E493G	30.9	24.2	27.1	115	98.5	2.9
Z05 D580G E493K	31.6	24.9	31.5	100	99.1	6.6
Z05 D580G E493R	31.0	24.2	27.9	115	97.6	3.7

5 **Average of 4 experiments**

This example demonstrates that the E493A, E493G, E493K, and E493R mutant enzymes have improved rare allele detection relative to both control polymerases, Z05 and Z05 D580G.

Example 3: Identification of additional mutant polymerases with increased 3'-mismatch discrimination

10 This example shows that polymerases having a mutation at position S488 of a *Thermus sp.* Z05 DNA polymerase have increased 3'-mismatch discrimination.

The control DNA polymerases and reaction conditions were as described above in Example 1. Table 7 shows that several amino acid substitutions at position S488 of Z05 DNA polymerase result in improved discrimination of primer mismatches in allele-selective PCR.

Table 7. Cps of Amplification of BRAF V600K mutant plasmid vs. Human genomic DNA

Enzyme	100 copies mutant plasmid	10,000 copies mutant plasmid	10,000 copies human gDNA	KCl Optimum	PCR Efficiency	High Copy delta Cp (gDNA- plasmid)
Z05	31.3	24.5	26.8	100	96.1	2.3
Z05 S488C	34.8	27.1	36.8	40	81.6	9.7
Z05 S488F	34.3	26.8	35.2	40	84.6	8.4
Z05 S488G	32.5	25.4	30.4	70	90.1	5.0
Z05 S488T	33.4	26.2	34.1	55	89.7	8.0
Z05 S488Y	33.8	26.4	34.6	55	86.1	8.2
Z05 D580G*	31.2	24.3	26.3	115	96.3	2.0
Z05 D580G S488A	31.0	24.2	26.9	100	96.3	2.7
Z05 D580G S488D	32.1	25.0	31.6	70	92.4	6.5
Z05 D580G S488F	32.3	25.2	30.6	85	91.3	5.4
Z05 D580G S488G	31.1	24.5	28.2	100	99.5	3.7
Z05 D580G S488K	31.4	24.6	29.9	100	95.8	5.4

* Average of 4 experiments

This example demonstrates that the S488C, S448F, S488G, S488T, S488Y, S488A, S488D, and S488K mutant enzymes have improved rare allele detection relative to the parental control enzymes, Z05 and Z05 D580G.

It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art.

INFORMAL SEQUENCE LISTING

SEQ ID NO:1 *Thermus sp.* Z05 DNA polymerase (Z05)

5 MKAMLPLFEPKGRVLLVDGHHLAYRTFFALKGLTTSRGEVPQAVYGF AKSLLKALKEDGYKAV
 FVVFDAKAPSF RHEAYEAYKAGRAPT PEDFPRQLALIKELVDLLGFTRLEVPGFEADDVLATL
 AKKAEREGYEVRILTADRDLYQLVSDRVAVLHPEGHLITPEWLWEKYGLKPEQWVDFRALVGD
 PSDNLPGVKGIG EKTALKLLKEWGSLENILKNLDRVKPESVRERIKAHLEDLKLSLELSRVRS
 DLPLEVDFA RRREPDREGLRAFLE RLEFGSLLHEFG LLEAPAPLEEAPWPPPEGAFVGVLSR
 10 PEPMWAE LKALAA CKEGRVHRAKDPLAGLKDLKEVRG LLA KD LAVLALREGLDLAPSDDPMLL
 AYLLDPSNTTPEGVARRYGG EWTE DAAHRALLAERLQQNLLERLK GEEKLLWLYQEVEKPLSR
 VLAHMEATGVRLDVAYLKALSLELAEEIRRLEEEVFRLAGHPFNLSRDQLERVLFDELRLPA
 LGKTQKTGKRSTSAAVLEALREAHPIVEKILQHRELTKLKNTYVDPLPGLVHPRTGRLHTRFN
 QTATATGRLSSSDPNLQNIPIRTPLGQRI RRA FVAEAGWALVALDYSQIELRVLAHLSGDENL
 15 IRVFQEGKDIHTQTASWMFGVSPEAVDPLMRRAAKTVNFGVLYGMSAHRLSQELAI PYEEAVA
 FIERYFQSFPKVR AWIEKTLEEGRKRGYVETLFGRRRYVPDLNARVKS VREAAERMAFNMPVQ
 GTAADLMKLAMVKLFPHLREMGARMMLLQVHDELLLEAPQARAE EVAALAKEAMEKAYPLAVPL
 EVEVGIGEDWLSAKG

20 SEQ ID NO:2 *Thermus aquaticus* DNA polymerase (Taq)

MRGMLPLFEPKGRVLLVDGHHLAYRTFFALKGLTTSRGEVPQAVYGF AKSLLKALKEDGDAVI
 VVVFDAKAPSF RHEAYGGYKAGRAPT PEDFPRQLALIKELVDLLGLARLEVPGYEADDVLASLA
 KKAKEKEGYEVRILTADKDLYQLLS DRIHVLHPEGYLITPAWLWEKYGLRPDQWADYRALTGDE
 25 SDNLPGVKGIG EKTARKLLEEWGSLEALLKNLDRLKPAIREKILAHMDDLKLSWDLAKVRTDL
 PLEVDFAKRREPDRE RERLRAFLE RLEFGSLLHEFG LLESPKALEEAPWPPPEGAFVGVLSRKE
 PMWADLLALAAARGGRVHRAPEPYKALRDLKEARGLLAKDLSVLALREGLGLPPGDDPMLLAY
 LLDPSNTTPEGVARRYGG EWTEEAGERAA L SERLFANLWGRLEGEERLLWLYREVERPLSAVL
 AHMEATGVRLDVAYLRALSLEVAEEIARLEAEVFRLAGHPFNLSRDQLERVLFDELGLPAIG
 30 KTEKTGKRSTSAAVLEALREAHPIVEKILQYRELTKLKSTYIDPLPDLIHPRTGRLHTRFNQT
 ATATGRLSSSDPNLQNIPIVRTPLGQRI RRA FIAEEGWLLVALDYSQIELRVLAHLSGDENLIR
 VFQEGRDIHTETASWMFGVPREAVDPLMRRAAKTINFGVLYGMSAHRLSQELAI PYEEAQAFI
 ERYFQSFPKVR AWIEKTLEEGRRRGYVETLFGRRRYVPDLEARVKS VREAAERMAFNMPVQGT
 AADLMKLAMVKLFPRLEEMGARMMLLQVHDELVL EAPKERA EAVARLAKEVMEGVYPLAVPLEV
 35 EVGIGEDWLSAKE

SEQ ID NO:3 *Thermus filiformis* DNA polymerase (Tfi)

MLPLLEPKGRVLLVDGHHLAYRTFFALKGLTTSRGEVPQAVYGF AKSLLKALKEDGEVAIVVF
 40 DAKAPSF RHEAYEAYKAGRAPT PEDFPRQLALIKELVDLLGLVRLEVPGFEADDVLATLARKA
 EREGYEVRILSADRDLYQLLS DRIHLLHPEGEVLT PGWLQERYGLSPERWVEYRALVGDPSDN
 LPGVPGIG EKTALKLLKEWGSLEAILKNLDQVKPERVWEAIRNNLDKLQMSLELSRLRTDLPL
 EVDFAKRREPTGKGLKAFLERLEFGSLLHEFG LLEAPKEAEEAPWPPPGGAFLGFLLSRPEPM
 WAELLALAGAKEGRVHRAEDPVGALKDLKEIRG LLA KDLSVLALREGREIPPGDDPMLLAYLL
 45 DPGNTNPEGVARRYGG EWKEDAAARALLSERLWQALYPRVAEEERLLWLYREVERPLAQVLAH
 MEATGVRLDVPYLEALSQEVAFELERLEAEVHRLAGHPFNLSRDQLERVLFDELGLPPIGKT
 EKTGKRSTSAAVLELLREAHPIVGRILEYRELMKLKSTYIDPLPRLVHPKTGRLHTRFNQTAT

ATGRLSSSDPNLQNI PVRTPLGQIRKAFIAEEGHLLVALDYSQIELRVLAHLSG DENLIRVF
 REGKDIHTETA AWMFGVPPEGVDGAMRRAAKTVNFGVLYGMSAHRLSQELSI PYEEAAAFIER
 YFQSFPKVRAWIAKTLEEGRKKGYVETLFGRRRYVPDLNARVKS VREAAERMAFNMPVQGTAA
 DLMK LAMVKLF PRLRPLGVRILLQVHDEL VLEAPKARAE EAAQLAKETMEGVYPLSVPLEVEV
 5 GMGEDWLSAKE

SEQ ID NO:4 *Thermus flavus* DNA polymerase (Tfl)

MAMLPLFEPKGRVLLVDGHHLAYRTFFALKGLTTSRGEPVQAVYGF AKSLLKALKEDGDV VVV
 10 VFDAKAPSFRHEAYEAYKAGRAPT PEDFPRQLALIKELVDLLGLVRLEVPGFEADDVLATLAK
 RAEKEGYEVRILTADRDLYQLLSERIAILHPEGYLITPAWLYEKYGLRPEQWVDYRALAGDPS
 DNIPGVKGIGEKTAQRLIREWGSLENLFQHLDQVKPSLREKLQAGMEALALSRKLSQVHTDLP
 LEVDFGRRRTPNLEGLRAFLERLEFGSLLHEFGLLEGPKAAEEAPWPPPEGAF LGFSFSRPEP
 MWAELLALAGAWEGRLHRAQDPLRGLRDLKGVRGILAKDLAVLALREGLDLFPEDDPMLLAYL
 15 LDPSNTTPEGVARRYGGEWTE DAGERALLAERLFQTLKERLKGEE RLLWL YEEVEKPLSRVLA
 RMEATGVRLDVAYLQALSLEVEAEVRQLEEEVFRLAGHPFNLNSRDQLERVL FDELGLPAIGK
 TEKTGKRSTSAAVLEALREAHPIVDRI LQYRELTKLKNTYIDPLPALVHPKTGRLHTRFNQTA
 TATGRLSSSDPNLQNI PVRTPLGQIRRAFVAEEGWVLVLDYSQIELRVLAHLSG DENLIRV
 FQEGRDIHTQTASWMFGVSPEGVDPLMRRAAKTINFGVLYGMSAHRLSGELSI PYEEAVAFIE
 20 RYFQSYPKVRAWIEGTLEEGRRRGYVETLFGRRRYVPDLNARVKS VREAAERMAFNMPVQGTAA
 ADLMK LAMVRLFPRLQELGARMLLQVHDEL VLEAPKDRAERVAALAKEVMEGVWPLQVPLEVE
 VGLGEDWLSAKE

SEQ ID NO:5 *Thermus sp.* Sps17 DNA polymerase (Sps17)

MLPLFEPKGRVLLVDGHHLAYRTFFALKGLTTSRGEPVQAVYGF AKSLLKALKEDGEVAI VVF
 25 DAKAPSFRHEAYEAYKAGRAPT PEDFPRQLALIKELVDLLGLVRLEVPGFEADDVLATLAKKA
 EREGYEVRILSADRDLYQLLSDRIHLLHPEGEVLT PGWLQERYGLSPERWVEYRALVGDP SDN
 LPGVPGIGEK TALKLLKEWGSLEAILKNLDQVKPERVREAIRNNLDKLQMSLELSRLRTDLPL
 30 EVDFAKRREP DWGLKAFLE RLEFGSLLHEFGLLEAPKEAEEAPWPPPGGAFLGFLLSRPEPM
 WAELLALAGAKEGRVHRAEDPVGALKDLKEIRGLLAKDLSVLALREGREIPPGDDPMLLAYLL
 DPGNTNPEGVARRYGGEWKEDAAARALLSERLWQALYPRVAEEERLLWL YREVERPLAQVLAH
 MEATGVRLDVPYLEALSQEVAFELE RLEAEVHRLAGHPFNLNSRDQLERVL FDELGLPPIGKT
 EKTGKRSTSAAVLELLREAHPIVGRILEYRELMKLKSTYIDPLPRLVHPKTGRLHTRFNQTAT
 35 ATGRLSSSDPNLQNI PVRTPLGQIRKAFIAEEGHLLVALDYSQIELRVLAHLSG DENLIRVF
 REGKDIHTETA AWMFGVPPEGVDGAMRRAAKTVNFGVLYGMSAHRLSQELSI PYEEAAAFIER
 YFQSFPKVRAWIAKTLEEGRKKGYVETLFGRRRYVPDLNARVKS VREAAERMAFNMPVQGTAA
 DLMK LAMVKLF PRLRPLGVRILLQVHDEL VLEAPKARAE EAAQLAKETMEGVYPLSVPLEVEV
 GMGEDWLSAKA
 40

SEQ ID NO:6 *Thermus thermophilus* DNA polymerase (Tth)

MEAMLPLFEPKGRVLLVDGHHLAYRTFFALKGLTTSRGEPVQAVYGF AKSLLKALKEDGYKAV
 45 FVVVFDAKAPSFRHEAYEAYKAGRAPT PEDFPRQLALIKELVDLLGFTRLEVPGYEADDVLATL
 AKKAEKEGYEVRILTADRDLYQLVSDRVAVLHPEGHLITPEWLWEKYGLRPEQWVDFRALVGD
 PSDNLPGVKGIGEK TALKLLKEWGSLENLLKNLDRVKPEN VREKIKAHLEDLRLSLELSRVRT
 DLPLEVDLAQGREPDREGLRAFLERLEFGSLLHEFGLLEAPAPLEEAPWPPPEGAFVGVFLSR
 PEPMWAE LKALAA CRDGRVHRAADPLAGL KDLKEVRGLLAKDLAVLASREGLDLVP GDDPMLL

AYLLDPSNTTPEGVARRYGGEWTEDAHRALLSERLHRNLLKRLEGEKLLWLHYHEVEKPLSR
 VLAHMEATGVRRDVAYLQALSLELAEEIRRLEEEVFRLAGHPFNLNSRDQLERVLFDELRLPA
 LGKTQKTGKRSTSAAVLEALREAHPIVEKILQHRELTKLKNTYVDPLPSLVHPRTGRLHTRFN
 QTATATGRLSSSDPNLQNI PVRTPLGQIRRAFVAEAGWALVALDYSQIELRVLAHLSGDENL
 5 IRVFQEGKDIHTQTASWMFGVPPEAVDPLMRRAAKTVNFGVLYGMSAHRLSQELAI PYEEAVA
 FIERYFQSFPKVRAWIEKTLEEGRKRGYVETLFGRRRYVPDLNARVKSVREAAERMAFNMPVQ
 GTAADLMK LAMVKLF PRLREMGARM L LQVHDELLLEAPQARAE EVAALAKEAMEKAYPLAVPL
 EVEVGMGEDWLSAKG

10 **SEQ ID NO:7 *Thermus caldophilus* DNA polymerase (Tca)**

MEAMLPLFEPKGRVLLVDGHHLAYRTFFALKGLTTSRGEVPQAVYGF AKSLLKALKEDGYKAV
 FVVFDAKAPSFRHEAYEAYKAGRPTPEDFPRQLALIKELVDLLGFTRLEVPGYEADDVLATL
 AKNPEKEGYEVRILTADRDL DQLVSDRVAVLHPEGHLITPEWLWQKYGLKPEQWVDFRALVGD
 15 PSDNLPGVKGIGEK TALKLLKEWGSLENLLKNLDRVKPENVREK IKAHLEDLRLSLELSRVRT
 DLPLEVDLAQGREPDREGLRAFLERLEFGSLLHEFGLLEAPAPLEEAPWPPPEGAFVGFVLSR
 PEPMWAELKALAACRDGRVHRAADPLAGLKDLKEVRGLLAKDLAVLASREGLDLVP GDDPMLL
 AYLLDPSNTTPEGVARRYGGEWTEDAHRALLSERLHRNLLKRLQGEKLLWLHYHEVEKPLSR
 VLAHMEATGVRLDVAYLQALSLELAEEIRRLEEEVFRLAGHPFNLNSRDQLERVLFDELRLPA
 20 LGKTQKTGKRSTSAAVLEALREAHPIVEKILQHRELTKLKNTYVDPLPSLVHPNTGRLHTRFN
 QTATATGRLSSSDPNLQNI PVRTPLGQIRRAFVAEAGWALVALDYSQIELRVLAHLSGDENL
 IRVFQEGKDIHTQTASWMFGVPPEAVDPLMRRAAKTVNFGVLYGMSAHRLSQELAI PYEEAVA
 FIERYFQSFPKVRAWIEKTLEEGRKRGYVETLFGRRRYVPDLNARVKSVREAAERMAFNMPVQ
 GTAADLMK LAMVKLF PRLREMGARM L LQVHDELLLEAPQAGAE EVAALAKEAMEKAYPLAVPL
 25 EVEVGMGEDWLSAKG

SEQ ID NO:8

30 A-G-X1-X2-F-X3-X4-X5-S-X7-X8-Q-X9-X10-X11-X12-L-X13-X14-X15-L
 wherein X1 is H, E or Q; X2 is P, T or E; X3 is N or H; X4 is L
 or I; X5 is N or R; X7 is R, P, or S; X8 is D, K or T; X9 is L
 or V; X10 is E, S, A or G; X11 is R, N, Y, T or V; X12 is V or
 I; X13 is an amino acid other than F or Y; X14 is D or E; and
 X15 is E or K.

35

SEQ ID NO:9

40 A-G-X1-P-F-N-X4-N-S-X7-X8-Q-X9-X10-X11-X12-L-X13-X14-X15-L
 wherein X1 is H or E; X4 is L or I; X7 is R or P; X8 is D or K;
 X9 is L or V; X10 is E or S; X11 is R or N; X12 is V or I; X13
 is any amino acid other than F; X14 is D or E; and X15 is E or
 K.

SEQ ID NO:10

45

A-G-H-P-F-N-L-N-S-R-D-Q-L-E-R-V-L-X13-D-E-L
 wherein X13 is any amino acid other than F (SEQ ID NO: 10).

SEQ ID NO:11

A-G-H-P-F-N-L-N-S-R-D-Q-L-E-R-V-L-X13-D-E-L
X13 is A, G, S, T, Y, D, or K (SEQ ID NO: 11).

5

SEQ ID NO:12 Z05

EEEVFRLAGHPFNLNSRDQLERVLFDELRLPALGKT

10

SEQ ID NO:13 Taq

EAEVFRLAGHPFNLNSRDQLERVLFDELGLPAIGKT

SEQ ID NO:14 Tfi

15

EAEVHRLAGHPFNLNSRDQLERVLFDELGLPPIGKT

SEQ ID NO:15 Tfi

20

EEEVFRLAGHPFNLNSRDQLERVLFDELGLPAIGKT

SEQ ID NO:16 Sps17

EAEVHRLAGHPFNLNSRDQLERVLFDELGLPPIGKT

25

SEQ ID NO:17 Tth

EEEVFRLAGHPFNLNSRDQLERVLFDELRLPALGKT

30

SEQ ID NO:18 Tca

EEEVFRLAGHPFNLNSRDQLERVLFDELRLPALGKT

SEQ ID NO:19 Tma

35

AEEIYRIAGEPFNINSPKQVSRILFEKLGKIPRGKT

SEQ ID NO:20 Tne

40

AEKIYQIAGEPFNINSPKQVSNILFEKLGKIPRGKT

SEQ ID NO:21 Taf

KEKVFEIAGETFNLSSTQVAYILFEKLNIAFY-KK

45

SEQ ID NO:23 Dra

ESQIHEYAGEEFHIRSPKQLETVLYDKLELASSKKT

SEQ ID NO:24 Bst

ERRIYELAGQEFNINSPKQLGTVLFDKLQLPVLKKT

5

SEQ ID NO:25 Bca

EQRIYELAGQEFNINSPKQLGVILFEKLQLPVLKKS

10 **SEQ ID NO:26 native consensus motif**

A-G-X1-X2-F-X3-X4-X5-X6-X7-X8-Q-X9-X10-X11-X12-L-X13-X14-X15-L
wherein X1 is H, E or Q; X2 is P, T or E; X3 is N or H; X4 is L
or I; X5 is N or R; X6 is S; X7 is R, P, or S; X8 is D, K or T;
15 X9 is L or V; X10 is E, S, A or G; X11 is R, N, Y, T or V; X12
is V or I; X13 is F or Y; X14 is D or E; and X15 is E or K.

SEQ ID NO:27 modified Z05 D580 motif

20 T-G-R-L-S-S-X7-X8-P-N-L-Q-N

wherein X7 is Ser (S) or Thr (T); and X8 is any amino acid
other than Asp (D), or Glu (E)

SEQ ID NO:28 Conserved DNA polymerase active site

25

DYSQIELR

SEQ ID NO:29 CS5 DNA polymerase

30 MKAMLPLFEPKGRVLLVDGHHLAYRTFFALKGLTTSRGEPVQAVYGFASLLKALKEDGYKAV
FVVFDAKAPSFRRHEAYEAYKAGRAPTPEDFPRQLALIKELVDLLGFTRLEVPGFEADDVLATL
AKKAEREGYEVRIILTADRDLYQLVSDRVAVLHPEGHLITPEWLWEKYGLKPEQWVDFRALVGD
PSDNLPGVKGIGEKTKALKLLKEWGSLENILKNLDRVKPESVRERIKAHLEDLKLSELSRVRS
DLPLEVDFARRREPDREGLRAFLEFGLSLLHEFGLLEESEPVGYRIVKDLVEFEKLEIEKLR
35 ESPSFAIDLETSSLDPFDCDIVGISVSFKPKEAYYIPLHHRNAQNLDEKEVLKKLKEILEDPG
AKIVGQNLKFDYKVLMMVGVEPVPPYFDTMIAAYLLEPNEKKFNLDDLALKFLGYKMTSYQEL
MSFSFPLFGFSFADVPVEKAANYSCEDADITYRLYKTLCLKLHEADLENVYKIEEMPLVNVLA
RMELNGVYVDTEFLKKLSEYGGKLEELAEIYRIAGEPFNINSPKQVSRILFEKLGKIKPRGK
TTKTGDYSTRIEVLEELAGEHEIIPILILEYRKIQKLKSTYIDALPKMVNPKTGRIHASFNQTG
40 TATGRLSSSDPNLQNLPTKSEEGKEIRKAIVPQDPNWWIVSADYSQIELRILAHLSGDNLLR
AFEEGIDVHTLTASRIFNVKPEEVTEEMRRAGKMNFSIIYGVTPYGLSVRLGVPVKEAEKMI
VNYFVLYPKVRDYYIQRVVSEAKEKGYVRTLFGRKRDIPQLMARDRNTQAEGERIAINTPIQGT
AADI IKLAMIEIDRELKERKMRSKMIIQVHDELVEVPNEEKDALVELVKDRMTNVVKLSVPL
EVDVTIGKTWS

45

SEQ ID NO:30 CS6 DNA polymerase

MKAMLPLFEPKGRVLLVDGHHLAYRTFFALKGLTTSRGEVPQAVYGFASLLKALKEDGYKAV
 FVVFDAKAPSFRRHEAYEAYKAGRPTPEDFPRQLALIKELVDLLGFTRLEVPGFEADDVLATL
 5 AKKAEREGYEVRIILTADRDLYQLVSDRVAVLHPEGHLITPEWLWEKYGLKPEQWVDFRALVGD
 PSDNLPGVKGIGEKTAALKLLKEWGSLENILKNLDRVKPESVRERIKAHLEDLKLSELSRVRS
 DLPLEVDFARRREPDREGLRAFLERLEFGSLLHEFGLEESEPVGYRIVKDLVEFEKLEIEKLR
 ESPSFAIALATSSLDPFDCDIVGISVSFKPKEAYYIPLHHRNAQNLDKEVLKKLKEILEDPG
 AKIVGQNLKFDYKVLMMVKGVPEVPPYFDTMIAAYLLEPNEKKFNLDLALKFLGYKMTSYQEL
 10 MSFSFPLFGFSFADVPVEKAANYSCEDADITYRLYKTLCLKHEADLENVFIEMPLVNVLA
 RMELNGVYVDTEFLKKLSEYGGKLEELAEIYRIAGEPFNINSPKQVSRILFEKLGKPRGK
 TTKTGDYSTRIEVLEELAGEHEIIPLEIYRKIQKLKSTYIDALPKMVNPKTGRIHASFNQTG
 TATGRLSSSDPNLQNLPTKSEEGKEIRKAIVPQDPNWWIVSADYSQIELRILAHLSGDENLLR
 AFEEGIDVHTLTASRIFNVKPEEVTEEMRRAGKMNFSIIYGVTPYGLSVRLGVPVKEAEKMI
 15 VNYFVLYPKVRDYIQRVVSEAKEKGYVRTLFGRKRDIPQLMARDRNTQAEGERIAINTPIQGT
 AADI IKLAMI EIDRELKERKMRSMIIQVHDELVEVPNEEKDALVELVKDRMTNVVKLSVPL
 EVDVTIGKTWS

SEQ ID NO:31 Forward Primer

20 5' - CTACCTCCTGGACCCCTCCAA-3'

SEQ ID NO:32 Reverse Primer

5' - ATAACCAACTGGTAGTGGCGTGTA-3'

25 SEQ ID NO:33 Polymerase domain of Z05 D580G DNA polymerase

CTACCTCCTGGACCCCTCCAAACACCACCCCCGAGGGGGTGGCCCCGGCGCTACGGGGGGGAGTG
 GACGGAGGACGCCGCCACCGGGCCCTCCTCGCTGAGCGGCTCCAGCAAAACCTCTTGGAACG
 CCTCAAGGGAGAGGAAAAGCTCCTTTGGCTCTACCAAGAGGTGGAAAAGCCCCCTCTCCCGGGT
 30 CCTGGCCCACATGGAGGCCACCGGGGTAAGGCTGGACGTGGCCTATCTAAAGGCCCTTTCCCT
 GGAGCTTGCGGAGGAGATTCGCCGCCTCGAGGAGGAGGTCTTCCGCCTGGCGGGCCACCCCTT
 CAACCTGAACTCCCGTGACCAGCTAGAGCGGGTGCTCTTTGACGAGCTTAGGCTTCCCGCCCT
 GGGCAAGACGCAAAAGACGGGGAAGCGCTCCACCAGCGCCGCGGTGCTGGAGGCCCTCAGGGA
 GGCCACCCCATCGTGGAGAAGATCCTCCAGCACCGGGAGCTCACCAAGCTCAAGAACACCTA
 35 CGTAGACCCCTCCCGGGCCTCGTCCACCCGAGGACGGGCGCCTCCACACCCGCTTCAACCA
 GACAGCCACGGCCACGGGAAGGCTCTCTAGCTCCGGGGCCCAACCTGCAGAACATCCCCATCCG
 CACCCCTTGGGCCAGAGGATCCGCCGGGCCTTCGTGGCCGAGGCGGGATGGGCGTTGGTGGC
 CCTGGACTATAGCCAGATAGAGCTCCGGGTCTTCGCCACCTCTCCGGGGACGAGAACCTGAT
 CAGGGTCTTCCAGGAGGGGAAGGACATCCACACCCAGACCGCAAGCTGGATGTTCCGGCGTCTC
 40 CCCGGAGGCCGTGGACCCCTGATGCGCCGGGCGGCCAAGACGGTGAACCTTCGGCGTCCTCTA
 CGGCATGTCCGCCCATAGGCTCTCCAGGAGCTTGCCATCCCCTACGAGGAGGCGGTGGCCTT
 TATAGAGCGCTACTTCCAAAGCTTCCCCAAGGTGCGGGCCTGGATAGAAAAGACCCTGGAGGA
 GGGGAGGAAGCGGGGCTACGTGGAAACCCTCTTCGGAAGAAGGCGCTACGTGCCCGACCTCAA
 CGCCCGGGTGAAGAGCGTCAGGGAGGCCGCGGAGCGCATGGCCTTCAACATGCCCGTCCAGGG
 45 CACCGCCGCGACCTCATGAAGCTCGCCATGGTGAAGCTCTTCCCCACCTCCGGGAGATGGG
 GGCCCGCATGCTCCTCCAGGTCCACGACGAGCTCCTCCTGGAGGCCCCCAAGCGCGGGCCGA
 GGAGGTGGCGGCTTTGGCCAAGGAGGCCATGGAGAAGGCCTATCCCCTCGCCGTGCCCTGGA
 GGTGGAGGTGGGGATCGGGGAGGACTGGCTTTCCGCCAAGGGCTGATATCAGATCTCCCTGAT

TATGCGTCAGTCTATGAAGAAAAATCGTATACAGATGGACGAAGAGAGAATCCTTGTGAATTT
AACAGAGGGTATAGGGATTACACGCCACTACCAGTTGGTTAT

SEQ ID NO:34 - BRAF wild type sequence

5

AGTAAAAATAGGTGATTTTGGTCTAGCTACAGTGAAATCTCGATGGAGTGGGTCCCATCAGTT
TGAACAGTTGTCTGGATCCATTTTGTGGATGGTAAGAATTGAGGCTA

SEQ ID NO:35 - BRAF V600R mutant sequence

10

AGTAAAAATAGGTGATTTTGGTCTAGCTACAAGGAAATCTCGATGGAGTGGGTCCCATCAGTT
TGAACAGTTGTCTGGATCCATTTTGTGGATGGTAAGAATTGAGGCTA

SEQ ID NO:36 *Deinococcus radiodurans* DNA polymerase (Dra)

15

MADASPDPSKPDALVLIDGHALAFRSYFALPPLNNSKGEMTDAIVGFMKLLLRLARQKSNQVI
VVFDPVPKTLRHEQYEGYKSGRAQTPEDLRGQINRIRALVDALGFPRLEEPGYEADDVIASLT
RMAEGKGYEVRIVTSRDAYQLLDEHVKVIANDFSLIGPAQVEEKYGVTVRQWVDYRALTGDA
SDNIPGAKGIGPKTAAKLLQEYGTLEKVYEAHAGTLKPDGTRKKLLDSEENVKFSHDLSCMV
20 TDLPLDIEFGVRRLPDNPLVTEDLLTELELHSLRPMILGLNGPEQDGHAPDDLLEREHAQTPE
EDEAAALPAFSAPELAEWQTPAEGAVWGYVLSREDDLTAALLAAATFEDGVARPARVSEPDEW
AQAEAPENLFGELLPSDKPLTKKEQKALEKAQKDAEKARAKLREQFPATVDEAEFVGQRTVTA
AAAKALAAHLSVRGTVVEPGDDPLLYAYLLDPANTNMPVVAKRYLDREWPADAPTRAAITGHL
VRELPLLDARRKMYDEMEKPLSGVLGRMEVRGVQVDSDFLQTLISIAGVRLADLESQIHEY
25 AGEEFHIRSPKQLETVLYDKLELASSKKTKLTGQRSTAVSALEPLRDAHPIIPLVLEFRELDK
LRGTYLDPIPNLVNPHTGRLHTTFAQTAVATGRLSSLNPNLQNIPIRSELGREIRKGFIAEDG
FTLIAADYSQIELRLLAHIADDPLMQQAFVEGADIHRRTAQAQVLGLDEATVDANQORRAAKTVN
FGVLYGMSAHRLSNDLGI PYAEAATFIEIYFATYPGIRRYINHTLDFGRTHGYVETLYGRRRY
VPGLSSRN RVQREAEERLAYNMPIQGTAADIMKLAMVQLDPQLDAIGARMLLQVHDELLIEAP
30 LDKAEQVAALTKKVMENVVQLKVPLAVEVGTGPNWFDTK

SEQ ID NO:37 *Thermosipho africanus* DNA polymerase (Taf)

35

MGKMFLFDGTGLVYRAFYAIDQSLQTSSGLHTNAVYGLTKMLIKFLKEHISIGKDACVFLDS
KGGSKKRKDILETYKANRPSTPDLLLEQIPYVEELVDALGIKVLKIEGFEADDIIATLSKKFE
SDFEKNVNIITGDKDLLQLVSDKVFWVRVERGITDLVLYDRNKVIEKYGIYPEQFKDYLSLVGD
QIDNIPGVKGIGKKTAVSLLKKYNSLENVLKNINLLTEKLRLLED SKEDLQKSIELVELIYD
VPMDVEKDEIIYRGYNPDKLLKVLKKYEFSSIIKELNLQEKLEKEYILVDNEDKLKKLAEIE
KYKTFSIDTETTS LDPFEAKLVGISISTMEGKAYYIPVSHFGAKNISKSLIDKFLKQILQEKD
40 YNIVGQNLKFDYEIFKSMGFSPNVPHFDTMIAAYLLNPDEKRFNLEELSLKYLGYKMISFDEL
VNENVPLFGNDFS YVPLERAVEYSCEDADV TYRIFRKLGRKIYENEMEKL FYEIEEMPLIDVLS
EMELNGVYFDEEY LKELSKKYQEKMDGIKEKVFEIAGETFNLSSTQVAYILFEKLNIAPYKK
TATGKFSTNAEVLEELSKEHEIAKLLLEYRKYQKLKSTYIDS IPLSINRKTNRVHTTFHQGT
STGRLSSSNPNLQNL PTRSEEGKEIRKAVRPQRQDWILGADYSQIELRVLAHVSKDENLLKA
45 FKEDLDIHTITAAKIFGVSEMFVSEQMRRVGMVNF AIIYGVSPYGLSKRIGLSVSETKKIID
NYFRYYKGVFEY LKRMKDEARKKGYVTTLFGRRRYIPQLRSKNGNRVQEGERIAVNTPIQGT
ADIIKIAMINIHNR LKKNLRSKMILQVHDEL VFEVPDNELEIVKDLVRDEMENA VKLDVPLK
VDVYYGKEWE

SEQ ID NO:38 *Thermotoga maritima* DNA polymerase (Tma)

5 MARLFLFDGTALAYRAYYALDRSLSTSTGIPTNATYGVARMLVRFIKDHIIVGKDYVAVAFDK
 KAATFRHKLLETYKAQRPKTPDLLIQQLPYIKKLVEALGMKVLEVEGYEADDIIATLAVKGLP
 LFDEIFIVTGDKMDLQLVNEKIKVWRIVKGISDLELYDAQKVKEYGVEPQQIPDLLALTGDE
 IDNIPGVTGIGEKTAVQLLEKYKDLEDILNHVRELPOKVRKALLRDRENAILSKKLAILETNV
 PIEINWEELRYQGYDREKLLPLLKELEFASIMKELQLYEESEPVGYRIVKDLVEFEKLEKLR
 ESPSFAIDLETSSLDPFDCDIVGISVSFKPKEAYYIPLHHRNAQNLDKEVLKKLKEILEDPG
 10 AKIVGQNLKFDYKVLVMVGVEPVPPYFDTMIAAYLLEPNEKKFNLLDLALKFLGYKMTSYQEL
 MSFSFPLFGFSFADVPVEKAANYSCEDADITYRLYKTLCLKHEADLENVFIKIEMPLVNVLA
 RMELNGVYVDTEFLKKLSEYGYKKLEELAEIYRIAGEPFNINSPKQVSRILFEKLGKPRGK
 TTKTGDYSTRIEVLEELAGEHEIIPILILEYRKIQKLKSTYIDALPKMVNPKTGRIHASFNQTG
 TATGRLSSSDPNLQNLPTKSEEGKEIRKAIVPQDPNWWIVSADYSQIELRILAHLSGDENLLR
 15 AFEEGIDVHTLTASRIFNVKPEEVTEEMRRAGKMNFSIIYGVTPYGLSVRLGVPVKEAEKMI
 VNYFVLYPKVRDYIQRVVSEAKEKGYVRTLFGRKRDIPQLMARDRNTQAEGERIAINTPIQGT
 AADI IKLAMIEIDRELKERKMRSMIIQVHDELVEVPNEEKDALVELVKDRMTNVVKLSVPL
 EVDVTIGKTWS

20 SEQ ID NO:39 *Thermotoga neopolitana* DNA polymerase (Tne)

MARLFLFDGTALAYRAYYALDRSLSTSTGIPTNAVYGVARMVLVKFIKEHIIPEKDYA AVAFDK
 KAATFRHKLLVSDKAQRPKTPALLVQQLPYIKRLIEALGFKVLELEGYEADDIIATLAVRAAR
 FLMRFSLITGDKMDLQLVNEKIKVWRIVKGISDLELYDSKKVKERYGVEPHQIPDLLALTGDD
 25 IDNIPGVTGIGEKTAVQLLGKYRNLEYILEHARELPQVRKALLRDREVAILSKKLATLVNTA
 PVEVDWEEMKYRGYDKRKLPLILKELEFASIMKELQLYEEAEPTGYEIVKDHKTFEDLIEKLR
 EVPSFALDLETSSLDPFNCEIVGISVSFKPKTAYYIPLHHRNAHNLDLTLVLSKLKEILEDPS
 SKIVGQNLKYDYKVLVMVGISPVPYPHFDTMIAAYLLEPNEKKFNLEDLSLKLFLGYKMTSYQEL
 MSFSSPLFGFSFADVPVDKAAEYSCEDADITYRLYKILSMKLHEAELENVFIYRIEMPLVNVLA
 30 RMEFNWVYVDTEFLKKLSEYGYKKLEELAEKIYQIAGEPFNINSPKQVSNILFEKLGKPRGK
 TTKTGDYSTRIEVLEEIANEHEIVPLILEFRKILKLKSTYIDTLPKLVNPKTGRFHASFHQTG
 TATGRLSSSDPNLQNLPTKSEEGKEIRKAIVPQDPDWWIVSADYSQIELRILAHLSGDENLVK
 AFEEGIDVHTLTASRIYNVKPEEVNEEMRRVGKMNFSIIYGVTPYGLSVRLGIPVKEAEKMI
 ISYFTLYPKVRSYIQQVVAEAKEKGYVRTLFGRKRDIPQLMARDKNTQSEGERIAINTPIQGT
 35 AADI IKLAMIDIDEELRKRNMKSRMIIQVHDELVEVPDEEKEELVDLVKNKMTNVVKLSVPL
 EVDISIGKSW

SEQ ID NO:40 *Bacillus stearothermophilus* DNA polymerase (Bst)

40 MKNKLVLDGNSVAYRAFFALPLLHNDKGIHTNAVYGFMMNLNKILAEEQPTHILVAFDAGKT
 TFRHETFQDYKGGRRQQTPELSEQFPLLRELLKAYRIPAYELDHYEADDII GTMAARAEREGF
 AVKVISGDRDLTQLASPQVTVEITKKGITDIESYTPETVVEKYGLTPEQIVDLKGLMGDKSDN
 IPGVPGIGEKTAVKLLKQFGTVENVLASIDEIKGEKLKENLRQYRDLALLSKQLAAICRDAPV
 ELTLDDIVYKGEDREKVVALFQELGFQSFLDKMAVQTDEGEKPLAGMDFAIADSVTDEMLADK
 45 AALVVEVVDNYHHAPIVGIALANERGRFFLRPETALADPKFLAWLGDETCKKTMFDSKRAAV
 ALKWKGIELRGVVFDDLAAAYLLDPAQAAGDVAAVAKMHQYEAVRSDEAVYGKGAKRTVPDEP
 TLAEHLARKAAAIWALEEPLMDELRRNEQDRLLTELEQPLAGILANMEFTGVKVDTKRLEQMG
 AELTEQLQAVERRIYELAGQEFNINSPKQLGTVLFDKLQLPVLKKTKTGYSTSADVLEKLAPH

HEIVEHILHYRQLGKLQSTYIEGLLKVVHPVTGKVHTMFNQALTQTGRLSSVEPNLQNIPIRL
 EEGRKIRQAFVPSEPDWLIFAADYSQIELRVLAHIAEDDNLIEAFRRGLDIHTKTAMDIFHVS
 EEDVTANMRRQAKAVNFGIVYGISDYGLAQNLNITRKEAAEFIERYFASFPGVKQYMDNIVQE
 AKQKGYVTLLHRRRYLPDITSRNFNVRSAERTAMNTPIQGSAAADIKKAMIDLSVRLREER
 5 LQARLLLQVHDELILEAPKEEIERLCRLVPEVMEQAVALRVPLKVDYHYGPTWYDAK

SEQ ID NO:41 *Bacillus caldotenax* DNA polymerase (Bca)

MKKKLVLIDGSSVAYRAFFALPLLHNDKGIHTNAVYGFTMMLNKILAEETPHMLVAFDAGKT
 10 TFRHEAFQEYKGGRRQQTPELSEQFPLLRELLRAYRIPAYELENYEADDIIGTLAARAEQEGF
 EVKVISGDRDLTQLASPHVTVDITKKGITDIEPYTPEAVREKYGLTPEQIVDLKGLMGDKSDN
 IPGVPGIGEKTAVKLLRQFGTVENVLASIDEIKGEKLETLRQHREMAALLSKKLAAIRRDAPV
 ELSLDDIAYQGEDREKVVALFKELGFQSFLEKMESPSSEEEKPLAKMAFTLADRVTEEMLADK
 AALVVEVVEENYHDAPIVGIADVNEHGRFFLRPETALADPQFVAWLGDETKKKSMFDSKRAAV
 15 ALKWKGIELCGVSFDLLLAAYLLDPAQGVDDVAAAAMKQYEAVRPDEAVYGGKAKRAVPDEP
 VLAEHLVRKAAAIWALERPFDELRRNEQDRLLVELEQPLSSILAEMEFAGVKVDTKRLEQMG
 EELAEQLRTVEQRIYELAGQEFNINSPKQLGVILFEKLQLPVLKKSKTGYSTSADVLEKLAPY
 HEIVENILQHYRQLGKLQSTYIEGLLKVVPRPDTKKVHTIFNQALTQTGRLSSTEPNLQNIPIR
 LEEGRKIRQAFVPSESDWLIFAADYSQIELRVLAHIAEDDNLMEAFRRDLDIHTKTAMDIFQV
 20 SEDEVTPNMRRQAKAVNFGIVYGISDYGLAQNLNISRKEAAEFIERYFESFPGVKRYMENIVQ
 EAKQKGYVTLLHRRRYLPDITSRNFNVRSAERMAMNTPIQGSAAADIKKAMIDLNARLKEE
 RLQARLLLQVHDELILEAPKEEMERLCRLVPEVMEQAVTLRVPLKVDYHYGSTWYDAK

SEQ ID NO:42

25 A-G-X1-X2-F-X3-X4-X5-X6-X7-X8-Q-X9-X10-X11-X12-L-X13-X14-X15-L
 wherein X1 is H, E or Q; X2 is P, T or E; X3 is N or H; X4 is L
 or I; X5 is N or R; X6 is any amino acid; X7 is R, P, or S; X8
 is D, K or T; X9 is L or V; X10 is any amino acid; X11 is R, N,
 30 Y, T or V; X12 is V or I; X13 is any amino acid other than F or
 Y; X14 is D or E; and X15 is E or K.

SEQ ID NO:43

35 A-G-X1-P-F-N-X4-N-X6-X7-X8-Q-X9-X10-X11-X12-L-X13-X14-X15-L
 wherein X1 is H or E; X4 is L or I; X6 is any amino acid; X7 is
 R or P; X8 is D or K; X9 is L or V; X10 is any amino acid; X11
 is R or N; X12 is V or I; X13 is any amino acid other than F;
 X14 is D or E; and X15 is E or K.

40

SEQ ID NO:44

A-G-X1-P-F-N-X4-N-S-X7-X8-Q-X9-X10-R-X12-L-X13-X14-X15-L
 wherein X1 is H or E; X4 is L or I; X7 is R or P; X8 is D or K;
 45 X9 is L or V; X10 is E or S; X12 is V or I; X13 is any amino
 acid other than F; X14 is D or E; and X15 is E or K.

SEQ ID NO:45

A-G-H-P-F-N-L-N-X6-R-D-Q-L-X10-R-V-L-X13-D-E-L

5 wherein X6 is any amino acid; X10 is any amino acid; and X13 is any amino acid other than F.

SEQ ID NO:46

A-G-H-P-F-N-L-N-X6-R-D-Q-L-X10-R-V-L-X13-D-E-L

10 wherein X6 is any amino acid other than S; X10 is any amino acid other than E; and X13 is any amino acid other than F.

SEQ ID NO:47

15 A-G-H-P-F-N-L-N-X6-R-D-Q-L-X10-R-V-L-X13-D-E-L

wherein X6 is S, G, A, D, F, K, C, T or Y; X10 is E, S, Q, A, G, K, or R; and X13 is A, G, S, T, Y, D or K.

SEQ ID NO:48

20 A-G-H-P-F-N-L-N-S-R-D-Q-L-E-R-V-L-X13-D-E-L

wherein X13 is A, G, S, T, D, or K.

SEQ ID NO:49

25 A-G-X₁-X₂-F-X₃-X₄-X₅-S-X₇-X₈-Q-X₉-X₁₀-X₁₁-X₁₂-L-X₁₃-X₁₄-X₁₅-L, wherein X₁ is H, E or Q; X₂ is P, T or E; X₃ is N or H; X₄ is L or I; X₅ is N or R; X₇ is R, P or S; X₈ is D, K or T; X₉ is L or V; X₁₀ is E, S, A or G; X₁₁ is R, N, Y, T or V; X₁₂ is V or I; X₁₃ is G, A, L, M, W, P, S, T, R, K, C, N, Q, D, E, V, I or H; and X₁₄ is D

30 or E; and X₁₅ is E or K.

WHAT IS CLAIMED IS:

1. A DNA polymerase having increased 3'-mismatch discrimination activity compared with a control DNA polymerase, wherein the amino acid of the DNA polymerase corresponding to position 497 of SEQ ID NO: 1 is any amino acid other than F or Y, and wherein the control DNA polymerase has the same amino acid sequence as the DNA polymerase except that the amino acid of the control DNA polymerase corresponding to position 497 of SEQ ID NO:1 is F or Y.
2. The DNA polymerase of claim 1, comprising a motif in the polymerase domain comprising
A-G-X₁-X₂-F-X₃-X₄-X₅-S-X₇-X₈-Q-X₉-X₁₀-X₁₁-X₁₂-L-X₁₃-X₁₄-X₁₅-L, wherein:
X₁ is H, E or Q;
X₂ is P, T or E;
X₃ is N or H;
X₄ is L or I;
X₅ is N or R;
X₇ is R, P, or S;
X₈ is D, K or T;
X₉ is L or V;
X₁₀ is E, S, A, or G;
X₁₁ is R, N, Y, T or V;
X₁₂ is V or I;
X₁₃ is any amino acid other than F or Y;
X₁₄ is D or E; and
X₁₅ is E or K (SEQ ID NO:8).
3. The DNA polymerase of claim 1, comprising a motif in the polymerase domain comprising
A-G-X₁-P-F-N-X₄-N-S-X₇-X₈-Q-X₉-X₁₀-X₁₁-X₁₂-L-X₁₃-X₁₄-X₁₅-L, wherein:
X₁ is H or E; and

X₄ is L or I;

X₇ is R or P;

X₈ is D or K;

X₉ is L or V;

5 X₁₀ is E or S;

X₁₁ is R or N;

X₁₂ is V or I;

X₁₃ is any amino acid other than F or Y;

X₁₄ is D or E; and

10 X₁₅ is E or K (SEQ ID NO:9).

4. The DNA polymerase of claim 1, comprising a motif in the polymerase domain comprising

A-G-H-P-F-N-L-N-S-R-D-Q-L-E-R-V-L-X₁₃-D-E-L, wherein:

X₁₃ is an amino acid other than F (SEQ ID NO:10).

- 15 5. The DNA polymerase of claim 4, wherein X₁₃ is A, G, S, T, Y, D, or K (SEQ ID NO:11).

6. The DNA polymerase of any one of claims 1 to 5, wherein the amino acid corresponding to position 580 of SEQ ID NO:1 is any amino acid other than D or E.

- 20 7. The DNA polymerase of any one of claims 1 to 6, wherein the amino acid corresponding to position 580 of SEQ ID NO:1 is selected from the group consisting of L, G, T, Q, A, S, N, R, and K.

8. The DNA polymerase of any one of claims 1 to 7, wherein the DNA polymerase comprises an amino acid sequence at least 80%, preferably at least 90%, more

preferably at least 95% sequence identical to SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 36, 37, 38, 39, 40 or 41.

- 5 9. The DNA polymerase of any one of claims 6 or 7, wherein the polymerase has at least 80%, preferably at least 90%, more preferably at least 95% sequence identity to SEQ ID NO:1.
- 10 10. The DNA polymerase of claim 9, wherein the DNA polymerase is a Z05 DNA polymerase, and the amino acid at position 580 is selected from the group consisting of L, G, T, Q, A, S, N, R, and K.
- 10 11. A recombinant nucleic acid encoding the DNA polymerase according to any one of claims 1 to 10.
12. A method for conducting primer extension, comprising:
contacting a DNA polymerase according to any one of claims 1 to 10 with a primer, a polynucleotide template, and nucleoside triphosphates under conditions suitable for extension of the primer, thereby producing an extended primer.
- 15 13. A kit for producing an extended primer, comprising:
at least one container providing a DNA polymerase according to any one of claims 1 to 10.
- 20 14. The kit according to claim 13 further comprising one or more additional containers selected from the group consisting of:
(a) a container providing a primer hybridizable, under primer extension conditions, to a predetermined polynucleotide template;
(b) a container providing nucleoside triphosphates; and
(c) a container providing a buffer suitable for primer extension.
15. A reaction mixture comprising a DNA polymerase according to any one of claims 1 to 10, at least one primer, a polynucleotide template, and nucleoside triphosphates.

Figure 1

Z05	EEVFRLAGH P FN L N S R D Q L E R V LF D E LRLPALGKT (SEQ ID NO:12)
Taq	EAEVFRLAGH P FN L N S R D Q L E R V LF D E LGLPAIGKT (SEQ ID NO:13)
Tfi	EAEVHRLAGH P FN L N S R D Q L E R V LF D E LGLPPIGKT (SEQ ID NO:14)
Tfl	EEVFRLAGH P FN L N S R D Q L E R V LF D E LGLPAIGKT (SEQ ID NO:15)
Sps17	EAEVHRLAGH P FN L N S R D Q L E R V LF D E LGLPPIGKT (SEQ ID NO:16)
Tth	EEVFRLAGH P FN L N S R D Q L E R V LF D E LRLPALGKT (SEQ ID NO:17)
Tca	EEVFRLAGH P FN L N S R D Q L E R V LF D E LRLPALGKT (SEQ ID NO:18)
Tma	AEEIYRIAGE P FN I N S P K Q V S R I LF E K LGIKPRGKT (SEQ ID NO:19)
Tne	AEKIQIAGE P FN I N S P K Q V S N I LF E K LGIKPRGKT (SEQ ID NO:20)
Taf	KEKVFEIAGE T FN L N S S T Q V A Y I LF E K LNIAPY-KK (SEQ ID NO:21)
Dra	ESQIHHEYAGE E FH I R S P K Q L E T V LY D K LELASSKKT (SEQ ID NO:23)
Bst	ERRIVELAQE E FN I N S P K Q L G T V LF D K LQLPVLKKT (SEQ ID NO:24)
Bca	EQRIVELAQE E FN I N S P K Q L G V I LF E K LQLPVLKKS (SEQ ID NO:25)
	-----AGX ₁ X ₂ FX ₃ X ₄ X ₅ X ₆ X ₇ X ₈ QX ₉ X ₁₀ X ₁₁ X ₁₂ LX ₁₃ X ₁₄ X ₁₅ L----- (SEQ ID NO:26)

FIGURE 2

A. Sequence identities over the entire polymerase I enzyme (corresponding to amino acids 1-834 of Z05)													
Name	Z05	Taq	Tfi	Tfl	Sps17	Tth	Tca	Dra	Tma	Tne	Taf	Bst	Bca
Z05		0.864	0.833	0.859	0.839	0.962	0.958	0.459	0.374	0.368	0.359	0.407	0.408
Taq	0.864		0.831	0.854	0.836	0.872	0.864	0.468	0.382	0.368	0.351	0.397	0.397
Tfi	0.833	0.831		0.82	0.991	0.829	0.824	0.45	0.371	0.375	0.353	0.405	0.397
Tfl	0.859	0.854	0.82		0.824	0.853	0.848	0.462	0.381	0.374	0.356	0.397	0.398
Sps17	0.839	0.836	0.991	0.824		0.835	0.83	0.452	0.375	0.377	0.355	0.407	0.399
Tth	0.962	0.872	0.829	0.853	0.835		0.989	0.463	0.373	0.367	0.358	0.406	0.406
Tca	0.958	0.864	0.824	0.848	0.83	0.989		0.46	0.371	0.365	0.356	0.404	0.404
Dra	0.459	0.468	0.45	0.462	0.452	0.463	0.46		0.334	0.325	0.314	0.338	0.339
Tma	0.374	0.382	0.371	0.381	0.375	0.373	0.371	0.334		0.854	0.567	0.37	0.377
Tne	0.368	0.368	0.375	0.374	0.377	0.367	0.365	0.325	0.854		0.558	0.377	0.376
Taf	0.359	0.351	0.353	0.356	0.355	0.358	0.356	0.314	0.567	0.558		0.356	0.364
Bst	0.407	0.397	0.405	0.397	0.407	0.406	0.404	0.338	0.37	0.377	0.356		0.881
Bca	0.408	0.397	0.397	0.398	0.399	0.406	0.404	0.339	0.377	0.376	0.364	0.881	
B. Sequence identities over polymerase sub domain only (corresponding to amino acids 420-834 of Z05)													
Name	Z05	Taq	Tfi	Tfl	Sps17	Tth	Tca	Dra	Tma	Tne	Taf	Bst	Bca
Z05		0.901	0.845	0.891	0.845	0.975	0.973	0.563	0.483	0.478	0.44	0.498	0.49
Taq	0.901		0.879	0.901	0.877	0.906	0.901	0.561	0.488	0.473	0.44	0.503	0.495
Tfi	0.845	0.879		0.857	0.997	0.853	0.853	0.566	0.495	0.49	0.449	0.512	0.49
Tfl	0.891	0.901	0.857		0.855	0.889	0.889	0.571	0.492	0.48	0.444	0.494	0.485
Sps17	0.845	0.877	0.997	0.855		0.853	0.853	0.566	0.495	0.49	0.449	0.512	0.49
Tth	0.975	0.906	0.853	0.889	0.853		0.99	0.563	0.478	0.473	0.437	0.496	0.488
Tca	0.973	0.901	0.853	0.889	0.853	0.99		0.563	0.478	0.473	0.437	0.496	0.488
Dra	0.563	0.561	0.566	0.571	0.566	0.563	0.563		0.45	0.448	0.426	0.474	0.454
Tma	0.483	0.488	0.495	0.492	0.495	0.478	0.478	0.45		0.883	0.622	0.474	0.475
Tne	0.478	0.473	0.49	0.48	0.49	0.473	0.473	0.448	0.883		0.615	0.476	0.473
Taf	0.44	0.44	0.449	0.444	0.449	0.437	0.437	0.426	0.622	0.615		0.46	0.473
Bst	0.498	0.503	0.512	0.494	0.512	0.496	0.496	0.474	0.474	0.476	0.46		0.898
Bca	0.49	0.495	0.49	0.485	0.49	0.488	0.488	0.454	0.475	0.473	0.473	0.898	

Figure 1

	*
Z05	EEEVFRL AGH P FN L N S R D QL E R V LF D E LRLPALGKT (SEQ ID NO:12)
Taq	EAEVFRL AGH P FN L N S R D QL E R V LF D E LGLPAIGKT (SEQ ID NO:13)
Tfi	EAEVHRL AGH P FN L N S R D QL E R V LF D E LGLPPIGKT (SEQ ID NO:14)
Tfl	EEEVFRL AGH P FN L N S R D QL E R V LF D E LGLPAIGKT (SEQ ID NO:15)
Sps17	EAEVHRL AGH P FN L N S R D QL E R V LF D E LGLPPIGKT (SEQ ID NO:16)
Tth	EEEVFRL AGH P FN L N S R D QL E R V LF D E LRLPALGKT (SEQ ID NO:17)
Tca	EEEVFRL AGH P FN L N S R D QL E R V LF D E LRLPALGKT (SEQ ID NO:18)
Tma	AEEIYRI AGE P FN I N S P K QV S R I LF E K LGIKPRGKT (SEQ ID NO:19)
Tne	AEKIYQI AGE P FN I N S P K QV S N I LF E K LGIKPRGKT (SEQ ID NO:20)
Taf	KEKVFEI AGE T FN L N S S T QV A Y I LF E K LNIAPY-KK (SEQ ID NO:21)
Dra	ESQIHEY AGE E FH I R S P K QL E T V LY D K LELASSKKT (SEQ ID NO:23)
Bst	ERRIYE LAGQ E FN I N S P K QL G T V LF D K LQLPVLKKT (SEQ ID NO:24)
Bca	EQRIYE LAGQ E FN I N S P K QL G V I LF E K LQLPVLKKKS (SEQ ID NO:25)
	----- AGX₁X₂FX₃X₄X₅X₆X₇X₈QX₉X₁₀X₁₁X₁₂LX₁₃X₁₄X₁₅L ----- (SEQ ID NO:26)