



(86) Date de dépôt PCT/PCT Filing Date: 2006/09/29

(87) Date publication PCT/PCT Publication Date: 2007/04/12

(45) Date de délivrance/Issue Date: 2014/05/27

(85) Entrée phase nationale/National Entry: 2008/03/12

(86) N° demande PCT/PCT Application No.: US 2006/038174

(87) N° publication PCT/PCT Publication No.: 2007/041358

(30) Priorités/Priorities: 2005/09/30 (US60/722,217);
2005/10/31 (US60/732,340); 2005/11/04 (US60/733,868)

(51) Cl.Int./Int.Cl. *C07D 403/12* (2006.01),
A61K 31/506 (2006.01), *A61P 35/00* (2006.01),
C07D 403/14 (2006.01), *C07D 405/14* (2006.01)


(72) Inventeurs/Inventors:
XIAO, XIAO-YI, US;
PATEL, DINISH V., US;
WARD, JOHN S., US;
BRAY, MARK R., CA;
AGOSTON, GREGORY E., US;
TRESTON, ANTHONY M., US

(73) Propriétaire/Owner:
MIKANA THERAPEUTICS, INC., US

(74) Agent: FINLAYSON & SINGLEHURST

(54) Titre : COMPOSES PYRAZOLE SUBSTITUES

(54) Title: SUBSTITUTED PYRAZOLE COMPOUNDS




(57) Abrégé/Abstract:

Disclosed are protein kinase inhibitors, compositions comprising such inhibitors, and methods of use thereof. More particularly, disclosed are inhibitors of Aurora A (Aurora-2) protein kinase. Wherein the compound of that which is disclosed has the formula of: (see above formula) Also disclosed are methods of treating diseases associated with protein kinases, especially diseases associated with Aurora-2, such as cancer.



ABSTRACT

Disclosed are protein kinase inhibitors, compositions comprising such inhibitors, and methods of use thereof. More particularly, disclosed are inhibitors of Aurora A (Aurora-2) protein kinase. Wherein the compound of that which is disclosed has the formula of:



Also disclosed are methods of treating diseases associated with protein kinases, especially diseases associated with Aurora-2, such as cancer.

5

SUBSTITUTED PYRAZOLE COMPOUNDS

10

15 FIELD OF THE INVENTION

This invention is directed to protein kinase inhibitors, compositions comprising such inhibitors, and methods of use thereof. More particularly, the invention relates to inhibitors of Aurora A (Aurora-2) protein kinase. The invention also relates to pharmaceutical compositions, as well as to methods of

20 treating diseases associated with protein kinases, especially diseases associated with Aurora A, such as cancer.

BACKGROUND OF THE INVENTION

The search for new therapeutic agents has been greatly aided in recent

25 years by better understanding of the structure of enzymes and other biomolecules associated with target diseases. One important class of enzymes that has been the subject of extensive study is the protein kinases.

Protein kinases mediate intracellular signal transduction by effecting a phosphoryl transfer from a nucleoside triphosphate to a protein acceptor that is

30 involved in a signaling pathway. There are a number of kinases and pathways through which extracellular and other stimuli cause a variety of cellular responses to occur inside the cell. Examples of such stimuli include environmental and chemical stress signals (e.g. osmotic shock, heat shock, ultraviolet radiation, bacterial endotoxin, H₂O₂), cytokines (e.g. interleukin-1 (IL-1) and tumor necrosis

factor alpha (TNF-alpha)), and growth factors (e.g. granulocyte macrophage-colony-stimulating factor (GM-CSF), and fibroblast growth factor (FGF). An extracellular stimulus may effect one or more cellular responses related to cell growth, migration, differentiation, secretion of hormones, activation of transcription factors, muscle contraction, glucose metabolism, control of protein synthesis and regulation of cell cycle.

Many diseases are associated with abnormal cellular responses triggered by protein kinase-mediated events. These diseases include autoimmune diseases, inflammatory diseases, neurological and neurodegenerative diseases, cancer, cardiovascular diseases, allergies and asthma, Alzheimer's disease or hormone-related diseases. Accordingly, there has been a substantial effort in medicinal chemistry to find protein kinase inhibitors that are effective as therapeutic agents.

In humans, there are three highly related Aurora kinases that are all serine/threonine protein kinases (see Andrews, P. D., et al., *Curr. Opin. Cell. Biol.* **2003**, *15*, 672-683; Carmena, M., Earnshaw, W. C., *Nat. Rev. Mol. Cell. Biol.* **2003**, *4*, 842-854; Brown, J. R., et al., *BMC Evol. Biol.* **2004**, *4*, 39, Andrews, P. D., *Oncogene* **2005**, *24*, 5005-5015). Despite the sequence relatedness of Aurora A, B and C, the localization and function of these kinases is quite distinct. As a result, overexpression or activation of each of these kinases can be associated with different disease states, including proliferative diseases such as cancer.

Members of the family demonstrate distinct subcellular localization during mitosis and are degraded by the proteosome following exit from mitosis (Graham *et al.* (2002) *J. Biol. Chem.* 277:42419-22). The kinases are often found complexed with other proteins, including cytoskeletal structures.

The Aurora kinases share a conserved C-terminal catalytic domain, with greater variation being observed at the N-terminus. Aurora A (Aurora-2) is unique in the presence of two lysine residues in the nucleotide-binding domain of the kinase (Warner *et al.* (2003) *Molecular Cancer Therapeutics* 2:589-95).

Maximum levels of the Aurora A polypeptide, and maximum Aurora A activity, are observed at the G₂/M transition leading into mitotic prophase, with the polypeptide localizing to the mitotic spindle pole (Graham *et al.* (2002) *J. Biol. Chem.* 277:42419-22; Sakai *et al.* (2002) *J. Biol. Chem.* 277:48714-23). Aurora A appears to regulate chromosome duplication with aberrant expression being associated with aneuploidy and an aggressive clinical phenotype,

particularly in solid tumors. Such observations, and additional experimental data, suggest that Aurora A disrupts the signaling cascade that regulates chromosome segregation (Sen *et al.* (2002) *J. Nat. Cancer. Inst.* 94:1320-29).

5 Aurora A also appears to function in meiosis, likely in separating homologous chromosomes and in spindle rotation. Injection of antibodies against Aurora A into *Xenopus* oocytes prevents first polar body extrusion and causes arrest at meiosis I (Castro *et al.* (2003) *J. Biol. Chem.* 2236-41). The *Xenopus* kinesin-like protein, Eg5, is known to be a substrate for Aurora-2 (Castro *et al.* (2003) *J. Biol. Chem.* 2236-41).

10 In addition, *in vitro* studies show that Aurora A is incorporated into chromatin and phosphorylates histone H3, and possibly histone H2B (Scrittore *et al.* (2001) *J. Biol. Chem.* 276:30002-10). H3 phosphorylation, *e.g.*, at serine-10, during chromosome assembly, appears to be a conserved event in eukaryotic cell division. Inhibition of H3 phosphorylation leads to chromosome condensation,
15 abnormal segregation, and the loss of chromosomes during mitosis and meiosis (Scrittore *et al.* (2001) *J. Biol. Chem.* 276:30002-10).

Accordingly, the emerging model for histone phosphorylation is analogous to that of histone acetylation, wherein partially redundant enzymatic activities are associated with histone modifications but different enzymes may function in
20 different cellular contexts. For example, some enzymes may modify histones in bulk, while other enzymes modify histones in a targeted manner, *i.e.*, in a sequence or domain-specific manner in the context of assembled chromatin (see, *e.g.*, Scrittore *et al.* (2001) *J. Biol. Chem.* 276:30002-10). According to this model, Aurora A would appear to be a kinase responsible for targeted histone
25 modification, in the context of assembled or assembling chromatin.

Other members of the Aurora kinase family are also associated with mitosis and meiosis. Aurora B, like Aurora A, is involved in distinct protein phosphorylation events that regulate the cell cycle. Unlike Aurora A, Aurora B is localized to inner-centromeric chromatin from prophase until the metaphase-anaphase transition, relocalizes to the microtubules in the spindle midzone during
30 telophase, and subsequently is found in the midbody throughout cytokinesis (See Andrews, P. D., *Oncogene* 2005, 24, 5005-5015, *loc. cit.*). The function of Aurora B is to ensure accurate chromosome segregation and appropriate cytokinesis. Aurora B appears to associate with a survivin, a polypeptide that

associates with the inner centromere and undergoes a significant degree of stretching during mitosis. Survivin appears to be involved with inhibition of apoptosis as well as cell cycle control. Interestingly, both Aurora B and survivin are delocalized during megakaryocyte endomitosis, a process by which late
5 anaphase and cytokinesis are skipped, leading to megakaryocyte polyploidy (Zhang *et al.* (2004) *Blood* 103:3717-26). Inhibitors of this function in a proliferative disease such as cancer would lead to stasis and cell death, making such inhibitors useful in cancer chemotherapy.

Aurora C (Aurora-3) is the least studied, known member of the family.
10 Aurora C localizes to centrosomes from anaphase until telophase (or even cytokinesis), and is highly expressed in the testis (Brown *et al.* (2004) *BMC Evolutionary Biology* 4:39).

As noted above, Aurora kinases are overexpressed in certain types of cancers, including colon, breast, and other solid-tumor cancers. The genes
15 encoding the Aurora B and A kinases tend to be amplified in certain types of cancers, while the gene encoding the Aurora C kinase resides in a region of the chromosome that is subject to rearrangement and deletion. Aurora A has been associated with a variety of malignancies, including primary colon, colorectal, breast, stomach, ovarian, prostate, and cervical cancer, neuroblastoma, and other
20 solid-tumor cancers (Warner *et al.* (2003) *Molecular Cancer Therapeutics* 2:589-95).

Inhibitors of Aurora A have been described. For example, Harrington
et al. ((2004) *Nat. Med.* 10:262-67) have described VX-680, a small-molecule inhibitor that blocks cell-cycle progression and induces apoptosis in certain types
25 of tumors in *in vivo* xenograft models. A pyrazole Aurora A kinase inhibitor is also described in U.S. Patent No. 6,653,301 (Bebbington *et al.*, issued November 25, 2003).

Hauf *et al.* ((2003) *J. Cell. Biol.* 161:281-294) identified the indolinone (Hesperadin) as an inhibitor of Aurora B, which causes cells to enter anaphase
30 with monooriented chromosomes, having both sister kinetochores attached to a single spindle pole (a condition known as syntelic attachment).

Ditchfield *et al.* ((2003) *J. Cell. Biol.* 161:267-280) described ZM447439 ((4-(4-(*N*-benzoylamino)anilino)-6-methoxy-7-(3-(1-morpholino)propoxy)quina-

zoline), an Aurora kinase inhibitor which interferes with chromosome alignment, segregation, and cytokinesis.

Accordingly, kinase inhibitors, particularly inhibitors of Aurora kinases, are of particular interest in treating certain disorders, including cancer.

5 Compounds exhibiting such inhibition are of particular value.

SUMMARY OF THE INVENTION

The present invention provides compounds or pharmaceutically acceptable derivatives or prodrugs thereof, compositions, and methods for treating diseases
10 mediated by kinases. Such diseases include primary, secondary, and metastatic cancers such as melanoma, lymphoma, leukemia, colon, colorectal, breast, lung, kidney, pancreatic, renal, CNS, stomach, ovarian, prostate, cervical, and neuroblastoma.

These compounds have the general formula I:



Formula I

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:

R^x and R^y are independently selected from the group consisting of $-T-R^3$
20 and $-L-Z-R^3$;

Q' is selected from the group consisting of $-CR^{6''}=CR^{6''}-$ and $\text{---}\equiv\text{---}$, wherein said $-CR^{6''}=CR^{6''}-$ may be a *cis* or *trans* double bond or a mixture thereof;

R^1 is $-T-(\text{Ring D})$;

25 Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from the group consisting of aryl, heteroaryl, heterocyclyl, and carbocyclyl, said heteroaryl or heterocyclyl ring having 1-4 ring heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, wherein each substitutable ring carbon of Ring D is independently substituted by oxo, $-T-R^5$, or

-V-Z-R⁵, and each substitutable ring nitrogen of Ring D is independently substituted by -R⁴;

T is a valence bond or -(C(R⁶)₂)-A-;

A is a valence bond or a C₁₋₃ alkylidene chain wherein a methylene unit
5 of said C₁₋₃ alkylidene chain is optionally replaced by -O-, -S-, -N(R⁴)-, -CO-, -CONH-, -NHCO-, -SO₂-, -SO₂NH-, -NHSO₂-, -CO₂-, -OC(O)-, -OC(O)NH-, or -NHCO₂-;

Z is a C₁₋₄ alkylidene chain;

L is selected from the group consisting of -O-, -S-, -SO-, -SO₂-, -N(R⁶)SO₂
10 -, -SO₂N(R⁶)-, -N(R⁶)-, -CO-, -CO₂-, -N(R⁶)CO-, -N(R⁶)C(O)O-, -N(R⁶)CON(R⁶)-, -N(R⁶)SO₂N(R⁶)-, -N(R⁶)N(R⁶)-, -C(O)N(R⁶)-, -OC(O)N(R⁶)-, -C(R⁶)₂O-, -C(R⁶)₂-, -C(R⁶)₂SO-, -C(R⁶)₂SO₂-, -C(R⁶)₂SO₂N(R⁶)-, -C(R⁶)₂N(R⁶)-, -C(R⁶)₂N(R⁶)C(O)-, -C(R⁶)₂N(R⁶)C(O)O-, -C(R⁶)=NN(R⁶)-, -C(R⁶)=N-O-, -C(R⁶)₂N(R⁶)N(R⁶)-, -C(R⁶)₂N(R⁶)SO₂N(R⁶)-, and -C(R⁶)₂N(R⁶)CON(R⁶)-;

15 R² and R^{2'} are independently selected from the group consisting of -R and -T-W-R⁶, or R² and R^{2'} taken together with their intervening atoms form a fused, 5-8 membered, unsaturated or partially unsaturated ring having 0-3 ring heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, wherein each substitutable ring carbon of said fused ring formed by R² and R^{2'} is
20 independently substituted by halo, oxo, -CN, -NO₂, R⁷, or -V-R⁶, and each substitutable ring nitrogen of said ring formed by R² and R^{2'} is independently substituted by -R⁴;

R³ is selected from the group consisting of -R, -halo, -OR, -C(=O)R, -CO₂R, -COCOR, -COCH₂COR, -NO₂, -CN, -S(O)R, -S(O)₂R, -SR, -N(R⁴)₂, -
25 CON(R⁷)₂, -SO₂N(R⁷)₂, -OC(=O)R, -N(R⁷)COR, -N(R⁷)CO₂(C₁₋₆ aliphatic), -N(R⁴)N(R⁴)₂, -C=NN(R⁴)₂, -C=N-OR, -N(R⁷)CON(R⁷)₂, -N(R⁷)SO₂N(R⁷)₂, -N(R⁴)SO₂R, and -OC(=O)N(R)₂;

each R is independently hydrogen or an optionally substituted group selected from the group consisting of C₁₋₆ aliphatic, C₆₋₁₀ aryl, a heteroaryl ring
30 having 5-10 ring atoms, and a heterocyclyl ring having 5-10 ring atoms;

each R⁴ is independently selected from the group consisting of -R⁷, -COR⁷, -CO₂(optionally substituted C₁₋₆ aliphatic), -CON(R⁷)₂, and -SO₂R⁷;

each R⁵ is independently selected from the group consisting of -R, halo, -OR, -C(=O)R, -CO₂R, -COCOR, -NO₂, -CN, -S(O)R, -SO₂R, -SR, -N(R⁴)₂, -

CON(R⁴)₂, -SO₂N(R⁴)₂, -OC(=O)R, -N(R⁴)COR, -N(R⁴)CO₂ (optionally substituted C₁₋₆ aliphatic), -N(R⁴)N(R⁴)₂, -C=NN(R⁴)₂, -C=N-OR, -N(R⁴)CON(R⁴)₂, -N(R⁴)SO₂N(R⁴)₂, -N(R⁴)SO₂R, and -OC(=O)N(R⁴)₂;

V is selected from the group consisting of -O-, -S-, -SO-, -SO₂-, -N(R⁶)SO₂-, -SO₂N(R⁶)-, -N(R⁶)-, -CO-, -CO₂-, -N(R⁶)CO-, -N(R⁶)C(O)O-, -N(R⁶)CON(R⁶)-, -N(R⁶)SO₂N(R⁶)-, -N(R⁶)N(R⁶)-, -C(O)N(R⁶)-, -OC(O)N(R⁶)-, -C(R⁶)₂O-, -C(R⁶)₂S-, -C(R⁶)₂SO-, -C(R⁶)₂SO₂-, -C(R⁶)₂SO₂N(R⁶)-, -C(R⁶)₂N(R⁶)-, -C(R⁶)₂N(R⁶)C(O)-, -C(R⁶)₂N(R⁶)C(O)O-, -C(R⁶)=NN(R⁶)-, -C(R⁶)=N-O-, -C(R⁶)₂N(R⁶)N(R⁶)-, -C(R⁶)₂N(R⁶)SO₂N(R⁶)-, and -C(R⁶)₂N(R⁶)CON(R⁶)-;

W is selected from the group consisting of -C(R⁶)₂O-, -C(R⁶)₂S-, -C(R⁶)₂SO-, -C(R⁶)₂SO₂-, -C(R⁶)₂SO₂N(R⁶)-, -C(R⁶)₂N(R⁶)-, -CO-, -CO₂-, -C(R⁶)OC(O)-, -C(R⁶)OC(O)N(R⁶)-, -C(R⁶)₂N(R⁶)CO-, -C(R⁶)₂N(R⁶)C(O)O-, -C(R⁶)=NN(R⁶)-, -C(R⁶)=N-O-, -C(R⁶)₂N(R⁶)N(R⁶)-, -C(R⁶)₂N(R⁶)SO₂N(R⁶)-, -C(R⁶)₂N(R⁶)CON(R⁶)-, and -CON(R⁶)-;

each R⁶ is independently selected from the group consisting of hydrogen and an optionally substituted C₁₋₄ aliphatic group, or two R⁶ groups on the same nitrogen atom may be taken together with the nitrogen atom to form a 3-6 membered heterocyclyl or heteroaryl ring;

each R^{6'} is independently selected from the group consisting of hydrogen and a C₁₋₄ aliphatic group, or two R^{6'} on the same carbon atom are taken together to form a 3-8 membered carbocyclic ring;

each R^{6''} is independently selected from the group consisting of hydrogen, a C₁₋₄ aliphatic group, halogen, optionally substituted aryl, and optionally substituted heteroaryl, or two R^{6''} on adjacent carbon atoms are taken together to form a 5-7 membered carbocyclic ring; and

each R⁷ is independently selected from the group consisting of hydrogen and an optionally substituted C₁₋₆ aliphatic group, or two R⁷ on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring.

DETAILED DESCRIPTION

As used herein, the following definitions shall apply unless otherwise indicated. The phrase "optionally substituted" is used interchangeably with the phrase "substituted or unsubstituted" or with the term "(un)substituted." Unless

otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group, and each substitution is independent of the other.

The term "acetamido" refers to the group -NHC(=O)CH_3 .

5 The term "aliphatic" as used herein means straight-chain, branched or cyclic $\text{C}_1\text{-C}_{12}$ hydrocarbons which are completely saturated or which contain one or more units of unsaturation but which are not aromatic. For example, suitable aliphatic groups include substituted or unsubstituted linear, branched or cyclic alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl. The terms "alkyl", "alkoxy", "hydroxyalkyl", "alkoxyalkyl", and "alkoxycarbonyl", used alone or as part of a larger moiety includes both straight and branched chains containing one to twelve carbon atoms. The terms "alkenyl" and "alkynyl" used alone or as part of a larger moiety shall include both straight and branched chains containing two to twelve carbon atoms. The term "cycloalkyl" used alone or as part of a larger moiety shall include cyclic $\text{C}_3\text{-C}_{12}$ hydrocarbons which are completely saturated or which contain one or more units of unsaturation, but which are not aromatic.

The term "amino" refers to an NH_2 group.

20 The term "alkylamino" refers to an amino group wherein one of the hydrogen atoms is replaced by an alkyl group.

The term "dialkylamino" refers to an amino group wherein the hydrogen atoms are replaced by alkyl groups, wherein the alkyl group may be the same or different.

25 The terms "haloalkyl", "haloalkenyl" and "haloalkoxy" means alkyl, alkenyl or alkoxy, as the case may be, substituted with one or more halogen atoms.

The term "halogen" means F, Cl, Br, or I.

30 The term "heteroatom" means nitrogen, oxygen, or sulfur and includes any oxidized form of nitrogen and sulfur, and the quaternized form of any basic nitrogen. Also the term "nitrogen" includes a substitutable nitrogen of a heterocyclic ring. As an example, in a saturated or partially unsaturated ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, the nitrogen may be N (as in 3,4-dihydro-2H-pyrrolyl), NH (as in pyrrolidinyl) or NR^+ (as in N-substituted pyrrolidinyl).

The terms "carbocycle", "carbocyclyl", "carbocyclo", or "carbocyclic" as used herein means an aliphatic ring system having three to fourteen members. The terms "carbocycle", "carbocyclyl", "carbocyclo", or "carbocyclic" whether saturated or partially unsaturated, also refers to rings that are optionally substituted. The terms "carbocycle", "carbocyclyl", "carbocyclo", or "carbocyclic" also include aliphatic rings that are fused to one or more aromatic or nonaromatic rings, such as in a decahydronaphthyl or tetrahydronaphthyl, where the radical or point of attachment is on the aliphatic ring.

The term "aryl" used alone or as part of a larger moiety as in "aralkyl", "aralkoxy", or "aryloxyalkyl", refers to aromatic ring groups having six to fourteen members, such as phenyl, benzyl, phenethyl, 1-naphthyl, 2-naphthyl, 1-anthracyl and 2-anthracyl. The term "aryl" also refers to rings that are optionally substituted. The term "aryl" may be used interchangeably with the term "aryl ring". "Aryl" also includes fused polycyclic aromatic ring systems in which an aromatic ring is fused to one or more rings. Examples include 1-naphthyl, 2-naphthyl, 1-anthracyl and 2-anthracyl. Also included within the scope of the term "aryl", as it is used herein, is a group in which an aromatic ring is fused to one or more non-aromatic rings, such as in an indanyl, phenanthridinyl, or tetrahydronaphthyl, where the radical or point of attachment is on the aromatic ring.

The term "heterocycle", "heterocyclyl", or "heterocyclic" as used herein includes non-aromatic ring systems having four to fourteen members, preferably five to ten, in which one or more ring carbons, preferably one to four, are each replaced by a heteroatom. Examples of heterocyclic rings include 3-1H-benzimidazol-2-one, (1-substituted)-2-oxo-benzimidazol-3-yl, 2-tetrahydrofuranyl, 3-tetrahydrofuranyl, 2-tetrahydropyranyl, 3-tetrahydropyranyl, 4-tetrahydropyranyl, [1,3]-dioxalanyl, [1,3]-dithiolanyl, [1,3]-dioxanyl, 2-tetrahydrothiophenyl, 3-tetrahydrothiophenyl, 2-morpholinyl, 3-morpholinyl, 4-morpholinyl, 2-thiomorpholinyl, 3-thiomorpholinyl, 4-thiomorpholinyl, 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 1-piperazinyl, 2-piperazinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 4-thiazolidinyl, diazolonyl, N-substituted diazolonyl, 1-phthalimidinyl, benzoxanyl, benzopyrrolidinyl, benzopiperidinyl, benzoxolanyl, benzothiolanyl, and benzothianyl. Also included within the scope of the term "heterocyclyl" or "heterocyclic", as it is used herein, is a group in

which a non-aromatic heteroatom-containing ring is fused to one or more aromatic or non-aromatic rings, such as in an indolinyl, chromanyl, phenanthridinyl, or tetrahydroquinolinyl, where the radical or point of attachment is on the non-aromatic heteroatom-containing ring. The term "heterocycle", "heterocyclyl", or
 5 "heterocyclic" whether saturated or partially unsaturated, also refers to rings that are optionally substituted.

The term "heteroaryl", used alone or as part of a larger moiety as in "heteroaralkyl" or "heteroarylalkoxy", refers to heteroaromatic ring groups having five to fourteen members. Examples of heteroaryl rings include 2-furanyl, 3-
 10 furanyl, 3-furazanyl, N-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-oxadiazolyl, 5-oxadiazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 1-pyrazolyl, 2-pyrazolyl, 3-pyrazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-pyrimidyl, 3-pyridazinyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 5-tetrazolyl, 2-
 15 triazolyl, 5-triazolyl, 2-thienyl, 3-thienyl, carbazolyl, benzimidazolyl, benzothienyl, benzofuranyl, indolyl, quinolinyl, benzotriazolyl, benzothiazolyl, benzooxazolyl, benzimidazolyl, isoquinolinyl, indazolyl, isoindolyl, acridinyl, and benzoisoxazolyl. Also included within the scope of the term "heteroaryl", as it is used herein, is a group in which a heteroatomic ring is fused to one or more
 20 aromatic or nonaromatic rings where the radical or point of attachment is on the heteroaromatic ring. Examples include tetrahydroquinolinyl, tetrahydroisoquinolinyl, and pyrido [3,4-d]pyrimidinyl. The term "heteroaryl" also refers to rings that are optionally substituted. The term "heteroaryl" may be used interchangeably with the term "heteroaryl ring" or the term "heteroaromatic".

25 An aryl (including aralkyl, aralkoxy, aryloxyalkyl and the like) or heteroaryl (including heteroaralkyl and heteroarylalkoxy and the like) group may contain one or more substituents. Examples of suitable substituents on any unsaturated carbon atom of an aryl, heteroaryl, aralkyl, or heteroaralkyl group include a halogen, $-R^0$, $-OR^0$, $-SR^0$, 1,2-methylene-dioxy, 1,2-ethylenedioxy, protected OH (such as acyloxy), phenyl (Ph), substituted Ph, $-O(Ph)$, substituted $-O(Ph)$, $-CH_2(Ph)$, substituted $-CH_2(Ph)$, $-CH_2CH_2(Ph)$, substituted $-CH_2CH_2(Ph)$, $-NO_2$, $-CN$, $-N(R^0)_2$, $-NR^0C(O)R^0$, $-NR^0C(O)N(R^0)_2$, $-NR^0CO_2R^0$, $-NR^0NR^0C(O)R^0$, $-NR^0NR^0C(O)N(R^0)_2$, $-NR^0NR^0C_2R^0$, $-C(O)C(O)R^0$, $-C(O)CH_2C(O)R^0$, $-CO_2R^0$, $-C(O)R^0$, $-C(O)N(R^0)_2$, $-OC(O)N(R^0)_2$, $-S(O)_2R^0$, $-$

$\text{SO}_2\text{N}(\text{R}^0)_2$, $-\text{S}(\text{O})\text{R}^0$, $-\text{NR}^0\text{SO}_2\text{N}(\text{R}^0)_2$, $-\text{NR}^0\text{SO}_2\text{R}^0$, $-\text{C}(=\text{S})\text{N}(\text{R}^0)_2$, $-\text{C}(=\text{NH})-\text{N}(\text{R}^0)_2$, $-(\text{CH}_2)_y\text{NHC}(\text{O})\text{R}^0$, and $-(\text{CH}_2)_y\text{NHC}(\text{O})\text{CH}(\text{V}-\text{R}^0)(\text{R}^0)$; wherein each R^0 is independently selected from hydrogen, a substituted or unsubstituted aliphatic group, an unsubstituted heteroaryl or heterocyclic ring, phenyl (Ph), substituted Ph, -O(Ph), substituted -O(Ph), -CH₂ (Ph), or substituted -CH₂(Ph); y is 0-6; and V is a linker group. Examples of substituents on the aliphatic group or the phenyl ring of R^0 include amino, alkylamino, dialkylamino, aminocarbonyl, halogen, alkyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminocarbonyloxy, dialkylaminocarbonyloxy, alkoxy, nitro, cyano, carboxy, alkoxycarbonyl, alkylcarbonyl, hydroxy, haloalkoxy, and haloalkyl.

An aliphatic group or a non-aromatic heterocyclic ring or a fused aryl or heteroaryl ring may contain one or more substituents. Examples of suitable substituents on any saturated carbon of an aliphatic group or of a non-aromatic heterocyclic ring or a fused aryl or heteroaryl ring include those listed above for the unsaturated carbon of an aryl or heteroaryl group and the following: =O, =S, =NNHR*, =NN(R*)₂, =N-, =NNHC(O)R*, =NNHCO₂(alkyl), =NNHSO₂ (alkyl), or =NR*, where each R* is independently selected from hydrogen, an unsubstituted aliphatic group, or a substituted aliphatic group. Examples of substituents on the aliphatic group include amino, alkylamino, dialkylamino, aminocarbonyl, halogen, alkyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminocarbonyloxy, dialkylaminocarbonyloxy, alkoxy, nitro, cyano, carboxy, alkoxycarbonyl, alkylcarbonyl, hydroxy, haloalkoxy, and haloalkyl.

Suitable substituents on the nitrogen of a non-aromatic heterocyclic ring include $-\text{R}^+$, $-\text{N}(\text{R}^+)_2$, $-\text{C}(\text{O})\text{R}^+$, $-\text{CO}_2\text{R}^+$, $-\text{C}(\text{O})\text{C}(\text{O})\text{R}^+$, $-\text{C}(\text{O})\text{CH}_2\text{C}(\text{O})\text{R}^+$, $-\text{SO}_2\text{R}^+$, $-\text{SO}_2\text{N}(\text{R}^+)_2$, $-\text{C}(=\text{S})\text{N}(\text{R}^+)_2$, $-\text{C}(=\text{NH})-\text{N}(\text{R}^+)_2$, and $-\text{NR}^+\text{SO}_2\text{R}^+$; wherein each R^+ is independently selected from hydrogen, an aliphatic group, a substituted aliphatic group, phenyl (Ph), substituted Ph, -O(Ph), substituted -O(Ph), -CH₂(Ph), substituted -CH₂(Ph), or an unsubstituted heteroaryl or heterocyclic ring. Examples of substituents on the aliphatic group or the phenyl ring include amino, alkylamino, dialkylamino, aminocarbonyl, halogen, alkyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminocarbonyloxy, dialkylaminocarbonyloxy, alkoxy, nitro, cyano, carboxy, alkoxycarbonyl, alkylcarbonyl, hydroxy, haloalkoxy, and haloalkyl.

The term "linker group" or "linker" means an organic moiety that connects two parts of a compound. Linkers are typically comprised of an atom such as oxygen or sulfur, a unit such as -NH-, -CH₂-, -C(O)-, -C(O)NH-, or a chain of atoms, such as an alkylidene chain. The molecular mass of a linker is typically in the range of about 14 to 200, preferably in the range of 14 to 96 with a length of up to about six atoms. Examples of linkers include a saturated or unsaturated C₁₋₆ alkylidene chain which is optionally substituted, and wherein one or two saturated carbons of the chain are optionally replaced by -C(O)-, -C(O)C(O)-, -CONH-, -CONHNH-, -CO₂-, -OC(O)-, -NHCO₂-, -O-, -NHCONH-, -OC(O)NH-, -NHNH-, -NHCO-, -S-, -SO-, -SO₂-, -NH-, -SO₂NH-, or -NHSO₂-.

The term "alkylidene chain" refers to an optionally substituted, straight or branched carbon chain, that may be fully saturated or have one or more units of unsaturation. The optional substituents are as described above for an aliphatic group.

A combination of substituents or variables is permissible only if such a combination results in a stable or chemically feasible compound. A stable compound or chemically feasible compound is one in which the chemical structure is not substantially altered when kept in the dark at a temperature of 40°C or less, in the absence of moisture or other chemically reactive conditions, for at least a week.

It is understood that in all substituted groups defined herein, polymers arrived at by defining substituents with further substituents to themselves (e.g., substituted phenyl having a substituted phenyl as a substituent which is itself substituted with a substituted phenyl, etc.) are not intended for inclusion herein. In such cases, the maximum number of such substituents is three. For example phenyl substituted with a substituted phenyl is limited to -substituted phenyl-(substituted phenyl)-(substituted phenyl).

Unless otherwise stated, structures depicted herein are also meant to include all stereochemical forms of the structure; i.e., the R and S configurations for each asymmetric center. Therefore, single stereochemical isomers as well as enantiomeric and diastereomeric mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, structures depicted herein are also meant to include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present

structures except for the replacement of a hydrogen by a deuterium or tritium, or the replacement of a carbon by a ^{13}C - or ^{14}C -enriched carbon are within the scope of this invention.

Compounds of formula I, or salts thereof, may be formulated into compositions. In a preferred embodiment, the composition is a pharmaceutical composition. In one embodiment, the composition comprises an amount of the protein kinase inhibitor effective to inhibit a protein kinase in a biological sample or in a patient. Compounds of this invention and pharmaceutical compositions thereof, which comprise an amount of the protein kinase inhibitor effective to treat or prevent a kinase mediated condition and a pharmaceutically acceptable carrier, adjuvant, or vehicle, may be formulated for administration to a patient.

Another aspect of this invention relates to a method of treating or preventing a kinase mediated disease. In one embodiment, the disease is a Aurora A-mediated disease, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I or a pharmaceutical composition thereof.

The term "Aurora A-mediated disease" or "Aurora A-mediated condition", as used herein, means any disease or other deleterious condition in which Aurora is thought to play a role. The terms "Aurora A-mediated disease" or "Aurora A-mediated condition" also mean those diseases or conditions that are alleviated by treatment with an Aurora A inhibitor. Such conditions include cancer.

The term "cancer" includes, but is not limited to, solid tumors and blood borne tumors and include, but is not limited to, the following cancers: breast, ovary, cervix, prostate, testis, genitourinary tract, esophagus, larynx, glioblastoma, neuroblastoma, stomach, skin, keratoacanthoma, lung, epidermoid carcinoma, large cell carcinoma, small cell carcinoma, lung adenocarcinoma, bone, colon, adenoma, pancreas, adenocarcinoma, thyroid, follicular carcinoma, undifferentiated carcinoma, papillary carcinoma, seminoma, melanoma, sarcoma, bladder carcinoma, liver carcinoma and biliary passages, kidney carcinoma, myeloid disorders, lymphoid disorders, Hodgkin's, hairy cells, buccal cavity and pharynx (oral), lip, tongue, mouth, pharynx, small intestine, colon-rectum, large intestine, rectum, brain and central nervous system, and leukemia. The term "cancer" includes primary cancer, cancers secondary to treatment, and metastatic cancers.

An aspect of the invention relates to compounds and compositions that are useful for treating cancer.

Another aspect of the invention relates to the treatment of the following cancers: breast, ovary, cervix, prostate, testis, genitourinary tract, esophagus, larynx, glioblastoma, neuroblastoma, stomach, skin, keratoacanthoma, lung, epidermoid carcinoma, large cell carcinoma, small cell carcinoma, lung adenocarcinoma, bone, colon, adenoma, pancreas, adenocarcinoma, thyroid, follicular carcinoma, undifferentiated carcinoma, papillary carcinoma, seminoma, melanoma, sarcoma, bladder carcinoma, liver carcinoma and biliary passages, kidney carcinoma, myeloid disorders, lymphoid disorders, Hodgkin's, hairy cells, buccal cavity and pharynx (oral), lip, tongue, mouth, pharynx, small intestine, colon-rectum, large intestine, rectum, brain and central nervous system, and leukemia.

Another aspect of the invention is a method for treating cancer comprising administering one or more of the compounds described herein to a patient with cancer.

Angiogenesis is characterized by the proliferation of endothelial cells to form new blood vessels (often called neovascularization). Inhibition of mitosis of endothelial cells results in inhibition of angiogenesis. Another aspect of this invention therefore relates to inhibition of undesirable mitosis, including undesirable angiogenesis. A mammalian disease characterized by undesirable cell mitosis, as defined herein, includes, but is not limited to, excessive or abnormal stimulation of endothelial cells (e.g., atherosclerosis), solid tumors and tumor metastasis, benign tumors, for example, hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, vascular malfunctions, abnormal wound healing, inflammatory and immune disorders, Bechet's disease, gout or gouty arthritis, abnormal angiogenesis accompanying rheumatoid arthritis, skin diseases, such as psoriasis, diabetic retinopathy and other ocular angiogenic diseases such as retinopathy of prematurity (retrolental fibroplastic), macular degeneration, corneal graft rejection, neovascular glaucoma and Osler Weber syndrome (Osler-Weber-Rendu disease).

Other undesired angiogenesis involves normal processes including ovulation and implantation of a blastula. The compositions described above can be used as a birth control agent by reducing or preventing uterine vascularization

required for embryo implantation. Accordingly, the compositions described above can be used to block ovulation and implantation of a blastula or to block menstruation (induce amenorrhea).

Diseases associated with undesirable mitosis including neovascularization can be treated according to the present invention. Such diseases include, but are not limited to, ocular neovascular disease, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, neovascular glaucoma and retrolental fibroplasias, epidemic keratoconjunctivitis, Vitamin A deficiency, contact lens overwear, atopic keratitis, superior limbic keratitis, pterygium keratitis sicca, Sjögren's syndrome, acne rosacea, phlyctenulosis, syphilis, *Mycobacteria* infections, lipid degeneration, chemical burns, bacterial ulcers, fungal ulcers, *Herpes simplex* infections, *Herpes zoster* infections, protozoan infections, Kaposi's sarcoma, Mooren's ulcer, Terrien's marginal degeneration, marginal keratolysis, trauma, rheumatoid arthritis, systemic lupus, polyarteritis, Wegener's sarcoidosis, Scleritis, Steven-Johnson disease, pemphigoid, radial keratotomy, and corneal graft rejection.

Other diseases associated with undesirable mitosis including neovascularization can be treated according to the present invention. Such diseases include, but are not limited to, sickle cell anemia, sarcoid, pseudoxanthoma elasticum, Paget's disease, vein occlusion, artery occlusion, carotid obstructive disease, chronic uveitis/vitritis, Lyme's disease, systemic lupus erythematosus, Eales' disease, Bechet's disease, infections causing a retinitis or choroiditis, presumed ocular histoplasmosis, Best's disease, myopia, optic pits, Stargart's disease, pars planitis, chronic retinal detachment, hyperviscosity syndromes, toxoplasmosis, and post-laser complications. Other diseases include, but are not limited to, diseases associated with rubeosis (neovascularization of the iris and the angle) and diseases caused by the abnormal proliferation of fibrovascular or fibrous tissue including all forms of proliferative vitreoretinopathy, whether or not associated with diabetes.

Another aspect of the invention relates to the treatment of inflammatory diseases including, but not limited to, excessive or abnormal stimulation of endothelial cells (e.g., atherosclerosis), solid tumors and tumor metastasis, benign tumors, for example, hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, vascular malfunctions, abnormal wound

healing, inflammatory and immune disorders, Bechet's disease, gout or gouty arthritis, abnormal angiogenesis accompanying rheumatoid arthritis, skin diseases, such as psoriasis, diabetic retinopathy and other ocular angiogenic diseases such as retinopathy of prematurity (retrolental fibroplastic), macular degeneration, corneal graft rejection, neovascular glaucoma and Osler Weber syndrome (Osler-Weber-Rendu disease). Other undesired angiogenesis involves normal processes including ovulation and implantation of a blastula. Accordingly, the compositions described above can be used to block ovulation and implantation of a blastula or to block menstruation (induce amenorrhea).

Another aspect of the invention relates to inhibiting Aurora A activity in a biological sample, which method comprises contacting the biological sample with the Aurora A inhibitor of formula I, or a composition thereof.

Another aspect of this invention relates to a method of inhibiting Aurora A activity in a patient, which method comprises administering to the patient a compound of formula I or a composition comprising said compound.

In another aspect of this invention, compounds of formula I are more potent inhibitors of Aurora A compared to Aurora B.

Another aspect of this invention relates to a method of treating or preventing a GSK-3-mediated disease with a GSK-3 inhibitor, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I or a pharmaceutical composition thereof.

The terms "GSK-3-mediated disease, or "GSK-3-mediated condition", as used herein, mean any disease or other deleterious condition or state in which GSK-3 is known to play a role. Such diseases or conditions include, without limitation, diabetes, Alzheimer's disease, Huntington's Disease, Parkinson's Disease, AIDS-associated dementia, amyotrophic lateral sclerosis (AML), multiple sclerosis (MS), schizophrenia, cardiomyocyte hypertrophy, reperfusion/ischemia, and baldness.

One aspect of this invention relates to a method of enhancing glycogen synthesis and/or lowering blood levels of glucose in a patient in need thereof, which method comprises administering to the patient a therapeutically effective amount of a compound of formula I or a pharmaceutical composition thereof. This method is especially useful for diabetic patients. Another method relates to

inhibiting the production of hyperphosphorylated Tau protein, which is useful in halting or slowing the progression of Alzheimer's disease. Another method relates to inhibiting the phosphorylation of .beta.-catenin, which is useful for treating schizophrenia.

5 Another aspect of the invention relates to inhibiting GSK-3 activity in a biological sample, which method comprises contacting the biological sample with a GSK-3 inhibitor of formula I.

 Another aspect of this invention relates to a method of inhibiting GSK-3 activity in a patient, which method comprises administering to the patient a
10 compound of formula I or a composition comprising said compound.

 Another aspect of this invention relates to a method of treating or preventing a CDK-2-mediated disease with a CDK-2 inhibitor, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I or a pharmaceutical composition
15 thereof.

 The terms "CDK-2-mediated disease" or "CDK-2-mediated condition", as used herein, mean any disease or other deleterious condition in which CDK-2 is known to play a role. The terms "CDK-2-mediated disease" or "CDK-2-mediated condition" also mean those diseases or conditions that are alleviated by treatment
20 with a CDK-2 inhibitor. Such conditions include, without limitation, cancer, Alzheimer's disease, restenosis, angiogenesis, glomerulonephritis, cytomegalovirus, HIV, herpes, psoriasis, atherosclerosis, alopecia, and autoimmune diseases such as rheumatoid arthritis, such as are described for example in Fischer, P. M. and Lane, D. P., Current Medicinal Chemistry, 7, 1213-
25 1245 (2000); Mani, S., Wang, C., Wu, K., Francis, R. and Pestell, R., Exp. Opin. Invest. Drugs, 9, 1849 (2000); Fry, D. W. and Garrett, M. D., Current Opinion in Oncologic, Endocrine & Metabolic Investigational Drugs, 2, 40-59 (2000).

 Another aspect of the invention relates to inhibiting CDK-2 activity in a biological sample or a patient, which method comprises administering to the
30 patient a compound of formula I, or a composition comprising said compound.

 Another aspect of this invention relates to a method of treating or preventing an ERK-2-mediated diseases with an ERK-2 inhibitor, which method comprises administering to a patient in need of such a treatment a therapeutically

effective amount of a compound of formula I or a pharmaceutical composition thereof.

The terms "ERK-mediated disease" or "ERK-mediated condition", as used herein mean any disease or other deleterious condition in which ERK may play a role. The terms "ERK-2-mediated disease" or "ERK-2-mediated condition" also mean those diseases or conditions that are alleviated by treatment with a ERK-2 inhibitor. Such conditions include, without limitation, cancer, stroke, diabetes, hepatomegaly, cardiovascular disease including cardiomegaly, Alzheimer's disease, cystic fibrosis, viral disease, autoimmune diseases, atherosclerosis, restenosis, psoriasis, allergic disorders including asthma, inflammation, neurological disorders and hormone-related diseases. ERK-2 protein kinase and its implication in various diseases has been described for example in Bokemeyer et al. 1996, *Kidney Int.* 49, 1187; Anderson et al., 1990, *Nature* 343, 651; Crews et al., 1992, *Science* 258, 478; Bjorbaek et al., 1995, *J. Biol. Chem.* 270, 18848; Rouse et al., 1994, *Cell* 78, 1027; Raingeaud et al., 1996, *Mol. Cell Biol.* 16, 1247; Raingeaud et al. 1996; Chen et al., 1993 *Proc. Natl. Acad. Sci. USA* 90, 10952; Oliver et al., 1995, *Proc. Soc. Exp. Biol. Med.* 210, 162; Moodie et al., 1993, *Science* 260, 1658; Frey and Mulder, 1997, *Cancer Res.* 57, 628; Sivaraman et al., 1997, *J Clin. Invest.* 99, 1478; Whelchel et al., 1997, *Am. J. Respir. Cell Mol. Biol.* 16, 589.

Another aspect of the invention relates to inhibiting ERK-2 activity in a biological sample or a patient, which method comprises administering to the patient a compound of formula I, or a composition comprising said compound.

Another aspect of this invention relates to a method of treating or preventing an AKT-mediated diseases with an AKT inhibitor, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I or a pharmaceutical composition thereof.

The terms "AKT-mediated disease" or "AKT-mediated condition", as used herein, mean any disease or other deleterious condition in which AKT is known to play a role. The terms "AKT-mediated disease" or "AKT-mediated condition" also mean those diseases or conditions that are alleviated by treatment with a AKT inhibitor. AKT-mediated diseases or conditions include, but are not limited to, proliferative disorders, cancer, and neurodegenerative disorders. The association

of AKT, also known as protein kinase B, with various diseases has been described for example in Khwaja, A., Nature, pp. 33-34, 1990; Zang, Q. Y., et al, Oncogene, 19 2000; Kazuhiko, N., et al, The Journal of Neuroscience, 20 2000.

Another aspect of the invention relates to inhibiting AKT activity in a
5 biological sample or a patient, which method comprises administering to the patient a compound of formula I, or a composition comprising said compound.

Another aspect of this invention relates to a method of treating or preventing a Src-mediated disease with a Src inhibitor, which method comprises administering to a patient in need of such a treatment a therapeutically effective
10 amount of a compound of formula I or a pharmaceutical composition thereof.

The terms "Src-mediated disease" or "Src-mediated condition", as used herein mean any disease or other deleterious condition in which Src is known to play a role. The terms "Src-mediated disease" or "Src-mediated condition" also mean those diseases or conditions that are alleviated by treatment with a Src
15 inhibitor. Such conditions include, without limitation, hypercalcemia, osteoporosis, osteoarthritis, cancer, symptomatic treatment of bone metastasis, and Paget's disease. Src protein kinase and its implication in various diseases has been described for example in Soriano, Cell, 69, 551 (1992); Soriano et al., Cell, 64, 693 (1991); Takayanagi, J. Clin. Invest., 104, 137 (1999); Boschelli, Drugs of
20 the Future 2000, 25(7), 717, (2000); Talamonti, J. Clin. Invest., 91, 53 (1993); Lutz, Biochem. Biophys. Res. 243, 503 (1998); Rosen, J. Biol. Chem., 261, 13754 (1986); Bolen, Proc. Natl. Acad. Sci. USA, 84, 2251 (1987); Masaki, Hepatology, 27, 1257 (1998); Biscardi, Adv. Cancer Res., 76, 61 (1999); Lynch, Leukemia, 7, 1416 (1993); Wiener, Clin. Cancer Res., 5, 2164 (1999); Staley, Cell Growth
25 Diff., 8, 269 (1997).

Another aspect of the invention relates to inhibiting Src activity in a biological sample or a patient, which method comprises administering to the patient a compound of formula I, or a composition comprising said compound.

Another aspect of this invention relates to a method of treating or preventing an Lck-mediated disease with an Lck inhibitor, which method
30 comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I, or a pharmaceutical composition thereof.

The terms "Lck-mediated disease" or "Lck-mediated condition", as used herein, mean any disease state or other deleterious condition in which Lck is known to play a role. The terms "Lck-mediated disease" or "Lck-mediated condition" also mean those diseases or conditions that are alleviated by treatment
5 with an Lck inhibitor. Lck-mediated diseases or conditions include, but are not limited to, autoimmune diseases such as transplant rejection, allergies, rheumatoid arthritis, and leukemia. The association of Lck with various diseases has been described for example in Molina et al., *Nature*, 357, 161 (1992).

Another aspect of the invention relates to inhibiting Lck activity in a
10 biological sample or a patient, which method comprises administering to the patient a compound of formula I, or a composition comprising said compound.

Another aspect of this invention relates to a method of treating or preventing an Abl-mediated disease with an Abl inhibitor, which method comprises administering to a patient in need of such a treatment a therapeutically
15 effective amount of a compound of formula I, or a pharmaceutical composition thereof.

The terms "Abl-mediated disease" or "Abl-mediated condition", as used herein, mean any disease state or other deleterious condition in which Abl is known to play a role. The terms "Abl-mediated disease" or "Abl-mediated
20 condition" also mean those diseases or conditions that are alleviated by treatment with an Abl inhibitor. Abl-mediated diseases or conditions include, but are not limited to, leukemias, particularly chronic myeloid leukemia. The association of Abl with various diseases has been described for example in Druker, *et al.*, *N. Engl. J. Med.* **2001**, 344, 1038-1042.

25 Another aspect of the invention relates to inhibiting Abl activity in a biological sample or a patient, which method comprises administering to the patient a compound of formula I, or a composition comprising said compound.

Another aspect of this invention relates to a method of treating or preventing a cKit-mediated disease with an cKit inhibitor, which method
30 comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I, or a pharmaceutical composition thereof.

The terms "cKit-mediated disease" or "cKit-mediated condition", as used herein, mean any disease state or other deleterious condition in which cKit is

known to play a role. The terms "cKit-mediated disease" or "cKit-mediated condition" also mean those diseases or conditions that are alleviated by treatment with an cKit inhibitor. cKit-mediated diseases or conditions include, but are not limited to, mastocytosis/mast cell leukemia, gastrointestinal stromal tumor, 5 sinonasal natural killer/T-cell lymphoma, seminoma/dysgerminoma, throid carcinoma, samll-cell lung carcinoma, malignant melanoma, adenoid cystic carcinoma, ovarian carcinoma, acute myelogenous leukemia, anaplastic large-cell lymphoma, angiosarcoma, endometrial carcinom, pediatric T-cell ALL/lymphoma, breast carcinoma and prostate carcinoma. The association of 10 cKit with various diseases has been described for example in Heinrich, *et al.*, *J. Clinical Oncology* 2002, 20, 1692-1703.

Another aspect of the invention relates to inhibiting cKit activity in a biological sample or a patient, which method comprises administering to the patient a compound of formula I, or a composition comprising said compound.

15 Another aspect of this invention relates to a method of treating or preventing a Flt3-mediated disease with an Flt3 inhibitor, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I, or a pharmaceutical composition thereof.

20 The terms "Flt3-mediated disease" or "Flt3-mediated condition", as used herein, mean any disease state or other deleterious condition in which Flt3 is known to play a role. The terms "Flt3-mediated disease" or "Flt3-mediated condition" also mean those diseases or conditions that are alleviated by treatment with an Flt3 inhibitor. Flt3-mediated diseases or conditions include, but are not 25 limited to, acute myelogenous leukemia, mixed lineage leukemia and acute lymphocytic leukemia. The association of Flt3 with various diseases has been described for example in Sternberg and Licht, *Curr. Opin Hematol.* 2004, 12, 7-13.

Another aspect of the invention relates to inhibiting Flt3 activity in a 30 biological sample or a patient, which method comprises administering to the patient a compound of formula I, or a composition comprising said compound.

Another aspect of this invention relates to a method of treating or preventing a KDR-mediated disease with a KDR inhibitor, which method comprises administering to a patient in need of such a treatment a therapeutically

effective amount of a compound of formula I, or a pharmaceutical composition thereof.

The terms "KDR-mediated disease" or "KDR-mediated condition", as used herein, mean any disease state or other deleterious condition in which KDR is known to play a role. The terms "KDR-mediated disease" or "KDR-mediated condition" also mean those diseases or conditions that are alleviated by treatment with an KDR inhibitor. KDR-mediated diseases or conditions include, but are not limited to, carcinoma of the lung, breast, gastrointestinal tract, kidney, bladder, ovary and endometrium, intracranial tumors including glioblastoma multiforme, sporadic capillary hemangioblastoma, hematological malignancies, including T cell lymphoma, acute lymphoblastic leukemia, Burkitt's lymphoma and promyelocytic leukemia, age-related macular degeneration, herpetic ocular disease, rheumatoid arthritis, cerebral ischemia and endometriosis. The association of KDR with various diseases has been described for example in Ferrara, *Endocrine Reviews* 2004, 25, 581-611.

Another aspect of the invention relates to inhibiting KDR activity in a biological sample or a patient, which method comprises administering to the patient a compound of formula I, or a composition comprising said compound.

The term "patient" includes human and veterinary subjects.

The term "biological sample", as used herein, includes, without limitation, cell cultures or extracts thereof; preparations of an enzyme suitable for *in vitro* assay; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

An amount effective to inhibit protein kinase, for example, Aurora A, is an amount that causes measurable inhibition of the kinase activity when compared to the activity of the enzyme in the absence of an inhibitor. Any method may be used to determine inhibition, such as, for example, the Biological Testing Examples described below.

The term "pharmaceutically acceptable carrier, adjuvant, or vehicle" refers to a non-toxic carrier, adjuvant, or vehicle that may be administered to a patient, together with a compound of this invention, and which does not destroy the pharmacological activity thereof.

Pharmaceutically acceptable carriers that may be used in these pharmaceutical compositions are generally known in the art. They include, but are

not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, solvents, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, silicates, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, oils, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

Pharmaceutically accepted vehicles can contain mixtures of more than one excipient in which the components and the ratios can be selected to optimize desired characteristics of the formulation including but not limited to shelf-life, stability, drug load, site of delivery, dissolution rate, self-emulsification, control of release rate and site of release, and metabolism.

The compositions of the present invention may be administered orally, parenterally, by inhalation, topically, rectally, nasally, buccally, vaginally, transdermally, or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally, subcutaneously, intraperitoneally or intravenously.

Formulations can be prepared by a variety of techniques known in the art. Examples of formulation techniques can be found in literature publications and in texts such as "Water-insoluble drug formulation", edited by Rong Liu, 2000, Interpharm Press.

Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose,

any bland fixed oil may be employed including synthetic mono-or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil
5 solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents which are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other surface-active emulsifying agents or bioavailability
10 enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

The pharmaceutical compositions of this invention may be prepared by techniques known in the art and may be orally administered in any orally
15 acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers commonly used include but are not limited to celluloses, lactose, or corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents or carriers include lactose and
20 dried cornstarch. When aqueous suspensions or solutions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

Alternatively, the pharmaceutical compositions of this invention may be administered in the form of suppositories for rectal administration. These can be
25 prepared using techniques known in the art including for example by mixing the agent with a suitable non-irritating excipient, which is solid at room temperature but liquid at rectal temperature, and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

The pharmaceutical compositions of this invention may also be
30 administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, the airways, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs using techniques known in the art. For example, topical application for the lower intestinal tract can be effected

in a rectal suppository formulation (see above) or in a suitable enema formulation.

Topically-transdermal patches may also be used.

For topical or transdermal applications, the pharmaceutical compositions may be formulated by techniques known in the art in a suitable ointment or base
5 containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this invention are well known in the art and include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax, and water. Alternatively, the pharmaceutical
10 compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

15 For ophthalmic use, the pharmaceutical compositions may be formulated by techniques known in the art as micronized or nanometer-sized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutical
20 compositions may be formulated in an ointment such as petrolatum.

The pharmaceutical compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as suspensions or solutions in saline, optionally employing
25 benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

The present invention can be used to treat inflammatory or immune mediated diseases in humans or animals, wherein the inflammatory or immune
30 mediated diseases include, but are not limited to, rheumatoid arthritis, osteoarthritis, ulcerative colitis, Crohn's disease, Mooren's ulcer, arthritis, sarcoidosis, inflammatory or immune mediated bowel disease, systemic lupus, Wegener's syndrome, Stevens-Johnson disease, Behcet's disease, pemphigoid, Lyme's disease, asthma or acquired immune deficiency syndrome.

The present invention can be used to treat infectious diseases in humans or animals, wherein the infectious diseases include, but are not limited to syphilis, a bacterial infection, a *Mycobacterial* infection, a bacterial ulcer, a fungal ulcer, a Herpes simplex infection, a Herpes zoster infection, a protozoan infection, a
5 Bartonellosis infection, or toxoplasmosis.

The present invention can be used to treat blood or blood vessel diseases in humans or animals, wherein the blood or blood vessel diseases include, but are not limited to, vein occlusion, artery occlusion, carotid obstructive disease, polyarteritis, atherosclerosis, Osler-Weber-Rendu disease, sickle cell anemia,
10 leukemia, acute or chronic neoplastic disease of the bone marrow, hemangiomas, hereditary hemorrhagic telangiectasia, disease of the bone marrow, anemia, impaired blood clotting or enlargement of the lymph nodes, liver, or spleen. The present invention can also be used to treat chronic neoplastic disease of the bone marrow, wherein those diseases include, but are not limited to, multiple myeloma
15 and myelo dysplastic syndrome.

The present invention can be used to treat skin conditions in a humans or an animals, wherein the skin conditions include, but are not limited to, abnormal wound healing, acne rosacea, chemical burns of the skin, dermatitis or psoriasis.

In addition, the invention can be used to treat a variety of post-menopausal
20 symptoms, osteoporosis, cardiovascular disease, myocardial angiogenesis, plaque neovascularization, hemophiliac joints, angiofibroma, wound granulation, intestinal adhesions, scleroderma, hypertrophic scars; *i.e.*, keloids. They are also useful in the treatment of diseases that have angiogenesis as a pathologic consequence, such as cat scratch disease, and *Helicobacter pylori* ulcers. The
25 invention can also be used to treat Alzheimer's disease, to reduce the incidence of stroke, and as an alternative to prior estrogen replacement therapies. The compounds of the present invention can work by estrogenic and non-estrogenic biochemical pathways.

Additionally, the compounds of the present invention can be used to treat
30 endometriosis. Endometriosis is the abnormal growth of endometrial cells; the same cells that line the uterus that are shed monthly in the menstrual process. Wayward endometrial cells can position themselves in the lower abdomen on areas such as the cul-de-sac, the recto-vaginal septum, the stomach, the fallopian tubes, the ovaries, and the bladder. During menstruation, the normal uterine

lining is sloughed off and expelled through the vagina, but transplanted endometrial tissue has no means of exiting the body; instead the endometrial tissue and cells adhere and grow where positioned. The results are internal bleeding, inflammation, and scarring. One of the serious consequences of endometrial scarring is infertility. The endometrial growths are generally not malignant or cancerous. Among other complications, the growths can rupture and can spread the endometriosis to new areas of the lower abdomen. Endometriosis is a progressive disease. The growths or lesions are first seen as clear vesicles, then become red, and finally progress to black lesions over a period of seven to ten years.

In addition, the compounds of this invention, can be formulated to increase the bioavailability of the compound by methods well known to those of ordinary skill in the art. Methods of formulating the compounds of this invention and examples of formulations are described in "Water-Insoluble Drug Formulation" Rong Liu editor, CRC Press LLC, 2000, which may be referred to for further details.

Formulations contemplated as part of this invention include nanoparticles formulations made by controlled precipitation methods and by methods disclosed in U.S. Patent Publication No. 2004/0033267 published February 19, 2004 and which may be referred to for further details. Common excipients for nanoparticles known in the art include water, surface active agents such as sugar polymers (modified celluloses) and detergents, and also optionally preservatives such as benzalkonium salts, benzoic acid or salts thereof, or parabens. By forming nanoparticles, the compositions disclosed herein have increased bioavailability. Preferably, the particles of the compounds of the present invention have an effective average particle size of less than about 2 microns, less than about 1900 nm, less than about 1800 nm, less than about 1700 nm, less than about 1600 nm, less than about 1500 nm, less than about 1400 nm, less than about 1300 nm, less than about 1200 nm, less than about 1100 nm, less than about 1000 nm, less than about 900 nm, less than about 800 nm, less than about 700 nm, less than about 600 nm, less than about 500 nm, less than about 400 nm, less than about 300 nm, less than about 250 nm, less than about 200 nm, less than about 150 nm, less than about 100 nm, less than about 75 nm, or less than about 50 nm, as measured by light-scattering methods, microscopy, or other appropriate methods

well known to those of ordinary skill in the art. Nanoparticle preparations can be incorporated into many of the formulation approaches described here, including for example suspensions or creams or ointments for topical or transdermal administration, suspensions or powders or tablets or capsules or pellets for suppositories or for oral administration, suspensions for sterile injectable formulations, and polymer formulations.

The compounds that make up this invention can be incorporated into biodegradable or non-biodegradable polymers allowing for sustained release of the compound. The polymers can be implanted so that the drug is delivered parenterally throughout the body or the polymers with the compounds that make up this invention can be implanted in the vicinity of the tumor. A review of polymers in controlled drug delivery can be found for example in "Biodegradable Polymers as Drug Delivery Systems, Chasin M and Langer R (eds), New York, Marcel Dekker, 1990, which may be referred to for further details.

Another review can be found in "Handbook of Biodegradable Polymers", D. Weseman, J. Kost and A. Domb, Taylor & Francis, 1998, which may be referred to for further details.

A "pharmaceutically acceptable derivative or prodrug" means any pharmaceutically acceptable salt, ester, amide, salt of an ester or amide, or other derivative of a compound of this invention which, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof. Particularly favored derivatives or prodrugs are those that increase the bioavailability of the compounds of this invention when such compounds are administered to a patient (e.g., by allowing an orally administered compound to be more readily absorbed into the blood) or which enhance delivery of the parent compound to a biological compartment (e.g., the brain or lymphatic system) relative to the parent species.

Pharmaceutically acceptable prodrugs of the compounds of this invention include, without limitation, the following derivatives of the present compounds: esters, amino acid esters, amino acid amides, phosphate esters, metal salts, sulfonate esters, carbamates, and amides.

Pharmaceutically acceptable salts of the compounds of this invention include those derived from pharmaceutically acceptable inorganic and organic acids and bases. Examples of suitable acid salts include acetate, adipate, alginate,

aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptanoate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-
5 hydroxyethanesulfonate, lactate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, palmoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, salicylate, succinate, sulfate, tartrate, thiocyanate, tosylate and undecanoate. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in
10 the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts.

Salts derived from appropriate bases include alkali metal (e.g., sodium and potassium), alkaline earth metal (e.g., magnesium), ammonium and $N^+(C_{1-4} \text{ alkyl})_4$ salts. This invention also envisions the quaternization of any basic nitrogen-
15 containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization.

The amount of the protein kinase inhibitor that may be combined with the carrier materials to produce a single dosage form will vary depending upon the patient treated and the particular mode of administration. Preferably, the
20 compositions should be formulated so that a dosage of between 0.01-100 mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the
25 activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of the inhibitor will also depend upon the particular compound in the composition.

30 Depending upon the particular protein kinase-mediated condition to be treated or prevented, additional therapeutic agents, which are normally administered to treat or prevent that condition, may be administered together with the inhibitors of this invention. For example, in the treatment of cancer, other kinase inhibitors, chemotherapeutic agents, anti-angiogenesis agents, anti-nausea


agents, colony-stimulating factors, or other anti-proliferative agents may be combined with the present compounds to treat cancer as is known in the art. These agents include, without limitation, bevacizumab, adriamycin, dexamethasone, vincristine, cyclophosphamide, fluorouracil, topotecan, taxanes, interferons, and
5 platinum derivatives.

Other examples of agents the inhibitors of this invention may also be combined with include, without limitation, agents for treating diabetes such as insulin or insulin analogues, in injectable or inhalation form, glitazones, alpha glucosidase inhibitors, biguanides, insulin sensitizers, and sulfonyl ureas; anti-
10 inflammatory agents such as corticosteroids, TNF blockers, IL-1 RA, azathioprine, cyclophosphamide, and sulfasalazine; immunomodulatory and immunosuppressive agents such as cyclosporin, tacrolimus, rapamycin, mycophenolate mofetil, interferons, corticosteroids, cyclophosphamide, azathioprine, and sulfasalazine; neurotrophic factors such as acetylcholinesterase
15 inhibitors, MAO inhibitors, interferons, anti-convulsants, ion channel blockers, riluzole, and anti-Parkinsonian agents; agents for treating cardiovascular disease such as beta-blockers, ACE inhibitors, diuretics, nitrates, calcium channel blockers, and statins; agents for treating liver disease such as corticosteroids, cholestyramine, interferons, and anti-viral agents; agents for treating blood
20 disorders such as corticosteroids, anti-leukemic agents, and growth factors; therapeutic antibodies such as bevacizumab; and agents for treating immunodeficiency disorders such as gamma globulin.


Those additional agents may be administered separately from the protein kinase inhibitor-containing composition, or as part of a multiple dosage regimen.
25 Alternatively, those agents may be part of a single dosage form, mixed together with the protein kinase inhibitor of this invention in a single composition.

Compounds of this invention may exist in alternative tautomeric forms, for example as in tautomers shown below. Unless otherwise indicated, the representation of any tautomer is meant to include any other tautomers.

30



In one embodiment, the present invention provides a compound of formula
 5 I or a pharmaceutically acceptable derivative or prodrug thereof,



Formula I

wherein:

10 R^x and R^y are independently selected from the group consisting of $-\text{T}-\text{R}^3$ and $-\text{L}-\text{Z}-\text{R}^3$;

Q' is selected from the group consisting of $-\text{CR}^{6''}=\text{CR}^{6''}-$ and $\text{---}\equiv\text{---}$, wherein said $-\text{CR}^{6''}=\text{CR}^{6''}-$ may be a *cis* or *trans* double bond or a mixture thereof;

15 R^1 is $-\text{T}-(\text{Ring D})$;

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from the group consisting of aryl, heteroaryl, heterocyclyl, and carbocyclyl, said heteroaryl or heterocyclyl ring having 1-4 ring heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, wherein each
 20 substitutable ring carbon of Ring D is independently substituted by oxo, $-\text{T}-\text{R}^5$, or $-\text{V}-\text{Z}-\text{R}^5$, and each substitutable ring nitrogen of Ring D is independently substituted by $-\text{R}^4$;

T is a valence bond or $-(\text{C}(\text{R}^6)_2)-\text{A}-$;

A is a valence bond or a $\text{C}_1\text{-C}_3$ alkylidene chain wherein a methylene unit
 25 of said C_{1-3} alkylidene chain is optionally replaced by $-\text{O}-$, $-\text{S}-$, $-\text{N}(\text{R}^4)-$, $-\text{CO}-$, -

CONH-, -NHCO-, -SO₂-, -SO₂NH-, -NHSO₂-, -CO₂-, -OC(O)-, -OC(O)NH-, or -NHCO₂-;

Z is a C₁₋₄ alkylidene chain;

L is selected from the group consisting of -O-, -S-, -SO-, -SO₂-, -N(R⁶)SO₂-,
 5 -, -SO₂N(R⁶)-, -N(R⁶)-, -CO-, -CO₂-, -N(R⁶)CO-, -N(R⁶)C(O)O-, -
 N(R⁶)CON(R⁶)-, -N(R⁶)SO₂N(R⁶)-, -N(R⁶)N(R⁶)-, -C(O)N(R⁶)-, -OC(O)N(R⁶)-, -
 C(R⁶)₂O-, -C(R⁶)₂-, -C(R⁶)₂SO-, -C(R⁶)₂SO₂-, -C(R⁶)₂SO₂N(R⁶)-, -C(R⁶)₂N(R⁶)-, -
 C(R⁶)₂N(R⁶)C(O)-, -C(R⁶)₂N(R⁶)C(O)O-, -C(R⁶)=NN(R⁶)-, -C(R⁶)=N-O-, -
 C(R⁶)₂N(R⁶)N(R⁶)-, -C(R⁶)₂N(R⁶)SO₂N(R⁶)-, and -C(R⁶)₂N(R⁶)CON(R⁶)-;

10 R² and R^{2'} are independently selected from the group consisting of -R and
 -T-W-R⁶, or R² and R^{2'} taken together with their intervening atoms form a fused,
 5-8 membered, unsaturated or partially unsaturated ring having 0-3 ring
 heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur,
 wherein each substitutable ring carbon of said fused ring formed by R² and R^{2'} is
 15 independently substituted by halo, oxo, -CN, -NO₂, R⁷, or -V-R⁶, and each
 substitutable ring nitrogen of said ring formed by R² and R^{2'} is independently
 substituted by -R⁴;

R³ is selected from the group consisting of -R, -halo, -OR, -C(=O)R, -
 CO₂R, -COCOR, -COCH₂COR, -NO₂, -CN, -S(O)R, -S(O)₂R, -SR, -N(R⁴)₂, -
 20 CON(R⁷)₂, -SO₂N(R⁷)₂, -OC(=O)R, -N(R⁷)COR, -N(R⁷)CO₂(C₁₋₆ aliphatic), -
 N(R⁴)N(R⁴)₂, -C=NN(R⁴)₂, -C=N-OR, -N(R⁷)CON(R⁷)₂, -N(R⁷)SO₂N(R⁷)₂, -
 N(R⁴)SO₂R, and -OC(=O)N(R)₂;

each R is independently hydrogen or an optionally substituted group
 selected from the group consisting of C₁₋₆ aliphatic, C₆₋₁₀ aryl, a heteroaryl ring
 25 having 5-10 ring atoms, and a heterocyclyl ring having 5-10 ring atoms;

each R⁴ is independently selected from the group consisting of -R⁷, -COR⁷,
 -CO₂(optionally substituted C₁₋₆ aliphatic), -CON(R⁷)₂, and -SO₂R⁷;

each R⁵ is independently selected from the group consisting of -R, halo, -
 OR, -C(=O)R, -CO₂R, -COCOR, -NO₂, -CN, -S(O)R, -SO₂R, -SR, -N(R⁴)₂, -
 30 CON(R⁴)₂, -SO₂N(R⁴)₂, -OC(=O)R, -N(R⁴)COR, -N(R⁴)CO₂ (optionally
 substituted C₁₋₆ aliphatic), -N(R⁴)N(R⁴)₂, -C=NN(R⁴)₂, -C=N-OR, -
 N(R⁴)CON(R⁴)₂, -N(R⁴)SO₂N(R⁴)₂, -N(R⁴)SO₂R, and -OC(=O)N(R⁴)₂;

V is selected from the group consisting of -O-, -S-, -SO-, -SO₂-, -
 N(R⁶)SO₂-, -SO₂N(R⁶)-, -N(R⁶)-, -CO-, -CO₂-, -N(R⁶)CO-, -N(R⁶)C(O)O-, -

$N(R^6)CON(R^6)-$, $-N(R^6)SO_2N(R^6)-$, $-N(R^6)N(R^6)-$, $-C(O)N(R^6)-$, $-OC(O)N(R^6)-$, $-C(R^6)_2O-$, $-C(R^6)_2S-$, $-C(R^6)_2SO-$, $-C(R^6)_2SO_2-$, $-C(R^6)_2SO_2N(R^6)-$, $-C(R^6)_2N(R^6)-$, $-C(R^6)_2N(R^6)C(O)-$, $-C(R^6)_2N(R^6)C(O)O-$, $-C(R^6)=NN(R^6)-$, $-C(R^6)=N-O-$, $-C(R^6)_2N(R^6)N(R^6)-$, $-C(R^6)_2N(R^6)SO_2N(R^6)-$, and $-C(R^6)_2N(R^6)CON(R^6)-$;

5 W is selected from the group consisting of $-C(R^6)_2O-$, $-C(R^6)_2S-$, $-C(R^6)_2SO-$, $-C(R^6)_2SO_2-$, $-C(R^6)_2SO_2N(R^6)-$, $-C(R^6)_2N(R^6)-$, $-CO-$, $-CO_2-$, $-C(R^6)OC(O)-$, $-C(R^6)OC(O)N(R^6)-$, $-C(R^6)_2N(R^6)CO-$, $-C(R^6)_2N(R^6)C(O)O-$, $-C(R^6)=NN(R^6)-$, $-C(R^6)=N-O-$, $-C(R^6)_2N(R^6)N(R^6)-$, $-C(R^6)_2N(R^6)SO_2N(R^6)-$, $-C(R^6)_2N(R^6)CON(R^6)-$, and $-CON(R^6)-$;


10 each R^6 is independently selected from the group consisting of hydrogen and an optionally substituted C_{1-4} aliphatic group, or two R^6 groups on the same nitrogen atom may be taken together with the nitrogen atom to form a 5-6 membered heterocyclyl or heteroaryl ring;

each R^6 is independently selected from the group consisting of hydrogen
15 and a C_{1-4} aliphatic group, or two R^6 on the same carbon atom are taken together to form a 3-6 membered carbocyclic ring;

each $R^{6''}$ is independently selected from the group consisting of hydrogen,
a C_{1-4} aliphatic group, halogen, optionally substituted aryl, and optionally
substituted heteroaryl, or two $R^{6''}$ on adjacent carbon atoms are taken together to
20 form a 5-7 membered carbocyclic or heterocyclic ring, or one of the $R^{6''}$ can be
taken together with a substituent on ring D to form a fused bicyclic carbocyclic or
heterocyclic ring; and

each R^7 is independently selected from the group consisting of hydrogen
and an optionally substituted C_{1-6} aliphatic group, or two R^7 on the same nitrogen
25 are taken together with the nitrogen to form a 5-8 membered heterocyclyl or
heteroaryl ring.

In one embodiment, the present invention provides a compound of formula
Ia or a pharmaceutically acceptable derivative or prodrug thereof,




Ia

wherein R^2 , $R^{2'}$, R^x , R^y , and R^1 is defined as in formula I; Q^a is *cis* or *trans* $-CR^{6''}=CR^{6''}-$ or a mixture thereof; and each $R^{6''}$ is independently selected from hydrogen, methyl, halogen, optionally substituted aryl, and optionally substituted heteroaryl.

5 In some embodiments, the invention provides compounds of formula Ia wherein Q^a is *trans* $-CR^{6''}=CR^{6''}-$.

In another embodiment, the present invention provides a compound of formula Ib or a pharmaceutically acceptable derivative or prodrug thereof,



Ib

10 wherein R^2 , $R^{2'}$, R^x , R^y , and R^1 is defined as in formula I, and Q^b is $\text{---}\equiv\text{---}$.


Preferred R^x groups in the compounds of formula I, Ia, Ib, II, IIa, and IIb include hydrogen, alkyl, amino, nitro, alkyl- or dialkylamino, acetamido, or a C_{1-4} aliphatic group such as methyl, ethyl cyclopropyl, or isopropyl, and most preferably hydrogen and amino.

Preferred R^y groups in the compounds of formula I, Ia, Ib, II, IIa, and IIb include $-T-R^3$ or $-L-Z-R^3$ wherein T is a valence bond or a alkyl (1-6 carbons in length, branched or unbranched) or alkene (1-6 carbons in length, branched or unbranched), L is $-O-$, $-S-$, $-C(R)_2O-$, $-CO-$, $C(O)N(R^6)-$, or $-N(R^4)-$, and R^3 is $-R$, $-N(R^4)_2$, or OR. Preferred R^y groups include 5-6 membered heteroaryl or non-aromatic heterocyclic rings, such as 2-pyridyl, 3-pyrididyl, 4-pyridyl, pyrrolidinyl, piperidinyl, morpholinyl, hydroxypiperidinyl, N-(4-hydroxypiperidin)-yl, O-(4-piperidinyl), piperazinyl, alkylpiperazinyl, or 4-methylpiperazinyl, N-acetylpiperiziny, N-alkylcarboxamidpiperiziny, N-(methylsulfone)piperiziny, thiophene, furan, tetrahydrofuran, cyclo[2.2.1]heptenyl; C_{1-6} aliphatic, such as methyl, ethyl, cyclopropyl, isopropyl, or t-butyl; alkoxyalkylamino such as methoxyethylamino; alkoxyalkyl such as methoxymethyl or methoxyethyl; amino, alkyl- or dialkylamino such as ethylamino or dimethylamino; alkyl- or dialkylaminoalkoxy such as dimethylaminopropoxy; acetamido; alkoxycarbonyl; alkyl- and

dialkylamidocarbonyl; and optionally substituted phenyl such as phenyl or halo-substituted phenyl. For amine nitrogens, the N can be in the free base form, a pharmaceutically acceptable salt or the quaternary salt. This invention envisions that R³ can be attached to L or T through either the heteroatom or any ring atom
5 where there is a hydrogen available for ring attachment.

R² and R^{2'} may be taken together to form a fused ring, thus providing a bicyclic ring system containing a pyrazole ring. Preferred fused rings include benzo, pyrido, pyrimido, a partially unsaturated 6-membered carbocyclo ring, wherein said fused ring is optionally substituted. Fused 5-membered rings are also
10 envisioned and include but are not limited to pyrrolo, tetrahydrofuran, tetrahydrothiofuran imidazolidine and pyrazolidine. These are exemplified in the following formula I compounds having a pyrazole-containing bicyclic ring system, but also apply to compounds of formula Ia and Ib:

15



Preferred substituents on the $R^2/R^{2'}$ fused rings include one or more of the following: -halo, $-N(R^4)_2$, $-C_{1-3}$ alkyl, $-C_{1-3}$ haloalkyl, $-NO_2$, $-O(C_{1-3}$ alkyl), $-CO_2(C_{1-3}$ alkyl), $-CN$, $-SO_2(C_{1-3}$ alkyl), $-SO_2NH_2$, $-OC(O)NH_2$, $-NH_2SO_2(C_{1-3}$

alkyl), -NHC(O)(C₁₋₃ alkyl), -C(O)NH₂, and -CO(C₁₋₃ alkyl), wherein the (C₁₋₃ alkyl) is most preferably methyl.

When the pyrazole ring system is monocyclic, preferred R² groups include hydrogen, C₁₋₄ aliphatic, alkoxy carbonyl, (un)substituted phenyl, hydroxyalkyl, alkoxyalkyl, aminocarbonyl, mono- or dialkylaminocarbonyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, phenylaminocarbonyl, and (N-heterocyclyl)carbonyl. Examples of such preferred R² substituents include methyl, cyclopropyl, ethyl, isopropyl, propyl, t-butyl, cyclopentyl, phenyl, CO₂H, CO₂CH₃, CH₂OH, CH₂OCH₃, CH₂CH₂CH₂OH, CH₂CH₂CH₂OCH₃, CH₂CH₂CH₂OCH₂Ph, CH₂CH₂CH₂NH₂, CH₂CH₂CH₂NHCOOC(CH₃)₃, CONHCH(CH₃)₂, CONHCH₂CH=CH₂, CONHCH₂CH₂OCH₃, CONHCH₂Ph, CONH(cyclohexyl), CON(Et)₂, CON(CH₃)CH₂Ph, CONH(n-C₃H₇), CON(Et)CH₂CH₂CH₃, CONHCH₂CH(CH₃)₂, CON(n-C₃H₇)₂, CO(3-methoxymethylpyrrolidin-1-yl), CONH(3-tolyl), CONH(4-tolyl), CONHCH₃, CO(morpholin-1-yl), CO(4-methylpiperazin-1-yl), CONHCH₂CH₂OH, CONH₂, and CO(piperidin-1-yl). A preferred R² group is hydrogen.

When Ring D of formula I, Ia, or Ib is monocyclic, preferred Ring D groups include optionally substituted phenyl, pyridyl, pyridazinyl, pyrimidinyl, and pyrazinyl.

When Ring D of formula I, Ia, or Ib is bicyclic, preferred optionally substituted bicyclic Ring D groups include naphthyl, tetrahydronaphthyl, indanyl, benzimidazolyl, quinolinyl, indolyl, isoindolyl, indolinyl, benzo[b]furyl, benzo[b]thiophenyl, indazolyl, benzothiazolyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxazolinyl, 1,8-naphthyridinyl and isoquinolinyl.

On Ring D of formula I, Ia, or Ib, preferred T-R⁵ or V-Z—R⁵ substituents include -halo, -CN, -NO₂, -N(R⁴)₂, optionally substituted C₁₋₆ aliphatic group, -OR, -C(O)R, -CO₂R, -CONH(R⁴), -N(R⁴)COR, -N(R⁴)CO₂R, -SO₂N(R⁴)₂, -N(R⁴)SO₂R, -N(R⁶)COCH₂N(R⁴)₂, -N(R⁶)COCH₂CH₂N(R⁴)₂, and -N(R⁶)COCH₂CH₂CH₂N(R⁴)₂, wherein R is selected from hydrogen, C₁₋₆ aliphatic, phenyl, a 5-6 membered heteroaryl ring, or a 5-6 membered heterocyclic ring. More preferred R⁵ substituents include -Cl, -Br, -F, -CN, -CF₃, -COOH, -CONHMe, -CONHEt, -NH₂, -NHAc, -NHSO₂Me, -NHSO₂Et, -NHSO₂(n-propyl), -NHSO₂(isopropyl), -NHCOEt, -NHCOCH₂NHCH₃, -NHCOCH₂N(CO₂

t-Bu)CH₃, -NHCOCH₂N(CH₃)₂, -NHCOCH₂CH₂ N(CH₃)₂, -NHCOCH₂CH₂CH₂N(CH₃)₂, -NHCO(cyclopropyl), -NHCO(isobutyl), -NHCOCH₂(morpholin-4-yl), -NHCOCH₂CH₂(morpholin-4-yl), -NHCOCH₂CH₂CH₂(morpholin-4-yl), -NHCO₂(t-butyl), -NH(C₁₋₄ aliphatic) such as -NHMe, -N(C₁₋₄ aliphatic)₂ such as -NMe₂, OH, -O(C₁₋₄ aliphatic) such as -OMe, C₁₋₄ aliphatic such as methyl, ethyl, cyclopropyl, isopropyl, or t-butyl, and -CO₂ (C₁₋₄ aliphatic).

Preferred formula I compounds have one, two, three, four, five, or all of the features selected from the group consisting of:

- (a) R^x is hydrogen, nitro, amino, alkyl- or dialkylamino, acetamido, or a C₁₋₄ aliphatic group;
- (b) R^y is -T-R³ or -L-Z-R³, wherein T is a valence bond or -(C(R⁶)₂)- and R³ is -R, -N(R⁴)₂, -OR, or -CO₂R;
- (c) R¹ is -T-(Ring D), wherein T is a valence bond or -(C(R⁶)₂)-;
- (d) Ring D is an optionally substituted 5-7 membered monocyclic or an 8-10 membered bicyclic aryl or heteroaryl ring;
- (e) R² is -R or -T-W-R⁶ and R^{2'} is hydrogen, or R² and R^{2'} are taken together to form an optionally substituted benzo ring; and
- (f) each R^{6"} is independently hydrogen, a C₁₋₄ aliphatic group, halogen, optionally substituted aryl, or optionally substituted heteroaryl.

More preferred formula I compounds have one, two, three, four, five, six or all of the features selected from the group consisting of:

- (a) R^y is -T-R³ or -L-Z-R³ wherein T is a valence bond or -(C(R⁶)₂)- and R³ is -R, -OR, -N(R⁴)₂ or -CO₂R, wherein R is hydrogen, C₁₋₆ aliphatic, 5-6 membered heterocyclyl, 6-membered aryl, or 5-6 membered heteroaryl;
- (b) R¹ is -T-(Ring D), wherein T is a valence bond or -(C(R⁶)₂)-;
- (c) Ring D is an optionally substituted 5-6 membered monocyclic or an 8-10 membered bicyclic aryl or heteroaryl ring;
- (d) R² is -R and R^{2'} is hydrogen, wherein -R is independently hydrogen or an optionally substituted group selected from the group consisting of C₁₋₆ aliphatic, C₆₋₁₀ aryl, a heteroaryl ring having 5-10 ring atoms, and a heterocyclyl ring having 5-10 ring atoms;
- (e) L is -O-, -S-, -N(R⁴)-, or -C(O)N(R⁶)-;
- (f) Q' is *trans* -CR^{6"}=CR^{6"}- or $\text{---}\equiv\text{---}$; and

(g) each $R^{6''}$ is independently hydrogen, methyl, halogen, optionally substituted aryl, or optionally substituted heteroaryl.

Even more preferred compounds of formula I have one, two, three, four, five, or all of the features selected from the group consisting of:

- 5 (a) R^x is hydrogen, methyl, ethyl, propyl, cyclopropyl, isopropyl, amino, dimethylamino, methylamino, nitro, or acetamido;
- (b) R^y is selected from 2-pyridyl, 4-pyridyl, pyrrolidinyl, piperidinyl, morpholinyl, hydroxypiperidinyl, N-(4-hydroxypiperidin)-yl, O-(4-piperidinyl), piperazinyl, alkylpiperazinyl, 4-alkylpiperazinyl, methyl, ethyl, cyclopropyl, isopropyl, t-butyl, alkoxyalkylamino, alkoxyalkyl, alkyl- or dialkylamino, alkyl- or dialkylaminoalkoxy, acetamido, optionally substituted phenyl, methoxymethyl, -CO₂R, and -C(O)N(R⁶)ZR;
- 10 (c) R^1 is -T-(Ring D), wherein T is a valence bond and Ring D is a 5-6 membered aryl or heteroaryl ring, wherein Ring D is optionally substituted with one to two groups selected from -halo, CF₃, -CN, -NO₂, -N(R⁴)₂, optionally substituted C₁₋₆ aliphatic group, -OR, -CO₂R, -CONH(R⁴), -N(R⁴)COR, -N(R⁴)SO₂R, -N(R⁴)COCH₂N(R⁶)₂, -N(R⁴)COCH₂CH₂N(R⁶)₂, and -N(R⁴)COCH₂CH₂CH₂N(R⁶)₂;
- (d) R^2 is hydrogen or a substituted or unsubstituted C₁₋₆ aliphatic;
- 20 (e) Q' is *trans* -CR^{6''}=CR^{6''}- or $\equiv$; and
- (f) each $R^{6''}$ is independently hydrogen, methyl, chloro, or fluoro.

Particularly more preferred compounds of formula I have one, two, three, four, five, or all of the features selected from the group consisting of:

- (a) R^x is hydrogen or amino;
- 25 (b) R^y is selected from 2-pyridyl, 4-pyridyl, pyrrolidinyl, piperidinyl, morpholinyl, hydroxypiperidinyl, N-(4-hydroxypiperidin)-yl, O-(4-piperidinyl), piperazinyl, alkylpiperazinyl, 4-alkylpiperazinyl, alkoxyalkylamino, alkoxyalkyl, alkyl- or dialkylamino, alkyl- or dialkylaminoalkoxy, optionally substituted phenyl, -CO₂R, and -C(O)NHZR;
- 30 (c) R^1 is T-(Ring D), wherein T is a valence bond and Ring D is a 5-6 membered aryl or heteroaryl ring, wherein Ring D is optionally substituted with one to two groups selected from -halogen, -CN, -CF₃, -NO₂, -N(R⁴)₂, optionally substituted C₁₋₆ aliphatic group, -OR, -CO₂R, -CONH(R⁴), -N(R⁴)COR, -

$N(R^4)SO_2R$, $-N(R^4)COCH_2N(R^6)_2$, $-N(R^4)COCH_2CH_2N(R^6)_2$, and $-N(R^4)COCH_2CH_2CH_2N(R^6)_2$;

(d) R^2 is hydrogen or a substituted or unsubstituted C_{1-6} aliphatic, and L is -O-, -S-, -NH- or $-C(O)NH-$;

5 (e) Q' is *trans* $-CR^{6''}=CR^{6''}-$; and

(f) each $R^{6''}$ is independently hydrogen or fluoro.

Preferred formula Ia or Ib compounds have one, two, three, four, five, or all of the features selected from the group consisting of:

(a) R^x is hydrogen, nitro, amino, alkyl- or dialkylamino, acetamido, or
10 a C_{1-4} aliphatic group;

(b) R^y is $-T-R^3$ or $-L-Z-R^3$, wherein T is a valence bond or $-(C(R^6)_2)-$ and R^3 is -R, $-N(R^4)_2$, -OR, or $-CO_2R$;

(c) R^1 is $-T-(\text{Ring D})$, wherein T is a valence bond or $-(C(R^6)_2)-$;

(d) Ring D is a 5-7 membered monocyclic or an 8-10 membered
15 bicyclic aryl or heteroaryl ring;

(e) R^2 is -R or $-T-W-R^6$ and $R^{2'}$ is hydrogen, or R^2 and $R^{2'}$ are taken together to form an optionally substituted benzo ring; and

(f) each $R^{6''}$ is independently hydrogen, a C_{1-4} aliphatic group, halogen, optionally substituted aryl, or optionally substituted heteroaryl.

20 More preferred formula Ia or Ib compounds have one, two, three, four, five, or all of the features selected from the group consisting of:

(a) R^y is $-T-R^3$ or $-L-Z-R^3$ wherein T is a valence bond or $-(C(R^6)_2)-$ and R^3 is -R, -OR, $-N(R^4)_2$ or $-CO_2R$, wherein R is hydrogen, C_{1-6} aliphatic, 5-6 membered heterocyclyl, 6-membered aryl, or 5-6 membered heteroaryl;

25 (b) R^1 is $-T-(\text{Ring D})$, wherein T is a valence bond or $-(C(R^6)_2)-$;

(c) Ring D is a 5-6 membered monocyclic or an 8-10 membered bicyclic aryl or heteroaryl ring;

(d) R^2 is -R and $R^{2'}$ is hydrogen, wherein -R is independently hydrogen or an optionally substituted group selected from the group consisting of C_{1-6}
30 aliphatic, C_{6-10} aryl, a heteroaryl ring having 5-10 ring atoms, and a heterocyclyl ring having 5-10 ring atoms;

(e) L is -O-, -S-, $-N(R^4)-$, or $-C(O)N(R^6)-$; and

(f) each $R^{6''}$ is independently hydrogen, methyl, halogen, optionally substituted aryl, or optionally substituted heteroaryl.

Even more preferred compounds of formula Ia or Ib have one, two, three, four, or all of the features selected from the group consisting of:


- (a) R^x is hydrogen, methyl, ethyl, propyl, cyclopropyl, isopropyl, amino, dimethylamino, methylamino, nitro, or acetamido;
- 5 (b) R^y is selected from 2-pyridyl, 4-pyridyl, pyrrolidinyl, piperidinyl, morpholinyl, hydroxypiperidinyl, N-(4-hydroxypiperidin)-yl, O-(4-piperidinyl), piperazinyl, alkylpiperazinyl, 4-alkylpiperazinyl, methyl, ethyl, cyclopropyl, isopropyl, t-butyl, alkoxyalkylamino, alkoxyalkyl, alkyl- or dialkylamino, alkyl- or dialkylaminoalkoxy, acetamido, optionally substituted phenyl, methoxymethyl, 10 $-\text{CO}_2\text{R}$, and $-\text{C}(\text{O})\text{N}(\text{R}^6)\text{ZR}$;
- (c) R^1 is $-\text{T}-(\text{Ring D})$, wherein T is a valence bond and Ring D is a 5-6 membered aryl or heteroaryl ring, wherein Ring D is optionally substituted with one to two groups selected from $-\text{halo}$, CF_3 , $-\text{CN}$, $-\text{NO}_2$, $-\text{N}(\text{R}^4)_2$, optionally substituted C_{1-6} aliphatic group, $-\text{OR}$, $-\text{CO}_2\text{R}$, $-\text{CONH}(\text{R}^4)$, $-\text{N}(\text{R}^4)\text{COR}$, 15 $\text{N}(\text{R}^4)\text{SO}_2\text{R}$, $-\text{N}(\text{R}^4)\text{COCH}_2\text{N}(\text{R}^6)_2$, $-\text{N}(\text{R}^4)\text{COCH}_2\text{CH}_2\text{N}(\text{R}^6)_2$, and $-\text{N}(\text{R}^4)\text{COCH}_2\text{CH}_2\text{CH}_2\text{N}(\text{R}^6)_2$;
- (d) R^2 is hydrogen or a substituted or unsubstituted C_{1-6} aliphatic; and
- (e) each $\text{R}^{6''}$ is independently hydrogen, methyl, chloro, or fluoro.

Particularly more preferred compounds of formula Ia or Ib have one, two, 20 three, four, or all of the features selected from the group consisting of:

- (a) R^x is hydrogen or amino;
- (b) R^y is selected from 2-pyridyl, 4-pyridyl, pyrrolidinyl, piperidinyl, morpholinyl, hydroxypiperidinyl, N-(4-hydroxypiperidin)-yl, O-(4-piperidinyl), piperazinyl, alkylpiperazinyl, 4-alkylpiperazinyl, alkoxyalkylamino, alkoxyalkyl, 25 alkyl- or dialkylamino, alkyl- or dialkylaminoalkoxy, optionally substituted phenyl, $-\text{CO}_2\text{R}$, and $-\text{C}(\text{O})\text{NHZR}$;
- (c) R^1 is $\text{T}-(\text{Ring D})$, wherein T is a valence bond and Ring D is a 5-6 membered aryl or heteroaryl ring, wherein Ring D is optionally substituted with one to two groups selected from $-\text{halogen}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{NO}_2$, $-\text{N}(\text{R}^4)_2$, optionally 30 substituted C_{1-6} aliphatic group, $-\text{OR}$, $-\text{CO}_2\text{R}$, $-\text{CONH}(\text{R}^4)$, $-\text{N}(\text{R}^4)\text{COR}$, $\text{N}(\text{R}^4)\text{SO}_2\text{R}$, $-\text{N}(\text{R}^4)\text{COCH}_2\text{N}(\text{R}^6)_2$, $-\text{N}(\text{R}^4)\text{COCH}_2\text{CH}_2\text{N}(\text{R}^6)_2$, and $-\text{N}(\text{R}^4)\text{COCH}_2\text{CH}_2\text{CH}_2\text{N}(\text{R}^6)_2$;
- (d) R^2 is hydrogen or a substituted or unsubstituted C_{1-6} aliphatic, and L is $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$ or $-\text{C}(\text{O})\text{NH}-$; and

(e) each $R^{6''}$ is independently hydrogen or fluoro.

In another embodiment, the present invention provides compounds of formula II or a pharmaceutically acceptable derivative or prodrug thereof:



wherein

R^x is hydrogen, nitro, amino, alkyl- or dialkylamino, acetamido, or a C_{1-4} aliphatic group;

10 R^y is 2-pyridyl, 4-pyridyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, 4-alkylpiperazinyl, alkoxyalkylamino, alkoxyalkyl, alkyl- or dialkylamino, alkyl- or dialkylaminoalkoxy, optionally substituted phenyl, $-CO_2R$, or $-C(O)NHZR$, wherein R is hydrogen, C_{1-6} aliphatic, 5-6 membered heterocyclyl, phenyl, or 5-6 membered heteroaryl and Z is a C_1-C_4 alkylidene chain;

15 Q' is $-CR^{6''}=CR^{6''}-$ or $-\equiv-$ wherein, the $-CR^{6''}=CR^{6''}-$ may be a *cis* or *trans* double bond; and each $R^{6''}$ is independently hydrogen, methyl, halogen, optionally substituted aryl, or optionally substituted heteroaryl; and

20 R^1 is phenyl optionally substituted with one to two groups selected from the group consisting of -halogen, -CN, $-CF_3$, $-NO_2$, $-N(R^4)_2$, optionally substituted C_{1-6} aliphatic group, -OR, $-CO_2R$, $-CONH(R^4)$, $-N(R^4)COR$, $-N(R^4)SO_2R$, $-N(R^4)COCH_2N(R^6)_2$, $-N(R^4)COCH_2CH_2N(R^6)_2$, and $-N(R^4)COCH_2CH_2CH_2N(R^6)_2$ wherein R, R^4 , and R^6 are defined as in formula I.

25 In some embodiments, the invention provides compounds of formula I or II wherein Q' is *trans* $-CH=CH-$. In other embodiments, Q' is $-\equiv-$.

In other embodiments of the compounds of formula I or II, R^x is hydrogen.


In still other embodiments, R^y is selected from 2-pyridyl, 4-pyridyl, pyrrolidinyl, piperidinyl, morpholinyl, hydroxypiperidinyl, N-(4-hydroxypiperidin)-yl, O-(4-piperidinyl), piperazinyl, alkylpiperazinyl, 4-

alkylpiperazinyl, alkoxyalkylamino, alkoxyalkyl, alkyl- or dialkylamino, alkyl- or dialkylaminoalkoxy, optionally substituted phenyl, $-\text{CO}_2\text{R}$, and $-\text{C}(\text{O})\text{NHZR}$.

In some embodiments, R^y is 4-alkylpiperazinyl. In other embodiments R^y is 4-methylpiperazinyl.

- 5 In some embodiments R^y is hydroxypiperidinyl. In other embodiments R^y is N-(4-hydroxypiperidin)-yl or O-(4-piperidinyl).

In one embodiment, the present invention provides compounds of formula IIa or a pharmaceutically acceptable derivative or prodrug thereof:




IIa

10 wherein R^x , R^y , and R^1 are as defined as in formula II; Q^a is *cis* or *trans* $-\text{CR}^{6''}=\text{CR}^{6''}-$ or a mixture thereof; and each $\text{R}^{6''}$ is independently selected from hydrogen, methyl, halogen, optionally substituted aryl, and optionally substituted heteroaryl.

- 15 In some embodiments of compound of formula IIa, Q^a is *cis* $-\text{CR}^{6''}=\text{CR}^{6''}-$.
In other embodiments, Q^a is *trans* $-\text{CR}^{6''}=\text{CR}^{6''}-$.

In one embodiment, the present invention provides compounds of formula IIa or a pharmaceutically acceptable derivative or prodrug thereof:




IIb

20 wherein R^x , R^y , and R^1 are as defined as in formula II and Q^b is $-\text{C}\equiv\text{C}-$.

In some embodiments, the invention provides compounds of formula I, Ia, Ib, II, IIa, or IIb wherein R^1 is selected from the following group:







where the line drawn through the side of the substituent indicates that the substituent can be joined to the linker at any ring atom where there is a hydrogen available for replacement.

- 5 In some embodiments of the compounds of formula I, Ia, Ib, II, IIa, or IIb, R^x is hydrogen.

In some embodiments of the compounds of formula I, Ia, Ib, II, IIa, or IIb, R^y is 4-methylpiperazinyl.

- 10 In some embodiments of the compounds of formula I, Ia, Ib, II, IIa, or IIb, R^y is N-(4-hydroxypiperidin)-yl or O-(4-piperidinyl).













In other embodiments of the compounds of formula I, Ia, Ib, II, IIa, or IIb, R^1 is phenyl.













In still other embodiments, the invention provides the compounds shown in Table 1, or a pharmaceutically acceptable salt, derivative or prodrug thereof.

15













Table 1













Example No.	Structure	Example No.	Structure
4		5	















6		7	
8		9	
10		11	
12		13	
14		15	
17		18	













19		20	
21		22	
23		24	
25		26	
29		30	
31		32	













34	 <chem>Xc1nc(C=Cc2ccccc2)n(c1Nc3cc[nH]3)X</chem> X = Cl, Br, I	37	 <chem>COC(=O)c1ccc(C=Cc2nc(Cl)c([N+](=O)[O-])n2Nc3cc[nH]3)cc1</chem>
38	 <chem>COC(=O)c1ccc(C=Cc2nc(NC3CCN(C)CC3)c([N+](=O)[O-])n2Nc4cc[nH]4)cc1</chem>	39	 <chem>COC(=O)c1ccc(C=Cc2nc(NC3CCN(C)CC3)c(N)n2Nc4cc[nH]4)cc1</chem>
40	 <chem>COc1ccc(C=Cc2nc(Nc3cc[nH]3)n(C4CCN(C)CC4)c2)cc1</chem>	41	 <chem>COc1ccc(C=Cc2nc(Nc3cc[nH]3)n(C4CCN(C)CC4)c2)cc1</chem>
42	 <chem>COc1ccc(C=Cc2nc(Nc3cc[nH]3)n(C4CCN(C)CC4)c2)cc1</chem>	43	 <chem>COc1ccc(C=Cc2nc(Nc3cc[nH]3)n(C4CCN(C)CC4)c2)cc1</chem>
44	 <chem>COc1ccc(C=Cc2nc(Nc3cc[nH]3)n(C4CCN(C)CC4)c2)cc1</chem>	45	 <chem>COc1ccc(C=Cc2nc(Nc3cc[nH]3)n(C4CCN(C)CC4)c2)cc1</chem>
46	 <chem>COc1ccc(C=Cc2nc(Nc3cc[nH]3)n(C4CCN(C)CC4)c2)cc1</chem>	47	 <chem>COc1ccc(C=Cc2nc(Nc3cc[nH]3)n(C4CCN(C)CC4)c2)cc1</chem>















48		49	
50		51	
52		53	
54		55	
56		57	
58		59	













60		61	
62		63	
64		65	
66		67	
68		69	
70		71	















72		73	
74		75	
76		77	
78		79	
80		81	
82		83	
84		85	













86		87	
88		89	
90		91	
92		93	
94		95	
96		97	

98		99	
100		101	
102		103	
104		105	
106		107	
108		109	


110		111	
112		113	
114		115	
116		117	
118		119	
120		121	
122		123	

124		125	
126		127	
128		129	
130		131	
132		133	
134		135	

136		137	
138		139	
140		141	
142		143	
144		145	
146		147	
148		149	

150		151	
152		153	
154		155	
156		157	
158		159	
160		161	

162		163	
164		165	
166		167	
168		169	
170		171	
172		173	
174		175	



In one embodiment, this invention provides a composition comprising a compound of formula I, and a pharmaceutically acceptable carrier. In some such embodiments, the composition is for treating or preventing a kinase mediated disorder.

In another embodiment, this invention relates to a method of treating or preventing a kinase mediated disease, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I, or a pharmaceutical composition thereof.

In some aspects of the aforementioned methods and compositions, the disorder is mediated by Aurora A, Aurora B, CDK-2, ERK-2, AKT, Src, Lck, Abl, cKit, Flt3, or KDR. In other aspects, the disorder is mediated by Aurora A, Src, Lck, Abl, cKit, Flt3, or KDR.

Another aspect of this invention relates to a method of inhibiting Aurora A activity in a patient, which method comprises administering to the patient a compound of formula I or a composition comprising said compound.

Another aspect of this invention relates to a method of treating or preventing a GSK-3-mediated disease with a GSK-3 inhibitor, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I or a pharmaceutical composition thereof.

One aspect of this invention relates to a method of enhancing glycogen synthesis and/or lowering blood levels of glucose in a patient in need thereof,

which method comprises administering to the patient a therapeutically effective amount of a compound of formula I, or a pharmaceutical composition thereof. This method is especially useful for diabetic patients. Another method relates to inhibiting the production of hyperphosphorylated Tau protein, which is useful in
5 halting or slowing the progression of Alzheimer's disease. Another method relates to inhibiting the phosphorylation of beta-catenin, which is useful for treating schizophrenia.

Another aspect of this invention relates to a method of inhibiting GSK-3 activity in a patient, which method comprises administering to the patient a
10 compound of formula I, or a composition comprising said compound.

Another aspect of this invention relates to a method of treating or preventing a Src-mediated disease with a Src inhibitor, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I or a pharmaceutical composition thereof.


15 Another aspect of the invention relates to inhibiting Src activity in a patient, which method comprises administering to the patient a compound of formula I, or a composition comprising said compound.

Another method relates to inhibiting Aurora A, GSK-3, or Src activity in a biological sample, which method comprises contacting the biological sample with
20 the Aurora A, GSK-3, or Src inhibitor of formula I, or a pharmaceutical composition thereof, in an amount effective to inhibit Aurora-2, GSK-3, or Src.

Each of the aforementioned methods directed to the inhibition of Aurora A, GSK-3, or Src, or the treatment of a disease alleviated thereby, is preferably carried out with a preferred compound of formula I, as described above.

25 The present invention also relates to the processes for preparing the compounds of the invention and to the synthetic intermediates useful in such process, as described below and in the Examples.

Scheme 1




In another aspect of this invention, preferred embodiments of Formula **1-K** can be synthesized as shown in Scheme 1 wherein the variable substituents are as described above and examples of which are indicated by Table 1 and $-\text{Ar}$ is an optionally substituted aryl or heteroaryl group as defined and examples of which are indicated by Table 1. Dihydroxypyrimidine **1-A** is dissolved in about one to 20 volumes of trifluoroacetic acid and treated with about one to five equivalents of nitric acid at about -20 to 30°C for about 30 minutes to 24 hours. The resulting nitropyrimidine **1-B** is treated with about one to 20 equivalents of an optionally substituted aromatic or heteroaromatic aldehyde **1-C** and about one to 20

equivalents of an organic base, preferably piperidine, at about 20 to 120 °C for about 30 minutes to 24 hours to provide a vinylpyrimidine **1-D**. A vinylpyrimidine **1-D** is treated with about two to 20 equivalents of phosphorous oxychloride and about two to five equivalents of a tertiary organic base, preferably diethylaniline, at about 0 to 200 °C for about 30 minutes to 24 hours to give a dichloropyrimidine **1-E**. A dichloropyrimidine **1-E** in a suitable solvent such as, but not limited to, tetrahydrofuran is treated with about one to 10 equivalents of a tertiary organic base such as, but not limited to, triethylamine and about one equivalent of an aminopyrazole **1-F** at about 0 to 65 °C for about 30 minutes to 24 hours to give a aminopyrimidine **1-G**. Subsequently, **1-G**, in a suitable solvent such as, but not limited to, tetrahydrofuran is treated with about one to 10 equivalents of a tertiary organic base such as, but not limited to, triethylamine and about one to five equivalents of a primary or secondary amine **1-H** at about 0 to 65 °C for about 30 minutes to 24 hours to give a diaminopyrimidine **1-J**. A diaminopyrimidines **1-J** is dissolved in a suitable solvent such as, but not limited to, methanol and added to about two to 10 equivalents of a suitable chemical reducing agent such as, but not limited to, tin (II) chloride or titanium (III) chloride in dilute hydrochloric acid at about 0 to 65 °C for about 30 minutes to 24 hours to give a triaminopyrimidine **1-K**.

In another aspect of this invention, preferred embodiments of Formula **2-F** are prepared as shown in Scheme 2 wherein the variable substituents are as described above and examples of which indicated by Table 1. A solution of about one to five equivalents of an aminopyrazole **2-B** in a suitable solvent such as, but not limited to, dioxane, is added to a solution of dichloropyrimidine **2-A** and about one to 10 equivalents of a suitable organic base, preferably 2,6-lutidine, in a suitable solvent such as, but not limited to, dioxane at from about 0 to 60 °C for about 15 minutes to 24 hours to give a chloropyrimidine **2-C**. A mixture of a chloropyrimidine **2-C**, about one to five equivalents of an inorganic base such as, but not limited to, sodium, potassium or cesium carbonate, about one to five equivalents of an optionally substituted *cis*- or *trans*-aryl- or heteroaryl-vinylboronic acid **2-D**, about 0.01 to 1 equivalents of a palladium catalyst such as, but not limited to, tetrakis(triphenylphosphine)-palladium in a suitable solvent such as, but not limited to, toluene or tetrahydrofuran at about 20 to 150 °C for about 30 minutes to 24 hours provides a vinylpyrimidine **2-E**. A solution of a

vinylpyrimidine **2-E** in a suitable solvent such as, but not limited to, dimethylsulfoxide, acetic acid or acetonitrile is treated with about two to 20 equivalents of a chemical reducing agent such as, but not limited to, zinc dust and ammonium formate, tin (II) chloride or titanium(III) chloride to give an aminopyrimidine **2-F**. The aminopyrimidine **2-F** will be either *cis* or *trans* depending upon whether the *cis* or *trans* isomer, respectively, of the aryl- or heteroaryl-vinylboronic acid is used in the reaction sequence.

Scheme 2




10

In another aspect of this invention, preferred embodiments **2-H** are prepared as shown in Scheme 2 whereby an aminopyrimidine **2-F** is treated with about one to 100 equivalents of an alcohol **2-G** alone or in a suitable solvent and about 0.1 to 5 equivalents of titanium(IV) isopropoxide or potassium cyanide at about 0 to 150 °C for about one to 48 hours.

15


In another aspect of this invention, preferred embodiments **2-K** are prepared as shown in Scheme 2 whereby an aminopyrimidine **2-F** is treated with about one to 20 equivalents of an amine **2-J** in a suitable solvent such as, but not limited to, dichloromethane, and about one to 20 equivalents of trimethylaluminum at 0 to 30 °C for about one to 48 hours.

Scheme 3



In another aspect of this invention, preferred embodiments of Formula **3-E** are prepared as shown in Scheme 3 wherein the variable substituents are as described above. A solution of about one to five equivalents of an aminopyrazole **3-B** in a suitable solvent such as, but not limited to, dioxane, is added to a solution of dichloropyrimidine **3-A** and about one to 10 equivalents of a suitable organic base, preferably 2,6-lutidine, in a suitable solvent such as, but not limited to, dioxane, at from about 0 to 60 °C for about 15 minutes to 24 hours to give a chloropyrimidine **3-C**. A mixture of a chloropyrimidine **3-C**, about one to five equivalents of an inorganic base such as, but not limited to, sodium, potassium or cesium carbonate, about one to five equivalents of an optionally substituted *cis*- or *trans*- aryl- or heteroaryl-vinylboronic acid **3-D**, about 0.01 to 1 equivalents of a palladium catalyst such as, but not limited to, tetrakis-(triphenyl-phosphine)-palladium in a suitable solvent such as, but not limited to, toluene or tetrahydrofuran, at about 20 to 150 °C for about 30 minutes to 24 hours provides a vinylpyrimidine **3-E**. The vinylpyrimidine **3-E** will be either *cis* or *trans* depending upon whether the *cis* or *trans* isomer, respectively, of the aryl- or heteroaryl-vinylboronic acid is used in the reaction.

Scheme 4




In another aspect of this invention, preferred embodiments **4-H** are prepared as shown in Scheme 4 wherein the variable substituents are as described above and examples are indicated by Table 1. A thiomethylpyrimidine **4-A** in a suitable solvent such as, but not limited to, dichloromethane is treated with about two to 10 equivalents of an oxidizing agent such as, but not limited to, 3-chlorobenzoylperoxide (mCPBA), at about 0 to 30 °C for about 30 minutes to 24 hours to give a sulfonylpyrimidine **4-B**. A sulfonylpyrimidine **4-B** in a suitable solvent such as, but not limited to, tetrahydrofuran is treated with about one to five equivalents of an optionally substituted aryl or heteroaryl-acetylenyl magnesium halide **4-C** at about -50 to 30 °C for about one to 24 hours to give a propargylpyrimidine **4-D**. A propargylpyrimidine **4-D** in a suitable solvent such as, but not limited to, dimethylacetamide is treated with about one to two equivalents of an aminopyrazole **4-E**, one to five equivalents of sodium iodide, and about one to five equivalents of a tertiary organic base such as, but not limited to, di-isopropylethylamine at about 0 to 140 °C for about one to 24 hours to give a monochloropyrimidine **4-F**. A monochloropyrimidine **4-F** in a suitable solvent

such as, but not limited to, dioxane is treated with one to five equivalents of a primary or secondary amine **4-G**, about 0.01 to one equivalents of 4-dimethylaminopyridine, and about one to five equivalents of a tertiary organic base such as, but not limited to, di-isopropylethylamine, at about 20 to 110 °C for about one to 24 hours to give an aminopyrimidine **4-H**.

In another aspect of this invention, preferred embodiments **4-J** are prepared as shown in Scheme 4 wherein an aminopyrimidine **4-H** in a suitable solvent such as, but not limited to, tetrahydrofuran, is treated with about 0.8 to 1.1 equivalents of lithium aluminum hydride at about -10 to 30 °C for about one to 24 hours to give a *trans*-styrylpyrimidine **4-J**.

In another aspect of this invention, preferred embodiments **4-K** are prepared as shown in Scheme 4 wherein an aminopyrimidine **4-H** in a suitable solvent such as, but not limited to, ethyl acetate or methanol is treated with about 0.1 to 1 weight equivalents of Lindlar's catalyst, about 0.1 to 2 weight equivalents of quinoline and hydrogen gas at about one atmosphere to give a *cis*-styrylpyrimidine **4-K**.


Scheme 5




In another aspect of this invention, preferred embodiments **5-D** and **5-E** are prepared as shown in Scheme 5 wherein the variable substituents are as described above and examples of which are indicated by Table 1. A
5 sulfonylpyrimidine **4-B** in a suitable solvent such as, but not limited to, tetrahydrofuran is treated with about one to two equivalents of an optionally substituted *cis*- or *trans*- 2-aryl- or heteroaryl-vinyl magnesium halide **5-A** at about -50 to 30 °C for about 30 minutes to 24 hours to give a dichloropyrimidine **5-B**. A dichloropyrimidine **5-B** in a suitable solvent such as, but not limited to,
10 dimethylacetamide is treated with about one to two equivalents of an aminopyrazole **4-E**, about one to five equivalents of sodium iodide and about one to five equivalents of a tertiary organic base such as, but not limited to, di-isopropylethylamine at about 20 to 140 °C for about one to 24 hours to give a monochloropyrimidine **5-C**. A monochloropyrimidine **5-C** in a suitable solvent
15 such as, but not limited to, dioxane is treated with about one to five equivalents of a primary or secondary amine **4-G**, 0.05 to 1 equivalents of 4-dimethylaminopyridine, and about one to five equivalents of a tertiary organic base such as, but not limited to, di-isopropylethylamine at about 20 to 110 °C for about one to 72 hours to give either styrylpyrimidine **5-D** or **5-E** depending upon
20 whether the vinyl magnesium halide **5-A** used in the reaction sequence is the *trans*- or *cis*- isomer, respectively.

Scheme 6


In another aspect of this invention, preferred embodiments **6-G** are prepared as shown in Scheme 6 wherein the variable substituents are as described above and examples of which are indicated in the accompanying figures. Selected
5 examples of **6-G** are indicated by Table 1.




R-X-H in Scheme 6 can for example be selected from the following structures:



The moiety denoted "Ar" in Scheme 6 can for example be selected from the following structures, where the line drawn through the side of the substituent indicates that the "Ar" substituent can be joined to the linker alkene or linker alkyne at any ring atom where there is a hydrogen available for replacement:





The product 6-G from Scheme 6 can be produced by selection of a nitrile with an Ar- substituent described above or selected from examples shown in the above figure. These nitriles are commercially available or can be produced by reactions known to one skilled in the art including for example by converting the appropriate aldehyde to the acrylonitrile using for examples but not limited to a Wittig reaction or related reactions such as the Peterson Olefination reaction. Generally the desired product will also be made by selecting an appropriate Pz-NH₂, chosen for example based on the descriptions above or selected from the figure above showing examples of Pz-NH₂, and selecting an R-X-H, chosen for example based on the descriptions above or selected from the figure above showing examples of R-X-H. Examples of some possible products are indicated in Table 1. The present invention also includes molecules 6-G for which reagents for both coupling reactions shown in Scheme 6 are chosen from Pz-NH₂.

A nitrile 6-A Ar-CR^{6''}=CR^{6''}-CN or Ar-CC-CN is dissolved in usually between 1 and 100 volumes of a solvent for example a mixture of anhydrous toluene and absolute ethanol. The solution is cooled to usually between 0°C and -70°C and dry HCl gas is bubbled for between 1 and 24 hours after which the reaction is closed and stirred for between 1 and 72 hours at a temperature usually between -20°C and room temperature. Workup of the reaction mixture provides the O-ethyl imidate 6-B or the alkynyl analog which can be isolated as the HCl salt. 6-B or the alkynyl analog is dissolved in a solvent for example including but not limited to ethanol and cooled to usually between room temperature and -20°C and solution of dry ammonia in an alcohol for example methanol or ethanol or a mixture is added. The dry ammonia solution can be purchased commercially or freshly prepared by bubbling ammonia gas through the appropriate solvent. The mixture is stirred at a temperature generally between 0°C and 50°C for usually between 1 and 24 hours. The mixture is worked up to provide 6-C or the alkynyl analog.

6-C or the alkynyl analog is dissolved in from 1 to 10 weight equivalents of a solvent including but not limited to methanol. A malonic ester derivative including but not limited to dimethyl malonate (generally from 0.75 to 5 molar equivalents) is added to the solution usually at a temperature between 0°C and 50°C. The mixture is cooled to usually -20°C to room temperature and a base for

example including but not limited to NaOCH_3 (usually from 2 to 10 equivalents) is added slowly over generally between 1 and 120 minutes. The resulting solution is heated to a temperature appropriate to the solvent chosen and heated or refluxed for generally between 1 and 24 hours. Work up provides 6-D or the alkynyl analog.

6-D or the alkynyl analog is slowly added in portions to an excess of a chlorinating reagent including but not limited to POCl_3 . One skilled in the art can optionally use the chlorinating agent for example POCl_3 as the solvent as well as reactant, or can use the chlorinating agent as reactant and optionally a solvent can be selected for example including but not limited to acetonitrile. One skilled in the art can use a base including for example but not limited to Hunig's base (diisopropylethylamine) or other amine bases to improve the effectiveness of the reaction. The mixture is stirred generally for from 1 - 24 hours at usually between 75°C and 130°C . The reaction mixture is worked up and purified to give 6-E or the alkynyl analog.

6-E or the alkynyl analog is then coupled to the desired substituents. As illustrated in the Scheme, if the chosen substituents are for example a Pz-NH₂ and an R-X-H chosen from the accompanying figures, either the Pz-NH₂ can be coupled first followed by the R-X-H proceeding through 6-F, or the R-X-H can be coupled first followed by the Pz-NH₂ proceeding through 6-H. If the desired product has two substituents selected from Pz-NH₂ the appropriate reaction conditions as described below are used.

To couple Pz-NH₂, to a solution of 6-E or 6-H or the alkynyl analog in a solvent including for example but not limited to anhydrous DMA (usually from 1 to 100 equivalents) is added usually between 0.5 and 5 equivalents of an amino-pyrazole which can be chosen from the descriptions in this invention, and optionally a reagent including for example NaI or a catalyst and also optionally a base including but not limited to DIPEA. The mixture is stirred at a temperature generally from 50°C to 90°C for usually between 2 and 36 hours. Workup and purification give 6-F or 6-G, respectively, or the alkynyl analog.

To couple R-X-H, 6-E or 6-F or the alkynyl analog is dissolved in the desired R-X-H and heated between generally 50°C to 110°C for usually from 0.25 to 12 hours. Optionally a solvent can be chosen including but not limited to anhydrous DMA and also optionally a base including but not limited to DIPEA or

TEA and also optionally a reagent including but not limited to NaI or a catalyst can be used to facilitate the reaction. Work up and purification provide the target compound 6-H or 6-G, respectively, or the alkynyl analog.

Alternative coupling methods to that shown in Scheme 6 may be used to increase the effectiveness or selectivity of reaction for certain RXH, for example the addition of a base including but not limited to TEA or DIPEA, and also for example replacement of one or both chlorine atoms by iodine or bromine, and also for example the use of catalysts including but not limited to heavy metal catalysts such as palladium complexes or Cu(I). These approaches are well known to one skilled in the art, and are reported for example in publications such as Gomtsyan et al (*J. Med. Chem.* **2002** 45, 3639) and US 5,453,414 which may be referred to for further details. Additional modifications can be accomplished by use of coupling reactions including but not limited to Suzuki, Stille, Grignard, and Buchwald, as known to one skilled in the art. Optionally, appropriate protection groups can be added and removed to facilitate the reaction, as is known to one skilled in the art. Examples of other coupling reactions can be found in the literature including "Strategic Applications of Named Reactions in Organic Synthesis" (L. Kurti and B. Czako, Elsevier Academic Press, New York, NY, 2005). Examples of protection groups can be found in the literature including "Protective Groups in Organic Synthesis – Third Edition" (T.W. Greene, P. G. M. Wuts, Wiley-Interscience, New York, NY, 1999).

The present invention will be understood more readily by reference to the following examples, which are provided by way of illustration and are not intended to be limiting of the present invention.


EXAMPLES

The following abbreviations are used in the examples:

ATP:	adenosine triphosphate
Brij-35:	polyoxyethyleneglycol dodecyl ether
°C:	degrees Celcius
DMEM:	Dulbecco's Modified Eagle's Medium
DMSO:	dimethylsulfoxide
DTT:	dithiothreitol

	g:	gram
	HEPES:	4-(2-hydroxyethyl)piperazine-1-ethanesulfonic acid
	hplc:	high performance liquid chromatography
	IC ₅₀ value:	concentration of an inhibitor that causes a 50% reduction
5		in a measured activity.
	mg:	milligram
	mL:	milliliter
	mmol:	millimole
	MS:	mass spectrum
10	<i>m/z</i> :	mass to charge ratio
	Pz:	optionally modified or substituted or fused pyrazole ring system
	Rf:	ratio to front (ratio of distance traveled by substance/distance traveled by solvent)
	SRB:	sulphorhodamine-B
15	TCA:	trichloroacetic acid
	THF:	tetrahydrofuran
	tlc	thin layer chromatography
	br	broad
	s	singlet
20	d	doublet
	t	triplet
	q	quartet
	dd	doublet of doublets
25	J	coupling constant

EXAMPLE 1




2-methyl-5-nitropyrimidine-4,6-diol

5 Powdered 4,6-dihydroxy-2-methylpyrimidine (9 g, 71 mmol) was dissolved in trifluoroacetic acid (54 mL). The reaction mixture was cooled in ice-water as nitric acid (4.3 mL) was added dropwise over a period of 30 minutes. During addition the internal temperature was maintained below 15 °C. After the addition, the cooling was removed and the reaction stirred overnight. Water (50
10 mL) was added to the reaction mixture and the resulting solid was filtered and dried under vacuum to obtain 2-methyl-5-nitropyrimidine-4,6-diol (5.8 g). ¹H NMR (400 MHz, DMSO-*d*₆): δ 12.97(br, 2H), 2.32 (s, 3H). MS (*m/z*) 172 (M + 1).


15

EXAMPLE 2

5-nitro-2-[(*E*)-2-phenylvinyl]pyrimidine-4,6-diol

A mixture of 2-methyl-5-nitropyrimidine-4,6-diol (4.7 g, 27.5 mmol) and
20 benzaldehyde (42 mL, 414 mmol) was treated with piperidine (21.3 mL, 215.3 mmol). The reaction mixture was heated to 90 °C for 2 hrs. The temperature was then increased to 120 °C for 30 min. After cooling the reaction mixture to ambient, methanol (10 mL) and diethyl ether (200 mL) were added sequentially. The resulting solid was collected by filtration and washed with 5% aqueous
25 hydrochloric acid to give 5-nitro-2-[(*E*)-2-phenylvinyl]pyrimidine-4,6-diol (4.5 g). ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.15 (d, 1H, *J*= 16.4Hz), 7.15 (m, 2H), 7.54 (m, 3H), 6.81 (d, 1H, *J*= 16.4Hz). MS (*m/z*) 260 (M + 1).


EXAMPLE 3

4,6-dichloro-5-nitro-2-[(*E*)-2-phenylvinyl]pyrimidine

5 5-Nitro-2-[(*E*)-2-phenylvinyl]pyrimidine-4,6-diol (2.4 g, 9.63 mmol) was treated with phosphorous oxychloride (10 mL, 106.6 mmol) followed by dropwise addition of diethylaniline (4 mL, 25.1 mmol; slight exotherm was observed during addition). The reaction mixture was slowly heated at 200 °C. After 1.5 hours, the reaction was cooled then poured into crushed ice with stirring. The resulting solid was collected by filtration and dried to give 4,6-dichloro-5-nitro-2-[(*E*)-2-phenylvinyl]pyrimidine (1.6 g). ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.15 (d, 1H, *J*= 16.4Hz), 7.89 (m, 2H), 7.47 (m, 3H), 7.37 (d, 1H, *J*= 16.4Hz). MS (*m/z*) 296 (*M* + 1).

15

EXAMPLE 4


6-(4-methylpiperazin-1-yl)-*N*-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine

20 To a solution of 4,6-dichloro-5-nitro-2-[(*E*)-2-phenylvinyl]pyrimidine (200 mg, 0.676 mmol) in THF (5 mL) was added triethylamine (204 mg, 2.03 mmol) and the reaction mixture was stirred for 15 minutes. 3-Amino-5-methylpyrazole (66 mg, 0.68 mmol) in THF (3 mL) was added dropwise to the reaction mixture. After 1 hour, *N*-methylpiperazine (74 mg, 0.74 mmol) in THF (3 mL) was added to the reaction mixture dropwise. The reaction mixture was stirred at

room temperature for 1 hour followed by addition of water (10 mL). After stirring for 15 minutes, ethyl acetate (50 mL) was added. The organic layer was separated, dried and the solvent evaporated under reduced pressure. The residue was triturated with chloroform and hexane to obtain 6-(4-methylpiperazin-1-yl)-
 5 *N*-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine (80 mg). ¹H NMR (400 MHz, DMSO-*d*₆): δ 12.35 (br, 1H), 10.52 (br, 1H), 7.95 (d, 1H, *J*= 16Hz), 7.83 (d, 2H, *J*= 6.8Hz), 7.5 (m, 3H), 7.11 (d, 1H, *J*= 16Hz), 6.8 (s, 1H), 3.65 (br, 4H), 2.5 (br, 4H), 2.36 (s, 3H), 2.29 (s, 3H). MS (*m/z*) 421 (*M* + 1).

10 Examples 5 to 9 were prepared in the same manner as Example 4 by replacing the *N*-methylpiperazine with the appropriate amine.


EXAMPLE 5



15 *N,N'*-bis(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-2-[(*E*)-2-phenylvinyl]pyrimidin-4,6-diamine

By replacing *N*-methylpiperazine with 3-amino-5-methylpyrazole was obtained *N,N'*-bis(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-2-[(*E*)-2-phenylvinyl]-
 20 pyrimidin-4,6-diamine in 44% yield. ¹H NMR (400 MHz, DMSO-*d*₆): δ 12.41 (br, 2H), 11.35 (br, 2H), 7.92 (d, 1H, *J*=16Hz), 7.79 (d, 1H, *J*= 6.8Hz), 7.76 (m, 3H), 7.16 (d, 1H, *J*= 16Hz), 6.80 (s, 2H), 2.31 (s, 6H). MS (*m/z*) 418 (*M* + 1).

EXAMPLE 6




N-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-6-(morpholin-4-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine

5

By replacing *N*-methylpiperazine with morpholine was obtained *N*-(3-methyl-1*H*-pyrazol-5-yl)-6-(morpholin-4-yl)-5-nitro-2-[(*E*)-2-phenylvinyl]-pyrimidin-4-amine in 15% yield after preparative hplc chromatography. ¹H NMR (400 MHz, CDCl₃): δ 10.56 (br, 1H), 7.88 (d, 1H, *J*=16Hz), 7.65 (d, 2H, *J*=6.8Hz), 7.40 (m, 3H), 7.06 (d, 1H, *J*=16Hz), 6.38 (s, 1H), 3.83 (m, 4H), 3.69 (m, 4H), 2.37 (s, 3H). MS (*m/z*) 408 (*M* + 1).

10

EXAMPLE 7




N-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-6-(piperazin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine

15

By replacing *N*-methylpiperazine with piperazine was obtained *N*-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-6-(piperazin-1-yl)-2-[(*E*)-2-phenylvinyl]-pyrimidin-4-amine in 17% yield after preparative hplc. ¹H NMR (400 MHz, CDCl₃): δ 10.58 (br, 1H), 7.87 (d, 1H, *J*=16Hz), 7.62 (m, 2H), 7.38 (m, 3H), 6.96 (d, 1H, *J*=16Hz), 6.37 (s, 1H), 3.74 (m, 4H), 3.58 (m, 4H), 2.36 (s, 3H). MS (*m/z*) 407 (*M* + 1).

20

EXAMPLE 8




N-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-6-(piperidin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine

5

By replacing *N*-methylpiperazine with piperidine was obtained *N*-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-6-(piperidin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine in 16% yield after preparative hplc. ¹H NMR (400 MHz, CDCl₃): δ 10.58(br, 1H), 7.87 (d, 1H, *J*=16Hz), 7.62 (m, 2H), 7.38(m, 3H), 6.95 (d, 1H, *J*=16Hz), 6.35 (s, 1H), 3.58 (m, 4H), 2.36 (s, 3H), 1.4-1.9(m, 6H). MS (*m/z*) 406 (*M* + 1).

10

EXAMPLE 9




N-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-6-(pyrrolidin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine

15

By replacing *N*-methylpiperazine with pyrrolidine was obtained *N*-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-6-(pyrrolidin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine in 14% yield after preparative chromatography. ¹H NMR (400 MHz, CDCl₃): δ 10.5 (br, 1H), 7.9 (d, 1H, *J*=16Hz), 7.65 (m, 2H), 7.39 (m, 3H), 7.13 (d, 1H, *J*= 16Hz), 6.29 (s, 1H), 2.5-3.5 (m, 4H), 2.37 (s, 3H), 2.02 (m, 4H). MS (*m/z*) 392 (*M* + 1).

20

EXAMPLE 10



6-(4-methylpiperazin-1-yl)-*N*-(3-methyl-1*H*-pyrazol-5-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4,5-diamine

5


A solution of tin (II) chloride dihydrate (600 mg, 4.43 mmol) in concentrated hydrochloric acid (1 mL, 9.04 mmol) was cooled below 10 °C. A solution of 6-(4-methylpiperazin-1-yl)-*N*-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine (Example 4, 400 mg, 0.95 mmol) in methanol (20 mL) was added dropwise to the reaction mixture. The reaction was allowed to come to ambient temperature and was then heated to 60 °C for 3 hours. Progress of the reaction was monitored by TLC. The reaction mixture was reduced to 1/3rd volume of the original volume under vacuum. The residue was diluted with ethyl acetate (40 mL) and 1N NaOH (20 mL) was added to the reaction mixture. The organic layer was separated, dried and the solvent evaporated under reduced pressure. The residue was then triturated with ethyl acetate and hexane to afford 6-(4-methylpiperazin-1-yl)-*N*-(3-methyl-1*H*-pyrazol-5-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4,5-diamine as a yellow solid (200 mg). ¹H NMR (400 MHz, DMSO-*d*₆): δ 7.94 (br, 1H), 7.70(d, 1H, *J*= 16Hz), 7.59 (m, 2H) 7.38 (m, 2H), 7.29 (m, 1H), 7.05 (d, 1H, *J*= 16Hz), 6.36 (br, 1H), 3.44 (br, 2H), 3.3 (br, 4H), 2.59 (br, 4H), 2.36 (s, 3H), 2.33 (s, 3H). MS (*m/z*) 391 (*M* + 1).

20

Examples 11 to 15 were prepared in the same manner as Example 10 from the appropriate starting material.

25

EXAMPLE 11




N,N'-bis(3-methyl-1*H*-pyrazol-5-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4,5,6-triamine

5

From Example 5, *N,N'*-bis(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-2-[(*E*)-2-phenylvinyl]pyrimidin-4,6-diamine, was obtained *N,N'*-bis(3-methyl-1*H*-pyrazol-5-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4,5,6-triamine in 43% yield. ¹H NMR (400 MHz, DMSO-*d*₆): δ 11.86 (br, 2H), 8.41 (br, 2H), 7.2-7.8 (m, 6H), 7.0-7.1 (br, 1H), 6.5 (br, 1H), 2.11 (s, 6H). MS (*m/z*) 388 (*M* + 1).

10

EXAMPLE 12




N-(3-methyl-1*H*-pyrazol-5-yl)-6-(morpholin-4-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4,5-diamine

15

From Example 6, *N*-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-6-(morpholin-4-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine, was obtained *N*-(3-methyl-1*H*-pyrazol-5-yl)-6-(morpholin-4-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4,5-diamine in 21% yield after preparative hplc. ¹H NMR (400 MHz, CDCl₃): δ 11.35 (br, 1H), 7.58 (m, 2H), 7.42 (m, 4H), 6.74 (d, 1H, *J*=16Hz), 5.82 (s, 1H), 3.67 (m, 4H), 3.27 (m, 4H), 2.17 (s, 3H). MS (*m/z*) 378 (*M* + 1).

20

EXAMPLE 13




N-(3-methyl-1*H*-pyrazol-5-yl)-6-(piperazin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4,5-diamine

5

From Example 7, *N*-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-6-(piperazin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine, was obtained *N*-(3-methyl-1*H*-pyrazol-5-yl)-6-(piperazin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4,5-diamine in 17% yield after preparative hplc. ¹H NMR (400 MHz, CDCl₃): δ 11.3 (br, 1H), 7.31-7.63 (m, 6H), 6.82 (d, 1H, *J*=16Hz), 5.85 (s, 1H), 3.28 (m, 4H), 2.22 (s, 3H), 1.65(m, 4H). MS (*m/z*) 377 (*M* + 1).

10

EXAMPLE 14




N-(3-methyl-1*H*-pyrazol-5-yl)-6-(piperidin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4,5-diamine

15

From Example 8, *N*-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-6-(piperidin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine, was obtained *N*-(3-methyl-1*H*-pyrazol-5-yl)-6-(piperidin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4,5-diamine in 18% yield. ¹H NMR (400 MHz, CDCl₃): δ 7.73 (d, 1H, *J*=16Hz), 7.62 (d, 2H, *J*=7.2Hz), 7.39 (t, 2H, *J*=7.2Hz), 7.32 (m, 1H), 7.09 (d, 1H, *J*=16Hz), 6.3 (s, 1H), 3.29 (br, 2H), 3.22 (br, 4H), 2.36 (s, 3H), 1.4-1.9 (m, 6H). MS (*m/z*) 376 (*M* + 1).

20

EXAMPLE 15




N-(3-methyl-1*H*-pyrazol-5-yl)-6-(pyrrolidin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4,5-diamine

5

From Example 9, *N*-(3-methyl-1*H*-pyrazol-5-yl)-5-nitro-6-(pyrrolidin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine, was obtained *N*-(3-methyl-1*H*-pyrazol-5-yl)-6-(pyrrolidin-1-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4,5-diamine in 20% yield. ¹H NMR (400 MHz, CDCl₃): δ 7.88 (br, 1H), 7.71 (d, 1H, *J*=16Hz), 7.59 (d, 2H, *J*=7.2Hz), 7.39 (t, 2H, *J*=7.2Hz), 7.29 (m, 1H), 7.00 (d, 1H, *J*=16Hz), 5.95 (s, 1H), 3.73 (m, 4H), 2.67(br, 2H), 2.30 (s, 3H), 1.97 (m, 4H). MS (*m/z*) 362 (*M* + 1).

10

EXAMPLE 16



15


ethyl 2-chloro-6-[(3-methyl-1*H*-pyrazol-5-yl)amino]-5-nitropyrimidine-4-carboxylate

To a solution of commercially available ethyl 2,6-dichloro-5-nitropyrimidine-4-carboxylate (Matrix Scientific, 1.00 g, 3.76 mmol, 1.0 equiv) and 2,6-lutidine (0.65 mL, 5.60 mmol, 1.5 equiv) in anhydrous 1,4-dioxane (5 mL) at room temperature was added drop-wise a solution of 5-methyl-3-aminopyrazole (0.38 g, 3.95 mmol, 1.05 equiv) in anhydrous 1,4-dioxane (5 mL). The solution was stirred at room temperature for 30 min, diluted with ethyl acetate (100 mL), washed with 1 N hydrochloric acid (50 mL x 3) and brine (50 mL x 1), dried over Na₂SO₄, and concentrated to dryness to yield a light-brown solid as

25

Example 16 (1.16 g, 95%): R_f 0.55 (60% ethyl acetate/hexane); MS m/z 327, calcd 327 ($C_{11}H_{11}ClN_6O_4 + H$).

EXAMPLE 17



5


ethyl 6-[(3-methyl-1*H*-pyrazol-5-yl)amino]-5-nitro-2-[(*E*)-2-phenylvinyl]pyrimidine-4-carboxylate

10 To a glass vessel containing Example 16 (100 mg, 0.31 mmol, 1.0 equiv), anhydrous potassium carbonate (64 mg, 0.46 mmol, 1.5 equiv), *trans*-2-phenylvinylboronic acid (69 mg, 0.46 mmol, 1.5 equiv), and tetrakis(triphenylphosphine)palladium(0) (18 mg, 0.015 mmol, 0.05 equiv) was added anhydrous toluene (3 mL) under nitrogen atmosphere. The suspension was

15 de-gassed by bubbling nitrogen through for 2-3 minutes. The suspension was then heated with rapid stirring at 70 °C for 8 hours, cooled to room temperature, diluted with water (50 mL), and extracted with dichloromethane (50 mL x 3). The combined extracts were and concentrated under reduced pressure. The residue was

20 purified on a silica gel flash column with 0-50% ethyl acetate/hexane to yield Example 17 as an orange solid (62 mg, 51%): R_f 0.40 (60% ethyl acetate/hexane); MS m/z 395, calcd 395 ($C_{19}H_{18}N_6O_4 + H$).

EXAMPLE 18




ethyl 5-amino-6-[(3-methyl-1H-pyrazol-5-yl)amino]-2-[(E)-2-phenylvinyl]pyrimidine-4-carboxylate

5

To a solution of Example 16 (0.40 g, 1.02 mmol, 1 equiv) in dimethylsulfoxide (5 mL) and acetonitrile (20 mL) was added zinc dust (0.66 g, 10.09 mmol, 10 equiv), followed by the addition of ammonium formate (0.63 g, 10.00 mmol, 10 equiv). The suspension was stirred at room temperature for 30 minutes. The solid was filtered off with a Celite® pad and the filtrate was concentrated under reduced pressure to yield the crude product that was purified by column chromatography on silica gel using 5 % methanol in chloroform, yielding Example 18 as a yellow solid (70 mg, 19 %): R_f 0.55 (10% methanol/dichloromethane); ^1H NMR (300 MHz, DMSO- d_6) δ 12.19 (br s, 1 H), 9.60 (br s, 1 H), 7.63 (d, $J = 7.5$ Hz, 12 H), 7.53 (d, $J = 15.9$ Hz, 1 H), 7.35-7.45 (m, 2 H), 7.26-7.34 (m, 1 H), 7.07 (d, $J = 16.0$ Hz, 1 H), 7.04 (br s, 2 H), 6.75 (s, 1 H), 4.38 (q, $J = 7.8$ Hz, 2 H), 2.32 (s, 3 H), 1.34 (t, $J = 7.8$ Hz, 3 H); MS m/z 365, calcd 365 ($C_{19}H_{20}N_6O_2 + H$).

20 Examples 19 to 23 were prepared in the same manner as Example 17 starting from Example 16 by substituting the appropriate vinylboronic acid for *trans*-2-phenylvinylboronic acid in the synthetic sequence.

EXAMPLE 19




ethyl 5-amino-2-[(*E*)-2-(4-chlorophenyl)vinyl]-6-[(3-methyl-1*H*-pyrazol-5-yl)amino]pyrimidine-4-carboxylate

5

By substituting *trans*-2-(4-chlorophenyl)vinylboronic acid for *trans*-2-phenylvinylboronic acid was obtained a light-yellow solid (14 mg): ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.12 (br s, 1 H), 9.52 (s, 1 H), 7.69 (d, *J* = 8.3 Hz, 1 H), 7.49 (d, *J* = 16.0 Hz, 1 H), 7.44 (d, *J* = 8.3 Hz, 2 H), 7.09 (d, *J* = 16.0 Hz, 1 H), 7.04 (s, 2 H), 6.75 (s, 1 H), 4.34 (q, *J* = 7.0 Hz, 2 H), 2.32 (s, 3 H), 1.34 (t, *J* = 7.0 Hz, 3 H); MS *m/z* 399, calcd 399 (C₁₉H₁₉ClN₆O₂+H).

10

EXAMPLE 20




ethyl 5-amino-6-[(3-methyl-1*H*-pyrazol-5-yl)amino]-2-[(*E*)-2-[4-(trifluoromethyl)phenyl]vinyl]pyrimidine-4-carboxylate

15

By substituting *trans*-2-(4-(trifluoromethyl)phenyl)vinylboronic acid for *trans*-2-phenylvinylboronic acid was obtained a light-yellow solid (60 mg): ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.20 (br s, 1 H), 7.89 (d, *J* = 8.0 Hz, 2 H), 7.77 (d, *J* = 8.0 Hz, 2 H), 7.66 (d, *J* = 16.0 Hz, 1 H), 7.41-7.37 (m, 3 H), 6.76 (s, 1 H), 4.40 (q, *J* = 6.9 Hz, 2 H), 2.34 (s, 3 H), 1.37 (t, *J* = 6.9 Hz, 3 H); MS *m/z* 433, calcd 433 (C₂₀H₁₉F₃N₆O₂+H).

20

EXAMPLE 21




ethyl 5-amino-2-[(*E*)-2-(4-methylphenyl)vinyl]-6-[(3-methyl-1*H*-pyrazol-5-yl)amino]pyrimidine-4-carboxylate

5

By substituting *trans*-2-(4-methylphenyl)vinylboronic acid for *trans*-2-phenylvinylboronic acid was obtained a light-yellow solid (22 mg): ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.13 (br s, 1 H), 9.49 (s, 1 H), 7.55-7.48 (m, 3 H), 7.21 (d, *J* = 7.9 Hz, 2 H), 7.02-7.00 (m, 3 H), 6.78 (s, 1 H), 4.34 (q, *J* = 7.1 Hz, 2 H), 2.33 (s, 6 H), 1.34 (t, *J* = 7.1 Hz, 3 H); MS *m/z* 379, calcd 379 (C₂₀H₂₂N₆O₂+H).

10

EXAMPLE 22




ethyl 5-amino-2-[(*E*)-2-(3-fluorophenyl)vinyl]-6-[(3-methyl-1*H*-pyrazol-5-yl)amino]pyrimidine-4-carboxylate

15

By substituting *trans*-2-(3-fluorophenyl)vinylboronic acid for *trans*-2-phenylvinylboronic acid was obtained a light-yellow solid (13 mg): ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.11 (br s, 1 H), 9.51 (s, 1 H), 7.56-7.38 (m, 4 H), 7.14-7.04 (m, 4 H), 6.76 (s, 1 H), 4.33 (q, *J* = 7.1 Hz, 2 H), 2.31 (s, 3 H), 1.32 (t, *J* = 7.1 Hz, 3 H); MS *m/z* 383, calcd 383 (C₁₉H₁₉FN₆O₂+H).

20

EXAMPLE 23




ethyl 5-amino-2-[(*E*)-2-(4-methoxyphenyl)vinyl]-6-[(3-methyl-1*H*-pyrazol-5-yl)amino]pyrimidine-4-carboxylate

5

By substituting *trans*-2-(4-methoxyphenyl)vinylboronic acid for *trans*-2-phenylvinylboronic acid was obtained a light-yellow solid (76 mg): ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.70 (d, *J* = 15.7 Hz, 1 H), 7.59 (d, *J* = 8.7 Hz, 2 H), 7.22 (d, *J* = 15.7 Hz, 1 H), 7.02 (d, *J* = 8.7 Hz, 2 H), 6.48 (br s, 4 H), 4.42 (q, *J* = 7.1 Hz, 2 H), 3.79 (s, 3 H), 2.32 (s, 3 H), 1.35 (t, *J* = 7.1 Hz, 3 H); MS *m/z* 395, calcd 395 (C₂₀H₂₂N₆O₃+H).

10

EXAMPLE 24



1-methylpiperidin-4-yl 5-amino-6-[(3-methyl-1*H*-pyrazol-5-yl)amino]-2-[(*E*)-2-phenylvinyl]pyrimidine-4-carboxylate


15

A mixture of Example 18 (4.5 g, 12.3 mmol, 1 equiv) and 4-hydroxy-*N*-methyl piperidine (40 mL) was slowly added titanium isopropoxide (1 mL). The reaction was heated to reflux overnight. The excess of the alcohol was distilled off and the resultant solid was purified by column chromatography on silica gel using 3 % methanol in dichloromethane to afford Example 24 as a yellow solid (1.55 g, 29 %): ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.11 (s, 1 H), 9.49 (s, 1 H), 7.64-7.62 (m, 2 H), 7.53 (d, *J* = 16.0 Hz, 1 H), 7.40-7.27 (m, 3 H), 7.05-6.98 (m, 3 H), 6.76

20

(s, 1 H), 4.88 (m, 1 H), 2.73-2.49 (m, 2 H), 2.30 (s, 3 H), 2.15 (m, 5 H), 1.97-1.93 (m, 2 H), 1.77-1.69 (m, 2 H); MS m/z 434, calcd 434 ($C_{23}H_{27}N_7O_2+H$).

EXAMPLE 25




5

N-(2-morpholin-4-ylethyl)-5-amino-6-[(3-methyl-1*H*-pyrazol-5-yl)amino]-2-[(*E*)-2-phenylvinyl]pyrimidine-4-carboxamide

To a mixture of Example 18 (2.5 g, 6.9 mmol, 1 equiv) in dry
 10 dichloromethane and 2-morpholin-4-ylethylamine (1.8 g, 13.7 mmol, 2 equiv) at 0
 °C was added trimethylaluminium (28 mL, 54.8 mmol, 8 equiv) dropwise in about
 one hour. The reaction was brought to room temperature and stirred for another 4
 hours. The reaction was carefully quenched with 1.5 N hydrochloric acid (20 mL).
 The resultant solid was collected and purified by column chromatography on
 15 silica gel using 4 % methanol in dichloromethane to yield Example 25 as a yellow
 solid (1.1 g, 36 %): ^1H NMR (400 MHz, DMSO- d_6) δ 12.08 (s, 1 H), 9.34 (s, 1
 H), 8.75 (s, 1 H), 7.64-7.60 (m, 3 H), 7.42-7.39 (m, 2 H), 7.30 (m, 1 H), 7.14 (s, 2
 H), 7.02 (d, $J = 16.0$ Hz, 1 H), 6.77 (s, 1 H), 3.61-3.59 (m, 4 H), 3.41-3.38 (m, 2
 H), 2.53-2.49 (m, 6 H), 2.29 (s, 3 H); MS m/z 449, clcad 449 ($C_{23}H_{28}N_8O_2+H$).

20

EXAMPLE 26




methyl 6-[(3-methyl-1*H*-pyrazol-5-yl)amino]-2-[(*E*)-2-phenylvinyl]pyrimidine-4-carboxylate

5

By substituting commercially available methyl 2,4-dichloropyrimidine-6-carboxylate for ethyl 2,6-dichloro-5-nitropyrimidine-4-carboxylate in the reaction sequence used to produce Example 17 was obtained Example 26 as a light-yellow solid (30 mg): ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.30 (br s, 1 H), 7.85 (d, *J* = 16.0 Hz, 1 H), 7.65-7.75 (m, 2 H), 7.33-7.48 (m, 4 H), 7.19 (d, *J* = 16.0 Hz, 1 H), 6.40 (br s, 1 H), 3.89 (s, 3 H), 2.27 (s, 3 H); MS *m/z* 336, calcd 336 (C₁₈H₁₇N₅O₂+H).

10

EXAMPLE 27



15

4,6-dichloro-2-(methylsulfonyl)pyrimidine


Commercially available 4,6-dichloro-2-methylthiopyrimidine (Aldrich, 21.0 g, 107.0 mmol, 1 equiv) was dissolved in dichloromethane and cooled with an ice-bath. 3-chlorobenzoyl peroxide (60.0 g, 77% wt, 268.0 mmol, 2.5 equiv) was added in small portions. The resulting white suspension was stirred at room temperature for 4 hours, washed with 1 M sodium thiosulfate/saturated sodium bicarbonate (1:1, v/v, 200 mL x 3) and saturated sodium bicarbonate (100 mL x 3), and brine (100 mL x 1). The organic solution was dried and concentrated under reduced pressure. After drying under high vacuum for overnight Example

20

25

27 was obtained as a white solid: (22.0 g, 91%): R_f 0.20 (20% ethyl acetate/hexane); MS m/z 227, calcd 227 ($C_5H_4Cl_2N_2O_2S+H$).

EXAMPLE 28




5

4,6-dichloro-2-(phenylethynyl)pyrimidine

Example 27 (4.54 g, 20.00 mmol, 1 equiv) was dissolved in anhydrous tetrahydrofuran (50 mL) and cooled to $-20\text{ }^{\circ}\text{C}$. Phenylacetylenylmagnesium bromide (22.00 mL, 1.0 M in tetrahydrofuran, 22.00 mmol, 1.1 equiv) was added dropwise with vigorous stirring. The solution was stirred from $-20\text{ }^{\circ}\text{C}$ to room temperature for overnight, diluted with ethyl acetate (300 mL), and added to 1 N hydrochloric acid (200 mL). The mixture was vigorously stirred for 5 minutes. The organic layer was collected, and the aqueous layer was extracted with more ethyl acetate (100 mL x 2). The combined organic solution was dried over Na_2SO_4 , concentrated under reduced pressure, and dried under high vacuum for overnight to afford the Example 28 as an off-white solid (5.00 g, quantitative): R_f 0.75 (20% ethyl acetate/hexane); MS m/z 249, calcd 249 ($C_{12}H_6N_2Cl_2+H$).

20

EXAMPLE 29


6-chloro-*N*-(3-methyl-1*H*-pyrazol-5-yl)-2-(phenylethynyl)pyrimidin-4-amine

Example 28 (5.00 g, 20.00 mmol, 1 equiv) was dissolved in anhydrous dimethylacetamide (20 mL). 5-Methyl-2-aminopyrazole (2.14 g, 22.00 mmol, 1.1

25

equiv), sodium iodide (3.60 g, 24.00 mmol, 1.2 equiv), and diisopropylethylamine (4.18 mL, 24.00 mmol, 1.2 equiv) were added. The solution was heated at 90 °C for overnight, cooled to room temperature, diluted with ethyl acetate (200 mL), washed with saturated sodium bicarbonate (200 mL x 3), dried
 5 and concentrated under reduced pressure. The resulting crude product was purified with flash chromatography on silica gel using 0%~4% methanol/dichloromethane to yield Example 29 as a yellow solid (5.51 g, 89%): R_f 0.40 (5% methanol/dichloromethane); ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 12.13 (br s, 1 H), 10.39 (s, 1 H), 7.63 (d, $J = 7.5$ Hz, 2 H), 7.740-7.60 (m, 4 H), 5.95 (br
 10 s, 1 H), 2.22 (s, 3 H); MS m/z 310, calcd 310 ($\text{C}_{16}\text{H}_{12}\text{ClN}_5 + \text{H}$).


EXAMPLE 30



15 6-(4-methylpiperazin-1-yl)-N-(3-methyl-1H-pyrazol-5-yl)-2-(phenylethynyl)pyrimidin-4-amine

Example 29 (2.80 g, 9.04 mmol, 1 equiv) was dissolved in anhydrous 1,4-dioxane (10 mL). *N*-methylpiperazine (1.10 mL, 9.92 mmol, 1.1 equiv), 4-dimethylaminopyridine (0.055 g, 0.45 mmol, 0.05 equiv), and diisopropylethylamine (1.89 mL, 10.85 mmol, 1.2 equiv) were added. The solution
 20 was stirred at 100 °C for 2 hours, cooled to room temperature, and concentrated under reduced pressure. The residue was purified with flash chromatography on silica gel using 0%~10% methanol/dichloromethane to give the Example 30 as a light-yellow solid (3.00 g, 89%): R_f 0.10 (5% methanol/dichloromethane); ^1H
 25 NMR (300 MHz, $\text{DMSO}-d_6$) δ 11.90 (br s, 1 H), 9.43 (s, 1 H), 7.57 (d, $J = 7.5$ Hz, 2 H), 7.40-7.50 (m, 3 H), 6.80 (br s, 1 H), 5.82 (s, 1 H), 3.51 (br s, 4 H), 2.43 (br s, 4 H), 2.27 (s, 3 H), 2.18 (s, 3 H); MS m/z 374, calcd 374 ($\text{C}_{21}\text{H}_{23}\text{N}_7 + \text{H}$).

EXAMPLE 31



6-(4-methylpiperazin-1-yl)-*N*-(3-methyl-1*H*-pyrazol-5-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine


5

Example 30 (2.54 g, 6.80 mmol, 1 equiv) was dissolved in anhydrous tetrahydrofuran (50 mL) and cooled with an ice-water bath. Lithium aluminum hydride (5.61 mL, 1.0 M in tetrahydrofuran, 5.61 mmol, 0.83 equiv) was added dropwise with rapid stirring. The solution was stirred from 0 °C to room temperature for 12 hours. The reaction was cooled to 0 °C and slowly quenched with methanol (5 mL). Saturated sodium potassium tartrate (500 mL) was added to the reaction mixture. The resulting suspension was rapidly stirred at room temperature until a clear solution was obtained. The solution was then extracted with ethyl acetate (100 mL x 5). The combined extracts were dried and concentrated under reduced pressure. Flash chromatography on silica gel with 0%~8% methanol/dichloromethane afforded Example 31 as an off-white solid (1.55 g, 61%): R_f 0.30 (10% methanol/dichloromethane); ^1H NMR (300 MHz, DMSO- d_6) δ 11.80 (br s, 1 H), 9.08 (s, 1 H), 7.72 (d, J = 15.9 Hz, 1 H), 7.63 (d, J = 7.2 Hz, 2 H), 7.28-7.45 (m, 3 H), 6.92 (d, J = 15.9 Hz, 1 H), 6.62 (br s, 1 H), 5.95 (s, 1 H), 3.56 (br s, 4 H), 2.45 (br s, 4 H), 2.26 (s, 3 H), 2.20 (s, 3 H); MS m/z 376, calcd 376 (C₂₁H₂₅N₇+H).

20

Example 31 was also made via Scheme 6 as described below.

EXAMPLE 32




6-(4-methylpiperazin-1-yl)-N-(3-methyl-1H-pyrazol-5-yl)-2-[(Z)-2-phenylvinyl]pyrimidin-4-amine

5

Example 30 (38 mg, 0.10 mmol) was dissolved in ethyl acetate/methanol (2:1, v/v, 1.5 mL). Quinoline (0.04 mL) and Lindlar's catalyst (20 mg) were added. The suspension was de-gassed with hydrogen and rapidly stirred under an hydrogen atmosphere (1 atm) for overnight. The catalyst was filtered off with a small celite pad. The filtrate was concentrated and the residue was purified with preparative thin-layer chromatography on silica gel using 10% methanol/dichloromethane to yield Example 32 as an off-white solid (32 mg, 85%); R_f 0.45 (10% methanol/dichloromethane); ^1H NMR (300 MHz, DMSO- d_6) δ 11.78 (br s, 1 H), 9.19 (s, 1 H), 7.38 (d, $J = 7.2$ Hz, 2 H), 7.15-7.30 (m, 3 H), 6.77 (d, $J = 12.5$ Hz, 1 H), 6.50 (br s, 1 H), 6.36 (d, $J = 12.5$ Hz, 1 H), 5.74 (s, 1 H), 3.35-3.50 (m, 4 H), 2.65-2.80 (m, 4 H), 2.56 (s, 3 H), 2.13 (s, 3 H); LRMS m/z 376, calcd 376 ($C_{21}H_{25}N_7+H$).

15

EXAMPLE 33



20

4,6-dichloro-2-[(E)-2-phenylvinyl]pyrimidine

To a rapidly stirred suspension of Rieke Mg (4.81 g, 200 mmol, 1.5 equiv) in tetrahydrofuran under nitrogen was added a tetrahydrofuran solution of β -bromostyrene (mixture of *cis* and *trans*, 29.00 g, 160 mmol, 1.2 equiv) dropwise in such a rate that the reaction temperature was maintained between 40-60 °C throughout the entire addition process. After the addition, the resulting deep-red


25

solution was stirred at room temperature for 30 minutes and slowly added *via* a double-ended needle into a cooled (-20 °C) tetrahydrofuran solution of Example 27 (30.00 g, 132 mmol, 1 equiv) within 45 minutes. The reaction was stirred from -20 °C to room temperature for 4 hours, cooled to -20 °C, and quenched with the dropwise addition of 1 N hydrochloric acid (200 mL). The mixture was concentrated at room temperature under reduced pressure and extracted with ethyl acetate (500 mL). The ethyl acetate layer was separated, washed with 1 N hydrochloric acid (200 mL x 3), dried, and concentrated under reduced pressure. The crude product was purified with flash chromatography on silica gel with 0-2% ethyl acetate/hexane to yield Example 33 as a light-yellow solid (14.20 g, ~9:1 *trans/cis*, 43%): R_f 0.60 (5% ethyl acetate/hexane); MS m/z 251, calcd 251 ($C_{12}H_8Cl_2N_2+H$).

Example 33 was also synthesized via Scheme 6 as indicated below.

15

EXAMPLE 34



6-chloro-*N*-(3-methyl-1*H*-pyrazol-5-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine

Example 33 (14.20 g, 56.55 mmol, 1 equiv) was dissolved in anhydrous dimethylacetamide (100 mL). 5-Methyl-3-aminopyrazole (6.60 g, 68.00 mmol, 1.2 equiv), sodium iodide (12.70 g, 84.73 mmol, 1.5 equiv), and diisopropylethylamine (14.76 mL, 84.73 mmol, 1.5 equiv) were added. The solution was stirred at 90 °C for 12 hours, cooled to room temperature, diluted with ethyl acetate (500 mL), washed with water (500 mL x 1), saturated sodium bicarbonate (200 mL x 3), and brine (100 mL x 1), dried, and concentrated under reduced pressure. The resulting semi-solid was suspended in minimum dichloromethane (15 mL) with rapid swirling, and the fine precipitates were collected by filtration, washed with dichloromethane/hexane (1:1, v/v, 10 mL x 4), and dried under high vacuum for overnight to yield Example 34 as an off-white solid (8.75 g, 50%, *trans* only): R_f 0.75 (10% methanol/dichloromethane); ^1H NMR (300 MHz,


30

DMSO-*d*₆) δ 12.20 (s, 1 H), 10.16 (s, 1 H), 7.83 (d, *J* = 12.0 Hz, 1 H), 7.73 (d, *J* = 5.1 Hz, 2 H), 7.36-7.48 (m, 4 H), 7.07 (d, *J* = 12.0 Hz, 1 H), 5.78 (s, 1 H), 2.24 (s, 3 H); MS *m/z* 312, calcd 312 (C₁₆H₁₄ClN₅+H).

Example 34 was also synthesized via Scheme 6 as indicated below.

5

EXAMPLE 35



6-(4-methylpiperazin-1-yl)-*N*-(3-methyl-1*H*-pyrazol-5-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine


10

Example 34 (7.79 g, 25.00 mmol, 1 equiv) was dissolved in 1,4-dioxane (15 mL). *N*-Methylpiperazine (4.16 mL, 37.50 mmol, 1.5 equiv), 4-dimethylaminopyridine (0.15 g, 1.25 mmol, 0.05 equiv), and diisopropylethylamine (6.97 mL, 40.00 mmol, 1.6 equiv) were added. The solution was stirred at 100 °C for 48 hours, cooled to room temperature, and concentrated under reduced pressure. The residue was purified with flash chromatography on silica gel using 0-8% methanol/dichloromethane to give Example 31 as a light-yellow solid (3.56 g, 38%): Example 35 was also synthesized via Scheme 6 as indicated below. All analytical data were identical to those of Example 31 prepared by the alternate routes described.

15


20

EXAMPLE 36

methyl 4-[(*E*)-2-(3,5-dihydroxy-4-nitrophenyl)vinyl]benzoate

5 To Example 1 (2 g, 11.69 mmol) was added 4-formyl-benzoic acid methyl ester (7.69 g, 46.76 mmol) followed by piperidine (10 mL, 93.52 mmol). The reaction mixture was heated at 90 °C for 4 hours. The reaction mixture was then cooled to room temperature and methanol (10 mL) was added followed by diethyl ether (100 mL). The solid obtained was filtered and washed with 5%
 10 hydrochloric acid to afford Example 36 (2.0 g). ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.3 (br, 1H), 8.0-8.2 (m, 3H), 7.4 (d, 2H, *J* = 8Hz), 6.9 (d, 1H, *J* = 16.4Hz), 3.88(s, 3H).

EXAMPLE 36A




15

methyl 4-[(*E*)-2-(3,5-dichloro-4-nitrophenyl)vinyl]benzoate

Example 36 (6 g, 19.8 mmol) was treated with phosphorous oxychloride (21 mL, 229 mmol) followed by drop-wise addition of diethylaniline (9.5 mL, 59.6 mmol). The reaction mixture was heated to 80 °C overnight. The reaction mixture was then poured into crushed ice and the resulting solid collected and dried. The compound was then purified by column chromatography. 10% ethyl acetate:hexane was used as the eluent to give Example 36A (1.7 g). ¹H NMR (400 MHz, CDCl₃): δ 8.0-8.2 (m, 3H), 7.70 (m, 2H), 7.47 (d, 2H, *J* = 8Hz), 7.22
 20 (d, 1H, *J* = 16.4Hz), 3.95(s, 3H).

25

EXAMPLE 37



methyl 4-((*E*)-2-{4-chloro-6-[(3-methyl-1*H*-pyrazol-5-yl)amino]-5-
nitropyrimidin-2-yl} vinyl)benzoate


5

To a solution of Example 36A (1.55 g, 4.39 mmol) in tetrahydrofuran (30 mL) was added triethylamine (1.1 mL, 7.9 mmol) under nitrogen atmosphere at room temperature (30 °C). After 15 minutes, 5-amino-2-methyl-pyrazzole (426 mg, 4.39 mmol) in tetrahydrofuran (10 mL) was added drop-wise to the reaction mixture. After 5 hours the reaction mixture was evaporated to dryness. The solid obtained was triturated with ethyl acetate and hexane to obtain Example 37 (1.8 g). ¹H NMR (400 MHz, DMSO-*d*₆): δ 12.1 (br, 1H), 9.75 (br, 1H), 8.08 (m, 3H), 7.69 (d, 1H, *J*= 8.4Hz), 7.13 (d, 1H, *J*= 16Hz), 6.66 (s, 1H), 3.94 (s, 3H), 2.17 (s, 3H). MS (*m/z*) 415 (M + 1).

10

15

EXAMPLE 38




methyl 4-((*E*)-2-{4-(4-methylpiperazin-1-yl)-6-[(3-methyl-1*H*-pyrazol-5-yl)amino]-5-nitropyrimidin-2-yl} vinyl)benzoate

20

To a solution of Example 37 (2.0 g, 4.83 mmol) in tetrahydrofuran (30 mL) was added triethylamine (1.0 mL, 7.3 mmol) under nitrogen atmosphere at room temperature (30 °C). After 15 min, N-methylpiperazine (483 mg, 4.83

mmol) in tetrahydrofuran (10 mL) was added drop-wise to the reaction mixture. After 0.5 hours, the reaction mixture was evaporated to dryness. The solid obtained was triturated with ethyl acetate and hexane to obtain Example 38 (2.2 g). A 500 mg of the crude compound was purified by preparative HPLC to give
 5 pure Example 38 (150mg). ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.53 (br, 1H), 10.00 (br, 1H), 7.9-8.1(m, 5H), 7.4 (m, 1H, *J*= 16 Hz), 6.75 (s, 1H), 3.88 (s, 3H), 3.0-3.2(m, 4H), 2.9(m, 4H), 2.3(s, 3H), 2.09(s, 3H). MS (*m/z*) 479 (M + 1)

EXAMPLE 39




10

methyl 4-((*E*)-2-{5-amino-4-(4-methylpiperazin-1-yl)-6-[(3-methyl-1*H*-pyrazol-5-yl)amino]-pyrimidin-2-yl}vinyl)benzoate

Tin (II) chloride dihydrate (1.0 g, 4.4 mmol) was dissolved in conc.
 15 hydrochloric acid (1 mL, 9.04 mmol). The reaction mixture was stirred for 10 minutes and was then cooled to <10 °C. Example 38 (250 mg, 0.52 mmol) in methanol (20 mL) was added drop-wise to the reaction mixture. The cooling was removed and the reaction was stirred overnight. The reaction mixture was evaporated to 1/3rd volume then diluted with 40 mL of ethyl acetate. A solution of
 20 1N NaOH (20 mL) was added to the reaction mixture. The organic layer was then separated, dried and evaporated under reduced pressure. The crude solid obtained was then purified by preparative HPLC to give 30mg of Example 39. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.89 (br, 1H), 8.93(br, 1H), 7.96(d, 2H, *J*=8Hz), 7.77(d, 2H, *J*= 8Hz), 7.57(d, 1H, *J*= 16Hz), 7.24(br, 1H), 7.17(d, 1H, *J*=16Hz), 7.11(br, 1H), 6.99(br, 1H), 6.58(s, 1H), 3.88 (s, 3H), 3.1-3.8 (m, 8H), 2.88 (s, 3H), 2.31 (s,
 25 3H). MS (*m/z*) 449 (M + 1).

EXAMPLE 31 VIA SCHEME 6



Synthesis of 6-(4-methylpiperazin-1-yl)-*N*-(3-methyl-1*H*-pyrazol-5-yl)-2-[(*E*)-2-phenylvinyl]pyrimidin-4-amine via Scheme 6

5

Following the general narrative described above for Scheme 6, commercially-available cinnamionitrile (70g) was dissolved in anhydrous toluene (1.19L) and absolute ethanol (287 ml, 0.91 mol, 9 eq). The clear solution obtained was cooled to -5°C and dry HCl gas was gently bubbled for 2 hours after which the reaction was closed to seal and stirred for 15 hours at 0°C , workup of the reaction mixture provided O-ethyl imidate HCl salt (100 gm); $^1\text{H-NMR}$ (200MHz, DMSO- D_6): 11.63 (2H, bs), 7.98 (1H, d, $J = 16.2\text{Hz}$), 7.73-7.32 (5H, m), 7.01 (1H, d, $J = 16.2\text{Hz}$), 4.46 (2H, q), 1.41 (3H, t, $J = 5.2\text{Hz}$). O-Ethyl imidate HCl salt (100g) in ethanol (500ml) was cooled to 0°C and a methanol solution of dry ammonia (204 ml, 7N, 0.30 mol) was added. The mixture was stirred at room temperature for 12 hours. The mixture was worked up to provide intermediate-1 (80g, 89%); $^1\text{H-NMR}$ (200MHz, DMSO- D_6): 9.32 (2H, bs), 8.84 (2H, bs), 7.97 (1H, d, $J = 16.2\text{Hz}$), 7.62-7.44 (5H, m), 6.81 (1H, d, $J = 16.2\text{Hz}$); M^+ 147 (100%, m/z).

Intermediate-1 (80g) in methanol (80ml) had dimethyl malonate (80ml, 1.1eq) added to it at room temperature. The mixture was cooled to 0°C and NaOCH₃ (44g, 4.4eq) was added slowly over 10 minutes. The light yellow-white solution was heated to 90°C and refluxed for 4 hours. Work up provided intermediate-2 (70g, 60%); $^1\text{H-NMR}$ (200MHz, DMSO- D_6): 11.62 (2H, bs), 7.87 (1H, d, $J = 16.2\text{Hz}$), 7.60-7.41 (5H, m), 6.85 (1H, d, $J = 16.2\text{Hz}$), 5.22 (1H, s); M^+ 215 (100%, m/z).


Intermediate-2 (70g) was slowly added in portions to 600 ml of POCl₃. The mixture was stirred for 4-6 hrs at 100°C . The reaction mixture was concentrated under reduced pressure at 60°C until dryness. Work up gave

intermediate-3 (70g, 85%); ¹H-NMR (200MHz, DMSO-D₆): 7.96 (1H, d, *J* = 16.2Hz), 7.80 (2H, m), 7.45-7.21 (5H, m); M+ 250.9, 252.9 (100%, m/z).

To a solution of intermediate-3 (70g, 0.2278 mole) in anhydrous DMA (400ml) was added 5-methyl-3-aminopyrazole (32.50g, 0.334 mol), NaI (62.33g, 0.418 mole) and DIPEA (54g, 1.48 mol) and the mixture was stirred at 90°C for 12h. Work up and chromatographic purification gave compound-4 (50g, 57%); ¹H-NMR (200MHz, DMSO-D₆): 12.12 (1H, s), 10.18 (1H, s), 7.83 (1H, d, *J* = 16.2Hz), 7.71 (3H, m), 7.45-7.33 (3H, m), 7.12 (1H, d, *J* = 16.2Hz), 6.22 (1H, bs), 2.21 (3H, s); M+ 312 (100%, m/z).

Intermediate-4 (50g) was dissolved in N-methyl piperazine (150 ml) and heated at 110°C for 1 hour. Work up and crystallization provided the target Example (25g, 41.4%); ¹H-NMR (200MHz, DMSO-D₆): 11.92 (1H, s), 9.18 (1H, s), 7.76 (1H, d, *J* = 16.2Hz), 7.62 (2H, m), 7.41-7.35 (3H, m), 6.91 (1H, d, *J* = 16.2Hz), 6.61 (1H, bs), 6.01 (1H, bs), 3.57 (4H, m), 2.41 (4H, m), 2.22 (3H, s), 2.20 (3H, s); HPLC (98.1% pure, RT 24.36min, Gradient, 0.1%TFA in Acetonitrile and 0.1%TFA in Water, Hypersil BDS C-18, 4.6x150mm, 5.0u).

EXAMPLE 49




2-(4-chlorostyryl)-N-(5-methyl-1H-pyrazol-3-yl)-6-(4-methylpiperazin-1-yl)pyrimidin-4-amine

20

Example 49 was synthesized via Scheme 6 according to the general scheme provided above with the appropriate starting materials 3-(4-chlorophenyl)acrylonitrile, 5-methyl-1H-pyrazole-3-amine, and 1-methylpiperazine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.

25

EXAMPLE 50




2-(3,5-dimethoxystyryl)-N-(5-methyl-1H-pyrazol-3-yl)-
6-(4-methylpiperazin-1-yl)pyrimidin-4-amine

Example 50 was synthesized via Scheme 6 according to the general
5 scheme provided above with the appropriate starting materials 3-(3,5-
dimethoxyphenyl)acrylonitrile, 5-methyl-1H-pyrazole-3-amine, and 1-methyl-
piperazine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is
attached.

10


EXAMPLE 60



2-((E)-2-(furan-2-yl)vinyl)-N-(5-methyl-1H-pyrazol-3-yl)-
6-(4-methylpiperazin-1-yl)pyrimidin-4-amine

Example 60 was synthesized via Scheme 6 according to the general
scheme provided above with the appropriate starting materials 3-(furan-2-
15 yl)acrylonitrile, 5-methyl-1H-pyrazole-3-amine, and 1-methylpiperazine.
Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.


EXAMPLE 64



N-(5-methyl-1*H*-pyrazol-3-yl)-6-morpholino-2-styrylpyrimidin-4-amine

Example 64 was synthesized via Scheme 6 according to the general
 5 scheme provided above with the appropriate starting materials cinnamionitrile, 5-
 methyl-1*H*-pyrazole-3-amine, and morpholine. Structure of the target was
 confirmed by ¹H-NMR. The ¹H-NMR is attached.


EXAMPLE 66



10 *N*⁴-(2-(dimethylamino)ethyl)-*N*⁶-(5-methyl-1*H*-pyrazol-3-yl)-2-styrylpyrimidine-4,6-diamine

Example 66 was synthesized via Scheme 6 according to the general
 scheme provided above with the appropriate starting materials cinnamionitrile, 5-
 methyl-1*H*-pyrazole-3-amine, and *N*¹,*N*¹-dimethylethane-1,2-diamine. Structure
 15 of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.

EXAMPLE 67




20 *N*-(5-methyl-1*H*-pyrazol-3-yl)-6-(piperidin-1-yl)-2-styrylpyrimidin-4-amine

Example 67 was synthesized via Scheme 6 according to the general
 scheme provided above with the appropriate starting materials cinnamionitrile, 5-

methyl-1H-pyrazole-3-amine, and piperidine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.


EXAMPLE 69



5 6-(2-(dimethylamino)ethoxy)-N-(5-methyl-1H-pyrazol-3-yl)-2-styrylpyrimidin-4-amine

Example 69 was synthesized via Scheme 6 according to the general scheme provided above with the appropriate starting materials cinnamonditrile, 5-methyl-1H-pyrazole-3-amine, and (2-dimethyl)aminoethanol. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.


EXAMPLE 71



N-(5-methyl-1H-pyrazol-3-yl)-6-(4-methylsulfonylpiperazin-1-yl)-2-styrylpyrimidin-4-amine

15 Example 71 was synthesized via Scheme 6 according to the general scheme provided above with the appropriate starting materials cinnamonditrile, 5-methyl-1H-pyrazole-3-amine, and 4-methylsulfonylpiperazine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.


EXAMPLE 76



N-(5-methyl-1*H*-pyrazol-3-yl)-6-(pyrrolidin-1-yl)-2-styrylpyrimidin-4-amine

Example 76 was synthesized via Scheme 6 according to the general
 5 scheme provided above with the appropriate starting materials cinnamionitrile, 5-methyl-1*H*-pyrazole-3-amine, and pyrrolidine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.

EXAMPLE 77




*N*⁴-(5-methyl-1*H*-pyrazol-3-yl)-*N*⁶-(2-(4-methylpiperazin-1-yl)ethyl)-
 2-styrylpyrimidine-4,6-diamine

10

Example 77 was synthesized via Scheme 6 according to the general
 scheme provided above with the appropriate starting materials cinnamionitrile, 5-methyl-1*H*-pyrazole-3-amine, and 2-(4'-methylpiperazin-1'-yl)-aminoethane.
 15 Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.

EXAMPLE 82




*N*⁴-(5-methyl-1*H*-pyrazol-3-yl)-*N*⁶-phenyl-2-styrylpyrimidine-4,6-diamine

20 Example 82 was synthesized via Scheme 6 according to the general
 scheme provided above with the appropriate starting materials cinnamionitrile, 5-

methyl-1H-pyrazole-3-amine, and aniline. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.


EXAMPLE 83



5 *N*-(5-methyl-1*H*-pyrazol-3-yl)-2-styryl-6-thiomorpholinopyrimidin-4-amine

Example 83 was synthesized via Scheme 6 according to the general scheme provided above with the appropriate starting materials cinnamitrile, 5-methyl-1H-pyrazole-3-amine, and thiomorpholine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.


EXAMPLE 86



*N*⁴-(4-fluorophenyl)-*N*⁶-(5-methyl-1*H*-pyrazol-3-yl)-2-styrylpyrimidine-4,6-diamine

15 Example 86 was synthesized via Scheme 6 according to the general scheme provided above with the appropriate starting materials cinnamitrile, 5-methyl-1H-pyrazole-3-amine, and 4-fluoroaniline. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.


EXAMPLE 87



*N*⁴-(2-(dimethylamino)ethyl)-*N*⁴-methyl-*N*⁶-(5-methyl-1*H*-pyrazol-3-yl)-2-styrylpyrimidine-4,6-diamine

Example 87 was synthesized via Scheme 6 according to the general scheme provided above with the appropriate starting materials cinnamionitrile, 5-methyl-1*H*-pyrazole-3-amine, and *N*¹,*N*¹,*N*²-trimethylethane-1,2-diamine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.


EXAMPLE 99



*N*⁴-(tetrahydro-2*H*-pyran-4-yl)-*N*⁶-(5-methyl-1*H*-pyrazol-3-yl)-2-styrylpyrimidine-4,6-diamine

Example 99 was synthesized via Scheme 6 according to the general scheme provided above with the appropriate starting materials cinnamionitrile, 5-methyl-1*H*-pyrazole-3-amine, and tetrahydro-2*H*-pyran-4-amine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.


EXAMPLE 101

1-(6-(5-methyl-1*H*-pyrazol-3-ylamino)-2-styrylpyrimidin-4-yl)piperidin-4-ol

Example 101 was synthesized via Scheme 6 according to the general
 5 scheme provided above with the appropriate starting materials cinnamionitrile, 5-methyl-1*H*-pyrazole-3-amine, and O-protected piperidin-4-ol. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.


EXAMPLE 103

10

2-(4-bromostyryl)-*N*-(5-methyl-1*H*-pyrazol-3-yl)-
6-(4-methylpiperazin-1-yl)pyrimidin-4-amine

Example 103 was synthesized via Scheme 6 according to the general
 scheme provided above with the appropriate starting materials 3-(4-
 15 bromophenyl)-acrylonitrile, 5-methyl-1*H*-pyrazole-3-amine, and 4-methyl-
 piperazine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is
 attached.

EXAMPLE 160




2-(4-methoxystyryl)-N-(5-methyl-1H-pyrazol-3-yl)-
6-(4-methylpiperazin-1-yl)pyrimidin-4-amine

5 Example 160 was synthesized via Scheme 6 according to the general scheme provided above with the appropriate starting materials 3-(4-methoxyphenyl)-acrylonitrile, 5-methyl-1H-pyrazole-3-amine, and 4-methylpiperazine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.

10


EXAMPLE 161



2-(4-methoxystyryl)-N-(5-methyl-1H-pyrazol-3-yl)-
6-(4-methylpiperazin-1-yl)pyrimidin-4-amine

15 Example 161 was synthesized via Scheme 6 according to the general scheme provided above with the appropriate starting materials 3-(4-methoxyphenyl)-acrylonitrile, 5-methyl-1H-pyrazole-3-amine, and 4-methylpiperazine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.

EXAMPLE 162




*N*⁴-((furan-2-yl)methyl)-*N*⁶-(5-methyl-1*H*-pyrazol-3-yl)-2-styrylpyrimidine-4,6-diamine

5 Example 162 was synthesized via Scheme 6 according to the general scheme provided above with the appropriate starting materials 3-(4-methoxyphenyl)acrylonitrile, 5-methyl-1*H*-pyrazole-3-amine, and (furan-2-yl)methanamine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.

10


EXAMPLE 163



*N*⁴-(2-methoxyethyl)-*N*⁶-(5-methyl-1*H*-pyrazol-3-yl)-2-styrylpyrimidine-4,6-diamine

15 Example 163 was synthesized via Scheme 6 according to the general scheme provided above with the appropriate starting materials cinnamionitrile, 5-methyl-1*H*-pyrazole-3-amine, and 2-methoxyethanamine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.

EXAMPLE 164



*N*⁴-((tetrahydrofuran-2-yl)methyl)-*N*⁶-(5-methyl-1*H*-pyrazol-3-yl)-2-styrylpyrimidine-4,6-diamine

5 Example 164 was synthesized via Scheme 6 according to the general scheme provided above with the appropriate starting materials cinnamionitrile, 5-methyl-1*H*-pyrazole-3-amine, and (tetrahydrofuran-2-yl)methanamine. Structure of the target was confirmed by ¹H-NMR. The ¹H-NMR is attached.

10

BIOLOGICAL TESTING

Biological Testing Example 1

Aurora A (Aurora 2) Inhibition Assay

15 Compounds were tested for their potency against recombinant Aurora A (Upstate, Lake Placid, NY) using the PanVera Z'-Lyte kinase assay kit – Ser/Thr 1 peptide (Invitrogen, Carlsbad, CA). Assays were carried out in kinase assay buffer (50 mM HEPES, pH 7.5, 10 mM MgCl₂, 5 mM EGTA, 0.05% Brij-35, 2 mM DTT). Test compounds were initially dissolved in DMSO at 100X the
20 highest tested concentration, then serially diluted to 4X test concentrations in kinase assay buffer. Next, Aurora A (final concentration 200-500 ng/mL), Z'-Lyte Ser/Thr 1 peptide (final concentration 2 μM) and ATP (final concentration 10 μM) were added according to the manufacturer's instructions. Assays were carried out in half-area 96-well white polystyrene assay plates (Corning, Corning,
25 NY) in a final volume of 20 μl. The reaction was allowed to proceed for 1 h at room temperature in the dark, at which point the development reagent and stop reagent were added according to the manufacturer's instructions. Coumarin (Ex. 400 nm, Em. 465 nm) and fluorescein (Ex. 400 nm, Em. 565 nm) fluorescence values were measured on a SpectraFluor Plus plate reader (Tecan, Durham, NC).

The emission ratio (coumarin/fluorescein) was determined and used to calculate the percent phosphorylation for each well. Wells containing substrate but no kinase and wells containing a phosphopeptide control were used to set 0% and 100% phosphorylation values, respectively. Typically 20-40% of the substrate was phosphorylated in wells without inhibitor. Dose-response curves of relative Aurora A activity vs. inhibitor concentration were plotted with Grafit (Erithacus Software, Horley, Surrey, UK).

Table 2 shows representative data for the inhibition of Aurora A by the compounds of this invention at a concentration of 100 μ M.

10

Table 2

Example No.	% Inhibition of Aurora A @ 100 μ M
4	91
10	100
11	98
12	100
13	93
14	98
15	100
17	62
18	100
19	99
20	77
21	100
22	97
23	99
24	100
25	100
26	99
30	100
31	100
32	100
34	99
37	80
38	99
39	100
49	100
50	100
60	100
64	100
66	100
67	100
69	100

71	100
77	100
82	100
83	100
86	100
99	100
100	100
103	100
160	100
161	100
162	100
163	100
164	100
169	95
170	100
171	89
172	99
173	100

Biological Testing Example 2

5

Aurora B (Aurora 1) Inhibition Assay

Assays for Aurora B kinase inhibition were carried out similarly to those for Aurora A kinase (see above) with the following modifications. Aurora B kinase (BPS Biosciences, San Diego, CA) was used as the enzyme, at a concentration was 2.5 $\mu\text{g/mL}$. The ATP concentration was 50 μM , and the kinase reaction was allowed to proceed for 16 h. Sodium orthovanadate (20 μM) was added to the buffer to inhibit contaminating phosphatases. Table 3 shows data for the inhibition of Aurora B by the compounds of this invention at a concentration of 100 μM .

Table 3

Example No.	% Inhibition of Aurora B @ 100 μ M
10	97
18	100
26	95
30	97
31	100
32	99
60	99
100	100
163	100
164	100

Biological Testing Example 3

5

Abl Kinase Inhibition Assay

Compounds were assayed for Abl kinase inhibitory activity using N-terminal His₆-tagged recombinant human Abl, residues 27-end (Upstate USA Inc, 706 Forest Street, Charlottesville, Virginia). Serial dilutions of compound were assayed in a final reaction volume of 25 μ L by incubating a solution of the above Abl kinase (5-10 mU), 8 mM MOPS (3-(N-morpholino) propanesulfonic acid) pH 7.0, 0.2 mM EDTA (ethylenediamine tetracetic acid), 50 μ M of amino acid sequence EAIYAAPFAKKK (Upstate USA Inc., Charlottesville, Virginia), and 10 mM magnesium acetate and [γ -³³P-ATP] (specific activity of about 500 cpm/pmol, concentration as required). The reaction was initiated by the addition of the magnesium acetate and [γ -³³P-ATP] mixture. After incubation for 40 minutes at room temperature, the reaction was stopped by the addition of 5 μ L of a 3% phosphoric acid solution. A 10 μ L aliquote of the reaction was then spotted onto a P30 filtermat (PerkinElmer, Wellesley, MA) and washed three times for five minutes in 75 mM phosphoric acid and once in methanol prior to drying and scintillation counting. Inhibition of Abl activity was determined by comparison to assays that contained no inhibitor. Under these conditions, Example 31 produced about 77% inhibition of Abl at a concentration of 1 μ M.

25

Biological Testing Example 4

cKit Kinase Inhibition Assay

5 Compounds were assayed for cKit kinase inhibitory activity using N-terminal GST-tagged recombinant human cKit, residues 544-end (Upstate USA Inc, 706 Forest Street, Charlottesville, Virginia). Serial dilutions of compound were assayed in a final reaction volume of 25 μ L by incubating a solution of the above cKit kinase (5-10 mU), 8 mM MOPS (3-(N-morpholino) propanesulfonic acid) pH 7.0, 0.2 mM EDTA (ethylenediamine tetracetic acid), 10 mM MnCl_2 , 0.1 mg/mL polyglutamic acid-tyrosine 4:1, and 10 mM magnesium acetate and [γ - ^{33}P -ATP] (specific activity of about 500 cpm/pmol, concentration as required). The reaction was initiated by the addition of the magnesium acetate and [γ - ^{33}P -ATP] mixture. After incubation for 40 minutes at room temperature, the reaction was stopped by the addition of 5 μ L of a 3% phosphoric acid solution. A 10 μ L aliquote of the reaction was then spotted onto a Filtermat A and washed three times for five minutes in 75 mM phosphoric acid and once in methanol prior to drying and scintillation counting. Inhibition of cKit activity was determined by comparison to assays that contained no inhibitor. Under these conditions, Example 31 produced about 86% inhibition of cKit at a concentration of 1 μ M.

Biological Testing Example 5

Src Kinase Inhibition Assay

25

Compounds were assayed for Src kinase inhibitory activity using N-terminal His-tagged human Src (Upstate USA Inc, 706 Forest Street, Charlottesville, Virginia). Serial dilutions of compound were assayed in a final reaction volume of 25 μ L by incubating a solution of the above Src kinase (5-10 mU), 8 mM MOPS (3-(N-morpholino) propanesulfonic acid) pH 7.0, 0.2 mM EDTA (ethylenediamine tetracetic acid), 250 μ M amino acid sequence KVEKIGEGTYGVVYK (Upstate USA Inc, 706 Forest Street, Charlottesville, Virginia), and 10 mM magnesium acetate and [γ - ^{33}P -ATP] (specific activity of about 500 cpm/pmol, concentration as required). The reaction was initiated by the

addition of the magnesium acetate and [γ -³³P-ATP] mixture. After incubation for 40 minutes at room temperature, the reaction was stopped by the addition of 5 μ L of a 3% phosphoric acid solution. A 10 μ L aliquote of the reaction was then spotted onto a P30 filtermat (PerkinElmer, Wellesley, MA) and washed three
5 times for five minutes in 75 mM phosphoric acid and once in methanol prior to drying and scintillation counting. Inhibition of Src activity was determined by comparison to assays that contained no inhibitor. Under these conditions, Example 31 produced about 95% inhibition of Src at a concentration of 1 μ M.

10 Biological Testing Example 6

Flt3 Kinase Inhibition Assay

Compounds were assayed for Flt3 kinase inhibitory activity using N-
15 terminal GST-tagged recombinant human Flt3, residues 564-end (Upstate USA Inc, 706 Forest Street, Charlottesville, Virginia). Serial dilutions of compound were assayed in a final reaction volume of 25 μ L by incubating a solution of the above Flt3 kinase (5-10 mU), 8 mM MOPS (3-(N-morpholino) propanesulfonic acid) pH 7.0, 0.2 mM EDTA (ethylenediamine tetracetic acid), 50 μ M amino acid
20 sequence EAIYAAPFAKKK (Upstate USA Inc, 706 Forest Street, Charlottesville, Virginia), and 10 mM magnesium acetate and [γ -³³P-ATP] (specific activity of about 500 cpm/pmol, concentration as required). The reaction was initiated by the addition of the magnesium acetate and [γ -³³P-ATP] mixture. After incubation for 40 minutes at room temperature, the reaction was stopped by
25 the addition of 5 μ L of a 3% phosphoric acid solution. A 10 μ L aliquote of the reaction was then spotted onto a P30 filtermat (PerkinElmer, Wellesley, MA) and washed three times for five minutes in 75 mM phosphoric acid and once in methanol prior to drying and scintillation counting. Inhibition of Flt3 activity was determined by comparison to assays that contained no inhibitor. Under these
30 conditions, Example 31 produced about 96% inhibition of Flt3 at a concentration of 1 μ M.

Biological Testing Example 7

KDR Kinase Inhibition Assay

5 Compounds were assayed for KDR kinase inhibitory activity using N-terminal His₆-tagged recombinant human KDR, residues 790-end (Upstate USA Inc, 706 Forest Street, Charlottesville, Virginia). Serial dilutions of compound were assayed in a final reaction volume of 25 μ L by incubating a solution of the above KDR kinase (5-10 mU), 8 mM MOPS (3-(N-morpholino) propanesulfonic acid) pH 7.0, 0.2 mM EDTA (ethylenediamine tetracetic acid), 0.33 mg/mL myelin basic protein (Upstate USA Inc, 706 Forest Street, Charlottesville, Virginia), and 10 mM magnesium acetate and [γ -³³P-ATP] (specific activity of about 500 cpm/pmol, concentration as required). The reaction was initiated by the addition of the magnesium acetate and [γ -³³P-ATP] mixture. After incubation for 15 40 minutes at room temperature, the reaction was stopped by the addition of 5 μ L of a 3% phosphoric acid solution. A 10 μ L aliquote of the reaction was then spotted onto a P30 filtermat (PerkinElmer, Wellesley, MA) and washed three times for five minutes in 75 mM phosphoric acid and once in methanol prior to drying and scintillation counting. Inhibition of KDR activity was determined by 20 comparison to assays that contained no inhibitor. Under these conditions, Example 31 produced about 94% inhibition of KDR at a concentration of 1 μ M.

Biological Testing Example 8

25 Lck Kinase Inhibition Assay

 Compounds were assayed for Lck kinase inhibitory activity using recombinant, full-length human N-terminal His-tagged Lck, residues 790-end (Upstate USA Inc, 706 Forest Street, Charlottesville, Virginia). Serial dilutions of 30 compound were assayed in a final reaction volume of 25 μ L by incubating a solution of the above Lck kinase (5-10 mU), 50 mM Tris- pH 7.0, 0.1 mM EGTA (ethylene glycol bis[2-aminoethyl ether]tetracetic acid), 0.1 mM Na₃VO₄, 0.1 % β -mercaptoethanol, 0.1 mg/mL poly-glutamate-tyrosine 4:1, and 10 mM magnesium acetate and [γ -³³P-ATP] (specific activity of about 500 cpm/pmol,

concentration as required). The reaction was initiated by the addition of the magnesium acetate and [γ - ^{33}P -ATP] mixture. After incubation for 40 minutes at room temperature, the reaction was stopped by the addition of 5 μL of a 3% phosphoric acid solution. A 10 μL aliquote of the reaction was then spotted onto
5 a Filtermat A and washed three times for five minutes in 75 mM phosphoric acid and once in methanol prior to drying and scintillation counting. Inhibition of Lck activity was determined by comparison to assays that contained no inhibitor. Under these conditions, Example 31 produced about 98% inhibition of Lck at a concentration of 1 μM .

10

Biological Testing Example 9

Kinase Selectivity Assay

15

Compounds were assayed for inhibitory activity against a panel of different kinase enzymes at a concentration of 1 μM . Where appropriate, full-length enzymes or catalytically active fragments, with N-terminal or C-terminal
20 His- or GST-tags, were used (Invitrogen, Carlsbad, CA). Test compounds were initially diluted to 100X test concentration in 100% DMSO. This 100X concentration was then diluted to a 4X working concentration in Kinase Buffer (50 mM HEPES pH 7.5, 0.01% Brij-35, 10 mM MgCl_2 , 1 mM EGTA) and 2.5 μL was then added to a low volume NBS, 384-well plate (Corning, Corning, NY).
25 Five μL of a 2X peptide/kinase mixture (Invitrogen, Carlsbad, CA), as appropriate (listed in Table 3), was then added and the reaction was then initiated by addition of 2.5 μL of a 4X working concentration of ATP in Kinase Buffer. Inhibition was determined at an ATP concentration equivalent to the apparent K_m for each individual kinase, or at an ATP concentration of 100 μM if the apparent K_m could
30 not be reached. Table 4 shows the percent inhibition values obtained for Example 10 and Example 31 under these conditions.

Table 4

	Percent Inhibition at 1 μ M (Example 10)	Percent Inhibition at 1 μ M (Example 31)	Peptide Substrate
ABL1	54	81	Tyr 2
ABL1 E255K	53	74	Tyr 2
ABL1 G250E	60	85	Tyr 2
ABL1 T315I	79	91	Tyr 2
ABL1 Y253F	57	80	Tyr 2
ABL2 (Arg)	32	74	Tyr 2
AKT1 (PKB alpha)	9	5	Ser/Thr 6
AURKB (Aurora B)	37	75	Ser/Thr 1
BLK	67	90	Tyr 1
BMX	19	61	Tyr 1
BTK	28	63	Tyr 1
CDK2/cyclin A	8	8	Ser/Thr 12
CSF1R (FMS)	80	97	Tyr 1
DAPK3 (ZIPK)	93	68	Ser/Thr 13
EPHA1	52	77	Tyr 2
EPHB1	32	61	Tyr 1
FGFR1	46	77	Tyr 4
FGFR2	53	86	Tyr 4
FLT1 (VEGFR1)	25	53	Tyr 4
FLT3	93	98	Tyr 2
FLT3 D835Y	80	98	Tyr 2
FLT4 (VEGFR3)	87	97	Tyr 4
FYN	50	89	Tyr 2
GSK3A (GSK3 alpha)	38	36	Ser/Thr 9
GSK3B (GSK3 beta)	26	28	Ser/Thr 9
IRAK4	14	63	Ser/Thr 7
JAK2	61	91	Tyr 4
KDR (VEGFR2)	88	96	Tyr 1
KIT	39	75	Tyr 6
KIT T670I	44	79	Tyr 6
LCK	81	94	Tyr 2
LYN A	63	84	Tyr 2
MAP2K1 (MEK1)	23	3	Ser/Thr 3
NTRK1 (TRKA)	92	96	Tyr 1
PDGFRA (PDGFR alpha)	58	83	Tyr 4
PDGFRA D842V	9	24	Tyr 4
PDGFRA T674I	65	88	Tyr 4
PTK2 (FAK)	81	94	Tyr 1
RET	95	98	Tyr 2
ROS1	54	65	Tyr 1
SRC	85	96	Tyr 2
STK6 (Aurora A)	83	96	Ser/Thr 1
SYK	88	86	Tyr 2
TBK1	73	80	Ser/Thr 5
YES1	80	88	Tyr 2

Biological Testing Example 10

Whole Cell Cytotoxicity

Assay: Sulforhodamine B

5 Reference: Developmental Therapeutics Program NCI/NIH

<http://dtp.nci.nih.gov/branches/btb/ivclsp.html>

Human tumor-derived cell lines, HCT116 or MCF7 (ATCC) were plated in a 96 well plate in DMEM containing 10% fetal bovine serum and 2mM L-glutamine at a density of 500 HCT116 cells or 1,000 MCF7 cells per well and
10 incubated at 37°C, 5% CO₂, for 24 hours prior to the addition of experimental compounds. Compounds were added using the dilution series indicated to duplicate plates and the cells were incubated in media plus compound for 96 hours. An additional plate was fixed in 10 % TCA at the time of the addition of compound to provide a measurement of the cell population at time zero, the time
15 of drug addition. Following the 96 hour incubation, cells were fixed in situ by gently aspirating off the culture media and then adding 50ul of ice cold 10% TCA per well and incubation at 4°C for 60 minutes. The plates were washed with tap water five times and allowed to air dry for 5 minute. 50ul of a 0.4% (w/v) Sulforhodamine B solution in 1% (v/v) acetic acid was added per well and the
20 cells were incubated for 30 minutes at room temperature. Following staining, plates were washed four times with 1% acetic acid to remove any unbound dye and then allowed to air dry for 5 minutes. The stain was solubilized with 100ul of 10mM Tris pH 10.5 per well and placed on an orbital rotator for 5 minutes. The absorbance was read at 570 nm. Percentage growth was calculated using the
25 absorbance readings from the time zero plate (Tz) and the dilution series plate (C) which included a column of cells grown in media without compound as a control (C) using the formulas:

$$[(Ti-Tz)/(C-Tz)] \times 100 \text{ for concentrations for which } Ti \geq Tz$$

$$[(Ti-Tz)/Tz] \times 100 \text{ for concentrations for which } Ti < Tz.$$

30 Three dose response parameters were calculated for each experimental agent. Growth inhibition of 50 % (GI50) was calculated from $[(Ti-Tz)/(C-Tz)] \times 100 = 50$, which was the drug concentration resulting in a 50% reduction in the net protein increase (as measured by SRB staining) in control cells during the drug incubation. The drug concentration resulting in total growth inhibition (TGI) was

calculated from $T_i = T_z$. The LC50 (concentration of drug resulting in a 50% reduction in the measured protein at the end of the drug treatment as compared to that at the beginning) indicating a net loss of cells following treatment was calculated from $[(T_i - T_z)/T_z] \times 100 = -50$. Values are calculated for each of these

5 three parameters if the level of activity was reached; however, if the effect was not reached or was exceeded, the value for that parameter is expressed as greater or less than the maximum or minimum concentration tested.

Table 5 shows representative values for the inhibition of HCT-116 cell growth by the compounds of this invention at a concentration of 100 μ M.

10


Table 5

Example No.	% Inhibition of HCT-116 cell growth @ 100 μ M
4	87
10	99
11	99
12	99
15	91
17	93
18	80
19	96
20	97
21	93
22	85
23	91
24	99
25	95
26	98
30	99
31	99
32	87
34	98
39	97
49	75
50	98
60	98
64	98
66	98
67	98
69	99
71	98
77	99
82	98
83	98

86	98
99	98
100	96
103	84
160	97
161	93
162	98
163	98
164	96
170	99
172	99
173	98

What is claimed is:

1. A compound of formula I:



Formula I


or a pharmaceutically acceptable salt thereof, wherein:

R^x is $-T-R^3$ or $-L-Z-R^3$;

R^y is $-T'-R^{3'}$ or $-L'-Z-R^{3'}$, wherein $R^{3'}$ is $-R'$, $-\text{halo}$, $-\text{OR}'$, $-\text{C}(=\text{O})\text{R}'$, $-\text{CO}_2\text{R}'$, $-\text{COCOR}'$, $-\text{COCH}_2\text{COR}'$, $-\text{NO}_2$, $-\text{CN}$, $-\text{S}(\text{O})\text{R}'$, $-\text{S}(\text{O})_2\text{R}'$, $-\text{SR}'$, $-\text{N}(\text{R}^{4'})_2$, $-\text{CON}(\text{R}^{7'})_2$, $-\text{SO}_2\text{N}(\text{R}^{7'})_2$, $-\text{OC}(=\text{O})\text{R}'$, $-\text{N}(\text{R}^{7'})\text{COR}'$, $-\text{N}(\text{R}^{7'})\text{CO}_2$ (C_{1-6} aliphatic), $-\text{N}(\text{R}^{4'})\text{N}(\text{R}^{4'})_2$, $-\text{C}=\text{NN}(\text{R}^{4'})_2$, $-\text{C}=\text{N}-\text{OR}'$, $-\text{N}(\text{R}^{7'})\text{CON}(\text{R}^{7'})_2$, $-\text{N}(\text{R}^{7'})\text{SO}_2\text{N}(\text{R}^{7'})_2$, $-\text{N}(\text{R}^{4'})\text{SO}_2\text{R}'$, or $-\text{OC}(=\text{O})\text{N}(\text{R}')_2$;

T' is a valence bond or $-(\text{C}(\text{R}^{6'})_2)-\text{A}'$; A' is a valence bond or a C_{1-3} alkylidene chain wherein a methylene unit of said C_{1-3} alkylidene chain is optionally replaced by $-\text{O}-$, $-\text{S}-$, $-\text{N}(\text{R}^{4'})-$, $-\text{CO}-$, $-\text{CONH}-$, $-\text{NHCO}-$, $-\text{SO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{NHSO}_2-$, $-\text{CO}_2-$, $-\text{OC}(\text{O})-$, $-\text{OC}(\text{O})\text{NH}-$, or $-\text{NHCO}_2-$;

L' is $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{N}(\text{R}^{6a})\text{SO}_2-$, $-\text{SO}_2\text{N}(\text{R}^{6a})-$, $-\text{CO}-$, $-\text{CO}_2-$, $-\text{N}(\text{R}^{6a})\text{CO}-$, $-\text{N}(\text{R}^{6a})\text{C}(\text{O})\text{O}-$, $-\text{N}(\text{R}^{6a})\text{CON}(\text{R}^{6a})-$, $-\text{N}(\text{R}^{6a})\text{SO}_2\text{N}(\text{R}^{6a})-$, $-\text{N}(\text{R}^{6a})\text{N}(\text{R}^{6a})-$, $-\text{C}(\text{O})\text{N}(\text{R}^{6a})-$,



each R^{6a} is independently hydrogen or an optionally substituted C_{1-4} aliphatic group, or two R^{6a} groups on the same nitrogen atom may be taken together with the nitrogen atom to form a 3-6 membered heterocyclyl or heteroaryl ring;

each R^4 is independently $-\text{R}^7$, $-\text{COR}^7$, $-\text{CO}_2$ (optionally substituted C_{1-6} aliphatic), $-\text{CON}(\text{R}^7)_2$, or $-\text{SO}_2\text{R}^7$;

each R^7 is independently hydrogen or an optionally substituted C_{1-6} aliphatic group, or two R^7 on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring; and each R' is independently hydrogen, optionally substituted C_{1-6} aliphatic, optionally substituted C_{6-10} aryl, an optionally substituted heteroaryl ring having 5-10 ring atoms, or a an optionally substituted heterocyclyl ring having 5-10 ring atoms;

Q' is $-\text{CR}^{6''}=\text{CR}^{6''}-$ or --- , wherein said $-\text{CR}^{6''}=\text{CR}^{6''}-$ may be a cis or trans double bond or a mixture thereof;

R^1 is $-\text{T}-(\text{Ring D})$;

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic aryl or carbocyclyl ring wherein each substitutable ring carbon of Ring D is independently substituted by oxo, $-\text{T}-\text{R}^5$, or $-\text{V}-\text{Z}-\text{R}^5$;

T is a valence bond or $-(\text{C}(\text{R}^6)_2)-\text{A}-$;

A is a valence bond or a C_1 - C_3 alkylidene chain wherein a methylene unit of said C_{1-3} alkylidene chain is optionally replaced by $-\text{O}-$, $-\text{S}-$, $-\text{N}(\text{R}^4)-$, $-\text{CO}-$, $-\text{CONH}-$, $-\text{NHCO}-$, $-\text{SO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{NHSO}_2-$, $-\text{CO}_2-$, $-\text{OC}(\text{O})-$, $-\text{OC}(\text{O})\text{NH}-$, or $-\text{NHCO}_2-$;

Z is a C₁₋₄ alkylidene chain;

L is —O—, —S—, —SO—, —SO₂—, —N(R⁶)SO₂—, —SO₂N(R⁶)—, —N(R⁶)—, —CO—, —CO₂—, —N(R⁶)CO—, —N(R⁶)C(O)O—, —N(R⁶)CON(R⁶)—, —N(R⁶)SO₂N(R⁶)—, —N(R⁶)N(R⁶)—, —C(O)N(R⁶)—, —OC(O)N(R⁶)—, —C(R⁶)₂O—, —C(R⁶)₂—, —C(R⁶)₂SO—, —C(R⁶)₂SO₂—, —C(R⁶)₂SO₂N(R⁶)—, —C(R⁶)₂N(R⁶)—, —C(R⁶)₂N(R⁶)C(O)—, —C(R⁶)₂N(R⁶)C(O)O—, —C(R⁶)=NN(R⁶)—, —C(R⁶)=N—O—, —C(R⁶)₂N(R⁶)N(R⁶)—, —C(R⁶)₂N(R⁶)SO₂N(R⁶)—, or —C(R⁶)₂N(R⁶)CON(R⁶)—;

R² and R^{2'} are independently —R or —T-W-R⁶, or R² and R^{2'} taken together with their intervening atoms form a fused, 5-8 membered, unsaturated or partially unsaturated ring wherein each substitutable ring carbon of said fused ring formed by R² and R^{2'} is independently substituted by halo, oxo, —CN, —NO₂, R⁷, or —V-R⁶; R³ is —R, —halo, —OR, —C(=O)R, —CO₂R, —COCOR, —COCH₂COR, —NO₂, —CN, —S(O)R, —S(O)₂R, —SR, —N(R⁴)₂, —CON(R⁷)₂, —SO₂N(R⁷)₂, —OC(=O)R, —N(R⁷)COR, —N(R⁷)CO₂(C₁₋₆ aliphatic), —N(R⁴)N(R⁴)₂, —C=NN(R⁴)₂, —C=N—OR, —N(R⁷)CON(R⁷)₂, —N(R⁷)SO₂N(R⁷)₂, —N(R⁴)SO₂R, or —OC(=O)N(R)₂;

each R is independently hydrogen, optionally substituted C₁₋₆ aliphatic, or optionally substituted C₆₋₁₀ aryl;

each R⁴ is independently —R⁷, —COR⁷, —CO₂ (optionally substituted C₁₋₆ aliphatic), —CON(R⁷)₂, or —SO₂R⁷;

each R⁵ is independently —R, halo, —OR, —C(=O)R, —CO₂R, —COCOR, —NO₂, —CN, —S(O)R, —SO₂R, —SR, —N(R⁴)₂, —CON(R⁴)₂, —SO₂N(R⁴)₂, —OC(=O)R, —N(R⁴)COR, —N(R⁴)CO₂(optionally substituted C₁₋₆ aliphatic), —N(R⁴)N(R⁴)₂, —C=NN(R⁴)₂, —C=N—OR, —N(R⁴)CON(R⁴)₂, —N(R⁴)SO₂N(R⁴)₂, —N(R⁴)SO₂R, or —OC(=O)N(R⁴)₂;

V is $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{N}(\text{R}^6)\text{SO}_2-$, $-\text{SO}_2\text{N}(\text{R}^6)-$, $-\text{N}(\text{R}^6)-$,
 $-\text{CO}-$, $-\text{CO}_2-$, $-\text{N}(\text{R}^6)\text{CO}-$, $-\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$, $-\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$,
 $-\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$, $-\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$, $-\text{C}(\text{O})\text{N}(\text{R}^6)-$, $-\text{OC}(\text{O})\text{N}(\text{R}^6)-$,
 $-\text{C}(\text{R}^6)_2\text{O}-$, $-\text{C}(\text{R}^6)_2\text{S}-$, $-\text{C}(\text{R}^6)_2\text{SO}-$, $-\text{C}(\text{R}^6)_2\text{SO}_2-$,
 $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})-$,
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$, $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$,
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$, or
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$;

W is $-\text{C}(\text{R}^6)_2\text{O}-$, $-\text{C}(\text{R}^6)_2\text{S}-$, $-\text{C}(\text{R}^6)_2\text{SO}-$, $-\text{C}(\text{R}^6)_2\text{SO}_2-$,
 $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$, $-\text{CO}-$, $-\text{CO}_2-$,
 $-\text{C}(\text{R}^6)\text{OC}(\text{O})-$, $-\text{C}(\text{R}^6)\text{OC}(\text{O})\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CO}-$,
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$, $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$,
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$,
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$, or $-\text{CON}(\text{R}^6)-$;







each R^6 is independently hydrogen or an optionally substituted C_{1-4} aliphatic group;

each $\text{R}^{6'}$ is independently hydrogen or a C_{1-4} aliphatic group, or two $\text{R}^{6'}$ on the same carbon atom are taken together to form a 3-8 membered carbocyclic ring;

each $\text{R}^{6''}$ is independently hydrogen, a C_{1-4} aliphatic group, halogen, or optionally substituted aryl, or two $\text{R}^{6''}$ on adjacent carbon atoms are taken together to form a 5-7 membered carbocyclic ring; and







each R^7 is independently hydrogen or an optionally substituted C_{1-6} aliphatic group.









2. A compound selected from the group consisting of:













Example No.	Structure	Example No.	Structure
4		5	
6		7	
8		9	













10		11	
12		13	
14		15	
17		18	
19		20	
21		22	













23		24	
25		26	
29		30	
31		32	
34		37	
38		39	

40		41	
	deleted		deleted
44			deleted
46			deleted
	deleted	49	
50			deleted

52		53	
	deleted	55	
56			deleted
58		59	
	deleted		deleted
62		63	













64		65	
66		67	
68		69	
70		71	
72		73	
74		75	






76		77	
78		79	
80		81	
82		83	
84		85	
86		87	





88		89	
90		91	
92		93	
94		95	
96		97	
98		99	













100		101	
102		103	
104		105	
106		107	
108		109	
110		111	







112		113	
114			deleted
	deleted	117	
118		119	
120		121	
122		123	

124		125	
126		127	
128		129	
130		131	
132		133	
134		135	

136		137	
138		139	
140			deleted
	deleted		deleted
	deleted		deleted
	deleted		deleted

	deleted		deleted
	deleted		deleted
	deleted		deleted
	deleted	155	
156		157	
158			deleted

160		161	
162		163	
164		165	
166		167	
168		169	
170		171	

172		173	
174		175	
176			deleted
178			deleted

and biologically acceptable salts thereof.


3. A composition comprising particles of a compound of claim 2, that are less than 2 microns average particle size and an excipient.

4. The compound of claim 2, wherein the compound is incorporated into a biodegradable or non-biodegradable polymer.

5. A composition comprising a compound of claim 2 and an additive.

6. The composition of claim 5, wherein the additive is selected from the group consisting of an anti-oxidant, a buffer, a bacteriostat, a liquid carrier, a solute, a suspending agent, a thickening agent, a flavoring agent, a gelatin, glycerin, a binder, a lubricant, an inert diluent, a preservative, a surface active agent, a dispersing agent, a biodegradable polymer, and combinations thereof.

7. The use of a compound for treating a patient with a disease wherein the compound is selected from a compound of Formula I:



Formula I

or a pharmaceutically acceptable salt thereof, wherein:

R^x is $-T-R^3$ or $-L-Z-R^3$;

R^y is $-T'-R^{3'}$ or $-L'-Z-R^{3'}$, wherein $R^{3'}$ is $-R'$, $-\text{halo}$, $-\text{OR}'$, $-\text{C}(=\text{O})\text{R}'$, $-\text{CO}_2\text{R}'$, $-\text{COCOR}'$, $-\text{COCH}_2\text{COR}'$, $-\text{NO}_2$, $-\text{CN}$, $-\text{S}(\text{O})\text{R}'$, $-\text{S}(\text{O})_2\text{R}'$, $-\text{SR}'$, $-\text{N}(\text{R}^{4'})_2$, $-\text{CON}(\text{R}^{7'})_2$, $-\text{SO}_2\text{N}(\text{R}^{7'})_2$, $-\text{OC}(=\text{O})\text{R}'$, $-\text{N}(\text{R}^{7'})\text{COR}'$, $-\text{N}(\text{R}^{7'})\text{CO}_2(\text{C}_{1-6} \text{ aliphatic})$, $-\text{N}(\text{R}^{4'})\text{N}(\text{R}^{4'})_2$,

$-\text{C}=\text{NN}(\text{R}^{4'})_2$, $-\text{C}=\text{N}-\text{OR}'$, $-\text{N}(\text{R}^{7'})\text{CON}(\text{R}^{7'})_2$, $-\text{N}(\text{R}^{7'})\text{SO}_2\text{N}(\text{R}^{7'})_2$,
 $-\text{N}(\text{R}^{4'})\text{SO}_2\text{R}'$, or $-\text{OC}(=\text{O})\text{N}(\text{R}')_2$;

T' is a valence bond or $-(\text{C}(\text{R}^{6'})_2)-\text{A}'$ -; A' is a valence bond or a C_1 - C_3 alkylidene chain wherein a methylene unit of said C_{1-3} alkylidene chain is optionally replaced by $-\text{O}-$, $-\text{S}-$, $-\text{N}(\text{R}^{4'})-$, $-\text{CO}-$, $-\text{CONH}-$, $-\text{NHCO}-$, $-\text{SO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{NHSO}_2-$, $-\text{CO}_2-$, $-\text{OC}(\text{O})-$, $-\text{OC}(\text{O})\text{NH}-$, or $-\text{NHCO}_2-$;

L' is $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{N}(\text{R}^{6a})\text{SO}_2-$, $-\text{SO}_2\text{N}(\text{R}^{6a})-$, $-\text{N}(\text{R}^{6a})-$, $-\text{CO}-$, $-\text{OC}_2-$, $-\text{N}(\text{R}^{6a})\text{CO}-$, $-\text{N}(\text{R}^{6a})\text{C}(\text{O})\text{O}-$, $-\text{N}(\text{R}^{6a})\text{CON}(\text{R}^{6a})-$, $-\text{N}(\text{R}^{6a})\text{SO}_2\text{N}(\text{R}^{6a})-$, $-\text{N}(\text{R}^{6a})\text{N}(\text{R}^{6a})-$, $-\text{C}(\text{O})\text{N}(\text{R}^{6a})-$, $-\text{OC}(\text{O})\text{N}(\text{R}^{6a})-$, $-\text{C}(\text{R}^{6a})_2\text{O}-$, $-\text{C}(\text{R}^{6a})_2-$, $-\text{C}(\text{R}^{6a})_2\text{SO}-$, $-\text{C}(\text{R}^{6a})_2\text{SO}_2-$, $-\text{C}(\text{R}^{6a})_2\text{SO}_2\text{N}(\text{R}^{6a})-$, $-\text{C}(\text{R}^{6a})_2\text{N}(\text{R}^{6a})-$, $-\text{C}(\text{R}^{6a})_2\text{N}(\text{R}^{6a})\text{C}(\text{O})-$, $-\text{C}(\text{R}^{6a})_2\text{N}(\text{R}^{6a})\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{R}^{6a})=\text{NN}(\text{R}^{6a})-$, $-\text{C}(\text{R}^{6a})=\text{N}-\text{O}-$, $-\text{C}(\text{R}^{6a})_2\text{N}(\text{R}^{6a})\text{N}(\text{R}^{6a})-$, $-\text{C}(\text{R}^{6a})_2\text{N}(\text{R}^{6a})\text{SO}_2\text{N}(\text{R}^{6a})-$, or $-\text{C}(\text{R}^{6a})_2\text{N}(\text{R}^{6a})\text{CON}(\text{R}^{6a})-$;

each R^{6a} is independently hydrogen or an optionally substituted C_{1-4} aliphatic group, or two R^{6a} groups on the same nitrogen atom may be taken together with the nitrogen atom to form a 3-6 membered heterocyclyl or heteroaryl ring;

each $\text{R}^{4'}$ is independently $-\text{R}^{7'}$, $-\text{COR}^{7'}$, $-\text{CO}_2$ (optionally substituted C_{1-6} aliphatic), $-\text{CON}(\text{R}^{7'})_2$, or $-\text{SO}_2\text{R}^{7'}$;

each $\text{R}^{7'}$ is independently hydrogen or an optionally substituted C_{1-6} aliphatic group, or two $\text{R}^{7'}$ on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring;
 and

each R' is independently hydrogen, optionally substituted C_{1-6} aliphatic, optionally substituted C_{6-10} aryl, an optionally substituted heteroaryl ring having 5-10 ring atoms, or a an optionally substituted heterocyclyl ring having 5-10 ring atoms;

Q' is $-\text{CR}^{6''}=\text{CR}^{6''}-$ or $-\equiv-$, wherein said $-\text{CR}^{6''}=\text{CR}^{6''}-$ may be a cis or trans double bond or a mixture thereof;

R¹ is -T-(Ring D);

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic aryl or carbocyclyl ring, wherein each substitutable ring carbon of Ring D is independently substituted by oxo, -T-R⁵, or -V-Z-R⁵;

T is a valence bond or $-(\text{C}(\text{R}^6)_2)\text{-A-}$;

A is a valence bond or a C₁-C₃ alkylidene chain wherein a methylene unit of said C₁₋₃ alkylidene chain is optionally replaced by -O-, -S-, -N(R⁴)-, -CO-, -CONH-, -NHCO-, -SO₂-, -SO₂NH-, -NHSO₂-, -CO₂-, -OC(O)-, -OC(O)NH-, or -NHCO₂-;

Z is a C₁₋₄ alkylidene chain;

L is -O-, -S-, -SO-, -SO₂-, -N(R⁶)SO₂-, -SO₂N(R⁶)-, -N(R⁶)-, -CO-, -CO₂-, -N(R⁶)CO-, -N(R⁶)C(O)O-, -N(R⁶)CON(R⁶)-, -N(R⁶)SO₂N(R⁶)-, -N(R⁶)N(R⁶)-, -C(O)N(R⁶)-, -OC(O)N(R⁶)-, -C(R⁶)₂O-, -C(R⁶)₂-, -C(R⁶)₂SO-, -C(R⁶)₂SO₂-C(R⁶)₂SO₂N(R⁶)-, -C(R⁶)₂N(R⁶)-, -C(R⁶)₂N(R⁶)C(O)-, -C(R⁶)₂N(R⁶)C(O)O-, -C(R⁶)=NN(R⁶)-, -C(R⁶)=N-O-, -C(R⁶)₂N(R⁶)N(R⁶)-, -C(R⁶)₂N(R⁶)SO₂N(R⁶)-, or -C(R⁶)₂N(R⁶)CON(R⁶)-;

R² and R^{2'} are independently -R or -T-W-R⁶, or R² and R^{2'} taken together with their intervening atoms form a fused, 5-8 membered, unsaturated or partially unsaturated ring wherein each substitutable ring carbon of said fused ring formed by R² and R^{2'} is independently substituted by halo, oxo, -CN, -NO₂, R⁷, or -V-R⁶;

R³ is -R, -halo, -OR, -C(=O)R, -CO₂R, -COCOR, -COCH₂COR, -NO₂, -CN, -S(O)R, -S(O)₂R, -SR, -N(R⁴)₂, -CON(R⁷)₂, -SO₂N(R⁷)₂, -OC(=O)R, -N(R⁷)COR, -N(R⁷)CO₂(C₁₋₆ aliphatic),

$-\text{N}(\text{R}^4)\text{N}(\text{R}^4)_2$, $-\text{C}=\text{NN}(\text{R}^4)_2$, $-\text{C}=\text{N}-\text{OR}$, $-\text{N}(\text{R}^7)\text{CON}(\text{R}^7)_2$, $-\text{N}(\text{R}^7)\text{SO}_2\text{N}(\text{R}^7)_2$, $-\text{N}(\text{R}^4)\text{SO}_2\text{R}$, or $-\text{OC}(=\text{O})\text{N}(\text{R})_2$;

each R is independently hydrogen, optionally substituted C_{1-6} aliphatic, or optionally substituted C_{6-10} aryl;

each R^4 is independently $-\text{R}^7$, $-\text{COR}'$, $-\text{CO}_2$ (optionally substituted C_{1-6} aliphatic), $-\text{CON}(\text{R}^7)_2$, or $-\text{SO}_2\text{R}^7$;

each R^5 is independently $-\text{R}$, halo, $-\text{OR}$, $-\text{C}(=\text{O})\text{R}$, $-\text{CO}_2\text{R}$, $-\text{COCOR}$, $-\text{NO}_2$, $-\text{CN}$, $-\text{S}(\text{O})\text{R}$, $-\text{SO}_2\text{R}$, $-\text{SR}$, $-\text{N}(\text{R}^4)_2$, $-\text{CON}(\text{R}^4)_2$, $-\text{SO}_2\text{N}(\text{R}^4)_2$, $-\text{OC}(=\text{O})\text{R}$, $-\text{N}(\text{R}^4)\text{COR}$, $-\text{N}(\text{R}^4)\text{CO}_2$ (optionally substituted C_{1-6} aliphatic), $-\text{N}(\text{R}^4)\text{N}(\text{R}^4)_2$, $-\text{C}=\text{NN}(\text{R}^4)_2$, $-\text{C}=\text{N}-\text{OR}$, $-\text{N}(\text{R}^4)\text{CON}(\text{R}^4)_2$, $-\text{N}(\text{R}^4)\text{SO}_2\text{N}(\text{R}^4)_2$, $-\text{N}(\text{R}^4)\text{SO}_2\text{R}$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^4)_2$;

V is $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{N}(\text{R}^6)\text{SO}_2-$, $-\text{SO}_2\text{N}(\text{R}^6)-$, $-\text{N}(\text{R}^6)-$, $-\text{CO}-$, $-\text{CO}_2-$, $-\text{N}(\text{R}^6)\text{CO}-$, $-\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$, $-\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$, $-\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$, $-\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$, $-\text{C}(\text{O})\text{N}(\text{R}^6)-$, $-\text{OC}(\text{O})\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{O}-$, $-\text{C}(\text{R}^6)_2\text{S}-$, $-\text{C}(\text{R}^6)_2\text{SO}-$, $-\text{C}(\text{R}^6)_2\text{SO}_2-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$, $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$, or $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$;

W is $-\text{C}(\text{R}^6)_2\text{O}-$, $-\text{C}(\text{R}^6)_2\text{S}-$, $-\text{C}(\text{R}^6)_2\text{SO}-$, $-\text{C}(\text{R}^6)_2\text{SO}_2-$, $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$, $-\text{CO}-$, $-\text{CO}_2-$, $-\text{C}(\text{R}^6)\text{OC}(\text{O})-$, $-\text{C}(\text{R}^6)\text{OC}(\text{O})\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CO}-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$, $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$, or $-\text{CON}(\text{R}^6)-$;

each R^6 is independently hydrogen or an optionally substituted C_{1-4} aliphatic group;

each $R^{6'}$ is independently hydrogen or a C_{1-4} aliphatic group, or two $R^{6'}$ on the same carbon atom are taken together to form a 3-8 membered carbocyclic ring;













each $R^{6''}$ is independently hydrogen, a C_{1-4} aliphatic group, halogen, or optionally substituted aryl, or two $R^{6''}$ on adjacent carbon atoms are taken together to form a 5-7 membered carbocyclic ring; and











each R^7 is independently hydrogen or an optionally substituted C_{1-6} aliphatic group; wherein the disease is selected from the group consisting of asthma, undesirable neovascularization and cancer, wherein the cancer is selected from the group consisting of: solid tumor, blood borne tumor, breast, ovary, cervix, prostate, testis, genitourinary tract, esophagus, larynx, glioblastoma, neuroblastoma, stomach, skin, keratoacanthoma, lung, epidermoid carcinoma, large cell carcinoma, small cell carcinoma, lung adenocarcinoma, bone, colon, adenoma, pancreas, adenocarcinoma, thyroid, follicular carcinoma, undifferentiated carcinoma, papillary carcinoma, seminoma, melanoma, sarcoma, bladder carcinoma, liver carcinoma and biliary passages, kidney carcinoma, myeloid disorders, lymphoid disorders, Hodgkin's, hairy cells, buccal cavity, pharynx, lip, tongue, mouth, pharynx, small intestine, colon-rectum, large intestine, rectum, brain and central nervous system, and leukemia.













8. The use of the compound or biologically acceptable salts of claim 2, for treating a patient with a disease; wherein the disease is selected from the group consisting of asthma, undesirable neovascularization, and cancer; wherein the cancer is selected from the group consisting of a solid tumor, blood borne tumor, breast, ovary, cervix, prostate, testis, genitourinary tract, esophagus, larynx, glioblastoma, neuroblastoma, stomach, skin, keratoacanthoma, lung, epidermoid carcinoma, large cell carcinoma, small cell carcinoma, lung adenocarcinoma, bone, colon, adenoma, pancreas, adenocarcinoma, thyroid, follicular carcinoma, undifferentiated carcinoma, papillary carcinoma, seminoma, melanoma, sarcoma,













bladder carcinoma, liver carcinoma and biliary passages, kidney carcinoma, myeloid disorders, lymphoid disorders, Hodgkin's, hairy cells, buccal cavity, pharynx, lip, tongue, mouth, pharynx, small intestine, colon-rectum, large intestine, rectum, brain and central nervous system, and leukemia.













9. The use of a compound for treating a patient with a disease, wherein the compound is selected from the group consisting of:


			
			
			
			
			
			


	
	
	
	
	


	
	
	
	
	
	


	
	
	
	
	
	













	
	
	
	
	
	












and biologically acceptable salts thereof, wherein the disease is selected from the group consisting of asthma, undesirable neovascularization, and cancer; wherein the cancer is selected from the group consisting of a solid tumor, blood borne tumor, breast, ovary, cervix, prostate, testis, genitourinary tract, esophagus, larynx, glioblastoma, neuroblastoma, stomach, skin, keratoacanthoma, lung, epidermoid carcinoma, large cell carcinoma, small cell carcinoma, lung adenocarcinoma, bone, colon, adenoma, pancreas, adenocarcinoma, thyroid, follicular carcinoma, undifferentiated carcinoma, papillary carcinoma, seminoma, melanoma, sarcoma, bladder carcinoma, liver carcinoma and biliary passages, kidney carcinoma, myeloid disorders, lymphoid disorders, Hodgkin's, hairy cells, buccal cavity, pharynx, lip, tongue, mouth, pharynx, small intestine, colon-rectum, large intestine, rectum, brain and central nervous system, and leukemia.

10. The use according to claim 7, wherein the disease associated with undesirable neovascularization is selected from the group consisting of ocular neovascular disease, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, neovascular glaucoma and retrolental fibroplasias, epidemic

keratoconjunctivitis, Vitamin A deficiency, contact lens overwear, atopic keratitis, superior limbic keratitis, pterygium keratitis sicca, Sjögren's syndrome, acne rosacea, phlyctenulosis, syphilis, *Mycobacteria* infections, lipid degeneration, chemical burns, bacterial ulcers, fungal ulcers, *Herpes simplex* infections, *Herpes zoster* infections, protozoan infections, Kaposi's sarcoma, Mooren's ulcer, Terrien's marginal degeneration, marginal keratolysis, trauma, rheumatoid arthritis, systemic lupus, polyarteritis, Wegener's sarcoidosis, Scleritis, Steven-Johnson disease, pemphigoid, radial keratotomy, or corneal graft rejection, sickle cell anemia, sarcoid, pseudoxanthoma elasticum, Paget's disease, vein occlusion, artery occlusion, carotid obstructive disease, chronic uveitis/vitritis, Lyme's disease, systemic lupus erythematosus, Eales' disease, Bechet's disease, infections causing a retinitis or choroiditis, presumed ocular histoplasmosis, Best's disease, myopia, optic pits, Stargart's disease, pars planitis, chronic retinal detachment, hyperviscosity syndromes, toxoplasmosis, and post-laser complications.

11. The use according to claim 7, wherein the compound is in an administrable form; wherein the administrable form is selected from the group consisting of a tablet, a capsule, a lozenge, a cachet, a solution, a suspension, an emulsion, a powder, an aerosol, a suppository, a spray, a pastille, an ointment, a cream, a paste, a foam, a gel, a tampon, a pessary, a granule, a bolus, a mouthwash, and a transdermal patch.

