



(86) Date de dépôt PCT/PCT Filing Date: 2014/10/17
(87) Date publication PCT/PCT Publication Date: 2016/02/25
(85) Entrée phase nationale/National Entry: 2017/02/13
(86) N° demande PCT/PCT Application No.: US 2014/061005
(87) N° publication PCT/PCT Publication No.: 2016/028328
(30) Priorité/Priority: 2014/08/19 (US62/039,128)

(51) Cl.Int./Int.Cl. *C07D 231/14* (2006.01),
A01N 43/46 (2006.01), *A61K 31/415* (2006.01)
(71) Demandeur/Applicant:
DOW AGROSCIENCES LLC, US
(72) Inventeurs/Inventors:
YANG, QIANG, US;
LORSBACH, BETH, US;
PODHOREZ, DAVID E., US
(74) Agent: SMART & BIGGAR

(54) Titre : PROCEDE DE PREPARATION DE 3-(3-CHLORO-1H-PYRAZOL-1-YL)PYRIDINE
(54) Title: PROCESS FOR THE PREPARATION OF 3-(3-CHLORO-1H-PYRAZOL-1-YL)PYRIDINE

(57) **Abrégé/Abstract:**

3-(3-chloro-1H-pyrazol-1-yl)pyridine is prepared by cyclizing 3-hydrazinopyridine•dihydrochloride with a dialkyl maleate to provide an alkyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate, by chlorinating to provide an alkyl 3-chloro-1-(pyridin-3-yl)-4,5-dihydro-1H-pyrazole-5-carboxylate, by oxidizing to provide an alkyl 3-chloro-1-(pyridin-3-yl)-1H-pyrazole-5-carboxylate, by converting the ester to the carboxylic acid by hydrolysis to provide 3-chloro-1-(pyridin-3-yl)-1H-pyrazole-5-carboxylic acid hydrochloride, and by removing the carboxylic acid by a decarboxylation reaction.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



WIPO | PCT



(10) International Publication Number

WO 2016/028328 A1

(43) International Publication Date
25 February 2016 (25.02.2016)

(51) International Patent Classification:

C07D 231/14 (2006.01) A61K 31/415 (2006.01)
A01N 43/46 (2006.01)

(21) International Application Number:

PCT/US2014/061005

(22) International Filing Date:

17 October 2014 (17.10.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/039,128 19 August 2014 (19.08.2014) US

(71) Applicant: DOW AGROSCIENCES LLC [US/US];
9330 Zionsville Road, Indianapolis, Indiana 46268 (US).

(72) Inventors: YANG, Qiang; 11437 Mears Drive, Zionsville,
Indiana 46077 (US). LORSBACH, Beth; 6034 Haverford
Avenue, Indianapolis, Indiana 46220 (US). PODHOREZ,
David E.; 5604 Berry Court, Midland, Michigan 48640
(US).

(74) Agents: ADDISON, Bradford G. et al.; Barnes & Thorn-
burg LLP, 11 South Meridian Street, Indianapolis, Indiana
46204 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))

(54) Title: PROCESS FOR THE PREPARATION OF 3-(3-CHLORO-1H-PYRAZOL-1-YL)PYRIDINE

(57) Abstract: 3-(3-chloro-1H-pyrazol-1-yl)pyridine is prepared by cyclizing 3-hydrazinopyridine•dihydrochloride with a dialkyl maleate to provide an alkyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate, by chlorinating to provide an alkyl 3-chloro-1-(pyridin-3-yl)-4,5-dihydro-1H-pyrazole-5-carboxylate, by oxidizing to provide an alkyl 3-chloro-1-(pyridin-3-yl)-1H-pyrazole-5-carboxylic acid hydrochloride, and by removing the carboxylic acid by a decarboxylation reaction.



WO 2016/028328 A1

PROCESS FOR THE PREPARATION OF 3-(3-CHLORO-1*H*-PYRAZOL-1-YL)PYRIDINE

CROSS REFERENCE TO RELATED APPLICATIONS

This Application claims the benefit of U.S. Provisional Application Serial No. 62/039,128, filed August 19, 2014, the entire disclosure of which is hereby expressly incorporated by reference in this Application.

BACKGROUND


The present invention concerns an improved process for preparing 3-(3-chloro-1*H*-pyrazol-1-yl)pyridine.

US 20130288893(A1) describes, *inter alia*, certain (3-halo-1-(pyridin-3-yl)-1*H*-pyrazol-4-yl)amides and carbamates and their use as pesticides. The route to prepare such compounds involved the preparation of 3-(3-chloro-1*H*-pyrazol-1-yl)pyridine by the direct coupling of 3-bromopyridine with 3-chloropyrazole. The 3-chloropyrazole was prepared by a) treating 1*H*-pyrazole with 2 dimethylsulfamoyl chloride and sodium hydride to provide *N,N*-dimethyl-1*H*-pyrazole-1-sulfonamide, b) treating the *N,N*-dimethyl-1*H*-pyrazole-1-sulfonamide with perchloroethane and *n*-butyl lithium to provide 3-chloro-*N,N*-dimethyl-1*H*-pyrazole-1-sulfonamide, and c) removing the *N,N*-dimethylsulfonamide from 3-chloro-*N,N*-dimethyl-1*H*-pyrazole-1-sulfonamide with trifluoroacetic acid to give the 3-chloropyrazole.

The disclosed process produces low yields, relies on a starting material that is difficult to prepare (3-chloropyrazole) and provides a product that is difficult to isolate in a pure form. It would be desirable to have a process for preparing 3-(3-chloro-1*H*-pyrazol-1-yl)pyridine that avoids these problems.


SUMMARY

The present invention provides such an alternative by cyclizing 3-hydrazinopyridine-dihydrochloride with a dialkyl maleate to provide alkyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate (10a), by chlorinating to provide alkyl 3-chloro-1-(pyridin-3-yl)-4,5-dihydro-1*H*-pyrazole-5-carboxylate (10b), by oxidizing to provide alkyl 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylate (10c), by converting the ester to the carboxylic acid by hydrolysis to provide 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride (10d), and by removing the carboxylic acid by a decarboxylation reaction. Thus, the present invention concerns a process for preparing 3-(3-chloro-1*H*-pyrazol-1-yl)pyridine (5b),




which comprises

- a) treating 3-hydrazinopyridine·dihydrochloride




with a dialkyl maleate



wherein


R represents (C₁-C₄) alkyl,

in a (C₁-C₄) aliphatic alcohol at a temperature of about 25 °C to about 100 °C in the presence of an alkali metal (C₁-C₄) alkoxide to provide an alkyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate (10a)




wherein R is as previously defined;

b) treating the alkyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate (10a) with a chlorinating reagent in an inert organic solvent at a temperature of about 25 °C to about 100 °C to provide an alkyl 3-chloro-1-(pyridin-3-yl)-4,5-dihydro-1H-pyrazole-5-carboxylate (10b)




wherein R is as previously defined;

c) treating the alkyl 3-chloro-1-(pyridin-3-yl)-4,5-dihydro-1*H*-pyrazole-5-carboxylate (10b) with an oxidant in an inert organic solvent at a temperature of about 25 °C to about 100 °C to provide an alkyl 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylate (10c)



wherein R is as previously defined;

d) treating the alkyl 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylate (10c) with aqueous hydrochloric acid at a temperature of about 25 °C to about 100 °C to provide 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride (10d)



e) treating 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride with copper(II) oxide in a polar aprotic solvent at a temperature of about 80 °C to about 140 °C.

DETAILED DESCRIPTION

The present invention provides an improved process for preparing 3-(3-chloro-1*H*-pyrazol-1-yl)pyridine (5b), by cyclizing 3-hydrazinopyridine·dihydrochloride with a dialkyl maleate to provide an alkyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate (10a), by chlorinating to provide an alkyl 3-chloro-1-(pyridin-3-yl)-4,5-dihydro-1*H*-pyrazole-5-carboxylate (10b), by oxidizing to provide an alkyl 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylate (10c), by converting the ester to the carboxylic acid by hydrolysis to provide 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride (10d), and by removing the carboxylic acid by a decarboxylation reaction.

In the first step, 3-hydrazinopyridine·dihydrochloride is treated with a dialkyl maleate in a (C₁-C₄) aliphatic alcohol at a temperature of about 25 °C to about 100 °C in the presence of an alkali metal (C₁-C₄) alkoxide to provide an alkyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate (10a). While stoichiometric amounts of 3-hydrazinopyridine·dihydrochloride and dialkyl maleate are required, it is often convenient to use about a 1.5 fold to about a 2 fold excess of dialkyl maleate. The cyclization is run in the presence of an alkali metal (C₁-C₄)

alkoxide base. It is often convenient to use about a 2 fold to about a 5 fold excess of base. The cyclization is performed in a (C₁-C₄) aliphatic alcohol. It is most convenient that the alkoxide base, the alcohol solvent and the ester of the maleate be the same, for example, sodium ethoxide in ethanol with diethyl maleate.

In a preferred reaction, sodium ethoxide in an anhydrous ethanol are introduced into a reaction vessel and 3-hydrazinopyridine·dihydrochloride is added. The mixture is stirred and diethyl maleate is added. The mixture is heated at about 60 °C until most of the 3-hydrazinopyridine has reacted. The mixture is allowed to cool and the excess base is neutralized with acid. The crude ethyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate (10a) is conveniently isolated and purified by standard techniques.

In the second step, the alkyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate (10a) is treated with a chlorinating reagent in an inert organic solvent at a temperature of about 25 °C to about 100 °C to provide alkyl 3-chloro-1-(pyridin-3-yl)-4,5-dihydro-1*H*-pyrazole-5-carboxylate (10b). Suitable chlorinating reagents include phosphoryl chloride (phosphorous oxychloride) and phosphorus pentachloride. Phosphoryl chloride is preferred. It is often convenient to use about a 1.1 fold to about a 10 fold excess of the chlorinating reagent. The chlorination is performed in an organic solvent that is inert to the chlorinating reagent. Suitable solvents include nitriles such as acetonitrile. With phosphoryl chloride as the chlorinating reagent, acetonitrile is a preferred solvent.

In a preferred reaction, ethyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate (10a) and acetonitrile are mixed with phosphoryl chloride and the mixture is heated to about 60 °C for 2 hours. After the reaction is determined to be complete, the reaction is cooled to room temperature and diluted with water. The reaction mixture is then neutralized with base and extracted. The alkyl 3-chloro-1-(pyridin-3-yl)-4,5-dihydro-1*H*-pyrazole-5-carboxylate (10b) can be purified by standard techniques.

In the third step, alkyl 3-chloro-1-(pyridin-3-yl)-4,5-dihydro-1*H*-pyrazole-5-carboxylate (10b) is treated with an oxidant in an organic solvent at a temperature of about 25 °C to about 100 °C to provide alkyl 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylate (10c). Suitable oxidants include manganese (IV) oxide and sodium persulfate/sulfuric acid. It is often convenient to use about a 1.5 fold to about a 15 fold excess of oxidant. The oxidation is performed in an organic solvent that is inert to the oxidant. Suitable solvents include nitriles such as acetonitrile. With manganese (IV) oxide or sodium persulfate/sulfuric acid as the oxidant, acetonitrile is a preferred solvent.

In a preferred reaction, ethyl 3-chloro-1-(pyridin-3-yl)-4,5-dihydro-1*H*-pyrazole-5-carboxylate (10b) and acetonitrile are mixed with manganese(IV) oxide and the mixture is

heated at about 60 °C until the reaction is completed. The ethyl 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylate (10c) is conveniently isolated and purified by standard techniques.

In the fourth step, alkyl 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylate (10c) is then converted to the desired 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride (10d) by treatment in aqueous hydrochloric acid at a temperature of about 25 °C to about 100 °C. While stoichiometric amounts of reagents are required, it is often convenient to use an excess of reagent with respect to the alkyl 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylate (10c). Thus, aqueous hydrochloric acid is used in large excess as the reaction medium.

In a preferred reaction, a mixture of ethyl 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylate (10c) and aqueous hydrochloric acid are mixed and heated to about 90 °C. After completion of the reaction, the mixture is cooled and diluted with an organic solvent. The resulting solution is concentrated. The 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride (10d) can be purified by standard techniques such as filtration.


3-Chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride (10d) is then converted to the desired 3-(3-chloro-1*H*-pyrazol-1-yl)pyridine (5b) by decarboxylation in the presence of copper (II) oxide in polar solvents at a temperature from about 80 °C to about 140 °C. It was surprisingly discovered that this decarboxylation only occurs in the presence of copper (II) oxide. Several known decarboxylation agents from the literature such as, for example, hydrochloric acid (See Example 4), sulfuric acid (See “CE-5”), and palladium (II) trifluoroacetate/trifluoroacetic acid (See “CE-5”) did not yield the desired product.

In a preferred reaction, a mixture of 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride (10d) and copper(II) oxide are mixed in an organic solvent and heated to about 120 °C. It is often convenient to use less than stoichiometric amounts of copper(II) oxide. The decarboxylation is performed in a polar aprotic organic solvent. Suitable solvents include *N,N'*-dimethylformamide. After completion of the reaction, the mixture is cooled, diluted with ammonium hydroxide, and extracted. The 3-(3-chloro-1*H*-pyrazol-1-yl)pyridine (5b) can be purified by standard techniques.

The following examples are presented to illustrate the invention.


Examples

1. Preparation of ethyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate (10a)



A 4-neck round bottomed flask (250 mL) was charged with sodium ethoxide (21 wt% in ethanol, 56 mL, 192 mmol). 3-Hydrazinopyridine·dihydrochloride (10.0 g, 55.0 mmol) was added, causing an exotherm from 20 °C to 32 °C. The reaction was allowed to cool to 20 °C and diethyl maleate (13.4 mL, 82.0 mmol) was added, and the reaction was heated at 60 °C for 3 hours (h). The reaction was cooled to 20 °C and quenched with acetic acid. The reaction mixture was diluted with water (100 mL) and extracted with ethyl acetate (3 × 100 mL). The combined organics were concentrated to dryness and the residue was purified by flash column chromatography using ethyl acetate as eluent to the title compound as a blue oil (6.60 g, 51%): ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.40 (s, 1H), 8.40 – 8.26 (m, 1H), 8.19 (dd, *J* = 4.4, 1.6 Hz, 1H), 7.47 – 7.21 (m, 2H), 4.77 (dd, *J* = 9.8, 2.1 Hz, 1H), 4.22 (qd, *J* = 7.1, 1.7 Hz, 2H), 3.05 (dd, *J* = 17.0, 9.8 Hz, 1H), 1.99 (s, 1H), 1.25 (t, *J* = 7.1 Hz, 3H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ 170.37, 146.60, 142.60, 137.28, 123.54, 121.94, 65.49, 61.32, 32.15, 20.72, 13.94; ESIMS *m/z* 236 ([M+H]⁺).


2. Preparation of ethyl 3-chloro-1-(pyridin-3-yl)-4,5-dihydro-1*H*-pyrazole-5-carboxylate (10b)



A 3-neck round bottomed flask (100 mL) was charged with ethyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate (8.50 g, 36.1 mmol) and acetonitrile (40 mL). Phosphoryl chloride (4.05 mL, 43.4 mmol) was charged and the reaction was heated at 60 °C for 2 h. The reaction was cooled to 20 °C and water (100 mL) was added. Sodium carbonate was added to adjust pH to 8 and the mixture was extracted with ethyl acetate (3 × 100 mL). The organic layers were concentrated to dryness and the residue was purified by flash column chromatography using 30–80% ethyl acetate/hexanes as eluent to provide the title compound as a yellow oil (7.30 g, 79%): ¹H NMR (400 MHz, CDCl₃) δ 8.30 (dd, *J* = 2.9, 0.8 Hz, 1H), 8.17

(dd, $J = 4.7, 1.4$ Hz, 1H), 7.38 (ddd, $J = 8.4, 2.8, 1.4$ Hz, 1H), 7.18 (ddd, $J = 8.4, 4.7, 0.7$ Hz, 1H), 4.79 (dd, $J = 12.4, 6.9$ Hz, 1H), 4.24 (qd, $J = 7.1, 1.1$ Hz, 2H), 3.55 (dd, $J = 17.7, 12.4$ Hz, 1H), 3.33 (dd, $J = 17.8, 6.9$ Hz, 1H), 1.25 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ 169.65, 141.90, 141.33, 141.09, 135.13, 123.53, 120.37, 62.89, 62.35, 42.45, 14.03; ESIMS m/z 254 ($[\text{M}+\text{H}]^+$).

3. Preparation of ethyl 3-chloro-1-(pyridin-3-yl)-1H-pyrazole-5-carboxylate (10c)




A 3-neck round bottomed flask (100 mL) was charged with ethyl 3-chloro-1-(pyridin-3-yl)-1H-dihydropyrazole-5-carboxylate (2.00 g, 7.88 mmol) and acetonitrile (20 mL). Manganese(IV) oxide (3.43 g, 39.4 mmol) was added. The reaction was stirred at 60 °C for 18 h. Additional manganese(IV) oxide (3.43 g, 39.4 mmol) was added and the reaction was stirred at 80 °C for 6 h. The mixture was filtered through a Celite[®] pad and the pad was rinsed with ethyl acetate (20 mL). The combined filtrates were concentrated to dryness and the residue was purified by flash column chromatography using 10–60% ethyl acetate/hexanes. The pure fractions were concentrated to dryness to afford a white solid after drying (1.84 g, 93%): ^1H NMR (400 MHz, CDCl_3) δ 8.75 – 8.64 (m, 2H), 7.79 (ddd, $J = 8.2, 2.6, 1.5$ Hz, 1H), 7.42 (ddd, $J = 8.2, 4.8, 0.8$ Hz, 1H), 6.98 (s, 1H), 4.27 (q, $J = 7.1$ Hz, 2H), 1.27 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ 157.90, 149.88, 147.01, 141.41, 136.24, 135.27, 133.34, 123.11, 111.97, 61.87, 13.98; ESIMS m/z 252 ($[\text{M}+\text{H}]^+$).

Alternate synthetic route to ethyl 3-chloro-1-(pyridin-3-yl)-1H-pyrazole-5-carboxylate (10c)


A vial (20 mL) was charged with ethyl 3-chloro-1-(pyridin-3-yl)-1H-dihydropyrazole-5-carboxylate (0.500 g, 1.97 mmol) and acetonitrile (5 mL). Sodium persulfate (0.799 g, 2.96 mmol) was added, followed by sulfuric acid (0.733 g, 7.88 mmol) (Exotherm!). The reaction was heated at 60 °C for 18 hours. The reaction was cooled to 20 °C and poured into water (20 mL). The mixture was treated with sodium carbonate to achieve pH 9 and extracted with ethyl acetate (2 x 20 mL). The organic layers were concentrated to a residue, which was purified by flash column chromatography using 50% ethyl acetate/hexanes as eluent to provide the title compound as a white solid (0.280 g, 56%).

4. Preparation of 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride (10d)



A 3-neck round bottomed flask (100 mL) was charged with ethyl 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylate (0.200 g, 0.795 mmol) and hydrochloric acid (37%, 4 mL). The reaction was heated at 90 °C for 18 hours and allowed to cool to 20 °C. Dioxane (5 mL) was added to the resulting suspension and was concentrated to dryness. Dioxane (5 mL) was added and the suspension was concentrated again to afford a white solid. Dioxane (5 mL) was added and the resulting suspension was stirred for 1 hour at 20 °C. The solid was filtered and the solid was rinsed with dioxane (2 mL). The filter cake was dried under vacuum at 20 °C to afford the title compound as a white solid (0.218 g, 100%): ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.05 (dd, *J* = 2.5, 0.7 Hz, 1H), 8.84 (dd, *J* = 5.3, 1.4 Hz, 1H), 8.41 (ddd, *J* = 8.3, 2.5, 1.4 Hz, 1H), 7.88 (ddd, *J* = 8.3, 5.2, 0.7 Hz, 1H), 7.26 (s, 1H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ 158.71, 146.00, 143.44, 140.36, 137.76, 137.00, 136.83, 125.19, 111.71.

4. Preparation of 3-(3-chloro-1*H*-pyrazol-1-yl)pyridine (5b)




3-Chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride (1.00 g, 3.65 mmol) was stirred in *N,N'*-dimethylformamide (10 mL). Copper (II) oxide (0.0580 mg, 0.730 mmol) was added and the reaction was heated at 120 °C for 16 hours, at which point the reaction was ~20% complete. Additional copper (II) oxide (0.112 g, 1.46 mmol) was added and the reaction was stirred for 5 hours. The mixture was diluted with ammonium hydroxide and water and extracted with ethyl acetate. The organic layer was washed with lithium chloride (15%) and concentrated to provide an orange solid. The residue was purified by flash column

chromatography using ethyl acetate as eluent and the pure fractions were concentrated to afford the desired product as a white solid (0.481 g, 69.7%): mp: 66–68 °C; ^1H NMR (400 MHz, CDCl_3) δ 8.93 (d, $J = 27$ Hz, 1H), 8.57 (dd, $J = 4.8, 1.4$ Hz, 1H), 8.02 (ddd, $J = 8.3, 2.7, 1.5$ Hz, 1H), 7.91 (d, $J = 2.6$ Hz, 1H), 7.47–7.34 (m, 1H), 6.45 (d, $J = 2.6$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ 148.01, 142.72, 140.12, 135.99, 128.64, 126.41, 124.01, 108.0.

COMPARATIVE EXAMPLES

Example CE-5 3-chloro-*N*-ethyl-1-(pyridin-3-yl)-1*H*-pyrazol-amine (5b):



Attempted decarboxylation with sulfuric acid: 3-Chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride (1.00 g, 2.50 mmol) was dissolved in warm sulfolane (12.5 mL). Sulfuric acid (1.35 mL, 25.0 mmol) was added and the reaction mixture was heated to 100 °C. After stirring for 1 hour, LCMS indicated that the reaction did not occur. The reaction was further heated at 130 °C for 2 hours, at which point LCMS indicated no change. Additional sulfuric acid (4 mL) was added and the reaction was heated at 150 °C for 2 hours, at which point LCMS showed a new major peak that did not correspond to desired product.

Attempted decarboxylation with palladium(II) trifluoroacetate/trifluoroacetic acid: 3-Chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride (1.00g, 2.50 mmol) was dissolved in a mixture of dimethylsulfoxide (0.625 mL) and *N,N'*-dimethylformamide (11.9 ml). Trifluoroacetic acid (1.93 ml, 25.0 mmol) was added followed by the addition of palladium(II) trifluoroacetate/trifluoroacetic acid (0.332 g, 1.00 mmol). The reaction was heated at 100 °C overnight, at which time LCMS indicated that a reaction had occurred but no desired product had been formed.

BIOLOGICAL EXAMPLES

Example A Bioassays on Green Peach Aphid (“GPA”) (*Myzus persicae*) (MYZUPE.)

GPA is the most significant aphid pest of peach trees, causing decreased growth, shriveling of leaves, and the death of various tissues. It is also hazardous because it acts as a vector for the transport of plant viruses, such as potato virus Y and potato leafroll virus to

members of the nightshade /potato family *Solanaceae*, and various mosaic viruses to many other food crops. GPA attacks such plants as broccoli, burdock, cabbage, carrot, cauliflower, daikon, eggplant, green beans, lettuce, macadamia, papaya, peppers, sweet potatoes, tomatoes, watercress and zucchini among other plants. GPA also attacks many ornamental crops such as carnations, chrysanthemum, flowering white cabbage, poinsettia and roses. GPA has developed resistance to many pesticides.

Several molecules disclosed herein were tested against GPA using procedures described below.

Cabbage seedling grown in 3-in pots, with 2-3 small (3-5 cm) true leaves, were used as test substrate. The seedlings were infested with 20-5- GPA (wingless adult and nymph stages) one day prior to chemical application. Four pots with individual seedlings were used for each treatment. Test compounds (2 mg) were dissolved in 2 mL of acetone/MeOH (1:1) solvent, forming stock solutions of 1000 ppm test compound. The stock solutions were diluted 5X with 0.025% Tween 20 in water to obtain the solution at 200 ppm test compound. A hand-held aspirator-type sprayer was used for spraying a solution to both sides of the cabbage leaves until runoff. Reference plants (solvent check) were sprayed with the diluent only containing 20% by volume acetone/MeOH (1:1) solvent. Treated plants were held in a holding room for three days at approximately 25 °C and ambient relative humidity (RH) prior to grading. Evaluation was conducted by counting the number of live aphids per plant under a microscope. Percent Control was measured by using Abbott's correction formula (W.S. Abbott, "A Method of Computing the Effectiveness of an Insecticide" J. Econ. Entomol 18 (1925), pp.265-267) as follows.

$$\text{Corrected \% Control} = 100 * (X - Y) / X$$

where

X = No. of live aphids on solvent check plants and

Y = No. of live aphids on treated plants

The results are indicated in the table entitled "Table 1: GPA (MYZUPE) and sweetpotato whitefly-crawler (BEMITA) Rating Table".

Example B Bioassays on Sweetpotato Whitefly Crawler (*Bemisia tabaci*) (BEMITA.)

The sweetpotato whitefly, *Bemisia tabaci* (Gennadius), has been recorded in the United States since the late 1800s. In 1986 in Florida, *Bemisia tabaci* became an extreme economic pest. Whiteflies usually feed on the lower surface of their host plant leaves. From the egg hatches a minute crawler stage that moves about the leaf until it inserts its microscopic, threadlike mouthparts to feed by sucking sap from the phloem. Adults and nymphs excrete honeydew (largely plant sugars from feeding on phloem), a sticky, viscous liquid in which dark

sooty molds grow. Heavy infestations of adults and their progeny can cause seedling death, or reduction in vigor and yield of older plants, due simply to sap removal. The honeydew can stick cotton lint together, making it more difficult to gin and therefore reducing its value. Sooty mold grows on honeydew-covered substrates, obscuring the leaf and reducing photosynthesis, and reducing fruit quality grade. It transmitted plant-pathogenic viruses that had never affected cultivated crops and induced plant physiological disorders, such as tomato irregular ripening and squash silverleaf disorder. Whiteflies are resistant to many formerly effective insecticides.

Cotton plants grown in 3-inch pots, with 1 small (3-5 cm) true leaf, were used at test substrate. The plants were placed in a room with whitely adults. Adults were allowed to deposit eggs for 2-3 days. After a 2-3 day egg-laying period, plants were taken from the adult whitefly room. Adults were blown off leaves using a hand-held Devilbliss sprayer (23 psi). Plants with egg infestation (100-300 eggs per plant) were placed in a holding room for 5-6 days at 82°F and 50% RH for egg hatch and crawler stage to develop. Four cotton plants were used for each treatment. Compounds (2 mg) were dissolved in 1 mL of acetone solvent, forming stock solutions of 2000 ppm. The stock solutions were diluted 10X with 0.025% Tween 20 in water to obtain a test solution at 200 ppm. A hand-held Devilbliss sprayer was used for spraying a solution to both sides of cotton leaf until runoff. Reference plants (solvent check) were sprayed with the diluent only. Treated plants were held in a holding room for 8-9 days at approximately 82°F and 50% RH prior to grading. Evaluation was conducted by counting the number of live nymphs per plant under a microscope. Pesticidal activity was measured by using Abbott's correction formula (see above) and presented in Table 1.

Table 1: GPA (MYZUPE) and sweetpotato whitefly-crawler (BEMITA) Rating Table


% Mortality	Rating
80-100	A
More than 0 – Less than 80	B
Not Tested	C
No activity observed in this bioassay	D

Table 2

Example Compound	BEMITA	MYZUPE
10a	B	B
10b	B	B


WHAT IS CLAIMED IS:

1. A process for preparing 3-(3-chloro-1*H*-pyrazol-1-yl)pyridine (5b),




which comprises

- a) treating 3-hydrazinopyridine·dihydrochloride




with a dialkyl maleate



wherein


R represents (C₁-C₄) alkyl,

in a (C₁-C₄) aliphatic alcohol at a temperature of about 25 °C to about 100 °C in the presence of an alkali metal (C₁-C₄) alkoxide to provide an alkyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate (10a)




wherein R is as previously defined;

- b) treating the alkyl 5-oxo-2-(pyridin-3-yl)pyrazolidine-3-carboxylate (10a) with a chlorinating reagent in an inert organic solvent at a temperature of about 25 °C to about 100 °C to provide an alkyl 3-chloro-1-(pyridin-3-yl)-4,5-dihydro-1*H*-pyrazole-5-carboxylate (10b)




wherein R is as previously defined;

c) treating the alkyl 3-chloro-1-(pyridin-3-yl)-4,5-dihydro-1*H*-pyrazole-5-carboxylate (10b) with an oxidant in an inert organic solvent at a temperature of about 25 °C to about 100 °C to provide an alkyl 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylate (10c)



wherein R is as previously defined;

d) treating the alkyl 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylate (10c) with aqueous hydrochloric acid at a temperature of about 25 °C to about 100 °C to provide 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride (10d)



e) treating 3-chloro-1-(pyridin-3-yl)-1*H*-pyrazole-5-carboxylic acid hydrochloride with copper(II) oxide in a polar aprotic solvent at a temperature of about 80 °C to about 140 °C.


2. The process of Claim 1 in which for step a) the dialkyl maleate is diethyl maleate, the (C₁-C₄) aliphatic alcohol is ethanol and the alkali metal (C₁-C₄) alkoxide is sodium ethoxide.

3. The process of Claim 1 in which for step b) the chlorinating reagent is phosphoryl chloride and the inert organic solvent is acetonitrile.

4. The process of Claim 1 in which for step c) the oxidant is manganese (IV) oxide and the inert organic solvent is acetonitrile.

5. The process of Claim 1 in which for step e) the polar aprotic solvent is *N, N'*-dimethylformamide.


6. A molecule having the following structure,



wherein

R represents (C₁-C₄) alkyl.

7. A molecule having the following structure,



wherein

R represents (C₁-C₄) alkyl.