




(86) Date de dépôt PCT/PCT Filing Date: 2016/02/16
(87) Date publication PCT/PCT Publication Date: 2016/08/18
(85) Entrée phase nationale/National Entry: 2017/08/10
(86) N° demande PCT/PCT Application No.: US 2016/018065
(87) N° publication PCT/PCT Publication No.: 2016/131052
(30) Priorités/Priorities: 2015/02/13 (US62/116,232);
2015/07/22 (US62/195,462); 2015/10/16 (US62/242,383);
2015/10/23 (US62/245,726); 2016/02/01 (US62/289,617)

(51) Cl.Int./Int.Cl. *C12N 15/11* (2006.01),
A61K 31/7115 (2006.01), *A61K 38/00* (2006.01),
A61K 38/17 (2006.01), *A61K 38/18* (2006.01),
A61K 38/20 (2006.01), *A61K 38/46* (2006.01),
A61M 37/00 (2006.01), *C12N 15/87* (2006.01)
(71) Demandeur/Applicant:
FACTOR BIOSCIENCE INC., US
(72) Inventeurs/Inventors:
ANGEL, MATTHEW, US;
ROHDE, CHRISTOPHER, US
(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : PRODUITS D'ACIDES NUCLEIQUES ET LEURS PROCEDES D'ADMINISTRATION
(54) Title: NUCLEIC ACID PRODUCTS AND METHODS OF ADMINISTRATION THEREOF

FIG. 39



(57) **Abrégé/Abstract:**

The present invention relates in part to nucleic acids, including nucleic acids encoding proteins, therapeutics and cosmetics comprising nucleic acids, methods for delivering nucleic acids to cells, tissues, organs, and patients, methods for inducing cells to express proteins using nucleic acids, methods, kits and devices for transfecting, gene editing, and reprogramming cells, and cells, organisms, therapeutics, and cosmetics produced using these methods, kits, and devices.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
18 August 2016 (18.08.2016)

WIPO | PCT

(10) International Publication Number
WO 2016/131052 A1

(51) International Patent Classification:

C12N 15/11 (2006.01) *A61K 38/46* (2006.01)
C12N 15/87 (2006.01) *A61K 38/00* (2006.01)
A61K 38/17 (2006.01) *A61K 31/7115* (2006.01)
A61K 38/18 (2006.01) *A61M 37/00* (2006.01)
A61K 38/20 (2006.01)

(21) International Application Number:

PCT/US2016/018065

(22) International Filing Date:

16 February 2016 (16.02.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/116,232	13 February 2015 (13.02.2015)	US
62/195,462	22 July 2015 (22.07.2015)	US
62/242,383	16 October 2015 (16.10.2015)	US
62/245,726	23 October 2015 (23.10.2015)	US
62/289,617	1 February 2016 (01.02.2016)	US

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(71) Applicant: **FACTOR BIOSCIENCE INC.** [US/US]; 8 Museum Way, #2204, Cambridge, MA 02141 (US).

(72) Inventors: **ANGEL, Matthew**; 8 Museum Way, #2204, Cambridge, MA 02141 (US). **ROHDE, Christopher**; 8 Museum Way, #2204, Cambridge, MA 02141 (US).


(74) Agents: **ALTIERI, Stephen L.** et al.; Morgan, Lewis & Bockius LLP, 1111 Pennsylvania Avenue, NW, Washington, DC 20004 (US).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: NUCLEIC ACID PRODUCTS AND METHODS OF ADMINISTRATION THEREOF

FIG. 39



(57) Abstract: The present invention relates in part to nucleic acids, including nucleic acids encoding proteins, therapeutics and cosmetics comprising nucleic acids, methods for delivering nucleic acids to cells, tissues, organs, and patients, methods for inducing cells to express proteins using nucleic acids, methods, kits and devices for transfecting, gene editing, and reprogramming cells, and cells, organisms, therapeutics, and cosmetics produced using these methods, kits, and devices.

NUCLEIC ACID PRODUCTS AND METHODS OF ADMINISTRATION THEREOF**PRIORITY**

This application claims the benefit of priority of US Patent Application Nos. 62/116,232, filed February 13, 2015, 62/195,462, filed July 22, 2015, 62/242,383, filed October 16, 2015, 62/245,726, filed October 23, 2015; and 62/289,617, filed February 1, 2016, the entire contents of all of which are incorporated by reference herein.

FIELD OF THE INVENTION

The present invention relates, in part, to methods, compositions, and products for producing and delivering nucleic acids to cells, tissues, organs, and patients, methods for expressing proteins in cells, tissues, organs, and patients, and cells, therapeutics, and cosmetics produced using these methods, compositions, and products.

DESCRIPTION OF THE TEXT FILE SUBMITTED ELECTRONICALLY

The contents of the text file submitted electronically herewith are incorporated herein by reference in their entirety: A computer readable format copy of the Sequence Listing (filename: FAB-009PC_Sequence.txt; date recorded: February 16, 2016; file size: 2.3 MB).

BACKGROUND*Synthetic RNA and Nucleic-Acid Therapeutics*

Ribonucleic acid (RNA) is ubiquitous in both prokaryotic and eukaryotic cells, where it encodes genetic information in the form of messenger RNA, binds and transports amino acids in the form of transfer RNA, assembles amino acids into proteins in the form of ribosomal RNA, and performs numerous other functions including gene expression regulation in the forms of microRNA and long non-coding RNA. RNA can be produced synthetically by methods including direct chemical synthesis and *in vitro* transcription, and can be administered to patients for therapeutic use. However, previously described synthetic RNA molecules are unstable and trigger a potent innate-immune response in human cells. In addition, methods for efficient non-viral delivery of nucleic acids to patients, organs, tissues, and cells *in vivo* have not been previously described. The many drawbacks of existing synthetic RNA technologies and methods for delivery of nucleic acids make them undesirable for therapeutic or cosmetic use.

Cell Reprogramming and Cell-Based Therapies

Cells can be reprogrammed by exposing them to specific extracellular cues and/or by ectopic expression of specific proteins, microRNAs, etc. While several reprogramming methods have been previously described, most that rely on ectopic expression require the introduction of exogenous DNA, which can carry mutation risks. DNA-free reprogramming methods based on direct delivery of reprogramming proteins have been reported. However, these methods are too inefficient and unreliable for commercial use. In addition, RNA-based reprogramming

methods have been described (see, e.g., Angel. MIT Thesis. 2008. 1-56; Angel *et al.* PLoS ONE. 2010. 5,107; Warren *et al.* Cell Stem Cell. 2010. 7,618-630; Angel. MIT Thesis. 2011. 1-89; and Lee *et al.* Cell. 2012. 151,547-558; the contents of all of which are hereby incorporated by reference). However, existing RNA-based reprogramming methods are slow, unreliable, and inefficient when performed on adult cells, require many transfections (resulting in significant expense and opportunity for error), can reprogram only a limited number of cell types, can reprogram cells to only a limited number of cell types, require the use of immunosuppressants, and require the use of multiple human-derived components, including blood-derived HSA and human fibroblast feeders. The many drawbacks of previously disclosed RNA-based reprogramming methods make them undesirable for research, therapeutic or cosmetic use.

10 *Gene Editing*

Several naturally occurring proteins contain DNA-binding domains that can recognize specific DNA sequences, for example, zinc fingers (ZFs) and transcription activator-like effectors (TALEs). Fusion proteins containing one or more of these DNA-binding domains and the cleavage domain of FokI endonuclease can be used to create a double-strand break in a desired region of DNA in a cell (see, e.g., US Patent Appl. Pub. No. US 2012/0064620, US Patent Appl. Pub. No. US 2011/0239315, US Patent No. 8,470,973, US Patent Appl. Pub. No. US 2013/0217119, US Patent No. 8,420,782, US Patent Appl. Pub. No. US 2011/0301073, US Patent Appl. Pub. No. US 2011/0145940, US Patent No. 8,450,471, US Patent No. 8,440,431, US Patent No. 8,440,432, and US Patent Appl. Pub. No. 2013/0122581, the contents of all of which are hereby incorporated by reference). Other gene-editing proteins include clustered regularly interspaced short palindromic repeat (CRISPR)-associated proteins. However, current methods for gene editing cells are inefficient and carry a risk of uncontrolled mutagenesis, making them undesirable for research, therapeutic or cosmetic use. Methods for DNA-free gene editing of somatic cells have not been previously explored, nor have methods for simultaneous or sequential gene editing and reprogramming of somatic cells. In addition, methods for directly gene editing cells in patients (*i.e., in vivo*) have not been previously explored, and the development of such methods has been limited by a lack of acceptable targets, inefficient delivery, inefficient expression of the gene-editing protein/proteins, inefficient gene editing by the expressed gene-editing protein/proteins, due in part to poor binding of DNA-binding domains, excessive off-target effects, due in part to non-directed dimerization of the FokI cleavage domain and poor specificity of DNA-binding domains, and other factors. Finally, the use of gene editing in anti-bacterial, anti-viral, and anti-cancer treatments has not been previously explored.

Accordingly, there remains a need for improved methods and compositions for the production and delivery of nucleic acids to cells, tissues, organs, and patients.

SUMMARY OF THE INVENTION

The present invention provides, in part, compositions, methods, articles, and devices for delivering nucleic acids to cells, tissues, organs, and patients, methods for inducing cells to express proteins, methods, articles, and

devices for producing these compositions, methods, articles, and devices, and compositions and articles, including cells, organisms, cosmetics and therapeutics, produced using these compositions, methods, articles, and devices. Unlike previously reported methods, certain embodiments of the present invention provide small doses of nucleic acids to achieve significant and lasting protein expression in humans.

- 5 In various aspects, the present invention is based on the surprising discovery of safe and effective doses and administration parameters for nucleic acid drugs in human subjects. In some aspects, there is provided a method for delivering a nucleic acid drug, comprising administering an effective dose of a nucleic acid drug to a human subject in need thereof. In various embodiments, the nucleic acid drug comprises a RNA comprising one or more non-canonical nucleotides (a/k/a "modified RNA"). In other embodiments, the effective dose is an amount
- 10 sufficient to substantially increase an amount of a protein encoded by the nucleic acid drug in the human subject and/or substantially avoid an immune reaction in a human subject, wherein the immune reaction is optionally mediated by the innate immune system.

- In some aspects, there is provided a method for expressing a protein of interest in a population of cells in a mammalian subject, comprising administering a non-viral transfection composition comprising an effective dose
- 15 of a RNA encoding the protein of interest to said cells, where the transfection composition is administered in an amount that allows for expression of said protein in said cells for at least about six hours to about five days without substantial cellular toxicity. In some embodiments, the RNA contains one or more non-canonical nucleotides that avoid substantial cellular toxicity.

- In some embodiments, the effective dose is about 100 ng to about 2000 ng (e.g., about, or no more than about,
- 20 100 ng, or 200 ng, or 300 ng, or 400 ng, or 500 ng, or 600 ng, or 700 ng, or 800 ng, or 900 ng, or 1000 ng, or 1100 ng, or 1200 ng, or 1300 ng, or 1400 ng, or 1500 ng, or 1600 ng, or 1700 ng, or 1800 ng, or 1900 ng, or 2000 ng). In other embodiments, the effective dose is less than about 100 ng. In certain embodiments, the effective dose is about 10 ng to about 100 ng (e.g., about, or no more than about, 10 ng, or 20 ng, or 30 ng, or 40 ng, or 50 ng, or 60 ng, or 70 ng, or 80 ng, or 90 ng, or 100 ng).

- 25 In some embodiments, the effective dose is about 1.4 ng/kg to about 30 ng/kg (e.g., about, or no more than about, 1.4 ng/kg, or 2.5 ng/kg, or 5 ng/kg, or 10 ng/kg, or 15 ng/kg, or 20 ng/kg, or 25 ng/kg, or 30 ng/kg. In other embodiments, the effective dose is less than about 1.5 ng/kg. In certain embodiments, the effective dose is about 0.14 ng/kg to about 1.4 ng/kg (e.g., about, or no more than about, 0.14 ng/kg, or 0.25 ng/kg, or 0.5 ng/kg, or 0.75 ng/kg, or 1 ng/kg, or 1.25 ng/kg, or 1.4 ng/kg).

- 30 In some embodiments, the effective dose is about 350 ng/cm² to about 7000 ng/cm² (e.g., about, or no more than about, 350 ng/cm², or 500 ng/cm², or 750 ng/cm², or 1000 ng/cm², or 2000 ng/cm², or 3000 ng/cm², or 4000 ng/cm², or 5000 ng/cm², or 6000 ng/cm², or 7000 ng/cm²). In other embodiments, the effective dose is less than about 350 ng/cm². In certain embodiments, the effective dose is about 35 ng/cm² to about 350 ng/cm² (e.g.,

about, or no more than about, 35 ng/cm², or 50 ng/cm², or 75 ng/cm², or 100 ng/cm², or 150 ng/cm², or 200 ng/cm², or 250 ng/cm², or 300 ng/cm², or 350 ng/cm².

In some embodiments, the effective dose is about 0.28 picomoles to about 5.7 picomoles (e.g., about, or no more than about, 0.28 picomoles, or 0.5 picomoles, or 0.75 picomoles, or 1 picomole, or 2 picomoles, or 3 picomoles, or 4 picomoles, or 5 picomoles, or 5.7 picomoles). In other embodiments, the effective dose is less than about 0.28 picomoles. In certain embodiments, the effective dose is about 0.028 picomoles to about 0.28 picomoles (e.g., about, or no more than about, 0.028 picomoles, or 0.05 picomoles, or 0.075 picomoles, or 0.1 picomoles, or 0.15 picomoles, or 0.2 picomoles, or 0.25 picomoles, or 0.28 picomoles).

In some embodiments, the effective dose is about 0.004 picomoles/kg to about 0.082 picomoles/kg (e.g., about, or no more than about, 0.004 picomoles/kg, or 0.01 picomoles/kg, or 0.02 picomoles/kg, or 0.03 picomoles/kg, or 0.04 picomoles/kg, or 0.05 picomoles/kg, or 0.06 picomoles/kg, or 0.07 picomoles/kg, or 0.08 picomoles/kg, or 0.082 picomoles/kg). In other embodiments, the effective dose is less than about 0.004 picomoles/kg. In certain embodiments, the effective dose is about 0.0004 picomoles/kg to about 0.004 picomoles/kg (e.g., about, or no more than about, 0.0004 picomoles/kg, or 0.001 picomoles/kg, or 0.002 picomoles/kg, or 0.003 picomoles/kg, or 0.004 picomoles/kg).

In some embodiments, the effective dose is about 1 picomole/cm² to about 20 picomoles/cm² (e.g., about, or no more than about, 1 picomole/cm², or 2 picomoles/cm², or 3 picomoles/cm², or 4 picomoles/cm², or 5 picomoles/cm², or 6 picomoles/cm², or 7 picomoles/cm², or 8 picomoles/cm², or 9 picomoles/cm², or 10 picomoles/cm², or 12 picomoles/cm², or 14 picomoles/cm², or 16 picomoles/cm², or 18 picomoles/cm², or 20 picomoles/cm²). In other embodiments, the effective dose is less than about 1 picomole/cm². In certain embodiments, the effective dose is about 0.1 picomoles/cm² to about 1 picomole/cm² (e.g., about, or no more than about, 0.1 picomoles/cm², or 0.2 picomoles/cm², or 0.3 picomoles/cm², or 0.4 picomoles/cm², or 0.5 picomoles/cm², or 0.6 picomoles/cm², or 0.7 picomoles/cm², or 0.8 picomoles/cm², or 0.9 picomoles/cm², or 1 picomole/cm²).

In various embodiments, the nucleic acid drug is administered in a pharmaceutically acceptable formulation, including a formulation suitable for one or more of injection (e.g. subcutaneous injection, intradermal injection (including to the dermis or epidermis), subdermal injection, intramuscular injection, intraocular injection, intravitreal injection, intra-articular injection, intracardiac injection, intravenous injection, epidural injection, intrathecal injection, intratumoral injection) and topical administration and/or for administration to the integumentary system (e.g. to one or more of the epidermis (optionally selected from the stratum corneum, stratum lucidum, stratum granulosum, stratum spinosum, and stratum germinativum), basement membrane, dermis (optionally selected from the papillary region and the reticular region), subcutis, and conjunctiva) and/or for administration to the eye (e.g., to one or more of the cornea, sclera, iris, lens, corneal limbus, optic nerve, choroid, ciliary body, anterior segment, anterior chamber, and retina).

In various embodiments, the nucleic acid drug is formulated with one or more lipids to enhance uptake of the nucleic acid drug by cells, the lipids optionally selected from **Table 1**. In other embodiments, the nucleic acid drug is formulated with one or more nanoparticles, optionally lipid or polymeric nanoparticles, to enhance uptake of the nucleic acid drug by cells, to enhance duration of protein expression, or to otherwise enhance the safety and/or efficacy of the nucleic acid drug.

In various embodiments, the nucleic acid drug is administered locally, optionally by one or more of subcutaneous injection, intradermal injection, subdermal injection and intramuscular injection, and the effective dose is administered to a surface area of about 4 mm² to about 1000 mm² (e.g. about, or no more than about, 4 mm², or 5 mm², or 10 mm², or 25 mm², or 50 mm², or 75 mm², or 100 mm², or 125 mm², or 150 mm², or 200 mm², or 500 mm², or 1000 mm²).

In various embodiments, the nucleic acid drug is administered in a treatment regimen, optionally with an additional agent or adjuvant therapy described herein, and the administration is about weekly to about once every 24 weeks (e.g. about, or not more than about, weekly, or once every 2 weeks, or once every 3 weeks, or once every 4 weeks, or once every 5 weeks, or once every 6 weeks, or once every 7 weeks, or once every 8 weeks, or once every 9 weeks, or once every 9 weeks, or once every 9 weeks, or once every 9 weeks, or once every 10 weeks, or once every 11 weeks, or once every 12 weeks, or once every 13 weeks, or once every 14 weeks, or once every 15 weeks, or once every 20 weeks, or once every 24 weeks). In other embodiments, the nucleic acid drug is administered in a treatment regimen, optionally with an additional agent or adjuvant therapy described herein, and the administration is about daily to about weekly (e.g., about, or not more than about, daily, or once every 2 days, or once every 3 days, or once every 4 days, or once every 5 days, or once every 6 days, or weekly).

In various embodiments the nucleic acid drug comprises RNA comprising one or more non-canonical nucleotides, optionally having one or more substitutions at the 2C and/or 4C and/or 5C positions for a pyrimidine or the 6C and/or 7N and/or 8C positions for a purine. In various embodiments, the non-canonical nucleotide is one or more of the non-canonical nucleotides described herein, including, for example, 5-hydroxycytidine, 5-methylcytidine, 5-hydroxymethylcytidine, 5-carboxycytidine, 5-formylcytidine, 5-methoxycytidine, pseudouridine, 5-hydroxyuridine, 5-hydroxypseudouridine, 5-methyluridine, 5-methylpseudouridine, 5-hydroxymethyluridine, 5-hydroxymethylpseudouridine, 5-carboxyuridine, 5-carboxypseudouridine, 5-formyluridine, 5-formylpseudouridine, 5-methoxyuridine, and 5-methoxypseudouridine. Further, the RNA comprising one or more non-canonical nucleotides may have one or more of a 5'-UTR comprising a Kozak consensus sequence, a 5'-UTR or 3'-UTR comprising a sequence that increases RNA stability *in vivo* (e.g. an alpha-globin or beta-globin 5'-UTR or an alpha-globin or beta-globin 3'-UTR), and a 3' poly(A) tail from about 20 nucleotides to about 250 nucleotides in length (e.g. about 20, or about 30, or about 40, or about 50, or about 60, or about 70, or about 80, or about 90, or about 100, or about 110, or about 120, or about 130, or about 140, or about 150, or about 160, or about 170, or

about 180, or about 190, or about 200, or about 210, or about 220, or about 230, or about 240, or about 250 nucleotides in length).

Further, some aspects of the methods described herein find use in various medical treatments, including, by way of illustration, treating a disease, disorder and/or condition of the integumentary system or altering, modifying
5 and/or changing the integumentary system (e.g. cosmetically).

Also contemplated are kits suitable for use in human therapy, comprising the nucleic acid drug described herein in a unit dosage form of about 10 ng to about 2000 ng (e.g. about, or no more than about, 10 ng, or 20 ng, or 50 ng, or 100 ng, or 200 ng, or 300 ng, or 400 ng, or 500 ng, or 600 ng, or 700 ng, or 800 ng, or 900 ng, or 1000 ng, or 1100 ng, or 1200 ng, or 1300 ng, or 1400 ng, or 1500 ng, or 1600 ng, or 1700 ng, or 1800 ng, or 1900 ng, or
10 2000 ng) and an injection needle.

Further, in some aspects, the present invention provides a pharmaceutical formulation comprising the nucleic acid drug described herein and one or more of the vehicles (a/k/a "transfection reagents", e.g., lipids) described herein, the formulation optionally being suitable for one or more of subcutaneous injection, intradermal injection, subdermal injection, intramuscular injection, intraocular injection, intravitreal injection, intra-articular injection, intracardiac injection, intravenous injection, epidural injection, intrathecal injection, intratumoral injection, and
15 topical administration.

In some aspects, nucleic acid delivery patches are provided. In one aspect, devices for delivering nucleic acids using electric fields are provided. Other aspects pertain to methods and compositions for delivery of nucleic acids to the skin. Still further aspects pertain to methods and compositions for expression of proteins in the skin.

In one aspect, the invention provides methods and compositions for treating diseases and conditions in humans, including, but not limited to, prophylactic treatments, treatments for rare diseases, including, but not limited to, dermatologic rare diseases, and treatments for use in medical dermatology and aesthetic medicine. In another aspect, the invention provides cosmetics comprising nucleic acids. Still further aspects relate to methods and compositions for altering pigmentation, for example, for the treatment of pigmentation disorders. Still further
25 aspects relate to methods and compositions for enhancing healing, including, but not limited to, healing in response to a wound or surgery. The compositions of the present invention may alter, modify and/or change the appearance of a member of the integumentary system of a subject such as, but not limited, to skin, hair and nails. Such alteration, modification and/or change may be in the context of treatment methods and/or therapeutic uses as described herein including, by way of non-limiting example, dermatological treatments and cosmetics
30 procedures.

Further, in various embodiments, the present invention relates to the targeting of various therapeutic proteins that are not limited to dermatological applications. For example, in various embodiments, the present compositions and methods find use in methods of treatment that are mediated by the increased expression of, for example, various soluble proteins as illustrated herein. In various embodiments, the nucleic acid drug encodes and/or

increases the expression of one or more of a circulating protein, an extracellular matrix protein, an engineered protein, a gene-editing protein, a protein or peptide hormone, an enzyme, erythropoietin, darbepoetin alfa, NOVEPOETIN, elastin, collagen, an antibody or antibody fragment (e.g., a neutralizing antibody or antibody fragment), an intracellular protein, telomerase reverse transcriptase, a membrane protein, a fusion protein, a receptor, a ligand binding domain, a protein inhibitor, or a biologically active fragment, analogue or variant thereof. In other embodiments, administration of the nucleic acid drug results in one or more of an increase in hematocrit, an increase in tissue elasticity, an increase in tissue strength, and increase in skin hydration and/or water retention, hair growth, fat reduction, an insertion, deletion or mutation in DNA, conversion of a prodrug to an active drug, a decrease in tumor size and/or number, a decrease in plaque size and/or number, an increase in vascularization, a decrease in vascularization, an increase in visual acuity, a decrease in pain, an increase in cardiac output (e.g., ejection fraction and stroke volume), a decrease in abnormal heart rhythm, a decrease in fibrosis, a decrease in one or more adverse neurological symptoms, conditions, or disorders (e.g., depression, dysregulation of appetite, polyphagia, anorexia, dementia, headache, fatigue, numbness, tremors, and dizziness), a decrease in erectile dysfunction, an increase in vitality, an increase in pulmonary function, an increase in kidney function, an increase in liver function, an increase in insulin sensitivity, a decrease in insulin sensitivity, a decrease in inflammation, an increase in tear production, an improvement in hearing, an increase in auditory perception, a decrease in tinnitus, a reduction in perspiration, partial or total clearance of an infection, an increase in fertility, a decrease in fertility, inhibition or neutralization of a protein, recruitment or stimulation of one or more components of the immune system, lengthening of telomeres, inhibition of cellular senescence, an increase in replicative potential, reprogramming, proliferation, differentiation, and an increase in differentiation potential.

In some aspects, RNA molecules with low toxicity and high translation efficiency are provided. In one aspect, a cell-culture medium for high-efficiency *in vivo* transfection, reprogramming, and gene editing of cells is provided. Other aspects pertain to methods for producing RNA molecules encoding reprogramming proteins. Still further aspects pertain to methods for producing RNA molecules encoding gene-editing proteins.

In one aspect, the invention provides high-efficiency gene-editing proteins comprising engineered nuclease cleavage domains. In another aspect, the invention provides high-fidelity gene-editing proteins comprising engineered nuclease cleavage domains. Other aspects relate to high-efficiency gene-editing proteins comprising engineered DNA-binding domains. Still further aspects pertain to high-fidelity gene-editing proteins comprising engineered DNA-binding domains. Still further aspects relate to gene-editing proteins comprising engineered repeat sequences. Some aspects relate to methods for altering the DNA sequence of a cell by transfecting the cell with or inducing the cell to express a gene-editing protein. Other aspects relate to methods for altering the DNA sequence of a cell that is present in an *in vitro* culture. Still further aspects relate to methods for altering the DNA sequence of a cell that is present *in vivo*.

In some aspects, the invention provides methods for treating cancer comprising administering to a patient a therapeutically effective amount of a gene-editing protein or a nucleic-acid encoding a gene-editing protein. In one aspect, the gene-editing protein is capable of altering the DNA sequence of a cancer associated gene. In another aspect, the cancer-associated gene is the BIRC5 gene. Still other aspects relate to therapeutics comprising nucleic acids and/or cells and methods of using therapeutics comprising nucleic acids and/or cells for the treatment of, for example, type 1 diabetes, heart disease, including ischemic and dilated cardiomyopathy, macular degeneration, Parkinson's disease, cystic fibrosis, sickle-cell anemia, thalassemia, Fanconi anemia, severe combined immunodeficiency, hereditary sensory neuropathy, xeroderma pigmentosum, Huntington's disease, muscular dystrophy, amyotrophic lateral sclerosis, Alzheimer's disease, cancer, and infectious diseases including hepatitis and HIV/AIDS. In some aspects, the nucleic acids comprise RNA. In other aspects, the RNA comprises one or more non-canonical nucleotides. In still other aspects, the nucleic acids are delivered to cells using a virus. In some aspects, the virus is a replication-competent virus. In other aspects, the virus is a replication-incompetent virus.

The details of the invention are set forth in the accompanying description below. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, illustrative methods and materials are now described. Other features, objects, and advantages of the invention will be apparent from the description and from the claims. In the specification and the appended claims, the singular forms also include the plural unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.

DETAILED DESCRIPTION OF THE FIGURES

FIG. 1 depicts primary adult human dermal fibroblasts transfected with RNA encoding green fluorescent protein ("GFP") and comprising the indicated nucleotides.

FIG. 2 depicts the result of a gene-expression analysis of the primary adult human dermal fibroblasts of **FIG. 1** using a one-step real-time RT-PCR and primers designed to amplify human interferon beta mRNA. Data were normalized to the untransfected sample ("Neg."). GAPDH was used as a loading control.

FIG. 3 depicts the results of a gene-expression analysis of cells transfected with RNA comprising the indicated nucleotides, conducted as in **FIG. 2**. Data were normalized to the untransfected sample ("Neg."). GAPDH was used as a loading control.

FIG. 4 depicts the results of a gene-expression analysis of cells transfected with RNA comprising the indicated nucleotides, conducted as in **FIG. 2**, and using primers designed to amplify the indicated transcripts. Data were normalized to the untransfected sample ("Neg."). GAPDH was used as a loading control.

FIG. 5 depicts the results of a gene-expression analysis of primary human epidermal keratinocytes transfected with RNA encoding NOVEPOETIN, and comprising the indicated nucleotides, conducted as in **FIG. 2**. Data were normalized to the untransfected sample ("Neg."). GAPDH was used as a loading control.

FIG. 6 depicts intradermal injection of a solution comprising RNA encoding GFP into the ventral forearm of a healthy, 33 year-old, 70 kg, male human subject.

FIG. 7 depicts a region of the ventral forearm of the subject shown in **FIG. 6** after treatment with RNA comprising 5-methoxyuridine and encoding GFP (injection sites 1-3) or COL7 (injection site 4). The image was taken immediately following the final injection.

FIG. 8 depicts the region of **FIG. 7**, 24 hours after injection.

FIG. 9 depicts the results of fluorescent imaging of the region of **FIG. 7**, using the indicated fluorescent channels. The dose at each injection site is also indicated. Images were taken 24 hours after injection.

FIG. 10 depicts the results of fluorescent imaging of the region of **FIG. 7**, using the FITC fluorescent channel. The dose at each injection site is indicated. Images were taken 48 hours after injection.

FIG. 11 depicts the results of quantitative fluorescent imaging of the region of **FIG. 7**, using the FITC fluorescent channel. The horizontal axis indicates time after injection.

FIG. 12 depicts the results of fluorescent imaging of an independent experiment in which a region of the ventral forearm of as the subject shown in **FIG. 6** treated with RNA comprising 5-methoxyuridine and encoding GFP. The image was taken 24 hours after injection.

FIG. 13 depicts the results of an ELISA designed to detect darbepoetin alfa in culture media of primary human epidermal keratinocytes transfected with RNA comprising the indicated nucleotides and encoding NOVEPOETIN.

FIG. 14 depicts the results of an ELISA designed to detect darbepoetin alfa in culture media of primary human epidermal keratinocytes transfected with RNA comprising the indicated nucleotides and encoding NOVEPOETIN.

FIG. 15 depicts the results of an ELISA designed to detect darbepoetin alfa in culture media of primary human epidermal keratinocytes transfected with RNA comprising the indicated nucleotides and encoding NOVEPOETIN.

FIG. 16 depicts primary human dermal fibroblasts transfected with RNA comprising 5-methoxyuridine and encoding hTERT. Cells were fixed and stained using an antibody targeting hTERT 24 hours after transfection.

FIG. 17 depicts primary adult human dermal fibroblasts transfected with RNA encoding green fluorescent protein ("GFP"), prepared and stored as indicated.

FIG. 18 depicts single administration of NOVECRIT induced a rapid increase and sustained level of NOVEPOIETIN in serum. The Y axis shows concentration of NOVEPOIETIN protein (mU/mL).

FIG. 19 depicts a single administration of NOVECRIT stimulated erythropoiesis, yielding elevated hematocrit for at least 14 days. The left panel shows % hematocrit on the Y axis, while the right panel shows % reticulocytes.

FIG. 20 depicts a table summarizing TNF α , IL-6, and IFN α cytokine levels in plasma samples collected from a maximum tolerated dose of NOVECRIT in male Sprague Dawley rats study of Example 35.

FIG. 21 depicts a SURVEYOR assay using the DNA of primary adult human dermal fibroblasts transfected with RNA TALENs targeting the sequence TGAGCAGAAAGTGGCTCAGTG (SEQ ID NO:467) and
5 TGGCTGTACAGCTACACCCC (SEQ ID NO: 468), located within the COL7A1 gene. The bands present in the +RNA lane indicate editing of a region of the gene that is frequently involved in dystrophic epidermolysis bullosa.

FIG. 22 depicts a SURVEYOR assay using the DNA of primary adult human dermal fibroblasts transfected with RNA TALENs targeting the sequence TTCCACTCCTGCAGGGCCCC (SEQ ID NO:469) and
10 TCGCCCTTCAGCCCGCGTTC (SEQ ID NO:470), located within the COL7A1 gene. The bands present in the +RNA lane indicate editing of a region of the gene that is frequently involved in dystrophic epidermolysis bullosa.

FIG. 23 shows the immunogenicity of various synthetic RNA constructs in the context of a gene-editing (*i.e.* unmodified nucleotides "A,G,U,C"; pseudouridine only "psU"; 5-methylcytidine only "5mC"; both pseudouridine and 5-methylcytidine "psU+5mC"; and a negative control "neg").

FIG. 24 shows the gene-editing activity in cells transfected with various synthetic RNA constructs (*i.e.* unmodified
15 nucleotides "A,G,U,C"; psuedouridine only "psU"; 5-methylcytidine only "5mC"; both psuedouridine and 5-methylcytidine "psU+5mC"; and a negative control "neg").

FIG. 25 depicts gene editing of the COL7A1 gene in primary human epidermal keratinocytes transfected with RNA encoding TALENs and a single-stranded DNA repair template ("RT") of the indicated length. The presence of bands at the locations shown by asterisks ("**") indicates successful gene editing.

FIG. 26 depicts gene editing of the COL7A1 gene in primary human epidermal keratinocytes transfected with
20 RNA encoding TALENs and an 80 nt single-stranded DNA repair template ("RT") at the indicated ratios of RNA to repair template. The presence of bands at the locations shown by asterisks ("**") indicates successful gene editing.

FIG. 27 depicts gene correction of the COL7A1 gene in primary human epidermal keratinocytes transfected with
25 RNA encoding TALENs and a single-stranded DNA repair template ("RT") of the indicated length. The presence of bands at the locations shown by asterisks ("**") indicates successful gene correction.

FIG. 28 depicts gene correction of the COL7A1 gene in primary human epidermal keratinocytes transfected with
30 RNA encoding TALENs and an 80 nt single-stranded DNA repair template ("RT") at the indicated ratios of RNA to repair template. The presence of bands at the locations shown by asterisks ("**") indicates successful gene correction.

FIG. 29 depicts gene editing ("T7E1") and correction ("Digestion") of the COL7A1 gene in primary human epidermal keratinocytes transfected with RNA encoding TALENs and an 80 nt single-stranded DNA repair

template ("RT"). The presence of bands at the locations shown by asterisks ("**") indicates successful gene editing ("T7E1") and correction ("Digestion").

FIG. 30 depicts the amount of BMP7 protein secreted by primary human dermal fibroblasts and primary human epidermal keratinocytes transfected with RNA encoding the indicated BMP7 variant, measured by ELISA.

5 Asterisks ("**") indicate saturation of the ELISA assay.

FIG. 31 depicts the amount of parathyroid hormone secreted by primary human epidermal keratinocytes transfected with RNA encoding PTH, measured by ELISA.

FIG. 32 demonstrates that repeated administration of NOVECRIT stimulated erythropoiesis, yielding elevated hematocrit for at least 14 days. For each data set, the order of histograms from left to right is: Group 1, Group 2,
10 Group 3, Group 4, and Group 5.

FIG. 33 provides an illustrative experimental design for studying the effects of RNAs encoding BMP7 variants for the prevention and treatment of diabetic nephropathy.

FIG. 34 depicts BMP7 protein levels in rats treated with RNAs encoding BMP7 variants as described in Example 42. Error bars indicate SEM (n=6).

15 **FIG. 35** depicts the urine volume, urine albumin, and urine creatinine levels in rats treated with RNAs encoding BMP7 variants as described in Example 42. For each data set, the order of histograms from left to right is: urine volume, urine creatinine, and urine albumin.

FIG. 36 depicts the effect of RNAs encoding BMP7 variants in treating diabetic nephropathy as described in Example 42. For each data set, the order of histograms from left to right is: Group 6 – vehicle and Group 7 –
20 FTB-2 (BMP7 variant A).

FIG. 37, panels A-I, depict the results of gene-expression analysis of human epidermal keratinocytes transfected with RNA encoding BDNF, BMP-2, BMP-6, IL-2, IL-6, IL-15, IL-22, LIF or FGF-21.

FIG. 38 depicts the results of gene-expression analysis of human epidermal keratinocytes transfected with RNA encoding IL-15 or IL-15 and IL-15RA.

25 **FIG. 39** depicts the serum levels of FGF21, IL15, IL6, IL22, and Novepoietin following a single intradermal injection of various RNAs encoding these proteins as described in Example 45. Three rats were analyzed for each RNA tested.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is based, in part, on the discovery of a safe and effective dosing strategy for nucleic acid
30 drugs, including RNA, such as RNA comprising non-canonical (or "modified") nucleotides, in humans. The inventors believe this to be the first report of safe and effective dosing of RNA molecules, including those comprising non-canonical nucleotides, in humans. Despite reports in the art that very large doses of RNA

molecules are needed for mammalian dosing, and minimal therapeutic effect is achieved despite high dosing (see, e.g. US Patent Publication No. 2013/0245103), the present inventors have surprisingly managed to dose synthetic RNA in a human and achieve significant target protein expression with minimal immunological or other side effects.

5 In various embodiments, the present invention provides improved doses, formulations, administration and methods of use of nucleic acid drugs, which include RNA, which may contain non-canonical nucleotides (e.g. a residue other than adenine, guanine, thymine, uracil, and cytosine or the standard nucleoside, nucleotide, deoxynucleoside or deoxynucleotide derivatives thereof). In various embodiments, the RNA comprising non-canonical nucleotides leads to the expression of a protein encoded by the RNA, the protein often being one of
10 therapeutic benefit (sometimes called the "target" or "protein of interest"). Further, this expression of therapeutic protein is achieved with minimal or negligible toxicity.

In various aspects, the present invention is based on the surprising discovery of safe and effective doses and administration parameters of nucleic acid drugs for human subjects. Nucleic acid drugs include a dsDNA molecule, a ssDNA molecule, a RNA molecule, a dsRNA molecule, a ssRNA molecule, a plasmid, an
15 oligonucleotide, a synthetic RNA molecule, a miRNA molecule, an mRNA molecule, and an siRNA molecule. In various embodiments, the RNA comprises non-canonical nucleotides.

In some aspects, there is provided a method for delivering a nucleic acid drug, comprising administering an effective dose of a nucleic acid drug to a human subject in need thereof, wherein the nucleic acid drug comprises a synthetic RNA. In various embodiments, the effective dose is an amount sufficient to substantially increase an
20 amount of a protein encoded by the nucleic acid drug in the human subject. For example, when the nucleic acid drug is a synthetic RNA comprising one or more modified nucleotides, the nucleic acid drug may result in higher protein expression than levels obtainable with a nucleic acid drug that does not comprise one or more modified nucleotides (e.g., RNA comprising the canonical nucleotides A, G, U, and C). In some embodiments, the nucleic acid drug results in about a 2-fold, or about a 3-fold, or about a 4-fold, or about a 5-fold, or about a 10-fold, or
25 about a 15-fold, or about a 20-fold, or about a 25-fold, or about a 30-fold, or about a 35-fold, or about a 40-fold, or about a 45-fold, or about a 50-fold, or about a 100-fold increase in protein expression as compared to levels obtainable with a nucleic acid drug that does not comprise one or more modified nucleotides.

In some embodiments, the nucleic acid drug provides a sustained therapeutic effect that is optionally mediated by a sustained expression of target protein. For instance, in some embodiments, the therapeutic effect is present
30 for over about 1 day, or over about 2 days, or over about 3 days, or over about 4 days, or over about 5 days, or over about 6 days, or over about 7 days, or over about 8 days, or over about 9 days, or over about 10 days, or over about 14 days after administration. In some embodiments, this sustained effect obviates the need for, or reduces the amount of, maintenance doses.

In some embodiments, the nucleic acid drug provides a sustained target protein level. For instance, in some embodiments, the target protein is present (e.g. in measurable amounts, e.g. in the serum of a patient to whom the nucleic acid drug has been administered) for over about 1 day, or over about 2 days, or over about 3 days, or over about 4 days, or over about 5 days, or over about 6 days, or over about 7 days, or over about 8 days, or over about 9 days, or over about 10 days, or over about 14 days after administration. In some embodiments, this sustained effect obviates the need for, or reduces the amount of, maintenance doses.

In various embodiments, the nucleic acid drug provides therapeutic action without sustained presence of the nucleic acid drug itself. In some embodiments, the nucleic acid drug is rapidly metabolized, for instance, within about 6 hours, or about 12 hours, or about 18 hours, or about 24 hours, or about 2 days, or about 3 days, or about 4 days, or about 5 days, or about 1 week from administration.

In various embodiments, the effective dose is an amount that substantially avoids cell toxicity *in vivo*. In various embodiments, the effective dose is an amount that substantially avoids an immune reaction in a human subject. For example, the immune reaction may be an immune response mediated by the innate immune system. Immune response can be monitored using markers known in the art (e.g. cytokines, interferons, TLRs). In some embodiments, the effective dose obviates the need for treatment of the human subject with immune suppressants agents (e.g. B18R) used to moderate the residual toxicity. Accordingly, in some embodiments, the present methods allow for dosing that provides increased protein expression and reduces toxicity.

In some embodiments, the immune response is reduced by about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 95%, about 99%, about 99.9%, or greater than about 99.9% as compared to the immune response induced by a corresponding unmodified nucleic acid. In some embodiments, upregulation of one or more immune response markers is reduced by about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 95%, about 99%, about 99.9%, or greater than about 99.9% as compared to the upregulation of the one or more immune response markers induced by a corresponding unmodified nucleic acid. In some embodiments, the immune response marker comprises an mRNA or protein product of an interferon gene, including an interferon alpha gene, IFNB1, TLR3, RARRES3, EIF2AK2, STAT1, STAT2, IFIT1, IFIT2, IFIT3, IFIT5, OAS1, OAS2, OAS3, OASL, ISG20 or a fragment, variant, analogue, or family-member thereof. In some embodiments, the immune response marker comprises an mRNA or protein product of a TNF gene, including a TNF alpha gene, TNFRSF1A; TNFRSF1B; LTBR; TNFRSF4; CD40; FAS; TNFRSF6B; CD27; TNFRSF8; TNFRSF9; TNFRSF10A; TNFRSF10B; TNFRSF10C; TNFRSF10D; TNFRSF11A; TNFRSF11B; TNFRSF12A; TNFRSF13B; TNFRSF13C; TNFRSF14; NGFR; TNFRSF17; TNFRSF18; TNFRSF19; TNFRSF21; TNFRSF25; and EDA2R or a fragment, variant, analogue, or family-member thereof. In some embodiments, the immune response marker comprises an mRNA or protein product of an interleukin gene, including an IL-6 gene, IL-1; IL-2; IL-3; IL-4; IL-5; IL-6; IL-7; IL-8 or CXCL8; IL-9; IL-10; IL-11; IL-12; IL-13; IL-14; IL-15; IL-16; IL-17; IL-18; IL-19;

IL-20; IL-21; IL-22; IL-23; IL-24; IL-25; IL-26; IL-27; IL-28; IL-29; IL-30; IL-31; IL-32; IL-33; IL-35; IL-36 or a fragment, variant, analogue, or family-member thereof.

In some embodiments, cell death is about 10%, about 25%, about 50%, about 75%, about 85%, about 90%, about 95%, or over about 95% less than the cell death observed with a corresponding unmodified nucleic acid.

5 Moreover, cell death may affect fewer than about 50%, about 40%, about 30%, about 20%, about 10%, about 5%, about 1%, about 0.1%, about 0.01% or fewer than about 0.01% of cells contacted with the modified nucleic acids.

In some embodiments, there is provided a method for expressing a protein of interest in a population of cells in a mammalian subject, comprising administering a non-viral transfection composition comprising an effective dose
10 of a RNA encoding the protein of interest to said cells, the RNA containing one or more non-canonical nucleotides that avoid substantial cellular toxicity, where the transfection composition is administered in an amount that allows for expression of said protein in said cells for at least about five days (e.g. about 5, or about 6, or about 7, about 8, or about 9, or about 10, or about 14 days) without substantial cellular toxicity. In some
15 embodiments, there is provided a method for expressing a protein of interest in a population of cells in a mammalian subject, comprising administering a non-viral transfection composition comprising an effective dose of a RNA encoding the protein of interest to said cells, the RNA containing one or more non-canonical nucleotides that avoid substantial cellular toxicity, where the transfection composition is administered in an amount that allows for expression of said protein in said cells for at least about six hours (e.g. about six hours, or
20 about 12 hours, or about 1 day, or about 2 days, or about 3 days, or about 4 days, or about 5 days) without substantial cellular toxicity.

In some embodiments, the effective dose of the nucleic acid drug, including synthetic RNA, is about 100 ng to about 2000 ng, or about 200 ng to about 1900 ng, or about 300 ng to about 1800 ng, or about 400 ng to about 1700 ng, or about 500 ng to about 1600 ng, or about 600 ng to about 1500 ng, or about 700 ng to about 1400 ng, or about 800 ng to about 1300 ng, or about 900 ng to about 1200 ng, or about 1000 ng to about 1100 ng, or
25 about 500 ng to about 2000 ng, or about 500 ng to about 1500 ng, or about 500 ng to about 1000 ng, or about 1000 ng to about 1500 ng, or about 1000 ng to about 2000 ng, or about 1500 ng to about 2000 ng, or about 100 ng to about 500 ng, or about 200 ng to about 400 ng, or about 10 ng to about 100 ng, or about 20 ng to about 90 ng, or about 30 ng to about 80 ng, or about 40 ng to about 70 ng, or about 50 ng to about 60 ng.

In some embodiments, the effective dose of the nucleic acid drug, including synthetic RNA, is no more than
30 about 50 ng, or about 100 ng, or about 200 ng, or about 300 ng, or about 400 ng, or about 500 ng, or about 600 ng, or about 700 ng, or about 800 ng, or about 900 ng, or about 1000 ng, or about 1100 ng, or about 1200 ng, or about 1300 ng, or about 1400 ng, or about 1500 ng, or about 1600 ng, or about 1700 ng, or about 1800 ng, or about 1900 ng, or about 2000 ng.

In some embodiments, the effective dose of the nucleic acid drug, including synthetic RNA, is about 50 ng, or about 100 ng, or about 200 ng, or about 300 ng, or about 400 ng, or about 500 ng, or about 600 ng, or about 700 ng, or about 800 ng, or about 900 ng, or about 1000 ng, or about 1100 ng, or about 1200 ng, or about 1300 ng, or about 1400 ng, or about 1500 ng, or about 1600 ng, or about 1700 ng, or about 1800 ng, or about 1900 ng, or about 2000 ng.

In some embodiments, the effective dose of the nucleic acid drug, including synthetic RNA, is about 0.028 pmol, or about 0.05 pmol, or about 0.1 pmol, or about 0.2 pmol, or about 0.3 pmol, or about 0.4 pmol, or about 0.5 pmol, or about 0.6 pmol, or about 0.7 pmol, or about 0.8 pmol, or about 0.9 pmol, or about 1.0 pmol, or about 1.2 pmol, or about 1.4 pmol, or about 1.6 pmol, or about 1.8 pmol, or about 2.0 pmol, or about 2.2 pmol, or about 2.4 pmol, or about 2.6 pmol, or about 2.8 pmol, or about 3.0 pmol, or about 3.2 pmol, or about 3.4 pmol, or about 3.6 pmol, or about 3.8 pmol, or about 4.0 pmol, or about 4.2 pmol, or about 4.4 pmol, or about 4.6 pmol, or about 4.8 pmol, or about 5.0 pmol, or about 5.5 pmol, or about 5.7 pmol.

In some embodiments, the nucleic acid drug, including synthetic RNA, is administered at a concentration of about 0.1 nM, or about 0.25 nM, or about 0.5 nM, or about 0.75 nM, or about 1 nM, or about 2.5 nM, or about 5 nM, or about 7.5 nM, or about 10 nM, or about 20 nM, or about 30 nM, or about 40 nM, or about 50 nM, or about 60 nM, or about 70 nM, or about 80 nM, or about 90 nM, or about 100 nM, or about 110 nM, or about 120 nM, or about 150 nM, or about 175 nM, or about 200 nM.

In some embodiments, the effective dose of the nucleic acid drug is about 350 ng/cm², or about 500 ng/cm², or about 750 ng/cm², or about 1000 ng/cm², or about 2000 ng/cm², or about 3000 ng/cm², or about 4000 ng/cm², or about 5000 ng/cm², or about 6000 ng/cm², or about 7000 ng/cm². In other embodiments, the effective dose is less than about 350 ng/cm². In certain embodiments, the effective dose is about 35 ng/cm², or about 50 ng/cm², or about 75 ng/cm², or about 100 ng/cm², or about 150 ng/cm², or about 200 ng/cm², or about 250 ng/cm², or about 300 ng/cm², or about 350 ng/cm².

In some embodiments, the effective dose of the nucleic acid drug is about 35 ng/cm² to about 7000 ng/cm², or about 50 ng/cm² to about 5000 ng/cm², or about 100 ng/cm² to about 3000 ng/cm², or about 500 ng/cm² to about 2000 ng/cm², or about 750 ng/cm² to about 1500 ng/cm², or about 800 ng/cm² to about 1200 ng/cm², or about 900 ng/cm² to about 1100 ng/cm².

In some embodiments, the effective dose of the nucleic acid drug is about 1 picomole/cm², or about 2 picomoles/cm², or about 3 picomoles/cm², or about 4 picomoles/cm², or about 5 picomoles/cm², or about 6 picomoles/cm², or about 7 picomoles/cm², or about 8 picomoles/cm², or about 9 picomoles/cm², or about 10 picomoles/cm², or about 12 picomoles/cm², or about 14 picomoles/cm², or about 16 picomoles/cm², or about 18 picomoles/cm², or about 20 picomoles/cm². In other embodiments, the effective dose is less than about 1 picomole/cm². In certain embodiments, the effective dose is about 0.1 picomoles/cm², or about 0.2 picomoles/cm², or about 0.3 picomoles/cm², or about 0.4 picomoles/cm², or about 0.5 picomoles/cm², or about

0.6 picomoles/cm², or about 0.7 picomoles/cm², or about 0.8 picomoles/cm², or about 0.9 picomoles/cm², or about 1 picomole/cm².

In some embodiments, the effective dose of the nucleic acid drug is about 0.1 picomoles/cm² to about 20 picomoles/cm², or about 0.2 picomoles/cm² to about 15 picomoles/cm², or about 0.5 picomoles/cm² to about 10 picomoles/cm², or about 0.8 picomoles/cm² to about 8 picomoles/cm², or about 1 picomole/cm² to about 5 picomoles/cm², or about 2 picomoles/cm² to about 4 picomoles/cm².

In various embodiments, the nucleic acid drug, including synthetic RNA, is administered in a pharmaceutically acceptable formulation. In various embodiments, the nucleic acid drug, including synthetic RNA, is formulated for one or more of injection and topical administration. By way of example, the nucleic acid drug, including synthetic RNA, may be formulated for injection to a tissue of interest, e.g. a disease site (by way of non-limiting example, a tumor). In various embodiments, injection includes delivery via a patch. In some embodiments, the delivery is mediated by electrical stimulation. In various embodiments, the nucleic acid drug, including synthetic RNA, is formulated for administration to one or more of the epidermis (optionally selected from the stratum corneum, stratum lucidum, stratum granulosum, stratum spinosum, and stratum germinativum), basement membrane, dermis (optionally selected from the papillary region and the reticular region), subcutis, conjunctiva cornea, sclera, iris, lens, corneal limbus, optic nerve, choroid, ciliary body, anterior segment, anterior chamber, and retina. In various embodiments, the nucleic acid drug, including synthetic RNA, is formulated for one or more of subcutaneous injection, intradermal injection, subdermal injection, intramuscular injection, intraocular injection, intravitreal injection, intra-articular injection, intracardiac injection, intravenous injection, epidural injection, intrathecal injection, intratumoral injection, and topical administration. In various embodiments, the nucleic acid drug, including synthetic RNA, is formulated for intradermal (ID) injection to one or more of the dermis or epidermis. In various embodiments, the nucleic acid drug, including synthetic RNA, is administered in a manner such that it effects one or more of keratinocytes and fibroblasts (e.g. causes these cells to express one or more therapeutic proteins).

Accordingly, the present invention provides various formulations as described herein. Further, in some embodiments, the formulations described herein find use in the various delivery and/or treatment methods of the present invention. For instance, formulations can comprise a vesicle, for instance, a liposome (see Langer, 1990, *Science* 249:1527-1533; Treat *et al.*, in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353-365 (1989). In various embodiments, the formulation comprises an aqueous suspension of liposomes. Illustrative liposome components are set forth in **Table 1**, and are given by way of example, and not by way of limitation. In various embodiments, one or more, or two or more, or three or more, or four or more, or five or more of the lipids of **Table 1** are combined in a formulation.

Table 1. Illustrative Biocompatible Lipids

1	3β-[N-(N',N'-dimethylaminoethane)-carbamoyl]cholesterol (DC-Cholesterol)
---	--

2	1,2-dioleoyl-3-trimethylammonium-propane (DOTAP / 18:1 TAP)
3	N-(4-carboxybenzyl)-N,N-dimethyl-2,3-bis(oleoyloxy)propan-1-aminium (DOBAQ)
4	1,2-dimyristoyl-3-trimethylammonium-propane (14:0 TAP)
5	1,2-dipalmitoyl-3-trimethylammonium-propane (16:0 TAP)
6	1,2-stearoyl-3-trimethylammonium-propane (18:0 TAP)
7	1,2-dioleoyl-3-dimethylammonium-propane (DODAP / 18:1 DAP)
8	1,2-dimyristoyl-3-dimethylammonium-propane (14:0 DAP)
9	1,2-dipalmitoyl-3-dimethylammonium-propane (16:0 DAP)
10	1,2-distearoyl-3-dimethylammonium-propane (18:0 DAP)
11	dimethyldioctadecylammonium (18:0 DDAB)
12	1,2-dilauroyl-sn-glycero-3-ethylphosphocholine (12:0 EthylPC)
13	1,2-dimyristoyl-sn-glycero-3-ethylphosphocholine (14:0 EthylPC)
14	1,2-dimyristoleoyl-sn-glycero-3-ethylphosphocholine (14:1 EthylPC)
15	1,2-dipalmitoyl-sn-glycero-3-ethylphosphocholine (16:0 EthylPC)
16	1,2-distearoyl-sn-glycero-3-ethylphosphocholine (18:0 EthylPC)
17	1,2-dioleoyl-sn-glycero-3-ethylphosphocholine (18:1 EthylPC)
18	1-palmitoyl-2-oleoyl-sn-glycero-3-ethylphosphocholine (16:1-18:1 EthylPC)
19	1,2-di-O-octadecenyl-3-trimethylammonium propane (DOTMA)
20	N1-[2-((1S)-1-[(3-aminopropyl)amino]-4-[di(3-amino-propyl)amino]butylcarboxamido)ethyl]-3,4-di[oleyloxy]-benzamide (MVL5)
21	2,3-dioleyloxy-N-[2-spermine carboxamide]ethyl-N,N-dimethyl-1-propanammonium trifluoroacetate (DOSPA)
22	1,3-di-oleyloxy-2-(6-carboxy-spermyl)-propylamid (DOSPER)
23	N-[1-(2,3-dimyristyloxy)propyl]-N,N-dimethyl-N-(2-hydroxyethyl)ammonium bromide (DMRIE)
24	LIPOFECTAMINE, LIPOFECTAMINE 2000, LIPOFECTAMINE RNAiMAX, LIPOFECTAMINE 3000, LIPOFECTAMINE MessengerMAX, TransIT mRNA
25	dioctadecyl amidoglycerol spermine (DOGS)
26	dioleoyl phosphatidyl ethanolamine (DOPE)

In some embodiments, the liposomes include LIPOFECTAMINE 3000. In some embodiments, the liposomes include one or more lipids described in US Patent Nos. 4,897,355 or 7,479,573 or in International Patent Publication No. WO/2015/089487, or in Felgner, P. L. *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:7413–7417, the entire contents of each is incorporated by reference in their entireties).

In some embodiments, the liposome comprises N-[1-(2,3-dioleyloxy)propyl]-N,N,N-trimethylammonium chloride (DOTMA). In some embodiments, the liposome comprises dioleoylphosphatidylethanolamine (DOPE).

In one embodiment, the liposomes include one or more polyethylene glycol (PEG) chains, optionally selected from PEG200, PEG300, PEG400, PEG600, PEG800, PEG1000, PEG1500, PEG2000, PEG3000, and PEG4000. In some embodiments, the PEG is PEG2000. In some embodiments, the liposomes include 1,2-distearoyl-sn-glycero-3-phosphoethanolamine (DSPE) or a derivative thereof.

In some embodiments, the formulation comprises one or more of N-(carbonyl-ethoxypolyethylene glycol 2000)-1,2-distearoyl-sn-glycero-3-phosphoethanolamine (MPEG2000-DSPE), fully hydrogenated phosphatidylcholine, cholesterol, LIPOFECTAMINE 3000, a cationic lipid, a polycationic lipid, and 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[folate(polyethylene glycol)-5000] (FA-MPEG5000-DSPE).

- 5 In one embodiment, the formulation comprises about 3.2 mg/mL N-(carbonyl-ethoxypolyethylene glycol 2000)-1,2-distearoyl-sn-glycero-3-phosphoethanolamine (MPEG2000-DSPE), about 9.6 mg/mL fully hydrogenated phosphatidylcholine, about 3.2 mg/mL cholesterol, about 2 mg/mL ammonium sulfate, and histidine as a buffer, with about 0.27 mg/mL 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[folate(polyethylene glycol)-5000] (FA-MPEG5000-DSPE) added to the lipid mixture. In another embodiment, the nucleic acids are complexed by
- 10 combining 1 μ L of LIPOFECTAMINE 3000 per about 1 μ g of nucleic acid and incubating at room temperature for at least about 5 minutes. In one embodiment, the LIPOFECTAMINE 3000 is a solution comprising a lipid at a concentration of about 1 mg/mL. In some embodiments, nucleic acids are encapsulated by combining about 10 μ g of the liposome formulation per about 1 μ g of nucleic acid and incubating at room temperature for about 5 minutes.
- 15 In some embodiments, the formulation comprises one or more nanoparticles. In one embodiment, the nanoparticle is a polymeric nanoparticle. In various embodiments, the formulation comprises one or more of a diblock copolymer, a triblock copolymer, a tetrablock copolymer, and a multiblock copolymer. In various embodiments, the formulation comprises one or more of polymeric nanoparticles comprising a polyethylene glycol (PEG)-modified polylactic acid (PLA) diblock copolymer (PLA-PEG) and PEG-polypropylene glycol-PEG-
- 20 modified PLA-tetrablock copolymer (PLA-PEG-PPG-PEG).

In some embodiments, the formulation comprises one or more lipids that are described in WO/2000/027795, the entire contents of which are hereby incorporated by reference.

- In one embodiment, the therapeutic comprises one or more ligands. In another embodiment, the therapeutic comprises at least one of: androgen, CD30 (TNFRSF8), a cell-penetrating peptide, CXCR, estrogen, epidermal
- 25 growth factor, EGFR, HER2, folate, insulin, insulin-like growth factor-I, interleukin-13, integrin, progesterone, stromal-derived-factor-1, thrombin, vitamin D, and transferrin or a biologically active fragment or variant thereof.

- The active compositions of the present invention may include classic pharmaceutical preparations. Administration of these compositions according to the present invention may be via any common route so long as the target tissue is available via that route. This includes oral, nasal, or buccal. Alternatively, administration
- 30 may be by intradermal, subcutaneous, intramuscular, intraperitoneal or intravenous injection, or by direct injection into diseased, e.g. cancer, tissue. The agents disclosed herein may also be administered by catheter systems. Such compositions would normally be administered as pharmaceutically acceptable compositions as described herein.

Administration of the compositions described herein may be, for example, by injection, topical administration, ophthalmic administration and intranasal administration. The injection, in some embodiments, may be linked to an electrical force (e.g. electroporation, including with devices that find use in electrochemotherapy (e.g. CLINIPORATOR, IGEA Srl, Carpi [MO], Italy)). The topical administration may be, but is not limited to, a cream, lotion, ointment, gel, spray, solution and the like. The topical administration may further include a penetration enhancer such as, but not limited to, surfactants, fatty acids, bile salts, chelating agents, non-chelating non-surfactants, polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether, fatty acids and/or salts in combination with bile acids and/or salts, sodium salt in combination with lauric acid, capric acid and UDCA, and the like. The topical administration may also include a fragrance, a colorant, a sunscreen, an antibacterial and/or a moisturizer. The compositions described herein may be administered to at least one site such as, but not limited to, forehead, scalp, hair follicles, hair, upper eyelids, lower eyelids, eyebrows, eyelashes, infraorbital area, periorbital areas, temple, nose, nose bridge, cheeks, tongue, nasolabial folds, lips, periobicular areas, jaw line, ears, neck, breast, forearm, upper arm, palm, hand, finger, nails, back, abdomen, sides, buttocks, thigh, calf, feet, toes and the like.

Routes of administration include, for example: intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, oral, sublingual, intranasal, intracerebral, intravaginal, transdermal, rectally, by inhalation, or topically, particularly to the ears, nose, eyes, or skin. In some embodiments, the administering is effected orally or by parenteral injection.

Upon formulation, solutions may be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective, as described herein. The formulations may easily be administered in a variety of dosage forms such as injectable solutions, drug release capsules and the like. For parenteral administration in an aqueous solution, for example, the solution generally is suitably buffered and the liquid diluent first rendered isotonic with, for example, sufficient saline or glucose. Such aqueous solutions may be used, for example, for intravenous, intramuscular, subcutaneous and intraperitoneal administration. Preferably, sterile aqueous media are employed as is known to those of skill in the art, particularly in light of the present disclosure.

In various embodiments, the nucleic acid drug, including RNA comprising one or more non-canonical nucleotides, and/or formulations comprising the same, is administered locally, optionally by one or more of subcutaneous injection, intradermal injection, subdermal injection and intramuscular injection, and the effective dose is administered to a surface area of about 4 mm² to about 150 mm² (e.g. about, or no more than about, 4 mm², or about 5 mm², or about 6 mm², or about 7 mm², or about 8 mm², or about 10 mm², or about 20 mm², or about 50 mm², or about 100 mm², or about 150 mm²). In various embodiments, the nucleic acid drug, including RNA comprising one or more non-canonical nucleotides, and/or formulations comprising the same, is administered locally, optionally by one or more of subcutaneous injection, intradermal injection, subdermal injection and intramuscular injection, and the effective dose administered to a surface area of no more than about

4 mm², or about 5 mm², or about 6mm², or about 7 mm², or about 8 mm², or about 10 mm², or about 20 mm², or about 50 mm², or about 100 mm², or about 150 mm². In various embodiments, the nucleic acid drug, including RNA comprising one or more non-canonical nucleotides, and/or formulations comprising the same, is administered locally, optionally by one or more of subcutaneous injection, intradermal injection, subdermal injection and intramuscular injection, and the effective dose administered to a surface area of about 4 mm², or about 5 mm², or about 6 mm², or about 7 mm², or about 8 mm², or about 10 mm², or about 20 mm², or about 50 mm², or about 100 mm², or about 150 mm².

In various embodiments, the nucleic acid drug, including RNA comprising one or more non-canonical nucleotides, and/or formulations comprising the same, is administered locally, optionally by one or more of subcutaneous injection, intradermal injection, subdermal injection and intramuscular injection, and the effective dose (weight RNA / surface area of injection) is about 35 ng/cm² to about 7000 ng/cm². In various embodiments, the nucleic acid drug, including RNA comprising one or more non-canonical nucleotides, and/or formulations comprising the same, is administered locally, optionally by one or more of subcutaneous injection, intradermal injection, subdermal injection and intramuscular injection, and the effective dose (weight RNA / surface area of injection) is no more than about 35 ng/cm², or about 50 ng/cm², or about 75 ng/cm², or about 100 ng/cm², or about 125 ng/cm², or about 150 ng/cm², or about 175 ng/cm², or about 200 ng/cm², or about 225 ng/cm², or about 250 ng/cm², or about 500 ng/cm², or about 1000 ng/cm², or about 2000 ng/cm², or about 5000 ng/cm², or about 7000 ng/cm². In various embodiments, the nucleic acid drug, including RNA comprising one or more non-canonical nucleotides, and/or formulations comprising the same, is administered locally, optionally by one or more of subcutaneous injection, intradermal injection, subdermal injection and intramuscular injection, and the effective dose (weight RNA / surface area of injection) is about 35 ng/cm², or about 50 ng/cm², or about 75 ng/cm², or about 100 ng/cm², or about 125 ng/cm², or about 150 ng/cm², or about 175 ng/cm², or about 200 ng/cm², or about 225 ng/cm², or about 250 ng/cm², or about 500 ng/cm², or about 1000 ng/cm², or about 2000 ng/cm², or about 5000 ng/cm², or about 7000 ng/cm².

Pharmaceutical preparations may additionally comprise delivery reagents (a.k.a. "transfection reagents", a.k.a. "vehicles", a.k.a. "delivery vehicles") and/or excipients. Pharmaceutically acceptable delivery reagents, excipients, and methods of preparation and use thereof, including methods for preparing and administering pharmaceutical preparations to patients (a.k.a. "subjects") are well known in the art, and are set forth in numerous publications, including, for example, in US Patent Appl. Pub. No. US 2008/0213377, the entirety of which is incorporated herein by reference.

For example, the present compositions can be in the form of pharmaceutically acceptable salts. Such salts include those listed in, for example, *J. Pharma. Sci.* 66, 2-19 (1977) and *The Handbook of Pharmaceutical Salts; Properties, Selection, and Use*. P. H. Stahl and C. G. Wermuth (eds.), Verlag, Zurich (Switzerland) 2002, which are hereby incorporated by reference in their entirety. Non-limiting examples of pharmaceutically acceptable salts include: sulfate, citrate, acetate, oxalate, chloride, bromide, iodide, nitrate, bisulfate, phosphate, acid phosphate,

isonicotinate, lactate, salicylate, acid citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucaronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate, camphorsulfonate, pamoate, phenylacetate, trifluoroacetate, acrylate, chlorobenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, methylbenzoate, o-acetoxybenzoate, naphthalene-2-benzoate, isobutyrate, phenylbutyrate, α -hydroxybutyrate, butyne-1,4-dicarboxylate, hexyne-1,4-dicarboxylate, caprate, caprylate, cinnamate, glycollate, heptanoate, hippurate, malate, hydroxymaleate, malonate, mandelate, mesylate, nicotinate, phthalate, teraphthalate, propiolate, propionate, phenylpropionate, sebacate, suberate, p-bromobenzenesulfonate, chlorobenzenesulfonate, ethylsulfonate, 2-hydroxyethylsulfonate, methylsulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, naphthalene-1,5-sulfonate, xylenesulfonate, tartarate salts, hydroxides of alkali metals such as sodium, potassium, and lithium; hydroxides of alkaline earth metal such as calcium and magnesium; hydroxides of other metals, such as aluminum and zinc; ammonia, and organic amines, such as unsubstituted or hydroxy-substituted mono-, di-, or tri-alkylamines, dicyclohexylamine; tributyl amine; pyridine; N-methyl, N-ethylamine; diethylamine; triethylamine; mono-, bis-, or tris-(2-OH-lower alkylamines), such as mono-, bis-, or tris-(2-hydroxyethyl)amine, 2-hydroxy-tert-butylamine, or tris-(hydroxymethyl)methylamine, N,N-di-lower alkyl-N-(hydroxyl-lower alkyl)-amines, such as N,N-dimethyl-N-(2-hydroxyethyl)amine or tri-(2-hydroxyethyl)amine; N-methyl-D-glucamine; and amino acids such as arginine, lysine, and the like.

The present pharmaceutical compositions can comprises excipients, including liquids such as water and oils, including those of petroleum, animal, vegetable, or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. The pharmaceutical excipients can be, for example, saline, gum acacia, gelatin, starch paste, talc, keratin, colloidal silica, urea and the like. In addition, auxiliary, stabilizing, thickening, lubricating, and coloring agents can be used. In one embodiment, the pharmaceutically acceptable excipients are sterile when administered to a subject. Suitable pharmaceutical excipients also include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. Any agent described herein, if desired, can also comprise minor amounts of wetting or emulsifying agents, or pH buffering agents.

Dosage forms suitable for parenteral administration (e.g. subcutaneous, intradermal, subdermal, intramuscular, intravenous, intraperitoneal, intra-articular, and infusion) include, for example, solutions, suspensions, dispersions, emulsions, and the like. They may also be manufactured in the form of sterile solid compositions (e.g. lyophilized composition), which can be dissolved or suspended in sterile injectable medium immediately before use. They may contain, for example, suspending or dispersing agents known in the art.

In some embodiments, the formulations described herein may comprise albumin and a nucleic acid molecule.

In some embodiments, the invention relates to a cosmetic composition. In one embodiment, the cosmetic composition comprises albumin. In another embodiment, the albumin is treated with an ion-exchange resin or charcoal. In yet another embodiment, the cosmetic composition comprises a nucleic acid molecule. In a further

embodiment, the cosmetic composition comprises both albumin and a nucleic acid molecule. Still other embodiments are directed to a cosmetic treatment article comprising a cosmetic composition contained in a device configured to deliver the composition to a patient. Still other embodiments are directed to a device configured to deliver a cosmetic composition to a patient. In one embodiment, the nucleic acid molecule encodes a member of the group: elastin, collagen, tyrosinase, melanocortin 1 receptor, keratin, filaggren, an antibody, and hyaluronan synthase or a biologically active fragment, variant, analogue or family member thereof.

In some embodiments, the present invention provides treatment regimens. The inventors have discovered that the doses and administration described herein can produce a substantial protein expression effect quickly (e.g. in about 6, or about 12, or about 24, or about 36, or about 48 hours). Further, these effects can be sustained for about 7 days, or longer. In some embodiments, the present methods provide for administration of a nucleic acid drug, including RNA comprising one or more non-canonical nucleotides, about weekly to about once every 20 weeks.

In some embodiments, the nucleic acid drug, including RNA comprising one or more non-canonical nucleotides, is administered about weekly, for at least 2 weeks (e.g. 3, or 4, or 5, or 6, or 7, or 8, or 9, or 10 weeks). In some embodiments, the nucleic acid drug, including RNA comprising one or more non-canonical nucleotides, is administered about every other week for at least one month (e.g. 1, or 2, or 3, or 4, or 5, or 6, or 12 months). In some embodiments, the nucleic acid drug, including RNA comprising one or more non-canonical nucleotides, is administered monthly or about every other month. In some embodiments, the nucleic acid drug, including RNA comprising one or more non-canonical nucleotides, is administered for at least two months, or at least 4 months, or at least 6 months, or at least 9 months, or at least one year.

In some embodiments, the nucleic acid drug, including RNA comprising one or more non-canonical nucleotides, is administered about weekly, or about once every 2 weeks, or about once every 3 weeks, or about once every 4 weeks, or about once every 5 weeks, or about once every 6 weeks, or about once every 7 weeks, or about once every 8 weeks, or about once every 9 weeks, or about once every 10 weeks, or about once every 11 weeks, or about once every 12 weeks, or about once every 13 weeks, or about once every 14 weeks, or about once every 15 weeks, or about once every 20 weeks, or about once every 24 weeks.


In some embodiments, the nucleic acid drug, including RNA comprising one or more non-canonical nucleotides, is administered no more than about weekly, or about once every 2 weeks, or about once every 3 weeks, or about once every 4 weeks, or about once every 5 weeks, or about once every 6 weeks, or about once every 7 weeks, or about once every 8 weeks, or about once every 9 weeks, or about once every 10 weeks, or about once every 11 weeks, or about once every 12 weeks, or about once every 13 weeks, or about once every 14 weeks, or about once every 15 weeks, or about once every 20 weeks, or about 24 weeks.

Certain proteins have long half-lives, and can persist in tissues for several hours, days, weeks, months or years. It has now been discovered that certain methods of treating a patient can result in accumulation of one or more

proteins, including, for example, one or more beneficial proteins. Certain embodiments are therefore directed to a method for treating a patient comprising delivering to a patient in a series of doses a nucleic acid encoding one or more proteins. In one embodiment the nucleic acid comprises RNA comprising one or more non-canonical nucleotides. In another embodiment, a first dose is given at a first time-point. In yet another embodiment, a
5 second dose is given at a second time-point. In a further embodiment, the amount of at least one of the one or more proteins in the patient at the second time-point is greater than the amount of said protein at the first time-point. In a still further embodiment, the method results in the accumulation of said protein in the patient.

In various embodiments, the present invention relates to nucleic acid drugs, which, in various embodiments are RNA comprising one or more non-canonical nucleotides. Certain non-canonical nucleotides, when incorporated
10 into RNA molecules, can reduce the toxicity of the RNA molecules, in part, without wishing to be bound by theory, by interfering with binding of proteins that detect exogenous nucleic acids, for example, protein kinase R, Rlg-1 and the oligoadenylate synthetase family of proteins. Non-canonical nucleotides that have been reported to reduce the toxicity of RNA molecules when incorporated therein include: pseudouridine, 5-methyluridine, 2-thiouridine, 5-methylcytidine, N6-methyladenosine, and certain combinations thereof. However, the chemical
15 characteristics of non-canonical nucleotides that can enable them to lower the *in vivo* toxicity of RNA molecules have, until this point, remained unknown. Furthermore, incorporation of large amounts of most non-canonical nucleotides, for example, 5-methyluridine, 2-thiouridine, 5-methylcytidine, and N6-methyladenosine, can reduce the efficiency with which RNA molecules can be translated into protein, limiting the utility of RNA molecules containing these nucleotides in applications that require protein expression. In addition, while pseudouridine can
20 be completely substituted for uridine in RNA molecules without reducing the efficiency with which the synthetic RNA molecules can be translated into protein, in certain situations, for example, when performing frequent, repeated transfections, synthetic RNA molecules containing only adenosine, guanosine, cytidine, and pseudouridine can exhibit excessive toxicity.

It has now been discovered that, and in some embodiments the invention pertains to, RNA molecules containing
25 one or more non-canonical nucleotides that include one or more substitutions at the 2C and/or 4C and/or 5C positions in the case of a pyrimidine or the 6C and/or 7N and/or 8C positions in the case of a purine can be less toxic than synthetic RNA molecules containing only canonical nucleotides, due in part to the ability of substitutions at these positions to interfere with recognition of synthetic RNA molecules by proteins that detect exogenous nucleic acids, and furthermore, that substitutions at these positions can have minimal impact on the
30 efficiency with which the synthetic RNA molecules can be translated into protein, due in part to the lack of interference of substitutions at these positions with base-pairing and base-stacking interactions.



Examples of non-canonical nucleotides that include one or more substitutions at the 2C and/or 4C and/or 5C positions in the case of a pyrimidine or the 6C and/or 7N and/or 8C positions in the case of a purine include, but are not limited to: 2-thiouridine, 5-azauridine, pseudouridine, 4-thiouridine, 5-methyluridine, 5-methylpseudouridine, 5-aminouridine, 5-aminopseudouridine, 5-hydroxyuridine, 5-hydroxypseudouridine, 5-methoxyuridine, 5-methoxypseudouridine, 5-hydroxymethyluridine, 5-hydroxymethylpseudouridine, 5-carboxyuridine, 5-carboxypseudouridine, 5-formyluridine, 5-formylpseudouridine, 5-methyl-5-azauridine, 5-amino-5-azauridine, 5-hydroxy-5-azauridine, 5-methylpseudouridine, 5-aminopseudouridine, 5-hydroxypseudouridine, 4-thio-5-azauridine, 4-thiopseudouridine, 4-thio-5-methyluridine, 4-thio-5-aminouridine, 4-thio-5-hydroxyuridine, 4-thio-5-methyl-5-azauridine, 4-thio-5-amino-5-azauridine, 4-thio-5-hydroxy-5-azauridine, 4-thio-5-methylpseudouridine, 4-thio-5-aminopseudouridine, 4-thio-5-hydroxypseudouridine, 2-thiocytidine, 5-azacytidine, pseudoisocytidine, N4-methylcytidine, N4-aminocytidine, N4-hydroxycytidine, 5-methylcytidine, 5-aminocytidine, 5-hydroxycytidine, 5-methoxycytidine, 5-hydroxymethylcytidine, 5-carboxycytidine, 5-formylcytidine, 5-methyl-5-azacytidine, 5-amino-5-azacytidine, 5-hydroxy-5-azacytidine, 5-methylpseudoisocytidine, 5-aminopseudoisocytidine, 5-hydroxypseudoisocytidine, N4-methyl-5-azacytidine, N4-methylpseudoisocytidine, 2-thio-5-azacytidine, 2-thiopseudoisocytidine, 2-thio-N4-methylcytidine, 2-thio-N4-aminocytidine, 2-thio-N4-hydroxycytidine, 2-thio-5-methylcytidine, 2-thio-5-aminocytidine, 2-thio-5-hydroxycytidine, 2-thio-5-methyl-5-azacytidine, 2-thio-5-amino-5-azacytidine, 2-thio-5-hydroxy-5-azacytidine, 2-thio-5-methylpseudoisocytidine, 2-thio-5-aminopseudoisocytidine, 2-thio-5-hydroxypseudoisocytidine, 2-thio-N4-methyl-5-azacytidine, 2-thio-N4-methylpseudoisocytidine, N4-methyl-5-methylcytidine, N4-methyl-5-aminocytidine, N4-methyl-5-hydroxycytidine, N4-methyl-5-methyl-5-azacytidine, N4-methyl-5-amino-5-azacytidine, N4-methyl-5-hydroxy-5-azacytidine, N4-methyl-5-methylpseudoisocytidine, N4-methyl-5-aminopseudoisocytidine, N4-methyl-5-hydroxypseudoisocytidine, N4-amino-5-azacytidine, N4-aminopseudoisocytidine, N4-amino-5-methylcytidine, N4-amino-5-aminocytidine, N4-amino-5-hydroxycytidine, N4-amino-5-methyl-5-azacytidine, N4-amino-5-amino-5-azacytidine, N4-amino-5-hydroxy-5-azacytidine, N4-amino-5-methylpseudoisocytidine, N4-amino-5-aminopseudoisocytidine, N4-amino-5-hydroxypseudoisocytidine, N4-hydroxy-5-azacytidine, N4-hydroxypseudoisocytidine, N4-hydroxy-5-methylcytidine, N4-hydroxy-5-aminocytidine, N4-hydroxy-5-hydroxycytidine, N4-hydroxy-5-methyl-5-azacytidine, N4-hydroxy-5-amino-5-azacytidine, N4-hydroxy-5-hydroxy-5-azacytidine, N4-hydroxy-5-methylpseudoisocytidine, N4-hydroxy-5-aminopseudoisocytidine, N4-hydroxy-5-hydroxypseudoisocytidine, 2-thio-N4-methyl-5-methylcytidine, 2-thio-N4-

methyl-5-aminocytidine, 2-thio-N4-methyl-5-hydroxycytidine, 2-thio-N4-methyl-5-methyl-5-azacytidine, 2-thio-N4-methyl-5-amino-5-azacytidine, 2-thio-N4-methyl-5-hydroxy-5-azacytidine, 2-thio-N4-methyl-5-methylpseudoisocytidine, 2-thio-N4-methyl-5-aminopseudoisocytidine, 2-thio-N4-methyl-5-hydroxypseudoisocytidine, 2-thio-N4-amino-5-azacytidine, 2-thio-N4-aminopseudoisocytidine, 2-thio-N4-amino-5-methylcytidine, 2-thio-N4-amino-5-aminocytidine, 2-thio-N4-amino-5-hydroxycytidine, 2-thio-N4-amino-5-methyl-5-azacytidine, 2-thio-N4-amino-5-amino-5-azacytidine, 2-thio-N4-amino-5-hydroxy-5-azacytidine, 2-thio-N4-amino-5-methylpseudoisocytidine, 2-thio-N4-amino-5-aminopseudoisocytidine, 2-thio-N4-amino-5-hydroxypseudoisocytidine, 2-thio-N4-hydroxy-5-azacytidine, 2-thio-N4-hydroxypseudoisocytidine, 2-thio-N4-hydroxy-5-methylcytidine, N4-hydroxy-5-aminocytidine, 2-thio-N4-hydroxy-5-hydroxycytidine, 2-thio-N4-hydroxy-5-methyl-5-azacytidine, 2-thio-N4-hydroxy-5-amino-5-azacytidine, 2-thio-N4-hydroxy-5-hydroxy-5-azacytidine, 2-thio-N4-hydroxy-5-methylpseudoisocytidine, 2-thio-N4-hydroxy-5-aminopseudoisocytidine, 2-thio-N4-hydroxy-5-hydroxypseudoisocytidine, N6-methyladenosine, N6-aminoadenosine, N6-hydroxyadenosine, 7-deazaadenosine, 8-azaadenosine, N6-methyl-7-deazaadenosine, N6-methyl-8-azaadenosine, 7-deaza-8-azaadenosine, N6-methyl-7-deaza-8-azaadenosine, N6-amino-7-deazaadenosine, N6-amino-8-azaadenosine, N6-amino-7-deaza-8-azaadenosine, N6-hydroxyadenosine, N6-hydroxy-7-deazaadenosine, N6-hydroxy-8-azaadenosine, N6-hydroxy-7-deaza-8-azaadenosine, 6-thioguanosine, 7-deazaguanosine, 8-azaguanosine, 6-thio-7-deazaguanosine, 6-thio-8-azaguanosine, 7-deaza-8-azaguanosine, 6-thio-7-deaza-8-azaguanosine, and 5-methoxyuridine.

In some embodiments, the invention relates to one or more non-canonical nucleotides selected from 5-hydroxycytidine, 5-methylcytidine, 5-hydroxymethylcytidine, 5-carboxycytidine, 5-formylcytidine, 5-methoxycytidine, 5-hydroxyuridine, 5-hydroxymethyluridine, 5-carboxyuridine, 5-formyluridine, 5-methoxyuridine, pseudouridine, 5-hydroxypseudouridine, 5-methylpseudouridine, 5-hydroxymethylpseudouridine, 5-carboxypseudouridine, 5-formylpseudouridine, and 5-methoxypseudouridine. In some embodiments, at least 50%, or at least 55%, or at least 60%, or at least 65%, or at least 70%, or at least 75%, or at least 80%, or at least 85%, or at least 90%, or at least 95%, or 100% of the non-canonical nucleotides are one or more of 5-hydroxycytidine, 5-methylcytidine, 5-hydroxymethylcytidine, 5-carboxycytidine, 5-formylcytidine, 5-methoxycytidine, 5-hydroxyuridine, 5-methyluridine, 5-hydroxymethyluridine, 5-carboxyuridine, 5-formyluridine, 5-methoxyuridine, pseudouridine, 5-hydroxypseudouridine, 5-methylpseudouridine, 5-hydroxymethylpseudouridine, 5-carboxypseudouridine, 5-formylpseudouridine, and 5-methoxypseudouridine.

In some embodiments, at least about 50%, or at least about 55%, or at least 60%, or at least 65%, or at least 70%, or at least 75%, or at least 80%, or at least 85%, or at least 90%, or at least 95%, or 100% of cytidine residues are non-canonical nucleotides selected from 5-hydroxycytidine, 5-methylcytidine, 5-hydroxymethylcytidine, 5-carboxycytidine, 5-formylcytidine, 5-methoxycytidine.

In some embodiments, at least about 20%, or about 30%, or about 40%, or about 50%, or at least about 55%, or at least 60%, or at least 65%, or at least 70%, or at least 75%, or at least 80%, or at least 85%, or at least 90%,

or at least 95%, or 100% of uridine residues are non-canonical nucleotides selected from 5-hydroxyuridine, 5-methyluridine, 5-hydroxymethyluridine, 5-carboxyuridine, 5-formyluridine, 5-methoxyuridine, pseudouridine, 5-hydroxypseudouridine, 5-methylpseudouridine, 5-hydroxymethylpseudouridine, 5-carboxypseudouridine, 5-formylpseudouridine, and 5-methoxypseudouridine.

- 5 In some embodiments, at least about 10% (e.g. 10%, or about 20%, or about 30%, or about 40%, or about 50%) of guanosine residues are non-canonical nucleotides, and the non-canonical nucleotide is optionally 7-deazaguanosine. In some embodiments, the RNA contains no more than about 50% 7-deazaguanosine in place of guanosine residues.

In some embodiments, the RNA does not contain non-canonical nucleotides in place of adenosine residues.

- 10 Note that alternative naming schemes exist for certain non-canonical nucleotides. For example, in certain situations, 5-methylpseudouridine can be referred to as “3-methylpseudouridine” or “N3-methylpseudouridine” or “1-methylpseudouridine” or “N1-methylpseudouridine”.

- Nucleotides that contain the prefix “amino” can refer to any nucleotide that contains a nitrogen atom bound to the atom at the stated position of the nucleotide, for example, 5-aminocytidine can refer to 5-aminocytidine, 5-methylaminocytidine, and 5-nitrocytidine. Similarly, nucleotides that contain the prefix “methyl” can refer to any nucleotide that contains a carbon atom bound to the atom at the stated position of the nucleotide, for example, 5-methylcytidine can refer to 5-methylcytidine, 5-ethylcytidine, and 5-hydroxymethylcytidine, nucleotides that contain the prefix “thio” can refer to any nucleotide that contains a sulfur atom bound to the atom at the given position of the nucleotide, and nucleotides that contain the prefix “hydroxy” can refer to any nucleotide that contains an oxygen atom bound to the atom at the given position of the nucleotide, for example, 5-hydroxyuridine can refer to 5-hydroxyuridine and uridine with a methyl group bound to an oxygen atom, wherein the oxygen atom is bound to the atom at the 5C position of the uridine.
- 15
- 20

- Certain embodiments are therefore directed to RNA comprising one or more non-canonical nucleotides, wherein the RNA molecule contains one or more nucleotides that includes one or more substitutions at the 2C and/or 4C and/or 5C positions in the case of a pyrimidine or the 6C and/or 7N and/or 8C positions in the case of a purine. Other embodiments are directed to a therapeutic, wherein the therapeutic contains one or more RNA molecules comprising one or more non-canonical nucleotides, and wherein the one or more RNA molecules comprising one or more non-canonical nucleotides contains one or more nucleotides that includes one or more substitutions at the 2C and/or 4C and/or 5C positions in the case of a pyrimidine or the 6C and/or 7N and/or 8C positions in the case of a purine. In one embodiment, the therapeutic comprises a transfection reagent. In another embodiment, the transfection reagent comprises a cationic lipid, liposome or micelle. In still another embodiment, the liposome or micelle comprises folate and the therapeutic composition has anti-cancer activity. In another embodiment, the one or more nucleotides includes at least one of pseudouridine, 2-thiouridine, 4-thiouridine, 5-azauridine, 5-hydroxyuridine, 5-methyluridine, 5-aminouridine, 2-thiopseudouridine, 4-thiopseudouridine, 5-
- 25
- 30

hydroxypseudouridine, 5-methylpseudouridine, 5-aminopseudouridine, pseudoisocytidine, N4-methylcytidine, 2-thiocytidine, 5-azacytidine, 5-hydroxycytidine, 5-aminocytidine, 5-methylcytidine, N4-methylpseudoisocytidine, 2-thiopseudoisocytidine, 5-hydroxypseudoisocytidine, 5-aminopseudoisocytidine, 5-methylpseudoisocytidine, 7-deazaadenosine, 7-deazaguanosine, 6-thioguanosine, and 6-thio-7-deazaguanosine. In another embodiment, the one or more nucleotides includes at least one of pseudouridine, 2-thiouridine, 4-thiouridine, 5-azauridine, 5-hydroxyuridine, 5-methyluridine, 5-aminouridine, 2-thiopseudouridine, 4-thiopseudouridine, 5-hydroxypseudouridine, 5-methylpseudouridine, and 5-aminopseudouridine and at least one of pseudoisocytidine, N4-methylcytidine, 2-thiocytidine, 5-azacytidine, 5-hydroxycytidine, 5-aminocytidine, 5-methylcytidine, N4-methylpseudoisocytidine, 2-thiopseudoisocytidine, 5-hydroxypseudoisocytidine, 5-aminopseudoisocytidine, and 5-methylpseudoisocytidine. In still another embodiment, the one or more nucleotides include at least one of pseudouridine, 2-thiouridine, 4-thiouridine, 5-azauridine, 5-hydroxyuridine, 5-methyluridine, 5-aminouridine, 2-thiopseudouridine, 4-thiopseudouridine, 5-hydroxypseudouridine, and 5-methylpseudouridine, 5-aminopseudouridine and at least one of pseudoisocytidine, N4-methylcytidine, 2-thiocytidine, 5-azacytidine, 5-hydroxycytidine, 5-aminocytidine, 5-methylcytidine, N4-methylpseudoisocytidine, 2-thiopseudoisocytidine, 5-hydroxypseudoisocytidine, 5-aminopseudoisocytidine, and 5-methylpseudoisocytidine and at least one of 7-deazaguanosine, 6-thioguanosine, 6-thio-7-deazaguanosine, and 5-methoxyuridine. In yet another embodiment, the one or more nucleotides includes: 5-methylcytidine and 7-deazaguanosine. In another embodiment, the one or more nucleotides also includes pseudouridine or 4-thiouridine or 5-methyluridine or 5-aminouridine or 4-thiopseudouridine or 5-methylpseudouridine or 5-aminopseudouridine. In a still another embodiment, the one or more nucleotides also includes 7-deazaadenosine. In another embodiment, the one or more nucleotides includes: pseudoisocytidine and 7-deazaguanosine and 4-thiouridine. In yet another embodiment, the one or more nucleotides includes: pseudoisocytidine or 7-deazaguanosine and pseudouridine. In still another embodiment, the one or more nucleotides includes: 5-methyluridine and 5-methylcytidine and 7-deazaguanosine. In a further embodiment, the one or more nucleotides includes: pseudouridine or 5-methylpseudouridine and 5-methylcytidine and 7-deazaguanosine. In another embodiment, the one or more nucleotides includes: pseudoisocytidine and 7-deazaguanosine and pseudouridine. In one embodiment, the RNA comprising one or more non-canonical nucleotides is present *in vivo*.

Certain non-canonical nucleotides can be incorporated more efficiently than other non-canonical nucleotides into RNA molecules by RNA polymerases that are commonly used for *in vitro* transcription, due in part to the tendency of these certain non-canonical nucleotides to participate in standard base-pairing interactions and base-stacking interactions, and to interact with the RNA polymerase in a manner similar to that in which the corresponding canonical nucleotide interacts with the RNA polymerase. As a result, certain nucleotide mixtures containing one or more non-canonical nucleotides can be beneficial in part because *in vitro*-transcription reactions containing these nucleotide mixtures can yield a large quantity of RNA. Certain embodiments are therefore directed to a nucleotide mixture containing one or more nucleotides that includes one or more substitutions at the 2C and/or 4C and/or 5C positions in the case of a pyrimidine or the 6C and/or 7N and/or 8C

positions in the case of a purine. Nucleotide mixtures include, but are not limited to (numbers preceding each nucleotide indicate an exemplary fraction of the non-canonical nucleotide triphosphate in an *in vitro*-transcription reaction, for example, 0.2 pseudoisocytidine refers to a reaction containing adenosine-5'-triphosphate, guanosine-5'-triphosphate, uridine-5'-triphosphate, cytidine-5'-triphosphate, and pseudoisocytidine-5'-triphosphate, wherein pseudoisocytidine-5'-triphosphate is present in the reaction at an amount approximately equal to 0.2 times the total amount of pseudoisocytidine-5'-triphosphate + cytidine-5'-triphosphate that is present in the reaction, with amounts measured either on a molar or mass basis, and wherein more than one number preceding a nucleoside indicates a range of exemplary fractions): 1.0 pseudouridine, 0.1 – 0.8 2-thiouridine, 0.1 – 0.8 5-methyluridine, 0.2 – 1.0 5-hydroxyuridine, 0.2 – 1.0 5-methoxyuridine, 0.1 – 1.0 5-aminouridine, 0.1 – 1.0 4-thiouridine, 0.1 – 1.0 2-thiopseudouridine, 0.1 – 1.0 4-thiopseudouridine, 0.1 – 1.0 5-hydroxypseudouridine, 0.2 – 1.0 5-methylpseudouridine, 0.2 – 1.0 5-methoxypseudouridine, 0.1 – 1.0 5-aminopseudouridine, 0.2 – 1.0 2-thiocytidine, 0.1 – 0.8 pseudoisocytidine, 0.2 – 1.0 5-methylcytidine, 0.2 – 1.0 5-hydroxycytidine, 0.2 – 1.0 5-hydroxymethylcytidine, 0.2 – 1.0 5-methoxycytidine, 0.1 – 1.0 5-aminocytidine, 0.2 – 1.0 N4-methylcytidine, 0.2 – 1.0 5-methylpseudoisocytidine, 0.2 – 1.0 5-hydroxypseudoisocytidine, 0.2 – 1.0 5-aminopseudoisocytidine, 0.2 – 1.0 N4-methylpseudoisocytidine, 0.2 – 1.0 2-thiopseudoisocytidine, 0.2 – 1.0 7-deazaguanosine, 0.2 – 1.0 6-thioguanosine, 0.2 – 1.0 6-thio-7-deazaguanosine, 0.2 – 1.0 8-azaguanosine, 0.2 – 1.0 7-deaza-8-azaguanosine, 0.2 – 1.0 6-thio-8-azaguanosine, 0.1 – 0.5 7-deazaadenosine, and 0.1 – 0.5 N6-methyladenosine.

In various embodiments, the RNA comprising one or more non-canonical nucleotides composition or synthetic polynucleotide composition (*e.g.*, which may be prepared by *in vitro* transcription) contains substantially or entirely the canonical nucleotide at positions having adenine or “A” in the genetic code. The term “substantially” in this context refers to at least 90%. In these embodiments, the RNA composition or synthetic polynucleotide composition may further contain (*e.g.*, consist of) 7-deazaguanosine at positions with “G” in the genetic code as well as the corresponding canonical nucleotide “G”, and the canonical and non-canonical nucleotide at positions with G may be in the range of 5:1 to 1:5, or in some embodiments in the range of 2:1 to 1:2. In these embodiments, the RNA composition or synthetic polynucleotide composition may further contain (*e.g.*, consist of) one or more (*e.g.*, two, three or four) of 5-hydroxycytidine, 5-methylcytidine, 5-hydroxymethylcytidine, 5-carboxycytidine, 5-formylcytidine, 5-methoxycytidine at positions with “C” in the genetic code as well as the canonical nucleotide “C”, and the canonical and non-canonical nucleotide at positions with C may be in the range of 5:1 to 1:5, or in some embodiments in the range of 2:1 to 1:2. In some embodiments, the level of non-canonical nucleotide at positions of “C” are as described in the preceding paragraph. In these embodiments, the RNA composition or synthetic polynucleotide composition may further contain (*e.g.*, consist of) one or more (*e.g.*, two, three, or four) of 5-hydroxyuridine, 5-methyluridine, 5-hydroxymethyluridine, 5-carboxyuridine, 5-formyluridine, 5-methoxyuridine, pseudouridine, 5-hydroxypseudouridine, 5-methylpseudouridine, 5-hydroxymethylpseudouridine, 5-carboxypseudouridine, 5-formylpseudouridine, and 5-methoxypseudouridine at positions with “U” in the genetic code as well as the canonical nucleotide “U”, and the canonical and non-canonical nucleotide at positions with “U” may be in the range of 5:1 to 1:5, or in some embodiments in the range

of 2:1 to 1:2. In some embodiments, the level of non-canonical nucleotide at positions of “U” are as described in the preceding paragraph.

It has now been discovered that combining certain non-canonical nucleotides can be beneficial in part because the contribution of non-canonical nucleotides to lowering the toxicity of RNA molecules can be additive. Certain
 5 embodiments are therefore directed to a nucleotide mixture, wherein the nucleotide mixture contains more than one of the non-canonical nucleotides listed above, for example, the nucleotide mixture contains both pseudoisocytidine and 7-deazaguanosine or the nucleotide mixture contains both N4-methylcytidine and 7-deazaguanosine, *etc.* In one embodiment, the nucleotide mixture contains more than one of the non-canonical nucleotides listed above, and each of the non-canonical nucleotides is present in the mixture at the fraction listed
 10 above, for example, the nucleotide mixture contains 0.1 – 0.8 pseudoisocytidine and 0.2 – 1.0 7-deazaguanosine or the nucleotide mixture contains 0.2 – 1.0 N4-methylcytidine and 0.2 – 1.0 7-deazaguanosine, *etc.*

In certain situations, for example, when it may not be necessary or desirable to maximize the yield of an *in vitro*-transcription reaction, nucleotide fractions other than those given above may be used. The exemplary fractions and ranges of fractions listed above relate to nucleotide-triphosphate solutions of typical purity (greater than 90%
 15 purity). Larger fractions of these and other nucleotides can be used by using nucleotide-triphosphate solutions of greater purity, for example, greater than about 95% purity or greater than about 98% purity or greater than about 99% purity or greater than about 99.5% purity, which can be achieved, for example, by purifying the nucleotide triphosphate solution using existing chemical-purification technologies such as high-pressure liquid chromatography (HPLC) or by other means. In one embodiment, nucleotides with multiple isomers are purified to
 20 enrich the desired isomer.

Other embodiments are directed to a method for inducing a cell *in vivo* to express a protein of interest by contacting the cell with a RNA molecule that contains one or more non-canonical nucleotides that includes one or more substitutions at the 2C and/or 4C and/or 5C positions in the case of a pyrimidine or the 6C and/or 7N and/or 8C positions in the case of a purine. Still other embodiments are directed to a method for transfecting,
 25 reprogramming, and/or gene-editing a cell *in vivo* by contacting the cell with a RNA molecule that contains one or more non-canonical nucleotides that includes one or more substitutions at the 2C and/or 4C and/or 5C positions in the case of a pyrimidine or the 6C and/or 7N and/or 8C positions in the case of a purine. In one embodiment, the RNA molecule is produced by *in vitro* transcription. In one embodiment, the RNA molecule encodes one or more reprogramming factors. In another embodiment, the one or more reprogramming factors includes Oct4
 30 protein. In another embodiment, the cell is also contacted with a RNA molecule that encodes Sox2 protein. In yet another embodiment, the cell is also contacted with a RNA molecule that encodes Klf4 protein. In yet another embodiment, the cell is also contacted with a RNA molecule that encodes c-Myc protein. In yet another embodiment, the cell is also contacted with a RNA molecule that encodes Lin28 protein.

Enzymes such as T7 RNA polymerase may preferentially incorporate canonical nucleotides in an *in vitro*-transcription reaction containing both canonical and non-canonical nucleotides. As a result, an *in vitro*-
 35

transcription reaction containing a certain fraction of a non-canonical nucleotide may yield RNA containing a different, often lower, fraction of the non-canonical nucleotide than the fraction at which the non-canonical nucleotide was present in the reaction. In certain embodiments, references to nucleotide incorporation fractions (for example, "a synthetic RNA molecule containing 50% pseudoisocytidine" or "0.1 – 0.8 pseudoisocytidine")
 5 therefore can refer both to RNA molecules containing the stated fraction of the nucleotide, and to RNA molecules synthesized in a reaction containing the stated fraction of the nucleotide (or nucleotide derivative, for example, nucleotide-triphosphate), even though such a reaction may yield RNA containing a different fraction of the nucleotide than the fraction at which the non-canonical nucleotide was present in the reaction.

Different nucleotide sequences can encode the same protein by utilizing alternative codons. In certain
 10 embodiments, references to nucleotide incorporation fractions therefore can refer both to RNA molecules containing the stated fraction of the nucleotide, and to RNA molecules encoding the same protein as a different RNA molecule, wherein the different RNA molecule contains the stated fraction of the nucleotide.


Certain embodiments are directed to a kit containing one or more materials needed to practice the present invention. In one embodiment, the kit contains one or more synthetic RNA molecules. In one embodiment, the kit
 15 contains one or more synthetic RNA molecules that encode one or more reprogramming factors and/or gene-editing proteins. In another embodiment, the one or more synthetic RNA molecules contain one or more non-canonical nucleotides that include one or more substitutions at the 2C and/or 4C and/or 5C positions in the case of a pyrimidine or the 6C and/or 7N and/or 8C positions in the case of a purine. In another embodiment, the kit contains one or more of: a transfection medium, a transfection reagent, a complexation medium, and a matrix
 20 solution. In one embodiment, the matrix solution contains fibronectin and/or vitronectin or recombinant fibronectin and/or recombinant vitronectin. In one embodiment, one or more of the components of the kit are present as a plurality of aliquots. In one embodiment, the kit contains aliquots of nucleic acid transfection-reagent complexes. In another embodiment, the kit contains aliquots of nucleic acid transfection-reagent complexes that are provided in a solid form, for example, as frozen or freeze-dried pellets. In yet another embodiment, the kit contains aliquots
 25 of medium, wherein each aliquot contains transfection reagent-nucleic acid complexes that are stabilized either by chemical treatment or by freezing.

Transfection, in general, and reprogramming, in particular, can be difficult and time-consuming techniques that can be repetitive and prone to error. However, these techniques are often performed manually due to the lack of automated transfection equipment. Certain embodiments are therefore directed to a system that can transfect,
 30 reprogram, and/or gene-edit cells *in vivo* in an automated or semi-automated manner.

It has now been discovered that the non-canonical nucleotide members of the 5-methylcytidine de-methylation pathway, when incorporated into synthetic RNA, can increase the efficiency with which the synthetic RNA can be translated into protein *in vivo*, and can decrease the toxicity of the synthetic RNA *in vivo*. These non-canonical nucleotides include, for example: 5-methylcytidine, 5-hydroxymethylcytidine, 5-formylcytidine, and 5-
 35 carboxycytidine (a.k.a. "cytidine-5-carboxylic acid"). Certain embodiments are therefore directed to a nucleic acid.

In some embodiments, the nucleic acid is present *in vivo*. In one embodiment, the nucleic acid is a synthetic RNA molecule. In another embodiment, the nucleic acid comprises one or more non-canonical nucleotides. In one embodiment, the nucleic acid comprises one or more non-canonical nucleotide members of the 5-methylcytidine de-methylation pathway. In another embodiment, the nucleic acid comprises at least one of: 5-methylcytidine, 5-hydroxymethylcytidine, 5-formylcytidine, and 5-carboxycytidine or a derivative thereof. In a further embodiment, the nucleic acid comprises at least one of: pseudouridine, 5-methylpseudouridine, 5-hydroxyuridine, 5-methyluridine, 5-methylcytidine, 5-hydroxymethylcytidine, N4-methylcytidine, N4-acetylcytidine, and 7-deazaguanosine or a derivative thereof.

5-methylcytidine De-Methylation Pathway



Certain embodiments are directed to a protein. Other embodiments are directed to a nucleic acid that encodes a protein. In one embodiment, the protein is a protein of interest. In another embodiment, the protein is selected from: a reprogramming protein and a gene-editing protein. In one embodiment, the nucleic acid is a plasmid. In another embodiment, the nucleic acid is present in a virus or viral vector. In a further embodiment, the virus or viral vector is replication incompetent. In a still further embodiment, the virus or viral vector is replication competent. In one embodiment, the virus or viral vector includes at least one of: an adenovirus, a retrovirus, a lentivirus, a herpes virus, an adeno-associated virus or a natural or engineered variant thereof, and an engineered virus.

It has also been discovered that certain combinations of non-canonical nucleotides can be particularly effective at increasing the efficiency with which synthetic RNA can be translated into protein *in vivo*, and decreasing the toxicity of synthetic RNA *in vivo*, for example, the combinations: 5-methyluridine and 5-methylcytidine, 5-hydroxyuridine and 5-methylcytidine, 5-hydroxyuridine and 5-hydroxymethylcytidine, 5-methyluridine and 7-deazaguanosine, 5-methylcytidine and 7-deazaguanosine, 5-methyluridine, 5-methylcytidine, and 7-deazaguanosine, and 5-methyluridine, 5-hydroxymethylcytidine, and 7-deazaguanosine. Certain embodiments are therefore directed to a nucleic acid comprising at least two of: 5-methyluridine, 5-methylcytidine, 5-

hydroxymethylcytidine, and 7-deazaguanosine or one or more derivatives thereof. Other embodiments are directed to a nucleic acid comprising at least three of: 5-methyluridine, 5-methylcytidine, 5-hydroxymethylcytidine, and 7-deazaguanosine or one or more derivatives thereof. Other embodiments are directed to a nucleic acid comprising all of: 5-methyluridine, 5-methylcytidine, 5-hydroxymethylcytidine, and 7-deazaguanosine or one or more derivatives thereof. In one embodiment, the nucleic acid comprises one or more 5-methyluridine residues, one or more 5-methylcytidine residues, and one or more 7-deazaguanosine residues or one or more 5-methyluridine residues, one or more 5-hydroxymethylcytidine residues, and one or more 7-deazaguanosine residues.

It has been further discovered that synthetic RNA molecules containing certain fractions of certain non-canonical nucleotides and combinations thereof can exhibit particularly high translation efficiency and low toxicity *in vivo*. Certain embodiments are therefore directed to a nucleic acid comprising at least one of: one or more uridine residues, one or more cytidine residues, and one or more guanosine residues, and comprising one or more non-canonical nucleotides. In one embodiment, between about 20% and about 80% of the uridine residues are 5-methyluridine residues. In another embodiment, between about 30% and about 50% of the uridine residues are 5-methyluridine residues. In a further embodiment, about 40% of the uridine residues are 5-methyluridine residues. In one embodiment, between about 60% and about 80% of the cytidine residues are 5-methylcytidine residues. In another embodiment, between about 80% and about 100% of the cytidine residues are 5-methylcytidine residues. In a further embodiment, about 100% of the cytidine residues are 5-methylcytidine residues. In a still further embodiment, between about 20% and about 100% of the cytidine residues are 5-hydroxymethylcytidine residues. In one embodiment, between about 20% and about 80% of the guanosine residues are 7-deazaguanosine residues. In another embodiment, between about 40% and about 60% of the guanosine residues are 7-deazaguanosine residues. In a further embodiment, about 50% of the guanosine residues are 7-deazaguanosine residues. In one embodiment, between about 20% and about 80% or between about 30% and about 60% or about 40% of the cytidine residues are N4-methylcytidine and/or N4-acetylcytidine residues. In another embodiment, each cytidine residue is a 5-methylcytidine residue. In a further embodiment, about 100% of the cytidine residues are 5-methylcytidine residues and/or 5-hydroxymethylcytidine residues and/or N4-methylcytidine residues and/or N4-acetylcytidine residues and/or one or more derivatives thereof. In a still further embodiment, about 40% of the uridine residues are 5-methyluridine residues, between about 20% and about 100% of the cytidine residues are N4-methylcytidine and/or N4-acetylcytidine residues, and about 50% of the guanosine residues are 7-deazaguanosine residues. In one embodiment, about 40% of the uridine residues are 5-methyluridine residues and about 100% of the cytidine residues are 5-methylcytidine residues. In another embodiment, about 40% of the uridine residues are 5-methyluridine residues and about 50% of the guanosine residues are 7-deazaguanosine residues. In a further embodiment, about 100% of the cytidine residues are 5-methylcytidine residues and about 50% of the guanosine residues are 7-deazaguanosine residues. In a further embodiment, about 100% of the uridine residues are 5-hydroxyuridine residues. In one embodiment, about 40% of the uridine residues are 5-methyluridine residues, about 100% of the cytidine

residues are 5-methylcytidine residues, and about 50% of the guanosine residues are 7-deazaguanosine residues. In another embodiment, about 40% of the uridine residues are 5-methyluridine residues, between about 20% and about 100% of the cytidine residues are 5-hydroxymethylcytidine residues, and about 50% of the guanosine residues are 7-deazaguanosine residues. In some embodiments, less than 100% of the cytidine residues are 5-methylcytidine residues. In other embodiments, less than 100% of the cytidine residues are 5-hydroxymethylcytidine residues. In one embodiment, each uridine residue in the synthetic RNA molecule is a pseudouridine residue or a 5-methylpseudouridine residue. In another embodiment, about 100% of the uridine residues are pseudouridine residues and/or 5-methylpseudouridine residues. In a further embodiment, about 100% of the uridine residues are pseudouridine residues and/or 5-methylpseudouridine residues, about 100% of the cytidine residues are 5-methylcytidine residues, and about 50% of the guanosine residues are 7-deazaguanosine residues.

Other non-canonical nucleotides that can be used in place of or in combination with 5-methyluridine include, but are not limited to: pseudouridine, 5-hydroxyuridine, 5-hydroxypseudouridine, 5-methoxyuridine, 5-methoxypseudouridine, 5-carboxyuridine, 5-carboxypseudouridine, 5-formyluridine, 5-formylpseudouridine, 5-hydroxymethyluridine, 5-hydroxymethylpseudouridine, and 5-methylpseudouridine (a.k.a. "1-methylpseudouridine", a.k.a. "N1-methylpseudouridine") or one or more derivatives thereof. Other non-canonical nucleotides that can be used in place of or in combination with 5-methylcytidine and/or 5-hydroxymethylcytidine include, but are not limited to: pseudoisocytidine, 5-methylpseudoisocytidine, 5-hydroxymethylcytidine, 5-formylcytidine, 5-carboxycytidine, 5-methoxycytidine, N4-methylcytidine, N4-acetylcytidine or one or more derivatives thereof. In certain embodiments, for example, when performing only a single transfection, injection or delivery or when the cells, tissue, organ or patient being transfected, injected or delivered to are not particularly sensitive to transfection-associated toxicity or innate-immune signaling, the fractions of non-canonical nucleotides can be reduced. Reducing the fraction of non-canonical nucleotides can be beneficial, in part, because reducing the fraction of non-canonical nucleotides can reduce the cost of the nucleic acid. In certain situations, for example, when minimal immunogenicity of the nucleic acid is desired, the fractions of non-canonical nucleotides can be increased.

Enzymes such as T7 RNA polymerase may preferentially incorporate canonical nucleotides in an *in vitro*-transcription reaction containing both canonical and non-canonical nucleotides. As a result, an *in vitro*-transcription reaction containing a certain fraction of a non-canonical nucleotide may yield RNA containing a different, often lower, fraction of the non-canonical nucleotide than the fraction at which the non-canonical nucleotide was present in the reaction. In certain embodiments, references to nucleotide incorporation fractions (for example, "50% 5-methyluridine") therefore can refer both to nucleic acids containing the stated fraction of the nucleotide, and to nucleic acids synthesized in a reaction containing the stated fraction of the nucleotide (or nucleotide derivative, for example, nucleotide-triphosphate), even though such a reaction may yield a nucleic acid containing a different fraction of the nucleotide than the fraction at which the non-canonical nucleotide was

present in the reaction. In addition, different nucleotide sequences can encode the same protein by utilizing alternative codons. In certain embodiments, references to nucleotide incorporation fractions therefore can refer both to nucleic acids containing the stated fraction of the nucleotide, and to nucleic acids encoding the same protein as a different nucleic acid, wherein the different nucleic acid contains the stated fraction of the nucleotide.

- 5 Certain embodiments are directed to a nucleic acid comprising a 5'-cap structure selected from Cap 0, Cap 1, Cap 2, and Cap 3 or a derivative thereof. In one embodiment, the nucleic acid comprises one or more UTRs. In another embodiment, the one or more UTRs increase the stability of the nucleic acid. In a further embodiment, the one or more UTRs comprise an alpha-globin or beta-globin 5'-UTR. In a still further embodiment, the one or more UTRs comprise an alpha-globin or beta-globin 3'-UTR. In a still further embodiment, the synthetic RNA
- 10 molecule comprises an alpha-globin or beta-globin 5'-UTR and an alpha-globin or beta-globin 3'-UTR. In one embodiment, the 5'-UTR comprises a Kozak sequence that is substantially similar to the Kozak consensus sequence. In another embodiment, the nucleic acid comprises a 3'-poly(A) tail. In a further embodiment, the 3'-poly(A) tail is between about 20 nt and about 250 nt or between about 120 nt and about 150 nt long. In a further embodiment, the 3'-poly(A) tail is about 20 nt, or about 30 nt, or about 40 nt, or about 50 nt, or about 60 nt, or
- 15 about 70 nt, or about 80 nt, or about 90 nt, or about 100 nt, or about 110 nt, or about 120 nt, or about 130 nt, or about 140 nt, or about 150 nt, or about 160 nt, or about 170 nt, or about 180 nt, or about 190 nt, or about 200 nt, or about 210 nt, or about 220 nt, or about 230 nt, or about 240 nt, or about 250 nt long.

Certain embodiments are directed to methods of making nucleic acid drugs, including RNA comprising one or more non-canonical nucleotides. Such methods yield substantially stable RNA.

- 20 In various embodiments, the present methods and compositions find use in methods of treating, preventing or ameliorating a disease, disorder and/or condition. For instance, in some embodiments, the described methods of in vivo delivery, including various effective doses, administration strategies and formulations are used in a method of treatment.

- In various embodiments, the present methods and compositions find use in methods of altering, modifying and/or
- 25 changing a tissue (e.g. cosmetically).

In various embodiments, the present methods and compositions include using a nucleic acid drug, including a synthetic RNA, in the diagnosing, treating, preventing or ameliorating of a disease, disorder and/or condition described herein. In various embodiments, the present methods and compositions include using a nucleic acid drug, including a synthetic RNA, in the altering, modifying and/or changing of a tissue (e.g. cosmetically).

- 30 Generally speaking, in various embodiments, a synthetic RNA as described herein is administered to a human at specific doses described herein and the synthetic RNA comprises a sequence, sometimes referred to as a target sequence that encodes a protein of interest, which may be a therapeutic protein.

Synthetic RNA comprising only canonical nucleotides can bind to pattern recognition receptors, can be recognized as a pathogen-associated molecular pattern, and can trigger a potent immune response in cells,

which can result in translation block, the secretion of inflammatory cytokines, and cell death. It has now been discovered that synthetic RNA comprising certain non-canonical nucleotides can evade detection by the innate immune system, and can be translated at high efficiency into protein, including in humans. It has been further discovered that synthetic RNA comprising at least one of the non-canonical nucleotides described herein, including, for example, a member of the group: 5-methylcytidine, 5-hydroxycytidine, 5-hydroxymethylcytidine, 5-carboxycytidine, 5-formylcytidine, 5-methoxycytidine, pseudouridine, 5-hydroxyuridine, 5-methyluridine, 5-hydroxymethyluridine, 5-carboxyuridine, 5-methoxyuridine, 5-formyluridine, 5-hydroxypseudouridine, 5-methylpseudouridine, 5-hydroxymethylpseudouridine, 5-carboxypseudouridine, 5-methoxypseudouridine, and 5-formylpseudouridine can evade detection by the innate immune system, and can be translated at high efficiency into protein, including in humans. Certain embodiments are therefore directed to a method for inducing a cell to express a protein of interest comprising contacting a cell with synthetic RNA. Other embodiments are directed to a method for transfecting a cell with synthetic RNA comprising contacting a cell with a solution comprising one or more synthetic RNA molecules. Still other embodiments are directed to a method for treating a patient comprising administering to the patient synthetic RNA. In one embodiment, the synthetic RNA comprises at least one of the non-canonical nucleotides described herein, including, for example, a member of the group: 5-methylcytidine, 5-hydroxycytidine, 5-hydroxymethylcytidine, 5-carboxycytidine, 5-formylcytidine, 5-methoxycytidine, pseudouridine, 5-hydroxyuridine, 5-methyluridine, 5-hydroxymethyluridine, 5-carboxyuridine, 5-methoxyuridine, 5-formyluridine, 5-hydroxypseudouridine, 5-methylpseudouridine, 5-hydroxymethylpseudouridine, 5-carboxypseudouridine, 5-methoxypseudouridine, and 5-formylpseudouridine. In another embodiment, the synthetic RNA encodes a protein of interest. Exemplary RNAs may contain combinations and levels of non-canonical and non-canonical nucleotides as described elsewhere herein, including with respect to the expression of any protein of interest described herein. In yet another embodiment, the method results in the expression of the protein of interest. In a further embodiment, the method results in the expression of the protein of interest in the patient's skin.

Other embodiments are directed to a method for delivering a nucleic acid to a cell *in vivo*. Still other embodiments are directed to a method for inducing a cell *in vivo* to express a protein of interest. Still other embodiments are directed to a method for treating a patient. In one embodiment, the method comprises disrupting the stratum corneum. In another embodiment, the method comprises contacting a cell with a nucleic acid. In yet another embodiment, the method results in the cell internalizing the nucleic acid. In a further embodiment, the method results in the cell expressing the protein of interest. In a still further embodiment, the method results in the expression of the protein of interest in the patient. In a still further embodiment, the method results in the amelioration of one or more of the patient's symptoms. In a still further embodiment, the patient is in need of the protein of interest. In a still further embodiment, the patient is deficient in the protein of interest.

Still other embodiments are directed to a method for treating a patient comprising delivering to a patient a composition. In one embodiment, the composition comprises albumin that is treated with an ion-exchange resin

or charcoal. In another embodiment, the composition comprises one or more nucleic acid molecules. In yet another embodiment, at least one of the one or more nucleic acid molecules encodes a protein of interest. In one embodiment, the method results in the expression of the protein in the patient's skin. In another embodiment, the method results in the expression of a therapeutically or cosmetically effective amount of the protein of interest in the patient. In yet another embodiment, the method comprises administering a steroid. In a further embodiment, the steroid is a member of the group: hydrocortisone and dexamethasone.

Some embodiments are directed to a therapeutic composition and/or methods of treatment comprising a nucleic acid molecule encoding one or more proteins, wherein at least one of the one or more proteins is an extracellular matrix protein. Still other embodiments are directed to a cosmetic composition comprising a nucleic acid molecule encoding one or more proteins, wherein at least one of the one or more proteins is an extracellular matrix protein.

Pigmentation disorders can cause severe symptoms in patients. It has now been discovered that pigmentation disorders can be treated by delivering to a patient a nucleic acid encoding tyrosinase. Certain embodiments are therefore directed to a method for treating a pigmentation disorder. Other embodiments are directed to a method for altering the pigmentation of a patient. In one embodiment, the method comprises delivering to a patient a nucleic acid encoding tyrosinase. Other embodiments are directed to a cosmetic composition comprising a nucleic acid encoding tyrosinase. Still other embodiments are directed to a therapeutic composition comprising a nucleic acid encoding tyrosinase. Still other embodiments are directed to a method for increasing the ultraviolet absorption of a patient's skin. In one embodiment the method comprises delivering to a patient a nucleic acid encoding tyrosinase. In another embodiment, the method results in an increase in the ultraviolet absorption of the patient's skin. Still other embodiments are directed to a method for reducing photodamage to a person's skin upon exposure to ultraviolet light. In one embodiment, the method results in the reduction of photodamage to the person's skin upon exposure to ultraviolet light. Still other embodiments are directed to a method for treating xeroderma pigmentosum. In one embodiment, the method comprises delivering to a patient a nucleic acid encoding tyrosinase. Still other embodiments are directed to a method for treating epidermolysis bullosa. In one embodiment, the method comprises delivering to a patient a nucleic acid encoding one or more of keratin 5, keratin 14, plectin, an integrin family member, laminin, a laminin subunit, collagen XVII, collagen VII or a biologically active fragment, variant, analogue or family-member thereof. In one embodiment, the method comprises delivering to a patient a nucleic acid encoding collagen type VII. In another embodiment, the method comprises delivering to a patient a nucleic acid encoding melanocortin 1 receptor. Still other embodiments are directed to a method for treating xerosis. In one embodiment, the method comprises delivering to a patient a nucleic acid encoding a hyaluronan synthase. In another embodiment, the patient is diagnosed with atopic dermatitis. In yet another embodiment, the patient is diagnosed with ichthyosis. Certain embodiments are directed to a method for treating a cosmetic condition. Other embodiments are directed to a method for inducing tissue healing. In one embodiment, the method comprises delivering to a patient a nucleic acid encoding a

hyaluronan synthase. In another embodiment, the cosmetic condition is a member of the group: wrinkles, sagging skin, thin skin, discoloration, and dry skin. In yet another embodiment, the patient has had cataract surgery. In some embodiments, the nucleic acid is synthetic RNA. In other embodiments, the method results in the amelioration of one or more of the patient's symptoms. Other embodiments are directed to a method for

5 treating an indication by delivering to a cell or a patient a nucleic acid encoding a protein or a peptide. Still other embodiments are directed to a composition comprising a nucleic acid encoding a protein or a peptide. Indications that can be treated using the methods and compositions of the present invention and proteins and peptides that can be encoded by compositions of the present invention are set forth in **Table 2A** and/or **Table 2B**, and are given by way of example, and not by way of limitation. In one embodiment, the indication is selected from **Table**

10 **2A** and/or **Table 2B**. In another embodiment the protein or peptide is selected from **Table 2A** and/or **Table 2B**. In yet another embodiment, the indication and the protein or peptide are selected from the same row of **Table 2A** and/or **Table 2B**. In a further embodiment, the protein of interest is a member of the group: UCP1, UCP2, and UCP3. Other embodiments are directed to methods for inducing a cell to express a plurality of proteins of interest. In one embodiment, the proteins of interest include at least two members of the group: a lipase, UCP1,

15 UCP2, and UCP3. In another embodiment, the proteins of interest include a lipase and a member of the group: UCP1, UCP2, and UCP3. In another embodiment, the protein is a gene-editing protein. In yet another embodiment, the gene-editing protein targets a gene that is at least partly responsible for a disease phenotype. In yet another embodiment, the gene-editing protein targets a gene that encodes a protein selected from **Table 2A** and/or **Table 2B**. In still another embodiment, the gene-editing protein corrects or eliminates, either alone or

20 in combination with one or more other molecules or gene-editing proteins, a mutation that is at least partly responsible for a disease phenotype.

In various embodiments, the present invention contemplates the targeting of the precursor forms and/or mature forms and/or isoforms and/or mutants of any of the proteins disclosed in **Table 2A** and/or **Table 2B** and such proteins. In some embodiments, any of the precursor forms and/or mature forms and/or isoforms and/or mutants

25 have enhanced secretion relative to the corresponding wild type proteins. In some embodiments, any of the precursor forms and/or mature forms and/or isoforms and/or mutants have altered half-lives (e.g. serum, plasma, intracellular) - for instance, longer or shorter half-lives. In some embodiments, this is relative to wild type.

Table 2A. Illustrative Indications

Illustrative Indication	Illustrative Protein / Peptide
Acne	Retinol Dehydrogenase 10
	Elastin sp P15502 ELN_HUMAN Elastin (isoform 3) (SEQ ID NO:486)
Aging	Collagen Type I P02452 CO1A1_HUMAN Collagen alpha-1(I) chain (SEQ ID NO:487)
Aging	

	P08123 CO1A2_HUMAN Collagen alpha-2(I) chain (SEQ ID NO:488)
Aging	Collagen Type III P02461 CO3A1_HUMAN Collagen alpha-1(III) chain (isoform 1) (SEQ ID NO:489)
Aging	Collagen Type VII Q02388 CO7A1_HUMAN Collagen alpha-1(VII) chain (SEQ ID NO:490)
Aging	Hyaluronan Synthase
Aging	Telomerase Reverse Transcriptase
Albinism	Tyrosinase P14679 TYRO_HUMAN Tyrosinase (isoform 1) (SEQ ID NO:491)
	Collagen Type IV P02462 CO4A1_HUMAN Collagen alpha-1(IV) chain (isoform 1) (SEQ ID NO:492) P08572 CO4A2_HUMAN Collagen alpha-2(IV) chain (SEQ ID NO:493) Q01955 CO4A3_HUMAN Collagen alpha-3(IV) chain (isoform 1) (SEQ ID NO:494) P53420 CO4A4_HUMAN Collagen alpha-4(IV) chain (SEQ ID NO:495) P29400 CO4A5_HUMAN Collagen alpha-5(IV) chain (isoform 1) (SEQ ID NO:496) Q14031 CO4A6_HUMAN Collagen alpha-6(IV) (isoform A) (SEQ ID NO:497)
Alport Syndrome	
Anemia	Erythropoietin
Atopic Dermatitis	Filaggrin
	Elastin sp P15502 ELN_HUMAN Elastin (isoform 3) (SEQ ID NO:486)
Cutis Laxa	
Dry Skin	Filaggrin
	Collagen Type VII Q02388 CO7A1_HUMAN Collagen alpha-1(VII) chain (SEQ ID NO:498)
Dystrophic Epidermolysis Bullosa	

Ehlers-Danlos Syndrome	Collagen Type V P20908 CO5A1_HUMAN Collagen alpha-1(V) chain (SEQ ID NO:499) P05997 CO5A2_HUMAN Collagen alpha-2(V) chain (SEQ ID NO:500) P25940 CO5A3_HUMAN Collagen alpha-3(V) chain (SEQ ID NO:501)
Ehlers-Danlos Syndrome	Collagen Type I P02452 CO1A1_HUMAN Collagen alpha-1(I) chain (SEQ ID NO:487) P08123 CO1A2_HUMAN Collagen alpha-2(I) chain (SEQ ID NO:488)
Epidermolysis bullosa, lethal acantholytic	ADAM17 P78536 ADA17_HUMAN Disintegrin and metalloproteinase domain-containing protein 17 (isoform A) (SEQ ID NO:502)
Epidermolysis bullosa, type IV	Collagen Type III P02461 CO3A1_HUMAN Collagen alpha-1(III) chain (isoform 1) (SEQ ID NO:489)
Erythropoietic Protoporphyria	Ferrochelatase P22830 HEMH_HUMAN Ferrochelatase, mitochondrial (isoform 1) (SEQ ID NO:503)
Eczema	Filaggrin
Excess Fat	Thermogenin P25874 UCP1_HUMAN Mitochondrial brown fat uncoupling protein 1 (SEQ ID NO:504)
Excess Fat	Lipase Lipoprotein lipase P06858 LIPL_HUMAN Lipoprotein lipase (SEQ ID NO:516) Hepatic lipase P11150 LIPC_HUMAN Hepatic triacylglycerol lipase (SEQ ID NO:517) Pancreatic lipase P16233 LIPP_HUMAN Pancreatic triacylglycerol lipase (SEQ ID NO:518) Endothelial lipase (isoform 1)

	Q9Y5X9 LIPE_HUMAN Endothelial lipase (SEQ ID NO:519) Lysosomal lipase P38571 LICH_HUMAN Lysosomal acid lipase/cholesteryl ester hydrolase (isoform 1) (SEQ ID NO:520) Hormone sensitive lipase Q05469 LIPS_HUMAN Hormone-sensitive lipas (isoform 1) (SEQ ID NO:521) Gastric lipase P07098 LIPG_HUMAN Gastric triacylglycerol lipase (isoform 1) (SEQ ID NO:522) Pancreatic Lipase-Related Protein 1) P54315 LIPR1_HUMAN Inactive pancreatic lipase- related protein 1 (isoform 1) (SEQ ID NO:523) Pancreatic Lipase-Related Protein 2 P54317 LIPR2_HUMAN Pancreatic lipase-related protein 2 (SEQ ID NO:524) Carboxyl Ester Lipase P19835 CEL_HUMAN Bile salt-activated lipase (isoform long) (SEQ ID NO:525)
Hypotrichosis	ADAM17 P78536 ADA17_HUMAN Disintegrin and metalloproteinase domain-containing protein 17 (isoform A) (SEQ ID NO:502)
Ichthyosis Vulgaris	Filaggrin
Infections	Genetic Antibiotics (e.g. Anti-Sigma Factors)
Inflammatory and Bullous Skin Bowel Syndrome	Desmoglein 2 Q14126 DSG2_HUMAN Desmoglein-2 (SEQ ID NO:505)
Keratosis Pilaris	Retinol Dehydrogenase 10
Oily Skin	Retinol Dehydrogenase 10
Osteoarthritis	Hyaluronan Synthase
Pemphigus Vulgaris	Plakophilin-1 Q13835 PKP1_HUMAN Plakophilin-1 (isoform 2) (SEQ ID NO:506)

Pseudoxanthoma elasticum	Elastin sp P15502 ELN_HUMAN Elastin (isoform 3) (SEQ ID NO:486)
Psoriasis	Retinol Dehydrogenase 10
Scar Treatment	Tyrosinase P14679 TYRO_HUMAN Tyrosinase (isoform 1) (SEQ ID NO:491)
Scarring	Elastin sp P15502 ELN_HUMAN Elastin (isoform 3) (SEQ ID NO:486)
Scarring	Collagen Type I P02452 CO1A1_HUMAN Collagen alpha-1(I) chain (SEQ ID NO:487)
Scarring	P08123 CO1A2_HUMAN Collagen alpha-2(I) chain (SEQ ID NO:488)
Scarring	Collagen Type III P02461 CO3A1_HUMAN Collagen alpha-1(III) chain (isoform 1) (SEQ ID NO:489)
Skin Cancer	Interferon
	Interferon, Alpha 1 P01562 IFNA1_HUMAN Interferon alpha-1/13 (SEQ ID NO:530)
	Interferon, Alpha 2 P01563 IFNA2_HUMAN Interferon alpha-2 (SEQ ID NO:531)
	Interferon, Alpha 4 P05014 IFNA4_HUMAN Interferon alpha-4 (SEQ ID NO:532)
	Interferon, Alpha 5 P01569 IFNA5_HUMAN Interferon alpha-5 (SEQ ID NO:533)
	Interferon, Alpha 6 P05013 IFNA6_HUMAN Interferon alpha-6 (SEQ ID NO:534)
	Interferon, Alpha 7 P01567 IFNA7_HUMAN Interferon alpha-7 (SEQ ID NO:535)
	Interferon, Alpha 8 P32881 IFNA8_HUMAN Interferon alpha-8 (SEQ ID NO:536)
	Interferon, Alpha 10

	P01566 IFN10_HUMAN Interferon alpha-10 (SEQ ID NO:537) Interferon, Alpha 14 P01570 IFN14_HUMAN Interferon alpha-14 OS (SEQ ID NO:538) Interferon, Alpha 16 P05015 IFN16_HUMAN Interferon alpha-16 (SEQ ID NO:539) Interferon, Alpha 17 P01571 IFN17_HUMAN Interferon alpha-17 (SEQ ID NO:540) Interferon, Alpha 21 P01568 IFN21_HUMAN Interferon alpha-21 (SEQ ID NO:541) Interferon, Gamma P01579 IFNG_HUMAN Interferon gamma (SEQ ID NO:542) Interferon, Beta P01574 IFNB_HUMAN Interferon beta (SEQ ID NO:543) Interferon, Kappa Q9P0W0 IFNK_HUMAN Interferon kappa (SEQ ID NO:544) Interferon, Epsilon Q86WN2 IFNE_HUMAN Interferon epsilon (SEQ ID NO:545)
Striate Palmoplantar Keratoderma	ADAM17 P78536 ADA17_HUMAN Disintegrin and metalloproteinase domain-containing protein 17 (isoform A) (SEQ ID NO:502)
Tanning	Tyrosinase P14679 TYRO_HUMAN Tyrosinase (isoform 1) (SEQ ID NO:491)
Vitiligo	Melanocyte-Stimulating Hormone Alpha-MSH P01189 138-150 (SEQ ID NO:526) Beta-MSH P01189 217-234 (SEQ ID NO:527)

	Gamma-MSH P01189 77-87 (SEQ ID NO:528) Proopiomelanocortin P01189 COLI_HUMAN Pro-opiomelanocortin (SEQ ID NO:529)
Vitiligo	Tyrosinase P14679 TYRO_HUMAN Tyrosinase (isoform 1) (SEQ ID NO:491) Interferon Interferon, Alpha 1 P01562 IFNA1_HUMAN Interferon alpha-1/13 (SEQ ID NO:530) Interferon, Alpha 2 P01563 IFNA2_HUMAN Interferon alpha-2 (SEQ ID NO:531) Interferon, Alpha 4 P05014 IFNA4_HUMAN Interferon alpha-4 (SEQ ID NO:532) Interferon, Alpha 5 P01569 IFNA5_HUMAN Interferon alpha-5 (SEQ ID NO:533) Interferon, Alpha 6 P05013 IFNA6_HUMAN Interferon alpha-6 (SEQ ID NO:534) Interferon, Alpha 7 P01567 IFNA7_HUMAN Interferon alpha-7 (SEQ ID NO:535) Interferon, Alpha 8 P32881 IFNA8_HUMAN Interferon alpha-8 (SEQ ID NO:536) Interferon, Alpha 10 P01566 IFN10_HUMAN Interferon alpha-10 (SEQ ID NO:537) Interferon, Alpha 14 P01570 IFN14_HUMAN Interferon alpha-14 OS (SEQ ID NO:538) Interferon, Alpha 16 P05015 IFN16_HUMAN Interferon alpha-16 (SEQ ID NO:539)
Warts	Interferon, Alpha 17

	P01571 IFN17_HUMAN Interferon alpha-17 (SEQ ID NO:540) Interferon, Alpha 21 P01568 IFN21_HUMAN Interferon alpha-21 (SEQ ID NO:541) Interferon, Gamma P01579 IFNG_HUMAN Interferon gamma (SEQ ID NO:542) Interferon, Beta P01574 IFNB_HUMAN Interferon beta (SEQ ID NO:543) Interferon, Kappa Q9P0W0 IFNK_HUMAN Interferon kappa (SEQ ID NO:544) Interferon, Epsilon Q86WN2 IFNE_HUMAN Interferon epsilon (SEQ ID NO:545)
Wound Healing	Elastin sp P15502 ELN_HUMAN Elastin (isoform 3) (SEQ ID NO:486)
Wound Healing	Collagen Type I P02452 CO1A1_HUMAN Collagen alpha-1(I) chain (SEQ ID NO:487)
Wound Healing	P08123 CO1A2_HUMAN Collagen alpha-2(I) chain (SEQ ID NO:488)
Wound Healing	Collagen Type III P02461 CO3A1_HUMAN Collagen alpha-1(III) chain (isoform 1) (SEQ ID NO:489)
Xeroderma Pigmentosum	DNA Polymerase Eta Q9Y253 POLH_HUMAN DNA polymerase eta (isoform 1) (SEQ ID NO:507)

Additional illustrative targets of the present invention include the cosmetic targets listed in Table 6 of International Patent Publication No. WO 2013/151671, the contents of which are hereby incorporated by reference in their entirety.

- 5 In various embodiments, the agents of the present invention are used in methods the effect the integumentary system of a human. The present compositions and methods may be used to alter a biological and/or physiological process to, for example, reduce skin sagging, increase skin thickness, increase skin volume, reduce the number of wrinkles, the length of wrinkles and/or the depth of wrinkles, increase skin tightness,

firmness, tone and/or elasticity, increase skin hydration and ability to retain moisture, water flow and osmotic balance, increase the levels of skin lipids; increase the extracellular matrix and/or adhesion and communication polypeptides; increase skin energy production; utilization and conservation; improve oxygen utilization; improve skin cell life; improve skin cell immunity defense, heat shock stress response, antioxidant defense capacity to
 5 neutralize free radicals, and/or toxic defense; improve the protection and recovery from ultraviolet rays; improve skin cell communication and skin cell innervations; improve cell cohesion/adhesion; improve calcium mineral and other mineral metabolism; improve cell turnover; and improve cell circadian rhythms.

Further still, in some embodiments, the present compositions may be used in the treatment, control, or prevention of a disease, disorder and/or condition and/or may alter, modify or change the appearance of a
 10 member of the integumentary system of a subject suffering from a disease, disorder and/or condition such as, but not limited to, acne vulgaris, acne aestivalis, acne conglobata, acne cosmetic, acne fulminans, acne keloidalis nuchae, acne mechanica, acne medicamentosa, acne miliaris necrotica, acne necrotica, acne rosacea, actinic keratosis, acne vulgaris, acne aestivalis, acne conglobata, acne cosmetic, acne fulminans, acne keloidalis nuchae, acne mechanica, acne medicamentosa, acne miliaris necrotica, acne necrotica, acne rosacea, acute
 15 urticaria, allergic contact dermatitis, alopecia areata, angioedema, athlete's foot, atopic dermatitis, autoeczematization, baby acne, balding, blastomycosis, blackheads, birthmarks and other skin pigmentation problems, boils, bruises, bug bites and stings, burns, cellulitis, chiggers, chloracne, cholinergic or stress urticaria, chronic urticaria, cold type urticaria, confluent and reticulated papillomatosis, corns, cysts, dandruff, dermatitis herpetiformis, dermatographism, dyshidrotic eczema, diaper rash, dry skin, dyshidrosis, ectodermal dysplasia
 20 such as, hypohidrotic ectodermal dysplasia and X-linked hypohidrotic ectodermal dysplasia, eczema, epidermaodysplasia verruciformis, erythema nodosum, excoriated acne, exercise-induced anaphylaxis folliculitis, excess skin oil, folliculitis, freckles, frostbite, fungal nails, hair density, hair growth rate, halogen acne, hair loss, heat rash, hematoma, herpes simplex infections (e.g. non-genital), hidradenitis suppurativa, hives, hyperhidrosis, hyperpigmentation, hypohidrotic ectodermal dysplasia, hypopigmentation, impetigo, ingrown hair, heat type
 25 urticaria, ingrown toenail, infantile acne or neonatal acne, itch, irritant contact dermatitis, jock itch, keloid, keratosis pilaris, lichen planus, lichen sclerosus, lupus miliaris disseminatus faciei, melasma, moles, molluscum contagiosum, nail growth rate, nail health, neurodermatitis, nummular eczema, occupational acne, oil acne, onychomycosis, physical urticaria, pilonidal cyst, pityriasis rosea, pityriasis versicolor, poison ivy, pomade acne, pseudofolliculitis barbae or acne keloidalis nuchae, psoriasis, psoriatic arthritis, pressure or delayed pressure
 30 urticaria, puncture wounds such as cuts and scrapes, rash, rare or water type urticaria, rhinoplasty, ringworm, rosacea, rothmund-thomson syndrome, sagging of the skin, scabies, scars, seborrhea, seborrheic dermatitis, shingles, skin cancer, skin tag, solar type urticaria, spider bite, stretch marks, sunburn, tar acne, tropical acne, thinning of skin, thrush, tinea versicolor, transient acantholytic dermatosis, tycoon's cap or acne necrotica miliaris, uneven skin tone, varicose veins, venous eczema, vibratory angioedema, vitiligo, warts, Weber-Christian
 35 disease, wrinkles, x-linked hypohidrotic ectodermal dysplasia, xerotic eczema, yeast infection and general signs of aging.

In some embodiments, there is provided methods of treating, controlling or preventing dry skin with the present compositions. In some embodiments profilaggrin (a protein which is converted to filaggrin) is a protein of interest (e.g. when treating ichthyosis vulgaris).

In some embodiments, there is provided methods of treating, controlling or preventing any one of the various types of psoriasis (e.g. plaque psoriasis, guttate psoriasis, pustular psoriasis, inverse psoriasis, and erythrodermic psoriasis). In various embodiments, the protein of interest is any of the products of the genes psoriasis susceptibility 1 through 9 (PSORS1 - PSORS9).

Various embodiments relate to the treatment, control, or prevention of eczema (e.g. atopic dermatitis, nummular eczema, dyshidrotic eczema, seborrheic dermatitis, irritant contact dermatitis, allergic contact dermatitis, dyshidrosis, venous eczema, dermatitis herpetiformis, neurodermatitis, autoeczematization and xerotic eczema) and, optionally, one or more of the following may be targeted: filaggrin; three genetic variants, ovo-like 1 (OVOL1), actin-like 9 (ACTL9) and kinesin family member 3 A (KIF3A) have been associated with eczema; and the genes brain-derived neurotrophic factor (BDNF) and tachykinin, precursor 1 (TAC1).

Hives, or urticaria, including, but not limited to, acute urticaria, chronic urticaria and angioedema, physical urticaria, pressure or delayed pressure urticaria, cholinergic or stress urticaria, cold type urticaria, heat type urticaria, solar type urticaria, rare or water type urticaria, vibratory angioedema, exercise-induced anaphylaxis and dermatographism may be treated with the present compositions by, for example, targeting PLCG-2.

Various embodiments relate to the treatment, control, or prevention of rosacea, which includes, but is not limited to, erythematotelangiectatic rosacea, papulopustular rosacea, phymatous rosacea, and ocular rosacea. Optionally, cathelicidin antimicrobial peptide (CAMP) and/or kallikrein-related peptidase 5 (also known as stratum corneum tryptic enzyme (SCTE)) are proteins of interest.

In some embodiments, there is provided methods of treating, controlling or preventing acne with the present compositions. For example, acne may include, but is not limited to, acneiform eruptions, acne aestivalis, acne conglobata, acne cosmetic, acne fulminans, acne keloidalis nuchae, acne mechanica, acne medicamentosa, acne miliaris necrotica, acne necrotica, acne rosacea, baby acne, blackheads, chloracne, excoriated acne, halogen acne, infantile acne or neonatal acne, lupus miliaris disseminatus faciei, occupational acne, oil acne, pomade acne, tar acne, tropical acne, tycoon's cap or acne necrotica miliaris, pseudofolliculitis barbae or acne keloidalis nuchae, and hidradenitis suppurativa. In these embodiments, the protein of interest may be one or more matrix metalloproteinases (MMP), e.g., matrix metalloproteinase-1 (MMP-1 or interstitial collagenase), matrix metalloproteinase-9 (MMP-9), and matrix metalloproteinase-13 (MMP-13).

In further embodiments, vitiligo is treated with the present compositions, e.g. wherein the NLR family, pyrin domain containing 1 gene (NALP1) gene is targeted.

In some embodiments, the present compositions find use in the treatment, control, or prevention of hypohidrotic ectodermal dysplasia (HED), *e.g.* via the ectodysplasin A gene (EDA), receptor (EDAR), and receptor associated death domain (EDARADD).

In some embodiments, the present compositions find use in the treatment, control, or prevention of balding, or hair thinning (*e.g.* male pattern baldness, or androgenetic alopecia (AGA)) and, optionally, one or more of the following may be the protein of interest: androgen receptor (AR), ectodysplasin A2 receptor (EDA2R) and lysophosphatidic acid receptor 6 (P2RY5).

The present compositions also find use in methods of treatment, control, or prevention of scars and stretch marks (*striae*), *e.g.* via collagen, ribosomal s6 kinase, secreted phosphoprotein 1 (also known as osteopontin), or transforming growth factor beta 3.

Epidermodysplasia verruciformis (also known as Lutz-Lewandowsky epidermodysplasia), a rare autosomal recessive genetic hereditary skin disorder, may also be treated with compositions of the present invention, *e.g.* by targeted transmembrane channel-like 6 (EVER1) or transmembrane channellike 8 (EVER2) genes.

In some embodiments, skin sagging, thinning or wrinkling may be treated, controlled or prevented with present composition, *e.g.* by targeting one or more of the proteins of interest such as collagen, elastin, fibroblast growth factor 7, TIMP metalloproteinase inhibitors, matrix metalloproteinases, superoxide dismutase and other extracellular matrix proteins and proteoglycans.

Further embodiments are used in tanning of the skin, such as via melanocyte-stimulating hormone and/or pro-opiomelanocortin.

In some embodiments, the present compositions may be used for wound treatment. In some embodiments, methods of treating, controlling or preventing wounds with the present compositions comprises additional steps of, for example, cleaning the wound bed to facilitate wound healing and closure, including, but not limited to: debridement, sharp debridement (surgical removal of dead or infected tissue from a wound), optionally including chemical debriding agents, such as enzymes, to remove necrotic tissue; wound dressings to provide the wound with a moist, warm environment and to promote tissue repair and healing (*e.g.*, wound dressings comprising hydrogels (*e.g.*, AQUASORB; DUODERM), hydrocolloids (*e.g.*, AQUACEL; COMFEEL), foams (*e.g.*, LYOFAM; SPYROSORB), and alginates (*e.g.*, ALGISITE; CURASORB); administration of growth factors to stimulate cell division and proliferation and to promote wound healing *e.g.* becaplermin; and (iv) soft-tissue wound coverage, a skin graft may be necessary to obtain coverage of clean, non-healing wounds (*e.g.*, autologous skin grafts, cadaveric skin graft, bioengineered skin substitutes (*e.g.*, APLIGRAF; DERMAGRAFT)).

In various embodiments, the nucleic acid drug described herein can be used in a variety of cosmetic/plastic surgery procedures, including, without limitation, a surgical procedure involving skin grafting and an aesthetic or cosmetic surgery (*e.g.* a facial plastic surgery procedure including, but not limited to blepharoplasty, rhinoplasty, rhytidectomy, genioplasty, facial implants, otoplasty, hair implantation, cleft lip and cleft palate repair, and/or a

body plastic surgery procedure including but not limited to abdominoplasty, brachioplasty, thigh lift, breast reduction, breast augmentation, body contouring, liposuction, hand surgery).

In various embodiments, a variety of cancers are treated, controlled or prevented with the present compositions (e.g., colorectal cancer, gallbladder cancer, lung cancer, pancreatic cancer, and stomach cancer). In some
 5 embodiments, skin cancer is treated with the present compositions. For instance, the skin cancer is one or more of actinic keratosis, basal cell carcinoma, melanoma, Kaposi's sarcoma, and squamous cell carcinoma. In some embodiments, the present compositions are used adjuvant to complete circumferential peripheral and deep margin assessment, Mohs surgery, radiation (e.g. external beam radiotherapy or brachytherapy), chemotherapy (including but not limited to topical chemotherapy, e.g. with imiquimod or 5-fluorouracil), and cryotherapy. The
 10 present compositions also find use in the treatment of various stages of cancers, including skin cancers (e.g. basal cell cancer (BCC), squamous cell cancer (SCC), and melanoma), such as a stage of the American Joint Committee on Cancer (AJCC) TNM system (e.g. one or more of TX, T0, Tis, T1, T1a, T1b, T2, T2A, T2B, T3, T3a, T3b, T4, T4a, T4b, NX, N0, N1, N2, N3, M0, M1a, M1b, M1c) and/or a staging system (e.g. Stage 0, Stage IA, Stage IB, Stage IIA, Stage IIB, Stage IIC, Stage IIIA, Stage IIIB, Stage IIIC, Stage IV).

15 Illustrative cancers and/or tumors of the present invention include, but are not limited to, a basal cell carcinoma, biliary tract cancer; bladder cancer; bone cancer; brain and central nervous system cancer; breast cancer; cancer of the peritoneum; cervical cancer; choriocarcinoma; colon and rectum cancer; connective tissue cancer; cancer of the digestive system; endometrial cancer; esophageal cancer; eye cancer; cancer of the head and neck; gastric cancer (including gastrointestinal cancer); glioblastoma; hepatic carcinoma; hepatoma; intra-
 20 epithelial neoplasm; kidney or renal cancer; larynx cancer; leukemia; liver cancer; lung cancer (e.g., small-cell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung, and squamous carcinoma of the lung); melanoma; myeloma; neuroblastoma; oral cavity cancer (lip, tongue, mouth, and pharynx); ovarian cancer; pancreatic cancer; prostate cancer; retinoblastoma; rhabdomyosarcoma; rectal cancer; cancer of the respiratory system; salivary gland carcinoma; sarcoma; skin cancer; squamous cell cancer; stomach cancer; testicular
 25 cancer; thyroid cancer; uterine or endometrial cancer; cancer of the urinary system; vulval cancer; lymphoma including Hodgkin's and non-Hodgkin's lymphoma, as well as B-cell lymphoma (including low grade/follicular non-Hodgkin's lymphoma (NHL); small lymphocytic (SL) NHL; intermediate grade/follicular NHL; intermediate grade diffuse NHL; high grade immunoblastic NHL; high grade lymphoblastic NHL; high grade small non-cleaved cell NHL; bulky disease NHL; mantle cell lymphoma; AIDS-related lymphoma; and Waldenstrom's
 30 Macroglobulinemia; chronic lymphocytic leukemia (CLL); acute lymphoblastic leukemia (ALL); Hairy cell leukemia; chronic myeloblastic leukemia; as well as other carcinomas and sarcomas; and post-transplant lymphoproliferative disorder (PTLD), as well as abnormal vascular proliferation associated with phakomatoses, edema (such as that associated with brain tumors), and Meigs' syndrome.

In various embodiments, one or more rare diseases are treated, controlled or prevented with the present
 35 compositions, including, by way of illustration, Erythropoietic Protoporphyrria, Hailey-Hailey Disease,

Epidermolysis Bullosa (EB), Xeroderma Pigmentosum, Ehlers-Danlos Syndrome, Cutis Laxa, Protein C & Protein S Deficiency, Alport Syndrome, Striate Palmoplantar Keratoderma, Lethal Acantholytic EB, Pseudoxanthoma Elasticum (PXE), Ichthyosis Vulgaris, Pemphigus Vulgaris, and Basal Cell Nevus Syndrome.

In various embodiments, the present compositions are used to treat, control or prevent one or more inflammatory diseases or conditions, such as inflammation, acute inflammation, chronic inflammation, respiratory disease, atherosclerosis, restenosis, asthma, allergic rhinitis, atopic dermatitis, septic shock, rheumatoid arthritis, inflammatory bowel disease, inflammatory pelvic disease, pain, ocular inflammatory disease, celiac disease, Leigh Syndrome, Glycerol Kinase Deficiency, Familial eosinophilia (FE), autosomal recessive spastic ataxia, laryngeal inflammatory disease, Tuberculosis, Chronic cholecystitis, Bronchiectasis, Silicosis and other pneumoconioses.

In various embodiments, the present compositions are used to treat, control or prevent one or more autoimmune diseases or conditions, such as multiple sclerosis, diabetes mellitus, lupus, celiac disease, Crohn's disease, ulcerative colitis, Guillain-Barre syndrome, scleroderms, Goodpasture's syndrome, Wegener's granulomatosis, autoimmune epilepsy, Rasmussen's encephalitis, Primary biliary sclerosis, Sclerosing cholangitis, Autoimmune hepatitis, Addison's disease, Hashimoto's thyroiditis, Fibromyalgia, Menier's syndrome; transplantation rejection (e.g., prevention of allograft rejection) pernicious anemia, rheumatoid arthritis, systemic lupus erythematosus, dermatomyositis, Sjogren's syndrome, lupus erythematosus, multiple sclerosis, myasthenia gravis, Reiter's syndrome, Grave's disease, and other autoimmune diseases.

In various embodiments, the present compositions are used to treat, control or prevent one or more neurologic diseases, including ADHD, AIDS—Neurological Complications, Absence of the Septum Pellucidum, Acquired Epileptiform Aphasia, Acute Disseminated Encephalomyelitis, Adrenoleukodystrophy, Agenesis of the Corpus Callosum, Agnosia, Aicardi Syndrome, Alexander Disease, Alpers' Disease, Alternating Hemiplegia, Alzheimer's Disease, Amyotrophic Lateral Sclerosis, Anencephaly, Aneurysm, Angelman Syndrome, Angiomatosis, Anoxia, Aphasia, Apraxia, Arachnoid Cysts, Arachnoiditis, Arnold-Chiari Malformation, Arteriovenous Malformation, Aspartame, Asperger Syndrome, Ataxia Telangiectasia, Ataxia, Attention Deficit-Hyperactivity Disorder, Autism, Autonomic Dysfunction, Back Pain, Barth Syndrome, Batten Disease, Behcet's Disease, Bell's Palsy, Benign Essential Blepharospasm, Benign Focal Amyotrophy, Benign Intracranial Hypertension, Bernhardt-Roth Syndrome, Binswanger's Disease, Blepharospasm, Bloch-Sulzberger Syndrome, Brachial Plexus Birth Injuries, Brachial Plexus Injuries, Bradbury-Eggleston Syndrome, Brain Aneurysm, Brain Injury, Brain and Spinal Tumors, Brown-Sequard Syndrome, Bulbospinal Muscular Atrophy, Canavan Disease, Carpal Tunnel Syndrome, Causalgia, Cavernomas, Cavernous Angioma, Cavernous Malformation, Central Cervical Cord Syndrome, Central Cord Syndrome, Central Pain Syndrome, Cephalic Disorders, Cerebellar Degeneration, Cerebellar Hypoplasia, Cerebral Aneurysm, Cerebral Arteriosclerosis, Cerebral Atrophy, Cerebral Beriberi, Cerebral Gigantism, Cerebral Hypoxia, Cerebral Palsy, Cerebro-Oculo-Facio-Skeletal Syndrome, Charcot-Marie-Tooth Disorder, Chiari Malformation, Chorea, Choreoacanthocytosis, Chronic Inflammatory Demyelinating

Polyneuropathy (CIDP), Chronic Orthostatic Intolerance, Chronic Pain, Cockayne Syndrome Type II, Coffin
 Lowry Syndrome, Coma, including Persistent Vegetative State, Complex Regional Pain Syndrome, Congenital
 Facial Diplegia, Congenital Myasthenia, Congenital Myopathy, Congenital Vascular Cavernous Malformations,
 Corticobasal Degeneration, Cranial Arteritis, Craniosynostosis, Creutzfeldt-Jakob Disease, Cumulative Trauma
 5 Disorders, Cushing's Syndrome, Cytomegalic Inclusion Body Disease (CIBD), Cytomegalovirus Infection,
 Dancing Eyes-Dancing Feet Syndrome, Dandy-Walker Syndrome, Dawson Disease, De Morsier's Syndrome,
 Dejerine-Klumpke Palsy, Dementia—Multi-Infarct, Dementia—Subcortical, Dementia With Lewy Bodies,
 Dermatomyositis, Developmental Dyspraxia, Devic's Syndrome, Diabetic Neuropathy, Diffuse Sclerosis, Dravet's
 Syndrome, Dysautonomia, Dysgraphia, Dyslexia, Dysphagia, Dyspraxia, Dystonias, Early Infantile Epileptic
 10 Encephalopathy, Empty Sella Syndrome, Encephalitis Lethargica, Encephalitis and Meningitis, Encephaloceles,
 Encephalopathy, Encephalotrigeminal Angiomatosis, Epilepsy, Erb's Palsy, Erb-Duchenne and Dejerine-
 Klumpke Palsies, Fabry's Disease, Fahr's Syndrome, Fainting, Familial Dysautonomia, Familial Hemangioma,
 Familial Idiopathic Basal Ganglia Calcification, Familial Spastic Paralysis, Febrile Seizures (e.g., GEFS and
 GEFS plus), Fisher Syndrome, Floppy Infant Syndrome, Friedreich's Ataxia, Gaucher's Disease, Gerstmann's
 15 Syndrome, Gerstmann-Straussler-Scheinker Disease, Giant Cell Arteritis, Giant Cell Inclusion Disease, Globoid
 Cell Leukodystrophy, Glossopharyngeal Neuralgia, Guillain-Barre Syndrome, HTLV-1 Associated Myelopathy,
 Hallervorden-Spatz Disease, Head Injury, Headache, Hemispheric Continuity, Hemifacial Spasm, Hemiplegia
 Alterans, Hereditary Neuropathies, Hereditary Spastic Paraplegia, Hereditary Ataxia Polyneuritis, Hereditary
 Herpes Zoster Oticus, Herpes Zoster, Hirayama Syndrome, Holoprosencephaly, Huntington's Disease,
 20 Hydranencephaly, Hydrocephalus—Normal Pressure, Hydrocephalus, Hydromyelia, Hypercortisolism,
 Hypersomnia, Hypertonia, Hypotonia, Hypoxia, Immune-Mediated Encephalomyelitis, Inclusion Body Myositis,
 Incontinentia Pigmenti, Infantile Hypotonia, Infantile Phytanic Acid Storage Disease, Infantile Refsum Disease,
 Infantile Spasms, Inflammatory Myopathy, Intestinal Lipodystrophy, Intracranial Cysts, Intracranial Hypertension,
 Isaac's Syndrome, Joubert Syndrome, Kearns-Sayre Syndrome, Kennedy's Disease, Kinsbourne syndrome,
 25 Kleine-Levin syndrome, Klippel Feil Syndrome, Klippel-Trenaunay Syndrome (KTS), Klüver-Bucy Syndrome,
 Korsakoff's Amnesic Syndrome, Krabbe Disease, Kugelberg-Welander Disease, Kuru, Lambert-Eaton
 Myasthenic Syndrome, Landau-Kleffner Syndrome, Lateral Femoral Cutaneous Nerve Entrapment, Lateral
 Medullary Syndrome, Learning Disabilities, Leigh's Disease, Lennox-Gastaut Syndrome, Lesch-Nyhan
 Syndrome, Leukodystrophy, Levine-Critchley Syndrome, Lewy Body Dementia, Lissencephaly, Locked-In
 30 Syndrome, Lou Gehrig's Disease, Lupus—Neurological Sequelae, Lyme Disease—Neurological Complications,
 Machado-Joseph Disease, Macrencephaly, Megalencephaly, Melkersson-Rosenthal Syndrome, Meningitis,
 Menkes Disease, Meralgia Paresthetica, Metachromatic Leukodystrophy, Microcephaly, Migraine, Miller Fisher
 Syndrome, Mini-Stroke, Mitochondrial Myopathies, Mobius Syndrome, Monomelic Amyotrophy, Motor Neuron
 Diseases, Moyamoya Disease, Mucopolysaccharidoses, Mucopolysaccharidoses, Multi-Infarct Dementia, Multifocal Motor
 35 Neuropathy, Multiple Sclerosis, Multiple System Atrophy with Orthostatic Hypotension, Multiple System Atrophy,
 Muscular Dystrophy, Myasthenia—Congenital, Myasthenia Gravis, Myelinoclastic Diffuse Sclerosis, Myoclonic

Encephalopathy of Infants, Myoclonus, Myopathy—Congenital, Myopathy—Thyrotoxic, Myopathy, Myotonia Congenita, Myotonia, Narcolepsy, Neuroacanthocytosis, Neurodegeneration with Brain Iron Accumulation, Neurofibromatosis, Neuroleptic Malignant Syndrome, Neurological Complications of AIDS, Neurological Manifestations of Pompe Disease, Neuromyelitis Optica, Neuromyotonia, Neuronal Ceroid Lipofuscinosis, Neuronal Migration Disorders, Neuropathy—Hereditary, Neurosarcoidosis, Neurotoxicity, Nevus Cavernosus, Niemann-Pick Disease, O'Sullivan-McLeod Syndrome, Occipital Neuralgia, Occult Spinal Dysraphism Sequence, Ohtahara Syndrome, Olivopontocerebellar Atrophy, Opsoclonus Myoclonus, Orthostatic Hypotension, Overuse Syndrome, Pain—Chronic, Paraneoplastic Syndromes, Paresthesia, Parkinson's Disease, Parnyotonia Congenita, Paroxysmal Choreoathetosis, Paroxysmal Hemicrania, Parry-Romberg, Pelizaeus-Merzbacher Disease, Pena Shokeir II Syndrome, Perineural Cysts, Periodic Paralyzes, Peripheral Neuropathy, Periventricular Leukomalacia, Persistent Vegetative State, Pervasive Developmental Disorders, Phytanic Acid Storage Disease, Pick's Disease, Piriformis Syndrome, Pituitary Tumors, Polymyositis, Pompe Disease, Porencephaly, Post-Polio Syndrome, Postherpetic Neuralgia, Postinfectious Encephalomyelitis, Postural Hypotension, Postural Orthostatic Tachycardia Syndrome, Postural Tachycardia Syndrome, Primary Lateral Sclerosis, Prion Diseases, Progressive Hemifacial Atrophy, Progressive Locomotor Ataxia, Progressive Multifocal Leukoencephalopathy, Progressive Sclerosing Poliodystrophy, Progressive Supranuclear Palsy, Pseudotumor Cerebri, Pyridoxine Dependent and Pyridoxine Responsive Seizure Disorders, Ramsay Hunt Syndrome Type I, Ramsay Hunt Syndrome Type II, Rasmussen's Encephalitis and other autoimmune epilepsies, Reflex Sympathetic Dystrophy Syndrome, Refsum Disease—Infantile, Refsum Disease, Repetitive Motion Disorders, Repetitive Stress Injuries, Restless Legs Syndrome, Retrovirus-Associated Myelopathy, Rett Syndrome, Reye's Syndrome, Riley-Day Syndrome, SUNCT Headache, Sacral Nerve Root Cysts, Saint Vitus Dance, Salivary Gland Disease, Sandhoff Disease, Schilder's Disease, Schizencephaly, Seizure Disorders, Septo-Optic Dysplasia, Severe Myoclonic Epilepsy of Infancy (SMEI), Shaken Baby Syndrome, Shingles, Shy-Drager Syndrome, Sjogren's Syndrome, Sleep Apnea, Sleeping Sickness, Soto's Syndrome, Spasticity, Spina Bifida, Spinal Cord Infarction, Spinal Cord Injury, Spinal Cord Tumors, Spinal Muscular Atrophy, Spinocerebellar Atrophy, Steele-Richardson-Olszewski Syndrome, Stiff-Person Syndrome, Striatonigral Degeneration, Stroke, Sturge-Weber Syndrome, Subacute Sclerosing Panencephalitis, Subcortical Arteriosclerotic Encephalopathy, Swallowing Disorders, Sydenham Chorea, Syncope, Syphilitic Spinal Sclerosis, Syringohydromyelia, Syringomyelia, Systemic Lupus Erythematosus, Tabes Dorsalis, Tardive Dyskinesia, Tarlov Cysts, Tay-Sachs Disease, Temporal Arteritis, Tethered Spinal Cord Syndrome, Thomsen Disease, Thoracic Outlet Syndrome, Thyrotoxic Myopathy, Tic Douloureux, Todd's Paralysis, Tourette Syndrome, Transient Ischemic Attack, Transmissible Spongiform Encephalopathies, Transverse Myelitis, Traumatic Brain Injury, Tremor, Trigeminal Neuralgia, Tropical Spastic Paraparesis, Tuberous Sclerosis, Vascular Erectile Tumor, Vasculitis including Temporal Arteritis, Von Economo's Disease, Von Hippel-Lindau disease (VHL), Von Recklinghausen's Disease, Wallenberg's Syndrome, Werdnig-Hoffman Disease, Wernicke-Korsakoff Syndrome, West Syndrome, Whipple's

Disease, Williams Syndrome, Wilson's Disease, X-Linked Spinal and Bulbar Muscular Atrophy, and Zellweger Syndrome.

In various embodiments, the present compositions are used to treat one or more respiratory diseases, such as asthma, chronic obstructive pulmonary disease (COPD), bronchiectasis, allergic rhinitis, sinusitis, pulmonary vasoconstriction, inflammation, allergies, impeded respiration, respiratory distress syndrome, cystic fibrosis, pulmonary hypertension, pulmonary vasoconstriction, emphysema, Hantavirus pulmonary syndrome (HPS), Loeffler's syndrome, Goodpasture's syndrome, Pleurisy, pneumonitis, pulmonary edema, pulmonary fibrosis, Sarcoidosis, complications associated with respiratory syncytial virus infection, and other respiratory diseases.

In various embodiments, the present compositions are used to treat, control or prevent cardiovascular disease, such as a disease or condition affecting the heart and vasculature, including but not limited to, coronary heart disease (CHD), cerebrovascular disease (CVD), aortic stenosis, peripheral vascular disease, atherosclerosis, arteriosclerosis, myocardial infarction (heart attack), cerebrovascular diseases (stroke), transient ischaemic attacks (TIA), angina (stable and unstable), atrial fibrillation, arrhythmia, valvular disease, and/or congestive heart failure.

In various embodiments, the present compositions are used to treat, control or prevent one or more metabolic-related disorders. In various embodiments, the present invention is useful for the treatment, controlling or prevention of diabetes, including Type 1 and Type 2 diabetes and diabetes associated with obesity. The compositions and methods of the present invention are useful for the treatment or prevention of diabetes-related disorders, including without limitation diabetic nephropathy, hyperglycemia, impaired glucose tolerance, insulin resistance, obesity, lipid disorders, dyslipidemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, low HDL levels, high LDL levels, atherosclerosis and its sequelae, vascular restenosis, irritable bowel syndrome, inflammatory bowel disease, including Crohn's disease and ulcerative colitis, other inflammatory conditions, pancreatitis, abdominal obesity, neurodegenerative disease, retinopathy, neoplastic conditions, adipose cell tumors, adipose cell carcinomas, such as liposarcoma, prostate cancer and other cancers, including gastric, breast, bladder and colon cancers, angiogenesis, Alzheimer's disease, psoriasis, high blood pressure, Metabolic Syndrome (e.g. a person has three or more of the following disorders: abdominal obesity, hypertriglyceridemia, low HDL cholesterol, high blood pressure, and high fasting plasma glucose), ovarian hyperandrogenism (polycystic ovary syndrome), and other disorders where insulin resistance is a component, such as sleep apnea. The compositions and methods of the present invention are useful for the treatment, control, or prevention of obesity, including genetic or environmental, and obesity-related disorders. The obesity-related disorders herein are associated with, caused by, or result from obesity. Examples of obesity-related disorders include obesity, diabetes, overeating, binge eating, and bulimia, hypertension, elevated plasma insulin concentrations and insulin resistance, dyslipidemia, hyperlipidemia, endometrial, breast, prostate, kidney and colon cancer, osteoarthritis, obstructive sleep apnea, gallstones, heart disease, abnormal heart rhythms and arrhythmias, myocardial infarction, congestive heart failure, coronary heart disease, sudden death, stroke, polycystic ovary disease,

craniopharyngioma, Prader-Willi Syndrome, Frohlich's syndrome, GH-deficient subjects, normal variant short stature, Turner's syndrome, and other pathological conditions showing reduced metabolic activity or a decrease in resting energy expenditure as a percentage of total fat-free mass, e.g, children with acute lymphoblastic leukemia. Further examples of obesity-related disorders are Metabolic Syndrome, insulin resistance syndrome, reproductive hormone abnormalities, sexual and reproductive dysfunction, such as impaired fertility, infertility, hypogonadism in males and hirsutism in females, fetal defects associated with maternal obesity, gastrointestinal motility disorders, such as obesity-related gastro-esophageal reflux, respiratory disorders, such as obesity-hypoventilation syndrome (Pickwickian syndrome), breathlessness, cardiovascular disorders, inflammation, such as systemic inflammation of the vasculature, arteriosclerosis, hypercholesterolemia, lower back pain, gallbladder disease, hyperuricemia, gout, and kidney cancer, and increased anesthetic risk. The compositions and methods of the present invention are also useful to treat Alzheimer's disease.

Nucleic acids, including liposomal formulations containing nucleic acids, when delivered in vivo, can accumulate in the liver and/or spleen. It has now been discovered that nucleic acids encoding proteins can modulate protein expression in the liver and spleen, and that nucleic acids used in this manner can constitute potent therapeutics for the treatment of liver and spleen diseases. Certain embodiments are therefore directed to a method for treating liver and/or spleen disease by delivering to a patient a nucleic acid encoding a protein of interest. Other embodiments are directed to a therapeutic composition comprising a nucleic acid encoding a protein of interest, for the treatment of liver and/or spleen disease. Diseases and conditions of the liver and/or spleen that can be treated include, but are not limited to: hepatitis, alcohol-induced liver disease, drug-induced liver disease, Epstein Barr virus infection, adenovirus infection, cytomegalovirus infection, toxoplasmosis, Rocky Mountain spotted fever, non-alcoholic fatty liver disease, hemochromatosis, Wilson's Disease, Gilbert's Disease, and cancer of the liver and/or spleen.

In some embodiments, the present compositions and methods relate to the treatment of type 1 diabetes, heart disease, including ischemic and dilated cardiomyopathy, macular degeneration, Parkinson's disease, cystic fibrosis, sickle-cell anemia, thalassemia, Fanconi anemia, severe combined immunodeficiency, hereditary sensory neuropathy, xeroderma pigmentosum, Huntington's disease, muscular dystrophy, amyotrophic lateral sclerosis, Alzheimer's disease, cancer, and infectious diseases including hepatitis and HIV/AIDS.

Further, in some embodiments, the present methods and compositions find use in targeting any of the proteins or in treatment of any of the diseases or disorders of **Table 2B**. In various embodiments, the present invention contemplates the targeting of the full-length and/or truncated forms of any of the proteins disclosed in Table 2B. In various embodiments, the present invention contemplates the targeting of the precursor forms and/or mature forms and/or isoforms of any of the proteins disclosed in **Table 2B**.

In various embodiments, the present invention contemplates the targeting of a protein having about 60% (e.g. about 60%, or about 61%, or about 62%, or about 63%, or about 64%, or about 65%, or about 66%, or about 67%, or about 68%, or about 69%, or about 70%, or about 71%, or about 72%, or about 73%, or about 74%, or

about 75%, or about 76%, or about 77%, or about 78%, or about 79%, or about 80%, or about 81%, or about 82%, or about 83%, or about 84%, or about 85%, or about 86%, or about 87%, or about 88%, or about 89%, or about 90%, or about 91%, or about 92%, or about 93%, or about 94%, or about 95%, or about 96%, or about 97%, or about 98%, or about 99%) sequence identity with any of the protein sequences disclosed herein (e.g. in **Table 2B**).

In various embodiments, the present invention contemplates the targeting of a protein comprising an amino acid sequence having one or more amino acid mutations relative to any of the protein sequences described herein (e.g. in **Table 2B**). For example, the present invention contemplates the targeting of a protein comprising an amino acid sequence having 1, or 2, or 3, or 4, or 5, or 6, or 7, or 8, or 9, or 10, or 11, or 12 amino acid mutations relative to any of the protein sequences described herein (e.g. in **Table 2B**). In some embodiments, the one or more amino acid mutations may be independently selected from substitutions, insertions, deletions, and truncations.

In some embodiments, the amino acid mutations are amino acid substitutions, and may include conservative and/or non-conservative substitutions.

"Conservative substitutions" may be made, for instance, on the basis of similarity in polarity, charge, size, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the amino acid residues involved. The naturally occurring amino acids can be grouped into the following six standard amino acid groups: (1) hydrophobic: Met, Ala, Val, Leu, Ile; (2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln; (3) acidic: Asp, Glu; (4) basic: His, Lys, Arg; (5) residues that influence chain orientation: Gly, Pro; and (6) aromatic: Trp, Tyr, Phe.

As used herein, "conservative substitutions" are defined as exchanges of an amino acid by another amino acid listed within the same group of the six standard amino acid groups shown above. For example, the exchange of Asp by Glu retains one negative charge in the so modified polypeptide. In addition, glycine and proline may be substituted for one another based on their ability to disrupt α -helices.

As used herein, "non-conservative substitutions" are defined as exchanges of an amino acid by another amino acid listed in a different group of the six standard amino acid groups (1) to (6) shown above.

In various embodiments, the substitutions may also include non-classical amino acids (e.g. selenocysteine, pyrrolysine, *N*-formylmethionine β -alanine, GABA and δ -Aminolevulinic acid, 4-aminobenzoic acid (PABA), D-isomers of the common amino acids, 2,4-diaminobutyric acid, α -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-amino butyric acid, γ -Abu, ϵ -Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, homocitrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, β -alanine, fluoro-amino acids, designer amino acids such as β methyl amino acids, C α -methyl amino acids, N α -methyl amino acids, and amino acid analogs in general).

In **Table 2B**, all Illustrative Identifiers (e.g. Gene Seq nos. and references are hereby incorporated by reference in their entireties).

Table 2B. *Illustrative Proteins, Illustrative Peptides, and Illustrative Indications*

Protein / Peptide Illustrative Identifier Reference	Description / Illustrative Indication(s)
Parathyroid hormone P01270 PTHY_HUMAN Parathyroid hormone (SEQ ID NO:508)	PTH is secreted by the chief cells of the parathyroid glands. PTH plays an important role in the regulation of serum calcium, serum phosphate, and vitamin D synthesis.
BMP-1 GeneSeq Accession P80618 WO8800205 P13497/BMP1_HUMAN Bone morphogenetic protein 1 (isoform BMP1-3) (SEQ ID NO:169) P13497-2 BMP1_HUMAN Isoform BMP1-1 of Bone morphogenetic protein 1 (isoform BMP1-1) (SEQ ID NO:509)	BMP1 belongs to the transforming growth factor-beta (TGFB) super- family. Bone morphogenic proteins induce cartilage and bone formation, play important role in nephrogenesis, and play an important role in the development of many organs, including lung, heart, teeth, gut, skin, and particularly the kidney. BMP-1 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
P13497-3 BMP1_HUMAN Isoform BMP1-4 of Bone morphogenetic protein 1 (isoform BMP1-4) (SEQ ID NO:510)	
P13497-4 BMP1_HUMAN Isoform BMP1-5 of Bone morphogenetic protein 1 (isoform BMP1-5) (SEQ ID NO:511)	
P13497-5 BMP1_HUMAN Isoform BMP1-6 of Bone morphogenetic protein 1 (isoform BMP1-6) (SEQ ID NO:512)	
P13497-6 BMP1_HUMAN Isoform BMP1-7 of Bone	

morphogenetic protein 1 (isoform BMP1-7) (SEQ ID NO:513)	
BMP-2 GeneSeq Accession P80619 WO8800205 P12643/BMP2_HUMAN Bone morphogenetic protein 2 (SEQ ID NO:170)	BMP-2 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenetic protein induces bone formation. BMP-2 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes. Infarction recovery. Bone repair. Osteoporosis.
BMP-3 P12645 BMP3_HUMAN Bone morphogenetic protein 3 (SEQ ID NO:514)	BMP-3 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenetic protein induces bone formation. BMP-3 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
BMP-2B GeneSeq Accession W24850 US Pat. No. 5,631,142 P12644/BMP4_HUMAN Bone morphogenetic protein 4 (SEQ ID NO:171)	BMP-2b belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenetic protein induces bone formation. BMP-2b activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
BMP-4 GeneSeq Accession B02796 WO0020591 P12644/BMP4_HUMAN Bone morphogenetic protein 4 (SEQ ID NO:172)	BMP-4 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenetic protein induces bone formation. BMP-4 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
BMP-5 GeneSeq Accession B02797 WO0020591 P22003/BMP5_HUMAN Bone morphogenetic protein 5 (isoform 1) (SEQ ID NO:173) P22003-2 BMP5_HUMAN Isoform 2 of Bone morphogenetic protein 5 (isoform 2)	BMP-5 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenetic protein induces bone formation. BMP-5 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes

(SEQ ID NO:515)	
BMP-6 GeneSeq Accession R32904 US Pat. No. 5,187,076 P22004/BMP6_HUMAN Bone morphogenetic protein 6	BMP-6 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenic protein induces bone formation. BMP-6 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes. Hemochromatosis.
(SEQ ID NO:174)	
Osteogenic Protein-1; OP-1; BMP-7 GeneSeq Accession W34783 WO973462 P18075/BMP7_HUMAN Bone morphogenetic protein 7	OP-1 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenic protein induces bone formation. OP-1 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84-8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
(SEQ ID NO:175)	
BMP7 Variant A (SEQ ID NO:579)	OP-1 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenic protein induces bone formation. OP-1 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84-8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
BMP7 Variant B (SEQ ID NO:580)	OP-1 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenic protein induces bone formation. OP-1 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84-8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
BMP7 Variant C (SEQ ID NO:581)	OP-1 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenic protein induces bone formation. OP-1 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84-8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
Osteogenic Protein-2 GeneSeq Accession R57973 WO9406399 P34820/BMP8B_HUMAN Bone morphogenetic protein 8B	OP-2 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenic protein induces bone formation. OP-2 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84-8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
(SEQ ID NO:176)	
GDF-1 GeneSeq Accession R60961 WO9406449 P27539/GDF1_HUMAN	Members of the TGF-beta family of proteins initiate cell signaling by binding to heteromeric receptor complexes of type I (TbetaRI) and type II (TbetaRII) serine/threonine kinase receptors (reviewed by Massague, J. <i>et al.</i> (1994) Trends Cell Biol. 4: 172-178; Miyazono, K. <i>et al.</i> (1994) Adv. Immunol. 55: 181-220). Activation of this heteromeric receptor complex occurs when TGF-beta binds to

Embryonic growth/differentiation factor 1 (SEQ ID NO:177)	TbetaRII, which then recruits and phosphorylates TbetaRI. Activated TbetaRI then propagates the signal to downstream targets (Chen, F. and Weinberg, R. A. (1995) PNA892: 1565-1569; Wrana, J. L. <i>et al.</i> (1994) Nature 370: 341-347). The effect of GDF-1 on signaling can be assayed by treating Primary BAECs transferred with a construct called p3TP-Lux, containing a TGF-beta responsive promoter fused to a reporter gene, and measuring luciferase gene expression (Wrana <i>et al.</i> , 1994, Nature 370: 341-347). Developmental disorders, Induction of Cartilage, Tissue and Bone Growth, and Diabetes
BMP-9 GeneSeq Accession R86903 WO9533830 Q9UK05/GDF2_HUMAN Growth/differentiation factor 2 (SEQ ID NO:178)	BMP-9 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenic protein induces bone formation. BMP-9 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
BMP-10 GeneSeq Accession R66202 WO9426893 O95393/BMP10_HUMAN Bone morphogenetic protein 10 (SEQ ID NO:179)	BMP-10 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenic protein induces bone formation. BMP-10 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
BMP-12 GeneSeq Accession R78734 WO9516035 Q7Z4P5/GDF7_HUMAN Growth/differentiation factor 7 (SEQ ID NO:180)	BMP-12 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenic protein induces bone formation. BMP-12 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
BMP-15 GeneSeq Accession W11261 W09636710 O95972/BMP15_HUMAN Bone morphogenetic protein 15 (SEQ ID NO:181)	BMP-15 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenic protein induces bone formation. BMP-15 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
BMP-17 GeneSeq Accession Y17870 WO9929718 SEQ ID NO:2 from US Patent No. 7151086 (SEQ ID NO:182)	BMP-17 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenic protein induces bone formation. BMP-17 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
BMP-18 GeneSeq	BMP-18 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone

Accession Y17871 WO9929718 SEQ ID NO:4 from US Patent No. 7151086 (SEQ ID NO:183)	morphogenic protein induces bone formation. BMP-18 activity can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Induction of Cartilage, Tissue and Bone Growth, and Diabetes
Inhibin alpha GeneSeq Accession B02806 WO0020591 P05111/INHA_HUMAN Inhibin alpha chain (SEQ ID NO:184)	The inhibin beta A subunit joins the alpha subunit to form a pituitary FSH secretion inhibitor. Inhibin has been shown to regulate gonadal stromal cell proliferation negatively and to have tumor-suppressor activity. In addition, serum levels of inhibin have been shown to reflect the size of granulosa-cell tumors and can therefore be used as a marker for primary as well as recurrent disease. Tumor suppressor activity of inhibin can be determined using assays known in the art: Matzuk <i>et al.</i>, Nature 1992 Nov. 26: 360 (6402); 313-9. Tumor suppression.
Inhibin beta GeneSeq Accession H02808 WO0020591 P08476/INHBA_HUMAN Inhibin beta A chain (SEQ ID NO:185) P09529/INHBB_HUMAN Inhibin beta B chain (SEQ ID NO:186)	The inhibin beta A subunit joins the alpha subunit to form a pituitary FSH secretion inhibitor. Inhibin has been shown to regulate gonadal stromal cell proliferation negatively and to have tumour-suppressor activity. In addition, serum levels of inhibin have been shown to reflect the size of granulosa-cell tumors and can therefore be used as a marker for primary as well as recurrent disease. Tumor suppressor activity of inhibin can be determined using assays known in the art: Matzuk <i>et al.</i>, Nature 1992 Nov. 26: 360 (6402); 313-9. Tumor suppression.
Cerberus Protein GeneSeq Accession W86032 WO9849296 O95813/CER1_HUMAN Cerberus (SEQ ID NO:187)	Cerebus is believed to be involved in the inhibition of BMP activity BMP activity, in the presence of the antagonist Cerebus, can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. BMP Antagonist useful for Osteosarcoma, abnormal bone growth.
Soluble BMP Receptor Kinase Protein- 3 GeneSeq Accession R95227 WO9614579 Q13873/BMPR2_HUMAN Bone morphogenetic protein receptor type-2 (SEQ ID NO:188)	Soluble BMP receptor kinase protein-3 is involved in the binding of BMPs. Soluble BMP receptor kinase protein-3 is useful as an antagonist for the inhibition of BMP activity. BMP activity, in the presence of the soluble antagonist BMP receptor kinase protein-3, can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84- 8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. BMP Antagonist useful for Osteosarcoma, abnormal bone growth.
BMP Processing Enzyme Furin GeneSeq Accession W36099 WO9741250 P09958/FURIN_HUMAN Furin	BMPs belong to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenic protein induces bone formation. BMP activity, in the presence of the Furin, can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84-8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 10897-10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Dev. 10, 1580-1594. Bone formation or Regeneration Abnormalities

(SEQ ID NO:189)	
TGF-beta 1 GeneSeq Accession R29657 WO9216228 P01137/TGFB1_HUMAN Transforming growth factor beta-1 (SEQ ID NO:190)	Members of the TGF-beta family of proteins initiate cell signaling by binding to heteromeric receptor complexes of type I (TbetaRI) and type II (TbetaRII) serine/threonine kinase receptors (reviewed by Massague, J. <i>et al.</i> (1994) Trends Cell Biol. 4: 172 178; Miyazono, K. <i>et al.</i> (1994) Adv. Immunol. 55: 181-220). Activation of this heteromeric receptor complex occurs when TGF-beta. binds to TbetaRII, which then recruits and phosphorylates TbetaRI. Activated TbetaRI then propagates the signal to downstream targets (Chen, F. and Weinberg. R. A. (1995) PNA892: 1565-1569; Wrana, J. L. <i>et al.</i> (1994) Nature 370: 341. The effect of TGF betas on signaling can be assayed by treating Primary BAECs transfected with a construct called p3TP-Lux, containing a TGF- beta responsive promoter fused to a reporter gene, and measuring luciferase gene expression (Wrana <i>et al.</i>, 1994, Nature 370: 341-347). Useful for treating cancer and to promote wound healing.
TGF-beta 2 GeneSeq Accession R39659 EP542679 P61812/TGFB2_HUMAN Transforming growth factor beta-2 (SEQ ID NO:191)	Members of the TGF-beta family of proteins initiate cell signaling by binding to heteromeric receptor complexes of type I (TbetaRI) and type II (TbetaRII) serine/threonine kinase receptors (reviewed by Massague, J. <i>et al.</i> (1994) Trends Cell Biol. 4: 172 178; Miyazono, K. <i>et al.</i> (1994) Adv. Immunol. 55: 181-220). Activation of this heteromeric receptor complex occurs when TGF-beta. binds to TbetaRII, which then recruits and phosphorylates TbetaRI. Activated TbetaRI then propagates the signal to downstream targets (Chen, F. and Weinberg. R. A. (1995) PNA892: 1565-1569; Wrana, J. L. <i>et al.</i> (1994) Nature 370: 341. The effect of TGF betas on signaling can be assayed by treating Primary BAECs transfected with a construct called p3TP-Lux, containing a TGF- beta responsive promoter fused to a reporter gene, and measuring luciferase gene expression (Wrana <i>et al.</i>, 1994, Nature 370: 341-347). Useful for treating cancer and to promote wound healing.
ZTGF-beta 9 GeneSeq Accession Y70654 WO0015798 SEQ ID NO:2 of WO0015798 (SEQ ID NO:192)	Members of the TGF-beta family of proteins initiate cell signaling by binding to heteromeric receptor complexes of type I (TbetaRI) and type II (TbetaRII) serine/threonine kinase receptors (reviewed by Massague, J. <i>et al.</i> (1994) Trends Cell Biol. 4: 172 178; Miyazono, K. <i>et al.</i> (1994) Adv. Immunol. 55: 181-220). Activation of this heteromeric receptor complex occurs when TGF-beta. binds to TbetaRII, which then recruits and phosphorylates TbetaRI. Activated TbetaRI then propagates the signal to downstream targets (Chen, F. and Weinberg. R. A. (1995) PNA892: 1565-1569; Wrana, J. L. <i>et al.</i> (1994) Nature 370: 341. The effect of TGF betas on signaling can be assayed by treating Primary BAECs transfected with a construct called p3TP-Lux, containing a TGF- beta responsive promoter fused to a reporter gene, and measuring luciferase gene expression (Wrana <i>et al.</i>, 1994, Nature 370: 341-347). Useful for treating cancer and to promote wound healing.
Anti-TGF beta family antibodies GB2305921	Members of the TGF-beta family of proteins initiate cell signaling by binding to heteromeric receptor complexes of type I (TbetaRI) and type II (TbetaRII) serine/threonine kinase receptors (reviewed by Massague, J. <i>et al.</i> (1994) Trends Cell Biol. 4: 172 178; Miyazono, K. <i>et al.</i> (1994) Adv. Immunol. 55: 181-220). Activation of this heteromeric receptor complex occurs when TGF-beta. binds to TbetaRII, which then recruits and phosphorylates TbetaRI. Activated TbetaRI then propagates the signal to downstream targets (Chen, F. and Weinberg. R. A. (1995) PNA892: 1565-1569; Wrana, J. L. <i>et al.</i> (1994) Nature 370: 341. The effect of TGF betas on signaling in the presence of an anti-TGF beta antibody, can be assayed by treating Primary BAECs transfected with a construct called p3TP-Lux, containing a TGF-beta responsive promoter fused to a reporter gene, and measuring luciferase gene expression (Wrana <i>et al.</i>, 1994, Nature 370: 341-347). Useful for control of fibrosis, immune, and inflammatory disease.
Latent TGF beta binding protein II GeneSeq Accession Y70552	Members of the TGF-beta family of proteins initiate cell signaling by binding to heteromeric receptor complexes of type I (TbetaRI) and type II (TbetaRII) serine/threonine kinase receptors (reviewed by Massague, J. <i>et al.</i> (1994) Trends

WO0012551 Q14767/LTBP2_HUMAN Latent-transforming growth factor beta-binding protein 2 (SEQ ID NO:193)	Cell Biol. 4: 172 178; Miyazono, K. <i>et al.</i> (1994) Adv. Immunol. 55: 181-220). Activation of this heteromeric receptor complex occurs when TGF-beta. binds to TbetaRII, which then recruits and phosphorylates TbetaRI. Activated TbetaRI then propagates the signal to downstream targets (Chen, F. and Weinberg. R. A. (1995) PNA892: 1565-1569; Wrana, J. L. <i>et al.</i> (1994) Nature 370: 341. The effect of TGF betas on signaling in the presence of a TGF beta binding protein, can be assayed by treating Primary BAECs transfected with a construct called p3TP-Lux, containing a TGF-beta responsive promoter fused to a reporter gene, and measuring luciferase gene expression (Wrana <i>et al.</i>, 1994, Nature 370: 341-347). Useful for inhibiting tissue or tumor growth.
MP52 GeneSeq Accession W36100 WO9741250 P43026/GDF5_HUMAN Growth/differentiation factor 5 (SEQ ID NO:194)	Members of the TGF-beta family of proteins initiate cell signaling by binding to heteromeric receptor complexes of type I (TbetaRI) and type II (TbetaRII) serine/threonine kinase receptors (reviewed by Massague, J. <i>et al.</i> (1994) Trends Cell Biol. 4: 172 178; Miyazono, K. <i>et al.</i> (1994) Adv. Immunol. 55: 181-220). Activation of this heteromeric receptor complex occurs when TGF-beta. binds to TbetaRII, which then recruits and phosphorylates TbetaRI. Activated TbetaRI then propagates the signal to downstream targets (Chen, F. and Weinberg. R. A. (1995) PNA892: 1565-1569; Wrana, J. L. <i>et al.</i> (1994) Nature 370: 341. The effect of TGF betas on signaling can be assayed by treating Primary BAECs transfected with a construct called p3TP-Lux, containing a TGF- beta responsive promoter fused to a reporter gene, and measuring luciferase gene expression (Wrana <i>et al.</i>, 1994, Nature 370: 341-347). Bone formation or Regeneration Abnormalities
b57 Protein GeneSeq Accession W69293 WO9837195 SEQ ID NO:2 of WO9837195 (SEQ ID NO:195)	BMPs are involved in the induction of bone formation. Specific antagonists are useful is preventing this activity from occurring. BMP activity, in the presence of b57 protein, can be determined using the following assays known in the art: Nat Genet. 2001 Jan; 27(1): 84-8; Eur J Biochem 1996 Apr 1; 237(1): 295-302; J Biol Chem, Vol. 274, Issue 16, 1089- 10902, Apr. 16, 1999; and Hogan, B. L. M. (1996) Genes Deve. 10, 1580- 1594. BMP Antagonist useful for Osteosarcoma, abnormal bone growth.
Resistin GeneSeq Accession W69293 WO0064920 Q9HD89/RETN_HUMAN Resistin (isoform 1) (SEQ ID NO:196)	This gene belongs to the family defined by mouse FIZZI and FIZZ3/Resistin genes. The characteristic feature of this family is the C-terminal stretch of 10 cys residues with identical spacing. The mouse homolog of this protein is secreted by adipocytes, may be the hormone potentially linking obesity to type II diabetes. Ability of resistin to influence type II diabetes can be determined using assays known in the art: Pontoglio <i>et al.</i>, J Clin Invest 1998 May 15; 101(10): 2215-22. Type II diabetes and Syndrome X.
Galectin-4 GeneSeq Accession W11841 WO9703190 P56470/LEG4_HUMAN Galectin-4 (SEQ ID NO:197)	Galectins are a family of carbohydrate-binding proteins characterized by an affinity for beta- galactoside containing glycoconjugates. Ability of Galectin-4 polypeptides to bind lactose can be determined using assays known in the art: Wada, <i>et al.</i>, J Biol Chem 1997 Feb 28; 272(9): 6078- 86. Lactose intolerance.
APM-I; ACRP- 30; Famoxin GeneSeq Accession Y71035 W00026363 Q15848/ADIPO_HUMAN Adiponectin	ACPR30 gene is exclusively expressed in adipose tissue. ACRP30 is thought to increase fatty acid oxidation by muscle tissue. Ability of ACRP30 polypeptides to influence obesity and fat oxidation can be determined using assays known in the art: Fruebis <i>et al.</i>, Proc Nat'l Acad Sci USA 2001 Feb 13; 98(4): 2005-10. Obesity, Metabolic disorders, Lipid Metabolism; Hormone Secretion.

(SEQ ID NO:198)	
ACRP-30 Homologue; Complement Component Clq C GeneSeq Accession B30234 WO0063376 P02747/C1QC_HUMAN Complement C1q subcomponent subunit C	ACPR30 gene is exclusively expressed in adipose tissue. ACRP30 is thought to increase fatty acid oxidation by muscle tissue. Ability of ACRP30 homologue polypeptides to influence obesity and fat oxidation can be determined using assays known in the art: Fruebis <i>et al.</i> , Proc Nat'l Acad Sci USA 2001 Feb 13; 98(4): 2005-10. Obesity, Metabolic disorders, Lipid Metabolism; Hormone Secretion.
(SEQ ID NO:199)	
Calpain-10a GeneSeq Accession Y79567 WO0023603 Q9HC96/CAN10_HUMAN Calpain-10 (Isoform A)	Calpain is believed to play a role in insulin secretion and insulin activity, and therefore may be useful in the treatment of type II diabetes. Ability of Calpain-10 to influence type II diabetes can be determined using assays known in the art: Pontoglio <i>et al.</i> , J Clin Invest 1998 May 15; 101(10): 2215-22. Diabetes mellitus; Regulation of Insulin secretory response; Insulin mediated glucose transport disorders.
(SEQ ID NO:200)	
Calpain-10b GeneSeq Accession Y79568 WO0023603 Q9HC96- 2/CAN10_HUMAN Isoform B of Calpain-10	Calpain is believed to play a role in insulin secretion and insulin activity, and therefore may be useful in the treatment of type II diabetes. Ability of Calpain-10 to influence type II diabetes can be determined using assays known in the art: Pontoglio <i>et al.</i> , J Clin Invest 1998 May 15; 101(10): 2215-22. Diabetes mellitus; Regulation of Insulin secretory response; Insulin mediated glucose transport disorders.
(SEQ ID NO:201)	
Calpain-10c GeneSeq Accession Y79569 WO0023603 Q9HC96- 3/CAN10_HUMAN Isoform C of Calpain-10	Calpain is believed to play a role in insulin secretion and insulin activity, and therefore may be useful in the treatment of type II diabetes. Ability of Calpain-10 to influence type II diabetes can be determined using assays known in the art: Pontoglio <i>et al.</i> , J Clin Invest 1998 May 15; 101(10): 2215-22. Diabetes mellitus; Regulation of Insulin secretory response; Insulin mediated glucose transport disorders.
(SEQ ID NO:202)	
PDGF-D GeneSeq Accession Y71130 WO0027879 Q9GZP0/PDGFD_HUMAN Platelet-derived growth factor D (isoform 1)	Vascular Endothelial Growth Factor. Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266). Wound Healing; Atherosclerosis.
(SEQ ID NO:203)	
FasL GeneSeq Accession Y28594 WO9936079 P48023/TNFL6_HUMAN	Activities associated with apoptosis and immune system functions. Activity can be determined using Apoptosis assays known in the art: Walczak <i>et al.</i> (1996) EMBOJ 16: 5386-5397. Apoptosis-related disorders; Auto-immune disorders; Graft v-Host disorders.

Tumor necrosis factor ligand superfamily member 6 (isoform 1) (SEQ ID NO:204)	
Chondro modulin-like protein GeneSeq Accession Y71262 WO0029579 SEQ ID NO:2 from WO0029579 (SEQ ID NO:370)	Chondromodulin proteins are cartilage proteins thought to confer resistance to angiogenesis, and thus are useful as anti-angiogenic agents that may have utility in combating cancer. Ability of Chondromodulin-like protein to inhibit vascularization can be determined using assays known in the art: Hirakie <i>et al.</i>, J Biol Chem 1997 Dec 19; 272(51): 32419-26. Antiangiogenic agent; Osteoblast proliferation stimulator; prevents vascularization of cartilage tissue; Useful to treat cancer.
Patched GeneSeq Accession W72969 US Pat. No. 5,837,538 Q13635/PTC1_HUMAN Protein patched homolog 1 (isoform L) (SEQ ID NO:205)	Patched is a tumour-suppressor receptor for Sonic hedgehog (shh), which is a protein that controls developmental patterning and growth. Ability of soluble Patched to bind to and inhibit the activities of shh can be determined using assays known in the art: Stone <i>et al.</i>, Nature 1996 Nov 14; 384(6605): 129-34. Receptor for Hedgehog cellular proliferation signaling molecule. This receptor is useful as a means of preventing cellular proliferation via the shh signaling pathway, thus useful for cancers.
Patched-2 GeneSeq Accession Y43261 WO9953058 Q9Y6C5/PTC2_HUMAN Protein patched homolog 2 (isoform 1) (SEQ ID NO:206)	Patched is a tumour-suppressor receptor for Sonic hedgehog (shh), which is a protein that controls developmental patterning and growth. Ability of soluble Patched to bind to and inhibit the activities of shh can be determined using assays known in the art: Stone <i>et al.</i>, Nature 1996 Nov 14; 384(6605): 129-34. Receptor for Hedgehog cellular proliferation signaling molecule. This receptor is useful as a means of preventing cellular proliferation via the shh signaling pathway, thus useful for cancers.
Maspin; Protease Inhibitor 5 GeneSeq Accession R50938 WO9405804 P36952/SPB5_HUMAN Serpin B5 (isoform 1) (SEQ ID NO:207)	Maspin is a member of the serpin family of serine protease inhibitors that is thought to suppress tumor metastasis. The inhibitory effects of Maspin and other protease inhibitors can be assayed using methods known in the art such as a labeled protease substrate, for example, Universal Protease Substrate (casein, resorufin-labeled): Roche Molecular Biochemicals, Cat. No. 1080733. Tumor suppressor which is down-regulated in breast cancers. The maspin protein has tumour suppressing and invasion suppressing activity.
Endostatin GeneSeq Accession B28399 WO0064946 P39060/COIA1_HUMAN Collagen alpha-1(XVIII) chain (isoform 1)	Endostatin is believed to inhibit effects of capillary endothelial cell proliferation. The inhibitory effects of endostatin can be assayed using assays disclosed by Cao <i>et al.</i> (1996) J. Biol. Chem. 271 29461-29467. Anti-angiogenic activity. Useful in the prevention and/ or treatment of cancers.

(SEQ ID NO:208)	
aFGF; FGF-1 GeneSeq Accession P94037 EP298723 P05230/FGF1_HUMAN Fibroblast growth factor 1 (isoform 1)	Fibroblast Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266); Examples 23 and 39 disclosed herein. Promotion of growth and proliferation of cells, such as epithelial cells and keratinocytes. Antagonists may be useful as anti-cancer agents. Diabetes, Metabolic Disease, Obesity.
(SEQ ID NO:209)	
bFGF; FGF-2 GeneSeq Accession R06685 FR2642086 P09038/FGF2_HUMAN Fibroblast growth factor 2 (isoform 1)	Fibroblast Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266); Examples 23 and 39 disclosed herein. Promotion of growth and proliferation of cells, such as epithelial cells and keratinocytes. Antagonists may be useful as anti-cancer agents.
(SEQ ID NO:210)	
FGF-3; INT-2 GeneSeq Accession R07824 WO9503831 P11487/FGF3_HUMAN Fibroblast growth factor 3	Fibroblast Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266); Examples 23 and 39 disclosed herein. Promotion of growth and proliferation of cells, such as epithelial cells and keratinocytes. Antagonists may be useful as anti-cancer agents.
(SEQ ID NO:211)	
FGF-4; HST-1; HBGF-4 GeneSeq Accession R07825 WO9503831 P08620/FGF4_HUMAN Fibroblast growth factor 4 (isoform 1)	Fibroblast Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266); Examples 23 and 39 disclosed herein. Promotion of growth and proliferation of cells, such as epithelial cells and keratinocytes. Antagonists may be useful as anti-cancer agents.
(SEQ ID NO:212)	
FGF-5 GeneSeq Accession W22600 WO9730155 P12034/FGF5_HUMAN Fibroblast growth factor 5 (isoform long)	Fibroblast Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266); Examples 23 and 39 disclosed herein. Promotion of growth and proliferation of cells, such as epithelial cells and keratinocytes. Antagonists may be useful as anti-cancer agents.
(SEQ ID NO:213)	
FGF-6; Heparin binding secreted transforming factor-2 GeneSeq Accession R58555 EP613946 P10767/FGF6_HUMAN Fibroblast growth factor 6	Fibroblast Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266); Examples 23 and 39 disclosed herein. Promotion of growth and proliferation of cells, such as epithelial cells and keratinocytes. Antagonists may be useful as anti-cancer agents.

(SEQ ID NO:214)	
FGF-8 GeneSeq Accession R80783 WO9524928 P55075/FGF8_HUMAN Fibroblast growth factor 8 (isoform 8E)	Fibroblast Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266); Examples 23 and 39 disclosed herein. Promotion of growth and proliferation of cells, such as epithelial cells and keratinocytes. Antagonists may be useful as anti-cancer agents.
(SEQ ID NO:215)	
FGF-9; Gila activating factor GeneSeq Accession R70822 WO9503831 P31371/FGF9_HUMAN Fibroblast growth factor 9	Fibroblast Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266); Examples 23 and 39 disclosed herein. Promotion of growth and proliferation of cells, such as epithelial cells and keratinocytes. Hair growth. Antagonists may be useful as anti-cancer agents.
(SEQ ID NO:216)	
FGF-12; Fibroblast growth factor homologous factor-1 GeneSeq Accession W06309 WO9635708 P61328/FGF12_HUMAN Fibroblast growth factor 12 (isoform 1)	Fibroblast Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266); Examples 23 and 39 disclosed herein. Promotion of growth and proliferation of cells, such as epithelial cells and keratinocytes. Antagonists may be useful as anti-cancer agents.
(SEQ ID NO:217)	
FGF-19 GeneSeq Accession Y08582 WO9927100 O95750/FGF19_HUMAN Fibroblast growth factor 19	Fibroblast Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266); Examples 23 and 39 disclosed herein. Promotion of growth and proliferation of cells, such as epithelial cells and keratinocytes. Chronic liver disease. Primary biliary cirrhosis. Bile acid-induced liver damage. Antagonists may be useful as anti-cancer agents.
(SEQ ID NO:218)	
FGF-16 GeneSeq Accession Y05474 WO9918128 O43320/FGF16_HUMAN Fibroblast growth factor 16	Fibroblast Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266); Examples 23 and 39 disclosed herein. Promotion of growth and proliferation of cells, such as epithelial cells and keratinocytes. Antagonists may be useful as anti-cancer agents.
(SEQ ID NO:219)	
FGF-18 GeneSeq Accession Y08590 WO9927100 O76093/FGF18_HUMAN Fibroblast growth factor 18	Fibroblast Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266); Examples 23 and 39 disclosed herein. Promotion of growth and proliferation of cells, such as epithelial cells and keratinocytes. Antagonists may be useful as anti-cancer agents.
(SEQ ID NO:220)	

flt-3 ligand GeneSeq Accession R67541 EP627487 P49771 FLT3L_HUMAN Fms-related tyrosine kinase 3 ligand (isoform 1) (SEQ ID NO:221)	Stem Cell Progenitor Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Promotion of immune cell growth and/or differentiation.
VEGF-110 GeneSeq Accession Y69417 WO0013702 SEQ ID NO:11 from WO0013702 (SEQ ID NO:222)	Promotes the growth and/or proliferation of endothelial cells. VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
VEGF-121 GeneSeq Accession B50432 WO0071713 SEQ ID NO:2 from WO0071713 (SEQ ID NO:223)	Promotes the growth and/or proliferation of endothelial cells. VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
VEGF-138 GeneSeq Accession Y43483 WO9940197 SEQ ID NO:4 of WO99/40197 (SEQ ID NO:371)	Promotes the growth and/or proliferation of endothelial cells. VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
VEGF-145 GeneSeq Accession Y69413 WO0013702 SEQ ID NO:4 from WO0013702 (SEQ ID NO:224)	Promotes the growth and/or proliferation of endothelial cells. VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
VEGF-162 GeneSeq Accession Y43484 WO9940197 SEQ ID NO:8 of WO99/40197 (SEQ ID NO:372)	Promotes the growth and/or proliferation of endothelial cells. VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
VEGF-165 GeneSeq Accession Y69414 WO0013702	Promotes the growth and/or proliferation of endothelial cells. VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of

SEQ ID NO:6 from WO0013702 (SEQ ID NO:225)	cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
VEGF-182 GeneSeq Accession Y43483 WO9940197 SEQ ID NO:6 of WO99/40197 (SEQ ID NO:373)	Promotes the growth and/or proliferation of endothelial cells. VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
VEGF-189 GeneSeq Accession Y69415 WO0013702 SEQ ID NO:8 from WO0013702 (SEQ ID NO:226)	Promotes the growth and/or proliferation of endothelial cells. VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
VEGF-206 GeneSeq Accession Y69416 WO0013702 SEQ ID NO:10 from WO0013702 (SEQ ID NO:227)	Promotes the growth and/or proliferation of endothelial cells. VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
VEGF-D GeneSeq Accession W53240 WO9807832 O43915/VEGFD_HUMAN Vascular endothelial growth factor D (SEQ ID NO:374)	Promotes the growth and/or proliferation of endothelial cells. VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
VEGF-E; VEGF-X GeneSeq Accession Y33679 WO9947677 SEQ ID NO:2 from WO9947677 (SEQ ID NO:228)	Promotes the growth and/or proliferation of endothelial cells. VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
VEGF Receptor; KDR; flk-1 GeneSeq Accession W69679 WO9831794 P35968/VGFR2_HUMAN Vascular endothelial growth factor receptor 2	Receptor for VEGF polypeptides VEGF activity, in the presence of flk-1 polypeptides, can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. VEGF Receptor. Fusion protein with the extracellular domain is useful as an anti-angiogenic agent. Antagonists may be useful in the promotion of angiogenesis.

(isoform 1)	
(SEQ ID NO:229)	
Soluble VEGF Receptor GeneSeq Accession W47037 US Pat. No. 5,712,380	
sVEGF-RI (Figure 3) of US Pat. No. 5,712,380	
(SEQ ID NO:442)	
sVEGF-RII (Figure 11) of US Pat. No. 5,712,380	Receptor for VEGF polypeptides VEGF activity, in the presence of VEGF Receptor polypeptides, can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. VEGF Receptor. Fusion protein with the extracellular domain is useful as an anti-angiogenic agent.
(SEQ ID NO:443)	Antagonists may be useful in the promotion of angiogenesis.
sVEGF-RTMI (Figure 15) of US Pat. No. 5,712,380	
(SEQ ID NO:444)	
sVEGF-RTMII (Figure 13) of US Pat. No. 5,712,380	
(SEQ ID NO:445)	
flt-1 GeneSeq Accession Y70751 WO0021560	
P17948/VGFR1_HUMAN Vascular endothelial growth factor receptor 1 (isoform 1)	Receptor for VEGF polypeptides VEGF activity, in the presence of flt-1 polypeptides, can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. VEGF Receptor. Fusion protein with the extracellular domain is useful as an anti-angiogenic agent.
(SEQ ID NO:230)	Antagonists may be useful in the promotion of angiogenesis.
VEGF R-3; flt-4 GeneSeq Accession B29047 WO0058511	
P35916/VGFR3_HUMAN Vascular endothelial growth factor receptor 3 (isoform 1)	Receptor for VEGF polypeptides VEGF activity, in the presence of flt-4 polypeptides, can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. VEGF Receptor. Fusion protein with the extracellular domain is useful as an anti-angiogenic agent.
(SEQ ID NO:231)	Antagonists may be useful in the promotion of angiogenesis.
Neuropilin-1 GeneSeq Accession Y06319 WO9929858	
O14786/NRP1_HUMAN Neuropilin-1 (isoform 1)	Vascular Endothelial Growth Factor VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.

(SEQ ID NO:232)	
Neuropilin-2 GeneSeq Accession Y03618 WO9929858	Vascular Endothelial Growth Factor VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
O60462/NRP2_HUMAN Neuropilin-2 (isoform A22)	
(SEQ ID NO:233)	
Human fast twitch skeletal muscle troponin C GeneSeq Accession W22597 W09730085	Troponins are contractile proteins that are thought to inhibit angiogenesis. High levels may contribute to the difficulty encountered in revascularizing the ischemic myocardium after cardiovascular injury. Ability of soluble Troponins to inhibit angiogenesis can be determined using assays known in the art.: Proc Natl Acad Sci USA 1999 Mar 16; 96(6): 2645-50. Anti-angiogenesis
P02585/TNNC2_HUMAN Troponin C, skeletal muscle	
(SEQ ID NO:234)	
Human fast twitch skeletal muscle troponin I GeneSeq Accession W18054 W09730085	Troponins are contractile proteins that are thought to inhibit angiogenesis. High levels may contribute to the difficulty encountered in revascularizing the ischemic myocardium after cardiovascular injury. Ability of soluble Troponins to inhibit angiogenesis can be determined using assays known in the art. Proc Natl Acad Sci USA 1999 Mar 16; 96(6): 2645-50. Anti-angiogenesis
P48788/TNNI2_HUMAN Troponin I, fast skeletal muscle (isoform 1)	
(SEQ ID NO:235)	
Human fast twitch skeletal muscle troponin T GeneSeq Accession W22599 W09730085	Troponins are contractile proteins that are thought to inhibit angiogenesis. High levels may contribute to the difficulty encountered in revascularizing the ischemic myocardium after cardiovascular injury. Ability of soluble Troponins to inhibit angiogenesis can be determined using assays known in the art.: Proc Natl Acad Sci USA 1999 Mar 16; 96(6): 2645-50. Anti-angiogenesis
SEQ ID NO:3 of WO9730085	
(SEQ ID NO:236)	
fragment. myofibrillar protein troponin I GeneSeq Accession W18053 W09719955	Troponins are contractile proteins that are thought to inhibit angiogenesis. High levels may contribute to the difficulty encountered in revascularizing the ischemic myocardium after cardiovascular injury. Ability of soluble Troponins to inhibit angiogenesis can be determined using assays known in the art.: Proc Natl Acad Sci USA 1999 Mar 16; 96(6): 2645-50. Anti-angiogenesis
SEQ ID NO:3 of WO9719955	
(SEQ ID NO:237)	
myofibrillar protein troponin I GeneSeq Accession W18054	Troponins are contractile proteins that are thought to inhibit angiogenesis. High levels may contribute to the difficulty encountered in revascularizing the ischemic myocardium after cardiovascular injury. Ability of soluble Troponins to inhibit

WO9719955	angiogenesis can be determined using assays known in the art.: Proc Natl Acad Sci USA 1999 Mar 16; 96(6): 2645-50. Anti-angiogenesis
SEQ ID NO:3 of WO9719955	
(SEQ ID NO:237)	
Troponin peptides GeneSeq Accessions Y29581, Y29582, Y29583, Y29584, Y29585, and Y29586 WO9933874	Troponins are contractile proteins that are thought to inhibit angiogenesis. High levels may contribute to the difficulty encountered in revascularizing the ischemic myocardium after cardiovascular injury. Ability of soluble Troponins to inhibit angiogenesis can be determined using assays known in the art.: Proc Natl Acad Sci USA 1999 Mar 16; 96(6): 2645-50. Anti-angiogenesis
Wildtype troponins provided as:	
Human fast twitch skeletal muscle troponin C GeneSeq Accession W22597 WO9730085	
P02585/TNNC2_HUMAN Troponin C, skeletal muscle	
(SEQ ID NO:234)	
Human fast twitch skeletal muscle troponin I GeneSeq Accession W18054 WO9730085	
P48788/TNNI2_HUMAN Troponin I, fast skeletal muscle (isoform 1)	
(SEQ ID NO:235)	
Human fast twitch skeletal muscle troponin T GeneSeq Accession W22599 WO9730085	
SEQ ID NO:3 of WO9730085	
(SEQ ID NO:236)	
fragment. myofibrillar protein troponin I GeneSeq Accession W18053 WO9719955	

SEQ ID NO:3 of WO9719955 (SEQ ID NO:237) Human fast twitch skeletal muscle Troponin subunit C GeneSeq Accession B00134 WO0054770 SEQ ID NO:1 of WO0054770 (SEQ ID NO:375) Human fast twitch skeletal muscle Troponin subunit I Protein GeneSeq Accession B00135 WO0054770 SEQ ID NO:2 of WO0054770 (SEQ ID NO:376) Human fast twitch skeletal muscle Troponin subunit T GeneSeq Accession B00136 WO0054770 SEQ ID NO:3 of WO0054770 (SEQ ID NO:377)	
Human fast twitch skeletal muscle Troponin subunit C GeneSeq Accession B00134 WO0054770 SEQ ID NO:1 of WO0054770 (SEQ ID NO:375)	Troponins are contractile proteins that are thought to inhibit angiogenesis. High levels may contribute to the difficulty encountered in revascularizing the ischemic myocardium after cardiovascular injury. Ability of soluble Troponins to inhibit angiogenesis can be determined using assays known in the art.: Proc Natl Acad Sci USA 1999 Mar 16; 96(6): 2645-50. Anti-angiogenesis
Human fast twitch skeletal muscle Troponin subunit I Protein GeneSeq Accession B00135 WO0054770 SEQ ID NO:2 of WO0054770	Troponins are contractile proteins that are thought to inhibit angiogenesis. High levels may contribute to the difficulty encountered in revascularizing the ischemic myocardium after cardiovascular injury. Ability of soluble Troponins to inhibit angiogenesis can be determined using assays known in the art.: Proc Natl Acad Sci USA 1999 Mar 16; 96(6): 2645-50. Anti-angiogenesis

(SEQ ID NO:376)	
Human fast twitch skeletal muscle Troponin subunit T GeneSeq Accession B00136 WO0054770 SEQ ID NO:3 of WO0054770	Troponins are contractile proteins that are thought to inhibit angiogenesis. High levels may contribute to the difficulty encountered in revascularizing the ischemic myocardium after cardiovascular injury. Ability of soluble Troponins to inhibit angiogenesis can be determined using assays known in the art. Proc Natl Acad Sci USA 1999 Mar 16; 96(6): 2645-50. Anti-angiogenesis
(SEQ ID NO:377)	
Activator Inhibitor-1; PAI-1 GeneSeq Accession R08411 WO9013648 P05121/PAI1_HUMAN Plasminogen activator inhibitor 1 (isoform 1)	PAIs are believed to play a role in cancer, and cardiovascular disease and blood-clotting disorders. Methods that measure plasminogen activator inhibitor (PAI) activity are known in the art, for example, assay the ability of PAI to inhibit tissue plasminogen activator (tPA) or urokinase (uPA): J Biochem Biophys Methods 2000 Sep 11; 45(2): 127-40, Breast Cancer Res Treat 1996; 41(2): 141-6. Methods that measure anti- angiogenesis activity are known in the art, for example, Proc Natl Acad Sci USA 1999 Mar 16; 96(6): 2645-50. Anti-angiogenesis; blood-clotting disorders.
(SEQ ID NO:238)	
Plasminogen Activator Inhibitor-2; PAI-2 GeneSeq Accession P94160 DE3722673 P05120/PAI2_HUMAN Plasminogen activator inhibitor 2	PAIs are believed to play a role in cancer, and cardiovascular disease and blood-clotting disorders. Methods that measure plasminogen activator inhibitor (PAI) activity are known in the art, for example, assay the ability of PAI to inhibit tissue plasminogen activator (tPA) or urokinase (uPA): J Biochem Biophys Methods 2000 Sep 11; 45(2): 127-40, Breast Cancer Res Treat 1996; 41(2): 141-6. Methods that measure anti- angiogenesis activity are known in the art, for example, Proc Natl Acad Sci USA 1999 Mar 16; 96(6): 2645-50. Anti-angiogenesis; blood-clotting disorders.
(SEQ ID NO:239)	
Activator Inhibitor-2; PAI-2 GeneSeq Accession R10921 WO9102057 P05120/PAI2_HUMAN Plasminogen activator inhibitor 2	PAIs are believed to play a role in cancer, and cardiovascular disease and blood-clotting disorders. Methods that measure plasminogen activator inhibitor (PAI) activity are known in the art, for example, assay the ability of PAI to inhibit tissue plasminogen activator (tPA) or urokinase (uPA): J Biochem Biophys Methods 2000 Sep 11; 45(2): 127-40, Breast Cancer Res Treat 1996; 41(2): 141-6. Methods that measure anti- angiogenesis activity are known in the art, for example, Proc Natl Acad Sci USA 1999 Mar 16; 96(6): 2645-50. Anti-angiogenesis; blood-clotting disorders.
(SEQ ID NO:239)	
Human PAI-1 mutants GeneSeq Accessions R11755, R11756, R11757, R11758, R11759, R11760, R11761, R11762 and R11763 WO9105048 Wildtype PAI-1 is provided as P05121/PAI1_HUMAN Plasminogen activator inhibitor 1 (isoform 1)	PAIs are believed to play a role in cancer, and cardiovascular disease and blood-clotting disorders. Methods that measure plasminogen activator inhibitor (PAI) activity are known in the art, for example, assay the ability of PAI to inhibit tissue plasminogen activator (tPA) or urokinase (uPA): J Biochem Biophys Methods 2000 Sep 11; 45(2): 127-40, Breast Cancer Res Treat 1996; 41(2): 141-6. Methods that measure anti- angiogenesis activity are known in the art, for example, Proc Natl Acad Sci USA 1999 Mar 16; 96(6): 2645-50. Anti-angiogenesis; blood-clotting disorders.

(SEQ ID NO:238)	
CXCR3; CXC GeneSeq Accession Y79372 WO0018431 P49682 CXCR3_HUMAN C-X-C chemokine receptor type 3 (isoform 1) (SEQ ID NO:240)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Soluble CXCR3 polypeptides may be useful for inhibiting chemokine activities and viral infection.
Modified Rantes GeneSeq Accession W38129 WO9737005 Wildtype Rantes provided herein as P13501/CCL5_HUMAN C- C motif chemokine 5 (SEQ ID NO:241)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders.
RANTES GeneSeq Accession Y05299 EP905240 P13501/CCL5_HUMAN C- C motif chemokine 5 (SEQ ID NO:241)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders.
MCP-1a GeneSeq Accession R73914 WO9509232 MCP-1 provided as P13500/CCL2_HUMAN C- C motif chemokine 2 (SEQ ID NO:337)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders.
MCP-1b GeneSeq Accession Y26176 WO9929728 MCP-1 provided as P13500/CCL2_HUMAN C- C motif chemokine 2 (SEQ ID NO:337)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders.

MCP-I receptor GeneSeq Accession R79165 WO9519436 MCP-1A SEQ ID NO:2 of WO9519436 (SEQ ID NO:446) MCP-1B SEQ ID NO:4 of WO9519436 (SEQ ID NO:447)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Soluble MCP-1 Receptor polypeptides may be useful for inhibiting chemokine activities and viral infection.
MCP-3 GeneSeq Accession R73915 W09509232 P80098/CCL7_HUMAN C- C motif chemokine 7 (SEQ ID NO:336)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders.
MCP-4 receptor GeneSeq Accession W56689 WO9809171 SEQ ID NO:2 of WO9809171 (SEQ ID NO:378)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Soluble MCP-4 Receptor polypeptides may be useful for inhibiting chemokine activities and viral infection.
RANTES receptor GeneSeq Accession W29588 US Pat. No. 5,652,133 SEQ ID NO:2 of US Pat. No. 5,652,133 (SEQ ID NO:379)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Soluble RANTES Receptor polypeptides may be useful for inhibiting chemokine activities and viral infection.
CCR5 variant GeneSeq Accession W88238 WO9854317 Variants of wildtype CCR5 which has the sequence	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which

of : P51681 CCR5_HUMAN C-C chemokine receptor type 5 (SEQ ID NO:448)	bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Soluble CCR5 polypeptides may be useful for inhibiting chemokine activities and viral infection.
CCR7 GeneSeq Accession B50859 US Pat. No. 6,153,441 P32248 CCR7_HUMAN C-C chemokine receptor type 7 (SEQ ID NO:243)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Soluble CCR7 polypeptides may be useful for inhibiting chemokine activities and viral infection.
CXC3 GeneSeq Accession W23345 WO9727299 P78423 X3CL1_HUMAN Fractalkine (SEQ ID NO:244)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders.
Eotaxin GeneSeq Accession W10099 WO9700960 P51671 CCL11_HUMAN Eotaxin (SEQ ID NO:245)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders.
Neurotactin GeneSeq Accessions Y77537, W34307, Y53259, and, Y77539 US Pat. No. 6,013,257 WO9742224 P78423 X3CL1_HUMAN Fractalkine (SEQ ID NO:244)	Neurotactin may play a role in chemotactic leukocyte migration and brain inflammation processes. Chemotactic leukocyte migration assays are known in the art, for example: J. Immunol. Methods 33, ((1980)); Nature 1997 Jun 5; 387(6633): 611-7. Immune disorders.
Human CKbeta-9 GeneSeq Accession B50860 US Pat. No. 6,153,441	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The

SEQ ID NO:2 of US Pat. No. 6,153,441 (SEQ ID NO:246)	chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders.
Lymphotactin GeneSeq Accession B50052 WO0073320 P47992/XCL1_HUMAN Lymphotactin (SEQ ID NO:247)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders.
MIP-3 alpha GeneSeq Accession W44398 WO9801557 P78556/CCL20_HUMAN C-C motif chemokine 20 (isoform 1) (SEQ ID NO:248)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders.
MIP-3 beta GeneSeq Accession W44399 WO9801557 Q99731/CCL19_HUMAN C-C motif chemokine 19 (SEQ ID NO:249)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders.
MIP-Gamma GeneSeq Accession R70798 WO2006135382 (SEQ ID NO:457)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders.
Stem Cell Inhibitory Factor GeneSeq Accession R11553 WO9104274 SCIF in Table I of WO9104274 (SEQ ID NO:380) SCIF in Table II of	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Hematopoietic growth factors.

WO9104274 (SEQ ID NO:381)	
Thrombopoietin GeneSeq Accession R79905 WO9521920 P40225 TPO_HUMAN Thrombopoietin (isoform 1) (SEQ ID NO:250)	Thrombopoietin is involved in the regulation of the growth and differentiation of megakaryocytes and preceptors thereof. Thrombopoietin (TPO) can be assayed to determine regulation of growth and differentiation of megakaryocytes. Mol Cell Biol 2001 Apr; 21(8): 2659-70; Exp Hematol 2001 Jan; 29(1): 51-8 and within. Hematopoietic growth factors.
c-kit ligand; SCF; Mast cell growth factor; MGF; Fibrosarcoma- derived stem cell factor GeneSeq Accession Y53284, R83978 and R83977 EP992579 and EP676470 P21583 SCF_HUMAN Kit ligand (isoform 1) (SEQ ID NO:251)	C-kit ligand is thought to stimulate the proliferation of mast cells, and is able to augment the proliferation of both myeloid and lymphoid hematopoietic progenitors in bone marrow culture. C-kit ligand is also thought to act synergistically with other cytokines. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Hematopoietic growth factors.
Platelet derived growth factor GeneSeq Accession B48653 WO0066736 PDGF-A P04085/PDGFA_HUMAN Platelet-derived growth factor subunit A (Isoform long) (SEQ ID NO:257) PDGF-B P01127/PDGFB_HUMAN Platelet-derived growth factor subunit B (isoform 1) (SEQ ID NO:258)	Vascular Endothelial Growth Factor VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
Melanoma inhibiting protein GeneSeq Accession R69811 WO9503328 (SEQ ID NO:458)	Melanoma inhibiting protein has melanoma-inhibiting activity and can be used to treat cancer (melanoma, glioblastoma, neuroblastoma, small cell lung cancer, neuroectodermal tumors) or as an immunosuppressant (it inhibits IL-2 or phytohaemagglutinin induced proliferation of peripheral blood lymphocytes. Tumor suppressor activity of melanoma inhibiting protein can be determined using assays known in the art: Matzuk <i>et al.</i> , Nature 1992 Nov 26; 360(6402): 313-9. Cancer; melanoma
Glioma-derived growth factor GeneSeq Accession	Vascular Endothelial Growth Factor VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No.

R08120 EP399816	WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
Platelet derived growth factor precursor A GeneSeq Accession R84759 EP682110 PDGF-A precursor (variant D1) (SEQ ID NO:382) PDGF-A precursor (variant 13-1) (SEQ ID NO:383)	Vascular Endothelial Growth Factor VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
Platelet derived growth factor precursor B GeneSeq Accession R84760 EP682110, Figure 1 or Figure 2 Wildtype PDGF-B provided as: PDGF-B P01127/PDGFB_HUMAN Platelet-derived growth factor subunit B (isoform 1) (SEQ ID NO:258)	Vascular Endothelial Growth Factor VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
Platelet derived growth factor Bvsis GeneSeq Accession P80595 and P80596 EP282317 Figure 1 of EP282317 (SEQ ID NO:384)	Vascular Endothelial Growth Factor VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
Placental Growth Factor GeneSeq Accessions R23059 and R23060 WO9206194 P49763-2/PLGF_HUMAN Isoform PIGF-1 of Placenta growth factor (isoform PIGF-1) (SEQ ID NO:252)	Vascular Endothelial Growth Factor VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer.
Placental Growth Factor-2 GeneSeq Accession Y08289 DE19748734	Vascular Endothelial Growth Factor VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and

P49763-3/PLGF_HUMAN Isoform PIGF-2 of Placenta growth factor (isoform PIGF-2) (SEQ ID NO:253)	may be applicable for cancer.
Thrombopoietin derivative1 GeneSeq Accession Y77244 WO0000612 (e.g. Table 3) Wildtype thrombopoietin provided as: P40225 TPO_HUMAN Thrombopoietin (isoform 1) (SEQ ID NO:250)	Thrombopoietin is involved in the regulation of the growth and differentiation of megakaryocytes and preceptors thereof. Thrombopoietin (TPO) can be assayed to determine regulation of growth and differentiation of megakaryocytes. Mol Cell Biol 2001 Apr; 21(8): 2659-70; Exp Hematol 2001 Jan; 29(1): 51-8 and within. Thrombocytopenia, cancer.
Thrombopoietin derivative2 GeneSeq Accession Y77255 WO0000612 (e.g. Table 3) Wildtype thrombopoietin provided as: P40225 TPO_HUMAN Thrombopoietin (isoform 1) (SEQ ID NO:250)	Thrombopoietin is involved in the regulation of the growth and differentiation of megakaryocytes and preceptors thereof. Thrombopoietin (TPO) can be assayed to determine regulation of growth and differentiation of megakaryocytes. Mol Cell Biol 2001 Apr; 21(8): 2659-70; Exp Hematol 2001 Jan; 29(1): 51-8 and within. Thrombocytopenia, cancer.
Thrombopoietin derivative 3 GeneSeq Accession Y77262 WO0000612 (e.g. Table 3) Wildtype thrombopoietin provided as: P40225 TPO_HUMAN Thrombopoietin (isoform 1) (SEQ ID NO:250)	Thrombopoietin is involved in the regulation of the growth and differentiation of megakaryocytes and preceptors thereof. Thrombopoietin (TPO) can be assayed to determine regulation of growth and differentiation of megakaryocytes. Mol Cell Biol 2001 Apr; 21(8): 2659-70; Exp Hematol 2001 Jan; 29(1): 51-8 and within. Thrombocytopenia, cancer.
Thrombopoietin derivative 4 GeneSeq Accession Y77267 WO0000612 (e.g. Table 3) Wildtype thrombopoietin provided as:	Thrombopoietin is involved in the regulation of the growth and differentiation of megakaryocytes and preceptors thereof. Thrombopoietin (TPO) can be assayed to determine regulation of growth and differentiation of megakaryocytes. Mol Cell Biol 2001 Apr; 21(8): 2659-70; Exp Hematol 2001 Jan; 29(1): 51-8 and within. Thrombocytopenia, cancer.

P40225 TPO_HUMAN Thrombopoietin (isoform 1) (SEQ ID NO:250)	
Thrombopoietin derivative 5 GeneSeq Accession Y77246 WO0000612 (e.g. Table 3) Wildtype thrombopoietin provided as: P40225 TPO_HUMAN Thrombopoietin (isoform 1) (SEQ ID NO:250)	Thrombopoietin is involved in the regulation of the growth and differentiation of megakaryocytes and preceptors thereof. Thrombopoietin (TPO) can be assayed to determine regulation of growth and differentiation of megakaryocytes. Mol Cell Biol 2001 Apr; 21(8): 2659-70; Exp Hematol 2001 Jan; 29(1): 51-8 and within. Thrombocytopenia, cancer.
Thrombopoietin derivative 6 GeneSeq Accession Y77253 WO0000612 (e.g. Table 3) Wildtype thrombopoietin provided as: P40225 TPO_HUMAN Thrombopoietin (isoform 1) (SEQ ID NO:250)	Thrombopoietin is involved in the regulation of the growth and differentiation of megakaryocytes and preceptors thereof. Thrombopoietin (TPO) can be assayed to determine regulation of growth and differentiation of megakaryocytes. Mol Cell Biol 2001 Apr; 21(8): 2659-70; Exp Hematol 2001 Jan; 29(1): 51-8 and within. Thrombocytopenia, cancer.
Thrombopoietin derivative 7 GeneSeq Accession Y77256 WO0000612 (e.g. Table 3) Wildtype thrombopoietin provided as: P40225 TPO_HUMAN Thrombopoietin (isoform 1) (SEQ ID NO:250)	Thrombopoietin is involved in the regulation of the growth and differentiation of megakaryocytes and preceptors thereof. Thrombopoietin (TPO) can be assayed to determine regulation of growth and differentiation of megakaryocytes. Mol Cell Biol 2001 Apr; 21(8): 2659-70; Exp Hematol 2001 Jan; 29(1): 51-8 and within. Thrombocytopenia, cancer.
Fractalkine GeneSeq Accession Y53255 US Pat. No. 6,043,086 P78423/X3CL1_HUMAN Fractalkine (SEQ ID NO:244)	Fractalkine is believed to play a role in chemotactic leukocyte migration and neurological disorders. Fractalkine activity can be determined using Chemotactic leukocyte migration assays known in the art, for example: J. Immunol. Methods 33, ((1980)); Nature 1997 Jun 5; 387(6633): 611-7. Immune disorders.

CXC3 GeneSeq Accession W23345 WO9757599 P78423/X3CL1_HUMAN Fractalkine (SEQ ID NO:244)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders.
CCR7 GeneSeq Accession B50859 US Pat. No. 6,153,441 P32248/CCR7_HUMAN C-C chemokine receptor type 7 (SEQ ID NO:243)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Soluble CCR7 polypeptides may be useful for inhibiting chemokine activities and viral infection.
Nerve Growth Factor-beta GeneSeq Accession R11474 EP414151 P01138/NGF_HUMAN Beta-nerve growth factor (SEQ ID NO:254)	Nerve Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266) Neurological disorders, cancer
Nerve Growth Factor-beta2 GeneSeq Accession W69725 EP859056 Fig 1 of EP859056 (SEQ ID NO:465)	Nerve Growth Factor Proliferation assay using NR6R 3T3 cells (Rizzino 1988 Cancer Res. 48: 4266 Neurological disorders, cancer
Neurotrophin-3 GeneSeq Accession W8889 WO9821234 P20783/NTF3_HUMAN Neurotrophin-3 (isoform 1) (SEQ ID NO:255)	Neurotrophins regulate neuronal cell survival and synaptic plasticity. Trk tyrosine kinase activation assays known in the art can be used to assay for neurotrophin activity, for example, Proc Natl Acad Sci USA 2001 Mar 13; 98(6): 3555-3560. Neurological disorders, cancer
Neurotrophin-4 GeneSeq Accession R47100 WO9325684 P34130/NTF4_HUMAN Neurotrophin-4 (SEQ ID NO:256)	Neurotrophins regulate neuronal cell survival and synaptic plasticity. Trk tyrosine kinase activation assays known in the art can be used to assay for neurotrophin activity, for example, Proc Natl Acad Sci USA 2001 Mar 13; 98(6): 3555-3560. Neurological disorders, cancer
Neurotrophin- 4a	Neurotrophins regulate neuronal cell survival and synaptic plasticity. Trk tyrosine

GeneSeq Accession R47101 WO9325684 Wildtype neurotrophin provided as: P34130/NTF4_HUMAN Neurotrophin-4 (SEQ ID NO:256)	kinase activation assays known in the art can be used to assay for neurotrophin activity, for example, Proc Natl Acad Sci USA 2001 Mar 13; 98(6): 3555-3560. Neurological disorders, cancer
Neurotrophin- 4b GeneSeq Accession R47102 WO9325684 P34130/NTF4_HUMAN Neurotrophin-4 (SEQ ID NO:256)	Neurotrophins regulate neuronal cell survival and synaptic plasticity. tyrosine kinases. Trk tyrosine kinase activation assays known in the art can be used to assay for neurotrophin activity, for example, Proc Natl Acad Sci USA 2001 Mar 13; 98(6): 3555-3560. Neurological disorders, cancer
Neurotrophin- 4c GeneSeq Accession R47103 WO9325684 P34130/NTF4_HUMAN Neurotrophin-4 (SEQ ID NO:256)	Neurotrophins regulate neuronal cell survival and synaptic plasticity. tyrosine kinases. Trk tyrosine kinase activation assays known in the art can be used to assay for neurotrophin activity, for example, Proc Natl Acad Sci USA 2001 Mar 13; 98(6): 3555-3560. Neurological disorders, cancer
Neurotrophin- 4d GeneSeq Accession R47102 WO9325684 P34130/NTF4_HUMAN Neurotrophin-4 (SEQ ID NO:256)	Neurotrophins regulate neuronal cell survival and synaptic plasticity. tyrosine kinases. Trk tyrosine kinase activation assays known in the art can be used to assay for neurotrophin activity, for example, Proc Natl Acad Sci USA 2001 Mar 13; 98(6): 3555-3560. Neurological disorders, cancer
Platelet-Derived Growth Factor A chain GeneSeq Accession R38918 US Pat. No. 5,219,739 P04085/PDGFA_HUMAN Platelet-derived growth factor subunit A (Isoform long) (SEQ ID NO:257)	Vascular Endothelial Growth Factor VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. W00045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Hematopoietic and immune disorders. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer
Platelet-Derived Growth Factor B chain GeneSeq Accession R38919 US Pat. No. 5,219,739 P01127/PDGFB_HUMAN Platelet-derived growth	Vascular Endothelial Growth Factor VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. W00045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Hematopoietic and immune disorders. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer

factor subunit B (isoform 1) (SEQ ID NO:258)	
Stromal Derived Factor- 1 alpha GeneSeq Accession Y39995 WO9948528 P48061-2/SDF1_HUMAN Isoform Alpha of Stromal cell-derived factor 1 (isoform alpha) (SEQ ID NO:259)	Stromal Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266) Hematopoietic, immune disorders, cancer
Stromal Derived Factor- 1 beta GeneSeq Accession R75420 CA2117953 P48061/SDF1_HUMAN Stromal cell-derived factor 1 (isoform beta) (SEQ ID NO:260)	Stromal Growth Factor Proliferation assay using NR6R-3T3 cells (Rizzino 1988 Cancer Res. 48: 4266) Hematopoietic, immune disorders, cancer
Tarc GeneSeq Accession W14917 WO9711969 Q92583/CCL17_HUMAN C-C motif chemokine 17 (SEQ ID NO:261)	Chemotactic for T lymphocytes. May play a role in T-cell development. Thought to bind CCR8 and CCR4 Chemotactic leukocyte migration assays are known in the art, for example: J. Immunol. Methods 33 ((1980)) Antiinflammatory. Immune disorders, cancer
Prolactin GeneSeq Accession R78691 WO9521625 P01236/PRL_HUMAN Prolactin (SEQ ID NO:262)	Prolactin is involved in immune cell proliferation and apoptosis. Immune cell proliferation and suppression of apoptosis by prolactin can be assayed by methods well-known in the art, for example, Buckley, AR and Buckley DJ, Ann N Y Acad Sci 2000; 917: 522-33, and within. Reproductive system disorders, cancer.
Prolactin2 GeneSeq Accession Y31764 US Pat. No. 5,955,346	Prolactin is involved in immune cell proliferation and apoptosis. Immune cell proliferation and suppression of apoptosis by prolactin can be assayed by methods well-known in the art, for example, Buckley, AR and Buckley DJ, Ann N Y Acad Sci 2000; 917: 522-33, and within. Reproductive system disorders, cancer.
Follicle stimulating hormone Alpha subunit GeneSeq Accession Y54160 EP974359 P01215/GLHA_HUMAN Glycoprotein hormones alpha chain (SEQ ID NO:263)	FSH stimulates secretion of interleukin-1 by cells isolated from women in the follicular phase FSH activities can be determined using assays known in the art; J Gend Specif Med 1999 Nov-Dec; 2(6): 30-4; Mol Cell Endocrinol. 1997 Nov 15; 134(2): 109-18. Reproductive system disorders, cancer.

Follicle stimulating hormone Beta subunit GeneSeq Accession Y54161 EP974359 P01225/FSHB_HUMAN Follitropin subunit beta (SEQ ID NO:264)	FSH stimulates secretion of interleukin-1 by cells isolated from women in the follicular phase FSH activities can be determined using assays known in the art; J Gend Specif Med 1999 Nov-Dec; 2(6): 30-4; Mol Cell Endocrinol. 1997 Nov 15; 134(2): 109-18. Reproductive system disorders, cancer.
Substance P (tachykinin) GeneSeq Accession B23027 WO0054053 (SEQ ID NO:385)	Substance P is associated with immunoregulation. Immunoregulation and bone marrow, cell proliferation by substance P can be assayed by methods well-known in the art, for example, Lai <i>et al.</i> Proc Natl Acad Sci USA 2001 Mar 27; 98(7): 3970-5; Jallat-Daloz <i>et al.</i> Allergy Asthma Proc 2001 Jan-Feb; 22(1): 17- 23; Kahler <i>et al.</i> Exp Lung Res 2001 Jan-Feb; 27(1): 25-46; and Adamus MA and Dabrowski ZJ. J Cell Biochem 2001; 81(3)499-506. diabetes mellitus, hypertension, cancer
Oxytocin (Neurophysin I) GeneSeq Accession B24085 and B24086 WO0053755 P01178/NEU1_HUMAN Oxytocin-neurophysin 1 (SEQ ID NO:265)	Oxytocin is involved in the induction of prostaglandin (E2) release as well as an increased amount of calcium release by smooth muscle cells. Oxytocin and prostaglandin E(2) release and Ocytocin (Ca2+) increase can be assayed by methods well-known in the art, for example, Pavan <i>et al.</i>, AM J Obset Gynecol 2000 Jul; 183(1): 76- 82 and Holda <i>et al.</i>, Cell Calcium 1996 Jul; 20(1): 43 51. inflammatory disorders immunologic disorders, cancer
Vasopressin (Neurophysin II) GeneSeq Accession B24085 and B24086 WO0053755 P01185/NEU2_HUMAN Vasopressin-neurophysin 2-copeptin (SEQ ID NO:266)	Vasopressinis believed to have a direct antidiuretic action on the kidney, and it is thought to cause vasoconstriction of the peripheral vessels. Vasopressin activity can be determined using assays known in the art, for example, Endocr Regul 1996 Mar; 30(1): 13-17. inflammatory disorders immunologic disorders, cancer
IL-1 GeneSeq Accession P60326 EP165654 IL-1 alpha P01583 IL1A_HUMAN Interleukin-1 alpha (SEQ ID NO:269) IL-1 beta P01584 IL1B_HUMAN Interleukin-1 beta (SEQ ID NO:267)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinardlio (1989) Cytokine 1, 14-20. inflammatory disorders immunologic disorders, cancer
IL-1 mature GeneSeq Accession R14855 EP456332	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity

(mature truncated form wherein the precursor is cleaved between amino acids 116-117) (SEQ ID NO:386)	can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinardio (1989) Cytokine 1, 14-20. inflammatory disorders immunologic disorders, cancer
IL-1 beta GeneSeq Accession Y08322 WO9922763 P01584 IL1B_HUMAN Interleukin-1 beta (SEQ ID NO:267)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinardio (1989) Cytokine 1, 14-20. inflammatory disorders immunologic disorders, cancer
IL-3 variants GeneSeq Accession P80382, P80383, P80384, and P80381 WO8806161 Variants of wildtype IL-3 which has the sequence: P08700 IL3_HUMAN Interleukin-3 (SEQ ID NO:449)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Kitamura <i>et al.</i> (1989) J Cell Physiol. 140 323-334. inflammatory disorders immunologic disorders, cancer
IL-4 GeneSeq Accession P70615 WO8702990 P05112 IL4_HUMAN Interleukin-4 (isoform 1) (SEQ ID NO:268)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Siegel & Mostowski (1990) J Immunol Methods 132, 287-295. inflammatory disorders immunologic disorders, cancer
IL-4 muteins GeneSeq Accession W52151 W52152 W52153 W52154 W52155 W52156 W52157 W52158 W52159 W52160 W52161 W52162 W52163 W52164 and W52165 WO9747744 Variants of wildtype IL-4 which has the sequence: P05112 IL4_HUMAN Interleukin-4 (isoform 1) (SEQ ID NO:268)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Siegel & Mostowski (1990) J Immunol Methods 132, 287-295. inflammatory disorders immunologic disorders, cancer
IL-1 alpha GeneSeq Accession P90108	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of

EP324447 P01583 IL1A_HUMAN Interleukin-1 alpha (SEQ ID NO:269)	immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinardio (1989) Cytokine 1, 14-20. inflammatory disorders immunologic disorders, cancer
IL-3 variants GeneSeq Accession R38561, R38562, R38563, R38564, R38565, R38566, R38567, R38568, R38569, R38570, R38571, and R38572 WO9307171 Variants of wildtype IL-3 which has the sequence: P08700 IL3_HUMAN Interleukin-3 (SEQ ID NO:449)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Aarden <i>et al</i> (1987) Eur. J. Immunol 17, 1411-16. inflammatory disorders immunologic disorders, cancer
IL-6 GeneSeq Accession R45717 and R45718 WO9402512 P05231 IL6_HUMAN Interleukin-6 (SEQ ID NO:270)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Aarden <i>et al</i> (1987) Eur. J. Immunol 17, 1411-16. inflammatory disorders immunologic disorders, cancer. Obesity. Metabolic Disease. Diabetes.
IL-13 GeneSeq Accession R48624 WO9404680 P35225 IL13_HUMAN Interleukin-13 (SEQ ID NO:271)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Boutelier <i>et al</i> (1995) J. Immunol. Methods 181, 29. inflammatory disorders immunologic disorders, cancer
IL-4 mutein GeneSeq Accession R47182 DE4137333 Variants of wildtype IL-4 which has the sequence: P05112 IL4_HUMAN Interleukin-4 (isoform 1) (SEQ ID NO:268)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Siegel & Mostowski (1990) J Immunol Methods 132, 287-295. inflammatory disorders immunologic disorders, cancer
IL-4 mutein Y124X GeneSeq Accession	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of

R47183 DE4137333 Variants of wildtype IL-4 which has the sequence: P05112/IL4_HUMAN Interleukin-4 (isoform 1) (SEQ ID NO:268))	immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Siegel & Mostowski (1990) J Immunol Methods 132, 287-295. inflammatory disorders immunologic disorders, cancer
IL-4 mutein Y124G GeneSeq Accession R47184 DE4137333 Variants of wildtype IL-4 which has the sequence: P05112/IL4_HUMAN Interleukin-4 (isoform 1) (SEQ ID NO:268)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Siegel & Mostowski (1990) J Immunol Methods 132, 287-295. inflammatory disorders immunologic disorders, cancer
Human Interleukin-10 (precursor) GeneSeq Accession R41664 WO9317698 P22301/IL10_HUMAN Interleukin-10 (precursor form is processed into a truncated mature form) (SEQ ID NO:272)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Thompson-Snipes <i>et al</i> (1991) J. Exp. Med. 173, 507-510. inflammatory disorders immunologic disorders, cancer
Human Interleukin-10 GeneSeq Accession R42642 WO9318783-A SEQ ID NO:3 of WO9318783-A (mature IL-10) (SEQ ID NO:273)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Thompson-Snipes <i>et al</i> (1991) J. Exp. Med. 173, 507-510. inflammatory disorders immunologic disorders, cancer
Human interleukin-1 beta precursor. GeneSeq Accession R42447 EP569042 P01584/IL1B_HUMAN Interleukin-1 beta (SEQ ID NO:274)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinardio (1989) Cytokine 1, 14-20. inflammatory disorders immunologic disorders, cancer

Interleukin- 1alpha GeneSeq Accession R45364 EP578278 P01583 IL1A_HUMAN Interleukin-1 alpha (SEQ ID NO:269)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225. inflammatory disorders immunologic disorders, cancer
Human interleukin-3 variant GeneSeq Accession R22814 JP04063595 Variants of wildtype IL-3 which has the sequence: P08700 IL3_HUMAN Interleukin-3 (SEQ ID NO:449)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; Kitamura <i>et al.</i> (1989) J Cell Physiol. 140 323-334. inflammatory disorders immunologic disorders, cancer
IL-1i fragments GeneSeq Accession R35484 and R35485 EP541920	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinardio (1989) Cytokine 1, 14-20. inflammatory disorders immunologic disorders, cancer
IL-1 inhibitor (IL-Ii) GeneSeq Accession R35486 and R35484 EPS541920	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinardio (1989) inflammatory disorders immunologic disorders, cancer
ICE 22 kD subunit. GeneSeq Accession R33780 EP533350 SEQ ID NO:16 of EP533350 (SEQ ID NO:450)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225. inflammatory disorders immunologic disorders, cancer
ICE 20 kD subunit. GeneSeq Accession R33781 EP533350 SEQ ID NO:17 of EP533350 (SEQ ID NO:451)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225. inflammatory disorders immunologic disorders, cancer

ICE 10 kD subunit GeneSeq Accession R33782 EP533350 SEQ ID NO:18 of EP533350 (SEQ ID NO:452)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225. inflammatory disorders immunologic disorders, cancer
Human Interleukin-10 (precursor) GeneSeq Accession R41664 WO9317698 P22301/IL10_HUMAN Interleukin-10 (precursor form is processed into a truncated mature form) (SEQ ID NO:272)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Thompson-Snipes <i>et al</i> (1991) J. Exp. Med. 173, 507-510. inflammatory disorders immunologic disorders, cancer
Human Interleukin-10 GeneSeq Accession R42642 WO9318783 SEQ ID NO:3 of WO9318783-A (mature IL-10) (SEQ ID NO:273)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Thompson-Snipes <i>et al</i> (1991) J. Exp. Med. 173, 507-510. inflammatory disorders immunologic disorders, cancer
Human Interleukin-1 beta precursor GeneSeq Accession R42447 EP569042 P01584/IL1B_HUMAN Interleukin-1 beta (SEQ ID NO:274)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; Kitamura <i>et al</i> (1989) J Cell Physiol. 140 323-334. inflammatory disorders immunologic disorders, cancer
Human interleukin-6 GeneSeq Accession R49041 WO9403492 P05231/IL6_HUMAN Interleukin-6 (SEQ ID NO:270)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Aarden <i>et al</i> (1987) Eur. J. Immunol 17, 1411-16. inflammatory disorders immunologic disorders, cancer
Mutant Interleukin 6 S176R GeneSeq Accession R54990 WO9411402 S176R variant of wildtype IL-6 which has the	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Aarden <i>et al</i> (1987) Eur. J. Immunol 17, 1411-16.

sequence: P05231/IL6_HUMAN Interleukin-6 (SEQ ID NO:270)	inflammatory disorders immunologic disorders, cancer
Interleukin 6 GeneSeq Accession R55256 JP06145063 P05231/IL6_HUMAN Interleukin-6 (SEQ ID NO:270)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Aarden et al (1987) Eur. J. Immunol 17, 1411-16. inflammatory disorders immunologic disorders, cancer
Interleukin 8 (IL-8) receptor GeneSeq Accession R53932 JP06100595 GenBank: AAA59159.1 (SEQ ID NO:275)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Holmes et al (1991) Science 253, 1278-80. Soluble IL-8 receptor polypep- tides may be useful for inhibiting interleukin activities.
Human interleukin-7 GeneSeq Accession R59919 US Pat. No. 5,328,988 P13232/IL7_HUMAN Interleukin-7 (isoform 1) (SEQ ID NO:276)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Park et al (1990) J. Exp. Med. 171, 1073-79. inflammatory disorders immunologic disorders, cancer
IL-3 containing fusion protein. GeneSeq Accession R79342 and R79344 WO9521254 Fusions of wildtype IL-3 which has the sequence: P08700/IL3_HUMAN Interleukin-3 (SEQ ID NO:449)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; Kitamura et al (1989) J Cell Physiol. 140 323-334. inflammatory disorders immunologic disorders, cancer
IL-3 mutant proteins GeneSeq Accession R79254, R79255, R79256, R79257, R79258, R79259, R79260, R79261, R79262, R79263, R79264, R79265, R79266, R79267, R79268, R79269, R79270, R79271, R79272, R79273, R79274,	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Giri et al (1994) EMBO J. 13 2822-2830. inflammatory disorders immunologic disorders, cancer

R79275, R79276, R79277, R79278, R79279, R79280, R79281, R79282, R79283, R79284, and R79285 ZA9402636 Variants of wildtype IL-3 which has the sequence: P08700 IL3_HUMAN Interleukin-3 (SEQ ID NO:449)	
IL-12 p40 subunit. GeneSeq Accession R63018 AU9466072 P2946 IL12B_HUMAN Interleukin-12 subunit beta (SEQ ID NO:277)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225. inflammatory disorders immunologic disorders, cancer
AGF GeneSeq Accession R64240 WO9429344 Q8NI99/ANGL6_HUMAN Angiopoietin-related protein 6 (SEQ ID NO:278)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225. inflammatory disorders immunologic disorders, cancer
Human interlaukin-12 40 kD subunit GeneSeq Accession R79187 WO9519786 P2946 IL12B_HUMAN Interleukin-12 subunit beta (SEQ ID NO:277)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Hori <i>et al</i> (1987), Blood 70, 1069- 1078. inflammatory disorders immunologic disorders, cancer
Human interleukin-15 receptor from clone P1 GeneSeq Accession R90843 WO9530695 Q13261 IL15RA_HUMAN Interleukin-15 receptor subunit alpha Isoform 1) (SEQ ID NO:453)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Giri <i>et al</i> (1994) EMBO J. 13 2822- 2830. Soluble IL-8 receptor polypep- tides may be useful for inhibiting interleukin activities. Obesity. Metabolic Disease. Diabetes. Enhancing secretion and stability of Interleukin 15
Human interleukin-7 GeneSeq Accession R92796 WO9604306	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity

P13232/IL7_HUMAN Interleukin-7 (isoform 1) (SEQ ID NO:276)	can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Park <i>et al</i> (1990) J. Exp. Med. 171, 1073-79. inflammatory disorders immunologic disorders, cancer
interleukin-9 GeneSeq Accession R92797 WO9604306 P15248/IL9_HUMAN Interleukin-9 (SEQ ID NO:279)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Yang <i>et al</i> (1989) Blood 74, 1880- 84. inflammatory disorders immunologic disorders, cancer
interleukin-3 GeneSeq Accession R92801 WO9604306 P08700/IL3_HUMAN Interleukin-3 (SEQ ID NO:280)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; Kitamura <i>et al</i> (1989) J Cell Physiol. 140 323-334. inflammatory disorders immunologic disorders, cancer
Human interleukin-5 GeneSeq Accession R92802 WO9604306 P05113/IL5_HUMAN Interleukin-5 (SEQ ID NO:281)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; Kitamura <i>et al</i> (1989) J Cell Physiol. 140 323-334. inflammatory disorders immunologic disorders, cancer
Recombinant interleukin-16 GeneSeq Accession W33373 DE19617202 Q14005/IL16_HUMAN Pro-interleukin-16 (isoform 1) (SEQ ID NO:282)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Lim <i>et al</i> (1996) J. Immunol. 156, 2566-70. inflammatory disorders immunologic disorders, cancer
Human IL-16 protein GeneSeq Accession W33234 DE19617202 Q14005/IL16_HUMAN Pro-interleukin-16 (isoform 1) (SEQ ID NO:282)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Lim <i>et al</i> (1996) J. Immunol. 156, 2566-70. inflammatory disorders immunologic disorders, cancer
Thrl 17 human interleukin 9 GeneSeq Accession W27521 WO9708321 P15248/IL9_HUMAN Interleukin-9	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington,

(SEQ ID NO:387)	D.C. 1987, pp. 221-225. inflammatory disorders immunologic disorders, cancer
Metl 17 human interleukin 9 GeneSeq Accession W27522 WO9708321 (SEQ ID NO:388)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Yang et al (1989) Blood 74, 1880- 84. inflammatory disorders immunologic disorders, cancer
Human intracellular IL- 1 receptor antagonist. GeneSeq Accession W77158 EP864585 (e.g. SEQ ID NOs:12 to 19, or 22 to 25 of this publication).	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinarello (1989) Cytokine 1, 14-20. inflammatory disorders immunologic disorders, cancer
Human interleukin-18 protein (IL-18) GeneSeq Accession W77158 EP864585 Q14116/IL18_HUMAN Interleukin-18 (isoform 1) (SEQ ID NO:283)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and USHIO et al (1996) J. Immunol. 156, 4274-79. inflammatory disorders immunologic disorders, cancer
Human interleukin-18 GeneSeq Accession W77077 EP861663 Q14116/IL18_HUMAN Interleukin-18 (isoform 1) (SEQ ID NO:283)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and USHIO et al (1996) J. Immunol. 156, 4274-79. inflammatory disorders immunologic disorders, cancer
Human interleukin 18 derivatives GeneSeq Accessions W77083, W77084, W77085, W77086, W77087, W77088, and W77089 EP861663 Variants of wildtype IL18 which is provided as: Q14116/IL18_HUMAN Interleukin-18 (isoform 1)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Ushio et al (1996) J. Immunol, 156, 4274-79. inflammatory disorders immunologic disorders, cancer

(SEQ ID NO:283) Interleukin-9 (IL-9) mature protein (Thr117 version). GeneSeq Accession W68158 WO9827997 Figure 2 of WO9827997 (SEQ ID NO:389)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Yang et al (1989) Blood 74, 1880- 84. inflammatory disorders immunologic disorders, cancer
IL-9 mature protein variant (Met117 version) GenSeq Accession W68157 WO9827997 Figure 3 of WO9827997 (SEQ ID NO:390)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Yang et al (1989) Blood 74, 1880- 84. inflammatory disorders immunologic disorders, cancer
Human IL-9 receptor protein variant #3. GeneSeq Accession W64058 WO9824904 Wildtype IL-9R is provided as: Q01113/IL9R_HUMAN Interleukin-9 receptor (isoform 1) (SEQ ID NO:303)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Yang et al (1989) Blood 74, 1880- 84. inflammatory disorders immunologic disorders, cancer
Human IL-9 receptor protein variant fragment GenSeq Accession W64060 WO9824904 Wildtype IL-9R is provided as: Q01113/IL9R_HUMAN Interleukin-9 receptor (isoform 1) (SEQ ID NO:303)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Yang et al (1989) Blood 74, 1880- 84. Soluble IL-9 receptor polypep- tides may be useful for inhibiting interleukin activities.
Human IL-9 receptor protein variant #3. GeneSeq Accession W64061 WO9824904 Wildtype IL-9R is provided as: Q01113/IL9R_HUMAN	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Yang et al (1989) Blood 74, 1880- 84. Soluble IL-9 receptor polypep- tides may be useful for inhibiting interleukin activities.

Interleukin-9 receptor (isoform 1)	
(SEQ ID NO:303)	
Human Interleukin-12 p40 protein GeneSeq Accession W51311 WO9817689	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Hori et al (1987), Blood 70, 1069- 1078. inflammatory disorders immunologic disorders, cancer
P2946/IL12B_HUMAN Interleukin-12 subunit beta	
(SEQ ID NO:277)	
Human Interleukin-12 p35 protein GeneSeq Accession W51312 WO9817689	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Hori et al (1987), Blood 70, 1069- 1078. inflammatory disorders immunologic disorders, cancer
P29459/IL12A_HUMAN Interleukin-12 subunit alpha	
(SEQ ID NO:284)	
Human protein with IL-16 activity GeneSeq Accession W63753 DE19649233-	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Lim et al (1996) J. Immunol. 156, 2566-70. inflammatory disorders immunologic disorders, cancer
Human protein with IL-16 activity GeneSeq Accession W59425 DE19649233-	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Lim et al (1996) J. Immunol. 156, 2566-70. inflammatory disorders immunologic disorders, cancer
Human interleukin-15 GeneSeq Accession W53878 US Pat. No. 5,747,024	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Giri et al (1994) EMBO J. 13 2822- 2830. inflammatory disorders immunologic disorders, cancer. Obesity. Metabolic Disease. Diabetes.
P40933/IL15_HUMAN Interleukin-15 (isoform IL15-S48AA)	
(SEQ ID NO:285)	
Human wild- type interleukin-4 (hIL-4) protein GeneSeq	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis

Accession W52149 WO9747744 P05112/IL4_HUMAN Interleukin-4 (isoform 1) (SEQ ID NO:286)	of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Siegel & Mostowski (1990) J Immunol Methods 132, 287-295. inflammatory disorders immunologic disorders, cancer
interleukin-4 muteins GeneSeq Accessions W52150, W52151, W52153, W52154, W52155, W52156, W52157, W52158, W52159, W52160, W52161, W52162, W52163, W52164, W52165, W52166, and W52167 WO9747744 Variants of wildtype IL-4 which has the sequence: P05112/IL4_HUMAN Interleukin-4 (isoform 1) (SEQ ID NO:268)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Siegel & Mostowski (1990) J Immunol Methods 132, 287-295. inflammatory disorders immunologic disorders, cancer
Human interleukin 1 delta GeneSeq Accession Y28408 WO9935268 SEQ ID NO:4 of WO9935268 (SEQ ID NO:287)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinarello (1989) Cytokine 1, 14-20. inflammatory disorders immunologic disorders, cancer
Human interleukin-1 receptor antagonist beta GeneSeq Accession Y24395 WO9935268	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinarello (1989) Cytokine 1, 14-20. inflammatory disorders immunologic disorders, cancer
Human EDIRF II protein sequence GeneSeq Accession Y22199 WO9932632 SEQ ID NO:6 of WO9932632 (SEQ ID NO:391)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225. inflammatory disorders immunologic disorders, cancer

Human EDIRF I protein sequence GeneSeq Accession Y22197 WO9932632 SEQ ID NO:2 of WO9932632 (SEQ ID NO:392)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225. inflammatory disorders immunologic disorders, cancer
Human IL- 1RD10 protein sequence GeneSeq Accession Y14131 WO9919480 SEQ ID NO:20 of WO9919480	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinarello (1989) Cytokine 1, 14-20. Soluble IL-1RD10 receptor polypep- tides may be useful for inhibiting interleukin activites.
Human IL- 1RD9 GeneSeq Accession Y14122 WO9919480 SEQ ID NOS:6, 8, 10 of WO9919480	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinarello (1989) Cytokine 1, 14-20. Soluble IL-1RD10 receptor polypep- tides may be useful for inhibiting interleukin activites.
Human DNAX interleukin- 40 GeneSeq Accession Y09196 WO9919491 SEQ ID NO: 2 or 4 of WO9919491 (SEQ ID NO:454)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225. inflammatory disorders immunologic disorders, cancer
(DIL-40) alternative sequence GeneSeq Accession Y09197 WO9919491 SEQ ID NO: 4 of WO9919491 (SEQ ID NO:455)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225. inflammatory disorders immunologic disorders, cancer
IL-11 GeneSeq Accession R50176 WO9405318 P2080/IL11_HUMAN Interleukin-11 (isoform 1) (SEQ ID NO:288)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Lu et al (1994) J immunol. Methods 173, 19. inflammatory disorders immunologic disorders, cancer
Human adipogenesis inhibitory factor GeneSeq	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of

Accession R43260 EP566410 (also known as IL-11) P2080/IL11_HUMAN Interleukin-11 (isoform 1) (SEQ ID NO:288)	immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225. inflammatory disorders immunologic disorders, cancer
IL-11 GeneSeq Accession W02202 JP08127539 P2080/IL11_HUMAN Interleukin-11 (isoform 1) (SEQ ID NO:288)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Lu et al (1994) J immunol. Methods 173, 19. inflammatory disorders immunologic disorders, cancer
IL-14 GeneSeq Accession R55800 WO9416074 P40222/TXLNA_HUMAN Alpha-taxilin (SEQ ID NO:289)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Ambrus et al (1993) PNAS 90, 63330-34. inflammatory disorders immunologic disorders, cancer
IL-17 receptor GeneSeq Accession B03807 US Pat. No. 6,072,033 Q96F46/I17RA_HUMAN Interleukin-17 receptor A (SEQ ID NO:290)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Yao et al (1995) J. Immunol. 155, 5483-86. Soluble IL-17 receptor polypep- tides may be useful for inhibiting interleukin activities.
IL-17 GeneSeq Accession R76573 WO9518826 Q16552/IL17_HUMAN Interleukin-17A (SEQ ID NO:291)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Yao et al (1995) J. Immunol. 155, 5483-86. inflammatory disorders immunologic disorders, cancer
CTLA-8 GeneSeq Accession W13651 WO9704097 (also known as IL-17) Q16552/IL17_HUMAN Interleukin-17A (SEQ ID NO:291)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225. inflammatory disorders immunologic disorders, cancer

IL-19 GeneSeq Accession W37935 WO9808870 Q9UHD0 IL19_HUMAN Interleukin-19 (isoform 1) (SEQ ID NO:292)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Gallagher et al (2000) Genes Immun. 1, 442-50. inflammatory disorders immunologic disorders, cancer
IL-21 (TIF) GeneSeq Accession Y92879 WO0024758 Q9HBE4 IL21_HUMAN Interleukin-21 (isoform 1) (SEQ ID NO:293)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Parrish-Novak et al (2000) Nature 408, 57-63. inflammatory disorders immunologic disorders, cancer
IL-8 receptor GeneSeq Accession R33420 WO9306229 IL-8RA P25024/CXCR1_HUMAN C-X-C chemokine receptor type 1 (SEQ ID NO:294) IL-8RB P25025/CXCR2_HUMAN C-X-C chemokine receptor type 2 (SEQ ID NO:295)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Holmes et al (1991) Science 253, 1278-80. Soluble IL-8 receptor polypep- tides may be useful for inhibiting interleukin activities.
Human type II interleukin- 1 receptor GeneSeq Accession R85480 US Pat. No. 5,464,937 P27930 IL1R2_HUMAN Interleukin-1 receptor type 2 (SEQ ID NO:296)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinarello (1989) Cytokine 1, 14-20. Soluble type II interleukin-1 receptor polypep- tides may be useful for inhibiting interleukin activities.
Human interleukin-12 receptor GeneSeq Accession R69632 EP638644 IL-12 receptor B1 P42701 IL12R1_HUMAN Interleukin-12 receptor	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Hori et al (1987), Blood 70, 1069- 1078. Soluble IL-12 receptor polypep- tides may be useful for inhibiting interleukin activities.

subunit beta-1 (isoform 1) (SEQ ID NO:393) IL-12 receptor B2 Q99665 I12R2_HUMAN Interleukin-12 receptor subunit beta-2 (isoform 1) (SEQ ID NO:394)	
Interleukin 8 receptor B GeneSeq Accession R80758 US Pat. No. 5,440,021 IL-8RB P25025/CXCR2_HUMAN C-X-C chemokine receptor type 2 (SEQ ID NO:295)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Holmes et al (1991) Science 253, 1278-80. Soluble IL-8 receptor B polypep- tides may be useful for inhibiting interleukin activities.
Human IL-8 receptor protein hIL8RA GeneSeq Accession B09989 JP08103276 IL-8RA P25024/CXCR1_HUMAN C-X-C chemokine receptor type 1 (SEQ ID NO:294)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Holmes et al (1991) Science 253, 1278-80. Soluble IL-8 receptor A polypep- tides may be useful for inhibiting interleukin activities.
Human IL-8 receptor protein hIL8R GeneSeq Accession B09990 JP08103276 IL-8RA P25024/CXCR1_HUMAN C-X-C chemokine receptor type 1 (SEQ ID NO:294) IL-8RB P25025/CXCR2_HUMAN C-X-C chemokine receptor type 2 (SEQ ID NO:295)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Holmes et al (1991) Science 253, 1278-80. Soluble IL-8 receptor polypep- tides may be useful for inhibiting interleukin activities.
Interleukin-2 receptor associated protein p43	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of

GeneSeq Accession R97569 WO9621732- SEQ ID NO:2 of WO9621732 (SEQ ID NO:395)	immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferons: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Gillis <i>et al.</i> (1978) J. Immunol. 120, 2027. Soluble IL-2 receptor polypeptides may be useful for inhibiting interleukin activities.
Human interleukin-17 receptor GeneSeq Accession W04185 WO9629408 Q96F46/I17RA_HUMAN Interleukin-17 receptor A (SEQ ID NO:290)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Yao <i>et al.</i> (1995) J. Immunol. 155, 5483-86. Soluble IL-17 receptor polypeptides may be useful for inhibiting interleukin activities.
Human interleukin-11 receptor GeneSeq Accession R99090 WO9619574 Q14626/I11RA_HUMAN Interleukin-11 receptor subunit alpha (SEQ ID NO:297)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Lu <i>et al.</i> (1994) J immunol. Methods 173, 19. Soluble IL-11 receptor polypeptides may be useful for inhibiting interleukin activities.
Human interleukin-1 receptor accessory protein GeneSeq Accession W01911 WO9623067 Human IL1R Acp SEQ ID NO:3 of WO9623067 (SEQ ID NO:396) Soluble Human IL1R Acp SEQ ID NO:9 of WO9623067 (SEQ ID NO:397)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinarello (1989) Cytokine 1, 14-20. Inflammatory disorders immunologic disorders, cancer
AGF Protein GeneSeq Accession R92749 US Pat. No. 5,488,032 Q8NI99/ANGL6_HUMAN Angiopoietin-related protein 6 (SEQ ID NO:278)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225. Inflammatory disorders immunologic disorders, cancer
Human interleukin-1 type-	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes,

3 receptor GeneSeq Accession R91064 WO9607739 SEQ ID NO:2 and 4 of WO9607739 (SEQ ID NO:398 and SEQ ID NO:399, respectively)	monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Orencole & Dinarello (1989) Cytokine 1, 14-20. Soluble IL-type-3 receptor polypep- tides may be useful for inhibiting interleukin activities
Human interleukin-13 beta receptor GeneSeq Accession W24972 WO9720926 SEQ ID NO:2 from WO9720926 (SEQ ID NO:400)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Boutelier et al (1995) J. Immunol. Methods 181, 29. Soluble IL-13 beta receptor polypep- tides may be useful for inhibiting interleukin activities.
Human interleukin-13 alpha receptor GeneSeq Accession W24973 WO9720926 IL-13RA1 P78552/I13R1_HUMAN Interleukin-13 receptor subunit alpha-1 (isoform 1) (SEQ ID NO:298) IL-13RA2 Q14627/I13R2_HUMAN Interleukin-13 receptor subunit alpha-2 (SEQ ID NO:299)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Boutelier et al (1995) J. Immunol. Methods 181, 29. Soluble IL-13 alpha receptor polypep- tides may be useful for inhibiting interleukin activities.
Human interleukin-4 receptor GeneSeq Accession W13499 US Pat. No. 5,599,905 P24394/IL4RA_HUMAN Interleukin-4 receptor subunit alpha (isoform 1) (SEQ ID NO:300)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Siegel & Mostowski (1990) J Immunol Methods 132, 287-295. Soluble IL-4 receptor polypep- tides may be useful for inhibiting interleukin activities.
Human interleukin-12 beta-2 receptor GeneSeq Accession W12771	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis

EP759466 Q9966/ I12R2_HUMAN Interleukin-12 receptor subunit beta-2 (isoform 1) (SEQ ID NO:301)	of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Hori et al (1987), Blood 70, 1069- 1078. Soluble IL-12 beta-2 receptor polypep- tides may be useful for inhibiting interleukin activities.
Human interleukin-12 beta-1 receptor. GeneSeq Accession W12772 EP759466 P4270/ I12R1_HUMAN Interleukin-12 receptor subunit beta-1 (isoform 1) (SEQ ID NO:302)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Hori et al (1987), Blood 70, 1069- 1078. Soluble IL-12 beta-1 receptor polypep- tides may be useful for inhibiting interleukin activities.
Human IL-9 receptor protein GeneSeq Accessions W64055, W64056, and W64057 WO9824904 Q01113/IL9R_HUMAN Interleukin-9 receptor (isoform 1) (SEQ ID NO:303)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Yang et al (1989) Blood 74, 1880- 84. Soluble IL-9 receptor polypep- tides may be useful for inhibiting interleukin activities.
IL-10 receptor GeneSeq Accession W41804 US Pat. No. 5,716,804 IL-10RA Q13651/I10R1_HUMAN Interleukin-10 receptor subunit alpha (SEQ ID NO:304) IL-10RB Q0833/I10R2_HUMAN Interleukin-10 receptor subunit beta (SEQ ID NO:305)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Thompson-Snipes et al (1991) J. Exp. Med. 173, 507- 510. Soluble IL-10 receptor polypep- tides may be useful for inhibiting interleukin activities.
Human IL-6 receptor GeneSeq Accession Y30938 JP11196867 P08887/IL6RA_HUMAN Interleukin-6 receptor	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington,

subunit alpha (isoform 1) (SEQ ID NO:306)	D.C. 1987, pp. 221-225; and Aarden et al (1987) Eur. J. Immunol 17, 1411-16. Soluble IL-6 receptor polypep- tides may be useful for inhibiting interleukin activities.
Il-17 receptor GeneSeq Accession Y97181 US Pat. No. 6,096,305 Q96F46/I17RA_HUMAN Interleukin-17 receptor A (SEQ ID NO:290)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Yao et al (1995) J. Immunol. 155, 5483-86. Soluble IL-17 receptor polypep- tides may be useful for inhibiting interleukin activities.
Il-17 receptor GeneSeq Accession Y97131 US Pat. No. 6,100,235 Q96F46/I17RA_HUMAN Interleukin-17 receptor A (SEQ ID NO:290)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Yao et al (1995) J. Immunol. 155, 5483-86. Soluble IL-17 receptor polypep- tides may be useful for inhibiting interleukin activities.
Human interleukin-3 receptor GeneSeq Accession R25300 EP509826 P26951/IL3RA_HUMAN Interleukin-3 receptor subunit alpha (isoform 1) (SEQ ID NO:307)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; Kitamura et al (1989) J Cell Physiol. 140 323-334. Soluble IL-3 receptor polypep- tides may be useful for inhibiting interleukin activities.
Human GM- CSF receptor GeneSeq Accession R10919 WO9102063 GM-CSF receptor A P15509/CSF2R_HUMAN Granulocyte-macrophage colony-stimulating factor receptor subunit alpha (isoform 1) (SEQ ID NO:308) GM-CSF receptor B P32927/IL3RB_HUMAN Cytokine receptor common subunit beta (isoform 1) (SEQ ID NO:309)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225. Soluble GM-CSF receptor polypep- tides may be useful for inhibiting interleukin activities.
Human IL-5 receptor alpha	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes,

chain GeneSeq Accession R25064 EP492214	monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; Kitamura <i>et al.</i> (1989) J Cell Physiol. 140 323-334. Soluble IL-5 receptor alpha polypeptides may be useful for inhibiting interleukin activities.
Q01344/IL5RA_HUMAN Interleukin-5 receptor subunit alpha (isoform 1) (SEQ ID NO:310)	
IL-5 receptor GeneSeq Accession W82842 WO9847923	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; Kitamura <i>et al.</i> (1989) J Cell Physiol. 140 323-334. Soluble IL-5 receptor polypep- tides may be useful for inhibiting interleukin activities.
Q01344/IL5RA_HUMAN Interleukin-5 receptor subunit alpha (isoform 1) (SEQ ID NO:310)	
IL-6 receptor GeneSeq Accession R37215 JP05091892	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Aarden <i>et al.</i> (1987) Eur. J. Immunol 17, 1411-16. Soluble IL-6 receptor polypep- tides may be useful for inhibiting interleukin activities.
P08887/IL6RA_HUMAN Interleukin-6 receptor subunit alpha (isoform 1) (SEQ ID NO:306)	
Human B cell stimulating factor-2 receptor GeneSeq Accession P90525 AU8928720	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225. Soluble B cell stimulating factor-2 receptor polypep- tides may be useful for inhibiting interleukin activities.
P08887/IL6RA_HUMAN Interleukin-6 receptor subunit alpha (isoform 1) (SEQ ID NO:306)	
IL-7 receptor clone GeneSeq Accession R08330 EP403114	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i> , in <i>Lymphokines and Interferens: A Practical Approach</i> , Clemens <i>et al.</i> , eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Park <i>et al.</i> (1990) J. Exp. Med. 171, 1073-79. Soluble IL-7 receptor polypep- tides may be useful for inhibiting interleukin activities.
P1687/IL7RA_HUMAN Interleukin-7 receptor subunit alpha (isoform 1) (SEQ ID NO:311)	
EPO receptor; EPOR GeneSeq Accession R06512 WO9008822	EPO Receptor is involved in the proliferation and differentiation of erythroblasts. EPO Receptor activity can be determined using assays known in the art, such as, J Biol Chem 2001 Mar 23; 276(12: 8995-9002; JAK2 protein tyrosine kinase activity: Blood

P19235/EPOR_HUMAN Erythropoietin receptor (isoform EPOR-F) (SEQ ID NO:312)	1994 Sep 1; 84(5): 1501-7 and Mol Cell Biol. 1994 Oct; 14(10): 6506-14. Inflammatory disorders, immunologic disorders, cancer, erythroblast proliferation and differentiation
IL-15 receptor GeneSeq Accession R90843 WO9530695 Q1326/IL15RA_HUMAN Interleukin-15 receptor subunit alpha (isoform 1) (SEQ ID NO:313)	Interleukins are a group of multi- functional cytokines synthesized by lymphocytes, monocytes, and macrophages. Known functions include stimulating proliferation of immune cells (e.g., T helper cells, B cells, eosinophils, and lymphocytes), chemotaxis of neutrophils and T lymphocytes, and/or inhibition of interferons. Interleukin activity can be determined using assays known in the art: Matthews <i>et al.</i>, in <i>Lymphokines and Interferens: A Practical Approach</i>, Clemens <i>et al.</i>, eds, IRL Press, Washington, D.C. 1987, pp. 221-225; and Giri et al (1994) EMBO J. 13 2822- 2830. Soluble IL-15 receptor polypep- tides may be useful for inhibiting interleukin activities. Obesity. Metabolic Disease. Diabetes. Enhancing secretion and stability of Interleukin 15
CD137; 4-1BB Receptor Protein GeneSeq Accession R70977 WO9507984 Q07011/TNR9_HUMAN Tumor necrosis factor receptor superfamily member 9 (SEQ ID NO:314)	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i>, 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i>, 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Soluble 4-1BB receptor polypep- tides may be useful for inhibiting apoptosis, NF-kB activation, and/or co-stimulation of immune cells such as B and T cells.
BCMA GeneSeq Accession Y71979 WO0068378 Q02223/TNR17_HUMAN Tumor necrosis factor receptor superfamily member 17 (isoform 1) (SEQ ID NO:315)	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i>, 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i>, 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Soluble BCMA receptor polypep- tides may be useful for inhibiting apoptosis, NF-kB activation, and/or co-stimulation of immune cells such as B and T cells.
CD27 GeneSeq Accession R20814 WO9201049 P26842/CD27_HUMAN CD27 antigen (SEQ ID NO:316)	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i>, 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i>, 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Soluble CD27 polypeptides may be useful for inhibiting apoptosis, NF-kB activation, and/or co- stimulation of immune cells such as B and T cells.
CD30 GeneSeq Accession R35478 DE4200043 P28908/TNR8_HUMAN Tumor necrosis factor receptor superfamily member 8 (isoform 1)	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i>, 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i>, 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Soluble CD30 polypeptides may be useful for inhibiting apoptosis, NF-kB activation, and/or co- stimulation of immune cells such as B and T cells.

(SEQ ID NO:317)	
CD40 GeneSeq Accession Y33499 WO9945944	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Soluble CD40 polypeptides may be useful for inhibiting apoptosis, NF-kB activation, and/or co- stimulation of immune cells such as B and T cells.
P25942/TNR5_HUMAN Tumor necrosis factor receptor superfamily member 5 (isoform 1)	
(SEQ ID NO:318)	
EDAR Genbank Accession AAD50077	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Immune Disorders, Lymphomas, X- linked hypohidrotic ectodermal dysplasia
Q9UNE0 EDAR_HUMAN Tumor necrosis factor receptor superfamily member EDAR (isoform 1)	
(SEQ ID NO:319)	
OX40; ACT-4 GeneSeq Accession R74737 WO9512673	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Immune Disorders, Lymphomas, T cell disorders
P43489/TNR4_HUMAN Tumor necrosis factor receptor superfamily member 4	
(SEQ ID NO:320)	
TACI GeneSeq Accession W75783 WO9839361	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Soluble TACI receptor polypep- tides may be useful for inhibiting apoptosis, NF-kB activation, and/or co-stimulation of immune cells such as B and T cells.
O14836/TR13B_HUMAN Tumor necrosis factor receptor superfamily member 13B (isoform 1)	
(SEQ ID NO:321)	
TNF-R GeneSeq Accession R10986 AU9058976	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Soluble TNF-R receptor polypep- tides may be useful for inhibiting apoptosis, NF-kB activation, and/or co-stimulation of immune cells such as B and T cells.
P19438/TNR1A_HUMAN Tumor necrosis factor receptor superfamily member 1A (isoform 1)	
(SEQ ID NO:322)	

TNF-RII; TNF p75 receptor; Death Receptor GeneSeq Accession R11141 EP418014 P20333/TNR1B_HUMAN Tumor necrosis factor receptor superfamily member 1B (isoform 1) (SEQ ID NO:323)	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i>, 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i>, 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Soluble TNFR-II receptor polypep- tides may be useful for inhibiting apoptosis, NF-kB activation, and/or co-stimulation of immune cells such as B and T cells.
hAPO-4; TROY GeneSeq Accession W93581 WO9911791 Q9NS68/TNR19_HUMAN Tumor necrosis factor receptor superfamily member 19 (isoform 1) (SEQ ID NO:324)	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i>, 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i>, 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Immune Disorders, Cancers
TNF-alpha precursor GeneSeq Accession P60074 EP205038	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i>, 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i>, 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer
Human TNF- alpha GeneSeq Accession R62463 EP619372 P01375/TNFA_HUMAN Tumor necrosis factor (SEQ ID NO:325)	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i>, 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i>, 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer
Human TNF- alpha GeneSeq Accession R42679 EP563714 P01375/TNFA_HUMAN Tumor necrosis factor (SEQ ID NO:325)	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i>, 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i>, 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer
Human TNF- beta (LT- alpha) GeneSeq Accession B37799 WO0064479 P01374/TNFB_HUMAN Lymphotoxin-alpha	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i>, 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i>, 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer

(SEQ ID NO:326)	
LT-alpha GeneSeq Accession P70107 EP250000	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer
P01374/TNFB_HUMAN Lymphotoxin-alpha	
(SEQ ID NO:326)	
LT-beta GeneSeq Accession R56869 WO9413808	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer
Q06643/TNFC_HUMAN Lymphotoxin-beta (isoform 1)	
(SEQ ID NO:327)	
OPGL GeneSeq Accession W83195 WO9846751	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer, loss of bone mass
O14788/TNF11_HUMAN Tumor necrosis factor ligand superfamily member 11 (isoform 1)	
(SEQ ID NO:328)	
FasL GeneSeq Accession W98071 WO9903999	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer
P48023/TNFL6_HUMAN Tumor necrosis factor ligand superfamily member 6 (isoform 1)	
(SEQ ID NO:329)	
FasL GeneSeq Accession W95041 WO9903998	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer
P48023/TNFL6_HUMAN Tumor necrosis factor ligand superfamily member 6 (isoform 1)	
(SEQ ID NO:329)	
CD27L GeneSeq Accession R50121 WO9405691	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18):

P32970/CD70_HUMAN CD70 antigen (isoform 1) (SEQ ID NO:330)	9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer
CD30 ligand GeneSeq Accession R45007 WO9324135 P32971/TNFL8_HUMAN Tumor necrosis factor ligand superfamily member 8 (SEQ ID NO:331)	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer
CD40L GeneSeq Accession R85486 WO9529935 P29965/CD40L_HUMAN CD40 ligand (SEQ ID NO:332)	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer
4-1BB ligand GeneSeq Accession W26657 US Pat. No. 5,674,704 P41273/TNFL9_HUMAN Tumor necrosis factor ligand superfamily member 9 (SEQ ID NO:333)	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer
FAS Ligand Inhibitory Protein (DcR3) GeneSeq Accession B19335 WO0058465 O95407/TNF6B_HUMAN Tumor necrosis factor receptor superfamily member 6B (SEQ ID NO:334)	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Soluble DcR3 polypeptides may be useful for inhibiting apoptosis, NF-kB activation, and/or co- stimulation of immune cells such as B and T cells.
OX40L GeneSeq Accession R79903 WO9521915 P23510/TNFL4_HUMAN Tumor necrosis factor ligand superfamily member 4 (isoform 1)	Activities associated with apoptosis, NF-kB activation, and co- stimulation of immune cells such as T and B cells. Apoptosis activity, NF-kB activation, and B and T cell co- stimulation can be determined using assays known in the art: Moore <i>et al.</i> , 1999, Science, 285(5425): 260-3; Song HY <i>et al.</i> , 1997 Proc Natl Acad Sci USA 94(18): 9792- 6; Epsevik and Nissen-Meyer, 1986, J. Immunol. Methods. Inflammatory disorders immunologic disorders, cancer

(SEQ ID NO:335)	
Protease inhibitor peptides GeneSeq Accessions R12435, R12436, R12437, R12438, R12439, R12440, and R1244 WO9106561	Peptides that inhibit the function/binding of HIV HIV protease activities are known in the art: HIV protease assays: EP0387231. One can modify the assay to look for inhibition using any of the disclosed protease inhibitor polypeptides. HIV, inflammatory disorders, immuno- logic disorders, cancer, viral infections
Retroviral protease inhibitors GeneSeq Accessions R06660, R06661, R06662, R06663, R06664, R06665, R06666, R06667, R06668, R06669, R06670, R06671, R06672, R06673, R06674, R06675, and R06676 EP387231	Peptides that inhibit the function/binding of HIV HIV protease activities are known in the art: HIV protease assays: EP0387231. One can modify the assay to look for inhibition using any of the disclosed protease inhibitor polypeptides. HIV, inflammatory disorders, immuno- logic disorders, cancer, viral infections
HIV protease inhibiting peptides GeneSeq Accessions R59293, R59294, R59295, R59296, R59297, R59298, R59299, R592300, R59301, R59302, R59301, R59302, R59303, R59304, R59305, R59306, R59307, R59308, R59309, R59310, R59311, R59312, R59313, R59314, R59315, R59316, R59317, R59318, R59319, R59320, R59321, R59322, R59323, R59324, R59325, R59326, R59327, R59328, R59329, R59330, R59331, R59332, R59333, R59334, R59335, R59336, R59337, R59338, R59339, R59340, R59341, R59342, R59343, R59344, R59345, R59346, R59347, R59348, R59349, and R59350 WO9301828	Peptides that inhibit the function/binding of HIV HIV protease activities are known in the art: HIV protease assays: EP0387231. One can modify the assay to look for inhibition using any of the disclosed protease inhibitor polypeptides. HIV, inflammatory disorders, immuno- logic disorders, cancer, viral infections
HIV-1 protease inhibitors GeneSeq Accessions R86326, R86327, R86328, R86329, R86330, R86331, R86332, R86333, R86334, R86335, R86336, R86337, R86338, R86339, R86340,	Peptides that inhibit the function/binding of HIV HIV protease activities are known in the art: HIV protease assays: EP0387231. One can modify the assay to look for inhibition using any of the disclosed protease inhibitor polypeptides. HIV, inflammatory disorders, immuno- logic disorders, cancer, viral infections

R86341, R86342, R86343, R86344, R86345, R86346, R86347, R86348, R86349, R86350, R86351, R86352, R86353, R86354, R86355, R86356, R86357, R86358, R86359, R86360, R86361, R86362, R86363, R86364, R86365, R86366, R86367, R86368, R86369, R86370, and R86371 DE4412174	
HIV Inhibitor Peptide GeneSeq Accession Y89687 WO9959615	Peptides that inhibit the function/binding of HIV HIV protease activities are known in the art: HIV protease assays: EP0387231. One can modify the assay to look for inhibition using any of the disclosed protease inhibitor polypeptides. HIV, inflammatory disorders, immuno- logic disorders, cancer, viral infections
HIV Inhibitor Peptide GenSeq Accession Y31955 WO9948513	Peptides that inhibit the function/binding of HIV HIV protease activities are known in the art: HIV protease assays: EP0387231. One can modify the assay to look for inhibition using any of the disclosed protease inhibitor polypeptides. HIV, inflammatory disorders, immuno- logic disorders, cancer, viral infections
HIV Inhibitor Peptide Science 291, 884 (2001); Published online 12 Jan. 2001; 10.1126/science.1 057453	Peptides that inhibit the function/binding of HIV HIV protease activities are known in the art: HIV protease assays: EP0387231. One can modify the assay to look for inhibition using any of the disclosed protease inhibitor polypeptides. HIV, inflammatory disorders, immuno- logic disorders, cancer, viral infections
Human monocyte chemoattractant factor hMCP-3 GeneSeq Accession R73915 WO9509232 P80098/CCL7_HUMAN C- C motif chemokine 7 (SEQ ID NO:336)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, particularly useful for treating bacterial and/or viral meningitis
Human monocyte chemoattractant factor hMCP-1 GeneSeq Accession R73914 WO9509232 P13500/CCL2_HUMAN C- C motif chemokine 2 (SEQ ID NO:337)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, particularly useful for treating bacterial and/or viral meningitis
Human gro-beta	Chemokines are a family of related small, secreted proteins involved in biological

chemokine GeneSeq Accessions R66699 and W17671 WO9429341 P19875/CXCL2_HUMAN C-X-C motif chemokine 2 (SEQ ID NO:338)	processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, inflammatory disorders, blood-related disorders, stem cell transplantation, cancer
Human gro-gamma chemokine GeneSeq Accessions R66700 and W17672 WO9429341 P19876/CXCL3_HUMAN C-X-C motif chemokine 3 (SEQ ID NO:339)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, inflammatory disorders, blood-related disorders, stem cell transplantation, cancer
Human gro-alpha chemokine GeneSeq Accessions R66698 and W18024 WO9429341 P09341/GROA_HUMAN Growth-regulated alpha protein (SEQ ID NO:340)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, inflammatory disorders, blood-related disorders, stem cell transplantation, cancer
Human eosinophil- expressed chemokine (EEC) GeneSeq Accession W05186 WO9632481 SEQ ID NO:2 of WO9632481 (SEQ ID NO:401)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, particularly treatment of eosinophilia, inflammation, allergies, asthma, leukaemia and lymphoma
Chemokine-like protein PF4-414 Full-Length and Mature GeneSeq Accessions R92318 and R99809 WO9613587 Figure 3C of WO9613587 (SEQ ID NO:402)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer and blood-related disorders, particularly

	myelosuppression
Chemokine-like protein IL-8M3 GeneSeq Accession R99812 WO9613587	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ; and Holmes et al (1991) Science 253, 1278-80. Cancer and blood- related disorders, particularly myelosuppression
Human interleukin-8 (IL-8) GeneSeq Accession R99814 WO9613587 P10145/IL8_HUMAN Interleukin-8 (isoform 1) (SEQ ID NO:341)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ; and Holmes et al (1991) Science 253, 1278-80. Cancer and blood- related disorders, particularly myelosuppression
Chemokine-like protein IL-8M1 Full-Length and Mature GeneSeq Accessions R99815 and R99803 WO9613587 Figure 4B of WO9613587 (SEQ ID NO:403)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ; and Holmes et al (1991) Science 253, 1278-80. Cancer and blood- related disorders, particularly myelosuppression
Chemokine-like protein IL-8M8 Full-Length and Mature GeneSeq Accessions R99816 and R99805 WO9613587 Figure 4C of WO9613587 (SEQ ID NO:404)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ; and Holmes et al (1991) Science 253, 1278-80. Cancer and blood- related disorders, particularly myelosuppression
Chemokine-like protein IL-8M8 Full-Length and Mature GeneSeq Accessions R99817 and R99806 WO9613587 Figure 4C of WO9613587	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138:

(SEQ ID NO:404)	Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ; and Holmes et al (1991) Science 253, 1278-80. Cancer and blood- related disorders, particularly myelosuppression.
Chemokine-like protein IL-8M8 Full-Length and Mature GeneSeq Accessions R99818 and R99804 WO9613587 Figure 4C of WO9613587 (SEQ ID NO:404)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ; and Holmes et al (1991) Science 253, 1278-80. Cancer and blood- related disorders, particularly myelosuppression.
Chemokine-like protein IL-8M8 Full-Length and Mature GeneSeq Accessions R99819 and R99807 WO9613587 Figure 4C of WO9613587 (SEQ ID NO:404)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer and blood- related disorders, particularly myelosuppression.
Chemokine-like protein IL-8M8 Full-Length and Mature GeneSeq Accessions R99822 and R9807 WO9613587 Figure 4C of WO9613587 (SEQ ID NO:404)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer and blood- related disorders, particularly myelosuppression.
Human foetal spleen expressed chemo-kine, FSEC GeneSeq Accession R98499 WO9622374 SEQ ID NO:2 of WO9622374 (SEQ ID NO:405)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders
Liver expressed chemokine- 1(LVEC-1) GeneSeq Accession R95689 WO9616979 SEQ ID NO:2 of WO9616979	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined

(SEQ ID NO:406)	using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inflammation of the liver
Liver expressed chemokine- 2(LVEC-2) GeneSeq Accession R95690 WO9616979 SEQ ID NO:4 of WO9616979 (SEQ ID NO:407)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inflammation of the liver
Pituitary expressed chemokine (PGEC) GeneSeq Accession R95691 WO9616979 SEQ ID NO:6 of WO9616979 (SEQ ID NO:408)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inflammation, particularly of the liver
Adenoid- expressed chemokine (ADEC) GeneSeq Accession R97664 WO9617868 SEQ ID NO:2 of WO9617868 (SEQ ID NO:409)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inflammation, angiogenesis, tumorigenesis, musculoskeletal disorders
Human chemokine CC-2 GeneSeq Accession W38170 WO9741230 Q16663/CCL15_HUMAN C-C motif chemokine 15 (SEQ ID NO:342)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, cell migration, proliferation, and differentiation disorders
Human chemokine HCC-1 GeneSeq Accession W38171 WO9741230 Q16627/CCL14_HUMAN C-C motif chemokine 14 (SEQ ID NO:343)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138:

	Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, cell migration, proliferation, and differentiation disorders
Human chemokine CC-3 GeneSeq Accession W38172 WO9741230 Q16627/CCL14_HUMAN C-C motif chemokine 14 (SEQ ID NO:343)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, cell migration, proliferation, and differentiation
Novel betachemokine designated PTEC GeneSeq Accession W27271 WO9739126 SEQ ID NO:2 of WO9739126 (SEQ ID NO:410)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, vascular disorders, cancer
Human CX3C 111 amino acid chemokine GeneSeq Accession W23344 WO9727299 SEQ ID NO:2 of WO9727299 (SEQ ID NO:411)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, inflammatory diseases, abnormal proliferation, regeneration, degeneration, and atrophy
Human CCF18 chemokine GeneSeq Accession W25942 WO9721812 SEQ ID NO:4 of WO9721812 (SEQ ID NO:412)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Abnormal physiology and development disorders, can also be used as an anti- viral agent
Human beta- chemokine H1305 (MCP-2) GeneSeq Accession W26655 WO9725427 P80075/CCL8_HUMAN C- C motif chemokine 8	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined

(SEQ ID NO:344)	using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Chemotaxis, blood- related disorders, viral infection, HIV, wound healing, cancer
Human eosinocyte CC type chemokine eotaxin GeneSeq Accession W14990 WO9712914 P51671/CCL11_HUMAN Eotaxin (SEQ ID NO:245)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inflammatory and immune disorders
Human thymus and activation regulated cytokine (TARC) GeneSeq Accession W14018 WO9711969 Q92583/CCL17_HUMAN C-C motif chemokine 17 (SEQ ID NO:261)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in molecular Biology, 2000, vol. 138: Chemokine Protocols, Edited by A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power Humana Press Inc., Totowa, NJ Inflammatory and immune disorders
Human chemokine beta-8 short forms GeneSeq Accession W16315 WO9712041 Wildtype chemokine beta-8 provided as: P55773/CCL23_HUMAN C-C motif chemokine 23 (SEQ ID NO:459)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer, wound healing, immune disorders
Microphage derived chemokine, MDC GeneSeq Accession W20058 WO9640923 O00626/CCL22_HUMAN C-C motif chemokine 22 (SEQ ID NO:345)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inflammatory diseases, wound healin, angiogenesis
Human chemokine ZSIG-35 GeneSeq Accession W30565 WO9844117 SEQ ID NO:2 of WO WO9844117	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined

(SEQ ID NO:413)	using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inflammatory and immune diseases
Primate CC chemokine "ILINCK" GeneSeq Accession W69990 WO98328658 SEQ ID NO:4 from WO9832858 (SEQ ID NO:414)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune and inflammatory disorders, abnormal proliferation, regeneration and atrophy disorders
Primate CXC chemokine "IBICK" GeneSeq Accession W69989 WO9832858 SEQ ID NO:2 from WO9832858 (SEQ ID NO:415)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune and inflammatory disorders, abnormal proliferation, regeneration and atrophy disorders
Human CC-type chemokine protein designated SLC (secondary lymphoid chemokine) GeneSeq Accession W69163 WO9831809 O00585/CCL21_HUMAN C-C motif chemokine 21 (SEQ ID NO:346)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune, inflammatory, and infectious disorders, cancer
Human CC chemokine ELC protein GeneSeq Accession W62542 WO9826071 Q99731/CCL19_HUMAN C-C motif chemokine 19 (SEQ ID NO:249)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer and infectious diseases, particularly herpes virus
Human DVic-1 C-C chemokine GeneSeq Accession W60649 WO9823750 SEQ ID NO:2 of WO9823750	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined

(SEQ ID NO:416)	using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Abnormal proliferation, regeneration, degeneration, and atrophy disorders, including cancer
Human C-C chemokine DGWCC GeneSeq Accession W60650 WO9823750 SEQ ID NO:6 of WO9823750 (SEQ ID NO:417)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, cell proliferation disorders, cancer
Human STCP-1 GeneSeq Accession W62783 WO9824907 O00626/CCL22_HUMAN C-C motif chemokine 22 (SEQ ID NO:345)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, particularly T cell related disorders, viral infection, and inflammation, especially joint
Exodus protein GeneSeq Accession W61279 WO9821330 P78556/CCL20_HUMAN C-C motif chemokine 20 (isoform 1) (SEQ ID NO:248)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune and inflammatory disorders, angiogenesis, cancer, and proliferation disorders, particularly myeloproliferative diseases
Human Chr19kine protein GeneSeq Accession W50887 WO9814581 SEQ ID NO:10 of WO9814581 (SEQ ID NO:418)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer and degenerative disorders
Human T cell mixed lymphocyte reaction expressed chemokine (TMEC) GeneSeq Accession W58703 US Pat. No. 5,780,268	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which

SEQ ID NO:2 of US Pat. No. 5,780,268 (SEQ ID NO:460)	bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune, inflammatory, and infectious disorders, cancer
Human 6CKine protein GeneSeq Accession W50885 WO9814581 SEQ ID NO:8 of WO9814581 (SEQ ID NO:419)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer and de- generative disorders
human liver and activation regulated chemokine (LARC) GeneSeq Accession W57475 WO9817800 P78556/CCL20_HUMAN C-C motif chemokine 20 (isoform 1) (SEQ ID NO:248)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune, inflammatory, and infectious disorders, cancer
RANTES peptide GeneSeq Accession W29538 WO9744462 Wildtype Rantes provided herien as P13501/CCL5_HUMAN C- C motif chemokine 5 (SEQ ID NO:241)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Infectious diseases, particularly HIV
RANTES 8-68 GeneSeq Accession W29529 WO9744462 Wildtype Rantes provided herien as P13501/CCL5_HUMAN C- C motif chemokine 5 (SEQ ID NO:241)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Infectious diseases, particularly HIV
RANTES 9-68 GeneSeq Accession W29528 WO9744462 Wildtype Rantes provided herien as P13501/CCL5_HUMAN C-	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined

C motif chemokine 5 (SEQ ID NO:241)	using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Infectious diseases, particularly HIV
Human chemokine protein 331D5 GeneSeq Accession W59433 WO9811226 SEQ ID NO:12 of WO9811226 (SEQ ID NO:420)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Abnormal proliferation, regeneration, degeneration, or atrophy, including cancer
Human chemokine protein 61164 GeneSeq Accession W59430 WO9811226 SEQ ID NO:6 of WO9811226 (SEQ ID NO:421)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Abnormal proliferation, regeneration, degeneration, or atrophy, including cancer
Chemokine MCP-4 GeneSeq Accession W56690 WO9809171 Q99616/CCL13_HUMAN C-C motif chemokine 13 (SEQ ID NO:347)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune, inflammatory, and infectious diseases
Human stromal cell-derived chemokine, SDF-1 GeneSeq Accession W50766 FR2751658 P48061/SDF1_HUMAN Stromal cell-derived factor 1 (isoform beta) (SEQ ID NO:260)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. HIV infections
Thymus expressed chemokine (TECK) GeneSeq Accession W44397 WO9801557 O15444/CCL25_HUMAN C-C motif chemokine 25	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined

(SEQ ID NO:348)	using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune and inflammatory disorders
Human chemokine MIP-3alpha GeneSeq Accession W44398 WO9801557 P78556/CCL20_HUMAN C-C motif chemokine 20 (isoform 1) (SEQ ID NO:248)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune and inflammatory disorders
Human chemokine MIP-3beta GeneSeq Accession W44399 WO9801557 Q99731/CCL19_HUMAN C-C motif chemokine 19 (SEQ ID NO:249)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune and inflammatory disorders
Human monocyte chemotactic proprotein (MCP) sequence GeneSeq Accession W42072 WO9802459 SEQ ID NO:1 of WO9802459 (SEQ ID NO:456)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, respiratory disorders, cancer
Macrophage- derived chemokine (MDC) GeneSeq Accessions W40811 and Y24414 US Pat. No. 5,688,927/US Pat. No. 5,932,703 O00626/CCL22_HUMAN C-C motif chemokine 22 (SEQ ID NO:345)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune, and inflammatory disorders, cancer
Macrophage derived chemokine analogue MDC-eyfy GeneSeq Accession Y24416 US Pat. No. 5,932,703 ("eyfy" disclosed as SEQ ID NO: 546) Wildtype MDC is SEQ ID	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power.

NO:2 of 5,932,703 (SEQ ID NO:422)	Humana Press Inc., Totowa, NJ. Immune and inflammatory disorders
Macrophage derived chemokine analogue MDC (n + 1) GeneSeq Accession Y24413 US Pat. No. 5,932,703	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune and inflammatory disorders
Macrophage derived chemokine analogue MDC-yl GeneSeq Accession Y24415 US Pat. No. 5,932,703	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune and inflammatory disorders
Human type CC chemokine eotaxin 3 protein sequence GeneSeq Accession Y43178 JP11243960 Q9Y258/CCL26_HUMAN C-C motif chemokine 26 (SEQ ID NO:349)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Allergic diseases and HIV infection
Human MCP-3 and human Muc-1 core epitope (VNT) fusion protein GeneSeq Accession Y29893 WO9946392 Wildtype MCP-3 has the sequence: P80098/CCL7_HUMAN C-C motif chemokine 7 (SEQ ID NO:336) Wildtype Muc-1 has the sequence: P15941 MUC1_HUMAN Mucin-1 (isoform 1)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer and immune disorders, particularly HIV infection

(SEQ ID NO:461)	
Human IP-10 and human Muc-1 core epitope (VNT) fusion protein GeneSeq Accession Y29894 WO9946392	
Wildtype IP10 has the sequence:	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking.
P02778/CXL10_HUMAN C-X-C motif chemokine 10	Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer and immune disorders, particularly HIV infection
(SEQ ID NO:242)	
Wildtype Muc-1 has the sequence:	
P15941 MUC1_HUMAN Mucin-1 (isoform 1)	
(SEQ ID NO:461)	
Human IP-10 and HIV-1 gp 120 hyper- variable region fusion protein GeneSeq Accession Y29897 WO9946392	
Wildtype IP10 has the sequence:	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking.
P02778/CXL10_HUMAN C-X-C motif chemokine 10	Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer and immune disorders, particularly HIV infection
(SEQ ID NO:242)	
Wildtype gp120 has the sequence:	
P03378 32-509 (cleaved product of gp160)	
(SEQ ID NO:462)	
Human mammary associated chemokine (MACK) protein Full-Length and Mature GeneSeq Accessions Y29092 and Y29093 WO9936540	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138:

Full-length: SEQ ID NO:1 of WO9936540 (SEQ ID NO:423) Mature Form: SEQ ID NO:2 of WO9936540 (SEQ ID NO:424)	Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Breast disease, including cancer
Tim-1 protein GeneSeq Accession Y28290 WO9933990 SEQ ID NO:2 of WO9933990 (SEQ ID NO:350)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inflammation due to stimuli such as heart attacks and stroke, infection, physical trauma, UV or ionizing radiation, burns, frostbite or corrosive chemicals
Human Lkn-1 Full-Length and Mature protein GeneSeq Accessions Y17280, Y17274, Y17281, and Y17275 WO9928473 and WO9928472 Q16663/CCL15_HUMAN C-C motif chemokine 15 (SEQ ID NO:342)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. HIV infection and cancer, particularly leukemia
N-terminal modified chemokine met- hSDF-1 alpha GeneSeq Accession Y05818 WO9920759 SEQ ID NO:10 of WO9920759 (SEQ ID NO:425)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inhibit or stimulate angiogenesis, inhibit the binding of HIV
N-terminal modified chemokine met- hSDF-1 beta GeneSeq Accession Y05819 WO9920759 SEQ ID NO:11 of WO9920759 (SEQ ID NO:426)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inhibit or stimulate angiogenesis, inhibit the binding

	of HIV, antiinflammatory; immunosuppressant
N-terminal modified chemokine GroHEK/hSDF- 1alpha GeneSeq Accession Y05820 WO9920759 SEQ ID NO:12 of WO9920759 (SEQ ID NO:427)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inhibit or stimulate angiogenesis, inhibit the binding of HIV, antiinflammatory; immunosuppressant
N-terminal modified chemokine GroHEK/hSDF- 1beta. GeneSeq Accession Y05821 WO9920759 SEQ ID NO:13 of WO9920759 (SEQ ID NO:428)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inhibit or stimulate angiogenesis, inhibit the binding of HIV, antiinflammatory; immunosuppressant
Chemokine Eotaxin GeneSeq Accession Y14230 WO9912968 P51671/CCL11_HUMAN Eotaxin (SEQ ID NO:245)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Increase or enhance an inflammatory response, an immune response or haematopoietic cell-associated activity; treat a vascular indication; Cancer; enhance wound healing, to prevent or treat asthma, organ transplant rejection, rheumatoid arthritis or allergy
Chemokine hMCP1a GeneSeq Accession Y14225 WO9912968	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, Vascular disorders, Wound healing, cancer, prevent organ transplant rejection, Increase or enhance an inflammatory response,
Chemokine hMCP1b GeneSeq Accession Y14226 WO9912968	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-

	protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, Vascular disorders, Wound healing, cancer, prevent organ transplant rejection, Increase or enhance an inflammatory response,
Chemokine hSDF1b GeneSeq Accession Y14228 WO9912968 P48061/SDF1_HUMAN Stromal cell-derived factor 1 (isoform beta) (SEQ ID NO:260)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, Vascular disorders, Wound healing, cancer, prevent organ transplant rejection, Increase or enhance an inflammatory response,
Chemokine hIL-8 GeneSeq Accession Y14229 WO9912968 P10145/IL8_HUMAN Interleukin-8 (isoform 1) (SEQ ID NO:341)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ; and Holmes et al (1991) Science 253, 1278-80. Immune disorders, Vascular disorders, Wound healing, cancer, prevent organ transplant rejection, Increase or enhance an inflammatory response,
Chemokine hMCP1 GeneSeq Accession Y14222 WO9912968 P13500/CCL2_HUMAN C- C motif chemokine 2 (SEQ ID NO:337)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, Vascular disorders, Wound healing, cancer, prevent organ transplant rejection, Increase or enhance an inflammatory response,
Chemokine hMCP2 GeneSeq Accession Y14223 WO9912968 P80075/CCL8_HUMAN C- C motif chemokine 8 (SEQ ID NO:344)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, Vascular disorders, Wound

	healing, cancer, prevent organ transplant rejection, Increase or enhance an inflammatory response,
Chemokine hMCP3 GeneSeq Accession Y14224 WO9912968 P80098/CCL7_HUMAN C- C motif chemokine 7 (SEQ ID NO:336)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, Vascular disorders, Wound healing, cancer, prevent organ transplant rejection, Increase or enhance an inflammatory response,
C-C chemokine, MCP2 GeneSeq Accession Y05300 EP905240 P80075/CCL8_HUMAN C- C motif chemokine 8 (SEQ ID NO:344)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inflammatory, Immune and infectious diseases; pulmonary diseases and skin disorders; tumours, and angiogenesis-and haematopoiesis- related diseases
Wild type monocyte chemotactic protein 2 GeneSeq Accession Y07233 EP906954 P80075/CCL8_HUMAN C- C motif chemokine 8 (SEQ ID NO:344)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inflammatory, Immune and infectious diseases; pulmonary diseases and skin disorders; tumours, and angiogenesis-and haematopoiesis- related diseases
Truncated monocyte chemotactic protein 2 (6- 76) GeneSeq Accession Y07234 EP906954 Figure 1 of EP905241 and EP906954 (SEQ ID NO:429)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power, Humana Press Inc., Totowa, NJ Inflammatory, Immune and infectious diseases; pulmonary diseases and skin disorders; tumours, and angiogenesis-and haematopoiesis- related diseases
Truncated RANTES protein (3-68) GeneSeq Accessions Y07236 and	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies

Y07232 EP905241; EP906954 Figure 1 of EP906954 (SEQ ID NO:430)	including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power, Humana Press Inc., Totowa, NJ Inflammatory, immune and infectious diseases; pulmonary diseases and skin disorders; tumours, and angiogenesis-and haematopoiesis- related diseases
Wild type monocyte chemotactic protein 2 GeneSeq Accession Y07237 EP905241 P80075/CCL8_HUMAN C- C motif chemokine 8 (SEQ ID NO:344)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power, Humana Press Inc., Totowa, NJ Inflammatory, immune and infectious diseases; pulmonary diseases and skin disorders; tumours, and angiogenesis-and haematopoiesis- related diseases
Truncated monocyte chemotactic protein 2 (6-76) GeneSeq Accession Y07238 EP905241 Figure 1 of EP905241 and EP906954 (SEQ ID NO:429)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power, Humana Press Inc., Totowa, NJ Inflammatory, immune and infectious diseases; pulmonary diseases and skin disorders; tumours, and angiogenesis-and haematopoiesis- related diseases
A partial CXCR4B protein GeneSeq Accession W97363 EP897980 SEQ ID NO:2 of EP897980 (sEQ ID NO:431)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power, Humana Press Inc., Totowa, NJ Soluble CXCR4B receptor polypeptides may be useful for inhibiting chemokine activities and viral infection.
Interferon gamma-inducible protein (IP-10) GeneSeq Accession W96709 US Pat. No. 5,871,723 P02778/CXL10_HUMAN C-X-C motif chemokine 10	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power,

(SEQ ID NO:242)	Humana Press Inc., Totowa, NJ Angiogenesis, Cancer, Inflammatory and Immune disorders, Cardio- Vascular disorders, Musco-skeletal disorders
A monokine induced by gamma- interferon (MIG) GeneSeq Accession W96710 US Pat. No. 5,871,723 Q07325/CXCL9_HUMAN C-X-C motif chemokine 9 (SEQ ID NO:351)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power, Humana Press Inc., Totowa, NJ Angiogenesis, Cancer, Inflammatory and Immune disorders, Cardio- Vascular disorders, Musco-skeletal disorders
Interleukin-8 (IL-8) protein. GeneSeq Accession W96711 US Pat. No. 5,871,723 P10145/IL8_HUMAN Interleukin-8 (isoform 1) (SEQ ID NO:341)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ; and Holmes et al (1991) Science 253, 1278-80. Angiogenesis, Cancer, Inflammatory and Immune disorders, Cardio- Vascular disorders, Musco-skeletal disorders
Epithelial neutrophil activating protein-78 (ENA-78) GeneSeq Accession W96712 US Pat. No. 5,871,723 P42830/CXCL5_HUMAN C-X-C motif chemokine 5 (SEQ ID NO:352)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power, Humana Press Inc., Totowa, NJ Angiogenesis, Cancer, Inflammatory and Immune disorders, Cardio- Vascular disorders, Musco-skeletal disorders
Growth related oncogene-alpha (GRO-alpha). GeneSeq Accession W96713 US Pat. No. 5,871,723 P09341/GROA_HUMAN Growth-regulated alpha protein (SEQ ID NO:340)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power, Humana Press Inc., Totowa, NJ Angiogenesis, Cancer, Inflammatory and Immune disorders, Cardio- Vascular disorders, Musco-skeletal disorders
Growth related oncogene-beta (GRO-beta). GeneSeq Accession W96714 US Pat. No. 5,871,723	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which

P19875/CXCL2_HUMAN C-X-C motif chemokine 2 (SEQ ID NO:338)	bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Angiogenesis, Cancer, Inflammatory and Immune disorders, Cardio- Vascular disorders, Musco-skeletal disorders
Growth related oncogene- gamma (GRO-gamma) GeneSeq Accession W96715 US Pat. No. 5,871,723 P19876/CXCL3_HUMAN C-X-C motif chemokine 3 (SEQ ID NO:339)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Angiogenesis, Cancer, Inflammatory and Immune disorders, Cardio- Vascular disorders, Musco-skeletal disorders
A platelet basic protein (PBP) GeneSeq Accession W96716 US Pat. No. 5,871,723 P02775/CXCL7_HUMAN Platelet basic protein (SEQ ID NO:353)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Angiogenesis, Cancer, Inflammatory and Immune disorders, Cardio- Vascular disorders, Musco-skeletal disorders
Connective tissue activating protein-III (CTAP-III) GeneSeqAc- cession S96717 US Pat. No. 5,871,723 SEQ ID NO:9 of U.S. Patent NO. 5,871,723 (SEQ ID NO:354)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Angiogenesis, Cancer, Inflammatory and Immune disorders, Cardio- Vascular disorders, Musco-skeletal disorders
Beta-thrombo- globulin protein (beta-TG) GeneSeq Accession W96718 US Pat. No. 5,871,723 SEQ ID NO:10 of U.S. Patent NO. 5,871,723 (SEQ ID NO:355)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Angiogenesis, Cancer, Inflammatory and Immune disorders, Cardio- Vascular disorders, Musco-skeletal disorders
Neutrophil activating peptide-2 (NAP-2) GeneSeq Accession W96719 US Pat. No.	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The

5,871,723 SEQ ID NO:11 of U.S. Patent NO. 5,871,723 (SEQ ID NO:356)	chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Angiogenesis, Cancer, Inflammatory and Immune disorders, Cardio- Vascular disorders, Musco-skeletal disorders
Granulocyte chemotactic protein-2 (GCP-2) GeneSeq Accession W96720 US Pat. No. 5,871,723 P80162/CXCL6_HUMAN C-X-C motif chemokine 6 (SEQ ID NO:357)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Angiogenesis, Cancer, Inflammatory and Immune disorders, Cardio- Vascular disorders, Musco-skeletal disorders
Human chemokine MIG-beta protein GeneSeq Accession W90124 EP887409 (SEQ ID NO:463)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, viral, parasitic, fungal or bacterial infections, Cancer; autoimmune diseases or transplant rejection
Human ZCHEMO-8 GeneSeq Accession W82716 WO9854326 SEQ ID NO:2 of WO9854326 (SEQ ID NO:432)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, cancer, myelopoietic disorders, autoimmune disorders and immunodeficiencies, Inflammatory and infectious diseases, Vascular disorders, wound healing
Human Act-2 protein GeneSeq Accession W82717 WO9854326 P13236/CCL4_HUMAN C-C motif chemokine 4 (SEQ ID NO:358)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, cancer, myelopoietic disorders, autoimmune disorders and immunodeficiencies, Inflammatory and infectious diseases, Vascular disorders, wound healing

Human SISD protein GeneSeq Accession W82720 WO9854326 P13501/CCL5_HUMAN C- C motif chemokine 5 (SEQ ID NO:241)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, cancer, myelopoietic disorders, autoimmune disorders and immunodeficiencies, Inflammatory and infectious diseases, Vascular disorders, wound healing
Human MI10 protein GeneSeq Accession W82721 WO9854326 SEQ ID NO:37 of WO9854326 (SEQ ID NO:433)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, cancer, myelopoietic disorders, autoimmune disorders and immunodeficiencies, Inflammatory and infectious diseases, Vascular disorders, wound healing
Human MI1A protein GeneSeq Accession W82722 WO9854326 SEQ ID NO:38 of WO9854326 (SEQ ID NO:434)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, cancer, myelopoietic disorders, autoimmune disorders and immunodeficiencies, Inflammatory and infectious diseases, Vascular disorders, wound healing
Human CCC3 protein GeneSeq Accession W82723 WO9854326 SEQ ID NO:39 of WO9854326 (SEQ ID NO:435)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune disorders, cancer, myelopoietic disorders, autoimmune disorders and immunodeficiencies, Inflammatory and infectious diseases, Vascular disorders, wound healing
A human L105 chemokine designated huL105_3. GeneSeq Accession W87588 WO9856818	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-

SEQ ID NO:2 of WO9856818 (SEQ ID NO:436)	protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer, wound healing
A human L105 chemokine designated huL105_7. GeneSeq Accession W87589 WO9856818 SEQ ID NO:4 of WO9856818 (SEQ ID NO:437)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer, wound healing
Human mature gro-alpha polypeptide used to treat sepsis GeneSeq Accession W81498 WO9848828 P09341/GROA_HUMAN Growth-regulated alpha protein (SEQ ID NO:340)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Infectious diseases, sepsis
Human mature gro-gamma polypeptide used to treat sepsis GeneSeq Accession W81500 WO9848828 P19876/CXCL3_HUMAN C-X-C motif chemokine 3 (SEQ ID NO:339)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Infectious diseases, sepsis
Human thymus expressed chemokine TECK and TECK variant GeneSeq Accessions B19607 and B19608 WO0053635 Wildtype TECK provided as: O15444/CCL25_HUMAN C-C motif chemokine 25 (SEQ ID NO:348)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Inflammatory disorders, cancer, Immune and vascular disorders
Human chemokine SDF1alpha GeneSeq Accession B15791 WO0042071	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The

P48061-2/SDF1_HUMAN Isoform Alpha of Stromal cell-derived factor 1 (isoform alpha) (SEQ ID NO:259)	chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
Human chemokine GRO- alpha GeneSeq Accession B15793 WO0042071 P09341/GROA_HUMAN Growth-regulated alpha protein (SEQ ID NO:340)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
Human chemokine eotaxin GeneSeq Accession B15794 WO0042071 P51671/CCL11_HUMAN Eotaxin (SEQ ID NO:245)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
Human chemokine MIG GeneSeq Accession B15803 WO0042071 Q07325/CXCL9_HUMAN C-X-C motif chemokine 9 (SEQ ID NO:351)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
Human chemokine PF4 GeneSeq Accession B15804 WO0042071 P02776/PLF4_HUMAN Platelet factor 4 (SEQ ID NO:359)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
Human chemokine I- 309	Chemokines are a family of related small, secreted proteins involved in biological

GeneSeq Accession B15805 WO0042071 P22362/CCL1_HUMAN C- C motif chemokine 1 (SEQ ID NO:360)	processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
Human chemokine HCC-1 GeneSeq Accession B15806 WO0042071 Q16627/CCL14_HUMAN C-C motif chemokine 14 (SEQ ID NO:361)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
Human chemokine C10 GeneSeq Accession B15807 WO0042071 SEQ ID NO:49 of WO0042071 (SEQ ID NO:438)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
Human chemokine CCR-2 GeneSeq Accession B15808 WO0042071 P41597/CCR2_HUMAN C-C chemokine receptor type 2 (isoform A) (SEQ ID NO:362)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
Human chemokine ENA- 78 GeneSeq Accession B15809 WO0042071 P42830/CXCL5_HUMAN C-X-C motif chemokine 5 (SEQ ID NO:352)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and

	Inflammatory disorders
Human chemokine GRObeta GeneSeq Accession B15810 WO0042071 P19875/CXCL2_HUMAN C-X-C motif chemokine 2 (SEQ ID NO:338)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
Human chemokine IP-10 GeneSeq Accession B15811 WO0042071 P02778/CXL10_HUMAN C-X-C motif chemokine 10 (SEQ ID NO:242)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
Human chemokine SDF1beta GeneSeq Accession B15812 WO0042071 P48061/SDF1_HUMAN Stromal cell-derived factor 1 (isoform beta) (SEQ ID NO:260)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
Human chemokine GRO alpha GeneSeq Accession B15813 WO0042071 P09341/GROA_HUMAN Growth-regulated alpha protein (SEQ ID NO:340)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
Human chemokine MIP1beta GeneSeq Accession B15831 WO0042071 P13236/CCL4_HUMAN C-C motif chemokine 4	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138:

(SEQ ID NO:358)	Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Autoimmune disorders, Immune, Vascular and Inflammatory disorders
A human C-C chemokine designated exodus GeneSeq Accession B07939 US Pat. No. 6,096,300 P78556/CCL20_HUMAN C-C motif chemokine 20 (isoform 1) (SEQ ID NO:248)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer
Human chemokine L105_7 GeneSeq Accession Y96922 US Pat. No. 6,084,071 SEQ ID NO:4 of WO9856818 (SEQ ID NO:437)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Chemotaxis, Gene Therapy, Wound healing
Human chemokine L105_3 GeneSeq Accession Y96923 US Pat. No. 6,084,071 SEQ ID NO:2 of WO9856818 (SEQ ID NO:436)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Chemotaxis, Gene Therapy, Wound healing
Human secondary lymphoid chemokine (SLC) GeneSeq Accession B01434 WO0038706 O00585/CCL21_HUMAN C-C motif chemokine 21 (SEQ ID NO:346)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer, Vascular and Immune disorders
Human non- ELR CXC chemokine H174 GeneSeq Accession Y96310 WO0029439 O14625/CXL11_HUMAN C-X-C motif chemokine 11 (SEQ ID NO:363)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power.

	Humana Press Inc., Totowa, NJ. Immune and Inflammatory disorders, Cancer, Haemostatic and thrombolytic activity
Human non- ELR CXC chemokine IP10 GeneSeq Accession Y96311 WO0029439 P02778/CXL10_HUMAN C-X-C motif chemokine 10 (SEQ ID NO:242)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune and Inflammatory disorders, Cancer, haemostatic and thrombolytic activity
Human non- ELR CXC chemokine Mig GeneSeq Accession Y96313 WO0029439 Q07325/CXCL9_HUMAN C-X-C motif chemokine 9 (SEQ ID NO:351)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Immune and Inflammatory disorders, Cancer, haemostatic and thrombolytic activity
Human chemokine Ckbeta-7 GeneSeq Accession Y96280 WO0028035 Figure1 of WO0028035 (SEQ ID NO:439)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer, wound healing, inflammatory and immunoregulatory disorders
Human chemokine MIP-1alpha GeneSeq Accession Y96281 WO0028035 P10147/CCL3_HUMAN C motif chemokine 3 (SEQ ID NO:364)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer, wound healing, inflammatory and immunoregulatory disorders
Human mature chemokine Ckbeta-7 (optionally truncated) GenSeq Accession Y96282 WO0028035 Figure1 of WO0028035	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined

(SEQ ID NO:440)	using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer, wound healing, inflammatory and immunoregulatory disorders
Human chemokine receptor CXCR3 GeneSeq Accession Y79372 WO0018431 P49682 CXCR3_HUMAN C-X-C chemokine receptor type 3 (isoform 1) (SEQ ID NO:240)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Soluble CXCR3 polypeptides may be useful for inhibiting
Human neurotactin chemokine like domain GeneSeq Accession Y53259 US Pat. No. 6,043,086 P78423/X3CL1_HUMAN Fractalkine (SEQ ID NO:244)	hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. chemokine activities and viral infection.
Human CC type chemokine interleukin C GeneSeq Accession Y57771 JP11302298	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer and infectious diseases
Human CKbeta- 9 GeneSeq Accession B50860 US Pat. No. 6,153,441 O00585/CCL21_HUMAN C-C motif chemokine 21 (SEQ ID NO:346)	Chemokines are a family of related small, secreted proteins involved in biological processes ranging from hematopoiesis, angiogenesis, and leukocyte trafficking. Members of this family are involved in a similarly diverse range of pathologies including inflammation, allergy, tissue rejection, viral infection, and tumor biology. The chemokines exert their effects by acting on a family of seven transmembrane G-protein coupled receptors. Over 40 human chemokines have been described, which bind to ~17 receptors thus far identified. Chemokine activities can be determined using assays known in the art: Methods in Molecular Biology, 2000, vol. 138: Chemokine Protocols. Edited by: A. E. I. Proudfoot, T. N. C. Wells, and C. A. Power. Humana Press Inc., Totowa, NJ. Cancer, Auto- immune and inflammatory disorders, Cardiovascular disorders
Preproapolipo- protein "paris" variant GeneSeq Accession W08602 WO9637608 (SEQ ID NO:466)	Apoa-1 participates in the reverse transport of cholesterol from tissues to the liver for excretion by promoting cholesterol efflux from tissues and by acting as a cofactor for the lecithin cholesterol acyltransferase (lcat). Lipid binding activity can be determined using assays known in the art, such as, for example, the Cholesterol Efflux Assays of Takahaski <i>et al.</i> , P. N. A. S., Vol. 96, Issue 20, 11358-11363, Sep. 28, 1999. Useful for cardio- vascular disorders, cholesterol disorders, and Hyperlipidaemia
Preproapolipo- protein	Apoa-1 participates in the reverse transport of cholesterol from tissues to the liver for

“milano” variant 5,721,114 SEQ ID NO:6 of U.S. Patent No. 5,721,114 (SEQ ID NO:441)	excretion by promoting cholesterol efflux from tissues and by acting as a cofactor for the lecithin cholesterol acyltransferase (Lcat). Lipid binding activity can be determined using assays known in the art, such as, for example, the Cholesterol Efflux Assays of Takahaski <i>et al.</i>, P. N. A. S., Vol. 96, Issue 20, 11358-11363, Sep. 28, 1999. Useful for cardio- vascular disorders, cholesterol disorders, and Hyperlipidaemia
Glycodelin-A; Progesterone- associated endometrial protein GeneSeq Accession W00289 WO9628169 P09466/PAEP_HUMAN Glycodelin (SEQ ID NO:365)	Naturally produced female contraceptive that is removed rapidly from the body following 2-3 days production. Uses include contraception Glycodelin-A activity can be determined using the hemizona assay as described in Oehninger, S., Coddington, C. C., Hodgen, G. D., and Seppala, M (1995) Fertil. Steril. 63, 377-383. Naturally derived contraceptive useful for the prevention of pregnancy.
NOGO-A Genbank Accession CAB99248 (SEQ ID NO:366)	NOGO polypeptides are potent inhibitors of neurite growth. Inhibition of Neurite outgrowth. Antagonists to NOGO polypeptides may promote the outgrowth of neurites, thus inducing regeneration of neurons. NOGO-A polypeptide antagonists are useful for the promotion of neural growth, which could be useful in the treatment of neural disorders and dysfunction due to degenerative diseases or trauma; useful in the treatment of neoplastic diseases of the CNS; induce regeneration of neurons or to promote the structural plasticity of the CNS.
NOGO-B Genbank Accession CAB99249 (SEQ ID NO:367)	NOGO polypeptides are potent inhibitors of neurite growth. Inhibition of Neurite outgrowth. Antagonists to NOGO polypeptides may promote the outgrowth of neurites, thus inducing regeneration of neurons. NOGO-B polypeptide antagonists are useful for the promotion of neural growth, which could be useful in the treatment of neural disorders and dysfunction due to degenerative diseases or trauma; useful in the treatment of neoplastic diseases of the CNS; induce regeneration of neurons or to promote the structural plasticity of the CNS.
NOGO-C Genbank Accession CAB99250 (SEQ ID NO:368)	NOGO polypeptides are potent inhibitors of neurite growth. Inhibition of Neurite outgrowth. Antagonists to NOGO polypeptides may promote the outgrowth of neurites, thus inducing regeneration of neurons. NOGO-C polypeptide antagonists are useful for the promotion of neural growth, which could be useful in the treatment of neural disorders and dysfunction due to degenerative diseases or trauma; useful in the treatment of neoplastic diseases of the CNS; induce regeneration of neurons or to promote the structural plasticity of the CNS.
NOGO-66 Receptor Genbank Accession AAG53612 (SEQ ID NO:369)	NOGO polypeptides are potent inhibitors of neurite growth, and are thought to mediate their effects through the NOGO-66 Receptor. Inhibition of Neurite outgrowth by mediating the biological effects of NOGO polypeptides. Soluble NOGO-66 receptor polypeptides may promote the outgrowth of neurites, thus inducing regeneration of neurons. NOGO-66 receptor polypeptides are useful for the promotion of neural growth, which could be useful in the treatment of neural disorders and dysfunction due to degenerative diseases or trauma; useful in the treatment of neoplastic diseases of the CNS; induce regeneration of neurons or to promote the structural plasticity of the CNS.
Antibodies specific for collapsin US Pat. No. 5,416,197 Wildtype collapsin has the sequence:	These antibodies are useful for the promotion of neurite outgrowth Collapsin activity, which is thought to inhibit the outgrowth of neurites, can be assayed in the presence of antibodies specific for collapsing using assays known in the art, such as, for example, the collapse assay disclosed by Luo <i>et al.</i>, Cell 1993 Oct 22; 75(2): 217-27 Useful for the promotion of neural growth, which could be useful in the treatment of neural disorders and dysfunction due to degenerative diseases or trauma.

SEQ ID NO:2 of 5,416,197 (SEQ ID NO:464)	
Humanized Anti-VEGF Antibodies, and fragments thereof WO9845331	These agents have anti-inflammatory and anti-cancer applications VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer
Humanized Anti-VEGF Antibodies, and fragments thereof WO0029584	These agents have anti-inflammatory and anti-cancer applications VEGF activity can be determined using assays known in the art, such as those disclosed in International Publication No. WO0045835, for example. Promotion of growth and proliferation of cells, such as vascular endothelial cells. Hematopoietic and immune disorders. Antagonists may be useful as anti-angiogenic agents, and may be applicable for cancer
Membrane bound proteins GeneSeq. Accession Y66631-Y66765 WO9963088	Cancer, Immune Disorders These proteins can be used for linking bioactive molecules to cells and for modulating biological activities of cells, using the polypeptides for specific targeting. The polypeptide targeting can be used to kill the target cells, e.g. for the treatment of cancers. These proteins are useful for the treatment of immune system disorders. Activities can be determined using assay known in the art, such as, for example, the assays disclosed in International Publication No.WO0121658.
Secreted and Transmembrane polypeptides GeneSeq Accession B44241-B44334 WO0053756	Cancer, Immune Disorders These proteins can be used for linking bioactive molecules to cells and for modulating biological activities of cells, using the polypeptides for specific targeting. The polypeptide targeting can be used to kill the target cells, e.g. for the treatment of cancers. These proteins are useful for the treatment of immune system disorders. Activities can be determined using assay known in the art, such as, for example, the assays disclosed in International Publication No.WO0121658.
Secreted and Transmembrane polypeptides GeneSeq Accession Y41685-Y41774 WO9946281	Cancer, Immune Disorders These proteins can be used for linking bioactive molecules to cells and for modulating biological activities of cells, using the polypeptides for specific targeting. The polypeptide targeting can be used to kill the target cells, e.g. for the treatment of cancers. These proteins are useful for the treatment of immune system disorders. Activities can be determined using assay known in the art, such as, for example, the assays disclosed in International Publication No. WO0121658.
Interleukin 2 (IL-2) SEQ ID NO:548	Metabolic Disease, Type 1 diabetes, Graft-versus-host disease.
Interleukin 15_vA (IL-15_vA) SEQ ID NO:549	Obesity, Metabolic Disease, Diabetes.
Interleukin 15_vB (IL-15_vB) SEQ ID NO:550	Obesity, Metabolic Disease, Diabetes.
Interleukin 15_vC (IL-15_vC)	Obesity, Metabolic Disease, Diabetes.

SEQ ID NO:551	
Interleukin 15_vD (IL15_vD) SEQ ID NO:552	Obesity, Metabolic Disease, Diabetes.
Interleukin 15_vE (IL15_vE) SEQ ID NO:553	Obesity, Metabolic Disease, Diabetes.
Interleukin 15_vF (IL15_vF) SEQ ID NO:565	Obesity, Metabolic Disease, Diabetes.
Interleukin 22 (IL22) SEQ ID NO:554	Metabolic Disease, Diabetic Ulcers, Inflammatory Bowel Diseases.
Fibroblast Growth Factor 1 (FGF1) SEQ ID NO:555	Diabetes, Metabolic Disease, Obesity.
Fibroblast Growth Factor 1_vA (FGF1_vA) SEQ ID NO:556	Diabetes, Metabolic Disease, Obesity. See Nature 513, 436–439 (18 September 2014) doi:10.1038/nature13540.
Fibroblast Growth Factor 1_vB (FGF1_vB) SEQ ID NO:557	Diabetes, Metabolic Disease, Obesity. See Proc Natl Acad Sci U S A. 1991 Apr 1; 88(7): 2893–2897.
Fibroblast Growth Factor 1_vC (FGF1_vC) SEQ ID NO:566	Diabetes, Metabolic Disease, Obesity.
Fibroblast Growth Factor 19_vA (FGF19_vA) SEQ ID NO:558	Chronic liver disease, primary biliary cirrhosis, bile acid-induced liver damage. See Cancer Res. 2014 Jun 15;74(12):3306-16. doi: 10.1158/0008-5472.CAN-14-0208. Epub 2014 Apr 11. Regulates bile acid metabolism without tumorigenicity.

Fibroblast Growth Factor 21 (FGF21) SEQ ID NO:559	Metabolic Disease, Fibrotic Diseases.
Fibroblast Growth Factor 23 (FGF23) SEQ ID NO:560	Hyperphosphatemic familial tumoral calcinosis.
Brain-Derived Neurotrophic Factor (BDNF) SEQ ID NO:561	Neurological diseases (including Alzheimer's Disease, Autism, Huntington's Disease, Parkinson's Disease, and Depression), obesity, metabolic disease.
Serpin Peptidase Inhibitor, Clade B (Ovalbumin), Member 1 (SERPINB1) SEQ ID NO:562	Diabetes.
CASPASE1 SEQ ID NO:563	Diabetes.
Leukemia Inhibitory Factor (LIF) SEQ ID NO:564	Muscular dystrophy, atherosclerosis, kidney disease
Proprotein Convertase Subtilisin/Kexin Type 1 (PCSK1) SEQ ID NO:567	Cardiovascular disease, hypercholesterolemia, heterozygous familial hypercholesterolemia (HeFH), atherosclerotic cardiovascular disease such as heart attacks or strokes, increasing the amount of a functional protein, polypeptide or peptide
Proprotein Convertase Subtilisin/Kexin Type 2 (PCSK2) SEQ ID NO:568	Cardiovascular disease, hypercholesterolemia, heterozygous familial hypercholesterolemia (HeFH), atherosclerotic cardiovascular disease such as heart attacks or strokes, increasing the amount of a functional protein, polypeptide or peptide
Proprotein Convertase Subtilisin/Kexin Type 3 (PCSK3) SEQ ID NO:569	Cardiovascular disease, hypercholesterolemia, heterozygous familial hypercholesterolemia (HeFH), atherosclerotic cardiovascular disease such as heart attacks or strokes, increasing the amount of a functional protein, polypeptide or peptide
Proprotein Convertase Subtilisin/Kexin Type 3 Sol (PCSK3_SOL) SEQ ID NO:570	Cardiovascular disease, hypercholesterolemia, heterozygous familial hypercholesterolemia (HeFH), atherosclerotic cardiovascular disease such as heart attacks or strokes, increasing the amount of a functional protein, polypeptide or peptide

Proprotein Convertase Subtilisin/Kexin Type 4 (PCSK4) SEQ ID NO:571	Cardiovascular disease, hypercholesterolemia, heterozygous familial hypercholesterolemia (HeFH), atherosclerotic cardiovascular disease such as heart attacks or strokes, increasing the amount of a functional protein, polypeptide or peptide
Proprotein Convertase Subtilisin/Kexin Type 5 (PCSK5) SEQ ID NO:572	Cardiovascular disease, hypercholesterolemia, heterozygous familial hypercholesterolemia (HeFH), atherosclerotic cardiovascular disease such as heart attacks or strokes, increasing the amount of a functional protein, polypeptide or peptide
Proprotein Convertase Subtilisin/Kexin Type 6 (PCSK6) SEQ ID NO:573	Cardiovascular disease, hypercholesterolemia, heterozygous familial hypercholesterolemia (HeFH), atherosclerotic cardiovascular disease such as heart attacks or strokes, increasing the amount of a functional protein, polypeptide or peptide
Proprotein Convertase Subtilisin/Kexin Type (PCSK7) SEQ ID NO:574	Cardiovascular disease, hypercholesterolemia, heterozygous familial hypercholesterolemia (HeFH), atherosclerotic cardiovascular disease such as heart attacks or strokes, increasing the amount of a functional protein, polypeptide or peptide
Proprotein Convertase Subtilisin/Kexin Type 8 (PCSK8) SEQ ID NO:575	Cardiovascular disease, hypercholesterolemia, heterozygous familial hypercholesterolemia (HeFH), atherosclerotic cardiovascular disease such as heart attacks or strokes, increasing the amount of a functional protein, polypeptide or peptide
Proprotein Convertase Subtilisin/Kexin Type 9 (PCSK9) SEQ ID NO:576	Cardiovascular disease, hypercholesterolemia, heterozygous familial hypercholesterolemia (HeFH), atherosclerotic cardiovascular disease such as heart attacks or strokes, increasing the amount of a functional protein, polypeptide or peptide
Membrane-Bound Transcription Factor Peptidase, Site 2 (MBTPS2) SEQ ID NO:577	IFAP syndrome, increasing the amount of a functional protein, polypeptide or peptide
Carboxypeptidase E (CPE) SEQ ID NO:578	Endocrine disorders, such as, for example, obesity and infertility; hyperproinsulinemia; metabolic syndrome, increasing the amount of a functional protein, polypeptide or peptide

In various embodiments, the nucleic acid drug, including synthetic RNA, is administered in a manner that it effects one or more of keratinocytes and fibroblasts (e.g. causes these cells to express one or more therapeutic proteins). For example, present methods allow for methods in which a patient's cells are used to generate a therapeutic protein and the levels of such protein are tailored by synthetic RNA dosing.

In a specific embodiment, the synthetic RNA targets a soluble protein. In some embodiments, the synthetic RNA targets a protein of one or more of the following families of proteins: transforming growth factor (TGF) beta, bone

morphogenetic proteins (BMPs), Fibroblast growth factors (FGFs), vascular endothelial growth factors (VEGFs), and interleukins. The terms “family”, “superfamily”, and “subfamily” can be used interchangeably

In a specific embodiment, the synthetic RNA targets a member of the TGF beta family. TGF- β superfamily proteins comprise cytokines characterized by six-conserved cysteine residues (Lander *et al.*, (2001) Nature, 409:860-921). The human genome contains at least about 42 open reading frames encoding TGF- β superfamily proteins. TGF- β superfamily proteins can at least be divided into the BMP subfamily and the TGF- β subfamily based on sequence similarity and the specific signaling pathways that they activate. In various embodiments, the synthetic RNA targets one or more of TGFs (e.g., TGF- β 1, TGF- β 2, and TGF- β 3), activins (e.g., activin A) and inhibins, macrophage inhibitory cytokine-1 (MIC-1), Mullerian inhibiting substance, anti-Mullerian hormone, and glial cell line derived neurotrophic factor (GDNF).

The TGF- β superfamily comprises a subset of the cysteine knot cytokine superfamily. Additional members of the cysteine knot cytokine superfamily include, but are not limited to, platelet derived growth factor (PDGF), vascular endothelial growth factor (VEGF), placenta growth factor (P1GF), Noggin, neurotrophins (BDNF, NT3, NT4, and β NGF), gonadotropin, follitropin, lutropin, interleukin-17, and coagulogen. This family of proteins is also among the targets encompassed by the present invention.

In various embodiments, the present invention relates to targeting a TGF beta family member for treatment or prevention of various immunological disorders, cancer, bronchial asthma, lung fibrosis, heart disease, diabetes, hereditary hemorrhagic telangiectasia, Marfan syndrome, Vascular Ehlers-Danlos syndrome, Loeys-Dietz syndrome, Parkinson's disease, chronic kidney disease, multiple sclerosis and AIDS.

In a specific embodiment, the synthetic RNA targets a member of the BMP family. The BMP subfamily includes, but is not limited to, BMP-2, BMP-3 (osteogenin), BMP-3b (GDF-10), BMP-4 (BMP-2b), BMP-5, BMP-6, BMP-7 (osteogenic protein-1 or OP-1), BMP-8 (OP-2), BMP-8B (OP-3), BMP-9 (GDF-2), BMP-10, BMP-11 (GDF-11), BMP-12 (GDF-7), BMP-13 (GDF-6, CDMP-2), BMP-15 (GDF-9), BMP-16, GDF-1, GDF-3, GDF-5 (CDMP-1), and GDF-8 (myostatin). In various embodiments, the synthetic RNA targets one or more of BMP-2, BMP-3 (osteogenin), BMP-3b (GDF-10), BMP-4 (BMP-2b), BMP-5, BMP-6, BMP-7 (osteogenic protein-1 or OP-1), BMP-8 (OP-2), BMP-8B (OP-3), BMP-9 (GDF-2), BMP-10, BMP-11 (GDF-11), BMP-12 (GDF-7), BMP-13 (GDF-6, CDMP-2), BMP-15 (GDF-9), BMP-16, GDF-1, GDF-3, GDF-5 (CDMP-1), and GDF-8 (myostatin). BMPs are sometimes referred to as Osteogenic Protein (OPs), Growth Differentiation Factors (GDFs), or Cartilage-Derived Morphogenetic Proteins (CDMPs). In a specific embodiment, the synthetic RNA targets one or more BMP fusions (e.g. as described in US Patent Publication No. 2009/0202638, the entire contents of which are hereby incorporated by reference) and/or one or more BMP mutants (e.g. as described in US Patent Publication No. 2011/0039773, the entire contents of which are hereby incorporated by reference).

In various embodiments, the present invention relates to targeting a BMP family member for regenerative medicine or metabolic applications, including without limitation orthopedic applications such as spinal fusions,

nonunions and oral surgery, metabolic disease, prediabetes, diabetes, thermogenesis, insulin sensitivity, insulin resistance, and adipogenesis, including brown-fat adipogenesis. Various other metabolic applications are described elsewhere herein. In various embodiments, the present invention relates to targeting a BMP family member, including BMP-7, for treatment of chronic kidney disease (CKD) and/or to reverse the loss of glomeruli due to sclerosis.

In various embodiments, the present invention relates to targeting a BMP family member to induce proliferation of bone and cartilage in a variety of locations in the body. For example, repair of joints such as knee, elbow, ankle, and finger joints are contemplated by the invention. For example, targeting a BMP family member may result in regenerating cartilage in patients suffering from arthritis or other cartilage degenerating diseases. Further, the invention pertains to treating tears in cartilage due to injury. In addition, the invention is useful for inducing bone growth in patients, for instance, by way of non-limitation, for use in treating patients suffering from bone fractures or breaks, osteoporosis, or patients in need of spinal fusion or for repair of the spine, vertebrae or the like.

In various embodiments, the present invention relates to targeting a BMP family member to induce a developmental cascade of bone morphogenesis and tissue morphogenesis for a variety of tissues in mammals different from bone or bone cartilage. This morphogenic activity includes the ability to induce proliferation and differentiation of progenitor cells, and the ability to support and maintain the differentiated phenotype through the progression of events that results in the formation of bone, cartilage, non-mineralized skeletal or connective tissues, and other adult tissues.

For example, the present invention may be used for treatment to prevent loss of and/or increase bone mass in metabolic bone diseases. General methods for treatment to prevent loss of and/or increase bone mass in metabolic bone diseases using osteogenic proteins are disclosed in US Pat. No. 5,674,844, the entire contents of which are hereby incorporated by reference. Further the present compositions and methods find use in replacing or repairing bone or cartilage at injury sites such as bone breaks, bone fractures, and cartilage tears, periodontal tissue regeneration (e.g. general methods for periodontal tissue regeneration using osteogenic proteins are disclosed in US Pat. No. 5,733,878, the entire contents of which are hereby incorporated by reference), liver regeneration, including following a partial hepatectomy (see, e.g., US Pat. No. 5,849,686, the entire contents of which are hereby incorporated by reference), treatment of chronic renal failure (see, e.g., US Pat. No. 6,861,404, the entire contents of which are hereby incorporated by reference), enhancing functional recovery following central nervous system ischemia or trauma (see, e.g. US Pat. No. 6,407,060, the entire contents of which are hereby incorporated by reference), inducing dendritic growth (see, e.g., US Pat. No. 6,949,505, the entire contents of which are hereby incorporated by reference), inducing neural cell adhesion (see, e.g., US Pat. No. 6,800,603, the entire contents of which are hereby incorporated by reference), and treatment and prevention of Parkinson's disease (see, e.g., US Pat. No. 6,506,729, the entire contents of which are hereby incorporated by reference). As another example, the present compositions and methods, including

when targeting one or more BMPs, can be used to induce dentinogenesis. To date, the unpredictable response of dental pulp tissue to injury is a basic clinical problem in dentistry. Using standard dental surgical procedures, small areas (e.g., 2 mm) of dental pulps can be surgically exposed by removing the enamel and dentin immediately above the pulp (by drilling) of sample teeth, performing a partial amputation of the coronal pulp tissue, inducing hemostasis, application of the pulp treatment, and sealing and filling the cavity by standard procedures.

In various embodiments, the present invention relates to targeting a BMP family member for the treatment of one or more metabolic-related disorders as described herein.

In a specific embodiment, the synthetic RNA targets a member of the FGF family. FGFs are a family of growth factors, with members involved in angiogenesis, wound healing, embryonic development and various endocrine signaling pathways. In various embodiments, any one of the following FGFs are targets of the invention: FGF1, FGF2, FGF3, FGF4, FGF5, FGF6, FGF7, FGF8, FGF9, FGF10, FGF11, FGF12, FGF13, FGF14 (FGF11, FGF12, FGF13, and FGF14 being FGF homologous factors 1-4 (FHF1-FHF4)), FGF16, FGF17, FGF18, FGF19 (aka FHF15/19), FGF20, FGF21, FGF22 and FGF22. In some embodiments, the synthetic RNA targets a cognate receptor of any of the above FGFs. In various embodiments, the present invention relates to targeting a FGF family member to a treat or prevent disease or disorder associated with abnormal function and/or expression of an FGF, a metabolic disease or disorder (e.g., diabetes, obesity, dyslipidemia, hyperglycemia, hyperinsulinemia, hypertension, hepatosteatosis such as non-alcoholic steatohepatitis (NASH) etc.), cancer, a disease or disorder associated with impaired lipid metabolism a disease or disorder associated with impaired renal function, a disease or disorder associated with impaired hepatic function, abnormal cell proliferation, a vascular disease or disorder (e.g., coronary artery disease, peripheral artery disease, atherosclerosis, abdominal aortic aneurysm, a blood clot, deep vein thrombosis, venous stasis disease, phlebitis, varicose veins etc.), angiogenesis, atherosclerosis, a cardiovascular disease or disorder, a disease or disorder associated with impaired blood vessel formation, a disease or disorder associated with impaired cell signaling, a disease or disorder associated with impaired kinase activity, and a disease or disorder associated with impaired uptake of glucose into adipocytes.

In a specific embodiment, the synthetic RNA targets a member of the VEGF family. In some embodiments, the target is one or more of VEGF-A (including all isoforms, e.g. VEGF121, VEGF165 and VEGF189), placenta growth factor (PGF, including all isoforms, e.g. PGF-1, PGF-2, and PGF-3), VEGF-B, VEGF-C and VEGF-D, and any variants thereof (see, e.g. US Patent No. 9,078,860, the entire contents of which are hereby incorporated by reference). In some embodiments, the synthetic RNA targets a cognate receptor of any of the above VEGFs. The present invention also encompasses as targets VEGF-related proteins including orf virus-encoded VEGF-like proteins referred to as VEGF-E and a series of snake venoms referred to as VEGF-F. VEGFs and VEGF-related proteins are members of the Platelet Derived Growth Factor (PDGF) supergene family of cystine knot growth factors. All members of the PDGF supergene family share a high degree of structural homology with PDGF.

In various embodiments, the present invention relates to targeting a VEGF family member to treat diseases and conditions associated with angiogenesis, including but not limited to, solid tumor cancers, hemangiomas, rheumatoid arthritis, osteoarthritis, septic arthritis, asthma, atherosclerosis, idiopathic pulmonary fibrosis, vascular restenosis, arteriovenous malformations, meningiomas, neovascular glaucoma, psoriasis, Kaposi's Syndrome, angiofibroma, hemophilic joints, hypertrophic scars, Osler-Weber syndrome, pyogenic granuloma, retrolental fibroplasias, scleroderma, trachoma, von Hippel-Lindau disease, vascular adhesion pathologies, synovitis, dermatitis, neurological degenerative diseases, preeclampsia, unexplained female infertility, endometriosis, unexplained male infertility, pterygium, wounds, sores, skin ulcers, gastric ulcers, and duodenal ulcers. In various embodiments, the present invention relates to targeting a VEGF family member to treat angiogenesis-associated eye diseases, including without limitation any eye disease associated with abnormal intraocular neovascularization, including but not limited to retinopathy of prematurity, diabetic retinopathy, retinal vein occlusion, and age-related macular degeneration, as well diabetic macular edema and retinal vein occlusion. In an embodiment, the present compositions and methods relate to the treatment of wet age-related macular degeneration.

In a specific embodiment, the synthetic RNA targets a member of the interleukin family. The interleukins represent a large group of cytokines with diverse functions and were first characterized by expression in leukocytes and have since been shown to be expressed in a wide variety of cells, for example macrophages, TH-1 and TH-2 cells, T- lymphocytes, monocytes and bone marrow stroma. Broadly, the function of the immune system depends in a large part on the expression and function of the interleukins. In some embodiments, the target is one or more of interleukins 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 35 and 36, and, within each species of interleukin, various isotypes and/or interleukin receptors (e.g., IL-1R, IL-2R, IL-3R, IL-4R, IL-5R, IL-6R, IL-7R, IL-8R, IL-9R, IL-10R, IL-11R, IL-12R, IL-13R, IL-14R, IL-15R, IL-16R, IL-17R, IL-18R, IL-19R, IL-20R, IL-21R, IL-22R, IL-23R, IL-24R, IL-25R, IL-26, IL-27R, IL-28R, IL-29R, IL-30R, IL-31R, IL-32R, IL-33R, IL-34R, IL-35R, and IL-36R). In a specific embodiment, both IL-15 and IL-15R (e.g., IL-15RA) are targeted. Without wishing to be bound by theory, it is believed that the targeting of both interleukin (e.g., IL-15) and its cognitive interleukin receptor (e.g., IL-15RA) results in synergistic beneficial effects. In various embodiments, the present invention relates to targeting a member of the interleukin family to treat cancer, inflammatory, respiratory, autoimmune, cardiovascular, neurological, metabolic, and/or proliferative diseases, disorders, and/or conditions in a subject or organism, as described herein. In a specific embodiment, the present invention relates to targeting a member of the interleukin family to treat cancer. In a specific embodiment, the present invention relates to targeting a member of the interleukin family to treat rheumatoid arthritis.

In a specific embodiment, the synthetic RNA targets an EPO gene or a derivative thereof (e.g. SEQ ID NO: 164, SEQ ID NO: 165, SEQ ID NO: 166, and SEQ ID NO: 167). Some embodiments are related to NOVEPOETIN protein (SEQ ID NO: 167). Other embodiments are related to NOVECRIT (SEQ ID NO: 168).

Erythropoietin can stimulate erythropoiesis in anemic patients with chronic renal failure in whom the endogenous production of erythropoietin is impaired. Without being bound by theory, because of the length of time often required for erythropoiesis (several days for erythroid progenitors to mature and be released into the circulation), a clinically significant increase in hemoglobin is usually not observed in less than two weeks and may require up to ten weeks in some patients. The present methods and compositions provide, in some embodiments, more rapid therapeutic effect. The present methods and compositions provide, in some embodiments, more rapid therapeutic effect, for instance, when compared to wild type EPO and/or EPO without non-canonical nucleotides and/or EPO delivered as a protein biologic. The present methods and compositions provide, in some embodiments, a sustained therapeutic effect. The present methods and compositions provide, in some
 5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55
 60
 65
 70
 75
 80
 85
 90
 95
 100
 105
 110
 115
 120
 125
 130
 135
 140
 145
 150
 155
 160
 165
 170
 175
 180
 185
 190
 195
 200
 205
 210
 215
 220
 225
 230
 235
 240
 245
 250
 255
 260
 265
 270
 275
 280
 285
 290
 295
 300
 305
 310
 315
 320
 325
 330
 335
 340
 345
 350
 355
 360
 365
 370
 375
 380
 385
 390
 395
 400
 405
 410
 415
 420
 425
 430
 435
 440
 445
 450
 455
 460
 465
 470
 475
 480
 485
 490
 495
 500
 505
 510
 515
 520
 525
 530
 535
 540
 545
 550
 555
 560
 565
 570
 575
 580
 585
 590
 595
 600
 605
 610
 615
 620
 625
 630
 635
 640
 645
 650
 655
 660
 665
 670
 675
 680
 685
 690
 695
 700
 705
 710
 715
 720
 725
 730
 735
 740
 745
 750
 755
 760
 765
 770
 775
 780
 785
 790
 795
 800
 805
 810
 815
 820
 825
 830
 835
 840
 845
 850
 855
 860
 865
 870
 875
 880
 885
 890
 895
 900
 905
 910
 915
 920
 925
 930
 935
 940
 945
 950
 955
 960
 965
 970
 975
 980
 985
 990
 995

For instance, for EPO, the present methods and compositions provide a clinically significant increase in hematocrit in less than about 6 weeks, or less than about 5 weeks, or less than about 4 weeks, or less than about 3 weeks, or less than about 2 weeks, or less than about 1 week. In some embodiments, the present methods and compositions provide a clinically significant increase in hematocrit in about 2 weeks, or about 10 days, or about 1 week, or about 3 days, or about 1 day. In various embodiments, the present methods and compositions accelerate the process by which erythroid progenitors mature and are released into the circulation.

In some embodiments, the present EPO-related compositions find use in decreasing the dose and/or frequency of administration when compared to wild type EPO and/or EPO without non-canonical nucleotides and/or EPO delivered as a protein biologic. For instance, the present EPO-related compositions may find use in treatment regimens for the diseases disclosed herein (including, without limitation, one or more anemias) that involve administration on a monthly, or biweekly, or weekly basis. In some embodiments, therefore, the present EPO-related compositions reduce the need for daily, or, in some embodiments, weekly, administration. In some embodiments, the present EPO-related compositions require lower maintenance doses as compared to wild type EPO and/or EPO without non-canonical nucleotides and/or EPO delivered as a protein biologic. Certain
 20
 25
 30
 35
 40
 45
 50
 55
 60
 65
 70
 75
 80
 85
 90
 95
 100
 105
 110
 115
 120
 125
 130
 135
 140
 145
 150
 155
 160
 165
 170
 175
 180
 185
 190
 195
 200
 205
 210
 215
 220
 225
 230
 235
 240
 245
 250
 255
 260
 265
 270
 275
 280
 285
 290
 295
 300
 305
 310
 315
 320
 325
 330
 335
 340
 345
 350
 355
 360
 365
 370
 375
 380
 385
 390
 395
 400
 405
 410
 415
 420
 425
 430
 435
 440
 445
 450
 455
 460
 465
 470
 475
 480
 485
 490
 495
 500
 505
 510
 515
 520
 525
 530
 535
 540
 545
 550
 555
 560
 565
 570
 575
 580
 585
 590
 595
 600
 605
 610
 615
 620
 625
 630
 635
 640
 645
 650
 655
 660
 665
 670
 675
 680
 685
 690
 695
 700
 705
 710
 715
 720
 725
 730
 735
 740
 745
 750
 755
 760
 765
 770
 775
 780
 785
 790
 795
 800
 805
 810
 815
 820
 825
 830
 835
 840
 845
 850
 855
 860
 865
 870
 875
 880
 885
 890
 895
 900
 905
 910
 915
 920
 925
 930
 935
 940
 945
 950
 955
 960
 965
 970
 975
 980
 985
 990
 995

In some embodiments, the present the present methods and compositions provide a fast and robust response that obviates the need for RBC transfusion. For instance, in some embodiments, the present methods and compositions allow for treatment of patients do not consent to transfusions.

In some embodiments, the present methods and compositions increase the rate of increase in hematocrit. In some embodiments, the present methods and compositions maintain elevated hematocrits (e.g. of 25%, or 30%, or 35%, or 40% or more) for a sustained period (e.g. about 1 month, or about 2 months, or about 3 months, or about 4 months, or about 5 months, or about 6 months, or about 9 months).

In some embodiments, the present methods and compositions stimulate red blood cell production. In some embodiments, the present methods and compositions stimulate division and differentiation of committed erythroid progenitors in the bone marrow.

In some embodiments, including, without limitation, when targeting EPO, the present invention relates to the treatment of one or more of anemia, including anemia resulting from resulting from chronic kidney disease (e.g. from dialysis) and/or chemotherapy and/or HIV treatment (e.g. Zidovudine (INN) or azidothymidine (AZT)), inflammatory bowel disease (e.g. Crohn's disease and ulcer colitis), anemia linked to inflammatory conditions (e.g. arthritis, lupus, IBD), anemia linked to diabetes, schizophrenia, cerebral malaria, as aplastic anemia, and myelodysplasia from the treatment of cancer (e.g. chemotherapy and/or radiation), and various myelodysplastic syndrome diseases (e.g. sickle cell anemia, hemoglobin SC disease, hemoglobin C disease, alpha- and beta-thalassemias, neonatal anemia after premature birth, and comparable conditions).

In some embodiments, including, without limitation, when targeting EPO, the present invention relates to the treatment of, or a patient having one or more of cancer, heart failure, autoimmune disease, sickle cell disease, thalassemia, blood loss, transfusion reaction, diabetes, vitamin B12 deficiency, collagen vascular disease, Shwachman syndrome, thrombocytopenic purpura, Celiac disease, endocrine deficiency state such as hypothyroidism or Addison's disease, autoimmune disease such as Crohn's Disease, systemic lupus erythematosus, rheumatoid arthritis or juvenile rheumatoid arthritis, ulcerative colitis immune disorders such as eosinophilic fasciitis, hypogammaglobulinemia, or thymoma/thymic carcinoma, graft vs. host disease, preleukemia, Nonhematologic syndrome (e.g. Down's, Dubowitz, Seckel), Felty syndrome, hemolytic uremic syndrome, myelodysplastic syndrome, nocturnal paroxysmal hemoglobinuria, osteomyelofibrosis, pancytopenia, pure red-cell aplasia, Schoenlein-Henoch purpura, malaria, protein starvation, menorrhagia, systemic sclerosis, liver cirrhosis, hypometabolic states, congestive heart failure, chronic infections such as HIV/ AIDS, tuberculosis, osteomyelitis, hepatitis B, hepatitis C, Epstein-barr virus or parvovirus, T cell leukemia virus, bacterial overgrowth syndrome, fungal or parasitic infections, and/or red cell membrane disorders such as hereditary spherocytosis, hereditary elliptocytosis, hereditary pyropoikilocytosis, hereditary stomatocytosis, red cell enzyme defects, hypersplenism, immune hemolysis or paroxysmal nocturnal hemoglobinuria.

In some embodiments, including, without limitation, when targeting EPO, the present invention relates to the treatment of, or a patient having anemia, *i.e.* a condition in which the number of red blood cells and/or the amount of hemoglobin found in the red blood cells is below normal. In various embodiments, the anemia may be acute or chronic. For example, the present anemias include but are not limited to iron deficiency anemia, renal anemia, anemia of chronic diseases/inflammation, pernicious anemia such as macrocytic achylic anemia, juvenile pernicious anemia and congenital pernicious anemia, cancer-related anemia, chemotherapy-related anemia, radiotherapy-related anemia, pure red cell aplasia, refractory anemia with excess of blasts, aplastic anemia, X-linked sideroblastic anemia, hemolytic anemia, sickle cell anemia, anemia caused by impaired

production of ESA, myelodysplasia syndromes, hypochromic anemia, microcytic anemia, sideroblastic anemia, autoimmune hemolytic anemia, Cooley's anemia, Mediterranean anemia, Diamond Blackfan anemia, Fanconi's anemia and drug-induced immune hemolytic anemia. Anemia may cause serious symptoms, including hypoxia, chronic fatigue, lack of concentration, pale skin, low blood pressure, dizziness and heart failure.

5 In some embodiments, the anemia is induced by chemotherapy is one cause of anemia. For instance, the chemotherapy may be any myelosuppressive chemotherapy. In some embodiments, the chemotherapy is one or more platinum-based drugs including cisplatin (e.g. PLATINOL) and carboplatin (e.g. PARAPLATIN). In some embodiments, the chemotherapy is any one of the agents described herein. In some embodiments, the chemotherapy is any agent described in Groopman *et al.* J Natl Cancer Inst (1999) 91 (19): 1616-1634, the
10 contents of which are hereby incorporated by reference in their entireties. In some embodiments, the present compositions and methods are used in the treatment of chemotherapy-related anemia in later stage cancer patients (e.g. a stage IV, or stage III, or stage II cancer). In some embodiments, the present compositions and methods are used in the treatment of chemotherapy-related anemia in cancer patients receiving dose-dense chemotherapy or other aggressive chemotherapy regimens.

15 In some embodiments, the present invention relates to the treatment of anemia in a patient having one or more blood-based cancers, such as leukemia, lymphoma, and multiple myeloma. Such cancers may affect the bone marrow directly. Further, the present invention relates to metastatic cancer that has spread to the bone or bone marrow. In some embodiments, the present invention relates to the treatment of anemia in a patient undergoing radiation therapy. Such radiation therapy may damage the bone marrow, lowering its ability to make red blood
20 cells. In further embodiments, the present invention relates to the treatment of anemia in a patient having a reduction or deficiency of one or more of iron, vitamin B12, and folic acid. In further embodiments, the present invention relates to the treatment of anemia in a patient having excessive bleeding including without limitation, after surgery or from a tumor that is causing internal bleeding. In further embodiments, the present invention relates to the treatment of anemia in a patient having anemia of chronic disease.

25 In some embodiments, the present invention relates to the treatment of anemia resulting from chronic renal failure. In some embodiments, the present invention relates to the treatment of anemia resulting from the use of one or more renal replacement therapies, inclusive of dialysis, hemodialysis, peritoneal dialysis, hemofiltration, hemodiafiltration, and renal transplantation.

In some embodiments, the present invention relates to the treatment of anemia in patients with chronic kidney
30 disease who are not on dialysis. For instance, the present invention relates to patients in stage 1 CKD, or stage 2 CKD, or stage 3 CKD, or stage 4 CKD, or stage 5 CKD. In some embodiments, the present patient is stage 4 CKD or stage 5 CKD. In some embodiments, the present patient has undergone a kidney transplant. In some embodiments, the present invention relates to the treatment of anemia is a patient having an acute kidney injury (AKI).

In various embodiments, the present compositions and methods are used to reduce or eliminate fatigue, dizziness, and shortness of breath in a patient.

In various embodiments, the present compositions and methods are used to treat a patient presenting with hyporesponse or resistance to erythropoiesis stimulating agent therapy. In some embodiments, hyporesponsiveness to erythropoietin or ESA-resistant anemia refers to the presence of at least one of the following conditions: i) a significant decrease in hemoglobin levels at a constant dose of ESA treatment, ii) a significant increase in the ESA dose requirement to achieve or maintain a certain hemoglobin level, iii) a failure to raise the hemoglobin level to the target range despite the ESA dose equivalent to erythropoietin greater than 150 IU/kg/week or 0.75 mg/kg/week of darbepoietin-alpha or continued need for such high dose of ESA to maintain the target hemoglobin level. For example, approximately 5-10% of patients with CDK demonstrate hyporesponsiveness to ESA, defined as a continued need for greater than 300 IU/kg per week erythropoietin or 1.5 ng/kg per week darbepoietin administered by the subcutaneous route.

In various embodiments, the present compositions and methods mitigate the need for dose-escalation of the erythropoiesis stimulating agent therapy and therefore, optionally, avoid side effects (e.g. flu-like symptoms such as joint pains, weakness, dizziness and tiredness, skin irritation, increased risk of adverse cardiovascular complications).

In various embodiments, the present compositions and methods are used to maintain a hemoglobin level of about 12.5 to 13 g/dL. In various embodiments, the present compositions and methods are used in patients having hemoglobin levels of below about 12 g/dL, or about 11 g/dL, or about 10 g/dL, or about 9 g/dL, or about 8 g/dL, or about 7 g/dL, or about 6 g/dL, or about 5 g/dL. In various embodiments, the present compositions and methods are used in patients having iron blood test scores that indicate blood pathology, e.g. a ferritin score of below about 200 ng/L and/or a transferrin saturation score below about 30%.

In various embodiments, the present compositions and methods are used to increase or maintain hemoglobin levels at a target level ranging from 9 to 10 g/dL, at a target level ranging from 9 g/dL to 11 g/dL, at a target level ranging from 9 g/dL to 12 g/dL, at a target level ranging from 9 g/dL to 14 g/dL, at a target level ranging from 10 g/dL to 14 g/dL, or at a target level ranging from 12 g/dL to 14 g/dL.

In various embodiments, the present compositions and methods are used to bring a patient's hemoglobin levels to normal. In various embodiments, normal hemoglobin ranges for humans are about 14-18 g/dl for men and 12-16 for women g/dl with the average hemoglobin value for men at about 16 g/dL and for women at about 14 g/dL.

In some embodiments, for instance when targeting EPO, the present invention relates to the treatment of anemia of one or more of the following toxicity grading criteria (e.g. NCI Common Toxicity Criteria): grade 1 (mild), 10.0 g hemoglobin/dL to within normal limits; grade 2 (moderate), 8.0-10.0 g of hemoglobin/dL; grade 3 (serious or severe), 6.5-7.9 g of hemoglobin/dL; and grade 4 (life threatening), less than 6.5 g of hemoglobin/dL. In various

embodiments, the present invention brings an increase in toxicity grading criteria by about 1 point, or about 2 points, or about 3 points, or about 4 points. In various embodiments, the present invention results in a patient having a level of 0 or 1. In various embodiments, the present compositions and methods improve anemia as assessed by one or more scales described in Groopman *et al.* J Natl Cancer Inst (1999) 91 (19): 1616-1634, the
 5 entire contents of which are hereby incorporated by reference in their entireties.

In some embodiments, when targeting EPO, the present invention relates to combination therapy with one or more EPOs. For example, the present compositions may provide a sustained effect that can supplement a fast action of another EPO. In some embodiments, the present compositions are used as an adjuvant to other EPOs. In some embodiments, the present compositions are used as a maintenance therapy to other EPOs. Other EPOs
 10 include the following: epoetin alfa, including without limitation, DARBEPOETIN (ARANESP), EPOCEPT (LUPIN PHARMA), NANOKINE (NANOGEN PHARMACEUTICAL), EPOFIT (INTAS PHARMA), EPOGEN (AMGEN), EPOGIN, EPREX, (JANSSEN-CILAG), BINOCRIT (SANDOZ), PROCRIT; epoetin beta, including without limitation, NEORECORMON (HOFFMANN-LA ROCHE), RECORMON, Methoxy polyethylene glycol-epoetin beta (MIRCERA, ROCHE); epoetin delta, including without limitation, DYNEPO (erythropoiesis stimulating
 15 protein, SHIRE PLC); epoetin omega, including without limitation, EPOMAX; epoetin zeta, including without limitation, SILAPO (STADA) and RETACRIT (HOSPIRA) and other EPOs, including without limitation, EPOCEPT (LUPIN PHARMACEUTICALS), EPOTRUST (PANACEA BIOTEC LTD), ERYPRO SAFE (BIOCON LTD.), REPOITIN (SERUM INSTITUTE OF INDIA LIMITED), VINTOR (EMCURE PHARMACEUTICALS), EPOFIT (INTAS PHARMA), ERYKINE (INTAS BIOPHARMACEUTICA), WEPOX (WOCKHARDT BIOTECH), ESPOGEN
 20 (LG LIFE SCIENCES), RELIPOIETIN (RELIANCE LIFE SCIENCES), SHANPOIETIN (SHANTHA BIOTECHNICS LTD), ZYROP (CADILA HEALTHCARE LTD.), EPIAO (RHUEPO) (SHENYANG SUNSHINE PHARMACEUTICAL CO. LTD), CINNAPOIETIN (CINNAGEN).

In some embodiments, when targeting EPO, the present invention relates to combination therapy with a blood transfusion. For instance, the present compositions may supplement a blood transfusion. In some embodiments,
 25 when targeting EPO, the present invention relates to combination therapy with iron supplements.

In some embodiments, the present invention also relates to the following protein targets, e.g. in the treatment of disease: growth hormone (GH) e.g. human and bovine growth hormone, growth hormone-releasing hormones; interferon including α -, β -, or γ -interferons, etc., interleukin-I; interleukin-II; erythropoietin including α - and β -erythropoietin (EPO), granulocyte colony stimulating factor (GCSF), granulocyte macrophage colony stimulating
 30 factor (GM-CSF), anti-angiogenic proteins (e.g., angiostatin, endostatin) PACAP polypeptide (pituitary adenylate cyclase activating polypeptide), vasoactive intestinal peptide (VIP), thyrotrophin releasing hormone (TRH), corticotropin releasing hormone (CRH), vasopressin, arginine vasopressin (AVP), angiotensin, calcitonin, atrial natriuretic factor, somatostatin, adrenocorticotropin, gonadotropin releasing hormone, oxytocin, insulin, somatotropin, plasminogen tissue activator, coagulation factors including coagulation factors VIII and IX,
 35 glucosylceramidase, sargramostim, lenograstin, filgrastin, dornase- α , molgramostim, PEG-L-asparaginase, PEG-

adenosine deaminase, hirudin, eptacog- α (human blood coagulation factor VIIa) nerve growth factors, transforming growth factor, epidermal growth factor, basic fibroblast growth factor, VEGF; heparin including low molecular weight heparin, calcitonin; antigens; monoclonal antibodies; vancomycin; desferrioxamine (DFO); parathyroid hormone, an immunogen or antigen, and an antibody such as a monoclonal antibody.

- 5 In some embodiments, the present methods allow for effective additional therapeutic agent (e.g. those described herein) activity and/or targeting to a cell and/or tissue of interest. For example, the present synthetic RNA can lead to increased expression of one or more targeting molecules that direct an additional therapeutic to the location of therapy. For example, the additional therapeutic agent may have a binding partner that the synthetic RNA encodes. For example, the synthetic RNA may induce the expression of an antigen that directs the
- 10 therapeutic activity of an antibody that may be used in combination (e.g. herceptin, rituxan, campath, gemtuzumab, herceptin, panorex, rituximab, bexxar, edrecolomab, alemtuzumab, mylotrag, IMC-C225, smartin 195, and mitomomab). In some embodiments, the synthetic RNA can be injected directly into one or more of the tumors described herein and home the therapeutic antibody to the tumor.

- In some embodiments, the present methods allow for effective additional therapeutic agent generation, especially
- 15 when the additional therapeutic agent is a prodrug, for example, to produce an active form of the drug. In some embodiments, the synthetic RNA can be injected directly into one or more of the tumors described herein and home the prodrug to the tumor. For instance, the synthetic RNA may encode an enzyme that catalyzes the localized conversion of a non-toxic, systemically delivered agent into a potent chemotherapeutic agent. By way of illustration (note that any of the prodrugs or drugs listed herein are additional agents as used herein):

Enzyme	Prodrug	Drug
aldehyde oxidases	5-Ethynyl-2(1H)-pyrimidinone	5-Ethynyluracil
aldehyde oxidases	IPdR	IUdR
aldehyde oxidases	5-FP	5-FU
amino acid oxidases	d-alanine	Hydrogen peroxide
amino acid oxidases	SeCys conjugates	Selenols and hydrogen peroxide
cytochrome P450 reductase	Menadione	Semiquinone radical
cytochrome P450 reductase	Mitomycin C	Quinone methide intermediate
cytochrome P450 reductase	Tirapazamine	Nitroxide radical
cytochrome P450 reductase	EO9	Unidentified semiquinone radical
DT-diaphorase	Streptonigrin	Unidentified
DT-diaphorase	Mitomycin C	Quinone methide intermediate
DT-diaphorase	CB 1954	5-(Aziridin-1-yl)-4-hydroxyl-amino-2-nitrobenzamide
DT-diaphorase	Diaziquone	Semiquinone radical ^a
cytochrome P450	Ipomeanol	Unidentified (possibly furan epoxide)
cytochrome P450	Ftorafur (tegafur)	5-FU
cytochrome P450	Dacarbazine	HHMTIC
cytochrome P450	Trofosfamide	Trofosfamide mustard
cytochrome P450	Ifosfamide	Isophosphamide mustard
cytochrome P450	Cyclophosphamide	Phosphoramidate mustard
cytochrome P450	AQ4N	AQ4
tyrosinase	2,4-Dihydroxyphenylalanine	6-Hydroxydopa

tyrosinase	4-S-CAP	BQ
tyrosinase	GHB	GBQ
tyrosinase	Substituted phenols	Orthoquinones
tyrosinase	Phenyl mustards	Phenol mustard
tyrosinase	Urea mustards	Unidentified
glutathione S-transferase	TER286	Aziridinium agent
glutathione S-transferase	S-CPHC-ethylsulfoxide	S-CPHC-glutathione
glutathione S-transferase	PTA	6-MP
carboxylesterase	CPT-11	SN-38
carboxylesterase	Paclitaxel-2-ethylcarbonate	Paclitaxel
carboxylesterase	Capecitabine	5'-Deoxy-5-fluorocytidine (5-FU)
carboxylesterase	Tertiary amidomethyl esters	Carboxylic acids and amines ^a
alkaline phosphatase	Amifostine	WR-1065
alkaline phosphatase	3-AP phosphate	3-AP
β -glucuronidase	BHAMG	pHAM
β -glucuronidase	Epirubicin-glucuronide	Epirubicin
β -glucuronidase	HMR 1826	Doxorubicin
β -glucuronidase	DNR-GA3	Daunorubicin
β -glucuronidase	DOX-GA3	Doxorubicin
β -glucuronidase	Paclitaxel glucuronide	Paclitaxel
β -glucuronidase	5-FU glucuronide	5-FU
cysteine conjugate β -lyase	PC	6-MP
cysteine conjugate β -lyase	GC	6-Thioguanine
cysteine conjugate β -lyase	SeCys conjugate	Selenol
Nitroreductase	CB 1954	5-(Aziridin-1-yl)-4- hydroxyl-amino-2-nitro- benzamide
Cytochrome P450	4-Ipomeanol	Unidentified (furan epoxide is speculated)
Cytochrome P450	Ifosfamide	Isophosphoramidate mustard
Cytochrome P450	Cyclophosphamide	Phosphoramidate mustard
Purine-nucleoside phosphorylase	Fludarabine	2-Fluoroadenine
Purine-nucleoside phosphorylase	MeP-dR	MeP
Thymidine kinase	Ganciclovir	Ganciclovir-triphosphate nucleotide
Alkaline phosphatase	Etoposide phosphate	Etoposide
Alkaline phosphatase	Mitomycin C phosphate	Mitomycin C
Alkaline phosphatase	POMP	POM
Alkaline phosphatase	N-(4-phosphonooxy)-phenylacetyl)doxorubicin	Doxorubicin
β -Glucuronidase	Glucuronidated Nor nitrogen mustard	Oxazolidinone
β -Glucuronidase	Glucuronidated 9-amino-camptothecin	9-Aminocamptothecin
β -Glucuronidase	Glucuronide mustard	Mustard
Carboxypeptidase	Methotrexate-amino acids	Methotrexate
Carboxypeptidase	CMDA	Benzoic acid mustard
Penicillin amidase	DPO	Doxorubicin
Penicillin amidase	MeIPO	Melphalan
Penicillin amidase	NHPAP	Palytoxin
Penicillin amidase	N-(phenylacetyl) doxorubicin	Doxorubicin
Penicillin amidase	N-(phenylacetyl) melphalan	Melphalan
β -Lactamase	LY 266070	DAVLBHYD

β -Lactamase	C-DOX	Doxorubicin
β -Lactamase	PRODOX	Doxorubicin
β -Lactamase	CM	Phenylenediamine mustard
β -Lactamase	CCM	Phenylenediamine mustard
β -Lactamase	Cephalosporin-DACCP	DACCP
β -Lactamase	PROTAX	Taxol
β -Lactamase	Cephalosporin mitomycin C	Mitomycin C
β -Lactamase	C-Mel	Melphalan
Cytosine deaminase	5-Fluorocytosine	5-Fluorouracil
Methionine γ -lyase	Selenomethionine	Methylselenol
Methionine γ -lyase	Trifluoromethionine	CSF ₂

In certain embodiments, the synthetic RNA may encode an enzyme that catalyzes the conversion of 5-FU and/or Doxorubicin from various prodrugs, as illustrated by the examples below:

5-FU Prodrugs and Enzymes

5-FP	Aldehyde oxidase
Ftorafur	P450
5'-DFUR	Thymidine phosphorylase
5-FU glucuronide	β -Glucuronidase
5-FC	Cytosine deaminase

Doxorubicin Prodrugs and Enzymes

N-(4-phosphono- oxy)-phenylacetyl) doxorubicin	Alkaline phosphatase
HMR 1826	β -Glucuronidase
DOX-GA3	β -Glucuronidase
DPO	Penicillin amidase
N-(phenylacetyl) doxorubicin	Penicillin amidase
C-DOX	β -Lactamase
PRODOX	β -Lactamase

- 5 In some embodiments, the nucleic acid drugs at the doses and regimens described herein may be used in combination with one or more additional agents (aka adjuvant therapy or combination agent). In some embodiments, the nucleic acid drugs at the doses and regimens described herein may be used in a human patient undergoing treatment with one or more additional agents. In some embodiments, the nucleic acid drug is used as an adjuvant or neoadjuvant to any of the additional agents described herein. In some embodiments, the invention pertains to co-administration and/or co-formulation. Any of the compositions described herein may be co-formulated and/or co-administered. In some embodiments, any nucleic acid drug described herein acts synergistically when co-administered with another agent and may be administered at doses that are lower than the doses commonly employed when such agents are used as monotherapy.

- 15 In some embodiments, any nucleic acid drug described herein may include derivatives that are modified, *i.e.*, by the covalent attachment of any type of molecule to the composition such that covalent attachment does not prevent the activity of the composition. For example, but not by way of limitation, derivatives include composition that have been modified by, *inter alia*, glycosylation, lipidation, acetylation, pegylation, phosphorylation,

amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications can be carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of turicamycin, etc. Additionally, the derivative can contain one or more non-classical amino acids. In various embodiments, one or
 5 more additional agents (aka adjuvant therapy or combination agent) may be conjugated to any nucleic acid drug described herein.

Contacting a cell with a steroid can suppress the innate immune response to foreign nucleic acids, and can increase the efficiency of nucleic acid delivery and translation. Certain embodiments are therefore directed to contacting a cell with a steroid. Other embodiments are directed to administering a steroid to a patient. Illustrative
 10 steroids include corticosteroid steroids. In some embodiments, the steroid is one or more of cortisone, hydrocortisone, prednisone, prednisolone, dexamethasone, triamcinolone, and betamethasone. In one embodiment, the steroid is hydrocortisone. In another embodiment, the steroid is dexamethasone.

Other embodiments are directed to administering to a patient a member of the group: an antibiotic, an antimycotic, and an RNase inhibitor.

15 Botulinum toxin type A has been approved by the US Food and Drug Administration (FDA) for the treatment of essential blepharospasm, strabismus and hemifacial spasm in patients over the age of twelve, cervical dystonia, glabellar line (facial) wrinkles and for treating hyperhidrosis and botulinum toxin type B has been approved for the treatment of cervical dystonia. The present compositions may be combined with these toxins in the treatment of these diseases and related diseases. Some embodiments are directed to a nucleic acid drug targeting a
 20 neurotoxin. In various embodiments, the neurotoxin is a botulinum toxin or a biologically active fragment, variant, analogue or family-member thereof.

Further the combination of any one of the aforementioned toxins may be used in combination with the present compositions for various cosmetic procedures, including, without limitation, facial wrinkles, hyperkinetic skin lines, glabellar lines, crow's feet, marionette lines, skin disorders, nasolabial folds, blepharospasm, strabismus,
 25 hemifacial spasms and sweating disorders. Alternatively, the present compositions may be used to in these cosmetic procedures as a monotherapy.

Certain embodiments are directed to a combination therapy comprising one or more of the therapeutic or cosmetic compositions of the present invention and one or more adjuvant therapies or cosmetic treatments. In certain embodiments, one or more of the therapeutic or cosmetic compositions of the present invention are
 30 administered to a subject which is undergoing treatment with one or more adjuvant therapies or cosmetic treatments. Example adjuvant therapies and cosmetic treatments are set forth in Table 3 and Table 5 of US Provisional Application No. 61/721,302, the contents of which are hereby incorporated by reference, and are given by way of example, and not by way of limitation.

Table 3. Illustrative Adjuvant Therapies

Therapy/Treatment Class	Disease/Condition	Example Therapy/Treatment
Acetylcholinesterase inhibitors	Myasthenia gravis, Glaucoma, Alzheimer's disease, Lewy body dementia, Postural tachycardia syndrome	Edrophonium
Angiotensin-converting-enzyme inhibitor	Hypertension, Congestive heart failure	Perindopril
Alkylating agents	Cancer	Cisplatin
Angiogenesis inhibitors	Cancer, Macular degeneration	Bevacizumab
Angiotensin II receptor antagonists	Hypertension, Diabetic nephropathy, Congestive heart failure	Valsartan
Antibiotics	Bacterial infection	Amoxicillin
Antidiabetic drugs	Diabetes	Metformin
Antimetabolites	Cancer, Infection	5-fluorouracil (5FU)
Antisense oligonucleotides	Cancer, Diabetes, Amyotrophic lateral sclerosis (ALS), Hypercholesterolemia	Mipomersen
Cytotoxic antibiotics	Cancer	Doxorubicin
Deep-brain stimulation	Chronic pain, Parkinson's disease, Tremor, Dystonia	N/A
Dopamine agonists	Parkinson's disease, Type II diabetes, Pituitary tumors	Bromocriptine
Entry/Fusion inhibitors	HIV/AIDS	Maraviroc
Glucagon-like peptide-1 agonists	Diabetes	Exenatide
Glucocorticoids	Asthma, Adrenal insufficiency, Inflammatory diseases, Immune diseases, Bacterial meningitis	Dexamethasone
Immunosuppressive drugs	Organ transplantation, Inflammatory diseases, Immune diseases	Azathioprine
Insulin/Insulin analogs	Diabetes	NPH insulin
Integrase inhibitors	HIV/AIDS	Raltegravir
MAO-B inhibitors	Parkinson's disease, Depression, Dementia	Selegiline
Maturation inhibitors	HIV/AIDS	Bevirimat
Nucleoside analog reverse-transcriptase inhibitors	HIV/AIDS, Hepatitis B	Lamivudine
Nucleotide analog reverse-transcriptase inhibitors	HIV/AIDS, Hepatitis B	Tenofovir
Non-nucleoside reverse-transcriptase inhibitors	HIV/AIDS	Rilpivirine
Pegylated interferon	Hepatitis B/C, Multiple sclerosis	Interferon beta-1a
Plant alkaloids/terpenoids	Cancer	Paclitaxel
Protease inhibitors	HIV/AIDS, Hepatitis C, Other viral infections	Telaprevir
Radiotherapy	Cancer	Brachytherapy
Renin inhibitors	Hypertension	Aliskiren
Statins	Hypercholesterolemia	Atorvastatin
Topoisomerase inhibitors	Cancer	Topotecan
Vasopressin receptor antagonist	Hyponatremia, Kidney disease	Tolvaptan
Dermal filler	Wrinkles, aged skin	Hyaluronic Acid

Botulinum toxin	Wrinkles, aged skin	botulinum toxin type A
Induction of skin repair	Acne scars, aged skin	Laser treatment, dermabrasion

In some embodiments, the additional agent is a cytotoxic agent, comprising, in illustrative embodiments, a toxin, a chemotherapeutic agent, a radioisotope, and an agent that causes apoptosis or cell death.

Illustrative cytotoxic agents include, but are not limited to, methotrexate, aminopterin, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine; alkylating agents such as mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU), mitomycin C, lomustine (CCNU), 1-methylnitrosourea, cyclophosphamide, mechlorethamine, busulfan, dibromomannitol, streptozotocin, mitomycin C, cis-dichlorodiamine platinum (II) (DDP) cisplatin and carboplatin (paraplatin); anthracyclines include daunorubicin (formerly daunomycin), doxorubicin (adriamycin), detorubicin, carminomycin, idarubicin, epirubicin, mitoxantrone and bisantrene; antibiotics include dactinomycin (actinomycin D), bleomycin, calicheamicin, mithramycin, and anthramycin (AMC); and antimetabolic agents such as the vinca alkaloids, vincristine and vinblastine. Other cytotoxic agents include paclitaxel (taxol), ricin, pseudomonas exotoxin, gemcitabine, cytochalasin B, gramicidin D, ethidium bromide, emetine, etoposide, teniposide, colchicin, dihydroxy anthracin dione, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, puromycin, procarbazine, hydroxyurea, asparaginase, corticosteroids, mytostane (O,P'-(DDD)), interferons, and mixtures of these cytotoxic agents.

Further cytotoxic agents include, but are not limited to, chemotherapeutic agents such as carboplatin, cisplatin, paclitaxel, gemcitabine, calicheamicin, doxorubicin, 5-fluorouracil, mitomycin C, actinomycin D, cyclophosphamide, vincristine, bleomycin, VEGF antagonists, EGFR antagonists, platins, taxols, irinotecan, 5-fluorouracil, gemcitabine, leucovorine, steroids, cyclophosphamide, melphalan, vinca alkaloids (e.g., vinblastine, vincristine, vindesine and vinorelbine), mustines, tyrosine kinase inhibitors, radiotherapy, sex hormone antagonists, selective androgen receptor modulators, selective estrogen receptor modulators, PDGF antagonists, TNF antagonists, IL-1 antagonists, interleukins (e.g. IL-12 or IL-2), IL-12R antagonists, Toxin conjugated monoclonal antibodies, tumor antigen specific monoclonal antibodies, Erbitux, Avastin, Pertuzumab, anti-CD20 antibodies, Rituxan, ocrelizumab, ofatumumab, DXL625, HERCEPTIN, or any combination thereof. Toxic enzymes from plants and bacteria such as ricin, diphtheria toxin and Pseudomonas toxin may be conjugated to the therapeutic agents (e.g. antibodies) to generate cell-type-specific-killing reagents (Youle, *et al.*, Proc. Nat'l Acad. Sci. USA 77:5483 (1980); Gilliland, *et al.*, Proc. Nat'l Acad. Sci. USA 77:4539 (1980); Krolick, *et al.*, Proc. Nat'l Acad. Sci. USA 77:5419 (1980)).

Other cytotoxic agents include cytotoxic ribonucleases as described by Goldenberg in US Pat. No. 6,653,104. Embodiments of the invention also relate to radioimmunoconjugates where a radionuclide that emits alpha or beta particles is stably coupled to the antibody, or binding fragments thereof, with or without the use of a

complex-forming agent. Such radionuclides include beta-emitters such as Phosphorus-32, Scandium-47, Copper-67, Gallium-67, Yttrium-88, Yttrium-90, Iodine-125, Iodine-131, Samarium-153, Lutetium-177, Rhenium-186 or Rhenium-188, and alpha-emitters such as Astatine-211, Lead-212, Bismuth-212, Bismuth-213 or Actinium-225.

5 Illustrative detectable moieties further include, but are not limited to, horseradish peroxidase, acetylcholinesterase, alkaline phosphatase, beta-galactosidase and luciferase. Further exemplary fluorescent materials include, but are not limited to, rhodamine, fluorescein, fluorescein isothiocyanate, umbelliferone, dichlorotriazinylamine, phycoerythrin and dansyl chloride. Further exemplary chemiluminescent moieties include, but are not limited to, luminol. Further exemplary bioluminescent materials include, but are not limited to, luciferin
10 and aequorin. Further exemplary radioactive materials include, but are not limited to, Iodine-125, Carbon-14, Sulfur-35, Tritium and Phosphorus-32.

The dosage of any additional agent described herein as well as the dosing schedule can depend on various parameters, including, but not limited to, the human patient's general health, and the administering physician's and/or human patient's discretion. Co-administration may be simultaneous or sequential. Any additional agent
15 described herein, can be administered prior to (e.g., about 5 minutes, about 15 minutes, about 30 minutes, about 45 minutes, about 1 hour, about 2 hours, about 4 hours, about 6 hours, about 12 hours, about 24 hours, about 48 hours, about 72 hours, about 96 hours, about 1 week, about 2 weeks, about 3 weeks, about 4 weeks, about 5 weeks, about 6 weeks, 8 weeks, or about 12 weeks before), concurrently with, or subsequent to (e.g., about 5 minutes, about 15 minutes, about 30 minutes, about 45 minutes, about 1 hour, about 2 hours, about 4 hours,
20 about 6 hours, about 12 hours, about 24 hours, about 48 hours, about 72 hours, about 96 hours, about 1 week, about 2 weeks, about 3 weeks, about 4 weeks, about 5 weeks, about 6 weeks, about 8 weeks, or about 12 weeks after) the administration of the nucleic acid drug to a human patient in need thereof. In various embodiments any agent described herein is administered about 1 minute apart, about 10 minutes apart, about 30 minutes apart, less than about 1 hour apart, about 1 hour apart, about 1 hour to about 2 hours apart, about 2
25 hours to about 3 hours apart, about 3 hours to about 4 hours apart, about 4 hours to about 5 hours apart, about 5 hours to about 6 hours apart, about 6 hours to about 7 hours apart, about 7 hours to about 8 hours apart, about 8 hours to about 9 hours apart, about 9 hours to about 10 hours apart, about 10 hours to about 11 hours apart, about 11 hours to about 12 hours apart, no more than about 24 hours apart or no more than about 48 hours apart.

30 In a specific embodiment, the combination regimen is designed to exploit the finding that the nucleic acid drug dosing and formulation of the present invention has potent effects quickly (e.g. in about 6, or about 12, or about 24, or about 30, or about 36, or about 48 hours) and the effects can last about 7 days, or longer.

The dose of a nucleic acid drug is disclosed herein. In general, the dose of any additional agent that is useful is known to those in the art. For example, doses may be determined with reference *Physicians' Desk Reference*,
35 66th Edition, PDR Network; 2012 Edition (December 27, 2011), the contents of which are incorporated by

reference in its entirety. In some embodiments, the present invention allows a patient to receive doses that exceed those determined with reference *Physicians' Desk Reference*. The dosage of any additional agent described herein can depend on several factors including the severity of the condition, whether the condition is to be treated or prevented, and the age, weight, and health of the human patient to be treated. Additionally, 5 pharmacogenomic (the effect of genotype on the pharmacokinetic, pharmacodynamic or efficacy profile of a therapeutic) information about a particular human patient may affect dosage used. Furthermore, the exact individual dosages can be adjusted somewhat depending on a variety of factors, including the specific combination of the agents being administered, the time of administration, the route of administration, the nature of the formulation, the rate of excretion, the particular disease being treated, the severity of the disorder, and the 10 anatomical location of the disorder. Some variations in the dosage can be expected.

Cells, tissues, organs, and organisms, including, but not limited to, humans, have several characteristics that can inhibit or prevent the delivery of nucleic acids, including, for example, the stratum corneum, which can serve as a barrier to foreign organisms and nucleic acids. These characteristics can thus inhibit the effects of therapeutics and cosmetics comprising nucleic acids. It has now been discovered that many of these characteristics can be 15 circumvented or overcome using a patch comprising a flexible membrane and a plurality of needles, and that such a patch can serve as an effective and safe article for the delivery of nucleic acids. Certain embodiments are therefore directed to a nucleic acid delivery patch. In one embodiment, the nucleic acid delivery patch comprises a flexible membrane. In another embodiment, the nucleic acid delivery patch comprises a plurality of needles. In yet another embodiment, the plurality of needles are attached to the flexible membrane. In some embodiments, 20 the patch comprises a nucleic acid. In one embodiment, the nucleic acid is present in solution. In one embodiment, the plurality of needles include one or more needles having a lumen. In another embodiment, the patch further comprises a second flexible membrane. In yet another embodiment, the flexible membrane and the second flexible membrane are arranged to form a cavity. In a further embodiment, the cavity contains a nucleic acid. In a still further embodiment, the membrane comprises one or more holes through which a nucleic acid can 25 pass. In a still further embodiment, one or more holes and one or more needles having a lumen are arranged to allow the passage of a solution containing a nucleic acid through at least one of the one or more holes and through at least one of the one or more needles having a lumen. In some embodiments, the patch is configured to deliver a solution to the skin. In one embodiment, the solution comprises a nucleic acid. In another embodiment, the solution comprises a vehicle. In yet another embodiment, the vehicle is a lipid or lipidoid. In a 30 still further embodiment, the vehicle is a lipid-based transfection reagent.

The cell membrane can serve as a barrier to foreign nucleic acids. It has now been discovered that combining the patch of the present invention with an electric field can increase the efficiency of nucleic acid delivery. Certain embodiments are therefore directed to a nucleic acid delivery patch comprising a plurality of needles, wherein at least two needles form part of a high-voltage circuit. Certain embodiments are directed to an implantable "tattoo" 35 for microneedle delivery (see, e.g. *Nature Materials* 12, pp 367–376 (2013), the contents of which are hereby

incorporated by reference in their entirety). In one embodiment, the high-voltage circuit generates a voltage greater than about 10V. In another embodiment, the high-voltage circuit generates a voltage greater than about 20V. In yet another embodiment, an electric field is produced between two of the needles. In a further embodiment, the magnitude of the electric field is at least about 100V/cm. In a still further embodiment, the magnitude of the electric field is at least about 200V/cm. In some embodiments, the patch is configured to deliver a nucleic acid to the epidermis. In other embodiments, the patch is configured to deliver a nucleic acid to the dermis. In still other embodiments, the patch is configured to deliver a nucleic acid to sub-dermal tissue. In still other embodiments, the patch is configured to deliver a nucleic acid to muscle. Certain embodiments are directed to a nucleic acid delivery patch comprising a plurality of electrodes. In one embodiment, the plurality of electrodes is attached to a flexible membrane. Other embodiments are directed to a nucleic acid delivery patch comprising a rigid structure. In one embodiment, a plurality of electrodes are attached to the rigid structure.

In some embodiments, the compositions described herein are administered using an array of needles covering an affected area of the subject. In some embodiments, the treatment area is mechanically massaged after administration. In some embodiments, the treatment area is exposed to electric pulses after administration. In some embodiments, the electric pulses are between about 10V and about 200V for from about 50 microseconds to about 1 second. In some embodiments, the electric pulses are generated around the treatment area by a multielectrode array.

In some embodiments, the present invention provides a patch delivery system, comprising a non-viral RNA transfection composition enclosed within a membrane, and an array of delivery needles delivering from about 10 ng to about 2000 ng of RNA per treatment area of about 100 cm² or less, or about 50 cm² or less, or about 10 cm² or less, or about 5 cm² or less, or about 1 cm² or less, or about 0.5 cm² or less, or about 0.2 cm² or less. In some embodiments, the non-viral transfection composition contains from about 10 ng to about 2000 ng per injection volume of about 20 μ L to about 1 ml. In some embodiments, each needle delivers an injection volume of between 1 μ L and 500 μ L.

In some embodiments, the delivery patch comprises an acrylic reservoir that holds the nucleic acid drug. In some embodiments, a silicon adhesive is added to create a semisolid suspension of microscopic, concentrated drug cells. Further, some embodiments provide a patch that is associated with one or more enhancers (these include, without limitation, iontophoresis, ultrasound, chemicals including gels, microneedles, sonophoresis, lasers, and electroporatic methods).

In some embodiments, the delivery is effected via a gel, optionally a hydro alcoholic gel containing a combination of enhancers (e.g. COMBIGEL (ANTARES PHARMA)).

In various embodiments, the RNA is delivered using needle arrays. Illustrative needle arrays include, but are not limited to AdminPen 600 and those described in U.S. Patent Nos. 7,658,728, 7,785,301, and 8,414,548, the entire disclosure of which are hereby incorporated by reference. Other examples of

needles include, for example, the 3M™ Hollow Microstructured Transdermal System and the 3M Solid Microstructured Transdermal Systems (sMTS). See, e.g. US Patent Nos. 3,034,507 and 3,675,766; Microneedles for Transdermal Drug Delivery. *Advanced Drug Delivery Reviews*. 56: 581-587 (2004); *Pharm Res*. 2011 Jan; 28(1): 31–40, the entire contents of which are hereby incorporated by reference in their entireties.

- 5 In some embodiments, microneedles and/or microneedle arrays may be used. In various embodiments, the microneedles and/or microneedle arrays may be, without limitation, solid, RNA-coated, dissolving, biodegradable, and/or hollow. In some embodiments, the delivery is effected via a microneedle system, optionally combined with an electronically controlled micropump that delivers the drug at specific times or upon demand. For example, the MACROFLUX (Alza) system may be used.
- 10 Other embodiments are directed to a method for delivering a nucleic acid to a cell *in vivo* comprising applying a nucleic acid to a tissue containing a cell *in vivo*. In one embodiment, the method further comprises applying a transient electric field in the vicinity of the cell. In another embodiment, the method results in the cell *in vivo* internalizing the nucleic acid. In yet another embodiment, the nucleic acid comprises synthetic RNA. In a further embodiment, the method further results in the cell internalizing a therapeutically or cosmetically effective amount
- 15 of the nucleic acid. In one embodiment, the cell is a skin cell. In another embodiment, the cell is a muscle cell. In yet another embodiment, the cell is a dermal fibroblast. In a further embodiment, the cell is a keratinocyte. In a still further embodiment, the cell is a myoblast. In some embodiments, the nucleic acid comprises a protein of interest. In one embodiment, the protein of interest is a fluorescent protein. In another embodiment, the protein of interest is an extracellular-matrix protein. In yet another embodiment, the protein of interest is a member of the
- 20 group: elastin, collagen, laminin, fibronectin, vitronectin, lysyl oxidase, elastin binding protein, a growth factor, fibroblast growth factor, transforming growth factor beta, granulocyte colony-stimulating factor, a matrix metalloproteinase, an actin, fibrillin, microfibril-associated glycoprotein, a lysyl-oxidase-like protein, platelet-derived growth factor, a lipase, an uncoupling protein, thermogenin, filaggrin, a fibroblast growth factor, an antibody, and a protein involved with pigment production. In some embodiments, the method further comprises
- 25 delivering the nucleic acid to the epidermis. In other embodiments, the method further comprises delivering the nucleic acid to the dermis. In still other embodiments, the method further comprises delivering the nucleic acid below the dermis. In one embodiment, the delivering is by injection. In another embodiment, the delivering is by injection using a micro-needle array. In yet another embodiment, the delivering is by topical administration. In a further embodiment, the delivering comprises disruption or removal of a part of the tissue. In a still further
- 30 embodiment, the delivering comprises disruption or removal of the stratum corneum. In some embodiments, the nucleic acid is present in solution. In one embodiment, the solution comprises a growth factor. In another embodiment, the growth factor is a member of the group: a fibroblast growth factor and a transforming growth factor. In yet another embodiment, the growth factor is a member of the group: basis fibroblast growth factor and transforming growth factor beta. In other embodiments, the solution comprises cholesterol.

In another embodiment, the method further comprises contacting the cell with one or more nucleic acid molecules. In yet another embodiment, at least one of the one or more nucleic acid molecules encodes a protein of interest. In a further embodiment, the method results in the cell expressing the protein of interest. In a still further embodiment, the method results in the cell expressing a therapeutically or cosmetically effective amount of the protein of interest.

In another embodiment, the cell is contacted with a nucleic acid molecule. In yet another embodiment, the method results in the cell internalizing the nucleic acid molecule. In a further embodiment, the method results in the cell internalizing a therapeutically or cosmetically effective amount of the nucleic acid molecule. In one embodiment, the nucleic acid encodes a protein of interest. In one embodiment, the nucleic acid molecule comprises a member of the group: a dsDNA molecule, a ssDNA molecule, a RNA molecule, a dsRNA molecule, a ssRNA molecule, a plasmid, an oligonucleotide, a synthetic RNA molecule, a miRNA molecule, an mRNA molecule, and an siRNA molecule. In various embodiments, the RNA comprises one or more non-canonical nucleotides.

In some embodiments, the present invention relates to one or more administration techniques described in US Patent Nos. 5,711,964; 5,891,468; 6,316,260; 6,413,544; 6,770,291; and 7,390,780, the entire contents of which are hereby incorporated by reference in their entireties.

The invention also provides kits that can simplify the administration of the nucleic acid drugs described herein and/or any additional agent described herein. An illustrative kit of the invention comprises a nucleic acid drug and/or any additional agent described herein in unit dosage form. In one embodiment, the unit dosage form is a container, such as a pre-filled syringe, which can be sterile, containing any agent described herein and a pharmaceutically acceptable carrier, diluent, excipient, or vehicle. The kit can further comprise a label or printed instructions instructing the use of any agent described herein. The kit or one or more components of the kit may be stored at room temperature, about 4°C, about -20°C, about -80°C, or about -196°C. The kit may also include a lid speculum, topical anesthetic, and a cleaning agent for the administration location. The kit can also further comprise one or more additional agent described herein. In one embodiment, the kit comprises a container containing an effective amount of a nucleic acid drug as disclosed herein and an effective amount of another composition, such as an additional agent as described herein. In some embodiments, the unit dosage form is a pre-loaded (a.k.a. pre-dosed or pre-filled) syringe or a pen needle injector (injection pen)). Such unit dosage forms may comprise the effective doses of nucleic acid drug described herein, e.g. about 10 ng to about 2000 ng, e.g. about 10 ng, or about 20 ng, or about 50 ng, or about 100 ng, or about 200 ng, or about 300 ng, or about 400 ng, or about 500 ng, or about 600 ng, or about 700 ng, or about 800 ng, or about 900 ng, or about 1000 ng, or about 1100 ng, or about 1200 ng, or about 1300 ng, or about 1400 ng, or about 1500 ng, or about 1600 ng, or about 1700 ng, or about 1800 ng, or about 1900 ng, or about 2000 ng.

Some embodiments are directed to synthetic RNA molecules with low toxicity and high translation efficiency.

Other embodiments are directed to a cell-culture medium for high-efficiency *in vivo* transfection, reprogramming,

and gene editing of cells. Other embodiments pertain to methods for producing synthetic RNA molecules encoding reprogramming proteins. Still further embodiments pertain to methods for producing synthetic RNA molecules encoding gene-editing proteins.

Some embodiments are directed to methods of gene-editing and/or gene correction. Some embodiments encompass synthetic RNA-based gene-editing and/or gene correction, e.g. with RNA comprising non-canonical nucleotides, e.g. RNA encoding one or more of a nuclease, a transcription activator-like effector nuclease (TALEN), a zinc-finger nuclease, a meganuclease, a nickase, a clustered regularly interspaced short palindromic repeat (CRISPR)-associated protein or a natural or engineered variant, family-member, orthologue, fragment or fusion construct thereof. In some embodiments, the efficiency of the gene-editing and/or gene correction is high, for example, higher than DNA-based gene editing and/or gene correction. In some embodiments, the present methods of gene-editing and/or gene correction are efficient enough for in vivo application. In some embodiments, the present methods of gene-editing and/or gene correction are efficient enough to not require cellular selection (e.g. selection of cells that have been edited). In various embodiments, the efficiency of gene-editing of the present methods is about 1%, or about 2%, or about 3%, or about 4%, or about 5%, or about 6%, or about 7%, or about 8%, or about 9%, or about 10%, or about 20%, or about 30%, or about 40%, or about 50%, or about 60%, or about 70%, or about 80%, or about 90%, or about 100%. In various embodiments, the efficiency of gene-correction of the present methods is about 1%, or about 2%, or about 3%, or about 4%, or about 5%, or about 6%, or about 7%, or about 8%, or about 9%, or about 10%, or about 20%, or about 30%, or about 40%, or about 50%, or about 60%, or about 70%, or about 80%, or about 90%, or about 100%

Some embodiments are directed to high-efficiency gene-editing proteins comprising engineered nuclease cleavage domains. Other embodiments are directed to high-fidelity gene-editing proteins comprising engineered nuclease cleavage domains. Various embodiments are directed to high-efficiency gene-editing proteins comprising engineered DNA-binding domains. Other embodiments are directed to high-fidelity gene-editing proteins comprising engineered DNA-binding domains. Still other embodiments are directed to gene-editing proteins comprising engineered repeat sequences. Some embodiments are directed to gene-editing proteins comprising one or more CRISPR associated family members. Some embodiments are directed to methods for altering the DNA sequence of a cell by transfecting the cell with or inducing the cell to express a gene-editing protein. Other embodiments are directed to methods for altering the DNA sequence of a cell that is present in an *in vitro* culture. Still further embodiments are directed to methods for altering the DNA sequence of a cell that is present *in vivo*.

Some embodiments are directed to methods of modulating the secretion or subcellular localization of a polypeptide. In such embodiments, the present invention provides an RNA encoding a protein comprising a signal peptide operably linked to a polypeptide having biological functionality. In an embodiment, the signal peptide modulates the secretion of the polypeptide. In another embodiment, the signal peptide modulates the subcellular localization of the polypeptide. For example, BMP7 comprising at least one FGF21 signal peptide

may result in increased secretion of BMP7. In another example, BMP7 comprising at least one FGF21 signal peptide in place of the endogenous BMP7 signal peptide may result in increased secretion of BMP7. In various embodiments, any of the proteins listed in **Table 2A** or **Table 2B** comprising at least one signal peptide listed in the table below may result in increased secretion of the proteins.

Protein	Signal Peptide
<i>Gaussia</i> luciferase	MGVKVLFALICIAVAEA
Human BMP7	MHVRSRLRAAAPHSFVALWAPLFLLRSALA
Human chymotrypsinogen B	MAFLWLLSCWALLGTTFG
Human chymotrypsinogen C	MLGITVLAALLACASS
Human EPO	MGVHECPAWLWLLLSLLSLPLGLPVLG
Human FGF19	MMSGCVVHVWILAGLWLAVAGRP
Human FGF21	MDSDETGFEHSGLWVSVLAGLLLGACQA
Human FGF23	MLGARLRLWVCALCSVCSMSVLRA
Human IL2	MYRMQLLSCIALSLALVTNS
Human IL22	MAALQKSVSSFLMGTLATSCLLLLALLVQGGAA
Human IL6	MNSFSTSAFGPVAFSLGLLLVLPAAFPAP
Human Interferon Alpha-2	MALTFALLVALLVLSCKSSCSVG
Human Interferon Beta	MTNKCLLQIALLLCFSTTALS
Human Interferon Gamma	MKYTSYILAFQLCIVLGSLGACYC
Human Trypsin-1	MNPLLILTFVAAALA

5

Some embodiments are directed to methods of modulating the serum half-life, secretion, bioavailability, and/or activity of a protein comprising administering a RNA encoding the protein and a RNA encoding a receptor for the protein. For example, IL15 may be administered with its receptor, e.g., IL15RA to enhance one or more of the serum half-life, secretion, bioavailability, and activity of IL15.

- 10 Many proteins and peptides, when translated from in vitro transcribed RNA, can exhibit reduced activity resulting from incomplete or inadequate post-translational processing. It has now been discovered that the amount of active protein, polypeptide or peptide produced following RNA transfection can be increased by contacting cells with and/or inducing cells to express a second protein that is capable of processing a polypeptide into the protein, peptide or polypeptide. Certain embodiments are therefore directed to a method for inducing a cell to
- 15 express an active protein, polypeptide or peptide comprising contacting the cell with a synthetic RNA molecule encoding a first polypeptide and contacting the cell with a second protein that is capable of processing the first polypeptide into an active protein, polypeptide or peptide.

In certain embodiments, the second protein is administered to a patient. In one embodiment, the second protein is a recombinant protein. In another embodiment, the cells are contacted with a synthetic RNA molecule

20 encoding the second protein. In a further embodiment, the cells are contacted with and/or induced to express an

inhibitor of a molecule that inhibits the second protein. In one embodiment, the inhibitor is a short interfering RNA molecule.

In certain embodiments, the second protein is a member of the PCSK family. In other embodiments, the second protein is a proprotein convertase. In still other embodiments, the second protein is a prohormone convertase. In still other embodiments, the second protein is a carboxypeptidase. In one embodiment, the second protein is PCSK3 (Furin/PACE). In another embodiment, the second protein is primarily secreted. In yet another embodiment, the second protein is primarily intracellular. In a further embodiment, the first polypeptide, protein, polypeptide or peptide is selected from the group: a product of proopiomelanocortin, renin, a product of enkephalin, a product of prodynorphin, somatostatin, insulin, agouti-related peptide, glucagon, parathyroid hormone, a member of the transforming growth factor beta superfamily, albumin, beta-secretase 1, nerve growth factor, caldesmon, the alpha-integrins, factor IX, α -melanocyte-stimulating hormone, adrenocorticotrophic hormone, β -endorphin, and met-enkefalin.

Glycation and glycosylation are processes by which one or more sugar molecules are bound to a protein. It has now been discovered that altering the number or location of glycation and glycosylation sites can increase or decrease the stability of a protein. Certain embodiments are therefore directed to a protein with one or more glycation or glycosylation sites. In one embodiment, the protein is engineered to have more glycation or glycosylation sites than a natural variant of the protein. In another embodiment, the protein is engineered to have fewer glycation or glycosylation sites than a natural variant of the protein. In yet another embodiment, the protein has increased stability. In yet another embodiment, the protein has decreased stability. In some embodiments, the protein is a circulating protein. In one embodiment, the protein is erythropoietin or a biologically active fragment, variant, analogue, or family-member thereof. In another embodiment, the protein is darbepoetin alfa or a biologically active fragment, variant, analogue, or family-member thereof. In another embodiment, the protein is NOVEPOETIN or a biologically active fragment, variant, analogue, or family-member thereof.

It has been further discovered that in certain situations, including one or more steroids and/or one or more antioxidants in the transfection medium can increase *in vivo* transfection efficiency, *in vivo* reprogramming efficiency, and *in vivo* gene-editing efficiency. Certain embodiments are therefore directed to contacting a cell or patient with a glucocorticoid, such as hydrocortisone, prednisone, prednisolone, methylprednisolone, dexamethasone or betamethasone. Other embodiments are directed to a method for inducing a cell to express a protein of interest by contacting a cell with a medium containing a steroid and contacting the cell with one or more nucleic acid molecules. In one embodiment, the nucleic acid molecule comprises synthetic RNA. In another embodiment, the steroid is hydrocortisone. In yet another embodiment, the hydrocortisone is present in the medium at a concentration of between about 0.1uM and about 10uM, or about 1uM. Other embodiments are directed to a method for inducing a cell *in vivo* to express a protein of interest by contacting the cell with a medium containing an antioxidant and contacting the cell with one or more nucleic acid molecules. In one embodiment, the antioxidant is ascorbic acid or ascorbic-acid-2-phosphate. In another embodiment, the ascorbic

acid or ascorbic-acid-2-phosphate is present in the medium at a concentration of between about 0.5mg/L and about 500mg/L, including about 50mg/L. Still other embodiments are directed to a method for reprogramming and/or gene-editing a cell *in vivo* by contacting the cell with a medium containing a steroid and/or an antioxidant and contacting the cell with one or more nucleic acid molecules, wherein the one or more nucleic acid molecules
 5 encodes one or more reprogramming and/or gene-editing proteins. In certain embodiments, the cell is present in an organism, and the steroid and/or antioxidant are delivered to the organism.

Adding transferrin to the complexation medium has been reported to increase the efficiency of plasmid transfection in certain situations. It has now been discovered that adding transferrin to the complexation medium can also increase the efficiency of *in vivo* transfection with synthetic RNA molecules. Certain embodiments are
 10 therefore directed to a method for inducing a cell *in vivo* to express a protein of interest by adding one or more synthetic RNA molecules and a transfection reagent to a solution containing transferrin. In one embodiment, the transferrin is present in the solution at a concentration of between about 1mg/L and about 100mg/L, such as about 5mg/L. In another embodiment, the transferrin is recombinant.

In certain situations, including pertaining to culturing, it may be desirable to replace animal-derived components
 15 with non-animal-derived and/or recombinant components, in part because non-animal-derived and/or recombinant components can be produced with a higher degree of consistency than animal-derived components, and in part because non-animal-derived and/or recombinant components carry less risk of contamination with toxic and/or pathogenic substances than do animal-derived components. Certain embodiments are therefore directed to a protein that is non-animal-derived and/or recombinant. Other embodiments are directed to a
 20 medium, wherein some or all of the components of the medium are non-animal-derived and/or recombinant.

Other embodiments are directed to a method for transfecting a cell *in vivo*. In one embodiment, a cell *in vivo* is transfected with one or more nucleic acids, and the transfection is performed using a transfection reagent, such as a lipid-based transfection reagent. In one embodiment, the one or more nucleic acids includes at least one RNA molecule. In another embodiment, the cell is transfected with one or more nucleic acids, and the one or
 25 more nucleic acids encodes at least one of: p53, TERT, an antibody, an extracellular matrix protein, a cytokine, a secreted protein, a membrane-bound protein, an enzyme, a gene-editing protein, a chromatin-modifying protein, a DNA-binding protein, a transcription factor, a histone deacetylase, a pathogen-associated molecular pattern, and a tumor-associated antigen or a biologically active fragment, analogue, variant or family-member thereof. In another embodiment, the cell is transfected repeatedly, such as at least about 2 times during about 10
 30 consecutive days, or at least about 3 times during about 7 consecutive days, or at least about 4 times during about 6 consecutive days. Some embodiments are directed to a method for increasing expression of telomerase in one of a fibroblast, a hematopoietic stem cell, a mesenchymal stem cells, a cardiac stem cell, a hair follicle stem cell, a neural stem cell, an intestinal stem cell, an endothelial stem cell, an olfactory stem cell, a neural crest stem cell, a testicular cell, and a keratinocyte. Some embodiments are directed to a method for increasing the
 35 length of telomeres in one of a fibroblast, a hematopoietic stem cell, a mesenchymal stem cells, a cardiac stem

cell, a hair follicle stem cell, a neural stem cell, an intestinal stem cell, an endothelial stem cell, an olfactory stem cell, a neural crest stem cell, a testicular cell, and a keratinocyte. Other embodiments are directed to a method for isolating a cell from a patient, contacting the cell with a nucleic acid drug encoding a component of telomerase (e.g., TERT), and reintroducing the cell to the patient. Various embodiments are directed to a method

5 for increasing the replicative potential of a cell.

Reprogramming can be performed by transfecting cells with one or more nucleic acids encoding one or more reprogramming factors. Examples of reprogramming factors include, but are not limited to: Oct4 protein, Sox2 protein, Klf4 protein, c-Myc protein, l-Myc protein, TERT protein, Nanog protein, Lin28 protein, Ulf1 protein, Aicda protein, miR200 micro-RNA, miR302 micro-RNA, miR367 micro-RNA, miR369 micro-RNA and biologically active

10 fragments, analogues, variants and family-members thereof. Certain embodiments are therefore directed to a method for reprogramming a cell *in vivo*. In one embodiment, the cell *in vivo* is reprogrammed by transfecting the cell with one or more nucleic acids encoding one or more reprogramming factors. In one embodiment, the one or more nucleic acids includes an RNA molecule that encodes Oct4 protein. In another embodiment, the one or more nucleic acids also includes one or more RNA molecules that encodes Sox2 protein, Klf4 protein, and c-Myc

15 protein. In yet another embodiment, the one or more nucleic acids also includes an RNA molecule that encodes Lin28 protein. In one embodiment, the cell is a human skin cell, and the human skin cell is reprogrammed to a pluripotent stem cell. In another embodiment, the cell is a human skin cell, and the human skin cell is reprogrammed to a glucose-responsive insulin-producing cell. Examples of other cells that can be reprogrammed and other cells to which a cell can be reprogrammed include, but are not limited to: skin cells, pluripotent stem

20 cells, mesenchymal stem cells, β -cells, retinal pigmented epithelial cells, hematopoietic cells, cardiac cells, airway epithelial cells, neural stem cells, neurons, glial cells, bone cells, blood cells, and dental pulp stem cells. In one embodiment, the cell is contacted with a medium that supports the reprogrammed cell. In one embodiment, the medium also supports the cell.

Importantly, infecting skin cells with viruses encoding Oct4, Sox2, Klf4, and c-Myc, combined with culturing the

25 cells in a medium that supports the growth of cardiomyocytes, has been reported to cause reprogramming of the skin cells to cardiomyocytes, without first reprogramming the skin cells to pluripotent stem cells (See Efs *et al* Nat Cell Biol. 2011;13:215-22, the contents of which are hereby incorporated by reference). In certain situations, direct reprogramming (reprogramming one somatic cell to another somatic cell without first reprogramming the somatic cell to a pluripotent stem cell, also known as "transdifferentiation") may be desirable, in part because

30 culturing pluripotent stem cells can be time-consuming and expensive, the additional handling involved in establishing and characterizing a stable pluripotent stem cell line can carry an increased risk of contamination, and the additional time in culture associated with first producing pluripotent stem cells can carry an increased risk of genomic instability and the acquisition of mutations, including point mutations, copy-number variations, and karyotypic abnormalities. Certain embodiments are therefore directed to a method for reprogramming a somatic

cell *in vivo*, wherein the cell is reprogrammed to a somatic cell, and wherein a characterized pluripotent stem-cell line is not produced.

It has been further discovered that, in certain situations, fewer total transfections may be required to reprogram a cell according to the methods of the present invention than according to other methods. Certain embodiments are therefore directed to a method for reprogramming a cell *in vivo*, wherein between about 1 and about 12 transfections are performed during about 20 consecutive days, or between about 4 and about 10 transfections are performed during about 15 consecutive days, or between about 4 and about 8 transfections are performed during about 10 consecutive days. It is recognized that when a cell is contacted with a medium containing nucleic acid molecules, the cell may likely come into contact with and/or internalize more than one nucleic acid molecule either simultaneously or at different times. A cell can therefore be contacted with a nucleic acid more than once, e.g. repeatedly, even when a cell is contacted only once with a medium containing nucleic acids.

Of note, nucleic acids can contain one or more non-canonical or "modified" residues as described herein. For instance, any of the non-canonical nucleotides described herein can be used in the present reprogramming methods. In one embodiment, pseudouridine-5'-triphosphate can be substituted for uridine-5'-triphosphate in an *in vitro*-transcription reaction to yield synthetic RNA, wherein up to 100% of the uridine residues of the synthetic RNA may be replaced with pseudouridine residues. *In vitro*-transcription can yield RNA with residual immunogenicity, even when pseudouridine and 5-methylcytidine are completely substituted for uridine and cytidine, respectively (see, e.g., Angel. Reprogramming Human Somatic Cells to Pluripotency Using RNA [Doctoral Thesis]. Cambridge, MA: MIT; 2011, the contents of which are hereby incorporated by reference). For this reason, it is common to add an immunosuppressant to the transfection medium when transfecting cells with RNA. In certain situations, adding an immunosuppressant to the transfection medium may not be desirable, in part because the recombinant immunosuppressant most commonly used for this purpose, B18R, can be expensive and difficult to manufacture. It has now been discovered that cells *in vivo* can be transfected and/or reprogrammed according to the methods of the present invention, without using B18R or any other immunosuppressant. It has been further discovered that reprogramming cells *in vivo* according to the methods of the present invention without using immunosuppressants can be rapid, efficient, and reliable. Certain embodiments are therefore directed to a method for transfecting a cell *in vivo*, wherein the transfection medium does not contain an immunosuppressant. Other embodiments are directed to a method for reprogramming a cell *in vivo*, wherein the transfection medium does not contain an immunosuppressant. In certain situations, for example when using a high cell density, it may be beneficial to add an immunosuppressant to the transfection medium. Certain embodiments are therefore directed to a method for transfecting a cell *in vivo*, wherein the transfection medium contains an immunosuppressant. Other embodiments are directed to a method for reprogramming a cell *in vivo*, wherein the transfection medium contains an immunosuppressant. In one embodiment, the immunosuppressant is B18R or a biologically active fragment, analogue, variant or family-member thereof or dexamethasone or a derivative thereof. In one embodiment, the transfection medium does not

contain an immunosuppressant, and the nucleic-acid dose is chosen to prevent excessive toxicity. In another embodiment, the nucleic-acid dose is less than about 1mg/cm² of tissue or less than about 1mg/100,000 cells or less than about 10mg/kg.

Reprogrammed cells produced according to certain embodiments of the present invention are suitable for therapeutic and/or cosmetic applications as they do not contain undesirable exogenous DNA sequences, and they are not exposed to animal-derived or human-derived products, which may be undefined, and which may contain toxic and/or pathogenic contaminants. Furthermore, the high speed, efficiency, and reliability of certain embodiments of the present invention may reduce the risk of acquisition and accumulation of mutations and other chromosomal abnormalities. Certain embodiments of the present invention can thus be used to generate cells that have a safety profile adequate for use in therapeutic and/or cosmetic applications. For example, reprogramming cells using RNA and the medium of the present invention, wherein the medium does not contain animal or human-derived components, can yield cells that have not been exposed to allogeneic material. Certain embodiments are therefore directed to a reprogrammed cell that has a desirable safety profile. In one embodiment, the reprogrammed cell has a normal karyotype. In another embodiment, the reprogrammed cell has fewer than about 5 copy-number variations (CNVs) relative to the patient genome, such as fewer than about 3 copy-number variations relative to the patient genome, or no copy-number variations relative to the patient genome. In yet another embodiment, the reprogrammed cell has a normal karyotype and fewer than about 100 single nucleotide variants in coding regions relative to the patient genome, or fewer than about 50 single nucleotide variants in coding regions relative to the patient genome, or fewer than about 10 single nucleotide variants in coding regions relative to the patient genome.

Endotoxins and nucleases can co-purify and/or become associated with other proteins, such as serum albumin. Recombinant proteins, in particular, can often have high levels of associated endotoxins and nucleases, due in part to the lysis of cells that can take place during their production. Endotoxins and nucleases can be reduced, removed, replaced or otherwise inactivated by many of the methods of the present invention, including, for example, by acetylation, by addition of a stabilizer such as sodium octanoate, followed by heat treatment, by the addition of nuclease inhibitors to the albumin solution and/or medium, by crystallization, by contacting with one or more ion-exchange resins, by contacting with charcoal, by preparative electrophoresis or by affinity chromatography. It has now been discovered that partially or completely reducing, removing, replacing or otherwise inactivating endotoxins and/or nucleases from a medium and/or from one or more components of a medium can increase the efficiency with which cells can be transfected and reprogrammed. Certain embodiments are therefore directed to a method for transfecting a cell *in vivo* with one or more nucleic acids, wherein the transfection medium is treated to partially or completely reduce, remove, replace or otherwise inactivate one or more endotoxins and/or nucleases. Other embodiments are directed to a medium that causes minimal degradation of nucleic acids. In one embodiment, the medium contains less than about 1EU/mL, or less than about 0.1EU/mL, or less than about 0.01EU/mL.

In certain situations, protein-based lipid carriers such as serum albumin can be replaced with non-protein-based lipid carriers such as methyl-beta-cyclodextrin. The medium of the present invention can also be used without a lipid carrier, for example, when transfection is performed using a method that may not require or may not benefit from the presence of a lipid carrier, for example, using one or more lipid-based transfection reagents, polymer-based transfection reagents or peptide-based transfection reagents or using electroporation. Many protein-associated molecules, such as metals, can be highly toxic to cells *in vivo*. This toxicity can cause decreased viability, as well as the acquisition of mutations. Certain embodiments thus have the additional benefit of producing cells that are free from toxic molecules.

The associated-molecule component of a protein can be measured by suspending the protein in solution and measuring the conductivity of the solution. Certain embodiments are therefore directed to a medium that contains a protein, wherein about a 10% solution of the protein in water has a conductivity of less than about 500 $\mu\text{mho/cm}$. In one embodiment, the solution has a conductivity of less than about 50 $\mu\text{mho/cm}$. In another embodiment, less than about 0.65% of the dry weight of the protein comprises lipids and/or less than about 0.35% of the dry weight of the protein comprises free fatty acids.

The amount of nucleic acid delivered to cells *in vivo* can be increased to increase the desired effect of the nucleic acid. However, increasing the amount of nucleic acid delivered to cells *in vivo* beyond a certain point can cause a decrease in the viability of the cells, due in part to toxicity of the transfection reagent. It has now been discovered that when a nucleic acid is delivered to a population of cells *in vivo* in a fixed volume (for example, cells in a region of tissue), the amount of nucleic acid delivered to each cell can depend on the total amount of nucleic acid delivered to the population of cells and to the density of the cells, with a higher cell density resulting in less nucleic acid being delivered to each cell. In certain embodiments, a cell *in vivo* is transfected with one or more nucleic acids more than once. Under certain conditions, for example when the cells are proliferating, the cell density may change from one transfection to the next. Certain embodiments are therefore directed to a method for transfecting a cell *in vivo* with a nucleic acid, wherein the cell is transfected more than once, and wherein the amount of nucleic acid delivered to the cell is different for two of the transfections. In one embodiment, the cell proliferates between two of the transfections, and the amount of nucleic acid delivered to the cell is greater for the second of the two transfections than for the first of the two transfections. In another embodiment, the cell is transfected more than twice, and the amount of nucleic acid delivered to the cell is greater for the second of three transfections than for the first of the same three transfections, and the amount of nucleic acid delivered to the cells is greater for the third of the same three transfections than for the second of the same three transfections. In yet another embodiment, the cell is transfected more than once, and the maximum amount of nucleic acid delivered to the cell during each transfection is sufficiently low to yield at least about 80% viability for at least two consecutive transfections.

It has now been discovered that modulating the amount of nucleic acid delivered to a population of proliferating cells *in vivo* in a series of transfections can result in both an increased effect of the nucleic acid and increased

viability of the cells. It has been further discovered that, in certain situations, when cells *in vivo* are contacted with one or more nucleic acids encoding one or more reprogramming factors in a series of transfections, the efficiency of reprogramming can be increased when the amount of nucleic acid delivered in later transfections is greater than the amount of nucleic acid delivered in earlier transfections, for at least part of the series of transfections.

- 5 Certain embodiments are therefore directed to a method for reprogramming a cell *in vivo*, wherein one or more nucleic acids is repeatedly delivered to the cell in a series of transfections, and the amount of the nucleic acid delivered to the cell is greater for at least one later transfection than for at least one earlier transfection. In one embodiment, the cell is transfected between about 2 and about 10 times, or between about 3 and about 8 times, or between about 4 and about 6 times. In another embodiment, the one or more nucleic acids includes at least
- 10 one RNA molecule, the cell is transfected between about 2 and about 10 times, and the amount of nucleic acid delivered to the cell in each transfection is the same as or greater than the amount of nucleic acid delivered to the cell in the most recent previous transfection. In yet another embodiment, the amount of nucleic acid delivered to the cell in the first transfection is between about 20ng/cm² and about 250ng/cm², or between 100ng/cm² and 600ng/cm². In yet another embodiment, the cell is transfected about 5 times at intervals of between about 12 and
- 15 about 48 hours, and the amount of nucleic acid delivered to the cell is about 25ng/cm² for the first transfection, about 50ng/cm² for the second transfection, about 100ng/cm² for the third transfection, about 200ng/cm² for the fourth transfection, and about 400ng/cm² for the fifth transfection. In yet another embodiment, the cell is further transfected at least once after the fifth transfection, and the amount of nucleic acid delivered to the cell is about 400ng/cm².
- 20 Certain embodiments are directed to a method for transfecting a cell *in vivo* with a nucleic acid, wherein the amount of nucleic acid is determined by measuring the cell density, and choosing the amount of nucleic acid to transfect based on the measurement of cell density. In one embodiment, the cell density is measured by optical means. In another embodiment, the cell is transfected repeatedly, the cell density increases between two transfections, and the amount of nucleic acid transfected is greater for the second of the two transfections than
- 25 for the first of the two transfections.

- It has now been discovered that, in certain situations, the amount of a circulating protein that is produced in a patient can be increased by administering to a patient a nucleic acid at a plurality of administration sites. In certain embodiments, the amount of a circulating protein is increased relative to the amount of the circulating protein that is produced in a patient by administering to the patient the nucleic acid at a single injection site. In
- 30 one embodiment, the administering is by injection. In another embodiment, the injection is intradermal injection. In still another embodiment, the injection is subcutaneous or intramuscular injection. In some embodiments, the plurality of administration sites comprise administration sites in the skin. In other embodiments, the plurality of administration sites are at least about 1 or at least about 2 or at least about 5 or at least about 10 or at least about 20 or at least about 50 or at least about 100 administration sites. In one embodiment, the administering is
- 35 performed within at least about 5 minutes or at least about 10 minutes or at least about 30 minutes or at least

about 1 hour or at least about 2 hours or at least about 5 hours or at least about 12 hours or at least about 1 day. In certain embodiments, the amount of a circulating protein is increased by at least about 10 percent or at least about 20 percent or at least about 50 percent or at least about 100 percent or at least about 3-fold or at least about 5-fold or at least about 10-fold or at least about 20-fold or at least about 50-fold or at least about 100-fold
5 or at least about 500-fold or at least about 1000-fold or greater than 1000-fold.

It has now been discovered that, in certain situations, the *in vivo* transfection efficiency and viability of cells contacted with the medium of the present invention can be improved by conditioning the medium. Certain embodiments are therefore directed to a method for conditioning a medium. Other embodiments are directed to a medium that is conditioned. In one embodiment, the feeders are fibroblasts, and the medium is conditioned for
10 approximately 24 hours. Other embodiments are directed to a method for transfecting a cell *in vivo*, wherein the transfection medium is conditioned. Other embodiments are directed to a method for reprogramming and/or gene-editing a cell *in vivo*, wherein the medium is conditioned. In one embodiment, the feeders are mitotically inactivated, for example, by exposure to a chemical such as mitomycin-C or by exposure to gamma radiation. In certain embodiments, it may be beneficial to use only autologous materials, in part to, for example and not
15 wishing to be bound by theory, avoid the risk of disease transmission from the feeders to the cell or the patient. Certain embodiments are therefore directed to a method for transfecting a cell *in vivo*, wherein the transfection medium is conditioned, and wherein the feeders are derived from the same individual as the cell being transfected. Other embodiments are directed to a method for reprogramming and/or gene-editing a cell *in vivo*, wherein the medium is conditioned, and wherein the feeders are derived from the same individual as the cell
20 being reprogrammed and/or gene-edited.

Several molecules can be added to media by conditioning. Certain embodiments are therefore directed to a medium that is supplemented with one or more molecules that are present in a conditioned medium. In one embodiment, the medium is supplemented with Wnt1, Wnt2, Wnt3, Wnt3a or a biologically active fragment, analogue, variant, agonist, or family-member thereof. In another embodiment, the medium is supplemented with
25 TGF- β or a biologically active fragment, analogue, variant, agonist, or family-member thereof. In yet another embodiment, a cell *in vivo* is reprogrammed according to the method of the present invention, wherein the medium is not supplemented with TGF- β for between about 1 and about 5 days, and is then supplemented with TGF- β for at least about 2 days. In yet another embodiment, the medium is supplemented with IL-6, IL-6R or a biologically active fragment, analogue, variant, agonist, or family-member thereof. In yet another embodiment,
30 the medium is supplemented with a sphingolipid or a fatty acid. In still another embodiment, the sphingolipid is lysophosphatidic acid, lysosphingomyelin, sphingosine-1-phosphate or a biologically active analogue, variant or derivative thereof.

In addition to mitotically inactivating cells, under certain conditions, irradiation can change the gene expression of cells, causing cells to produce less of certain proteins and more of certain other proteins than non-irradiated cells,
35 for example, members of the Wnt family of proteins. In addition, certain members of the Wnt family of proteins

can promote the growth and transformation of cells. It has now been discovered that, in certain situations, the efficiency of reprogramming can be greatly increased by contacting a cell *in vivo* with a medium that is conditioned using irradiated feeders instead of mitomycin-c-treated feeders. It has been further discovered that the increase in reprogramming efficiency observed when using irradiated feeders is caused in part by Wnt proteins that are secreted by the feeders. Certain embodiments are therefore directed to a method for reprogramming a cell *in vivo*, wherein the cell is contacted with Wnt1, Wnt2, Wnt3, Wnt3a or a biologically active fragment, analogue, variant, family-member or agonist thereof, including agonists of downstream targets of Wnt proteins, and/or agents that mimic one or more of the biological effects of Wnt proteins, for example, 2-amino-4-[3,4-(methylenedioxy)benzylamino]-6-(3-methoxyphenyl)pyrimidine.

Because of the low efficiency of many DNA-based reprogramming methods, these methods may be difficult or impossible to use with cells derived from patient samples, which may contain only a small number of cells. In contrast, the high efficiency of certain embodiments of the present invention can allow reliable reprogramming of a small number of cells, including single cells. Certain embodiments are directed to a method for reprogramming a small number of cells. Other embodiments are directed to a method for reprogramming a single cell. In one embodiment, the cell is contacted with one or more enzymes. In another embodiment, the enzyme is collagenase. In yet another embodiment, the collagenase is animal-component free. In one embodiment, the collagenase is present at a concentration of between about 0.1mg/mL and about 10mg/mL, or between about 0.5mg/mL and about 5mg/mL. In another embodiment, the cell is a blood cell. In yet another embodiment, the cell is contacted with a medium containing one or more proteins that is derived from the patient's blood. In still another embodiment, the cell is contacted with a medium comprising: DMEM/F12 + 2mM L-alanyl-L-glutamine + between about 5% and about 25% patient-derived serum, or between about 10% and about 20% patient-derived serum, or about 20% patient-derived serum.

It has now been discovered that, in certain situations, transfecting cells *in vivo* with a mixture of RNA encoding Oct4, Sox2, Klf4, and c-Myc using the medium of the present invention can cause the rate of proliferation of the cells to increase. When the amount of RNA delivered to the cells is too low to ensure that all of the cells are transfected, only a fraction of the cells may show an increased proliferation rate. In certain situations, such as when generating a personalized therapeutic, increasing the proliferation rate of cells may be desirable, in part because doing so can reduce the time necessary to generate the therapeutic, and therefore can reduce the cost of the therapeutic. Certain embodiments are therefore directed to a method for transfecting a cell *in vivo* with a mixture of RNA encoding Oct4, Sox2, Klf4, and c-Myc. In one embodiment, the cell exhibits an increased proliferation rate. In another embodiment, the cell is reprogrammed.

Many diseases are associated with one or more mutations. Mutations can be corrected by contacting a cell with a nucleic acid that encodes a protein that, either alone or in combination with other molecules, corrects the mutation (an example of gene-editing). Examples of such proteins include: a nuclease, a transcription activator-like effector nuclease (TALEN), a zinc-finger nuclease, a meganuclease, a nickase, a clustered regularly

interspaced short palindromic repeat (CRISPR)-associated protein or a natural or engineered variant, family-member, orthologue, fragment or fusion construct thereof. Certain embodiments are therefore directed to a method for transfecting a cell *in vivo* with a nucleic acid, wherein the nucleic acid encodes a protein that, either alone or in combination with other molecules, creates a single-strand or double-strand break in a DNA molecule.

5 In a one embodiment, the protein is a zinc finger nuclease or a TALEN. In another embodiment, the nucleic acid is an RNA molecule. In yet another embodiment, the single-strand or double-strand break is within about 5,000,000 bases of the transcription start site of a gene selected from the group: CCR5, CXCR4, GAD1, GAD2, CFTR, HBA1, HBA2, HBB, HBD, FANCA, XPA, XPB, XPC, ERCC2, POLH, HTT, DMD, SOD1, APOE, PRNP, BRCA1, and BRCA2 or an analogue, variant or family-member thereof. In one embodiment, the present
10 invention relates to gene-editing of the MYC protein (e.g. correcting one or more mutations that may be linked to cancer), optionally with a TALEN. In yet another embodiment, the cell is transfected with a nucleic acid that acts as a repair template by either causing the insertion of a DNA sequence in the region of the single-strand or double-strand break or by causing the DNA sequence in the region of the single-strand or double-strand break to otherwise change. In yet another embodiment, the cell is reprogrammed, and subsequently, the cell is gene-
15 edited. In yet another embodiment, the cell is gene-edited, and subsequently, the cell is reprogrammed. In yet another embodiment, the gene-editing and reprogramming are performed within about 7 days of each other. In yet another embodiment, the gene-editing and reprogramming occur simultaneously or on the same day. In yet another embodiment, the cell is a skin cell, the skin cell is gene-edited to disrupt the CCR5 gene, the skin cell is reprogrammed to a hematopoietic stem cell, thus producing a therapeutic for HIV/AIDS, and the therapeutic is
20 used to treat a patient with HIV/AIDS. In yet another embodiment, the skin cell is derived from the same patient whom the therapeutic is used to treat.

Genes that can be edited according to the methods of the present invention to produce therapeutics of the present invention include genes that can be edited to restore normal function, as well as genes that can be edited to reduce or eliminate function. Such genes include, but are not limited to beta globin (HBB), mutations in
25 which can cause sickle cell disease (SCD) and β -thalassemia, breast cancer 1, early onset (BRCA1) and breast cancer 2, early onset (BRCA2), mutations in which can increase susceptibility to breast cancer, C-C chemokine receptor type 5 (CCR5) and C-X-C chemokine receptor type 4 (CXCR4), mutations in which can confer resistance to HIV infection, cystic fibrosis transmembrane conductance regulator (CFTR), mutations in which can cause cystic fibrosis, dystrophin (DMD), mutations in which can cause muscular dystrophy, including Duchenne
30 muscular dystrophy and Becker's muscular dystrophy, glutamate decarboxylase 1 and glutamate decarboxylase 2 (GAD1, GAD2), mutations in which can prevent autoimmune destruction of β -cells, hemoglobin alpha 1, hemoglobin alpha 2, and hemoglobin delta (HBA1, HBA2, and HBD), mutations in which can cause thalassemia, desmoplakin, keratin 5, keratin 14, plectin, integrin alpha-6, integrin beta-4, laminin subunit alpha-3, laminin subunit beta-3, laminin subunit gamma-2, collagen type VII alpha 1, collagen type XVII alpha 1, and matrix
35 metalloproteinase-1 (DSP, KRT5, KRT14, PLEC1, ITGA6, ITGB4, LAMA3, LAMB3, LAMC2, COL7A1, COL17A1, and MMP1), mutations in which can cause epidermolysis bullosa, Huntington (HTT), mutations in which can

5

10

25

30

35

binding efficiency and excessive off-target activity, which can introduce undesired mutations in the DNA of the cell, severely limiting their use *in vivo*, for example in therapeutic and cosmetic applications, in which the introduction of undesired mutations in a patient's cells could lead to the development of cancer. It has now been discovered that gene-editing proteins that comprise the Stsl endonuclease cleavage domain (SEQ ID NO: 1) can exhibit substantially lower off-target activity *in vivo* than previously disclosed gene-editing proteins, while maintaining a high level of on-target activity *in vivo*. Other novel engineered proteins have also been discovered that can exhibit high on-target activity *in vivo*, low off-target activity *in vivo*, small size, solubility, and other desirable characteristics when they are used as the nuclease domain of a gene-editing protein: Stsl-HA (SEQ ID NO: 2), Stsl-HA2 (SEQ ID NO: 3), Stsl-UHA (SEQ ID NO: 4), Stsl-UHA2 (SEQ ID NO: 5), Stsl-HF (SEQ ID NO: 6), and Stsl-UHF (SEQ ID NO: 7). Stsl-HA, Stsl-HA2 (high activity), Stsl-UHA, and Stsl-UHA2 (ultra-high activity) can exhibit higher on-target activity *in vivo* than both wild-type Stsl and wild-type FokI, due in part to specific amino-acid substitutions within the N-terminal region at the 34 and 61 positions, while Stsl-HF (high fidelity) and Stsl-UHF (ultra-high fidelity) can exhibit lower off-target activity *in vivo* than both wild-type Stsl and wild-type FokI, due in part to specific amino-acid substitutions within the C-terminal region at the 141 and 152 positions.

Certain embodiments are therefore directed to a protein. In some embodiments, the protein is present *in vivo*. In other embodiments, the protein comprises a nuclease domain. In one embodiment, the nuclease domain comprises one or more of: the cleavage domain of FokI endonuclease (SEQ ID NO: 53), the cleavage domain of Stsl endonuclease (SEQ ID NO: 1), Stsl-HA (SEQ ID NO: 2), Stsl-HA2 (SEQ ID NO: 3), Stsl-UHA (SEQ ID NO: 4), Stsl-UHA2 (SEQ ID NO: 5), Stsl-HF (SEQ ID NO: 6), and Stsl-UHF (SEQ ID NO: 7) or a biologically active fragment or variant thereof.

It has also been discovered that engineered gene-editing proteins that comprise DNA-binding domains comprising certain novel repeat sequences can exhibit lower off-target activity *in vivo* than previously disclosed gene-editing proteins, while maintaining a high level of on-target activity *in vivo*. Certain of these engineered gene-editing proteins can provide several advantages over previously disclosed gene-editing proteins, including, for example, increased flexibility of the linker region connecting repeat sequences, which can result in increased binding efficiency. Certain embodiments are therefore directed to a protein comprising a plurality of repeat sequences. In one embodiment, at least one of the repeat sequences contains the amino-acid sequence: GabG, where "a" and "b" each represent any amino acid. In one embodiment, the protein is a gene-editing protein. In another embodiment, one or more of the repeat sequences are present in a DNA-binding domain. In a further embodiment, "a" and "b" are each independently selected from the group: H and G. In a still further embodiment, "a" and "b" are H and G, respectively. In one embodiment, the amino-acid sequence is present within about 5 amino acids of the C-terminus of the repeat sequence. In another embodiment, the amino-acid sequence is present at the C-terminus of the repeat sequence. In some embodiments, one or more G in the amino-acid sequence GabG is replaced with one or more amino acids other than G, for example A, H or GG. In one embodiment, the repeat sequence has a length of between about 32 and about 40 amino acids or between about

33 and about 39 amino acids or between about 34 and 38 amino acids or between about 35 and about 37 amino acids or about 36 amino acids or greater than about 32 amino acids or greater than about 33 amino acids or greater than about 34 amino acids or greater than about 35 amino acids. Other embodiments are directed to a protein comprising one or more transcription activator-like effector domains. In one embodiment, at least one of the transcription activator-like effector domains comprises a repeat sequence. Other embodiments are directed to a protein comprising a plurality of repeat sequences generated by inserting one or more amino acids between at least two of the repeat sequences of a transcription activator-like effector domain. In one embodiment, one or more amino acids is inserted about 1 or about 2 or about 3 or about 4 or about 5 amino acids from the C-terminus of at least one repeat sequence. Still other embodiments are directed to a protein comprising a plurality of repeat sequences, wherein about every other repeat sequence has a different length than the repeat sequence immediately preceding or following the repeat sequence. In one embodiment, every other repeat sequence is about 36 amino acids long. In another embodiment, every other repeat sequence is 36 amino acids long. Still other embodiments are directed to a protein comprising a plurality of repeat sequences, wherein the plurality of repeat sequences comprises at least two repeat sequences that are each at least 36 amino acids long, and wherein at least two of the repeat sequences that are at least 36 amino acids long are separated by at least one repeat sequence that is less than 36 amino acids long. Some embodiments are directed to a protein that comprises one or more sequences selected from, for example, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, and SEQ ID NO: 60.

Other embodiments are directed to a protein that comprises a DNA-binding domain. In some embodiments, the DNA-binding domain comprises a plurality of repeat sequences. In one embodiment, the plurality of repeat sequences enables high-specificity recognition of a binding site in a target DNA molecule. In another embodiment, at least two of the repeat sequences have at least about 50%, or about 60%, or about 70%, or about 80%, or about 90%, or about 95%, or about 98%, or about 99% homology to each other. In a further embodiment, at least one of the repeat sequences comprises one or more regions capable of binding to a binding site in a target DNA molecule. In a still further embodiment, the binding site comprises a defined sequence of between about 1 to about 5 bases in length. In one embodiment, the DNA-binding domain comprises a zinc finger. In another embodiment, the DNA-binding domain comprises a transcription activator-like effector (TALE). In a further embodiment, the plurality of repeat sequences includes at least one repeat sequence having at least about 50% or about 60% or about 70% or about 80% or about 90% or about 95% or about 98%, or about 99% homology to a TALE. In a still further embodiment, the gene-editing protein comprises a clustered regularly interspaced short palindromic repeat (CRISPR)-associated protein. In one embodiment, the gene-editing protein comprises a nuclear-localization sequence. In another embodiment, the nuclear-localization sequence comprises the amino-acid sequence: PKKKRKV (SEQ ID NO:471). In one embodiment, the gene-editing protein comprises a mitochondrial-localization sequence. In another embodiment, the mitochondrial-localization sequence comprises the amino-acid sequence: LGRVIPRKIASRLM (SEQ ID NO:472). In one embodiment, the gene-editing protein comprises a linker. In another embodiment, the linker connects a DNA-

binding domain to a nuclease domain. In a further embodiment, the linker is between about 1 and about 10 amino acids long. In some embodiments, the linker is about 1, about 2, or about 3, or about 4, or about 5, or about 6, or about 7, or about 8, or about 9, or about 10 amino acids long. In one embodiment, the gene-editing protein is capable of generating a nick or a double-strand break in a target DNA molecule.

- 5 Certain embodiments are directed to a method for modifying the genome of a cell *in vivo*, the method comprising introducing into a cell *in vivo* a nucleic acid molecule encoding a non-naturally occurring fusion protein comprising an artificial transcription activator-like (TAL) effector repeat domain comprising one or more repeat units 36 amino acids in length and an endonuclease domain, wherein the repeat domain is engineered for recognition of a predetermined nucleotide sequence, and wherein the fusion protein recognizes the
- 10 predetermined nucleotide sequence. In one embodiment, the cell is a eukaryotic cell. In another embodiment, the cell is an animal cell. In a further embodiment, the cell is a mammalian cell. In a still further embodiment, the cell is a human cell. In one embodiment, the cell is a plant cell. In another embodiment, the cell is a prokaryotic cell. In some embodiments, the fusion protein introduces an endonucleolytic cleavage in a nucleic acid of the cell, whereby the genome of the cell is modified.
- 15 Certain embodiments are directed to a composition for altering the DNA sequence of a cell *in vivo* comprising a nucleic acid, wherein the nucleic acid encodes a gene-editing protein. Other embodiments are directed to a composition for altering the DNA sequence of a cell *in vivo* comprising a nucleic-acid mixture, wherein the nucleic-acid mixture comprises: a first nucleic acid that encodes a first gene-editing protein, and a second nucleic acid that encodes a second gene-editing protein. In one embodiment, the binding site of the first gene-editing
- 20 protein and the binding site of the second gene-editing protein are present in the same target DNA molecule. In another embodiment, the binding site of the first gene-editing protein and the binding site of the second gene-editing protein are separated by less than about 50 bases, or less than about 40 bases, or less than about 30 bases or less than about 20 bases, or less than about 10 bases, or between about 10 bases and about 25 bases or about 15 bases. In one embodiment, the nuclease domain of the first gene-editing protein and the nuclease
- 25 domain of the second gene-editing protein are capable of forming a dimer. In another embodiment, the dimer is capable of generating a nick or double-strand break in a target DNA molecule.

Certain embodiments are directed to a therapeutic composition. Other embodiments are directed to a cosmetic composition. In some embodiments, the composition comprises a repair template. In a further embodiment, the repair template is a single-stranded DNA molecule or a double-stranded DNA molecule.

- 30 Other embodiments are directed to an article of manufacture for synthesizing a protein or a nucleic acid encoding a protein. In one embodiment, the article is a nucleic acid. In another embodiment, the protein comprises a DNA-binding domain. In a further embodiment, the nucleic acid comprises a nucleotide sequence encoding a DNA-binding domain. In one embodiment, the protein comprises a nuclease domain. In another embodiment, the nucleic acid comprises a nucleotide sequence encoding a nuclease domain. In one embodiment, the protein
- 35 comprises a plurality of repeat sequences. In another embodiment, the nucleic acid encodes a plurality of repeat

sequences. In a further embodiment, the nuclease domain is selected from: FokI, Stsl, Stsl-HA, Stsl-HA2, Stsl-UHA, Stsl-UHA2, Stsl-HF, and Stsl-UHF or a natural or engineered variant or biologically active fragment thereof. In one embodiment, the nucleic acid comprises an RNA-polymerase promoter. In another embodiment, the RNA-polymerase promoter is a T7 promoter or a SP6 promoter. In a further embodiment, the nucleic acid
 5 comprises a viral promoter. In one embodiment, the nucleic acid comprises an untranslated region. In another embodiment, the nucleic acid is an *in vitro*-transcription template.

Certain embodiments are directed to a method for inducing a cell to express a protein *in vivo*. Other embodiments are directed to a method for altering the DNA sequence of a cell *in vivo* comprising transfecting the cell *in vivo* with a gene-editing protein or inducing the cell to express a gene-editing protein *in vivo*. Still other
 10 embodiments are directed to a method for reducing the expression of a protein of interest in a cell *in vivo*. In one embodiment, the cell is induced to express a gene-editing protein, wherein the gene-editing protein is capable of creating a nick or a double-strand break in a target DNA molecule. In another embodiment, the nick or double-strand break results in inactivation of a gene. Still other embodiments are directed to a method for generating an inactive, reduced-activity or dominant-negative form of a protein *in vivo*. In one embodiment, the protein is
 15 survivin. Still other embodiments are directed to a method for repairing one or more mutations in a cell *in vivo*. In one embodiment, the cell is contacted with a repair template. In another embodiment, the repair template is a DNA molecule. In a further embodiment, the repair template does not contain a binding site of the gene-editing protein. In a still further embodiment, the repair template encodes an amino-acid sequence that is encoded by a DNA sequence that comprises a binding site of the gene-editing protein.

In various embodiments, the repair template is about 20 nucleotides, or about 30 nucleotides, or about 40 nucleotides, or about 50 nucleotides, or about 60 nucleotides, or about 70 nucleotides, or about 80 nucleotides, or about 90 nucleotides, or about 100 nucleotides, or about 150 nucleotides, or about 200 nucleotides, or about 300 nucleotides, or about 400 nucleotides, or about 500 nucleotides, or about 750 nucleotides, or about 1000 nucleotides. In various embodiments, the repair template is about 20-1000 nucleotides, or about 20-500
 25 nucleotides, or about 20-400 nucleotides, or about 20-200 nucleotides, or about 20-100 nucleotides, or about 80-100 nucleotides, or about 50-100 nucleotides.

In various embodiments, the mass ratio of RNA (*e.g.* synthetic RNA encoding gene-editing protein) to repair template is about 1:10, or about 1:9, or about 1:8, or about 1:7, or about 1:6, or about 1:5, or about 1:4, or about 1:3, or about 1:2, or about 1:1, or about 2:1, or about 3:1, or about 4:1, or about 5:1, or about 6:1, or about 7:1, or
 30 about 8:1, or about 9:1, or about 10:1.

In various embodiments, the molar ratio of RNA (*e.g.* synthetic RNA encoding gene-editing protein) to repair template is about 1:10, or about 1:9, or about 1:8, or about 1:7, or about 1:6, or about 1:5, or about 1:4, or about 1:3, or about 1:2, or about 1:1, or about 2:1, or about 3:1, or about 4:1, or about 5:1, or about 6:1, or about 7:1, or about 8:1, or about 9:1, or about 10:1.

In various embodiments, the repair template has a dual function, causing a repair to a gene-edited target sequence and preventing further binding of a gene-editing protein, thereby reducing or eliminating further gene-editing (e.g. via the repair template causing a repair that renders what was the gene-editing protein binding site no longer suitable for gene-editing protein binding). Accordingly, in some embodiments, the present gene-editing methods are tunable to ensure a single gene-edit per target site.

Other embodiments are directed to a method for treating a patient comprising administering to the patient a therapeutically or cosmetically effective amount of a protein or a nucleic acid encoding a protein. In one embodiment, the treatment results in one or more of the patient's symptoms being ameliorated. Certain embodiments are directed to a method for treating a patient comprising: a. inducing a cell to express a protein of interest by transfecting the cell *in vivo* with a nucleic acid encoding the protein of interest and/or b. reprogramming the cell *in vivo*. In one embodiment, the cell is reprogrammed to a less differentiated state. In another embodiment, the cell is reprogrammed by transfecting the cell with one or more synthetic RNA molecules encoding one or more reprogramming proteins. In a further embodiment, the cell is differentiated. In a still further embodiment, the cell is differentiated into one of: a skin cell, a glucose-responsive insulin-producing cell, a hematopoietic cell, a cardiac cell, a retinal cell, a renal cell, a neural cell, a stromal cell, a fat cell, a bone cell, a muscle cell, an oocyte, and a sperm cell. Other embodiments are directed to a method for treating a patient comprising: a. inducing a cell to express a gene-editing protein by transfecting the cell *in vivo* with a nucleic acid encoding a gene-editing protein and/or b. reprogramming the cell *in vivo*.

Other embodiments are directed to a complexation medium. In one embodiment, the complexation medium has a pH greater than about 7, or greater than about 7.2, or greater than about 7.4, or greater than about 7.6, or greater than about 7.8, or greater than about 8.0, or greater than about 8.2, or greater than about 8.4, or greater than about 8.6, or greater than about 8.8, or greater than about 9.0. In another embodiment, the complexation medium comprises transferrin. In a further embodiment, the complexation medium comprises DMEM. In a still further embodiment, the complexation medium comprises DMEM/F12. Still other embodiments are directed to a method for forming nucleic-acid-transfection-reagent complexes. In one embodiment, the transfection reagent is incubated with a complexation medium. In another embodiment, the incubation occurs before a mixing step. In a further embodiment, the incubation step is between about 5 seconds and about 5 minutes or between about 10 seconds and about 2 minutes or between about 15 seconds and about 1 minute or between about 30 seconds and about 45 seconds. In one embodiment, the transfection reagent is selected from **Table 1**. In another embodiment, the transfection reagent is a lipid or lipidoid. In a further embodiment, the transfection reagent comprises a cation. In a still further embodiment, the cation is a multivalent cation. In a still further embodiment, the transfection reagent is N1-[2-((1S)-1-[(3-aminopropyl)amino]-4-[di(3-aminopropyl)amino]butylcarboxamido)ethyl]-3,4-di[oleyloxy]-benzamide (a.k.a. MVL5) or a derivative thereof.

Certain embodiments are directed to a method for inducing a cell to express a protein by contacting the cell with a nucleic acid *in vivo*. In one embodiment, the cell is a mammalian cell. In another embodiment, the cell is a

human cell or a rodent cell. Other embodiments are directed to a cell produced using one or more of the methods of the present invention. In one embodiment, the cell is present in a patient. In another embodiment, the cell is isolated from a patient. Other embodiments are directed to a screening library comprising a cell produced using one or more of the methods of the present invention. In one embodiment, the screening library is used for at least one of: toxicity screening, including: cardiotoxicity screening, neurotoxicity screening, and hepatotoxicity screening, efficacy screening, high-throughput screening, high-content screening, and other screening.

Other embodiments are directed to a kit containing a nucleic acid. In one embodiment, the kit contains a delivery reagent (a.k.a. "transfection reagent"). In another embodiment, the kit is a reprogramming kit. In a further embodiment, the kit is a gene-editing kit. Other embodiments are directed to a kit for producing nucleic acids. In one embodiment, the kit contains at least two of: pseudouridine-triphosphate, 5-methyluridine triphosphate, 5-methylcytidine triphosphate, 5-hydroxymethylcytidine triphosphate, N4-methylcytidine triphosphate, N4-acetylcytidine triphosphate, and 7-deazaguanosine triphosphate or one or more derivatives thereof. Other embodiments are directed to a therapeutic or cosmetic comprising a nucleic acid. In one embodiment, the therapeutic or cosmetic is a pharmaceutical composition. In another embodiment, the pharmaceutical composition is formulated. In a further embodiment, the formulation comprises an aqueous suspension of liposomes. Example liposome components are set forth in **Table 1**, and are given by way of example, and not by way of limitation. In one embodiment, the liposomes include one or more polyethylene glycol (PEG) chains. In another embodiment, the PEG is PEG2000. In a further embodiment, the liposomes include 1,2-distearoyl-sn-glycero-3-phosphoethanolamine (DSPE) or a derivative thereof. In one embodiment, the therapeutic comprises one or more ligands. In another embodiment, the therapeutic comprises at least one of: androgen, CD30 (TNFRSF8), a cell-penetrating peptide, CXCR, estrogen, epidermal growth factor, EGFR, HER2, folate, insulin, insulin-like growth factor-I, interleukin-13, integrin, progesterone, stromal-derived-factor-1, thrombin, vitamin D, and transferrin or a biologically active fragment or variant thereof. Still other embodiments are directed to a therapeutic or cosmetic comprising a cell generated using one or more of the methods of the present invention.

In one embodiment, the therapeutic is administered to a patient for the treatment of any of the diseases or disorders described herein, including by way of non-limitation, type 1 diabetes, heart disease, including ischemic and dilated cardiomyopathy, macular degeneration, Parkinson's disease, cystic fibrosis, sickle-cell anemia, thalassemia, Fanconi anemia, severe combined immunodeficiency, hereditary sensory neuropathy, xeroderma pigmentosum, Huntington's disease, muscular dystrophy, amyotrophic lateral sclerosis, Alzheimer's disease, cancer, and infectious diseases including: hepatitis and HIV/AIDS.

Other embodiments are directed to a method for reprogramming a cell *in vivo*. In one embodiment, the cell is reprogrammed by contacting the cell with one or more nucleic acids. In one embodiment, the cell is contacted with a plurality of nucleic acids encoding at least one of: Oct4 protein, Sox2 protein, Klf4 protein, c-Myc protein, Lin28 protein or a biologically active fragment, variant or derivative thereof. In another embodiment, the cell is contacted with a plurality of nucleic acids encoding a plurality of proteins including: Oct4 protein, Sox2 protein,

Klf4 protein, and c-Myc protein or one or more biologically active fragments, variants or derivatives thereof. Still other embodiments are directed to a method for gene editing a cell *in vivo*. In one embodiment, the cell is gene-edited by contacting the cell with one or more nucleic acids.

Certain embodiments are directed to a method for inducing a cell *in vivo* to express a protein of interest comprising contacting a cell *in vivo* with a solution comprising albumin that is treated with an ion-exchange resin or charcoal and one or more nucleic acid molecules, wherein at least one of the one or more nucleic acid molecules encodes a protein of interest. In one embodiment, the method results in the cell expressing the protein of interest. In another embodiment, the one or more nucleic acid molecules comprise a synthetic RNA molecule. In one embodiment, the cell is a skin cell. In another embodiment, the cell is a muscle cell. In yet another embodiment, the cell is a dermal fibroblast. In yet another embodiment, the cell is a myoblast. In one embodiment, the protein of interest is an extracellular matrix protein. In another embodiment, the protein of interest is selected from: elastin, collagen, laminin, fibronectin, vitronectin, lysyl oxidase, elastin binding protein, a growth factor, fibroblast growth factor, transforming growth factor beta, granulocyte colony-stimulating factor, a matrix metalloproteinase, an actin, fibrillin, microfibril-associated glycoprotein, a lysyl-oxidase-like protein, and platelet-derived growth factor. In one embodiment, the solution is delivered to the dermis. In another embodiment, the delivering is by injection. In yet another embodiment, the delivering is by injection using a microneedle array. In one embodiment, the solution further comprises a growth factor. In another embodiment, the growth factor is selected from: fibroblast growth factor and transforming growth factor beta. In yet another embodiment, the solution further comprises cholesterol. Other embodiments are directed a method for inducing a cell *in vivo* to express a protein of interest comprising contacting a cell *in vivo* with a solution comprising cholesterol and one or more nucleic acid molecules, wherein at least one of the one or more nucleic acid molecules encodes a protein of interest. In one embodiment, the method results in the cell expressing the protein of interest. Still other embodiments are directed to a method for transfecting a cell *in vivo* with a nucleic acid molecule comprising contacting a cell *in vivo* with a solution comprising albumin that is treated with an ion-exchange resin or charcoal and a nucleic acid molecule. In one embodiment, the method results in the cell being transfected with the nucleic acid molecule. In another embodiment, the nucleic acid molecule is one of: a dsDNA molecule, a ssDNA molecule, a dsRNA molecule, a ssRNA molecule, a plasmid, an oligonucleotide, a synthetic RNA molecule, a miRNA molecule, an mRNA molecule, an siRNA molecule. Still other embodiments are directed to a method for treating a patient comprising delivering to a patient a composition comprising albumin that is treated with an ion-exchange resin or charcoal and one or more nucleic acid molecules, wherein at least one of the one or more nucleic acid molecules encodes a protein of interest. In one embodiment, the method results in the expression of the protein of interest in the patient. In another embodiment, the method results in the expression of the protein of interest in the dermis of the patient.

Certain embodiments are directed to a cosmetic composition comprising albumin that is treated with an ion-exchange resin or charcoal and a nucleic acid molecule. Other embodiments are directed to a cosmetic

treatment article. In one embodiment, the cosmetic treatment article comprises a device configured to deliver a composition to a patient. In another embodiment, the nucleic acid molecule encodes elastin protein or collagen protein. Still other embodiments are directed to a solution for transfecting a cell *in vivo* comprising cholesterol or a cholesterol analog and one or more nucleic acid molecules. In one embodiment, the cholesterol or cholesterol analog is covalently bound to at least one of the one or more nucleic acid molecules. In another embodiment, the cholesterol analog is an oxysterol. In yet another embodiment, the cholesterol analog includes one or more of: an A-ring substitution, a B-ring substitution, a D-ring substitution, a side-chain substitution, a cholestanoic acid, a cholestenoic acid, a polyunsaturated moiety, a deuterated moiety, a fluorinated moiety, a sulfonated moiety, a phosphorylated moiety, and a fluorescent moiety. In yet another embodiment, the method comprises treating the patient with one or more of: a dermal filler, a neurotoxin (by way of illustration sodium channel inhibitors (e.g., tetrodotoxin), potassium channel inhibitors (e.g., tetraethylammonium), chloride channel inhibitors (e.g., chlorotoxin and curare), calcium channel inhibitors (e.g., conotoxin), synaptic vesicle release inhibitors (e.g., botulinum toxin and tetanus toxin) and blood brain barrier inhibitor (e.g., aluminum and mercury)) and a repair-inducing treatment.

Despite the tendency of transfection reagent nucleic acid complexes to precipitate, form clumps or otherwise degrade when stored for more than a few minutes, the present inventors have surprisingly discovered that transfection reagent nucleic acid complexes produced according to some embodiments of the present invention can be frozen and/or can be stored at various temperatures, including room temperature, about 4°C, about -20°C, about -80°C, and about -196°C for an extended period of time, for example, for several hours, about 1 day, about 1 week, about 1 month, about 1 year, and longer than about 1 year. Some embodiments are therefore directed to a pharmaceutical formulation comprising synthetic RNA and a transfection reagent, wherein the pharmaceutical formulation is provided in solid form. Other embodiments are directed to a pharmaceutical formulation comprising synthetic RNA transfection reagent complexes, wherein the synthetic RNA transfection reagent complexes are provided in solid form. In various embodiments, the synthetic RNA transfection reagent complexes are provided in frozen form. Various embodiments are directed to a method for stabilizing nucleic acid transfection reagent complexes comprising forming nucleic acid transfection reagent complexes and contacting the nucleic acid transfection reagent complexes or vessel in which such are contained with a cryogenic liquid to produce stabilized nucleic acid transfection reagent complexes. In one embodiment, the nucleic acid transfection reagent complexes are stabilized for shipment or storage.

Illustrative subjects or patients refers to any vertebrate including, without limitation, humans and other primates (e.g., chimpanzees and other apes and monkey species), farm animals (e.g., cattle, sheep, pigs, goats, and horses), domestic mammals (e.g., dogs and cats), laboratory animals (e.g., rodents such as mice, rats, and guinea pigs), and birds (e.g., domestic, wild and game birds such as chickens, turkeys and other gallinaceous birds, ducks, geese, and the like). In some embodiments, the subject is a mammal. In some embodiments, the subject is a human.

Definitions

By “molecule” is meant a molecular entity (molecule, ion, complex, *etc.*).

By “RNA molecule” is meant a molecule that comprises RNA.

By “synthetic RNA molecule” is meant an RNA molecule that is produced outside of a cell or that is produced inside of a cell using bioengineering, by way of non-limiting example, an RNA molecule that is produced in an *in vitro*-transcription reaction, an RNA molecule that is produced by direct chemical synthesis or an RNA molecule that is produced in a genetically-engineered *E.coli* cell.

By “transfection” is meant contacting a cell with a molecule, wherein the molecule is internalized by the cell.

By “upon transfection” is meant during or after transfection.

By “transfection reagent” is meant a substance or mixture of substances that associates with a molecule and facilitates the delivery of the molecule to and/or internalization of the molecule by a cell, by way of non-limiting example, a cationic lipid, a charged polymer or a cell-penetrating peptide.

By “reagent-based transfection” is meant transfection using a transfection reagent.

By “medium” is meant a solvent or a solution comprising a solvent and a solute, by way of non-limiting example, Dulbecco’s Modified Eagle’s Medium (DMEM), DMEM + 10% fetal bovine serum (FBS), saline or water.

By “complexation medium” is meant a medium to which a transfection reagent and a molecule to be transfected are added and in which the transfection reagent associates with the molecule to be transfected.

By “transfection medium” is meant a medium that can be used for transfection, by way of non-limiting example, Dulbecco’s Modified Eagle’s Medium (DMEM), DMEM/F12, saline or water.

By “recombinant protein” is meant a protein or peptide that is not produced in animals or humans. Non-limiting examples include human transferrin that is produced in bacteria, human fibronectin that is produced in an *in vitro* culture of mouse cells, and human serum albumin that is produced in a rice plant.

By “Oct4 protein” is meant a protein that is encoded by the POU5F1 gene, or a natural or engineered variant, family-member, orthologue, fragment or fusion construct thereof, by way of non-limiting example, human Oct4 protein (SEQ ID NO: 8), mouse Oct4 protein, Oct1 protein, a protein encoded by POU5F1 pseudogene 2, a DNA-binding domain of Oct4 protein or an Oct4-GFP fusion protein. In some embodiments the Oct4 protein comprises an amino acid sequence that has at least 70% identity with SEQ ID NO: 8, or in other embodiments, at least 75%, 80%, 85%, 90%, or 95% identity with SEQ ID NO: 8. In some embodiments, the Oct4 protein comprises an amino acid sequence having from 1 to 20 amino acid insertions, deletions, or substitutions (collectively) with respect to SEQ ID NO: 8. Or in other embodiments, the Oct4 protein comprises an amino acid sequence having from 1 to 15 or from 1 to 10 amino acid insertions, deletions, or substitutions (collectively) with respect to SEQ ID NO: 8.

By "Sox2 protein" is meant a protein that is encoded by the SOX2 gene, or a natural or engineered variant, family-member, orthologue, fragment or fusion construct thereof, by way of non-limiting example, human Sox2 protein (SEQ ID NO: 9), mouse Sox2 protein, a DNA-binding domain of Sox2 protein or a Sox2-GFP fusion protein. In some embodiments the Sox2 protein comprises an amino acid sequence that has at least 70% identity with SEQ ID NO: 9, or in other embodiments, at least 75%, 80%, 85%, 90%, or 95% identity with SEQ ID NO: 9. In some embodiments, the Sox2 protein comprises an amino acid sequence having from 1 to 20 amino acid insertions, deletions, or substitutions (collectively) with respect to SEQ ID NO: 9. Or in other embodiments, the Sox2 protein comprises an amino acid sequence having from 1 to 15 or from 1 to 10 amino acid insertions, deletions, or substitutions (collectively) with respect to SEQ ID NO: 9.

By "Klf4 protein" is meant a protein that is encoded by the KLF4 gene, or a natural or engineered variant, family-member, orthologue, fragment or fusion construct thereof, by way of non-limiting example, human Klf4 protein (SEQ ID NO: 10), mouse Klf4 protein, a DNA-binding domain of Klf4 protein or a Klf4-GFP fusion protein. In some embodiments the Klf4 protein comprises an amino acid sequence that has at least 70% identity with SEQ ID NO: 10, or in other embodiments, at least 75%, 80%, 85%, 90%, or 95% identity with SEQ ID NO: 10. In some embodiments, the Klf4 protein comprises an amino acid sequence having from 1 to 20 amino acid insertions, deletions, or substitutions (collectively) with respect to SEQ ID NO: 10. Or in other embodiments, the Klf4 protein comprises an amino acid sequence having from 1 to 15 or from 1 to 10 amino acid insertions, deletions, or substitutions (collectively) with respect to SEQ ID NO: 10.

By "c-Myc protein" is meant a protein that is encoded by the MYC gene, or a natural or engineered variant, family-member, orthologue, fragment or fusion construct thereof, by way of non-limiting example, human c-Myc protein (SEQ ID NO: 11), mouse c-Myc protein, l-Myc protein, c-Myc (T58A) protein, a DNA-binding domain of c-Myc protein or a c-Myc-GFP fusion protein. In some embodiments the c-Myc protein comprises an amino acid sequence that has at least 70% identity with SEQ ID NO: 11, or in other embodiments, at least 75%, 80%, 85%, 90%, or 95% identity with SEQ ID NO: 11. In some embodiments, the c-Myc protein comprises an amino acid having from 1 to 20 amino acid insertions, deletions, or substitutions (collectively) with respect to SEQ ID NO: 11. Or in other embodiments, the c-Myc protein comprises an amino acid sequence having from 1 to 15 or from 1 to 10 amino acid insertions, deletions, or substitutions (collectively) with respect to SEQ ID NO: 11.

By "erythropoietin" or "erythropoietin protein" is meant a protein that is encoded by the EPO gene, or a natural or engineered variant, family-member, orthologue, fragment or fusion construct thereof, by way of non-limiting example, human erythropoietin (SEQ ID NO: 164), mouse erythropoietin, darbepoetin, darbepoetin alfa, NOVEPOETIN, a binding domain of erythropoietin or an erythropoietin-GFP fusion protein. In some embodiments the erythropoietin comprises an amino acid sequence that has at least 70% identity with SEQ ID NO: 164, or in other embodiments, at least 75%, 80%, 85%, 90%, or 95% identity with SEQ ID NO: 164. In some embodiments, the erythropoietin comprises an amino acid sequence having from 1 to 20 amino acid insertions, deletions, or substitutions (collectively) with respect to SEQ ID NO: 164. Or in other embodiments, the

erythropoietin comprises an amino acid sequence having from 1 to 15 or from 1 to 10 amino acid insertions, deletions, or substitutions (collectively) with respect to SEQ ID NO: 164.

By “reprogramming” is meant causing a change in the phenotype of a cell, by way of non-limiting example, causing a β -cell progenitor to differentiate into a mature β -cell, causing a fibroblast to dedifferentiate into a pluripotent stem cell, causing a keratinocyte to transdifferentiate into a cardiac stem cell, causing the telomeres of a cell to lengthen or causing the axon of a neuron to grow.

By “reprogramming factor” is meant a molecule that, when a cell is contacted with the molecule and/or the cell expresses the molecule, can, either alone or in combination with other molecules, cause reprogramming, by way of non-limiting example, Oct4 protein, Tert protein or erythropoietin.

10 By “germ cell” is meant a sperm cell or an egg cell.

By “pluripotent stem cell” is meant a cell that can differentiate into cells of all three germ layers (endoderm, mesoderm, and ectoderm) *in vivo*.

By “somatic cell” is meant a cell that is not a pluripotent stem cell or a germ cell, by way of non-limiting example, a skin cell.

15 By “hematopoietic cell” is meant a blood cell or a cell that can differentiate into a blood cell, by way of non-limiting example, a hematopoietic stem cell or a white blood cell.

By “cardiac cell” is meant a heart cell or a cell that can differentiate into a heart cell, by way of non-limiting example, a cardiac stem cell or a cardiomyocyte.

20 By “retinal cell” is meant a cell of the retina or a cell that can differentiate into a cell of the retina, by way of non-limiting example, a retinal pigmented epithelial cell.

By “skin cell” is meant a cell that is normally found in the skin, by way of non-limiting example, a fibroblast, a keratinocyte, a melanocyte, an adipocyte, a mesenchymal stem cell, an adipose stem cell or a blood cell.

By “immunosuppressant” is meant a substance that can suppress one or more aspects of an immune system, and that is not normally present in a mammal, by way of non-limiting example, B18R or dexamethasone.

25 By “single-strand break” is meant a region of single-stranded or double-stranded DNA in which one or more of the covalent bonds linking the nucleotides has been broken in one of the one or two strands.

By “double-strand break” is meant a region of double-stranded DNA in which one or more of the covalent bonds linking the nucleotides has been broken in each of the two strands.

30 By “nucleotide” is meant a nucleotide or a fragment or derivative thereof, by way of non-limiting example, a nucleobase, a nucleoside, a nucleotide-triphosphate, *etc.*

By “nucleoside” is meant a nucleotide or a fragment or derivative thereof, by way of non-limiting example, a nucleobase, a nucleoside, a nucleotide-triphosphate, *etc.*

By “gene editing” is meant altering the DNA sequence of a cell, by way of non-limiting example, by transfecting the cell with a protein that causes a mutation in the DNA of the cell.

By “gene-editing protein” is meant a protein that can, either alone or in combination with one or more other molecules, alter the DNA sequence of a cell, by way of non-limiting example, a nuclease, a transcription
 5 activator-like effector nuclease (TALEN), a zinc-finger nuclease, a meganuclease, a nickase, a clustered regularly interspaced short palindromic repeat (CRISPR)-associated protein or a natural or engineered variant, family-member, orthologue, fragment or fusion construct thereof.

By “repair template” is meant a nucleic acid containing a region of at least about 70% homology with a sequence that is within 10kb of a target site of a gene-editing protein.

10 By “repeat sequence” is meant an amino-acid sequence that is present in more than one copy in a protein, to within at least about 10% homology, by way of non-limiting example, a monomer repeat of a transcription activator-like effector.

By “DNA-binding domain” is meant a region of a molecule that is capable of binding to a DNA molecule, by way of non-limiting example, a protein domain comprising one or more zinc fingers, a protein domain comprising one
 15 or more transcription activator-like (TAL) effector repeat sequences or a binding pocket of a small molecule that is capable of binding to a DNA molecule.

By “binding site” is meant a nucleic-acid sequence that is capable of being recognized by a gene-editing protein, DNA-binding protein, DNA-binding domain or a biologically active fragment or variant thereof or a nucleic-acid
 20 sequence for which a gene-editing protein, DNA-binding protein, DNA-binding domain or a biologically active fragment or variant thereof has high affinity, by way of non-limiting example, an about 20-base-pair sequence of DNA in exon 1 of the human BIRC5 gene.

By “target” is meant a nucleic acid that contains a binding site.

Other definitions are set forth in US Application No. 13/465,490, US Provisional Application No. 61/664,494, US
 25 Provisional Application No. 61/721,302, International Application No. PCT/US12/67966, US Provisional Application No. 61/785,404, US Provisional Application No. 61/842,874, International Application No. PCT/US13/68118, US Provisional Application No. 61/934,397, US Application No. 14/296,220, US Provisional Application No. 62/038,608, US Provisional Application No. 62/069,667, and International Application No. PCT/US2015/013949, the contents of which are hereby incorporated by reference in their entireties.

Selected Sequences

SEQ ID NO	Description
1	Stsl
2	Stsl-HA
3	Stsl-HA2
4	Stsl-UHA

5	Stsl-UHA2
6	Stsl-HF
7	Stsl-UHF
8	Oct4
9	Sox2
10	Klf4
11	c-Myc
12	BIRC5_exon1
13	BIRC5_exon2
14	BIRC5_exon3
15	BIRC5_exon4
16	BIRC5-1.1-L
17	BIRC5-1.1-R
18	BIRC5-1.2-L
19	BIRC5-1.2-R
20	BIRC5-1.3-L
21	BIRC5-1.3-R
22	BIRC5-2.1-L
23	BIRC5-2.1-R
24	BIRC5-2.2-L
25	BIRC5-2.2-R
26	BIRC5-3.1-L
27	BIRC5-3.1-R
28	CDK1
29	CDK2
30	CDK3
31	CDK4
32	CDK5
33	CDK6
34	BIRC5
35	HIF1A
36	RRM2
37	KRAS
38	EGFR
39	MYC
40	PKN3
41	KIF11
42	APC
43	BRCA1
44	BRCA2
45	TP53
46	APP
47	HTT
48	IAPP

49	MAPT
50	PRNP
51	SNCA
52	SOD1
53	FokI
54	Repeat1
55	Repeat2
56	Repeat3
57	EO-GHGG-FokI ("GHGG" disclosed as SEQ ID NO: 547)
58	GHGG-FokI ("GHGG" disclosed as SEQ ID NO: 547)
59	EO-GHGG-StaI ("GHGG" disclosed as SEQ ID NO: 547)
60	GHGG-StaI ("GHGG" disclosed as SEQ ID NO: 547)
61	collagen alpha-1(I) chain preproprotein
62	collagen alpha-2(I) chain precursor
63	collagen alpha-1(II) chain isoform 1 precursor
64	collagen alpha-1(II) chain isoform 2 precursor
65	collagen alpha-1(III) chain preproprotein
66	collagen alpha-1(IV) chain preproprotein
67	collagen alpha-2(IV) chain preproprotein
68	collagen alpha-3(IV) chain precursor
69	collagen alpha-4(IV) chain precursor
70	collagen alpha-5(IV) chain isoform 1 precursor
71	collagen alpha-6(IV) chain isoform A precursor
72	collagen alpha-1(V) chain isoform 1 preproprotein
73	collagen alpha-2(V) chain preproprotein
74	collagen alpha-3(V) chain preproprotein
75	collagen alpha-1(VI) chain precursor
76	collagen alpha-2(VI) chain isoform 2C2 precursor
77	collagen alpha-3(VI) chain isoform 1 precursor
78	collagen alpha-1(VII) chain precursor
79	elastin isoform a precursor
80	elastin isoform b precursor
81	elastin isoform c precursor
82	elastin isoform d precursor
83	elastin isoform e precursor
84	elastin isoform f precursor
85	elastin isoform g precursor
86	elastin isoform h precursor
87	elastin isoform i precursor
88	elastin isoform j precursor
89	elastin isoform k precursor
90	elastin isoform l precursor
91	elastin isoform m precursor
92	protein-lysine 6-oxidase isoform 1 preproprotein

93	protein-lysine 6-oxidase isoform 2
94	telomerase reverse transcriptase isoform 1
95	telomerase reverse transcriptase isoform 2
96	fibronectin isoform 1 preproprotein
97	fibronectin isoform 3 preproprotein
98	fibronectin isoform 4 preproprotein
99	fibronectin isoform 5 preproprotein
100	fibronectin isoform 6 preproprotein
101	fibronectin isoform 7 preproprotein
102	vitronectin precursor
103	nidogen-1 precursor
104	laminin subunit alpha-1 precursor
105	insulin-like growth factor I isoform 1 preproprotein
106	fibroblast growth factor 1 isoform 1 precursor
107	fibroblast growth factor 2
108	transforming growth factor beta-1 precursor
109	transforming growth factor beta-2 isoform 1 precursor
110	transforming growth factor beta-2 isoform 2 precursor
111	actin, alpha skeletal muscle
112	actin, aortic smooth muscle
113	actin, cytoplasmic 1
114	actin, alpha cardiac muscle 1 proprotein
115	actin, cytoplasmic 2
116	actin, gamma-enteric smooth muscle isoform 1 precursor
117	actin, gamma-enteric smooth muscle isoform 2 precursor
118	granulocyte colony-stimulating factor isoform a precursor
119	granulocyte colony-stimulating factor isoform b precursor
120	granulocyte colony-stimulating factor isoform c precursor
121	granulocyte colony-stimulating factor isoform d precursor
122	platelet-derived growth factor subunit A isoform 1 preproprotein
123	platelet-derived growth factor subunit A isoform 2 preproprotein
124	platelet-derived growth factor subunit B isoform 1 preproprotein
125	platelet-derived growth factor subunit B isoform 2 preproprotein
126	platelet-derived growth factor C precursor
127	platelet-derived growth factor D isoform 1 precursor
128	platelet-derived growth factor D isoform 2 precursor
129	interstitial collagenase isoform 1 preproprotein
130	interstitial collagenase isoform 2
131	neutrophil collagenase preproprotein
132	stromelysin-2 preproprotein
133	macrophage metalloelastase preproprotein
134	fibrillin-1 precursor
135	fibrillin-2 precursor
136	lysyl oxidase homolog 1 preproprotein

137	lysyl oxidase homolog 2 precursor
138	lysyl oxidase homolog 3 isoform 1 precursor
139	lysyl oxidase homolog 3 isoform 2 precursor
140	lysyl oxidase homolog 3 isoform 3
141	lysyl oxidase homolog 4 precursor
142	microfibrillar-associated protein 2 isoform a precursor
143	microfibrillar-associated protein 2 isoform b precursor
144	microfibrillar-associated protein 5 precursor
145	disintegrin and metalloproteinase domain-containing protein 17 preprotein
146	desmoglein-2 preproprotein
147	DNA polymerase eta isoform 1
148	DNA polymerase eta isoform 2
149	DNA polymerase eta isoform 3
150	ferrochelatase, mitochondrial isoform a precursor
151	ferrochelatase, mitochondrial isoform b precursor
152	filaggrin
153	hyaluronan synthase 1 isoform 1
154	hyaluronan synthase 1 isoform 2
155	hyaluronan synthase 2
156	hyaluronan synthase 3 isoform a
157	hyaluronan synthase 3 isoform b
158	proopiomelanocortin
159	plakophilin-1 isoform 1a
160	plakophilin-1 isoform 1b
161	retinol dehydrogenase 10
162	mitochondrial brown fat uncoupling protein 1
163	tyrosinase precursor
164	erythropoietin
165	epoetin alfa
166	darbepoetin alfa
167	NOVEPOETIN
168	NOVECRIT

This invention is further illustrated by the following non-limiting examples.

EXAMPLES

Example 1 RNA Synthesis

- RNA encoding green fluorescent protein ("GFP"), NOVEPOETIN ("EPO"), elastin ("ELN"), tyrosinase ("TYR"), melanocortin-1-receptor ("MC1R"), HAS1, HAS2, HAS3, COL3A1, COL7A1, COL1A1, COL1A2, hTERT, Holly GFP, Fresno RFP, Blitzen Blue, RIBOSLICE gene-editing proteins, TALENs, Cas9, Oct4, Sox2, Klf4, c-Myc-2 (T58A), Lin28, IL2, IL6, IL15, IL22, BMP2, BDNF, LIF, BMP6, IL15RA, FGF21, LIF, and PTH, and comprising various combinations of canonical and non-canonical nucleotides, was synthesized from DNA templates using

the T7 High Yield RNA Synthesis Kit and the Vaccinia Capping System kit with mRNA Cap 2'-O-Methyltransferase (all from New England Biolabs, Inc.), according to the manufacturer's instructions and the present inventors' previously disclosed inventions (US Application No. 13/465,490 (now US Patent No. 8,497,124), International Application No. PCT/US12/67966, US Application No. 13/931,251, International Application No. PCT/US13/68118, and International Application No. PCT/US2015/013949, the contents of all of which are hereby incorporated by reference in their entirety) (**Table 4**). The RNA was then diluted with nuclease-free water to between 100ng/μL and 2000ng/μL. For certain experiments, an RNase inhibitor (Suprase-In, Life Technologies Corporation) was added at a concentration of 1μL/100μg of RNA. RNA solutions were stored at room temperature, 4C, -20C or -80C. For reprogramming experiments, RNA encoding Oct4, Sox2, Klf4, c-Myc-2 (T58A), and Lin28 was mixed at a molar ratio of 3:1:1:1:1.

Table 4. RNA Synthesis

Template	Nucleotides	Reaction Volume/μL	ivT Yield/μg
ELN	A, 0.5 7dG, 0.4 5mU, 5mC	20	34.1
ELN	A, 0.5 7dG, 0.4 5mU, 5mC	20	67.6
GFP	A, 0.5 7dG, 0.4 5mU, 5mC	10	60.5
GFP	A, 0.5 7dG, 0.4 5mU, 5hmC	10	25.5
GFP	A, G, U, 5hmC	10	58.3
GFP	A, 0.5 7dG, U, 5hmC	10	47.3
GFP	A, 0.5 7dG, 0.4 5mU, 5cC	10	33.8
GFP	A, G, U, 5hmC	15	30.3
GFP	A, G, U, 5hmC	15	44.6
GFP	A, G, U, 5hmC	15	24.7
TYR	A, G, U, 5hmC	15	45.4
MC1R	A, G, U, 5hmC	15	47.5
TYR	A, G, U, C	20	67.0
TYR	A, G, psU, C	20	93.7
TYR	A, G, 0.4 5mU, C	20	85.7
TYR	A, G, U, 5mC	20	73.4
TYR	A, G, U, 5hmC	20	72.7
TYR	A, 0.5 7dG, U, C	20	62.7
TYR	A, G, psU, 5mC	20	116.3
TYR	A, G, psU, 5hmC	20	102.4
TYR	A, 0.5 7dG, psU, C	20	87.3
TYR	A, G, 0.4 5mU, 5mC	20	106.5
TYR	A, G, 0.4 5mU, 5hmC	20	85.0
TYR	A, 0.5 7dG, 0.4 5mU, C	20	70.9
TYR	A, 0.5 7dG, U, 5mC	20	88.5
TYR	A, 0.5 7dG, U, 5hmC	20	59.1
TYR	A, 0.5 7dG, psU, 5mC	20	7.8
TYR	A, 0.5 7dG, psU, 5hmC	20	98.0

TYR	A, 0.5 7dG, 0.4 5mU, 5mC	20	106.5
TYR	A, 0.5 7dG, 0.4 5mU, 5hmC	20	82.3
HAS1	A, G, U, 5hmC	20	178.4
HAS2	A, G, U, 5hmC	20	59.3
HAS3	A, G, U, 5hmC	20	102.6
TYR	A, G, 0.4 5mU, 5hmC	100	377.3
COL3A1	A, G, 0.4 5mU, 5hmC	20	108.3
COL7A1	A, G, 0.4 5mU, 5hmC	20	94.6
COL1A1 (20 μ L)	A, G, 0.4 5mU, 5hmC	20	114.0
COL1A2 (10 μ L)	A, G, 0.4 5mU, 5hmC	10	31.3
TYR	A, G, 0.4 5mU, 5hmC	100	249.9
GFP	A, G, 0.4 5mU, 5hmC	100	264.0
hTERT	A, G, 0.4 5mU, 5hmC	100	349.2
GFP	A, G, U, 5hC	20	81.7
GFP	A, G, U, 0.5 5hC	20	65.4
GFP	A, sG, U, C	20	34.7
GFP	A, 0.5 sG, U, C	20	47.5
GFP	A, G, 5hmU, C	20	22.1
GFP	A, G, 0.5 5hmU, C	20	28.4
GFP	A, G, 5cU, C	20	24.4
GFP	A, G, 0.5 5cU, C	20	28.4
GFP	A, G, 5moU, C	20	39.2
GFP	A, G, 0.5 5moU, C	20	34.2
GFP	A, G, U, C	20	42.0
GFP	A, G, 5moU, C	20	53.8
GFP	A, G, 5moU, 5hmC	20	101.5
GFP	A, G, 0.4 5mU, 0.6 5moU, C	20	98.6
GFP	A, G, 0.4 5mU, C	20	99.6
GFP	A, G, U, 5mC	20	106.1
GFP	A, G, U, C	20	85.7
GFP	A, G, 5moU, C	100	398.4
hTERT	A, G, 5moU, C	20	82.6
COL7A1	A, G, 5moU, C	20	34.9
COL7A1	A, G, 5moU, C	100	342.0
Holly GFP	A, G, 5moU, C	20	36.7
Fresno RFP	A, G, 5moU, C	20	72.0
Blitzen Blue	A, G, 5moU, C	20	30.3
hTERT	A, G, 5moU, C	20	49.6
Cas9	A, G, 5moU, C	20	31.6
EPO	A, G, U, C	20	101.0
EPO	A, G, 5moU, C	20	52.9
EPO	A, G, psU, C	20	106.0
COL7A1	A, G, 5moU, C	20	80.2
Oct4 (SEQ ID NO: 8)	A, G, 5moU, C	300	1925.5

Sox2 (SEQ ID NO: 9)	A, G, 5moU, C	100	641.8
Klf4 (SEQ ID NO: 10)	A, G, 5moU, C	100	739.0
c-Myc-2 (T58A)	A, G, 5moU, C	100	574.0
Lin28	A, G, 5moU, C	100	556.0
IL2	A, G, 5moU, C	20	62.4
IL6	A, G, 5moU, C	20	22.2
IL15	A, G, 5moU, C	20	50.4
IL22	A, G, 5moU, C	20	63.6
BMP2	A, G, 5moU, C	20	83.2
BDNF	A, G, 5moU, C	20	45.0
LIF	A, G, 5moU, C	20	54.0
BMP6	A, G, 5moU, C	20	92.2
IL15RA	A, G, 5moU, C	20	91.4
FGF21	A, G, 5moU, C	20	79.2
GFP	A, G, 5moU, C	40	181.0
IL2	A, G, 5moU, C	30	99.4
IL6	A, G, 5moU, C	30	31.2
IL15	A, G, 5moU, C	30	89.8
IL22	A, G, 5moU, C	30	104.0
BDNF	A, G, 5moU, C	30	95.9
BMP2	A, G, 5moU, C	30	112.0
LIF	A, G, 5moU, C	30	116.0
PTH	A, G, 5moU, C	30	88.4
EPO	A, G, 5moU, C	30	83.3

“A” refers to adenosine-5'-triphosphate, “G” refers to guanosine-5'-triphosphate, “U” refers to uridine-5'-triphosphate, “C” refers to cytidine-5'-triphosphate, “7dG” refers to 7-deazaguanosine-5'-triphosphate, “sG” refers to thienoguanosine-5'-triphosphate, “5mC” refers to 5-methylcytidine-5'-triphosphate, “5hmC” refers to 5-hydroxymethylcytidine-5'-triphosphate, “5cC” refers to 5-carboxycytidine-5'-triphosphate, “5fC” refers to 5-formylcytidine-5'-triphosphate, “5hC” refers to 5-hydroxycytidine-5'-triphosphate, “psU” refers to 5-pseudouridine-5'-triphosphate, “5mU” refers to 5-methyluridine-5'-triphosphate, “5hmU” refers to 5-hydroxymethyluridine-5'-triphosphate, “5cU” refers to 5-carboxyuridine-5'-triphosphate, and “5moU” refers to 5-methoxyuridine-5'-triphosphate.

10 Example 2 Preparation of RNA-Transfection-Reagent Complexes

For each microgram of RNA, 1μg RNA and 1μL transfection reagent (LIPOFECTAMINE 3000, Life Technologies Corporation) were first diluted separately in complexation medium (Opti-MEM, Life Technologies Corporation or DMEM/F12 + 10μg/mL insulin + 5.5μg/mL transferrin + 6.7ng/mL sodium selenite + 2μg/mL ethanolamine) to a total volume of between 5μL and 100μL each. Diluted RNA and transfection reagent were then mixed and incubated for 10min at room temperature, according to the transfection reagent-manufacturer's instructions.

Example 3 Transfection of Cells with Synthetic RNA

Complexes were prepared according to Example 2, and were then added directly to cells in culture. For transfection in 6-well plates, between 10 μ L and 250 μ L of complexes were added to each well of the 6-well plate, which already contained 2mL of transfection medium per well. Plates were shaken gently to distribute the complexes throughout the well. Cells were incubated with complexes for 4 hours to overnight, before replacing the medium with fresh transfection medium (2mL/well). Alternatively, the medium was not replaced. Volumes were scaled for transfection in 24-well and 96-well plates.

Example 4 Toxicity of and Protein Translation from Synthetic RNA Containing Non-Canonical Nucleotides

Primary human fibroblasts were transfected according to Example 2, using RNA synthesized according to Example 1. Cells were fixed and stained 20-24h after transfection using an antibody against Oct4. The relative toxicity of the RNA was determined by assessing cell density at the time of fixation.

Example 5 Delivery of Synthetic RNA to the Skin

The complexation reaction shown in **Table 5** was prepared using RNA encoding green fluorescent protein (GFP) or collagen, type VII, alpha1 (COL7), synthesized according to Example 1. The concentration of the RNA stock solution was 500 μ g/mL.

Table 5. RNA Complexation Reaction

RNA solution tube	Volume
GFP or COL7 RNA	8 μ L
FactorPlex™ complexation buffer	42 μ L
Transfection reagent solution tube	
LIPOFECTAMINE 3000 (LIFE TECHNOLOGIES)	4 μ L
FactorPlex™ complexation buffer	46 μ L

Each tube was mixed by pipetting, and the transfection reagent solution tube was incubated for 30s at room temperature. The transfection reagent solution was then transferred to the RNA solution, and the contents were mixed by rapidly pipetting up and down 10 times. Following a 10min incubation, dilutions were prepared according to

Table 6. Injection Solutions

Site	RNA	Complexation Volume	FactorPlex™ Volume	RNA amount
1	GFP	7.5 μ L	22.5 μ L	0.3 μ g
2	GFP	15 μ L	15 μ L	0.6 μ g
3	GFP	30 μ L	0 μ L	1.2 μ g
4	COL7	30 μ L	0 μ L	1.2 μ g

For each injection, the corresponding solution was drawn into a 3cc insulin syringe with an 8mm, 31 gauge needle (Becton, Dickinson and Company, Part Number: 328291) and air bubbles were removed. A clear field was selected on the left forearm of a healthy 33 year-old male human subject, and was disinfected with 70% isopropanol and allowed to dry. The needle was positioned at an angle of approximately 10° to the anterior (palmar) forearm with bevel facing up, and was inserted until the bevel was just covered. 30μL of the RNA solution was injected intradermally over the course of approximately 10sec. A distinct wheal appeared during the injection process. The needle was withdrawn, the wheal remained for approximately 1 minute, and no fluid escaped from the injection site. A total of 4 injections were performed according to **Table 6**, and all of the injections were performed between 11 and 28 minutes following the preparation of the RNA complexation reaction. No swelling, redness, or soreness occurred as a result of the injections. A small amount of bleeding occurred when the needle was removed from sites 2 and 4, resulting in the appearance of a small red spot at these sites.

The injection sites were imaged according to the schedule of **Table 7**, and every 24 hours thereafter for 6 days. Fluorescence images were acquired using an inverted microscope (Nikon Eclipse TS100) equipped with an EXFO X-Cite™ 120 fluorescence illumination system and the filter sets shown in **Table 7**. Fluorescence images were captured using a Sony NEX-7 digital camera (**FIGs. 9-12**).

Table 7. Measurement Parameters

Site	Time	Image Type	Exposure Time
All	0h	Brightfield	Automatic
1	0h	FITC	1/10s
2	0h	FITC	1/10s
3	0h	FITC	1/10s
4	0h	FITC	1/10s
1	12h	FITC	1/10s
2	12h	FITC	1/10s
3	12h	FITC	1/10s
4	12h	FITC	1/10s
1	12h	FITC	1/20s
2	12h	FITC	1/20s
3	12h	FITC	1/20s
4	12h	FITC	1/20s
All	24h	Brightfield	Automatic
1	24h	FITC	1/20s
2	24h	FITC	1/20s
3	24h	FITC	1/20s
4	24h	FITC	1/20s
1	24h	Cy3.5	1/5s
2	24h	Cy3.5	1/5s
3	24h	Cy3.5	1/5s

4	24h	Cy3.5	1/5s
1	24h	Cy3	1/5s
2	24h	Cy3	1/5s
3	24h	Cy3	1/5s
4	24h	Cy3	1/5s
1	36h	FITC	1/20s
2	36h	FITC	1/20s
3	36h	FITC	1/20s
4	36h	FITC	1/20s
1	48h	FITC	1/20s
2	48h	FITC	1/20s
3	48h	FITC	1/20s
4	48h	FITC	1/20s

An independent experiment was carried out using the 1.2µg dose of GFP RNA, with similar results (**FIG. 12**).

Table 8. Filter Sets

Image Type	Filter Set
Cy3	Chroma SP102V2
Cy3.5	Chroma SP103V2
FITC	Chroma SP101

5 *Example 6 Transfection of Human Keratinocytes with RNA Encoding NOVEPOETIN*

RNA encoding NOVEPOETIN was synthesized according to Example 1 with three nucleotide combinations: 1) U, G, U, C, 2) U, G, 5moU, C, and 3) U, G, psU, C. Sub-confluent layers of primary human keratinocytes cultured in EpiLife medium were transfected in wells of a 6 well plate according to Example 3 with 1 µg of RNA per well. 12, 24, 36 and 48h following transfection, 0.5 mL of medium was removed and 0.5 mL of fresh EpiLife medium was added to the plate. After the final medium sampling, the cells were harvested by trypsinization, and total RNA was isolated using the RNeasy mini kit (Qiagen). Genomic DNA was digested using DNase I and RNA was purified. Expression of interferon-β and GAPDH was measured by RT-PCR (**FIG. 5**).

Example 7 Transfection of Human Cells with RNA Encoding hTERT

RNA encoding human telomerase reverse transcriptase (hTERT) was synthesized according to Example 1 with the following nucleotides: U, G, 5moU, C. A sub-confluent layer of primary human dermal fibroblasts cultured in DMEM+10%FBS were transfected in wells of a 24 well plate according to Example 3 with 0.25 µg of RNA per well. 12h after transfection, cells were fixed and stained using a 1:50 dilution of rabbit anti-hTERT antibody (Millipore, Part Number: MABE14) (**FIG. 16**).

Example 8 High-Efficiency Gene Editing by Repeated Transfection with RIBOSLICE

Primary human fibroblasts were plated in 6-well plates coated with recombinant human fibronectin and recombinant human vitronectin (each diluted in DMEM/F12 to a concentration of 1µg/mL, 1mL/well, and incubated at room temperature for 1h) at a density of 10,000 cells/well in transfection medium. The following day, the cells were transfected as in Example 2 with RNA synthesized according to Example 1. The following day, cells in one of the wells were transfected a second time. Two days after the second transfection, the efficiency of gene editing was measured using a mutation-specific nuclease assay.

Example 9 Transfection of Cells with Synthetic RNA Containing Non-Canonical Nucleotides and DNA Encoding a Repair Template

For transfection in 6-well plates, 1µg RNA encoding gene-editing proteins targeting exon 16 of the human APP gene, 1µg single-stranded repair template DNA containing a PstI restriction site that was not present in the target cells, and 6µL transfection reagent (LIPOFECTAMINE RNAiMAX, Life Technologies Corporation) were first diluted separately in complexation medium (Opti-MEM, Life Technologies Corporation) to a total volume of 120µL. Diluted RNA, repair template, and transfection reagent were then mixed and incubated for 15min at room temperature, according to the transfection reagent-manufacturer's instructions. Complexes were added to cells in culture. Approximately 120µL of complexes were added to each well of a 6-well plate, which already contained 2mL of transfection medium per well. Plates were shaken gently to distribute the complexes throughout the well. Cells were incubated with complexes for 4 hours to overnight, before replacing the medium with fresh transfection medium (2mL/well). The next day, the medium was changed to DMEM + 10% FBS. Two days after transfection, genomic DNA was isolated and purified. A region within the APP gene was amplified by PCR, and the amplified product was digested with PstI and analyzed by gel electrophoresis.

Example 10 in vivo RIBOSLICE Safety Study

40 female NCr nu/nu mice were injected subcutaneously with 5×10^6 MDA-MB-231 tumor cells in 50% Matrigel (BD Biosciences). Cell injection volume was 0.2mL/mouse. The age of the mice at the start of the study was 8 to 12 weeks. A pair match was conducted, and animals were divided into 4 groups of 10 animals each when the tumors reached an average size of 100-150mm³, and treatment was begun. Body weight was measured every day for the first 5 days, and then biweekly to the end of the study. Treatment consisted of RIBOSLICE BIRC5-1.2 complexed with a vehicle (LIPOFECTAMINE 2000, Life Technologies Corporation). To prepare the dosing solution for each group, 308µL of complexation buffer (Opti-MEM, Life Technologies Corporation) was pipetted into each of two sterile, RNase-free 1.5mL tubes. 22µL of RIBOSLICE BIRC5-1.2 (500ng/µL) was added to one of the two tubes, and the contents of the tube were mixed by pipetting. 22µL of vehicle was added to the second tube. The contents of the second tube were mixed, and then transferred to the first tube, and mixed with the contents of the first tube by pipetting to form complexes. Complexes were incubated at room temperature for 10min. During the incubation, syringes were loaded. Animals were injected either intravenously or intratumorally with a total dose of 1µg RNA/animal in 60µL total volume/animal. A total of 5 treatments were given, with injections performed every other day. Doses were not adjusted for body weight. Animals were followed for 17

days. No significant reduction in mean body weight was observed, demonstrating the *in vivo* safety of RIBOSLICE gene-editing RNA.

Example 11 Screening of Reagents for Delivery of Nucleic Acids to Cells

Delivery reagents including polyethyleneimine (PEI), various commercial lipid-based transfection reagents, a peptide-based transfection reagent (N-TER, Sigma-Aldrich Co. LLC.), and several lipid-based and sterol-based delivery reagents were screened for transfection efficiency and toxicity *in vitro*. Delivery reagents were complexed with RIBOSLICE BIRC5-1.2, and complexes were delivered to HeLa cells in culture. Toxicity was assessed by analyzing cell density 24h after transfection. Transfection efficiency was assessed by analyzing morphological changes. The tested reagents exhibited a wide range of toxicities and transfection efficiencies. Reagents containing a higher proportion of ester bonds exhibited lower toxicities than reagents containing a lower proportion of ester bonds or no ester bonds.

Example 12 High-Concentration Liposomal RIBOSLICE

High-Concentration Liposomal RIBOSLICE was prepared by mixing 1 μ g RNA at 500ng/ μ L with 3 μ L of complexation medium (Opti-MEM, Life Technologies Corporation), and 2.5 μ L of transfection reagent (LIPOFECTAMINE 2000, Life Technologies Corporation) per μ g of RNA with 2.5 μ L of complexation medium. Diluted RNA and transfection reagent were then mixed and incubated for 10min at room temperature to form High-Concentration Liposomal RIBOSLICE. Alternatively, a transfection reagent containing DOSPA or DOSPER is used.

Example 13 In Vivo RIBOSLICE Efficacy Study – Subcutaneous Glioma Model

40 female NCr nu/nu mice were injected subcutaneously with 1×10^7 U-251 tumor cells. Cell injection volume was 0.2mL/mouse. The age of the mice at the start of the study was 8 to 12 weeks. A pair match was conducted, and animals were divided into 4 groups of 10 animals each when the tumors reached an average size of 35-50mm³, and treatment was begun. Body weight was measured every day for the first 5 days, and then biweekly to the end of the study. Caliper measurements were made biweekly, and tumor size was calculated. Treatment consisted of RIBOSLICE BIRC5-2.1 complexed with a vehicle (LIPOFECTAMINE 2000, Life Technologies Corporation). To prepare the dosing solution, 294 μ L of complexation buffer (Opti-MEM, Life Technologies Corporation) was pipetted into a tube containing 196 μ L of RIBOSLICE BIRC5-1.2 (500ng/ μ L), and the contents of the tube were mixed by pipetting. 245 μ L of complexation buffer was pipetted into a tube containing 245 μ L of vehicle. The contents of the second tube were mixed, and then transferred to the first tube, and mixed with the contents of the first tube by pipetting to form complexes. Complexes were incubated at room temperature for 10min. During the incubation, syringes were loaded. Animals were injected intratumorally with a total dose of either 2 μ g or 5 μ g RNA/animal in either 20 μ L or 50 μ L total volume/animal. A total of 5 treatments were given, with injections performed every other day. Doses were not adjusted for body weight. Animals were followed for 25 days.

Example 14 Liposome Formulation and Nucleic-Acid Encapsulation

Liposomes are prepared using the following formulation: 3.2mg/mL N-(carbonyl-ethoxypolyethylene glycol 2000)-1,2-distearoyl-sn-glycero-3-phosphoethanolamine (MPEG2000-DSPE), 9.6mg/mL fully hydrogenated phosphatidylcholine, 3.2mg/mL cholesterol, 2mg/mL ammonium sulfate, and histidine as a buffer. pH is controlled using sodium hydroxide and isotonicity is maintained using sucrose. To form liposomes, lipids are mixed in an organic solvent, dried, hydrated with agitation, and sized by extrusion through a polycarbonate filter with a mean pore size of 800nm. Nucleic acids are encapsulated by combining 10µg of the liposome formulation per 1µg of nucleic acid and incubating at room temperature for 5 minutes.

Example 15 Folate-Targeted Liposome Formulation

Liposomes are prepared using the following formulation: 3.2mg/mL N-(carbonyl-ethoxypolyethylene glycol 2000)-1,2-distearoyl-sn-glycero-3-phosphoethanolamine (MPEG2000-DSPE), 9.6mg/mL fully hydrogenated phosphatidylcholine, 3.2mg/mL cholesterol, 2mg/mL ammonium sulfate, and histidine as a buffer, with 0.27mg/mL 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[folate(polyethylene glycol)-5000] (FA-MPEG5000-DSPE) added to the lipid mixture. pH is controlled using sodium hydroxide and isotonicity is maintained using sucrose. To form liposomes, lipids are mixed in an organic solvent, dried, hydrated with agitation, and sized by extrusion through a polycarbonate filter with a mean pore size of 800nm. Nucleic acids are encapsulated by combining 10µg of the liposome formulation per 1µg of nucleic acid and incubating at room temperature for 5 minutes.

Example 16 Therapy Comprising Liposomal Protein-Encoding RNA

Liposomes encapsulating synthetic RNA encoding a therapeutic protein, synthesized according to Example 1, are prepared according to Example 14 or Example 15. The liposomes are administered by injection or intravenous infusion.

Example 17 Generation of Elastin ivT-RNA template

Total RNA was extracted from neonatal human dermal fibroblasts using the RNeasy mini kit (QIAGEN GmbH), according to the manufacturer's instructions. cDNA encoding human elastin was prepared using MonsterScript™ Reverse Transcriptase (Epicentre Biotechnologies) and the primer: AAAAAAACCGGTTCATTTTCTCTTCCGGCCAC (SEQ ID NO:483). An *in vitro* transcription (ivT) template was prepared from the cDNA by PCR amplification of the elastin coding sequence (CDS) using the primers: F: AAAAAAGCTAGCATGGCGGGTCTGACG (SEQ ID NO:484), and R: AAAAAAACCGGTTCATTTTCTCTTCCGGCCAC (SEQ ID NO:485). The PCR product was then purified using agarose gel electrophoresis and the QIAquick Gel Extraction Kit (QIAGEN GmbH) and was cloned into a vector containing the human beta globin (HBB) 5' and 3' untranslated regions and a strong Kozak sequence. The vector was amplified, purified, and linearized prior to RNA synthesis.

Example 18 Generation of Tyrosinase ivT-RNA Template

Total RNA was extracted from human epidermal melanocytes using the RNeasy mini kit (QIAGEN GmbH), according to the manufacturer's instructions. cDNA encoding human tyrosinase was prepared using MonsterScript™ Reverse Transcriptase (Epicentre Biotechnologies). An *in vitro* transcription (ivT) template was prepared from the cDNA by PCR amplification of the tyrosinase coding sequence (CDS). The PCR product was then purified using agarose gel electrophoresis and the QIAquick Gel Extraction Kit (QIAGEN GmbH) and was cloned into a vector containing the human beta globin (HBB) 5' and 3' untranslated regions and a strong Kozak sequence. The vector was amplified, purified, and linearized prior to RNA synthesis.

Example 19 Synthesis of Tyrosinase RNA

RNA encoding human tyrosinase was synthesized according to Example 1, using the DNA template of Example 18 and the T7 High Yield RNA Synthesis Kit (New England Biolabs, Inc.), according to the manufacturer's instructions (**Table 4**). Samples of the RNA were analyzed by agarose gel electrophoresis to assess the quality of the RNA. The RNA was then diluted to 1 µg/µL. The RNA solution was stored at 4°C.

Example 20 Increasing Melanin Production in Skin by Transdermal Injection via Syringe of RNA Encoding Tyrosinase

The RNA of Example 19 was loaded into a syringe and delivered to the dermis of the ventral forearm of a healthy 33 year-old male patient over the course of approximately 30 seconds.

Example 21 Increasing Melanin Production in Skin by Combined Delivery of RNA Encoding Tyrosinase and Electroporation

The area of skin treated in Example 20 was exposed to electrical pulses of between 10V and 155V and between approximately 10 milliseconds and approximately 1 second using a two-electrode array electrically connected to a capacitor. The patient reported a tingling sensation at all voltages and penetration depths. The treated area became darker after 24-48 hours. The experiment was repeated several times, with similar results.

Example 22 Increasing Melanin Production in Skin by Topical or Intradermal Application of RNA Encoding Tyrosinase

The RNA of Example 19 or the liposomes of Example 16 are applied directly to the skin, with or without disruption of the stratum corneum or injected intradermally using a dose of one microgram or less per square centimeter. Optionally, an electric field is applied as in Example 21 or using a surface-contact patch to enhance delivery of the RNA.

Example 23 Increasing Elastin Production in Skin by Transdermal Delivery of RNA Encoding Elastin

RNA encoding elastin was prepared according to Example 1. The RNA is delivered as in Example 20, 21, or 22.

Example 24 Increasing Collagen Production in Skin by Transdermal Delivery of RNA Encoding Collagen

RNA encoding collagen was prepared according to Example 1. The RNA is delivered as in Example 20, 21, or 22.

Example 25 Anemia Therapy Comprising Delivery of RNA Encoding NOVEPOETIN

RNA encoding NOVEPOETIN was prepared according to Example 1. The RNA is delivered as in Example 20,
5 21, or 22.

Example 26 Increasing Production of Actin in Skeletal Muscle by Intramuscular Delivery of RNA Encoding Actin

RNA encoding actin is prepared according to Example 1. The RNA is delivered to the patient via intramuscular injection with or without the use of an electric field as in Example 20, 21, or 22.

Example 27 Wound Healing Treatment

10 RNA encoding basic fibroblast growth factor is prepared according to Example 1. The RNA is delivered as in Example 20, 21, or 22.

Example 28 Anti-Scarring Treatment

RNA encoding collagenase is prepared according to Example 1. The RNA is delivered as in Example 20, 21, or 22.

15 *Example 29 Production of Botulinum Toxin*

RNA encoding botulinum toxin is prepared according to Example 1. The RNA is delivered as in Example 20, 21, or 22.

Example 30 Increasing Collagen Production in Skin Cells by Transfection with RNA Encoding Collagen I

RNA comprising the coding sequence of the human COL1A1 gene was synthesized according to Example 1.
20 Primary human dermal fibroblasts were plated in wells of a 24-well plate, and were transfected according to Example 2. Between 24 and 72 hours after transfection, the cells were fixed and stained using an antibody targeting collagen I. Many extracellular deposits of collagen were visible in the transfected wells.

Example 31 Increasing Collagen Production in Skin Cells by Transfection with RNA Encoding Collagen VII

RNA comprising the coding sequence of the human COL7 gene was synthesized according to Example 1.
25 Primary human dermal fibroblasts were plated in wells of a 24-well plate, and were transfected according to Example 2. Between 24 and 72 hours after transfection, the cells were fixed and stained using an antibody targeting collagen VII. Transfected cells exhibited high levels of collagen VII, compared to an un-transfected control.

*Example 32 Increasing Collagen Production in Skin by Transdermal Injection via Syringe of RNA Encoding
30 Collagen I or Collagen VII*

RNA comprising the coding sequence of the human COL1A1 gene or the human COL7 gene was synthesized according to Example 1. The RNA is loaded into a syringe and delivered to the dermis of a patient over the course of approximately 30 seconds or as in Example 20, 21, or 22.

5 *Example 33 Increasing Collagen Production in Skin by Combined Delivery of RNA Encoding Collagen I or Collagen VII and Electroporation*

The area of skin treated in Example 32 is exposed to electrical pulses of between 10V and 155V and between approximately 50 microseconds and approximately 1 second using a multi-electrode array electrically connected to a power source.

Example 34 Storage and Stability of Synthetic RNA Complexes

10 A complexation reaction using RNA encoding GFP was prepared according to Example 5. Following the 10 min incubation, the complexation reaction was divided into three equal parts, one of which was diluted 1:10 in FactorPlex™ complexation medium, one of which was diluted 1:10 in sterile, nuclease-free water, and one of which was left undiluted. Each of the three parts was then further divided into four equal parts, one of which was applied to primary human dermal fibroblasts according to Example 3, one of which was left at room temperature
15 for six hours before applying to primary human dermal fibroblasts according to Example 3, one of which was placed at 4°C for six hours before applying to primary human dermal fibroblasts according to Example 3, and one of which was snap frozen in liquid nitrogen and placed at -80°C for six hours before applying to primary human dermal fibroblasts according to Example 3. The cells were imaged using a fluorescence microscope approximately 24 hours after the first transfection (**FIG. 17**). All wells contained GFP-positive cells, demonstrating
20 that the synthetic RNA complexes were stable and maintained activity in all of the storage conditions tested.

Example 35: In Vivo Analysis of NOVECRIT

RNA encoding NOVEPOETIN was synthesized according to Example 1 with the nucleotide combination A, G, 5moU, C. In this Example, NOVECRIT was formulated with a lipid delivery vehicle, specifically LIPOFECTAMINE 3000. In this Example, NOVECRIT encoded NOVEPOETIN, a novel, high-stability erythropoiesis-stimulating
25 agent.

A 15-day maximum tolerated dose (MTD) study was performed to evaluate safety (rat, n=62). *In vivo* toxicology and biodistribution, as well as pharmacodynamics and dose response and therapeutic effect, specifically on erythropoiesis, were evaluated. Furthermore, the immune response was monitored by analysis of cytokines in treated animals.

30 Sixty-nine, naïve, 8 week old male Sprague-Dawley rats (*Rattus norvegicus*) weighing 253 to 274 grams at receipt were used (HARLAN LABORATORIES). Animals were acclimated to the study room for at least seven days prior to dosing. On Day -2 all animals were shaved in the dorsal lumbar area and four intradermal sites (upper left, upper right, lower right and lower left) were designated on the dorsal back using a permanent marker.

Sixty-four animals were randomly assigned to four treatment groups, with the remaining five animals serving as spares. Two animals were dropped from the study because of incomplete dosing.

The following study design was used (total dose volumes (μL) were constant):

Group	Dose Level (μg)	Dose Route	Conc. ($\mu\text{g/mL}$)	Dose Vol. (μL)	Number of Males
1	0	Intradermal	0	50	$4^a + 2^b + 2^h$
2	0.25	Intradermal	5.0	50	$4^a + 2^b + 2^c + 2^d + 2^e + 2^f + 2^g + 2^h$
3	1.0	Intradermal	20	50	$4^a + 2^b + 2^c + 2^d + 2^e + 2^f + 2^g + 2^h$
4	4 X 1.0 μg (4.0 μg total)	Intradermal	20	50	$4^a + 2^b + 3^c + 3^d + 2^e + 2^f + 2^g + 2^h$

- 5 ID: ^a Toxicology animals necropsy on Day 15, TK animals with terminal tissue collections on Days 6 (^b) and 24 (^c) hours postdose, and on Days 3 (^d), 4 (^e), 6 (^f), 8 (^g), and 15 (^h)

Blood was collected from the vena cava from anesthetized animals prior to necropsy or terminal tissue collection. Whenever possible, blood was collected via a single draw and then divided appropriately. Blood specimens for toxicokinetic analysis were collected from two animals per time point for Group 1 at 6 hours postdose and on Day 10 15. For Groups 2-4, blood specimens were collected at 6 and 24 hours postdose, and on Days 3, 4, 6, 8, and 15.

Blood specimens for hematology assessment were collected from all toxicology animals on Day 15 following an overnight fast, and all TK animals scheduled for terminal tissue collections at 6 and 24 hours postdose and on Days 8 and 15. Whole blood (1.3 mL) was deposited into K2EDTA tubes and analyzed using an Advia 120 automated analyzer.

- 15 Blood specimens for coagulation assessment were collected from all toxicology animals on Day 15 following an overnight fast. Coagulation specimens (1.8 mL) were collected into 3.2% sodium citrate tubes, processed according to standard procedures and analyzed using a STA Compact automated analyzer.

- 20 Blood specimens for serum chemistry assessments were collected from all toxicology animals on Day 15 following an overnight fast. Serum chemistry specimens (1 mL) were collected into serum separator tubes, processed according to standard procedures and analyzed using an AU680 analyzer.

- 25 Blood specimens for cytokine analysis were collected from animals scheduled for terminal tissue collection at 5 and 24 hours postdose and Day 8. Cytokine specimens (1 mL) were collected into K2EDTA tubes and processed to plasma according to standard procedures. TNF α , IL-6, and IFN α cytokine levels in plasma samples were studied. The study measured the production of TNF α , IL-6, and IFN α cytokines in plasma samples following dosing with the test article on Day 1 (6 hours post-dose), Day 2 (24 hours post-dose) and Day 8.

Study Outline for Cytokine Analysis Samples:

Group	NOVECRIT Dose Level (μg)	Dose Concentration ($\mu\text{g/mL}$)	Dose Volume (μL)	Number of Rats (Males)	Endpoints
1	0	0	50	2 ^b	Plasma TNF α , IL-6, and IFN α cytokine analysis
2	0.25	5.0	50	2 ^b + 2 ^c + 2 ^g	
3	1.0	20	50	2 ^b + 2 ^c + 2 ^g	
4	4 x 1.0 μg (4.0 μg total)	20	50	2 ^b + 2 ^c + 2 ^g	

b = plasma samples collected 6 hours post-dose, c = plasma samples collected 24 hours post-dose, g = plasma samples collected 8 days post-dose.

For TNF α analysis, plasma samples were analyzed using a commercially-available assay kit from R&D
 5 SYSTEMS (cat. no. RTA00). The rat TNF α ELISA is a solid phase enzyme-linked immunosorbent assay. The ELISA kit employs a TNF α -specific anti-rat monoclonal antibody for solid phase immobilization (pre-coated on the microtiter plate), and HRP conjugated to an anti-rat TNF α polyclonal antibody for detection. The reference standard provided with the kit is a recombinant rat TNF α . For each assay, plasma samples and standards were diluted with the diluent provided in each respective kit. For the TNF α ELISA, plasma samples were diluted 1:2.
 10 The standard curve consisted of six (6) serial 2-fold concentrations ranging from 800 to 12.5 pg/mL. Controls were reconstituted with diluent, and blanks contained diluent only.

Following sample, standard, control, or diluent addition (50 μL per well, each in duplicate), plates were incubated for 2 hours at room temperature. Following a wash step to remove unbound substances, HRP conjugate was added to each well (100 μL /well), and plates were incubated for 2 hours at room temperature. Following a
 15 second wash step, 100 μL /well TMB substrate was added to the plate. The plate was incubated for 30 minutes at room temperature, protected from light, to allow the color reaction to develop. The reaction was stopped following the addition of HCl Stop Solution (100 μL /well). The optical density was read on a SpectraMax 340 (MOLECULAR DEVICES) plate reader at 450 nm, within 30 minutes following addition of Stop Solution. The intensity of the color measured was in proportion to the amount of rat TNF α bound in the initial step. A standard
 20 curve was generated for each assay plate, and test sample TNF α concentrations were determined by interpolation of absorbance A_{450} values from the standard curve and dilution factor. The assay range for the TNF α kit is 12.5 - 800 pg/mL, with a minimum detectable concentration of less than 10 pg/mL (with the 1:2 minimum required dilution for plasma samples).

For IL-6 analysis, plasma IL-6 samples were analyzed using a commercially-available assay kit from R&D
 25 SYSTEMS (cat. no. R6000B – IL-6). The rat IL-6 ELISA is a solid phase enzyme-linked immunosorbent assay.

The ELISA kit employs anti-rat IL-6 monoclonal antibody for solid phase immobilization (pre-coated on the microtiter plate), and HRP conjugated to IL-6-specific anti-rat polyclonal antibody for detection. The reference standard provided with the kit is a recombinant rat IL-6. Plasma samples were used undiluted (as provided), and standards were diluted with the diluent provided in the kit. The standard curve consisted of six (6) serial 2-fold concentrations ranging from 4,000 to 62.5 pg/mL. Controls were reconstituted with diluent, and blanks contained diluent only. Following sample, standard, control, or diluent addition (50 μ L per well, each in duplicate), plates were incubated for 2 hours at room temperature. Following a wash step to remove unbound substances, HRP conjugate was added to each well (100 μ L/well), and plates were incubated for 2 hours at room temperature. Following a second wash step, 100 μ L/well TMB substrate was added to the plate. The plate was incubated for 30 minutes at room temperature, protected from light, to allow the color reaction to develop. The reaction was stopped following the addition of HCl Stop Solution (100 μ L/well). The optical density was read on a SpectraMax 340 (MOLECULAR DEVICES) plate reader at 450 nm, within 30 minutes following addition of Stop Solution. The intensity of the color measured was in proportion to the amount of rat IL-6 bound in the initial step. A standard curve was generated for each assay plate, and test sample IL-6 concentrations were determined by interpolation of absorbance A450 values from the standard curve and dilution factor. The assay range for the IL-6 kit is 62.5 – 4,000 pg/mL, with a minimum detectable concentration of less than 21 pg/mL.

For IFN α analysis, plasma IFN α samples were analyzed using a commercially-available assay kit from NOVATEINBIO (cat. no. BG-RAT11380). The ELISA is a solid phase enzyme-linked immunosorbent assay. The ELISA kit employs anti-rat IFN α monoclonal antibody for solid phase immobilization (pre-coated on the microtiter plate), and HRP conjugated antibody specific for IFN α for detection. Plasma samples were diluted 1:4, and the standards were used as provided in the kit. Standards consisted of six (6) serial 2-fold concentrations ranging from 100 to 3.1 pg/mL. Blanks contained diluent only. Following sample, standard, or diluent addition (50 μ L per well, each in duplicate), HRP conjugate was added to each well (100 μ L/well), and plates were incubated for 1 hours at 37°C. Following a wash step, 50 μ L/well each of Chromogen Solution A and Chromogen Solution B were added to the plate. The plate was incubated for 15 minutes at 37°C, protected from light, to allow the color reaction to develop. The reaction was stopped following the addition of Stop Solution (50 μ L/well). The optical density was read on a SpectraMax 340 (MOLECULAR DEVICES) plate reader at 450 nm. The intensity of the color measured was in proportion to the amount of rat IFN α bound in the initial step. A standard curve was generated for each assay plate, and test IFN α concentrations were determined by interpolation of absorbance A450 values from the standard curve and dilution factor. The assay range for the IFN α kit is 3.1 – 100 pg/mL, with a minimum detectable concentration of less than 1 pg/mL (4 pg/mL, with the 1:4 dilution required for plasma samples).

The results of this study were, *inter alia*, that NOVECRIT was not detected at the injection site 24h after dosing (as assayed by RT-PCR). More specifically, NOVECRIT was not detected in any of the following samples (RT-PCR): serum (6h, 24h, 48h and 72h after dosing), liver (6h and 24h after dosing), and kidney (6h and 24h after

dosing). Further, positive controls (spiked with NOVECRIT) yielded a robust signal in all tissues (as assayed RT-PCR). **FIG. 18** depicts single administration of NOVECRIT induced a rapid increase and sustained level of NOVEPOIETIN in serum. The Y axis shows concentration of NOVEPOIETIN protein (mU/mL). This suggests, without wishing to be bound by theory, that the studied RNA therapeutic is able to provide therapeutically important pharmacodynamic properties. In fact, this PD behavior is vastly improved relative to wild type EPO (which is known in the art to have a half-life of about 4-12 hours).

Furthermore, serum spiked with NOVECRIT showed near-instant degradation of the RNA (as assayed by RT-PCR). Without wishing to be bound by theory, this data indicates that the present RNA therapeutics are safe and have little chance of toxicity limitations, for example, liver or kidney toxicities.

FIG. 19 depicts a single administration of NOVECRIT stimulated erythropoiesis, yielding elevated hematocrit for at least 14 days. The left panel shows % hematocrit on the Y axis, while the right panel shows % reticulocytes. Accordingly, the studied RNA therapeutic is fully functional.

With respect to cytokines, IL-6 and TNF α cytokine levels were below assay detection thresholds for all animals at each of the timepoints evaluated in this study. Low levels of IFN- γ were only detectable in the Day 8 samples from animals in Groups 3 and 4. **FIG. 20** depicts a table summarizing TNF α , IL-6, and IFN α cytokine levels in plasma samples collected from a maximum tolerated dose of NOVECRIT in male Sprague Dawley rats. Without wishing to be bound by theory, this data indicates that the present RNA therapeutics do not stimulate an unfavorable immunogenicity which has limited the therapeutic utility of certain RNA therapeutics.

Example 36: Gene Editing of the COL7A Gene

The present RNA-based gene editing approaches were applied to the COL7A1 gene. This gene is of interest because, *inter alia*, it is frequently involved in dystrophic epidermolysis bullosa. **FIG. 21** depicts a SURVEYOR assay using the DNA of primary adult human dermal fibroblasts transfected with RNA TALENs targeting the sequence TGAGCAGAAGTGGCTCAGTG (SEQ ID NO:467) and TGGCTGTACAGCTACACCCC (SEQ ID NO:468), located within the COL7A1 gene. The bands present in the +RNA lane indicate editing of a region of the gene that is frequently involved in dystrophic epidermolysis bullosa. **FIG. 22** depicts another SURVEYOR assay using the DNA of primary adult human dermal fibroblasts transfected with RNA TALENs, now targeting the sequence TTCCACTCCTGCAGGGCCCC (SEQ ID NO:469) and TCGCCCTTCAGCCCGCGTTC (SEQ ID NO:470), located within the COL7A1 gene. The bands present in the +RNA lane indicate editing of a region of the gene that is frequently involved in dystrophic epidermolysis bullosa. This data points to, among others, a gene editing approach to the treatment of certain genetic disorders such as dystrophic epidermolysis bullosa.

Example 37: Gene-editing of the MYC gene using a Synthetic RNA with Non-Canonical Nucleotides

Experiments were conducted with *in vitro* transcribed synthetic RNA molecules containing non-canonical nucleotides and encoding gene-editing proteins. The immunogenicity and the gene-editing efficiency of *in vitro* transcribed synthetic RNA molecules having (1) only pseudouridine (psU) as a non-canonical nucleotide; (2) only

5-methylcytidine (5mC) as a non-canonical nucleotide; and (3) both of pseudouridine and 5-methylcytidine as non-canonical nucleotides was evaluated (as well as controls).

Specifically, mRNA containing the following nucleotide combinations: (i) A,G,U,C, (ii) A,G,psU,C, (iii) A,G,U,5mC, and (iv) A,G,psU,5mC, and encoding TALEN pairs targeting the following DNA sequences, which can be found within the MYC gene: TCGGCCGCCGCCAAGCTCGT (SEQ ID NO:474) and TGCGCGCAGCCTGGTAGGAG (SEQ ID NO:475), were synthesized according to the methods described herein. Human dermal fibroblasts (MA001SK) were plated in 6-well and 24-well tissue culture plates in DMEM with 10% FBS at 100,000 and 10,000 cells per well, respectively. The next day, the cells were transfected in the 6-well plate with 2 µg of RNA (1 µg for each component of the TALEN pair) and the cells were transfected in the 24-well plate with 0.2 µg of RNA (0.1 µg for each component of the TALEN pair) according to the methods described herein. 24 hours after transfection, the total RNA from the cells in the 24-well culture plate was isolated using an RNeasy Mini Kit (74106; QIAGEN), including isolating the total RNA from a sample of cells that had not been transfected with RNA (negative control; "Neg." in **FIG. 23**). The genomic DNA was removed by a 15 minute digestion with DNase I (RNase-Free) (M0303L; NEW ENGLAND BIOLABS) and the reaction purified using an RNeasy Mini Kit. 1 µL of total RNA was used to assess gene expression by real-time RT-PCR using TAQMAN gene-expression assays (APPLIED BIOSYSTEMS) designed to detect expression of the immunogenicity markers TLR3, IFIT1, and IFIT2 (**FIG. 23**). The data were normalized to both the positive experimental control sample ("A,G,U,C") and to a loading control (GAPDH).

48 hours after transfection, the genomic DNA was isolated from the cells in the 6-well culture plate using a DNeasy Blood and Tissue Kit (69506; QIAGEN), including from a sample of cells that had not been transfected with RNA (negative control, "Neg." in **FIG. 24**). A 970 bp region of the MYC gene surrounding the predicted TALEN cut location was amplified using a 35 cycle 2-step PCR reaction containing the following primers: TAACTCAAGACTGCCTCCCGCTTT (SEQ ID NO: 476) and AGCCCAAGGTTTCAGAGGTGATGA (SEQ ID NO: 477). 160 ng was hybridized in 5 µL of amplified sequence from RNA-treated cells to 160 ng in 5 µL of amplified sequences from untreated MA001SK cells by mixing the two sequences with 0.5 µL of 1M KCl and 0.5 µL of 25mM MgCl₂ and running the following program in a thermocycler: 95°C for 10 minutes; 95°C to 85°C at 0.625C/s; 85°C to 25°C at 0.125C/s. The SURVEYOR assay was performed by adding 0.5µL of SURVEYOR nuclease and 0.5µL of Enhancer from the SURVEYOR Mutation Detection Kit (706020I; INTEGRATED DNA TECHNOLOGIES) to the hybridized product, mixing, and incubating at 42°C for 25 minutes. The protocol above was also used to process the positive control DNA sample provided with the SURVEYOR Mutation Detection Kit as a positive experimental control for the SURVEYOR Assay ("Assay Pos." in **FIG. 24**). Samples were analyzed by agarose gel electrophoresis (**FIG. 24**). For each sample, gene-editing efficiency was calculated as a ratio of the intensity of the digested bands (indicated by "*" in **FIG. 24**) to that of the undigested band.

As shown in **FIG. 23** below, the samples from cells transfected with the positive control RNA (A,G,U,C), and the samples from cells transfected with RNA containing either pseudouridine or 5-methylcytidine exhibited

upregulation of all three of the immunogenicity markers TLR3, IFIT1, and IFIT2. The sample from cells transfected with RNA containing both pseudouridine and 5-methylcytidine exhibited negligible upregulation of the immunogenicity markers (less than 0.01-fold of the positive control), demonstrating that *in vitro* transcribed synthetic RNA with both pseudouridine and 5-methylcytidine and encoding a gene-editing protein can evade
5 detection by the innate-immune system of mammalian cells.

Further, as shown in **FIG. 24** below, the sample from cells transfected with RNA containing both pseudouridine and 5-methylcytidine exhibited highly efficient gene editing (41.7%), which was greater than the efficiency exhibited by samples from cells transfected with RNA containing pseudouridine alone (35.2%), demonstrating that *in vitro* transcribed synthetic RNA comprising both pseudouridine and 5-methylcytidine and encoding a gene-
10 editing protein can both (i) gene-edit mammalian cells at high efficiency, and (ii) gene-edit mammalian cells at higher efficiency than *in vitro* transcribed synthetic RNA comprising pseudouridine and not comprising 5-methylcytidine.

Example 38: COL7A1 Gene Editing and Repair in Human Cells

RNA encoding gene editing proteins targeting the following sequences in the COL7A1 gene was synthesized
15 according to Example 1: TGAGCAGAAAGTGGCTCAGTG (SEQ ID NO:473) and TGGCTGTACAGCTACACCCC (SEQ ID NO:468) (see also table below).

RNA Synthesis

Template	Nucleotides	Reaction Volume/ μ L	ivT Yield/ μ g
COL7A1 TALEN 1L	A, G, 5moU, C	20	120.528
COL7A1 TALEN 1R	A, G, 5moU, C	20	120.204
COL7A1 TALEN 1L	A, G, 5moU, C	15	81.94
COL7A1 TALEN 1R	A, G, 5moU, C	15	61.88

50,000 primary human epidermal keratinocytes (HEKn, Gibco) were plated in wells of 6-well plates in EpiLife +
20 Supplement S7. The next day, cells were transfected according to Example 3 with 1 μ g of RNA encoding each component of the gene editing pair and 2 μ g of a single-stranded DNA repair template having a length of 60, 70, 80, 90 or 100 nucleotides ("nt"). 48 hours after transfection, genomic DNA was purified. A segment of the COL7A1 gene was amplified using the primers GCATCTGCCCTGCGGGAGATC (SEQ ID NO:478) and CCACGTTCTCCTTTCTCTCCCGTTC (SEQ ID NO:479), which produce a 535 bp amplicon. The efficiency of
25 gene editing was assessed using T7 Endonuclease I ("T7E1", New England Biolabs) according to the manufacturer's instructions. Bands of approximately 385 bp and 150 bp indicate successful gene editing. **FIG. 25** and **FIG. 29** show the result of digestion with T7E1, analyzed by agarose gel electrophoresis. **FIG. 27** and **FIG. 29**

show the result of digestion with Mlul-HF, analyzed by agarose gel electrophoresis. Because the repair template contains the sequence ACGCGT (SEQ ID NO:480), digestion of the amplified product with Mlul-HF (New England Biolabs) produces bands of approximately 385 bp and 150 bp in the case of successful gene repair.

RNA encoding gene editing proteins targeting the following sequences in the COL7A1 gene was synthesized according to Example 1: TGAGCAGAAAGTGGCTCAGTG (SEQ ID NO:473) and TGGCTGTACAGCTACACCCC (SEQ ID NO:468). 50,000 primary human epidermal keratinocytes (HEKn, Gibco) were plated in wells of 6-well plates in EpiLife + Supplement S7. The next day, cells were transfected according to Example 3 with 1 µg of RNA encoding each component of the gene editing pair and 1-4 µg of an 80 nucleotide single-stranded DNA repair template. 48 hours after transfection, genomic DNA was purified. A segment of the COL7A1 gene was amplified using the primers GCATCTGCCCTGCGGGAGATC (SEQ ID NO:481) and CCACGTTCTCCTTTCTCTCCCGTTC (SEQ ID NO:482), which produce a 535 bp amplicon. The efficiency of gene editing was assessed using T7 Endonuclease I ("T7E1", New England Biolabs) according to the manufacturer's instructions. Bands of approximately 385 bp and 150 bp indicate successful gene editing. **FIG. 26** show the result of digestion with T7E1, analyzed by agarose gel electrophoresis. **FIG. 28** show the result of digestion with Mlul-HF, analyzed by agarose gel electrophoresis. Because the repair template contains the sequence ACGCGT (SEQ ID NO:480), digestion of the amplified product with Mlul-HF (New England Biolabs) produces bands of approximately 385 bp and 150 bp in the case of successful gene repair.

Example 39: Expression of BMP7 Variants in Human Cells

RNA encoding wild type BMP7 and RNA encoding variants of BMP7 was synthesized according to Example 1 (see also table below).

RNA Synthesis

Template	Nucleotides	Reaction Volume/µL	ivT Yield/µg
BMP7 Wild Type	A, G, 5moU, C	15	125.8
BMP7 Variant A	A, G, 5moU, C	15	120.36
BMP7 Variant B	A, G, 5moU, C	15	143.14
BMP7 Variant C	A, G, 5moU, C	15	106.42

50,000 primary human dermal fibroblasts (MA001SK, Factor Bioscience) or 100,000 primary human epidermal keratinocytes (HEKn, Gibco) were plated in DMEM+10% FBS or EpiLife + Supplement S7, respectively. Cells were transfected according to Example 3 with 1 µg of RNA encoding wild type BMP7 or a variant thereof. 24 hours after transfection, the medium was sampled and secreted BMP7 levels were measured with a human BMP7 ELISA kit (ab99985, Abcam) using medium diluted 10-fold according to the manufacturer's instructions.

Secreted BMP7 levels were determined by measuring 450 nm absorbance using a microplate reader (EMax Plus, Molecular Devices). Secreted BMP7 levels are shown in **FIG. 30**.

Example 40: Expression of Parathyroid Hormone (PTH) in Human Cells

RNA encoding PTH was synthesized according to Example 1 (see also table below).

5 *RNA Synthesis*

Template	Nucleotides	Reaction Volume/ μ L	ivT Yield/ μ g
PTH	A, G, 5moU, C	15	51.68

100,000 human epidermal keratinocytes (HEKn, Gibco) were plated in EpiLife + Supplement S7. Cells were transfected according to Example 3 with 1 μ g of RNA encoding PTH. 24 hours after transfection, the medium was sampled and secreted PTH levels were measured using a human PTH ELISA kit (EIA-PTH-1, Ray Biotech) according the manufacturer's instructions. Secreted PTH levels were determined by measuring 450 nm absorbance using a microplate reader (EMax Plus, Molecular Devices). Secreted PTH levels are shown in **FIG. 31**.

Example 41: Intradermal, Subcutaneous, Rectal and Nasal Administration of NOVECRIT for the Treatment of Anemia

15 A repeat dose toxicity study of Novecrit was conducted for the treatment of anemia. Specifically, 8-10 weeks old male Sprague Dawley rats were administered with Novecrit via intradermal, subcutaneous, rectal, or nasal routes once per day on days 1, 8, and 15. The animals were assigned to groups and treated as indicated in the table below:

Group	Group Color	Test Article	Dose Route	Dose Level (μ g)	Dose Concentration (μ g/mL)	Dose Volume (μ L)	Number of Animals
							Males
1	White	Control	Intradermal	0.0	0.0	50	3 ^a + 8 ^b
2	Yellow	Novecrit	Intradermal	4X (0.25 μ g /injection; 1.0 μ g total)	5.0	50	3 ^a + 8 ^b
3	Green	Novecrit	Subcutaneous	4.0	20.0	200	3 ^a + 8 ^b
4	Blue	Novecrit	Rectal	4.0	20.0	200	3 ^a + 8 ^b
5	Red	Novecrit	Nasal	4.0	20.0	200	3 ^a + 8 ^b

20 Note: Total dose volume (μ L/per animal) are constant

^a Toxicity animals (also called main study animals), necropsy on Day 44

^b TK animals, euthanized as n = 2/time point on Days 2, 16, 23 and 44

More particularly, Group 1 was dosed via intradermal injection. Each dose was administered in four intradermal
 5 injections of 50 μ L/injection for a total of 200 μ L per animal. Injections were carried out on previously marked
 sites near the midline of the dorsal lumbar area (upper left, upper right, lower left and lower right quadrants).
 Group 2 was dosed four intradermal injections at 0.25 μ g each (1.0 μ g total) into previously marked sites near
 the midline of the dorsal lumbar area (upper left, upper right, lower left and lower right quadrants). Group 3 was
 dosed by subcutaneous injections into an area of the back located in the lower dorsal thoracic/lumbar region.
 10 Group 4 was dosed by rectal administration. The animal was manually restrained and the abdomen of each rat
 was manually palpated to remove any fecal matter. If deemed necessary, the rectum was lavaged with up to 2 ml
 of saline for enema followed by manual palpation of the abdomen to remove the fecal matter, if any. Novecrit was
 drawn into a syringe and an appropriately sized gavage needle with a rounded tip (ball) was attached to the
 syringe. A lubricant jelly was applied to the insertion device to aid with insertion; it was advanced approximately
 15 1cm into the lumen of the colon and the Novecrit instilled. The rat was then maintained in a head-down position
 for approximately 20 - 30 seconds to limit the expulsion of solution. Group 5 was dosed via nasal route. The
 animal was anesthetized (per SNBL USA SOP). The animal laid on its back with the head elevated. The dose
 was dispensed slowly into the nares. Approximately half the dose volume was administered to one nare. The
 remaining dose volume was administered to another nare. The first dose was administered on Day 1, and the
 20 last dose on Day 15.

The rats were clinically monitored including their food consumption and body weight. In addition, blood was
 collected for hematology, coagulation, and serum chemistry analysis as indicated below:

Toxicity Group Specimen collection frequency

Time Point	Hematology	Coagulation	Serum Chemistry
Day 44	1X (n = 3/Group)	1X (n = 3/Group)	1X (n = 3/Group)

25 x = Number of times procedure performed

Analysis was done on the toxicokinetic group as follows:

In-text Table 3: Toxicokinetic Group Specimen collection frequency

Time Point	Hematology^a	TK	Cytokines
Acclimation	1X (n = 8/Group)	-	-
Day 2	-	1X (n = 2/Group, 24 hours postdose)	1X (n = 2/Group, 24 hours postdose)
Day 8	Pre-dose (n = 6/Group)	-	-
Day 15	Pre-dose (n = 6/Group)	-	-
Day 16	-	1X (n = 2/Group, 24 hours postdose)	1X (n = 2/Group, 24 hours postdose)
Day 22	1X (n = 4/Group)	-	-
Day 23	-	1X (n = 2/Group)	1X (n = 2/Group)
Day 29	1X (n = 2/Group)	-	-
Day 36	1X (n = 2/Group)	-	-
Day 43	1X (n = 2/Group)	-	-
Day 44	-	1X (n = 2/Group)	1X (n = 2/Group)

x = Number of times procedure performed

TK = toxicokinetics

5

Terminal necropsy for toxicity animals occurred on Day 44. TK animals were euthanized on Days 2, 16, 23, and 44. Pathological analysis was conducted on the animals.

As shown in **FIG. 32**, administration of NOVECRIT stimulated erythropoiesis resulting in elevated hematocrit for at least 14 days in all four study groups compared to control.

10 *Example 42: Intradermal Administration of RNA Encoding BMP7 Variants for the Treatment of Diabetic Nephropathy*

A ZDSD rat model was utilized to study the effects of RNAs encoding BMP7 variants for the prevention and treatment of diabetic nephropathy. Specifically, ZDSD rats were treated with RNAs encoding BMP7 variants administered intradermally. A schematic of the study design is provided in **FIG. 33**. The animals were assigned to groups and treated as indicated in the table below:

15

Group Number	Treatment	No. of Animals	Dose Level (µg)	Dose Conc. (µg /ml)	Dose Volume (ml/kg)	Frequency	Route of Admin.
1	Vehicle	24	0	0	0.1	2x/wk	I.D.
2	FTB-F1 (wt BMP7)	6	1	5	0.1	2x/wk	I.D.
3	FTB-F2 (BMP7 variant A)	6	1	5	0.1	2x/wk	I.D.
4	FTB-F3 (BMP7 variant B)	6	1	5	0.1	2x/wk	I.D.
5	Lisinopril	6	Standard	Standard	5	Continuous	Diet
6 ^a	Vehicle	20	0	0	0.2/rat	3x/wk	I.D.
7 ^a	FTB-F2 (BMP7 variant A)	10	2	10	0.2/rat	3x/wk	I.D.

^a Group 1 animals at the end of the “Prevention” arm were randomized into Groups 6 and 7 (n=20 or 10 each, respectively) to define the “Treatment” arm. The remaining 6 animals from Group 1 were euthanized for kidney collection and blood draw at the end of the “Prevention” arm.

- 5 To study the effect of the RNAs on prevention of diabetic nephropathy, animals were delivered from the barrier (PCO-5638) to the vivarium (PCO-Rm C) and allowed 3 days for acclimation. All animals were placed on 5SCA diabetogenic diet for 3 weeks (weekly body weight measurements were taken). Animals were then returned to regular 5008 diet for the duration of the study (weekly body weight measurements were taken). After 2 weeks on 5008 diet, body weight measurement, blood draw (500 ul) and a 24 hour baseline urine collection (urinary albumin and creatinine measurements) were performed on all animals. 36 rats were randomized to groups 1, 2 and 3 based on body weight, glucose and urinary albumin. Groups 1, 2, 3 and 4 received either vehicle or test article (i.d.) via intradermal administration (2x/week, Mon and Thur). Group 5 was administered Lisinopril admixed in the 5008 diet (food consumption will begin on this group of animals). Blood draws (500 ul, tail vein) were taken from Groups 2, 3 and 4, 24 hours after each dose for the first 2 weeks and then 1x/week thereafter.
- 10 After 4 weeks of dosing, blood sample via tail vein (500 ul) was taken from all animals (Groups 1-5). At the end of 8 weeks of dosing, a 24 hour urine collection was performed (urinary albumin and creatinine measurements) together with a tail vein (500 ul) blood sample from Group 1 (n=18 going into the Treatment arm). Results for urine volume, creatinine, and albumin is shown in **FIG. 35**. As shown in **FIG. 35**, treatment with RNA encoding BMP7 Variant A resulted in reduced levels of urine albumin in rats afflicted with diabetic nephropathy compared

to the control, indicating superior kidney function in the treated animals compared to the control animals (placebo group). To terminate the study, all animals were euthanized with CO₂ asphyxiation and a blood draw was performed via cardiac puncture to collect serum for BUN, creatinine and glucose, and plasma for compound exposure and biomarker analysis. Both kidneys were dissected and fixed for histology. **FIG. 34** shows the serum level of BMP7 protein in Groups 1, 2, 3, and 4. Results indicate that the serum levels of BMP7 protein in these groups were as follows: Group 4, Group 3 > Group 2, Group 1.

A study was also conducted to analyze the effect of the RNAs encoding BMP7 variants on treating diabetic nephropathy. Specifically, the vehicle animals from Group 1 (Prevention arm, n=18) were randomized based on body weight, glucose and urine albumin (Groups 6, n=20; 7, n=10;). Animals received either Vehicle or BMP7 variant-encoding RNAs (intradermal, 2x/week, Mon and Thur), groups 6 and 7, respectively. After 4 weeks of treatment, animals were euthanized with CO₂ asphyxiation and a blood draw will be performed via cardiac puncture to collect serum for BUN, creatinine and glucose, and plasma for compound exposure and biomarker analysis. Both kidneys were dissected and fixed for histology. As shown in **FIG. 36**, treatment with RNA encoding BMP7 Variant A decreased urine albumin level in rats afflicted with diabetic nephropathy, indicating superior kidney function in the treated animals. This result suggested that BMP7 variant-encoding RNA effectively treated diabetic nephropathy and promoted superior kidney function compared to the control animals (placebo group).

An illustrative study protocol is provided below:

Week	Day	Procedure
15 weeks old	-7	Animals arrive in Room C from the barrier
	0	BW*, Stat Strip, and Place animals on 5SCA for 3 wks
1	7	BW*, Stat Strip
2	14	BW*, Stat Strip
3	21	BW*, Stat Strip, put animals on 5008
4	28	BW*, Stat Strip
	34	BW*, Tail vein blood draw and randomize for Prevention arm (Grps 1, 2, 3, 4, 5)
5	35	Begin 24 hr urine collection
	36	Collect 24 hr urine
	39	- BW* - Intradermal dosing Grps 1,2, 3 and 4, and put Grp 5 on admixed 5008 diet - Begin food consumption on group 5
	40	Collect 24 blood sample for exposure (gps 2, 3, & 4)
	42	Dosing

	43	Collect 24 blood sample for exposure (gps 2, 3, & 4)
6 (1 P)	46	- BW*, Dosing - Food consumption group 5
	47	Collect 24 blood sample for exposure (gps 2, 3, & 4)
	49	Dosing
	50	Collect 24 blood sample for exposure (gps 2, 3, & 4)
7 (2 P)	53	- BW*, Dosing - Food consumption group 5
	54	Collect 24 blood sample for exposure (gps 2, 3, & 4)
	56	Dosing
8 (3 P)	60	- BW*, Dosing - Food consumption group 5
	61	Collect 24 blood sample for exposure (gps 2, 3, & 4)
	63	Dosing
9 (4 P)	67	- BW*, Dosing - Food consumption group 5 - Blood draw (tail vein)
	68	- Blood draw (all groups) - Collect 24 blood sample for exposure (gps 2, 3, & 4)
	70	Dosing
10 (5 P)	74	- BW*, Dosing - Food consumption group 5
	75	Collect 24 blood sample for exposure (gps 2, 3, & 4)
	77	Dosing
11 (6 P)	81	- BW*, Dosing - Food consumption group 5
		Collect 24 blood sample for exposure (gps 2, 3, & 4)
	84	Dosing
12 (7 P)	88	- BW*, Dosing - Food consumption group 5
	89	Collect 24 blood sample for exposure (gps 2, 3, & 4)
	91	Dosing
13 (8 P)	95	- BW*, dosing - Food consumption group 5 - Blood draw (tail vein) (all groups)
	96	Begin 24 hr urine collection

	97	- Collect 24 hr urine - Terminate Groups 1 (n=6), 2, 3, 4 and 5 cardiac puncture blood draw, and collect kidneys, weigh and fix.
	98	- BW* and randomize (BW*, glucose, albumin) Group 1 (n=18) rats into Groups 6, 7 and 8 - Dose - Food Consumption group 8
	103	Dose
14 (1 T)	106	- BW*, Dosing - Food Consumption group 8
	110	Dosing
15 (2 T)	113	- BW*, Dosing - Food Consumption group 8
	117	Dosing
16 (3 T)	120	- BW*, Dosing - Food Consumption group 8
	124	Dosing
17 (4T)	127	BW*, Terminate, cardiac puncture blood draw, and collect kidneys, weigh and fix.

*BW: body weight measurement

In an illustrative study, the protocol was amended from above towards the end of the study as follows:

Treatment Arm	Day	- BW - Begin 24 hr urine collection
	81	Collect 24 hr urine
		- BW, Blood draw - Randomize (BW, glucose, albumin) rats into Groups 6 and 7
1 T		- BW, Dose 12 hr PK sample
		Dose
		Dose and Begin 24 hr Urine collection
	89	Collect 24 hr Urine and Dose
2 T		- BW, Dose 12 hr PK sample
		Dose
		Begin 24 hr Urine collection
	96	Collect 24 hr Urine and Dose
3 T		- BW, Dose 12 hr PK sample
		Dose
		Begin 24 hr Urine collection
		Collect 24 hr Urine and Dose (last dose)
4 T		BW
		No activity

		Begin 24 hr urine collection
		- Collect 24 hr urine - BW, Terminate, cardiac puncture blood draw, and collect kidneys, weigh and fix.

During both studies, pathological analysis is conducted as follows:

Specimen to be Collected	Method of Collection	Anti-Coagulant	Specimen to be Analyzed	Clinical Chemistry	Method of Analysis
Urine	24hr Metabolism Cages	NA	Urine	Creatinine/ Albumin	AU480/ ICL Elisa
Whole Blood	Tail Vein	None	Serum	BUN Creatinine Glucose/	AU480/ Sponsor
Whole Blood	Cardiac Puncture (Grps 1, 2, 3, 4 and 5; and 6, 7, 8)	None	Serum	BUN Creatinine Glucose/ Cmpd Exp & Biomarker	AU480/ Sponsor

Altogether, these results suggested that RNAs encoding BMP-7 variants effectively prevented the development of diabetic nephropathy in rats. In addition, these RNAs also effectively treated and restored kidney functions in rats already afflicted with diabetic nephropathy. RNA encoding BMP7 Variant A was particularly effective at preventing and treating diabetic nephropathy in rats.

Example 43: Transfection of Human Keratinocytes with RNA Encoding BDNF, BMP-2, BMP-6, IL-2, IL-6, IL-15, IL-22, LIF or FGF-21

- 100,000 human epidermal keratinocytes (HEK, Gibco) were plated in EpiLife + Supplement S7. Cells were transfected according to Example 3 with 2 µg of RNA encoding BDNF, BMP-2, BMP-6, IL-2, IL-6, IL-15, IL-22, LIF or FGF-21. 24 hours after transfection, the medium was sampled and secreted protein levels were measured using a human ELISA kit (see Table below) according the manufacturer's instructions. Secreted protein levels were determined by measuring 450 nm absorbance using a microplate reader (EMax Plus, Molecular Devices).
- Secreted protein levels are shown in **FIG. 34**, panels A-I.

Protein	Part Number	Vendor
BDNF	DBNT00	R&D Systems

BMP-2	DBP200	R&D Systems
BMP-6	ab99984	Abcam
IL-2	D2050	R&D Systems
IL-6	D6050	R&D Systems
IL-15	D1500	R&D Systems
IL-22	D2200	R&D Systems
LIF	DLF00	R&D Systems
FGF-21	DF2100	R&D Systems

Example 44: Transfection of Human Keratinocytes with RNA Encoding IL-15 and/or IL-15RA

100,000 human epidermal keratinocytes (HEK, Gibco) were plated in EpiLife + Supplement S7. Cells were transfected according to Example 3 with 2 µg of RNA encoding IL-15 or 1 µg each of RNA encoding IL-15 and RNA encoding IL-15RA. 24 hours after transfection, the medium was sampled and secreted IL-15 levels were measured using a human IL-15 ELISA kit (D1500, R&D Systems) according the manufacturer's instructions. Secreted IL-15 levels were determined by measuring 450 nm absorbance using a microplate reader (EMax Plus, Molecular Devices). Secreted IL-15 levels are shown in **FIG. 35**. As demonstrated in **FIG. 35**, co-transfection of IL-15 and IL-15RA significantly increased secreted IL-15 levels compared to transfection with IL-15 alone.

Example 45: Pharmacokinetic Study Via Intradermal Injection in Rats

Studies were conducted to evaluate the responses of Sprague Dawley rats to intradermal administration of various RNAs. Specifically, 8-10 weeks old, female Sprague Dawley rats weighing about 200 g to about 350 g were used for this study. A total of 33 rats were tested, and the animals were assigned to study groups and treated as indicated in the Table below:

Group	Group Color	Test Article	Dose Route	Dose Level (µg)	Dose Concentration (µg/mL)	Dose Volume (µL/per injection) ^a	Number of Animals
							Females
1	White	Control (NOVEP OEITIN)	ID	4.0	20.0	200 (4 x 50)	3 ^b
2	Yellow	TA1 (IL2)	ID	4.0	20.0	200 (4 x 50)	3 ^b
3	Green	TA2 (IL6)	ID	4.0	20.0	200 (4 x 50)	3 ^b
4	Blue	TA3 (IL15)	ID	4.0	10.0	200 (4 x 50)	3 ^b
5	Red	TA4 (IL15 + IL15RA)	ID	4.0	20.0 (10.0 each)	200 (4 x 50)	3 ^b
6	Dark Grey	TA5 (IL22)	ID	4.0	20.0	200 (4 x 50)	3 ^b

7	Purple	TA6 (BMP2)	ID	4.0	20.0	200 (4 x 50)	3 ^b
8	Black	TA7 (BDNF)	ID	4.0	20.0	200 (4 x 50)	3 ^b
9	White/ Yellow	TA8 (LIF)	ID	4.0	20.0	200 (4 x 50)	3 ^b
10	Green/Blue	TA9 (PTH)	ID	4.0	20.0	200 (4 x 50)	3 ^b
11	Red/Dark Grey	TA10 (FGF21)	ID	4.0	20.0	200 (4 x 50)	3 ^b

^aTotal dose volume (μL /per animal) was constant. Each animal received four intradermal injections of 50 μL /per injection for a total of 200 μL per animal.

^bAnimals, euthanized on Day 3 without examination or necropsy

Intradermal = ID

- 5 For the study, the animals were treated with either 4 μg of RNA. All groups were dosed via intradermal injection. Each dose was administered in four intradermal injections of 50 μL /injection for a total of 200 μL per animal. Injections occurred into previously marked sites near the midline of the dorsal lumbar area (upper left, upper right, lower left and lower right quadrants). Dose time (after the last injection) was recorded. Additional markings were made as needed to allow for identification of the dose site. Animals were administered with the RNAs on
- 10 day 1 and euthanized on day 3. Clinical observations were made on the rats twice daily. Food consumption and body weight were also monitored.

During the study, approximately 1 ml of blood samples was collected from the jugular vein for pharmacokinetic analysis as follows:

Time Point	PK ^a
Acclimation	-
Day 1	12 hours postdose
Day 2	24 hours postdose
Day 3	48 hours postdose

^aTime points for blood collection (n = 3 animals/Group/Time point)

PK = Pharmacokinetics

15

Results indicate that following administration of RNAs encoding FGF21, IL15, IL15 and IL15R, IL6, IL22, and NOVEPOEITIN, these proteins were readily detected in the blood with protein levels peaking at approximately 12 hours post injection (**FIG. 39**). Of note, the proteins tested in this study can be taken up by cells and tissues

20 and/or can exert an effect near the site of expression without appreciable accumulation in systemic circulation.

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain, using no more than routine experimentation, numerous equivalents to the specific embodiments described specifically herein. Such equivalents are intended to be encompassed in the scope of the following claims.

INCORPORATION BY REFERENCE

- 5 All patents and publications referenced herein are hereby incorporated by reference in their entireties.

CLAIMS

What is claimed is:

1. A method for expressing a protein of interest in a population of cells in a mammalian subject, comprising: administering a composition comprising an effective dose of a RNA encoding the protein of interest to the mammalian subject, the RNA containing a combination of non-canonical nucleotides that avoid substantial cellular toxicity, wherein:
 - the effective dose is about 100 ng to about 2000 ng and
 - the composition is administered intradermally.
2. The method of claim 1, wherein the protein of interest is an interleukin, optionally selected from IL2, IL15, IL16, and IL22, or a variant or mutant thereof.
3. The method of claim 1, wherein the protein of interest is selected from LIF, FGF1, FGF9, FGF15, FGF19, FGF21, FGF23, BDNF, SERPINB1, caspase-1, BMP2, and BMP6
4. The method of claim 1, wherein the non-canonical nucleotides have one or more substitutions at positions selected from the 2C, 4C, and 5C positions for a pyrimidine, or selected from the 6C, 7N and 8C positions for a purine.
5. The method of claim 4, wherein the non-canonical nucleotides comprise one or more of 5-hydroxycytidine, 5-methylcytidine, 5-hydroxymethylcytidine, 5-carboxycytidine, 5-formylcytidine, 5-methoxycytidine, pseudouridine, 5-hydroxyuridine, 5-methyluridine, 5-hydroxymethyluridine, 5-carboxyuridine, 5-formyluridine, 5-methoxyuridine, 5-hydroxypseudouridine, 5-methylpseudouridine, 5-hydroxymethylpseudouridine, 5-carboxypseudouridine, 5-formylpseudouridine, and 5-methoxypseudouridine, optionally at an amount of at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or 100% of the non-canonical nucleotides.
6. The method of any of the above claims, wherein at least about 50% of cytidine residues are non-canonical nucleotides, and which are selected from 5-hydroxycytidine, 5-methylcytidine, 5-hydroxymethylcytidine, 5-carboxycytidine, 5-formylcytidine, and 5-methoxycytidine.
7. The method of any of the above claims, wherein at least about 75% or at least about 90% of cytidine residues are non-canonical nucleotides, and the non-canonical nucleotides are selected from 5-hydroxycytidine, 5-methylcytidine, 5-hydroxymethylcytidine, 5-carboxycytidine, 5-formylcytidine, and 5-methoxycytidine.
8. The method of any of the above claims, wherein at least about 20% of uridine, or at least about 40%, or at least about 50%, or at least about 75%, or at about least 90% of uridine residues are non-canonical nucleotides, and the non-canonical are selected from pseudouridine, 5-hydroxyuridine, 5-methyluridine, 5-hydroxymethyluridine, 5-carboxyuridine, 5-formyluridine, 5-methoxyuridine, 5-hydroxypseudouridine,

5-methylpseudouridine, 5-hydroxymethylpseudouridine, 5-carboxypseudouridine, 5-formylpseudouridine, and 5-methoxypseudouridine.

9. The method of any of the above claims, wherein at least about 40%, or at least about 50%, or at least about 75%, or at about least 90% of uridine residues are non-canonical nucleotides, and the non-canonical nucleotides are selected from pseudouridine, 5-hydroxyuridine, 5-methyluridine, 5-hydroxymethyluridine, 5-carboxyuridine, 5-formyluridine, 5-methoxyuridine, 5-hydroxypseudouridine, 5-methylpseudouridine, 5-hydroxymethylpseudouridine, 5-carboxypseudouridine, 5-formylpseudouridine, and 5-methoxypseudouridine.
10. The method of any of the above claims, wherein at least about 10% of guanine residues are non-canonical nucleotides, and the non-canonical nucleotide is optionally 7-deazaguanosine.
11. The method of any of the above claims, wherein the RNA contains no more than about 50% 7-deazaguanosine in place of guanosine residues.
12. The method of any of the above claims, wherein the RNA does not contain non-canonical nucleotides in place of adenosine residues.
13. The method of any of the above claims, wherein the RNA comprises a 5' cap structure.
14. The method of any of the above claims, wherein the RNA 5'-UTR comprises a Kozak consensus sequence.
15. The method of any of the above claims, wherein the RNA 5'-UTR comprises a sequence that increases RNA stability in vivo, and the 5'-UTR may comprise an alpha-globin or beta-globin 5'-UTR.
16. The method of any of the above claims, wherein the RNA 3'-UTR comprises a sequence that increases RNA stability in vivo, and the 3'-UTR may comprise an alpha-globin or beta-globin 3'-UTR.
17. The method of any of the above claims, wherein the RNA comprises a 3' poly(A) tail.
18. The method of any of the above claims, wherein the RNA 3' poly(A) tail is from about 20 nucleotides to about 250 nucleotides in length.
19. The method of any of the above claims, wherein the RNA is from about 200 nucleotides to about 5000 nucleotides in length.
20. The method of any of the above claims, wherein the RNA is from about 500 to about 2000 nucleotides in length, or about 500 to about 1500 nucleotides in length, or about 500 to about 1000 nucleotides in length.
21. The method of any of the above claims, wherein the RNA is prepared by in vitro transcription.
22. The method of any of the above claims, wherein the composition is administered by subcutaneous injection, intradermal injection, subdermal injection, intramuscular injection, and topical administration.

23. The method of any of the above claims, wherein the composition is administered as one or more injections containing about 10 ng to about 2000 ng of RNA.
24. The method of any of the above claims, wherein the composition is administered as one or more injections each containing no more than about 10 ng, or no more than about 20 ng, or no more than about 50 ng, or no more than about 100 ng, or no more than about 200 ng, or no more than about 300 ng, or no more than about 400 ng, or no more than about 500 ng, or no more than about 600 ng, or no more than about 700 ng, or no more than about 800 ng, or no more than about 900 ng, or no more than about 1000 ng, or no more than about 1100 ng, or no more than about 1200 ng, or no more than about 1300 ng, or no more than about 1400 ng, or no more than about 1500 ng, or no more than about 1600 ng, or no more than about 1700 ng, or no more than about 1800 ng, or no more than about 1900 ng, or no more than about 2000 ng.
25. The method of any of the above claims, wherein the composition is administered as one or more injections each containing about 10 ng, or about 20 ng, or about 50 ng, or about 100 ng, or about 200 ng, or about 300 ng, or about 400 ng, or about 500 ng, or about 600 ng, or about 700 ng, or about 800 ng, or about 900 ng, or about 1000 ng, or about 1100 ng, or about 1200 ng, or about 1300 ng, or about 1400 ng, or about 1500 ng, or about 1600 ng, or about 1700 ng, or about 1800 ng, or about 1900 ng, or about 2000 ng.
26. The method of any of the above claims, wherein the composition contains one or more lipids to enhance uptake of RNA by cells.
27. The method of any of the above claims, wherein the composition is a cationic liposome formulation and the lipids are optionally selected from **Table 1**.
28. The method of any of the above claims, wherein the subject is a human.
29. The method of any of the above claims, wherein said population of cells comprises cells of the integumentary system.
30. The method of any of the above claims, wherein said population of cells comprises one or more of cells of the epidermis, cells of the basement membrane, cells of the dermis, and/or cells of the subcutis.
31. The method of any of the above claims, wherein said population of cells are cells of the epidermis, and comprise one more of cells of the stratum corneum, stratum lucidum, stratum granulosum, stratum spinosum, and/or stratum germinativum.
32. The method of any of the above claims, wherein said population of cells are cells of the dermis, and comprise one or more of cells from the papillary region and the reticular region.
33. The method of any of the above claims, wherein the composition is administered by intradermal injection to one or more of the dermis or epidermis.

34. The method of any of the above claims, wherein the composition containing from 10 ng to about 2000 ng is administered per treatment area of about 10 cm² or less, or about 5 cm² or less, or about 1 cm² or less, or about 0.5 cm² or less, or about 0.2 cm² or less.
- 5 35. The method of any of the above claims, wherein about 10 ng, or about 20 ng, or about 50 ng, or about 100 ng, or about 200 ng, or about 300 ng, or about 400 ng, or about 500 ng, or about 600 ng, or about 700 ng, or about 800 ng, or about 900 ng, or about 1000 ng, or about 1100 ng, or about 1200 ng, or about 1300 ng, or about 1400 ng, or about 1500 ng, or about 1600 ng, or about 1700 ng, or about 1800 ng, or about 1900 ng, or about 2000 ng of RNA is administered per treatment area of about 10 cm² or less, or about 5 cm² or less, or about 1 cm² or less, or about 0.5 cm² or less, or about 0.2 cm² or less.
- 10 36. The method of any of the above claims, wherein the composition is administered using an array of needles covering an affected area of the subject.
37. The method of any of the above claims, wherein the treatment area is mechanically massaged after administration.
- 15 38. The method of any of the above claims, wherein the treatment area is exposed to electric pulses after administration.
39. The method of claim 38, wherein the electric pulses are between about 10V and about 200V for from about 50 microseconds to about 1 second.
40. The method of claims 38 or 39, wherein the electric pulses are generated around the treatment area by a multielectrode array.
- 20 41. The method of any of the above claims, wherein the composition is administered about weekly, for at least 2 weeks.
42. The method of any of the above claims, wherein the composition is administered about every other week for at least one month.
- 25 43. The method of any of the above claims, wherein the composition is administered monthly or about every other month.
44. The method of any one of claims 41 to 43, wherein the composition is administered for at least two months, or at least 4 months, or at least 6 months, or at least 9 months, or at least one year.
45. The method of any of the above claims, wherein the subject has a disease, disorder, or condition of the integumentary system.
- 30 46. The method of claim 45, wherein the disease, disorder, or condition is selected from acne, rosacea, keratosis, urticaria, allergic dermatitis, atopic dermatitis, hair loss, hair growth, pigmentation disorder, burn, wound, corns, cysts, dry skin, dermatitis herpetiformis, eczema, ectodermal dysplasia, oily skin,

folliculitis, itch, psoriasis, hyperhidrosis, scars, wrinkles, uneven skin tone, and signs and symptoms of aging.

47. The method of claim 45, wherein the disease, disorder or condition is one or more of acne vulgaris, acne aestivalis, acne conglobata, acne cosmetic, acne fulminans, acne keloidalis nuchae, acne mechanica, acne medicamentosa, acne miliaris necrotica, acne necrotica, acne rosacea, actinic keratosis, acute urticaria, allergic contact dermatitis, alopecia areata, angioedema, athlete's foot, atopic dermatitis, autoeczemazation, baby acne, balding, blastomycosis, blackheads, birthmarks, skin pigmentation disorder, boil, bruise, bug bite, bug sting, burn, cellulitis, chiggers, chloracne, cholinergic or stress urticaria, chronic urticaria, cold type urticaria, confluent and reticulated papillomatosis, corns, cysts, dandruff, dermatitis herpetiformis, dermatographism, dyshidrotic eczema, diaper rash, dry skin, dyshidrosis, ectodermal dysplasia, hypohidrotic ectodermal dysplasia, X-linked hypohidrotic ectodermal dysplasia, eczema, epidermaodysplasia verruciformis, erythema nodosum, excoriated acne, exercise-induced anaphylaxis folliculitis, excess skin oil, folliculitis, freckles, frostbite, fungal nails, hair density, hair growth rate, halogen acne, hair loss, heat rash, hematoma, herpes simplex infections, hidradenitis suppurativa, hives, hyperhidrosis, hyperpigmentation, hypohidrotic ectodermal dysplasia, hypopigmentation, impetigo, ingrown hair, heat type urticaria, ingrown toenail, infantile acne or neonatal acne, itch, irritant contact dermatitis, jock itch, keloid, keratosis pilaris, lichen planus, lichen sclerosus, lupus miliaris disseminatus faciei, melasma, moles, molluscum contagiosum, nail growth rate, nail health, neurodermatitis, nummular eczema, occupational acne, oil acne, onychomycosis, physical urticaria, pilonidal cyst, pityriasis rosea, pityriasis versicolor, poison ivy, pomade acne, pseudofolliculitis barbae or acne keloidalis nuchae, psoriasis, psoriatic arthritis, pressure or delayed pressure urticaria, puncture wounds such as cuts and scrapes, rash, rare or water type urticaria, rhinoplasty, ringworm, rosacea, rothmund-thomson syndrome, sagging of the skin, scabis, scars, seborrhea, seborrheic dermatitis, shingles, skin cancer, skin tag, solar type urticaria, spider bite, stretch marks, sunburn, tar acne, tropical acne, thinning of skin, thrush, tinea versicolor, transient acantholytic dermatosis, tycoon's cap, uneven skin tone, varicose veins, venous eczema, vibratory angioedema, vitiligo, warts, Weber-Christian disease, wrinkles, x-linked hypohidrotic ectodermal dysplasia, xerotic eczema, yeast infection, and general signs of aging.
48. The method of any of the above claims wherein the disease, disorder or condition of the integumentary system is one or more of actinic keratosis, basal cell carcinoma, melanoma, Kaposi's sarcoma, and squamous cell carcinoma.
49. The method of any of the above claims, wherein the method alters, modifies and/or changes the appearance of a member of the integumentary system.
50. A patch delivery system, comprising a non-viral RNA transfection composition enclosed within a membrane, and an array of delivery needles delivering from 10 ng to about 2000 ng of RNA per

treatment area of about 100 cm² or less, or about 50 cm² or less, or about 20 cm² or less, or about 10 cm² or less, or about 5 cm² or less, or about 1 cm² or less, or about 0.5 cm² or less, or about 0.2 cm² or less.

51. The patch of claim 50, wherein the non-viral transfection composition contains from 10 ng to about 2000 ng per injection volume of about 20 µl to about 1 ml.

52. The patch of claim 50 or 51, wherein each needle delivers an injection volume of between 5 µl and 500 µl.

53. A kit suitable for use in human therapy, comprising:

(a) the synthetic RNA of any of the above claims in a unit dosage form of about 10 ng to about 2000 ng and

(b) an injection needle.

54. The kit of claim 53, wherein the unit dose form is about 10 ng, or about 20 ng, or about 50 ng, or about 100 ng, or about 200 ng, or about 300 ng, or about 400 ng, or about 500 ng, or about 600 ng, or about 700 ng, or about 800 ng, or about 900 ng, or about 1000 ng, or about 1100 ng, or about 1200 ng, or about 1300 ng, or about 1400 ng, or about 1500 ng, or about 1600 ng, or about 1700 ng, or about 1800 ng, or about 1900 ng, or about 2000 ng.

55. The kit of any of claims 53 or 54, wherein the kit further comprises instructions for use.

56. A pharmaceutical formulation comprising:

(a) the synthetic RNA of any of the above claims and

(b) the lipids of any of the above claims.

57. The pharmaceutical formulation of claim 56, wherein the pharmaceutical formulation is suitable for one or more of subcutaneous injection, intradermal injection, subdermal injection, intramuscular injection, and topical administration.

58. A pharmaceutical formulation comprising:

(a) the synthetic RNA of any of the above claims and

(b) a transfection reagent,

wherein the pharmaceutical formulation is provided in solid form.

59. A pharmaceutical formulation comprising synthetic RNA transfection reagent complexes, wherein the synthetic RNA transfection reagent complexes are provided in solid form.

60. The pharmaceutical formulation of claim 59, wherein the synthetic RNA transfection reagent complexes are provided in frozen form.

61. A method for stabilizing nucleic acid transfection reagent complexes comprising:
- (a) forming nucleic acid transfection reagent complexes
 - (b) contacting the nucleic acid transfection reagent complexes or vessel in which such are contained with a cryogenic liquid,
- 5 to produce stabilized nucleic acid transfection reagent complexes.
62. The method of claim 61, wherein the nucleic acid transfection reagent complexes are stabilized for shipment or storage.
63. A pharmaceutical formulation comprising the protein of SEQ ID NO: 167.
64. A pharmaceutical formulation comprising the nucleic acid of SEQ ID NO: 168.

10

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6



FIG. 7




FIG. 8



FIG. 9




FIG. 10




FIG. 11




FIG. 12






FIG. 13



A1, A2	EPO Standard, 100 mU/mL
B1, B2	EPO Standard, 50 mU/mL
C1, C2	EPO Standard, 25 mU/mL
D1, D2	EPO Standard, 12.5 mU/mL
E1, E2	EPO Standard, 6.25 mU/mL
F1, F2	EPO Standard, 3.12 mU/mL
G1, G2	EPO Standard, 1.56 mU/mL
H1, H2	EPO Standard, 0 mU/mL
A3, A4	EPO RNA (A, G, U, C), 24h
B3, B4	EPO RNA (A, G, 5 moU, C), 24h
C3, C4	EPO RNA (A, G, psU, C), 24h
D3, D4	GFP RNA, 24h
E3, E4	untransfected, 24h
F3, F4	1:100 EPO RNA (A, G, U, C), 24h
G3, G4	1:100 EPO RNA (A, G, 5 moU, C), 24h
H3, H4	1:100 EPO RNA (A, G, psU, C), 24h

FIG. 14



A1	EPO Standard, 100 mU/mL
B1	EPO Standard, 50 mU/mL
C1	EPO Standard, 25 mU/mL
D1	EPO Standard, 12.5 mU/mL
E1	EPO Standard, 6.25 mU/mL
F1	EPO Standard, 3.12 mU/mL
G1	EPO Standard, 1.56 mU/mL
H1	EPO Standard, 0 mU/mL
A2	1:100 EPO RNA (A, G, 5 moU, C), 24h
B2	1:500 EPO RNA (A, G, 5 moU, C), 24h
C2	1:2500 EPO RNA (A, G, 5 moU, C), 24h
D2	1:12500 EPO RNA (A, G, 5 moU, C), 24h
E2	1:62500 EPO RNA (A, G, 5 moU, C), 24h
F2	1:312500 EPO RNA (A, G, 5 moU, C), 24h
G2	1:1562500 EPO RNA (A, G, 5 moU, C), 24h
H2	1:7812500 EPO RNA (A, G, 5 moU, C), 24h

FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20

Animal #	Group	Day	Time Point	Sex	IL-6 (pg/mL)	TNFα (pg/mL)	IFNα (pg/mL)
5	1	D1	6H	M	ND	ND	ND
6	1	D1	6H	M	ND	ND	ND
13	2	D1	6H	M	ND	ND	ND
14	2	D1	6H	M	ND	ND	ND
31	3	D1	6H	M	ND	ND	ND
32	3	D1	6H	M	ND	ND	ND
49	4	D1	6H	M	ND	ND	ND
50	4	D1	6H	M	ND	ND	ND
15	2	D2	24H	M	ND	ND	ND
16	2	D2	24H	M	ND	ND	ND
33	3	D2	24H	M	ND	ND	ND
34	3	D2	24H	M	ND	ND	ND
52	4	D2	24H	M	ND	ND	ND
63	4	D2	24H	M	ND	ND	ND
23	2	D8		M	ND	ND	ND
24	2	D8		M	ND	ND	ND
41	3	D8		M	ND	ND	11.95
42	3	D8		M	ND	ND	9.13
59	4	D8		M	ND	ND	7.66
60	4	D8		M	ND	ND	12.51

ND = Not Detectable (<10 pg/mL for TNF-alpha; < 21 pg/mL for IL-6; <4 pg/mL for IFN-alpha)

FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27




FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32




FIG. 33




FIG. 34




FIG. 35




FIG. 36






FIG. 37

A.



B.



C.




FIG. 37 (continued)




FIG. 37 (continued)




FIG. 38




FIG. 39




FIG. 39

