



(43) International Publication Date
27 November 2014 (27.11.2014)

(10) International Publication Number
WO 2014/190273 A1

(51) International Patent Classification:

C07K 16/28 (2006.01) *G01N 33/53* (2006.01)
C07K 16/46 (2006.01) *C12N 5/07* (2010.01)
C12Q 1/04 (2006.01) *C12N 15/85* (2006.01)

(21) International Application Number:

PCT/US2014/039365

(22) International Filing Date:

23 May 2014 (23.05.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/827,312 24 May 2013 (24.05.2013) US

(71) Applicant: **BOARD OF REGENTS, THE UNIVERSITY OF TEXAS SYSTEM** [US/US]; 201 West 7th St., Austin, TX 78701 (US).

(72) Inventors: **COOPER, Laurence, J.**; 1515 Holcombe Blvd., Unit 907, Houston, TX 77030 (US). **JENA, Bipulendu**; 7900 Cambridge St., Houston, TX 77054 (US). **SINGH, Harjeet**; 9851 Meadowglen Lane, Houston, TX 77042 (US). **RUSHWORTH, David**; 9215 Portal Dr., Houston, TX 77031 (US).

(74) Agent: **BYRD, Marshall, P.**; Parker Highlander PLLC, 1120 So. Capital of Texas Highway, Building One, Suite 200, Austin, TX 78746 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: CHIMERIC ANTIGEN RECEPTOR-TARGETING MONOCLONAL ANTIBODIES

(57) Abstract: Provided are monoclonal antibodies that detect CD 19 CAR-modified immune cells and CAR-modified immune cells irrespective of the tumor associated antigen they target. Methods of using these functional monoclonal antibodies include, but are not limited to, detection, quantification, activation, and selective propagation of CAR-modified immune cells.



WO 2014/190273 A1

DESCRIPTION

CHIMERIC ANTIGEN RECEPTOR-TARGETING MONOCLONAL ANTIBODIES

5 **[0001]** This application claims the benefit of United States Provisional Patent Application No. 61/827,312, filed May 24, 2013, the entirety of which is incorporated herein by reference.

[0002] The invention was made with government support under Grant Nos. CA16672, CA124782, CA120956, CA141303, CA116127, and CA148600 awarded by the
10 National Institutes of Health. The government has certain rights in the invention.

INCORPORATION OF SEQUENCE LISTING

[0003] The sequence listing that is contained in the file named “UTFC.P1221WO_ST25.txt”, which is 23 KB (as measured in Microsoft Windows®) and was created on May 23, 2014, is filed herewith by electronic submission and is incorporated
15 by reference herein.

BACKGROUND OF THE INVENTION

1. Field of the Invention

[0004] The present invention relates generally to the fields of medicine, molecular biology, immunology, and gene therapy. More particularly, it concerns CD19
20 chimeric-antigen receptor (CAR)-targeting and universal CAR-targeting monoclonal antibodies and methods of using the same.

2. Description of Related Art

[0005] Genetically modified T cells engineered to express a tumor-specific chimeric antigen receptor (CAR) have been infused initially with modest (Till *et al.*, 2008; Pule *et al.*, 2008; Morgan *et al.*, 2010) and recently significant anti-tumor effects
25 (Kochenderfer *et al.*, 2010; Kalos *et al.*, 2011; Brentjens *et al.*, 2011; Kochenderfer *et al.*, 2012; O’Connor *et al.*, 2012). The prototypical CAR uses an extracellular domain to directly dock to a cell surface molecule, which is usually a tumor-associated antigen (TAA). The specificity of a CAR is typically derived from a scFv region assembled from the antigen-

binding region of a TAA-specific monoclonal antibody (mAb). The components of a second generation CD19-specific CAR currently in use in clinical trials, designated CD19RCD28, are shown in FIG. 1A. The scFv from mouse mAb clone FMC63 (Nicholson *et al.*, 1997) is fused in frame to an extracellular scaffold (*e.g.*, human immunoglobulin hinge and Fc regions or hinge and constant regions from human CD8 α) to promote oligomerization after binding TAA, which contributes to CAR-dependent activation via one or more signaling motifs in the endodomain.

[0006] Investigators measure the persistence of adoptively transferred T cells *in vivo* to assess the therapeutic potential of CAR⁺ T cells (Kalos *et al.*, 2011; June *et al.*, 2009). The most common approach to assessing survival of infused CAR⁺ T cells is quantitative PCR using CAR-specific primers (Kochenderfer *et al.*, 2010; Morgan *et al.*, 2006). However, this technique does not allow retrieval of genetically modified T cells from the recipient for multi-parameter analyses. What is needed, and is provided herein, is a mAb to specifically detect and isolate CD19-specific CAR⁺ genetically modified T cells. In addition to informing on persistence, a CAR-specific mAb can be used for in-process testing during manufacture to assess expression of the transgene, such as after electro-transfer of *Sleeping Beauty* (SB) or *piggyBac* DNA plasmids coding for CD19RCD28 (Singh *et al.*, 2008; Manuri *et al.*, 2009; Hackett *et al.*, 2010; Singh *et al.*, 2011).

[0007] Furthermore, two different reagents for the detection of CAR-modified T cells are known. Brentjens *et al.* (2013) describes the use of a biotinylated goat anti-mouse IgG (Fab')₂ (Jackson ImmunoResearch) for detection of CAR-modified T cells. As the sensitivity of this polyclonal antibody is low, in the setting of low lymphocyte numbers in patient samples, detection of CAR-modified T cells was accomplished only after non-specific expansion of T cells using Dynabeads. This prevents a direct assessment of circulating CAR-modified T cell *in vivo* after infusion. Zheng *et al.* (2012) describes the use of Protein L for detection of CAR-modified T cells by flow cytometry. This reagent has restricted use in terms of detection and sensitivity in a multi-parameter flow cytometry assay. Its use in other assay formats has not been shown. Importantly, both of these reagents are used for detection purposes only and their functionality has not been shown. Thus, there remains a need for reagents that specifically target CAR expressing cells for analysis and manipulation of the cell populations.

SUMMARY OF THE INVENTION

[0008] Embodiments of the invention provide monoclonal antibodies that specifically bind to a CAR polypeptide. For example, in some aspects, an antibody of the embodiments specifically binds to a constant domain of a CAR polypeptide. In alternative
5 aspects, the antibody specifically binds to the antigen recognition domain of a CAR polypeptide (*e.g.*, a CD-19-binding CAR polypeptide).

[0009] In a first embodiment an antibody is provided that binds specifically to a CD19-targeted CAR polypeptide (*e.g.*, the polypeptide of SEQ ID NO: 17). In some aspects, an isolated and/or recombinant antibody binds to a CD19-specific scFv CAR
10 polypeptide and competes for binding of the polypeptide with the 136.20.1 monoclonal antibody. In certain aspects, the antibody comprises all or part of the heavy chain variable region and/or the light chain variable region of the 136.20.1 monoclonal antibody. Thus, in some aspects, the antibody comprises an amino acid sequence that corresponds to a first, second, and/or third complementarity determining region (CDR) from the light variable
15 and/or heavy variable chain of the 136.20.1 monoclonal antibody of the present embodiment. In one aspect, the antibody may be the 136.20.1 antibody.

[0010] In certain aspects, the isolated antibody comprises CDR sequences at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the CDRs of the 136.20.1 monoclonal antibody heavy and light chain amino acid sequences (*e.g.*,
20 between 80% and 99% identical). In further aspects, an antibody comprises CDRs identical to the 136.20.1 CDRs, except for one, two or three amino acid substitutions, deletions, or insertions at one or more of the CDRs. For example, the antibody can comprise CDRs wherein the CDR sequences comprise 1 or 2 amino acid substitutions in the V_H CDR1, V_H CDR2, V_H CDR3, V_L CDR1, V_L CDR2 and/or V_L CDR3 relative to the CDRs of a 136.20.1
25 monoclonal antibody. Thus in some specific aspects, an antibody of the embodiment comprises a first V_H, a second V_H, a third V_H, a first V_L, a second V_L, and a third V_L CDR sequence at least 80% identical to the corresponding CDR sequence of monoclonal antibody 136.20.1, which are represented by SEQ ID NOs: 2, 3, 4, 6, 7, and 8, respectively. In some aspects, the isolated antibody comprises CDR sequences that are identical to the CDR
30 sequences of monoclonal antibody 136.20.1.

[0011] In another aspect, the isolated antibody comprises a V_H domain at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the V_H domain of 136.20.1 (SEQ ID NO: 1) and/or a V_L domain at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the V_L domain of 136.20.1 (SEQ ID NO: 5). In other aspects, the isolated antibody comprises a V_H domain between 80% and 99% identical to the V_H domain of 136.20.1 (SEQ ID NO: 1) or a V_L domain between 80% and 99% identical to the V_L domain of 136.20.1 (SEQ ID NO: 5). In a further aspect, the isolated antibody comprises V_H and V_L domains identical to those of monoclonal antibody 136.20.1.

[0012] In a further embodiment, there is provided a recombinant polypeptide comprising an antibody V_H domain comprising CDRs 1-3 of the V_H domain of 136.20.1 (SEQ ID NOs: 2, 3 and 4). In another embodiment, there is provided a recombinant polypeptide comprising an antibody V_L domain comprising CDRs 1-3 of the V_L domain of 136.20.1 (SEQ ID NOs: 6, 7 and 8). In still further embodiments, there is provided an isolated polynucleotide molecule comprising nucleic acid sequence encoding an antibody or a polypeptide comprising an antibody V_H or V_L domain disclosed herein.

[0013] In yet further embodiments, there is provided a method for selecting a cell comprising a specific CAR polypeptide comprising contacting the cell with an antibody specifically binds to the antigen recognition domain of the CAR polypeptide and selecting a cell comprising the specific CAR based on binding of the antibody. For example, the antibody may be contacted with the cell to inhibit the effector activity of the cell (*e.g.*, to block *in vitro* or *in vivo* activity). In further aspects, the antibody may be conjugated to a toxin for use in selectively inhibiting or killing a cell comprising a specific CAR polypeptide. In certain aspects, the specific CAR polypeptide can be a CD-19-binding CAR polypeptide. Thus, in some aspects, the antibody can be an antibody that binding the antigen recognition domain of a CD-19-binding CAR polypeptide, such as one of the specific antibodies disclosed herein.

[0014] In further embodiments, a method is provided for purifying or enriching cells comprising a specific CAR polypeptide. In some aspects, the antibody comprises a tag, such as a reporter, and the method comprises selecting cells comprising a specific CAR polypeptide (*e.g.*, by fluorescence-activated cell sorting (FACS) and/or paramagnetic beads). In further aspect, an antibody may be bound to a support. The support

may be, without limitation, a bead, a surface, a column, or an antigen presenting cell (APC) (*e.g.*, an aAPC or an activating and propagating cell (AAPC)). In some aspects, a method for purifying, enriching or stimulating proliferation of cells comprises use of cells that express a specific CAR, such as CD19-specific scFv CAR. In some aspects, a method of the
5 embodiments is further defined as a method for preventing or modifying the ability of a specific CAR (*e.g.*, a CD-19-binding CAR) to activate a T cell by contacting the cell with an antibody that binds to the antigen recognition domain of the CAR. In a further aspect, a method is further defined as a method for controlling toxicity of a cell (*e.g.*, controlling toxicity of a CAR T-cell therapy in a human subject).

10 **[0015]** Thus, in a further embodiment, there is provided a cell culture system comprising mammalian T-cells expressing a specific CAR and an antibody in accordance with the embodiments. In yet another embodiment, there is provided a kit comprising an antibody or cell of the embodiments and at least a first agent for increasing proliferation of mammalian T cells expressing a specific CAR (*e.g.*, a cytokine or inactivated aAPCs). In
15 still a further embodiment there is provided a recombinant cell expressing an antibody that specifically binds to the antigen recognition domain of a CAR. In some aspects, the antibody is a membrane-bound antibody. In still further aspects, the antibody is a scFv antibody. For example, in some specific embodiments cell (*e.g.*, an APC) comprises a membrane-bound antibody comprising one, two, three, four or more of: (i) a GM-CSF leader peptide (*e.g.*,
20 amino acids 1-22 of SEQ ID NO: 22); (ii) the scFv coding sequence for an antibody that specifically binds to the antigen recognition domain of a CAR (*e.g.*, a CD-19-binding CAR); (iii) a CD8 α linker sequence (*e.g.*, amino acids 136-182 of SEQ ID NO: 23); (iv) a CD28 transmembrane domain (*e.g.*, amino acids 56-123 of SEQ ID NO: 24); and/or (v) a CD3 ξ intracellular domain (*e.g.*, amino acids 48-163 of SEQ ID NO: 25).

25 **[0016]** In yet a further embodiment, the present invention provides an isolated or recombinant monoclonal antibody that specifically binds to the constant domain of a CAR polypeptide. For example, the antibody can compete for binding of the polypeptide with the 2D3 monoclonal antibody. In certain aspects, the antibody may comprise all or part of the heavy chain variable region and/or the light chain variable region of the 2D3 monoclonal
30 antibody. In a further aspect, the antibody may comprise an amino acid sequence that corresponds to a first, second, and/or third complementarity determining region (CDR) from the light variable and/or heavy variable chain of the 2D3 monoclonal antibody of the present

embodiment. In one aspect, the antibody may be the 2D3 antibody. In still further aspects, an antibody of the embodiments (*e.g.*, a 2D3 monoclonal antibody) binds to a polypeptide comprising the sequence of SEQ ID NO: 18.

[0017] Thus, in certain aspects, the isolated antibody comprises CDR sequences at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the CDRs of the 2D3 monoclonal antibody heavy and light chain amino acid sequences. In further aspects, an antibody comprises CDRs identical to the 2D3 CDRs, except for one, two or three amino acid substitutions, deletions, or insertions at one or more of the CDRs. For example, the antibody can comprise CDRs wherein the CDR sequences comprise 1 or 2 amino acid substitutions in the V_H CDR1, V_H CDR2, V_H CDR3, V_L CDR1, V_L CDR2 and/or V_L CDR3 relative to the CDRs of a 2D3 monoclonal antibody. Thus, in some specific aspects, an antibody of the embodiment comprises a first V_H, a second V_H, a third V_H, a first V_L, a second V_L, and a third V_L CDR sequence at least 80% identical to the corresponding CDR sequence of monoclonal antibody 2D3, which are represented by SEQ ID NOs: 10, 11, 12, 14, 15, and 16, respectively. In certain aspects, the antibody comprises CDR sequences that are identical to the CDR sequences of monoclonal antibody 2D3. In certain aspects, such an antibody is a humanized or de-immunized antibody comprising the foregoing CDRs on a human IgG (*e.g.*, IgG1 or IgG2) backbone.

[0018] In another aspect, the isolated antibody comprises a V_H domain at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the V_H domain of 2D3 (SEQ ID NO: 9) and a V_L domain at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the V_L domain of 2D3 (SEQ ID NO: 13). In certain aspects, the isolated antibody comprises a V_H domain between 80% and 99% identical to the V_H domain of 2D3 (SEQ ID NO: 9) or a V_L domain between 80% and 99% identical to the V_L domain of 2D3 (SEQ ID NO: 13). In some aspects, the isolated antibody comprises V_H and V_L domains identical to those of monoclonal antibody 2D3.

[0019] In a further embodiment, there is provided a recombinant polypeptide comprising an antibody V_H domain comprising CDRs 1-3 of the V_H domain of 2D3 (SEQ ID NOs: 10, 11 and 12) or a recombinant polypeptide comprising an antibody V_L domain comprising CDRs 1-3 of the V_L domain of 2D3 (SEQ ID NOs: 14, 15 and 16). In still further embodiments, there is provided an isolated polynucleotide molecule comprising nucleic acid

sequence encoding an antibody or a polypeptide comprising an antibody V_H or V_L domain disclosed herein.

[0020] In yet further embodiments, there is provided a method for selecting a cell comprising a CAR polypeptide comprising contacting the cell with an antibody (e.g.,
5 such as those disclosed herein) that binds to the constant domain of a CAR polypeptide and selecting a cell comprising the CAR based on binding of the antibody. In certain aspects, the antibody may be conjugated to a toxin for use in selectively inhibiting or killing a cell comprising a CAR. Thus, in some aspect, a method is provided for inhibiting or killing a CAR-expressing cell *in vivo*. In some aspects, a method of the embodiments is further
10 defined as a method for preventing or modifying the ability of a CAR to activate a T cell by contacting the cell with an antibody that binds to the constant domain of the CAR. In a further aspect, a method is further defined as a method for controlling toxicity of a cell (e.g., controlling toxicity of a CAR T-cell therapy in a human subject).

[0021] In still a further embodiment there is provided a recombinant cell
15 expressing an antibody that binds to the constant domain of a CAR. In some aspects, the antibody is a membrane-bound antibody. In still further aspects, the antibody is a scFv antibody. For example, in some specific embodiments cell (e.g., an APC) comprises a membrane-bound antibody comprising one, two, three, four or more of: (i) a GM-CSF leader peptide (e.g., amino acids 1-22 of SEQ ID NO: 22); (ii) the scFv coding sequence for an
20 antibody that binds to the constant domain of a CAR (e.g., a 2D3 scFv coding sequence); (iii) a CD8 α linker sequence (e.g., amino acids 136-182 of SEQ ID NO: 23); (iv) a CD28 transmembrane domain (e.g., amino acids 56-123 of SEQ ID NO: 24); and/or (v) a CD3 ζ intracellular domain (e.g., amino acids 48-163 of SEQ ID NO: 25).

[0022] In certain aspects, a method for purifying or enriching cells comprising
25 CAR using an antibody of the embodiments is provided. In one aspect, the antibody comprises a tag, such as a reporter, and the method comprises selecting cells comprising CAR (e.g., by FACS). In further aspects, the antibody may be bound to a support. The support may be, without limitation, a bead, a surface, a column, or an aAPC or AAPC. In some aspect, purifying or enriching cells comprising CAR comprises stimulating
30 proliferation of the cells. Thus, in a further embodiment, there is provided a cell culture system comprising mammalian T-cells expressing a CAR and an antibody in accordance with the embodiments. In a further embodiment, there is provided a kit comprising an antibody of

the embodiments and at least a first agent for increasing proliferation of mammalian T cells expressing a CAR. In one aspect, the kit may comprise inactivated aAPCs.

[0023] In still further aspects, CAR-binding antibody of the embodiments (e.g., a CAR constant domain-binding or CAR antigen recognition domain-binding antibody) is a humanized or de-immunized antibody comprising the foregoing CDRs on a human IgG (e.g., IgG1 or IgG2) backbone. In one aspect, the antibody may be recombinant. In further aspects, an antibody of the embodiments may be an IgG (e.g., IgG1, IgG2, IgG3 or IgG4), IgM, IgA, or an antigen binding fragment thereof. The antibody may be, for example, a Fab', a F(ab')₂, a F(ab')₃, a monovalent scFv, a bivalent scFv, or a single domain antibody. In some aspects, the antibody may be a human, humanized, or de-immunized antibody. In still further aspects, the antibody may be conjugated to an imaging agent, a chemotherapeutic agent, a toxin, a reporter, or a radionuclide.

[0024] In further embodiments, a host cell is provided that comprises one or more polynucleotide molecule(s) of the embodiments and/or that produces a monoclonal antibody or recombinant polypeptide of the embodiments. In some aspects, the host cell is a mammalian cell, a yeast cell, a bacterial cell, a ciliate cell, or an insect cell. In certain aspects, the host cell is a hybridoma cell. In certain aspects, the host cell is an artificial antigen presenting cell.

[0025] In still further embodiments, there is provided a method of manufacturing an antibody of the present invention comprising expressing one or more polynucleotide molecule(s) encoding a V_L or V_H chain of an antibody disclosed herein in a cell and purifying the antibody from the cell.

[0026] Certain embodiments of the invention are directed to an antibody or recombinant polypeptide composition comprising an isolated and/or recombinant antibody or polypeptide that specifically binds CAR or CD19-specific scFv CAR. In certain aspects the antibody or polypeptide has a sequence that is, is at least, or is at most 80, 85, 90, 95, 96, 97, 98, 99, or 100% identical (or any range derivable therein) to all or part of any monoclonal antibody provided herein. In still further aspects the isolated and/or recombinant antibody or polypeptide has, has at least, or has at most 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72,

73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 or more contiguous amino acids from any of the sequences provided herein or a combination of such sequences.

[0027] In still further aspects, an antibody or polypeptide of the embodiments
 5 comprises one or more amino acid segments of the any of the amino acid sequences disclosed herein. For example, the antibody or polypeptide can comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acid segments comprising about, at least or at most 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 to 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61,
 10 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162,
 15 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199 or 200 amino acids in length, including all values and ranges there between, that are at least 80, 85, 90, 95, 96, 97, 98, 99, or 100% identical to any of the amino acid sequences disclosed herein. In certain aspects the amino segment(s) are selected from one of the amino acid
 20 sequences of a CAR or CD19-specific scFv CAR binding antibody as provided in Tables 2 and 3.

[0028] In yet further aspects, an antibody or polypeptide of the embodiments comprises an amino acid segment that is at least 80, 85, 90, 95, 96, 97, 98, 99, or 100% identical (or any range derivable therein) to a V, VJ, VDJ, D, DJ, J or CDR domain of a CAR
 25 or CD19-specific scFv CAR antibody (as provided in Tables 2 and 3). For example, a polypeptide may comprise 1, 2 or 3 amino acid segment that are at least 80, 85, 90, 95, 96, 97, 98, 99, or 100% identical (or any range derivable therein) to CDRs 1, 2, and/or 3 a CAR or CD19-specific scFv CAR antibody as provided in Tables 2 and 3.

[0029] Embodiments discussed in the context of methods and/or compositions
 30 of the invention may be employed with respect to any other method or composition described herein. Thus, an embodiment pertaining to one method or composition may be applied to other methods and compositions of the invention as well.

[0030] As used herein the terms “encode” or “encoding” with reference to a nucleic acid are used to make the invention readily understandable by the skilled artisan; however, these terms may be used interchangeably with “comprise” or “comprising,” respectively.

5 [0031] As used herein the specification, “a” or “an” may mean one or more. As used herein in the claim(s), when used in conjunction with the word “comprising”, the words “a” or “an” may mean one or more than one.

[0032] The use of the term “or” in the claims is used to mean “and/or” unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive,
10 although the disclosure supports a definition that refers to only alternatives and “and/or.” As used herein “another” may mean at least a second or more.

[0033] Throughout this application, the term “about” is used to indicate that a value includes the inherent variation of error for the device, the method being employed to determine the value, or the variation that exists among the study subjects.

15 [0034] Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those
20 skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

[0035] The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the
25 detailed description of specific embodiments presented herein.

[0036] **FIG. 1.** Schematic of DNA plasmids used for electroporation and expression of transgenes. (A) Constructs showing components of CD19-specific CAR (CD19RCD28). The scFv is derived from mAb clone FMC63 that binds human CD19 and was generated by fusing the V_L and V_H regions via a "Whitlow" linker peptide. The scFv was

attached to modified human IgG₄ hinge and CH₂-CH₃ regions that was fused to the CD28 (transmembrane and cytoplasmic) and CD3 ζ (cytoplasmic) domains (Kowolik *et al.*, 2006). (B) DNA plasmid designed CD19scFv-mCD8 α used to express the transgene on L cells for immunization and screening. The CD19-specific scFv was generated as described, mCD8 α EC and TM represents mouse CD8 α extracellular and transmembrane domains. (C) Map of destination vector pIRESneo2 (Clontech) where the cDNA for the fusion protein CD19scFv-mCD8 α is cloned into *NheI* and *NotI* restriction enzyme sites. Expression cassette shows ColE1, colicin E1 (origin of replication); pCMV IE, human cytomegalovirus promoter/enhancer; ECMV IRES, encephalomyocarditis virus internal ribosome entry site which permits translation of two open reading frames from one messenger RNA, and poly A (polyadenylation) signal of the bovine growth hormone. Plasmid confers resistance through ampicillin and after electroporation the cells expressing the transgene of interest are selected on a cytotoxic concentration of drug neomycin sulfate (G418).

[0037] FIG. 2. Specificity of anti-CD19scFv mAb.

(A) Solid phase ELISA shows specificity of mAb (clone no. 136.20.1) as it binds to parental monoclonal antibody (FMC63) with background binding to other antibodies (*e.g.*, CD20-specific mAb or a different CD19-specific mAb). Purified human IgG served as a negative control. (B) Western blot shows clone 136.20.1 detects CAR protein in T cells genetically modified to express CD19RCD28. Lane A: Unmodified control T cells show endogenous CD3 ζ (14 kDa) as detected by commercial CD3 ζ -specific antibody and absence of CAR. Lane B: CAR⁺ T cells show CAR-specific band at 75 kDa as detected by mAb (clone 136.20.1). Lane C: CAR⁺ T cells show absence of CAR-specific band when the blot is treated with primary antibody after blocking (clone 136.20.1 was blocked with molar excess (1:5) of parental antibody FMC63). Lane D: CAR⁺ T cells show the presence of CAR (~75 kDa) and endogenous CD3 ζ (14 kDa) as detected by commercial CD3 ζ -specific antibody.

[0038] FIGS. 3A-C. Detection of CD19-specific CAR on the surface of

genetically modified cells. (A) CAR⁺ T cells were electroporated with SB system and propagated on aAPC. Upper row: Unmodified T cells as back ground control; Bottom row: CD19RCD28⁺ T cells (labeled as CAR⁺ T cells) detected by flow cytometry using clone no. 136.20.1, a commercially-available antibody (clone H10104, Invitrogen) that binds to the CAR hinge/Fc scaffold, and the inventor's own scaffold-specific mAb (clone 2D3). (B) Jurkat cells genetically modified to express CD19RCD28 were detected by clone no.

136.20.1 conjugated to Alexa-Fluor 488 and Alexa-Fluor 647 similar to commercial antibody (clone H10104). (C) Multi-parameter flow cytometry analysis of T cells that do and do not express CD19RCD28 as detected by clone no. 136.20.1. Sequential staining was performed, first with clone no. 136.20.1 and then other antibodies.

5 **[0039]** **FIG. 4.** Sensitivity of clone no. 136.20.1 to detect CD19-specific CAR⁺ T cells in PBMC. CD19-specific T cells expressing CD19RCD28 were serially diluted in PBMC (1:10 to 1:10,000). (A) The upper panel shows background level signal in PBMC control. (B) Middle panel shows staining of CAR⁺ T cells in undiluted sample. (C) Bottom panel shows lowest detection level (1:1,000) of CAR⁺ T cells in PBMC. Data acquisition by
10 FACSCaliber and plots were analyzed using FlowJo software. Shown in picture are representative flow cytometry data obtained from two independent experiments.

[0040] **FIG. 5.** Localization of CD19RCD28 CAR on the surface of genetically modified T cells. (A) Genetically-modified CAR⁺ T cells (expressing CD19RCD28) and unmodified control T cells were fixed using paraformaldehyde, stained
15 with AlexaFluor 647-conjugated clone no. 136.20.1 mAb and then spread on glass slides using cytospin. Confocal images were acquisitioned by Leica microscope (60X magnification). Upper panels (i and ii) show surface distribution of CAR molecules and bottom panels (iii and iv) show no staining in unmodified control T cells. Nuclear staining was by DAPI (pseudo-color green). The bar (1 μM) on images provides scale. (B) TEM
20 images showing staining of gold-labeled nanoparticles conjugated to clone no. 136.20.1 mAb. Upper row: Unmodified control T cells (i) 120 nm thin section; 15k magnification stained with uranyl acetate, (ii) 120 nm thin section; 75k magnification, and (iii) 120 nm thin section; 200k magnification. No gold particles were appreciated in these CAR^{neg} T cells. Bottom row: CAR⁺ T cells with enforced expression of CD19RCD28 (iv) 120 nm thin
25 section stained with uranyl acetate; 20k magnification, (v) 120 nm thin section; 200k magnification, and (vi) 80 nm thin section 200K magnification. Arrow heads on plates v and vi indicate surface distribution of gold-labeled nanoparticles attached to CAR molecules. The bar indicates scale.

[0041] **FIGS. 6A-C.** Clone no. 136.20.1 inhibits CAR⁺ T-cell effector
30 function. (A) Data from CRA shows inhibition of tumor cell lysis in CD19⁺ EL4 cells when CD19-specific CAR⁺T cells were blocked with clone no. 136.20.1 (B) Dose response curve showing clone no. 136.20.1-mediated inhibition of lysis of CD19⁺ EL4 target cells. (C)

Images obtained from VTLM showing CD19-specific CAR⁺ T cells co-cultured with CD19⁺ Daudiβ₂m expressing eGFP and stained with either clone no. 136.20.1 conjugated to AlexaFluor-647 or Fc mAb that recognizes CAR scaffold conjugated to PE. Images show two separate focal planes (at 2 and 90 minutes, respectively). Upper panels (i-iii) reveal coculture of Daudiβ₂m target cells with CAR⁺ T cells in the presence of Fc-specific antibody. Lower panels (iv-vi) reveal coculture of Daudiβ₂m target cells with CAR⁺ T cells in the presence of clone no. 136.20.1. Panels i-iv show phase contrast images of CAR⁺ T cells along with Daudiβ₂m. Panels ii-v show formation of immunological synapse (yellow arrow heads) when CD19-specific CAR⁺ T cells attack tumor cells. Panel iii shows killing of CD19⁺ tumor cell by CD19-specific CAR⁺ T cells as observed by formation of green fluorescence blub. Panel vi shows intact Daudiβ₂m cells when the CAR⁺ T cells were blocked by clone no. 136.20.1 and no killing was observed even when effector T cells engaged the tumor cells for more than 6 hours. Shown are data from 3 independent experiments.

[0042] **FIG. 7.** Schematic describes the steps for generating clone no. 136.20.1 mAb by L-cell immunization. DNA plasmid encoding the scFv immunogen was introduced into L cells by an electroporator device (Lonza). L cells stably expressed the transgenes under G418 drug selection. BALB/c mice were immunized in the foot pad and draining lymph nodes were harvested to collect lymphocytes. Candidate hybridoma clones were isolated and expanded for isolation of mAb that could detect CD19-specific CAR that employs scFv region derived from FMC63.

[0043] **FIG. 8.** Specificity of anti-CD19scFv mAb (clone no. 136.20.1) towards CD19-specific CAR⁺ T cells. A panel of CAR⁺ T cells were collected and stained with a commercial Fc-specific antibody (goat Fab₂ anti-human Fc gamma-PE) followed by staining with anti-CD19scFv mAb-Alexa 488. Shown are (A) Isotype control, (B) CD19RCD28 CAR⁺ T cells, (C) CD123-specific T cells (expressing CD123RCD28mZ-CAR), (D) CD33-specific T cells (expressing CD33RCD28z/Neo-CAR), (E) ROR1-specific T cells (expressing ROR1RCD137mz-CAR), and (F) HERV-K-specific T cells (expressing HERV-K-CD28z CAR). All cells were co-stained with Fc-specific antibody as well as clone no. 136.20.1. Only T cells expressing the CD19RCD28 CAR co-stained with both Fc-specific antibody and clone no. 136.20.1.

[0044] **FIG. 9.** Flow cytometry detection sensitivity of CD19-specific CAR⁺ T cells mixed with PBMC from healthy volunteers. Shown are plots obtained from entire

dilution range (1:10-1:10,000) of CAR⁺ T cells mixed with PBMC. Cells after mixing, were gated on live Aqua stain (Life Tech) lymphocytes (primary gate), followed by positive gating on CD3⁺, CD4⁺, CD8⁺ T cells (panels i, ii and iii, respectively) (secondary gate), and then identified as co-staining with Alexa-Fluor 647-conjugated clone no. 136.20.1 mAb (tertiary gate). On left panel are CAR⁺ T cells within specific lymphocyte population. Right panel shows back-gating within original lymphocyte population.

[0045] FIG. 10. Immunohistochemical staining of a panel of T cells (CAR modified and unmodified control) in formaldehyde fixed and paraffin embedded sections. (A) H&E staining of CAR⁺ T cells. (B) Absence of specific staining was observed in sections of CAR^{neg} control T cells stained with clone no. 136.20.1 mAb, HRP-conjugated detection antibody, and counter stained with hematoxyline. (C) CD19-specific CAR⁺ T cells expressing CD19RCD28 stained with clone no. 136.20.1 mAb and peroxidase-labeled secondary antibodies. Deposition of DAB on T cells indicating localization of CAR protein on the cell surface (arrow).

[0046] FIG. 11. Inhibition of specific lysis mediated by CD19-specific CAR⁺ T effector cells in a CRA. CD19-specific CAR⁺ T cells were incubated with clone no. 136.20.1 mAb at 250 µg/mL and then washed to remove unbound antibody. The effector cells were then co-cultured with ⁵¹Cr-labeled CD19⁺ tumor targets (NALM-6 and Daudiβ2m). Percentage specific lysis were calculated for effector cells bound by clone no. 136.20.1 and effector cell only at different Effector:Target (E:T) ratios.

[0047] FIG. 12. FACS plots showing detection CAR modified T cells by all CAR mAb (2D3-FITC). T cells were genetically modified to express a CD19-specific CAR that contains an IgG₄ Fc domain and expanded *ex vivo* on clone 4.0 (K562-artificial antigen presenting cells). Shown in picture are detection of CAR-modified T cells by 2D3 along with *ex vivo* expanded unmodified control T cells.

[0048] FIG. 13. Proof-of-principle showing all-CAR mAb (2D3) can impart proliferative signal upon CAR modified T cells through CD3-zeta *via* ITAM motif. A panel of CAR⁺ T cells targeting variety of tumor associated antigens viz., CD19, CD33, GD2 were made on a CAR backbone that contains an IgG₄ immunoglobulin constant region. scFv derived from 2D3 were expressed on artificial antigen presenting cells (K562), which were then irradiated and used in co-culture with PBMC-derived T cells. (A) Shown in the image

are *ex vivo* expansion of genetically-modified T cells over a period of 21 days. All CAR⁺ T cells, irrespective of antigen specificity, selectively proliferated in culture over time. Unmodified control T cells lacking IgG₄ Fc domain received no proliferative signal and died out in the same time. (B) Shown in image are total T cell counts, of CAR modified T cells
 5 over 21 day co-culture period as described.

[0049] FIG. 14. Study design to compare ability of chimeric antigen receptor (CAR) ligand (CARL) versus CD19 TAA on K562 cells for the selective propagation of CAR⁺ T cells (CART). A) Artificial antigen presenting cells (aAPC) demonstrated in I) were derived from parental K562 cells following transgene transfer, stable integration, and clonal
 10 selection. Each aAPC clone expresses either CARL, a scFv derived from 2D3 mAb that binds IgG₄ exodomain of CAR, or truncated human CD19. II) CART used to evaluate specificity towards CARL or CD19 are shown. SB-derived DNA plasmids coding for a panel of CARs were individually electro-transferred into PBMC and recursively stimulated with CD19⁺ K562 or CARL⁺ K562 in the presence of soluble recombinant human IL-2. Each CAR
 15 follows a modular design. 19G4CAR contains the IgG₄ scaffold and targets CD19 through the same scFv as 19CAR which lacks IgG₄ scaffold and instead uses CD8 α hinge and extracellular domain. GD2G4CAR contains the IgG₄ scaffold and targets GD2. G4CAR contains the IgG₄ scaffold, but has no scFv. All CARs employ of a 2nd generation design containing CD28 and CD3 ζ signaling endodomains. B) On Day 0, synchronous
 20 electroporation of PBMC was undertaken with DNA plasmid coding for SB transposase (SB11) and SB DNA plasmids coding for CAR species. To achieve outgrowth of T cells stably expressing CARs, the genetically modified cells were co-cultured, beginning on Day 1, upon γ -irradiated CD19⁺ or CARL⁺ K562 in the presence of 50 IU/mL IL-2. Cytokine was added with stimulation or during media change. Restimulation of CAR with aAPC occurred
 25 every 7 days until Day 21. C) Diagram of docking between CARL⁺ K562 cells and 19G4CAR⁺ T cells as compared with CD19⁺ K562 cells with 19G4CAR⁺ T cells.

[0050] FIG. 15. Characterization of aAPC and CAR⁺ T cells. A) CD19 and CARL as SB transposons were integrated into parental K562 cells using SB11 transposase and clonally expanded for homogeneous expression of CD19 or CARL. Dot plots depict the
 30 expression of CD19 and CARL on parental K562 and derived clones. The stable expression of CARL is shown using antibody that detects mouse Fab. B) The expression level of CAR species as determined by flow cytometry is shown on Days 1 and 21 of co-culture with

aAPC. Expression of chimeric IgG4 revealed CAR expression in all constructs except 19CAR which was determined using an antibody against human Fc. The percentage of cells in each flow plot quadrant is provided as an inset. C) The effect of aAPC design on abundance of CAR expression was assessed on Day 21 by measuring mean fluorescent intensity (MFI) of IgG4 signal by flow cytometry. The experiments are designated [CAR & aAPC] with unmodified mock electroporated T cells (No DNA plasmid) used as a control. Each experimental group contained 4 or 5 separate donor derived PBMC. Statistical comparison was undertaken by One-way ANOVA followed by unpaired t-tests between each experiment (* = $p < 0.05$).

[0051] FIG. 16. Comparison of CAR⁺ T cells propagated on CD19⁺ or CARL⁺ aAPC. A) Total inferred T-cell number and B) CAR (IgG4) expression for each CART was measured every 7 days for 5 donors. Top panel: 19CAR⁺ or G4CAR⁺ T cells were numerically expanded on either CD19⁺ or CARL⁺ aAPC. Bottom panel: 19G4CAR⁺ T cells were propagated on either CD19⁺ or CARL⁺ aAPC. C) After 21 days of co-culture on CD19⁺ or CARL⁺ aAPC, 19G4CAR⁺ T cells from 5 donors were assessed for expression of markers associated with memory (top panel) or T cell co-receptors (bottom panel). D) Specific killing by electroporated/propagated T cells expressing 19CAR, G4CAR, and 19G4CAR, by CRA at a ratio of 5 effectors to 1 target cell. The tumor targets were EL-4 (murine thymoma- GD2⁺, CD19^{neg}), NALM-6 (human B cell ALL- GD2^{neg}, CD19⁺), and K562 (a human CML- GD2^{neg}, CD19^{neg}). Up to 5 donors were tested in 4 independent experiments. ns- No significance, * = $p < 0.05$, *** = $p < 0.001$.

[0052] FIG. 17. Numeric expansion of CAR⁺ T cells using CARL⁺ aAPC. A) Total inferred T-cell number and B) CAR (IgG4) expression for each CART was measured every 7 days from 4 to 5 donors for 21 days of co-culture on aAPC. The differences between Day 21 total T-cell number and percent CAR expression was assessed using One-way ANOVA. C) The specific killing by panel of T cells expressing GD2G4CAR, 19G4CAR, and G4CAR, were tested using CRA at a ratio of 5 effectors to 1 target cell. The targets were EL-4 (GD2⁺, CD19^{neg}), NALM-6 (GD2^{neg}, CD19⁺), and parental K562 (GD2^{neg}, CD19^{neg}). Two-way ANOVA followed by unpaired t-tests was performed for 4 to 5 donors tested in 4 independent tests on Day 21 of co-culture on aAPC. ns- No significance, * = $p < 0.05$.

[0053] FIG. 18. Comparison of TCR repertoire changes induced by CAR-mediated expansion on aAPC. A) TCR repertoire was measured for 111 TCR α , β , γ , and δ

alleles using DTEA. 24 TCR abundance was organized from the most to the least frequently occurring transcripts based on sorted CD3⁺CD56^{neg} cells from Day 0. The set is visually represented next to TCR repertoire expressed by T cells at Day 21 of co-culture on CARL⁺ K562 cells and CD19⁺ K562 cells. Analysis was performed on 2 donors and a representative plot of one donor is shown.

[0054] FIG. 19. Vector maps for expression of transgenes. Each DNA plasmid expresses a transgene of interest under promoter human Elongation Factor 1 alpha (pHEF-1 α), using the beta hemoglobin poly-adenylation signal (BGH) to terminate transcription. The indirect repeats / direct repeats (IR/DR) allow for transgene transposition into the genome using SB11. All plasmids were propagated in bacteria using the origin of replication ColE1 and Kanamycin resistance (KanR) under the promoter pKan. A) 19G4CAR demonstrates the original plasmid design used in these studies and shows NheI and XmnI restriction enzyme (RE) sites used to generate B) GD2G4CAR from PCR-directed truncation of CD19- specific scFv on 19G4CAR which led to the generation of C) G4CAR and final ligation using ApaI and KpnI REs. D) 19CAR was designed without an IgG4 exodomain, instead expressing the CD8 α hinge and exodomain. E) Zeo-2ACARL, expressing CARL, and F) CD19-2A-Neo, expressing truncated human CD19 (tCD19), were designed to express CARL or CD19 on aAPC under drug selection conditions.

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

[0055] Therapeutics that employ CAR-expressing cells for targeting or specific antigens (*e.g.*, tumor-associated antigens) are currently being investigated for the treatment of a variety of disease from cancers to infectious disease. However, the complexity of these agents requires specialized tools for the purification, propagation and control of CAR-expressing cells, which were previously not available. Embodiments of the present invention provide antibodies that specifically bind to CAR T-cells and can be used to selectively purify, propagate or target CAR T-cells. For example, in one aspect, antibodies are provided that specifically bind to a region of the heavy chain constant domain shared by a wide array of CAR T-cells. For example, the 2D3 monoclonal antibody specifically binds to CAR expressing T-cell and can be used to analyze, isolate or propagate these cells. Likewise, specific anti-idiotypic (anti-Id) mAbs are provided with specificity for CD19-specific CARs. The mouse mAb (clone no. 136.20.1) recognizes CD19-specific CARs derived from anti-human CD19 mAb (clone FMC63). The specificity of this mAb for CD19-specific CAR id

demonstrated using (i) flow cytometry, (ii) western blotting, (iii) transmission electron microscopy (TEM), and (iv) immunocytochemistry. Furthermore, the binding of this anti-Id mAb to the scFv region of CD19RCD28 was demonstrated using a panel of CAR⁺ T cells and inhibition of CD19-dependent cytotoxicity as observed by chromium release assay (CRA) and video time lapse microscopy (VTLM).

[0056] Previously, CARs expressed on the T-cell surface were detected by antibodies directed against epitope tags or spacer domains incorporated in the extracellular domain. However, such tags would preferably not be employed in therapeutic CAR T-cells. Other commercially-available antibodies detecting Fc regions can, in some case, detect CARs, but cross-reactivity against immunoglobulin, especially when genetically modified T cells are to be assayed from patient-derived PBMC remains an obstacle. Likewise, alternative CAR designs that are non-immunogenic and avoid using a CH₂-CH₃ or employ scaffolding domains that are smaller than the Fc region, may improve the therapeutic potential of genetically modified T cells, but render Fc-binding antibodies useless as reagents. Thus, reagents that can directly bind to and detect the scFv region of the CAR, such as the mAbs provided here, represent the most effective reagents for manipulation of CAR T-cell populations. Indeed, the anti-idiotypic mAb developed for detection of CD19-specific CAR⁺ T cells provides a methodology to generate reagents to detect CARs targeting TAAs alternative to CD19.

[0057] Thus, the mAbs that binds to the scFv derived CAR T cells provided herein can serve as new and effective systems for T-cell manipulation. Antibodies specific for CAR T-cells (or in some cases CD-19 targeted CAR T-cells particularly) can be used to analyze and characterize CAR reagents. Perhaps even more importantly the CAR-binding agents provided herein can be used to specifically select, purify or propagate CAR T-cells. For example, antibodies of the embodiments can be used as part of a system for selectively growing CAR cells, such as by expressing the antibodies (or portions thereof) in aAPCs. In the case of antibodies that bind CARs in general, the antibodies can be used to generate lines of aAPCs as standardized reagents for expansion of CAR T-cell populations, a significant advantage over the systems previously available that must be individualized for particular CAR therapeutics. Likewise, the antibodies can be used to isolate and target CAR cells *in vivo* to monitor therapy and, if needed, target and inhibit CAR-expressing cells in a subject. These reagents will likely be instrumental in developing and implementing gene therapy trials

infusing CD-19 CAR⁺ T cells as well as CAR⁺ T cells in general (*i.e.*, CARs containing a generic human Fc gamma chain derived scaffold in its polypeptide chain).

I. Chimeric Antigen Receptors

[0058] The term “chimeric antigen receptors (CARs),” as used herein, may refer to artificial T-cell receptors, chimeric T-cell receptors, or chimeric immunoreceptors, for example, and encompass engineered receptors that graft an artificial specificity onto a particular immune effector cell. CARs may be employed to impart the specificity of a monoclonal antibody onto a T cell, thereby allowing a large number of specific T cells to be generated, for example, for use in adoptive cell therapy. In specific embodiments, CARs direct specificity of the cell to a tumor associated antigen, for example. In some embodiments, CARs comprise an intracellular activation domain, a transmembrane domain, and an extracellular domain comprising a tumor associated antigen binding region. In particular aspects, CARs comprise fusions of single-chain variable fragments (scFv) derived from monoclonal antibodies, fused to CD3 ζ a transmembrane domain and endodomain. The specificity of other CAR designs may be derived from ligands of receptors (*e.g.*, peptides) or from Dectins. In particular embodiments, one can target malignant B cells by redirecting the specificity of T cells by using a CAR specific for the B-lineage molecule, CD19. In certain cases, CARs comprise domains for additional co-stimulatory signaling, such as CD3 ζ , FcR, CD27, CD28, CD137, DAP10, and/or OX40. In some cases, molecules can be co-expressed with the CAR, including co-stimulatory molecules, reporter genes for imaging (*e.g.*, for positron emission tomography), gene products that conditionally ablate the T cells upon addition of a pro-drug, homing receptors, cytokines, and cytokine receptors.

[0059] As used herein, the term “antigen” is a molecule capable of being bound by an antibody or T-cell receptor. An antigen is additionally capable of inducing a humoral immune response and/or cellular immune response leading to the production of B and/or T lymphocytes.

[0060] The present invention involves antibodies that have specificity towards antigen-specific and universal chimeric antigen receptor (CAR) polypeptides, including a CAR that has been humanized to reduce immunogenicity (hCAR). In certain embodiments, the CAR may recognize an epitope comprised of the overlapping sequence between one or more antigens.

[0061] In certain aspects, the invention includes a method of making and/or expanding the antigen-specific redirected TCR^{neg} cells that comprises transfecting TCR^{neg} cells with an expression vector containing a DNA construct encoding the hCAR, then stimulating the cells with an antibody to the receptor to cause the cells to proliferate.

5 II. Antibodies of the Embodiments

[0062] In certain embodiments, an antibody or a fragment thereof that binds to at least a portion of CD19scFv-specific CAR protein and interferes with the killing of CD19⁺ target cells and its associated use in treatment of diseases is contemplated. In other embodiments, an antibody or a fragment thereof that binds to at least the CH₂-CH₃ hinge of
10 CAR protein (all-CAR) is provided. As used herein, the term “antibody” is intended to refer broadly to any immunologic binding agent, such as IgG, IgM, IgA, IgD, and IgE as well as polypeptides comprising antibody CDR domains that retain antigen binding activity. The antibody may be selected from the group consisting of a chimeric antibody, an affinity matured antibody, a polyclonal antibody, a monoclonal antibody, a humanized antibody, a
15 human antibody, or an antigen-binding antibody fragment or a natural or synthetic ligand. Preferably, the anti-CD19-specific CAR or all-CAR antibody is a monoclonal antibody or a humanized antibody. By known means and as described herein, polyclonal or monoclonal antibodies, antibody fragments, and binding domains and CDRs (including engineered forms of any of the foregoing) may be created that are specific to CD19-specific CAR or all-CAR
20 protein, one or more of its respective epitopes, or conjugates of any of the foregoing, whether such antigens or epitopes are isolated from natural sources or are synthetic derivatives or variants of the natural compounds.

[0063] Examples of antibody fragments suitable for the present embodiments include, without limitation: (i) the Fab fragment, consisting of V_L, V_H, C_L, and C_{H1} domains;
25 (ii) the “Fd” fragment consisting of the V_H and C_{H1} domains; (iii) the “Fv” fragment consisting of the V_L and V_H domains of a single antibody; (iv) the “dAb” fragment, which consists of a V_H domain; (v) isolated CDR regions; (vi) F(ab')₂ fragments, a bivalent fragment comprising two linked Fab fragments; (vii) single chain Fv molecules (“scFv”), wherein a V_H domain and a V_L domain are linked by a peptide linker that allows the two
30 domains to associate to form a binding domain; (viii) bi-specific single chain Fv dimers (see U.S. Pat. No. 5,091,513); and (ix) diabodies, multivalent or multispecific fragments constructed by gene fusion (US Patent App. Pub. 20050214860). Fv, scFv, or diabody

molecules may be stabilized by the incorporation of disulphide bridges linking the V_H and V_L domains. Minibodies comprising a scFv joined to a CH3 domain may also be made (Hu *et al.*, 1996).

5 **[0064]** Antibody-like binding peptidomimetics are also contemplated in embodiments. Liu *et al.* (2003) describe “antibody like binding peptidomimetics” (ABiPs), which are peptides that act as pared-down antibodies and have certain advantages of longer serum half-life as well as less cumbersome synthesis methods.

10 **[0065]** Animals may be inoculated with an antigen in order to produce antibodies specific for CD19scFv-specific CAR or all-CAR protein. Frequently an antigen is bound or conjugated to another molecule to enhance the immune response. As used herein, a conjugate is any peptide, polypeptide, protein, or non-proteinaceous substance bound to an antigen that is used to elicit an immune response in an animal. Antibodies produced in an animal in response to antigen inoculation comprise a variety of non-identical molecules (polyclonal antibodies) made from a variety of individual antibody producing B lymphocytes.

15 A polyclonal antibody is a mixed population of antibody species, each of which may recognize a different epitope on the same antigen. Given the correct conditions for polyclonal antibody production in an animal, most of the antibodies in the animal's serum will recognize the collective epitopes on the antigenic compound to which the animal has been immunized. This specificity is further enhanced by affinity purification to select only

20 those antibodies that recognize the antigen or epitope of interest.

25 **[0066]** A monoclonal antibody is a single species of antibody wherein every antibody molecule recognizes the same epitope because all antibody producing cells are derived from a single B-lymphocyte cell line. The methods for generating monoclonal antibodies (MAbs) generally begin along the same lines as those for preparing polyclonal antibodies. In some embodiments, rodents such as mice and rats are used in generating monoclonal antibodies. In some embodiments, rabbit, sheep, or frog cells are used in generating monoclonal antibodies. The use of rats is well known and may provide certain advantages. Mice (*e.g.*, BALB/c mice) are routinely used and generally give a high percentage of stable fusions.

30 **[0067]** Hybridoma technology involves the fusion of a single B lymphocyte from a mouse previously immunized with a CD19scFv-specific CAR or all-CAR antigen

with an immortal myeloma cell (usually mouse myeloma). This technology provides a method to propagate a single antibody-producing cell for an indefinite number of generations, such that unlimited quantities of structurally identical antibodies having the same antigen or epitope specificity (monoclonal antibodies) may be produced.

5 **[0068]** In one embodiment, the antibody is a chimeric antibody, for example, an antibody comprising antigen binding sequences from a non-human donor grafted to a heterologous non-human, human, or humanized sequence (*e.g.*, framework and/or constant domain sequences). Methods have been developed to replace light and heavy chain constant domains of the monoclonal antibody with analogous domains of human origin, leaving the
10 variable regions of the foreign antibody intact. Alternatively, “fully human” monoclonal antibodies are produced in mice transgenic for human immunoglobulin genes. Methods have also been developed to convert variable domains of monoclonal antibodies to more human form by recombinantly constructing antibody variable domains having both rodent, for example, mouse, and human amino acid sequences. In “humanized” monoclonal antibodies,
15 only the hypervariable CDR is derived from mouse monoclonal antibodies, and the framework and constant regions are derived from human amino acid sequences (see U.S. Pat. Nos. 5,091,513 and 6,881,557). It is thought that replacing amino acid sequences in the antibody that are characteristic of rodents with amino acid sequences found in the corresponding position of human antibodies will reduce the likelihood of adverse immune
20 reaction during therapeutic use. A hybridoma or other cell producing an antibody may also be subject to genetic mutation or other changes, which may or may not alter the binding specificity of antibodies produced by the hybridoma.

[0069] Methods for producing polyclonal antibodies in various animal species, as well as for producing monoclonal antibodies of various types, including
25 humanized, chimeric, and fully human, are well known in the art and highly predictable. For example, the following U.S. patents and patent applications provide enabling descriptions of such methods: U.S. Patent Application Nos. 2004/0126828 and 2002/0172677; and U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,196,265; 4,275,149; 4,277,437; 4,366,241; 4,469,797; 4,472,509; 4,606,855; 4,703,003; 4,742,159; 4,767,720; 4,816,567;
30 4,867,973; 4,938,948; 4,946,778; 5,021,236; 5,164,296; 5,196,066; 5,223,409; 5,403,484; 5,420,253; 5,565,332; 5,571,698; 5,627,052; 5,656,434; 5,770,376; 5,789,208; 5,821,337; 5,844,091; 5,858,657; 5,861,155; 5,871,907; 5,969,108; 6,054,297; 6,165,464; 6,365,157;

6,406,867; 6,709,659; 6,709,873; 6,753,407; 6,814,965; 6,849,259; 6,861,572; 6,875,434; and 6,891,024. All patents, patent application publications, and other publications cited herein and therein are hereby incorporated by reference in the present application.

[0070] Substitutional variants typically contain the exchange of one amino acid for another at one or more sites within the protein, and may be designed to modulate one or more properties of the polypeptide, with or without the loss of other functions or properties. Substitutions may be conservative, that is, one amino acid is replaced with one of similar shape and charge. Conservative substitutions are well known in the art and include, for example, the changes of: alanine to serine; arginine to lysine; asparagine to glutamine or histidine; aspartate to glutamate; cysteine to serine; glutamine to asparagine; glutamate to aspartate; glycine to proline; histidine to asparagine or glutamine; isoleucine to leucine or valine; leucine to valine or isoleucine; lysine to arginine; methionine to leucine or isoleucine; phenylalanine to tyrosine, leucine or methionine; serine to threonine; threonine to serine; tryptophan to tyrosine; tyrosine to tryptophan or phenylalanine; and valine to isoleucine or leucine. Alternatively, substitutions may be non-conservative such that a function or activity of the polypeptide is affected. Non-conservative changes typically involve substituting a residue with one that is chemically dissimilar, such as a polar or charged amino acid for a nonpolar or uncharged amino acid, and vice versa.

[0071] Proteins may be recombinant, or synthesized *in vitro*. Alternatively, a non-recombinant or recombinant protein may be isolated from bacteria. It is also contemplated that a bacteria containing such a variant may be implemented in compositions and methods. Consequently, a protein need not be isolated.

[0072] An antibody or preferably an immunological portion of an antibody, can be chemically conjugated to, or expressed as, a fusion protein with other proteins. For purposes of this specification and the accompanying claims, all such fused proteins are included in the definition of antibodies or an immunological portion of an antibody.

III. Chimeric Antigen Receptors

[0073] Embodiments of the present invention involve nucleic acids, including nucleic acids encoding an antigen-specific chimeric antigen receptor (CAR) polypeptide, including a CAR that has been humanized to reduce immunogenicity (hCAR), comprising an intracellular signaling domain, a transmembrane domain, and an extracellular domain

comprising one or more signaling motifs. In certain embodiments, the CAR may recognize an epitope comprised of the shared space between one or more antigens. Pattern recognition receptors, such as Dectin-1, may be used to derive specificity to a carbohydrate antigen. In certain embodiments, the binding region can comprise complementary determining regions of a monoclonal antibody, variable regions of a monoclonal antibody, and/or antigen binding fragments thereof. In another embodiment, that specificity is derived from a peptide (*e.g.*, cytokine) that binds to a receptor. A complementarity determining region (CDR) is a short amino acid sequence found in the variable domains of antigen receptor (*e.g.*, immunoglobulin and T-cell receptor) proteins that complements an antigen and therefore provides the receptor with its specificity for that particular antigen. Each polypeptide chain of an antigen receptor contains three CDRs (CDR1, CDR2, and CDR3). Since the antigen receptors are typically composed of two polypeptide chains, there are six CDRs for each antigen receptor that can come into contact with the antigen -- each heavy and light chain contains three CDRs. Because most sequence variation associated with immunoglobulins and T-cell receptors are found in the CDRs, these regions are sometimes referred to as hypervariable domains. Among these, CDR3 shows the greatest variability as it is encoded by a recombination of the VJ (VDJ in the case of heavy chain and TCR $\alpha\beta$ chain) regions.

[0074] It is contemplated that the human CAR nucleic acids are human genes to enhance cellular immunotherapy for human patients. In a specific embodiment, the invention includes a full length CAR cDNA or coding region. The antigen binding regions or domain can comprise a fragment of the V_H and V_L chains of a single-chain variable fragment (scFv) derived from a particular human monoclonal antibody, such as those described in U.S. Patent 7,109,304, incorporated herein by reference. The fragment can also be any number of different antigen binding domains of a human antigen-specific antibody. In a more specific embodiment, the fragment is an antigen-specific scFv encoded by a sequence that is optimized for human codon usage for expression in human cells.

[0075] The arrangement could be multimeric, such as a diabody or multimers. The multimers are most likely formed by cross pairing of the variable portion of the light and heavy chains into what has been referred to by Winters as a diabody. The hinge portion of the construct can have multiple alternatives from being totally deleted, to having the first cysteine maintained, to a proline rather than a serine substitution, to being truncated up to the first cysteine. The Fc portion can be deleted. Any protein that is stable and/or dimerizes can

serve this purpose. One could use just one of the Fc domains, *e.g.*, either the CH2 or CH3 domain from human immunoglobulin. One could also use the hinge, CH2 and CH3 region of a human immunoglobulin that has been modified to improve dimerization. One could also use just the hinge portion of an immunoglobulin. One could also use portions of CD8 α .

5 **[0076]** The intracellular signaling domain of the chimeric receptor of the invention is responsible for activation of at least one of the normal effector functions of the immune cell in which the chimeric receptor has been placed. The term “effector function” refers to a specialized function of a differentiated cell. Effector function of a T cell, for example, may be cytolytic activity or helper activity including the secretion of cytokines.

10 Effector function in a naive, memory, or memory-type T cell includes antigen-dependent proliferation. Thus the term “intracellular signaling domain” refers to the portion of a protein that transduces the effector function signal and directs the cell to perform a specialized function. While usually the entire intracellular signaling domain will be employed, in many cases it will not be necessary to use the entire intracellular polypeptide. To the extent that a

15 truncated portion of the intracellular signaling domain may find use, such truncated portion may be used in place of the intact chain as long as it still transduces the effector function signal. The term intracellular signaling domain is thus meant to include any truncated portion of the intracellular signaling domain sufficient to transduce the effector function signal. Examples include the zeta chain of the T-cell receptor or any of its homologs (*e.g.*, eta, delta,

20 gamma, or epsilon), MB1 chain, B29, Fc RIII, Fc RI, and combinations of signaling molecules, such as CD3 ζ and CD28, CD27, 4-1BB, DAP-10, OX40, and combinations thereof, as well as other similar molecules and fragments. Intracellular signaling portions of other members of the families of activating proteins can be used, such as Fc γ RIII and Fc ϵ RI. See Gross *et al.* (1992), Stancovski *et al.* (1993), Moritz *et al.* (1994), Hwu *et al.* (1995),

25 Weijtens *et al.* (1996), and Hekele *et al.* (1996) for disclosures of cTCR's using these alternative transmembrane and intracellular domains. In a preferred embodiment, the human CD3 ζ intracellular domain was taken for activation.

[0077] The antigen-specific extracellular domain and the intracellular signaling-domain may be linked by a transmembrane domain, such as the human IgG₄Fc

30 hinge and Fc regions. Alternatives include the human CD4 transmembrane domain, the human CD28 transmembrane domain, the transmembrane human CD3 ζ domain, or a cysteine

mutated human CD3 ζ domain, or other transmembrane domains from other human transmembrane signaling proteins, such as CD16 and CD8 and erythropoietin receptor.

[0078] In some embodiments, the CAR nucleic acid comprises a sequence encoding other costimulatory receptors, such as a transmembrane domain and a modified
5 CD28 intracellular signaling domain. Other costimulatory receptors include, but are not limited to one or more of CD28, CD27, OX-40 (CD134), DAP10, and 4-1BB (CD137). In addition to a primary signal initiated by CD3 ζ , an additional signal provided by a human costimulatory receptor inserted in a human CAR is important for full activation of T cells and could help improve *in vivo* persistence and the therapeutic success of the adoptive
10 immunotherapy.

[0079] In particular embodiments, the invention concerns isolated nucleic acid segments and expression cassettes incorporating DNA sequences that encode the CAR. Vectors of the present invention are designed, primarily, to deliver desired genes to immune cells, preferably T cells under the control of regulated eukaryotic promoters, for example,
15 MNDU3 promoter, CMV promoter, EF1alpha promoter, or Ubiquitin promoter. Also, the vectors may contain a selectable marker, if for no other reason, to facilitate their manipulation *in vitro*. In other embodiments, the CAR can be expressed from mRNA *in vitro* transcribed from a DNA template.

[0080] Chimeric antigen receptor molecules are recombinant and are
20 distinguished by their ability to both bind antigen and transduce activation signals *via* immunoreceptor activation motifs (ITAM's) present in their cytoplasmic tails. Receptor constructs utilizing an antigen-binding moiety (for example, generated from single chain antibodies (scFv)) afford the additional advantage of being "universal" in that they bind native antigen on the target cell surface in an HLA-independent fashion. For example,
25 several laboratories have reported on scFv constructs fused to sequences coding for the intracellular portion of the CD3 complex's zeta chain (ζ), the Fc receptor gamma chain, and sky tyrosine kinase (Eshhar *et al.*, 1993; Fitzer-Attas *et al.*, 1998). Re-directed T cell effector mechanisms including tumor recognition and lysis by CTL have been documented in several murine and human antigen-scFv: ζ systems (Eshhar, 1997; Altenschmidt *et al.*, 1997;
30 Brocker *et al.*, 1998).

[0081] To date non-human antigen binding regions are typically used in constructing a chimeric antigen receptor. A potential problem with using non-human antigen binding regions, such as murine monoclonal antibodies, is the lack of human effector functionality and inability to penetrate into tumor masses. In other words, such antibodies may be unable to mediate complement-dependent lysis or lyse human target cells through antibody-dependent cellular toxicity or Fc-receptor mediated phagocytosis to destroy cells expressing CAR. Furthermore, non-human monoclonal antibodies can be recognized by the human host as a foreign protein, and therefore, repeated injections of such foreign antibodies can lead to the induction of immune responses leading to harmful hypersensitivity reactions. For murine-based monoclonal antibodies, this is often referred to as a Human Anti-Mouse Antibody (HAMA) response. Therefore, the use of human antibodies is more preferred because they do not elicit as strong a HAMA response as murine antibodies. Similarly, the use of human sequences in the CAR can avoid immune-mediated recognition and therefore elimination by endogenous T cells that reside in the recipient and recognize processed antigen in the context of HLA.

[0082] In some embodiments, the chimeric antigen receptor comprises: a) an intracellular signaling domain, b) a transmembrane domain, and c) an extracellular domain comprising an antigen binding region.

[0083] In specific embodiments, intracellular receptor signaling domains in the CAR include those of the T cell antigen receptor complex, such as the zeta chain of CD3, also Fcγ RIII costimulatory signaling domains, CD28, CD27, DAP10, CD137, OX40, CD2, alone or in a series with CD3zeta, for example. In specific embodiments, the intracellular domain (which may be referred to as the cytoplasmic domain) comprises part or all of one or more of TCR zeta chain, CD28, CD27, OX40/CD134, 4-1BB/CD137, FcεRIγ, ICOS/CD278, IL-2Rβ/CD122, IL-2Rα/CD132, DAP10, DAP12, and CD40. In some embodiments, one employs any part of the endogenous T cell receptor complex in the intracellular domain. One or multiple cytoplasmic domains may be employed, as so-called third generation CARs have at least two or three signaling domains fused together for additive or synergistic effect, for example.

[0084] In certain embodiments of the chimeric antigen receptor, the antigen-specific portion of the receptor (which may be referred to as an extracellular domain comprising an antigen binding region) comprises a tumor associated antigen or a pathogen-

specific antigen binding domain including carbohydrate antigen recognized by pattern-recognition receptors, such as Dectin-1. A tumor associated antigen may be of any kind so long as it is expressed on the cell surface of tumor cells. Exemplary embodiments of tumor associated antigens include CD19, CD20, carcinoembryonic antigen, alphafetoprotein, CA-125, MUC-1, CD56, EGFR, c-Met, AKT, Her2, Her3, epithelial tumor antigen, melanoma-associated antigen, mutated p53, mutated ras, and so forth. In certain embodiments, the CAR can be co-expressed with a membrane-bound cytokine to improve persistence when there is a low amount of tumor-associated antigen. For example, CAR can be co-expressed with membrane-bound IL-15.

[0085] In certain embodiments intracellular tumor associated antigens may be targeted, such as HA-1, survivin, WT1, and p53. This can be achieved by a CAR expressed on a universal T cell that recognizes the processed peptide described from the intracellular tumor associated antigen in the context of HLA. In addition, the universal T cell may be genetically modified to express a T-cell receptor pairing that recognizes the intracellular processed tumor associated antigen in the context of HLA.

[0086] The pathogen may be of any kind, but in specific embodiments the pathogen is a fungus, bacteria, or virus, for example. Exemplary viral pathogens include those of the families of Adenoviridae, Epstein-Barr virus (EBV), Cytomegalovirus (CMV), Respiratory Syncytial Virus (RSV), JC virus, BK virus, HSV, HHV family of viruses, Picornaviridae, Herpesviridae, Hepadnaviridae, Flaviviridae, Retroviridae, Orthomyxoviridae, Paramyxoviridae, Papovaviridae, Polyomavirus, Rhabdoviridae, and Togaviridae. Exemplary pathogenic viruses cause smallpox, influenza, mumps, measles, chickenpox, ebola, and rubella. Exemplary pathogenic fungi include *Candida*, *Aspergillus*, *Cryptococcus*, *Histoplasma*, *Pneumocystis*, and *Stachybotrys*. Exemplary pathogenic bacteria include *Streptococcus*, *Pseudomonas*, *Shigella*, *Campylobacter*, *Staphylococcus*, *Helicobacter*, *E. coli*, *Rickettsia*, *Bacillus*, *Bordetella*, *Chlamydia*, *Spirochetes*, and *Salmonella*. In one embodiment the pathogen receptor Dectin-1 can be used to generate a CAR that recognizes the carbohydrate structure on the cell wall of fungi. T cells genetically modified to express the CAR based on the specificity of Dectin-1 can recognize *Aspergillus* and target hyphal growth. In another embodiment, CARs can be made based on an antibody recognizing viral determinants (e.g., the glycoproteins from CMV and Ebola) to interrupt viral infections and pathology.

[0087] In some embodiments, the pathogenic antigen is an *Aspergillus* carbohydrate antigen for which the extracellular domain in the CAR recognizes patterns of carbohydrates of the fungal cell wall, such as via Dectin-1.

[0088] It is contemplated that the chimeric construct can be introduced into
5 the subject's own T cells as naked DNA or in a suitable vector. Methods of stably transfecting T cells by electroporation using naked DNA are known in the art. See, e.g., U.S. Pat. No. 6,410,319. Naked DNA generally refers to the DNA encoding a chimeric receptor of the present invention contained in a plasmid expression vector in proper orientation for expression. Advantageously, the use of naked DNA reduces the time required to produce T
10 cells expressing the chimeric receptor of the present invention.

[0089] Alternatively, a viral vector (e.g., a retroviral vector, adenoviral vector, adeno-associated viral vector, or lentiviral vector) can be used to introduce the chimeric construct into T cells. Suitable vectors for use in accordance with the method of the present invention are non-replicating in the subject's T cells. A large number of vectors are known
15 that are based on viruses, where the copy number of the virus maintained in the cell is low enough to maintain the viability of the cell. Illustrative vectors include the pFB-neo vectors (STRATAGENE®) disclosed herein as well as vectors based on HIV, SV40, EBV, HSV, or BPV.

[0090] Once it is established that the transfected or transduced T cell is
20 capable of expressing the chimeric receptor as a surface membrane protein with the desired regulation and at a desired level, it can be determined whether the chimeric receptor is functional in the host cell to provide for the desired signal induction. Subsequently, the transduced T cells are reintroduced or administered to the subject to activate anti-tumor responses in the subject. To facilitate administration, the transduced T cells according to the
25 invention can be made into a pharmaceutical composition or made into an implant appropriate for administration *in vivo*, with appropriate carriers or diluents, which further can be pharmaceutically acceptable. The means of making such a composition or an implant have been described in the art (see, for instance, Remington's Pharmaceutical Sciences, 16th Ed., Mack, ed. (1980)). Where appropriate, the transduced T cells can be formulated into a
30 preparation in semisolid or liquid form, such as a capsule, solution, injection, inhalant, or aerosol, in the usual ways for their respective route of administration. Means known in the art can be utilized to prevent or minimize release and absorption of the composition until it

reaches the target tissue or organ, or to ensure timed-release of the composition. Desirably, however, a pharmaceutically acceptable form is employed that does not ineffectuate the cells expressing the chimeric receptor. Thus, desirably the transduced T cells can be made into a pharmaceutical composition containing a balanced salt solution, preferably Hanks' balanced salt solution, or normal saline.

IV. Artificial Antigen Presenting Cells

[0091] In some cases, aAPCs are useful in preparing CAR-based therapeutic compositions and cell therapy products. For general guidance regarding the preparation and use of antigen-presenting systems, see, *e.g.*, U.S. Pat. Nos. 6,225,042, 6,355,479, 6,362,001 and 6,790,662; U.S. Patent Application Publication Nos. 2009/0017000 and 2009/0004142; and International Publication No. WO2007/103009). As further detailed herein, in some aspects, aAPCs are provided, which comprise an antibody of the embodiments expressed on the surface. For instance, in some aspects, the antibody can be provided in addition to or instead of the antigen that is recognized by the CAR in order to mediate CAR T-cell expansion.

[0092] Thus, aAPCs, the cells can express an antibody and/or an antigen of interest. In addition to antibody or antigen of interest, the aAPC systems may also comprise at least one exogenous assisting molecule. Any suitable number and combination of assisting molecules may be employed. The assisting molecule may be selected from assisting molecules such as co-stimulatory molecules and adhesion molecules. Exemplary co-stimulatory molecules include CD70 and B7.1 (B7.1 was previously known as B7 and also known as CD80), which among other things, bind to CD28 and/or CTLA-4 molecules on the surface of T cells, thereby affecting, for example, T-cell expansion, Th1 differentiation, short-term T-cell survival, and cytokine secretion such as interleukin (IL)-2 (see Kim *et al.*, 2004, *Nature*, Vol. 22(4), pp. 403-410). Adhesion molecules may include carbohydrate-binding glycoproteins such as selectins, transmembrane binding glycoproteins such as integrins, calcium-dependent proteins such as cadherins, and single-pass transmembrane immunoglobulin (Ig) superfamily proteins, such as intercellular adhesion molecules (ICAMs), that promote, for example, cell-to-cell or cell-to-matrix contact. Exemplary adhesion molecules include LFA-3 and ICAMs, such as ICAM-1. Techniques, methods, and reagents useful for selection, cloning, preparation, and expression of exemplary assisting

molecules, including co-stimulatory molecules and adhesion molecules, are exemplified in, *e.g.*, U.S. Pat. Nos. 6,225,042, 6,355,479, and 6,362,001.

[0093] Cells selected to become aAPCs, preferably have deficiencies in intracellular antigen-processing, intracellular peptide trafficking, and/or intracellular MHC Class I or Class II molecule-peptide loading, or are poikilothermic (*i.e.*, less sensitive to temperature challenge than mammalian cell lines), or possess both deficiencies and poikilothermic properties. Preferably, cells selected to become aAPCs also lack the ability to express at least one endogenous counterpart (*e.g.*, endogenous MHC Class I or Class II molecule and/or endogenous assisting molecules as described above) to the exogenous MHC Class I or Class II molecule and assisting molecule components that are introduced into the cells. Furthermore, aAPCs preferably retain the deficiencies and poikilothermic properties that were possessed by the cells prior to their modification to generate the aAPCs. Exemplary aAPCs either constitute or are derived from a transporter associated with antigen processing (TAP)-deficient cell line, such as an insect cell line. An exemplary poikilothermic insect cells line is a *Drosophila* cell line, such as a Schneider 2 cell line (see, *e.g.* Schneider, J. Embryol. Exp. Morph. 1972 Vol 27, pp. 353-365). Illustrative methods for the preparation, growth, and culture of Schneider 2 cells, are provided in U.S. Pat. Nos. 6,225,042, 6,355,479, and 6,362,001.

[0094] In one embodiment, aAPCs are also subjected to a freeze-thaw cycle. In an exemplary freeze-thaw cycle, the aAPCs may be frozen by contacting a suitable receptacle containing the aAPCs with an appropriate amount of liquid nitrogen, solid carbon dioxide (*i.e.*, dry ice), or similar low-temperature material, such that freezing occurs rapidly. The frozen aAPCs are then thawed, either by removal of the aAPCs from the low-temperature material and exposure to ambient room temperature conditions, or by a facilitated thawing process in which a lukewarm water bath or warm hand is employed to facilitate a shorter thawing time. Additionally, aAPCs may be frozen and stored for an extended period of time prior to thawing. Frozen aAPCs may also be thawed and then lyophilized before further use. Preferably, preservatives that might detrimentally impact the freeze-thaw procedures, such as dimethyl sulfoxide (DMSO), polyethylene glycols (PEGs), and other preservatives, are absent from media containing aAPCs that undergo the freeze-thaw cycle, or are essentially removed, such as by transfer of aAPCs to media that is essentially devoid of such preservatives.

[0095] In other preferred embodiments, xenogenic nucleic acid and nucleic acid endogenous to the aAPCs, may be inactivated by crosslinking, so that essentially no cell growth, replication or expression of nucleic acid occurs after the inactivation. In one embodiment, aAPCs are inactivated at a point subsequent to the expression of exogenous MHC and assisting molecules, presentation of such molecules on the surface of the aAPCs, and loading of presented MHC molecules with selected peptide or peptides. Accordingly, such inactivated and selected peptide loaded aAPCs, while rendered essentially incapable of proliferating or replicating, retain selected peptide presentation function. Preferably, the crosslinking also yields aAPCs that are essentially free of contaminating microorganisms, such as bacteria and viruses, without substantially decreasing the antigen-presenting cell function of the aAPCs. Thus crosslinking maintains the important APC functions of aAPCs while helping to alleviate concerns about safety of a cell therapy product developed using the aAPCs. For methods related to crosslinking and aAPCs, see for example, U.S. Patent Application Publication No. 20090017000, which is incorporated herein by reference.

V. Kits of the Invention

[0096] Any of the compositions described herein may be comprised in a kit. In some embodiments, CAR-binding antibodies are provided in the kit, which also may include reagents suitable for expanding the cells, such as media, aAPCs, growth factors, antigens, other antibodies (*e.g.*, for sorting or characterizing CAR T-cells) and/or plasmids encoding CARs or transposase.

[0097] In a non-limiting example, a CAR-binding antibody, a chimeric receptor expression construct (or reagents to generate a chimeric receptor expression construct), reagents for transfection of the expression construct, and/or one or more instruments to obtain allogeneic cells for transfection of the expression construct (such an instrument may be a syringe, pipette, forceps, and/or any such medically approved apparatus) are provided in a kit. In some aspects, the kit comprises reagents or apparatuses for electroporation of cells.

[0098] The kits may comprise one or more suitably aliquoted compositions of the present invention or reagents to generate compositions of the invention. The components of the kits may be packaged either in aqueous media or in lyophilized form. The container means of the kits may include at least one vial, test tube, flask, bottle, syringe, or other container means, into which a component may be placed, and preferably, suitably aliquoted.

Where there is more than one component in the kit, the kit also will generally contain a second, third, or other additional container into which the additional components may be separately placed. However, various combinations of components may be comprised in a vial. The kits of the present invention also will typically include a means for containing the
5 chimeric receptor construct and any other reagent containers in close confinement for commercial sale. Such containers may include injection or blow molded plastic containers into which the desired vials are retained, for example.

VI. Examples

[0099] The following examples are included to demonstrate preferred
10 embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific
15 embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Example 1 – CD19 CAR-targeting monoclonal antibody - Methods

[00100] **Ethics Statement.** All experimental procedures pertaining to the use of animals were performed according to the U.T. MD Anderson Cancer Center Institutional
20 Animal Care and Use Committee (IACUC) guidelines and were carried out by the MD Anderson Hybridoma core lab (IACUC approval no. 11-071-2533). All efforts were made to reduce suffering to the animals caused by any experimental procedure. Isoflurane (2-5%) were used as anesthesia whenever required. Details of animal care, anesthesia use and sample collection are provided in separate paragraph. For the peripheral blood mononuclear cells
25 (PBMC) used in the study, blood samples were obtained from healthy volunteers under the protocol title “Acquisition of peripheral blood from healthy volunteers” that aimed to investigate the immunobiology of lymphocytes such as T cells in general and genetically modified T cells in particular (approval obtained from U.T. MD Anderson Cancer Center Institutional Review Board protocol no. LAB07-0296). Healthy volunteers donated blood
30 only after written consent was obtained and the study objectives were explained.

[00101] Cells. Cell lines were obtained from American Type Culture Collection (ATCC) unless otherwise stated. L cells, (a mouse adherent fibroblast cell line derived from C3H/An strain; ATCC no. CRL-2648), Jurkat cells (ATCC no. CRL-1990), NS0 cells (Sigma Aldrich no. 85110503), NALM-6 pre-B ALL cell (DSMZ no. ACC 128), EL4 murine lymphoma T cell line (ATCC no. TIB-39), CD19⁺ EL4 (genetically modified to express truncated human CD19) (Torikai *et al.*, 2012), and Daudi cells co-expressing EGFP and β_2 -microglobulin (Daudi β_2 m; Rabinovich *et al.* 2008), were all cultured in complete media (CM) defined as RPMI 1640 (Hyclone) supplemented with heat inactivated 10% fetal bovine serum (FBS) (Hyclone) and 2 mM L-Glutamine (Gibco-Invitrogen). Genetically modified cells were selected in various cytotoxic concentrations of neomycin sulfate G418 (Invivogen) at 0.8 mg/mL for NS0 cells, 0.9 mg/mL for L cells, and 1 mg/mL for Jurkat cells (Singh *et al.*, 2008). K562 were transduced with lentivirus to co-express CD64, CD86, CD137L and a membrane-bound IL-15 (mIL-15) to generate artificial antigen presenting cells (aAPC clone no. 4) (O'Connor *et al.*, 2012; Singh *et al.*, 2008; Manuri *et al.*, 2009; Singh *et al.*, 2011; Maiti *et al.*, 2013). Primary human T cells from PBMC were genetically modified to express CD19RC28 CAR using the *Sleeping Beauty* (SB) transposon/transposase system and propagated *ex vivo* in a CAR-dependent manner on CD19⁺ aAPC (K562, clone no. 4) in CM supplemented with soluble recombinant IL-2 (Novartis/Chiron) at 50 IU/mL and IL-21 at 30 ng/mL (Peprotech) (Singh *et al.*, 2011; Huls *et al.*, 2012).

[00102] Cell surface expression of CD19-specific scFv. V_L (amino terminus) and V_H (carboxyl terminus) regions were derived from anti-human CD19-specific mAb clone FMC63 (Nicholson *et al.*, 1997). The scFv was formed by joining the V_L and V_H with the 18 amino acid (AA) Whitlow peptide linker (GSTSGSGKPGSGEGSTKG) (Whitlow *et al.*, 1993). This binding domain, denoted as CD19scFv, was then fused in frame with mouse CD8 α extra-cellular domain (AA 28-196) and transmembrane domain (AA 197-217) (Swiss-Prot No.P01731 www.expasy.org) to create the fusion protein CD19scFvmCD8 α . The cDNA representing the fusion protein was mouse codon optimized and synthesized by a commercial vendor (FIG. 1B, GENEART). The nucleotide sequence of CD19scFvmCD8 α was verified and the cDNA was cloned into plasmid pIRESneo2 (Clontech cat no. 6938-1) to generate DNA vector CD19scFvmCD8 α pIRESneo2 (FIG. 1C). Purified vector CD19scFvmCD8 α pIRESneo2 DNA (endotoxin level <4 EU/mL) at a concentrations of 1 μ g per 10⁶ of L cells and 5 μ g per 10⁶ Jurkat and NS0 cells, were electro-transferred by a

Nucleofector device (model no. AAD-1001, Lonza) employing Nucleofector kit R and program X-005 for L cells and human T cell Nucleofector kit and program U-14 for Jurkat and NS0 cells. Electroporated cells were plated in 6-well culture plates (Corning) containing complete media (CM), incubated for 4 hours, and then added with drug neomycin sulfate (G418) as mentioned. Cells were cultured in an incubator (5% CO₂ and 37°C) and at 24 to 48 hours, the transient expression of CD19scFvmCD8α was assessed by flow cytometry using PE-conjugated anti-mouse CD8α antibody (BD Biosciences). Drug-resistant clones of L cells were propagated to confluence and harvested by trypsinization using 0.05% Trypsin-EDTA (Gibco-Invitrogen). Cells with highest expression level of CD19scFvmCD8α were sorted on a FACS Aria sorter (BD Biosciences) using anti-mouse CD8α-PE (BD Biosciences) as detection antibody. Sorted cells were numerically expanded further in the presence of neomycin sulfate (G418). Transgene expression was confirmed by flow cytometry. One L cell (clone no. 12) with the highest stable expression of transgene was chosen and expanded further for immunization purposes. Similarly, Jurkat and NS0 cells expressing CD19scFv were generated to screen hybridoma clones.

[00103] Animal care, immunization, lymphocyte collection. All the experiments pertaining to the animal use were performed as per the recommendations of the U.T. MD Anderson Cancer Center Institutional Animal Care and Use Committee (IACUC approval no. 11-071-2533). Two BALB/c mice (NCI, Frederick) were immunized in the footpads, each with 5x10⁶ L cells expressing CD19scFv-mCD8α (clone no. 12) suspended in 50 μL PBS. No adjuvant was used for immunization. The mice were restrained manually (for less than 2 minutes) for cell injection or anesthetized by isoflurane (2-5%) whenever required. Each mouse received a total of 6 injections at 3 day intervals. At the end of 5th immunization, mice were anesthetized by isoflurane (2-5%) inhalation and then blood was drawn from the tail vein. Sera were used to evaluate the antibody titer in an indirect ELISA. On Day 18 a mouse with highest titer was chosen for lymphocyte collection and fusion. Mice were euthanized by CO₂ inhalation and then popliteal lymph nodes were dissected for collection of lymphocytes.

[00104] Generation of hybridomas. A schematic outlining the steps involved for development and isolation of a mAb with specificity for the CD19scFv region in CD19RCD28 CAR is shown in FIG. 7. In brief, lymphocytes were washed with RPMI 1640 and then fused with a myeloma partner SP2/0 (ATCC no. CRL1581) at a lymphocyte to

myeloma ratio of 1:0.8. Hybridomas were obtained by standard polyethylene glycol (PEG-1450, Sigma-Aldrich) mediated fusion technology. Fused cells were grown in 96-well tissue-culture plates for 10 days in HAT selection media (Sigma-Aldrich) supplemented with 10% hybridoma cloning supplement (HCS) (GE/PAA Lab Inc.). Hybridoma clones were screened
 5 on day 10 to evaluate for the presence of antibody.

[00105] Screening hybridoma clones. A solid phase ELISA was used to determine the titer for the immune sera and to screen for hybridoma clones producing mAb specific for CD19scFv. Monoclonal antibodies were purified by standard protein G or protein A columns. One hybridoma clone (no. 136.20.1) was selected for characterization.

10 **[00106] Flow Cytometry.** For detecting surface expression of CAR, about 0.25×10^6 cells were suspended in 0.1mL of FACS buffer (2% FBS in 1X PBS) and stained with antibodies conjugated to fluorochromes (Table 1). Following blocking for 30 minutes at 4°C with 2% normal mouse sera (Jackson ImmunoResearch) in FACS buffer and washing
 15 twice with FACS buffer, the samples were incubated on ice with primary antibodies or conjugated antibodies for 30 minutes. Data were acquired on a FACSCalibur (BD Biosciences) using CellQuest software, version 3.3 (BD Biosciences). Dot plot analyses or MFI (median fluorescence intensity) was calculated by either FCS Express version 3.00.007 (Thornhill) or FlowJo software (version 7.6). To determine the sensitivity of anti-CD19scFv mAb (clone no. 136.20.1), CD19-specific CAR⁺ T cells were mixed with PBMC obtained
 20 from healthy donors (CAR^{neg} cells) at varying cell ratios (1:10-1:10,000 CAR⁺ T cell to PBMC). AlexaFluor-647 conjugated mAb (clone no. 136.20.1) was used to detect CAR⁺ T cells by flow cytometry.

Table 1: Antibodies used for flow cytometry analysis

Antibody	Amount (μL)	Cat no.	Vendor
Rat Anti-mouse CD8α	1	553033	Becton and Dickinson, San Jose, CA
Goat anti-mouse IgG Fcg fragment specific PE	1	115-096-071	Jackson ImmunoResearch, West Grove, PA
Goat F(Ab)2 anti-human IgGg R-PE	2.5	H10104	Invitrogen, Carlsbad, CA
Anti-CD19 ⁺ CAR Fc specific (2D3)	1	NA	Cooper Lab, MDACC
Anti-CD19 ⁺ CAR Idiotypic 136.20.1-AlexaFluor-	1	NA	Cooper Lab, MDACC

Antibody	Amount (μL)	Cat no.	Vendor
647AlexaFluor-488			
Anti-CD3 PE	1	555333	Becton and Dickinson, San Jose, CA
Anti-CD4 APC	2.5	555349	Becton and Dickinson, San Jose, CA
Anti-CD4 PercP Cy5.5	2	341654	Becton and Dickinson, San Jose, CA
Anti-CD8 PercP Cy5.5	2	341051	Becton and Dickinson, San Jose, CA
Anti-CD8 Pacific blue	1	558207	Becton and Dickinson, San Jose, CA

[00107] Binding assay. Binding activity of purified anti-CD19scFv mAb (clone no. 136.20.1) was determined by an indirect ELISA. In brief, wells of a 96-well plate (Thermo Scientific Nunc) were coated with 100 ng of one of the following mAbs: anti-human CD19 mAb (clone FMC63, Millipore), anti-human CD19 mAb (clone MHCD1900, Gibco-Invitrogen), anti-human CD20 mAb (clone no. 1F5.1D4.1 derived by us from clone 1F5, clone no. HB-9645, ATCC), or human IgG (Jackson ImmunoResearch), in 0.1 mL of coating buffer (100 mM NaHCO₃ pH 8.6). The plates were incubated at 37°C for 1 h and then washed three times with PBST (1X PBS with 0.05% tween-20). Blocking was achieved with 300 μL of 5% BSA in PBST. A dose curve was generated using different concentrations of purified anti-CD19scFv primary mAbs (2-30 μg/mL) and detected by secondary antibody goat anti-mouse IgG Fc HRP (Sigma-Aldrich). Absorbance was read at 450nm using a microplate reader (Victor 2030, Perkin Elmer).

[00108] Western blot. Reactivity of anti-CD19ScFv mAb (clone no. 136.20.1) to SDS-denatured protein was tested by western blot analysis. Ten million CAR modified or unmodified control T cells were lysed using RIPA buffer (50 mM Tris, pH 7.5, 150 mM NaCl, 0.1% SDS, 0.5% Sodium Deoxycholate, 1% TritonX100, 1 mM PMSF) containing protease inhibitor tablets as per the manufacturer's instructions (Roche Applied Science). Protein concentration was determined by a BCA kit (Thermo Scientific Pierce). Ten micrograms of total protein obtained from each cell lysates were electrophoresed under denaturing condition on a 4-20% gradient gel (Biorad) followed by electroblotting onto a PVDF membrane using a Criterion blotter (Biorad). PVDF membrane was blocked using 5% skim milk in PBST, and then incubated with primary anti-CD19ScFv mAb (clone no. 136.20.1). Binding was detected by goat anti-mouse IgG Fc HRP (Sigma-Aldrich) enhanced

with ECL Westfemto™ substrate (Thermo Scientific Pierce). Images of blots were acquired on a gel doc image system using versa doc Quantityone™ software (Biorad).

[00109] **Microscopy.** For confocal microscopy, genetically modified T cells expressing CD19RCD28 CAR were fixed in 4% paraformaldehyde (EMS Inc.) and stained with AlexaFluor 647 conjugated anti-CD19scFv mAb (clone 136.20.1) at 1:500 using a standard protocol. Labeled cells were added with ProLong gold anti-fade reagent with DAPI (Gibco-Invitrogen) and viewed under a confocal microscope (Leica Microsystems). For TEM, CAR⁺ T cells were fixed in 1% glutaraldehyde (Sigma-Aldrich) followed by washing with PBS. The cells were then incubated in 20 mM glycine-PBS followed by washing with PBS-BSA buffer (20 mM phosphate, 150 mM NaCl, pH 7.4 containing 0.5% BSA and 0.1% gelatin). NANOGOLD antibody conjugates were diluted 1:50 in PBS-BSA buffer and incubated with CD19-specific CAR⁺ T cells for 15 min followed by washing with the same buffer to remove unbound conjugates. Finally, NANOGOLD bound cells were fixed with 1% glutaraldehyde and stored at 4°C until further use. Before sectioning, cells were subjected to an alcohol dehydration steps (5% to 100% graded alcohol separated in 5 steps) followed by embedding in epoxy resin. The resin embedded cells were then cut into sections (1 μm to 100 nm thickness) in an ultra-microtome (Leica Microsystem). Sections were drawn onto a formavar coated copper grid. Staining was done by 2% uranyl acetate and then air dried. Unstained samples underwent silver enhancement (Nanoprobes) so that surface bound NANOGOLD particles could be visualized. High magnification images were acquired by a 1 k × 1 k CCD camera attached to a JEM-1200 EX60KV electron microscope (JEOL).

[00110] **Chromium Release Assay (CRA).** The functionality of anti-CD19scFv mAb was assessed in a CD19-specific CAR⁺ T cell-mediated cytotoxicity in a standard 4 h chromium (⁵¹Cr) release assay (CRA) employing parental CD19^{neg} EL4, CD19⁺ EL4, CD19⁺ Daudiβ₂m, and CD19⁺ NALM-6 as targets (Singh *et al.*, 2008; Singh *et al.*, 2011). Effector cells (*ex vivo* expanded CD19RCD28 CAR⁺ T cells harvested on day 28) were incubated with various concentrations of purified mAb (clone 136.20.1) at 10-fold serial dilution (maximum dose at 250 μg/mL and minimum at 1.25 μg/mL) for 20 min at 4°C, followed by washing with RPMI 1640 to remove unbound antibodies. The antibody-bound effector cells were incubated with ⁵¹Cr labeled target cells in a clear v-bottom 96-well plate (Corning) at 37°C. Release of ⁵¹Cr was quantified in a micro-plate scintillation counter TopCount NXT (Perkin-Elmer). Specific lysis was calculated for effector to target cell (E:T)

ratios in the presence and absence of anti-CD19scFv mAb. Data were reported as mean $\pm$ standard deviation (SD).

[00111] Video time lapse microscopy (VTLM). CD19-specific T cells expressing CD19RCD28 CAR were washed in sterile 1X PBS and stained for 30 min at room temperature using AlexaFluor 647-conjugated to anti-CD19ScFv mAb (clone no. 136.20.1) at 2 μ g mAb per 100,000 cells and then washed three times with 1X PBS. The CAR⁺ T cells were mixed together with EGFP⁺ Daudi β_2 m at an effector to target ratio of 2:1. A cell pellet was made by brief centrifugation and then re-suspended in media RPMI 1640 and put in one of the chamber of a 35 mm culture dish (Nikon). This was kept in the specimen chamber of Biostation IM (Nikon) and allowed to settle for 10 min. CCD camera focal planes were fixed and the effector to target cell dynamics were assessed for 2 to 12 hours using imaging software (IM-Q, v2.1.2.136, Nikon). As control, APC-conjugated goat anti-human IgG Fc γ chain specific F(ab')₂ fragment antibody (Jackson ImmunoResearch) was used to bind CAR⁺ T cells and the co-culture was set up as described.

[00112] Indirect ELISA to detect antibody titer and screen positive hybridoma clones. The assay was performed by coating flat-bottom well of a 96-well Medisorp plate (Thermo Scientific Nunc) with parental L cells or L cells transfected with CD19scFvmCD8 α . Approximately 250,000 cells were coated per well in 50 μ L of sodium phosphate buffer (pH 7.4). Plates were left overnight in laminar airflow hood for cell binding and drying and stored at -20°C until use. To block non-specific binding, 2% BSA in PBST (1X PBS containing 0.05% Tween-20) was dispensed at 300 μ L per well. Unbound cells were washed with PBST. Various dilutions of sera (1:100 to 1:3,000) or antibody supernatant (1:2-1:10) were allowed to bind with the coated cells/antigen for 1 h at room temperature (RT). After washing with PBST, secondary antibody anti-mouse IgG Fc HRP (Sigma-Aldrich) was added at 1:5,000 dilution and incubated for 45 min at room temperature. Detection of the bound antibody was performed after addition of 1X TMB/H₂O₂ substrate (Sigma-Aldrich). Absorbance was measured at 450 nm using a microplate reader (Victor 2030, Perkin Elmer). Cross reaction of immune sera and hybridoma supernatant with parental L cells was measured at OD₄₅₀ and used as background controls. For transfected cells, wells showing OD₄₅₀ values 3 times above background controls were marked positive. Random wells were viewed by microscopy before and after washing to ensure no loss of cells occurred during the process. The isotype of mAbs were determined using a mouse

immunoglobulin isotyping kit (BD Biosciences). Positive clones were identified and expanded further in 1X HT (Sigma-Aldrich) supplement media. Sub-cloning was performed by standard limiting dilution method to obtain a monoclonal population. Monoclonal antibodies were purified using protein G or protein A columns (GE Life Science).

5 **[00113] Antibody purification and conjugation.** One CD19scFv specific clone (no. 136.20.1) was grown and expanded further *in vitro* or injected in mice for ascites production (Harlan Inc.). Monoclonal antibodies from the culture supernatant or ascites were obtained by standard protein G or A (GE Life Science) purification using 0.1 M glycine (pH 2.7) as elution buffer. Affinity eluted and neutralized antibodies were dialyzed extensively
10 against PBS and subjected to flow cytometry analysis for final validation and use in cell culture assays. Purified antibodies were conjugated to different fluorochromes *viz.*, Alexa Fluor 488 and Alexa Fluor 647 (Molecular Probes-Invitrogen) using a commercially available kit that exploits the principle of conjugation through succinimidyl ester moiety to primary amines of proteins/ immunoglobulins to form stable dye-protein conjugates.
15 Fluorochrome-conjugated antibodies were subjected to gel filtration chromatography to remove unbound dye. For transmission electron microscope (TEM) studies, antibodies were conjugated to gold nanoparticles. In brief, antibodies were reduced by adding 1 mL of purified mAb (2 mg/mL) to 6 mg of mercaptoethylamine hydrochloride (MEA) (Thermoscientific-Pierce) and then purified on a gel filtration column MatrexGH25
20 (Millipore, Billerica, MA). Reduced antibodies were labeled with monomaleimido NanogoldTM (Nanoprobes), as per the manufacturer's instructions, and purified on a gel filtration column Superdex-75 (Sigma-Aldrich) to remove unbound NanogoldTM. The final concentration of the purified antibodies was measured by a BCA kit (Thermoscientific Pierce).

25 **[00114] Immunohistochemistry.** CD19-specific CAR⁺ T cells were propagated as described. Cells were harvested, washed in 1X PBS and then fixed in formalin. Five micron paraffin-embedded sections were made and stained. In brief, sections were blocked initially with 3% hydrogen peroxide and whole goat sera blocking reagent for 5 minutes, followed by incubation with primary antibody anti-Id CAR idotype for 30 minutes.
30 Biotinylated goat anti-mouse IgG-HRP was used as secondary antibody at 1:500 for 30 minutes, followed by incubation with streptavidin-HRP for 15 minutes. Cells were localized with DAB (diaminobenzidine) for 5 minutes. Slides were counterstained with Mayer's

hematoxylin for 5 minutes, dehydrated and mounted with a cover slip. Photographs were taken using a Leica microscope.

[00115] Methodology to derive antibody variable region sequence. The antibody variable heavy chain (VH) and light chain sequences (VL) were derived after amplifying the VH and VL using sets of degenerate primers annealed to either variable heavy or light chain constant regions of monoclonal antibody (clone 136.20.1) cDNA. The PCR-amplified product was cloned into PCR4 TOPO vector (Invitrogen) and then sequenced. Consensus sequences were identified after aligning sequences obtained with T3 forward and T7 reverse primed pCR4 TOPO TA vector inserted with 136.20.1 VH or VL PCR product. Sequence alignment was performed in Vector NTI 11 (Invitrogen). CDR sequences of the monoclonal antibody were analyzed from the cDNA amplified sequences using IMGT software (on the world wide web at imgt.org/IMGT_vquest/vquest?livret=0&Option=mouseIg).

Example 2 – CD19 CAR-targeting monoclonal antibody - Results

[00116] Generating anti-ScFv mAb recognizing CD19RCD28 CAR. BALB/c mice were immunized using L cells (derived from C3H/An mouse strain) genetically modified to express the scFv region in a CD19-specific CAR (FIG. 7). The scFv was derived from FMC63 (encoded in DNA plasmid CD19scFv-mCD8 α) and stably expressed (>90% homogenous expression after single-cell sorting) on the surface of neomycin-resistant genetically modified L cells, which was used to immunize allogeneic mice. The murine CD8 α trans-membrane and extra-cellular region were used to display scFv on the surface and to detect expression (FIGS. 1B and 1C). Genetically modified L cells were injected in mice at 3 day intervals and high titer sera (OD values ~5 times above background at 1:2,700 dilution) was obtained after five successive immunizations. Hybridomas were generated and ELISA was used to initially select 23 “high binders” (OD₄₅₀>1.5) and 15 “moderate binders” (OD₄₅₀~1.00-1.5) to advance to the second round of screening. Flow cytometry using Jurkat and NS0 cells genetically modified with CD19scFv-mCD8 α was used to cull the number of candidate hybridomas to 12. One clone was selected (no. 136.20.1; Table 2) that could efficiently bind CD19RCD28 expressing cells by flow cytometry. The absence of adjuvant at the time of immunization did not affect the antibody affinity maturation process as the obtained mAb clones were predominantly of IgG_{2a} sub-class. Immunization by genetically

modified L cells provides an efficient method for eliciting desirable anti-idiotypic immune response.

Table 2: Sequence of CD19scFv-specific mAb clone 136.20.1

mAb	Heavy Chain Sequence			Light Chain Sequence		
	CDR1	CDR2	CDR3	CDR1	CDR2	CDR3
	Amino acid sequence			Amino acid sequence		
136.20.1	GFDFSRYW (SEQ ID NO: 2)	INLDSSTI (SEQ ID NO: 3)	ARRYDAMD Y (SEQ ID NO: 4)	ESVDDYGI SF (SEQ ID NO: 6)	AAP (SEQ ID NO: 7)	QQSKD (SEQ ID NO: 8)
	LKPREVKLVESGGGLVQPGGSLKLSCA ASGFDFSRYWMSWVRQAPGKGLEWIGE INLDSSTINYTPSLKDKFIIISRDNAKN TLYLQMSKVRSEDTALYYCARRYDAMD YWGQGTSVTVSSAKTTAPSVYPLAPVC GDTTGSSVTLGCLVKASQ (SEQ ID NO: 1)			ASDIVLTQSPASLAVSLGQRATIS CRASESVDDYGISFMNWFQQKPGQ PPKLLIYAAPNQGSQVAPRFSQSG SGTDFSLNIHPMEEDDTAMYFCQQ SKDVRWRHQAGDQTG (SEQ ID NO: 5)		

5 **[00117] Specificity of mAb (clone no. 136.20.1) for CD19scFv.** The specificity of mAb (clone no. 136.20.1) towards CD19scFv was initially evaluated by an indirect ELISA with wells coated with purified mAbs. Except for parental anti-human CD19 mAb (FMC63) all other antibodies exhibited negligible cross reactivity to CD19scFv specific mAb (FIG. 2A). Anti-CD19scFv binds to its parental mAb at concentrations as low as 2

10 μg/mL with binding saturated at 4 μg/mL. The specificity of this mAb towards clone FMC63 was confirmed as it did not cross react to another anti-human CD19 antibody (Invitrogen, clone MHCD 1900). Further, it did not bind to purified human IgG and CD20-specific mAb. The ability of mAb (clone no. 136.20.1) to detect CD19⁺ CAR protein was verified by western blot using whole protein lysates obtained from *ex vivo*-propagated CD19-specific

15 CAR⁺ T cells. MAb (clone no. 136.20.1) detected a ~75 kDa protein in CD19RCD28 modified cell lysates run in reducing SDS-PAGE and electroblotted onto PVDF membrane (FIG. 2B, lane B). The specificity was verified in parallel by a commercially available CD3ζ antibody which detects the CD3ζ signaling domain in CD19RCD28-expressing CAR (FIG. 2B, lane D) and endogenous CD3ζ in unmodified control T cells (FIG. 2B, lane A). The

20 ability of this mAb to detect the surface expression of CD19⁺ CAR was verified by flow

cytometry analysis using CD19RCD28-expressing primary T cells and Jurkat cells (FIGS. 3A and 3B, respectively). MAb (clone 136.20.1) detected expression of CD19scFv on the surface of genetically modified T cells (85%) as well as in Jurkat cells (99%) which matches with detection level of anti-Fc antibodies that binds to the CH₂-CH₃ stalk of the CAR. Indeed, the staining pattern was compared with commercial IgG- γ chain-specific antibodies (Fc-PE Invitrogen) that bind to the CH₂-CH₃ region of the CAR and by mAb clone no. 2D3 developed in-house which binds to the CH₂-CH₃ hinge region of the CAR (Fc-FITC) (Singh *et al.*, 2008). A panel of CAR⁺ T cells that target different TAAs (CD33, CD123, ROR1, HERV-K) were used to verify if clone no. 136.20.1 cross reacted with any of these CAR transgenes. Each CAR construct was expressed from the *Sleeping Beauty* vector backbone and contained different signaling endodomains in addition to the common IgG₄-derived hinge/CH₂-CH₃ linker. It did not cross react with any of other CARs tested other than those with specificity for CD19 (FIG. 8). No background binding to mock-electroporated control T cells was observed. Additional evidence supporting specificity of mAb (clone no. 136.20.1) are provided (FIGS. 9 and 10). In both cases CAR scaffold differs either by the presence of human CD4 transmembrane region or IgG₁ Fc stalk that connects the scFv with the signaling endodomains. Furthermore, the mAb has been used to detect CD19-specific CAR⁺ T cells that employ an extracellular domain derived from CD8 α (Kalos *et al.*, 2011; Porter *et al.*, 2011). These results demonstrate that, irrespective of extra-cellular scaffolding and transmembrane domains, the mAb (clone no. 136.20.1) detects CD19scFv within CAR expressing scFv derived from FMC63.

[00118] The anti-CD19scFv mAb (clone no. 136.20.1) could be used in multi-parameter flow cytometry analysis to characterize CAR⁺ cells selectively propagated on K562-derived aAPC (Jena *et al.* 2013). CAR-modified T cells were numerically expanded in the presence of irradiated aAPC and cytokines (IL-2 and IL-21) for 4 weeks as previously reported (Singh *et al.*, 2011). The anti-CD19scFv mAb (clone 136.20.1) could detect both CD4⁺ and CD8⁺ CAR⁺ T cells (FIG. 3C). The inventors assessed the limit of detection using a spiking experiment. As shown in FIG. 4, the inventors were able to detect one CD19-specific CAR⁺ T cell in 1,000 PBMC using anti-CD19scFv tagged to a AlexaFluor 647 dye. The assay was validated in PBMC obtained from two individual donors by an independent laboratory (Immune Monitoring Core lab) at MDACC (Representative data along with dilution schematic and gating strategy are provided in FIG. 9 (i-iii). The anti-CD19scFv mAb could help in detecting the presence of CAR⁺ T cells in patients after infusion

(demonstrating the detection of CAR⁺ T cells with central memory immunophenotype in PBMC).

[00119] Persistence of adoptively transferred CAR⁺ T cells in patients correlates with efficacy in eradicating tumor. The inventors desired that clone no. 136.20.1 be used in correlative studies pertaining to clinical trials that infuse CAR⁺ T cells and thus wanted to check if this mAb could detect CAR⁺ cells preserved in various harsh organic solvent fixatives. CARs could be detected by immunocytochemistry and TEM. At day 28 of culture CD19-specific CAR⁺ T cells were harvested and fixed by paraformaldehyde. FIG. 5A shows that anti-CD19scFv mAb, when used at a concentration of 1:500, can establish surface localization of CARs in genetically modified T cells by immunocytochemistry. The inventors have also established the utility of this mAb in immunohistochemistry using CAR⁺ T cells fixed in neutral buffer formalin (FIG. 10) and are currently working to detect CAR⁺ T cells in bone marrow of patients infused with CD19-specific CAR⁺ T cells. To detect CAR molecules on surface of genetically engineered T cells by electron microscopy, mAb (clone no. 136.20.1) was tagged to gold-labeled nanoparticles. Control unmodified T-cell sections are shown in FIG. 5B (i-iii). The size of the conjugated nanogold (1.4 nm) precluded detection within thick sections of CAR⁺ T cells (FIG. 5B (iv)). Thus, the inventors performed a silver enhancement process and could view CARs as detected by clone no. 136.20.1 conjugated nanoparticles within 100 nm thin sections (FIG. 5B (v-vi)). The silver enhancement was performed on non-stained samples as uranyl acetate staining prevented distinguishing nanoparticles from other cellular granules. These studies establish the utility of clone no. 136.20.1 to detect CAR⁺ T cells in fixed sections.

[00120] **Anti-CD19scFv mAb can block the effector function of CAR⁺ T cells.** The inventors evaluated functional properties of anti-CD19scFv mAb (clone no. 136.20.1) based on its ability to prevent lysis mediated by CD19-specific CAR⁺ T cells. Previously, the inventors developed and validated an approach for *ex vivo* expansion of genetically modified CAR⁺ T cells on artificial antigen presenting cells (K562 clone #4). CAR⁺ T cells receive proliferative signals from the TAA (CD19) coordinated with co-stimulatory molecules (CD86, CD137L, membrane bound IL15) cloned onto the surface of K562 cells (FIG. 3C) (Singh *et al.*, 2008; Manuri *et al.*, 2009; Singh *et al.*, 2011). The CD19-specific CAR⁺ T cells used in this study were propagated on the same K562-derived aAPC. A standard 4 h CRA was performed using mouse T-cell lines EL4 (parental) and CD19⁺ EL4 as

the tumor target. The percentage specific lysis was calculated using CD19-specific CAR⁺ T cells bound to CD19scFv mAb or CD19-specific CAR⁺ T cells alone. FIG. 6A shows that anti-CD19scFv mAb completely inhibits CAR-dependent killing (at 250 µg/mL) by blocking the ability of CAR to dock to the TAA. However, at 25 µg/mL the anti-Id inhibited killing at 30% maximum and there was little or no effect when the mAb concentration was reduced to 5 µg/mL (FIG. 6B). The inventors repeated the experiment in two other B-cell lines (Daudiβ₂m and NALM-6) to validate that inhibition of lysis could be achieved in alternate B-cell tumors. The application of anti-CD19scFv at 250 µg/mL reduced the cytolytic activity to 18% in Daudiβ₂m and 32% in NALM-6 (FIG. 11). However, the mechanism of lysis may not be same in all these tumor cell lines and hence the degree of inhibition differed. To investigate the inhibitory effect of anti-CD19scFv mAb in real time, the inventors used VTLM to serially monitor the killing process in a co-culture experiment involving Daudiβ₂m as a tumor target and CD19-specific CAR⁺ T cells as effectors. As a control, the inventors used a Fc-specific F(ab')₂ fragment of an antibody that bound to the scaffolding of CD19RCD28. CAR⁺ T cells continued to lyse tumor cells when the effector cells were bound by this control antibody (FIG. 6i-iii). In contrast, CAR⁺ T cells were unable to kill target cells when bound by anti-CD19scFv mAb (FIG. 6iv-vi). Thus, blocking by clone no. 136.20.1 prevented triggering of T cells, which supports the claim that this mAb binds within the scFv domain of the CAR.

Example 3 – A novel CAR-targeting functional monoclonal antibody (2D3) amenable for CAR-mediated gene therapy

[00121] Targeting tumor antigens through gene modified immune cells has increasingly been adopted by clinical oncologists, thanks to the ability of these cells to eradicate tumors that have been otherwise refractory to conventional therapies. The clinical trials conducted so far showed encouraging results. There is an increasing demand to place specific reagents so that isolation, activation, and functional characterization of these gene modified cells are possible. In this context, the inventors developed a Chimeric Antigen Receptor (CAR)-specific monoclonal antibody (2D3) that possesses unique ability to detect all chimeric antigen receptor (all CAR)-modified immune cells (FIG. 12). The 2D3 antibody employs the immunoglobulin heavy chain constant region to cross-link the T cell signaling endodomain with antigen binding domain of a CAR. This antibody has multiple functions, such as detection (FIG. 12), quantification, activation, and selective propagation (FIG. 13) of

CAR modified cells, and hence will likely be an important reagent suitable for CAR-modified gene therapy.

[00122] A novel and promising form of cancer therapy genetically modifies a patient's own T cells in peripheral blood to target and kill their cancer using a chimeric antigen receptor (CAR) directed against a tumor associated antigen (TAA). Growing a therapeutic number of CAR⁺ T cells to treat cancer patients requires an artificial antigen presenting cell (aAPC) to express the TAA that the CAR⁺ T cell targets. Antigen-specific growth using a cell based system requires at least a month of intensive monitoring for outgrowth of unwanted cell types. Here, the inventors provide a system that decreases the high cost and time associated with manufacturing a different aAPC for each TAA and decreases the time and effort needed to develop, validate, and manufacture large numbers of CAR⁺ T cells. An HLA-null K562 (HnK) was modified to surface express a chimeric antibody receptor that could bind to spacer domain present in all to all other CARs produced by the inventor's lab which are G4CAR. In another aspect, the specified spacer domain (Hinge-CH2-CH3) may also be used for CAR expression. When peripheral blood mononuclear cells (PBMC) are genetically modified to express a CAR containing this domain, those CAR⁺ PBMC can be co-cultured with G4CAR⁺ HnK (G4HnK) to specifically grow CAR⁺ T cells independent of TAA. Utilizing G4HnKs to grow CAR⁺ T cells generates a sufficient number of CAR⁺ T cells for therapy in half the amount of time typically needed utilizing antigen-specific aAPC. The T cells consistently grow without competition to a pure population within 2 weeks (>80% CAR⁺ T cells), and this level of infusion product purity is rarely achieved using current aAPC before one month of T cell co-culture with standard aAPC. Also unique to this system is the capacity to expand antigen non-specific CAR⁺ T cells as a means of antibody-independent T cell growth without tissue targeting. This provides a novel method of expanding CAR modified T cells developed on a common backbone but targeted to alternate tumor antigens. In conclusion, the inventors have made a single aAPC cell line capable of expanding any CAR⁺ T cell independent of a CAR-specific antigen.

Table 3: Sequence of All CAR mAb 2D3

mAb	Heavy Chain Sequence			Light Chain Sequence		
	CDR1	CDR2	CDR3	CDR1	CDR2	CDR3
	Amino acid sequence			Amino acid sequence		
2D3	GYTFTSYW (SEQ ID NO: 10)	IYPGNSDT (SEQ ID NO: 11)	TRVNWDGY STMDY (SEQ ID NO: 12)	QSLVHSNG NTY (SEQ ID NO: 14)	KVS (SEQ ID NO: 15)	CSQTT HVPYT F (SEQ ID NO: 16)
	PREXKLEQSGTVLARPGASVKMSCKTSGYT FTSYWMHWVIQRPQGQLEWIGAIYPGNSDT NYNQKFKVKAKLTAVTSTSTAYMELSSLTD EDSAVYYCTRVNWDGYSTMDYWGQGSVTV SSAKTTPPSVYASQRANSRP (SEQ ID NO: 9)			STLSLPVSLGDQASISCRSSQSLVH SNGNTYFHWYLQKPGQSPKLLIYKV SNRFSGVDPDRFSGSGSGTNFTLKIS RVEAEDLGVIYFCSQTTHVPYTFGGG TELEIKRADAAPTVS (SEQ ID NO: 13)		

Example 4 – Universal artificial antigen presenting cells to selectively propagate T cells expressing chimeric antigen receptor independent of specificity

- 5 **[00123] Cells and culture conditions.** K562 cells (European Collection of Cell Cultures through Sigma-Aldrich, St. Louis, MO; Cat. No. 89121407), noted for expression of desired endogenous adhesion molecules and the absence of most HLA class I and all class II molecules, (Suhoski *et al.* 2007) were used to derive CD19⁺ and CARL⁺ K562 that served as aAPC. Immortalized tumor targets CD19^{neg}, GD2⁺ EL-4 murine thymoma
- 10 (Cat. No. TIB-40) and CD19⁺, GD2^{neg} NALM-6 (pre-B cell leukemia, Cat. No. CRL-1567) were purchased from American Type Culture Collection (ATCC, Manassas, VA). Identity of cell lines was validated by the MDACC Cancer Center Support Grant Characterized Cell Line Core using short tandem repeat DNA fingerprinting. Peripheral blood was donated by consenting healthy volunteer adults at Gulf Coast Regional Blood Center (Houston, TX).
- 15 Peripheral blood mononuclear cells (PBMC) were isolated using Ficoll-Paque Plus density centrifugation (GE Healthcare Biosciences, Piscataway Township, NJ; Cat. No. 17-1440-02) before freezing in a mixture of 10% DMSO (Sigma, Allentown, PA; Cat. No. D2650), 50% heat-inactivated fetal bovine serum (FBS-Thermo Scientific Hyclone, Bridgewater, NJ, Cat. No. SH30070.03), and 40% RPMI 1640 (Thermo Scientific Hyclone; Cat. No. SH30096.01).

All cells were cultured in a 37° C humidified incubator with complete media (CM) prepared from RPMI 1640, 10% heatinactivated FBS, and 2 mM GlutaMAX supplement (Life Technologies, Grand Island, NY; Cat. No. 35050061).

[00124] DNA expression plasmids. Codon-optimized

5 CD19RCD28mZ(CoOp)/pSBSO, (Singh *et al.* 2008) which codes for CAR between transposition sites mediated by SB transposase, (Davies *et al.* 2010) was used as the vector backbone for cloning of the following transgenes. The DNA plasmid 19G4CAR (also designated CD19RCD28 [Kowolik *et al.* 2006] FIG. 19A) codes for a 2nd generation CD19-specific CAR containing a modified IgG4 exodomain, CD28 transmembrane, and
 10 CD28/CD3 ζ endodomain. The synthesis of DNA plasmid GD2G4CAR (FIG. 19B), specific to sphingolipid GD2, utilized the same 19G4CAR backbone. The GD2-specific scFv derived from murine mAb (clone 14G2a) (Alvarez-Rueda *et al.* 2011) was commercially synthesized (Geneart, Life Technologies) as codonoptimized cDNA with NheI and XmnI restriction enzyme (RE) sites flanking the scFv. The 19G4CAR plasmid backbone and GD2-specific
 15 scFv cDNA were excised using these REs and ligated to replace CD19-specific scFv with GD2- specific scFv. A DNA plasmid (FIG. 19C) coding for a control CAR that contains no scFv region, designated G4CAR, but does contain an Ig κ -FLAG peptide sequence (METDTLLLWVLLLWVPGSTGDYKDEGTS; SEQ ID NO: 19), was derived from 19G4CAR using primer-directed PCR amplification from the beginning of the IgG4 domain
 20 hinge (primer 5'GGTACCTCTGGGGGGCAGGGCCTGCATG3'; SEQ ID NO: 20) to the terminus of the CD3 ζ domain (primer 5'GGGCCCAGCGCTGAGAGCAAGTACGGCCCTCCC3'; SEQ ID NO: 21) and sequence verified. The G4CAR was ligated into the 19G4CAR backbone ApaI and KpnI RE sites. DNA plasmid coding for a CD19-specific CAR with no IgG4 (FIG. 19D), designated
 25 19CAR, encodes from amino to carboxyl termini a GM-CSF (amino acid 1-22; NP_758452.1 (SEQ ID NO: 22)), CD19-specific scFv (245 amino acids), CD8 α extracellular domain and hinge (amino acids 136-203; NP_001759 (SEQ ID NO: 23)) followed by the same CD28 transmembrane and CD28 and CD3 ζ domains as other CARs. (Kowolik *et al.* 2006) The full length of this transgene was synthesized by GeneArt, cut with ClaI and SpeI REs, and ligated
 30 into the 19G4CAR backbone replacing the 19G4CAR codon, which had been excised using EcoRV and SpeI. The scFv sequence of CARL was derived from the cDNA library of the 2D3 hybridoma. (Singh *et al.* 2008) This was achieved by extracting RNA from 5x10⁶ cells using the RNeasy Mini Kit (Qiagen, Gaithersburg, MD; Cat No. 74104) according to

manufacturer's protocol. A cDNA library was generated by reverse transcription using oligo-dT primers per the protocol provided in the Superscript III First Strand kit (Invitrogen; Cat No. 18080-051). PCR (using Amplitaq Gold) was performed on the cDNA using the degenerate primers for the FR1 region (Wang *et al.* 2000) to amplify the mouse VH and VL regions. The VH and VL amplified products were ligated into the TOPO TA vector and sequenced. The CARL construct for surface expression on aAPC was composed of GM-CSF leader peptide (amino acid 1-22; NP_758452.1 (SEQ ID NO: 22)) fused to the 2D3-derived scFv, and tethered by CD8 α (amino acid 136-182; NP_001759.3 (SEQ ID NO: 23)) to the transmembrane and intracellular portions of CD28 (amino acid 56-123; NP_001230006.1 (SEQ ID NO: 24)) followed by CD3 ζ (amino acid. 48-163; NP_000725.1 (SEQ ID NO: 25)) intracellular domain. Design of all transgenes utilized Vector NTI Advance™ 11 software (Invitrogen). All transgenes were human codon optimized before synthesis at GeneArt. The CARL construct was excised and ligated into a SB expression plasmid, designated Zeo-2A-CARL (FIG. 19E), to co-express. Before freezing, the aAPC were routinely tested for the presence of transgenes, absence of mycoplasma, and absence of endotoxin.

[00125] Propagation of CAR⁺ T cells (CART). The designs of each CAR and antigen (CARL and CD19) as expressed on the respective T cell or aAPC are shown in FIG. 14A. The propagation of CART is depicted in FIG. 14B. Each CAR from FIG. 14A.II was co-cultured with aAPC from FIG. 14A.I. At the initiation of the experiment (defined as Day 0), thawed PBMC were washed twice, and maintained in CM for 3 to 4 hours before electroporation using the Amaxa 2D Nucleofector under program U-14 with human T cell Nucleofector kit (Lonza Biosciences; Cat No. VPA-1002). After resting overnight in CM, viable PBMC (counted by exclusion of 0.1% Trypan Blue) were resuspended in CM and mixed at a 1:2 ratio (mononuclear cell to γ -irradiated aAPC) using thawed aAPC that were washed twice and counted. The co-culture contained 10^6 total cells/ mL in CM and 50 IU/mL recombinant human IL-2 (Proleukin, Prometheus Labs, San Diego, CA). The live-cell counts were determined by Trypan Blue exclusion on Days 1, 7, 14, and 21 of co-culture. Flow cytometry for CD3, CD4, CD8, and human IgG (to assess CAR expression) occurred on Days 1, 7, 14, and 21, and for CD45RO, CD62L, and CD28 occurred on Days 14 and 21. Irradiated aAPC were re-added to co-cultures on Days 7 and 14 by re-stimulating mononuclear cells with γ -irradiated aAPC at 1:2 ratio. On Day 21 products of propagation were assessed for specific killing, and DNA and RNA were extracted. Each experiment was repeated at least 4 times using 5 donors.

[00126] Flow cytometry. FACSCalibur (BD Biosciences, Billerica, MA) was used to acquire samples prepared in FACS staining solution as previously described. (Deniger *et al.* 2013) After washing once in FACS staining solution, cells were stained for 30 minutes at 4° C without blocking in FACS staining solution containing a 1:33 dilution of antibody. When anti-human Fc antibody was used, the anti-Fc stained sample was washed and re-stained for alternative surface markers before re-suspension in FACS buffer for flow cytometer analysis. Measurement of intracellular cytokine used the same protocol for cell surface staining followed by 20 min. fixation using BD cytofix/cytoperm kit fixative (BD Biosciences; Cat No. 554714), followed by washing twice in perm/wash buffer containing 20% FBS and staining with a 1:33 dilution of antibody in perm/wash buffer. Antibody incubation lasted 30 minutes at 4°C before samples were washed in perm/wash buffer and resuspended in FACS staining solution for acquisition. FlowJo v 10.0.5 (Tree Star Inc., Ashland, OR) was used for analysis of flow cytometry data.

[00127] Chromium release assay (CRA). CRA was performed as previously described. (Singh *et al.* 2008) In brief, on Day 21 of T-cell co-culture on aAPC, the tumor targets (i) EL-4, (ii) NALM-6, and (iii) K562 were loaded with ⁵¹Cr for 3 hours, and, after washing, co-cultured with effector T cells for 6 hours at 37° C using a ratio of 5 T cells to 1 target cell.

[00128] Abundance and diversity of TCR repertoire. The direct TCR expression assay (DTEA), as previously reported, (Zhang *et al.* 2012) was used to measure the abundance of mRNA transcripts coding for 45 TCR α alleles, 46 TCR β alleles, 13 TCR γ alleles, and 5 TCR δ alleles from RNA obtained on Day 0 (T cells in PBMC before electroporation) and Day 21 (from T cells after electroporation/ propagation). Day 0 samples were negatively sorted for CD56 (Miltenyi Biotec, Cambridge, MA; Cat. No. 130-050-401) and then positively sorted for CD3 (Miltenyi Biotec; Cat. No. 130-050-101). The resulting CD3⁺CD56^{neg} T cells (2 to 3 x 10⁶ from each sample) were snap frozen as were 2 x 10⁶ cells directly harvested at Day 21 of co-culture. RNA was extracted from thawed samples using the ALLprep DNA/RNA mini kit (Qiagen; Cat. No. 80204).

[00129] Statistics. Statistical analysis was performed using Prism v6.0 (Graph Pad Software Inc.). Student's t-test (unpaired) was used to perform two sample comparisons. One- or two-way ANOVA F-test was used to perform group comparisons, and if found significant ($p < 0.05$); a t-test (unpaired) was undertaken to assess and report differences.

Spearman's nonparametric correlation was performed on housekeeping gene normalized DTEA transcript counts to assess the divergence of the TCR repertoire in T cells from an experimental group and autologous Day 0 PBMC. If the Spearman correlation coefficient was greater than or equal to 0.8 ($p \geq 0.8$) within the 95% confidence interval of the correlation coefficient, then the two TCR repertoires were considered to be highly correlated.

Results

[00130] Numeric expansion of CAR+ T cells upon K562 cells expressing CARL or CD19. Mouse mAb clone 2D3 was obtained by repeated footpad injections of NSO cells expressing 19G4CAR into BALB/c mice and blocking studies defined the specificity of the mAb to the human IgG4 exodomain of 19G4CAR. (Singh *et al.* 2008) It was hypothesized that this mAb may be used to cross-link CAR and activate genetically modified T cells for sustained proliferation. Therefore, the scFv of 2D3 (designated CARL) was expressed on the cell surface to compare with human truncated CD19 TAA on K562 cells. The CARL and CD19 transgenes were cloned into DNA plasmids for co-expression with drug-selection genes between SB transposable elements. The SB transposon DNA plasmids for 2D3-derived scFv or CD19 were electro-transferred with SB11 transposase DNA plasmid into K562 cells in separate experiments. Genetically modified cells were propagated under drug selection from a single cell for homogeneous expression of CARL (as detected by antibody against mouse Fab) or CD19 (FIG. 15A). A comparison of the γ -irradiated K562-derived aAPC to selectively propagate CART was undertaken following electroporation (defined as Day 0) of the panel of CAR constructs (FIG. 14A.II) into PBMC using SB system. On Day 1, initial expression of CAR in T cells was evaluated by flow cytometry using antibody specific for human Fc (FIG. 15B). The expression of CARs and number of total viable T cells were measured weekly for 21 days of co-culture with CD19+ K562 or CARL+ K562 with the following immunoreceptors on T cells; (i) 19G4CAR with specificity for CD19 and containing the IgG4 exodomain, (ii) 19CAR with specificity for CD19 and absence of IgG4 exodomain, (iii) G4CAR without scFv, but containing an IgG4 exodomain, and (iv) GD2G4CAR with specificity for GD2 and containing the IgG4 exodomain. All CAR species contained the same transmembrane and intracellular domains (CD28/CD3 ζ) as the 2nd generation 19G4CAR. (Singh *et al.* 2008) The co-cultures of CART with the two types of aAPC were found to have significantly different amounts of T cells by Day 21 depending on the choice of aAPC ($p < 0.05$) using two-way ANOVA followed by un-

paired t-tests. The 19CAR⁺ T cells proliferated upon co-culture with CD19⁺ K562 and G4CAR⁺ T cells proliferated upon co-culture with CARL⁺ K562 in an exponential fashion, whereas 19CAR on CARL⁺ K562 and G4CAR on CD19⁺ K562 did not numerically expand (FIG. 16A top panel). These data indicate that, as expected, the CD19 TAA on aAPC selectively supports the outgrowth of CD19-specific CART. Furthermore, they demonstrate that CARL can activate T cells to proliferate that contain a CAR species with an IgG4 exodomain. Next, the ability of CD19⁺ K562 and CARL⁺ K562 were assessed for ability to sustain the proliferation of T cells expressing 19G4CAR to evaluate how two modes of crosslinking (FIG. 14C) can activate T cells. There were no significant differences in the accumulated number of viable T cells on Day 21 of co-culture based on the type of aAPC used (FIG. 16A bottom panel), the expression of CAR as a percent of the population (FIG. 16B bottom panel), or the mean fluorescence intensity (MFI). A trend ($p = 0.09$) towards a difference in MFI of CAR expression resulting from aAPC employed to expand 19G4CAR was noted (FIG. 15C). There were no significant differences (FIG. 16C & D) between the two aAPC types for propagating 19G4CAR⁺ T cells co-expressing cell-surface proteins associated with memory phenotype ($p = 0.82$), 25 or other co-receptors ($p = 0.26$), as well as the specific lysis by 19G4CAR⁺ T cells ($p = 0.16$). Therefore, CD19-specific CAR⁺ T cells can be propagated in similar quantity and quality by K562-derived aAPC expressing CAR or CD19.

[00131] CARL⁺ K562 can numerically expand CAR⁺ T cells independent of specificity. The 2D3-derived scFv on aAPC was evaluated for ability to propagate not just CD19-specific T cells, but CAR⁺ T cells of alternative specificities. The CD19 and GD226 TAAs are not present on parental K562 cells to propagate T cells expressing GD2G4CAR, 19G4CAR, and G4CAR. Nonetheless, T cells bearing these three CARs numerically expanded on CARL⁺ K562. The number of total viable T cells on Day 21 of co-culture with CARL⁺ K562 cells did not significantly differ between 19G4CAR, G4CAR, and GD2G4CAR ($p = 0.16$, FIG. 17A). Similarly, the percentage of each CAR expressed on T cells at Day 21 did not significantly differ among the three populations of genetically modified T cells ($p = 0.68$, FIG. 17B). Finally, the electroporated and propagated T cells exhibited specific lysis of CD19 and GD2 TAAs recognized by CD19-specific and GD2-specific CARs. EL-4 cells, previously reported to express GD2 (Alvarez-Rueda *et al.* 2011) were specifically killed by GD2G4CAR⁺ T cells and not with T cells expressing G4CAR or 19G4CAR. As anticipated, CD19⁺ NALM-6 cells were targeted by T cells expressing

19G4CAR (FIG. 17C). In summary, genetically modified T cells can be selectively propagated by CARL⁺ K562 cells resulting in T cells that retain specificity for TAA and stable expression of CAR.

[00132] The choice of aAPC does not skew the TCR repertoire for

numerically expanded CART. Each T cell in peripheral blood bears a distinct pair of $\alpha\beta$ or $\gamma\delta$ TCRs which can be analyzed using the direct TCR expression assay (DTEA) to determine the abundance of TCR chains. This assay was employed to determine whether CARL⁺ or CD19⁺ K562 influenced the distribution of TCR alleles after 21 days co-culture on aAPC. TCR variants were assayed on the nCounter Analysis System using a set of 111 TCR α , β , γ , and δ transcripts. (Deniger *et al.* 2013; Zhang *et al.* 2012) By measuring the distribution of TCR alleles we could determine if the aAPC design preferentially supported the numeric expansion of some, but not all genetically modified T cells. The starting TCR distribution of T cells on Day 0 was ranked from the most to least frequent TCR usage and the rank order compared for T cells harvested on Day 21 of co-culture with aAPC (FIG. 18A). This revealed no apparent monoclonal or oligoclonal outgrowth of electroporated T cells propagated on CARL⁺ or CD19⁺ K562 cells. The ranks of TCR frequencies on Day 0 and Day 21 from each experiment were compared using Spearman's rank correlation test and found to significantly correlate ($p < 0.0001$; Table 4). The statistical comparison of TCR abundance and type from Day 0 and 21 indicated that all correlation coefficients (ρ) had values greater than 0.8 within the 95% confidence interval of ρ which is consistent with a strong correlation, indicating no change in TCR frequency. The measurement of TCR abundance demonstrates that CARL or TAA on aAPC do not skew the outgrowth of subpopulations of propagated T cells, but rather that both 2D3-derived scFv and CD19 on K562 cells can sustain the outgrowth of CAR⁺ T cells that maintain a polyclonal repertoire.

[00133] Table 4: Comparison of TCR abundance harvested from T cells before versus after propagation on aAPC*.

		Day 0	Day 21			
			19G4CAR & CARL ⁺ K562	G4CAR & CARL ⁺ K562	GD2G4CAR & CARL ⁺ K562	19G4CAR & CD19 ⁺ K562
Day 0			0.748 (0.65-0.82)	0.857 (0.80-0.90)	0.867 (0.81-0.91)	0.912 (0.87-0.94)

		Day 0	Day 21			
Day 21	19G4CAR & CARL ⁺ K562	0.752 (0.65-0.83)		0.805 (0.72-0.86)	0.706 (0.59-0.79)	0.734 (0.63-0.81)
	G4CAR & CARL ⁺ K562	0.899 (0.85-0.93)	0.71 (0.61-0.79)		0.816 (0.74-0.87)	0.839 (0.77-0.89)
	GD2G4CAR & CARL ⁺ K562	0.825 (0.75-0.88)	0.72 (0.61-0.80)	0.801 (0.72-0.86)		0.881 (0.83-0.92)
	19G4CAR & CD19 ⁺ K562	0.916 (0.87-0.94)	0.69 (0.57-0.78)	0.887 (0.84-0.92)	0.808 (0.73-0.87)	

* Analysis DTEA data from two donors was normalized using housekeeping genes and assessed using Spearman correlation coefficient to compare distributions of TCR usage for two donors. The upper right of the table contains the correlation between experimental groups for one donor. The lower left of the table contains the second donor subjected to the same analysis. Each cell in the table contains the Spearman correlation coefficient (ρ) and within the brackets the 95% confidence interval. (A strong correlation is considered to be $\rho \geq 0.8$.)

* * *

10 **[00134]** All of the methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the methods and in the steps or in the sequence of steps of the method described herein

15 without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the

20 appended claims.

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

Alvarez-Rueda N, Desselte A, Cochonneau D, et al. A monoclonal antibody to O-acetyl-GD2 ganglioside and not to GD2 shows potent antitumor activity without peripheral nervous system cross-reactivity. *PloS One*. 2011;6(9):e25220.

10 Brentjens *et al.*, Safety and persistence of adoptively transferred autologous CD19-targeted T cells in patients with relapsed or chemotherapy refractory B-cell leukemias. *Blood*, 118:4817-4828, 2011.

Brentjens *et al.*, CD19-targeted T cells rapidly induce molecular remissions in adults with chemotherapy-refractory acute lymphoblastic leukemia *Sci. Transl. Med.*, 15 5:177ra38, 2013.

Cooper *et al.*, T-cell clones can be rendered specific for CD19: toward the selective augmentation of the graft-versus-B-lineage leukemia effect. *Blood*, 101:1637-1644, 2003.

Cooper *et al.*, Good T cells for bad B cells. *Blood*, 119:2700-2702, 2012.

20 Davies JK, Singh H, Huls H, et al. Combining CD19 redirection and alloanergization to generate tumor-specific human T cells for allogeneic cell therapy of B-cell malignancies. *Can Res*. May 15 2010;70(10):3915-3924.

Deniger DC, Switzer K, Mi T, et al. Bispecific T-cells expressing polyclonal repertoire of endogenous gamma delta T-cell receptors and introduced CD19-specific chimeric antigen receptor. *Mol Ther*. Mar 2013;21(3):638-647.

Hackett *et al.*, A transposon and transposase system for human application. *Mol. Ther.*, 18:674-683, 2010.

Hombach *et al.*, Adoptive immunotherapy with genetically engineered T cells: modification of the IgG1 Fc 'spacer' domain in the extracellular moiety of chimeric antigen receptors avoids 'off-target' activation and unintended initiation of an innate immune response. *Gene Ther.*, 17:1206-1213, 2010.

30 Huls *et al.*, Clinical Application of Sleeping Beauty and Artificial Antigen Presenting Cells to Genetically Modify T Cells from Peripheral and Umbilical Cord Blood. *J. Vis. Exp.*, doi:10.3791/50070, 2012.

- Jena *et al.*, Redirecting T-cell specificity by introducing a tumor-specific chimeric antigen receptor. *Blood*, 116:1035-1044, 2010.
- Jena *et al.* Chimeric Antigen Receptor (CAR)-specific monoclonal antibody to detect CD19-specific T cells in clinical trial. *PLoS One*, 8(3):e57838, 2013
- 5 June *et al.*, Engineering lymphocyte subsets: tools, trials and tribulations. *Nat. Rev. Immunol.*, 9:704-716, 2009.
- Kalos *et al.*, T cells with chimeric antigen receptors have potent antitumor effects and can establish memory in patients with advanced leukemia. *Sci. Transl. Med.*, 3:95ra73, 2011.
- 10 Kochenderfer *et al.*, Eradication of B-lineage cells and regression of lymphoma in a patient treated with autologous T cells genetically engineered to recognize CD19. *Blood*, 116:4099-4102, 2010.
- Kochenderfer *et al.*, B-cell depletion and remissions of malignancy along with cytokine-associated toxicity in a clinical trial of anti-CD19 chimeric-antigen-receptor-transduced T cells. *Blood*, 119:2709-2720, 2012.
- 15 Kowolik *et al.*, CD28 costimulation provided through a CD19-specific chimeric antigen receptor enhances in vivo persistence and antitumor efficacy of adoptively transferred T cells. *Cancer Res.*, 66:10995-11004, 2006.
- Lamers *et al.* Immune responses to transgene and retroviral vector in patients treated with ex vivo-engineered T cells. *Blood*, 117:72-82, 2011.
- 20 Ma *et al.*, Anti-prostate specific membrane antigen designer T cells for prostate cancer therapy. *Prostate*, 61:12-25, 2004.
- Maiti *et al.* *J Immunother.* Sleeping beauty system to redirect T-cell specificity for human applications 36(2):112-23, 2013.
- 25 Manuri *et al.*, piggyBac transposon/transposase system to generate CD19-specific T cells for the treatment of B-lineage malignancies. *Hum. Gene Ther.*, 21:427-437, 2009.
- Moeller *et al.*, Adoptive transfer of gene-engineered CD4+ helper T cells induces potent primary and secondary tumor rejection. *Blood*, 106:2995-3003, 2005.
- Morgan *et al.*, Cancer regression in patients after transfer of genetically engineered lymphocytes. *Science*, 314:126-129, 2006.
- 30 Morgan, Adoptive cell therapy: genetic modification to redirect effector cell specificity. *Cancer J.*, 16:336-341, 2010.

- Nicholson *et al.* Construction and characterisation of a functional CD19 specific single chain Fv fragment for immunotherapy of B lineage leukaemia and lymphoma. *Mol. Immunol.*, 34:1157-1165, 1997.
- O'Connor *et al.*, Adoptive T-cell therapy improves treatment of canine non-Hodgkin
5 lymphoma post chemotherapy. *Scientific Reports*, 2, 2012.
- Park *et al.*, Adoptive transfer of chimeric antigen receptor re-directed cytolytic T lymphocyte clones in patients with neuroblastoma. *Mol. Ther.*, 15:825-833, 2007.
- Porter *et al.*, Chimeric Antigen Receptor-Modified T Cells in Chronic Lymphoid Leukemia. *New England Journal of Medicine*, 365:725-733, 2011.
- 10 Pule *et al.*, Virus-specific T cells engineered to coexpress tumor-specific receptors: persistence and antitumor activity in individuals with neuroblastoma. *Nat. Med.*, 14:1264-1270, 2008.
- Rabinovich *et al.* A role for the MHC class I-like Moll molecules in nutrient metabolism and wound healing. *Immunol. Cell Biol.* 86: 489-496, 2008
- 15 Rossig *et al.*, Targeting of G(D2)-positive tumor cells by human T lymphocytes engineered to express chimeric T-cell receptor genes. *Int. J. Cancer*, 94:228-236, 2001.
- Rossig *et al.*, Epstein-Barr virus-specific human T lymphocytes expressing antitumor chimeric T-cell receptors: potential for improved immunotherapy. *Blood*, 99:2009-2016, 2002.
- 20 Savoldo *et al.*, CD28 costimulation improves expansion and persistence of chimeric antigen receptor-modified T cells in lymphoma patients. *J. Clin. Invest.*, 121:1822-1826, 2011.
- Singh *et al.*, Redirecting specificity of T-cell populations for CD19 using the Sleeping Beauty system. *Cancer Res.*, 68:2961-2971, 2008.
- 25 Singh *et al.*, Reprogramming CD19-specific T cells with IL-21 signaling can improve adoptive immunotherapy of B-lineage malignancies. *Cancer Res.*, 71:3516-3527, 2011.
- Till *et al.*, Adoptive immunotherapy for indolent non-Hodgkin lymphoma and mantle cell lymphoma using genetically modified autologous CD20-specific T cells. *Blood*,
30 112:2261-2271, 2008.
- Torikai *et al.*, A foundation for universal T-cell based immunotherapy: T cells engineered to express a CD19-specific chimeric-antigen-receptor and eliminate expression of endogenous TCR. *Blood*, 119:5697-5705, 2012.

- Vera *et al.*, T lymphocytes redirected against the kappa light chain of human immunoglobulin efficiently kill mature B lymphocyte-derived malignant cells. *Blood*, 108:3890-3897, 2006.
- 5 Wang Z, Raifu M, Howard M, et al. Universal PCR amplification of mouse immunoglobulin gene variable regions: the design of degenerate primers and an assessment of the effect of DNA polymerase 3' to 5' exonuclease activity. *J Immuno Methods*. Jan 13 2000;233(1-2):167-177.
- Whitlow *et al.*, An improved linker for single-chain Fv with reduced aggregation and enhanced proteolytic stability. *Protein Engineering*, 6:989-995, 1993.
- 10 Zhang M, Maiti S, Bernatchez C, et al. A new approach to simultaneously quantify both TCR alpha- and beta-chain diversity after adoptive immunotherapy. *Clin Can Res*. Sep 1 2012;18(17):4733-4742.
- Zheng *et al.*, Protein L: a novel reagent for the detection of chimeric antigen receptor (CAR) expression by flow cytometry *J. Transl. Med.*, 10:29, 2012.

CLAIMS

1. An isolated monoclonal antibody, wherein the antibody specifically binds to a chimeric antigen receptor (CAR) polypeptide and wherein the antibody competes for binding
5 of the polypeptide with the 2D3 monoclonal antibody.
2. The isolated antibody of claim 1, wherein the antibody comprises:
 - (a) a first V_H CDR at least 80% identical to V_H CDR1 of 2D3 (SEQ ID NO: 10);
 - (b) a second V_H CDR at least 80% identical to V_H CDR2 of 2D3 (SEQ ID NO: 11);
 - (c) a third V_H CDR at least 80% identical to V_H CDR3 of 2D3 (SEQ ID NO: 12);
 - 10 (d) a first V_L CDR at least 80% identical to V_L CDR1 of 2D3 (SEQ ID NO: 14);
 - (e) a second V_L CDR at least 80% identical to V_L CDR2 of 2D3 (SEQ ID NO: 15);
 - and
 - (f) a third V_L CDR at least 80% identical to V_L CDR3 of 2D3 (SEQ ID NO: 16).
3. The isolated antibody of claim 2, wherein the antibody comprises:
 - 15 (a) a first V_H CDR is identical to SEQ ID NO: 10;
 - (b) a second V_H CDR is identical to SEQ ID NO: 11;
 - (c) a third V_H CDR is identical to SEQ ID NO: 12;
 - (d) a first V_L CDR is identical to SEQ ID NO: 14;
 - (e) a second V_L CDR is identical to SEQ ID NO: 15; and
 - 20 (f) a third V_L CDR is identical to SEQ ID NO: 16.
4. The isolated antibody of claim 1, wherein the antibody comprises a V_H domain at least about 80% identical to the V_H domain of 2D3 (SEQ ID NO: 9) and a V_L domain at least about 80% identical to the V_L domain of 2D3 (SEQ ID NO: 13).
5. The isolated antibody of claim 4, wherein the antibody comprises a V_H domain
25 identical to the V_H domain of 2D3 (SEQ ID NO: 9) and a V_L domain identical to the V_L domain of 2D3 (SEQ ID NO: 13).
6. The isolated antibody of any one of claims 1-3, wherein the antibody is a membrane-bound antibody.

7. The isolated antibody of claim 6, wherein the membrane-bound antibody comprises one or more of (i) a GM-CSF leader peptide; (ii) 2D3 scFv coding sequence; (iii) a CD8 α linker sequence; (iv) a CD28 transmembrane domain; and/or (v) a CD3 ξ intracellular domain.
8. The isolated antibody of claim 4, wherein the antibody comprises a V_H domain
5 between 80% and 99% identical to the V_H domain of 2D3 (SEQ ID NO: 9) or a V_L domain between 80% and 99% identical to the V_L domain of 2D3 (SEQ ID NO: 13).
9. The isolated antibody of any one of claims 1-5, wherein the antibody is the 2D3 antibody.
10. The isolated antibody of any one of claims 1-5, wherein the antibody is recombinant.
- 10 11. The isolated antibody of any one of claims 1-5, wherein the antibody is an IgG, IgM, IgA or an antigen binding fragment thereof.
12. The isolated antibody of any one of claims 1-5, wherein the antibody is a Fab', a F(ab')₂, a F(ab')₃, a monovalent scFv, a bivalent scFv, or a single domain antibody.
13. The isolated antibody of any one of claims 1-5, wherein the antibody is a human,
15 humanized antibody or de-immunized antibody.
14. The isolated antibody of any one of claims 1-5, wherein the antibody is conjugated to a toxin or a reporter.
15. An isolated polynucleotide molecule comprising a nucleic acid sequence encoding an antibody of any one of claims 1-5.
- 20 16. A recombinant polypeptide comprising an antibody V_H domain comprising CDRs 1-3 of the V_H domain of 2D3 (SEQ ID NOs: 10, 11 and 12).
17. A recombinant polypeptide comprising an antibody V_L domain comprising CDRs 1-3 of the V_L domain of 2D3 (SEQ ID NOs: 14, 15 and 16).
18. An isolated polynucleotide molecule comprising a nucleic acid sequence encoding a
25 polypeptide of claim 16 or 17.
19. A host cell comprising one or more polynucleotide molecule(s) encoding an antibody that binds to the constant domain of a chimeric antigen receptor (CAR) polypeptide.

20. The host cell of claim 19, wherein the antibody competes for binding of constant domain with the 2D3 monoclonal antibody.
21. The host cell of claim 19, wherein the one or more polynucleotide molecule(s) encode an antibody of any one of claims 1-5 or a recombinant polypeptide of claim 16 or 17.
- 5 22. The host cell of claim 19, wherein the host cell is a mammalian cell, a yeast cell, a bacterial cell, a ciliate cell or an insect cell.
23. The host cell of claim 22, wherein the host cell is an antigen presenting cell.
24. The host cell of claim 23, wherein the host cell is an artificial antigen presenting cell (aAPC) or an activating and propagating cell (AAPC).
- 10 25. A method of manufacturing an antibody comprising:
(a) expressing one or more polynucleotide molecule(s) encoding a V_L and V_H chain of an antibody of any one of claims 1-5 in a cell; and
(b) purifying the antibody from the cell.
26. A method of manufacturing an antibody comprising:
15 (a) expressing one or more polynucleotide molecule(s) encoding a V_L and V_H chain of an antibody of any one of claims 1-5 in an in vitro expression system; and
(b) purifying the antibody therefrom.
27. A method for selecting a cell comprising a chimeric antigen receptor (CAR) polypeptide and/or an endogenous or introduced T-cell receptor (TCR) comprising:
20 (a) contacting the cell with an antibody in accordance with claim 1; and
(b) selecting a cell comprising the CAR or TCR based on binding of the antibody.
28. A method for selecting a cell comprising a chimeric antigen receptor (CAR) polypeptide comprising:
(a) contacting the cell with an antibody that binds to the constant domain of a
25 chimeric antigen receptor (CAR) polypeptide; and
(b) selecting a cell comprising the CAR based on binding of the antibody.

29. The method of claim 28, wherein the antibody competes for binding of constant domain with the 2D3 monoclonal antibody.
30. The method of claim 29, wherein the antibody is an antibody in accordance with claim 2.
- 5 31. The method of claim 29, wherein the antibody is an antibody in accordance with claim 3.
32. The method of claim 27 or 28, wherein the antibody is conjugated to a toxin and wherein the method is further defined as a method for selectively inhibiting or killing a cell comprising a CAR.
- 10 33. The method of claim 32, wherein the cell is *in vivo*.
34. The method of claim 27 or 28, wherein the method is further defined as a method for preventing or modifying the ability of a CAR to activate a T cell by contacting the cell with the antibody.
35. The method of claim 34, wherein the method is further defined as a method for
15 controlling toxicity of a cell.
36. The method of claim 27 or 28, wherein the method is further defined as a method for purifying or enriching cells comprising a CAR.
37. The method of claim 27 or 28, wherein contacting the cell with an antibody comprises inducing activation or growth in the cell comprising the CAR or TCR.
- 20 38. The method of claim 36, wherein the antibody comprises a reporter and the method comprises selecting cells comprising a CAR by fluorescence-activated cell sorting (FACS).
39. The method of claim 27 or 28, wherein selecting cells comprises selecting using paramagnetic beads.
40. The method of claim 27 or 28, wherein the antibody is bound to a support.
- 25 41. The method of claim 40, wherein the support is a bead, a surface, a column or an aAPC.

42. The method of claim 36, wherein purifying or enriching cells comprising a CAR comprises stimulating proliferation of the cells.

43. A cell culture system comprising mammalian T-cells expressing a CAR and an antibody that binds to the constant domain of a CAR polypeptide.

5 44. The cell culture system of claim 43, wherein the antibody is an antibody in accordance with claim 1.

45. The cell culture system of claim 43, wherein the antibody is expressed at the surface of an antigen presenting cell (APC).

10 46. A recombinant cell comprising an expression cassette comprising the coding sequence for an antibody that binds to the constant domain of a CAR polypeptide.

47. The cell of claim 46, wherein the antibody is a membrane-bound antibody.

48. The cell of claim 46, wherein the antibody is an antibody in according to any one of claims 1-4, wherein the antibody is a membrane-bound antibody.

15 49. The cell of claim 48, wherein the membrane-bound antibody comprises a 2D3 scFv coding sequence.

50. The cell of claim 49, wherein the membrane-bound antibody comprises one or more of (i) a GM-CSF leader peptide; (ii) 2D3 scFv coding sequence; (iii) a CD8 α linker sequence; (iv) a CD28 transmembrane domain; and/or (v) a CD3 ξ intracellular domain.

51. The cell of claim 46, wherein the antibody is a membrane-bound scFv antibody.

20 52. The cell of claim 51, wherein the membrane-bound antibody comprises one or more of (i) a GM-CSF leader peptide; (ii) the scFv coding sequence; (iii) a CD8 α linker sequence; (iv) a CD28 transmembrane domain; and/or (v) a CD3 ξ intracellular domain.

25 53. The cell of claim 51, wherein the membrane-bound antibody comprises (i) a GM-CSF leader peptide; (ii) the scFv coding sequence; (iii) a CD8 α linker sequence; (iv) a CD28 transmembrane domain; and (v) a CD3 ξ intracellular domain.

54. A kit comprising an antibody of claim 1 or a cell of claim 46 and at least a first agent for increasing proliferation of mammalian T-cells expressing a CAR.

55. The kit of claim 54, comprising inactivated aAPCs.

56. An isolated monoclonal antibody, wherein the antibody specifically binds to a CD19-specific scFv chimeric antigen receptor (CAR) polypeptide and wherein the antibody competes for binding of the polypeptide with the 136.20.1 monoclonal antibody.

5 57. The isolated antibody of claim 56, wherein the antibody comprises:

- (a) a first V_H CDR at least 80% identical to V_H CDR1 of 136.20.1 (SEQ ID NO: 2);
- (b) a second V_H CDR at least 80% identical to V_H CDR2 of 136.20.1 (SEQ ID NO: 3);
- (c) a third V_H CDR at least 80% identical to V_H CDR3 of 136.20.1 (SEQ ID NO: 4);
- 10 (d) a first V_L CDR at least 80% identical to V_L CDR1 of 136.20.1 (SEQ ID NO: 6);
- (e) a second V_L CDR at least 80% identical to V_L CDR2 of 136.20.1 (SEQ ID NO: 7); and
- (f) a third V_L CDR at least 80% identical to V_L CDR3 of 136.20.1 (SEQ ID NO: 8).

58. The isolated antibody of claim 57, wherein the antibody comprises:

- 15 (a) a first V_H CDR is identical to SEQ ID NO: 2;
- (b) a second V_H CDR is identical to SEQ ID NO: 3;
- (c) a third V_H CDR is identical to SEQ ID NO: 4;
- (d) a first V_L CDR is identical to SEQ ID NO: 6;
- (e) a second V_L CDR is identical to SEQ ID NO: 7; and
- 20 (f) a third V_L CDR is identical to SEQ ID NO: 8.

59. The isolated antibody of claim 56, wherein the antibody comprises a V_H domain at least about 80% identical to the V_H domain of 136.20.1 (SEQ ID NO: 1) and a V_L domain at least about 80% identical to the V_L domain of 136.20.1 (SEQ ID NO: 5).

60. The isolated antibody of claim 59, wherein the antibody comprises a V_H domain
25 identical to the V_H domain of 136.20.1 (SEQ ID NO: 1) and a V_L domain identical to the V_L domain of 136.20.1 (SEQ ID NO: 5).

61. The isolated antibody of claim 56, wherein the antibody comprises a V_H domain between 80% and 99% identical to the V_H domain of 136.20.1 (SEQ ID NO: 1) or a V_L domain between 80% and 99% identical to the V_L domain of 136.20.1 (SEQ ID NO: 5).

62. The isolated antibody of any one of claims 56-58, wherein the antibody is a membrane-bound antibody.
63. The isolated antibody of claim 62, wherein the membrane-bound antibody comprises one or more of (i) a GM-CSF leader peptide; (ii) 136.20.1 scFv coding sequence; (iii) a CD8 α linker sequence; (iv) a CD28 transmembrane domain; and/or (v) a CD3 ζ intracellular domain.
64. The isolated antibody of any one of claims 56-60, wherein the antibody is the 136.20.1 antibody.
65. The isolated antibody of any one of claims 56-60, wherein the antibody is recombinant.
66. The isolated antibody of any one of claims 56-60, wherein the antibody is an IgG, IgM, IgA or an antigen binding fragment thereof.
67. The isolated antibody of any one of claims 56-60, wherein the antibody is a Fab', a F(ab')₂, a F(ab')₃, a monovalent scFv, a bivalent scFv, or a single domain antibody.
68. The isolated antibody of any one of claims 56-60, wherein the antibody is a human, humanized antibody or de-immunized antibody.
69. The isolated antibody of any one of claims 56-60, wherein the antibody is conjugated to a toxin.
70. An isolated polynucleotide molecule comprising a nucleic acid sequence encoding an antibody of any one of claims 56-60.
71. A recombinant polypeptide comprising an antibody V_H domain comprising CDRs 1-3 of the V_H domain of 136.20.1 (SEQ ID NOs: 2, 3 and 4).
72. A recombinant polypeptide comprising an antibody V_L domain comprising CDRs 1-3 of the V_L domain of 136.20.1 (SEQ ID NOs: 6, 7 and 8).
73. An isolated polynucleotide molecule comprising a nucleic acid sequence encoding a polypeptide of claim 71 or 72.

74. A host cell comprising one or more polynucleotide molecule(s) encoding an antibody that specifically binds to the antigen recognition domain of a Chimeric Antigen Receptor (CAR) polypeptide.

75. The host cell of claim 74, wherein the CAR polypeptide is a CD-19-binding CAR polypeptide.

76. The host cell of claim 74, wherein the one or more polynucleotide molecule(s) encode an antibody of claims 56-60 or a recombinant polypeptide of claim 71 or 72.

77. The host cell of claim 74, wherein the host cell is a mammalian cell, a yeast cell, a bacterial cell, a ciliate cell or an insect cell.

78. The host cell of claim 77, wherein the host cell is an antigen presenting cell (APC).

79. The host cell of claim 77, wherein the host cell is an artificial antigen presenting cell (aAPC) or an activating and propagating cell (AAPC).

80. A method of manufacturing an antibody comprising:

- (a) expressing one or more polynucleotide molecule(s) encoding a V_L and V_H chain of an antibody of any one of claims 56-60 in a cell; and
- (b) purifying the antibody from the cell.

81. A method of manufacturing an antibody comprising:

- (a) expressing one or more polynucleotide molecule(s) encoding a V_L and V_H chain of an antibody of any one of claims 56-60 in an *in vitro* expression system; and
- (b) purifying the antibody therefrom.

82. A method for selecting a cell comprising a specific chimeric antigen receptor (CAR) polypeptide comprising:

- (a) contacting the cell with an antibody that specifically binds to the antigen recognition domain of a CAR polypeptide; and
- (b) selecting a cell comprising the specific CAR polypeptide based on binding of the antibody.

83. The method of claim 82, wherein the CAR polypeptide is a CD-19-binding CAR polypeptide.

84. The method of claim 82, wherein the antibody is an antibody in accordance with claim 56.

85. The method of claim 82, wherein the antibody is conjugated to a toxin and wherein the method is further defined as a method for selectively inhibiting or killing a cell
5 comprising a specific CAR polypeptide.

86. The method of claim 85, wherein the cell is *in vivo*.

87. The method of claim 82, wherein the method is further defined as a method for preventing or modifying the ability of a specific CAR to activate a T cell by contacting the cell with the antibody.

10 88. The method of claim 87, wherein the method is further defined as a method for controlling toxicity of a cell.

89. The method of claim 82, wherein the method is further defined as a method for purifying or enriching cells comprising the specific CAR polypeptide.

90. The method of claim 89, wherein the antibody comprises a reporter and the method
15 comprises selecting cells comprising the specific CAR polypeptide by fluorescence-activated cell sorting (FACS).

91. The method of claim 89, wherein selecting cells comprises selecting using paramagnetic beads.

92. The method of claim 89, wherein the antibody is bound to a support.

20 93. The method of claim 92, wherein the support is a bead, a surface, a column or an APC.

94. The method of claim 89, wherein purifying or enriching cells comprising the specific CAR comprises stimulating proliferation of the cells.


95. A cell culture system comprising mammalian T-cells expressing a specific CAR
25 polypeptide and an antibody that specifically binds to the antigen recognition domain of the CAR polypeptide.

96. The cell culture system of claim 95, wherein the CAR polypeptide is a CD-19-binding CAR polypeptide.
97. The cell culture system of claim 95, wherein the antibody is an antibody in accordance with claim 56.
- 5 98. The cell culture system of claim 95, wherein the antibody is expressed at the surface of an antigen presenting cell (APC).
99. A recombinant cell comprising an expression cassette comprising the coding sequence for an antibody that specifically binds to the antigen recognition domain of a CAR polypeptide.
- 10 100. The cell of claim 99, wherein the antibody binds to the antigen recognition domain of a CD-19-binding CAR polypeptide.
101. The cell of claim 100, wherein the antibody is an antibody in according to any one of claims 56-60.
102. The cell of claim 99, wherein the antibody is a membrane-bound antibody.
- 15 103. The cell of claim 102, wherein the membrane-bound antibody comprises a 136.20.1 scFv coding sequence.
104. The cell of claim 103, wherein the membrane-bound antibody comprises one or more of (i) a GM-CSF leader peptide; (ii) 136.20.1 scFv coding sequence; (iii) a CD8 α linker sequence; (iv) a CD28 transmembrane domain; and/or (v) a CD3 ξ intracellular domain.
- 20 105. The cell of claim 99, wherein the antibody is a membrane-bound scFv antibody.
106. The cell of claim 105, wherein the membrane-bound antibody comprises one or more of (i) a GM-CSF leader peptide; (ii) the scFv coding sequence; (iii) a CD8 α linker sequence; (iv) a CD28 transmembrane domain; and/or (v) a CD3 ξ intracellular domain.
- 25 107. The cell of claim 106, wherein the membrane-bound antibody comprises (i) a GM-CSF leader peptide; (ii) the scFv coding sequence; (iii) a CD8 α linker sequence; (iv) a CD28 transmembrane domain; and (v) a CD3 ξ intracellular domain.


108. A kit comprising an antibody of claim 56 or a cell of claim 99 and at least a first agent for increasing proliferation of mammalian T-cells expressing a specific CAR polypeptide.

109. The kit of claim 108, comprising inactivated aAPCs.

A



B



C




FIG. 1




FIG. 2




FIG. 3A




FIG. 3B




FIG. 3C





FIG. 4

A



B




FIG. 5




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 7




FIG. 8



FIG. 9




i) CAR⁺ T Cell (HcE) (40X)ii) CAR^{+/CD3} T Cell (Anti-CD19scFv) (40X)iii) CAR⁺ T Cell (Anti-CD19scFv) (40X)

FIG. 10




FIG. 11





FIG. 12

A



B




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17

A




FIG. 18




FIG. 19

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2014/039365**A. CLASSIFICATION OF SUBJECT MATTER****C07K 16/28(2006.01)i, C07K 16/46(2006.01)i, C12Q 1/04(2006.01)i, G01N 33/53(2006.01)i, C12N 5/07(2010.01)i, C12N 15/85(2006.01)i**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07K 16/28; C07H 21/04; A61K 39/00; C07K 16/46; C12Q 1/04; G01N 33/53; C12N 5/07; C12N 15/85

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean utility models and applications for utility models

Japanese utility models and applications for utility models

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKOMPASS(KIPO internal) & Keywords: chimeric antigen receptor, monoclonal antibodies, detection

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JENA et al., `Chimeric antigen receptor (CAR)-specific monoclonal antibody to detect CD19-specific T cells in clinical trials` PloS One, Vol.8, Issue 3, Article No.e57838 (internal pages 1-12) (01 March 2013) See abstract; pages 1-6 and 11; and figure S1.	1, 6-7, 9-15, 19-29 , 32-56, 62-70 , 74-109
A		2-5, 8, 16-18, 30-31 , 57-61, 71-73
A	WO 2012-079000 A1 (THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA) 14 June 2012 See abstract and claims 1-20.	1-109
A	DE OLIVEIRA et al., `A CD19/Fc fusion protein for detection of anti-CD19 chimeric antigen receptors` Journal of Translational Medicine, Vol.11, Article No.23 (internal pages 1-9) (29 January 2013) See abstract.	1-109



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

29 September 2014 (29.09.2014)

Date of mailing of the international search report

01 October 2014 (01.10.2014)

Name and mailing address of the ISA/KR

International Application Division
Korean Intellectual Property Office
189 Cheongsu-ro, Seo-gu, Daejeon Metropolitan City, 302-701,
Republic of Korea

Facsimile No. +82-42-472-7140

Authorized officer

KIM, Seung Beom

Telephone No. +82-42-481-3371



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2014/039365

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ZHENG et al., `Protein L: a novel reagent for the detection of chimeric antigen receptor (CAR) expression by flow cytometry` Journal of Translational Medicine, Vol.10, Article No.29 (internal pages 1-6) (2012) See abstract.	1-109
A	NCBI, PDB accession no. 1UZ6_F (18 October 2012) See the whole document.	1-109
A	NCBI, Genbank accession no. CBK46760.1 (13 May 2010) See the whole document.	1-109
A	NCBI, Genbank accession no. AAT73716.1 (20 October 2004) See the whole document.	1-109
A	NCBI, Genbank accession no. AAU14166.1 (16 March 2005) See the whole document.	1-109

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2014/039365Patent document
cited in search reportPublication
datePatent family
member(s)Publication
date

WO 2012-079000 A1

14/06/2012

AU 2011-338200 A1

04/07/2013

EP 2649086 A1

16/10/2013

JP 2014-507118 A

27/03/2014

KR 10-2013-0124521 A

14/11/2013

US 2013-0287748 A1

31/10/2013

US 2013-0288368 A1

31/10/2013

US 2013-0309258 A1

21/11/2013

US 2014-0106449 A1

17/04/2014