



(51) International Patent Classification:

A61K 31/164 (2006.01) A61K 9/00 (2006.01)
A61K 31/20 (2006.01) A61P 17/00 (2006.01)
A61K 31/201 (2006.01) A61P 29/00 (2006.01)
A61K 31/202 (2006.01)

(21) International Application Number:

PCT/IB2015/055885

(22) International Filing Date:

3 August 2015 (03.08.2015)

(25) Filing Language:

Italian

(26) Publication Language:

English

(30) Priority Data:

MI2014A001474 8 August 2014 (08.08.2014) IT

(72) Inventor; and

(71) Applicant : MIGLIACCIO, Raffaele [IT/IT]; Via S. Gottardo, 67, I-20900 Monza (MB) (IT).

(72) Inventors: SARDEI, Antonella; Via della Fiore, I-36034 Molina di Malo (VI) (IT). MIGLIACCIO, Carmela; Via S. Gottardo, 67, I-20090 Monza (MB) (IT).

(74) Agents: FINETTI, Claudia et al.; c/o Dragotti & Associati S.R.L., Via Nino Bixio, 7, I-20129 Milano (IT).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:


— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

(54) Title: MIXTURE OF FATTY ACIDS AND PALMITOYLETHANOLAMIDE FOR USE IN THE TREATMENT OF INFLAMMATORY AND ALLERGIC PATHOLOGIES

Figure 1



(57) Abstract: The present invention relates to a mixture containing up to two fatty acids selected from palmitic acid, oleic acid, stearic acid, linoleic acid, alpha-linolenic acid, gamma-linolenic acid, eicosapentaenoic acid, docosahexaenoic acid, azelaic acid and myristic acid and palmitoylethanolamide. In one embodiment of the present invention said mixture is characterized in that at least one of said up to two fatty acids is saturated. The present invention also relates to the use of the aforesaid mixture in the treatment of inflammatory and allergic pathologies.

TITLE:

Mixture of fatty acids and palmitoylethanolamide for use in the treatment of inflammatory and allergic pathologies.

The present invention relates to a mixture containing up to two fatty acids chosen from palmitic acid, oleic acid, stearic acid, linoleic acid, alpha-linolenic acid, gamma-linolenic acid, eicosapentaenoic acid, docosahexaenoic acid, azelaic acid and myristic acid and palmitoylethanolamide.

The present invention also relates to the use of the aforesaid mixture in the treatment of inflammatory and allergic pathologies.

PRIOR ART

Fatty acids are aliphatic carboxylic acids and represent the constituent ingredients of almost all complex lipids and vegetable and animal fats. These compounds can be classified on the basis of the length of the carbon chain and/or on the basis of the presence of double bonds in the aforesaid chain (saturated, unsaturated fatty acids).

In particular, some unsaturated fatty acids are considered to be essential because they are not synthesised by the organism but have to be supplied by the diet. Bergstrom, Samuelsson et al. (Nutrition classics. Biochimica et Biophysica Acta 90:207-10, 1964. The enzymatic formation of prostaglandin E2 from arachidonic acid. Prostaglandins and related factors 32) illustrated in 1964 the role and the biological effects of lipids in the inflammatory process and in other pathologies..

In 1979 the first phospholipid, phosphatidylinositol, was discovered as a factor in activating cellular response processes that is able to activate and control reactivity through messaging elements.

Lipids thus have different roles in the organism, including the roles of acting as chemical messengers that can cause changes to the role of the individual cell or

determine the actions that are able to modify the microenvironment, as in the case of response processes to inflammatory insults.

Essential fatty acids like linoleic acid and linolenic acid are known for being precursors of the arachidonic acid found in membrane phospholipids (such as, for example, cell membrane) and of many different types of eicosanoids, substances that are involved in the organism's inflammatory response, including hydroxyeicosatetraenes, prostanoids (prostaglandins, thromboxanes and prostacyclins), leukotrienes, lipoxins and resolvins, cell signalling that play an important role in pain, fevers, oedemas, blot clotting and more in general in inflammation.

As is known, inflammatory response is a multifactor physiological reaction characterised by the participation of different cells of the immune system such as, for example, mastocytes, macrophages, basophils and/or lymphocytes having different intervention times.

The first cell to intervene in the inflammatory process is the mastocyte, which is capable of responding to trigger the inflammatory process within microseconds. Activation of the mastocyte generates a series of reactions following the release of preformed mediators contained within the cytoplasm of the mastocyte; in rapid succession, the macrophages are contacted and activated.

The biological systems are based on receptor control: following stimulation of the pathogenic agent, the cells express specific receptors that are saturated by self-produced mediators, i.e. mediators that are formed by fatty acids constituting the membranes of the same cells. The expression of the receptors is the system through which the cells that are involved in the inflammatory process succeed in "transferring" to the microenvironment growth factors, interleukins, cytokines, etc. Saturation of these receptors first enables the degranulation of the mediators inside the cytoplasm of the cells involved in the inflammatory process, which are

mainly mastocytes, to be reduced and then modulated until the stimulus induced by the presence of the pathogenic agent is stopped.

However, this regulatory system is exhausted at the moment in which continuing to strip the cellular membranes of fatty acids causes the cell to suffer. In this condition, the receptors remain overexpressed and for the cell this is a degranulation signal of the mediators that trigger defence phenomena that are no longer necessary.

It is thus clear that if the receptor control did not occur, the cells would degranulate everything found in the cytoplasm with the resulting contact of other cells in the microenvironment and this would exasperate the system that by remaining active could become a source of harm and give rise to chronic and autoimmune inflammatory diseases such as, for example, rheumatoid arthritis, multiple sclerosis and systemic lupus erythematosus.

In these pathological conditions, it is thus very important for the organism to be able to control the hyperactivated inflammatory process by forming the receptor antagonist consisting of fatty acids removed from the membrane of the cells.

In order to respond to this need, diet strategies have been devised to reduce the synthesis of pro-inflammatory chemical mediators such as prostaglandins through the diminution of the consumption of vegetable oils and fatty meats and promoting the consumption of fish and certain particular oils such as linseed oil and hemp seed oil. This was done because it was thought that greater quantities of eicosapentaenoic acid and docosahexaenoic acid (omega-3) would be absorbed by the membrane phospholipids in place of arachidonic acid.

It is nevertheless known that these diet strategies are not sufficient to eradicate important inflammatory complaints such as, for example, rheumatoid arthritis, chronic ulcerous colitis, systemic lupus erythematosus, pelvic inflammatory disease or also atherosclerosis and pharmacological treatments must therefore be

resorted to.

The pharmacological treatments currently used for inflammatory complaints are corticosteroids (such as for example cortisone or the like) or NSAIDS (nonsteroidal anti-inflammatory drugs), which are drugs that act at different levels of the arachidonic acid cascade.

The function of this cascade is to trigger the immune response, maintaining it until the danger is eliminated and then moderate the immune response until it in turn becomes harmful (as occurs for example in chronic inflammations or in autoimmune diseases). In particular, corticosteroids prevent the cellular processes that lead to the synthesis of proinflammatory and immunostimulant substances and activate those cellular processes that lead to the synthesis of anti-inflammatory immunosuppressive substances so as to reduce the symptoms of the disease.

The side effects of synthetic corticosteroids mostly depend on the fact that in addition to the antiinflammatory/immunosuppressive effect, they interface with the organism's homeostatic systems and can thus cause: hypertension, water retention, hyperglycaemia, potassium loss, osteoporosis, muscular hypotrophia, capillary fragility, delayed healing of wounds, hyperlipidaemia, accumulation of adipose tissue in the face, neck and abdomen, gastroduodenal ulcers, increased blood clotting, haematological changes, euphoria and insomnia.

With prolonged treatment, these drugs tend to inhibit the production of similar natural hormones by the adrenal glands, thus causing adrenal insufficiency, which manifests itself with also serious consequences above all when treatment is suspended. Further, the protracted use of corticosteroids is linked to their immunosuppressive action that increases susceptibility to infection.

NSAIDS by contrast interfere at a different level with the arachidonic acid cascade, inhibiting the cyclooxygenase COX1 and 2 involved in the inflammatory

processes. Some of the most common side effects affect the gastroenteric apparatus, and in particular the stomach: pain or burning or nausea, ulceration of the gastric mucose with possible bleeding; skin reactions (rashes, itches) in predisposed subjects.

Palmitoylethanolamide is an endogenous compound belonging to the class of fatty acid amides and is known chemically as N-(2-Hydroxyethyl)hexadecanamide. This compound is a key element in regulating paths connected to the inflammation process, in particular to the process of degranulation downregulation of the mastocyte and also of itch and pain-based processes.

The need is therefore felt to identify one or more compounds for treating the inflammation that is able to block effectively hyperactivation of the inflammatory process, reducing the side effects associated with traditional treatments.

DESCRIPTON

It has been surprisingly found that a mixture of specific fatty acids in association with palmitoylethanolamide is able to treat the inflammatory process efficiently, with improved control of the regulatory system and with an absence of side effects through the biological control of the cells involved in the inflammatory process.

One object of the present invention is accordingly a mixture containing up to two fatty acids containing between 8 and 24 carbon atoms; said fatty acids can be saturated, unsaturated or be mixtures thereof, in association with palmitoylethanolamide.

Said fatty acids are preferably selected from palmitic acid, oleic acid, stearic acid, linoleic acid, alpha-linolenic acid, gamma-linolenic acid, eicosapentaenoic acid, docosahexaenoic acid, azelaic acid, myristic acid or a mixture thereof.

According to one embodiment of the present invention, the mixture is

characterized in that at least one of said up to two fatty acids is saturated.

Preferably, said saturated fatty acid is selected from stearic acid, palmitic acid, azelaic acid or myristic acid, more preferably, is selected from palmitic acid.

According to a further embodiment of the present invention, the mixture contains a fatty acid selected from the aforesaid fatty acids in association with palmitoylethanolamide.

Preferably, said fatty acid is present in the aforesaid mixture in a weight ratio comprised between 15% and 55%, more preferably between 20% and 53%.

In the aforesaid mixture, the palmitoylethanolamide is contained in a weight quantity comprised between 45% and 85%, preferably between 47% and 80%, of the total weight of the mixture.

Preferably, the aforesaid mixture contains palmitic acid in association with palmitoylethanolamide.

In the mixture of the present invention, the palmitic acid is contained in a weight quantity comprised between 20% and 52%, preferably between 30% and 48%, more preferably about 45%, of the total weight of the mixture.

Preferably, the aforesaid mixture contains the oleic acid in association with palmitoylethanolamide.

In the mixture of the present invention, the oleic acid is contained in a weight quantity comprised between 35% and 48%, preferably between 38% and 45%, more preferably about 40%, of the total weight of the mixture.

Preferably, the aforesaid mixture contains the stearic acid in association with palmitoylethanolamide.

In the mixture of the present invention, the stearic acid is contained in a weight quantity comprised between 30% and 55%, preferably between 38% and 53%, more preferably about 50%, of the total weight of the mixture.

Preferably, the aforesaid mixture contains the linoleic acid in association with

palmitoylethanolamide.

In the mixture of the present invention, the linoleic acid is contained in a weight quantity comprised between 35% and 55%, preferably between 40% and 50%, more preferably about 40%, of the total weight of the mixture.

Preferably, the aforesaid mixture contains the alpha-linolenic acid in association with palmitoylethanolamide.

In the mixture of the present invention, the alpha-linolenic acid is contained in a weight quantity comprised between 35% and 48%, preferably between 38% and 42%, more preferably about 40%, of the total weight of the mixture.

Preferably, the aforesaid mixture contains the gamma-linolenic acid in association with palmitoylethanolamide.

In the mixture of the present invention, the gamma-linolenic acid is contained in a weight quantity comprised between 30% and 40%, preferably between 32% and 38%, more preferably about 35%, of the total weight of the mixture.

Preferably, the aforesaid mixture contains the eicosapentaenoic acid in association with palmitoylethanolamide.

In the mixture of the present invention, the eicosapentaenoic acid is contained in a weight quantity comprised between 30% and 55%, preferably between 36% and 52%, more preferably about 50%, of the total weight of the mixture.

Preferably, the aforesaid mixture contains the docosahexaenoic acid in association with palmitoylethanolamide.

In the mixture of the present invention, the docosahexaenoic acid is contained in a weight quantity comprised between 30% and 55%, preferably between 30% and 40%, more preferably about 35%, of the total weight of the mixture.

Preferably, the aforesaid mixture contains the azelaic acid in association with palmitoylethanolamide.

In the mixture of the present invention, the azelaic acid is contained in a weight

quantity comprised between 15% and 45%, preferably between 20% and 30%, more preferably about 25%, of the total weight of the mixture.

Preferably, the aforesaid mixture contains the myristic acid in association with palmitoylethanolamide.

In the mixture of the present invention, the myristic acid is contained in a weight quantity comprised between 15% and 35%, preferably between 20% and 30%, more preferably about 25%, of the total weight of the mixture.

According to another particularly preferred embodiment, the mixture contains a fatty acid selected from eicosapentaenoic acid and docosahexaenoic acid in association with palmitoylethanolamide.

According to another embodiment of the present invention, the mixture contains two fatty acids selected from the above fatty acids in association with palmitoylethanolamide.

Preferably, the two fatty acids are present in the aforesaid mixture in a weight ratio to one another comprised between 0.25 and 2.33, more preferably between 0.33 and 1.5, still more preferably about 1.

Furthermore, in the aforesaid mixture the two fatty acids are present in a weight ratio to the palmitoylethanolamide comprised between 0.25 and 1.5, preferably between 0.43 and 1, more preferably about 0.67.

When two fatty acids are present in the mixture of the present invention, the palmitoylethanolamide is present in a weight quantity comprised between 25% and 70%, preferably between 35% and 60%, of the total weight of the mixture.

The two fatty acids are present in the mixture of the invention in a weight quantity comprised between 30% and 75%, preferably between 40% and 65%, of the total weight of the mixture.

According to one embodiment, the mixture of the invention contains palmitic acid, preferably in a weight quantity comprised between 10% and 48%, more

preferably between 25% and 35%, still more preferably about 30%, of the total weight of the mixture.

Alternatively, the mixture of the invention can contain oleic acid, preferably in a weight quantity comprised between 15% and 38%, more preferably between 18% and 25%, still more preferably about 20%, of the total weight of the mixture.

According to another embodiment of the invention, the aforesaid mixture can contain stearic acid, preferably in a weight quantity comprised between 20% and 45%, more preferably comprised between 25% and 35%, still more preferably about 30%, of the total weight of the mixture.

Otherwise, the linoleic acid can be contained in the mixture of the present invention, preferably in a weight quantity comprised between 15% and 40%, more preferably between 20% and 38%, still more preferably about 35%, of the total weight of the mixture.

According to another embodiment of the present invention, the mixture contains the alpha-linolenic acid, preferably in a weight quantity comprised between 25% and 38%, more preferably comprised between 30% and 36%, still more preferably about 35%, of the total weight of the mixture.

According to the present invention, the mixture can contain gamma-linolenic acid, preferably in a weight quantity comprised between 15% and 35%, more preferably between 20% and 34%, still more preferably about 30%, of the total weight of the mixture.

According to another embodiment of the invention, the aforesaid mixture can contain eicosapentaenoic acid, preferably in a weight quantity comprised between 10% and 35%, more preferably comprised between 20% and 30%, still more preferably about 25%, of the total weight of the mixture.

Alternatively, the docosahexaenoic acid can be contained in the mixture, preferably in a weight quantity comprised between 8% and 35%, more preferably

between 18% and 30%, still more preferably about 20%, of the total weight of the mixture.

According to the present invention, the aforesaid mixture can contain azelaic acid, preferably in a weight quantity comprised between 5% and 35%, more preferably between 10% and 20%, still more preferably about 15%, of the total weight of the mixture.

According to another embodiment of the present invention, the mixture contains the myristic acid, preferably in a weight quantity comprised between 15% and 30%, more preferably comprised between 15% and 20%, still more preferably about 18%, of the total weight of the mixture.

According to another particularly preferred embodiment, the mixture of the invention contains two fatty acids selected from eicosapentaenoic acid and docosahexaenoic acid in association with palmitoylethanolamide.

In the aforesaid mixture, the eicosapentaenoic acid is present in a weight quantity comprised between 20% and 40%, preferably about 30%.

The docosahexaenoic acid is present in a quantity comprised between 25% and 50%, preferably about 30%.

Palmitoylethanolamide is present in the aforesaid preferred mixture in a quantity comprised between 30% and 50%, preferably about 40%.

It has been surprisingly noticed that by making up a pool of the aforesaid fatty acids, in particular one or two fatty acids according to the present invention, it is possible to determine the control of cells involved in the inflammatory processes such as mastocytes, macrophages, basophils and lymphocytes.

This system for administering the fatty acids with palmitoylethanolamide has two important advantages:

- 1) a more rapid and reactive response to the hyperstimulation induced by the inflammatory agent: the cell does not have to be deprived of the membrane lipids

with consequent temporal "dispersal";

2) significant energy saving: the cell does not have to use energy to recover the lipids from the membranes and to replace the lipids.

It has in fact been observed that by supplying the pool of fatty acids according to the present invention, improved control of the inflammatory process is obtained as well as rapid restoration of physiological conditions and of normal conditions of the microenvironment affected compared with what is known in the prior art.

A further object of the present invention is a pharmaceutical composition comprising the aforesaid mixture and at least one pharmaceutically acceptable excipient.

The term "pharmaceutically acceptable excipient" refers to a substance that is devoid of any pharmacological effect of its own and does not produce adverse reactions when administered to a mammal, preferably to a human being. Pharmaceutically acceptable excipients are well known in the prior art and are disclosed, for example in the *Handbook of Pharmaceutical Excipients, sixth edition 2009*, which is included here for reference.

Excipients are normally classified according to the function that they have in the final pharmaceutical composition. Preferably, suitable excipients according to the present invention are for example diluent, adsorbent, glidant, binder, lubricant, surfactant, disintegrating, preservatives, antioxidant or mixtures thereof.

According to one embodiment of the present invention, the aforesaid composition comprises a mixture containing up to two fatty acids, or containing one or two fatty acids, selected from palmitic acid, oleic acid, stearic acid, linoleic acid, alpha-linolenic acid, gamma-linolenic acid, eicosapentaenoic acid, docosahexaenoic acid, azelaic acid, myristic acid, palmitoylethanolamide and at least one pharmaceutically acceptable excipient.

When the palmitic acid is contained in the composition of the invention it is

present in a quantity in weight comprised between 10% and 48%, more preferably comprised between 20% and 30%, still more preferably about 25%.

When the oleic acid is contained in the composition of the invention it is present in a quantity in weight comprised between 15% and 38%, more preferably comprised between 15% and 25%, still more preferably about 20%.

When the stearic acid is contained in the composition of the invention it is present in a quantity in weight comprised between 20% and 45%, more preferably comprised between 25% and 35%, still more preferably about 30%.

When the linoleic acid is contained in the composition of the invention it is present in a quantity in weight comprised between 15% and 40%, more preferably comprised between 25% and 35%, still more preferably about 30%.

When the alpha-linolenic acid is contained in the composition of the invention, it is present in a quantity in weight comprised between 25% and 38%, more preferably comprised between 28% and 32%, still more preferably about 30%.

When the gamma-linolenic acid is contained in the composition of the invention, it is present in a quantity in weight comprised between 15% and 30%, more preferably comprised between 20% and 30%, still more preferably about 25%.

When the eicosapentaenoic acid is contained in the composition of the invention it is present in a quantity in weight comprised between 10% and 35%, more preferably comprised between 15% and 25%, still more preferably about 20%.

When the docosahexaenoic acid is contained in the composition of the invention, it is present in a weight quantity comprised between 8% and 35%, more preferably comprised between 15% and 25%, still more preferably about 20%.

When the azelaic acid is contained in the composition of the invention it is present in a quantity in weight comprised between 5% and 35%, more preferably comprised between 10% and 20%, still more preferably about 15%.

When the myristic acid is contained in the composition of the invention it is

present in a quantity in weight comprised between 15% and 30%, more preferably comprised between 15% and 25%, still more preferably about 20%.

The composition of the invention further contains a quantity of palmitoylethanolamide comprised between 35% and 55%, preferably comprised between 35% and 45%, more preferably about 40%.

The aforesaid percentage quantities are weight quantities expressed as a percentage the total weight of the composition.

The composition of the present invention can be formulated in a form that is suitable for oral, topical, rectal, vaginal, ophthalmic or parenteral administration.

According to one preferred embodiment of the present invention, said oral form is selected from a tablet, capsule, granule, oily perl, solution, suspension, aerosol, more preferably is selected from a capsule or tablet.

The capsule can be a soft gelatine capsule, a hard capsule or a capsule containing granules.

According to another preferred embodiment of the present invention, said topical form is selected from cream, ointment, gel, salve, solution, suspension, eyewash, drops, spray, or powder, more preferably it is selected from cream, gel, spray, ointment, drops, eyewash.

According to another preferred embodiment of the present invention, said form that is suitable for vaginal administration is a pessary, cannula, douche or cream.

According to another preferred embodiment of the present invention, said form that is suitable for rectal administration is a suppository, an enema or a cream.

According to another preferred embodiment of the present invention, said form that is suitable for ophthalmic administration is an eyewash, a bath or a cream.

According to another preferred embodiment of the present invention, said form that is suitable for parenteral administration is selected from a water buffer solution or oily suspension.

The mixture of fatty acids of the present invention is preferably contained in the aforesaid formulations in a weight quantity that varies from 3% to 60%, preferably from 5% to 35%, of the total weight of the formulation.

According to one embodiment of the present invention, the composition containing a mixture of a fatty acid and palmitoylethanolamide is preferably formulated in topical form.

According to one embodiment of the present invention, the composition containing a mixture of one or two fatty acids and palmitoylethanolamide is preferably formulated in topical or oral form.

According to one embodiment of the present invention, the composition containing a mixture of two fatty acids and palmitoylethanolamide is preferably formulated in oral form, more preferably in the form of a tablet, of a suspension, in rectal form or in topical form, more preferably in the form of a cream, gel, ointment, drops, eyewash, spray, solution.

According to the invention, the composition of the present invention can be administered to animals and humans, defined as adults and as "paediatric population", wherein with the term "paediatric population" is indicated the part of the population from birth to eighteen years of age.

A further object of the present invention is to use a mixture containing up to two fatty acids selected from palmitic acid, oleic acid, stearic acid, linoleic acid, alpha-linolenic acid, gamma-linolenic acid, eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA), azelaic acid, myristic acid and palmitoylethanolamide and/or a composition containing said mixture in the treatment of inflammatory and allergic pathologies.

Said inflammatory and allergic pathologies are acute or chronic and are selected from dermatological pathologies such as, for example, atopic dermatitis, dermatomyositis, scleroderma, psoriasis, polymyositis, pemphigus, pemphigus

epidermolysis bullosa; ophthalmic pathologies such as, for example, Sjorgen's syndrome, sympathetic ophthalmia, uveitis, uveoretinitis; mucosal pathologies such as inflammation of the gastrointestinal mucous membranes (Crohn's disease), inflammation of the oral and genital mucous membrane; joint and connective tissue pathologies such as, for example, rheumatoid arthritis, psoriatic arthritis, arthritis from lupus erythematosus, discoid and systemic lupus; chronic pathological inflammations such as for example from chronic solar dermatitis, asthma and intestinal and pulmonary fibrosis, chronic arthritis, degenerative pathologies of the peripheral nervous system (PNS) and of the central nervous system (CNS) such as, for example, multiple sclerosis, neurodegenerative pathologies, not only autoimmune inflammatory processes connected to CNS like Parkinson's disease, tinnitus, senile dementia, bacterial meningitis, HIV infections and trauma injuries, and PNS pathologies such as, for example, radiculopathy of inflammatory origin; pathologies of the peripheral and central system where the inflammatory processes follow the first insult of ischaemic origin, such as, for example, compression and traumatic neuropathies, cerebral strokes and traumatic brain injury; heart disease arising from perfusion phenomena as a consequence of ischaemic damage; inflammatory pathologies associated with fibrosis, such as, for example, allergic conjunctivitis, giant papillary conjunctivitis, dietary allergies, scarring anomalies such as, for example, hypertrophic scars, keloids and cicatricial pemphigoid; pathologies in which renal function is altered following renal inflammation.

Preferred pathologies according to the present invention are the inflammatory pathologies linked to hyperactivity of the cells involved in the inflammatory processes on a neurogenic basis, more preferably they are neuroinflammatory pathologies of the central and peripheral nervous system, such as, for example, tinnitus, polyneuropathies, myasthenia, myopathies.

The composition of the present invention is preferably administered daily, from one to four doses a day, in which said dose contains from 0.1 to 50 mg of composition/kg of body weight of patient, more preferably from 0.5 to 20 mg/kg of body weight of patient, for at least 4 weeks.

EXAMPLES

The following mixtures according to the present invention were prepared.

Example 1: Formulation in topical cream

Eicosapentaenoic acid	40%
Palmitoylethanolamide	60%

Example 2: Formulation in vaginal pessary

Docosahexaenoic acid	45%
Palmitoylethanolamide	55%

Example 3: Formulation in otological drops

Eicosapentaenoic acid	30%
Docosahexaenoic acid	30%
Palmitoylethanolamide	40%

Example 4: Formulation in cream for periophthalmic use

Eicosapentaenoic acid	35%
Docosahexaenoic acid	30%
Palmitoylethanolamide	35%

Example 5

Evaluation of the anti-inflammatory activity in vitro

A mixture of fatty acids containing: palmitoylethanolamide (33%), linoleic acid (33%) and palmitic acid (33%) (defined in the tables as Fatty Acids) was tested in vitro on a human cell line THP1 (monocyte-macrophage) previously sensitized with LPS (1 µg/ml), a known inflammatory agent, to evaluate the activity of the mixture to reduce the release of pro-inflammatory mediators (IL-6 and IL-8).

Results

All doses of the mixture analyzed have proved to be able to inhibit the release of both cytokines checked IL-6 and IL-8 (Figure 1 and 2, Table 1 and 2). The tests performed have identified two effective doses 0.06 mg/ml and 0.03 mg/ml: at these concentrations in fact the anti-inflammatory effectiveness of fatty acids mixture determines a strong reduction of IL-6 release equal to 100% and 58% respectively, and at 0.06 mg/ml it causes a IL-8 release inhibition equal to 42.7%.

Table 1.

Acidi Grassi+LPS	pg/ml IL8	% Inhib. IL8
0,06 mg/ml	323,70	41,06
0,03 mg/ml	488,02	11,13
0,01 mg/ml	526,12	4,19
Quality Control	369,76	32,67
Pos. Control	549	0

Table 2.

Acidi Grassi+LPS	pg/ml IL6	% Inhib. IL6
0,06 mg/ml	0	100
0,03 mg/ml	15,74	58
0,01 mg/ml	39,62	0
Quality Control	0	100
Pos. Control	38	0

CLAIMS

1. Mixture containing up to two fatty acids selected from palmitic acid, oleic acid, stearic acid, linoleic acid, alpha-linolenic acid, gamma-linolenic acid, eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA), azelaic acid and myristic acid and palmitoylethanolamide.
2. Mixture according to claim 1, characterized in that at least one of said up to two fatty acids is saturated.
3. Mixture according to claim 1/2, characterised in that it comprises a fatty acid and palmitoylethanolamide.
4. Mixture according to claim 1/2, characterised in that it comprises two fatty acids and palmitoylethanolamide.
5. Mixture according to claim 3, characterised in that said fatty acid is present in the aforesaid mixture in a quantity in weight comprised between 15% and 55%, preferably between 20% and 53%, of the total weight of the mixture.
6. Mixture according to claim 3, characterised in that said palmitoylethanolamide is present in the aforesaid mixture in a quantity in weight comprised between 45% and 85%, preferably between 47% and 80%, of the total weight of the mixture.
7. Mixture according to claim 4, characterised in that the two fatty acids are present in the aforesaid mixture in a weight ratio from one to another comprised between 0.25 and 2.33, preferably between 0.33 and 1.5, more preferably about 1.
8. Mixture according to claim 4, characterised in that the two fatty acids are present in a weight ratio with respect to the palmitoylethanolamide comprised between 0.25 and 1.5, preferably between 0.43 and 1, more preferably about 0.67.
9. Mixture according to claim 4, characterised in that the palmitoylethanolamide is present in said mixture in a weight quantity comprised between 25% and 70%, preferably between 35% and 60%, more preferably about

40%, of the total weight of the mixture.

10. Mixture according to claim 4, characterised in that the two fatty acids are present in said mixture in a weight quantity comprised between 30% and 75%, preferably between 40% and 65%, more preferably about 60%, of the total weight of the mixture.

11. Composition comprising a mixture according to any of preceding claims and at least one pharmaceutically acceptable excipient.

12. Composition according to claim 11, formulated in an oral, topical, rectal, vaginal, ophthalmic or parenteral form.

13. Composition according to claim 12, characterised in that said oral form is selected from a tablet, capsule, granule, oily perl, solution, suspension, aerosol, preferably said oral form is a tablet, capsule or solution.

14. Composition according to claim 12, characterised in that said topical form is selected from cream, ointment, gel, salve, solution, suspension, eyewash, spray, drops or powder, preferably said topical form is a cream, gel, spray, otological drops or ointment.

15. Composition according to claim 12, characterised in that said parenteral form is selected from a water buffer solution or oily suspension, preferably said parental form is an oily suspension.

16. Composition according to claim 12, characterised in that said rectal form is a suppository, a cream or a enema.

17. Composition according to claim 12, characterised in that said vaginal form is a pessary, a cream, a vaginal douche or it is administered with a cannula.

18. Composition according to claim 12, characterised in that said ophthalmic form is an eyewash, a cream or a bath.

19. Composition according to any one of claims 11-18, characterised in that the mixture according to anyone of claims 1-9 is present in said composition in a

weight quantity comprised between 15% and 70%, preferably comprised between 20% and 65% of the total weight of the composition.

20. Mixture and/or composition according to any one of claims 1-19, for use in the treatment of inflammatory and allergic pathologies, preferably in the treatment of neuroinflammatory pathologies of the peripheral nervous system.

21. Composition according to claim 11, characterised in that it is administered daily, from one to four doses a day, wherein said dose contains from 0.1 to 50 mg of composition/kg of patient body weight, preferably from 0.5 to 20 mg/kg of patient body weight, for at least 4 weeks.

Figure 1





Figure 2



INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2015/055885

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K31/164 A61K31/20 A61K31/201 A61K31/202 A61K9/00
A61P17/00 A61P29/00

ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 679 667 A (DELLA VALLE FRANCESCO [IT] ET AL) 21 October 1997 (1997-10-21) the whole document example 9	1-3,5, 11-14, 16,17, 19-21
X	WO 2008/075978 A2 (SEPEREX NUTRITIONALS LTD [NZ]; WILLIAMS CHARLES EDWARD [NZ]; SANSOM AN) 26 June 2008 (2008-06-26) the whole document claims 1-43; table 2	1,3-5,7, 10-17, 19-21
X	EP 1 844 784 A1 (EPITECH GROUP S R L [IT]) 17 October 2007 (2007-10-17) claim 19; examples 1,5 the whole document ----- -/-	1-3,5,6, 11-13, 19-21



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

2 October 2015

Date of mailing of the international search report

09/10/2015

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Jakobs, Andreas

INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2015/055885

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Anonymous: "12 Current Essentials - Physiogel AI Cream Very Sensitive", 26 May 2012 (2012-05-26), XP055179372, Retrieved from the Internet: URL:https://verysensitive.wordpress.com/2012/05/26/12-current-essentials-physiogel-ai-cream/ [retrieved on 2015-03-25] the whole document	1-4, 11-16, 19-21
X,P	----- WO 2014/135529 A1 (AGAIN LIFE ITALIA SRL [IT]) 12 September 2014 (2014-09-12) the whole document page 14, paragraph 5-6 example 10	1-21
A	----- WO 01/24645 A1 (NESTLE SA [CH]; BERGER ALVIN [CH]; CROZIER GAYLE [CH]) 12 April 2001 (2001-04-12) the whole document	1-20
A	----- WO 02/053108 A2 (UNILEVER PLC [GB]; UNILEVER NV [NL]; LEVER HINDUSTAN LTD [IN]) 11 July 2002 (2002-07-11) the whole document	1-20
A	----- DE 102 17 131 A1 (KUHS KOSMETIK GMBH & CO KG [DE]) 13 February 2003 (2003-02-13) the whole document -----	1-20

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2015/055885

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5679667	A	21-10-1997	NONE
WO 2008075978	A2	26-06-2008	CA 2710054 A1 26-06-2008 CN 101641089 A 03-02-2010 CN 103638003 A 19-03-2014 DE 112007003136 T5 19-11-2009 GB 2458599 A 30-09-2009 HK 1136218 A1 03-06-2011 KR 20090103918 A 01-10-2009 NZ 552238 A 31-07-2009 TW 200833318 A 16-08-2008 WO 2008075978 A2 26-06-2008
EP 1844784	A1	17-10-2007	AT 464890 T 15-05-2010 CA 2582027 A1 28-09-2007 EP 1844784 A1 17-10-2007 ES 2344920 T3 09-09-2010 HK 1110016 A1 23-07-2010 JP 5253750 B2 31-07-2013 JP 2007262068 A 11-10-2007 US 2007270496 A1 22-11-2007 US 2011288037 A1 24-11-2011
WO 2014135529	A1	12-09-2014	AU 2014224710 A1 10-09-2015 WO 2014135529 A1 12-09-2014
WO 0124645	A1	12-04-2001	AT 282330 T 15-12-2004 AU 7518700 A 10-05-2001 CA 2386362 A1 12-04-2001 DE 60016007 D1 23-12-2004 DE 60016007 T2 02-03-2006 EP 1221867 A1 17-07-2002 ES 2234665 T3 01-07-2005 PT 1221867 E 29-04-2005 US 8323680 B1 04-12-2012 WO 0124645 A1 12-04-2001
WO 02053108	A2	11-07-2002	AU 2002229642 B2 06-01-2005 CA 2431539 A1 11-07-2002 CN 1482899 A 17-03-2004 EP 1349538 A2 08-10-2003 JP 3792651 B2 05-07-2006 JP 2004522728 A 29-07-2004 KR 20030074687 A 19-09-2003 KR 20080067386 A 18-07-2008 MX PA03005704 A 06-10-2003 US 2002143059 A1 03-10-2002 WO 02053108 A2 11-07-2002 ZA 200303936 A 21-05-2004
DE 10217131	A1	13-02-2003	NONE