



(51) International Patent Classification:
C12Q 1/68 (2006.01)

(21) International Application Number:
PCT/DK2010/050029

(22) International Filing Date:
2 February 2010 (02.02.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2009 00156 2 February 2009 (02.02.2009) DK
PA 2009 01038 17 September 2009 (17.09.2009) DK

(71) Applicant (for all designated States except US): **EX-IQON A/S** [DK/DK]; Skelstedet 16, DK-2950 Vedbaek (DK).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **BUSK, Peter, Kamp** [DK/DK]; Mars Allé 4, DK-2860 Søborg (DK).

(74) Common Representative: **EXIQON A/S**; Jan Skouv, Skelstedet 16, DK-2950 Vedbaek (DK).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: METHOD FOR QUANTIFICATION OF SMALL RNA SPECIES

(57) Abstract: The present invention relates to a method for amplification and quantification of microRNA molecules using quantitative reverse transcription polymerase chain reaction (q RT- PC R) technology. The method comprise the steps of (a) producing cDNA molecules complementary to microRNAs in a sample using polyadenylation and primer extension by reverse transcription, and (b) amplification and quantification of the cDNAs by qPCR using microRNA specific primer sets of forward and reverse primers containing LNA monomers.



WO 2010/085966 A2

METHOD FOR QUANTIFICATION OF SMALL RNA SPECIES

FIELD OF THE INVENTION

The present invention relates to a method for amplification and quantification of small non-coding RNA molecules using quantitative reverse transcription polymerase chain reaction (qRT-PCR) technology.

BACKGROUND OF THE INVENTION

MicroRNAs are an abundant class of approximately 22-nucleotide-noncoding RNAs, which play important regulatory roles in animal, plant and virus development. The awareness of microRNAs was initiated almost 15 years ago by the discovery of *lin-4*, which encode a small RNA involved in the timing and progression of the nematode in *C. elegans* life cycle and larval development (Lee et al. 1993 Cell 75:843-854, Wightman et al. 1993 Cell 75:855-862), but it was only recently recognized that microRNAs form a major class of ribo-regulators that have broad regulatory functions in animals (Lagos-Quintana et al. 2001 Science 294:853-858, Lau et al. 2001 Science 294:858-862, Lee and Ambros. 2001 Science 294:862-864). Since then, a revolution in the study of microRNAs have taken place, and today the miRBase database version 12.0 (<http://microrna.sanger.ac.uk/>) include 866 human microRNAs and the PubMed database (<http://www.ncbi.nlm.nih.gov/pubmed/>) encompass 3900 microRNA related articles, reflecting the interest and importance of microRNAs.

MicroRNAs are involved in the regulation of gene expression at the posttranscriptional level by degrading or blocking translation of messenger RNA targets, and it has been speculated that approximately 30 % of the human genome could be regulated by microRNAs. The importance of microRNAs is also obvious due to their involvement in various cellular processes including development, growth and proliferation, apoptosis, differentiation, and various human diseases (<http://www.mir2disease.org/>) such as cancer and diabetes.

The importance of microRNAs in cancer is highlighted in a recent article (Barbarotto et al 2008 Int. J. Cancer. 122:969-977), which summarizes the main paradigms for the miRNA involvement in human cancers: Thus, "(i) miRNAs are altered in every type of analyzed human cancer; (ii) miRNAs act as oncogenes and tumor suppressors; (iii) miRNAs alterations may cause cancer predisposition; (iv) miRNAs profiling is a new diagnostic tool for cancer patients and (v) miRNA profiling represents prognostic tools for cancer patients.". Accordingly, methods for expression profiling and quantification of microRNAs in cells and body-fluids from

cancer patients are of great importance. To address this requirement, the present invention describes the development of a new robust and reliable qRT-PCR assay for microRNA measurements.

Quantification of microRNAs by qRT-PCR procedures is very challenging due to the small size
5 of microRNAs of only 21 to 25 nucleotides, which is the size of primers normally used for PCR. Solutions to this problem have been published in Raymond et al. RNA. 2005 Nov;11(11):1737-44, Gilad et al. PLoS ONE. 2008 Sep 5;3(9):e3148 and Sharbati-Tehrani et al. BMC Molecular Biology. 2008, 9:34. Raymond et al. describes a qRT-PCR assay that involves a gene-specific reverse transcription step followed by a SYBR[®] green qPCR step
10 using a gene-specific forward primer containing locked nucleic acid (LNA) molecules and a universal reverse primer. Gilad et al. reports a qRT-PCR assay that involves a polyadenylation step, an unspecific reverse transcription step, and a qPCR step involving a gene-specific forward primer, a gene-specific TaqMan primer and a universal reverse primer. Sharbati-Tehrani et al. developed a qRT-PCR assay that involves a gene-specific reverse transcription
15 step followed by a SYBR[®] green qPCR step using a gene-specific forward primer and 2 universal primers.

However, the existing techniques for quantification of microRNAs by qRT-PCR do not fulfil the present need for microRNA assays, which requires high specificity that allows discrimination between closely related microRNAs, high sensitivity, low background and a relatively simple
20 procedure.

The present invention is characterised by only one reverse transcription reaction for all microRNAs in a sample and furthermore provides an extremely sensitive PCR method with an unmatched specificity that can be used for accurate quantification of small RNA molecules such as microRNAs.

25

SUMMARY OF THE INVENTION

Establishing and understanding the microRNA dysregulation patterns associated with various human diseases such as cancers, call for novel, improved technologies for detection and quantification of microRNAs in human cells and body fluids. The present invention introduces
30 a new highly sensitive and specific assay for this purpose.

In one aspect, the present invention provides a procedure for amplifying and quantifying microRNA molecules in a sample: In the first step of the procedure complementary DNAs (cDNAs) of the microRNAs in a sample are produced by the concerted action of two enzymes in a single-tube reaction. Firstly, poly-A tails are added to the 3'-terminus of the microRNAs using a poly(A) polymerase, and secondly an extension primer is hybridized to the poly-A tail and the cDNAs are produced by a reverse transcriptase using the microRNA as template. The first step is unspecific and produces cDNAs of all microRNAs present in a sample. In the second step of the procedure specific cDNAs are amplified and quantified in a qPCR reaction using microRNA specific primer sets of forward and reverse primers containing LNA monomers.

In another aspect, the invention provides oligo nucleotide primers listed in Table 18 (SEQ ID NO 1 to SEQ ID NO 128).

The primers of the invention can be used for detecting mammalian microRNAs using the method of the invention.

In another aspect, the invention provides kits for detection of mammalian microRNAs, the kits comprising a universal extension primer and microRNA-specific forward and reverse primers sets for quantification of at least one microRNA, a subset of microRNAs or all known microRNAs.

The present invention is useful for reliable and specific quantitative microRNA assays, including assays for diagnosing and prognosing diseases such as cancer using single assays or high through-put applications on robot platforms. RNA containing samples extracted from various cell types from living organisms, such as mammals and plants and including virus infected cells may be analysed using the method of the invention.

Although the present invention mainly aims at providing a method for quantification of microRNAs the method can be used for detection and/or quantification of all types of RNA in particular all types of small noncoding RNAs.

FIGURES

FIGURE 1 shows the steps involved in specific qRT-PCR of the present invention. To illustrate the principle the qRT-PCR of a microRNA serves as an example, the RNA to be analysed by the method may as well be any other small RNA molecule or even a mRNA. Step 1 is a one-tube-reaction for all microRNAs present in a sample. Step 2 is a microRNA specific qPCR

using forward and reverse primer pairs for a specific microRNA. An oval indicate insertion of LNAs in forward and reverse primers. When the method is carried out in practice the miRNAs present in a sample are firstly poly-A-tailed using a poly(A) polymerase, which adds adenine residues to the 3'-end of RNA molecules. Secondly, an extension primer, which has a poly-T-
 5 core nucleotide sequence, a 3'-end VN-degenerate motif and a 5'-end tail, is annealed to the poly-A-tailed miRNA through hybridisation with the VN-poly-T-sequence of the extension primer, (N=C, G, A and T; V= C, G, and A). This primer may be referred to as the Universal RT primer. Subsequently, the extension primer is extended in a reverse transcription reaction using the miRNA as template. All of these reaction are performed in a one-tube reaction. The
 10 resulting primary extension product is composed of the extension primer and the newly synthesized DNA, which is cDNA complementary to all the miRNAs in the sample. In the next step a miRNA-specific PCR is carried out. A miRNA-specific forward primer is annealed to 3'-end of the newly synthesized cDNA and the upper-strand synthesis is carried out by extending the forward primer in a DNA-polymerization reaction using the primary extension
 15 product as template. A miRNA-specific reverse primer composed of a miRNA-specific 3'-end sequence, a poly-T-stretch and a 5'-end tail is then hybridized to the upper-strand and the lower-strand is synthesized by extension of the reverse primer. Various demonstrations of the method of the invention are described in the EXAMPLES section.

FIGURE 2 shows amplification of hsa-miR-197 (Example 1) with the forward primer 5'
 20 ttmCaccaccttctcca (SEQ ID NO 1) and the reverse primer 5'-cttttttttttttGctgggt (SEQ ID NO 2). **Synthetic template:** 10⁷ copies of synthetic hsa-miR-197 were used for polyA tailing/RT. An amount corresponding to 10⁵ copies was used for PCR. **Total human RNA:** 100 ng of total human RNA was used for polyA-tailing/RT (step 1). An amount corresponding to 1 ng was used for PCR. **Negative controls:** water control and a total human RNA control without
 25 PolyA Polymerase in the RT reaction. The black line designates the result obtained with the synthetic template whereas the grey line show the result obtained with total human RNA as template. The melting curves of the result obtained with the synthetic template and the total human RNA are almost identical.

FIGURE 3 shows the amplification curves and the first derivative of the melting curves of
 30 Example 2

FIGURE 4 shows amplification plots of Example 4 wherein discrimination between *hsa-let-7a* and the variants *hsa-let-7f*, *hsa-let-7c* and *hsa-let-7e* are demonstrated. The amplification-signal from negative controls are also indicated.

FIGURE 5A, 5B and 5C shows the method and processes that enables PCR amplification and discrimination between mature miRs, pre-miRs and pri-miRs. See example 6, 7 and 8. A) Polyadenylation of all RNA 3'-ends. B) Reverse transcription of polyadenylated RNA to cDNA using the Universal RT primer (N=C, G, A and T; V= C, G, and A). C) MicroRNA specific PCR amplification using gene specific primers. Note that the site of polyadenylation in combination with the gene specific reverse primers ensures specific detection of each of the individual molecules mature miRs, pre-miRs and pri-miRs. Fig 5A illustrate the assay used for assaying mature miRs. Fig 5B illustrate the assay used for assaying pre-miRs and Fig 5C illustrate the assay used for assaying pri-miRs.

FIGURE 6 shows amplification curves of (A) the pre-mir-203 assay on pre-miR-203, miR-203 and non-template controls (NTC). (B) Amplification curves of (A) the miR-203 assay on pre-miR-203, miR-203 and non-template controls (NTC).

FIGURE 7 shows the different expression of a panel of selected miRNA genes in heart and liver. The data is presented as the difference in the crossing point (Cp) value (Δ Cp) between heart and liver, see table 15.

FIGURE 8 shows the detection of each of four listed miRs assayed by the present method (black lines with diamonds) and compared with a commercial available method wherein the Reverse Transcription is performed as a one-tube-reaction for all microRNAs present in a sample (gray lines with spheres). The figure presents the crossing point (Cp) values as a function of template concentration. Each value is in triplicate.

DEFINITIONS

"small RNA molecules" refer to tiny RNA molecules in a living cell such as small "non-coding RNA molecules" i.e. molecules that are not translated into proteins. Non-coding RNA molecules include RNAs such as microRNAs (miRNAs), small interfering RNAs (siRNAs), small nuclear RNAs (snRNAs), small nucleolar RNAs (snoRNAs), small temporal RNAs (stRNAs), antigene RNAs (agRNAs) and piwi-interacting RNAs (piRNAs).

The terms "microRNA", "miRNA" and "miR" are used synonymously and refer to 21-25 nucleotides non-coding RNAs derived from endogenous genes of living organisms such as animals and plants. These so-called mature miRNAs are biologically active and processed from longer hairpin-like precursors termed pre-miRNAs (pre-miRs) having a length of approximately 75 nucleotides. The precursors of pre-miRNAs are pri-miRNAs (pri-miRs) with a length of 500 to 3000 nucleotides. miRNAs assemble in complexes termed miRNPs and act as major

regulators on important biological processes by binding to messenger RNA and interfering with translation efficiency. Target microRNAs of the present invention refer to all known microRNAs, such as microRNAs known from the scientific literature and public databases such as the miRBase database (<http://microrna.sanger.ac.uk/>) which is the home of microRNA data on the web administrated by the Sanger Institute, UK. The miRBase release 12 is hereby incorporated by reference, including all the mature miRNA and pre-mature miRNA sequences disclosed therein. "microRNA profiling" describes a large-scale analysis in which the expression levels of all microRNAs in a sample such as a tumor sample are determined to build microRNA signatures for a particular disease such as a cancer disease.

5 "Adding poly-A tails", "Poly-A tailing" and "polyadenylation" refers to the synthesis of a poly(A) tail, a stretch of RNA where all the bases are adenines, at the 3'-termini of an RNA molecule. Polyadenylation is a natural biological process in living organisms, but it can also be carried out in vitro using various polymerases such as commercial available *E. coli* Poly(A) Polymerase I (E-PAP).

15 "Extension primer" and "RT-primer" refers to an oligonucleotide primer, comprising a recognition sequence, complementary to a sequence in the target deoxyribonucleic and/or ribonucleic acid sequence, e.g. to the 3'-end of the mature microRNA or small noncoding RNA in the target ribonucleic acid sequence, and an anchor sequence essential for subsequent amplification by PCR. The said extension primer is used as an anchored primer in a reverse transcription reaction to generate a primer extension product or cDNA.

25 A "cDNA" refers to a complementary DNA produced by reverse transcription of an RNA template using a reverse transcriptase enzyme. Any reverse transcriptase can be used to synthesize the cDNA molecules, such as reverse transcriptases derived from moloney murine leukemia virus (M-MuLV) reverse transcriptase, avian myeloblastosis virus (AMV) reverse transcriptase, bovine leukemia virus (BLV) reverse transcriptase, Rous sarcoma virus (RSV) reverse transcriptase and human immunodeficiency virus (HIV) reverse transcriptase.

30 "Primers" refers to short, chemically synthesized oligonucleotides, usually with a length of about twenty to thirty bases. They are hybridized to a target DNA, which is then copied by a DNA polymerase to produce a complementary DNA strand. A "forward primer" and a "reverse primer" constitute a "PCR primer set" used in PCR, where they hybridise to complementary DNA strands and direct replication towards each other producing the upper-strand and the lower-strand, respectively, leading to an exponential increase in the target DNA segment. The template derived extension of PCR primers can be carried out by any DNA polymerases, such as bacterial thermostable DNA polymerases, including Taq DNA polymerase from *Thermus*

aquaticus, Pfu DNA polymerase from *Pyrococcus furiosus*, Vent DNA polymerase from *Thermococcus litoralis* or recombinant DNA polymerases such as Phusion DNA polymerase.

The terms "amplification", "PCR", "PCR reaction" and "PCR amplification", are interchangeable terms used to signify use of a nucleic acid amplification system, which multiplies the target
5 nucleic acids using the polymerase chain reaction (PCR).

"qPCR" and "real-time quantitative PCR" refers to the use of PCR to amplify and simultaneously quantify a targeted DNA molecule. qPCR enables both detection and quantification of the number of copies or relative amount of a specific sequence in a DNA sample, when normalized to DNA input. Amplified DNA is quantified as it accumulates in the
10 reaction in real time after each amplification cycle. Quantification is accomplished using various assay chemistries including fluorescent dyes, such as SYBR® green, that intercalate with double-stranded DNA, and fluorescent reporter oligonucleotide probes such as Taqman probes, that release fluorescence signal during the amplification process.

"qRT-PCR" refers to quantitative reverse transcription polymerase chain reaction in which a
15 cDNA produced in a reverse transcription reaction serves as the initial DNA template for the amplification process, which is then combined with qPCR to quantify low abundance of RNA molecules such as microRNAs in a sample of a particular cell or tissue type. Methods for qPCR and qRT-PCR is described in "A-Z of Quantitative PCR." (Bustin, SA (ed.) International University Line (La Jolla, CA, USA), 2004) which is hereby incorporated by reference in its
20 entirety.

"Hybridisation" refers to the bonding of two complementary single stranded nucleic acid polymers (such as oligonucleotides), such as RNA, DNA or polymers comprising or consisting of nucleotide analogues (such as LNA oligonucleotides). Hybridisation is highly specific, and may be controlled by regulation of the concentration of salts and temperature. Hybridisation
25 occurs between complementary sequences, but may also occur between sequences which comprise some mismatches. The oligonucleotides used in the methods of the present invention may, therefore be 100% complementary to the target molecule. Alternatively, in one embodiment the oligonucleotides may comprise at least one or two mismatches.

The term " T_m " or "melting temperature" of an oligonucleotide is in the present context a
30 measure of the stability of a duplex formed between the oligonucleotide and its perfect complement DNA strand determined at 115 mM Na^+ , no formamide. In general T_m is defined as the temperature at which 50 % of the duplexes formed between the oligonucleotide and a

complement nucleotide strand are dissociated into single strands. The length and nucleotide composition, such as the sequence of nucleotides and content of G and C nucleotides, of the oligonucleotide are important factors affecting T_m . Substitution of the normal A, G, C and T nucleotides with the corresponding LNA molecules in an oligonucleotide increases T_m .
5 Similarly, hybridisation conditions defined by salt concentration, oligonucleotide concentration, and the presence of denaturants (such as formamide or DMSO) affects T_m . Those skilled in the art of molecular biology know that several useful formulas for calculation of theoretical T_m 's have been developed to evaluate the T_m of an oligonucleotide for PCR, Southern and Northern blots, and *in situ hybridization*. Examples of T_m calculators are
10 OligoCalc (W. A. Kibbe (2007) *Nucleic Acids Res* Volume 35, Web Server issue W43-W46) and LNA Probe T_m Predictor at <http://www.exigon.com/oligo-tools>.

The term "base", as used herein covers the naturally occurring nucleobases adenine (A), guanine (G), cytosine (C), thymine (T) and uracil (U) as well as non-naturally occurring nucleobases such as xanthine, diaminopurine, 8-oxo- N^6 -methyladenine, 7-deazaxanthine, 7-
15 deazaguanine, N^4,N^4 -ethanocytosine, N^6,N^6 -ethano-2,6-diaminopurine, 5-methylcytosine, 5-(C^3-C^6)-alkynyl-cytosine, 5-fluorouracil, 5-bromouracil, pseudoisocytosine, 2-hydroxy-5-methyl-4-triazolopyridin, isocytosine, isoguanine, inosine and the "non-naturally occurring" nucleobases described in Benner et al., U.S. Patent No. 5,432,272 and Susan M. Freier and Karl-Heinz Altmann, *Nucleic Acid Research*, 25: 4429-4443, 1997. The term "nucleobase"
20 thus includes not only the known purine and pyrimidine heterocycles, but also heterocyclic analogues and tautomers thereof. Further naturally and non naturally occurring nucleobases include those disclosed in U.S. Patent No. 3,687,808; in chapter 15 by Sanghvi, in *Antisense Research and Application*, Ed. S. T. Crooke and B. Lebleu, CRC Press, 1993; in Englisch, et al., *Angewandte Chemie, International Edition*, 30: 613-722, 1991 (see, especially pages 622
25 and 623, and in the *Concise Encyclopedia of Polymer Science and Engineering*, J. I. Kroschwitz Ed., John Wiley & Sons, pages 858-859, 1990, Cook, *Anti-Cancer DrugDesign* 6: 585-607, 1991, each of which are hereby incorporated by reference in their entirety).

Nucleotides incorporated into oligonucleotides are referred to as nucleotide residues.

The term "nucleosidic base" or "nucleobase analogue" is further intended to include
30 heterocyclic compounds that can serve as like nucleosidic bases including certain "universal bases" that are not nucleosidic bases in the most classical sense but serve as nucleosidic bases. Especially mentioned as a universal base is 3-nitropyrrole or a 5-nitroindole. Other


preferred compounds include pyrene and pyridyloxazole derivatives, pyrenyl, pyrenylmethylglycerol derivatives and the like. Other preferred universal bases include, pyrrole, diazole or triazole derivatives, including those universal bases known in the art.

By "locked nucleic acid", "LNA", "LNA monomer" or "LNA molecule" (e.g., an LNA nucleoside or LNA nucleotide) or an LNA oligomer (e.g., an oligonucleotide or nucleic acid) is meant a nucleoside or nucleotide analogue that includes at least one LNA monomer.

To distinguish between LNA and natural occurring nucleotide residues in LNA containing oligonucleotide sequences presented herein, the LNAs are indicated by capital letters, whereas natural occurring nucleotide residues are indicated by lowercase letters: mC denotes LNA methyl cytosine.

LNA monomers as disclosed in PCT Publication WO 99/14226 are in general particularly desirable modified nucleic acids for incorporation into an oligonucleotide of the invention. Additionally, the nucleic acids may be modified at either the 3' and/or 5' end by any type of modification known in the art. For example, either or both ends may be capped with a protecting group, attached to a flexible linking group, attached to a reactive group to aid in attachment to the substrate surface, etc. Desirable LNA monomers and their method of synthesis also are disclosed in WO 98/39352.

Preferred LNA monomers, also referred to as "oxy-LNA" are LNA monomers which include bicyclic compounds as disclosed in PCT Publication WO 03/020739 wherein the bridge between R^{4'} and R^{2'} as shown in formula (I) below together designate -CH₂-O- or -CH₂-CH₂-O-.



It is understood that references herein to a nucleic acid unit, nucleic acid residue, LNA monomer, or similar term are inclusive of both individual nucleoside units and nucleotide units and nucleoside units and nucleotide units within an oligonucleotide.

A "modified base" or other similar terms refer to a composition (e.g., a non-naturally occurring nucleotide or nucleosidic base), which can pair with a natural base (e.g., adenine,

guanine, cytosine, uracil, and/or thymine) and/or can pair with a non-naturally occurring nucleotide or nucleosidic base. Desirably, the modified base provides a T_m differential of 15, 12, 10, 8, 6, 4, or 2°C or less as described herein. Exemplary modified bases are described in EP 1 072 679 and WO 97/12896.

- 5 The term "chemical moiety" refers to a part of a molecule. "Modified by a chemical moiety" thus refer to a modification of the standard molecular structure by inclusion of an unusual chemical structure. The attachment of said structure can be covalent or non-covalent.

The term "inclusion of a chemical moiety" in an oligonucleotide probe thus refers to attachment of a molecular structure. Such as chemical moiety include but are not limited to covalently
10 and/or non-covalently bound minor groove binders (MGB) and/or intercalating nucleic acids (INA) selected from a group consisting of asymmetric cyanine dyes, DAPI, SYBR[®] Green I, SYBR[®] Green II, SYBR[®] Gold, PicoGreen[®], thiazole orange, Hoechst 33342, Ethidium Bromide, 1-O-(1-pyrenylmethyl)glycerol and Hoechst 33258. Other chemical moieties include the modified nucleotides, nucleosidic bases or LNA modified oligonucleotides.

- 15 "Oligonucleotide analogue" refers to a nucleic acid binding molecule capable of recognizing a particular target nucleotide sequence. A particular oligonucleotide analogue is peptide nucleic acid (PNA) in which the sugar phosphate backbone of an oligonucleotide is replaced by a protein like backbone. In PNA, nucleotides are attached to the uncharged polyamide backbone yielding a chimeric pseudopeptide-nucleic acid structure, which is homomorphous
20 to nucleic acid forms.

"High affinity nucleotide analogue" or "affinity-enhancing nucleotide analogue" refers to a non-naturally occurring nucleotide analogue that increases the "binding affinity" of an oligonucleotide probe to its complementary recognition sequence when substituted with at least one such high-affinity nucleotide analogue.

- 25 As used herein, a probe with an increased "binding affinity" for a recognition sequence compared to a probe which comprises the same sequence but does not comprise a stabilizing nucleotide, refers to a probe for which the association constant (K_a) of the probe recognition segment is higher than the association constant of the complementary strands of a double-stranded molecule. In another preferred embodiment, the association constant of the probe
30 recognition segment is higher than the dissociation constant (K_d) of the complementary strand of the recognition sequence in the target sequence in a double stranded molecule.

Monomers are referred to as being "complementary" if they contain nucleotides that can form hydrogen bonds according to Watson-Crick base-pairing rules (e.g. G with C, A with T or A with U) or other hydrogen bonding motifs such as for example diaminopurine with T, 5-methyl C with G, 2-thiothymidine with A, inosine with C, pseudoisocytosine with G, etc.

- 5 By "oligonucleotide," "oligomer," or "oligo" is meant a successive chain of monomers (e.g., glycosides of heterocyclic bases) connected via internucleoside linkages. The linkage between two successive monomers in the oligo consist of 2 to 4, desirably 3, groups/atoms selected from $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{NR}^{\text{H}}-$, $>\text{C}=\text{O}$, $>\text{C}=\text{NR}^{\text{H}}$, $>\text{C}=\text{S}$, $-\text{Si}(\text{R}'')_2-$, $-\text{SO}-$, $-\text{S}(\text{O})_2-$, $-\text{P}(\text{O})_2-$, $-\text{PO}(\text{BH}_3)-$, $-\text{P}(\text{O},\text{S})-$, $-\text{P}(\text{S})_2-$, $-\text{PO}(\text{R}'')-$, $-\text{PO}(\text{OCH}_3)-$, and $-\text{PO}(\text{NHR}^{\text{H}})-$, where R^{H} is
- 10 selected from hydrogen and C_{1-4} -alkyl, and R'' is selected from C_{1-6} -alkyl and phenyl. Illustrative examples of such linkages are $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$, $-\text{CH}_2-\text{CO}-\text{CH}_2-$, $-\text{CH}_2-\text{CHOH}-\text{CH}_2-$, $-\text{O}-\text{CH}_2-\text{O}-$, $-\text{O}-\text{CH}_2-\text{CH}_2-$, $-\text{O}-\text{CH}_2-\text{CH}=\text{}$ (including R^{S} (see formula I) when used as a linkage to a succeeding monomer), $-\text{CH}_2-\text{CH}_2-\text{O}-$, $-\text{NR}^{\text{H}}-\text{CH}_2-\text{CH}_2-$, $-\text{CH}_2-\text{CH}_2-\text{NR}^{\text{H}}-$, $-\text{CH}_2-\text{NR}^{\text{H}}-\text{CH}_2-$, $-\text{O}-\text{CH}_2-\text{CH}_2-\text{NR}^{\text{H}}-$, $-\text{NR}^{\text{H}}-\text{CO}-\text{O}-$, $-\text{NR}^{\text{H}}-\text{CO}-\text{NR}^{\text{H}}-$, $-\text{NR}^{\text{H}}-\text{CS}-\text{NR}^{\text{H}}-$, $-\text{NR}^{\text{H}}-\text{C}(=\text{NR}^{\text{H}})-\text{NR}^{\text{H}}-$, $-\text{NR}^{\text{H}}-\text{CO}-\text{CH}_2-\text{NR}^{\text{H}}-$, $-\text{O}-\text{CO}-\text{O}-$, $-\text{O}-\text{CO}-\text{CH}_2-\text{O}-$, $-\text{O}-\text{CH}_2-\text{CO}-\text{O}-$, $-\text{CH}_2-\text{CO}-\text{NR}^{\text{H}}-$, $-\text{O}-\text{CO}-\text{NR}^{\text{H}}-$, $-\text{NR}^{\text{H}}-\text{CO}-\text{CH}_2-$, $-\text{O}-\text{CH}_2-\text{CO}-\text{NR}^{\text{H}}-$, $-\text{O}-\text{CH}_2-\text{CH}_2-\text{NR}^{\text{H}}-$, $-\text{CH}=\text{N}-\text{O}-$, $-\text{CH}_2-\text{NR}^{\text{H}}-\text{O}-$, $-\text{CH}_2-\text{O}-\text{N}=\text{}$ (including R^{S} when used as a linkage to a succeeding monomer), $-\text{CH}_2-\text{O}-\text{NR}^{\text{H}}-$, $-\text{CO}-\text{NR}^{\text{H}}-\text{CH}_2-$, $-\text{CH}_2-\text{NR}^{\text{H}}-\text{O}-$, $-\text{CH}_2-\text{NR}^{\text{H}}-\text{CO}-$, $-\text{O}-\text{NR}^{\text{H}}-\text{CH}_2-$, $-\text{O}-\text{NR}^{\text{H}}-$, $-\text{O}-\text{CH}_2-\text{S}-$, $-\text{S}-\text{CH}_2-\text{O}-$, $-\text{CH}_2-\text{CH}_2-\text{S}-$, $-\text{O}-\text{CH}_2-\text{CH}_2-\text{S}-$, $-\text{S}-\text{CH}_2-\text{CH}=\text{}$ (including R^{S} when used as a linkage to a succeeding monomer), $-\text{S}-\text{CH}_2-\text{CH}_2-$, $-\text{S}-\text{CH}_2-\text{CH}_2-\text{O}-$, $-\text{S}-\text{CH}_2-\text{CH}_2-\text{S}-$, $-\text{CH}_2-\text{S}-\text{CH}_2-$, $-\text{CH}_2-\text{SO}-\text{CH}_2-$, $-\text{CH}_2-\text{SO}_2-\text{CH}_2-$, $-\text{O}-\text{SO}-\text{O}-$, $-\text{O}-\text{S}(\text{O})_2-\text{O}-$, $-\text{O}-\text{S}(\text{O})_2-\text{CH}_2-$, $-\text{O}-\text{S}(\text{O})_2-\text{NR}^{\text{H}}-$, $-\text{NR}^{\text{H}}-\text{S}(\text{O})_2-\text{CH}_2-$, $-\text{O}-\text{S}(\text{O})_2-\text{CH}_2-$, $-\text{O}-\text{P}(\text{O})_2-\text{O}-$, $-\text{O}-\text{P}(\text{O},\text{S})-\text{O}-$, $-\text{O}-\text{P}(\text{S})_2-\text{O}-$, $-\text{S}-\text{P}(\text{O})_2-\text{O}-$, $-\text{S}-\text{P}(\text{O},\text{S})-\text{O}-$, $-\text{S}-\text{P}(\text{S})_2-\text{O}-$, $-\text{O}-\text{P}(\text{O})_2-\text{S}-$, $-\text{O}-\text{P}(\text{O},\text{S})-\text{S}-$, $-\text{O}-\text{P}(\text{S})_2-\text{S}-$, $-\text{S}-\text{P}(\text{O})_2-\text{S}-$, $-\text{S}-\text{P}(\text{O},\text{S})-\text{S}-$, $-\text{S}-\text{P}(\text{S})_2-\text{S}-$, $-\text{O}-\text{PO}(\text{R}'')-\text{O}-$, $-\text{O}-\text{PO}(\text{OCH}_3)-\text{O}-$, $-\text{O}-\text{PO}(\text{OCH}_2\text{CH}_3)-\text{O}-$, $-\text{O}-\text{PO}(\text{OCH}_2\text{CH}_2\text{S}-\text{R})-\text{O}-$, $-\text{O}-\text{PO}(\text{BH}_3)-\text{O}-$, $-\text{O}-\text{PO}(\text{NHR}^{\text{H}})-\text{O}-$, $-\text{O}-\text{P}(\text{O})_2-\text{NR}^{\text{H}}-$, $-\text{NR}^{\text{H}}-\text{P}(\text{O})_2-\text{O}-$, $-\text{O}-\text{P}(\text{O},\text{NR}^{\text{H}})-\text{O}-$, $-\text{CH}_2-\text{P}(\text{O})_2-\text{O}-$, $-\text{O}-\text{P}(\text{O})_2-\text{CH}_2-$, and $-\text{O}-\text{Si}(\text{R}'')_2-\text{O}-$; among which $-\text{CH}_2-\text{CO}-\text{NR}^{\text{H}}-$, $-\text{CH}_2-\text{NR}^{\text{H}}-\text{O}-$, $-\text{S}-\text{CH}_2-\text{O}-$, $-\text{O}-\text{P}(\text{O})_2-\text{O}-$, $-\text{O}-\text{P}(\text{O},\text{S})-\text{O}-$, $-\text{O}-\text{P}(\text{S})_2-\text{O}-$, $-\text{NR}^{\text{H}}-\text{P}(\text{O})_2-\text{O}-$, $-\text{O}-\text{P}(\text{O},\text{NR}^{\text{H}})-\text{O}-$, $-\text{O}-\text{PO}(\text{R}'')-\text{O}-$, $-\text{O}-\text{PO}(\text{CH}_3)-\text{O}-$, and $-\text{O}-\text{PO}(\text{NHR}^{\text{H}})-\text{O}-$, where R^{H} is selected from hydrogen and C_{1-4} -alkyl, and R'' is selected from C_{1-6} -alkyl and phenyl, are especially desirable. Further illustrative examples are given in
- 30 Mesmaeker *et. al.*, Current Opinion in Structural Biology 1995, 5, 343-355 and Susan M. Freier and Karl-Heinz Altmann, Nucleic Acids Research, 1997, vol 25, pp 4429-4443. The left-hand side of the internucleoside linkage is bound to the 5-membered ring as substituent P^* at the 3'-position, whereas the right-hand side is bound to the 5'-position of a preceding monomer.

The term "succeeding monomer" relates to the neighbouring monomer in the 5'-terminal direction and the "preceding monomer" relates to the neighbouring monomer in the 3'-terminal direction. It should be noted that whereas the natural synthesis of DNA and RNA proceed in the direction 5' to 3', many chemical synthesis-schemes proceed in the direction of
 5 3' to 5'.

When performing quantitative PCR, the cycle at which the fluorescence from a sample crosses the threshold is called the "cycle threshold" or Ct. Ct is used for quantification of a template.

The "crossing point" or Cp value is a slightly different, but related value that can be used for
 10 quantification of a template somewhat similar to the use of Ct. The LightCycler® 480 Software calculates the second derivatives of entire amplification curves and determines where this value is at its maximum. This value (crossing point, Cp) represents the cycle at which the increase of fluorescence is highest and where the logarithmic phase of a PCR begins. Both terms Ct and Cp are further explained in Bustin, SA (ed.) "A-Z of Quantitative
 15 PCR." International University Line (La Jolla, California, USA), 2004, which is included herein by reference.

Suitably, the "target" or "target nucleic acid" or "target ribonucleic acid" refers to any relevant nucleic acid of a single specific sequence, e. g., a biological nucleic acid, e. g., derived from a subject or human being. Within the context of the oligonucleotides and detection probes used
 20 in the invention to detect miRNAs, the "target" is a human miRNA or precursor thereof, or in one embodiment, a molecule which retains the genetic sequence information contained therein – such as all or (a sufficient) part of the sequence of nucleotides or reverse complement thereof.

"Target sequence" refers to a specific nucleic acid sequence (or corresponding nucleobase
 25 sequence) within any target nucleic acid.

The term "primer design" refers to methods such as those provided below. The primer design is an systematic approach used for designing the sequence of nucleotides in forward and reverse primers thus ensuring probe specificity and binding efficiency to target microRNAs. The following rules have been used for design of primers for miR-specific qPCR:


30 **Forward primer design:**

Preferably, the forward primers are designed to be identical of from 12 to 18 bases of the 5' end of the miR sequence. Preferably, the T_m of the forward primers should be in the range of

between 55 °C and 65 °C, however, a T_m below 55 °C and above 65 °C may also be acceptable. To ensure that the T_m of the primers preferably is in the range of between 55 °C and 65 °C, one or more LNA monomers may be inserted into the sequence substituting the natural nucleotide. An artificial nucleotide sequence may also be added to the 5'-end of the forward primer to make certain that the T_m is in the range of between 55 °C and 65 °C.

Reverse primer design:

The reverse primer is according to formula II:



wherein R^3 is a 5'-end nucleotide sequence, $(T)_x$ is a central part of x consecutive thymine residues in the amount of x, wherein x is an integer of from 1 to 100, and R^4 is a 3'-end nucleotide sequence that specifically hybridizes to a part of a microRNA molecule.

R^4 is preferably a nucleotide sequence of from 1 to 10 nucleotides designed from the 3'-end of a specific miRNA. R^4 is able to hybridize specifically to the DNA strand (i.e. the upper-strand) produced by extension of the corresponding miRNA specific forward primer. To ensure miRNA specificity and make certain that the T_m of the primer preferably is in the range of between 55 °C and 65 °C, one or more LNA monomer may be inserted into the R^4 sequence substituting the corresponding natural nucleotide.

The $(T)_x$ central part is preferably a stretch of approximately 15 consecutive thymine nucleotide residues that hybridizes to the poly-A-tail part of the DNA strand produced from the corresponding miRNA specific forward primer.

The R^3 sequence of the reverse primer typically has a length of from 1 to 20 nucleotides. For example R^3 can be 17 nucleotides long, 8, 7 or 6 nucleotides long or even only one nucleotide long. In certain embodiments of the present invention, the R^3 sequence is 5'-TGACACGGAGGTACTAG-3' (SEQ ID NO 3). The length of the R^3 sequence may be reduced

from the 5'-end to adjust the T_m of the reverse primer to the preferable range of between 55 °C and 65 °C. The R^3 sequence is identical to or at least has overlap to part of the R^1 sequence (formula (III)) of the extension primer.

The procedure for designing the reverse primer is described below.

5 **A.** Design a number of primers that fulfil the following conditions:

- 1) Delete all adenine residues from 3'-end of miR nucleotide sequence. They will form part of the poly-A tail.
- 2) After deleting adenine residues start with reverse complement to 3' end base of the miR and continue until one base overlap with forward primer.
- 10 3) Delete one base from the 3' end of the reverse primer if the sequence is a cytosine residue and the 3' end of the forward primer is a guanine residue.
- 4) Delete one base from the 3' end of the primer if the sequence is a guanine residue and the 3' end of the forward primer is a cytosine residue.
- 15 5) Delete one base from the 3' end of the primer if the last two bases are overlapping with the last two bases of the forward primer (from the 3' end).
- 6) Repeat this process until the conditions are fulfilled.
- 7) If no primer fulfils the conditions disregard rules #2 and 3 and allow two bases overlap (but not more) for sequences ending with AA, AT, TA or TT.
- 8) If no primer fulfils the conditions try another forward primer design.

20 **B.** Choose the longest reverse primer that fulfils the following conditions:

- 1) At least four bases long.
- 2) Less than five cytosine residues and guanine residues in the last six bases at the 3' end of the primer.

3) Less than four cytosine residues and guanine residues in the last five bases at the 3' end of the primer except if the last or second last base is an adenine residue or thymine residue .

4) If possible not more than two adenine residues or thymine residues in the 3' end.

5) If no primer fulfils the conditions try another forward primer design.

C. Optionally insert one LNA monomer according to the following rules:

1) Insert LNA at first cytosine residue or guanine residue from 5' end of the R^4 part of the reverse primer.

2) No LNA in sequences with three or more consecutive cytosine residues or guanine residues.

3) LNA should be in position four or higher from the 3' end of the primer.

4) If the conditions cannot be fulfilled insert LNA at first adenine residue or thymine residue from 5' end of the R^4 part of the reverse primer.

5) LNA should be in position four or higher from the 3' end of the primer.

6) If the conditions cannot be fulfilled insert LNA-thymine (LNA-T) at 5' end of the R^4 part of the primer.

D. Tailing:

Add a sequence of the formula $R^3-(T)_x$ to the 5' end of the reverse primer. I certain embodiments of the present invention the nucleotide sequence:

5'-TGACACGGAGGTACTAGTTTTTTTTTTTTTTT-3' (SEQ ID NO 4) is added at the 5' end of the reverse primer.

If an LNA-T was added to the miR-specific part of the reverse primer (step C6) delete one thymine residue from the tail as there should never be more than 15 consecutive thymine residues.

Eventually, nucleotides are removed from the 5'-end of the reverse primer to fine-tune the T_m of the primer.

"Sample" of RNA refers to RNA comprising compositions obtained from cells, tissues or fluids from an organism according to conventional procedures described e.g. in RNA Isolation and Characterization Protocols (Rapley, Ralph; Manning, David L. (Eds.) 1998) known to the skilled artisan or by using commercial kits such as miRNeasy (QIAGEN GmbH, Hilden, Germany)) or miRVana (Ambion Inc., Austin, TX, US). Sources for isolation of RNA fractions are samples of cells, or tissue or fluid isolated from an organism or organisms, including but not limited to, for example, skin, plasma, serum, spinal fluid, lymph fluid, synovial fluid, urine, tears, blood cells, organs, tumours, and also to samples of *in vitro* cell culture constituents (including but not limited to conditioned medium resulting from the growth of cells in cell culture medium, recombinant cells and cell components). "Sample" may also refer to cells or fluids that can be used directly in a qRT-PCR process without previous enrichment of the RNA fraction or even RNA comprising compositions comprising artificially synthesised RNA.

Cells or cell types also refer to any cells of Eukaryote, Prokaryote and Archaea origin.

A "living organism" refers to a living entity, including but not limited to, for example, human, mouse, rat, *Drosophila*, *C. elegans*, yeast, *Arabidopsis thaliana*, maize, rice, zebra fish, primates, domestic animals, etc.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

Detailed outline of the qRT-PCR reaction and the primers involved:

The present invention provides methods for amplification of microRNA molecules in a sample, the method comprising the steps described schematically in Fig. 1, and comprise:

(a) adding poly-A tails to a population of RNA molecules in a sample;

- (b) producing cDNA molecules of the poly-A-tailed RNA molecules using an extension primer in a reverse transcription reaction; and
- (c) amplifying the cDNA molecules by PCR using a forward primer and a reverse primer both of which are specific for said RNA molecule.

- 5 When performing the method in practice step (a) and (b) is typically performed as one concerted reaction which is universal for all RNAs to be analysed – hence its nickname: 'Universal RT'. The advantage of only one first-strand cDNA synthesis reaction (or RT reaction) to be used as template for multiple real-time PCR assays is that it saves precious sample, reduces technical variation and reduces time spent in the laboratory.
- 10 During step (c) of the method individual (or groups of individual) RNAs are specifically PCR amplified using specific forward and reverse primers. Typically the primers are optimised by introducing one or more LNA nucleotide analogues into the sequence of the primers, and typically the PCR is quantitative Real-Time PCR. As can be seen from the examples the method result in 1) uniquely specific assays that enable discrimination between highly related
- 15 RNA sequences and 2) an extremely low background which enables accurate quantitation of very low RNA levels.


The method has been extensively used to quantify small RNAs. In a preferred embodiment the small RNA, comprises small non-coding RNAs such as short interfering RNAs (siRNAs), mature microRNAs and pre-microRNAs. Also larger RNAs, e.g. precursors of pre-microRNAs

20 the pri-miRNAs (pri-miRs) and mRNAs may be assessed by method.

Most preferably the small RNA is a microRNA.

The extension primer may be of a length in the range from 10 to 100 nucleotides, such as a length in the range from 15 to 45 nucleotides. Preferably, the extension primer has a length of 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 nucleotides.

- 25 In a preferred embodiment, the extension primer is of the formula III:



wherein R^1 is a 5'-end nucleotide sequence, $(T)_y$ is a central part of consecutive thymine residues in the amount of y , wherein y is an integer of from 1 to 100, and R^2 is a 3'-end nucleotide sequence.

Typically, the 5'-end part R^1 is a nucleotide sequence with a length of from 1 to 30
 5 nucleotides. For example the R^1 sequence may be 6, 7, 8, 9, or 10 nucleotides long. The R^1 sequence includes at least a hybridisation sequence for the R^3 sequence in the reverse primer of the formula II (see below) to be used in the subsequent miRNA-specific qPCR.

Preferably, y in formula III is in the interval of from 5 to 50, more preferably y is in the interval of from 5 to 21. For example y is 12, 13, 14, 15, 16, 17 or 18. Most preferably y is
 10 15.

In one especially preferred embodiment y of formula (III) equals x of formula (II).

In a preferred embodiment, R^2 is a degenerated anchor sequence motif VN, comprising the two 3'-end terminal nucleotide residues, wherein V designate a selection of residues comprising bases which can base pair to all but one of the natural bases (e.g. base pair to
 15 guanine, cytosine, uracil and thymine but not to adenine), typically V designate that the base in any primer molecule is a base randomly selected between adenine, guanine and cytosine and wherein N designate a base which can base pair to any of the natural bases (e.g. adenine, guanine, cytosine, uracil and thymine), typically N may be a base randomly selected between adenine, guanine, cytosine or thymine residue.

20 In another preferred embodiment R^2 is a degenerated sequence motif VNN, comprising the three 3'-end terminal nucleotide residues, wherein V designate a selection of residues comprising bases which can base pair to all but one of the natural bases (e.g. base pair to guanine, cytosine, uracil and thymine but not to adenine), typically V designate that the base is a base randomly selected between adenine, guanine and cytosine and wherein N designate
 25 a base which can base pair to any of the natural bases (e.g. adenine, guanine, cytosine, uracil and thymine), typically N may be a base randomly selected between adenine, guanine, cytosine or thymine residue.

In a preferable embodiment, the extension primer comprises at least one LNA.


In a certain embodiment of the present invention the extension primer has the sequence 5'-GGTACTAGTTTTTTTTTTTTTTVN-3' (SEQ ID NO 5).

Preferably, the forward primer has a length in the range of from 10 to 100 nucleotides, such as 12 to 22 or 13 to 20 or 14,15, 16, 17, 18 or 19 nucleotides. See table 18 for specific embodiments.

The forward primer may comprise one, two or even more LNAs.

In a preferred embodiment, the nucleotide sequence of the forward primer is designed to specifically hybridize to the complement DNA molecule of a specific microRNA molecule using the primer design rules described in the definition of 'primer design' (see DEFINITIONS section)

Preferably, the reverse primer is of the formula II:



wherein R^3 is a 5'-end nucleotide sequence $(T)_x$ is a central part of consecutive thymine nucleotides in the amount of x, wherein x is an integer of from 1 to 100, and R^4 is a 3'-end nucleotide sequence that specifically hybridizes to a nucleotide sequence of a target RNA molecule.

Preferably, the 5'-end nucleotide sequence R^3 of the reverse primer of formula II is a nucleotide sequence with a length of from 1 to 30 nucleotides.

Preferably, x in formula II is in the interval of from 5 to 50, more preferably x is in the interval of from 5 to 21. For example x is 12, 13, 14, 15, 16, 17 or 18. Most preferably x is 15.

In one especially preferred embodiment x of formula (II) equals y of formula (III).

In a preferred embodiment, the 3'-end nucleotide sequence R^4 of the reverse primer of formula II has a length in the range of from 1 to 10 nucleotides.

As illustrated in example 2 LNA has a profound effect. Accordingly, in a preferred embodiment, the 3'-end part, R^4 , of the reverse primer of formula II comprises at least one

5 LNA. More preferably, the 3'-end part, R^4 , of the reverse primer of formula II contains only one LNA. In a particularly preferred embodiment the LNA is situated in the 5' position or the position adjacent to the 5' position of the R^4 part of the reverse primer.

In a preferred embodiment the nucleotide sequence of the reverse primer is designed to specifically hybridize to a specific microRNA molecule using the primer design rules described
10 in the definition of "primer design" (see DEFINITIONS section)

Design of primers

The invention also provides methods for designing the nucleotide sequence of the forward primers and the reverse primers used in the invention. Systematic approaches and experimental evaluation of the primers are presented in the EXAMPLES 3 section.

15 Examples of applications of the present invention

The present invention is useful for amplification and quantification of small RNA molecules such as microRNAs or siRNA's in various cells or tissues originating from humans or other organisms as described in EXAMPLES 5 and 9. As illustrated in example 10 the present method is superior with respect to sensitivity and specificity when compared with a
20 somewhat similar single tube cDNA synthesis approach which is commercially available.

Thus one aspect of the invention is a method for measuring the amount of a target microRNA in a sample from a living organism, the method comprising the steps of:

- a) amplifying the target microRNA according to the method of any one of claims 1 to 28
- b) measuring the amount of the amplified DNA molecules.

The amount of the amplified DNA molecules is typically measured using fluorescence-based quantitative real-time PCR by e.g. monitoring SYBR[®] green fluorescence as a function of PCR cycle number.

Accordingly, the present invention can be used as a tool for microRNA profiling of different cell and tissue types within a living organism. Thus, the number of specific microRNAs and the amounts of specific microRNAs present in different cell and tissue types can be determined using the present invention. Similarly, the invention can be used to discriminate between the levels of mature microRNAs, pre-microRNAs and pri-microRNAs in different cell and tissue types by using appropriate primers targeting the mature microRNA or its precursors (see fig 5).

In another aspect, the present invention can be used as a tool for microRNA profiling of patients suffering from various diseases such as cancer. By way of example, the microRNAomes of various cancer tissues can be established using the invention.

In yet another aspect the present invention can be used for diagnosis of various diseases such as cancer by measuring microRNA expression in tissues and body fluids from normal individuals and from individuals having a disease, and subsequently analyse for differences in the microRNA profiles.

In another embodiment the present invention may be used for measuring changes in the amount of specific microRNAs in response to treatment of a disease affected individual by means of pharmaceuticals, such as chemotherapeutical agents, and by means of surgery.

In another embodiment the present invention may be used for measuring specific predictive microRNA biomarkers that correlate with re-currence free survival of patients having a disease, such as a cancer disease.

Due to its simplicity: one "universal-RT" step and one discriminatory PCR step, the present invention is also suitable for high-throughput methods on robot platforms directed to the quantification of a single microRNA or a collection of microRNAs. Thus, the method is particularly suitable for multiple subsequent PCR reactions of individual microRNAs as described in EXAMPLE 5, 9 and 10.

In a certain embodiment the method of the invention may be used to quantitatively determine the amount of mature microRNA in a sample without interference from the

corresponding pre-miR, and vice versa the amount of a pre-miR can be determined without interference from the mature microRNA as described in EXAMPLES 6, 7 and 8.

In another embodiment the method of the invention may be used for discrimination between targets with single nucleotide differences as described in EXAMPLE 4.

- 5 In another aspect, the present invention provides kits for detecting at least one target microRNA, the kits comprising one or more primer sets specific for the detection of a target microRNA, each primer set comprising firstly, a universal extension primer for producing a cDNA molecule complementary to a target microRNA, and secondly, a PCR primer set comprising a forward primer specific for the 5'-end of the target miRNA and a reverse primer
10 specific for the 3'-end of the target miRNA for amplifying the cDNA molecule.

Accordingly in one aspect the invention a kit for detecting at least one target RNA is provided. The kit comprises at least one primer set specific for the detection of a target RNA, said primer set comprises:

a) an extension primer of formula I: $R^1-(T)_y-R^2$ (III)

- 15 wherein R^1 is a 5'-end nucleotide sequence, $(T)_y$ is a central part of consecutive thymine residues in the amount of y, wherein y is an integer of from 1 to 100, and R^2 is a 3'-end nucleotide sequence.

b) a reverse primer of formula II: $R^3-(T)_x-R^4$ (II)

- wherein R^3 is a 5'-end nucleotide sequence, $(T)_x$ is a central part of consecutive thymine
20 residues in the amount of x, wherein x is an integer of from 1 to 100, and R^4 is a 3'-end nucleotide sequence that specifically hybridizes to a nucleotide sequence of a target RNA molecule, and

c) a forward primer

- In one embodiment the kit is designed for detecting at least one mammalian target microRNA
25 comprising at least one primer set specific for the detection of a target microRNA.

Preferably, the extension, the reverse and the forward primers are designed following the design rules given in the "primer design"-part of the DEFINITIONS section and elsewhere and forward and reverse primers which includes at least one LNA molecule are especially preferred embodiments.

- 5 A non-limiting example of an extension primer is shown as SEQ ID NO 5.

Forward and reverse primers included in the kit may be designed to detect any mammalian target microRNA in accordance with the methods described herein. Non-limiting examples of forward and reverse primers are listed in TABLE 16.

- 10 In certain embodiments, the kit includes multiple primer sets that may be used to detect various mammalian microRNA targets, such as two microRNA targets up to several hundred microRNA targets.

A kit of the invention may also provide an array of primers delivered in microtiter qPCR plates such as 96, 768, 869, 1536 or even 3456-well microtiter qPCR plates suitable for robot handling as indicated in EXAMPLE 9.

- 15 A kit of the invention may also include reagents required for the poly-A-tailing, primer extension and PCR reactions, such as buffers, salts, reducing agents, deoxy nucleoside triphosphates, nucleoside triphosphates, and enzymes. A detection reagent for the qPCR such as SYBR® green may also be included. Similarly a kit for RNA isolation may also be included.

- 20 A further aspect of the invention is a high-throughput method for measuring the amount of specific target microRNAs in a sample from a living organism by using the method of the present invention and integrating an automated wherein the combined poly-A-tailing and reverse transcription reaction is aliquoted into individual wells of a microtiter plate containing microRNA specific primer sets of forward and reverse primers, resulting in the steps of:

- 25 a) adding poly-A tails to a population of small RNA molecules in a sample and producing cDNA molecules of the poly-A-tailed small RNA molecules using an extension primer in a reverse transcription reaction; and
- b) pipetting aliquots of the combined poly-A-tailing and reverse transcription reaction into individual wells of a microtiter plate

- c) amplifying specific target microRNAs in individual wells of a microtiter plate containing microRNA specific primer sets
- d) measuring the amount of specific microRNA molecules in individual wells.

being performed in a fashion compatible with a high-throughput experimental set-up. Such a
5 set-up will typically comprise one or more pipetting robots.

EXAMPLES

EXAMPLE 1: PRODUCTION OF SPECIFIC DNA MOLECULES USING MIR-SPECIFIC QPCR.

In this example hsa-miR-197 was amplified from a human RNA sample using the miR specific quantitative reverse transcription polymerase chain reaction (qRT-PCR) of this invention.

5 Mixed on ice:

- 1 µl 10x Poly(A) polymerase buffer (New England Biolabs)
- 1 µl 1 mM ATP
- 1 µl 10 µM RT-primer (=extension primer) (L2TA: 5'-ggtagctagttttttttttttvn (SEQ ID NO 5), v designates cytosine, guanine and adenine residues, n designates cytosine, guanine, adenine and thymine residues)).
- 1 µl mix of 1 mM dATP, 1 mM dCTP, 1 mM dGTP, 1 mM dTTP
- 0,5 µl (200U/µl) MuLV reverse transcriptase (New England Biolabs Inc., Ipswich, MA, US)
- 0,2 µl (5U/µl) Poly(A) polymerase (New England Biolabs)
- 1 µl RNA sample (total human RNA sample =100 ng of a mixture of 25 ng human heart RNA, 25 ng human brain RNA, 25 ng human liver RNA and 25 ng human lung RNA all from Ambion, Austin, TX, US. Synthetic templates were prepared in 10ng/µl phage MS2 RNA in TE, approximately 10⁷ copies are added to the reaction. Synthetic template is obtained from Integrated DNA technologies Inc., Coralville, IA, US.)
- Water to 10 µl

Negative controls: water control and a total human RNA control without PolyA Polymerase in the RT reaction.

The mixture was incubated at 42 °C, 1 hour followed by 95 °C, 5 minutes.

For qPCR 1 µl or less of the polyA-tailing/RT reaction (step 1 on fig 1) was used for each PCR reaction with gene-specific forward and reverse primers for hsa-miR-197; forward primer 5' ttmCaccaccttctcca (SEQ ID NO 1) and the reverse primer 5'-cttttttttttttGctgggt (SEQ ID NO 2) (in the nucleotide sequences lower case designates a natural occurring nucleotide, upper case designates LNA, mC denotes LNA methyl cytosine).

Real-time PCR was done on a ABI 7500[®] thermocycler (Applied Biosystems Inc, Foster City, CA, US) by monitoring SYBR[®] green fluorescence as a function of PCR cycle number. A typical PCR reaction mixture contained:

- 10 µl 2× PCR mastermix (Roche cat#04 673 484 001, Roche Diagnostics A/S, Hvidovre, Denmark)
 - 1 µl or less of the RT reaction
 - 1 µl TE buffer with 2.5 µM miR-specific forward primer and 2.5 µM miR-specific reverse primer (TE buffer: 10 mM Tris/HCl (pH 8.0), 1mM EDTA).
 - Water to 20 µl.
- 10 The mixture was incubated at 95°C for 10 min followed 40 cycles of 95°C for 5 sec; 60°C for 60 sec with measurement of the fluorescence.

Amplification of the correct product was measured by comparing the melting curve profile of the PCR product to the melting curve profile of a PCR product obtained by amplification of a synthetic template (Fig. 2). No signal was obtained with the negative controls. The experiment shows that the same and correct hsa-miR-197 PCR product was obtained both using the total human RNA sample and a c hsa-miR-197 as template in step 1.

EXAMPLE 2: EFFECT OF LNA IN THE REVERSE PRIMER.

The effect of different designs of miR-specific PCR primers of step 2 can be tested on artificial DNA templates with the same sequence as the product of the reverse transcription reaction. An important advantage of using an artificial DNA template for PCR is that experimental variations in the efficiency of the reverse transcription step are eliminated.

Hsa-let-7a DNA template:

5'-tgcggtgacacggaggtactagtttttttttttaactatacaacctactaccta-3' (SEQ ID NO 6)

Salmon DNA: 2 ng/µl in TE buffer

25 MiR-specific forward primer:

F7a: 5'-tGaGgtagtaggttg (SEQ ID NO 7) (lower case designates a natural occurring nucleotide, upper case designates LNA).

MiR-specific reverse primers:

r7a.2: 5'-cggtgacacggagggtactagtttttttttttaactata (SEQ ID NO 8)

5 r7a.7: 5'-cggtgacacggagggtactagtttttttttttaamCtata (SEQ ID NO 9)

A PCR mix was prepared:

- 550 µl 2× PCR mastermix (Roche cat#04 673 484 001)
- 440 µl water, and as described i tab 1.

TABLE 1: Four PCR reactions were prepared:

Sample	Template	10 µM forward primer	10 µM reverse primer	PCR Mix
#1	1 µl salmon DNA	0,5 µl F7a	0,5 µl r7a.2	18 µl
#2	10000 copies of hsa- let-7a DNA template in salmon DNA	0,5 µl F7a	0,5 µl r7a.2	18 µl
#3	1 µl salmon DNA	0,5 µl F7a	0,5 µl r7a.7	18 µl
#4	10000 copies of hsa- let-7a DNA template in salmon DNA	0,5 µl F7a	0,5 µl r7a.7	18 µl

10 Real-time PCR was carried out on a ABI 7500[®] thermocycler by monitoring SYBR[®] green fluorescence as a function of PCR cycle number.

The mixture was incubated at 95°C for 10 min followed by 40 cycles of 95°C for 5 sec; 60°C for 60 sec with measurement of the fluorescence.

15 Amplification of the correct product was measured according to the recommendations from the manufacturer of the thermocycler (Applied Biosystems, Foster City, CA, US). Briefly, the PCR reaction was incubated at 60°C for 1 minute and fluorescence was measured during a

slow increase of the temperature to 95°C. The first derivative of the melting curves is shown in figure 3:

The results show (Figure 3) that the reverse primer without LNA (r7a.2) gives a positive signal in the negative control whereas the reverse primer with LNA (r7a.7) does not. Thus the
5 LNA base in r7a.7 is necessary in order to avoid an unspecific PCR product developed in the negative control reaction. Both primers amplify the positive control template.

As can be seen from table 2, it was found that the inclusion of one LNA in the miR-specific sequence of the reverse primer reduced background signal in PCR.

TABLE 2: Effect of LNA in primer pairs for miR specific qPCR

<i>hsa</i> - miR	Forward primer 5' → 3	Reverse primer 5' → 3	C _T Nega- tive control	C _T Tem- plate
let- 7a	tGaGgtagtaggttg (SEQ ID NO 7)	cggtgacacggaggtagtagtttttttttttaactata (SEQ ID NO 8)	28	25
		cggtgacacggaggtagtagttttttttttaamCta a (SEQ ID NO 9)	nd	24
miR -16	tAgmCagcacgtaaa (SEQ ID NO 10)	cacggaggtagtagttttttttttcgccaat (SEQ ID NO 11)	27	24
		cacggaggtagtagttttttttttmCgccaat (SEQ ID NO 12)	nd	26
miR -21	tAgmCttatcagactg atg (SEQ ID NO 13)	gcggtgacacggaggtagtagttttttttttcaac (SEQ ID NO 14)	32	25
		gcggtgacacggaggtagtagttttttttttmCaac (SEQ ID NO 15)	38	24
miR -23a	atmCamCattgccag (SEQ ID NO 16)	tgacacggaggtagtagttttttttttggaaatc (SEQ ID NO 17)	34	25
		tgacacggaggtagtagttttttttttGgaaatc (SEQ ID NO 18)	39	26
miR - 150	tmCtmCccaacccttg (SEQ ID NO 19)	tgacacggaggtagtagttttttttttmCactggta (SEQ ID NO 20)	28	23
		tgacacggaggtagtagttttttttttcactggta (SEQ ID NO 21)	38	25
miR - 208 a	ataaGamCgAgcaa aaag (SEQ ID NO 22)	mCggtgacacggaggtagtagtttttttttttacaag c (SEQ ID NO 23)	35	25
		cggtgacacggaggtagtagttttttttttAcaagc (SEQ ID NO 24)	nd	26
		cggtgacacggaggtagtagttttttttttTacaagc (SEQ ID NO 25)	nd	26

Lower case designates natural occurring nucleotides, upper case designates LNA, : mC denotes LNA methyl cytosine, C_T designates cycle threshold, nd, not detectable.

EXAMPLE 3: MANUAL PRIMER DESIGN AND VALIDATION OF PRIMERS FOR MIRS.

Primers were manually designed following the design-rules presented in the "primer design"-part of the DEFINITIONS section.

Primer validation:

- 5 By using the primer design rules it is possible to achieve PCR primers that have >70 % success rate in real-time PCR experiments according to the following validation criteria:

Prepare Rtmix for four reverse transcription (RT) samples:

1. Water
2. 100 ng total human RNA mix* (as default we use a mix of 25 ng of each heart, brain,
10 liver and lung total RNA from Ambion).

*RNA mix: 1µl 1µg/µl heart RNA, 1µl 1µg/µl brain RNA, 1µl 1µg/µl liver RNA, 1µl 1µg/µl lung RNA, 36µl TE. Store in 1µl aliquots at -80°C

3. 100 ng total human RNA mix without PolyA tailing.
4. 10⁷ copies of the synthetic miR in 10 ng phage MS2 RNA in TE (synthetic miRs were
15 obtained from Integrated DNA technologies Inc., Coralville, IA, US.). The miR can be added as a pool comprising up to at least 20 miRs.

RTmix	Total
10x PAP buffer	4,4µl
1mM ATP	4,4µl
1µM L6TA RTprimer	4,4µl
4x 1mM dNTP	4,4µl
Transcriptor 1:10	2,2µl
Water	11µl

PAP buffer = Poly(A) polymerase buffer (New England Biolabs).

L6TA = 5'-tgccgtgacacggaggtactagtttttttttttVN (SEQ ID NO 129).

- 20 Transcriptor = Reverse Transcriptase. (Roche Diagnostics A/S, Hvidovre, Denmark, cat# 03 531 295 001).

25

30

Reverse transcription reactions:

sample	0.1 µg/µl RNAmix*	1µl	water	0,5U/µl PAP ^a (NEB)	RTmix
#1			1µl	2µl	7µl
#2	1µl			2µl	7µl
#3	1µl		2µl		7µl
#4		1e7miR		2µl	7µl

Incubate 42°C, 1h > 95°C, 5'

Add 90µl TE to each sample and store at -20°C

a) Remember to dilute the poly(A) polymerase (PAP) 1:10 before use!

miR-specific qPCR:

Run PCR with the forward and reverse primers on 1 µl of each reverse transcription (RT) reaction:

5

Sample	1µl RT reaction	5µM forward primer	5µM reverse primer	2xfaststart	Water
#1	#1	0,5µl	0,5µl	10µl	8µl
#2	#2	0,5µl	0,5µl	10µl	8µl
#3	#3	0,5µl	0,5µl	10µl	8µl
#4	#4	0,5µl	0,5µl	10µl	8µl

Real-time PCR on the ABI 7500:

95°C, 10 min

95°C, 15 sec; 60°C, 60 sec; 40 cycles

10 Run melting curve analysis

Acceptance criteria:

Sample 1: Cycle threshold (C_t) above 40 compared to exponential area for #2 and 4.
Melting peak derivative below 5000.

Sample 2: C_t below 35

15 One peak on bell shaped melting curve, unless otherwise justified the peak should be between 69 °C and 80 °C. Peak at same temperature (+/- 0.5 °C) as in sample 4.

In the exponential area the crossing point at 10 times delta Rn minus the crossing point at delta Rn should be between 3.2 and 4.4:

$$3.2 < (C_{10T} - C_T) < 4.4$$

Sample 3: C_t above 40 compared to exponential area for #2 and 4.

5 Melting peak derivative below 5000.

Sample 4: One peak on bell shaped melting curve, unless otherwise justified the peak should be between 69 °C and 80 °C. Peak at same temperature (+/- 0.5 °C) as in sample 2.

Table 3: Design and validation of 17 primer pairs

miR	Forward primer 5' → 3'	SEQ ID NO	Reverse primer 5' → 3'	SEQ ID NO	Assay valid?
<i>hsa-miR-124</i>	taaGgcacgcggtga	26	tgacacggaggtagtctttttttttTggcat	43	Yes
<i>hsa-miR-9</i>	ctgimCtttGgttatctag	27	tgacacggaggtagtctttttttttmCatacag	44	Yes
<i>hsa-miR-181a</i>	aamCattmCaacgctgt	28	tgacacggaggtagtctttttttttamCtcac	45	Yes
<i>hsa-miR-128</i>	tmCamCagtgaaccggt	29	tgacacggaggtagtctttttttttAaagaga	46	Yes
<i>hsa-miR-488</i>	ttGaaaGgctatttc	30	tgacacggaggtagtctttttttttGacca	47	No
<i>hsa-miR-328</i>	ctggccctctctgcc	31	tgacacggaggtagtctttttttttAcggaa	48	Yes
<i>hsa-miR-324</i>	cgcatcccttagggcat	32	tgacacggaggtagtctttttttttAcacca	49	Yes
<i>hsa-miR-331</i>	cTaGgtatggtccca	33	tgacacggaggtagtctttttttttGgatc	50	No
<i>hsa-miR-34a</i>	tGgcAgtgtcttagc	34	tgacacggaggtagtctttttttttamCaacca	51	Yes
<i>hsa-miR-130a</i>	caGtGcaatgttaaag	35	tgacacggaggtagtctttttttttAtgc	52	No
<i>hsa-miR-1</i>	tGgAatgtaaagaagt	36	tgacacggaggtagtctttttttttAtacata	53	No
<i>hsa-miR-23b</i>	atmCamCattgccag	37	tgacacggaggtagtctttttttttggTaatac	54	Yes
<i>hsa-miR-133a</i>	tTtGgtcccttcaa	38	tgacacggaggtagtctttttttttGagctg	55	Yes
<i>hsa-miR-133b</i>	tTtGgtcccttcaa	39	tgacacggaggtagtctttttttttAgctg	56	Yes
<i>hsa-miR-93</i>	cAaaGtgctgtctgtg	40	tgacacggaggtagtctttttttttmCtacctg	57	No
<i>hsa-miR-181b</i>	aamCattmCattgctgtc	41	tgacacggaggtagtctttttttttAccca	58	Yes
<i>hsa-miR-24</i>	tGgctmCagtcagca	42	tgacacggaggtagtctttttttttmCtgttc	59	Yes
					Total: 12/17 = 71%

Lower case designates natural occurring nucleotides, upper case designates LNA, mC denotes LNA methyl cytosine.

- 5 Result (see tab 3): Seventeen primers pairs were designed according to the design rules. Twelve of the 17 assays were successfully validated according to the validation protocol corresponding to a success rate of 71%.

Table 4: Design and validation of 15 primer pairs. The theoretical T_m of the reverse primer was optimised to 59 °C:

miR	Forward primer 5' → 3	SEQ ID NO	Reverse primer 5' → 3	SEQ ID NO	Assay valid?
<i>hsa-miR-146b</i>	ctgtGagaactgaattcca	60	gtactagtttttttttttAgcct	75	Yes
<i>hsa-miR-146b-3p</i>	tgccctgttgactca	61	gtactagttttttttttmCcagaac	76	Yes
<i>hsa-miR-429</i>	ctgtaatamCtGcttgga	62	gtactagtttttttttttAcggtt	77	Yes
<i>hsa-miR-193a-3p</i>	aactggcctacaaagtc	63	gtactagttttttttttamCtgg	78	Yes
<i>hsa-miR-193a</i>	gtctttgcgggcca	64	ggtactagttttttttttmCaict	79	Yes
<i>hsa-miR-193b</i>	aamCtggccctcaaag	65	gaggiactagttttttttttmCaict	80	Yes
<i>hsa-miR-149</i>	tmCtggcTcogtgcctca	66	tagtttttttttttgggAggtg	81	Yes
<i>hsa-miR-452</i>	aamCtGttgcagaggaa	67	gtactagttttttttttmCagtt	82	Yes
<i>hsa-miR-452*</i>	ctmCaimCtgcaaagaagta	68	gtactagttttttttttmCacttt	83	No
<i>hsa-miR-30b</i>	tGtaaamCatactactc	69	gtactagtttttttttttaGciga	84	No
<i>hsa-miR-34b</i>	ctgcaatcactaactccact	70	gaggtagtagtttttttttttAtgg	85	Yes
<i>hsa-miR-34c</i>	aggcaGttagtagctg	71	tactagtttttttttttGcaatca	86	Yes
<i>hsa-miR-134</i>	tgtgactggfaccaga	72	gtactagttttttttttTccct	87	Yes
<i>hsa-miR-671</i>	ctgaGgaagccctgga	73	gtactagttttttttttmCtcca	88	No
<i>hsa-miR-769-3p</i>	ctgggatctccgggtct	74	aggiactagttttttttttAacca	89	Yes
					Total: 12/15 = 80%

Lower case designates natural occurring nucleotides, upper case designates LNA, mC denotes LNA methyl cytosine.

5

Result (see tab 4): 15 primers pairs were designed according to the design rules. 12 of the 15 assays were successfully validated according to the validation protocol corresponding to a success rate of 80%.

10 EXAMPLE 4: DISCRIMINATION BETWEEN TARGETS WITH A SINGLE NUCLEOTIDE DIFFERENCE.

There are three miRs that only differ by one nucleotide from the sequence of *hsa-let-7a* (table below).

Table 5: Nucleotide sequence of let-7 family miRNAs

miR	Sequence ^{a)}
<i>hsa-let-7a</i>	5'-ugagguaguagguuguauaguu-3' (SEQ ID NO 90)
<i>hsa-let-7f</i>	5'-ugagguaguag g uuguauaguu-3' (SEQ ID NO 91)
<i>hsa-let-7c</i>	5'-ugagguaguagguuguau g guu-3' (SEQ ID NO 92)
<i>hsa-let-7e</i>	5'-ugagguag g agguuguauaguu-3' (SEQ ID NO 93)

a): The difference between the four closely related members of the let-7 family is indicated by bold letters.

To test if the qPCR primers for *hsa-let-7a* detect the miRs with a single nucleotide difference the following miR-specific qPCR experiment was performed:

Mixed on ice:

RT mix:

- 1 µl 10x PAP buffer (Epicentre Biotechnologies, Madison, WI, US.)
- 1 µl 1 mM ATP
- 10 • 1 µl 10 µM RT-primer (L2TA: 5'-ggtagtagtttttttttttttn (SEQ ID NO 5), v designates c, g and a, n designates c, g, a and t))
- 1 µl mix of 1 mM dATP, 1 mM dCTP, 1 mM dGTP, 1 mM dTTP
- 0,5 µl 2 units/µl Transcriptor (Roche Diagnostics A/S, Hvidovre, Denmark, cat# 03 531 295 001).
- 15 • 0,2 µl (5U/µl) Poly(A) polymerase (Epicentre)
- 4.5 µl water

The following samples were prepared and incubated at 42 °C, 1 hour followed by 95 °C, 5 minutes:

Synthetic template	RT mix
1 μ l 10ng/ μ l phage MS2 RNA in TE	9 μ l
1 μ l 10^6 copies of <i>hsa-let-7a</i> ¹	9 μ l
1 μ l 10^8 copies of <i>hsa-let-7f</i> ¹	9 μ l
1 μ l 10^8 copies of <i>hsa-let-7c</i> ¹	9 μ l
1 μ l 10^6 copies of <i>hsa-let-7e</i> ¹	9 μ l

¹The synthetic templates were prepared in 10ng/μl phage MS2 RNA in TE. The synthetic templates are obtained from Integrated DNA technologies Inc., Coralville, IA, US.

For qPCR 1 µl of the polyA tailing/RT reaction was used for each PCR reaction with the *hsc*-
5 let-7a forward primer 5'-tGaGgtagtaggttg (SEQ ID NO 7) and reverse primer 5'-
cggaggtagctagttttttttttAactat (SEQ ID NO 94)

Real-time PCR was done on a ABI 7500 thermocycler by monitoring SYBR® green fluorescence as a function of PCR cycle number. The PCR reaction mixture contained:

- 10 µl 2× PCR mastermix (Roche cat#04 673 484 001)
- 1 µl or less of the RT reaction
- 1 µl TE with 2.5 µM miR-specific forward primer and 2.5 µM miR-specific reverse primer
- water to 20 µl.

The mixture was incubated at 95°C for 10 min followed 40 cycles of 95°C for 5 sec; 60°C for 15 60 sec with measurement of the fluorescence.

Amplification of the correct product was measured by comparing the melting curve profile of the PCR product to the melting curve profile of a PCR product obtained by amplification of a synthetic template.

The result of the real-time PCR experiment was analysed according to standard methods (Bustin, SA (ed.) "A-Z of Quantitative PCR," International University Line (La Jolla, California, USA), 2004).

The result is shown in table 6 and as amplification plots in Figure 4.

Table 6:

Template	C _T	% of signal on <i>hsa-let-7a</i> template ²
10ng/μl phage MS2 RNA	nd ¹	0
<i>hsa-let-7a</i>	25	100
<i>hsa-let-7f</i>	38	0.01
<i>hsa-let-7c</i>	30	4
<i>hsa-let-7e</i>	nd	0

¹nd, not detected²The percent signal was calculated as $100/2^{(C_T(\text{template}) - C_T(\text{hsa-let-7a}))}$ assuming an amplification efficiency of 100.

5 EXAMPLE 5: QUANTIFICATION OF MIRS IN HUMAN BRAIN TOTAL RNA.

In this example the copy number of miRs *hsa-let-7a*, *hsa-miR-21*, *hsa-miR-27b* and *hsa-miR-195* in total human brain RNA (Ambion) was determined.

Mixed on ice:

RT mix:

- 10
- 1 μl 10x PAP buffer (New England Biolabs)
 - 1 μl 1 mM ATP
 - 1 μl 10 μM RT-primer (L2TA: 5'-ggtagtagttttttttttttvn (SEQ ID NO 5), v designates cytosine, guanine and adenine residues, n designates cytosine, guanine, adenine and thymine residues)).
- 15
- 1 μl mix of 1 mM dATP, 1 mM dCTP, 1 mM dGTP, 1 mM dTTP
 - 0,5 μl 2 units/μl Transcriptor (Roche cat# 03 531 295 001, Roche Diagnostics A/S, Hvidovre, Denmark).
 - 0,5 μl (5U/μl) Poly(A) polymerase (New England Biolabs Inc., Ipswich, MA, US)
 - 3 μl water
- 20
- The following samples were prepared and incubated at 42 °C, 1 hour followed by 95 °C, 5 minutes:

Synthetic template/water	Phage MS2 RNA in TE (10ng/ μ l)	Human brain RNA (10ng/ μ l)	RT mix
1 μ l water	1 μ l		8 μ l
1 μ l $\sim 10^2$ copies	1 μ l		8 μ l
1 μ l $\sim 10^3$ copies	1 μ l		8 μ l
1 μ l $\sim 10^4$ copies	1 μ l		8 μ l
1 μ l $\sim 10^5$ copies	1 μ l		8 μ l
1 μ l $\sim 10^6$ copies	1 μ l		8 μ l
1 μ l $\sim 10^7$ copies	1 μ l		8 μ l
1 μ l $\sim 10^8$ copies	1 μ l		8 μ l
1 μ l $\sim 10^9$ copies	1 μ l		8 μ l
1 μ l water		1 μ l	8 μ l

Synthetic template: Equal amounts (number of copies) of synthetic *hsa-let-7a*, *hsa-miR-21*, *hsa-miR-27b* and *hsa-miR-195* in 10ng/ μ l phage MS2 RNA in TE (obtained from Integrated DNA technologies Inc).

- 5 TE: 10mM Tris/HCl (pH 8.0), 1mM EDTA.

For qPCR 1 μ l of the polyA tailing/RT reaction was used for each PCR reaction with the gene-specific primers:

Table 7:

miR	Forward primer 5' $\rightarrow$ 3'	Reverse primer 5' $\rightarrow$ 3'
<i>hsa-let-7a</i>	tGaGgtagtaggttg (SEQ ID NO 7)	tgacacggagggtactagtttttttttttAactat (SEQ ID NO 94)
<i>hsa-miR-21</i>	tAgmCttatcagactgatg (SEQ ID NO 13)	gcggtgacacggagggtactagttttttttttmCaac (SEQ ID NO 15)
<i>hsa-miR-27b</i>	ttmCamCagtggttaag (SEQ ID NO 95)	tgacacggagggtactagtttttttttttGcaga (SEQ ID NO 96)
<i>hsa-miR-195</i>	ctgtaGcaGcacagaa (SEQ ID NO 97)	tgacacggagggtactagtttttttttttGccaat (SEQ ID NO 98)

Real-time PCR was done on a ABI 7500® thermocycler by monitoring SYBR® green fluorescence as a function of PCR cycle number. The PCR reaction mixture contained:

- 10 µl 2× PCR mastermix (Roche cat#04 673 484 001)
- 1 µl or less of the RT reaction
- 5 • 1 µl TE with 2.5 µM miR-specific forward primer and 2.5 µM miR-specific reverse primer
- water to 20 µl.

The mixture was incubated at 95°C for 10 min followed 40 cycles of 95°C for 5 sec; 60°C for 60 sec with measurement of the fluorescence.

- 10 Amplification of the correct product was measured by comparing the melting curve profile of the PCR product to the melting curve profile of a PCR product obtained by amplification of a synthetic template.

- 15 For each miR, the result of the real-time PCR experiment was analysed according to standard methods (Bustin, SA (ed.) "A-Z of Quantitative PCR." International University Line (La Jolla, California, USA),, 2004) and the Ct values from the samples without human brain RNA was used to construct standard curves.

The C_t of the sample with human brain RNA was compared to the standard curve to determine the number of miRs in the sample (Bustin, SA (ed.) "A-Z of Quantitative PCR." International University Line (La Jolla, California, USA), 2004).

- 20 Results:

Table 8:

miR	Copies per 10 pg brain RNA
<i>hsa-let-7a</i>	16000
<i>hsa-miR-21</i>	400
<i>hsa-miR-27b</i>	400
<i>hsa-miR-195</i>	1000

EXAMPLE 6: DESIGN OF PREMIR QPCR ASSAYS

This example shows that the *hsa*-miR-10a assay does not detect the corresponding premiR, *hsa*-premiR-10a. Similarly, it is possible to use the primer design to make an assay that detects *hsa*-premiR-10a but not *hsa*-miR-10a.

5 Mixed on ice:

RT mix:

- 1 μ l 10x PAP buffer (New England Biolabs)
- 1 μ l 1 mM ATP
- 10 • 1 μ l 10 μ M RT-primer (L2TA: 5'-ggtagtagtntttttttttttvn (SEQ ID NO 5), v designates cytosine, guanine and adenine residues, n designates cytosine, guanine, adenine and thymine residues)).
- 1 μ l mix of 1 mM dATP, 1 mM dCTP, 1 mM dGTP, 1 mM dTTP
- 0,5 μ l 2 units/ μ l Transcriptor (Roche cat# 03 531 295 001).
- 0,5 μ l (5U/ μ l) Poly(A) polymerase (New England Biolabs)
- 15 • 3 μ l water

The following samples were prepared and incubated at 42 °C, 1 hour followed by 95 °C, 5 minutes:

RNA sample	RT mix
1 μ l ~ water	9 μ l
1 μ l ~ 10 ⁶ copies <i>hsa</i> -miR-10a	9 μ l
1 μ l ~ 10 ⁶ copies <i>hsa</i> -premiR-10a	9 μ l

20 For qPCR 1 μ l of the polyA tailing/RT reaction was used for each PCR reaction with the specific primers in Table 9.

Table 9:

Target	Forward primer (5'→3')	Reverse primer (5'→3')
<i>hsa-miR-10a</i>	tAccctGtagatccgaa (SEQ ID NO 99)	tgacacggaggtagttagttttttttttttCacaaat (SEQ ID NO 100)
<i>hsa-premiR-10a</i>	tgtGtaaggaattttgtggt (SEQ ID NO 101)	gaggtagtagtttttttttttttttAttc (SEQ ID NO 102)

Real-time PCR was done on a ABI 7500 thermocycler by monitoring SYBR® green fluorescence as a function of PCR cycle number. The PCR reaction mixture contained:

- 5
 - 10 µl 2× PCR mastermix (Roche cat#04 673 484 001)
 - 1 µl or less of the RT reaction
 - 1 µl TE with 2.5 µM miR-specific forward primer and 2.5 µM miR-specific reverse primer
 - water to 20 µl.
- 10 The mixture was incubated at 95°C for 10 min followed 40 cycles of 95°C for 5 sec; 60°C for 60 sec with measurement of the fluorescence.

For each miR, the result of the real-time PCR experiment was analysed according to standard methods (Bustin, SA (ed.) "A-Z of Quantitative PCR." International University Line, 2004).

Table 10:

RNA	<i>hsa-miR10a</i> assay	<i>hsa-premiR10a</i> assay
negative control	below detection	below detection
<i>hsa-miR10a</i>	C _T = 29.53	below detection
<i>hsa-premiR10a</i>	below detection	C _T = 30.82

15

The result shows that both the *hsa-miR-10a* and the *hsa-premiR-10a* assays detect the correct target and that there is no cross reaction to premiR-10a or miR -10a, respectively.

EXAMPLE 7: DETECTION OF PRE-MIR BY QPCR

In this example *hsa-miR-10a* and the corresponding pre-miR, *hsa-pre-miR-10a* were detected in total human RNA.

Mixed on ice:

5 **RT mix:**

- 1 µl 10x PAP buffer (New England Biolabs Inc., Ipswich, MA, US)
- 1 µl 1 mM ATP
- 1 µl 10 µM RT-primer (L2TA: 5'-ggtagctgtttttttttttttvn (SEQ ID NO 5), v designates cytosine, guanine and adenine residues, n designates cytosine, guanine, adenine and thymine residues)).
- 1 µl mix of 1 mM dATP, 1 mM dCTP, 1 mM dGTP, 1 mM dTTP
- 0,5 µl 2 units/µl Transcriptor (Roche cat# 03 531 295 001).
- 0,5 µl (5U/µl) Poly(A) polymerase (New England Biolabs)
- 3 µl water

- 15 The following samples were prepared and incubated at 42 °C, 1 hour followed by 95 °C, 5 minutes:

RNA sample	RT mix
1 µl water	9 µl
1 µl ~ 100 ng/µl RNA mix ^a	9 µl
1 µl ~ 10 ⁶ copies <i>hsa-miR-10a</i>	9 µl
1 µl ~ 10 ⁶ copies <i>hsa-premiR-10a</i>	9 µl

- ^aMixture of 1 µl of 1 µg/µl heart RNA, 1 µl of 1 µg/µl brain RNA, 1 µl of 1 µg/µl liver RNA, 1 µl of 1 µg/µl lung RNA, 1 µl of 1 µg/µl kidney RNA, 1 µl of 1 µg/µl lymph RNA, 1 µl of 1 µg/µl jejunum RNA, 1 µl of 1 µg/µl colon RNA, 1 µl of 1 µg/µl breast RNA and 1 µl of 1 µg/µl leukemia RNA in TE.

TE buffer: 10mM Tris/HCl (pH 8.0), 1mM EDTA.

For qPCR 1 μ l of the polyA tailing/RT reaction was used for each PCR reaction with the specific primers in Table 11.

Table 11:

Target	Forward primer 5' $\rightarrow$ 3'	reverse primer 5' $\rightarrow$ 3'
<i>hsa-miR-10a</i>	tAccctGtagatccgaa (SEQ ID NO 99)	tgacacggaggtagtattttttttttmCacaaat (SEQ ID NO 100)
<i>hsa-premiR-10a</i>	tgtGtaaggaattttgtggt (SEQ ID NO 101)	gaggtagtagtttttttttttttAttc (SEQ ID NO 102)

5 Real-time PCR was done on a ABI 7500 thermocycler by monitoring SYBR® green fluorescence as a function of PCR cycle number. The PCR reaction mixture contained:

- 10 μ l 2 $\times$ PCR mastermix (Roche cat#04 673 484 001)
- 1 μ l or less of the RT reaction
- 1 μ l TE with 2.5 μ M miR-specific forward primer and 2.5 μ M miR-specific reverse primer
- water to 20 μ l.

The mixture was incubated at 95°C for 10 min followed 40 cycles of 95°C for 5 sec; 60°C for 60 sec with measurement of the fluorescence.

15 For each miR, the result of the real-time PCR experiment was analysed according to standard methods (Bustin, SA (ed.) "A-Z of Quantitative PCR." International University Line, 2004). The result is shown in tab 12.

Table 12:

RNA	<i>hsa-miR10a</i> assay	<i>hsa-premiR10a</i> assay
negative control	below detection	below detection
total RNA	$C_T = 26.22$	$C_T = 37.25$
<i>hsa-miR10a</i>	$C_T = 28.83$	not testet
<i>hsa-premiR10a</i>	not testet	$C_T = 32.59$

20 The result shows that the total RNA sample contains more than 10^6 copies of *hsa-miR-10a* because the C_T is below the C_T of the sample with 10^6 copies of *hsa-miR-10a* whereas the

total RNA sample contains less than 10^6 copies of *hsa*-pre-miR-10a because the C_T is below the C_T of the sample with 10^6 copies of *hsa*-pre-miR-10a.

EXAMPLE 8: SPECIFIC DETECTION OF PRE-MIRNAS

Aim of the experiment: To determine if the universal reverse transcriptase quantitative PCR methodology (UniRT qPCR) described in this application could be used to detect pre-miR specifically without co-detection of the corresponding mature miR.

Materials: Synthetic miR 203 RNA (5'- gugaaauguuuaggaccacuag) (SEQ ID NO 103) and pre-miR 203 RNA (5'- agugguucuaacagucaacaguucugu-agcgcaauugugaaauguuuaggaccacuag) (SEQ ID NO 104) were selected as test subjects. The synthetic RNAs were synthesized by Integrated DNA technologies Inc., Coralville, IA, US. The RNAs were diluted to 1×10^6 molecules/ μ L in TEMS2 (TE buffer (10 mM Tris HCl pH 8, and 0.1 mM EDTA mixed with 10 ng/ μ L MS2 viral RNA (Roche Applied Science Inc). Primers used were miR-203.Rev (5'- tgacacggaggtagttagtttttttttttCtag) (SEQ ID NO 105), miR-203.Fwd (5'- gtGaaatGtttaggacca) (SEQ ID NO 106) and pre-miR- 203.Fwd (5'-cagttcaacagttctgtagc) (SEQ ID NO 107). The pre-miR-203 Fwd primer was designed in the loop structure of the pre-miR-203 molecule. Mature miR-203 template and pre-miR-203 were subject to reverse transcription using Universal cDNA synthesis kit (Exiqon AS, Cat.no #203300).

Mix:

- Synthetic RNA 1×10^6 molecules
- 5 x UniRT reaction buffer 2 μ L
- 10x Enzyme mixture 1 μ L
- Water up to 10 μ L

Incubate at 42°C for 60 minutes and heat denature at 85°C for 5 minutes. Dilute 10 x in water.

- 25 qPCR with SYBR Green Master Mix, UniRT (Exiqon AS, Cat.No. 203400)

Mix:

- Primer mix (3uM each) 1 μ L
- SYBR green master mix 5 μ L
- cDNA template 1 μ L
- Water 3 μ L

Two primer mixes were used: 1) miR-203 ((miR-203.Fwd and miR-203.Rev primer) and 2) pre-miR-203 (pre-miR- 203.Fwd and miR-203.Rev primer). Templates used were mir-203 and pre-mir-203. Non template control (NTC) qPCR was also run for each PCR assay. All qPCR were done in duplicates.

- 5 q-RT-PCR reactions were performed in a 384 well plate in a LightCycler 480 (Roche Diagnostics) instrument using the following PCR protocol

1. 95°C for 10 minutes
2. 95°C for 10 seconds
3. 60°C for 1 minute

- 10 Signal detection with SYBR green (HRM dye) setup. Steps 2-3 repeated for 45 times followed by melting curve analysis.

Results and discussion

- 15 Standard miR-203 primers detect the miR-203 template well with a Cp value of 28,86. Also the miR-203 assay, partially detects the pre-miR-203(Cp of 30,125) since the miR is located on the 3' end of the pre-mir-203. Pre-miR-203 assay design detects the pre-miR with the Cp values of 25,7 however the mature miR-203 is not detected with the pre-miR-203 specific assay (Cp=40). This data clearly shows that pre-mir specific assays can be designed to specifically target the pre-miR molecules.

- 20 Table 13: Detection of pre-miR-203 and miR-203 using miR specific primer and pre-mir specific primers.

Template	Assay	Mean Cp	Percent detection
NTC	miR-203		
miR-203	miR-203	28,86	100,0
pre-miR-203	miR-203	30,125	41,6
NTC	Pre-miR-203		
miR-203	Pre-miR-203	40	0,0
pre-miR-203	Pre-miR-203	25,675	100

Mean Cp represents a mean value of a duplicate qPCR reaction. Percent detection is estimated as a $=100/\text{POWER}(2;(\text{Cp_test}-\text{Cp_ref}))$ where the Cp_ref is the corresponding template and primer set Cp value.

EXAMPLE 9: MICRO-RNAS DIFFERENTIALLYLY EXPRESSED IN HEART AND LIVER TISSUES

Aim of the experiment is to determine if well expressed miRNAs that are differently expressed between heart and liver tissue can be discriminated using a qPCR array based on the SYBR green universal reverse transcriptase quantitative PCR (UniRT qPCR) method of the present invention.

Materials and methods

Total RNA from liver and whole-heart was obtained from Ambion Inc., and diluted in nuclease free water to a concentration of 10 ng/ μ L and stored at -80°C. We selected a few miRNA that are known to be differentially expressed in heart and liver tissue samples from the literature (see for example: Liang, Y., et al. (2007) BMC Genomics. 8: pp166 and Landgraf P., et al. (2007) Nature Biotechnol. (9): pp 996-7). The miR selected were hsa-miR-1, hsa-miR-126 and hsa-miR-133b (heart) and hsa-miR-192, hsa-miR-122*, hsa-miR-194 and hsa-miR-122 (liver).

Table 14: Primer sequences for the seven miR assays. Capital letters represent LNA nucleotides.

Assay	Reverse primer (5 $\rightarrow$ 3)	SEQ ID NO	Forward primer (5 $\rightarrow$ 3)	SEQ ID NO
hsa-miR-122	tgacacggaggtactagttttttttttCaaacac	108	tGgaGtgtgacaatg	115
hsa-miR-194	gtactagtttttttttttCcaca	109	tGtaaCagcaactcca	116
hsa-miR-122*	gaggtagtagtttttttttttAtttag	110	aAcgccAtTatcacact	117
hsa-miR-192	gtactagtttttttttttGgct	111	tgactGacCtatgaattgac	118
hsa-miR-126	gtactagttttttttttTcgcat	112	ctgtCgtaccgtgagta	119
hsa-miR-133b	tgacacggaggtactagttttttttttAgctg	113	tTtGgtccccttcaa	120
hsa-miR-1	tgacacggaggtactagttttttttttAtacata	114	ctgtGgAatgtaaagaagt	121

Capital letters represent LNA nucleotides.

RT reactions were performed in triplicate for each tissue and non enzyme control (NEC)) with Universal cDNA synthesis kit (Exiqon AS, Cat.no #203300)

- total RNA 20 ng
- 5 x UniRT reaction buffer 4 μ L
- Enzyme mixture 2 μ L
- Water up to 20 μ L

Incubate at 42°C for 60 minutes and heat denature at 85°C for 5 minutes. Dilute the cDNA 1/100 in nuclease free water.

Quantitative polymerase chain reaction (q-PCR), step 2 of the protocol, was performed as follows:

- 5 5 µL diluted cDNA was mixed with 5 µL SYBR Green Master Mix, UniRT (Exiqon AS, Cat.No. 203400) into the 384 well plate with desiccated primer sets of the seven miR assays described below. The plate was sealed and directly put on to the LightCycler for amplification and analysis.

- 10 q-RT-PCR reactions were performed in a LightCycler 480 (Roche Diagnostics) using the following PCR protocol:

1. 95°C for 10 minutes
2. 95°C for 10 seconds
3. 60°C for 1 minute

- 15 Signal detection with SYBR green (HRM dye) setup. Steps 2-3 repeated for 45 times followed by melting curve analysis.

Standard data analysis was performed on the LC480 raw data using the supplied data analysis software (Roche Diagnostics). Cp values were collected as Abs Quant/second derivatives max.

- 20 For this experiment of looking at examples of well expressed miRNA genes from both liver and heart, we used mean of raw data Cp values without normalization or calibration. This is done since normalization between tissues is not very accurate way of estimating miRs from different sources. Then we compare the difference between the two tissues in terms of Δ Cp values. Note that a difference of 1 in Cp value represents approximately a two fold difference in expression.

25 **Results and discussion**

We selected total of 7 miRNAs that are known previously from the literature, to be differentially expressed between heart and liver tissue. The results are shown in table 15 and fig. 7. The data obtained showed that the three genes miR-1, miR-133b and 126 are all showing much higher expression in heart samples than liver samples. Similarly the miR-192,

miR-194 and miR-122 and 122* are showing much higher expression in liver than heart. The difference is ranging from 2.6 to 12.9 Cp which corresponds to a range of from 5 fold to over 1000 fold difference in expression. In general the differentially expressed miRs described from the literature are easily distinguished by the UniRT expression platform using the assays described here.

Table 15: Expression values for the 7 miR selected and tested on the UniRT platform.

miR name	Heart (Mean Cp)	Liver (Mean Cp)	Δ Cp
hsa-miR-1	20,9	33,9	12,9
hsa-miR-133b	22,0	34,7	12,7
hsa-miR-126	22,5	25,1	2,6
hsa-miR-192	31,6	25,3	-6,3
hsa-miR-122*	37,6	30,0	-7,5
hsa-miR-194	32,3	25,9	-6,4
hsa-miR-122	31,8	21,1	-10,8

Positive Δ Cp values represent excess expression in heart and negative Cps represents excess expression in liver.

10

EXAMPLE 10: COMPARISON OF miR SPECIFIC ASSAYS DESIGNED IN PRESENT INVENTION TO COMPETING METHOD USING PURE DNA PRIMERS

The LNA based design described in the present invention was compared to a commercial DNA based product, the miScript Reverse Transcription Kit (Qiagen, Cat. no. 218060, QIAGEN GmbH, Hilden, Germany). This DNA based product also relies on a miRNA 3'- polyadenylation step followed by reverse transcription with a DNA based poly dT primer where both reactions occur in a one-tube reaction. Because both these methods use the same enzymatic steps a comparison very well illustrate the surprising advantages of the present LNA based method since the miScript does not include LNA in the primers. Another difference is that the miScript uses a reverse primer specific to the universal tag added with the RT primer whereas the LNA based reverse primer of the present method is specific to the miRNA being detected.

20

Table 16: Nucleotide sequences of compared miRNAs

miR	Sequence	SEQ ID NO	miR gc%	miR Tm
hsa-let-7a	5'-ugagguaguagguuguauaguu-3'	90	36	50
hsa-miR-143	5'-ugagaugaagcacuguagcuc-3'	122	39	54
hsa-miR-155	5'-uuaaugcuaaucgugauaggggu-3'	123	47	54
hsa-mir-1	5'-uggaauguaaagaaguauguau-3'	124	27	47

Table 17: Primer sets used in comparison

miR	Forward primer (5'→3')	SEQ ID NO	Reverse primer (5'→3')	SEQ ID NO	corres- ponding Qiagen assay cat. no.
<i>hsa-let-7a</i>	tGaGgtagtaggttg	7	cggaggtagtagtttttttttttAactat	94	MS 00006482
<i>hsa-miR-143</i>	tGaGatgaagcactg	125	tgacacggaggtagtttttttttttGag cta	126	MS 00003514
<i>hsa-miR-155</i>	gacttaaTgCtaatcgtga t	127	gtactagtttttttttttAccccta	128	MS 00003605
<i>hsa-mir-1</i>	ctgtGgAatgtaaagaagt	121	tgacacggaggtagtttttttttttAta cata	114	MS000083 58

FIRST STEP

Reverse transcription was performed on a dilution series of synthetic miRNA target (obtained from Integrated DNA technologies Inc., Coralville, IA, US.), with a background of 10 ng/μl MS2 bacteriophage RNA (Roche Applied Science, Catalog Number 10165948001), using either Universal cDNA Synthesis Kit (Exiqon A/S, Vedbaek, Denmark., Prod. No. 203300) or miScript Reverse Transcription Kit (Qiagen, Cat. no. 218060, QIAGEN GmbH, Hilden, Germany) according to the manufacturer's instructions.

10 SECOND STEP

On the cDNA obtained using the Universal cDNA Synthesis Kit (Exiqon, Prod. No. 203300), qPCR was performed using SYBR Green master mix, Universal RT (Exiqon, Prod. No. 203400) and the primer sets described in **Table 17**. On the cDNA obtained using miScript Reverse Transcription Kit (Qiagen, Cat. no. 218060), qPCR was performed using miScript SYBR Green PCR Kit (Qiagen, Cat. no. 218073). In both cases the amplification and detection was performed on a Roche LC480 LightCycler (Roche Diagnostics A/S, Hvidovre, Denmark), using the cycling conditions instructed by the manufacturer. Each cDNA/assay combination was run in triplicate.

RESULTS

20 Figure 8 shows the triplicate Cp values versus template concentration obtained in the experiment for each of the four tested miRNAs. In all four compared assays, the method of the present invention was more sensitive (black lines connecting diamonds) than the miScript

assay (gray lines connecting spheres) as indicated by the lower Cp values consistently obtained with the assay of the present invention. In the case of *hsa-let-7a*, there was a 10-fold improved sensitivity of the assay of the present invention, measured as the lowest copy-number detected quantitatively. For *hsa-miR-143* and *hsa-miR-155* the difference in

5 sensitivity was 100-fold better with the assay of the present invention. For *hsa-miR-1* with a very low gc-content (**see table 16**) and thus low melting temperature, the alternative assay was unable to detect the template even at highest concentration of RNA. The assay designed in the present invention quantitatively detected an equivalent of as little as 10 miRNA copies in the PCR reaction. This data clearly shows that the assay of the present invention is

10 surprisingly more sensitive. We ascribe the improvement in sensitivity partly to the design of the extension primer which comprise both a 5' "tag-sequence" (R^1) as well as a 3' "anchor

sequence" (R^2) and partly to the design of template-specific (e.g. miR-specific) forward and reverse primers of the qPCR reaction which includes LNA. Our results indicate that the gene specific design of the forward and reverse primers allow more sensitive detection than with

15 pure DNA primers using a universal reverse primer.

TABLE 18: Nucleotide sequences

Lower case designates natural occurring nucleotides, upper case designates LNA, mC denotes LNA methyl cytosine. v is either an adenine residue, a guanine residue, or a cytosine residue and n is either an adenine residue, a guanine residue, a cytosine residue or a thymine residue.

Primer/Sequence name	Nucleotide sequence (5' → 3')	SEQ ID NO
<i>hsa-miR-197</i> ; forward primer	tTmCaccaccttctcca	1
<i>hsa-miR-197</i> reverse primer	cttttttttttttGctgggt	2
Reverse primer R ³ sequence	Tgacacggagggtactag	3
Reverse primer tail	Tgacacggagggtactagttttttttttt	4
L2TA extension primer	Ggtactagtttttttttttvn	5
<i>hsa-let-7a</i> DNA template	tgcggtgacacggagggtactagtttttttttttaactatacaacctactaccta	6
<i>hsa-let-7a</i> forward primer	tGaGgtagtaggttg	7
<i>hsa-let-7a</i> reverse primer	Cggtgacacggagggtactagtttttttttttaactata	8
<i>hsa-let-7a</i> reverse primer	cggtgacacggagggtactagtttttttttttaamCtata	9
<i>hsa-miR-16</i> forward primer	tAgmCagcacgtaaa	10
<i>hsa-miR-16</i> reverse primer	cacggagggtactagtttttttttttcgccaat	11
<i>hsa-miR-16</i> reverse primer	cacggagggtactagtttttttttttmCgccaat	12
<i>hsa-miR-21</i> forward primer	tAgmCttatcagactgatg	13
<i>hsa-miR-21</i> reverse primer	gcggtgacacggagggtactagtttttttttttcaac	14
<i>hsa-miR-21</i> reverse primer	gcggtgacacggagggtactagtttttttttttmCaac	15
<i>hsa-miR-23a</i> forward primer	atmCamCattgccag	16
<i>hsa-miR-23a</i> reverse primer	tgacacggagggtactagtttttttttttggaatc	17
<i>hsa-miR-23a</i> reverse primer	tgacacggagggtactagtttttttttttGgaatc	18
<i>hsa-miR-150</i> forward primer	tmCtmCccaacccttg	19

<i>hsa</i> -miR-150 reverse primer	tgacacggaggtagtTTTTTTTTTTmCactggta	20
<i>hsa</i> -miR-150 reverse primer	tgacacggaggtagtTTTTTTTTTTcactggta	21
<i>hsa</i> -miR-208a forward primer	ataaGmCgAgcaaaaag	22
<i>hsa</i> -miR-208a reverse primer	mCggtgacacggaggtagtTTTTTTTTTacaagc	23
<i>hsa</i> -miR-208a reverse primer	cggtagacacggaggtagtTTTTTTTTTAcagc	24
<i>hsa</i> -miR-208a reverse primer	cggtagacacggaggtagtTTTTTTTTTTacaagc	25
<i>hsa</i> -miR-124 forward primer	taaGgcacgcggtga	26
<i>hsa</i> -miR-9 forward primer	ctgtmCtttGgttatctag	27
<i>hsa</i> -miR-181a forward primer	aamCattmCaacgctgt	28
<i>hsa</i> -miR-128 forward primer	tmCamCagtgaaccggt	29
<i>hsa</i> -miR-488 forward primer	ttGaaaGgctatttc	30
<i>hsa</i> -miR-328 forward primer	ctggccctctctgcc	31
<i>hsa</i> -miR-324 forward primer	cgcacccctagggcat	32
<i>hsa</i> -miR-331 forward primer	cTaGgtatggtccca	33
<i>hsa</i> -miR-34a forward primer	tGgcAgtgtcttagc	34
<i>hsa</i> -miR-130a forward primer	caGtGcaatgttaaag	35
<i>hsa</i> -miR-1 forward primer	tGgAatgtaaagaagt	36
<i>hsa</i> -miR-23b forward primer	atmCamCattgccag	37
<i>hsa</i> -miR-133a forward primer	tTtGgtccccttcaa	38
<i>hsa</i> -miR-133b forward primer	tTtGgtccccttcaa	39
<i>hsa</i> -miR-93 forward primer	cAaaGtgctgttcgtg	40
<i>hsa</i> -miR-181b forward primer	aamCattmCattgctgtc	41
<i>hsa</i> -miR-24 forward primer	tGgctmCagttcagca	42
<i>hsa</i> -miR-124 reverse primer	tgacacggaggtagtTTTTTTTTTTTggcat	43
<i>hsa</i> -miR-9 reverse primer	tgacacggaggtagtTTTTTTTTTTmCatacag	44

<i>hsa</i> -miR-181a reverse primer	tgacacggagggtactagttttttttttttamCtcac	45
<i>hsa</i> -miR-128 reverse primer	tgacacggagggtactagttttttttttttAaagaga	46
<i>hsa</i> -miR-488 reverse primer	tgacacggagggtactagttttttttttttGacca	47
<i>hsa</i> -miR-328 reverse primer	tgacacggagggtactagttttttttttttAcggaa	48
<i>hsa</i> -miR-324 reverse primer	tgacacggagggtactagttttttttttttAcacca	49
<i>hsa</i> -miR-331 reverse primer	tgacacggagggtactagttttttttttttGgatc	50
<i>hsa</i> -miR-34a reverse primer	tgacacggagggtactagttttttttttttamCaacca	51
<i>hsa</i> -miR-130a reverse primer	tgacacggagggtactagttttttttttttAtgc	52
<i>hsa</i> -miR-1 reverse primer	tgacacggagggtactagttttttttttttAtacata	53
<i>hsa</i> -miR-23b reverse primer	tgacacggagggtactagttttttttttttggTaatac	54
<i>hsa</i> -miR-133a reverse primer	tgacacggagggtactagttttttttttttGagctg	55
<i>hsa</i> -miR-133b reverse primer	tgacacggagggtactagttttttttttttAgctg	56
<i>hsa</i> -miR-93 reverse primer	tgacacggagggtactagttttttttttttmCtacctg	57
<i>hsa</i> -miR-181b reverse primer	tgacacggagggtactagttttttttttttAccca	58
<i>hsa</i> -miR-24 reverse primer	tgacacggagggtactagttttttttttttmCtggtc	59
<i>hsa</i> -miR-146b forward primer	ctgtGagaactgaattcca	60
<i>hsa</i> -miR-146b-3p forward primer	tgcctgtggactca	61
<i>hsa</i> -miR-429 forward primer	ctgtaatamCtGtctggta	62
<i>hsa</i> -miR-193a-3p forward primer	aactggcctacaaagtcc	63
<i>hsa</i> -miR-193a forward primer	gtctttgcgggcga	64
<i>hsa</i> -miR-193b forward primer	aamCtggccctcaaag	65
<i>hsa</i> -miR-149 forward primer	tmCtggcTccgtgtcttca	66
<i>hsa</i> -miR-452 forward primer	aamCtGttgcagaggaa	67
<i>hsa</i> -miR-452* forward primer	ctmCatmCtgcaaagaagta	68
<i>hsa</i> -miR-30b forward primer	tGtaaamCatcctacactc	69

<i>hsa</i> -miR-34b forward primer	ctgcaatcactaactccact	70
<i>hsa</i> -miR-34c forward primer	aggcaGtgtagttagctg	71
<i>hsa</i> -miR-134 forward primer	tgtgactggttgaccaga	72
<i>hsa</i> -miR-671 forward primer	ctgaGgaagccctgga	73
<i>hsa</i> -miR-769-3p forward primer	ctgggatctccggggctct	74
<i>hsa</i> -miR-146b reverse primer	cgtactagttttttttttttAgcct	75
<i>hsa</i> -miR-146b-3p reverse primer	gtactagttttttttttttmCcagaac	76
<i>hsa</i> -miR-429 reverse primer	gtactagttttttttttttAcggtt	77
<i>hsa</i> -miR-193a-3p reverse primer	gtactagttttttttttttamCtgg	78
<i>hsa</i> -miR-193a reverse primer	ggtactagttttttttttttmCatct	79
<i>hsa</i> -miR-193b reverse primer	gaggtactagttttttttttttmCatct	80
<i>hsa</i> -miR-149 reverse primer	tagtttttttttttttgggAgtg	81
<i>hsa</i> -miR-452 reverse primer	gtactagttttttttttttmCagtt	82
<i>hsa</i> -miR-452* reverse primer	gtactagttttttttttttmCactt	83
<i>hsa</i> -miR-30b reverse primer	gtactagttttttttttttaGctga	84
<i>hsa</i> -miR-34b reverse primer	gaggtactagtttttttttttAtgg	85
<i>hsa</i> -miR-34c reverse primer	tactagttttttttttttGcaatca	86
<i>hsa</i> -miR-134 reverse primer	gtactagttttttttttttTcccct	87
<i>hsa</i> -miR-671 reverse primer	gtactagttttttttttttmCtcca	88
<i>hsa</i> -miR-769-3p reverse primer	aggtactagttttttttttttAacca	89
<i>hsa</i> -let-7a	ugagguaguagguuguauaguu	90
<i>hsa</i> -let-7f	ugagguaguagauuguauaguu	91
<i>hsa</i> -let-7c	ugagguaguagguuguauaguu	92
<i>hsa</i> -let-7e	ugagguaggagguuguauaguu	93
<i>hsa</i> -let-7a reverse primer	cggaggtactagttttttttttttAactat	94

<i>hsa</i> -miR-27b forward primer	ttnCamCagtggctaag	95
<i>hsa</i> -miR-27b reverse primer	tgacacggaggtagtagttttttttttGcaga	96
<i>hsa</i> -miR-195 forward primer	ctgtaGcaGcacagaa	97
<i>hsa</i> -miR-195 reverse primer	tgacacggaggtagtagttttttttttGccaat	98
<i>hsa</i> -miR-10a forward primer	tAccctGtagatccgaa	99
<i>hsa</i> -miR-10a reverse primer	tgacacggaggtagtagttttttttttmCacaaat	100
<i>hsa</i> -pre-miR-10a forward primer	tgtGtaaggaattttgtggt	101
<i>hsa</i> -pre-miR-10a reverse primer	gaggtagtagttttttttttttAttc	102
<i>Has</i> -miR-203 RNA	gugaaauguuuaggaccacuag	103
Pre-miR-203 RNA	agugguucuuuacaguucaacaguucugu- agcgcaauugugaaauguuuaggaccacuag	104
<i>hsa</i> -miR-203 reverse primer	tgacacggaggtagtagttttttttttCtag	105
<i>hsa</i> -miR-203 forward primer	gtGaaatGtttaggacca	106
<i>hsa</i> -pre-miR-203 forward primer	cagttcaacagttctgtagc	107
<i>hsa</i> -miR-122 reverse primer	tgacacggaggtagtagttttttttttCaaacac	108
<i>hsa</i> -miR-194 reverse primer	gtactagttttttttttCcac	109
<i>hsa</i> -miR-122* reverse primer	gaggtagtagttttttttttAtttag	110
<i>hsa</i> -miR-192 reverse primer	gtactagttttttttttGgct	111
<i>hsa</i> -miR-126 reverse primer	gtactagttttttttttTcgcat	112
<i>hsa</i> -miR-133b reverse primer	tgacacggaggtagtagttttttttttAgctg	113
<i>hsa</i> -miR-1 reverse primer	tgacacggaggtagtagttttttttttAtacata	114
<i>hsa</i> -miR-122 forward primer	tGgaGgtgacaatg	115
<i>hsa</i> -miR-194 forward primer	tGtaaCagcaactcca	116
<i>hsa</i> -miR-122* forward primer	aAcgccAtTatcacact	117
<i>hsa</i> -miR-192 forward primer	tgactGacCtatgaattgac	118
<i>hsa</i> -miR-126 forward primer	ctgtCgtaccgtgagta	119

<i>hsa</i> -miR-133b forward primer	tTtGgtccccttcaa	120
<i>hsa</i> -miR-1 forward primer	ctgtGgAatgtaaagaagt	121
<i>hsa</i> -miR-143	5'-ugagaugaagcacuguagcuc-3'	122
<i>hsa</i> -miR-155	5'-uuaaugcuaaucgugauaggggu-3'	123
<i>hsa</i> -mir-1	5'-uggaauguaaagaaguauguau-3'	124
<i>hsa</i> -miR-143 forward primer	tGaGatgaagcactg	125
<i>hsa</i> -miR-143 reverse primer	tgacacggaggtagtagttttttttttttGagcta	126
<i>hsa</i> -miR-155 forward primer	gacttaaTgCtaatcgtgat	127
<i>hsa</i> -miR-155 reverse primer	gtactagtttttttttttttAccccta	128
L6TA extension primer	tgcggtgacacggaggtagtagttttttttttttVN	129

CLAIMS

1. A method for amplification of a specific RNA molecule in a sample, the method comprising the steps of:

- a) adding poly-A tails to a population of RNA molecules in a sample;
- 5 b) producing cDNA molecules of the poly-A-tailed RNA molecules using an primer in a reverse transcription reaction; and
- c) amplifying the cDNA molecules by PCR using a forward primer and a reverse primer both of which are specific for said specific RNA molecule.


2. The method of claim 1, wherein the specific RNA molecule is a small, non-coding RNA.

10 3. The method of claims 1 or 2, wherein the specific small RNA molecule is a microRNA.

4. The method according to of any of claims 1, 2 or 3, wherein the extension primer has a length in the range from 10 to 100 nucleotides.

5. The method according claim 4, wherein the extension primer has a length in the range from 15 to 45 nucleotides.

15 6. The method according to any one of claims 1-5, wherein said extension primer is a nucleotide sequence according to formula III:




wherein R^1 is a 5'-end nucleotide sequence, $(T)_y$ is a central part of y consecutive thymine residues, wherein y is an integer of from 1 to 100, and R^2 is a 3'-end nucleotide sequence.

20 7. The method according to claim 6, wherein R^1 is a nucleotide sequence with a length of from 1 to 30 nucleotides.

8. The method according to claim 7, wherein R^1 is a nucleotide sequence with a length of from 6 to 10 nucleotides.
9. The method according to claim 8, wherein R^1 is a nucleotide sequence with a length of 8 nucleotides.
- 5 10. The method according to any of claims 6 to 9, wherein y is an integer from 5 to 50.
11. The method according to claim 10, wherein y is an integer from 5 to 21.
12. The method according to claim 11, wherein y is 12, 13, 14, 15, 16, 17 or 18.
13. The method according to claim 12, wherein y is 15.
- 10 14. A method according to any one of claims 6-13, wherein R^2 is a sequence motif VN, consisting of two 3'-end terminal nucleotide residues, wherein V is either an adenine residue, a guanine residue, or a cytosine residue and N is either an adenine residue, a guanine residue, a cytosine residue or a thymine residue.
15. The method according to any one of claims 6-13, wherein R^2 is a sequence motif VNN, consisting of the three 3'-end terminal nucleotides, wherein V is either an adenine residue, a guanine residue, or a cytosine residue and N is either an adenine residue, a guanine residue, a cytosine residue or a thymine residue.
- 15 16. The method according to any one of claims 1-15, wherein the extension primer comprises at least one LNA.
17. The method according to any one of claims 1-16, wherein the forward primer has a length in the range of from 10 to 100 nucleotides.
- 20 18. The method according to any one of claims 1-17, wherein the forward primer comprises at least one LNA.
19. The method according to any one of claims 1-18, wherein the forward primer is designed to specifically hybridize to the complement DNA molecule of a target RNA molecule.

20. The method according to any one of claims 1-19, wherein the reverse primer is a nucleotide sequence according to formula II:



wherein R^3 is a 5'-end nucleotide sequence, $(T)_x$ is a central part of x consecutive thymine residues, wherein x is an integer of from 1 to 100, and R^4 is a 3'-end nucleotide sequence that specifically hybridizes to a nucleotide sequence of a target RNA molecule.

21. The method according to claim 20, wherein R^3 is a nucleotide sequence with a length of from 1 to 30 nucleotides.

22. The method according to any of claims 20 or 21, wherein x is 5 to 50.

23. The method according to claim 22, wherein x is 5 to 21.

24. The method according to claim 23, wherein x is 15.

25. The method according to any of the preceding claims wherein x of formula (II) equals y of formula (III).

26. The method according to any one of claims 20-25, wherein R^4 has a length in the range of from 1 to 10 nucleotides.

27. The method according to any one of claims 20-26, wherein R^4 comprises at least one LNA.

28. The method according to claim 27, wherein R^4 contains only one LNA.

29. The method according to any one of claims 20-28, wherein R^4 is designed to specifically hybridize to the 3'-end of a target RNA.

30. The method according to any one of claims 27-29, wherein the LNA is situated in the 5' position or the position adjacent to the 5' position of the R4 part of the reverse primer.

31. A method for measuring the amount of a target microRNA in a sample from a living organism, the method comprising the steps of:

- 5 a) amplifying the target microRNA according to the method of any one of claims 1 to 30.
- b) measuring the amount of the amplified DNA molecules.

32. The method according to claim 31, wherein the amount of the amplified DNA molecules is measured using a fluorescence-based quantitative real-time PCR method.

33. The method according to any of claims 31 or 32, wherein the amount of the amplified
10 DNA molecules is measured using SYBR[®] green dye.

34. A kit for detecting at least one target RNA comprising at least one primer set specific for the detection of a the target RNA, the primer set comprising:

a) an extension primer of formula III: $R^1-(T)_y-R^2$ (III)

wherein R^1 is a 5'-end nucleotide sequence, $(T)_y$ is a central part of y consecutive thymine
15 residues, wherein y is an integer of from 1 to 100, and R^2 is a 3'-end nucleotide sequence.

b) a reverse primer of formula II: $R^3-(T)_x-R^4$ (II)

wherein R^3 is a 5'-end nucleotide sequence, $(T)_x$ is a central part of x consecutive thymine
residues, wherein x is an integer of from 1 to 100, and R^4 is a 3'-end nucleotide sequence
that specifically hybridizes to a nucleotide sequence of a target RNA molecule.

20 c) a forward primer

35. The kit according to claim 34 for detecting at least one mammalian target microRNA comprising at least one primer set specific for the detection of a target microRNA.

36. The kit according to claim 34 or 35, wherein the forward primer has a length in the range from 10 to 100 nucleotides.
37. The kit according to any one of claim 34 to 36, wherein the forward primer comprises at least one LNA.
- 5 38. The kit according to any one of claim 34 to 37, wherein the reverse primer comprises at least one LNA.
39. The kit according to any one of claims 34 to 38, wherein the forward primer is designed to specifically hybridize to the complement DNA molecule of a target microRNA molecule.
40. The kit according to any of claims 34 to 39, wherein the reverse primer is a reverse
10 primer according to any of claims 27 to 30.
41. The kit according any of claims 34 to 40, wherein the at least one mammalian target microRNA is a human microRNA.
42. The kit according to any one of claims 34 to 41, wherein the at least one mammalian target microRNA is selected from the group consisting of all known human microRNAs
15 available from the miRBase (<http://microrna.sanger.ac.uk/>).
43. The kit according to any one of claims 34 to 41, wherein the at least one mammalian target microRNA is a murine microRNA.
44. A high-throughput method for detecting or measuring the amount of specific target microRNAs in a sample from a living organism, according to any one of claims 1 to 33, the
20 method further comprising a distribution step in which the combined poly-A-tailing and reverse transcription reaction is aliquoted into individual wells of a microtiter plate containing microRNA specific primer sets of forward and reverse primers, resulting in the steps of:
- 25 c) adding poly-A tails to a population of small RNA molecules in a sample and producing cDNA molecules of the poly-A-tailed small RNA molecules using an extension primer in a reverse transcription reaction; and
- d) pipetting aliquots of the combined poly-A-tailing and reverse transcription reaction into individual wells of a microtiter plate

- e) amplifying specific target microRNAs in individual wells of a microtiter plate containing microRNA specific primer sets
- f) measuring the amount of specific microRNA molecules in individual wells.


being performed in a fashion compatible with a high-throughput experimental set-up.

- 5 45. An oligonucleotide selected from the group of oligonucleotides disclosed in table 18.

STEP 1

PolyA-tailing
Annealing of
Extension primer

Synthesis of cDNA



STEP 2
microRNA
specific qPCR




Fig. 1




Fig. 2




Fig. 3




Fig. 4

5/10

Assay for mature miR

Poly-
Adenylation

5' - AAA..... AAAAAA-3'

Reverse
transcription
to cDNA

5' - AAA.....AAAAA-3'

<- <- NVTTT.....TTTTT

Universal RT primer

-5'

PCR with
gene
specific
primers

3' - TTT.....TTTTT

→ ---AAA.....AAAAA

<-<- ← TTT.....TTTTT

-5'


-3'

-5'


Reverse primer – specific to the 3'-end of the mature miR

Fig. 5A

6/10



7/10



8/10




Fig. 6




Fig. 7




Fig. 8