

(19)



(11)

EP 2 944 638 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

18.11.2015 Bulletin 2015/47

(51) Int Cl.:

C07D 471/04 ^(2006.01)

C07D 491/052 ^(2006.01)

C07D 519/00 ^(2006.01)

A61K 31/496 ^(2006.01)

A61K 31/437 ^(2006.01)

(21) Application number: **15167892.7**

(22) Date of filing: **26.05.2010**

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO SE SI SK SM TR**

Designated Extension States:

BA ME RS

(30) Priority: **26.05.2009 US 181203 P**

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:

10721246.6 / 2 435 432

(71) Applicant: **AbbVie Bahamas Limited**

New Providence, Nassau (BS)

(72) Inventors:

- **BRUNCKO,, Milan**
Green Oaks, IL Illinois 60048 (US)
- **DING,, Hong**
Gurnee, IL Illinois 60031 (US)
- **DOHERTY,, George**
Libertyville, IL Illinois 60048 (US)
- **ELMORE,, Steven**
Northbrook, IL Illinois 60062 (US)
- **HASVOLD,, Lisa**
Grayslake, IL Illinois 60030 (US)
- **HEXAMER,, Laura**
Grayslake, IL Illinois 60030 (US)
- **KUNZER,, Aaron R.**
Arlington, IL Illinois 60004 (US)
- **SONG,, Xiaohong**
Grayslake, IL Illinois 60030 (US)

- **SOUERS,, Andrew J.**
Libertyville, IL 60048 (US)
- **SULLIVAN,, Gerald**
Lake Villa, IL Illinois 60046 (US)
- **TAO,, Zhi-Fu**
Vernon Hills, IL 60061 (US)
- **WANG,, Gary T.**
Libertyville, IL Illinois 60048 (US)
- **WANG,, Le**
Vernon Hills, IL Illinois 60061 (US)
- **WANG,, Xilu**
Libertyville, IL Illinois 60048 (US)
- **WENDT,, Michael**
Vernon Hills, IL Illinois 60061 (US)
- **MANTEL,, Robert**
Franklin, WI Wisconsin 53132 (US)
- **HANSEN,, Todd M.**
Grayslake, IL Illinois 60030 (US)

(74) Representative: **Miller, David James**

Mathys & Squire LLP
The Shard
32 London Bridge Street
London SE1 9SG (GB)

Remarks:

- A request for correction of the description has been filed pursuant to Rule 139 EPC. A decision on the request will be taken during the proceedings before the Examining Division (Guidelines for Examination in the EPO, A-V, 3.).
- This application was filed on 15-05-2015 as a divisional application to the application mentioned under INID code 62.

(54) **APOPTOSIS-INDUCING AGENTS FOR THE TREATMENT OF CANCER AND IMMUNE AND AUTOIMMUNE DISEASES**

(57) Disclosed are compounds which inhibit the activity of anti-apoptotic Bcl-2 proteins, compositions containing the compounds and methods of treating diseases during which is expressed anti-apoptotic Bcl-2 protein.

EP 2 944 638 A1

Description

[0001] This application claims priority to United States Provisional Application Serial No. 61/181,203 filed May 26, 2009, which is incorporated by reference in its entirety.

FIELD OF THE INVENTION

[0002] This invention pertains to compounds which inhibit the activity of Bcl-2 anti-apoptotic proteins, compositions containing the compounds, and methods of treating diseases during which anti-apoptotic Bcl-2 proteins are expressed.

BACKGROUND OF THE INVENTION

[0003] Anti-apoptotic Bcl-2 proteins are associated with a number of diseases. There is therefore an existing need in the therapeutic arts for compounds which inhibit the activity of anti-apoptotic Bcl-2 proteins.


[0004] Overexpression of Bcl-2 proteins correlates with resistance to chemotherapy, clinical outcome, disease progression, overall prognosis or a combination thereof in various cancers and disorders of the immune system.

[0005] Involvement of Bcl-2 proteins in bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, prostate cancer, small cell lung cancer, spleen cancer, and the like is described in commonly-owned PCT US 2004/36770, published as WO 2005/049593, and PCT US 2004/37911, published as WO 2005/024636.

[0006] Involvement of Bcl-2 proteins in immune and autoimmune diseases is described in Current Allergy and Asthma Reports 2003, 3, 378-384; British Journal of Haematology 2000, 110(3), 584-90; Blood 2000, 95(4), 1283-92; and New England Journal of Medicine 2004, 351(14), 1409-1418. Involvement of Bcl-2 proteins in arthritis is disclosed in commonly-owned United States Provisional Patent Application Serial No. 60/988,479. Involvement of Bcl-2 proteins in bone marrow transplant rejection is disclosed in commonly-owned United States Patent Application Serial No. 11/941,196.

SUMMARY OF THE INVENTION

[0007] One embodiment of this invention, therefore, pertains to compounds or therapeutically acceptable salts, which are useful as inhibitors of anti-apoptotic Bcl-2 proteins, the compounds having Formula (I)



(I),

wherein

A^1 is N or $\text{C}(\text{A}^2)$;

A^2 is H, R^1 , OR^1 , SR^1 , $\text{S}(\text{O})\text{R}^1$, SO_2R^1 , $\text{C}(\text{O})\text{R}^1$, $\text{C}(\text{O})\text{OR}^1$, $\text{OC}(\text{O})\text{R}^1$, NHR^1 , $\text{N}(\text{R}^1)_2$, $\text{C}(\text{O})\text{NHR}^1$, $\text{C}(\text{O})\text{N}(\text{R}^1)_2$, $\text{NHC}(\text{O})\text{R}^1$, $\text{NR}^1\text{C}(\text{O})\text{R}^1$, $\text{NHC}(\text{O})\text{OR}^1$, $\text{NR}^1\text{C}(\text{O})\text{OR}^1$, $\text{NHC}(\text{O})\text{NH}_2$, $\text{NHC}(\text{O})\text{NHR}^1$, $\text{NHC}(\text{O})\text{N}(\text{R}^1)_2$, $\text{NR}^1\text{C}(\text{O})\text{NHR}^1$, $\text{NR}^1\text{C}(\text{O})\text{N}(\text{R}^1)_2$, SO_2NH_2 , SO_2NHR^1 , $\text{SO}_2\text{N}(\text{R}^1)_2$, NHSO_2R^1 , $\text{NR}^1\text{SO}_2\text{R}^1$, $\text{NHSO}_2\text{NHR}^1$, $\text{NHSO}_2\text{N}(\text{R}^1)_2$, $\text{NR}^1\text{SO}_2\text{NHR}^1$, $\text{NR}^1\text{SO}_2\text{N}(\text{R}^1)_2$, $\text{C}(\text{O})\text{NHNOH}$, $\text{C}(\text{O})\text{NHNOR}^1$, $\text{C}(\text{O})\text{NHSO}_2\text{R}^1$, $\text{C}(\text{NH})\text{NH}_2$, $\text{C}(\text{NH})\text{NHR}^1$, $\text{C}(\text{NH})\text{N}(\text{R}^1)_2$, $\text{NHSO}_2\text{NHR}^1$, $\text{NHSO}_2\text{N}(\text{CH}_3)\text{R}^1$, $\text{N}(\text{CH}_3)\text{SO}_2\text{N}(\text{CH}_3)\text{R}^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $\text{C}(\text{O})\text{H}$, CHNOH , $\text{CH}(\text{NOCH}_3)$, CF_3 , $\text{C}(\text{O})\text{OH}$, $\text{C}(\text{O})\text{NH}_2$ or $\text{C}(\text{O})\text{OR}^{1A}$;

B^1 is H, R^1 , OR^1 , SR^1 , $\text{S}(\text{O})\text{R}^1$, SO_2R^1 , $\text{C}(\text{O})\text{R}^1$, $\text{C}(\text{O})\text{OR}^1$, $\text{OC}(\text{O})\text{R}^1$, NHR^1 , $\text{N}(\text{R}^1)_2$, $\text{C}(\text{O})\text{NHR}^1$, $\text{C}(\text{O})\text{N}(\text{R}^1)_2$, $\text{NHC}(\text{O})\text{R}^1$, $\text{NR}^1\text{C}(\text{O})\text{R}^1$, $\text{NHC}(\text{O})\text{OR}^1$, $\text{NR}^1\text{C}(\text{O})\text{OR}^1$, $\text{NHC}(\text{O})\text{NH}_2$, $\text{NHC}(\text{O})\text{NHR}^1$, $\text{NHC}(\text{O})\text{N}(\text{R}^1)_2$, $\text{NR}^1\text{C}(\text{O})\text{NHR}^1$, $\text{NR}^1\text{C}(\text{O})\text{N}(\text{R}^1)_2$, SO_2NH_2 , SO_2NHR^1 , $\text{SO}_2\text{N}(\text{R}^1)_2$, NHSO_2R^1 , $\text{NR}^1\text{SO}_2\text{R}^1$, $\text{NHSO}_2\text{NHR}^1$, $\text{NHSO}_2\text{N}(\text{R}^1)_2$, $\text{NR}^1\text{SO}_2\text{NHR}^1$, $\text{NR}^1\text{SO}_2\text{N}(\text{R}^1)_2$, $\text{C}(\text{O})\text{NHNOH}$, $\text{C}(\text{O})\text{NHNOR}^1$, $\text{C}(\text{O})\text{NHSO}_2\text{R}^1$, $\text{C}(\text{NH})\text{NH}_2$, $\text{C}(\text{NH})\text{NHR}^1$, $\text{C}(\text{NH})\text{N}(\text{R}^1)_2$, $\text{NHSO}_2\text{NHR}^1$, $\text{NHSO}_2\text{N}(\text{CH}_3)\text{R}^1$, $\text{N}(\text{CH}_3)\text{SO}_2\text{N}(\text{CH}_3)\text{R}^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $\text{C}(\text{O})\text{H}$, CHNOH , $\text{CH}(\text{NOCH}_3)$, CF_3 , $\text{C}(\text{O})\text{OH}$, $\text{C}(\text{O})\text{NH}_2$ or $\text{C}(\text{O})\text{OR}^{1A}$;

D^1 is H, R^1 , OR^1 , SR^1 , $\text{S}(\text{O})\text{R}^1$, SO_2R^1 , $\text{C}(\text{O})\text{R}^1$, $\text{C}(\text{O})\text{OR}^1$, $\text{OC}(\text{O})\text{R}^1$, NHR^1 , $\text{N}(\text{R}^1)_2$, $\text{C}(\text{O})\text{NHR}^1$, $\text{C}(\text{O})\text{N}(\text{R}^1)_2$,

HNC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; E¹ is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; and Y¹ is H, CN, NO₂, C(O)OH, F, Cl, Br, I, CF₃, OCF₃, CF₂CF₃, OCF₂CF₃, R¹⁷, OR¹⁷, C(O)R¹⁷, C(O)OR¹⁷, SR¹⁷, SO₂R¹⁷, NH₂, NHR¹⁷, N(R¹⁷)₂, NHC(O)R¹⁷, C(O)NH₂, C(O)NHR¹⁷, C(O)N(R¹⁷)₂, NHS(O)R¹⁷ or NHSO₂R¹⁷; or E¹ and Y¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and A², B¹, and D¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; or Y¹ and B¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and A² D¹, and E¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; or A² and B¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and D¹, E¹, and Y¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; or A² and D¹, together with the atoms to which

- NHC(O)NHR⁷, NHC(O)CH(CH₃)NHC(O)CH(CH₃)NH₂, NHC(O)CH(CH₃)NHC(O)CH(CH₃)NHR⁷, OH, (O), C(O)OH, N₃, CN, NH₂, CF₃, CF₂CF₃, F, Cl, Br or I;
R⁶ is C₂-C₅-spiroalkyl, each of which is unsubstituted or substituted with OH, (O), N₃, CN, CF₃, CF₂CF₃, F, Cl, Br, I, NH₂, NH(CH₃) or N(CH₃)₂;
5 R^{6A} and R^{6B} are independently selected alkyl or, together with the N to which they are attached, R^{6C};
R^{6C} is aziridin-1-yl, azetidin-1-yl, pyrrolidin-1-yl or piperidin-1-yl, each having one CH₂ moiety unreplaced or replaced with O, C(O), CNOH, CNOCH₃, S, S(O), SO₂ or NH;
R⁷ is R⁸, R⁹, R¹⁰ or R¹¹;
R⁸ is phenyl, which is unfused or fused with R^{8A}; R^{8A} is benzene, heteroarene, cycloalkane, cycloalkene, hetero-
10 cycloalkane or heterocycloalkene;
R⁹ is heteroaryl, which is unfused or fused with R^{9A}; R^{9A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
R¹⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{10A};
R^{10A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
15 R¹¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R¹², OR¹², SR¹², S(O)R¹², SO₂R¹², C(O)R¹², CO(O)R¹², OC(O)R¹², OC(O)OR¹², NH₂, NH¹², N(R¹²)₂, NHC(O)R¹², NR¹²C(O)R¹², NHS(O)₂R¹², NR¹²S(O)₂R¹², NHC(O)OR¹², NR¹²C(O)OR¹², NHC(O)NH₂, NHC(O)NHR¹², NHC(O)N(R¹²)₂, NR¹²C(O)NHR¹², NR¹²C(O)N(R¹²)₂, C(O)NH₂, C(O)NHR¹², C(O)N(R¹²)₂, C(O)NHOH, C(O)NHOR¹², C(O)NHSOR¹², C(O)NR¹²SO₂R¹², SO₂NH₂, SO₂NHR¹², SO₂N(R¹²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR¹², C(N)N(R¹²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;
20 R¹² is R¹³, R¹⁴, R¹⁵ or R¹⁶;
R¹³ is phenyl, which is unfused or fused with R^{13A}; R^{13A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
R¹⁴ is heteroaryl, which is unfused or fused with R^{14A}; R^{14A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
25 R¹⁵ is cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene, each of which is unfused or fused with R^{15A}; R^{15A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
R¹⁶ is alkyl, alkenyl or alkynyl;
30 R¹⁷ is R¹⁸, R¹⁹, R²⁰ or R²¹;
R¹⁸ is phenyl, which is unfused or fused with R^{18A}; R^{18A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
R¹⁹ is heteroaryl, which is unfused or fused with R^{19A}; R^{19A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
35 R²⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl each of which is unfused or fused with R^{20A}; R^{20A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
R²¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R²², OR²², SR²², S(O)R²², SO₂R²², C(O)R²², CO(O)R²², OC(O)R²², OC(O)OR²², NH₂, NHR²², N(R²²)₂, NHC(O)R²², NR²²C(O)R²², NHS(O)₂R²², NR²²S(O)₂R²², NHC(O)OR²², NR²²C(O)OR²², NHC(O)NH₂, NHC(O)NHR²², NHC(O)N(R²²)₂, NR²²C(O)NHR²², NR²²C(O)N(R²²)₂, C(O)NH₂, C(O)NHR²², C(O)N(R²²)₂, C(O)NHOH, C(O)NHOR²², C(O)NHSO₂R²², C(O)NR²²SO₂R²², SO₂NH₂, SO₂NHR²², SO₂N(R²²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR²², C(N)N(R²²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;
40 R²² is R²³, R²⁴ or R²⁵;
R²³ is phenyl, which is unfused or fused with R^{23A}; R^{23A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
R²⁴ is heteroarene, which is unfused or fused with R^{24A}; R^{24A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
R²⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{25A};
50 R^{25A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
Z¹ is R²⁶ or R²⁷;
Z² is R²⁸, R²⁹ or R³⁰;
Z^{1A} and Z^{2A} are both absent or are taken together to form CH₂, CH₂CH₂ or Z^{12A};
Z^{12A} is C₂-C₆-alkylene having one or two CH₂ moieties replaced by NH, N(CH₃), S, S(O) or SO₂;
55 L¹ is a R³⁷, OR³⁷, SR³⁷, S(O)R³⁷, SO₂R³⁷, C(O)R³⁷, CO(O)R³⁷, OC(O)R³⁷, OC(O)OR³⁷, NHR³⁷, C(O)NH, C(O)NR³⁷, C(O)NHOR³⁷, C(O)NHSO₂R³⁷, SO₂NH, SO₂NHR³⁷, C(N)NH, C(N)NHR³⁷;
R²⁶ is phenylene, which is unfused or fused with R^{26A}; R^{26A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

- R²⁷ is heteroarylene, which is unfused or fused with R^{27A}; R^{27A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
- R²⁸ is phenylene, which is unfused or fused with R^{28A}; R^{28A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
- 5 R²⁹ is heteroarylene, which is unfused or fused with R^{29A}; R^{29A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
- R³⁰ is cycloalkylene, cycloalkenylene, heterocycloalkylene or heterocycloalkenylene, each of which is unfused or fused with R^{30A}; R^{30A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
- R³⁷ is a bond or R^{37A};
- 10 R^{37A} is alkylene, alkenylene, or alkynylene, each of which is unsubstituted or substituted with one or two or three independently selected R^{37B}, OR^{37B}, SR^{37B}, S(O)R^{37B}, SO₂R^{37B}, C(O)R^{37B}, CO(O)R^{37B}, OC(O)R^{37B}, OC(O)OR^{37B}, NH₂, NHR^{37B}, N(R^{37B})₂, NHC(O)R^{37B}, NR^{37B}C(O)R^{37B}, NHS(O)₂R^{37B}, NR^{37B}S(O)₂R^{37B}, NHC(O)OR^{37B}, NR^{37B}C(O)OR^{37B}, NHC(O)NH₂, NHC(O)NHR^{37B}, NHC(O)N(R^{37B})₂, NR^{37B}C(O)NHR^{37B}, NR^{37B}C(O)N(R^{37B})₂, C(O)NH₂, C(O)NHR^{37B}, C(O)N(R^{37B})₂, C(O)NHOH, C(O)NHOR^{37B}, C(O)NHSO₂R^{37B}, C(O)NR^{37B}SO₂R^{37B}, SO₂NH₂, SO₂NHR^{37B}, SO₂N(R^{37B})₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR^{37B}, C(N)N(R^{37B})₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I substituents;
- 15 R^{37B} is alkyl, alkenyl, alkynyl, or R^{37C};
- R^{37C} is phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl;
- Z³ is R³⁸, R³⁹ or R⁴⁰;
- 20 R³⁸ is phenyl, which is unfused or fused with R^{38A}; R^{38A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
- R³⁹ is heteroaryl, which is unfused or fused with R^{39A}; R^{39A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
- R⁴⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{40A};
- 25 R^{40A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
- wherein the moieties represented by R²⁶ and R²⁷ are substituted (i.e., if Z^{1A} and Z^{2A} are absent) or further substituted (i.e., if Z^{1A} and Z^{2A} are present) with R⁴¹, OR⁴¹, SR⁴¹, S(O)R⁴¹, SO₂R⁴¹, C(O)R⁴¹, CO(O)R⁴¹, OC(O)R⁴¹, OC(O)OR⁴¹, NHR⁴¹, N(R⁴¹)₂, NHC(O)R⁴¹, NR⁴¹C(O)R⁴¹, NHS(O)₂R⁴¹, NR⁴¹S(O)₂R⁴¹, NHC(O)OR⁴¹, NR⁴¹C(O)OR⁴¹, NHC(O)NHR⁴¹, NHC(O)N(R⁴¹)₂, NR⁴¹C(O)NHR⁴¹, NR⁴¹C(O)N(R⁴¹)₂, C(O)NHR⁴¹, C(O)N(R⁴¹)₂, C(O)NHOR⁴¹, C(O)NHSO₂R⁴¹, C(O)NR⁴¹SO₂R⁴¹, SO₂NHR⁴¹, SO₂N(R⁴¹)₂, C(N)NHR⁴¹, or C(N)N(R⁴¹)₂;
- 30 R⁴¹ is heteroaryl, which is fused with R^{43A}; R^{43A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; which is unfused or fused with benzene, heteroarene or R^{43B}; R^{43B} is cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
- wherein the cyclic moieties represented by E¹ and Y¹ together, Y¹ and B¹ together, A² and B¹ together, A² and D¹ together, R^{1A}, R², R^{2A}, R³, R^{3A}, R⁴, R^{4A}, R⁶, R^{6C}, R⁸, R^{8A}, R⁹, R^{9A}, R¹⁰, R^{10A}, R¹³, R^{13A}, R¹⁴, R^{14A}, R¹⁵, R^{15A}, R¹⁸, R^{18A}, R¹⁹, R^{19A}, R²⁰, R^{20A}, R²³, R^{23A}, R²⁴, R^{24A}, R²⁵, R^{25A}, R²⁶, R^{26A}, R²⁷, R^{27A}, R²⁸, R^{28A}, R²⁹, R^{29A}, R³⁰, R^{30A}, R^{37B}, R³⁸, R^{38A}, R³⁹, R^{39A}, R⁴⁰, and R^{40A} are independently unsubstituted, further unsubstituted, substituted or further substituted with one or two or three or four or five of independently selected R^{57A}, R⁵⁷, OR⁵⁷, SR⁵⁷, S(O)R⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, OC(O)R⁵⁷, OC(O)OR⁵⁷, NH₂, NHR⁵⁷, N(R⁵⁷)₂, NHC(O)R⁵⁷, NR⁵⁷C(O)R⁵⁷, NHS(O)₂R⁵⁷, NR⁵⁷S(O)₂R⁵⁷, NHC(O)OR⁵⁷, NR⁵⁷C(O)OR⁵⁷, NHC(O)NH₂, NHC(O)NHR⁵⁷, NHC(O)N(R⁵⁷)₂, NR⁵⁷C(O)NHR⁵⁷, NR⁵⁷C(O)N(R⁵⁷)₂, C(O)NH₂, C(O)NHR⁵⁷, C(O)N(R⁵⁷)₂, C(O)NHOH, C(O)NHOR⁵⁷, C(O)NHSO₂R⁵⁷, C(O)NR⁵⁷SO₂R⁵⁷, SO₂NH₂, SO₂NHR⁵⁷, SO₂N(R⁵⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁵⁷, C(N)N(R⁵⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;
- 40 R^{57A} is spiroalkyl, or spiroheteroalkyl;
- 45 R⁵⁷ is R⁵⁸, R⁵⁹, R⁶⁰ or R⁶¹;
- R⁵⁸ is phenyl, which is unfused or fused with R^{58A}; R^{58A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
- R⁵⁹ is heteroaryl, which is unfused or fused with R^{59A}; R^{59A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
- 50 R⁶⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{60A};
- R^{60A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
- R⁶¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶², OR⁶², SR⁶², S(O)R⁶², SO₂R⁶², C(O)R⁶², CO(O)R⁶², OC(O)R⁶², OC(O)OR⁶², NH₂, NHR⁶², N(R⁶²)₂, NHC(O)R⁶², NR⁶²C(O)R⁶², NHS(O)₂R⁶², NR⁶²S(O)₂R⁶², NHC(O)OR⁶², NR⁶²C(O)OR⁶², NHC(O)NH₂, NHC(O)NHR⁶², NHC(O)N(R⁶²)₂, NR⁶²C(O)NHR⁶², NR⁶²C(O)N(R⁶²)₂, C(O)NH₂, C(O)NHR⁶², C(O)N(R⁶²)₂, C(O)NHOH, C(O)NHOR⁶², C(O)NHSO₂R⁶², C(O)NR⁶²SO₂R⁶², SO₂NH₂, SO₂NHR⁶², SO₂N(R⁶²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶², C(N)N(R⁶²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R⁶² is R⁶³, R⁶⁴, R⁶⁵ or R⁶⁶;

R⁶³ is phenyl, which is unfused or fused with R^{63A}; R^{63A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁴ is heteroaryl, which is unfused or fused with R^{64A}; R^{64A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl, each of which is unfused or fused with R^{65A}; R^{65A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁶ is alkyl, alkenyl or alkenyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶⁷, OR⁶⁷, SR⁶⁷, S(O)R⁶⁷, SO₂R⁶⁷, C(O)R⁶⁷, CO(O)R⁶⁷, OC(O)R⁶⁷, OC(O)OR⁶⁷, NH₂, NHR⁶⁷, N(R⁶⁷)₂, NHC(O)R⁶⁷, NR⁶⁷C(O)R⁶⁷, NHS(O)₂R⁶⁷, NR⁶⁷S(O)₂R⁶⁷, NHC(O)OR⁶⁷, NR⁶⁷C(O)OR⁶⁷, NHC(O)NH₂, NHC(O)NHR⁶⁷, NHC(O)N(R⁶⁷)₂, NR⁶⁷C(O)NHR⁶⁷, NR⁶⁷C(O)N(R⁶⁷)₂, C(O)NH₂, C(O)NHR⁶⁷, C(O)N(R⁶⁷)₂, C(O)NHOH, C(O)NHOR⁶⁷, C(O)NHSO₂R⁶⁷, C(O)NR⁶⁷SO₂R⁶⁷, SO₂NH₂, SO₂NHR⁶⁷, SO₂N(R⁶⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁷, C(N)N(R⁶⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I substituents;

R⁶⁷ is alkyl, alkenyl, alkynyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; wherein the cyclic moieties represented by R^{57A}, R⁵⁸, R⁵⁹, R⁶⁰, R⁶³, R⁶⁴, R⁶⁵, and R⁶⁷ are unsubstituted or substituted with one or two or three or four of independently selected R⁶⁸, OR⁶⁸, SR⁶⁸, S(O)R⁶⁸, SO₂R⁶⁸, C(O)R⁶⁸, CO(O)R⁶⁸, OC(O)R⁶⁸, OC(O)OR⁶⁸, NH₂, NHR⁶⁸, N(R⁶⁸)₂, NHC(O)R⁶⁸, NR⁶⁸C(O)R⁶⁸, NHS(O)₂R⁶⁸, NR⁶⁸S(O)₂R⁶⁸, NHC(O)OR⁶⁸, NR⁶⁸C(O)OR⁶⁸, NHC(O)NH₂, NHC(O)NHR⁶⁸, NHC(O)N(R⁶⁸)₂, NR⁶⁸C(O)NHR⁶⁸, NR⁶⁸C(O)N(R⁶⁸)₂, C(O)NH₂, C(O)NHR⁶⁸, C(O)N(R⁶⁸)₂, C(O)NHOH, C(O)NHOR⁶⁸, C(O)NHSO₂R⁶⁸, C(O)NR⁶⁸SO₂R⁶⁸, SO₂NH₂, SONHR⁶⁸, SO₂N(R⁶⁸)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁸, C(N)N(R⁶⁸)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R⁶⁸ is R⁶⁹, R⁷⁰, R⁷¹ or R⁷²;

R⁶⁹ is phenyl, which is unfused or fused with R^{69A}; R^{69A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;


R⁷⁰ is heteroaryl, which is unfused or fused with R^{70A}; R^{70A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁷¹ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{71A}; R^{71A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁷² is alkyl, alkenyl or alkenyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁷³, OR⁷³, SR⁷³, S(O)R⁷³, SO₂R⁷³, C(O)R⁷³, CO(O)R⁷³, OC(O)R⁷³, OC(O)OR⁷³, NH₂, NHR⁷³, N(R⁷³)₂, NHC(O)R⁷³, NR⁷³C(O)R⁷³, NHS(O)₂R⁷³, NR⁷³S(O)₂R⁷³, NHC(O)OR⁷³, NR⁷³C(O)OR⁷³, NHC(O)NH₂, NHC(O)NHR⁷³, NHC(O)N(R⁷³)₂, NR⁷³C(O)NHR⁷³, NR⁷³C(O)N(R⁷³)₂, C(O)NH₂, C(O)NHR⁷³, C(O)N(R⁷³)₂, C(O)NHOH, C(O)NHOR⁷³, C(O)NHSO₂R⁷³, C(O)NR⁷³SO₂R⁷³, SO₂NH₂, SO₂NHR⁷³, SO₂N(R⁷³)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁷³, C(N)N(R⁷³)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R⁷³ is alkyl, alkenyl, alkenyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; and wherein the moieties represented by R⁶⁹, R⁷⁰, and R⁷¹ are unsubstituted or substituted with one or two or three or four of independently selected NH₂, C(O)NH₂, C(O)NHOH, SO₂NH₂, CF₃, CF₂CF₃, C(O)H, C(O)OH, C(N)NH₂, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I.

[0008] Another embodiment of this invention pertains to compounds or therapeutically acceptable salts, which are useful as inhibitors of anti-apoptotic Bcl-2 proteins, the compounds having Formula (II)



(II),

or a therapeutically acceptable salt thereof, wherein

R^{100} is as described for substituents on R^{26} ;

n is 0, 1, 2, or 3;

A^1 is N or C(A^2);

A^2 is H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$;

B^1 is H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$;

D^1 is H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$;

E^1 is H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$; and

Y^1 is H, CN, NO_2 , $C(O)OH$, F, Cl, Br, I, CF_3 , OCF_3 , CF_2CF_3 , OCF_2CF_3 , R^{17} , OR^{17} , $C(O)R^{17}$, $C(O)OR^{17}$, SR^{17} , SO_2R^{17} , NH_2 , NHR^{17} , $N(R^{17})_2$, $NHC(O)R^{17}$, $C(O)NH_2$, $C(O)NHR^{17}$, $C(O)N(R^{17})_2$, $NHS(O)R^{17}$ or $NHSO_2R^{17}$; or E^1 and Y^1 , together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

A^2 , B^1 , and D^1 are independently selected H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$; or

Y^1 and B^1 , together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

A^2 , D^1 , and E^1 are independently selected H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$; or
 A^2 and B^1 , together with the atoms to which they are attached, are benzene, naphthylene, heteroarene cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and
 D^1 , E^1 , and Y^1 are independently selected H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$; or
 A^2 and D^1 , together with the atoms to which they are attached, are benzene, naphthalene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and
 B^1 , E^1 , and Y^1 are independently selected H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$;
 R^1 is R^2 , R^3 , R^4 or R^5 ;
 R^{1A} is cycloalkyl, cycloalkenyl or cycloalkynyl;
 R^2 is phenyl, which is unfused or fused with R^{2A} ; R^{2A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^3 is heteroaryl, which is unfused or fused with R^{3A} ; R^{3A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^4 is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{4A} ;
 R^{4A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^5 is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R^6 , $NC(R^{6A})(R^{6B})$, R^7 , OR^7 , SR^7 , $S(O)R^7$, SO_2R^7 , NHR^7 , $N(R^7)_2$, $C(O)R^7$, $C(O)NH_2$, $C(O)NHR^7$, $C(O)N(R^7)_2$, $NHC(O)R^7$, $NR^7C(O)R^7$, $NHSO_2R^7$, $NHC(O)OR^7$, SO_2NH_2 , SO_2NHR^7 , $SO_2N(R^7)_2$, $NHC(O)NH_2$, $NHC(O)NHR^7$, $NHC(O)CH(CH_3)NHC(O)CH(CH_3)NH_2$, $NHC(O)CH(CH_3)NHC(O)CH(CH_3)NHR^7$, OH, (O), $C(O)OH$, N_3 , CN, NH_2 , CF_3 , CF_2CF_3 , F, Cl, Br or I;
 R^6 is C_2 - C_5 -spiroalkyl, each of which is unsubstituted or substituted with OH, (O), N_3 , CN, CF_3 , CF_2CF_3 , F, Cl, Br, I, NH_2 , $NH(CH_3)$ or $N(CH_3)_2$;
 R^{6A} and R^{6B} are independently selected alkyl or, together with the N to which they are attached, R^{6C} ;
 R^{6C} is aziridin-1-yl, azetidin-1-yl, pyrrolidin-1-yl or piperidin-1-yl, each having one CH_2 moiety unreplaced or replaced with O, $C(O)$, $CNOH$, $CNOCH_3$, S, $S(O)$, SO_2 or NH;
 R^7 is R^8 , R^9 , R^{10} or R^{11} ;
 R^8 is phenyl, which is unfused or fused with R^{8A} ; R^{8A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^9 is heteroaryl, which is unfused or fused with R^{9A} ; R^{9A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^{10} is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{10A} ;
 R^{10A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^{11} is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R^{12} , OR^{12} , SR^{12} , $S(O)R^{12}$, SO_2R^{12} , $C(O)R^{12}$, $CO(O)R^{12}$, $OC(O)R^{12}$, $OC(O)OR^{12}$, NH_2 , NHR^{12} , $N(R^{12})_2$, $NHC(O)R^{12}$, $NR^{12}C(O)R^{12}$, $NHS(O)_2R^{12}$, $NR^{12}S(O)_2R^{12}$, $NHC(O)OR^{12}$, $NR^{12}C(O)OR^{12}$, $NHC(O)NH_2$, $NHC(O)NHR^{12}$, $NHC(O)N(R^{12})_2$, $NR^{12}C(O)NHR^{12}$, $NR^{12}C(O)N(R^{12})_2$, $C(O)NH_2$, $C(O)NHR^{12}$, $C(O)N(R^{12})_2$, $C(O)NHNOH$, $C(O)NHNOR^{12}$, $C(O)NHSO_2R^{12}$, $C(O)NR^{12}SO_2R^{12}$, SO_2NH_2 , SO_2NHR^{12} , $SO_2N(R^{12})_2$, $C(O)H$, $C(O)OH$, $C(N)NH_2$, $C(N)NHR^{12}$, $C(N)N(R^{12})_2$, $CNOH$, $CNOCH_3$, OH, (O), CN, N_3 , NO_2 , CF_3 , CF_2CF_3 , OCF_3 , OCF_2CF_3 , F, Cl, Br or I;
 R^{12} is R^{13} , R^{14} , R^{15} or R^{16} ;
 R^{13} is phenyl, which is unfused or fused with R^{13A} ; R^{13A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^{14} is heteroaryl, which is unfused or fused with R^{14A} ; R^{14A} is benzene, heteroarene, cycloalkane, cycloalkene,

heterocycloalkane or heterocycloalkene;

R¹⁵ is cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene, each of which is unfused or fused with R^{15A}; R^{15A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R¹⁶ is alkyl, alkenyl or alkynyl;

5 R¹⁷ is R¹⁸, R¹⁹, R²⁰ or R²¹;

R¹⁸ is phenyl, which is unfused or fused with R^{18A}; R^{18A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R¹⁹ is heteroaryl, which is unfused or fused with R^{19A}; R^{19A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

10 R²⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl each of which is unfused or fused with R^{20A}; R^{20A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R²², OR²², SR²², S(O)R²², SO₂R²², C(O)R²², CO(O)R²², OC(O)R²², OC(O)OR²², NH₂, NHR²², N(R²²)₂, NHC(O)R²², NR²²C(O)R²², NHS(O)₂R²², NR²²S(O)₂R²², NHC(O)OR²², NR²²C(O)OR²², NHC(O)NH₂, NHC(O)NHR²², NHC(O)N(R²²)₂, NR²²C(O)NHR²², NR²²C(O)N(R²²)₂, C(O)NH₂, C(O)NHR²², C(O)N(R²²)₂, C(O)NHOH, C(O)NHOR²², C(O)NHSO₂R²², C(O)NR²²SO₂R²², SO₂NH₂, SO₂NHR²², SO₂N(R²²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR²², C(N)N(R²²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R²² is R²³, R²⁴ or R²⁵;

20 R²³ is phenyl, which is unfused or fused with R^{23A}; R^{23A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁴ is heteroarene, which is unfused or fused with R^{24A}; R^{24A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{25A};

25 R^{25A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

Z² is R²⁸, R²⁹ or R³⁰;

L¹ is a R³⁷, OR³⁷, SR³⁷, S(O)R³⁷, SO₂R³⁷, C(O)R³⁷, CO(O)R³⁷, OC(O)R³⁷, OC(O)OR³⁷, NHR³⁷, C(O)NH, C(O)NR³⁷, C(O)NHOR³⁷, C(O)NHSO₂R³⁷, SO₂NH, SO₂NHR³⁷, C(N)NH, C(N)NHR³⁷;

30 R²⁸ is phenylene, which is unfused or fused with R^{28A}; R^{28A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁹ is heteroarylene, which is unfused or fused with R^{29A}; R^{29A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁰ is cycloalkylene, cycloalkenylene, heterocycloalkylene or heterocycloalkenylenylene, each of which is unfused or fused with R^{30A}; R^{30A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

35 R³⁷ is a bond or R^{37A};

R^{37A} is alkylene, alkenylene, or alkynylene, each of which is unsubstituted or substituted with one or two or three independently selected R^{37B}, OR^{37B}, SR^{37B}, S(O)R^{37B}, SO₂R^{37B}, C(O)R^{37B}, CO(O)R^{37B}, OC(O)R^{37B}, OC(O)OR^{37B}, NH₂, NHR^{37B}, N(R^{37B})₂, NHC(O)R^{37B}, NR^{37B}C(O)R^{37B}, NHS(O)₂R^{37B}, NR^{37B}S(O)₂R^{37B}, NHC(O)OR^{37B}, NR^{37B}C(O)OR^{37B}, NHC(O)NH₂, NHC(O)NHR^{37B}, NHC(O)N(R^{37B})₂, NR^{37B}C(O)NHR^{37B}, NR^{37B}C(O)N(R^{37B})₂, C(O)NH₂, C(O)NHR^{37B}, C(O)N(R^{37B})₂, C(O)NHOH, C(O)NHOR^{37B}, C(O)NHSO₂R^{37B}, C(O)NR^{37B}SO₂R^{37B}, SO₂NH₂, SO₂NHR^{37B}, SO₂N(R^{37B})₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR^{37B}, C(N)N(R^{37B})₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I substituents;

R^{37B} is alkyl, alkenyl, alkynyl, or R^{37C};

R^{37C} is phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl;

45 Z³ is R³⁸, R³⁹ or R⁴⁰;

R³⁸ is phenyl, which is unfused or fused with R^{38A}; R^{38A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁹ is heteroaryl, which is unfused or fused with R^{39A}; R^{39A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

50 R⁴⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{40A}; R^{40A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

wherein the cyclic moieties represented by E¹ and Y¹ together, Y¹ and B¹ together, A² and B¹ together, A² and D¹ together, R^{1A}, R², R^{2A}, R³, R^{3A}, R⁴, R^{4A}, R⁶, R^{6C}, R⁸, R^{8A}, R⁹, R^{9A}, R¹⁰, R^{10A}, R¹³, R^{13A}, R¹⁴, R^{14A}, R¹⁵, R^{15A}, R¹⁸, R^{18A}, R¹⁹, R^{19A}, R²⁰, R^{20A}, R²³, R^{23A}, R²⁴, R^{24A}, R²⁵, R^{25A}, R²⁶, R^{26A}, R²⁷, R^{27A}, R²⁸, R^{28A}, R²⁹, R^{29A}, R³⁰, R^{30A}, R^{37B}, R³⁸, R^{38A}, R³⁹, R^{39A}, R⁴⁰, and R^{40A} are independently unsubstituted, further unsubstituted, substituted or further substituted with one or two or three or four or five of independently selected R^{57A}, R⁵⁷, OR⁵⁷, SR⁵⁷, S(O)R⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, OC(O)R⁵⁷, OC(O)OR⁵⁷, NH₂, NHR⁵⁷, N(R⁵⁷)₂, NHC(O)R⁵⁷, NR⁵⁷C(O)R⁵⁷, NHS(O)₂R⁵⁷, NR⁵⁷S(O)₂R⁵⁷, NHC(O)OR⁵⁷, NR⁵⁷C(O)OR⁵⁷, NHC(O)NH₂, NHC(O)NHR⁵⁷, NHC(O)N(R⁵⁷)₂,

NR⁵⁷C(O)NHR⁵⁷, NR⁵⁷C(O)N(R⁵⁷)₂, C(O)NH₂, C(O)NHR⁵⁷, C(O)N(R⁵⁷)₂, C(O)NHOH, C(O)NHR⁵⁷, C(O)NHSO₂R⁵⁷, C(O)NR⁵⁷SO₂R⁵⁷, SO₂NH₂, SO₂NHR⁵⁷, SO₂N(R⁵⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁵⁷, C(N)N(R⁵⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R^{57A} is spiroalkyl or heterospiroalkyl;

5 R⁵⁷ is R⁵⁸, R⁵⁹, R⁶⁰ or R⁶¹;

R⁵⁸ is phenyl, which is unfused or fused with R^{58A}; R^{58A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁵⁹ is heteroaryl, which is unfused or fused with R^{59A}; R^{59A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

10 R⁶⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{60A}; R^{60A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶², OR⁶², SR⁶², S(O)R⁶², SO₂R⁶², C(O)R⁶², CO(O)R⁶², OC(O)R⁶², OC(O)OR⁶², NH₂, NHR⁶², N(R⁶²)₂, NHC(O)R⁶², NR⁶²C(O)R⁶², NHS(O)₂R⁶², NR⁶²S(O)₂R⁶², NHC(O)OR⁶², NR⁶²C(O)OR⁶², NHC(O)NH₂, NHC(O)NHR⁶², NHC(O)N(R⁶²)₂, NR⁶²C(O)NHR⁶², NR⁶²C(O)N(R⁶²)₂, C(O)NH₂, C(O)NHR⁶², C(O)N(R⁶²)₂, C(O)NHOH, C(O)NHR⁶², C(O)NHSO₂R⁶², C(O)NR⁶²SO₂R⁶², SO₂NH₂, SO₂NHR⁶², SO₂N(R⁶²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶², C(N)N(R⁶²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

15 R⁶² is R⁶³, R⁶⁴, R⁶⁵ or R⁶⁶;

20 R⁶³ is phenyl, which is unfused or fused with R^{63A}; R^{63A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁴ is heteroaryl, which is unfused or fused with R^{64A}; R^{64A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{65A}; R^{65A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

25 R⁶⁶ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶⁷, OR⁶⁷, SR⁶⁷, S(O)R⁶⁷, SO₂R⁶⁷, C(O)R⁶⁷, CO(O)R⁶⁷, OC(O)R⁶⁷, OC(O)OR⁶⁷, NH₂, NHR⁶⁷, N(R⁶⁷)₂, NHC(O)R⁶⁷, NR⁶⁷C(O)R⁶⁷, NHS(O)₂R⁶⁷, NR⁶⁷S(O)₂R⁶⁷, NHC(O)OR⁶⁷, NR⁶⁷C(O)OR⁶⁷, NHC(O)NH₂, NHC(O)NHR⁶⁷, NHC(O)N(R⁶⁷)₂, NR⁶⁷C(O)NHR⁶⁷, NR⁶⁷C(O)N(R⁶⁷)₂, C(O)NH₂, C(O)NHR⁶⁷, C(O)N(R⁶⁷)₂, C(O)NHOH, C(O)NHR⁶⁷, C(O)NHSO₂R⁶⁷, C(O)NR⁶⁷SO₂R⁶⁷, SO₂NH₂, SO₂NHR⁶⁷, SO₂N(R⁶⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁷, C(N)N(R⁶⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I substituents;

30 R⁶⁷ is alkyl, alkenyl, alkynyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; wherein the cyclic moieties represented by R^{57A}, R⁵⁸, R⁵⁹, R⁶⁰, R⁶³, R⁶⁴, R⁶⁵, and R⁶⁷ are unsubstituted or substituted with one or two or three or four of independently selected R⁶⁸, OR⁶⁸, SR⁶⁸, S(O)R⁶⁸, SO₂R⁶⁸, C(O)R⁶⁸, CO(O)R⁶⁸, OC(O)R⁶⁸, OC(O)OR⁶⁸, NH₂, NHR⁶⁸, N(R⁶⁸)₂, NHC(O)R⁶⁸, NR⁶⁸C(O)R⁶⁸, NHS(O)₂R⁶⁸, NR⁶⁸S(O)₂R⁶⁸, NHC(O)OR⁶⁸, NR⁶⁸C(O)OR⁶⁸, NHC(O)NH₂, NHC(O)NHR⁶⁸, NHC(O)N(R⁶⁸)₂, NR⁶⁸C(O)NHR⁶⁸, NR⁶⁸C(O)N(R⁶⁸)₂, C(O)NH₂, C(O)NHR⁶⁸, C(O)N(R⁶⁸)₂, C(O)NHOH, C(O)NHR⁶⁸, C(O)NHSO₂R⁶⁸, C(O)NR⁶⁸SO₂R⁶⁸, SO₂NH₂, SO₂NHR⁶⁸, SO₂N(R⁶⁸)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁸, C(N)N(R⁶⁸)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

35 R⁶⁸ is R⁶⁹, R⁷⁰, R⁷¹ or R⁷²;

R⁶⁹ is phenyl, which is unfused or fused with R^{69A}; R^{69A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁷⁰ is heteroaryl, which is unfused or fused with R^{70A}; R^{70A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

45 R⁷¹ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{71A}; R^{71A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁷² is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁷³, OR⁷³, SR⁷³, S(O)R⁷³, SO₂R⁷³, C(O)R⁷³, CO(O)R⁷³, OC(O)R⁷³, OC(O)OR⁷³, NH₂, NHR⁷³, N(R⁷³)₂, NHC(O)R⁷³, NR⁷³C(O)R⁷³, NHS(O)₂R⁷³, NR⁷³S(O)₂R⁷³, NHC(O)OR⁷³, NR⁷³C(O)OR⁷³, NHC(O)NH₂, NHC(O)NHR⁷³, NHC(O)N(R⁷³)₂, NR⁷³C(O)NHR⁷³, NR⁷³C(O)N(R⁷³)₂, C(O)NH₂, C(O)NHR⁷³, C(O)N(R⁷³)₂, C(O)NHOH, C(O)NHR⁷³, C(O)NHSO₂R⁷³, C(O)NR⁷³SO₂R⁷³, SO₂NH₂, SO₂NHR⁷³, SO₂N(R⁷³)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁷³, C(N)N(R⁷³)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

55 R⁷³ is alkyl, alkenyl, alkynyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; and wherein the moieties represented by R⁶⁹, R⁷⁰, and R⁷¹ are unsubstituted or substituted with one or two or three or four of independently selected NH₂, C(O)NH₂, C(O)NHOH, SO₂NH₂, CF₃, CF₂CF₃, C(O)H, C(O)OH, C(N)NH₂, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I.

Another embodiment pertains to compounds of Formula (I) wherein A¹ is N or C(A²); A² is H, and G¹ is H.

Another embodiment pertains to compounds of Formula (I) wherein A¹ is N or C(A²); A² is H; G¹ is H, and B¹ is NHR¹.

Another embodiment pertains to compounds of Formula (I) wherein A¹ is N or C(A²); A² is H; G¹ is H, B¹ is NHR¹; and D¹ is H.

5 Another embodiment pertains to compounds of Formula (I) wherein A¹ is N or C(A²); A² is H; G¹ is H, B¹ is NHR¹; D¹ is H; and E¹ is H.

[0009] Another embodiment pertains to compounds of Formula (I) wherein A¹ is N or C(A²); A² is H; G¹ is H, B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂.

10 **[0010]** Still another embodiment pertains to compounds having Formula (I) or Formula (II), which are

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({4-[(3-morpholin-4-ylpropyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(1-methylpiperidin-4-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

20 4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(4-methylpiperazin-1-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

25 Trans-4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(2-methoxyethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(3S)-tetrahydro-2H-pyran-3-ylmethyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

30 4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(3R)-tetrahydro-2H-pyran-3-ylmethyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

35 4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-(2-naphthylsulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(2-methoxyethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

40 4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

45 4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(1,4-dioxan-2-ylmethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(2,2,2-trifluoroethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(3,3,3-trifluoropropyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

50 4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(2S)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

Cis-4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(4-methoxycyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(2R)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

55 4-{4-[(4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl)methyl]piperazin-1-yl}-N-({4-[(1,4-dioxan-2-ylmethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl)methyl]piperazin-1-yl}-N-({4-[(1,4-dioxan-2-yl-

methoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[(4-methoxycyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-(aminocarbonyl)-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-([5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-(trifluoromethyl)phenyl]sulfonyl)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl)benzamide;
 Trans-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(1-methylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5-([4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino)sulfonyl)-2-(tetrahydro-2H-pyran-4-ylmethoxy)nicotinamide;
 N-([5-bromo-6-[(1-methylpiperidin-4-yl)methoxy]pyridin-3-yl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1-methylpiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-cyano-6-(1,4-dioxan-2-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-bromo-6-(1,4-dioxan-2-ylmethoxy)pyridin-3-yl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2,2-dimethyltetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-5-cyano-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({4-[(1-acetyl)piperidin-4-yl]amino]-3-nitrophenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([2-chloro-5-fluoro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(3-morpholin-4-ylpropyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-cyano-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-cyano-6-(2-morpholin-4-ylethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-[[2-(2-methoxyethoxy)ethyl]sulfonyl]phenyl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[2-(2-methoxyethoxy)ethyl]sulfonyl]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)oxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-({5-bromo-6-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino}pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2-cyanoethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 Cis-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-yl)cyclohexyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-N-({4-({4-[bis(cyclopropylmethyl)amino]cyclohexyl)amino}-3-nitrophenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[[1-methylpiperidin-4-yl]methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(morpholin-3-ylmethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-methylpiperazin-1-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-yl)but-2-ynyl]oxy}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-ethynyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-oxo-3,4-dihydroquinazolin-6-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 Trans-4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-yl)cyclohexyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[[4-methoxycyclohexyl]methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-({4-[[4-methoxycyclohexyl]methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 tert-butyl 3-({4-({4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl}amino)sulfonyl)-2-nitrophenoxymethylmorpholine-4-carboxylate;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-(morpholin-3-ylmethoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[[1-(methylsulfonyl)piperidin-4-yl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[[1,1-dioxidotetrahydro-2H-thiopyran-4-yl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({4-chloro-3-nitrophenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[[1-(2,2,2-trifluoroethyl)piperidin-4-yl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-cyano-6-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]oxy}pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-isopropyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 N-({3-chloro-5-fluoro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

and therapeutically acceptable salts, and metabolites thereof.

55

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-([1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl]amino)-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yl oxy)benzamide;
 N-({5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-

dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[1-(2,2-difluoroethyl)piperidin-4-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[[1-(cyclopropyl)piperidin-4-yl]amino]-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[[1-(morpholin-4-yl)cyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[[4-(dicyclopropylamino)cyclohexyl]amino]-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-6,6-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-bromo-6-[(4-ethylmorpholin-3-yl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[(4-ethylmorpholin-3-yl)methoxy]-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(4-tetrahydro-2H-pyran-4-ylmorpholin-3-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([3-nitro-4-[[[(3S)-1-tetrahydro-2H-pyran-4-yl]piperidin-3-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({5-cyano-6-[(tetrahydro-2H-pyran-4-ylmethyl)amino]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[[1,1-dioxidothiomorpholin-4-yl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([4-[[[(4-aminotetrahydro-2H-pyran-4-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({5-cyano-6-[(4-morpholin-4-yl)cyclohexyl]amino]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-cyano-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([4-[[[(1S,3R)-3-morpholin-4-yl]cyclopentyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([4-[[[(1R,3S)-3-morpholin-4-yl]cyclopentyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[(morpholin-2-ylmethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydrofuran-3-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([4-[[1-[cis-3-fluorotetrahydro-2H-pyran-4-yl]piperidin-4-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(1-tetrahydro-2H-pyran-4-ylazetidin-3-yl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(1-tetrahydrofuran-3-ylazetidin-3-yl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([3-nitro-4-[[[(3R)-1-tetrahydro-2H-pyran-4-yl]pyrrolidin-3-yl]methyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[[4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-4-((trans-4-hydroxycyclohexyl)methoxy)-3-nitrophenyl)sulfonyl)benzamide;
 50 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-4-((cis-4-methoxycyclohexyl)methoxy)-3-nitrophenyl)sulfonyl)benzamide;
 Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([4-[[4-(cyclopropylamino)cyclohexyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([3-nitro-4-[[4-(tetrahydro-2H-pyran-4-ylamino)cyclohexyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 Trans-N-({5-bromo-6-[(4-morpholin-4-yl)cyclohexyl]oxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[[4-methoxycy-

clohexyl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 tert-butyl 4-{[4-({[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino}sulfonyl]-2-nitrophenyl)methyl]-4-fluoropiperidine-1-carboxylate;
 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-[(4-fluoropiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 Trans-4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[4-(4-tetrahydro-2H-pyran-4-yl)piperazin-1-yl]cyclohexyl}amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl]methoxy}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(3S)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3S)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-methylmorpholin-2-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(2-methoxyethyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(4-{[(4-acetylmorpholin-2-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[trans-4-(fluoromethyl)-1-oxetan-3-ylpyrrolidin-3-yl]methoxy}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-fluorotetrahydro-2H-pyran-4-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(1-oxetan-3-yl)piperidin-4-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[1-cyclobutylpiperidin-4-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3S)-1-cyclopropylpyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(1-tetrahydrofuran-3-yl)piperidin-4-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-cyclopropylpyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(3S)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]methyl}amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3-hydroxy-2,2-dimethylpropyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(1-(methylsulfonyl)piperidin-3-yl]methyl}amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(4-{[(1-acetyl)piperidin-3-yl]methyl}amino)-3-nitrophenyl)sulfonyl]-4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(methylsulfonyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{[2-(4-chlorophenyl)-3,3-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[1-[2-fluoro-1-(fluoromethyl)ethyl]azetidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(1-(methylsulfonyl)pyrrolidin-3-yl]methyl}amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(4-{[(1-acetylpyrrolidin-3-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

clohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(4-[[[(3R)-1-acetylpyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[3-methoxy-2,2-dimethylpropyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(1R,3R)-3-hydroxycyclopentyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(1S,3S)-3-hydroxycyclopentyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(1S,3R)-3-hydroxycyclopentyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(1R,3S)-3-hydroxycyclopentyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[(3-nitro-4-[[[(3S)-2-oxopiperidin-3-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(1-[2-fluoro-1-(fluoromethyl)ethyl]azetidin-3-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[(3-nitro-4-[[[(1-oxetan-3-yl)azetidin-3-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[(3-nitro-4-[[[(1-oxetan-3-yl)piperidin-4-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(1-cyclopropyl)piperidin-4-yl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(2-fluoroethyl)morpholin-2-yl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(2-difluoroethyl)morpholin-2-yl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(4-fluoro-1-oxetan-3-yl)piperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(2S)-4,4-difluoro-1-oxetan-3-yl]pyrrolidin-2-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[(3-nitro-4-[[[(4-tetrahydro-2H-pyran-4-yl)morpholin-3-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(4-cyclobutyl)morpholin-3-yl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[(3-nitro-4-[[[(4-tetrahydrofuran-3-yl)morpholin-3-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(1-cyclopropyl-4-fluoropiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(4-methoxybenzyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[(3-nitro-4-[[[(3-(trifluoromethoxy)benzyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(3-methoxybenzyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(4-difluoromethoxy)benzyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[(1,4-dioxaspiro[4.5]dec-8-ylamino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-N-[(4-[[[(4-(acetylamino)cyclohexyl]amino)-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(3S)-1-(2-fluoroethyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[[[(3S)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

thyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3R)-1-(2-fluoroethyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3S)-1-oxetan-3-yl]pyrrolidin-3-yl]methoxy]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-hydroxybenzyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[3-hydroxybenzyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[3-(difluoromethoxy)benzyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[cis-3-morpholin-4-yl]cyclopentyl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-[(methylsulfonyl)amino]cyclohexyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-(cyclopropylpiperidin-4-yl)amino]-3-[[trifluoromethyl]sulfonyl]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[1-oxetan-3-yl]piperidin-4-yl]methoxy]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-fluoro-1-tetrahydro-2H-pyran-4-yl]piperidin-4-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-fluoro-1-tetrahydrofuran-3-yl]piperidin-4-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-fluoro-1-(methylsulfonyl)piperidin-4-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3R)-1-oxetan-3-yl]pyrrolidin-3-yl]methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-hydroxycyclohexyl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-[3-(dimethylamino)propoxy]benzyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(2-morpholin-4-ylethoxy)benzyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[4-(E)-4-hydroxy-1-adamantyl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[4-(Z)-4-hydroxy-1-adamantyl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[4-[[[1S,4S)-bicyclo[2.2.1]hept-5-en-2-yl]methoxy]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-methyl-5-oxopyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(1R,4R,SR,6S)-5,6-dihydroxybicyclo[2.2.1]hept-2-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(1R,4R,5S,6R)-5,6-dihydroxybicyclo[2.2.1]hept-2-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[3-oxocyclohexyl]methoxy]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohexa-1,3-dien-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[tetrahydro-2H-pyran-4-yl]methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3R)-1-[2-fluoro-1-(fluoromethyl)ethyl]pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[6-[[4-fluorotetrahydro-2H-pyran-4-yl]methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3S)-1-oxetan-3-yl]pyrrolidin-3-yl]methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-N-[[5-chloro-6-[[4-methoxycyclohexyl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-N-[[5-chloro-6-[[4-methoxycyclohexyl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

clohex-1-en-1-yl)methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[(3-nitro-4-[(3S)-1-oxetan-3-ylpyrrolidin-3-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{4-[(4-{2-(2-methoxy-
 5 yethoxy)ethyl)morpholin-2-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[4-[(4-(cyanomethyl)morpholin-2-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[[4-[(N,N-dimethylgly-
 10 cyl)morpholin-2-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 (2-[[[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino)methyl)morpholin-4-yl)acetic acid;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[(3-nitro-4-[(4-(oxetan-3-yl)morpholin-2-yl)methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[(4-[(4-cyclopropylmorpholin-2-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[(5-(methylsulfonyl)-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{4-[(4-fluorotetrahydro-2H-
 20 pyran-4-yl)methoxy]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{4-[(4-methyltetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 ethyl 4-(4-{[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)piperazine-1-carboxylate;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[(4-[4-(morpholin-4-yl)piperidin-1-yl]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[(3-nitro-4-[(3R)-1-(oxetan-3-yl)pyrrolidin-3-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[(4-[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]amino)-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{4-[(1-isopropylpiperidin-4-yl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{4-[(1-tert-butylpiperidin-4-yl)amino]-3-nitrophenyl)sulfonyl}-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[[4-[(1-(2-methoxyethyl)piperidin-3-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[[4-[(1-(cyanomethyl)piperidin-3-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]amino)pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 tert-butyl 4-[(4-{[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)amino]piperazine-1-carboxylate;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{3-(pentafluoro- λ^6 -sulfanyl)-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{4-[(4-methoxytetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[(4-[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[(3-nitro-4-[(4-(oxetan-3-yl)piperazin-1-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[(3-nitro-4-[(4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{3-nitro-4-[(3R)-tetrahydrofuran-3-ylamino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[(4-[(4,4-difluorocy-

clohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({4-[(1-tert-butylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl}-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({4-[(4-oxetan-3-yl)morpholin-2-yl]methyl]amino)-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 N-[(5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methyl]amino)pyridin-3-yl)sulfonyl]-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(1-cyclopropylpiperidin-4-yl)amino]pyridin-3-yl)sulfonyl}-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 N-[(5-chloro-6-[(2S)-4-(cyanomethyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(2R)-4-(cyanomethyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 N-[(5-chloro-6-[(2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({5-fluoro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 N-({5-chloro-6-[3-hydroxy-2-(hydroxymethyl)-2-methylpropoxy]pyridin-3-yl)sulfonyl}-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({4-[(1,3-difluoropropan-2-yl)morpholin-2-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 N-[(5-chloro-6-[(1-(cyanomethyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({4-[(3R)-1-(2-(2-methoxyethoxy)ethyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({4-[(3R)-1-(N,N-dimethylglycyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-[(3-nitro-4-[(1-oxetan-3-yl)azetididin-3-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(1-(cyanomethyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({4-[(2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({4-[(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 N-[(5-chloro-6-[(1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({4-[(3R)-1-(cyanomethyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-methoxypiperidin-1-yl}-N-({5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-methoxypiperidin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-{4-[[9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[[9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl}-N-({5-chloro-6-(tetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-{4-[[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl}-N-({5-chloro-6-(tetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl}-4-{4-[[2-(4-chlorophenyl)-4,4-

dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[1-(N,N-dimethylglycyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-fluoropiperidin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-fluoropiperidin-1-yl}-N-({5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}-3-(trifluoromethyl)phenyl)sulfonyl)benzamide;
 N-[(5-chloro-6-[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[2-(tetrahydrofuran-3-yloxy)ethoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[(trans-4-cyanocyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({5-cyano-6-[2-(tetrahydro-2H-pyran-4-yl)ethoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-(3-furylmethoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({3-chloro-4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([3-cyano-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[1-(2,2-difluoroethyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({3-chloro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(4,4-difluorocyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([6-[[1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl]methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({5-chloro-6-[2-(tetrahydrofuran-2-yl)ethoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]-3-methylpiperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([4-[(3-(cyclopropylamino)propyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-(2-methoxyethoxy)pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([5-fluoro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-[[1-(methoxyacetyl)piperidin-4-yl]methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-[[1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohexyl]methyl]piperidin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([6-(tetrahydro-2H-pyran-4-ylmethoxy)-5-(trifluoromethyl)pyridin-3-yl]sulfonyl)benzamide;

N-({5-chloro-6-[(trans-4-hydroxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-cyano-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({6-[(trans-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({6-[(cis-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(4,4-difluoro-1-hydroxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperidin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 N-([3-chloro-4-[(trans-4-(morpholin-4-yl)cyclohexyl)methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({3-[cyclopropyl(1,3-thiazol-5-yl)methyl]amino}propyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 N-({3-chloro-4-[(trans-4-hydroxycyclohexyl)methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-chloro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-(trifluoromethyl)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({3-[cyclopropyl(2,2,2-trifluoroethyl)amino]propyl}amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-({1-(oxetan-3-yl)piperidin-4-yl)methoxy}phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3,5-difluoro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({3-[cyclopropyl(oxetan-3-yl)amino]propyl}amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-({1-(1-methyl-L-prolyl)piperidin-4-yl)methoxy}phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3,4-difluoro-5-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-chloro-6-({(2S)-4-cyclopropylmorpholin-2-yl)methoxy}pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperidin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperidin-1-yl)-N-([3-chloro-4-(tetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 methyl 2-([4-([4-([4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl)-2-nitrophenyl)amino]methyl)morpholine-4-carboxylate;
 40 2-([4-([4-([4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl)-2-nitrophenyl)amino]methyl)-N-ethyl-N-methylmorpholine-4-carboxamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({4-(methylsulfonyl)morpholin-2-yl)methyl}amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({3-[cyclobutyl(cyclopropyl)amino]propyl}amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-([4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-chloro-4-(tetrahydrofuran-3-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({(trans-4-hydroxycyclohexyl)methyl}amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({3-chloro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-4-(4-[[9-(4-chlorophenyl)-3-(oxetan-3-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({(2R)-4-cyclopropylmorpholin-2-yl)methyl}amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(2S)-4-cyclopropylmorpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl}piperazin-1-yl)-N-{3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 N-{[5-chloro-6-({4-[cyclopropyl(oxetan-3-yl)amino]cyclohexyl}methoxy)pyridin-3-yl]sulfonyl}-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl}piperazin-1-yl)-N-{[4-({[(4-cyclopropylmorpholin-2-yl)methyl]amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 N-{[3-chloro-4-[(4-cyclopropylmorpholin-2-yl)methoxy]phenyl]sulfonyl}-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; N-{[3-chloro-4-({[(4-cyclopropylmorpholin-2-yl)methyl]amino}phenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 2-({[2-chloro-4-({4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl}phenyl)amino)methyl}-N-ethyl-N-methylmorpholine-4-carboxamide;
- 15 (2S)-2-({[3-chloro-5-({4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl}pyridin-2-yl)oxy)methyl}-N-ethyl-N-methylmorpholine-4-carboxamide;
- N-{[5-chloro-6-({(4-cyclopropylmorpholin-2-yl)methyl}amino)pyridin-3-yl]sulfonyl}-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 2-({[3-chloro-5-({4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl}pyridin-2-yl)amino)methyl}-N-ethyl-N-methylmorpholine-4-carboxamide;
- 20 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(cis-4-hydroxy-4-methylcyclohexyl)methyl]amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 25 N-{[5-chloro-6-({[(1R,2R,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({4-[(2-cyanoethyl)(cyclopropyl)amino]cyclohexyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 30 N-{[5-chloro-6-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl]sulfonyl}-4-(4-{[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[5-chloro-6-(5,6,7,8-tetrahydroimidazo[1,2-a]pyridin-6-ylmethoxy)pyridin-3-yl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 N-{[5-chloro-6-({[(1R,2S,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(cis-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 N-{[5-chloro-6-({4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({4-(3,3-difluoropyrrolidin-1-yl)cyclohexyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-{[5-chloro-6-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl]sulfonyl}-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 N-{[5-chloro-6-[(cis-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl]sulfonyl}-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({4-[(2,2-difluorocyclopropyl)amino]cyclohexyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 N-{[5-chloro-6-[(cis-1-fluoro-4-hydroxycyclohexyl)methoxy]pyridin-3-yl]sulfonyl}-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[3-nitro-4-(2-oxaspiro[3.5]non-7-ylmethoxy)phenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 4-(4-{[2-(4-chlorophenyl)-5,5-bis(fluoromethyl)cyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({4-cyclopropylmorpholin-2-

yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(trans-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(cis-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-cyano-4-{[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(trans-4-ethyl-4-hydroxycyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(cis-4-ethyl-4-hydroxycyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-5-(methoxymethyl)-5-methylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-[(2S)-4-(oxetan-3-yl)morpholin-2-yl]methyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({3-chloro-4-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]phenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[2-(4-cyanoethyl)(cyclopropyl)amino]-1-fluorocyclohexyl)methoxy}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({5-nitro-6-[(tetrahydro-2H-pyran-4-yl)methyl]amino}pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(2-oxaspiro[3.5]non-7-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-cyano-4-methylcyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 {[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]}{4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl}amino)methyl pivalate;
 {[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]}{4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl}amino)methyl butyrate;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-[(3-(tetrahydro-2H-pyran-4-yl)methyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-6-yl)sulfonyl]benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(6-{[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino}-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(5-cyano-6-{[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy}pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-4-{[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl}-2-nitrophenyl)morpholine-4-carboxamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-(methoxymethyl)cyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(5-chloro-6-{[1-(1,3-thiazol-2-yl)piperidin-4-yl]methoxy}pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(6-{[(cis-4-hydroxy-4-methylcyclohexyl)methyl]amino}-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; and therapeutically acceptable salts, and metabolites thereof.

[0011] Another embodiment pertains to the compound 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

[0012] Another embodiment pertains to the compound Trans-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

[0013] Another embodiment pertains to the compound Cis-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-methoxycyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

[0014] Another embodiment pertains to the compound Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

[0015] Another embodiment pertains to the compound 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

[0016] Another embodiment pertains to the compound 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([trans-4-hydroxy-4-methylcyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

[0017] Another embodiment pertains to the compound 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([cis-4-hydroxy-4-methylcyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

[0018] Another embodiment pertains to the compound N-([5-chloro-6-([4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

[0019] Another embodiment pertains to the compound N-([5-bromo-6-([1-tetrahydro-2H-pyran-4-yl]piperidin-4-yl)amino)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0020] Another embodiment pertains to the compound 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3R)-1-(methylsulfonyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0021] Another embodiment pertains to the compound 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-cyclopropyl(oxetan-3-yl)amino]propyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0022] Another embodiment pertains to the compound 4-(4-([2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([tetrahydro-2H-pyran-4-ylmethyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0023] Still another embodiment pertains to compounds having Formula (I) or Formula (II), which are 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([trans-4-hydroxy-4-methylcyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3R)-1-(methylsulfonyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-cyclopropyl(oxetan-3-yl)amino]propyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0024] Still another embodiment pertains to compounds having Formula (I) or Formula (II), which are N-([5-chloro-6-([1-(cyanomethyl)piperidin-4-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-cyclopropyl(oxetan-3-yl)amino]propyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0025] Still another embodiment pertains to compounds having Formula (I) or Formula (II), which are 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([trans-4-hydroxy-4-methylcyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([cis-4-hydroxy-4-methylcyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

[0026] Another embodiment pertains to the compound 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-cyclopropyl(oxetan-3-yl)amino]propyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0027] Still another embodiment pertains to compounds having Formula (I) or Formula (II), which are 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([3R)-1-tetrahydro-2H-pyran-4-yl]pyrrolidin-3-yl)methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methylmorpholin-2-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0028] Still another embodiment pertains to compounds having Formula (I) or Formula (II), which are

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([3S)-tetrahydro-2H-pyran-3-ylmethyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-({5-bromo-6-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino}pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(4-{[(4-aminotetrahydro-2H-pyran-4-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[3-nitro-4-({[(3R)-1-tetrahydro-2H-pyran-4-yl]pyrrolidin-3-yl)methyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-methylmorpholin-2-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(methylsulfonyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-methoxytetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-{[(1-cyanomethyl)piperidin-4-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3-cyclopropyl(oxetan-3-yl)amino]propyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-{[(1R,2S,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(6-{[(cis-4-hydroxy-4-methylcyclohexyl)methyl]amino}-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0029] Another embodiment pertains to the compound N-[(5-chloro-6-{[(1-cyanomethyl)piperidin-4-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0030] Still another embodiment pertains to compounds having Formula (I) or Formula (II), which are N-[(4-{[(4-aminotetrahydro-2H-pyran-4-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-methoxytetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0031] Another embodiment pertains to the compound 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-methoxytetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

[0032] Another embodiment pertains to a composition for treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, chronic lymphocytic leukemia, myeloma, prostate cancer, small cell lung cancer or spleen cancer, said composition comprising an excipient and a therapeutically effective amount of a compound of Formula (I) or Formula (II).

[0033] Another embodiment pertains to a method of treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, chronic lymphocytic leukemia, myeloma, prostate cancer, small cell lung cancer or spleen cancer in a patient, said method comprising administering to the patient a therapeutically effective amount of a compound of Formula (I) or Formula (II).

[0034] Another embodiment pertains to a method of treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, chronic lymphocytic leukemia, myeloma, prostate cancer, small cell lung cancer or spleen cancer in a patient, said method comprising administering to the patient therapeutically effective amount of the compound of Formula (I) or Formula (II) and a therapeutically effective amount of one additional therapeutic agent or more than one additional therapeutic agent.

DETAILED DESCRIPTION OF THE INVENTION

[0035] Variable moieties herein are represented by identifiers (capital letters with numerical and/or alphabetical superscripts) and may be specifically embodied.

[0036] It is meant to be understood that proper valences are maintained for all moieties and combinations thereof, that monovalent moieties having more than one atom are drawn from left to right and are attached through their left ends, and that divalent moieties are also drawn from left to right.

[0037] It is also meant to be understood that a specific embodiment of a variable moiety herein may be the same or different as another specific embodiment having the same identifier.

The term "alkenyl" as used herein, means a straight or branched hydrocarbon chain containing from 2 to 10 carbons and containing at least one carbon-carbon double bond. The term " C_x-C_y alkyl" means a straight or branched hydrocarbon chain containing at least one carbon-carbon double bond containing x to y carbon atoms. The term " C_3-C_6 alkenyl" means an alkenyl group containing 3-6 carbon atoms. Representative examples of alkenyl include, but are not limited to, buta-2,3-dienyl, ethenyl, 2-propenyl, 2-methyl-2-propenyl, 3-butenyl, 4-pentenyl, 5-hexenyl, 2-heptenyl, 2-methyl-1-heptenyl, and 3-decenyl.

[0038] The term "alkenylene" means a divalent group derived from a straight or branched chain hydrocarbon of 2 to 4 carbon atoms and contains at least one carbon-carbon double bond. The term " C_x-C_y alkenylene" means a divalent group derived from a straight or branched hydrocarbon chain containing at least one carbon-carbon double bond and containing x to y carbon atoms. Representative examples of alkenylene include, but are not limited to, $-CH=CH-$ and $-CH_2CH=CH-$.

[0039] The term "alkyl" as used herein, means a straight or branched, saturated hydrocarbon chain containing from 1 to 10 carbon atoms. The term " C_x-C_y alkyl" means a straight or branched chain, saturated hydrocarbon containing x to y carbon atoms. For example " C_1-C_6 alkyl" means a straight or branched chain, saturated hydrocarbon containing 2 to 6 carbon atoms. Representative examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, 3-methylhexyl, 2,2-dimethylpentyl, 2,3-dimethylpentyl, n-heptyl, n-octyl, n-nonyl, and n-decyl.

[0040] The term "alkylene" means a divalent group derived from a straight or branched, saturated hydrocarbon chain of 1 to 10 carbon atoms, for example, of 1 to 4 carbon atoms. The term " C_x-C_y alkylene" means a divalent group derived from a straight or branched chain, saturated hydrocarbon containing x to y carbon atoms. For example " C_2-C_6 alkylene" means a straight or branched chain, saturated hydrocarbon containing 2 to 6 carbon atoms. Representative examples of alkylene include, but are not limited to, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-CH_2CH_2CH_2CH_2-$, and $-CH_2CH(CH_3)CH_2-$. The term "alkynyl" as used herein, means a straight or branched chain hydrocarbon group containing from 2 to 10 carbon atoms and containing at least one carbon-carbon triple bond. The term " C_x-C_y alkynyl" means a straight or branched chain hydrocarbon group containing from x to y carbon atoms. For example " C_3-C_6 alkynyl" means a straight or branched chain hydrocarbon group containing from 3 to 6 carbon atoms and containing at least one carbon-carbon triple bond. Representative examples of alkynyl include, but are not limited to, acetylenyl, 1-propynyl, 2-propynyl, 3-butylnyl, 2-pentylnyl, and 1-butylnyl.

[0041] The term "alkynylene," as used herein, means a divalent radical derived from a straight or branched chain hydrocarbon group containing from 2 to 10 carbon atoms and containing at least one carbon-carbon triple bond.

[0042] The term "aryl" as used herein, means phenyl.

[0043] The term "cyclic moiety," as used herein, means benzene, phenyl, phenylene, cycloalkane, cycloalkyl, cycloalkylene, cycloalkene, cycloalkenyl, cycloalkenylene, cycloalkyne, cycloalkynyl, cycloalkynylene, heteroarene, heteroaryl, heterocycloalkane, heterocycloalkyl, heterocycloalkene, heterocycloalkenyl and spiroalkyl.

[0044] The term "cycloalkylene" or cycloalkyl" or "cycloalkane" as used herein, means a monocyclic or bridged hydrocarbon ring system. The monocyclic cycloalkyl is a carbocyclic ring system containing three to ten carbon atoms, zero heteroatoms and zero double bonds. Examples of monocyclic ring systems include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. The monocyclic ring may contain one or two alkylene bridges, each consisting of one, two, or three carbon atoms, each linking two non-adjacent carbon atoms of the ring system. Representative examples of such bridged cycloalkyl ring systems include, but are not limited to, bicyclo[3.1.1]heptane, bicyclo[2.2.1]heptane, bicyclo[2.2.2]octane, bicyclo[3.2.2]nonane, bicyclo[3.3.1]nonane, bicyclo[4.2.1]nonane, tricyclo[3.3.1.0^{3,7}]nonane (octahydro-2,5-methanopentalene or noradamantane), and tricyclo[3.3.1.1^{3,7}]decane (adamantane). The monocyclic and bridged cycloalkyl can be attached to the parent molecular moiety through any substitutable atom contained within the ring system.

[0045] The term "cycloalkenylene," or "cycloalkenyl" or "cycloalkene" as used herein, means a monocyclic or a bridged hydrocarbon ring system. The monocyclic cycloalkenyl has four to ten carbon atoms and zero heteroatoms. The four-membered ring systems have one double bond, the five- or six-membered ring systems have one or two double bonds, the seven- or eight-membered ring systems have one, two, or three double bonds, and the nine- or ten-membered rings have one, two, three, or four double bonds. Representative examples of monocyclic cycloalkenyl groups include, but

are not limited to, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, and cyclooctenyl. The monocyclic cycloalkenyl ring may contain one or two alkylene bridges, each consisting of one, two, or three carbon atoms, each linking two non-adjacent carbon atoms of the ring system. Representative examples of the bridged cycloalkenyl groups include, but are not limited to, 4,5,6,7-tetrahydro-3aH-indene, octahydronaphthalenyl, and 1,6-dihydro-pentalene. The monocyclic and bridged cycloalkenyl can be attached to the parent molecular moiety through any substitutable atom contained within the ring systems.

[0046] The term "cycloalkyne," or "cycloalkynyl," or "cycloalkynylene," as used herein, means a monocyclic or a bridged hydrocarbon ring system. The monocyclic cycloalkynyl has eight or more carbon atoms, zero heteroatoms, and one or more triple bonds. The monocyclic cycloalkynyl ring may contain one or two alkylene bridges, each consisting of one, two, or three carbon atoms, each linking two non-adjacent carbon atoms of the ring system. The monocyclic and bridged cycloalkynyl can be attached to the parent molecular moiety through any substitutable atom contained within the ring systems.

[0047] The term "heteroarene," or "heteroaryl," or "heteroarylene," as used herein, means a five-membered or six-membered aromatic ring having at least one carbon atom and one or more than one independently selected nitrogen, oxygen or sulfur atom. The heteroarenes of this invention are connected through any adjacent atoms in the ring, provided that proper valences are maintained. Representative examples of heteroaryl include, but are not limited to, furanyl (including, but not limited thereto, furan-2-yl), imidazolyl (including, but not limited thereto, 1H-imidazol-1-yl), isoxazolyl, isothiazolyl, oxadiazolyl, 1,3-oxazolyl, pyridinyl (e.g. pyridin-4-yl, pyridin-2-yl, pyridin-3-yl), pyridazinyl, pyrimidinyl, pyrazinyl, pyrazolyl, pyrrolyl, tetrazolyl, thiadiazolyl, 1,3-thiazolyl, thienyl (including, but not limited thereto, thien-2-yl, thien-3-yl), triazolyl, and triazinyl.

[0048] The term "heterocycloalkane," or "heterocycloalkyl," or "heterocycloalkylene," as used herein, means monocyclic or bridged three-, four-, five-, six-, seven-, or eight-membered ring containing at least one heteroatom independently selected from the group consisting of O, N, and S and zero double bonds. The monocyclic and bridged heterocycloalkane are connected to the parent molecular moiety through any substitutable carbon atom or any substitutable nitrogen atom contained within the rings. The nitrogen and sulfur heteroatoms in the heterocycle rings may optionally be oxidized and the nitrogen atoms may optionally be quarternized. Representative examples of heterocycloalkane groups include, but are not limited to, morpholinyl, tetrahydropyranyl, pyrrolidinyl, piperidinyl, dioxolanyl, tetrahydrofuranyl, thiomorpholinyl, 1,4-dioxanyl, tetrahydrothienyl, tetrahydrothiopyranyl, oxetanyl, piperazinyl, imidazolidinyl, azetidyl, azepanyl, aziridinyl, diazepanyl, dithiolanyl, dithianyl, isoxazolidinyl, isothiazolidinyl, oxadiazolidinyl, oxazolidinyl, pyrazolidinyl, tetrahydrothienyl, thiadiazolidinyl, thiazolidinyl, thiomorpholinyl, trithianyl, and trithianyl.

[0049] The term "heterocycloalkene," or "heterocycloalkenyl," or "heterocycloalkenylene," as used herein, means monocyclic or bridged three-, four-, five-, six-, seven-, or eight-membered ring containing at least one heteroatom independently selected from the group consisting of O, N, and S and one or more double bonds. The monocyclic and bridged heterocycloalkene are connected to the parent molecular moiety through any substitutable carbon atom or any substitutable nitrogen atom contained within the rings. The nitrogen and sulfur heteroatoms in the heterocycle rings may optionally be oxidized and the nitrogen atoms may optionally be quarternized. Representative examples of heterocycloalkene groups include, but are not limited to, 1,4,5,6-tetrahydropyridazinyl, 1,2,3,6-tetrahydropyridinyl, dihydropyranyl, imidazolyl, isothiazolyl, oxadiazolyl, isoxazolyl, oxazolyl, pyranyl, pyrazolyl, pyrrolinyl, thiadiazolyl, thiazolyl, and thiopyranyl.

[0050] The term "phenyl," as used herein, means a monovalent radical formed by removal of a hydrogen atom from benzene.

[0051] The term "phenylene," as used herein, means a divalent radical formed by removal of a hydrogen atom from phenyl.

[0052] The term "spiroalkyl," as used herein, means alkylene, both ends of which are attached to the same carbon atom and is exemplified by C₂-spiroalkyl, C₃-spiroalkyl, C₄-spiroalkyl, C₅-spiroalkyl, C₆-spiroalkyl, C₇-spiroalkyl, C₈-spiroalkyl, C₉-spiroalkyl and the like.

[0053] The term "spiroheteroalkyl," as used herein, means spiroalkyl having one or two CH₂ moieties replaced with independently selected O, C(O), CNOH, CNOCH₃, S, S(O), SO₂ or NH and one or two CH moieties unreplaced or replaced with N.

[0054] The term "spiroheteroalkenyl," as used herein, means spiroalkenyl having one or two CH₂ moieties replaced with independently selected O, C(O), CNOH, CNOCH₃, S, S(O), SO₂ or NH and one or two CH moieties unreplaced or replaced with N and also means spiroalkenyl having one or two CH₂ moieties unreplaced or replaced with independently selected O, C(O), CNOH, CNOCH₃, S, S(O), SO₂ or NH and one or two CH moieties replaced with N.

[0055] The term "C₂-C₅-spiroalkyl," as used herein, means C₂-spiroalkyl, C₃-spiroalkyl, C₄-spiroalkyl, and C₅-spiroalkyl.

[0056] The term "C₂-spiroalkyl," as used herein, means eth-1,2-ylene, both ends of which replace hydrogen atoms of the same CH₂ moiety.

[0057] The term "C₃-spiroalkyl," as used herein, means prop-1,3-ylene, both ends of which replace hydrogen atoms

of the same CH₂ moiety.

[0058] The term "C₄-spiroalkyl," as used herein, means but-1,4-ylene, both ends of which replace hydrogen atoms of the same CH₂ moiety.

[0059] The term "C₅-spiroalkyl," as used herein, means pent-1,5-ylene, both ends of which replace hydrogen atoms of the same CH₂ moiety.

[0060] The term "C₆-spiroalkyl," as used herein, means hex-1,6-ylene, both ends of which replace hydrogen atoms of the same CH₂ moiety.

[0061] The term "NH protecting group," as used herein, means trichloroethoxycarbonyl, tribromoethoxycarbonyl, benzyloxycarbonyl, para-nitrobenzylcarbonyl, ortho-bromobenzyloxycarbonyl, chloroacetyl, dichloroacetyl, trichloroacetyl, trifluoroacetyl, phenylacetyl, formyl, acetyl, benzoyl, tert-amylloxycarbonyl, tert-butoxycarbonyl, para-methoxybenzyloxycarbonyl, 3,4-dimethoxybenzyl-oxycarbonyl, 4-(phenylazo)benzyloxycarbonyl, 2-furfuryl-oxycarbonyl, diphenylmethoxycarbonyl, 1,1-dimethylpropoxy-carbonyl, isopropoxycarbonyl, phthaloyl, succinyl, alanyl, leucyl, 1-adamantylloxycarbonyl, 8-quinolyloxycarbonyl, benzyl, diphenylmethyl, triphenylmethyl, 2-nitrophenylthio, methanesulfonyl, para-toluenesulfonyl, N,N-dimethylaminomethylene, benzylidene, 2-hydroxybenzylidene, 2-hydroxy-5-chlorobenzylidene, 2-hydroxy-1-naphthyl-methylene, 3-hydroxy-4-pyridylmethylene, cyclohexylidene, 2-ethoxycarbonylcyclohexylidene, 2-ethoxycarbonylcyclopentylidene, 2-acetylcyclohexylidene, 3,3-dimethyl-5-oxycyclohexylidene, diphenylphosphoryl, dibenzylphosphoryl, 5-methyl-2-oxo-2H-1,3-dioxol-4-yl-methyl, trimethylsilyl, triethylsilyl, and triphenylsilyl.

[0062] The term "C(O)OH protecting group," as used herein, means methyl, ethyl, n-propyl, isopropyl, 1,1-dimethylpropyl, n-butyl, tert-butyl, phenyl, naphthyl, benzyl, diphenylmethyl, triphenylmethyl, para-nitrobenzyl, para-methoxybenzyl, bis(para-methoxyphenyl)methyl, acetylmethyl, benzoylmethyl, para-nitrobenzoylmethyl, para-bromobenzoylmethyl, para-methanesulfonylbenzoylmethyl, 2-tetrahydropyranyl 2-tetrahydrofuran, 2,2,2-trichloroethyl, 2-(trimethylsilyl)ethyl, acetoxymethyl, propionyloxymethyl, pivaloyloxymethyl, phthalimidomethyl, succinimidomethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, methoxymethyl, methoxyethoxymethyl, 2-(trimethylsilyl)ethoxymethyl, benzyloxymethyl, methylthiomethyl, 2-methylthioethyl, phenylthiomethyl, 1,1-dimethyl-2-propenyl, 3-methyl-3-butenyl, allyl, trimethylsilyl, triethylsilyl, triisopropylsilyl, diethylisopropylsilyl, tert-butyl dimethylsilyl, tert-butyl diphenylsilyl, diphenylmethylsilyl, and tert-butylmethoxyphenylsilyl.

[0063] The term "OH or SH protecting group," as used herein, means benzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 3,4-dimethoxybenzyloxycarbonyl, methoxycarbonyl, ethoxycarbonyl, tert-butoxycarbonyl, 1,1-dimethylpropoxycarbonyl, isopropoxycarbonyl, isobutyloxycarbonyl, diphenylmethoxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2,2,2-tribromoethoxycarbonyl, 2-(trimethylsilyl)ethoxycarbonyl, 2-(phenylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphonio)ethoxycarbonyl, 2-furfuryloxycarbonyl, 1-adamantylloxycarbonyl, vinylloxycarbonyl, allyloxycarbonyl, S-benzylthiocarbonyl, 4-ethoxy-1-naphthylloxycarbonyl, 8-quinolyloxycarbonyl, acetyl, formyl, chloroacetyl, dichloroacetyl, trichloroacetyl, trifluoroacetyl, methoxyacetyl, phenoxyacetyl, pivaloyl, benzoyl, methyl, tert-butyl, 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 1,1-dimethyl-2-propenyl, 3-methyl-3-butenyl, allyl, benzyl (phenylmethyl), para-methoxybenzyl, 3,4-dimethoxybenzyl, diphenylmethyl, triphenylmethyl, tetrahydrofuryl, tetrahydropyranyl, tetrahydrothiopyranyl, methoxymethyl, methylthiomethyl, benzyloxymethyl, 2-methoxyethoxymethyl, 2,2,2-trichloroethoxymethyl, 2-(trimethylsilyl)ethoxymethyl, 1-ethoxyethyl, methanesulfonyl, para-toluenesulfonyl, trimethylsilyl, triethylsilyl, triisopropylsilyl, diethylisopropylsilyl, tert-butyl dimethylsilyl, tert-butyl diphenylsilyl, diphenylmethylsilyl, and tert-butylmethoxyphenylsilyl.

Compounds

[0064] Geometric isomers may exist in the present compounds. Compounds of this invention may contain carbon-carbon double bonds or carbon-nitrogen double bonds in the E or Z configuration, wherein the term "E" represents higher order substituents on opposite sides of the carbon-carbon or carbon-nitrogen double bond and the term "Z" represents higher order substituents on the same side of the carbon-carbon or carbon-nitrogen double bond as determined by the Cahn-Ingold-Prelog Priority Rules. The compounds of this invention may also exist as a mixture of "E" and "Z" isomers. Substituents around a cycloalkyl or heterocycloalkyl are designated as being of cis or trans configuration. Furthermore, the invention contemplates the various isomers and mixtures thereof resulting from the disposal of substituents around an adamantane ring system. Two substituents around a single ring within an adamantane ring system are designated as being of Z or E relative configuration. For examples, see C. D. Jones, M. Kaselj, R. N. Salvatore, W. J. le Noble J. Org. Chem. 1998, 63, 2758-2760.

[0065] Compounds of this invention may contain asymmetrically substituted carbon atoms in the R or S configuration, in which the terms "R" and "S" are as defined by the IUPAC 1974 Recommendations for Section E, Fundamental Stereochemistry, Pure Appl. Chem. (1976) 45, 13-10. Compounds having asymmetrically substituted carbon atoms with equal amounts of R and S configurations are racemic at those carbon atoms. Atoms with an excess of one configuration over the other are assigned the configuration present in the higher amount, preferably an excess of about 85%-90%, more preferably an excess of about 95%-99%, and still more preferably an excess greater than about 99%. Accordingly,

this invention includes racemic mixtures, relative and absolute stereoisomers, and mixtures of relative and absolute stereoisomers.

Isotope Enriched or Labeled Compounds

[0066] Compounds of the invention can exist in isotope-labeled or -enriched form containing one or more atoms having an atomic mass or mass number different from the atomic mass or mass number most abundantly found in nature. Isotopes can be radioactive or nonradioactive isotopes. Isotopes of atoms such as hydrogen, carbon, phosphorous, sulfur, fluorine, chlorine, and iodine include, but are not limited to, ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{32}P , ^{35}S , ^{18}F , ^{36}Cl , and ^{125}I . Compounds that contain other isotopes of these and/or other atoms are within the scope of this invention.

[0067] In another embodiment, the isotope-labeled compounds contain deuterium (^2H), tritium (^3H) or ^{14}C isotopes. Isotope-labeled compounds of this invention can be prepared by the general methods well known to persons having ordinary skill in the art. Such isotope-labeled compounds can be conveniently prepared by carrying out the procedures disclosed in the Examples disclosed herein and Schemes by substituting a readily available isotope-labeled reagent for a non-labeled reagent. In some instances, compounds may be treated with isotope-labeled reagents to exchange a normal atom with its isotope, for example, hydrogen for deuterium can be exchanged by the action of a deuterio acid such as $\text{D}_2\text{SO}_4/\text{D}_2\text{O}$. In addition to the above, relevant procedures and intermediates are disclosed, for instance, in Lizondo, J et al., *Drugs Fut*, 21(11), 1116 (1996); Brickner, S J et al., *J Med Chem*, 39(3), 673 (1996); Mallesham, B et al., *Org Lett*, 5(7), 963 (2003); PCT publications WO1997010223, WO2005099353, WO1995007271, WO2006008754; US Patent Nos. 7538189; 7534814; 7531685; 7528131; 7521421; 7514068; 7511013; and US Patent Application Publication Nos. 20090137457; 20090131485; 20090131363; 20090118238; 20090111840; 20090105338; 20090105307; 20090105147; 20090093422; 20090088416; and 20090082471, the methods are hereby incorporated by reference.


[0068] The isotope-labeled compounds of the invention may be used as standards to determine the effectiveness of Bcl-2 inhibitors in binding assays. Isotope containing compounds have been used in pharmaceutical research to investigate the *in vivo* metabolic fate of the compounds by evaluation of the mechanism of action and metabolic pathway of the nonisotope-labeled parent compound (Blake et al. *J. Pharm. Sci.* 64, 3, 367-391 (1975)). Such metabolic studies are important in the design of safe, effective therapeutic drugs, either because the *in vivo* active compound administered to the patient or because the metabolites produced from the parent compound prove to be toxic or carcinogenic (Foster et al., *Advances in Drug Research* Vol. 14, pp. 2-36, Academic press, London, 1985; Kato et al., *J. Labelled Comp. Radiopharmaceut.*, 36(10):927-932 (1995); Kushner et al., *Can. J. Physiol. Pharmacol.*, 77, 79-88 (1999).

[0069] In addition, non-radio active isotope containing drugs, such as deuterated drugs called "heavy drugs," can be used for the treatment of diseases and conditions related to Bcl-2 activity. Increasing the amount of an isotope present in a compound above its natural abundance is called enrichment. Examples of the amount of enrichment include from about 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 16, 21, 25, 29, 33, 37, 42, 46, 50, 54, 58, 63, 67, 71, 75, 79, 84, 88, 92, 96, to about 100 mol %. Replacement of up to about 15% of normal atom with a heavy isotope has been effected and maintained for a period of days to weeks in mammals, including rodents and dogs, with minimal observed adverse effects (Czajka D M and Finkel A J, *Ann. N.Y. Acad. Sci.* 1960 84: 770; Thomson J F, *Ann. New York Acad. Sci.* 1960 84: 736; Czajka D M et al., *Am. J. Physiol.* 1961 201: 357). Acute replacement of as high as 15%-23% in human fluids with deuterium was found not to cause toxicity (Blagojevic N et al. in "Dosimetry & Treatment Planning for Neutron Capture Therapy", Zamenhof R, Solares G and Harling O Eds. 1994. Advanced Medical Publishing, Madison Wis. pp.125-134; *Diabetes Metab.* 23: 251 (1997)).

[0070] Stable isotope labeling of a drug can alter its physico-chemical properties such as pKa and lipid solubility. These effects and alterations can affect the pharmacodynamic response of the drug molecule if the isotopic substitution affects a region involved in a ligand-receptor interaction. While some of the physical properties of a stable isotope-labeled molecule are different from those of the unlabeled one, the chemical and biological properties are the same, with one important exception: because of the increased mass of the heavy isotope, any bond involving the heavy isotope and another atom will be stronger than the same bond between the light isotope and that atom. Accordingly, the incorporation of an isotope at a site of metabolism or enzymatic transformation will slow said reactions potentially altering the pharmacokinetic profile or efficacy relative to the non-isotopic compound.

[0071] Suitable groups for A^1 , B^1 , D^1 , E^1 , G^1 , Y^1 , L^1 , Z^1A , Z^2A , Z^1 , Z^2 , and Z^3 in compounds of Formula (I) are independently selected. The described embodiments of the present invention may be combined. Such combination is contemplated and within the scope of the present invention. For example, it is contemplated that embodiments for any of A^1 , B^1 , D^1 , E^1 , G^1 , Y^1 , L^1 , Z^1A , Z^2A , Z^1 , Z^2 , and Z^3 can be combined with embodiments defined for any other of A^1 , B^1 , D^1 , E^1 , G^1 , Y^1 , L^1 , Z^1A , Z^2A , Z^1 , Z^2 , and Z^3 .

[0072] One embodiment of this invention, therefore, pertains to compounds or therapeutically acceptable salts, which are useful as inhibitors of anti-apoptotic Bcl-2 proteins, the compounds having Formula (I)



(I),

wherein

A¹ is N or C(A²);

A² is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};

B¹ is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};

D¹ is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSON(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};

E¹ is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; and

Y¹ is H, CN, NO₂, C(O)OH, F, Cl, Br, I, CF₃, OCF₃, CF₂CF₃, OCF₂CF₃, R¹⁷, OR¹⁷, C(O)R¹⁷, C(O)OR¹⁷, SR¹⁷, SO₂R¹⁷, NH₂, NHR¹⁷, N(R¹⁷)₂, NHC(O)R¹⁷, C(O)NH₂, C(O)NHR¹⁷, C(O)N(R¹⁷)₂, NHS(O)R¹⁷ or NHSO₂R¹⁷; or E¹ and Y¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

A², B¹, and D¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; or

Y¹ and B¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

A², D¹, and E¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; or

A² and B¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

D¹, E¹, and Y¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹,

$\text{NHC(O)N(R}^1\text{)}_2$, $\text{NR}^1\text{C(O)NHR}^1$, $\text{NR}^1\text{C(O)N(R}^1\text{)}_2$, SO_2NH_2 , SO_2NHR^1 , $\text{SO}_2\text{N(R}^1\text{)}_2$, NHSO_2R^1 , $\text{NR}^1\text{SO}_2\text{R}^1$, $\text{NHSO}_2\text{NHR}^1$, $\text{NHSO}_2\text{N(R}^1\text{)}_2$, $\text{NR}^1\text{SO}_2\text{NHR}^1$, $\text{NR}^1\text{SO}_2\text{N(R}^1\text{)}_2$, C(O)NHNOH , C(O)NHNOR^1 , $\text{C(O)NHSO}_2\text{R}^1$, C(NH)NH_2 , C(NH)NHR^1 , $\text{C(NH)N(R}^1\text{)}_2$, $\text{NHSO}_2\text{NHR}^1$, $\text{NHSO}_2\text{N(CH}_3\text{)R}^1$, $\text{N(CH}_3\text{)SO}_2\text{N(CH}_3\text{)R}^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, C(O)H , CHNOH , $\text{CH(NOCH}_3\text{)}$, CF_3 , C(O)OH , C(O)NH_2 or C(O)OR^{1A} ; or
 5 A^2 and D^1 , together with the atoms to which they are attached, are benzene, naphthalene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and
 B^1 , E^1 , and Y^1 are independently selected H, R^1 , OR^1 , SR^1 , S(O)R^1 , SO_2R^1 , C(O)R^1 , C(O)OR^1 , OC(O)R^1 , NHR^1 , $\text{N(R}^1\text{)}_2$, C(O)NHR^1 , $\text{C(O)N(R}^1\text{)}_2$, NHC(O)R^1 , $\text{NR}^1\text{C(O)R}^1$, NHC(O)OR^1 , $\text{NR}^1\text{C(O)OR}^1$, NHC(O)NH_2 , NHC(O)NHR^1 , $\text{NHC(O)N(R}^1\text{)}_2$, $\text{NR}^1\text{C(O)NHR}^1$, $\text{NR}^1\text{C(O)N(R}^1\text{)}_2$, SO_2NH_2 , SO_2NHR^1 , $\text{SO}_2\text{N(R}^1\text{)}_2$, NHSO_2R^1 , $\text{NR}^1\text{SO}_2\text{R}^1$, $\text{NHSO}_2\text{NHR}^1$, $\text{NHSO}_2\text{N(R}^1\text{)}_2$, $\text{NR}^1\text{SO}_2\text{NHR}^1$, $\text{NR}^1\text{SO}_2\text{N(R}^1\text{)}_2$, C(O)NHNOH , C(O)NHNOR^1 , $\text{C(O)NHSO}_2\text{R}^1$, C(NH)NH_2 , C(NH)NHR^1 , $\text{C(NH)N(R}^1\text{)}_2$, $\text{NHSO}_2\text{NHR}^1$, $\text{NHSO}_2\text{N(CH}_3\text{)R}^1$, $\text{N(CH}_3\text{)SO}_2\text{N(CH}_3\text{)R}^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, C(O)H , CHNOH , $\text{CH(NOCH}_3\text{)}$, CF_3 , C(O)OH , C(O)NH_2 or C(O)OR^{1A} ;
 10 G^1 is H, or C(O)OR ;
 R is alkyl;
 15 R^1 is R^2 , R^3 , R^4 or R^5 ;
 R^{1A} is cycloalkyl, cycloalkenyl or cycloalkynyl;
 R^2 is phenyl, which is unfused or fused with R^{2A} ; R^{2A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^3 is heteroaryl, which is unfused or fused with R^{3A} ; R^{3A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 20 R^4 is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{4A} ; R^{4A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^5 is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R^6 , $\text{NC(R}^{6A}\text{)(R}^{6B}\text{)}$, R^7 , OR^7 , SR^7 , S(O)R^7 , SO_2R^7 , NHR^7 , $\text{N(R}^7\text{)}_2$, C(O)R^7 , C(O)NH_2 , C(O)NHR^7 , $\text{C(O)N(R}^7\text{)}_2$, NHC(O)R^7 , $\text{NR}^7\text{C(O)R}^7$, NHSO_2R^7 , NHC(O)OR^7 , SO_2NH_2 , SO_2NHR^7 , $\text{SO}_2\text{N(R}^7\text{)}_2$, NHC(O)NH_2 , NHC(O)NHR^7 , $\text{NHC(O)CH(CH}_3\text{)NHC(O)CH(CH}_3\text{)NH}_2$, $\text{NHC(O)CH(CH}_3\text{)NHC(O)CH(CH}_3\text{)NHR}^7$, OH, (O), C(O)OH , N_3 , CN, NH_2 , CF_3 , CF_2CF_3 , F, Cl, Br or I;
 R^6 is $\text{C}_2\text{-C}_5\text{-spiroalkyl}$, each of which is unsubstituted or substituted with OH, (O), N_3 , CN, CF_3 , CF_2CF_3 , F, Cl, Br, I, NH_2 , $\text{NH(CH}_3\text{)}$ or $\text{N(CH}_3\text{)}_2$;
 30 R^{6A} and R^{6B} are independently selected alkyl or, together with the N to which they are attached, R^{6C} ;
 R^{6C} is aziridin-1-yl, azetidin-1-yl, pyrrolidin-1-yl or piperidin-1-yl, each having one CH_2 moiety unreplaced or replaced with O, C(O) , CNOH , CNOCH_3 , S, S(O) , SO_2 or NH;
 R^7 is R^8 , R^9 , R^{10} or R^{11} ;
 R^8 is phenyl, which is unfused or fused with R^{8A} ; R^{8A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 35 R^9 is heteroaryl, which is unfused or fused with R^{9A} ; R^{9A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^{10} is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{10A} ; R^{10A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 40 R^{11} is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R^{12} , OR^{12} , SR^{12} , S(O)R^{12} , SO_2R^{12} , C(O)R^{12} , CO(O)R^{12} , OC(O)R^{12} , OC(O)OR^{12} , NH_2 , NHR^{12} , $\text{N(R}^{12}\text{)}_2$, NHC(O)R^{12} , $\text{NR}^{12}\text{C(O)R}^{12}$, $\text{NHS(O)}_2\text{R}^{12}$, $\text{NR}^{12}\text{S(O)}_2\text{R}^{12}$, NHC(O)OR^{12} , $\text{NR}^{12}\text{C(O)OR}^{12}$, NHC(O)NH_2 , NHC(O)NHR^{12} , $\text{NHC(O)N(R}^{12}\text{)}_2$, $\text{NR}^{12}\text{C(O)NHR}^{12}$, $\text{NR}^{12}\text{C(O)N(R}^{12}\text{)}_2$, C(O)NH_2 , C(O)NHR^{12} , $\text{C(O)N(R}^{12}\text{)}_2$, C(O)NHOH , C(O)NHOR^{12} , $\text{C(O)NHSO}_2\text{R}^{12}$, $\text{C(O)NR}^{12}\text{SOR}^{12}$, SO_2NH_2 , $\text{SO}_2\text{NHR}^{12}$, $\text{SO}_2\text{N(R}^{12}\text{)}_2$, C(O)H , C(O)OH , C(N)NH_2 , C(N)NHR^{12} , $\text{C(N)N(R}^{12}\text{)}_2$, CNOH , CNOCH_3 , OH, (O), CN, N_3 , NO_2 , CF_3 , CF_2CF_3 , OCF_3 , OCF_2CF_3 , F, Cl, Br or I;
 R^{12} is R^{13} , R^{14} , R^{15} or R^{16} ;
 R^{13} is phenyl, which is unfused or fused with R^{13A} ; R^{13A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 50 R^{14} is heteroaryl, which is unfused or fused with R^{14A} ; R^{14A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^{15} is cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene, each of which is unfused or fused with R^{15A} ; R^{15A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^{16} is alkyl, alkenyl or alkynyl;
 55 R^{17} is R^{18} , R^{19} , R^{20} or R^{21} ;
 R^{18} is phenyl, which is unfused or fused with R^{18A} ; R^{18A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^{19} is heteroaryl, which is unfused or fused with R^{19A} ; R^{19A} is benzene, heteroarene, cycloalkane, cycloalkene,

heterocycloalkane or heterocycloalkene;

R²⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl each of which is unfused or fused with R^{20A}; R^{20A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R²², OR²², SR²², S(O)R²², SO₂R²², C(O)R²², CO(O)R²², OC(O)R²², OC(O)OR²², NH₂, NHR²², N(R²²)₂, NHC(O)R²², NR²²C(O)R²², NHS(O)₂R²², NR²²S(O)₂R²², NHC(O)OR²², NR²²C(O)OR²², NHC(O)NH₂, NHC(O)NHR²², NHC(O)N(R²²)₂, NR²²C(O)NHR²², NR²²C(O)N(R²²)₂, C(O)NH₂, C(O)NHR²², C(O)N(R²²)₂, C(O)NHOH, C(O)NHOR²², C(O)NHSO₂R²², C(O)NR²²SO₂R²², SO₂NH₂, SO₂NHR²², SO₂N(R²²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR²², C(N)N(R²²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R²² is R²³, R²⁴ or R²⁵;

R²³ is phenyl, which is unfused or fused with R^{23A}; R^{23A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁴ is heteroarene, which is unfused or fused with R^{24A}; R^{24A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{25A}; R^{25A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

Z¹ is R²⁶ or R²⁷;

Z² is R²⁸, R²⁹ or R³⁰;

Z^{1A} and Z^{2A} are both absent or are taken together to form CH₂, CH₂CH₂ or Z^{12A};

Z^{12A} is C₂-C₆-alkylene having one or two CH₂ moieties replaced by NH, N(CH₃), S, S(O) or SO₂;

L¹ is a R³⁷, OR³⁷, SR³⁷, S(O)R³⁷, SO₂R³⁷, C(O)R³⁷, CO(O)R³⁷, OC(O)R³⁷, OC(O)OR³⁷, NHR³⁷, C(O)NH, C(O)NR³⁷, C(O)NHOR³⁷, C(O)NHSO₂R³⁷, SO₂NH, SO₂NHR³⁷, C(N)NH, C(N)NHR³⁷;

R²⁶ is phenylene, which is unfused or fused with R^{26A}; R^{26A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁷ is heteroarylene, which is unfused or fused with R^{27A}; R^{27A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁸ is phenylene, which is unfused or fused with R^{28A}; R^{28A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁹ is heteroarylene, which is unfused or fused with R^{29A}; R^{29A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁰ is cycloalkylene, cycloalkenylene, heterocycloalkylene or heterocycloalkenylene, each of which is unfused or fused with R^{30A}; R^{30A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R³⁷ is a bond or R^{37A};

R^{37A} is alkylene, alkenylene, or alkynylene, each of which is unsubstituted or substituted with one or two or three independently selected R^{37B}, OR^{37B}, SR^{37B}, S(O)R^{37B}, SO₂R^{37B}, C(O)R^{37B}, CO(O)R^{37B}, OC(O)R^{37B}, OC(O)OR^{37B}, NH₂, NHR^{37B}, N(R^{37B})₂, NHC(O)R^{37B}, NR^{37B}C(O)R^{37B}, NHS(O)₂R^{37B}, NR^{37B}S(O)₂R^{37B}, NHC(O)OR^{37B}, NR^{37B}C(O)OR^{37B}, NHC(O)NH₂, NHC(O)NHR^{37B}, NHC(O)N(R^{37B})₂, NR^{37B}C(O)NHR^{37B}, NR^{37B}C(O)N(R^{37B})₂, C(O)NH₂, C(O)NHR^{37B}, C(O)N(R^{37B})₂, C(O)NHOH, C(O)NHOR^{37B}, C(O)NHSO₂R^{37B}, C(O)NR^{37B}SO₂R^{37B}, SO₂NH₂, SO₂NHR^{37B}, SO₂N(R^{37B})₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR^{37B}, C(N)N(R^{37B})₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I substituents;

R^{37B} is alkyl, alkenyl, alkynyl, or R^{37C};

R^{37C} is phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl;

Z³ is R³⁸, R³⁹ or R⁴⁰;

R³⁸ is phenyl, which is unfused or fused with R^{38A}; R^{38A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁹ is heteroaryl, which is unfused or fused with R^{39A}; R^{39A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁴⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{40A}; R^{40A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

wherein the moieties represented by R²⁶ and R²⁷ are substituted (i.e., if Z^{1A} and Z^{2A} are absent) or further substituted (i.e., if Z^{1A} and Z^{2A} are present) with R⁴¹, OR⁴¹, SR⁴¹, S(O)R⁴¹, SO₂R⁴¹, C(O)R⁴¹, CO(O)R⁴¹, OC(O)R⁴¹, OC(O)OR⁴¹, NHR⁴¹, N(R⁴¹)₂, NHC(O)R⁴¹, NR⁴¹C(O)R⁴¹, NHS(O)₂R⁴¹, NR⁴¹S(O)₂R⁴¹, NHC(O)OR⁴¹, NR⁴¹C(O)OR⁴¹, NHC(O)NHR⁴¹, NHC(O)N(R⁴¹)₂, NR⁴¹C(O)NHR⁴¹, NR⁴¹C(O)N(R⁴¹)₂, C(O)NHR⁴¹, C(O)N(R⁴¹)₂, C(O)NHOR⁴¹, C(O)NHSO₂R⁴¹, C(O)NR⁴¹SO₂R⁴¹, SO₂NHR⁴¹, SO₂N(R⁴¹)₂, C(N)NHR⁴¹, or C(N)N(R⁴¹)₂;

R⁴¹ is heteroaryl, which is fused with R^{43A}; R^{43A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; which is unfused or fused with benzene, heteroarene or R^{43B}; R^{43B} is cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

wherein the cyclic moieties represented by E¹ and Y¹ together, Y¹ and B¹ together, A² and B¹ together, A² and D¹ together, R^{1A}, R², R^{2A}, R³, R^{3A}, R⁴, R^{4A}, R⁶, R^{6C}, R⁸, R^{8A}, R⁹, R^{9A}, R¹⁰, R^{10A}, R¹³, R^{13A}, R¹⁴, R^{14A}, R¹⁵, R^{15A}, R¹⁸, R^{18A}, R¹⁹, R^{19A}, R²⁰, R^{20A}, R²³, R^{23A}, R²⁴, R^{24A}, R²⁵, R^{25A}, R²⁶, R^{26A}, R²⁷, R^{27A}, R²⁸, R^{28A}, R²⁹, R^{29A}, R³⁰, R^{30A}, R^{37B}, R³⁸, R^{38A}, R³⁹, R^{39A}, R⁴⁰, and R^{40A} are independently unsubstituted, further unsubstituted, substituted or further substituted with one or two or three or four or five of independently selected R^{57A}, R⁵⁷, OR⁵⁷, SR⁵⁷, S(O)R⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, OC(O)R⁵⁷, OC(O)OR⁵⁷, NH₂, NHR⁵⁷, N(R⁵⁷)₂, NHC(O)R⁵⁷, NR⁵⁷C(O)R⁵⁷, NHS(O)₂R⁵⁷, NR⁵⁷S(O)₂R⁵⁷, NHC(O)OR⁵⁷, NR⁵⁷C(O)OR⁵⁷, NHC(O)NH₂, NHC(O)NHR⁵⁷, NHC(O)N(R⁵⁷)₂, NR⁵⁷C(O)NHR⁵⁷, NR⁵⁷C(O)N(R⁵⁷)₂, C(O)NH₂, C(O)NHR⁵⁷, C(O)N(R⁵⁷)₂, C(O)NHOH, C(O)NHOR⁵⁷, C(O)NHSO₂R⁵⁷, C(O)NR⁵⁷SO₂R⁵⁷, SO₂NH₂, SO₂NHR⁵⁷, SO₂N(R⁵⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁵⁷, C(N)N(R⁵⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I; R^{57A} is spiroalkyl, or spiroheteroalkyl; R⁵⁷ is R⁵⁸, R⁵⁹, R⁶⁰ or R⁶¹; R⁵⁸ is phenyl, which is unfused or fused with R^{58A}; R^{58A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁵⁹ is heteroaryl, which is unfused or fused with R^{59A}; R^{59A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁶⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{60A}; R^{60A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁶¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶², OR⁶², SR⁶², S(O)R⁶², SO₂R⁶², C(O)R⁶², CO(O)R⁶², OC(O)R⁶², OC(O)OR⁶², NH₂, NHR⁶², N(R⁶²)₂, NHC(O)R⁶², NR⁶²C(O)R⁶², NHS(O)R⁶², NR⁶²S(O)₂R⁶², NHC(O)OR⁶², NR⁶²BC(O)OR⁶², NHC(O)NH₂, NHC(O)NHR⁶², NHC(O)N(R⁶²)₂, NR⁶²C(O)NHR⁶², NR⁶²C(O)N(R⁶²)₂, C(O)NH₂, C(O)NHR⁶², C(O)N(R⁶²)₂, C(O)NHOH, C(O)NHOR⁶², C(O)NHSOR⁶², C(O)NR⁶²SO₂R⁶², SO₂NH₂, SO₂NHR⁶², SO₂N(R⁶²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶², C(N)N(R⁶²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I; R⁶² is R⁶³, R⁶⁴, R⁶⁵ or R⁶⁶; R⁶³ is phenyl, which is unfused or fused with R^{63A}; R^{63A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁶⁴ is heteroaryl, which is unfused or fused with R^{64A}; R^{64A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁶⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl, each of which is unfused or fused with R^{65A}; R^{65A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁶⁶ is alkyl, alkenyl or alkenyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶⁷, OR⁶⁷, SR⁶⁷, S(O)R⁶⁷, SO₂R⁶⁷, C(O)R⁶⁷, CO(O)R⁶⁷, OC(O)R⁶⁷, OC(O)OR⁶⁷, NH₂, NHR⁶⁷, N(R⁶⁷)₂, NHC(O)R⁶⁷, NR⁶⁷C(O)R⁶⁷, NHS(O)₂R⁶⁷, NR⁶⁷S(O)₂R⁶⁷, NHC(O)OR⁶⁷, NR⁶⁷C(O)OR⁶⁷, NHC(O)NH₂, NHC(O)NHR⁶⁷, NHC(O)N(R⁶⁷)₂, NR⁶⁷C(O)NHR⁶⁷, NR⁶⁷C(O)N(R⁶⁷)₂, C(O)NH₂, C(O)NHR⁶⁷, C(O)N(R⁶⁷)₂, C(O)NHOH, C(O)NHOR⁶⁷, C(O)NHSO₂R⁶⁷, C(O)NR⁶⁷SO₂R⁶⁷, SO₂NH₂, SO₂NHR⁶⁷, SO₂N(R⁶⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁷, C(N)N(R⁶⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I substituents; R⁶⁷ is alkyl, alkenyl, alkynyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; wherein the cyclic moieties represented by R^{57A}, R⁵⁸, R⁵⁹, R⁶⁰, R⁶³, R⁶⁴, R⁶⁵, and R⁶⁷ are unsubstituted or substituted with one or two or three or four of independently selected R⁶⁸, OR⁶⁸, SR⁶⁸, S(O)R⁶⁸, SO₂R⁶⁸, C(O)R⁶⁸, CO(O)R⁶⁸, OC(O)R⁶⁸, OC(O)OR⁶⁸, NH₂, NHR⁶⁸, N(R⁶⁸)₂, NHC(O)R⁶⁸, NR⁶⁸C(O)R⁶⁸, NHS(O)₂R⁶⁸, NR⁶⁸S(O)₂R⁶⁸, NHC(O)OR⁶⁸, NR⁶⁸C(O)OR⁶⁸, NHC(O)NH₂, NHC(O)NHR⁶⁸, NHC(O)N(R⁶⁸)₂, NR⁶⁸C(O)NHR⁶⁸, NR⁶⁸C(O)N(R⁶⁸)₂, C(O)NH₂, C(O)NHR⁶⁸, C(O)N(R⁶⁸)₂, C(O)NHOH, C(O)NHOR⁶⁸, C(O)NHSO₂R⁶⁸, C(O)NR⁶⁸SO₂R⁶⁸, SO₂NH₂, SONHR⁶⁸, SO₂N(R⁶⁸)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁸, C(N)N(R⁶⁸)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I; R⁶⁸ is R⁶⁹, R⁷⁰, R⁷¹ or R⁷²; R⁶⁹ is phenyl, which is unfused or fused with R^{69A}; R^{69A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁷⁰ is heteroaryl, which is unfused or fused with R^{70A}; R^{70A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁷¹ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{71A}; R^{71A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁷² is alkyl, alkenyl or alkenyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁷³, OR⁷³, SR⁷³, S(O)R⁷³, SO₂R⁷³, C(O)R⁷³, CO(O)R⁷³, OC(O)R⁷³, OC(O)OR⁷³, NH₂, NHR⁷³, N(R⁷³)₂, NHC(O)R⁷³, NR⁷³C(O)R⁷³, NHS(O)₂R⁷³, NR⁷³S(O)₂R⁷³, NHC(O)OR⁷³, NR⁷³C(O)OR⁷³, NHC(O)NH₂, NHC(O)NHR⁷³, NHC(O)N(R⁷³)₂, NR⁷³C(O)NHR⁷³, NR⁷³C(O)N(R⁷³)₂, C(O)NH₂, C(O)NHR⁷³, C(O)N(R⁷³)₂,

C(O)NHOH, C(O)NHOR⁷³, C(O)NHSO₂R⁷³, C(O)NR⁷³SO₂R⁷³, SO₂NH₂, SO₂NHR⁷³, SO₂N(R⁷³)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁷³, C(N)N(R⁷³)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R⁷³ is alkyl, alkenyl, alkenyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; and
 5 wherein the moieties represented by R⁶⁹, R⁷⁰, and R⁷¹ are unsubstituted or substituted with one or two or three or four of independently selected NH₂, C(O)NH₂, C(O)NHOH, SO₂NH₂, CF₃, CF₂CF₃, C(O)H, C(O)OH, C(N)NH₂, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I.

[0073] Another embodiment of this invention pertains to compounds of Formula (I), wherein

A¹ is N or C(A²);

A² is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};

B¹ is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};

D¹ is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};

E¹ is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; and

Y¹ is H, CN, NO₂, C(O)OH, F, Cl, Br, I, CF₃, OCF₃, CF₂CF₃, OCF₂CF₃, R¹⁷, OR¹⁷, C(O)R¹⁷, C(O)OR¹⁷, SR¹⁷, SO₂R¹⁷, NH₂, NHR¹⁷, N(R¹⁷)₂, NHC(O)R¹⁷, C(O)NH₂, C(O)NHR¹⁷, C(O)N(R¹⁷)₂, NHS(O)R¹⁷ or NHSO₂R¹⁷; or
 E¹ and Y¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

A², B¹, and D¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; or

Y¹ and B¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

A², D¹, and E¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; or

A² and B¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

D¹, E¹, and Y¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹,

$\text{NHSO}_2\text{NHR}^1$, $\text{NHSO}_2\text{N(R}^1)_2$, $\text{NR}^1\text{SO}_2\text{NHR}^1$, $\text{NR}^1\text{SO}_2\text{N(R}^1)_2$, C(O)NHNOH , C(O)NHNOR^1 , $\text{C(O)NHSO}_2\text{R}^1$,
 C(NH)NH_2 , C(NH)NHR^1 , $\text{C(NH)N(R}^1)_2$, $\text{NHSO}_2\text{NHR}^1$, $\text{NHSO}_2\text{N(CH}_3\text{)R}^1$, $\text{N(CH}_3\text{)SO}_2\text{N(CH}_3\text{)R}^1$, F, Cl, Br, I, CN,
 NO_2 , N_3 , OH, C(O)H , CHNOH , $\text{CH(NOCH}_3\text{)}$, CF_3 , C(O)OH , C(O)NH_2 or C(O)OR^{1A} ; or
 A^2 and D^1 , together with the atoms to which they are attached, are benzene, naphthalene, heteroarene, cycloalkane,
5 cycloalkene, heterocycloalkane or heterocycloalkene; and
 B^1 , E^1 , and Y^1 are independently selected H, R^1 , OR^1 , SR^1 , S(O)R^1 , SO_2R^1 , C(O)R^1 , C(O)OR^1 , OC(O)R^1 , NHR^1 ,
 $\text{N(R}^1)_2$, C(O)NHR^1 , $\text{C(O)N(R}^1)_2$, NHC(O)R^1 , $\text{NR}^1\text{C(O)R}^1$, NHC(O)OR^1 , $\text{NR}^1\text{C(O)OR}^1$, NHC(O)NH_2 , NHC(O)NHR^1 ,
 $\text{NHC(O)N(R}^1)_2$, $\text{NR}^1\text{C(O)NHR}^1$, $\text{NR}^1\text{C(O)N(R}^1)_2$, SO_2NH_2 , SO_2NHR^1 , $\text{SON(R}^1)_2$, NHSO_2R^1 , $\text{NR}^1\text{SO}_2\text{R}^1$,
 $\text{NHSO}_2\text{NHR}^1$, $\text{NHSO}_2\text{N(R}^1)_2$, $\text{NR}^1\text{SO}_2\text{NHR}^1$, $\text{NR}^1\text{SO}_2\text{N(R}^1)_2$, C(O)NHNOH , C(O)NHNOR^1 , $\text{C(O)NHSO}_2\text{R}^1$,
10 C(NH)NH_2 , C(NH)NHR^1 , $\text{C(NH)N(R}^1)_2$, $\text{NHSO}_2\text{NHR}^1$, $\text{NHSO}_2\text{N(CH}_3\text{)R}^1$, $\text{N(CH}_3\text{)SO}_2\text{N(CH}_3\text{)R}^1$, F, Cl, Br, I, CN,
 NO_2 , N_3 , OH, C(O)H , CHNOH , $\text{CH(NOCH}_3\text{)}$, CF_3 , C(O)OH , C(O)NH_2 or C(O)OR^{1A} ;
 G^1 is H, or C(O)OR ;
R is alkyl;
 R^1 is R^2 , R^3 , R^4 or R^5 ;
15 R^{1A} is cycloalkyl, cycloalkenyl or cycloalkynyl;
 R^2 is phenyl, which is unfused or fused with R^{2A} ; R^{2A} is benzene, heteroarene, cycloalkane, cycloalkene, hetero-
cycloalkane or heterocycloalkene;
 R^3 is heteroaryl, which is unfused or fused with R^{3A} ; R^{3A} is benzene, heteroarene, cycloalkane, cycloalkene, hete-
rocycloalkane or heterocycloalkene;
20 R^4 is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{4A} ;
 R^{4A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^5 is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently
selected R^6 , $\text{NC(R}^{6A}\text{)(R}^{6B}\text{)}$, R^7 , OR^7 , SR^7 , S(O)R^7 , SO_2R^7 , NHR^7 , $\text{N(R}^7)_2$, C(O)R^7 , C(O)NH_2 , C(O)NHR^7 ,
 $\text{C(O)N(R}^7)_2$, NHC(O)R^7 , $\text{NR}^7\text{C(O)R}^7$, NHSO_2R^7 , NHC(O)OR^7 , SO_2NH_2 , SO_2NHR^7 , $\text{SO}_2\text{N(R}^7)_2$, NHC(O)NH_2 ,
25 NHC(O)NHR^7 , $\text{NHC(O)CH(CH}_3\text{)NHC(O)CH(CH}_3\text{)NH}_2$, $\text{NHC(O)CH(CH}_3\text{)NHC(O)CH(CH}_3\text{)NHR}^7$, OH, (O), C(O)OH ,
 N_3 , CN, NH_2 , CF_3 , CF_2CF_3 , F, Cl, Br or I;
 R^6 is C_2 - C_5 -spiroalkyl, each of which is unsubstituted or substituted with OH, (O), N_3 , CN, CF_3 , CF_2CF_3 , F, Cl, Br,
I, NH_2 , $\text{NH(CH}_3\text{)}$ or $\text{N(CH}_3\text{)}_2$;
 R^{6A} and R^{6B} are independently selected alkyl or, together with the N to which they are attached, R^{6C} ;
30 R^{6C} is aziridin-1-yl, azetidin-1-yl, pyrrolidin-1-yl or piperidin-1-yl, each having one CH_2 moiety unreplaced or replaced
with O, C(O) , CNOH , CNOCH_3 , S, S(O) , SO_2 or NH;
 R^7 is R^8 , R^9 , R^{10} or R^{11} ;
 R^8 is phenyl, which is unfused or fused with R^{8A} ; R^{8A} is benzene, heteroarene, cycloalkane, cycloalkene, hetero-
cycloalkane or heterocycloalkene;
35 R^9 is heteroaryl, which is unfused or fused with R^{9A} ; R^{9A} is benzene, heteroarene, cycloalkane, cycloalkene, hete-
rocycloalkane or heterocycloalkene;
 R^{10} is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{10A} ;
 R^{10A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^{11} is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently
40 selected R^{12} , OR^{12} , SR^{12} , S(O)R^{12} , SO_2R^{12} , C(O)R^{12} , CO(O)R^{12} , OC(O)R^{12} , OC(O)OR^{12} , NH_2 , NHR^{12} , $\text{N(R}^{12}\text{)}_2$,
 NHC(O)R^{12} , $\text{NR}^{12}\text{C(O)R}^{12}$, $\text{NHS(O)}_2\text{R}^{12}$, $\text{NR}^{12}\text{S(O)}_2\text{R}^{12}$, NHC(O)OR^{12} , $\text{NR}^{12}\text{C(O)OR}^{12}$, NHC(O)NH_2 ,
 NHC(O)NHR^{12} , $\text{NHC(O)N(R}^{12}\text{)}_2$, $\text{NR}^{12}\text{C(O)NHR}^{12}$, $\text{NR}^{12}\text{C(O)N(R}^{12}\text{)}_2$, C(O)NH_2 , C(O)NHR^{12} , $\text{C(O)N(R}^{12}\text{)}_2$,
 C(O)NHOH , C(O)NHOR^{12} , $\text{C(O)NHSO}_2\text{R}^{12}$, $\text{C(O)NR}^{12}\text{SO}_2\text{R}^{12}$, SO_2NH_2 , $\text{SO}_2\text{NHR}^{12}$, $\text{SO}_2\text{N(R}^{12}\text{)}_2$, C(O)H ,
 C(O)OH , C(N)NH_2 , C(N)NHR^{12} , $\text{C(N)N(R}^{12}\text{)}_2$, CNOH , CNOCH_3 , OH, (O), CN, N_3 , NO_2 , CF_3 , CF_2CF_3 , OCF_3 ,
45 OCF_2CF_3 , F, Cl, Br or I;
 R^{12} is R^{13} , R^{14} , R^{15} or R^{16} ;
 R^{13} is phenyl, which is unfused or fused with R^{13A} ; R^{13A} is benzene, heteroarene, cycloalkane, cycloalkene, hete-
rocycloalkane or heterocycloalkene;
 R^{14} is heteroaryl, which is unfused or fused with R^{14A} ; R^{14A} is benzene, heteroarene, cycloalkane, cycloalkene,
50 heterocycloalkane or heterocycloalkene;
 R^{15} is cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene, each of which is unfused or fused with
 R^{15A} ; R^{15A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R^{16} is alkyl, alkenyl or alkynyl;
 R^{17} is R^{18} , R^{19} , R^{20} or R^{21} ;
55 R^{18} is phenyl, which is unfused or fused with R^{18A} ; R^{18A} is benzene, heteroarene, cycloalkane, cycloalkene, hete-
rocycloalkane or heterocycloalkene;
 R^{19} is heteroaryl, which is unfused or fused with R^{19A} ; R^{19A} is benzene, heteroarene, cycloalkane, cycloalkene,
heterocycloalkane or heterocycloalkene;

R²⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl each of which is unfused or fused with R^{20A}; R^{20A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R²², OR²², SR²², S(O)R²², SO₂R²², C(O)R²², CO(O)R²², OC(O)R²², OC(O)OR²², NH₂, NHR²², N(R²²)₂, NHC(O)R²², NR²²C(O)R²², NHS(O)₂R²², NR²²S(O)₂R²², NHC(O)OR²², NR²²C(O)OR²², NHC(O)NH₂, NHC(O)NHR²², NHC(O)N(R²²)₂, NR²²C(O)NHR²², NR²²C(O)N(R²²)₂, C(O)NH₂, C(O)NHR²², C(O)N(R²²)₂, C(O)NHOH, C(O)NHOR²², C(O)NHSO₂R²², C(O)NR²²SO₂R²², SO₂NH₂, SO₂NHR²², SO₂N(R²²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR²², C(N)N(R²²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R²² is R²³, R²⁴ or R²⁵;

R²³ is phenyl, which is unfused or fused with R^{23A}; R^{23A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁴ is heteroarene, which is unfused or fused with R^{24A}; R^{24A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{25A}; R^{25A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

Z¹ is R²⁶ or R²⁷;

Z² is R²⁸, R²⁹ or R³⁰;

Z^{1A} and Z^{2A} are both absent or are taken together to form CH₂, CH₂CH₂ or Z^{12A};

Z^{12A} is C₂-C₆-alkylene having one or two CH₂ moieties replaced by NH, N(CH₃), S, S(O) or SO₂;

L¹ is a R³⁷, OR³⁷, SR³⁷, S(O)R³⁷, SO₂R³⁷, C(O)R³⁷, CO(O)R³⁷, OC(O)R³⁷, OC(O)OR³⁷, NHR³⁷, C(O)NH, C(O)NR³⁷, C(O)NHOR³⁷, C(O)NHSO₂R³⁷, SO₂NH, SO₂NHR³⁷, C(N)NH, C(N)NHR³⁷;

R²⁶ is phenylene, which is unfused or fused with R^{26A}; R^{26A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁷ is heteroarylene, which is unfused or fused with R^{27A}; R^{27A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁸ is phenylene, which is unfused or fused with R^{28A}; R^{28A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁹ is heteroarylene, which is unfused or fused with R^{29A}; R^{29A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁰ is cycloalkylene, cycloalkenylene, heterocycloalkylene or heterocycloalkenylene, each of which is unfused or fused with R^{30A}; R^{30A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R³⁷ is a bond or R^{37A};

R^{37A} is alkylene, alkenylene, or alkynylene, each of which is unsubstituted or substituted with one or two or three independently selected R^{37B}, OR^{37B}, SR^{37B}, S(O)R^{37B}, SO₂R^{37B}, C(O)R^{37B}, CO(O)R^{37B}, OC(O)R^{37B}, OC(O)OR^{37B}, NH₂, NHR^{37B}, N(R^{37B})₂, NHC(O)R^{37B}, NR^{37B}C(O)R^{37B}, NHS(O)₂R^{37B}, NR^{37B}S(O)₂R^{37B}, NHC(O)OR^{37B}, NR^{37B}C(O)OR^{37B}, NHC(O)NH₂, NHC(O)NHR^{37B}, NHC(O)N(R^{37B})₂, NR^{37B}C(O)NHR^{37B}, NR^{37B}C(O)N(R^{37B})₂, C(O)NH₂, C(O)NHR^{37B}, C(O)N(R^{37B})₂, C(O)NHOH, C(O)NHOR^{37B}, C(O)NHSO₂R^{37B}, C(O)NR^{37B}SO₂R^{37B}, SO₂NH₂, SO₂NHR^{37B}, SO₂N(R^{37B})₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR^{37B}, C(N)N(R^{37B})₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I substituents;

R^{37B} is alkyl, alkenyl, alkynyl, or R^{37C};

R^{37C} is phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl;

Z³ is R³⁸, R³⁹ or R⁴⁰;

R³⁸ is phenyl, which is unfused or fused with R^{38A}; R^{38A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁹ is heteroaryl, which is unfused or fused with R^{39A}; R^{39A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁴⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{40A}; R^{40A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

wherein the moieties represented by R²⁶ and R²⁷ are substituted (i.e., if Z^{1A} and Z^{2A} are absent) or further substituted (i.e., if Z^{1A} and Z^{2A} are present) with R⁴¹, OR⁴¹, SR⁴¹, S(O)R⁴¹, SO₂R⁴¹, C(O)R⁴¹, CO(O)R⁴¹, OC(O)R⁴¹, OC(O)OR⁴¹, NHR⁴¹, N(R⁴¹)₂, NHC(O)R⁴¹, NR⁴¹C(O)R⁴¹, NHS(O)₂R⁴¹, NR⁴¹S(O)₂R⁴¹, NHC(O)OR⁴¹, NR⁴¹C(O)OR⁴¹, NHC(O)NHR⁴¹, NHC(O)N(R⁴¹)₂, NR⁴¹C(O)NHR⁴¹, NR⁴¹C(O)N(R⁴¹)₂, C(O)NH₂, C(O)NHR⁴¹, C(O)N(R⁴¹)₂, C(O)NHOH, C(O)NHOR⁴¹, C(O)NHSO₂R⁴¹, C(O)NR⁴¹SO₂R⁴¹, SO₂NH₂, SO₂NHR⁴¹, SO₂N(R⁴¹)₂, C(N)NH₂, C(N)NHR⁴¹, or C(N)N(R⁴¹)₂;

R⁴¹ is heteroaryl, which is fused with R^{43A}; R^{43A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; which is unfused;

wherein the cyclic moieties represented by E¹ and Y¹ together, Y¹ and B¹ together, A² and B¹ together, A² and D¹ together, R^{1A}, R², R^{2A}, R³, R^{3A}, R⁴, R^{4A}, R⁶, R^{6C}, R⁸, R^{8A}, R⁹, R^{9A}, R¹⁰, R^{10A}, R¹³, R^{13A}, R¹⁴, R^{14A}, R¹⁵, R^{15A}, R¹⁸,

R^{18A}, R¹⁹, R^{19A}, R²⁰, R^{20A}, R²³, R^{23A}, R²⁴, R^{24A}, R²⁵, R^{25A}, R²⁶, R^{26A}, R²⁷, R^{27A}, R²⁸, R^{28A}, R²⁹, R^{29A}, R³⁰, R^{30A}, R^{37B}, R³⁸, R^{38A}, R³⁹, R^{39A}, R⁴⁰, and R^{40A} are independently unsubstituted, further unsubstituted, substituted or further substituted with one or two or three or four or five of independently selected R^{57A}, R⁵⁷, OR⁵⁷, SR⁵⁷, S(O)R⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, OC(O)R⁵⁷, OC(O)OR⁵⁷, NH₂, NHR⁵⁷, N(R⁵⁷)₂, NHC(O)R⁵⁷, NR⁵⁷C(O)R⁵⁷, NHS(O)₂R⁵⁷, NR⁵⁷S(O)₂R⁵⁷, NHC(O)OR⁵⁷, NR⁵⁷C(O)OR⁵⁷, NHC(O)NH₂, NHC(O)NHR⁵⁷, NHC(O)N(R⁵⁷)₂, NR⁵⁷C(O)NHR⁵⁷, NR⁵⁷C(O)N(R⁵⁷)₂, C(O)NH₂, C(O)NHR⁵⁷, C(O)N(R⁵⁷)₂, C(O)NHOH, C(O)NHOR⁵⁷, C(O)NHSO₂R⁵⁷, C(O)NR⁵⁷SO₂R⁵⁷, SO₂NH₂, SO₂NHR⁵⁷, SO₂N(R⁵⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁵⁷, C(N)N(R⁵⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I; R^{57A} is spiroalkyl, or spiroheteroalkyl;

R⁵⁷ is R⁵⁸, R⁵⁹, R⁶⁰ or R⁶¹;

R⁵⁸ is phenyl, which is unfused or fused with R^{58A}; R^{58A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁵⁹ is heteroaryl, which is unfused or fused with R^{59A}; R^{59A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{60A}; R^{60A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶², OR⁶², SR⁶², S(O)R⁶², SO₂R⁶², C(O)R⁶², CO(O)R⁶², OC(O)R⁶², OC(O)OR⁶², NH₂, NHR⁶², N(R⁶²)₂, NHC(O)R⁶², NR⁶²C(O)R⁶², NHS(O)₂R⁶², NR⁶²S(O)₂R⁶², NHC(O)OR⁶², NR⁶²C(O)OR⁶², NHC(O)NH₂, NHC(O)NHR⁶², NHC(O)N(R⁶²)₂, NR⁶²C(O)NHR⁶², NR⁶²C(O)N(R⁶²)₂, C(O)NH₂, C(O)NHR⁶², C(O)N(R⁶²)₂, C(O)NHOH, C(O)NHOR⁶², C(O)NHSO₂R⁶², C(O)NR⁶²SO₂R⁶², SO₂NH₂, SO₂NHR⁶², SO₂N(R⁶²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶², C(N)N(R⁶²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R⁶² is R⁶³, R⁶⁴, R⁶⁵ or R⁶⁶;

R⁶³ is phenyl, which is unfused or fused with R^{63A}; R^{63A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁴ is heteroaryl, which is unfused or fused with R^{64A}; R^{64A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{65A}; R^{65A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁶ is alkyl, alkenyl or alkenyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶⁷, OR⁶⁷, SR⁶⁷, S(O)R⁶⁷, SO₂R⁶⁷, C(O)R⁶⁷, CO(O)R⁶⁷, OC(O)R⁶⁷, OC(O)OR⁶⁷, NH₂, NHR⁶⁷, N(R⁶⁷)₂, NHC(O)R⁶⁷, NR⁶⁷C(O)R⁶⁷, NHS(O)₂R⁶⁷, NR⁶⁷S(O)₂R⁶⁷, NHC(O)OR⁶⁷, NR⁶⁷C(O)OR⁶⁷, NHC(O)NH₂, NHC(O)NHR⁶⁷, NHC(O)N(R⁶⁷)₂, NR⁶⁷C(O)NHR⁶⁷, NR⁶⁷C(O)N(R⁶⁷)₂, C(O)NH₂, C(O)NHR⁶⁷, C(O)N(R⁶⁷)₂, C(O)NHOH, C(O)NHOR⁶⁷, C(O)NHSO₂R⁶⁷, C(O)NR⁶⁷SO₂R⁶⁷, SO₂NH₂, SO₂NHR⁶⁷, SO₂N(R⁶⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁷, C(N)N(R⁶⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I substituents;

R⁶⁷ is alkyl, alkenyl, alkynyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; wherein the cyclic moieties represented by R^{57A}, R⁵⁸, R⁵⁹, R⁶⁰, R⁶³, R⁶⁴, R⁶⁵, and R⁶⁷ are unsubstituted or substituted with one or two or three or four of independently selected R⁶⁸, OR⁶⁸, SR⁶⁸, S(O)R⁶⁸, SO₂R⁶⁸, C(O)R⁶⁸, CO(O)R⁶⁸, OC(O)R⁶⁸, OC(O)OR⁶⁸, NH₂, NHR⁶⁸, N(R⁶⁸)₂, NHC(O)R⁶⁸, NR⁶⁸C(O)R⁶⁸, NHS(O)₂R⁶⁸, NR⁶⁸S(O)₂R⁶⁸, NHC(O)OR⁶⁸, NR⁶⁸C(O)OR⁶⁸, NHC(O)NH₂, NHC(O)NHR⁶⁸, NHC(O)N(R⁶⁸)₂, NR⁶⁸C(O)NHR⁶⁸, NR⁶⁸C(O)N(R⁶⁸)₂, C(O)NH₂, C(O)NHR⁶⁸, C(O)N(R⁶⁸)₂, C(O)NHOH, C(O)NHOR⁶⁸, C(O)NHSO₂R⁶⁸, C(O)NR⁶⁸SO₂R⁶⁸, SO₂NH₂, SONHR⁶⁸, SO₂N(R⁶⁸)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁸, C(N)N(R⁶⁸)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R⁶⁸ is R⁶⁹, R⁷⁰, R⁷¹ or R⁷²;

R⁶⁹ is phenyl, which is unfused or fused with R^{69A}; R^{69A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁷⁰ is heteroaryl, which is unfused or fused with R^{70A}; R^{70A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁷¹ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{71A}; R^{71A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁷² is alkyl, alkenyl or alkenyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁷³, OR⁷³, SR⁷³, S(O)R⁷³, SO₂R⁷³, C(O)R⁷³, CO(O)R⁷³, OC(O)R⁷³, OC(O)OR⁷³, NH₂, NHR⁷³, N(R⁷³)₂, NHC(O)R⁷³, NR⁷³C(O)R⁷³, NHS(O)₂R⁷³, NR⁷³S(O)₂R⁷³, NHC(O)OR⁷³, NR⁷³C(O)OR⁷³, NHC(O)NH₂, NHC(O)NHR⁷³, NHC(O)N(R⁷³)₂, NR⁷³C(O)NHR⁷³, NR⁷³C(O)N(R⁷³)₂, C(O)NH₂, C(O)NHR⁷³, C(O)N(R⁷³)₂, C(O)NHOH, C(O)NHOR⁷³, C(O)NHSO₂R⁷³, C(O)NR⁷³SO₂R⁷³, SO₂NH₂, SO₂NHR⁷³, SO₂N(R⁷³)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁷³, C(N)N(R⁷³)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃,

OCF₂CF₃, F, Cl, Br or I;

R⁷³ is alkyl, alkenyl, alkenyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; and wherein the moieties represented by R⁶⁹, R⁷⁰, and R⁷¹ are unsubstituted or substituted with one or two or three or four of independently selected NH₂, C(O)NH₂, C(O)NHOH, SO₂NH₂, CF₃, CF₂CF₃, C(O)H, C(O)OH, C(N)NH₂, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I.

[0074] Another embodiment of this invention pertains to compounds of Formula (I), wherein

A¹ is N or C(A²);

A² is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R²)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};

B¹ is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};

D¹ is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};

E¹ is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; and

Y¹ is H, CN, NO₂, C(O)OH, F, Cl, Br, I, CF₃, OCF₃, CF₂CF₃, OCF₂CF₃, R¹⁷, OR¹⁷, C(O)R¹⁷, C(O)OR¹⁷, SR¹⁷, SO₂R¹⁷, NH₂, NHR¹⁷, N(R¹⁷)₂, NHC(O)R¹⁷, C(O)NH₂, C(O)NHR¹⁷, C(O)N(R¹⁷)₂, NHS(O)R¹⁷ or NHSO₂R¹⁷; or E¹ and Y¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

A², B¹, and D¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; or

Y¹ and B¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

A, D¹, and E¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; or

A² and B¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

D¹, E¹, and Y¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNHOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN,

NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; or
 A² and D¹, together with the atoms to which they are attached, are benzene, naphthalene, heteroarene, cycloalkane,
 cycloalkene, heterocycloalkane or heterocycloalkene; and
 B¹, E¹, and Y¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹,
 5 N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹,
 NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹,
 NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNH₂, C(O)NHNOR¹, C(O)NHSO₂R¹,
 C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN,
 10 NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};
 G¹ is H, or C(O)OR;
 R is alkyl;
 R¹ is R², R³, R⁴ or R⁵;
 R^{1A} is cycloalkyl, cycloalkenyl or cycloalkynyl;
 R² is phenyl, which is unfused or fused with R^{2A}; R^{2A} is benzene, heteroarene, cycloalkane, cycloalkene, hetero-
 15 cycloalkane or heterocycloalkene;
 R³ is heteroaryl, which is unfused or fused with R^{3A}; R^{3A} is benzene, heteroarene, cycloalkane, cycloalkene, hete-
 rocycloalkane or heterocycloalkene;
 R⁴ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{4A};
 R^{4A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 20 R⁵ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently
 selected R⁶, NC(R^{6A})(R^{6B}), R⁷, OR⁷, SR⁷, S(O)R⁷, SO₂R⁷, NHR⁷, N(R⁷)₂, C(O)R⁷, C(O)NH₂, C(O)NHR⁷,
 C(O)N(R⁷)₂, NHC(O)R⁷, NR⁷C(O)R⁷, NHSO₂R⁷, NHC(O)OR⁷, SO₂NH₂, SO₂NHR⁷, SO₂N(R⁷)₂, NHC(O)NH₂,
 NHC(O)NHR⁷, NHC(O)CH(CH₃)NHC(O)CH(CH₃)NH₂, NHC(O)CH(CH₃)NHC(O)CH(CH₃)NHR⁷, OH, (O), C(O)OH,
 N₃, CN, NH₂, CF₃, CF₂CF₃, F, Cl, Br or I;
 25 R⁶ is C₂-C₅-spiroalkyl, each of which is unsubstituted or substituted with OH, (O), N₃, CN, CF₃, CF₂CF₃, F, Cl, Br,
 I, NH₂, NH(CH₃) or N(CH₃)₂;
 R^{6A} and R^{6B} are independently selected alkyl or, together with the N to which they are attached, R^{6C};
 R^{6C} is aziridin-1-yl, azetidin-1-yl, pyrrolidin-1-yl or piperidin-1-yl, each having one CH₂ moiety unreplaced or replaced
 with O, C(O), CNOH, CNOCH₃, S, S(O), SO₂ or NH;
 30 R⁷ is R⁸, R⁹, R¹⁰ or R¹¹;
 R⁸ is phenyl, which is unfused or fused with R^{8A}; R^{8A} is benzene, heteroarene, cycloalkane, cycloalkene, hetero-
 cycloalkane or heterocycloalkene;
 R⁹ is heteroaryl, which is unfused or fused with R^{9A}; R^{9A} is benzene, heteroarene, cycloalkane, cycloalkene, hete-
 rocycloalkane or heterocycloalkene;
 35 R¹⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{10A};
 R^{10A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R¹¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently
 selected R¹², OR¹², SR¹², S(O)R¹², SO₂R¹², C(O)R¹², CO(O)R¹², OC(O)R¹², OC(O)OR¹², NH₂, NHR¹², N(R¹²)₂,
 40 NHC(O)R¹², NR¹²C(O)R¹², NHS(O)₂R¹², NR¹²S(O)₂R¹², NHC(O)OR¹², NR¹²C(O)OR¹², NHC(O)NH₂,
 NHC(O)NHR¹², NHC(O)N(R¹²)₂, NR¹²C(O)NHR¹², NR¹²C(O)N(R¹²)₂, C(O)NH₂, C(O)NHR¹², C(O)N(R¹²)₂,
 C(O)NHOH, C(O)NHOR¹², C(O)NHSO₂R¹², C(O)NR¹²SO₂R¹², SO₂NH₂, SO₂NHR¹², SO₂N(R¹²)₂, C(O)H,
 C(O)OH, C(N)NH₂, C(N)NHR¹², C(N)N(R¹²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃,
 OCF₂CF₃, F, Cl, Br or I;
 R¹² is R¹³, R¹⁴, R¹⁵ or R¹⁶;
 45 R¹³ is phenyl, which is unfused or fused with R^{13A}; R^{13A} is benzene, heteroarene, cycloalkane, cycloalkene, hete-
 rocycloalkane or heterocycloalkene;
 R¹⁴ is heteroaryl, which is unfused or fused with R^{14A}; R^{14A} is benzene, heteroarene, cycloalkane, cycloalkene,
 heterocycloalkane or heterocycloalkene;
 R¹⁵ is cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene, each of which is unfused or fused with
 50 R^{15A}; R^{15A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R¹⁶ is alkyl, alkenyl or alkynyl;
 R¹⁷ is R¹⁸, R¹⁹, R²⁰ or R²¹;
 R¹⁸ is phenyl, which is unfused or fused with R^{18A}; R^{18A} is benzene, heteroarene, cycloalkane, cycloalkene, hete-
 rocycloalkane or heterocycloalkene;
 55 R¹⁹ is heteroaryl, which is unfused or fused with R^{19A}; R^{19A} is benzene, heteroarene, cycloalkane, cycloalkene,
 heterocycloalkane or heterocycloalkene;
 R²⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl each of which is unfused or fused with R^{20A};
 R^{20A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R²² OR²², SR²², S(O)R²², SO₂R²², C(O)R²², CO(O)R²², OC(O)R²², OC(O)OR²², NH₂, NHR²², N(R²²)₂, NHC(O)R²², NR²²C(O)R²², NHS(O)₂R²², NR²²S(O)₂R²², NHC(O)OR²², NR²²C(O)OR²², NHC(O)NH₂, NHC(O)NHR²², NHC(O)N(R²²)₂, NR²²C(O)NHR²², NR²²C(O)N(R²²)₂, C(O)NH₂, C(O)NHR²², C(O)N(R²²)₂, C(O)NHOH, C(O)NHOR²², C(O)NHSO₂R²², C(O)NR²²SO₂R²², SO₂NH₂, SO₂NHR²², SO₂N(R²²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR²², C(N)N(R²²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R²² is R²³, R²⁴ or R²⁵;

R²³ is phenyl, which is unfused or fused with R^{23A}; R^{23A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁴ is heteroarene, which is unfused or fused with R^{24A}; R^{24A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{25A}; R^{25A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

Z¹ is R²⁶ or R²⁷;

Z² is R²⁸, R²⁹ or R³⁰;

Z^{1A} and Z^{2A} are both absent or are taken together to form CH₂, CH₂CH₂ or Z^{12A};

Z^{12A} is C₂-C₆-alkylene having one or two CH₂ moieties replaced by NH, N(CH₃), S, S(O) or SO₂;

L¹ is a R³⁷, OR³⁷, SR³⁷, S(O)R³⁷, SO₂R³⁷, C(O)R³⁷, CO(O)R³⁷, OC(O)R³⁷, OC(O)OR³⁷, NHR³⁷, C(O)NH, C(O)NR³⁷, C(O)NHOR³⁷, C(O)NHSO₂R³⁷, SO₂NH, SO₂NHR³⁷, C(N)NH, C(N)NHR³⁷;

R²⁶ is phenylene, which is unfused or fused with R^{26A}; R^{26A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁷ is heteroarylene, which is unfused or fused with R^{27A}; R^{27A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁸ is phenylene, which is unfused or fused with R^{28A}; R^{28A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁹ is heteroarylene, which is unfused or fused with R^{29A}; R^{29A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁰ is cycloalkylene, cycloalkenylene, heterocycloalkylene or heterocycloalkenylene, each of which is unfused or fused with R^{30A}; R^{30A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁷ is a bond or R^{37A};

R^{37A} is alkylene, alkenylene, or alkynylene, each of which is unsubstituted or substituted with one or two or three independently selected R^{37B}, OR^{37B}, SR^{37B}, S(O)R^{37B}, SO₂R^{37B}, C(O)R^{37B}, CO(O)R^{37B}, OC(O)R^{37B}, OC(O)OR^{37B}, NH₂, NHR^{37B}, N(R^{37B})₂, NHC(O)R^{37B}, NR^{37B}C(O)R^{37B}, NHS(O)₂R^{37B}, NR^{37B}S(O)₂R^{37B}, NHC(O)OR^{37B}, NR^{37B}C(O)OR^{37B}, NHC(O)NH₂, NHC(O)NHR^{37B}, NHC(O)N(R^{37B})₂, NR^{37B}C(O)NHR^{37B}, NR^{37B}C(O)N(R^{37B})₂, C(O)NH₂, C(O)NHR^{37B}, C(O)N(R^{37B})₂, C(O)NHOH, C(O)NHOR^{37B}, C(O)NHSO₂R^{37B}, C(O)NR^{37B}SO₂R^{37B}, SO₂NH₂, SO₂NHR^{37B}, SO₂N(R^{37B})₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR^{37B}, C(N)N(R^{37B})₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I substituents;

R^{37B} is alkyl, alkenyl, alkynyl, or R^{37C};

R^{37C} is phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl;

Z³ is R³⁸, R³⁹ or R⁴⁰;

R³⁸ is phenyl, which is unfused or fused with R^{38A}; R^{38A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁹ is heteroaryl, which is unfused or fused with R^{39A}; R^{39A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁴⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{40A}; R^{40A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

wherein the moieties represented by R²⁶ and R²⁷ are substituted (i.e., if Z^{1A} and Z^{2A} are absent) or further substituted (i.e., if Z^{1A} and Z^{2A} are present) with R⁴¹, OR⁴¹, SR⁴¹, S(O)R⁴¹, SO₂R⁴¹, C(O)R⁴¹, CO(O)R⁴¹, OC(O)R⁴¹, OC(O)OR⁴¹, NHR⁴¹, N(R⁴¹)₂, NHC(O)R⁴¹, NR⁴¹C(O)R⁴¹, NHS(O)₂R⁴¹, NR⁴¹S(O)₂R⁴¹, NHC(O)OR⁴¹, NR⁴¹C(O)OR⁴¹, NHC(O)NHR⁴¹, NHC(O)N(R⁴¹)₂, NR⁴¹C(O)NHR⁴¹, NR⁴¹C(O)N(R⁴¹)₂, C(O)NHR⁴¹, C(O)N(R⁴¹)₂, C(O)NHOR⁴¹, C(O)NHSO₂R⁴¹, C(O)NR⁴¹SO₂R⁴¹, SO₂NHR⁴¹, SO₂N(R⁴¹)₂, C(N)NHR⁴¹, or C(N)N(R⁴¹)₂;

R⁴¹ is heteroaryl, which is fused with R^{43A}; R^{43A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; which is fused with benzene, heteroarene or R^{43B}; R^{43B} is cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

wherein the cyclic moieties represented by E¹ and Y¹ together, Y¹ and B¹ together, A² and B¹ together, A² and D¹ together, R^{1A}, R², R^{2A}, R³, R^{3A}, R⁴, R^{4A}, R⁶, R^{6C}, R⁸, R^{8A}, R⁹, R^{9A}, R¹⁰, R^{10A}, R¹³, R^{13A}, R¹⁴, R^{14A}, R¹⁵, R^{15A}, R¹⁸, R^{18A}, R¹⁹, R^{19A}, R²⁰, R^{20A}, R²³, R^{23A}, R²⁴, R^{24A}, R²⁵, R^{25A}, R²⁶, R^{26A}, R²⁷, R^{27A}, R²⁸, R^{28A}, R²⁹, R^{29A}, R³⁰, R^{30A},

- R^{37B}, R³⁸, R^{38A}, R³⁹, R^{39A}, R⁴⁰, and R^{40A} are independently unsubstituted, further unsubstituted, substituted or further substituted with one or two or three or four or five of independently selected R^{57A}, R⁵⁷, OR⁵⁷, SR⁵⁷, S(O)R⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, OC(O)R⁵⁷, OC(O)OR⁵⁷, NH₂, NHR⁵⁷, N(R⁵⁷)₂, NHC(O)R⁵⁷, NR⁵⁷C(O)R⁵⁷, NHS(O)₂R⁵⁷, NR⁵⁷S(O)₂R⁵⁷, NHC(O)OR⁵⁷, NR⁵⁷C(O)OR⁵⁷, NHC(O)NH₂, NHC(O)NHR⁵⁷, NHC(O)N(R⁵⁷)₂, NR⁵⁷C(O)NHR⁵⁷, NR⁵⁷C(O)N(R⁵⁷)₂, C(O)NH₂, C(O)NHR⁵⁷, C(O)N(R⁵⁷)₂, C(O)NHOH, C(O)NHOR⁵⁷, C(O)NHSO₂R⁵⁷, C(O)NR⁵⁷SO₂R⁵⁷, SO₂NH₂, SO₂NHR⁵⁷, SO₂N(R⁵⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁵⁷, C(N)N(R⁵⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I; R^{57A} is spiroalkyl, or spiroheteroalkyl; R⁵⁷ is R⁵⁸, R⁵⁹, R⁶⁰ or R⁶¹;
- R⁵⁸ is phenyl, which is unfused or fused with R^{58A}; R^{58A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁵⁹ is heteroaryl, which is unfused or fused with R^{59A}; R^{59A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁶⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{60A}; R^{60A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁶¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶², OR⁶², SR⁶², S(O)R⁶², SO₂R⁶², C(O)R⁶², CO(O)R⁶², OC(O)R⁶², OC(O)OR⁶², NH₂, NHR⁶², N(R⁶²)₂, NHC(O)R⁶², NR⁶²C(O)R⁶², NHS(O)₂R⁶², NR⁶²S(O)₂R⁶², NHC(O)OR⁶², NR⁶²C(O)OR⁶², NHC(O)NH₂, NHC(O)NHR⁶², NHC(O)N(R⁶²)₂, NR⁶²C(O)NHR⁶², NR⁶²C(O)N(R⁶²)₂, C(O)NH₂, C(O)NHR⁶², C(O)N(R⁶²)₂, C(O)NHOH, C(O)NHOR⁶², C(O)NHSO₂R⁶², C(O)NR⁶²SO₂R⁶², SO₂NH₂, SO₂NHR⁶², SO₂N(R⁶²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶², C(N)N(R⁶²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I; R⁶² is R⁶³, R⁶⁴, R⁶⁵ or R⁶⁶;
- R⁶³ is phenyl, which is unfused or fused with R^{63A}; R^{63A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁶⁴ is heteroaryl, which is unfused or fused with R^{64A}; R^{64A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁶⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl, each of which is unfused or fused with R^{65A}; R^{65A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁶⁶ is alkyl, alkenyl or alkenyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶⁷, OR⁶⁷, SR⁶⁷, S(O)R⁶⁷, SO₂R⁶⁷, C(O)R⁶⁷, CO(O)R⁶⁷, OC(O)R⁶⁷, OC(O)OR⁶⁷, NH₂, NHR⁶⁷, N(R⁶⁷)₂, NHC(O)R⁶⁷, NR⁶⁷C(O)R⁶⁷, NHS(O)₂R⁶⁷, NR⁶⁷S(O)₂R⁶⁷, NHC(O)OR⁶⁷, NR⁶⁷C(O)OR⁶⁷, NHC(O)NH₂, NHC(O)NHR⁶⁷, NHC(O)N(R⁶⁷)₂, NR⁶⁷C(O)NHR⁶⁷, NR⁶⁷C(O)N(R⁶⁷)₂, C(O)NH₂, C(O)NHR⁶⁷, C(O)N(R⁶⁷)₂, C(O)NHOH, C(O)NHOR⁶⁷, C(O)NHSO₂R⁶⁷, C(O)NR⁶⁷SO₂R⁶⁷, SO₂NH₂, SO₂NHR⁶⁷, SO₂N(R⁶⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁷, C(N)N(R⁶⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I substituents; R⁶⁷ is alkyl, alkenyl, alkynyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; wherein the cyclic moieties represented by R^{57A}, R⁵⁸, R⁵⁹, R⁶⁰, R⁶³, R⁶⁴, R⁶⁵, and R⁶⁷ are unsubstituted or substituted with one or two or three or four of independently selected R⁶⁸, OR⁶⁸, SR⁶⁸, S(O)R⁶⁸, SO₂R⁶⁸, C(O)R⁶⁸, CO(O)R⁶⁸, OC(O)R⁶⁸, OC(O)OR⁶⁸, NH₂, NHR⁶⁸, N(R⁶⁸)₂, NHC(O)R⁶⁸, NR⁶⁸C(O)R⁶⁸, NHS(O)₂R⁶⁸, NR⁶⁸S(O)₂R⁶⁸, NHC(O)OR⁶⁸, NR⁶⁸C(O)OR⁶⁸, NHC(O)NH₂, NHC(O)NHR⁶⁸, NHC(O)N(R⁶⁸)₂, NR⁶⁸C(O)NHR⁶⁸, NR⁶⁸C(O)N(R⁶⁸)₂, C(O)NH₂, C(O)NHR⁶⁸, C(O)N(R⁶⁸)₂, C(O)NHOH, C(O)NHOR⁶⁸, C(O)NHSO₂R⁶⁸, C(O)NR⁶⁸SO₂R⁶⁸, SO₂NH₂, SONHR⁶⁸, SO₂N(R⁶⁸)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁸, C(N)N(R⁶⁸)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I; R⁶⁸ is R⁶⁹, R⁷⁰, R⁷¹ or R⁷²;
- R⁶⁹ is phenyl, which is unfused or fused with R^{69A}; R^{69A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁷⁰ is heteroaryl, which is unfused or fused with R^{70A}; R^{70A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁷¹ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{71A}; R^{71A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; R⁷² is alkyl, alkenyl or alkenyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁷³, OR⁷³, SR⁷³, S(O)R⁷³, SO₂R⁷³, C(O)R⁷³, CO(O)R⁷³, OC(O)R⁷³, OC(O)OR⁷³, NH₂, NHR⁷³, N(R⁷³)₂, NHC(O)R⁷³, NR⁷³C(O)R⁷³, NHS(O)₂R⁷³, NR⁷³S(O)₂R⁷³, NHC(O)OR⁷³, NR⁷³C(O)OR⁷³, NHC(O)NH₂, NHC(O)NHR⁷³, NHC(O)N(R⁷³)₂, NR⁷³C(O)NHR⁷³, NR⁷³C(O)N(R⁷³)₂, C(O)NH₂, C(O)NHR⁷³, C(O)N(R⁷³)₂, C(O)NHOH, C(O)NHOR⁷³, C(O)NHSO₂R⁷³, C(O)NR⁷³SO₂R⁷³, SO₂NH₂, SO₂NHR⁷³, SO₂N(R⁷³)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁷³, C(N)N(R⁷³)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R⁷³ is alkyl, alkenyl, alkenyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; and wherein the moieties represented by R⁶⁹, R⁷⁰, and R⁷¹ are unsubstituted or substituted with one or two or three or four of independently selected NH₂, C(O)NH₂, C(O)NHOH, SO₂NH₂, CF₃, CF₂CF₃, C(O)H, C(O)OH, C(N)NH₂, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I.

5

[0075] In one embodiment of Formula (I), A¹ is N, and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²) and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H, F, Cl, Br, or I; and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H; and G¹ is H.

10

[0076] In one embodiment of Formula (I), B¹ is R¹, OR¹, NHR¹, NHC(O)R¹, F, Cl, Br, or I. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is NHR¹, and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is OR¹, and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is Cl; and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is R¹; and G¹ is H. In another embodiment of Formula (I), A¹ is N; B¹ is NHR¹; and G¹ is H. In another embodiment of Formula (I), A¹ is N; B¹ is OR¹; and G¹ is H. In another embodiment of Formula (I), A¹ is N; B¹ is Cl; and G¹ is H. In another embodiment of Formula (I), A¹ is N; B¹ is R¹; and G¹ is H.

15

[0077] In one embodiment of Formula (I), D¹ is H or Cl. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is Cl; and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is OR¹; D¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is Cl; D¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is R¹; D¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is N; B¹ is NHR¹; D¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is N; B¹ is OR¹; D¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is N; B¹ is Cl; D¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is N; B¹ is R¹; D¹ is H; and G¹ is H.

20

[0078] In one embodiment of Formula (I), E¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is NHR¹; E¹ is H; D¹ is Cl; and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is OR¹; D¹ is H; E¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is Cl; D¹ is H; E¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is R¹; D¹ is H; E¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is N; B¹ is NHR¹; E¹ is H; D¹ is Cl, and G¹ is H. In another embodiment of Formula (I), A¹ is N; B¹ is OR¹; D¹ is H; E¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is N; B¹ is Cl; D¹ is H; E¹ is H; and G¹ is H. In another embodiment of Formula (I), A¹ is N; B¹ is R¹; D¹ is H; E¹ is H; and G¹ is H.

25

30

[0079] In one embodiment of Formula (I), Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SO₂R¹⁷, or C(O)NH₂. In another embodiment of Formula (I), Y¹ is H. In another embodiment of Formula (I), Y¹ is CN. In another embodiment of Formula (I), Y¹ is F, Cl, Br, or I. In another embodiment of Formula (I), Y¹ is CF₃. In another embodiment of Formula (I), Y¹ is SR¹⁷. In another embodiment of Formula (I), Y¹ is OR¹⁷. In another embodiment of Formula (I), Y¹ is NO₂. In another embodiment of Formula (I), Y¹ is SO₂R¹⁷; wherein R¹⁷ is as defined herein. In another embodiment of Formula (I), Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl. In another embodiment of Formula (I), Y¹ is R¹⁷; wherein R¹⁷ is alkynyl. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; G¹ is H; and Y¹ is NO₂ or SO₂R¹⁷; wherein R¹⁷ is alkyl or alkynyl. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; G¹ is H; and Y¹ is NO₂. In another embodiment of Formula (I), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; G¹ is H; and Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl substituted with three F. In another embodiment of Formula (I), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; G¹ is H; and Y¹ is NO₂ or SO₂R¹⁷; wherein R¹⁷ is alkyl or alkynyl. In another embodiment of Formula (I), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; G¹ is H; and Y¹ is NO₂. In another embodiment of Formula (I), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; G¹ is H; and Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl substituted with three F.

35

40

45

[0080] In one embodiment of Formula (I), G¹ is H; A¹ is N or C(A²); and A² is H. In another embodiment of Formula (I), G¹ is H; A¹ is N or C(A²); A² is H; and B¹ is NHR¹. In another embodiment of Formula (I), G¹ is H; A¹ is N or C(A²); A² is H; B¹ is NHR¹; and D¹ is H. In another embodiment of Formula (I), G¹ is H; A¹ is N or C(A²); A² is H; B¹ is NHR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (I), G¹ is H; A¹ is N or C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂.

50

[0081] In one embodiment of Formula (I), Y¹ and B¹, together with the atoms to which they are attached, are benzene. In another embodiment of Formula (I), Y¹ and B¹, together with the atoms to which they are attached, are heteroarene. In another embodiment of Formula (I), A¹ is C(A²); A², G¹, E¹, and D¹ are independently selected H; and Y¹ and B¹, together with the atoms to which they are attached, are benzene. In another embodiment of Formula (I), A¹ is C(A²); A², G¹, E¹, and D¹ are independently selected H; and Y¹ and B¹, together with the atoms to which they are attached, are heteroarene.

55

[0082] In one embodiment of Formula (I), R¹ is R⁴ or R⁵. In one embodiment of Formula (I), R¹ is R⁴. In one embodiment of Formula (I), R¹ is R⁵. In one embodiment of Formula (I), R¹ is R⁴; and R⁴ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl. In one embodiment of Formula (I), R¹ is R⁴; and R⁴ is cycloalkyl or heterocycloalkyl. In one

embodiment of Formula (I), R¹ is R⁴; and R⁴ is cycloalkyl. In one embodiment of Formula (I), R¹ is R⁴; and R⁴ is heterocycloalkyl.

[0083] In one embodiment of Formula (I), R¹ is R⁴; and R⁴ is cycloalkyl; wherein R⁴ is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted as defined herein. In another embodiment of Formula (I), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted with R⁵⁷ or N(R⁵⁷)₂. In another embodiment of Formula (I), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; and R⁵⁷ is R⁶⁰. In another embodiment of Formula (I), R¹ is R⁴; R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; and R⁶⁰ is heterocycloalkyl. In another embodiment of Formula (I), R¹ is R⁴; R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; R⁶⁰ is heterocycloalkyl; wherein the heterocycloalkyl ring is morpholinyl. In another embodiment of Formula (I), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted with N(R⁵⁷)₂. In another embodiment of Formula (I), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with N(R⁵⁷)₂. In another embodiment of Formula (I), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁶². In another embodiment of Formula (I), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is unsubstituted or substituted with R⁶²; R⁶² is R⁶⁵; and R⁶⁵ is cycloalkyl. In another embodiment of Formula (I), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is substituted with R⁶²; R⁶² is R⁶⁵; and R⁶⁵ is cyclopropyl.

[0084] In one embodiment of Formula (I), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein R⁴ is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is substituted as defined herein. In another embodiment of Formula (I), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is substituted with R⁵⁷. In another embodiment of Formula (I), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperidinyl or piperizinyl; and wherein the heterocycloalkyl ring is substituted with R⁵⁷; and R⁵⁷ is R⁶⁰ or R⁶¹. In another embodiment of Formula (I), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperidinyl or piperizinyl; and wherein the piperidinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰ or R⁶¹; R⁶⁰ is heterocycloalkyl; and R⁶¹ is alkyl. In another embodiment of Formula (I), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperidinyl or piperizinyl; and wherein the piperidinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; R⁶⁰ is heterocycloalkyl; wherein the heterocycloalkyl is tetrahydropyranyl. In another embodiment of Formula (I), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperidinyl or piperizinyl; and wherein the piperidinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; R⁶¹ is alkyl; and the alkyl is methyl. In another embodiment of Formula (I), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperidinyl or piperizinyl; and wherein the piperidinyl or piperizinyl ring is substituted with C(O)OR⁵⁷; R⁵⁷ is R⁶¹; R⁶¹ is alkyl; and the alkyl is methyl.

[0085] In one embodiment of Formula (I), R¹ is R⁵; and R⁵ is alkyl which is unsubstituted or substituted. In one embodiment of Formula (I), R¹ is R⁵; and R⁵ is alkyl which is unsubstituted or substituted with R⁷, OR⁷, OH, CN, or F. In another embodiment of Formula (I), R¹ is R⁵; and R⁵ is alkyl which is substituted with R⁷, OR⁷, NHR⁷, or N(R⁷)₂.

[0086] In one embodiment of Formula (I), R⁷ is R⁸, R⁹, R¹⁰ or R¹¹ which are unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R⁷ is R⁸ which is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R⁷ is R⁹ which is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R⁷ is R¹⁰ which is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R⁷ is R¹¹ which is unsubstituted or substituted as defined herein.

[0087] In one embodiment of Formula (I), R⁹ is phenyl which is unsubstituted or substituted as defined herein.

[0088] In one embodiment of Formula (I), R⁹ is heteroaryl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R⁹ is furanyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R⁹ is furanyl; which is unsubstituted.

[0089] In one embodiment of Formula (I), R¹⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, heterocycloalkyl which are unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R¹⁰ is heterocycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R¹⁰ is tetrahydrofuranyl, tetrahydropyranyl, morpholinyl, dioxanyl, oxetanyl, piperidinyl, or pyrrolidinyl, which are unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R¹⁰ is tetrahydropyranyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R¹⁰ is morpholinyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R¹⁰ is cycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R¹⁰ is cycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R¹⁰ is cyclohexyl, cyclopropyl, cyclobutyl, or bicyclo[2.2.1]heptanyl, which is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R¹⁰ is heterocycloalkyl which is fused with R^{10A}; and R^{10A} is heteroarene. In another embodiment of Formula (I), R¹⁰ is 5,6,7,8-tetrahydroimidazo[1,2-a]pyridinyl.

[0090] In one embodiment of Formula (I), R¹¹ is alkyl, alkenyl or alkynyl which are unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R¹¹ is alkyl. In another embodiment of Formula (I), R¹¹ is methyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (I), R¹¹ is alkyl; which is substituted as defined herein. In another embodiment of Formula (I), R¹¹ is alkyl; which is substituted with R¹², OR¹², or CF₃. In another embodiment of Formula (I), R¹¹ is alkyl; which is substituted with OR¹², R¹² is R¹⁶; and R¹⁶ is alkyl. In another embodiment of Formula (I), R¹¹ is alkyl; which is substituted with CF₃. In another embodiment of Formula (I), R¹¹ is alkyl; which is substituted with R¹²; R¹² is R¹⁴; and R¹⁴ is heteroaryl.

[0091] In one embodiment of Formula (I), A¹ is N or C(A²);

- 10 A² is H, F, Br, I, or Cl;
B¹ is R¹, OR¹, NHR¹, NHC(O)R¹, F, Br, I or Cl;
D¹ is H, F, Br, I, or Cl;
E¹ is H; and
Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SO₂R¹⁷, or C(O)NH₂; or
- 15 Y¹ and B¹, together with the atoms to which they are attached, are benzene, heteroarene, or heterocycloalkene; and
A², D¹, and E¹ are independently selected H;
G¹ is H, or C(O)OR;
R is alkyl;
R¹ is R⁴ or R⁵;
- 20 R⁴ is cycloalkyl, or heterocycloalkyl;
R⁵ is alkyl, or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected
R⁷, OR⁷, NHR⁷, N(R⁷)₂, CN, OH, F, Cl, Br or I;
R⁷ is R⁸, R⁹, R¹⁰, or R¹¹;
R⁸ is phenyl;
- 25 R⁹ is heteroaryl;
R¹⁰ is cycloalkyl, cycloalkenyl, or heterocycloalkyl; each of which is unfused or fused with R^{10A}, R^{10A} is heteroarene;
R¹¹ is alkyl each of which is unsubstituted or substituted with one or two or three of independently selected R¹²,
OR¹² or CF₃;
R¹² is R¹⁴ or R¹⁶;
- 30 R¹⁴ is heteroaryl;
R¹⁶ is alkyl;
R¹⁷ is R²¹;
R²¹ is alkyl, or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently
selected R²², F, Cl, Br or I;
- 35 R²² is R²⁵;
R²² is R²⁵;
R²⁵ is heterocycloalkyl;
Z¹ is R²⁶;
Z² is R³⁰;
- 40 Z^{1A} and Z^{2A} are both absent;
L¹ is a R³⁷;
R²⁶ is phenylene;
R³⁰ is heterocycloalkylene;
R³⁷ is R^{37A};
- 45 R^{37A} is alkylene;
Z³ is R³⁸, or R⁴⁰;
R³⁸ is phenyl;
R⁴⁰ is cycloalkyl, cycloalkenyl, or heterocycloalkenyl;
wherein the moiety represented by R²⁶ is substituted with OR⁴¹;
- 50 R⁴¹ is heteroaryl, which is fused with R^{43A}; R^{43A} is heteroarene; which is unfused or fused with benzene;
wherein the cyclic moieties represented by Y¹ and B¹ together, R⁴, R⁸, R¹⁰, R²⁵, R³⁰, R³⁸, and R⁴⁰ are independently
unsubstituted, further unsubstituted, substituted or further substituted with one or two or three or four or five of
independently selected R^{57A}, R⁵⁷, OR⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, C(O)N(R⁵⁷)₂, NH₂, NHR⁵⁷, N(R⁵⁷)₂,
NHC(O)R⁵⁷, NHS(O)₂R⁵⁷, OH, CN, (O), F, Cl, Br or I;
- 55 R^{57A} is spiroalkyl, or spiroheteroalkyl;
R⁵⁷ is R⁵⁸, R⁵⁹, R⁶⁰ or R⁶¹;
R⁵⁸ is phenyl;
R⁵⁹ is heteroaryl;

R⁶⁰ is cycloalkyl, or heterocycloalkyl;

R⁶¹ is alkyl which is unsubstituted or substituted with one or two or three of independently selected R⁶², OR⁶², N(R⁶²)₂, C(O)OH, CN, F, Cl, Br or I;

R⁶² is R⁶⁵, or R⁶⁶;

5 R⁶⁵ is cycloalkyl, or heterocycloalkyl;

R⁶⁶ is alkyl which is unsubstituted or substituted with OR⁶⁷;

R⁶⁷ is alkyl;

wherein the cyclic moieties represented by R^{57A}, R⁵⁸, and R⁶⁰ are unsubstituted or substituted with one or two or three or four of independently selected R⁶⁸, F, Cl, Br or I;

10 R⁶⁸ is R⁷¹ or R⁷²;

R⁷¹ is heterocycloalkyl; and

R⁷² is alkyl, which is unsubstituted or substituted with one or two F.

[0092] Still another embodiment pertains to compounds having Formula (I), which are

15

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({4-[(3-morpholin-4-ylpropyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

20

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(1-methylpiperidin-4-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

25

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(4-methylpiperazin-1-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

30

2-(9H-carbazol-4-yloxy)-4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(1-methylpiperidin-4-yl)amino]-3-nitrophenyl)sulfonyl)benzamide;

2-(9H-carbazol-4-yloxy)-4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(3-pyrrolidin-1-ylpropyl)amino]phenyl)sulfonyl)benzamide;

35

Trans-4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(4-morpholin-4-yl)cyclohexyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(2-methoxyethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

40

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(3S)-tetrahydro-2H-pyran-3-ylmethyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

45

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(3R)-tetrahydro-2H-pyran-3-ylmethyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({2-naphthylsulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

50

4-{4-[(4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(2-methoxyethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

55

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(1,4-dioxan-2-ylmethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

60

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(2,2,2-trifluoroethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(3,3,3-trifluoropropyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(2S)-1,4-dioxan-2-ylmeth-

oxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 -(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((2R)-1,4-dioxan-2-ylmethoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-([1,4-dioxan-2-ylmethyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-([4-morpholin-4-ylcyclohexyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-fluorotetrahydro-2H-pyran-4-yl]methoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-(aminocarbonyl)-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-morpholin-4-ylcyclohexyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([4-([tetrahydro-2H-pyran-4-ylmethyl]amino)-3-(trifluoromethyl)phenyl]sulfonyl)benzamide;
 4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([4-([tetrahydro-2H-pyran-4-ylmethyl]amino)-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl)benzamide;
 Trans-4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-([4-morpholin-4-ylcyclohexyl]amino)-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-([1-methylpiperidin-4-yl]amino)-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5-([4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino)sulfonyl)-2-(tetrahydro-2H-pyran-4-ylmethoxy)nicotinamide;
 N-([5-bromo-6-([1-methylpiperidin-4-yl]methoxy)pyridin-3-yl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-methylpiperidin-4-yl]methoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(1,4-dioxan-2-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-bromo-6-(1,4-dioxan-2-ylmethoxy)pyridin-3-yl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([2,2-dimethyltetrahydro-2H-pyran-4-yl]methoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-5-cyano-4-([tetrahydro-2H-pyran-4-ylmethyl]amino)phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([4-([1-acetyl]piperidin-4-yl]amino)-3-nitrophenyl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([2-chloro-5-fluoro-4-([tetrahydro-2H-pyran-4-ylmethyl]amino)phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-morpholin-4-yl]propyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-([4-fluorotetrahydro-2H-pyran-4-yl]methoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(2-morpholin-4-ylethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-([2-(2-methoxyethoxy)ethyl]sulfonyl)phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([2-(2-methoxyethoxy)ethyl]sulfonyl)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-morpholin-4-ylcyclohexyl]oxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-bromo-6-([1-tetrahydro-2H-pyran-4-yl]piperidin-4-yl)amino]pyridin-3-yl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([2-cyanoethyl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-([4-morpholin-4-ylcyclohexyl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 Trans-N-([4-([4-bis(cyclopropylmethyl)amino]cyclohexyl]amino)-3-nitrophenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-methylpiperidin-4-yl]methyl)amino]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-morpholin-3-ylmethyl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-([4-methylpiperazin-1-yl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-morpholin-4-ylbut-2-ynyl]oxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-ethynyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-oxo-3,4-dihydroquinazolin-6-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-([8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl)piperazin-1-yl)-N-([4-([4-morpholin-4-ylcyclohexyl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 Cis-4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-([4-methoxycyclohexyl]methyl)amino]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl)piperazin-1-yl)-N-([4-([4-fluorotetrahydro-2H-pyran-4-yl]methoxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 Trans-4-(4-([8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl)piperazin-1-yl)-N-([4-([4-methoxycyclohexyl]methyl)amino]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([tetrahydro-2H-pyran-4-ylmethyl]amino)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 tert-butyl 3-([4-([4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino)sulfonyl)-2-nitrophenoxymethylmorpholine-4-carboxylate;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(morpholin-3-ylmethoxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-([8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([tetrahydro-2H-pyran-4-ylmethyl]amino)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-(methylsulfonyl)piperidin-4-yl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1,1-dioxidotetrahydro-2H-thiopyran-4-yl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 N-([4-chloro-3-nitrophenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([1-(2,2,2-trifluoroethyl)piperidin-4-yl]amino)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-([1-tetrahydro-2H-pyran-4-yl]piperidin-4-yl)oxy]pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-isopropyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-({3-chloro-5-fluoro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

and therapeutically acceptable salts, and metabolites thereof.

5

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yl oxy)benzamide;

N-({5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

10

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({1-(2,2-difluoroethyl)piperidin-4-yl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1-cyclopropylpiperidin-4-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

15

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(1-morpholin-4-ylcyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(4-(dicyclopropylamino)cyclohexyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-6,6-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

20

N-({5-bromo-6-[(4-ethylmorpholin-3-yl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-ethylmorpholin-3-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

25

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(4-tetrahydro-2H-pyran-4-ylmorpholin-3-yl)methoxy]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(3S)-1-tetrahydro-2H-pyran-4-ylpiperidin-3-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-cyano-6-[(tetrahydro-2H-pyran-4-ylmethyl)amino]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

30

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1,1-dioxidothiomorpholin-4-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[[4-[(4-aminotetrahydro-2H-pyran-4-yl)methyl]amino]-3-nitrophenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

35

Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-cyano-6-[(4-morpholin-4-ylcyclohexyl)amino]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-cyano-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(1S,3R)-3-morpholin-4-ylcyclopentyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

40

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(1R,3S)-3-morpholin-4-ylcyclopentyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(morpholin-2-ylmethyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

45

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydrofuran-3-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({1-[cis-3-fluorotetrahydro-2H-pyran-4-yl]piperidin-4-yl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(1-tetrahydro-2H-pyran-4-ylazetidin-3-yl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

50

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(1-tetrahydrofuran-3-ylazetidin-3-yl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

55

4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-4-((trans-4-hydroxycyclohexyl)methoxy)-3-nitrophenyl)sulfonyl)benzamide;

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-

yl)-N-(4-((cis-4-methoxycyclohexyl)methoxy)-3-nitrophenylsulfonyl)benzamide;
 Cis-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((4-(cyclopropylamino)cyclohexyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-((4-(tetrahydro-2H-pyran-4-ylamino)cyclohexyl)amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-N-([5-bromo-6-([4-morpholin-4-ylcyclohexyl)oxy]pyridin-3-yl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((4-methoxycyclohexyl)methoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 tert-butyl 4-([4-([4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino)sulfonyl]-2-nitrophenoxymethyl)-4-fluoropiperidine-1-carboxylate;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-fluoropiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-((4-(4-tetrahydro-2H-pyran-4-yl)piperazin-1-yl)cyclohexyl)amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([[(3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(3R)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([[(3S)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(3S)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(4-methylmorpholin-2-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-(2-methoxyethyl)morpholin-2-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([4-([[(4-acetylmorpholin-2-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([trans-4-(fluoromethyl)-1-oxetan-3-ylpyrrolidin-3-yl]methoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-fluorotetrahydro-2H-pyran-4-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([1-oxetan-3-yl]piperidin-4-yl)amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-cyclobutyl]piperidin-4-yl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(3S)-1-cyclopropylpyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([1-tetrahydrofuran-3-yl]piperidin-4-yl)amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(3R)-1-cyclopropylpyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([[(3S)-1-tetrahydro-2H-pyran-4-yl]pyrrolidin-3-yl)methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(3-hydroxy-2,2-dimethylpropyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(1-(methylsulfonyl)piperidin-3-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([4-([[(1-acetyl]piperidin-3-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(3R)-1-(methylsulfonyl)pyr-

rolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-3,3-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-
 pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({1-[2-fluoro-1-(fluorome-
 5 thyl)ethyl]azetidin-3-yl}amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({1-(methylsulfonyl)pyrrolidin-3-yl}methyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[4-({1-(1-acetylpyrrolidin-3-yl)methyl}amino)-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 N-[[4-({(3R)-1-acetylpyrrolidin-3-yl}amino)-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(3-methoxy-2,2-dimethylpropyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(1R,3R)-3-hydroxycyclopentyl]methyl}amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(1S,3S)-3-hydroxycyclopentyl]methyl}amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(1S,3R)-3-hydroxycyclopentyl]methyl}amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(1R,3S)-3-hydroxycyclopentyl]methyl}amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(1R,3S)-3-hydroxycyclopentyl]methyl}amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(3S)-2-oxopiperidin-3-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1-[2-fluoro-1-(fluoromethyl)ethyl]azetidin-3-yl)methyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(1-oxetan-3-yl)azetidin-3-yl]methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(1-oxetan-3-yl)piperidin-4-yl]methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(1-cyclopropylpiperidin-4-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({4-(2-fluoroethyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({4-(2,2-difluoroethyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluoro-1-oxetan-3-yl)piperidin-4-yl]methoxy}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(2S)-4,4-difluoro-1-oxetan-3-ylpyrrolidin-2-yl]methoxy}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(4-tetrahydro-2H-pyran-4-yl)morpholin-3-yl]methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(4-cyclobutyl)morpholin-3-yl]methyl}amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(4-tetrahydrofuran-3-yl)morpholin-3-yl]methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl)methyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1-cyclopropyl-4-fluoropiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methoxybenzyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(3-(trifluoromethoxy)benzyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(3-methoxybenzyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(4-(difluoromethoxy)benzyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-(1,4-dioxaspiro[4.5]dec-8-

ylamino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-N-[[4-[(4-(acetylamino)cyclohexyl)amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3S)-1-(2-fluoroethyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3S)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3R)-1-(2-fluoroethyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3S)-1-oxetan-3-yl]pyrrolidin-3-yl]methoxy]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(4-hydroxybenzyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3-hydroxybenzyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3-difluoromethoxy)benzyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(cis-3-morpholin-4-yl)cyclopentyl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(methylsulfonyl)amino]cyclohexyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(1-cyclopropylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[(1-oxetan-3-yl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-fluoro-1-tetrahydro-2H-pyran-4-yl]piperidin-4-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-fluoro-1-tetrahydrofuran-3-yl]piperidin-4-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-fluoro-1-(methylsulfonyl)piperidin-4-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3R)-1-oxetan-3-yl]pyrrolidin-3-yl]methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-hydroxycyclohexyl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-[(3-(dimethylamino)propoxy]benzyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(2-morpholin-4-ylethoxy)benzyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(E)-4-hydroxy-1-Adamantyl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(Z)-4-hydroxy-1-Adamantyl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[4-[(1S,4S)-bicyclo[2.2.1]hept-5-en-2-yl]methoxy]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-methyl-5-oxopyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(1R,4R,5R,6S)-5,6-dihydroxybicyclo[2.2.1]hept-2-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(1R,4R,5S,6R)-5,6-dihydroxybicyclo[2.2.1]hept-2-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[(3-oxocyclohexyl)methoxy]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohexa-1,3-dien-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3R)-1-[2-fluoro-1-(fluor-

omethyl)ethyl]pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[6-[[4-fluorotetrahydro-2H-
 pyran-4-yl)methoxy]-5-(trifluoromethyl)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3S)-1-oxetan-3-
 5 ylpyrrolidin-3-yl]methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-N-([5-chloro-6-[[4-methoxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethyl-
 cyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-N-([5-chloro-6-[[4-methoxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcy-
 clohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3S)-1-oxetan-3-
 ylpyrrolidin-3-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[[2-(2-methox-
 yethoxy)ethyl]morpholin-2-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benza-
 mide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[4-(cyanomethyl)morpho-
 lin-2-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[4-(N,N-dimethylgly-
 cyl)morpholin-2-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 (2-[[[4-[[4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyrid-
 20 in-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino)methyl]morpholin-4-yl)acetic acid;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[4-(oxetan-3-
 yl)morpholin-2-yl]methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[4-(cyclopropylmorpholin-2-
 yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-(methylsulfonyl)-6-(tetrahy-
 dro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[[4-fluorotetrahydro-2H-
 pyran-4-yl)methoxy]-3-[[trifluoromethyl]sulfonyl]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[[4-methyltetrahydro-2H-
 30 pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 ethyl 4-(4-[[4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]py-
 ridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)piperazine-1-carboxylate;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[4-(morpholin-4-yl)piperidin-
 1-yl]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3R)-1-(oxetan-3-
 yl)pyrrolidin-3-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3R)-1-(1,3-difluoropropan-
 2-yl)pyrrolidin-3-yl]amino]-3-[[trifluoromethyl]sulfonyl]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benza-
 mide;
 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[[1-isopropylpiperidin-4-
 yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([4-[[[1-tert-butylpiperidin-4-yl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-
 en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[1-(2-methoxyethyl)piperi-
 45 din-3-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[1-(cyanomethyl)piperidin-
 3-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[[4-fluoro-1-methylpiperidin-
 4-yl)methoxy]-3-[[trifluoromethyl]sulfonyl]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 N-[[5-chloro-6-[[[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]amino]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophe-
 nyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 tert-butyl 4-[[4-[[4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-
 b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino]piperazine-1-carboxylate;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-(pentafluoro- λ^6 -sulfanyl)-
 55 4-[[tetrahydro-2H-pyran-4-ylmethyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[[4-methoxytetrahydro-2H-
 pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[[(3R)-1-(1,3-difluoropropan-

2-yl)pyrrolidin-3-yl]oxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(4-(oxetan-3-yl)piperazin-1-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(3R)-tetrahydrofuran-3-ylamino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[4-(4,4-difluorocyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(4-[(1-tert-butylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(4-(oxetan-3-yl)morpholin-2-yl)methyl]amino)-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[4-(4-fluorotetrahydro-2H-pyran-4-yl)methyl]amino]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(1-cyclopropylpiperidin-4-yl)amino]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(2S)-4-(cyanomethyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(2R)-4-(cyanomethyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(5-fluoro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[3-hydroxy-2-(hydroxymethyl)-2-methylpropoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(1,3-difluoropropan-2-yl)morpholin-2-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(1-(cyanomethyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(3R)-1-[2-(2-methoxyethoxy)ethyl]pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(3R)-1-(N,N-dimethylglycyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(1-(oxetan-3-yl)azetidin-3-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(1-(cyanomethyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(3R)-1-(cyanomethyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[(4'-chlorobiphenyl-2-yl)methyl]-4-methoxypiperidin-1-yl)-N-[(5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[(4'-chlorobiphenyl-2-yl)methyl]-4-methoxypiperidin-1-yl)-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl)-N-[(3-

nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-[[5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 4-(4-[[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-[[5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 N-[[5-chloro-6-[[1-(N,N-dimethylglycyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4'-chlorobiphenyl-2-yl)methyl]-4-fluoropiperidin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4'-chlorobiphenyl-2-yl)methyl]-4-fluoropiperidin-1-yl)-N-({5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-(trifluoromethyl)phenyl)sulfonyl)benzamide;
 N-[[5-chloro-6-[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[2-(tetrahydrofuran-3-yloxy)ethoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[(trans-4-cyanocyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[5-chloro-6-[(4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({5-cyano-6-[2-(tetrahydro-2H-pyran-4-yl)ethoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-(3-furylmethoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 N-[[5-chloro-6-[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[5-chloro-6-[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[5-chloro-6-[(1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 N-({3-chloro-4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[3-cyano-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 N-[[5-chloro-6-[(1-(2,2-difluoroethyl)-4-fluoropiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({3-chloro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(4,4-difluorocyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[6-[[1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl]methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({5-chloro-6-[2-(tetrahydrofuran-2-yl)ethoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]-3-methylpiperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[(3-cyclopropylamino)propyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 N-[[5-chloro-6-(2-methoxyethoxy)pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[5-fluoro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- N-[(3-chloro-4-[[1-(methoxyacetyl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(3-chloro-4-[[1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohexyl]methyl]piperidin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-[[6-(tetrahydro-2H-pyran-4-ylmethoxy)-5-(trifluoromethyl)pyridin-3-yl]sulfonyl]benzamide;
- 10 N-({5-chloro-6-[(trans-4-hydroxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-cyano-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({6-[(trans-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({6-[(cis-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-({5-chloro-6-[(4,4-difluoro-1-hydroxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperidin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 20 N-[(3-chloro-4-[[trans-4-(morpholin-4-yl)cyclohexyl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({3-[cyclopropyl(1,3-thiazol-5-ylmethyl)amino]propyl)amino]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-({3-chloro-4-[(trans-4-hydroxycyclohexyl)methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-chloro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-(trifluoromethyl)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({3-[cyclopropyl(2,2,2-trifluoroethyl)amino]propyl)amino]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 30 N-[(3-chloro-4-[[1-(oxetan-3-yl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3,5-difluoro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({3-[cyclopropyl(oxetan-3-yl)amino]propyl)amino]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(3-chloro-4-[[1-(1-methyl-L-prolyl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3,4-difluoro-5-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 N-[(5-chloro-6-{{(2S)-4-cyclopropylmorpholin-2-yl}methoxy}pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperidin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperidin-1-yl)-N-({3-chloro-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- methyl 2-[[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino]methyl]morpholine-4-carboxylate;
- 2-[[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
- 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[4-(methylsulfonyl)morpholin-2-yl]methyl]amino]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({3-[cyclobutyl(cyclopropyl)amino]propyl]amino]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 4-(4-[[2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(3-chloro-4-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-chloro-4-(tetrahydrofuran-3-ylmethoxy)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[trans-4-hydroxycyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 N-[[3-chloro-4-[[4-fluorotetrahydro-2H-pyran-4-yl]methoxy]phenyl]sulfonyl]-4-(4-[[9-(4-chlorophenyl)-3-(oxetan-3-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(2R)-4-cyclopropylmorpholin-2-yl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(2S)-4-cyclopropylmorpholin-2-yl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[tetrahydro-2H-pyran-4-yl)methyl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[5-chloro-6-[[4-[[cyclopropyl(oxetan-3-yl)amino]cyclohexyl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-N-[[4-[[[(4-cyclopropylmorpholin-2-yl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[3-chloro-4-[[4-cyclopropylmorpholin-2-yl]methoxy]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[3-chloro-4-[[[(4-cyclopropylmorpholin-2-yl)methyl]amino]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 20 2-[[[(2-chloro-4-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]phenyl]amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
- (2S)-2-[[[(3-chloro-5-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]pyridin-2-yl)oxy]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
- 25 N-[[5-chloro-6-[[[(4-cyclopropylmorpholin-2-yl)methyl]amino]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 2-[[[(3-chloro-5-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]pyridin-2-yl)amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
- 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(cis-4-hydroxy-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[5-chloro-6-[[[(1R,2R,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(4-cyanoethyl)(cyclopropyl)amino]cyclohexyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[5-chloro-6-[[[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-chloro-6-(5,6,7,8-tetrahydroimidazo[1,2-a]pyridin-6-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[5-chloro-6-[[[(1R,2S,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(cis-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[5-chloro-6-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(3,3-difluoropyrrolidin-1-yl)cyclohexyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[5-chloro-6-[[[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[5-chloro-6-[[[(cis-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-[[2-(2,2-difluorocyclopropyl)amino]cyclohexyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[5-chloro-6-[[[(cis-1-fluoro-4-hydroxycyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-

dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[3-nitro-4-(2-oxaspiro[3.5]non-7-ylmethoxy)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-{{4-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 4-(4-[[2-(4-chlorophenyl)-5,5-bis(fluoromethyl)cyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-{{3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-{{4-[(4-cyclopropylmorpholin-2-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 N-{{5-chloro-6-[(trans-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{{5-chloro-6-[(cis-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(3-cyano-4-{{4-fluoro-1-(oxetan-3-yl)piperidin-4-yl}methoxy}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-{{(trans-4-ethyl-4-hydroxycyclohexyl)methyl}amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-{{(cis-4-ethyl-4-hydroxycyclohexyl)methyl}amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[2-(4-chlorophenyl)-5-(methoxymethyl)-5-methylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-{{3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[3-nitro-4-{{(2S)-4-(oxetan-3-yl)morpholin-2-yl}methyl}amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{{3-chloro-4-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]phenyl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[4-{{4-[(2-cyanoethyl)(cyclopropyl)amino]-1-fluorocyclohexyl}methoxy}-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-{{5-nitro-6-[(tetrahydro-2H-pyran-4-ylmethyl)amino]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-{{3-nitro-4-[(2-oxaspiro[3.5]non-7-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-{{(4-cyano-4-methylcyclohexyl)methyl}amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 {{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl}{{4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl}sulfonyl)amino}methyl pivalate;
 35 {{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl}{{4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl}sulfonyl)amino}methyl butyrate;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-[[3-(tetrahydro-2H-pyran-4-ylmethyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-6-yl]sulfonyl]benzamide;
 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[6-{{(trans-4-hydroxy-4-methylcyclohexyl)methyl}amino}-5-nitropyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(5-cyano-6-{{4-fluoro-1-(oxetan-3-yl)piperidin-4-yl}methoxy}pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-4-[[4-4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)morpholine-4-carboxamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[4-{{(4-(methoxymethyl)cyclohexyl)methyl}amino}-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[5-chloro-6-{{1-(1,3-thiazol-2-yl)piperidin-4-yl}methoxy}pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[6-{{(cis-4-hydroxy-4-methylcyclohexyl)methyl}amino}-5-nitropyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-{{(trans-4-hydroxy-4-methylcyclohexyl)methyl}amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-{{3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; and therapeutically acceptable salts, and metabolites thereof.

[0093] Still another embodiment pertains to 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piper-

azin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; and therapeutically acceptable salts, and metabolites thereof.

[0094] Still another embodiment pertains to Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; and therapeutically acceptable salts, and metabolites thereof.

[0095] Still another embodiment pertains to Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methoxycyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; and therapeutically acceptable salts, and metabolites thereof.

[0096] Still another embodiment pertains to Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methoxycyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; and therapeutically acceptable salts, and metabolites thereof.

[0097] Still another embodiment pertains to 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; and therapeutically acceptable salts, and metabolites thereof.

[0098] Another embodiment pertains to the compound N-([5-chloro-6-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

[0099] Another embodiment pertains to the compound 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[trans-4-hydroxy-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0100] Another embodiment pertains to the compound 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[cis-4-hydroxy-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0101] Another embodiment pertains to the compound 4-[4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-(1-{3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0102] Another embodiment pertains to the compound N-([5-bromo-6-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0103] Another embodiment pertains to the compound 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[3-(3R)-1-(methylsulfonyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0104] Another embodiment pertains to the compound 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[3-(cyclopropyl(oxetan-3-yl)amino)propyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0105] Another embodiment pertains to the compound 4-(4-[[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0106] Still another embodiment pertains to compounds having Formula (I) or Formula (II), which are 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[trans-4-hydroxy-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[3-(3R)-1-(methylsulfonyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[3-(cyclopropyl(oxetan-3-yl)amino)propyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0107] Still another embodiment pertains to compounds having Formula (I) or Formula (II), which are N-([5-chloro-6-[[1-(cyanomethyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[3-(cyclopropyl(oxetan-3-yl)amino)propyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0108] Still another embodiment pertains to compounds having Formula (I) or Formula (II), which are 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[trans-4-hydroxy-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[cis-4-hydroxy-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

[0109] Another embodiment pertains to the compound 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[3-(cyclopropyl(oxetan-3-yl)amino)propyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0110] Still another embodiment pertains to compounds having Formula (I) or Formula (II), which are 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-((3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl)methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methylmorpholin-2-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0111] Still another embodiment pertains to compounds having Formula (I) or Formula (II), which are

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-((3S)-tetrahydro-2H-pyran-3-ylmethyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-([5-bromo-6-([1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-([4-([4-aminotetrahydro-2H-pyran-4-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-((3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl)methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methylmorpholin-2-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((3R)-1-(methylsulfonyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methoxytetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-([5-chloro-6-([1-(cyanomethyl)piperidin-4-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-(cyc1opropyl(oxetan-3-yl)amino)propyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-([5-chloro-6-([1R,2S,4R,5R]-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([6-([cis-4-hydroxy-4-methylcyclohexyl)methyl]amino)-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-([2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([tetrahydro-2H-pyran-4-ylmethyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0112] Another embodiment pertains to the compound N-([5-chloro-6-([1-(cyanomethyl)piperidin-4-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

[0113] Still another embodiment pertains to compounds having Formula (I) or Formula (II), which are N-([4-([4-aminotetrahydro-2H-pyran-4-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methoxytetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.


[0114] Another embodiment pertains to the compound 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methoxytetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

[0115] In another aspect, the present invention provides compounds of Formula (II)

5

10

15



(II)

and therapeutically acceptable salts, and metabolites thereof,

wherein A¹, B¹, D¹, E¹, Y¹, Z², L¹, and Z³ are as described herein for Formula (II); n is 0, 1, 2, or 3; describing the number of substituents on Z¹; and R¹⁰⁰ is as described for substituents on R²⁶.

[0116] In one embodiment of Formula (II), n is 0 or 1. In another embodiment of Formula (II), n is 0.

[0117] In one embodiment of Formula (II), A¹ is N. In another embodiment of Formula (II), A¹ is C(A²). In another embodiment of Formula (II), A¹ is C(A²); and A² is H, F, Cl, Br, or I. In another embodiment of Formula (II), A¹ is C(A²); and A² is H.

[0118] In one embodiment of Formula (II), B¹ is R¹, OR¹, NHR¹, NHC(O)R¹, F, Cl, Br, or I. In another embodiment of Formula (II), A¹ is C(A²); A² is H; and B¹ is NHR¹. In another embodiment of Formula (II), A¹ is C(A²); A² is H; and B¹ is OR¹. In another embodiment of Formula (II), A¹ is C(A²); A² is H; and B¹ is Cl. In another embodiment of Formula (II), A¹ is C(A²); A² is H; and B¹ is R¹. In another embodiment of Formula (II), A¹ is N; and B¹ is NHR¹. In another embodiment of Formula (II), A¹ is N; and B¹ is OR¹. In another embodiment of Formula (II), A¹ is N; and B¹ is Cl. In another embodiment of Formula (II), A¹ is N; and B¹ is R¹.

[0119] In one embodiment of Formula (II), D¹ is H or Cl. In another embodiment of Formula (II), A¹ is C(A²); A² is H; B¹ is NHR¹; and D¹ is H. In another embodiment of Formula (II), A¹ is C(A²); A² is H; B¹ is NHR¹; and D¹ is Cl. In another embodiment of Formula (II), A¹ is C(A²); A² is H; B¹ is OR¹; and D¹ is H. In another embodiment of Formula (II), A¹ is C(A²); A² is H; B¹ is Cl; and D¹ is H. In another embodiment of Formula (II), A¹ is C(A²); A² is H; B¹ is R¹; and D¹ is H. In another embodiment of Formula (II), A¹ is N; B¹ is NHR¹; and D¹ is H. In another embodiment of Formula (II), A¹ is N; B¹ is NHR¹; and D¹ is Cl. In another embodiment of Formula (II), A¹ is N; B¹ is OR¹; and D¹ is H. In another embodiment of Formula (II), A¹ is N; B¹ is Cl; and D¹ is H. In another embodiment of Formula (II), A¹ is N; B¹ is R¹; and D¹ is H.

[0120] In one embodiment of Formula (II), E¹ is H. In another embodiment of Formula (II), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (II), A¹ is C(A²); A² is H; B¹ is NHR¹; E¹ is H; and D¹ is Cl. In another embodiment of Formula (II), A¹ is C(A²); A² is H; B¹ is OR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (II), A¹ is C(A²); A² is H; B¹ is Cl; D¹ is H; and E¹ is H. In another embodiment of Formula (II), A¹ is C(A²); A² is H; B¹ is R¹; D¹ is H; and E¹ is H. In another embodiment of Formula (II), A¹ is N; B¹ is NHR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (II), A¹ is N; B¹ is NHR¹; E¹ is H; and D¹ is Cl. In another embodiment of Formula (II), A¹ is N; B¹ is OR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (II), A¹ is N; B¹ is Cl; D¹ is H; and E¹ is H. In another embodiment of Formula (II), A¹ is N; B¹ is R¹; D¹ is H; and E¹ is H.

[0121] In one embodiment of Formula (II), Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SO₂R¹⁷, or C(O)NH₂. In another embodiment of Formula (II), Y¹ is H. In another embodiment of Formula (II), Y¹ is CN. In another embodiment of Formula (II), Y¹ is F, Cl, Br, or I. In another embodiment of Formula (II), Y¹ is CF₃. In another embodiment of Formula (II), Y¹ is SR¹⁷. In another embodiment of Formula (II), Y¹ is OR¹⁷. In another embodiment of Formula (II), Y¹ is NO₂. In another embodiment of Formula (II), Y¹ is SO₂R¹⁷; wherein R¹⁷ is as defined herein. In another embodiment of Formula (II), Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl. In another embodiment of Formula (II), Y¹ is R¹⁷; wherein R¹⁷ is alkynyl. In another embodiment of Formula (II), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂ or SO₂R¹⁷; wherein R¹⁷ is alkyl or alkynyl. In another embodiment of Formula (II), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂. In another embodiment of Formula (II), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl substituted with three F. In another embodiment of Formula (II), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂ or SO₂R¹⁷; wherein R¹⁷ is alkyl or alkynyl. In another embodiment of Formula (II), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂. In another embodiment of Formula (II), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl substituted with three F.

[0122] In one embodiment of Formula (II), G^1 is H; A^1 is N or $C(A^2)$; and A^2 is H. In another embodiment of Formula (II), G^1 is H; A^1 is N or $C(A^2)$; A^2 is H; and B^1 is NHR^1 . In another embodiment of Formula (II), G^1 is H; A^1 is N or $C(A^2)$; A^2 is H; B^1 is NHR^1 ; D^1 is H; and E^1 is H. In another embodiment of Formula (II), G^1 is H; A^1 is N or $C(A^2)$; A^2 is H; B^1 is NHR^1 ; D^1 is H; and E^1 is H. In another embodiment of Formula (II), G^1 is H; A^1 is N or $C(A^2)$; A^2 is H; B^1 is NHR^1 ; D^1 is H; and E^1 is H; and Y^1 is NO_2 .

[0123] In one embodiment of Formula (II), Y^1 and B^1 , together with the atoms to which they are attached, are benzene. In another embodiment of Formula (II), Y^1 and B^1 , together with the atoms to which they are attached, are heteroarene. In another embodiment of Formula (II), A^1 is $C(A^2)$; A^2 , G^1 , E^1 , and D^1 are independently selected H; and Y^1 and B^1 , together with the atoms to which they are attached, are benzene. In another embodiment of Formula (II), A^1 is $C(A^2)$; A^2 , G^1 , E^1 , and D^1 are independently selected H; and Y^1 and B^1 , together with the atoms to which they are attached, are heteroarene.

[0124] In one embodiment of Formula (II), R^1 is R^4 or R^5 . In one embodiment of Formula (II), R^1 is R^4 . In one embodiment of Formula (II), R^1 is R^5 . In one embodiment of Formula (II), R^1 is R^4 ; and R^4 is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl. In one embodiment of Formula (II), R^1 is R^4 ; and R^4 is cycloalkyl or heterocycloalkyl. In one embodiment of Formula (II), R^1 is R^4 ; and R^4 is cycloalkyl. In one embodiment of Formula (II), R^1 is R^4 ; and R^4 is heterocycloalkyl.

[0125] In one embodiment of Formula (II), R^1 is R^4 ; and R^4 is cycloalkyl; wherein R^4 is unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R^1 is R^4 ; and R^4 is cycloalkyl; wherein the cycloalkyl ring is substituted as defined herein. In another embodiment of Formula (II), R^1 is R^4 ; and R^4 is cycloalkyl; wherein the cycloalkyl ring is substituted with R^{57} or $N(R^{57})_2$. In another embodiment of Formula (II), R^1 is R^4 ; and R^4 is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R^{57} ; and R^{57} is R^{60} . In another embodiment of Formula (II), R^1 is R^4 ; R^4 is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R^{57} ; R^{57} is R^{60} ; and R^{60} is heterocycloalkyl. In another embodiment of Formula (II), R^1 is R^4 ; R^4 is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R^{57} ; R^{57} is R^{60} ; R^{60} is heterocycloalkyl; wherein the heterocycloalkyl ring is morpholinyl. In another embodiment of Formula (II), R^1 is R^4 ; and R^4 is cycloalkyl; wherein the cycloalkyl ring is substituted with $N(R^{57})_2$. In another embodiment of Formula (II), R^1 is R^4 ; and R^4 is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with $N(R^{57})_2$. In another embodiment of Formula (II), R^1 is R^4 ; and R^4 is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with $N(R^{57})_2$; R^{57} is R^{61} ; and R^{61} is alkyl which is unsubstituted or substituted with R^{62} . In another embodiment of Formula (II), R^1 is R^4 ; and R^4 is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R^{57} ; R^{57} is R^{61} ; and R^{61} is alkyl which is unsubstituted or substituted with R^{62} ; R^{62} is R^{65} ; and R^{65} is cycloalkyl. In another embodiment of Formula (II), R^1 is R^4 ; and R^4 is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R^{57} ; R^{57} is R^{61} ; and R^{61} is alkyl which is substituted with R^{62} ; R^{62} is R^{65} ; and R^{65} is cyclopropyl.

[0126] In one embodiment of Formula (II), R^1 is R^4 ; and R^4 is heterocycloalkyl; wherein R^4 is unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R^1 is R^4 ; and R^4 is heterocycloalkyl; wherein the heterocycloalkyl ring is substituted as defined herein. In another embodiment of Formula (II), R^1 is R^4 ; and R^4 is heterocycloalkyl; wherein the heterocycloalkyl ring is substituted with R^{57} . In another embodiment of Formula (II), R^1 is R^4 ; and R^4 is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the heterocycloalkyl ring is substituted with R^{57} ; and R^{57} is R^{60} or R^{61} . In another embodiment of Formula (II), R^1 is R^4 ; R^4 is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with R^{57} ; R^{57} is R^{60} or R^{61} ; R^{60} is heterocycloalkyl; and R^{61} is alkyl. In another embodiment of Formula (II), R^1 is R^4 ; R^4 is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with R^{57} ; R^{57} is R^{60} ; R^{60} is heterocycloalkyl; wherein the heterocycloalkyl is tetrahydropyranyl. In another embodiment of Formula (II), R^1 is R^4 ; R^4 is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with R^{57} ; R^{57} is R^{61} ; R^{61} is alkyl; and the alkyl is methyl. In another embodiment of Formula (II), R^1 is R^4 ; R^4 is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with $C(O)OR^{57}$; R^{57} is R^{61} ; R^{61} is alkyl; and the alkyl is methyl.

[0127] In one embodiment of Formula (II), R^1 is R^5 ; and R^5 is alkyl which is unsubstituted or substituted. In one embodiment of Formula (II), R^1 is R^5 ; and R^5 is alkyl which is unsubstituted or substituted with R^7 , OR^7 , OH , CN , or F . In another embodiment of Formula (II), R^1 is R^5 ; and R^5 is alkyl which is substituted with R^7 , OR^7 , NHR^7 , or $N(R^7)_2$.

[0128] In one embodiment of Formula (II), R^7 is R^8 , R^9 , R^{10} or R^{11} which are unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R^7 is R^8 which is unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R^7 is R^9 which is unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R^7 is R^{10} which is unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R^7 is R^{11} which is unsubstituted or substituted as defined herein.

[0129] In one embodiment of Formula (II), R^8 is phenyl which is unsubstituted or substituted as defined herein.

[0130] In one embodiment of Formula (II), R^9 is heteroaryl which is unsubstituted or substituted as defined herein. In

another embodiment of Formula (II), R⁹ is furanyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R⁹ is furanyl; which is unsubstituted.

[0131] In one embodiment of Formula (II), R¹⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, heterocycloalkyl which are unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R¹⁰ is heterocycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R¹⁰ is tetrahydrofuranyl, tetrahydropyranyl, morpholinyl, dioxanyl, oxetanyl, piperdinyl, or pyrrolidinyl, which are unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R¹⁰ is tetrahydropyranyl, which is unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R¹⁰ is morpholinyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R¹⁰ is cycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R¹⁰ is cycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R¹⁰ is cyclohexyl, cyclopropyl, cyclobutyl, or bicyclo[2.2.1]heptanyl, which is unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R¹⁰ is heterocycloalkyl which is fused with R^{10A}; and R^{10A} is heteroarene. In another embodiment of Formula (II), R¹⁰ is 5,6,7,8-tetrahydroimidazo[1,2-a]pyridinyl.

[0132] In one embodiment of Formula (II), R¹¹ is alkyl, alkenyl or alkynyl which are unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R¹¹ is alkyl. In another embodiment of Formula (II), R¹¹ is methyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (II), R¹¹ is alkyl; which is substituted as defined herein. In another embodiment of Formula (II), R¹¹ is alkyl; which is substituted with R¹², OR¹², or CF₃. In another embodiment of Formula (II), R¹¹ is alkyl; which is substituted with OR¹²; R¹² is R¹⁶; and R¹⁶ is alkyl. In another embodiment of Formula (II), R¹¹ is alkyl; which is substituted with CF₃. In another embodiment of Formula (II), R¹¹ is alkyl; which is substituted with R¹², R¹² is R¹⁴, and R¹⁴ is heteroaryl.

[0133] In one embodiment of Formula (II),

n is 0;

A¹ is N or C(A²);

A² is H, F, Br, I, or Cl;

B¹ is R¹, OR¹, NHR¹, NHC(O)R¹, F, Br, I or Cl;

D¹ is H, F, Br, I, or Cl;

E¹ is H; and

Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SOR¹⁷, or C(O)NH₂; or

Y¹ and B¹, together with the atoms to which they are attached, are benzene, heteroarene, or heterocycloalkene; and

A², D¹, and E¹ are independently selected H;

R¹ is R⁴ or R⁵;

R⁴ is cycloalkyl, or heterocycloalkyl;

R⁵ is alkyl, or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁷, OR⁷, NHR⁷, N(R⁷)₂, CN, OH, F, Cl, Br or I;

R⁷ is R⁸, R⁹, R¹⁰, or R¹¹;

R⁸ is phenyl;

R⁹ is heteroaryl;

R¹⁰ is cycloalkyl, cycloalkenyl, or heterocycloalkyl; each of which is unfused or fused with R^{10A}, R^{10A} is heteroarene;

R¹¹ is alkyl each of which is unsubstituted or substituted with one or two or three of independently selected R¹², OR¹² or CF₃;

R¹² is R¹⁴ or R¹⁶;

R¹⁴ is heteroaryl;

R¹⁶ is alkyl;

R¹⁷ is R²¹;

R²¹ is alkyl, or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R²², F, Cl, Br or I;

R²² is R²⁵;

R²⁵ is heterocycloalkyl;

Z² is R³⁰;

Z^{1A} and Z^{2A} are both absent;

L¹ is a R³⁷;

R³⁰ is heterocycloalkylene;

R³⁷ is R^{37A};

R^{37A} is alkylene;

Z³ is R³⁸ or R⁴⁰;

R³⁸ is phenyl;

R⁴⁰ is cycloalkyl, cycloalkenyl, or heterocycloalkenyl;

wherein the cyclic moieties represented by Y¹ and B¹ together, R⁴, R⁸, R¹⁰, R²⁵, R³⁰, R³⁸, and R⁴⁰ are independently unsubstituted, further unsubstituted, substituted or further substituted with one or two or three or four or five of independently selected R^{57A}, R⁵⁷, OR⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, C(O)N(R⁵⁷)₂, NH₂, NHR⁵⁷, N(R⁵⁷)₂, NHC(O)R⁵⁷, NHS(O)₂R⁵⁷, OH, CN, (O), F, Cl, Br or I;

R^{57A} is spiroalkyl, or spiroheteroalkyl;

R⁵⁷ is R⁵⁸, R⁶⁰ or R⁶¹;

R⁵⁸ is phenyl;

R⁶⁰ is cycloalkyl, or heterocycloalkyl;

R⁶¹ is alkyl which is unsubstituted or substituted with one or two or three of independently selected R⁶², OR⁶², N(R⁶²)₂, C(O)OH, CN, F, Cl, Br or I;

R⁶² is R⁶⁵, or R⁶⁶;

R⁶⁵ is cycloalkyl, or heterocycloalkyl;

R⁶⁶ is alkyl which is unsubstituted or substituted with OR⁶⁷;

R⁶⁷ is alkyl;

wherein the cyclic moieties represented by R^{57A}, R⁵⁸, and R⁶⁰ are unsubstituted or substituted with one or two or three or four of independently selected R⁶⁸, F, Cl, Br or I;

R⁶⁸ is R⁷¹ or R⁷²;

R⁷¹ is heterocycloalkyl; and

R⁷² is alkyl, which is unsubstituted or substituted with one or two F.

[0134] Still another embodiment pertains to compounds having Formula (II), which are

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({4-[(3-morpholin-4-ylpropyl)amino]-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(1-tetrahydro-2H-pyran-4-ylpiperidin-4-yl)amino]phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(1-methylpiperidin-4-yl)amino]-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(4-methylpiperazin-1-yl)amino]-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

Trans-4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(2-methoxyethyl)amino]-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(3S)-tetrahydro-2H-pyran-3-ylmethyl]amino]phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(3R)-tetrahydro-2H-pyran-3-ylmethyl]amino]phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({2-naphthylsulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(2-methoxyethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl}benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({4-[(1,4-dioxan-2-ylmethyl)amino]-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(2,2,2-trifluoro-

thyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(3,3,3-trifluoropropyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2S)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[4-methoxycyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2R)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(1,4-dioxan-2-ylmethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-([4-(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[4-methoxycyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-(aminocarbonyl)-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-([5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-(trifluoromethyl)phenyl)sulfonyl)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)benzamide;
 35 Trans-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(1-methylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 5-([4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino)sulfonyl)-2-(tetrahydro-2H-pyran-4-ylmethoxy)nicotinamide;
 N-([5-bromo-6-[(1-methylpiperidin-4-yl)methoxy]pyridin-3-yl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1-methylpiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-cyano-6-(1,4-dioxan-2-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-bromo-6-(1,4-dioxan-2-ylmethoxy)pyridin-3-yl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2,2-dimethyltetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-5-cyano-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 N-({4-[(1-acetyl)piperidin-4-yl]amino]-3-nitrophenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({2-chloro-5-fluoro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(3-morpholin-4-ylpropyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-cyano-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-(2-morpholin-4-ylethoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(3-chloro-4-[[2-(2-methoxyethoxy)ethyl]sulfonyl]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[2-(2-methoxyethoxy)ethyl]sulfonyl]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)oxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-({5-bromo-6-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino}pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[[2-(cyanoethyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Cis-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-N-([4-({4-[bis(cyclopropylmethyl)amino]cyclohexyl)amino]-3-nitrophenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([1-methylpiperidin-4-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(morpholin-3-ylmethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 25 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-methylpiperazin-1-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylbut-2-ynyl)oxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-ethynyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-oxo-3,4-dihydroquinazolin-6-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 Cis-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-([4-([4-methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-([4-([4-methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 tert-butyl 3-[[4-([4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino)sulfonyl]-2-nitrophenoxymethyl]morpholine-4-carboxylate;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-(morpholin-3-ylmethoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([1-(methylsulfonyl)piperidin-4-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-([1,1-dioxidotetrahydro-2H-thiopyran-4-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 N-[(4-chloro-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-([1-(2,2,2-trifluoroethyl)piperidin-4-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{5-cyano-6-[[1-tetrahydro-2H-pyran-4-yl]piperidin-4-yl]oxy}pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{5-isopropyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 N-{{3-chloro-5-fluoro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

and therapeutically acceptable salts, and metabolites thereof.

10 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[[4-{{1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl}amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yl oxy)benzamide;
 N-{{5-chloro-6-[[4-fluorotetrahydro-2H-pyran-4-yl]methoxy]pyridin-3-yl)sulfonyl)-4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[[4-{{1-(2,2-difluoroethyl)piperidin-4-yl}amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{4-[[1-(1-cyclopropylpiperidin-4-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[[4-[[1-(1-morpholin-4-yl)cyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 Trans-4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[[4-[[4-(dicyclopropylamino)cyclohexyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{{2-(4-chlorophenyl)-6,6-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{{5-bromo-6-[[4-ethylmorpholin-3-yl]methoxy]pyridin-3-yl)sulfonyl)-4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{4-[[4-ethylmorpholin-3-yl]methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{3-nitro-4-[[4-(tetrahydro-2H-pyran-4-ylmorpholin-3-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[[3-nitro-4-[[[(3S)-1-tetrahydro-2H-pyran-4-yl]piperidin-3-yl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{5-cyano-6-[[tetrahydro-2H-pyran-4-ylmethyl]amino]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{4-[[1,1-dioxidothiomorpholin-4-yl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 N-[[4-[[[(4-aminotetrahydro-2H-pyran-4-yl)methyl]amino]-3-nitrophenyl)sulfonyl)-4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{5-cyano-6-[[4-morpholin-4-ylcyclohexyl]amino]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{3-cyano-4-[[tetrahydro-2H-pyran-4-ylmethyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[[4-[[[(1S,3R)-3-morpholin-4-ylcyclopentyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[[4-[[[(1R,3S)-3-morpholin-4-ylcyclopentyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{4-[[morpholin-2-ylmethyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{3-nitro-4-[[tetrahydrofuran-3-ylmethyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[[4-{{1-[cis-3-fluorotetrahydro-2H-pyran-4-yl]piperidin-4-yl}amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{3-nitro-4-[[1-tetrahydro-2H-pyran-4-ylazetidin-3-yl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{3-nitro-4-[[1-tetrahydrofuran-3-ylazetidin-3-yl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-[[3-nitro-4-[[[(3R)-1-tetrahydro-2H-pyran-4-yl]pyrrolidin-3-yl]methyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{{2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl)methyl}piperazin-1-yl)-N-{{4-[[4-fluorotetrahydro-2H-

pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-(4-((trans-4-hydroxycyclohexyl)methoxy)-3-nitrophenyl)sulfonyl)benzamide;
 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-(4-((cis-4-methoxycyclohexyl)methoxy)-3-nitrophenyl)sulfonyl)benzamide;
 5 Cis-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-(cyclopropylamino)cyclohexyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(4-(tetrahydro-2H-pyran-4-ylamino)cyclohexyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 Trans-N-([5-bromo-6-([4-(morpholin-4-yl)cyclohexyl]oxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methoxycyclohexyl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 tert-butyl 4-([4-([4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino)sulfonyl]-2-nitrophenoxymethyl]-4-fluoropiperidine-1-carboxylate;
 15 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-fluoropiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([4-(4-tetrahydro-2H-pyran-4-yl)piperazin-1-yl)cyclohexyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([[(3R)-1-tetrahydro-2H-pyran-4-yl]pyrrolidin-3-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(3R)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([[(3S)-1-tetrahydro-2H-pyran-4-yl]pyrrolidin-3-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(3S)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(4-methylmorpholin-2-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(4-(2-methoxyethyl)morpholin-2-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 N-([4-([[(4-acetylmorpholin-2-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([trans-4-(fluoromethyl)-1-oxetan-3-yl]pyrrolidin-3-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(4-fluorotetrahydro-2H-pyran-4-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([1-oxetan-3-yl]piperidin-4-yl)amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-cyclobutyl]piperidin-4-yl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(3S)-1-cyclopropyl]pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([1-tetrahydrofuran-3-yl]piperidin-4-yl)amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(3R)-1-cyclopropyl]pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([[(3S)-1-tetrahydro-2H-pyran-4-yl]pyrrolidin-3-yl)methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(3-hydroxy-2,2-dimethylpropyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-(methylsulfonyl)piperidin-

in-3-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(4-[[[1-acetyl]piperidin-3-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[3R)-1-(methylsulfonyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 4-(4-{[2-(4-chlorophenyl)-3,3-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[1-[2-fluoro-1-(fluoromethyl)ethyl]azetidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[1-(methylsulfonyl)pyrrolidin-3-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(4-[[[1-acetyl]pyrrolidin-3-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(4-[[[3R)-1-acetyl]pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[3-methoxy-2,2-dimethylpropyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[1R,3R)-3-hydroxycyclopentyl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[1S,3S)-3-hydroxycyclopentyl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[1S,3R)-3-hydroxycyclopentyl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[1R,3S)-3-hydroxycyclopentyl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(3-nitro-4-[[[3S)-2-oxopiperidin-3-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[1-[2-fluoro-1-(fluoromethyl)ethyl]azetidin-3-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(3-nitro-4-[[[1-oxetan-3-yl]azetidin-3-yl]methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(3-nitro-4-[[[1-oxetan-3-yl]piperidin-4-yl]methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[1-cyclopropyl]piperidin-4-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[2-fluoroethyl]morpholin-2-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[2,2-difluoroethyl]morpholin-2-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[4-fluoro-1-oxetan-3-yl]piperidin-4-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[2S)-4,4-difluoro-1-oxetan-3-yl]pyrrolidin-2-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(3-nitro-4-[[[4-tetrahydro-2H-pyran-4-yl]morpholin-3-yl]methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[4-cyclobutyl]morpholin-3-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(3-nitro-4-[[[4-tetrahydrofuran-3-yl]morpholin-3-yl]methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[1-cyclopropyl-4-fluoropiperidin-4-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[4-methoxybenzyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(3-nitro-4-[[[3-(trifluoromethoxy)benzyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[3-methoxybenzyl]amino]-

3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((difluoromethoxy)benzyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-(1,4-dioxaspiro[4.5]dec-8-ylamino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 Trans-N-[[4-((acetylamino)cyclohexyl)amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-(((3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-(((3S)-1-(2-fluoroethyl)pyrrolidin-3-yl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-(((3S)-1-(2,2-difluoroethyl)pyrrolidin-3-yl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-(((3R)-1-(2-fluoroethyl)pyrrolidin-3-yl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-(((3S)-1-oxetan-3-ylpyrrolidin-3-yl)methoxy)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((4-hydroxybenzyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((3-hydroxybenzyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((3-(difluoromethoxy)benzyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((cis-3-morpholin-4-ylcyclopentyl)methyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((4-(methylsulfonyl)amino)cyclohexyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((1-cyclopropylpiperidin-4-yl)amino)-3-((trifluoromethyl)sulfonyl)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-((1-oxetan-3-ylpiperidin-4-yl)methoxy)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((4-fluoro-1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((4-fluoro-1-tetrahydrofuran-3-yl)piperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((4-fluoro-1-(methylsulfonyl)piperidin-4-yl)methoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-(((3R)-1-oxetan-3-ylpyrrolidin-3-yl)methyl)amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((4-hydroxycyclohexyl)methoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((4-[3-(dimethylamino)propoxy]benzyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((4-(2-morpholin-4-ylethoxy)benzyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-(((E)-4-hydroxy-1-Adamantyl)methyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-(((Z)-4-hydroxy-1-Adamantyl)methyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 N-[[4-((1S,4S)-bicyclo[2.2.1]hept-5-en-2-ylmethoxy)-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((1-methyl-5-oxopyrrolidin-3-yl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-(((1R,4R,5R,6S)-5,6-dihydroxybicyclo[2.2.1]hept-2-yl)methoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-(((1R,4R,5S,6R)-5,6-dihydroxybicyclo[2.2.1]hept-2-yl)methoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-((3-oxocyclohexyl)methyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

clohexyl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohex-1,3-dien-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(3R)-1-[2-fluoro-1-(fluoromethyl)ethyl]pyrrolidin-3-yl}amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-({[(3S)-1-oxetan-3-yl]pyrrolidin-3-yl}methyl)amino}phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 Trans-N-({5-chloro-6-[(4-methoxycyclohexyl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-N-({5-chloro-6-[(4-methoxycyclohexyl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-({[(3S)-1-oxetan-3-yl]pyrrolidin-3-yl}amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[[2-(2-methoxyethoxy)ethyl]morpholin-2-yl]methyl}amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[(4-cyanomethyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[(N,N-dimethylglycyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 (2-[[[4-({4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino)methyl)morpholin-4-yl]acetic acid;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-({[(4-oxetan-3-yl)morpholin-2-yl]methyl}amino)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[(1-cyclopropyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-(methylsulfonyl)-6-(tetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[(4-methyltetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 ethyl 4-(4-{{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)piperazine-1-carboxylate;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[(4-morpholin-4-yl)piperidin-1-yl]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-({[(3R)-1-(oxetan-3-yl]pyrrolidin-3-yl}amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[(3R)-1-(1,3-difluoropropan-2-yl]pyrrolidin-3-yl}amino)-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[(1-isopropylpiperidin-4-yl]amino)-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{{4-[(1-tert-butylpiperidin-4-yl)amino]-3-nitrophenyl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[(1-(2-methoxyethyl)piperidin-3-yl]methyl}amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[(1-(cyanomethyl)piperidin-3-yl]methyl}amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[5-chloro-6-({[(3R)-1-(1,3-difluoropropan-2-yl]pyrrolidin-3-yl}amino)pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 tert-butyl 4-[[4-{{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino]piperazine-1-carboxylate;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{3-(pentafluoro- λ^6 -sulfanyl)-

4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-[(4-methoxytetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{(3-nitro-4-{[4-(oxetan-3-yl)piperazin-1-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{(3-nitro-4-{[4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{(3-nitro-4-{(3R)-tetrahydrofuran-3-ylamino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-[(4,4-difluorocyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{4-[(1-tert-butylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-[(4-(oxetan-3-yl)morpholin-2-yl)methyl]amino}-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{(5-chloro-6-{[(2S)-4-(4-fluorotetrahydro-2H-pyran-4-yl)methyl]amino}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{(5-chloro-6-{(1-cyclopropylpiperidin-4-yl)amino}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{(5-chloro-6-{[(2S)-4-(cyanomethyl)morpholin-2-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{(5-chloro-6-{[(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{(5-chloro-6-{[(2R)-4-(cyanomethyl)morpholin-2-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{(5-chloro-6-{[(2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{5-fluoro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{(5-chloro-6-[3-hydroxy-2-(hydroxymethyl)-2-methylpropoxy]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-[(1,3-difluoropropan-2-yl)morpholin-2-yl]methyl}amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{(5-chloro-6-{1-(cyanomethyl)piperidin-4-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-[(3R)-1-[2-(2-methoxyethoxy)ethyl]pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-[(3R)-1-(N,N-dimethylglycyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{(3-nitro-4-{[1-(oxetan-3-yl)azetididin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{(5-chloro-6-{1-(cyanomethyl)-4-fluoropiperidin-4-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-[(2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl}amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-[(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl}amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{(5-chloro-6-{1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{(5-chloro-6-{[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]oxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-[(3R)-1-(cyanomethyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-methoxypiperidin-1-yl)-N-{5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-


yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-methoxypiperidin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-({5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-({5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[[1-(N,N-dimethylglycyl)-4-fluoropiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-fluoropiperidin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-fluoropiperidin-1-yl)-N-({5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-(trifluoromethyl)phenyl)sulfonyl)benzamide;
 N-({5-chloro-6-[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[2-(tetrahydrofuran-3-yloxy)ethoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[(trans-4-cyanocyclohexyl)methyl]amino-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({5-cyano-6-[2-(tetrahydro-2H-pyran-4-yl)ethoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-(3-furylmethoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({3-chloro-4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]phenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-cyano-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[[1-(2,2-difluoroethyl)-4-fluoropiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({3-chloro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(4,4-difluorocyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({6-[[1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl)methoxy]-5-(trifluoromethyl)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({5-chloro-6-[2-(tetrahydrofuran-2-yl)ethoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]-3-methylpiperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[(3-(cyclopropylamino)propyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- N-[[5-chloro-6-(2-methoxyethoxy)pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-fluoro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 N-[(3-chloro-4-[[1-(methoxyacetyl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(3-chloro-4-[[1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohexyl]methyl]piperidin-1-yl)-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-[[6-(tetrahydro-2H-pyran-4-ylmethoxy)-5-(trifluoromethyl)pyridin-3-yl]sulfonyl]benzamide;
 N-[[5-chloro-6-[(trans-4-hydroxycyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-cyano-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[6-[(trans-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[6-[(cis-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 N-[[5-chloro-6-[(4,4-difluoro-1-hydroxycyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperidin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(3-chloro-4-[[trans-4-(morpholin-4-yl)cyclohexyl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-2-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(3-cyclopropyl(1,3-thiazol-5-ylmethyl)amino)propyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(3-chloro-4-[(trans-4-hydroxycyclohexyl)methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-chloro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-(trifluoromethyl)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(3-cyclopropyl(2,2,2-trifluoroethyl)amino)propyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 N-[(3-chloro-4-[[1-(oxetan-3-yl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3,5-difluoro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(3-cyclopropyl(oxetan-3-yl)amino)propyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 N-[(3-chloro-4-[[1-(1-methyl-L-prolyl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3,4-difluoro-5-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 N-[[5-chloro-6-[(2S)-4-cyclopropylmorpholin-2-yl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperidin-1-yl)-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperidin-1-yl)-N-[(3-chloro-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 methyl 2-[[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino]methyl]morpholine-4-carboxylate;
 2-[[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(4-(methylsulfonyl)morpholin-2-yl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(3-cyclobutyl(cyclopropyl)amino)propyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-((3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(3-chloro-4-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-chloro-4-(tetrahydrofuran-3-ylmethoxy)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[trans-4-hydroxycyclohexyl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(3-chloro-4-[[4-fluorotetrahydro-2H-pyran-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[9-(4-chlorophenyl)-3-(oxetan-3-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[(2R)-4-cyclopropylmorpholin-2-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[(2S)-4-cyclopropylmorpholin-2-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(4-cyclopropyl(oxetan-3-yl)amino)cyclohexyl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-N-[(4-[[[(4-cyclopropylmorpholin-2-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 N-[(3-chloro-4-[(4-cyclopropylmorpholin-2-yl)methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(3-chloro-4-[[[(4-cyclopropylmorpholin-2-yl)methyl]amino]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 2-[[[(3-chloro-4-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]phenyl)amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
 (2S)-2-[[[(3-chloro-5-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]pyridin-2-yl)oxy]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
 N-[(5-chloro-6-[(4-cyclopropylmorpholin-2-yl)methyl]amino)pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 2-[[[(3-chloro-5-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]pyridin-2-yl)amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[trans-4-hydroxy-4-methylcyclohexyl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[cis-4-hydroxy-4-methylcyclohexyl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[[(1R,2R,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[(2-cyanoethyl)(cyclopropyl)amino]cyclohexyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(5-chloro-6-(5,6,7,8-tetrahydroimidazo[1,2-a]pyridin-6-ylmethoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[[(1R,2S,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[cis-4-hydroxy-4-methylcyclohexyl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[4-(3,3-difluoropyrrolidin-1-yl)cyclohexyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 N-[(5-chloro-6-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(cis-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-

dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[4-[[2-(2,2-difluorocyclopropyl)amino]cyclohexyl]amino]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(cis-1-fluoro-4-hydroxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-
 5 dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[3-nitro-4-(2-oxaspiro[3.5]non-
 7-ylmethoxy)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[(trans-4-hydroxy-4-methyl-
 cyclohexyl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-5,5-bis(fluoromethyl)cyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-
 2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({4-[(4-cyclopropylmorpholin-2-
 yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(trans-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophe-
 15 nyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(cis-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophe-
 nyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[3-cyano-4-[[4-fluoro-1-(oxe-
 tan-3-yl)piperidin-4-yl]methoxy]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[4-[[[(trans-4-ethyl-4-hydroxycy-
 clohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[4-[[[(cis-4-ethyl-4-hydroxycy-
 clohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5-(methoxymethyl)-5-methylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({3-nitro-4-[(tet-
 25 rahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[3-nitro-4-[[[(2S)-4-(oxetan-3-
 yl)morpholin-2-yl]methyl]amino]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({3-chloro-4-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]phenyl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-
 dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[4-[[2-(2-cyanoethyl)(cyclopro-
 pyl)amino]-1-fluorocyclohexyl]methoxy]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({5-nitro-6-[(tetrahydro-2H-
 pyran-4-ylmethyl)amino]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({3-nitro-4-[(2-oxaspiro[3.5]non-
 35 7-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[4-[[[(4-cyano-4-methylcy-
 clohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-
 yloxy)-N-[[3-(tetrahydro-2H-pyran-4-ylmethyl)-3H-1,2,3]triazolo[4,5-b]pyridin-6-yl]sulfonyl]benzamide;
 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[6-[[[(trans-4-hydroxy-4-methyl-
 cyclohexyl)methyl]amino]-5-nitropyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[5-cyano-6-[[4-fluoro-1-(oxe-
 tan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-4-[[4-4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-
 45 5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)morpholine-4-carboxamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[4-[[[(4-(methoxymethyl)cy-
 clohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[5-chloro-6-[[1-(1,3-thiazol-2-
 yl)piperidin-4-yl]methoxy]pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[6-[[[(cis-4-hydroxy-4-methylcy-
 clohexyl)methyl]amino]-5-nitropyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-[[4-[[[(trans-4-hydroxy-4-
 methylcyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-
 2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; and therapeutically
 55 acceptable salts, and metabolites thereof.

[0135] In another aspect, the present invention provides compounds of Formula (III)



(III);

and therapeutically acceptable salts, and metabolites thereof, wherein A¹, B¹, D¹, E¹, and Y¹ are as described herein for Formula (I); R¹⁰⁰ is as described for substituents on R²⁶; n is 0, 1, 2, or 3; R¹⁰¹ is as described for substituents on R³⁰; p is 0, 1, 2, 3, 4, 5, or 6; R¹⁰² is as described for substituents on R⁴⁰; q is 0, 1, 2, 3, 4, 5, or 6; R¹⁰³ is as described for substituents on R⁵⁸; and r is 0, 1, 2, 3, or 4.

[0136] In one embodiment of Formula (III), n, p, r, and q are each 0.

[0137] In one embodiment of Formula (III), A¹ is N. In another embodiment of Formula (III), A¹ is C(A²). In another embodiment of Formula (III), A¹ is C(A²); and A² is H, F, Cl, Br, or I. In another embodiment of Formula (III), A¹ is C(A²); and A² is H.

[0138] In one embodiment of Formula (III), B¹ is R¹, OR¹, NHR¹, NHC(O)R¹, F, Cl, Br, or I. In another embodiment of Formula (III), A¹ is C(A²); A² is H; and B¹ is NHR¹. In another embodiment of Formula (III), A¹ is C(A²); A² is H; and B¹ is OR¹. In another embodiment of Formula (III), A¹ is C(A²); A² is H; and B¹ is Cl. In another embodiment of Formula (III), A¹ is C(A²); A² is H; and B¹ is R¹. In another embodiment of Formula (III), A¹ is N; and B¹ is NHR¹. In another embodiment of Formula (III), A¹ is N; and B¹ is OR¹. In another embodiment of Formula (III), A¹ is N; and B¹ is Cl. In another embodiment of Formula (III), A¹ is N; and B¹ is R¹.

[0139] In one embodiment of Formula (III), D¹ is H or Cl. In another embodiment of Formula (III), A¹ is C(A²); A² is H; B¹ is NHR¹; and D¹ is H. In another embodiment of Formula (III), A¹ is C(A²); A² is H; B¹ is NHR¹; and D¹ is Cl. In another embodiment of Formula (III), A¹ is C(A²); A² is H; B¹ is OR¹; and D¹ is H. In another embodiment of Formula (III), A¹ is C(A²); A² is H; B¹ is Cl; and D¹ is H. In another embodiment of Formula (III), A¹ is C(A²); A² is H; B¹ is R¹; and D¹ is H. In another embodiment of Formula (III), A¹ is N; B¹ is NHR¹; and D¹ is H. In another embodiment of Formula (III), A¹ is N; B¹ is NHR¹; and D¹ is Cl. In another embodiment of Formula (III), A¹ is N; B¹ is OR¹; and D¹ is H. In another embodiment of Formula (III), A¹ is N; B¹ is Cl; and D¹ is H. In another embodiment of Formula (III), A¹ is N; B¹ is R¹; and D¹ is H.

[0140] In one embodiment of Formula (III), E¹ is H. In another embodiment of Formula (III), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (III), A¹ is C(A²); A² is H; B¹ is NHR¹; E¹ is H; and D¹ is Cl. In another embodiment of Formula (III), A¹ is C(A²); A² is H; B¹ is OR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (III), A¹ is C(A²); A² is H; B¹ is Cl; D¹ is H; and E¹ is H. In another embodiment of Formula (III), A¹ is C(A²); A² is H; B¹ is R¹; D¹ is H; and E¹ is H. In another embodiment of Formula (III), A¹ is N; B¹ is NHR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (III), A¹ is N; B¹ is NHR¹; E¹ is H; and D¹ is Cl. In another embodiment of Formula (III), A¹ is N; B¹ is OR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (III), A¹ is N; B¹ is Cl; D¹ is H; and E¹ is H. In another embodiment of Formula (III), A¹ is N; B¹ is R¹; D¹ is H; and E¹ is H.

[0141] In one embodiment of Formula (III), Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SO₂R¹⁷, or C(O)NH₂. In another embodiment of Formula (III), Y¹ is H. In another embodiment of Formula (III), Y¹ is CN. In another embodiment of Formula (III), Y¹ is F, Cl, Br, or I. In another embodiment of Formula (III), Y¹ is CF₃. In another embodiment of Formula (III), Y¹ is SR¹⁷. In another embodiment of Formula (III), Y¹ is OR¹⁷. In another embodiment of Formula (III), Y¹ is NO₂. In another embodiment of Formula (III), Y¹ is SO₂R¹⁷; wherein R¹⁷ is as defined herein. In another embodiment of

Formula (III), Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl. In another embodiment of Formula (III), Y¹ is R¹⁷; wherein R¹⁷ is alkynyl. In another embodiment of Formula (III), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂ or SO₂R¹⁷; wherein R¹⁷ is alkyl or alkynyl. In another embodiment of Formula (III), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂. In another embodiment of Formula (III), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl substituted with three F. In another embodiment of Formula (III), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂ or SO₂R¹⁷; wherein R¹⁷ is alkyl or alkynyl. In another embodiment of Formula (III), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂. In another embodiment of Formula (III), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl substituted with three F.

[0142] In one embodiment of Formula (III), Y¹ and B¹, together with the atoms to which they are attached, are benzene. In another embodiment of Formula (III), Y¹ and B¹, together with the atoms to which they are attached, are heteroarene. In another embodiment of Formula (III), A¹ is C(A²); A², G¹, E¹, and D¹ are independently selected H; and Y¹ and B¹, together with the atoms to which they are attached, are benzene. In another embodiment of Formula (III), A¹ is C(A²); A², G¹, E¹, and D¹ are independently selected H; and Y¹ and B¹, together with the atoms to which they are attached, are heteroarene.

[0143] In one embodiment of Formula (III), R¹ is R⁴ or R⁵. In one embodiment of Formula (III), R¹ is R⁴. In one embodiment of Formula (III), R¹ is R⁵. In one embodiment of Formula (III), R¹ is R⁴; and R⁴ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl. In one embodiment of Formula (III), R¹ is R⁴; and R⁴ is cycloalkyl or heterocycloalkyl. In one embodiment of Formula (III), R¹ is R⁴; and R⁴ is cycloalkyl. In one embodiment of Formula (III), R¹ is R⁴; and R⁴ is heterocycloalkyl.

[0144] In one embodiment of Formula (III), R¹ is R⁴; and R⁴ is cycloalkyl; wherein R⁴ is unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted as defined herein. In another embodiment of Formula (III), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted with R⁵⁷ or N(R⁵⁷)₂. In another embodiment of Formula (III), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; and R⁵⁷ is R⁶⁰. In another embodiment of Formula (III), R¹ is R⁴; R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; and R⁶⁰ is heterocycloalkyl. In another embodiment of Formula (III), R¹ is R⁴; R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; R⁶⁰ is heterocycloalkyl; wherein the heterocycloalkyl ring is morpholinyl. In another embodiment of Formula (III), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted with N(R⁵⁷)₂. In another embodiment of Formula (III), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with N(R⁵⁷)₂. In another embodiment of Formula (III), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with N(R⁵⁷)₂; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is unsubstituted or substituted with R⁶². In another embodiment of Formula (III), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is unsubstituted or substituted with R⁶²; R⁶² is R⁶⁵; and R⁶⁵ is cycloalkyl. In another embodiment of Formula (III), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is substituted with R⁶²; R⁶² is R⁶⁵; and R⁶⁵ is cyclopropyl.

[0145] In one embodiment of Formula (III), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein R⁴ is unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is substituted as defined herein. In another embodiment of Formula (III), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is substituted with R⁵⁷. In another embodiment of Formula (III), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the heterocycloalkyl ring is substituted with R⁵⁷; and R⁵⁷ is R⁶⁰ or R⁶¹. In another embodiment of Formula (III), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰ or R⁶¹; R⁶⁰ is heterocycloalkyl; and R⁶¹ is alkyl. In another embodiment of Formula (III), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; R⁶⁰ is heterocycloalkyl; wherein the heterocycloalkyl is tetrahydropyranyl. In another embodiment of Formula (III), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; R⁶¹ is alkyl; and the alkyl is methyl. In another embodiment of Formula (III), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with C(O)OR⁵⁷; R⁵⁷ is R⁶¹; R⁶¹ is alkyl; and the alkyl is methyl.

[0146] In one embodiment of Formula (III), R¹ is R⁵; and R⁵ is alkyl which is unsubstituted or substituted. In one embodiment of Formula (III), R¹ is R⁵; and R⁵ is alkyl which is unsubstituted or substituted with R⁷, OR⁷, OH, CN, or F. In another embodiment of Formula (III), R¹ is R⁵; and R⁵ is alkyl which is substituted with R⁷, OR⁷, NHR⁷, or N(R⁷)₂.

[0147] In one embodiment of Formula (III), R⁷ is R⁸, R⁹, R¹⁰ or R¹¹ which are unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R⁷ is R⁸ which is unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R⁷ is R⁹ which is unsubstituted or substituted as defined herein. In another em-

bodiment of Formula (III), R⁷ is R¹⁰ which is unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R⁷ is R¹¹ which is unsubstituted or substituted as defined herein.

[0148] In one embodiment of Formula (III), R⁸ is phenyl which is unsubstituted or substituted as defined herein.

[0149] In one embodiment of Formula (III), R⁹ is heteroaryl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R⁹ is furanyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R⁹ is furanyl; which is unsubstituted.

[0150] In one embodiment of Formula (III), R¹⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, heterocycloalkyl which are unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R¹⁰ is heterocycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R¹⁰ is tetrahydrofuranyl, tetrahydropyranyl, morpholinyl, dioxanyl, oxetanyl, piperdiny, or pyrrolidinyl, which are unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R¹⁰ is tetrahydropyranyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R¹⁰ is morpholinyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R¹⁰ is cycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R¹⁰ is cycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R¹⁰ is cyclohexyl, cyclopropyl, cyclobutyl, or bicyclo[2.2.1]heptanyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R¹⁰ is heterocycloalkyl which is fused with R^{10A}; and R^{10A} is heteroarene. In another embodiment of Formula (III), R¹⁰ is 5,6,7,8-tetrahydroimidazo[1,2-a]pyridinyl.

[0151] In one embodiment of Formula (III), R¹¹ is alkyl, alkenyl or alkynyl which are unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R¹¹ is alkyl. In another embodiment of Formula (III), R¹¹ is methyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (III), R¹¹ is alkyl; which is substituted as defined herein. In another embodiment of Formula (III), R¹¹ is alkyl; which is substituted with R¹², OR¹², or CF₃. In another embodiment of Formula (III), R¹¹ is alkyl; which is substituted with OR¹²; R¹² is R¹⁶; and R¹⁶ is alkyl. In another embodiment of Formula (III), R¹¹ is alkyl; which is substituted with CF₃. In another embodiment of Formula (III), R¹¹ is alkyl; which is substituted with R¹²; R¹² is R¹⁴; and R¹⁴ is heteroaryl.

[0152] In one embodiment of Formula (III),

n, p, r, and q are each 0;
 A¹ is N or C(A²);
 A² is H, F, Br, I, or Cl;
 B¹ is R¹, OR¹, NHR¹, NHC(O)R¹, F, Br, I or Cl;
 D¹ is H, F, Br, I, or Cl;
 E¹ is H; and
 Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SO₂R¹⁷, or C(O)NH₂; or
 Y¹ and B¹, together with the atoms to which they are attached, are benzene, heteroarene, or heterocycloalkene; and
 A², D¹, and E¹ are independently selected H;
 R¹ is R⁴ or R⁵;
 R⁴ is cycloalkyl, or heterocycloalkyl;
 R⁵ is alkyl, or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected
 R⁷, OR⁷, NHR⁷, N(R⁷)₂, CN, OH, F, Cl, Br or I;
 R⁷ is R⁸, R⁹, R¹⁰, or R¹¹;
 R⁸ is phenyl;
 R⁹ is heteroaryl;
 R¹⁰ is cycloalkyl, cycloalkenyl, or heterocycloalkyl; each of which is unfused or fused with R^{10A} R^{10A} is heteroarene;
 R¹¹ is alkyl each of which is unsubstituted or substituted with one or two or three of independently selected R¹²,
 OR¹² or CF₃;
 R¹² is R¹⁴ or R¹⁶;
 R¹⁴ is heteroaryl;
 R¹⁶ is alkyl;
 R¹⁷ is R²¹;
 R²¹ is alkyl, or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently
 selected R²², F, Cl, Br or I;
 R²² is R²⁵;
 R²⁵ is heterocycloalkyl;
 wherein the cyclic moieties represented by Y¹ and B¹ together, R⁴, R⁸, R¹⁰, and R²⁵, are independently unsubstituted,
 further unsubstituted, substituted or further substituted with one or two or three or four or five of independently
 selected R^{57A}, R⁵⁷, OR⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, C(O)N(R⁵⁷)₂, NH₂, NHR⁵⁷, N(R⁵⁷)₂, NHC(O)R⁵⁷,
 NHS(O)₂R⁵⁷, OH, CN, (O), F, Cl, Br or I;

R^{57A} is spiroalkyl, or spiroheteroalkyl;

R⁵⁷ is R⁵⁸, R⁶⁰ or R⁶¹;

R⁵⁸ is phenyl;

R⁶⁰ is cycloalkyl, or heterocycloalkyl;

5 R⁶¹ is alkyl which is unsubstituted or substituted with one or two or three of independently selected R⁶², OR⁶², N(R⁶²)₂, C(O)OH, CN, F, Cl, Br or I;

R⁶² is R⁶⁵, or R⁶⁶;

R⁶⁵ is cycloalkyl, or heterocycloalkyl;

R⁶⁶ is alkyl which is unsubstituted or substituted with OR⁶⁷;

10 R⁶⁷ is alkyl;

wherein the cyclic moieties represented by R^{57A}, R⁵⁸, and R⁶⁰ are unsubstituted or substituted with one or two or three or four of independently selected R⁶⁸, F, Cl, Br or I;

R⁶⁸ is R⁷¹ or R⁷²;

R⁷¹ is heterocycloalkyl; and

15 R⁷² is alkyl, which is unsubstituted or substituted with one or two F.

[0153] Still another embodiment pertains to compounds having Formula (III), which are

20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1-methylpiperidin-4-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methylpiperazin-1-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2-methoxyethyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(3S)-tetrahydro-2H-pyran-3-ylmethyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(3R)-tetrahydro-2H-pyran-3-ylmethyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-(2-naphthylsulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2-methoxyethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1,4-dioxan-2-ylmethyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(2,2,2-trifluoroethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(3,3,3-trifluoropropyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2S)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(4-methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2R)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(4-methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[[4-fluorotetrahydro-2H-pyran-4-yl]methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 N-[[3-(aminocarbonyl)-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[[4-morpholin-4-ylcyclohexyl]amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 5-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino]sulfonyl)-2-(tetrahydro-2H-pyran-4-ylmethoxy)nicotinamide;
- N-{{5-bromo-6-[[1-methylpiperidin-4-yl]methoxy]pyridin-3-yl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[[1-methylpiperidin-4-yl]methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-(1,4-dioxan-2-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[5-bromo-6-(1,4-dioxan-2-ylmethoxy)pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[[2,2-dimethyltetrahydro-2H-pyran-4-yl]methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-{{3-chloro-5-cyano-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 25 N-[[4-[[1-acetyl]piperidin-4-yl]amino]-3-nitrophenyl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[2-chloro-5-fluoro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[[3-morpholin-4-ylpropyl]amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{5-cyano-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-(2-morpholin-4-ylethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 N-[[3-chloro-4-[[2-(2-methoxyethoxy)ethyl]sulfonyl]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[2-(2-methoxyethoxy)ethyl]sulfonyl]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[[4-morpholin-4-ylcyclohexyl]oxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 N-{{5-bromo-6-[[1-tetrahydro-2H-pyran-4-yl]piperidin-4-yl]amino]pyridin-3-yl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[[2-cyanoethyl]amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 N-[[4-{{4-[[bis(cyclopropylmethyl)amino]cyclohexyl]amino]-3-nitrophenyl}sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[1-methylpiperidin-4-yl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[[4-morpholin-3-ylmethyl]amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-{{4-[[4-morpholin-4-ylbut-2-ynyl]oxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-ethynyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-oxo-3,4-dihydroquinazolin-6-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- tert-butyl 3-[[4-{{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino]sulfonyl]-2-nitrophenoxymethyl]morpholine-4-carboxylate;

- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-(morpholin-3-ylmethoxy)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-(methylsulfonyl)piperidin-4-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1,1-dioxidotetrahydro-2H-thiopyran-4-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[4-(chloro-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[1-(2,2,2-trifluoroethyl)piperidin-4-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(5-cyano-6-[[1-tetrahydro-2H-pyran-4-yl]piperidin-4-yl]oxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-isopropyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 N-[(3-chloro-5-fluoro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(5-chloro-6-[[4-fluorotetrahydro-2H-pyran-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-(2,2-difluoroethyl)piperidin-4-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-(1-cyclopropyl)piperidin-4-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-(morpholin-4-yl)cyclohexyl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(dicyclopropylamino)cyclohexyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(5-bromo-6-[[4-ethylmorpholin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-ethylmorpholin-3-yl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[4-tetrahydro-2H-pyran-4-yl]morpholin-3-yl]methoxy)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[3-(3S)-1-tetrahydro-2H-pyran-4-yl]piperidin-3-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(5-cyano-6-[(tetrahydro-2H-pyran-4-ylmethyl)amino]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1,1-dioxidothiomorpholin-4-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 N-[[4-[[[4-aminotetrahydro-2H-pyran-4-yl]methyl]amino]-3-nitrophenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(5-cyano-6-[[4-morpholin-4-yl]cyclohexyl]amino)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-cyano-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[1S,3R]-3-morpholin-4-yl]cyclopentyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[1R,3S]-3-morpholin-4-yl]cyclopentyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[morpholin-2-yl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(tetrahydrofuran-3-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[1-cis-3-fluorotetrahydro-2H-pyran-4-yl]piperidin-4-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[1-tetrahydro-2H-pyran-4-yl]azetidin-3-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(1-tetrahydrofuran-3-ylazetidin-3-yl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-(((3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl)methyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-(4-((trans-4-hydroxycyclohexyl)methoxy)-3-nitrophenyl)sulfonyl)benzamide;
 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-(4-((cis-4-methoxycyclohexyl)methoxy)-3-nitrophenyl)sulfonyl)benzamide;
 10 Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-((4-(cyclopropylamino)cyclohexyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-((4-(tetrahydro-2H-pyran-4-ylamino)cyclohexyl)amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-N-({5-bromo-6-[(4-morpholin-4-ylcyclohexyl)oxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methoxycyclohexyl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 tert-butyl 4-[4-({4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino)sulfonyl)-2-nitrophenoxymethyl]-4-fluoropiperidine-1-carboxylate;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluoropiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-((4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl)cyclohexyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl)methoxy}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-(((3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl)amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-(((3R)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-(((3S)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl)amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-(((3S)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-((4-methylmorpholin-2-yl)methyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-((4-(2-methoxyethyl)morpholin-2-yl)methyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([4-(((4-acetylmorpholin-2-yl)methyl)amino)-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({trans-4-(fluoromethyl)-1-oxetan-3-ylpyrrolidin-3-yl)methoxy}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-(((4-fluorotetrahydro-2H-pyran-4-yl)methyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(1-oxetan-3-yl)piperidin-4-yl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1-cyclobutyl)piperidin-4-yl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-(((3S)-1-cyclopropylpyrrolidin-3-yl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(1-tetrahydrofuran-3-yl)piperidin-4-yl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-(((3R)-1-cyclopropylpyrrolidin-3-yl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-(((3S)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl)methyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[3-hydroxy-2,2-dimethylpropyl]amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[1-(methylsulfonyl)piperidin-3-yl]methyl}amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 N-{4-{[1-(1-acetyl)piperidin-3-yl]methyl}amino}-3-nitrophenyl)sulfonyl}-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[3-(3R)-1-(methylsulfonyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[1-[2-fluoro-1-(fluoromethyl)ethyl]azetidin-3-yl]amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[1-(methylsulfonyl)pyrrolidin-3-yl]methyl}amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-{4-{[1-(1-acetylpyrrolidin-3-yl)ethyl]amino}-3-nitrophenyl)sulfonyl}-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 N-{4-{[3-(3R)-1-acetylpyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl}-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[3-(3-methoxy-2,2-dimethylpropyl)amino]-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[1-(1R,3R)-3-hydroxycyclopentyl]methyl}amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[1-(1S,3S)-3-hydroxycyclopentyl]methyl}amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[1-(1S,3R)-3-hydroxycyclopentyl]methyl}amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[1-(1R,3S)-3-hydroxycyclopentyl]methyl}amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[3-(3S)-2-oxopiperidin-3-yl]amino}phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[1-[2-fluoro-1-(fluoromethyl)ethyl]azetidin-3-yl]methyl}amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[3-nitro-4-{[1-(1-oxetan-3-yl)azetidin-3-yl]methyl}amino]phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[3-nitro-4-{[1-(1-oxetan-3-yl)piperidin-4-yl]methyl}amino]phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[1-(1-cyclopropyl)piperidin-4-yl]methyl}amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[4-(2-fluoroethyl)morpholin-2-yl]methyl}amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[4-(2,2-difluoroethyl)morpholin-2-yl]methyl}amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[4-fluoro-1-oxetan-3-yl]piperidin-4-yl]methoxy}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[4-{[2-(2S)-4,4-difluoro-1-oxetan-3-yl]pyrrolidin-2-yl]methoxy}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[3-nitro-4-{[4-tetrahydro-2H-pyran-4-yl]morpholin-3-yl]methyl}amino]phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[4-(4-cyclobutylmorpholin-3-yl]methyl}amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[3-nitro-4-{[4-tetrahydrofuran-3-yl]morpholin-3-yl]methyl}amino]phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl]methyl}amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[1-(1-cyclopropyl-4-fluoropiperidin-4-yl]methoxy}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[4-(2-methoxybenzyl)amino]-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{4-{[3-(trifluoromethoxy)benzyl]amino}phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(3-methoxybenzyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[4-(difluoromethoxy)benzyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{1,4-dioxaspiro[4.5]dec-8-ylamino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-N-[(4-{[4-(acetylamino)cyclohexyl]amino}-3-nitrophenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3S)-1-(2-fluoroethyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3S)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(2-fluoroethyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(3S)-1-oxetan-3-ylpyrrolidin-3-yl]methoxy}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-hydroxybenzyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 20 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(3-hydroxybenzyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3-difluoromethoxy)benzyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 25 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(cis-3-morpholin-4-ylcyclopentyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(methylsulfonyl)amino]cyclohexyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1-cyclopropylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 30 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(1-oxetan-3-ylpiperidin-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-fluoro-1-tetrahydro-2H-pyran-4-ylpiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-fluoro-1-tetrahydrofuran-3-ylpiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-fluoro-1-(methylsulfonyl)piperidin-4-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(3R)-1-oxetan-3-ylpyrrolidin-3-yl]methyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 Trans-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-hydroxycyclohexyl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[4-(3-(dimethylamino)propoxy]benzyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[4-(2-morpholin-4-ylethoxy)benzyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(E)-4-hydroxy-1-Adamantyl]methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(Z)-4-hydroxy-1-Adamantyl]methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 N-({4-[(1S,4S)-bicyclo[2.2.1]hept-5-en-2-ylmethoxy]-3-nitrophenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1-methyl-5-oxopyrrolidin-3-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(1R,4R,5R,6S)-5,6-dihydroxybicyclo[2.2.1]hept-2-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(1R,4R,5S,6R)-5,6-dihydroxybicyclo[2.2.1]hept-2-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(3-oxocyclohexyl)methoxy]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(3R)-1-[2-fluoro-1-(fluoromethyl)ethyl]pyrrolidin-3-yl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-({(3S)-1-oxetan-3-ylpyrrolidin-3-yl}methyl)amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-N-({5-chloro-6-[(4-methoxycyclohexyl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 Cis-N-({5-chloro-6-[(4-methoxycyclohexyl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-({(3S)-1-oxetan-3-ylpyrrolidin-3-yl}amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-[2-(2-methoxyethoxy)ethyl]morpholin-2-yl)methyl]amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(4-cyanomethyl)morpholin-2-yl)methyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(N,N-dimethylglycyl)morpholin-2-yl)methyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 (2-[[4-({4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)amino)methyl]morpholin-4-yl)acetic acid;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-({(4-oxetan-3-yl)morpholin-2-yl)methyl]amino)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(4-cyclopropylmorpholin-2-yl)methyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-(methylsulfonyl)-6-(tetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methyltetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 ethyl 4-(4-[[4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)piperazine-1-carboxylate;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[4-(morpholin-4-yl)piperidin-1-yl]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-({(3R)-1-(oxetan-3-yl)pyrrolidin-3-yl}amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl}amino)-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1-isopropylpiperidin-4-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 N-({4-[(1-tert-butylpiperidin-4-yl)amino]-3-nitrophenyl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(1-(2-methoxyethyl)piperidin-3-yl)methyl]amino)-3-nitrophenyl}sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({(1-(cyanomethyl)piperidin-3-yl)methyl]amino)-3-nitrophenyl}sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-({(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl}amino)pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 tert-butyl 4-([4-([4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)amino]piperazine-1-carboxylate;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-(pentafluoro- λ^6 -sulfanyl)-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino)phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methoxytetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl}oxy)-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(4-oxetan-3-yl)piperazin-1-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(4-tetrahydro-2H-pyran-4-yl)piperazin-1-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(3R)-tetrahydrofuran-3-ylamino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({(4,4-difluorocyclohexyl)methyl}amino)-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({4-[(1-tert-butylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-oxetan-3-yl)morpholin-2-yl]methyl}amino)-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methyl]amino)pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(1-cyclopropylpiperidin-4-yl)amino]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 N-[(5-chloro-6-[(2S)-4-(cyanomethyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 N-[(5-chloro-6-[(2R)-4-(cyanomethyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-fluoro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[3-hydroxy-2-(hydroxymethyl)-2-methylpropoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({(1,3-difluoropropan-2-yl)morpholin-2-yl]methyl}amino)-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(1-(cyanomethyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({(3R)-1-(2-methoxyethoxy)ethyl}pyrrolidin-3-yl)amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({(3R)-1-(N,N-dimethylglycyl)pyrrolidin-3-yl}amino)-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(4-oxetan-3-yl)azetididin-3-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 N-[(5-chloro-6-[(1-(cyanomethyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({(2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 N-[(5-chloro-6-[(1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-({(3R)-1-(cyanomethyl)pyrrolidin-3-yl}amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;


N-[(5-chloro-6-[(1-(N,N-dimethylglycyl)-4-fluoropiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[2-(tetrahydrofuran-3-yloxy)ethoxy]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(trans-4-cyanocyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-cyano-6-[2-(tetrahydro-2H-pyran-4-yl)ethoxy]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-(3-furylmethoxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 N-[(5-chloro-6-[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 N-({3-chloro-4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]phenyl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-cyano-4-(tetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 N-[(5-chloro-6-[(1-(2,2-difluoroethyl)-4-fluoropiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({3-chloro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(4,4-difluorocyclohexyl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([6-[(1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl)methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-chloro-6-[2-(tetrahydrofuran-2-yl)ethoxy]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[(3-(cyclopropylamino)propyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-(2-methoxyethoxy)pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-fluoro-6-(tetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(3-chloro-4-[[1-(methoxyacetyl)piperidin-4-yl)methoxy]phenyl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(3-chloro-4-[(1-(N,N-dimethylglycyl)piperidin-4-yl)methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([6-(tetrahydro-2H-pyran-4-yl)methoxy]-5-(trifluoromethyl)pyridin-3-yl)sulfonyl)benzamide;
 N-({-chloro-6-[(trans-4-hydroxycyclohexyl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-cyano-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([6-[(trans-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([6-[(cis-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 N-[(3-chloro-4-[(trans-4-(morpholin-4-yl)cyclohexyl)methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[(3-[cyclopropyl(1,3-thiazol-

5-ylmethyl)amino]propyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({3-chloro-4-[(trans-4-hydroxycyclohexyl)methoxy]phenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-chloro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-(trifluoromethyl)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-([4-({3-[cyclopropyl(2,2,2-trifluoroethyl)amino]propyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 N-([3-chloro-4-{{1-(oxetan-3-yl)piperidin-4-yl}methoxy}phenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3,5-difluoro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-([4-({3-[cyclopropyl(oxetan-3-yl)amino]propyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 N-([3-chloro-4-{{1-(1-methyl-L-prolyl)piperidin-4-yl}methoxy}phenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3,4-difluoro-5-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 N-([5-chloro-6-{{(2S)-4-cyclopropylmorpholin-2-yl}methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 methyl 2-[[[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino]methyl]morpholine-4-carboxylate;
 2-[[[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
 25 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-([4-{{(4-methylsulfonyl)morpholin-2-yl}methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-([4-{{3-[cyclobutyl(cyclopropyl)amino]propyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 N-([3-chloro-4-{{[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy}phenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-([3-chloro-4-(tetrahydrofuran-3-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-([4-{{[(2R)-4-cyclopropylmorpholin-2-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-([4-{{[(2S)-4-cyclopropylmorpholin-2-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 N-([5-chloro-6-{{[4-[cyclopropyl(oxetan-3-yl)amino]cyclohexyl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl}piperazin-1-yl)-N-([4-{{[4-cyclopropylmorpholin-2-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({3-chloro-4-[(4-cyclopropylmorpholin-2-yl)methoxy]phenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 N-([3-chloro-4-{{[4-cyclopropylmorpholin-2-yl]methyl]amino]phenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 2-[[[2-chloro-4-{{[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]phenyl]amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
 50 (2S)-2-[[[3-chloro-5-{{[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]pyridin-2-yl]oxy]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
 N-([5-chloro-6-{{[4-cyclopropylmorpholin-2-yl]methyl]amino]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 2-[[[3-chloro-5-{{[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]pyridin-2-yl]amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
 55 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-([4-{{[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(cis-4-hydroxy-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
N-[[5-chloro-6-[[[(1R,2R,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(2-cyanoethyl)(cyclopropyl)amino]cyclohexyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
N-[[5-chloro-6-[[[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-chloro-6-(5,6,7,8-tetrahydroimidazo[1,2-a]pyridin-6-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
N-[[5-chloro-6-[[[(1R,2S,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(cis-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
N-[[5-chloro-6-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(3,3-difluoropyrrolidin-1-yl)cyclohexyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
N-[[5-chloro-6-[[[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 25 N-[[5-chloro-6-[[[(cis-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(2,2-difluorocyclopropyl)amino]cyclohexyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 30 N-[[5-chloro-6-[[[(cis-1-fluoro-4-hydroxycyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-(2-oxaspiro[3.5]non-7-ylmethoxy)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(cyclopropylmorpholin-2-yl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 N-[[5-chloro-6-[[[(trans-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
N-[[5-chloro-6-[[[(cis-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-cyano-4-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(trans-4-ethyl-4-hydroxycyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(cis-4-ethyl-4-hydroxycyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(2S)-4-(oxetan-3-yl)morpholin-2-yl]methyl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 N-[[3-chloro-4-[[[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(2-cyanoethyl)(cyclopropyl)amino]-1-fluorocyclohexyl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-nitro-6-[(tetrahydro-2H-pyran-4-ylmethyl)amino]pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[2-oxaspiro[3.5]non-7-ylmethyl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(4-cyano-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-[[3-(tetrahydro-2H-pyran-4-ylmethyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-6-yl]sulfonyl]benzamide;

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(6-([(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino)-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(5-cyano-6-([4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 N-(4-([4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl)sulfamoyl]-2-nitrophenyl)morpholine-4-carboxamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(4-([(4-(methoxymethyl)cyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(5-chloro-6-([1-(1,3-thiazol-2-yl)piperidin-4-yl]methoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(6-([(cis-4-hydroxy-4-methylcyclohexyl)methyl]amino)-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; and therapeutically acceptable salts, and metabolites thereof.

[0154] In another aspect, the present invention provides compounds of Formula (IV)



and therapeutically acceptable salts, and metabolites thereof,

wherein A¹, B¹, D¹, E¹, and Y¹ are as described herein for Formula (I); R¹⁰⁰ is as described for substituents on R²⁶; n is 0, 1, 2, or 3; R¹⁰¹ is as described for substituents on R³⁰; p is 0, 1, 2, 3, 4, 5, or 6; R¹⁰⁴ is as described for substituents on R³⁸; s is 0, 1, 2, 3, 4, 5, or 6; R¹⁰³ is as described for substituents on R⁵⁸; and r is 0, 1, 2, 3, or 4.

[0155] In one embodiment of Formula (IV), n, p, r, and s are each 0.

[0156] In one embodiment of Formula (IV), A¹ is N. In another embodiment of Formula (IV), A¹ is C(A²). In another embodiment of Formula (IV), A¹ is C(A²); and A² is H, F, Cl, Br, or I. In another embodiment of Formula (IV), A¹ is C(A²); and A² is H.

[0157] In one embodiment of Formula (IV), B¹ is R¹, OR¹, NHR¹, NHC(O)R¹, F, Cl, Br, or I. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; and B¹ is NHR¹. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; and B¹ is OR¹. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; and B¹ is Cl. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; and B¹ is R¹. In another embodiment of Formula (IV), A¹ is N; and B¹ is NHR¹. In another embodiment of Formula (IV), A¹ is N; and B¹ is OR¹. In another embodiment of Formula (IV), A¹ is N; and B¹ is Cl. In another embodiment of Formula (IV), A¹ is N; and B¹ is R¹.

[0158] In one embodiment of Formula (IV), D¹ is H or Cl. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; B¹ is NHR¹; and D¹ is H. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; B¹ is NHR¹; and D¹ is Cl. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; B¹ is OR¹; and D¹ is H. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; B¹ is Cl; and D¹ is H. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; B¹ is R¹;

and D¹ is H. In another embodiment of Formula (IV), A¹ is N; B¹ is NHR¹; and D¹ is H. In another embodiment of Formula (IV), A¹ is N; B¹ is NHR¹; and D¹ is Cl. In another embodiment of Formula (IV), A¹ is N; B¹ is OR¹; and D¹ is H. In another embodiment of Formula (IV), A¹ is N; B¹ is Cl; and D¹ is H. In another embodiment of Formula (IV), A¹ is N; B¹ is R¹; and D¹ is H.

[0159] In one embodiment of Formula (IV), E¹ is H. In another embodiment of Formula (IV), A¹ is C(A²); A¹ is H; B¹ is NHR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; B¹ is NHR¹; E¹ is H; and D¹ is Cl. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; B¹ is OR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; B¹ is Cl; D¹ is H; and E¹ is H. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; B¹ is R¹; D¹ is H; and E¹ is H. In another embodiment of Formula (IV), A¹ is N; B¹ is NHR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (IV), A¹ is N; B¹ is NHR¹; E¹ is H; and D¹ is Cl. In another embodiment of Formula (IV), A¹ is N; B¹ is OR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (IV), A¹ is N; B¹ is Cl; D¹ is H; and E¹ is H. In another embodiment of Formula (IV), A¹ is N; B¹ is R¹; D¹ is H; and E¹ is H.

[0160] In one embodiment of Formula (IV), Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SO₂R¹⁷, or C(O)NH₂. In another embodiment of Formula (IV), Y¹ is H. In another embodiment of Formula (IV), Y¹ is CN. In another embodiment of Formula (IV), Y¹ is F, Cl, Br, or I. In another embodiment of Formula (IV), Y¹ is CF₃. In another embodiment of Formula (IV), Y¹ is SR¹⁷. In another embodiment of Formula (IV), Y¹ is OR¹⁷. In another embodiment of Formula (IV), Y¹ is NO₂. In another embodiment of Formula (IV), Y¹ is SO₂R¹⁷; wherein R¹⁷ is as defined herein. In another embodiment of Formula (IV), Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl. In another embodiment of Formula (IV), Y¹ is R¹⁷; wherein R¹⁷ is alkynyl. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂ or SO₂R¹⁷; wherein R¹⁷ is alkyl or alkynyl. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂. In another embodiment of Formula (IV), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl substituted with three F. In another embodiment of Formula (IV), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂ or SO₂R¹⁷; wherein R¹⁷ is alkyl or alkynyl. In another embodiment of Formula (IV), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂. In another embodiment of Formula (IV), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl substituted with three F.

[0161] In one embodiment of Formula (IV), Y¹ and B¹, together with the atoms to which they are attached, are benzene. In another embodiment of Formula (IV), Y¹ and B¹, together with the atoms to which they are attached, are heteroarene. In another embodiment of Formula (IV), A¹ is C(A²); A², G¹, E¹, and D¹ are independently selected H; and Y¹ and B¹, together with the atoms to which they are attached, are benzene. In another embodiment of Formula (IV), A¹ is C(A²); A², G¹, E¹, and D¹ are independently selected H; and Y¹ and B¹, together with the atoms to which they are attached, are heteroarene.

[0162] In one embodiment of Formula (IV), R¹ is R⁴ or R⁵. In one embodiment of Formula (IV), R¹ is R⁴. In one embodiment of Formula (IV), R¹ is R⁵. In one embodiment of Formula (IV), R¹ is R⁴; and R⁴ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl. In one embodiment of Formula (IV), R¹ is R⁴; and R⁴ is cycloalkyl or heterocycloalkyl. In one embodiment of Formula (IV), R¹ is R⁴; and R⁴ is cycloalkyl. In one embodiment of Formula (IV), R¹ is R⁴; and R⁴ is heterocycloalkyl.

[0163] In one embodiment of Formula (IV), R¹ is R⁴; and R⁴ is cycloalkyl; wherein R⁴ is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted as defined herein. In another embodiment of Formula (IV), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted with R⁵⁷ or N(R⁵⁷)₂. In another embodiment of Formula (IV), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; and R⁵⁷ is R⁶⁰. In another embodiment of Formula (IV), R¹ is R⁴; R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; and R⁶⁰ is heterocycloalkyl. In another embodiment of Formula (IV), R¹ is R⁴; R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; R⁶⁰ is heterocycloalkyl; wherein the heterocycloalkyl ring is morpholinyl. In another embodiment of Formula (IV), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted with N(R⁵⁷)₂. In another embodiment of Formula (IV), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with N(R⁵⁷)₂. In another embodiment of Formula (IV), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with N(R⁵⁷)₂; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is unsubstituted or substituted with R⁶². In another embodiment of Formula (IV), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is unsubstituted or substituted with R⁶², R⁶² is R⁶⁵; and R⁶⁵ is cycloalkyl. In another embodiment of Formula (IV), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is substituted with R⁶²; R⁶² is R⁶⁵; and R⁶⁵ is cyclopropyl.

[0164] In one embodiment of Formula (IV), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein R⁴ is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is substituted as defined herein. In another embodiment of Formula (IV), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is substituted with R⁵⁷. In another embodiment of Formula (IV), R¹

is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the heterocycloalkyl ring is substituted with R⁵⁷; and R⁵⁷ is R⁶⁰ or R⁶¹. In another embodiment of Formula (IV), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰ or R⁶¹; R⁶⁰ is heterocycloalkyl; and R⁶¹ is alkyl. In another embodiment of Formula (IV), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; R⁶⁰ is heterocycloalkyl; wherein the heterocycloalkyl is tetrahydropyranyl. In another embodiment of Formula (IV), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; R⁶¹ is alkyl; and the alkyl is methyl. In another embodiment of Formula (IV), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with C(O)OR⁵⁷; R⁵⁷ is R⁶¹; R⁶¹ is alkyl; and the alkyl is methyl.

[0165] In one embodiment of Formula (IV), R¹ is R⁵; and R⁵ is alkyl which is unsubstituted or substituted. In one embodiment of Formula (IV), R¹ is R⁵; and R⁵ is alkyl which is unsubstituted or substituted with R⁷, OR⁷, OH, CN, or F. In another embodiment of Formula (IV), R¹ is R⁵; and R⁵ is alkyl which is substituted with R⁷, OR⁷, NHR⁷, or N(R⁷)₂.

[0166] In one embodiment of Formula (IV), R⁷ is R⁸, R⁹, R¹⁰ or R¹¹ which are unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R⁷ is R⁸ which is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R⁷ is R⁹ which is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R⁷ is R¹⁰ which is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R⁷ is R¹¹ which is unsubstituted or substituted as defined herein.

[0167] In one embodiment of Formula (IV), R⁸ is phenyl which is unsubstituted or substituted as defined herein.

[0168] In one embodiment of Formula (IV), R⁹ is heteroaryl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R⁹ is furanyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R⁹ is furanyl; which is unsubstituted.

[0169] In one embodiment of Formula (IV), R¹⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, heterocycloalkyl which are unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R¹⁰ is heterocycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R¹⁰ is tetrahydrofuranyl, tetrahydropyranyl, morpholinyl, dioxanyl, oxetanyl, piperdinyl, or pyrrolidinyl, which are unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R¹⁰ is tetrahydropyranyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R¹⁰ is morpholinyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R¹⁰ is cycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R¹⁰ is cycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R¹⁰ is cyclohexyl, cyclopropyl, cyclobutyl, or bicyclo[2.2.1]heptanyl, which is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R¹⁰ is heterocycloalkyl which is fused with R^{10A}; and R^{10A} is heteroarene. In another embodiment of Formula (IV), R¹⁰ is 5,6,7,8-tetrahydroimidazo[1,2-a]pyridinyl.

[0170] In one embodiment of Formula (IV), R¹¹ is alkyl, alkenyl or alkynyl which are unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R¹¹ is alkyl. In another embodiment of Formula (IV), R¹¹ is methyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (IV), R¹¹ is alkyl; which is substituted as defined herein. In another embodiment of Formula (IV), R¹¹ is alkyl; which is substituted with OR¹²; R¹² is R¹⁶; and R¹⁶ is alkyl. In another embodiment of Formula (IV), R¹¹ is alkyl; which is substituted with CF₃. In another embodiment of Formula (IV), R¹¹ is alkyl; which is substituted with R¹², R¹² is R¹⁴, and R¹⁴ is heteroaryl.

[0171] In one embodiment of Formula (IV),

n, p, r, and s are each 0;

A¹ is N or C(A²);

A¹ is H, F, Br, I, or Cl;

B¹ is R¹, OR¹, NHR¹, NHC(O)R¹, F, Br, I or Cl;

D¹ is H, F, Br, I, or Cl;

E¹ is H; and

Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SOR¹⁷, or C(O)NH₂; or

Y¹ and B¹, together with the atoms to which they are attached, are benzene, heteroarene, or heterocycloalkene; and A², D¹, and E¹ are independently selected H;

R¹ is R⁴ or R⁵;

R⁴ is cycloalkyl, or heterocycloalkyl;

R⁵ is alkyl, or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected

R⁷, OR⁷, NHR⁷, N(R⁷)₂, CN, OH, F, Cl, Br or I;

R⁷ is R⁸, R⁹, R¹⁰, or R¹¹;

R⁸ is phenyl;

R⁹ is heteroaryl;

R¹⁰ is cycloalkyl, cycloalkenyl, or heterocycloalkyl; each of which is unfused or fused with R^{10A}, R^{10A} is heteroarene;

R¹¹ is alkyl each of which is unsubstituted or substituted with one or two or three of independently selected R¹², OR¹² or CF₃;

R¹² is R¹⁴ or R¹⁶;

R¹⁴ is heteroaryl;

R¹⁶ is alkyl;

R¹⁷ is R²¹;

R²¹ is alkyl, or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R²², F, Cl, Br or I;

R²² is R²⁵;

R²⁵ is heterocycloalkyl;

wherein the cyclic moieties represented by Y¹ and B¹ together, R⁴, R⁸, R¹⁰, and R²⁵, are independently unsubstituted, further unsubstituted, substituted or further substituted with one or two or three or four or five of independently selected R^{57A}, R⁵⁷, OR⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, C(O)N(R⁵⁷)₂, NH₂, NHR⁵⁷, N(R⁵⁷)₂, NHC(O)R⁵⁷, NHS(O)₂R⁵⁷, OH, CN, (O), F, Cl, Br or I;

R^{57A} is spiroalkyl, or spiroheteroalkyl;

R⁵⁷ is R⁵⁸, R⁶⁰ or R⁶¹;

R⁵⁸ is phenyl;

R⁶⁰ is cycloalkyl, or heterocycloalkyl;

R⁶¹ is alkyl which is unsubstituted or substituted with one or two or three of independently selected R⁶², OR⁶², N(R⁶²)₂, C(O)OH, CN, F, Cl, Br or I;

R⁶² is R⁶⁵, or R⁶⁶;

R⁶⁵ is cycloalkyl, or heterocycloalkyl;

R⁶⁶ is alkyl which is unsubstituted or substituted with OR⁶⁷;

R⁶⁷ is alkyl;

wherein the cyclic moieties represented by R^{57A}, R⁵⁸, and R⁶⁰ are unsubstituted or substituted with one or two or three or four of independently selected R⁶⁸, F, Cl, Br or I;


R⁶⁸ is R⁷¹ or R⁷²;

R⁷¹ is heterocycloalkyl; and

R⁷² is alkyl, which is unsubstituted or substituted with one or two F.

[0172] Still another embodiment pertains to compounds having Formula (IV), which are 4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({4-[(3-morpholin-4-ylpropyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; and therapeutically acceptable salts, and metabolites thereof.

[0173] In another aspect, the present invention provides compounds of Formula (V)



and therapeutically acceptable salts, and metabolites thereof,
 wherein A¹, B¹, D¹, E¹, and Y¹ are as described herein for Formula (I); R¹⁰⁰ is as described for substituents on R²⁶; n is 0, 1, 2, or 3; R¹⁰¹ is as described for substituents on R³⁰; p is 0, 1, 2, 3, 4, 5, or 6; R¹⁰⁵ is as described for substituents on R⁴⁰; t is 0, 1, 2, 3, or 4; R¹⁰³ is as described for substituents on R⁵⁸; and r is 0, 1, 2, 3, or 4.

[0174] In one embodiment of Formula (V), n, p, r, and t are each 0.

[0175] In one embodiment of Formula (V), A¹ is N. In another embodiment of Formula (V), A¹ is C(A²). In another embodiment of Formula (V), A¹ is C(A²); and A² is H, F, Cl, Br, or I. In another embodiment of Formula (V), A¹ is C(A²); and A² is H.

[0176] In one embodiment of Formula (V), B¹ is R¹, OR¹, NHR¹, NHC(O)R¹, F, Cl, Br, or I. In another embodiment of Formula (V), A¹ is C(A²); A² is H; and B¹ is NHR¹. In another embodiment of Formula (V), A¹ is C(A²); A² is H; and B¹ is OR¹. In another embodiment of Formula (V), A¹ is C(A²); A² is H; and B¹ is Cl. In another embodiment of Formula (V), A¹ is C(A²); A² is H; and B¹ is R¹. In another embodiment of Formula (V), A¹ is N; and B¹ is NHR¹. In another embodiment of Formula (V), A¹ is N; and B¹ is OR¹. In another embodiment of Formula (V), A¹ is N; and B¹ is Cl. In another embodiment of Formula (V), A¹ is N; and B¹ is R¹.

[0177] In one embodiment of Formula (V), D¹ is H or Cl. In another embodiment of Formula (V), A¹ is C(A²); A² is H; B¹ is NHR¹; and D¹ is H. In another embodiment of Formula (V), A¹ is C(A²); A² is H; B¹ is NHR¹; and D¹ is Cl. In another embodiment of Formula (V), A¹ is C(A²); A² is H; B¹ is OR¹; and D¹ is H. In another embodiment of Formula (V), A¹ is C(A²); A² is H; B¹ is Cl; and D¹ is H. In another embodiment of Formula (V), A¹ is C(A²); A² is H; B¹ is R¹; and D¹ is H. In another embodiment of Formula (V), A¹ is N; B¹ is NHR¹; and D¹ is H. In another embodiment of Formula (V), A¹ is N; B¹ is NHR¹; and D¹ is Cl. In another embodiment of Formula (V), A¹ is N; B¹ is OR¹; and D¹ is H. In another embodiment of Formula (V), A¹ is N; B¹ is Cl; and D¹ is H. In another embodiment of Formula (V), A¹ is N; B¹ is R¹; and D¹ is H.

[0178] In one embodiment of Formula (V), E¹ is H. In another embodiment of Formula (V), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (V), A¹ is C(A²); A² is H; B¹ is NHR¹; E¹ is H; and D¹ is Cl. In another embodiment of Formula (V), A¹ is C(A²); A² is H; B¹ is OR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (V), A¹ is C(A²); A² is H; B¹ is Cl; D¹ is H; and E¹ is H. In another embodiment of Formula (V), A¹ is C(A²); A² is H; B¹ is R¹; D¹ is H; and E¹ is H. In another embodiment of Formula (V), A¹ is N; B¹ is NHR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (V), A¹ is N; B¹ is NHR¹; E¹ is H; and D¹ is Cl. In another embodiment of Formula (V), A¹ is N; B¹ is OR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (V), A¹ is N; B¹ is Cl; D¹ is H; and E¹ is H. In another embodiment of Formula (V), A¹ is N; B¹ is R¹; D¹ is H; and E¹ is H.

[0179] In one embodiment of Formula (V), Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SO₂R¹⁷, or C(O)NH₂. In another embodiment of Formula (V), Y¹ is H. In another embodiment of Formula (V), Y¹ is CN. In another embodiment of Formula (V), Y¹ is F, Cl, Br, or I. In another embodiment of Formula (V), Y¹ is CF₃. In another embodiment of Formula (V), Y¹ is SR¹⁷. In another embodiment of Formula (V), Y¹ is OR¹⁷. In another embodiment of Formula (V), Y¹ is NO₂. In another embodiment of Formula (V), Y¹ is SO₂R¹⁷; wherein R¹⁷ is as defined herein. In another embodiment of

Formula (V), Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl. In another embodiment of Formula (V), Y¹ is R¹⁷; wherein R¹⁷ is alkynyl. In another embodiment of Formula (V), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂ or SO₂R¹⁷; wherein R¹⁷ is alkyl or alkynyl. In another embodiment of Formula (V), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂. In another embodiment of Formula (V), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl substituted with three F. In another embodiment of Formula (V), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂ or SO₂R¹⁷; wherein R¹⁷ is alkyl or alkynyl. In another embodiment of Formula (V), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂. In another embodiment of Formula (V), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl substituted with three F.

[0180] In one embodiment of Formula (V), Y¹ and B¹, together with the atoms to which they are attached, are benzene. In another embodiment of Formula (V), Y¹ and B¹, together with the atoms to which they are attached, are heteroarene. In another embodiment of Formula (V), A¹ is C(A²); A², G¹, E¹, and D¹ are independently selected H; and Y¹ and B¹, together with the atoms to which they are attached, are benzene. In another embodiment of Formula (V), A¹ is C(A²); A², G¹, E¹, and D¹ are independently selected H; and Y¹ and B¹, together with the atoms to which they are attached, are heteroarene.

[0181] In one embodiment of Formula (V), R¹ is R⁴ or R⁵. In one embodiment of Formula (V), R¹ is R⁴. In one embodiment of Formula (V), R¹ is R⁵. In one embodiment of Formula (V), R¹ is R⁴; and R⁴ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl. In one embodiment of Formula (V), R¹ is R⁴; and R⁴ is cycloalkyl or heterocycloalkyl. In one embodiment of Formula (V), R¹ is R⁴; and R⁴ is cycloalkyl. In one embodiment of Formula (V), R¹ is R⁴; and R⁴ is heterocycloalkyl.

[0182] In one embodiment of Formula (V), R¹ is R⁴; and R⁴ is cycloalkyl; wherein R⁴ is unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted as defined herein. In another embodiment of Formula (V), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted with R⁵⁷ or N(R⁵⁷)₂. In another embodiment of Formula (V), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; and R⁵⁷ is R⁶⁰. In another embodiment of Formula (V), R¹ is R⁴; R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; and R⁶⁰ is heterocycloalkyl. In another embodiment of Formula (V), R¹ is R⁴; R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; R⁶⁰ is heterocycloalkyl; wherein the heterocycloalkyl ring is morpholinyl. In another embodiment of Formula (V), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted with N(R⁵⁷)₂. In another embodiment of Formula (V), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with N(R⁵⁷)₂. In another embodiment of Formula (V), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with N(R⁵⁷)₂; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is unsubstituted or substituted with R⁶². In another embodiment of Formula (V), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is unsubstituted or substituted with R⁶²; R⁶² is R⁶⁵; and R⁶⁵ is cycloalkyl. In another embodiment of Formula (V), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is substituted with R⁶²; R⁶² is R⁶⁵; and R⁶⁵ is cyclopropyl.

[0183] In one embodiment of Formula (V), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein R⁴ is unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is substituted as defined herein. In another embodiment of Formula (V), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is substituted with R⁵⁷. In another embodiment of Formula (V), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the heterocycloalkyl ring is substituted with R⁵⁷; and R⁵⁷ is R⁶⁰ or R⁶¹. In another embodiment of Formula (V), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰ or R⁶¹; R⁶⁰ is heterocycloalkyl; and R⁶¹ is alkyl. In another embodiment of Formula (V), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; R⁶⁰ is heterocycloalkyl; wherein the heterocycloalkyl ring is tetrahydropyranyl. In another embodiment of Formula (V), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; R⁶¹ is alkyl; and the alkyl is methyl. In another embodiment of Formula (V), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperdinyl or piperizinyl; and wherein the piperdinyl or piperizinyl ring is substituted with C(O)OR⁵⁷; R⁵⁷ is R⁶¹; R⁶¹ is alkyl; and the alkyl is methyl.

[0184] In one embodiment of Formula (V), R¹ is R⁵; and R⁵ is alkyl which is unsubstituted or substituted. In one embodiment of Formula (V), R¹ is R⁵; and R⁵ is alkyl which is unsubstituted or substituted with R⁷, OR⁷, OH, CN, or F. In another embodiment of Formula (V), R¹ is R⁵; and R⁵ is alkyl which is substituted with R⁷, OR⁷, NHR⁷, or N(R⁷)₂.

[0185] In one embodiment of Formula (V), R⁷ is R⁸, R⁹, R¹⁰ or R¹¹ which are unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R⁷ is R⁸ which is unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R⁷ is R⁹ which is unsubstituted or substituted as defined herein. In another embodiment of

Formula (V), R⁷ is R¹⁰ which is unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R⁷ is R¹¹ which is unsubstituted or substituted as defined herein.

[0186] In one embodiment of Formula (V), R⁸ is phenyl which is unsubstituted or substituted as defined herein.

[0187] In one embodiment of Formula (V), R⁹ is heteroaryl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R⁹ is furanyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R⁹ is furanyl; which is unsubstituted.

[0188] In one embodiment of Formula (V), R¹⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, heterocycloalkyl which are unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R¹⁰ is heterocycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R¹⁰ is tetrahydrofuranyl, tetrahydropyranyl, morpholinyl, dioxanyl, oxetanyl, piperdinyl, or pyrrolidinyl, which are unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R¹⁰ is tetrahydropyranyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R¹⁰ is morpholinyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R¹⁰ is cycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R¹⁰ is cycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R¹⁰ is cyclohexyl, cyclopropyl, cyclobutyl, or bicyclo[2.2.1]heptanyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R¹⁰ is heterocycloalkyl which is fused with R^{10A}; and R^{10A} is heteroarene. In another embodiment of Formula (V), R¹⁰ is 5,6,7,8-tetrahydroimidazo[1,2-a]pyridinyl.

[0189] In one embodiment of Formula (V), Rⁿ is alkyl, alkenyl or alkynyl which are unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R¹¹ is alkyl. In another embodiment of Formula (V), R¹¹ is methyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (V), R¹¹ is alkyl; which is substituted as defined herein. In another embodiment of Formula (V), R¹¹ is alkyl; which is substituted with R¹², OR¹², or CF₃. In another embodiment of Formula (V), R¹¹ is alkyl; which is substituted with OR¹²; R¹² is R¹⁶; and R¹⁶ is alkyl. In another embodiment of Formula (V), R¹¹ is alkyl; which is substituted with CF₃. In another embodiment of Formula (V), R¹¹ is alkyl; which is substituted with R¹²; R¹² is R¹⁴; and R¹⁴ is heteroaryl.

[0190] In one embodiment of Formula (V),

n, p, r, and t are each 0;

A¹ is N or C(A²);

A² is H, F, Br, I, or Cl;

B¹ is R¹, OR¹, NHR¹, NHC(O)R¹, F, Br, I or Cl;

D¹ is H, F, Br, I, or Cl;

E¹ is H; and

Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SO₂R¹⁷, or C(O)NH₂; or

Y¹ and B¹, together with the atoms to which they are attached, are benzene, heteroarene, or heterocycloalkene; and A², D¹, and E¹ are independently selected H;

R¹ is R⁴ or R⁵;

R⁴ is cycloalkyl, or heterocycloalkyl;

R⁵ is alkyl, or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected

R⁷, OR⁷, NHR⁷, N(R⁷)₂, CN, OH, F, Cl, Br or I;

R⁷ is R⁸, R⁹, R¹⁰, or R¹¹;

R⁸ is phenyl;

R⁹ is heteroaryl;

R¹⁰ is cycloalkyl, cycloalkenyl, or heterocycloalkyl; each of which is unfused or fused with R^{10A}, R^{10A} is heteroarene;

R¹¹ is alkyl each of which is unsubstituted or substituted with one or two or three of independently selected R¹², OR¹² or CF₃;

R¹² is R¹⁴ or R¹⁶;

R¹⁴ is heteroaryl;

R¹⁶ is alkyl;

R¹⁷ is R²¹;

R²¹ is alkyl, or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R²², F, Cl, Br or I;

R²² is R²⁵;

R²⁵ is heterocycloalkyl;

wherein the cyclic moieties represented by Y¹ and B¹ together, R⁴, R⁸, R¹⁰, and R²⁵, are independently unsubstituted, further unsubstituted, substituted or further substituted with one or two or three or four or five of independently selected R^{57A}, R⁵⁷, OR⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, C(O)N(R⁵⁷)₂, NH₂, NHR⁵⁷, N(R⁵⁷)₂, NHC(O)R⁵⁷, NHS(O)₂R⁵⁷, OH, CN, (O), F, Cl, Br or I;

R^{57A} is spiroalkyl, or spiroheteroalkyl;

R⁵⁷ is R⁵⁸, R⁶⁰ or R⁶¹;

R⁵⁸ is phenyl;

R⁶⁰ is cycloalkyl, or heterocycloalkyl;

5 R⁶¹ is alkyl which is unsubstituted or substituted with one or two or three of independently selected R⁶², OR⁶², N(R⁶²)₂, C(O)OH, CN, F, Cl, Br or I;

R⁶² is R⁶⁵, or R⁶⁶;

R⁶⁵ is cycloalkyl, or heterocycloalkyl;

R⁶⁶ is alkyl which is unsubstituted or substituted with OR⁶⁷;

10 R⁶⁷ is alkyl;

wherein the cyclic moieties represented by R^{57A}, R⁵⁸, and R⁶⁰ are unsubstituted or substituted with one or two or three or four of independently selected R⁶⁸, F, Cl, Br or I;

R⁶⁸ is R⁷¹ or R⁷²;

R⁷¹ is heterocycloalkyl; and

15 R⁷² is alkyl, which is unsubstituted or substituted with one or two F.

[0191] Still another embodiment pertains to a compound having Formula (V), which is

20 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(1,4-dioxan-2-ylmethyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-[[4-(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl}sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 Trans-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl}sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-[[5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl}sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-(trifluoromethyl)phenyl}sulfonyl)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)benzamide;
 35 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(1-methylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-methylpiperazin-1-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-methoxycyclohexyl)methyl]amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;


45

and therapeutically acceptable salts, and metabolites thereof.

[0192] In another aspect, the present invention provides compounds of Formula (VI)

50

55



(VI);

and therapeutically acceptable salts, and metabolites thereof,

wherein A¹, B¹, D¹, E¹, and Y¹ are as described herein for Formula (I); R¹⁰⁰ is as described for substituents on R²⁶; n is 0, 1, 2, or 3; R¹⁰¹ is as described for substituents on R³⁰; p is 0, 1, 2, 3, 4, 5, or 6; R¹⁰² is as described for substituents on R⁴⁰; q is 0, 1, 2, 3, 4, 5, or 6; R¹⁰³ is as described for substituents on R⁵⁸; and r is 0, 1, 2, 3, or 4.

[0193] In one embodiment of Formula (VI), n, p, r, and q are each 0.

[0194] In one embodiment of Formula (VI), A¹ is N. In another embodiment of Formula (VI), A¹ is C(A²). In another embodiment of Formula (VI), A¹ is C(A²); and A² is H, F, Cl, Br, or I. In another embodiment of Formula (VI), A¹ is C(A²); and A² is H.

[0195] In one embodiment of Formula (VI), B¹ is R¹, OR¹, NHR¹, NHC(O)R¹, F, Cl, Br, or I. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; and B¹ is NHR¹. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; and B¹ is OR¹. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; and B¹ is Cl. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; and B¹ is R¹. In another embodiment of Formula (VI), A¹ is N; and B¹ is NHR¹. In another embodiment of Formula (VI), A¹ is N; and B¹ is OR¹. In another embodiment of Formula (VI), A¹ is N; and B¹ is Cl. In another embodiment of Formula (VI), A¹ is N; and B¹ is R¹.

[0196] In one embodiment of Formula (VI), D¹ is H or Cl. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; B¹ is NHR¹; and D¹ is H. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; B¹ is NHR¹; and D¹ is Cl. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; B¹ is OR¹; and D¹ is H. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; B¹ is Cl; and D¹ is H. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; B¹ is R¹; and D¹ is H. In another embodiment of Formula (VI), A¹ is N; B¹ is NHR¹; and D¹ is H. In another embodiment of Formula (VI), A¹ is N; B¹ is NHR¹; and D¹ is Cl. In another embodiment of Formula (VI), A¹ is N; B¹ is OR¹; and D¹ is H. In another embodiment of Formula (VI), A¹ is N; B¹ is Cl; and D¹ is H. In another embodiment of Formula (VI), A¹ is N; B¹ is R¹; and D¹ is H.

[0197] In one embodiment of Formula (VI), E¹ is H. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; B¹ is NHR¹; E¹ is H; and D¹ is Cl. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; B¹ is OR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; B¹ is Cl; D¹ is H; and E¹ is H. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; B¹ is R¹; D¹ is H; and E¹ is H. In another embodiment of Formula (VI), A¹ is N; B¹ is NHR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (VI), A¹ is N; B¹ is NHR¹; E¹ is H; and D¹ is Cl. In another embodiment of Formula (VI), A¹ is N; B¹ is OR¹; D¹ is H; and E¹ is H. In another embodiment of Formula (VI), A¹ is N; B¹ is Cl; D¹ is H; and E¹ is H. In another embodiment of Formula (VI), A¹ is N; B¹ is R¹; D¹ is H; and E¹ is H.

[0198] In one embodiment of Formula (VI), Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SO₂R¹⁷, or C(O)NH₂. In another embodiment of Formula (VI), Y¹ is H. In another embodiment of Formula (VI), Y¹ is CN. In another embodiment of Formula (VI), Y¹ is F, Cl, Br, or I. In another embodiment of Formula (VI), Y¹ is CF₃. In another embodiment of Formula

(VI), Y¹ is SR¹⁷. In another embodiment of Formula (VI), Y¹ is OR¹⁷. In another embodiment of Formula (VI), Y¹ is NO₂. In another embodiment of Formula (VI), Y¹ is SO₂R¹⁷; wherein R¹⁷ is as defined herein. In another embodiment of Formula (VI), Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl. In another embodiment of Formula (VI), Y¹ is R¹⁷; wherein R¹⁷ is alkynyl. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂ or SO₂R¹⁷; wherein R¹⁷ is alkyl or alkynyl. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂. In another embodiment of Formula (VI), A¹ is C(A²); A² is H; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl substituted with three F. In another embodiment of Formula (VI), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂ or SO₂R¹⁷; wherein R¹⁷ is alkyl or alkynyl. In another embodiment of Formula (VI), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is NO₂. In another embodiment of Formula (VI), A¹ is N; B¹ is NHR¹; D¹ is H; E¹ is H; and Y¹ is SO₂R¹⁷; wherein R¹⁷ is alkyl substituted with three F.

[0199] In one embodiment of Formula (IV), Y¹ and B¹, together with the atoms to which they are attached, are benzene. In another embodiment of Formula (IV), Y¹ and B¹, together with the atoms to which they are attached, are heteroarene. In another embodiment of Formula (IV), A¹ is C(A²); A², G¹, E¹, and D¹ are independently selected H; and Y¹ and B¹, together with the atoms to which they are attached, are benzene. In another embodiment of Formula (IV), A¹ is C(A²); A², G¹, E¹, and D¹ are independently selected H; and Y¹ and B¹, together with the atoms to which they are attached, are heteroarene.

[0200] In one embodiment of Formula (VI), R¹ is R⁴ or R⁵. In one embodiment of Formula (VI), R¹ is R⁴. In one embodiment of Formula (VI), R¹ is R⁵. In one embodiment of Formula (VI), R¹ is R⁴; and R⁴ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl. In one embodiment of Formula (VI), R¹ is R⁴; and R⁴ is cycloalkyl or heterocycloalkyl. In one embodiment of Formula (VI), R¹ is R⁴; and R⁴ is cycloalkyl. In one embodiment of Formula (VI), R¹ is R⁴; and R⁴ is heterocycloalkyl.

[0201] In one embodiment of Formula (VI), R¹ is R⁴; and R⁴ is cycloalkyl; wherein R⁴ is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted as defined herein. In another embodiment of Formula (VI), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted with R⁵⁷ or N(R⁵⁷)₂. In another embodiment of Formula (VI), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; and R⁵⁷ is R⁶⁰. In another embodiment of Formula (VI), R¹ is R⁴; R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; and R⁶⁰ is heterocycloalkyl. In another embodiment of Formula (VI), R¹ is R⁴; R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; R⁶⁰ is heterocycloalkyl; wherein the heterocycloalkyl ring is morpholinyl. In another embodiment of Formula (VI), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is substituted with N(R⁵⁷)₂. In another embodiment of Formula (VI), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with N(R⁵⁷)₂. In another embodiment of Formula (VI), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with N(R⁵⁷)₂; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is unsubstituted or substituted with R⁶². In another embodiment of Formula (VI), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is unsubstituted or substituted with R⁶²; R⁶² is R⁶⁵; and R⁶⁵ is cycloalkyl. In another embodiment of Formula (VI), R¹ is R⁴; and R⁴ is cycloalkyl; wherein the cycloalkyl ring is cyclohexyl; and wherein the cyclohexyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; and R⁶¹ is alkyl which is substituted with R⁶²; R⁶² is R⁶⁵; and R⁶⁵ is cyclopropyl.

[0202] In one embodiment of Formula (VI), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein R⁴ is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is substituted as defined herein. In another embodiment of Formula (VI), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is substituted with R⁵⁷. In another embodiment of Formula (VI), R¹ is R⁴; and R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperidinyl or piperizinyl; and wherein the heterocycloalkyl ring is substituted with R⁵⁷; and R⁵⁷ is R⁶⁰ or R⁶¹. In another embodiment of Formula (VI), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperidinyl or piperizinyl; and wherein the piperidinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰ or R⁶¹; R⁶⁰ is heterocycloalkyl; and R⁶¹ is alkyl. In another embodiment of Formula (VI), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperidinyl or piperizinyl; and wherein the piperidinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶⁰; R⁶⁰ is heterocycloalkyl; wherein the heterocycloalkyl is tetrahydropyranyl. In another embodiment of Formula (VI), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperidinyl or piperizinyl; and wherein the piperidinyl or piperizinyl ring is substituted with R⁵⁷; R⁵⁷ is R⁶¹; R⁶¹ is alkyl; and the alkyl is methyl. In another embodiment of Formula (VI), R¹ is R⁴; R⁴ is heterocycloalkyl; wherein the heterocycloalkyl ring is piperidinyl or piperizinyl; and wherein the piperidinyl or piperizinyl ring is substituted with C(O)OR⁵⁷; R⁵⁷ is R⁶¹; R⁶¹ is alkyl; and the alkyl is methyl.

[0203] In one embodiment of Formula (VI), R¹ is R⁵; and R⁵ is alkyl which is unsubstituted or substituted. In one embodiment of Formula (VI), R¹ is R⁵; and R⁵ is alkyl which is unsubstituted or substituted with R⁷, OR⁷, OH, CN, or F. In another embodiment of Formula (VI), R¹ is R⁵; and R⁵ is alkyl which is substituted with R⁷, OR⁷, NHR⁷, or N(R⁷)₂.

[0204] In one embodiment of Formula (VI), R⁷ is R⁸, R⁹, R¹⁰ or R¹¹ which are unsubstituted or substituted as defined

herein. In another embodiment of Formula (VI), R⁷ is R⁸ which is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R⁷ is R⁹ which is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R⁷ is R¹⁰ which is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R⁷ is R¹¹ which is unsubstituted or substituted as defined herein.

[0205] In one embodiment of Formula (VI), R⁸ is phenyl which is unsubstituted or substituted as defined herein.

[0206] In one embodiment of Formula (VI), R⁹ is heteroaryl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R⁹ is furanyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R⁹ is furanyl; which is unsubstituted.

[0207] In one embodiment of Formula (VI), R¹⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, heterocycloalkyl which are unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R¹⁰ is heterocycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R¹⁰ is tetrahydrofuranyl, tetrahydropyranyl, morpholinyl, dioxanyl, oxetanyl, piperdinyl, or pyrrolidinyl; which are unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R¹⁰ is tetrahydropyranyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R¹⁰ is morpholinyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R¹⁰ is cycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R¹⁰ is cycloalkyl which is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R¹⁰ is cyclohexyl, cyclopropyl, cyclobutyl, or bicyclo[2.2.1]heptanyl, which is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R¹⁰ is heterocycloalkyl which is fused with R^{10A}; and R^{10A} is heteroarene. In another embodiment of Formula (VI), R¹⁰ is 5,6,7,8-tetrahydroimidazo[1,2-a]pyridinyl.

[0208] In one embodiment of Formula (VI), R¹¹ is alkyl, alkenyl or alkynyl which are unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R¹¹ is alkyl. In another embodiment of Formula (VI), R¹¹ is methyl; which is unsubstituted or substituted as defined herein. In another embodiment of Formula (VI), R¹¹ is alkyl; which is substituted as defined herein. In another embodiment of Formula (VI), R¹¹ is alkyl; which is substituted with R¹², OR¹², or CF₃. In another embodiment of Formula (VI), R¹¹ is alkyl, which is substituted with OR¹²; R¹² is R¹⁶; and R¹⁶ is alkyl. In another embodiment of Formula (VI), R¹¹ is alkyl; which is substituted with CF₃. In another embodiment of Formula (VI), R¹¹ is alkyl, which is substituted with R¹²; R¹² is R¹⁴; and R¹⁴ is heteroaryl.

[0209] In one embodiment of Formula (VI),

n, p, r, and q are each 0;

A¹ is N or C(A²);

A² is H, F, Br, I, or Cl;

B¹ is R¹, OR¹, NHR¹, NHC(O)R¹, F, Br, I or Cl;

D¹ is H, F, Br, I, or Cl;

E¹ is H; and

Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SO₂R¹⁷, or C(O)NH₂; or

Y¹ and B¹, together with the atoms to which they are attached, are benzene, heteroarene, or heterocycloalkene; and A², D¹, and E¹ are independently selected H;

R¹ is R⁴ or R⁵;

R⁴ is cycloalkyl, or heterocycloalkyl;

R⁵ is alkyl, or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁷, OR⁷, NHR⁷, N(R⁷)₂, CN, OH, F, Cl, Br or I;

R⁷ is R⁸, R⁹, R¹⁰, or R¹¹;

R⁸ is phenyl;

R⁹ is heteroaryl;

R¹⁰ is cycloalkyl, cycloalkenyl, or heterocycloalkyl; each of which is unfused or fused with R^{10A}, R^{10A} is heteroarene;

R¹¹ is alkyl each of which is unsubstituted or substituted with one or two or three of independently selected R¹², OR¹² or CF₃;

R¹² is R¹⁴ or R¹⁶;

R¹⁴ is heteroaryl;

R¹⁶ is alkyl;

R¹⁷ is R²¹;

R²¹ is alkyl, or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R²², F, Cl, Br or I;

R²² is R²⁵;

R²⁵ is heterocycloalkyl;

wherein the cyclic moieties represented by Y¹ and B¹ together, R⁴, R⁸, R¹⁰, and R²⁵, are independently unsubstituted, further unsubstituted, substituted or further substituted with one or two or three or four or five of independently

selected R^{57A} , R^{57} , OR^{57} , SO_2R^{57} , $C(O)R^{57}$, $CO(O)R^{57}C(O)N(R^{57})_2$, NH_2 , NHR^{57} , $N(R^{57})_2$, $NHC(O)R^{57}$, $NHS(O)_2R^{57}$, OH , CN , (O) , F , Cl , Br or I ;

R^{57A} is spiroalkyl, or spiroheteroalkyl;

R^{57} is R^{58} , R^{60} or R^{61} ;

5 R^{58} is phenyl;

R^{60} is cycloalkyl, or heterocycloalkyl;

R^{61} is alkyl which is unsubstituted or substituted with one or two or three of independently selected R^{62} , OR^{62} , $N(R^{62})_2$, $C(O)OH$, CN , F , Cl , Br or I ;

R^{62} is R^{65} , or R^{66} ;

10 R^{65} is cycloalkyl, or heterocycloalkyl;

R^{66} is alkyl which is unsubstituted or substituted with OR^{67} ;

R^{67} is alkyl;

wherein the cyclic moieties represented by R^{57A} , R^{58} , and R^{60} are unsubstituted or substituted with one or two or three or four of independently selected R^{68} , F , Cl , Br or I ;

15 R^{68} is R^{71} or R^{72} ;

R^{71} is heterocycloalkyl; and

R^{72} is alkyl, which is unsubstituted or substituted with one or two F .

[0210] Still another embodiment pertains to a compound having Formula (VI), which is

20

4-(4-([2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-([2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

25

4-(4-([2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

and therapeutically acceptable salts, and metabolites thereof.

30

Pharmaceutical Compositions, Combination Therapies, Methods of Treatment, and Administration

[0211] Another embodiment comprises pharmaceutical compositions comprising a compound having Formula (I) and an excipient.

35

[0212] Still another embodiment comprises methods of treating cancer in a mammal comprising administering thereto a therapeutically acceptable amount of a compound having Formula (I).

[0213] Still another embodiment comprises methods of treating autoimmune disease in a mammal comprising administering thereto a therapeutically acceptable amount of a compound having Formula (I).

40

[0214] Still another embodiment pertains to compositions for treating diseases during which anti-apoptotic Bcl-2 proteins are expressed, said compositions comprising an excipient and a therapeutically effective amount of the compound having Formula (I).

[0215] Still another embodiment pertains to methods of treating disease in a patient during which anti-apoptotic Bcl-2 proteins are expressed, said methods comprising administering to the patient a therapeutically effective amount of a compound having Formula (I).

45

[0216] Still another embodiment pertains to compositions for treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, prostate cancer, small cell lung cancer or spleen cancer, said compositions comprising an excipient and a therapeutically effective amount of the compound having Formula (I).

50

[0217] Still another embodiment pertains to methods of treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, prostate cancer, small cell lung cancer or spleen cancer in a patient, said methods comprising administering to the patient a therapeutically effective amount of a compound having Formula (I).

55

[0218] Still another embodiment pertains to compositions for treating diseases during which are expressed anti-apoptotic Bcl-2 proteins, said compositions comprising an excipient and a therapeutically effective amount of the compound having Formula (I) and a therapeutically effective amount of one additional therapeutic agent or more than one additional

therapeutic agent.

[0219] Still another embodiment pertains to methods of treating disease in a patient during which are expressed anti-apoptotic Bcl-2 proteins, said methods comprising administering to the patient a therapeutically effective amount of a compound having Formula (I) and a therapeutically effective amount of one additional therapeutic agent or more than one additional therapeutic agent.

[0220] Still another embodiment pertains to compositions for treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, chronic lymphocytic leukemia, myeloma, prostate cancer, small cell lung cancer or spleen cancer, said compositions comprising an excipient and a therapeutically effective amount of the compound having Formula (I) and a therapeutically effective amount of one additional therapeutic agent or more than one additional therapeutic agent.

[0221] Still another embodiment pertains to methods of treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, chronic lymphocytic leukemia, myeloma, prostate cancer, small cell lung cancer or spleen cancer in a patient, said methods comprising administering to the patient a therapeutically effective amount of the compound having Formula (I) and a therapeutically effective amount of one additional therapeutic agent or more than one additional therapeutic agent.

[0222] Metabolites of compounds having Formula (I), produced by in vitro or in vivo metabolic processes, may also have utility for treating diseases associated with anti-apoptotic Bcl-2 proteins.

[0223] Certain precursor compounds which may be metabolized in vitro or in vivo to form compounds having Formula (I) may also have utility for treating diseases associated with expression of anti-apoptotic Bcl-2 proteins.

[0224] Compounds having Formula (I) may exist as acid addition salts, basic addition salts or zwitterions. Salts of the compounds are prepared during isolation or following purification of the compounds. Acid addition salts of the compounds are those derived from the reaction of the compounds with an acid. For example, the acetate, adipate, alginate, bicarbonate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, formate, fumarate, glycerophosphate, glutamate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, lactobionate, lactate, maleate, mesitylenesulfonate, methanesulfonate, naphthylenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, phosphate, picrate, propionate, succinate, tartrate, thiocyanate, trichloroacetic, trifluoroacetic, para-toluenesulfonate, and undecanoate salts of the compounds are contemplated as being embraced by this invention. Basic addition salts of the compounds are those derived from the reaction of the compounds with the hydroxide, carbonate or bicarbonate of cations such as lithium, sodium, potassium, calcium, and magnesium.

[0225] The compounds having Formula (I) may be administered, for example, buccally, ophthalmically, orally, osmotically, parenterally (intramuscularly, intraperitoneally, intrasternally, intravenously, subcutaneously), rectally, topically, transdermally or vaginally.

[0226] Therapeutically effective amounts of compounds having Formula (I) depend on the recipient of the treatment, the disorder being treated and the severity thereof, the composition containing the compound, the time of administration, the route of administration, the duration of treatment, the compound potency, its rate of clearance and whether or not another drug is co-administered. The amount of a compound of this invention having Formula (I) used to make a composition to be administered daily to a patient in a single dose or in divided doses is from about 0.03 to about 200 mg/kg body weight. Single dose compositions contain these amounts or a combination of submultiples thereof.

[0227] Compounds having Formula (I) may be administered with or without an excipient. Excipients include, for example, encapsulating materials or additives such as absorption accelerators, antioxidants, binders, buffers, coating agents, coloring agents, diluents, disintegrating agents, emulsifiers, extenders, fillers, flavoring agents, humectants, lubricants, perfumes, preservatives, propellants, releasing agents, sterilizing agents, sweeteners, solubilizers, wetting agents and mixtures thereof.

[0228] Excipients for preparation of compositions comprising a compound having Formula (I) to be administered orally in solid dosage form include, for example, agar, alginic acid, aluminum hydroxide, benzyl alcohol, benzyl benzoate, 1,3-butylene glycol, carbomers, castor oil, cellulose, cellulose acetate, cocoa butter, corn starch, corn oil, cottonseed oil, cross-povidone, diglycerides, ethanol, ethyl cellulose, ethyl laureate, ethyl oleate, fatty acid esters, gelatin, germ oil, glucose, glycerol, groundnut oil, hydroxypropylmethyl cellulose, isopropanol, isotonic saline, lactose, magnesium hydroxide, magnesium stearate, malt, mannitol, monoglycerides, olive oil, peanut oil, potassium phosphate salts, potato starch, povidone, propylene glycol, Ringer's solution, safflower oil, sesame oil, sodium carboxymethyl cellulose, sodium phosphate salts, sodium lauryl sulfate, sodium sorbitol, soybean oil, stearic acids, stearyl fumarate, sucrose, surfactants, talc, tragacanth, tetrahydrofurfuryl alcohol, triglycerides, water, and mixtures thereof. Excipients for preparation of compositions comprising a compound of this invention having Formula (I) to be administered ophthalmically or orally in liquid dosage forms include, for example, 1,3-butylene glycol, castor oil, corn oil, cottonseed oil, ethanol, fatty acid esters of

sorbitan, germ oil, groundnut oil, glycerol, isopropanol, olive oil, polyethylene glycols, propylene glycol, sesame oil, water and mixtures thereof. Excipients for preparation of compositions comprising a compound of this invention having Formula (I) to be administered osmotically include, for example, chlorofluorohydrocarbons, ethanol, water and mixtures thereof. Excipients for preparation of compositions comprising a compound of this invention having Formula (I) to be administered parenterally include, for example, 1,3-butanediol, castor oil, corn oil, cottonseed oil, dextrose, germ oil, groundnut oil, liposomes, oleic acid, olive oil, peanut oil, Ringer's solution, safflower oil, sesame oil, soybean oil, U.S.P. or isotonic sodium chloride solution, water and mixtures thereof. Excipients for preparation of compositions comprising a compound of this invention having Formula (I) to be administered rectally or vaginally include, for example, cocoa butter, polyethylene glycol, wax and mixtures thereof.

[0229] Compounds having Formula (I) are expected to be useful when used with alkylating agents, angiogenesis inhibitors, antibodies, antimetabolites, antimetotics, antiproliferatives, antivirals, aurora kinase inhibitors, other apoptosis promoters (for example, Bcl-xL, Bcl-w and Bfl-1) inhibitors, activators of death receptor pathway, Bcr-Abl kinase inhibitors, BiTE (Bi-Specific T cell Engager) antibodies, antibody drug conjugates, biologic response modifiers, cyclin-dependent kinase inhibitors, cell cycle inhibitors, cyclooxygenase-2 inhibitors, DVDs, leukemia viral oncogene homolog (ErbB2) receptor inhibitors, growth factor inhibitors, heat shock protein (HSP)-90 inhibitors, histone deacetylase (HDAC) inhibitors, hormonal therapies, immunologicals, inhibitors of inhibitors of apoptosis proteins (IAPs), intercalating antibiotics, kinase inhibitors, kinesin inhibitors, Jak2 inhibitors, mammalian target of rapamycin inhibitors, microRNA's, mitogen-activated extracellular signal-regulated kinase inhibitors, multivalent binding proteins, non-steroidal anti-inflammatory drugs (NSAIDs), poly ADP (adenosine diphosphate)-ribose polymerase (PARP) inhibitors, platinum chemotherapeutics, polo-like kinase (Plk) inhibitors, phosphoinositide-3 kinase (PI3K) inhibitors, proteasome inhibitors, purine analogs, pyrimidine analogs, receptor tyrosine kinase inhibitors, etinoids/deltoids plant alkaloids, small inhibitory ribonucleic acids (siRNAs), topoisomerase inhibitors, ubiquitin ligase inhibitors, and the like, and in combination with one or more of these agents.

[0230] BiTE antibodies are bi-specific antibodies that direct T-cells to attack cancer cells by simultaneously binding the two cells. The T-cell then attacks the target cancer cell. Examples of BiTE antibodies include adecatumumab (Micromet MT201), blinatumomab (Micromet MT103) and the like. Without being limited by theory, one of the mechanisms by which T-cells elicit apoptosis of the target cancer cell is by exocytosis of cytolytic granule components, which include perforin and granzyme B. In this regard, Bcl-2 has been shown to attenuate the induction of apoptosis by both perforin and granzyme B. These data suggest that inhibition of Bcl-2 could enhance the cytotoxic effects elicited by T-cells when targeted to cancer cells (V.R. Sutton, D.L. Vaux and J.A. Trapani, J. of Immunology 1997, 158 (12), 5783).

[0231] siRNAs are molecules having endogenous RNA bases or chemically modified nucleotides. The modifications do not abolish cellular activity, but rather impart increased stability and/or increased cellular potency. Examples of chemical modifications include phosphorothioate groups, 2'-deoxynucleotide, 2'-OCH₃-containing ribonucleotides, 2'-F-ribonucleotides, 2'-methoxyethyl ribonucleotides, combinations thereof and the like. The siRNA can have varying lengths (e.g., 10-200 bps) and structures (e.g., hairpins, single/double strands, bulges, nicks/gaps, mismatches) and are processed in cells to provide active gene silencing. A double-stranded siRNA (dsRNA) can have the same number of nucleotides on each strand (blunt ends) or asymmetric ends (overhangs). The overhang of 1-2 nucleotides can be present on the sense and/or the antisense strand, as well as present on the 5'- and/ or the 3'-ends of a given strand. For example, siRNAs targeting Mcl-1 have been shown to enhance the activity of ABT-263, (i.e., N-(4-(4-((2-(4-chlorophenyl)-5,5-dimethyl-1-cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(morpholin-4-yl)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide) or ABT-737 (i.e., N-(4-(4-((4'-chloro(1,1'-biphenyl)-2-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(dimethylamino)-1-((phenylsulfanyl)methyl)propyl)amino)-3-nitrobenzenesulfonamide) in multiple tumor cell lines (Tse et. al, Cancer Research 2008, 68(9), 3421 and references therein).

[0232] Multivalent binding proteins are binding proteins comprising two or more antigen binding sites. Multivalent binding proteins are engineered to have the three or more antigen binding sites and are generally not naturally occurring antibodies. The term "multispecific binding protein" means a binding protein capable of binding two or more related or unrelated targets. Dual variable domain (DVD) binding proteins are tetravalent or multivalent binding proteins binding proteins comprising two or more antigen binding sites. Such DVDs may be monospecific (i.e., capable of binding one antigen) or multispecific (i.e., capable of binding two or more antigens). DVD binding proteins comprising two heavy chain DVD polypeptides and two light chain DVD polypeptides are referred to as DVD Ig's. Each half of a DVD Ig comprises a heavy chain DVD polypeptide, a light chain DVD polypeptide, and two antigen binding sites. Each binding site comprises a heavy chain variable domain and a light chain variable domain with a total of 6 CDRs involved in antigen binding per antigen binding site.

[0233] Alkylating agents include altretamine, AMD-473, AP-5280, apaziquone, bendamustine, brostallicin, busulfan, carboquone, carmustine (BCNU), chlorambucil, CLORETAZINE® (laromustine, VNP 40101M), cyclophosphamide, decarbazine, estramustine, fotemustine, glufosfamide, ifosfamide, KW-2170, lomustine (CCNU), mafosfamide, melphalan, mitobronitol, mitolactol, nimustine, nitrogen mustard N-oxide, ranimustine, temozolomide, thiotepa, TREANDA® (bendamustine), treosulfan, rofosfamide and the like.

- [0234] Angiogenesis inhibitors include endothelial-specific receptor tyrosine kinase (Tie-2) inhibitors, epidermal growth factor receptor (EGFR) inhibitors, insulin growth factor-2 receptor (IGFR-2) inhibitors, matrix metalloproteinase-2 (MMP-2) inhibitors, matrix metalloproteinase-9 (MMP-9) inhibitors, platelet-derived growth factor receptor (PDGFR) inhibitors, thrombospondin analogs, vascular endothelial growth factor receptor tyrosine kinase (VEGFR) inhibitors and the like.
- 5 [0235] Antimetabolites include ALIMTA® (pemetrexed disodium, LY231514, MTA), 5-azacitidine, XELODA® (capecitabine), carmofur, LEUSTAT® (cladribine), clofarabine, cytarabine, cytarabine ocfosfate, cytosine arabinoside, decitabine, deferoxamine, doxifluridine, eflornithine, EICAR (5-ethynyl-1-β-D-ribofuranosylimidazole-4-carboxamide), enocitabine, ethynylcytidine, fludarabine, 5-fluorouracil alone or in combination with leucovorin, GEMZAR® (gemcitabine), hydroxyurea, ALKERAN® (melphalan), mercaptopurine, 6-mercaptopurine riboside, methotrexate, mycophenolic acid, nelarabine, nolatrexed, ocfosfate, pelitrexol, pentostatin, raltitrexed, Ribavirin, triapine, trimetrexate, S-1, tiazofurin, tegafur, TS-1, vidarabine, UFT and the like.
- 10 [0236] Antivirals include ritonavir, hydroxychloroquine and the like.
- [0237] Aurora kinase inhibitors include ABT-348, AZD-1152, MLN-8054, VX-680, Aurora A-specific kinase inhibitors, Aurora B-specific kinase inhibitors and pan-Aurora kinase inhibitors and the like.
- 15 [0238] Bcl-2 protein inhibitors include AT-101 ((-)-gossypol), GENASENSE® (G3139 or oblimersen (Bcl-2-targeting antisense oligonucleotide)), IPI-194, IPI-565, N-(4-(4-((4'-chloro(1,1'-biphenyl)-2-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(dimethylamino)-1-((phenylsulfanyl)methyl)propyl)amino)-3-nitrobenzenesulfonamide (ABT-737), N-(4-(4-((2-(4-chlorophenyl)-5,5-dimethyl-1-cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(morpholin-4-yl)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide (ABT-263), GX-070 (obatoclax) and the like.
- 20 [0239] Bcr-Abl kinase inhibitors include DASATINIB® (BMS-354825), GLEEVEC® (imatinib) and the like.
- [0240] CDK inhibitors include AZD-5438, BMI-1040, BMS-032, BMS-387, CVT-2584, flavopyridol, GPC-286199, MCS-5A, PD0332991, PHA-690509, seliciclib (CYC-202, R-roscovitine), ZK-304709 and the like.
- 25 [0241] COX-2 inhibitors include ABT-963, ARCOXIA® (etoricoxib), BEXTRA® (valdecoxib), BMS347070, CELEBREX® (celecoxib), COX-189 (lumiracoxib), CT-3, DERAMAXX® (deracoxib), JTE-522, 4-methyl-2-(3,4-dimethylphenyl)-1-(4-sulfamoylphenyl)-1H-pyrrole), MK-663 (etoricoxib), NS-398, parecoxib, RS-57067, SC-58125, SD-8381, SVT-2016, S-2474, T-614, VIOXX® (rofecoxib) and the like.
- [0242] EGFR inhibitors include ABX-EGF, anti-EGFR immunoliposomes, EGF-vaccine, EMD-7200, ERBITUX® (cetuximab), HR3, IgA antibodies, IRESSA® (gefitinib), TARCEVA® (erlotinib or OSI-774), TP-38, EGFR fusion protein, TYKERB® (lapatinib) and the like.
- 30 [0243] ErbB2 receptor inhibitors include CP-724-714, CI-1033 (canertinib), HERCEPTIN® (trastuzumab), TYKERB® (lapatinib), OMNITARG® (2C4, pertuzumab), TAK-165, GW-572016 (ionafarnib), GW-282974, EKB-569, PI-166, dHER2 (HER2 vaccine), APC-8024 (HER-2 vaccine), anti-HER/2neu bispecific antibody, B7.her2IgG3, AS HER2 trifunctional bispecific antibodies, mAB AR-209, mAB 2B-1 and the like.
- 35 [0244] Histone deacetylase inhibitors include depsipeptide, LAQ-824, MS-275, trapoxin, suberoylanilide hydroxamic acid (SAHA), TSA, valproic acid and the like.
- [0245] HSP-90 inhibitors include 17-AAG-nab, 17-AAG, CNF-101, CNF-1010, CNF-2024, 17-DMAG, geldanamycin, IPI-504, KOS-953, MYCOGRAB® (human recombinant antibody to HSP-90), NCS-683664, PU24FCI, PU-3, radicicol, SNX-2112, STA-9090 VER49009 and the like.
- 40 [0246] Inhibitors of inhibitors of apoptosis proteins include HGS1029, GDC-0145, GDC-0152, LCL-161, LBW-242 and the like.
- [0247] Antibody drug conjugates include anti-CD22-MC-MMAF, anti-CD22-MC-MMAE, anti-CD22-MCC-DM1, CR-011-vcMMAE, PSMA-ADC, MEDI-547, SGN-19Am SGN-35, SGN-75 and the like
- 45 [0248] Activators of death receptor pathway include TRAIL, antibodies or other agents that target TRAIL or death receptors (e.g., DR4 and DR5) such as Apomab, conatumumab, ETR2-ST01, GDC0145, (lexatumumab), HGS-1029, LBY-135, PRO-1762 and trastuzumab.
- [0249] Kinesin inhibitors include Eg5 inhibitors such as AZD4877, ARRY-520; CENPE inhibitors such as GSK923295A and the like.
- 50 [0250] JAK-2 inhibitors include CEP-701 (lesaurtinib), XL019 and INCB018424 and the like.
- [0251] MEK inhibitors include ARRY-142886, ARRY-438162 PD-325901, PD-98059 and the like.
- [0252] mTOR inhibitors include AP-23573, CCI-779, everolimus, RAD-001, rapamycin, temsirolimus, ATP-competitive TORC1/TORC2 inhibitors, including PI-103, PP242, PP30, Torin 1 and the like.
- 55 [0253] Non-steroidal anti-inflammatory drugs include AMIGESIC® (salsalate), DOLOBID® (diflunisal), MOTRIN® (ibuprofen), ORUDIS® (ketoprofen), RELAFEN® (nabumetone), FELDENE® (piroxicam), ibuprofen cream, ALEVE® (naproxen) and NAPROSYN® (naproxen), VOLTAREN® (diclofenac), INDOCIN® (indomethacin), CLINORIL® (sulindac), TOLECTIN® (tolmetin), LODINE® (etodolac), TORADOL® (ketorolac), DAYPRO® (oxaprozin) and the like.
- [0254] PDGFR inhibitors include C-451, CP-673, CP-868596 and the like.
- [0255] Platinum chemotherapeutics include cisplatin, ELOXATIN® (oxaliplatin) eptaplatin, lobaplatin, nedaplatin, PAR-

APLATIN® (carboplatin), satraplatin, picoplatin and the like.

[0256] Polo-like kinase inhibitors include BI-2536 and the like.

[0257] Phosphoinositide-3 kinase (PI3K) inhibitors include wortmannin, LY294002, XL-147, CAL-120, ONC-21, AEZS-127, ETP-45658, PX-866, GDC-0941, BGT226, BEZ235, XL765 and the like.

5 **[0258]** Thrombospondin analogs include ABT-510, ABT-567, ABT-898, TSP-1 and the like.

[0259] VEGFR inhibitors include AVASTIN® (bevacizumab), ABT-869, AEE-788, ANGIOZYME™ (a ribozyme that inhibits angiogenesis (Ribozyme Pharmaceuticals (Boulder, CO.) and Chiron, (Emeryville, CA)), axitinib (AG-13736), AZD-2171, CP-547,632, IM-862, MACUGEN (pegaptamib), NEXAVAR® (sorafenib, BAY43-9006), pazopanib (GW-786034), vatalanib (PTK-787, ZK-222584), SUTENT® (sunitinib, SU-11248), VEGF trap, ZACTIMA™ (vandetanib, ZD-6474) and the like.

10 **[0260]** Antibiotics include intercalating antibiotics aclarubicin, actinomycin D, amrubicin, annamycin, adriamycin, BLENOXANE® (bleomycin), daunorubicin, CAELYX® or MYOCET® (liposomal doxorubicin), elsamitrucin, epirubicin, glarubicin, ZAVEDOS® (idarubicin), mitomycin C, nemorubicin, neocarzinostatin, peplomycin, pirarubicin, rebeccamycin, stimalamer, streptozocin, VALSTAR® (valrubicin), zinostatin and the like.

15 **[0261]** Topoisomerase inhibitors include aclarubicin, 9-aminocamptothecin, amonafide, amsacrine, becatecarin, belotecan, BN-80915, CAMPTOSAR® (irinotecan hydrochloride), camptothecin, CARDIOXANE® (dexrazoxine), diflomotecan, edotecarin, ELLENCE® or PHARMORUBICIN® (epirubicin), etoposide, exatecan, 10-hydroxycamptothecin, gimatecan, lurtotecan, mitoxantrone, orathecin, pirarubicin, pixantrone, rubitecan, sobuzoxane, SN-38, tafluposide, topotecan and the like.

20 **[0262]** Antibodies include AVASTIN® (bevacizumab), CD40-specific antibodies, chTNT-1/B, denosumab, ERBITUX® (cetuximab), HUMAX-CD4® (zanolimumab), IGF1R-specific antibodies, lintuzumab, PANOREX® (edrecolomab), RENCAREX® (WX G250), RITUXAN® (rituximab), ticilimumab, trastuzumab, CD20 antibodies types I and II and the like.

[0263] Hormonal therapies include ARIMIDEX® (anastrozole), AROMASIN® (exemestane), arzoxifene, CASODEX® (bicalutamide), CETROTIDE® (cetorelix), degarelix, deslorelin, DESOPAN® (trilostane), dexamethasone, DROGENIL® (flutamide), EVISTA® (raloxifene), AFEMA™ (fadrozole), FARESTON® (toremifene), FASLODEX® (fulvestrant), FEMARA® (letrozole), formestane, glucocorticoids, HECTOROL® (doxercalciferol), RENAGEL® (sevelamer carbonate), lasofoxifene, leuprolide acetate, MEGACE® (megesterol), MIFEPREX® (mifepristone), NILANDRON™ (nilutamide), NOLVADEX® (tamoxifen citrate), PLENAXIS™ (abarelix), prednisone, PROPECIA® (finasteride), rilostane, SUPREFACT® (buserelin), TRELSTAR® (luteinizing hormone releasing hormone (LHRH)), VANTAS® (Histrelin implant), VETORYL® (trilostane or modrastane), ZOLADEX® (fosreltin, goserelin) and the like.

30 **[0264]** Deltoids and retinoids include seocalcitol (EB 1089, CB 1093), lexacalcitol (KH1060), fenretinide, PANRETIN® (aliretinoin), ATRAGEN® (liposomal tretinoin), TARGRETIN® (bexarotene), LGD-1550 and the like.

[0265] PARP inhibitors include ABT-888 (veliparib), olaparib, KU-59436, AZD-2281, AG-014699, BSI-201, BGP-15, INO-1001, ONO-2231 and the like.

35 **[0266]** Plant alkaloids include, but are not limited to, vincristine, vinblastine, vindesine, vinorelbine and the like.

[0267] Proteasome inhibitors include VELCADE® (bortezomib), MG132, NPI-0052, PR-171 and the like.

[0268] Examples of immunologicals include interferons and other immune-enhancing agents. Interferons include interferon alpha, interferon alpha-2a, interferon alpha-2b, interferon beta, interferon gamma-1a, ACTIMMUNE® (interferon gamma-1b) or interferon gamma-n1, combinations thereof and the like. Other agents include ALFAFERONE® (IFN-α), BAM-002 (oxidized glutathione), BEROMUN® (tasonermin), BEXXAR® (tositumomab), CAMPATH® (alemtuzumab), CTLA4 (cytotoxic lymphocyte antigen 4), decarbazine, denileukin, epratuzumab, GRANOCYTE® (lenograstim), lentinan, leukocyte alpha interferon, imiquimod, MDX-010 (anti-CTLA-4), melanoma vaccine, mitumomab, molgramostim, MYLOTARG™ (gemtuzumab ozogamicin), NEUPOGEN® (filgrastim), OncoVAC-CL, OVAREX® (oregovomab), pemtumorab (Y-muHMF1), PROVENGE® (sipuleucel-T), sargaramostim, sizofilan, teceleukin, THERACYS® (Bacillus Calmette-Guerin), ubenimex, VIRULIZIN® (immunotherapeutic, Lorus Pharmaceuticals), Z-100 (Specific Substance of Maruyama (SSM)), WF-10 (Tetrachlorodecaoxide (TCDO)), PROLEUKIN® (aldesleukin), ZADAXIN® (thymalfasin), ZENAPAX® (daclizumab), ZEVALIN® (90Y-Ibritumomab tiuxetan) and the like.

[0269] Biological response modifiers are agents that modify defense mechanisms of living organisms or biological responses, such as survival, growth or differentiation of tissue cells to direct them to have anti-tumor activity and include krestin, lentinan, sizofiran, picibanil PF-3512676 (CpG-8954), ubenimex and the like.

50 **[0270]** Pyrimidine analogs include cytarabine (ara C or Arabinoside C), cytosine arabinoside, doxifluridine, FLUDARA® (fludarabine), 5-FU (5-fluorouracil), floxuridine, GEMZAR® (gemcitabine), TOMUDEX® (ratitrexed), TROXATYL™ (triacetyluridine troxacitabine) and the like.

[0271] Purine analogs include LANVIS® (thioguanine) and PURI-NETHOL® (mercaptopurine).

55 **[0272]** Antimitotic agents include batabulin, epothilone D (KOS-862), N-(2-((4-hydroxyphenyl)amino)pyridin-3-yl)-4-methoxybenzenesulfonamide, ixabepilone (BMS 247550), paclitaxel, TAXOTERE® (docetaxel), PNU100940 (109881), patupilone, XRP-9881 (larotaxel), vinflunine, ZK-EPO (synthetic epothilone) and the like.

[0273] Ubiquitin ligase inhibitors include MDM2 inhibitors, such as nutlins, NEDD8 inhibitors such as MLN4924 and

the like.

[0274] Compounds of this invention can also be used as radiosensitizers that enhance the efficacy of radiotherapy. Examples of radiotherapy include external beam radiotherapy, teletherapy, brachytherapy and sealed, unsealed source radiotherapy and the like.

[0275] Additionally, compounds having Formula (I) may be combined with other chemotherapeutic agents such as ABRAXANE™ (ABI-007), ABT-100 (farnesyl transferase inhibitor), ADVEXIN® (Ad5CMV-p53 vaccine), ALTOCOR® or MEVACOR® (lovastatin), AMPLIGEN® (poly I:poly C12U, a synthetic RNA), APTOSYN® (exisulind), AREDIA® (pamidronic acid), arglabin, L-asparaginase, atamestane (1-methyl-3,17-dione-androsta-1,4-diene), AVAGE® (tazarotene), AVE-8062 (combrestatin derivative) BEC2 (mitumomab), cachectin or cachexin (tumor necrosis factor), canvaxin (vaccine), CEAVAC® (cancer vaccine), CELEUK® (celmoleukin), CEPLENE® (histamine dihydrochloride), CERVARIX® (human papillomavirus vaccine), CHOP® (C: CYTOXAN® (cyclophosphamide); H: ADRIAMYCIN® (hydroxydoxorubicin); O: Vincristine (ONCOVIN®); P: prednisone), CYPAT™ (cyproterone acetate), combrestatin A4P, DAB(389)EGF (catalytic and translocation domains of diphtheria toxin fused via a His-Ala linker to human epidermal growth factor) or TransMID-107R™ (diphtheria toxins), dacarbazine, dactinomycin, 5,6-dimethylxanthenone-4-acetic acid (DMXAA), eniluracil, EVI-ZON™ (squalamine lactate), DIMERICINE® (T4N5 liposome lotion), discodermolide, DX-8951f (exatecan mesylate), enzastaurin, EPO906 (epithilone B), GARDASIL® (quadrivalent human papillomavirus (Types 6, 11, 16, 18) recombinant vaccine), GASTRIMMUNE®, GENASENSE®, GMK (ganglioside conjugate vaccine), GVAX® (prostate cancer vaccine), halofuginone, histerelin, hydroxycarbamide, ibandronic acid, IGN-101, IL-13-PE38, IL-13-PE38QQR (cintredekin besudotox), IL-13-pseudomonas exotoxin, interferon- α , interferon- γ , JUNOVAN™ or MEPACT™ (mifamurtide), lonafarnib, 5,10-methylenetetrahydrofolate, miltefosine (hexadecylphosphocholine), NEOVASTAT® (AE-941), NEUTREXIN® (trimetrexate glucuronate), NIPENT® (pentostatin), ONCONASE® (a ribonuclease enzyme), ONCOPHAGE® (melanoma vaccine treatment), ONCOVAX® (IL-2 Vaccine), ORATHECIN™ (rubitecan), OSIDEM® (antibody-based cell drug), OVAREX® Mab (murine monoclonal antibody), paclitaxel, PANDIMEX™ (aglycone saponins from ginseng comprising 20(S)protopanaxadiol (aPPD) and 20(S)protopanaxatriol (aPTT)), panitumumab, PANVAC®-VF (investigational cancer vaccine), pegaspargase, PEG Interferon A, phenoxodiol, procarbazine, rebimastat, REMOVAB® (catumaxomab), REV-LIMID® (lenalidomide), RSR13 (efaproxiral), SOMATULINE® LA (lanreotide), SORIATANE® (acitretin), staurosporine (Streptomyces staurospores), talabostat (PT100), TARGRETIN® (bexarotene), TAXOPREXIN® (DHA-paclitaxel), TEL-CYTA® (canfosfamide, TLK286), temilifene, TEMODAR® (temozolomide), tesmilifene, thalidomide, THERATOPE® (STn-KLH), thymitaq (2-amino-3,4-dihydro-6-methyl-4-oxo-5-(4-pyridylthio)quinazoline dihydrochloride), TNFERADE™ (adenovector: DNA carrier containing the gene for tumor necrosis factor- α), TRACLEER® or ZAVESCA® (bosentan), tretinoin (Retin-A), tetrandrine, TRISENOX® (arsenic trioxide), VIRULIZIN®, ukrain (derivative of alkaloids from the greater celandine plant), vitaxin (anti-alphavbeta3 antibody), XCYTRIN® (motexafin gadolinium), XINLAY™ (atrasentan), XYOTAX™ (paclitaxel poliglumex), YONDELIS® (trabectedin), ZD-6126, ZINECARD® (dexrazoxane), ZOMETA® (zoledronic acid), zorubicin and the like.

Data

[0276] Determination of the utility of compounds having Formula (I) as binders to and inhibitors of anti-apoptotic Bcl-2 proteins was performed using the Time Resolved-Fluorescence Resonance Energy Transfer (TR-FRET) Assay. Tb-anti-GST antibody was purchased from Invitrogen (Catalog No. PV4216).

Probe Synthesis

[0277] All reagents were used as obtained from the vendor unless otherwise specified. Peptide synthesis reagents including diisopropylethylamine (DIEA), dichloromethane (DCM), N-methylpyrrolidone (NMP), 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU), N-hydroxybenzotriazole (HOBt) and piperidine were obtained from Applied Biosystems, Inc. (ABI), Foster City, CA or American Bioanalytical, Natick, MA. Preloaded 9-Fluorenylmethyloxycarbonyl (Fmoc) amino acid cartridges (Fmoc-Ala-OH, Fmoc-Cys(Trt)-OH, Fmoc-Asp(tBu)-OH, Fmoc-Glu(tBu)-OH, Fmoc-Phe-OH, Fmoc-Gly-OH, Fmoc-His(Trt)-OH, Fmoc-Ile-OH, Fmoc-Leu-OH, Fmoc-Lys(Boc)-OH, Fmoc-Met-OH, Fmoc-Asn(Trt)-OH, Fmoc-Pro-OH, Fmoc-Gln(Trt)-OH, Fmoc-Arg(Pbf)-OH, Fmoc-Ser(tBu)-OH, Fmoc-Thr(tBu)-OH, Fmoc-Val-OH, Fmoc-Trp(Boc)-OH, Fmoc-Tyr(tBu)-OH) were obtained from ABI or Anaspec, San Jose, CA. The peptide synthesis resin (Fmoc-Rink amide MBHA resin) and Fmoc-Lys(Mtt)-OH were obtained from Novabiochem, San Diego, CA. Single-isomer 6-carboxyfluorescein succinimidyl ester (6-FAM-NHS) was obtained from Anaspec. Trifluoroacetic acid (TFA) was obtained from Oakwood Products, West Columbia, SC. Thioanisole, phenol, triisopropylsilane (TIS), 3,6-dioxo-1,8-octanedithiol (DODT) and isopropanol were obtained from Aldrich Chemical Co., Milwaukee, WI. Matrix-assisted laser desorption ionization mass-spectra (MALDI-MS) were recorded on an Applied Biosystems Voyager DE-PRO MS. Electrospray mass-spectra (ESI-MS) were recorded on Finnigan SSQ7000 (Finnigan Corp., San Jose, CA) in both positive and negative ion mode.

General Procedure For Solid-Phase Peptide Synthesis (SPPS)

5 [0278] Peptides were synthesized with, at most, 250 μ mol preloaded Wang resin/vessel on an ABI 433A peptide synthesizer using 250 μ mol scale Fastmoc™ coupling cycles. Preloaded cartridges containing 1 mmol standard Fmoc-amino acids, except for the position of attachment of the fluorophore, where 1 mmol Fmoc-Lys(Mtt)-OH was placed in the cartridge, were used with conductivity feedback monitoring. N-terminal acetylation was accomplished by using 1 mmol acetic acid in a cartridge under standard coupling conditions.

10 *Removal Of 4-Methyltrityl (Mtt) From Lysine*

[0279] The resin from the synthesizer was washed thrice with DCM and kept wet. 150 mL of 95:4:1 dichloromethane:triisopropylsilane:trifluoroacetic acid was flowed through the resin bed over 30 minutes. The mixture turned deep yellow then faded to pale yellow. 100 mL of DMF was flowed through the bed over 15 minutes. The resin was then washed thrice with DMF and filtered. Ninhydrin tests showed a strong signal for primary amine.

15 *Resin Labeling With 6-Carboxyfluorescein-NHS (6-FAM-NHS)*

[0280] The resin was treated with 2 equivalents 6-FAM-NHS in 1% DIEA/DMF and stirred or shaken at ambient temperature overnight. When complete, the resin was drained, washed thrice with DMF, thrice with (1 $\times$ DCM and 1 $\times$ methanol) and dried to provide an orange resin that was negative by ninhydrin test.

General Procedure For Cleavage And Deprotection Of Resin-Bound Peptide

25 [0281] Peptides were cleaved from the resin by shaking for 3 hours at ambient temperature in a cleavage cocktail consisting of 80% TFA, 5% water, 5% thioanisole, 5% phenol, 2.5% TIS, and 2.5% EDT (1 mL/0.1 g resin). The resin was removed by filtration and rinsing twice with TFA. The TFA was evaporated from the filtrates, and product was precipitated with ether (10 mL/0.1 g resin), recovered by centrifugation, washed twice with ether (10 mL/0.1 g resin) and dried to give the crude peptide.

30 *General Procedure For Purification Of Peptides*

[0282] The crude peptides were purified on a Gilson preparative HPLC system running Unipoint® analysis software (Gilson, Inc., Middleton, WI) on a radial compression column containing two 25 $\times$ 100 mm segments packed with Delta-Pak™ C18 15 μ m particles with 100 Å pore size and eluted with one of the gradient methods listed below. One to two milliliters of crude peptide solution (10 mg/mL in 90% DMSO/water) was purified per injection. The peaks containing the product(s) from each run were pooled and lyophilized. All preparative runs were run at 20 mL/min with eluents as buffer A: 0.1% TFA-water and buffer B: acetonitrile.

40 *General Procedure For Analytical HPLC*

[0283] Analytical HPLC was performed on a Hewlett-Packard 1200 series system with a diode-array detector and a Hewlett-Packard 1046A fluorescence detector running HPLC 3D ChemStation software version A.03.04 (Hewlett-Packard, Palo Alto, CA) on a 4.6 $\times$ 250 mm YMC column packed with ODS-AQ 5 μ m particles with a 120 Å pore size and eluted with one of the gradient methods listed below after preequilibrating at the starting conditions for 7 minutes. Eluents were buffer A: 0.1% TFA-water and buffer B: acetonitrile. The flow rate for all gradients was 1 mL/min.

F-Bak: Peptide Probe Acetyl-(SEQ ID NO: 1)GQVGRQLAIIQDK(6-FAM)-(SEQ ID NO: 2)INR-NH₂

50 [0284] Fmoc-Rink amide MBHA resin was extended using the general peptide synthesis procedure to provide the protected resin-bound peptide (1.020 g). The Mtt group was removed, labeled with 6-FAM-NHS and cleaved and deprotected as described hereinabove to provide the crude product as an orange solid (0.37 g). This product was purified by RP-HPLC. Fractions across the main peak were tested by analytical RP-HPLC, and the pure fractions were isolated and lyophilized, with the major peak providing the title compound (0.0802 g) as a yellow solid; MALDI-MS m/z = 2137.1 [(M+H)⁺].

55

Alternative Synthesis of Peptide Probe F-Bak: Acetyl-(SEQ ID NO: 1)GQVGRQLAIIGDK(6-FAM)-(SEQ ID NO:2)INR-NH₂

[0285] The protected peptide was assembled on 0.25 mmol Fmoc-Rink amide MBHA resin (Novabiochem) on an Applied Biosystems 433A automated peptide synthesizer running Fastmoc™ coupling cycles using pre-loaded 1 mmol amino acid cartridges, except for the fluorescein(6-FAM)-labeled lysine, where 1 mmol Fmoc-Lys(4-methyltrityl) was weighed into the cartridge. The N-terminal acetyl group was incorporated by putting 1 mmol acetic acid in a cartridge and coupling as described hereinabove. Selective removal of the 4-methyltrityl group was accomplished with a solution of 95:4:1 DCM:TIS:TFA (v/v/v) flowed through the resin over 15 minutes, followed by quenching with a flow of dimethylformamide. Single-isomer 6-carboxyfluorescein-NHS was reacted with the lysine side-chain in 1% DIEA in DMF and confirmed complete by ninhydrin testing. The peptide was cleaved from the resin and side-chains deprotected by treating with 80:5:5:5:2.5:2.5 TFA/water/phenol/ thioanisole/triisopropylsilane: 3,6-dioxa-1,8-octanedithiol (v/v/v/v/v/v), and the crude peptide was recovered by precipitation with diethyl ether. The crude peptide was purified by reverse-phase high-performance liquid chromatography, and its purity and identity were confirmed by analytical reverse-phase high-performance liquid chromatography and matrix-assisted laser-desorption mass-spectrometry ($m/z = 2137.1 ((M+H)^+)$).

Time Resolved-Fluorescence Resonance Energy Transfer (TR-FRET) Assay

[0286] Representative compounds were serially diluted in dimethyl sulfoxide (DMSO) starting at 50 μ M (2x starting concentration; 10% DMSO) and 10 μ L were transferred into a 384-well plate. Then 10 μ L of a protein/probe/antibody mix was added to each well at final concentrations listed in TABLE 1. The samples are then mixed on a shaker for 1 minute and incubated for an additional 3 hours at room temperature. For each assay, the probe/antibody and protein/probe/antibody were included on each assay plate as negative and positive controls, respectively. Fluorescence was measured on the Envision (Perkin Elmer) using a 340/35 nm excitation filter and 520/525 (F-Bak peptide) and 495/510 nm (Tb-labeled anti-Histidine antibody) emission filters. Dissociation constants (K_d) are shown in TABLE 2 below and were determined using Wang's equation (Wang Z.-X., An Exact Mathematical Expression For Describing Competitive Binding Of Two Different Ligands To A Protein Molecule. FEBS Lett. 1995,360:111-4).

TABLE 1. Protein, Probe And Antibody Used For TR-FRET Assays

Protein	Probe	Protein (nM)	Probe (nM)	Antibody	Antibody (nM)
GST-Bcl-2	F-Bak Peptide Probe Acetyl-(SEQ ID NO: 1 GQVGRQLAIIGDK(6-FAM) SEQ ID NO: 2 INR-amide)	1	100	Tb-anti-GST	1
6-FAM = 6- carboxyfluorescein.; Tb = terbium; GST = glutathione S-transferase					

[0287] The samples were then mixed on a shaker for 1 minute and incubated for an additional 3 hours at room temperature. For each assay, the probe/antibody and protein/probe/antibody were included on each assay plate as negative and positive controls, respectively. Fluorescence was measured on the Envision (Perkin Elmer) using a 340/35 nm excitation filter and 520/525 (F-Bak peptide) and 495/510 nm (Tb-labeled anti-Histidine antibody) emission filters.

[0288] Inhibition constants (K_i) for compounds according to the invention are shown in TABLE 2 below. Where the K_i for a compound is represented as "<" (less than) a certain numerical value, it is intended to mean that the binding affinity value (e.g., for Bcl-2) is lower than the limit of detection of the assay used. Inhibition constants were determined using Wang's equation (Wang Zx., An Exact Mathematical Expression For Describing Competitive Binding Of Two Different Ligands To A Protein Molecule. FEBS Lett. 1995, 360:111-4).

TABLE 2. TR-FRET Bcl-2 Binding K_i (μ M)

EXAMPLE #	K_i	EXAMPLE #	K_i
1	0.000225	190	0.000026
2	< 0.000010	191	< 0.000010
3	0.000013	192	< 0.000010
4	< 0.000010	193	< 0.000010
5	< 0.000010	194	< 0.000010
6	0.000018	195	< 0.000010

EP 2 944 638 A1

(continued)

EXAMPLE #	Ki	EXAMPLE #	Ki
7	0.00492	196	< 0.000010
8	0.000153	197	< 0.000010
9	< 0.000010	198	< 0.000010
10	< 0.000010	199	< 0.000010
11	0.000016	200	< 0.000010
12	< 0.000010	201	0.000014
13	< 0.000010	202	< 0.000010
14	0.002798	203	< 0.000010
15	< 0.000010	204	< 0.000010
16	0.000219	205	< 0.000010
17	0.00009	206	0.000036
18	0.000017	207	0.00003
19	0.000226	208	0.000104
20	0.000181	209	< 0.000010
21	0.000912	210	0.000011
22	0.000291	211	0.000058
23	0.000083	212	0.0001330
24	< 0.000010	213	< 0.000010
25	< 0.000010	214	< 0.000010
26	0.000011	215	< 0.000010
27	0.000134	216	< 0.000010
28	< 0.000010	217	< 0.000010
29	< 0.000010	218	0.000013
30	< 0.000010	219	0.001192
31	< 0.000010	220	0.000988
32	< 0.000010	221	0.000049
33	< 0.000010	222	0.000938
34	0.00001	223	0.000053
35	< 0.000010	224	< 0.000010
36	0.000017	225	0.000196
37	< 0.000010	226	0.000139
38	0.0003	227	< 0.000010
39	0.000012	228	0.026761
40	< 0.000010	229	0.002109
41	< 0.000010	230	0.000031
42	0.000439	231	0.000770
43	0.000012	232	0.001631
44	< 0.000010	233	0.001654

EP 2 944 638 A1

(continued)

EXAMPLE #	Ki	EXAMPLE #	Ki
45	< 0.000010	234	0.000115
46	0.000935	235	0.000023
47	< 0.000010	236	0.000033
48	< 0.000010	237	0.000024
49	0.000074	238	< 0.000010
50	0.000021	239	0.000026
51	< 0.000010	240	< 0.000010
52	0.000114	241	< 0.000010
53	< 0.000010	242	0.000057
54	0.002071	243	0.000546
55	< 0.000010	244	0.000281
56	0.000037	245	0.000015
57	0.000063	246	0.000144
58	< 0.000010	247	0.000019
59	0.000203	248	0.000029
60	< 0.000010	250	0.000412
61	0.000091	251	0.000571
62	< 0.000010	252	< 0.000010
63	< 0.000010	253	0.000052
64	< 0.000010	254	< 0.000010
65	< 0.000010	255	< 0.000010
66	< 0.000010	256	< 0.000010
67	< 0.000010	257	0.000052
68	0.000012	258	< 0.000010
69	0.001157	259	< 0.000010
70	0.003964	260	0.000016
71	0.00001	261	0.000134
72	< 0.000010	262	< 0.000010
73	< 0.000010	263	0.000156
74	0.000029	264	0.000036
75	< 0.000010	265	< 0.000010
76	0.000196	266	< 0.000010
77	0.000213	267	0.000035
78	< 0.000010	268	< 0.000010
79	< 0.000010	269	0.000016
80	< 0.000010	270	< 0.000010
81	< 0.000010	271	0.000039
82	0.000328	272	0.000031

EP 2 944 638 A1

(continued)

EXAMPLE #	Ki	EXAMPLE #	Ki
83	0.000071	273	0.000035
84	0.000123	274	0.000040
85	0.000391	275	< 0.000010
86	0.000498	276	< 0.000010
87	0.000618	277	< 0.000010
88	0.000672	278	0.000252
89	0.000073	279	0.000035
90	0.000013	280	0.000071
91	0.000487	281	0.000145
92	0.000128	282	< 0.000010
93	0.003461	283	< 0.000010
94	0.000678	284	0.000024
95	0.000014	285	< 0.000010
96	0.000014	286	< 0.000010
97	0.000017	287	0.000081
98	< 0.000010	288	0.000251
99	0.000233	289	0.000090
100	< 0.000010	290	< 0.000010
101	0.000021	291	< 0.000010
102	0.000094	292	0.000190
103	< 0.000010	293	0.000093
104	0.000016	294	0.000046
105	< 0.000010	295	< 0.000010
106	0.000895	296	0.000512
107	0.000035	297	0.000174
108	< 0.000010	298	< 0.000010
109	0.000127	299	0.000039
110	0.000557	300	0.001627
111	< 0.000010	301	0.002065
112	< 0.000010	302	0.000332
113	< 0.000010	303	0.000044
114	< 0.000010	304	nd
115	< 0.000010	305	0.000033
116	< 0.000010	306	0.002067
117	< 0.000010	307	0.000130
118	< 0.000010	308	0.000141
119	< 0.000010	309	0.000023
120	< 0.000010	310	0.000165

EP 2 944 638 A1

(continued)

5

10

15

20

25

30

35

40

45

50

55

EXAMPLE #	Ki	EXAMPLE #	Ki
121	< 0.000010	311	< 0.000010
122	< 0.000010	312	< 0.000010
123	< 0.000010	313	0.001102
124	< 0.000010	314	0.000042
125	< 0.000010	315	0.000052
126	< 0.000010	316	0.000601
127	< 0.000010	317	< 0.000010
128	< 0.000010	318	< 0.000010
129	0.000002	319	< 0.000010
130	< 0.000010	320	< 0.000010
131	< 0.000010	321	< 0.000010
132	< 0.000010	322	< 0.000010
133	< 0.000010	323	0.000104
134	< 0.000010	324	< 0.000010
135	< 0.000010	325	< 0.000010
136	< 0.000010	326	< 0.000010
137	< 0.000010	327	< 0.000010
138	< 0.000010	328	< 0.000010
139	< 0.000010	329	0.000030
140	< 0.000010	330	< 0.000010
141	< 0.000010	331	0.001086
142	0.00013	332	0.000621
143	< 0.000010	333	0.000511
144	< 0.000010	334	0.000572
145	< 0.000010	335	0.000150
146	< 0.000010	336	0.000198
147	< 0.000010	337	< 0.000010
148	< 0.000010	338	0.000013
149	< 0.000010	339	0.000036
150	< 0.000010	340	< 0.000010
151	0.000017	341	< 0.000010
152	< 0.000010	342	< 0.000010
153	< 0.000010	343	< 0.000010
154	< 0.000010	344	< 0.000010
155	0.000059	345	< 0.000010
156	< 0.000010	346	0.000042
157	< 0.000010	347	0.000013
158	< 0.000010	348	0.000034

EP 2 944 638 A1

(continued)

EXAMPLE #	K _i	EXAMPLE #	K _i
159	< 0.000010	349	0.000023
160	< 0.000010	350	< 0.000010
161	< 0.000010	351	< 0.000010
162	< 0.000010	352	0.000014
163	< 0.000010	353	< 0.000010
164	< 0.000010	354	0.000010
165	< 0.000010	355	0.000014
166	< 0.000010	356	0.000039
167	< 0.000010	357	< 0.000010
168	< 0.000010	358	< 0.000010
169	0.000021	359	< 0.000010
170	0.000022	360	< 0.000010
171	< 0.000010	361	< 0.000010
172	< 0.000010	362	0.000016
173	< 0.000010	363	0.000017
174	< 0.000010	364	< 0.000010
175	0.000119	365	< 0.000010
176	0.000023	366	0.000024
177	0.000111	367	nd
178	0.000076	368	nd
179	< 0.000010	369	< 0.000010
180	< 0.000010	370	0.000285
181	0.000017	371	< 0.0000010
182	0.000068	372	nd
183	< 0.000010	373	< 0.0000010
184	< 0.000010	374	< 0.0000010
185	0.000022	375	0.00010999
186	0.000047	376	< 0.0000010
187	0.00008	377	< 0.0000010
188	< 0.000010	378	< 0.0000010
189	0.000018		
nd = not determined			

The inhibition constant (K_i) is the dissociation constant of an enzyme-inhibitor complex or a protein/small molecule complex, wherein the small molecule is inhibiting binding of one protein to another protein or peptide. So a large K_i value indicates a low binding affinity and a small K_i value indicates a high binding affinity.

[0289] TABLE 2 shows inhibition constants for the inhibition of a Bak BH3 peptide probe to Bcl-2 protein and indicate that compounds according to the invention have high binding affinities for anti-apoptotic Bcl-2 protein. The compounds are therefore expected to have utility in treatment of diseases during which anti-apoptotic Bcl-2 protein is expressed.

RS4;11 Cell Viability Assay

[0290] The acute lymphoblastic leukemia (ALL) cell line RS4;11 was used as the primary human cell line to assess

the cellular activity of Bcl-2 selective agents in vitro and their efficacy in vivo. Previous studies have shown by BH3 profiling, a mitochondrial assay that classifies blocks in the intrinsic apoptotic pathway, that RS4;11 cells were highly dependant on BCL-2 for survival and sensitive to the Bcl-2 family member inhibitor ABT-737 (Blood, 2008, Vol. 111, 2300-2309). The prevalence of Bcl-2 complexed to the proapoptotic BH3 protein Bim in RS4;11 suggests that these cells are "primed" or more susceptible to cell death by antagonism of the antiapoptotic protein Bcl-2 for which they depend on for survival.

[0291] RS4;11 cells were cultured in RPMI-1640 supplemented with 2 mM L-glutamine, 10% FBS, 1 mM sodium pyruvate, 2 mM HEPES, 1% penicillin/streptomycin (Invitrogen), 4.5 g/L glucose and maintained at 37 C containing 5% CO₂. To test for the cellular activity of compounds in vitro, cells were treated at 50,000 cells per well in 96-well microtiter plates in the presence of 10% human serum for 48 hours in a humidified chamber with 5% CO₂. Cell cytotoxicity EC₅₀ values were assessed using CellTiter Glo (Promega) according to the manufacturer's recommendations. The EC₅₀ values were determined as a percentage of viable cells following treatment compared to the untreated control cells.

Table 3. RS4;11 EC₅₀ Values (μM)

EXAMPLE #	EC50	EXAMPLE #	EC50
1	0.712	190	0.55068
2	0.783	191	0.00691
3	0.0142	192	0.00241
4	0.01854	193	0.00076
5	0.01241	194	0.00819
6	0.03487	195	0.00207
7	0.192	196	0.00172
8	0.158	197	0.0125
9	0.01476	198	0.03619
10	0.05202	199	0.00506
11	0.01393	200	0.01099
12	0.03471	201	0.59132
13	0.0232	202	0.0438
14	3.8947	203	0.02208
15	0.01276	204	0.16475
16	1.2098	205	0.01059
17	0.475	206	0.05291
18	0.086	207	0.00376
19	0.465	208	0.12121
20	0.191	209	0.0045
21	0.062	210	0.06022
22	0.085	211	0.3073
23	0.045	212	0.01283
24	0.00983	213	0.0060976
25	0.007	214	0.0043751
26	0.05888	215	0.00056038
27	0.33237	216	0.68263

EP 2 944 638 A1

(continued)

	EXAMPLE #	EC50	EXAMPLE #	EC50
5	28	0.0419	217	0.0015528
	29	0.02047	218	0.0072907
	30	0.01529	219	> 1
	31	0.01565	220	> 1
10	32	0.08147	221	0.094771
	33	0.00711	222	> 1
	34	0.00748	223	0.18208
	35	0.29147	224	0.013887
15	36	0.18137	225	0.56001
	37	0.00118	226	0.1178
	38	3.5092	227	0.0073566
20	39	0.01974	228	> 1
	40	0.09974	229	> 1
	41	0.05801	230	0.052821
25	42	0.53412	231	0.52301
	43	0.27208	232	> 1
	44	0.05309	233	> 1
	45	0.00992	234	0.13532
30	46	> 5	235	0.03232
	47	0.03265	236	0.04292
	48	0.00333	237	0.05316
35	49	0.35161	238	0.10303
	50	0.31264	239	0.023699
	51	0.02308	240	0.017266
	52	0.19964	241	0.11377
40	53	0.06674	242	0.22275
	54	1.9158	243	0.80718
	55	0.0132	244	0.79378
45	56	0.08654	245	0.083614
	57	0.42611	246	0.40218
	58	> 5	247	0.092976
	59	0.7215	248	0.099588
50	60	0.05948	250	> 1
	61	0.18337	251	0.91782
	62	0.02506	252	0.003475
55	63	0.00751	253	0.049586
	64	0.00025	254	0.019908
	65	0.00025	255	0.009004

EP 2 944 638 A1

(continued)

	EXAMPLE #	EC50	EXAMPLE #	EC50
5	66	0.01893	256	0.017997
	67	0.04954	257	0.026002
	68	0.10846	258	0.00055345
	69	1.7243	259	0.00038795
10	70	> 5	260	0.0054323
	71	0.09165	261	0.18366
	72	0.00751	262	0.016346
15	73	0.02369	263	> 1
	74	0.057	264	0.68866
	75	0.01509	265	0.0071718
	76	0.51131	266	0.0072924
20	77	0.76196	267	0.06944
	78	0.01252	268	0.048792
	79	0.0649	269	0.0072346
25	80	0.06863	270	0.0025216
	81	0.04814	271	0.43657
	82	0.68383	272	0.84006
	83	0.197	273	0.20925
30	84	0.158	274	0.21418
	85	1.95	275	0.14303
	86	1.02	276	0.0035006
35	87	1.18	277	0.0081845
	88	0.447	278	0.79393
	89	0.06446	279	0.22492
	90	0.06299	280	0.45923
40	91	0.18296	281	0.65371
	92	0.08089	282	0.032187
	93	> 5	283	0.013096
45	94	1.6946	284	0.16213
	95	0.02954	285	0.057413
	96	0.04356	286	0.034464
	97	0.05557	287	0.59312
50	98	0.0229	288	0.39042
	99	1.3923	289	0.6687
	100	0.13666	290	0.10663
55	101	0.2991	291	0.016079
	102	0.62178	292	0.88938
	103	0.03917	293	0.28715

EP 2 944 638 A1

(continued)

	EXAMPLE #	EC50	EXAMPLE #	EC50
5	104	0.07125	294	0.12525
	105	0.05357	295	0.014803
	106	0.82639	296	0.76869
	107	0.06117	297	0.59157
10	108	0.02407	298	0.070305
	109	0.18339	299	0.067981
	110	0.53638	300	0.76334
15	111	0.01451	301	> 1
	112	0.02063	302	0.38106
	113	0.00136	303	0.04776
	114	0.01078	304	0.29755
20	115	0.01184	305	0.032539
	116	0.02853	306	0.55348
	117	0.0182	307	0.12767
25	118	0.01294	308	0.257
	119	0.01138	309	0.052421
	120	0.00147	310	> 1
	121	0.05972	311	0.035835
30	122	0.00185	312	0.016178
	123	0.00333	313	> 1
	124	0.21224	314	0.66006
35	125	0.00838	315	0.21027
	126	0.05359	316	> 1
	127	0.00975	317	0.013313
	128	0.00589	318	0.011566
40	129	0.01484	319	0.0044972
	130	0.01059	320	0.050974
	131	0.01266	321	0.0188
45	132	0.02209	322	0.012367
	133	0.03186	323	0.71689
	134	0.00251	324	0.0045254
	135	0.00237	325	0.012319
50	136	0.00296	326	0.023133
	137	0.01272	327	0.0027224
	138	0.00152	328	0.0098808
55	139	0.01681	329	0.42369
	140	0.01275	330	0.0097843
	141	0.02044	331	0.92638

EP 2 944 638 A1

(continued)

	EXAMPLE #	EC50	EXAMPLE #	EC50
5	142	0.34531	332	0.45738
	143	0.01914	333	0.46292
	144	0.0212	334	> 1
	145	0.004	335	0.26951
10	146	0.01916	336	0.35134
	147	0.02618	337	0.001759
	148	0.00938	338	0.003399
15	149	0.01347	339	0.45016
	150	0.05103	340	0.05646
	151	0.03372	341	0.031652
	152	0.02037	342	0.050891
20	153	0.01723	343	0.12664
	154	0.02647	344	0.0066616
	155	0.59421	345	0.0092536
25	156	0.00805	346	0.19003
	157	0.01086	347	0.018849
	158	0.01793	348	0.050263
	159	0.01179	349	0.023086
30	160	0.08363	350	0.0058378
	161	0.03465	351	0.0020618
	162	0.01297	352	0.0011961
35	163	0.00432	353	0.0050512
	164	0.01476	354	0.053231
	165	0.0051	355	0.018771
	166	0.01185	356	0.026623
40	167	0.00093	357	0.013235
	168	0.08867	358	0.0038131
	169	0.07626	359	0.0059243
45	170	0.12515	360	0.0098968
	171	0.05272	361	0.00053755
	172	0.02053	362	0.031726
	173	0.00516	363	0.02643
50	174	0.12621	364	0.011244
	175	> 1	365	0.0030168
	176	0.13353	366	0.016548
55	177	0.15936	367	nd
	178	0.20234	368	nd
	179	0.04273	369	0.0079974

(continued)

EXAMPLE #	EC50	EXAMPLE #	EC50
180	0.0118	370	nd
181	0.10612	371	0.007165
182	0.1234	372	nd
183	0.01753	373	nd
184	0.02323	374	0.015475
185	0.02747	375	0.56013
186	0.06443	376	0.008765
187	0.21494	377	0.002377
188	0.01638	378	0.006764
189	0.14397		
nd = not determined			

[0292] TABLE 3 shows the utility of compounds having Formula I to functionally inhibit anti-apoptotic Bcl-2 protein in a cellular context. The acute lymphoblastic leukemia (ALL) cell line RS4;11 has been shown by BH3 profiling, a mitochondrial assay that classifies blocks in the intrinsic apoptotic pathway, to be highly dependant on Bcl-2 for survival and is sensitive to the Bcl-2 family member inhibitor ABT-737 (Blood, 2008, Vol. 111, 2300-2309). The ability of compounds to kill RS4;11 cells is a direct measure of the compounds ability to inhibit anti-apoptotic Bcl-2 protein function. Compounds of Formula I are very effective in killing RS4;11 cells as demonstrated by low EC₅₀ values.

[0293] Compounds taught in U.S. Patent Application No. 12/631404, entitled "BCL-2-SELECTIVE APOPTOSIS-INDUCING AGENTS FOR THE TREATMENT OF CANCER AND IMMUNE DISEASES," filed on December 4, 2009, have utility for the treatment of various cancers and auto-immune diseases due to their activity against Bcl-2 family proteins, and more specifically Bcl-2. These compounds bind to Bcl-2 with high affinity in a FRET based assay described in 12/631404. The administration of a one or more of these compounds to cells that are dependant on Bcl-2 or Bcl-2 family proteins for survival, such as the RS4:11 B-cell leukemia human tumor cell line, results in apoptosis, also known as programmed cell death. The amount of apoptosis caused by administration of the compound is represented by the EC50 in the cell viability assay, which is a measure of the number of living cells after administration of compound.

[0294] TABLE 4 identifies certain compounds (described below in Examples in 19, 20, 23 and 92 and described more fully in 12/631404, the disclosure of which is incorporated herein by reference) with the various substituents being defined by R, X and Y as set forth. As can be seen from TABLE 4, these compounds exhibit a trend of increasing binding affinity (K_i) for Bcl-2 with increasing levels of apoptosis, or cell death, in the Bcl-2 dependant tumor cell line RS4;11. On this basis, the inventors expect that compounds with even greater affinity towards Bcl-2 than those compounds shown in Table 4 will exhibit a similar trend, potentially eliciting even greater levels of apoptosis, when administered to cells dependent on Bcl-2 for survival.

Table 4. Selected compounds in US Patent Application No. 12/631404

Chemical structure of the Bcl-2 FRET probe, showing a central benzene ring substituted with a 4-fluoro-2-iodophenyl group (X), a 4-(4-chlorophenyl)-4,4-dimethyl-1,2,3,4-tetrahydropyran-2-yl group (Y), and a 4-(4-(2-(4-((4-chlorophenyl)methyl)piperidin-1-yl)phenoxy)-5-nitrophenyl)butan-1-yl group (R).

EXAMPLE	R	X	Y	Bcl-2 FRET K_i (μ M)	RS4;11 EC50 (μ M)
(23)		C	O	0.000083	0.045
(92)		C	C	0.000128	0.081
(20)		C	C	0.000181	0.191
(19)		C	C	0.000226	0.465

[0295] To this end, binding affinity and cellular activity for compounds according to the present invention were compared with structurally similar indole compounds. In particular, the compounds of the present invention, in which a nitrogen is contained at a specific position within the heteroarene fused to the heteroaryl ring were compared with the corresponding indole compounds, which latter compounds lack only the specific nitrogen substitution included in the compounds of the present invention.

[0296] As can be seen in TABLE 5, compounds of the present invention having the specific nitrogen substitutions shown (i.e., compounds of Examples 1, 2, 3, 4, 5, 6, 9, 10, 11, 12, 13, 15, 16, and 17, where Z = N) in fact achieve relatively greater levels of apoptosis when administered to cells that depend on Bcl-2 for survival and have increased affinity towards Bcl-2 relative to the corresponding structural analogs lacking the specific nitrogen substitution (i.e., compounds of Examples 87, 88, 89, 90, 91, 19, 20, 21, 92, 22, 23, 93, and 94, respectively, where Z = C, taught in 9696USL2, the disclosure of which is incorporated herein by reference).

[0297] Specifically, the seventh column of TABLE 5 compares binding affinity of compounds of the present invention (the compound identified by the designated substituents in the uppermost row in each pair of rows set apart by blank rows) to corresponding compounds lacking the described nitrogen substitution. In each comparison, compounds of the present application (upper row of each pair of rows separated by a blank row) bind Bcl-2 with greater affinity to Bcl-2 than the corresponding analogs (lower row of each pair of rows separated by a blank row).

[0298] Further, column 8 of TABLE 5 compares the amount of apoptosis in the Bcl-2 dependant RS4;11 cell line






achieved using compounds of the present invention (again the compound identified by the designated substituents in the uppermost row in each pair of rows set apart by blank rows) to that achieved using compounds of Examples 87, 88, 89, 90, 91, 19, 20, 21, 92, 22, 23, 93, and 94, where Z = C. In each comparison, compounds of the present invention (upper row of each pair of rows separated by a blank row) achieve greater extent of apoptosis in Bcl-2 dependent RS4;11 cells than the corresponding analogs (lower row of each pair of rows separated by a blank row).

[0299] The increase in binding affinity between the compounds of the present invention and corresponding analogs ranges from 2.7x to greater than 100x, and the increased potency in RS4;11 cells ranges from a 1.65x increase to greater than 10x increase.





















[0300] As detailed below, a specific substitution of a nitrogen atom for a carbon atom leads to unexpected increase in binding affinity to antiapoptotic Bcl-2 and increase in potency in cell viability assays assessing apoptosis in Bcl-2 dependent cell lines.

[0301] This invention therefore comprises a series of compounds that demonstrate unexpected properties with respect to their binding to and inhibiting the activity of anti-apoptotic Bcl-2 protein to a significantly greater extent than corresponding analog compounds.

TABLE 5. Direct comparison of compounds of the present invention with corresponding analogs.









							
EXAMPLE	R	W	X	Ring	Z	Bcl-2 FRET ki (μM)	RS4;11 EC50 (μM)
(1)		NH	NO ₂		N	0.000225	0.712
(87)		NH	NO ₂		C	0.000618	1.180
(2)		NH	NO ₂		N	< 0.000010	0.783
(88)		NH	NO ₂		C	0.672	0.447

(continued)

EXAMPLE	R	W	X	Ring	Z	Bcl-2 FRET k_i (μ M)	RS4;11 EC ₅₀ (μ M)
5 (3)		NH	NO ₂		N	0.000013	0.0142
10 (89)		NH	NO ₂		C	0.000074	0.064
15 (4)		NH	NO ₂		N	< 0.00001	0.019
20 (90)		NH	NO ₂		C	0.000013	0.063
25 (5)		NH	NO ₂		N	< 0.00001	0.012
30 (18)		NH	NO ₂		C	0.000017	0.086
35 (6)		NH	NO ₂		N	0.000018	0.035
40 (91)		NH	NO ₂		C	0.000487	0.183
45 (9)		NH	NO ₂		N	< 0.00001	0.015
50 (19)		NH	NO ₂		C	0.000226	0.465

EP 2 944 638 A1


(continued)

EXAMPLE	R	W	X	Ring	Z	Bcl-2 FRET k_i (μ M)	RS4;11 EC ₅₀ (μ M)
5							
(10)		NH	NO ₂		N	< 0.00001	0.052
10							
(20)		NH	NO ₂		C	0.000181	0.191
15							
(11)		NH	NO ₂		N	0.000016	0.014
20							
(21)		NH	NO ₂		C	0.000912	0.062
25							
(12)		O	NO ₂		N	< 0.00001	0.035
30							
(92)		O	NO ₂		C	0.000128	0.081
35							
(13)		NH	NO ₂		N	< 0.00001	0.023
40							
(22)		NH	NO ₂		C	0.000291	0.085
45							
(15)		NH	NO ₂		N	< 0.00001	0.013
50							
(23)		NH	NO ₂		C	0.000083	0.045
55							


(continued)

EXAMPLE	R	W	X	Ring	Z	Bcl-2 FRET k_i (μ M)	RS4;11 EC ₅₀ (μ M)
(16)		NH	SO ₂ CF ₃		N	0.000219	1.210
(93)		NH	SO ₂ CF ₃		C	0.035	> 5.000
(17)		NH	SO ₂ CF ₃		N	0.000090	0.475
(94)		NH	SO ₂ CF ₃		C	0.000678	1.690


[0302] More specifically, compounds of the present invention contain a substitution pattern shown in the diagram below.




[0303] Other compounds that contain isomeric ring systems to that shown above, such as those rings systems containing a nitrogen adjacent to the oxygenated carbon within the ring, as shown below, are compromised by instability.



[0304] Specifically, this was discovered by the inventors in the following compound preparation. The intermediate structure F, that directly precedes the final product of the unstable compound, was prepared according to the route below. All intermediates A-F were stable and isolable using techniques known to those skilled in the art.



[0305] Intermediate F, shown in the scheme above, was reacted with intermediate G using standard coupling conditions that are known to those skilled in the art. The reaction mixture was analyzed via HPLC/MS to monitor the formation of a peak corresponding to the compound H. While this peak formed within hours of initiating the reaction below, the peak progressively disappeared during workup and chromatography, until it no longer was present. The lack of stability of the putative compound originates from the position of the nitrogen within the fused ring-system described above. This position, which is adjacent to the oxygen-bearing carbon in the 5-substituted-1H-pyrrolo[2,3-c]pyridine ring system shown below and described above, makes the compound H unstable.



[0306] It is expected that a compound containing the fused 5-substituted-1H-pyrrolo[3,2-b]pyridine ring system below would be similarly unstable, since the position of the nitrogen is adjacent to the oxygen-bearing carbon within the ring.



[0307] Therefore, compounds with the 5-substituted-1H-pyrrolo[2,3-b]pyridines are preferred over the isomeric compounds.

[0308] It is expected that, because compounds having Formula (I) bind to Bcl-2, they would also have utility as binders

to anti-apoptotic proteins having close structural homology to Bcl-2, such as, for example, anti-apoptotic Bcl-X_L, Bcl-w, Mcl-1 and Bfl-1/A1 proteins.

[0309] Involvement of Bcl-2 proteins in bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, prostate cancer, small cell lung cancer, chronic lymphocytic leukemia, myeloma, prostate cancer spleen cancer, and the like is described in commonly-owned PCT US 2004/36770, published as WO 2005/049593, and PCT US 2004/37911, published as WO 2005/024636.

[0310] Involvement of Bcl-2 proteins in immune and autoimmune diseases is described in Current Allergy and Asthma Reports 2003, 3, 378-384; British Journal of Haematology 2000, 110(3), 584-90; Blood 2000, 95(4), 1283-92; and New England Journal of Medicine 2004, 351(14), 1409-1418.

[0311] Involvement of Bcl-2 proteins in arthritis is disclosed in commonly-owned United States Provisional Patent Application Serial No. 60/988,479.

[0312] Involvement of Bcl-2 proteins in bone marrow transplant rejection is disclosed in commonly-owned United States Patent Application Serial No. 11/941,196.

[0313] Overexpression of Bcl-2 proteins correlates with resistance to chemotherapy, clinical outcome, disease progression, overall prognosis or a combination thereof in various cancers and disorders of the immune system. Cancers include, but are not limited to, hematologic and solid tumor types such as acoustic neuroma, acute leukemia, acute lymphoblastic leukemia, acute myelogenous leukemia (monocytic, myeloblastic, adenocarcinoma, angiosarcoma, astrocytoma, myelomonocytic and promyelocytic), acute t-cell leukemia, basal cell carcinoma, bile duct carcinoma, bladder cancer, brain cancer, breast cancer (including estrogen-receptor positive breast cancer), bronchogenic carcinoma, Burkitt's lymphoma, cervical cancer, chondrosarcoma, chordoma, choriocarcinoma, chronic leukemia, chronic lymphocytic leukemia, chronic myelocytic (granulocytic) leukemia, chronic myelogenous leukemia, colon cancer, colorectal cancer, craniopharyngioma, cystadenocarcinoma, dysproliferative changes (dysplasias and metaplasias), embryonal carcinoma, endometrial cancer, endotheliosarcoma, ependymoma, epithelial carcinoma, erythroleukemia, esophageal cancer, estrogen-receptor positive breast cancer, essential thrombocythemia, Ewing's tumor, fibrosarcoma, gastric carcinoma, germ cell testicular cancer, gestational trophoblastic disease, glioblastoma, head and neck cancer, heavy chain disease, hemangioblastoma, hepatoma, hepatocellular cancer, hormone insensitive prostate cancer, leiomyosarcoma, liposarcoma, lung cancer (including small cell lung cancer and non-small cell lung cancer), lymphangioendotheliosarcoma, lymphangiosarcoma, lymphoblastic leukemia, lymphoma (lymphoma, including diffuse large B-cell lymphoma, follicular lymphoma, Hodgkin's lymphoma and non-Hodgkin's lymphoma), malignancies and hyperproliferative disorders of the bladder, breast, colon, lung, ovaries, pancreas, prostate, skin and uterus, lymphoid malignancies of T-cell or B-cell origin, leukemia, medullary carcinoma, medulloblastoma, melanoma, meningioma, mesothelioma, multiple myeloma, myelogenous leukemia, myeloma, myxosarcoma, neuroblastoma, oligodendroglioma, oral cancer, osteogenic sarcoma, ovarian cancer, pancreatic cancer, papillary adenocarcinomas, papillary carcinoma, peripheral T-cell lymphoma, pinealoma, polycythemia vera, prostate cancer (including hormone-insensitive (refractory) prostate cancer), rectal cancer, renal cell carcinoma, retinoblastoma, rhabdomyosarcoma, sarcoma, sebaceous gland carcinoma, seminoma, skin cancer, small cell lung carcinoma, solid tumors (carcinomas and sarcomas), stomach cancer, squamous cell carcinoma, synovioma, sweat gland carcinoma, testicular cancer (including germ cell testicular cancer), thyroid cancer, Waldenström's macroglobulinemia, testicular tumors, uterine cancer, Wilms' tumor and the like.

[0314] It is also expected that compounds having Formula (I) would inhibit growth of cells expressing Bcl-2 proteins derived from a pediatric cancer or neoplasm including embryonal rhabdomyosarcoma, pediatric acute lymphoblastic leukemia, pediatric acute myelogenous leukemia, pediatric alveolar rhabdomyosarcoma, pediatric anaplastic ependymoma, pediatric anaplastic large cell lymphoma, pediatric anaplastic medulloblastoma, pediatric atypical teratoid/rhabdoid tumor of the central nervous system, pediatric biphenotypic acute leukemia, pediatric Burkitt's lymphoma, pediatric cancers of Ewing's family of tumors such as primitive neuroectodermal tumors, pediatric diffuse anaplastic Wilm's tumor, pediatric favorable histology Wilm's tumor, pediatric glioblastoma, pediatric medulloblastoma, pediatric neuroblastoma, pediatric neuroblastoma-derived myelocytomatosis, pediatric pre-B-cell cancers (such as leukemia), pediatric psteosarcoma, pediatric rhabdoid kidney tumor, pediatric rhabdomyosarcoma, and pediatric T-cell cancers such as lymphoma and skin cancer and the like.

[0315] Autoimmune disorders include acquired immunodeficiency disease syndrome (AIDS), autoimmune lymphoproliferative syndrome, hemolytic anemia, inflammatory diseases, and thrombocytopenia, acute or chronic immune disease associated with organ transplantation, Addison's disease, allergic diseases, alopecia, alopecia areata, atheromatous disease/arteriosclerosis, atherosclerosis, arthritis (including osteoarthritis, juvenile chronic arthritis, septic arthritis, Lyme arthritis, psoriatic arthritis and reactive arthritis), autoimmune bullous disease, abetalipoproteinemia, acquired immunodeficiency-related diseases, acute immune disease associated with organ transplantation, acquired acrocyanosis, acute and chronic parasitic or infectious processes, acute pancreatitis, acute renal failure, acute rheumatic fever, acute transverse myelitis, adenocarcinomas, aerial ectopic beats, adult (acute) respiratory distress syndrome, AIDS

dementia complex, alcoholic cirrhosis, alcohol-induced liver injury, alcohol-induced hepatitis, allergic conjunctivitis, allergic contact dermatitis, allergic rhinitis, allergy and asthma, allograft rejection, alpha-1- antitrypsin deficiency, Alzheimer's disease, amyotrophic lateral sclerosis, anemia, angina pectoris, ankylosing spondylitis associated lung disease, anterior horn cell degeneration, antibody mediated cytotoxicity, antiphospholipid syndrome, anti-receptor hypersensitivity reactions, aortic and peripheral aneurysms, aortic dissection, arterial hypertension, arteriosclerosis, arteriovenous fistula, arthropathy, asthenia, asthma, ataxia, atopic allergy, atrial fibrillation (sustained or paroxysmal), atrial flutter, atrioventricular block, atrophic autoimmune hypothyroidism, autoimmune haemolytic anaemia, autoimmune hepatitis, type-1 autoimmune hepatitis (classical autoimmune or lupoid hepatitis), autoimmune mediated hypoglycaemia, autoimmune neutropaenia, autoimmune thrombocytopaenia, autoimmune thyroid disease, B cell lymphoma, bone graft rejection, bone marrow transplant (BMT) rejection, bronchiolitis obliterans, bundle branch block, burns, cachexia, cardiac arrhythmias, cardiac stun syndrome, cardiac tumors, cardiomyopathy, cardiopulmonary bypass inflammation response, cartilage transplant rejection, cerebellar cortical degenerations, cerebellar disorders, chaotic or multifocal atrial tachycardia, chemotherapy associated disorders, chlamydia, choleosarcoma, chronic alcoholism, chronic active hepatitis, chronic fatigue syndrome, chronic immune disease associated with organ transplantation, chronic eosinophilic pneumonia, chronic inflammatory pathologies, chronic mucocutaneous candidiasis, chronic obstructive pulmonary disease (COPD), chronic salicylate intoxication, colorectal common varied immunodeficiency (common variable hypogammaglobulinaemia), conjunctivitis, connective tissue disease associated interstitial lung disease, contact dermatitis, Coombs positive haemolytic anaemia, cor pulmonale, Creutzfeldt-Jakob disease, cryptogenic autoimmune hepatitis, cryptogenic fibrosing alveolitis, culture negative sepsis, cystic fibrosis, cytokine therapy associated disorders, Crohn's disease, dementia pugilistica, demyelinating diseases, dengue hemorrhagic fever, dermatitis, dermatitis scleroderma, dermatologic conditions, dermatomyositis/polymyositis associated lung disease, diabetes, diabetic arteriosclerotic disease, diabetes mellitus, Diffuse Lewy body disease, dilated cardiomyopathy, dilated congestive cardiomyopathy, discoid lupus erythematosus, disorders of the basal ganglia, disseminated intravascular coagulation, Down's Syndrome in middle age, drug-induced interstitial lung disease, drug-induced hepatitis, drug-induced movement disorders induced by drugs which block CNS dopamine receptors, drug sensitivity, eczema, encephalomyelitis, endocarditis, endocrinopathy, enteropathic synovitis, epiglottitis, Epstein-Barr virus infection, erythromelalgia, extrapyramidal and cerebellar disorders, familial hematophagocytic lymphohistiocytosis, fetal thymus implant rejection, Friedreich's ataxia, functional peripheral arterial disorders, female infertility, fibrosis, fibrotic lung disease, fungal sepsis, gas gangrene, gastric ulcer, giant cell arteritis, glomerular nephritis, glomerulonephritides, Goodpasture's syndrome, goitrous autoimmune hypothyroidism (Hashimoto's disease), gouty arthritis, graft rejection of any organ or tissue, graft versus host disease, gram negative sepsis, gram positive sepsis, granulomas due to intracellular organisms, group B streptococci (GBS) infection, Grave's disease, haemosiderosis associated lung disease, hairy cell leukemia, hairy cell leukemia, Hallerorden-Spatz disease, Hashimoto's thyroiditis, hay fever, heart transplant rejection, hemachromatosis, hematopoietic malignancies (leukemia and lymphoma), hemolytic anemia, hemolytic uremic syndrome/thrombolytic thrombocytopenic purpura, hemorrhage, Henoch-Schoenlein purpura, Hepatitis A, Hepatitis B, Hepatitis C, HIV infection/HIV neuropathy, Hodgkin's disease, hypoparathyroidism, Huntington's chorea, hyperkinetic movement disorders, hypersensitivity reactions, hypersensitivity pneumonitis, hyperthyroidism, hypokinetic movement disorders, hypothalamic-pituitary-adrenal axis evaluation, idiopathic Addison's disease, idiopathic leucopaenia, idiopathic pulmonary fibrosis, idiopathic thrombocytopaenia, idiosyncratic liver disease, infantile spinal muscular atrophy, infectious diseases, inflammation of the aorta, inflammatory bowel disease, insulin dependent diabetes mellitus, interstitial pneumonitis, iridocyclitis/uveitis/optic neuritis, ischemia-reperfusion injury, ischemic stroke, juvenile pernicious anaemia, juvenile rheumatoid arthritis, juvenile spinal muscular atrophy, Kaposi's sarcoma, Kawasaki's disease, kidney transplant rejection, legionella, leishmaniasis, leprosy, lesions of the corticospinal system, linear IgA disease, lipidema, liver transplant rejection, Lyme disease, lymphedema, lymphocytic infiltrative lung disease, malaria, male infertility idiopathic or NOS, malignant histiocytosis, malignant melanoma, meningitis, meningococcemia, microscopic vasculitis of the kidneys, migraine headache, mitochondrial multisystem disorder, mixed connective tissue disease, mixed connective tissue disease associated lung disease, monoclonal gammopathy, multiple myeloma, multiple systems degenerations (Mencel Dejerine-Thomas Shi-Drager and Machado-Joseph), myalgic encephalitis/Royal Free Disease, myasthenia gravis, microscopic vasculitis of the kidneys, mycobacterium avium intracellulare, mycobacterium tuberculosis, myelodysplastic syndrome, myocardial infarction, myocardial ischemic disorders, nasopharyngeal carcinoma, neonatal chronic lung disease, nephritis, nephrosis, nephrotic syndrome, neurodegenerative diseases, neurogenic muscular atrophies, neutropenic fever, Non-alcoholic Steatohepatitis, occlusion of the abdominal aorta and its branches, occlusive arterial disorders, organ transplant rejection, orchitis/epididymitis, orchitis/vasectomy reversal procedures, organomegaly, osteoarthritis, osteoporosis, ovarian failure, pancreas transplant rejection, parasitic diseases, parathyroid transplant rejection, Parkinson's disease, pelvic inflammatory disease, pemphigus vulgaris, pemphigus foliaceus, pemphigoid, perennial rhinitis, pericardial disease, peripheral atherosclerotic disease, peripheral vascular disorders, peritonitis, pernicious anemia, phacogenic uveitis, pneumocystis carinii pneumonia, pneumonia, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal gammopathy, and skin changes syndrome), post perfusion syndrome, post pump syndrome, post-MI cardiomyopathy syndrome, postinfectious interstitial lung disease,


premature ovarian failure, primary biliary cirrhosis, primary sclerosing hepatitis, primary myxoedema, primary pulmonary hypertension, primary sclerosing cholangitis, primary vasculitis, Progressive supranucleo Palsy, psoriasis, psoriasis type 1, psoriasis type 2, psoriatic arthropathy, pulmonary hypertension secondary to connective tissue disease, pulmonary manifestation of polyarteritis nodosa, post-inflammatory interstitial lung disease, radiation fibrosis, radiation therapy, Raynaud's phenomenon and disease, Raynoud's disease, Refsum's disease, regular narrow QRS tachycardia, Reiter's disease, renal disease NOS, renovascular hypertension, reperfusion injury, restrictive cardiomyopathy, rheumatoid arthritis associated interstitial lung disease, rheumatoid spondylitis, sarcoidosis, Schmidt's syndrome, scleroderma, senile chorea, Senile Dementia of Lewy body type, sepsis syndrome, septic shock, seronegative arthropathies, shock, sickle cell anemia, Sjögren's disease associated lung disease, Sjörgren's syndrome, skin allograft rejection, skin changes syndrome, small bowel transplant rejection, sperm autoimmunity, multiple sclerosis (all subtypes), spinal ataxia, spinocerebellar degenerations, spondyloarthropathy, spondyloarthropathy, sporadic, polyglandular deficiency type I sporadic, polyglandular deficiency type II, Still's disease, streptococcal myositis, stroke, structural lesions of the cerebellum, Subacute sclerosing panencephalitis, sympathetic ophthalmia, Syncope, syphilis of the cardiovascular system, systemic anaphylaxis, systemic inflammatory response syndrome, systemic onset juvenile rheumatoid arthritis, systemic lupus erythematosus, systemic lupus erythematosus-associated lung disease, systemic sclerosis, systemic sclerosis-associated interstitial lung disease, T-cell or FAB ALL, Takayasu's disease/arteritis, Telangiectasia, Th2 Type and Th1 Type mediated diseases, thromboangitis obliterans, thrombocytopenia, thyroiditis, toxicity, toxic shock syndrome, transplants, trauma/hemorrhage, type-2 autoimmune hepatitis (anti-LKM antibody hepatitis), type B insulin resistance with acanthosis nigricans, type III hypersensitivity reactions, type IV hypersensitivity, ulcerative colitic arthropathy, ulcerative colitis, unstable angina, uremia, urosepsis, urticaria, uveitis, valvular heart diseases, varicose veins, vasculitis, vasculitic diffuse lung disease, venous diseases, venous thrombosis, ventricular fibrillation, vitiligo acute liver disease, viral and fungal infections, vital encephalitis/aseptic meningitis, vital-associated hemaphagocytic syndrome, Wegener's granulomatosis, Wernicke-Korsakoff syndrome, Wilson's disease, xenograft rejection of any organ or tissue, yersinia and salmonella-associated arthropathy and the like.

Schemes and Experimentals

[0316] The following abbreviations have the meanings indicated. ADDP means 1,1'-(azodicarbonyl)dipiperidine; AD-mix- β means a mixture of $(\text{DHQD})_2\text{PHAL}$, $\text{K}_3\text{Fe}(\text{CN})_6$, K_2CO_3 , and K_2SO_4 ; 9-BBN means 9-borabicyclo(3.3.1)nonane; Boc means tert-butoxycarbonyl; $(\text{DHQD})_2\text{PHAL}$ means hydroquinidine 1,4-phthalazinediyl diethyl ether; DBU means 1,8-diazabicyclo[5.4.0]undec-7-ene; DIBAL means diisobutylaluminum hydride; DIEA means diisopropylethylamine; DMAP means N,N-dimethylaminopyridine; DMF means N,N-dimethylformamide; dmpe means 1,2-bis(dimethylphosphino)ethane; DMSO means dimethylsulfoxide; dppb means 1,4-bis(diphenylphosphino)-butane; dppe means 1,2-bis(diphenylphosphino)ethane; dppf means 1,1'-bis(diphenylphosphino)ferrocene; dppm means 1,1-bis(diphenylphosphino)methane; EDAC-HCl means 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride; Fmoc means fluorenylmethoxycarbonyl; HATU means O-(7-azabenzotriazol-1-yl)-N,N',N'-tetramethyluronium hexafluorophosphate; HM-PA means hexamethylphosphoramide; IPA means isopropyl alcohol; MP-BH₃ means macroporous triethylammonium methylpolystyrene cyanoborohydride; TEA means triethylamine; TFA means trifluoroacetic acid; THF means tetrahydrofuran; NCS means N-chlorosuccinimide; NMM means N-methylmorpholine; NMP means N-methylpyrrolidine; PPh₃ means triphenylphosphine.


[0317] The following schemes are presented to provide what is believed to be the most useful and readily understood description of procedures and conceptual aspects of this invention. Compounds of this invention may be made by synthetic chemical processes, examples of which are shown herein. It is meant to be understood that the order of the steps in the processes may be varied, that reagents, solvents and reaction conditions may be substituted for those specifically mentioned, and that vulnerable moieties may be protected and deprotected, as necessary.

SCHEME 1




[0318] Compounds of Formula (4) can be prepared as shown in SCHEME 1, and can be used as described in SCHEME 8 to prepare compounds of Formula (I), which are representative of the compounds of the present invention. Compounds of Formula (1) wherein R is alkyl, can be converted to compounds of Formula (2) using $Z^3L^1MgX^1$, wherein X^1 is a halide, in a solvent such as but not limited to ether or tetrahydrofuran. Compounds of Formula (3) can be prepared from compounds of Formula (2) using a strong base such as NaH and $R^{57}X^2$, wherein X^2 is a halide and R^{57} is as described herein. Compounds of Formula (3), when treated with aqueous NaOH or LiOH, will provide compounds of Formula (4).

SCHEME 2




[0319] As shown in SCHEME 2, compounds of Formula (5) can be reacted with compounds of Formula (6) and a reducing agent to provide compounds of Formula (7). Examples of reducing agents include sodium borohydride, sodium cyanoborohydride, sodium triacetoxyborohydride, polymer supported cyanoborohydride, and the like. The reaction is typically performed in a solvent such as but not limited to methanol, tetrahydrofuran, and dichloromethane or mixtures thereof. Compounds of Formula (8) can be prepared from compounds of Formula (7) as described in SCHEME 1, and can be used as described in SCHEME 8 to prepare compounds of Formula (I).

SCHEME 3




[0320] Compounds of Formula (9), when reacted with a compound a Formula (10) wherein X is a halide or triflate, and a base will provide a compound of Formula (11). Bases useful in the reaction include triethylamine, diisopropylethylamine and the like. Compounds of Formula (13), wherein Y is as described herein for substituents on Z^3 , can be prepared from compounds of Formula (11) and compounds of Formula (12) using Suzuki coupling conditions known to those skilled in the art and readily available in the literature. Compounds of Formula (14) can be prepared from compounds of Formula (13) as described in SCHEME 1, and can be used as described in SCHEME 8 to prepare compounds of Formula (I).

SCHEME 4



[0321] As shown in SCHEME 4, compounds of Formula (17) can be prepared from compounds of Formula (15) and compounds of Formula (16), wherein R is alkyl and R³⁸ is as described herein, using Suzuki coupling conditions known to those skilled in the art and readily available in the literature. Compounds of Formula (17) can be reduced to compounds of Formula (18) using a reducing agent such as LiAlH₄ in a solvent such as but not limited to diethyl ether or THF. Compounds of Formula (19) can be prepared from compounds of Formula (18) using Dess-Martin periodinane or Swern oxidation conditions known to those skilled in the art and readily available in the literature. Compounds of Formula (19) can be reacted with a compound of Formula (5) and a reducing agent to provide compounds of Formula (20). Examples of reducing agents include sodium borohydride, sodium cyanoborohydride, sodium triacetoxyborohydride, polymer supported cyanoborohydride, and the like. The reaction is typically performed in a solvent such as but not limited to methanol, tetrahydrofuran, 1,2-dichloroethane, and dichloromethane or mixtures thereof. Compounds of Formula (21) can be prepared from compounds of Formula (20) as described in SCHEME 1, and can be used as described in SCHEME 8 to prepare compounds of Formula (I).


SCHEME 5



[0322] As shown in SCHEME 5, compounds of Formula (22), wherein R is alkyl, may be converted to compounds of Formula (23) by reacting the former, wherein X¹ is Cl, Br, I, or CF₃SO₃⁻, and compounds of Formula R⁴¹-OH and a catalyst, with or without a first base. Examples of catalysts include copper(I) trifluoromethanesulfonate toluene complex, PdCl₂, Pd(OAc)₂, and Pd₂(dba)₃. Examples of first bases include triethylamine, N,N-diisopropylethylamine, Cs₂CO₃, Na₂CO₃, K₃PO₄, and mixtures thereof.


[0323] Compounds of Formula (22) may also be converted to compounds of Formula (23) by reacting the former, when X¹ is Cl, F, or NO₂, and compounds of Formula R⁴¹-OH with a first base. Examples of first bases include triethylamine, N,N-diisopropylethylamine, Cs₂CO₃, Na₂CO₃, K₃PO₄, and mixtures thereof.

SCHEME 6




[0324] Compounds of Formula (18) can be reacted with mesyl chloride and a base such as but not limited to triethylamine, followed by N-t-butoxycarbonylpiperazine, to provide compounds of Formula (24). Compounds of Formula (25) can be prepared by reacting compounds of Formula (24) with triethylsilane and trifluoroacetic acid. Compounds of Formula (25) can be reacted with compounds of Formula (26) and HK₂PO₄ to provide compounds of Formula (27) in a solvent such as but not limited to dimethylsulfoxide. Compounds of Formula (28) can be prepared from compounds of Formula (27) as described in SCHEME 1, and can be used as described in SCHEME 8 to prepare compounds of Formula (I).

SCHEME 7



[0325] As shown in SCHEME 7, compounds of Formula (1) can be reacted with an appropriate triphenylphosphonium bromide of Formula (29) and a base such as but not limited to sodium hydride or n-butyllithium to provide compounds of Formula (30). The reaction is typically performed in a solvent such as THF or DMSO. Compounds of Formula (31) can be prepared from compounds of Formula (30) as described in SCHEME 1, and can be used as described in SCHEME 8 to prepare compounds of Formula (I).


SCHEME 8



10 **[0326]** As shown in SCHEME 8, compounds of Formula (32), which can be prepared as described herein, may be converted to compounds of Formula (33) by reacting the former with ammonia. Compounds of Formula (33) may be converted to compounds of Formula (I) by reacting the former and compounds of Formula (4), (8), (14), (21), (28), (31), or (38) and a coupling agent, with or without a first base. Examples of coupling agents include 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride, 1,1'-carbonyldiimidazole, and benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate. Examples of first bases include triethylamine, N,N-diisopropylethylamine, 4-(dimethylamino)pyridine, and mixtures thereof.

15


SCHEME 9



30 **[0327]** Compounds of Formula (33), prepared as described in SCHEME 8, may also be converted to compounds of Formula (I) by reacting the former and compounds of Formula (34) and a first base. Examples of first bases include but are not limited to sodium hydride, triethylamine, N,N-diisopropylethylamine, 4-(dimethylamino)pyridine, and mixtures thereof.

35


SCHEME 10



[0328] As shown in SCHEME 10, compounds of Formula (35), wherein L is a bond, alkyl, O, S, S(O), S(O)₂, NH, etc., can be reacted with compounds of Formula (36), to provide compounds of Formula (37). The reaction is typically performed at elevated temperatures in a solvent such as but not limited to dimethylsulfoxide, and may require the use of a base such as but not limited to potassium phosphate, potassium carbonate, and the like. Compounds of Formula (38) can be prepared from compounds of Formula (37) as described in SCHEME 1, and can be used as described in SCHEME 8 to prepare compounds of Formula (I).


55

SCHEME 11



[0329] Compounds of Formula (39), wherein Y is as described herein for substituents on Z³, can be prepared from compounds of Formula (39A) wherein X is a halide or triflate, and Y-B(OH)₂ using Suzuki coupling conditions known to those skilled in the art and readily available in the literature. Compounds of Formula (39) can be reacted with tert-butyl piperazine-1-carboxylate and a reducing agent such as sodium triacetoxyborohydride to provide compounds of Formula (40). The reaction is typically performed in a solvent such as but not limited to methylene chloride. Compounds of Formula (41) can be prepared from compounds of Formula (40) by reacting the latter with R⁵⁷X, wherein X is a halide, and NaH in a solvent such as N,N-dimethylformamide, and then the resulting material can be treated with triethylsilane and trifluoroacetic acid in dichloromethane. Compounds of Formula (41) can be used as described in Scheme 10 wherein L¹-Z³ is as shown in Formula (41).

SCHEME 12



[0330] As shown in SCHEME 12, substituted piperazin-2-ones wherein R⁵⁷ is alkyl, can be reacted with compounds of Formula (6) and a reducing agent such as sodium triacetoxyborohydride in dichloromethane to provide compounds of Formula (42). Compounds of Formula (42) can be reduced to compounds of Formula (43) using a reducing agent such as but not limited to lithium aluminum hydride in a solvent such as but not limited to tetrahydrofuran. Compounds of Formula (43) can be used as described in Scheme 10 wherein L¹-Z³ is as shown in Formula (43).

[0331] The following examples are presented to provide what is believed to be the most useful and readily understood description of procedures and conceptual aspects of this invention. The exemplified compounds were named using ACD/ChemSketch Version 5.06 (05 June 2001, Advanced Chemistry Development Inc., Toronto, Ontario), ACD/ChemSketch Version 12.01 (13 May 2009), Advanced Chemistry Development Inc., Toronto, Ontario), or ChemDraw® Ver. 9.0.5 (CambridgeSoft, Cambridge, MA). Intermediates were named using ChemDraw® Ver. 9.0.5 (CambridgeSoft, Cambridge, MA).

EXAMPLE 1

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 1A

tert-butyl 4-[(4'-chlorobiphenyl-2-yl)methyl]piperazine-1-carboxylate

[0332] 4'-Chlorobiphenyl-2-carboxaldehyde (4.1 g), tert-butyl piperazine-1-carboxylate (4.23 g), and sodium triace-

toxyborohydride (5.61 g) in CH_2Cl_2 (60 mL) were stirred for 24 hours. The reaction was quenched with methanol and poured into ether. The solution was washed with water and brine, concentrated, and chromatographed on silica gel with 2-25% ethyl acetate/hexanes.

5 EXAMPLE 1B

1-((4'-chlorobiphenyl-2-yl)methyl)piperazine

10 **[0333]** EXAMPLE 1A (3.0 g) and triethylsilane (1 mL) were stirred in CH_2Cl_2 (30 mL) and trifluoroacetic acid (30 mL) for 2 hours, and the reaction was concentrated, and then taken up in ether and concentrated again. The material was taken up in dichloromethane (200 mL) and NaHCO_3 solution (100 mL), and partitioned. The organic layer was dried over Na_2SO_4 , and condensed to give the title compound.

15 EXAMPLE 1C

tert-butyl 4-(4-((4'-chlorobiphenyl-2-yl)methyl)piperazin-1-yl)-2-fluorobenzoate

20 **[0334]** Tert-butyl 4-bromo-2-fluorobenzoate (14.0 g), EXAMPLE 1B (16.05 g), $\text{Pd}_2(\text{dba})_3$ (tris(dibenzylideneacetone)dipalladium(0)) (1.40 g), 2-(di-tert-butylphosphino)biphenyl (1.82 g), and K_3PO_4 (16.2 g) were stirred in 1,2-dimethoxyethane (300 mL) at 80°C for 24 hours. The reaction was cooled and concentrated. The crude product was chromatographed on silica gel with 10-20% ethyl acetate/hexanes.

EXAMPLE 1D

25 tert-butyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((4'-chlorobiphenyl-2-yl)methyl)piperazin-1-yl)benzoate

30 **[0335]** 1H-Pyrrolo[2,3-B]pyridine-5-ol (167 mg), EXAMPLE 1C (500 mg), and Cs_2CO_3 (508 mg) were stirred in dimethylsulfoxide (5 mL) at 130°C for 24 hours. The mixture was cooled, diluted with ethyl acetate, washed three times with water, and brine, and dried (Na_2SO_4), filtered and concentrated. The crude product was chromatographed on silica gel with 25% ethyl acetate/hexanes.

EXAMPLE 1E

35 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((4'-chlorobiphenyl-2-yl)methyl)piperazin-1-yl)benzoic acid

[0336] EXAMPLE 1D (200 mg) and triethylsilane (1 mL) were stirred in dichloromethane (15 mL) and trifluoroacetic acid (15 mL) for 1 hour. The mixture was concentrated, taken up in ethyl acetate, washed twice with NaH_2PO_4 , and brine, and dried (Na_2SO_4), filtered and concentrated.

40 EXAMPLE 1F

3-nitro-4-((tetrahydro-2H-pyran-4-yl)methylamino)benzenesulfonamide

45 **[0337]** 4-Fluoro-3-nitrobenzenesulfonamide (2.18 g), 1-(tetrahydropyran-4-yl)methylamine (1.14 g), and triethylamine (1 g) were stirred in tetrahydrofuran (30 mL) for 24 hours. The solution was diluted with ethyl acetate, washed with NaH_2PO_4 solution and brine, and dried (Na_2SO_4), filtered and concentrated. The product was triturated from ethyl acetate.

EXAMPLE 1G

50 4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0338] EXAMPLE 1E (115 mg), EXAMPLE 1F (67 mg), 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (82 mg), and 4-dimethylaminopyridine (26 mg) were stirred in CH_2Cl_2 (3 mL) for 24 hours. The reaction was cooled and chromatographed on silica gel with 0-5% methanol/ethyl acetate. ^1H NMR (300MHz, dimethylsulfoxide- d_6) δ 11.48 (brs, 1H), 8.34 (br s, 1H), 8.31 (m, 1H), 7.90 (d, 1H), 7.68 (m, 1H), 7.58 (m, 2H), 7.46 (m, 4H), 7.35 (m, 2H), 7.21 (dd, 1H), 6.76 (m, 4H), 6.28 (m, 2H), 3.02 (m, 2H), 2.89 (m, 4H), 2.80 (m, 4H), 2.40 (m, 3H), 1.59 (m, 2H), 1.25 (m, 4H), 0.87 (m, 2H).

EXAMPLE 2

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({4-[(3-morpholin-4-ylpropyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 2A

4-(3-morpholinopropylamino)-3-nitrobenzenesulfonamide

[0339] This EXAMPLE was prepared by substituting 3-(N-morpholinyl)-propylamine for 1-(tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 2B

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({4-[(3-morpholin-4-ylpropyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0340] This EXAMPLE was prepared by substituting EXAMPLE 2A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.60 (brs, 1H), 8.60 (m, 1H), 8.43 (d, 1H), 7.94 (d, 1H), 7.64 (m, 2H), 7.54 (d, 1H), 7.45 (m, 4H), 7.33 (m, 2H), 7.23 (dd, 1H), 6.96 (d, 1H), 6.85 (m, 2H), 6.32 (d, 1H), 6.26 (d, 1H), 3.60 (m, 4H), 3.10 (m, 4H), 3.05 (m, 10H), 2.40 (m, 2H), 2.33 (m, 2H), 1.77 (m, 2H).

EXAMPLE 3

4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({3-nitro-4-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 3A

methyl 4,4-dimethyl-2-(trifluoromethylsulfonyloxy)cyclohex-1-enecarboxylate

[0341] To a suspension of hexane washed NaH (17 g) in dichloromethane (700 mL) was added 5,5-dimethyl-2-methoxycarbonylcyclohexanone (38.5 g) dropwise at 0°C. After stirring for 30 minutes, the mixture was cooled to -78°C and trifluoroacetic anhydride (40 mL) was added. The reaction mixture was warmed to room temperature and stirred for 24 hours. The organic layer was washed with brine, dried (Na₂SO₄), filtered, and concentrated to give the product.

EXAMPLE 3B

methyl 2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enecarboxylate

[0342] EXAMPLE 3A (62.15g), 4-chlorophenylboronic acid (32.24 g), CsF (64 g) and tetrakis(triphenylphosphine)palladium(0) (2g) in 2:1 dimethoxyethane /methanol (600 mL) were heated to 70°C for 24 hours. The mixture was concentrated. Ether (4x 200 mL) was added and the mixture was filtered. The combined ether solution was concentrated to give the product.

EXAMPLE 3C

(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methanol

[0343] To a mixture of LiBH₄ (13g), EXAMPLE 3B (53.8 g) and ether (400 mL), was added methanol (25 mL) slowly by syringe. The mixture was stirred at room temperature for 24 hours. The reaction was quenched with 1N HCl with ice-cooling. The mixture was diluted with water and extracted with ether (3x 100 mL). The extracts were dried (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 0-30% ethyl acetate/hexanes.

EXAMPLE 3D

tert-butyl 4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazine-1-carboxylate

- 5 **[0344]** Mesyl Chloride (7.5 mL) was added via syringe to EXAMPLE 3C (29.3 g) and triethylamine (30 mL) in CH₂Cl₂ (500 mL) at 0°C, and the mixture was stirred for 1 minute. N-t-butoxycarbonylpiperazine (25 g) was added and the mixture was stirred at room temperature for 24 hours. The suspension was washed with brine, dried, (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 10-20% ethyl acetate/hexanes.

10 EXAMPLE 3E

1-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazine

- 15 **[0345]** EXAMPLE 3D (1 g) was stirred in dichloromethane (10 mL), trifluoroacetic acid (10 mL), and triethylsilane (1 mL) for 1 hour. The mixture was concentrated, taken up in a mixture of dichloromethane (100 mL) and saturated aqueous Na₂CO₃ solution (20 mL) and stirred for 10 minutes. The layers were separated, and the organic layer was dried over Na₂SO₄, filtered, and concentrated to give the product.

EXAMPLE 3F

20

5-bromo-1-(triisopropylsilyl)-1H-pyrrolo[2,3-b]pyridine

- 25 **[0346]** To a mixture of 5-bromo-1H-pyrrolo[2,3-b]pyridine (15.4 g) in tetrahydrofuran (250 mL) was added 1M lithium hexamethyldisilazide in tetrahydrofuran (86 mL), and after 10 minutes, TIPS-Cl (triisopropylchlorosilane) (18.2 mL) was added. The mixture was stirred at room temperature for 24 hours. The reaction was diluted with ether, and the resulting solution was washed twice with water. The extracts were dried (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 10% ethyl acetate/hexanes.

EXAMPLE 3G

30

1-(triisopropylsilyl)-1H-pyrrolo[2,3-b]pyridin-5-ol

- 35 **[0347]** To a mixture of EXAMPLE 3F (24.3 g) in tetrahydrofuran (500 mL) at -78°C was added 2.5M BuLi (30.3 mL). After 2 minutes, trimethylborate (11.5 mL) was added, and the mixture was allowed to warm to room temperature over 1 hour. The reaction was poured into water, extracted three times with ethyl acetate, and the combined extracts were washed with brine and concentrated. The crude product was taken up in tetrahydrofuran (200 mL) at 0°C, and 1M NaOH (69 mL) was added, followed by 30% H₂O₂ (8.43 mL), and the solution was stirred for 1 hour. Na₂S₇O₃ (10 g) was added, and the pH was adjusted to 4-5 with concentrated HCl and solid NaH₂PO₄. The solution was extracted twice with ethyl acetate, and the combined extracts were washed with brine, dried (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 5-25% ethyl acetate/hexanes.

EXAMPLE 3H

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-fluorobenzoate

45

- [0348]** A mixture of EXAMPLE 3G (8.5 g), methyl 2,4-difluorobenzoate (7.05 g), and K₃PO₄ (9.32 g) in diglyme (40 mL) at 115°C was stirred for 24 hours. The reaction was cooled, diluted with ether (600 mL), and washed twice with water, and brine, and concentrated. The crude product was chromatographed on silica gel with 2-50% ethyl acetate/hexanes.

50

EXAMPLE 3I

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoate

55

- [0349]** A mixture of EXAMPLE 3H (1.55 g), EXAMPLE 3E (2.42 g), and HK₂PO₄ (1.42 g) in dimethylsulfoxide (20 mL) at 135°C was stirred for 24 hours. The reaction was cooled, diluted with ether (400 mL), and washed with 3x 1M NaOH, and brine, and concentrated. The crude product was chromatographed on silica gel with 10-50% ethyl acetate/hexanes.

EXAMPLE 3J

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoic acid

[0350] EXAMPLE 3I (200 mg) in dioxane (10 mL) and 1M NaOH (6 mL) at 50°C was stirred for 24 hours. The reaction was cooled, added to NaH₂PO₄ solution, and extracted three times with ethyl acetate. The combined extracts were washed with brine, and concentrated to give the pure product.

EXAMPLE 3K

tert-butyl 1-(tetrahydro-2H-pyran-4-yl)piperidin-4-ylcarbamate

[0351] Tert-butyl piperidin-4-ylcarbamate (45.00 g, 225 mmol) and dihydro-2H-pyran-4(3H)-one (24.74 g, 247 mmol) were added to dichloromethane (1000 mL). Sodium triacetoxymethylborohydride (61.90 g, 292 mmol) was added, and the solution was stirred at room temperature for 16 hours. The solution was extracted with 1M sodium hydroxide and dried over anhydrous sodium sulfate. The solution was filtered and concentrated and purified by flash column chromatography on silica gel with 10% methanol (in dichloromethane) increasing to 20% methanol (in dichloromethane).

EXAMPLE 3L

1-(tetrahydro-2H-pyran-4-yl)piperidin-4-amine dihydrochloride

[0352] A solution of EXAMPLE 3K (52.57 g, 185 mmol) in dichloromethane (900 mL) was treated with 4M aqueous HCl (462 mL), and the solution was mixed vigorously at room temperature for 16 hours. Solvent was removed under vacuum to give crude product as the dihydrochloride salt, which was used without further purification.

EXAMPLE 3M

3-nitro-4-(1-(tetrahydro-2H-pyran-4-yl)piperidin-4-ylamino)benzenesulfonamide

[0353] EXAMPLE 3L (22.12 g, 86 mmol) was added to 1,4-dioxane (300 mL) and water (43 mL). Triethylamine (43.6 mL, 31.6 g, 313 mmol) was added, and the mixture was stirred at room temperature until EXAMPLE 3L had completely dissolved. 4-chloro-3-nitrobenzenesulfonamide was added and the mixture was heated at 90°C for 16 hours. The mixture was cooled, and the solvents were removed under vacuum. 10% methanol (in dichloromethane) was added and the solution was stirred vigorously at room temperature until a fine suspension was obtained. The solid was isolated by vacuum filtration and washed with dichloromethane to give pure product.

EXAMPLE 3N

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([1-(tetrahydro-2H-pyran-4-yl)piperidin-4-yl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0354] This EXAMPLE was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 3M for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.65 (br s, 1H), 8.53 (br s, 1H), 8.18 (m, 1H), 8.00 (br s, 1H), 7.63 (m, 1H), 7.49 (m, 3H), 7.34 (d, 2H), 7.12 (m, 1H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.37 (d, 1H), 6.20 (d, 1H), 3.95 (m, 2H), 3.05 (m, 10H), 2.73 (m, 4H), 2.17 (m, 10H), 1.95 (m, 2H), 1.80 (m, 2H), 1.63 (m, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 4

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-(1-methylpiperidin-4-yl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 4A

4-(1-methylpiperidin-4-ylamino)-3-nitrobenzenesulfonamide

[0355] This EXAMPLE was prepared by substituting 4-amino-N-methylpiperidine for 1-(tetrahydropyran-4-yl)methyl-

amine in EXAMPLE 1F.

EXAMPLE 4B

5 4-(4-{ [2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1-methylpiperidin-4-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 **[0356]** This EXAMPLE was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 4A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.65 (brs, 1H), 8.55 (br s, 1H), 8.17 (m, 1H), 8.02 (d, 1H), 7.85 (dd, 1H), 7.51 (m, 3H), 7.35 (m, 2H), 7.18 (dd, 1H), 7.05 (d, 2H), 6.68 (dd, 1H), 6.38 (d, 1H), 6.20 (d, 1H), 3.90 (m, 1H), 3.09 (m, 8H), 2.77 (m, 2H), 2.05-2.30 (m, 10H), 1.95 (s, 3H), 1.39 (t, 2H), 1.24 (m, 2H), 0.93 (s, 6H).

EXAMPLE 5

15 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 5A

20 3-nitro-4-((tetrahydro-2H-pyran-4-yl)methylamino)benzenesulfonamide

25 **[0357]** A mixture of 4-fluoro-3-nitrobenzenesulfonamide (2.18 g), 1-(tetrahydropyran-4-yl)methylamine (1.14 g), and triethylamine (1 g) in tetrahydrofuran (30 mL) were stirred overnight, neutralized with concentrated HCl and concentrated. The residue was suspended in ethyl acetate and the precipitates were collected, washed with water and dried to provide the title compound.

EXAMPLE 5B

30 methyl 4,4-dimethyl-2-(trifluoromethylsulfonyloxy)cyclohex-1-enecarboxylate

35 **[0358]** To a suspension of hexane washed NaH (17 g) in dichloromethane (700 mL) was added 5,5-dimethyl-2-methoxycarbonylcyclohexanone (38.5 g) dropwise at 0°C. After stirring for 30 minutes, the mixture was cooled to -78°C and trifluoroacetic anhydride (40 mL) was added. The reaction mixture was warmed to room temperature and stirred for 24 hours. The organic layer was washed with brine, dried (Na₂SO₄), filtered, and concentrated to give the product.

EXAMPLE 5C

methyl 2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enecarboxylate

40 **[0359]** EXAMPLE 5B (62.15g), 4-chlorophenylboronic acid (32.24 g), CsF (64 g) and tetrakis(triphenylphosphine)palladium(0) (2g) in 2:1 dimethoxyethane /methanol (600 mL) were heated to 70°C for 24 hours. The mixture was concentrated. Ether (4x 200 mL) was added and the mixture was filtered. The combined ether solution was concentrated to give the product.

EXAMPLE 5D

(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methanol

50 **[0360]** To a mixture of LiBH₄ (13g), EXAMPLE 5C (53.8 g) and ether (400 mL), was added methanol (25 mL) slowly by syringe. The mixture was stirred at room temperature for 24 hours. The reaction was quenched with 1N HCl with ice-cooling. The mixture was diluted with water and extracted with ether (3x 100 mL). The extracts were dried (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 0-30% ethyl acetate/hexanes.

55

EXAMPLE 5E

tert-butyl 4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazine-1-carboxylate

- 5 **[0361]** Mesyl Chloride (7.5 mL) was added via syringe to EXAMPLE 5D (29.3 g) and triethylamine (30 mL) in CH₂Cl₂ (500 mL) at 0°C, and the mixture was stirred for 1 minute. N-t-butoxycarbonylpiperazine (25 g) was added and the mixture was stirred at room temperature for 24 hours. The suspension was washed with brine, dried, (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 10-20% ethyl acetate/hexanes.

10 EXAMPLE 5F

1-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazine

- 15 **[0362]** EXAMPLE 5E (200 mg) and triethylsilane (1 mL) were stirred in dichloromethane (15 mL) and trifluoroacetic acid (15 mL) for 1 hour. The mixture was concentrated, taken up in ethyl acetate, washed twice with NaH₂PO₄, and brine, and dried (Na₂SO₄), filtered and concentrated.

EXAMPLE 5G

- 20 5-bromo-1-(triisopropylsilyl)-1H-pyrrolo[2,3-b]pyridine

- [0363]** To a mixture of 5-bromo-1H-pyrrolo[2,3-b]pyridine (15.4 g) in tetrahydrofuran (250 mL) was added 1M lithium hexamethyldisilazide in tetrahydrofuran (86 mL), and after 10 minutes, TIPS-Cl (triisopropylchlorosilane) (18.2 mL) was added. The mixture was stirred at room temperature for 24 hours. The reaction was diluted with ether, and the resulting solution was washed twice with water. The extracts were dried (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 10% ethyl acetate/hexanes.

EXAMPLE 5H

- 30 1-(triisopropylsilyl)-1H-pyrrolo[2,3-b]pyridin-5-ol

- [0364]** To a mixture of EXAMPLE 5G (24.3 g) in tetrahydrofuran (500 mL) at -78°C was added 2.5M BuLi (30.3 mL). After 2 minutes, trimethylborate (11.5 mL) was added, and the mixture was allowed to warm to room temperature over 1 hour. The reaction was poured into water, extracted three times with ethyl acetate, and the combined extracts were washed with brine and concentrated. The crude product was taken up in tetrahydrofuran (200 mL) at 0°C, and 1M NaOH (69 mL) was added, followed by 30% H₂O₂ (8.43 mL), and the solution was stirred for 1 hour. Na₂S₂O₃ (10 g) was added, and the pH was adjusted to 4-5 with concentrated HCl and solid NaH₂PO₄. The solution was extracted twice with ethyl acetate, and the combined extracts were washed with brine, dried (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 5-25% ethyl acetate/hexanes.

EXAMPLE 5I

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-fluorobenzoate

- 45 **[0365]** A mixture of EXAMPLE 5H (8.5 g), methyl 2,4-difluorobenzoate (7.05 g), and K₃PO₄ (9.32 g) in diglyme (40 mL) at 115°C was stirred for 24 hours. The reaction was cooled, diluted with ether (600 mL), and washed twice with water, and brine, and concentrated. The crude product was chromatographed on silica gel with 2-50% ethyl acetate/hexanes.

50 EXAMPLE 5J

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoate

- 55 **[0366]** A mixture of EXAMPLE 5I (1.55 g), EXAMPLE 5F (2.42 g), and HK₂PO₄ (1.42 g) in dimethylsulfoxide (20 mL) at 135°C was stirred for 24 hours. The reaction was cooled, diluted with ether (400 mL), and washed with 3x 1M NaOH, and brine, and concentrated. The crude product was chromatographed on silica gel with 10-50% ethyl acetate/hexanes.

EXAMPLE 5K

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoic acid

5

[0367] EXAMPLE 5J (200 mg) in dioxane (10 mL) and 1M NaOH (6 mL) at 50°C was stirred for 24 hours. The reaction was cooled, added to NaH₂PO₄ solution, and extracted three times with ethyl acetate. The combined extracts were washed with brine, and concentrated to give the pure product.

10 EXAMPLE 5L

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

15 **[0368]** EXAMPLE 5K (3.39 g), EXAMPLE 5A (1.87 g), 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (2.39 g), and 4-dimethylaminopyridine (1.09 g) were stirred in CH₂Cl₂ (40 mL) for 24 hours. The reaction was cooled and chromatographed on silica gel with 25-100% ethyl acetate/hexanes, then 10% methanol/ethyl acetate with 1% acetic acid, to give the product (1.62 g, 32%) as a white solid. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.65 (brs, 1H), 8.55 (br s, 1H), 8.04 (d, 1H), 7.89 (dd, 1H), 7.51 (m, 3H), 7.33 (d, 2H), 7.08 (m, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.39 (d, 1H), 20 6.19 (d, 1H), 3.84 (m, 1H), 3.30 (m, 4H), 3.07 (m, 4H), 2.73 (m, 2H), 2.18 (m, 6H), 1.95 (m, 2H), 1.61 (dd, 2H), 1.38 (m, 2H), 1.24 (m, 4H), 0.92 (s, 6H).

EXAMPLE 6

25 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({4-[(4-methylpiperazin-1-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 6A

30 4-(4-methylpiperazin-1-ylamino)-3-nitrobenzenesulfonamide

[0369] A 50 mL round-bottomed flask was charged with 4-chloro-3-nitrobenzenesulfonamide (1 g, 4.23 mmol), 4-methylpiperazin-1-amine dihydrochloride (1 g, 5.32 mmol), and N¹,N¹,N²,N²-tetramethylethane-1,2-diamine (3 mL, 20.01 mmol) in dioxane (10 mL). The reaction mixture was refluxed for 12 hours. After this time, the reaction mixture was cooled to room temperature, the salt filtered off via a Buchner funnel, and the solvent removed in vacuo. The crude product was added to a silica gel column (Analogix, SF65-200g) and purified by eluting with 0-5% methanol in dichloromethane.

35

EXAMPLE 6B

40

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({4-[(4-methylpiperazin-1-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0370] This EXAMPLE was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 6A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.65 (brs, 1H), 9.09 (br s, 1H), 8.47 (d, 1H), 8.24 (dd, 1H), 7.99 (d, 1H), 7.50 (m, 4H), 7.34 (d, 2H), 7.04 (d, 2H), 6.64 (dd, 1H), 6.35 (d, 1H), 6.20 (d, 1H), 3.04 (m, 4H), 2.89 (m, 4H), 2.73 (m, 2H), 2.34 (s, 3H), 2.17 (m, 6H), 1.95 (br s, 2H), 1.38 (t, 2H), 1.05 (m, 4H), 0.93 (s, 6H).

45

EXAMPLE 7

50

2-(9H-carbazol-4-yloxy)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({4-[(1-methylpiperidin-4-yl)amino]-3-nitrophenyl}sulfonyl)benzamide

EXAMPLE 7A

55

ethyl 2-(9H-carbazol-4-yloxy)-4-fluorobenzoate

[0371] This EXAMPLE was prepared by substituting ethyl 2,4-difluorobenzoate for methyl 2,4-difluorobenzoate and

4-hydroxycarbazole for EXAMPLE 3G in EXAMPLE 3H.

EXAMPLE 7B

ethyl 2-(9H-carbazol-4-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoate

[0372] This EXAMPLE was prepared by substituting EXAMPLE 7A for EXAMPLE 3H in EXAMPLE 3I.

EXAMPLE 7C

10

2-(9H-carbazol-4-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoic acid

[0373] This EXAMPLE was prepared by substituting EXAMPLE 7B for EXAMPLE 3I in EXAMPLE 3J, except here upon completion of the reaction, water and 2N HCl were added to adjust the pH to 2, and the HCl salt of the product was extracted using $\text{CHCl}_3/\text{CH}_3\text{OH}$.

15

EXAMPLE 7D

2-(9H-carbazol-4-yloxy)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(1-methylpiperidin-4-yl)amino]-3-nitrophenyl)sulfonyl)benzamide

20

[0374] This EXAMPLE was prepared by substituting EXAMPLE 7C for EXAMPLE 1E and EXAMPLE 4A for EXAMPLE 1F in EXAMPLE 1G, except here the purification was done by preparative HPLC using a C18 column, 250 x 50 mm, 10 μ , and eluting with a gradient of 20-100% CH_3CN vs. 0.1% trifluoroacetic acid in water, giving the product as a bistrifluoroacetate salt. ^1H NMR (300 MHz, dimethylsulfoxide- d_6) δ 11.82 (br s, 1H), 11.40 (s, 1H), 9.70, 9.40 (both v br s, total 2H), 8.40 (d, 1H), 8.10 (br d, 1H), 7.90 (br d, 1H), 7.72 (dd, 1H), 7.60 (d, 1H), 7.48 (d, 1H), 7.38 (m, 3H), 7.22 (m, 2H), 7.07 (m, 4H), 6.78 (dd, 1H), 6.43 (dd, 1H), 6.19 (s, 1H), 3.97 (m, 1H), 3.80 (m, 2H), 3.60, 3.30, 3.10, 2.80 (all br m, total 11H), 2.20, 2.10, 2.00 (all br m, total 8H), 1.78 (m, 2H), 1.42 (m, 2H), 1.25 (m, 2H), 0.92 (s, 6H).

25

EXAMPLE 8

30

2-(9H-carbazol-4-yloxy)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-[(3-pyrrolidin-1-yl)propyl]amino]phenyl)sulfonyl)benzamide

EXAMPLE 8A

35

3-nitro-4-(3-(pyrrolidin-1-yl)propylamino)benzenesulfonamide

[0375] This EXAMPLE was prepared by substituting 3-(pyrrolidin-1-yl)propan-1-amine for 1-(tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

40

EXAMPLE 8B

2-(9H-carbazol-4-yloxy)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-[(3-pyrrolidin-1-yl)propyl]amino]phenyl)sulfonyl)benzamide

45

[0376] This EXAMPLE was prepared by substituting EXAMPLE 7C for EXAMPLE 1E and EXAMPLE 8A for EXAMPLE 1F in EXAMPLE 1G, except here the purification was done by preparative HPLC using a C18 column, 250 x 50 mm, 10 μ , and eluting with a gradient of 20-100% CH_3CN vs. 0.1% trifluoroacetic acid in water, giving the product as a bistrifluoroacetate salt. ^1H NMR (300 MHz, dimethylsulfoxide- d_6) δ 11.80 (br s, 1H), 11.42 (s, 1H), 9.50, 9.25 (both v br s, total 2H), 8.58 (br t, 1H), 8.43 (d, 1H), 7.91 (d, 1H), 7.72 (dd, 1H), 7.60 (d, 1H), 7.50 (d, 1H), 7.38 (m, 3H), 7.23 (m, 2H), 7.07 (m, 3H), 6.93 (d, 1H), 6.78 (dd, 1H), 6.44 (dd, 1H), 6.18 (s, 1H), 3.70, 3.60, 3.20, 3.00 (all br m, total 18H), 2.18 (br m, 2H), 2.00-1.80 (envelope, 8H), 1.42 (m, 2H), 0.92 (s, 6H).

50

55

EXAMPLE 9

Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 9A

Trans-tert-butyl 4-morpholinocyclohexylcarbamate

[0377] A solution of tert-butyl-4-aminocyclohexylcarbamate (20.32 g, 95 mmol), bis(2-bromoethyl) ether (14.30 ml, 114 mmol) and triethylamine (33.0 ml, 237 mmol) in N,N-dimethylformamide (200 ml) was stirred for 16 hours at 70 °C. The reaction mixture was cooled down to room temperature, concentrated and the product was extracted with ethyl acetate. The organic layer was washed with sodium carbonate solution (15% aq.), dried and concentrated. The product was used in next step without purification.

EXAMPLE 9B

Trans-4-morpholinocyclohexanamine dihydrochloride

[0378] To a solution of trans-tert-butyl-4-morpholinocyclohexylcarbamate (19.2 g, 67.5 mmol) in dichloromethane (100 ml) was added HCl (100 ml, 400 mmol) (4M in dioxane) and the reaction mixture was stirred for 16 hours at room temperature. The reaction mixture was diluted with ether and solid salt was filtered off, and dried in an oven.

EXAMPLE 9C

Trans-4-(4-morpholinocyclohexylamino)-3-nitrobenzenesulfonamide

[0379] A solution of trans-4-morpholinocyclohexanamine dihydrochloride (5 g, 19.44 mmol), 4-fluoro-3-nitrobenzenesulfonamide (4.32 g, 19.63 mmol) and triethylamine (20 ml, 143 mmol) in tetrahydrofuran (60 ml) was stirred for 16 hours at room temperature. The solid product was filtered off, washed with tetrahydrofuran, ether, dichloromethane (3 x) and dried under vacuum.

EXAMPLE 9D

Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0380] This EXAMPLE was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 9C for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.61 (brs, 1H), 8.49 (br s, 1H), 8.12 (m, 1H), 7.99 (br s, 1H), 7.71 (m, 1H), 7.50 (m, 3H), 7.35 (d, 2H), 7.04 (d, 2H), 7.01 (m, 1H), 6.65 (dd, 1H), 6.36 (d, 1H), 6.21 (d, 1H), 3.60 (m, 4H), 3.04 (m, 4H), 2.73 (m, 2H), 2.57 (m, 2H), 2.42 (m, 1H), 2.18 (m, 6H), 2.05 (m, 2H), 1.95 (m, 2H), 1.90 (m, 2H), 1.38 (m, 6H), 1.15 (m, 3H), 0.92 (s, 6H).

EXAMPLE 10

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({4-[(2-methoxyethyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 10A

4-(2-methoxyethylamino)-3-nitrobenzenesulfonamide

[0381] This EXAMPLE was prepared by substituting 2-methoxyethylamine for 1-(tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 10B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((2-methoxyethyl)amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0382] This EXAMPLE was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 10A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.65 (brs, 1H), 8.58-8.49 (m, 1H), 8.55 (d, 1H), 8.03 (d, 1H), 7.79 (m, 1H), 7.49 (m, 3H), 7.34 (m, 2H), 7.06 (m, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.38 (m, 1H), 6.20 (d, 1H), 3.61-3.51 (m, 4H), 3.31 (s, 3H), 3.07 (m, 4H), 2.74 (m, 2H), 2.17 (m, 6H), 1.95 (br s, 2H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 11

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(3-nitro-4-([(3S)-tetrahydro-2H-pyran-3-ylmethyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 11A

(S)-3-nitro-4-((tetrahydro-2H-pyran-3-yl)methylamino)benzenesulfonamide and (R)-3-nitro-4-((tetrahydro-2H-pyran-3-yl)methylamino)benzenesulfonamide

[0383] This EXAMPLE was prepared by substituting (tetrahydro-2H-pyran-3-yl)methanamine for 1-(tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 11B

(S)-3-nitro-4-((tetrahydro-2H-pyran-3-yl)methylamino)benzenesulfonamide

[0384] The racemic mixture of EXAMPLE 11A was resolved by chiral SFC on an AD column (21mm i.d.x 250 mm in length) using a gradient of 10-30% 0.1% diethylamine methanol in CO₂ over 15 minutes (oven temperature: 40°C; flow rate: 40 mL/minute) to provide the title compound.

EXAMPLE 11C

(R)-3-nitro-4-((tetrahydro-2H-pyran-3-yl)methylamino)benzenesulfonamide

[0385] The racemic mixture of EXAMPLE 11A was resolved by chiral SFC on an AD column (21mm i.d.x 250 mm in length) using a gradient of 10-30% 0.1% diethylamine methanol in CO₂ over 15 minutes (oven temperature: 40°C; flow rate: 40 mL/minute) to provide the title compound.

EXAMPLE 11D

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(3-nitro-4-([(3S)-tetrahydro-2H-pyran-3-ylmethyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0386] To a mixture of EXAMPLE 3J (59.8 mg, 0.105 mmol), EXAMPLE 11B (33mg, 0.105 mmol) and N,N-dimethylpyridin-4-amine (38.4 mg, 0.314 mmol) in dichloromethane (5 ml) was added 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (24.07 mg, 0.13 mmol). The reaction mixture was stirred at room temperature overnight and concentrated. The residue was purified by reverse phase HPLC on a C18 column using a gradient of 40-60% acetonitrile/0.1% trifluoroacetic acid in water to give the title compound as the trifluoroacetate salt. The trifluoroacetic acid salt was dissolved in dichloromethane (6 ml) and washed with 50% aqueous NaHCO₃. The organic layer was dried over anhydrous Na₂SO₄ and concentrated to give the title compound. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 11.40 (s, br, 1H), 8.53 - 8.58 (m, 2H), 8.04 (d, 1H), 7.80 (dd, 1H), 7.47 - 7.54 (m, 3H), 7.34 (d, 2H), 7.02 - 7.09 (m, 3H), 6.67 (dd, 1H), 6.39 (dd, 1H), 6.19 (d, 1H), 3.79 (dd, 1H), 3.69 - 3.73 (m, 1H), 3.22 - 3.37 (m, 3H), 3.16 - 3.21 (m, 1H), 3.07 (s, 4H), 2.74 (s, 2H), 2.09 - 2.24 (m, 6H), 1.95 (s, 2H), 1.86 - 1.93 (m, 1H), 1.79 - 1.85 (m, 1H), 1.58 - 1.64 (m, 1H), 1.42 - 1.51 (m, 1H), 1.38 (t, 2H), 1.25 - 1.34 (m, 1H), 0.92 (s, 6H).

EXAMPLE 12

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 12A

4-((1,4-dioxan-2-yl)methoxy)-3-nitrobenzenesulfonamide

[0387] (1,4-Dioxan-2-yl)methanol (380 mg, 3.22 mmol) in tetrahydrofuran (30 ml) was treated with sodium hydride (60%) (245 mg, 6.13 mmol) at room temperature for 30 minutes. The reaction mixture was cooled in an ice bath and 4-fluoro-3-nitrobenzenesulfonamide (675 mg, 3.06 mmol) was added. The resulting mixture was stirred at room temperature for 2 hours and another portion of sodium hydride (60%) (245 mg, 6.13 mmol) was added. The reaction mixture was stirred overnight and quenched with ice water (3 ml). The cloudy mixture was filtered and the filtrate was concentrated. The residue was triturated with methanol to give the title compound.

EXAMPLE 12B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0388] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 12A in place of EXAMPLE 11B. ¹H NMR (500 MHz, dimethylsulfoxide -d₆) δ 11.67 (s, 1 H), 11.42 (s, br, 1 H), 8.34 (s, 1 H), 8.03 (d, 2 H), 7.48 - 7.55 (m, 3 H), 7.41 (d, 1 H), 7.35 (d, 2 H), 7.04 (d, 2 H), 6.67 (dd, 1 H), 6.39 (dd, 1 H), 6.21 (d, 1 H), 4.20 - 4.28 (m, 2 H), 3.85 - 3.91 (m, 1 H), 3.82 (dd, 1 H), 3.74 - 3.78 (m, 1 H), 3.59 - 3.69 (m, 2 H), 3.41 - 3.51 (m, 2 H), 3.05 - 3.17 (m, 4 H), 2.83 (s, br, 2 H), 2.27 (s, br, 4 H), 2.15 (s, 2 H), 1.96 (s, 2 H), 1.39 (t, 2 H), 0.93 (s, 6 H).

EXAMPLE 13

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[3-nitro-4-[[3R]-tetrahydro-2H-pyran-3-ylmethyl]amino}phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0389] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 11C in place of EXAMPLE 11B. The proton NMR spectra of EXAMPLE 13 and EXAMPLE 11D are identical. ¹H NMR (500 MHz, dimethylsulfoxide -d₆) δ 11.68 (s, 1 H), 11.40 (s, br, 1 H), 8.53 - 8.58 (m, 2 H), 8.04 (d, 1 H), 7.80 (dd, 1 H), 7.47 - 7.54 (m, 3 H), 7.34 (d, 2 H), 7.02 - 7.09 (m, 3 H), 6.67 (dd, 1 H), 6.39 (dd, 1 H), 6.19 (d, 1 H), 3.79 (dd, 1 H), 3.69 - 3.73 (m, 1 H), 3.22 - 3.37 (m, 3 H), 3.16 - 3.21 (m, 1 H), 3.07 (s, 4 H), 2.74 (s, 2 H), 2.09 - 2.24 (m, 6 H), 1.95 (s, 2 H), 1.86 - 1.93 (m, 1 H), 1.79 - 1.85 (m, 1 H), 1.58 - 1.64 (m, 1 H), 1.42 - 1.51 (m, 1 H), 1.38 (t, 2 H), 1.25 - 1.34 (m, 1 H), 0.92 (s, 6 H).

EXAMPLE 14

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-(2-naphthylsulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0390] The title compound was prepared as described in EXAMPLE 11D using naphthalene-2-sulfonamide (47 mg, 0.227 mmol) in place of EXAMPLE 11B. ¹H NMR (500 MHz, dimethylsulfoxide -d₆) δ 11.82 (s, 1 H), 11.69 (s, 1 H), 8.51 (s, 1 H), 8.08 (d, 1 H), 8.05 (d, 1 H), 7.97 (dd, 2 H), 7.82 (dd, 1 H), 7.66 - 7.71 (m, 1 H), 7.63 (t, 1 H), 7.54 (d, 1 H), 7.47 - 7.52 (m, 2 H), 7.34 (d, 2 H), 7.04 (d, 2 H), 6.65 (dd, 1 H), 6.39 (dd, 1 H), 6.18 (s, 1 H), 3.04 (s, 4 H), 2.72 (s, 2 H), 2.10 - 2.20 (m, 6 H), 1.95 (s, 2 H), 1.38 (t, 2 H), 0.92 (s, 6 H).

EXAMPLE 15

4-(4-{[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl}piperazin-1-yl)-N-({[3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 15A

methyl 6,6-dimethyl-4-oxotetrahydro-2H-pyran-3-carboxylate

[0391] To a suspension of hexane-washed NaH (0.72 g, 60% in mineral oil) in tetrahydrofuran (30 mL) was added a solution of 2,2-dimethyldihydro-2H-pyran-4(3H)-one (2.0 g) in tetrahydrofuran (20 mL). The suspension was stirred at room temperature for 30 minutes. The dimethylcarbonate (6.31 mL) was added dropwise by syringe. The mixture was heated to reflux for 4h. LC/MS showed the expected product as the major product. The mixture was acidified with 5% HCl and extracted with dichloromethane (100 mL x 3) and washed with water, brine and dried over Na₂SO₄. After evaporation, the crude product was loaded on a column and eluted with 10% ethyl acetate in hexane to give the product.

EXAMPLE 15B

methyl 6,6-dimethyl-4-(trifluoromethylsulfonyloxy)-5,6-dihydro-2H-pyran-3-carboxylate

[0392] To a cooled (0°C) stirring suspension of NaH (0.983 g, 60% in mineral oil) in ether (50 mL) was added EXAMPLE 15A (3.2 g). The mixture was stirred at 0°C for 30 minutes before the addition of Tf₂O (4.2 mL). The mixture was then stirred at room temperature overnight. The mixture was diluted with ether (200 mL) and washed with 5% HCl, water and brine. After drying over Na₂SO₄, evaporation of solvent gave the crude product which was used in the next step without further purification.

EXAMPLE 15C

methyl 4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-carboxylate

[0393] To a solution of EXAMPLE 15B (2.88 g), 4-chlorophenylboronic acid (1.88 g) and Pd(Ph₃P)₄ (0.578 g) in toluene (40 mL) and ethanol (10 mL) was added 2N Na₂CO₃ (10 mL). The mixture was stirred at reflux overnight. The mixture was diluted ether (300 mL) and washed with water, brine and dried over Na₂SO₄. After evaporation of solvent, the residue was loaded on a column and eluted with 3% ethyl acetate in hexane to give the product.

EXAMPLE 15D

(4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl)methanol

[0394] To a solution of EXAMPLE 15C (1.6 g) in ether (20 mL) was added LiAlH₄ (1.2 g). The mixture was stirred for 4 hours. The mixture was acidified carefully with 5% HCl and extracted with ethyl acetate (100 mL x 3) and washed with water, brine and dried over Na₂SO₄. After concentration, the crude product was loaded on a column and eluted with 10% ethyl acetate in hexane to give the product.

EXAMPLE 15E

4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-carbaldehyde

[0395] To a solution of oxalyl chloride (1.1 g) in dichloromethane (30 mL) at -78°C was added dimethylsulfoxide (6.12 mL). The mixture was stirred at the temperature for 30 minutes, and then a solution of EXAMPLE 15D (1.2 g) in dichloromethane (10 mL) was added. The mixture was stirred at -78°C for 2 hours before the addition of triethylamine (10 mL). The mixture was stirred overnight and the temperature was allowed to rise to room temperature. The mixture was diluted with ether (300 mL) and washed with water, brine and dried over Na₂SO₄. Concentration of the solvent and column purification (5% ethyl acetate in hexane) gave the product.

EXAMPLE 15F

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(piperazin-1-yl)benzoate

- 5 **[0396]** A mixture of EXAMPLE 3H (20.5 g) and piperazine (37.0 g) in dimethylsulfoxide (200 mL) was heated to 110°C for 24 hours, and the mixture was allowed to cool to room temperature. The mixture was poured into water (1 L), extracted three times with dichloromethane, and the combined extracts were washed with 2x water, and brine and filtered and concentrated to give the pure product.

10 EXAMPLE 15G

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-((4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl)methyl)piperazin-1-yl)benzoate

- 15 **[0397]** To a solution of EXAMPLE 15E (100 mg) and EXAMPLE 15F (177 mg) in dichloromethane (10 mL) was added sodium triacetoxymethylborohydride (154 mg). The mixture was stirred overnight. The mixture was diluted with ethyl acetate (200 mL) and washed with 2% NaOH, water and brine. After drying over Na₂SO₄, the mixture was filtered and the solvent was evaporated under vacuum. The residue was loaded on a column and eluted with 30% ethyl acetate in hexane to give the pure product.

20

EXAMPLE 15H

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-((4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl)methyl)piperazin-1-yl)benzoic acid

25

- [0398]** To a solution of EXAMPLE 15G (254 mg) in tetrahydrofuran (4 mL), methanol (2 mL) and water (2 mL) was added LiOH H₂O (126 mg). The mixture was stirred overnight. The mixture was then neutralized with 5% HCl and diluted with ethyl acetate (200 mL). After washing with brine, it was dried over Na₂SO₄. Filtration and evaporation of solvent gave the product.

30

EXAMPLE 15I

4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

35

- [0399]** The title compound was prepared as described in EXAMPLE 1G, substituting EXAMPLE 1E with EXAMPLE 15H. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.68 (br s, 1H), 11.42 (s, 1H), 8.60 (m, 1H), 8.57 (d, 1H), 8.05 (d, 1H), 7.80 (dd, 1H), 7.48-7.54 (m, 3H), 7.38 (d, 2H), 7.12 (m, 3H), 6.68 (dd, 1H), 6.40 (dd, 1H), 6.20 (s, 1H), 4.11 (s, 2H), 3.85 (m, 2H), 3.27 (m, 6H), 3.07 (m, 2H), 2.84 (m, 2H), 2.14 (m, 5H), 1.92 (m, 1H), 1.42 (m, 2H), 1.24 (m, 2H), 1.10 (s, 6H).

40

EXAMPLE 16

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[(2-methoxyethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

45

EXAMPLE 16A

4-(2-methoxyethylamino)-3-(trifluoromethylsulfonyl)benzenesulfonamide

- 50 **[0400]** 4-Fluoro-3-(trifluoromethylsulfonyl)benzenesulfonamide (1.536 g, 5 mmol), 2-methoxyethanamine (0.376 g, 5 mmol), and triethylamine (1.939 g, 15 mmol) in anhydrous tetrahydrofuran (30 mL) solution was heated at 55°C for 3 hours. The solution was diluted with ethyl acetate, washed with water and brine, and dried (Na₂SO₄), filtered and the filtrate was concentrated. The crude material was used in the next step without further purification.

55

EXAMPLE 16B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(2-methoxyethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0401] This EXAMPLE was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 16A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.67 (brs, 1H), 8.14 (m, 1H), 8.03 (d, 1H), 7.91 (d, 1H), 7.50 (m, 3H), 7.34 (d, 2H), 7.19 (s, 1H), 7.04 (m, 3H), 6.67 (dd, 1H), 6.39 (m, 1H), 6.19 (d, 1H), 3.51 (m, 4H), 3.28 (s, 3H), 3.06 (m, 4H), 2.75 (m, 2H), 2.17 (m, 6H), 1.95 (m, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 17

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)benzamide

EXAMPLE 17A

4-((tetrahydro-2H-pyran-4-yl)methylamino)-3-(trifluoromethylsulfonyl)benzenesulfonamide

[0402] This EXAMPLE was prepared by substituting 1-(tetrahydropyran-4-yl)methylamine for 2-methoxyethanamine in EXAMPLE 16A.

EXAMPLE 17B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)benzamide

[0403] This EXAMPLE was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 17A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.70 (brs, 1H), 8.15 (m, 1H), 8.04 (d, 1H), 7.92 (d, 1H), 7.51 (m, 3H), 7.34 (d, 2H), 7.19 (s, 1H), 7.05 (m, 3H), 6.68 (dd, 1H), 6.40 (m, 1H), 6.18 (d, 1H), 3.85 (m, 2H), 3.25 (m, 4H), 3.07 (m, 4H), 2.77 (m, 2H), 2.17 (m, 6H), 1.95 (m, 2H), 1.84 (m, 1H), 1.54 (m, 2H), 1.39 (t, 2H), 1.24 (m, 2H), 0.93 (s, 6H).

EXAMPLE 18

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-indol-5-yloxy)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)benzamide

EXAMPLE 18A

methyl 2-(1H-indol-5-yloxy)-4-fluorobenzoate

[0404] A mixture of 5-hydroxyindole (8.5 g), methyl 2,4-difluorobenzoate (7.05 g), and K₃PO₄ (9.32 g) in diglyme (40 mL) at 115°C was stirred for 24 hours. The reaction was cooled, diluted with ether (600 mL), and washed twice with water, and brine, and concentrated. The crude product was chromatographed on silica gel with 2-50% ethyl acetate/hexanes.

EXAMPLE 18B

methyl 2-(1H-indol-5-yloxy)-4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoate

[0405] A mixture of EXAMPLE 18A (1.7 g), EXAMPLE 3E (1.8 g), and HK₂PO₄ (1.21 g) in dimethylsulfoxide (20 mL) at 135°C was stirred for 24 hours. The reaction was cooled, diluted with ether (400 mL), and washed with 3x 1M NaOH, and brine, and concentrated. The crude product was chromatographed on silica gel with 10-50% ethyl acetate/hexanes.

EXAMPLE 18C

2-(1H-indol-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoic acid

- 5 **[0406]** EXAMPLE 18B (200 mg) in dioxane (10 mL) and 1M NaOH (6 mL) at 50°C was stirred for 24 hours. The reaction was cooled, added to NaH₂PO₄ solution, and extracted three times with ethyl acetate. The combined extracts were washed with brine, and concentrated to give the pure product.

EXAMPLE 18D

10

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-indol-5-yloxy)-N-({3-nitro-4-((tetrahydro-2H-pyran-4-ylmethyl)amino)phenyl)sulfonyl)benzamide

- 15 **[0407]** The title compound was prepared as described in EXAMPLE 11D by replacing EXAMPLE 3J with EXAMPLE 18C, and EXAMPLE 1F for EXAMPLE 11B. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.18 (s, 2H), 8.59-8.64 (m, 2H), 7.80 (dd, 1H), 7.52 (d, 1H), 7.39-7.42 (m, 2H), 7.33 (d, 2H), 7.16 (d, 1H), 7.10 (d, 1H), 7.03 (d, 2H), 6.8 (dd, 1H), 6.65 (dd, 1H), 6.40 (s, 1H), 6.14 (d, 1H), 3.85 (dd, 2H), 3.24-3.32 (m, 4H), 3.03 (s, 3H), 2.73 (s, 2H), 2.12-2.17 (m, 5H), 1.68-1.94 (m, 3H), 1.61 (d, 2H), 1.37 (t, 2H), 1.24-1.27 (m, 2H), 0.92 (s, 6H).

20 EXAMPLE 19

Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-indol-5-yloxy)-N-({4-((4-morpholin-4-ylcyclohexyl)amino)-3-nitrophenyl)sulfonyl)benzamide

- 25 **[0408]** The title compound was prepared as described in EXAMPLE 11D by replacing EXAMPLE 11B with EXAMPLE 9B and EXAMPLE 3J with EXAMPLE 18C. ¹H NMR (500 MHz, pyridine-d₅) δ 12.29 (s, 1H), 9.29 (d, J=2.1 Hz, 1H), 8.37 (d, J=7.6 Hz, 1H), 8.32 (dd, J=9.3, 2.3 Hz, 1H), 8.18 (d, J=8.8 Hz, 1H), 7.52 - 7.57 (m, 2H), 7.39 - 7.47 (m, 3H), 7.10 (dd, J=8.7, 2.3 Hz, 1H), 7.05 - 7.08 (m, 2H), 6.90 (d, J=9.5 Hz, 1H), 6.74 (dd, J=9.0, 2.3 Hz, 1H), 6.59 - 6.63 (m, 1H), 6.55 (d, J=2.4 Hz, 1H), 3.72 - 3.78 (m, 4H), 3.33 - 3.43 (m, 1H), 2.99 - 3.09 (m, 4H), 2.76 (s, 2H), 2.46 - 2.54 (m, 4H), 30 2.16 - 2.29 (m, 3H), 2.09 - 2.14 (m, 4H), 2.05 (d, J=11.9 Hz, 2H), 1.97 (d, J=1.8 Hz, 2H), 1.87 (d, J=11.6 Hz, 2H), 1.19 - 1.42 (m, 6H), 0.93 (s, 6H).

EXAMPLE 20

- 35 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-indol-5-yloxy)-N-({4-((2-methoxyethyl)amino)-3-nitrophenyl)sulfonyl)benzamide

- [0409]** The title compound was prepared as described in EXAMPLE 11D by replacing EXAMPLE 11B with EXAMPLE 10A and EXAMPLE 3J with EXAMPLE 18C. ¹H NMR (400 MHz, dimethylsulfoxide - d₆) δ 11.20 (br. s, 1H) 11.15 (s, 1H) 8.59 (m, 2H) 7.81 (dd, 1H) 7.50 (d, 1H) 7.36 (m, 4H) 7.08 (m, 4H) 6.85 (dd, 1H) 6.65 (dd, 1H) 6.38 (m, 1H) 6.14 (m, 1H) 40 3.58 (m, 4H) 3.30 (s, 3H) 3.03 (m, 4H) 2.73 (s, 2H) 2.15 (m, 6H) 1.96 (s, 2H) 1.38 (t, 2H) 0.92 (s, 6H)

EXAMPLE 21

- 45 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-indol-5-yloxy)-N-([3-nitro-4-((3S)-tetrahydro-2H-pyran-3-ylmethyl)amino]phenyl)sulfonyl)benzamide

- [0410]** The title compound was prepared as described in EXAMPLE 11D by replacing EXAMPLE 3J with EXAMPLE 18C. ¹H NMR (400 MHz, dimethylsulfoxide -d₆) δ 11.17 (s, 2 H), 8.53 - 8.65 (m, 2 H), 7.80 (d, 1 H), 7.51 (d, 1 H), 7.38 - 7.44 (m, 2 H), 7.33 (d, 2 H), 7.15 (s, 1 H), 7.02 - 7.09 (m, 3 H), 6.82 - 6.92 (m, 1 H), 6.65 (d, 1 H), 6.39 (s, 1 H), 6.14 (s, 1 H), 3.68 - 3.82 (m, 2 H), 3.22 - 3.32 (m, 2 H), 3.13 - 3.22 (m, 1 H), 3.03 (s, 4 H), 2.72 (s, 2 H), 2.09 - 2.23 (m, 6 H), 50 1.78 - 1.98 (m, 4 H), 1.56 - 1.66 (m, 1 H), 1.43 - 1.51 (m, 1 H), 1.37 (t, 2 H), 1.22 - 1.33 (m, 1 H), 0.92 (s, 6 H).

55

EXAMPLE 22

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-indol-5-yloxy)-N-[(3-nitro-4-[[[(3R)-tetrahydro-2H-pyran-3-ylmethyl]amino]phenyl)sulfonyl]benzamide

5

[0411] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 11C in place of EXAMPLE 11B, and EXAMPLE 18C in place of EXAMPLE 3J. ¹H NMR (400 MHz, dimethylsulfoxide -d₆) δ 11.17 (s, 2 H), 8.53 - 8.65 (m, 2 H), 7.80 (d, 1 H), 7.51 (d, 1 H), 7.38 - 7.44 (m, 2 H), 7.33 (d, 2 H), 7.15 (s, 1 H), 7.02 - 7.09 (m, 3 H), 6.82 - 6.92 (m, 1 H), 6.65 (d, 1 H), 6.39 (s, 1 H), 6.14 (s, 1 H), 3.68 - 3.82 (m, 2 H), 3.22 - 3.32 (m, 2 H), 3.13 - 3.22 (m, 1 H), 3.03 (s, 4 H), 2.72 (s, 2 H), 2.09 - 2.23 (m, 6 H), 1.78 - 1.98 (m, 4 H), 1.56 - 1.66 (m, 1 H), 1.43 - 1.51 (m, 1 H), 1.37 (t, 2 H), 1.22 - 1.33 (m, 1 H), 0.92 (s, 6 H).

10

EXAMPLE 23

15 4-(4-{[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl}piperazin-1-yl)-2-(1H-indol-5-yloxy)-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]benzamide

EXAMPLE 23A

20 methyl 2-(1H-indol-5-yloxy)-4-(piperazin-1-yl)benzoate

[0412] The title compound was prepared as described in EXAMPLE 15F by replacing EXAMPLE 3H with EXAMPLE 18A.

25 EXAMPLE 23B

methyl 2-(1H-indol-5-yloxy)-4-(4-((4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl)methyl)piperazin-1-yl)benzoate

30 **[0413]** The title compound was prepared as described in EXAMPLE 15G by replacing EXAMPLE 15F with EXAMPLE 23A.

EXAMPLE 23C

35 2-(1H-indol-5-yloxy)-4-(4-((4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl)methyl)piperazin-1-yl)benzoic acid

[0414] The title compound was prepared as described in EXAMPLE 15H by replacing EXAMPLE 15G with EXAMPLE 23B.

40

EXAMPLE 23D

4-(4-{[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl}piperazin-1-yl)-2-(1H-indol-5-yloxy)-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]benzamide

45

[0415] The title compound was prepared as described in EXAMPLE 11D by replacing EXAMPLE 11B with EXAMPLE 1F, and EXAMPLE 3J with EXAMPLE 23C. ¹H NMR (400 MHz, dimethylsulfoxide -d₆) δ 11.20 (br s, 1H), 11.17(s, 1H), 8.63 (t, 1 H), 8.59 (d, 1 H), 7.79 (dd, 1 H), 7.51 (d, 1 H), 7.36 (m, 3 H), 7.13 (m, 2 H), 6.86 (dd, 1 H), 6.66 (dd, 1 H), 6.39 (s, 1 H), 6.15 (d, 1 H), 4.10 (s, 2 H), 3.85 (m, 3 H), 3.50 (m, 2 H), 3.42 (m, 2 H), 3.24 (m, 4 H), 3.02 (m, 4 H), 2.82 (m, 2 H), 2.16 (m, 2 H), 1.61 (m, 3 H), 1.25 (m, 4 H), 1.17 (s, 6 H).

50

55

EXAMPLE 24

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[3-nitro-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 24A

[0416] 3-nitro-4-((tetrahydro-2H-pyran-4-yl)methoxy)benzenesulfonamide

[0417] (Tetrahydro-2H-pyran-4-yl)methanol (2.0 g) in tetrahydrofuran (20 mL) was treated with 60% NaH (1.377 g). The solution was stirred for 20 minutes at the room temperature. To this solution was added 4-fluoro-3-nitrobenzenesulfonamide (2.84 g) portion-wise. The reaction was stirred for another 2 hours. The mixture was poured into water, neutralized with 10% HCl, and extracted with ethyl acetate three times. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified with flash column chromatography on silica gel eluting with 20-60% ethyl acetate in hexanes.

EXAMPLE 24B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[3-nitro-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0418] The title compound was prepared by substituting EXAMPLE 24A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 8.33 (s, 1H), 8.00-8.02 (m, 2H), 7.50-7.53 (m, 3H), 7.34-7.36 (m, 3H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.38 (d, 1H), 6.21 (s, 1H), 4.06 (d, 2H), 3.88 (dd, 2H), 3.08 (s, 4H), 2.80 (s, 2H), 2.25 (s, 4H), 2.15 (s, 2H), 1.96 (s, 2H), 1.63-1.66 (m, 2H), 1.52-1.55 (m, 1 H), 1.33-1.40 (m, 4H), 0.92 (s, 6H).

EXAMPLE 25

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1,4-dioxan-2-ylmethyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 25A

4-((1,4-dioxan-2-yl)methylamino)-3-nitrobenzenesulfonamide

[0419] The title compound was prepared as described in EXAMPLE 1F using (1,4-dioxan-2-yl)methanamine in place of (tetrahydropyran-4-yl)methanamine.

EXAMPLE 25B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1,4-dioxan-2-ylmethyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0420] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 25A in place of EXAMPLE 11B. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1H), 11.38 (s, 1H), 8.53 - 8.59 (m, 2H), 8.03 (d, 1H), 7.81 (dd, 1H), 7.46 - 7.54 (m, 3H), 7.34 (d, 2H), 7.09 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.38 (dd, 1H), 6.19 (d, 1H), 3.75 - 3.86 (m, 3H), 3.58 - 3.68 (m, 2H), 3.45 - 3.52 (m, 2H), 3.35 - 3.43 (m, 2H), 3.07 (s, 4H), 2.75 (s, 2H), 2.17 (d, 6H), 1.95 (s, 2H), 1.38 (t, 2H), 0.92 (s, 6H)

EXAMPLE 26

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(2,2,2-trifluoroethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 26A

3-nitro-4-(2,2,2-trifluoroethylamino)benzenesulfonamide

[0421] The title compound was prepared by substituting 2,2,2-trifluoroethanamine for (tetrahydropyran-4-yl)methyl-

amine in EXAMPLE 1F.

EXAMPLE 26B

5 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(2,2,2-trifluoroethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 **[0422]** The title compound was prepared by substituting EXAMPLE 26A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.48 (s, 1H), 8.40 (m, 2H), 7.90 (d, 1H), 7.71 (dd, 1H), 7.59 (d, 1H), 7.40 (t, 1H), 7.34 (d, 2H), 7.25 (d, 1H), 7.06 (m, 3H), 6.61 (dd, 1H), 6.26 (m, 2H), 4.32 (m, 2H), 3.00 (m, 4H), 2.73 (s, 2H), 2.19 (m, 6H), 1.96 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 27

15 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(3,3,3-trifluoropropyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 27A

20 3-nitro-4-(3,3,3-trifluoropropylamino)benzenesulfonamide

[0423] The title compound was prepared by substituting 3,3,3-trifluoropropan-1-amine for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 27B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(3,3,3-trifluoropropyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

30 **[0424]** The title compound was prepared by substituting EXAMPLE 27A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.47 (s, 1H), 8.37 (d, 1H), 8.29 (m, 1H), 7.89 (d, 1H), 7.61 (m, 2H), 7.39 (t, 1H), 7.35 (d, 2H), 7.22 (d, 1H), 7.05 (d, 2H), 6.75 (d, 1H), 6.62 (dd, 1H), 6.27 (m, 2H), 3.59 (q, 2H), 3.00 (m, 4H), 2.73 (s, 2H), 2.66 (m, 2H), 2.18 (m, 6H), 1.96 (s, 2H), 1.39 (t, 2H), 0.93 (m, 6H).

EXAMPLE 28

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(2S)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 28A

(S)-4-((1,4-dioxan-2-yl)methoxy)-3-nitrobenzenesulfonamide

[0425] The racemic mixture of EXAMPLE 12A was resolved on a SFC chiral AD column to provide the title compound.

45

EXAMPLE 28B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(2S)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50

[0426] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 28A in place of EXAMPLE 11B. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 2H), 8.35 (s, 1H), 8.03 (d, 2H), 7.48 - 7.57 (m, 3H), 7.42 (d, 1H), 7.35 (d, 2H), 7.05 (d, 2H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.21 (s, 1H), 4.19 - 4.30 (m, 2H), 3.85 - 3.92 (m, 1H), 3.73 - 3.85 (m, 2H), 3.58 - 3.70 (m, 2H), 3.40 - 3.52 (m, 2H), 3.10 (s, 4H), 2.85 (s, 2H), 2.18 - 2.39 (m, 3H), 2.15 (s, 2H), 1.96 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

55

EXAMPLE 29

Cis-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 29A

Cis-4-([4-methoxycyclohexyl)methylamino)-3-nitrobenzenesulfonamide

[0427] 4-Fluoro-3-nitrobenzenesulfonamide (1.098 g) and EXAMPLE 34A (1 g) in tetrahydrofuran (20 mL) was treated with N,N-diisopropylethylamine (0.871 mL) overnight. The reaction mixture was concentrated and the residue was purified by reverse phase chromatography, eluted with 40-55% acetonitrile in 0.1% trifluoroacetic acid in water over 25 min to give the *cis* isomer EXAMPLE 29A and *trans* isomer EXAMPLE 34B.

EXAMPLE 29B

Cis-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0428] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 29A in place of EXAMPLE 11B. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 11.36 (s, 1H), 8.53 - 8.63 (m, 2H), 8.04 (d, 1H), 7.79 (dd, 1H), 7.47 - 7.56 (m, 3H), 7.34 (d, 2H), 7.00 - 7.12 (m, 3H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.19 (d, 1H), 3.37 (s, 1H), 3.26 (t, 2H), 3.20 (s, 3H), 3.07 (s, 4H), 2.75 (s, 2H), 2.17 (d, 6H), 1.95 (s, 2H), 1.81 (dd, 2H), 1.64-1.74 (m, 1H), 1.48 (dd, 2H), 1.23 - 1.42 (m, 6H), 0.92 (s, 6H).

EXAMPLE 30

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-[(2R)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 30A

(R)-4-((1,4-dioxan-2-yl)methoxy)-3-nitrobenzenesulfonamide

[0429] The racemic mixture of EXAMPLE 12A was resolved on a SFC chiral AD column to provide the title compound.

EXAMPLE 30B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-[(2R)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0430] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 30A in place of EXAMPLE 11B. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 2H), 8.35 (s, 1H), 8.03 (d, 2H), 7.48 - 7.57 (m, 3H), 7.42 (d, 1H), 7.35 (d, 2H), 7.05 (d, 2H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.21 (s, 1H), 4.19 - 4.30 (m, 2H), 3.85 - 3.92 (m, 1H), 3.73 - 3.85 (m, 2H), 3.58 - 3.70 (m, 2H), 3.40 - 3.52 (m, 2H), 3.10 (s, 4H), 2.85 (s, 2H), 2.18 - 2.39 (m, 3H), 2.15 (s, 2H), 1.96 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 31

4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-[(1,4-dioxan-2-yl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0431] The title compound was prepared as described in EXAMPLE 1G by replacing EXAMPLE 1E and EXAMPLE 1F with EXAMPLE 15H and EXAMPLE 25A, respectively. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1H), 11.46 (m, 1H), 8.54 (m, 2H), 8.45 (m, 1H), 8.03 (d, 1H), 7.83 (m, 2H), 7.50 (m, 3H), 7.34 (m, 3H), 7.12 (m, 2H), 6.68 (dd, 1H), 6.38 (dd, 1H), 6.20 (d, 1H), 4.11 (s, 2H), 3.79 (m, 4H), 3.51 (m, 6H), 3.05 (m, 4H), 2.17 (m, 3H), 1.17 (s, 6H).

EXAMPLE 32

4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0432] The title compound was prepared as described in EXAMPLE 1G by replacing EXAMPLE 1E and EXAMPLE 1F with EXAMPLE 15H and EXAMPLE 12A, respectively. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 8.37 (d, 1H), 8.03 (m, 2H), 7.50 (m, 3H), 7.37 (d, 2H), 7.13 (d, 2H), 6.68 (dd, 1H), 6.40 (dd, 1H), 6.20 (d, 1H), 4.25 (m, 2H), 4.12 (s, 2H), 3.84 (m, 3H), 3.63 (m, 2H), 3.45 (m, 2H), 3.06 (m, 4H), 2.86 (m, 2H), 2.24 (m, 6H), 1.20 (m, 6H).

EXAMPLE 33

Trans-4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-([4-morpholin-4-ylcyclohexyl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0433] The title compound was prepared as described in EXAMPLE 1G by replacing EXAMPLE 1E and EXAMPLE 1F with EXAMPLE 15H and EXAMPLE 9C, respectively. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.63 (s, 1H), 8.51 (d, 1H), 8.15 (d, 1H), 8.01 (d, 1H), 7.76 (dd, 1H), 7.48 (m, 3H), 7.38 (d, 2H), 7.13 (d, 2H), 7.06 (d, 1H), 6.66 (dd, 1H), 6.36 (dd, 1H), 6.21 (d, 1H), 4.11 (s, 2H), 3.63 (m, 5H), 3.05 (m, 4H), 2.83 (s, 2H), 2.64 (m, 4H), 2.17 (m, 6H), 2.05 (m, 2H), 1.91 (s, 2H), 1.43 (m, 6H), 1.17 (m, 6H).

EXAMPLE 34

Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methoxycyclohexyl]methyl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 34A

(4-methoxycyclohexyl)methanamine

[0434] (4-Methoxyphenyl)methanamine (1 g, 1.29 mmol) in ethanol (10 ml) was treated with 5% Rh-Al₂O₃ (99.8 mg, 0.048 mmol) under H₂ atmosphere (500 psi) at 50°C for 16 hours. Additional 5% Rh-Al₂O₃ (0.4 g) was added. The resulting mixture was stirred under H₂ atmosphere (500 psi) at 60°C for 2 hours. The insoluble material was filtered off and the filtrate was concentrated to provide a mixture of *cis* and *trans* product as an oil, which was used in the next step without further purification.

EXAMPLE 34B

Trans-4-((4-methoxycyclohexyl)methylamino)-3-nitrobenzenesulfonamide

[0435] 4-Fluoro-3-nitrobenzenesulfonamide (1.098 g) and EXAMPLE 34A (1 g) in tetrahydrofuran (20 mL) was treated with N,N-diisopropylethylamine (0.871 mL) overnight. The reaction mixture was concentrated and the residue was purified by reverse phase chromatography, and was eluted with 40-55% acetonitrile in 0.1% trifluoroacetic acid in water over 25 minutes.

EXAMPLE 34C

Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-methoxycyclohexyl]methyl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0436] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 34B in place of EXAMPLE 11B. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1 H), 11.37 (s, 1 H), 8.52 - 8.62 (m, 2 H), 8.04 (d, 1 H), 7.79 (dd, 1 H), 7.47 - 7.55 (m, 3 H), 7.34 (d, 2 H), 7.02 - 7.09 (m, 3 H), 6.68 (dd, 1 H), 6.39 (dd, 1 H), 6.19 (d, 1 H), 3.21 - 3.27 (m, 5 H), 3.02 - 3.12 (m, 5 H), 2.75 (s, 2 H), 2.20 (s, 4 H), 2.14 (s, 2 H), 1.93 - 2.04 (m, 4 H), 1.79 (d, 2 H), 1.55 - 1.65 (m, 1 H), 1.38 (t, 2 H), 0.97 - 1.12 (m, 4 H), 0.92 (s, 6 H).

EXAMPLE 35

4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0437] The title compound was prepared as described in EXAMPLE 1G by replacing EXAMPLE 1E and EXAMPLE 1F with EXAMPLE 15H and EXAMPLE 36C, respectively. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1H), 8.78 (s, 1H), 8.58 (s, 1H), 8.00 (d, 1H), 7.51 (m, 3H), 7.38 (d, 2H), 7.14 (d, 2H), 6.68 (dd, 1H), 6.37 (dd, 1H), 6.23 (d, 1H), 4.31 (d, 2H), 4.13 (s, 2H), 3.88 (dd, 2H), 3.11 (m, 5H), 2.16 (m, 6H), 1.65 (m, 2H), 1.35 (m, 2H), 1.19 (s, 6H).

EXAMPLE 36

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 36A

[0438] 5-Bromo-6-chloropyridine-3-sulfonyl chloride (8.2 g) in methanol (20 mL) was cooled to 0 °C. To this solution was added 7N NH₃ in methanol (80 mL). The reaction mixture was stirred overnight. The solvent was removed at low temperature, and the residue was partitioned between ethyl acetate and water. The aqueous layer was extracted with ethyl acetate three times. The combined organic layers were washed with brine, dried (MgSO₄), filtered, and concentrated. The solid was purified by flash column chromatography on silica gel using 20-100% ethyl acetate in hexanes to give the title compound.

EXAMPLE 36B

[0439] The title compound was prepared by substituting EXAMPLE 36A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 36C

[0440] A mixture of EXAMPLE 36B (0.702 g), dicyanozinc (0.129 g), and tetrakis(triphenylphosphine)palladium(0) (0.231 g) in N,N-dimethylformamide (2 mL) was degassed via vacuum/nitrogen cycle three times. The reaction mixture was heated at 120 °C for 3 hours. After cooling, it was poured into water and extracted with ethyl acetate three times. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified with flash column chromatography on silica gel eluting with 20%-60% ethyl acetate in hexanes to give the title compound.

EXAMPLE 36D

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0441] The title compound was prepared by substituting EXAMPLE 36C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.56 (s, 1H), 8.66 (s, 1H), 8.44 (s, 1H), 7.94 (d, 1H), 7.55 (d, 1H), 7.44 (t, 1H), 7.34-7.35 (m, 3H), 7.04 (d, 2H), 6.65 (dd, 1H), 6.32 (s, 1H), 6.24 (s, 1H), 4.26 (d, 2H), 3.86 (dd, 2H), 3.10 (s, 4H), 2.75 (s, 2H), 2.31-2.35 (m, 2H), 2.01-2.05 (m, 1H), 2.15 (s, 2H), 1.96 (s, 2H), 1.63-1.66 (m, 2H), 1.33-1.40 (m, 4H), 0.92 (s, 6H).

EXAMPLE 37

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 37A

1,6-dioxaspiro[2.5]octane-2-carbonitrile

[0442] A mixture of tetrahydropyran-4-one (10 mL) and chloroacetonitrile (6.4 mL) in tert-butanol (10 mL) was stirred

for 10 minutes. To this solution was added a solution of potassium tert-butoxide (12.11 g) in 200 mL of tert-butanol at room temperature over 40 minutes. The reaction mixture was stirred for 16 hours, diluted with water and quenched slowly with 1 N HCl. The solvent was partially removed by rotary evaporation. It was then extracted with ether (5x 200 mL). The combined extracts was washed with brine, dried over MgSO₄, filtered, and the filtrate was concentrated and purified by flash chromatography on silica with 3:7 to 1:1 ethyl acetate:hexanes to provide the title compound.

EXAMPLE 37B

2-(4-fluorotetrahydro-2H-pyran-4-yl)-2-hydroxyacetonitrile

[0443] EXAMPLE 37A (11.5 g) in dichloromethane (40 mL) in a polypropylene bottle was treated with 70% hydrogen fluoride-pyridine (10.4 mL) dropwise at 0 °C. The solution was allowed to warm to room temperature over 3 hours, and stirred for an additional 1.5 hours. The reaction mixture was diluted with ethyl acetate (200 mL) and poured into saturated aqueous NaHCO₃. Additional solid NaHCO₃ was used carefully until bubbling ceased. The organic layer was isolated, and the aqueous layer was extracted with additional ethyl acetate three times (150 mL each). The combined organic layers were washed with 5% HCl (50 mL each, twice), brine, dried over MgSO₄, filtered and concentrated to give the desired product which was used directly in the next step.

EXAMPLE 37C

(4-fluorotetrahydro-2H-pyran-4-yl)methanol

[0444] EXAMPLE 37B (11.7 g, 74 mmol) in 2-propanol (150 mL) and water (37.5 mL) was cooled to 0°C. To this solution was added NaBH₄ (4.20 g, 111 mmol). The solution was stirred and allowed to warm to room temperature over 3 hours. It was quenched with acetone, and stirred for another 1 hour. The clear liquid was separated from solid by decanting. Additional ethyl acetate (2x 100 mL) was used to wash the solid, and the mixture was decanted. The combined organic solutions were concentrated. The residue was purified by flash chromatography, eluting with 1:1 ethyl acetate:hexanes to provide the title compound.

EXAMPLE 37D

4-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)-3-nitrobenzenesulfonamide

[0445] The title compound was prepared by substituting EXAMPLE 37C for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 37E

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0446] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 37D in place of EXAMPLE 11B. ¹H NMR (dimethylsulfoxide-d₆). 11.64 (s, 2H), 8.33 (s, 1H), 8.00-8.01 (m, 2H), 7.39-7.57 (m, 4H), 7.33 (d, J = 8.24 Hz, 2H), 7.03 (d, J = 8.54 Hz, 2H), 6.65 (dd, J = 9, 1.98 Hz, 1H), 6.37-6.38 (m, 1H), 6.19 (d, J = 1.53 Hz, 1H), 4.35 (d, J = 20.75 Hz, 2H), 3.74-3.78 (m, 2H), 3.55-3.60 (m, 2H), 3.07 (br, 4H), 2.80 (br, 2H), 2.25 (br, 4H), 2.13 (br, 2H), 1.81-1.94 (m, 6H), 1.38 (t, J = 6.26 Hz, 2H), 0.91 (s, 6H).

EXAMPLE 38

N-([3-(aminocarbonyl)-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 38A

3-cyano-4-((tetrahydro-2H-pyran-4-yl)methoxy)benzenesulfonamide

[0447] The title compound was prepared by substituting 3-cyano-4-fluorobenzenesulfonamide for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 38B

5-sulfamoyl-2-((tetrahydro-2H-pyran-4-yl)methoxy)benzamide

5 **[0448]** To a solution of EXAMPLE 38A (0.455 g) in ethanol (3 mL) and tetrahydrofuran (1 mL) was added hydrogen peroxide (30% in water, 2 mL) followed by 1 N aqueous NaOH (1.024 ml) and heated to 35°C for 3 hours. The reaction was poured into dichloromethane (50 mL) and 1N aqueous HCl (25 mL). The aqueous layer was extracted with dichloromethane (3 x 50 mL). The precipitate contained in the combined organic layers was collected by filtration to give the title compound.

10

EXAMPLE 38C

N-([3-(aminocarbonyl)-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

15

[0449] The title compound was prepared by substituting EXAMPLE 38B for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.79 - 11.70 (m, 1H), 11.66 - 11.54 (m, 1H), 9.29 - 9.08 (m, 1H), 8.27 (d, 1H), 8.08 (d, 1H), 7.97 - 7.90 (m, 1H), 7.76 - 7.72 (m, 1H), 7.62 (s, 1H), 7.54 (s, 1H), 7.50 (d, 1H), 7.39 (d, 1H), 7.23 (d, 1H), 7.08 (d, 1H), 6.74 - 6.67 (m, 1H), 6.44 (s, 1H), 6.22 (s, 1H), 4.03 (d, 6H), 3.74 - 3.52 (m, 4H), 3.33 (s, 4H), 3.11 - 2.90 (m, 2H), 2.01 (s, 4H), 1.79 - 1.58 (m, 2H), 1.24 (s, 5H), 0.94 (s, 6H).

20

EXAMPLE 39

Cis-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

25

EXAMPLE 39A

Cis-tert-butyl-4-morpholinocyclohexylcarbamate

30

[0450] To a solution of morpholine (4.08 g) and tert-butyl 4-oxocyclohexylcarbamate (10 g) stirred for 24 hours at room temperature in titanium (IV) isopropoxide (27.5 mL), methanol (10 mL) was added followed by careful addition of sodium borohydride (3.55 g). The reaction mixture was quenched with water/NaOH solution, extracted with ether, dried over magnesium sulfate, filtered, and concentrated. The product was separated from the trans isomer and purified by flash chromatography (silica gel, 50% - 100% acetone in hexanes) to provide the title compound.

35

EXAMPLE 39B

cis-4-morpholinocyclohexanamine bis(2,2,2-trifluoroacetate)

40

[0451] To a solution of EXAMPLE 39A (2.43 g) in dichloromethane (15 ml) was added trifluoroacetic acid (5 ml) and the reaction mixture was stirred for 16 hours at room temperature. The reaction mixture was concentrated and the crude product was used without purification.

45

EXAMPLE 39C

4-(cis-4-morpholinocyclohexylamino)-3-nitrobenzenesulfonamide

50

[0452] A solution of EXAMPLE 39B (0.40 g), 4-fluoro-3-nitrobenzenesulfonamide (0.478 g) and triethylamine (2 mL) in tetrahydrofuran (10 mL) was stirred for 3 days at room temperature. The reaction mixture was concentrated and purified by flash chromatography (silica gel, 0-30% methanol / dichloromethane) providing the product.

EXAMPLE 39D

55

Cis-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0453] The title compound was prepared by substituting EXAMPLE 39C for EXAMPLE 11B in EXAMPLE 11D. ¹H

NMR (500 MHz, pyridine-d₅) δ 13.07 (s, 1H), 9.30 (d, 1H), 8.64 (d, 1H), 8.43 (d, 1H), 8.38 (dd, 1H), 8.11 (d, 1H), 7.67 (t, 2H), 7.44 (d, 2H), 7.06 (d, 2H), 6.91 (d, 1H), 6.74 (dd, 1H), 6.48 - 6.55 (m, 2H), 3.65 - 3.73 (m, 5H), 3.02 - 3.09 (m, 4H), 2.76 (s, 2H), 2.41 - 2.48 (m, 4H), 2.25 (t, 2H), 2.09 - 2.16 (m, 5H), 1.97 (s, 2H), 1.77 - 1.86 (m, 2H), 1.55 - 1.63 (m, 6H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 40

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 40A

5,6-dichloropyridine-3-sulfonamide

[0454] The title compound was prepared by substituting 5,6-dichloropyridine-3-sulfonyl chloride for 5-bromo-6-chloropyridine-3-sulfonyl chloride in EXAMPLE 36A.

EXAMPLE 40B

5-chloro-6-((tetrahydro-2H-pyran-4-yl)methoxy)pyridine-3-sulfonamide

[0455] The title compound was prepared by substituting EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 40C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0456] The title compound was prepared by substituting EXAMPLE 40B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 8.52 (s, 1H), 8.39 (d, 1H), 8.03 (d, 1H), 7.54 (d, 1H), 7.52 (d, 1H), 7.50 (dd, 1H), 7.35 (d, 2H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.39 (m, 1H), 6.21 (d, 1H), 4.25 (d, 2H), 3.87 (dd, 2H), 3.30 (m, 2H), 3.10 (v br s, 4H), 2.90 (v br s, 2H), 2.35 (v br s, 4H), 2.17 (br m, 2H), 2.05 (m, 1H), 1.96 (s, 2H), 1.64 (d, 2H), 1.40 (t, 2H), 1.35 (ddd, 2H), 0.93 (s, 6H).

EXAMPLE 41

4-(4-{[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl}piperazin-1-yl)-N-{[5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0457] The title compound was prepared by substituting EXAMPLE 15H for EXAMPLE 3J and EXAMPLE 40B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 8.55 (d, 1H), 8.41 (d, 1H), 8.04 (d, 1H), 7.54 (m, 2H), 7.50 (dd, 1H), 7.38 (d, 2H), 7.14 (d, 2H), 6.68 (dd, 1H), 6.40 (m, 1H), 6.20 (d, 1H), 4.25 (d, 2H), 4.12 (s, 2H), 3.87 (dd, 2H), 3.30 (m, 2H), 3.10 (v br s, 4H), 2.90 (v br s, 2H), 2.27 (v br s, 4H), 2.17 (br m, 2H), 2.05 (m, 1H), 1.96 (s, 2H), 1.64 (d, 2H), 1.35 (ddd, 2H), 0.97 (s, 6H).

EXAMPLE 42

4-(4-{[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-{[4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-(trifluoromethyl)phenyl]sulfonyl}benzamide

EXAMPLE 42A

4-((tetrahydro-2H-pyran-4-yl)methylamino)-3-(trifluoromethyl)benzenesulfonamide

[0458] A mixture of 4-fluoro-3-(trifluoromethyl)benzenesulfonamide (1.056 g), (tetrahydro-2H-pyran-4-yl)methanamine (0.5 g) and N,N-diisopropylethylamine (1.68 g) in anhydrous dimethylsulfoxide (15 mL) solution was heated at 90°C overnight. The reaction mixture was cooled to room temperature and diluted with ethyl acetate. The organic phase was

washed with water, brine, dried over anhydrous sodium sulfate, filtered and concentrated to afford the title compound.

EXAMPLE 42B

5 4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([4-([tetrahydro-2H-pyran-4-ylmethyl]amino)-3-(trifluoromethyl)phenyl]sulfonyl)benzamide

[0459] The title compound was prepared as described in EXAMPLE 1G by replacing EXAMPLE 1E and EXAMPLE 1F with EXAMPLE 15H and EXAMPLE 42A, respectively. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.73 (s, 1H), 11.25 (s, 1H), 8.08 (d, 1H), 7.89 (d, 1H), 7.77 (m, 1H), 7.61 (d, 1H), 7.51 (m, 2H), 7.37 (d, 2H), 7.13 (d, 2H), 6.88 (d, 1H), 6.67 (dd, 1H), 6.53 (m, 1H), 6.43 (m, 1H), 6.15 (d, 1H), 4.11 (s, 2H), 3.82 (dd, 2H), 3.19 (m, 5H), 3.05 (m, 4H), 2.82 (s, 2H), 2.20 (m, 7H), 1.85 (m, 1H), 1.56 (m, 2H), 1.18 (s, 6H).

EXAMPLE 43

15 4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([4-([tetrahydro-2H-pyran-4-ylmethyl]amino)-3-(trifluoromethyl)sulfonyl]phenyl)sulfonyl)benzamide

[0460] The title compound was prepared as described in EXAMPLE 1G by replacing EXAMPLE 1E and EXAMPLE 1F with EXAMPLE 15H and EXAMPLE 17A, respectively. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 11.48 (m, 1H), 8.16 (d, 1H), 8.05 (d, 1H), 7.92 (dd, 1H), 7.52 (m, 3H), 7.37 (d, 2H), 7.27 (m, 1H), 7.11 (m, 3H), 6.68 (dd, 1H), 6.41 (dd, 1H), 6.18 (d, 1H), 4.11 (s, 2H), 3.84 (dd, 2H), 3.25 (m, 4H), 3.07 (m, 4H), 2.84 (m, 2H), 2.23 (m, 5H), 1.84 (m, 1H), 1.55 (m, 2H), 1.25 (m, 3H), 1.18 (s, 6H).

EXAMPLE 44

Trans-4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-([4-morpholin-4-ylcyclohexyl]amino)-3-([trifluoromethyl]sulfonyl)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 44A

Trans-4-(4-morpholinocyclohexylamino)-3-(trifluoromethylsulfonyl)benzenesulfonamide

[0461] The title compound was prepared as described in EXAMPLE 16A by replacing 2-methoxyethanamine with EXAMPLE 9B.

EXAMPLE 44B

40 Trans-4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-([4-morpholin-4-ylcyclohexyl]amino)-3-([trifluoromethyl]sulfonyl)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0462] The title compound was prepared as described in EXAMPLE 1G by replacing EXAMPLE 1E and EXAMPLE 1F with EXAMPLE 15H and EXAMPLE 44A, respectively. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.62 (s, 1H), 8.08 (s, 1H), 8.00 (d, 1H), 7.85 (d, 1H), 7.47 (m, 3H), 7.38 (d, 2H), 7.14 (d, 2H), 6.98 (d, 1H), 6.65 (dd, 1H), 6.55 (m, 1H), 6.37 (m, 1H), 6.21 (d, 1H), 4.12 (s, 2H), 3.54 (m, 6H), 3.04 (m, 4H), 2.83 (s, 2H), 2.57 (m, 3H), 2.24 (m, 6H), 1.91 (m, 5H), 1.34 (m, 4H), 1.20 (s, 6H).

EXAMPLE 45

50 4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-([1-methylpiperidin-4-yl]amino)-3-([trifluoromethyl]sulfonyl)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 45A

55 4-(1-methylpiperidin-4-ylamino)-3-(trifluoromethylsulfonyl)benzenesulfonamide

[0463] The title compound was prepared as described in EXAMPLE 16A by replacing 2-methoxyethanamine with 1-methyl-4-aminopiperidine.

EXAMPLE 45B

4-(4-{[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl}piperazin-1-yl)-N-({4-[(1-methylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0464] The title compound was prepared as described in EXAMPLE 1G by replacing EXAMPLE 1E and EXAMPLE 1F with EXAMPLE 15H and EXAMPLE 45A, respectively. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.59 (s, 1H), 8.10 (s, 1H), 7.98 (d, 1H), 7.90 (dd, 1H), 7.49 (m, 3H), 7.39 (m, 3H), 7.14 (d, 2H), 7.02 (d, 1H), 6.65 (dd, 2H), 6.36 (dd, 1H), 6.22 (d, 1H), 4.12 (s, 2H), 3.75 (m, 1H), 3.16 (m, 4H), 2.98 (m, 5H), 2.88 (m, 5H), 2.67 (s, 2H), 2.22 (m, 6H), 1.68 (m, 1H), 1.18 (s, 6H).

EXAMPLE 46

5-({[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino}sulfonyl)-2-(tetrahydro-2H-pyran-4-ylmethoxy)nicotinamide

5-sulfamoyl-2-((tetrahydro-2H-pyran-4-yl)methoxy)nicotinamide

[0465] To EXAMPLE 36C (0.025 g) in ethanol (1 mL) and tetrahydrofuran (1 mL) was added hydrogen peroxide (30% in water, 0.5 mL) followed by 1M aqueous sodium hydroxide (0.056 mL) then another 1 mL of tetrahydrofuran. The reaction was heated to 45°C for 2 hours, cooled, quenched with 1N aqueous HCl (5 mL), and the product extracted into dichloromethane (10 mL). The organic layer was dried over magnesium sulfate, filtered and concentrated to give the title compound.

EXAMPLE 46B

5-({[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino}sulfonyl)-2-(tetrahydro-2H-pyran-4-ylmethoxy)nicotinamide

[0466] The title compound was prepared by substituting EXAMPLE 46A for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, CDCl₃) δ 10.31 - 10.09 (m, 1H), 9.09 (s, 2H), 8.93 - 8.81 (m, 1H), 8.28 - 8.18 (m, 1H), 8.03 - 7.87 (m, 1H), 7.77 - 7.68 (m, 1H), 7.59 - 7.51 (m, 1H), 7.48 - 7.41 (m, 1H), 6.91 (d, 2H), 6.59-6.48 (m, 2H), 5.97 (s, 2H), 4.50 (d, 2H), 4.08 - 3.98 (m, 2H), 3.45 (s, 4H), 3.13 - 2.99 (m, 4H), 2.82 - 2.68 (m, 2H), 2.19 (s, 4H), 1.86 (s, 5H), 1.61 - 1.35 (m, 4H), 0.94 (s, 6H).

EXAMPLE 47

N-({5-bromo-6-[(1-methylpiperidin-4-yl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 47A

5-bromo-6-((1-methylpiperidin-4-yl)methoxy)pyridine-3-sulfonamide

[0467] To (1-methylpiperidin-4-yl)methanol (0.109 g) in tetrahydrofuran (2 mL) was added sodium hydride (0.136 g). After 30 minutes, EXAMPLE 36A (0.230 g) was added as a solution in tetrahydrofuran (1 mL) and the reaction was heated to 50°C. After 4 hours, the reaction was cooled, poured into water (10 mL) and dichloromethane (50 mL), and the pH was adjusted to pH~8. The aqueous layer was extracted with dichloromethane (3 x 50 mL), and the organic layers were combined, washed with brine (30 mL), dried over magnesium sulfate, filtered, and concentrated to give the title compound.

EXAMPLE 47B

N-({5-bromo-6-[(1-methylpiperidin-4-yl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0468] The title compound was prepared by substituting EXAMPLE 47A for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.51 (s, 1H), 8.35 (d, 1H), 8.17 (d, 1H),

7.93 (d, 1H), 7.60 (d, 1H), 7.44 - 7.40 (m, 1H), 7.33 (dd, 3H), 7.05 (d, 2H), 6.61 (d, 1H), 6.31 (dd, 1H), 6.24 (s, 1H), 4.25 (d, 2H), 3.40 (s, 4H), 3.01 (s, 4H), 2.73 (d, J = 8.2, 5H), 2.20 (s, 6H), 1.93 (d, 4H), 1.54 (s, 1H), 1.39 (s, 2H), 1.24 (s, 2H), 0.93 (s, 6H).

5 EXAMPLE 48

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-methylpiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 EXAMPLE 48A

4-([1-methylpiperidin-4-yl)methoxy]-3-nitrobenzenesulfonamide

15 **[0469]** The title compound was prepared by substituting (1-methylpiperidin-4-yl)methanol for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 48B

20 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-methylpiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

25 **[0470]** The title compound was prepared by substituting EXAMPLE 48A for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.54 (s, 1H), 8.17 (s, 1H), 7.92 (s, 1H), 7.87 - 7.77 (m, 1H), 7.58 (d, 1H), 7.43 (s, 1H), 7.40 - 7.00 (m, 7H), 6.70 - 6.56 (m, 1H), 6.31 (s, 1H), 6.24 (s, 1H), 4.05 (s, 2H), 3.46 - 3.33 (m, 2H), 3.02 (s, 6H), 2.72 (d, 5H), 2.21 (s, 6H), 1.96 (s, 5H), 1.70 - 1.48 (m, 2H), 1.39 (s, 2H), 0.93 (s, 6H).

EXAMPLE 49

30 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(1,4-dioxan-2-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 49A

35 6-([1, 4-dioxan-2-yl)methoxy)-5-bromopyridine-3-sulfonamide

[0471] The title compound was prepared by substituting (1, 4-dioxan-2-yl)methanol for (tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 36A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 49B

40 6-([1,4-dioxan-2-yl)methoxy)-5-cyanopyridine-3-sulfonamide

[0472] The title compound was prepared by substituting EXAMPLE 49A for EXAMPLE 36B in EXAMPLE 36C.

45 EXAMPLE 49C

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(1,4-dioxan-2-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50 **[0473]** The title compound was prepared by substituting EXAMPLE 49B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.50 (s, 1H), 8.60 (s, 1H), 8.40 (s, 1H), 7.91 (d, 1H), 7.58 (d, 1H), 7.42 (t, 1H), 7.35 (d, 2H), 7.28 (s, 1H), 7.06 (d, 2H), 6.64 (dd, 1H), 6.29 (m, 2H), 4.40 (d, 2H), 3.90 (m, 1H), 3.79 (m, 2H), 3.63 (m, 2H), 3.46 (m, 4H), 3.07 (s, 4H), 2.85 (m, 2H), 2.34 (m, 4H), 2.16 (m, 2H), 1.40 (t, 2H), 0.93 (s, 6H).

55

EXAMPLE 50

N-([5-bromo-6-(1,4-dioxan-2-ylmethoxy)pyridin-3-yl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0474] The title compound was prepared by substituting EXAMPLE 49A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.60 (s, 1H), 8.46 (s, 1H), 8.27 (s, 1H), 7.99 (d, 1H), 7.56 (d, 1H), 7.46 (m, 2H), 7.35 (d, 2H), 7.05 (d, 2H), 6.65 (dd, 1H), 6.36 (dd, 1H), 6.22 (d, 1H), 4.34 (m, 2H), 3.88 (m, 1H), 3.79 (m, 2H), 3.63 (m, 2H), 3.46 (m, 2H), 3.06 (s, 4H), 2.81 (s, 2H), 2.26 (m, 4H), 2.15 (s, 2H), 1.96 (s, 2H), 1.38 (m, 2H), 0.93 (s, 6H).

EXAMPLE 51

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([2,2-dimethyltetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 51 A

[0475] The title compound was prepared as described in EXAMPLE 12A by replacing (1,4-dioxan-2-yl)methanol with (2,2-dimethyltetrahydro-2H-pyran-4-yl)methanol.

EXAMPLE 51B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([2,2-dimethyltetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0476] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 51A in place of EXAMPLE 11B. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 2H), 8.35 (s, 2H), 8.03 (d, 4H), 7.47 - 7.58 (m, 6H), 7.31 - 7.42 (m, 6H), 7.04 (d, 4H), 6.68 (dd, 2H), 6.40 (s, 2H), 6.20 (d, 2H), 3.96 - 4.09 (m, 2H), 3.54 - 3.68 (m, 2H), 3.09 (s, 4H), 2.83 (s, 2H), 2.09 - 2.37 (m, 7H), 1.96 (s, 2H), 1.55 - 1.69 (m, 2H), 1.39 (t, 2H), 1.19 (m, 8H), 0.92 (s, 6H).

EXAMPLE 52

N-([3-chloro-5-cyano-4-((tetrahydro-2H-pyran-4-ylmethyl)amino)phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 52A

3-cyano-4-fluorobenzenesulfonamide

[0477] 3-Cyano-4-fluorobenzene-l-sulfonyl chloride (1.1 g) in 1,4-dioxane (10 mL) at 0 °C was treated dropwise with a 7 M ammonia solution in methanol (3.57 mL) and stirred for 30 minutes. A small amount of solid was removed by filtration and discarded. The filtrate was concentrated, diluted with ethyl acetate, washed with water and brine, dried (MgSO₄), filtered, concentrated and triturated with diethyl ether to give the product.

EXAMPLE 52B

3-cyano-4-((tetrahydro-2H-pyran-4-yl)methylamino)benzenesulfonamide

[0478] The title compound was prepared by substituting EXAMPLE 52A for 4-chloro-3-nitrobenzenesulfonamide and (tetrahydro-2H-pyran-4-yl)methanamine for 4-methylpiperazin-1-amine dihydrochloride in EXAMPLE 6A.

EXAMPLE 52C

3-chloro-5-cyano-4-((tetrahydro-2H-pyran-4-yl)methylamino)benzenesulfonamide

[0479] EXAMPLE 52B (0.148 g) in acetonitrile (5 mL) was treated with N-chlorosuccinimide (0.080 g), heated at 60 °C for 3 hours and filtered to remove a small amount of solid. The filtrate was concentrated and chromatographed on silica gel with 3-15% ethyl acetate in dichloromethane as eluent. The obtained solid was slurried in water, filtered, rinsed

with additional water and dried under vacuum to give the product.

EXAMPLE 52D

5 N-({3-chloro-5-cyano-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0480] The title compound was prepared by substituting EXAMPLE 52C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.70 (s, 1H), 11.41 (br s, 1H), 8.07 (d, 1H), 7.89 (s, 2H), 7.61 (m, 1H), 7.53 (m, 2H), 7.35 (d, 2H), 7.18 (m, 1H), 7.05 (d, 2H), 6.69 (m, 1H), 6.42 (dd, 1H), 6.18 (dd, 1H), 3.83 (m, 2H), 3.55 (t, 2H), 3.23 (m, 3H), 3.06 (m, 4H), 2.15 (m, 4H), 1.92 (m, 4H), 1.60 (m, 2H), 1.40 (m, 2H), 1.19 (m, 4H), 0.93 (s, 6H).

EXAMPLE 53

15 N-({4-[(1-acetyl)piperidin-4-yl]amino}-3-nitrophenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 53A

20 N-[(4-chloro-3-nitrophenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0481] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and 4-chloro-3-nitrobenzenesulfonamide for EXAMPLE 1F in EXAMPLE 1G.

EXAMPLE 53B

30 N-({4-[(1-acetyl)piperidin-4-yl]amino}-3-nitrophenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0482] A 5 mL round-bottomed flask was charged with EXAMPLE 53A (120 mg), 1-acetyl piperidin-4-amine (28 mg), and triethylamine (0.064 mL) in dioxane (2 mL). The reaction mixture was heated to 90°C for 24 hours. The reaction mixture was cooled to room temperature, and added to a silica gel column and purified by eluting with 0-5% methanol in dichloromethane. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.66 (br s, 1H), 8.65 (d, 1H), 8.24 (d, 1H), 8.03 (d, 1H), 7.83 (dd, 1H), 7.54-7.46 (m, 3H), 7.35 (d, 2H), 7.19 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.39 (m, 1H), 6.20 (d, 1H), 4.28 (d, 1H), 3.97-3.75 (m, 2H), 3.07 (br s, 4H), 2.87-2.70 (m, 4H), 2.29-2.10 (m, 6H), 2.02 (s, 3H), 2.00-1.89 (m, 4H), 1.66-1.54 (m, 2H), 1.39 (t, 2H), 0.92 (s, 6H).

EXAMPLE 54

40 N-({2-chloro-5-fluoro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 54A

45 2-chloro-5-fluoro-4-((tetrahydro-2H-pyran-4-yl)methylamino)benzenesulfonamide

[0483] The title compound was prepared by substituting 2-chloro-4,5-difluorobenzenesulfonamide for 4-chloro-3-nitrobenzenesulfonamide and (tetrahydro-2H-pyran-4-yl)methanamine for 4-methylpiperazin-1-amine dihydrochloride in EXAMPLE 6A.

EXAMPLE 54

55 N-({2-chloro-5-fluoro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0484] The title compound was prepared by substituting EXAMPLE 54A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.76 (s, 1H), 11.31 (s, 1H), 8.08 (d, 1H), 7.69 (d, 1H), 7.60 (d, 1H), 7.55 (m,

2H), 7.34 (d, 2H), 7.04 (d, 2H), 6.90 (s, 1H), 6.84 (d, 1H), 6.69 (dd, 1H), 6.45 (dd, 1H), 6.13 (d, 1H), 3.82 (dd, 2H), 3.24 (t, 2H), 3.05 (m, 6H), 2.73 (s, 2H), 2.14 (m, 6H), 1.95 (s, 2H), 1.81 (m, 1H), 1.61 (m, 2H), 1.38 (t, 2H), 1.17 (m, 2H), 0.92 (s, 6H).

5 EXAMPLE 55

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((3-morpholin-4-yl)propyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 **[0485]** The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 2A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.66 (br s, 1H), 8.75 (t, 1H), 8.54 (d, 1H), 8.03 (d, 1H), 7.79 (dd, 1H), 7.54-7.48 (m, 3H), 7.35 (d, 2H), 7.08-7.02 (m, 3H), 6.67 (dd, 1H), 6.38 (m, 1H), 6.20 (d, 1H), 3.61 (t, 4H), 3.43 (q, 2H), 3.29 (m, 2H), 3.06 (br s, 4H), 2.73 (br s, 2H), 2.47 (br s, 4H), 2.18 (m, 6H), 1.95 (br s, 2H), 1.80 (m, 2H), 1.38 (t, 2H), 0.92 (s, 6H).

15

EXAMPLE 56

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

20

EXAMPLE 56A

5-bromo-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridine-3-sulfonamide

25 **[0486]** The title compound was prepared by substituting EXAMPLE 37C for tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 36A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 56B

30 **[0487]** The title compound was prepared by substituting EXAMPLE 56A for EXAMPLE 36B in EXAMPLE 36C.

EXAMPLE 56C

35 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0488] The title compound was prepared by substituting EXAMPLE 56B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.58 (s, 1H), 8.70 (s, 1H), 8.51 (s, 1H), 7.96 (d, 1H), 7.57 (d, 1H), 7.45 (t, 1H), 7.35-7.37 (m, 3H), 7.06 (d, 2H), 6.67 (dd, 1H), 6.33 (d, 1H), 6.26 (s, 1H), 4.56 (d, 2H), 3.76-3.80 (s, 2H), 3.56-3.62 (m, 2H), 3.01-3.10 (m, 4H), 2.14-2.18 (m, 2H), 1.96 (s, 2H), 1.80-1.87 (m, 4H), 1.41 (t, 2H), 0.93 (s, 6H).

40

EXAMPLE 57

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(2-morpholin-4-ylethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

45

EXAMPLE 57A

5-bromo-6-(2-morpholinoethoxy)pyridine-3-sulfonamide

50

[0489] The title compound was prepared by substituting 2-morpholinoethanol for tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 36A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 57B

55

5-cyano-6-(2-morpholinoethoxy)pyridine-3-sulfonamide

[0490] The title compound was prepared by substituting EXAMPLE 57A for EXAMPLE 36A in EXAMPLE 36B.

EXAMPLE 57C

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(2-morpholin-4-ylethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0491] The title compound was prepared by substituting EXAMPLE 57B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.56 (s, 1H), 8.64 (s, 1H), 8.41 (s, 1H), 7.92 (d, 1H), 7.58 (d, 1H), 7.44 (t, 1H), 7.36 (d, 2H), 7.31 (s, 1H), 7.06 (d, 2H), 6.65 (dd, 1H), 6.31 (d, 1H), 6.27 (d, 1H), 4.59 (t, 2H), 3.59 (s, 4H), 3.08 (s, 4H), 2.89 (s, 2H), 2.65 (s, 4H), 2.16-2.18 (m, 2H), 1.97 (s, 2H), 1.41 (t, 2H), 0.93 (s, 6H).

EXAMPLE 58

N-([3-chloro-4-([2-(2-methoxyethoxy)ethyl]sulfonyl)phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 58A

3-chloro-4-(2-(2-methoxyethoxy)ethylthio)benzenesulfonamide

[0492] In a 25 mL microwave tube was added sodium hydride (0.6 g) in tetrahydrofuran (10 mL) to give a suspension. 2-(2-Methoxyethoxy)ethanethiol (1 g) was added slowly. After stirring for 30 minutes, 3-chloro-4-fluorobenzenesulfonamide (1.54 g) dissolved in 10 mL tetrahydrofuran was added slowly. The mixture was heated at 110°C for 30 minutes in a Biotage Initiator microwave reactor. Water was added, the product was extracted with ether (20 mL x3), dried over Na₂SO₄, filtered, and the solvent was removed under reduced pressure. The crude product was purified by flash chromatography on silica eluting with 0-25% ethyl acetate in hexane.

EXAMPLE 58B

3-chloro-4-(2-(2-methoxyethoxy)ethylsulfonyl)benzenesulfonamide

[0493] EXAMPLE 58A (0.15 g) was suspended in acetic acid (3 mL). Peracetic acid (0.4 mL) was added slowly. The mixture was stirred at room temperature overnight, then poured into Na₂S₂O₃ solution, and the product precipitated. After filtration and washing with water, the product was dried under vacuum.

EXAMPLE 58C

N-([3-chloro-4-([2-(2-methoxyethoxy)ethyl]sulfonyl)phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0494] The title compound was prepared by substituting EXAMPLE 58B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.52 (s, 1H), 7.92 (d, 1H), 7.84 (m, 2H), 7.68 (m, 1H), 7.62 (d, 1H), 7.42 (t, 1H), 7.35 (d, 2H), 7.29 (m, 1H), 7.05 (d, 2H), 6.62 (dd, 1H), 6.32 (m, 1H), 6.26 (d, 1H), 3.74 (t, 2H), 3.68 (t, 2H), 3.24 (m, 2H), 3.06 (m, 5H), 3.01 (m, 4H), 2.74 (s, 2H), 2.19 (m, 6H), 1.96 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 59

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([2-(2-methoxyethoxy)ethyl]sulfonyl)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 59A

4-(2-(2-methoxyethoxy)ethylthio)-3-nitrobenzenesulfonamide

[0495] The title compound was prepared by substituting 4-fluoro-3-nitrobenzenesulfonamide for 3-chloro-4-fluorobenzenesulfonamide in EXAMPLE 58A.

EXAMPLE 59B

4-(2-(2-methoxyethoxy)ethylsulfonyl)-3-nitrobenzenesulfonamide

5 **[0496]** The title compound was prepared by substituting EXAMPLE 59A for EXAMPLE 58A in EXAMPLE 58B.

EXAMPLE 59C

10 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([2-(2-methoxyethoxy)ethyl]sulfonyl)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0497] The title compound was prepared by substituting EXAMPLE 59B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.51 (s, 1H), 8.17 (m, 1H), 7.94 (m, 3H), 7.64 (d, 1H), 7.42 (m, 1H), 7.35 (d, 2H), 7.28 (d, 1H), 7.05 (d, 2H), 6.62 (m, 1H), 6.28 (m, 2H), 3.83 (m, 4H), 3.16 (m, 2H), 3.08 (s, 3H), 3.01 (m, 4H), 2.73 (s, 2H), 2.18 (m, 6H), 1.96 (m, 4H), 1.39 (t, 2H), 0.93 (s, 6H)

EXAMPLE 60

20 Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-morpholin-4-ylcyclohexyl]oxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3b]pyridin-5-yloxy)benzamide

EXAMPLE 60A

25 Trans-4-(4-aminocyclohexyloxy)-3-nitrobenzenesulfonamide

[0498] To a solution of *tert*-butyl 4-hydroxycyclohexylcarbamate (0.250 g) in tetrahydrofuran (5 mL) was added sodium hydride (0.186 g). After stirring for 15 minutes, 4-fluoro-3-nitrobenzenesulfonamide (0.256 g) was added as a solution in tetrahydrofuran (1 mL). The reaction was heated to 60°C for 1.5 hours, cooled, and poured into a mixture of dichloromethane (100 mL) and water (25 mL). The aqueous layer was adjusted to pH~4 with 1N aqueous HCl and the organic layer was separated, washed with brine (50 mL), dried over magnesium sulfate, filtered, and concentrated. The residue was loaded onto silica gel (GraceResolv 40 g) and eluted using a gradient of 0.5% to 7.5% methanol/dichloromethane over 30 minutes. This solid was treated with HCl (4.0M in dioxane, 5 mL) at room temperature for 1 hour and concentrated to give the title compound.

35 EXAMPLE 60B

4-(trans-4-morpholinocyclohexyloxy)-3-nitrobenzenesulfonamide

[0499] To EXAMPLE 60A (0.220 g) and 1-bromo-2-(2-bromoethoxy)ethane (0.177 g) in N,N-dimethylformamide (3 mL) was added triethylamine (0.338 mL) and the reaction heated to 70°C for 5 hours. The reaction was cooled and the resulting precipitate was removed by filtration. The reaction was concentrated and loaded onto silica gel and was eluted using a gradient of 0.5% to 7.5% methanol/dichloromethane to give the title compound.

EXAMPLE 60C

45 Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-morpholin-4-ylcyclohexyl]oxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0500] The title compound was prepared by substituting EXAMPLE 60B for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.62 (s, 1H), 8.23 (s, 1H), 7.99 (s, 1H), 7.96 - 7.88 (m, 1H), 7.54 (d, 1H), 7.48 (s, 2H), 7.34 (d, 3H), 7.04 (d, 2H), 6.72 - 6.58 (m, 1H), 6.37 (s, 1H), 6.21 (s, 1H), 4.69 - 4.47 (m, 1H), 3.66 (s, 4H), 3.05 (s, 4H), 2.76 (s, 6H), 2.22 (s, 9H), 1.96 (s, 4H), 1.39 (s, 6H), 0.92 (s, 6H).

55

EXAMPLE 61

N-({5-bromo-6-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino}pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 61 A

5-bromo-6-(1-(tetrahydro-2H-pyran-4-yl)piperidin-4-ylamino)pyridine-3-sulfonamide

[0501] A mixture of EXAMPLE 36A (1.0 g), EXAMPLE 3L (0.95 g) and triethylamine (3.08 mL) in anhydrous dioxane (20 mL) was heated at 110°C overnight. The organic solvent was removed under vacuum. The residue was purified with flash column chromatography on silica gel eluting with 2%-8% methanol/dichloromethane to give the title compound.

EXAMPLE 61B

N-({5-bromo-6-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino}pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0502] The title compound was prepared by substituting EXAMPLE 61A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.59 (s, 1H), 8.35 (s, 1H), 8.00 (s, 2H), 7.55 (d, 1H), 7.46 (m, 2H), 7.34 (d, 2H), 7.04 (d, 2H), 6.63 (dd, 1H), 6.49 (m, 1H), 6.36 (s, 1H), 6.20 (s, 1H), 4.05 (m, 1H), 3.94 (d, 2H), 3.28 (m, 6H), 3.01 (s, 4H), 2.72 (s, 2H), 2.16 (m, 6H), 1.93 (m, 4H), 1.80 (m, 4H), 1.57 (m, 2H), 1.38 (t, 2H), 1.17 (t, 2H), 0.90 (s, 6H).

EXAMPLE 62

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(2-cyanoethyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 62A

4-(2-cyanoethylamino)-3-nitrobenzenesulfonamide

[0503] The title compound was prepared by substituting 3-aminopropanenitrile for EXAMPLE 39B in EXAMPLE 39C.

EXAMPLE 62B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(2-cyanoethyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0504] The title compound was prepared by substituting EXAMPLE 62A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (501 MHz, pyridine-d₅) δ 13.04 (s, 1H), 9.24 (d, 1H), 9.04 (t, 1H), 8.43 (d, 1H), 8.38 (dd, 1H), 8.13 (d, 1H), 7.64 - 7.68 (m, 2H), 7.44 (ddd, 2H), 7.07 (ddd, 2H), 7.02 (d, 1H), 6.76 (dd, 1H), 6.55 (d, 1H), 6.48 (dd, 1H), 3.83 (q, 2H), 3.07 (d, 4H), 2.98 (t, 2H), 2.77 (s, 2H), 2.26 (s, 2H), 2.11 - 2.17 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 63

Cis-4-(4-{[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl}piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0505] The title compound was prepared by substituting EXAMPLE 15H for EXAMPLE 3J and EXAMPLE 39C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (501 MHz, pyridine-d₅) δ 13.09 (s, 1H), 9.30 (d, 1H), 8.64 (d, 1H), 8.43 (d, 1H), 8.38 (dd, 1H), 8.10 (d, 1H), 7.68 (dt, 2H), 7.46 (ddd, 2H), 7.12 (ddd, 2H), 6.91 (d, 1H), 6.72 (dd, 1H), 6.51 (dd, 1H), 6.49 (d, 1H), 5.69 (s, 2H), 4.40 (s, 2H), 3.69 - 3.73 (m, 4H), 3.68 (s, 1H), 2.95 - 3.02 (m, 4H), 2.84 (s, 2H), 2.40 - 2.46 (m, 4H), 2.21 (s, 2H), 2.08 - 2.15 (m, 5H), 1.76 - 1.84 (m, 2H), 1.55 - 1.63 (m, 6H), 1.29 (s, 6H).

EXAMPLE 64

Trans-N- [4-({4-[bis(cyclopropylmethyl)amino]cyclohexyl}amino)-3-nitrophenyl]sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 64A

tert-butyl (trans)-4-(bis(cyclopropylmethyl)amino)cyclohexylcarbamate

[0506] The title compound was prepared by substituting cyclopropanecarbaldehyde for 4'-chlorobiphenyl-2-carboxaldehyde and tert-butyl (trans)-4-aminocyclohexylcarbamate for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A.

EXAMPLE 64B

(trans)-N¹,N¹-bis(cyclopropylmethyl)cyclohexane-1,4-diamine dihydrochloride

[0507] To a solution of EXAMPLE 64A (1.4 g) in dichloromethane (10 ml) was added hydrogen chloride (10 ml, 4M in dioxane) and the reaction was stirred for 16 hours at room temperature. The reaction mixture was diluted with ether and pure product was filtered off.

EXAMPLE 64C

Trans-4-(4-(bis(cyclopropylmethyl)amino)cyclohexylamino)-3-nitrobenzenesulfonamide

[0508] The title compound was prepared by substituting EXAMPLE 64B for EXAMPLE 39B in EXAMPLE 39C.

EXAMPLE 64D

Trans-N- {[4-({4-[bis(cyclopropylmethyl)amino]cyclohexyl}amino)-3-nitrophenyl]sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0509] The title compound was prepared by substituting EXAMPLE 64C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, pyridine-d₅) δ 13.06 (s, 1H), 9.30 (d, 1H), 8.44 (d, 1H), 8.41 (dd, 1H), 8.37 (d, 1H), 8.12 (d, 1H), 7.67 (d, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 7.00 (d, 1H), 6.75 (dd, 1H), 6.53 (d, 1H), 6.49 (dd, 1H), 3.36 - 3.43 (m, 1H), 3.02 - 3.09 (m, 4H), 2.87 - 2.94 (m, 1H), 2.77 (s, 2H), 2.47 (d, 4H), 2.25 (t, 2H), 2.11 - 2.16 (m, 4H), 2.08 (d, 2H), 1.97 (s, 2H), 1.84 (d, 2H), 1.39 (t, 2H), 1.26 - 1.35 (m, 4H), 0.90 - 0.98 (m, 8H), 0.50 - 0.56 (m, 4H), 0.18 - 0.23 (m, 4H).

EXAMPLE 65

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(1-methylpiperidin-4-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 65A

4-((1-methylpiperidin-4-yl)methylamino)-3-nitrobenzenesulfonamide

[0510] The title compound was prepared by substituting 4-aminomethyl-1-methyl piperidine for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 65B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(1-methylpiperidin-4-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0511] The title compound was prepared by substituting EXAMPLE 65A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, dichloromethane-d₂) δ 9.57 (bs, 1H), 8.78 (d, 1H), 8.41 (d, 1H), 8.14 (d, 1H), 7.90 (m, 2H), 7.64 (d, 1H), 7.45 (d, 1H), 7.23 (d, 2H), 6.95 (d, 2H), 6.76 (d, 1H), 6.59 (dd, 1H), 6.51 (d, 1H), 6.09 (d, 1H), 3.21 (m, 2H), 3.08 (m, 4H), 3.02 (m, 2H), 2.74 (s, 2H), 2.33 (s, 3H), 2.21-2.17 (m, 6H), 2.16-2.02 (m, 3H), 1.97 (br.s, 2H), 1.78 (m, 4H), 1.41

(t, 2H), 0.94 (s, 6H).

EXAMPLE 66

5 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(morpholin-3-ylmethyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 66A

10 tert-butyl 3-((2-nitro-4-sulfamoylphenylamino)methyl)morpholine-4-carboxylate

[0512] The title compound was prepared by substituting tert-butyl 3-(aminomethyl)morpholine-4-carboxylate for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 66B

tert-butyl 3-((4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)sulfamoyl)-2-nitrophenylamino)methyl)morpholine-4-carboxylate

20 **[0513]** The title compound was prepared by substituting EXAMPLE 66A for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1F, with the exception that the product was purified on a silica gel column eluted with 4% methanol in dichloromethane.

EXAMPLE 66C

25

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(morpholin-3-ylmethyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

30 **[0514]** A solution of EXAMPLE 66B in 50% trifluoroacetic acid and dichloromethane mixture was stirred at ambient temperature for 2 hours. The solvents were evaporated and the residue was purified on a reverse phase HPLC using a gradient of 20-80% acetonitrile in water containing 10 mM ammonium acetate. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.61 (s, 1H), 8.52 (bs, 1H), 8.49 (d, 1H), 7.98 (d, 1H), 7.78 (d, 1H), 7.54 (d, 1H), 7.46 (s, 1H), 7.42 (s, 1H), 7.34 (d, 2H), 7.04 (m, 3H), 6.65 (dd, 1H), 6.34 (s, 1H), 6.21 (d, 1H), 3.89 (d, 1H), 3.76 (d, 1H), 3.55-3.46 (m, 2H), 3.40-3.35 (m, 4H), 3.04 (m, 4H), 2.91 (t, 1H), 2.73 (s, 2H), 2.20-2.12 (m, 6H), 1.95 (s, 2H), 1.38 (t, 2H), 0.92 (s, 6H).

35

EXAMPLE 67

4-(4-{[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl}piperazin-1-yl)-N-({4-[(4-methylpiperazin-1-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

40

[0515] The title compound was prepared as described in EXAMPLE 1G by replacing EXAMPLE 1E and EXAMPLE 1F with EXAMPLE 15H and EXAMPLE 6A, respectively. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.58 (s, 1H), 9.04 (s, 1H), 8.44 (d, 1H), 7.97 (d, 1H), 7.76 (dd, 1H), 7.49 (m, 4H), 7.38 (d, 2H), 7.14 (d, 2H), 6.64 (dd, 1H), 6.34 (d, 1H), 6.21 (d, 1H), 4.12 (s, 2H), 3.03 (m, 6H), 2.85 (m, 5H), 2.29 (m, 4H), 2.18 (m, 6H), 1.20 (s, 6H).

45

EXAMPLE 68

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-morpholin-4-ylbut-2-ynyl)oxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50

EXAMPLE 68A

4-morpholinobut-2-yn-1-ol

55 **[0516]** To a solution of morpholine (4.36 g) in toluene (15 mL) was added 4-chlorobut-2-yn-1-ol (2.09 g) in toluene (5 mL). The solution was stirred at 85 °C for 3 hours. After cooling, the solid was filtered off. The filtrate was subjected to vacuum distillation to give the pure title compound.

EXAMPLE 68B

4-(4-morpholinobut-2-ynyloxy)-3-nitrobenzenesulfonamide

- 5 **[0517]** The title compound was prepared by substituting EXAMPLE 68A for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 68C

- 10 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-morpholin-4-ylbut-2-ynyloxy)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0518]** The title compound was prepared by substituting EXAMPLE 68B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 8.36 (s, 1H), 8.08 (d, 1H), 8.03 (d, 1H), 7.47-7.53 (m, 4H), 7.35 (d, 2H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.40 (dd, 1H), 6.20 (d, 1H), 5.15 (s, 2H), 3.52-3.55 (m, 4H), 3.09 (s, 4H), 2.84 (br s, 2H), 2.23-2.40 (m, 6H), 2.12-2.18 (m, 2H), 1.96 (s, 2H), 1.39 (t, 2H), 0.92 (s, 6H).

EXAMPLE 69

- 20 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-([5-ethynyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 69A

- 25 6-((tetrahydro-2H-pyran-4-yl)methoxy)-5-((triisopropylsilyl)ethynyl)pyridine-3-sulfonamide

- [0519]** EXAMPLE 36B (0.176 g), bis(triphenylphosphine)palladium(II) chloride (0.176 g), copper(I) iodide (0.010 g), N,N-dimethylacetamide (2.5 mL) and triethylamine (0.105 mL) were combined, flushed with nitrogen and stirred for 2 minutes. (Triisopropylsilyl)acetylene (0.135 mL) was added and the reaction mixture was flushed with nitrogen again, heated at 60 °C overnight, diluted with ethyl acetate, washed with water and brine, dried (MgSO₄), filtered, concentrated and chromatographed on silica gel with 10-30% ethyl acetate in hexanes as the eluent to give the product.

EXAMPLE 69B

- 35 5-ethynyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)pyridine-3-sulfonamide

- [0520]** EXAMPLE 69A (0.205 g) in tetrahydrofuran (3 mL) at ambient temperature was treated with tetrabutyl ammonium fluoride (1 M in tetrahydrofuran, 0.906 mL) and stirred at ambient temperature for 4 hours. Additional tetrabutyl ammonium fluoride (1 M in tetrahydrofuran, 1.8 mL) was added and the mixture was heated at 40 °C for 45 minutes. Solid tetrabutyl ammonium fluoride (0.253 g) was added and heating was continued for 30 minutes. The reaction mixture was concentrated and then chromatographed on silica gel using 0-2% methanol in dichloromethane as the eluent to give the product.

EXAMPLE 69

- 45 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-([5-ethynyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0521]** The title compound was prepared by substituting EXAMPLE 69B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 11.41 (s, 1H), 8.58 (d, 1H), 8.19 (d, 1H), 8.05 (d, 1H), 7.53 (m, 3H), 7.35 (d, 2H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.41 (dd, 1H), 6.18 (d, 1H), 4.56 (s, 1H), 4.24 (d, 2H), 3.87 (dd, 2H), 3.38 (m, 3H), 3.07 (m, 4H), 2.86 (m, 2H), 2.29 (m, 5H), 2.04 (m, 3H), 1.64 (dd, 2H), 1.34 (m, 4H), 0.93 (s, 6H).

55

EXAMPLE 70

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-oxo-3,4-dihydroquinazolin-6-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 70A

4-amino-3-cyanobenzenesulfonamide

[0522] 3-Cyano-4-fluorobenzene-1-sulfonyl chloride (1.1 g) was dissolved in dioxane (4 mL). The solution was cooled to 0°C and 7 mL of an ammonia (7N in methanol) solution was added. After the addition was complete, the ice bath was removed and the reaction was stirred at room temperature for 24 hours. After concentration of the reaction mixture, the crude material was purified by flash chromatography eluting with a gradient of 30-100% ethyl acetate/hexanes.

EXAMPLE 70B

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-(4-amino-3-cyanophenylsulfonyl)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzamide

[0523] The title compound was prepared by substituting EXAMPLE 70A for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G.

EXAMPLE 70C

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-(4-amino-3-carbamoylphenylsulfonyl)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzamide

[0524] To a solution of EXAMPLE 70B (90 mg) in ethanol (2 mL) was added tetrahydrofuran (2 mL), hydrogen peroxide (30%, 1 mL) and 1M sodium hydroxide solution (0.48 mL), followed by an additional 2 mL of tetrahydrofuran. The reaction was heated to 45°C for 30 minutes, cooled, and then quenched with 5% HCl solution and extracted twice with dichloromethane. The extracts were combined and concentrated to obtain the product.

EXAMPLE 70D

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-oxo-3,4-dihydroquinazolin-6-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0525] EXAMPLE 70C (80 mg) was combined with trimethyl orthoformate (2.3 mL) and trifluoroacetic acid (0.03 mL) and the resulting solution was stirred at room temperature for 4 hours. The mixture was purified by flash chromatography, eluting with a gradient of 3-10% methanol/dichloromethane. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 12.61 (s, 1H), 11.71 (s, 1H), 8.65 (d, 1H), 8.24 (s, 1H), 8.17 (dd, 1H), 8.04 (m, 1H), 7.73 (d, 1H), 7.57 (d, 1H), 7.51 (m, 2H), 7.39 (d, 2H), 7.07 (d, 2H), 6.70 (dd, 1H), 6.40 (m, 1H), 6.24 (br s, 1H), 3.61 (m, 6H), 3.03 (m, 2H), 2.75 (m, 2H), 2.17 (m, 2H), 2.01 (m, 2H), 1.44 (m, 2H), 0.94 (s, 6H).

EXAMPLE 71

Trans-4-(4-([8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl)methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 71 A

8-chlorospiro[4.5]dec-7-ene-7-carbaldehyde

[0526] To a solution of N,N-dimethylformamide (2.81 mL) in dichloromethane (40 mL) was added dropwise POCl₃ (2.78 mL) at 0°C. The reaction mixture was warmed up to room temperature and spiro[4.5]decan-8-one (3.95 g) in dichloromethane (5 mL) was added dropwise. The mixture was stirred overnight. The reaction was quenched with cold aqueous sodium acetate and the resulting mixture was extracted with ether and the organic layer was dried over Na₂SO₄, filtered, and concentrated to provide the title compound.

EXAMPLE 71B

8-(4-chlorophenyl)spiro[4.5]dec-7-ene-7-carbaldehyde

- 5 **[0527]** To a suspension of EXAMPLE 71A (3 g) in water (50 mL) was added 4-chlorophenylboronic acid (2.83 g), tetrabutylammonium (4.87 g), potassium carbonate (6.26 g) and palladium(II) acetate (0.169 g). The reaction mixture was stirred at 45°C for 5 hours and extracted with dichloromethane. The organic layer was concentrated and the residue was loaded onto a silica gel column, and eluted with 5-20% ethyl acetate in hexane to give the title compound.

10 EXAMPLE 71C

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl)methyl)piperazin-1-yl)benzoate

- 15 **[0528]** To a solution of EXAMPLE 71B (274 mg) in dichloroethane (3.5 mL) was added EXAMPLE 15F (387 mg) and sodium triacetoxyborohydride (317 mg). The reaction mixture was stirred overnight. Sodium cyanoborohydride (37.6 mg) was added and the resulting mixture stirred overnight. The reaction was quenched with water and diluted with dichloromethane. The mixture was washed with water extensively and the organic layer was dried over Na₂SO₄, filtered, and concentrated to provide the title compound.

20

EXAMPLE 71D

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl)methyl)piperazin-1-yl)benzoic acid

25

- [0529]** The title compound was prepared as described in EXAMPLE 3J using EXAMPLE 71C in place of EXAMPLE 3I.

EXAMPLE 71E

- 30 Trans-4-(4-([8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl)piperazin-1-yl)-N-([4-([4-(morpholin-4-yl)cyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0530]** The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 71D and EXAMPLE 9C in place of EXAMPLE 3J and EXAMPLE 11B, respectively. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.64 (s, 1H), 8.51 (s, 1H), 8.15 (d, 1H), 8.01 (d, 1H), 7.76 (d, 1H), 7.44 - 7.53 (m, 3H), 7.34 (d, 2H), 7.07 (d, 3H), 6.66 (dd, 1H), 6.37 (dd, 1H), 6.20 (d, 1H), 3.50 - 3.70 (m, 5H), 3.04 (s, 4H), 2.55 - 2.76 (m, 5H), 2.34 - 2.39 (m, 1H), 2.20 (d, 6H), 2.03 (s, 4H), 1.91 (s, 2H), 1.61 (q, 4H), 1.51 (t, 2H), 1.36 - 1.46 (m, 8H).

35

EXAMPLE 72

40

Cis-4-(4-([4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl)piperazin-1-yl)-N-([4-([4-(methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0531]** The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 15H and 29A in place of EXAMPLE 3J and EXAMPLE 11B, respectively. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 11.45 (s, 1H), 8.59 (t, 1H), 8.56 (d, 1H), 8.04 (d, 1H), 7.79 (dd, 1H), 7.54 (d, 1H), 7.47 - 7.52 (m, 2H), 7.37 (d, 2H), 7.13 (d, 2H), 7.08 (d, 1H), 6.68 (dd, 1H), 6.35 - 6.42 (m, 1H), 6.19 (d, 1H), 4.11 (s, 2H), 3.37 (s, 1H), 3.26 (t, 2H), 3.20 (s, 3H), 3.07 (s, 4H), 2.83 (s, 2H), 2.17 (d, 6H), 1.81 (dd, 2H), 1.64 - 1.73 (m, 1H), 1.48 (dd, 2H), 1.23 - 1.41 (m, 4H), 1.18 (s, 6H).

45

50 EXAMPLE 73

4-(4-([8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl)piperazin-1-yl)-N-([4-([4-(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- 55 **[0532]** The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 71D and EXAMPLE 37D in place of EXAMPLE 3J and EXAMPLE 11B, respectively. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 8.37 (s, 1H), 7.98 - 8.11 (m, 2H), 4.38 (d, 2H), 3.74 - 3.82 (m, 2H), 3.54 - 3.64 (m, 2H), 3.44 (s, 1H), 3.08 (s, 3H), 2.58 - 2.89 (m, 2H), 2.13 - 2.35 (m, 4H), 2.04 (s, 2H), 1.78 - 1.93 (m, 4H), 1.57 - 1.65 (m, 4H), 1.52 (t, 2H), 1.36 - 1.47 (m, 4H).

EXAMPLE 74

Trans-4-(4-([8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl)methyl]piperazin-1-yl)-N-[(4-((4-methoxycyclohexyl)methyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

5

[0533] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 71D and EXAMPLE 34B in place of EXAMPLE 3J and EXAMPLE 11B, respectively. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 11.39 (s, 1H), 8.58 (t, 1H), 8.56 (d, 1H), 8.04 (d, 1H), 7.79 (dd, 1H), 7.47 - 7.55 (m, 3H), 7.34 (d, 2H), 7.07 (d, 3H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.19 (d, 1H), 3.25 (t, 2H), 3.22 (s, 3H), 3.06 (s, 5H), 2.71 (s, 2H), 2.21 (s, 6H), 1.94 - 2.06 (m, 4H), 1.79 (d, 2H), 1.57 - 1.65 (m, 5H), 1.51 (t, 2H), 1.39 (t, 4H), 0.95 - 1.11 (m, 4H).

10

EXAMPLE 75

4-(4-([2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([3-nitro-4-((tetrahydro-2H-pyran-4-ylmethyl)amino)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

15

EXAMPLE 75A

methyl 5,5-dimethyl-2-(trifluoromethylsulfonyloxy)cyclohex-1-enecarboxylate

20

[0534] The title compound was prepared by substituting 4,4-dimethyl-2-methoxycarbonylcyclohexanone for 5,5-dimethyl-2-methoxycarbonylcyclohexanone in EXAMPLE 3A.

EXAMPLE 75B

25

methyl 2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-enecarboxylate

[0535] The title compound was prepared by substituting EXAMPLE 75A for EXAMPLE 3A in EXAMPLE 3B.

30

EXAMPLE 75C

(2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-enyl)methanol

[0536] The title compound was prepared by substituting EXAMPLE 75B for EXAMPLE 3B in EXAMPLE 3C.

35

EXAMPLE 75D

2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-enecarbaldehyde

40

[0537] To a solution of EXAMPLE 75C (2.8 g) in dichloromethane (50 mL) was added Dess-Martin Periodinane (5.68 g). The reaction mixture was stirred at room temperature for 3 hours and diluted with ether and washed with 5% NaOH and brine. The organic layer was dried over Na₂SO₄, filtered, and concentrated. The residue was purified by flash chromatography using 20% ethyl acetate in hexanes to provide the title compound.

45

EXAMPLE 75E

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoate

50

[0538] The title compound was prepared by replacing 4'-chlorobiphenyl-2-carboxaldehyde with EXAMPLE 75D and tert-butyl piperazine-1-carboxylate with EXAMPLE 15F in EXAMPLE 1A.

EXAMPLE 75F

55

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoic acid

[0539] The title compound was prepared as described in EXAMPLE 15H by replacing EXAMPLE 15G with EXAMPLE

75E.

EXAMPLE 75G

5 4-(4-([2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-((tetrahydro-2H-pyran-4-ylmethyl)amino)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0540] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 75F and EXAMPLE 1F in place of EXAMPLE 3J and EXAMPLE 11B, respectively. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 11.38 (s, 1H), 8.60 (t, 1H), 8.56 (d, 1H), 8.04 (d, 1H), 7.80 (dd, 1H), 7.47 - 7.55 (m, 3H), 7.31 - 7.36 (m, 2H), 7.05 - 7.13 (m, 3H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.18 (d, 1H), 3.85 (dd, 2H), 3.22 - 3.31 (m, 4H), 3.07 (s, 4H), 2.67 - 2.78 (m, 2H), 2.19 (s, 6H), 1.82 - 1.98 (m, 3H), 1.56 - 1.66 (m, 2H), 1.39 (t, 2H), 1.17 - 1.33 (m, 3H), 0.93 (s, 6H).

EXAMPLE 76

15 4-(4-([2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0541] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 75F and EXAMPLE 36C in place of EXAMPLE 3J and EXAMPLE 11B, respectively. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.62 (s, 1H), 8.73 (s, 1H), 8.52 (s, 1H), 7.98 (d, 1H), 7.56 (d, 1H), 7.45 - 7.51 (m, 1H), 7.43 (s, 1H), 7.37 (d, 2H), 7.10 (d, 2H), 6.68 (dd, 1H), 6.35 (dd, 1H), 6.25 (s, 1H), 4.29 (d, 2H), 3.88 (dd, 2H), 3.12 (d, 4H), 2.21 (s, 2H), 2.00 - 2.11 (m, 1H), 1.95 (s, 2H), 1.64 (dd, 2H), 1.27 - 1.46 (m, 4H), 0.95 (s, 6H)

EXAMPLE 77

tert-butyl 3-{[4-([4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino)sulfonyl]-2-nitrophenoxy)methyl}morpholine-4-carboxylate

EXAMPLE 77A

tert-butyl 3-((2-nitro-4-sulfamoylphenoxy)methyl)morpholine-4-carboxylate

[0542] The title compound was prepared as described in EXAMPLE 12A by replacing (1,4-dioxan-2-yl)methanol with tert-butyl 3-(hydroxymethyl)morpholine-4-carboxylate.

EXAMPLE 77B

40 tert-butyl 3-{[4-([4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino)sulfonyl]-2-nitrophenoxy)methyl}morpholine-4-carboxylate

[0543] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 77A in place of EXAMPLE 11B. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 8.36 (s, 1H), 8.01 - 8.11 (m, 2H), 7.47 - 7.61 (m, 4H), 7.35 (d, 2H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.39 (d, 1H), 6.20 (s, 1H), 4.41 - 4.52 (m, 2H), 4.15 - 4.28 (m, 1H), 3.59 - 3.95 (m, 3H), 3.51 (d, 1H), 3.34 - 3.43 (m, 1H), 3.10 (s, 5H), 2.84 (s, 2H), 2.28 (s, 4H), 2.15 (s, 2H), 1.96 (s, 2H), 1.20 - 1.45 (m, 12H), 0.92 (s, 6H).

EXAMPLE 78

50 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(morpholin-3-ylmethoxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0544] EXAMPLE 77B (100 mg) in dichloromethane (10 mL) at 0°C was treated with trifluoroacetic acid (5 mL) for 20 minutes. The reaction mixture was concentrated. The residue was purified by reverse phase HPLC on a C18 column using a gradient of 35-60% acetonitrile in 0.1% trifluoroacetic acid water to give the title compound as a trifluoroacetate salt. The trifluoroacetic acid salt was dissolved in dichloromethane (10 mL) and washed with 50% aqueous NaHCO₃. The organic layer was dried over anhydrous Na₂SO₄ and concentrated to give the title compound. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.56 (s, 1H), 8.23 (d, 1H), 7.94 (d, 1H), 7.90 (dd, 1H), 7.57 (d, 1H), 7.42 - 7.46 (m, 1H), 7.31 -

7.37 (m, 3H), 7.25 (d, 1H), 7.01 - 7.09 (m, 2H), 6.64 (dd, 1H), 6.29 - 6.37 (m, 1H), 6.24 (d, 1H), 4.17 - 4.31 (m, 2H), 3.90 - 4.05 (m, 1H), 3.77 - 3.85 (m, 1H), 3.45 - 3.59 (m, 4H), 2.94 - 3.13 (m, 6H), 2.76 (s, 2H), 2.18 (d, 6H), 1.96 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

5 EXAMPLE 79

4-(4-([8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl)piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 **[0545]** The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 71D and EXAMPLE 1F in place of EXAMPLE 3J and EXAMPLE 11B, respectively. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 11.38 (s, 1H), 8.60 (t, 1H), 8.56 (d, 1H), 8.04 (d, 1H), 7.77 - 7.84 (m, 1H), 7.45 - 7.56 (m, 3H), 7.34 (d, 2H), 7.04 - 7.13 (m, 3H), 6.68 (dd, 1H), 6.39 (d, 1H), 6.19 (d, 1H), 3.85 (dd, 2H), 3.22 - 3.31 (m, 4H), 3.07 (s, 4H), 2.71 (s, 2H), 2.21 (s, 6H), 2.03 (s, 2H), 1.81 - 1.94 (m, 1H), 1.56 - 1.68 (m, 6H), 1.51 (t, 2H), 1.34 - 1.45 (m, 4H), 1.20 - 1.33 (m, 2H).

15 EXAMPLE 80

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-(methylsulfonyl)piperidin-4-yl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

20 **[0546]** The title compound was prepared by substituting 1-(methylsulfonyl)piperidin-4-amine for 1-acetyl-piperidin-4-amine in EXAMPLE 53B. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.67 (br s, 1H), 8.57 (d, 1H), 8.25 (d, 1H), 8.04 (d, 1H), 7.83 (dd, 1H), 7.54-7.46 (m, 3H), 7.35 (d, 2H), 7.17 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.39 (m, 1H), 6.20 (d, 1H), 3.80 (m, 1H), 3.57 (m, 2H), 3.08 (br s, 4H), 2.95 (td, 2H), 2.92 (s, 3H), 2.85-2.72 (m, 2H), 2.30-2.10 (m, 6H), 2.07-1.93 (m, 4H), 1.70 (m, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 81

30 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({4-[(1,1-dioxotetrahydro-2H-thiopyran-4-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 81 A

1,1-Dioxotetrahydro-2H-thiopyran-4-amine

35 **[0547]** N-Benzyl-1,1-dioxotetrahydro-2H-thiopyran-4-amine (2.00 g) was added to ethanol (40 mL) in a pressure bottle. Palladium hydroxide on carbon (0.587 g,) was added and the solution was stirred under 30 psi of hydrogen at room temperature for 2 hours. The mixture was filtered through a nylon membrane and the solvent was removed under vacuum.

40 EXAMPLE 81B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({4-[(1,1-dioxotetrahydro-2H-thiopyran-4-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

45 **[0548]** The title compound was prepared by substituting EXAMPLE 81A for 1-acetyl-piperidin-4-amine in EXAMPLE 53B. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.65 (br s, 1H), 8.55 (d, 1H), 8.25 (d, 1H), 8.03 (d, 1H), 7.86 (dd, 1H), 7.52-7.47 (m, 3H), 7.35 (d, 2H), 7.17 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.39 (m, 1H), 6.21 (d, 1H), 4.05 (m, 1H), 3.22-3.00 (m, 8H), 2.79 (br s, 2H), 2.31-2.11 (m, 10H), 1.96 (br s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

50 EXAMPLE 82

N-([4-chloro-3-nitrophenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

55 **[0549]** The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and 4-chloro-3-nitrobenzenesulfonamide for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.60 (br s, 1H), 8.38 (br s, 1H), 7.96 (d, 1H), 7.91 (d, 1H), 7.68 (d, 1H), 7.58 (d, 1H), 7.46 (t, 1H), 7.39-7.35 (m, 3H), 7.07 (d, 2H), 6.67 (dd, 1H), 6.34 (m, 1H), 6.28 (d, 1H), 3.31 (br s, 2H), 3.17 (br s, 8H), 2.18 (m, 2H), 1.98 (br s, 2H), 1.42 (t, 2H), 0.94 (s, 6H).

EXAMPLE 83

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-[[1-(2,2,2-trifluoroethyl)piperidin-4-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 83A

3-Nitro-4-[1-(2,2,2-trifluoro-ethyl)-piperidin-4-ylamino]-benzenesulfonamide

[0550] The title compound was prepared by substituting 1-(2,2,2-trifluoroethyl)piperidin-4-amine hydrochloride for (tetrahydropyran-4-yl)methylamine in EXAMPLE 6A.

EXAMPLE 83B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-[[1-(2,2,2-trifluoroethyl)piperidin-4-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0551] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 82A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.66 (br s, 1H), 8.56 (d, 1H), 8.24 (d, 1H), 8.04 (d, 1H), 7.81 (dd, 1H), 7.52 (dd, 2H), 7.48 (d, 1H), 7.35 (d, 2H), 7.15 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.38 (m, 1H), 6.20 (d, 1H), 3.68 (m, 1H), 3.22 (q, 2H), 3.07 (br s, 4H), 2.90 (m, 2H), 2.75 (br s, 2H), 2.29-2.12 (m, 8H), 1.97-1.86 (m, 4H), 1.63 (m, 2H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 84

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({5-cyano-6-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]oxy}pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 84A

1-(tetrahydro-2H-pyran-4-yl)piperidin-4-ol

[0552] Piperidin-4-ol (7.8 g) and dihydro-2H-pyran-4(3H)-one (5.0 g) were dissolved in titanium(IV) isopropoxide (30 mL) and the reaction was stirred at room temperature overnight. Methanol (40 mL) was added and the reaction was cooled to 0°C. Then NaBH₄ (3.8 g) was added in portions over one hour. After 2 hours 1N aqueous NaOH was added, followed by ethyl acetate addition. After filtration through celite the layers were separated, the aqueous layer extracted with ethyl acetate, and the combined organic layers were dried over Na₂SO₄. The crude was purified by column chromatography using dichloromethane having 5-10% 7N NH₃ in methanol.

EXAMPLE 84B

5-bromo-6-(1-(tetrahydro-2H-pyran-4-yl)piperidin-4-yloxy)pyridine-3-sulfonamide

[0553] The title compound was prepared by substituting EXAMPLE 84A for (tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 36A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 84C

5-cyano-6-(1-(tetrahydro-2H-pyran-4-yl)piperidin-4-yloxy)pyridine-3-sulfonamide

[0554] The title compound was prepared by substituting EXAMPLE 84B for EXAMPLE 36B in EXAMPLE 36C.

EXAMPLE 84D

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({5-cyano-6-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]oxy}pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0555] The title compound was prepared by substituting EXAMPLE 84C for EXAMPLE 11B in EXAMPLE 11D. ¹H

NMR (500MHz, dimethylsulfoxide-d₆) δ 11.50 (s, 1H), 8.60 (d, 1H), 8.37 (d, 1H), 7.90 (d, 1H), 7.60 (d, 1H), 7.42 (dd, 1H), 7.35 (d, 2H), 7.25 (d, 1H), 7.04 (d, 2H), 6.63 (dd, 1H), 6.28 (m, 1H), 6.24 (d, 1H), 5.30 (br s, 1H), 4.50 (d, 2H), 3.95 (dd, 2H), 3.30 (m, 5H), 3.02 (br s, 4H), 2.95 (br s, 2H), 2.24 (br s, 4H), 2.17 (br m, 4H), 1.96 (s, 2H), 1.90 (br m, 4H), 1.60 (br m, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

5

EXAMPLE 85

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-isopropyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10

EXAMPLE 85A

5-isopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)pyridine-3-sulfonamide

15

[0556] EXAMPLE 36B (0.176 g), 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (0.041 g), and palladium(II) acetate (0.011 g) were combined in a 10 mL oven-dried flask. Tetrahydrofuran (1 mL) was added and the mixture was flushed with nitrogen and stirred at ambient temperature for 5 minutes. 2-Propylzinc bromide solution (0.5 M in tetrahydrofuran) (1.5 mL) was added and stirring was continued under nitrogen overnight. Additional 2-2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (0.041 g) and palladium(II) acetate (0.011 g) were added. The mixture was flushed with nitrogen and stirred at ambient temperature for 5 minutes. 2-Propylzinc bromide solution (0.5 M in tetrahydrofuran) (1.5 mL) was added and stirring was continued under nitrogen for 2.5 days. The reaction mixture was diluted with ethyl acetate, washed with water and brine, dried (MgSO₄), filtered, concentrated, and chromatographed on silica gel with 0 to 3% methanol in CH₂Cl₂ as the eluent. The obtained material was chromatographed on silica gel a second time with 10-40% ethyl acetate in CH₂Cl₂ as the eluent, triturated with diethyl ether and dried under vacuum at 45 °C to give the product.

20

25

EXAMPLE 85B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-isopropyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

30

[0557] The title compound was prepared by substituting EXAMPLE 85A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.70 (s, 1H), 8.49 (m, 1H), 8.04 (d, 1H), 7.90 (m, 1H), 7.57 (m, 1H), 7.52 (t, 1H), 7.48 (dd, 1H), 7.34 (d, 2H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.41 (dd, 1H), 6.17 (s, 1H), 4.19 (m, 2H), 3.88 (m, 2H), 3.30 (m, 2H), 3.05 (m, 5H), 2.77 (s, 2H), 2.21 (s, 4H), 2.14 (s, 2H), 2.03 (m, 1H), 1.95 (s, 2H), 1.64 (m, 2H), 1.34 (m, 4H), 1.12 (d, 6H), 0.92 (s, 6H).

35

EXAMPLE 86

N-([3-chloro-5-fluoro-4-((tetrahydro-2H-pyran-4-ylmethyl)amino)phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

40

EXAMPLE 86A

3-fluoro-4-((tetrahydro-2H-pyran-4-yl)methylamino)benzenesulfonamide

45

[0558] The title compound was prepared by substituting 3,4-difluorobenzenesulfonamide for 4-chloro-3-nitrobenzenesulfonamide and (tetrahydro-2H-pyran-4-yl)methanamine for 4-methylpiperazin-1-amine dihydrochloride in EXAMPLE 6A.

50

EXAMPLE 86B

3-chloro-5-fluoro-4-((tetrahydro-2H-pyran-4-yl)methylamino)benzenesulfonamide

[0559] The title compound was prepared by substituting EXAMPLE 86A for EXAMPLE 52B in EXAMPLE 52C.

55

EXAMPLE 86C

N-({3-chloro-5-fluoro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0560] The title compound was prepared by substituting EXAMPLE 86B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.72 (s, 1H), 11.20 (s, 1H), 8.08 (d, 1H), 7.61 (m, 2H), 7.50 (m, 3H), 7.34 (d, 2H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.42 (dd, 1H), 6.16 (d, 1H), 6.09 (m, 1H), 3.81 (dd, 2H), 3.25 (m, 4H), 3.07 (m, 4H), 2.76 (s, 2H), 2.18 (m, 6H), 1.95 (s, 2H), 1.72 (m, 1H), 1.53 (d, 2H), 1.38 (t, 2H), 1.16 (m, 2H), 0.92 (s, 6H).

EXAMPLE 87

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-2-(1H-indol-5-yloxy)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)benzamide

EXAMPLE 87A

methyl 2-(1H-indol-5-yloxy)-4-fluorobenzoate

[0561] The title compound was prepared by substituting 5-hydroxyindole for EXAMPLE 3G in EXAMPLE 3H.

EXAMPLE 87B

methyl 2-(1H-indol-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoate

[0562] The title compound was prepared by substituting EXAMPLE 87A for EXAMPLE 3H in EXAMPLE 3I.

EXAMPLE 87C

2-(1H-indol-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoic acid

[0563] The title compound was prepared by substituting EXAMPLE 87B for EXAMPLE 3I in EXAMPLE 3J.

EXAMPLE 87D

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-2-(1H-indol-5-yloxy)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)benzamide

[0564] The title compound was prepared by substituting EXAMPLE 87C for EXAMPLE 1E in EXAMPLE 1G, except here the crude was purified by preparative HPLC using a 250 x 50 mm C18 column and eluting with 20-100% CH₃CN vs. 0.1% trifluoroacetic acid in water, giving the product as a trifluoroacetate salt. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.40 (br s, 1H), 11.17 (s, 1H), 9.50 (v br s, 1H), 8.61 (t, 1H), 8.57 (d, 1H), 7.77 (dd, 1H), 7.70 (br s, 1H), 7.50 (m, 5H), 7.36 (m, 5H), 7.10 (s, 1H), 7.08 (d, 1H), 6.83 (dd, 1H), 6.69 (dd, 1H), 6.37 (m, 1H), 6.21 (d, 1H), 4.30 (br s, 1H), 3.84 (dd, 2H), 3.70 (br s, 1H), 3.30 (m, 6H), 3.20, 2.95, 2.80 (all br s, total 6H), 1.86 (m, 1H), 1.60 (m, 2H), 1.25 (m, 2H).

EXAMPLE 88

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-2-(1H-indol-5-yloxy)-N-({4-[(3-morpholin-4-ylpropyl)amino]-3-nitrophenyl}sulfonyl)benzamide

[0565] The title compound was prepared by substituting EXAMPLE 87C for EXAMPLE 1E and EXAMPLE 2A for EXAMPLE 1F in EXAMPLE 1G, except here the crude was purified by preparative HPLC using a 250 x 50 mm C18 column and eluting with 20-100% CH₃CN vs. 0.1% trifluoroacetic acid in water, giving the product as a trifluoroacetate salt. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.40 (br s, 1H), 11.19 (s, 1H), 9.60 (v br s, 1H), 8.69 (t, 1H), 8.60 (d, 1H), 7.83 (dd, 1H), 7.65 (br s, 1H), 7.50 (m, 5H), 7.38 (m, 5H), 7.12 (m, 2H), 6.83 (dd, 1H), 6.69 (dd, 1H), 6.39 (m, 1H), 6.20 (d, 1H), 4.38 (br s, 1H), 4.00 (m, 2H), 3.80 (br s, 1H), 3.40 (m, 4H), 3.30-2.80 (envelope, 10H), 3.20 (m, 4H), 1.96 (m, 2H).

EXAMPLE 89

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-indol-5-yloxy)-N-({3-nitro-4-([1-tetrahydro-2H-pyran-4-yl]piperidin-4-yl)amino]phenyl sulfonyl)benzamide

[0566] This EXAMPLE was prepared by substituting EXAMPLE 87C for EXAMPLE 1E and EXAMPLE 3M for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.15 (s, 1H), 8.56 (d, 1H), 8.20 (d, 1H), 7.84 (dd, 1H), 7.52 (d, 1H), 7.39 - 7.31 (m, 4H), 7.12 (d, 2H), 7.04 (d, 2H), 6.84 (dd, 1H), 6.65 (dd, 1H), 6.38 (t, 1H), 6.14 (d, 1H), 3.94 (m, 2H), 3.84 (m, 1H), 3.02 (m, 8H), 2.79 (m, 3H), 2.72 (s, 2H), 2.20-2.02 (m, 8H), 1.85 (m, 6H), 1.60 (m, 2H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 90

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-indol-5-yloxy)-N-({4-([1-methylpiperidin-4-yl]amino)-3-nitrophenyl} sulfonyl)benzamide

[0567] The title compound was prepared by substituting EXAMPLE 87C for EXAMPLE 1E and EXAMPLE 4A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.08 (s, 1H), 8.51 (d, 1H), 8.13 (d, 1H), 7.78 (dd, 1H), 7.52 (d, 1H), 7.37 - 7.31 (m, 4H), 7.06 - 7.00 (m, 4H), 6.79 (dd, 1H), 6.59 (dd, 1H), 6.35 (t, 1H), 6.14 (d, 1H), 3.73 (m, 1H), 3.05 - 2.95 (m, 6H), 2.71 (s, 2H), 2.60 (m, 2H), 2.48 (s, 3H), 2.16 (m, 6H), 2.01 (m, 2H), 1.95 (s, 2H), 1.70 (m, 2H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 91

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-indol-5-yloxy)-N-({4-([4-methylpiperazin-1-yl]amino)-3-nitrophenyl} sulfonyl)benzamide

[0568] The title compound was prepared by substituting EXAMPLE 6A for EXAMPLE 11B and EXAMPLE 87C for EXAMPLE 3J in EXAMPLE 11D. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.14 (s, 1H), 9.18 (s, 1H), 8.53 (d, 1H), 7.84 (dd, 1H), 7.56 (d, 1H), 7.51 (d, 1H), 7.39 (m, 2H), 7.33 (d, 2H), 7.12 (d, 1H), 7.03 (d, 2H), 6.84 (dd, 1H), 6.62 (dd, 1H), 6.38 (m, 1H), 6.13 (d, 1H), 3.00 (m, 4H), 2.90 (m, 4H), 2.71 (s, 2H), 2.33 (s, 3H), 2.15 (m, 6H), 1.94 (s, 2H), 1.37 (t, 2H), 0.92 (s, 6H).

EXAMPLE 92

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl]sulfonyl)-2-(1H-indol-5-yloxy)benzamide

[0569] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 87C and EXAMPLE 12A in place of EXAMPLE 3J and EXAMPLE 11B, respectively. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.16 (s, 2H), 8.39 (d, 1H), 8.06 (dd, 1H), 7.51 (d, 1H), 7.38 - 7.43 (m, 3H), 7.34 (d, 2H), 7.15 (d, 1H), 7.04 (d, 2H), 6.85 (dd, 1H), 6.64 (dd, 1H), 6.39 (s, 1H), 6.15 (d, 1H), 4.20 - 4.28 (m, 2H), 3.85 - 3.91 (m, 1H), 3.82 (dd, 1H), 3.74 - 3.78 (m, 1H), 3.59 - 3.69 (m, 2H), 3.40 - 3.51 (m, 2H), 3.05 (s, 4H), 2.78 (s, 2H), 2.23 (s, 4H), 2.14 (s, 2H), 1.95 (s, 2H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 93

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-indol-5-yloxy)-N-({4-([2-methoxyethyl]amino)-3-[(trifluoromethyl)sulfonyl]phenyl} sulfonyl)benzamide

[0570] The title compound was prepared by substituting EXAMPLE 87C for EXAMPLE 3J and EXAMPLE 16A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.17 (s, 1H), 8.18 (d, 1H), 7.92 (dd, 1H), 7.49 (d, 1H), 7.40 (m, 2H), 7.33 (d, 2H), 7.26 (m, 1H), 7.17 (d, 1H), 7.04 (m, 3H), 6.86 (dd, 1H), 6.65 (dd, 1H), 6.40 (s, 1H), 6.14 (d, 1H), 3.51 (m, 4H), 3.28 (s, 3H), 3.03 (s, 4H), 2.74 (s, 2H), 2.16 (m, 6H), 1.95 (s, 2H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 94

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-indol-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)benzamide

5

[0571] The title compound was prepared by substituting EXAMPLE 87C for EXAMPLE 3J and EXAMPLE 17A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.20 (s, 1H), 8.19 (d, 1H), 7.90 (dd, 1H), 7.53 (d, 1H), 7.40 (m, 4H), 7.33 (t, 1H), 7.17 (d, 1H), 7.07 (m, 3H), 6.86 (dd, 1H), 6.70 (dd, 1H), 6.41 (s, 1H), 6.21 (d, 1H), 3.84 (dd, 2H), 3.59 (m, 2H), 3.25 (m, 6H), 3.00 (m, 2H), 2.74 (s, 2H), 2.54 (m, 2H), 2.18 (s, 2H), 2.01 (s, 2H), 1.83 (m, 1H), 1.54 (m, 2H), 1.45 (t, 2H), 1.23 (m, 2H), 0.94 (s, 6H).

10

EXAMPLE 95

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl}amino)-3-nitrophenyl}sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

15

EXAMPLE 95A

1-(1,3-difluoropropan-2-yl)piperidin-4-amine

20

[0572] Tert-butyl piperidin-4-ylcarbamate (0.212 g), 1,3-difluoropropan-2-one (0.149 g) and sodium triacetoxyborohydride (0.337 g) were stirred together in dichloroethane at room temperature. After stirring overnight the reaction was quenched with water (10 mL) and extracted into dichloromethane (2 x 20 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated. The residue was treated with hydrogen chloride (4.0M in dioxane, 1.323 ml) for 1 hour to give the title compound as the HCl salt after concentration.

25

EXAMPLE 95B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl}amino)-3-nitrophenyl}sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

30

[0573] EXAMPLE 95A (0.057 g) and EXAMPLE 53A (0.162 g) were suspended in dioxane (3 mL) and heated to 105°C overnight. The reaction was concentrated, loaded onto silica gel (GraceResolv 12 g) and eluted with a gradient of 0.5% to 4% methanol/dichloromethane. The product containing fractions were concentrated and loaded onto C18 (SF25-75g analogix column) and eluted using a gradient of 30% to 60% acetonitrile/water. The product was partitioned between dichloromethane (20 mL) and saturated aqueous NaHCO₃ solution (20 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated to give the title compound. ¹H NMR (300 MHz, CDCl₃) δ 10.10 (s, 1H), 8.88 (d, 2H), 8.45 (d, 1H), 8.20 (s, 1H), 8.18 - 8.09 (m, 1H), 7.95 (d, 1H), 7.68 (d, 1H), 7.44 (s, 1H), 7.23 - 7.19 (m, 1H), 6.91 (d, 3H), 6.53 (d, 2H), 5.98 (d, 1H), 4.64 (dd, 4H), 3.68 - 3.50 (m, 1H), 3.01 (d, 6H), 2.72 (d, 4H), 2.19 (s, 11H), 1.69 (s, 2H), 1.41 (s, 2H), 0.94 (s, 6H).

40

EXAMPLE 96

N-({5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

45

EXAMPLE 96A

5-chloro-6-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)pyridine-3-sulfonamide

50

[0574] The title compound was prepared by substituting EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide and EXAMPLE 37C for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

55

EXAMPLE 96B

N-({5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-4-({2-[(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0575] The title compound was prepared by substituting EXAMPLE 96A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 8.52 (s, 1H), 8.41 (s, 1H), 8.03 (d, 1H), 7.56 (d, 1H), 7.50 (m, 2H), 7.35 (d, 2H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.39 (m, 1H), 6.22 (s, 1H), 4.50 (d, 2H), 3.78 (m, 2H), 3.60 (m, 2H), 3.12 (v br s, 4H), 2.93 (v br s, 2H), 2.38 (v br s, 4H), 2.17 (br m, 2H), 1.96 (s, 2H), 1.86 (m, 4H), 1.40 (t, 2H), 0.93 (s, 6H).

EXAMPLE 97

4-({2-[(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{{1-(2,2-difluoroethyl)piperidin-4-yl}amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 97A

tert-butyl 4-(2-nitro-4-sulfamoylphenylamino)piperidine-1-carboxylate

[0576] The title compound was prepared by substituting tert-butyl 4-aminopiperidine-1-carboxylate for 4-methylpiperazin-1-amine dihydrochloride in EXAMPLE 6A.

EXAMPLE 97B

3-nitro-4-(piperidin-4-ylamino)benzenesulfonamide

[0577] Tert-butyl 4-(2-nitro-4-sulfamoylphenylamino)piperidine-1-carboxylate was dissolved in dichloromethane (3 mL) and treated with 1N HCl in ether (4 mL). The reaction was stirred overnight then concentrated to give the title compound.

EXAMPLE 97C

4-(1-(2,2-difluoroethyl)piperidin-4-ylamino)-3-nitrobenzenesulfonamide

[0578] 3-nitro-4-(piperidin-4-ylamino)benzenesulfonamide hydrochloride (0.100g), 1,1-difluoro-2-iodoethane (0.063 mL) and diisopropylamine (0.156 mL) were stirred together in N,N-dimethylformamide (3 mL) and heated to 85°C. The reaction was diluted with dichloromethane (50 mL) and washed with water (50 mL), brine (50 mL), dried over magnesium sulfate, filtered, and concentrated. The residue was loaded onto silica gel (GraceResolve 12g) and eluted using a gradient of 0.5% methanol/dichloromethane to 3% methanol/dichloromethane over 30 minutes to give the title compound.

EXAMPLE 97D

4-({2-[(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{{1-(2,2-difluoroethyl)piperidin-4-yl}amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0579] The title compound was prepared by substituting EXAMPLE 97B for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 11.54 - 11.27 (m, 1H), 8.55 (d, 1H), 8.24 (d, 1H), 8.03 (d, 1H), 7.81 (d, 1H), 7.50 (dd, 3H), 7.34 (d, 2H), 7.13 (d, 1H), 7.04 (d, 2H), 6.68 (d, 1H), 6.38 (dd, 1H), 6.15 (dt, 2H), 3.64 (s, 1H), 3.07 (s, 4H), 2.79 (ddd, 6H), 2.41 (t, 2H), 2.17 (d, 6H), 1.92 (d, 4H), 1.61 (d, 2H), 1.38 (s, 2H), 0.92 (s, 6H).

EXAMPLE 98

4-({2-[(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1-cyclopropylpiperidin-4-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0580] The title compound was prepared as described in EXAMPLE 53B by replacing 1-acetyl piperidin-4-amine with 4-amino-1-cyclopropylpiperidine. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1H), 8.54 (d, 1H), 8.22 (d, 1H),

8.02 (d, 1H), 7.80 (dd, 1H), 7.49 (m, 3H), 7.34 (d, 2H), 7.11 (d, 1H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.38 (dd, 1H), 6.19 (d, 1H), 3.69 (m, 1H), 3.06 (m, 4H), 2.92 (m, 2H), 2.74 (s, 2H), 2.23 (m, 7H), 1.93 (m, 5H), 1.77 (m, 1H), 1.55 (m, 3H), 1.38 (t, 2H), 0.92 (s, 6H), 0.43 (m, 4H).

5 EXAMPLE 99

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-morpholin-4-ylcyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 **[0581]** The title compound was prepared as described in EXAMPLE 53B by replacing 1-acetylpiperidin-4-amine with 1-(4-morpholino)cyclohexanemethylamine. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.70 (s, 1H), 9.06 (s, 1H), 8.59 (d, 1H), 8.06 (d, 1H), 7.83 (dd, 1H), 7.57 (d, 1H), 7.50 (m, 2H), 7.34 (m, 3H), 7.19 (d, 1H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.41 (dd, 1H), 6.17 (d, 1H), 3.56 (m, 6H), 3.44 (m, 2H), 3.07 (m, 5H), 2.57 (m, 5H), 2.24 (m, 6H), 1.95 (s, 3H), 1.45 (m, 6H), 1.23 (m, 3H), 0.92 (s, 6H).

15

EXAMPLE 100

Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-(dicyclopropylamino)cyclohexyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

20

EXAMPLE 100A

Trans-tert-butyl-4-(dicyclopropylamino)cyclohexylcarbamate

25 **[0582]** A suspension of trans-tert-butyl-4-aminocyclohexylcarbamate (1 g), molecular sieves 3A (1 g), acetic acid (2.67 ml), (1-ethoxycyclopropoxy)trimethylsilane (3.74 ml) and sodium cyanoborohydride (0.880 g) in dry methanol (10 ml) was heated at reflux for 3 hours. The insolubles were filtered off, the resulting solution was basified with aqueous NaOH (6 M) to pH 14, and extracted with ether. The combined extracts were washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue was purified by flash chromatography (silica gel 80 g, 30 - 100% acetone/hexanes) to provide the title compound.

30

EXAMPLE 100B

(trans)-N¹,N¹-dicyclopropylcyclohexane-1,4-diamine bis(2,2,2-trifluoroacetate)

35

[0583] The title compound was prepared by substituting EXAMPLE 100A for EXAMPLE 39A in EXAMPLE 39B.

EXAMPLE 100C

40 Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-(dicyclopropylamino)cyclohexyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0584] A suspension of EXAMPLE 53A (0.14 g), EXAMPLE 100B (0.112 g) and N,N-diisopropylethylamine (0.310 mL) in dioxane (10 mL) was stirred for 3 days at 100 °C. The product was concentrated and purified by RP HPLC(C₈, 30% - 100% CH₃CN/water/0.1% trifluoroacetic acid). ¹H NMR (500 MHz, pyridine-d₅) δ 13.07 (s, 1H), 9.28 (d, 1H), 8.41 - 8.45 (m, 2H), 8.37 (d, 1H), 8.12 (d, 1H), 7.67 (d, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 7.01 (d, 1H), 6.75 (dd, 1H), 6.53 (d, 1H), 6.48 - 6.51 (m, 1H), 3.43 (ddd, 1H), 3.03 - 3.09 (m, 4H), 2.72 - 2.79 (m, 3H), 2.22 - 2.28 (m, 2H), 2.11 - 2.16 (m, 4H), 2.10 (s, 2H), 2.00 - 2.05 (m, 2H), 1.97 (s, 2H), 1.89 (s, 1H), 1.86 (s, 3H), 1.62 - 1.71 (m, 2H), 1.39 (t, 2H), 1.19 - 1.29 (m, 2H), 0.93 (s, 6H), 0.48 (d, 8H).

50

55

EXAMPLE 101

4-(4-{[2-(4-chlorophenyl)-6,6-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 101A

Ethyl 2-hydroxy-6,6-dimethylcyclohex-1-enecarboxylate

[0585] Into a 500 mL flame dried round-bottomed flask was added copper(I) iodide (18 g) in ether (200 mL) to give a suspension. After cooling to -5°C, methyllithium (120 mL, 1.6M in ether) was added dropwise. After stirring at -5°C for 1 hour, 3-methylcyclohex-2-enone (5.15 mL) in 15 mL ether was added dropwise, and the mixture was stirred at -5°C for 1 hour. After cooling to -78°C, hexamethylphosphoramide (60 mL) was added dropwise. Ethyl carbonocyanidate (23.74 mL) was added. After stirring at -78 °C for 20 minutes, the mixture was warmed up to room temperature, and stirred for 1 hour. The mixture was poured into cold water, and the layers were separated. The aqueous layer was extracted with ether (3x 20 mL). The combined organic layers were washed with saturated aqueous NH₄Cl (3x 20 mL), dried over Na₂SO₄, filtered, and dried under vacuum. The crude product was added to a silica gel column and purified by eluting with 0-10% ethyl acetate in hexane.

EXAMPLE 101B

Ethyl 6,6-dimethyl-2-(trifluoromethylsulfonyloxy)cyclohex-1-enecarboxylate

[0586] Into a 500 mL round-bottomed flask was added hexane-washed sodium hydride (0.5 g) in dichloromethane (100 mL) to give a suspension. After cooling to -5 °C, EXAMPLE 101A (2.0 g) was added. After stirring at -5 °C for 30 minutes, the mixture was cooled to -78°C. Trifluoromethanesulfonic anhydride (2.2 mL) was added. The mixture was warmed to room temperature and stirred overnight. Water was added slowly to the mixture, the aqueous layer was then extracted by dichloromethane (2x 20 mL). The combined organic layers were washed with saturated NH₄Cl and brine, dried over Na₂SO₄, filtered, and concentrated.

EXAMPLE 101C

ethyl 2-(4-chlorophenyl)-6,6-dimethylcyclohex-1-enecarboxylate

[0587] Into a 25 mL microwave tube was added EXAMPLE 101B (2.9 g), 4-chlorophenylboronic acid (2.2 g), and tetrakis(triphenylphosphine)palladium (0.05 g) in 1,2-dimethoxyethane/methanol(2:1, 10 mL) to give a solution. Cesium fluoride (4 g) was then added. The reaction mixture was stirred at 150°C under (100W) in a Biotage Initiator microwave reactor for 30 minutes. After removing the solvents, water was added, and the mixture was extracted with ethyl acetate (2x). The combined organic layers were dried by MgSO₄. After filtering, the crude product was purified by reverse phase chromatography eluting with 50-100% acetonitrile/water with 0.1% trifluoroacetic acid.

EXAMPLE 101D

(2-(4-chlorophenyl)-6,6-dimethylcyclohex-1-enyl)methanol

[0588] In a 100 mL round-bottomed flask was placed lithium aluminum hydride (1 g) in ether (20 mL) to give a suspension. EXAMPLE 101C (1 g) dissolved in ether (5 mL) was added slowly by syringe. The mixture was stirred at room temperature overnight. After cooling to 0°C, the reaction was quenched by water. Ether (2x 10 mL) was used to extract the product. The crude product was purified by flash chromatography on silica by eluting with 0-15% ethyl acetate in hexane.

EXAMPLE 101E

Methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-6,6-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoate

[0589] To a 0°C solution of EXAMPLE 101D (0.43 g) in dichloromethane (5 mL) was added triethylamine (1 mL). Methanesulfonyl chloride (0.134 mL) was then added slowly. After 5 minutes, EXAMPLE 15F (0.61 g) was added. The

mixture was stirred at room temperature overnight. The crude product was purified by flash chromatography on silica with 0 to 25% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 101F

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-6,6-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoic acid

[0590] In a 5 mL microwave tube was added lithium hydroxide hydrate (15 mg) and EXAMPLE 101E (45 mg) in dioxane/water(2:1) (2 mL) to give a suspension. The mixture was heated to 130°C in a Biotage Initiator microwave reactor for 20 minutes. After cooling and neutralization by HCl, the crude product was added to a Prep HPLC column and was eluted with 20-80% acetonitrile/water with 0.1% trifluoroacetic acid.

EXAMPLE 101G

4-(4-([2-(4-chlorophenyl)-6,6-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0591] The title compound was prepared by substituting EXAMPLE 101F for EXAMPLE 3J and EXAMPLE 1F for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 11.47 (s, 1H), 8.58 (m, 2H), 8.03 (m, 1H), 7.79 (m, 1H), 7.51 (m, 3H), 7.31 (d, 2H), 7.10 (m, 1H), 7.02 (d, 2H), 6.65 (m, 1H), 6.39 (m, 1H), 6.15 (m, 1H), 3.85 (m, 2H), 3.27 (m, 4H), 2.97 (m, 4H), 2.76 (s, 2H), 2.14 (m, 6H), 1.70 (m, 2H), 1.61 (m, 2H), 1.44 (m, 2H), 1.26 (m, 3H), 1.16 (m, 6H)

EXAMPLE 102

N-({5-bromo-6-[(4-ethylmorpholin-3-yl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 102A

(4-ethylmorpholin-3-yl)methanol

[0592] Morpholin-3-ylmethanol (500 mg) and iodoethane (666 mg) in N,N-dimethylformamide was treated with K₂CO₃ (1.1 g) overnight. The reaction mixture was diluted with water and extracted with ethyl acetate. The organic layer was dried over Na₂SO₄, filtered, and concentrated to provide the title compound.

EXAMPLE 102B

5-bromo-6-((4-ethylmorpholin-3-yl)methoxy)pyridine-3-sulfonamide

[0593] The title compound was prepared as described in EXAMPLE 12A by replacing 4-fluoro-3-nitrobenzenesulfonamide and (1,4-dioxan-2-yl)methanol with 5-bromo-6-fluoropyridine-3-sulfonamide and EXAMPLE 102A, respectively.

EXAMPLE 102C

N-({5-bromo-6-[(4-ethylmorpholin-3-yl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0594] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 102B in place of EXAMPLE 11B. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.62 (s, 1H), 8.51 (s, 1H), 8.30 (s, 1H), 8.00 (d, 1H), 7.55 (d, 1H), 7.45 - 7.50 (m, 2H), 7.35 (d, 2H), 7.05 (d, 2H), 6.66 (dd, 1H), 6.37 (s, 1H), 6.21 (d, 1H), 4.58 (dd, 1H), 4.39 - 4.50 (m, 1H), 3.78 - 3.90 (m, 1H), 3.67 - 3.77 (m, 1H), 3.50 - 3.65 (m, 2H), 3.08 (s, 4H), 2.59 - 3.00 (m, 4H), 2.20 - 2.39 (m, 2H), 2.15 (s, 2H), 1.96 (s, 2H), 1.39 (t, 2H), 0.99 - 1.11 (m, 3H), 0.93 (s, 6H)

EXAMPLE 103

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-ethylmorpholin-3-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 103A

4-((4-ethylmorpholin-3-yl)methoxy)-3-nitrobenzenesulfonamide

[0595] The title compound was prepared as described in EXAMPLE 12A by replacing (1,4-dioxan-2-yl)methanol with EXAMPLE 102A.

EXAMPLE 103B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-ethylmorpholin-3-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0596] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 103A in place of EXAMPLE 11B. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1H), 8.33 (s, 1H), 7.99 - 8.06 (m, 2H), 7.47 - 7.57 (m, 3H), 7.45 (d, 1H), 7.35 (d, 2H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.38 (dd, 1H), 6.20 (d, 1H), 4.42 (dd, 1H), 4.23 (dd, 1H), 3.81 (d, 1H), 3.69 (d, 1H), 3.49 - 3.63 (m, 2H), 3.08 (s, 4H), 2.92 (s, 1H), 2.81 (s, 4H), 2.54 (s, 1H), 2.25 (s, 4H), 2.15 (s, 2H), 1.96 (s, 2H), 1.39 (t, 2H), 1.00 (t, 3H), 0.92 (s, 6H)

EXAMPLE 104

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(4-tetrahydro-2H-pyran-4-ylmorpholin-3-yl)methoxy]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0597] EXAMPLE 78 (20 mg) and dihydro-2H-pyran-4(3H)-one (10 mg) in dichloroethane (2 mL) was treated with NaCNBH₃ (9.74 mg) overnight. Additional dihydro-2H-pyran-4(3H)-one (20 mg) and titanium (IV) isopropoxide (0.05 mL) were added. The resulting mixture was stirred at room temperature overnight and concentrated. The residue was purified by reverse phase HPLC on a C18 column using a gradient of 35-60% acetonitrile in 0.1% trifluoroacetic acid water to give the title compound as a trifluoroacetate salt. The trifluoroacetic acid salt was dissolved in dichloromethane (6 mL) and washed with 50% aqueous NaHCO₃. The organic layer was dried over anhydrous Na₂SO₄ and concentrated to give the title compound. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 8.35 (s, 1H), 8.04 (s, 2H), 7.44 - 7.58 (m, 4H), 7.35 (d, 2H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.40 (s, 1H), 6.20 (s, 1H), 4.44 (s, 1H), 4.28 (s, 1H), 3.85 (d, 2H), 3.71 (d, 1H), 3.61 (s, 3H), 3.20 - 3.29 (m, 2H), 3.08 (s, 5H), 2.54 - 2.96 (m, 5H), 2.06 - 2.42 (m, 5H), 1.96 (s, 2H), 1.77 (d, 1H), 1.53 - 1.66 (m, 1H), 1.29 - 1.51 (m, 4H), 0.92 (s, 6H)

EXAMPLE 105

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-[(3S)-1-tetrahydro-2H-pyran-4-ylpiperidin-3-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 105A

(S)-tert-butyl 1-(tetrahydro-2H-pyran-4-yl)piperidin-3-ylcarbamate

[0598] The title compound was prepared by substituting (S)-tert-butyl piperidin-3-ylcarbamate for tert-butyl piperazine-1-carboxylate and dihydro-2H-pyran-4(3H)-one for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 105B

(S)-1-(tetrahydro-2H-pyran-4-yl)piperidin-3-amine

[0599] The title compound was prepared by substituting EXAMPLE 105A for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 105C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-[(3S)-1-tetrahydro-2H-pyran-4-yl]piperidin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0600] The title compound was prepared by substituting EXAMPLE 105B for 1-acetylpiperidin-4-amine in EXAMPLE 53B. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 8.68 (br s, 1H), 8.54 (br s, 1H), 8.02 (d, 1H), 7.77 (m, 1H), 7.50 (m, 3H), 7.34 (d, 2H), 7.03 (m, 3H), 6.67 (dd, 1H), 6.38 (m, 1H), 6.19 (d, 1H), 3.98 (m, 2H), 3.90 (m, 2H), 3.52 (m, 2H), 3.09 (s, 2H), 3.05 (m, 4H), 2.77 (m, 2H), 2.60 (m, 2H), 2.16 (m, 6H), 1.95 (m, 2H), 1.65 (m, 5H), 1.50 (m, 3H), 1.38 (m, 2H), 0.94 (s, 6H).

EXAMPLE 106

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({5-cyano-6-[(tetrahydro-2H-pyran-4-yl)methyl]amino}pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 106 A

5-Bromo-6-((tetrahydro-2H-pyran-4-yl)methylamino)pyridine-3-sulfonamide

[0601] The title compound was prepared by substituting (tetrahydro-2H-pyran-4-yl)methanamine for EXAMPLE 3L in EXAMPLE 61A.

EXAMPLE 106B

5-cyano-6-((tetrahydro-2H-pyran-4-yl)methylamino)pyridine-3-sulfonamide

[0602] The title compound was prepared by substituting EXAMPLE 106A for EXAMPLE 36B in EXAMPLE 36C.

EXAMPLE 106C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({5-cyano-6-[(tetrahydro-2H-pyran-4-yl)methyl]amino}pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0603] The title compound was prepared by substituting EXAMPLE 106B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.62 (s, 1H), 8.55 (s, 1H), 8.14 (s, 1H), 8.01 (d, 1H), 7.87 (s, 1H), 7.56 (d, 1H), 7.48 (d, 2H), 7.34 (d, 2H), 7.04 (d, 2H), 6.64 (m, 1H), 6.37 (s, 1H), 6.19 (d, 1H), 3.81 (dd, 2H), 3.25 (m, 4H), 3.04 (s, 4H), 2.74 (s, 2H), 2.17 (m, 6H), 1.95 (s, 2H), 1.87 (m, 1H), 1.53 (m, 2H), 1.37 (t, 2H), 1.18 (m, 2H), 0.91 (s, 6H).

EXAMPLE 107

Trans-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1,1-dioxidothiomorpholin-4-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 107A

3-nitro-4-(4-aminothiomorpholine-1,1-dioxide)benzenesulfonamide

[0604] The title compound was prepared by substituting 4-aminothiomorpholine-1,1-dioxide for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 107B

Trans-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1,1-dioxidothiomorpholin-4-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0605] The title compound was prepared by substituting EXAMPLE 107A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.64 (s, 1H), 9.58 (s, 1H), 8.50 (s, 1H), 8.02 (d, 1H), 7.78 (m, 2H), 7.50 (m,

3H), 7.34 (d, 2H), 7.04 (d, 2H), 6.66 (dd, 1H), 6.38 (s, 1H), 6.19 (d, 1H), 3.48 (m, 4H), 3.23 (m, 4H), 3.05 (s, 4H), 2.73 (d, 2H), 2.16 (m, 6H), 1.95 (s, 2H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 108

N-[[4-[[[4-aminotetrahydro-2H-pyran-4-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 108A

4-((4-aminotetrahydro-2H-pyran-4-yl)methylamino)-3-nitrobenzenesulfonamide

[0606] The title compound was prepared by substituting 4-(aminomethyl)tetrahydro-2H-pyran-4-amine for (tetrahydro-2H-pyran-4-yl)methanamine in EXAMPLE 1F.

EXAMPLE 108B

N-[[4-[[[4-aminotetrahydro-2H-pyran-4-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0607] The title compound was prepared by substituting EXAMPLE 108A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.55 (s, 1H), 8.45 (s, 2H), 7.95 (d, 1H), 7.75-7.77 (m, 1H), 7.57 (d, 2H), 7.44 (s, 1H), 7.34 (d, 2H), 7.09 (d, J = 8.85 Hz, 1H), 7.05 (d, 2H), 6.69 (dd, 1H), 6.33 (d, 1H), 6.22 (d, 1H), 3.59-3.71 (m, 6H), 3.01 (s, 4H), 2.73 (s, 2H), 2.15-2.19 (m, 6H), 1.95 (s, 2H), 1.71-1.74 (m, 2H), 1.59-1.61 (m, 1H), 1.38 (t, 2H), 0.93 (s, 6H).

EXAMPLE 109

Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({5-cyano-6-[(4-morpholin-4-yl)cyclohexyl]amino}pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 109A

Trans-5-bromo-6-(4-morpholinocyclohexyloxy)pyridine-3-sulfonamide

[0608] The title compound was prepared by substituting EXAMPLE 9B for EXAMPLE 3L in EXAMPLE 61 A.

EXAMPLE 109B

Trans-5-cyano-6-(4-morpholinocyclohexylamino)pyridine-3-sulfonamide

[0609] The title compound was prepared by substituting EXAMPLE 109A for EXAMPLE 36B in EXAMPLE 36C.

EXAMPLE 109C

Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({5-cyano-6-[(4-morpholin-4-yl)cyclohexyl]amino}pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0610] The title compound was prepared by substituting EXAMPLE 109B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.59 (s, 1H), 8.56 (d, 1H), 8.13 (s, 1H), 8.00 (d, 1H), 7.55 (d, 1H), 7.47 (m, 2H), 7.34 (d, 2H), 7.04 (d, 2H), 6.64 (dd, 1H), 6.36 (d, 1H), 6.19 (d, 1H), 4.00 (m, 1H), 3.65 (m, 4H), 3.28 (m, 4H), 3.03 (m, 4H), 2.73 (m, 4H), 2.16 (m, 6H), 1.90 (m, 6H), 1.40 (m, 6H), 0.93 (s, 6H).

EXAMPLE 110

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-cyano-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0611] The title compound was prepared by substituting EXAMPLE 52B for EXAMPLE 11B in EXAMPLE 11D. ¹H

EP 2 944 638 A1

NMR (300 MHz, dimethylsulfoxide- d_6) δ 11.71 (s, 1H), 11.23 (s, 1H), 8.08 (d, 1H), 7.91 (d, 1H), 7.74 (dd, 1H), 7.60 (d, 1H), 7.52 (m, 2H), 7.34 (m, 2H), 7.16 (s, 1H), 7.04 (m, 2H), 6.83 (d, 1H), 6.68 (dd, 1H), 6.43 (dd, 1H), 6.16 (d, 1H), 3.83 (dd, 2H), 3.23 (m, 2H), 3.12 (t, 2H), 3.06 (m, 4H), 2.73 (m, 2H), 2.15 (m, 6H), 1.95 (s, 2H), 1.82 (m, 1H), 1.58 (m, 2H), 1.38 (m, 2H), 1.18 (m, 2H), 0.92 (s, 6H).

EXAMPLE 111

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([(1S,3R)-3-morpholin-4-ylcyclopentyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 111A

benzyl (1S,3R)-3-(tert-butoxycarbonylamino)cyclopentylcarbamate

[0612] (1S,3R)-3-(tert-butoxycarbonylamino)cyclopentanecarboxylic acid (1.03 g), diphenylphosphoryl azide (DPPA, 1.00 mL), triethylamine (0.929 mL), and benzyl alcohol (0.931 mL) were combined in toluene (10 mL) and stirred at 100°C for 24 hours. The solution was cooled and chromatographed on silica gel using 10% ethyl acetate/ hexanes to give the pure product.

EXAMPLE 111B

benzyl (1S,3R)-3-aminocyclopentylcarbamate

[0613] The title compound was prepared by substituting EXAMPLE 111A for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 111C

benzyl (1S,3R)-3-morpholinocyclopentylcarbamate

[0614] A solution of EXAMPLE 111B (400 mg), 1-bromo-2-(2-bromoethoxy)ethane (0.246 mL), and triethylamine (0.595 mL) in N,N-dimethylformamide (6 mL) was stirred at 70°C for 24 hours. The solution was cooled and poured into ethyl acetate (200 mL). The solution was extracted with 3x water, washed with brine, concentrated, and chromatographed on silica gel using 10% methanol/ethyl acetate to give the pure product.

EXAMPLE 111D

(1S,3R)-3-morpholinocyclopentanamine

[0615] EXAMPLE 111C (300 mg) and ethanol (20 mL) were added to wet 20% Pd(OH)₂-C (60.0 mg) in a 50 mL pressure bottle and stirred for 8 hours at 30 psi. The mixture was filtered through a nylon membrane and condensed to give the product.

EXAMPLE 111E

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([(1S,3R)-3-morpholin-4-ylcyclopentyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0616] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 111D for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (dimethylsulfoxide- d_6) δ 11.65 (s, 1H), 8.45 (d, 1H), 8.28 (dd, 1H), 7.97 (d, 1H), 7.68 (d, 1H), 7.52 (d, 1H), 7.44 (d, 2H), 7.35 (d, 2H), 7.05 (d, 2H), 6.92 (dd, 1H), 6.85 (dd, 1H), 6.33 (s, 1H), 6.22 (s, 1H), 4.08 (m, 1H), 3.60 (br s, 4H), 3.06 (br s, 4H), 2.73 (br s, 3H), 2.48 (m, 4H), 2.28 (m, 1H), 2.18 (m, 6H), 2.07 (m, 1H), 1.95 (s, 2H), 1.79 (m, 2H), 1.63 (m, 2H), 1.38 (t, 2H), 0.93 (s, 6H).

EXAMPLE 112

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([(1R,3S)-3-morpholin-4-ylcyclopentyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 112A

tert-butyl (1R,3S)-3-aminocyclopentylcarbamate

[0617] The title compound was prepared by substituting EXAMPLE 111A for EXAMPLE 111C in EXAMPLE 111D.

EXAMPLE 112B

tert-butyl (1R,3S)-3-morpholinocyclopentylcarbamate

[0618] The title compound was prepared by substituting EXAMPLE 112A for EXAMPLE 111B in EXAMPLE 111C.

EXAMPLE 112C

(1R,3S)-3-morpholinocyclopentanamine

[0619] The title compound was prepared by substituting EXAMPLE 112B for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 112D

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([(1R,3S)-3-morpholin-4-ylcyclopentyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0620] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 112C for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (dimethylsulfoxide-d₆) δ 11.35 (s, 1H), 8.51 (d, 1H), 8.44 (dd, 1H), 8.00 (d, 1H), 7.77 (d, 1H), 7.50 (d, 1H), 7.48 (s, 2H), 7.34 (d, 2H), 7.04 (d, 2H), 7.02 (dd, 1H), 6.67 (dd, 1H), 6.37 (dd, 1H), 6.21 (d, 1H), 4.11 (m, 1H), 3.61 (br s, 4H), 3.06 (br s, 4H), 2.73 (br s, 3H), 2.50 (m, 4H), 2.28 (m, 1H), 2.18 (m, 6H), 2.06 (m, 1H), 1.95 (s, 2H), 1.77 (m, 2H), 1.66 (m, 2H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 113

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([(morpholin-2-ylmethyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 113A

tert-butyl 2-((2-nitro-4-sulfamoylphenylamino)methyl)morpholine-4-carboxylate

[0621] The title compound was prepared by substituting tert-butyl 2-(aminomethyl)morpholine-4-carboxylate for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 113B

tert-butyl 2-((4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)sulfamoyl)-2-nitrophenylamino)methyl)morpholine-4-carboxylate

[0622] The title compound was prepared by substituting EXAMPLE 113A for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G, with the exception that the product was purified on a silica gel column eluted with 4% methanol in dichloromethane.

EXAMPLE 113C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(morpholin-2-ylmethyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

5

[0623] The title compound was prepared by substituting EXAMPLE 113B for EXAMPLE 66B in EXAMPLE 66C. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.60 (s, 1H), 8.55 (br, s, 1H), 8.51 (s, 1H), 8.00 (d, 1H), 7.80 (d, 1H), 7.52 (d, 1H), 7.49-7.46 (m, 2H), 7.34 (d, 2H), 7.07 (d, 1H), 7.04(d, 2H), 6.66 (dd, 1H), 6.36 (s, 1H), 6.20 (d, 1H), 4.00 (dd, 1H), 3.91 (m, 1H), 3.70 (t, 1H), 3.60 (m, 1H), 3.58 (m, 1H), 3.32 (m, 1H), 3.16 (d, 1H), 3.05 (m, 4H), 2.98 (td, 1H), 2.86 (t, 1H), 2.73 (s, 2H), 2.20-2.12 (m, 6H), 1.95 (s, 2H), 1.38 (t, 2H), 0.92 (s, 6H).

10

EXAMPLE 114

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydrofuran-3-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

15

EXAMPLE 114A

3-nitro-4-[(tetrahydrofuran-3-yl)methylamino]benzenesulfonamide

20

[0624] The title compound was prepared by substituting 3-aminomethyl-tetrahydrofuran for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 114B

25

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydrofuran-3-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

30

[0625] The title compound was prepared by substituting EXAMPLE 114A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 11.42 (bs, 1H), 8.63 (t, 1H), 8.56 (d, 1H), 8.04 (d, 1H), 7.80 (dd, 1H), 7.53-7.48 (m, 3H), 7.34 (d, 2H), 7.10 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.39 (m, 1H), 6.19 (d, 1H), 3.82-3.79 (m, 1H), 3.71 (t, 1H), 3.62 (dd, 1H), 3.50 (dd, 1H), 3.38 (m, 1H), 3.32 (m, 1H), 3.07 (m, 4H), 2.76 (s, 2H), 2.58 (m, 1H), 2.25-2.00 (m, 6H), 1.98 (m, 1H), 1.95 (s, 2H), 1.65 (m, 1H), 1.38 (t, 2H), 0.92 (s, 6H).

35

EXAMPLE 115

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-({1-[cis-3-fluorotetrahydro-2H-pyran-4-yl]piperidin-4-yl}amino)-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

40

EXAMPLE 115A

Cis-tert-butyl 1-(3-fluorotetrahydro-2H-pyran-4-yl)piperidin-4-ylcarbamate

45

[0626] The title compound was prepared as a racemate of the cis diastereomer by substituting tert-butyl piperidin-4-ylcarbamate for piperidin-4-ol and 3-fluorodihydro-2H-pyran-4(3H)-one (prepared by the method described in US2005/0101628A1) for dihydro-2H-pyran-4(3H)-one in EXAMPLE 84A.

EXAMPLE 115B

50

[0627] Cis-1-(3-fluorotetrahydro-2H-pyran-4-yl)piperidin-4-amine EXAMPLE 115A (0.29 g) was dissolved in CH₂Cl₂ (9 mL), then 4N HCl in dioxane (4 mL) was added and the reaction stirred at room temperature for 16 hours. The reaction was diluted with CH₂Cl₂ (30 mL), then 4N aqueous NaOH (5 mL) was added. After shaking and separating the layers the aqueous layer was saturated with solid NaCl and extracted with more CH₂Cl₂ (10 mL). The combined organic layers were dried over Na₂SO₄. After filtration and concentration the amine was used with no further purification.

55

EXAMPLE 115C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({[4-({1-[cis-3-fluorotetrahydro-2H-pyran-4-yl]piperidin-4-yl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

5

[0628] The title compound was prepared by substituting EXAMPLE 115B for 1-acetylpiperidin-4-amine in EXAMPLE 53B. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.64 (s, 1H), 8.54 (d, 1H), 8.43 (br d, 1H), 8.03 (d, 1H), 7.80 (dd, 1H), 7.50 (m, 3H), 7.35 (d, 2H), 7.11 (d, 1H), 7.05 (d, 2H), 6.66 (dd, 1H), 6.38 (m, 1H), 6.20 (d, 1H), 4.92 (d, 1H), 3.95 (m, 2H), 3.70 (v br m, 1H), 3.50, 3.40, 3.30 (all m, total 5H), 3.05, 3.00 (both v br m, total 5H), 2.74 (s, 2H), 2.55 (v br m, 1H), 2.18 (br m, 6H), 1.95 (m, 4H), 1.88 (ddd, 1H), 1.63 (v br m, 3H), 1.38 (t, 2H), 0.92 (s, 6H).

10

EXAMPLE 116

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(1-tetrahydro-2H-pyran-4-yl)azetidin-3-yl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

15

EXAMPLE 116A

1-(tetrahydro-2H-pyran-4-yl)azetidin-3-amine

20

[0629] Tert-butyl azetidin-3-ylcarbamate (0.46 g), dihydro-2H-pyran-4(3H)-one (0.29 g) and sodium triacetoxymethylborohydride (0.85 g) were stirred together in dichloromethane (5 mL) overnight. The reaction was poured into dichloromethane (50 mL) and saturated aqueous NaHCO₃ solution (25 mL). The organic layer was separated, washed with brine (25 mL), dried over Na₂SO₄, filtered, and concentrated. Silica gel chromatography (GraceResolv 12 g) eluting with a gradient of 0.75% to 7.5% methanol/dichloromethane over 20 minutes gave the Boc-protected intermediate. Treatment with HCl (4.0M in dioxane, 2 mL) and methanol (1 mL) for 1 hour gave the title compound after concentration as the di-HCl salt.

25

EXAMPLE 116B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(1-tetrahydro-2H-pyran-4-yl)azetidin-3-yl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

30

[0630] A suspension of 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-(4-chloro-3-nitrophenyl)sulfonyl)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzamide (0.180 g), 1-(tetrahydro-2H-pyran-4-yl)azetidin-3-amine (0.078 g), and triethylamine (0.159 mL) in dioxane (2 mL) was degassed with nitrogen for 30 seconds then sealed. The reaction was heated to 110°C. After stirring for 16 hours, more triethylamine (10 equivalents total) and dimethylsulfoxide (1 mL) were added and the reaction stirred for an additional 18 hours at 110°C. The reaction was cooled, diluted with water (50 mL) and extracted with dichloromethane (2 x 150 mL). The organic layer was dried over magnesium sulfate, filtered and concentrated. Silica gel chromatography (GraceResolv 12 g) eluting with a gradient of 0.75% to 7.5% methanol/dichloromethane (Flow = 36 mL/minutes) gave the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.59 (s, 1H), 8.49 (d, 1H), 8.40 (s, 1H), 7.97 (d, 1H), 7.77 (s, 1H), 7.47 (dd, 3H), 7.34 (d, 2H), 7.04 (d, 2H), 6.90 - 6.78 (m, 1H), 6.65 (d, 1H), 6.35 (s, 1H), 6.21 (s, 1H), 4.47 - 4.23 (m, 1H), 3.83 (s, 3H), 3.05 (s, 6H), 2.73 (s, 2H), 2.18 (s, 8H), 1.95 (s, 2H), 1.68 (s, 2H), 1.38 (s, 2H), 1.24 (s, 4H), 0.92 (s, 6H).

35

40

EXAMPLE 117

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(1-tetrahydrofuran-3-yl)azetidin-3-yl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50

EXAMPLE 117A

1-(tetrahydrofuran-3-yl)azetidin-3-amine

[0631] Tert-butyl azetidin-3-ylcarbamate (0.550 g), dihydrofuran-3(2H)-one (0.412 g) and sodium triacetoxymethylborohydride (1.015 g) were stirred together in dichloromethane (5 mL). After stirring overnight, the reaction was poured into saturated aqueous NaHCO₃ solution (25 mL) and extracted with dichloromethane (50 mL). The organic layer was washed with brine (25 mL), dried over magnesium sulfate, filtered, and concentrated. Silica gel chromatography (GraceResolv 12 g) eluting with a gradient of 0.5% to 7.5% methanol/dichloromethane over 30 minutes gave tert-butyl 1-(tetrahydro-

55

furan-3-yl)azetidin-3-ylcarbamate. The resulting material was treated with HCl/dioxane for 1 hour, and then concentrated to give the title compound.

EXAMPLE 117B

3-nitro-4-(1-(tetrahydrofuran-3-yl)azetidin-3-ylamino)benzenesulfonamide 4-Fluoro-3-nitrobenzenesulfonamide (0.084 g), 1-(tetrahydrofuran-3-yl)azetidin-3-amine

[0632] (0.090 g) and triethylamine (0.266 ml) in tetrahydrofuran (3 mL) was heated to 60°C. After stirring for 4 hours, the reaction was cooled, the tetrahydrofuran was removed and the residue was partitioned between dichloromethane (200 mL) and water (20 mL). The organic layer was separated, washed with brine (25 mL), dried over magnesium sulfate, filtered, and concentrated to give the title compound.

EXAMPLE 117C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({[3-nitro-4-[(1-tetrahydrofuran-3-yl)azetidin-3-yl]amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0633] The title compound was prepared by substituting EXAMPLE 117B for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, CDCl₃) δ 10.39 - 9.79 (m, 1H), 9.17 (s, 1H), 8.87 (d, 1H), 8.51 (d, 1H), 8.15 (dd, 2H), 7.94 (d, 1H), 7.68 (d, 1H), 7.48 - 7.42 (m, 1H), 7.23 (d, 2H), 6.91 (d, 2H), 6.69 (d, 1H), 6.54 (dd, 2H), 5.99 (d, 1H), 4.29 (d, 1H), 4.01 - 3.73 (m, 4H), 3.66 (d, 2H), 3.08 (s, 6H), 2.76 (s, 2H), 2.21 (s, 6H), 2.03 - 1.83 (m, 3H), 1.64 (s, 2H), 1.42 (d, 2H), 0.93 (s, 6H).

EXAMPLE 118

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({[3-nitro-4-({[(3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]methyl}amino)phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 118A

(R)-tert-butyl (1-(tetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl)methylcarbamate

[0634] The title compound was prepared by substituting (S)-tert-butyl pyrrolidin-3-ylmethylcarbamate for tert-butyl piperazine-1-carboxylate and dihydro-2H-pyran-4(3H)-one for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 118B

(R)-(1-(tetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl)methanamine

[0635] The title compound was prepared by substituting EXAMPLE 118A for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 118C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({[3-nitro-4-({[(3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]methyl}amino)phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0636] The title compound was prepared by substituting EXAMPLE 118B for 1-acetylpiperidin-4-amine in EXAMPLE 53B. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.57 (s, 1H), 8.59 (br s, 1H), 8.45 (br s, 1H), 8.02 (d, 1H), 7.95 (m, 1H), 7.71 (m, 1H), 7.56 (d, 1H), 7.45 (m, 1H), 7.35 (m, 3H), 7.05 (m, 2H), 6.90 (br s, 1H), 6.64 (d, 1H), 6.33 (m, 1H), 6.22 (m, 1H), 3.90 (m, 2H), 3.44 (m, 2H), 3.27 (m, 4H), 3.02 (m, 5H), 2.73 (m, 3H), 2.59 (m, 2H), 2.19 (m, 6H), 1.95 (m, 2H), 1.85 (m, 2H), 1.64 (m, 1H), 1.50 (m, 2H), 1.39 (m, 2H), 1.23 (m, 1H), 0.94 (s, 6H).

EXAMPLE 119

4-(4-([2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

5

[0637] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 75F and EXAMPLE 37D in place of EXAMPLE 3J and EXAMPLE 11B, respectively. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 8.39 (s, 1H), 8.08 (d, 1H), 8.04 (d, 1H), 7.41 - 7.59 (m, 4H), 7.35 (d, 2H), 7.08 (d, 2H), 6.68 (dd, 1H), 6.37 - 6.43 (m, 1H), 6.20 (s, 1H), 4.38 (d, 2H), 3.73 - 3.82 (m, 2H), 3.54 - 3.63 (m, 2H), 3.09 (s, 4H), 2.81 (s, 2H), 2.16 - 2.39 (m, 5H), 1.94 (s, 2H), 1.79 - 1.93 (m, 4H), 1.40 (t, 2H), 0.94 (s, 6H).

10

EXAMPLE 120

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)-N-(4-((trans-4-hydroxycyclohexyl)methoxy)-3-nitrophenylsulfonyl)benzamide

15

EXAMPLE 120A

Trans-4-(aminomethyl)cyclohexanol

20

[0638] Tert-butyl ((1r,4r)-4-hydroxycyclohexyl)methylcarbamate (1 g) in dichloromethane (10 mL) was treated with trifluoroacetic acid (5 mL) at 0°C for 10 minutes and at room temperature for 30 minutes. The reaction mixture was concentrated and dried in vacuo to provide the title compound as a trifluoroacetic acid salt.

25

EXAMPLE 120B

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)-N-(4-((trans-4-hydroxycyclohexyl)methoxy)-3-nitrophenylsulfonyl)benzamide

30

[0639] A mixture of EXAMPLE 53A (211 mg), EXAMPLE 120A (104 mg) and N-ethyl-N-isopropylpropan-2-amine (0.3 mL) in dimethylsulfoxide (2 mL) was heated at 150°C in a Biotage Initiator microwave synthesizer for 1.5 hours and concentrated. The residue was purified by reverse phase HPLC on a C18 column using a gradient of 40-60% acetonitrile in 0.1% trifluoroacetic acid water to give the title compound as a trifluoroacetate salt. The trifluoroacetic acid salt was dissolved in dichloromethane (30 mL) and washed with 50% aqueous NaHCO₃. The organic layer was dried over anhydrous Na₂SO₄, filtered, and concentrated to give the title compound. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 11.41 (s, 1H), 8.61 (t, 1H), 8.53 - 8.58 (m, 1H), 8.04 (d, 1H), 7.76 - 7.83 (m, 1H), 7.47 - 7.56 (m, 3H), 7.34 (d, 2H), 7.07 - 7.11 (m, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.19 (d, 1H), 4.82 - 4.99 (m, 1H), 4.50 (d, 1H), 3.26 - 3.31 (m, 2H), 3.23 (t, 1H), 3.07 (s, 4H), 2.76 (s, 2H), 2.10 - 2.28 (m, 6H), 2.05 (dd, 1H), 1.95 (s, 2H), 1.84 (t, 2H), 1.52 - 1.76 (m, 2H), 1.41 - 1.51 (m, 1H), 1.38 (t, 2H), 0.95 - 1.25 (m, 4H), 0.92 (s, 6H)

35

40

EXAMPLE 121

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)-N-(4-((cis-4-methoxycyclohexyl)methoxy)-3-nitrophenylsulfonyl)benzamide

45

EXAMPLE 121A

(4-methoxycyclohexyl)methanol

50

[0640] 4-Methoxycyclohexanecarboxylic acid (7 g) in tetrahydrofuran (20 mL) was treated with 1 M (in tetrahydrofuran) borane-tetrahydrofuran complex (100 mL) overnight. The mixture was concentrated and the residue was dissolved in methanol (100 mL) and concentrated HCl (10 mL). The resulting mixture was stirred for 1 hour and concentrated. The residue was dissolved in dichloromethane and washed with water. The organic layer was dried over Na₂SO₄, filtered, and concentrated to give the title compound.

55

EXAMPLE 121B

4-((4-methoxycyclohexyl)methoxy)-3-nitrobenzenesulfonamide

- 5 **[0641]** The title compound was prepared as described in EXAMPLE 12A by replacing (1,4-dioxan-2-yl)methanol with EXAMPLE 121A.

EXAMPLE 121C

- 10 4-((cis-4-methoxycyclohexyl)methoxy)-3-nitrobenzenesulfonamide

[0642] Separation of the cis and trans mixture of EXAMAPLE 121B on a reverse phase HPLC (gradient: 40-55% acetonitrile in 0.1% TFA in water over 25 minutes) provided the title compound.

- 15 EXAMPLE 121D

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-(4-((cis-4-methoxycyclohexyl)methoxy)-3-nitrophenylsulfonyl)benzamide

- 20 **[0643]** The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 121C in place of EXAMPLE 11B. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 11.39 (s, 1H), 8.34 (s, 1H), 7.96 - 8.07 (m, 2H), 7.48 - 7.56 (m, 3H), 7.31 - 7.42 (m, 3H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.40 (dd, 1H), 6.20 (d, 1H), 4.02 (d, 2H), 3.39 (s, 1H), 3.20 (s, 3H), 3.09 (s, 4H), 2.82 (s, 2H), 2.09 - 2.34 (m, 6H), 1.96 (s, 2H), 1.78 - 1.86 (m, 3H), 1.54 (dd, 2H), 1.28 - 1.46 (m, 6H), 0.92 (s, 6H)

25

EXAMPLE 122

Cis-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-(cyclopropylamino)cyclohexyl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

30

EXAMPLE 122A

Cis-tert-butyl-4-(cyclopropylamino)cyclohexylcarbamate

- 35 **[0644]** The title compound was prepared by substituting tert-butyl 4-oxocyclohexylcarbamate for 4'-chlorobiphenyl-2-carboxaldehyde and cyclopropylamine for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A.

EXAMPLE 122B

- 40 cis-N¹-cyclopropylcyclohexane-1,4-diamine bis(2,2,2-trifluoroacetate)

[0645] The title compound was prepared by substituting EXAMPLE 122A for EXAMPLE 39A in EXAMPLE 39B.

EXAMPLE 122C

45

Cis-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-(cyclopropylamino)cyclohexyl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- 50 **[0646]** The title compound was prepared by substituting EXAMPLE 122B for EXAMPLE 100B in EXAMPLE 100C. ¹H NMR (500 MHz, pyridine-d₅) δ 13.06 (s, 1H), 9.28 (d, 1H), 8.59 (d, 1H), 8.44 (d, 1H), 8.37 (dd, 1H), 8.12 (d, 1H), 7.67 (t, 2H), 7.43 (t, 2H), 7.07 (d, 2H), 6.90 (d, 1H), 6.75 (dd, 1H), 6.53 (d, 1H), 6.50 (dd, 1H), 3.56 - 3.63 (m, 1H), 3.02 - 3.08 (m, 4H), 2.77 (s, 3H), 2.26 (t, 2H), 2.10 - 2.16 (m, 4H), 2.06 (ddd, 1H), 1.97 (s, 2H), 1.74 - 1.82 (m, 2H), 1.61 - 1.71 (m, 5H), 1.39 (t, 2H), 0.93 (s, 6H), 0.39 - 0.44 (m, 4H).

55

EXAMPLE 123

Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(3-nitro-4-([4-(tetrahydro-2H-pyran-4-ylamino)cyclohexyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 123A

Trans-tert-butyl-4-(tetrahydro-2H-pyran-4-ylamino)cyclohexylcarbamate

[0647] The title compound was prepared by substituting trans-tert-butyl-4-aminocyclohexylcarbamate for tert-butyl piperazine-1-carboxylate and dihydro-2H-pyran-4(3H)-one for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 123B

trans-N1-(tetrahydro-2H-pyran-4-yl)cyclohexane-1,4-diamine bis(2,2,2-trifluoroacetate)

[0648] The title compound was prepared by substituting EXAMPLE 123A for EXAMPLE 39A in EXAMPLE 39B.

EXAMPLE 123C

Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(3-nitro-4-([4-(tetrahydro-2H-pyran-4-ylamino)cyclohexyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0649] The title compound was prepared by substituting EXAMPLE 123B for EXAMPLE 100B in EXAMPLE 100C. ¹H NMR (500 MHz, pyridine-d₅) δ 13.01 (s, 1H), 9.28 (d, 1H), 8.48 (d, 1H), 8.38 (dd, 1H), 8.32 (d, 1H), 8.24 (d, 1H), 7.67 - 7.69 (m, 2H), 7.44 (d, 2H), 7.08 (d, 2H), 6.91 (d, 1H), 6.78 (dd, 1H), 6.59 (d, 1H), 6.48 (dd, 1H), 4.01 (d, 2H), 3.44 - 3.49 (m, 1H), 3.37 - 3.43 (m, 2H), 3.01 - 3.09 (m, 5H), 2.85 (t, 1H), 2.78 (s, 2H), 2.27 (t, 2H), 2.13 - 2.18 (m, 4H), 2.05 (t, 4H), 1.97 (s, 2H), 1.93 (d, 2H), 1.52 - 1.60 (m, 2H), 1.44 - 1.50 (m, 2H), 1.39 (t, 2H), 1.25 - 1.34 (m, 2H), 0.94 (s, 6H).

EXAMPLE 124

Trans-N-({5-bromo-6-([4-morpholin-4-ylcyclohexyl]oxy)pyridin-3-yl}sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 124A

Trans-4-morpholinocyclohexanol

[0650] Trans-4-Aminocyclohexanol (0.5 g), 1-bromo-2-(2-bromoethoxy)ethane (1.07 g) and triethylamine (2.42 mL) were dissolved in anhydrous acetonitrile (20 mL). The reaction mixture was heated at 60°C overnight. The organic solvent was removed under vacuum. The residue was purified with flash column chromatography on silica gel eluting with 7%-10% methanol in dichloromethane to give the title compound.

EXAMPLE 124B

Trans-5-bromo-6-(4-morpholinocyclohexyloxy)pyridine-3-sulfonamide

[0651] The title compound was prepared by substituting EXAMPLE 124A for (tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 36A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 124C

Trans-N-({5-bromo-6-([4-morpholin-4-ylcyclohexyl]oxy)pyridin-3-yl}sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0652] The title compound was prepared by substituting EXAMPLE 124B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 8.56 (m, 2H), 8.03 (d, 1H), 7.80 (m, 1H), 7.50 (m, 3H), 7.34 (d, 2H), 7.12 (m, 1H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.39 (dd, 1H), 6.19 (d, 1H), 3.99 (m, 1H), 3.67 (m, 1H), 3.37 (m, 2H), 3.24

(m, 2H), 3.07 (m, 4H), 2.89 (m, 1H), 2.71 (m, 2H), 2.16 (m, 6H), 1.96 (s, 3H), 1.80 (m, 4H), 1.38 (t, 2H), 1.27 (m, 2H), 0.92 (s, 6H).

EXAMPLE 125

Trans-4-({[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-methoxycyclohexyl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 125A

4-(((trans)-4-methoxycyclohexyl)methoxy)-3-nitrobenzenesulfonamide

[0653] Separation of the *cis* and *trans* mixture of EXAMAPLE 121B on a reverse phase HPLC provided the title compound.

EXAMPLE 125B

Trans-4-({[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-methoxycyclohexyl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0654] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 125A in place of EXAMPLE 11B. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 8.34 (s, 1H), 7.96 - 8.09 (m, 2H), 7.51 (dd, 3H), 7.32 - 7.39 (m, 3H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.20 (d, 1H), 4.02 (d, 2H), 3.24 (s, 3H), 3.00 - 3.15 (m, 5H), 2.83 (s, 2H), 2.09 - 2.36 (m, 6H), 2.03 (d, 2H), 1.96 (s, 2H), 1.77 - 1.86 (m, 2H), 1.73 (s, 1H), 1.39 (t, 2H), 1.02 - 1.17 (m, 4H), 0.92 (s, 6H)

EXAMPLE 126

tert-butyl 4-({[4-({[4-(4-({[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino}sulfonyl)-2-nitrophenoxy]methyl}-4-fluoropiperidine-1-carboxylate

EXAMPLE 126A

tert-butyl 4-fluoro-4-(hydroxymethyl)piperidine-1-carboxylate

[0655] 1-Tert-butyl 4-ethyl 4-fluoropiperidine-1,4-dicarboxylate (1.0 g) in tetrahydrofuran (5 mL) was treated with 1.0 N LiAlH₄ in THF (2.54 mL) at 0 °C. The reaction mixture was stirred at room temperature for 2 hours. Water (0.6 mL) was added to the reaction mixture drop-wise, followed by 2 N aqueous NaOH (0.2 mL). The reaction was stirred for another 1 hour. The solid was removed by filtration via a pack of Celite and washed with ethyl acetate. The filtrate was washed with brine, dried over MgSO₄, filtered, and concentrated to give the product.

EXAMPLE 126B

tert-butyl 4-fluoro-4-((2-nitro-4-sulfamoylphenoxy)methyl)piperidine-1-carboxylate

[0656] The title compound was prepared by substituting EXAMPLE 126A for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 126C

tert-butyl 4-({[4-({[4-(4-({[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino}sulfonyl)-2-nitrophenoxy]methyl}-4-fluoropiperidine-1-carboxylate)oxy]benzamide

[0657] The title compound was prepared by substituting EXAMPLE 126B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 8.36 (s, 2H), 8.02-8.06 (m, 2H), 7.49-7.53 (m, 3H), 7.40 (d, 1H), 7.35 (d, 2H), 7.04 (d, 1H), 6.67 (dd, 1H), 6.39 (dd, 1H), 6.21 (d, 1H), 4.36 (d, 2H), 3.83-3.85 (m, 2H), 3.09 (s, 4H), 2.33 (s, 2H), 2.27-2.32 (m, 4H), 2.13-2.16 (m, 2H), 1.96 (s, 2H), 1.83-1.92 (m, 2H), 1.67-1.75 (m, 2H), 1.38-1.41 (m, 11H), 0.92 (s, 6H).

EXAMPLE 127

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-fluoropiperidin-4-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0658] The title compound was prepared by substituting EXAMPLE 126C for EXAMPLE 1A in EXAMPLE 1B. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.50 (s, 1H), 8.14 (d, 1H), 7.90 (d, 2H), 7.80 (dd, 1H), 7.60 (d, 1H), 7.40 (t, 1H), 7.35 (d, 2H), 7.25 (t, 1H), 7.13 (d, 1H), 7.05 (d, 2H), 6.61 (dd, 1H), 6.30 (dd, 1H), 6.26 (d, 1H), 4.28 (d, 2H), 3.10-3.13 (m, 2H), 2.91-3.00 (m, 6H), 2.73 (s, 2H), 1.96-2.02 (m, 4H), 1.77-1.89 (m, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 128

Trans-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[4-(4-tetrahydro-2H-pyran-4-yl)piperazin-1-yl]cyclohexyl}amino)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 128A

tert-butyl 4-(tetrahydro-2H-pyran-4-yl)piperazine-1-carboxylate

[0659] The title compound was prepared by substituting tert-butyl piperazine-1-carboxylate for morpholine and dihydro-2H-pyran-4(3H)-one for tert-butyl 4-oxocyclohexylcarbamate in EXAMPLE 39A.

EXAMPLE 128B

1-(tetrahydro-2H-pyran-4-yl)piperazine dihydrochloride

[0660] To a solution of EXAMPLE 128A (3.92 g) in ether was added HCl (25 ml, 2M in ether) and the reaction mixture was stirred for 16 hours at room temperature. The solid product was filtered off, dried and used in next step without further purification.

EXAMPLE 128C

Trans-tert-butyl-4-(4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl)cyclohexylcarbamate

[0661] The title compound was prepared by substituting EXAMPLE 128B for morpholine in EXAMPLE 39A.

EXAMPLE 128D

trans-4-(4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl)cyclohexanamine tris(2,2,2-trifluoroacetate)

[0662] The title compound was prepared by substituting EXAMPLE 128C for EXAMPLE 39A in EXAMPLE 39B.

EXAMPLE 128E

Trans-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[4-(4-tetrahydro-2H-pyran-4-yl)piperazin-1-yl]cyclohexyl}amino)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0663] The title compound was prepared by substituting EXAMPLE 128D for EXAMPLE 100B in EXAMPLE 100C. ¹H NMR (500 MHz, pyridine-d₅) δ 13.07 (s, 1H), 9.28 - 9.32 (m, 1H), 8.44 (t, 1H), 8.34 - 8.39 (m, 2H), 8.10 - 8.14 (m, 1H), 7.66 - 7.69 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.92 (t, 1H), 6.73 - 6.77 (m, 1H), 6.52 - 6.55 (m, 1H), 6.49 - 6.52 (m, 1H), 3.99 - 4.06 (m, 2H), 3.29 - 3.36 (m, 2H), 3.03 - 3.09 (m, 4H), 2.77 (s, 2H), 2.62 (s, 8H), 2.24 - 2.29 (m, 3H), 2.10 - 2.16 (m, 5H), 2.05 (s, 2H), 1.97 (s, 2H), 1.92 (s, 2H), 1.70 (d, 2H), 1.57 (td, 2H), 1.34 - 1.43 (m, 4H), 1.20 - 1.30 (m, 2H), 0.93 (s, 6H).

EXAMPLE 129

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl]methoxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 129A

(1-(1,3-difluoropropan-2-yl)piperidin-4-yl)methanol

[0664] A suspension of piperidin-4-ylmethanol (0.250 g), sodium triacetoxyborohydride (0.690 g) and 1,3-difluoropropan-2-one (0.245 g) were stirred together in dichloromethane. After stirring overnight the reaction was poured into saturated aqueous NaHCO₃ solution (10 mL) and stirred for 15 minutes. The reaction was extracted with dichloromethane (3 x 25 mL), dried over magnesium sulfate, filtered, and concentrated. Silica gel chromatography (GraceResolv 12 g) eluting with a gradient of 0.75% to 3% methanol/dichloromethane gave the title compound.

EXAMPLE 129B

4-([1-(1,3-difluoropropan-2-yl)piperidin-4-yl]methoxy)-3-nitrobenzenesulfonamide

[0665] To a solution of (1-(1,3-difluoropropan-2-yl)piperidin-4-yl)methanol (0.068 g) in tetrahydrofuran (1 mL) was added sodium hydride (0.056 g) and the reaction stirred for 30 minutes at room temperature. 4-Fluoro-3-nitrobenzenesulfonamide (0.077 g) was added in one portion and stirring was continued for 1 hour. The reaction was poured into water (20 mL) and extracted with dichloromethane. The pH of the aqueous layer was adjusted to pH~8 and it was extracted with dichloromethane (50 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated to give the title compound.

EXAMPLE 129C

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl]methoxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0666] The title compound was prepared by substituting EXAMPLE 129B for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 11.47 - 10.98 (m, 1H), 8.33 (d, 1H), 8.03 (d, 2H), 7.50 (dd, 3H), 7.36 (t, 3H), 7.04 (d, 2H), 6.67 (d, 1H), 6.39 (dd, 1H), 6.20 (s, 1H), 4.62 (dd, 4H), 4.06 (d, 2H), 3.18 - 2.71 (m, 11H), 2.20 (d, 6H), 1.96 (s, 2H), 1.73 (d, 3H), 1.35 (d, 4H), 0.92 (s, 6H).

EXAMPLE 130

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([(3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]amino)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 130A

(R)-tert-butyl 1-(tetrahydro-2H-pyran-4-yl)pyrrolidin-3-ylcarbamate

[0667] The title compound was prepared by substituting dihydro-2H-pyran-4(3H)-one for 4'-chlorobiphenyl-2-carboxaldehyde and (R)-tert-butyl pyrrolidin-3-ylcarbamate for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A.

EXAMPLE 130B

(R)-1-(tetrahydro-2H-pyran-4-yl)pyrrolidin-3-amine

[0668] A solution of EXAMPLE 130A (550 mg) in dichloromethane (25 mL) was cooled in an ice bath under nitrogen. 2,2,2-Trifluoroacetic acid (8.333 mL) was added and the reaction was stirred for 2 hours. The product was obtained by concentration and high vacuum drying.

EXAMPLE 130C

(R)-3-nitro-4-(1-(tetrahydro-2H-pyran-4-yl)pyrrolidin-3-ylamino)benzenesulfonamide

- 5 **[0669]** The title compound was prepared by substituting EXAMPLE 130B for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 130D

- 10 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0670]** To a solution of EXAMPLE 3J (90 mg), EXAMPLE 130C (64.2 mg), triethylamine (0.077 ml), N,N-dimethylpyridin-4-amine (38.5 mg) in a mixture of dichloromethane (5 ml) and N,N-dimethylformamide (0.5 ml) was added N¹-(ethyl-
15 imino)methylene)-N³,N³-dimethylpropane-1,3-diamine, hydrochloric acid (60.4 mg) and the mixture was stirred 18 hours. This was concentrated on high vacuum and the crude was purified by reverse phase chromatography with ammonium acetate buffer / acetonitrile. ¹H NMR (500 MHz, pyridine-d₅) δ 13.03 (s, 1H), 9.27 (d, 1H), 8.59 (d, 1H), 8.43 (d, 1H), 8.37 (dd, 1H), 8.11 (d, 1H), 7.65 - 7.67 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.88 (d, 1H), 6.76 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 4.06 (m, 1H), 3.98 (d, 2H), 3.35 (t, 2H), 3.07 (m, 4H), 2.73 - 2.80 (m, 4H), 2.68 - 2.72 (m, 1H), 2.36 (q, 1H),
20 2.11 - 2.30 (m, 9H), 1.97 (m, 2H), 1.62 - 1.71 (m, 3H), 1.48 - 1.58 (m, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 131

- 25 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 131A

tert-butyl (3R)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-ylcarbamate

- 30 **[0671]** The title compound was prepared by substituting 2,2-dimethyldihydro-2H-pyran-4(3H)-one for 4'-chlorobiphenyl-2-carboxaldehyde and (R)-tert-butyl pyrrolidin-3-ylcarbamate for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A.

EXAMPLE 131B

- 35 (3R)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-amine

[0672] The title compound was prepared by substituting EXAMPLE 131A for EXAMPLE 130A in EXAMPLE 130B.

EXAMPLE 131C

4-((3R)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-ylamino)-3-nitrobenzenesulfonamide

- 45 **[0673]** The title compound was prepared by substituting EXAMPLE 131B for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 131D

- 50 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0674]** The title compound was prepared by substituting EXAMPLE 131C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500 MHz, pyridine-d₅) δ 13.03 (d, 1H), 9.28 (m, 1H), 8.61 (m, 1H), 8.44 (d, 1H), 8.38 (dd, 1H), 8.11 (d, 1H), 7.64 - 7.68 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.89 (m, 1H), 6.76 (dd, 1H), 6.54 (m, 1H), 6.49 (m, 1H), 4.08 (m, 1H),
55 3.78 (m, 1H), 3.61 (m, 1H), 3.07 (m, 4H), 2.71 - 2.82 (m, 5H), 2.37 - 2.44 (m, 2H), 2.19 - 2.29 (m, 3H), 2.14 (m, 5H), 1.97 (s, 2H), 1.76 (m, 1H), 1.66 (m, 2H), 1.32 - 1.49 (m, 4H), 1.28 (d, 3H), 1.20 (s, 3H), 0.94 (s, 6H).

EXAMPLE 132

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(3-nitro-4-((3S)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl)amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 132A

(S)-tert-butyl 1-(tetrahydro-2H-pyran-4-yl)pyrrolidin-3-ylcarbamate

[0675] The title compound was prepared by substituting dihydro-2H-pyran-4(3H)-one for 4'-chlorobiphenyl-2-carboxaldehyde and (S)-tert-butyl pyrrolidin-3-ylcarbamate for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A.

EXAMPLE 132B

(S)-1-(tetrahydro-2H-pyran-4-yl)pyrrolidin-3-amine

[0676] The title compound was prepared by substituting EXAMPLE 132A for EXAMPLE 130A in EXAMPLE 130B.

EXAMPLE 132C

(S)-3-nitro-4-(1-(tetrahydro-2H-pyran-4-yl)pyrrolidin-3-ylamino)benzenesulfonamide

[0677] The title compound was prepared by substituting EXAMPLE 132B for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 132D

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(3-nitro-4-((3S)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl)amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0678] The title compound was prepared by substituting EXAMPLE 132C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500 MHz, pyridine-*d*₅) δ 13.04 (m, 1H), 9.27 (d, 1H), 8.58 (d, 1H), 8.43 (d, 1H), 8.37 (dd, 1H), 8.11 (d, 1H), 7.64 - 7.68 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.88 (d, 1H), 6.75 (dd, 1H), 6.54 (m, 1H), 6.49 (m, 1H), 4.06 (m, 1H), 3.98 (d, 2H), 3.36 (t, 2H), 3.07 (m, 4H), 2.68 - 2.80 (m, 5H), 2.36 (m, 1H), 2.09 - 2.29 (m, 9H), 1.97 (s, 2H), 1.62 - 1.72 (m, 3H), 1.48 - 1.60 (m, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 133

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(4-((3S)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 133A

tert-butyl (3S)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-ylcarbamate

[0679] The title compound was prepared by substituting 2,2-dimethyldihydro-2H-pyran-4(3H)-one for 4'-chlorobiphenyl-2-carboxaldehyde and (S)-tert-butyl pyrrolidin-3-ylcarbamate for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A.

EXAMPLE 133B

(3S)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-amine

[0680] The title compound was prepared by substituting EXAMPLE 133A for EXAMPLE 130A in EXAMPLE 130B.

EXAMPLE 133C

4-(3S)-(1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-ylamino)-3-nitro benzenesulfonamide

- 5 **[0681]** The title compound was prepared by substituting EXAMPLE 133B for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 133D

- 10 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3S)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0682]** The title compound was prepared by substituting EXAMPLE 133C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500 MHz, pyridine-*d*₅) δ 13.03 (d, 1H), 9.28 (m, 1H), 8.61 (m, 1H), 8.43 (d, 1H), 8.38 (dd, 1H), 8.11 (d, 1H), 7.64 - 7.68 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.89 (m, 1H), 6.76 (dd, 1H), 6.54 (m, 1H), 6.49 (m, 1H), 4.08 (m, 1H), 3.78 (m, 1H), 3.61 (m, 1H), 3.07 (m, 4H), 2.71 - 2.82 (m, 5H), 2.37 - 2.44 (m, 2H), 2.19 - 2.29 (m, 3H), 2.14 (m, 5H), 1.97 (s, 2H), 1.76 (m, 1H), 1.66 (m, 2H), 1.33 - 1.48 (m, 4H), 1.28 (d, 3H), 1.20 (s, 3H), 0.94 (s, 6H).

EXAMPLE 134

- 20 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-methylmorpholin-2-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 134A

- 25 4-(morpholin-2-ylmethylamino)-3-nitrobenzenesulfonamide

- [0683]** A solution of EXAMPLE 113A (0.8 g) in dichloromethane (10 mL) and trifluoroacetic acid (10 mL) was stirred at room temperature for 2 hours. The solvents were evaporated and the residue triturated with diethyl ether. The resulting solid was dissolved in 5% aqueous sodium carbonate solution (20 mL). The solution was concentrated to dryness and the resulting solid was triturated with a solution of 10% methanol in dichloromethane several times. Evaporation of the organic solvents gave the title compound.

EXAMPLE 134B

- 35 4-((4-methylmorpholin-2-yl)methylamino)-3-nitrobenzenesulfonamide

- [0684]** To a solution of EXAMPLE 134A (158 mg) in anhydrous *N,N*-dimethylformamide (4 mL) was added sodium carbonate (64 mg) and methyl iodide (78 mg). After stirring overnight at room temperature, the mixture was evaporated to dryness. The crude product was then absorbed on silica gel (6 g) and purified on a silica gel column eluting with 10% methanol in dichloromethane to give the title compound.

EXAMPLE 134C

- 45 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-methylmorpholin-2-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0685]** The title compound was prepared by substituting EXAMPLE 134B for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-*d*₅) δ 13.00 (s, 1H), 9.27 (d, 1H), 8.87 (t, 1H), 8.43 (d, 1H), 8.32 (dd, 1H), 8.11 (d, 1H), 7.65 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.91 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 3.92 (m, 1H), 3.86 (d, 1H), 3.67 (dt, 1H), 3.49-3.39 (m, 2H), 3.07 (m, 4H), 2.77 (s, 2H), 3.71 (m, 1H), 2.49 (d, 1H), 2.26 (m, 2H), 2.16 (s, 3H), 2.14 (m, 4H), 2.03 (dt, 1H), 1.97 (s, 2H), 1.90 (t, 1H), 1.39 (t, 2H), 0.94 (s, 6H).

55

EXAMPLE 135

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[4-(2-methoxyethyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 135A

4-((4-(2-methoxyethyl)morpholin-2-yl)methylamino)-3-nitrobenzenesulfonamide

[0686] The title compound was prepared by substituting 2-methoxyethyl bromide for methyl iodide in EXAMPLE 134B.

EXAMPLE 135B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[4-(2-methoxyethyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0687] The title compound was prepared by substituting EXAMPLE 135A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 12.98 (s, 1H), 9.26 (d, 1H), 8.88 (t, 1H), 8.43 (d, 1H), 8.32 (dd, 1H), 8.11 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.91 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 3.93 (m, 1H), 3.87 (d, 1H), 3.70 (dt, 1H), 3.51 (t, 2H), 3.48-3.38 (m, 2H), 3.27 (s, 3H), 3.07 (m, 4H), 2.95 (d, 1H), 2.77 (s, 2H), 2.70 (m, 1H), 2.57 (t, 2H), 2.27-2.07 (m, 8H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 136

N-[[4-({[4-(4-acetylmorpholin-2-yl)methyl]amino)-3-nitrophenyl]sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 136A

4-((4-(4-acetylmorpholin-2-yl)methylamino)-3-nitrobenzenesulfonamide

[0688] The title compound was prepared by substituting acetic anhydride for methyl iodide in EXAMPLE 134B.

EXAMPLE 136B

N-[[4-({[4-(4-acetylmorpholin-2-yl)methyl]amino)-3-nitrophenyl]sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0689] The title compound was prepared by substituting EXAMPLE 136A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.26 (s, 1H), 8.85 (s, 1H), 8.43 (d, 1H), 8.32 (dd, 1H), 8.10 (d, 1H), 7.65 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.91 (dd, 1H), 6.75 (dd, 1H), 6.54 (s, 1H), 6.48 (s, 1H), 4.73 (dd, 1H), 3.93-3.65 (m, 2H), 3.60-3.40 (m, 4H), 3.12 (m, 1H), 3.07 (m, 4H), 2.77 (s, 2H), 2.70 (m, 1H), 2.57 (t, 2H), 2.14 (s, 3H), 2.27-2.07 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 137

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[trans-4-(fluoromethyl)-1-oxetan-3-yl]pyrrolidin-3-yl]methoxy}-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 137A

ethyl 4-fluorobut-2-enoate

[0690] Ethyl 2-fluoroacetate (21.0 g) in CH₂Cl₂ (200 mL) at -78 °C was treated dropwise over 45 min with a 1.0 M solution of diisobutylaluminum hydride in CH₂Cl₂ (200 mL) maintaining the internal temperature below -70 °C. Stirring was continued at -78 °C for 30 minutes and then (carbethoxymethylene)triphenylphosphorane (70.0 g) was added in one portion. The reaction mixture was allowed to slowly reach room temperature while stirring overnight. It was then quenched with methanol, filtered and concentrated to give the product as a mixture of isomers (E/Z = 3:1).

EXAMPLE 137B

Trans-ethyl 1-benzyl-4-(fluoromethyl)pyrrolidine-3-carboxylate

- 5 **[0691]** A mixture of N-benzyl-1-methoxy-N-((trimethylsilyl)methyl)methanamine (4.5 g) and EXAMPLE 137A (2.5 g) in dichloromethane (50 mL) was cooled to 0 °C, treated dropwise with trifluoroacetic acid (0.15 mL), stirred for 4 hours at 0 °C and neutralized with saturated aqueous Na₂CO₃ solution. The mixture was poured into a separatory funnel and the layers separated. The organic layer was washed with water and brine, dried (MgSO₄), filtered, concentrated and chromatographed on silica gel with 0-20% ethyl acetate in hexanes as eluent to give both the cis and trans isomers of the product. Only the trans diastereomers were carried on in the following steps.

EXAMPLE 137C

Trans-ethyl 4-(fluoromethyl)pyrrolidine-3-carboxylate

- 15 **[0692]** EXAMPLE 137B (0.83 g) in ethanol (9 mL) was treated with 10% Pd/C (0.208 g) and ammonium formate (1.97 g), refluxed for 1.5 hours, concentrated, dissolved in dichloromethane, filtered through a pad of celite rinsing with dichloromethane, and concentrated to give the product.

20 EXAMPLE 137D

- [0693]** Trans-1-benzyl 3-ethyl 4-(fluoromethyl)pyrrolidine-1,3-dicarboxylate EXAMPLE 137C (0.44 g) in dioxane (4 mL) and water (4 mL) at 0 °C was treated sequentially with Na₂CO₃ (0.89 g) and benzyl chloroformate (0.48 mL). The reaction mixture was stirred at 0 °C for 3 hours and was then allowed to slowly warm to room temperature over 1.5 hours. The reaction mixture was diluted with ethyl acetate, washed with water and brine, dried (MgSO₄), filtered, concentrated and chromatographed on silica gel with 10-25% ethyl acetate in hexanes as eluent to give the product.

EXAMPLE 137E

- 30 Trans-1-(benzyloxycarbonyl)-4-(fluoromethyl)pyrrolidine-3-carboxylic acid

[0694] The title compound was prepared by substituting EXAMPLE 137D for EXAMPLE 15G in EXAMPLE 15H.

EXAMPLE 137F

- 35 Trans-benzyl 3-(fluoromethyl)-4-(hydroxymethyl)pyrrolidine-1-carboxylate

- [0695]** EXAMPLE 137E (0.563 g) in tetrahydrofuran (10 mL) at 0 °C was treated dropwise with a 1 M solution of borane in tetrahydrofuran (4 mL), stirred for 3 hours and then slowly quenched with saturated aqueous NH₄Cl solution. The reaction mixture was diluted with ethyl acetate, washed with water and brine, dried (MgSO₄), filtered and concentrated to give the product.

EXAMPLE 137G

- 45 Trans-benzyl 3-(fluoromethyl)-4-((2-nitro-4-sulfamoylphenoxy)methyl)pyrrolidine-1-carboxylate

[0696] The title compound was prepared by substituting EXAMPLE 137F for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

50 EXAMPLE 137H

Trans-4-((4-(fluoromethyl)pyrrolidin-3-yl)methoxy)-3-nitrobenzenesulfonamide

- 55 **[0697]** EXAMPLE 137G (0.232g) in acetic acid (2.5 ml) was treated with hydrobromic acid (33 wt % in acetic acid) (0.875 mL) at ambient temperature, stirred for 1 hour and concentrated. The product was free-based using a MEGA BE-SCX column with 1:1 CH₂Cl₂/methanol as eluent for the hydrobromic acid and acetic acid. The product was released from the column with 10% (7 M ammonia in methanol) in CH₂Cl₂ as eluent.

EXAMPLE 137I

Trans-4-((4-(fluoromethyl)-1-(oxetan-3-yl)pyrrolidin-3-yl)methoxy)-3-nitrobenzenesulfonamide

- 5 **[0698]** The title compound was prepared by substituting EXAMPLE 137H for tert-butyl piperazine-1-carboxylate and 3-oxetanone for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 137J

- 10 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl}methyl}piperazin-1-yl)-N-[(4-{{trans-4-(fluoromethyl)-1-oxetan-3-ylpyrrolidin-3-yl}methoxy}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0699]** The title compound was prepared by substituting EXAMPLE 137I for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 8.35 (d, 1H), 8.03 (m, 2H), 7.51 (m, 3H), 7.37 (m, 3H), 7.04 (m, 2H), 6.67 (dd, 1H), 6.39 (dd, 1H), 6.21 (d, 1H), 4.45 (m, 6H), 4.21 (d, 2H), 3.62 (m, 1H), 3.08 (m, 4H), 2.72 (m, 5H), 2.31 (m, 9H), 1.96 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 138

- 20 4-(4-{{2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl}methyl}piperazin-1-yl)-N-[(4-{{(4-fluorotetrahydro-2H-pyran-4-yl)methyl}amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 138A

- 25 (4-fluorotetrahydro-2H-pyran-4-yl)methyl methanesulfonate

- [0700]** A mixture of EXAMPLE 37C (1.4 g), methanesulfonyl chloride (1.054 mL), triethylamine (2.99 mL), and 4-dimethylaminopyridine (0.051 g) in CH₂Cl₂ (20 mL) was stirred at 0 °C for 2 hours, concentrated and chromatographed on silica gel eluting with 30% ethyl acetate in hexanes to give the product.

EXAMPLE 138B

2-((4-fluorotetrahydro-2H-pyran-4-yl)methyl)isoindoline-1,3-dione

- 35 **[0701]** A mixture of EXAMPLE 138A (1.8 g) and potassium phthalimide (2.356 g) in N,N-dimethylformamide (30 mL) was heated at 150 °C overnight, diluted with ethyl acetate, washed with water and brine, dried (MgSO₄), filtered, concentrated and chromatographed on silica gel eluting with 30% ethyl acetate in hexanes to give the product.

EXAMPLE 138C

- 40 (4-fluorotetrahydro-2H-pyran-4-yl)methanamine

- [0702]** A mixture of EXAMPLE 138B (1.4 g) and hydrazine (1.548 mL) in ethanol (40 mL) was heated at 70 °C overnight, cooled to room temperature, slurried with CH₂Cl₂ (200 mL) and the solid removed by filtration. The filtrate was concentrated and chromatographed on silica gel eluting with 100:5:1 ethyl acetate/methanol/NH₄OH to give the product.

EXAMPLE 138D

4-((4-fluorotetrahydro-2H-pyran-4-yl)methylamino)-3-nitrobenzenesulfonamide

- 50 **[0703]** A mixture of 4-fluoro-3-nitrobenzenesulfonamide (0.44 g), EXAMPLE 138C (0.266 g), and triethylamine (1.11 mL) in tetrahydrofuran (10 mL) was heated at 70 °C overnight, diluted with ethyl acetate, washed with water and brine, dried (MgSO₄), filtered, concentrated and chromatographed on silica gel eluting with 50% ethyl acetate in hexanes to give the product.

EXAMPLE 138E

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-fluorotetrahydro-2H-pyran-4-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0704] The title compound was prepared by substituting EXAMPLE 138D for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 8.62 (t, 1H), 8.56 (d, 1H), 8.04 (d, 1H), 7.82 (dd, 1H), 7.48-7.54 (m, 3H), 7.34 (d, 2H), 7.24 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.19 (d, 1H), 3.70-3.77 (m, 4H), 3.50-3.55 (m, 2H), 3.07 (s, 4H), 2.76 (s, 2H), 2.14-2.20 (m, 6H), 1.76-1.84 (m, 4H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 139

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(1-oxetan-3-yl)piperidin-4-yl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 139A

tert-butyl 4-(4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)sulfamoyl)-2-nitrophenylamino)piperidine-1-carboxylate

[0705] The title compound was prepared as described in EXAMPLE 53B by replacing 1-acetylpiperidin-4-amine with 4-amino-piperidine-1-carboxylic acid tert-butyl ester.

EXAMPLE 139B

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-(3-nitro-4-(piperidin-4-ylamino)phenyl)sulfonyl)benzamide

[0706] To a cooled (0°C) solution of EXAMPLE 139A (960 mg) in dichloromethane (10 mL) was added dropwise trifluoroacetic acid (5 mL). The mixture was stirred at the temperature for 3 hours. Then, the mixture was concentrated under vacuum and the residue was dissolved in dichloromethane (200 mL) and washed with aqueous NaHCO₃ and brine. After drying over Na₂SO₄, the mixture was filtered, and evaporation of the solvent from the filtrate gave the title compound.

EXAMPLE 139C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(1-oxetan-3-yl)piperidin-4-yl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0707] To a solution of EXAMPLE 139B (120 mg) in tetrahydrofuran (3 mL) and acetic acid (1 mL) was added oxetan-3-one (50.8 mg) and MP-cyanoborohydride (2.15 mmol/g, 150 mg). The mixture was stirred at room temperature overnight. The mixture was filtered. The filtrate was concentrated and the residue was loaded on a silica gel cartridge and eluted with 5-10% 7N NH₃ in methanol in dichloromethane to give the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.62 (s, 1H), 8.51 (d, 1H), 8.20 (d, 1H), 7.99 (d, 1H), 7.74 (m, 1H), 7.48 (m, 3H), 7.35 (d, 2H), 7.05 (d, 2H), 6.66 (dd, 1H), 6.36 (dd, 1H), 6.20 (d, 1H), 4.54 (t, 2H), 4.43 (t, 2H), 3.66 (m, 1H), 3.44 (m, 3H), 3.04 (m, 5H), 2.73 (s, 2H), 2.61 (m, 2H), 2.12 (m, 11H), 1.61 (m, 2H), 1.38 (t, 2H), 0.93 (m, 6H).

EXAMPLE 140

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1-cyclobutyl)piperidin-4-yl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0708] The title compound was prepared as described in EXAMPLE 139C by replacing oxetan-3-one with cyclobutanone. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.58 (s, 1H), 8.47 (d, 1H), 8.12 (d, 1H), 7.97 (d, 1H), 7.74 (d, 1H), 7.53 (d, 1H), 7.45 (m, 1H), 7.36 (m, 3H), 7.02 (m, 3H), 6.64 (dd, 1H), 6.33 (m, 1H), 6.22 (d, 1H), 3.74 (m, 1H), 2.97 (m, 6H), 2.73 (s, 3H), 2.15 (m, 15H), 1.67 (m, 4H), 1.38 (t, 2H), 0.93 (s, 6H).

EXAMPLE 141

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0709] The title compound was prepared as described in EXAMPLE 139C by replacing oxetan-3-one with 2,2-dimethyltetrahydropyran-4-one. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.60 (s, 1H), 8.50 (d, 1H), 8.15 (m, 1H), 7.99 (d, 1H), 7.78 (m, 1H), 7.62 (m, 1H), 7.47 (m, 3H), 7.34 (m, 3H), 7.05 (m, 3H), 6.65 (m, 2H), 6.35 (dd, 1H), 6.21 (d, 1H), 4.56 (d, 3H), 3.89 (m, 3H), 3.67 (m, 6H), 3.45 (m, 2H), 3.04 (m, 3H), 2.75 (m, 3H), 2.14 (m, 3H), 1.71 (m, 5H), 1.16 (s, 9H).

EXAMPLE 142

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([(3S)-1-cyclopropylpyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 142A

(S)-tert-butyl 1-cyclopropylpyrrolidin-3-ylcarbamate

[0710] (S)-tert-butyl pyrrolidin-3-ylcarbamate (415 mg), (1-ethoxycyclopropoxy)trimethylsilane (1.8 mL) and molecular sieves (500 mg) were combined in methanol (4.5 mL). Acetic acid (1.3 mL) was added, followed by sodium cyanoborohydride (420 mg). The resulting mixture was heated to reflux for 4 hours. Insoluble material was filtered off and reaction was made basic to pH 14 with addition of 6M aqueous NaOH solution. The solution was extracted three times with diethyl ether, and the combined extracts were dried over MgSO₄, filtered and concentrated to obtain an oil, which was purified by flash chromatography, eluting first with 100% dichloromethane, followed by 5% methanol/dichloromethane and 10% methanol/dichloromethane.

EXAMPLE 142B

(S)-1-cyclopropylpyrrolidin-3-amine

[0711] The title compound was prepared by substituting EXAMPLE 142A for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 142C

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([(3S)-1-cyclopropylpyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0712] The title compound was prepared by substituting EXAMPLE 142B for 1-acetylpiperidin-4-amine in EXAMPLE 53B. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.64 (s, 1H), 8.51 (m, 2H), 8.30 (m, 1H), 8.00 (br s, 1H), 7.77 (m, 1H), 7.49 (m, 3H), 7.34 (d, 2H), 7.04 (d, 2H), 6.97 (br s, 1H), 6.67 (dd, 1H), 6.36 (m, 1H), 6.21 (m, 1H), 4.19 (m, 1H), 3.00 (m, 5H), 2.74 (m, 3H), 2.64 (m, 1H), 2.36 (m, 1H), 2.15 (m, 6H), 1.95 (s, 2H), 1.78 (br s, 1H), 1.68 (m, 1H), 1.38 (t, 2H), 1.23 (m, 1H), 0.92 (s, 6H), 0.39 (m, 4H).

EXAMPLE 143

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-[(1-tetrahydrofuran-3-yl)piperidin-4-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0713] The title compound was prepared as described in EXAMPLE 139C by replacing oxetan-3-one with 3-oxotetrahydrofuran. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1H), 8.53 (d, 1H), 8.21 (m, 1H), 8.02 (m, 1H), 7.80 (dd, 1H), 7.49 (m, 3H), 7.34 (m, 3H), 7.05 (m, 3H), 6.67 (dd, 1H), 6.37 (m, 1H), 6.19 (d, 1H), 4.29 (m, 3H), 3.73 (m, 6H), 3.09 (m, 4H), 2.76 (m, 2H), 2.05 (m, 8H), 1.68 (m, 2H), 1.37 (m, 2H), 0.94 (s, 6H).

EXAMPLE 144

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((3R)-1-cyclopropylpyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 144A

(R)-tert-butyl 1-cyclopropylpyrrolidin-3-ylcarbamate

[0714] The title compound was prepared by substituting (R)-tert-butyl pyrrolidin-3-ylcarbamate for (S)-tert-butyl pyrrolidin-3-ylcarbamate in EXAMPLE 142A.

EXAMPLE 144B

(R)-1-cyclopropylpyrrolidin-3-amine

[0715] The title compound was prepared by substituting EXAMPLE 144A for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 144C

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((3R)-1-cyclopropylpyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0716] The title compound was prepared by substituting EXAMPLE 144B for 1-acetylpiperidin-4-amine in EXAMPLE 53B. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1H), 8.53 (d, 2H), 8.32 (d, 1H), 8.02 (d, 1H), 7.81 (m, 1H), 7.49 (m, 3H), 7.34 (d, 2H), 7.03 (m, 3H), 6.67 (dd, 1H), 6.37 (m, 1H), 6.20 (d, 1H), 4.21 (m, 1H), 3.00 (m, 5H), 2.74 (m, 3H), 2.64 (m, 1H), 2.36 (m, 1H), 2.15 (m, 6H), 1.95 (s, 2H), 1.74 (br s, 1H), 1.66 (m, 1H), 1.38 (t, 2H), 1.23 (m, 1H), 0.92 (s, 6H), 0.39 (m, 4H).

EXAMPLE 145

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-((3S)-1-tetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 145A

(S)-tert-butyl (1-(tetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl)methylcarbamate

[0717] The title compound was prepared by substituting (R)-tert-butyl pyrrolidin-3-ylmethylcarbamate for tert-butyl piperazine-1-carboxylate and dihydro-2H-pyran-4(3H)-one for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 145B

(S)-(1-(tetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl)methanamine

[0718] The title compound was prepared by substituting EXAMPLE 145A for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 145C

(S)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-(3-nitro-4-((1-(tetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl)methylamino)phenyl)sulfonyl)benzamide

[0719] The title compound was prepared by substituting EXAMPLE 145B for 1-acetylpiperidin-4-amine in EXAMPLE 53B. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.58 (s, 1H), 8.61 (br s, 1H), 8.46 (s, 1H), 7.96 (d, 1H), 7.72 (m, 1H), 7.54 (d, 1H), 7.45 (t, 1H), 7.37 (br s, 2H), 7.34 (d, 2H), 7.04 (m, 2H), 6.94 (m, 1H), 6.64 (dd, 1H), 6.34 (m, 1H), 6.22 (d, 1H), 3.89 (m, 2H), 3.38 (m, 4H), 3.27 (m, 4H), 3.02 (m, 5H), 2.73 (s, 2H), 2.61 (m, 1H), 2.18 (m, 6H), 2.05 (m, 1H), 1.95 (m, 2H), 1.85 (m, 2H), 1.64 (m, 1H), 1.50 (m, 2H), 1.38 (m, 2H), 0.94 (s, 6H).

EXAMPLE 146

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(3-hydroxy-2,2-dimethylpropyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0720] The title compound was prepared as described in EXAMPLE 120B using 3-amino-2,2-dimethylpropan-1-ol in place of EXAMPLE 120A. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 11.35 (s, 1H), 8.96 (t, 1H), 8.56 (d, 1H), 8.05 (d, 1H), 7.79 (dd, 1H), 7.46 - 7.56 (m, 3H), 7.34 (d, 2H), 7.10 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.19 (d, 1H), 5.10 (t, 1H), 3.29 (d, 1H), 3.24 (d, 1H), 3.07 (s, 4H), 2.75 (s, 2H), 2.17 (d, 6H), 1.95 (s, 2H), 1.38 (t, 2H), 0.93 (d, 12H).

EXAMPLE 147

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-([1-(methylsulfonyl)piperidin-3-yl]methyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 147A

tert-butyl (1-(methylsulfonyl)piperidin-3-yl)methylcarbamate

[0721] tert-Butyl piperidin-3-ylmethylcarbamate (500 mg) was dissolved in anhydrous dichloromethane (10 mL), and methanesulfonyl chloride (0.181 mL) was added followed by the addition of triethylamine (1.3 mL). The reaction mixture was stirred at room temperature overnight. The organic solvent was removed under vacuum. The residue was purified with flash column chromatography on silica gel eluting with 0-70% ethyl acetate in hexane to give the title compound.

EXAMPLE 147B

(1-(methylsulfonyl)piperidin-3-yl)methanamine

[0722] EXAMPLE 147A (400 mg) was suspended in 4N HCl in dioxane (10 mL) followed by the addition of anhydrous methanol (1 mL). The clear solution was stirred at room temperature for 2 hours. The organic solvent was removed under vacuum. The solid residue was used in the next step without further purification.

EXAMPLE 147C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-([1-(methylsulfonyl)piperidin-3-yl]methyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0723] EXAMPLE 53A (50 mg), EXAMPLE 147B (26 mg) and triethylamine (0.088 mL) were dissolved in anhydrous dioxane (1 mL) and N,N-dimethylformamide (0.2 mL). The reaction vial was heated in a Biotage Initiator microwave reactor at 130°C for 25 minutes. The solvent was removed under vacuum. The residue was purified by reverse phase HPLC on a C18 column using a gradient of 20-80% acetonitrile/0.1% trifluoroacetic acid in water to give the title compound as the trifluoroacetate salt. The trifluoroacetic acid salt was dissolved in dichloromethane (6 mL) and washed with 50% aqueous NaHCO₃. The organic layer was dried over anhydrous Na₂SO₄ and concentrated to give the title compound. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1H), 8.56 (m, 2H), 8.03 (d, 1H), 7.80 (m, 1H), 7.50 (m, 3H), 7.35 (d, 2H), 7.12 (m, 1H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.38 (dd, 1H), 6.19 (d, 1H), 3.52 (m, 1H), 3.40 (m, 2H), 3.06 (m, 4H), 2.84 (s, 3H), 2.75 (m, 2H), 2.75 (m, 4H), 2.58 (m, 1H), 2.16 (m, 6H), 1.95 (s, 3H), 1.76 (m, 2H), 1.52 (m, 1H), 1.37 (m, 2H), 0.92 (s, 6H).

EXAMPLE 148

N-[(4-{[(1-acetylpiperidin-3-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 148A

tert-butyl (1-acetylpiperidin-3-yl)methylcarbamate

[0724] The title compound was prepared by substituting acetyl chloride for methanesulfonyl chloride in EXAMPLE 147A.

EXAMPLE 148B

1-(3-(aminomethyl)piperidin-1-yl)ethanone

[0725] The title compound was prepared by substituting EXAMPLE 148A for EXAMPLE 147A in EXAMPLE 147B.

EXAMPLE 148C

N-[(4-{[(1-acetylpiperidin-3-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0726] The title compound was prepared by substituting EXAMPLE 148B for EXAMPLE 147B in EXAMPLE 147C. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 8.56 (m, 2H), 8.03 (d, 1H), 7.80 (m, 1H), 7.50 (m, 3H), 7.34 (d, 2H), 7.12 (m, 1H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.39 (dd, 1H), 6.19 (d, 1H), 3.99 (m, 1H), 3.67 (m, 1H), 3.37 (m, 2H), 3.24 (m, 2H), 3.07 (m, 4H), 2.89 (m, 1H), 2.71 (m, 2H), 2.16 (m, 6H), 1.96 (s, 3H), 1.80 (m, 4H), 1.38 (t, 2H), 1.27 (m, 2H), 0.92 (s, 6H).

EXAMPLE 149

4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(methylsulfonyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 149A

(R)-tert-butyl 1-(methylsulfonyl)pyrrolidin-3-ylcarbamate

[0727] The title compound was prepared by substituting (R)-tert-butyl pyrrolidin-3-ylcarbamate for tert-butyl piperidin-3-ylmethylcarbamate in EXAMPLE 147A.

EXAMPLE 149B

(R)-1-(methylsulfonyl)pyrrolidin-3-amine

[0728] The title compound was prepared by substituting EXAMPLE 149A for EXAMPLE 147A in EXAMPLE 147B.

EXAMPLE 149C

4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(methylsulfonyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0729] The title compound was prepared by substituting EXAMPLE 149B for EXAMPLE 147B in EXAMPLE 147C. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1H), 8.55 (d, 1H), 8.29 (d, 1H), 8.02 (d, 1H), 7.86 (dd, 1H), 7.49 (m, 3H), 7.33 (d, 2H), 7.17 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.38 (dd, 1H), 6.20 (d, 1H), 4.41 (m, 1H), 3.69 (m, 1H), 3.39 (m, 3H), 3.06 (m, 4H), 2.97 (s, 3H), 2.76 (m, 2H), 2.27 (m, 8H), 1.93 (m, 2H), 1.54 (m, 1H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 150

4-(4-{[2-(4-chlorophenyl)-3,3-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 150A

Ethyl 2-hydroxy-3,3-dimethylcyclohex-1-enecarboxylate

[0730] Into a 500 mL round-bottomed flask was added diisopropylamine (3.5 mL) in ether (200 mL). After cooling to -30°C, butyllithium (16 mL) (1.6M in hexane) was added slowly. After stirring 30 minutes, the temperature was cooled to -5 °C. 2,2-Dimethylcyclohexanone (3 g) was added slowly. The mixture was warmed up to 0°C and stirred for 1 hour. After cooling to -5 °C, hexamethylphosphoramide (8 mL) and ethyl carbonocyanidate (2.5 mL) were added. After stirring at -5 °C for 20 minutes, and warming to room temperature, the reaction was stirred for 1 hour. The mixture was poured into cold water, and the layers were separated. The aqueous layer was extracted with ether (3x 20 mL). The combined organic layers were washed with saturated aqueous NH₄Cl (3x 20 mL). After drying over Na₂SO₄, the mixture was filtered and the filtrate was concentrated. The crude product was purified by flash chromatography on silica with 0-10% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 150B

Ethyl 3,3-dimethyl-2-(trifluoromethylsulfonyloxy)cyclohex-1-enecarboxylate

[0731] The title compound was prepared by substituting EXAMPLE 150A for EXAMPLE 101A in EXAMPLE 101B.

EXAMPLE 150C

Ethyl 2-(4-chlorophenyl)-3,3-dimethylcyclohex-1-enecarboxylate

[0732] The title compound was prepared by substituting EXAMPLE 150B for EXAMPLE 101B in EXAMPLE 101C.

EXAMPLE 150D

(2-(4-chlorophenyl)-3,3-dimethylcyclohex-1-enyl)methanol

[0733] In a 200 mL round-bottomed flask was added EXAMPLE 150C (0.97 g) and lithium borohydride (0.47 g) in ether (20 mL) to give a suspension. Methanol (2.2 mL) was added slowly. The mixture was refluxed overnight. The reaction was then cooled, and methanol was added to quench the reaction. 1N aqueous HCl was then added until the pH<7, and ether (3x 30 mL) was used to extract the product. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated. The crude material was purified by flash chromatography on silica with 0-25% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 150E

2-(4-chlorophenyl)-3,3-dimethylcyclohex-1-enecarbaldehyde

[0734] Into a 100 mL round-bottomed flask was added EXAMPLE 150D (0.3 g) and Dess-Martin Periodinane (0.6 g) in dichloromethane (10 mL) to give a suspension. The mixture was stirred at room temperature overnight. After filtration, the mixture was washed with saturated aqueous NaHCO₃ (2x 20 mL), dried over Na₂SO₄, filtered, and concentrated. The crude product was purified by flash chromatography on silica with 0-25% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 150F

Methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-3,3-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoate

[0735] The title compound was prepared by substituting EXAMPLE 150E for 4'-chlorobiphenyl-2-carboxaldehyde and

EP 2 944 638 A1

EXAMPLE 15F for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A.

EXAMPLE 150G

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-3,3-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoic acid

[0736] The title compound was prepared by substituting EXAMPLE 150F for EXAMPLE 101E in EXAMPLE 101F.

EXAMPLE 150H

4-(4-([2-(4-chlorophenyl)-3,3-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0737] The title compound was prepared by substituting EXAMPLE 150G for EXAMPLE 3J and EXAMPLE 1F for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, dimethyl sulfoxide-d₆) δ 11.50 (s, 1H), 8.36 (m, 1H), 8.32 (m, 1H), 7.91 (d, 1H), 7.59 (m, 2H), 7.40 (t, 1H), 7.35 (d, 2H), 7.25 (m, 1H), 6.94 (d, 2H), 6.79 (d, 1H), 6.60 (m, 1H), 6.29 (m, 1H), 6.24 (d, 1H), 3.83 (m, 2H), 3.25 (m, 4H), 2.98 (m, 4H), 2.42 (s, 2H), 2.14 (m, 6H), 1.60 (m, 6H), 1.25 (m, 3H), 0.86 (s, 6H)

20

EXAMPLE 151

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-({1-[2-fluoro-1-(fluoromethyl)ethyl]azetidin-3-yl}amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

25

EXAMPLE 151A

1-(1,3-difluoropropan-2-yl)azetidin-3-amine

[0738] To a solution of tert-butyl azetidin-3-ylcarbamate (0.256 g) and 1,3-difluoropropan-2-one (0.154 g) in dichloromethane (2 mL) was added sodium triacetoxymethylborohydride (0.473 g) and the reaction was allowed to stir at room temperature. After 16 hours, the reaction was quenched with saturated NaHCO₃ solution (10 mL) and extracted into dichloromethane (25 mL). The organic layer was dried and concentrated. Silica gel chromatography (GraceResolv 12 g) eluting with a gradient of 0.5% to 3.5% methanol/dichloromethane followed by treatment with HCl (4.0M in dioxane, 3 mL) and methanol (0.5 mL) for 2 hours gave the title compound after concentration.

35

EXAMPLE 151B

4-(1-(1,3-difluoropropan-2-yl)azetidin-3-ylamino)-3-nitrobenzenesulfonamide

40

[0739] To a suspension of 4-chloro-3-nitrobenzenesulfonamide (0.225 g) and 1-(1,3-difluoropropan-2-yl)azetidin-3-amine (0.193 g) in dioxane (5 mL) was added diisopropylamine (0.832 mL). The reaction was sonicated and then heated to 100°C. After stirring overnight, the reaction was concentrated and loaded onto silica gel (GraceResolv 12 g) and eluted with a gradient of 0.5% to 3.5% methanol/dichloromethane to give the title compound.

45

EXAMPLE 151C

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-({1-[2-fluoro-1-(fluoromethyl)ethyl]azetidin-3-yl}amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50

[0740] The title compound was prepared by substituting EXAMPLE 151B for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1H), 11.54 - 11.28 (m, 1H), 8.54 (d, 1H), 8.45 (s, 1H), 8.01 (d, 1H), 7.82 (d, 1H), 7.48 (d, 3H), 7.34 (d, 2H), 7.04 (d, 2H), 6.90 (d, 1H), 6.67 (d, 1H), 6.37 (s, 1H), 6.20 (s, 1H), 4.64 - 4.23 (m, 6H), 3.81 (s, 2H), 3.08 (s, 4H), 2.75 (s, 3H), 2.15 (s, 7H), 1.95 (s, 2H), 1.38 (s, 2H), 0.92 (s, 6H).

55

EXAMPLE 152

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{[4-({[1-(methylsulfonyl)pyrrolidin-3-yl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 152A

tert-butyl (1-(methylsulfonyl)pyrrolidin-3-yl)methylcarbamate

[0741] The title compound was prepared by substituting tert-butyl pyrrolidin-3-ylmethylcarbamate for tert-butyl piperidin-3-ylmethylcarbamate in EXAMPLE 147A.

EXAMPLE 152B

(1-(methylsulfonyl)pyrrolidin-3-yl)methanamine

[0742] The title compound was prepared by substituting EXAMPLE 152A for EXAMPLE 147A in EXAMPLE 147B.

EXAMPLE 152C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{[4-({[1-(methylsulfonyl)pyrrolidin-3-yl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0743] The title compound was prepared by substituting EXAMPLE 152B for EXAMPLE 147B in EXAMPLE 147C. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.60 (s, 1H), 8.49 (m, 2H), 7.99 (s, 1H), 7.73 (m, 1H), 7.53 (d, 1H), 7.47 (s, 1H), 7.42 (m, 1H), 7.34 (d, 2H), 7.04 (m, 3H), 6.65 (m, 1H), 6.35 (s, 1H), 6.22 (s, 1H), 3.41 (m, 4H), 3.22 (m, 2H), 3.03 (m, 4H), 2.89 (s, 3H), 2.73 (m, 2H), 2.59 (m, 1H), 2.17 (m, 6H), 2.00 (m, 4H), 1.68 (m, 1H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 153

N-[[4-({[1-(acetylpyrrolidin-3-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 153A

tert-butyl (1-acetylpyrrolidin-3-yl)methylcarbamate

[0744] The title compound was prepared by substituting tert-butyl pyrrolidin-3-ylmethylcarbamate for tert-butyl piperidin-3-ylmethylcarbamate and acetyl chloride for methanesulfonyl chloride in EXAMPLE 147A.

EXAMPLE 153B

1-(3-(aminomethyl)pyrrolidin-1-yl)ethanone

[0745] The title compound was prepared by substituting EXAMPLE 153A for EXAMPLE 147A in EXAMPLE 147B.

EXAMPLE 153C

N-[[4-({[1-(acetylpyrrolidin-3-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0746] The title compound was prepared by substituting EXAMPLE 153B for EXAMPLE 147B in EXAMPLE 147C. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1H), 8.62 (m, 1H), 8.54 (s, 1H), 8.03 (m, 1H), 7.78 (d, 1H), 7.50 (m, 3H), 7.35 (t, 2H), 7.09 (s, 1H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.37 (d, 1H), 6.20 (s, 1H), 3.56 (m, 1H), 3.42 (m, 4H), 3.43 (m, 4H), 3.23 (m, 1H), 3.07 (m, 4H), 2.74 (m, 2H), 2.16 (m, 6H), 1.93 (m, 5H), 1.38 (t, 2H), 0.93 (s, 6H).

EXAMPLE 154

N-[(4-[[[(3R)-1-acetylpyrrolidin-3-yl]amino]-3-nitrophenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 154A

(R)-tert-butyl 1-acetylpyrrolidin-3-ylcarbamate

[0747] The title compound was prepared by substituting (R)-tert-butyl pyrrolidin-3-ylcarbamate for tert-butyl piperidin-3-ylmethylcarbamate and acetyl chloride for methanesulfonyl chloride in EXAMPLE 147A.

EXAMPLE 154B

(R)-1-(3-aminopyrrolidin-1-yl)ethanone

[0748] The title compound was prepared by substituting EXAMPLE 154A for EXAMPLE 147A in EXAMPLE 147B.

EXAMPLE 154C

N-[(4-[[[(3R)-1-acetylpyrrolidin-3-yl]amino]-3-nitrophenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0749] The title compound was prepared by substituting EXAMPLE 154B for EXAMPLE 147B in EXAMPLE 147. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.61 (s, 1H), 8.50 (s, 1H), 8.17 (d, 1H), 7.98 (s, 1H), 7.78 (s, 1H), 7.49 (m, 3H), 7.34 (d, 2H), 7.10 (m, 1H), 7.04 (d, 2H), 6.66 (dd, 1H), 6.35 (s, 1H), 6.22 (s, 1H), 4.34 (m, 1H), 3.81 (m, 1H), 3.58 (m, 1H), 3.43 (m, 1H), 3.05 (m, 4H), 2.74 (s, 2H), 2.19 (m, 9H), 1.96 (m, 5H), 1.38 (t, 2H), 0.94 (s, 6H).

EXAMPLE 155

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(3-methoxy-2,2-dimethylpropyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0750] The title compound was prepared as described in EXAMPLE 120B using 3-methoxy-2,2-dimethylpropan-1-amine in place of EXAMPLE 120A. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 11.32 (s, 1H), 8.92 (t, 1H), 8.57 (d, 1H), 8.04 (d, 1H), 7.80 (dd, 1H), 7.46 - 7.55 (m, 3H), 7.34 (d, 2H), 7.08 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.36 - 6.42 (m, 1H), 6.19 (d, 1H), 3.25 - 3.30 (m, 5H), 3.19 (s, 2H), 3.07 (s, 4H), 2.76 (s, 2H), 2.17 (d, 6H), 1.95 (s, 2H), 1.38 (t, 2H), 0.96 (s, 6H), 0.92 (s, 6H).

EXAMPLE 156

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([[(1R,3R)-3-hydroxycyclopentyl]methyl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 156A

4-(((1R,3R)-3-hydroxycyclopentyl)methylamino)-3-nitrobenzenesulfonamide

[0751] The title compound was prepared by substituting (1R,3R)-3-hydroxycyclopentyl)methylamine for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 156B

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([[(1R,3R)-3-hydroxycyclopentyl]methyl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0752] The title compound was prepared by substituting EXAMPLE 156A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.05 (s, 1H), 9.29 (s, 1H), 8.62 (t, 1H), 8.43 (d, 1H), 8.32 (dd, 1H), 8.10 (d, 1H), 7.66 (m,

2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.85 (d, 1H), 6.74 (dd, 1H), 6.54 (s, 1H), 6.49 (m, 1H), 4.60 (m, 1H), 3.19 (dd, 2H), 3.06 (m, 4H), 2.77 (s, 2H), 2.70 (m, 1H), 2.26 (t, 2H), 2.20-2.07 (m, 6H), 2.00 (m, 1H), 1.97 (s, 2H), 1.90 (m, 1H), 1.56 (m, 1H), 1.39 (t, 2H), 1.34 (m, 1H), 0.93 (s, 6H).

5 EXAMPLE 157

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(1S,3S)-3-hydroxycyclopentyl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 EXAMLE 157A

4-(((1S,3S)-3-hydroxycyclopentyl)methylamino)-3-nitrobenzenesulfonamide

15 **[0753]** The title compound was prepared by substituting (1S, 3S)-3-hydroxycyclopentyl)methylamine for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 157B

20 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(1S,3S)-3-hydroxycyclopentyl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

25 **[0754]** The title compound was prepared by substituting EXAMPLE 157A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.03 (s, 1H), 9.29 (s, 1H), 8.60 (t, 1H), 8.44 (d, 1H), 8.32 (dd, 1H), 8.14 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.83 (d, 1H), 6.75 (dd, 1H), 6.55 (s, 1H), 6.49 (m, 1H), 4.60 (m, 1H), 3.19 (dd, 2H), 3.06 (m, 4H), 2.77 (s, 2H), 2.70 (m, 1H), 2.26 (t, 2H), 2.20-2.07 (m, 6H), 2.00 (m, 1H), 1.97 (s, 2H), 1.90 (m, 1H), 1.56 (m, 1H), 1.39 (t, 2H), 1.34 (m, 1H), 0.93 (s, 6H).

EXAMPLE 158

30 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(1S,3R)-3-hydroxycyclopentyl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 158A

35 4-(((1S,3R)-3-hydroxycyclopentyl)methylamino)-3-nitrobenzenesulfonamide

[0755] The title compound was prepared by substituting (1S, 3R)-3-hydroxycyclopentyl)methylamine for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

40 EXAMPLE 158B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(1S,3R)-3-hydroxycyclopentyl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

45 **[0756]** The title compound was prepared by substituting EXAMPLE 158A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 12.94 (s, 1H), 9.25 (d, 1H), 8.59 (t, 1H), 8.48 (d, 1H), 8.27 (m, 2H), 7.66 (m, 2H), 7.45 (d, 2H), 7.08 (d, 2H), 6.77 (dd, 1H), 6.72 (d, 1H), 6.60 (d, 1H), 6.47 (m, 1H), 4.53 (m, 1H), 3.30 (m, 2H), 3.06 (m, 4H), 2.78 (s, 2H), 2.27 (m, 3H), 2.19-2.10 (m, 5H), 1.98 (m, 3H), 1.85-1.66 (m, 4H), 1.39 (t, 2H), 0.94 (s, 6H).

50

55

EXAMPLE 159

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[(1R,3S)-3-hydroxycyclopentyl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 159A

4-(((1R,3S)-3-hydroxycyclopentyl)methylamino)-3-nitrobenzenesulfonamide

[0757] The title compound was prepared by substituting (1R, 3S)-3-hydroxycyclopentyl)methylamine for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 159B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[(1R,3S)-3-hydroxycyclopentyl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0758] The title compound was prepared by substituting EXAMPLE 158A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.02 (s, 1H), 9.28 (d, 1H), 8.59 (t, 1H), 8.44 (d, 1H), 8.29 (d, 1H), 8.13 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.08 (d, 2H), 6.82 (dd, 1H), 6.74 (d, 1H), 6.55 (d, 1H), 6.48 (m, 1H), 4.53 (m, 1H), 3.34 (m, 2H), 3.06 (m, 4H), 2.77 (s, 2H), 2.27 (m, 3H), 2.19-2.10 (m, 5H), 1.97 (m, 3H), 1.85-1.66 (m, 4H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 160

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-({[(3S)-2-oxopiperidin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0759] The title compound was prepared by substituting (S)-3-aminopiperidin-2-one for 1-acetylpiperidin-4-amine in EXAMPLE 53B. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.68 (br s, 1H), 8.88 (d, 1H), 8.57 (d, 1H), 8.04 (d, 1H), 7.95 (br s, 1H), 7.83 (dd, 1H), 7.55-7.46 (m, 3H), 7.35 (d, 2H), 7.16 (d, 1H), 7.05 (d, 2H), 6.68 (dd, 1H), 6.38 (m, 1H), 6.21 (d, 1H), 4.41 (m, 1H), 3.22 (m, 2H), 3.09 (br s, 4H), 2.78 (br s, 2H), 2.35-2.09 (m, 8H), 1.96 (br s, 2H), 1.86 (m, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 161

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[[{1-[2-fluoro-1-(fluoromethyl)ethyl]azetidin-3-yl}methyl]amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 161A

tert-butyl 3-((4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)sulfamoyl)-2-nitrophenylamino)methyl)azetidine-1-carboxylate

[0760] EXAMPLE 82 (305 mg), tert-butyl 3-(aminomethyl)azetidine-1-carboxylate (86 mg) and diisopropylamine (0.202 mL) in dioxane (3 mL) were heated to 110°C. After stirring overnight, the reaction was concentrated. Silica gel chromatography (Reveleris, 12 g) eluting with a gradient of 0.5% to 3% methanol/dichloromethane (Flow = 36 ml/minute) gave the title compound.

EXAMPLE 161B

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-(4-(azetidin-3-ylmethylamino)-3-nitrophenylsulfonyl)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzamide

[0761] To a solution of EXAMPLE 161A (0.257 g) in dichloromethane (5 mL) was added trifluoroacetic acid (0.211 mL). After 30 minutes an additional 0.2 mL of trifluoroacetic acid was added. After 3 hours, the reaction was concentrated to give the title compound.

EXAMPLE 161C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[[{1-[2-fluoro-1-(fluoromethyl)ethyl]azetidin-3-yl}methyl]amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0762] A solution of EXAMPLE 161B (0.118 g), sodium triacetoxyborohydride (0.035 g) and 1,3-difluoropropan-2-one (0.012 g) were stirred together in dichloromethane (1 mL) overnight. The reaction was quenched with saturated aqueous NaHCO₃ solution (10 mL) and extracted into dichloromethane (30 mL). The organic layer was dried and concentrated. Silica gel chromatography (Reveleris 12 g) eluting with a gradient of 0.5% to 3.5% methanol/dichloromethane over 30 minutes (Flow = 36 ml/min) gave the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 11.47 - 11.21 (m, 1H), 8.85 (s, 1H), 8.55 (d, 1H), 8.03 (d, 1H), 7.80 (dd, 1H), 7.54 - 7.45 (m, 3H), 7.33 (s, 2H), 7.04 (d, 3H), 6.67 (d, 1H), 6.38 (dd, 1H), 6.20 (d, 1H), 4.43 (dt, 4H), 3.56 (t, 2H), 3.46 (s, 2H), 3.12 (m, 6H), 2.74 (m, 3H), 2.17 (m, 7H), 1.95 (s, 2H), 1.39 (d, 2H), 0.92 (s, 6H).

EXAMPLE 162

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-[[1-oxetan-3-yl]azetidin-3-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0763] The title compound was prepared by substituting oxetan-3-one for 1,3-difluoropropan-2-one in EXAMPLE 161C. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1H), 11.51 - 11.03 (m, 1H), 8.81 (s, 1H), 8.54 (d, 1H), 8.02 (d, 1H), 7.79 (dd, 1H), 7.50 (dd, 3H), 7.34 (d, 2H), 7.04 (d, 3H), 6.67 (d, 1H), 6.38 (dd, 1H), 6.20 (d, 1H), 4.57 (s, 2H), 4.43 - 4.35 (m, 2H), 3.82 (s, 1H), 3.59 (t, 2H), 3.44 (t, 2H), 3.20 (s, 2H), 3.06 (s, 4H), 2.73 (s, 3H), 2.18 (s, 6H), 1.95 (s, 2H), 1.39 (d, 2H), 0.92 (s, 6H).

EXAMPLE 163

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-[[1-oxetan-3-yl]piperidin-4-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 163A

tert-butyl 4-((4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)sulfamoyl)-2-nitrophenylamino)methyl)piperidine-1-carboxylate

[0764] The title compound was prepared by substituting tert-butyl 4-(aminomethyl)piperidine-1-carboxylate for 1-acetyl-piperidin-4-amine in EXAMPLE 53B.

EXAMPLE 163B

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-(3-nitro-4-(piperidin-4-ylmethylamino)phenyl)sulfonyl]benzamide

[0765] The title compound was prepared by substituting EXAMPLE 163A for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 163C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-[[1-oxetan-3-yl]piperidin-4-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0766] The title compound was prepared by substituting EXAMPLE 163B for EXAMPLE 161B and oxetan-3-one for 1,3-difluoropropan-2-one in EXAMPLE 161C. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 8.60 (t, 1H), 8.54 (d, 1H), 8.03 (d, 1H), 7.79 (dd, 1H), 7.50 (m, 3H), 7.34 (d, 2H), 7.09 (d, 1H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.39 (m, 1H), 6.19 (d, 1H), 4.55 (t, 2H), 4.46 (t, 2H), 3.52 (br s, 1H), 3.28 (m, 2H), 3.17 (d, 1H), 3.06 (m, 4H), 2.82 (m, 2H), 2.74 (m, 2H), 2.17 (m, 6H), 1.95 (m, 3H), 1.72 (m, 3H), 1.38 (t, 2H), 1.28 (m, 2H), 0.92 (s, 6H).

EXAMPLE 164

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[1-cyclopropylpiperidin-4-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

5

[0767] The title compound was prepared by substituting EXMAPLE 163B for (S)-tert-butyl pyrrolidin-3-ylcarbamate in EXAMPLE 142A. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.96 (br s, 1H), 11.62 (br s, 1H), 8.50 (m, 2H), 7.98 (d, 1H), 7.72 (m, 1H), 7.52 (d, 1H), 7.45 (m, 2H), 7.34 (d, 2H), 7.04 (m, 2H), 6.94 (m, 1H), 6.64 (dd, 1H), 6.34 (m, 1H), 6.22 (d, 1H), 3.28 (m, 3H), 3.04 (m, 5H), 2.72 (s, 2H), 2.64 (m, 1H), 2.64 (m, 1H), 2.36 (m, 1H), 2.16 (m, 7H), 1.95 (s, 2H), 1.68 (m, 3H), 1.38 (t, 2H), 1.18 (m, 3H), 0.94 (s, 6H), 0.35 (m, 3H).

10

EXAMPLE 165

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[4-(2-fluoroethyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

15

EXAMPLE 165A

4-((4-(2-fluoroethyl)morpholin-2-yl)methylamino)-3-nitrobenzenesulfonamide

20

[0768] The title compound was prepared by substituting 2-fluoro-ethyl bromide for methyl iodide in EXAMPLE 134B.

EXAMPLE 165B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[4-(2-fluoroethyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

25

[0769] The title compound was prepared by substituting EXAMPLE 165A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.26 (d, 1H), 8.87 (t, 1H), 8.43 (d, 1H), 8.32 (dd, 1H), 8.11 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.92 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (dd, 1H), 3.93 (m, 1H), 4.63, 4.51 (dt, 2H), 3.95-3.85 (m, 2H), 3.68 (dt, 1H), 3.43-3.37 (m, 2H), 3.07 (m, 4H), 2.92 (d, 1H), 2.77 (s, 2H), 2.65 (m, 2H), 2.59 (m, 1H), 2.26 (m, 2H), 2.17-2.08 (m, 5H), 1.97 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

30

EXAMPLE 166

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[4-(2,2-difluoroethyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

35

EXAMPLE 166A

4-((4-(2,2-difluoroethyl)morpholin-2-yl)methylamino)-3-nitrobenzenesulfonamide

40

[0770] The title compound was prepared by substituting 2,2-difluoro-ethyl bromide for methyl iodide in EXAMPLE 134B.

EXAMPLE 166B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[4-(2,2-difluoroethyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

45

[0771] The title compound was prepared by substituting EXAMPLE 166A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.01 (s, 1H), 9.26 (d, 1H), 8.86 (t, 1H), 8.43 (d, 1H), 8.34 (dd, 1H), 8.11 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.93 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 6.31, 6.20, 6.09 (tt, 1H), 3.90 (m, 1H), 3.85 (d, 1H), 3.67 (dt, 1H), 3.49-3.30 (m, 2H), 3.07 (m, 4H), 2.84 (d, 1H), 2.82-2.75 (m, 4H), 2.69 (d, 1H), 2.33 (dt, 1H), 2.27-2.20 (m, 3H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

50

55

EXAMPLE 167

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-fluoro-1-oxetan-3-yl)piperidin-4-yl]methoxy}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 167A

4-((4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methoxy)-3-nitrobenzenesulfonamide

[0772] The title compound was prepared by substituting EXAMPLE 173A for tert-butyl piperazine-1-carboxylate and 3-oxetanone for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 167B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-fluoro-1-oxetan-3-yl)piperidin-4-yl]methoxy}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0773] The title compound was prepared by substituting EXAMPLE 167A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 8.39 (s, 1H), 8.09 (d, 1H), 8.04 (d, 1H), 7.52 (m, 4H), 7.35 (d, 2H), 7.05 (m, 2H), 6.68 (dd, 1H), 6.40 (dd, 1H), 6.20 (s, 1H), 4.57 (t, 2H), 4.48 (m, 2H), 4.38 (d, 2H), 4.02 (m, 1H), 3.63 (m, 2H), 3.08 (m, 4H), 2.74 (m, 4H), 2.17 (m, 6H), 1.88 (m, 6H), 1.40 (t, 2H), 0.93 (s, 6H).

EXAMPLE 168

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(2S)-4,4-difluoro-1-oxetan-3-ylpyrrolidin-2-yl]methoxy}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 168A

(S)-methyl 4,4-difluoropyrrolidine-2-carboxylate

[0774] (S)-1-tert-butyl 2-methyl 4,4-difluoropyrrolidine-1,2-dicarboxylate (0.472 g) in CH₂Cl₂ (1 mL) was treated with trifluoroacetic acid (1.4 mL), stirred at ambient temperature for 4 hours, and concentrated. The product was free-based using a MEGA BE-SCX column with 1:1 CH₂Cl₂/methanol as eluent for the trifluoroacetic acid. The product was released from the column with 5% (7 M ammonia in methanol) in CH₂Cl₂ as eluent.

EXAMPLE 168B

(S)-methyl 4,4-difluoro-1-(oxetan-3-yl)pyrrolidine-2-carboxylate

[0775] The title compound was prepared by substituting EXAMPLE 168A for tert-butyl piperazine-1-carboxylate and 3-oxetanone for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 168C

(S)-(4,4-difluoro-1-(oxetan-3-yl)pyrrolidin-2-yl)methanol

[0776] EXAMPLE 168B (0.180 g) in tetrahydrofuran (3 mL) was treated sequentially with a solution of calcium chloride (0.245 g) in ethanol (3 mL) and NaBH₄ (0.167 g) and then stirred at ambient temperature for 7 hours. The reaction was quenched with saturated aqueous NH₄Cl solution and extracted with ethyl acetate. The combined extracts were washed with brine, dried (MgSO₄), filtered, concentrated and chromatographed on silica gel with 50% ethyl acetate in hexanes as eluent to give the product.

EXAMPLE 168D

(S)-4-((4,4-difluoro-1-(oxetan-3-yl)pyrrolidin-2-yl)methoxy)-3-nitrobenzenesulfonamide

[0777] The title compound was prepared by substituting EXAMPLE 168C for (tetrahydro-2H-pyran-4-yl)methanol in

EXAMPLE 24A.

EXAMPLE 168E

5 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([(2S)-4,4-difluoro-1-oxetan-3-ylpyrrolidin-2-yl]methoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 **[0778]** The title compound was prepared by substituting EXAMPLE 168D for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 8.38 (s, 1H), 8.06 (m, 2H), 7.49 (m, 4H), 7.35 (d, 2H), 7.05 (d, 2H), 6.68 (dd, 1H), 6.40 (m, 1H), 6.21 (s, 1H), 4.54 (m, 3H), 4.43 (t, 1H), 4.23 (m, 1H), 4.12 (m, 2H), 3.44 (m, 2H), 3.12 (m, 7H), 2.58 (m, 1H), 2.29 (m, 7H), 1.97 (s, 2H), 1.40 (t, 2H), 0.93 (s, 6H).

EXAMPLE 169

15 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([(4-tetrahydro-2H-pyran-4-ylmorpholin-3-yl)methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 169A

20 tert-butyl 3-((4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)sulfamoyl)-2-nitrophenylamino)methyl)morpholine-4-carboxylate

25 **[0779]** The title compound was prepared as described in EXAMPLE 53B by replacing 1-acetylpiperidin-4-amine with tert-butyl 3-(aminomethyl)morpholine-4-carboxylate.

EXAMPLE 169B

30 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-(4-(morpholin-3-ylmethylamino)-3-nitrophenyl)sulfonylbenzamide

[0780] The title compound was prepared as described in EXAMPLE 139B by replacing EXAMPLE 139A with EXAMPLE 169A.

EXAMPLE 169C

35 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([(4-tetrahydro-2H-pyran-4-ylmorpholin-3-yl)methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

40 **[0781]** The title compound was prepared as described in EXAMPLE 139C by replacing EXAMPLE 139B and oxetan-3-one with EXAMPLE 169B and tetrahydropyran-4-one, respectively. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 8.77 (m, 1H), 8.57 (d, 1H), 8.05 (d, 1H), 7.84 (dd, 1H), 7.52 (m, 3H), 7.34 (m, 2H), 7.03 (m, 3H), 6.68 (dd, 1H), 6.40 (dd, 1H), 6.18 (d, 1H), 3.86 (m, 2H), 3.72 (m, 2H), 3.11 (m, 6H), 2.74 (m, 4H), 2.20 (m, 6H), 1.95 (m, 3H), 1.51 (m, 7H), 0.92 (s, 6H).

45 EXAMPLE 170

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([(4-cyclobutylmorpholin-3-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50 **[0782]** The title compound was prepared as described in EXAMPLE 139C by replacing EXAMPLE 139B and oxetan-3-one with EXAMPLE 169B and cyclobutanone. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 8.72 (s, 1H), 8.57 (d, 1H), 8.04 (d, 1H), 7.84 (dd, 1H), 7.52 (m, 3H), 7.34 (m, 3H), 7.03 (m, 4H), 6.67 (dd, 1H), 6.39 (dd, 1H), 6.18 (d, 1H), 3.47 (m, 3H), 3.10 (m, 6H), 2.72 (m, 6H), 2.25 (m, 8H), 1.95 (m, 4H), 1.56 (m, 3H), 1.38 (m, 2H), 0.92 (s, 6H).

55

EXAMPLE 171

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-[(4-tetrahydrofuran-3-yl)morpholin-3-yl)methyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0783] The title compound was prepared as described in EXAMPLE 139C by replacing EXAMPLE 139B and oxetan-3-one with EXAMPLE 169B and 3-oxotetrahydrofuran, respectively. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.64 (s, 1H), 8.66 (s, 1H), 8.53 (d, 1H), 8.01 (d, 1H), 7.80 (d, 1H), 7.50 (m, 3H), 7.35 (d, 2H), 7.05 (d, 2H), 6.98 (d, 1H), 6.66 (dd, 1H), 6.37 (d, 1H), 6.19 (d, 1H), 3.68 (m, 8H), 3.05 (m, 6H), 2.85 (m, 3H), 2.73 (s, 2H), 2.25 (m, 6H), 1.91 (m, 3H), 1.37 (m, 3H), 0.95 (m, 6H).

EXAMPLE 172

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl)methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0784] The title compound was prepared by substituting EXAMPLE 163B for tert-butyl piperazine-1-carboxylate and 1,3-difluoropropan-2-one for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 11.40 (br s, 1H), 8.57 (m, 2H), 8.03 (d, 1H), 7.78 (d, 1H), 7.50 (m, 3H), 7.34 (d, 2H), 7.07 (d, 1H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.38 (m, 1H), 6.19 (d, 1H), 4.63 (d, 2H), 4.53 (d, 2H), 3.28 (m, 2H), 3.07 (m, 4H), 2.89 (m, 2H), 2.74 (m, 2H), 2.40 (m, 2H), 2.16 (m, 6H), 1.95 (s, 2H), 1.67 (m, 3H), 1.38 (t, 2H), 1.23 (m, 3H), 0.94 (s, 6H).

EXAMPLE 173

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1-cyclopropyl-4-fluoropiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 173A

4-((4-fluoropiperidin-4-yl)methoxy)-3-nitrobenzenesulfonamide

[0785] The title compound was prepared by substituting EXAMPLE 126B for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 173B

4-((1-cyclopropyl-4-fluoropiperidin-4-yl)methoxy)-3-nitrobenzenesulfonamide

[0786] To EXAMPLE 173A (0.24 g) in methanol (3 mL) was added 3Å molecular sieves (0.1 g), followed sequentially by acetic acid (0.31 mL), (1-ethoxycyclopropoxy)trimethylsilane (0.64 mL), and sodium cyanoborohydride (0.148 g). The reaction was heated under reflux overnight. After cooling, the reaction mixture was loaded onto a silica gel column. After drying, the column was eluted with 100:2:0.2 ethyl acetate/methanol/NH₄OH to give the title compound.

EXAMPLE 173C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1-cyclopropyl-4-fluoropiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0787] The title compound was prepared by substituting EXAMPLE 173B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1H), 8.33 (s, 1H), 8.01 (m, 2H), 7.53 (d, 1H), 7.48-7.49 (m, 2H), 7.34-7.38 (m, 3H), 7.04 (d, 2H), 6.66 (dd, 1H), 6.38 (dd, 1H), 6.21 (d, 1H), 4.32 (d, 2H), 3.70-3.77 (m, 2H), 3.07 (s, 4H), 2.92 (s, 2H), 2.80 (s, 2H), 2.58 (s, 2H), 2.25 (s, 4H), 2.13-2.16 (m, 2H), 1.38 (t, 2H), 0.92 (s, 6H), 0.40-0.49 (m, 4H).

EXAMPLE 174

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-methoxybenzyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0788] A suspension of EXAMPLE 53A (120 mg), (4-methoxyphenyl)methanamine (31 mg) and Hunig's Base (0.159

mL) in dimethylsulfoxide (2 mL) was heated for 2 hours at 150 °C in a Biotage Initiator microwave reactor. The reaction mixture was diluted with methanol (2 mL) and purified by reverse phase HPLC (C8, 30% - 100% CH₃CN/water/0.1% trifluoroacetic acid). ¹H NMR (500 MHz, pyridine-d₅) δ 13.07 (s, 1H), 9.32 (d, 1H), 9.17 (t, 1H), 8.43 (d, 1H), 8.28 (dd, 1H), 8.08 (d, 1H), 7.64 - 7.68 (m, 2H), 7.44 (d, 2H), 7.38 (d, 2H), 7.07 (d, 2H), 6.97 - 7.02 (m, 2H), 6.90 (d, 1H), 6.74 (dd, 1H), 6.52 (d, 1H), 6.49 (dd, 1H), 4.55 (d, 2H), 3.68 (s, 3H), 3.03 - 3.09 (m, 4H), 2.77 (s, 2H), 2.26 (t, 2H), 2.10 - 2.17 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 175

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[3-(trifluoromethoxy)benzyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0789] The title compound was prepared by substituting (3-trifluoromethoxyphenyl)methanamine for (4-methoxyphenyl)methanamine in EXAMPLE 174. ¹H NMR (500 MHz, pyridine-d₅) δ 13.06 (s, 1H), 9.38 (t, 1H), 9.31 (d, 1H), 8.42 (d, 1H), 8.28 (dd, 1H), 8.08 (d, 1H), 7.65 (ddd, 2H), 7.41 - 7.46 (m, 3H), 7.36 - 7.40 (m, 2H), 7.07 (d, 2H), 6.88 (d, 1H), 6.74 (dd, 1H), 6.52 (d, 1H), 6.49 (d, 1H), 4.73 (d, 2H), 3.02 - 3.08 (m, 4H), 2.77 (s, 2H), 2.22 - 2.28 (m, 2H), 2.09 - 2.16 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 176

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(3-methoxybenzyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0790] The title compound was prepared by substituting (3-methoxyphenyl)methanamine for (4-methoxyphenyl)methanamine in EXAMPLE 174. ¹H NMR (500 MHz, pyridine-d₅) δ 13.06 (s, 1H), 9.27 - 9.32 (m, 2H), 8.42 (d, 1H), 8.26 (dd, 1H), 8.08 (d, 1H), 7.64 - 7.67 (m, 2H), 7.44 (d, 2H), 7.32 (t, 1H), 7.14 (s, 1H), 7.04 - 7.09 (m, 3H), 6.88 - 6.94 (m, 2H), 6.74 (dd, 1H), 6.52 (d, 1H), 6.48 - 6.50 (m, 1H), 4.64 (d, 2H), 3.68 (s, 3H), 3.03 - 3.09 (m, 4H), 2.77 (s, 2H), 2.26 (t, 2H), 2.10 - 2.18 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 177

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[4-(difluoromethoxy)benzyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0791] The title compound was prepared by substituting (4-difluoromethoxyphenyl)methanamine for (4-methoxyphenyl)methanamine in EXAMPLE 174. ¹H NMR (500 MHz, pyridine-d₅) δ 13.06 (s, 1H), 9.32 (d, 1H), 9.28 (t, 1H), 8.42 (d, 1H), 8.28 (dd, 1H), 8.07 (d, 1H), 7.66 (t, 1H), 7.64 (d, 1H), 7.58 (s, 1H), 7.44 (s, 2H), 7.26 (s, 1H), 7.25 (d, 1H), 7.07 (d, 2H), 6.87 (d, 1H), 6.74 (dd, 1H), 6.52 (d, 1H), 6.49 (dd, 1H), 4.64 (d, 2H), 3.03 - 3.10 (m, 4H), 2.77 (s, 2H), 2.26 (t, 2H), 2.11 - 2.17 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 178

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-(1,4-dioxaspiro[4.5]dec-8-ylamino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0792] The title compound was prepared by substituting 1,4-dioxaspiro[4.5]dec-8-ylamine for 1-acetylpiperidin-4-amine in EXAMPLE 53B. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.67 (br s, 1H), 8.55 (d, 1H), 8.26 (d, 1H), 8.04 (d, 1H), 7.81 (dd, 1H), 7.54-7.46 (m, 3H), 7.35 (d, 2H), 7.15 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.38 (m, 1H), 6.19 (d, 1H), 3.89 (s, 4H), 3.78 (m, 1H), 3.07 (br s, 4H), 2.78 (br s, 2H), 2.28-2.11 (m, 6H), 2.00-1.88 (m, 4H), 1.75-1.57 (m, 4H), 1.54-1.35 (m, 4H), 0.92 (s, 6H).

EXAMPLE 179

Trans-N-[(4-[[4-(acetylamino)cyclohexyl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 179A

tert-butyl trans-4-acetamidocyclohexylcarbamate

[0793] Tert-butyl (trans)-4-aminocyclohexylcarbamate (1.500 g) and triethylamine (2.93 mL, 2.125 g) were added to dichloromethane and stirred until the tert-butyl (trans)-4-aminocyclohexylcarbamate had dissolved completely. Acetyl chloride (0.577 g) was added slowly, and the solution was stirred at room temperature for 16 hours. The solvent was removed, and the residue taken up in ethyl acetate, washed with pH 4 buffer, washed with brine, dried with anhydrous sodium sulfate, and filtered. The filtrate was concentrated under vacuum.

EXAMPLE 179B

N-(trans-4-aminocyclohexyl)acetamide

[0794] The title compound was prepared by substituting EXAMPLE 179A for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 179C

Trans-N-[(4-[[4-(acetylamino)cyclohexyl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0795] The title compound was prepared by substituting EXAMPLE 179B for 1-acetyl piperidin-4-amine in EXAMPLE 53B. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.67 (br s, 1H), 8.55 (d, 1H), 8.20 (d, 1H), 8.04 (d, 1H), 7.82-7.76 (m, 2H), 7.53-7.46 (m, 3H), 7.35 (d, 2H), 7.16 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.39 (m, 1H), 6.19 (d, 1H), 3.57 (m, 2H), 3.07 (br s, 4H), 2.75 (br s, 2H), 2.28-2.10 (m, 6H), 2.03-1.94 (m, 4H), 1.83 (d, 2H), 1.80 (s, 3H), 1.55-1.24 (m, 6H), 0.92 (s, 6H).

EXAMPLE 180

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 180A

(R)-tert-butyl 1-(2,2-difluoroethyl)pyrrolidin-3-ylcarbamate

[0796] To a solution of (R)-tert-butyl pyrrolidin-3-ylcarbamate (500 mg) and 1,1-difluoro-2-iodoethane (618 mg) in N,N-dimethylformamide (6 mL) was added N-ethyl-N-isopropylpropan-2-amine (1.403 mL) and the mixture was stirred at 70 °C for 72 hours. The reaction mixture was concentrated and the crude product was purified on silica gel with methanol / dichloromethane.

EXAMPLE 180B

(R)-1-(2,2-difluoroethyl)pyrrolidin-3-amine

[0797] To a solution of EXAMPLE 180A (525 mg) in a mixture of dichloromethane (3 mL) and methanol (4.0 mL) was added hydrogen chloride, 4M in dioxane (5.24 mL) and the reaction was stirred for 1.5 hours. The reaction was concentrated and the crude material was taken up in dichloromethane and the solvent evaporated, then taken up in ether and the solvent evaporated, and then dried on high vacuum.

EXAMPLE 180C

(R)-4-(1-(2,2-difluoroethyl)pyrrolidin-3-ylamino)-3-nitrobenzenesulfonamide

- 5 **[0798]** The title compound was prepared by substituting EXAMPLE 180B for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 180D

- 10 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0799]** The title compound was prepared by substituting EXAMPLE 180C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500 MHz, pyridine- d₅) δ 13.02 (m, 1H), 9.27 (d, 1H), 8.55 (d, 1H), 8.43 (d, 1H), 8.35 (dd, 1H), 8.10 (d, 1H), 7.64 - 7.68 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.83 (d, 1H), 6.75 (dd, 1H), 6.54 (m, 1H), 6.48 (m, 1H), 6.04 - 6.29 (m, 1H), 4.06 (m, 1H), 3.07 (m, 4H), 2.83 - 2.95 (m, 4H), 2.74 - 2.82 (m, 3H), 2.47 (m, 1H), 2.09 - 2.30 (m, 8H), 1.97 (s, 2H), 1.67 (m, 1H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 181

- 20 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((3S)-1-(2-fluoroethyl)pyrrolidin-3-yl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 181A

- 25 (S)-tert-butyl 1-(2-fluoroethyl)pyrrolidin-3-ylcarbamate

- [0800]** The title compound was prepared by substituting 1-fluoro-2-iodoethane for 1,1-difluoro-2-iodoethane and (S)-tert-butyl pyrrolidin-3-ylcarbamate for (R)-tert-butyl pyrrolidin-3-ylcarbamate in EXAMPLE 180A.

EXAMPLE 181B

(S)-1-(2-fluoroethyl)pyrrolidin-3-amine

- 35 **[0801]** The title compound was prepared by substituting EXAMPLE 181A for EXAMPLE 180A in EXAMPLE 180B.

EXAMPLE 181C

(S)-4-(1-(2-fluoroethyl)pyrrolidin-3-ylamino)-3-nitrobenzenesulfonamide

- 40 **[0802]** The title compound was prepared by substituting EXAMPLE 181B for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 181D

- 45 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((3S)-1-(2-fluoroethyl)pyrrolidin-3-yl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0803]** The title compound was prepared by substituting EXAMPLE 181C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500 MHz, pyridine- d₅) δ 13.00 (m, 1H), 9.26 (d, 1H), 8.56 (d, 1H), 8.43 (d, 1H), 8.34 (dd, 1H), 8.10 (d, 1H), 7.63 - 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.82 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 4.60 (t, 1H), 4.51 (t, 1H), 4.05 (m, 1H), 3.07 (m, 4H), 2.84 (m, 1H), 2.66 - 2.79 (m, 6H), 2.39 (q, 1H), 2.20 - 2.29 (m, 3H), 2.15 (m, 5H), 1.97 (s, 2H), 1.66 (m, 1H), 1.39 (t, 2H), 0.94 (s, 6H).

55

EXAMPLE 182

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3S)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 182A

(S)-tert-butyl 1-(2,2-difluoroethyl)pyrrolidin-3-ylcarbamate

[0804] The title compound was prepared by substituting (S)-tert-butyl pyrrolidin-3-ylcarbamate for (R)-tert-butyl pyrrolidin-3-ylcarbamate in EXAMPLE 180A.

EXAMPLE 182B

(S)-1-(2,2-difluoroethyl)pyrrolidin-3-amine

[0805] The title compound was prepared by substituting EXAMPLE 182A for EXAMPLE 180A in EXAMPLE 180B.

EXAMPLE 182C

(S)-4-(1-(2,2-difluoroethyl)pyrrolidin-3-ylamino)-3-nitrobenzenesulfonamide

[0806] The title compound was prepared by substituting EXAMPLE 182B for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 182D

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3S)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0807] The title compound was prepared by substituting EXAMPLE 182C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500 MHz, pyridine- d₅) δ 13.02 (m, 1H), 9.27 (d, 1H), 8.54 (d, 1H), 8.43 (d, 1H), 8.35 (dd, 1H), 8.11 (d, 1H), 7.64 - 7.68 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.82 (d, 1H), 6.75 (dd, 1H), 6.54 (m, 1H), 6.48 (m, 1H), 6.04 - 6.29 (m, 1H), 4.06 (m, 1H), 3.07 (m, 4H), 2.83 - 2.95 (m, 4H), 2.74 - 2.82 (m, 3H), 2.47 (m, 1H), 2.09 - 2.30 (m, 8H), 1.97 (s, 2H), 1.67 (m, 1H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 183

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(2-fluoroethyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,b]pyridin-5-yloxy)benzamide

EXAMPLE 183A

(R)-tert-butyl 1-(2-fluoroethyl)pyrrolidin-3-ylcarbamate

[0808] The title compound was prepared by substituting 1-fluoro-2-iodoethane for 1,1-difluoro-2-iodoethane in EXAMPLE 180A.

EXAMPLE 183B

(R)-1-(2-fluoroethyl)pyrrolidin-3-amine

[0809] The title compound was prepared by substituting EXAMPLE 183A for EXAMPLE 180A in EXAMPLE 180B.

EXAMPLE 183C

(R)-4-(1-(2-fluoroethyl)pyrrolidin-3-ylamino)-3-nitrobenzenesulfonamide

- 5 **[0810]** The title compound was prepared by substituting EXAMPLE 183B for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 183D

- 10 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(2-fluoroethyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0811]** The title compound was prepared by substituting EXAMPLE 183C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500 MHz, pyridine-*d*₅) δ 13.00 (m, 1H), 9.26 (d, 1H), 8.56 (d, 1H), 8.43 (d, 1H), 8.34 (dd, 1H), 8.10 (d, 1H), 7.63 - 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.83 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 4.60 (t, 1H), 4.50 (t, 1H), 4.04 (m, 1H), 3.07 (m, 4H), 2.84 (m, 1H), 2.66 - 2.79 (m, 6H), 2.39 (q, 1H), 2.19 - 2.28 (m, 3H), 2.14 (m, 5H), 1.97 (s, 2H), 1.66 (m, 1H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 184

- 20 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(3S)-1-oxetan-3-yl]pyrrolidin-3-yl}methoxy}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 184A

- 25 (S)-tert-butyl 3-((2-nitro-4-sulfamoylphenoxy)methyl)pyrrolidine-1-carboxylate

- [0812]** To a solution of (S)-tert-butyl 3-(hydroxymethyl)pyrrolidine-1-carboxylate (0.300 g) in tetrahydrofuran (5 mL) was added sodium hydride (0.238 g). After stirring for 15 minutes, 4-fluoro-3-nitrobenzenesulfonamide (0.295 g) was added and reaction stirred at room temperature. After 1 hour, the reaction was partitioned between water (25 mL) and dichloromethane (50 mL) and the reaction quenched with 1N aqueous HCl (5.96 mL). The organic layer was separated, dried over magnesium sulfate, filtered, and concentrated. Silica gel chromatography (Reveleris 12 g) eluting with a gradient of 0.2% to 2% methanol/dichloromethane over 30 minutes (flow = 36 mL/minute) gave the title compound.

EXAMPLE 184B

(S)-3-nitro-4-((1-(oxetan-3-yl)pyrrolidin-3-yl)methoxy)benzenesulfonamide

- [0813]** To (S)-tert-butyl 3-((2-nitro-4-sulfamoylphenoxy)methyl)pyrrolidine-1-carboxylate (0.433 g) was added hydrogen chloride (4.0M in dioxane, 1.0 mL). After stirring for 1 hour, the reaction was concentrated and partitioned between dichloromethane (50 mL) and saturated aqueous NaHCO₃ solution (50 mL). The aqueous layer was separated and concentrated. The residue was triturated with methanol (100 mL), filtered and concentrated and treated with sodium cyanoborohyde (0.068 g) and cyclobutanone (0.078 g) and stirred overnight. The reaction was partitioned between dichloromethane (50 mL) and water (25 mL) and saturated aqueous NaHCO₃ (10 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated to give the title compound.

EXAMPLE 184C

- 50 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(3S)-1-oxetan-3-yl]pyrrolidin-3-yl}methoxy}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0814]** The title compound was prepared by substituting EXAMPLE 184B for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-*d*₆) δ 11.64 (s, 1H), 11.45 - 11.01 (m, 1H), 8.30 (d, 1H), 7.98 (dd, 2H), 7.60 - 7.43 (m, 3H), 7.33 (t, 3H), 7.04 (d, 2H), 6.74 - 6.59 (m, 1H), 6.37 (dd, 1H), 6.21 (d, 1H), 4.49 (td, 2H), 4.33 (s, 1H), 4.13 (dd, 2H), 3.79 (s, 2H), 3.44 (dd, 2H), 3.07 (s, 4H), 2.74 (d, 6H), 2.19 (d, 6H), 1.98 (d, 2H), 1.74 - 1.52 (m, 1H), 1.39 (t, 2H), 0.92 (s, 6H).

EXAMPLE 185

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-hydroxybenzyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

5

[0815] The title compound was prepared by substituting (4-hydroxyphenyl)methanamine for (4-methoxyphenyl)methanamine in EXAMPLE 174. ¹H NMR (500 MHz, pyridine-d₅) δ 13.06 (s, 1H), 11.67 (bs, 1H), 9.32 (d, 1H), 9.14 (s, 1H), 8.44 (d, 1H), 8.28 (dd, 1H), 8.09 (d, 1H), 7.65 - 7.68 (m, 2H), 7.44 (d, 2H), 7.37 - 7.41 (m, 2H), 7.19 (s, 2H), 7.07 (d, 2H), 6.93 (d, 1H), 6.75 (dd, 1H), 6.53 (d, 1H), 6.49 (dd, 1H), 4.54 (d, 2H), 3.02 - 3.09 (m, 4H), 2.77 (s, 2H), 2.22 - 2.29 (m, 2H), 2.10 - 2.17 (m, 4H), 1.97 (d, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

10

EXAMPLE 186

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(3-hydroxybenzyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

15

[0816] The title compound was prepared by substituting (3-hydroxyphenyl)methanamine for (4-methoxyphenyl)methanamine in EXAMPLE 174. ¹H NMR (500 MHz, pyridine-d₅) δ 13.06 (s, 1H), 11.67 (bs, 1H), 9.27 - 9.32 (m, 2H), 8.43 (d, 1H), 8.20 (dd, 1H), 8.08 (d, 1H), 7.66 (t, 2H), 7.44 (d, 2H), 7.33 (t, 1H), 7.25 (s, 1H), 7.13 (dd, 1H), 7.07 (d, 2H), 6.98 (d, 1H), 6.88 (d, 1H), 6.74 (dd, 1H), 6.52 (d, 1H), 6.49 (dd, 1H), 4.64 (d, 2H), 3.02 - 3.09 (m, 4H), 2.77 (s, 2H), 2.22 - 2.28 (m, 2H), 2.09 - 2.16 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

20

EXAMPLE 187

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(3-(difluoromethoxy)benzyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

25

[0817] The title compound was prepared by substituting (3-difluoromethoxyphenyl)methanamine for (4-methoxyphenyl)methanamine in EXAMPLE 174. ¹H NMR (500 MHz, pyridine-d₅) δ 13.06 (s, 1H), 9.34 (t, 1H), 9.30 (d, 1H), 8.42 (d, 1H), 8.26 (dd, 1H), 8.08 (d, 1H), 7.66 (ddd, 2H), 7.40 - 7.45 (m, 3H), 7.36 (t, 1H), 7.27 - 7.30 (m, 2H), 7.19 (d, 1H), 7.07 (d, 2H), 6.87 (d, 1H), 6.74 (dd, 1H), 6.52 (d, 1H), 6.49 (dd, 1H), 4.69 (d, 2H), 3.02 - 3.08 (m, 4H), 2.77 (s, 2H), 2.26 (t, 2H), 2.09 - 2.16 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

30

EXAMPLE 188

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-([cis-3-morpholin-4-ylcyclopentyl]methyl)amino)-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

35

EXAMPLE 188A

Cis-methyl 3-morpholinocyclopentanecarboxylate

40

[0818] The title compound was prepared by substituting methyl 3-oxocyclopentanecarboxylate for 4'-chlorobiphenyl-2-carboxaldehyde and morpholine for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A.

45

EXAMPLE 188B

Cis-3-morpholinocyclopentyl)methanol

50

[0819] The title compound was prepared by substituting EXAMPLE 188A for EXAMPLE 101C in EXAMPLE 101D.

EXAMPLE 188C

4-((Cis-3-morpholinocyclopentyl)methoxy)-3-nitrobenzenesulfonamide

55

[0820] The title compound was prepared by substituting EXAMPLE 188B for (1,4-dioxan-2-yl)methanol in EXAMPLE 12A.

EXAMPLE 188D

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[cis-3-morpholin-4-ylcyclopentyl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0821] The title compound was prepared by substituting EXAMPLE 188C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.57 (s, 1H), 8.17 (m, 1H), 7.94 (m, 1H), 7.82 (m, 1H), 7.56 (d, 1H), 7.44 (t, 1H), 7.34 (m, 3H), 7.16 (m, 1H), 7.05 (d, 2H), 6.64 (dd, 1H), 6.33 (m, 1H), 6.24 (d, 1H), 4.06 (m, 2H), 3.62 (m, 4H), 3.03 (m, 4H), 2.75 (s, 2H), 2.35 (m, 2H), 2.19 (m, 6H), 2.03 (m, 2H), 1.96 (s, 2H), 1.78 (m, 2H), 1.51 (m, 4H), 1.39 (t, 2H), 0.93 (s, 6H)

EXAMPLE 189

Trans-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(methylsulfonyl)amino]cyclohexyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 189A

Trans-(4-Methanesulfonylamino-cyclohexyl)-carbamic acid tert-butyl ester

[0822] The title compound was prepared by substituting methanesulfonyl chloride for acetyl chloride in EXAMPLE 179A.

EXAMPLE 189B

Trans-N-(4-Aminocyclohexyl)-methanesulfonamide

[0823] The title compound was prepared by substituting EXAMPLE 189A for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 189C

Trans-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(methylsulfonyl)amino]cyclohexyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0824] The title compound was prepared by substituting EXAMPLE 189B for 1-acetyl piperidin-4-amine in EXAMPLE 53B. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.68 (br s, 1H), 8.55 (d, 1H), 8.18 (d, 1H), 8.04 (d, 1H), 7.84 (d, 1H), 7.79 (dd, 1H), 7.56-7.47 (m, 3H), 7.34 (d, 2H), 7.16 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.39 (m, 1H), 6.19 (d, 1H), 3.56 (m, 1H), 3.17 (m, 1H), 3.07 (br s, 4H), 2.93 (s, 3H), 2.75 (br s, 2H), 2.28-2.10 (m, 6H), 2.05-1.90 (m, 6H), 1.55-1.32 (m, 6H), 0.92 (s, 6H).

EXAMPLE 190

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-[(1-cyclopropylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 190A

4-(1-cyclopropylpiperidin-4-ylamino)-3-(trifluoromethylsulfonyl)benzenesulfonamide

[0825] The title compound was prepared as described in EXAMPLE 17A by replacing (tetrahydropyran-4-yl)methylamine with 4-amino-1-cyclopropylpiperidine.

EXAMPLE 190B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-[(1-cyclopropylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0826] The title compound was prepared as described in EXAMPLE 1G by replacing EXAMPLE 1E and EXAMPLE 1F with EXAMPLE 3J and EXAMPLE 190A, respectively. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1H), 8.13

(d, 1H), 8.02 (d, 1H), 7.91 (m, 1H), 7.48 (m, 3H), 7.34 (d, 2H), 7.04 (d, 2H), 6.67 (m, 2H), 6.38 (dd, 1H), 6.19 (d, 1H), 3.64 (m, 1H), 3.13 (m, 5H), 2.73 (m, 5H), 2.22 (m, 6H), 1.92 (m, 5H), 1.70 (m, 1H), 1.41 (m, 5H), 0.94 (s, 6H), 0.41 (m, 4H).

EXAMPLE 191

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([1-oxetan-3-yl]piperidin-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 191 A

3-nitro-4-(piperidin-4-ylmethoxy)benzenesulfonamide

[0827] To a solution of tert-butyl 4-(hydroxymethyl)piperidine-1-carboxylate (0.300 g) in tetrahydrofuran (5 mL) was added sodium hydride (0.223 g). After stirring for 15 minutes, 4-fluoro-3-nitrobenzenesulfonamide (0.276 g) was added and reaction stirred at room temperature. After 1 hour the reaction was partitioned between water (25 mL) and dichloromethane (50 mL) and the reaction quenched with 1N aqueous HCl (5.57 mL). The organic layer was separated, dried over magnesium sulfate, filtered, and concentrated. Treatment with HCl (4.0M in dioxane, 2 mL) and methanol (2 mL) for 1 hour, followed by concentration, trituration with dichloromethane and filtration gave the title compound.

EXAMPLE 191B

3-nitro-4-([1-(oxetan-3-yl)piperidin-4-yl]methoxy)benzenesulfonamide

[0828] To a suspension of 3-nitro-4-(piperidin-4-ylmethoxy)benzenesulfonamide (0.100 g) and cyclobutanone (0.030 g) in methanol (1 mL) was added sodium cyanoborohydride (0.027 g). After stirring overnight, the reaction was quenched with saturated NaHCO_3 (5 mL) and extracted into dichloromethane (2 x 10 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated to give the title compound.

EXAMPLE 191C

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([1-oxetan-3-yl]piperidin-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0829] The title compound was prepared by substituting EXAMPLE 191B for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ^1H NMR (300 MHz, dimethylsulfoxide- d_6) δ 11.64 (s, 1H), 11.46 - 10.46 (m, 1H), 8.29 (s, 1H), 8.00 (d, 2H), 7.61 - 7.41 (m, 3H), 7.35 (d, 3H), 7.04 (d, 2H), 6.66 (d, 1H), 6.37 (s, 1H), 6.21 (s, 1H), 4.67 - 4.40 (m, 4H), 4.08 (d, 2H), 3.06 (s, 4H), 2.78 (s, 4H), 2.19 (m, 6H), 1.96 (s, 4H), 1.79 (m, 4H), 1.39 (s, 4H), 0.93 (s, 6H).

EXAMPLE 192

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-fluoro-1-tetrahydro-2H-pyran-4-yl]piperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 192A

4-([4-fluoro-1-(tetrahydro-2H-pyran-4-yl)piperidin-4-yl]methoxy)-3-nitrobenzenesulfonamide

[0830] A mixture of EXAMPLE 173A (0.4 g), dihydro-2H-pyran-4(3H)-one (0.179 g), sodium cyanoborohydride (0.112 g), and acetic acid (0.5 mL) in tetrahydrofuran (3 mL) was stirred overnight. The solvents were removed under reduced pressure. The residue was purified with flash column chromatography on silica gel eluting with 100:5:0.5 ethyl acetate/methanol/ NH_4OH to give the desired product.

EXAMPLE 192B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-fluoro-1-tetrahydro-2H-pyran-4-yl]piperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0831] The title compound was prepared by substituting EXAMPLE 192A for EXAMPLE 11B in EXAMPLE 11D. ^1H

NMR (500MHz, dimethylsulfoxide- d_6) δ 11.58 (s, 1H), 8.25 (s, 1H), 7.96 (d, 1H), 7.93 (d, 1H), 7.57 (d, 1H), 7.45 (t, 1H), 7.34-7.37 (m, 3H), 7.26 (d, 1H), 7.05 (d, 2H), 6.64 (dd, 1H), 6.34 (dd, 1H), 6.23 (d, 1H), 4.34 (d, 2H), 3.93 (dd, 2H), 3.03 (s, 6H), 2.76 (s, 4H), 2.09-2.22 (m, 6H), 1.96 (s, 2H), 1.52-1.27 (m, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

5 EXAMPLE 193

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-fluoro-1-tetrahydrofuran-3-yl]piperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 EXAMPLE 193A

4-([4-fluoro-1-(tetrahydrofuran-3-yl)piperidin-4-yl)methoxy]-3-nitrobenzenesulfonamide

15 **[0832]** The title compound was prepared by substituting dihydrofuran-3(2H)-one for dihydro-2H-pyran-4(3H)-one in EXAMPLE 192A.

EXAMPLE 193B

20 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-fluoro-1-tetrahydrofuran-3-yl]piperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

25 **[0833]** The title compound was prepared by substituting EXAMPLE 193A for EXAMPLE 11B in EXAMPLE 11D. ^1H NMR (500MHz, dimethylsulfoxide- d_6) δ 11.63 (s, 1H), 8.31 (s, 1H), 7.99-8.00 (m, 2H), 7.54 (d, 1H), 7.46-7.48 (m, 2H), 7.34-7.35 (m, 3H), 7.05 (d, 2H), 6.66 (dd, 1H), 6.37 (dd, 1H), 6.21 (d, 1H), 4.34 (d, 2H), 3.76-3.83 (m, 3H), 3.62-3.65 (m, 2H), 3.03 (s, 4H), 2.79 (s, 4H), 2.24 (s, 2H), 2.15 (s, 2H), 1.84-1.99 (m, 8H), 1.52-1.27 (m, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 194

30 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-fluoro-1-(methylsulfonyl)piperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 194A

35 4-([4-fluoro-1-(methylsulfonyl)piperidin-4-yl)methoxy]-3-nitrobenzenesulfonamide

[0834] A mixture of EXAMPLE 173A (0.4 g), methanesulfonyl chloride (0.113 g), and triethylamine (0.64 mL) in dichloromethane (5 mL) was stirred overnight. The reaction mixture was loaded onto a silica gel column and eluted with 100:1 ethyl acetate:methanol to give the clean product.

40 EXAMPLE 194B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-fluoro-1-(methylsulfonyl)piperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

45 **[0835]** The title compound was prepared by substituting EXAMPLE 194A for EXAMPLE 11B in EXAMPLE 11D. ^1H NMR (500MHz, dimethylsulfoxide- d_6) δ 11.66 (s, 1H), 8.37 (s, 1H), 8.06 (d, 1H), 8.02 (d, 1H), 7.49-7.53 (m, 3H), 7.42 (d, 1H), 7.35 (d, 2H), 7.05 (d, 2H), 6.67 (dd, 1H), 6.38-6.39 (m, 1H), 6.21 (d, 1H), 4.40 (d, 2H), 3.51-3.54 (m, 2H), 3.09 (s, 4H), 2.96-3.01 (m, 4H), 2.92 (s, 3H), 2.82 (s, 2H), 2.25-2.34 (m, 4H), 2.13-2.16 (m, 6H), 2.01-2.07 (m, 2H), 1.99 (s, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

50

55

EXAMPLE 195

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[3-nitro-4-({(3R)-1-oxetan-3-ylpyrrolidin-3-yl]methyl}amino)phenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 195A

(R)-tert-butyl 3-((4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)sulfamoyl)-2-nitrophenylamino)methyl)pyrrolidine-1-carboxylate

[0836] The title compound was prepared by substituting (R)-tert-butyl 3-(aminomethyl)pyrrolidine-1-carboxylate for 1-acetylpiperidin-4-amine in EXAMPLE 53B.

EXAMPLE 195B

(S)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-(3-nitro-4-(pyrrolidin-3-ylmethylamino)phenylsulfonyl)benzamide

[0837] The title compound was prepared by substituting EXAMPLE 195A for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 195C

(R)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-(3-nitro-4-((1-oxetan-3-yl)pyrrolidin-3-yl)methylamino)phenylsulfonyl)benzamide

[0838] The title compound was prepared by substituting EXAMPLE 195B for tert-butyl piperazine-1-carboxylate and oxetan-3-one for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A. ¹H NMR (500MHz, dimethylsulfoxide-d₆) 11.67 (s, 1H), 8.81 (t, 1H), 8.55 (d, 1H), 8.02 (d, 1H), 7.79 (dd, 1H), 7.50 (m, 3H), 7.35 (m, 2H), 7.04 (m, 3H), 6.67 (dd, 1H), 6.39 (m, 1H), 6.19 (d, 1H), 4.57 (m, 2H), 4.48 (m, 2H), 3.68 (m, 2H), 3.30 (m, 2H), 3.06 (m, 4H), 2.74 (m, 3H), 2.56 (m, 3H), 2.44 (m, 1H), 2.18 (m, 5H), 1.95 (m, 3H), 1.58 (m, 1H), 1.36 (m, 2H), 0.94 (s, 6H).

EXAMPLE 196

Trans-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-hydroxycyclohexyl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 196A

Trans-4-(4-(tert-butyldimethylsilyloxy)cyclohexyl)methoxy)-3-nitrobenzenesulfonamide

[0839] The title compound was prepared as described in EXAMPLE 12A by replacing (1,4-dioxan-2-yl)methanol with trans-4-(tert-butyldimethylsilyloxy)cyclohexyl)methanol (made according to the procedures in WO 2008/124878).

EXAMPLE 196B

Trans-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-(4-(((1r,4r)-4-(tert-butyldimethylsilyloxy)cyclohexyl)methoxy)-3-nitrophenylsulfonyl)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzamide

[0840] The title compound was prepared as described in EXAMPLE 1G using EXAMPLE 196A in place of EXAMPLE 1F and EXAMPLE 3J in place of EXAMPLE 1E.

EXAMPLE 196C

Trans-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-hydroxycyclohexyl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0841] EXAMPLE 196B (150 mg) in dichloromethane (5 mL) and methanol (2 mL) was treated with 10% aqueous HCl (3 mL) for 1 hour and concentrated. The residue was purified by reverse phase HPLC on a C18 column using a gradient

of 40-60% acetonitrile in 0.1% trifluoroacetic acid water to give the title compound as a trifluoroacetate salt. The trifluoroacetic acid salt was dissolved in dichloromethane (30 mL) and washed with 50% aqueous NaHCO₃. The organic layer was dried over anhydrous Na₂SO₄, filtered, and concentrated to give the title compound. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 11.27 (s, 1H), 8.34 (d, 1H), 7.95 - 8.08 (m, 2H), 7.47 - 7.55 (m, 3H), 7.32 - 7.40 (m, 3H), 7.01 - 7.07 (m, 2H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.20 (d, 1H), 4.54 (d, 1H), 3.96 - 4.06 (m, 2H), 3.10 (s, 4H), 2.84 (s, 2H), 2.05 - 2.39 (m, 6H), 1.96 (s, 2H), 1.46 - 1.93 (m, 5H), 1.39 (t, 2H), 0.98 - 1.29 (m, 4H), 0.92 (s, 6H)

EXAMPLE 197

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[3-(dimethylamino)propoxy]benzyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 197A

3-(4-(aminomethyl)phenoxy)-N,N-dimethylpropan-1-amine

[0842] 4-(3-(Dimethylamino)propoxy)benzonitrile (300 mg) in methanol (20 mL) was treated with Raney nickel (wet, 1.5 g) under H₂ (30 psi) for 4 hour. The insoluble material was filtered off and the filtrate was concentrated to provide the title compound.

EXAMPLE 197B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[3-(dimethylamino)propoxy]benzyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0843] The title compound was prepared as described in EXAMPLE 120B using EXAMPLE 197A in place of EXAMPLE 120A. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.56 (s, 1H), 8.80 (t, 1H), 8.42 (d, 1H), 7.93 (d, 1H), 7.52 - 7.61 (m, 2H), 7.41 - 7.47 (m, 1H), 7.26 - 7.36 (m, 5H), 7.03 - 7.08 (m, 2H), 6.89 (d, 2H), 6.73 (d, 1H), 6.61 (dd, 1H), 6.31 (dd, 1H), 6.22 (d, 1H), 4.52 (d, 2H), 3.99 (t, 2H), 2.90 - 3.05 (m, 7H), 2.72 (s, 2H), 2.61 (s, 6H), 2.09 - 2.24 (m, 6H), 1.89 - 2.04 (m, 5H), 1.38 (t, 2H), 0.92 (s, 6H)

EXAMPLE 198

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[4-(2-morpholin-4-ylethoxy)benzyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 198A

(4-(2-morpholinoethoxy)phenyl)methanamine

[0844] The title compound was prepared as described in EXAMPLE 197A using 4-(2-morpholinoethoxy)benzonitrile in place of 4-(3-(dimethylamino)propoxy)benzonitrile.

EXAMPLE 198B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[4-(2-morpholin-4-ylethoxy)benzyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0845] The title compound was prepared as described in EXAMPLE 120B using EXAMPLE 198A in place of EXAMPLE 120A. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 9.00 (t, 1H), 8.56 (d, 1H), 8.02 (d, 1H), 7.72 (dd, 1H), 7.46 - 7.54 (m, 3H), 7.27 - 7.36 (m, 4H), 7.01 - 7.07 (m, 2H), 6.89 - 6.95 (m, 3H), 6.66 (dd, 1H), 6.38 (dd, 1H), 6.18 (d, 1H), 4.56 (d, 2H), 4.07 (t, 2H), 3.54 - 3.61 (m, 4H), 3.06 (s, 4H), 2.71 - 2.78 (m, 4H), 2.07 - 2.24 (m, 6H), 1.95 (s, 2H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 199

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(*E*)-4-hydroxy-1-adamantyl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 199A

4-({[(*E*)-4-Hydroxy-adamantan-1-ylmethyl]-amino}-3-nitro-benzenesulfonamide

[0846] 4-Fluoro-3-nitrobenzenesulfonamide (0.5 g) and 5-(aminomethyl)adamantan-2-ol (0.6 g) in tetrahydrofuran (10 mL) were treated with triethylamine (1 mL) overnight. The reaction mixture was concentrated and the residue was purified by reverse phase HPLC, eluting 40-60% acetonitrile in 0.1 trifluoroacetic acid water to give two isomers, which were temporarily assigned as EXAMPLE 199A and EXAMPLE 199B, respectively.

EXAMPLE 199B

4-({[(*Z*)-4-Hydroxy-adamantan-1-ylmethyl]-amino}-3-nitro-benzenesulfonamide

[0847] 4-Fluoro-3-nitrobenzenesulfonamide (0.5 g) and 5-(aminomethyl)adamantan-2-ol (0.6 g) in tetrahydrofuran (10 mL) were treated with triethylamine (1 mL) overnight. The reaction mixture was concentrated and the residue was purified by reverse phase HPLC, eluting 40-60% acetonitrile in 0.1 trifluoroacetic acid water to give two isomers, which were temporarily assigned as EXAMPLE 199A and EXAMPLE 199B, respectively.

EXAMPLE 199C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(*E*)-4-hydroxy-1-adamantyl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0848] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 199A in place of EXAMPLE 11B. ¹H NMR (400 MHz, dimethylsulfoxide-*d*₆) δ 11.68 (s, 1H), 11.40 (s, 1H), 8.55 (d, 1H), 8.50 (t, 1H), 8.03 (d, 1H), 7.77 (dd, 1H), 7.46 - 7.54 (m, 3H), 7.31 - 7.38 (m, 2H), 7.14 (d, 1H), 7.01 - 7.06 (m, 2H), 6.68 (dd, 1H), 6.38 (dd, 1H), 6.19 (d, 1H), 4.61 (d, 1H), 3.63 (d, 1H), 3.02 - 3.16 (m, 6H), 2.75 (s, 2H), 2.17 (d, 6H), 2.04 (d, 2H), 1.95 (s, 2H), 1.76 - 1.88 (m, 3H), 1.49 - 1.61 (m, 6H), 1.38 (t, 2H), 1.29 (d, 2H), 0.92 (s, 6H).

EXAMPLE 200

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[(*Z*)-4-hydroxy-1-adamantyl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0849] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 199B in place of EXAMPLE 11B. ¹H NMR (400 MHz, dimethylsulfoxide-*d*₆) δ 11.68 (s, 1H), 11.39 (s, 1H), 8.55 (d, 1H), 8.51 (t, 1H), 8.04 (d, 1H), 7.77 (dd, 1H), 7.46 - 7.55 (m, 3H), 7.31 - 7.37 (m, 2H), 7.14 (d, 1H), 7.01 - 7.06 (m, 2H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.19 (d, 1H), 4.61 (d, 1H), 3.61 (d, 1H), 3.08 (d, 6H), 2.75 (s, 2H), 2.17 (d, 6H), 1.79 - 1.99 (m, 7H), 1.55 - 1.69 (m, 4H), 1.49 (s, 2H), 1.38 (t, 2H), 1.22 (d, 2H), 0.92 (s, 6H).

EXAMPLE 201

N-({4-[(1*S*,4*S*)-bicyclo[2.2.1]hept-5-en-2-ylmethoxy]-3-nitrophenyl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 201 A

4-(1*S*,4*S*)-bicyclo[2.2.1]hept-5-en-2-ylmethoxy-3-nitrobenzenesulfonamide

[0850] The title compound was prepared as described in EXAMPLE 12A by replacing (1,4-dioxan-2-yl)methanol with (1*S*,4*S*)-bicyclo[2.2.1]hept-5-en-2-ylmethanol.

EXAMPLE 201B

N-({4-([1S,4S]-bicyclo[2.2.1]hept-5-en-2-ylmethoxy)-3-nitrophenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0851] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 201A in place of EXAMPLE 11B. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 8.35 (d, 1H), 7.95 - 8.10 (m, 2H), 7.47 - 7.58 (m, 3H), 7.30 - 7.45 (m, 3H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.40 (d, 1H), 5.92 - 6.23 (m, 3H), 3.65 - 4.39 (m, 3H), 3.00 - 3.22 (m, 4H), 2.76 - 2.98 (m, 4H), 2.28 (s, 4H), 2.15 (s, 2H), 1.96 (s, 2H), 1.71 - 1.91 (m, 1H), 1.33 - 1.47 (m, 3H), 1.20 - 1.32 (m, 2H), 0.92 (s, 6H), 0.50 - 0.66 (m, 1H).

EXAMPLE 202

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1-methyl-5-oxopyrrolidin-3-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0852] EXAMPLE 82 (140 mg) was dissolved in dioxane (3.0 mL), and 4-amino-1-methylpyrrolidin-2-one hydrochloride (30 mg) and triethylamine (0.100 mL) were added. The reaction mixture was heated at 110°C for 40 hours. The reaction was concentrated and the crude material was purified by preparative HPLC using a C18 column, 250 x 50 mm, 10μ, and eluting with a gradient of 20-100% CH₃CN vs. 0.1% trifluoroacetic acid in water, giving the product as a trifluoroacetate salt. The salt was dissolved in dichloromethane (6 mL) and washed with 50% aqueous NaHCO₃. The organic layer was dried over anhydrous Na₂SO₄ and concentrated to give the title compound. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1H), 8.74 (d, 1H), 8.37 (br d, 1H), 8.02 (d, 1H), 7.83 (dd, 1H), 7.49 (m, 3H), 7.34 (d, 2H), 7.07 (d, 1H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.38 (m, 1H), 6.21 (d, 1H), 4.46 (m, 1H), 3.81 (dd, 1H), 3.38 (dd, 1H), 3.08 (br m, 4H), 2.82 (dd, 1H), 2.75 (s, 5H), 2.43 (dd, 1H), 2.21 (br m, 4H), 2.16 (br t, 2H), 1.95 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 203

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(1R,4R,5R,6S)-5,6-dihydroxybicyclo[2.2.1]hept-2-yl]methoxy}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 203A

4-(((1R,4R,5R,6S)-5,6-dihydroxybicyclo[2.2.1]heptan-2-yl)methoxy)-3-nitrobenzenesulfonamide

[0853] To a solution of EXAMPLE 201A (340 mg) in tetrahydrofuran (10 mL) and water (1 mL) was added N-methylmorpholine N-oxide (184 mg) and OsO₄ (2.5% in 2-methyl-2-propanol) (1.05 mL). The reaction mixture was stirred overnight and purified by reverse phase HPLC to provide two isomers, which were temporarily assigned as EXAMPLE 203A and EXAMPLE 203B, respectively.

EXAMPLE 203B

4-(((1R,4R,5S,6R)-5,6-dihydroxybicyclo[2.2.1]heptan-2-yl)methoxy)-3-nitrobenzenesulfonamide

[0854] To a solution of EXAMPLE 201A (340 mg) in tetrahydrofuran (10 mL) and water (1 mL) was added N-methylmorpholine N-oxide (184 mg) and OsO₄ (2.5% in 2-methyl-2-propanol) (1.05 mL). The reaction mixture was stirred overnight and purified by reverse phase HPLC to provide two isomers, which were temporarily assigned as EXAMPLE 203A and EXAMPLE 203B, respectively.

EXAMPLE 203C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(1R,4R,5R,6S)-5,6-dihydroxybicyclo[2.2.1]hept-2-yl]methoxy}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0855] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 203A in place of EXAMPLE 11B. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 8.33 (s, 1H), 7.97 - 8.07 (m, 2H), 7.48 - 7.55 (m, 3H), 7.41 (d, 1H), 7.32 - 7.37 (m, 2H), 7.02 - 7.07 (m, 2H), 6.67 (dd, 1H), 6.40 (dd, 1H), 6.20 (d, 1H), 4.58 (dd, 2H), 4.07 - 4.19 (m, 2H), 3.82 (t, 1H), 3.51 (t, 1H), 3.09 (s, 4H), 2.81 (s, 2H), 2.09 - 2.34 (m, 8H), 2.04 - 2.09 (m, 2H), 1.93 - 2.01

(m, 3H), 1.62 - 1.77 (m, 2H), 1.39 (t, 2H), 1.11 (d, 1H), 0.92 (s, 6H), 0.67 - 0.76 (m, 1H).

EXAMPLE 204

5 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([1R,4R,5S,6R)-5,6-dihydroxy-bicyclo[2.2.1]hept-2-yl]methoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0856] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 203B in place of EXAMPLE 11B. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 8.33 (s, 1H), 7.98 - 8.07 (m, 2H), 7.49 - 7.54 (m, 3H),
10 7.41 (d, 1H), 7.32 - 7.36 (m, 2H), 7.02 - 7.07 (m, 2H), 6.67 (dd, 1H), 6.40 (dd, 1H), 6.20 (d, 1H), 4.58 (dd, 2H), 4.13 (dd, 2H), 3.82 (t, 1H), 3.51 (t, 1H), 3.09 (s, 4H), 2.81 (s, 2H), 2.09 - 2.35 (m, 8H), 2.07 (s, 2H), 1.93 - 2.02 (m, 3H), 1.61 - 1.80 (m, 2H), 1.39 (t, 2H), 1.11 (d, 1H), 0.92 (s, 6H), 0.66 - 0.78 (m, 1H).

EXAMPLE 205

15 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-((3-oxocyclohexyl)methoxy)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 205A

20 Methyl 1,4-dioxaspiro[4.5]decane-7-carboxylate

[0857] To a solution of trimethylsilyltrifluoromethanesulfonate (0.034 mL) in dry dichloromethane (5 mL) was added 1,2-bis(trimethylsiloxy)ethane (4.55 mL) followed by methyl 3-oxocyclohexanecarboxylate (2.9 g). The reaction mixture
25 was stirred for 3 hours at -78 °C. The reaction mixture was quenched with dry pyridine (0.5 mL), poured into saturated aqueous NaHCO₃, and extracted with ether. The ether layer was dried over Na₂CO₃ / Na₂SO₄. The reaction mixture was concentrated and purified by flash chromatography on silica with 5 to 30% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 205B

1,4-dioxaspiro[4.5]decan-7-ylmethanol

[0858] The title compound was prepared by substituting EXAMPLE 205A for EXAMPLE 101C in EXAMPLE 101D.

EXAMPLE 205C

3-nitro-4-((3-oxocyclohexyl)methoxy)benzenesulfonamide

40 **[0859]** Into a 250 mL round-bottomed flask was added sodium hydride (0.5 g) in tetrahydrofuran (10 mL) and then 1,4-dioxaspiro[4.5]decan-7-ylmethanol (0.5 g) was added. After the mixture stirred at room temperature for 20 minutes, 4-fluoro-3-nitrobenzenesulfonamide (0.65 g) was added. The mixture was stirred at room temperature for overnight. Water (20 mL) was added slowly. The aqueous layer was extracted by dichloromethane (3x 20 mL). The combined
45 organic layers were dried over Na₂SO₄. After filtration, and concentration of the filtrate, the residue was purified by reverse phase chromatography, eluting with 30-60% acetonitrile in water with 0.1% trifluoroacetic acid.

EXAMPLE 205D

50 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-((3-oxocyclohexyl)methoxy)phenyl]sulfonyl)-2-(1 H-pyrrolo [2,3-b]pyridin-5-yloxy)benzamide

[0860] The title compound was prepared by substituting EXAMPLE 205C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.59 (s, 1H), 8.22 (s, 1H), 7.96 (d, 1H), 7.87 (m, 1H), 7.55 (d, 1H), 7.45 (t, 1H),
55 7.35 (m, 3H), 7.20 (m, 1H), 7.04 (d, 2H), 6.64 (dd, 1H), 6.34 (m, 1H), 6.23 (d, 1H), 4.07 (d, 2H), 3.04 (m, 4H), 2.76 (s, 2H), 2.35 (m, 2H), 2.20 (m, 8H), 1.96 (m, 4H), 1.58 (m, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 206

4-(4-{[2-(4-chlorophenyl)-5,5-dimethylcyclohexa-1,3-dien-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 206A

2-chloro-5,5-dimethylcyclohexa-1,3-dienecarbaldehyde

[0861] In a 250 mL round-bottomed flask was added N,N-dimethylformamide (3.5 mL) in dichloromethane (30 mL), and the mixture was cooled to -10 °C. Phosphoryl trichloride (4 mL) was added dropwise, and the solution was warmed up to room temperature. 4,4-Dimethylcyclohex-2-enone (5.5 mL) was then added slowly, and the mixture was heated to reflux overnight. The reaction mixture was cooled and quenched with a 0°C solution of sodium acetate (25 g in 50 mL water). The aqueous layer was extracted with diethyl ether (200 mL x3). The combined organic layers were dried over Na₂SO₄, filtered, and concentrated to give the product.

EXAMPLE 206B

2-(4-chlorophenyl)-5,5-dimethylcyclohexa-1,3-dienecarbaldehyde

[0862] Into a 1 L round-bottomed flask was added EXAMPLE 206A (6.8 g), 4-chlorophenylboronic acid (6.5 g), and palladium (II) acetate (0.2 g) in water (100 mL) to give a suspension. Potassium carbonate (15 g) and tetrabutylammonium bromide (10 g) were added. After degassing, the mixture was stirred at 45°C for 4 hours. After cooling and filtering through silica gel in a funnel, diethyl ether (4x 200mL) was used to extract the product. The combined organic layers were dried over Na₂SO₄, and filtered. The filtrate was concentrated and purified by flash chromatography on silica with 0-10% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 206C

Methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5,5-dimethylcyclohexa-1,3-dienyl)methyl)piperazin-1-yl)benzoate

[0863] The title compound was prepared by substituting EXAMPLE 206B for 4'-chlorobiphenyl-2-carboxaldehyde and EXAMPLE 15F for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A.

EXAMPLE 206D

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5,5-dimethylcyclohexa-1,3-dienyl)methyl)piperazin-1-yl)benzoic acid

[0864] The title compound was prepared by substituting EXAMPLE 206C for EXAMPLE 101E in EXAMPLE 101F.

EXAMPLE 206E

4-(4-{[2-(4-chlorophenyl)-5,5-dimethylcyclohexa-1,3-dien-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0865] The title compound was prepared by substituting EXAMPLE 206D for EXAMPLE 3J and EXAMPLE 1F for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.61 (s, 1H), 8.49 (m, 2H), 7.99 (m, 1H), 7.72 (m, 1H), 7.53 (d, 1H), 7.41 (m, 4H), 7.12 (d, 2H), 6.99 (m, 1H), 6.66 (dd, 1H), 6.35 (m, 1H), 6.23 (d, 1H), 5.74 (d, 1H), 5.58 (d, 1H), 3.84 (m, 2H), 3.26 (m, 4H), 3.06 (m, 4H), 2.88 (s, 2H), 2.24 (m, 6H), 1.61 (m, 2H), 1.26 (m, 3H), 1.00 (s, 6H).

EXAMPLE 207

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({(3R)-1-[2-fluoro-1-(fluoromethyl)ethyl]pyrrolidin-3-yl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 207A

(R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-amine

[0866] The title compound was prepared by substituting (R)-tert-butyl pyrrolidin-3-ylcarbamate for tert-butyl azetidin-3-ylcarbamate in EXAMPLE 151A.

EXAMPLE 207B

(R)-4-(1-(1,3-difluoropropan-2-yl)pyrrolidin-3-ylamino)-3-nitrobenzenesulfonamide

[0867] The title compound was prepared by substituting EXAMPLE 207A for EXAMPLE 151A in EXAMPLE 151B.

EXAMPLE 207C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({(3R)-1-[2-fluoro-1-(fluoromethyl)ethyl]pyrrolidin-3-yl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0868] The title compound was prepared by substituting EXAMPLE 207B for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 11.52 - 11.24 (m, 1H), 8.55 (d, 1H), 8.37 (d, 1H), 8.03 (d, 1H), 7.83 (dd, 1H), 7.57 - 7.45 (m, 3H), 7.34 (d, 2H), 7.06 (t, 3H), 6.67 (d, 1H), 6.38 (dd, 1H), 6.20 (d, 1H), 4.70 (d, 2H), 4.54 (d, 2H), 4.23 (s, 1H), 3.11 - 2.87 (m, 7H), 2.74 (dd, 4H), 2.35 - 2.13 (m, 7H), 1.95 (s, 2H), 1.70 (s, 1H), 1.39 (d, 2H), 0.92 (s, 6H).

EXAMPLE 208

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 208A

2-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-5-iodo-3-(trifluoromethyl)pyridine

[0869] A mixture of EXAMPLE 37C (0.537 g), 5-iodo-3-(trifluoromethyl)pyridin-2-ol (1.156 g), and triphenylphosphine (1.574 g) in tetrahydrofuran (20 mL) was cooled to 0 °C. To this solution was added (E)-di-tert-butyl diazene-1,2-dicarboxylate (0.921 g). The reaction mixture was stirred overnight. The solvent was removed, and the residue was purified with column flash chromatography on silica gel eluting with 4:1 hexanes/ethyl acetate to give the desired product.

EXAMPLE 208B

6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-5-(trifluoromethyl)pyridine-3-sulfonamide

[0870] EXAMPLE 207A (1.3 g) in tetrahydrofuran (10 mL) was cooled to -42 °C with a cold bath of CH₃CN/dry ice. To this solution was added 2.0 M isopropylmagnesium chloride (1.6 mL) dropwise over 5 minutes. The reaction mixture was stirred for 30 minutes at -42 °C, then allowed to warm to 0 °C over 10 minutes. The reaction mixture was cooled again to -42 °C, and SO₂ was bubbled through it for 10 minutes. The reaction mixture was stirred for another 30 minutes. To this solution was sulfuric dichloride (0.433 g). On warming to room temperature, concentrated NH₄OH (10 mL) was added and the reaction mixture was stirred for another 2 hours. The reaction mixture was partitioned between water and ethyl acetate. The aqueous layer was extracted with additional ethyl acetate three times. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified by flash column chromatography on silica gel eluting with 3:1 hexanes/ethyl acetate to give the title compound.

EXAMPLE 208C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0871] The title compound was prepared by substituting EXAMPLE 208B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.61 (s, 1H), 8.72 (s, 1H), 8.36 (s, 1H), 7.98 (d, 1H), 7.55 (d, 1H), 7.42-7.47 (m, 2H), 7.36 (d, 2H), 7.05 (d, 2H), 6.66 (dd, 1H), 6.35 (s, 1H), 6.23 (s, 1H), 4.56 (d, 2H), 3.75-3.79 (m, 2H), 3.56-3.61 (m, 2H), 3.09 (s, 4H), 2.32-2.37 (m, 2H), 2.16(s, 2H), 1.97-1.99 (m, 2H), 1.79-1.86 (m, 4H), 1.40 (t, 2H), 0.93 (s, 6H).

EXAMPLE 209

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[3-nitro-4-(((3S)-1-oxetan-3-ylpyrrolidin-3-yl)methyl)amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 209A

(S)-tert-butyl (1-(oxetan-3-yl)pyrrolidin-3-yl)methylcarbamate

[0872] The title compound was prepared by substituting (R)-tert-butyl pyrrolidin-3-ylmethylcarbamate for tert-butyl piperazine-1-carboxylate and 3-oxetanone for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 209B

(S)-(1-(oxetan-3-yl)pyrrolidin-3-yl)methanamine

[0873] The title compound was prepared by substituting EXAMPLE 209A for (S)-1-tert-butyl 2-methyl 4,4-difluoropyrrolidine-1,2-dicarboxylate in EXAMPLE 168A.

EXAMPLE 209C

(S)-3-nitro-4-((1-(oxetan-3-yl)pyrrolidin-3-yl)methylamino)benzenesulfonamide

[0874] The title compound was prepared by substituting 4-fluoro-3-nitrobenzenesulfonamide for 4-chloro-3-nitrobenzenesulfonamide and EXAMPLE 209B for 4-methylpiperazin-1-amine dihydrochloride in EXAMPLE 6A.

EXAMPLE 209D

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[3-nitro-4-(((3S)-1-oxetan-3-ylpyrrolidin-3-yl)methyl)amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0875] The title compound was prepared by substituting EXAMPLE 209C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, pyridine-d₅) δ 13.05 (s, 1H), 9.30 (d, 1H), 9.02 (t, 1H), 8.42 (d, 1H), 8.34 (dd, 1H), 8.10 (d, 1H), 7.67 (dd, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.82 (d, 1H), 6.75 (m, 1H), 6.52 (m, 2H), 4.82 (t, 1H), 4.75 (t, 1H), 4.67 (t, 2H), 3.57 (m, 1H), 3.24 (t, 2H), 3.07 (m, 4H), 2.75 (m, 3H), 2.57 (dd, 1H), 2.45 (s, 1H), 2.36 (t, 1H), 2.26 (s, 2H), 2.18 (m, 5H), 1.93 (m, 3H), 1.56 (m, 1H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 210

Trans-N-({5-chloro-6-[(4-methoxycyclohexyl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 210A

(4-methoxycyclohexyl)methanol

[0876] The title compound was prepared by substituting 4-methoxycyclohexanecarboxylic acid for 1-tert-butyl 4-ethyl 4-fluoropiperidine-1,4-dicarboxylate in EXAMPLE 126A.

EXAMPLE 210B

Trans-5-chloro-6-((4-methoxycyclohexyl)methoxy)pyridine-3-sulfonamide

- 5 **[0877]** The title compound was prepared by substituting EXAMPLE 210A for tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 210C

- 10 Trans-N-({5-chloro-6-[(4-methoxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0878]** The title compound was prepared by substituting EXAMPLE 210C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 8.50 (s, 1H), 8.17 (s, 1H), 8.02 (d, 1H), 7.49-7.54 (m, 3H), 7.35 (d, 2H), 7.05 (d, 2H), 6.67 (dd, J 1H), 6.39 (s, 1H), 6.21 (s, 1H), 4.20 (d, 2H), 3.23 (s, 3H), 3.06-3.09 (m, 4H), 2.15-2.37 (m, 4H), 1.96-2.03 (m, 4H), 1.74-1.84 (m, 2H), 1.40 (t, 2H), 1.04-1.13 (m, 4H), 0.93 (s, 6H).

EXAMPLE 211

- 20 Cis-N-({5-chloro-6-[(4-methoxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 211 A

- 25 Cis-5-chloro-6-((4-methoxycyclohexyl)methoxy)pyridine-3-sulfonamide

[0879] The title compound was isolated as a by-product in the synthesis of EXAMPLE 210B.

EXAMPLE 211B

- 30 Cis-N-({5-chloro-6-[(4-methoxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0880]** The title compound was prepared by substituting EXAMPLE 211A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 8.51 (s, 1H), 8.17 (s, 1H), 8.03 (d, 1H), 7.49-7.54 (m, 3H), 7.35 (d, 2H), 7.05 (d, 2H), 6.68 (dd, 1H), 6.39 (s, 1H), 6.21 (s, 1H), 4.21 (d, 2H), 3.20 (s, 3H), 3.06 (s, 4H), 2.15-2.37 (m, 4H), 1.96 (s, 2H), 1.80-1.84 (m, 2H), 1.50-1.54 (m, 2H), 1.34-1.44 (m, 6H), 0.93 (s, 6H).

EXAMPLE 212

- 40 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(3S)-1-oxetan-3-yl]pyrrolidin-3-yl}amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 212A

- 45 (S)-tert-butyl 1-(oxetan-3-yl)pyrrolidin-3-ylcarbamate

- [0881]** The title compound was prepared by substituting (S)-tert-butyl pyrrolidin-3-ylcarbamate for tert-butyl piperazine-1-carboxylate and 3-oxetanone for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 212B

(S)-1-(oxetan-3-yl)pyrrolidin-3-amine

- 55 **[0882]** The title compound was prepared by substituting EXAMPLE 212A for (S)-1-tert-butyl 2-methyl 4,4-difluoropyrrolidine-1,2-dicarboxylate in EXAMPLE 168A.

EXAMPLE 212C

(S)-3-nitro-4-(1-(oxetan-3-yl)pyrrolidin-3-ylamino)benzenesulfonamide

- 5 **[0883]** The title compound was prepared by substituting 4-fluoro-3-nitrobenzenesulfonamide for 4-chloro-3-nitrobenzenesulfonamide and EXAMPLE 212B for 4-methylpiperazin-1-amine dihydrochloride in EXAMPLE 6A.

EXAMPLE 212D

- 10 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-[[[(3S)-1-oxetan-3-ylpyrrolidin-3-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0884]** The title compound was prepared by substituting EXAMPLE 212C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, pyridine-d₅) δ 13.03 (s, 1H), 9.27 (d, 1H), 8.58 (d, 1H), 8.42 (d, 1H), 8.37 (dd, 1H), 8.09 (d, 1H), 7.67 (t, 1H), 7.64 (d, 1H), 7.44 (d, 2H), 7.07 (d, 2H), 6.86 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (dd, 1H), 4.67 (m, 4H), 4.09 (m, 1H), 3.59 (m, 1H), 3.07 (m, 4H), 2.77 (s, 2H), 2.69 (m, 2H), 2.62 (dd, 1H), 2.28 (m, 4H), 2.14 (m, 4H), 1.97 (s, 2H), 1.68 (m, 1H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 213

- 20 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[[{4-[2-(2-methoxyethoxy)ethyl]morpholin-2-yl]methyl}amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 213A

- 25 4-((4-(2-(2-methoxyethoxy)ethyl)morpholin-2-yl)methylamino)-3-nitrobenzenesulfonamide

- [0885]** The title compound was prepared by substituting 2-(2'-methoxyethoxy)ethyl bromide for methyl iodide in EXAMPLE 134B.

30 EXAMPLE 213B

- 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[[{4-[2-(2-methoxyethoxy)ethyl]morpholin-2-yl]methyl}amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- 35 **[0886]** The title compound was prepared by substituting EXAMPLE 213A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 12.98 (s, 1H), 9.26 (d, 1H), 8.87 (t, 1H), 8.43 (d, 1H), 8.32 (dd, 1H), 8.11 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.91 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 3.96-3.86 (m, 2H), 3.72 (dd, 1H), 3.67-3.61 (m, 4H), 3.51 (t, 2H), 3.48-3.38 (m, 2H), 3.28 (s, 3H), 3.07 (m, 4H), 2.95 (d, 1H), 2.77 (s, 2H), 2.70 (m, 1H), 2.60 (t, 2H), 2.30-2.05 (m, 8H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 214

- 45 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[4-(cyanomethyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 214A

- 50 4-((4-(cyanomethyl)morpholin-2-yl)methylamino)-3-nitrobenzenesulfonamide

- [0887]** The title compound was prepared by substituting 2-bromoacetonitrile for methyl iodide in EXAMPLE 134B.

EXAMPLE 214B

- 55 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[4-(cyanomethyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0888]** The title compound was prepared by substituting EXAMPLE 214A for EXAMPLE 130C in EXAMPLE 130D. ¹H

NMR (500MHz, pyridine- d_5) δ 13.01 (s, 1H), 9.26 (d, 1H), 8.86 (t, 1H), 8.43 (d, 1H), 8.35 (dd, 1H), 8.11 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.94 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 3.93 (m, 1H), 3.87 (d, 1H), 3.77 (s, 2H), 3.65 (dt, 1H), 3.51-3.40 (m, 2H), 3.07 (m, 4H), 2.87 (d, 1H), 2.77 (s, 2H), 2.60 (d, 1H), 2.50 (m, 1H), 2.38 (t, 1H), 2.26 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

5

EXAMPLE 215

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-(N,N-dimethylglycyl)morpholin-2-yl]methyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10

EXAMPLE 215A

4-([4-(2-(dimethylamino)acetyl)morpholin-2-yl]methylamino)-3-nitrobenzenesulfonamide

15

[0889] The title compound was prepared by substituting 2-dimethylaminoacetyl chloride hydrochloride for methyl iodide in EXAMPLE 134B.

EXAMPLE 215B

20

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-(N,N-dimethylglycyl)morpholin-2-yl]methyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

25

[0890] The title compound was prepared by substituting EXAMPLE 215A for EXAMPLE 130C in EXAMPLE 130D. ^1H NMR (500MHz, pyridine- d_5) δ 13.00 (s, 1H), 9.27 (d, 1H), 8.87 (bs, 1H), 8.43 (d, 1H), 8.34 (dd, 1H), 8.10 (dd, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.94 (dd, 1H), 6.75 (d, 1H), 6.54 (s, 1H), 6.48 (s, 1H), 4.55 (dd, 1H), 4.20 (dd, 1H), 3.95-3.76 (m, 2H), 3.60-3.40 (m, 3H), 3.32 (dd, 1H), 3.25-3.12 (m, 2H), 3.07 (m, 4H), 2.80 (m, 1H), 2.77 (s, 2H), 2.26 (s, 6H), 2.23 (s, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 216

30

(2-([4-([4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl)sulfamoyl]-2-nitrophenyl)amino)methyl)morpholin-4-yl)acetic acid

EXAMPLE 216A

35

tert-butyl 2-(2-((2-nitro-4-sulfamoylphenylamino)methyl)morpholino)acetate

[0891] The title compound was prepared by substituting *tert*-butyl 2-bromoacetate for methyl iodide in EXAMPLE 134B.

40

EXAMPLE 216B

tert-butyl 2-(2-((4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)benzoyl)sulfamoyl)-2-nitrophenylamino)methyl)morpholino)acetate

45

[0892] The title compound was prepared by substituting EXAMPLE 216A for EXAMPLE 130C in EXAMPLE 130D.

EXAMPLE 216C

50

(2-([4-([4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl)sulfamoyl]-2-nitrophenyl)amino)methyl)morpholin-4-yl)acetic acid

55

[0893] The title compound was prepared by treating EXAMPLE 216B with 50% trifluoroacetic acid in dichloromethane. ^1H NMR (500MHz, pyridine- d_5) δ 12.97 (s, 1H), 9.26 (d, 1H), 8.87 (t, 1H), 8.43 (d, 1H), 8.30 (dd, 1H), 8.12 (d, 1H), 7.69 (t, 1H), 7.64 (d, 1H), 7.43 (d, 2H), 7.08 (d, 2H), 6.88 (d, 1H), 6.76 (dd, 1H), 6.55 (d, 1H), 6.47 (m, 1H), 4.05-4.00 (m, 1H), 3.91 (d, 1H), 3.79 (dt, 1H), 3.50 (s, 2H), 3.45 (m, 2H), 3.13 (d, 1H), 3.07 (m, 4H), 2.88 (d, 1H), 2.78 (s, 2H), 2.57 (dt, 1H), 2.43 (t, 1H), 2.26 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 217

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[3-nitro-4-({[4-(oxetan-3-yl)morpholin-2-yl]methyl}amino)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 217A

[0894] The title compound was prepared by substituting EXAMPLE 134A for tert-butyl piperazine-1-carboxylate and 3-oxetanone for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 217B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[3-nitro-4-({[4-(oxetan-3-yl)morpholin-2-yl]methyl}amino)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0895] The title compound was prepared by substituting EXAMPLE 217A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.26 (d, 1H), 8.87 (t, 1H), 8.44 (d, 1H), 8.34 (dd, 1H), 8.12 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.94 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 4.64 (m, 4H), 3.93 (m, 1H), 3.89 (d, 1H), 3.68 (dt, 1H), 3.53-3.35 (m, 3H), 3.07 (m, 4H), 2.77 (s, 2H), 2.72 (d, 1H), 2.44 (d, 1H), 2.26 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.85 (t, 1H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 218

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[4-(cyclopropylmorpholin-2-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 218A

4-({[4-(cyclopropylmorpholin-2-yl)methylamino]-3-nitrobenzenesulfonamide

[0896] The title compound was prepared by substituting EXAMPLE 134A for EXAMPLE 173A in EXAMPLE 173B.

EXAMPLE 218B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[4-(cyclopropylmorpholin-2-yl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0897] The title compound was prepared by substituting EXAMPLE 218A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.26 (d, 1H), 8.88 (t, 1H), 8.44 (d, 1H), 8.34 (dd, 1H), 8.12 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.94 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 3.88 (d, 1H), 3.84-3.81 (m, 1H), 3.59 (dt, 1H), 3.50-3.40 (m, 2H), 3.07 (m, 4H), 2.93 (d, 1H), 2.77 (s, 2H), 2.69 (d, 1H), 2.34 (dt, 1H), 2.26 (m, 2H), 2.21 (t, 1H), 2.14 (m, 4H), 1.97 (s, 2H), 1.58 (m, 1H), 1.39 (t, 2H), 0.94 (s, 6H), 0.45-0.39 (m, 4H).

EXAMPLE 219

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[5-(methylsulfonyl)-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 219A

5-(methylthio)-6-((tetrahydro-2H-pyran-4-yl)methoxy)pyridine-3-sulfonamide

[0898] A mixture of EXAMPLE 36B (0.1 g) and sodium methanethiolate (0.04 g) in N,N-dimethylformamide (2 mL) was heated at 80°C overnight. After cooling, the reaction mixture was partitioned between water and ethyl acetate. The organic layer was separated, and the aqueous layer was extracted with additional ethyl acetate three times. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified by flash column chromatography on silica gel using 10-50% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 219B

5-(methylsulfonyl)-6-((tetrahydro-2H-pyran-4-yl)methoxy)pyridine-3-sulfonamide

5 **[0899]** A mixture of EXAMPLE 219A (0.15 g) and 75% meta-chloroperoxybenzoic acid (0.217 g) in chloroform (4 mL) was stirred at room temperature. The reaction mixture was stirred overnight. The reaction mixture was then partitioned between ethyl acetate and water. The organic layer was separated, and the aqueous layer was extracted with additional ethyl acetate three times. The combined organic layers were washed with brine, dried over MgSO_4 , filtered, and concentrated. The residue was purified by flash column chromatography on silica gel using 10-50% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 219C

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-(methylsulfonyl)-6-(tetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

15 **[0900]** The title compound was prepared by substituting EXAMPLE 219B for EXAMPLE 11B in EXAMPLE 11D. ^1H NMR (500MHz, dimethylsulfoxide- d_6) δ 11.65 (s, 1H), 8.81 (s, 1H), 8.55 (d, 1H), 8.01 (d, 1H), 7.55 (d, 1H), 7.49-7.50 (m, 2H), 7.37 (d, 2H), 7.05 (d, 2H), 6.67 (dd, 1H), 6.38 (dd, 1H), 6.21 (d, 1H), 4.36 (d, 2H), 3.88 (dd, 2H), 3.13 (s, 4H), 2.95 (s, 2H), 2.36-2.38 (m, 2H), 2.03-2.16 (m, 4H), 1.97 (s, 3H), 1.66-1.69 (m, 2H), 1.38-1.402 (m, 4H), 0.93 (s, 6H).

EXAMPLE 220

25 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 220A

30 4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-(trifluoromethylsulfonyl)benzenesulfonamide

[0901] To a solution of EXAMPLE 37C (0.500 g) in tetrahydrofuran (5 mL) was added sodium hydride (0.596 g). Additional tetrahydrofuran (25 mL) was added and the mixture stirred for 30 minutes, then 4-fluoro-3-(trifluoromethylsulfonyl)benzenesulfonamide (1.145 g) was added as a solution in tetrahydrofuran (5 mL). After stirring for 2 hours, the reaction mixture was partitioned between 1N aqueous HCl (50 mL) and dichloromethane (200 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated. The resulting solid was chromatographed over silica gel (Reveleris 80g) eluting with a gradient of 0.5% to 7.5% methanol/dichloromethane over 30 minutes (flow = 40 mL/min) to provide the title compound.

EXAMPLE 220B

40 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

45 **[0902]** The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 220A for EXAMPLE 1F in EXAMPLE 1G. ^1H NMR (300 MHz, dimethylsulfoxide- d_6) δ 11.66 (s, 1H), 8.42 (s, 1H), 8.35 - 8.22 (m, 1H), 8.01 (s, 1H), 7.49 (d, 4H), 7.35 (d, 2H), 7.05 (d, 2H), 6.68 (s, 1H), 6.38 (s, 1H), 6.21 (s, 1H), 4.42 (d, 2H), 3.76 (s, 2H), 3.59 (s, 2H), 3.10 (s, 6H), 2.15 (s, 6H), 2.02 - 1.74 (m, 6H), 1.40 (s, 2H), 0.93 (s, 6H).

EXAMPLE 221

50 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-[(4-methyltetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 221 A

55 4-[(4-methyltetrahydro-2H-pyran-4-yl)methoxy]-3-nitrobenzenesulfonamide

[0903] The title compound was prepared by substituting (4-methyltetrahydro-2H-pyran-4-yl)methanol for (tetrahydro-

2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 221B

5 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-methyltetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0904] The title compound was prepared by substituting EXAMPLE 221A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 8.36 (s, 1H), 8.04-8.06 (m, 2H), 7.50-7.53 (m, 3H), 7.41 (d, 1H),
10 7.35 (d, 2H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.40 (dd, 1H), 6.20 (d, 1H), 4.00 (s, 2H), 3.63-3.67 (m, 2H), 3.53-3.58 (m, 2H), 3.09 (s, 4H), 2.82 (s, 2H), 2.27 (s, 2H), 2.15 (s, 2H), 1.58-1.63 (m, 2H), 1.39 (t, 2H), 1.30-1.34 (m, 2H), 1.09 (s, 3H), 0.92 (s, 6H).

EXAMPLE 222

15 ethyl 4-(4-{[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl}-2-nitrophenyl)piperazine-1-carboxylate

EXAMPLE 222A

20 ethyl 4-(2-nitro-4-sulfamoylphenyl)piperazine-1-carboxylate

[0905] The title compound was prepared by substituting ethyl piperazine-1-carboxylate for 1-(tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 222B

30 ethyl 4-(4-{[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl}-2-nitrophenyl)piperazine-1-carboxylate

[0906] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 222A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.52 (br. s, 1H), 8.08 (d, 1H), 7.89 (d, 1H), 7.59 (m, 2H), 7.43 (t, 1H), 7.35 (d, 2H), 7.23 (d, 1H), 7.05 (d, 2H), 6.94 (d, 1H), 6.63 (dd, 1H), 6.29 (m, 2H), 4.07 (q, 2H), 3.47 (m, 4H), 3.17 (d, 2H), 3.00 (m, 8H), 2.73 (s, 2H), 2.18 (m, 6H), 1.96 (s, 2H), 1.39 (t, 2H), 1.20 (t, 3H), 0.93 (s, 6H).

EXAMPLE 223

40 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[4-(morpholin-4-yl)piperidin-1-yl]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 223A

[0907] The title compound was prepared by substituting 4-(piperidin-4-yl)morpholine for 1-(tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 223B

50 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[4-(morpholin-4-yl)piperidin-1-yl]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0908] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 223A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.53 (br. s, 1H), 8.05 (d, 1H), 7.91 (d, 1H), 7.58 (m, 2H), 7.43 (t, 1H), 7.35 (d, 2H), 7.26 (d, 1H), 7.05 (d, 2H), 6.91 (d, 1H), 6.62 (dd, 1H), 6.29 (m, 2H), 5.76 (s, 1H), 3.57 (m, 4H), 3.20 (m, 2H), 3.01 (m, 4H), 2.80 (t, 2H), 2.73 (s, 2H), 2.47 (m, 4H), 2.32 (m, 1H), 2.18 (m, 6H), 1.96 (m, 3H), 1.82 (m, 2H), 1.44 (m, 4H), 0.93 (s, 6H).

EXAMPLE 224

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(3R)-1-(oxetan-3-yl)pyrrolidin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 224A

(R)-tert-butyl 1-(oxetan-3-yl)pyrrolidin-3-ylcarbamate

[0909] The title compound was prepared by substituting (R)-tert-butyl pyrrolidin-3-ylcarbamate for tert-butyl piperazine-1-carboxylate and 3-oxetanone for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 224B

(R)-1-(oxetan-3-yl)pyrrolidin-3-amine

[0910] The title compound was prepared by substituting EXAMPLE 224A for (S)-1-tert-butyl 2-methyl 4,4-difluoropyrrolidine-1,2-dicarboxylate in EXAMPLE 168A.

EXAMPLE 224C

(R)-3-nitro-4-(1-(oxetan-3-yl)pyrrolidin-3-ylamino)benzenesulfonamide

[0911] The title compound was prepared by substituting 4-fluoro-3-nitrobenzenesulfonamide for 4-chloro-3-nitrobenzenesulfonamide and EXAMPLE 224B for 4-methylpiperazin-1-amine dihydrochloride in EXAMPLE 6A.

EXAMPLE 224D

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[(3R)-1-(oxetan-3-yl)pyrrolidin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0912] The title compound was prepared by substituting EXAMPLE 224C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400 MHz, pyridine-d₅) δ 13.03 (s, 1H), 9.26 (d, 1H), 8.57 (d, 1H), 8.42 (d, 1H), 8.36 (dd, 1H), 8.09 (d, 1H), 7.66 (m, 1H), 7.64 (d, 1H), 7.44 (m, 2H), 7.07 (m, 2H), 6.86 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (dd, 1H), 4.67 (m, 4H), 3.58 (m, 1H), 3.07 (m, 4H), 2.77 (m, 2H), 2.68 (m, 2H), 2.61 (m, 1H), 2.28 (m, 4H), 2.14 (m, 4H), 1.97 (m, 2H), 1.67 (m, 1H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 225

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]amino}-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 225A

(R)-4-(1-(1,3-difluoropropan-2-yl)pyrrolidin-3-ylamino)-3-(trifluoromethylsulfonyl)benzenesulfonamide

[0913] To EXAMPLE 207A (0.217 g) and 4-fluoro-3-(trifluoromethylsulfonyl)benzenesulfonamide (0.281 g) in tetrahydrofuran (5 mL) was added diisopropylethylamine (0.559 mL) and the reaction was allowed to stir at room temperature for 1 hour and was then heated to 50°C for 1 hour. The reaction was concentrated, the residue was loaded onto silica gel (Reveleris 40 g) and eluted with a gradient of 0.75% methanol/dichloromethane to 7.5% methanol/dichloromethane to provide the title compound.

EXAMPLE 225B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]amino}-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0914] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 225A for

EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 11.52 - 11.23 (m, 1H), 8.17 (d, 1H), 8.04 (d, 1H), 7.95 (d, 1H), 7.54 (d, 1H), 7.53 - 7.50 (m, 1H), 7.48 (d, 1H), 7.34 (d, 2H), 7.10 - 6.97 (m, 4H), 6.67 (d, 1H), 6.40 (dd, 1H), 6.18 (d, 1H), 4.60 (dd, 4H), 4.20 (s, 1H), 3.11 - 2.63 (m, 12H), 2.19 (d, 6H), 1.95 (s, 2H), 1.58 (s, 1H), 1.40 (d, 2H), 0.92 (s, 6H).

EXAMPLE 226

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1-isopropylpiperidin-4-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 226A

tert-butyl 4-(4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)sulfamoyl)-2-nitrophenylamino)piperidine-1-carboxylate

[0915] To a solution of EXAMPLE 82 (800 mg) and tert-butyl 4-aminopiperidine-1-carboxylate (203 mg) in dioxane (10 mL) was added Hunig's Base (1 mL). The mixture was stirred at 120°C overnight. The mixture was diluted with ethyl acetate (200 mL) and washed with water, brine and dried over Na₂SO₄. After filtration and evaporation of solvent, the residue was loaded on a silica gel cartridge and eluted with 3% methanol in dichloromethane to give the title compound.

EXAMPLE 226B

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-(3-nitro-4-(piperidin-4-ylamino)phenyl)sulfonylbenzamide

[0916] To a solution of EXAMPLE 226A (902 mg) in dichloromethane (10 mL) was added trifluoroacetic acid (5 mL). The mixture was stirred at room temperature for 3 hours. The mixture was concentrated under vacuum and co-concentrated with dichloromethane twice to afford the crude product which was used in the next step without further purification.

EXAMPLE 226C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(1-isopropylpiperidin-4-yl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0917] To a solution of EXAMPLE 226B (79 mg) in tetrahydrofuran (3 mL) and acetic acid (1 mL) was added acetone (54 mg) and MP-cyanoborohydride (150 mg, 2.25mmol/g). The mixture was stirred overnight. The mixture was filtered. The filtrate was concentrated and the residue was loaded on a silica gel cartridge and eluted with 5 to 10% 7N NH₃ in methanol in dichloromethane to provide the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 12.09 (s, 1H), 8.34 (m, 1H), 7.93 (m, 2H), 7.66 (m, 4H), 7.35 (d, 2H), 7.06 (d, 2H), 6.89 (m, 1H), 6.74 (dd, 1H), 6.59 (dd, 1H), 6.50 (d, 1H), 3.11 (m, 6H), 2.73 (m, 4H), 2.26 (m, 9H), 1.97 (s, 3H), 1.40 (t, 2H), 1.23 (s, 8H), 0.94 (s, 6H).

EXAMPLE 227

N-({4-[(1-tert-butylpiperidin-4-yl)amino]-3-nitrophenyl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 227A

1-tert-butylpiperidin-4-amine

[0918] To a solution of 1-tert-butylpiperidin-4-one (5.0 g) in methanol (100 mL) and water (10 mL) was added ammonium formate (20.3 g) and 0.5 g of Pd/C (10%). The mixture was stirred overnight. The mixture was filtered and the filtrate was concentrated under vacuum and the residue was diluted with ethyl acetate (500 mL) and washed with water and brine. After drying over Na₂SO₄ and filtration, the solvent was evaporated under vacuum to provide the title compound.

EXAMPLE 227B

4-(1-tert-butylpiperidin-4-ylamino)-3-nitrobenzenesulfonamide

- 5 **[0919]** To a mixture of 4-fluoro-3-nitrobenzenesulfonamide (2.2 g) and EXAMPLE 227A (1.56 g) in tetrahydrofuran (20 mL) was added Hunig's Base (6 mL). The mixture was stirred for 3 days. The mixture was diluted with ethyl acetate (300 mL) and water (100 mL) and stirred until the solid disappeared into the solution. The layers were separated and the organic phase was washed with water and brine, dried over Na₂SO₄, and filtered. The combined aqueous layers were extracted again with ethyl acetate and the combined organic phases were dried over Na₂SO₄. After filtration, the solvent was evaporated to provide the title compound.

EXAMPLE 227C

- 15 N-({4-[(1-tert-butylpiperidin-4-yl)amino]-3-nitrophenyl}sulfonyl)-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0920]** The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 227B for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.51 (s, 1 H), 8.43 (d, 1 H), 8.04 (m, 1 H), 7.93 (d, 1 H), 7.72 (m, 1 H), 7.56 (dd, 1 H), 7.42 (m, 1 H), 7.34 (m, 3 H), 7.05 (d, 2 H), 6.93 (dd, 1 H), 6.62 (dd, 1 H), 6.28 (m, 1 H), 3.04 (m, 6 H), 2.73 (s, 3 H), 2.25 (m, 9 H), 1.95 (s, 2 H), 1.68 (m, 2 H), 1.32 (m, 9 H), 0.93 (s, 6 H).

EXAMPLE 228

- 25 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({4-[(1-(2-methoxyethyl)piperidin-3-yl]methyl}amino)-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 228A

- 30 tert-butyl 3-((2-nitro-4-sulfamoylphenylamino)methyl)piperidine-1-carboxylate

[0921] The title compound was prepared by substituting *tert-butyl* 3-(aminomethyl)piperidine-1-carboxylate for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 228B

- 35 3-nitro-4-(piperidin-3-ylmethylamino)benzenesulfonamide

[0922] The title compound was prepared by substituting EXAMPLE 228A for EXAMPLE 113A in EXAMPLE 134A.

EXAMPLE 228C

4-((1-(2-methoxyethyl)piperidin-3-yl)methylamino)-3-nitrobenzenesulfonamide

- 45 **[0923]** The title compound was prepared by substituting EXAMPLE 228B for EXAMPLE 134A and 2-methoxyethyl bromide for methyl iodide in EXAMPLE 134B.

EXAMPLE 228D

- 50 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({4-[(1-(2-methoxyethyl)piperidin-3-yl]methyl}amino)-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [0924]** The title compound was prepared by substituting EXAMPLE 228C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅, 90 °C) δ 12.40 (s, 1H), 8.52 (s, 1H), 8.43 (s, 1H), 8.20 (m, 2H), 7.95 (bs, 1H), 7.80 (s, 1H), 7.46 (d, 1H), 7.36 (d, 2H), 7.07 (d, 2H), 7.05 (s, 1H), 6.75 (d, 1H), 6.59 (s, 1H), 6.47 (s, 1H), 3.65-3.50 (m, 5H), 3.20 (s, 3H), 3.04 (m, 5H), 2.81 (s, 3H), 2.74 (m, 1H), 2.24 (m, 7H), 2.06 (s, 2H), 2.00 (s, 2H), 1.75 (m, 1H), 1.57 (m, 2H), 1.42 (t, 2H), 1.15 (m, 1H), 0.95 (s, 6H).

EXAMPLE 229

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{4-([1-(cyanomethyl)piperidin-3-yl]methyl)amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 229A

4-([1-(cyanomethyl)piperidin-3-yl)methylamino]-3-nitrobenzenesulfonamide

[0925] The title compound was prepared by substituting EXAMPLE 228B for EXAMPLE 134A and 2-bromoacetonitrile for methyl iodide in EXAMPLE 134B.

EXAMPLE 229B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{4-([1-(cyanomethyl)piperidin-3-yl]methyl)amino}-3-nitrophenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0926] The title compound was prepared by substituting EXAMPLE 229A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.06 (s, 1H), 9.03 (s, 1H), 8.42 (s, 1H), 8.30 (d, 1H), 8.10 (d, 1H), 7.68 (m, 2H), 7.44 (d, 2H), 7.08 (m, 3H), 6.99 (d, 1H), 6.75 (d, 1H), 6.51 (m, 2H), 3.78 (m, 2H), 3.43 (d, 1H), 3.13 (m, 1H), 3.04 (m, 4H), 2.76 (s, 2H), 2.71-2.65 (m, 3H), 2.52 (m, 1H), 2.25 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.84 (m, 1H), 1.68 (m, 1H), 1.50 (m, 2H), 1.39 (t, 2H), 1.07-0.99 (m, 1H), 0.93 (s, 6H).

EXAMPLE 230

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 230A

4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]-3-(trifluoromethylsulfonyl)benzenesulfonamide

[0927] To a solution of (4-fluoro-1-methylpiperidin-4-yl)methanol (0.315 g) in tetrahydrofuran (5 mL) was added sodium hydride (0.342 g). After stirring for 15 minutes, 4-fluoro-3-(trifluoromethylsulfonyl)benzenesulfonamide (0.658 g) was added as a solution in tetrahydrofuran (2 mL) followed by additional tetrahydrofuran (5 mL). After stirring for 1 hour, the reaction was poured in dichloromethane (50 mL) and water (25 mL) and the pH of the water layer was adjusted to 8. The organic layer was dried over magnesium sulfate, filtered, and concentrated. The resulting oil was chromatographed over silica gel (Reveleris 40 g) eluting with a gradient of 1.0% to 10% 7N NH₃ in methanol/dichloromethane over 20 minutes then maintaining 10% 7N NH₃ in methanol/dichloromethane for 5 minutes (flow = 30 mL/min) to provide the title compound.

EXAMPLE 230B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0928] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 230A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.63 - 11.57 (m, 1H), 8.40 - 8.36 (m, 1H), 8.28 - 8.17 (m, 1H), 7.97 (s, 1H), 7.53 (d, 1H), 7.50 - 7.32 (m, 5H), 7.05 (d, 1H), 7.05 (d, 1H), 6.68 - 6.61 (m, 1H), 6.35 (s, 1H), 6.22 (s, 1H), 4.55 - 4.40 (m, 2H), 3.06 (s, 8H), 2.79 (s, 4H), 2.06 (d, 13H), 1.39 (s, 2H), 0.93 (s, 6H).

EXAMPLE 231

N-[[[5-chloro-6-[[[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]amino]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 231 A

(R)-5-chloro-6-(1-(1,3-difluoropropan-2-yl)pyrrolidin-3-ylamino)pyridine-3-sulfonamide

[0929] To EXAMPLE 207A (0.051 g) and EXAMPLE 40A (0.049 g) in dioxane (5 mL) was added diisopropylethylamine (0.131 mL) and the reaction was heated to 75°C for 1 hour then 85°C for 2 days. The reaction was concentrated, loaded onto silica gel (Reveleris 12 g) and eluted with a gradient of 0.75% methanol/dichloromethane to 7.5% methanol/dichloromethane to provide the title compound.

EXAMPLE 231B

N-[[[5-chloro-6-[[[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]amino]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0930] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 231A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-*d*₆) δ 11.71 (s, 1H), 11.44 - 11.11 (m, 1H), 8.44 (d, 1H), 8.07 (d, 1H), 7.90 (d, 1H), 7.61 (d, 1H), 7.52 (dd, 2H), 7.34 (d, 2H), 7.19 (s, 1H), 7.04 (d, 2H), 6.67 (d, 1H), 6.42 (dd, 1H), 6.16 (s, 1H), 4.77 - 4.39 (m, 5H), 3.19 - 2.63 (m, 11H), 2.19 (s, 7H), 1.91 (d, 3H), 1.38 (s, 2H), 0.92 (s, 6H).

EXAMPLE 232

tert-butyl 4-[[[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino]piperazine-1-carboxylate

EXAMPLE 232A

tert-butyl 4-nitrosopiperazine-1-carboxylate

[0931] In a 500 mL round-bottomed flask, 6N aqueous HCl (30 mL) was cooled to -10 °C, and tert-butyl piperazine-1-carboxylate (10 g) was added. Sodium nitrite (4.5 g) dissolved in 35 mL water was added slowly. NaOH (10 g in 20 mL water) was used to neutralize the solution. Dichloromethane (3x 50 mL) was used to extract the product. After drying over Na₂SO₄ and filtration, the solution was concentrated. The crude product was added to a silica gel column (Analogix, SF65-400 g,) and purified by eluting with 0-30% ethyl acetate in hexane.

EXAMPLE 232B

tert-butyl 4-aminopiperazine-1-carboxylate

[0932] In a 100 mL round-bottomed flask was added EXAMPLE 232A (0.15 g) and zinc (1 g) in water/methanol (1:1, 10 mL) to give a suspension. The mixture was cooled to 0°C. 12N Aqueous HCl (2 mL) was added slowly, and the mixture was stirred at 0 °C for 30 minutes. 2N Aqueous NaOH solution was used to adjust the mixture to basic pH. The mixture was filtered, and extracted with ether (3x 30 mL). After drying over Na₂SO₄, filtration, and concentration, the crude product was added to a silica gel column (Analogix, SF15-12g,) and purified by eluting with 0-25% ethyl acetate in hexane.

EXAMPLE 232C

tert-butyl 4-(2-nitro-4-sulfamoylphenylamino)piperazine-1-carboxylate

[0933] The title compound was prepared by substituting EXAMPLE 232B for 1-(tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 232D

tert-butyl 4-[[4-[[4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino]piperazine-1-carboxylate

[0934] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 232C for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.47 (br. s, 1H), 8.86 (s, 1H), 8.34 (d, 1H), 7.90 (d, 1H), 7.59 (m, 2H), 7.36 (m, 4H), 7.23 (m, 1H), 7.05 (d, 2H), 6.61 (dd, 1H), 6.27 (m, 2H), 2.99 (m, 5H), 2.76 (m, 6H), 2.19 (m, 6H), 1.96 (s, 2H), 1.41 (m, 11H), 1.24 (m, 4H), 0.93 (s, 6H).

EXAMPLE 233

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-(pentafluoro-λ⁶-sulfonyl)-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 233A

2-(5-bromo-2-nitrophenyl)sulfur pentafluoride

[0935] To a solution of 3-bromophenylsulfur pentafluoride (2.18 g) in concentrated H₂SO₄ (5 mL) was added KNO₃ (780 mg). The mixture was stirred overnight. The mixture was diluted with diethyl ether (100 mL) and washed with water and brine. After drying over Na₂SO₄ and filtration, the solvent was evaporated under vacuum to provide the title compound.

EXAMPLE 233B

2-(5-bromo-2-aminophenyl)sulfur pentafluoride

[0936] EXAMPLE 233A (6.4g) and tetrahydrofuran (300 mL) were added to Ra-Ni, (12.80 g) in a 50 mL pressure bottle and the mixture stirred for 2 hours at 30 psi and room temperature. The mixture was filtered through a nylon membrane and the filtrate was concentrated under vacuum to provide the title compound.

EXAMPLE 233C

4-bromo-2-pentafluorosulfonyl-N-(tetrahydro-2H-pyran-4-ylmethyl)aniline

[0937] To a solution of EXAMPLE 233B (4.4 g) in methanol (50 mL) was added tetrahydro-2H-pyran-4-carbaldehyde (1.68 g) and decaborane (1.1 g). The mixture was stirred and monitored by thin layer chromatography. More tetrahydro-2H-pyran-4-carbaldehyde (500 mg) was added to the stirring mixture to drive the reaction to completion. The reaction mixture was concentrated under vacuum and ethyl acetate (500 mL) and brine (200 mL) were added. The organic phase was dried over Na₂SO₄. Filtration and evaporation of the solvent and flash chromatography (20% ethyl acetate in hexane) gave the title compound.

EXAMPLE 233D

4-thioacetoxyl-2-pentafluorosulfonyl-N-(tetrahydro-2H-pyran-4-ylmethyl)aniline

[0938] To a solution of EXAMPLE 233C (456 mg) and potassium ethanethioate (197 mg) in dioxane (4 mL) was added tris(dibenzylideneacetone)dipalladium(0) (27 mg) and xantphos (33 mg) followed by N,N-diisopropylethylamine (0.5 mL). The mixture was purged with argon, sealed and stirred under microwave irradiation for 60 minutes at 120°C. The mixture was dissolved in ethyl acetate (300 mL) and water (100 mL). The organic phase was washed with brine and dried over Na₂SO₄. Filtration and evaporation of the solvent followed by flash chromatography (20% ethyl acetate in hexane) provided the title compound.

EXAMPLE 233E

3-pentafluorosulfonyl-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenylsulfonamide

[0939] N-chlorosuccinimide (527 mg) was added to a mixture of 2N aqueous HCl (1.5 mL) and acetonitrile (12 mL)

and then cooled to 0°C. A solution of EXAMPLE 233D (386 mg) in acetonitrile (3 mL) was added to the mixture which was then stirred at 0°C for 2 hours, and then diluted with ethyl acetate (300 mL) and washed with brine and dried over Na₂SO₄. After filtration and evaporation of solvent, the residue was dissolved in isopropyl alcohol (20 mL) and cooled to 0°C with stirring. Then, ammonium hydroxide (conc. 10 mL) was added to mixture. After stirring for 2 hours, the mixture was concentrated under vacuum and the residue was added to ethyl acetate (400 mL) and water (150 mL). The organic layer was washed with brine and dried over Na₂SO₄. After filtration and evaporation of solvent, the residue was purified by flash column (20% ethyl acetate in dichloromethane) to provide the title compound.

EXAMPLE 233F

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({3-(pentafluoro-λ⁶-sulfonyl)-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0940] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 233E for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.71 (s, 1 H), 11.33 (m, 1 H), 8.12 (m, 2 H), 7.72 (d, 1 H), 7.54 (m, 3 H), 7.33 (m, 2 H), 7.02 (m, 3 H), 6.67 (m, 2 H), 6.42 (m, 1 H), 6.16 (d, 1 H), 3.82 (m, 2 H), 3.21 (m, 4 H), 3.05 (m, 4 H), 2.73 (s, 2 H), 2.21 (m, 8 H), 1.97 (m, 3 H), 1.29 (m, 4 H), 0.92 (s, 6 H).

EXAMPLE 234

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({4-[(4-methoxytetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 234A

4-vinyltetrahydro-2H-pyran-4-ol

[0941] Dihydro-2H-pyran-4(3H)-one (8.01 g) in anhydrous ethyl ether (50 mL) was treated with 1.0 M vinylmagnesium bromide (104 mL) over 20 minutes at 0°C. The reaction mixture was stirred at room temperature overnight. The reaction was quenched with saturated NH₄Cl, and the organic layer was separated. The aqueous layer was extracted with additional ethyl ether three times. The combined organic layers were washed with brine, dried, filtered, and concentrated. The residue was purified by flash column chromatography on silica gel using 20% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 234B

4-methoxy-4-vinyltetrahydro-2H-pyran

[0942] To a solution of EXAMPLE 234A (9.4 g) in tetrahydrofuran (150 mL) was added 60 % sodium hydride (5.28 g) at 0°C portionwise. After the addition was complete, the solution was heated under reflux for three hours. After cooling, to this suspension was added dimethyl sulfate (8.41 mL) slowly. The solution was heated under reflux overnight, cooled to room temperature, and hydrolyzed with cool saturated aqueous NH₄Cl. After extraction with diethyl ether several times, the combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified by flash column chromatograph on silica gel using 1-10% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 234C

4-methoxytetrahydro-2H-pyran-4-carbaldehyde

[0943] EXAMPLE 234B (4.3 g) in tetrahydrofuran (200 mL) and water (67 mL) was treated with 4% osmium tetroxide in water (9.24 mL). To this solution was added potassium periodate (13.91 g) portionwise over 2 hours. The solution was stirred overnight at room temperature. Water was added to the mixture followed by repeat extractions with diethyl ether. The combined organic layers were dried over MgSO₄, filtered, and concentrated. The residue was purified by flash column chromatography on silica gel using 5-20% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 234D

(4-methoxytetrahydro-2H-pyran-4-yl)methanol

5 **[0944]** EXAMPLE 234C (1.8 g) in 2-propanol (28 mL) and water (7 mL) was cooled to 0 °C. To this solution was added sodium borohydride (0.709 g). The solution was stirred and allowed to warm to room temperature over 3 hours. The reaction was quenched with acetone, and stirred for another 1 hour. The clear liquid was separated from solid by decanting. Additional ethyl acetate was used to wash the solid, and the mixture was decanted. The combined organic solutions were concentrated. The residue was purified by flash chromatography on silica gel eluting 1:1 ethyl
10 acetate: hexane to provide the title compound.

EXAMPLE 234E

4-((4-methoxytetrahydro-2H-pyran-4-yl)methoxy)-3-nitrobenzenesulfonamide

15 **[0945]** The title compound was prepared by substituting EXAMPLE 234D for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 234F

20 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((4-methoxytetrahydro-2H-pyran-4-yl)methoxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

25 **[0946]** The title compound was prepared by substituting EXAMPLE 234E for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 8.36 (s, 1H), 8.04-8.07 (m, 2H), 7.50-7.53 (m, 3H), 7.45 (d, 1H), 7.35 (d, 2H), 7.05 (d, 2H), 6.68 (dd, 1H), 6.40 (dd, 1H), 6.20 (d, 1H), 4.21 (s, 2H), 3.65-3.67 (m, 2H), 3.53-3.56 (m, 2H), 3.19 (s, 3H), 3.10 (s, 4H), 2.86 (s, 2H), 2.30 (s, 4H), 2.15 (s, 2H), 1.96 (s, 2H), 1.61-1.74 (m, 4H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 235

30 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 235A

35 (R)-*tert*-butyl 3-(2-nitro-4-sulfamoylphenoxy)pyrrolidine-1-carboxylate

[0947] The title compound was prepared by substituting (R)-*tert*-butyl 3-hydroxypyrrolidine-1-carboxylate for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

40

EXAMPLE 235B

(R)-*tert*-butyl 3-(4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)sulfamoyl)-2-nitrophenoxy)pyrrolidine-1-carboxylate

45

[0948] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 235A for EXAMPLE 1F in EXAMPLE 1G.

EXAMPLE 235C

50

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

55 **[0949]** To a solution of EXAMPLE 235B (0.230 g) in dichloromethane (3 mL) was added trifluoroacetic acid (0.377 mL). After stirring for 4 hours, the reaction was concentrated then dissolved in dichloromethane (3 mL) and treated with 1,3-difluoropropan-2-one (0.028 g) followed by sodium triacetoxymethylborohydride (0.078 g). After stirring for 4 hours, the reaction was quenched by the addition of saturated aqueous NaHCO₃ and dichloromethane (5 mL). The reaction was diluted with dichloromethane (250 mL) and saturated aqueous NaHCO₃ (100 mL) was added. The organic layer was

separated, washed with brine (100 mL), dried over magnesium sulfate, filtered and concentrated. Trituration with acetonitrile gave the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-*d*₆) δ 11.67 (s, 1H), 8.34 (s, 1H), 8.03 (s, 2H), 7.52 (d, 3H), 7.35 (d, 3H), 7.04 (d, 2H), 6.75 - 6.60 (m, 1H), 6.40 (s, 1H), 6.20 (s, 1H), 5.17 - 5.06 (m, 1H), 4.60 (d, 4H), 2.98 (d, 12H), 2.37 - 2.02 (m, 6H), 1.96 (s, 3H), 1.39 (s, 2H), 0.93 (s, 6H).

EXAMPLE 236

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(3-nitro-4-[(4-(oxetan-3-yl)piperazin-1-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-*b*]pyridin-5-yloxy)benzamide

EXAMPLE 236A

2-(1H-pyrrolo[2,3-*b*]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-(3-nitro-4-(piperazin-1-ylamino)phenyl)sulfonylbenzamide

[0950] The title compound was prepared by substituting EXAMPLE 232D for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 236B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(3-nitro-4-[(4-(oxetan-3-yl)piperazin-1-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-*b*]pyridin-5-yloxy)benzamide

[0951] The title compound was prepared by substituting EXAMPLE 236A for tert-butyl piperazine-1-carboxylate and oxetan-3-one for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A. ¹H NMR (400 MHz, dimethylsulfoxide-*d*₆) δ 11.67 (br. s, 1H), 9.20 (s, 1H), 8.53 (d, 1H), 8.04 (d, 1H), 7.83 (dd, 1H), 7.53 (m, 4H), 7.34 (d, 2H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.39 (m, 1H), 6.18 (d, 1H), 4.55 (t, 2H), 4.44 (t, 2H), 3.47 (m, 1H), 3.06 (m, 4H), 2.88 (m, 4H), 2.74 (m, 4H), 2.09 (m, 11H), 1.38 (t, 2H), 0.91 (s, 6H).

EXAMPLE 237

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(3-nitro-4-[(4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-*b*]pyridin-5-yloxy)benzamide

[0952] The title compound was prepared by substituting EXAMPLE 236A for tert-butyl piperazine-1-carboxylate and dihydro-2H-pyran-4(3H)-one for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A. ¹H NMR (400 MHz, pyridine-*d*₅) δ 13.05 (br. s, 1H), 9.27 (d, 1H), 9.23 (s, 1H), 8.44 (m, 2H), 8.12 (d, 1H), 7.68 (m, 3H), 7.44 (m, 2H), 7.06 (m, 2H), 6.75 (dd, 1H), 6.51 (m, 2H), 4.02 (m, 2H), 3.31 (m, 2H), 3.06 (m, 4H), 2.91 (m, 5H), 2.76 (s, 2H), 2.38 (m, 2H), 2.26 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.65 (m, 2H), 1.39 (m, 7H), 0.93 (s, 6H).

EXAMPLE 238

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(3-nitro-4-[(3R)-tetrahydrofuran-3-ylamino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-*b*]pyridin-5-yloxy)benzamide

EXAMPLE 238A

(R)-3-nitro-4-(tetrahydrofuran-3-ylamino)benzenesulfonamide

[0953] The title compound was prepared by substituting (R)-tetrahydrofuran-3-amine for 4-methylpiperazin-1-amine dihydrochloride in EXAMPLE 6A.

EXAMPLE 238B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(3-nitro-4-[(3R)-tetrahydrofuran-3-ylamino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-*b*]pyridin-5-yloxy)benzamide

[0954] The title compound was prepared by substituting EXAMPLE 238A for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (400 MHz, dimethylsulfoxide-*d*₆) δ 11.59 (s, 1H), 8.47 (d, 1H), 8.19 (m, 2H),

7.97 (d, 1H), 7.74 (m, 1H), 7.52 (d, 1H), 7.46 (t, 1H), 7.34 (m, 2H), 7.05 (m, 2H), 6.96 (d, 1H), 6.89 (d, 1H), 6.65 (dd, 1H), 6.33 (m, 1H), 6.22 (d, 1H), 4.31 (m, 1H), 3.92 (m, 1H), 3.87 (m, 1H), 3.76 (m, 1H), 3.69 (m, 1H), 3.04 (m, 4H), 2.73 (m, 2H), 2.33 (m, 1H), 2.18 (m, 6H), 1.95 (m, 2H), 1.88 (m, 1H), 1.39 (t, 2H), 0.92 (s, 6H).

5 EXAMPLE 239

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(4-((4,4-difluorocyclohexyl)methyl)amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 EXAMPLE 239A

tert-butyl (4,4-difluorocyclohexyl)methylcarbamate

[0955] Tert-butyl (4-oxocyclohexyl)methylcarbamate (5 g) and diethylaminosulfur trifluoride (7.45 g) were stirred in dichloromethane (100 mL) for 24 hours. The mixture was quenched with pH 7 buffer (100 mL), and poured into ether (400 mL). The resulting solution was separated, and the organic layer was washed twice with water, and once with brine, and then concentrated to give the crude product and fluoroolefin by-product in a 3:2 ratio. The crude material was taken up in tetrahydrofuran (70 mL) and water (30 mL), and N-methylmorpholine-N-oxide (1.75 g), and OsO₄ (2.5 wt % solution in t-butanol) were added, and the mixture was stirred for 24 hours. Na₂S₂O₃ (10 g) was then added, and the mixture was stirred for 30 minutes. The mixture was then diluted with ether (300 mL), and the resulting solution was separated, and rinsed twice with water, and once with brine, and concentrated. The crude product was chromatographed on silica gel using 5-10% ethyl acetate in hexanes to provide the title compound.

25 EXAMPLE 239B

(4,4-difluorocyclohexyl)methanamine

[0956] A solution of EXAMPLE 239A (3 g) in dichloromethane (35 mL), trifluoroacetic acid (15 mL), and triethylsilane (1 mL) was stirred for 2 hours. The solution was concentrated, then concentrated from toluene, and left on high vacuum for 24 hours. The semi-solid was taken up in ether/hexane and filtered to provide the title compound as its trifluoroacetic acid salt.

EXAMPLE 239C

35 4-((4,4-difluorocyclohexyl)methylamino)-3-nitrobenzenesulfonamide

[0957] The title compound was prepared by substituting EXAMPLE 239B for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

40 EXAMPLE 239D

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[(4-((4,4-difluorocyclohexyl)methyl)amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0958] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 239C for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 12.40 (s, 1H), 11.61 (br s, 1H), 8.53 (m, 1H), 8.50 (d, 1H), 7.99 (d, 1H), 7.73 (d, 1H), 7.49 (m, 2H), 7.32 (d, 2H), 7.04 (d, 2H), 7.00 (d, 1H), 6.65 (d, 1H), 6.32 (s, 1H), 6.21 (s, 1H), 3.37 (m, 4H), 3.06 (m, 4H), 2.73 (m, 2H), 2.18 (m, 4H), 1.97 (m, 4H), 1.81 (m, 4H), 1.38 (m, 2H), 1.20 (m, 4H), 0.92 (s, 6H).

50

55

EXAMPLE 240

N-({4-[(1-tert-butylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 240A

4-(1-tert-butylpiperidin-4-ylamino)-3-(trifluoromethylsulfonyl)benzenesulfonamide

[0959] To a mixture of 4-fluoro-3-(trifluoromethylsulfonyl)benzenesulfonamide (307 mg) and EXAMPLE 227A (156 mg) in tetrahydrofuran (4 mL) was added Hunig's Base (1 mL). The mixture was stirred for 3 days. The mixture was diluted with ethyl acetate (300 mL) and water (100 mL) and stirred until the solid disappeared into the solution. The layers were separated and the organic phase was washed with water, brine and dried over Na₂SO₄. After filtration, the combined aqueous layers were extracted again with ethyl acetate and the combined organic phase was dried over Na₂SO₄. After filtration, the solvent was evaporated to provide the title compound.

EXAMPLE 240B

N-({4-[(1-tert-butylpiperidin-4-yl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl}sulfonyl)-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0960] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 240A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.53 (s, 1 H), 8.04 (s, 1 H), 7.94 (d, 1 H), 7.86 (m, 1 H), 7.55 (d, 2 H), 7.44 (d, 1 H), 7.33 (m, 3 H), 7.05 (d, 2 H), 6.92 (m, 1 H), 6.62 (dd, 1 H), 6.43 (m, 1 H), 6.29 (d, 2 H), 3.79 (m, 1 H), 3.05 (m, 6 H), 2.73 (s, 3 H), 2.19 (m, 8 H), 1.96 (s, 3 H), 1.27 (m, 12 H), 0.92 (s, 6 H).

EXAMPLE 241

4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({4-[(4-(oxetan-3-yl)morpholin-2-yl)methyl]amino}-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 241 A

tert-butyl 2-((4-sulfamoyl-2-(trifluoromethylsulfonyl)phenylamino)methyl)morpholine-4-carboxylate

[0961] The title compound was prepared by substituting tert-butyl 2-(aminomethyl)morpholine-4-carboxylate for (tetrahydropyran-4-yl)methylamine and 4-fluoro-3-(trifluoromethylsulfonyl)benzenesulfonamide for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 1F.

EXAMPLE 241B

tert-butyl 2-((4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)benzoyl)sulfamoyl)-2-(trifluoromethylsulfonyl)phenylamino)methyl)morpholine-4-carboxylate

[0962] The title compound was prepared by substituting EXAMPLE 241A for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G.

EXAMPLE 241C

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-{4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl}-N-(4-(morpholin-2-ylmethylamino)-3-(trifluoromethylsulfonyl)phenylsulfonyl)benzamide

[0963] The title compound was prepared by substituting EXAMPLE 241B for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 241D

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({4-([4-(oxetan-3-yl)morpholin-2-yl]methyl)amino}-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

5

[0964] The title compound was prepared by substituting EXAMPLE 241C for tert-butyl piperazine-1-carboxylate and oxetan-3-one for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 8.15 (d, 1H), 8.04 (d, 1H), 7.92 (dd, 1H), 7.54 (d, 1H), 7.51 (t, 1H), 7.48 (d, 1H), 7.34 (d, 2H), 7.25 (m, 1H), 7.04 (m, 3H), 6.68 (dd, 1H), 6.41 (m, 1H), 6.19 (d, 1H), 4.54 (t, 2H), 4.43 (m, 2H), 3.85 (m, 1H), 3.69 (m, 1H), 3.52 (m, 1H), 3.48 (m, 1H), 3.39 (m, 2H), 3.07 (m, 4H), 2.77 (br s, 2H), 2.69 (d, 1H), 2.56 (d, 1H), 2.21 (br s, 4H), 2.15 (t, 2H), 1.94 (m, 3H), 1.76 (t, 1H), 1.38 (t, 2H), 0.92 (s, 6H).

10

EXAMPLE 242

N-([5-chloro-6-([4-fluorotetrahydro-2H-pyran-4-yl)methyl]amino)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

15

EXAMPLE 242A

5-chloro-6-([4-fluorotetrahydro-2H-pyran-4-yl)methylamino)pyridine-3-sulfonamide

20

[0965] The title compound was prepared by substituting EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 138D.

25

EXAMPLE 242B

N-([5-chloro-6-([4-fluorotetrahydro-2H-pyran-4-yl)methyl]amino)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

30

[0966] The title compound was prepared by substituting EXAMPLE 242A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.71 (s, 1H), 8.41 (d, 1H), 8.07 (d, 1H), 7.93 (d, 1H), 7.60 (d, 1H), 7.51-7.53 (m, 2H), 7.40 (s, 1H), 7.33-7.35 (m, 2H), 7.03-7.05 (m, 2H), 6.68 (dd, 1H), 6.42 (dd, 1H), 6.16 (d, 1H), 3.77 (d, 1H), 3.69-3.71 (m, 3H), 3.48-3.53 (m, 2H), 3.07 (s, 4H), 2.76 (s, 2H), 2.14-2.20 (m, 6H), 1.96 (s, 2H), 1.65-1.76 (m, 4H), 1.38 (t, 2H), 0.93 (s, 6H).

35

EXAMPLE 243

N-([5-chloro-6-([1-cyclopropylpiperidin-4-yl]amino)pyridin-3-yl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

40

EXAMPLE 243A

5-chloro-6-(1-cyclopropylpiperidin-4-ylamino)pyridine-3-sulfonamide

45

[0967] The title compound was prepared by substituting EXAMPLE 40A for 4-chloro-3-nitrobenzenesulfonamide, 1-cyclopropylpiperidin-4-amine for 4-methylpiperazin-1-amine dihydrochloride and Hunig's base for N¹,N¹,N²,N²-tetramethylethane-1,2-diamine in EXAMPLE 6A.

EXAMPLE 243B

N-([5-chloro-6-([1-cyclopropylpiperidin-4-yl]amino)pyridin-3-yl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50

[0968] The title compound was prepared by substituting EXAMPLE 243A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 8.40 (d, 1H), 8.05 (d, 1H), 7.88 (d, 1H), 7.56 (d, 1H), 7.50 (m, 2H), 7.34 (d, 2H), 7.03 (d, 2H), 6.97 (br d, 1H), 6.66 (dd, 1H), 6.40 (m, 1H), 6.16 (d, 1H), 4.04 (m, 1H), 3.03 (br m, 6H), 2.73 (s, 2H), 2.42 (br m, 2H), 2.18 (br m, 6H), 1.95 (s, 2H), 1.80 (m, 3H), 1.62 (m, 2H), 1.38 (t, 2H), 0.91 (s, 6H), 0.47 (m, 2H), 0.40 (br m, 2H).

EXAMPLE 244

N-[(5-chloro-6-([(2S)-4-(cyanomethyl)morpholin-2-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 244A

(S)-tert-butyl 2-((3-chloro-5-sulfamoylpyridin-2-yloxy)methyl)morpholine-4-carboxylate

[0969] The title compound was prepared by substituting (S)-tert-butyl 2-(hydroxymethyl)-morpholine-4-carboxylate for tetrahydro-2H-pyran-4-yl-methanol and EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 244B

(S)-5-chloro-6-(morpholin-2-ylmethoxy)pyridine-3-sulfonamide

[0970] The title compound was prepared by substituting EXAMPLE 244A for EXAMPLE 113A in EXAMPLE 134A.

EXAMPLE 244C

(S)-5-chloro-6-((4-(cyanomethyl)morpholin-2-yl)methoxy)pyridine-3-sulfonamide The title compound was prepared by substituting EXAMPLE 244B for EXAMPLE 134A and 2-bromoacetonitrile for methyl iodide in EXAMPLE 134B.

EXAMPLE 244D

N-[(5-chloro-6-([(2S)-4-(cyanomethyl)morpholin-2-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0971] The title compound was prepared by substituting EXAMPLE 244C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 12.99 (s, 1H), 9.09 (d, 1H), 8.70 (d, 1H), 8.42 (d, 1H), 8.12 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.75 (dd, 1H), 6.53 (d, 1H), 6.48 (m, 1H), 4.55 (dd, 1H), 4.43 (dd, 1H), 4.05 (m, 1H), 3.85 (d, 1H), 3.76 (s, 2H), 3.63 (dt, 1H), 3.06 (m, 4H), 2.91 (d, 1H), 2.77 (s, 2H), 2.58 (d, 1H), 2.51-2.44 (m, 2H), 2.26 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 245

N-[(5-chloro-6-([(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 245A

(S)-5-chloro-6-((4-(2-(dimethylamino)acetyl)morpholin-2-yl)methoxy)pyridine-3-sulfonamide

[0972] The title compound was prepared by substituting EXAMPLE 244B for EXAMPLE 134A and 2-(dimethylamino)acetyl chloride hydrochloride for methyl iodide in EXAMPLE 134B.

EXAMPLE 245B

N-[(5-chloro-6-([(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0973] The title compound was prepared by substituting EXAMPLE 245A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.09 (d, 1H), 8.69 (s, 1H), 8.42 (s, 1H), 8.11 (t, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.76 (s, 1H), 6.54 (s, 1H), 6.49 (s, 1H), 4.85-4.46 (m, 3H), 4.45-3.87 (m, 3H), 3.50 (m, 1H), 3.37 (dd, 1H), 3.21 (m, 2H), 3.07 (m, 4H), 2.86 (t, 1H), 2.77 (s, 2H), 2.27 (m, 8H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 246

N-[(5-chloro-6-([(2R)-4-(cyanomethyl)morpholin-2-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 246A

(R)-tert-butyl 2-((3-chloro-5-sulfamoylpyridin-2-yloxy)methyl)morpholine-4-carboxylate

[0974] The title compound was prepared by substituting (R)-tert-butyl 2-(hydroxymethyl)-morpholine-4-carboxylate for tetrahydro-2H-pyran-4-yl-methanol and EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 246B

(R)-5-chloro-6-(morpholin-2-ylmethoxy)pyridine-3-sulfonamide

[0975] The title compound was prepared by substituting EXAMPLE 246A for EXAMPLE 113A in EXAMPLE 134A.

EXAMPLE 246C

(R)-5-chloro-6-((4-(cyanomethyl)morpholin-2-yl)methoxy)pyridine-3-sulfonamide

[0976] The title compound was prepared by substituting EXAMPLE 246B for EXAMPLE 134A and 2-bromoacetonitrile for methyl iodide in EXAMPLE 134B.

EXAMPLE 246D

N-[(5-chloro-6-([(2R)-4-(cyanomethyl)morpholin-2-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0977] The title compound was prepared by substituting EXAMPLE 246C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 12.99 (s, 1H), 9.09 (d, 1H), 8.70 (d, 1H), 8.42 (d, 1H), 8.12 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.75 (dd, 1H), 6.53 (d, 1H), 6.48 (m, 1H), 4.55 (dd, 1H), 4.43 (dd, 1H), 4.05 (m, 1H), 3.85 (d, 1H), 3.76 (s, 2H), 3.63 (dt, 1H), 3.06 (m, 4H), 2.91 (d, 1H), 2.77 (s, 2H), 2.58 (d, 1H), 2.51-2.44 (m, 2H), 2.26 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 247

N-[(5-chloro-6-([(2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 247A

(R)-5-chloro-6-((4-(2-(dimethylamino)acetyl)morpholin-2-yl)methoxy)pyridine-3-sulfonamide

[0978] The title compound was prepared by substituting EXAMPLE 246B for EXAMPLE 134A and 2-(dimethylamino)acetyl chloride hydrochloride for methyl iodide in EXAMPLE 134B.

EXAMPLE 247B

N-[(5-chloro-6-([(2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0979] The title compound was prepared by substituting EXAMPLE 247A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.09 (d, 1H), 8.69 (s, 1H), 8.42 (s, 1H), 8.11 (t, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.76 (s, 1H), 6.54 (s, 1H), 6.49 (s, 1H), 4.85-4.46 (m, 3H), 4.45-3.87 (m, 3H), 3.50 (m, 1H), 3.37 (dd, 1H), 3.21 (m, 2H), 3.07 (m, 4H), 2.86 (t, 1H), 2.77 (s, 2H), 2.27 (m, 8H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 248

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({5-fluoro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 248A

5-bromo-3-fluoro-2-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)pyridine

[0980] The title compound was prepared by substituting 5-bromo-2,3-difluoropyridine for 4-fluoro-3-nitrobenzenesulfonamide and EXAMPLE 37C for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 248B

tert-butyl 5-fluoro-6-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)pyridin-3-ylcarbamate

[0981] EXAMPLE 248A (0.308 g), tert-butyl carbamate (0.141 g), palladium(II) acetate (0.011 g), Xantphos (0.043 g) and cesium carbonate (0.489 g) were combined with dioxane (5.0 mL) in a 20-mL vial equipped with a magnetic stir bar. The vial was flushed with nitrogen, capped and stirred at 100°C overnight. Additional palladium(II) acetate (0.011 g), Xantphos (0.043 g) and tert-butyl carbamate (0.141 g) were added and heating was continued at 100°C for 8 hours. The cooled reaction mixture was diluted with ethyl acetate, washed with water and brine, dried (MgSO₄) filtered and concentrated. The concentrate was chromatographed on silica gel with 7-25% ethyl acetate in hexanes as the eluent.

EXAMPLE 248C

5-fluoro-6-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)pyridine-3-sulfonyl chloride

[0982] Under ice-cooling, thionyl chloride (1.563 mL) was added dropwise over 20 minutes to water (9 mL). The mixture was stirred for 12 hours to give a SO₂-containing solution. Separately, EXAMPLE 248B (0.295 g) was added to a mixture of 1,4-dioxane (3.2 mL) and concentrated HCl (8 mL) at 0°C. After stirring for 15 minutes, a solution of sodium nitrite (0.065 g) in water (2 mL) was added dropwise and stirring was continued at 0°C for 3 hours. Copper(I) chloride (0.042 g) and then the freshly prepared solution of diazotized material were added sequentially to the previously prepared SO₂-containing solution. The resulting solution was stirred for 30 minutes and then extracted with ethyl acetate (2 x 125 mL). The combined extracts were dried (MgSO₄) filtered and concentrated. The concentrate was chromatographed on silica gel with 5% ethyl acetate in hexanes as the eluent.

EXAMPLE 248D

5-fluoro-6-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)pyridine-3-sulfonamide

[0983] EXAMPLE 248C (0.08 g) in isopropanol (2 mL) at 0°C was treated with ammonium hydroxide (1.697 mL), stirred overnight and then concentrated to dryness. The obtained solid was slurried in water, filtered, rinsed with water and dried under high vacuum to provide the title compound.

EXAMPLE 248E

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({5-fluoro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0984] The title compound was prepared by substituting EXAMPLE 248D for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, pyridine-d₅) δ 13.05 (s, 1H), 9.03 (d, 1H), 8.44 (dd, 1H), 8.41 (d, 1H), 8.10 (d, 1H), 7.67 (m, 1H), 7.65 (d, 1H), 7.44 (m, 2H), 7.07 (m, 2H), 6.77 (dd, 1H), 6.53 (d, 1H), 6.49 (dd, 1H), 4.55 (d, 2H), 3.80 (m, 4H), 3.08 (m, 4H), 2.77 (s, 2H), 2.26 (t, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.88 (m, 4H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 250

N-({5-chloro-6-[3-hydroxy-2-(hydroxymethyl)-2-methylpropoxy]pyridin-3-yl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 250A

5-chloro-6-((3-methyloxetan-3-yl)methoxy)pyridine-3-sulfonamide

[0985] The title compound was prepared by substituting (3-methyloxetan-3-yl)methanol for (tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 250B

N-({5-chloro-6-[3-hydroxy-2-(hydroxymethyl)-2-methylpropoxy]pyridin-3-yl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0986] The title compound was prepared by substituting EXAMPLE 250A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, pyridine-d₅) δ 13.05 (s, 1H), 9.22 (d, 1H), 8.51 (d, 1H), 8.42 (d, 1H), 8.09 (d, 1H), 7.66 (t, 2H), 7.43 - 7.46 (m, 2H), 7.04 - 7.09 (m, 2H), 6.75 (dd, 1H), 6.45 - 6.54 (m, 2H), 4.47 (s, 2H), 3.81 - 3.84 (m, 2H), 3.74 (d, 2H), 3.03 - 3.11 (m, 4H), 2.77 (s, 2H), 2.26 (t, 2H), 2.10 - 2.17 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 1.16 (s, 3H), 0.94 (s, 6H).

EXAMPLE 251

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 251 A

5-bromo-6-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)pyridine-3-sulfonamide

[0987] The title compound was prepared by substituting (4-fluorotetrahydro-2H-pyran-4-yl)methanol for (tetrahydro-2H-pyran-4-yl)methanol and 5-bromo-6-chloropyridine-3-sulfonamide for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 251B

6-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)pyridine-3-sulfonamide

[0988] To a suspension of 5-bromo-6-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)pyridine-3-sulfonamide (200 mg) and cyclohexene (0.549 mL) in ethyl acetate (10 mL) was added 10% palladium on carbon (57.6 mg). The suspension was stirred for 60 minutes at 120°C. The reaction mixture was filtered and concentrated. The product was purified by reverse-phase flash chromatography (C18, 150g, 10%-100% acetonitrile/H₂O/trifluoroacetic acid 0.1%).

EXAMPLE 251C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0989] The title compound was prepared by substituting EXAMPLE 251B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, pyridine-d₅) δ 13.06 (s, 1H), 9.29 (d, 1H), 8.50 (dd, 1H), 8.41 (d, 1H), 8.07 (d, 1H), 7.66 - 7.70 (m, 1H), 7.64 (d, 1H), 7.44 (d, 2H), 7.07 (d, 2H), 6.84 (d, 1H), 6.75 (dd, 1H), 6.52 (d, 1H), 6.49 (dd, 1H), 4.50 (d, 2H), 3.81 - 3.89 (m, 2H), 3.70 - 3.81 (m, 2H), 3.02 - 3.12 (m, 4H), 2.77 (s, 2H), 2.26 (t, 2H), 2.10 - 2.18 (m, 4H), 1.97 (s, 2H), 1.77 - 1.94 (m, 4H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 252

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{[4-({[4-(1,3-difluoropropan-2-yl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 252A

tert-butyl 4-(1,3-difluoropropan-2-yl)morpholin-2-yl)methylcarbamate

[0990] The title compound was prepared by substituting 1,3-difluoropropan-2-one for 4'-chlorobiphenyl-2-carboxaldehyde and tert-butyl morpholin-2-ylmethylcarbamate for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A.

EXAMPLE 252B

4-(1,3-difluoropropan-2-yl)morpholin-2-yl)methanamine

[0991] A solution of EXAMPLE 252A (538 mg) in dioxane (4 mL) was treated with 4.0M HCl in dioxane solution (1.8 mL). The reaction was stirred at room temperature overnight. The reaction mixture was concentrated under vacuum and used without further purification.

EXAMPLE 252C

4-((4-(1,3-difluoropropan-2-yl)morpholin-2-yl)methylamino)-3-nitrobenzenesulfonamide

[0992] The title compound was prepared by substituting EXAMPLE 252B for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 252D

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{[4-({[4-(1,3-difluoropropan-2-yl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[0993] The title compound was prepared by substituting EXAMPLE 252C for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.64 (s, 1H), 8.59 (t, 1H), 8.57 (d, 1H), 8.04 (d, 1H), 7.83 (dd, 1H), 7.51 (m, 3H), 7.33 (d, 2H), 7.07 (d, 1H), 7.03 (d, 2H), 6.66 (dd, 1H), 6.39 (m, 1H), 6.19 (d, 1H), 4.69 (t, 2H), 4.57 (t, 2H), 3.85 (m, 1H), 3.70 (m, 1H), 3.52 (m, 2H), 3.41 (m, 2H), 3.07 (br s, 4H), 2.91 (d, 1H), 2.74 (m, 3H), 2.59 (m, 1H), 2.43 (m, 1H), 2.20 (m, 4H), 2.15 (m, 2H), 1.95 (br s, 2H), 1.39 (t, 2H), 0.92 (s, 6H).

EXAMPLE 253

N-{{[5-chloro-6-{{[1-(cyanomethyl)piperidin-4-yl]methoxy}pyridin-3-yl]sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 253A

tert-butyl 4-((3-chloro-5-sulfamoylpyridin-2-yloxy)methyl)piperidine-1-carboxylate

[0994] The title compound was prepared by substituting tert-butyl 4-(hydroxymethyl)piperidine-1-carboxylate for tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 253B

5-chloro-6-(piperidin-4-ylmethoxy)pyridine-3-sulfonamide ditrifluoroacetic acid

[0995] The title compound was prepared by substituting EXAMPLE 253A for EXAMPLE 39A in EXAMPLE 39B.

EXAMPLE 253C

5-chloro-6-((1-(cyanomethyl)piperidin-4-yl)methoxy)pyridine-3-sulfonamide

- 5 **[0996]** EXAMPLE 253B (0.061 g), 2-chloroacetonitrile (0.017 g), sodium carbonate (0.025 g) and N,N-dimethylformamide (1 mL) were combined in a 4-mL vial and heated at 60°C overnight. The cooled reaction mixture was diluted with ethyl acetate, washed with water and brine, dried (MgSO₄) filtered and concentrated. The concentrate was chromatographed on silica gel with 2-10% methanol in CH₂Cl₂ as the eluent.

10 EXAMPLE 253D

N-[(5-chloro-6-[[1-(cyanomethyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- 15 **[0997]** The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 253C for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (400 MHz, pyridine-d₅) δ 13.04 (s, 1H), 9.14 (d, 1H), 8.41 (d, 1H), 8.10 (d, 1H), 7.66 (t, 2H), 7.44 (m, 2H), 7.07 (m, 2H), 6.74 (dd, 1H), 6.50 (m, 2H), 4.18 (d, 2H), 3.64 (s, 2H), 3.05 (s, 4H), 2.77 (m, 4H), 2.24 (m, 4H), 2.13 (m, 4H), 1.97 (s, 2H), 1.69 (m, 3H), 1.41 (m, 4H), 0.93 (s, 6H).

20 EXAMPLE 254

4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-[[4-((3R)-1-[2-(2-methoxyethoxy)ethyl]pyrrolidin-3-yl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

25 EXAMPLE 254A

(R)-tert-butyl 3-(2-nitro-4-sulfamoylphenylamino)pyrrolidine-1-carboxylate

- 30 **[0998]** The title compound was prepared by substituting (R)-tert-butyl 3-aminopyrrolidine-1-carboxylate for 1-(tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 254B

(R)-3-nitro-4-(pyrrolidin-3-ylamino)benzenesulfonamide

- 35 **[0999]** The title compound was prepared by substituting EXAMPLE 254A for EXAMPLE 113A in EXAMPLE 134A.

EXAMPLE 254C

- 40 (R)-4-{1-[2-(2-methoxyethoxy)ethyl]pyrrolidin-3-ylamino}-3-nitrobenzenesulfonamide

- [1000]** To a solution of (R)-3-nitro-4-(pyrrolidin-3-ylamino)benzenesulfonamide (440 mg) in N,N-dimethylformamide (10 mL) was added sodium carbonate (132 mg) and 1-bromo-2-(2-methoxyethoxy)ethane (0.155 mL). The reaction mixture was heated at 60°C for 18 hours and after an aqueous workup, the crude product was purified on silica gel with a 2.5 - 10% methanol in methylene chloride gradient to provide the title compound.

EXAMPLE 254D

- 50 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-[[4-((3R)-1-[2-(2-methoxyethoxy)ethyl]pyrrolidin-3-yl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yl-oxy)benzamide

- [1001]** The title compound was prepared by substituting EXAMPLE 254C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500 MHz, pyridine-d₅) δ 12.96 (m, 1H), 9.25 (m, 1H), 8.57 (d, 1H), 8.43 (d, 1H), 8.34 (dd, 1H), 8.11 (d, 1H), 7.66 (t, 1H), 7.64 (d, 1H), 7.44 (d, 2H), 7.07 (d, 2H), 6.82 (d, 1H), 6.76 (dd, 1H), 6.55 (m, 1H), 6.47 (m, 1H), 5.26 (br s, 1H), 4.02 (m, 1H), 3.63 (m, 4H), 3.53 (m, 2H), 3.28 (s, 3H), 3.07 (m, 4H), 2.89 - 2.81 (m, 2H), 2.78 (s, 2H), 2.75 - 2.66 (m, 3H), 2.37 (m, 1H), 2.26 (m, 2H), 2.24 - 2.18 (m, 1H), 2.15 (m, 4H), 1.97 (s, 2H), 1.65 (m, 1H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 255

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(N,N-dimethylglycyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 255A

(R)-4-(1-(2-(dimethylamino)acetyl)pyrrolidin-3-ylamino)-3-nitrobenzenesulfonamide

[1002] The title compound was prepared by substituting 2-(dimethylamino)acetyl chloride, hydrochloric acid for 1-bromo-2-(2-methoxyethoxy)ethane in EXAMPLE 254C except the reaction was stirred at ambient temperature for 18 hours.

EXAMPLE 255B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(N,N-dimethylglycyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1003] The title compound was prepared by substituting EXAMPLE 255A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500 MHz, pyridine - d₅) δ 13.01 (d, 1H), 9.26 (m, 1H), 8.46 - 8.33 (m, 3H), 8.14 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 7.01 - 6.89 (m, 1H), 6.76 (dd, 1H), 6.55 (m, 1H), 6.48 (m, 1H), 5.32 (br s, 1H), 4.27 - 4.14 (m, 1H), 4.05-3.95 (m, 1H), 3.82 - 3.62 (m, 3H), 3.27 - 3.15 (m, 2H), 3.07 (m, 4H), 2.77 (s, 2H), 2.34 (2, 3H), 2.32 (s, 3H), 2.30 - 2.20 (m, 3H), 2.15 (m, 4H), 1.97 (s, 2H), 1.87 - 1.81 (m, 1H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 256

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[1-(oxetan-3-yl)azetidin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 256A

tert-butyl 3-(2-nitro-4-sulfamoylphenylamino)azetidine-1-carboxylate

[1004] The title compound was prepared by substituting tert-butyl 3-aminoazetidine-1-carboxylate for 4-methylpiperazin-1-amine dihydrochloride in EXAMPLE 6A.

EXAMPLE 256B

4-(azetidin-3-ylamino)-3-nitrobenzenesulfonamide

[1005] The title compound was prepared by substituting EXAMPLE 256A for (S)-1-tert-butyl 2-methyl 4,4-difluoropyrrolidine-1,2-dicarboxylate in EXAMPLE 168A.

EXAMPLE 256C

3-nitro-4-(1-(oxetan-3-yl)azetidin-3-ylamino)benzenesulfonamide

[1006] The title compound was prepared by substituting EXAMPLE 256B for tert-butyl piperazine-1-carboxylate and 3-oxetanone for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 256D

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-nitro-4-{[1-(oxetan-3-yl)azetidin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1007] The title compound was prepared by substituting EXAMPLE 256C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, pyridine-d₅) δ 13.04 (s, 1H), 9.27 (d, 1H), 8.62 (d, 1H), 8.42 (d, 1H), 8.35 (dd, 1H), 8.09 (d, 1H), 7.67 (m, 1H), 7.63 (d, 1H), 7.44 (m, 2H), 7.07 (m, 2H), 6.76 (dd, 1H), 6.67 (d, 1H), 6.55 (d, 1H), 6.48 (dd, 1H), 4.66 (t, 2H),

4.58 (m, 2H), 4.23 (m, 1H), 3.71 (m, 3H), 3.12 (dd, 2H), 3.07 (m, 4H), 2.77 (s, 2H), 2.26 (t, 2H), 2.14 (t, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 257

N-[[5-chloro-6-[[1-(cyanomethyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 257A

tert-butyl 4-((3-chloro-5-sulfamoylpyridin-2-yloxy)methyl)-4-fluoropiperidine-1-carboxylate

[1008] The title compound was prepared by substituting EXAMPLE 126A for (tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 257B

5-chloro-6-((4-fluoropiperidin-4-yl)methoxy)pyridine-3-sulfonamide ditrifluoroacetic acid

[1009] The title compound was prepared by substituting EXAMPLE 257A for EXAMPLE 39A in EXAMPLE 39B.

EXAMPLE 257C

5-chloro-6-((1-(cyanomethyl)-4-fluoropiperidin-4-yl)methoxy)pyridine-3-sulfonamide

[1010] EXAMPLE 257B (0.166 g) in acetonitrile (3 mL) was treated with 2-chloroacetonitrile (0.027 g) and sodium carbonate (0.064 g), heated at 60°C overnight, cooled to room temperature and chromatographed on silica gel with 0 to 3% methanol in CH₂Cl₂ as the eluent. The obtained solid was slurried in water, filtered, rinsed with water and diethyl ether, and dried in a vacuum oven at 80°C.

EXAMPLE 257D

N-[[5-chloro-6-[[1-(cyanomethyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1011] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 257C for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (400 MHz, pyridine-d₅) δ 13.05 (s, 1H), 9.12 (d, 1H), 8.72 (d, 1H), 8.41 (d, 1H), 8.10 (d, 1H), 7.66 (m, 2H), 7.44 (m, 2H), 7.07 (m, 2H), 6.75 (dd, 1H), 6.50 (m, 2H), 4.49 (d, 2H), 3.72 (s, 2H), 3.06 (m, 4H), 2.77 (s, 2H), 2.73 (m, 4H), 2.26 (t, 2H), 2.13 (m, 4H), 2.07 (m, 2H), 1.90 (m, 4H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 258

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 258A

(S)-tert-butyl 2-(tosyloxymethyl)morpholine-4-carboxylate

[1012] To a solution of (S)-tert-butyl 2-(hydroxymethyl)morpholine-4-carboxylate (1 g) in dichloromethane (50 mL) was added triethylamine (1.604 mL) and 4-methylbenzene-1-sulfonyl chloride (1.097 g). The mixture was stirred at ambient temperature under nitrogen for 72 hours. The reaction was diluted with methylene chloride (50 mL) and brine (100 mL). The brine layer was extracted with methylene chloride (75 mL). The combined organics were dried over sodium sulfate, filtered and concentrated. The crude material was purified on a silica gel column eluting with a 15-65 % ethyl acetate in hexane gradient to provide the title compound.

EXAMPLE 258B

(S)-tert-butyl 2-(azidomethyl)morpholine-4-carboxylate

- 5 **[1013]** A solution of EXAMPLE 258A (1.66 g) and sodium azide (0.581 g) in anhydrous N,N-dimethylformamide (10 mL) was stirred at 90 °C for 4 hours. The mixture was cooled and concentrated to dryness. The residue was taken up in 5% aqueous sodium carbonate solution and extracted with methylene chloride. The organic solution was dried (MgSO₄) filtered and concentrated to give a solid.

10 EXAMPLE 258C

(R)-tert-butyl 2-(aminomethyl)morpholine-4-carboxylate

- 15 **[1014]** This compound was obtained by hydrogenation of EXAMPLE 258B under 60 psi of hydrogen over 10% palladium on carbon in methanol for 24 hours, followed by filtration and evaporation of the solvent.

EXAMPLE 258D

(R)-tert-butyl 2-((2-nitro-4-sulfamoylphenylamino)methyl)morpholine-4-carboxylate

- 20 **[1015]** The title compound was prepared by substituting EXAMPLE 258C for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 258E

(S)-4-(morpholin-2-ylmethylamino)-3-nitrobenzenesulfonamide

- 25 **[1016]** The title compound was prepared by substituting EXAMPLE 258D for EXAMPLE 113A in EXAMPLE 134A.

30 EXAMPLE 258F

(R)-4-((4-(2-(dimethylamino)acetyl)morpholin-2-yl)methylamino)-3-nitrobenzenesulfonamide

- 35 **[1017]** The title compound was prepared by substituting EXAMPLE 258E for EXAMPLE 134A and 2-(dimethylamino)acetyl chloride hydrochloride for methyl iodide in EXAMPLE 134B.

EXAMPLE 258G

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- 40 **[1018]** The title compound was prepared by substituting EXAMPLE 258F for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.27 (d, 1H), 8.87 (bs, 1H), 8.43 (d, 1H), 8.34 (dd, 1H), 8.10 (dd, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.94 (dd, 1H), 6.75 (d, 1H), 6.54 (s, 1H), 6.48 (s, 1H), 4.55 (dd, 1H), 4.20 (dd, 1H), 3.95-3.76 (m, 2H), 3.60-3.40 (m, 3H), 3.32 (dd, 1H), 3.25-3.12 (m, 2H), 3.07 (m, 4H), 2.80 (m, 1H), 2.77 (s, 2H), 2.26 (s, 6H), 2.23 (s, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 259

50 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([[(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 259A

55 (R)-tert-butyl 2-(tosyloxymethyl)morpholine-4-carboxylate

- [1019]** The title compound was prepared by substituting (R)-tert-butyl 2-(hydroxymethyl)morpholine-4-carboxylate for (S)-tert-butyl 2-(hydroxymethyl)morpholine-4-carboxylate in EXAMPLE 258A.

EXAMPLE 259B

(R)-tert-butyl 2-(azidomethyl)morpholine-4-carboxylate

5 [1020] The title compound was prepared by substituting EXAMPLE 259A for EXAMPLE 258A in EXAMPLE 258B.

EXAMPLE 259C

(S)-tert-butyl 2-(aminomethyl)morpholine-4-carboxylate

10

[1021] The title compound was prepared by substituting EXAMPLE 259B for EXAMPLE 258B in EXAMPLE 258C.

EXAMPLE 259D

15

(S)-tert-butyl 2-((2-nitro-4-sulfamoylphenylamino)methyl)morpholine-4-carboxylate

[1022] The title compound was prepared by substituting EXAMPLE 259C for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

20

EXAMPLE 259E

(R)-4-(morpholin-2-ylmethylamino)-3-nitrobenzenesulfonamide

25

[1023] The title compound was prepared by substituting EXAMPLE 259D for EXAMPLE 113A in EXAMPLE 134A.

EXAMPLE 259F

(S)-4-((4-(2-(dimethylamino)acetyl)morpholin-2-yl)methylamino)-3-nitrobenzenesulfonamide

30

[1024] The title compound was prepared by substituting EXAMPLE 259E for EXAMPLE 134A and 2-(dimethylamino)acetyl chloride hydrochloride for methyl iodide in EXAMPLE 134B.

EXAMPLE 259G

35

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(([(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl)amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

40

[1025] The title compound was prepared by substituting EXAMPLE 259F for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.27 (d, 1H), 8.87 (bs, 1H), 8.43 (d, 1H), 8.34 (dd, 1H), 8.10 (dd, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.94 (dd, 1H), 6.75 (d, 1H), 6.54 (s, 1H), 6.48 (s, 1H), 4.55 (dd, 1H), 4.20 (dd, 1H), 3.95-3.76 (m, 2H), 3.60-3.40 (m, 3H), 3.32 (dd, 1H), 3.25-3.12 (m, 2H), 3.07 (m, 4H), 2.80 (m, 1H), 2.77 (s, 2H), 2.26 (s, 6H), 2.23 (s, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 260

45

N-([5-chloro-6-([1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy)pyridin-3-yl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 260A

50

5-chloro-6-((1-(2-(dimethylamino)acetyl)piperidin-4-yl)methoxy)pyridine-3-sulfonamide

55

[1026] EXAMPLE 253B (0.061 g), 2-(dimethylamino)acetyl chloride, hydrochloric acid (0.061 g), and sodium carbonate (0.032 g) were combined in a 4-mL vial with N,N-dimethylformamide (2 mL). The mixture was stirred at ambient temperature for 3 days. Additional 2-(dimethylamino)acetyl chloride, hydrochloric acid (0.037 g), sodium carbonate (0.032 g) and N,N-dimethylformamide (1 mL) were added and stirring was continued for 24 hours. The reaction mixture was diluted with ethyl acetate, washed with water and brine, dried (MgSO₄), filtered, concentrated and chromatographed on silica gel with 0 to 20% methanol in CH₂Cl₂ as the eluent.

EXAMPLE 260B

N-[(5-chloro-6-[[1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1027] The title compound was prepared by substituting EXAMPLE 260A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400 MHz, pyridine-d₅) δ 12.91 (s, 1H), 9.16 (d, 1H), 8.75 (d, 1H), 8.51 (d, 1H), 8.33 (d, 1H), 7.70 (d, 1H), 7.62 (d, 1H), 7.45 (m, 2H), 7.09 (m, 2H), 6.77 (dd, 1H), 6.60 (d, 1H), 6.45 (d, 1H), 4.81 (d, 1H), 4.15 (m, 3H), 3.24 (m, 2H), 3.04 (m, 4H), 2.89 (m, 1H), 2.79 (s, 2H), 2.53 (m, 1H), 2.29 (m, 6H), 2.26 (m, 2H), 2.18 (m, 4H), 1.98 (m, 2H), 1.91 (m, 1H), 1.71 (m, 2H), 1.39 (t, 2H), 1.25 (m, 2H), 0.94 (s, 6H).

EXAMPLE 261

N-[(5-chloro-6-[[[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 261 A

(R)-tert-Butyl 3-(3-chloro-5-sulfamoylpyridin-2-yloxy)pyrrolidine-1-carboxylate

[1028] The title compound was prepared by substituting EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide and (R)-tert-butyl 3-hydroxypyrrolidine-1-carboxylate for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 261B

(R)-5-Chloro-6-(pyrrolidin-3-yloxy)pyridine-3-sulfonamide

[1029] The title compound was prepared by substituting EXAMPLE 261A for tert-butyl (4-(1,3-difluoropropan-2-yl)morpholin-2-yl)methylcarbamate in EXAMPLE 252B.

EXAMPLE 261C

(R)-5-chloro-6-(1-(2,2-difluoroethyl)pyrrolidin-3-yloxy)pyridine-3-sulfonamide

[1030] A mixture of EXAMPLE 261B (353 mg), 1, 1-difluoro-2-iodoethane (268 mg), sodium carbonate (283 mg) in N,N-dimethylformamide (10mL) was heated at 80°C overnight. The reaction mixture was cooled to room temperature and diluted with ethyl acetate. The organic phase was washed with water, brine, dried over magnesium sulfate, filtered, and concentrated. The residue was loaded onto silica gel column and eluted using a gradient of 0.5 to 3 % methanol in dichloromethane to provide the title compound.

EXAMPLE 261D

N-[(5-chloro-6-[[[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1031] The title compound was prepared by substituting EXAMPLE 261C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.55 (s, 1 H), 8.04 (s, 1 H), 7.95 (d, 1 H), 7.58 (d, 1 H), 7.44 (t, 1 H), 7.35 (m, 3 H), 7.05 (d, 2 H), 6.64 (dd, 1 H), 6.33 (m, 1 H), 6.24 (d, 1 H), 6.25-5.97 (m, 1 H), 5.39 (m, 1 H), 2.98 (m, 6 H), 2.86 (m, 6 H), 2.55 (m, 2 H), 2.24 (m, 7 H), 1.96 (s, 2 H), 1.83 (m, 1 H), 1.39 (t, 2 H), 0.93 (s, 6 H)

EXAMPLE 262

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(cyanomethyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 262A

(R)-4-(1-(cyanomethyl)pyrrolidin-3-ylamino)-3-nitrobenzenesulfonamide

[1032] The title compound was prepared by substituting 2-bromoacetonitrile for 1-bromo-2-(2-methoxyethoxy)ethane in EXAMPLE 254C.

EXAMPLE 262B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(3R)-1-(cyanomethyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1033] The title compound was prepared by substituting EXAMPLE 262A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (400 MHz, pyridine-*d*₅) δ 13.03 (s, 1H), 9.27 (d, 1H), 8.53 (d, 1H), 8.43 (d, 1H), 8.35 (dd, 1H), 8.10 (d, 1H), 7.67 - 7.64 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.81 (d, 1H), 6.76 (dd, 1H), 6.54 (m, 1H), 6.48 (m, 1H), 5.15 (br s, 1H), 4.10 (m, 1H), 3.89 (s, 2H), 3.07 (m, 4H), 2.93 - 2.86 (m, 2H), 2.80 - 2.77 (m, 3H), 2.61 - 2.53 (m, 1H), 2.31 - 2.21 (m, 3H), 2.14 (m, 4H), 1.97 (s, 2H), 1.75 - 1.68 (m, 1H), 1.39 (t, 2H), 0.94 (m, 6H).

EXAMPLE 263

4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-methoxypiperidin-1-yl}-N-[(5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 263A

tert-butyl 1-oxa-6-azaspiro[2.5]octane-6-carboxylate

[1034] Sodium hydride (6.63 g, 60% in mineral oil) was added to trimethylsulfoxonium iodide (36.5 g) in dimethyl sulfoxide (150 mL) and tetrahydrofuran (150 mL), was and stirred for 30 minutes. tert-Butyl 4-oxopiperidine-1-carboxylate (25.4 g) was added and the reaction was stirred for 3 hours. The reaction was poured into water (800 mL) and extracted three times with ether. The combined extracts were washed three times with water, and brine, dried over Na₂SO₄, filtered, and concentrated to yield the crude product which was used without further purification.

EXAMPLE 263B

tert-butyl 4-(2-(benzyloxy)benzyl)-4-hydroxypiperidine-1-carboxylate

[1035] (2-(Benzyloxy)phenyl)magnesium bromide (33.8 mL, 1M) was added to a solution of EXAMPLE 263A (6.0 g) and CuI (1.07 g) in tetrahydrofuran (220 mL) at 0°C over 10 minutes. The reaction was quenched with pH 7 buffer (20 mL), extracted twice with ether, and the combined extracts were washed with brine, dried over Na₂SO₄, filtered, and concentrated. The crude product was chromatographed on silica gel using 2-20% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 263C

tert-butyl 4-hydroxy-4-(2-hydroxybenzyl)piperidine-1-carboxylate

[1036] EXAMPLE 263B (11.5 g) and methanol (120 mL) were added to Raney Nickel (1.150 g) in a 250 mL SS pressure bottle and stirred for 1 hour at 30 psi under hydrogen. The mixture was filtered through a nylon membrane and the solution was concentrated to yield the title compound.

EXAMPLE 263D

tert-butyl 4-hydroxy-4-(2-(trifluoromethylsulfonyloxy)benzyl)piperidine-1-carboxylate

- 5 **[1037]** A mixture of EXAMPLE 263C (4.6 g), N-phenylbis(trifluoromethanesulfonimide) (5.88 g), and Hunig's base (2.88 mL) in dichloromethane (100 mL) was stirred for 24 hours. The mixture was concentrated and chromatographed on silica gel using 5-50% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 263E

10

tert-butyl 4-((4'-chlorobiphenyl-2-yl)methyl)-4-hydroxypiperidine-1-carboxylate

- 15 **[1038]** A mixture of EXAMPLE 263D (4.3 g), 4-chlorophenylboronic acid (1.84 g), K_3PO_4 (2.91 g), and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (0.36 g) in 2-methyltetrahydrofuran (50 mL) was stirred at 70°C for 24 hours. The reaction was cooled and quenched with water (50 mL), extracted twice with ether, and the combined extracts were washed with brine, dried over Na_2SO_4 , filtered, and concentrated. The crude product was chromatographed on silica gel using 5-30% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 263F

20

tert-butyl 4-((4'-chlorobiphenyl-2-yl)methyl)-4-methoxypiperidine-1-carboxylate

- 25 **[1039]** Sodium hydride (0.36 g, 60% in mineral oil) was added to EXAMPLE 263E (4.3 g), in tetrahydrofuran (40 mL) and the reaction was stirred for 10 minutes. Hexamethylphosphoramide (5 mL) and CH_3I (2.34 mL) were added and the reaction was stirred at 50°C for 18 hours. The reaction was cooled and quenched with water (50 mL), extracted twice with ether, and the combined extracts were washed with brine, dried over Na_2SO_4 , filtered, and concentrated. The crude product was chromatographed on silica gel using 5-25% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 263G

30

4-((4'-chlorobiphenyl-2-yl)methyl)-4-methoxypiperidine

- [1040]** The title compound was prepared by substituting EXAMPLE 263F for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 263H

35

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((4'-chlorobiphenyl-2-yl)methyl)-4-methoxypiperidin-1-yl)benzoate

- 40 **[1041]** A solution of EXAMPLE 263G (1.4 g), EXAMPLE 3H (1.06 g) and Hunig's base (0.75 mL) in dimethylsulfoxide (20 mL) was stirred at 120°C for 18 hours. The reaction was cooled and quenched with water (200 mL), extracted three times with ether, and the combined extracts were washed three times with water, and brine, dried over Na_2SO_4 , filtered, and concentrated. The crude product was chromatographed on silica gel using 5-50% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 263I

45

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((4'-chlorobiphenyl-2-yl)methyl)-4-methoxypiperidin-1-yl)benzoic acid

- 50 **[1042]** The title compound was prepared by substituting EXAMPLE 263H for EXAMPLE 3I in EXAMPLE 3J.

EXAMPLE 263J

55

4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-methoxypiperidin-1-yl}-N-[(5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [1043]** The title compound was prepared by substituting EXAMPLE 263I for EXAMPLE 1E and EXAMPLE 96A for EXAMPLE 1F in EXAMPLE 1G. 1H NMR (300MHz, dimethylsulfoxide- d_6) δ 11.71 (s, 1H), 11.58 (br s, 1H), 8.58 (d, 1H), 8.28 (d, 1H), 8.05 (d, 1H), 7.56 (d, 1H), 7.52 (m, 1H), 7.46 (d, 1H), 7.44 (d, 2H), 7.28 (m, 5H), 7.11 (dd, 1H), 6.62 (dd,

1H), 6.41 (dd, 1H), 6.11 (d, 1H), 4.54 (d, 2H), 3.75 (m, 2H), 3.59 (m, 2H), 3.20 (m, 2H), 2.97 (s, 3H), 2.81 (m, 2H), 2.74 (m, 2H), 1.89 (m, 2H), 1.83 (m, 2H), 1.36 (m, 2H), 1.09 (m, 2H).

EXAMPLE 264

4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-methoxypiperidin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1044] The title compound was prepared by substituting EXAMPLE 263I for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.71 (s, 1H), 11.40 (br s, 1H), 8.62 (t, 1H), 8.58 (d, 1H), 8.03 (d, 1H), 7.81 (dd, 1H), 7.54 (m, 2H), 7.44 (m, 3H), 7.28 (m, 5H), 7.13 (dd, 1H), 6.62 (dd, 1H), 6.41 (dd, 1H), 6.11 (d, 1H), 3.85 (dd, 2H), 3.31 (m, 4H), 3.20 (m, 2H), 2.97 (s, 3H), 2.81 (m, 2H), 2.73 (m, 2H), 1.89 (m, 1H), 1.62 (m, 2H), 1.38 (m, 2H), 1.25 (m, 2H), 1.09 (m, 2H).

EXAMPLE 265

4-{4-[[9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 265A

benzyl 4-(piperidin-1-ylmethylene)piperidine-1-carboxylate

[1045] To a solution of benzyl 4-formylpiperidine-1-carboxylate (12.5g) in toluene (120 mL) was added piperidine (6.46g). The mixture was stirred at reflux under a Dean-Stark trap overnight. The mixture was then concentrated under vacuum and the residue was used directly in the next step.

EXAMPLE 265B

benzyl 9-oxo-3-azaspiro[5.5]undec-7-ene-3-carboxylate

[1046] To a solution of EXAMPLE 265A (15.88g) in ethanol (300 mL) was added but-3-enone (3.89 g). The mixture was stirred at reflux overnight. Then acetic acid (30 mL) was added to the mixture which was stirred at reflux again overnight. The mixture was then concentrated under vacuum and the residue was diluted with ethyl acetate (400 mL) and washed with water and brine and dried over Na₂SO₄. After filtration and evaporation of the solvent, column purification gave the title compound.

EXAMPLE 265C

benzyl 9-hydroxy-3-azaspiro[5.5]undecane-3-carboxylate

[1047] EXAMPLE 265B (21 g) and tetrahydrofuran (160 mL) were added to 5% Pt-C wet (3.15 g) in a 250 mL pressure bottle and stirred for 1 hour at 30 psi and room temperature. The mixture was filtered through a nylon membrane and the filtrate was concentrated under vacuum to provide the title compound.

EXAMPLE 265D

benzyl 9-oxo-3-azaspiro[5.5]undecane-3-carboxylate

[1048] To a solution of EXAMPLE 265C (8.0 g) in dichloromethane (200 mL) was added Dess-Martin Periodinane (11.2 g). The mixture was stirred overnight. The mixture was diluted with ethyl acetate (400 mL) and washed with 2N aqueous NaOH, water, and brine. After drying over Na₂SO₄ and filtration, concentration of the solvent gave the crude product which was used directly in the next reaction without further purification.

EXAMPLE 265E

benzyl 9-chloro-8-formyl-3-azaspiro[5.5]undec-8-ene-3-carboxylate

- 5 **[1049]** Phosphorus oxychloride (2.33 mL) was added dropwise to a cooled (0°C) solution of EXAMPLE 265D (7.5 g) in N,N-dimethylformamide (10 mL) and dichloromethane (30 mL). The mixture was then stirred overnight before it was diluted with ethyl acetate (300 mL) and washed with aqueous sodium acetate, water (3x), and brine and dried over Na₂SO₄. After filtration and concentration, the crude product was used directly in the next reaction without further purification.

10

EXAMPLE 265F

benzyl 9-(4-chlorophenyl)-8-formyl-3-azaspiro[5.5]undec-8-ene-3-carboxylate

- 15 **[1050]** To a mixture of 4-chlorophenylboronic acid (5.94 g), EXAMPLE 265E (11.01 g), palladium(II) acetate (142 mg), K₂CO₃ (13.2 g) and tetrabutylammonium bromide (10.2 g) was added water (120 mL). The mixture was stirred at 50°C overnight. The mixture was diluted with ethyl acetate (400 mL) and washed with water (3x) and brine and dried over Na₂SO₄. After filtration and concentration, the residue was loaded on a column and eluted with 5 to 20% ethyl acetate in hexane to provide the title compound.

20

EXAMPLE 265G

benzyl 8-((4-(3-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(methoxycarbonyl)phenyl)piperazin-1-yl)methyl)-9-(4-chlorophenyl)-3-azaspiro[5.5]undec-8-ene-3-carboxylate

25

- [1051]** To a solution of EXAMPLE 15F (1.37 g) and EXAMPLE 265F (1.65 g) in dichloromethane (20 mL) was added sodium triacetoxymethylborohydride (1.24 g). The mixture was stirred overnight. The mixture was diluted with ethyl acetate (200 mL) and washed with 2N aqueous NaOH, water and brine. After drying over Na₂SO₄, the mixture was filtered and the solvent was evaporated under vacuum to provide the title compound.

30

EXAMPLE 265H

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((9-(4-chlorophenyl)-3-azaspiro[5.5]undec-8-en-8-yl)methyl)piperazin-1-yl)benzoate

35

- [1052]** EXAMPLE 265G (2 g) and tetrahydrofuran (10 mL) were added to 20% Pd(OH)₂-C, wet (0.400 g) in a 50 mL pressure bottle and stirred for 16 hours at 30 psi and room temperature. The mixture was filtered through a nylon membrane and evaporation of the solvent gave the title compound.

40

EXAMPLE 265I

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl)methyl)piperazin-1-yl)benzoate

45

- [1053]** To a solution of EXAMPLE 265H (320 mg) in dichloromethane (5 mL) was added 1, 3-difluoroacetone (139 mg) and sodium triacetoxymethylborohydride (157 mg). The mixture was stirred overnight. The mixture was diluted with ethyl acetate (200 mL) and washed with 2N aqueous NaOH, water and brine. After drying over Na₂SO₄, the mixture was filtered and the solvent was evaporated under vacuum to provide the title compound.

50

EXAMPLE 265J

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl)methyl)piperazin-1-yl)benzoic acid

55

- [1054]** To a solution of EXAMPLE 265I (320 mg) in tetrahydrofuran (4 mL) and methanol (2 mL) was added LiOH H₂O (120 mg) and the solution was stirred overnight. The reaction was cooled, carefully neutralized with 1N aqueous HCl and extracted with dichloromethane (3x 50 mL). The combined organic layers were washed with brine (25 mL), dried over sodium sulfate, filtered and concentrated under vacuum to provide the title compound.

EXAMPLE 265K

4-(4-([9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1055] The title compound was prepared by substituting EXAMPLE 265J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.61 (s, 1 H), 8.49 (d, 2 H), 7.72 (m, 1 H), 7.49 (m, 2 H), 7.32 (d, 2 H), 7.07 (m, 3 H), 6.65 (dd, 1 H), 6.35 (d, 1 H), 6.20 (m, 1 H), 4.66 (m, 2 H), 4.50 (m, 2 H), 3.84 (m, 2 H), 3.04 (m, 5 H), 2.70 (m, 6 H), 2.23 (m, 6 H), 2.00 (m, 4 H), 1.35 (m, 12 H).

EXAMPLE 266

4-(4-([9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 266A

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl)methyl)piperazin-1-yl)benzoate

[1056] To a solution of EXAMPLE 265H (320 mg) in dichloromethane (5 mL) was added acetone (143 mg) and sodium triacetoxymethylborohydride (157 mg). The mixture was stirred overnight. The mixture was diluted with ethyl acetate (200 mL) and washed with 2N aqueous NaOH, water and brine. After drying over Na₂SO₄, the mixture was filtered and the solvent was evaporated under vacuum to provide the title compound.

EXAMPLE 266B

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl)methyl)piperazin-1-yl)benzoic acid

[1057] The title compound was prepared by substituting EXAMPLE 266A for EXAMPLE 265I in EXAMPLE 265J.

EXAMPLE 266C

4-(4-([9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1058] The title compound was prepared by substituting EXAMPLE 266B for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.54 (s, 1 H), 8.38 (m, 2 H), 7.93 (d, 1 H), 7.60 (m, 3 H), 7.39 (m, 4 H), 7.09 (d, 2 H), 6.85 (d, 1 H), 6.63 (dd, 1 H), 6.27 (dd, 2 H), 3.84 (m, 3 H), 3.08 (m, 8 H), 2.71 (s, 3 H), 2.15 (m, 8 H), 1.71 (m, 9 H), 1.24 (m, 11 H)

EXAMPLE 267

4-(4-([9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-([5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1059] The title compound was prepared by substituting EXAMPLE 265J for EXAMPLE 1E and EXAMPLE 40B for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.56 (s, 1 H), 8.38 (s, 1 H), 8.06 (m, 1 H), 7.57 (d, 1 H), 7.38 (m, 5 H), 7.07 (m, 3 H), 6.64 (dd, 1 H), 6.33 (d, 1 H), 6.23 (m, 1 H), 4.68 (d, 2 H), 4.52 (d, 2 H), 4.21 (d, 2 H), 3.86 (dd, 2 H), 3.08 (m, 8 H), 2.71 (m, 6 H), 2.10 (m, 12 H), 1.42 (m, 7 H).

EXAMPLE 268

4-(4-([9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-([5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1060] The title compound was prepared by substituting EXAMPLE 266B for EXAMPLE 1E and EXAMPLE 40B for

EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.49 (s, 1 H), 8.28 (d, 1 H), 7.94 (dd, 2 H), 7.60 (d, 1 H), 7.35 (m, 4 H), 7.08 (m, 2 H), 6.61 (dd, 1 H), 6.28 (dd, 2 H), 4.18 (d, 2 H), 3.85 (m, 2 H), 3.05 (m, 7 H), 2.71 (s, 3 H), 2.25 (m, 6 H), 2.02 (m, 2 H), 1.63 (m, 8 H), 1.30 (m, 9 H).

5 EXAMPLE 269

N-({5-chloro-6-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 EXAMPLE 269A

5-chloro-6-((4-fluoro-1-methylpiperidin-4-yl)methoxy)pyridine-3-sulfonamide

15 **[1061]** EXAMPLE 257B (0.131 g) in N,N-dimethylformamide (3.0 mL) was treated with iodomethane (0.043 g) and sodium carbonate (0.079 g) and stirred at ambient temperature for 3 days. The N,N-dimethylformamide was removed on high vacuum and the concentrate was chromatographed on amine functionalized silica gel with 0 to 2% methanol in CH₂Cl₂ as the eluent.

20 EXAMPLE 269B

N-({5-chloro-6-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

25 **[1062]** The title compound was prepared by substituting EXAMPLE 269A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400 MHz, pyridine-d₅) δ 13.01 (s, 1H), 9.11 (d, 1H), 8.71 (d, 1H), 8.44 (d, 1H), 8.16 (d, 1H), 7.66 (m, 2H), 7.44 (m, 2H), 7.07 (m, 2H), 6.76 (dd, 1H), 6.54 (d, 1H), 6.49 (dd, 1H), 4.49 (d, 2H), 3.06 (m, 4H), 2.77 (s, 2H), 2.68 (m, 2H), 2.38 (m, 2H), 2.26 (m, 5H), 2.14 (t, 4H), 1.97 (m, 6H), 1.39 (t, 2H), 0.94 (s, 6H).

30 EXAMPLE 270

N-[(5-chloro-6-[[1-(N,N-dimethylglycyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

35 EXAMPLE 270A

5-chloro-6-((1-(2-(dimethylamino)acetyl)-4-fluoropiperidin-4-yl)methoxy)pyridine-3-sulfonamide

40 **[1063]** EXAMPLE 257B (0.131 g), 2-(dimethylamino)acetyl chloride, hydrochloric acid (0.139 g), and sodium carbonate (0.048 g) were combined in a 5-mL vial with N,N-dimethylformamide (3.0 mL) and stirred overnight at ambient temperature. Additional sodium carbonate (0.048 g) was added followed by 2-(dimethylamino)acetyl chloride, hydrochloric acid (0.139 g) and stirring was continued over a second night. The reaction mixture was concentrated under high vacuum, slurried in CH₂Cl₂, filtered, concentrated and chromatographed on amine functionalized silica gel with 0 to 4% methanol in CH₂Cl₂ as the eluent.

45 EXAMPLE 270B

N-[(5-chloro-6-[[1-(N,N-dimethylglycyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50 **[1064]** The title compound was prepared by substituting EXAMPLE 270A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, pyridine-d₅) δ 13.04 (s, 1H), 9.12 (d, 1H), 8.73 (d, 1H), 8.42 (d, 1H), 8.11 (d, 1H), 7.66 (m, 2H), 7.44 (m, 2H), 7.07 (m, 2H), 6.76 (dd, 1H), 6.52 (d, 1H), 6.49 (dd, 1H), 4.66 (d, 1H), 4.52 (dd, 2H), 4.07 (d, 1H), 3.46 (m, 1H), 3.40 (m, 1H), 3.30 (m, 1H), 3.11 (m, 1H), 3.06 (m, 4H), 2.77 (s, 2H), 2.35 (s, 6H), 2.26 (t, 2H), 2.14 (m, 4H), 2.05 (m, 2H), 1.97 (s, 2H), 1.81 (m, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

55

EXAMPLE 271

4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-fluoropiperidin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 271 A

tert-butyl 4-((4'-chlorobiphenyl-2-yl)methyl)-4-fluoropiperidine-1-carboxylate

[1065] A solution of EXAMPLE 263E (2.0 g) and diethylaminosulfur trifluoride (1.39 mL) in dichloromethane (40 mL) was stirred for 24 hours. The reaction was quenched with water (30 mL), extracted twice with ether, and the combined extracts were washed with water, and brine, dried over Na₂SO₄, filtered, and concentrated. The crude product was chromatographed on silica gel using 5% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 271B

4-((4'-chlorobiphenyl-2-yl)methyl)-4-fluoropiperidine

[1066] The title compound was prepared by substituting EXAMPLE 271A for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 271C

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((4'-chlorobiphenyl-2-yl)methyl)-4-fluoropiperidin-1-yl)benzoate

[1067] The title compound was prepared by substituting EXAMPLE 271B for EXAMPLE 263G in EXAMPLE 263H.

EXAMPLE 271D

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((4'-chlorobiphenyl-2-yl)methyl)-4-fluoropiperidin-1-yl)benzoic acid

[1068] The title compound was prepared by substituting EXAMPLE 271C for EXAMPLE 3I in EXAMPLE 3J.

EXAMPLE 271E

4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-fluoropiperidin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1069] The title compound was prepared by substituting EXAMPLE 271D for EXAMPLE 1E EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1H), 11.46 (br s, 1H), 8.62 (t, 1H), 8.56 (d, 1H), 8.03 (d, 1H), 7.81 (dd, 1H), 7.52 (m, 3H), 7.44 (d, 2H), 7.28 (m, 5H), 7.14 (m, 1H), 6.68 (dd, 1H), 6.40 (dd, 1H), 6.19 (d, 1H), 3.84 (dd, 2H), 3.31 (m, 9H), 2.95 (d, 2H), 2.81 (m, 2H), 1.91 (m, 1H), 1.62 (m, 2H), 1.45 (m, 2H), 1.29 (m, 2H).

EXAMPLE 272

4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-fluoropiperidin-1-yl}-N-({5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1070] The title compound was prepared by substituting EXAMPLE 271D for EXAMPLE 1E and EXAMPLE 96A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1H), 11.64 (br s, 1H), 8.58 (m, 1H), 8.25 (m, 1H), 8.03 (d, 1H), 7.70 (dd, 1H), 7.50 (m, 4H), 7.43 (m, 3H), 7.28 (m, 4H), 7.15 (m, 1H), 6.68 (dd, 1H), 6.40 (dd, 1H), 6.19 (d, 1H), 4.54 (d, 2H), 4.04 (m, 1H), 3.75 (m, 2H), 3.58 (m, 2H), 2.95 (d, 2H), 2.80 (m, 2H), 1.88 (m, 2H), 1.82 (m, 2H), 1.48 (m, 2H), 1.28 (m, 2H), 0.85 (m, 2H).

EXAMPLE 273

4-(4-{[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-(trifluoromethyl)phenyl}sulfonyl)benzamide

[1071] The title compound was prepared by substituting EXAMPLE 266B for EXAMPLE 1E and EXAMPLE 42A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.57 (s, 1 H), 7.97 (d, 1 H), 7.77 (s, 1 H), 7.55 (m, 2 H), 7.45 (m, 1 H), 7.36 (m, 3 H), 7.08 (d, 2 H), 6.62 (dd, 2 H), 6.35 (dd, 1 H), 6.21 (d, 1 H), 3.82 (m, 3 H), 3.06 (m, 9 H), 2.72 (m, 3 H), 2.25 (m, 8 H), 2.09 (m, 2 H), 1.56 (m, 9 H), 1.20 (m, 10 H).

EXAMPLE 274

N-[(5-chloro-6-[[[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 274A

(R)-5-chloro-6-(1-(3-fluoro-2-(fluoromethyl)propyl)pyrrolidin-3-yloxy)pyridine-3-sulfonamide

[1072] The title compound was prepared by substituting 1,3-difluoropropan-2-one for 4'-chlorobiphenyl-2-carboxaldehyde and EXAMPLE 261B for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A.

EXAMPLE 274B

N-[(5-chloro-6-[[[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1073] The title compound was prepared by substituting EXAMPLE 274A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.52 (s, 1 H), 8.32 (d, 1 H), 8.01 (d, 1 H), 7.93 (d, 1 H), 7.59 (d, 1 H), 7.42 (m, 1 H), 7.33 (m, 3 H), 7.05 (d, 2 H), 6.63 (dd, 1 H), 6.31 (dd, 1 H), 6.25 (d, 1 H), 5.38 (m, 1 H), 4.65 (t, 2 H), 4.53 (t, 2 H), 3.02 (s, 4 H), 2.94 (m, 5 H), 2.75 (s, 2 H), 2.66 (m, 1 H), 2.23 (m, 7 H), 1.96 (s, 2 H), 1.82 (m, 1 H), 1.39 (t, 2 H), 0.93 (s, 6 H).

EXAMPLE 275

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[2-(tetrahydrofuran-3-yloxy)ethoxy]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 275A

3-(2-(benzyloxy)ethoxy)tetrahydrofuran

[1074] Tetrahydrofuran-3-ol (0.881 g) in tetrahydrofuran (15 mL) was treated with 60% sodium hydride (0.8 g). After 10 minutes, ((2-bromoethoxy)methyl)benzene (3.23 g) was added. The solution was stirred for 16 hours. The reaction mixture was partitioned between water and ethyl acetate. The aqueous layer was separated, and was extracted with additional ethyl acetate twice. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography on silica gel eluting with 1:1 ethyl acetate: hexane to provide the title compound.

EXAMPLE 275B

2-(tetrahydrofuran-3-yloxy)ethanol

[1075] EXAMPLE 275A (0.85 g) and 5% palladium on carbon (0.1 g) in ethanol (10 mL) was treated with a balloon of hydrogen. The reaction was stirred overnight. The solid was filtered off, and the filtrate was concentrated to give the title compound.

EXAMPLE 275C

3-nitro-4-(2-(tetrahydrofuran-3-yloxy)ethoxy)benzenesulfonamide

- 5 **[1076]** The title compound was prepared by substituting EXAMPLE 275B for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 275D

- 10 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[2-(tetrahydrofuran-3-yloxy)ethoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [1077]** The title compound was prepared by substituting EXAMPLE 275C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1H), 8.32 (s, 1H), 8.00-8.02 (m, 2H), 7.49-7.52 (m, 2H), 7.39-7.41 (m, 1H), 7.38 (d, 2H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.21 (d, 1H), 4.33-4.35 (m, 2H), 4.18-4.21 (m, 1H), 3.62-3.67 (m, 4H), 3.09 (s, 4H), 2.83 (s, 2H), 2.26 (s, 2H), 2.15 (s, 2H), 1.96 (s, 2H), 1.85-1.94 (m, 2H), 1.39 (t, 2H), 0.92 (s, 6H).

EXAMPLE 276

- 20 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(trans-4-cyanocyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 276A

- 25 trans-4-(aminomethyl)cyclohexanecarbonitrile

[1078] To a solution of tert-butyl (trans-4-(cyanomethyl)cyclohexyl)methylcarbamate (500 mg) in dichloromethane (10 mL) was slowly added trifluoroacetic acid (2 mL) at 0°C. The reaction mixture was warmed to room temperature, stirred for 1 hour and concentrated to provide the title compound.

EXAMPLE 276B

4-((trans-4-cyanocyclohexyl)methylamino)-3-nitrobenzenesulfonamide

- 35 **[1079]** A mixture of 4-fluoro-3-nitrobenzenesulfonamide (347 mg) and EXAMPLE 276A (300 mg) in tetrahydrofuran (20 mL) was treated with triethylamine (1.4 mL) overnight and concentrated. The residue was triturated with ethyl acetate to provide the title compound.

EXAMPLE 276C

- 40 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(trans-4-cyanocyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [1080]** The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 276B in place of EXAMPLE 11B. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1 H), 11.36 (s, 1 H), 8.60 (t, 1 H), 8.56 (d, 1 H), 8.04 (d, 1 H), 7.79 (dd, 1 H), 7.47 - 7.54 (m, 3 H), 7.34 (d, 2 H), 7.01 - 7.09 (m, 3 H), 6.68 (dd, 1 H), 6.39 (dd, 1 H), 6.19 (d, 1 H), 3.25 (t, 2 H), 3.07 (s, 4 H), 2.76 (s, 2 H), 2.57 - 2.68 (m, 1 H), 2.17 (d, 6 H), 1.92 - 2.06 (m, 4 H), 1.78 (d, 2 H), 1.66 (s, 1 H), 1.35 - 1.53 (m, 4 H), 0.96 - 1.10 (m, 2 H), 0.92 (s, 6 H).

50

55

EXAMPLE 277

N-[(5-chloro-6-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 277A

methyl 4,4-dimethyl-2-(trifluoromethylsulfonyloxy)cyclohex-1-enecarboxylate

[1081] To a suspension of hexane washed NaH (17 g) in dichloromethane (700 mL) was added 5,5-dimethyl-2-methoxycarbonylcyclohexanone (38.5 g) dropwise at 0°C. After stirring for 30 minutes, the mixture was cooled to -78°C and trifluoroacetic anhydride (40 mL) was added. The reaction mixture was warmed to room temperature and stirred for 24 hours. The organic layer was washed with brine, dried (Na₂SO₄), filtered, and concentrated to give the product.

EXAMPLE 277B

methyl 2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enecarboxylate

[1082] EXAMPLE 277A (62.15g), 4-chlorophenylboronic acid (32.24 g), CsF (64 g) and tetrakis(triphenylphosphine)palladium(0) (2g) in 2:1 dimethoxyethane /methanol (600 mL) were heated to 70°C for 24 hours. The mixture was concentrated. Ether (4x 200 mL) was added and the mixture was filtered. The combined ether solution was concentrated to give the product.

EXAMPLE 277C

(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methanol

[1083] To a mixture of LiBH₄ (13g), EXAMPLE 277B (53.8 g) and ether (400 mL), was added methanol (25 mL) slowly by syringe. The mixture was stirred at room temperature for 24 hours. The reaction was quenched with 1N HCl with ice-cooling. The mixture was diluted with water and extracted with ether (3x 100 mL). The extracts were dried (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 0-30% ethyl acetate/hexanes.

EXAMPLE 277D

tert-butyl 4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazine-1-carboxylate

[1084] Mesyl Chloride (7.5 mL) was added via syringe to EXAMPLE 277C (29.3 g) and triethylamine (30 mL) in CH₂Cl₂ (500 mL) at 0°C, and the mixture was stirred for 1 minute. N-t-butoxycarbonylpiperazine (25 g) was added and the mixture was stirred at room temperature for 24 hours. The suspension was washed with brine, dried, (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 10-20% ethyl acetate/hexanes.

EXAMPLE 277E

1-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazine

[1085] EXAMPLE 277D (1 g) was stirred in dichloromethane (10 mL), trifluoroacetic acid (10 mL), and triethylsilane (1 mL) for 1 hour. The mixture was concentrated, taken up in a mixture of dichloromethane (100 mL) and saturated aqueous Na₂CO₃ solution (20 mL) and stirred for 10 minutes. The layers were separated, and the organic layer was dried over Na₂SO₄, filtered, and concentrated to give the product.

EXAMPLE 277F

5-bromo-1-(triisopropylsilyl)-1H-pyrrolo[2,3-b]pyridine

[1086] To a mixture of 5-bromo-1H-pyrrolo[2,3-b]pyridine (15.4 g) in tetrahydrofuran (250 mL) was added 1M lithium hexamethyldisilazide in tetrahydrofuran (86 mL), and after 10 minutes, TIPS-Cl (triisopropylchlorosilane) (18.2 mL) was added. The mixture was stirred at room temperature for 24 hours. The reaction was diluted with ether, and the resulting solution was washed twice with water. The extracts were dried (Na₂SO₄), filtered, and concentrated. The crude product

was chromatographed on silica gel with 10% ethyl acetate/hexanes.

EXAMPLE 277G

5 1-(triisopropylsilyl)-1H-pyrrolo[2,3-b]pyridin-5-ol

[1087] To a mixture of EXAMPLE 277F (24.3 g) in tetrahydrofuran (500 mL) at -78°C was added 2.5M BuLi (30.3 mL). After 2 minutes, trimethylborate (11.5 mL) was added, and the mixture was allowed to warm to room temperature over 1 hour. The reaction was poured into water, extracted three times with ethyl acetate, and the combined extracts were washed with brine and concentrated. The crude product was taken up in tetrahydrofuran (200 mL) at 0°C, and 1M NaOH (69 mL) was added, followed by 30% H₂O₂ (8.43 mL), and the solution was stirred for 1 hour. Na₂S₂O₃ (10 g) was added, and the pH was adjusted to 4-5 with concentrated HCl and solid NaH₂PO₄. The solution was extracted twice with ethyl acetate, and the combined extracts were washed with brine, dried (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 5-25% ethyl acetate/hexanes.

15

EXAMPLE 277H

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-fluorobenzoate

[1088] A mixture of EXAMPLE 277G (8.5 g), methyl 2,4-difluorobenzoate (7.05 g), and K₃PO₄ (9.32 g) in diglyme (40 mL) at 115°C was stirred for 24 hours. The reaction was cooled, diluted with ether (600 mL), and washed twice with water, and brine, and concentrated. The crude product was chromatographed on silica gel with 2-50% ethyl acetate/hexanes.

25

EXAMPLE 277I

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoate

[1089] A mixture of EXAMPLE 277H (1.55 g), EXAMPLE 277E (2.42 g), and HK₂PO₄ (1.42 g) in dimethylsulfoxide (20 mL) at 135°C was stirred for 24 hours. The reaction was cooled, diluted with ether (400 mL), and washed three times with 1M NaOH, and brine, and concentrated. The crude product was chromatographed on silica gel with 10-50% ethyl acetate/hexanes.

35

EXAMPLE 277J

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoic acid

40

[1090] EXAMPLE 277I (200 mg) in dioxane (10 mL) and 1M NaOH (6 mL) at 50°C was stirred for 24 hours. The reaction was cooled, added to NaH₂PO₄ solution, and extracted three times with ethyl acetate. The combined extracts were washed with brine, and concentrated to give the pure product.

EXAMPLE 277K

45

5,6-dichloropyridine-3-sulfonamide

[1091] To a solution of 5,6-dichloropyridine-3-sulfonyl chloride (32.16 g) in isopropyl alcohol (300 mL) at 0 °C was added a 30% aqueous solution of NH₄OH (50.8 mL). After stirring overnight, the solvent was reduced to 1/3 of the original volume. It was then partitioned between water and ethyl acetate. The aqueous layer was extracted with additional ethyl acetate. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was chromatographed on silica gel. The material was then slurried in 1:9 ethyl acetate/hexanes, filtered and dried under vacuum to give the title compound.

55

EXAMPLE 277L

tert-butyl 4-fluoro-4-(hydroxymethyl)piperidine-1-carboxylate

- 5 **[1092]** 1-Tert-butyl 4-ethyl 4-fluoropiperidine-1,4-dicarboxylate (1.0 g) in tetrahydrofuran (5 mL) was treated with 1.0 N LiAlH₄ in tetrahydrofuran (2.54 mL) at 0 °C. The reaction mixture was stirred at room temperature for 2 hours. Water (0.6 mL) was added to the reaction mixture drop-wise, followed by 2 N aqueous NaOH (0.2 mL). The reaction was stirred for another 1 hour. The solid was removed by filtration via a pack of diatomaceous earth and washed with ethyl acetate. The filtrate was washed with brine, dried over MgSO₄, filtered, and concentrated to give the product.

10

EXAMPLE 277M

tert-butyl 4-((3-chloro-5-sulfamoylpyridin-2-yl)oxy)methyl)-4-fluoropiperidine-1-carboxylate

- 15 **[1093]** To a solution of EXAMPLE 277L (1 g) in tetrahydrofuran (15 mL) was added NaH (60% dispersion in mineral oil, 685 mg), and the solution was stirred for 10 minutes. EXAMPLE 277K (1 g) was added and the reaction stirred for 24 hours. The mixture was poured into water, neutralized with 10% HCl, and extracted with ethyl acetate three times. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified with flash column chromatography on silica gel eluting with 30% ethyl acetate in hexanes.

20

EXAMPLE 277N

5-chloro-6-((4-fluoropiperidin-4-yl)methoxy)pyridine-3-sulfonamide ditrifluoroacetic acid EXAMPLE 277M (13 mL) was treated with trifluoroacetic acid (2.363 mL), stirred at ambient temperature for 2 hours, concentrated and dried to give the title compound.

25

EXAMPLE 277O

5-chloro-6-((4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methoxy)pyridine-3-sulfonamide

30

- [1094]** EXAMPLE 277N (0.088 g) and oxetan-3-one (0.014 g) were combined in dichloromethane (2.0 mL) and dimethylformamide (1.0 mL) and stirred at ambient temperature for 45 minutes. Sodium triacetoxymethylborohydride (0.064 g) was added in portions. Stirring was continued overnight at ambient temperature. Additional oxetan-3-one (0.014 g) was added and stirring was continued for 30 minutes at ambient temperature before more sodium triacetoxymethylborohydride (0.064 g) was added. The reaction mixture was stirred for 72 hours at ambient temperature, concentrated, chromatographed on silica gel with 0 to 5% methanol in dichloromethane as the eluent, and dried in a vacuum oven at 80 °C to give the title compound.

35

EXAMPLE 277P

40

N-[(5-chloro-6-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yl)benzamide

45

- [1095]** EXAMPLE 277J (0.063 g), EXAMPLE 277O (0.042 g), 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (0.032 g), and 4-dimethylaminopyridine (0.027 g) were combined in a 4-mL vial with dichloromethane (1.0 mL) and stirred overnight at ambient temperature. The reaction mixture was chromatographed directly without aqueous workup on silica gel with 0-4% methanol in dichloromethane as the eluent. Fractions containing the desired product were concentrated, slurried in acetonitrile, concentrated and dried overnight in a vacuum oven at 80 °C to give the title compound. ¹H NMR (500 MHz, pyridine-d₅) δ 13.05 (s, 1H), 9.13 (d, 1H), 8.72 (d, 1H), 8.41 (d, 1H), 8.10 (d, 1H), 7.67 (m, 1H), 7.66 (d, 1H), 7.44 (m, 2H), 7.07 (m, 2H), 6.76 (dd, 1H), 6.51 (m, 2H), 4.63 (m, 4H), 4.53 (d, 2H), 3.39 (m, 1H), 3.07 (m, 4H), 2.77 (s, 2H), 2.51 (m, 2H), 2.25 (m, 2H), 2.18 (m, 2H), 2.13 (m, 4H), 2.06 (t, 2H), 1.97 (s, 2H), 1.89 (m, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

50

55

EXAMPLE 278

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({5-cyano-6-[2-(tetrahydro-2H-pyran-4-yl)ethoxy]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 278A

5-bromo-6-(2-(tetrahydro-2H-pyran-4-yl)ethoxy)pyridine-3-sulfonamide

[1096] The title compound was prepared by substituting 2-(tetrahydro-2H-pyran-4-yl)ethanol for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 36B.

EXAMPLE 278B

5-cyano-6-(2-(tetrahydro-2H-pyran-4-yl)ethoxy)pyridine-3-sulfonamide

[1097] The title compound was prepared by substituting EXAMPLE 278A for EXAMPLE 36B in EXAMPLE 36C.

EXAMPLE 278C

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({5-cyano-6-[2-(tetrahydro-2H-pyran-4-yl)ethoxy]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1098] The title compound was prepared by substituting EXAMPLE 278B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.60 (s, 1H), 8.70 (s, 1H), 8.48 (s, 1H), 7.96 (d, 1H), 7.56 (d, 1H), 7.45-7.47 (m, 1H), 7.40 (s, 1H), 7.36 (d, 2H), 7.06 (d, 2H), 6.67 (dd, 1H), 6.34 (dd, 1H), 6.25 (d, 1H), 4.47 (d, 2H), 3.80-3.84 (m, 2H), 3.24-3.28 (m, 2H), 3.12 (s, 2H), 2.16 (s, 2H), 1.97 (s, 2H), 1.61-1.71 (m, 4H), 1.40 (t, 2H), 1.21-1.25 (m, 2H), 0.93 (s, 6H).

EXAMPLE 279

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[[4-(3-furylmethoxy)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 279A

4-(furan-3-ylmethoxy)-3-nitrobenzenesulfonamide

[1099] The title compound was prepared by substituting furan-3-ylmethanol for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 279B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[[4-(3-furylmethoxy)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1100] The title compound was prepared by substituting EXAMPLE 279A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 8.34 (s, 1H), 8.03-8.06 (m, 2H), 7.83 (s, 1H), 7.69 (t, 1H), 7.51-7.53 (m, 4H), 7.34-7.36 (m, 2H), 7.04-7.06 (m, 2H), 6.68 (dd, 1H), 6.57 (s, 1H), 6.40 (dd, 1H), 6.20 (d, 1H), 5.23 (s, 2H), 3.10 (s, 4H), 2.83 (s, 2H), 2.15-2.32 (m, 6H), 1.39 (t, 2H), 0.92 (s, 6H).

EXAMPLE 280

N-[[5-chloro-6-[[[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 280A

(R)-tert-butyl 3-((3-chloro-5-sulfamoylpyridin-2-yloxy)methyl)pyrrolidine-1-carboxylate

[1101] The title compound was prepared by substituting EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide and (R)-tert-butyl 3-(hydroxymethyl)pyrrolidine-1-carboxylate for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 280B

(R)-5-chloro-6-(pyrrolidin-3-ylmethoxy)pyridine-3-sulfonamide

[1102] The title compound was prepared by substituting EXAMPLE 280A for tert-butyl (4-(1,3-difluoropropan-2-yl)morpholin-2-yl)methylcarbamate in EXAMPLE 252B.

EXAMPLE 280C

(R)-5-chloro-6-((1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl)methoxy)pyridine-3-sulfonamide

[1103] The title compound was prepared by substituting 1,3-difluoropropan-2-one for 4'-chlorobiphenyl-2-carboxaldehyde and EXAMPLE 280B for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A.

EXAMPLE 280D

N-[[5-chloro-6-[[[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1104] The title compound was prepared by substituting EXAMPLE 280C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.57 (s, 1 H), 8.38 (d, 1 H), 8.07 (d, 1 H), 7.96 (d, 1 H), 7.57 (d, 1 H), 7.44 (m, 1 H), 7.35 (m, 3 H), 7.05 (d, 2 H), 6.64 (dd, 1 H), 6.33 (dd, 1 H), 6.23 (d, 1 H), 4.65 (d, 2 H), 4.53 (dd, 2 H), 2.92 (m, 8 H), 2.75 (m, 4 H), 2.58 (m, 2 H), 2.20 (m, 6 H), 1.96 (m, 4 H), 1.53 (m, 1 H), 1.39 (t, 2 H), 0.89 (s, 6 H).

EXAMPLE 281

N-[[5-chloro-6-[[[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 281 A

(R)-5-chloro-6-((1-(2,2-difluoroethyl)pyrrolidin-3-yl)methoxy)pyridine-3-sulfonamide

[1105] The title compound was prepared by substituting EXAMPLE 280B for EXAMPLE 261B in EXAMPLE 261C.

EXAMPLE 281B

N-[[5-chloro-6-[[[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1106] The title compound was prepared by substituting EXAMPLE 281A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.59 (s, 1 H), 8.41 (d, 1 H), 8.10 (d, 1 H), 7.98 (d, 1 H), 7.56 (d, 1 H), 7.46 (m, 1 H), 7.41 (d, 1 H), 7.34 (d, 2 H), 7.04 (d, 2 H), 6.65 (dd, 1 H), 6.35 (dd, 1 H), 6.23 (m, 1 H), 6.03 (m, 1 H), 3.06 (s, 4 H), 2.84 (m, 6 H), 2.63 (m, 4 H), 2.20 (m, 6 H), 1.94 (m, 3 H), 1.53 (m, 1 H), 1.39 (t, 2 H), 0.91 (s, 6 H).

EXAMPLE 282

N-[(5-chloro-6-[[1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 282A

5-chloro-6-((1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl)methoxy)pyridine-3-sulfonamide

[1107] EXAMPLE 257B (0.088 g) and 1,3-difluoropropan-2-one (0.028 g) were combined in dichloromethane (2 mL) and N,N-dimethylformamide (0.500 mL) and stirred at ambient temperature for 45 minutes. Sodium triacetoxyborohydride (0.064 g) was added in portions and then the reaction mixture was stirred overnight at ambient temperature. Additional 1,3-difluoropropan-2-one (0.028 g) was added, followed 30 minutes later by the addition of more sodium triacetoxyborohydride (0.064 g). The reaction mixture was stirred at ambient temperature for 72 hours. Additional 1,3-difluoropropan-2-one (0.028 g) was again added, followed 30 minutes later by the addition of more sodium triacetoxyborohydride (0.064 g). The reaction mixture was stirred overnight at ambient temperature. Additional 1,3-difluoropropan-2-one (0.028 g) was again added, followed 30 minutes later by the addition of more sodium triacetoxyborohydride (0.064 g). The reaction mixture was stirred overnight at ambient temperature. The reaction mixture was concentrated under high vacuum to remove N,N-dimethylformamide and then chromatographed on silica gel with 0 to 4% methanol in CH₂Cl₂ as the eluent.

EXAMPLE 282B

N-[(5-chloro-6-[[1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1108] The title compound was prepared by substituting EXAMPLE 282A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400 MHz, pyridine-d₅) δ 13.05 (s, 1H), 9.12 (t, 1H), 8.72 (d, 1H), 8.41 (d, 1H), 8.10 (d, 1H), 7.66 (m, 2H), 7.44 (m, 2H), 7.07 (m, 2H), 6.75 (dd, 1H), 6.50 (m, 2H), 4.77 (dd, 1H), 4.65 (dd, 1H), 4.52 (dd, 2H), 3.06 (m, 4H), 2.93 (t, 1H), 2.80 (m, 5H), 2.52 (m, 1H), 2.26 (t, 2H), 2.13 (m, 4H), 2.04 (m, 2H), 1.97 (s, 2H), 1.85 (m, 2H), 1.39 (t, 2H), 1.28 (m, 2H), 0.93 (s, 6H).

EXAMPLE 283

N-({3-chloro-4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]phenyl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 283A

3-chloro-4-((4-fluoro-1-methylpiperidin-4-yl)methoxy)benzenesulfonamide

[1109] To a solution of (4-fluoro-1-methylpiperidin-4-yl)methanol (0.265 g) in tetrahydrofuran (2 mL) was added sodium hydride (0.288 g). After 15 minutes, 3-chloro-4-fluorobenzenesulfonamide (0.377 g) was added as a solution in tetrahydrofuran (1 mL). The reaction was stirred for 2 hours, quenched with water (5 mL), adjusted to pH~7 with 1N aqueous HCl, and extracted with dichloromethane (2 x 25 mL). The organic layer was washed with brine (25 mL), dried over magnesium sulfate, filtered and concentrated. Silica gel chromatography (Reveleris 40 g) eluting with a gradient of 0.1% to 10% methanol containing 2N NH₃/dichloromethane over 30 minutes gave the title compound.

EXAMPLE 283B

N-({3-chloro-4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]phenyl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1110] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 283A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide- d₆) δ 11.60 (s, 1H), 10.68 - 9.84 (m, 1H), 7.99 (d, 1H), 7.79 (d, 1H), 7.63 (t, 1H), 7.54 (d, 1H), 7.50 - 7.38 (m, 2H), 7.34 (d, 2H), 7.04 (d, 3H), 6.64 (dd, 1H), 6.36 (dd, 1H), 6.22 (s, 1H), 4.23 (d, 2H), 3.03 (s, 6H), 2.71 (m, 4H), 2.07 (m, 12H), 1.38 (s, 3H), 1.24 (s, 2H), 0.92 (s, 6H).

EXAMPLE 284

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-cyano-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 284A

3-cyano-4-((tetrahydro-2H-pyran-4-yl)methoxy)benzenesulfonamide

[1111] To a solution of (tetrahydro-2H-pyran-4-yl)methanol (0.258 g) in tetrahydrofuran (5 mL) was added sodium hydride (0.355 g) and the reaction stirred at room temperature for 15 minutes. EXAMPLE 52A (0.400 g) was added and the reaction stirred for an additional 1 hour. The reaction was poured into ethyl acetate (50 mL) and 1N aqueous HCl (35 mL). The organic layer was washed with brine (35 mL) dried over magnesium sulfate, filtered, and concentrated. Silica gel chromatography (Reveleris 40 g) eluting with a gradient of 10% to 100% ethyl acetate/hexanes over 30 minutes gave the title compound.

EXAMPLE 284B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-cyano-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1112] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 284A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide- d₆) δ 11.69 (s, 1H), 11.60 - 11.16 (m, 1H), 8.15 (s, 1H), 8.08 - 8.01 (m, 2H), 7.58 - 7.46 (m, 3H), 7.35 (d, J = 8.4, 2H), 7.29 (d, 1H), 7.04 (d, 2H), 6.68 (d, 1H), 6.40 (dd, 1H), 6.20 (s, 1H), 4.05 (d, 2H), 3.89 (d, 2H), 3.37 (d, 4H), 3.09 (s, 4H), 2.81 (s, 2H), 2.21 (d, 7H), 1.96 (s, 2H), 1.67 (d, 2H), 1.39 (s, 2H), 0.92 (s, 6H).

EXAMPLE 285

N-([5-chloro-6-([1-(2,2-difluoroethyl)-4-fluoropiperidin-4-yl]methoxy)pyridin-3-yl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 285A

5-chloro-6-((1-(2,2-difluoroethyl)-4-fluoropiperidin-4-yl)methoxy)pyridine-3-sulfonamide

[1113] EXAMPLE 257B (0.263 g), 1,1-difluoro-2-iodoethane (0.23 g), and sodium carbonate (0.254 g) were combined in a 20-mL vial with N,N-dimethylformamide (6 mL) and stirred at 70 °C overnight. The reaction mixture was concentrated under high vacuum and then chromatographed on silica gel with 0 to 5% methanol in CH₂Cl₂ as the eluent.

EXAMPLE 285B

N-([5-chloro-6-([1-(2,2-difluoroethyl)-4-fluoropiperidin-4-yl]methoxy)pyridin-3-yl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1114] The title compound was prepared by substituting EXAMPLE 285A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400 MHz, pyridine-d₅) δ 13.05 (s, 1H), 9.12 (d, 1H), 8.72 (d, 1H), 8.41 (d, 1H), 8.10 (d, 1H), 7.66 (m, 2H), 7.43 (m, 2H), 7.06 (m, 2H), 6.75 (dd, 1H), 6.50 (m, 2H), 6.18 (tt, 2H), 4.51 (d, 2H), 3.07 (m, 4H), 2.80 (m, 6H), 2.60 (td, 2H), 2.25 (t, 2H), 2.13 (m, 4H), 2.03 (t, 2H), 1.97 (s, 2H), 1.93 (m, 1H), 1.85 (m, 1H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 286

N-({3-chloro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 286A

3-chloro-4-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)benzenesulfonamide

[1115] The title compound was prepared by substituting EXAMPLE 37C for (4-fluoro-1-methylpiperidin-4-yl)methanol in EXAMPLE 283A.

EXAMPLE 286B

N-({3-chloro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1116] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 286A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-*d*₆) δ 11.71 (s, 1H), 11.56 - 11.16 (m, 1H), 8.06 (d, 1H), 7.90 (d, 1H), 7.81 (d, 1H), 7.64 - 7.45 (m, 3H), 7.34 (d, 2H), 7.26 (d, 1H), 7.04 (d, 2H), 6.68 (d, 1H), 6.42 (dd, 1H), 6.18 (s, 1H), 4.28 (d, 2H), 3.78 (d, 2H), 3.61 (dd, 2H), 3.07 (s, 4H), 2.76 (s, 2H), 2.17 (d, 6H), 1.87 (dd, 6H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 287

N-({5-chloro-6-[(4,4-difluorocyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 287A

(4,4-difluorocyclohexyl)methanol

[1117] Ethyl 4,4-difluorocyclohexanecarboxylate (1.0 g, 5.20 mmol) in diethyl ether (2 mL) was added dropwise to lithium aluminium hydride (0.24 g) in diethyl ether (15 mL), and heated under reflux for 4 hours. The reaction was then cooled to 0°C, and water was added (0.24 mL), followed by 5N aqueous NaOH (0.24 mL) and water (0.72 mL). Then Na₂SO₄ and more diethyl ether (40 mL) were added, and the mixture was stirred for 30 minutes, then filtered through celite. After concentration, the residue was diluted with CH₂Cl₂ and Na₂SO₄ was added, and the mixture was filtered and concentrated to provide the title compound.

EXAMPLE 287B

5-chloro-6-((4,4-difluorocyclohexyl)methoxy)pyridine-3-sulfonamide

[1118] The title compound was prepared by substituting EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide and EXAMPLE 287A for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 287C

N-({5-chloro-6-[(4,4-difluorocyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1119] The title compound was prepared by substituting EXAMPLE 287B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-*d*₆) δ 11.63 (s, 1H), 8.46 (d, 1H), 8.14 (d, 1H), 8.00 (d, 1H), 7.56 (d, 1H), 7.47 (m, 2H), 7.35 (d, 2H), 7.03 (d, 2H), 6.66 (dd, 1H), 6.37 (m, 1H), 6.21 (d, 1H), 4.25 (d, 2H), 3.07 (br m, 4H), 2.82 (br s, 2H), 2.30 (br m, 4H), 2.16 (br m, 2H), 2.00, 1.95, 1.85 (all m, total 9H), 1.40 (t, 2H), 1.37 (m, 2H), 0.92 (s, 6H).

EXAMPLE 288

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([6-([1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl]methoxy)-5-(trifluoromethyl)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 288A

5-Nitro-3-(trifluoromethyl)pyridin-2-ol

[1120] 3-(Trifluoromethyl)pyridin-2-ol (2.3 g) was added to concentrated sulfuric acid (15 mL) at 0 °C. The mixture was stirred at 0°C for 5 minutes. To this solution was added fuming nitric acid (6 mL) dropwise over 5 minutes. The reaction mixture was stirred at room temperature for 2 hours, and then heated at 50°C for 3 hours. After cooling, the reaction mixture was poured onto ice (200 g), and the mixture was extracted with ethyl acetate three times. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated under reduced pressure to provide the title compound.

EXAMPLE 288B

2-Chloro-5-nitro-3-(trifluoromethyl)pyridine

[1121] A mixture of EXAMPLE 288A (1.69 g), phosphorus pentachloride (2.03 g), and phosphoryl trichloride (0.97 mL) was heated at 90°C for 3 hours. After cooling, the reaction mixture was poured into ice, and extracted with ethyl acetate three times. The extract was washed with brine, dried over MgSO₄, filtered, and concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel eluting with 10% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 288C

6-Chloro-5-(trifluoromethyl)pyridin-3-amine

[1122] A mixture of iron (1.5 g) and ammonium chloride (2.38 g) in water (40 mL) was stirred at room temperature for 5 minutes. To this suspension was added EXAMPLE 288B in methanol (40 mL). The reaction mixture was stirred at room temperature for 1 hour. More iron (1.8 g) was added to the reaction mixture, and it was stirred for another 3 hours. The solid from the reaction mixture was filtered off, and the filtrate was partitioned between water and ethyl acetate. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel eluting with 20% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 288D

6-chloro-5-(trifluoromethyl)pyridine-3-sulfonyl chloride

[1123] Under ice-cooling, thionyl chloride (4 mL) was added dropwise over 20 minutes to water (27 mL). The mixture was stirred overnight for 12 hours to give a SO₂ containing solution. Separately, EXAMPLE 288C (1.14 g) in dioxane (5 mL) was added to concentrated HCl (20 mL) at 0°C. The solution was stirred for 5 minutes. To this suspension/solution was added sodium nitrite (0.44 g) in water (6 mL) dropwise at 0°C. The solution was stirred at 0°C for 3 hours. During this time, any solid formed was crushed with a glass rod to make sure that EXAMPLE 288C was completely reacted. To the SO₂ containing solution was added copper(I) chloride (0.115 g). Then, to this solution was added the diazotized EXAMPLE 288C at 0°C. The solution was stirred for 30 minutes. The reaction mixture was extracted with ethyl acetate. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel eluting with 5% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 288E

6-chloro-5-(trifluoromethyl)pyridine-3-sulfonamide

[1124] EXAMPLE 288D (2.03 g) in dioxane (20 mL) solution was cooled to 0°C. Ammonium hydroxide solution was

added dropwise. The reaction mixture was stirred at 0°C for 2 hours followed by room temperature over night. The solvent was partially removed, and the residue was partitioned between water and ethyl acetate. The organic phase was washed with brine, dried over MgSO₄, filtered, and concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel eluting with 0-3% methanol in dichloromethane to afford the title compound.

EXAMPLE 288F

tert-butyl 4-fluoro-4-((5-sulfamoyl-3-(trifluoromethyl)pyridin-2-yloxy)methyl)piperidine-1-carboxylate

[1125] The title compound was prepared by substituting EXAMPLE 288E for 4-fluoro-3-nitrobenzenesulfonamide and EXAMPLE 322A for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 288G

6-((4-fluoropiperidin-4-yl)methoxy)-5-(trifluoromethyl)pyridine-3-sulfonamide

[1126] The title compound was prepared by substituting EXAMPLE 288F for tert-butyl (4-(1,3-difluoropropan-2-yl)morpholin-2-yl)methylcarbamate in EXAMPLE 252B.

EXAMPLE 288H

6-((1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl)methoxy)-5-(trifluoromethyl)pyridine-3-sulfonamide

[1127] The title compound was prepared by substituting 1,3-difluoropropan-2-one for 4'-chlorobiphenyl-2-carboxaldehyde and EXAMPLE 288G for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A.

EXAMPLE 288I

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([6-([1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl]methoxy)-5-(trifluoromethyl)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1128] The title compound was prepared by substituting EXAMPLE 288H for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.50 (s, 1 H), 8.57 (s, 1 H), 8.27 (d, 1 H), 7.91 (d, 1 H), 7.58 (d, 1 H), 7.40 (m, 1 H), 7.35 (d, 2 H), 7.28 (d, 1 H), 7.05 (d, 2 H), 6.61 (dd, 1 H), 6.29 (dd, 1 H), 6.24 (d, 1 H), 4.67 (d, 2 H), 4.55 (d, 2 H), 4.50 (s, 1 H), 4.44 (s, 1 H), 3.06 (m, 5 H), 2.73 (m, 6 H), 2.19 (d, 6 H), 1.90 (m, 7 H), 1.39 (t, 2 H), 0.93 (s, 6 H).

EXAMPLE 289

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-chloro-6-[2-(tetrahydrofuran-2-yl)ethoxy]pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 289A

5-chloro-6-(2-(tetrahydrofuran-2-yl)ethoxy)pyridine-3-sulfonamide

[1129] The title compound was prepared by substituting 2-(tetrahydro-2H-pyran-4-yl)ethanol for (tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 40A for EXAMPLE 36A in EXAMPLE 36B.

EXAMPLE 289B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([5-chloro-6-[2-(tetrahydrofuran-2-yl)ethoxy]pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1130] The title compound was prepared by substituting EXAMPLE 289A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1H), 8.52 (d, 1H), 8.18 (s, 1H), 8.02 (s, 1H), 7.50-7.55 (m, 3H), 7.35 (d, 2H), 7.05 (d, 2H), 6.68 (dd, 1H), 6.38 (dd, 1H), 6.21 (d, 1H), 4.39-4.51 (m, 4H), 3.87-3.94 (m, 1H), 3.73-3.78 (m, 1H), 3.57-3.62 (m, 1H), 3.11 (s, 4H), 2.89 (s, 2H), 2.33 (s, 4H), 2.15 (s, 2H), 1.77-2.01 (m, 7H), 1.45-1.54 (m, 1H), 1.40 (t, 2H), 0.93 (s, 6H).

EXAMPLE 290

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl})-3-methylpiperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 290A

2-chloro-4,4-dimethylcyclohex-1-enecarbaldehyde

[1131] Into a 250 ml round-bottomed flask was added N,N-dimethylformamide (3.5 mL) in dichloromethane (30 mL). The mixture was cooled to -10°C, and phosphoryl trichloride (4 mL) was added dropwise. The solution was warmed up to room temperature and 3,3-dimethylcyclohexanone (5.5 mL) was added slowly. The mixture was heated to reflux overnight. The reaction mixture was quenched by 0°C solution of sodium acetate (25 g in 50 mL water). The aqueous layer was extracted with ether (3 x 200 mL). The organic layers were combined, dried over Na₂SO₄, filtered, and dried under vacuum.

EXAMPLE 290B

2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enecarbaldehyde

[1132] Into a 1 L round-bottomed flask was added EXAMPLE 290A (6.8 g), 4-chlorophenylboronic acid (6.5 g) and palladium(II) acetate (0.2 g) in water (100 mL) to give a suspension. Potassium carbonate (15 g) and tetrabutylammonium bromide (10 g) were added. After degassing after subjecting to vacuum and nitrogen, the mixture was stirred at 45°C for 4 hours. After filtering through silica gel, diethyl ether (4x 200 mL) was used to extract the product. The combined organic layers were dried over Na₂SO₄ and filtered. The filtrate was concentrated and purified by flash chromatography on silica with 0-10% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 290C

tert-butyl 4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)-3-methylpiperazine-1-carboxylate

[1133] To a solution of *tert*-butyl 3-methylpiperazine-1-carboxylate (0.256 g) and EXAMPLE 290B (0.350 g) in dichloromethane (2 mL) was added sodium triacetoxyborohydride (0.406 g) and the reaction was stirred at room temperature overnight. The reaction was quenched with NaHCO₃ solution (50 mL) and extracted with dichloromethane (50 mL). The organic layer was dried over magnesium sulfate, filtered and concentrated. Silica gel chromatography (Reveleris 40 g) eluting with a gradient of 0.5% to 2.5% methanol/dichloromethane gave the title compound.

EXAMPLE 290D

1-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)-2-methylpiperazine

[1134] A solution of EXAMPLE 290C (0.298 g) and HCl (4.0M in dioxane, 2 mL) were stirred for 1 hour. The reaction was concentrated and partitioned between dichloromethane (100 mL) and NaHCO₃ (100 mL). The organic layer was washed with brine (50 mL), dried over magnesium sulfate, filtered and concentrated to provide the title compound.

EXAMPLE 290E

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)-3-methylpiperazin-1-yl)benzoate

[1135] The title compound was prepared by substituting EXAMPLE 290D for EXAMPLE 3E in EXAMPLE 3I.

EXAMPLE 290F

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)-3-methylpiperazin-1-yl)benzoic acid

[1136] The title compound was prepared by substituting EXAMPLE 290E for EXAMPLE 15G in EXAMPLE 15H.

EXAMPLE 290G

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)-3-methylpiperazin-1-yl)-N-([3-nitro-4-((tetrahydro-2H-pyran-4-ylmethyl)amino)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1137] The title compound was prepared by substituting EXAMPLE 290F for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide- *d*₆) δ 11.66 (s, 1H), 11.54 - 11.30 (m, 1H), 8.62 - 8.53 (m, 2H), 8.03 (d, 1H), 7.78 (d, 1H), 7.48 (d, 3H), 7.34 (d, 2H), 7.06 (t, 3H), 6.68 (d, 1H), 6.38 (dd, 1H), 6.21 (s, 1H), 3.84 (d, 2H), 3.23 (s, 4H), 2.75 (s, 4H), 1.64 (s, 8H), 1.62 (d, 2H), 1.42 - 1.17 (m, 6H), 0.92 (s, 6H), 0.87 (s, 3H).

EXAMPLE 291

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-(cyclopropylamino)propyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 291 A

tert-butyl 2-cyanoethyl(cyclopropyl)carbamate

[1138] To a solution of 3-(cyclopropylamino)propanenitrile (5.0g) in tetrahydrofuran (30 mL) was added di-*tert*-butyl dicarbonate (9.91g) and a catalytic amount of 4-dimethylaminopyridine. The mixture was stirred overnight. The mixture was diluted with ethyl acetate (400 mL) and washed with 5% aqueous HCl, water and brine. After drying over Na₂SO₄, the mixture was filtered, and the solvent was evaporated under vacuum to provide the title compound.

EXAMPLE 291B

tert-butyl 3-aminopropyl(cyclopropyl)carbamate

[1139] EXAMPLE 291A (9.75 g) and 7M NH₃-methanol (25 mL) were added to a Ra-Ni 2800, water slurry (19.50 g, 332 mmol) in a 250 mL pressure bottle and stirred for 2 hours at 30 psi and room temperature. The mixture was filtered through a nylon membrane and evaporation of the solvent gave the title compound.

EXAMPLE 291C

tert-butyl cyclopropyl(3-(2-nitro-4-sulfamoylphenylamino)propyl)carbamate

[1140] To a solution of 4-chloro-3-nitrobenzenesulfonamide (2.5 g), and EXAMPLE 291B (2.26 g) in dioxane (20 mL) was added N,N-diisopropylethylamine (5 mL). The mixture was stirred at reflux overnight. The mixture was diluted with ethyl acetate (400 mL) and washed with water and brine. After drying over Na₂SO₄, the mixture was filtered, and the solvent was evaporated under vacuum to provide the title compound.

EXAMPLE 291D

tert-butyl 3-(4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)sulfamoyl)-2-nitrophenylamino)propyl(cyclopropyl)carbamate

[1141] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 291C for EXAMPLE 1F in EXAMPLE 1G.

EXAMPLE 291E

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-(cyclopropylamino)propyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1142] To a solution of EXAMPLE 291D (2.56 g) in dichloromethane (10 mL) was added trifluoroacetic acid (10 mL). The mixture was stirred for 2 hours. The mixture was concentrated under vacuum and the residue was dissolved in dichloromethane (300 mL) and washed with aqueous NaHCO₃, water, and brine and dried over Na₂SO₄. Filtration and evaporation of the solvent gave the crude product. The title compound was obtained by dissolving 200 mg of the crude

material in dimethylsulfoxide/methanol (1:1, 10 mL) and loaded on Gilson, C18(100A) 250x121.2 mm(10micron), eluting with 30% acetonitrile to 65% acetonitrile over 40 minutes. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.54 (s, 1 H), 8.43 (m, 2 H), 7.94 (d, 1 H), 7.71 (dd, 1 H), 7.57 (d, 1 H), 7.43 (m, 1 H), 7.34 (m, 3 H), 7.05 (d, 2 H), 6.90 (d, 1 H), 6.63 (dd, 1 H), 6.29 (d, 2 H), 3.43 (m, 2 H), 2.96 (m, 6 H), 2.73 (m, 2 H), 2.22 (m, 7 H), 1.87 (m, 4 H), 1.38 (m, 3 H), 0.94 (m, 6 H), 0.62 (m, 4 H).

EXAMPLE 292

N-[[5-chloro-6-(2-methoxyethoxy)pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 292A

5-chloro-6-(2-methoxyethoxy)pyridine-3-sulfonamide

[1143] The title compound was prepared by substituting EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide and 2-methoxyethanol for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 292B

N-[[5-chloro-6-(2-methoxyethoxy)pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1144] The title compound was prepared by substituting EXAMPLE 292A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.63 (s, 1H), 8.48 (d, 1H), 8.17 (d, 1H), 8.01 (d, 1H), 7.56 (d, 1H), 7.49 (m, 2H), 7.35 (d, 2H), 7.04 (d, 2H), 6.66 (dd, 1H), 6.37 (m, 1H), 6.21 (d, 1H), 4.52 (m, 2H), 3.70 (m, 2H), 3.28 (s, 3H), 3.13 (br m, 4H), 2.88 (br s, 2H), 2.34 (br m, 4H), 2.16 (br m, 2H), 1.97 (s, 2H), 1.40 (t, 2H), 0.92 (s, 6H).

EXAMPLE 293

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-fluoro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 293A

[1145] The title compound was prepared by substituting 5-bromo-2,3-difluoropyridine for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 293B

tert-butyl 5-fluoro-6-((tetrahydro-2H-pyran-4-yl)methoxy)pyridin-3-ylcarbamate

[1146] The title compound was prepared by substituting EXAMPLE 293A for EXAMPLE 248A in EXAMPLE 248B.

EXAMPLE 293C

5-fluoro-6-((tetrahydro-2H-pyran-4-yl)methoxy)pyridine-3-sulfonyl chloride

[1147] The title compound was prepared by substituting EXAMPLE 293B for EXAMPLE 248B in EXAMPLE 248C.

EXAMPLE 293D

5-fluoro-6-((tetrahydro-2H-pyran-4-yl)methoxy)pyridine-3-sulfonamide

[1148] The title compound was prepared by substituting EXAMPLE 293C for EXAMPLE 248C in EXAMPLE 248D.

EXAMPLE 293E

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[5-fluoro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1149] The title compound was prepared by substituting EXAMPLE 293D for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400 MHz, pyridine-d₅) δ 13.07 (s, 1H), 9.05 (d, 1H), 8.44 (dd, 1H), 8.41 (d, 1H), 8.09 (d, 1H), 7.67 (t, 1H), 7.65 (d, 1H), 7.44 (m, 2H), 7.07 (m, 2H), 6.76 (dd, 1H), 6.53 (d, 1H), 6.49 (dd, 1H), 4.21 (d, 2H), 3.96 (dd, 2H), 3.31 (td, 2H), 3.07 (m, 4H), 2.77 (s, 2H), 2.26 (t, 2H), 2.14 (m, 4H), 1.97 (m, 3H), 1.58 (dd, 2H), 1.38 (m, 4H), 0.94 (s, 6H).

EXAMPLE 294

N-[(3-chloro-4-[[1-(methoxyacetyl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 294A

tert-butyl 4-((2-chloro-4-sulfamoylphenoxy)methyl)piperidine-1-carboxylate

[1150] The title compound was prepared by substituting *tert-butyl*-4-(hydroxymethyl)piperidine-1-carboxylate for (4-fluoro-1-methylpiperidin-4-yl)methanol in EXAMPLE 283A.

EXAMPLE 294B

tert-butyl 4-((4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)sulfamoyl)-2-chlorophenoxy)methyl)piperidine-1-carboxylate

[1151] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 294A for EXAMPLE 1F in EXAMPLE 1G.

EXAMPLE 294C

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-(3-chloro-4-(piperidin-4-ylmethoxy)phenylsulfonyl)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzamide

[1152] To EXAMPLE 294B (0.286 g) in dichloromethane (3 mL) was added trifluoroacetic acid (1 mL) and the reaction stirred at room temperature. After 3 hours the reaction was concentrated to provide the title compound.

EXAMPLE 294D

N-[(3-chloro-4-[[1-(methoxyacetyl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1153] To EXAMPLE 294C (0.75 g) as a solution in dichloromethane (1 mL) was added N,N-diisopropylethylamine (0.055 mL) followed by 2-methoxyacetyl chloride (6 μl). After stirring for 10 minutes the reaction was loaded onto silica gel (Reveleris 40 g) and eluted using a gradient of 0.5% to 3.5% methanol/dichloromethane over 30 minutes (flow = 40 mL/minutes) to provide the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.71 (s, 1H), 11.55 - 11.24 (m, 1H), 8.06 (d, 1H), 7.88 (d, 1H), 7.78 (d, 1H), 7.57 (s, 1H), 7.51 (s, 1H), 7.48 (d, 1H), 7.34 (d, 2H), 7.21 (d, 1H), 7.04 (d, 2H), 6.67 (d, 1H), 6.42 (dd, 1H), 6.18 (s, 1H), 4.42 - 4.32 (m, 1H), 4.03 (dd, 4H), 3.86 - 3.74 (m, 1H), 3.28 (s, 3H), 3.07 (s, 5H), 2.77 (s, 3H), 2.30 - 1.92 (m, 9H), 1.77 (s, 2H), 1.31 (d, 4H), 0.92 (s, 6H).

EXAMPLE 295

N-[(3-chloro-4-[[1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1154] The title compound was prepared by substituting 2-(dimethylamino)acetyl chloride for 2-methoxyacetyl chloride in EXAMPLE 294D. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.58 (s, 1H), 10.35 - 9.94 (m, 1H), 7.96 (d, 1H), 7.74

(d, 1H), 7.55 (d, 2H), 7.45 (s, 1H), 7.41 - 7.29 (m, 3H), 7.05 (d, 3H), 6.63 (d, 1H), 6.37 - 6.32 (m, 1H), 6.22 (d, 1H), 4.39 (d, 1H), 3.94 (s, 6H), 3.01 (s, 6H), 2.73 (m, 4H), 2.55 (m, 5H), 2.19 (s, 6H), 1.95 (m, 2H), 1.82 (m, 2H), 1.38 (s, 4H), 0.93 (s, 6H).

5 EXAMPLE 296

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohexyl)methyl]piperidin-1-yl)-N-([3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 EXAMPLE 296A

tert-butyl 4-((4,4-dimethyl-2-oxocyclohexyl)methyl)piperidine-1-carboxylate

15 **[1155]** 3,3-Dimethylcyclohexanone (5.60 mL) was added to sodium bis(trimethylsilyl)amide (45.3 mL, 1M in tetrahydrofuran), and the reaction was stirred for 1 hour. tert-Butyl 4-(bromomethyl)piperidine-1-carboxylate (11.1 g) in dimethylsulfoxide (30 mL) was added, and the reaction was stirred at 50°C for 24 hours. The reaction was cooled, poured into water (300 mL), extracted three times with ether, and the combined extracts were washed three times with water, and brine, dried over Na₂SO₄, filtered, and concentrated. The crude product was chromatographed on silica gel using 5-20% ethyl acetate in hexanes to provide the title compound.

20

EXAMPLE 296B

tert-butyl 4-((2-(4-chlorophenyl)-2-hydroxy-4,4-dimethylcyclohexyl)methyl)piperidine-1-carboxylate

25 **[1156]** (4-Chlorophenyl)magnesium bromide (14.1 mL, 1M in ether) was added to EXAMPLE 296A (3.25 g) in tetrahydrofuran (40 mL) at -78°C, and the reaction was stirred for 20 minutes, and then allowed to warm to room temperature overnight. The reaction was quenched with pH 7 buffer (20 mL), extracted with 2x ether, and the combined extracts were washed with brine, dried over Na₂SO₄, filtered, and concentrated. The crude product was chromatographed on silica gel using 1-20% ethyl acetate in hexanes to provide the title compound.

30

EXAMPLE 296C

trans-4-((2-(4-chlorophenyl)-4,4-dimethylcyclohexyl)methyl)piperidine

35 **[1157]** The title compound was prepared by substituting EXAMPLE 296B for EXAMPLE 1A in EXAMPLE 1B.

EXAMPLE 296D

40 Trans-methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohexyl)methyl)piperidin-1-yl)benzoate

[1158] The title compound was prepared by substituting EXAMPLE 296C for EXAMPLE 263G in EXAMPLE 263H.

EXAMPLE 296E

45

Trans-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohexyl)methyl)piperidin-1-yl)benzoic acid

50

[1159] The title compound was prepared by substituting EXAMPLE 296D for EXAMPLE 3I in EXAMPLE 3J.

EXAMPLE 296F

Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohexyl)methyl]piperidin-1-yl)-N-([3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

55

[1160] The title compound was prepared by substituting EXAMPLE 296E for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 11.36 (br s, 1H), 8.60 (t, 1H), 8.55 (d, 1H), 8.03 (d, 1H), 7.78 (dd, 1H), 7.52 (m, 3H), 7.27 (d, 2H), 7.16 (d, 2H), 7.09 (m, 1H), 6.63 (dd, 1H), 6.38 (dd, 1H), 6.11 (d, 1H), 3.83 (dd, 2H), 3.52 (m,

2H), 3.26 (m, 4H), 2.61 (m, 2H), 2.35 (m, 1H), 1.89 (m, 2H), 1.76 (m, 1H), 1.62 (m, 2H), 1.38 (m, 4H), 1.25 (m, 6H), 1.12 (m, 2H), 0.95 (m, 2H), 0.94 (s, 3H), 0.88 (s, 3H).

EXAMPLE 297

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([6-(tetrahydro-2H-pyran-4-ylmethoxy)-5-(trifluoromethyl)pyridin-3-yl]sulfonyl)benzamide

EXAMPLE 297A

6-((tetrahydro-2H-pyran-4-yl)methoxy)-5-(trifluoromethyl)pyridine-3-sulfonamide

[1161] The title compound was prepared by substituting EXAMPLE 288E for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 297B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([6-(tetrahydro-2H-pyran-4-ylmethoxy)-5-(trifluoromethyl)pyridin-3-yl]sulfonyl)benzamide

[1162] The title compound was prepared by substituting EXAMPLE 297A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.49 (s, 1 H), 8.56 (d, 1 H), 8.23 (d, 1 H), 7.90 (d, 1 H), 7.58 (d, 1 H), 7.40 (m, 1 H), 7.34 (m, 2 H), 7.26 (d, 1 H), 7.05 (d, 2 H), 6.61 (dd, 1 H), 6.28 (dd, 1 H), 6.24 (d, 1 H), 4.24 (d, 2 H), 3.86 (dd, 2 H), 3.30 (m, 4 H), 3.00 (s, 4 H), 2.73 (s, 2 H), 2.16 (m, 6 H), 1.97 (m, 2 H), 1.61 (dd, 2 H), 1.33 (m, 4 H), 0.93 (s, 6 H).

EXAMPLE 298

N-([5-chloro-6-((trans-4-hydroxycyclohexyl)methoxy)pyridin-3-yl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 298A

6-((trans-4-(tert-butyldimethylsilyloxy)cyclohexyl)methoxy)-5-chloropyridine-3-sulfonamide

[1163] The title compound was prepared by substituting (trans-4-(tert-butyldimethylsilyloxy)cyclohexyl)methanol for (tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 40A for EXAMPLE 36A in EXAMPLE 36B.

EXAMPLE 298B

N-([5-chloro-6-((trans-4-hydroxycyclohexyl)methoxy)pyridin-3-yl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1164] The title compound was prepared by substituting EXAMPLE 298A for EXAMPLE 11B in EXAMPLE 11D. After the reaction was over, the solvent was removed, and the residue was treated with 1:1 trifluoroacetic acid/dichloromethane for two hours. The solvents were removed, and the residue was purified by reverse phase Gilson Prep HPLC system with a Phenomenex prep column (Luna, 5 μm, C18(2), 250X21.20 mm, 5 Å) eluting with 20-80% acetonitrile in water with 0.1% trifluoroacetic acid to provide the title compound. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1H), 8.47 (s, 1H), 8.15 (s, 1H), 8.01 (d, 1H), 7.54 (d, 1H), 7.48-7.49 (m, 1H), 7.35 (d, 2H), 7.05 (d, 2H), 6.67 (dd, 1H), 6.37 (d, 1H), 6.21 (s, 1H), 4.53 (t, 1H), 4.18 (d, 2H), 3.08 (s, 4H), 2.84 (s, 2H), 2.29 (s, 4H), 2.15 (s, 2H), 1.96 (s, 2H), 1.79-1.83 (m, 5H), 1.39 (t, 2H), 1.08-1.13 (m, 5H), 0.93 (s, 6H).

EXAMPLE 299

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-cyano-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 299A

3-cyano-4-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)benzenesulfonamide

[1165] The title compound was prepared by substituting EXAMPLE 37C for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 284A.

EXAMPLE 299B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-cyano-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1166] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 299A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide- d₆) δ 11.72 (s, 1H), 10.24 - 9.27 (m, 1H), 8.21 (d, 1H), 8.12 (dd, 1H), 8.05 (d, 1H), 7.63 - 7.46 (m, 3H), 7.45 - 7.31 (m, 3H), 7.07 (d, 2H), 6.70 (dd, 1H), 6.42 (s, 1H), 6.23 (s, 1H), 4.38 (d, 2H), 3.91 - 3.73 (m, 2H), 3.68 - 3.51 (m, 2H), 3.22 - 2.96 (m, 10H), 2.31 - 2.12 (m, 2H), 1.99 (s, 6H), 1.43 (t, 2H), 0.93 (s, 6H).

EXAMPLE 300

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({6-[(trans-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 300A

6-((trans-4-methoxycyclohexyl)methoxy)-5-(trifluoromethyl)pyridine-3-sulfonamide

[1167] The title compound was prepared by substituting EXAMPLE 288E for 4-fluoro-3-nitrobenzenesulfonamide and EXAMPLE 121A for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 300B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({6-[(trans-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1168] The title compound was prepared by substituting EXAMPLE 300A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.50 (s, 1 H), 8.56 (d, 1 H), 8.23 (d, 1 H), 7.90 (d, 1 H), 7.58 (d, 1 H), 7.40 (m, 1 H), 7.35 (d, 2 H), 7.27 (d, 1 H), 7.05 (d, 2 H), 6.61 (dd, 1 H), 6.28 (dd, 1 H), 6.24 (d, 1 H), 4.20 (d, 2 H), 3.23 (s, 3 H), 3.03 (m, 5 H), 2.73 (s, 2 H), 2.18 (m, 6 H), 1.98 (m, 5 H), 1.80 (m, 3 H), 1.39 (t, 2 H), 1.09 (m, 4 H), 0.93 (s, 6 H).

EXAMPLE 301

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({6-[(cis-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 301A

6-((cis-4-methoxycyclohexyl)methoxy)-5-(trifluoromethyl)pyridine-3-sulfonamide

[1169] The title compound was prepared by substituting EXAMPLE 288E for 4-fluoro-3-nitrobenzenesulfonamide and EXAMPLE 121A for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 301B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({6-[(cis-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3b]pyridin-5-yloxy)benzamide

[1170] The title compound was prepared by substituting EXAMPLE 301A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.49 (m, 1 H), 8.54 (m, 1 H), 8.23 (d, 1 H), 7.91 (d, 1 H), 7.59 (d, 1 H), 7.40 (m, 1 H), 7.34 (m, 2 H), 7.27 (d, 1 H), 7.04 (d, 2 H), 6.61 (dd, 1 H), 6.29 (dd, 1 H), 6.24 (d, 1 H), 4.20 (d, 2 H), 3.37 (m, 2 H), 3.19 (s, 3 H), 3.00 (s, 4 H), 2.73 (s, 2 H), 2.18 (m, 6 H), 1.96 (s, 2 H), 1.80 (m, 3 H), 1.50 (dd, 2 H), 1.37 (m, 6 H), 0.93 (s, 6 H).

EXAMPLE 302

N-({5-chloro-6-[(4,4-difluoro-1-hydroxycyclohexyl)methoxy]pyridin-3-yl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperidin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 302A

4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperidine

[1171] EXAMPLE 296B (1.0 g) was stirred in dichloromethane (15 mL) and trifluoroacetic acid (15 mL) at 35°C for 48 hours. The mixture was concentrated, taken up in dichloromethane (100 mL), and stirred, and saturated Na₂CO₃ solution (20 mL) was added slowly. The solution was separated and the organic layer was dried over Na₂SO₄, filtered, and concentrated to provide the title compound.

EXAMPLE 302B

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperidin-1-yl)benzoate

[1172] The title compound was prepared by substituting EXAMPLE 302A for EXAMPLE 263G in EXAMPLE 263H.

EXAMPLE 302C

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperidin-1-yl)benzoic acid

[1173] The title compound was prepared by substituting EXAMPLE 302B for EXAMPLE 3I in EXAMPLE 3J.

EXAMPLE 302D

1,1-difluoro-4-methylenecyclohexane

[1174] Butyllithium (12.32 mL, 2.5 M solution in hexanes) was added to a solution of methyltriphenylphosphonium chloride (9.63 g) in tetrahydrofuran (50 mL) at 0°C, and the reaction was stirred for 5 minutes. 4,4-Difluorocyclohexanone (3.76 g) in dioxane (150 mL) was then added, and the reaction was stirred for 30 minutes. Water (3 mL) was added, and then hexane (150 mL) was slowly added, the reaction was filtered, and the solution carried on.

EXAMPLE 302E

4,4-difluoro-1-(hydroxymethyl)cyclohexanol

[1175] To the solution from EXAMPLE 302D was added water (75 mL), then N-methylmorpholine-N-oxide (6.4 mL, 50% solution in water) and OsO₄ (14.2 g, 2.5 wt % solution in tert-butanol) were added, and the reaction was stirred for 96 hours at 50°C. The solution was cooled to room temperature, treated with saturated aqueous Na₂S₂O₃ solution (100 mL) for 30 minutes, and then acidified with concentrated aqueous HCl. The solution was then extracted three times with ethyl acetate, and the organic layers were combined, washed with 1M HCl, and brine, and concentrated. The crude mixture was chromatographed on silica gel using 10-100% ethyl acetate in hexanes, and then 5% methanol in ethyl acetate to give the product.

EXAMPLE 302F

5-chloro-6-((4,4-difluoro-1-hydroxycyclohexyl)methoxy)pyridine-3-sulfonamide

- 5 **[1176]** This example was prepared by substituting EXAMPLE 302E for tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

EXAMPLE 302G

- 10 N-({5-chloro-6-((4,4-difluoro-1-hydroxycyclohexyl)methoxy)pyridin-3-yl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperidin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [1177]** The title compound was prepared by substituting EXAMPLE 302C for EXAMPLE 1E and EXAMPLE 302F for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-d₆) δ 11.65 (br s, 2H), 8.51 (s, 1H), 8.18 (s, 1H), 8.02 (d, 1H), 7.53 (m, 3H), 7.35 (d, 2H), 7.04 (d, 2H), 6.69 (dd, 1H), 6.39 (dd, 1H), 6.21 (d, 1H), 4.88 (s, 1H), 4.27 (s, 2H), 3.10 (m, 4H), 2.88 (m, 1H), 2.33 (m, 2H), 2.15 (m, 4H), 1.97 (s, 2H), 1.91 (m, 2H), 1.73 (m, 4H), 1.52 (m, 1H), 1.40 (m, 2H), 1.31 (m, 1H), 0.93 (s, 3H), 0.91 (m, 2H).

EXAMPLE 303

- 20 N-([3-chloro-4-([trans-4-(morpholin-4-yl)cyclohexyl)methoxy]phenyl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 303A

- 25 trans-4-morpholinocyclohexyl)methanol

- [1178]** To *tert*-butyl trans-4-(hydroxymethyl)cyclohexylcarbamate (0.500 g) was added hydrogen chloride (4.0M in dioxane, 2.2 mL) and the reaction was stirred for 1 hour and concentrated. The resulting solid was dissolved in acetonitrile (4 mL) and treated with N,N-diisopropylethylamine (1.523 mL) followed by 1-bromo-2-(2-bromoethoxy)ethane (0.556 g) and heated to 60°C. After stirring overnight the reaction was concentrated, loaded onto silica gel (Reveleris 40 g) and eluted using a gradient of 1% to 10% methanol/dichloromethane over 30 minutes (flow = 40 mL/min) to provide the title compound.

EXAMPLE 303B

3-chloro-4-(((1*r*,4*r*)-4-morpholinocyclohexyl)methoxy)benzenesulfonamide

- 40 **[1179]** The title compound was prepared by substituting EXAMPLE 303A for (4-fluoro-1-methylpiperidin-4-yl)methanol in EXAMPLE 283A.

EXAMPLE 303C

- 45 N-([3-chloro-4-([trans-4-(morpholin-4-yl)cyclohexyl)methoxy]phenyl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- [1180]** The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 303B for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1H), 10.96 - 10.59 (m, 1H), 8.02 (d, 1H), 7.82 (d, 1H), 7.69 (s, 1H), 7.50 (dd, 3H), 7.38 - 7.30 (m, 2H), 7.15 - 6.99 (m, 3H), 6.65 (dd, 1H), 6.39 (dd, 1H), 6.20 (d, 1H), 3.91 (d, 2H), 3.64 (s, 4H), 3.04 (s, 4H), 2.73 (s, 7H), 2.18 (s, 6H), 1.93 (m, 6H), 1.80 - 1.65 (m, 1H), 1.32 (m, 6H), 0.92 (s, 6H).

EXAMPLE 304

- 55 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([4-({3-[cyclopropyl(1,3-thiazol-5-ylmethyl)amino]propyl}amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1181] To a solution of EXAMPLE 291E (95 mg) in dichloromethane (2 mL) and acetic acid (0.5 mL) was added

thiazole-5-carbaldehyde (13 mg) followed by sodium triacetoxymethylborohydride (35 mg). The mixture was stirred overnight. The mixture was diluted with dichloromethane (300 mL) and washed with aqueous NaHCO₃, water, and brine and dried over Na₂SO₄. Filtration and evaporation of the solvent gave crude product which was dissolved in dimethylsulfoxide/methanol (6 mL, 1:1) and loaded on Gilson, C18(100A) 250x121.2 mm(10micron), with 30% acetonitrile to 65% acetonitrile over 40 minutes. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1 H), 8.95 (s, 1 H), 8.57 (m, 2 H), 8.03 (d, 1 H), 7.78 (m, 2 H), 7.49 (m, 3 H), 7.35 (m, 2 H), 7.02 (m, 3 H), 6.67 (dd, 1 H), 6.38 (dd, 1 H), 6.19 (d, 1 H), 4.00 (s, 2 H), 3.05 (d, 4 H), 2.73 (m, 2 H), 2.60 (m, 2 H), 2.18 (m, 7 H), 1.95 (s, 2 H), 1.79 (m, 3 H), 1.37 (m, 3 H), 0.92 (s, 6 H), 0.45 (m, 4 H).

EXAMPLE 305

N-({3-chloro-4-[(trans-4-hydroxycyclohexyl)methoxy]phenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 305A

3-chloro-4-((trans-4-hydroxycyclohexyl)methoxy)benzenesulfonamide

[1182] (Trans-4-(tert-butyldimethylsilyloxy)cyclohexyl)methanol (275 mg, prepared according to a procedures in WO 2008/124878) and 3-chloro-4-fluorobenzenesulfonamide (259 mg) in tetrahydrofuran (15 mL) were treated with sodium hydride (180 mg, 60%) overnight. The reaction was quenched with water (1 mL) and trifluoroacetic acid (4 mL) was added. The resulting mixture was stirred for 1 hour and concentrated. The residue was triturated with water and methanol to provide the title compound.

EXAMPLE 305B

N-({3-chloro-4-[(trans-4-hydroxycyclohexyl)methoxy]phenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1183] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 305A in place of EXAMPLE 11B. ¹H NMR (500 MHz, dimethylsulfoxide-d₆) δ 11.71 (s, 1 H), 11.38 (s, 1 H), 8.06 (d, 1 H), 7.87 (d, 1 H), 7.76 (dd, 1 H), 7.57 (d, 1 H), 7.51 - 7.55 (m, 1 H), 7.49 (d, 1 H), 7.34 (d, 2 H), 7.18 (d, 1 H), 7.04 (d, 2 H), 6.67 (dd, 1 H), 6.42 (dd, 1 H), 6.18 (d, 1 H), 4.54 (d, 1 H), 3.91 (d, 2 H), 3.07 (s, 4 H), 2.75 (s, 2 H), 2.17 (d, 6 H), 1.95 (s, 2 H), 1.78 - 1.90 (m, 4 H), 1.63 - 1.75 (m, 1 H), 1.38 (t, 2 H), 1.00 - 1.25 (m, 4 H), 0.92 (s, 6 H).

EXAMPLE 306

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-chloro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 306A

3-chloro-4-((tetrahydro-2H-pyran-4-yl)methylamino)benzenesulfonamide

[1184] The title compound was prepared by substituting 4-fluoro-3-chlorobenzenesulfonamide for 4-chloro-3-nitrobenzenesulfonamide, (tetrahydro-2H-pyran-4-yl)methanamine for 4-methylpiperazin-1-amine dihydrochloride and Hunig's base for N¹,N¹,N²,N²-tetramethylethane-1,2-diamine in EXAMPLE 6A.

EXAMPLE 306B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-chloro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1185] The title compound was prepared by substituting EXAMPLE 306A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400MHz, dimethylsulfoxide-d₆) δ 11.80 (s, 1H), 11.17 (br s, 1H), 8.09 (d, 1H), 7.71 (d, 1H), 7.63 (d, 1H), 7.58 (dd, 1H), 7.53 (dd, 1H), 7.50 (d, 1H), 7.34 (d, 2H), 7.03 (d, 2H), 6.74 (d, 1H), 6.66 (dd, 1H), 6.42 (m, 1H), 6.40 (t, 1H), 6.16 (d, 1H), 3.83 (m, 2H), 3.24 (m, 2H), 3.10 (m, 2H), 3.06 (br m, 4H), 2.72 (s, 2H), 2.17 (br m, 6H), 1.95 (s, 2H), 1.83 (m, 1H), 1.59 (br m, 2H), 1.38 (t, 2H), 1.20 (ddd, 2H), 0.92 (s, 6H).

EXAMPLE 307

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-(trifluoromethyl)phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 307A

4-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)-3-(trifluoromethyl)benzenesulfonamide

[1186] The title compound was prepared by substituting 4-fluoro-3-(trifluoromethyl)benzenesulfonamide for 4-fluoro-3-nitrobenzenesulfonamide and EXAMPLE 37C for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 307B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-(trifluoromethyl)phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1187] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 307A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (400 MHz, pyridine-d₅) δ 13.07 (s, 1H), 8.78 (d, 1H), 8.58 (dd, 1H), 8.42 (d, 1H), 8.09 (d, 1H), 7.67 (t, 1H), 7.65 (d, 1H), 7.43 (m, 2H), 7.16 (d, 1H), 7.06 (m, 2H), 6.74 (dd, 1H), 6.51 (m, 2H), 4.21 (d, 2H), 3.87 (m, 2H), 3.78 (td, 2H), 3.06 (m, 4H), 2.76 (s, 2H), 2.25 (t, 2H), 2.13 (m, 4H), 1.95 (m, 6H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 308

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-({3-[cyclopropyl(2,2,2-trifluoroethyl)amino]propyl}amino)-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 308A

4-(3-(cyclopropylamino)propylamino)-3-nitrobenzenesulfonamide

[1188] To a solution of EXAMPLE 291C (4.14 g) in dichloromethane (10 mL) was added trifluoroacetic acid (10 mL). The mixture was stirred for 2 hours. The mixture was concentrated under vacuum and the residue was dissolved in dichloromethane (300 mL) and washed with aqueous NaHCO₃, water, and brine and dried over Na₂SO₄. Filtration and evaporation of solvent gave the title compound.

EXAMPLE 308B

4-(3-(cyclopropyl(2,2,2-trifluoroethyl)amino)propylamino)-3-nitrobenzenesulfonamide

[1189] To a solution of EXAMPLE 308A (314 mg) in dichloromethane (6 mL) was added 2,2,2-trifluoroethyl trifluoromethanesulfonate (255 mg) and N,N-diisopropylethylamine (258 mg). The mixture was stirred overnight. The mixture was diluted with dichloromethane (300 mL) and washed with aqueous NaHCO₃, water, brine and dried over Na₂SO₄. Filtration and evaporation of solvent gave the title compound.

EXAMPLE 308C

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-({3-[cyclopropyl(2,2,2-trifluoroethyl)amino]propyl}amino)-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1190] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 308B for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 11.38 (m, 1H), 8.55 (d, 2H), 8.03 (d, 1H), 7.81 (dd, 1H), 7.50 (m, 3H), 7.34 (d, 2H), 7.05 (m, 3H), 6.67 (dd, 1H), 6.38 (dd, 1H), 6.19 (d, 1H), 3.07 (m, 4H), 2.82 (m, 4H), 2.18 (m, 7H), 1.38 (m, 2H), 0.92 (s, 6H), 0.44 (m, 4H).

EXAMPLE 309

N-[(3-chloro-4-[[1-(oxetan-3-yl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

5

[1191] To a solution of EXAMPLE 294B (0.150 g) in dichloromethane (2 mL) was added trifluoroacetic acid (1 mL). After stirring for 1 hour the reaction was concentrated and dried under high vacuum. The residue was dissolved in dichloromethane (2 mL) and treated with sodium triacetoxymethylborohydride (0.050 g) and oxetan-3-one (0.017 g) and stirred overnight at room temperature. The reaction was quenched with saturated aqueous NaHCO₃ (20 mL) and extracted into dichloromethane (50 mL). The organic layer was separated, washed with brine (25 mL), dried over Na₂SO₄, filtered, and concentrated. Silica gel chromatography (Reveleris 40 g) eluting with a gradient of 0.5% to 5% methanol/dichloromethane over 30 minutes (flow = 40 mL/min) provided the title compound. ¹H NMR (300 MHz, dimethylsulfoxide- d₆) δ 11.70 (s, 1H), 11.21 (s, 1H), 8.05 (d, 1H), 7.87 (d1H), 7.75 (dd, 1H), 7.61 - 7.42 (m, 3H), 7.42 - 7.26 (m, 2H), 7.18 (d, 1H), 7.14 - 6.97 (m, 2H), 6.67 (dd, 1H), 6.41 (dd, 1H), 6.18 (d, 1H), 4.51 (dt, 4H), 3.99 (d, 2H), 3.56 - 3.32 (m, 1H), 3.06 (s, 4H), 2.89 - 2.68 (m, 4H), 2.16 (d, 6H), 2.01 - 1.69 (m, 7H), 1.50 - 1.07 (m, 4H), 0.92 (s, 6H).

15

EXAMPLE 310

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3,5-difluoro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

20

EXAMPLE 310A

3,5-difluoro-4-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)benzenesulfonamide

25

[1192] EXAMPLE 37C (0.423 g) in tetrahydrofuran (30 mL) was treated with NaH (60% oil dispersion) (0.480 g), stirred 20 minutes at ambient temperature, treated with 3,4,5-trifluorobenzenesulfonamide (0.633 g) and stirred 30 minutes. N,N-Dimethylacetamide (15 mL) was added to increase solubility of the reactants and stirring was continued overnight at ambient temperature. Additional NaH (60% oil dispersion) (0.480 g) and N,N-dimethylacetamide (15 mL) were added and the mixture was heated overnight at 50°C. The reaction was quenched with saturated aqueous NH₄Cl solution and then partitioned between saturated aqueous NH₄Cl solution and ethyl acetate. The organic layer was washed with water and brine, dried (MgSO₄), filtered and concentrated. The concentrate was chromatographed on amine functionalized silica gel with 0 to 2% methanol in CH₂Cl₂ as the eluent. The residue was further purified by reverse phase HPLC on a C18 column using a gradient of 10-70% acetonitrile/0.1% trifluoroacetic acid in water to provide the title compound.

30

35

EXAMPLE 310B

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3,5-difluoro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

40

[1193] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 310A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (500 MHz, pyridine-d₅) δ 13.06 (s, 1H), 8.41 (d, 1H), 8.11 (m, 2H), 8.08 (d, 1H), 7.66 (m, 2H), 7.44 (m, 2H), 7.07 (m, 2H), 6.75 (dd, 1H), 6.52 (d, 1H), 6.50 (dd, 1H), 4.26 (d, 2H), 3.85 (dd, 1H), 3.83 (dd, 1H), 3.74 (m, 2H), 3.07 (m, 4H), 2.77 (s, 2H), 2.26 (t, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.87 (m, 4H), 1.39 (t, 2H), 0.94 (s, 6H).

45

EXAMPLE 311

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-({3-[cyclopropyl(oxetan-3-yl)amino]propyl}amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50

EXAMPLE 311 A

4-(3-(cyclopropyl(oxetan-3-yl)amino)propylamino)-3-nitrobenzenesulfonamide

55

[1194] To a solution of EXAMPLE 308A (314 mg) in dichloromethane (5 mL) was added oxetan-3-one (72 mg) followed by sodium triacetoxymethylborohydride (318 mg). The mixture was stirred overnight. The mixture was diluted with dichloromethane (300 mL) and washed with aqueous NaHCO₃, water and brine and dried over Na₂SO₄. After filtration, evaporation

of the solvent gave the crude title compound.

EXAMPLE 311B

5 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-(cyclopropyl(oxetan-3-yl)amino]propyl)amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 **[1195]** The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 311A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1 H), 11.37 (s, 1 H), 8.68 (s, 1 H), 8.54 (d, 1 H), 8.02 (d, 1 H), 7.79 (d, 1 H), 7.49 (m, 3 H), 7.34 (d, 2 H), 7.03 (m, 3 H), 6.67 (dd, 1 H), 6.38 (dd, 1 H), 6.19 (d, 1 H), 4.62 (m, 2 H), 4.48 (t, 2 H), 3.98 (m, 1 H), 3.37 (m, 2 H), 3.06 (m, 4 H), 2.73 (d, 2 H), 2.59 (m, 2 H), 2.23 (m, 6 H), 1.95 (s, 2 H), 1.74 (m, 3 H), 1.38 (t, 2 H), 0.92 (s, 6 H), 0.41 (m, 4 H).

EXAMPLE 312

15 N-([3-chloro-4-([1-(1-methyl-L-prolyl)piperidin-4-yl]methoxy)phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

20 **[1196]** To EXAMPLE 294B (0.065 g) was added hydrogen chloride (4.0M in dioxane, 0.339 mL) and a few drops of methanol. After 30 minutes, the reaction was concentrated, and (S)-1-methylpyrrolidine-2-carboxylic acid (0.013 g), N¹-((ethylimino)methylene)-N³,N³-dimethylpropane-1,3-diamine hydrochloride (0.026 g), suspended in dichloromethane (0.5 mL) were added followed by diisopropylethylamine (0.036 mL). The mixture stirred at room temperature. After stirring overnight, the reaction mixture was loaded onto silica gel (Reveleris 40g) and eluted using a gradient of 1% to 10% methanol (containing 1N NH₃)/dichloromethane over 30 minutes (flow = 40 mL/minutes) to provide the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.51 (s, 1H), 10.00 - 9.22 (m, 1H), 7.92 (d, 1H), 7.68 (d, 1H), 7.57 (d, 1H), 7.47 (dd, 1H), 7.44 - 7.38 (m, 1H), 7.38 - 7.31 (m, 2H), 7.29 (d, 1H), 7.12 - 7.01 (m, 2H), 6.90 (d, 1H), 6.61 (dd, 1H), 6.31 (dd, 1H), 6.25 (d, 1H), 5.85 (d, 1H), 4.40 (s, 1H), 3.92 (s, 4H), 3.17 - 2.89 (m, 8H), 2.73 (s, 4H), 2.38 (s, 3H), 2.18 (m, 6H), 1.96 (s, 2H), 1.80 (m, 2H), 1.57 (s, 2H), 1.39 (s, 2H), 1.22 (m, 2H), 0.96 (m, 6H).

EXAMPLE 313

30 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3,4-difluoro-5-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 313A

35 3,4-difluoro-5-((4-fluorotetrahydro-2H-pyran-4-yl)methoxy)benzenesulfonamide

40 **[1197]** The title compound was obtained as a side product in EXAMPLE 310A.

EXAMPLE 313B

45 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3,4-difluoro-5-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50 **[1198]** The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 313A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (400 MHz, pyridine-d₅) δ 13.05 (s, 1H), 8.41 (d, 1H), 8.10 (d, 1H), 7.98 (m, 2H), 7.66 (m, 1H), 7.63 (d, 1H), 7.44 (m, 2H), 7.07 (m, 2H), 6.77 (dd, 1H), 6.54 (d, 1H), 6.48 (dd, 1H), 4.12 (d, 2H), 3.83 (m, 2H), 3.75 (m, 2H), 3.08 (m, 4H), 2.77 (s, 2H), 2.26 (t, 2H), 2.15 (m, 4H), 1.97 (s, 2H), 1.82 (m, 4H), 1.39 (t, 2H), 0.94 (s, 6H).

55

EXAMPLE 314

N-[[5-chloro-6-[[[(2S)-4-cyclopropylmorpholin-2-yl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 314A

(S)-5-chloro-6-((4-cyclopropylmorpholin-2-yl)methoxy)pyridine-3-sulfonamide

[1199] A solution of EXAMPLE 244B (250 mg), anhydrous methanol (6 mL), (1-ethoxycyclopropoxy)trimethylsilane (0.474 mL), and acetic acid (0.509 mL) was heated at 70 °C for 30 minutes. After cooling to ambient temperature, sodium cyanoborohydride (112 mg) was added and the mixture was stirred for 18 hours. Additional sodium cyanoborohydride (75 mg) was added and stirring was continued 18 hours. The reaction was concentrated and the residue was partitioned between methylene chloride and saturated sodium bicarbonate solution. The crude product was isolated from the dried methylene chloride layer and was purified on silica gel and was eluted with a 1, 2.5, 5, 10% methanol in methylene chloride step gradient to provide the title compound.

EXAMPLE 314B

N-[[5-chloro-6-[[[(2S)-4-cyclopropylmorpholin-2-yl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1200] The title compound was prepared by substituting EXAMPLE 314A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (400 MHz, pyridine-*d*₅) δ 12.98 (s, 1H), 9.09 (d, 1H), 8.69 (d, 1H), 8.41 (d, 1H), 8.11 (d, 1H), 7.66 - 7.64 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.75 (dd, 1H), 6.53 (m, 1H), 6.48 (m, 1H), 5.72 (br s, 1H), 4.62 - 4.57 (m, 1H), 4.51 - 4.47 (m, 1H), 3.99 (m, 1H), 3.85 (m, 1H), 3.57 (m, 1H), 3.08 - 3.01 (m, 5H), 2.77 (s, 2H), 2.69 (m, 1H), 2.39 - 2.24 (m, 4H), 2.14 (m, 4H), 1.97 (s, 2H), 1.57 (m, 1H), 1.39 (t, 2H), 0.94 (m, 6H), 0.48 - 0.3 (m, 4H).

EXAMPLE 315

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperidin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1201] The title compound was prepared by substituting EXAMPLE 302C for EXAMPLE 1E and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-*d*₆) δ 11.70 (s, 1H), 11.35 (br s, 1H), 8.61 (m, 1H), 8.57 (d, 1H), 8.04 (d, 1H), 7.82 (dd, 1H), 7.45-7.57 (m, 3H), 7.33 (d, 2H), 7.15 (d, 1H), 7.01 (d, 2H), 6.65 (dd, 1H), 6.40 (dd, 1H), 6.11 (d, 1H), 3.85 (dd, 2H), 3.53 (m, 2H), 3.27 (m, 4H), 2.63 (m, 2H), 2.04 (m, 2H), 1.91 (s, 2H), 1.77 (m, 2H), 1.62 (m, 4H), 1.45 (m, 2H), 1.38 (m, 2H), 1.27 (m, 1H), 1.23 (m, 4H), 0.92 (s, 6H).

EXAMPLE 316

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperidin-1-yl)-N-([3-chloro-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 316A

3-chloro-4-((tetrahydro-2H-pyran-4-yl)methoxy)benzenesulfonamide

[1202] The title compound was prepared by substituting (tetrahydro-2H-pyran-4-yl)methanol for (4-fluoro-1-methylpiperidin-4-yl)methanol in EXAMPLE 283A.

EXAMPLE 316B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperidin-1-yl)-N-([3-chloro-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1203] The title compound was prepared by substituting EXAMPLE 302C for EXAMPLE 1E and EXAMPLE 316A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300MHz, dimethylsulfoxide-*d*₆) δ 11.77 (s, 1H), 11.35 (br s, 1H), 8.06 (m, 1H),

7.88 (d, 1H), 7.79 (dd, 1H), 7.58 (s, 1H), 7.53 (t, 1H), 7.46 (d, 1H), 7.34 (d, 2H), 7.22 (d, 1H), 7.01 (d, 2H), 6.66 (dd, 1H), 6.42 (dd, 1H), 6.11 (d, 1H), 3.99 (d, 2H), 3.88 (dd, 2H), 3.52 (m, 2H), 3.34 (m, 4H), 2.62 (m, 2H), 2.04 (m, 4H), 1.76 (m, 2H), 1.68 (m, 2H), 1.46 (m, 2H), 1.38 (m, 4H), 0.92 (s, 6H), 0.75 (m, 2H).

5 EXAMPLE 317

methyl 2-([(4-{[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl}-2-nitrophenyl)amino)methyl]morpholine-4-carboxylate

10 EXAMPLE 317A

methyl 2-((2-nitro-4-sulfamoylphenylamino)methyl)morpholine-4-carboxylate

[1204] The title compound was prepared by substituting methyl chloroformate for methyl iodide in EXAMPLE 134B.

15

EXAMPLE 317B

methyl 2-([(4-{[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl}-2-nitrophenyl)amino)methyl]morpholine-4-carboxylate

20

[1205] The title compound was prepared by substituting EXAMPLE 317A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.26 (d, 1H), 8.84 (t, 1H), 8.43 (d, 1H), 8.35 (d, 1H), 8.12 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.91 (bs, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 4.29-4.03 (m, 1H), 3.89-3.70 (m, 3H), 3.71 (s, 3H), 3.55-3.38 (m, 3H), 3.07 (m, 4H), 2.96 (dt, 1H), 2.86 (dd, 1H), 2.77 (s, 2H), 2.26 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

25

EXAMPLE 318

2-([(4-{[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl}-2-nitrophenyl)amino)methyl]-N-ethyl-N-methylmorpholine-4-carboxamide

30

EXAMPLE 318A

N-ethyl-N-methyl-2-((2-nitro-4-sulfamoylphenylamino)methyl)morpholine-4-carboxamide

35

[1206] The title compound was prepared by substituting *N*-methyl-*N*-ethyl carbamyl chloride for methyl iodide in EXAMPLE 134B.

EXAMPLE 318B

40

2-([(4-{[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl}-2-nitrophenyl)amino)methyl]-N-ethyl-N-methylmorpholine-4-carboxamide

[1207] The title compound was prepared by substituting EXAMPLE 318A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.26 (d, 1H), 8.86 (t, 1H), 8.44 (d, 1H), 8.33 (dd, 1H), 8.12 (d, 1H), 7.67 (t, 1H), 7.64 (d, 1H), 7.44 (d, 2H), 7.07 (d, 2H), 6.91 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 3.92-3.85 (m, 2H), 3.75 (d, 1H), 3.62 (dt, 1H), 3.55-3.48 (m, 1H), 3.45-3.39 (m, 2H), 3.21 (q, 2H), 3.07 (m, 4H), 2.99 (dt, 1H), 2.90 (dd, 1H), 2.77 (s, 2H), 2.76 (s, 3H), 2.26 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 1.06 (t, 3H), 0.93 (s, 6H).

45

50

55

EXAMPLE 319

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{[4-({[4-(methylsulfonyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 319A

4-({[4-(methylsulfonyl)morpholin-2-yl]methylamino)-3-nitrobenzenesulfonamide

[1208] The title compound was prepared by substituting methanesulfonyl chloride for methyl iodide in EXAMPLE 134B.

EXAMPLE 319B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{[4-({[4-(methylsulfonyl)morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1209] The title compound was prepared by substituting EXAMPLE 319A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.25 (d, 1H), 8.84 (t, 1H), 8.43 (d, 1H), 8.32 (dd, 1H), 8.13 (d, 1H), 7.67 (t, 1H), 7.65 (d, 1H), 7.44 (d, 2H), 7.07 (d, 2H), 6.92 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 3.99 (m, 1H), 3.92-3.88 (m, 2H), 3.64 (m, 2H), 3.56 (m, 1H), 3.50 (m, 1H), 3.07 (m, 4H), 3.04 (s, 3H), 2.95-2.88 (m, 2H), 2.77 (s, 2H), 2.26 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 320

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{[4-({[3-(cyclobutyl(cyclopropyl)amino]propyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 320A

4-(3-(cyclobutyl(cyclopropyl)amino)propylamino)-3-nitrobenzenesulfonamide

[1210] To a solution of EXAMPLE 308A (314 mg) in dichloromethane (5 mL) was added cyclobutanone (70 mg) followed by sodium triacetoxyborohydride (318 mg). The mixture was stirred overnight. The mixture was diluted with dichloromethane (300 mL) and washed with aqueous NaHCO₃, water and brine and dried over Na₂SO₄. After filtration, evaporation of solvent gave the title compound.

EXAMPLE 320B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{[4-({[3-(cyclobutyl(cyclopropyl)amino]propyl}amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1211] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 320A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1 H), 8.70 (m, 1 H), 8.54 (d, 1 H), 8.02 (d, 1 H), 7.79 (dd, 1 H), 7.49 (m, 3 H), 7.34 (d, 2 H), 7.03 (m, 3 H), 6.66 (dd, 1 H), 6.38 (dd, 1 H), 6.19 (d, 1 H), 3.37 (q, 2 H), 3.06 (m, 4 H), 2.73 (s, 2 H), 2.63 (m, 2 H), 2.21 (m, 8 H), 1.82 (m, 3 H), 1.53 (m, 2 H), 1.38 (t, 2 H), 0.94 (m, 6 H), 0.41 (m, 4 H).

EXAMPLE 321

4-(4-{[2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({[3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 321A

ethyl 5,5-difluoro-2-oxocyclohexanecarboxylate

[1212] To a solution of diethyl 4,4-difluoroheptanedioate (4.3 g) in toluene (50 mL) was added potassium 2-methylpropan-2-olate (2.87 g) and the reaction stirred overnight at room temperature. The reaction was quenched with 1N

aqueous HCl (100 mL) and extracted with diethyl ether (150 mL). The ether layer was washed with brine (50 mL), dried over magnesium sulfate, filtered, and concentrated. Silica gel chromatography (Reveleris 40 g) eluting with a gradient of 1% to 5% ethyl acetate/hexanes gave the title compound.

5 EXAMPLE 321B

ethyl 5,5-difluoro-2-(trifluoromethylsulfonyloxy)cyclohex-1-enecarboxylate

10 **[1213]** To a solution of EXAMPLE 321A (2.37 g) in dichloromethane (40 mL) at 0°C was added N,N-diisopropylethylamine (5.02 mL) followed by trifluoromethanesulfonic anhydride (2.33 mL) and the reaction was allowed to slowly warm to room temperature. After stirring overnight the reaction was quenched with 10 ml of water then 1N aqueous HCl (100 mL). The reaction was extracted with dichloromethane (3 x 75 mL), and the combined organics were washed with brine (50 mL) and concentrated. Silica gel chromatography (Reveleris 40 g) eluting with a gradient of 1% to 25% ethyl acetate/hexanes gave the title compound.

15

EXAMPLE 321C

ethyl 2-(4-chlorophenyl)-5,5-difluorocyclohex-1-enecarboxylate

20 **[1214]** A solution of EXAMPLE 321B (3.47 g), 4-chlorophenylboronic acid (1.925 g) and cesium fluoride (3.43 g) in 30 ml of 1,2-dimethoxyethane and 15 ml of ethanol was degassed with nitrogen for 5 minutes. Tetrakis(triphenylphosphine)palladium(0) (0.237 g) was added and the reaction was heated to 70°C. The reaction was diluted with ether (200 mL) and washed with 1N aqueous HCl (100 mL) and brine (100 mL), dried over magnesium sulfate, filtered, and concentrated. Silica gel chromatography (Reveleris 40 g) eluting with a gradient of 1% to 8% ethyl acetate/hexanes over 40 minutes gave the title compound.

25

EXAMPLE 321D

(2-(4-chlorophenyl)-5,5-difluorocyclohex-1-enyl)methanol

30

[1215] To a solution of EXAMPLE 321C (1.84 g) in diethyl ether (25 mL) at 0°C was added lithium aluminum hydride (1.0M, 4.28 mL). The reaction was quenched with the dropwise addition of water, then 1N aqueous HCl (50 mL) was added and the reaction diluted with diethyl ether (100 mL). The organic layer was separated, washed with brine (50 mL) dried over magnesium sulfate, filtered and concentrated to provide the title compound.

35

EXAMPLE 321E

2-(4-chlorophenyl)-5,5-difluorocyclohex-1-enecarbaldehyde

40 **[1216]** To a solution of EXAMPLE 321D (1.38 g) in dichloromethane (25 mL) was added Dess-Martin periodinane (2.489 g) and the reaction stirred for 1 hour at room temperature. The reaction was quenched with 1N aqueous NaOH solution (75 mL) and the product was extracted into dichloromethane (2 x 100 mL). The combined organics were washed with brine (75 mL), dried over magnesium sulfate, filtered, and concentrated. Silica gel chromatography (Reveris 80 g) eluting with a gradient of 1% to 10% ethyl acetate/hexanes over 40 minutes gave the title compound.

45

EXAMPLE 321F

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5,5-difluorocyclohex-1-enyl)methyl)piperazin-1-yl)benzoate

50

[1217] The title compound was prepared by substituting EXAMPLE 321E for EXAMPLE 15E in EXAMPLE 15G.

EXAMPLE 321G

55 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5,5-difluorocyclohex-1-enyl)methyl)piperazin-1-yl)benzoic acid

[1218] The title compound was prepared by substituting EXAMPLE 321F for EXAMPLE 15G in EXAMPLE 15H.

EXAMPLE 321H

4-(4-([2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([tetrahydro-2H-pyran-4-ylmethyl]amino)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

5

[1219] The title compound was prepared by substituting EXAMPLE 321G for EXAMPLE 1E and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide- d₆) δ 11.74 - 11.63 (m, 1H), 11.53 - 11.29 (m, 1H), 8.57 (d, 2H), 8.05 (d, 1H), 7.85 - 7.77 (m, 1H), 7.49 (d, 3H), 7.38 (d, 2H), 7.16 - 7.06 (m, 3H), 6.73 - 6.64 (m, 1H), 6.43 - 6.36 (m, 1H), 6.21 - 6.14 (m, 1H), 3.93 - 3.77 (m, 2H), 3.29 (d, 4H), 3.07 (s, 4H), 2.79 - 2.57 (m, 4H), 2.45 (dd, 2H), 2.19 (s, 6H), 1.99 - 1.80 (m, 1H), 1.70 - 1.54 (m, 2H), 1.38 - 1.13 (m, 2H).

10

EXAMPLE 322

N-([3-chloro-4-([4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy)phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

15

EXAMPLE 322A

tert-butyl 4-fluoro-4-(hydroxymethyl)piperidine-1-carboxylate

20

[1220] 1-Tert-butyl 4-ethyl 4-fluoropiperidine-1,4-dicarboxylate (2 g) was taken up in tetrahydrofuran (20 ml) and cooled in an ice bath. Lithium aluminum hydride (1.0M in dioxane, 5.09 mL) was added dropwise. The reaction was stirred at room temperature for 2 hours. The reaction was quenched with water and with 1M aqueous NaOH solution and then stirred another 1 hour at room temperature. The mixture was extracted with ethyl acetate, and the extracts were combined and washed with water and with brine, dried over MgSO₄, filtered and concentrated under vacuum. The crude product was used without further purification.

25

EXAMPLE 322B

tert-butyl 4-((2-chloro-4-sulfamoylphenoxy)methyl)-4-fluoropiperidine-1-carboxylate

30

[1221] The title compound was prepared by substituting EXAMPLE 322A for (tetrahydro-2H-pyran-4-yl)methanol and 3-chloro-4-fluorobenzenesulfonamide for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

35

EXAMPLE 322C

3-chloro-4-((4-fluoropiperidin-4-yl)methoxy)benzenesulfonamide

[1222] The title compound was prepared by substituting EXAMPLE 322B for EXAMPLE 1A in EXAMPLE 1B.

40

EXAMPLE 322D

3-chloro-4-((4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methoxy)benzenesulfonamide

45

[1223] To a solution of EXAMPLE 322C (830 mg) in tetrahydrofuran (15 mL) and acetic acid (5 mL) was added oxetan-3-one (163 mg) and MP-cyanoborohydride (2.38 mmol/g, 1.9 g). The mixture was stirred at room temperature overnight. The reaction was then filtered and the filtrate was concentrated under vacuum. The residue was slurried in ether and the solid product was collected by filtration.

50

EXAMPLE 322E

N-([3-chloro-4-([4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy)phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

55

[1224] The title compound was prepared by substituting EXAMPLE 322D for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.71 (s, 1H), 8.06 (d, 1H), 7.89 (d, 1H), 7.79 (m, 1H), 7.58 (d, 1H), 7.52 (t, 1H), 7.49 (d, 1H), 7.34 (d, 2H), 7.25 (d, 1H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.42 (m, 1H), 6.18 (d, 1H), 4.55 (t, 2H), 4.44 (t, 2H), 4.24 (d, 2H), 3.44 (m, 2H), 3.07 (br s, 4H), 2.74 (m, 2H), 2.59 (m, 2H), 2.14 (m,

7H), 1.95 (m, 4H), 1.78 (m, 2H), 1.38 (t, 2H), 0.92 (s, 6H).

EXAMPLE 323

5 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-chloro-4-(tetrahydrofuran-3-yl-methoxy)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 323A

10 3-chloro-4-((tetrahydrofuran-3-yl)methoxy)benzenesulfonamide

[1225] The title compound was prepared by substituting 4-fluoro-3-chlorobenzenesulfonamide for 4-fluoro-3-nitrobenzenesulfonamide and (tetrahydrofuran-3-yl)methanol for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A, except here, dimethylformamide was used in place of tetrahydrofuran and the reaction was heated at 70° C for two days.

15

EXAMPLE 323B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-chloro-4-(tetrahydrofuran-3-yl-methoxy)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

20

[1226] The title compound was prepared by substituting EXAMPLE 323A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.73 (s, 1H), 8.07 (d, 1H), 7.89 (d, 1H), 7.80 (dd, 1H), 7.59 (d, 1H), 7.51 (dd, 1H), 7.49 (d, 1H), 7.34 (d, 2H), 7.23 (d, 1H), 7.03 (d, 2H), 6.66 (dd, 1H), 6.42 (m, 1H), 6.19 (d, 1H), 4.07 (m, 2H), 3.80 (m, 2H), 3.68 (m, 1H) 3.56 (m, 1H), 3.10 (br m, 4H), 2.85 (br s, 2H), 2.69 (m, 1H), 2.32 (br m, 4H), 2.17 (br m, 2H), 2.02 (m, 1H), 1.96 (s, 2H), 1.69 (m, 1H), 1.40 (t, 2H), 0.92 (s, 6H).

25

EXAMPLE 324 4-(4-([2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((trans-4-hydroxycyclohexyl)methyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

30

EXAMPLE 324A

4-((trans-4-hydroxycyclohexyl)methylamino)-3-nitrobenzenesulfonamide

[1227] The title compound was prepared by substituting EXAMPLE 120A for EXAMPLE 39B in EXAMPLE 39C.

35

EXAMPLE 324B

4-(4-([2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((trans-4-hydroxycyclohexyl)methyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

40

[1228] The title compound was prepared by substituting EXAMPLE 321G for EXAMPLE 1E and EXAMPLE 324A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide- d₆) δ 11.69 (s, 1H), 11.41 (s, 1H), 8.65 - 8.50 (m, 2H), 8.05 (d, 1H), 7.80 (dd, 1H), 7.60 - 7.44 (m, 3H), 7.41 - 7.34 (m, 2H), 7.14 - 7.02 (m, 3H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.17 (d, 1H), 4.50 (d, 1H), 3.23 (t, 2H), 3.06 (s, 4H), 2.70 (d4H), 2.44 (s, 2H), 2.33 - 1.94 (m, 6H), 1.78 (dd, 4H), 1.51 (d, 2H), 1.23 (s, 2H), 1.16 - 0.92 (m, 2H).

45

EXAMPLE 325

N-([3-chloro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl]sulfonyl)-4-(4-([9-(4-chlorophenyl)-3-(oxetan-3-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50

EXAMPLE 325A

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((9-(4-chlorophenyl)-3-(oxetan-3-yl)-3-azaspiro[5.5]undec-8-en-8-yl)methyl)piperazin-1-yl)benzoate

55

[1229] The title compound was prepared by substituting oxetan-3-one for 1,3-difluoropropan-2-one in EXAMPLE 265G.

EXAMPLE 325B

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((9-(4-chlorophenyl)-3-(oxetan-3-yl)-3-azaspiro[5.5]undec-8-en-8-yl)methyl)piperazin-1-yl)benzoic acid

[1230] The title compound was prepared by substituting EXAMPLE 325A for EXAMPLE 15G in EXAMPLE 15H.

EXAMPLE 325C

N-({3-chloro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-4-(4-{[9-(4-chlorophenyl)-3-(oxetan-3-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1231] The title compound was prepared by substituting EXAMPLE 325B for EXAMPLE 1E and EXAMPLE 286A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide- d₆) δ 11.67 (s, 1H), 11.13 (s, 1H), 8.05 (d, 1H), 7.87 (d, 1H), 7.80 - 7.70 (m, 1H), 7.59 - 7.46 (m, 3H), 7.34 (d, 2H), 7.21 (d, 1H), 7.11 - 7.03 (m, 2H), 6.66 (d, 1H), 6.41 (dd, 1H), 6.18 (d, 1H), 4.50 (dd, 4H), 4.26 (d, 2H), 3.85 - 3.69 (m, 2H), 3.61 (d, 3H), 3.05 (s, 4H), 2.69 (s, 2H), 2.37 (s, 4H), 2.17 (s, 6H), 2.04 (s, 2H), 1.87 (d, 4H), 1.49 (d, 6H).

EXAMPLE 326

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[(2R)-4-cyclopropylmorpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 326A

(R)-4-((4-cyclopropylmorpholin-2-yl)methylamino)-3-nitrobenzenesulfonamide

[1232] The title compound was prepared by substituting EXAMPLE 258E for EXAMPLE 173A in EXAMPLE 173B.

EXAMPLE 326B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[(2R)-4-cyclopropylmorpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1233] The title compound was prepared by substituting EXAMPLE 326A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.26 (d, 1H), 8.88 (t, 1H), 8.44 (d, 1H), 8.34 (dd, 1H), 8.12 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.94 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 3.88 (d, 1H), 3.84-3.81 (m, 1H), 3.59 (dt, 1H), 3.50-3.40 (m, 2H), 3.07 (m, 4H), 2.93 (d, 1H), 2.77 (s, 2H), 2.69 (d, 1H), 2.34 (dt, 1H), 2.26 (m, 2H), 2.21 (t, 1H), 2.14 (m, 4H), 1.97 (s, 2H), 1.58 (m, 1H), 1.39 (t, 2H), 0.94 (s, 6H), 0.45-0.39 (m, 4H).

EXAMPLE 327

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[(2S)-4-cyclopropylmorpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 327A

(S)-4-((4-cyclopropylmorpholin-2-yl)methylamino)-3-nitrobenzenesulfonamide

[1234] The title compound was prepared by substituting EXAMPLE 259E for EXAMPLE 173A in EXAMPLE 173B.

EXAMPLE 327B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({[(2S)-4-cyclopropylmorpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1235] The title compound was prepared by substituting EXAMPLE 327A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.26 (d, 1H), 8.88 (t, 1H), 8.44 (d, 1H), 8.34 (dd, 1H), 8.12 (d, 1H), 7.66 (m,

2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.94 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 3.88 (d, 1H), 3.84-3.81 (m, 1H), 3.59 (dt, 1H), 3.50-3.40 (m, 2H), 3.07 (m, 4H), 2.93 (d, 1H), 2.77 (s, 2H), 2.69 (d, 1H), 2.34 (dt, 1H), 2.26 (m, 2H), 2.21 (t, 1H), 2.14 (m, 4H), 1.97 (s, 2H), 1.58 (m, 1H), 1.39 (t, 2H), 0.94 (s, 6H), 0.45-0.39 (m, 4H).

5 EXAMPLE 328

4-(4-([5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 EXAMPLE 328A

spiro[2.5]octan-5-one

[1236] To a solution of 3-ethoxycyclohex-2-enone (48.1 mL) in ether (1000 mL) was added titanium(IV) isopropoxide (110 mL) followed by addition of ethylmagnesium bromide (357 mL) at ambient temperature. The reaction mixture was stirred for 2 hours at ambient temperature and was then quenched with water (500 mL). The organic layer was separated (decanted) and the water layer was extracted with ether (3 x 300 mL). The combined extracts were partially concentrated to approximately 300 mL. p-Toluenesulfonic acid monohydrate (3.0 g) was added and the reaction mixture was stirred overnight at ambient temperature. The reaction mixture was then washed with saturated aqueous NaHCO₃ solution., dried (MgSO₄), filtered and concentrated. The concentrate was purified by fractional distillation (1st fraction b.p. 27°C at 23 torr (not product), 2nd fraction (product) b.p. 75°C at 8 torr).

EXAMPLE 328B

25 5-chlorospiro[2.5]oct-5-ene-6-carbaldehyde

[1237] N,N-dimethylformamide (2.1 mL) in dichloromethane (3.2 mL) at -5 °C was treated slowly with POCl₃ (2.33 mL) keeping the bath temperature less than 0°C. The cooling bath was removed and the mixture was stirred at ambient temperature for 30 minutes. The reaction mixture was returned to the cooling bath and EXAMPLE 328A (2.484 g) in dichloromethane (4 mL) was added slowly to the reaction mixture. The reaction mixture was heated at 45°C for 15 hours, cooled to room temperature and then poured into a mixture of ice and saturated aqueous sodium acetate solution. After the ice melted, the mixture was extracted with diethyl ether. The combined extracts were washed with saturated aqueous NaHCO₃ solution and brine, dried (MgSO₄), filtered and concentrated. The concentrate was chromatographed with 0 to 10% CH₂Cl₂ in hexanes, then 25% CH₂Cl₂ in hexanes and then 100% CH₂Cl₂ as the eluents.

35 EXAMPLE 328C

5-(4-chlorophenyl)spiro[2.5]oct-5-ene-6-carbaldehyde

[1238] EXAMPLE 328B (2.9 g), 4-chlorophenylboronic acid (2.87 g), palladium(II) acetate (0.103 g), K₂CO₃ (5.28 g) and tetrabutylammonium bromide (4.93 g) were combined in a 100-mL round bottomed flask with water (17.0 mL). The flask was flushed with nitrogen and stirred at 45°C for 14 hours. The reaction mixture was partitioned between brine and diethyl ether. The organic layer was washed with brine, dried (MgSO₄), filtered through a plug of celite, concentrated and chromatographed on silica gel with 0 to 2% ethyl acetate in hexanes as the eluent.

45 EXAMPLE 328D

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl)methyl)piperazin-1-yl)benzoate

[1239] The title compound was prepared by substituting EXAMPLE 15F for tert-butyl piperazine carboxylate and EXAMPLE 328C for 4-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 328E

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl)methyl)piperazin-1-yl)benzoic acid hydrochloride

[1240] EXAMPLE 328D (0.85 g) in a mixture of tetrahydrofuran (4.8 mL), methanol (2.4 mL) and water (2.4 mL) was treated with LiOH·H₂O (0.184 g) and heated overnight at 50°C. The reaction mixture was cooled to room temperature, concentrated to remove tetrahydrofuran and methanol and acidified with 1 N aqueous HCl causing precipitation of the product. The solid was collected by filtration, rinsed with water and dried overnight in a vacuum oven at 80°C to provide the title compound.

EXAMPLE 328F

4-(4-([5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl)piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1241] The title compound was prepared by substituting EXAMPLE 328E for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (500 MHz, pyridine-d₅) δ 13.07 (s, 1H), 9.32 (d, 1H), 8.68 (t, 1H), 8.44 (d, 1H), 8.38 (dd, 1H), 8.10 (d, 1H), 7.68 (m, 1H), 7.66 (d, 1H), 7.43 (m, 2H), 7.10 (m, 2H), 6.91 (d, 1H), 6.75 (dd, 1H), 6.51 (m, 2H), 3.97 (dd, 2H), 3.30 (td, 2H), 3.16 (t, 2H), 3.06 (m, 4H), 2.81 (s, 2H), 2.37 (t, 2H), 2.16 (m, 4H), 2.11 (s, 2H), 1.81 (m, 1H), 1.58 (dd, 2H), 1.45 (t, 2H), 1.32 (qd, 2H), 0.38 (s, 4H).

EXAMPLE 329

N-[[5-chloro-6-({4-[cyclopropyl(oxetan-3-yl)amino]cyclohexyl}methoxy)pyridin-3-yl]sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 329A

ethyl 4-(cyclopropylamino)cyclohexanecarboxylate

[1242] To a solution of ethyl 4-oxocyclohexanecarboxylate (3.4 g) in dichloromethane (30 mL) was added cyclopropanamine (1.14 g) followed by sodium triacetoxymethylborohydride (4.24 g). The mixture was stirred overnight. The mixture was diluted with dichloromethane (300 mL) and washed with 2N NaOH, water, brine and dried over Na₂SO₄. Filtration and evaporation of the solvent gave the title compound.

EXAMPLE 329B

ethyl 4-(cyclopropyl(oxetan-3-yl)amino)cyclohexanecarboxylate

[1243] To a solution of EXAMPLE 329A (1.05g) in dichloromethane (10 mL) was added oxetan-3-one (0.358 g) followed by sodium triacetoxymethylborohydride (1.05 g). The mixture was stirred overnight. The mixture was diluted with dichloromethane (300 mL) and washed with 2N aqueous NaOH, water, brine and dried over Na₂SO₄. Filtration and evaporation of the solvent gave the title compound.

EXAMPLE 329C

(4-(cyclopropyl(oxetan-3-yl)amino)cyclohexyl)methanol

[1244] To a solution of EXAMPLE 329B (1.2 g) in tetrahydrofuran (20 mL) was added lithium aluminum hydride (0.681 g). The mixture was stirred overnight. 2N aqueous NaOH solution was added dropwise to the reaction mixture. The mixture was then diluted with ethyl acetate (300 mL) and washed with water, brine and dried over Na₂SO₄. Filtration and evaporation of the solvent gave the title compound.

EXAMPLE 329D

5-chloro-6-((4-(cyclopropyl(oxetan-3-yl)amino)cyclohexyl)methoxy)pyridine-3-sulfonamide

5 **[1245]** To a solution of EXAMPLE 329C (706 mg) in N,N-dimethylformamide (6 mL) was added NaH (60% in mineral oil, 300 mg). The mixture was stirred for 30 minutes, and then 5,6-dichloropyridine-3-sulfonamide (706 mg) was added. The mixture was stirred overnight. The mixture was poured over aqueous NH₄Cl and extracted with ethyl acetate (3x 200 mL). The combined organic layers were washed with water, brine and dried over Na₂SO₄. After filtration and evaporation of the solvent the residue was loaded on a silica gel cartridge and eluted with 5 to 10% 7N NH₃ in methanol in dichloromethane to provide the title compound.

EXAMPLE 329E

15 N-[[5-chloro-6-((4-(cyclopropyl(oxetan-3-yl)amino)cyclohexyl)methoxy)pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1246] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 329D for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.64 (s, 1 H), 8.50 (m, 1 H), 8.16 (s, 1 H), 8.02 (d, 1 H), 7.51 (m, 3 H), 7.35 (d, 2 H), 7.04 (d, 2 H), 6.67 (dd, 1 H), 6.38 (m, 1 H), 6.21 (s, 1 H), 4.70 (m, 2 H), 4.43 (t, 3 H), 4.19 (m, 2 H), 3.12 (m, 4 H), 2.84 (m, 2 H), 2.19 (m, 6 H), 1.96 (s, 3 H), 1.77 (m, 3 H), 1.38 (m, 7 H), 0.93 (s, 6 H), 0.44 (m, 4 H).

EXAMPLE 330

25 4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(cyclopropylmorpholin-2-yl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1247] The title compound was prepared by substituting EXAMPLE 328E for EXAMPLE 3J and EXAMPLE 218A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (400 MHz, pyridine-d₅) δ 13.01 (s, 1H), 9.26 (d, 1H), 8.88 (t, 1H), 8.43 (d, 1H), 8.34 (dd, 1H), 8.11 (d, 1H), 7.66 (m, 2H), 7.42 (m, 2H), 7.09 (m, 2H), 6.95 (d, 1H), 6.75 (dd, 1H), 6.53 (d, 1H), 6.49 (dd, 1H), 3.84 (m, 2H), 3.58 (td, 1H), 3.45 (m, 2H), 3.06 (m, 4H), 2.93 (d, 1H), 2.81 (s, 2H), 2.69 (d, 1H), 2.35 (m, 3H), 2.19 (m, 5H), 2.11 (s, 2H), 1.58 (m, 1H), 1.45 (t, 2H), 0.42 (m, 8H).

EXAMPLE 331

35 N-((3-chloro-4-[[4-(cyclopropylmorpholin-2-yl)methoxy]phenyl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 331A

40 *tert*-butyl 2-((2-chloro-4-sulfamoylphenoxy)methyl)morpholine-4-carboxylate

[1248] To a solution of *tert*-butyl 2-(hydroxymethyl)morpholine-4-carboxylate (0.478 g) in anhydrous N,N-dimethylformamide (5 mL) was added sodium hydride (0.280 g). The mixture was stirred at room temperature for 30 minutes, followed by addition of 3-chloro-4-fluorobenzenesulfonamide (0.419 g). The mixture was stirred at 40°C overnight. The reaction was quenched with water (10 mL), and the mixture was adjusted to ~pH 7 and extracted with ethyl acetate. The crude product was purified on a silica gel column eluting with 60% ethyl acetate in hexane to provide the title compound.

EXAMPLE 331B

3-chloro-4-(morpholin-2-ylmethoxy)benzenesulfonamide

[1249] The title compound was prepared by substituting EXAMPLE 331A for EXAMPLE 113A in EXAMPLE 134A.

EXAMPLE 331C

3-chloro-4-((4-cyclopropylmorpholin-2-yl)methoxy)benzenesulfonamide

5 **[1250]** The title compound was prepared by substituting EXAMPLE 331B for EXAMPLE 173A in EXAMPLE 173B.

EXAMPLE 331D

10 N-({3-chloro-4-[(4-cyclopropylmorpholin-2-yl)methoxy]phenyl)sulfonyl)-4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

15 **[1251]** The title compound was prepared by substituting EXAMPLE 331C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.04 (s, 1H), 8.54 (d, 1H), 8.43 (d, 1H), 8.27 (dd, 1H), 8.09 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 7.05 (d, 1H), 6.75 (dd, 1H), 6.52 (d, 1H), 6.50 (m, 1H), 4.20 (dd, 1H), 4.10 (dd, 1H), 3.94 (m, 1H), 3.86 (d, 1H), 3.58 (dt, 1H), 3.06 (m, 5H), 2.77 (s, 2H), 2.69 (d, 1H), 2.40-2.20 (m, 4H), 2.14 (m, 4H), 1.97 (s, 2H), 1.60 (m, 1H), 1.39 (t, 2H), 0.94 (s, 6H), 0.41 (m, 4H).

EXAMPLE 332

20 N-[(3-chloro-4-[(4-cyclopropylmorpholin-2-yl)methyl]amino)phenyl]sulfonyl]-4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 332A

25 tert-butyl 2-((2-chloro-4-sulfamoylphenylamino)methyl)morpholine-4-carboxylate

30 **[1252]** A solution of 3-chloro-4-fluorobenzenesulfonamide (1.0 g), tert-butyl 2-(aminomethyl)morpholine-4-carboxylate (1.135 g) and N-ethyl-N-isopropylpropan-2-amine (1.246 mL) in dimethylsulfoxide (15 mL) was stirred at 115°C for 72 hours. The mixture was concentrated, and the residue was purified on a silica gel column eluting with 60% ethyl acetate to provide the title compound.

EXAMPLE 332B

35 3-chloro-4-(morpholin-2-ylmethylamino)benzenesulfonamide

[1253] The title compound was prepared by substituting EXAMPLE 332A for EXAMPLE 113A in EXAMPLE 134A.

EXAMPLE 332C

40 **[1254]** The title compound was prepared by substituting EXAMPLE 332B for EXAMPLE 173A in EXAMPLE 173B.

EXAMPLE 332D

45 N-[(3-chloro-4-[(4-cyclopropylmorpholin-2-yl)methyl]amino)phenyl]sulfonyl]-4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50 **[1255]** The title compound was prepared by substituting EXAMPLE 332C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.05 (s, 1H), 8.45 (m, 2H), 8.21 (dd, 1H), 8.12 (d, 1H), 7.69 (d, 1H), 7.67 (t, 1H), 7.44 (d, 2H), 7.07 (d, 2H), 6.78 (d, 1H), 6.74 (dd, 1H), 6.52 (d, 1H), 6.50 (m, 1H), 6.37 (m, 1H), 3.84 (d, 1H), 3.77 (m, 1H), 3.54 (dt, 1H), 3.35 (m, 2H), 3.05 (m, 4H), 2.94 (d, 1H), 2.77 (s, 2H), 2.68 (d, 1H), 2.32 (dt, 1H), 2.26 (m, 2H), 2.18-2.12 (m, 5H), 1.97 (s, 2H), 1.55 (m, 1H), 1.39 (t, 2H), 0.94 (s, 6H), 0.41 (m, 4H).

55

EXAMPLE 333

2-[[[(2-chloro-4-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]phenyl)amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide

EXAMPLE 333A

2-((2-chloro-4-sulfamoylphenylamino)methyl)-N-ethyl-N-methylmorpholine-4-carboxamide

[1256] The title compound was prepared by substituting EXAMPLE 332B for EXAMPLE 134A and *N*-methyl-*N*-ethyl carbamyl chloride for methyl iodide in EXAMPLE 134B.

EXAMPLE 333B

2-[[[(2-chloro-4-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]phenyl)amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide

[1257] The title compound was prepared by substituting EXAMPLE 333A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.05 (s, 1H), 8.46 (s, 1H), 8.45 (s, 1H), 8.20 (dd, 1H), 8.10 (d, 1H), 7.69 (d, 1H), 7.67 (t, 1H), 7.44 (d, 2H), 7.07 (d, 2H), 6.79 (d, 1H), 6.73 (dd, 1H), 6.52 (dd, 1H), 6.49 (d, 1H), 6.43 (m, 1H), 3.83 (d, 2H), 3.73 (d, 1H), 3.59 (dt, 1H), 3.41-3.35 (m, 3H), 3.20 (q, 2H), 3.05 (m, 4H), 2.95 (t, 1H), 2.84 (dd, 1H), 2.76 (s, 2H), 2.73 (s, 3H), 2.25 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 1.04 (t, 3H), 0.94 (s, 6H).

EXAMPLE 334

(2S)-2-[[[(3-chloro-5-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]pyridin-2-yl)oxy]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide

EXAMPLE 334A

(S)-2-((3-chloro-5-sulfamoylpyridin-2-yloxy)methyl)-N-ethyl-N-methylmorpholine-4-carboxamide

[1258] The title compound was prepared by substituting EXAMPLE 244B for EXAMPLE 134A and *N*-methyl-*N*-ethyl carbamyl chloride for methyl iodide in EXAMPLE 134B.

EXAMPLE 334B

(2S)-2-[[[(3-chloro-5-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]pyridin-2-yl)oxy]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide

[1259] The title compound was prepared by substituting EXAMPLE 334A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 12.98 (s, 1H), 9.08 (d, 1H), 8.70 (d, 1H), 8.42 (d, 1H), 8.11 (d, 1H), 7.67 (t, 1H), 7.64 (d, 1H), 7.44 (d, 2H), 7.07 (d, 2H), 6.75 (dd, 1H), 6.53 (d, 1H), 6.48 (m, 1H), 4.58 (dd, 1H), 4.47 (dd, 1H), 4.03 (m, 1H), 3.84 (m, 2H), 3.63 (dt, 1H), 3.45 (d, 1H), 3.22 (q, 2H), 3.07 (m, 4H), 3.05-2.95 (m, 2H), 2.78 (s, 3H), 2.77 (s, 2H), 2.26 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 1.07 (t, 3H), 0.94 (s, 6H).

EXAMPLE 335

N-[[[5-chloro-6-[[[4-(4-cyclopropylmorpholin-2-yl)methyl]amino]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 335A

tert-butyl 2-((3-chloro-5-sulfamoylpyridin-2-ylamino)methyl)morpholine-4-carboxylate

[1260] The title compound was prepared by substituting EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide and tert-butyl 2-(aminomethyl)morpholine-4-carboxylate for (tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 335B

5-chloro-6-(morpholin-2-ylmethylamino)pyridine-3-sulfonamide

5 **[1261]** The title compound was prepared by substituting EXAMPLE 335A for EXAMPLE 113A in EXAMPLE 134A.

EXAMPLE 335C

5-chloro-6-((4-cyclopropylmorpholin-2-yl)methylamino)pyridine-3-sulfonamide

10

[1262] The title compound was prepared by substituting EXAMPLE 335B for EXAMPLE 173A in EXAMPLE 173B.

EXAMPLE 335D

15 N-[[[5-chloro-6-[[[4-(4-cyclopropylmorpholin-2-yl)methyl]amino]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

20 **[1263]** The title compound was prepared by substituting EXAMPLE 335C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.02 (s, 1H), 9.15 (d, 1H), 8.49 (d, 1H), 8.43 (d, 1H), 8.11 (d, 1H), 7.80 (t, 1H), 7.69 (d, 1H), 7.65 (t, 1H), 7.44 (d, 2H), 7.07 (d, 2H), 6.73 (dd, 1H), 6.52 (m, 1H), 6.49 (d, 1H), 3.92 (m, 1H), 3.84 (m, 2H), 3.70 (m, 1H), 3.54 (dt, 1H), 3.05 (m, 4H), 2.99 (d, 1H), 2.76 (s, 2H), 2.68 (d, 1H), 2.32 (dt, 1H), 2.25 (m, 2H), 2.12 (m, 5H), 1.97 (s, 2H), 1.53 (m, 1H), 1.39 (t, 2H), 0.93 (s, 6H), 0.40 (m, 4H).

EXAMPLE 336

25

2-[[[3-chloro-5-[[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]pyridin-2-yl)amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide

EXAMPLE 336A

30

2-((3-chloro-5-sulfamoylpyridin-2-ylamino)methyl)-N-ethyl-N-methylmorpholine-4-carboxamide

[1264] The title compound was prepared by substituting EXAMPLE 335B for EXAMPLE 134A and *N*-methyl-*N*-ethyl carbamyl chloride for methyl iodide in EXAMPLE 134B.

35

EXAMPLE 336B

2-[[[3-chloro-5-[[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]pyridin-2-yl)amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide

40

45 **[1265]** The title compound was prepared by substituting EXAMPLE 336A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.03 (s, 1H), 9.14 (d, 1H), 8.51 (d, 1H), 8.43 (d, 1H), 8.11 (d, 1H), 7.89 (m, 1H), 7.69 (d, 1H), 7.66 (t, 1H), 7.44 (d, 2H), 7.07 (d, 2H), 6.74 (dd, 1H), 6.51 (m, 1H), 6.48 (d, 1H), 3.96 (m, 1H), 3.90-3.70 (m, 4H), 3.59 (dt, 1H), 3.43 (d, 1H), 3.17 (q, 2H), 3.05 (m, 4H), 2.95 (dt, 1H), 2.81 (dd, 1H), 2.76 (s, 2H), 2.72 (s, 3H), 2.25 (m, 2H), 2.13 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 1.03 (t, 3H), 0.93 (s, 6H).

EXAMPLE 337

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50

EXAMPLE 337A

methyl 4,4-dimethyl-2-(trifluoromethylsulfonyloxy)cyclohex-1-enecarboxylate

55

[1266] To a suspension of hexane washed NaH (17 g) in dichloromethane (700 mL) was added 5,5-dimethyl-2-methoxycarbonylcyclohexanone (38.5 g) dropwise at 0°C. After stirring for 30 minutes, the mixture was cooled to -78°C and trifluoroacetic anhydride (40 mL) was added. The reaction mixture was warmed to room temperature and stirred for

24 hours. The organic layer was washed with brine, dried (Na_2SO_4), filtered, and concentrated to give the product.

EXAMPLE 337B

5 methyl 2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enecarboxylate

[1267] EXAMPLE 337A (62.15g), 4-chlorophenylboronic acid (32.24 g), CsF (64 g) and tetrakis(triphenylphosphine)palladium(0) (2g) in 2:1 dimethoxyethane /methanol (600 mL) were heated to 70°C for 24 hours. The mixture was concentrated. Ether (4x 200 mL) was added and the mixture was filtered. The combined ether solution was concentrated to give the product.

EXAMPLE 337C

15 (2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methanol

[1268] To a mixture of LiBH_4 (13g), EXAMPLE 337B (53.8 g) and ether (400 mL), was added methanol (25 mL) slowly by syringe. The mixture was stirred at room temperature for 24 hours. The reaction was quenched with 1N HCl with ice-cooling. The mixture was diluted with water and extracted with ether (3x 100 mL). The extracts were dried (Na_2SO_4), filtered, and concentrated. The crude product was chromatographed on silica gel with 0-30% ethyl acetate/hexanes.

EXAMPLE 337D

tert-butyl 4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazine-1-carboxylate

25 [1269] Mesyl Chloride (7.5 mL) was added via syringe to EXAMPLE 337C (29.3 g) and triethylamine (30 mL) in CH_2Cl_2 (500 mL) at 0°C, and the mixture was stirred for 1 minute. N-butoxycarbonylpiperazine (25 g) was added and the mixture was stirred at room temperature for 24 hours. The suspension was washed with brine, dried, (Na_2SO_4), filtered, and concentrated. The crude product was chromatographed on silica gel with 10-20% ethyl acetate/hexanes.

EXAMPLE 337E

1-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazine

35 [1270] EXAMPLE 337D (1 g) was stirred in dichloromethane (10 mL), trifluoroacetic acid (10 mL), and triethylsilane (1 mL) for 1 hour. The mixture was concentrated, taken up in a mixture of dichloromethane (100 mL) and saturated aqueous Na_2CO_3 solution (20 mL) and stirred for 10 minutes. The layers were separated, and the organic layer was dried over Na_2SO_4 , and concentrated to give the product.

EXAMPLE 337F

40

5-bromo-1-(triisopropylsilyl)-1H-pyrrolo[2,3-b]pyridine

[1271] To a mixture of 5-bromo-1H-pyrrolo[2,3-b]pyridine (15.4 g) in tetrahydrofuran (250 mL) was added 1M lithium hexamethyldisilazide in tetrahydrofuran (86 mL), and after 10 minutes, TIPS-Cl (triisopropylchlorosilane) (18.2 mL) was added. The mixture was stirred at room temperature for 24 hours. The reaction was diluted with ether, and the resulting solution was washed twice with water. The extracts were dried (Na_2SO_4), filtered, and concentrated. The crude product was chromatographed on silica gel with 10% ethyl acetate/hexanes.

EXAMPLE 337G

50

1-(triisopropylsilyl)-1H-pyrrolo[2,3-b]pyridin-5-ol

[1272] To a mixture of EXAMPLE 337F (24.3 g) in tetrahydrofuran (500 mL) at -78°C was added 2.5M BuLi (30.3 mL). After 2 minutes, trimethylborate (11.5 mL) was added, and the mixture was allowed to warm to room temperature over 1 hour. The reaction was poured into water, extracted three times with ethyl acetate, and the combined extracts were washed with brine and concentrated. The crude product was taken up in tetrahydrofuran (200 mL) at 0°C, and 1M aqueous NaOH (69 mL) was added, followed by 30% aqueous H_2O_2 (8.43 mL), and the solution was stirred for 1 hour. $\text{Na}_2\text{S}_2\text{O}_3$ (10 g) was added, and the pH was adjusted to 4-5 with concentrated HCl and solid NaH_2PO_4 . The solution

was extracted twice with ethyl acetate, and the combined extracts were washed with brine, dried (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 5-25% ethyl acetate/hexanes.

EXAMPLE 337H

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-fluorobenzoate

[1273] A mixture of EXAMPLE 337G (8.5 g), methyl 2,4-difluorobenzoate (7.05 g), and K₃PO₄ (9.32 g) in diglyme (40 mL) at 115°C was stirred for 24 hours. The reaction was cooled, diluted with ether (600 mL), and washed twice with water, and brine, and concentrated. The crude product was chromatographed on silica gel with 2-50% ethyl acetate/hexanes.

EXAMPLE 337I

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoate

[1274] A mixture of EXAMPLE 337H (1.55 g), EXAMPLE 337E (2.42 g), and HK₂PO₄ (1.42 g) in dimethylsulfoxide (20 mL) at 135°C was stirred for 24 hours. The reaction was cooled, diluted with ether (400 mL), and washed three times with 1M aqueous NaOH, and brine, and concentrated. The crude product was chromatographed on silica gel with 10-50% ethyl acetate/hexanes.

EXAMPLE 337J

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoic acid

[1275] EXAMPLE 337I (200 mg) in dioxane (10 mL) and 1M aqueous NaOH (6 mL) at 50°C was stirred for 24 hours. The reaction was cooled, added to NaH₂PO₄ solution, and extracted three times with ethyl acetate. The combined extracts were washed with brine, and concentrated to give the pure product.

EXAMPLE 337K

tert-butyl (4-hydroxy-4-methylcyclohexyl)methylcarbamate

[1276] To a vigorous stirring solution of tert-butyl (4-oxocyclohexyl)methylcarbamate (1.7 g) in tetrahydrofuran (40 mL) at -78°C was dropwise added 1.6 M methyllithium (14.02 mL) in ether. After completion of the addition, the mixture was stirred at -78°C for 1.2 hours and poured into a cold NH₄Cl aqueous solution. The resulting mixture was extracted with dichloromethane (100 mL, three times) and the organic layer was dried over Na₂SO₄, filtered, and concentrated. The residue was dissolved in dichloromethane and loaded onto an Analogix purification system, and it was eluted with 0 - 50% ethyl acetate in dichloromethane to provide the title compound.

EXAMPLE 337L

4-(aminomethyl)-1-methylcyclohexanol

[1277] EXAMPLE 337K (1.3 g) in dichloromethane (5 mL) at 0°C was treated with trifluoroacetic acid (2.1 mL) and a few drops of water for 1 hour. The reaction mixture was concentrated and the residue was directly used for next step.

EXAMPLE 337M

4-((trans-4-hydroxy-4-methylcyclohexyl)methylamino)-3-nitrobenzenesulfonamide

[1278] EXAMPLE 337L (732 mg) and 4-fluoro-3-nitrobenzenesulfonamide (1.1 g) in tetrahydrofuran (15 mL) was treated with triethylamine overnight. The reaction mixture was concentrated and the residue was purified by a reverse phase chromatography, eluting with 30% - 50% acetonitrile in 0.1% trifluoroacetic acid water solution to isolate the title compound.

EXAMPLE 337N

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1279] A mixture of EXAMPLE 337J (3.0 g), EXAMPLE 337M (1.98 g), N, N-dimethylpyridin-4-amine (1.93 g) and N¹-((ethylimino)methylene)-N³,N³-dimethylpropane-1,3-diamine hydrochloride (1.31 g) in dichloromethane (50 ml) was stirred overnight and concentrated. The residue was purified by reverse chromatography, eluted with 40%-70% acetonitrile in 0.1% TFA water. The desired fractions were concentrated to remove acetonitrile, neutralized with NaHCO₃ and extracted with dichloromethane. The organic layer was dried over Na₂SO₄, concentrated and dried to provide the title compound. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1 H), 8.52 - 8.58 (m, 2 H), 8.04 (d, 1 H), 7.79 (dd, 1 H), 7.53 (d, 1 H), 7.47 - 7.52 (m, 2 H), 7.30 - 7.37 (m, 2 H), 7.07 (d, 1 H), 7.01 - 7.06 (m, 2 H), 6.68 (dd, 1 H), 6.39 (dd, 1 H), 6.19 (d, 1 H), 4.25 (s, 1 H), 3.25 - 3.32 (m, 4 H), 3.07 (s, 4 H), 2.75 (s, 2 H), 2.09 - 2.24 (m, 6 H), 1.95 (s, 2 H), 1.50 - 1.73 (m, 5 H), 1.28 - 1.43 (m, 4 H), 1.06 - 1.18 (m, 5 H), 0.92 (s, 6 H).

EXAMPLE 338

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(cis-4-hydroxy-4-methylcyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 338A

methyl 4,4-dimethyl-2-(trifluoromethylsulfonyloxy)cyclohex-1-enecarboxylate

[1280] To a suspension of hexane washed NaH (17 g) in dichloromethane (700 mL) was added 5,5-dimethyl-2-methoxycarbonylcyclohexanone (38.5 g) dropwise at 0°C. After stirring for 30 minutes, the mixture was cooled to -78°C and trifluoroacetic anhydride (40 mL) was added. The reaction mixture was warmed to room temperature and stirred for 24 hours. The organic layer was washed with brine, dried (Na₂SO₄), filtered, and concentrated to give the product.

EXAMPLE 338B

methyl 2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enecarboxylate

[1281] EXAMPLE 338A (62.15g), 4-chlorophenylboronic acid (32.24 g), CsF (64 g) and tetrakis(triphenylphosphine)palladium(0) (2g) in 2:1 dimethoxyethane /methanol (600 mL) were heated to 70°C for 24 hours. The mixture was concentrated. Ether (4x 200 mL) was added and the mixture was filtered. The combined ether solution was concentrated to give the product.

EXAMPLE 338C

(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methanol

[1282] To a mixture of LiBH₄ (13g), EXAMPLE 338B (53.8 g) and ether (400 mL), was added methanol (25 mL) slowly by syringe. The mixture was stirred at room temperature for 24 hours. The reaction was quenched with 1N aqueous HCl with ice-cooling. The mixture was diluted with water and extracted with ether (3x 100 mL). The extracts were dried (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 0-30% ethyl acetate/hexanes.

EXAMPLE 338D

tert-butyl 4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazine-1-carboxylate

[1283] Mesyl Chloride (7.5 mL) was added via syringe to EXAMPLE 338C (29.3 g) and triethylamine (30 mL) in CH₂Cl₂ (500 mL) at 0°C, and the mixture was stirred for 1 minute. N-t-butoxycarbonylpiperazine (25 g) was added and the mixture was stirred at room temperature for 24 hours. The suspension was washed with brine, dried, (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 10-20% ethyl acetate/hexanes.

EXAMPLE 338E

1-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazine

- 5 **[1284]** EXAMPLE 338D (1 g) was stirred in dichloromethane (10 mL), trifluoroacetic acid (10 mL), and triethylsilane (1 mL) for 1 hour. The mixture was concentrated, taken up in a mixture of dichloromethane (100 mL) and saturated aqueous Na₂CO₃ solution (20 mL) and stirred for 10 minutes. The layers were separated, and the organic layer was dried over Na₂SO₄, filtered, and concentrated to give the product.

10 EXAMPLE 338F

5-bromo-1-(triisopropylsilyl)-1H-pyrrolo[2,3-b]pyridine

- 15 **[1285]** To a mixture of 5-bromo-1H-pyrrolo[2,3-b]pyridine (15.4 g) in tetrahydrofuran (250 mL) was added 1M lithium hexamethyldisilazide in tetrahydrofuran (86 mL), and after 10 minutes, TIPS-Cl (triisopropylchlorosilane) (18.2 mL) was added. The mixture was stirred at room temperature for 24 hours. The reaction was diluted with ether, and the resulting solution was washed twice with water. The extracts were dried (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 10% ethyl acetate/hexanes.

20 EXAMPLE 338G

1-(triisopropylsilyl)-1H-pyrrolo[2,3-b]pyridin-5-ol

- 25 **[1286]** To a mixture of EXAMPLE 338F (24.3 g) in tetrahydrofuran (500 mL) at -78°C was added 2.5M BuLi (30.3 mL). After 2 minutes, trimethylborate (11.5 mL) was added, and the mixture was allowed to warm to room temperature over 1 hour. The reaction was poured into water, extracted three times with ethyl acetate, and the combined extracts were washed with brine and concentrated. The crude product was taken up in tetrahydrofuran (200 mL) at 0°C, and 1M aqueous NaOH (69 mL) was added, followed by 30% aqueous H₂O₂ (8.43 mL), and the solution was stirred for 1 hour. Na₂S₂O₃ (10 g) was added, and the pH was adjusted to 4-5 with concentrated HCl and solid NaH₂PO₄. The solution
30 was extracted twice with ethyl acetate, and the combined extracts were washed with brine, dried (Na₂SO₄), filtered, and concentrated. The crude product was chromatographed on silica gel with 5-25% ethyl acetate/hexanes.

EXAMPLE 338H

- 35 methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-fluorobenzoate

- [1287]** A mixture of EXAMPLE 338G (8.5 g), methyl 2,4-difluorobenzoate (7.05 g), and K₃PO₄ (9.32 g) in diglyme (40 mL) at 115°C was stirred for 24 hours. The reaction was cooled, diluted with ether (600 mL), and washed twice with water, and brine, and concentrated. The crude product was chromatographed on silica gel with 2-50% ethyl acetate/hexanes.
40

EXAMPLE 338I

- 45 methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoate

- [1288]** A mixture of EXAMPLE 338H (1.55 g), EXAMPLE 338E (2.42 g), and HK₂PO₄ (1.42 g) in dimethylsulfoxide (20 mL) at 135°C was stirred for 24 hours. The reaction was cooled, diluted with ether (400 mL), and washed three times with 1M aqueous NaOH, and brine, and concentrated. The crude product was chromatographed on silica gel with 10-50% ethyl acetate/hexanes.
50

EXAMPLE 338J

- 55 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoic acid

- [1289]** EXAMPLE 338I (200 mg) in dioxane (10 mL) and 1M NaOH (6 mL) at 50°C was stirred for 24 hours. The reaction was cooled, added to NaH₂PO₄ solution, and extracted three times with ethyl acetate. The combined extracts

were washed with brine, and concentrated to give the pure product.

EXAMPLE 338K

5 tert-butyl (4-hydroxy-4-methylcyclohexyl)methylcarbamate

[1290] To a vigorous stirring solution of tert-butyl (4-oxocyclohexyl)methylcarbamate (1.7 g) in tetrahydrofuran (40 mL) at -78°C was dropwise added 1.6 M methyllithium (14.02 mL) in ether. After completion of the addition, the mixture was stirred at -78°C for 1.2 hours and poured into a cold NH₄Cl aqueous solution. The resulting mixture was extracted with dichloromethane (100 ml, three times) and the organic layer was dried over Na₂SO₄, filtered, and concentrated. The residue was dissolved in dichloromethane and loaded onto an Analogix purification system, and it was eluted with 0 - 50% ethyl acetate in dichloromethane to provide the title compound.

EXAMPLE 338L

15 4-(aminomethyl)-1-methylcyclohexanol

[1291] EXAMPLE 338K (1.3 g) in dichloromethane (5 mL) at 0°C was treated with trifluoroacetic acid (2.1 mL) and a few drops of water for 1 hour. The reaction mixture was concentrated and the residue was directly used for next step.

EXAMPLE 338M

4-((cis-4-hydroxy-4-methylcyclohexyl)methylamino)-3-nitrobenzenesulfonamide

25 **[1292]** EXAMPLE 338L (732 mg) and 4-fluoro-3-nitrobenzenesulfonamide (1.1 g) in tetrahydrofuran (15 mL) was treated with triethylamine overnight. The reaction mixture, was concentrated and the residue was purified by a reverse phase chromatography, eluting with 30% - 50% acetonitrile in 0.1% trifluoroacetic acid water solution to isolate the title compound.

EXAMPLE 338N

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((cis-4-hydroxy-4-methylcyclohexyl)methyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

35 **[1293]** A mixture of EXAMPLE 338J (144 mg), EXAMPLE 338M (95 mg), N, N-dimethylpyridin-4-amine (123 mg) and N¹-((ethylimino)methylene)-N³,N³-dimethylpropane-1,3-diamine hydrochloride (62.7 mg) in dichloromethane (7 ml) was stirred overnight and concentrated. The residue was purified by reverse chromatography, eluted with 40%-70% acetonitrile in 0.1% TFA water. The desired fractions were concentrated, neutralized with NaHCO₃ and extracted with dichloromethane. The organic layer was dried over Na₂SO₄, filtered, concentrated and dried to provide the title compound.

40 ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1 H), 11.38 (s, 1 H), 8.59 (t, 1 H), 8.55 (d, 1 H), 8.04 (d, 1 H), 7.79 (dd, 1 H), 7.54 (d, 1 H), 7.46 - 7.52 (m, 2 H), 7.30 - 7.38 (m, 2 H), 7.00 - 7.10 (m, 3 H), 6.68 (dd, 1 H), 6.39 (dd, 1 H), 6.19 (d, 1 H), 3.95 (s, 1 H), 3.25 (t, 4 H), 3.07 (s, 4 H), 2.75 (s, 2 H), 2.10 - 2.26 (m, 6 H), 1.95 (s, 2 H), 1.29 - 1.62 (m, 8 H), 1.16 - 1.30 (m, 2 H), 1.08 (s, 3 H), 0.92 (s, 6 H).

EXAMPLE 339

N-([5-chloro-6-([1R,2R,4R,5R]-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 339A

(1R,4S)-methyl spiro[bicyclo[2.2.1]heptane-2,2'-[1,3]dioxolane]-5-carboxylate

55 **[1294]** A reaction mixture of 1,4-dioxaspiro[4.4]non-6-ene (5 g), methyl acrylate (10.24 g), and hydroquinone (0.13 g) was heated at 100°C in acetonitrile (12 mL) for three days. After cooling, the solvent was removed, and residue was purified by flash chromatography on silica gel eluting with 4:1 hexane/ethyl acetate to provide the title compound as a mixture of two isomers.

EXAMPLE 339B

(1R,4S)-spiro[bicyclo[2.2.1]heptane-2,2'-[1,3]dioxolane]-5-ylmethanol

- 5 **[1295]** EXAMPLE 339A (1.0 g) in tetrahydrofuran was cooled to 0°C. To this solution was added 1.0 N lithium aluminum hydride (2.8 mL) dropwise. The reaction mixture was stirred for 2 hours. Water (0.4 mL) was added followed by 2 N aqueous NaOH (0.2 mL). The solid was filtered off, and the filtrate was concentrated. Toluene was added, and it was then distilled to remove any trace amount of water. The title compound was used for the next reaction without further purification.

10

EXAMPLE 339C

5-chloro-6-(((1S,2R,4R)-5-oxobicyclo[2.2.1]heptan-2-yl)methoxy)pyridine-3-sulfonamide

- 15 **[1296]** The title compound was prepared by substituting EXAMPLE 339B for (tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 40A for EXAMPLE 36A in EXAMPLE 36B. The two stereoisomers at the 5 position were isolated by reverse phase Gilson Prep HPLC system with a Phenomenex prep column (Luna, 5 μ , C18(2), 250X21.20 mm, 5 Å) eluting with 20-80% acetonitrile in water with 0.1% trifluoroacetic acid. The desired fractions were collected, and the solvents were removed under reduced vacuum at 60°C. During this process, a lot of solid formed. It was then partitioned between water and ethyl acetate. The organic layer was separated, and the aqueous layer was extracted with additional ethyl acetate. The combined organic layers were washed with brine, dried over MgSO₄, filtered and concentrated to give the title compound.
- 20

EXAMPLE 339D

25

5-chloro-6-(((1S,2R,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]heptan-2-yl)methoxy)pyridine-3-sulfonamide

- [1297]** EXAMPLE 339C (0.44 g) in tetrahydrofuran (15 mL) was treated with 3.0 M methylmagnesium bromide (5.3 mL) at 0 °C. The solution was stirred for 16 hours. The reaction mixture was then partitioned between ethyl acetate and 0.05 N aqueous HCl (20 mL). The organic layer was separated, and the aqueous layer was extracted with additional ethyl acetate three times. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified by flash column chromatography on silica gel using 10-50% ethyl acetate in hexanes to provide the title compound.
- 30

EXAMPLE 339E

35

N-[(5-chloro-6-(((1R,2R,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl)methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- 40 **[1298]** The title compound was prepared by substituting EXAMPLE 339D for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1H), 8.50 (s, 1H), 8.16 (s, 1H), 8.02 (d, 1H), 7.49-7.55 (m, 3H), 7.35 (d, 2H), 7.05 (d, 2H), 6.67 (dd, 1H), 6.38 (dd, 1H), 6.20 (s, 1H), 4.40-4.48 (m, 2H), 4.31 (s, 1H), 3.09 (s, 4H), 2.83 (s, 2H), 2.15-2.33 (m, 7H), 1.96 (s, 2H), 1.87 (d, 1H), 1.65-1.69 (m, 1H), 1.54-1.56 (m, 2H), 1.36-1.47 (m, 6H), 1.26-1.30 (m, 1H), 1.19 (s, 3H), 0.93 (s, 6H).

45

EXAMPLE 340

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([2-(2-cyanoethyl)(cyclopropyl)amino]cyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

50

EXAMPLE 340A

4-(1,4-dioxaspiro[4.5]decan-8-ylamino)-3-nitrobenzenesulfonamide

- 55 **[1299]** To a solution of 4-fluoro-3-nitrobenzenesulfonamide (1.4 g) in tetrahydrofuran (30 mL) was added 1,4-dioxaspiro[4.5]decan-8-amine (1.0 g) and diisopropylethylamine (5 mL). The mixture was stirred overnight. The mixture was diluted with ethyl acetate (300 mL) and washed with water, brine and dried over Na₂SO₄. Filtration and evaporation of the solvent gave the title compound.

EXAMPLE 340B

N-(4-(1,4-dioxaspiro[4.5]decan-8-ylamino)-3-nitrophenylsulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzamide

[1300] To a solution of EXAMPLE 3J (617 mg) and EXAMPLE 340A (386 mg) in dichloromethane (10 mL) was added 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (288 mg) and 4-(dimethylamino)pyridine (183 mg). The mixture was stirred overnight. The mixture was diluted with dichloromethane (300 mL) and washed with aqueous NaHCO₃, water, brine and dried over Na₂SO₄. Filtration and evaporation of the solvent gave the title compound.

EXAMPLE 340C

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)-N-(3-nitro-4-(4-oxocyclohexylamino)phenylsulfonyl)benzamide

[1301] To a solution of EXAMPLE 340B (386 mg) in acetone (10 mL) and water (5 mL) was added para-toluenesulfonic acid monohydrate (50 mg). The mixture was stirred at 120°C in a Biotage Initiator microwave reactor for 30 minutes. The mixture was diluted with dichloromethane (300 mL) and washed with aqueous NaHCO₃, water, brine and dried over Na₂SO₄. Filtration and evaporation of the solvent gave the title compound.

EXAMPLE 340D

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((2-cyanoethyl)(cyclopropyl)amino)cyclohexyl]amino)-3-nitrophenylsulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1302] To a solution of EXAMPLE 340C (240 mg) and 3-(cyclopropylamino)propanenitrile (62 mg) in tetrahydrofuran (10 mL) was added acetic acid (2 mL) and MP-cyanoborohydride (300 mg, 2.15mmol/g). The mixture was stirred overnight. The mixture was filtered and concentrated under vacuum and the residue was dissolved in dimethylsulfoxide/methanol (1:1, 10 mL) and loaded on Gilson, C18(100A) 250x121.2 mm(10 micron), with 30% acetonitrile to 65% acetonitrile over 40 minutes. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1 H), 8.55 (dd, 1 H), 8.17 (d, 1 H), 8.03 (d, 1 H), 7.79 (d, 1 H), 7.49 (m, 3 H), 7.34 (d, 2 H), 7.11 (m, 1 H), 7.04 (d, 2 H), 6.67 (dd, 1 H), 6.38 (d, 1 H), 6.19 (d, 1 H), 4.01 (m, 1 H), 3.56 (m, 1 H), 3.06 (m, 4 H), 2.88 (t, 2 H), 2.65 (m, 6 H), 2.19 (m, 6 H), 2.00 (m, 7 H), 1.51 (m, 6 H), 0.92 (s, 6 H), 0.42 (m, 4 H).

EXAMPLE 341

N-([5-chloro-6-((trans-4-hydroxy-4-methylcyclohexyl)methoxy)pyridin-3-yl]sulfonyl)-4-(4-([5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 341A

ethyl 1,4-dioxaspiro[4.5]decan-8-carboxylate

[1303] To a solution of ethyl 4-oxocyclohexanecarboxylate (31.8 g) in toluene (100 mL) was added ethylene glycol (36.5 mL) and p-toluenesulfonic acid monohydrate (0.426 g). The two phase mixture was stirred rapidly at ambient temperature for 72 hours. The reaction was diluted with water (900 mL) and extracted with ether (900 mL). The organic layer was washed with saturated sodium bicarbonate solution and brine, and then dried over anhydrous sodium sulfate. After filtration, the title compound was obtained by concentration under high vacuum.

EXAMPLE 341B

1,4-dioxaspiro[4.5]decan-8-ylmethanol

[1304] To a suspension of lithium aluminum hydride (8.19 g) in tetrahydrofuran (400 mL) was added dropwise a solution of EXAMPLE 341A (37.8 g) in tetrahydrofuran (75 mL). The mixture was then heated at reflux for 2 hours. The reaction mixture was cooled in an ice bath and quenched very slowly with water (8 mL). Then added sequentially were 4N sodium hydroxide (8 mL), ether (200 mL), water (24 mL), ether (500 mL) and anhydrous sodium sulfate (250 g). The resulting mixture was stirred rapidly for 2 hours and was filtered. The title compound was isolated by concentration of the filtrate.

EXAMPLE 341C

8-(benzyloxymethyl)-1,4-dioxaspiro[4.5]decane

5 **[1305]** To a suspension of sodium hydride (60% oil dispersion, 8.86 g) in tetrahydrofuran (170 mL) was added a solution of EXAMPLE 341B (30.52 g) in tetrahydrofuran (100 mL). This mixture was stirred for 30 minutes and benzyl bromide (24 mL) was added. After stirring for 72 hours, the reaction was quenched with saturated ammonium chloride solution (400 mL) and diluted with ether (500 mL). The layers were separated and the aqueous layer was extracted with ether (2 X 150 mL). The combined organics were dried over sodium sulfate, filtered and concentrated. The crude product
10 was purified on silica gel eluting with a 0, 10, 15, 75 % ethyl acetate in hexanes step gradient to provide the title compound.

EXAMPLE 341D

4-(benzyloxymethyl)cyclohexanone

15 **[1306]** To a solution of EXAMPLE 341C (43.02 g) in dioxane (500 mL) was added water (125 mL) and 2M hydrochloric acid (90 mL). The mixture was heated at 85°C for 18 hours. Upon cooling, the reaction mixture was diluted with brine (1500 mL), saturated sodium bicarbonate solution (300 mL) and ether (1000 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. The crude product was purified on silica gel eluting with a 5- 50% ethyl acetate
20 in hexanes step gradient to provide the title compound.

EXAMPLE 341E

trans-4-(benzyloxymethyl)-1-methylcyclohexanol

25 **[1307]** To 2,6-di-*t*-butyl-4-methylphenol (83.4 g) in toluene (1100 mL) was added 2.0M (in hexanes) trimethylaluminum (95 mL) somewhat carefully to control methane evolution and a small exotherm. The reaction mixture was stirred at ambient temperature under N₂ for 75 minutes and was then cooled to -77° C. A solution of EXAMPLE 341D (14 g) in toluene (15 mL) was added dropwise, keeping the temperature below -74 °C. Methylolithium (1.6M in diethyl ether, 120
30 mL) was then added dropwise, keeping the temperature below -65 °C. The resulting mixture was stirred at -77 °C under N₂ for 2 hours. The reaction mixture was then poured into 1N aqueous HCl (1600 mL), rinsing the flask with toluene. The organic layer was washed with brine and the combined aqueous layers were extracted with diethyl ether. The combined organic layers were dried (Na₂SO₄), filtered and concentrated. The concentrate was chromatographed on 650 g of spherical silica gel using 2.5 L of 80/20 hexanes/ethyl acetate, then 3.0 L of 75/25 hexanes/ethyl acetate, and
35 finally 4.0 L of 70/30 hexanes/ethyl acetate as the eluents to provide the title compound.

EXAMPLE 341F

Trans-4-(hydroxymethyl)-1-methylcyclohexanol

40 **[1308]** EXAMPLE 341E (12.6 g) and ethanol (120 mL) were added to 20% Pd(OH)₂/C, wet (1.260 g) in a 500 mL SS pressure bottle. The reaction mixture was stirred at ambient temperature under 30 psi hydrogen gas. Hydrogen uptake ceased at 5 minutes. The mixture was filtered through a nylon membrane rinsing with ethanol. The filtrate was concentrated and then azeotroped with toluene (100 mL) to remove any remaining ethanol. The concentrate was dried under
45 high vacuum for 40 minutes to provide the title compound.

EXAMPLE 341G

5-chloro-6-((trans-4-hydroxy-4-methylcyclohexyl)methoxy)pyridine-3-sulfonamide

50 **[1309]** The title compound was prepared by substituting EXAMPLE 40A for 4-fluoro-3-nitrobenzenesulfonamide and EXAMPLE 341F for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

55

EXAMPLE 341H

N-({5-chloro-6-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-{4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1310] The title compound was prepared by substituting EXAMPLE 328E for EXAMPLE 3J and EXAMPLE 341G for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, pyridine-d₅) δ 13.09 (s, 1H), 9.18 (d, 1H), 8.74 (d, 1H), 8.41 (d, 1H), 8.09 (d, 1H), 7.67 (m, 2H), 7.42 (m, 2H), 7.09 (m, 2H), 6.74 (dd, 1H), 6.52 (dd, 1H), 6.49 (d, 1H), 4.29 (d, 2H), 3.05 (m, 4H), 2.80 (s, 2H), 2.37 (t, 2H), 2.15 (m, 4H), 2.11 (s, 2H), 1.89 (m, 6H), 1.75 (m, 2H), 1.45 (t, 2H), 1.41 (s, 3H), 1.32 (m, 2H), 0.37 (m, 4H).

EXAMPLE 342

4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-[[5-chloro-6-(5,6,7,8-tetrahydroimidazo[1,2-a]pyridin-6-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 342A

methyl 5,6,7,8-tetrahydroimidazo[1,2-a]pyridine-6-carboxylate

[1311] To a 50 ml pressure bottle were placed methyl imidazo[1,2-a]pyridine-6-carboxylate (0.26 g), acetic acid (10 ml), and wet 5% palladium on carbon (0.052 g). The reaction mixture was stirred for 16 hours at 30 psi and 50 °C. The solid was filtered off, and the filtrate was concentrated. The residue was taken up in ethyl acetate. It was then washed with saturated sodium bicarbonate, brine, dried over MgSO₄, filtered, and concentrated. The residue was purified by flash column chromatography on silica gel using 10-100% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 342B

(5,6,7,8-tetrahydroimidazo[1,2-a]pyridin-6-yl)methanol

[1312] The title compound was prepared by substituting EXAMPLE 342A for EXAMPLE 339A in EXAMPLE 339B.

EXAMPLE 342C

5-chloro-6-((5,6,7,8-tetrahydroimidazo[1,2-a]pyridin-6-yl)methoxy)pyridine-3-sulfonamide The title compound was prepared by substituting EXAMPLE 342B for (tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 40A for EXAMPLE 36A in EXAMPLE 36B.

EXAMPLE 342D

4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-[[5-chloro-6-(5,6,7,8-tetrahydroimidazo[1,2-a]pyridin-6-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1313] The title compound was prepared by substituting EXAMPLE 342C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.54 (s, 1H), 8.36 (s, 1H), 8.06 (s, 1H), 7.93 (s, 1H), 7.58 (d, 1H), 7.41-7.44 (m, 2H), 7.2-7.36 (m, 4H), 7.05 (d, 2H), 6.63 (dd, 1H), 6.32 (dd, 1H), 6.24 (d, 1H), 4.42-4.51 (m, 1H), 4.37-4.40 (m, 1H), 4.29 (dd, 1H), 3.91 (dd, 1H), 3.03 (s, 4H), 2.90-2.95 (m, 2H), 2.77 (s, 2H), 2.51-2.52 (m, 1H), 2.07-2.23 (m, 7H), 1.96 (s, 2H), 1.76-1.82 (m, 1H), 1.65-1.69 (m, 2H), 1.54-1.56 (m, 2H), 1.39 (t, 2H), 0.93 (s, 6H).

EXAMPLE 343

N-[[5-chloro-6-[[[1R,2S,4R,5R]-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 343A

5-chloro-6-(((1S,2S,4R)-5-oxobicyclo[2.2.1]heptan-2-yl)methoxy)pyridine-3-sulfonamide The title compound was isolated as another isomer in EXAMPLE 339C.

EXAMPLE 343B

5-chloro-6-(((1S,2S,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]heptan-2-yl)methoxy)pyridine-3-sulfonamide

[1314] The title compound was prepared by substituting EXAMPLE 343A for EXAMPLE 339B in EXAMPLE 339C.

EXAMPLE 343C

N-([5-chloro-6-(((1R,2S,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl)methoxy)pyridin-3-yl)sulfonyl]-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1315] The title compound was prepared by substituting EXAMPLE 343B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1H), 8.51 (s, 1H), 8.17 (s, 1H), 8.03 (s, 1H), 7.49-7.55 (m, 3H), 7.35 (d, 2H), 7.05 (d, 2H), 6.67 (dd, 1H), 6.39 (dd, 1H), 6.20 (d, 1H), 4.27 (s, 1H), 4.11-4.19 (m, 2H), 3.11 (s, 4H), 2.87 (s, 2H), 1.96-2.23 (m, 10H), 1.88 (d, 1H), 1.50 (dd, 1H), 1.33-1.44 (m, 2H), 1.13-1.19 (m, 4H), 0.88-0.93 (m, 8H).

EXAMPLE 344

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([cis-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 344A

4-([cis-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrobenzenesulfonamide

[1316] EXAMPLE 347A (732 mg) and 4-fluoro-3-nitrobenzenesulfonamide (1.2 g) in tetrahydrofuran (40 mL) were treated with 60% sodium hydride (1.6 g) for 3 days. The reaction was quenched with water. The resulting mixture was neutralized with diluted aqueous HCl, and extracted with ethyl acetate. The organic layer was dried over Na₂SO₄, filtered, and concentrated. The residue was purified by a reverse phase chromatography, eluting with 30-50% CH₃CN in 0.1% trifluoroacetic acid water to provide the title compound as a single enantiomer.

EXAMPLE 344B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([cis-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1317] The title compound was prepared by substituting EXAMPLE 344A for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1H), 8.34 (d, 1H), 8.04 (m, 2H), 7.52 (m, 3H), 7.40 (d, 1H), 7.35 (d, 2H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.40 (m, 1H), 6.20 (d, 1H), 4.02 (d, 2H), 3.96 (s, 1H), 3.10 (br s, 4H), 2.85 (m, 2H), 2.29 (m, 3H), 2.15 (t, 2H), 1.96 (br s, 2H), 1.68 (m, 1H), 1.55 (m, 4H), 1.42 (m, 4H), 1.27 (m, 2H), 1.10 (s, 3H), 0.92 (s, 6H).

EXAMPLE 345

N-([5-chloro-6-([4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy)pyridin-3-yl)sulfonyl]-4-(4-([5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1318] The title compound was prepared by substituting EXAMPLE 328E for EXAMPLE 3J and EXAMPLE 2770 for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500 MHz, pyridine-d₅) δ 13.07 (s, 1H), 9.13 (d, 1H), 8.41 (d, 1H), 8.09 (d, 1H), 7.68 (t, 1H), 7.66 (d, 1H), 7.42 (m, 2H), 7.09 (m, 2H), 6.75 (dd, 1H), 6.51 (m, 2H), 4.64 (d, 4H), 4.53 (d, 2H), 3.39 (m, 1H), 3.06 (m, 4H), 2.81 (s, 2H), 2.51 (m, 2H), 2.37 (m, 2H), 2.12 (m, 10H), 1.90 (m, 2H), 1.45 (t, 2H), 0.38 (s, 4H).

EXAMPLE 346

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(3,3-difluoropyrrolidin-1-yl)cyclohexyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

5

[1319] The title compound was prepared by substituting 3,3-difluoropyrrolidine hydrochloride for 3-(cyclopropylamino)propanenitrile in EXAMPLE 340D. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1 H), 11.38 (m, 1 H), 8.55 (m, 1 H), 8.36 (d, 1 H), 8.03 (d, 1 H), 7.80 (m, 1 H), 7.50 (m, 3 H), 7.34 (d, 2 H), 7.13 (d, 1 H), 7.04 (d, 2 H), 6.83 (m, 1 H), 6.68 (m, 1 H), 6.38 (d, 1 H), 6.19 (s, 1 H), 4.02 (s, 1 H), 3.83 (m, 1 H), 3.06 (m, 4 H), 2.96 (m, 2 H), 2.73 (m, 4 H), 2.26 (m, 8 H), 1.97 (m, 4 H), 1.68 (m, 4 H), 1.37 (m, 2 H), 0.92 (s, 6 H).

10

EXAMPLE 347

N-([5-chloro-6-((trans-4-hydroxy-4-methylcyclohexyl)methoxy)pyridin-3-yl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

15

EXAMPLE 347A

4-(hydroxymethyl)-1-methylcyclohexanol

20

[1320] 4-(Hydroxymethyl)cyclohexanone (800 mg) in tetrahydrofuran (15 mL) was treated with 3 M methylmagnesium chloride in tetrahydrofuran (6.24 mL) at 0°C. The reaction was warmed to room temperature over 2 hours and quenched with methanol and water. The resulting mixture was concentrated and the residue was suspended in ethyl acetate. The precipitates were filtered off and the filtrate was concentrated. The residue was purified by chromatography, eluting with 0-100% ethyl acetate in hexane to provide the title compound.

25

EXAMPLE 347B

5-chloro-6-((trans-4-hydroxy-4-methylcyclohexyl)methoxy)pyridine-3-sulfonamide

30

[1321] EXAMPLE 347A (970 mg) and EXAMPLE 40A (1.6 g) in N,N-dimethylformamide (8 mL) were treated with sodium hydride (1.8 g, 60%) at room temperature for 2 days. The reaction was quenched with water. The resulting mixture was neutralized with diluted aqueous HCl, and extracted with ethyl acetate. The organic layer was dried over Na₂SO₄, filtered, and concentrated. The residue was purified by a reverse phase chromatography, eluting with 30-45% acetonitrile in 0.1% trifluoroacetic acid water to isolate the title compound.

35

EXAMPLE 347C

5-chloro-6-((cis-4-hydroxy-4-methylcyclohexyl)methoxy)pyridine-3-sulfonamide The title compound was prepared and isolated as described in Example 347B.

40

EXAMPLE 347D

N-([5-chloro-6-((trans-4-hydroxy-4-methylcyclohexyl)methoxy)pyridin-3-yl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

45

[1322] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 347B in place of EXAMPLE 11B. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1 H), 8.51 (d, 1 H), 8.18 (d, 1 H), 8.03 (d, 1 H), 7.48 - 7.56 (m, 3 H), 7.35 (d, 2 H), 7.05 (d, 2 H), 6.68 (dd, 1 H), 6.39 (dd, 1 H), 6.21 (d, 1 H), 4.17 - 4.34 (m, 3 H), 3.11 (s, 4 H), 2.89 (s, 2 H), 2.24 - 2.42 (m, 4 H), 2.15 (s, 2 H), 1.96 (s, 2 H), 1.66 - 1.82 (m, 3 H), 1.55 (d, 2 H), 1.31 - 1.44 (m, 4 H), 1.12 - 1.27 (m, 2 H), 1.10 (s, 3 H), 0.93 (s, 6 H).

50

EXAMPLE 348

N-([5-chloro-6-((cis-4-hydroxy-4-methylcyclohexyl)methoxy)pyridin-3-yl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

55

[1323] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 347C in place of EXAMPLE

11B. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1 H), 8.51 (d, 1 H), 8.18 (d, 1 H), 8.03 (d, 1 H), 7.47 - 7.58 (m, 3 H), 7.35 (d, 2 H), 7.05 (d, 2 H), 6.68 (dd, 1 H), 6.39 (dd, 1 H), 6.21 (d, 1 H), 4.21 (d, 2 H), 3.95 (s, 1 H), 3.11 (s, 4 H), 2.89 (s, 2 H), 2.33 (d, 4 H), 2.15 (s, 2 H), 1.96 (s, 2 H), 1.63 - 1.77 (m, 1 H), 1.48 - 1.60 (m, 4 H), 1.35 - 1.48 (m, 4 H), 1.20 - 1.33 (m, 2 H), 1.09 (s, 3 H), 0.93 (s, 6 H).

EXAMPLE 349

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{[4-({[2,2-difluorocyclopropyl]amino]cyclohexyl)amino)-3-nitrophenyl]sulfonyl}-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1324] The title compound was prepared by substituting 2,2-difluorocyclopropanamine hydrochloride for 3-(cyclopropylamino)propanenitrile in EXAMPLE 340D. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.60 (s, 1 H), 8.47 (m, 2 H), 8.12 (m, 1 H), 7.98 (m, 1 H), 7.72 (m, 2 H), 7.47 (m, 3 H), 7.34 (m, 3 H), 7.05 (m, 3 H), 6.65 (dd, 1 H), 6.35 (m, 1 H), 6.22 (d, 1 H), 3.54 (m, 2 H), 3.08 (m, 4 H), 2.74 (m, 4 H), 2.25 (m, 4 H), 2.01 (m, 4 H), 1.38 (m, 4 H), 0.92 (s, 6 H).

EXAMPLE 350

N-{[5-chloro-6-[(cis-1-fluoro-4-hydroxycyclohexyl)methoxy]pyridin-3-yl]sulfonyl}-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 350A

ethyl spiro[benzo[d][1,3]dioxole-2,1'-cyclohexane]-4'-carboxylate

[1325] To a solution of ethyl 4-oxocyclohexanecarboxylate (22.75 g) and pyrocatechol (14.75 g) in toluene (200 mL) was added catalytic amount of para-toluenesulfonic acid monohydrate and the mixture was stirred under reflux and a Dean-Stark trap overnight. The mixture was diluted with diethyl ether (600 mL) and washed with aqueous NaHCO₃, water and brine. After drying over Na₂SO₄, the mixture was filtered and the solvent was evaporated under vacuum to provide the title compound.

EXAMPLE 350B

ethyl 4'-fluorospiro[benzo[d][1,3]dioxole-2,1'-cyclohexane]-4'-carboxylate

[1326] A solution of EXAMPLE 350A (5.25g) in tetrahydrofuran (40 mL) was added dropwise to a solution of lithium diisopropylamide (12 mL, 2.0M in tetrahydrofuran/heptane/ethylbenzene) at 0°C. The solution was stirred at 0°C for 30 minutes, and then was transferred by cannula to a pre-cooled (0°C) stirring solution of N-fluorobenzenesulfonimide (7.89g) in dry tetrahydrofuran (20 mL). The reaction mixture was stirred at 0°C for 30 minutes, and then at 20°C for 18 hours. The reaction mixture was poured over aqueous NH₄Cl and extracted with diethyl ether (3x200 mL). The combined organic layers were washed with water, brine and dried over Na₂SO₄. Filtration and evaporation of the solvent gave the crude product.

EXAMPLE 350C

(4'-fluorospiro[benzo[d][1,3]dioxole-2,1'-cyclohexane]-4'-yl)methanol

[1327] To a solution of EXAMPLE 350B (23 g) in tetrahydrofuran (150 mL) was added lithium aluminum hydride (3.11 g). The mixture was stirred overnight. Aqueous 2N NaOH solution was added dropwise to the reaction mixture. The mixture was then diluted with ethyl acetate (600 mL) and washed with water, brine and dried over Na₂SO₄. Filtration and evaporation of the solvent gave the crude product which was loaded on a 600 g analogics column and eluted with 10% to 20% ethyl acetate in hexane to provide the title compound.

EXAMPLE 350D

5-chloro-6-((4'-fluorospiro[benzo[d][1,3]dioxole-2,1'-cyclohexane]-4'-yl)methoxy)pyridine-3-sulfonamide

[1328] To a solution of EXAMPLE 350C (89 mg) in N,N-dimethylformamide (3 mL) was added NaH (65% in mineral oil, 36 mg). The mixture was stirred for 30 minutes, and then 5, 6-dichloropyridine-3-sulfonamide (85 mg) was added.

The mixture was stirred overnight. The mixture was poured over aqueous NH_4Cl and extracted with ethyl acetate (100 mL). The combined organic layers were washed with water, brine and dried over Na_2SO_4 . After filtration and evaporation of the solvent, the residue was loaded on a silica gel cartridge and eluted with 30% ethyl acetate in hexane to provide the title compound.

5

EXAMPLE 350E

5-chloro-6-((1-fluoro-4-oxocyclohexyl)methoxy)pyridine-3-sulfonamide

10

[1329] To a solution of EXAMPLE 350D (1.6g) and pyridinium p-toluenesulfonate (1.2 g) in acetone (10 mL) was added water (2 mL) and the mixture was stirred under microwave irradiation at 100°C for 10 minutes. The mixture was diluted with dichloromethane (300 mL) and washed with aqueous NaHCO_3 , water, brine and dried over Na_2SO_4 . Filtration and evaporation of the solvent gave the title compound.

15

EXAMPLE 350F

5-chloro-6-((cis-1-fluoro-4-hydroxycyclohexyl)methoxy)pyridine-3-sulfonamide

20

[1330] To a solution of EXAMPLE 350E (336 mg) in tetrahydrofuran (10 mL) was added NaBH_4 (75 mg). The mixture was stirred for 45 minutes. The mixture was diluted with ethyl acetate (300 mL) and washed with 2N aqueous NaOH , water, and brine. After drying over Na_2SO_4 , the mixture was filtered and the solvent was evaporated to give the crude product.

EXAMPLE 350G

25

N-({5-chloro-6-((cis-1-fluoro-4-hydroxycyclohexyl)methoxy)pyridin-3-yl}sulfonyl)-4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

30

[1331] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 350F for EXAMPLE 1F in EXAMPLE 1G. ^1H NMR (300 MHz, dimethylsulfoxide- d_6) δ 11.63 (s, 1 H), 8.48 (s, 1 H), 8.18 (s, 1 H), 8.01 (d, 1 H), 7.50 (m, 3 H), 7.35 (d, 2 H), 7.05 (d, 2 H), 6.67 (dd, 1 H), 6.37 (dd, 1 H), 6.21 (d, 1 H), 4.62 (d, 1 H), 4.47 (s, 1 H), 4.40 (s, 1 H), 3.46 (m, 1 H), 3.06 (m, 4 H), 2.88 (m, 1 H), 2.25 (m, 6 H), 1.99 (m, 4 H), 1.58 (m, 8 H), 0.93 (s, 6 H).

EXAMPLE 351

35

4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-[3-nitro-4-(2-oxaspiro[3.5]non-7-yl-methoxy)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 351A

40

diethyl 1,4-dioxaspiro[4.5]decane-8,8-dicarboxylate

45

[1332] A 500 mL round-bottomed flask was charged with diisopropylamine (16 mL) and tetrahydrofuran (311 mL). The solution was cooled to -78°C under N_2 and n-BuLi (2.5 M in hexanes, 44.8 mL) was added. The reaction was stirred for 30 minutes at -78°C and ethyl 1,4-dioxaspiro[4.5]decane-8-carboxylate (20 g) was added as a tetrahydrofuran solution (ca. 10 mL). The solution was stirred at -78°C for 1 hour and ethyl chloroformate (9 mL) was added neat. After stirring at -78°C for 10 minutes, the reaction was warmed to room temperature over 2 hours. The reaction was quenched with saturated aqueous NH_4Cl and was diluted with diethyl ether. The layers were separated, the aqueous layer was extracted with diethyl ether and the combined organics were dried (Na_2SO_4), filtered and concentrated by rotary evaporation. The residue was purified by regular phase flash column chromatography (Analoxig, 0-65% hexanes / ethyl acetate).

50

EXAMPLE 351B

1,4-dioxaspiro[4.5]decane-8,8-diyl dimethanol

55

[1333] To a 1 L round-bottomed flask was added EXAMPLE 351A (26.6 g) and tetrahydrofuran (310 mL) to give a colorless solution. The solution was cooled to 0°C and lithium aluminum hydride (2M in tetrahydrofuran, 62 mL) was added via syringe. The reaction was allowed to warm to room temperature and stirred overnight. The mixture was cooled

back down to 0 °C and quenched slowly with 4.7 mL water, 4.7 mL 10% aqueous NaOH and 14 mL water. The mixture was allowed to stir until salts were formed and was then filtered through a Supelco 90 mm silica gel Buchner funnel. The filtrate was concentrated by rotary evaporation and the residue was purified by regular phase flash column chromatography (Analogix, 0-80% hexanes / ethyl acetate).

5

EXAMPLE 351C

2,8,11-trioxa-dispiro[3.2.4]tridecane

10

[1334] To a 1 L round-bottomed flask was added EXAMPLE 351B (13 g) in tetrahydrofuran (321 mL). The solution was cooled to -78 °C under N₂ and n-BuLi (25.7 mL) was added dropwise via syringe. After addition was complete, the mixture stirred for 30 minutes and a tetrahydrofuran solution of 4-toluenesulfonyl chloride (12.25 g) was added via addition funnel. The reaction was allowed to stir overnight, and gradually warm to room temperature. The reaction mixture was cooled to -78 °C and n-BuLi (25.7 mL) was added. The mixture was warmed to room temperature and stirred for 3

15

hours. The reaction was quenched with sat aqueous NH₄Cl and diluted with diethyl ether. The layers were separated, the aqueous layers extracted with diethyl ether and the combined organics were dried (Na₂SO₄), filtered and concentrated by rotary evaporation. The residue was purified by regular phase flash column chromatography (Analogix, 0-20% acetone / hexanes).

20

EXAMPLE 351D

2-oxaspiro[3.5]nonan-7-one

25

[1335] To a 500 mL round-bottomed flask was added EXAMPLE 351C (11 g) in 80 % aqueous acetic acid (200 mL). The reaction was heated to 65°C and stirred for about 4 hours. Most of the acetic acid and water were removed by rotary evaporation and the residue was purified by regular phase flash column chromatography (Analogix, 0-65% hexanes/ethyl acetate).

EXAMPLE 351E

30

7-methylene-2-oxaspiro[3.5]nonane

35

[1336] To a 250 mL round-bottomed flask was added methyltriphenylphosphonium iodide (4.33 g) in tetrahydrofuran (35.7 mL) to give a suspension. The suspension was cooled to -15 °C. n-BuLi (2.5 M in hexanes, 4.28 mL) was added dropwise and the mixture was stirred at -15 °C for 40 minutes and EXAMPLE 351D (1 g) was added as a tetrahydrofuran (ca. 5 mL) solution. The mixture was stirred at -15 °C for about 15 minutes and warmed to room temperature. After 1.5 hours, the reaction was complete and was quenched with saturated aqueous NH₄Cl and diluted with diethyl ether. The layers were separated and the aqueous layer was extracted (2x) with diethyl ether. The combined organics were washed with brine, dried (Na₂SO₄), filtered and concentrated by rotary evaporation. The residue was purified by regular phase

40

chromatography (Analogix, 80 g Grace silica gel column, 0-50% hexanes/ethyl acetate).

EXAMPLE 351F

2-oxaspiro[3.5]nonan-7-ylmethanol

45

[1337] To a 25 mL round-bottomed flask was added EXAMPLE 351E (568 mg) and EXAMPLE 351F tetrahydrofuran (4.11 mL) to give a colorless solution. 9-Borabicyclo[3.3.1]nonane (0.5 M in tetrahydrofuran, 24.7 mL) was added and the reaction was allowed to stir for 2 hours at room temperature. Ethanol (11 mL) was added followed by aqueous NaOH (5M, 4.11 mL) and then hydrogen peroxide (2.1 mL) was added. The reaction was heated at 50 °C for 2 hours. The mixture was concentrated by rotary evaporation, and diluted with water and ethyl acetate. The aqueous layer was extracted with ethyl acetate (3x) and the combined organics were dried (Na₂SO₄), filtered and concentrated by rotary evaporation. The residue was purified by regular phase flash column chromatography (Analogix, 80 g Grace, 0-70 % hexanes / ethyl acetate).

50

55

EXAMPLE 351G

4-(2-oxaspiro[3.5]nonan-7-ylmethoxy)-3-nitrobenzenesulfonamide EXAMPLE 351G was prepared substituting EXAMPLE 351F for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 351H

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-(4-(2-oxaspiro[3.5]nonan-7-ylmethoxy)-3-nitrophenylsulfonyl)-4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzamide

[1338] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 351G for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1 H) 8.34 (s, 1 H) 8.03 (d, 2 H) 7.45 - 7.57 (m, 3 H) 7.30 - 7.40 (m, 3 H) 7.04 (d, 2 H) 6.67 (dd, 1 H) 6.39 (dd, 1 H) 6.17 - 6.23 (m, 1 H) 4.29 (s, 2 H) 4.20 (s, 2 H) 4.00 (d, 2 H) 3.08 (s, 4 H) 2.73 - 2.90 (m, 2 H) 2.72 (s, 1 H) 2.01 - 2.32 (m, 6 H) 1.96 (s, 2 H) 1.64 - 1.78 (m, 4 H) 1.33 - 1.50 (m, 6 H) 0.96 - 1.15 (m, 2 H) 0.92 (s, 6 H).

EXAMPLE 352

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 352A

4-((trans-4-hydroxy-4-methylcyclohexyl)methoxy)-3-nitrobenzenesulfonamide

[1339] The title compound was prepared by substituting EXAMPLE 341F for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 352B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1340] The title compound was prepared by substituting EXAMPLE 352A for EXAMPLE 1F and EXAMPLE 3J for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1H), 8.31 (br s, 1H), 8.01 (m, 2H), 7.49 (m, 3H), 7.33 (m, 3H), 7.03 (m, 2H), 6.66 (dd, 1H), 6.37 (m, 1H), 6.19 (d, 1H), 4.27 (s, 1H), 4.05 (d, 2H), 3.40 (m, 2H), 3.17 (s, 1H), 3.07 (m, 3H), 2.79 (m, 1H), 2.24 (m, 3H), 2.14 (m, 2H), 1.94 (m, 2H), 1.71 (m, 3H), 1.52 (m, 2H), 1.38 (m, 4H), 1.22 (m, 2H), 1.09 (s, 3H), 0.91 (s, 6H).

EXAMPLE 353

4-(4-([2-(4-chlorophenyl)-5,5-bis(fluoromethyl)cyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 353A

1,4-dioxaspiro[4.5]decane-8,8-diylbis(methylene) bis(4-methylbenzenesulfonate)

[1341] To a 500 mL round-bottomed flask was added EXAMPLE 351B (10 g) and dichloromethane (165 mL) to give a colorless solution. Triethylamine (24.1 mL) and toluene-2-sulfonyl chloride (19.8 g) were added followed by 4-dimethylaminopyridine (0.604 g). The reaction was refluxed overnight. Saturated aqueous NH₄Cl was added followed by dilution with water and additional dichloromethane. The aqueous layer was extracted with dichloromethane (2x) and the combined organics were dried (MgSO₄), filtered and concentrated by rotary evaporation. The residue was purified by regular phase flash column chromatography (Analogix, 0-55% hexanes / ethyl acetate).

EXAMPLE 353B

8,8-bis(fluoromethyl)-1,4-dioxaspiro[4.5]decane

5 **[1342]** To a 500 mL round-bottomed flask was added EXAMPLE 353A (20 g). tetra-*n*-Butylammonium fluoride (1M in tetrahydrofuran, 200 mL) was added and the resulting solution was refluxed for 6 days. The reaction was cooled, diluted with diethyl ether and washed with water (3x). The organics were dried (Na₂SO₄), filtered and concentrated by rotary evaporation. The residue was purified by regular phase flash column chromatography (Analogix, 0-30% hexanes / ethyl acetate).

10

EXAMPLE 353C

4,4-bis(fluoromethyl)cyclohexanone

15 **[1343]** To a 250 mL round bottom flask was added EXAMPLE 353B (1.1 g) and 80% aqueous acetic acid (50 mL). The reaction was heated at 65 °C for 3 hours, cooled and concentrated by rotary evaporation to remove most of the acetic acid and water. The residue was purified by regular phase flash column chromatography (Analogix, 0-50% hexanes / ethyl acetate).

20 EXAMPLE 353D

2-chloro-5,5-bis(fluoromethyl)cyclohex-1-enecarbaldehyde

25 **[1344]** To a 100 mL pear flask was added N,N-dimethylformamide (498 µl) and dichloromethane (8.9 mL) to give a colorless solution. The solution was cooled to 0 °C and POCl₃ (550 µl) was added dropwise and then the mixture was warmed to room temperature for 30 minutes. In the meantime, to a 100 mL pear shaped flask was added EXAMPLE 353C (870 mg, 5.36 mmol) in dichloromethane (8941 µl) to give a colorless solution. The Vilsmeier reagent was then taken up in a syringe and added dropwise to the 4,4-bis(fluoromethyl)cyclohexanone (870 mg) solution at room temperature. The resulting solution was stirred overnight. The reaction was poured into saturated aqueous NaHCO₃ and ice, warmed to room temperature and extracted with dichloromethane (3 x 30 mL). The organics were combined, dried over MgSO₄, filtered and concentrated by rotary evaporation. The residue was purified by regular phase flash column chromatography (Analogix (0-60% hexanes / ethyl acetate).

30

EXAMPLE 353E

35

2-(4-chlorophenyl)-5,5-bis(fluoromethyl)cyclohex-1-enecarbaldehyde

40 **[1345]** To a 20 mL vial was added EXAMPLE 353D (460 mg), 4-chlorophenylboronic acid (414 mg), potassium carbonate (762 mg), tetrabutylammonium bromide (711 mg), palladium (II) acetate (14.85 mg) and water (2450 µl) to give a suspension which was degassed with N₂ for 2 minutes. The reaction was stirred at 45 °C overnight, cooled, and poured over a Supelco silica gel Buchner funnel, washing with ethyl acetate several times. The filtrate was concentrated by rotary evaporation and the residue was purified by regular phase flash column chromatography (Analogix, 0-60% hexanes / ethyl acetate).

45 EXAMPLE 353F

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5,5-bis(fluoromethyl)cyclohex-1-enyl)methyl)piperazin-1-yl)benzoate

50 **[1346]** To a 20 mL vial was added EXAMPLE 353E (240 mg), EXAMPLE 15F (297 mg) and dichloromethane (4.2 mL). Sodium triacetoxyborohydride (268 mg) was added and the reaction was stirred overnight at room temperature. The reaction was loaded directly onto silica gel and purified by regular phase flash column chromatography (Analogix, 0-80% hexanes / ethyl acetate).

55

EXAMPLE 353G

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5,5-bis(fluoromethyl)cyclohex-1-enyl)methyl)piperazin-1-yl)benzoic acid

5

[1347] The title compound was prepared by substituting EXAMPLE 353F for EXAMPLE 15G in EXAMPLE 15 H.

EXAMPLE 353H

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5,5-bis(fluoromethyl)cyclohex-1-enyl)methyl)piperazin-1-yl)-N-(3-nitro-4-((tetrahydro-2H-pyran-4-yl)methylamino)phenylsulfonyl)benzamide

10

[1348] EXAMPLE 353H was prepared by replacing EXAMPLE 3J with EXAMPLE 353G and EXAMPLE 11B with EXAMPLE 1F in EXAMPLE 11D. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1 H) 11.44 (s, 1 H) 8.48 - 8.70 (m, 1 H) 8.05 (d, 2 H) 7.81 (dd, 1 H) 7.46 - 7.59 (m, 3 H) 7.35 (d, 2 H) 7.12 (d, 2 H) 6.68 (dd, 1 H) 6.40 (dd, 1 H) 6.16 (d, 1 H) 4.39 - 4.49 (m, 2 H) 4.23 - 4.35 (m, 2 H) 3.85 (dd, J=11.87, 2.71 Hz, 2 H) 3.20 - 3.30 (m, 4 H) 2.98 - 3.10 (m, 4 H) 2.66 - 2.77 (m, 2 H) 2.11 - 2.30 (m, 6 H) 2.02 - 2.12 (m, 3 H) 1.99 (s, 1 H) 1.82 - 1.97 (m, 1 H) 1.54 - 1.67 (m, 4 H) 1.20 - 1.34 (m, 2 H).

15

20 EXAMPLE 354

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(4-cyclopropylmorpholin-2-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

25 EXAMPLE 354A

tert-butyl 2-((2-nitro-4-sulfamoylphenoxy)methyl)morpholine-4-carboxylate

[1349] The title compound was prepared by substituting tert-butyl 2-(hydroxymethyl)-morpholine-4-carboxylate for tetrahydro-2H-pyran-4-yl-methanol EXAMPLE 24A.

30

EXAMPLE 354B

4-(morpholin-2-ylmethoxy)-3-nitrobenzenesulfonamide

35

[1350] The title compound was prepared by substituting EXAMPLE 354A for EXAMPLE 113A in EXAMPLE 134A.

EXAMPLE 354C

4-((4-cyclopropylmorpholin-2-yl)methoxy)-3-nitrobenzenesulfonamide

40

[1351] The title compound was prepared by substituting EXAMPLE 354B for EXAMPLE 173A in EXAMPLE 173B.

EXAMPLE 354D

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-(4-cyclopropylmorpholin-2-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

45

[1352] The title compound was prepared by substituting EXAMPLE 354C for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 12.98 (s, 1H), 9.06 (d, 1H), 8.50 (dd, 1H), 8.41 (d, 1H), 8.09 (d, 1H), 7.66 (t, 1H), 7.62 (d, 1H), 7.44 (d, 2H), 7.26 (d, 1H), 7.07 (d, 2H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 4.31 (dd, 1H), 4.22 (dd, 1H), 3.92 (m, 1H), 3.83 (d, 1H), 3.56 (dt, 1H), 3.07 (m, 5H), 2.77 (s, 2H), 2.68 (d, 1H), 2.35 (m, 2H), 2.26 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.59 (m, 1H), 1.39 (t, 2H), 0.94 (s, 6H), 0.40 (m, 4H).

50

55

EXAMPLE 355

N-({5-chloro-6-[(trans-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 355A

5-chloro-6-((trans-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy)pyridine-3-sulfonamide

[1353] To a cooled (0°C) solution of EXAMPLE 350E (1.2 g) in tetrahydrofuran (30 mL) was added dropwise a solution of methylmagnesium bromide (5 mL, 3.0M in ether). Upon addition, the reaction mixture solidified. More tetrahydrofuran (10 mL) was added to the mixture and stirring was continued for 1 hour. The mixture was poured over aqueous NH₄Cl and extracted with ethyl acetate (3x 150 mL). The combined organic layers were washed with water, brine and dried over Na₂SO₄. The mixture was filtered and concentrated. The residue was dissolved in dimethylsulfoxide/methanol (20 mL, 1:1) and loaded on Gilson, C18(100A) 250x121.2 mm (10 micron), with 30% acetonitrile to 65% acetonitrile over 40 minutes to separate the two isomers and isolate the title compound.

EXAMPLE 355B

N-({5-chloro-6-[(trans-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1354] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 355A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.63 (s, 1 H), 8.47 (s, 1 H), 8.17 (s, 1 H), 7.54 (d, 1 H), 7.48 (m, 2 H), 7.35 (d, 2 H), 7.05 (d, 2 H), 6.67 (dd, 1 H), 6.37 (d, 1 H), 6.22 (d, 1 H), 4.49 (s, 1 H), 4.42 (s, 1 H), 4.15 (s, 1 H), 3.06 (m, 4 H), 2.84 (m, 1 H), 2.25 (m, 6 H), 1.96 (s, 3 H), 1.83 (m, 4 H), 1.44 (m, 6 H), 1.14 (s, 3 H), 0.93 (s, 6 H).

EXAMPLE 356

[1355] N-({5-chloro-6-[(cis-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 356A

5-chloro-6-((cis-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy)pyridine-3-sulfonamide The title compound was prepared as described in EXAMPLE 355A.

EXAMPLE 356B

N-({5-chloro-6-[(cis-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1356] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 356A for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1 H), 8.52 (s, 1 H), 8.20 (s, 1 H), 8.03 (d, 1 H), 7.51 (m, 3 H), 7.35 (d, 2 H), 7.05 (d, 2 H), 6.67 (dd, 1 H), 6.39 (dd, 1 H), 6.21 (d, 1 H), 4.55 (s, 1 H), 4.48 (s, 1 H), 4.34 (s, 1 H), 3.08 (m, 4 H), 2.89 (d, 2 H), 2.27 (m, 5 H), 1.93 (m, 4 H), 1.66 (m, 4 H), 1.43 (m, 4 H), 1.11 (s, 3 H), 0.93 (s, 6 H).

EXAMPLE 357

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-cyano-4-{[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 357A

ethyl 4-fluoro-1-(oxetan-3-yl)piperidine-4-carboxylate

[1357] To 1-*tert*-butyl 4-ethyl 4-fluoropiperidine-1,4-dicarboxylate (1.000 g) was added HCl (4.0M in dioxane, 4.54 mL). After 1 hour the reaction was concentrated and dried under high vacuum. The resulting solid was dissolved in dichloromethane (5 mL) and treated with sodium triacetoxyborohydride (1.155 g) and oxetan-3-one (0.262 g) and stirred overnight. The reaction was quenched with saturated NaHCO₃ solution (20 mL) and extracted into dichloromethane (2 x 25 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated. Silica gel chromatography (Reveleris 80 g) eluting with a gradient of 0.5% to 3.75% methanol/dichloromethane over 40 minutes (flow = 30 mL/min) gave the title compound.

EXAMPLE 357B

(4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methanol

[1358] To a solution of EXAMPLE 357A (0.59 g) in tetrahydrofuran (5 mL) was added lithium aluminum hydride (1.80 mL) at 0°C. The reaction was removed from the ice bath and allowed to warm to room temperature. The reaction was quenched by the dropwise addition of 0.6 mL of water followed by 0.2 mL of 2N aqueous NaOH. The reaction was filtered through celite and rinsed with ethyl acetate (50 mL). The mixture and the residue was loaded onto silica gel (Reveleris 40 g) and eluted using a gradient of 0.75% to 7.5% methanol/dichloromethane over 30 minutes (flow = 40 mL/minutes) to provide the title compound.

EXAMPLE 357C

3-cyano-4-((4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methoxy)benzenesulfonamide

[1359] The title compound was prepared by substituting EXAMPLE 357B for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 284A.

EXAMPLE 357D

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(3-cyano-4-{[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1360] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 357C for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-*d*₆) δ 11.67 (s, 1H), 11.49 - 11.14 (m, 1H), 8.17 (d, 1H), 8.03 (d, 2H), 7.51 (dd, 3H), 7.43 - 7.26 (m, 3H), 7.12 - 6.96 (m, 2H), 6.67 (dd, 1H), 6.40 (dd, 1H), 6.20 (d, 1H), 4.55 (t, 2H), 4.45 (t, 2H), 4.34 (d, 2H), 3.49 (s, 1H), 3.09 (s, 8H), 2.39 - 1.66 (m, 14H), 1.39 (s, 2H), 0.92 (s, 6H).

EXAMPLE 358

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(trans-4-ethyl-4-hydroxycyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 358A

benzyl (4-ethyl-4-hydroxycyclohexyl)methylcarbamate

[1361] To a vigorous stirring solution of benzyl (4-oxocyclohexyl)methylcarbamate (1 g) in tetrahydrofuran (20 mL) at -78°C was slowly added 1 M ethylmagnesium bromide (11.48 mL, 11.48 mmol) in ether. After completion of the addition, the mixture was stirred at -78°C for 2 hours and was warmed to 0°C, and stirred in an ice bath for 30 minutes. The reaction was quenched with a cold NH₄Cl aqueous solution. The precipitates were filtered off and washed with ethyl

acetate. The filtrate was concentrated. The residue was dissolved in dichloromethane and loaded onto Analogix purification system, and was eluted with 0 - 50% ethyl acetate in dichloromethane to provide the title compound.

EXAMPLE 358B

4-(aminomethyl)-1-ethylcyclohexanol

[1362] A mixture of EXAMPLE 358A (500 mg) and 10% Pd/C (100 mg) in tetrahydrofuran (15 mL) was stirred under H₂ for 3 hours. The insoluble material was removed by filtration, and the filtrate was concentrated to provide the title compound.

EXAMPLE 358C

4-((trans-4-ethyl-4-hydroxycyclohexyl)methylamino)-3-nitrobenzenesulfonamide

[1363] EXAMPLE 358B (270 mg) and 4-fluoro-3-nitrobenzenesulfonamide (417 mg) in tetrahydrofuran were treated with triethylamine (0.8 mL) overnight. The reaction was quenched with water. The resulting mixture was neutralized with diluted aqueous HCl, and extracted with ethyl acetate. The organic layer was dried over Na₂SO₄, filtered and concentrated. The residue was purified by a reverse phase chromatography, eluting with 40-55% acetonitrile in 0.1% trifluoroacetic acid water to isolate the title compound.

EXAMPLE 358D

4-((cis-4-ethyl-4-hydroxycyclohexyl)methylamino)-3-nitrobenzenesulfonamide The title compound was prepared and isolated as described in Example 358C.

EXAMPLE 358E

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((trans-4-ethyl-4-hydroxycyclohexyl)methyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1364] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 358C in place of EXAMPLE 11B. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1 H), 11.35 (s, 1 H), 8.56 (d, 2 H), 8.05 (d, 1 H), 7.80 (dd, 1 H), 7.45 - 7.57 (m, 3 H), 7.34 (d, 2 H), 7.00 - 7.10 (m, 3 H), 6.68 (dd, 1 H), 6.39 (dd, 1 H), 6.19 (d, 1 H), 3.98 (s, 1 H), 3.24-3.31 (m, 4 H), 3.07 (s, 4 H), 2.75 (s, 2 H), 2.17 (d, 6 H), 1.95 (s, 2 H), 1.54 - 1.73 (m, 5 H), 1.35 - 1.47 (m, 4 H), 1.20 - 1.32 (m, 2 H), 1.03 - 1.18 (m, 2 H), 0.92 (s, 6 H), 0.81 (t, 3 H).

EXAMPLE 359

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-((cis-4-ethyl-4-hydroxycyclohexyl)methyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1365] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 358D in place of EXAMPLE 11B. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.69 (s, 1 H), 11.34 (s, 1 H), 8.60 (t, 1 H), 8.56 (d, 1 H), 8.05 (d, 1 H), 7.80 (dd, 1 H), 7.54 (d, 1 H), 7.47 - 7.52 (m, 2 H), 7.34 (d, 2 H), 7.01 - 7.10 (m, 3 H), 6.68 (dd, 1 H), 6.39 (dd, 1 H), 6.19 (d, 1 H), 3.77 (s, 1 H), 3.26 (t, 2 H), 3.07 (s, 4 H), 2.76 (s, 2 H), 2.10 - 2.26 (m, 6 H), 1.95 (s, 2 H), 1.46 - 1.61 (m, 5 H), 1.28 - 1.46 (m, 6 H), 1.12 - 1.24 (m, 2 H), 0.92 (s, 6 H), 0.82 (t, 3 H).

EXAMPLE 360

4-(4-([2-(4-chlorophenyl)-5-(methoxymethyl)-5-methylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-((tetrahydro-2H-pyran-4-ylmethyl)amino)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 360A

ethyl 8-methyl-1,4-dioxaspiro[4.5]decane-8-carboxylate

[1366] Into a 500 mL round-bottomed flask was added diisopropylamine (7.98 mL) in tetrahydrofuran (233 mL) to give

a colorless solution. The mixture was cooled to -78 °C under N₂ and n-BuLi (2.5 M in hexanes, 22.40 mL) was added. The reaction was stirred for 30 minutes and ethyl 1,4-dioxaspiro[4.5]decane-8-carboxylate (10 g) was added. The reaction was allowed to stir for 1.5 hours upon which time CH₃I (4.38 mL) was added. The reaction was allowed to warm to room temperature overnight with stirring. Water was added and the aqueous layer was extracted with ethyl acetate. The combined organics were dried (Na₂SO₄), filtered and concentrated by rotary evaporation. The residue was purified by normal phase flash column chromatography (Analogix, 0-50% hexanes / ethyl acetate).

EXAMPLE 360B

(8-methyl-1,4-dioxaspiro[4.5]decan-8-yl)methanol

[1367] In a 500 mL round-bottomed flask was lithium aluminum hydride (1.772 g) in tetrahydrofuran (234 mL) to give a suspension. This suspension was cooled to 0 °C and ethyl 8-methyl-1,4-dioxaspiro[4.5]decane-8-carboxylate (10.66 g) was added via addition funnel. The reaction was stirred overnight at room temperature and then cooled back down to 0 °C. The excess lithium aluminum hydride was slowly quenched with 1.8 mL water, 1.8 mL aqueous NaOH (5N) and 5.6 mL water. The suspension was stirred until the salts turned white and was then filtered through a plug of silica gel. The filtrate was concentrated by rotary evaporation and the residue was purified by regular phase flash column chromatography (Analogix, 0-75% hexanes / ethyl acetate).

EXAMPLE 360C

8-(methoxymethyl)-8-methyl-1,4-dioxaspiro[4.5]decane

[1368] To a 250 mL round-bottomed flask was added NaH (0.902 g) and tetrahydrofuran (37.6 mL) to give a suspension. EXAMPLE 360B was added as a tetrahydrofuran solution at room temperature. The suspension was stirred for 30 minutes and then CH₃I (0.611 mL) was added. The reaction was stirred under N₂ overnight, carefully quenched with brine and diluted with water and ether. The aqueous layer was extracted with ether (2x) and the combined organics were dried (Na₂SO₄), filtered and concentrated by rotary evaporation. The residue was purified by flash column chromatography (Analogix, 0-60% hexanes / ethyl acetate).

EXAMPLE 360D

4-(methoxymethyl)-4-methylcyclohexanone

[1369] The title compound was prepared by substituting EXAMPLE 360C for EXAMPLE 353B in EXAMPLE 353C.

EXAMPLE 360E

2-chloro-5-(methoxymethyl)-5-methylcyclohex-1-enecarbaldehyde

[1370] The title compound was prepared by substituting EXAMPLE 360D for EXAMPLE 353C in EXAMPLE 353D.

EXAMPLE 360F

2-(4-chlorophenyl)-5-(methoxymethyl)-5-methylcyclohex-1-enecarbaldehyde

[1371] The title compound was prepared by substituting EXAMPLE 360E for EXAMPLE 353D in EXAMPLE 353E.

EXAMPLE 360G

methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5-(methoxymethyl)-5-methylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoate

[1372] The title compound was prepared by substituting EXAMPLE 360F for EXAMPLE 353E in EXAMPLE 353F.

EXAMPLE 360H

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5-(methoxymethyl)-5-methylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoic acid

[1373] The title compound was prepared by substituting EXAMPLE 360G for EXAMPLE 15G in EXAMPLE 15H.

EXAMPLE 360I

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5-(methoxymethyl)-5-methylcyclohex-1-enyl)methyl)piperazin-1-yl)-N-(3-nitro-4-((tetrahydro-2H-pyran-4-yl)methylamino)phenylsulfonyl)benzamide

[1374] The title compound was prepared by replacing EXAMPLE 3J with EXAMPLE 360H and EXAMPLE 11B with EXAMPLE 1F in EXAMPLE 11D. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.68 (s, 1 H) 11.43 (s, 1 H) 8.45 - 8.72 (m, 2 H) 8.04 (d, 1 H) 7.80 (dd, 1 H) 7.44 - 7.61 (m, 3 H) 7.34 (d, 2 H) 6.99 - 7.20 (m, 3 H) 6.68 (dd, 1 H) 6.39 (dd, 1 H) 6.18 (d, 1 H) 3.85 (dd, 2 H) 3.25 - 3.30 (m, 4 H) 3.24 (s, 3 H) 3.02 - 3.17 (m, 6 H) 2.72 (dd, 2 H) 2.18 (s, 5 H) 2.03 - 2.13 (m, 2 H) 1.81 - 1.93 (m, 2 H) 1.57 - 1.67 (m, 2 H) 1.47 - 1.56 (m, 1 H) 1.17 - 1.41 (m, 3 H) 0.91 (s, 3 H).

EXAMPLE 361

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-(((2S)-4-(oxetan-3-yl)morpholin-2-yl)methyl)amino)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 361A

[1375] (S)-3-nitro-4-((4-(oxetan-3-yl)morpholin-2-yl)methylamino)benzenesulfonamide The title compound was prepared by substituting EXAMPLE 259E for tert-butyl piperazine-1-carboxylate and 3-oxetanone for 4'-chlorobiphenyl-2-carboxaldehyde in EXAMPLE 1A.

EXAMPLE 361B

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-(((2S)-4-(oxetan-3-yl)morpholin-2-yl)methyl)amino)phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1376] The title compound was prepared by substituting EXAMPLE 361A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.00 (s, 1H), 9.26 (d, 1H), 8.87 (t, 1H), 8.43 (d, 1H), 8.35 (dd, 1H), 8.11 (d, 1H), 7.66 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.96 (d, 1H), 6.75 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 4.64 (m, 4H), 3.93 (m, 1H), 3.89 (d, 1H), 3.68 (dt, 1H), 3.53-3.35 (m, 3H), 3.07 (m, 4H), 2.77 (s, 2H), 2.72 (d, 1H), 2.44 (d, 1H), 2.26 (m, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.85 (t, 1H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 362

N-([3-chloro-4-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]phenyl]sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 362A

3-chloro-4-(((1*r*, 4*r*)-4-hydroxy-4-methylcyclohexyl)methoxy)benzenesulfonamide

[1377] To a solution of EXAMPLE 341F (300 mg) in N,N-dimethylformamide (10 mL) was added sodium hydride (416 mg) portionwise. The resulting suspension was stirred for 15 minutes. 3-Chloro-4-fluorobenzenesulfonamide (425 mg) was added and stirring was continued for 72 hours. The reaction was quenched with water and the pH was adjusted to ca. 7. The mixture was diluted with brine (75 mL) and extracted with methylene chloride. The crude product was isolated from the dried methylene chloride layer by concentration and was purified on silica gel eluted with a 10, 25, 50 % ethyl acetate in methylene chloride step gradient to provide the title compound.

EXAMPLE 362B

N-({3-chloro-4-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]phenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1378] The title compound was prepared by substituting EXAMPLE 362A for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (400 MHz, pyridine- d₅) δ 13.07 (m, 1H), 8.58 (d, 1H), 8.45 (d, 1H), 8.31 (dd, 1H), 8.11 (d, 1H), 7.69 - 7.67 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.97 (d, 1H), 6.74 (dd, 1H), 6.52 (m, 2H), 5.34 (br s, 2H), 3.82 (d, 2H), 3.06 (m, 4H), 2.77 (s, 2H), 2.25 (m, 2H), 2.13 (m, 4H), 1.97 - 1.85 (m, 7H), 1.82 - 1.73 (m, 2H), 1.44 - 1.32 (m, 7H), 0.94 (m, 6H).

EXAMPLE 363

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{4-[(2-cyanoethyl)(cyclopropyl)amino]-1-fluorocyclohexyl}methoxy}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 363A

4-((4'-fluorospiro[benzo[d][1,3]dioxole-2,1'-cyclohexane]-4'-yl)methoxy)-3-nitrobenzenesulfonamide

[1379] To a solution of EXAMPLE 350C (495 mg) in N,N-dimethylformamide (6 mL) was added NaH (65% in mineral oil, 320 mg). The mixture was stirred for 30 minutes, and then 4-fluoro-3-nitrobenzenesulfonamide (457 mg) was added. The mixture was stirred overnight. The mixture was poured over aqueous NH₄Cl and extracted with ethyl acetate (300 mL). The combined organic layers were washed with water, brine and dried over Na₂SO₄. After filtration and evaporation of the solvent, the residue was loaded on a silica gel cartridge and was eluted with 30% ethyl acetate in hexane to provide the title compound.

EXAMPLE 363B

4-((1-fluoro-4-oxocyclohexyl)methoxy)-3-nitrobenzenesulfonamide

[1380] To a solution of EXAMPLE 363A (860 mg) in ethanol (30 mL) was added concentrated HCl (10 mL) and the mixture was stirred at 100°C for 3 hours. The mixture was neutralized with solid Na₂CO₃ and extracted with dichloromethane (300 mL) and washed with aqueous NaHCO₃, water, brine and dried over Na₂SO₄. Filtration and evaporation of the solvent gave the title compound.

EXAMPLE 363C

4-((4-((2-cyanoethyl)(cyclopropyl)amino)-1-fluorocyclohexyl)methoxy)-3-nitrobenzenesulfonamide

[1381] To a solution of EXAMPLE 363B (200 mg) in dichloromethane (6 mL) was added 3-(cyclopropylamino)propanenitrile (64 mg) followed by sodium triacetoxyborohydride (184 mg). The mixture was stirred overnight. The reaction mixture was diluted with dichloromethane (400 mL) and washed with 2N aqueous NaOH, water, and brine. After drying over Na₂SO₄, the mixture was filtered and evaporation of the solvent gave the title compound.

EXAMPLE 363D

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-{{4-[(2-cyanoethyl)(cyclopropyl)amino]-1-fluorocyclohexyl}methoxy}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1382] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 363C for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.66 (s, 1H), 8.35 (s, 1H), 8.02 (d, 2H), 7.51 (m, 3H), 7.40 (m, 1H), 7.35 (d, 2H), 7.04 (d, 2H), 6.67 (dd, 1H), 6.39 (d, 1H), 6.20 (s, 1H), 4.27 (d, 2H), 3.13 (m, 4H), 2.88 (m, 3H), 2.67 (m, 4H), 2.09 (m, 10H), 1.49 (m, 9H), 0.93 (s, 6H), 0.45 (m, 4H).

EXAMPLE 364

4-(4-{ [2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({5-nitro-6-[(tetrahydro-2H-pyran-4-ylmethyl)amino]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 364A

6-amino-5-nitropyridine-3-sulfonic acid

[1383] 6-Aminopyridine-3-sulfonic acid (20 g) in concentrated H₂SO₄ (80 mL) was heated at 50 °C until it was completely dissolved. To this solution was added fuming HNO₃ slowly over 20 minutes, so the internal temperature did not exceed 55 °C. After the addition was complete, the reaction mixture was heated at 50 °C for 1 hour. After it was cooled to room temperature, it was poured into 150 g of ice. The mixture was stirred for another hour. The flask was cooled to 0 °C, and was kept at 0 °C for another 2 hours. The solid was collected by filtration, and washed with cold 1:1 water/ethanol (20 mL), followed by diethyl ether (10 mL). The solid was dried in a vacuum oven overnight to provide the title compound.

EXAMPLE 364B

6-hydroxy-5-nitropyridine-3-sulfonic acid

[1384] EXAMPLE 364A (4.0 g) in aqueous HCl (37%, 12 mL) and water (50 mL) was treated with sodium nitrite (1.19 g) in water (8 mL) dropwise at 0 °C. After the addition was complete, the reaction mixture was stirred at 0 °C for 1 hour. The mixture was heated at reflux for 2 hours. Water was distilled off to give a dry residue. After the residue was cooled to room temperature, a solution of 1:1 ethano/water (20 mL) was added. The resulting suspension was cooled to 0 °C, and kept at 0 °C for 1 hour. The solid was collected by filtration to provide the title compound.

EXAMPLE 364C

6-chloro-5-nitropyridine-3-sulfonyl chloride

[1385] A mixture of EXAMPLE 364B (2.6 g), PCl₅ (5.91 g), and POCl₃ (10 mL) was heated at 120 °C for 4 hours. The initial suspension became a clear solution. The excess of POCl₃ was distilled off. After it was cooled to room temperature, the residue was poured into 50 g of crushed ice. The solid was extracted into ethyl acetate. The aqueous layer was extracted with additional ethyl acetate. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated to give crude product that was used in the next step without further purification.

EXAMPLE 364D

6-chloro-5-nitropyridine-3-sulfonamide

[1386] EXAMPLE 364C in tetrahydrofuran (10 mL) was cooled to -10 °C. To this solution was added concentrated ammonium hydroxide (0.82 mL) dropwise. The solution was stirred at -10 °C for 10 minutes. The solvent was removed under pressure at room temperature. The residue was partitioned between water and ethyl acetate. The aqueous layer was extracted with additional ethyl acetate. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified by flash column chromatography on silica gel using 5-50% ethyl acetate in hexanes to provide the title compound.

EXAMPLE 364E

5-nitro-6-((tetrahydro-2H-pyran-4-yl)methylamino)pyridine-3-sulfonamide

[1387] The title compound was prepared by substituting EXAMPLE 364D for 4-fluoro-3-nitrobenzenesulfonamide and (tetrahydro-2H-pyran-4-yl)methanamine for (4-fluorotetrahydro-2H-pyran-4-yl)methanamine in EXAMPLE 138D.

EXAMPLE 364F

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({5-nitro-6-[(tetrahydro-2H-pyran-4-ylmethyl)amino]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1388] The title compound was prepared by substituting EXAMPLE 364E for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.63 (s, 1H), 8.93 (s, 1H), 8.73 (d, 1H), 8.69 (d, 1H), 8.00 (d, 1H), 7.54 (d, 1H), 7.47-7.48 (m, 2H), 7.35 (d, 2H), 7.05 (d, 2H), 6.68 (dd, 1H), 6.35 (dd, 1H), 6.22 (d, 1H), 3.83 (dd, 2H), 3.51 (t, 2H), 3.21-3.27 (m, 2H), 3.10 (s, 4H), 2.83 (s, 2H), 1.90-2.27 (m, 12H), 1.58 (dd, 2H), 1.39 (t, 2H), 1.18-1.28 (m, 2H), 0.88-0.93 (m, 8H).

EXAMPLE 365

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(2-oxaspiro[3.5]nonan-7-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 365A

7-(azidomethyl)-2-oxaspiro[3.5]nonane

[1389] To a 250 mL round-bottomed flask was EXAMPLE 351F (350 mg) in tetrahydrofuran (75.0 mL) to give a colorless solution. The solution was cooled to 0 °C, triphenylphosphine (2.94 g), diisopropyl azodicarboxylate (2.18 mL) and diphenyl phosphorazidate (2.32 mL) were added and the reaction was stirred for 30 minutes at room temperature. The mixture was concentrated and purified the residue by regular phase flash column chromatography (Analogix, 0-20% hexanes / ethyl acetate).

EXAMPLE 365B

2-oxaspiro[3.5]nonan-7-ylmethanamine

[1390] To a 50 mL round-bottomed flask was added 10% palladium on carbon (58.7 mg). The flask was flushed with N₂ and EXAMPLE 365A (400 mg) was added as a methanol solution (10.5 mL). The flask was then flushed several times with H₂ (via balloon) and heated to 45 °C for 2 hours. The reaction was cooled to room temperature, filtered through celite and the filtrate was concentrated by rotary evaporation. The residue was used in the next step without further purification.

EXAMPLE 365C

4-(2-oxaspiro[3.5]nonan-7-ylmethylamino)-3-nitrobenzenesulfonamide

[1391] The title compound was prepared by substituting EXAMPLE 365B for 1-(tetrahydropyran-4-yl)methylamine in EXAMPLE 1F.

EXAMPLE 365D

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-(4-(2-oxaspiro[3.5]nonan-7-ylmethylamino)-3-nitrophenylsulfonyl)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzamide

[1392] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 365C for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1 H) 11.25 - 11.49 (m, 1 H) 8.48 - 8.66 (m, 2 H) 8.03 (d, 1 H) 7.79 (dd, 1 H) 7.41 - 7.61 (m, 3 H) 7.27 - 7.40 (m, 2 H) 7.05 (t, 3 H) 6.67 (dd, 1 H) 6.39 (dd, 1 H) 6.18 (d, 1 H) 4.29 (s, 2 H) 4.19 (s, 2 H) 3.17 - 3.27 (m, 2 H) 2.99 - 3.14 (m, 4 H) 2.69 - 2.79 (m, 2 H) 2.09 - 2.28 (m, 6 H) 2.04 (d, 2 H) 1.95 (s, 2 H) 1.66 (d, 2 H) 1.49 - 1.61 (m, 1 H) 1.29 - 1.45 (m, 4 H) 0.93 - 1.05 (m, 2 H) 0.92 (s, 6 H).

EXAMPLE 366

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-cyano-4-methylcyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 366A

tert-butyl (4-cyano-4-methylcyclohexyl)methylcarbamate

[1393] To a cooled (-78°C) solution of tert-butyl (4-cyanocyclohexyl)methylcarbamate (500 mg) in tetrahydrofuran (10 mL) was added lithium diisopropylamide (2.0 mL, 2M in heptane). The mixture was stirred at -78°C for 30 minutes before the addition of CH₃I (1 mL). The mixture was then stirred and the temperature was allowed to warm to room temperature. The reaction was quenched with aqueous NH₄Cl and the mixture was extracted with ethyl acetate (300 mL) and washed with water, brine and dried over Na₂SO₄. Filtration and evaporation of the solvent and silica gel chromatography (40% ethyl acetate in hexane) of the crude material gave the title compound.

EXAMPLE 366B

4-(aminomethyl)-1-methylcyclohexanecarbonitrile

[1394] To a solution of EXAMPLE 366A (480 mg) in dichloromethane (10 mL) was added trifluoroacetic acid (10 mL). The mixture was stirred for 3 hours. The mixture was then concentrated under vacuum and was used directly in the next reaction without further purification.

EXAMPLE 366C

4-((4-cyano-4-methylcyclohexyl)methylamino)-3-nitrobenzenesulfonamide

[1395] To a solution of 4-fluoro-3-nitrobenzenesulfonamide (362 mg) in tetrahydrofuran (10 mL) was added EXAMPLE 366B (250 mg) and N,N-diisopropylethylamine (2 mL). The mixture was stirred overnight. The mixture was diluted with ethyl acetate (300 mL) and washed with water, brine and dried over Na₂SO₄. Filtration and evaporation of the solvent gave the title compound.

EXAMPLE 366D

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(4-cyano-4-methylcyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1396] The title compound was prepared by substituting EXAMPLE 3J for EXAMPLE 1E and EXAMPLE 366C for EXAMPLE 1F in EXAMPLE 1G. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.67 (s, 1 H), 11.37 (m, 1 H), 8.59 (m, 2 H), 8.04 (d, 1 H), 7.80 (d, 1 H), 7.51 (m, 3 H), 7.34 (d, 2 H), 7.10 (d, 1 H), 7.04 (d, 2 H), 6.68 (dd, 1 H), 6.39 (m, 1 H), 6.19 (s, 1 H), 3.07 (m, 4 H), 2.75 (m, 2 H), 2.17 (m, 7 H), 1.76 (m, 9 H), 1.32 (m, 9 H), 0.92 (s, 6 H).

EXAMPLE 367

{[4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)amino}methyl pivalate

[1397] This example was prepared by substituting chloromethyl pivalate for chloromethyl butyrate in EXAMPLE 368. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.72 (s, 1H), 8.43 (d, 1H), 8.22 (dd, 1H), 8.01 (d, 1H), 7.55 (m, 3H), 7.36 (m, 3H), 7.03 (d, 2H), 6.68 (dd, 1H), 6.41 (m, 1H), 6.17 (d, 1H), 5.83 (s, 2H), 4.40 (d, 2H), 3.78 (m, 2H), 3.59 (m, 2H), 3.08 (br m, 4H), 2.73 (br s, 2H), 2.18 (br m, 6H), 1.96 (s, 2H), 1.84 (m, 4H), 1.39 (m, 2H), 1.00 (s, 9H), 0.92 (s, 6H).

EXAMPLE 368

{[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)amino}methyl butyrate

[1398] EXAMPLE 37E (500 mg) was dissolved in acetonitrile (3.7 mL) and chloromethyl butyrate (77 mg) and Hunig's base (73 mg) were added. The reaction was heated under reflux for one day. After cooling and dilution with dimethylsulfoxide (4 mL) the reaction was purified by preparative HPLC using a 250 x 50 mm C18 column and eluting with 20-100% CH₃CN vs. 0.1% trifluoroacetic acid in water, giving the product as a trifluoroacetate salt. The trifluoroacetic acid salt was dissolved in dichloromethane (6 ml) and washed with 50% aqueous NaHCO₃. The organic layer was dried over anhydrous Na₂SO₄, filtered, and concentrated to provide the title compound. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.72 (s, 1H), 8.43 (d, 1H), 8.22 (dd, 1H), 8.01 (d, 1H), 7.55 (m, 3H), 7.36 (m, 3H), 7.03 (d, 2H), 6.68 (dd, 1H), 6.41 (m, 1H), 6.17 (d, 1H), 5.83 (s, 2H), 4.40 (d, 2H), 3.78 (m, 2H), 3.59 (m, 2H), 3.08 (br m, 4H), 2.73 (br s, 2H), 2.18 (m, 8H), 1.96 (s, 2H), 1.84 (m, 4H), 1.39 (m, 4H), 0.92 (s, 6H), 0.75 (t, 3H).

EXAMPLE 369

4-[4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]({²H₈)piperazin-1-yl]-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 369A

methyl 4-[[2,2,3,3,5,5,6,6-²H₈]piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoate

[1399] Into a 40 mL vial were added EXAMPLE 3H (1.55 g) and piperazine-d₈ (2.040 g) in dimethylsulfoxide (13 mL). The solution was heated to 85°C for 2.5 hours, and was then allowed to cool to room temperature overnight. The mixture was transferred to a 120 mL flask and was cooled to 5-10 °C. Dichloromethane (30 mL) was added, then water (10 mL) was added via syringe over 5 minutes maintaining temp at no more than 15°C. The layers were separated and the organic layer was washed with water (4x 10-15 mL) until pH of aqueous layer was 8-9. The organic layer was filtered through Na₂SO₄ and rinsed with dichloromethane (5 mL), and concentrated to provide the title compound.

EXAMPLE 369B

methyl 4-[4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]({²H₈)piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoate

[1400] In a 100 mL round-bottomed flask, EXAMPLE 369A (3.4 g), EXAMPLE 290B (1.321 g) and dichloromethane (3 mL) were added to a 100 mL round bottom flask at room temperature. To a separate 50 mL 3 neck round bottom flask, sodium triacetoxymethylborohydride (1.330 g) and dichloromethane (12 mL) were added to give a slurry. After cooling the 50 mL round bottom flask to 18-20 °C, the piperazine adduct/aldehyde solution was added via syringe over 5 minutes. The triacetoxymethylborohydride gradually dissolved to give a clear solution after ~5 minutes. After an additional 10 minutes, the solution became hazy. After 16 hours, the reaction was cooled to 5-10 °C. Saturated aqueous NaHCO₃ (12 mL) was added over 5 minutes maintaining the temperature at no more than 10 °C. The layers were separated and the organic layer was washed with saturated aqueous NaHCO₃, and 10% NaCl (12 mL), and then filtered through Na₂SO₄ and rinsed with dichloromethane (4 mL). The solution was concentrated on a rotovap, and chase concentrated with methanol (40 mL). The resulting solution was cooled to 5-10 °C, and the product precipitated. The solution was mixed at room temperature for 30 minutes, then filtered and rinsed with methanol (5 mL), and the product was air dried.

EXAMPLE 369C

4-[4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]({²H₈)piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoic acid

[1401] The title compound was prepared by substituting EXAMPLE 369B for EXAMPLE 15G in EXAMPLE 15H.

EXAMPLE 369D

4-[4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}(²H₈)piperazin-1-yl]-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

[1402] To a mixture of EXAMPLE 369C (2.0g), EXAMPLE 1F (1.1g) and N,N-dimethylpyridin-4-amine (0.7 g) in dichloromethane (20 mL) was added 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (0.8 g). The reaction mixture was stirred at room temperature overnight. The reaction was quenched with N,N-dimethylethane-1,2-diamine (0.6g) and stirred at room temperature for 3 hours. The mixture was extracted with 20% aqueous acetic acid and washed with 5% aqueous NaCl. Methanol (2 mL) and ethyl acetate (18 mL) were added and the precipitate was collected by filtration to provide the title compound. ¹H NMR (400 MHz, dimethylsulfoxide-d₆) δ 11.71 (s, 1H), 11.37 (s, br, 1H), 8.60 (t, 1H), 8.55 (d, 1H), 8.04 (d, 1H), 7.80 (dd, 1H), 7.47 - 7.54 (m, 3H), 7.31 - 7.34 (m, 2H), 7.09 (d, 1H), 7.01 - 7.03 (m, 2H), 6.67 (dd, 1H), 6.39 (dd, 1H), 6.19 (d, 1H), 3.83 (dd, 2H), 3.21 - 3.30 (m, 4H), 3.00 - 3.10 (s, 4H), 2.75 (s, 2H), 2.05 - 2.24 (m, 6H), 1.95 (s, 2H), 1.80 - 1.93 (m, 1H), 1.55 - 1.64 (m, 2H), 1.37 (t, 2H), 1.18 - 1.31 (m, 2H), 0.90 (s, 6H).

EXAMPLE 370

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([3-(tetrahydro-2H-pyran-4-ylmethyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-6-yl]sulfonyl)benzamide

EXAMPLE 370A

5-amino-6-((tetrahydro-2H-pyran-4-yl)methylamino)pyridine-3-sulfonamide

[1403] A mixture of EXAMPLE 364E (0.16 g) and 5% palladium on carbon (0.025 g) in ethanol (5 mL) was treated with a balloon of hydrogen. The reaction mixture was stirred overnight. The solid was filtered off. The filtrate was concentrated. The residue was purified by flash chromatography on silica gel to give the title compound.

EXAMPLE 370B

3-((tetrahydro-2H-pyran-4-yl)methyl)-3H-[1,2,3]triazolo[4,5-b]pyridine-6-sulfonamide

[1404] EXAMPLE 370A (0.085 g) in water (10 mL) was treated with concentrated H₂SO₄ (0.5 mL). The solution was cooled to 0 °C. To this solution was added NaNO₂ (0.023 g) in water (1 mL) dropwise. The solution was stirred for 1 hour at 0 °C. The reaction mixture was poured into saturated NaHCO₃ solution, and extracted with ethyl acetate three times. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated to give the title compound.

EXAMPLE 370C

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-N-3-((tetrahydro-2H-pyran-4-yl)methyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-6-ylsulfonyl)benzamide

[1405] This example was prepared by substituting EXAMPLE 370B for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.60 (s, 1H), 9.11 (s, 1H), 8.92 (d, 1H), 7.96 (d, 1H), 7.55 (d, 1H), 7.45-7.46 (m, 1H), 7.42 (s, 1H), 7.36 (d, 2H), 7.05 (d, 2H), 6.66 (dd, 1H), 6.32 (s, 1H), 6.22 (s, 1H), 4.63 (d, 2H), 3.80 (dd, 2H), 3.21-3.30 (m, 2H), 3.16 (s, 4H), 2.83 (s, 2H), 2.19-2.29 (m, 6H), 1.97 (s, 2H), 1.33-1.41 (m, 6H), 0.93 (s, 2H).

EXAMPLE 371

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-([6-([trans-4-hydroxy-4-methylcyclohexyl)methyl]amino)-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 371A

6-((trans-4-hydroxy-4-methylcyclohexyl)methylamino)-5-nitropyridine-3-sulfonamide

[1406] This example was prepared by substituting EXAMPLE 364D for 4-fluoro-3-nitrobenzenesulfonamide and EX-

AMPLE 376B for EXAMPLE 138C in EXAMPLE 138D. The title compound was isolated by reverse phase Gilson Prep HPLC system with a Phenomenex prep column (Luna, 5 μ , C18(2), 250X21.20 mm, 5 Å) eluting with 20-80% acetonitrile in water with 0.1% TFA.

5 EXAMPLE 371B

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(6-{[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino}-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

10 **[1407]** The title compound was prepared by substituting EXAMPLE 371A for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (300 MHz, dimethylsulfoxide-*d*₆) δ 11.63 (s, 1H), 11.53 - 10.99 (m, 1H), 8.91 (s, 1H), 8.71 (dd, 2H), 8.01 (d, 1H), 7.61 - 7.44 (m, 3H), 7.44 - 7.28 (m, 2H), 7.12 - 6.97 (m, 2H), 6.76 - 6.61 (m, 1H), 6.36 (dd, 1H), 6.21 (d, 1H), 3.92 (s, 1H), 3.48 (t, 2H), 3.10 (s, 4H), 2.83 (s, 2H), 2.24 (dd, 6H), 1.96 (s, 2H), 1.37 (ddd, 11H), 1.07 (s, 3H), 0.93 (s, 6H).

15 EXAMPLE 372

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(5-cyano-6-{[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy}pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

20 EXAMPLE 372A

ethyl 4-fluoro-1-(oxetan-3-yl)piperidine-4-carboxylate

25 **[1408]** To 1-*tert*-butyl 4-ethyl 4-fluoropiperidine-1,4-dicarboxylate (1.00 g) was added HCl (4.0M in dioxane, 4.54 mL). After 1 hour the reaction was concentrated and dried under high vacuum. The resulting solid was dissolved in dichloromethane (5 ml) and treated with sodium triacetoxyborohydride (1.155 g) and oxetan-3-one (0.262 g) and stirred overnight. The reaction was quenched with saturated NaHCO₃ solution (20 mL) and extracted into dichloromethane (2 x 25 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated. Silica gel chromatography (Reveleris 80 g) eluting with a gradient of 0.5% to 3.75% methanol/dichloromethane over 40 minutes (flow = 30 mL/minute) gave the title compound.

EXAMPLE 372B

(4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methanol

35 **[1409]** To a solution of EXAMPLE 372A (0.59 g) in tetrahydrofuran (5 mL) was added lithium aluminum hydride (1.80 mL) at 0°C. The reaction was removed from the ice bath and allowed to warm to room temperature. The reaction was quenched by the dropwise addition of 0.6 mL of water followed by 0.2 ml of 2N aqueous NaOH. The reaction was filtered through diatomaceous earth and rinsed with ethyl acetate (50 mL). The organics were concentrated and loaded onto silica gel (Reveleris 40 g) and eluted using a gradient of 0.75% to 7.5% methanol/dichloromethane over 30 minute (flow = 40 mL/minutes) to give the title compound.

EXAMPLE 372C

45 5-bromo-6-((4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methoxy)pyridine-3-sulfonamide

[1410] This example was prepared by substituting EXAMPLE 372B for (tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 36A for 4-fluoro-3-nitrobenzenesulfonamide in EXAMPLE 24A.

50 EXAMPLE 372D

5-cyano-6-((4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methoxy)pyridine-3-sulfonamide

[1411] This example was prepared by substituting EXAMPLE 372C for EXAMPLE 36B in EXAMPLE 36C.

55

EXAMPLE 372E

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)-N-(5-cyano-6-((4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methoxy)pyridin-3-ylsulfonyl)benzamide

[1412] The title compound was prepared by substituting EXAMPLE 372D for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ 11.58 (s, 1H), 8.71 (s, 1H), 8.52 (s, 1H), 7.96 (d, 1H), 7.57 (d, 1H), 7.48 - 7.30 (m, 4H), 7.06 (d, 2H), 6.68 (d, 1H), 6.37 - 6.22 (m, 2H), 4.65 - 4.40 (m, 6H), 3.58 (s, 1H), 3.12 (s, 6H), 2.84 - 2.59 (m, 4H), 2.17 (s, 6H), 1.96 (d, 6H), 1.41 (s, 2H), 0.93 (s, 6H).

EXAMPLE 373

N-(4-([4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl)-2-nitrophenyl)morpholine-4-carboxamide

EXAMPLE 373A

morpholine-4-carboxamide

[1413] A solution of morpholine-4-carbonyl chloride (2.0 g) in methanol (10 mL) and 7 N NH₃ in methanol (5 mL) was stirred at 45 °C overnight. The mixture was concentrated to give a solid, which was dried under vacuum.

EXAMPLE 373B

N-(2-nitro-4-sulfamoylphenyl)morpholine-4-carboxamide

[1414] This example was prepared by substituting EXAMPLE 373A for (tetrahydro-2H-pyran-4-yl)methanol in EXAMPLE 24A.

EXAMPLE 373C

N-(4-(N-(2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoyl)sulfamoyl)-2-nitrophenyl)morpholine-4-carboxamide

[1415] This example was prepared by substituting EXAMPLE 373B for EXAMPLE 130C in EXAMPLE 130D. ¹H NMR (500MHz, pyridine-d₅) δ 13.02 (s, 1H), 10.41 (s, 1H), 9.27 (d, 1H), 8.81 (d, 1H), 8.50 (dd, 1H), 8.40 (d, 1H), 8.09 (d, 1H), 7.65 (m, 2H), 7.44 (d, 2H), 7.07 (d, 2H), 6.76 (dd, 1H), 6.54 (d, 1H), 6.48 (m, 1H), 3.67 (m, 4H), 3.58 (m, 4H), 3.07 (m, 4H), 2.77 (s, 2H), 2.26 (t, 2H), 2.14 (m, 4H), 1.97 (s, 2H), 1.39 (t, 2H), 0.94 (s, 6H).

EXAMPLE 374

4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-(methoxymethyl)cyclohexyl]methyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 374A

(4,4-diethoxycyclohexyl)methanol

[1416] Ethyl 4, 4-diethoxycyclohexanecarboxylate (6.67 g) synthesized according to a literature procedure (European Journal of Organic Chemistry, 2008, 5, 895) in tetrahydrofuran (60 mL) was treated with 2 M lithium aluminum hydride in tetrahydrofuran (14.5 mL) at 0 °C for 1 hour. Water (3 mL) was slowly added to quench the reaction. The precipitates were filtered off and washed with ethyl acetate. The filtrate was dried over Na₂SO₄, filtered, and concentrated to provide the title compound.

EXAMPLE 374B

1,1-diethoxy-4-(methoxymethyl)cyclohexane

5 [1417] EXAMPLE 374A (665 mg) in tetrahydrofuran (20 mL) was treated with NaH (394 mg) for 30 minutes and then CH_3I (0.267 mL) was slowly added. The resulting mixture was stirred overnight and the reaction was quenched with a few drops of water. The mixture was concentrated and the residue was suspended in water and extracted with dichloromethane. The organic layer was dried over Na_2SO_4 , filtered and concentrated. The residue was purified by flash chromatography, and was eluted with 0-15% ethyl acetate in dichloromethane to provide the title compound.

10

EXAMPLE 374C

4-(methoxymethyl)cyclohexanone

15 [1418] EXAMPLE 374B (2.2 g) in a mixture of water (3 mL) and acetic acid (12 mL) was heated at 65°C for 2 hours. The reaction mixture was concentrated. The residue was mixed with water and saturated aqueous NaHCO_3 and extracted with dichloromethane. The dichloromethane layer was dried over Na_2SO_4 , filtered, and concentrated to provide the title compound.

20 EXAMPLE 374D

4-(methoxymethyl)cyclohexanecarbonitrile

25 [1419] To a cold (-10°C) solution of EXAMPLE 374C (1.18 g) and toluenesulfonylmethyl isocyanide (2.268 g) in dimethoxyethane (3 mL) and absolute ethanol (0.1 mL) was added (in small portions) potassium tert-butoxide (2.235 g). The reaction mixture was continued to stir at <5°C for 30 minutes, warmed to room temperature, heated at 35°C for 30 minutes and then at room temperature for 2 hours. The reaction mixture was concentrated and the residue was dissolved in water-brine, and extracted with dichloromethane. The dichloromethane layer was purified by flash chromatography, and was eluted with 5% ethyl acetate in dichloromethane to provide the title compound.

30

EXAMPLE 374E

(4-(methoxymethyl)cyclohexyl)methanamine

35 [1420] To a solution of EXAMPLE 374D (460 mg) in tetrahydrofuran (15 mL) was added 2M lithium aluminum hydride in tetrahydrofuran (2.252 mL) slowly. The reaction mixture was stirred at room temperature for 1 hour, refluxed for 1 hour and cooled. 2 mL of 2M aqueous NaOH and water (5 mL) was added. The solid was filtered off and washed with ether. The filtrate was concentrated. The residue was mixed with dichloromethane (50 mL) and the resulting mixture was dried over Na_2SO_4 and concentrated to provide the title compound.

40

EXAMPLE 374F

4-((4-(methoxymethyl)cyclohexyl)methylamino)-3-nitrobenzenesulfonamide

45 [1421] EXAMPLE 374E (450 mg) and 4-fluoro-3-nitrobenzenesulfonamide (693 mg) in tetrahydrofuran (10 mL) were stirred overnight. The reaction mixture was concentrated and the residue was suspended in a mixture of CH_3CN , methanol and water. The precipitates were collected, washed with water and dried to give the title compound.

EXAMPLE 374G

50

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)-N-(4-(4-(methoxymethyl)cyclohexyl)methylamino)-3-nitrophenylsulfonyl)benzamide

55 [1422] The title compound was prepared as described in EXAMPLE 11D using EXAMPLE 374F in place of EXAMPLE 11B. ^1H NMR (400 MHz, dimethylsulfoxide- d_6) δ 11.69 (s, 1H), 11.40 (s, 1H), 8.53 - 8.61 (m, 2H), 8.04 (d, 1H), 7.77 - 7.82 (m, 1H), 7.47 - 7.55 (m, 3H), 7.34 (d, 2H), 7.02 - 7.09 (m, 3H), 6.68 (dd, 1H), 6.39 (dd, 1H), 6.19 (d, 1H), 3.18 - 3.27 (m, 5H), 3.04 - 3.14 (m, 5H), 2.75 (s, 2H), 2.11 - 2.24 (m, 6H), 1.95 (s, 2H), 1.69 - 1.84 (m, 3H), 1.33 - 1.63 (m, 7H), 0.84 - 1.05 (m, 9H).

EXAMPLE 375

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(5-chloro-6-[[1-(1,3-thiazol-2-yl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 375A

methyl 1-(thiazol-2-yl)piperidine-4-carboxylate

[1423] A mixture of methyl piperidine-4-carboxylate (2.045 g), 2-bromothiazole (1.64 g), and Cs₇CO₃ (5.86 g) in dimethylformamide (15 mL) was heated at 100 °C overnight. After it cooled to room temperature, the reaction mixture was partitioned between water and ethyl acetate. The organic layer was separated, and the aqueous layer was extracted with additional ethyl acetate three times. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography on silica gel to give the title compound.

EXAMPLE 375B

(1-(thiazol-2-yl)piperidin-4-yl)methanol

[1424] This example was prepared by substituting EXAMPLE 375A for EXAMPLE 339A in EXAMPLE 339B.

EXAMPLE 375C

5-chloro-6-((1-(thiazol-2-yl)piperidin-4-yl)methoxy)pyridine-3-sulfonamide

[1425] This example was prepared by substituting EXAMPLE 375B for (tetrahydro-2H-pyran-4-yl)methanol and EXAMPLE 40A for EXAMPLE 36A in EXAMPLE 36B.

EXAMPLE 375D

2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-(5-chloro-6-((1-(thiazol-2-yl)piperidin-4-yl)methoxy)pyridin-3-ylsulfonyl)-4-(4-(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enyl)methyl)piperazin-1-yl)benzamide

[1426] This example was prepared by substituting EXAMPLE 375C for EXAMPLE 11B in EXAMPLE 11D. ¹H NMR (500MHz, dimethylsulfoxide-d₆) δ 11.65 (s, 1H), 8.49 (s, 1H), 8.17 (s, 1H), 8.01 (d, 1H), 7.54 (d, 1H), 7.48-7.49 (m, 2H), 7.35 (d, 2H), 7.14 (d, 1H), 7.05 (d, 2H), 6.80 (d, 1H), 6.67 (dd, 1H), 6.38 (dd, 1H), 6.21 (d, 1H), 4.28 (d, 2H), 3.92 (d, 2H), 2.98-3.10 (m, 6H), 2.86 (s, 2H), 2.30 (m, 4H), 2.03-2.15 (m, 3H), 1.96 (s, 2H), 1.96 (s, 2H), 1.82-1.86 (m, 2H), 1.33-1.44 (m, 4H), 0.93 (s, 6H).

EXAMPLE 376

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(6-[[[cis-4-hydroxy-4-methylcyclohexyl)methyl]amino]-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

EXAMPLE 376A

tert-butyl (4-hydroxy-4-methylcyclohexyl)methylcarbamate

[1427] A solution of tert-butyl (4-oxocyclohexyl)methylcarbamate (1.00 g) was dissolved in tetrahydrofuran (20 mL) and cooled to -78°C. Methylmagnesium bromide (4.40 mL) was added dropwise. The reaction was stirred for 2 hours at -78°C then allowed to warm to 0°C and stirred for 30 minutes. The resulting suspension was quenched with water (10 mL), diluted with ether (50 mL), washed with ammonium chloride (25 mL), washed with brine (25 mL), dried over magnesium sulfate, filtered, and concentrated. Silica gel chromatography (Reveleris 80 g) eluting using a gradient of 5% to 50% ethyl acetate/dichloromethane over 30 minutes (flow = 60 mL/min) gave the title compound as a ~2:1 mixture of cis and trans isomers.

EXAMPLE 376B

4-(aminomethyl)-1-methylcyclohexanol

- 5 **[1428]** To a solution of EXAMPLE 376A (0.75 g) in dichloromethane (3 mL) was added a few drops of water followed by trifluoroacetic acid (1.19 mL) and the reaction stirred at room temperature. After stirring for 2h added additional trifluoroacetic acid (0.5 mL). After an additional 4 h the reaction was concentrated and dried under high vacuum. The resulting oily solid was triturated with diethyl ether with sonication. Filtration and washing with diethyl ether gave the title compound as a trifluoroacetic acid salt and a mixture of cis and trans isomers.

10

EXAMPLE 376C

6-((cis-4-hydroxy-4-methylcyclohexyl)methylamino)-5-nitropyridine-3-sulfonamide

- 15 **[1429]** This example was prepared by substituting EXAMPLE 364D for 4-fluoro-3-nitrobenzenesulfonamide and EXAMPLE 376B for (4-fluorotetrahydro-2H-pyran-4-yl)methanamine in EXAMPLE 138D. The title compound was isolated by reverse phase Gilson Prep HPLC system with a Phenomenex prep column (Luna, 5 μ , C18(2), 250X21.20 mm, 5 $\text{\AA}$) eluting with 20-80% acetonitrile in water with 0.1% TFA.

20 EXAMPLE 376D

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(6-{[(cis-4-hydroxy-4-methylcyclohexyl)methyl]amino}-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- 25 **[1430]** This example was prepared by substituting EXAMPLE 376C for EXAMPLE 11B in EXAMPLE 11D. ^1H NMR (500MHz, dimethylsulfoxide- d_6) δ 11.64 (s, 1H), 8.91 (s, 1H), 8.72 (d, 1H), 8.70 (d, 1H), 8.01 (d, 1H), 7.47-7.54 (m, 3H), 7.35 (d, 2H), 7.04 (d, 2H), 6.68 (dd, 1H), 6.36 (dd, 1H), 6.21 (d, 1H), 3.93 (s, 1H), 3.48 (t, 2H), 3.10 (s, 4H), 2.83 (s, 2H), 2.15-2.33 (m, 6H), 1.96 (s, 1H), 1.34-1.59 (m, 9H), 1.17-1.24 (m, 2H), 1.07 (s, 2H), 0.92 (s, 6H).

30 EXAMPLE 377

4-(4-{[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[(4-{[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

- 35 **[1431]** The title compound was prepared by substituting EXAMPLE 378D for EXAMPLE 1E and EXAMPLE 337M for EXAMPLE 1F in EXAMPLE 1G. ^1H NMR (500 MHz, pyridine- d_5) δ 13.07 (s, 1H), 9.31 (d, 1H), 8.68 (t, 1H), 8.44 (d, 1H), 8.37 (dd, 1H), 8.10 (d, 1H), 7.68 (m, 1H), 7.66 (d, 1H), 7.41 (m, 2H), 7.09 (m, 2H), 6.92 (d, 1H), 6.74 (dd, 1H), 6.52 (d, 1H), 6.50 (dd, 1H), 3.20 (m, 5H), 3.06 (t, 4H), 2.77 (m, 2H), 2.57 (d, 1H), 2.49 (m, 1H), 2.17 (m, 6H), 1.86 (m, 5H), 1.69 (m, 4H), 1.40 (s, 3H), 1.23 (m, 5H).

40

EXAMPLE 378

4-(4-{[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide

45

EXAMPLE 378A

2-chloro-5-methoxy-5-methylcyclohex-1-enecarbaldehyde

- 50 **[1432]** Dimethylformamide (1.298 mL) in dichloromethane (2.0 mL) at -10°C was treated dropwise with POCl_3 (1.426 mL) to give a colorless solution. The mixture was stirred 5 minutes and then warmed to room temperature and stirred 30 minutes. The solution was cooled to -10°C , treated dropwise with a solution of 4-methoxy-4-methylcyclohexanone (1.74 g) in dichloromethane (2.5 mL), and stirred for 4 hours at ambient temperature. The reaction mixture was poured over a mixture of ice and 25% aqueous sodium acetate solution. After the ice melted, the reaction mixture was poured into a separatory funnel and extracted with diethyl ether (4 x 125 mL). The diethyl ether extracts were washed with NaHCO_3 solution and brine, dried (MgSO_4), filtered and concentrated. The concentrate was chromatographed on silica gel with 0 to 5% ethyl acetate in hexanes as the eluent.
- 55

EXAMPLE 378B

2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-enecarbaldehyde

5 **[1433]** EXAMPLE 378A (1.55 g), 4-chlorophenylboronic acid (1.542 g), PdOAc₂ (0.055 g), K₂CO₃ (2.84 g) and tetrabutylammonium bromide (2.65 g) were combined in a 50-mL round-bottomed flask equipped with a magnetic stir bar. Water (9.13 mL) was added. The vial was flushed with nitrogen, capped and stirred at 45 °C for 14 hours. The reaction mixture was cooled to room temperature and partitioned between brine and diethyl ether. The organic layer was washed with
10 brine, dried (MgSO₄), filtered through a plug of celite, concentrated and chromatographed on silica gel with 5 to 20% ethyl acetate in hexanes as the eluent.

EXAMPLE 378C

15 methyl 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoate

[1434] The title compound was prepared by substituting EXAMPLE 378B for 4'-chlorobiphenyl-2-carboxaldehyde and EXAMPLE 15F for tert-butyl piperazine-1-carboxylate in EXAMPLE 1A except that a small amount of DMSO was added to the reaction mixture.

20

EXAMPLE 378D

25 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-enyl)methyl)piperazin-1-yl)benzoic acid

25

[1435] The title compound was prepared by substituting EXAMPLE 378C for EXAMPLE 15G in EXAMPLE 15H.

EXAMPLE 378E

30 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-enyl)methyl)piperazin-1-yl)-N-(3-nitro-4-((tetrahydro-2H-pyran-4-yl)methylamino)phenylsulfonyl)benzamide


[1436] The title compound was prepared by substituting EXAMPLE 378D for EXAMPLE 1E in EXAMPLE 1G. ¹H NMR (500 MHz, pyridine-d₅) δ 13.07 (s, 1H), 9.31 (d, 1H), 8.68 (t, 1H), 8.43 (d, 1H), 8.37 (dd, 1H), 8.09 (d, 1H), 7.68 (m, 1H),
35 7.66 (d, 1H), 7.41 (m, 2H), 7.09 (m, 2H), 6.90 (d, 1H), 6.74 (dd, 1H), 6.52 (d, 1H), 6.50 (dd, 1H), 3.97 (dd, 2H), 3.30 (td, 2H), 3.21 (s, 3H), 3.15 (m, 2H), 3.06 (t, 4H), 2.77 (m, 2H), 2.57 (d, 1H), 2.50 (m, 1H), 2.16 (m, 6H), 1.81 (m, 2H), 1.63 (m, 1H), 1.57 (dd, 2H), 1.32 (m, 2H), 1.21 (s, 3H).

[1437] Preferred embodiments of the present invention are described below and are referred to as embodiments E1 to E32.

40

E1. A compound having Formula (I)

45



50

(I),

or a therapeutically acceptable salt thereof, wherein

55 A¹ is N or C(A²);
A² is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂,

$C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$;
 B^1 is H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$,
5 $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$;
 D^1 is H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$,
10 $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$;
 E^1 is H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$,
15 $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$; and
 Y^1 is H, CN, NO_2 , $C(O)OH$, F, Cl, Br, I, CF_3 , OCF_3 , CF_2CF_3 , OCF_2CF_3 , R^{17} , OR^{17} , $C(O)R^{17}$, $C(O)OR^{17}$, SR^{17} , SO_2R^{17} , NH_2 , NHR^{17} , $N(R^{17})_2$, $NHC(O)R^{17}$, $C(O)NH_2$, $C(O)NHR^{17}$, $C(O)N(R^{17})_2$, $NHS(O)R^{17}$ or $NHSO_2R^{17}$; or
20 E^1 and Y^1 , together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and
 A^2 , B^1 , and D^1 are independently selected H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2NCR^1$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$,
30 F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$; or
 Y^1 and B^1 , together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and
 A^2 , D^1 , and E^1 are independently selected H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$,
35 $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$; or
 A^2 and B^1 , together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and
40 D^1 , E^1 , and Y^1 are independently selected H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$,
45 $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$; or
 A^2 and D^1 , together with the atoms to which they are attached, are benzene, naphthalene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and
 B^1 , E^1 , and Y^1 are independently selected H, R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOH$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$,
50 $N(CH_3)SO_2N(CH_3)R^1$, F, Cl, Br, I, CN, NO_2 , N_3 , OH, $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$;
 G^1 is H, or $C(O)OR$;
R is alkyl;

R¹ is R², R³, R⁴ or R⁵;

R^{1A} is cycloalkyl, cycloalkenyl or cycloalkynyl;

R² is phenyl, which is unfused or fused with R^{2A}; R^{2A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

5 R³ is heteroaryl, which is unfused or fused with R^{3A}; R^{3A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁴ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{4A}; R^{4A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

10 R⁵ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶, NC(R^{6A})(R^{6B}), R⁷, OR⁷, SR⁷, S(O)R⁷, SO₂R⁷, NHR⁷, N(R⁷)₂, C(O)R⁷, C(O)NH₂, C(O)NHR⁷, C(O)N(R⁷)₂, NHC(O)R⁷, NR⁷C(O)R⁷, NHSO₂R⁷, NHC(O)OR⁷, SO₂NH₂, SO₂NHR⁷, SO₂N(R⁷)₂, NHC(O)NH₂, NHC(O)NHR⁷, NHC(O)CH(CH₃)NHC(O)CH(CH₃)NH₂, NHC(O)CH(CH₃)NHC(O)CH(CH₃)NHR⁷, OH, (O), C(O)OH, N₃, CN, NH₂, CF₃, CF₂CF₃, F, Cl, Br or I;

15 R⁶ is C₂-C₅-spiroalkyl, each of which is unsubstituted or substituted with OH, (O), N₃, CN, CF₃, CF₂CF₃, F, Cl, Br, I, NH₂, NH(CH₃) or N(CH₃)₂;

R^{6A} and R^{6B} are independently selected alkyl or, together with the N to which they are attached, R^{6C};

R^{6C} is aziridin-1-yl, azetidin-1-yl, pyrrolidin-1-yl or piperidin-1-yl, each having one CH₂ moiety unreplaced or replaced with O, C(O), CNOH, CNOCH₃, S, S(O), SO₂ or NH;

R⁷ is R⁸, R⁹, R¹⁰ or R¹¹;

20 R⁸ is phenyl, which is unfused or fused with R^{8A}; R^{8A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁹ is heteroaryl, which is unfused or fused with R^{9A}; R^{9A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

25 R¹⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{10A}; R^{10A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

30 R¹¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R¹², OR¹², SR¹², S(O)R¹², SO₂R¹², C(O)R¹², CO(O)R¹², OC(O)R¹², OC(O)OR¹², NH₂, NHR¹², N(R¹²)₂, NHC(O)R¹², NR¹²C(O)R¹², NHS(O)₂R¹², NR¹²S(O)₂R¹², NHC(O)OR¹², NR¹²C(O)OR¹², NHC(O)NH₂, NHC(O)NHR¹², NHC(O)N(R¹²)₂, NR¹²C(O)NHR¹², NR¹²C(O)N(R¹²)₂, C(O)NH₂, C(O)NHR¹², C(O)N(R¹²)₂, C(O)NHOH, C(O)NHOR¹², C(O)NHSO₂R¹², C(O)NR¹²SO₂R¹², SO₂NH₂, SO₂NHR¹², SO₂N(R¹²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR¹², C(N)N(R¹²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R¹² is R¹³, R¹⁴, R¹⁵ or R¹⁶;

35 R¹³ is phenyl, which is unfused or fused with R^{13A}; R^{13A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R¹⁴ is heteroaryl, which is unfused or fused with R^{14A}; R^{14A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R¹⁵ is cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene, each of which is unfused or fused with R^{15A}; R^{15A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

40 R¹⁶ is alkyl, alkenyl or alkynyl;

R¹⁷ is R¹⁸, R¹⁹, R²⁰ or R²¹;

R¹⁸ is phenyl, which is unfused or fused with R^{18A}; R^{18A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

45 R¹⁹ is heteroaryl, which is unfused or fused with R^{19A}; R^{19A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl each of which is unfused or fused with R^{20A}; R^{20A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

50 R²¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R²², OR²², SR²², S(O)R²², SO₂R²², C(O)R²², CO(O)R²², OC(O)R²², OC(O)OR²², NH₂, NHR²², N(R²²)₂, NHC(O)R²², NR²²C(O)R²², NHS(O)₂R²², NR²²S(O)₂R²², NHC(O)OR²², NR²²C(O)OR²², NHC(O)NH₂, NHC(O)NHR²², NHC(O)N(R²²)₂, NR²²C(O)NHR²², NR²²C(O)N(R²²)₂, C(O)NH₂, C(O)NHR²², C(O)N(R²²)₂, C(O)NHOH, C(O)NHOR²², C(O)NHSO₂R²², C(O)NR²²SO₂R²², SO₂NH₂, SO₂NHR²², SO₂N(R²²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR²², C(N)N(R²²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

55 R²² is R²³, R²⁴ or R²⁵;

R²³ is phenyl, which is unfused or fused with R^{23A}; R^{23A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁴ is heteroarene, which is unfused or fused with R^{24A}; R^{24A} is benzene, heteroarene, cycloalkane, cycloalkene,

heterocycloalkane or heterocycloalkene;

R²⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{25A}; R^{25A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

Z¹ is R²⁶ or R²⁷;

Z² is R²⁸, R²⁹ or R³⁰;

Z^{1A} and Z^{2A} are both absent or are taken together to form CH₂, CH₂CH₂ or Z^{12A};

Z^{12A} is C₂-C₆-alkylene having one or two CH₂ moieties replaced by NH, N(CH₃), S, S(O) or SO₂;

L¹ is a R³⁷, OR³⁷, SR³⁷, S(O)R³⁷, SO₂R³⁷, C(O)R³⁷, CO(O)R³⁷, OC(O)R³⁷, OC(O)OR³⁷, NHR³⁷, C(O)NH, C(O)NR³⁷, C(O)NHOR³⁷, C(O)NHSO₂R³⁷, SO₂NH, SO₂NHR³⁷, C(N)NH, C(N)NHR³⁷;

R²⁶ is phenylene, which is unfused or fused with R^{26A}; R^{26A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁷ is heteroarylene, which is unfused or fused with R^{27A}; R^{27A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁸ is phenylene, which is unfused or fused with R^{28A}; R^{28A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁹ is heteroarylene, which is unfused or fused with R^{29A}; R^{29A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁰ is cycloalkylene, cycloalkenylene, heterocycloalkylene or heterocycloalkenylene, each of which is unfused or fused with R^{30A}; R^{30A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁷ is a bond or R^{37A};

R^{37A} is alkylene, alkenylene, or alkynylene, each of which is unsubstituted or substituted with one or two or three independently selected R^{37B}, OR^{37B}, SR^{37B}, S(O)R^{37B}, SO₂R^{37B}, C(O)R^{37B}, CO(O)R^{37B}, OC(O)R^{37B}, OC(O)OR^{37B}, NH₂, NBR^{37B}, N(R^{37B})₂, NHC(O)R^{37B}, NR^{37B}C(O)R^{37B}, NHS(O)₂R^{37B}, NR^{37B}SO₂R^{37B}, NHC(O)OR^{37B}, NR^{37B}C(O)OR^{37B}, NHC(O)NH₂, NHC(O)NHR^{37B}, NHC(O)N(R^{37B})₂, NR^{37B}C(O)NHR^{37B}, NR^{37B}C(O)N(R^{37B})₂, C(O)NH₂, C(O)NHR^{37B}, C(O)N(R^{37B})₂, C(O)NHOH, C(O)NHOR^{37B}, C(O)NHSO₂R^{37B}, C(O)NR^{37B}SO₂R^{37B}, SO₂NH₂, SO₂NHR^{37B}, SO₂N(R^{37B})₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR^{37B}, C(N)N(R^{37B})₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I substituents;

R^{37B} is alkyl, alkenyl, alkynyl, or R^{37C};

R^{37C} is phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl;

Z³ is R³⁸, R³⁹ or R⁴⁰;

R³⁸ is phenyl, which is unfused or fused with R^{38A}; R^{38A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁹ is heteroaryl, which is unfused or fused with R^{39A}; R^{39A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁴⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{40A}; R^{40A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

wherein the moieties represented by R²⁶ and R²⁷ are substituted (i.e., if Z^{1A} and Z^{2A} are absent) or further substituted (i.e., if Z^{1A} and Z^{2A} are present) with R⁴¹, OR⁴¹, SR⁴¹, S(O)R⁴¹, SO₂R⁴¹, C(O)R⁴¹, CO(O)R⁴¹, OC(O)R⁴¹, OC(O)OR⁴¹, NHR⁴¹, N(R⁴¹)₂, NHC(O)R⁴¹, NR⁴¹C(O)R⁴¹, NHS(O)₂R⁴¹, NR⁴¹SO₂R⁴¹, NHC(O)OR⁴¹, NR⁴¹C(O)OR⁴¹, NHC(O)NHR⁴¹, NHC(O)N(R⁴¹)₂, NR⁴¹C(O)NHR⁴¹, NR⁴¹C(O)N(R⁴¹)₂, C(O)NHR⁴¹, C(O)N(R⁴¹)₂, C(O)NHOR⁴¹, C(O)NHSO₂R⁴¹, C(O)NR⁴¹SO₂R⁴¹, SO₂NHR⁴¹, SO₂N(R⁴¹)₂, C(N)NHR⁴¹, or C(N)N(R⁴¹)₂;

R⁴¹ is heteroaryl, which is fused with R^{43A}; R^{43A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; which is unfused or fused with benzene, heteroarene or R^{43B}; R^{43B} is cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

wherein the cyclic moieties represented by E¹ and Y¹ together, Y¹ and B¹ together, A² and B¹ together, A² and D¹ together, R^{1A}, R², R^{2A}, R³, R^{3A}, R⁴, R^{4A}, R⁶, R^{6C}, R⁸, R^{8A}, R⁹, R^{9A}, R¹⁰, R^{10A}, R¹³, R^{13A}, R¹⁴, R^{14A}, R¹⁵, R^{15A}, R¹⁸, R^{18A}, R¹⁹, R^{19A}, R²⁰, R^{20A}, R²³, R^{23A}, R²⁴, R^{24A}, R²⁵, R^{25A}, R²⁶, R^{26A}, R²⁷, R^{27A}, R²⁸, R^{28A}, R²⁹, R^{29A}, R³⁰, R^{30A}, R^{37B}, R³⁸, R^{38A}, R³⁹, R^{39A}, R⁴⁰, and R^{40A} are independently unsubstituted, further unsubstituted, substituted or further substituted with one or two or three or four or five of independently selected R^{57A}, R⁵⁷, OR⁵⁷, SR⁵⁷, S(O)R⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, OC(O)R⁵⁷, OC(O)OR⁵⁷, NH₂, NHR⁵⁷, N(R⁵⁷)₂, NHC(O)R⁵⁷, NR⁵⁷C(O)R⁵⁷, NHS(O)₂R⁵⁷, NR⁵⁷SO₂R⁵⁷, NHC(O)OR⁵⁷, NR⁵⁷C(O)OR⁵⁷, NHC(O)NH₂, NHC(O)NHR⁵⁷, NHC(O)N(R⁵⁷)₂, NR⁵⁷C(O)NHR⁵⁷, NR⁵⁷C(O)N(R⁵⁷)₂, C(O)NH₂, C(O)NHR⁵⁷, C(O)N(R⁵⁷)₂, C(O)NHOH, C(O)NHOR⁵⁷, C(O)NHSO₂R⁵⁷, C(O)NR⁵⁷SO₂R⁵⁷, SO₂NH₂, SO₂NHR⁵⁷, SO₂N(R⁵⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁵⁷, C(N)N(R⁵⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R^{57A} is spiroalkyl, or spiroheteroalkyl;

R⁵⁷ is R⁵⁸, R⁵⁹, R⁶⁰ or R⁶¹;

R⁵⁸ is phenyl, which is unfused or fused with R^{58A}; R^{58A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

5 R⁵⁹ is heteroaryl, which is unfused or fused with R^{59A}; R^{59A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{60A}; R^{60A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

10 R⁶¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶², OR⁶², SR⁶², S(O)R⁶², SO₂R⁶², C(O)R⁶², CO(O)R⁶², OC(O)R⁶², OC(O)OR⁶², NH₂, NHR⁶², N(R⁶²)₂, NHC(O)R⁶², NR⁶²C(O)R⁶², NHS(O)₂R⁶², NR⁶²S(O)₂R⁶², NHC(O)OR⁶², NR⁶²C(O)OR⁶², NHC(O)NH₂, NHC(O)NHR⁶², NHC(O)N(R⁶²)₂, NR⁶²C(O)NHR⁶², NR⁶²C(O)N(R⁶²)₂, C(O)NH₂, C(O)NHR⁶², C(O)N(R⁶²)₂, C(O)NHOH, C(O)NHOR⁶², C(O)NHSO₂R⁶², C(O)NR⁶²SO₂R⁶², SO₂NH₂, SO₂NHR⁶², SO₂N(R⁶²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶², C(N)N(R⁶²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂,
15 CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R⁶² is R⁶³, R⁶⁴, R⁶⁵ or R⁶⁶;

R⁶³ is phenyl, which is unfused or fused with R^{63A}; R^{63A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

20 R⁶⁴ is heteroaryl, which is unfused or fused with R^{64A}; R^{64A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl, each of which is unfused or fused with R^{65A}; R^{65A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

25 R⁶⁶ is alkyl, alkenyl or alkenyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶⁷, OR⁶⁷, SR⁶⁷, S(O)R⁶⁷, SO₂R⁶⁷, C(O)R⁶⁷, CO(O)R⁶⁷, OC(O)R⁶⁷, OC(O)OR⁶⁷, NH₂, NHR⁶⁷, N(R⁶⁷)₂, NHC(O)R⁶⁷, NR⁶⁷C(O)R⁶⁷, NHS(O)₂R⁶⁷, NR⁶⁷S(O)₂R⁶⁷, NHC(O)OR⁶⁷, NR⁶⁷C(O)OR⁶⁷, NHC(O)NH₂, NHC(O)NHR⁶⁷, NHC(O)N(R⁶⁷)₂, NR⁶⁷C(O)NHR⁶⁷, NR⁶⁷C(O)N(R⁶⁷)₂, C(O)NH₂, C(O)NHR⁶⁷, C(O)N(R⁶⁷)₂, C(O)NHOH, C(O)NHOR⁶⁷, C(O)NHSO₂R⁶⁷, C(O)NR⁶⁷SO₂R⁶⁷, SO₂NH₂, SO₂NHR⁶⁷, SO₂N(R⁶⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁷, C(N)N(R⁶⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂,
30 CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I substituents;

30 R⁶⁷ is alkyl, alkenyl, alkynyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; wherein the cyclic moieties represented by R^{57A}, R⁵⁸, R⁵⁹, R⁶⁰, R⁶³, R⁶⁴, R⁶⁵, and R⁶⁷ are unsubstituted or substituted with one or two or three or four of independently selected R⁶⁸, OR⁶⁸, SR⁶⁸, S(O)R⁶⁸, SO₂R⁶⁸, C(O)R⁶⁸, CO(O)R⁶⁸, OC(O)R⁶⁸, OC(O)OR⁶⁸, NH₂, NHR⁶⁸, N(R⁶⁸)₂, NHC(O)R⁶⁸, NR⁶⁸C(O)R⁶⁸, NHS(O)₂R⁶⁸, NR⁶⁸S(O)₂R⁶⁸, NHC(O)OR⁶⁸, NR⁶⁸C(O)OR⁶⁸, NHC(O)NH₂, NHC(O)NHR⁶⁸, NHC(O)N(R⁶⁸)₂,
35 NR⁶⁸C(O)NHR⁶⁸, NR⁶⁸C(O)N(R⁶⁸)₂, C(O)NH₂, C(O)NHR⁶⁸, C(O)N(R⁶⁸)₂, C(O)NHOH, C(O)NHOR⁶⁸, C(O)NHSO₂R⁶⁸, C(O)NR⁶⁸SO₂R⁶⁸, SO₂NH₂, SO₂NHR⁶⁸, SO₂N(R⁶⁸)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁸, C(N)N(R⁶⁸)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R⁶⁸ is R⁶⁹, R⁷⁰, R⁷¹ or R⁷²;

40 R⁶⁹ is phenyl, which is unfused or fused with R^{69A}; R^{69A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁷⁰ is heteroaryl, which is unfused or fused with R^{70A}; R^{70A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;


R⁷¹ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{71A}; R^{71A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

45 R⁷² is alkyl, alkenyl or alkenyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁷³, OR⁷³, SR⁷³, S(O)R⁷³, SO₂R⁷³, C(O)R⁷³, CO(O)R⁷³, OC(O)R⁷³, OC(O)OR⁷³, NH₂, NHR⁷³, N(R⁷³)₂, NHC(O)R⁷³, NR⁷³C(O)R⁷³, NHS(O)₂R⁷³, NR⁷³S(O)₂R⁷³, NHC(O)OR⁷³, NR⁷³C(O)OR⁷³, NHC(O)NH₂, NHC(O)NHR⁷³, NHC(O)N(R⁷³)₂, NR⁷³C(O)NHR⁷³, NR⁷³C(O)N(R⁷³)₂, C(O)NH₂, C(O)NHR⁷³, C(O)N(R⁷³)₂, C(O)NHOH, C(O)NHOR⁷³, C(O)NHSO₂R⁷³, C(O)NR⁷³SO₂R⁷³, SO₂NH₂, SO₂NHR⁷³, SO₂N(R⁷³)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁷³, C(N)N(R⁷³)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂,
50 CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R⁷³ is alkyl, alkenyl, alkenyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; and

55 wherein the moieties represented by R⁶⁹, R⁷⁰, and R⁷¹ are unsubstituted or substituted with one or two or three or four of independently selected NH₂, C(O)NH₂, C(O)NHOH, SO₂NH₂, CF₃, CF₂CF₃, C(O)H, C(O)OH, C(N)NH₂, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I.

E 2. A compound having Formula (II)



(II),

or a therapeutically acceptable salt thereof, wherein

R¹⁰⁰ is as described for substituents on R²⁶;

n is 0, 1, 2, or 3;

A¹ is N or C(A²);

A² is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNH₂, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};

B¹ is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNH₂, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};

D¹ is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNH₂, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};

E¹ is H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNH₂, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; and

Y¹ is H, CN, NO₂, C(O)OH, F, Cl, Br, I, CF₃, OCF₃, CF₂CF₃, OCF₂CF₃, R¹⁷, OR¹⁷, C(O)R¹⁷, C(O)OR¹⁷, SR¹⁷, SO₂R¹⁷, NH₂, NHR¹⁷, N(R¹⁷)₂, NHC(O)R¹⁷, C(O)NH₂, C(O)NHR¹⁷, C(O)N(R¹⁷)₂, NHS(O)R¹⁷ or NHSO₂R¹⁷; or E¹ and Y¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

A², B¹, and D¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNH₂, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; or

Y¹ and B¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cyclo-

alkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

A², D¹, and E¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; or

A² and B¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

D¹, E¹, and Y¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A}; or

A² and D¹, together with the atoms to which they are attached, are benzene, naphthalene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

B¹, E¹, and Y¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ or C(O)OR^{1A};

R¹ is R², R³, R⁴ or R⁵;

R^{1A} is cycloalkyl, cycloalkenyl or cycloalkynyl;

R² is phenyl, which is unfused or fused with R^{2A}; R^{2A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³ is heteroaryl, which is unfused or fused with R^{3A}; R^{3A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁴ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{4A}; R^{4A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁵ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R⁶, NC(R^{6A})(R^{6B}), R⁷, OR⁷, SR⁷, S(O)R⁷, SO₂R⁷, NHR⁷, N(R⁷)₂, C(O)R⁷, C(O)NH₂, C(O)NHR⁷, C(O)N(R⁷)₂, NHC(O)R⁷, NR⁷C(O)R⁷, NHSO₂R⁷, NHC(O)OR⁷, SO₂NH₂, SO₂NHR⁷, SO₂N(R⁷)₂, NHC(O)NH₂, NHC(O)NHR⁷, NHC(O)CH(CH₃)NHC(O)CH(CH₃)NH₂, NHC(O)CH(CH₃)NHC(O)CH(CH₃)NHR⁷, OH, (O), C(O)OH, N₃, CN, NH₂, CF₃, CF₂CF₃, F, Cl, Br or I;

R⁶ is C₂-C₅-spiroalkyl, each of which is unsubstituted or substituted with OH, (O), N₃, CN, CF₃, CF₂CF₃, F, Cl, Br, I, NH₂, NH(CH₃) or N(CH₃)₂;

R^{6A} and R^{6B} are independently selected alkyl or, together with the N to which they are attached, R^{6C};

R^{6C} is aziridin-1-yl, azetidin-1-yl, pyrrolidin-1-yl or piperidin-1-yl, each having one CH₂ moiety unreplaced or replaced with O, C(O), CNOH, CNOCH₃, S, S(O), SO₂ or NH;

R⁷ is R⁸, R⁹, R¹⁰ or R¹¹;

R⁸ is phenyl, which is unfused or fused with R^{8A}; R^{8A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁹ is heteroaryl, which is unfused or fused with R^{9A}; R^{9A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R¹⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{10A}; R^{10A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R¹¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R¹², OR¹², SR¹², S(O)R¹², SO₂R¹², C(O)R¹², CO(O)R¹², OC(O)R¹², OC(O)OR¹², NH₂, NHR¹², N(R¹²)₂, NHC(O)R¹², NR¹²C(O)R¹², NHS(O)₂R¹², NR¹²S(O)₂R¹², NHC(O)OR¹², NR¹²C(O)OR¹², NHC(O)NH₂, NHC(O)NHR¹², NHC(O)N(R¹²)₂, NR¹²C(O)NHR¹², NR¹²C(O)N(R¹²)₂, C(O)NH₂, C(O)NHR¹², C(O)N(R¹²)₂, C(O)NHOH, C(O)NHOR¹², C(O)NHSO₂R¹², C(O)NR¹²SO₂R¹², SO₂NH₂, SO₂NHR¹², SO₂N(R¹²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR¹², C(N)N(R¹²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R¹² is R¹³, R¹⁴, R¹⁵ or R¹⁶;

R¹³ is phenyl, which is unfused or fused with R^{13A}; R^{13A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R¹⁴ is heteroaryl, which is unfused or fused with R^{14A}; R^{14A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R¹⁵ is cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene, each of which is unfused or fused with R^{15A}; R^{15A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R¹⁶ is alkyl, alkenyl or alkynyl;

R¹⁷ is R¹⁸, R¹⁹, R²⁰ or R²¹;

R¹⁸ is phenyl, which is unfused or fused with R^{18A}; R^{18A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R¹⁹ is heteroaryl, which is unfused or fused with R^{19A}; R^{19A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl each of which is unfused or fused with R^{20A}; R^{20A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of independently selected R²², OR²², SR²², S(O)R²², SO₂R²², C(O)R²², CO(O)R²², OC(O)R²², OC(O)OR²², NH₂, NHR²², N(R²²)₂, NHC(O)R²², NR²²C(O)R²², NHS(O)₂R²², NR²²S(O)₂R²², NHC(O)OR²², NR²²C(O)OR²², NHC(O)NH₂, NHC(O)NHR²², NHC(O)N(R²²)₂, NR²²C(O)NHR²², NR²²C(O)N(R²¹)₂, C(O)NH₂, C(O)NHR²², C(O)N(R²²)₂, C(O)NHOH, C(O)NHOR²², C(O)NHSO₂R²², C(O)NR²²SO₂R²², SO₂NH₂, SO₂NHR²², SO₂N(R²²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR²², C(N)N(R²²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R²² is R²³, R²⁴ or R²⁵;

R²³ is phenyl, which is unfused or fused with R^{23A}; R^{23A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁴ is heteroarene, which is unfused or fused with R^{24A}; R^{24A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{25A}; R^{25A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

Z² is R²⁸, R²⁹ or R³⁰;

L¹ is a R³⁷, OR³⁷, SR³⁷, S(O)R³⁷, SO₂R³⁷, C(O)R³⁷, CO(O)R³⁷, OC(O)R³⁷, OC(O)OR³⁷, NHR³⁷, C(O)NH, C(O)NR³⁷, C(O)NHOR³⁷, C(O)NHSO₂R³⁷, SO₂NH, SO₂NHR³⁷, C(N)NH, C(N)NHR³⁷;

R²⁸ is phenylene, which is unfused or fused with R^{28A}; R^{28A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R²⁹ is heteroarylene, which is unfused or fused with R^{29A}; R^{29A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁰ is cycloalkylene, cycloalkenylene, heterocycloalkylene or heterocycloalkenylenylene, each of which is unfused or fused with R^{30A}; R^{30A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁷ is a bond or R^{37A};

R^{37A} is alkylene, alkenylene, or alkynylene, each of which is unsubstituted or substituted with one or two or three independently selected R^{37B}, OR^{37B}, SR^{37B}, S(O)R^{37B}, SO₂R^{37B}, C(O)R^{37B}, CO(O)R^{37B}, OC(O)R^{37B}, OC(O)OR^{37B}, NH₂, NHR^{37B}, N(R^{37B})₂, NHC(O)R^{37B}, NR^{37B}C(O)R^{37B}, NHS(O)₂R^{37B}, NR^{37B}S(O)₂R^{37B}, NHC(O)OR^{37B}, NR^{37B}C(O)OR^{37B}, NHC(O)NH₂, NHC(O)NHR^{37B}, NHC(O)N(R^{37B})₂, NR^{37B}C(O)NHR^{37B}, NR^{37B}C(O)N(R^{37B})₂, C(O)NH₂, C(O)NHR^{37B}, C(O)N(R^{37B})₂, C(O)NHOH, C(O)NHOR^{37B}, C(O)NHSO₂R^{37B}, C(O)NR^{37B}SO₂R^{37B}, SO₂NH₂, SO₂NHR^{37B}, SO₂N(R^{37B})₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR^{37B}, C(N)N(R^{37B})₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I substituents;

R^{37B} is alkyl, alkenyl, alkynyl, or R^{37C};

R^{37C} is phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl;

Z³ is R³⁸, R³⁹ or R⁴⁰;

R³⁸ is phenyl, which is unfused or fused with R^{38A}; R^{38A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³⁹ is heteroaryl, which is unfused or fused with R^{39A}; R^{39A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁴⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{40A}; R^{40A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

wherein the cyclic moieties represented by E¹ and Y¹ together, Y¹ and B¹ together, A² and B¹ together, A² and D¹ together, R^{1A}, R², R^{2A}, R³, R^{3A}, R⁴, R^{4A}, R⁶, R^{6C}, R⁸, R^{8A}, R⁹, R^{9A}, R¹⁰, R^{10A}, R¹³, R^{13A}, R¹⁴, R^{14A}, R¹⁵,

R^{15A}, R¹⁸, R^{18A}, R¹⁹, R^{19A}, R²⁰, R^{20A}, R²³, R^{23A}, R²⁴, R^{24A}, R²⁵, R^{25A}, R²⁶, R^{26A}, R²⁷, R^{27A}, R²⁸, R^{28A}, R²⁹,
 R^{29A}, R³⁰, R^{30A}, R^{37B}, R³⁸, R^{38A}, R³⁹, R^{39A}, R⁴⁰, and R^{40A} are independently unsubstituted, further unsubstituted,
 substituted or further substituted with one or two or three or four or five of independently selected R^{57A}, R⁵⁷,
 OR⁵⁷, SR⁵⁷, S(O)R⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, OC(O)R⁵⁷, OC(O)OR⁵⁷, NH₂, NHR⁵⁷, N(R⁵⁷)₂,
 5 NHC(O)R⁵⁷, NR⁵⁷C(O)R⁵⁷, NHS(O)₂R⁵⁷, NR⁵⁷S(O)₂R⁵⁷, NHC(O)OR⁵⁷, NR⁵⁷C(O)OR⁵⁷, NHC(O)NH₂,
 NHC(O)NHR⁵⁷, NHC(O)N(R⁵⁷)₂, NR⁵⁷C(O)NHR⁵⁷, NR⁵⁷C(O)N(R⁵⁷)₂, C(O)NH₂, C(O)NHR⁵⁷, C(O)N(R⁵⁷)₂,
 C(O)NHOH, C(O)NHOR⁵⁷, C(O)NHSO₂R⁵⁷, C(O)NR⁵⁷SO₂R⁵⁷, SO₂NH₂, SO₂NHR⁵⁷, SO₂N(R⁵⁷)₂, C(O)H,
 C(O)OH, C(N)NH₂, C(N)NHR⁵⁷, C(N)N(R⁵⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃,
 OCF₂CF₃, F, Cl, Br or I;
 10 R^{57A} is spiroalkyl or heterospiroalkyl;
 R⁵⁷ is R⁵⁸, R⁵⁹, R⁶⁰ or R⁶¹;
 R⁵⁸ is phenyl, which is unfused or fused with R^{58A}; R^{58A} is benzene, heteroarene, cycloalkane, cycloalkene,
 heterocycloalkane or heterocycloalkene;
 R⁵⁹ is heteroaryl, which is unfused or fused with R^{59A}; R^{59A} is benzene, heteroarene, cycloalkane, cycloalkene,
 15 heterocycloalkane or heterocycloalkene;
 R⁶⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with
 R^{60A}; R^{60A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R⁶¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three of inde-
 pendentlly selected R⁶², OR⁶², SR⁶², S(O)R⁶², SO₂R⁶², C(O)R⁶², CO(O)R⁶², OC(O)R⁶², OC(O)OR⁶², NH₂,
 20 NHR⁶², N(R⁶²)₂, NHC(O)R⁶², NR⁶²C(O)R⁶², NHS(O)₂R⁶², NR⁶²S(O)₂R⁶², NHC(O)OR⁶², NR⁶²C(O)OR⁶²,
 NHC(O)NH₂, NHC(O)NHR⁶², NHC(O)N(R⁶²)₂, NR⁶²C(O)NHR⁶², NR⁶²C(O)N(R⁶²)₂, C(O)NH₂, C(O)NHR⁶²,
 C(O)N(R⁶²)₂, C(O)NHOH, C(O)NHOR⁶², C(O)NHSO₂R⁶², C(O)NR⁶²SO₂R⁶², SO₂NH₂, SO₂NHR⁶²,
 SO₂N(R⁶²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶², C(N)N(R⁶²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂,
 CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;
 25 R⁶² is R⁶³, R⁶⁴, R⁶⁵ or R⁶⁶;
 R⁶³ is phenyl, which is unfused or fused with R^{63A}; R^{63A} is benzene, heteroarene, cycloalkane, cycloalkene,
 heterocycloalkane or heterocycloalkene;
 R⁶⁴ is heteroaryl, which is unfused or fused with R^{64A}; R^{64A} is benzene, heteroarene, cycloalkane, cycloalkene,
 heterocycloalkane or heterocycloalkene;
 30 R⁶⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with
 R^{65A}; R^{65A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R⁶⁶ is alkyl, alkenyl or alkenyl, each of which is unsubstituted or substituted with one or two or three of inde-
 pendentlly selected R⁶⁷, OR⁶⁷, SR⁶⁷, S(O)R⁶⁷, SO₂R⁶⁷, C(O)R⁶⁷, CO(O)R⁶⁷, OC(O)R⁶⁷, OC(O)OR⁶⁷, NH₂,
 35 NHR⁶⁷, N(R⁶⁷)₂, NHC(O)R⁶⁷, NR⁶⁷C(O)R⁶⁷, NHS(O)₂R⁶⁷, NR⁶⁷S(O)₂R⁶⁷, NHC(O)OR⁶⁷, NR⁶⁷C(O)OR⁶⁷,
 NHC(O)NH₂, NHC(O)NHR⁶⁷, NHC(O)N(R⁶⁷)₂, NR⁶⁷C(O)NHR⁶⁷, NR⁶⁷C(O)N(R⁶⁷)₂, C(O)NH₂, C(O)NHR⁶⁷,
 C(O)N(R⁶⁷)₂, C(O)NHOH, C(O)NHOR⁶⁷, C(O)NHSO₂R⁶⁷, C(O)NR⁶⁷SO₂R⁶⁷, SO₂NH₂, SO₂NHR⁶⁷,
 SO₂N(R⁶⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁷, C(N)N(R⁶⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂,
 CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I substituents;
 40 R⁶⁷ is alkyl, alkenyl, alkynyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl;
 wherein the cyclic moieties represented by R^{57A}, R⁵⁸, R⁵⁹, R⁶⁰, R⁶³, R⁶⁴, R⁶⁵, and R⁶⁷ are unsubstituted or
 substituted with one or two or three or four of independently selected R⁶⁸, OR⁶⁸, SR⁶⁸, S(O)R⁶⁸, SO₂R⁶⁸,
 C(O)R⁶⁸, CO(O)R⁶⁸, OC(O)R⁶⁸, OC(O)OR⁶⁸, NH₂, NHR⁶⁸, N(R⁶⁸)₂, NHC(O)R⁶⁸, NR⁶⁸C(O)R⁶⁸, NHS(O)₂R⁶⁸,
 NR⁶⁸S(O)₂R⁶⁸, NHC(O)OR⁶⁸, NR⁶⁸C(O)OR⁶⁸, NHC(O)NH₂, NHC(O)NHR⁶⁸, NHC(O)N(R⁶⁸)₂,
 45 NR⁶⁸C(O)NHR⁶⁸, NR⁶⁸C(O)N(R⁶⁸)₂, C(O)NH₂, C(O)NHR⁶⁸, C(O)N(R⁶⁸)₂, C(O)NHOH, C(O)NHOR⁶⁸,
 C(O)NHSO₂R⁶⁸, C(O)NR⁶⁸SO₂R⁶⁸, SO₂NH₂, SO₂NHR⁶⁸, SO₂N(R⁶⁸)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁸,
 C(N)N(R⁶⁸)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;
 R⁶⁸ is R⁶⁹, R⁷⁰, R⁷¹ or R⁷²;
 R⁶⁹ is phenyl, which is unfused or fused with R^{69A}; R^{69A} is benzene, heteroarene, cycloalkane, cycloalkene,
 heterocycloalkane or heterocycloalkene;
 50 R⁷⁰ is heteroaryl, which is unfused or fused with R^{70A}; R^{70A} is benzene, heteroarene, cycloalkane, cycloalkene,
 heterocycloalkane or heterocycloalkene;
 R⁷¹ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with
 R^{71A}; R^{71A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R⁷² is alkyl, alkenyl or alkenyl, each of which is unsubstituted or substituted with one or two or three of inde-
 55 pendentlly selected R⁷³, OR⁷³, SR⁷³, S(O)R⁷³, SO₂R⁷³, C(O)R⁷³, CO(O)R⁷³, OC(O)R⁷³, OC(O)OR⁷³, NH₂,
 NHR⁷³, N(R⁷³)₂, NHC(O)R⁷³, NR⁷³C(O)R⁷³, NHS(O)₂R⁷³, NR⁷³S(O)₂R⁷³, NHC(O)OR⁷³, NR⁷³C(O)OR⁷³,
 NHC(O)NH₂, NHC(O)NHR⁷³, NHC(O)N(R⁷³)₂, NR⁷³C(O)NHR⁷³, NR⁷³C(O)N(R⁷³)₂, C(O)NH₂, C(O)NHR⁷³,
 C(O)N(R⁷³)₂, C(O)NHOH, C(O)NHOR⁷³, C(O)NHSO₂R⁷³, C(O)NR⁷³SO₂R⁷³, SO₂NH₂, SO₂NHR⁷³,

SO₂N(R⁷³)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁷³, C(N)N(R⁷³)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I;

R⁷³ is alkyl, alkenyl, alkenyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; and

5 wherein the moieties represented by R⁶⁹, R⁷⁰, and R⁷¹ are unsubstituted or substituted with one or two or three or four of independently selected NH₂, C(O)NH₂, C(O)NHOH, SO₂NH₂, CF₃, CF₂CF₃, C(O)H, C(O)OH, C(N)NH₂, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br or I.

E 3. The compound of E1, wherein

10

G¹ is H;
A¹ is N or C(A²); and
A² is H.

15

E 4. The compound of E1, wherein

G¹ is H;
A¹ is N or C(A²);
A² is H; and
20 B¹ is NHR¹.

20

E5. The compound of E 1, wherein

G¹ is H;
25 A¹ is N or C(A²);
A² is H;
B¹ is NHR¹; and
D¹ is H.

25

30

E 6. The compound of E1, wherein

G¹ is H;
A¹ is N or C(A²);
A² is H;
35 B¹ is NHR¹;
D¹ is H; and
E¹ is H.

35

E 7. The compound of E1, wherein

G¹ is H;
A¹ is N or C(A²);
A² is H;
B¹ is NHR¹;
45 D¹ is H;
E¹ is H; and
Y¹ is NO₂.

40

45

E 8. A compound of E 1 or E2, wherein the compound is chosen from:

50

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({4-[(3-morpholin-4-ylpropyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
55 4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({3-nitro-4-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({4-[(1-methylpiperidin-4-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

55

- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methylpiperazin-1-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2-methoxyethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({(3S)-tetrahydro-2H-pyran-3-ylmethylamino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({(3R)-tetrahydro-2H-pyran-3-ylmethylamino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-(2-naphthylsulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2-methoxyethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1,4-dioxan-2-ylmethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(2,2,2-trifluoroethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(3,3,3-trifluoropropyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2S)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methoxycyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2R)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(1,4-dioxan-2-ylmethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 Trans-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methoxycyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 N-({3-(aminocarbonyl)-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-[[4-[[tetrahydro-2H-pyran-4-ylmethyl]amino]-3-(trifluoromethyl)phenyl]sulfonyl]benzamide;
- 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-[[4-[[tetrahydro-2H-pyran-4-ylmethyl]amino]-3-[[trifluoromethyl]sulfonyl]phenyl]sulfonyl]benzamide;
- Trans-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-[[4-[[4-morpholin-4-ylcyclohexyl]amino]-3-[[trifluoromethyl]sulfonyl]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-[[4-[[1-methylpiperidin-4-yl]amino]-3-[[trifluoromethyl]sulfonyl]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino]sulfonyl]-2-(tetrahydro-2H-pyran-4-ylmethoxy)nicotinamide;
- N-[[5-bromo-6-[[1-methylpiperidin-4-yl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-methylpiperidin-4-yl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-(1,4-dioxan-2-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[5-bromo-6-(1,4-dioxan-2-ylmethoxy)pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[2,2-dimethyltetrahydro-2H-pyran-4-yl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[3-chloro-5-cyano-4-[[tetrahydro-2H-pyran-4-ylmethyl]amino]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[4-[[1-acetyl piperidin-4-yl]amino]-3-nitrophenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[2-chloro-5-fluoro-4-[[tetrahydro-2H-pyran-4-ylmethyl]amino]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[3-morpholin-4-ylpropyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-[[4-fluorotetrahydro-2H-pyran-4-yl]methoxy]pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-(2-morpholin-4-ylethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[3-chloro-4-[[2-(2-methoxyethoxy)ethyl]sulfonyl]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[2-(2-methoxyethoxy)ethyl]sulfonyl]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-morpholin-4-ylcyclohexyl]oxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[5-bromo-6-[[1-tetrahydro-2H-pyran-4-ylpiperidin-4-yl]amino]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[2-(cyanoethyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Cis-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-[[4-[[4-morpholin-4-ylcyclohexyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-N-[[4-[[4-bis(cyclopropylmethyl)amino]cyclohexyl]amino]-3-nitrophenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[1-methylpiperidin-4-yl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-morpholin-3-ylmethyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-[[4-[[4-methylpiperazin-1-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-morpholin-4-ylbut-2-ynyl]oxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-ethynyl-6-(tetrahydro-

2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-oxo-3,4-dihydroquina-
 zolin-6-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-[(4-[(4-morpholin-4-ylcy-
 5 clohexyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-[(4-[(4-meth-
 oxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-[(4-[(4-fluorotetrahydro-2H-pyran-4-
 yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 Trans-4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-[(4-[(4-methoxycy-
 clohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(tetrahydro-2H-
 pyran-4-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(5-cyano-6-(tetrahydro-2H-
 15 pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 tert-butyl 3-[[4-[(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrro-
 lo[2,3-b]pyridin-5-yloxy)benzoyl]amino]sulfonyl]-2-nitrophenoxymethyl]morpholine-4-carboxylate;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-(morpholin-3-ylmeth-
 oxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-
 ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(1-(methylsulfonyl)pip-
 eridin-4-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(1,1-dioxidotetrahydro-
 25 2H-thiopyran-4-yl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(4-chloro-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piper-
 azin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[1-(2,2-trifluor-
 oethyl)piperidin-4-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(5-cyano-6-[(1-tetrahydro-
 2H-pyran-4-yl)piperidin-4-yl]oxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(5-isopropyl-6-(tetrahydro-
 2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(3-chloro-5-fluoro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-
 35 4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

and therapeutically acceptable salts, and metabolites thereof.

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(1-[2-fluoro-1-(fluor-
 40 omethyl)ethyl]piperidin-4-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-
 dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(1-(2,2-difluoro-
 ethyl)piperidin-4-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(1-cyclopropyl)piperidin-
 4-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(1-morpholin-4-ylcy-
 clohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(4-(dicyclopro-
 50 pylamino)cyclohexyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-6,6-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(tetrahydro-2H-
 pyran-4-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-bromo-6-[(4-ethylmorpholin-3-yl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcy-
 clohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(4-ethylmorpholin-3-
 yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(4-tetrahydro-
 2H-pyran-4-ylmorpholin-3-yl)methoxy]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(3S)-1-tetrahydro-2H-pyran-4-ylpiperidin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(5-cyano-6-[(tetrahydro-2H-pyran-4-ylmethyl)amino]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(1,1-dioxidothio-moipholin-4-yl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(4-[(4-aminotetrahydro-2H-pyran-4-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(5-cyano-6-[(4-morpholin-4-ylcyclohexyl)amino]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-cyano-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(1S,3R)-3-moipholin-4-ylcyclopentyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(1R,3S)-3-morpholin-4-ylcyclopentyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(morpholin-2-ylmethyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(tetrahydrofuran-3-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(1-cis-3-fluorotetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(1-tetrahydro-2H-pyran-4-ylazetidid-3-yl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(1-tetrahydrofuran-3-ylazetidid-3-yl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl)methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[(trans-4-hydroxycyclohexyl)methoxy]-3-nitrophenyl)sulfonyl]benzamide;
 35 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-[(2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[(cis-4-methoxycyclohexyl)methoxy]-3-nitrophenyl)sulfonyl]benzamide;
 Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(4-(cyclopropylamino)cyclohexyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(4-(tetrahydro-2H-pyran-4-ylamino)cyclohexyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 Trans-N-[(5-bromo-6-[(4-morpholin-4-ylcyclohexyl)oxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(4-methoxycyclohexyl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 tert-butyl 4-[(4-[(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino)sulfonyl]-2-nitrophenoxymethyl]-4-fluoropiperidine-1-carboxylate;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(4-fluoropiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(4-(tetrahydro-2H-pyran-4-ylpiperazin-1-yl)cyclohexyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(3R)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-

yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[[(3S)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[(3S)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[4-methylmorpholin-2-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[4-(2-methoxyethyl)morpholin-2-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(4-[[[4-(acetylmorpholin-2-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[trans-4-(fluoromethyl)-1-oxetan-3-ylpyrrolidin-3-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[4-fluorotetrahydro-2H-pyran-4-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(1-oxetan-3-ylpiperidin-4-yl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[1-(cyclobutyl)piperidin-4-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[(3S)-1-cyclopropylpyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(1-tetrahydrofuran-3-ylpiperidin-4-yl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[(3R)-1-cyclopropylpyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[[(3S)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[3-hydroxy-2,2-dimethylpropyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[1-(methylsulfonyl)piperidin-3-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(4-[[[1-(acetylpiperidin-3-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[(3R)-1-(methylsulfonyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-3,3-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[1-[2-fluoro-1-(fluoromethyl)ethyl]azetidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[1-(methylsulfonyl)pyrrolidin-3-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(4-[[[1-(acetylpyrrolidin-3-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(4-[[[(3R)-1-acetylpyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[3-methoxy-2,2-dimethylpropyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[(1R,3R)-3-hydroxycyclopentyl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[(1S,3S)-3-hydroxycyclopentyl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[[(1S,3R)-3-hydroxycyclopentyl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([[(1R,3S)-3-hydroxycyclopentyl]methyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[[(3S)-2-oxopiperidin-3-yl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([[(1-[2-fluoro-1-(fluoromethyl)ethyl]azetidin-3-yl)methyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(1-oxetan-3-ylazetidin-3-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(1-oxetan-3-ylpiperidin-4-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([[(1-cyclopropyl)piperidin-4-yl]methyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([[(2-fluoroethyl)morpholin-2-yl]methyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([[(2,2-difluoroethyl)morpholin-2-yl]methyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([[(4-fluoro-1-oxetan-3-ylpiperidin-4-yl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([[(2S)-4,4-difluoro-1-oxetan-3-ylpyrrolidin-2-yl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[[4-tetrahydro-2H-pyran-4-yl]morpholin-3-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([[(4-cyclobutyl)morpholin-3-yl]methyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[[4-tetrahydrofuran-3-yl]morpholin-3-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([[(1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl)methyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([[(1-cyclopropyl-4-fluoropiperidin-4-yl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([[(4-methoxybenzyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[[3-(trifluoromethoxy)benzyl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([[(3-methoxybenzyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([[(difluoromethoxy)benzyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([[(1,4-dioxaspiro[4.5]deca-8-ylamino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-N-[[4-([[(4-(acetyl)amino)cyclohexyl]amino)-3-nitrophenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([[(3S)-1-(2-fluoroethyl)pyrrolidin-3-yl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([[(3S)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([[(3R)-1-(2-fluoroethyl)pyrrolidin-3-yl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[[(3S)-1-oxetan-3-ylpyrrolidin-3-yl]methoxy]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([[(4-hydroxybenzyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([[(3-hydroxybenzyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[3-(difluoromethoxy)benzyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[cis-3-morpholin-4-yl-cyclopentyl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-[(methylsulfonyl)amino]cyclohexyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-cyclopropylpiperidin-4-yl]amino]-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[(1-oxetan-3-yl-piperidin-4-yl)methoxy]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-fluoro-1-tetrahydro-2H-pyran-4-yl]piperidin-4-yl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-fluoro-1-tetrahydro-furan-3-yl]piperidin-4-yl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-fluoro-1-(methylsulfonyl)piperidin-4-yl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3R)-1-oxetan-3-yl]pyrrolidin-3-yl]methyl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-hydroxycyclohexyl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(3-(dimethylamino)propoxy]benzyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(2-morpholin-4-ylethoxy)benzyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(E)-4-hydroxy-1-adamantyl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(Z)-4-hydroxy-1-adamantyl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[4-[[[(1S,4S)-bicyclo[2.2.1]hept-5-en-2-yl]methoxy]-3-nitrophenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-methyl-5-oxopyrrolidin-3-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(1R,4R,5R,6S)-5,6-dihydroxybicyclo[2.2.1]hept-2-yl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(1R,4R,5S,6R)-5,6-dihydroxybicyclo[2.2.1]hept-2-yl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[(3-oxocyclohexyl)methoxy]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohexa-1,3-dien-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3R)-1-[2-fluoro-1-(fluoromethyl)ethyl]pyrrolidin-3-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[6-[[4-fluorotetrahydro-2H-pyran-4-yl]methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3S)-1-oxetan-3-yl]pyrrolidin-3-yl]methyl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 Trans-N-[[5-chloro-6-[[4-methoxycyclohexyl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Cis-N-[[5-chloro-6-[[4-methoxycyclohexyl]methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3S)-1-oxetan-3-yl]pyrrolidin-3-yl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-[2-(2-methoxyethoxy)ethyl]morpholin-2-yl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([4-(cyanomethyl)morpholin-2-yl]methyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([4-(N,N-dimethylglycyl)morpholin-2-yl]methyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 (2-[[4-([4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)amino]methyl)morpholin-4-yl]acetic acid;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-([4-(oxetan-3-yl)morpholin-2-yl]methyl)amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([4-cyclopropylmorpholin-2-yl]methyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-(methylsulfonyl)-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([4-fluorotetrahydro-2H-pyran-4-yl]methoxy)-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([4-methyltetrahydro-2H-pyran-4-yl]methoxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 ethyl 4-(4-[[4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)piperazine-1-carboxylate;
 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([4-(morpholin-4-yl)piperidin-1-yl]-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-([3(R)-1-(oxetan-3-yl)pynolidin-3-yl]amino)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([3(R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]amino)-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([1-isopropylpiperidin-4-yl]amino)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([4-([1-tert-butylpiperidin-4-yl]amino)-3-nitrophenyl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([1-(2-methoxyethyl)piperidin-3-yl]methyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([1-(cyanomethyl)piperidin-3-yl]methyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([4-fluoro-1-methylpiperidin-4-yl]methoxy)-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[5-chloro-6-([3(R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]amino)pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 tert-butyl 4-([4-([4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)amino]piperazine-1-carboxylate;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-(pentafluoro- λ^6 -sulfonyl)-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-([4-methoxytetrahydro-2H-pyran-4-yl]methoxy)-3-nitrophenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([3(R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-([4-(oxetan-3-yl)piperazin-1-yl]amino)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-([4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]amino)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-([3(R)-tetrahydrofuran-3-ylamino]phenyl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([4,4-difluorocyclohexyl]methyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([4-([1-tert-butylpiperidin-4-yl]amino)-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-

- 4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-({[4-(oxetan-3-yl)morpholin-2-yl)methyl]amino}-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 N-[(5-chloro-6-[[[4-fluorotetrahydro-2H-pyran-4-yl)methyl]amino]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(1-cyclopropylpiperidin-4-yl)amino]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 N-[(5-chloro-6-[[[(2S)-4-(cyanomethyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[[(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 N-[(5-chloro-6-[[[(2R)-4-(cyanomethyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[[(2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({5-fluoro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[3-hydroxy-2-(hydroxymethyl)-2-methylpropoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-({[4-(1,3-difluoropropan-2-yl)morpholin-2-yl)methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 30 N-[(5-chloro-6-[[1-(cyanomethyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-({[3R]-1-[2-(2-methoxyethoxy)ethyl]pyrrolidin-3-yl)amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-({[3R]-1-(N,N-dimethylglycyl)pyrrolidin-3-yl)amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[[1-(oxetan-3-yl)azetidin-3-yl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 N-[(5-chloro-6-[[1-(cyanomethyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-({[(2R)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-({[(2S)-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 N-[(5-chloro-6-[[[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-({[3R]-1-(cyanomethyl)pyrrolidin-3-yl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 4-(4-[[4-(4'-chlorobiphenyl-2-yl)methyl]-4-methoxypiperidin-1-yl]-N-({5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4'-chlorobiphenyl-2-yl)methyl]-4-methoxypiperidin-1-yl]-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl)-N-({5-chloro-6-(tetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl)-N-({5-chloro-6-(tetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-({5-chloro-6-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[1-(N,N-dimethylglycyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-fluoropiperidin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-fluoropiperidin-1-yl}-N-({5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}-3-(trifluoromethyl)phenyl)sulfonyl)benzamide;
 N-[(5-chloro-6-[[3-(1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[2-(tetrahydrofuran-3-yloxy)ethoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[(trans-4-cyanocyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[(4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({5-cyano-6-[2-(tetrahydro-2H-pyran-4-yl)ethoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-(3-furylmethoxy)-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[3-(1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[3-(1-(2,2-difluoroethyl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({3-chloro-4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]phenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-cyano-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[1-(2,2-difluoroethyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({3-chloro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(4,4-difluorocyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({6-[[1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl]methoxy]-5-(trifluoromethyl)pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({5-chloro-6-[2-(tetrahydrofuran-2-yl)ethoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}-3-methylpiperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({4-[[3-(cyclopropylamino)propyl]amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-(2-methoxyethoxy)pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({5-fluoro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(3-chloro-4-[[1-(methoxyacetyl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(3-chloro-4-[[1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-

dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohexyl)methyl]piperidin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({6-(tetrahydro-2H-pyran-4-ylmethoxy)-5-(trifluoromethyl)pyridin-3-yl)sulfonyl}benzamide;
 5 N-({5-chloro-6-[(trans-4-hydroxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-cyano-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({6-[(trans-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({6-[(cis-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({5-chloro-6-[(4,4-difluoro-1-hydroxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperidin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 N-([3-chloro-4-([trans-4-(morpholin-4-yl)cyclohexyl)methoxy]phenyl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-([3-(cyclopropyl(1,3-thiazol-5-ylmethyl)amino)propyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 N-({3-chloro-4-[(trans-4-hydroxycyclohexyl)methoxy]phenyl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-chloro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-(trifluoromethyl)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-([3-(cyclopropyl(2,2,2-trifluoroethyl)amino)propyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-([1-(oxetan-3-yl)piperidin-4-yl)methoxy]phenyl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3,5-difluoro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({4-([3-(cyclopropyl(oxetan-3-yl)amino)propyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-([1-(1-methyl-L-prolyl)piperidin-4-yl)methoxy]phenyl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3,4-difluoro-5-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-chloro-6-([2-(2S)-4-cyclopropylmorpholin-2-yl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperidin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperidin-1-yl)-N-([3-chloro-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 methyl 2-([4-([4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl)sulfamoyl)-2-nitrophenyl]amino)methyl)morpholine-4-carboxylate;
 45 2-([4-([4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl)sulfamoyl)-2-nitrophenyl]amino)methyl)-N-ethyl-N-methylmorpholine-4-carboxamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([4-([4-(methylsulfonyl)morpholin-2-yl)methyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([4-([3-(cyclobutyl(cyclopropyl)amino)propyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 N-([3-chloro-4-([4-fluoro-1-(oxetan-3-yl)piperidin-4-yl)methoxy]phenyl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-([3-chloro-4-(tetrahydrofuran-3-ylmethoxy)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- 4-(4-[[2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[trans-4-hydroxycyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-[[4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-4-(4-[[9-(4-chlorophenyl)-3-(oxetan-3-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[2R)-4-cyclopropylmorpholin-2-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[2S)-4-cyclopropylmorpholin-2-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-[[tetrahydro-2H-pyran-4-yl)methyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-chloro-6-[[4-[[cyclopropyl(oxetan-3-yl)amino]cyclohexyl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-N-[[4-[[4-cyclopropylmorpholin-2-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-[[4-cyclopropylmorpholin-2-yl)methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 20 N-([3-chloro-4-[[4-cyclopropylmorpholin-2-yl)methyl]amino]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 2-[[2-chloro-4-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]phenyl)amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
- 25 (2S)-2-[[[3-chloro-5-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]pyridin-2-yl)oxy]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
- 30 N-([5-chloro-6-[[4-cyclopropylmorpholin-2-yl)methyl]amino]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 2-[[[3-chloro-5-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]pyridin-2-yl)amino]methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;
- 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[trans-4-hydroxy-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[cis-4-hydroxy-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 N-([5-chloro-6-[[1R,2R,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[2-cyanoethyl](cyclopropyl)amino]cyclohexyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-chloro-6-[[trans-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-chloro-6-(5,6,7,8-tetrahydroimidazo[1,2-a]pyridin-6-yl)methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 N-([5-chloro-6-[[1R,2S,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[cis-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 N-([5-chloro-6-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(3,3-difluoropyrrolidin-1-yl)cyclohexyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 N-([5-chloro-6-[[trans-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-chloro-6-[[cis-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[2,2-difluorocyclo-

propyl)amino]cyclohexyl)amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-chloro-6-[(cis-1-fluoro-4-hydroxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-(2-oxaspiro[3.5]non-7-ylmethoxy)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5,5-bis(fluoromethyl)cyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[(4-cyclopropylmorpholin-2-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-chloro-6-[(trans-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-chloro-6-[(cis-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-cyano-4-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[(trans-4-ethyl-4-hydroxycyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[(cis-4-ethyl-4-hydroxycyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5-(methoxymethyl)-5-methylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-[(2S)-4-(oxetan-3-yl)morpholin-2-yl]methyl]amino)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[(4-[(2-cyanoethyl)(cyclopropyl)amino]-1-fluorocyclohexyl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-nitro-6-[(tetrahydro-2H-pyran-4-ylmethyl)amino]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([3-nitro-4-[(2-oxaspiro[3.5]non-7-ylmethoxy)phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[(4-cyano-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 {[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]}{4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl}amino)methyl pivalate;
 {[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]}{4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl}amino)methyl butyrate;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([3-(tetrahydro-2H-pyran-4-ylmethyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-6-yl)sulfonyl)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([6-[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino]-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-cyano-6-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([4-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)morpholine-4-carboxamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[(4-(methoxymethyl)cyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-chloro-6-[[1-(1,3-thiazol-2-yl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([6-[(cis-4-hydroxy-4-methylcyclohexyl)methyl]amino]-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino]-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

droxy-4-methylcyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

and therapeutically acceptable salts, and metabolites thereof.

E 9. The compound 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

E10. The compound Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

E11. The compound Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

E 12. The compound Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

E 13. The compound 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

E 14. The compound 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

E 15. The compound 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(cis-4-hydroxy-4-methylcyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

E16. The compound N-[(5-chloro-6-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharmaceutically acceptable salt thereof.

E17. The compound N-({5-bromo-6-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino}pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

E 18. The compound 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(3R)-1-(methylsulfonyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

E 19. The compound 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(3-cyclopropyl(oxetan-3-yl)amino)propyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

E 20. The compound 4-(4-[[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically acceptable salt thereof.

E 21. A compound of E1 or E 2, wherein the compound is chosen from:

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(trans-4-hydroxy-4-

methylcyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3R)-1-(methylsulfo-
 5 nyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-[[2-(4-
 chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[3-[cyclopropyl(oxetan-3-yl)ami-
 no]propyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically
 acceptable salt thereof.

E 22. A compound of E1 or E2, wherein the compound is chosen from:

10 N-[[5-chloro-6-[[1-(cyanomethyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-
 dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[3-[cyclopropyl(oxetan-
 3-yl)amino]propyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharma-
 15 ceutically acceptable salt thereof.

E23. A compound of E1 or E2, wherein the compound is chosen from:

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(trans-4-hydroxy-4-
 methylcyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-
 20 [2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(cis-4-hydroxy-4-methylcy-
 clohexyl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide or a pharma-
 ceutically acceptable salt thereof.

E 24. The compound 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[3-[cy-
 25 clopropyl(oxetan-3-yl)amino]propyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 or a pharmaceutically acceptable salt thereof.

E 25. A compound of E1 or E2, wherein the compound is chosen from:

30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3R)-1-tetrahy-
 dro-2H-pyran-4-yl]pyrrolidin-3-yl]methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benza-
 mide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[4-methylmorpholin-2-
 yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically
 35 acceptable salt thereof.

E 26. A compound of E1 or E2, wherein the compound is chosen from:

40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3S)-tetrahy-
 dro-2H-pyran-3-yl]methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1,4-dioxan-2-yl]meth-
 oxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[5-bromo-6-[[1-tetrahydro-2H-pyran-4-yl]piperidin-4-yl]amino]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophe-
 45 nyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[4-[[[4-aminotetrahydro-2H-pyran-4-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-
 dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3R)-1-tetrahy-
 dro-2H-pyran-4-yl]pyrrolidin-3-yl]methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benza-
 50 mide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[4-methylmorpholin-2-
 yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3R)-1-(methylsulfo-
 nyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-methoxytetrahydro-
 2H-pyran-4-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 N-[[5-chloro-6-[[1-(cyanomethyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-
 dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[3-[cyclopropyl(oxetan-

3-yl)amino]propyl]amino)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[(5-chloro-6-[[[1R,2S,4R,5R]-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy]pyridin-3-yl)sulfonyl]-
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-
 yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(6-[[[cis-4-hydroxy-4-meth-
 ylcyclohexyl]methyl]amino]-5-nitropyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(tetrahy-
 dro-2H-pyran-4-yl)methyl]amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

or a pharmaceutically acceptable salt thereof.

E 27. The compound N-[(5-chloro-6-[[[1-(cyanomethyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlo-
 rophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or
 a pharmaceutically acceptable salt thereof.

E 28. A compound of E 1 or E 2, wherein the compound is chosen from:

N-[(4-[[[4-aminotetrahydro-2H-pyran-4-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-
 dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; 4-(4-[[2-(4-
 chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[4-methoxytetrahydro-2H-pyran-4-
 yl]methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; or a pharmaceutically ac-
 ceptable salt thereof.

E 29. The compound 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[[4-
 methoxytetrahydro-2H-pyran-4-yl]methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide
 or a pharmaceutically acceptable salt thereof.

E 30. A composition for treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, chronic lymphocytic leukemia, myeloma, prostate cancer, small cell lung cancer or spleen cancer, said composition comprising an excipient and a therapeutically effective amount of a compound of E 1 or E 2.

E31. A method of treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, chronic lymphocytic leukemia, myeloma, prostate cancer, small cell lung cancer or spleen cancer in a patient, said method comprising administering to the patient a therapeutically effective amount of a compound of E1 or E 2.


E 32. A method of treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, chronic lymphocytic leukemia, myeloma, prostate cancer, small cell lung cancer or spleen cancer in a patient, said method comprising administering to the patient therapeutically effective amount of the compound of E1 or E2 and a therapeutically effective amount of one additional therapeutic agent or more than one additional therapeutic agent.

SEQUENCE LISTING

<110> AbbVie Bahamas Ltd.
 5 <120> APOPTOSIS-INDUCING AGENTS FOR THE TREATMENT OF CANCER AND IMMUNE AND
 AUTOIMMUNE DISEASES
 <130> P37808EP-D1-PCT
 10 <140> Not yet assigned
 <141> 2010-05-26
 <150> US 61/181,203
 <151> 2009-05-26
 15 <160> 1
 <170> FastSEQ for Windows Version 4.0
 <210> 1
 20 <211> 16
 <212> PRT
 <213> Artificial Sequence
 <220>
 25 <223> Probe Peptide
 <220>
 <221> MOD_RES
 <222> 1
 30 <223> Gly is modified with acetyl
 <220>
 <221> MOD_RES
 <222> 13
 35 <223> Lys is modified with 6-FAM
 <220>
 <221> MOD_RES
 <222> 16
 40 <223> Arg is modified with NH2
 <400> 1
 Gly Gln Val Gly Arg Gln Leu Ala Ile Ile Gly Asp Lys Ile Asn Arg
 1 5 10 15

Claims

1. A combination comprising
 50 a compound or a therapeutically acceptable salt thereof, and
 one or more than one additional therapeutic agent(s),
 wherein the compound has Formula (II)
 55



(II),

wherein

R^{100} is R^{57A} , R^{57} , OR^{57} , SR^{57} , $S(O)R^{57}$, SO_2R^{57} , $C(O)R^{57}$, $CO(O)R^{57}$, $OC(O)R^{57}$, $OC(O)OR^{57}$, NHR^{57} , $N(R^{57})_2$, $NHC(O)R^{57}$, $NR^{57}C(O)R^{57}$, $NHS(O)_2R^{57}$, $NR^{57}S(O)_2R^{57}$, $NHC(O)OR^{57}$, $NR^{57}C(O)OR^{57}$, $NHC(O)NH_2$, $NHC(O)NHR^{57}$, $NHC(O)N(R^{57})_2$, $NR^{57}C(O)NHR^{57}$, $NR^{57}C(O)N(R^{57})_2$, $C(O)NH_2$, $C(O)NHR^{57}$, $C(O)N(R^{57})_2$, $C(O)NHOH$, $C(O)NHOR^{57}$, $C(O)NHSO_2R^{57}$, $C(O)NR^{57}SO_2R^{57}$, SO_2NH_2 , SO_2NHR^{57} , $SO_2N(R^{57})_2$, $C(O)H$, $C(O)OH$, $C(N)NH_2$, $C(N)NHR^{57}$, $C(N)N(R^{57})_2$, $CNOH$, $CNOCH_3$, OH , (O) , CN , N_3 , NO_2 , CF_3 , CF_2CF_3 , OCF_3 , OCF_2CF_3 , F , Cl , Br or I ;

n is 0, 1, 2, or 3;

A^1 is N or $C(A^2)$;

A^2 is H , R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F , Cl , Br , I , CN , NO_2 , N_3 , OH , $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$;

B^1 is H , R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F , Cl , Br , I , CN , NO_2 , N_3 , OH , $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$;

D^1 is H , R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F , Cl , Br , I , CN , NO_2 , N_3 , OH , $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$;

E^1 is H , R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 , $C(O)R^1$, $C(O)OR^1$, $OC(O)R^1$, NHR^1 , $N(R^1)_2$, $C(O)NHR^1$, $C(O)N(R^1)_2$, $NHC(O)R^1$, $NR^1C(O)R^1$, $NHC(O)OR^1$, $NR^1C(O)OR^1$, $NHC(O)NH_2$, $NHC(O)NHR^1$, $NHC(O)N(R^1)_2$, $NR^1C(O)NHR^1$, $NR^1C(O)N(R^1)_2$, SO_2NH_2 , SO_2NHR^1 , $SO_2N(R^1)_2$, $NHSO_2R^1$, $NR^1SO_2R^1$, $NHSO_2NHR^1$, $NHSO_2N(R^1)_2$, $NR^1SO_2NHR^1$, $NR^1SO_2N(R^1)_2$, $C(O)NHNOR^1$, $C(O)NHSO_2R^1$, $C(NH)NH_2$, $C(NH)NHR^1$, $C(NH)N(R^1)_2$, $NHSO_2NHR^1$, $NHSO_2N(CH_3)R^1$, $N(CH_3)SO_2N(CH_3)R^1$, F , Cl , Br , I , CN , NO_2 , N_3 , OH , $C(O)H$, $CHNOH$, $CH(NOCH_3)$, CF_3 , $C(O)OH$, $C(O)NH_2$ or $C(O)OR^{1A}$; and

Y^1 is H , CN , NO_2 , $C(O)OH$, F , Cl , Br , I , CF_3 , OCF_3 , CF_2CF_3 , OCF_2CF_3 , R^{17} , OR^{17} , $C(O)R^{17}$, $C(O)OR^{17}$, SR^{17} , SO_2R^{17} , NH_2 , NHR^{17} , $N(R^{17})_2$, $NHC(O)R^{17}$, $C(O)NH_2$, $C(O)NHR^{17}$, $C(O)N(R^{17})_2$, $NHS(O)R^{17}$ or $NHSO_2R^{17}$; or E^1 and Y^1 , together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

A^2 , B^1 , and D^1 are independently selected from the group consisting of H , R^1 , OR^1 , SR^1 , $S(O)R^1$, SO_2R^1 ,

C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ and C(O)OR^{1A}; or

Y¹ and B¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

A², D¹, and E¹ are independently selected from the group consisting of H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ and C(O)OR^{1A}; or

A² and B¹, together with the atoms to which they are attached, are benzene, naphthylene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

D¹, E¹, and Y¹ are independently selected H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ and C(O)OR^{1A}; or

A² and D¹, together with the atoms to which they are attached, are benzene, naphthalene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene; and

B¹, E¹, and Y¹ are independently selected from the group consisting of H, R¹, OR¹, SR¹, S(O)R¹, SO₂R¹, C(O)R¹, C(O)OR¹, OC(O)R¹, NHR¹, N(R¹)₂, C(O)NHR¹, C(O)N(R¹)₂, NHC(O)R¹, NR¹C(O)R¹, NHC(O)OR¹, NR¹C(O)OR¹, NHC(O)NH₂, NHC(O)NHR¹, NHC(O)N(R¹)₂, NR¹C(O)NHR¹, NR¹C(O)N(R¹)₂, SO₂NH₂, SO₂NHR¹, SO₂N(R¹)₂, NHSO₂R¹, NR¹SO₂R¹, NHSO₂NHR¹, NHSO₂N(R¹)₂, NR¹SO₂NHR¹, NR¹SO₂N(R¹)₂, C(O)NHNOH, C(O)NHNOR¹, C(O)NHSO₂R¹, C(NH)NH₂, C(NH)NHR¹, C(NH)N(R¹)₂, NHSO₂NHR¹, NHSO₂N(CH₃)R¹, N(CH₃)SO₂N(CH₃)R¹, F, Cl, Br, I, CN, NO₂, N₃, OH, C(O)H, CHNOH, CH(NOCH₃), CF₃, C(O)OH, C(O)NH₂ and C(O)OR^{1A};

R¹ is R², R³, R⁴ or R⁵;

R^{1A} is cycloalkyl, cycloalkenyl or cycloalkynyl;

R² is phenyl, which is unfused or fused with R^{2A}; R^{2A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R³ is heteroaryl, which is unfused or fused with R^{3A}; R^{3A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁴ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{4A}; R^{4A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁵ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three substituents independently selected from the group consisting of R⁶, NC(R^{6A})(R^{6B}), R⁷, OR⁷, SR⁷, S(O)R⁷, SO₂R⁷, NHR⁷, N(R⁷)₂, C(O)R⁷, C(O)NH₂, C(O)NHR⁷, C(O)N(R⁷)₂, NHC(O)R⁷, NR⁷C(O)R⁷, NHSO₂R⁷, NHC(O)OR⁷, SO₂NH₂, SO₂NHR⁷, SO₂N(R⁷)₂, NHC(O)NH₂, NHC(O)NHR⁷, NHC(O)CH(CH₃)NHC(O)CH(CH₃)NH₂, NHC(O)CH(CH₃)NHC(O)CH(CH₃)NHR⁷, OH, (O), C(O)OH, N₃, CN, NH₂, CF₃, CF₂CF₃, F, Cl, Br and I;

R⁶ is C₂-C₅-spiroalkyl, each of which is unsubstituted or substituted with OH, (O), N₃, CN, CF₃, CF₂CF₃, F, Cl, Br, I, NH₂, NH(CH₃) or N(CH₃)₂;

R^{6A} and R^{6B} are independently selected alkyl;

R⁷ is R⁸, R⁹, R¹⁰ or R¹¹;

R⁸ is phenyl, which is unfused or fused with R^{8A}; R^{8A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁹ is heteroaryl, which is unfused or fused with R^{9A}; R^{9A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R¹⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{10A}; R^{10A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R¹¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three substituents independently selected from the group consisting of R¹², OR¹², SR¹², S(O)R¹², SO₂R¹², C(O)R¹², CO(O)R¹²,

OC(O)R¹², OC(O)OR¹², NH₂, NHR¹², N(R¹²)₂, NHC(O)R¹², NR¹²C(O)R¹², NHS(O)₂R¹², NR¹²S(O)₂R¹²,
 NHC(O)OR¹², NR¹²C(O)OR¹², NHC(O)NH₂, NHC(O)NHR¹², NHC(O)N(R¹²)₂, NR¹²C(O)NHR¹²,
 NR¹²C(O)NR¹²)₂, C(O)NH₂, C(O)NHR¹², C(O)N(R¹²)₂, C(O)NHOH, C(O)NHOR¹², C(O)NHSO₂R¹²,
 C(O)NR¹²SO₂R¹², SO₂NH₂, SO₂NHR¹², SO₂N(R¹²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR¹², C(N)N(R¹²)₂,
 5 CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I;
 R¹² is R¹³, R¹⁴, R¹⁵ or R¹⁶;
 R¹³ is phenyl, which is unfused or fused with R^{13A}; R^{13A} is benzene, heteroarene, cycloalkane, cycloalkene,
 heterocycloalkane or heterocycloalkene;
 R¹⁴ is heteroaryl, which is unfused or fused with R^{14A}; R^{14A} is benzene, heteroarene, cycloalkane, cycloalkene,
 10 heterocycloalkane or heterocycloalkene;
 R¹⁵ is cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene, each of which is unfused or fused
 with R^{15A}; R^{15A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R¹⁶ is alkyl, alkenyl or alkynyl;
 R¹⁷ is R¹⁸, R¹⁹, R²⁰ or R²¹;
 15 R¹⁸ is phenyl, which is unfused or fused with R^{18A}; R^{18A} is benzene, heteroarene, cycloalkane, cycloalkene,
 heterocycloalkane or heterocycloalkene;
 R¹⁹ is heteroaryl, which is unfused or fused with R^{19A}; R^{19A} is benzene, heteroarene, cycloalkane, cycloalkene,
 heterocycloalkane or heterocycloalkene;
 R²⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl each of which is unfused or fused with
 R^{20A}; R^{20A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 R²¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three substituents
 independently selected from the group consisting of R²², OR²², SR²², S(O)R²², SO₂R²², C(O)R²², CO(O)R²²,
 OC(O)R²², OC(O)OR²², NH₂, NHR²², N(R²²)₂, NHC(O)R²², NR²²C(O)R²², NHS(O)₂R²², NR²²S(O)₂R²²,
 NHC(O)OR²², NR²²C(O)OR²², NHC(O)NH₂, NHC(O)NHR²², NHC(O)N(R²²)₂, NR²²C(O)NHR²²,
 25 NR²²C(O)N(R²²)₂, C(O)NH₂, C(O)NHR²², C(O)N(R²²)₂, C(O)NHOH, C(O)NHOR²², C(O)NHSO₂R²²,
 C(O)NR²²SO₂R²², SO₂NH₂, SO₂NHR²², SO₂N(R²²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR²², C(N)N(R²²)₂,
 CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I;
 R²² is R²³, R²⁴ or R²⁵;
 R²³ is phenyl, which is unfused or fused with R^{23A}; R^{23A} is benzene, heteroarene, cycloalkane, cycloalkene,
 30 heterocycloalkane or heterocycloalkene;
 R²⁴ is heteroarene, which is unfused or fused with R^{24A}; R^{24A} is benzene, heteroarene, cycloalkane, cycloalkene,
 heterocycloalkane or heterocycloalkene;
 R²⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with
 R^{25A}; R^{25A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;
 35 Z² is R²⁸, R²⁹ or R³⁰;
 L¹ is a R³⁷, OR³⁷, SR³⁷, S(O)R³⁷, SO₂R³⁷, C(O)R³⁷, CO(O)R³⁷, OC(O)R³⁷, OC(O)OR³⁷, NHR³⁷, C(O)NH,
 C(O)NR³⁷, C(O)NHOR³⁷, C(O)NHSO₂R³⁷, SO₂NH, SO₂NHR³⁷, C(N)NH, or C(N)NHR³⁷;
 R²⁸ is phenylene, which is unfused or fused with R^{28A}; R^{28A} is benzene, heteroarene, cycloalkane, cycloalkene,
 heterocycloalkane or heterocycloalkene;
 40 R²⁹ is heteroarylene, which is unfused or fused with R^{29A}; R^{29A} is benzene, heteroarene, cycloalkane, cy-
 cloalkene, heterocycloalkane or heterocycloalkene;
 R³⁰ is cycloalkylene, cycloalkenylene, heterocycloalkylene or heterocycloalkenylene, each of which is unfused
 or fused with R^{30A}; R^{30A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocyc-
 cloalkene;
 45 R³⁷ is a bond or R^{37A};
 R^{37A} is alkylene, alkenylene, or alkynylene, each of which is unsubstituted or substituted with one or two or
 three substituents independently selected from the group consisting of R^{37B}, OR^{37B}, SR^{37B}, S(O)R^{37B}, SO₂R^{37B},
 C(O)R^{37B}, CO(O)R^{37B}, OC(O)R^{37B}, OC(O)OR^{37B}, NH₂, NHR^{37B}, N(R^{37B})₂, NHC(O)R^{37B}, NR^{37B}C(O)R^{37B},
 NHS(O)₂R^{37B}, NR^{37B}S(O)₂R^{37B}, NHC(O)OR^{37B}, NR^{37B}C(O)OR^{37B}, NHC(O)NH₂, NHC(O)NHR^{37B},
 50 NHC(O)N(R^{37B})₂, NR^{37B}C(O)NHR^{37B}, NR^{37B}C(O)N(R^{37B})₂, C(O)NH₂, C(O)NHR^{37B}, C(O)N(R^{37B})₂,
 C(O)NHOH, C(O)NHOR^{37B}, C(O)NHSO₂R^{37B}, C(O)NR^{37B}SO₂R^{37B}, SO₂NH₂, SO₂NHR^{37B}, SO₂N(R^{37B})₂,
 C(O)H, C(O)OH, C(N)NH₂, C(N)NHR^{37B}, C(N)N(R^{37B})₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃,
 OCF₃, OCF₂CF₃, F, Cl, Br and I;
 R^{37B} is alkyl, alkenyl, alkynyl, or R^{37C};
 55 R^{37C} is phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl;
 Z³ is R³⁸, R³⁹ or R⁴⁰;
 R³⁸ is phenyl, which is unfused or fused with R^{38A}; R^{38A} is benzene, heteroarene, cycloalkane, cycloalkene,
 heterocycloalkane or heterocycloalkene;

R³⁹ is heteroaryl, which is unfused or fused with R^{39A}; R^{39A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁴⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{40A}; R^{40A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

wherein the cyclic moieties represented by E¹ and Y¹ together, Y¹ and B¹ together, A² and B¹ together, A² and D¹ together, R^{1A}, R², R^{2A}, R³, R^{3A}, R⁴, R^{4A}, R⁶, R⁸, R^{8A}, R⁹, R^{9A}, R¹⁰, R^{10A}, R¹³, R^{13A}, R¹⁴, R^{14A}, R¹⁵, R^{15A}, R¹⁸, R^{18A}, R¹⁹, R^{19A}, R²⁰, R^{20A}, R²³, R^{23A}, R²⁴, R^{24A}, R²⁵, R^{25A}, R²⁸, R^{28A}, R²⁹, R^{29A}, R³⁰, R^{30A}, R³⁸, R^{38A}, R³⁹, R^{39A}, R⁴⁰, and R^{40A} are independently unsubstituted, substituted or further substituted with one or two or three or four or five substituents independently selected from the group consisting of R^{57A}, R⁵⁷, OR⁵⁷, SR⁵⁷, S(O)R⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, OC(O)R⁵⁷, OC(O)OR⁵⁷, NH₂, NHR⁵⁷, N(R⁵⁷)₂, NHC(O)R⁵⁷, NR⁵⁷C(O)R⁵⁷, NHS(O)₂R⁵⁷, NR⁵⁷S(O)₂R⁵⁷, NHC(O)OR⁵⁷, NR⁵⁷C(O)OR⁵⁷, NHC(O)NH₂, NHC(O)NHR⁵⁷, NHC(O)N(R⁵⁷)₂, NR⁵⁷CONHR⁵⁷, NR⁵⁷C(O)N(R⁵⁷)₂, C(O)NH₂, C(O)NHR⁵⁷, C(O)N(R⁵⁷)₂, C(O)NHOH, C(O)NHOR⁵⁷, C(O)NHSO₂R⁵⁷, C(O)NR⁵⁷SO₂R⁵⁷, SO₂NH₂, SO₂NHR⁵⁷, SO₂N(R⁵⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁵⁷, C(N)N(R⁵⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I;

R^{57A} is spiroalkyl or heterospiroalkyl;

R⁵⁷ is R⁵⁸, R⁵⁹, R⁶⁰ or R⁶¹;

R⁵⁸ is phenyl, which is unfused or fused with R^{58A}; R^{58A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁵⁹ is heteroaryl, which is unfused or fused with R^{59A}; R^{59A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁰ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{60A}; R^{60A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶¹ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three substituents independently selected from the group consisting of R⁶², OR⁶², SR⁶², S(O)R⁶², SO₂R⁶², C(O)R⁶², CO(O)R⁶², OC(O)R⁶², OC(O)OR⁶², NH₂, NHR⁶², N(R⁶²)₂, NHC(O)R⁶², NR⁶²C(O)R⁶², NHS(O)₂R⁶², NR⁶²S(O)₂R⁶², NHC(O)OR⁶², NR⁶²C(O)OR⁶², NHC(O)NH₂, NHC(O)NHR⁶², NHC(O)N(R⁶²)₂, NR⁶²C(O)NHR⁶², NR⁶²C(O)N(R⁶²)₂, C(O)NH₂, C(O)NHR⁶², C(O)N(R⁶²)₂, C(O)NHOH, C(O)NHOR⁶², C(O)NHSO₂R⁶², C(O)NR⁶²SO₂R⁶², SO₂NH₂, SO₂NHR⁶², SO₂N(R⁶²)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶², C(N)N(R⁶²)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I;

R⁶² is R⁶³, R⁶⁴, R⁶⁵ or R⁶⁶;

R⁶³ is phenyl, which is unfused or fused with R^{63A}; R^{63A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁴ is heteroaryl, which is unfused or fused with R^{64A}; R^{64A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁵ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{65A}; R^{65A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁶⁶ is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three substituents independently selected from the group consisting of R⁶⁷, OR⁶⁷, SR⁶⁷, S(O)R⁶⁷, SO₂R⁶⁷, C(O)R⁶⁷, CO(O)R⁶⁷, OC(O)R⁶⁷, OC(O)OR⁶⁷, NH₂, NHR⁶⁷, N(R⁶⁷)₂, NHC(O)R⁶⁷, NR⁶⁷C(O)R⁶⁷, NHS(O)₂R⁶⁷, NR⁶⁷S(O)₂R⁶⁷, NHC(O)OR⁶⁷, NR⁶⁷C(O)OR⁶⁷, NHC(O)NH₂, NHC(O)NHR⁶⁷, NHC(O)N(R⁶⁷)₂, NR⁶⁷C(O)NHR⁶⁷, NR⁶⁷C(O)N(R⁶⁷)₂, C(O)NH₂, C(O)NHR⁶⁷, C(O)N(R⁶⁷)₂, C(O)NHOH, C(O)NHOR⁶⁷, C(O)NHSO₂R⁶⁷, C(O)NR⁶⁷SO₂R⁶⁷, SO₂NH₂, SO₂NHR⁶⁷, SO₂N(R⁶⁷)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁷, C(N)N(R⁶⁷)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I;

R⁶⁷ is alkyl, alkenyl, alkynyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; wherein the cyclic moieties represented by R^{57A}, R⁵⁸, R⁵⁹, R⁶⁰, R⁶³, R⁶⁴, R⁶⁵, and R⁶⁷ are unsubstituted or substituted with one or two or three or four substituents independently selected from the group consisting of R⁶⁸, OR⁶⁸, SR⁶⁸, S(O)R⁶⁸, SO₂R⁶⁸, C(O)R⁶⁸, CO(O)R⁶⁸, OC(O)R⁶⁸, OC(O)OR⁶⁸, NH₂, NHR⁶⁸, N(R⁶⁸)₂, NHC(O)R⁶⁸, NR⁶⁸C(O)R⁶⁸, NHS(O)₂R⁶⁸, NR⁶⁸S(O)₂R⁶⁸, NHC(O)OR⁶⁸, NR⁶⁸C(O)OR⁶⁸, NHC(O)NH₂, NHC(O)NHR⁶⁸, NHC(O)N(R⁶⁸)₂, NR⁶⁸C(O)NHR⁶⁸, NR⁶⁸C(O)N(R⁶⁸)₂, C(O)NH₂, C(O)NHR⁶⁸, C(O)N(R⁶⁸)₂, C(O)NHOH, C(O)NHOR⁶⁸, C(O)NHSO₂R⁶⁸, C(O)NR⁶⁸SO₂R⁶⁸, SO₂NH₂, SO₂NHR⁶⁸, SO₂N(R⁶⁸)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁶⁸, C(N)N(R⁶⁸)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I;

R⁶⁸ is R⁶⁹, R⁷⁰, R⁷¹ or R⁷²;

R⁶⁹ is phenyl, which is unfused or fused with R^{69A}; R^{69A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁷⁰ is heteroaryl, which is unfused or fused with R^{70A}; R^{70A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁷¹ is cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl, each of which is unfused or fused with R^{71A}; R^{71A} is benzene, heteroarene, cycloalkane, cycloalkene, heterocycloalkane or heterocycloalkene;

R⁷² is alkyl, alkenyl or alkynyl, each of which is unsubstituted or substituted with one or two or three substituents independently selected from the group consisting of R⁷³, OR⁷³, SR⁷³, S(O)R⁷³, SO₂R⁷³, C(O)R⁷³, CO(O)R⁷³, OC(O)R⁷³, OC(O)OR⁷³, NH₂, NHR⁷³, N(R⁷³)₂, NHC(O)R⁷³, NR⁷³C(O)R⁷³, NHS(O)₂R⁷³, NR⁷³S(O)₂R⁷³, NHC(O)OR⁷³, NR⁷³C(O)OR⁷³, NHC(O)NH₂, NHC(O)NHR⁷³, NHC(O)N(R⁷³)₂, NR⁷³C(O)NHR⁷³, NR⁷³C(O)N(R⁷³)₂, C(O)NH₂, C(O)NHR⁷³, C(O)N(R⁷³)₂, C(O)NHOH, C(O)NHOR⁷³, C(O)NHSO₂R⁷³, C(O)NR⁷³SO₂R⁷³, SO₂NH₂, SO₂NHR⁷³, SO₂N(R⁷³)₂, C(O)H, C(O)OH, C(N)NH₂, C(N)NHR⁷³, C(N)N(R⁷³)₂, CNOH, CNOCH₃, OH, (O), CN, N₃, NO₂, CF₃, CF₂CF₃, OCF₃, OCF₂CF₃, F, Cl, Br and I;

R⁷³ is alkyl, alkenyl, alkynyl, phenyl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl or heterocycloalkenyl; and

wherein the moieties represented by R⁶⁹, R⁷⁰, and R⁷¹ are unsubstituted or substituted with one or two or three or four substituents independently selected from the group consisting of NH₂, C(O)NH₂, C(O)NHOH, SO₂NH₂, CF₃, CF₂CF₃, C(O)H, C(O)OH, C(N)NH₂, OH, (O), CN, N₃, NO₂, OCF₃, OCF₂CF₃, F, Cl, Br and I,

and

wherein the one or more than one additional therapeutic agent(s) are selected from the group consisting of an alkylating agent, an angiogenesis inhibitor, an antibody, an antimetabolite, an antimitotic, an antiproliferative, an antiviral, an aurora kinase inhibitor, an apoptosis promoter, an activator of death receptor pathway, a Bcr-Abl kinase inhibitor, a BiTE (Bi-Specific T cell Engager) antibody, an antibody drug conjugate, a biologic response modifier, a cyclin-dependent kinase inhibitor, a cell cycle inhibitor, a cyclooxygenase-2 inhibitor, a DVD, a leukemia viral oncogene homolog (ErbB2) receptor inhibitor, a growth factor inhibitor, a heat shock protein (HSP)-90 inhibitor, a histone deacetylase (HDAC) inhibitor, a hormonal therapy, an immunological, an inhibitor of inhibitor of apoptosis proteins (IAPs), an intercalating antibiotic, a kinase inhibitor, a kinesin inhibitor, a Jak2 inhibitor, a mammalian target of rapamycin inhibitor, a microRNA, a mitogen-activated extracellular signal-regulated kinase inhibitor, a multivalent binding protein, a non-steroidal anti-inflammatory drug (NSAID), a poly ADP (adenosine diphosphate)-ribose polymerase (PARP) inhibitor, a platinum chemotherapeutic, a polo-like kinase (Plk) inhibitor, a phosphoinositide-3 kinase (PI3K) inhibitor, a proteasome inhibitor, a purine analog, a pyrimidine analog, a receptor tyrosine kinase inhibitor, an etinoids/deltoids plant alkaloid, a small inhibitory ribonucleic acid (siRNA), a topoisomerase inhibitor, and a ubiquitin ligase inhibitor.

2. The combination of claim 1, wherein for the compound having Formula (II)

n is 0;

A¹ is N or C(A²);

A² is H, ; F, Cl, Br, or I;

B¹ is R¹, OR¹, NHR¹, NHC(O)R¹, F, Br, I, or Cl;

D¹ is H, F, Cl, Br, or I;

E¹ is H; and

Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SO₂R¹⁷, NH₂, or C(O)NH₂; or

Y¹ and B¹, together with the atoms to which they are attached, are benzene, heteroarene, heterocycloalkene; and A², D¹ and E¹ are each H;

R¹ is R⁴ or R⁵;

R⁴ is cycloalkyl or heterocycloalkyl;

R⁵ is alkyl or alkynyl, each of which is unsubstituted or substituted with one or two or three substituents independently selected from the group consisting of R⁷, OR⁷, NHR⁷, N(R⁷)₂, CN, OH, F, Cl, Br and I;

R⁷ is R⁸, R⁹, R¹⁰ or R¹¹;

R⁸ is phenyl;

R⁹ is heteroaryl;

R¹⁰ is cycloalkyl, cycloalkenyl, or heterocycloalkyl, each of which is unfused or fused with R^{10A};

R^{10A} is a heteroarene;

R¹¹ is alkyl, which is unsubstituted or substituted with one or two or three substituents independently selected from the group consisting of R¹², OR¹² and CF₃;

R¹² is R¹⁴ or R¹⁶

R¹⁴ is heteroaryl;

R¹⁶ is alkyl;

R¹⁷ is alkyl, alkynyl, each of which is unsubstituted or substituted with one or two or three substituents independently selected from the group consisting of R²², F, Cl, Br and I;

R²² is heterocycloalkyl;

Z² is R³⁰;

L¹ is a R³⁷;

R³⁰ is heterocycloalkylene;

R³⁷ is R^{37A};

R^{37A} is alkylene;

Z³ is R³⁸, or R⁴⁰;

R³⁸ is phenyl;

R⁴⁰ is cycloalkyl, cycloalkenyl, or heterocycloalkenyl;

wherein the cyclic moieties represented Y¹ and B¹ together, R⁴, R⁸, R¹⁰, R²², R³⁰, R³⁸, and R⁴⁰ are independently unsubstituted, substituted or further substituted with one or two or three or four or five substituents independently selected from the group consisting of R^{57A}, R⁵⁷, OR⁵⁷, S(O)R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, NH₂, NHR⁵⁷, N(R⁵⁷)₂, NHC(O)R⁵⁷, NHS(O)₂R⁵⁷, OH, CN, (O), F, Cl, Br and I;

R^{57A} is spiroalkyl or heterospiro alkyl;

R⁵⁷ is R⁵⁸, R⁶⁰ or R⁶¹;

R⁵⁸ is phenyl;

R⁶⁰ is cycloalkyl or heterocycloalkyl;

R⁶¹ is alkyl which is unsubstituted or substituted with one or two or three substituents independently selected from the group consisting of R⁶², OR⁶², N(R⁶²)₂, C(O)OH, CN, F, Cl, Br and I;

R⁶² is R⁶⁵ or R⁶⁶;

R⁶⁵ is cycloalkyl or heterocycloalkyl;

R⁶⁶ is alkyl, which is unsubstituted or substituted with OR⁶⁷;

R⁶⁷ is alkyl;


wherein the cyclic moieties represented by R^{57A}, R⁵⁸, and R⁶⁰ are unsubstituted or substituted with one or two or three or four substituents independently selected from the group consisting of R⁶⁸, F, Cl, Br and I;

R⁶⁸ is R⁷¹ or R⁷²;

R⁷¹ is heterocycloalkyl; and

R⁷² is alkyl, which is unsubstituted or substituted with one or two F.

3. The combination of claim 1, wherein the compound has Formula (III)



(III),

wherein

n, p, r, and q are each 0, and R¹⁰⁰, R¹⁰¹, R¹⁰², and R¹⁰³ are each absent;

A¹ is N or C(A²);

A² is H, F, Cl, Br or I;

B¹ is R¹, OR⁷, NHR¹, NHC(O)R¹, F, Br, I or Cl;

D¹ is H, F, Cl, Br, or I;

E¹ is H; and

Y¹ is H, CN, NO₂, F, Cl, Br, I, CF₃, R¹⁷, OR¹⁷, SR¹⁷, SO₂R¹⁷, or C(O)NH₂; or

Y¹ and B¹, together with the atoms to which they are attached, are benzene, heteroarene, or heterocycloalkene; and

A², D¹, and E¹ are each H;

R¹ is R⁴ or R⁵;

R⁴ is cycloalkyl or heterocycloalkyl;

R⁵ is alkyl or alkynyl, each of which is unsubstituted or substituted with one or two or three substituents independently selected from the group consisting of R⁷, OR⁷, NHR⁷, N(R⁷)₂, CN, OH, F, Cl, Br and I;

R⁷ is R⁸, R⁹, R¹⁰ or R¹¹;

R⁸ is phenyl;

R⁹ is heteroaryl;

R¹⁰ is cycloalkyl, cycloalkenyl, or heterocycloalkyl, each of which is unfused or fused with R^{10A};

R^{10A} is a heteroarene;

R¹¹ is alkyl, which is unsubstituted or substituted with one or two or three substituents independently selected from the group consisting of R¹², OR¹² and CF₃;

R¹² is R¹⁴ or R¹⁶

R¹⁴ is heteroaryl;

R¹⁶ is alkyl;

R¹⁷ is alkyl, alkynyl, each of which is unsubstituted or substituted with one or two or three substituents independently selected from the group consisting of R²², F, Cl, Br and I;

R²² is heterocycloalkyl;

wherein the cyclic moieties represented by Y¹ and B¹ together, R⁴, R⁸, R¹⁰, and R²², are independently unsubstituted, substituted or further substituted with one or two or three or four or five substituents independently selected from the group consisting of R^{57A}, R⁵⁷, OR⁵⁷, SO₂R⁵⁷, C(O)R⁵⁷, CO(O)R⁵⁷, C(O)N(R⁵⁷)₂, NH₂, NHR⁵⁷, N(R⁵⁷)₂, NHC(O)R⁵⁷, NHS(O)₂R⁵⁷, OH, CN, (O), F, Cl, Br, or I;

R^{57A} is spiroalkyl or spiroheteroalkyl;

R⁵⁷ is R⁵⁸, R⁶⁰ or R⁶¹;

R⁵⁸ is phenyl;

R⁶⁰ is cycloalkyl or heterocycloalkyl;

R⁶¹ is alkyl which is unsubstituted or substituted with one or two or three substituents independently selected from the group consisting of R⁶², OR⁶², N(R⁶²)₂, C(O)OH, CN, F, Cl, Br and I;

R⁶² is R⁶⁵ or R⁶⁶;

R⁶⁵ is cycloalkyl or heterocycloalkyl;

R⁶⁶ is alkyl, which is unsubstituted or substituted with OR⁶⁷;

R⁶⁷ is alkyl;

wherein the cyclic moieties represented by R^{57A}, R⁵⁸, and R⁶⁰ are unsubstituted or substituted with one or two or three or four substituents independently selected from the group consisting of R⁶⁸, F, Cl, Br and I;

R⁶⁸ is R⁷¹ or R⁷²;

R⁷¹ is heterocycloalkyl; and

R⁷² is alkyl, which is unsubstituted or substituted with one or two F.

4. The combination of claim 1, wherein the compound is selected from the group consisting of:

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[(4'-chloro-1,1'-biphenyl-2-yl)methyl]piperazin-1-yl}-N-({4-[(3-morpholin-4-ylpropyl)amino]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({3-nitro-4-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-{4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl}-N-({4-[(1-methylpiperidin-4-

yl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methylpiperazin-1-yl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2-methoxyethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(3S)-tetrahydro-2H-pyran-3-ylmethyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(3R)-tetrahydro-2H-pyran-3-ylmethyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-(2-naphthylsulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2-methoxyethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(1,4-dioxan-2-ylmethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(2,2,2-trifluoroethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(3,3,3-trifluoropropyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2S)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(4-methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(2R)-1,4-dioxan-2-ylmethoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(1,4-dioxan-2-ylmethyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(1,4-dioxan-2-ylmethoxy)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(4-[(4-methoxycyclohexyl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-(aminocarbonyl)-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([5-chloro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-([5-chloro-6-(tet-

- rahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-[[4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-(trifluoromethyl)phenyl]sulfonyl]benzamide;
 5 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-[[4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]-3-(trifluoromethyl)sulfonyl]phenyl]sulfonyl]benzamide;
 Trans-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-[[4-[(4-morpholin-4-ylcyclohexyl)amino]-3-(trifluoromethyl)sulfonyl]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-[[4-[(1-methylpiperidin-4-yl)amino]-3-(trifluoromethyl)sulfonyl]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino]sulfonyl]-2-(tetrahydro-2H-pyran-4-ylmethoxy)nicotinamide;
 15 N-[[5-bromo-6-[(1-methylpiperidin-4-yl)methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(1-methylpiperidin-4-yl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-(1,4-dioxan-2-ylmethoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 N-[[5-bromo-6-(1,4-dioxan-2-ylmethoxy)pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(2,2-dimethyltetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 N-[[3-chloro-5-cyano-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[4-[(1-acetyl)piperidin-4-yl]amino]-3-nitrophenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[2-chloro-5-fluoro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(3-morpholin-4-ylpropyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-(2-morpholin-4-ylethoxy)pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[3-chloro-4-[[2-(2-methoxyethoxy)ethyl]sulfonyl]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[2-(2-methoxyethoxy)ethyl]sulfonyl]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(4-morpholin-4-ylcyclohexyl)oxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[5-bromo-6-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]amino]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(2-cyanoethyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Cis-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-[[4-[(4-morpholin-4-ylcyclohexyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-N-[[4-[[4-bis(cyclopropylmethyl)amino]cyclohexyl]amino]-3-nitrophenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(1-methylpiperidin-4-yl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(morpholin-3-ylmethyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-[[4-[(4-methylpiperazin-1-yl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(4-morpholin-4-ylbut-2-ynyl)oxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-ethynyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-oxo-3,4-dihydroquinazolin-6-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 5 Trans-4-(4-[[4-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-[[4-[[4-morpholin-4-ylcyclohexyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Cis-4-(4-[[4-(4-chlorophenyl)-6,6-dimethyl-5,6-dihydro-2H-pyran-3-yl]methyl]piperazin-1-yl)-N-[[4-[[4-methoxycyclohexyl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-[[4-[[4-fluorotetrahydro-2H-pyran-4-yl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 Trans-4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-[[4-[[4-methoxycyclohexyl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- tert-butyl 3-[[4-((4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino]sulfonyl]-2-nitrophenoxymethyl]morpholine-4-carboxylate;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-(morpholin-3-ylmethoxy)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 20 4-(4-[[8-(4-chlorophenyl)spiro[4.5]dec-7-en-7-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-(methylsulfonyl)piperidin-4-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1,1-dioxidotetrahydro-2H-thiopyran-4-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[4-chloro-3-nitrophenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[1-(2,2,2-trifluoroethyl)piperidin-4-yl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-cyano-6-[(1-tetrahydro-2H-pyran-4-yl)piperidin-4-yl]oxy]pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-isopropyl-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 N-[[3-chloro-5-fluoro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[[5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-(2,2-difluoroethyl)piperidin-4-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-(cyclopropyl)piperidin-4-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-(morpholin-4-ylcyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(dicyclopropylamino)cyclohexyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-6,6-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 N-[[5-bromo-6-[(4-ethylmorpholin-3-yl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-ethylmorpholin-3-yl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[(4-tetrahydro-2H-pyran-4-ylmorpholin-3-yl)methoxy]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[3S]-1-tetrahydro-2H-pyran-4-ylpiperidin-3-yl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-cyano-6-[(tetrahydro-2H-pyran-4-ylmethyl)amino]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[1,1-dioxidothiomorpholin-4-yl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 5 N-[[4-[[[4-aminotetrahydro-2H-pyran-4-yl]methyl]amino]-3-nitrophenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-cyano-6-[(4-morpholin-4-ylcyclohexyl)amino]pyridin-3-yl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-cyano-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[1S,3R]-3-morpholin-4-ylcyclopentyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[1R,3S]-3-morpholin-4-ylcyclopentyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(morpholin-2-yl)methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(tetrahydrofuran-3-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({1-[cis-3-fluorotetrahydro-2H-pyran-4-yl]piperidin-4-yl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(1-tetrahydro-2H-pyran-4-ylazetidin-3-yl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[(1-tetrahydrofuran-3-ylazetidin-3-yl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-({[(3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]methyl}amino)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 30 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-4-((trans-4-hydroxycyclohexyl)methoxy)-3-nitrophenyl)sulfonyl)benzamide;
 2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-4-((cis-4-methoxycyclohexyl)methoxy)-3-nitrophenyl)sulfonyl)benzamide;
 35 Cis-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(cyclopropylamino)cyclohexyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(4-tetrahydro-2H-pyran-4-ylamino)cyclohexyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 40 Trans-N-({5-bromo-6-[(4-morpholin-4-ylcyclohexyl)oxy]pyridin-3-yl}sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-methoxycyclohexyl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 tert-butyl 4-[[4-({[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]amino}sulfonyl)-2-nitrophenoxymethyl]-4-fluoropiperidine-1-carboxylate;
 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(4-fluoropiperidin-4-yl)methoxy]-3-nitrophenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[(4-tetrahydro-2H-pyran-4-ylpiperazin-1-yl)cyclohexyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl}methoxy)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-({[(3R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-([[(3R)-1-(2,2-dimethyltetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-([[(3S)-1-tetrahydro-

dro-2H-pyran-4-ylpyrrolidin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3S)-1-(2,2-dimethyl-
 tetrahydro-2H-pyran-4-yl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-
 yloxy)benzamide;
 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[4-methylmorpholin-2-
 yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[4-(2-methoxye-
 thyl)morpholin-2-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 10 N-[[4-[[[4-acetylmorpholin-2-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethyl-
 cyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[trans-4-(fluoromethyl)-
 1-oxetan-3-ylpyrrolidin-3-yl]methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[4-fluorotetrahydro-
 2H-pyran-4-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[(1-oxetan-3-yl-
 piperidin-4-yl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[(1-cyclobutylpiperidin-
 4-yl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-(2,2-dimethyltetrahy-
 20 dro-2H-pyran-4-yl)piperidin-4-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benza-
 mide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3S)-1-cyclopropylpyr-
 rolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[(1-tetrahydro-
 25 furan-3-ylpiperidin-4-yl)amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3R)-1-cyclopropylpyr-
 rolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(3S)-1-tetrahy-
 30 dro-2H-pyran-4-ylpyrrolidin-3-yl]methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benza-
 mide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[3-hydroxy-2,2-dimeth-
 ylpropyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[1-(methylsulfonyl)pip-
 eridin-3-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 35 N-[[4-[[[1-acetyl]piperidin-3-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcy-
 clohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(3R)-1-(methylsulfo-
 nyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-3,3-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[(tetrahydro-2H-
 40 pyran-4-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[1-[2-fluoro-1-(fluor-
 omethyl)ethyl]azetidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[1-(methylsulfonyl)pyr-
 rolidin-3-yl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 45 N-[[4-[[[1-acetyl]pyrrolidin-3-yl)methyl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcy-
 clohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[4-[[[(3R)-1-acetyl]pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcy-
 clohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[3-methoxy-2,2-
 50 dimethylpropyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(1R,3R)-3-hydroxycy-
 clopentyl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(1S,3S)-3-hydroxycy-
 clopentyl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(1S,3R)-3-hydroxycy-
 clopentyl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[[(1R,3S)-3-hydroxycy-
 clopentyl]methyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[{(3S)-2-oxopiperidin-3-yl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[[{1-[2-fluoro-1-(fluoromethyl)ethyl]azetid-3-yl}methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[{(1-oxetan-3-ylazetid-3-yl)methyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[{(1-oxetan-3-ylpiperidin-4-yl)methyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{1-cyclopropylpiperidin-4-yl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{4-(2-fluoroethyl)morpholin-2-yl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{4-(2,2-difluoroethyl)morpholin-2-yl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[[4-fluoro-1-oxetan-3-ylpiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{(2S)-4,4-difluoro-1-oxetan-3-ylpyrrolidin-2-yl)methoxy]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[{(4-tetrahydro-2H-pyran-4-yl)morpholin-3-yl)methyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{4-cyclobutylmorpholin-3-yl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[{(4-tetrahydrofuran-3-yl)morpholin-3-yl)methyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{1-[2-fluoro-1-(fluoromethyl)ethyl]piperidin-4-yl}methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[[{1-cyclopropyl-4-fluoropiperidin-4-yl)methoxy]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
30 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[[4-methoxybenzyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[{3-(trifluoromethoxy)benzyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[[{3-methoxybenzyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{4-(difluoromethoxy)benzyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[1,4-dioxaspiro[4.5]dec-8-ylamino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
40 Trans-N-([4-[[{4-(acetylaminocyclohexyl)amino]-3-nitrophenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{(3S)-1-(2-fluoroethyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{(3S)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{(3R)-1-(2-fluoroethyl)pyrrolidin-3-yl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[(3-nitro-4-[[{(3S)-1-oxetan-3-ylpyrrolidin-3-yl]methoxy}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{4-(hydroxybenzyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{3-(hydroxybenzyl)amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-([4-[[{3-(difluoromethoxy)benzyl]amino]-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[cis-3-morpholin-4-yl-cyclopentyl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[methylsulfo-
 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({[4-({[1-cyclopropylpiperidin-4-yl]amino)-3-[(trifluoromethyl)sulfonyl]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({[3-nitro-4-[(1-oxetan-3-yl-piperidin-4-yl)methoxy]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({[4-[(4-fluoro-1-tetrahydro-
 10 2H-pyran-4-yl]piperidin-4-yl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({[4-[(4-fluoro-1-tetrahydro-furan-3-yl]piperidin-4-yl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[4-fluoro-1-(methylsulfonyl)piperidin-4-yl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-({[(3R)-1-oxetan-3-ylpyrrolidin-3-yl]methyl}amino)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({[4-[(4-hydroxycyclohexyl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[3-(dimethylami-
 20 no)propoxy]benzyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[4-(2-morpholin-4-ylethoxy)benzyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[(E)-4-hydroxy-1-ada-
 25 mantyl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[(Z)-4-hydroxy-1-ada-mantyl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-({[4-({[1S,4S]-bicyclo[2.2.1]hept-5-en-2-ylmethoxy]-3-nitrophenyl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-
 dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({[4-({[1-methyl-5-oxopyrroli-
 30 din-3-yl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[(1R,4R,5R,6S)-5,6-di-hydroxybicyclo[2.2.1]hept-2-yl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benza-
 mide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[(1R,4R,5S,6R)-5,6-di-
 35 hydroxybicyclo[2.2.1]hept-2-yl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benza-mide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({[3-nitro-4-[(3-oxocyclohexyl)methoxy]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5,5-dimethylcyclohexa-1,3-dien-1-yl]methyl]piperazin-1-yl)-N-({[3-nitro-4-[(tetrahy-
 40 dro-2H-pyran-4-yl)methyl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[(3R)-1-[2-fluoro-1-(fluoromethyl)ethyl]pyrrolidin-3-yl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benza-
 mide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[6-[(4-fluorotetrahydro-2H-
 45 pyran-4-yl)methoxy]-5-(trifluoromethyl)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-({[(3S)-1-oxetan-3-ylpyrrolidin-3-yl]methyl}amino)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 Trans-N-({[5-chloro-6-[(4-methoxycyclohexyl)methoxy]pyridin-3-yl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-
 dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 50 Cis-N-({[5-chloro-6-[(4-methoxycyclohexyl)methoxy]pyridin-3-yl]sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimeth-ylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-({[(3S)-1-oxetan-3-ylpyrrolidin-3-yl]amino)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({[4-({[4-(2-methox-
 55 yethoxy)ethyl]morpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benza-mide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-({[4-(cyanomethyl)mor-pholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-((4-(N,N-dimethylglycyl)morpholin-2-yl)methyl)amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 (2-[[4-[[4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino)methyl]morpholin-4-yl)acetic acid;
- 5 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[4-(oxetan-3-yl)morpholin-2-yl]methyl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-cyclopropylmorpholin-2-yl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 10 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[5-(methylsulfonyl)-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-[[trifluoromethyl]sulfonyl]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 15 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-methyltetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 ethyl 4-(4-[[4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)piperazine-1-carboxylate;
- 20 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(morpholin-4-yl)piperidin-1-yl]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[3(R)-1-(oxetan-3-yl)pyrrolidin-3-yl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 25 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[3(R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]amino]-3-[[trifluoromethyl]sulfonyl]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-isopropylpiperidin-4-yl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 30 N-[[4-[[1-tert-butylpiperidin-4-yl]amino]-3-nitrophenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-(2-methoxyethyl)piperidin-3-yl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 35 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[1-(cyanomethyl)piperidin-3-yl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-fluoro-1-methylpiperidin-4-yl)methoxy]-3-[[trifluoromethyl]sulfonyl]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 40 N-[[5-chloro-6-[[3(R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]amino]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 tert-butyl 4-[[4-[[4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl]amino]piperazine-1-carboxylate;
- 45 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-(pentafluoro- λ^6 -sulfonyl)-4-[[tetrahydro-2H-pyran-4-yl)methyl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-methoxytetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 50 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[3(R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[4-(oxetan-3-yl)piperazin-1-yl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 55 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[3-nitro-4-[[3(R)-tetrahydrofuran-3-ylamino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4,4-difluorocyclohexyl]methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[4-[[1-tert-butylpiperidin-4-yl]amino]-3-[[trifluoromethyl]sulfonyl]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-[[4-[[4-(oxetan-3-yl)mor-

pholin-2-yl)methyl}amino)-3-[(trifluoromethyl)sulfonyl]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(5-chloro-6-[[[4-fluorotetrahydro-2H-pyran-4-yl)methyl]amino]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(5-chloro-6-[[1-cyclopropylpiperidin-4-yl]amino]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(5-chloro-6-[[[2S]-4-(cyanomethyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(5-chloro-6-[[[2S]-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(5-chloro-6-[[[2R]-4-(cyanomethyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(5-chloro-6-[[[2R]-4-(N,N-dimethylglycyl)morpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(5-fluoro-6-[[4-fluorotetrahydro-2H-pyran-4-yl]methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(5-chloro-6-[3-hydroxy-2-(hydroxymethyl)-2-methylpropoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(6-[[4-fluorotetrahydro-2H-pyran-4-yl]methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[1,3-difluoropropan-2-yl]morpholin-2-yl)methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(5-chloro-6-[[1-(cyanomethyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[3R]-1-[2-(2-methoxyethoxy)ethyl]pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[3R]-1-(N,N-dimethylglycyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(3-nitro-4-[[1-(oxetan-3-yl)azetidin-3-yl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(5-chloro-6-[[1-(cyanomethyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[2R]-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[2S]-4-(N,N-dimethylglycyl)morpholin-2-yl]methyl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(5-chloro-6-[[1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(5-chloro-6-[[[3R]-1-(2,2-difluoroethyl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[(4-[[[3R]-1-(cyanomethyl)pyrrolidin-3-yl]amino)-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[4'-chlorobiphenyl-2-yl)methyl]-4-methoxypiperidin-1-yl)-N-[(5-chloro-6-[[4-fluorotetrahydro-2H-pyran-4-yl]methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[4'-chlorobiphenyl-2-yl)methyl]-4-methoxypiperidin-1-yl)-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-[(3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino]phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-[(5-chloro-6-[(tetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[9-(4-chlorophenyl)-3-(1,3-difluoropropan-2-yl)-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-[(5-chloro-6-[(tetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-[[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl)methyl]piperazin-1-yl)-N-[(5-chloro-6-(tetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-[(5-chloro-6-[[4-fluoro-1-methylpiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

- N-[(5-chloro-6-[[1-(N,N-dimethylglycyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-fluoropiperidin-1-yl}-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-{4-[(4'-chlorobiphenyl-2-yl)methyl]-4-fluoropiperidin-1-yl}-N-({5-chloro-6-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[9-(4-chlorophenyl)-3-isopropyl-3-azaspiro[5.5]undec-8-en-8-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-({4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}-3-(trifluoromethyl)phenyl)sulfonyl)benzamide;
- N-[(5-chloro-6-[[[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]oxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-nitro-4-[2-(tetrahydrofuran-3-yloxy)ethoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(trans-4-cyanocyclohexyl)methyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(5-chloro-6-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-cyano-6-[2-(tetrahydro-2H-pyran-4-yl)ethoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-(3-furylmethoxy)-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(5-chloro-6-[[[(3R)-1-(1,3-difluoropropan-2-yl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(5-chloro-6-[[[(3R)-1-(2,2-difluoroethyl)pyrrolidin-3-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(5-chloro-6-[[1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-({3-chloro-4-[(4-fluoro-1-methylpiperidin-4-yl)methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({3-cyano-4-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(5-chloro-6-[[1-(2,2-difluoroethyl)-4-fluoropiperidin-4-yl]methoxy]pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-({3-chloro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-({5-chloro-6-[[4,4-difluorocyclohexyl]methoxy]pyridin-3-yl)sulfonyl)-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({6-[[1-(1,3-difluoropropan-2-yl)-4-fluoropiperidin-4-yl]methoxy]-5-(trifluoromethyl)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-chloro-6-[2-(tetrahydrofuran-2-yl)ethoxy]pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]-3-methylpiperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({4-[(3-(cyclopropylamino)propyl]amino}-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(5-chloro-6-(2-methoxyethoxy)pyridin-3-yl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-N-({5-fluoro-6-(tetrahydro-2H-pyran-4-ylmethoxy)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(3-chloro-4-[[1-(methoxyacetyl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- N-[(3-chloro-4-[[1-(N,N-dimethylglycyl)piperidin-4-yl]methoxy]phenyl)sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
- Trans-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohexyl]methyl]piperidin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-

pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([6-(tetrahydro-2H-pyran-4-ylmethoxy)-5-(trifluoromethyl)pyridin-3-yl)sulfonyl]benzamide;
 N-([5-chloro-6-([trans-4-hydroxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-cyano-4-([4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([6-([trans-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([6-([cis-4-methoxycyclohexyl)methoxy]-5-(trifluoromethyl)pyridin-3-yl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-chloro-6-([4,4-difluoro-1-hydroxycyclohexyl)methoxy]pyridin-3-yl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperidin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-([trans-4-(morpholin-4-yl)cyclohexyl)methoxy]phenyl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-[cyclopropyl(1,3-thiazol-5-ylmethyl)amino]propyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-([trans-4-hydroxycyclohexyl)methoxy]phenyl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-chloro-4-([tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-(trifluoromethyl)phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-[cyclopropyl(2,2,2-trifluoroethyl)amino]propyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-([1-(oxetan-3-yl)piperidin-4-yl]methoxy]phenyl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3,5-difluoro-4-([4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-[cyclopropyl(oxetan-3-yl)amino]propyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-([1-(1-methyl-L-prolyl)piperidin-4-yl]methoxy]phenyl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3,4-difluoro-5-([4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([5-chloro-6-([[(2S)-4-cyclopropylmorpholin-2-yl]methoxy]pyridin-3-yl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperidin-1-yl)-N-([3-nitro-4-([tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperidin-1-yl)-N-([3-chloro-4-([tetrahydro-2H-pyran-4-ylmethoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 methyl 2-([4-([4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl)sulfamoyl)-2-nitrophenyl)amino)methyl)morpholine-4-carboxylate;
 2-([4-([4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl)sulfamoyl)-2-nitrophenyl)amino)methyl)-N-ethyl-N-methylmorpholine-4-carboxamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([4-(methylsulfonyl)morpholin-2-yl]methyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([3-[cyclobutyl(cyclopropyl)amino]propyl)amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-nitro-4-([tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-([3-chloro-4-([4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]phenyl)sulfonyl)-4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([3-chloro-4-([tetrahydrofuran-3-ylmethoxy]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-([2-(4-chlorophenyl)-5,5-difluorocyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-([4-([trans-4-hydroxycyclohexyl)methyl]amino]-3-nitrophenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-({3-chloro-4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]phenyl}sulfonyl)-4-(4-{[9-(4-chlorophenyl)-3-(oxetan-3-yl)-3-azaspiro[5.5]undec-8-en-8-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({(2R)-4-cyclopropylmorpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({(2S)-4-cyclopropylmorpholin-2-yl]methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-{[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-([5-chloro-6-({4-[cyclopropyl(oxetan-3-yl)amino]cyclohexyl)methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-{[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl}piperazin-1-yl)-N-[[4-({(4-cyclopropylmorpholin-2-yl)methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-({3-chloro-4-[(4-cyclopropylmorpholin-2-yl)methoxy]phenyl}sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-([3-chloro-4-({(4-cyclopropylmorpholin-2-yl)methyl}amino)phenyl]sulfonyl)-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

2-([2-chloro-4-({4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl}phenyl)amino)methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;

(2S)-2-([3-chloro-5-({4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl}pyridin-2-yl)oxy)methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;

N-([5-chloro-6-({(4-cyclopropylmorpholin-2-yl)methyl}amino)pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

2-([3-chloro-5-({4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl}pyridin-2-yl)amino)methyl]-N-ethyl-N-methylmorpholine-4-carboxamide;

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({(trans-4-hydroxy-4-methylcyclohexyl)methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({(cis-4-hydroxy-4-methylcyclohexyl)methyl}amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-([5-chloro-6-({(1R,2R,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({(2-cyanoethyl)(cyclopropyl)amino}cyclohexyl)amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-([5-chloro-6-({(trans-4-hydroxy-4-methylcyclohexyl)methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[5-chloro-6-(5,6,7,8-tetrahydroimidazo[1,2-a]pyridin-6-yl)methoxy}pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-([5-chloro-6-({(1R,2S,4R,5R)-5-hydroxy-5-methylbicyclo[2.2.1]hept-2-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-([4-({(cis-4-hydroxy-4-methylcyclohexyl)methoxy}-3-nitrophenyl)sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-([5-chloro-6-({4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[5-(4-chlorophenyl)spiro[2.5]oct-5-en-6-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({(3,3-difluoropyrrolidin-1-yl)cyclohexyl]amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-([5-chloro-6-({(trans-4-hydroxy-4-methylcyclohexyl)methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-([5-chloro-6-({(cis-4-hydroxy-4-methylcyclohexyl)methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-[[4-({(2,2-difluorocyclopropyl)amino}cyclohexyl)amino)-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;

N-([5-chloro-6-({(cis-1-fluoro-4-hydroxycyclohexyl)methoxy}pyridin-3-yl)sulfonyl]-4-(4-{[2-(4-chlorophenyl)-4,4-

dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[3-nitro-4-(2-oxaspiro[3.5]non-7-ylmethoxy)phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[4-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5,5-bis(fluoromethyl)cyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[4-[(4-cyclopropylmorpholin-2-yl)methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[5-chloro-6-[(trans-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[5-chloro-6-[(cis-1-fluoro-4-hydroxy-4-methylcyclohexyl)methoxy]pyridin-3-yl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[3-cyano-4-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[4-[[[(trans-4-ethyl-4-hydroxycyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[4-[[[(cis-4-ethyl-4-hydroxycyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5-(methoxymethyl)-5-methylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[3-nitro-4-[[[(2S)-4-(oxetan-3-yl)morpholin-2-yl)methyl]amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[3-chloro-4-[(trans-4-hydroxy-4-methylcyclohexyl)methoxy]phenyl]sulfonyl]-4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[4-[[4-[(2-cyanoethyl)(cyclopropyl)amino]-1-fluorocyclohexyl]methoxy]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[5-nitro-6-[(tetrahydro-2H-pyran-4-ylmethyl)amino]pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[3-nitro-4-[(2-oxaspiro[3.5]non-7-ylmethyl)amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[4-[[[(4-cyano-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-[[3-[(tetrahydro-2H-pyran-4-ylmethyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-6-yl]sulfonyl]benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[6-[[[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino]-5-nitropyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[5-cyano-6-[[4-fluoro-1-(oxetan-3-yl)piperidin-4-yl]methoxy]pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 N-[[4-[[4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]sulfamoyl]-2-nitrophenyl)morpholine-4-carboxamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[4-[[4-(methoxymethyl)cyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[5-chloro-6-[[1-(1,3-thiazol-2-yl)piperidin-4-yl]methoxy]pyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[6-[[[(cis-4-hydroxy-4-methylcyclohexyl)methyl]amino]-5-nitropyridin-3-yl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide;
 4-(4-[[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[4-[[[(trans-4-hydroxy-4-methylcyclohexyl)methyl]amino]-3-nitrophenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide; and
 4-(4-[[2-(4-chlorophenyl)-5-methoxy-5-methylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide.

5. The combination of claim 1, wherein the compound is 4-(4-[[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl]piperazin-1-yl)-N-[[3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl]sulfonyl]-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide.

din-5-yloxy)benzamide.

6. The combination of claim 1, wherein the one or more than one additional therapeutic agent(s) comprise a Bcr-Abl kinase inhibitor, wherein the Bcr-Abl kinase inhibitor is BMS-354825 or imatinib.
7. The combination of claim 1, wherein the one or more than one additional therapeutic agent(s) comprise a PI3K inhibitor, wherein the PI3K inhibitor is selected from the group consisting of wortmannin, LY294002, XL-147, CAL-120, ONC-21, AEZS-127, ETP-45658, PX-866, GDC-0941, BGT226, BEZ235 and XL765.
8. The combination of claim 1, wherein the one or more than one additional therapeutic agent(s) comprise an antibody, wherein the antibody is selected from the group consisting of AVASTIN® (bevacizumab), a CD40-specific antibody, chTNT-1/B, denosumab, ERBITUX® (cetuximab), HUMAX-CD4® (zanolimumab), an IGF1R-specific antibody, linituzumab, PANOREX® (edrecolomab), RENCAREX® (WX G250), RITUXAN® (rituximab), ticilimumab, trastuzumab, and a CD20 antibody type I or II.
9. The combination of claim 1, wherein the one or more than one additional therapeutic agent(s) comprise an alkylating agent, wherein the alkylating agent is selected from the group consisting of altretamine, AMD-473, AP-5280, apaziquone, bendamustine, brostallicin, busulfan, carboquone, carmustine (BCNU), chlorambucil, CLORETAZINE® (laromustine, VNP 40101M), cyclophosphamide, dacarbazine, estramustine, fotemustine, glufosfamide, ifosfamide, KW-2170, lomustine (CCNU), mafosfamide, melphalan, mitobronitol, mitolactol, nimustine, nitrogen mustard N-oxide, ranimustine, temozolomide, thiotepa, TREANDA® (bendamustine), treosulfan, trofosfamide.
10. The combination of claim 1, wherein the one or more than one additional therapeutic agent(s) comprise an antimetabolite, wherein the antimetabolite is selected from the group consisting of ALIMTA® (pemetrexed disodium, LY231514, MTA), 5-azacitidine, XELODA® (capecitabine), carmofur, LEUSTAT® (cladribine), clofarabine, cytarabine, cytarabine ocfosfate, cytosine arabinoside, decitabine, deferoxamine, doxifluridine, eflornithine, EICAR (5-ethynyl-1-P-D-ribofuranosylimidazole-4-carboxamide), enocitabine, ethynylcytidine, fludarabine, 5-fluorouracil alone or in combination with leucovorin, GEMZAR® (gemcitabine), hydroxyuracil, ALKERAN® (melphalan), mercaptopurine, 6-mercaptopurine riboside, methotrexate, mycophenolic acid, nelarabine, nolatrexed, ocfosfate, pelitrexol, pentostatin, raltitrexed, Ribavirin, triapine, trimetrexate, S-1, tiazofurin, tegafur, TS-1, vidarabine, and UFT.
11. The combination of claim 1, wherein the one or more than one additional therapeutic agent(s) comprise a proteasome inhibitor, wherein the proteasome inhibitor is selected from the group consisting of VELCADE® (bortezomib), MG132, NPI-0052 and PR 171.
12. The combination of claim 1, wherein the one or more than one additional therapeutic agent(s) comprise a hormonal therapy, wherein the hormonal therapy is selected from the group consisting of ARIMIDEX® (anastrozole), AROMASIN (exemestane), arzoxifene, CASODEX® (bicalutamide), CETROTIDE® (cetorelix), degarelix, deslorelin, DESOPAN® (trilostane), dexamethasone, DROGENIL® (flutamide), EVISTA® (raloxifene), AFEMA™ (fadrozole), FARESTON® (toremifene), FASLODEX® (fulvestrant), FEMARA® (letrozole), formestane, glucocorticoids, HECTOROL® (doxercalciferol), RENAGEL® (sevelamer carbonate), lasofoxifene, leuprolide acetate, MEGACE® (megestrol), MIFEPREX® (mifepristone), NILANDRON™ (nilutamide), NOLVADEX® (tamoxifen citrate), PLENAXIS™ (abarelix), prednisone, PROPECIA® (finasteride), trilostane, SUPREFACT® (buserelin), TRELSTAR® (luteinizing hormone releasing hormone (LHRH)), VANTAS® (Histrelin implant), VETORYL® (trilostane or modrastane), and ZOLADEX® (fosreltin, goserelin).
13. A combination, comprising 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide, or a pharmaceutically acceptable salt thereof, and a Bcr-Abl kinase inhibitor.
14. The combination of claim 13, wherein the Bcr-Abl kinase inhibitor is BMS-354825 or imatinib.
15. A combination, comprising 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide, or a pharmaceutically acceptable salt thereof, and a phosphoinositide-3 kinase (PI3K) inhibitor.
16. The combination of claim 15, wherein the PI3K inhibitor is selected from the group consisting of wortmannin, LY294002, XL-147, CAL-120, ONC-21, AEZS-127, ETP-45658, PX-866, GDC-0941, BGT226, BEZ235 and XL765.

17. A combination, comprising 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide, or a pharmaceutically acceptable salt thereof, and an antibody.
- 5 18. The combination of claim 17, wherein the antibody is selected from the group consisting of AVASTIN® (bevacizumab), a CD40-specific antibody, chTNT-1/B, denosumab, ERBITUX® (cetuximab), HUMAX-CD4® (zanolimumab), an IGF1R-specific antibody, lintuzumab, PANOREX® (edrecolomab), RENCAREX® (WX G250), RITUXAN® (rituximab), ticilimumab, trastuzumab, and a CD20 antibody type I or II.
- 10 19. The combination of any one of claims 1 to 19, for use in treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, a lymphoid malignancy of T cell or B cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non small cell lung cancer, prostate cancer, small cell lung cancer or spleen cancer in a patient.
- 15 20. A compound or a therapeutically acceptable salt thereof, wherein the compound is selected from the group consisting of
- 20 { [4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzoyl]({4-[(4-fluorotetrahydro-2H-pyran-4-yl)methoxy]-3-nitrophenyl)sulfonyl)amino)methyl butyrate, and
- 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)-N-([3-(tetrahydro-2H-pyran-4-ylmethyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-6-yl]sulfonyl)benzamide.
- 25
- 30
- 35
- 40
- 45
- 50
- 55



EUROPEAN SEARCH REPORT

Application Number
EP 15 16 7892

5

10

15

20

25

30

35

40

45

2

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
A	US 2007/015787 A1 (BRUNCKO MILAN [US] ET AL) 18 January 2007 (2007-01-18) * paragraphs [0001], [0856]; claims 1, 5 * -----	1-20	INV. C07D471/04 C07D491/052 C07D519/00 A61K31/496 A61K31/437
			TECHNICAL FIELDS SEARCHED (IPC)
			C07D
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 10 August 2015	Examiner Moriggi, J
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03/02 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 15 16 7892

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-08-2015

10

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2007015787	A1	18-01-2007	NONE

15

20

25

30

35

40

45

50

55

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61181203 B [0001]
- US 200436770 W [0005] [0309]
- WO 2005049593 A [0005] [0309]
- US 200437911 W [0005] [0309]
- WO 2005024636 A [0005] [0309]
- US 60988479 B [0006] [0311]
- US 11941196 B [0006] [0312]
- WO 1997010223 A [0067]
- WO 2005099353 A [0067]
- WO 1995007271 A [0067]
- WO 2006008754 A [0067]
- US 7538189 B [0067]
- US 7534814 B [0067]
- US 7531685 B [0067]
- US 7528131 B [0067]
- US 7521421 B [0067]
- US 7514068 B [0067]
- US 7511013 B [0067]
- US 20090137457 A [0067]
- US 20090131485 A [0067]
- US 20090131363 A [0067]
- US 20090118238 A [0067]
- US 20090111840 A [0067]
- US 20090105338 A [0067]
- US 20090105307 A [0067]
- US 20090105147 A [0067]
- US 20090093422 A [0067]
- US 20090088416 A [0067]
- US 20090082471 A [0067]
- US 63140409 A [0293]
- WO 12631404 A [0294]
- US 20050101628 A1 [0626]
- WO 2008124878 A [0839] [1182]

Non-patent literature cited in the description

- *Current Allergy and Asthma Reports*, 2003, vol. 3, 378-384 [0006] [0310]
- *British Journal of Haematology*, 2000, vol. 110 (3), 584-90 [0006] [0310]
- *Blood*, 2000, vol. 95 (4), 1283-92 [0006] [0310]
- *New England Journal of Medicine*, 2004, vol. 351 (14), 1409-1418 [0006] [0310]
- C. D. JONES ; M. KASELJ ; R. N. SALVATORE ; W. J. LE NOBLE. *J. Org. Chem.*, 1998, vol. 63, 2758-2760 [0064]
- IUPAC 1974 Recommendations for Section E, Fundamental Stereochemistry. *Pure Appl. Chem.*, 1976, vol. 45, 13-10 [0065]
- LIZONDO, J et al. *Drugs Fut*, 1996, vol. 21 (11), 1116 [0067]
- BRICKNER, S J et al. *J Med Chem*, 1996, vol. 39 (3), 673 [0067]
- MALLESHAM, B et al. *Org Lett*, 2003, vol. 5 (7), 963 [0067]
- BLAKE et al. *J. Pharm. Sci.*, 1975, vol. 64 (3), 367-391 [0068]
- FOSTER et al. *Advances in Drug Research*. Academic press, 1985, vol. 14, 2-36 [0068]
- KATO et al. *J. Labelled Comp. Radiopharmaceut.*, 1995, vol. 36 (10), 927-932 [0068]
- KUSHNER et al. *Can. J. Physiol. Pharmacol.*, 1999, vol. 77, 79-88 [0068]
- CZAJKA D M ; FINKEL A J. *Ann. N.Y. Acad. Sci.*, 1960, vol. 84, 770 [0069]
- THOMSON J F. *Ann. New York Acad. Sci.*, 1960, vol. 84, 736 [0069]
- CZAKJA D M et al. *Am. J. Physiol.*, 1961, vol. 201, 357 [0069]
- Dosimetry & Treatment Planning for Neutron Capture Therapy. *BLAGOJEVIC N et al.* Advanced Medical Publishing. Madison Wis, 1994, 125-134 [0069]
- *Diabetes Metab.*, 1997, vol. 23, 251 [0069]
- V.R. SUTTON ; D.L. VAUX ; J.A. TRAPANI. *J. of Immunology*, 1997, vol. 158 (12), 5783 [0230]
- TSE. *Cancer Research*, 2008, vol. 68 (9), 3421 [0231]
- WANG Z.-X. An Exact Mathematical Expression For Describing Competitive Binding Of Two Different Ligands To A Protein Molecule. *FEBS Lett.*, 1995, vol. 360, 111-4 [0286]
- WANG ZX. An Exact Mathematical Expression For Describing Competitive Binding Of Two Different Ligands To A Protein Molecule. *FEBS Lett.*, 1995, vol. 360, 111-4 [0288]
- *Blood*, 2008, vol. 111, 2300-2309 [0290] [0292]
- *European Journal of Organic Chemistry*, 2008, vol. 5, 895 [1416]

摘要

本发明涉及抑制 Bcl-2 抗凋亡蛋白的活性的化合物、含有该化合物的组合物和治疗疾病的方法，在该疾病期间，抗凋亡 Bcl-2 蛋白被表达。