

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 April 2006 (06.04.2006)

PCT

(10) International Publication Number
WO 2006/036445 A2

(51) International Patent Classification:
A61K 38/17 (2006.01) C07K 14/715 (2006.01)

(21) International Application Number:
PCT/US2005/031100

(22) International Filing Date: 31 August 2005 (31.08.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/612,836 24 September 2004 (24.09.2004) US
60/681,782 17 May 2005 (17.05.2005) US

(71) Applicant (for all designated States except US):
TRUSTEES OF DARTMOUTH COLLEGE [US/US];
Technology Transfer Office, 11 Rope Ferry Road, Hanover,
NH 03755-1404 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ZHANG, Tong**
[CN/US]; 30 Wolf Road, Unit 101, Lebanon, NH 03766
(US). **SENTMAN, Charles, L.** [US/US]; 11 Pearl Street,
West Lebanon, NH 03784 (US).

(74) Agents: **LICATA, Jane, Massey** et al.; Licata & Tyrrel
P.C., 66 E. Main Street, Marlton, NJ 08053 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CHIMERIC NK RECEPTOR AND METHODS FOR TREATING CANCER



(57) Abstract: The present invention relates to chimeric immune receptor molecules for reducing or eliminating tumors. The chimeric receptors are composed a C-type lectin-like natural killer cell receptor, or a protein associated therewith, fused to an immune signaling receptor containing an immunoreceptor tyrosine-based activation motif. Methods for using the chimeric receptors are further provided.

-1-

CHIMERIC NK RECEPTOR AND METHODS FOR TREATING CANCER**Introduction**

5 This invention was made in the course of research sponsored by the National Cancer Institute (Grant No. CA 101748). The U.S. government may have certain rights in this invention.

 This application claims the benefit of priority from
10 U.S. provisional patent application Serial Nos. 60/612,836, filed September 24, 2004 and 60/681,782, filed May 17, 2005, whose contents are incorporated herein by reference in their entireties.

Background of the Invention

15 T cells, especially cytotoxic T cells, play important roles in anti-tumor immunity (Rossing and Brenner (2004) *Mol. Ther.* 10:5-18). Adoptive transfer of tumor-specific T cells into patients provides a means to treat cancer (Sadelain, et al. (2003) *Nat. Rev. Cancer* 3:35-45).
20 However, the traditional approaches for obtaining large numbers of tumor-specific T cells are time-consuming, laborious and sometimes difficult because the average frequency of antigen-specific T cells in periphery is extremely low (Rosenberg (2001) *Nature* 411:380-384; Ho, et al. (2003) *Cancer Cell* 3:431-437; Crowley, et al. (1990) *Cancer Res.* 50:492-498). In addition, isolation and expansion of T cells that retain their antigen specificity and function can also be a challenging task (Sadelain, et al. (2003) *supra*). Genetic modification of primary T cells
25 with tumor-specific immunoreceptors, such as full-length T cell receptors or chimeric T cell receptor molecules can be used for redirecting T cells against tumor cells (Stevens, et al. (1995) *J. Immunol.* 154:762-771; Oelke, et al. (2003) *Nat. Med.* 9:619-624; Stancovski, et al. (1993) *J. Immunol.*

30

-2-

151:6577-6582; Clay, et al. (1999) *J. Immunol.* 163:507-153). This strategy avoids the limitation of low frequency of antigen-specific T cells, allowing for facilitated expansion of tumor-specific T cells to therapeutic doses.

5 Natural killer (NK) cells are innate effector cells serving as a first line of defense against certain viral infections and tumors (Biron, et al. (1999) *Annu. Rev. Immunol.* 17:189-220; Trinchieri (1989) *Adv. Immunol.* 47:187-376). They have also been implicated in the
10 rejection of allogeneic bone marrow transplants (Lanier (1995) *Curr. Opin. Immunol.* 7:626-631; Yu, et al. (1992) *Annu. Rev. Immunol.* 10:189-214). Innate effector cells recognize and eliminate their targets with fast kinetics, without prior sensitization. Therefore, NK cells need to
15 sense if cells are transformed, infected, or stressed to discriminate between abnormal and healthy tissues. According to the missing self phenomenon (Kärre, et al. (1986) *Nature (London)* 319:675-678), NK cells accomplish this by looking for and eliminating cells with aberrant
20 major histocompatibility complex (MHC) class I expression; a concept validated by showing that NK cells are responsible for the rejection of the MHC class I-deficient lymphoma cell line RMA-S, but not its parental MHC class I-positive line RMA.

25 Inhibitory receptors specific for MHC class I molecules have been identified in mice and humans. The human killer cell Ig-like receptors (KIR) recognize HLA-A, -B, or -C; the murine Ly49 receptors recognize H-2K or H-2D; and the mouse and human CD94/NKG2 receptors are
30 specific for Qa1^b or HLA-E, respectively (Long (1999) *Annu. Rev. Immunol.* 17:875-904; Lanier (1998) *Annu. Rev. Immunol.* 16:359-393; Vance, et al. (1998) *J. Exp. Med.* 188:1841-1848).

-3-

Activating NK cell receptors specific for classic MHC class I molecules, nonclassic MHC class I molecules or MHC class I-related molecules have been described (Bakker, et al. (2000) *Hum. Immunol.* 61:18-27). One such receptor is
5 NKG2D (natural killer cell group 2D) which is a C-type lectin-like receptor expressed on NK cells, $\gamma\delta$ -TcR⁺ T cells, and CD8⁺ $\alpha\beta$ -TcR⁺ T cells (Bauer, et al. (1999) *Science* 285:727-730). NKG2D is associated with the transmembrane adapter protein DAP10 (Wu, et al. (1999) *Science* 285:730-
10 732), whose cytoplasmic domain binds to the p85 subunit of the PI-3 kinase.

In humans, two families of ligands for NKG2D have been described (Bahram (2000) *Adv. Immunol.* 76:1-60; Cerwenka and Lanier (2001) *Immunol. Rev.* 181:158-169). NKG2D binds
15 to the polymorphic MHC class I chain-related molecules (MIC)-A and MICB (Bauer, et al. (1999) *supra*). These are expressed on many human tumor cell lines, on several freshly isolated tumor specimens, and at low levels on gut epithelium (Groh, et al. (1999) *Proc. Natl. Acad. Sci. USA*
20 96:6879-6884). NKG2D also binds to another family of ligands designated the UL binding proteins (ULBP)-1, -2, and -3 molecules (Cosman, et al. (2001) *Immunity* 14:123-133; Kubin, et al. (2001) *Eur. J. Immunol.* 31:1428-1437). Although similar to class I MHC molecules in their $\alpha 1$ and
25 $\alpha 2$ domains, the genes encoding these proteins are not localized within the MHC. Like MIC (Groh, et al. (1996) *supra*), the ULBP molecules do not associate with β_2 -microglobulin or bind peptides. The known murine NKG2D-binding repertoire encompasses the retinoic acid early
30 inducible-1 gene products (RAE-1) and the related H60 minor histocompatibility antigen (Cerwenka, et al. (2000) *Immunity* 12:721-727; Diefenbach, et al. (2000) *Nat. Immunol.* 1:119-126). RAE-1 and H60 were identified as

-4-

ligands for mouse NKG2D by expression cloning these cDNA from a mouse transformed lung cell line (Cerwenka, et al. (2000) *supra*). Transcripts of RAE-1 are rare in adult tissues but abundant in the embryo and on many mouse tumor
5 cell lines, indicating that these are oncofetal antigens.

Recombinant receptors containing an intracellular domain for activating T cells and an extracellular antigen-binding domain, which is typically a single-chain fragment of a monoclonal antibody and is specific for a tumor-
10 specific antigen, are known in the art for targeting tumors for destruction. See, e.g., U.S. Patent No. 6,410,319.

Baba et al. ((2000) *Hum. Immunol.* 61:1202-18) teach KIR2DL1-CD3 zeta chain chimeric proteins. Further, WO 02/068615 suggests fusion proteins of NKG2D containing the
15 external domain of NKG2D with a distinct DAP10 interacting domain or with cytoplasmic domains derived from other signaling molecules, for example CD28, for use in engineering cells that respond to NKG2D ligands and potentially create a system with enhanced signaling
20 capabilities.

U.S. Patent No. 5,359,046 discloses a chimeric DNA sequence encoding a membrane bound protein, wherein the chimeric DNA comprises a DNA sequence encoding a signal sequence which directs the membrane bound protein to the
25 surface membrane; a DNA sequence encoding a non-MHC restricted extracellular binding domain of a surface membrane protein selected from the group consisting of CD4, CD8, IgG and single-chain antibody that binds specifically to at least one ligand, wherein said ligand is a protein on
30 the surface of a cell or a viral protein; a transmembrane domain from a protein selected from the group consisting of CD4, CD8, IgG, single-chain antibody, the CD3 zeta chain, the CD3 gamma chain, the CD3 delta chain and the CD3

-5-

epsilon chain; and a cytoplasmic signal-transducing domain of a protein that activates an intracellular messenger system selected from the group consisting of the CD3 zeta chain, the CD3 gamma chain, the CD3 delta chain and the CD3 epsilon chain, wherein the extracellular domain and cytoplasmic domain are not naturally joined together and the cytoplasmic domain is not naturally joined to an extracellular ligand-binding domain, and when the chimeric DNA is expressed as a membrane bound protein in a selected host cell under conditions suitable for expression, the membrane bound protein initiates signaling in the host cell.

Summary of the Invention

The present invention is a nucleic acid construct for expressing a chimeric receptor to reduce or eliminate a tumor. The nucleic acid construct contains a first nucleic acid sequence encoding a promoter operably linked to a second nucleic acid sequence encoding a chimeric receptor protein comprising a C-type lectin-like natural killer cell receptor, or a protein associated therewith, fused to an immune signaling receptor having an immunoreceptor tyrosine-based activation motif of SEQ ID NO:1. In one embodiment, the nucleic acid construct is in a vector. In particular embodiments, the nucleic acid construct further contains a suicide gene.

The present invention also relates to a method for reducing or eliminating tumors. The method involves introducing into an isolated T cell of a patient having or suspected of having a tumor a nucleic acid construct containing a first nucleic acid sequence encoding a promoter operably linked to a second nucleic acid sequence encoding a chimeric receptor protein comprising a C-type lectin-like natural killer cell receptor, or a protein

-6-

associated therewith, fused to an immune signaling receptor having an immunoreceptor tyrosine-based activation motif of SEQ ID NO:1. The T cell is subsequently injected back into the patient so that the chimeric receptor is expressed on the surface of the T cell to activate anti-tumor immunity in the patient thereby reducing or eliminating the tumor.

Brief Description of the Drawings

Figure 1 illustrates chimeric NK receptors exemplified herein. Extracellular (EC), transmembrane (TM), and cytoplasmic (Cyp) portions are indicated. Wild-type (WT) and chimeric (CH) forms of the receptors are indicated, wherein NH₂ denotes the N-terminus and COOH denotes the C-terminus.

Figure 2 shows specific lysis of target cells by gene-modified primary T cells. Effector T cells modified with vector only (shaded diamond), wild-type NKG2D (open square), murine chimeric NKG2D (shaded square), wild-type DAP10 (open triangle), murine chimeric DAP10 (shaded triangle), or wild-type DAP12 (open circle) were co-cultured with target cells RMA (Panel A), RMA/Rae-1 β (Panel B), RMA/H60 (Panel C), YAC-1 (Panel D), or EG7 (Panel E) cells, respectively, at ratios from 1:1 to 25:1 in a 4 hour ⁵¹Cr release assay. The data are presented as mean $\pm$ SD and representative of 3 to 5 independent experiments.

Detailed Description of the Invention

The present invention relates to a chimeric receptor molecule composed of a natural killer cell receptor and an immune signaling receptor expressed on the surface of a T cell to activate killing of a tumor cell. Nucleic acid sequences encoding the chimeric receptor molecule are introduced into a patient's T-cells *ex vivo* and T-cells that express the chimeric receptor molecule are subsequently injected back into the patient. In this

-7-

manner, the chimeric receptor molecules provide a means for the patient's own immune cells to recognize and activate anti-tumor immunity and establish long-term, specific, anti-tumor responses for treating tumors or preventing regrowth of dormant or residual tumor cells. To prevent potential side effects that may occur from uncontrolled inflammation or response against non-tumor tissue, suicide genes are further introduced into the T-cells expressing the chimeric receptor molecule. The suicide gene is activated by administering an agent, specific for the suicide gene, to the patient thereby eliminating all cells expressing the chimeric receptor molecule.

By way of illustration, murine chimeric receptor molecules composed of NKG2D or Dap10 in combination with a N-terminally attached CD3 ζ were generated and expressed in murine T-cells. NKG2D is a type II protein, in which the N-terminus is located intracellularly (Raulet (2003) *Nat. Rev. Immunol.* 3:781-790), whereas the CD3 ζ chain is type I protein with the C-terminus in the cytoplasm (Weissman, et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:9709-9713). To generate a chimeric NKG2D-CD3 ζ fusion protein, an initiation codon ATG was placed ahead of the coding sequence for the cytoplasmic region of the CD3 ζ chain (without a stop codon TAA) followed by a wild-type NKG2D gene. Upon expression, the orientation of the CD3 ζ portion is reversed inside the cells. The extracellular and transmembrane domains are derived from NKG2D. A second chimeric gene encoding the Dap10 gene followed by a fragment coding for the CD3 ζ cytoplasmic domain was also constructed. The structures of the chimeric and wild-type receptors used are diagrammed in Figure 1.

To determine whether murine chimeric NKG2D or murine chimeric Dap10 receptors could be expressed in a similar

-8-

manner as wild-type murine NKG2D or Dapl0, a NKG2D gene with an adaptor protein gene (Dapl0/Dapl2) were co-transfected into Bosc23 cells and NKG2D expression was determined by flow cytometry. To analyze those cells that were transfected, a bicistronic vector with a green fluorescent protein (GFP) gene controlled by an internal ribosome entry site (IRES) was used. NKG2D surface expression was normalized by gating on the GFP⁺ cell population. Like many NK receptors, such as CD94/NKG2C, Ly49D, and Ly49H, NKG2D needs to be associated with adaptor proteins (*i.e.*, Dapl0 and/or Dapl2) for surface expression (Raulet (2003) *supra*; Lanier (2003) *Curr. Opin. Immunol.* 15:308-314). Packaging cell Bosc23 did not express either NKG2D or Dapl0/Dapl2, and transfection with only one of the two components did not give rise to surface expression of NKG2D. However, co-transfection of a NKG2D gene along with an adaptor protein gene led to significant membrane expression of NKG2D. Compared with Dapl2, Dapl0 transfection resulted in higher NKG2D surface expression. Surface expression of NKG2D after association with chimeric Dapl0 adaptor was higher than that with wild-type DAP10. Higher surface expression of NKG2D was also observed after transfection with chimeric NKG2D than with wild-type NKG2D genes, especially when pairing with the Dapl2 gene (>5-fold increase in MFI).

Concentrated, high-titer, retroviral vectors (ecotropic) were used to infect C57BL/6 spleen cells, and NKG2D surface expression was determined by flow cytometry seven days after retroviral transduction. Genetic modification of T cells with wild-type Dapl0, Dapl2 and NKG2D did not significantly increase the surface expression of NKG2D (10-20%) compared to vector alone. In contrast, significantly higher NKG2D expression was observed in T

-9-

cells modified with either chimeric NKG2D (42%) or chimeric Dap10 (64%). In chimeric Dap10-transduced T cells, the surface-expressed NKG2D molecules were only due to endogenous molecules, whereas both endogenous and exogenous NKG2D molecules were responsible for surface expression in chimeric NKG2D-modified T cells. Taken together, these data indicate that chimeric NKG2D and chimeric Dap10 molecules are expressed in a similar manner as the wild-type molecules and that they increase NKG2D expression on T cells.

To assess whether the murine chimeric DAP10 or murine chimeric NKG2D-transduced T cells were capable of recognizing NKG2D ligands, NKG2D ligand-positive tumor cells (RMA/Rae-1 β , RMA/H60 and YAC-1) were used as targets for chimeric NKG2D-bearing T cells. Chimeric DAP10 or chimeric NKG2D-transduced T cells produced high amounts of IFN- γ (20-30 ng/mL) after co-culture with RMA/Rae-1 β , RMA/H60 or YAC-1 cells (Table 1) but not with RMA cells (no ligands), indicating that these chimeric NKG2D-modified T cells could functionally recognize NKG2D ligand-bearing tumor cells.

TABLE 1

Construct	IFN- γ (ng/mL $\pm$ SD)				
	Media	RMA	RMA/Rae-1 β	RMA/H60	YAC-1
Vector Only	0.03 $\pm$ 0.03	0.09 $\pm$ 0.18	0.02 $\pm$ 0.03	0.11 $\pm$ 0.63	0.84 $\pm$ 0.29
Wild-type NKG2D*	0.01 $\pm$ 0.01	0.10 $\pm$ 0.21	0.04 $\pm$ 0.06	0.05 $\pm$ 0.00	1.08 $\pm$ 1.48
Chimeric NKG2D	0.07$\pm$0.08	0.37$\pm$0.34	4.70$\pm$0.78	8.40$\pm$1.60	17.80$\pm$4.60
Wild-type DAP10*	0.01 $\pm$ 0.10	0.11 $\pm$ 0.11	0.04 $\pm$ 0.03	0.09 $\pm$ 0.03	1.43 $\pm$ 1.72
Chimeric DAP10	0.49$\pm$0.55	0.82$\pm$0.52	7.50$\pm$4.40	18.60$\pm$7.60	28.70$\pm$8.30
Wild-Type DAP12 [#]	0.00 $\pm$ 0.01	0.00 $\pm$ 0.01	0.53 $\pm$ 0.67	0.13 $\pm$ 0.10	0.73 $\pm$ 0.09

*p=0.74; [#]p=0.56.

Data are representative of 3 experiments.

-10-

Similarly, chimeric human NKG2D-bearing CD8⁺ T cells secrete IFN- γ when brought into contact with human tumor cells from breast cancer (MCF-7, T47D), prostate cancer (DU145), pancreatic cancer (Pan-1), and melanoma cancer (A375) (Table 2). T cells were cultured with irradiated tumor cells at a 4:1 ratio for 72 hours and IFN- γ was measured by ELISA. T cells cultured without tumor cells functioned as a media only control which produced no detectable IFN- γ . The specificity of the interaction was evident by comparing chimeric NKG2D transduced T cells to vector only.

TABLE 2

Construct	IFN- γ (pg/mL $\pm$ SD)				
	T47D	MCF-7	Panc-1	DU-145	A375
Vector Only	28.9 ($\pm$ 12.5)	53.5 ($\pm$ 3.6)	97.2 ($\pm$ 8.0)	61.4 ($\pm$ 4.2)	262.4 ($\pm$ 44.2)
Wild-type NKG2D*	35.2 ($\pm$ 30.0)	43.6 ($\pm$ 9.2)	115.8 ($\pm$ 89.8)	84.4 ($\pm$ 47.1)	200.5 ($\pm$ 79.5)
Chimeric NKG2D	130.3 ($\pm$70.4)	2928.3 ($\pm$251.1)	5028.1 ($\pm$407.2)	4427.9 ($\pm$470.1)	2609.2 ($\pm$293.2)
Tumor Alone	23.4 ($\pm$ 26.9)	17.7 ($\pm$ 8.7)	26.6 ($\pm$ 7.5)	27.4 ($\pm$ 10.2)	11.4 ($\pm$ 17.7)

In addition, upon NKG2D ligation, chimeric DAP10 or chimeric NKG2D-modified T cells also released significant amounts of proinflammatory chemokines (CCL3 and CCL5), as well as Th1 cytokines, GM-CSF and IL-3, but not Th2 cytokines IL-5 and IL-10. In contrast, wild-type Dap10, Dap12 or NKG2D alone-modified T cells did not show any significant response to the stimulation by RMA/Rae-1 β , RMA/H60 or YAC-1 cells. These data demonstrate that the chimeric molecules led to the direct activation of T cells.

The cytotoxic activity of murine chimeric NKG2D-modified splenic T cells against tumor cells was also determined. Chimeric Dap10 or chimeric NKG2D-transduced T

-11-

cells were able to lyse NKG2D ligand-expressing target cells (RMA/Rae-1 β , RMA/H60, EG7 and YAC-1) *in vitro* (Figure 2, Panels B-E). The specificity of the interaction was apparent from the absence of lysis of YAC-1, EG7, RMA/Rae-1 β and RMA/H60 cells by vector only-transduced T cells, and the lack of lysis of RMA cells by chimeric Dap10 or chimeric NKG2D-modified T cells (Figure 2, Panel A). Similar to cytokine production, no significant specific lysis of tumor cells was observed by wild-type Dap10 or wild-type NKG2D-modified T cells. T cells transduced with wild-type Dap12 were able to kill target cells that expressed ligands for NKG2D. Activated murine CD8+ T cells express NKG2D (associated with Dap10), so expression of Dap12 would allow the endogenous NKG2D to associate with Dap12 and provide a primary activation signal. It is noteworthy that T cells transduced with Dap12 were three- to five-fold less efficient than T cells transduced with chimeric NK receptors at killing tumor cells. The killing of YAC-1 and EG7 tumor cells demonstrates that chimeric NK receptors provide the T cells with a means to kill tumor cells that express endogenous NKG2D ligands.

These data demonstrated the need for NKG2D ligand expression on the target cells. To investigate the role of the NKG2D receptor, it was determined whether blocking antibodies to NKG2D would diminish cytotoxic activity. Chimeric NKG2D-transduced T cells killed RMA/Rae-1 β and EG7 tumor cells and this activity was reduced when anti-NKG2D antibodies were included in the assay. Vector only-transduced T cells were unable to kill the target cells and the activity was not changed with the addition of anti-NKG2D antibodies. While the data indicate that the NKG2D receptor was responsible for the activity in these assays, the chimeric receptors may have, in some way, altered the T

-12-

cells to kill via their T cell receptor. To address this, the ability of chimeric NKG2D-transduced T cells to kill RMA/Rae-1 β tumor cells was examined. RMA-S cells are deficient in TAP genes and express very low levels of MHC class I molecules on the cell surface and no MHC class II molecules (Aldrich, et al. (1992) *J. Immunol.* 149:3773-3777). Chimeric NKG2D-bearing T cells killed RMA/Rae-1 β tumor cells but not RMA-S cells. Vector-transduced T cells did not kill either RMA-S cell line. Thus, these data indicate that chimeric NKG2D functions via direct NKG2D recognition of its ligand on target cells.

Having shown that chNKG2D-modified T cells could react against NKG2D ligand-positive tumor cells *in vitro*, the therapeutic potential of chimeric NKG2D-modified T lymphocytes was determined *in vivo*. Chimeric NKG2D-bearing T cells (10^6) were co-injected with RMA/Rae-1 β tumor cells (10^5) subcutaneously to C57BL/6 mice. T cells transduced with the chimeric NKG2D construct significantly ($P < 0.05$ at days 5-15) inhibited the growth of RMA/Rae-1 β tumors compared with vector-transduced T cells or tumor alone (Table 3). Approximately 36% (4/11) of chimeric NKG2D-bearing T cell-treated mice were tumor-free after 30 days. Chimeric NKG2D-bearing T cells did not show any significant inhibition effects on the growth of wild-type RMA cells, indicating that inhibition of RMA/Rae-1 β tumor growth by chimeric NKG2D T cells was mediated by chimeric NKG2D-Rae-1 β engagement.

TABLE 3

Day	Tumor Area (Mean mm ² $\pm$ SEM)		
	T Cells transduced with chimeric NKG2D + RMA/Rae-1 β	T cells transduced with vector only + RMA/Rae-1 β	RMA/Rae-1 β
0	0.00 $\pm$ 0.00	0.00 $\pm$ 0.00	0.00 $\pm$ 0.00
5	0.00 $\pm$ 0.00	16.14 $\pm$ 2.86	6.79 $\pm$ 1.47

-13-

7	3.10±1.40	38.45±3.79	28.69±5.49
9	8.11±3.09	57.40±6.43	42.22±6.38
11	11.84±5.24	90.31±11.64	60.60±12.10
13	14.73±7.24	127.30±16.85	82.67±19.44
15	20.60±8.32	N.D.	110.51±29.07

Results are a summary of three experiments.

In a second and more stringent model, transduced T cells (10^7) were adoptively transferred *i.v.* into B6 mice one day before *s.c.* tumor inoculation in the right flank. These chimeric NKG2D-bearing T cells significantly ($P < 0.05$ at days 9-17) suppressed the growth of RMA/Rae-1 β tumors (*s.c.*) compared with control vector-modified T cells (Table 4). As for the toxicity of treatment with chimeric NKG2D-modified T cells, the animals treated with chimeric NKG2D-bearing T cells did not show any overt evidence of inflammatory damage (*i.e.*, ruffled hair, hunchback or diarrhea, etc.) indicating there was no overt toxicity.

TABLE 4

Day	Tumor Area (Mean mm ² ± SEM)	
	T Cells transduced with chimeric NKG2D	Control T cells with Vector Only
5	3.06±1.97	4.41±2.20
7	12.14±3.06	17.81±1.75
9	13.94±2.85	30.58±3.87
11	25.92±4.77	45.13±3.27
13	32.11±5.84	64.83±10.45
15	34.39±9.77	80.72±13.34
17	37.81±11.68	96.30±14.15

Results are a summary of three experiments.

15

Because the immune system can select for tumor variants, the most effective immunotherapies for cancer are likely going to be those that induce immunity against multiple tumor antigens. Thus, it was tested whether

-14-

treatment with chimeric NKG2D-bearing T cells could induce host immunity against wild-type tumor cells. Mice that were treated with chimeric NKG2D-bearing T cells and RMA/Rae-1 β tumor cells, and were tumor-free after 30 days, were challenged with RMA tumor cells. These tumor-free mice were resistant to a subsequent challenge of wild-type RMA cells (10^4), whereas all control naïve mice had aggressive tumors (tumor area: ~100 mm²) after 2 weeks (Table 5). This observation indicates that adoptive transfer of chimeric NKG2D-bearing T cells allows hosts to generate T cell memory.

TABLE 5

Day	Tumor Area (Mean mm ² $\pm$ SEM)	
	Mice treated with T Cells transduced with chimeric NKG2D + RMA/Rae-1 β	Naïve Mice
5	0.00 $\pm$ 0.00	1.33 $\pm$ 2.31
7	0.00 $\pm$ 0.00	11.65 $\pm$ 10.20
9	0.00 $\pm$ 0.00	38.75 $\pm$ 8.84
11	0.00 $\pm$ 0.00	60.17 $\pm$ 6.10
13	0.00 $\pm$ 0.00	91.10 $\pm$ 5.59
15	0.00 $\pm$ 0.00	102.81 $\pm$ 17.94
19	0.00 $\pm$ 0.00	146.71 $\pm$ 45.72

Results are a summary of three experiments.

In similar experiments, human chimeric receptor molecules composed of NKG2D or Dap10 in combination with a N-terminally attached CD3 ζ were generated and expressed in Bosc23 cells. Surface expression of NKG2D was not observed when either human Dap10 or human chimeric NKG2D were transfected alone. However, co-transfection of a human chimeric NKG2D or human chimeric NKG2D-GFP gene along with a wild-type human DAP10 gene or mouse DAP10-GFP construct led to significant membrane expression of NKG2D.

-15-

Binding of a human NKG2D fusion protein, composed of NKG2D with an N-terminally attached murine IgG1 Fc portion, to human NKG2D ligand on various tumor cell lines was assessed. Human NKG2D ligand was found to be present on
 5 Jurkat (T lymphocyte origin), RPMI8866 (B cell origin), K562 (erythroid origin), Daubi (B cell origin), and U937 (monocyte origin) tumor cell lines. Therefore, like the mouse chimeras, a human chimeric NKG2D construct can functionally recognize NKG2D ligand-bearing tumor cells.

10 The cytotoxic activity of human chimeric NKG2D-modified T cells against tumor cells was also determined. Human chimeric NKG2D-transduced primary human T cells were able to lyse mastocytoma cell line P815 transduced with human MIC-A (P815/MICA-A) *in vitro* (Table 6). The
 15 specificity of the interaction was apparent from the absence of lysis of wild-type P815 tumor cells and the absence of lysis by vector only-transduced T cells.

TABLE 6

Tumor Cell Line	Specific Lysis (%)					
	P815/MIC-A			P815		
<i>Effector:Target Ratio</i>	<i>1</i>	<i>5</i>	<i>25</i>	<i>1</i>	<i>5</i>	<i>25</i>
Vector Only	0.0	0.0	0.6	0.0	-0.6	0.0
Human chimeric NKG2D	5.4	17.3	35.4	-2.5	-0.3	1.1

20 Further, blocking of human chimeric NKG2D with an anti-NKG2D antibody prevented killing of K562 and RPMI8866 tumor cells by chimeric NKG2D-transduced human T cells (Table 7). These data demonstrate receptor specificity because a control antibody could not prevent killing.

-16-

TABLE 7

Tumor Cell Line	Specific Lysis (%)					
	K562			RPMI8866		
Effector:Target Ratio	1	5	25	1	5	25
Vector + control antibody	0.0	0.3	2.3	0.0	1.0	0.0
Vector + anti-NKG2D antibody	0.0	0.7	2.1	0.0	1.0	1.3
Human chimeric NKG2D + control antibody	3.1	17.6	53.8	9.3	27.4	41.6
Human chimeric NKG2D + anti-NKG2D antibody	1.6	10.1	27.0	1.0	2.5	7.5

Similar to the mouse studies, human chimeric NKG2D-transduced T cells produced high amounts of IFN- γ (~150-2250 pg/mL) after a 24 hour co-culture with with tumor cells that express ligands for NKG2D (i.e., Jurkat, RPMI8866, K652, ECC-01 and P815/MIC-A tumor cells) compared with tumor cells that do not express NKG2D ligands (P815) or T cells incubated alone (Table 8). Vector only-transduced T cells did not produce IFN- γ , except against RPMI8866, indicating another ligand on this cell type for these activated T cells; however, IFN- γ production was almost 10-times as high with the human chimeric NKG2D-bearing T cells. Tumor cells alone produce no detectable IFN- γ .

TABLE 8

Tumor cell Type	IFN- γ (Mean pg/mL $\pm$ SD)		
	Chimeric NKG2D	Vector Only	Tumor cell control
P815	30.32 $\pm$ 1.31	15.11 $\pm$ 0.94	5.53 $\pm$ 1.31
P815/MIC-A	140.19 $\pm$ 5.91	9.19 $\pm$ 3.30	2.85 $\pm$ 1.50
Jurkat	182.66 $\pm$ 18.31	7.64 $\pm$ 1.04	2.83 $\pm$ 1.33
RPMI8866	2239.95 $\pm$ 19.59	280.41 $\pm$ 13.84	2.47 $\pm$ 2.47
K652	2305.46 $\pm$ 75.84	1.91 $\pm$ 0.57	1.82 $\pm$ 1.39
ECC-1	469.97 $\pm$ 18.79	2.67 $\pm$ 2.67	0.00 $\pm$ 0.00
T Cells only	13.67 $\pm$ 2.55	0.94 $\pm$ 0.54	N.D.

Amounts represent the average of three experiments

-17-

Thus, having demonstrated the activation of host anti-tumor immunity and tumor elimination using chimeric NK cell receptors expressed in the T cells of an animal model of cancer and likewise demonstrated human tumor cell killing with human chimeric NK cell receptors, the present invention relates to a nucleic acid construct for expressing a chimeric receptor in host T cells to reduce or eliminate a tumor. The nucleic acid construct contains a first nucleic acid sequence encoding a promoter operably linked to a second nucleic acid sequence encoding a chimeric receptor protein containing a C-type lectin-like natural killer cell receptor, or a protein associated therewith, fused to an immune signaling receptor having an immunoreceptor tyrosine-based activation motif of SEQ ID NO:1. In general, the C-type lectin-like NK cell type II receptor (or a protein associated therewith) is located at the C-terminus of the chimeric receptor protein of the present invention whereas the immune signaling receptor is at the N-terminus, thereby facilitating intracellular signal transduction from the C-type lectin-like NK cell type II receptor.

A C-type lectin-like NK cell receptor protein particularly suitable for use in the chimeric receptor of the present invention includes a receptor expressed on the surface of natural killer cells, wherein upon binding to its cognate ligand(s) it alters NK cell activation. The receptor can work alone or in concert with other molecules. Ligands for these receptors are generally expressed on the surface of one or more tumor cell types, e.g., tumors associated with cancers of the colon, lung, breast, kidney, ovary, cervix, and prostate; melanomas; myelomas; leukemias; and lymphomas (Wu, et al. (2004) *J. Clin. Invest.* 114:60-568; Groh, et al. (1999) *Proc. Natl. Acad.*

-18-

Sci. USA 96:6879-6884; Pende, et al. (2001) *Eur. J. Immunol.* 31:1076-1086) and are not widely expressed on the surface of cells of normal tissues. Examples of such ligands include, but are not limited to, MIC-A, MIC-B, heat shock proteins, ULBP binding proteins (e.g., ULPBs 1-4), and non-classical HLA molecules such as HLA-E and HLA-G, whereas classical MHC molecules such as HLA-A, HLA-B, or HLA-C and alleles thereof are not generally considered strong ligands of the C-type lectin-like NK cell receptor protein of the present invention. C-type lectin-like NK cell receptors which bind these ligands generally have a type II protein structure, wherein the N-terminal end of the protein is intracellular. Exemplary NK cell receptors of this type include, but are not limited to, Dectin-1 (GENBANK accession number AJ312373 or AJ312372), Mast cell function-associated antigen (GENBANK accession number AF097358), HNKRP-1A (GENBANK accession number U11276), LLT1 (GENBANK accession number AF133299), CD69 (GENBANK accession number NM_001781), CD69 homolog, CD72 (GENBANK accession number NM_001782), CD94 (GENBANK accession number NM_002262 or NM_007334), KLRF1 (GENBANK accession number NM_016523), Oxidised LDL receptor (GENBANK accession number NM_002543), CLEC-1, CLEC-2 (GENBANK accession number NM_016509), NKG2D (GENBANK accession number BC039836), NKG2C (GENBANK accession number AJ001684), NKG2A (GENBANK accession number AF461812), NKG2E (GENBANK accession number AF461157), WUGSC:H_DJ0701016.2, or Myeloid DAP12-associating lectin (MDL-1; GENBANK accession number AJ271684). In particular embodiments, the NK cell receptor is human NKG2D (SEQ ID NO:2) or human NKG2C (SEQ ID NO:3).

Similar type I receptors which would be useful in the chimeric receptor of the present invention include NKp46 (e.g., GENBANK accession number AJ001383), NKp30 (e.g.,

-19-

GENBANK accession number AB055881), or NKp44 (e.g., GENBANK accession number AJ225109).

As an alternative to the C-type lectin-like NK cell receptor protein, a protein associated with a C-type
5 lectin-like NK cell receptor protein can be used in the chimeric receptor protein of the present invention. In general, proteins associated with C-type lectin-like NK cell receptor are defined as proteins that interact with the receptor and transduce signals therefrom. Suitable
10 human proteins which function in this manner include, but are not limited to DAP10 (e.g., GENBANK accession number AF072845; SEQ ID NO:4) and DAP12 (e.g., GENBANK accession number AF019562; SEQ ID NO:5).

To the N-terminus of the C-type lectin-like NK cell
15 receptor is fused an immune signaling receptor having an immunoreceptor tyrosine-based activation motif (ITAM), (Asp/Glu)-Xaa-Xaa-Tyr*-Xaa-Xaa-(Ile/Leu)-Xaa₆₋₈-Tyr*-Xaa-Xaa-(Ile/Leu) (SEQ ID NO:1) which is involved in the activation of cellular responses via immune receptors.
20 Similarly, when employing a protein associated with a C-type lectin-like NK cell receptor, an immune signaling receptor can be fused to the C-terminus of said protein (Figure 1). Suitable immune signaling receptors for use in the chimeric receptor of the present invention include, but
25 are not limited to, the zeta chain of the T-cell receptor, the eta chain which differs from the zeta chain only in its most C-terminal exon as a result of alternative splicing of the zeta mRNA, the delta, gamma and epsilon chains of the T-cell receptor (CD3 chains) and the gamma subunit of the
30 FcR1 receptor. In particular embodiments, the immune signaling receptor is CD3-zeta (CD3ζ) (e.g., GENBANK accession number human NM_198053; SEQ ID NO:6), or human Fc epsilon receptor-gamma chain (e.g., GENBANK accession

-20-

number M33195; SEQ ID NO:7) or the cytoplasmic domain or a splicing variant thereof.

In particular embodiments, a chimeric receptor of the present invention is a fusion between NKG2D and CD3 ζ or
5 Dap10 and CD3 ζ .

As used herein, a nucleic acid construct or nucleic acid sequence is intended to mean a DNA molecule which can be transformed or introduced into a T cell and be transcribed and translated to produce a product (e.g., a
10 chimeric receptor or a suicide protein).

In the nucleic acid construct of the present invention, the promoter is operably linked to the nucleic acid sequence encoding the chimeric receptor of the present invention, i.e., they are positioned so as to promote
15 transcription of the messenger RNA from the DNA encoding the chimeric receptor. The promoter can be of genomic origin or synthetically generated. A variety of promoters for use in T cells are well-known in the art (e.g., the CD4 promoter disclosed by Marodon, et al. (2003) *Blood*
20 101(9):3416-23). The promoter can be constitutive or inducible, where induction is associated with the specific cell type or a specific level of maturation. Alternatively, a number of well-known viral promoters are also suitable. Promoters of interest include the β -actin promoter, SV40
25 early and late promoters, immunoglobulin promoter, human cytomegalovirus promoter, retrovirus promoter, and the Friend spleen focus-forming virus promoter. The promoters may or may not be associated with enhancers, wherein the enhancers may be naturally associated with the particular
30 promoter or associated with a different promoter.

The sequence of the open reading frame encoding the chimeric receptor can be obtained from a genomic DNA source, a cDNA source, or can be synthesized (e.g., via

-21-

PCR), or combinations thereof. Depending upon the size of the genomic DNA and the number of introns, it may be desirable to use cDNA or a combination thereof as it is found that introns stabilize the mRNA or provide T cell-specific expression (Barthel and Goldfeld (2003) *J. Immunol.* 171(7):3612-9). Also, it may be further advantageous to use endogenous or exogenous non-coding regions to stabilize the mRNA.

For expression of a chimeric receptor of the present invention, the naturally occurring or endogenous transcriptional initiation region of the nucleic acid sequence encoding N-terminal component of the chimeric receptor can be used to generate the chimeric receptor in the target host. Alternatively, an exogenous transcriptional initiation region can be used which allows for constitutive or inducible expression, wherein expression can be controlled depending upon the target host, the level of expression desired, the nature of the target host, and the like.

Likewise, the signal sequence directing the chimeric receptor to the surface membrane can be the endogenous signal sequence of N-terminal component of the chimeric receptor. Optionally, in some instances, it may be desirable to exchange this sequence for a different signal sequence. However, the signal sequence selected should be compatible with the secretory pathway of T cells so that the chimeric receptor is presented on the surface of the T cell.

Similarly, a termination region can be provided by the naturally occurring or endogenous transcriptional termination region of the nucleic acid sequence encoding the C-terminal component of the chimeric receptor. Alternatively, the termination region can be derived from a

-22-

different source. For the most part, the source of the termination region is generally not considered to be critical to the expression of a recombinant protein and a wide variety of termination regions can be employed without adversely affecting expression.

As will be appreciated by one of skill in the art, in some instances, a few amino acids at the ends of the C-type lectin-like natural killer cell receptor (or protein associated therewith) or immune signaling receptor can be deleted, usually not more than 10, more usually not more than 5 residues. Also, it may be desirable to introduce a small number of amino acids at the borders, usually not more than 10, more usually not more than 5 residues. The deletion or insertion of amino acids will usually be as a result of the needs of the construction, providing for convenient restriction sites, ease of manipulation, improvement in levels of expression, or the like. In addition, the substitute of one or more amino acids with a different amino acid can occur for similar reasons, usually not substituting more than about five amino acids in any one domain.

The chimeric construct, which encodes the chimeric receptor according to this invention can be prepared in conventional ways. Since, for the most part, natural sequences are employed, the natural genes are isolated and manipulated, as appropriate (e.g., when employing a Type II receptor, the immune signaling receptor component may have to be inverted), so as to allow for the proper joining of the various components. Thus, the nucleic acid sequences encoding for the N-terminal and C-terminal proteins of the chimeric receptor can be isolated by employing the polymerase chain reaction (PCR), using appropriate primers which result in deletion of the undesired portions of the

-23-

gene. Alternatively, restriction digests of cloned genes can be used to generate the chimeric construct. In either case, the sequences can be selected to provide for restriction sites which are blunt-ended, or have
5 complementary overlaps.

The various manipulations for preparing the chimeric construct can be carried out *in vitro* and in particular embodiments the chimeric construct is introduced into vectors for cloning and expression in an appropriate host
10 using standard transformation or transfection methods. Thus, after each manipulation, the resulting construct from joining of the DNA sequences is cloned, the vector isolated, and the sequence screened to insure that the sequence encodes the desired chimeric receptor. The
15 sequence can be screened by restriction analysis, sequencing, or the like.

The chimeric constructs of the present invention find application in subjects having or suspected of having cancer by reducing the size of a tumor or preventing the
20 growth or regrowth of a tumor in these subjects. Accordingly, the present invention further relates to a method for reducing growth or preventing tumor formation in a subject by introducing a chimeric construct of the present invention into an isolated T cell of the subject
25 and reintroducing into the subject the transformed T cell thereby effecting anti-tumor responses to reduce or eliminate tumors in the subject. Suitable T cells which can be used include, cytotoxic lymphocytes (CTL), tumor-infiltrating-lymphocytes (TIL) or other cells which are
30 capable of killing target cells when activated. As is well-known to one of skill in the art, various methods are readily available for isolating these cells from a subject. For example, using cell surface marker expression or using

-24-

commercially available kits (e.g., ISOCELL™ from Pierce, Rockford, IL).

While the present invention relates to the elimination of tumors, the chimeric NK receptors of the present invention may also be useful for treatment of other diseases where these ligands may be present. For example, the immune response can be down-modulated during autoimmune disease or transplantation by expressing these type of chimeric NK receptors in T regulatory or T suppressor cells. Thus, these cells would mediate their regulatory/suppressive function only in the location where the body has upregulated one of the ligands for these receptors. This ligand upregulation may occur during stress or inflammatory responses.

It is contemplated that the chimeric construct can be introduced into the subject's own T cells as naked DNA or in a suitable vector. Methods of stably transfecting T cells by electroporation using naked DNA are known in the art. See, e.g., U.S. Patent No. 6,410,319. Naked DNA generally refers to the DNA encoding a chimeric receptor of the present invention contained in a plasmid expression vector in proper orientation for expression. Advantageously, the use of naked DNA reduces the time required to produce T cells expressing the chimeric receptor of the present invention.

Alternatively, a viral vector (e.g., a retroviral vector, adenoviral vector, adeno-associated viral vector, or lentiviral vector) can be used to introduce the chimeric construct into T cells. Suitable vectors for use in accordance with the method of the present invention are non-replicating in the subject's T cells. A large number of vectors are known which are based on viruses, where the copy number of the virus maintained in the cell is low

-25-

enough to maintain the viability of the cell. Illustrative vectors include the pFB-neo vectors (STRATAGENE®) disclosed herein as well as vectors based on HIV, SV40, EBV, HSV or BPV.

5 Once it is established that the transfected or transduced T cell is capable of expressing the chimeric receptor as a surface membrane protein with the desired regulation and at a desired level, it can be determined whether the chimeric receptor is functional in the host
10 cell to provide for the desired signal induction (e.g., production of Rantes, Mip1-alpha, GM-CSF upon stimulation with the appropriate ligand).

Subsequently, the transduced T cells are reintroduced or administered to the subject to activate anti-tumor
15 responses in said subject. To facilitate administration, the transduced T cells according to the invention can be made into a pharmaceutical composition or made implant appropriate for administration *in vivo*, with appropriate carriers or diluents, which further can be pharmaceutically
20 acceptable. The means of making such a composition or an implant have been described in the art (see, for instance, Remington's Pharmaceutical Sciences, 16th Ed., Mack, ed. (1980)). Where appropriate, the transduced T cells can be formulated into a preparation in semisolid or liquid form,
25 such as a capsule, solution, injection, inhalant, or aerosol, in the usual ways for their respective route of administration. Means known in the art can be utilized to prevent or minimize release and absorption of the composition until it reaches the target tissue or organ, or
30 to ensure timed-release of the composition. Desirably, however, a pharmaceutically acceptable form is employed which does not ineffectuate the cells expressing the chimeric receptor. Thus, desirably the transduced T cells

-26-

can be made into a pharmaceutical composition containing a balanced salt solution, preferably Hanks' balanced salt solution, or normal saline.

A pharmaceutical composition of the present invention
5 can be used alone or in combination with other well-established agents useful for treating cancer. Whether delivered alone or in combination with other agents, the pharmaceutical composition of the present invention can be delivered via various routes and to various sites in a
10 mammalian, particularly human, body to achieve a particular effect. One skilled in the art will recognize that, although more than one route can be used for administration, a particular route can provide a more immediate and more effective reaction than another route.
15 For example, intradermal delivery may be advantageously used over inhalation for the treatment of melanoma. Local or systemic delivery can be accomplished by administration comprising application or instillation of the formulation into body cavities, inhalation or insufflation of an
20 aerosol, or by parenteral introduction, comprising intramuscular, intravenous, intraportal, intrahepatic, peritoneal, subcutaneous, or intradermal administration.

A composition of the present invention can be provided in unit dosage form wherein each dosage unit, e.g., an
25 injection, contains a predetermined amount of the composition, alone or in appropriate combination with other active agents. The term unit dosage form as used herein refers to physically discrete units suitable as unitary dosages for human and animal subjects, each unit containing
30 a predetermined quantity of the composition of the present invention, alone or in combination with other active agents, calculated in an amount sufficient to produce the desired effect, in association with a pharmaceutically

-27-

acceptable diluent, carrier, or vehicle, where appropriate. The specifications for the novel unit dosage forms of the present invention depend on the particular pharmacodynamics associated with the pharmaceutical composition in the particular subject.

Desirably an effective amount or sufficient number of the isolated transduced T cells is present in the composition and introduced into the subject such that long-term, specific, anti-tumor responses are established to reduce the size of a tumor or eliminate tumor growth or regrowth than would otherwise result in the absence of such treatment. Desirably, the amount of transduced T cells reintroduced into the subject causes a 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 98%, or 100% decrease in tumor size when compared to otherwise same conditions wherein the transduced T cells are not present.

Accordingly, the amount of transduced T cells administered should take into account the route of administration and should be such that a sufficient number of the transduced T cells will be introduced so as to achieve the desired therapeutic response. Furthermore, the amounts of each active agent included in the compositions described herein (e.g., the amount per each cell to be contacted or the amount per certain body weight) can vary in different applications. In general, the concentration of transduced T cells desirably should be sufficient to provide in the subject being treated at least from about 1×10^6 to about 1×10^9 transduced T cells, even more desirably, from about 1×10^7 to about 5×10^8 transduced T cells, although any suitable amount can be utilized either above, e.g., greater than 5×10^8 cells, or below, e.g., less than 1×10^7 cells. The dosing schedule can be based on well-established cell-based therapies (see, e.g., Topalian and

-28-

Rosenberg (1987) *Acta Haematol.* 78 Suppl 1:75-6; U.S. Patent No. 4,690,915) or an alternate continuous infusion strategy can be employed.

5 These values provide general guidance of the range of transduced T cells to be utilized by the practitioner upon optimizing the method of the present invention for practice of the invention. The recitation herein of such ranges by no means precludes the use of a higher or lower amount of a component, as might be warranted in a particular
10 application. For example, the actual dose and schedule can vary depending on whether the compositions are administered in combination with other pharmaceutical compositions, or depending on interindividual differences in pharmacokinetics, drug disposition, and metabolism. One
15 skilled in the art readily can make any necessary adjustments in accordance with the exigencies of the particular situation.

In particular embodiments, the chimeric nucleic acid construct further contains a suicide gene such as thymidine
20 kinase (TK) of the HSV virus (herpesvirus) type I (Bonini, et al. (1997) *Science* 276:1719-1724), a Fas-based "artificial suicide gene" (Thomis, et al. (2001) *Blood* 97:1249-1257), or *E. coli* cytosine deaminase gene which are activated by gancyclovir, AP1903, or 5-fluorocytosine,
25 respectively. The suicide gene is advantageously included in the nucleic acid construct of the present invention to provide for the opportunity to ablate the transduced T cells in case of toxicity and to destroy the chimeric construct once a tumor has been reduced or eliminated. The
30 use of suicide genes for eliminating transformed or transduced cells is well-known in the art. For example, Bonini, et al. ((1997) *Science* 276:1719-1724) teach that donor lymphocytes transduced with the HSV-TK suicide gene

-29-

provide antitumor activity in patients for up to one year and elimination of the transduced cells is achieved using ganciclovir. Further, Gonzalez, et al. ((2004) *J. Gene Med.* 6:704-711) describe the targeting of neuroblastoma with cytotoxic T lymphocyte clones genetically modified to express a chimeric scFvFc:ζ immunoreceptor specific for an epitope on L1-CAM, wherein the construct further expresses the hygromycin thymidine kinase (HyTK) suicide gene to eliminate the transgenic clones.

10 It is contemplated that the suicide gene can be expressed from the same promoter as the chimeric receptor or from a different promoter. Generally, however, nucleic acid sequences encoding the suicide protein and chimeric receptor reside on the same construct or vector. Expression
15 of the suicide gene from the same promoter as the chimeric receptor can be accomplished using any well-known internal ribosome entry site (IRES). Suitable IRES sequences which can be used in the nucleic acid construct of the present invention include, but are not limited to, IRES from EMCV,
20 c-myc, FGF-2, poliovirus and HTLV-1. By way of illustration, a nucleic acid construct for expressing a chimeric receptor can have the following structure: promoter -> chimeric receptor -> IRES - > suicidal gene. Alternatively, the suicide gene can be expressed from a
25 different promoter than that of the chimeric receptor (e.g., promoter 1 -> chimeric receptor -> promoter 2 -> suicidal gene).

The following non-limiting examples are presented to better illustrate the invention.

-30-

Example 1: Mice and Cell Lines

C57BL/6 mice were purchased from the National Cancer Institute, and all animal work was conducted in accordance with standard guidelines.

5 Cell lines Bosc23, PT67, GP+E86, EG7 (H-2^b), and YAC-1 were obtained from the American Type Culture Collection (ATCC, Rockville, MD). RMA cells (H-2^b) originated from a Rauscher virus-induced C57BL/6 T-cell lymphoma (Ljunggren and Karre (1985) *J. Exp. Med.* 162:1745-1759). RMAS-S is a
10 sub-line of RMA which lacks MHC class-I surface expression (Kärre, et al. (1986) *Nature* 319:675-678). All packaging cells were grown in Dulbecco's modified Eagle medium (DMEM) with a high glucose concentration (4.5 gram/liter) supplemented with 10% heat-inactivated fetal bovine serum
15 (FBS; Hyclone, Logan, UT), 20 U/mL penicillin, 20 µg/mL streptomycin, 1 mM pyruvate, 10 mM HEPES, 0.1 mM non-essential amino acids, 50 µM 2-mercaptoethanol. RMA, EG7, RMA-S and YAC-1 cells were cultured in RPMI plus the same supplements described above.

20 **Example 2: Retroviral Vector Construction**

The full-length murine NKG2D cDNA was purchased from Open Biosystems (Huntsville, AL). Murine CD3ζ chain, Dap10 and Dap12 cDNAs were cloned by RT-PCR using RNAs from ConA- or IL-2 (1000 U/mL)-activated spleen cells as templates.
25 Mouse NKG2D ligands Rae-1β and H60 were cloned from YAC-1 cells by RT-PCR. All PCR reactions were performed using high-fidelity enzyme Pfu or PFUULTRA™ (STRATAGENE®, La Jolla, CA). The oligonucleotides employed in these PCR reactions are listed in Table 9.

-31-

TABLE 9

No.	Primer	Sequence	SEQ ID NO:
1	5' wtNKG2D	GCGAATTCGCCACCATGGCATTGATTCGTGATCGA	8
2	3' wtNKG2D	GGCGCTCGAGTTACACCGCCCTTTTCATGCAGAT	9
3	5' chNKG2D	GGCGAATTCGCATTGATTCGTGATCGAAAGTCT	10
4	5' wtDAP10	GCAAGTCGACGCCACCATGGACCCCCCAGGCTACC	11
5	3' wtDAP10	GGCGAATTCTCAGCCTCTGCCAGGCATGTTGAT	12
6	3' chDAP10	GGCAGAATTCGCCTCTGCCAGGCATGTTGATGTA	13
7	5' wtDAP12	GTTAGAATTCGCCACCATGGGGGCTCTGGAGCCCT	14
8	3' wtDAP12	GCAACTCGAGTCATCTGTAATATTGCCTCTGTG	15
9	5' ATG-CD3 ζ	GGCGTCGACACCATGAGAGCAAAATTCAGCAGGAG	16
10	3' ATG-CD3 ζ	GCTTGAATTCGCGAGGGGCCAGGGTCTGCATAT	17
11	5' CD3 ζ -TAA	GCAGAATTCAGAGCAAAATTCAGCAGGAGTGC	18
12	3' CD3 ζ -TAA	GCTTTCTCGAGTTAGCGAGGGGCCAGGGTCTGCAT	19
13	5' Rae-1	GCATGTCGACGCCACCATGGCCAAGGCAGCAGTGA	20
14	3' Rae-1	GCGGCTCGAGTCACATCGCAAATGCAAATGC	21
15	5' H60	GTTAGAATTCGCCACCATGGCAAAGGGAGCCACC	22
16	3' H60	GCGCTCGAGTCATTTTTTCTTCAGCATAACCAAG	23

Restriction sites inserted for cloning purposes are underlined.

5 Chimeric NKG2D was created by fusing the murine CD3 ζ chain cytoplasmic region coding sequence (CD3 ζ -CYP) to the full-length gene of murine NKG2D. Briefly, the *SalI*-*EcoRI* fragment of CD3 ζ -CYP (with the initiation codon ATG at the 5' end, primer numbers 9 and 10) and the *EcoRI*-*XhoI* fragment of NKG2D (without ATG, primer numbers 2 and 3) were ligated into the *SalI*/*XhoI*-digested pFB-neo retroviral vector (STRATAGENE®, La Jolla, CA). Similarly, chimeric Dap10 was generated by fusing the *SalI*-*EcoRI* fragment of full-length Dap10 (primer numbers 4 and 6) to the *EcoRI*-*XhoI* fragment of CD3 ζ -CYP (primer numbers 11 and 12). Wild-

-32-

type NKG2D (primer numbers 2 and 3), Dapl0 (primer numbers 4 and 5) and Dapl2 (primer numbers 7 and 8) fragments were inserted between the *EcoRI* and *XhoI* sites in pFB-neo. In some cases, a modified vector pFB-IRES-GFP was used to allow co-expression of green fluorescent protein (GFP) with genes of interest. pFB-IRES-GFP was constructed by replacing the 3.9 kb *AvrUScaI* fragment of pFB-neo with the 3.6kb *AvrII/ScaI* fragment of a plasmid GFP-RV(Ouyang, et al. (1998) *Immunity* 9:745-755). Rae-1 β (primer numbers 13 and 14) and H60 (primer numbers 15 and 16) cDNAs were cloned into pFB-neo. Constructs containing human NKD2D and human CD3 ζ or murine Fc were prepared in the same manner using the appropriate cDNAs as templates.

Example 3: Retrovirus Production and Transduction

Eighteen hours before transfection, Bosc23 cells were plated in 25 cm² flasks at a density of 4 x 10⁶ cells per flask in 6 mL of DMEM-10. Transfection of retroviral constructs into Bosc23 cells was performed using LIPOFECTAMINE™ 2000 (INVITROGEN™, Carlsbad, CA) according to the manufacturer's instruction. Viral supernatants were collected 48 and 72 hours post-transfection and filtered (0.45 μ m) before use. For generation of large scale, high-titer ecotropic vectors, the ecotropic viruses produced above were used to transduce the dualtropic packaging cell PT67 in the presence of polybrene (8 μ g/mL). After three rounds of transduction, PT67 cells were selected in G418 (1 mg/mL) for 7 days. Dualtropic vectors were then used to transduce ecotropic cell line GP+E86. Through this process, the virus titer from pooled GP+E86 cells generally was over 1x10⁶ CFU/mL. Concentration of retroviruses by polyethylene glycol (PEG) was performed according to standard methods (Zhang, et al. (2004) *Cancer Gene Ther.* 11:487-496; Zhang, et al. (2003) *J. Hemetother. Stem Cell Res.* 12:123-130).

-33-

Viral stocks with high titer ($1\sim 2 \times 10^7$ CFU/mL) were used for transduction of T cells. Primary T cells from spleens of C57BL/6 (B6) mice were infected 18-24 hours after concanavalin A (ConA, 1 μ g/mL) stimulation based on a well-established protocol (Sentman, et al. (1994) *J. Immunol.* 153:5482-5490). Two days after infection, transduced primary T cells ($0.5\sim 1 \times 10^6$ /mL) were selected in RPMI-10 media containing G418 (0.5 mg/mL) plus 25 U/mL rHuIL-2 for an additional 3 days. Viable cells were isolated using HISTOPAQUE®-1083 (Sigma, St. Louise, MO) and expanded for 2 days without G418 before functional analyses. NKG2D ligand-expressing RMA (RMA/Rae-1 β and RMA/H60) or RMA-S (RMA-S/Rae-1 β) cells were established by retroviral transduction with dualtropic vectors from PT67.

15 **Example 4: Cytokine Production by Gene-Modified T Cells**

Gene-modified primary T cells (10^5) were co-cultured with an equal number of RMA, RMA/Rae-1 β , RMA/H60 or YAC-1 cells in 96-well plates in complete media. After twenty-four hours, cell-free supernatants were collected. IFN- γ was assayed by ELISA using a DUOSET® ELISA kits (R&D, Minneapolis, MN). In some cases, T cells were cultured with equal numbers of irradiated (100 Gys) tumor cells for 3 days. Detection of other cytokines in culture was performed using a BIO-PLEX® kit (BIO-RAD®, Hercules, CA) based on the manufacturer's protocol.

25 **Example 5: Flow Cytometry**

For FACS analysis of NKG2D ligand expression, tumor cells were stained with mouse NKG2D-Ig fusion protein (R&D systems) according to manufacturer's instruction. Cell-surface phenotyping of transduced primary T cells was determined by direct staining with APC-anti-CD3 ϵ (clone 145-2C11; Pharmingen, San Diego, CA), PE-anti-NKG2D (clone 16-10A1; eBioscience, San Diego, CA) and FITC-anti-CD4

-34-

(Clone RM4-5; Caltag, Burlingame, CA) monoclonal antibodies. Cell fluorescence was monitored using a FACSCALIBER™ cytometer. Sorting of NKG2D ligand-expressing cells was performed on a FACSTARM™ cell sorter (Becton Dickinson, San Jose, CA).

Example 6: Cytotoxicity Assay

Three or four days after G418 selection (0.5 mg/mL), retroviral vector-transduced primary T cells were cultured in complete RPMI media containing 25 U/mL human IL-2 for an additional 2-3 days. Viable lymphocytes were recovered by centrifugation over HISTOPAQUE®-1083 (Sigma, St. Louis, MO) and used as effector cells. Lysis of target cells (RMA, RMA/Rae-1β, RMA/H60, EG7, RMA-S, RMA-S/Rae-1β, and YAC-1) was determined by a 4-hour ⁵¹Cr release assay (Sentman, et al. (1994) *supra*). To block NKG2D receptors, anti-NKG2D (clone: CX5, 20 μg/mL) was included in those assays. The percentage of specific lysis was calculated as follows:

$$\% \text{ Specific lysis} = \frac{(\text{Specific } ^{51}\text{Cr release} - \text{spontaneous } ^{51}\text{Cr release})}{(\text{Maximal } ^{51}\text{Cr release} - \text{spontaneous } ^{51}\text{Cr release})} \times 100.$$
Example 7: Treatment of Mice with Genetically Modified T Cells

For the determination of direct effects of chimeric NKG2D-bearing T cells (10⁶) on the growth of RMA or RMA/Rae-1β tumors, chimeric NKG2D- or vector-transduced T cells were mixed with tumor cells (10⁵) and then injected *s.c.* into the shaved right flank of recipient mice. Tumors were then measured using a caliper, and tumor areas were calculated. Animals were regarded as tumor-free when no tumor was found four weeks after inoculation. For the rechallenge experiments, mice were inoculated with 10⁴ RMA cells on the shaved left flank. In other experiments, transduced T cells were injected intravenously the day

-35-

before s.c. inoculation of tumor cells. Mice were monitored for tumor size every two days and were sacrificed when tumor burden became excessive.

Example 8: Statistical Analysis

- 5 Differences between groups were analyzed using the student's *t*-test. *p* values < 0.05 were considered significant.

-36-

What is claimed is:

1. A nucleic acid construct for expressing a chimeric receptor to reduce or eliminate a tumor comprising:

5 a first nucleic acid sequence encoding a promoter operably linked to



 a second nucleic acid sequence encoding a chimeric receptor comprising a C-type lectin-like natural killer cell receptor, or a protein associated therewith, fused to
10 an immune signaling receptor containing an immunoreceptor tyrosine-based activation motif of SEQ ID NO:1.

2. The nucleic acid construct of claim 1, wherein the construct is in a vector.

15

3. The nucleic acid construct of claim 1 further comprising a suicide gene.

4. A method for reducing or preventing tumors
20 comprising introducing a nucleic acid construct of claim 1 into an isolated T cell of a patient having or suspected of having a tumor and injecting the T cell into the patient so that the chimeric receptor is expressed on the surface of the T cell to activate anti-tumor immunity in the patient
25 thereby reducing or preventing the tumor.

**FIG. 1****FIG. 2**

SEQUENCE LISTING

<110> Trustees of Dartmouth College
Zhang, Tong
Sentman, Charles L.

<120> CHIMERIC NK RECEPTOR AND METHODS FOR TREATING CANCER

<130> DC0321WO

<150> US 60/612,836
<151> 2004-09-24

<150> US 60/681,782
<151> 2005-05-17

<160> 23

<170> PatentIn version 3.3

<210> 1
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Consensus immunoreceptor tyrosine-based activation motif

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> Xaa denotes Asp or Glu

<220>
<221> MISC_FEATURE
<222> (2)..(3)
<223> Xaa denotes any amino acid residue

<220>
<221> MISC_FEATURE
<222> (5)..(6)
<223> Xaa denotes any amino acid residue

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> Xaa denotes Ile or Leu

<220>
<221> MISC_FEATURE
<222> (8)..(12)
<223> Xaa denotes any amino acid residue

<220>
<221> MISC_FEATURE
<222> (13)..(13)
<223> Xaa denotes 1 to 3 amino acid residues, wherein the amino acid
can be any amino acid residue

- 2 -

<220>
 <221> MISC_FEATURE
 <222> (15)..(16)
 <223> Xaa denotes any amino acid residue

<220>
 <221> MISC_FEATURE
 <222> (17)..(17)
 <223> Xaa denotes Ile or Leu

<400> 1

Xaa	Xaa	Xaa	Tyr	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Tyr	Xaa	Xaa
1			5					10					15		

Xaa

<210> 2
 <211> 1575
 <212> DNA
 <213> Homo sapiens

<400> 2

tgaggacata	tctaaat	ctagttttat	agaaggcttt	tatccacaag	aatcaagatc	60
ttccctctct	gagcaggaat	cctttgtgca	ttgaagactt	tagattcctc	tctgcggtag	120
acgtgcactt	ataagtat	gatggggtgg	attcgtggtc	ggagggtctcg	acacagctgg	180
gagatgagtg	aatttcataa	ttataacttg	gatctgaaga	agagtgattt	ttcaacacga	240
tggcaaaagc	aaagatgtcc	agtagtcaaa	agcaaagtga	gagaaaatgc	atctccattt	300
tttttctgct	gcttcatcgc	tgtagccatg	ggaatccggt	tcattattat	ggtagcaata	360
tggagtgtctg	tattcctaaa	ctcattattc	aaccaagaag	ttcaaattcc	cttgaccgaa	420
agttactgtg	gcccattgtc	taaaaactgg	atatgttaca	aaaataactg	ctaccaattt	480
tttgatgaga	gtaaaaactg	gtatgagagc	caggcttctt	gtatgtctca	aaatgccagc	540
cttctgaaag	tatacagcaa	agaggaccag	gatttactta	aactgggtgaa	gtcatatcat	600
tggatgggac	tagtacacat	tccaacaaat	ggatcttggc	agtgggaaga	tggctccatt	660
ctctcacc	acctaactaac	aataattgaa	atgcagaagg	gagactgtgc	actctatgcc	720
tcgagcttta	aaggctatat	agaaaactgt	tcaactccaa	atacatacat	ctgcatgcaa	780
aggactgtgt	aaagatgatc	aaccatctca	ataaaagcca	ggaacagaga	agagattaca	840
ccagcggtaa	cactgccaac	cgagactaaa	ggaaacaaac	aaaaacagga	caaatgacc	900
aaagactgtc	agatttctta	gactccacag	gaccaaacca	tagaacaatt	tactgcaaaa	960
catgcatgat	tctccaagac	aaaagaagag	agatcctaaa	ggcaattcag	atatcccaa	1020

- 3 -

```

ggctgcctct cccaccacaa gccagagtg gatgggctgg gggaggggtg ctgttttaat 1080
ttctaaaggt aggaccaaca cccaggggat cagtgaagga agagaaggcc agcagatcag 1140
tgagagtgca accccaccct ccacaggaaa ttgcctcatg ggcagggcca cagcagagag 1200
acacagcatg ggcagtgcct tccctgcctg tgggggtcat gctgccactt ttaatgggtc 1260
ctccacccaa cggggtcagg gaggtggtgc tgccctagtg ggccatgatt atcttaaagg 1320
cattattctc cagccttaag atcttaggac gtttcctttg ctatgatttg tacttgcttg 1380
agtcccatga ctgtttctct tcctctcttt ctcccttttg gaatagtaat atccatccta 1440
tgtttgctcc actattgtat ttggaagca cataacttgt ttggtttcac aggttcacag 1500
ttaagaagga attttgcctc tgaataaata gaatcttgag tctcatgcaa aaaaaaaaaa 1560
aaaaaaaaaa aaaaaa 1575

```

```

<210> 3
<211> 6098
<212> DNA
<213> Homo sapiens

```

```

<400> 3
gcagttatca tagagcacag tccctcacat cacacagctg cagagatgag taaacaaaga 60
ggaaccttct cagaagtgag tctggcccag gacccaaagc ggcagcaaag gaaacctaaa 120
ggcaataaaaa gctccatttc aggaaccgaa caggaaatat tccaagtaga attaaatctt 180
caaaatcctt ccctgaatca tcaagggatt gataaaatat atgactgcca aggtaaaaca 240
ttaaatatat cttcaatatt attgttctag gatgtgcagt tgaatgcaga agggtgagga 300
aagattaggg aatattttgc acttgtgaga atcggagttc ataattggga tctaaaattc 360
taatatgaaa tcagaagact aattttattc gggcattggt caactgtaat ctgcggtcca 420
ctcatggaac attatattta ctgaaaatga aatggatat tctgagagaa agattactag 480
agtagatgta gatthagagg ccagagttta tcattatggt tccctgtgca tgtgggttct 540
ctagtatgta attctctagt atgtaatcct aatcaactct ctatctcccc tctctcagt 600
cctctatttc tctccctgca ggtttactgc cacctccaga gaagctcact gccgaggtcc 660
taggaatcat ttgcattgtc ctgatggcca ctgtgttaaa aacaatagtt cttattcctt 720
gtaagcatat tcttgaaaga ttagaaggga acgttttact ttaatgcttg gaagtgcctc 780
aaaatatttc atactgttga agaatagaac tcttatttta ctgtttcttt caaagatcta 840
ttacttcatt tatttttata gaaaaagtta attttattaa agattgtccc cattttaaat 900
aacacacaaa gtttcaaagt aagaaactaa actcattatg gtttatctaa atattacttt 960

```

- 4 -

ttataaaaaat cattttaatt tttctgttac agtcctggaa cagaacaatt cttcccaaaa	1020
tacaagaacc cagaaaagta cttttttatt ttcaaagttc tgatattagt acaatttgga	1080
acaaaaagta atatggttat tctgaatfff tcacaacata aataacaaaa tcattgtaga	1140
gaacatgtgt ttatftttttg tgtgtaatct atatatatgt atatacatat acacacaaaag	1200
atattttctg atttcataat tcaaaggcat gctatagaag aaaagtatft agaaaaacaa	1260
attaattftft gaaagtggft acatcaaata ctacaagaga tgggtgaagt tgtgctaaag	1320
tctfttaaaaa tgtftattft aaaggtctat tactfttatat attftftatag aaaaagttaa	1380
ftfttattaaa gattctcccc attfttaata acacacaaaag ftftcaaagta agaaactaaa	1440
ctcgttatgg ftcatctaga tatcagftft tataaaaaatc attfttaatt ftctattaca	1500
gtcctggagc agaacaattc ftccccgaat acaagaacgc agaaaggtag attftftatft	1560
tcaatgtftc gatattagta caattftatat ftftgtgtctg ftfttaaggca tgtaaaagaa	1620
tagtggcatt ftftgcagaaa ataagccata aattcagcca taaatatttg taaagaaaaga	1680
ttatgaggca gcattftcct ftctccagt agtagaaaata ctactftaaa atcattctac	1740
cctctftctc ccaatttaaca gaggtftcct actgctgtga gatgatacca aataaataat	1800
fttactattc taaaaaagca gttgtgtatc agcgatgttc aacacatgtg tagagtgtat	1860
ftftgtftgt tcatttgctt tatatgggaa cacaattagg gaggagaggc taacccttgt	1920
ctgtgcatgt gtgtatgact gactcagfta ftaaaaatat acattftata gcctgtaagg	1980
atgcgtaaat atgttaagca catatatgtt tatactgttg aaatatgtga actaattftc	2040
attftftaaaa attcatattg gtctaaatag taattcatat cftftattagc acgtcattgt	2100
ggccattgtc ctgaggagt gattacatat tccaacagft gttattacat tggtaaggaa	2160
agaagaactt gggaagagag ftftgctggc tgtacttcga agaactccag tctgctftct	2220
atagataatg aagaagaaat ggtaagatgt aaatgtftca aacattfttat gaaaagcttc	2280
cttcagtga taatacatft gtagaaaaca tccatatgtg tgtacatata ftftatctcat	2340
atattftcaa gtgtatgtaa tattcaattg attgactfta taatgtftft aaagttatat	2400
actgctaattg tacattftatt ftcagftftft gftftftcaag gaaaaccatg cftctataag	2460
tgctftgaat ccacaataaa ftftgctatc taattftatc gggcatgata tcactgtgtc	2520
atgcagattg atcacaaaag gaatgaatgc atgtgatata agtcagatca tgaaataaaaa	2580
gftftccagct ctagcagftc caccctgtg tatgccctca tcactftatc tgactcctct	2640
ccaaaacgca gtcttgactt ttaatattat aaataatgat tgctgtftct tgaattftatt	2700
tatataaagg gaatcaaaca gtgtgaattt catgtcftft tcaatcctat ctgatatttg	2760

- 5 -

tgcaattcct ccatattatt gcagttatca gtagtatgtt actgttcact gctgtactat	2820
gtacaaagaa cagtaagaat ccattgagtc cttgtctctg gatggggaag tgggtctcat	2880
gccctcaggg acaaagagga ccctaggtgg tttagcgtgc actgttagtc atgggggtccc	2940
tttgctgata ctctcatcc acagccatcc tgggtgtctt tggatatgaga aggaagcact	3000
ttctctagct ccatattggt agcaggtctc ctggtagatc atccttgcca gtggcaccag	3060
ccttgccctgg tattgtggag gggactctcc ttcgataccc tctctctatt gccaggttgg	3120
gtgtagggaa acagcaggcc taggtcacct tcttctgtcg tgtggaggac ttaacatgct	3180
cacttggaaca cttggttgat ccctgatgct aggggtcccag acaatttcat ctttctcttt	3240
ccacctttca gagttctcca ttgcttttgt ctttcattaa tcccagagtt tatagttggt	3300
tttagtaggg agtagcagag agagacgagt ctacaccacc tggccaggac ccctgttatt	3360
ccgcaaaaac cgaatcggat aaaaattgag ggcttatcta gttaaagaat ggtgtggtac	3420
ccagaaaacc caatctgtag cttccatgtc atctatttct gaatgacaac ccctcaattc	3480
ccttctaaat ctccaactct gagaaatata gcacaaaaat agattgattt agtcacagta	3540
tctggagaaa tgaatgcaca gtatcaggaa acttattaaa acccttctctg tgttttattct	3600
gttaattgga gtaactatta cattgcaaga attaaaatgt ctttattaac atgagaataa	3660
gaatgaaagt actaagtata aacgttgaag agttcattta aataaaaaat tcaaacattt	3720
atgaaagttt ttggcactgc aaatagtggg ttccaacttt aatatattgt ttttgtaatg	3780
ttttcataat tattatttaa gtgaaaatta tttcttttct tttagaaatt tctggccagc	3840
attttacctt cctcatggat tgggtgtgtt cgtaacagca gtcacatcc atgggtgaca	3900
ataaatggtt tggctttcaa acataagtaa gttcttttgt atggcgctat ataaaaaata	3960
tatataaagg ataaattcag aagaataata tgaataaatt tatgtggaat cattgacatg	4020
aagaaagatg tggaaagtta gtgaaatgtt gatataaata ttttacaata gaccatagta	4080
gtccatatat ttcaaccgct cattggtctg ctagtaacct tcttggttat cagatggacc	4140
aggggtgtcc catctttggc ttctgtgggc caggttagaa gacgaatagt cttggcccac	4200
acatagaata cactaacact aacgatagct gacgagctaa aaaaaaaaaa aaatcacaga	4260
atgttttaag aaagtttacg tatttgtgtt gggccgcatt caaagctgtc ctgggtcacg	4320
tgcgggccat gggcagcgag ttggacaacc tcgagctgga ctatcaggga actgcagtgc	4380
ttgtttttat taaaaagcca cgcttacttt ttactttaag aatatcctca aagcacaata	4440
atagtgtctg ttggcatattg ctataatttt ttattacta gttattgttg tcaatctctt	4500
attgtgccta atttataaat taaactttat cacagttatg aatgtgtaga gaaaacataa	4560

- 6 -

tctctctata ggttctgcac tatctgccat ttcaggcatc cactggggtc ttgaaacata 4620
 tccctcgtgg atgaagaggg actactctgt tgagtgttca gaataatgac tcttactaat 4680
 attatgaaaa atttaattac ccttttccat gaaattcttt tcttacagta catggaaaat 4740
 gctttcgtct catgaatcat ttgcttaaaa tgtaacagaa tatggatttt tctccattac 4800
 aggataaaaag actcagataa tgctgaactt aactgtgcag tgctacaagt aaatcgactt 4860
 aaatcagccc agtgtggatc ttcaatgata tatcattgta agcataagct ttagaagtaa 4920
 agcatttgcg tttacagtgc atcagataca ttttatattt cttaaaatag aaatattatg 4980
 attgcataaa tctgaaaatg aattatgtta tttgctctaa tacaaaaatt ctaaataat 5040
 tattgaaata ggatgcacac aattactaaa gtacagacat cctagcattt gtgtcgggct 5100
 cattttgctc aacatgggat ttgtggtttt cagcctttct aaaagttgca tgttatgtga 5160
 gtcagcttat aggaagtacc aagaacagtc aaacccatgg agacagaaag tagaatagt 5220
 gttgccaatg tctcagggag gttgaaatag gagatgacca ctaattgata gaacgtttct 5280
 ttgtgtcgtg atgaaaactt tctaaatttc agtaatgggtg atgggtgtga ctttgcaat 5340
 atactaaaca tcattgattt ttaatcattt taagtgcag aaatgtatgc tttgtacatg 5400
 acacttcaat aaagctatcc agaaaaaaaa aagcctctga tgggattgtt tatgactgca 5460
 tttatctcta aagtaatttt aaagattagc ttctttataa tattgacttt tctaatacgt 5520
 ataaagtgtt tccttcaatg tactgtgtta tctttaattt ctctctcttg tattttgtat 5580
 tttgggggat tgaagtcata cagaaatgta ggtattttac atttatgctt ttgtaaattg 5640
 catcctgatt ctaaaattcc ctttagtaat tttgttgtt ataaatagaa atacaactga 5700
 tgtctgcatt ttgattttat atctacttat tccactgatt ttatatattt aaatctatta 5760
 tgtcaactat tgatttattt ctgggtgttc tatataacga gcaattttat ctgcaaatga 5820
 tcacactttt atttttttta atccatgtgc tataacttag ttttattttc atttattttc 5880
 actggctaag gttttatacc catagttgaa tagaaggcac aatcaaagtt ctttgtggat 5940
 catatgcac attttctggt tttggcaaaa aatacttcaa catgttatac atatttataa 6000
 agcttgggtg tttttgcac ctatctttct catatcgaag cagttttata atcctatttt 6060
 ctaatagatt ttatcaattg taacaatttt tattaatt 6098

<210> 4

<211> 3000

<212> DNA

<213> Homo sapiens

<400> 4

aattcccagc cctggagctg gcattccagt gggaggccac tctcagtttc acttgggtgac 60

- 7 -

ctttcacagc actgaccatg ttggccctat ttctcccctg cttgcttgct tttctatttt	120
attttattat tacattttta ttgttagaga gaggggtctca ttctgtcgcc caggctggag	180
tgcagtggca aagtgggtgag atctcggctc actgcaacct ccacttgccct cagtctccca	240
agtagctggg attacaggag cctgacacca tgcccgggta atttttgtat tttttagagag	300
acgggggttc accatattgg cttgaactcc tgatctcagg tgatcccccc accttggcct	360
cccaaagtgc tgggattaca ggcatgagcc actgcggtgg cctctcccct gctttcaaga	420
tgccatgctc tcaggggtcc cctccctctt tctccatttc cctggcaaag ttctctctct	480
tcccccatte agtgtgtgtt gtgatagggg cagaatcctg tctgactca ctctcttggg	540
gatctcacc agtcttgtgg ctttaagtac catccataag ccatcaacc ccaaatttac	600
atctccagac cagccttata ccctgaactc ctaaattgag tgaggttatt cagcatctcc	660
acaggggagat tgtcaggcat ttccaacct gtatgccc aaacctcgtcac tttccccgca	720
aaccacttc cctaccttcc atctctgcca gcagacactc ccatcttctc agcgtttcat	780
gccagaaggc ttggctgtct aggatccctc tcaaacacac ccacattcat ttaatcagca	840
aattttcttg gccctacctc caaaatattt ccagatctcc ctagcctgca cacccttgcc	900
acctgtcatt ccacttgga ccaggccagc agcctccctg gtctctctga ccctcccct	960
gagttcggtc accaaaggca gtaacggaga caccctctca acacacacag gaagcagatg	1020
gccttgacac cagcagggtg acatccgcta ttgctacttc tctgtctccc cacagttcct	1080
ctggacttct ctggaccaca gtctctgccc agaccctgc cagaccccag tccaccatga	1140
tccatctggg tcacatcctc ttctgtctt ttgtcccagg tgaagccagt ggttacaggg	1200
gatggtaggc agagcgtttg tgagatgggt gcttgggtga cgtctgcagg gacgggtgat	1260
gaaagtgggg ttcttctccc tgcacccctt cccttctggg agatccattc tgcttcaggg	1320
cctgggtcct tgggggcgga aggggtgag acaggagtt ctggaggggc tgcctgtag	1380
cgtccccctc tcatggctgg gtctctgctg ccacttccaa tttcttgtca ctctccatgt	1440
ctctgggagt ccccttccca tgtggctctg ttccatctct ccagcctgga gattacttct	1500
caggacacta cctttccttc tctacacct attttttggg ttgtttattt tgagatgggg	1560
tcttgtctg ttgtccaggc tggagtgcag tggcacaatc acggctcacg gcagccttga	1620
cttctgggc tcaggatgat ctcccagctc agcctcccga gtaactggga ttacagggtg	1680
gaaccaacac ttccagctaa tttttgtatt tctttagag acgaggctc actatgttgc	1740
ccaggctggt ctggaactcc tgggtcaag cgatcttctt gcctcggcct cccaaagtgc	1800
tgggatgaca ggcgtgagcc acggtgccag gctgagcatt ctgttttgtg gaccttctct	1860

- 8 -

```

ccacctcat ccaccttctt tctctttcca cagtggctgc agctcagacg actccaggag 1920
agagatcatc actccctgcc ttttaccctg gcacttcagg tatcacttcc accccagaag 1980
cttggccaga ggctcccaga acaccccagt ggttctccag gtcaccatcc cacctcccgt 2040
ccccaaatca gaggatccgt gtcctttctc gagtcccaga atcagcgcacc cccagcctgt 2100
gttcaggagc acccctgtgt ccgcgcgcac agccccgagg gtccctgggac accccagcct 2160
ctctgcatct gtctcccgtt tcattcccca agcgcaactc caaggaacct gggaccgcgc 2220
ccctcgcagg ggacttcctc tctgcctgtg gccaaagcac agccccagga cgcagagctt 2280
gagttgtctc cctgttccgg ccccaactct ccaggctctt gttccggatg tgggtccctc 2340
tctctgccgc tcctggcagg cctcgtggct gctgatgcgg tggcatcgct gctcatcggt 2400
ggggcggtgt tcctgtgcgc acgcccacgc cgcagccccg cccaagggtga gggcgagat 2460
ggggcgggcc tggaagggtg atagtgtccc tagggagggg gtcccaggga gggggccctt 2520
ggggaagccc tggaggaggt gctggggaaa ccctggggga ggtgcctggg ggaaccctg 2580
aggaaacccc tgaagcaggg ggtccccagg gaagtggaga tatgggtggt caagcttcat 2640
gctttctctc ccctatcccc agaagatggc aaagtctaca tcaacatgcc aggcaggggc 2700
tgaccctcct gcagcttgga cctttgactt ctgaccctct catcctggat ggtgtgtggt 2760
ggcacaggaa cccccgcccc aacttttgga ttgtaataaa acaattgaaa cacctgtagt 2820
cgtattcttt ctcaaagaac ccagagttc ccaaagcctc cctcccatga actgtttctg 2880
gatccaaggc ccctcagaa cccccacatg tcccatccc atcagcccaa ggatctggca 2940
taatgttttt gtgcttcatg tttattttag gagagtattg gggagcggtc tggctctca 3000

```

<210> 5
<211> 604
<212> DNA
<213> Homo sapiens

```

<400> 5
ccacgcgtcc gcgctgcgcc acatcccacc ggcccttaca ctgtggtgtc cagcagcatc 60
cggcttcatg gggggacttg aacctgcag caggctcctg ctctgcctc tcctgctggc 120
tgtaagtggc ctccgtcctg tccaggccca ggcccagagc gattgcagtt gctctacggt 180
gagccccggc gtgctggcag ggatcgtgat gggagacctg gtgctgacag tgctcattgc 240
cctggccgtg tacttcctgg gccggctggt ccctcggggg cgaggggctg cggaggcagc 300
gacccggaaa cagcgtatca ctgagaccga gtcgccttat caggagctcc agggtcagag 360
gtcggatgtc tacagcgacc tcaacacaca gaggccgtat tacaaatgag cccgaatcat 420

```

- 9 -

gacagtcagc aacatgatac ctggatccag ccattcctga agcccaccct gcacctcatt	480
ccaactccta ccgcgataca gaccacaga gtgccatccc tgagagacca gaccgctccc	540
caatactctc ctaaaataaa catgaagcac aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	600
aaaa	604

<210> 6
 <211> 1677
 <212> DNA
 <213> Homo sapiens

<400> 6	
gtcctccact tcctggggag gtagctgcag aataaaacca gcagagactc cttttctcct	60
aaccgtcccg gccaccgctg cctcagcctc tgcctccag cctctttctg agggaaagga	120
caagatgaag tggaaggcgc ttttcaccgc ggccatcctg caggcacagt tgccgattac	180
agaggcacag agctttggcc tgctggatcc caaactctgc tacctgctgg atggaatcct	240
cttcatctat ggtgtcattc tcaactgcctt gttcctgaga gtgaagttca gcaggagcgc	300
agacgcccc gcgtaccagc agggccagaa ccagctctat aacgagctca atctaggacg	360
aagagaggag tacgatgttt tggacaagag acgtggccgg gaccctgaga tggggggaaa	420
gccgcagaga aggaagaacc ctcaggaagg cctgtacaat gaactgcaga aagataagat	480
ggcggaggcc tacagtgaga ttgggatgaa aggcgagcgc cggaggggca aggggcacga	540
tggcctttac cagggtctca gtacagccac caaggacacc tacgacgcc ttcatatgca	600
ggccctgccc cctcgctaac agccagggga tttcaccact caaaggccag acctgcagac	660
gccagatta tgagacacag gatgaagcat ttacaaccgg gttcactctt ctcagccact	720
gaagtattcc cctttatgta caggatgctt tgggttatatt tagctccaaa cttcacaca	780
cagactgttg tccctgcact ctttaaggga gtgtactccc agggcttacg gccctggcct	840
tgggccctct gggttgccgg tgggtgcagg agacctgtct cctggcggtt cctcgttctc	900
cctgggaggg gggcgcaact cctctcacag ctgagttggt gagtctgttt tgtaaagtcc	960
ccagagaaa cgcagatgct agcacatgcc ctaatgtctg tatcactctg tgtctgagt	1020
gcttactcc tgctgtaaat ttggcttctg ttgtcacctt cacctccttt caaggtaact	1080
gtactgggcc atgttgtgcc tccctggtga gagggccggg cagaggggca gatggaaagg	1140
agcctaggcc aggtgcaacc agggagctgc aggggcatgg gaagggtggc gggcagggga	1200
gggtcagcca gggcctgcga gggcagcggg agcctccctg cctcaggcct ctgtgccgca	1260
ccattgaact gtaccatgtg ctacaggggc cagaagatga acagactgac cttgatgagc	1320
tgtgcacaaa gtggcataaa aaacatgtgg ttacacagtg tgaataaagt gctgcggagc	1380

- 10 -

```

aagaggaggc cggttgattca cttcacgctt tcagcgaatg acaaaatcat ctttgtgaag 1440
gcctcgcagg aagacccaac acatgggacc tataactgcc cagcggacag tggcaggaca 1500
ggaaaaaccc gtcaatgtac taggatactg ctgcgtcatt acagggcaca ggccatggat 1560
ggaaaacgct ctctgctctg ctttttttct actgttttaa tttatactgg catgctaaag 1620
ccttcctatt ttgcataata aatgcttcag tgaaaaaaaa aaaaaaaaaa aaaaaaa 1677

```

```

<210> 7
<211> 591
<212> DNA
<213> Homo sapiens

```

```

<400> 7
cagaacggcc gatctccagc ccaagatgat tccagcagtg gtcttgctct tactcctttt 60
ggttgaacaa gcagcggccc tgggagagcc tcagctctgc tatatcctgg atgccatcct 120
gtttctgtat ggaattgtcc tcaccctcct ctactgtcga ctgaagatcc aagtgcgaaa 180
ggcagctata accagctatg agaaatcaga tgggtgtttac acgggcctga gcaccaggaa 240
ccaggagact tacgagactc tgaagcatga gaaaccacca cagtagcttt agaatagatg 300
cggtcataatt cttcttttggc ttctggttct tccagccctc atggttggca tcacatatgc 360
ctgcatgcca ttaacaccag ctggccctac ccctataatg atcctgtgtc ctaaattaat 420
atacaccagt ggttcctcct cctgtttaa gactaatgct cagatgctgt ttacggatat 480
ttatatctta gtctcactct cttgtccac cttcttctc ttccccattc ccaactccag 540
ctaaaatatg ggaagggaga accccaata aaactgccat ggactggact c 591

```

```

<210> 8
<211> 35
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Synthetic oligonucleotide

```

```

<400> 8
gcgaattcgc caccatggca ttgattcgtg atcga 35

```

```

<210> 9
<211> 34
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Synthetic oligonucleotide

```

```

<400> 9

```

- 11 -

ggcgctcgag ttacaccgcc cttttcatgc agat 34

<210> 10
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 10
ggcgaattcg cattgattcg tgatcgaaag tct 33

<210> 11
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 11
gcaagtcgac gccaccatgg accccccagg ctacc 35

<210> 12
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 12
ggcgaattct cagcctctgc caggcatggt gat 33

<210> 13
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 13
ggcagaattc gcctctgcc ggcattgtga tgta 34

<210> 14
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 14

- 12 -

gttagaattc gccaccatgg gggctctgga gccct 35

<210> 15
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 15
gcaactcgag tcatctgtaa tattgcctct gtg 33

<210> 16
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 16
ggcgtcgaca ccatgagagc aaaattcagc aggag 35

<210> 17
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 17
gcttgaattc gcgagggggcc agggctctgca tat 33

<210> 18
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 18
gcagaattca gagcaaaatt cagcaggagt gc 32

<210> 19
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 19

- 13 -

gctttctcga gtttagcgagg ggccagggtc tgcac 35

<210> 20

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 20

gcatgtcgcac gccaccatgg ccaaggcagc agtga 35

<210> 21

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 21

gcggctcgag tcacatcgca aatgcaaag c 31

<210> 22

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 22

gttagaattc gccaccatgg caaagggagc cacc 34

<210> 23

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 23

gcgctcgagt cattttttct tcagcatata ccaag 35