




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 233/70, A61K 31/415 C07D 403/06, C07F 9/6506	A1	(11) International Publication Number: WO 92/04331 (43) International Publication Date: 19 March 1992 (19.03.92)
(21) International Application Number: PCT/GB91/01543 (22) International Filing Date: 10 September 1991 (10.09.91) (30) Priority data: 9019839.1 11 September 1990 (11.09.90) GB (71) Applicant (for all designated States except US): SMITH KLINE & FRENCH LABORATORIES LIMITED [GB/GB]; Mundells, Welwyn Garden City, Hertfordshire AL7 1EY (GB). (72) Inventors; and (75) Inventors/Applicants (for US only) : HICKEY, Deirdre, Mary, Bernadette [IE/GB]; SmithKline Beecham Pharmaceuticals, The Frythe, Welwyn, Hertfordshire AL6 9AR (GB). COOPER, David, Gwynn [GB/GB]; SmithKline Beecham Pharmaceuticals, Coldharbour Road, The Pinnacles, Harlow, Essex CM19 5AD (GB). JAXA-CHAMIEC, Albert, Andrzej [GB/GB]; 22 Beacon Way, Rickmansworth, Hertfordshire WD3 2PE (GB).		(74) Agent: GIDDINGS, Peter, J.; Corporate Patents, Smith-Kline Beecham, Mundells, Welwyn Garden City, Hertfordshire AL7 1EY (GB). (81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, KR, LU (European patent), NL (European patent), SE (European patent), US. Published <i>With international search report.</i>
(54) Title: SUBSTITUTED 2 IMIDAZOLONE DERIVATIVES, THEIR PREPARATION AND PHARMACEUTICAL COMPOSITIONS  (57) Abstract Substituted 4,5-diaryl-2-imidazolones of structure (I), in which each group Ar is the same or different and is optionally substituted phenyl or optionally substituted heteroaryl; R¹ is hydrogen, alkyl, optionally substituted phenyl or optionally substituted heteroaryl; n is 4 to 12; and X is 5-tetrazolyl, SO₃H, P(O)(OR)₂, P(O)(H)₂ or P(O)(R)(OR) in which R is hydrogen acceptable salt thereof, processes for preparing them, pharmaceutical compositions containing them and their use in therapy, inter alia, in the treatment of cardiovascular disorders.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU ⁺	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE*	Germany	MC	Monaco	US	United States of America
DK	Denmark				


⁺ Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

-1-

SUBSTITUTED 2-IMIDAZOLONE DERIVATIVES, THEIR PREPARATION AND
PHARMACEUTICAL COMPOSITIONS

The present invention relates to novel substituted
2-imidazolone derivatives, processes for their
5 preparation, pharmaceutical compositions containing them
and their use in therapy.

The present invention therefore provides in a first
aspect compounds of structure (I):



in which,

each group Ar is the same or different and is optionally
substituted phenyl or optionally substituted
20 heteroaryl;

R¹ is hydrogen, C₁₋₄alkyl, optionally substituted
phenyl or optionally substituted heteroaryl;

n is 4 to 12; and

X is 5-tetrazolyl, SO₃H, P(O)(OR)₂, P(O)(OH)₂ or

25 P(O)(R)(OR) in which R is hydrogen or C₁₋₄alkyl,
or a pharmaceutically acceptable salt thereof.

Suitably, each group Ar is the same and is
optionally substituted phenyl or optionally substituted
30 heteroaryl. More suitably, each group Ar is the same
and is optionally substituted phenyl. Preferably each
group Ar is the same and is unsubstituted phenyl.

Suitably, R¹ is hydrogen, C₁₋₄alkyl, optionally
35 substituted phenyl or optionally substituted heteroaryl.

-2-

Preferably R^1 is optionally substituted phenyl; most preferably unsubstituted phenyl.

Suitably, n is 4 to 12; preferably n is 4 to 8,
5 most preferably n is 6 or 7.

Suitably, X is 5-tetrazolyl, SO_3H , $P(O)(OR)_2$,
 $P(O)(OH)_2$ or $P(O)(R)(OR)$ in which R is hydrogen or
 C_{1-4} alkyl. Preferably X is SO_3H .

10

Suitable substituents for phenyl groups Ar and R^1
include, for example, 1-3 groups which may be the same
or different and are selected from C_{1-4} alkyl,
halo C_{1-4} alkyl such as CF_3 , halogen, hydroxy and
15 C_{1-4} alkoxy.

Suitable heteroaryl groups include, for example,
saturated or unsaturated 5- or 6-membered rings
comprising 1 to 3 heteroatoms selected from nitrogen,
20 oxygen and sulphur.


Preferred such rings include, for example, thienyl
and furyl rings.

25 Particularly preferred compounds of structure (I)
include:

sodium 7-(3,4,5-triphenyl-2-oxo-2,3-dihydroimidazol-1-yl)-
heptanesulphonate;
30 diethyl 7-(3,4,5-triphenyl-2-oxo-2,3-dihydroimidazol-1-
yl)-heptanephosphonate;
7-(3,4,5-triphenyl-2-oxo-2,3-dihydroimidazol-1-yl)-
heptanephosphonic acid; and
ethyl 7-(3,4,5-triphenyl-2-oxo-2,3-dihydroimidazol-1-yl)-
35 methylphosphinate.

-3-

The compounds of structure (I) can be prepared using procedures analogous to those known in the art. The present invention therefore provides in a further aspect a process for the preparation of compounds of structure (I) in which X is other than 5-tetrazolyl which comprises reaction of a compound of structure (II):



in which

15 Ar, R¹ and n are as described for structure (I) and L is a leaving group, with a suitable source of the group X; and optionally thereafter forming a pharmaceutically acceptable salt thereof. Compounds of structure (I) in which X is 5-tetrazolyl, can be prepared from compounds of structure (II) by standard techniques, for example,

20 when L is bromine, by reaction with sodium cyanide in a suitable solvent such as dimethylsulphoxide, to form the intermediate compound in which L is cyano; followed by reaction with tri-n-butyl tin azide in, for example,

25 tetrahydrofuran to form the desired compound of structure (I).

Suitable leaving groups L will be apparent to those skilled in the art and include, for example, halogen,

30 such as bromine.


Suitable sources of the group X will again be apparent to those skilled in the art and include, for example, where X is SO₃Na, sodium sulphite.

35

-4-

The reaction between the compounds of structure (II) and the source of X is carried out in a solvent at elevated temperature. Preferably, for example where X is SO_3Na the reaction is carried out in aqueous ethanol at reflux temperature for a suitable period to allow the reaction to go to completion; and where X is a phosphorus containing group the reaction is carried out in an organic solvent such as toluene or xylene.

The compounds of structure (II) can be prepared from compounds of structure (III):



(III)

in which

Ar and R^1 are as described for structure (I) by reaction with, for example, a compound of formula $\text{L}^1(\text{CH}_2)_n\text{L}$, in which L and L^1 are suitable leaving groups, in the presence of a base such as potassium carbonate and a suitable solvent such as butanone. Suitable groups L are as described for structure (II). Suitable groups L^1 will be apparent to those skilled in the art, and include halogen, in particular bromine.

Compounds of structure (III) are known or can be prepared by standard techniques.

The compounds of structure (I) and their pharmaceutically acceptable salts have been found to be PGI_2 agonists and as such are useful in therapy for the treatment of disease conditions in which such an effect is beneficial.

-5-

More specifically, the compounds are expected to have utility as antithrombotic, vasodilatory, anti-atherosclerotic, antiinflammatory and cytoprotective agents. In particular, as antithrombotic and
5 vasodilatory agents, the compounds are expected to be useful in the treatment of cardiovascular occlusive disorders including spasmodic and thrombotic disorders; coronary heart disease (primary and secondary prevention); stroke; post-operative thrombosis utilisation including
10 post-angioplasty; deep vein thrombosis; peripheral vascular disease and Reynaud's disease. As antiatherosclerotic agents the compounds would be expected to reduce atherosclerotic plaque formation; and as cytoprotective agents the compounds would be expected to
15 protect liver and gastric mucosa, protect against mucosal and ulcerative damage and reduce infarct size in myocardial infarct.

In addition to the foregoing utilities the compounds
20 have antihyperlipidaemic properties and as such are expected to be of use as lipid lowering agents, and in the treatment of atherosclerosis and its sequelae.

In therapeutic use, the compounds of the present
25 invention are usually administered in a standard pharmaceutical composition. The present invention therefore provides in a further aspect pharmaceutical compositions comprising a compound of structure (I) or a pharmaceutically acceptable salt thereof and a
30 pharmaceutically acceptable carrier.

The compounds of structure (I) and their pharmaceutically acceptable salts which are active when given orally can be formulated as liquids, for example
35

-6-

syrups, suspensions or emulsions, tablets, capsules and lozenges.

5 A liquid formulation will generally consist of a suspension or solution of the compound or pharmaceutically acceptable salt in a suitable liquid carrier(s) for example, ethanol, glycerine, non-aqueous solvent, for example polyethylene glycol, oils, or water with a suspending agent, preservative, flavouring or colouring agent.

A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations.

15 Examples of such carriers include magnesium stearate, starch, lactose, sucrose and cellulose.

A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloscs, silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule.

Typical parenteral compositions consist of a solution or suspension of the compound or pharmaceutically acceptable salt in a sterile aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration.

-7-

A typical suppository formulation comprises a compound of structure (I) or a pharmaceutically acceptable salt thereof which is active when administered in this way, with a binding and/or lubricating agent such as polymeric glycols, gelatins or cocoa butter or other low melting vegetable or synthetic waxes or fats.

Preferably the composition is in unit dose form such as a tablet or capsule.

10

Each dosage unit for oral administration contains preferably from 1 to 250 mg (and for parenteral administration contains preferably from 0.1 to 25 mg) of a compound of the structure (I) or a pharmaceutically acceptable salt thereof calculated as the free base.

15

The present invention also provides a method of mimicking the effects of PGI₂ which comprises administering to a mammal in need thereof an effective amount of a compound of the structure (I) or a pharmaceutically acceptable salt thereof; and a method of treatment of cardiovascular disorders which comprises administering to a mammal in need thereof an effective amount of a compound of the structure (I) or a pharmaceutically acceptable salt thereof.

20

25

The pharmaceutically acceptable compounds of the invention will normally be administered to a subject in a daily dosage regimen. For an adult patient this may be, for example, an oral dose of between 1 mg and 500 mg, preferably between 1 mg and 250 mg, or an intravenous, subcutaneous, or intramuscular dose of between 0.1 mg and 100 mg, preferably between 0.1 mg and 25 mg, of the compound of the structure (I) or a pharmaceutically

35

-8-

acceptable salt thereof calculated as the free base, the compound being administered 1 to 4 times per day. Suitably the compounds will be administered for a period of continuous therapy, for example for a week or more.

5

The following Examples serve to illustrate the invention. Temperatures are recorded in degrees centigrade.

10

EXAMPLE 1Sodium 7-(3,4,5-triphenyl-2-oxo-2,3-dihydroimidazol-1-yl)-heptanesulphonate

5

A mixture of 1,4,5-triphenylimidazol-2-one (15.3g), dibromoheptane (50.6g) and potassium carbonate (13.8g) was heated at reflux temperature in dry butanone (750ml) for 20 hours. The mixture was cooled, filtered and the filtrate evaporated to an oil which was chromatographed on silica gel (hexane/ethyl acetate) to give 1,4,5-triphenyl-3-(7-bromoheptyl)imidazole-2-one (11.1g, 46%) as an oil.

10
15 NMR δ (CDCl₃) 1.2-1.9 (10H, m, 5 x CH₂), 3.4 (2H, t, -CH₂Br), 3.7 (2H, t, -CH₂N), 6.8-7.4 (15H, m, 3 x Ph) ppm.

A solution of 1,4,5-triphenyl-3-(7-bromoheptyl)-imidazol-2-one (2.0g) in ethanol (10ml) was refluxed with a solution of sodium sulphite (0.55g) in water (5ml) for 20 hours. More sodium sulphite (0.2g) was added and refluxing continued for a further 20 hours. The mixture was evaporated to dryness, boiled in ethanol, filtered hot and evaporated to an oil. This was taken up in a small volume of ethanol, excess diethyl ether added and the precipitated solid filtered off and chromatographed on silica gel (dichloromethane/methanol 5:1). The resulting oil in methanol/water 1:1 was passed down an Amberlyst 15 ion exchange resin (Na form) and evaporated to a solid. This was taken up in ethanol and precipitated with diethyl ether giving sodium 7-(3,4,5-triphenyl-2-oxo-2,3-dihydroimidazol-1-yl)heptanesulphonate (0.49g, 23%) as a white solid, m.p. 160°C.

30 Found: C, 63.47; H, 5.69; N, 5.04; S, 5.63%

C₂₈H₂₉N₂NaO₄S + 3.5% water

35 Requires: C, 63.31; H, 5.89; N, 5.28; S, 6.04%

-10-

EXAMPLE 2Diethyl 7-(3,4,5-triphenyl-2-oxo-2,3-dihydro-imidazol-1-yl)heptanephosphonate

5

A solution of 1,4,5-triphenyl-3-(7-bromoheptyl)-imidazol-2-one (1.0g) and triethylphosphite (1.66g) in xylene (5ml) was heated at reflux temperature for 40 hours. The solution was evaporated to an oil and chromatographed on silica gel (ethyl acetate/ethanol). The resulting oil was taken up in diethyl ether, filtered and evaporated to give diethyl 7-(3,4,5-triphenyl-2-oxo-2,3-dihydroimidazol-1-yl)heptanephosphonate as a clear oil (0.84g, 75%).

15 Found: C, 70.11; H, 7.37; N, 4.94%

 $C_{32}H_{39}N_2O_4P$

Requires: C, 70.31; H, 7.19; N, 5.12%

EXAMPLE 3

20

7-(3,4,5-Triphenyl-2-oxo-2,3-dihydroimidazol-1-yl)-heptanephosphonic acid

Diethyl 7-(3,4,5-triphenyl-2-oxo-2,3-dihydroimidazol-1-yl)heptanephosphonate (0.58g) was dissolved in dry chloroform, cooled to -40°C and to it was added trimethylsilyl iodide (1.05g) over 2 mins under an atmosphere of nitrogen. The cooling bath was removed and the reaction mixture was stirred for 2.5 hours at room temperature then evaporated to an oil and re-evaporated from methanol, treated with excess aqueous sodium bicarbonate, evaporated to an oil and re-evaporated from methanol, water and ethanol respectively. The oil was taken up in ethanol, treated

35

-11-

with excess aqueous sodium bicarbonate, evaporated to dryness then taken up in ethanol, filtered and the filtrate evaporated to an oil which solidified under ether. The solid was dissolved in water and passed down
5 a Dowex 1 x 2-200 ion exchange resin in the formate form. Elution with aqueous formic acid gave on evaporation of the solvent 7-(3,4,5-triphenyl-2-oxo-2,3-dihydroimidazol-1-yl)heptanephosphonic acid (0.078g, 15%) as a light brown foam.

10 Found: C, 66.50; H, 6.08; N, 5.39%
 $C_{28}H_{31}N_2O_4P$ + 3% water
Requires: C, 66.50; H, 6.52; N, 5.54%

EXAMPLE 4

15

Ethyl 7-(3,4,5-triphenyl-2-oxo-2,3-dihydroimidazol-1-yl)methylphosphinate

A solution of 1,4,5-triphenyl-3-(7-bromoheptyl)-
20 imidazol-2-one (2.0g) and diethyl methylphosphonite (2.17g) in toluene (15 ml) was heated at reflux temperature for 48 hours with the addition of more diethyl methylphosphonite (0.5g) after 24 hours. Water (5ml) was added and the solution was evaporated to an oil which was
25 chromatographed on silica gel (ethyl acetate/ethanol). The resulting oil was taken up in diethyl ether, filtered and evaporated to give ethyl 7-(3,4,5-triphenyl-2-oxo-2,3-dihydroimidazol-1-yl)methylphosphinate (1.34g), 65%) as a clear oil.

30 Found: C, 70.70; H, 7.53; N, 5.35%
 $C_{31}H_{37}N_2O_3P$ + 0.9% Et_2O + 2% H_2O
Requires: C, 70.56; H, 7.36; N, 5.26%.

-12-

BIOLOGICAL DATA**METHOD FOR MEASUREMENT OF AGGREGATION OF WASHED HUMAN PLATELETS**

5

Platelets were prepared from freshly drawn human blood. Blood was collected into acid citrate anticoagulant, centrifuged (5 min at 500g), and the upper layer of platelet-rich plasma was removed. This platelet-rich plasma was incubated with aspirin (100 μ M) for 10 min at 37°C and then centrifuged (15 min at 200g). The platelet pellet was resuspended (at approx. 1.5×10^8 cells/ml) in medium containing NaCl (145mM), KCl (5mM), MgCl₂ (1mM), CaCl₂ (0.2mM), Hepes (10mM, pH 7.4 at 37°C), glucose (10mM), apyrase (10 μ g/ml). Aggregation was monitored (as a change in optical density) at 37°C in a 4 channel aggregometer (PAP-4 from Biodata Corp.). Fibrinogen (1mg/ml) and CaCl₂ (1mM) were added to aliquots of platelets that were continuously stirred. The test compound (or 0.1% DMSO vehicle) was added 2 min before the aggregatory stimulus (1 μ M U46619). The extent of aggregation was assessed 4 min after addition of the stimulus, and was calculated as % of the control response in the absence of test compound. Dose-response curves were constructed for measurement of IC₅₀ values for each compound.

25

METHOD FOR MEASUREMENT OF K_I FOR INHIBITION OF [³H]ILOPROST BINDING TO HUMAN PLATELET MEMBRANES

30

Membranes were prepared from outdated platelet-rich plasma concentrates obtained from the Blood Transfusion Service. The platelets were homogenised in buffer containing Tris-Cl (5mM, pH 7.4 at 20°C) and EDTA

35

-13-

(0.25mM), and then centrifuged (10 min at 26,000g). The membrane pellet was washed twice by homogenisation in buffer containing Tris-Cl (50mM, pH 7.4 at 20°C) and EDTA (0.25 mM), followed by centrifugation. For measurement of [³H]iloprost binding, membranes (0.4-0.8mg) were incubated in the presence of Tris-Cl (50mM, pH 7.4 at 20°C), MgCl₂ (4mM), EDTA (40μM), [³H]iloprost (10nM), DMSO (1.85%), and varying concentrations of the test compounds. For determination of non-specific binding, 20μM iloprost was included. The tubes (triplicates for each condition) were set up on ice, and then incubated for 30 min at 37°C. The incubations were terminated by rapid filtration on Whatman GF/B filters using a Brandel Harvester. The filters were washed and then counted for radioactivity. The K_i of the test compounds for inhibition of binding of [³H]iloprost to human platelet membranes was calculated from the IC₅₀ for displacement of [³H]iloprost binding.

20 RESULTS

25


30

EXAMPLE NO.	AGGREGATION IC ₅₀ (μM)	K _i (μM)
EXAMPLE 1	0.25	8
EXAMPLE 2	5.50	30
EXAMPLE 3	43	1.29
EXAMPLE 4	21	19

-14-

Claims:

1. A compound of structure (I):



in which,

each group Ar is the same or different and is optionally substituted phenyl or optionally substituted heteroaryl;


15 R¹ is hydrogen, C₁₋₄alkyl, optionally substituted phenyl or optionally substituted heteroaryl;

n is 4 to 12; and

X is 5-tetrazolyl, SO₃H, P(O)(OR)₂, P(O)(OH)₂ or P(O)(R)(OR) in which R is hydrogen or C₁₋₄alkyl,
20 or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1 in which both groups R and R¹ are all phenyl, and n is 7.

25 3. A process for preparing a compound according to claim 1 in which X is other than 5-tetrazolyl which comprises reaction of a compound of structure (II):



-15-

in which

Ar, R¹ and n are as described for structure (I) and L
is a leaving group, with a suitable source of the group X;
and optionally thereafter forming a pharmaceutically
5 acceptable salt thereof.

4. A pharmaceutical composition comprising a
compound according to claim 1 or a pharmaceutically
acceptable salt thereof and a pharmaceutical carrier.
10

5. A compound according to claim 1 for use in
therapy.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 91/01543

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all)⁶

According to International Patent Classification (IPC) or to both National Classification and IPC

Int.Cl. 5 C07D233/70; A61K31/415; C07D403/06; C07F9/6506

II. FIELDS SEARCHED

Minimum Documentation Searched⁷

Classification System	Classification Symbols
Int.Cl. 5	C07D ; A61K ; C07F

Documentation Searched other than Minimum Documentation
to the Extent that such Documents are Included in the Fields Searched⁸

III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹

Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	EP,A,0 051 829 (A.NATTERMANN & CIE.GMBH) 19 May 1982 ---	

* Special categories of cited documents :¹⁰

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search 27 NOVEMBER 1991	Date of Mailing of this International Search Report 19. 12. 91
International Searching Authority EUROPEAN PATENT OFFICE	Signature of Authorized Officer DE BUYSER I.A.F.

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO. G**

GB 9101543
SA 51169

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 27/11/91

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0051829	19-05-82	DE-A- 3042466	16-06-82
		AT-T- 10626	15-12-84
		JP-A- 57108073	05-07-82
		US-A- 4349684	14-09-82
