




(86) Date de dépôt PCT/PCT Filing Date: 1995/05/04
(87) Date publication PCT/PCT Publication Date: 1995/11/16
(45) Date de délivrance/Issue Date: 2006/11/21
(85) Entrée phase nationale/National Entry: 1996/10/09
(86) N° demande PCT/PCT Application No.: US 1995/005219
(87) N° publication PCT/PCT Publication No.: 1995/030670
(30) Priorités/Priorities: 1994/05/06 (US238,817);
1994/12/02 (US349,361)

(51) Cl.Int./Int.Cl. *C07D 309/36* (2006.01),
C07D 309/32 (2006.01), *C07D 309/38* (2006.01),
C07D 311/02 (2006.01), *C07D 311/56* (2006.01),
C07D 311/74 (2006.01), *C07D 311/94* (2006.01),
C07D 405/12 (2006.01), *C07D 405/14* (2006.01),
C07D 407/12 (2006.01), *C07D 417/12* (2006.01),
C07D 473/00 (2006.01)


(72) Inventeurs/Inventors:
SKALETZKY, LOUIS L., US;
BUNDY, GORDON L., US;
LUKE, GEORGE P., US;
GAMMILL, RONALD B., US;
...

(73) Propriétaire/Owner:


(54) Titre : COMPOSES DE PYRANONE UTILES POUR TRAITER DES INFECTIONS A RETROVIRUS
(54) Title: PYRANONE COMPOUNDS USEFUL TO TREAT RETROVIRAL INFECTIONS




(I)



(II)



(III)



(IV)

(57) Abrégé/Abstract:

The present invention relates to compounds of formulae (I) and (II) which are pyran-2-ones, 5,6-dihydro-pyran-2-ones, 4-hydroxy-benzopyran-2-ones, 4-hydroxy-cycloalkyl[b]pyran-2-ones, and derivatives thereof, useful for inhibiting a retrovirus in a mammalian cell infected with said retrovirus, wherein R₁₀ and R₂₀ taken together are formulae (III) and (IV).



(72) **Inventeurs(suite)/Inventors(continued):** MORRIS, JOEL, US; ANDERSON, DAVID JOHN, US;
ROMINES, KAREN RENE, US; SKULNICK, HARVEY IRVING, US; ARISTOFF, PAUL ADRIAN, US;
JOHNSON, PAUL D., US; THAISRIVONGS, SUVIT, US; TURNER, STEVEN RONALD, US;
STROHBACH, JOSEPH WALTER, US; TOMMASI, RUBEN A., US; SCHWARTZ, THERESA M., US

(73) **Propriétaires(suite)/Owners(continued):** PHARMACIA & UPJOHN COMPANY, US

(74) **Agent:** MACRAE & CO.




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 309/32, 309/38, 401/12, 403/12, 417/12, A61K 31/365, 31/44, 31/47, 31/415, 31/505, 31/54		A3	(11) International Publication Number: WO 95/30670
			(43) International Publication Date: 16 November 1995 (16.11.95)
(21) International Application Number: PCT/US95/05219		Road, Whitehouse Station, NJ 07960 (US). STROHBACH, Joseph, Walter [US/US]; 54490 M-66, Mendon, MI 49072 (US). TURNER, Steven, Ronald [US/US]; 1713 Banbury Road, Kalamazoo, MI 49001 (US). THAISRIVONGS, Suvit [US/US]; 5695 Swallow, Kalamazoo, MI 49002 (US). ARISTOFF, Paul, Adrian [US/US]; 1650 Brookmoor Lane, Kalamazoo, MI 49002 (US). JOHNSON, Paul, D. [US/US]; 6960 Towhee, Portage, MI 49002 (US). SKULNICK, Harvey, Irving [US/US]; 1745 Old Deer Run, Kalamazoo, MI 49009 (US). SKALETZKI, Louis, L. [US/US]; 3731 Greenleaf Circle, Kalamazoo, MI 49008 (US). ANDERSON, David, John [GB/US]; 3809 Middlebury, Kalamazoo, MI 49006 (US). MORRIS, Joel [US/US]; 3001 Applelane, Kalamazoo, MI 49008 (US). GAMMILL, Ronald, B. [US/US]; 6704 Pleasantview Drive, Portage, MI 49002 (US). LUKE, George, P. [US/US]; Apartment 3 B, 6365 Bravo Court, Kalamazoo, MI 49002 (US). (74) Agent: GAMMILL, Martha, A.; The Upjohn Company, Corporate Intellectual Property Law, 301 Henrietta Street, Kalamazoo, MI 49001 (US). (81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).	
(22) International Filing Date: 4 May 1995 (04.05.95)			
(30) Priority Data: 08/238,817 6 May 1994 (06.05.94) US 08/349,361 2 December 1994 (02.12.94) US			
(60) Parent Application or Grant (63) Related by Continuation US 08/349,361 (CIP) Filed on 2 December 1994 (02.12.94)			
(71) Applicant (for all designated States except US): THE UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): ROMINES, Karen, Rene [US/US]; 62181 41st Street, Paw Paw, MI 49079 (US). BUNDY, Gordon, L. [US/US]; 7622 Ravenswood Drive, Portage, MI 49002 (US). SCHWARTZ, Theresa, M. [US/US]; 8921 West C Avenue, Kalamazoo, MI 49009 (US). TOMMASI, Ruben, A. [US/US]; 1 Plantation			
2187523			
		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. (88) Date of publication of the international search report: 25 April 1996 (25.04.96)	


(54) Title: PYRANONE COMPOUNDS USEFUL TO TREAT RETROVIRAL INFECTIONS

(57) Abstract


The present invention relates to compounds of formulae (I) and (II) which are pyran-2-ones, 5,6-dihydro-pyran-2-ones, 4-hydroxy-benzopyran-2-ones, 4-hydroxy-cycloalkyl[b]pyran-2-ones, and derivatives thereof, useful for inhibiting a retrovirus in a mammalian cell infected with said retrovirus, wherein R₁₀ and R₂₀ taken together are formulae (III) and (IV).




(I)



(II)



(III)



(IV)

PYRANONE COMPOUNDS USEFUL TO TREAT RETROVIRAL INFECTIONS

The present invention relates to compounds useful for inhibiting a retrovirus in a human cell infected with said retrovirus. More particularly, the present invention provides pyran-2-ones, 5,6-dihydropyran-2-ones, 4-hydroxy-benzopyran-2-ones, 4-hydroxy-cycloalkyl[b]pyran-2-ones, and derivatives thereof as HIV-proteinase inhibitors.

BACKGROUND OF THE INVENTION

During the past decade, acquired immunodeficiency syndrome (AIDS) has progressed from having the status of a medical curiosity afflicting only a small number of individuals to a problem of major proportions, both medically and economically. John Saunders and Richard Storer, "New Developments in RT Inhibitors," DN&P 5(3), April 1992, pages 153-169. WHO figures reveal that more than 360,000 cases of AIDS have been reported worldwide, including nearly 175,000 cases in the U.S.A. Of these, approximately 100,000 worldwide (50,000 in the U.S.A.) were reported in the preceding 12-month period. In the U.S.A., the number of seropositive individuals is thought to be approximately two million, and estimates suggest that 5-10 million people worldwide may be seropositive. Saunders and Storer, page 153.

Since the first description of the malady in the early part of this decade, acquired immunodeficiency disease syndrome (AIDS) and its devastating consequences have been subjects of continuous and intense coverage in both the lay and scientific press. Indeed, an edition of Scientific American was entirely devoted to AIDS (Scientific American 289, #4 (1988)), and the literature on the disease and the virus is already so vast as to defy thorough citation.

On March 20, 1987, the FDA approved the use of the compound, zidovudine (AZT), to treat AIDS patients with a recent initial episode of pneumocystis carinii pneumonia, AIDS patients with conditions other than pneumocystis carinii pneumonia or patients infected with the virus with an absolute CD4 lymphocyte count of less than $200/\text{mm}^3$ in the peripheral blood. AZT is a known inhibitor of viral reverse transcriptase, an enzyme necessary for human immunodeficiency virus replication. U.S. Patent 4,724,232 claims a method of treating humans having acquired immunodeficiency syndrome utilizing 3'-azido-3'-deoxy-thymidine (azidothymidine, AZT).

Following the discovery of the anti-HIV activity of AZT, much effort has been focused on a wide variety of other dideoxynucleoside analogues in the search for

superior agents. In the case of the 2'.3'-dideoxy series, ddC and ddI have shown potent activity against HIV *in vitro* and have been evaluated in clinical trials. Saunders and Storer, page 160. The compound ddC is currently being developed by Hoffman-La Roche Co. as a potential anti-AIDS drug. Its limiting toxicity in

5 humans is peripheral neuropathy which is reversible at low doses. Raymond R. Schinazi, Jan R. Mead and Paul M. Feorino, "Insights Into HIV Chemotherapy," AIDS Research and Human Retroviruses, Vol. 8, Number 6, 1992, pages 963-990. It has been approved by the FDA for AIDS therapy in combination with AZT. The compound ddI has also been evaluated in clinical trials. Its limiting toxicities are
10 peripheral neuropathy and pancreatitis. It has also been shown to stimulate hepatic glycolysis leading to irreversible liver damage. Schinazi, Mead and Feorino, page 966. It has recently been approved by the FDA for the treatment of HIV-1 infections in adults and pediatric patients who are intolerant to or whose health has significantly deteriorated while on AZT treatment. Schinazi, Mead and Feorino,
15 page 966.

Among these approved drugs, AZT is currently the only drug that has been shown to decrease the mortality and frequency of opportunistic infections associated with AIDS. Schinazi, Mead and Feorino, page 963.

Human immunodeficiency virus (HIV) has long been recognized as the
20 causative agent in AIDS, although a minority opinion to the contrary has been expressed (e.g., P. Duesberg, Proc. Natl. Acad. Sci., USA, 86:755-764 (1989)). Sequence analysis of the complete genomes from several infective and non-infective HIV-isolates has shed considerable light on the make-up of the virus and the types of molecules that are essential for its replication and maturation to an infective
25 species. The HIV protease is essential for the processing of the viral gag and gag-pol polypeptides into mature virion proteins. L. Ratner, et al., Nature, 313:277-284 (1985); L.H. Pearl and W.R. Taylor, Nature, 329:351 (1987). HIV exhibits the same gag/pol/env organization seen in other retroviruses. L. Ratner, et al., above; S. Wain-Hobson, et al., Cell, 40:9-17 (1985); R. Sanchez-Pescador, et al., Science, 227:484-492
30 (1985); and M.A. Muesing, et al., Nature, 313:450-458 (1985).

Reverse transcriptase (RT) is an enzyme unique to retroviruses that catalyzes the conversion of viral RNA into double stranded DNA. Blockage at any point during the transcription process, by AZT or any other aberrant deoxynucleoside triphosphate incapable of elongation, should have dramatic consequences relative to
35 viral replication. Much work on the RT target is in progress based, in large measure, upon the fact that nucleosides like AZT are easily delivered to cells.

However, the inefficiency of phosphorylation steps to the triphosphate, and the lack of specificity and consequent toxicity, constitute major drawbacks to use of AZT and similar nucleosides having a blocked, or missing, 3'hydroxyl group.

The T4 cell receptor for HIV, the so-called CD4 molecule, has also been targeted as an intervention point in AIDS therapy. R.A. Fisher, et al., *Nature*, 331:76-78 (1988); R.E. Hussey, et al., *Nature*, 331:78-81 (1988); and K.C. Deen, et al., *Nature*, 331:82-84 (1988). The exterior portion of this transmembrane protein, a molecule of 371 amino acids (sCD4) has been expressed in Chinese hamster ovary (CHO) cells and Genentech (D.H. Smith, et al., *Science*, 238:1704-1707 (1987)) has had a product in clinical trials since the fall of 1987. CD4 has been shown to have a narrow spectrum of activity against wild-type virus and so far has failed to control HIV infection in humans. Schinazi, Mead and Feorino, page 963. The idea behind CD4 based therapy is that the molecules can neutralize HIV by interfering with viral attachment to T4, and other cells which express CD4 on their surfaces. A variant on this theme is to attach cell toxins to CD4 for specific binding and delivery to infected cells which display glycoprotein gp-120 on their surfaces. M.A. Till, et al., *Science*, 242:1166-1168 (1988); and V.K. Chaudhary, et al., *Nature*, 335:369-372 (1988).

Another therapeutic target in AIDS involves inhibition of the viral protease (or proteinase) that is essential for processing HIV-fusion polypeptide precursors. In HIV and several other retroviruses, the proteolytic maturation of the gag and gag/pol fusion polypeptides (a process indispensable for generation of infective viral particles) has been shown to be mediated by a protease that is, itself, encoded by the pol region of the viral genome. Y. Yoshinaka, et al., *Proc. Natl. Acad. Sci. USA*, 82:1618-1622 (1985); Y. Yoshinaka, et al., *J. Virol.*, 55:870-873 (1985); Y. Yoshinaka, et al., *J. Virol.*, 57:826-832 (1986); and K. von der Helm, *Proc. Natl. Acad. Sci., USA*, 74:911-915 (1977). Inhibition of the protease has been shown to inhibit the processing of the HIV p55 in mammalian cell and HIV replication in T lymphocytes. T.J. McQuade, et al., *Science*, 247:454 (1990).

The protease (or proteinase), consisting of only 99 amino acids, is among the smallest enzymes known, and its demonstrated homology to aspartyl proteases such as pepsin and renin (L.H. Pearl and W.R. Taylor, *Nature*, 329:351-354 (1987); and I. Katoh, et al., *Nature*, 329:654-656 (1987)), led to inferences regarding the three-dimensional structure and mechanism of the enzyme (L.H. Pearl and W.R. Taylor, above) that have since been borne out experimentally. Active HIV protease has been expressed in bacteria (see, e.g., P.L. Darke, et al., *J. Biol. Chem.*, 264:2307-2312

(1989)) and chemically synthesized (J. Schneider and S.B. Kent, Cell, 54:363-368 (1988); and R.F. Nutt, et al., Proc. Natl. Acad. Sci., USA, 85:7129-7133 (1988)). Site directed mutagenesis (P.L. Darke, et al., above); and N.E. Kohl, et al., Proc. Natl. Acad. Sci., USA, 85:4686-4690 (1988)) and pepstatin inhibition (P.L. Darke, et al., J. Biol. Chem., 264:2307-2312 (1989); S. Seelmeier, et al., Proc. Natl. Acad. Sci., USA, 85:6612-6616 (1988); C.-Z. Giam and I. Borsos, J. Biol. Chem., 263:14617-14720 (1988); and J. Hansen, et al., EMBO J., 7:1785-1791 (1988)) have provided evidence for HIV protease's mechanistic function as an aspartyl protease. A study has demonstrated that the protease cleaves at the sites expected in peptides modeled after the regions actually cleaved by the enzyme in the gag and pol precursor proteins during viral maturation. P.L. Darke, et al., Biochem. Biophys. Res. Commun., 156:297-303 (1988). X-ray crystallographic analysis of the HIV-protease (M.A. Navia, et al., Nature, 337:615-620 (1989)) and a related retroviral enzyme from Rous sarcoma virus (M. Miller, et al., Nature, 337:576-579 (1989)) reveal an active site in the protease dimer that is identical to that seen in other aspartyl proteases, thus supporting the supposition (L.H. Pearl and W.R. Taylor, above) that the HIV enzyme is active as a dimer. See also Joseph A. Martin, "Recent Advances in the Design of HIV Proteinase Inhibitors," Antiviral Research, 17 (1992) 265-278.

To date, the scientific search for a fully effective and safe means of inhibiting retroviruses in a human hosting such a virus, and thereby effectively treating diseases caused by such a virus, such as acquired immunodeficiency syndrome (AIDS), continues.

INFORMATION DISCLOSURE

JO 3227-923-A (Sawai Seiyaku KK) discloses the use of 4-hydroxy-coumarins as therapeutic agents for HIV-infected patients; however, unsubstituted 4-hydroxy-coumarin is the only compound specifically disclosed for this use.

WO 91/04663 (Univ. of Calif. at Oakland) discloses 6-amino-1,2-benzopyrones which are useful for treating viral diseases.

WO 91/12804 (Kabi Pharmaceutical), published 5 September 1991, discloses the use of N-phenyl-N-methyl-1,2-dihydro-4-hydroxy-1-methyl-2-oxo-quinoline-3-carboxamide also known as Linomide®, for the treatment of retrovirus infections.

International Publication No. WO 89/07939, published 8 September 1989, discloses specific coumarin compounds which are reverse transcriptase inhibitors.

U.S. Patents Nos. 3,489,774 and 3,493,586 disclose 3-(beta-aryl-beta-(arylothio) (or aryl seleno) propionyl-coumarin and pyrone products useful as parasiticides.

Biochemical and Biophysical Research Communications, Vol. 188, No. 2,

1992, pages 631-637, discloses chromones bearing hydroxyl substituents and a phenolic group at the 2-position (flavones) as having anti-HIV-1 proteinase activity.

Antimicrobial Patent Fast-Alert, Week Ending 4 September 1992, disclose gamma-pyrones, gamma-pyridones and gamma-thio-pyrones as antiviral agents.

5 International Publication Nos. WO 92/04326, 92/04327 and 92/04328, all published 19 March 1992, disclose antiviral heterocyclic derivatives, such as quinolinones and benzopyranones, as replication inhibitors for treating herpes simples 1 and 2, cytomegalovirus and Epstein-Barr virus.

C.A. Selects: Antitumor Agents, Issue 19, 1992, page 25, No. 117: 90147q
10 (PCT International Application WO 92 06,687) discloses the preparation of 5-iodo-5-amino-1,2-benzopyrones and analogs as cytostatic and antiviral agents.

Nowhere do these references teach or suggest the use of 4-hydroxy- α -pyrones as HIV protease inhibitors or as having antiviral activity.

Phytochemistry, 31(3):953-956 (1992), discloses compounds, such as 4-
15 hydroxy- α -(4-methoxyphenyl)-6-[2-(4-methoxyphenyl)ethenyl]-2-oxo-, methyl ester, (E)-(-)-2H-pyran-3-acetic acid.

Tetrahedron, 48(9):1695-1706 (1992), (see also Tetrahedron Lett., 30(23):3109-
12 (1989)), discloses compounds, such as 3-[1-(4-chlorophenyl)-3-(4-nitrophenyl)-2-propenyl]-4-hydroxy-6-methyl-2H-pyran-2-one; 3-[3-(4-chlorophenyl)-1-(4-
20 nitrophenyl)-2-propenyl]-4-hydroxy-6-methyl-2H-pyran-2-one; 4-hydroxy-3-[3-(4-methoxyphenyl)-1-(4-nitrophenyl)-2-propenyl]-6-methyl-2H-pyran-2-one; and 4-hydroxy-3-[1-(4-methoxyphenyl)-3-(4-nitrophenyl)-2-propenyl]-6-methyl-2H-pyran-2-one.

Tennen Yuki Kagobutsu Toronkai Koen Yoshishu, 30:17-24 (1988), discloses
25 compounds, such as 4-hydroxy- β -(4-methoxyphenyl)-6-[2-(4-methoxyphenyl)ethenyl]-2-oxo-, methyl ester, (E)-(-)-2H-pyran-3-propanoic acid.

Chem. Absts. 53:15072f discloses compounds, such as α -1,3-dihydroxy-2-butenylidene- β -ethyl-, δ -lactone, hydrocinnamic acid.

Chem. Absts. 53:15072c discloses compounds, such as α -1,3-dihydroxy-2-
30 butenylidene- β -isopropyl-, δ -lactone, hydrocinnamic acid.

Arch. Pharm. (Weinheim, Ger.), 316(12):988-94 (1983), discloses compounds, such as 3-[1-(4-chlorophenyl)-3-oxobutyl]-4-hydroxy-6-methyl-2H-pyran-2-one; and 3-[1-(4-chlorophenyl)propyl]-4-hydroxy-6-methyl-2H-pyran-2-one.

Chem. Ber., 110(3):1047-57 (1977), discloses compounds, such as 6-(3,4-
35 dimethoxyphenyl)-3-[2-(3,4-dimethoxyphenyl)-1-(4-methoxy-2-oxo-2H-pyran-6-yl)ethyl]-4-hydroxy-2H-pyran-2-one; and 3-[2-(3,4-dimethoxyphenyl)-1-(4-methoxy-2-

oxo-2H-pyran-6-yl)ethyl]-4-hydroxy-6-[2-(4-methoxyphenyl)ethyl]-2H-pyran-2-one.

J. Heterocycl. Chem., 23(2):413-16 (1986), discloses compounds, such as 3-[(4-chlorophenyl)-1-piperidinylmethyl]-4-hydroxy-6-methyl-2H-pyran-2-one.

The following published PCT applications disclose peptides useful as
5 retroviral protease inhibitors: International Publication No. WO 91/06561, published 16 May 1991; and International Publication No. WO 92/17490, published 15 October 1992.

The following references disclose pyrone compounds which are believed to be representative of those known in the art:

- 10 EP-443449 (German language) discloses 3-hexyl-5,6-dihydro-6-pentyl-2H-pyran-2-one and 3-ethyl-6-hexadecyl-5,6-dihydro-4-hydroxy-2H-pyran-2-one. Pestic. Sci., 27(1):45-63 (1989), discloses 5,6-dihydro-4-hydroxy-6-methyl-6-(1-methyl-1-propenyl)-3-(1-oxobutyl)-2H-pyran-2-one; and 6-cyclopropyl-5,6-dihydro-4-hydroxy-6-methyl-3-(1-oxobutyl)-2H-pyran-2-one. Acta. Chem. Scand., 43(2):193-95 (1989),
15 discloses 4-(acetyloxy)-5,6-dihydro-3,6-dimethyl-2H-pyran-2-one. J. Org. Chem., 54(14):3383-9 (1989), discloses 5,6-dihydro-4-hydroxy-3,6,6-trimethyl-2H-pyran-2-one. J. Org. Chem., 53(6):1218-21 (1988); and Tetrahedron Lett., 34(2):277-80 (1993), discloses 3-hexyldihydro-6-undecyl-2H-pyran-2,4 (3H)-dione, (6R)-. J. Chem. Soc. Perkins Trans., 1(6):1157-9 (1985), discloses dihydro-3-methyl-6-nonyl-6-
20 [[(tetrahydro-2H-pyran-2-yl)oxy]methyl]-2H-pyran-2,4 (3H)-dione. J. Chem. Ecol., 9(6):703-14 (1983), discloses 5,6-dihydro-4-hydroxy-3,6-dimethyl-2H-pyran-2-one. J. Org. Chem., 48(7):1123-5 (1983), discloses 6-(2-chloro-1-methylethenyl)-5,6-dihydro-4-hydroxy-3-methyl-2H-pyran-2-one, (Z)-(.+.-)-. Acta. Chem. Scand., 43(2):193-95 (1989); and Tetrahedron Lett., 21(6):551-4 (1980), discloses 5,6-dihydro-4-hydroxy-
25 3,6-dimethyl-2H-pyran-2-one. Helv. Chem. Acta, 59(7):2393-2401 (1976), discloses 4-[(3,6-dihydro-4-hydroxy-5-methyl-6-oxo-2H-pyran-2-yl)methyl]-2,6-piperidinedione. Acta. Chem. Scand., 30(7):613-18 (1976); and Tetrahedron Lett., 22:1903-4 (1976), discloses 5,6-dihydro-4-hydroxy-3-methyl-6-(1-methyl-1-propenyl)-2H-pyran-2-one, (E)-. 3,3'-[(4-nitrophenyl)methylene]bis[5,6-dihydro-4-hydroxy-6-methyl-2H-pyran-2-
30 one; and 3,3'-(phenylmethylene)bis[5,6-dihydro-4-hydroxy-6-methyl-2H-pyran-2-one are disclosed in Synth. Commun., 20(18):2827-2836, 1990.

WO 93/07868, published 29 April 1993, discloses new nitroso-benzopyrone, -benzamide and -isoquinolinone derivatives as adenosine di-phospho:ribose transferase inhibitors for treating viral infections and cancer.

35 WO 93/07128, published 15 April 1993, relates to substituted cyclic carbonyls and derivatives thereof useful as retroviral protease inhibitors.

J. Indian Chem. Soc., 69:397-398 (July 1992), discloses that coumarin-4-acetic acids were screen for their anticancer and anti-AIDS activities and were found to be inactive.

The Journal of Antibiotics, 46(7):1126 (July 1993), discloses germicidin, which is 6-(2-butyl)-3-ethyl-4-hydroxy-2-pyrone, to be an autoregulative germination inhibitor of *Streptomyces viridochromogenes* NRRL B-1551.

Derwent Abstracts, 93-168920/21 of EP 543201 discloses the use of coumarin derivatives, such as 1-(N-morpholyl)-6-(4-hydroxybenzoic acid ethyl ester) hexane, for the treatment of viral infections, such as influenza or acute rhinitis.

J. Org. Chem., 48(22):3945-7 (1983); and Chem. Pharm. Bull., 29(10):2762-8 (1981); disclose compounds such as 4-hydroxy-6-(3-pyridinyl)-2H-pyran-2-one.

J. Labelled Compd. Radiopharm., 28(10):1143-8 (1990), discloses compounds such as 4-hydroxy-6-methyl-2H-pyran-2-one.

J. Am. Chem. Soc., 113(25):9585-95 (1991), discloses compounds such as 3-(3-phenyl-2-propen-1-yl)-6-methyl-4-hydroxy-2H-pyran-2-one.

CA 54:14239d and CA 53:4272c disclose compounds such as α -(α , γ -dihydroxycinnamylidene)-, δ -lactone hydrocinnamic acid.

CA 53:15072f discloses compounds such as α -1,3-dihydroxy-2-butenylidene- β -ethyl-, δ -lactone hydrocinnamic acid.

Synth. Commun., 20(18):2827-36 (1990), discloses compounds such as 3,3'-[(4-nitrophenyl)methylene]bis[5,6-dihydro-4-hydroxy-6-methyl-2H-pyran-2-one, and 3,3'-(phenylmethylene)bis[5,6-dihydro-4-hydroxy-6-methyl-2H-pyran-2-one.

J. Org. Chem., 54(14):3383-9 (1989), discloses compounds such as 5,6-dihydro-4-hydroxy-3,6,6-trimethyl-2H-pyran-2-one.

Derwent Abstract, 92-166863/20, of EP 553248 discloses new optionally substituted 5-iodo-6-amino-1,2-benzopyrone derivatives, which are adenosine di:phospho-ribose inhibitors, for treatment and prevention of viruses and tumors associated with AIDS.

Synthesis of Heterocycles. XV. 4-Hydroxy-2-pyronecycloenes. E. Ziegler, H. Junek, and E. Nolken, Monatsh., 89: 678-82 (1958) (CA 53:12283-4) discloses compounds such as the following: 4-hydroxy-3-benzyl-5,6-octamethylene-2-pyrone; 4-hydroxy-3-benzyl-5,6-pentamethylene-2-pyrone; 4-hydroxy-3-benzyl-5,6-heptamethylene-2-pyrone; 4-hydroxy-3-benzyl-5,6-hexamethylene-2-pyrone; and 4-hydroxy-3-benzyl-5,6-tridecamethylene-2-pyrone.

R. Effenberger, T. Ziegler, K.-H. Schonwalder, T. Kesmarszky, B. Bauer, Chem. Ber 119: 3394-3404 (1986), discloses pyrone intermediates, such as those of

formula J-1 (wherein n is 4; refer to Chart J below).

Monatsh. Chem., 119(6-7): 727-37 (1988) (CA 110(13):114430k) discloses the compounds 8H-acenaphtho[1,2-b]pyran-8-one, 10-hydroxy-9-(phenylmethyl)-; and indeno[2,1-b]pyran-3(5H)-one, 1-hydroxy-2-(phenylmethyl)-.

5 CA 54:14239b discloses the compound 3-benzyl-4-hydroxy-2-oxoindeno-[1,2-b]pyran.

Monatsh. Chem., 113(4): 475-84 (1982) discloses compounds such as 6,7-dihydro-4-hydroxy-6-(3-methylphenyl)-7-phenyl-3-(phenylmethyl)-pyrano[2,3-c]pyrrole-2,5-dione; and 6,7-dihydro-4-hydroxy-6,7-diphenyl-3-(phenylmethyl)-
10 pyrano[2,3-c]pyrrole-2,5-dione.

Monatsh. Chem. 90: 594-9 (1959) (CA 54:14238g,h) discloses compounds such as 5H-benzocycloheptene-8-acrylic acid, α -benzyl-6,7-dihydro- β ,9-dihydroxy- δ -lactone; and 3-benzyl-5,6,7,8-tetrahydro-4-hydroxy-8-isopropyl-5-methyl-coumarin.

Bull. Soc. Chim. Fr. 5: 1719-23 (1069) (Fr) (CA 71(21): 101655p) discloses the
15 compound 3-benzyl-5,6,7,8-tetrahydro-4-hydroxy-coumarin.

WO 8804652 (equivalent AU 8810440 (Jap.)) discloses the compound 3-(4-chloro-2-nitrobenzoyl)-5,6,7,8-tetrahydro-4-hydroxy-2H-1-benzopyran-2-one.

Monatsh. 92: 246-53 (1961) (Gr) (CA 55:27296d) discloses the compound 3-(3,5-dimethylsalicyl)-5,6,7,8-tetrahydro-4-hydroxy-coumarin.

20 CA 94(9): 65472r discloses 5,6,7,8-hexahydro-3-phenyl-2-H-cycloocta[b]pyran-2-one; and 6,7,8,9-tetrahydro-4-hydroxy-3-phenyl-cyclohepta[b]pyran-2(5H)-one.

J. Org. Chem. 28(11): 3112-14 (1963) (CA 59:15185e) discloses the compound hexanedioic acid, 2-[hydroxy(2-hydroxy-1-cyclopenten-1-yl)methylene]-, δ -lactone.

Antimicrobial Patent Fast-Alert, Week Ending 30 April 1993, discloses cyclic
25 ureas and analogues useful as retroviral protease inhibitors.

Many 4-hydroxy-coumarin type compounds are known. For example, these references -- CA 54:577e,g,h (1960); U.S. Patent 2,872,457 (CA 53:12305e (1959)); CA 51:14826f,h (1957); U.S. Patent 2,723,276 (CA 52:5480g,h (1958)); CA 51:14827a,b (1957); CA 51:16453a (1957); CA 54:5699d (1960); CA 54:16450f (1960); CA
30 53:22454a (1959); and CA 53:20046a-- disclose compounds such as the following: 4-hydroxy-3-(1-phenylbutyl)-coumarin; 4-hydroxy-3-(1-phenylpentyl)-coumarin; 3-(cyclohexylphenylmethyl)-4-hydroxycoumarin; 4-hydroxy-3-(2-methyl-1-phenylpropyl)-coumarin; 4-hydroxy-3-(2-phenylpropyl) coumarin; 4-hydroxy-3-(1,3-diphenylpropyl)-coumarin; 4-hydroxy-3-(1-(4-methylphenyl)-butyl)-coumarin; 4-
35 hydroxy-3-(1-(1-naphthyl)-propyl)-coumarin; 4-hydroxy-7-methyl-3-(1-phenylpropyl)-coumarin; 7-chloro-4-hydroxy-3-(1-phenylpropyl)-coumarin; 4-hydroxy-3-[1-(4-

methoxyphenyl)propyl]-coumarin; 3-(.alpha.-ethyl-p-fluorobenzyl)-4-hydroxy-coumarin; 3-(.alpha.-ethyl-p-methoxybenzyl)-4-hydroxy-coumarin; and 3-(1-phenyl-propenyl)-4-hydroxy-coumarin.

To the best of our knowledge, from our review, these references do not
5 disclose the use of these compounds as HIV protease inhibitors. They are disclosed as being useful as: rodenticides, lowering the prothrombin level of the blood, blood anticoagulants, and pesticides.

Additional 4-hydroxy-coumarin compounds with similar uses have been disclosed in the following references:

10 Indian J. Chem., Sect. B, 25B: 1167-70 (1986) (CA 107(17):154201f) and CA 93(23):220546t discloses the compound 4-Hydroxy-3-(1-phenyl-2-propenyl)-coumarin.

CA 96(19):157432x; CA 90(1):1707f; CA 84(9):55338f; CA 79(13):74969a; and CA 71(15):69677j disclose the compound 4-hydroxy-3-[1-(1,2,3,4-tetrahydro)naphthyl]-coumarin; CA 54:579e discloses the compound 4-hydroxy-3-[1-
15 indanyl]-coumarin; CA 63:14743c discloses the compound 4-hydroxy-3-(1-naphthylmethyl)-coumarin; CA 63:5589c discloses the compound 3-(1'-(2-methoxy,3-methyl,5-chloro-phenyl)propyl)-4-hydroxy-coumarin; CA 64:12969b discloses the compound 3-(.alpha.-acetyl-.alpha.-acetylbenzyl)-4-hydroxy-coumarin.

CA 79(13):74969a; Chim. Ther. 7(4): 300-6 (1972) (Fr) (CA 78(7):38016h); CA
20 52:5399b; CA 54:5699e; CA 54:579e; and CA 72(15):78882v disclose 4-hydroxycoumarin compounds substituted at the 6- or 7-position by, e.g., methyl, methoxy and chloro.

J. M. Mulder, U.S. Patent No. 3,835,161, 10 September 1974, discloses the compound 3-[1-[4-(2-bromoethyl)phenyl]ethyl]-4-hydroxy-2H-1-benzopyran-2-one.

25 Merck Index, Eleventh Edition, (1989), Entry 9950, discusses Warfarin, its chemical name -- 3- α -phenyl- β -acetyethyl-4-hydroxycoumarin -- and its uses as a rodenticide and an anticoagulant. J. Med. Chem., 1978, Vol. 21, No. 2: 231-234, discloses the antivitamin K activity of warfarin and discusses the anticoagulant activity of several 3-substituted 4-hydroxycoumarins such as 4-Hydroxy-3-(1-
30 phenylbutyl)-coumarin; and 4-hydroxy-3-(.alpha.-methylbenzyl)-coumarin. J. Am. Chem. Soc. 83: 2676-9 (1961) (CA 55:22306e (1961)) discusses the resolution and absolute configuration of warfarin and discloses the preparation of compounds such as 4-hydroxy-3-(1-phenylbutyl)-coumarin.

Journal of Labelled Compounds and Radiopharmaceuticals Vol. XXIII, No. 2:
35 137-148 (1986), discloses several deuterium labelled metabolites of warfarin and phenprocoumon, such as the deuterium labelled analog of the compound 4-hydroxy-

2187523

7-methoxy-3-(1-phenylpropyl)-coumarin.

J48023942 discloses compounds, such as 4-hydroxy-3-(α -methylbenzyl)-coumarin; 4-hydroxy-3-(3-methyl-1-phenylbutyl)-coumarin; and 2H-1-benzopyran-2-one, 4-hydroxy-7-methoxy-3-(1-phenylpropyl)- (also cited in preceding reference) and
5 their use as rodenticides.

Tr. Voronezh. Tekhnol. Inst. 19(2): 27-30 (1971), Abstract No. 1zh274 (Russian language), discloses the compound 4-hydroxy-3-phenethylcoumarin. This reference and Helv. Chim. Acta 74(7): 1451-8 (1991) disclose the compound of 4-hydroxy-3-(3-phenylpropyl)coumarin.

10 J. Org. Chem. 33(1): 437-8 (1968); and Eur. J. Med. Chem. - Chim Ther. 12(2): 125-30 (1977) disclose compounds such as 4-hydroxy-3-diphenylmethylcoumarin.

U.S. 3,764,693 discloses the compound 4-hydroxy-3-(3-hydroxy-1-phenylbutyl)-coumarin and its anticoagulating and rodenticidal activity.

15 J. Med. Chem. 18(5): 513-19 (1975) (CA 83(5):37913q); J. Chromatogr. 338(2): 325-34 (1985); J. Chromatogr. 562 (1-2): 31-8 (1991); J. Labelled Compds. Radiopharm. 23(2): 137-48 (1986) (cited previously); and J. Chromatogr. 529(2): 479-85 (1990) disclose compounds such as 4-hydroxy-3-[1-[3-(phenylmethoxy)phenyl]propyl]-2H-1-benzopyran-2-one; 4-hydroxy-8-
20 (phenylmethoxy)-3-(1-phenylpropyl)-2H-1-benzopyran-2-one; 4-hydroxy-3-[1-(4-hydroxyphenyl)propyl]-coumarin; 4-hydroxy-6-methoxy-3-(1-phenylpropyl)-coumarin; 4,7-dihydroxy-3-(1-phenylpropyl)-coumarin; 4,6-dihydroxy-3-(1-phenylpropyl)-coumarin; 4-hydroxy-3-[1-(3-hydroxyphenyl)propyl]-coumarin; and p-chlorophenprocoumon.

25 AIDS 1993, Vol. 7, No. 1, pages 129-130, discusses the effect of warfarin on HIV-1 replication and spread.

CA Selects:AIDS & Related Immunodeficiencies, Issue 24, 1993, Abstract 119:195147j discloses the inhibitory effect of a single dose of coumarin derivatives, warfarin, 4-hydroxy-coumarin, umbelliferone, on HIV-1 replication and cell-mediated
30 or cell-free viral transmission.

At the First National Conference on Human Retroviruses and Related Infections, 12-16 December 1993, Washington, D.C., it was disclosed that coumarins, such as warfarin, and pyrones, such as 3-(thiophenyl)-6-phenyl-4-hydroxy-pyrone, displayed HIV protease inhibition in an assay.

35 Biochemical and Biophysical Research Communications, Vol. 201, No. 1, pages 290-294 (30 May 1994) discloses that warfarin and structurally related

coumarin analogs are HIV-1 protease inhibitors.

J. Med. Chem. 37:2664-2677 (1994) discloses 4-hydroxy-3-(3-phenoxypropyl)-2H-1-benzopyran-2-one and structural analogs, especially 4,7-dihydroxy-3-[4-(2-methoxyphenyl)butyl]-2H-1-benzopyran-2-one, as HIV-1 protease inhibitors.

5 Biochemical and Biophysical Research Communications, Vol. 200, No. 3, pages 1658-1664 (16 May 1994) discloses that 4-hydroxy-3-(3-phenoxypropyl)-1-benzopyran-2-one and 4-hydroxy-6-phenyl-3-(phenylthio)-pyran-2-one, and structural analogs of these compounds, are inhibitors of HIV-1 protease.

J. Am. Chem. Soc. 116:6989-6990 (1994) discloses 4-hydroxy-6-phenyl-3-
10 (phenylthio)pyran-2-one, and structural analogs thereof, are HIV-1 protease inhibitors.

Acta. Virol. 37:241-250 (1993) discloses the anti-HIV activity of coumarin derivatives, warfarin, 4-hydroxy-coumarin and umbelliferone.

Antiviral Research 24:275-288 (1994) discloses bicyclic imidazo derivatives
15 (imidazothiazoles and imidazopyridines) which inhibit HIV-1 through interaction with reverse transcriptase (RT).

U.S. Patent No. 3,325,515 (J. Schmitt, et al.) discloses coumarin derivatives, such as 3-(4-hydroxy-3-coumarinyl)-3-phenyl-1-propionic acid methyl ester, as exhibiting anti-coagulant activity.

20 U.S. Patent No. 2,723,277 (A. Grussner, et al.) discloses malonic acid derivatives, such as 3-[1'-(p-chloro-phenyl)-propyl]-4-hydroxy-coumarin, as anti-coagulant agents.

FR, A, 1276654 discloses 4-hydroxy-coumarins, such as (2'-hydroxy)-3-benzyl-4-hydroxycoumarin, which have anti-coagulant, anti-bacterial or anti-fungal
25 properties.

BE, A, 674997 discloses 4-hydroxycoumarin derivatives, such as 3-(5-methoxytetralyl-(1))-4-hydroxycoumarin as agents for fighting rodents.

GB, A, 734142 discloses the preparation of 3-substituted-4-hydroxycoumarins, such as 3-(1-phenyl-2-acetyl)-ethyl-4-hydroxycoumarin and 3-(1-furan-2-acetyl)-ethyl-
30 4-hydroxycoumarin, which are effective as anti-coagulants and rodenticides.

"The Application of Computer-Assisted Drug Design in the discovery of Nonpeptide HIV-1 Protease Inhibitors", Parke-Davis Pharm. Res., Keystone Symposia, 5-11 March 1994, Santa Fe, NM, discloses 4-hydroxy-3-(3-phenoxypropyl)-1-benzopyran-2-one as an HIV protease inhibitor.

35 Structural Biology, 1(1):199-200 (April 1994) discloses that the rat poison warfarin was a useful lead in the search for HIV proteinase inhibitors.

CA 85:78002b (1976) discloses 3-(2,4,6-trihydroxybenzyl)-4-hydroxy-2H-pyran-2-one derivatives as having anti-bacterial activity.

FR, A, 1092278 (Hoffman) (1955) discloses the preparation of coumarin derivatives, such as 3-[1'-phenyl-propene-(1')-yl]-4-hydroxycoumarin.

5 International Publication No. WO 94/11361, published 26 May 1994, discloses pyran-2-ones and 5,6-dihydroxypyran-2-ones as retroviral protease inhibitors.

International Publication No. WO 94/18188, published 18 August 1994, discloses 4-hydroxy-benzopyran-2-ones and 4-hydroxy-cycloalkyl[b]pyran-2-ones as retroviral protease inhibitors.

10 The following references were cited against the immediate parent application as disclosing the state of the art:

U.S. Patent No. 3,651,091 (Boschetti, et al.); U.S. Patent No. 4,262,013 (Mistui, et al.); U.S. Patent No. 4,900,754 (Regan, et al.); U.S. Patent No. 5,294,724 (Jendralla, et al.); Australian Patent Specification 219,371 (Enders, et al.); Canadian
15 Patent No. 1,171,424 (Willard, et al.); British Patent Specification 836,740 (Bayer); European Patent Application 0 024 348 (Willard, et al.); European Patent Application 0 588 137 (Fischer, et al.); French Patent No. 1,276,654 (Molho) (cited above); and International Publication No. WO 94/1136 (Thaisrivongs, et al.) (cited above).

20 "Collaborative Structure-Based Design of Small Organic Molecules as Inhibitors of HIV Proteases," Keystone Symposia, Santa Fe, NM (5-11 March 1994), discloses the crystallographic complexes of HIV-1 and HIV-2 protease with compounds, such as 3-(α -ethylbenzyl)-6-(α -ethylphenethyl)-4-hydroxy-2H-pyran-2-one.

25 "Discovery and Properties of Small Organic Molecules Inhibiting HIV-1 Protease," Keystone Symposia, Santa Fe, NM (5-11 March 1994), discloses an assay for determining inhibitory activity of compounds, such as 3-(α -ethylbenzyl)-6-(α -ethylphenethyl)-4-hydroxy-2H-pyran-2-one.

"Structure-based Design of Non-peptide HIV Protease Inhibitors," 35th
30 Annual Buffalo Medicinal Chemistry Symposium, Buffalo, NY (22-25 May 1994), discloses compounds, such as 3-(α -ethylbenzyl)-6-(α -ethylphenethyl)-4-hydroxy-2H-pyran-2-one, as potential anti-HIV therapeutic agents.

In Hruby et. al. (J. Org. Chem., 58 (26):7567 (1993), a description of the copper catalyzed addition of an aryl Grignard to an unsaturated chiral amide, 3-(2-butenoyl)-4-phenyl-2-oxazolidinone, is given. In Evans et. al. (J. Am. Chem. Soc.,
35 112:8215 (1990), the reaction between a chiral amide and 2-methoxy-2-methyl-1,3-

dioxoline is described. The preparation of 2-methoxy-2-methyl-1,3-dioxoline is found in Santry et. al. (J. Am. Chem. Soc., 110 (9):2910 (1988). For references on the reaction between an ester enolate and a ketone, refer to Dongala et. al., Tetrahedron Letters, 4983 (1973), and Mitsui et. al., Tetrahedron, 23:4271 (1967). For references on the reaction between an amide enolate and a ketone, refer to Viteva et. al., Tetrahedron 50:7193 (1994); Oare et. al., J. Org. Chem. 55:132 (1990); Hullot et. al., Can. J. Chem. 55:266 (1977); Woodbury et. al., J. Org. Chem. 42:1688 (1977); Stefanovsky et. al., Tetrahedron 42:5355 (1986); and Mathew et. al., U.S. Patent No. 5,284,975.

- 10 G. Carganico, P. Cozzi, G. Orsini, J. Med. Chem., 26:1767-1769 (1983), discloses synthesized compounds with a methyl and a hydroxyl group at the 4-position of the dihydropyrone ring and no substitution at the 3-position. The compounds of the present invention have a ketone at the 4-position (which may be in enol form) and substitution at the 3-position.
- 15 D.T. Witiak et al., J. Med. Chem., 31:1437-1445 (1988), discloses benzopyran-2-ones with a hydroxy group at the 3-position. The compounds of the present invention have alkyl substitution at that position.

B. Tait, Winter Conference on Bioorganic Medicinal Chemistry, 29 Jan.-2 Feb. 1995, Steamboat Springs, Colorado, disclosed a dihydropyrone having a phenyl group and a pentyl group at the 6-position and a -S-CH₂-CH₂-phenyl group at the 3-position in the HIV protease area.

J.V.N. Vara Prasad, et al., J. Med. Chem., 38:898-905 (1995), discloses 4-hydroxy-6-phenyl-2-oxo-2H-pyran-3-yl)thiomethanes, such as (+)-3-[cyclopentyl(cyclopentylthio)methyl]-4-hydroxy-6-phenyl-2H-pyran-2-one, as HIV-1 protease inhibitors.

25 SUMMARY OF THE INVENTION

The present invention provides:

A compound of the formula I

wherein R₁ is H-;

wherein R₂ is

- 30 a) C₃-C₅ alkyl,
b) phenyl-(CH₂)₂-,
c) het-SO₂NH-(CH₂)₂-,
d) cyclopropyl-(CH₂)₂-,
e) F-phenyl-(CH₂)₂-,
35 f) het-SO₂NH-phenyl-, or
g) F₃C-(CH₂)₂;

or wherein R_1 and R_2 taken together are a double bond;

wherein R_3 is the moiety of formula X

wherein R_4 is

- a) phenyl,
- 5 b) het,
- c) cyclopropyl,
- d) $H_3C-[O(CH_2)_2]_2-$,
- e) het-SO₂NH-,
- f) Br-,
- 10 g) N₃-, or
- h) $HO_3S(CH_2)_2-N(CH_3)-C(O)-(CH_2)_6-C(O)-NH-$;

wherein R_5 is -H;

wherein R_6 is

- a) $R_4-(CH_2)_n-CH(R_8)-$,
- 15 b) $H_3C-[O(CH_2)_2]_2-CH_2-$,
- c) C₃-C₅ alkyl,
- d) phenyl-(CH₂)₂-,
- e) het-SO₂NH-(CH₂)₂-,
- f) $(HOCH_2)_3C-NH-C(O)-NH-(CH_2)_3-$,
- 20 g) $(HO_2C)(H_2N)CH-(CH_2)_2-C(O)-NH-(CH_2)_3-$,
- h) piperazin-1-yl-C(O)-NH-(CH₂)₃,
- i) $HO_3S(CH_2)_2-N(CH_3)-C(O)-(CH_2)_6-C(O)-NH-(CH_2)_3-$,
- j) cyclopropyl-(CH₂)₂-,
- k) F-phenyl-(CH₂)₂-,
- 25 l) het-SO₂NH-phenyl, or
- m) $F_3C-(CH_2)_2-$;

wherein n is zero (0), one (1) or two (2);

wherein R_7 is

- a) cyclopropyl,
- 30 b) CH_3-CH_2- , or
- c) t-butyl;

wherein R_8 is

- a) -CH₂-CH₃, or
- b) -CH₂-cyclopropyl;

35 wherein R_9 is

- a) -NR₁₂SO₂-het,

- b) $-\text{NR}_{12}\text{SO}_2$ -phenyl substituted by zero (0) or one (1) R_{11} ,
- c) $-\text{CH}_2\text{-SO}_2$ -phenyl substituted by zero (0) or one (1) R_{11} , or
- d) $-\text{CH}_2\text{-SO}_2$ -het;

wherein het is a 5-, 6- or 7-membered saturated or unsaturated ring
 5 containing from one (1) to three (3) heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur; and including any bicyclic group in which any of the above heterocyclic rings is fused to a benzene ring or another heterocycle; substituted by zero (0) or one (1) R_{10} ;

wherein R_{10} is

- 10 a) $-\text{CH}_3$,
- b) $-\text{CN}$,
- c) $-\text{OH}$,
- d) $-\text{C}(\text{O})\text{OC}_2\text{H}_5$,
- e) $-\text{CF}_3$,
- 15 f) $-\text{NH}_2$, or
- g) $-\text{C}(\text{O})\text{-NH}_2$;

wherein R_{11} is

- a) $-\text{CN}$,
- b) $-\text{F}$,
- 20 c) $-\text{OH}$, or
- d) $-\text{NO}_2$;

wherein R_{12} is

- a) $-\text{H}$, or
- b) $-\text{CH}_3$;

25 or a pharmaceutically acceptable salt thereof.

The present invention more particularly provides:

A compound of the formula I

wherein R_1 is H- ;

wherein R_2 is

- 30 a) $\text{C}_3\text{-C}_5$ alkyl,
- b) phenyl- $(\text{CH}_2)_2\text{-}$, or
- c) het- $\text{SO}_2\text{NH-}(\text{CH}_2)_2\text{-}$;

or wherein R_1 and R_2 taken together are a double bond;

wherein R_3 is the moiety of formula X

35 wherein R_4 is

- a) phenyl,

- b) het,
- c) cyclopropyl,
- d) $\text{H}_3\text{C}-[\text{O}(\text{CH}_2)_2]_2-$,
- e) het- $\text{SO}_2\text{NH}-$,
- 5 f) $\text{Br}-$,
- g) N_3- , or
- h) $\text{HO}_3\text{S}(\text{CH}_2)_2-\text{N}(\text{CH}_3)-\text{C}(\text{O})-(\text{CH}_2)_6-\text{C}(\text{O})-\text{NH}-$;

wherein R_5 is $-\text{H}$;

wherein R_6 is

- 10 a) $\text{R}_4-(\text{CH}_2)_n-\text{CH}(\text{R}_8)-$,
- b) $\text{H}_3\text{C}-[\text{O}(\text{CH}_2)_2]_2-\text{CH}_2-$,
- c) C_3-C_5 alkyl,
- d) phenyl- $(\text{CH}_2)_2-$,
- e) het- $\text{SO}_2\text{NH}-(\text{CH}_2)_2-$,
- 15 f) $(\text{HOCH}_2)_3\text{C}-\text{NH}-\text{C}(\text{O})-\text{NH}-(\text{CH}_2)_3-$,
- g) $(\text{HO}_2\text{C})(\text{H}_2\text{N})\text{CH}-(\text{CH}_2)_2-\text{C}(\text{O})-\text{NH}-(\text{CH}_2)_3-$,
- h) piperazin-1-yl- $\text{C}(\text{O})-\text{NH}-(\text{CH}_2)_3$, or
- i) $\text{HO}_3\text{S}(\text{CH}_2)_2-\text{N}(\text{CH}_3)-\text{C}(\text{O})-(\text{CH}_2)_6-\text{C}(\text{O})-\text{NH}-(\text{CH}_2)_3-$;

wherein n is zero (0), one (1) or two (2);

20 wherein R_7 is

- a) cyclopropyl,
- b) CH_3-CH_2- , or
- c) t -butyl;

wherein R_8 is

- 25 a) $-\text{CH}_2-\text{CH}_3$, or
- b) $-\text{CH}_2$ -cyclopropyl;

wherein R_9 is

- a) $-\text{NR}_{12}\text{SO}_2$ -het,
- b) $-\text{NR}_{12}\text{SO}_2$ -phenyl substituted by zero (0) or one (1) R_{11} ,
- 30 c) $-\text{CH}_2-\text{SO}_2$ -phenyl substituted by zero (0) or one (1) R_{11} , or
- d) $-\text{CH}_2-\text{SO}_2$ -het;

wherein het is a 5-, 6- or 7-membered saturated or unsaturated ring containing from one (1) to three (3) heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur; and including any bicyclic group in which any of the above heterocyclic rings is fused to a benzene ring or another heterocycle; substituted by zero (0) or one (1) R_{10} ;

2187523

wherein R_{10} is

- 5
- a) $-\text{CH}_3$,
 - b) $-\text{CN}$,
 - c) $-\text{OH}$, or
 - d) $-\text{C}(\text{O})\text{OC}_2\text{H}_5$;

wherein R_{11} is

- 10
- a) $-\text{CN}$,
 - b) $-\text{F}$,
 - c) $-\text{OH}$, or
 - d) $-\text{NO}_2$;

wherein R_{12} is

- a) $-\text{H}$, or
- b) $-\text{CH}_3$;

or a pharmaceutically acceptable salt thereof.

15 The present invention provides for such compounds wherein het is the following, substituted by zero (0) or one (1) R_{10} ,

- 20
- a) 2-pyridinyl,
 - b) imidazol-2-yl,
 - c) imidazol-4-yl,
 - d) benzimidazol-2-yl,
 - e) quinolin-8-yl,
 - f) quinolin-2-yl,
 - g) pyrimidin-2-yl,
 - h) quinazolin-2-yl,
 - 25 i) purin-6-yl,
 - j) thiazol-2-yl,
 - k) thiazol-4-yl,
 - l) 2-pyrazolyl,
 - m) 2-pyrazinyl,
 - 30 n) tetrahydropyran-4-yl, or
 - o) tetrahydropyran-3-yl.

Also more particularly, the present invention provides for the compound of the formula I

wherein R_1 is H-;

35 wherein R_2 is

- a) $\text{H}_3\text{C}-(\text{CH}_2)_2-$,

2187523

- b) phenyl-(CH₂)₂-
- c) (CH₃)₂CH-CH₂-, or
- d) pentyl;

or wherein R₁ and R₂ taken together are a double bond;

5 wherein R₃ is the moiety of formula X

wherein R₄ is

- a) phenyl,
- b) het,
- c) cyclopropyl,
- 10 d) H₃C-[O(CH₂)₂]₂-,
- e) het-SO₂NH-,
- f) Br-,
- g) N₃-, or
- h) HO₃S(CH₂)₂-N(CH₃)-C(O)-(CH₂)₆-C(O)-NH-;

15 wherein R₅ is -H;

wherein R₆ is

- a) R₄-(CH₂)_n-CH(R₈)-,
- b) H₃C-[O(CH₂)₂]₂-CH₂-,
- c) H₃C-(CH₂)₂-,
- 20 d) phenyl-(CH₂)₂-,
- e) (CH₃)₂CH-CH₂-, or
- f) pentyl;

wherein n is zero (0), one (1) or two (2);

wherein R₇ is

- 25 a) cyclopropyl, or
- b) CH₃-CH₂-;

wherein R₈ is

- a) -CH₂-CH₃, or
- b) -CH₂-cyclopropyl;

30 wherein R₉ is

- a) -NHSO₂-het, or
- b) -NHSO₂-phenyl substituted by zero (0) or one (1) R₁₁;

wherein het is the following, substituted by zero (0) or one (1) R₁₀,

- a) 2-pyridinyl,
- 35 b) imidazol-2-yl,
- c) imidazol-4-yl,

- d) quinolin-8-yl,
- e) tetrahydropyran-4-yl,
- f) tetrahydropyran-3-yl, or
- g) benzimidazol-2-yl;

5 wherein R_{10} is

- a) $-\text{CH}_3$;

wherein R_{11} is

- a) $-\text{CN}$,
- b) $-\text{F}$ or
- 10 c) $-\text{NO}_2$;

or a pharmaceutically acceptable salt thereof.

Most particularly, the present invention provides for the compound of the formula VI

wherein R_2 is

- 15 a) $\text{H}_3\text{C}-(\text{CH}_2)_2-$,
- b) phenyl- $(\text{CH}_2)_2-$,
- c) $(\text{CH}_3)_2\text{CH}-\text{CH}_2-$, or
- d) pentyl;

wherein R_3 is the moiety of formula X

20 wherein R_6 is

- a) $\text{H}_3\text{C}-(\text{CH}_2)_2-$,
- b) phenyl- $(\text{CH}_2)_2-$,
- c) $(\text{CH}_3)_2\text{CH}-\text{CH}_2-$, or
- d) pentyl;

25 wherein R_7 is

- a) CH_3-CH_2- , or
- b) cyclopropyl;

wherein R_9 is

- a) $-\text{NHSO}_2$ -phenyl substituted by one (1) R_{11} , or
- 30 b) $-\text{NHSO}_2$ -het;

wherein het is the following, substituted by zero (0) or one (1) R_{10} ,

- a) imidazol-4-yl, or
- b) quinolin-8-yl;

wherein R_{10} is $-\text{CH}_3$;

35 wherein R_{11} is

- a) $-\text{CN}$, or

b) -F.

Also, most particularly, the present invention provides for the compound of the formula VII

wherein R_3 is the moiety of formula X

5 wherein R_4 is

- a) phenyl,
- b) het,
- c) cyclopropyl,
- d) $H_3C-[O(CH_2)_2]_2-$,
- 10 e) het- SO_2NH- ,
- f) Br-,
- g) N_3- , or
- h) $HO_3S(CH_2)_2-N(CH_3)-C(O)-(CH_2)_6-C(O)-NH-$;

wherein R_6 is

- 15 a) $R_4-(CH_2)_n-CH(R_8)-$, or
- b) $H_3C-[O(CH_2)_2]_2-CH_2-$;

wherein R_7 is cyclopropyl;

wherein R_8 is

- a) $-CH_2-CH_3$, or
- 20 b) $-CH_2-$ cyclopropyl;

wherein R_9 is

- a) $-NHSO_2$ -het, or
- b) $-NHSO_2$ -phenyl substituted by one (1) R_{11} ;

wherein n is zero (0), one (1) or two (2);

25 wherein het is the following, substituted by zero (0) or one (1) R_{10} ,

- a) imidazol-4-yl,
- b) imidazol-2-yl,
- c) quinolin-8-yl,
- d) tetrahydropyran-3-yl,
- 30 e) tetrahydropyran-4-yl,
- f) 2-pyridinyl, or
- g) benzimidazol-2-yl;

wherein R_{10} is $-CH_3$;

wherein R_{11} is

- 35 a) $-NO_2$,
- b) -F, or

c) -CN;

or a pharmaceutically acceptable salt thereof.

The present invention also provides:

A compound of the formula II

5 wherein R_{10} and R_{20} taken together are

a) the moiety of formula III, or

b) the moiety of formula IV;

wherein p is four (4);

wherein R_1 is -H;

10 wherein R_2 is

a) H-,

b) $\text{CH}_3\text{O}-$, or

c) $\text{CH}_3\text{O}-[(\text{CH}_2)_2\text{O}]_3-$;

wherein R_3 is the moiety of formula V

15 wherein R_4 is

a) cyclopropyl, or

b) $-\text{CH}_2-\text{CH}(\text{CH}_3)_2$;

wherein R_5 is

a) $-\text{NR}_9\text{SO}_2$ -phenyl substituted by zero (0) or one (1) R_6 ,

20 b) $-\text{NR}_9\text{SO}_2$ -het,

c) $-\text{CH}_2-\text{SO}_2$ -phenyl substituted by zero (0) or one (1) R_6 , or

d) $-\text{CH}_2-\text{SO}_2$ -het;

wherein R_6 is

a) -CN,

25 b) -F,

c) $-\text{CH}_3$,

d) $-\text{COOH}$, or

e) -OH;

wherein het is a 5-, 6- or 7-membered saturated or unsaturated ring

30 containing from one (1) to three (3) heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur; and including any bicyclic group in which any of the above heterocyclic rings is fused to a benzene ring or another heterocycle; substituted by zero (0), one (1) or two (2) R_7 ;

wherein R_7 is

35 a) $-\text{CH}_3$,

b) -CN,

c) $-\text{C}(\text{O})\text{OC}_2\text{H}_5$, or

d) $-\text{OH}$;

wherein R_8 is

a) $-\text{H}$,

5 b) $-(\text{CH}_2)_2-\text{CH}_3$,

c) $-\text{CH}_2$ -cyclopropyl, or

d) $-\text{CH}_2$ -phenyl;

wherein R_9 is

a) $-\text{H}$, or

10 b) $-\text{CH}_3$;

or a pharmaceutically acceptable salt thereof.

The present invention provides for such compounds wherein het is the following, substituted by zero (0) or one (1) R_7 ,

a) 2-pyridinyl,

15 b) imidazol-2-yl,

c) imidazol-4-yl,

d) benzimidazol-2-yl,

e) quinolin-8-yl,

f) quinolin-2-yl,

20 g) pyrimidin-2-yl,

h) quinazolin-2-yl,

i) purin-6-yl,

j) thiazol-2-yl,

k) thiazol-4-yl,

25 l) 2-pyrazolyl,

m) 2-pyrazinyl,

n) tetrahydropyran-4-yl, or

o) tetrahydropyran-3-yl.

30 More particularly, the present invention provides for the compound of the formula II

wherein R_{10} and R_{20} taken together are

a) the moiety of formula III, or

b) the moiety of formula IV;

wherein p is four (4);

35 wherein R_1 is $-\text{H}$;

wherein R_2 is

- a) $\text{CH}_3\text{O}-$, or
- b) $\text{CH}_3\text{O}[(\text{CH}_2)_2\text{O}]_3-$;

wherein R_3 is the moiety of formula V

wherein R_4 is

- 5
- a) cyclopropyl, or
 - b) $-\text{CH}_2-\text{CH}(\text{CH}_3)_2$;

wherein R_5 is

- 10
- a) $-\text{NR}_9\text{SO}_2$ -phenyl substituted by zero (0) or one (1) R_6 ,
 - b) $-\text{NR}_9\text{SO}_2$ -het,
 - c) $-\text{CH}_2-\text{SO}_2$ -phenyl substituted by zero (0) or one (1) R_6 , or
 - d) $-\text{CH}_2-\text{SO}_2$ -het;

wherein R_6 is

- 15
- a) $-\text{CN}$,
 - b) $-\text{F}$,
 - c) $-\text{CH}_3$, or
 - d) $-\text{COOH}$;

wherein het is the following, substituted by zero (0) or one (1) R_7 ,

- 20
- a) imidazol-4-yl,
 - b) quinolin-8-yl,
 - c) 2-pyridinyl, or
 - d) 4-pyridinyl;

wherein R_7 is $-\text{CH}_3$;

wherein R_8 is

- 25
- a) $-\text{H}$, or
 - b) $-(\text{CH}_2)_2-\text{CH}_3$;

wherein R_9 is

- a) $-\text{H}$, or
- b) $-\text{CH}_3$;

or a pharmaceutically acceptable salt thereof.

30 Most particularly, the present invention provides for the compound of the formula VIII

wherein R_3 is the moiety of formula V

wherein R_4 is

- 35
- a) cyclopropyl,
 - b) $-\text{CH}_2-\text{CH}(\text{CH}_3)_2$;

wherein R_5 is

- a) $-\text{NR}_9\text{SO}_2$ -phenyl substituted by zero (0) or one (1) R_6 ,
- b) $-\text{NR}_9\text{SO}_2$ -het, or
- c) $-\text{CH}_2\text{-SO}_2$ -phenyl;

wherein R_6 is

- a) $-\text{CN}$, or
- b) $-\text{F}$;

wherein het is the following, substituted by zero (0) or one (1) R_7 ,

- a) 2-pyridinyl,
- b) 4-pyridinyl, or
- c) imidazol-4-yl;

wherein R_7 is $-\text{CH}_3$;

wherein R_8 is

- a) $-\text{H}$, or
- b) $-(\text{CH}_2)_2\text{-CH}_3$;

wherein R_9 is

- a) $-\text{H}$, or
- b) $-\text{CH}_3$;

or a pharmaceutically acceptable salt thereof.

Also, most particularly, the present invention provides for the compound of the formula IX

wherein R_1 is H :-

wherein R_2 is

- a) $\text{CH}_3\text{O-}$, or
- b) $\text{CH}_3\text{O-}[(\text{CH}_2)_2\text{O}]_3\text{-}$;

wherein R_3 is the moiety of formula V

wherein R_4 is cyclopropyl;

wherein R_5 is $-\text{NHSO}_2$ -het;

wherein het is the following, substituted by zero (0) or one (1) R_7 ,

- a) imidazol-4-yl,
- b) 2-pyridinyl, or
- c) quinolin-8-yl;

wherein R_7 is $-\text{CH}_3$.

The present invention also provides for the compound of the formula VI

wherein R_2 is

- a) $\text{H}_3\text{C-CH}_2\text{-}$,
- b) $\text{H}_3\text{C-}(\text{CH}_2)_2\text{-}$,

- 5
- c) cyclopropyl-(CH₂)₂-,
 - d) F-phenyl-(CH₂)₂-,
 - e) het-SO₂NH-phenyl-,
 - f) (H₃C)₂HC-CH₂,
 - g) phenyl-(CH₂)₂-, or
 - h) F₃C-(CH₂)₂-;

wherein R₃ is the moiety of formula X

wherein R₆ is

- 10
- a) H₃C-CH₂-,
 - b) H₃C-(CH₂)₂-,
 - c) cyclopropyl-(CH₂)₂-,
 - d) F-phenyl-(CH₂)₂-,
 - e) het-SO₂NH-phenyl,
 - f) (H₃C)₂HC-CH₂,
 - 15 g) phenyl-(CH₂)₂-, or
 - h) F₃C-(CH₂)₂-;

wherein R₇ is

- a) H₃C-CH₂-,
- b) t-butyl, or
- 20 c) cyclopropyl

wherein R₉ is

- a) -NHSO₂-het, or
- b) -NHSO₂-phenyl substituted by one (1) R₁₁;

wherein het is the following, substituted by zero (0) or one (1) R₁₀,

- 25
- a) imidazol-4-yl,
 - b) 2-pyridinyl, or
 - c) quinolin-8-yl;

wherein R₁₀ is,

- 30
- a) -CH₃,
 - b) -CN,
 - c) -CF₃,
 - d) -NH₂, or
 - e) -C(O)-NH₂;

wherein R₁₁ is CN.

35 The present invention also provides:

A compound of the formula XI

wherein R_1 is $-(CH_2)_p-CH(R_2)-(CH_2)_o-Ar_1$;

wherein R_2 is

- a) $-C_1-C_5$ alkyl, or
- b) $-(CH_2)_q$ -cycloalkyl;

5 wherein Ar_1 is

- a) phenyl substituted by zero (0) or one (1) R_3 , or
- b) phenyl substituted by -meta-NHSO₂Ar₂;

wherein Ar_2 is

- a) phenyl substituted by zero (0) or one (1) R_3 , or
- 10 b) het;

wherein het is a 5-, 6- or 7-membered saturated or unsaturated ring containing from one (1) to three (3) heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur; and including any bicyclic group in which any of the above heterocyclic rings is fused to a benzene ring or another heterocycle; substituted by

15 zero (0) or one (1) R_4 ;

wherein R_3 is

- a) -CN,
- b) -F,
- c) -OH, or
- 20 d) -NO₂;

wherein R_4 is

- a) -CH₃,
- b) -CN,
- c) -OH,
- 25 d) -C(O)OC₂H₅,
- e) -CF₃, or
- f) -NH₂;

wherein n is zero (0) to eight (8), inclusive;

wherein o is zero (0) to three (3), inclusive;

30 wherein p is zero (0) to three (3), inclusive;

wherein q is zero (0) to three (3), inclusive; or

a pharmaceutically acceptable salt thereof.

More particularly, the present invention provides:

The compound wherein R_1 is $-CH(R_2)-Ar_1$;

35 wherein R_2 is

- a) $-CH_2-CH_3$, or

b) -t-butyl;

wherein Ar_1 is phenyl substituted by -meta-NHSO₂Ar₂;

wherein Ar_2 is 2-pyridinyl substituted by one (1) R₄;

wherein R₄ is

5 a) -CN, or

b) -CF₃;

wherein n is two (2) to four (4) inclusive.

The present invention also provides:

A process for producing a compound of the formula W-10

10 wherein R₁ is

a) n-propyl, or

b) phenethyl;

which comprises the steps of:

a) treating a compound of the formula W-9

15 wherein X_A is as defined above, with TiCl₄;

b) treating the product of step a) with an amine base; and

c) reacting the product of step b) with 4-heptanone or propylphenethylketone to yield the compound of formula W-10;

The process which further comprises the steps of:

20 d) treating the compound of formula W-10 with sodium hydride or potassium t-butoxide to obtain a compound of formula W-11

wherein R₁ is

a) n-propyl, or

b) phenethyl;

25 e) hydrogenating the compound of formula W-11 to obtain the compound of formula W-12

wherein R₁ is as defined above;

f) treating the compound of formula W-12 with a sulfonyl chloride of formula D-7

30 wherein R₄ is 5-trifluoromethyl-2-pyridinyl, in an organic solvent in the presence of an organic base to obtain a compound of the formula W-13

wherein R₁ is as defined above.

The present invention also provides:

A process for producing a compound of the formula X-10

35 wherein R₁ is

a) n-propyl, or

b) phenethyl;

which comprises the steps of:

a) treating a compound of the formula X-9

wherein X_A is as defined above, with $TiCl_4$;

5 b) treating the product of step a) with an amine base; and

c) reacting the product of step b) with 4-heptanone or propylphenethylketone to yield the compound of formula X-10;

The process which further comprises the steps of:

d) treating the compound of formula X-10 with sodium hydride or

10 potassium t-butoxide to obtain a compound of formula X-11

wherein R_1 is

a) n-propyl, or

b) phenethyl;

e) hydrogenating the compound of formula X-11 to obtain a compound of

15 formula X-12

wherein R_1 is as defined above;

f) treating the compound of formula X-12 with a sulfonyl chloride of formula D-7

wherein R_4 is 5-trifluoromethyl-2-pyridinyl, in an organic solvent in the presence of

20 an organic base to obtain a compound of the formula X-13

wherein R_1 is as defined above.

The present invention also provides:

A process for producing a compound of the formula GGG-10

wherein R_1 is

25 a) n-propyl, or

b) phenethyl;

which comprises the steps of:

a) treating a compound of the formula GGG-9

wherein X_A is as defined above, with $TiCl_4$;

30 b) treating the product of step a) with an amine base; and

c) reacting the product of step b) with 4-heptanone or 1-phenyl-3-hexanone to yield the compound of formula GGG-10.

The process which further comprises the steps of:

d) treating the compound of formula GGG-10 with sodium hydride or

35 potassium t-butoxide to obtain a compound of formula GGG-11

wherein R_1 is

- a) n-propyl, or
- b) phenethyl;
- e) hydrogenating the compound of formula GGG-11 to obtain a compound of formula GGG-12

5 wherein R_1 is as defined above;

- f) treating the compound of formula GGG-12 with a sulfonyl chloride of formula D-7

wherein R_4 is

- a) 5-trifluoromethyl-2-pyridinyl, or
 - 10 b) 5-cyano-2-pyridinyl, in an organic solvent in the presence of an organic base to obtain a compound of the formula GGG-13A
- wherein R_1 is as defined above.

A process for producing a compound of the formula HHH-10
wherein R_1 is

- 15 a) n-propyl, or
- b) phenethyl;

which comprises the steps of:

- a) treating a compound of the formula HHH-9
- wherein X_A is as defined above, with $TiCl_4$;
- 20 b) treating the product of step a) with an amine base; and
 - c) reacting the product of step b) with 4-heptanone or 1-phenyl-3-hexanone to yield the compound of formula HHH-10.

The process which further comprises the steps of:

- d) treating the compound of formula HHH-10 with sodium hydride or
- 25 potassium t-butoxide to obtain a compound of formula HHH-11

wherein R_1 is

- a) n-propyl, or
 - b) phenethyl;
 - e) hydrogenating the compound of formula HHH-11 to obtain a
- 30 compound of formula HHH-12

wherein R_1 is as defined above;

- f) treating the compound of formula HHH-12 with a sulfonyl chloride of formula D-7

wherein R_4 is

- 35 a) 5-trifluoromethyl-2-pyridinyl, or
- b) 5-cyano-2-pyridinyl, in an organic solvent in the presence of an organic

base to obtain a compound of the formula HHH-13A

wherein R_1 is as defined above.

A process for producing a compound of the formula III-10

wherein R_1 is

- 5 a) n-propyl, or
 b) phenethyl;

which comprises the steps of:

- a) treating a compound of the formula III-9

wherein X_A is as defined above, with $TiCl_4$;

- 10 b) treating the product of step a) with an amine base; and
 c) reacting the product of step b) with 4-heptanone or 1-phenyl-3-hexanone to yield the compound of formula III-10.

The process which further comprises the steps of:

- d) treating the compound of formula III-10 with sodium hydride or
15 potassium t-butoxide to obtain a compound of formula III-11

wherein R_1 is

- a) n-propyl, or
 b) phenethyl;
 e) hydrogenating the compound of formula III-11 to obtain a compound of
20 formula III-12.

wherein R_1 is as defined above;

- f) treating the compound of formula III-12 with a sulfonyl chloride of
 formula D-7

wherein R_4 is

- 25 a) 5-trifluoromethyl-2-pyridinyl, or
 b) 5-cyano-2-pyridinyl, in an organic solvent in the presence of an organic
base to obtain a compound of the formula III-13A

wherein R_1 is as defined above.

A process for producing a compound of the formula JJJ-10

30 wherein R_1 is

- a) n-propyl, or
 b) phenethyl;

which comprises the steps of:

- a) treating a compound of the formula JJJ-9

35 wherein X_A is as defined above, with $TiCl_4$;

- b) treating the product of step a) with an amine base; and

c) reacting the product of step b) with 4-heptanone or 1-phenyl-3-hexanone to yield the compound of formula JJJ-10.

The process which further comprises the steps of:

d) treating the compound of formula JJJ-10 with sodium hydride or
5 potassium t-butoxide to obtain a compound of formula JJJ-11

wherein R_1 is

- a) n-propyl, or
- b) phenethyl;
- e) hydrogenating the compound of formula JJJ-11 to obtain a compound
10 of formula JJJ-12

wherein R_1 is as defined above;

f) treating the compound of formula JJJ-12 with a sulfonyl chloride of
formula D-7

wherein R_4 is

- 15 a) 5-trifluoromethyl-2-pyridinyl, or
- b) 5-cyano-2-pyridinyl, in an organic solvent in the presence of an organic
base to obtain a compound of the formula JJJ-13A

wherein R_1 is as defined above.

The present invention most preferably provides:

20 The compound of formula VI wherein R_2 is

- a) $H_3C-(CH_2)_2-$, or
- b) phenyl- $(CH_2)_2-$;

wherein R_3 is the moiety of formula X;

wherein R_6 is

- 25 a) $H_3C-(CH_2)_2-$, or
- b) phenyl- $(CH_2)_2-$;

wherein R_7 is

- a) H_3C-CH_2- , or
- b) t-butyl;

30 wherein R_9 is $-NHSO_2$ -het;

wherein het is the following, substituted by one (1) R_{10} ,

- a) imidazol-4-yl, or
- b) 2-pyridinyl;

wherein R_{10} is,

- 35 a) $-CH_3$,
- b) $-CN$, or

c) $-\text{CF}_3$.

The compounds of the present invention are named according to the IUPAC or CAS nomenclature system.

The carbon atoms content of various hydrocarbon-containing moieties is indicated by a prefix designating the minimum and maximum number of carbon atoms in the moiety, i.e., the prefix $\text{C}_i\text{-C}_j$ indicates a moiety of the integer "i" to the integer "j" carbon atoms, inclusive. Thus, for example, $\text{C}_1\text{-C}_3$ alkyl refers to alkyl of one to three carbon atoms, inclusive, or methyl, ethyl, propyl, and isopropyl, straight and branched forms thereof.

Also, the carbon atom content of various hydrocarbon-containing moieties of the present invention is indicated by a subscripted integer representing the number of carbon and hydrogen atoms in the moiety, e.g., " C_nH_{2n} " indicates a moiety of the integer "n" carbon atoms, inclusive, and the integer "2n" hydrogen atoms, inclusive. Thus, for example, " C_nH_{2n} " wherein n is one to three carbon atoms, inclusive, and two to six hydrogen atoms, inclusive, or methyl, ethyl, propyl and isopropyl, and all isomeric, straight and branched forms thereof.

Examples of alkyl of one to nine carbon atoms, inclusive, are methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, and nonyl, and all isomeric forms thereof and straight and branched forms thereof.

Examples of alkenyl of one to five carbon atoms, inclusive, are ethenyl, propenyl, butenyl, pentenyl, all isomeric forms thereof, and straight and branched forms thereof.

By "halo" is meant the typical halogen atoms, such as fluorine, chlorine, bromine, and iodine.

The compounds of formula I and II of the present invention inhibit retroviral proteinases and thus inhibit the replication of the virus. They are useful for treating patients infected with human immunodeficiency virus (HIV) which results in acquired immunodeficiency syndrome (AIDS) and related diseases.

More particularly, the compounds of the present invention are useful as novel human retroviral protease inhibitors. Therefore, the compounds inhibit retroviral proteases and thus inhibit the replication of the virus. They are useful for treating human patients infected with a human retrovirus, such as human immunodeficiency virus (strains of HIV-1 or HIV-2) or human T-cell leukemia viruses (HTLV-I or HTLV-II) which results in acquired immunodeficiency syndrome (AIDS) and/or related diseases.

The capsid and replicative enzymes (i.e. protease, reverse transcriptase,

integrase) of retroviruses are translated from the viral gag and pol genes as polyproteins that are further processed by the viral protease (PR) to the mature proteins found in the viral capsid and necessary for viral functions and replication. If the PR is absent or nonfunctional, the virus cannot replicate. The retroviral PR, such as HIV-1 PR, has been found to be an aspartic protease with active site characteristics similar to those exhibited by the more complex aspartic protease, renin.

The term human retrovirus (HRV) includes human immunodeficiency virus type I, human immunodeficiency virus type II, or strains thereof, as well as human T cell leukemia virus 1 and 2 (HTLV-1 and HTLV-2) or strains apparent to one skilled in the art, which belong to the same or related viral families and which create similar physiological effects in humans as various human retroviruses.

Patients to be treated would be those individuals: 1) infected with one or more strains of a human retrovirus as determined by the presence of either measurable viral antibody or antigen in the serum and 2) in the case of HIV, having either an asymptomatic HIV infection or a symptomatic AIDS defining infection such as i) disseminated histoplasmosis, ii) isopsoriasis, iii) bronchial and pulmonary candidiasis including pneumocystic pneumonia iv) non-Hodgkin's lymphoma or v) Kaposi's sarcoma and being less than sixty years old; or having an absolute CD4+ lymphocyte count of less than $500/\text{mm}^3$ in the peripheral blood. Treatment would consist of maintaining an inhibitory level of the compound used according to this invention in the patient at all times and would continue until the occurrence of a second symptomatic AIDS defining infection indicates alternate therapy is needed.

More specifically, an example of one such human retrovirus is the human immunodeficiency virus (HIV, also known as HTLV-III or LAV) which has been recognized as the causative agent in human acquired immunodeficiency syndrome (AIDS), P. Duesberg, Proc. Natl. Acad. Sci. USA, 86:755 (1989). HIV contains a retro viral encoded protease, HIV-I protease, that cleaves the fusion polypeptides into the functional proteins of the mature viral particle, E.P. Lillehoj, et al., J. Virology, 62:3053 (1988); C. Debuck, et al., Proc. Natl. Acad. Sci., 84:8903 (1987). This enzyme, HIV-I protease, has been classified as an aspartyl protease and has a demonstrated homology to other aspartyl proteases such as renin, L.H. Pearl, et al., Nature 329:351 (1987); I. Katoh, et al., Nature 329:654 (1987). Inhibition of HIV-I protease blocks the replication of HIV and thus is useful in the treatment of human AIDS, E.D. Clerq, J. Med. Chem. 29:1561 (1986). Inhibitors of HIV-I protease are useful in the treatment of HIV-infected individuals who are asymptomatic or

symptomatic of AIDS.

Pepstatin A, a general inhibitor of aspartyl proteases, has been disclosed as an inhibitor of HIV-I protease, S. Seelmeier, et al., Proc. Natl. Acad. Sci. USA, 85:6612 (1986). Other substrate derived inhibitors containing reduced bond

5 isosteres or statine at the scissile position have also been disclosed, M.L. Moore, et al., Biochem. Biophys. Res. Commun. 159:420 (1989); S. Billich, et al., J. Biol. Chem. 263:17905 (1988); Sandoz, D.E. 3812-576-A.

Thus, the compounds of the present invention are useful for treating diseases caused by retroviruses, such as human acquired immunodeficiency disease syndrome
10 (AIDS).

The compounds are also useful for treating non-human animals infected with a retrovirus, such as cats infected with feline leukemia virus. Other viruses that infect cats include, for example, feline infectious peritonitis virus, calicivirus, rabies virus, feline immunodeficiency virus, feline parvovirus (panleukopenia virus), and
15 feline chlamydia. Exact dosages, forms and modes of administration of the compounds of the present invention to non-human animals would be apparent to one of ordinary skill in the art, such as a veterinarian.

The compounds of formula I and II of the present invention are prepared as described in the Charts, Preparations and Examples below, or are prepared by
20 methods analogous thereto, which are readily known and available to one of ordinary skill in the art of organic synthesis.

CHART A

Nitration of the cyclopropylphenyl ketone of formula A-1, which is commercially available, with fuming nitric acid at -40 °C produces a ca. 2:1 mixture
25 of isomers. The desired *m*-nitro compound of formula A-2 is easily separated from the crude mixture by recrystallization from methanol. Catalytic hydrogenation of the cyclopropyl-(3-nitrophenyl)methanone of formula A-2 with 10% platinum on carbon in methanol gives the aniline of formula A-3. The aniline is then coupled with benzenesulfonyl chloride using pyridine in methylene chloride to give the
30 sulfonamide derivative of formula A-4. Reduction of the ketone with sodium borohydride in tetrahydrofuran and ethanol then produces the carbinol of formula A-5.

The dianion of the cyclooctylpyranone of formula A-6, prepared as described in Chart B, is formed using lithium diisopropyl amide in tetrahydrofuran at 0 °C,
35 and then alkylated with iodopropane to give the 10-propyl-cyclooctylpyranone of formula A-7. The cyclooctylpyranone of formula A-7 and the carbinol of the formula

A-5 are then coupled using *p*-toluenesulfonic acid in methylene chloride to give the sulfonamide derivative of formula A-8.

CHART B

The commercially available amine of the formula B-1 is protected using
5 benzyl chloroformate and sodium bicarbonate in THF/water solution to give the compound of formula B-2. The aldehyde of formula B-2 is then reacted with a Grignard reagent to give the secondary alcohol of formula B-3, wherein, e.g., R₁ is isobutyl. The known cyclooctylpyranone of formula B-4 is prepared by acylation of the trimethylsilyl enol ether of cyclooctanone with malonyl dichloride as described in
10 R. Effenberger, T. Ziegler, K.-H. Schonwalder, T. Kesmarszky, B. Bauer Chem. Ber. 119:3394-3404 (1986). The alcohol of formula B-3 is then used to alkylate the cyclooctylpyranone of formula B-4 in refluxing toluene and *p*-toluenesulfonic acid to obtain the compound of the formula B-5, wherein, e.g., R₁ is isobutyl. At this point, the enantiomers of formula B-5 are separated using a chiral HPLC column. The
15 benzyloxy protecting group is then cleaved using 10% Pd/C in cyclohexene to give the amine of formula B-6, wherein, e.g., R₁ is isobutyl, which is reacted with aryl sulfonyl chlorides to give the compounds of the formula B-7, wherein, e.g., R₁ is isobutyl and R₂ is 1-methylimidazole.

CHART C

20 3-Bromobenzyl alcohol of formula C-1, which is commercially available, in tetrahydrofuran is treated with methyllithium, *n*-butyllithium and cyclopropanecarboxaldehyde in sequence at -78°C. The resulting solution is gradually warmed to room temperature and then heated at reflux affording the alcohol of formula C-2. The resulting alcohol, in dichloromethane, in the presence of molecular
25 sieves, is treated with 4-hydroxy-5,6,7,8,9,10-hexahydrocycloocta[b]pyran-2-one of formula C-8, prepared as described in Chart B, and *p*-toluenesulfonic acid. The solution is heated at reflux to afford the alcohol of formula C-3. The benzyl alcohol is treated with carbon tetrabromide and triphenylphosphine in dichloromethane at 0°C to afford compounds of formula C-4 and C-5 as an inseparable mixture after an
30 aqueous brine workup. The mixture is then treated with any thiol (e.g., thiophenol) and an organic base and heated at reflux to afford sulfides of the formula C-6. Finally treatment of the compounds of the formula C-6 with oxone in a mixture of tetrahydrofuran, methanol and water gives sulfones of formula C-7.

CHART D

35 This chart describes a generic procedure for the preparation of C-3 α branched 5,6-dihydropyrones via aluminum chloride (AlCl₃) mediated condensation with 3-

nitrobenzaldehyde. Thus, the AlCl_3 catalyzed reaction of the compound of formula D-1, prepared as described below in the Preparations, (e.g., wherein R_1 is phenethyl or propyl; R_2 is phenethyl or propyl) with 3-nitrobenzaldehyde (formula D-2), which is commercially available, provides compounds of formula D-3 (e.g., wherein R_1 is phenethyl or propyl; R_2 is phenethyl or propyl). Subsequent reaction with trialkyl aluminums or Grignard reagents in the presence of cuprous bromide-dimethylsulfide complex ($\text{CuBr} \cdot \text{Me}_2\text{S}$) provides compounds of formula D-4 (e.g., wherein R_1 is phenethyl or propyl; R_2 is phenethyl or propyl; R_3 is ethyl or cyclopropyl). Transfer hydrogenation with Pd/C and ammonium formate provides compounds of formula D-5 (e.g., wherein R_1 is phenethyl or propyl; R_2 is phenethyl or propyl; R_3 is ethyl or cyclopropyl). Treatment of the compound of formula D-5 with sulfonyl chlorides of formula D-7, wherein R_4 is defined below, and pyridine in methylene chloride (CH_2Cl_2) provides compounds of formula D-6 (e.g., wherein R_1 is phenethyl or propyl; R_2 is phenethyl or propyl; R_3 is ethyl or cyclopropyl; R_4 is 4-cyanophenyl, 4-fluorophenyl, 1-methylimidazol-4-yl, quinolin-8-yl, 2-pyridyl, 4-cyano-2-pyridyl, quinolin-2-yl, 2-hydroxyphenyl, 2-pyrimidyl, 2-quinazoline, 7H-purin-6-yl, 1H-imidazol-2-yl, 1H-benzimidazol-2-yl or thiazol-2-yl).

CHART E

Treatment of commercially available 4-hydroxy-6-methyl-2-pyrone of formula E-1 with three equivalents of lithium diisopropylamide in tetrahydrofuran and hexamethylphosphoramide is followed by bromomethylcyclopropane to afford the compound of formula E-2. Reaction between the compound of formula E-2 and the compound of formula F-5, prepared as described in Chart F, in benzene with p-toluenesulfonic acid catalyst in the presence of molecular sieves affords the compound of formula E-3. Hydrogenolysis of the compound of formula E-3 in methanol with hydrogen and palladium on charcoal gives the free amine of formula E-4. Treatment of the compound of formula E-4 with two equivalents of pyridine in dichloromethane followed by one equivalent of 4-fluorobenzenesulfonyl chloride gives the compound of formula E-5 (wherein, e.g., R is 4-fluorophenyl) which is the compound: N-(3-{cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-4-fluoro-benzenesulfonamide.

Under similar conditions, compounds of general formula E-5 are obtained by reacting amine E-4 with alkyl, aryl and heteroaryl sulfonyl chlorides in the presence of pyridine to give compounds of formula E-5 wherein R is alkyl, aryl or heteroaryl. Also, for example, the enantiomers of the compound of formula E-9 are separated chromatographically by chiral HPLC to give compounds of formula E-10 and E-11.

Additional final compounds of the present invention of formula E-6, E-7, E-8, and E-12 - E-16 are prepared using similar conditions.

CHART F

Nitration of commercially available cyclopropyl phenyl ketone of formula F-1
5 with fuming nitric acid affords the compound of formula F-2. Reduction of the compound of formula F-2 in methanol with hydrogen catalyzed by platinum on carbon gives the amine of formula F-3. The compound of formula F-3 is treated with benzylchloroformate and diisopropylethylamine in dichloromethane to give the compound of formula F-4. Reduction of the compound of formula F-4 with sodium
10 borohydride in tetrahydrofuran and ethanol gives the compound of formula F-5.

CHART G

The dianion of commercially available 4-hydroxy-6-methyl-2-pyrone of formula G-0 is generated by deprotonation with two equivalents of lithium diisopropylamide in tetrahydrofuran and hexamethylphosphoramide. Alkylation
15 with 2-(2-methoxy-ethoxy)-ethyl iodide, which is prepared from the commercially available alcohol by standard procedures, gives the compound of formula G-1. Reaction between the compound of formula G-1 and *meta*-benzyloxycarbonylamino-phenyl cyclopropyl carbinol, the compound of formula F-5, prepared as described in Chart F, in dichloromethane with *p*-toluenesulfonic acid catalyst in the presence of
20 molecular sieves gives the compound of formula G-2. Hydrogenolysis of the compound of formula G-2 in ethanol with hydrogen and palladium on charcoal gives the free amine of formula G-3. Treatment of the free amine of formula G-3 with two equivalents of pyridine in dichloromethane followed by one equivalent of 1-methylimidazole-4-sulfonyl chloride gives the compound of formula G-4, which is the
25 compound: N-(3-{cyclopropyl-[4-hydroxy-6-(3-{2-methoxy-ethoxy}-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide.

CHART H

Reaction between commercially available 4-hydroxy-6-methyl-2-pyrone of formula H-0 and *meta*-benzyloxycarbonylaminophenyl cyclopropyl carbinol, the title
30 compound of formula F-5, prepared as described in Chart F, in dichloromethane with *p*-toluenesulfonic acid catalyst in the presence of molecular sieves gives the compound of formula H-1. Alkylation of trianion of the compound of formula H-1 generated from three equivalents of lithium diisopropylamide in tetrahydrofuran with ethyl bromide affords the compound of formula H-2. Treatment of the
35 compound of formula H-2 with lithium diisopropylamide in tetrahydrofuran and 2-(2-methoxy-ethoxy)-ethyl iodide gives the compound of formula H-3. Hydrogenolysis

of the compound of formula H-3 in ethanol with hydrogen and palladium on charcoal gives the free amine of formula H-4. Treatment of the free amine of formula H-4 with two equivalents of pyridine in dichloromethane followed by one equivalent of 1-methylimidazole-4-sulfonyl chloride gives the compound of formula H-5, which is the compound: N-(3-{cyclopropyl-[6-(1-ethyl-3-{2-methoxy-ethoxy}-propyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide. Under similar conditions, compounds of the present invention are obtained by reacting the amine of formula H-4 with alkyl, aryl and heteroaryl sulfonyl chlorides in the presence of pyridine to give additional sulfonamides of formula H-5.

10

CHART I

Treatment of the compound of formula H-2, prepared as described in Chart H, with three equivalents of lithium diisopropylamide in tetrahydrofuran and ethylene oxide gives the compound of formula I-1. Reaction of the compound of formula I-1 with triphenylphosphine and carbon tetrabromide in tetrahydrofuran gives the compound of formula I-2. Treatment of the compound of formula I-2 with sodium azide in aqueous ethanol gives the compound of formula I-3. Reaction of the compound of formula I-3 with hydrogen and palladium on charcoal in ethanol gives the compound of formula I-4. Treatment of the compound of formula I-4 with diisopropylethylamine in dichloromethane followed by 1-methylimidazole-4-sulfonyl chloride gives the compound of formula I-5. Reaction of the compound of formula I-5 with ammonia in methanol gives the compound of formula I-6, which is the compound: N-(3-{cyclopropyl-[6-(1-ethyl-3-{1-methyl-1H-imidazole-4-sulfonylamino}-propyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide.

25

CHART J

Hydrogenolysis of the compound of formula I-1, prepared as described in Chart I, in ethanol with hydrogen and palladium on charcoal gives the compound of formula J-1. Treatment of the compound of formula J-1 with triphenylphosphine and carbon tetrabromide in tetrahydrofuran gives the compound of formula J-2. Reaction of the compound of formula J-2 with pyridine in dichloromethane followed by 1-methylimidazole-4-sulfonyl chloride gives the compound of formula J-3, which is the compound: N-(3-{[6-(3-bromo-1-ethyl-propyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-cyclopropyl-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide. Treatment of the compound of formula J-3 with sodium azide in aqueous ethanol gives the compound of formula J-4, which is the compound: N-(3-{[6-(3-azido-1-ethyl-propyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-cyclopropyl-methyl}-phenyl)-1-methyl-1H-imidazole-4-

35

2187523

- sulfonamide. Reaction of the compound of formula J-4 with hydrogen and palladium on charcoal in ethanol gives the compound of formula J-5. Treatment of the compound of formula J-5 with the triethylamine salt of suleptanic acid (Anderson, B.D.; Conradi, R.A; Knuth, K.E.; J. Pharm. Sci. 74:365 (1985)) and 1,3-
- 5 diisopropylcarbodiimide gives the compound of formula J-6, which is the compound: N-(3-{cyclopropyl-[6-(1-ethyl-3-{N-[8-(methyl-{2-sulfoethyl}-amino)-1,8-dioxooctyl]-amino-propyl)]-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide, sodium salt.

CHART K

- 10 The preparation of the compound of formula K-8, which is the compound: N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide is shown in Chart K. Reduction of commercially available tetrahydropyran-4-carboxylic acid of formula K-1 with borane in tetrahydrofuran provides the compound of formula K-2. The
- 15 compound of formula K-2 is treated with *p*-toluenesulfonyl chloride to afford the corresponding tosylate of formula K-3, which is converted to the iodide of formula K-4 by treatment with potassium iodide in refluxing acetone. Alkylation of the dianion of commercially available 4-hydroxy-6-methyl-2-pyrone of formula K-10 with ethyl bromide in tetrahydrofuran and hexamethylphosphoric triamide gives the propyl
- 20 derivative of formula K-9. The compound of formula K-4 is used to alkylate the compound of formula K-9 at the 6 α position, giving the compound of formula K-5. The compound of formula K-5 is further alkylated at the 3 position, using carbinol of formula F-5, prepared as described in Chart F, giving the compound of formula K-6. Removal of the benzyloxycarbonyl protecting group is accomplished using catalytic
- 25 transfer hydrogenation, giving the amine of formula K-7. Treatment of the amine of formula K-7 with 1-methylimidazole-4-sulfonyl chloride in the presence of pyridine provides the compound of formula K-8.

CHART L

- As shown in Chart L, the dianion of commercially available 4-hydroxy-6-
- 30 methyl-2-pyrone of formula L-1 is generated by deprotonation with two equivalents of lithium diisopropylamide in tetrahydrofuran and hexamethylphosphoramide. Alkylation with benzyl bromide gives the compound of formula L-2, which is then treated with two equivalents of lithium diisopropylamide in tetrahydrofuran and hexamethylphosphoramide, followed by ethyl iodide to give the compound of formula
- 35 L-3. Reaction between the compound of formula L-2 and the compound of formula F-5, prepared as described in Chart F, in benzene with *p*-toluenesulfonic acid

catalyst in the presence of molecular sieves affords the compound of formula L-4, which is 3-[(3-benzyloxycarbonylaminophenyl)-cyclopropyl-methyl]-6-(1-ethylphenethyl)-4-hydroxy-2H-pyran-2-one. Hydrogenolysis of the compound of formula L-4 in methanol using catalytic palladium on charcoal and ammonium formate or hydrogen gas gives the free amine of formula L-5, which is 3-[(3-aminophenyl)-cyclopropyl-methyl]-6-(1-ethylphenethyl)-4-hydroxy-2H-pyran-2-one. Reacting the compound of formula L-5 and the appropriate sulfonyl chloride gives the final compounds of the present invention.

CHART M

As shown in Chart M, commercially available triethylene glycol monomethyl ether is treated with *p*-toluenesulfonyl chloride and pyridine to provide the tosylate of formula M-2, which is then used to alkylate commercially available 2,4-dihydroxyacetophenone to give the compound of formula M-3. Condensation with diethyl carbonate yields the compound of formula M-4. Ring closure of the compound of formula M-4 to the compound of formula M-5 is accomplished by refluxing in acetic acid. The compound of formula M-5 is alkylated at the 3-position using the carbinol of formula F-5, prepared as described in Chart F, and catalytic *p*-toluenesulfonic acid to give the compound of formula M-6. Removal of the benzyloxycarbonyl protecting group is accomplished using catalytic transfer hydrogenation, giving the amine of formula M-7. Treatment of the amine with 1-methylimidazole-4-sulfonyl chloride in the presence of pyridine provides the final compound of formula M-8, which is N-(3-{Cyclopropyl-[7-(2-(2-(2-methoxyethoxy)-ethoxy)ethoxy)-4-hydroxycoumarin-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide.

CHART N

Nitration of cyclopropylphenyl ketone of formula N-1, which is commercially available, with fuming nitric acid at -40 °C produces a ca. 2:1 mixture of isomers. The desired meta-nitro compound of formula N-2 is easily separated from the crude mixture by recrystallization from methanol. Catalytic hydrogenation of cyclopropyl-(3-nitrophenyl)methanone of formula N-2 with 10% platinum on carbon in methanol at 0 °C provides the aniline of formula N-3. The product is isolated by filtration and concentration. The amino group is then protected using benzyl chloroformate and diisopropylethylamine in methylene chloride to give the ketone of formula N-4. The ketone is then reduced with sodium borohydride in 5:1 THF and ethanol to give the alcohol of formula N-5.

The compound of formula N-5 is then used to alkylate 4-hydroxy-5,6,7,8,9,10-

hexahydrocycloocta[b]pyran-2-one, which is prepared as described in R. Effenberger, T. Ziegler, K.-H. Schönzoalder, T. Kesmarsky, B. Bauer, Chem. Ber. 119:3394-3404 (1986), to give the compound of formula N-6. The preferred conditions for this alkylation reaction are *p*-toluene-sulfonic acid in refluxing methylene chloride with a Soxhlet extractor containing molecular sieves. Finally, the compound of formula N-7 is obtained by cleaving the benzyl protective group in a transfer hydrogenation. Best results for this reactions are achieved with 10% Pd/C in neat cyclohexene.

CHART O

Treatment of the amine of formula O-1, prepared as described in Chart N, with sulfonyl chlorides and a base such as pyridine in dichloromethane gives the sulfonamides of formula O-2 wherein R_{60} is, for example, 4-nitrophenyl. These sulfonamides are further modified by standard literature procedures as is apparent to those of ordinary skill in the art to give sulfonamides of formula O-3 wherein R_{61} is, for example, 4-aminophenyl and other functional groups that are not readily available from readily available sulfonyl chlorides. For example, the nitro group of N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-4-nitro-benzenesulfonamide is reduced by catalytic hydrogenation in ethyl acetate with palladium on carbon to give the amine in 4-amino-N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-benzenesulfonamide. Also, the carboxylic acid of 3-[[[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]amino]sulfonyl]-benzoic acid is esterified with methanol and catalytic sulfuric acid to give the methyl ester in 3-[[[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]amino]sulfonyl]-benzoic acid, methyl ester. Sulfonamides of formula O-3 are also obtained from compounds of formula O-2 by further elaboration of reactive functional groups. For example, the amine of 3-amino-N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-benzenesulfonamide is reacted with benzoyl chloride and a base such as pyridine to give the benzamide in N-[3-[[[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]amino]sulfonyl]henyl]-benzamide. Using commonly available sulfonyl chlorides, additional compounds of the present invention of formula II, wherein R_{10} and R_{20} is the moiety of formula IV, are prepared.

The sulfonyl chlorides used to make the compounds of the present invention are readily prepared by methods described in the literature by those skilled in the art, as the following examples illustrate: Reaction of a suitable thiol with KHF_2 in

water/methanol with chlorine gas gives the sulfonyl fluoride (D.J. Brown, J.A. Hoskins, Aust. J. Chem. 25:2641 (1972)) which is then converted into the desired sulfonyl chloride (T. Norris, J. Chem. Soc., Perkin Trans. 1(11):1378 (Eng.) (1978)). Oxidation of a suitable thiol with chlorine in water with ferric chloride (FeCl_3) added gives the desired sulfonyl chloride (G. Pala, Ed. Sci. 13:461 (1958); W.J. Close, J. Amer. Chem. Soc. 82:1132 (1960)). Reaction of the heteroaromatic compound with fuming sulfuric acid gives a heteroaromatic sulfonic acid followed by treatment with phosphorous-orychloride (POCl_3) and phosphorous chloride (PCl_5) gives the desired sulfonyl chloride (V. Georgian, R.J. Harrison, L.L. Skaletzky, J. Org. Chem. 27:4571 (1962)). Reaction of a heteroaromatic compound with manganese dioxide (MnO_2) and sodium sulfite (Na_2SO_3) in water gives the desired sulfonic acid followed by treatment with POCl_3 and PCl_5 gives the desired sulfonyl chloride (N.A. Androva, Izvest. 455 (1972); J.O. Morley, J. Chem. Comm. 88 (1976)). Treatment of the appropriate heteroaromatic chloride with sodium sulfate and HCl in water gives the desired sulfonic acid followed by treatment with POCl_3 and PCl_5 gives the desired sulfonyl chloride (T.R. Norton, J. Amer. Chem. Soc. 68:1330 (1946)). Treatment of the appropriate hydroxy compound with N,N-dimethylthiocarbonyl chloride (M.S. Newman, F.W. Hetzel, Org. Synth. Coll. Vol. IV:824 (1988); M.S. Newman, H.A. Karnes, J. Org. Chem. 31:3980 (1966)) followed by treatment of the resulting thiol, as described above, gives the desired sulfonyl chloride. Treatment of the appropriately protected thio-heteroaromatic compound with chlorine in acetic acid gives the desired sulfonyl chloride (Can. J. Chem. 55:421 (1977)). Using the literature procedures described above, the heteroaromatic sulfonyl chlorides of the present invention are prepared.

25

CHART P

The preferred procedure for the preparation of the heteroaryl sulfonamides of formula P-2 is described in Chart P. Sulfonation of the amine of formula P-1, prepared in Chart N, P-1 with various heteroarylsulfonyl chlorides of formula P-3 wherein R is, e.g., 2-pyridyl, 4-pyridyl, 5-cyanopyridin-2-yl, 2-pyrazinyl, 2-pyrimidinyl, 4,6-dimethylpyrimidin-2-yl, 4-methylpyrimidin-2-yl gives the sulfonamides of formula P-2 wherein R is the corresponding substituent.

30

CHART Q

Generated by sequential deprotonation with sodium hydride and *n*-butyl lithium in tetrahydrofuran at 0 °C, the dianion of commercially available methyl acetoacetate is reacted with ketone of formula Q-1, prepared as described in Chart S (formula S-4). The resulting intermediate hydroxy-ester is cyclized with dilute

35

aqueous hydroxide followed by aqueous hydrochloric acid to give the compound of formula Q-2. The compound of formula Q-2 is condensed with commercially available 3-nitrobenzaldehyde in tetrahydrofuran using aluminum trichloride as a catalyst followed by reaction of the intermediate benzylidene adduct with triethyl aluminum in the presence of copper bromide-dimethyl sulfide to provide the compound of formula Q-3. Catalytic transfer hydrogenation with Pd/C and ammonium formate in methanol affords the compound of formula Q-4. Treatment of the compound of formula Q-4 with the appropriate sulfonyl chloride and pyridine in dichloromethane provides the desired compound of formula Q-5 (wherein, e.g., R₁ is 5-cyano-2-pyridyl or 1-methylimidazol-4-yl).

CHART R

Catalytic hydrogenation of commercially available 3-nitropropiophenone of formula R-1 affords the amine of formula R-2. The amine of formula R-2 is treated with diisopropylethylamine and benzyl bromide to give the compound of formula R-3. The dianion of methyl acetoacetate, generated by treatment of commercially available methyl acetoacetate with sodium hydride and *n*-butyl lithium in tetrahydrofuran at 0 °C, is reacted with the ketone of formula R-3. The intermediate hydroxy-ester is cyclized with dilute aqueous hydroxide followed by aqueous hydrochloric acid to give the compound of formula R-4. The compound of formula R-4 is condensed with 3-nitrobenzaldehyde in tetrahydrofuran using aluminum trichloride as a catalyst followed by reaction of the intermediate benzylidene adduct with triethyl aluminum in the presence of copper bromide-dimethyl sulfide to provide the compound of formula R-5. Catalytic hydrogenation with Pd/C affords the diamine of formula R-6. Treatment of the compound of formula R-6 with the appropriate sulfonyl chloride and pyridine in dichloromethane provides the desired compound of formula R-7 (wherein, e.g., R₁ is 5-cyano-2-pyridyl or 1-methylimidazol-4-yl).

CHART S

Commercially available 4-pentenoic acid of formula S-1 is coupled with *N,O*-dimethylhydroxylamine using bis(2-oxo-3-oxazolidinyl)phosphinic chloride to afford the amide of formula S-2. The amide of formula S-2 is reacted with 3-butenyl magnesium bromide in tetrahydrofuran to give the ketone of formula S-3. The ketone of formula S-3 is treated with zinc metal, cuprous chloride and diiodomethane to provide the ketone of formula S-4 (also formula Q-1, see Chart Q above).

CHART T

The compound of formula T-2 (also formula D-1) (whose preparation is specifically described in Chart D and Preparation 17 above from commercially available methyl acetoacetate and 1-phenyl-3-hexanone (formula T-1)) is condensed with 3-nitrobenzaldehyde in tetrahydrofuran using aluminum trichloride as a catalyst followed by reaction of the intermediate benzylidene adduct with *t*-butylCu(CN)ZnI, (the organometallic reagent derived from zinc metal, 2-iodo-2-methyl-propane, copper cyanide and lithium chloride) to provide the compound of formula T-3. (The preparation of the organometallic reagent is further described in the text corresponding to Preparation J above). Catalytic transfer hydrogenation with Pd/C and ammonium formate in methanol affords the compound of formula T-4. Treatment of the compound of formula T-4 with the appropriate sulfonyl chloride and pyridine in dichloromethane provides the desired compound of formula T-5 (wherein, e.g., R₁ is 5-cyano-2-pyridyl or 1-methylimidazol-4-yl).

CHART U

Commercially available 4-fluorohydrocinnamic acid of formula U-1 is coupled with *N,O*-dimethylhydroxylamine using diethyl cyanophosphonate to provide the amide of formula U-2. Treatment of the amide with *n*-propylmagnesium chloride yields the ketone of formula U-3. Condensation of the ketone with the dianion of methyl acetoacetate, followed by hydrolysis of the intermediate ester and ring closure, provides the dihydropyrone of formula U-4. Reaction of the dihydropyrone with the aldehyde of formula B-2, prepared as described in Chart B above, in the presence of AlCl₃ provides the benzylidene compound of formula U-5; subsequent reaction with Grignard reagents or trialkyl aluminums in the presence of cuprous bromide-dimethyl sulfide complex affords compounds of formula U-6 (wherein, e.g., R₁ is ethyl, *tert*-butyl, or cyclopropyl). Removal of the benzyloxy-carbonyl (CBZ) protecting group is accomplished using ammonium formate and palladium on charcoal to give the amines of formula U-7 (wherein, e.g., R₁ is ethyl, *tert*-butyl, or cyclopropyl). Treatment of the amines with sulfonyl chlorides and pyridine in methylene chloride provides the sulfonamides of formula U-8 (wherein, e.g., R₁ is ethyl, *tert*-butyl, or cyclopropyl and R₂ is alkyl, aryl, or heteroaryl).

CHART V

Commercially available 4-fluorobenzaldehyde of formula V-1 is condensed with acetone, under basic conditions, to provide 1,5-Bis-(4-fluorophenyl)-penta-1,4-dien-3-one of formula V-2. The dienone is reduced with magnesium in methanol to provide the ketone of formula V-3. The ketone of formula V-3 is converted to dihydropyrone products of formula V-8 using chemistry analogous to that described

in Chart U for the sequence of reactions from U-3 to U-8.

CHART W

Commerically available *trans* 2-pentenoic acid of formual W-1 is converted to the corresponding acid chloride using oxalyl chloride in methylene chloride to afford
5 the product of formula W-2. The lithium amide of formula W-3, readily available from the treatment of commerically available (S)-(+)-4-phenyl-2-oxazolidinone with *n*-butyl lithium in tetrahydrofuran at -78°C, is treated with the acid chloride of formula W-2, to give the unsaturated amide of formula W-4. Addition of the amide of formula W-4 to a tetrahydrofuran solution containing commerically available
10 CuBr/(CH₃)₂S and 3-[bis(trimethylsilyl)amino]phenylmagnesium chloride at -20 °C affords the compound of formula W-5 upon acid workup (Hruby et al., J. Org. Chem., 58(26):7567 (1993)). Treatment of the aniline of formula W-5 with benzyl bromide and sodium carbonate in a water/methylene chloride mixture at reflux; or, potassium carbonate in refluxing acetonitrile, affords the compound of formula W-6. Treatment
15 of the amide of formula W-6 with TiCl₄ followed by an amine base in a solvent such as methylene chloride at below -20°C, preferably at -78 °C, then addition of the 2-methoxy-2-methyl-1,3-dioxoline of formula W-7 (prepared as described in Santry et al., J. Am. Chem. Soc., 110(9):2910 (1988)) affords the compound of formula W-8. Brief treatment of the compound of formula W-8 with a protic acid affords the β-
20 ketoamide of formula W-9. Further treatment of the compound of formula W-9 with TiCl₄ followed by an amine base, then 4-heptanone or propylphenethylketone, affords the compound of formula W-10 wherein R₁ is *n*-propyl or phenethyl, respectively. Treatment of the compound of formula W-10 with sodium hydride or preferably potassium *t*-butoxide, in an ether solvent then affords the pyrone of
25 formula W-11. Hydrogenation of the compound of formula W-11 using, e.g., a Pd on carbon as the catalyst, affords the compound of formula W-12. Finally, treatment of the compound of formula W-12 with a sulfonyl chloride of formula D-7, wherein R₄ is 5-trifluoromethyl-2-pyridinyl, in an organic solvent, such as methylene chloride, in the presence of an organic base, such as pyridine, provides the final compound of
30 formula W-13, wherein R₁ is *n*-propyl or phenethyl (when R₁ is phenethyl, it is a pair of diastereomers).

CHART X

The final (R) enantiomer of formula X-13, wherein R₁ is *n*-propyl or phenethyl, is prepared according to the procedures of Chart W.

35

CHART Y

Acetyl chloride of formula Y-1 is added to the lithium amide of formula Y-2

(also X-3), readily available from the treatment of commercially available (R)-(-)-4-phenyl-2-oxazolidinone with n-butyl lithium in tetrahydrofuran at -78°C, to afford the product of formula Y-3. The compound of formula Y-3 is treated first with TiCl_4 in methylene chloride below room temperature, followed by the addition of a tertiary amine base with subsequent addition of the aldehyde of formula Y-4 (aldehyde of the formula Y-4 is readily available from the reaction of commercially available 3-aminobenzaldehyde with benzyl bromide and potassium or sodium carbonate in either acetonitrile or a water/methylene chloride mixture) to yield the compound of formula Y-5. Addition of the amide of formula Y-5 to a tetrahydrofuran solution containing commercially available $\text{CuBr}/(\text{CH}_3)_2\text{S}$ and ethylmagnesium chloride at -20°C affords the compound of formula Y-6. Alternatively, the commercially available compound of formula Y-7 is treated with oxalyl chloride to afford the compound of formula Y-8. The compound of formula Y-8 is then added to a THF solution of the compound of formula Y-2 (also X-3), readily available from the treatment of commercially available (R)-(-)-4-phenyl-2-oxazolidinone with n-butyl lithium in tetrahydrofuran at -78°C, to yield the compound of formula Y-9. Reduction of the compound of formula Y-9 with iron metal in an alcohol/water mixture then affords the compound of formula Y-10. Treatment of the compound of formula Y-10 with benzyl bromide and potassium or sodium carbonate in either acetonitrile or methylene chloride/water then affords the compound of formula Y-5 which, as described above, is converted to the compound of formula Y-6. The compound of formula Y-6 is converted to final product as described for the conversion of the compound of the formula W-6 to the compound of the formula W-13 (wherein R_1 is propyl or phenethyl) in Chart W.

25

CHART Z

Preparation of the (3S) amide of formula Z-6 is accomplished in the same manner as outlined in Chart Y above, except using the compound of formula Z-2 (also W-3). The compound of the formula Z-6 is converted to final product as described for the conversion of the compound of formula X-6 to the compound of the formula X-13 (wherein R_1 is propyl or phenethyl) in Chart Z.

30

CHART AA

Preparation of the 3(S), 6(S) Diastereomers AA-12 and AA-14: Addition of the unsaturated amide of formula AA-1 (also Y-5) to a tetrahydrofuran solution containing commercially available $\text{CuBr}/(\text{CH}_3)_2\text{S}$ and ethylmagnesium chloride at -20 °C affords the compound of formula AA-2 (same as Y-6). Reduction of the compound of formula AA-2 with a metal hydride (sodium borohydride, lithium

35

aluminum hydride) affords the compound of formula AA-3. Oxidation of the compound of formula AA-3 (Swern oxidation) affords the aldehyde of formula AA-4 which is treated with trimethylsilylcyanide to yield the trimethylsilyl protected cyanohydrin of formula AA-5. Alternatively, the compound of formula AA-2 is
5 treated with trimethyl aluminum followed by N-methyl-O-methyl hydroxyl amine to yield the amide of formula AA-6 which is treated with lithium aluminum hydride to yield the aldehyde of formula AA-4. The trimethylsilyl cyanohydrin of formula AA-5 is reacted with a strong base (e.g. n-butyl lithium) followed by the addition of chiral epoxide of formula AA-7 (also BB-12; the synthesis of which is described in Chart
10 BB) to yield the compound of formula AA-8. The compound of formula AA-8 is dissolved in methylene chloride and cooled to -78°C and TiCl_4 is added followed by a tertiary amine base. To that solution is added trimethylorthoformate followed by additional TiCl_4 which yields the compound of formula AA-9. Treatment of the compound of formula AA-9 with base followed by trimethylsilyl chloride, then
15 treatment with an oxidizing agent (ozone), followed by treatment with tetrabutyl ammonium fluoride and then either potassium tert. butoxide or sodium hydride in an ether solvent, then affords the compound of formula AA-10. Hydrogenation of the compound of formula AA-10 then affords the compound of formula AA-11. Finally, treatment of the compound of formula AA-11 with a sulfonyl chloride of formula D-7
20 in Chart D, wherein R_4 is, e.g., 5-trifluoromethyl-2-pyridinyl, in an organic solvent, such as methylene chloride, in the presence of an organic base, such as pyridine, provides the final compound of formula AA-12.

Furthermore, addition of the compound of formula AA-1 to a tetrahydrofuran solution containing commercially available $\text{CuBr}/(\text{CH}_3)_2\text{S}$ and tertiary
25 butylmagnesium chloride at -20°C affords the compound of formula AA-13. The compound of formula AA-13 is converted to the final product, the compound of formula AA-14, using the chemistry described for the synthesis of AA-12.

CHART BB

Chart BB describes the asymmetric synthesis of epoxides of formula BB-7
30 and BB-12. Alkylation of 2-methyl-2-propen-1-ol (BB-1) with commercially available benzyl bromide provides the allylic alcohol of formula BB-2 (see Lipshutz, B.H. et al.; Synthesis 1992, 191). Catalytic Sharpless epoxidation using commercially available (+) diethyl L-tartrate provides the epoxy alcohol of formula BB-8 (see: (a) Pfenniger, A.; Synthesis 1986, 89. (b) Johnson, R.A.; Sharpless, K.B. In Catalytic
35 Asymmetric Synthesis; Ojima, I., Ed.; VCH: New York, 1993; Chapter 4.1, 103.). Alkylation of the compound of formula BB-8 with benzyl bromide (see: Lipshutz,

B.H. et al.; Synthesis 1992, 191) gives the compound of formula BB-9. Reaction of the compound of formula BB-9 with commercially available ethylmagnesium bromide affords the tertiary alcohol of formula BB-10 (see: Hanson, R.M. Chem. Rev. 1991, 91, 437). Catalytic hydrogenolysis of the compound of formula BB-10 provides the diol of formula BB-11. The compound of formula BB-11 is converted to the chiral epoxide of formula BB-12 by standard methodology (for a discussion of the conversion of vicinal diols to epoxides see: Mitsunobu, O. In Comprehensive Organic Synthesis; Trost, B.M. Ed.; Pergamon Press: Oxford, 1991; Vol. 6; Chapter 1.1, 1).

In an analogous manner, the epoxide of formula BB-7 is ultimately derived from the epoxy alcohol of formula BB-3, which in turn is prepared by Sharpless epoxidation of allylic alcohol BB-2 using commercially available (-) diethyl D-tartrate.

Alternatively, reaction of the epoxy alcohol of formula BB-8 with commercially available 4-toluenesulfonyl chloride under standard conditions affords the tosylate of formula BB-13. Reaction of the compound of the formula BB-13 with ethylmagnesium bromide under conditions similar to those described for the nucleophilic opening of arenesulfonate derivatives of glycidol (see: Klunder, J.M; Onami, T.; Sharpless, K.B. J. Org. Chem. 1989, 54, 1295) affords a mixture of the desired epoxide of formula BB-12 and hydroxytosylate of formula BB-14. The hydroxytosylate of formula BB-14 is readily converted to epoxide BB-12 by the action of K_2CO_3 in methanol.

CHART CC

Preparation of the 3(S), 6(R) Diastereomers CC-12 and CC-14: These diastereomers are prepared in a manner identical to that described in Chart AA with the exception that the epoxide of formula CC-7 (same as BB-7) is used.

CHART DD

Preparation of the 3(R), 6(S) Diastereomers DD-12 and DD14: These diastereomers are prepared in a manner identical to that described in Chart AA with the exception that the amide of formula DD-1 (same as Z-5) is used.

CHART EE

Preparation of the 3(R), 6(R) Diastereomers EE-12 and EE-14: These diastereomers are prepared in a manner identical to that described in Chart DD with the exception that the epoxide of formula EE-7 (same as BB-7) is used.

CHART FF

The lithium amide of formula FF-2, readily available from the treatment of commercially available (S)-(+)-4-phenyl-2-oxazolidinone with n-butyl lithium in tetrahydrofuran at $-78^\circ C$, is treated with acetyl chloride of formula FF-1 to give

the amide of formula FF-3. Treatment of the compound of formula FF-3 with TiCl_4 followed by treatment with a trialkylamine followed by the addition of commercially available trimethylacetaldehyde affords the compound of formula FF-4. Addition of the amide of formula FF-4 to a tetrahydrofuran solution containing commercially available $\text{CuBr}/(\text{CH}_3)_2\text{S}$ and 3-[bis(trimethylsilyl)amino]phenylmagnesium chloride at -20°C affords the compound of formula FF-5 upon acid workup. Treatment of the aniline of formula FF-5 with benzyl bromide and sodium carbonate in a water/methylene chloride mixture at reflux; or, potassium carbonate in refluxing acetonitrile, affords the compound of formula FF-6.

10 The lithium amide of formula FF-7, readily available from the treatment of commercially available (S)-(-)-4-benzyl-2-oxazolidinone with n-butyl lithium in tetrahydrofuran at -78°C , is treated with acetyl chloride of formula FF-1 to give the amide of formula FF-8. Treatment of the compound of formula FF-8 with TiCl_4 followed by treatment with a trialkylamine followed by the addition of commercially available trimethylacetaldehyde affords the compound of formula FF-9. Addition of the amide of formula FF-9 to a tetrahydrofuran solution containing commercially available $\text{CuBr}/(\text{CH}_3)_2\text{S}$ and 3-[bis(trimethylsilyl)amino]phenylmagnesium chloride at -20°C affords a mixture of compounds of formulae FF-10a and FF-10b. Treatment of the aniline of formula FF-10b with benzyl bromide and sodium carbonate in a water/methylene chloride mixture at reflux; or, potassium carbonate in refluxing acetonitrile, affords the compound of formula FF-11. Treatment of the compound of formula FF-11 with TiCl_4 in methylene chloride followed by the addition of a tertiary amine base then addition of 2-methyl-2-methoxy-1,3-dioxolane affords an intermediate dioxolane (see W-8 in Chart W) which is treated with mild acid to give the compound of formula FF-12. Treatment of the compound of formula FF-12 with TiCl_4 , then a tertiary amine base, followed by addition of either 4-heptanone or 1-phenyl-3-hexanone, affords the aldol product of formula FF-13. Treatment of the compound of formula FF-13 with either sodium hydride or potassium tert. butoxide in an ether solvent then affords the compound of formula FF-14. The compound of formula FF-14 is then hydrogenated under an atmosphere of hydrogen in the presence of a Pd on carbon catalyst to give the compound of formula FF-15. Finally, treatment of the compound of formula FF-15 with a sulfonyl chloride of formula D-7 in Chart D, wherein R_4 is, e.g., 5-trifluoromethyl-2-pyridinyl, in an organic solvent, such as methylene chloride, in the presence of an organic base, such as pyridine, provides the final compound of formula FF-16, wherein R_1 is, e.g., propyl or phenethyl.

CHART GG

Intermediate of formula GG-6 and final products of formula GG-16 are prepared as described in Chart FF with the exception that the (R)-(-)-4-phenyl-2-oxazolidinone and the (R)-(+)-4-benzyl-2-oxazolidinone chiral auxiliaries are used.

5

CHART HH

The compound of formula HH-1 (W-6), prepared as described in Chart W, is converted to the ester of formula HH-2 wherein R is *t*-Bu by addition of potassium *t*-butoxide to a solution of the compound of formula HH-1 in tetrahydrofuran at 0 °C. The compound of formula HH-2 wherein R is *t*-Bu may also be prepared from HH-1 in two steps. First, the oxazolidinone group is cleaved by treatment of the compound of formula HH-1 with lithium hydroxide and hydrogen peroxide at 0 °C in tetrahydrofuran and water. Next, the acid intermediate is treated with N,N-dimethylformamide *t*-butylacetal in refluxing benzene to produce the ester of formula HH-2 (R is *t*-Bu). The ester of formula HH-2 wherein R is Me is prepared by heating a mixture of titanium tetrachloride and HH-1 in methanol. The compound of formula HH-3 is prepared by treatment of the ester of formula HH-2 with lithium diisopropylamide or sodium hexamethyldisilylazide to form an enolate, which is then trapped by ethyl formate to give the compound of formula HH-3. Treatment of this intermediate with tosyl chloride in 1,2-dimethoxyethane gives the compound of formula HH-4, which is then converted to the sulfur derivative of formula HH-5 by treatment with a mixture of potassium hydride and thiophenol in tetrahydrofuran. The compound of formula HH-5 is then deprotonated using *t*-butyllithium in tetrahydrofuran at low temperature. Addition of the epoxide of formula HH-6 (BB-7), prepared as described in Chart BB, and an equivalent of boron trifluoride diethyl etherate affords the compound of formula HH-7. This intermediate is cyclized to the compound of formula HH-8 *in situ*, or it is isolated and treated with sodium hydride in tetrahydrofuran to produce the cyclic compound of the formula HH-8. The sulfur group is then hydrolyzed using either sodium hydroxide in acetonitrile or aqueous copper chloride to give the dihydropyrone derivative of formula HH-9. The benzyl protecting groups are then removed by catalytic hydrogenation using 10% palladium on carbon in ethyl acetate. The resulting amine of formula HH-10 is converted to the desired sulfonamide derivative of formula HH-11 by treatment with 5-cyanopyridine-2-sulfonyl chloride, prepared using the methods described in Chart O, and pyridine in dichloromethane.

35

CHARTS II-OO

The diastereomer of formula II-7 is prepared according to Chart II by

procedures analogous to those described for the preparation of the diastereomeric product in Chart HH. Likewise, stereoisomers of formulae JJ-11, KK-7, LL-11, MM-7, NN-11, and OO-7 are prepared according to Charts JJ, KK, LL, MM, NN, and OO, respectively, by procedures analogous to those described in Chart HH.

5

CHART PP

The compound of formula PP-4 (HH-8) is also generated as described in Chart PP. The acid of formula PP-2 is prepared by treatment of the *t*-butyl ester of formula PP-1 (HH-5), prepared as described in Chart HH, with aqueous acid. The compound of formula PP-2 is then treated with *t*-butyllithium in tetrahydrofuran at low temperature to produce a dianionic intermediate, which is treated with the epoxide of formula PP-3 (BB-7), prepared as described in Chart BB, and an equivalent of boron trifluoride diethyl etherate to afford the compound of formula PP-4 (HH-8).

10

CHARTS QQ-WW

The diastereomer of formula QQ-3 (II-4) is prepared according to Chart QQ by procedures analogous to those described for the preparation of the diastereomeric product in Chart PP. Likewise, stereoisomers of formulae RR-4 (JJ-8), SS-3 (KK-4), TT-4 (LL-8), UU-3 (MM-4), VV-4 (NN-8), and WW-3 (OO-4) are prepared according to Charts RR, SS, TT, UU, VV, and WW, respectively, by procedures analogous to those described in Chart PP.

20

CHART XX

The compound of formula XX-6 (HH-9) is also generated as described in Chart XX. The compound of formula XX-1 (HH-2), prepared as described in Chart HH, is heated neat in commercially-available tris(dimethylamino)methane, bis-(dimethylamino)-methoxymethane or *t*-butoxy-bis(dimethylamino)methane to generate the intermediate of formula XX-2. One equivalent of *t*-butyllithium is added to a solution of this ester in tetrahydrofuran at low temperature to produce an anionic intermediate, which is treated with the epoxide of formula XX-3 (BB-7), prepared as described in Chart BB, and an equivalent of boron trifluoride diethyl etherate to afford the compound of formula XX-4. The intermediate of formula XX-4 is cyclized to the dihydropyrone intermediate XX-5 *in situ*, or XX-4 is isolated and cyclized by treatment with potassium *t*-butoxide or sodium hydride in tetrahydrofuran. Likewise, intermediate XX-5 is hydrolyzed *in situ* to form the compound of formula XX-6 (HH-9), or it is isolated and converted to the dihydropyrone of formula XX-6 (HH-9) by treatment with aqueous acid or aqueous base.

25

30

35

CHARTS YY-EEE

The diastereomer of formula YY-5 (II-5) is prepared according to Chart YY by procedures analogous to those described for the preparation of the diastereomeric product in Chart XX. Likewise, stereoisomers of formulae ZZ-6 (JJ-9), AAA-5 (KK-5), BBB-6 (LL-9), CCC-5 (MM-5), DDD-6 (NN-9), and EEE-5 (OO-5) are prepared according to Charts ZZ, AAA, BBB, CCC, DDD, and EEE, respectively, by procedures analogous to those described in Chart XX.

CHART FFF

The diastereomers of formulae FFF-5 and FFF-7 are also prepared by separation of a diastereomeric intermediate. The diastereomeric mixture of formula FFF-1 (W-11), prepared as described in Chart W, is separated into the single diastereomers of formulae FFF-2 (less polar diastereomer) and FFF-3 (more polar diastereomer) using a preparative chiral HPLC column. The benzyl protecting groups of compounds FFF-2 and FFF-3 are then removed by catalytic hydrogenation using 10% palladium on carbon in ethyl acetate to form the amines of formulae FFF-4 and FFF-6, respectively. The amine intermediates are then converted to the desired sulfonamide derivatives of formulae FFF-5 (HH-11) and FFF-7 (II-7), respectively, by treatment with 5-cyanopyridine-2-sulfonyl chloride, prepared using the methods described in Chart O, and pyridine in dichloromethane.

CHART GGG

The m-nitrocinnamic acid chloride (available from the treatment of the commercially available acid with oxylal chloride) of formula GGG-1 is added to an ether solution of the lithiooxazolidinone of formula GGG-2 (readily available from the treatment of commercially available (R)-(+)-4-benzyl-2-oxazolidinone with n-butyl lithium) to afford the compound of formula GGG-3. The compound of formula GGG-3 is treated with either $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ in ethanol or iron powder in a mixture of ethanol/water and containing ammonium chloride, to effect the reduction of the nitro group to the corresponding amine found in the compound of formula GGG-4. The compound of formula GGG-4 is treated with excess benzyl bromide in the presence of potassium or sodium carbonate in an organic solvent (with methylene chloride/water also being added) to yield the compound of formula GGG-5. Addition of a THF solution of the compound of formula GGG-5 to a THF/dimethylsulfide mixture containing the cuprate reagent prepared from ethyl magnesium bromide and copper bromide/dimethyl sulfide complex affords the compound of formula GGG-6. The compound of GGG-6 is then treated with TiCl_4 , then a tertiary amine, followed by the addition of 2-methyl-2-methoxy-1,3-dioxolane of formula GGG-7 to

2187523

yield the compound of formula GGG-8. Treatment of the compound of formula GGG-8 with perchloric acid then yields the compound of formula GGG-9. Alternately, the compound of formula GGG-6 is treated with a strong base such lithium diisopropylamide in an ether solvent below room temperature and added to a solution of acetyl chloride (also in an ether solvent and cooled to below room temperature) to yield the compound of formula GGG-9. The compound of formula GGG-9 is treated with TiCl_4 in methylene chloride followed by the addition of a tertiary amine, then addition of either 4-heptanone or 1-phenyl-3-hexanone to yield the compound of formula GGG-10. The compound of formula GGG-10 is then treated with either sodium hydride or potassium tert-butoxide in an ether solvent to yield the compound of formula GGG-11. The compound of formula GGG-11 is then hydrogenated to yield the compound of formula GGG-12. The compound of formula GGG-12 is then converted to the final title compound by treatment with a sulfonyl chloride of formula D-7 in Chart D, wherein R_4 is, e.g., 5-trifluoromethyl-2-pyridinyl, in an organic solvent, such as methylene chloride, in the presence of an organic base, such as pyridine, provide the final compound of formula GGG-13, wherein R_1 is, e.g., n-propyl or phenethyl.

Alternatively, addition of the compound of formula GGG-5 to a THF/dimethylsulfide solution containing a mixture of tert-butyl magnesium chloride and copper bromide/dimethylsulfide complex at below 0°C yields a mixture of compounds of formulae GGG-14a and GGG-14b. Both the compounds of formula GGG-14a and GGG-14b are converted to the final products GGG-19 and GGG-20 using the methodology described in Chart GGG for the synthesis of the C-3 ethyl compound of formula GGG-13.

25

CHART HHH

The final compounds of formula HHH-13, HHH-19 and HHH-20 are prepared in the same manner as described for the final compounds in Chart GGG.

CHART III

The commercially available acid of formula III-1 is converted to the compound of formula III-2 by treatment with oxalyl chloride. The acid chloride of formula III-3 is then coupled to the lithio oxazolidinone of formula III-3 (readily available from the treatment of commercially available (S)-(-)-4-benzyl-2-oxazolidinone with n-butyl lithium in an ether solvent) to yield the compound of formula III-4. Addition of the amide of formula III-4 to a tetrahydrofuran solution containing commercially available copper bromide/dimethyl sulfide complex and 3-[bis(trimethylsilyl)-amino]phenylmagnesium chloride at -20°C affords the compounds of formula III-5a

30
35

and III-5b upon acid workup. These compounds are separable by silica gel chromatography. The compound of formula III-5a is treated with benzyl bromide in either acetonitrile or a methylene chloride/water mixture in the presence of either potassium or sodium carbonate to yield the compound of formula III-6. The
5 compound of formula III-6 is treated with TiCl_4 in methylene chloride followed by the addition of a tertiary amine and then 2-methyl-2-methoxy-1,3-dioxolane of formula III-7 is added to yield the compound of formula III-8. Treatment of the compound of the formula III-8 with an acid such as perchloric acid then yields the compound of formula III-9. Treatment of the compound of formula III-9 with TiCl_4
10 in methylene chloride then addition of a tertiary amine, followed by the addition of either 4-heptanone or 1-phenyl-3-hexanone then affords the compound of formula III-10. Treatment of the compound of formula III-10 with either sodium hydride or potassium tert. butoxide then affords the compound of formula III-11. The compound of formula III-11 is hydrogenated to afford the compound of formula III-
15 12. Finally, treatment of the compound of formula III-12 with a sulfonyl chloride of formula D-7 in Chart D, wherein R_4 is, e.g., 5-trifluoromethyl-2-pyridinyl, in an organic solvent, such as methylene chloride, in the presence of an organic base, such as pyridine, provides the final compound of formula III-13, where in R_1 is, e.g., propyl or phenethyl.

20 In an analogous fashion, starting with the compound of formula III-5b, the final compound of formula III-14 is also prepared.

CHART JJJ

The final compounds of formula JJJ-13 and JJJ-14 are prepared using the methodology described in Chart III.

25

CHART KKK

The compound of formula KKK-1 (same as JJJ-9) is treated with TiCl_4 in methylene chloride followed by the addition of a tertiary amine. To that solution is added commercially available hydrocinnamaldehyde to afford the compound of formula KKK-2. The compound of formula KKK-2 is oxidized (e.g. $\text{Me}_2\text{SO}-$
30 $\text{SO}_3/\text{pyridine}$) to yield the compound of formula KKK-3. The compound of formula KKK-3 is treated with propylmagnesium chloride (where R_1 is, e.g., phenyl) to yield the compounds of formula KKK-4a and KKK-4b. Depending on the specific reaction conditions, the ratio of KKK-4a/KKK-4b varies. Alternatively, addition of allylzinc bromide or allylsilane in the presence of TiCl_4 or $n\text{-Bu}_4\text{NF}$ (see Taniguchi et. al.
35 Chemistry Letters 2135, 1992) to the compound of formula KKK-3, followed by hydrogenation, also yields the compounds of formula KKK-4a and KKK-4b.

Depending on the specific reaction conditions the ratio of KKK-4a and KKK-4b vary. The compound of KKK-4a is treated with either sodium hydride or potassium tert. butoxide to yield the compound of formula KKK-5. It is also possible that upon treatment of KKK-3 with allyl zinc bromide, allyl silane or propylmagnesium chloride the intermediate metal alkoxide (metals being magnesium, zinc and titanium) will undergo spontaneous cyclization to yield an unsaturated intermediate which upon hydrogenation leads directly to KKK-5 without the isolation of KKK-4a. The compound of formula KKK-5 is hydrogenated to yield the compound of formula KKK-6. Finally, treatment of the compound of formula KKK-6 with a sulfonyl chloride of formula D-7 in Chart D, wherein R_4 is, e.g., 5-trifluoromethyl-2-pyridinyl, in an organic solvent, such as methylene chloride, in the presence of an organic base, such as pyridine, provides the final compound of formula KKK-7a, wherein, e.g., R_1 and R_2 are phenyl or propyl, respectively.

In an analogous manner to that described for the conversion of the compound of formula KKK-4a to the compound of formula KKK-7a, the compound of formula KKK-4b is converted to the final product of formula KKK-7b.

In an analogous manner to that described for the conversion of the compound of formula KKK-1 to final products of the formula KKK-7a and KKK-7b, the compounds of formula KKK-14a and KKK-14b, wherein R_1 and R_2 are, e.g., methyl or phenethyl, respectively, are prepared by starting with the compound of formula KKK-8 (same as III-6).

In an analogous manner to that described for the conversion of the compound of formula KKK-1 and the compound of formula KKK-8 (each containing the 4-benzyl-2-oxazolidinone auxillary) to the final products of the formulae KKK-7a and KKK-7b, and the final formulae KKK-14a and KKK-14b respectively, the compounds of the formula KKK-15 and the compound of the formula KKK-19 (each containing the 4-phenyl-2-oxazolidinone auxillary) are converted to the final products of the formula KKK-7a and KKK-7b, and the final products of formula KKK-14a and KKK-14b, respectively, wherein R_1 and R_2 are, e.g., methyl or phenethyl, respectively.

CHART LLL

The compound of formula LLL-1 (same as: wherein R is phenyl, AA-1; wherein R is benzyl, GGG-5) is added to a THF solution of commercially available copper bromide/dimethylsulfide complex and tert. butylmagnesium chloride below 0°C to afford the compound of formula LLL-2 as the major diastereomeric product. Where R is defined as benzyl in the compound of formula LLL-2, that compound is

2187523

treated with TiCl_4 in methylene chloride below 0°C followed by the addition of a tertiary amine, then the addition of 2-methyl-2-methoxy-1,3-dioxolane to yield the compound of formula LLL-3. The compound of formula LLL-3 is treated with a protic acid to afford the compound of formula LLL-4. The compound of formula LLL-4 is treated with TiCl_4 in methylene chloride below 0°C followed by the addition of an amine base, then addition of either 4-heptanone or 1-phenyl-3-hexanone affords the compound of formula LLL-5 wherein R_1 is, e.g., n-propyl or phenethyl, respectively. Treatment of the compound of formula LLL-5 with either sodium hydride or potassium tert. butoxide in an ether solvent affords the pyrone of formula LLL-6. Hydrogenation of the compound of formula LLL-6 using, e.g. a Pd on carbon as the catalyst, affords the compound of formula LLL-7. Finally, treatment of the compound of formula LLL-7 with a sulfonyl chloride of formula D-7 in Chart D, wherein R_4 is, e.g., 5-trifluoromethyl-2-pyridinyl, in an organic solvent, such as methylene chloride, in the presence of an organic base, such as pyridine, provides the final compound of formula LLL-8, wherein R_1 is, e.g., propyl or phenethyl.

The compound of formula LLL-2, where R is phenyl, is treated with TiCl_4 in methanol to yield the compound of formula LLL-9. The compound of formula LLL-9 is treated with a base to effect hydrolysis to give the compound of formula LLL-10. The acid of formula LLL-10 is treated with methyl lithium in an ether solvent to yield the compound of formula LLL-11. The ketone of formula LLL-11 is treated with TiCl_4 in methylene chloride below 0°C followed by the addition of an amine base, then addition of either 4-heptanone or 1-phenyl-3-hexanone, to give the compound of formula LLL-12 wherein R_1 is, e.g., n-propyl or phenethyl, respectively. The compound of formula LLL-12 is treated with TiCl_4 in methylene chloride below 0°C followed by the addition of an amine base, then the addition of trimethyl orthoformate to yield the compound of formula LLL-13. The compound of formula LLL-13, in an organic solvent such as THF or methylene chloride, is treated with a base followed by the addition of trimethylsilyl chloride. The solvent is removed from the aforementioned reaction and the resulting protected tertiary alcohol is oxidized (e.g. Ru cat./t-BuOH (see Murahashi et. al. Chemistry Letters 2237, 1992); tritylperchlorate/methylene chloride (see Mukaiyama et. al. Chemistry Letters 1255, 1985), ozone/methylene chloride (see Can. J. Chem. 49, 2465, 1971)) to afford the lactone LLL-6 directly or in a two step sequence where the intermediate ester is lactonized with the aid of either sodium hydride, potassium tert. butoxide or n-Bu₄NF in an ether solvent. The conversion of the compound of formula LLL-6 to

the final product is described above.

Following the same strategy the compound of formula LLL-16 is converted to the final products of formula LLL-23 wherein R_1 is propyl or phenethyl.

CHART MMM

5 The diastereomers of formulae MMM-5 and MMM-7 are also prepared by separation of a diastereomeric mixture of these two compounds. Alternatively, the diastereomeric mixture of formula MMM-1 (X-11 where R_1 is, e.g., phenethyl) prepared as described in Chart X, is separated into the single diastereomers of formulae MMM-2 and MMM-3 using a preparative chiral HPLC column. The benzyl
10 protecting group groups of compounds MMM-2 (less polar diastereomer) and MMM-3 (more polar diastereomer) are then removed by catalytic hydrogenation using 10% palladium on carbon in ethyl acetate to form the amines of formulae MMM-4 and MMM-6, respectively. The amine intermediates are then converted to the desired sulfonamide derivatives of formulae MMM-5 and MMM-7, respectively, by treatment
15 with 5-trifluoromethyl-2-pyridinylsulfonyl chloride, prepared using the methods described in Chart O, and pyridine in methylene chloride.

CHART NNN

 The commercially available (1R, 2S)-(-)-ephedrine of formula NNN-2 is treated with triethylamine and the acid chloride of formula NNN-1 (W-2), prepared as
20 described in Chart W, to afford the amide of formula NNN-3. A *t*-butyl methyl ether solution of this amide at 0 °C is treated sequentially with 1.1 equivalents of propyl magnesium chloride and 2.0 equivalents of 3-[bis(trimethylsilyl)amino]phenyl magnesium chloride, stirred for 3 hours at 0 °C, washed with ammonium chloride solution and concentrated *in vacuo*. The residue is then stirred with silica gel in
25 chloroform to afford the compound of formula NNN-4. Alternatively, the above reaction mixture may be washed with 1 N hydrochloric acid solution during the workup instead of ammonium chloride solution to generate the compound of formula NNN-4. The amine is then converted to the derivative of formula NNN-5 by heating a mixture of the compound of formula NNN-4, 2.2 equivalents of benzyl bromide and
30 2.2 equivalents of sodium carbonate in acetonitrile. The intermediate of formula NNN-5 is then treated with 2 equivalents of lithium diisopropylamide in tetrahydrofuran to form the lithium enolate, which is trapped with acetyl chloride to afford the β -ketoamide of formula NNN-6. A solution of this amide in methylene chloride at low temperature may be treated with 1 equivalent of titanium
35 tetrachloride and 1 equivalent of diisopropylethylamine, followed by 4-heptanone to generate the compound of formula NNN-7. Conversion of the amide of formula

NNN-7 to the dihydropyrone of formula NNN-8 may be accomplished with either sodium hydride in tetrahydrofuran or with aqueous acid. The benzyl protecting groups may then be removed by catalytic hydrogenation using 10% palladium on carbon in ethyl acetate. The resulting amine of formula NNN-9 is converted to the
5 desired sulfonamide derivative of formula NNN-10 (W-12) by treatment with 5-trifluoromethylpyridine-2-sulfonyl chloride, prepared using the methods described in Chart O, and pyridine in dichloromethane.

CHART OOO

The compound of formula OOO-7 (NNN-8) may also be generated as
10 described in Chart OOO. The amide of formula OOO-1 (NNN-5) is treated with aqueous acid to afford the compound of formula OOO-2. The methyl ester of formula OOO-3 is formed from the compound of formula OOO-2 using catalytic acid in methanol. Treatment of the methyl ester of formula OOO-3 with lithium diisopropylamide, followed by trimethylsilyl chloride gives the compound of formula
15 OOO-4. Treatment of this intermediate with either 2-methoxy-2-methyl-1,3-dioxolane followed by hydrolysis or treatment with acetyl chloride affords the β -keto ester of formula OOO-5. This β -keto ester is converted to the compound of formula OOO-6 by treatment of either the titanium enolate (formed using 1 equivalent of titanium tetrachloride and 1 equivalent of diisopropylethylamine in methylene
20 chloride at low temperature) or the lithium dianion (formed using 2 equivalents of lithium diisopropylamide in tetrahydrofuran at low temperature) with 4-heptanone. The dihydropyrone of formula OOO-7 (NNN-8) is formed by treatment of the compound of formula OOO-6 with either sodium hydride in tetrahydrofuran or aqueous base.

25

CHART PPP

Reduction of the commercially available ethyl 4,4,4-trifluorobutyrate of formula PPP-1, with DiBAL-H followed by in situ alkylation with 2-phenethyl magnesium bromide or chloride produces the alcohol of formula PPP-2. Swern oxidation of the alcohol gives the ketone of formula PPP-3. The ketone is converted
30 to the dihydropyrone of formula PPP-4 by alkylation with the dianion of methyl acetoacetate followed by saponification to the acid and lactonization with base.

CHART QQQ

The aluminum trichloride catalyzed reaction of the dihydropyrone of formula QQQ-1 (PPP-4), prepared as described in Chart PPP, with the CBZ-protected 3-
35 aminobenzaldehyde (which is available from the reaction of benzyl chloroformate with commercially available 3-aminobenzaldehyde) of formula QQQ-2 and

2187523

subsequent reaction with trialkyl aluminums or Grignard reagents in the presence of cuprous bromide-dimethylsulfide complexes provides compounds of formula QQQ-3. The individual stereoisomers are separated by HPLC using a chiral stationary phase to give the four possible stereoisomers of formula QQQ-4, QQQ-5, QQQ-6, and QQQ-7. Transfer hydrogenation of each stereoisomer with Pd/C and ammonium formate gives the amines of formula QQQ-8, QQQ-9, QQQ-10, and QQQ-11. Treatment of the amines with sulfonyl chlorides of general formula QQQ-12 and pyridine in methylene chloride provides compounds of general formula QQQ-13, QQQ-14, QQQ-15, and QQQ-16, wherein R_2 is, e.g., 5-cyano-2-pyridinyl, 1-methyl-4-imidazolyl, or 5-amino-2-pyridinyl.

CHART RRR

The procedure for the preparation of compounds of formula RRR-11 to RRR-15 is described in Chart RRR. The pyrone RRR-A is coupled to the Cbz protected benzaldehyde RRR-B in THF with $AlCl_3$ followed by treatment of the resulting intermediate with R_1MgX where $X = Br$ or Cl in THF with added $CuBr.Me_2S$ to give RRR-1. De-protection of the resulting intermediate with 10% Pd/C in methanol with added ammonium formate gives RRR-2. Separation of the racemic compound RRR-1 into its 4 enantiomers gives RRR-3 to RRR-6. De-protection of the resulting intermediates with 10% Pd/C in methanol with added ammonium formate gives the free amines RRR-7 to RRR-10. Treatment of the amines RRR-7 to RRR-10 and RRR-2 with an appropriate sulfonyl chloride gives the sulfonamides RRR-11 to RRR-14 and RRR-15, respectively.

CHART SSS

The procedure for the preparation of compounds of formula SSS-7 to SSS-9 is described in Chart SSS. The pyrone SSS-A is coupled to the Cbz protected benzaldehyde SSS-B in THF with $AlCl_3$ followed by treatment of the resulting intermediate with R_1MgX where $X = Br$ or Cl in THF with added $CuBr.Me_2S$ to give SSS-1. De-protection of the resulting intermediate with 10% Pd/C in methanol with added ammonium formate gives SSS-2. Separation of the racemic compound SSS-1 into its 2 enantiomers gives SSS-3 to SSS-4. De-protection of the resulting intermediates with 10% Pd/C in methanol with added ammonium formate gives the free amines SSS-5 to SSS-6. Treatment of the amine with an appropriate sulfonyl chloride gives the sulfonamides SSS-7 to SSS-9.

CHART TTT

The procedure for the preparation of compounds of formula TTT-6 and TTT-7 is described in Chart TTT. The pyrone TTT-A is coupled to the Cbz protected

benzaldehyde TTT-B in THF with AlCl_3 followed by treatment of the resulting intermediate with R_1MgX where $\text{X} = \text{Br}$ or Cl in THF with added $\text{CuBr} \cdot \text{Me}_2\text{S}$ to give TTT-1. Separation of the racemic compound TTT-1 into its 2 enantiomers gives TTT-2 and TTT-3. De-protection of the resulting intermediates with 10% Pd/C in methanol with added ammonium formate gives the free amines TTT-4 and TTT-5. Treatment of the amine with an appropriate sulfonyl chloride gives the sulfonamides TTT-6 and TTT-7.

CHART UUU

Reaction between commercially available thiourea in hot ethanol with commercially available 2-chloro-5-nitropyridine of formula UUU-1 affords the isothiurea compound of formula UUU-2. Treatment of the compound of formula UUU-2 with aqueous sodium carbonate and sodium hydroxide provides the thiol compound of formula UUU-3. Oxidation of the compound of formula UUU-3 with chlorine gas provides the sulfonyl chloride compound of formula UUU-4. Treatment of the compound of formula D-5 (e.g., the compound of formula T-4 wherein R_1 is 2-phenylethyl, R_2 is propyl, R_3 is *tert*-butyl) in dichloromethane with two equivalents of pyridine followed by one equivalent of the compound of formula UUU-4 gives the sulfonamide compound of formula UUU-5 (wherein R_1 is 2-phenylethyl, R_2 is propyl, R_3 is *tert*-butyl). Reduction of the compound of formula UUU-5 with palladium on carbon and ammonium formate affords the compound of formula UUU-6, which is the compound: 5-amino-N-[3-(1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide (Formula UUU-6: R_1 is 2-phenylethyl, R_2 is propyl, R_3 is *tert*-butyl).

CHART VVV

The compound of Formula VVV-1, which is 2-mercapto-5-carbamoylpyridine, is prepared via published procedure (J. Chem. Soc. 1948, 1939-1945). Treatment of a suspension of this compound in dilute hydrochloric acid with chlorine gas at 0° provides the sulfonyl chloride of Formula VVV-2.

CHART WWW

Amines of the generic formula WWW-1 are reacted with benzyl chloroformate to provide CBZ derivatives WWW-2. The individual stereoisomers of formula WWW-2 are generally separated by chiral HPLC methods, and then converted back to the free amines WWW-3 *via* hydrogenolysis. Sulfonation of the amines in the usual manner known to one of ordinary skill in the art provides the final compounds of formula WWW-4, in stereochemically pure form.

CHART XXX

Dihydropyrone XXX-1, which is prepared by procedures analogous to those in described in Preparations 17 and 84, is condensed with *meta*-nitrobenzaldehyde in the presence of aluminum trichloride to provide the benzyldiene intermediate XXX-2. Conjugate reduction of the double bond using sodium cyanoborohydride, followed by
5 reduction of the nitro group *via* catalytic hydrogenation, affords amine of formula XXX-4, which is converted to the sulfonamides XXX-5 by treatment with the appropriate sulfonyl chloride in dichloromethane and pyridine.

CHART YYY

Dihydropyrones of Formula YYY-1, wherein R₁ and R₂ are propyl or
10 phenethyl, and which are synthesized as described in Preparation 84, are condensed with the aldehyde of Formula B-2 using aluminum trichloride to provide the benzyldiene intermediates of formula YYY-2. Conjugate addition of *tert*-butylmagnesium chloride in the presence of copper (I) bromide-dimethyl sulfide provides compounds of Formula YYY-3. Hydrogenolytic deprotection affords amines
15 of formula YYY-4, which are converted to the sulfonamides of formula YYY-5 using the appropriate sulfonyl chloride in dichloromethane with added pyridine. The procedures used are analogous to those described for Chart D.

CHART ZZZ

Polymeric *meta*-aminobenzaldehyde is protected by treating with benzyl
20 bromide and potassium carbonate in acetonitrile at reflux to yield the compound of formula ZZZ-2. A vinyl anion is generated from 2-bromovinyltrimethylsilane of formula ZZZ-3 by treatment with *t*-butyl lithium at -78°C to -20°C. The vinyl anion so generated is cooled to -78°C and the diprotected *meta*-aminobenzaldehyde of formula ZZZ-2 is added to afford the desired allylic alcohol of formula ZZZ-3. The
25 alcohol is easily converted to the acetate or carbonate of formula ZZZ-5 by standard means (e.g., CH₃COCl, pyridine, CH₂Cl₂, 0°C). These substrates participate in palladium catalyzed allylic substitutions as delineated in Charts AAAA-CCCC (C.G. Frost; J. Howarth; J.M.J. Williams, Tetrahedron: Asymmetry (1992) 3:1089-1122).

CHART AAAA

30 The sodium salt of methyl acetoacetate of formula AAAA-1 generated by treating methyl acetoacetate with sodium hydride at 0°C in either DMF or THF acts as the nucleophile in a palladium catalyzed allylic substitution. If this reaction is run in the presence of palladium allyl chloride dimer of formula AAAA-3 as the palladium source and a chiral phosphine ligand (P. von Matt; A. Pfaltz, Angew.
35 Chem. Int. Ed. Engl. (1993) 32:566-568), a kinetic resolution of the starting allylic acetate or carbonate results in the synthesis of optically enriched allylated product

of formula AAAA-4. If the reaction with nucleophile is slow, the acetate generated from formation of the *pi*-allyl palladium intermediate isomerizes the two possible diastereomeric *pi*-allyl complexes so that a stereoselective synthesis of the allylated product occurs (B.M. Trost; P.E.J. Strege, Am. Chem. Soc. (1977) 99:1649). Treatment of the resulting vinyl silane of formula AAAA-4 with *para*-toluenesulfonic acid in acetonitrile at reflux affords the desilylated olefin of formula AAAA-5. The dihydropyrone product of formula AAAA-7 is formed by generating the dianion of the β -ketoester under standard conditions (J.R. Peterson; T.J. Winger; C.P. Miller, Syn. Comm. (1988) 18(9):949-963), (NaH, *n*-butyllithium, THF) and quenching with an appropriate symmetrical ketone of formula AAAA-6 (such as 4-heptanone). Hydrolysis of the ester (0.1 N NaOH/THF) and acidic work-up provide the dihydropyrone product of formula AAAA-7. Standard hydrogenation conditions reduce the olefin and deprotect the amine. Subsequent treatment of the amino compound with the appropriate sulfonyl chloride of formula AAAA-8 (pyridine, CH₂Cl₂) provides the desired sulfonamide protease inhibitor of formula AAAA-9.

CHART BBBB

Alternatively, the palladium catalyzed allylic substitution may be performed with the sodium anion of the requisite dihydropyrone J.R. (Peterson; T.J. Winger; C.P. Miller, Syn. Comm. (1988) 18(9):949-963) of formula BBBB-1 (dihhydropyrone, NaH, THF or DMF, 0°C) as the nucleophilic partner. Once again, if palladium allyl chloride dimer of formula BBBB-3 and a chiral phosphine ligand (P. von Matt; A. Pfaltz, Angew. Chem. Int. Ed. Engl. (1993) 32:566-568) are employed as catalyst, a kinetic resolution results in the synthesis of optically pure allylated dihydropyrone of formula BBBB-4; and a stereoselective synthesis of the allylated product will occur if the reaction with nucleophile is slow relative to isomerization of the two possible diastereomeric *pi*-allyl complexes by acetate generated from formation of the *pi*-allyl palladium intermediate. Subsequent, desilylation (*p*-TsOH, CH₃CN), olefin reduction and amine deprotection (H₂/Pd/C), and sulfonylation of the amine (ArSO₂Cl, pyridine, CH₂Cl₂) with a compound of the formula BBBB-5 provides the desired dihydropyrone protease inhibitor of formula BBBB-6.

CHART CCCC

Treatment of *m*-bis(benzyl)aminobenzoic acid of formula CCCC-1 with oxalyl chloride to form the acid chloride and reaction with bis(trimethylsilyl)acetylene and AlCl₃ in methylene chloride affords the propargylic ketone of formula CCCC-2. Asymmetric reduction of the ketone with a chiral borane (H.C. Brown; Beeraraghavan Ramachandran, P. Acc. Chem. Res. (1992) 25:16-24) such as DIP chloride [(+) or (-)-

β -chlorodiisopinocampheylborane] and acetylene reduction with REDAL^{*} provides the allylic alcohol of formula CCCC-3, primarily as a single enantiomer. Formation of the carbonate of formula CCCC-4 (methyl chloroformate, pyridine, CH₂Cl₂, 0°C) and subjection to palladium catalyzed allylic substitution with the desired dihydropyrone of formula CCCC-5 as nucleophile affords primarily one enantiomer of the allylated dihydropyrone of formula CCCC-6 (retention of configuration) (T. Hayashi; T. Hagihara; M. Konishi; M.J. Kumada, Am. Chem. Soc. (1983) 105:7768-7770). This product is transformed into the desired protease inhibitor of formula CCCC-7 as previously described in Chart BBBB.

10

CHART DDDD

The known cycloalkylpyranones of formula DDDD-1 are prepared by acylation of the trimethylsilyl enol ether of the corresponding cycloalkyl ketone with malonyl dichloride as described in R. Effenberger, T. Ziegler, K.-H. Schonwalder, T. Kesmarszky, B. Bauer Chem. Ber. 119: 3394-3404 (1986). Catalytic hydrogenation of the cycloalkylpyranones of formula DDDD-1 with platinum oxide (PtO₂) in acetic acid produces the cycloalkyldihydropyrones of Formula DDDD-2. The intermediate of formula DDDD-3 is then formed by aluminum chloride (AlCl₃) catalyzed condensation of the compound of formula DDDD-2 with 3-nitrobenzaldehyde, which is commercially available. Subsequent reaction of the intermediate of formula DDDD-3 with trialkyl aluminums in the presence of copper bromide-dimethyl sulfide complex (CuBr-Me₂S) or zinc reagents generated from zinc metal, alkyl halide, cuprous cyanide (CuCN) and lithium chloride (LiCl) provides compounds of formula DDDD-4 which contain a C-3 α branched substituent. Catalytic hydrogenation of compounds of the formula DDDD-4 with Pd/C in ethanol (EtOH) provides the amine derivatives of the formula DDDD-5. Treatment of the compounds of formula DDDD-5 with sulfonyl chlorides of formula DDDD-6 and pyridine in methylene chloride (CH₂Cl₂) provides compounds of the formula DDDD-7 (e.g., wherein n is 1, 2, or 3; R₁ is ethyl or *t*-butyl; R₂ is 4-cyanophenyl or 5-cyano-2-pyridyl).

Procedures by which the compounds of the present invention are prepared are also described in International application, (WO 94/11361, published 26 May 1994), and International application, (WO 94/18188, published 18 August 1994).

As is apparent to those of ordinary skill in the art, the compounds of the present invention can occur in several diastereomeric forms, depending on the configuration around the asymmetric carbon atoms. All such diastereomeric forms

35

*Trade-mark

are included within the scope of the present invention.

Also, the dihydropyrones of the present invention can be separated into individual stereoisomers or prepared as individual diastereomers. A diastereomeric pair can be prepared wherein C-3 α is a homogeneous center and C-6 is a mixture.

- 5 All such enantiomeric and diastereomeric forms, and mixtures thereof, are included within the scope of the present invention.

The compounds of the present invention of formula I can exist in several tautomeric forms, including the particular enol forms as depicted by formula I and IA and the keto form of formula IB. (For formulas I, IA and IB, the dashed line
10 indicates that a double bond may be present or absent.) All such tautomeric forms are included within the scope of the present invention. For compounds of the present invention which are 4-hydroxy-pyran-2-ones of formula VII, the enol form predominates. For compounds of the present invention which are 5,6-dihydro-4-hydroxy-pyran-2-ones of formula VI, a mixture of the enol and keto forms is
15 commonly expected.

Also, the compounds of the present invention of formula II can exist in several tautomeric forms of the 4-hydroxy-pyrone ring, including the particular enol forms depicted by formulas II and IIA, and the particular keto form depicted by formula IIB, and mixtures thereof. All such tautomeric forms are included within
20 the scope of the present invention.

The compounds of the present invention may be in either free form or in protected form at one or more of the remaining (not previously protected) carboxyl, amino, hydroxy, or other reactive groups. The protecting groups may be any of those known in the art. Examples of nitrogen and oxygen protecting groups are set forth
25 in T. W. Greene, Protecting Groups in Organic Synthesis, Wiley, New York, (1981); J. F. W. McOmie, ed. Protective Groups in Organic Chemistry, Plenum Press (1973); and J. Fuhrhop and G. Benzlin, Organic Synthesis, Verlag Chemie (1983). Included among the nitrogen protective groups are t-butoxycarbonyl (BOC), benzyloxycarbonyl, acetyl, allyl, phthalyl, benzyl, benzoyl, trityl and the like.

30 The present invention provides for compounds of formula I and II or pharmacologically acceptable salts and/or hydrates thereof. Pharmacologically acceptable salts refers to those salts which would be readily apparent to a manufacturing pharmaceutical chemist to be equivalent to the parent compound in properties such as formulation, stability, patient acceptance and bioavailability. Examples of salts
35 of the compounds of formula I include acidic salts, such as sodium, potassium, lysine, arginine and calcium salts, and basic salts, such as the hydrochloride salt,

wherein the R substituents in formula I contain a basic moiety. Examples of salts of the compounds of formula II include the hydrohalide salts, such as the hydrochloride and hydroiodide salts; and the sodium, potassium, calcium, lysine and arginine salts.

Also included as salts of the compounds of formulae I and II of the present invention are the bis-salts, such as the bis-arginine, bis-lysine, bis-sodium, bis-potassium and bis-calcium salts, provided that the compound contains, for example, -NHSO₂⁻, -SO₃H, -CONH-, -OH or COOH. The bis-sodium salt is most preferred.

The compounds of the present invention are useful for treating patients infected with human immunodeficiency virus (HIV) which results in acquired immunodeficiency syndrome (AIDS) and related diseases. For this indication, these compounds may be administered by oral, intranasal, transdermal, subcutaneous and parenteral (including intramuscular and intravenous) routes in doses of 0.1 mg to 100 mg/kg of body weight per day.

Those skilled in the art would know how to formulate the compounds of this invention into appropriate pharmaceutical dosage forms. Examples of the dosage forms include oral formulations, such as tablets or capsules, or parenteral formulations, such as sterile solutions.

When the compounds in this invention are administered orally, an effective amount is from about 0.1 mg to 100 mg per kg of body weight per day. Either solid or fluid dosage forms can be prepared for oral administration. Solid compositions, such as compressed tablets, are prepared by mixing the compounds of this invention with conventional ingredients such as talc, magnesium stearate, dicalcium phosphate, magnesium aluminum silicate, calcium sulfate, starch, lactose, acacia, methyl cellulose, or functionally similar pharmaceutical diluents and carriers. Capsules are prepared by mixing the compounds of this invention with an inert pharmaceutical diluent and placing the mixture into an appropriately sized hard gelatin capsule. Soft gelatin capsules are prepared by machine encapsulation of a slurry or solution of the compounds of this invention with an acceptable inert oil such as vegetable oil or light liquid petrolatum.

Pharmaceutically acceptable formulations of the disodium salts of the compounds of the present invention include: soft elastic capsules (SEC) containing a suspension of the salt; salt tablets; salt spray coated sucrose beads; or salt spray dried matrix with an enteric or non-enteric polymer.

Formulations of the compounds of the present invention, which present the compounds in free acid form, preferably contain the free acid in non-crystalline form. Examples of such formulations include: soft elastic capsules containing free acid

solution; non-crystalline spray dried matrix of the free acid with an enteric or non-enteric polymer; or a solid non-crystalline matrix of free acid in polyethyleneglycol (PEG) or Gelucire^{*} 44/14 (Gattefosse, Saint Priest, France).

Syrups are prepared by dissolving the compounds of this invention in an aqueous vehicle and adding sugar, aromatic flavoring agents and preservatives. Elixirs are prepared using a hydroalcoholic vehicle such as ethanol, suitable sweeteners such as sugar or saccharin and an aromatic flavoring agent. Suspensions are prepared with an aqueous vehicle and a suspending agent such as acacia, tragacanth, or methyl cellulose.

When the compounds of this invention are administered parenterally, they can be given by injection or by intravenous infusion. An effective amount is from about 0.1 mg to 100 mg per kg of body weight per day. Parenteral solutions are prepared by dissolving the compounds of this invention in liquid vehicle and filter sterilizing the solution before placing in a suitable sealable vial or ampule. Parenteral suspensions are prepared in substantially the same way except a sterile suspension vehicle is used and the compounds of this invention are sterilized with ethylene oxide or suitable gas before it is suspended in the vehicle.

The exact route of administration, dose, or frequency of administration would be readily determined by those skilled in the art and is dependant on the age, weight, general physical condition, or other clinical symptoms specific to the patient to be treated.

Patients to be treated would be those individuals: 1) infected with one or more than one strain of a human immunodeficiency virus as determined by the presence of either measurable viral antibody or antigen in the serum and 2) having either an asymptomatic HIV infection or a symptomatic AIDS defining infection such as i) disseminated histoplasmosis, ii) isoporiasis, iii) bronchial and pulmonary candidiasis including pneumocystis pneumonia, iv) non-Hodgkin's lymphoma, or v) Kaposi's sarcoma and being less than sixty years old; or having an absolute CD4+ lymphocyte count of less than 500/mm³ in the peripheral blood. Treatment would consist of maintaining an inhibitory level of the compounds of this invention in the patient at all times and would continue until the occurrence of a second symptomatic AIDS defining infection indicates alternate therapy is needed.

The utility of representative compounds of the present invention has been demonstrated in the biological tests described below:

The HIV protease screening assay is based on fluorescently labeled substrate which can be resolved from nonlabeled cleavage product using special beads coated

*Trade-mark

with streptavidin. The substrate is biotinylated at the amino terminal arginine and fluorescently labeled with fluorescein isothiocyanate (FITC) at the carboxyl terminal lysine. This assay has been employed to detect novel, nonpeptidic inhibitors of HIV-1 protease. Substrate (20 μ l of 0.2 μ M), sample (10 μ l of desired concentraion), and
 5 enzyme (10 μ l of 0.1 μ M) are added to a 96 well pandex plate. The assay is run in 0.1 M sodium acetate buffer at pH 5.5 in the presence of 1.0 M sodium chloride and 0.05% NP-40 with incubated in the dark for one hour at room temperature. Streptavidin coated polystyrene beads (40 μ l of 0.1% (w/v)) are added and the plate is incubated in the dark for an additional half hour. The labeled cleavage product is
 10 separated from the unreacted substrate via filtration and is read on the Idexx^{*} screen machine. The data are analyzed by appropriate computer algorithms to ascertain percent inhibition values.

Determination of K_i values utilizes the same materials and equipment employed for percent inhibition studies. Two-fold serial dilutions are made for a
 15 given inhibitor from 2, 3 or 4 starting concentrations with a total of 24, 36 or 48 individual inhibitor concentrations. These dilutions are performed utilizing the BioMek^{*} robotics system. The assay consists of 10 μ L of 40 nM HIV-1 protease, 10 μ L of the various inhibitor concentrations, and 20 μ L of 200 μ M substrate (40 μ L total). The reaction is allowed to proceed for 90 min at room temperature,
 20 terminated with 40 μ L of avidin beads and processed (*supra vide*). An inhibitor with a known K_i is run in parallel to verify the validity of the assay. The data is processed utilizing a computer program employing a nonlinear least square analysis of the data to generate the K_i values.

The % inhibition values and/or K_i values of representative compounds of the
 25 present invention tested in the HIV protease screening assay are listed in Table I below.

In the enzyme inhibition assay described above, the sensitivity of K_i value determination is in part limited by the ability to continue to lower the enzyme concentration for compounds with high binding affinity. To prevent de-dimerization
 30 at low enzyme concentration, a tandemly linked enzyme is prepared in which the two monomers are covalently linked by an appropriate stretch of amino acid residues. Using the latter enzyme, the sensitivity of the inhibition assay is improved since much lower enzyme concentration can be utilized, as compared to the condition using the wild-type enzyme.

35 Protocol for K_i value determination with tandem HIV protease: Due to the greater stability (no dedimerization) of the single chain tethered (tandem) HIV

*Trade-mark

protease enzyme, in which the two monomeric units are engineered to be linked by a polypeptide stretch, the method for the determination of K_i values for inhibitors uses very low concentrations of enzyme (0.2 nM) and increased incubation times (96 hours) at room temperature to improve the sensitivity in the measurement of K_i values for very potent inhibitors. The starting inhibitor concentrations are determined based on preliminary enzyme inhibition screening results which estimate the expected potency of the inhibitor. Inhibitor concentrations are then prepared using the Biomek 1000 (Beckman) and the Quadra 96 (Tomtec)*. Substrate (biotinylated at the amino terminal arginine and fluorescently labeled with fluorescein at the carboxyl terminal lysine), inhibitor and the tandem enzyme are allowed to react in solution at pH 5.5 (buffers identical to those used with the native dimeric enzyme) in the dark for 96 hours. Streptavidin coated polystyrene beads are added to stop the reaction. The labeled cleavage product is separated from unreacted substrate via filtration. Residual fluorescence is quantitated with the Idexx SM2000 (Idexx) and the resulting data are analyzed using the NLLSF* program.

The % inhibition values and/or K_i values of representative compounds of the present invention tested in the HIV protease screening assay and/or tandem HIV protease assay are listed in Table II below.

Several compounds of the present invention, such as N-[3-(1-[5,6-Dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide were tested in known human cell lines, such as human T-cell lines, e.g., MT4 and H9, which were infected with HIV-1_{IIIB}, and certain of these compounds were further tested in peripheral blood mononuclear cells (PBMC), which were infected with HIV-1_{JRCSF} (a clinical isolate). The compounds were found to inhibit retroviral replication.

The following compounds of the present invention are preferred:

5-Cyano-N-[3-(1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide

N-[3-(1-(4-Hydroxy-5,6-dihydro-2-oxo-6-phenethyl-6-propyl-2H-pyran-3-yl)propyl)phenyl]-5-cyanopyridine-2-sulfonamide

N-[3-(1-[5,6-Dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide

N-[3-[1-(4-hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide

N-[3-(1-(4-Hydroxy-5,6-dihydro-2-oxo-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-

*Trade-mark

- pyran-3-yl)-2,2-dimethylpropyl]phenyl]-5-cyanopyridine-2-sulfonamide
- N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide
- N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide
- 5 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide
- N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide
- 10 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide
- N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-(4-fluorophenyl)ethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide
- N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-(4-fluorophenyl)ethyl)-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide
- 15 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-(4-fluorophenyl)ethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide
- N-[3-(1-[6,6-Bis-(2-cyclopropyl-ethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-yl]-2,2-dimethyl-propyl)phenyl]-5-cyano-2-pyridinesulfonamide
- 20 N-[3-(1-[6,6-Bis-(2-cyclopropyl-ethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-yl]propyl)phenyl]-5-cyano-2-pyridinesulfonamide
- N-[3-(1-[6,6-Bis-(2-cyclopropyl-ethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-yl]-2,2-dimethyl-propyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide
- 5-cyano-N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R or S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,
- 25 N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(R or S)-(2-phenethyl)-6-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
- 30 5-amino-N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R or S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridine-sulfonamide,
- N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-[6-(R or S)-propyl]-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,
- 35 5-Trifluoromethyl-N-[3-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,

5-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide, or (3R or S)-N-[3-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6,6-dipropyl-2H-pyran-3-yl)propyl]phenyl]-5-(trifluoromethyl)-2-pyridinesulfonamide

5 5-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-phenethyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,

5-Trifluoromethyl-N-[3-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,

10 5-Trifluoromethyl-N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,

5-Trifluoromethyl-N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide, or (3R or S, 6R or S)-N-[3-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6-propyl-6-phenethyl-2H-pyran-3-yl)propyl]phenyl]-5-(trifluoromethyl)-2-pyridinesulfonamide

15 5-Amino-N-[3-(1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,

5-Cyano-N-[3(R or S)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,

20 5-Cyano-N-[3-(1-[5,6-dihydro-6,6-diisobutyl-4-hydroxy-2-oxo-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,

N-[3(R or S)-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6(R or S)-[2-phenylethyl]-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

25 5-Cyano-N-[3-(1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]cyclopropylmethyl)phenyl]-2-pyridinesulfonamide,

5-Amino-N-[3(R or S)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,

30 5-Amino-N-[3(R or S)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]propyl)phenyl]-2-pyridinesulfonamide,

5-Amino-N-[3(R or S)-(1-[6(R or S)-(2-[4-fluorophenyl]ethyl)-5,6-dihydro-4-hydroxy-2-oxo-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,

35 N-[3(R or S)-(1-[5,6-Dihydro-6,6-dipropyl-4-hydroxy-2-oxo-2H-pyran-3-yl]-2,2-

- dimethylpropyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
 5-Amino-N-[3(*R* or *S*)-(1-[5,6-dihydro-6,6-dipropyl-4-hydroxy-2-oxo-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,
 5-Cyano-N-[3(*R* or *S*)-(1-[5,6-dihydro-6,6-dipropyl-4-hydroxy-2-oxo-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,
 5 N-[3(*R* or *S*)-(1-[6,6-Bis(2-phenylethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-yl]propyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
 N-[3(*R* or *S*)-(1-[6,6-Bis(2-phenylethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-yl]propyl)phenyl]-5-cyano-2-pyridinesulfonamide,
 10 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide,
 N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R* or *S*)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide,
 N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S* or *R*)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
 15 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
 N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R* or *S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide,
 20 N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R* or *S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
 N-[3-{1(*S* or *R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-aminopyridine-2-sulfonamide
 25 N-[3-{1(*S* or *R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide
 N-[3-{1(*S* or *R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide
 30 N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide
 The following compounds of the present invention are more preferred:
 5-Cyano-N-[3-(1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide
 35 5-cyano-N-[3-(*R* or *S*)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(*R* or *S*)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-

pyridinesulfonamide,

5-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide, or (3R or S)-N-[3-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6,6-dipropyl-2H-pyran-3-yl)propyl]phenyl]-5-

5 (trifluoromethyl)-2-pyridinesulfonamide

5-Trifluoromethyl-N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide, or (3R or S, 6R or S)-N-[3-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6-propyl-6-phenethyl-2H-pyran-3-yl)propyl]phenyl]-5-(trifluoromethyl)-2-pyridinesulfonamide

10 5-Cyano-N-[3(R or S)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridine-sulfonamide,

N-[3(R or S)-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6(R or S)-[2-phenylethyl]-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-
15 sulfonamide,

N-[3-[1(R or S)-(4-Hydroxy-5,6-dihydro-2-oxo-6(S or R)-phenethyl-6-propyl-2H-pyran-3-yl)propyl]phenyl]-5-cyanopyridine-2-sulfonamide,

N-[3-[1(R or S)-(4-Hydroxy-5,6-dihydro-2-oxo-6(R or S)-(2-(4-fluorophenyl)-ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl]phenyl]-5-cyanopyridine-2-
20 sulfonamide,

N-[3-[1(R or S)-(4-Hydroxy-5,6-dihydro-2-oxo-6(R or S)-(2-(4-fluorophenyl)-ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide, and

N-[3-[1(R or S)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-propyl]phenyl]-5-cyanopyridine-2-sulfonamide.
25

The following compounds of the present invention are most preferred (see Chart EEE):

N-[3-[1(R or S)-(4-Hydroxy-5,6-dihydro-2-oxo-6(S or R)-phenethyl-6-propyl-2H-pyran-3-yl)propyl]phenyl]-5-cyanopyridine-2-sulfonamide of formula EEE-1,

30 N-[3-[1(R or S)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-propyl]phenyl]-5-cyanopyridine-2-sulfonamide of formula EEE-2,

N-[3(R or S)-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6(R or S)-[2-phenylethyl]-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide of formula EEE-3,

35 5-Trifluoromethyl-N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide

of formula EEE-4, or (3R or S, 6R or S)-N-[3-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6-propyl-6-phenethyl-2H-pyran-3-yl)propyl]phenyl]-5-(trifluoromethyl)-2-pyridinesulfonamide

5 5-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide of formula EEE-5, or (3R or S)-N-[3-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6,6-dipropyl-2H-pyran-3-yl)propyl]phenyl]-5-(trifluoromethyl)-2-pyridinesulfonamide.

Also the following compounds of the present invention, which are readily prepared by the synthetic procedures set out herein, are most preferred:

10 (3R)-N-[3-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6,6-dipropyl-2H-pyran-3-yl)propyl]phenyl]-5-(trifluoromethyl)-2-pyridinesulfonamide

(3R)-N-[3-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6-propyl-6-phenethyl-2H-pyran-3-yl)propyl]phenyl]-5-(trifluoromethyl)-2-pyridinesulfonamide.

2187523

DESCRIPTION OF THE PREFERRED EMBODIMENTS

In the Preparations and Examples below and throughout this document:

	$^{\circ}\text{C}$	is	degrees Centigrade.
	$^1\text{H-NMR}$	is	proton nuclear magnetic resonance spectrum.
5	$^{13}\text{C-NMR}$	is	carbon nuclear magnetic resonance spectrum.
	δ	is	chemical shift (parts per million) relative to TMS.
	AlCl_3	is	aluminum chloride.
	Anal.	is	analytical data.
	Br	is	benzyl.
10	CBZ	is	benzyloxycarbonyl.
	CDCl_3	is	deuterio-chloroform.
	CD_3OD	is	deuterio-methanol.
	CH_2Cl_2	is	methylene chloride.
	cm^{-1}	is	reciprocal centimeters.
15	CuBr_2	is	cupric bromide.
	DMSO	is	dimethylsulfoxide.
	DMSO-D_6	is	deuterio dimethylsulfoxide.
	EI MS	is	electron impact mass spectroscopy.
	EtOAc	is	ethyl acetate.
20	Et_3Al	is	triethyl aluminum.
	FAB MS	is	fast-atom-bombardment mass spectroscopy.
	HCl	is	hydrochloric acid.
	H_2O	is	water.
	HOBT	is	1-hydroxybenzotriazole hydrate.
25	HRMS	is	high-resolution mass spectroscopy.
	KOH	is	potassium hydroxide.
	M	is	molar (concentration).
	MeOH	is	methanol.
	Me_2S	is	dimethyl sulfide.
30	mg	is	milligram.
	MgSO_4	is	magnesium sulfate.
	mL	is	milliliter.
	mmHg	is	millimeter of mercury.
	MP	is	melting point.
35	N	is	normal (concentration).

	NaCl	is	sodium chloride.
	NaOH	is	sodium hydroxide.
	NaH	is	sodium hydride.
	NaHCO ₃	is	sodium bicarbonate.
5	Na ₂ CO ₃	is	sodium carbonate.
	Na ₂ SO ₄	is	sodium sulfate.
	NH ₄ Cl	is	ammonium chloride.
	Pd/C	is	palladium on charcoal.
	R _f	is	chromatographic movement relative to solvent front.
10	TFA	is	trifluoroacetic acid.
	THF	is	tetrahydrofuran.
	TMS	is	tetramethyl silane.

The following Preparations and Examples illustrate the present invention:

15 PREPARATION 1 Cyclopropyl-(3-nitrophenyl)methanone (Formula A-2) Refer to Chart A.

A 500-mL, three-necked, round-bottomed flask with a gas outlet and a 250-mL pressure-equalizing addition funnel is charged with cyclopropyl phenyl ketone of formula A-1 (30 mL) and cooled to -40 °C. The addition funnel is charged with nitric acid (180 mL), which is added to the reaction mixture dropwise over 2 h. The
20 reaction mixture is stirred another 3.5 h at -40 - 0 °C, and then quenched by pouring onto 500 mL of ice. The mixture is extracted with three 150-mL portions of ethyl acetate. The organic layers are combined, washed with two 250-mL portions of saturated sodium bicarbonate, dried over magnesium sulfate, filtered, and concentrated to give 41.117 g of yellow solid in an orange oil. Recrystallization from
25 65 mL of methanol yields 20.664 g of the title product as light yellow crystals.

Physical characteristics are as follows:

¹H NMR (CDCl₃) δ 8.85, 8.43, 8.33, 7.70, 2.72, 1.36-1.31, 1.20-1.14 ppm.

PREPARATION 2 Cyclopropyl-(3-aminophenyl)methanone (Formula A-3) Refer to Chart A.

30 A 500-mL Parr hydrogenation flask is charged with 2.1 g of 10% platinum on carbon and a solution of the title product of Preparation 1 (20.6 g) in 250 mL of methanol. The reaction mixture is shaken for 50 min under 44 psi of hydrogen, then filtered through Celite* twice. The light green solution is then concentrated to give
15.744 g of the title product as a green oil.

35 Physical characteristics are as follows:

*Trade-mark

^1H NMR (CDCl_3) δ 7.42, 7.30-7.23, 6.88, 3.83, 2.63, 1.24-1.19, 1.05-0.99 ppm.

PREPARATION 3 N-[3-cyclopropylmethanone]benzenesulfonamide (Formula A-4)

Refer to Chart A.

A 500-mL, three-necked, round-bottomed flask with a nitrogen inlet is
5 charged with the title product of Preparation 2 (15.7 g) and 200 mL of methylene
chloride. Benzenesulfonyl chloride (12 mL) and pyridine (7.8 mL) are added, and
the reaction mixture is stirred at room temperature for 45 min. 10% HCl (200 mL) is
added to quench the reaction. The organic layer is separated, dried over magnesium
sulfate, filtered, and concentrated to give 28.638 g of orange solid. Recrystallization
10 from 75 mL of hot methylene chloride yields the title product (22.264 g) as a pink
solid.

Physical characteristics are as follows:

MP 98-101 °C.

^1H NMR (CDCl_3) δ 7.81-7.73, 7.62, 7.55-7.35, 2.60, 1.30-1.25, 1.10-1.03 ppm.

15 ^{13}C NMR (CDCl_3) δ 200.4, 138.8, 137.2, 133.0, 129.5, 129.0, 127.0, 125.1,
124.7, 120.5, 17.3, 12.1 ppm.

IR (mineral oil) 3239, 3222, 1653, 1449, 1339, 1259, 1176, 1165, 1093, 939,
687 cm^{-1} .

Elemental analysis, found: C, 63.70; H, 5.01; N, 4.78.

20 MS (EI) m/e 301, 260, 160, 141, 77.

For high resolution, found: 301.0772.

PREPARATION 4 N-[3-cyclopropylmethanol]benzenesulfonamide (Formula A-5)

Refer to Chart A.

A 500-mL, three-necked, round-bottomed flask with a nitrogen inlet is
25 charged with the title compound of Preparation 3 (21.133 g), 200 mL of
tetrahydrofuran, and 100 mL of ethanol. The flask is cooled to 0 °C in an ice bath,
and sodium borohydride (10.6 g) is added in small portions over 20 minutes. The
reaction mixture is stirred at room temperature for ca. 18 h, and then cooled again
in an ice bath to 0 °C. 10% HCl (100 mL) is added dropwise over 45 min, and the
30 mixture is stirred another 1 h at 0 °C. The reaction mixture is then extracted with
three 100-mL portions of methylene chloride. The organic layers are combined,
dried over magnesium sulfate, filtered and concentrated to give 25.015 g of pale
yellow oil. Column chromatography on 150 g of silica gel (elution with 50-65% ether
in hexane followed by 2-5% methanol in methylene chloride) yields 18.692 g of the
35 title product as a white solid.

Physical characteristics are as follows:

MP 112-114 °C.

^1H NMR (CDCl_3) δ 7.69, 7.42, 7.32, 7.25, 7.12, 7.05-6.96, 3.82, 3.19, 1.03-0.94, 0.51-0.46, 0.39-0.29, 0.19-0.16 ppm.

5 ^{13}C NMR (DMSO) δ 147.0, 139.7, 137.4, 132.9, 129.3, 128.6, 126.8, 121.8, 118.5, 117.8, 75.0, 19.2, 3.1, 2.3 ppm.

IR (mineral oil) 3523, 3249, 1449, 732 cm^{-1} .

Elemental analysis, found: C, 63.41; H, 5.79; N, 4.86.

MS (EI) m/e 303, 275, 262, 77.

10 For high resolution, found: 303.0935.

PREPARATION 5 4-Hydroxy-10-propyl-2H-cycloocta[b]pyran-2-one (Formula A-7)

Refer to Chart A.

A 250-mL, three-necked, round-bottomed flask with a nitrogen inlet and a 125-mL pressure-equalizing addition funnel is charged with diisopropyl amine (3.6 mL) and 15 mL of tetrahydrofuran. The addition funnel is charged with 4-hydroxy-2H-cycloocta[b]pyran-2-one of formula A-6 (2.292 g) and 35 mL of tetrahydrofuran. The flask is cooled to 0 °C in an ice bath, *n*-butyllithium (16.3 mL of 1.6 M solution in hexanes) is added dropwise over 3 min, and the reaction mixture is stirred another 15 min at 0 °C. The solution of 4-hydroxy-2H-cycloocta[b]pyran-2-one in THF is added dropwise over 35 min, and the reaction mixture is stirred for another 25 min at 0 °C. Hexamethylphosphoramide (4 mL) is added in one portion, and iodopropane (1.3 mL) is added dropwise over 2 min. The reaction mixture is allowed to warm to room temperature and stirred for ca. 18 h. 30 mL of 10% HCl is added and the aqueous layer is separated. The pH of the aqueous layer is lowered from 10 to 2 with concentrated HCl, and the aqueous layer is extracted with two 50-mL portions of methylene chloride. The organic layers are combined, dried over magnesium sulfate, filtered, and concentrated to give an orange oil, which is partitioned between 100 mL of 1 N sodium hydroxide and 50 mL of ether. The aqueous layer pH is adjusted from 14 to 1 with concentrated hydrochloric acid, and is then extracted with two 50-mL portions of methylene chloride. The organic layers are then combined, dried over magnesium sulfate, and concentrated to give an orange oil, which is diluted with 100 mL of ether and washed with three 25-mL portions of 10% HCl. The organic layer is then dried over magnesium sulfate, filtered, and concentrated to give 1.829 g of orange solid. Column chromatography on 100 g of silica gel (elution with 0-10% methanol in methylene chloride) gives

2187523

1.358 g of a pale orange solid. An additional column chromatography on 150 g of silica gel (elution with 10% ether and 1% acetic acid in methylene chloride) gives 0.705 g of the title product as a yellow solid.

Physical characteristics are as follows:

5 ^1H NMR (CDCl_3) δ 11.38, 5.68, 3.02-2.93, 2.20, 1.98-1.82, 1.73-1.58, 1.46-1.25, 1.24-1.08, 0.89 ppm.

^{13}C NMR (CDCl_3) δ 172.3, 168.3, 165.3, 114.8, 89.7, 38.6, 36.0, 33.3, 30.1, 27.2, 25.5, 22.9, 21.0, 13.9 ppm.

IR (mineral oil) 1679, 1641, 1617, 1492 cm^{-1} .

10 Elemental analysis, found: C, 70.90; H, 8.36.

MS (EI) m/e 236, 208, 166.

For high resolution, found: 236.1414.

EXAMPLE 1 N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-10-propyl-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]benzenesulfonamide (Formula A-

15 8) Refer to Chart A.

A 100-mL, three-necked, round-bottomed flask with a 35-mL pressure-equalizing addition funnel filled with 3 A molecular sieves and fitted with a reflux condenser and a nitrogen inlet is charged with the title compound of Preparation 5 (0.196 g), *p*-toluenesulfonic acid (0.040 g), and 30 mL of methylene chloride. The title product of Preparation 4 (0.252 g) is added, and the reaction mixture is heated to reflux for 2 h, then stirred at room temperature for an additional hour. The reaction mixture is then diluted with 20 mL of methylene chloride and washed with 60 mL of 1:1 saturated sodium bicarbonate and brine, 30 mL of water, and 30 mL of brine. The aqueous layers are combined and extracted with 30 mL of methylene chloride. The organic layers are then combined, dried over magnesium sulfate, filtered, and concentrated to give 0.576 g of crude material. Column chromatography on 35 g of silica gel (elution with 20-80% ether in hexane) yields 0.096 g of the title compound as a white solid.

Physical characteristics are as follows:

30 MP 87-90 $^{\circ}\text{C}$ (decomposition).

MS (EI) m/e 521, 493, 380, 275, 262, 249, 144, 77.

For high resolution, found: 521.2236.

EXAMPLES 2 - 7

Following procedures analogous to those described above, the following additional compounds of the present invention are prepared:

- 2) 4-Cyano-N-[3-(R or S)-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-(R or S)-10-propyl-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]benzenesulfonamide
- 3) 4-Cyano-N-[3-(R or S)-[cyclopropyl(5,6,7,8,9,10-hexahydro-(R or S)-10-cyclopropylmethyl-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]benzenesulfonamide
- 5) N-[3-(R or S)-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-(R or S)-10-propyl-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide
- 6) N-[3-(R or S)-[cyclopropyl(5,6,7,8,9,10-hexahydro-(R or S)-10-cyclopropylmethyl-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide
- 7) N-[3-(R or S)-[cyclopropyl(5,6,7,8,9,10-hexahydro-(R or S)-10-benzyl-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide

PREPARATION 6 (3-Benzaldehyde)-carbamic acid, phenylmethyl ester (Formula B-2) Refer to Chart B.

A flask with a nitrogen inlet is charged with sodium bicarbonate (10.4 g) in 200 mL of THF and 200 mL of water, and *m*-aminobenzaldehyde of formula B-1 (10.0 g) and benzyl chloroformate (13.6 mL) are added sequentially. The mixture is stirred at room temperature for 40 min. Ether is then added, and the organic layer is separated, washed with saturated sodium bicarbonate, dried over sodium sulfate, filtered and concentrated to give a brown oil. Column chromatography on 300 g of silica gel yields 16.3 g of the title compound as a pale yellow oil. An analytical sample is crystallized from ethyl acetate-hexane.

Physical characteristics are as follows:

MP 100-104 °C.

¹H NMR (CDCl₃) δ 9.98, 7.91, 7.69, 7.59, 7.43-7.35, 6.83, 5.23 ppm.

¹³C NMR (CDCl₃) δ 191.8, 153.0, 138.6, 137.1, 135.6, 129.7, 128.6, 128.4, 128.3, 124.6, 124.2, 119.1, 67.2 ppm.

IR (mineral oil) 3269, 2954, 2925, 2868, 2855, 1729, 1682, 1597, 1560, 1465, 1455, 1326, 1294, 1237, 1229, 1170, 1155, 1048, 695 cm⁻¹.

Elemental analysis, found: C, 70.74; H, 5.14; N, 5.33.

MS (EI) m/e 255, 211, 91.

For high resolution, found 255.0900.

PREPARATION 7 [3-(1-Hydroxy-3-methylbutyl)phenyl]-carbamic acid, phenylmethyl ester (Formula B-3 wherein R₁ is isobutyl) Refer to Chart B.

5 A flask with a nitrogen inlet is charged with the title compound of Preparation 6 (4.0 g) and 60 mL of dry tetrahydrofuran. The mixture is cooled to 0 °C, and isobutyl magnesium chloride (17.2 mL) is added. The reaction mixture is then allowed to warm to room temperature and stir for 2 hours. Saturated ammonium chloride is added to quench the reaction, and the mixture is partitioned
10 between ether and water. The organic layer is washed with water and concentrated to give 5.78 g of pale yellow oil. The crude material is crystallized from ethyl acetate-hexane to yield 4.13 g of the title compound as white crystals.

Physical characteristics are as follows:

MP 73-77°C.

15 ¹H NMR (CDCl₃) δ 7.41-7.33, 7.25, 7.05, 6.74, 5.19, 4.73-4.65, 1.91, 1.73-1.65, 1.47, 0.93 ppm.

IR (Nujol) 3400, 3249, 3085, 2953, 2925, 2869, 2855, 1697, 1615, 1602, 1563, 1450, 1283, 1245, 1177, 1067, 1017, 798, 773, 740, 696 cm⁻¹.

Elemental analysis, found: C, 72.58; H, 7.25; N, 4.55.

20 MS (EI) m/z 313, 257, 213, 91.

PREPARATION 8 [3-[1-(5,6,7,8,9,10-Hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)-3-methylbutyl]phenyl]-carbamic acid, phenylmethyl ester (Formula B-5 wherein R₁ is isobutyl) Refer to Chart B.

25 A 200-mL, three-necked flask with a Dien-Stark trap and a nitrogen inlet is charged with *p*-toluenesulfonic acid (0.66 g) and toluene (100 mL) and warmed to reflux to collect 20 mL in the Dien-Stark trap. The reaction mixture is cooled to room temperature, and the trap is emptied. 4-Hydroxy-2H-cycloocta[b]pyran-2-one of formula B-4 (2.48 g) and the title compound of Preparation 7 (4.0 g) are added to
30 the reaction mixture and then heated to reflux for 6.5 h. The reaction mixture is allowed to stand at room temperature overnight, then poured into 350 mL of ethyl acetate, washed with two 25-mL portions of water, 25 mL of saturated sodium bicarbonate, and 25 mL of water. The organic layer is concentrated to give 7.9 g of yellow oil. Column chromatography on 150 g of silica gel (elution with 10-50% ethyl
35 acetate in hexane) gives 0.217 g of the title product as an off-white foam.

Physical characteristics are as follows:

MP 73-78°C (decomposition).

¹H NMR (CDCl₃) δ 7.38-7.25, 7.13, 6.72, 6.01, 5.19, 4.48, 2.58, 2.41, 1.93, 1.74, 1.62-1.33, 0.96 ppm.

5 PREPARATION 9 (R or S)-[3-[1-(5,6,7,8,9,10-Hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)-3-methylbutyl]phenyl]-carbamic acid, phenylmethyl ester (Formula B-5 wherein R₁ is isobutyl) Refer to Chart B.

A stock solution of the title compound of Preparation 8 (32 mg/mL) in 30% isopropyl alcohol and 0.1% acetic acid in hexane is chromatographed on a 2.0 x 25
10 cm (R, R) ^{*}Whelk-O 1 column at 2 mL per injection using an automated chromatographic system. The eluant is monitored at 310 nm, the flow rate was 10 mL/min and appropriate fractions from multiple injections combined and concentrated in vacuo to give snowy white solids.

15 Physical characteristics are as follows:

The retention time of the title compound is 18.8 min.

PREPARATION 10 (R or S)-[3-[1-(5,6,7,8,9,10-Hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)-3-methylbutyl]phenyl]-carbamic acid, phenylmethyl ester (Formula B-5 wherein R₁ is isobutyl) Refer
20 to Chart B.

The title compound of Preparation 8 is separated as described in Preparation 9 above.

Physical characteristics are as follows:

The retention time of the title compound is 22.1 min.

25 PREPARATION 11 (R or S)-3-[1-(3-Aminophenyl)-3-methylbutyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyranone (Formula B-6 wherein R₁ is isobutyl) Refer to Chart B.

A flask with a nitrogen inlet is charged with a solution of the title compound of Preparation 9 (0.637 g) in 6 mL of ethanol. Cyclohexene (6 mL) and 10%
30 palladium on carbon (0.16 g) are added, and the reaction mixture is heated at reflux for 2 h. The mixture is then filtered through Celite and concentrated to give 0.205 g of the title compound as an off-white foam.

Physical characteristics are as follows:

MP 158-162 °C

35 MS (EI) m/z 355, 312, 299, 161, 106

*Trade-mark

For high resolution, found: 355.2144.

EXAMPLE 8 (R or S)-N-[3-1-(5,6,7,8,9,10-Hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)-3-methylbutyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula B-7 wherein R₁ is isobutyl and R₂ is 1-methylimidazole) Refer to Chart B.

A flask with a nitrogen inlet is charged with the title compound of Preparation 11 (0.095 g), 1-methylimidazole-4-sulfonyl chloride (0.048 g), and 5 mL of methylene chloride (CH₂Cl₂). Pyridine (0.53 mL) is added, and the reaction mixture is stirred at room temperature for ca. 18 h. A precipitate forms, which is filtered to give 0.097 g of a white solid. Recrystallization from methanol-chloroform yields 0.065 g of the title compound as a white powder.

Physical characteristics are as follows:

MP 207-210 °C

¹H NMR (CDCl₃) δ 10.4, 10.0, 7.70, 7.11, 7.05, 6.92, 4.21, 3.64, 2.54, 2.16, 1.62, 1.53, 1.43, 1.34, 0.85 ppm.

MS (EI) m/z 499, 456, 443, 306, 251, 160, 145

For high resolution, found: 499.2151

PREPARATION 12 (R or S)-3-[1-(3-Aminophenyl)-3-methylbutyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyranone (Formula B-6 wherein R₁ is isobutyl) Refer to Chart B.

Following the general procedure of Preparation 11, and making non-critical variations, but substituting the title product of Preparation 10 for the title product of Preparation 9, 0.189 g of the title compound is obtained as a grey solid.

Physical characteristics are as follows:

MS (EI) m/z 355, 312, 299, 161

For high resolution, found: 355.2135

EXAMPLE 9 (R or S)-N-[3-1-(5,6,7,8,9,10-Hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)-3-methylbutyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula B-7 wherein R₁ is isobutyl and R₂ is 1-methylimidazole) Refer to Chart B.

Following the general procedure of Example 8, and making non-critical variations, but substituting the title product of Preparation 12 for the title product of Preparation 11, 0.047 g of the title compound is obtained as a white solid.

Physical characteristics are as follows:

¹H NMR (CDCl₃) δ 10.45, 10.06, 7.70, 7.11, 7.05, 6.94, 4.21, 3.64, 2.55, 2.16,

2187523

1.62, 1.53, 1.42, 1.35, 0.86 ppm.

MS (EI) m/z 499, 456, 443, 354, 306, 160, 145

For high resolution, found: 499.2146

PREPARATION 13 [3-(Cyclopropyl-hydroxymethyl)-phenyl]-methanol (Formula C-2)

5 Refer to Chart C.

To a solution of 6.5 mL of 3-bromobenzylalcohol of formula C-1 in 900 mL of tetrahydrofuran under nitrogen at -78°C is added 46 mL of a 1.4 M solution of methyllithium in diethyl ether. The solution is stirred for 20 min and then 66 mL of a 1.6 M solution of n-butyllithium in hexane is added. The solution is stirred 25
10 min and then 6 mL of cyclopropanecarboxaldehyde is added. The solution is stirred 1.5 h, warmed to 0°C and stirred for 40 min. Next the solution is warmed to room temperature and stirred for 30 min. Finally the solution is heated at reflux for 1h. The solution is poured onto 800 mL of water and acidified with concentrated HCl followed by 5% aqueous HCl to adjust the pH to approximately 6. The layers are
15 separated and the aqueous extracted with two portions of ethyl acetate. The combined organics are dried (Na₂SO₄) and concentrated to afford a yellow oil which is chromatographed over 900 g 230-400 mesh silica gel (2:1 ethyl acetate: hexane) to afford a 6.61g (68%) of the desired alcohol as a yellow oil.

Physical characteristics are as follows:

20 ¹H NMR (CDCl₃) δ 7.41-7.26, 4.67, 3.99-3.96, 2.18, 1.28-1.14, 0.68

PREPARATION 14 3-[cyclopropyl [3-[hydroxymethyl]phenyl]methyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one (Formula C-3)

Refer to Chart C.

To a solution of 501 mg of the title product of Preparation 13 in 50 mL of
25 dichloromethane in the presence of molecular sieves 3A under nitrogen is added 492 mg of 4-hydroxy-5,6,7,8,9,10-hexahydrocycloocta[b]pyran-2-one followed by 49 mg of p-toluenesulfonic acid monohydrate. The solution is heated at reflux for 2 h and then an additional 105 mg p-toluenesulfonic acid monohydrate is added and heating continued for a further hour. The solution is concentrated in vacuo to afford a white
30 foam which is treated with water and then 1 N KOH and extracted with one portion of ethyl acetate. The organic layer is washed with one portion of 1 N KOH. The combined aqueous layers are acidified with 5% aqueous HCl and extracted with three portions of ethyl acetate. The combined organics are dried (Na₂SO₄) and concentrated in vacuo to afford a yellow oil which is chromatographed over 180 g of
35 230-400 mesh silica gel (2:1 ethyl acetate: hexane) to afford 436 mg of the desired

2187523

benzyl alcohol as a white foam.

Physical characteristics are as follows:

MP 65-70°C

¹H NMR (CDCl₃) δ 7.25- 7.03, 4.36, 3.70-3.67, 2.41-2.37, 2.24-2.23, 1.53-1.50,
5 1.35-1.05, 0.54-0.43, 0.42-0.21, 0.07-0.02.

PREPARATION 15 3-[Cyclopropyl [3-[bromomethyl]phenyl]methyl]-5,6,7,8,9,10-
hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one; and 3-
[cyclopropyl [3-[chloromethyl]phenyl]methyl]-5,6,7,8,9,10-
hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one (Formulas C-
10 4,5) Refer to Chart C.

To a solution of 1.01g of the title product of Preparation 14 in 70 mL of
dichloromethane under nitrogen at 0°C is added 2.00 g of triphenylphosphine and
2.58 g of carbon tetrabromide in sequence. The solution is stirred 1 h and then
poured onto brine. The layers are separated and the aqueous extracted with three
15 portions of ethyl acetate. The combined organics are dried (Na₂SO₄) and
concentrated to afford a yellow oil which is triturated with ether. The solid is
filtered off and the filtrate concentrated and chromatographed over 180 g of 230-400
mesh silica gel (1:1 hexane: ethyl acetate) to afford 374 mg of the desired title
product as a mixture of bromide and chloride. The solids isolated from the filtration
20 are chromatographed as above to afford an additional 699 mg of the title product as
a mixture of bromide and chloride.

Physical characteristics are as follows:

Mass Spectrum *m/e* 418, 416 (M⁺ for Br), 388, 374, 372 (M⁺ for Cl), 337, 246,
233, 220, 207, 195, 179, 153, 143, 129.

25 PREPARATION 16 3-[Cyclopropyl[3-[(phenylthio)methyl]phenyl]methyl]-
5,6,7,8,9,10-hexahydro-4-hydroxy-2H-Cycloocta[b]pyran-2-one
(Formula C-6) Refer to Chart C.

To a solution of 138 mg of the title products of Preparation 15 in 5 mL of
dichloromethane is added 0.04 mL of thiolphenol and 0.17 mL of diisopropylethyl-
30 amine in sequence. The solution is heated at reflux for 1h and then allowed to
stand at room temperature overnight. The solution is poured onto brine and treated
with 5% aqueous hydrochloric acid. The layers are separated and the aqueous
extracted with three portions of ethyl acetate. The combined organics are dried
(Na₂SO₄) and concentrated to afford a yellow oil which is chromatographed over 80
35 g of 230-400 mesh silica gel (2:1 hexane: ethyl acetate) to afford 111 mg of the

desired sulfide as a white foam.

Physical characteristics are as follows:

MP 137-139°C

Mass Spectrum m/e 446 (M^+), 418, 337, 295, 233, 220, 207, 185, 145, 128,
5 109, 91, 79, 55, 40.

EXAMPLE 10 3-[Cyclopropyl[3-[(phenylsulfonyl)methyl]phenyl]methyl]-
5,6,7,8,9,10-hexahydro-4-hydroxy-2H-Cycloocta[b]pyran-2-one
(Formula C-7) Refer to Chart C.

To a solution of 119 mg of the title product of Preparation 16 in 5 mL of
10 tetrahydrofuran and 5 mL of methanol at 0°C is added a solution of 279 mg of oxone
in 5 mL of water. The solution is stirred 2.5 h and then warmed to room
temperature and stirred 2 h. The solution is filtered and the solids washed with
chloroform. The filtrate is diluted with water and the layers are separated. The
aqueous is extracted with three portions of ethyl acetate. The combined organics are
15 dried (Na_2SO_4) and concentrated to afford a clear oil which is chromatographed over
80 g of 230-400 mesh silica gel (1:1 hexane: ethyl acetate) to afford 78 mg of the title
product as a white foam.

Physical characteristics are as follows:

MP 80-85°C

20 Mass Spectrum m/e 479 (M^++1), 463, 450, 391, 337, 309, 207, 161, 149, 127,
115, 71, 57, 41.

Exact mass found: 479.1885.

EXAMPLES 11-39

The following compounds of the present invention are prepared by an
25 analogous synthetic route to that described above:

11) 3-[cyclopropyl[3-[(4-cyanophenylsulfonyl)methyl]phenyl]methyl]-
5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one

The starting material, 4-cyanobenzenethiol, is prepared from 4-
cyanobenzenesulfonyl chloride according to a general literature procedure: Wagner,
30 A.W. Ber Deutsch Chem Ges, 99:375 (1966).

Physical characteristics are as follows:

MP 100-105°C

Mass Spectrum m/e 504 (M^++1), 337, 247, 207, 143.

Exact mass found 504.1843.

35 12) 3-[cyclopropyl[3-[(4-fluorophenylsulfonyl)methyl]phenyl]methyl]-

2187523

5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one

Physical characteristics are as follows:

MP 95-100°C

¹H NMR (CDCl₃) δ 7.61-7.57, 7.40-7.37, 7.27-7.20, 7.13-7.07, 7.02-6.99, 6.42, 4.30, 3.88-3.85, 2.64-2.61, 2.51-2.47, 1.83-1.40, 1.40-1.27, 0.69-0.58, 0.48-0.43, 0.19-0.14.

13) 3-[cyclopropyl[3-[(4-methylphenylsulfonyl)methyl]phenyl]methyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one

Physical characteristics are as follows:

MP 100-105°C

¹H NMR (CDCl₃) δ 7.37-7.34, 7.25-7.22, 7.17-7.05, 6.86-6.84, 4.15, 3.60-3.58, 2.52-2.42, 2.42-2.30, 2.28, 1.70-1.14, 0.57-0.32, 0.32-0.20, 0.06-(-)0.16.

14) 3-[cyclopropyl[3-[(4-carboxyphenylsulfonyl)methyl]phenyl]methyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one

Physical characteristics are as follows:

MP 90-95°C

Mass Spectrum *m/e* 523 (M⁺+1), 337, 247, 207, 143.

Exact mass found 523.1785.

15) 3-[cyclopropyl[3-[(2-(1-methylimidazolyl)sulfonyl)methyl]phenyl]methyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one

Physical characteristics are as follows:

MP 95-103°C

¹H NMR (CDCl₃) δ 7.36-7.34, 7.29-7.27, 7.14, 7.06-7.03, 6.98 (s, 1H), 6.86, 4.30, 3.73-3.70, 3.20, 2.67-2.54, 1.90-1.36, 0.71-0.50, 0.46-0.33, 0.18-0.03.

16) 3-[cyclopropyl[3-[(2-pyrimidinylsulfonyl)methyl]phenyl]methyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one

17) 3-[cyclopropyl[3-[(2-pyridinylsulfonyl)methyl]phenyl]methyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one

18) 3-[cyclopropyl[3-[(1-methyl-4-imidazolylsulfonyl)methyl]phenyl]methyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one

19) 3-[cyclopropyl[3-[(5-cyano-2-pyridinylsulfonyl)methyl]phenyl]methyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one

20) 3-[cyclopropyl[3-[(2-benzimidazolylsulfonyl)methyl]phenyl]methyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one

21) 3-[cyclopropyl[3-[(2-quinolinylsulfonyl)methyl]phenyl]methyl]-

2187523

- 5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one
- 22) 3-[cyclopropyl[3-[(4-cyanophenylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-coumarin
- 23) 3-[cyclopropyl[3-[(2-pyridinylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-coumarin
- 24) 3-[cyclopropyl[3-[(1-methyl-4-imidazolylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-coumarin
- 25) 3-[cyclopropyl[3-[(5-cyano-2-pyridinylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-coumarin
- 26) 3-[cyclopropyl[3-[(2-benzimidazolylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-coumarin
- 27) 3-[cyclopropyl[3-[(2-quinolinylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-coumarin
- 28) 3-[cyclopropyl[3-[(4-cyanophenylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-6-[1-(phenylmethyl)propyl]-2H-pyran-2-one
- 29) 3-[cyclopropyl[3-[(2-pyridinylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-6-[1-(phenylmethyl)propyl]-2H-pyran-2-one
- 30) 3-[cyclopropyl[3-[(1-methyl-4-imidazolylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-6-[1-(phenylmethyl)propyl]-2H-pyran-2-one
- 31) 3-[cyclopropyl[3-[(5-cyano-2-pyridinylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-6-[1-(phenylmethyl)propyl]-2H-pyran-2-one
- 32) 3-[cyclopropyl[3-[(2-benzimidazolylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-6-[1-(phenylmethyl)propyl]-2H-pyran-2-one
- 33) 3-[cyclopropyl[3-[(2-quinolinylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-6-[1-(phenylmethyl)propyl]-2H-pyran-2-one
- 34) 3-[cyclopropyl[3-[(4-cyanophenylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-6-(2-phenylethyl)-6-(1-propyl)-5,6-dihydro-2H-pyran-2-one
- 35) 3-[cyclopropyl[3-[(2-pyridinylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-6-(2-phenylethyl)-6-(1-propyl)-5,6-dihydro-2H-pyran-2-one
- 36) 3-[cyclopropyl[3-[(1-methyl-4-imidazolylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-6-(2-phenylethyl)-6-(1-propyl)-5,6-dihydro-2H-pyran-2-one
- 37) 3-[cyclopropyl[3-[(5-cyano-2-pyridinylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-6-(2-phenylethyl)-6-(1-propyl)-5,6-dihydro-2H-pyran-2-one
- 38) 3-[cyclopropyl[3-[(2-benzimidazolylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-6-(2-phenylethyl)-6-(1-propyl)-5,6-dihydro-2H-pyran-2-one

39) 3-[cyclopropyl[3-[(2-quinolinylsulfonyl)methyl]phenyl]methyl]-4-hydroxy-6-(2-phenylethyl)-6-(1-propyl)-5,6-dihydro-2H-pyran-2-one

PREPARATION 17 5,6-Dihydro-4-Hydroxy-6-phenethyl-6-propyl-2H-pyran-2-one

(Formula D-1: R₁ is phenethyl, R₂ is propyl) Refer to Chart D.

5 Methyl acetoacetate (1.47 mL) is added to a suspension of sodium hydride (567 mg, 60% dispersion in mineral oil) in THF (30 mL) at 0 °C. After 15 minutes, *n*-butyl lithium (8.5 mL, 1.6 M solution in hexane) is added dropwise and the reaction is stirred 15 minutes. 1-Phenyl-3-hexanone (2.0g) is then added via syringe all at once to the reaction mixture. The reaction is stirred an additional hour, then
10 poured into a saturated ammonium chloride solution. It is extracted with EtOAc, dried over anhydrous sodium sulfate and evaporated *in vacuo*. The material obtained is dissolved in THF (25 mL) and a 0.1 N sodium hydroxide (113 mL) solution is added. After stirring three hours, the mixture is extracted with ethyl acetate (1 X). The aqueous layer is adjusted to pH 3 with hydrochloric acid, then
15 extracted with CH₂Cl₂ (3 x 25 mL), dried over anhydrous magnesium sulfate and evaporated to afford the title product as a white solid.

Physical characteristics are as follows:

¹H NMR (300 MHz, CDCl₃): δ 0.96, 1.21, 1.48, 1.72, 1.98, 2.73, 3.43, 7.15-7.32.

20 Anal. Found: C, 73.77; H, 7.96.

PREPARATION 18 4-Hydroxy -3-[1-(3-nitrophenyl)-propyl]-5,6-dihydro-6-phenethyl-6-propyl-2H-pyran-2-one (Formula D-4: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl) Refer to Chart D.

To a solution of the title product of Preparation 17 (Formula D-1: R₁ is phenethyl, R₂ is propyl) (1 g) and 3-nitrobenzaldehyde (Formula D-2) (581 mg) in dry THF at 0°C is added AlCl₃ (1.0 g) as one solid portion. The cooling bath is removed and the yellow solution is allowed to stir at room temperature for 2 hrs. The reaction mixture is quenched by the addition of solid Na₂CO₃·10H₂O (2.2 g) and vigorously stirred for 5 min. The mixture is filtered through celite with ether and
30 the filtrate is evaporated to dryness *in vacuo*. The benzylidene intermediate of formula D-3 and CuBr·Me₂S (237 mg) are dissolved in dry THF and a solution of Et₃Al (4.23 mL; 1 M in hexane) is added at room temperature, dropwise over 5 min. When the reaction is complete (as determined by tlc), it is quenched by the addition of water and the reaction mixture is transferred to a separatory funnel with ether.
35 The aqueous layer is extracted with ether (3 x 15 mL) and the combined organic

layers are washed with brine, dried (MgSO_4), filtered and evaporated *in vacuo* to provide an oil. Flash chromatography on silica gel with Hexanes/EtOAc (3:1) provides 1.1 g of the title product as a light yellow foam.

Physical characteristics are as follows:

5 ^1H NMR complicated by presence of diastereomers.

^1H NMR (300 MHz, CD_3OD) δ 0.93, 1.37, 1.74, 1.82-2.14, 2.29, 2.52-2.71, 4.19, 6.98-7.24, 7.44, 7.72, 8.02, 8.26.

10 PREPARATION 19 3-[Cyclopropyl-(3-nitrophenyl)-methyl]-4-hydroxy-5,6-dihydro-6-phenethyl-6-propyl-2H-pyran-2-one (Formula D-4: R_1 is phenethyl, R_2 is propyl, R_3 is cyclopropyl) Refer to Chart D.

To a solution of the title product of Preparation 17 (Formula D-1: R_1 is phenethyl, R_2 is propyl) (1 g) and 3-nitrobenzaldehyde (Formula D-2) (581 mg) in dry THF at 0°C is added AlCl_3 (1.0 g) as one solid portion. The cooling bath is removed and the yellow solution is allowed to stir at room temperature for 2 hrs.

15 The reaction mixture is quenched by the addition of solid $\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$ (2.2 g) and vigorously stirred for 5 min. The mixture is filtered through celite with ether and the filtrate is evaporated to dryness *in vacuo*. The benzylidene intermediate of formula D-3 and $\text{CuBr} \cdot \text{Me}_2\text{S}$ (237 mg) are dissolved in dry THF and cooled to -78°C . A solution of cyclopropylmagnesium bromide (15.6 mL; 0.25 M in THF) is added

20 dropwise over 10 min and the reaction mixture is stirred for 30 min. The reaction is quenched by the addition of water and neutralized by the addition of 1N HCl. The reaction mixture is transferred to a separatory funnel with ether and the aqueous layer is extracted with ether (3 x 15 mL). The combined organic layers are washed with brine, dried (MgSO_4), filtered and evaporated *in vacuo* to provide an oil. Flash

25 chromatography on silica gel with Hexanes/EtOAc (3:1) provides 0.9 g of the title product as a light yellow foam.

Physical characteristics are as follows:

^1H NMR complicated by presence of diastereomers.

30 ^1H NMR (300 MHz, CD_3OD) δ 0.25, 0.53, 0.74, 0.94, 1.41, 1.68-2.13, 2.57-2.72, 3.38, 7.04-7.23, 7.46, 7.82, 8.03, 8.30.

PREPARATION 20 3-[1-(3-Aminophenyl)-propyl]-4-hydroxy -5,6-dihydro-6-phenethyl-6-propyl-2H-pyran-2-one (Formula D-5: R_1 is phenethyl, R_2 is propyl, R_3 is ethyl) Refer to Chart D.

To a solution of the title product of Preparation 18 (Formula D-4: R_1 is phenethyl, R_2 is propyl, R_3 is ethyl) (350 mg) in MeOH at room temperature is

35

2187523

added 10% Pd/C (35 mg) and ammonium formate (521 mg). The resulting mixture is stirred for 2 hrs. and then filtered through celite with CH₂Cl₂. The filtrate is evaporated *in vacuo* and the residue is triturated with CH₂Cl₂ (3 x 10 mL). The combined organic solution is filtered and evaporated *in vacuo* to provide the 325 mg of the title compound as a light yellow foam.

Physical characteristics are as follows:

¹H NMR complicated by presence of diastereomers.

¹H NMR (300 MHz, CDCl₃) δ 0.89, 1.40, 1.64-2.07, 2.20, 2.62, 3.94, 6.54, 6.72-7.25.

The compounds of formula D-5, wherein R₁ is propyl, R₂ is propyl and R₃ is ethyl or t-butyl are prepared by analogous procedures.

Physical characteristics of the compound of the formula D-5, wherein R₁ is propyl and R₂ are propyl and R₃ is ethyl, are as follows:

¹H NMR: 0.9, 1.3, 1.5-1.8, 2.0, 2.2, 2.5, 3.9, 4.5, 6.5, 6.8, 7.0 ppm

TLC R_f: 0.32 (10% ethyl acetate in dichloromethane).

EXAMPLE 40 4-Cyano-N-[3-[1-(4-hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-benzenesulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 4-cyanophenyl) Refer to Chart D.

To a solution of the title product of Preparation 20 (Formula D-5: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl) (30 mg) and 4-cyanobenzenesulfonyl chloride of formula D-7, wherein R₄ is cyanophenyl, (16.1 mg) in CH₂Cl₂ (1 mL) at room temperature is added pyridine (13 μL) via syringe. The resulting solution is stirred for 3 hrs, after which the starting amine is consumed. The mixture is flash chromatographed on silica gel with the 5% EtOAc in CH₂Cl₂ to provide 21 mg of the title product as a white foam.

Physical characteristics are as follows:

¹H NMR complicated by presence of diastereomers.

¹H NMR (300 MHz, CDCl₃) δ 0.6-1.1, 1.2-2.2, 2.4-2.7, 3.86-4.01, 6.89-7.45, 7.66-7.92.

HRMS found: 559.2267.

EXAMPLE 41 N-[3-[1-(4-hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.

2187523

The title compound is prepared from the amine of Preparation 20 (Formula D-5; R₁ is phenethyl, R₂ is propyl, R₃ is ethyl) and 1-methyl-imidazole-4-sulfonyl chloride of formula D-7 wherein R₄ is 1-methylimidazol-4-yl, using the general procedure for sulfonylation of Example 40 to yield the title compound as an off-white amorphous solid after flash chromatography with 4% MeOH/EtOAc.

Physical characteristics are as follows:

¹H NMR complicated by presence of diastereomers and tautomerism.

¹H NMR (300 MHz, CDCl₃) δ 0.75-0.96, 1.17-1.43, 1.45-2.11, 2.43-2.68, 3.24, 3.64, 3.94, 6.72-7.51.

HRMS found: 538.2383.

EXAMPLE 42 N-[3-[1-(4-hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-8-quinolinesulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is quinolin-8-yl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula D-5: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl) and 8-quinolinesulfonyl chloride of formula D-7 wherein R₄ is quinolin-8-yl, using the general procedure for sulfonylation of Example 40 to yield the title compound as a white amorphous solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

¹H NMR complicated by presence of diastereomers and tautomerism.

¹H NMR (300 MHz, CDCl₃) δ 0.66, 0.90, 1.17-1.44, 1.58-2.03, 2.38-2.64, 3.77, 6.68-7.27, 7.35-7.69, 8.02, 8.26, 9.14.

HRMS found: 585.2402.

EXAMPLE 43 N-[3-[1-(4-hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-8-quinolinesulfonamide (Formula D-6: R₁ is propyl, R₂ is propyl, R₃ is ethyl, R₄ is quinolin-8-yl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula D-5: R₁ is propyl, R₂ is propyl, R₃ is ethyl) and 8-quinolinesulfonyl chloride of formula D-7 wherein R₄ is quinolin-8-yl, using the general procedure for sulfonylation of Example 40 to yield the title compound as a white amorphous solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

¹H NMR complicated by presence of diastereomers and tautomerism.

¹H NMR (300 MHz, CD₃OD) δ 0.67, 0.85, 1.27, 1.54, 2.01, 3.73, 6.78, 6.90,

2187523

7.04, 7.57, 8.12, 8.29, 8.38, 9.13.

HRMS found: 523.2276.

Anal. found: C, 66.09; H, 6.60; N, 5.13.

EXAMPLE 44 N-[3-[1-(4-hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1-methyl-1H-imidazolesulfonamide (Formula D-6: R₁ is propyl, R₂ is propyl, R₃ is ethyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula D-5: R₁ is propyl, R₂ is propyl, R₃ is ethyl) and 1-methyl-imidazole-4-sulfonyl chloride of formula D-7 wherein R₄ is 1-methylimidazol-4-yl, using the general procedure for sulfonylation of Example 40 to yield the title compound as an off-white amorphous solid after flash chromatography with 4% MeOH/EtOAc.

Physical characteristics are as follows:

¹H NMR (300 MHz, CD₃OD) δ 0.88, 1.32, 1.64, 1.93, 2.16, 2.56, 3.68, 3.91, 6.87, 7.03, 7.14, 7.53, 7.64.

HRMS found: 476.2223.

EXAMPLE 45 4-Fluoro-N-[3-[1-(4-hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-benzenesulfonamide (Formula D-6: R₁ is propyl, R₂ is propyl, R₃ is ethyl, R₄ is 4-fluorophenyl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula D-5: R₁ is propyl, R₂ is propyl, R₃ is ethyl) and 4-fluoro-benzenesulfonyl chloride of formula D-7 wherein R₄ is 4-fluorophenyl, using the general procedure for sulfonylation of Example 40 to yield the title compound as a white amorphous solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

¹H NMR complicated by presence of diastereomers and tautomerism.

¹H NMR (300 MHz, CDCl₃) δ 0.51-1.03, 1.15-1.73, 1.81-2.48, 2.73, 3.91, 6.69, 6.88, 7.09, 7.78.

HRMS found: 490.2085.

EXAMPLE 46 4-Cyano-N-[3-[1-(4-hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-benzenesulfonamide (Formula D-6: R₁ is propyl, R₂ is propyl, R₃ is ethyl, R₄ is 4-cyanophenyl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula

D-5: R₁ is propyl, R₂ is propyl, R₃ is ethyl) and 4-cyano-benzenesulfonyl chloride of formula D-7 wherein R₄ is 4-cyanophenyl, using the general procedure for sulfonylation of Example 40 to yield the title compound as a white amorphous solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

5 Physical characteristics are as follows:

¹H NMR complicated by presence of diastereomers and tautomerism.

¹H NMR (300 MHz, CDCl₃) δ 0.68-0.96, 1.15-1.42, 1.44-1.76, 1.83-2.12, 3.18, 3.88, 6.69-7.18, 7.71, 7.85.

HRMS found: 497.2126.

10 **EXAMPLE 47** N-[3-[1-(4-hydroxy-6,6-diisobutyl-2-oxo-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide
(Formula D-6: R₁ is isobutyl, R₂ is isobutyl, R₃ is ethyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.

15 The title compound is prepared from the amine of Preparation 20 (Formula D-5: R₁ is propyl, R₂ is isobutyl, R₃ is ethyl) and 1-methyl-imidazole-4-sulfonyl chloride of formula D-7 wherein R₄ is 1-methylimidazol-4-yl, using the general procedure for sulfonylation of Example 40 to yield the title compound as an off-white amorphous solid after flash chromatography with 4% MeOH/EtOAc.

Physical characteristics are as follows:

20 ¹H NMR (300 MHz, CD₃OD) δ 0.82-0.94, 1.52-1.83, 1.86-2.03, 2.06-2.22, 2.60, 3.68, 3.92, 6.87, 7.03, 7.16, 7.56, 7.65.

HRMS found: 504.2531.

Anal. found: C, 62.03; H, 7.43; N, 8.20.

25 **EXAMPLE 48** N-[3-[1-(4-hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-pyran-3-yl)-cyclopropylmethyl]-phenyl]-1-methyl-1H-imidazolesulfonamide (Formula D-6: R₁ is propyl, R₂ is propyl, R₃ is cyclopropyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.

30 The title compound is prepared from the amine of Preparation 20 (Formula D-5: R₁ is propyl, R₂ is propyl, R₃ is cyclopropyl) and 1-methyl-imidazole-4-sulfonyl chloride of formula D-7 wherein R₄ is 1-methylimidazol-4-yl, using the general procedure for sulfonylation of Example 40 to yield the title compound as an off-white amorphous solid after flash chromatography with 4% MeOH/EtOAc.

Physical characteristics are as follows:

35 ¹H NMR (300 MHz, CD₃OD) δ 0.12, 0.43, 0.68, 0.90-0.97, 1.36, 1.71, 2.60, 3.12, 3.67, 6.88, 7.06, 7.24, 7.51, 7.65.

2187523

HRMS found: 488.2225.

Anal. found: C, 61.25; H, 6.94; N, 8.42.

EXAMPLE 49 N-[3-[1-(4-Hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-pyran-3-yl)-cyclopropylmethyl]-phenyl]-8-quinolinesulfonamide (Formula D-6: R₁ is propyl, R₂ is propyl, R₃ is cyclopropyl, R₄ is quinolin-8-yl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula D-5: R₁ is propyl, R₂ is propyl, R₃ is cyclopropyl) and 8-quinolinesulfonyl chloride of formula D-7 wherein R₄ is quinolin-8-yl, using the general procedure for sulfonylation of Example 40 to yield the title compound as a white amorphous solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

¹H NMR (300 MHz, CD₃OD) δ -0.14, 0.01, 0.35, 0.89, 1.35, 1.63, 2.52, 2.94, 6.79, 6.94, 7.09, 7.64, 8.12, 8.28, 8.41, 9.13.

HRMS found: 535.2256

Anal. found: C, 67.58; H, 6.53; N, 5.11.

EXAMPLE 50 4-Cyano-N-[3-[1-(4-hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-pyran-3-yl)-cyclopropylmethyl]-phenyl]-benzenesulfonamide (Formula D-6: R₁ is propyl, R₂ is propyl, R₃ is cyclopropyl, R₄ is 4-cyanophenyl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula D-5: R₁ is propyl, R₂ is propyl, R₃ is cyclopropyl) and 4-cyano-benzenesulfonyl chloride of formula D-7 wherein R₄ is 4-cyanophenyl, using the general procedure for sulfonylation of Example 40 to yield the title compound as a white amorphous solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

¹H NMR (300 MHz, CD₃OD) δ 0.13, 0.44, 0.62, 0.91, 1.19, 1.67, 2.57, 3.14, 6.80, 7.12, 7.25, 7.83.

HRMS found: 509.2096

Anal. found: C, 65.86; H, 6.39; N, 5.48.

EXAMPLE 51 4-Fluoro-N-[3-[1-(4-hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-pyran-3-yl)-cyclopropylmethyl]-phenyl]-benzenesulfonamide (Formula D-6: R₁ is propyl, R₂ is propyl, R₃ is cyclopropyl, R₄ is 4-fluorophenyl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula

D-5: R₁ is propyl, R₂ is propyl, R₃ is cyclopropyl) and 4-fluoro-benzenesulfonyl chloride of formula D-7 wherein R₄ is 4-fluorophenyl, using the general procedure for sulfonylation of Example 40 to yield the title compound as a white amorphous solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

5 Physical characteristics are as follows:

¹H NMR (300 MHz, CD₃OD) δ 0.11, 0.43, 0.62, 0.92, 1.34, 1.65, 2.57, 3.13, 6.79, 7.03-7.24, 7.75.

HRMS found: 502.2063.

Anal. found: C, 63.96; H, 6.29; N, 2.71.

10 PREPARATION 21 Chiral HPLC resolution of 4-Hydroxy -3-[1-(3-nitrophenyl)-propyl]-5,6-dihydro-6,6-dipropyl-2H-pyran-2-one (Formula D-4: R₁ is propyl, R₂ is propyl, R₃ is ethyl) Refer to Chart D.

A solution of the title product of Preparation 18 (Formula D-4: R₁ is propyl, R₂ is propyl, R₃ is ethyl) (30 mg/mL) in 15% isopropyl alcohol in hexane is
15 chromatographed on a 2.0 x 25 cm (R,R) Whelk-O 1 (Regis technologies, Inc., Morton Grove, IL 60053) column at 1 mL per injection using an automated chromatographic system. The eluant is monitored at 270 nM and appropriate fractions from multiple injections combined and concentrated *in vacuo* to give tan oils. Fractions from multiple injections are analyzed on a 0.46 x 25 cm (S,S) Whelk-O 1 column with the
20 same solvent at 1.0 mL/min. The first peak from the 1.0 cm column is >99% ee (Rt is min) and the latter peak is 92% ee (Rt is min). Prior to further use, the resolved materials are subjected to flash chromatography on silica gel with 3:1 hexanes/EtOAc. The resolved materials are converted to the amines of Formula D-5 using the conditions described in Preparation 20.

25 Physical characteristics are as follows:

The resolved materials were found to exhibit identical ¹H NMR and tlc behavior as the racemic material.

EXAMPLE 52 (R or S)-N-[3-[1-(4-Hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-8-quinolinesulfonamide (Formula D-
30 6: R₁ is propyl, R₂ is propyl, R₃ is ethyl, R₄ is quinolin-8-yl)
Refer to Chart D.

The title compound is prepared from the amine of Preparation 21 (Formula D-5: R₁ is propyl, R₂ is propyl, R₃ is ethyl) and 8-quinolinesulfonyl chloride of formula D-7 wherein R₄ is quinolin-8-yl, using the general procedure for
35 sulfonylation of Example 40 to yield the title compound as a white amorphous solid

after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

¹H NMR and tlc behavior is identical to that of racemic mixture.

¹H NMR (300 MHz, CD₃OD) δ 0.67, 0.85, 1.27, 1.54, 2.01, 3.73, 6.78, 6.90,
5 7.04, 7.57, 8.12, 8.29, 8.38, 9.13.

MS m/e (rel%): 523 (100), 524 (34), 129 (11), 525 (11), 522 (10), 130 (7), 139
(5), 134 (4).

EXAMPLE 53 (R or S)-N-[3-[1-(4-hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-
pyran-3-yl)-propyl]-phenyl]-1-methyl-1H-imidazole-4-
10 sulfonamide (Formula D-6: R₁ is propyl, R₂ is propyl, R₃ is
ethyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 21 (Formula
D-5: R₁ is propyl, R₂ is propyl, R₃ is ethyl) and 1-methyl-imidazole-4-sulfonyl
chloride of formula D-7 wherein R₄ is 1-methylimidazol-4-yl, using the general
15 procedure for sulfonylation of Example 40 to yield the title compound as an off-white
amorphous solid after flash chromatography with 4% MeOH/EtOAc.

Physical characteristics are as follows:

¹H NMR and tlc behavior is identical to racemic mixture.

¹H NMR (300 MHz, CD₃OD) δ 0.88, 1.32, 1.64, 1.93, 2.16, 2.56, 3.68, 3.91,
20 6.87, 7.03, 7.14, 7.53, 7.64.

MS m/e (rel%): 476 (100), 477 (28), 139 (14), 492 (12), 134 (11), 278 (10), 478
(10), 83 (9), 552 (8), 145 (7).

EXAMPLE 54 (S or R)-N-[3-[1-(4-Hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-
pyran-3-yl)-propyl]-phenyl]-8-quinolinesulfonamide (Formula D-
25 6: R₁ is propyl, R₂ is propyl, R₃ is ethyl, R₄ is quinolin-8-yl)
Refer to Chart D.

The title compound is prepared from the amine of Preparation 21 (Formula
D-5: R₁ is propyl, R₂ is propyl, R₃ is ethyl) and 8-quinolinesulfonyl chloride of
formula D-7 wherein R₄ is quinolin-8-yl, using the general procedure for
30 sulfonylation of Example 40 to yield the title compound as a white amorphous solid
after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

¹H NMR and tlc behavior is identical to that of racemic mixture.

¹H NMR (300 MHz, CD₃OD) δ 0.67, 0.85, 1.27, 1.54, 2.01, 3.73, 6.78, 6.90,
35 7.04, 7.57, 8.12, 8.29, 8.38, 9.13.

2187523

MS m/e (rel%): 523 (100), 524 (34), 522 (24), 539 (13), 525 (10), 129 (10), 130 (5), 134 (5), 128 (5), 540 (5).

EXAMPLE 55 (S or R)-N-[3-[1-(4-hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula D-6: R₁ is propyl, R₂ is propyl, R₃ is ethyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 21 (Formula D-5: R₁ is propyl, R₂ is propyl, R₃ is ethyl) and 1-methyl-imidazole-4-sulfonyl chloride of formula D-7 wherein R₄ is 1-methylimidazol-4-yl, using the general procedure for sulfonylation of Example 40 to yield the title compound as an off-white amorphous solid after flash chromatography with 4% MeOH/EtOAc.

Physical characteristics are as follows:

¹H NMR and tlc behavior is identical to racemic mixture.

¹H NMR (300 MHz, CD₃OD) δ 0.88, 1.32, 1.64, 1.93, 2.16, 2.56, 3.68, 3.91, 6.87, 7.03, 7.14, 7.53, 7.64.

MS m/e (rel%): 476 (100), 477 (28), 139 (19), 490 (15), 498 (14), 83 (12), 478 (9), 55 (9), 145 (9), 134 (7).

EXAMPLE 56 (R or S)-N-[3-[1-(4-Hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-pyran-3-yl)-cyclopropylmethyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula D-6: R₁ is propyl, R₂ is propyl, R₃ is cyclopropyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 21 (Formula D-5: R₁ is propyl, R₂ is propyl, R₃ is cyclopropyl) and 1-methyl-imidazole-4-sulfonyl chloride of formula D-7 wherein R₄ is 1-methylimidazol-4-yl, using the general procedure for sulfonylation of Example 40 to yield the title compound as an off-white amorphous solid after flash chromatography with 4% MeOH/EtOAc.

Physical characteristics are as follows:

¹H NMR and tlc behavior is identical to racemic mixture.

¹H NMR (300 MHz, CD₃OD) δ 0.12, 0.43, 0.68, 0.90-0.97, 1.36, 1.71, 2.60, 3.12, 3.67, 6.88, 7.06, 7.24, 7.51, 7.65.

MS m/e (rel%): 488 (100), 489 (30), 139 (18), 145 (14), 490 (10), 55 (10), 83 (9), 564 (7), 146 (7), 510 (7).

EXAMPLE 57 (S or R)-N-[3-[1-(4-Hydroxy-2-oxo-6,6-dipropyl-5,6-dihydro-2H-pyran-3-yl)-cyclopropylmethyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula D-6: R₁ is propyl, R₂ is propyl, R₃ is

2187523

cyclopropyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.

The title compound is prepared from the amine Preparation 21 (Formula D-5: R₁ is propyl, R₂ is propyl, R₃ is cyclopropyl) and 1-methyl-imidazole-4-sulfonyl chloride of formula D-7 wherein R₄ is 1-methylimidazol-4-yl, using the general
5 procedure for sulfonylation of Example 40 to yield the title compound as an off-white amorphous solid after flash chromatography with 4% MeOH/EtOAc.

Physical characteristics are as follows:

¹H NMR and tlc behavior is identical to racemic mixture.

¹H NMR (300 MHz, CD₃OD) δ 0.12, 0.43, 0.68, 0.90-0.97, 1.36, 1.71, 2.60,
10 3.12, 3.67, 6.88, 7.06, 7.24, 7.51, 7.65.

MS m/e (rel%): 488 (100), 489 (29), 139 (18), 145 (16), 83 (10), 55 (10), 490
(10), 510 (8), 146 (8), 144 (7).

EXAMPLE 58 4-Cyano-N-[3-[1-(4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-
dihydro-2H-pyran-3-yl)-cyclopropylmethyl]-phenyl]-benzene-
15 sulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is
cyclopropyl, R₄ is 4-cyanophenyl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula
D-5: R₁ is phenethyl, R₂ is propyl, R₃ is cyclopropyl) and 4-fluoro-benzenesulfonyl
chloride of formula D-7 wherein R₄ is 4-cyanophenyl, using the general procedure for
20 sulfonylation of Example 40 to yield the title compound as a white amorphous solid
after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

¹H NMR (300 MHz, CD₃OD) δ 0.11, 0.42, 0.61, 0.95, 1.24, 1.74-2.00, 2.61-
2.73, 3.30, 6.83-7.23, 7.71-7.84.

25 HRMS found: 571.2267

EXAMPLE 59 4-Fluoro-N-[3-[1-(4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-
dihydro-2H-pyran-3-yl)-cyclopropylmethyl]-phenyl]-
benzenesulfonamide (Formula D-6: R₁ is phenethyl, R₂ is
propyl, R₃ is cyclopropyl, R₄ is 4-fluorophenyl) Refer to Chart
30 D.

The title compound is prepared from the amine of Preparation 20 (Formula
D-5: R₁ is phenethyl, R₂ is propyl, R₃ is cyclopropyl) and 4-fluoro-benzenesulfonyl
chloride of formula D-7 wherein R₄ is 4-fluorophenyl, using the general procedure
for sulfonylation of Example 40 to yield the title compound as a white amorphous
35 solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

¹H NMR complicated by presence of diastereomers.

¹H NMR (300 MHz, CD₃OD) δ 0.11, 0.43, 0.67, 0.96, 1.41, 1.67-2.13, 2.62, 3.16, 6.84, 7.02-7.31, 7.72.

5 HRMS found: 564.2211.

EXAMPLE 60 N-[3-[1-(4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-cyclopropylmethyl]-phenyl]-8-quinolinesulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is cyclopropyl, R₄ is quinolin-8-yl) Refer to Chart D.

10 The title compound is prepared from the amine of Preparation 20 (Formula D-5: R₁ is phenethyl, R₂ is propyl, R₃ is cyclopropyl) and 8-quinolinesulfonyl chloride of formula D-7 wherein R₄ is quinolin-8-yl, using the general procedure for sulfonylation of Example 40 to yield the title compound as a white amorphous solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

15 Physical characteristics are as follows:

¹H NMR complicated by presence of diastereomers.

¹H NMR (300 MHz, CD₃OD) δ -0.13, 0.01, 0.35, 0.93, 1.46, 1.54, 1.58-2.06, 2.56, 2.96, 6.81-7.23, 7.50-7.68, 8.08, 8.24, 8.37, 9.12.

HRMS found: 597.2398

20 EXAMPLE 61 N-[3-[1-(4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-cyclopropylmethyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is cyclopropyl, R₄ is 1-methyl-imidazol-4-yl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula D-5: R₁ is phenethyl, R₂ is propyl, R₃ is cyclopropyl) and 1-methyl-imidazole-4-sulfonyl chloride of formula D-7 wherein R₄ is 1-methylimidazol-4-yl using the general procedure for sulfonylation of Example 40 to yield the title compound as a white amorphous solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

30 ¹H NMR complicated by presence of diastereomers.

¹H NMR (300 MHz, CD₃OD) δ 0.13, 0.42, 0.67, 0.95, 1.44, 1.68-2.13, 2.56, 3.17, 6.91, 7.01-7.33, 7.52, 7.63.

HRMS found: 550.2370.

EXAMPLE 62 N-[3-[1-(4-Hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-cyclopropylmethyl]-phenyl]-1-methyl-1H-imidazole-4-

sulfonamide (Formula D-6: R_1 is phenethyl, R_2 is phenethyl, R_3 is cyclopropyl, R_4 is 1-methyl-imidazol-4-yl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula D-5: R_1 is phenethyl, R_2 is phenethyl, R_3 is cyclopropyl) and 1-methyl-imidazole-4-sulfonyl chloride of formula D-7 wherein R_4 is 1-methylimidazol-4-yl, using the general procedure for sulfonylation of Example 40 to yield the title compound as a white amorphous solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

¹H NMR (300 MHz, CD₃OD) δ 0.13, 0.42, 0.68, 1.73, 1.88-2.17, 2.68, 3.19, 3.64, 6.93, 7.02-7.31, 7.52, 7.64.

HRMS found: 612.2530.

EXAMPLE 63 N-[3-[1-(4-Hydroxy-2-oxo-6,6-dipentyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula D-6: R_1 is pentyl, R_2 is pentyl, R_3 is ethyl, R_4 is 1-methyl-imidazol-4-yl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula D-5: R_1 is pentyl, R_2 is pentyl, R_3 is ethyl) and 1-methyl-imidazole-4-sulfonyl chloride of formula D-7 wherein R_4 is 1-methylimidazole-4-yl, using the general procedure for sulfonylation of Example 40 to yield the title compound as a white amorphous solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

¹H NMR (300 MHz, CD₃OD) δ 0.87, 1.25, 1.55-1.68, 1.92, 2.13, 2.57, 3.66, 3.93, 6.86, 7.03, 7.16, 7.55, 7.63.

EXAMPLE 64 4-Cyano-N-[3-[1-(4-Hydroxy-2-oxo-6,6-dipentyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-benzenesulfonamide (Formula D-6: R_1 is pentyl, R_2 is pentyl, R_3 is ethyl, R_4 is 4-cyanophenyl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula D-5: R_1 is pentyl, R_2 is pentyl, R_3 is ethyl) and 1-methyl-imidazole-4-sulfonyl chloride of formula D-7 wherein R_4 is 4-cyanophenyl, using the general procedure for sulfonylation of Example 40 to yield the title compound as a white amorphous solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

¹H NMR (300 MHz, CD₃OD) δ 0.86, 1.23, 1.52-1.67, 1.93, 2.14, 2.56, 3.93, 6.80, 7.05, 7.18, 7.80, 7.86.

EXAMPLES 65-93

Using the general procedure of Example 40, but substituting the appropriate reactants, the following compounds of the present invention are prepared:

- EXAMPLE 65 N-[3-[1(*R* or *S*)-(6(*R* or *S*)-4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.
- EXAMPLE 66 N-[3-[1(*R* or *S*)-(6(*S* or *R*)-4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.
- EXAMPLE 67 N-[3-[1(*S* or *R*)-(6(*R* or *S*)-4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.
- EXAMPLE 68 N-[3-[1(*S* or *R*)-(6(*S* or *R*)-4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.
- EXAMPLE 69 N-[3-[*t*-Butyl-(4-hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-methyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is *t*-butyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.
- EXAMPLE 70 4-Cyano-N-[3-[*t*-butyl-(4-hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-methyl]-phenyl]-benzenesulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is *t*-butyl, R₄ is 4-cyanophenyl) Refer to Chart D.
- EXAMPLE 71 4-Fluoro-N-[3-[*t*-butyl-(4-hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-methyl]-phenyl]-benzenesulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is *t*-butyl, R₄ is 4-fluorophenyl) Refer to Chart D.

- EXAMPLE 72 N-[3-[*t*-Butyl-(4-hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-methyl]-phenyl]-8-quinolinesulfonamide
(Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is *t*-butyl, R₄ is quinolin-8-yl) Refer to Chart D.
- 5 EXAMPLE 73 N-[3-[1-(6-(2-(1-Methyl-1H-imidazole-4-sulfonylamino)-ethyl)-4-hydroxy-2-oxo-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula D-6: R₁ is 2-(1-methylimidazole-4-sulfonylamino)-ethyl, R₂ is propyl, R₃ is ethyl, R₄ is 1-methylimidazol-4-yl) Refer to Chart D.
- 10 EXAMPLE 74 N-[3-[1-(4-Hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-2-pyridinesulfonamide (Formula D-6: R₁ is phenethyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 2-pyridyl) Refer to Chart D.
- 15 EXAMPLE 75 4-Cyano-N-[3-[1-(4-hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-2-pyridinesulfonamide (Formula D-6: R₁ is phenethyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 4-cyano-2-pyridyl) Refer to Chart D.
- 20 EXAMPLE 76 N-[3-[1-(4-Hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-2-quinolinesulfonamide (Formula D-6: R₁ is phenethyl, R₂ is phenethyl, R₃ is ethyl, R₄ is quinolin-2-yl) Refer to Chart D.
- 25 EXAMPLE 77 2-Hydroxy-N-[3-[1-(4-hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-benzenesulfonamide (Formula D-6: R₁ is phenethyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 2-hydroxyphenyl) Refer to Chart D.
- EXAMPLE 78 N-[3-[1-(4-Hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-2-pyrimidinesulfonamide (Formula D-6: R₁ is phenethyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 2-pyrimidyl) Refer to Chart D.
- 30 EXAMPLE 79 N-[3-[1-(4-Hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-2-quinazolinesulfonamide (Formula D-6: R₁ is phenethyl, R₂ is phenethyl, R₃ is ethyl, R₄ is quinazolin-2-yl) Refer to Chart D.
- 35 EXAMPLE 80 N-[3-[1-(4-Hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-7H-purine-6-sulfonamide (Formula D-6: R₁

is phenethyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 7H-purin-6-yl)
Refer to Chart D.

EXAMPLE 81

5

N-[3-[1-(4-Hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1H-imidazole-2-sulfonamide (Formula D-6: R₁ is phenethyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 1H-imidazol-2-yl) Refer to Chart D.

EXAMPLE 82

10

N-[3-[1-(4-Hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1H-benzimidazole-2-sulfonamide (Formula D-6: R₁ is phenethyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 1H-benzimidazol-2-yl) Refer to Chart D.

EXAMPLE 83

15 EXAMPLE 84

N-[3-[1-(4-Hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-thiazole-4-sulfonamide (Formula D-6: R₁ is phenethyl, R₂ is phenethyl, R₃ is ethyl, R₄ is thiazol-2-yl) Refer to Chart D.

EXAMPLE 85

20

4-Cyano-N-[3-[1-(4-hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-2-pyridinesulfonamide (Formula D-6: R₁ is propyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 4-cyano-2-pyridyl) Refer to Chart D.

EXAMPLE 86

25

N-[3-[1-(4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-2-quinolinesulfonamide (Formula D-6: R₁ is propyl, R₂ is phenethyl, R₃ is ethyl, R₄ is quinolin-2-yl) Refer to Chart D.

EXAMPLE 87

30

2-Hydroxy-N-[3-[1-(4-hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-benzenesulfonamide (Formula D-6: R₁ is propyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 2-hydroxyphenyl) Refer to Chart D.

EXAMPLE 88

35 EXAMPLE 89

N-[3-[1-(4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-2-pyrimidinesulfonamide (Formula D-6: R₁ is propyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 2-pyrimidyl) Refer to Chart D.

N-[3-[1-(4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-

pyran-3-yl)-propyl]-phenyl]-2-quinazolinesulfonamide (Formula D-6: R₁ is propyl, R₂ is phenethyl, R₃ is ethyl, R₄ is quinazolin-2-yl) Refer to Chart D.

EXAMPLE 90

5

N-[3-[1-(4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-7H-purine-6-sulfonamide (Formula D-6: R₁ is propyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 7H-purin-6-yl) Refer to Chart D.

EXAMPLE 91

10

N-[3-[1-(4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1H-imidazole-2-sulfonamide (Formula D-6: R₁ is propyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 1H-imidazol-2-yl) Refer to Chart D.

EXAMPLE 92

15

N-[3-[1-(4-Hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-1H-benzimidazole-2-sulfonamide (Formula D-6: R₁ is propyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 1H-benzimidazol-2-yl) Refer to Chart D.

EXAMPLE 93

20

EXAMPLE 93A

4-Fluoro-N-[3-[1-(4-hydroxy-2-oxo-6-phenethyl-6-propyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-benzenesulfonamide (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 4-fluorophenyl) Refer to Chart D.

The title compound is prepared from the amine of Preparation 20 (Formula D-5: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl) and 4-fluoro benzenesulfonyl chloride using the general procedure for sulfonylation of Example 40 to yield the title compound as an off-white amorphous solid after flash chromatography with 5% EtOAc/CH₂Cl₂.

Physical characteristics are as follows:

30

¹H NMR complicated by presence of diastereomers.

¹H NMR (300 MHz, CD₃OD) δ 0.75-0.96, 1.31-1.48, 1.57-2.01, 2.09-2.22, 2.48-2.71, 3.92, 3.94, 6.86-7.24, 7.72.

PREPARATION 22 6-(2-Cyclopropyl-1-cyclopropylmethyl-ethyl)-

4-hydroxy-2H-pyran-2-one (Formula E-2) Refer to Chart E.

35

To a cold (-78°C) stirred solution of 1.5 ml of diisopropylamine in 9 ml of dry

tetrahydrofuran, under argon, is added 6.2 ml of a 1.6 M solution of *n*-butyllithium in hexane. The solution is warmed to 0°C and then treated with a solution of 378 mg of commercially available 4-hydroxy-6-methyl-2-pyrone of formula E-1 in 8 ml of hexamethylphosphoramide. After 30 minutes at 0°C, 0.32 ml of
5 bromomethylcyclopropane is added; after another ten minutes, a second portion of the same amount is added. The reaction is stirred, allowed to warm to room temperature overnight, and is then partitioned between ethyl acetate and excess dilute hydrochloric acid. The organic phase is washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The residue is flash
10 chromatographed on silica gel 60 (230-400 mesh) using 25% ethyl acetate in dichloromethane containing 1% acetic acid to provide 371 mg of the title compound, along with 206 mg of monoalkylated material.

Physical characteristics are as follows:

¹H NMR δ 0.0, 0.4, 0.6, 1.5, 1.6, 2.2, 5.6, 6.1, 7.2-7.3, 11.5;

15 EI MS *m/z*=234;

TLC *R_f* 0.29 (25% ethyl acetate in dichloromethane containing 1% acetic acid).

PREPARATION 23 3-(α-Cyclopropyl-*meta*-(benzyloxycarbonylamino)benzyl)-6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2H-pyran-2-one (Formula E-3) Refer to Chart E.
20

A mixture of 367 mg of the title compound of Preparation 22, 470 mg of the title compound of Preparation F-5, 60 mg of *p*-toluenesulfonic acid monohydrate, and 1 g of 3 Å molecular sieves in 5 ml of benzene is heated with stirring overnight under argon. The mixture is diluted with dichloromethane and ether and filtered
25 through a pad of sodium sulfate. The solvent is removed under reduced pressure and the residue is flash chromatographed on silica gel 60 (230-400 mesh) using 5-20% ethyl acetate in dichloromethane to afford 399 mg of the title compound.

Physical characteristics are as follows:

¹H NMR δ -0.06, 0.3, 0.5, 1.4, 1.5, 2.5, 3.5, 5.1, 7.2-7.4;

30 EI HRMS *m/z*=513.2513;

TLC *R_f* 0.28 (5% ethyl acetate in dichloromethane).

PREPARATION 24 3-(α-Cyclopropyl-*meta*-aminobenzyl)-6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2H-pyran-2-one (Formula E-4) Refer to Chart E.

35 A mixture of 391 mg of the title compound of Preparation 23 and 100 mg of

2187523

5% palladium on carbon in 10 ml of methanol is shaken overnight under 40 psi of hydrogen. The mixture is then filtered through Celite, and the filtrate is concentrated under reduced pressure to provide 280 mg of the title compound.

Physical characteristics are as follows:

5 ^1H NMR δ 0.0, 0.2-0.7, 1.4, 1.6, 1.8, 2.6, 6.8, 7.2-7.4;

TLC R_f 0.38 (30% ethyl acetate in dichloromethane.)

EXAMPLE 94 N-(3-{Cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-4-fluorobenzenesulfonamide (Formula E-6) Refer to Chart E.

10 To a mixture of 57 mg of the title compound of Preparation 24 and 24 μL of pyridine in 0.5 mL of dichloromethane is added 29 mg of 4-fluorobenzenesulfonyl chloride. After stirring overnight, the solution is diluted with ethyl acetate and washed with dilute aqueous hydrochloric acid, brine, dried over sodium sulfate, and concentrated under reduced pressure. The residue is purified by flash column
15 chromatography on silica gel 60 (230-400 mesh) using 10% ethyl acetate in dichloromethane to give 56 mg of the title compound.

Physical characteristics are as follows:

^1H -NMR (CDCl_3) δ -0.07, 0.13, 0.33, 0.54, 1.39, 1.51, 1.72, 2.55, 3.39, 6.12, 6.87, 7.00, 7.08, 7.19, 7.27, 7.72, 9.72;

20 EI-MS: $[M^+]=537.1977$ found.

EXAMPLES 95-97

Following the procedure described above and using starting materials and reagents known and available to one of ordinary skill in organic synthesis, the following additional compounds are prepared:

25 EXAMPLE 95 4-Cyano-N-(3-{cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-benzenesulfonamide (Formula E-7) Refer to Chart E.

Physical characteristics are as follows:

30 ^1H -NMR (CDCl_3) δ -0.03, 0.13, 0.23, 0.36, 0.44, 0.57, 1.41, 1.58, 1.75, 2.57, 3.32, 5.98, 6.89, 7.11, 7.21, 7.68, 7.82;

EI-MS: $[M^+]=544.2035$ found.

EXAMPLE 96 N-(3-{Cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-8-quinolinesulfonamide (Formula E-8) Refer to Chart E.

35 Physical characteristics are as follows:

¹H-NMR (CDCl₃) δ -0.07, 0.18, 0.37, 0.54, 1.37, 1.51, 2.53, 3.31, 5.96, 6.87, 7.00, 7.13, 7.48, 7.54, 7.92, 8.23, 9.07;

EI-MS: [M+]=570.2188 found.

5 EXAMPLE 97 N-(3-{Cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide (Formula E-9) Refer to Chart E.

Physical characteristics are as follows:

¹H-NMR (CDCl₃-CD₃OD) δ -0.08, 0.13, 0.33, 0.56, 1.37, 1.51, 1.73, 2.54, 3.21, 3.60, 5.95, 6.82, 7.0, 7.19, 7.37, 7.5;

10 EI-MS: [M+]=523.2142 found.

15 EXAMPLE 98 Chiral HPLC Separation of N-(3-{Cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide (Formula E-9) to give (*R* or *S*)-N-(3-{Cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide (Formula E-10) and (*R* or *S*)-N-(3-{Cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide (Formula E-11) Refer to Chart E.

20 A stock sample of the title compound of Example 97 (3 mg/ml) in 5.0 mL each of mobile phase (30% isopropanol, 0.1% acetic acid, and 0.2% water in hexane) and isopropanol is prepared. The stock sample is filtered through a 0.45 micron syringe filter and washed with ethanol to give 14.0 mL of clear filtrate. This solution is
25 chromatographed on a 2.0 x 2.5 cm (R,R) Whelk-O 1 (Regis Technologies, Inc., Morton Grove, IL 60053) column at 3.50 mL per injection using an automated chromatographic system. The eluant is monitored and the pools corresponding to the desired peaks from multiple injections are combined, concentrated under reduced pressure and azeotroped with toluene. The residues are dissolved in methanol,
30 filtered through a syringe filter and the filtrates concentrated under reduced pressure to give the title compounds (>95% pure):

(*R* or *S*)-N-(3-{Cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide (Formula E-10)

35 Physical characteristics are as follows:

2187523

$^1\text{H-NMR}$ ($\text{CDCl}_3\text{-CD}_3\text{OD}$) δ -0.07, 0.14, 0.34, 0.57, 1.32, 1.55, 1.75, 2.51, 3.24, 3.60, 5.87, 6.85, 7.03, 7.15, 7.27, 7.37;

EI-MS: $[\text{M}^+]=523.2149$ found.

(*R* or *S*)-N-(3-{Cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide (Formula E-11)

Physical characteristics are as follows:

$^1\text{H-NMR}$ ($\text{CDCl}_3\text{-CD}_3\text{OD}$) δ -0.07, 0.14, 0.33, 0.55, 1.33, 1.56, 1.75, 2.51, 3.23, 3.60, 5.88, 6.86, 7.03, 7.14, 7.27, 7.38;

EI-MS: $[\text{M}^+]=523.2137$ found.

EXAMPLES 99-103

Following the procedure described above and using starting materials and reagents known and available to one of ordinary skill in organic synthesis, the following additional compounds are prepared:

EXAMPLE 99 N-(3-{Cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-2-pyridine-sulfonamide (Formula E-12)

Physical characteristics are as follows:

$^1\text{H-NMR}$ ($\text{CDCl}_3\text{-CD}_3\text{OD}$) δ -0.05, 0.07, 0.17, 0.34, 0.55, 1.35, 1.55, 1.7, 2.5, 3.24, 5.86, 6.90, 7.03, 7.15, 7.39, 7.78, 8.60;

EI-MS: $[\text{M}^+]=520$;

TLC R_f 0.35 (25% ethyl acetate in dichloromethane).

EXAMPLE 100 N-(3-{Cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-2-sulfonamide (Formula E-13)

Physical characteristics are as follows:

$^1\text{H-NMR}$ ($\text{CDCl}_3\text{-CD}_3\text{OD}$) δ -0.05, 0.15, 0.35, 0.56, 1.35, 1.55, 1.75, 2.53, 3.23, 3.39, 5.89, 6.81, 6.90, 6.97, 7.09, 7.25;

EI-MS: $[\text{M}^+]=523$;

TLC R_f 0.31 (5% methanol in dichloromethane).

EXAMPLE 101 N-(3-{Cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1H-benzoimidazole-2-sulfonamide (Formula E-14)

Physical characteristics are as follows:

$^1\text{H-NMR}$ ($\text{CDCl}_3\text{-CD}_3\text{OD}$) δ -0.07, 0.1, 0.15, 0.35, 0.56, 1.38, 1.58, 1.65, 2.55,

3.28, 5.95, 6.73, 6.96, 7.10, 7.28, 7.58;

FAB-MS: [M+H]=560.2220 found.

EXAMPLE 102 N-(3-{Cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1H-imidazole-2-sulfonamide (Formula E-15)

Physical characteristics are as follows:

¹H-NMR (CDCl₃-CD₃OD) δ 0.0, 0.25, 0.4, 0.6, 1.4, 1.6, 1.65, 2.6, 3.35, 6.0, 6.8, 7.0, 7.2, 7.4;

EI-MS: [M+]=509;

TLC R_f 0.25 (5% methanol in dichloromethane).

EXAMPLE 103 N-(3-{Cyclopropyl-[6-(2-cyclopropyl-1-cyclopropylmethyl-ethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-2-quinolinesulfonamide (Formula E-16)

Physical characteristics are as follows:

¹H-NMR (CDCl₃) δ 0.0, 0.2, 0.4, 0.6, 1.4, 1.6, 1.7, 2.6, 3.3, 6.0, 7.0-7.2, 7.3, 7.7, 7.8-8.0, 8.2, 8.3;

EI-MS: [M+]=570;

TLC R_f 0.53 (5% methanol in dichloromethane).

PREPARATION 25 Cyclopropyl *meta*-nitrophenyl ketone (Formula F-2) Refer to Chart F.

A 250 ml three necked flask fitted with thermometer and addition funnel is charged with 130 ml of fuming 90% nitric acid and cooled to -10°C. Into the stirred liquid is added dropwise 21 ml of commercially available cyclopropyl phenyl ketone of formula F-1. The rate of addition is regulated to maintain the reaction temperature at about -10°C. Upon completion of addition, the resulting clear yellow solution is stirred for another 10 minutes at -10°C, then poured into 1 L of crushed ice. The precipitated solid is extracted with 700 ml of toluene, and the extract is washed twice with 5% sodium hydroxide solution, once with brine, and dried over magnesium sulfate. The solvent is removed under reduced pressure and the residue is recrystallized from methanol at -25°C to give 14.6 g of the title compound as dense, pale yellow prisms. The mother liquor contained substantial amounts of the *ortho* isomer.

Physical characteristics are as follows:

¹H NMR δ 1.2, 1.3, 2.7, 7.70, 8.3, 8.4, 8.85;

IR 1664, 1529, 1352, 1225, 1082, 1017, 852, 689 cm⁻¹;

Anal. Found: C, 62.89; H, 4.73; N, 7.32;

EI MS m/z 191;

TLC R_f 0.32 (25% ethyl acetate in hexane).

PREPARATION 26 *meta*-Aminophenyl cyclopropyl ketone (Formula F-3) Refer to Chart F.

A solution of 5.76 g of the title compound of Preparation 25 is prepared with the aid of heat in 100 ml of methanol. To this is added 450 mg of 5% platinum on carbon catalyst, and the mixture is stirred vigorously under 1 atmosphere of hydrogen. After 5 hours, the mixture is filtered through a pad of Celite and the filtrate concentrated under reduced pressure to afford 4.89 g of the title compound as a greenish oil.

Physical characteristics are as follows:

^1H NMR δ 1.0, 1.2, 2.6, 3.9, 6.8, 7.2, 7.4;

TLC R_f 0.50 (80% ethyl acetate in hexane).

PREPARATION 27 *meta*-Benzyloxycarbonylaminophenyl cyclopropyl ketone (Formula F-4) Refer to Chart F.

To a cold (0°C), stirred solution of 4.89 g of the title compound of Preparation 26 and 6.3 ml of diisopropylethylamine in 90 ml of dichloromethane is added dropwise 4.7 ml of benzyl chloroformate. The completed solution is allowed to warm to room temperature. After 4 hours, the mixture is washed with dilute hydrochloric acid, and the aqueous phase extracted with two additional portions of dichloromethane. The combined organic phase is dried over magnesium sulfate and concentrated under reduced pressure to a yellow solid. This is triturated with two 30 ml portions of hexane, these being discarded, and the remaining solid is dried under vacuum to afford 8.74 g of the title compound.

Physical characteristics are as follows:

TLC R_f 0.45 (5% ethyl acetate in dichloromethane).

PREPARATION 28 *meta*-Benzyloxycarbonylaminophenyl cyclopropyl carbinol (Formula F-5) Refer to Chart F.

To a stirred solution of 8.74 g of compound F-4 of Preparation 27 in 100 ml of tetrahydrofuran and 100 ml of ethanol is added, in portions, 4.5 g of sodium borohydride. After 3 hours at room temperature, the mixture is cooled in ice for the addition of 100 ml of 1N hydrochloric acid. The mixture is thrice extracted with dichloromethane, and the combined extract dried over magnesium sulfate. Solvent is removed under reduced pressure and the residue flash chromatographed on silica gel

60 (230-400 mesh) using 40% ethyl acetate in hexane to provide 8.48 g of the title compound as a white crystalline solid. This is optionally recrystallized from ethyl acetate-hexane.

Physical characteristics are as follows:

5 ^1H NMR δ 0.3-0.6, 1.1, 2.35, 3.92, 5.17, 7.1, 7.2-7.4;

IR 1693, 1599, 1559, 1449, 1235, 1054, 697 cm^{-1} ;

Anal. Found: C, 72.57; H, 6.51; N, 4.61;

PREPARATION 29 4-Hydroxy-6-[3-(2-methoxy-ethoxy)-propyl]-pyran-2-one
(Formula G-1) Refer to Chart G.

10 To a flame-dried flask under an argon atmosphere is added 2.80 mL of diisopropylamine and 20.0 mL of dry tetrahydrofuran. The solution is cooled to -78°C and treated with 12.5 mL (1.6 M in hexane) of *n*-butyllithium. The solution is warmed to 0°C for 30 minutes, then treated with 5.0 mL of dry hexamethylphosphoramide. The lithium diisopropylamide solution is then treated
15 with 1.20 g of commercially available 4-hydroxy-6-methyl-2-pyrone of formula G-0 as a solution in 16 mL of dry tetrahydrofuran and 14 mL of dry hexamethylphosphoramide. After 30 minutes the mixture is treated with 2.30 g of 2-(2-methoxy-ethoxy)-ethyl iodide as a solution in 12 mL of dry tetrahydrofuran. The mixture is stirred 1 hour at 0°C and then warmed to room temperature. After 1 hour
20 the reaction is quenched with excess 1 N aqueous hydrochloric acid. The mixture is concentrated under reduced pressure and partitioned between dichloromethane and water. The aqueous phase is extracted with sufficient volumes of dichloromethane to remove the title compound. The combined organic extracts are washed with brine, dried over magnesium sulfate and concentrated under reduced pressure. The residue
25 is purified by flash column chromatography on silica gel 60 (230-400 mesh) eluting with 30% ethyl acetate in dichloromethane containing 3% acetic acid to 80% ethyl acetate in dichloromethane containing 5% acetic acid to give 1.34 g of the title compound.

Physical characteristics are as follows:

30 ^1H -NMR (CDCl_3) δ 1.93, 2.54, 3.39, 3.55, 5.55, 5.90;

TLC R_f 0.26 (50% ethyl acetate in dichloromethane containing 5% acetic acid).

PREPARATION 30 3-(α -Cyclopropyl-*meta*-(benzyloxycarbonylamino)benzyl)-4-hydroxy-6-[3-(2-methoxy-ethoxy)-propyl]-pyran-2-one (Formula G-2) Refer to Chart G.

35

A mixture of 146 mg of the title compound of Preparation 29, 340 mg of the title compound of Preparation 28, prepared as described in Chart F, 25 mg of *p*-toluenesulfonic acid monohydrate, and 0.5 g of 3 Å molecular sieves in 5 mL of dichloromethane is heated overnight with stirring. The mixture is cooled and the solvent removed under reduced pressure. The residue is purified by flash column chromatography on silica gel 60 (230-400 mesh) eluting with 5% to 10% methanol in ethyl acetate to give 129 mg of the title compound.

Physical characteristics are as follows:

¹H-NMR (CDCl₃) δ 0.25, 0.45, 0.67, 1.77, 2.39, 3.38, 3.51, 5.13, 5.84, 7.17, 7.32, 7.42; EI-MS: [M+]=507.2257 found;

TLC R_f 0.28 (50% ethyl acetate in dichloromethane).

PREPARATION 31 3-(α-Cyclopropyl-*meta*-aminobenzyl)-4-hydroxy-6-[3-(2-methoxyethoxy)-propyl]-pyran-2-one (Formula G-3) Refer to Chart G.

A mixture of 124 mg of the title compound of Preparation 30 and 35 mg of 5% palladium on charcoal in 5 mL of ethanol is shaken overnight under 50 psi of hydrogen. The mixture is filtered through Celite with ethanol washes of the filter cake. The filtrates are combined and the solvent removed under reduced pressure to give 92 mg of the title compound. Physical characteristics are as follows:

TLC R_f 0.12 (ethyl acetate).

EXAMPLE 104 N-(3-{Cyclopropyl-[4-hydroxy-6-(3-(2-methoxyethoxy)-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide (Formula G-4) Refer to Chart G.

To a mixture of 37 mg of the title compound of Preparation 31 and 18 μL of pyridine in 0.5 mL of dichloromethane is added 20 mg of 1-methylimidazole-4-sulfonyl chloride. After stirring overnight, the solvent is removed under reduced pressure. The residue is azeotroped with toluene and is then purified by flash column chromatography on silica gel 60 (230-400 mesh) using 2% to 8% methanol in dichloromethane to give 32 mg of the title compound.

Physical characteristics are as follows:

¹H-NMR (CDCl₃) δ 0.1, 0.24, 0.45, 0.65, 1.75, 1.85, 2.46, 3.30, 3.34, 3.50, 5.98, 6.98, 7.08, 7.19, 7.29, 7.42;

EI-MS: [M+]=517.1874 found;

TLC R_f 0.22 (5% methanol in dichloromethane).

PREPARATION 32 3-(α-Cyclopropyl-*meta*-(benzyloxycarbonylamino)benzyl)-4-hydroxy-6-methyl-pyran-2-one (Formula H-1) Refer to Chart H.

2187523

A mixture of 493 mg of commercially available 4-hydroxy-6-methyl-2-pyrone of formula H-0, 592 mg of the title compound of Preparation 28, prepared as described in Chart F and 56 mg of *p*-toluenesulfonic acid monohydrate in 20 mL of dichloromethane is heated to reflux through an addition funnel containing 3 Å
5 molecular sieves for 6 hours. The mixture is cooled and the solvent removed under reduced pressure. The residue is purified by flash column chromatography on silica gel 60 (230-400 mesh) eluting with 60% to 100% ethyl acetate in dichloromethane to give 470 mg of the title compound.

Physical characteristics are as follows:

10 $^1\text{H-NMR}$ (CDCl_3) δ 0.23, 0.43, 0.66, 1.78, 3.41, 5.09, 5.89, 7.00, 7.14, 7.29, 7.37, 10.1;

EI-MS: $[\text{M}^+]=405$;

TLC R_f 0.52 (ethyl acetate).

PREPARATION 33 3-(α -Cyclopropyl-*meta*-(benzyloxycarbonylamino)benzyl)-4-
15 hydroxy-6-propyl-pyran-2-one (Formula H-2) Refer to Chart H.

To a flame-dried flask under an argon atmosphere is added 0.45 mL of diisopropylamine and 3.0 mL of dry tetrahydrofuran. The solution is cooled to -78°C and treated with 2.0 mL (1.6 M in hexane) of *n*-butyllithium. The solution is warmed to 0°C for 15 minutes, then cooled to -78°C . The lithium diisopropylamide
20 solution is treated with 405 mg of the title compound of Preparation 32 as a solution in 4 mL of dry tetrahydrofuran. After 1 hour at -78°C the mixture is treated with 85 μL of ethyl bromide. The mixture is then stirred at -78°C for 3 hours. The reaction is quenched with excess 1 N aqueous hydrochloric acid. The mixture is warmed and partitioned between ethyl acetate and phosphate buffer. The aqueous phase is
25 extracted twice with ethyl acetate. The combined organic extracts are dried over magnesium sulfate and concentrated under reduced pressure. The residue is purified by flash column chromatography on silica gel 60 (230-400 mesh) eluting with 10% to 20% ethyl acetate in dichloromethane to give 277 mg of the title compound.

Physical characteristics are as follows:

30 $^1\text{H-NMR}$ (CDCl_3) δ 0.24, 0.45, 0.65, 0.88, 1.55, 1.79, 2.28, 3.42, 5.10, 5.95, 6.89, 7.15, 7.3, 10.0;

EI-MS: $[\text{M}^+]=433$;

TLC R_f 0.33 (10% ethyl acetate in dichloromethane).

PREPARATION 34 3-(α -Cyclopropyl-*meta*-(benzyloxycarbonylamino)benzyl)-6-[1-
35 ethyl-3-(2-methoxy-ethoxy)-propyl]-4-hydroxy-pyran-2-one

2187523

(Formula H-3) Refer to Chart H.

To a flame-dried flask under an argon atmosphere is added 0.30 mL of diisopropylamine and 2.0 mL of dry tetrahydrofuran. The solution is cooled to -78°C and treated with 1.3 mL (1.6 M in hexane) of *n*-butyllithium. The solution is warmed to 0°C for 15 minutes, then cooled to -78°C. The lithium diisopropylamide solution is treated with 277 mg of the title compound of Preparation 33 as a solution in 3 mL of dry tetrahydrofuran. After 1 hour at -78°C, the mixture is treated with 180 mg of 2-(2-methoxy-ethoxy)-ethyl iodide in 3 mL of tetrahydrofuran. The mixture is then stirred at -78°C for 3 hours. The reaction is quenched with excess 1 N aqueous hydrochloric acid. The mixture is warmed and partitioned between ethyl acetate and phosphate buffer. The aqueous phase is extracted thrice with ethyl acetate. The combined organic extracts are washed with brine, dried over magnesium sulfate and concentrated under reduced pressure. The residue is purified by flash column chromatography on silica gel 60 (230-400 mesh) eluting with 25% to 40% ethyl acetate in dichloromethane to give 198 mg of the title compound.

Physical characteristics are as follows:

¹H-NMR (CDCl₃) δ 0.25, 0.46, 0.66, 0.75, 1.5, 1.76, 2.36, 3.4, 5.15, 5.96, 7.2, 7.3, 7.42, 10.0;

EI-MS: [M+]=535;

TLC R_f 0.29 (25% ethyl acetate in dichloromethane).

PREPARATION 35 3-(α-Cyclopropyl-*meta*-aminobenzyl)-6-[1-ethyl-3-(2-methoxy-ethoxy)-propyl]-4-hydroxy-pyran-2-one (Formula H-4) Refer to Chart H.

A mixture of 180 mg of the title compound of Preparation 34 and 50 mg of 5% palladium on charcoal in 2 mL of ethanol is shaken overnight under 50 psi of hydrogen. The mixture is filtered through Celite with ethanol washes of the filter cake. The filtrates are combined and the solvent removed under reduced pressure to give 127 mg of the title compound.

Physical characteristics are as follows:

TLC R_f 0.19 (ethyl acetate).

EXAMPLE 105 N-(3-{Cyclopropyl-[6-(1-ethyl-3-(2-methoxy-ethoxy)-propyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide (Formula H-5) Refer to Chart H.

To a mixture of 32 mg of the title compound of Preparation 35 and 13 μL of pyridine in 0.8 mL of dichloromethane is added 14.5 mg of 1-methylimidazole-4-

sulfonyl chloride. After stirring overnight, the solvent is removed under reduced pressure. The residue is azeotroped with toluene and is then purified by flash column chromatography on silica gel 60 (230-400 mesh) using 1% to 4% methanol in dichloromethane to give 38 mg of the title compound.

5 Physical characteristics are as follows:

$^1\text{H-NMR}$ (CDCl_3) δ 0.05, 0.25, 0.45, 0.65, 0.83, 1.5-2.0, 2.45, 3.3-3.5, 3.62, 6.00, 6.99, 7.1-7.3, 7.48;

EI-MS: $[\text{M}^+]=545.2186$ found;

TLC R_f 0.24 (5% methanol in dichloromethane).

10 EXAMPLE 106 4-Cyano-N-(3-{cyclopropyl-[6-(1-ethyl-3-{2-methoxy-ethoxy}-propyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-benzenesulfonamide

Following the procedure described above and using starting materials and reagents known and available to one of ordinary skill in organic synthesis, the title
15 compound is prepared.

Physical characteristics are as follows:

$^1\text{H-NMR}$ (CDCl_3) δ 0.15, 0.25, 0.45, 0.65, 0.78, 1.2-1.8, 2.4, 3.3-3.6, 3.54, 5.89, 6.95, 7.1-7.3, 7.6-7.9;

FAB-MS: $[\text{M}+\text{H}]=567.2176$ found;

20 TLC R_f 0.40 (50% ethyl acetate in dichloromethane).

PREPARATION 36 3-(α -Cyclopropyl-*meta*-(benzyloxycarbonylamino)benzyl)-6-(1-ethyl-3-hydroxy-propyl)-4-hydroxy-pyran-2-one (Formula I-1)
Refer to Chart I.

To a flame-dried flask under an argon atmosphere is added 0.46 mL of
25 diisopropylamine and 3.5 mL of dry tetrahydrofuran. The solution is cooled to -78°C and treated with 2.0 mL (1.6 M in hexane) of *n*-butyllithium. The solution is warmed to 0°C for 20 minutes, then cooled to -78°C . The lithium diisopropylamide solution is treated with 433 mg of the title compound of Preparation 33 as a solution in 4 mL of dry tetrahydrofuran. After 1 hour at -78°C the mixture is treated with
30 gaseous ethylene oxide for 5 minutes. The mixture is then stirred at -78°C for 15 minutes. The reaction is quenched with excess 1 N aqueous hydrochloric acid. The mixture is warmed and partitioned between dichloromethane and phosphate buffer. The aqueous phase is extracted twice with dichloromethane. The combined organic extracts are washed with brine, dried over magnesium sulfate and concentrated
35 under reduced pressure. The residue is purified by flash column chromatography on

silica gel 60 (230-400 mesh) eluting with 2% to 8% methanol in dichloromethane to give 144 mg of the title compound.

Physical characteristics are as follows:

$^1\text{H-NMR}$ (CDCl_3) δ 0.22, 0.45, 0.65, 0.7, 1.3-1.7, 1.8, 2.25, 3.4, 5.1, 5.91, 7.1-

5 7.4;

FAB-MS: $[\text{M}+\text{H}]=478$;

TLC R_f 0.29 (5% methanol in dichloromethane).

PREPARATION 37 6-(3-Bromo-1-ethyl-propyl)-3-(α -cyclopropyl-*meta*-
(benzyloxycarbonylamino)benzyl)-4-hydroxy-pyran-2-one
10 (Formula I-2) Refer to Chart I.

To a stirring solution of 114 mg of the title compound of Preparation 36 in 3 mL of tetrahydrofuran is added 160 mg of triphenylphosphine and 200 mg of carbon tetrabromide. After 2 hours, the solvent is removed under reduced pressure and the residue is purified by flash column chromatography on silica gel 60 (230-400 mesh)
15 eluting with 70% to 100% diethyl ether in hexane to give 113 mg of the title compound.

Physical characteristics are as follows:

$^1\text{H-NMR}$ (CDCl_3) δ 0.25, 0.35, 0.55, 0.65, 0.84, 1.5-2.2, 3.2, 3.35, 3.52, 5.16,
5.95, 6.79, 7.1-7.4;

20 FAB-MS: $[\text{M}+\text{H}]=504.1404$ found;

TLC R_f 0.29 (75% diethyl ether in hexane).

PREPARATION 38 6-(3-Azido-1-ethyl-propyl)-3-(α -cyclopropyl-*meta*-
(benzyloxycarbonylamino)benzyl)-4-hydroxy-pyran-2-one
(Formula I-3) Refer to Chart I.

25 To a stirring solution of 113 mg of the title compound of Preparation 37 in 2.0 mL of ethanol is added 55 mg of sodium azide and 0.5 mL of water. The reaction mixture is heated overnight and then cooled. The solvent is removed under reduced pressure and the residue is purified by flash column chromatography on silica gel 60 (230-400 mesh) eluting with diethyl ether to give 89 mg of the title compound.

30 Physical characteristics are as follows:

$^1\text{H-NMR}$ (CDCl_3) δ 0.23, 0.33, 0.51, 0.68, 0.82, 1.4-2.0, 2.33, 3.1-3.3, 3.5, 5.15,
5.94, 6.84, 7.1-7.4;

EI-MS: $[\text{M}+]=502$;

TLC R_f 0.52 (10% ethyl acetate in dichloromethane).

35 PREPARATION 39 6-(3-Amino-1-ethyl-propyl)-3-(α -cyclopropyl-*meta*-aminobenzyl)-

4-hydroxy-pyran-2-one (Formula I-4) Refer to Chart I.

A mixture of 87 mg of the title compound of Preparation 38 and 35 mg of 5% palladium on charcoal in 4 mL of ethanol is shaken for 4 hours under 40 psi of hydrogen. The mixture is filtered through Celite with ethanol washes of the filter cake. The filtrates are combined and the solvent removed under reduced pressure to give 70 mg of the title compound as a mixture with 6-(3-amino-1-ethyl-propyl)-3-(α -cyclopropyl-*meta*-(benzyloxycarbonylamino)benzyl)-4-hydroxy-pyran-2-one.

Physical characteristics are as follows:

TLC R_f 0.05 (5% methanol in dichloromethane).

10 PREPARATION 40 3-(Cyclopropyl-{3-[1-methyl-1H-imidazole-4-sulfonylamino]-phenyl}-methyl)-6-({1-ethyl-3-[1-methyl-1H-imidazole-4-sulfonylamino]}-propyl)-2-oxo-2H-pyran-4-yl 1-methyl-1H-imidazole-4-sulfonate (Formula I-5) Refer to Chart I.

To a mixture of 70 mg of the title compound of Preparation 39 and 6-(3-amino-1-ethyl-propyl)-3-(α -cyclopropyl-*meta*-(benzyloxycarbonylamino)benzyl)-4-hydroxy-pyran-2-one, also from Preparation 39, in 1.5 mL of dichloromethane is added 120 μ L of diisopropylethylamine and 92 mg of 1-methylimidazole-4-sulfonyl chloride. After stirring overnight, the solvent is removed under reduced pressure. The residue is azeotroped with toluene and is then purified by flash column chromatography on silica gel 60 (230-400 mesh) using 2% to 6% methanol in dichloromethane to give 49 mg of the title compound.

Physical characteristics are as follows:

$^1\text{H-NMR}$ (CDCl_3) δ 0.2-0.5, 0.75, 0.90, 1.4-2.0, 2.55, 3.0-3.4, 3.6-3.7, 6.63, 7.0-7.7;

25 FAB-MS: $[\text{M}+\text{H}]=775$;

TLC R_f 0.14 (5% methanol in dichloromethane).

EXAMPLE 107 N-(3-{Cyclopropyl-[6-(1-ethyl-3-{1-methyl-1H-imidazole-4-sulfonylamino}-propyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide (Formula I-6)

30 Refer to Chart I.

A solution of 49 mg of the title compound of Preparation 40 in 4 mL methanol containing ammonia is cooled to 0°C and treated with gaseous ammonia. After 5 minutes ammonia introduction is ceased, the flask is tightly capped and warmed to room temperature. After standing overnight the solvent is removed under reduced pressure and the residue is purified by flash column chromatography on

silica gel 60 (230-400 mesh) eluting with 3% to 9% methanol in dichloromethane to give 32 mg of the title compound.

Physical characteristics are as follows:

$^1\text{H-NMR}$ (CDCl_3) δ 0.2-0.5, 0.75, 0.90, 1.4-2.0, 2.55, 3.0-3.4, 3.6-3.7, 6.63, 7.0-

5 7.7;

EI-MS: $[\text{M}^+]=523$;

TLC R_f 0.33 (5% methanol in dichloromethane).

PREPARATION 41 3-(α -Cyclopropyl-*meta*-aminobenzyl)-6-(1-ethyl-3-hydroxy-propyl)-4-hydroxy-pyran-2-one (Formula J-1) Refer to Chart J.

10 A mixture of 477 mg of the title compound of Preparation 36 and 150 mg of 5% palladium on charcoal in 10 mL of ethanol is shaken overnight under 45 psi of hydrogen. The mixture is filtered through Celite with ethanol washes of the filter cake. The filtrates are combined and the solvent is removed under reduced pressure to give 340 mg of the title compound.

15 Physical characteristics are as follows:

TLC R_f 0.10 (5% methanol in dichloromethane).

PREPARATION 42 6-(3-Bromo-1-ethyl-propyl)-3-(α -cyclopropyl-*meta*-aminobenzyl)-4-hydroxy-pyran-2-one (Formula J-2) Refer to Chart J.

20 To a stirring solution of 340 mg of the title compound of Preparation 41 in 7 mL of tetrahydrofuran is added 525 mg of triphenylphosphine and 663 mg of carbon tetrabromide. After 30 minutes, the solvent is removed under reduced pressure and the residue is purified by flash column chromatography on silica gel 60 (230-400 mesh) eluting with 30% to 50% ethyl acetate in dichloromethane to give 228 mg of the title compound.

25 Physical characteristics are as follows:

$^1\text{H-NMR}$ (CDCl_3) δ 0.19, 0.42, 0.58, 0.75, 1.4-2.4, 3.14, 3.3, 5.26, 6.15, 6.47, 6.91, 7.00;

TLC R_f 0.45 (5% methanol in dichloromethane).

30 EXAMPLE 108 N-(3-[[6-(3-Bromo-1-ethyl-propyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-cyclopropyl-methyl]-phenyl)-1-methyl-1H-imidazole-4-sulfonamide (Formula J-3) Refer to Chart J.

To a mixture of 102 mg of the title compound of Preparation 42 and 40 μL of pyridine in 1.0 mL of dichloromethane is added 45 mg of 1-methylimidazole-4-sulfonyl chloride. After stirring overnight, the solvent is removed under reduced pressure. The residue is azeotroped with toluene and is then purified by flash

35

column chromatography on silica gel 60 (230-400 mesh) using 2% to 5% methanol in dichloromethane to give 86 mg of the title compound.

Physical characteristics are as follows:

¹H-NMR (CDCl₃-CD₃OD) δ 0.2, 0.44, 0.60, 0.82, 1.4-2.2, 2.5, 3.1-3.4, 3.62, 5.93, 6.92, 7.07, 7.19, 7.30, 7.40;

FAB-MS: [M+H]=550.1037 found;

TLC R_f 0.36 (5% methanol in dichloromethane).

EXAMPLE 109 N-(3-[[6-(3-Azido-1-ethyl-propyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-cyclopropyl-methyl]-phenyl)-1-methyl-1H-imidazole-4-sulfonamide (Formula J-4) Refer to Chart J.

To a stirring solution of 113 mg of the title compound of Example 108 in 1.2 mL of ethanol is added 50 mg of sodium azide and 0.4 mL of water. The reaction mixture is heated overnight and then cooled. The solvent is removed under reduced pressure and the residue is purified by flash column chromatography on silica gel 60 (230-400 mesh) eluting with 3% to 6% methanol in dichloromethane to give 57 mg of the title compound.

Physical characteristics are as follows:

¹H-NMR (CDCl₃-CD₃OD) δ 0.25, 0.48, 0.66, 0.90, 1.3-1.8, 2.42, 2.9-3.2, 3.68, 5.94, 6.93, 7.12, 7.19, 7.23, 7.35, 7.46;

FAB-MS: [M+H]=550.1037 found;

TLC R_f 0.36 (5% methanol in dichloromethane).

PREPARATION 43 N-(3-[[6-(3-Amino-1-ethyl-propyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-cyclopropyl-methyl]-phenyl)-1-methyl-1H-imidazole-4-sulfonamide (Formula J-5) Refer to Chart J.

A mixture of 104 mg of the title compound of Example 109 and 30 mg of 5% palladium on charcoal in 2 mL of each of methanol and ethanol is shaken overnight under 45 psi of hydrogen. The mixture is filtered through Celite with methanol washes of the filter cake. The filtrates are combined and the solvent is removed under reduced pressure to give 69 mg of the title compound.

Physical characteristics are as follows:

TLC R_f 0.05 (5% methanol in dichloromethane).

EXAMPLE 110 2-[[8-[[3-[3-[Cyclopropyl[3-[(1-methyl-1H-imidazol-4-yl)sulfonyl]amino]phenyl]methyl]-4-hydroxy-2-oxo-2H-pyran-6-yl]pentyl]amino]-1,8-dioxooctyl]methylamino]-ethane sulfonic acid, monosodium salt (Formula J-6) Refer to Chart J.

A suspension of 69 mg of the title compound of Preparation 43 in 1.0 mL of dichloromethane is treated with 0.22 mL (0.65 M in acetonitrile) of the triethylamine salt of suleptanic acid and 25 μ L of diisopropylcarbodiimide. After 1 hour the mixture is treated with 0.5 mL of dimethylformamide. After stirring overnight the solvent is removed under reduced pressure and the residue is purified by flash column chromatography on silica gel 60 (230-400 mesh) eluting with 10% to 30% methanol in dichloromethane. The crude product is dissolved in water saturated *n*-butanol and partitioned with saturated aqueous sodium sulfate. The aqueous phase is extracted twice with additional portions of water saturated *n*-butanol. The combined *n*-butanol layers are filtered through a pad of sodium sulfate and concentrated under reduced pressure to give 94 mg of the title compound.

Physical characteristics are as follows:

$^1\text{H-NMR}$ ($\text{CDCl}_3\text{-CD}_3\text{OD}$) δ 0.05-0.6, 0.83, 1.1-2.5, 2.9-3.7, 3.68, 5.84, 6.8-7.6;

FAB-MS: $[\text{M}+\text{H}]=786.2838$ found;

TLC R_f 0.21 (20% methanol in dichloromethane).

EXAMPLES 111-134

Utilizing procedures described above and using starting materials and reagents known and available to one of ordinary skill in organic synthesis, the following additional compounds are prepared:

111) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-[(2-hydroxy-1,1-bis(hydroxymethyl)-ethyl)-amino]-carbonyl)-amino-propyl]-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-benzenesulfonamide

112) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-[(2-hydroxy-1,1-bis(hydroxymethyl)-ethyl)-amino]-carbonyl)-amino-propyl]-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-2-pyridinesulfonamide

113) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-[(2-hydroxy-1,1-bis(hydroxymethyl)-ethyl)-amino]-carbonyl)-amino-propyl]-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1H-imidazole-2-sulfonamide

114) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-[(2-hydroxy-1,1-bis(hydroxymethyl)-ethyl)-amino]-carbonyl)-amino-propyl]-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1H-benzoimidazole-2-sulfonamide

115) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-[(2-hydroxy-1,1-bis(hydroxymethyl)-ethyl)-amino]-carbonyl)-amino-propyl]-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide

116) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-[(2-hydroxy-1,1-bis(hydroxymethyl)-

ethyl)-amino]-carbonyl)-amino-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-2-sulfonamide

- 117) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-{γ-L-glutamyl}-amino-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-benzenesulfonamide
- 5 118) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-{γ-L-glutamyl}-amino-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-2-pyridinesulfonamide
- 119) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-{γ-L-glutamyl}-amino-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1H-imidazole-sulfonamide
- 120) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-{γ-L-glutamyl}-amino-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1H-benzoimidazole-sulfonamide
- 10 121) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-{γ-L-glutamyl}-amino-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide
- 122) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-{γ-L-glutamyl}-amino-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-2-sulfonamide
- 15 123) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-{[piperazin-1-yl]-carbonyl}-amino-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-benzenesulfonamide
- 124) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-{[piperazin-1-yl]-carbonyl}-amino-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-2-pyridinesulfonamide
- 125) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-{[piperazin-1-yl]-carbonyl}-amino-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1H-imidazole-2-sulfonamide
- 20 126) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-{[piperazin-1-yl]-carbonyl}-amino-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1H-benzoimidazole-2-sulfonamide
- 127) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-{[piperazin-1-yl]-carbonyl}-amino-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide
- 25 128) N-(3-{Cyclopropyl-[4-hydroxy-6-(3-{[piperazin-1-yl]-carbonyl}-amino-propyl)-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-2-sulfonamide
- 129) 2-[[8-[[3-[3-[Cyclopropyl[3-[[phenylsulfonyl]amino]phenyl]methyl]-4-hydroxy-2-oxo-2H-pyran-6-yl]propyl]amino]-1,8-dioxooctyl]methylamino]-ethanesulfonic acid, monosodium salt
- 30 130) 2-[[8-[[3-[3-[Cyclopropyl[3-[(2-pyridyl)sulfonyl]amino]phenyl]methyl]-4-hydroxy-2-oxo-2H-pyran-6-yl]propyl]amino]-1,8-dioxooctyl]methylamino]-ethane sulfonic acid, monosodium salt
- 131) 2-[[8-[[3-[3-[Cyclopropyl[3-[(1H-benzimidazol-2-yl)sulfonyl]amino]phenyl]methyl]-4-hydroxy-2-oxo-2H-pyran-6-yl]propyl]amino]-1,8-dioxooctyl]methyl amino]-ethanesulfonic acid, monosodium salt
- 35

132) 2-[[8-[[3-[3-[Cyclopropyl[3-[(1H-imidazol-2-yl)sulfonyl]amino]phenyl]methyl]-4-hydroxy-2-oxo-2H-pyran-6-yl]propyl]amino]-1,8-dioxooctyl]methylamino]-ethane sulfonic acid, monosodium salt

133) 2-[[8-[[3-[3-[Cyclopropyl[3-[(1-methyl-1H-imidazol-4-yl)sulfonyl]amino]phenyl]methyl]-4-hydroxy-2-oxo-2H-pyran-6-yl]propyl]amino]-1,8-dioxooctyl]methylamino]-ethanesulfonic acid, monosodium salt

134) 2-[[8-[[3-[3-[Cyclopropyl[3-[(1-methyl-1H-imidazol-2-yl)sulfonyl]amino]phenyl]methyl]-4-hydroxy-2-oxo-2H-pyran-6-yl]propyl]amino]-1,8-dioxooctyl]methylamino]-ethane sulfonic acid, monosodium salt

10 PREPARATION 44 (Tetrahydropyran-4-yl)-methanol (Formula K-2) Refer to Chart K.

To a cold (0°), stirred solution of 651 mg of tetrahydropyran-4-carboxylic acid in 2.5 ml of dry tetrahydrofuran, under argon, is added dropwise 10 ml of a 1.0 M solution of borane in tetrahydrofuran. After 18 hours at room temperature, the solution is recooled to 0° and quenched with 1 ml of 1M KOH. The mixture is acidified with 1M aqueous hydrochloric acid and extracted four times with dichloromethane. The extract is dried over magnesium sulfate and concentrated carefully under reduced pressure to afford 0.72 g of the alcohol as a colorless liquid.

Physical characteristics are as follows:

20 ¹H NMR δ 1.2-1.4, 1.6, 1.8, 3.3-3.4, 3.6, 4.0 ppm.

PREPARATION 45 (Tetrahydropyran-4-yl)-methyl *p*-toluenesulfonate (Formula K-3) Refer to Chart K.

To a cold (0°), stirred solution of 5 mmol of the title compound of Preparation 44 and 0.81 ml of pyridine in 5 ml of dichloromethane is added 1.05 g of *p*-toluenesulfonyl chloride, and the solution is allowed to warm to room temperature. After 18 hours the mixture is partitioned between ethyl acetate and dilute aqueous hydrochloric acid, and the organic phase is washed with brine and dried over magnesium sulfate. Following removal of solvent under reduced pressure, the residue is flash chromatographed on silica using 50% ethyl acetate in hexane to afford 1.23 g of the title compound as a colorless liquid.

Physical characteristics are as follows:

¹H NMR δ 1.2-1.4, 1.6, 1.9-2.0, 2.46, 3.34, 3.85, 3.95, 7.3, 7.8 ppm.

MS: 270

35 PREPARATION 46 (Tetrahydropyran-4-yl)-methyl iodide (Formula K-4) Refer to Chart K.

A solution of 800 mg of tosylate of Preparation 45 and 887 mg of sodium iodide in 6 ml of acetone is refluxed under nitrogen for six hours, then partitioned between ether and dilute aqueous sodium thiosulfate. The organic phase is washed with brine, dried over magnesium sulfate, and concentrated carefully under atmospheric pressure to give 648 mg of the iodide as a colorless liquid.

Physical characteristics are as follows:

^1H NMR δ 1.2-1.4, 1.6-1.9, 3.1, 3.37, 3.97 ppm.

PREPARATION 47 6-(1-(Tetrahydropyran-4-ylmethyl)-propyl)-4-hydroxypyran-2-one (Formula K-5) Refer to Chart K.

To a cold (-78°) stirred solution of 0.90 ml of diisopropylamine in 5 ml of tetrahydrofuran, under argon, is added *via* syringe 3.7 ml of a 1.6 M solution of *n*-butyllithium in hexane. The solution is warmed to 0° , and after ten minutes, a solution of 431 mg of the title compound of Preparation 50 in 3 ml of hexamethylphosphoramide is added *via* cannula. After 20 minutes, the deep red solution is cooled to -50° , and 605 mg of iodide of Preparation 46 in 1 ml of tetrahydrofuran is added *via* cannula. The reaction is allowed to warm slowly to 0° and then quenched by addition of pH 7 phosphate buffer. Following removal of tetrahydrofuran under reduced pressure, the residual liquid is acidified with dilute aqueous hydrochloric acid and the resulting precipitate extracted with two portions of ethyl acetate. The organic is washed with dilute aqueous hydrochloric acid and brine, dried over magnesium sulfate, and concentrated under reduced pressure. Flash chromatography of the residue on silica using 5% acetic acid and 30-40% ethyl acetate in dichloromethane provides 553 mg of the title compound as a thick yellow gum.

Physical characteristics are as follows:

TLC R_f 0.36 (5% acetic acid, 65% ethyl acetate in dichloromethane)

^1H NMR δ 0.85, 1.2-1.8, 2.45, 3.34, 3.9, 5.56, 5.94 ppm.

MS: 252

PREPARATION 48 3-[(3-Benzoyloxycarbonylaminophenyl)-cyclopropylmethyl]-6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-hydroxy-pyran-2-one (Formula K-6) Refer to Chart K.

A solution of 549 mg of alkylation product of Preparation 47, 970 mg of 3-benzoyloxycarbonylaminophenyl cyclopropyl carbinol, and 60 mg of *p*-toluenesulfonic acid monohydrate in 5 ml of dichloromethane is refluxed through 10 ml of 3Å sieves for 18 hours. Following removal of solvent under reduced pressure, the residue is

2187523

flash chromatographed on silica using 25-100% ethyl acetate in dichloromethane to 5% methanol in ethyl acetate, providing 511 mg of the title compound as a tan solid.

Physical characteristics are as follows:

TLC R_f 0.32 (30% ethyl acetate in dichloromethane)

5 ^1H NMR δ 0.2, 0.5, 0.7, 0.8, 1.3-1.7, 3.27, 3.42, 3.86, 5.13, 5.96, 7.1-7.4 ppm.

MS: 531

PREPARATION 49 3-[(3-Aminophenyl)-cyclopropylmethyl]-6-(1-(tetrahydropyran-4-ylmethyl)-propyl-4-hydroxy-pyran-2-one (Formula K-7) Refer to Chart K.

10 A mixture of 510 mg of the title compound of Preparation 48, 605 mg of ammonium formate, and 100 mg of 5% palladium on carbon in 8 ml of methanol is stirred under argon for three hours, then filtered through diatomaceous earth. The filtrate is concentrated under reduced pressure, and the residue flash chromatographed on silica using 2-4% methanol in dichloromethane to afford 280 mg
15 of the title amine as a white solid.

Physical characteristics are as follows:

TLC R_f 0.33 (5% methanol in dichloromethane)

^1H NMR δ 0.24, 0.42, 0.53, 0.68, 0.84, 1.1-1.7, 2.35, 3.33, 3.6, 3.9, 5.82, 6.5, 6.83, 6.9, 7.11 ppm.

20 EXAMPLE 135 N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide (Formula K-8) Refer to Chart K.

To a stirred solution of 60 mg of the amine of Preparation 49 and 24 μL of pyridine in 0.5 ml of dichloromethane is added 27 mg of 1-methylimidazole-4-sulfonyl chloride. After 18 hours the reaction is flashed on silica using 3-6%
25 methanol in dichloromethane to afford 70 mg of the title compound as a white solid.

Physical characteristics are as follows:

TLC R_f 0.24 (5% methanol in dichloromethane)

^1H NMR δ 0.12, 0.26, 0.45, 0.60, 0.82, 1.1-1.9, 2.3, 3.3, 3.58, 3.9, 6.00, 6.9-7.5
30 ppm.

HRMS: 541.2238

PREPARATION 50 4-Hydroxy-6-propylpyran-2-one (Formula K-9) Refer to Chart K.

To a cold (-78°), stirred solution of 6.3 ml of diisopropylamine in 40 ml of dry tetrahydrofuran, under argon, is added 27.5 ml of a 1.6 M solution of butyllithium in
35 hexane. The solution is brought to 0° , and into this is cannulated a solution of 2.52

- g of 4-hydroxy-6-methyl-2-pyrone of formula K-10 in 20 ml of hexamethylphosphoric triamide. The deep red solution is stirred 30 minutes at 0°, then cooled to -45° for the addition of 1.5 ml of ethyl bromide. The solution is warmed to 0° and quenched with 60 ml of 1N aqueous hydrochloric acid. Tetrahydrofuran is removed under reduced pressure and the residue extracted five times with ethyl acetate. The organic phase is washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. Flash chromatography of the residue on silica gel using 4% acetic acid and 16% ethyl acetate in dichloromethane provides 2.34 g of the title compound as a waxy yellow solid.
- Physical characteristics are as follows:
TLC R_f 0.29 (5% acetic acid and 15% ethyl acetate in dichloromethane)
¹H NMR δ 0.98, 1.6, 2.4, 5.63, 6.05
- PREPARATION 51 4-Hydroxy-6-phenethyl-2H-pyran-2-one (Formula L-2) Refer to Chart L.
- To a flame-dried flask containing a stirred solution of 0.90 mL of diisopropylamine in 6 mL of anhydrous tetrahydrofuran at -78°C under an argon atmosphere is added 4.0 mL of a 1.6 M solution of n-butyllithium in hexane. The resulting solution is allowed to warm to 0°C for 20 min, and is then treated via cannula with a solution of 378 mg of commercially available 4-hydroxy-6-methyl-2-pyrone of formula L-1 in 15 mL of tetrahydrofuran. The resulting red, thick slurry is slowly treated with 6.0 mL of distilled hexamethylphosphoramide and allowed to stir for 30 min. The red, cloudy solution is then treated with 0.36 mL of benzyl bromide. The reaction quickly becomes a deep orange solution and is allowed to stir at 0°C for an additional 60 min. The mixture is quenched with excess 1 N aqueous hydrochloric acid and the resulting yellow, biphasic mixture is concentrated to remove the tetrahydrofuran. The resulting mixture is partitioned between dichloromethane and water and the acidic aqueous phase is further extracted with additional portions of dichloromethane. The combined organic phase is dried over magnesium sulfate and then concentrated under reduced pressure. The resulting material is diluted with a large volume of diethyl ether and washed with dilute aqueous hydrochloric acid. The ethereal phase is washed with two additional portions of aqueous hydrochloric acid, once with brine, dried over magnesium sulfate, and finally concentrated under reduced pressure. The residue is flash column chromatographed on silica gel 60 (230-400 mesh) eluting with 1% acetic acid and 20% to 40% ethyl acetate in dichloromethane to give 440 mg of the title compound as a tan solid.

Physical characteristics are as follows:

^1H NMR δ 2.7, 3.0, 5.46, 5.84, 7.1-7.3.

TLC R_f 0.38 (1% acetic acid and 25% ethyl acetate in dichloromethane.)

MP 137-138 °C.

5 PREPARATION 52 6-(α -Ethyl-phenethyl)-4-hydroxy-2H-pyran-2-one (Formula L-3)

Refer to Chart L.

To a cold (-78°C) stirred solution of 0.29 ml of diisopropylamine in 4 ml of dry tetrahydrofuran, under argon, is added 1.2 ml of a 1.6 M solution of n-butyllithium in hexane. The solution is warmed to 0°C, kept at that temperature for ten minutes,
10 then cooled to -30°C. Into this solution is cannulated a solution of 189 mg of the title compound of Preparation 51 in 4 ml of tetrahydrofuran. The resulting heterogeneous mixture is warmed to 0°, and sufficient hexamethylphosphoramide (ca 1 ml) is added to render the mixture mostly homogeneous. After the mixture is stirred for 30 minutes at 0°C, 77 μL of ethyl iodide is added dropwise. After another 90 minutes,
15 the reaction is quenched with excess 1N aqueous hydrochloric acid, and tetrahydrofuran is removed under reduced pressure. The residue is extracted with three portions of ethyl acetate, and the combined organic extract washed with dilute aqueous hydrochloric acid, dried over magnesium sulfate, and concentrated under reduced pressure. The residue is flash chromatographed on silica gel 60 (230-400
20 mesh) using 1% acetic acid and 25% ethyl acetate in dichloromethane to provide 182 mg of the title compound.

Physical characteristics are as follows:

^1H NMR δ 0.85, 1.6, 2.6, 2.9, 5.59, 5.86, 7.0-7.3.

FAB MS $[m+H]=245.1185$.

25 TLC R_f 0.33 (1% acetic acid and 25% ethyl acetate in dichloromethane.)

PREPARATION 53 3-(α -Cyclopropyl-*meta*-(benzyloxycarbonylamino)benzyl)-6-(α -ethyl-phenethyl)-4-hydroxy-2H-pyran-2-one (Formula L-4) Refer to Chart L.

A mixture of 181 mg of the title compound of Preparation 52, 220 mg of the
30 compound of formula F-5, 28 mg of *p*-toluenesulfonic acid monohydrate, and 600 mg of 3Å molecular sieves in 2 ml of benzene is refluxed under argon for 21 hours, then cooled and filtered through Celite. The filtrate is concentrated under reduced pressure, and the residue flash chromatographed on silica gel 60 (230-400 mesh) using 50-100% ethyl acetate in hexane to provide 250 mg of a mixture of materials.
35 This is re-subjected to silica gel chromatography, using 5-20% ethyl acetate in

dichloromethane, to afford 154 mg (40%) of the title compound.

Physical characteristics are as follows:

^1H NMR δ 0.26, 0.48, 0.67, 0.81, 1.6, 1.8, 2.5, 2.7, 2.9, 3.48, 5.14, 5.86, 6.81, 7.0-7.5, 9.46.

5 EI HRMS m/z =523.2350.

TLC R_f 0.27 (5% ethyl acetate in dichloromethane.)

PREPARATION 54 3-(α -Cyclopropyl-*meta*-aminobenzyl)-6-(α -ethyl-phenethyl)-4-hydroxy-2H-pyran-2-one (Formula L-5) Refer to Chart L.

10 A mixture of 146 mg of the title compound of Preparation 53 and 50 mg of 5% palladium on carbon in 2 ml of methanol is shaken under 40 psi of hydrogen for two hours, then filtered through Celite. The filtrate is concentrated under reduced pressure to give 105 mg (96%) of the title compound.

Physical characteristics are as follows:

^1H NMR δ 0.25, 0.5, 0.65, 0.81, 1.6, 2.5, 2.7, 2.9, 3.4, 5.79, 6.5, 6.8-7.3.

15 TLC R_f 0.38 (30% ethyl acetate in dichloromethane).

EXAMPLES 136-150

Utilizing procedures analogous to those described above, and reacting the compound of formula L-5 with the appropriate sulfonyl chloride, the following additional compounds of the present invention are prepared. Individual
20 stereoisomers are prepared by chiral HPLC resolution of intermediates such as the compounds of formulas L-3, L-4, L-5 and L-6. (Refer to Chart L).

136) N-(3-{Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide

Physical characteristics are as follows:

25 TLC R_f 0.29 (5% methanol in dichloromethane)

^1H NMR δ 0.2, 0.5, 0.65, 0.86, 1.63, 1.80, 2.51, 2.8, 3.3, 3.62, 5.7, 6.8-7.4 ppm.

HRMS: 533.1998

137) N-(3-(R or S)-{Cyclopropyl-[6-(1-(R)-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-4-sulfonamide

30 Physical characteristics are as follows:

TLC R_f 0.32 (5% methanol in dichloromethane)

^1H NMR δ 0.18, 0.43, 0.63, 0.83, 1.6, 1.75, 2.5, 2.7-2.9, 3.3, 3.55, 5.76, 6.9-7.4 ppm.

HRMS: 533.1983

35 138) N-(3-(R or S)-{Cyclopropyl-[6-(1-(R)-ethylphenethyl)-4-hydroxy-2-oxo-

2H-pyran-3-yl]-methyl)-phenyl)-1-methyl-1H-imidazole-4-sulfonamide

Physical characteristics are as follows:

TLC R_f 0.30 (5% methanol in dichloromethane)

^1H NMR δ 0.2, 0.5, 0.65, 0.86, 1.63, 1.80, 2.51, 2.8, 3.3, 3.62, 5.7, 6.8-7.4 ppm.

5 HRMS: 533.1993

139) N-(3-(R or S)-(Cyclopropyl-[6-(1-(S)-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl)-phenyl)-1-methyl-1H-imidazole-4-sulfonamide

Physical characteristics are as follows:

TLC R_f 0.30 (5% methanol in dichloromethane)

10 ^1H NMR δ 0.2, 0.5, 0.65, 0.86, 1.63, 1.80, 2.51, 2.8, 3.3, 3.62, 5.7, 6.8-7.4 ppm.

HRMS: 533.1993

140) N-(3-(R or S)-(Cyclopropyl-[6-(1-(S)-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl)-phenyl)-1-methyl-1H-imidazole-4-sulfonamide

Physical characteristics are as follows:

15 TLC R_f 0.30 (5% methanol in dichloromethane)

^1H NMR δ 0.17, 0.44, 0.62, 0.83, 1.6, 1.75, 2.50, 2.7-3.0, 3.3, 3.53, 5.80, 6.9-7.4 ppm.

HRMS: 533.1990

20 141) N-(3-(R or S)-(Cyclopropyl-[6-(1-(R)-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl)-phenyl)-2-pyridinesulfonamide

Physical characteristics are as follows:

TLC R_f 0.34 (30% ethyl acetate in dichloromethane)

^1H NMR δ 0.2, 0.45, 0.6, 0.86, 1.5-1.9, 2.5, 2.8-3.0, 3.2, 5.7, 6.9-7.4, 7.8, 8.6 ppm.

25 MS: 530

142) N-(3-(R or S)-(Cyclopropyl-[6-(1-(R)-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl)-phenyl)-2-pyridinesulfonamide

Physical characteristics are as follows:

TLC R_f 0.35 (30% ethyl acetate in dichloromethane)

30 ^1H NMR δ 0.11, 0.20, 0.43, 0.58, 0.85, 1.5-1.8, 2.5, 2.7-3.0, 3.3, 5.69, 6.9-7.4, 7.8, 8.6 ppm.

MS: 530

143) N-(3-(R or S)-(Cyclopropyl-[6-(1-(R)-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl)-phenyl)-1-methyl-1H-imidazole-2-sulfonamide

35 Physical characteristics are as follows:

TLC R_f 0.34 (5% methanol in dichloromethane)

¹H NMR δ 0.19, 0.5, 0.65, 0.89, 1.6-1.9, 2.5, 2.8-3.0, 3.3, 3.40, 5.70, 6.8-7.4 ppm.

MS: 533

- 5 144) N-(3-(R or S)-{Cyclopropyl-[6-(1-(R)-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-2-sulfonamide

Physical characteristics are as follows:

TLC R_f 0.34 (5% methanol in dichloromethane)

10 ¹H NMR δ 0.20, 0.44, 0.65, 0.88, 1.6-1.8, 2.5, 2.8-3.0, 3.3, 3.42, 5.73, 6.8-7.4 ppm.

MS: 533

- 145) N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-2-sulfonamide

Physical characteristics are as follows:

15 TLC R_f 0.22 (5% methanol in dichloromethane)

¹H NMR δ 0.16, 0.24, 0.47, 0.64, 0.86, 1.2-1.9, 3.2-3.4, 3.47, 3.7-4.0, 5.89, 6.9-7.4 ppm.

MS: 541

- 20 145A) N-(3-[Cyclopropyl[4-hydroxy-2-oxo-6-[1-[(tetrahydro-2H-pyran-3-yl)methyl]propyl]-2H-pyran-3-yl]methyl]phenyl]-8-quinolinesulfonamide

Physical characteristics are as follows:

MW Found: m/z 588.

- 25 145B) N-(3-[Cyclopropyl[4-hydroxy-2-oxo-6-[1-[(tetrahydro-2H-pyran-3-yl)methyl]propyl]-2H-pyran-3-yl]methyl]phenyl)-1-methyl-1H-imidazole-4-sulfonamide

Physical characteristics are as follows:

MW Found: m/z 541.

- 30 146) N-(3-{Cyclopropyl-[6-(1-(R)-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-benzimidazole-2-sulfonamide

Physical characteristics are as follows:

TLC R_f 0.40 (50% ethyl acetate in dichloromethane)

¹H NMR δ 0.1-0.6, 0.85, 1.5-1.7, 2.5, 2.7-3.0, 3.3, 5.74, 6.7-7.3, 7.5-7.7 ppm.

HRMS: 570.2054

- 35 147) N-(3-{Cyclopropyl-[6-(1-(R)-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1H-imidazole-2-sulfonamide

Physical characteristics are as follows:

TLC R_f 0.31 (5% methanol in dichloromethane)

^1H NMR δ 0.2, 0.4, 0.6, 0.87, 1.5-1.8, 2.5, 2.8-3.0, 3.3, 5.54, 6.8, 6.9-7.4 ppm.

148) N-(3-(R or S)-[Cyclopropyl-[6-(1-(R)-ethylphenethyl)-4-hydroxy-2-oxo-
5 2H-pyran-3-yl]-methyl]-phenyl)-4-cyanobenzenesulfonamide

Physical characteristics are as follows:

TLC R_f 0.47 (20% ethyl acetate in dichloromethane)

^1H NMR δ 0.1, 0.2, 0.4, 0.6, 0.84, 1.5-1.8, 2.5, 2.7-3.0, 3.3, 5.70, 6.9, 7.0-7.3,
7.6, 7.8 ppm.

10 HRMS: 554.1886

149) N-(3-(R or S)-[Cyclopropyl-[6-(1-(R)-ethylphenethyl)-4-hydroxy-2-oxo-
2H-pyran-3-yl]-methyl]-phenyl)-4-cyanobenzenesulfonamide

Physical characteristics are as follows:

TLC R_f 0.35 (15% ethyl acetate in dichloromethane)

15 ^1H NMR δ 0.1, 0.2, 0.4, 0.6, 0.85, 1.5-1.9, 2.5, 2.7-3.0, 3.3, 5.7, 6.9-7.3, 7.6, 7.8
ppm.

HRMS: 554.1876

150) N-(3-[Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-
yl]-methyl]-phenyl)-4-nitrobenzenesulfonamide

20 Physical characteristics are as follows:

TLC R_f 0.28 (10% ethyl acetate in dichloromethane)

^1H NMR δ 0.1, 0.2, 0.4, 0.6, 0.83, 1.5-1.9, 2.5, 2.7-3.0, 3.3, 5.70, 6.9-7.3, 7.9,
8.2 ppm.

HRMS: 574.1773

25 PREPARATION 55 (2-(2-(2-Methoxyethoxy)-ethoxy)-ethoxy)-*p*-toluenesulfonate
(Formula M-2) Refer to Chart M.

To a stirred suspension of 19.1 g of *p*-toluenesulfonyl chloride in 100 ml of
dichloromethane is added a mixture of 16 ml of triethylene glycol monomethyl ether
and 10 ml of pyridine, followed by 200 mg of dimethylaminopyridine. After three
30 days the mixture is concentrated under reduced pressure, and the residue
partitioned between ethyl acetate and dilute aqueous hydrochloric acid. The organic
phase is washed with water, aqueous sodium bicarbonate, and brine, and dried over
magnesium sulfate. After removal of solvent under reduced pressure, the residue is
flash chromatographed on silica using 25% ethyl acetate in dichloromethane to
35 afford 18.25 g of the title compound as a colorless liquid.

Physical characteristics are as follows:

TLC R_f 0.27 (20% ethyl acetate in dichloromethane)

^1H NMR δ 2.45, 3.38, 3.5-3.8, 4.15, 7.35, 7.8 ppm.

IR 2879, 1357, 1190, 1177, 1108, 1099, 924, 665 cm^{-1}

5 MS: 318

PREPARATION 56 2-Hydroxy-4-(2-(2-(2-methoxyethoxy)-ethoxy)-ethoxy)-
acetophenone (Formula M-3) Refer to Chart M.

A mixture of 1.52 g of 2,4-dihydroxyacetophenone, 3.82 g of the tosylate of
Preparation 55, 3.26 g of cesium carbonate, and 0.2 g of potassium iodide in 20 ml of
10 dioxane is heated overnight at 100°, then cooled and partitioned between
dichloromethane and dilute aqueous hydrochloric acid. The aqueous phase is
extracted with two additional portions of dichloromethane, and the combined organic
phase dried over magnesium sulfate and then concentrated under reduced pressure.
Flash chromatography of the residue on silica gel using 80-100% ethyl acetate in
15 hexane provides 2.91 g of the title compound as a nearly colorless liquid.

Physical characteristics are as follows:

TLC R_f 0.35 (80% ethyl acetate in hexane)

^1H NMR δ 2.56, 3.38, 3.5-3.9, 4.2, 6.4-6.5, 7.6 ppm.

IR 1635, 1372, 1257, 1133 cm^{-1}

20 MS: 298

PREPARATION 57 3-(2-Hydroxy-4-(2-(2-(2-methoxyethoxy)-ethoxy)-ethoxy)-phenyl)-
3-oxopropionic acid ethyl ester (Formula M-4) Refer to Chart M.

To a stirred solution of 1.49 g of the title compound of Preparation 56 in 10
ml of diethyl carbonate is added, in portions, 600 mg of 60% sodium hydride
25 dispersion in mineral oil. The resulting mixture is heated at 80° for two hours, then
cooled and partitioned between dichloromethane and dilute aqueous hydrochloric
acid. The organic phase is dried over magnesium sulfate and concentrated under
reduced pressure, and the residue purified by flash chromatography on silica gel
using 20-30% ethyl acetate in dichloromethane to afford 0.91 g of the title compound
30 as a yellow oil.

Physical characteristics are as follows:

TLC R_f 0.44 (3% acetic acid and 30% ethyl acetate in dichloromethane)

^1H NMR δ 1.3, 3.38, 3.5-4.0, 4.2, 6.4-6.5, 7.6 ppm.

MS: 370

35 PREPARATION 58 4-Hydroxy-(2-(2-(2-methoxyethoxy)-ethoxy)-ethoxy)-coumarin

(Formula M-5) Refer to Chart M.

A solution of 789 mg of the title compound of Preparation 57 in 10 ml of acetic acid is refluxed for two hours, then concentrated under reduced pressure. Flash chromatography of the residue on silica using 5-10% acetic acid in ethyl acetate provides 634 mg of the title compound as a buff colored solid.

Physical characteristics are as follows:

TLC R_f 0.31 (10% acetic acid in ethyl acetate)

^1H NMR δ 3.37, 3.5-3.9, 4.1, 5.67, 6.6, 6.7, 7.6 ppm.

MS: 324

10 PREPARATION 59 3-[(3-Benzoyloxycarbonylamino-phenyl)-cyclopropylmethyl]-4-hydroxy-7-{2-[2-(2-methoxyethoxy)-ethoxy]-ethoxy}-coumarin
(Formula M-6) Refer to Chart M.

A mixture of 704 mg of the title compound of Preparation 58, 775 mg of *meta*-benzyloxycarbonylamino-phenyl cyclopropyl carbinol of formula F-5, and 62 mg of *p*-toluenesulfonic acid monohydrate in 8 ml of dichloromethane is refluxed for 18 hours through ca. 10 ml of 3Å sieves. The solution is then concentrated under reduced pressure and the residue flash chromatographed on silica gel using 10-20% of (10% acetic acid in ethyl acetate) in dichloromethane to afford 760 mg of the title compound.

20 Physical characteristics are as follows:

TLC R_f 0.33 (2% acetic acid and 20% ethyl acetate in dichloromethane)

^1H NMR δ 0.27, 0.46, 0.71, 1.61, 3.33, 3.5-3.9, 4.1, 5.13, 6.6, 6.7, 7.1-7.6 ppm.

PREPARATION 60 3-[(3-Amino-phenyl)-cyclopropylmethyl]-4-hydroxy-7-{2-[2-(2-methoxyethoxy)-ethoxy]-ethoxy}-coumarin (Formula M-7) Refer
25 to Chart M.

A solution of 760 mg of the title compound of Preparation 59, 800 mg of ammonium formate, and 200 mg of 5% palladium on charcoal catalyst in 8 ml of methanol is stirred under argon for one hour, then filtered through a pad of diatomaceous earth. The filtrate is concentrated under reduced pressure and the
30 residue triturated with dichloromethane. Removal of solvent under reduced pressure provides 591 mg of the title amine.

Physical characteristics are as follows:

TLC R_f 0.29 (5% methanol in dichloromethane)

EXAMPLE 151 N-(3-{Cyclopropyl-[7-(2-(2-(2-methoxyethoxy)-ethoxy)ethoxy)-4-hydroxycoumarin-3-yl]-methyl}-phenyl)-1-methyl-1H-imidazole-
35

4-sulfonamide (Formula M-8) Refer to Chart M.

To a stirred solution of 70 mg of the title compound of Preparation 60 and 24 μ L of pyridine in 0.5 ml of dichloromethane is added 27 mg of 1-methylimidazole-4-sulfonyl chloride. After 18 hours, the solution is flash chromatographed on silica gel using 5-15% methanol in dichloromethane, affording 76 mg of the title sulfonamide as a pink amorphous foam.

Physical characteristics are as follows:

TLC R_f 0.21 (5% methanol in dichloromethane)

^1H NMR δ 0.16, 0.29, 0.45, 0.61, 1.71, 3.34, 3.4-3.9, 4.1, 6.6-6.8, 7.0-7.4, 7.7 ppm.

HRMS: 614.2179

EXAMPLES 152-154

Utilizing procedures analogous to those described above, the following additional compounds of the present invention are prepared:

152) N-(3-(Cyclopropyl-[7-methoxy-4-hydroxycoumarin-3-yl]-methyl)-phenyl)-1-methyl-1H-imidazole-4-sulfonamide

Physical characteristics are as follows:

TLC R_f 0.29 (5% methanol in dichloromethane)

^1H NMR δ 0.18, 0.35, 0.50, 0.63, 1.61, 3.51, 3.7, 3.84, 6.7-6.8, 7.1-7.4, 7.7 ppm.

HRMS: 481.1301

153) N-(3-(Cyclopropyl-[7-(2-(2-(2-methoxyethoxy)-ethoxy)ethoxy)-4-hydroxycoumarin-3-yl]-methyl)-phenyl)-8-quinolinesulfonamide

Physical characteristics are as follows:

TLC R_f 0.41 (5% methanol in dichloromethane)

^1H NMR δ -0.03, 0.31, 0.47, 1.30, 3.36, 3.5-3.8, 3.9, 4.2, 6.6-7.6, 7.8, 8.0, 8.2 ppm.

HRMS: 661.2219

154) N-(3-(Cyclopropyl-[7-(2-(2-(2-methoxyethoxy)-ethoxy)ethoxy)-4-hydroxycoumarin-3-yl]-methyl)-phenyl)-2-pyridinesulfonamide

Physical characteristics are as follows:

TLC R_f 0.31 (5% methanol in dichloromethane)

^1H NMR δ 0.13, 0.34, 0.49, 0.63, 1.6, 3.36, 3.5-3.9, 4.1, 6.68, 6.8, 7.1-7.4, 7.6-7.8, 8.5 ppm.

HRMS: 611.2051

EXAMPLES 155-190

The following additional compounds of the present invention are prepared by procedures analogous to those described above:

- 155) N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-
5 hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-2-pyridinesulfonamide
- 156) N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-
hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-4-cyano-2-pyridinesulfonamide
- 157) N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-
hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-2-quinolinesulfonamide
- 10 158) N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-
hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-2-hydroxybenzenesulfonamide
- 159) N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-
hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-2-pyrazolesulfonamide
- 160) N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-
15 hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-2-quinazolinesulfonamide
- 161) N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-
hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-7H-purine-6-sulfonamide
- 162) N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-
hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1H-imidazole-2-sulfonamide
- 20 163) N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-
hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-benzimidazole-2-sulfonamide
- 164) N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-
hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-thiazole-4-sulfonamide
- 25 165) N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-
hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-4-ethoxycarbonyl-1H-imidazole-2-
sulfaonamide
- 166) N-(3-{Cyclopropyl-[6-(1-(tetrahydropyran-4-ylmethyl)-propyl)-4-
hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-3-hydroxy-2-pyridinesulfonamide
- 167) N-(3-{Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-
30 yl]-methyl}-phenyl)-2-pyridinesulfonamide
- 168) N-(3-{Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-
yl]-methyl}-phenyl)-4-cyano-2-pyridinesulfonamide
- 169) N-(3-{Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-
yl]-methyl}-phenyl)-2-quinolinesulfonamide
- 35 170) N-(3-{Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-

- yl]-methyl}-phenyl)-2-hydroxybenzenesulfonamide
- 171) N-(3-{Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-2-pyrazolesulfonamide
- 172) N-(3-{Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-2-quinazolinesulfonamide
- 5 173) N-(3-{Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-7H-purine-6-sulfonamide
- 174) N-(3-{Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-1H-imidazole-2-sulfonamide
- 10 175) N-(3-{Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-benzimidazole-2-sulfonamide
- 176) N-(3-{Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-thiazole-4-sulfonamide
- 177) N-(3-{Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-4-ethoxycarbonyl-1H-imidazole-2-sulfonamide
- 15 178) N-(3-{Cyclopropyl-[6-(1-ethylphenethyl)-4-hydroxy-2-oxo-2H-pyran-3-yl]-methyl}-phenyl)-3-hydroxy-2-pyridinesulfonamide
- 179) N-(3-{Cyclopropyl-[4-hydroxycoumarin-3-yl]-methyl}-phenyl)-2-pyridinesulfonamide
- 20 180) N-(3-{Cyclopropyl-[4-hydroxycoumarin-3-yl]-methyl}-phenyl)-4-cyano-2-pyridinesulfonamide
- 181) N-(3-{Cyclopropyl-[4-hydroxycoumarin-3-yl]-methyl}-phenyl)-2-quinolinesulfonamide
- 182) N-(3-{Cyclopropyl-[4-hydroxycoumarin-3-yl]-methyl}-phenyl)-2-hydroxybenzenesulfonamide
- 25 183) N-(3-{Cyclopropyl-[4-hydroxycoumarin-3-yl]-methyl}-phenyl)-2-pyrazolesulfonamide
- 184) N-(3-{Cyclopropyl-[4-hydroxycoumarin-3-yl]-methyl}-phenyl)-2-quinazolinesulfonamide
- 30 185) N-(3-{Cyclopropyl-[4-hydroxycoumarin-3-yl]-methyl}-phenyl)-7H-purine-6-sulfonamide
- 186) N-(3-{Cyclopropyl-[4-hydroxycoumarin-3-yl]-methyl}-phenyl)-1H-imidazole-2-sulfonamide
- 187) N-(3-{Cyclopropyl-[4-hydroxycoumarin-3-yl]-methyl}-phenyl)-benzimidazole-2-sulfonamide
- 35

188) N-(3-{Cyclopropyl-[4-hydroxycoumarin-3-yl]-methyl}-phenyl)-thiazole-4-sulfonamide

189) N-(3-{Cyclopropyl-[4-hydroxycoumarin-3-yl]-methyl}-phenyl)-4-ethoxycarbonyl-1H-imidazole-2-sulfonamide

5 190) N-(3-{Cyclopropyl-[4-hydroxycoumarin-3-yl]-methyl}-phenyl)-3-hydroxy-2-pyridinesulfonamide

PREPARATION 61 Cyclopropyl-(3-nitrophenyl)methanone (Formula N-2) Refer to Chart N.

Charge a jacketed 1 L three neck round bottom flask equipped with stirrer
10 and addition funnel under nitrogen with 580 mL fuming nitric acid and cool to -40°C. Slowly, over 1.5 hours, add cyclopropyl phenyl ketone of formula N-1 (100 g) keeping the temperature below -35°C. Stir 3 hours, monitoring reaction by TLC. Pour reaction mixture into 3 kg ice/water. Extract with 3X500 mL ethyl acetate. Wash combined organic phase with 2X1.5 L saturated aqueous sodium bicarbonate,
15 dry over magnesium sulfate, filter and concentrate to 138 g. Dissolve residue in 270 mL methanol, cool to -20°C for 18 hours, filter and wash cake with cold methanol. Dry product under reduced pressure for 72 hours, obtaining 63.86 g. GC analysis (15 m. DB-1, $T_0 = 100^\circ\text{C}$, $10^\circ\text{C}/\text{min.}$, RT - 6.0 min.) indicates material to be >98% pure.

20 Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 8.86, 8.43, 8.34, 7.70, 2.72, 1.33, 1.17 ppm.

IR (Nujol) 2954, 2925, 1664, 1614, 1529, 1442, 1386, 1352, 1225, 1082, 1047, 852, 720, 689 cm^{-1} .

Elemental analysis, Found: C, 62.89; H, 4.73; N, 7.32.

25 MS (EI) 191, 150, 104, 69 m/z.

PREPARATION 62 Cyclopropyl-(3-aminophenyl)methanone (Formula N-3) Refer to Chart N.

Charge platinum on carbon (8.7 g) to Paar bottle. Charge a flask with cyclopropyl(3-nitrophenyl)methanone of Preparation 61 (86.7 g) and methanol (1.56
30 L) and warm to dissolve, then cool with ice bath to 9°C . Hydrogenate for 50 minutes, keeping temperature below 35°C and monitoring reaction by TLC. Filter reaction mixture through solka floc, and concentrate under reduced pressure to 70 g.

Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 7.99, 7.47-7.19, 6.84, 3.84, 2.60, 1.23-1.15, 1.03-0.96 ppm.

35 ^{13}C NMR (CDCl_3) δ 200.9, 146.8, 139.1, 129.4, 119.3, 118.4, 113.9, 17.2, 11.6

ppm.

PREPARATION 63 Cyclopropyl-(3-aminocarbobenzoxyphenyl)methanone (Formula N-4) Refer to Chart N.

Charge a 3 L round bottom flask equipped with mechanical stirrer and addition funnel under nitrogen with cyclopropyl-(3-aminophenyl)methanone of Preparation 62 (70.0 g), diisopropylethylamine (DIPEA, 90.2 mL) and methylene chloride (CH_2Cl_2) (1.3 L). Cool reaction mixture to 0°C . Dilute the benzylchloroformate (67.5 mL) with methylene chloride (186 mL) and add to the substrate solution over one hour keeping temperature at $0-5^\circ\text{C}$. A heavy precipitate will form. Allow to warm with stirring for 1.5 hours monitoring reaction by TLC. Pour reaction mixture into 600 mL 1N HCl/600 g ice/4.2 L methylene chloride and stir to dissolve. Separate phases and dry organic phase over magnesium sulfate, filter and concentrate to a dryness. Slurry solids in 3 mL/g hexane, filter, and vacuum dry for 125 g.

Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 8.01, 7.76-7.69, 7.43-7.33, 7.18, 5.21, 2.64, 1.25-1.20, 1.03-0.97 ppm.

^{13}C NMR (CDCl_3) δ 200.6, 153.4, 138.7, 138.5, 135.9, 129.3, 128.6, 128.4, 123.1, 122.8, 118.1, 67.2, 17.3, 12.0 ppm.

PREPARATION 64 Cyclopropyl-(3-aminocarbobenzoxyphenyl)methanol (Formula N-5) Refer to Chart N.

Charge a 2 L three neck round bottom flask equipped with overhead stirrer under nitrogen with cyclopropyl-(3-aminocarbobenzoxyphenyl)methanone of Preparation 63 (25 g), tetrahydrofuran (THF) (450 mL) and ethanol (90 mL). Cool reaction mixture to $0-5^\circ\text{C}$ and add the sodium borohydride pellets (12.4 g) in three equal portions over 30 minutes. Allow to warm to 23°C and stir for 20 hours, monitoring reaction by TLC. Recool reaction mixture to $0-5^\circ\text{C}$ and slowly quench by adding 90 mL 1N hydrochloric acid, keeping the temperature below 10°C . Pour with stirring into methylene chloride (600 mL) and 1N hydrochloric acid (400 mL). Separate the phases and wash the organic phase with saturated sodium chloride solution (1 L). Dry over magnesium sulfate, filter, and concentrate to 23.7 g.

Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 7.41-7.35, 7.33, 7.17, 7.10, 5.17, 3.93, 2.36, 1.16-1.12, 0.60-0.32 ppm.

^{13}C NMR (CDCl_3) δ 153.5, 145.0, 137.9, 136.1, 129.0, 128.6, 128.3, 121.2,

117.9, 116.5, 67.9, 67.0, 19.1, 3.6, 2.8 ppm.

PREPARATION 65 Carbamic acid, [3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-, phenylmethyl ester (Formula N-6) Refer to Chart N.

5 A 12-L, three-necked, round-bottomed flask with a Soxhlet extractor containing 3 Å molecular sieves (180 g) and nitrogen inlet is charged with cyclooctene-1-acrylic acid, β , 2-dihydroxy- δ -lactone (59.6 g), *p*-toluenesulfonic acid (14.9 g), and methylene chloride (7.2 L). The title compound of Preparation 64 (90.0 g) is added, and the reaction mixture is warmed to reflux for 1 h. The reaction
10 mixture is then cooled to 20° C and washed with 1:1 saturated sodium chloride/saturated sodium bicarbonate (3 L), water (3 L), and saturated sodium chloride (3 L), backwashing each aqueous phase with methylene chloride (2 x 1.5 L). The organic layers are then combined, dried over magnesium sulfate, filtered and concentrated to ca. 1.5 L. The reaction mixture is cooled to -20° C for 72 h, filtered,
15 and dried under reduced pressure to give 103.5 g. The crude product is then slurried with 12.5 mL/g of hexane, filtered, and dried to give 102.4 g of the title compound. An additional 10.9 g of the title compound is obtained by concentrating the mother liquors from the crystallization and recrystallizing the residue from ethyl acetate.

20 Physical characteristics are as follows:

MP 113-115°C (decomposition).

¹H NMR (CDCl₃) δ 7.48, 7.38-7.26, 7.17, 6.70, 6.29, 5.20, 3.95, 2.64-2.60, 2.47-2.43, 1.76-1.72, 1.61-1.42, 0.88, 0.73-0.72, 0.63-0.55, 0.29-0.26 ppm.

¹³C NMR (CDCl₃) δ 165.6, 164.0, 161.3, 142.2, 138.5, 129.9, 128.5, 128.3,
25 128.2, 122.9, 118.0, 117.9, 117.6, 110.7, 106.0, 67.0, 43.7, 30.7, 29.1, 28.8, 26.2, 25.8, 22.1, 13.0, 4.9, 3.8 ppm.

IR (Nujol) 3304, 2995, 2953, 2923, 2855, 1734, 1698, 1665, 1666, 1633, 1610, 1595, 1553, 1491, 1463, 1455, 1445, 1406, 1377, 1313, 1222, 1175, 1085, 1068, 740, 696 cm⁻¹.

30 MS (EI) *m/z* 473, 445, 382, 338, 91.

For high resolution, Found: 473.2202.

PREPARATION 66 3-[(3-Aminophenyl)cyclopropylmethyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one (Formula N-7) Refer to
Chart N.

35 In a 100-mL, three-necked, round-bottomed flask with a reflux condensor and

nitrogen inlet, 10% palladium on carbon (1.0 g) is added to a mixture of the title product of formula N-6, prepared in Preparation 65 (1.95 g) in cyclohexene (50 mL) and the mixture is refluxed for 4h. The mixture is then filtered through Celite, washed with methylene chloride (CH_2Cl_2), and concentrated to give 1.25 g of the title compound as a white solid.

Physical characteristics are as follows:

MP 75-79 °C

IR (Nujol) 2995, 2951, 2921, 2868, 1660, 1619, 1605, 1590, 1551, 1491, 1460, 1447, 1428, 1404, 1247, 1226, 1202, 1191, 1172, 1126 cm^{-1} .

MS (EI) m/z 339, 310, 213, 187, 159.

^1H NMR (CDCl_3) δ 7.16, 6.96, 6.84, 6.63, 5.67, 3.87, 2.61, 2.48-2.37, 1.98, 1.75, 1.63-1.26, 0.74-0.65, 0.61-0.53, 0.28-0.22 ppm.

^{13}C NMR (CDCl_3) δ 164.2, 161.1, 142.8, 130.2, 117.7, 117.6, 114.7, 114.6, 114.5, 110.9, 106.2, 43.5, 30.6, 29.1, 28.8, 26.2, 25.8, 22.0, 12.8, 4.7, 3.7 ppm.

For high resolution, Found: 339.1845.

PREPARATION 67 4-Cyano-N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-benzenesulfonamide (Formula O-3 wherein R_{61} is 4-cyanophenyl) Refer to Chart O.

A solution of the title product of Preparation 66 (660 mg), pyridine (320 μL), and 4-cyanobenzenesulfonyl chloride (440 mg) in dichloromethane (40 mL) is stirred at room temperature for 18 hr. The crude reaction mixture is evaporated to a volume of 5 ml and chromatographed on silica gel using 50% ethyl acetate in hexane as eluent to give the title compound (641mg) as a white amorphous solid. This amorphous solid is alternatively crystallized from acetone:hexane to give 499mg.

Physical characteristics are as follows:

White solid mp: 183-183.5°C

Elemental analysis: found, C, 66.76; H, 5.68; N, 5.38; s, 6.30.

MS(EI): 504, 476, 463, 338, 309, 233, 220, 207, 195, 186, 153, 144, 130, 117, 102.

HRMS: 504.1710.

TLC(silica gel GF): R_f = 0.4 in 50% ethyl acetate in hexane.

EXAMPLE 191 Disodium-4-cyano-N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-benzenesulfonamide

To 12.6 g of the title product of Preparation 67 is added 500 ml of methanol and, with rapid stirring, 50 ml of a 1N aqueous NaOH solution. The reaction solution is allowed to stir at room temperature for 1 hour. The yellow solution is evaporated to dryness at 35°C and the resulting amorphous residue is dissolved in absolute ethanol and re-evaporated to dryness. The yellow residue is kept under high vacuum at room temperature for 18 hours to yield 14g of a yellow amorphous solid.

Physical characteristics are as follows:

TLC(silica gel GF): R_f =0.8 streak from the origin (20% ethylacetate in methylene chloride)

K.F. Water: 6.16%

Melt Solvate: 4.2% ethanol

Weight Loss at Room Temperature: 4.99%

Ash: found: 7.83% ; Calc'd: 7.50% (corrected for 6.16% water and 4.2% ethanol)

PREPARATION 68 N-methyl-3[(3-aminophenyl)cyclopropylmethyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one

To 678 mg of the title product of Preparation 66 is added 100 ml of absolute ethanol and 330 mg of 10% Pd/C. 183 microliters of a 35% $\text{CH}_2\text{O}/\text{H}_2\text{O}$ solution is added and the mixture allowed to shake on a Paar apparatus, under 50 lbs of hydrogen, for 2 hours at room temperature. The reaction is filtered over celite and the filter cake is washed well with ethanol. The resulting amber solution is evaporated to dryness. The resulting residue is chromatographed using 10% ethyl acetate in methylene chloride to give 110 mg of the title product. This material is used without further purification in the synthesis of the following sulfonamides.

Physical characteristics are as follows:

TLC(silica gel GF): R_f =0.5 in 10% ethyl acetate in methylene chloride.

^1H NMR (CDCl_3) δ 7.19, 6.90, 6.71, 6.54-6.52, 3.90, 2.80, 2.63-2.59, 2.43-2.39, 1.75-1.26, .70-.53, .28-.22 ppm.

EXAMPLE 192 4-Cyano-N-methyl-N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]benzenesulfonamide

A solution of the title compound of Preparation 68 (35 mg), pyridine (16 μL), and 4-cyanobenzenesulfonyl chloride (20.1 mg) in dichloromethane (2 mL) is stirred at room temperature for 18 hr. The crude reaction mixture is chromatographed on

silica gel using 10% ethyl acetate in methylene chloride as eluent to give 27mg of the title compound as a white amorphous solid.

Physical characteristics are as follows:

MS(EI): 518, 490, 352, 233, 207, 172, 158, 143, 129, 115, 102, 81, 54, 43.

5 TLC(silica gel GF): R_f =0.7 in 10% ethyl acetate in methylene chloride.

^1H NMR (CDCl_3) δ 7.75-7.72, 7.63-7.60, 7.38-7.19, 6.97-6.94, 6.62, 3.86, 3.19, 2.66-2.62, 2.54-2.50, 1.76-1.20, .70-.59, .47-.42, .24-.19 ppm.

EXAMPLE 193 4-Fluoro-N-methyl-N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-benzenesulfonamide

10 A solution of the title compound of Preparation 68 (20 mg), pyridine (11 μL), and 4-fluorobenzenesulfonyl chloride (10.7 mg) in dichloromethane (2 mL) is stirred at room temperature for 18 hr. The crude reaction mixture is chromatographed on silica gel using 10% ethyl acetate in methylene chloride as eluent to give 19mg of
15 the title compound as a white amorphous solid.

Physical characteristics are as follows:

MS(EI): 512, 483, 470, 366, 352, 324, 247, 227, 207, 172, 158, 147, 118, 55.

HRMS: Found: 512.1915

TLC(silica gel GF): R_f =0.7 in 10% ethyl acetate in methylene chloride.

20 ^1H NMR (CDCl_3) δ 7.53-7.48, 7.33-7.23, 7.13-7.07, 6.99-6.97, 6.38, 3.93, 3.16, 2.63-2.61, 2.49-2.46, 1.76-1.25, .78-.61, .51-.45, .30-.17 ppm.

EXAMPLE 194 N-methyl-N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-benzenesulfonamide

25 A solution of the title compound of Preparation 68 (33.4 mg), pyridine (16 μL), and benzenesulfonyl chloride (16.6 mg) in dichloromethane (2 mL) is stirred at room temperature for 18 hr. The crude reaction mixture was chromatographed on silica gel using 10% ethyl acetate in methylene chloride as eluent to give 20mg of the title compound as a white amorphous solid.

30 Physical characteristics are as follows:

TLC(silica gel GF): R_f =0.7 in 10% ethyl acetate in methylene chloride.

^1H NMR (CDCl_3) δ 7.59-7.41, 7.33-7.23, 6.98-6.96, 6.44, 3.90, 3.16, 2.64-2.60, 2.50-2.48, 1.75-1.20, .67-.40, .23-.20 ppm.

EXAMPLE 195 N-methyl-N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-1H-Imidazole-1-

methyl-sulfonamide

A solution of the title compound of Preparation 68 (33.4 mg), pyridine (16 μ L), and N-methyl-imidazole-3-sulfonyl chloride (16 mg) in dichloromethane (2 mL) is stirred at room temperature for 18 hr. The crude reaction mixture is chromatographed on silica gel using 50% ethyl acetate in methylene chloride as eluent to give 28mg of the title compound as a white amorphous solid.

Physical characteristics are as follows:

TLC(silica gel GF): R_f =0.5 in 50% ethyl acetate in methylene chloride.

^1H NMR (CDCl_3) δ 7.43, 7.33, 7.27-7.15, 3.84-3.81, 3.69, 3.35, 2.63-2.59, 2.50-2.46, 1.75-1.26, .68, .55, .47-.42, .24-.20 ppm.

Utilizing procedures analogous to those described above, the following compounds of the present invention are prepared:

196) 5-cyano-N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-2-pyridinesulfonamide

197) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-2-quinolinesulfonamide

198) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-2-imidazolesulfonamide

199) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-2-pyrimidinesulfonamide

200) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-2-benzimidazolesulfonamide

201) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-2-quinazolinesulfonamide

202) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-6-purinesulfonamide

203) 5-cyano-N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-N-methyl-2-pyridinesulfonamide

204) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-N-methyl-2-quinolinesulfonamide

205) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-N-methyl-2-imidazolesulfonamide

206) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-N-methyl-2-pyrimidinesulfonamide

207) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-

- cycloocta[b]pyran-3-yl)methyl]phenyl]-2-benzimidazolesulfonamide
- 208) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-2-quinazolinesulfonamide
- 209) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-N-methyl-6-purinesulfonamide
- 5 210) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-N-methyl-4-thiazolesulfonamide
- 211) N-[3-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-N-methyl-2-pyridinesulfonamide
- 10 212) 5-cyano-N-[6-(1'-benzylpropyl)-4-hydroxy-3-(1'-cyclopropylmethylphenyl)-2-pyrone]-N-methyl-2-pyridinesulfonamide
- 213) N-[6-(1'-benzylpropyl)-4-hydroxy-3-(1'-cyclopropylmethylphenyl)-2-pyrone]-N-methyl-2-quinolinesulfonamide
- 214) N-[6-(1'-benzylpropyl)-4-hydroxy-3-(1'-cyclopropylmethylphenyl)-2-pyrone]-N-methyl-2-imidazolesulfonamide
- 15 215) N-[6-(1'-benzylpropyl)-4-hydroxy-3-(1'-cyclopropylmethylphenyl)-2-pyrone]-N-methyl-2-pyrimidinesulfonamide
- 216) N-[6-(1'-benzylpropyl)-4-hydroxy-3-(1'-cyclopropylmethylphenyl)-2-pyrone]-N-methyl-2-benzimidazolesulfonamide
- 20 217) N-[6-(1'-benzylpropyl)-4-hydroxy-3-(1'-cyclopropylmethylphenyl)-2-pyrone]-N-methyl-2-quinazolinesulfonamide
- 218) N-[6-(1'-benzylpropyl)-4-hydroxy-3-(1'-cyclopropylmethylphenyl)-2-pyrone]-N-methyl-6-purinesulfonamide
- 219) N-[6-(1'-benzylpropyl)-4-hydroxy-3-(1'-cyclopropylmethylphenyl)-2-pyrone]-N-methyl-4-thiazolesulfonamide
- 25 220) N-[6-(1'-benzylpropyl)-4-hydroxy-3-(1'-cyclopropylmethylphenyl)-2-pyrone]-N-methyl-2-pyridinesulfonamide
- 221) 5-cyano-N-[3-(1'-cyclopropylmethylphenyl)-4-hydroxycoumarin]-N-methyl-2-pyridinesulfonamide
- 30 222) N-[3-(1'-cyclopropylmethylphenyl)-4-hydroxycoumarin]-N-methyl-2-quinolinesulfonamide
- 223) N-[3-(1'-cyclopropylmethylphenyl)-4-hydroxycoumarin]-N-methyl-2-imidazolesulfonamide
- 224) N-[3-(1'-cyclopropylmethylphenyl)-4-hydroxycoumarin]-N-methyl-2-pyrimidinesulfonamide
- 35

225) N-[3-(1'-cyclopropylmethylphenyl)-4-hydroxycoumarin]-N-methyl-2-benzimidazolesulfonamide

226) N-[3-(1'-cyclopropylmethylphenyl)-4-hydroxycoumarin]-N-methyl-2-quinazolinesulfonamide

5 227) N-[3-(1'-cyclopropylmethylphenyl)-4-hydroxycoumarin]-N-methyl-6-purinesulfonamide

228) N-[3-(1'-cyclopropylmethylphenyl)-4-hydroxycoumarin]-N-methyl-4-thiazolesulfonamide

10 229) N-[3-(1'-cyclopropylmethylphenyl)-4-hydroxycoumarin]-N-methyl-2-pyridinesulfonamide

230) 5-cyano-N-[3-[1-(4-hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-N-methyl-2-pyridinesulfonamide

231) N-[3-[1-(4-hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-N-methyl-2-quinolinesulfonamide

15 232) N-[3-[1-(4-hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-N-methyl-2-imidazolesulfonamide

233) N-[3-[1-(4-hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-N-methyl-2-pyrimidinesulfonamide

20 234) N-[3-[1-(4-hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-N-methyl-2-benzimidazolesulfonamide

235) N-[3-[1-(4-hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-N-methyl-2-quinazolinesulfonamide

236) N-[3-[1-(4-hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-N-methyl-6-purinesulfonamide

25 237) N-[3-[1-(4-hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-N-methyl-4-thiazolesulfonamide

238) N-[3-[1-(4-hydroxy-2-oxo-6,6-diphenethyl-5,6-dihydro-2H-pyran-3-yl)-propyl]-phenyl]-N-methyl-2-pyridinesulfonamide

30 EXAMPLE 239 2-Pyridylsulfonamide, N-[4-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-
(Formula P-2, R is 2-pyridyl) Refer to Chart P.

35 3-[(3-Aminophenyl)cyclopropylmethyl]-5,6,7,8,9,10-hexahydro-4-hydroxy-2H-cycloocta[b]pyran-2-one of Preparation 66 (100 mg) is dissolved in methylene chloride (3 mL) and pyridine (70 μ L) added. 2-Pyridylsulfonyl chloride (52 mg) is added and the solution stirred for 2 hr at 25°C. Chloroform (25 mL) is added and

the combined extracts washed with 1N-HCl (20 mL) and dried over sodium sulfate. Removal of the solvent gives a pink gum which is chromatographed over silica gel using the flash column technique eluting with 60% ethyl acetate-hexane. The title compound is obtained as a white solid (80 mg).

5 Physical characteristics are as follows:

MS m/z 480, 339, 338, 186, 145, 144, 132, 130, 78, 55.

EXAMPLE 240 4-Pyridylsulfonamide, N-[4-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]-
(Formula P-2, R is 4-pyridyl) Refer to Chart P.

10 Using procedures described in Example 239, the title compound is obtained as a white solid.

Physical characteristics are as follows:

MS m/z 480, 338, 207, 186, 145, 144, 117, 79, 78, 55

15 EXAMPLE 241 5-Cyanopyridin-2-yl-sulfonamide, N-[4-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]- (Formula P-2, R is 5-cyanopyridin-2-yl) Refer to Chart P.

The title compound is prepared using procedures described in Example 239.

20 EXAMPLE 242 2-Pyrazinylsulfonamide, N-[4-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]- (Formula P-2, R is 2-pyrazinyl) Refer to Chart P.

The title compound is prepared using procedures described in Example 239.

25 EXAMPLE 243 2-Pyrimidinylsulfonamide, N-[4-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]- (Formula P-2, R is 2-pyrimidinyl) Refer to Chart P.

The title compound is prepared using procedures described in Example 239.

30 EXAMPLE 244 4-6-Dimethylpyrimidin-2-yl-sulfonamide, N-[4-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)methyl]phenyl]- (Formula P-2, R is 4,6-dimethylpyrimidin-2-yl) Refer to Chart P.

The title compound is prepared using procedures described in Example 239.

35 EXAMPLE 245 4-Methylpyrimidin-2-yl-sulfonamide, N-[4-[cyclopropyl(5,6,7,8,9,10-hexahydro-4-hydroxy-2-oxo-2H-

cycloocta[b]pyran-3-yl)methyl]phenyl]- (Formula P-2, R is 4-methylpyrimidin-2-yl) Refer to Chart P.

The title compound is prepared using procedures described in Example 239.

PREPARATION 69 6,6-Bis-(2-cyclopropyl-ethyl)-dihydro-pyran-2,4-dione (Formula Q-2) Refer to Chart Q.

To a suspension of 150 mg of sodium hydride (60% dispersion in mineral oil) in 4 ml of dry THF under argon atmosphere at 0 °C is added dropwise 0.38 ml of methyl acetoacetate. After 10 minutes 2.3 ml of butyllithium (1.6 M in hexanes) is added. After 10 minutes a solution of 0.48 g of the compound of formula Q-1 (prepared as described in Preparation 79 (Formula S-4, refer to Chart S)) in 3 ml of tetrahydrofuran is added. The reaction mixture is stirred for 1 hour, then partitioned between ethyl acetate and dilute aqueous hydrogen chloride. The aqueous phase is extracted with two additional portions of ethyl acetate. The organic phases are combined, washed with brine, dried over magnesium sulfate and concentrated under reduced pressure. The residue is diluted with 5 mL of methanol and the resulting solution treated with 12 mL of water followed by 3.0 ml of 1 M aqueous sodium hydroxide. After 2 hours of vigorous stirring the methanol is removed under reduced pressure. The aqueous phase is washed once with diethyl ether; the ether phase is discarded. The aqueous phase is cooled to 0 °C, then acidified with dilute aqueous hydrogen chloride. The resulting precipitate is extracted with four portions of dichloromethane. The combined dichloromethane extracts are dried over magnesium sulfate and concentrated under reduced pressure. The residue is dissolved in diethyl ether-hexane and the solution is chilled to provide 0.42 g of the title compound as a pale yellow solid.

Physical characteristics are as follows:

^1H NMR δ 0.0, 0.4, 0.6, 1.2, 1.7, 2.6, 3.4.

PREPARATION 70 6,6-Bis-(2-cyclopropyl-ethyl)-5,6-dihydro-4-hydroxy-3-[1-(3-nitrophenyl)-propyl]-pyran-2-one (Formula Q-3) Refer to Chart Q .

To a stirred solution of 0.41 g of the title compound of Preparation 69 (Formula Q-2) and 0.25 g of the 3-nitrobenzaldehyde in 5 ml of dry tetrahydrofuran is added a solution of 0.44 g of aluminum trichloride in 4.5 ml of tetrahydrofuran. After 2 hours, the reaction mixture is treated with 1.0 g of sodium carbonate decahydrate, stirred 10 minutes, diluted with diethyl ether and finally charged with magnesium sulfate. The resulting mixture is filtered through a pad of Celite with diethyl ether rinses. The filtrates are combined and concentrated under reduced

2187523

pressure. The resulting residue is charged with 103 mg of copper (I) bromide-dimethyl sulfide complex and 5 ml of dry tetrahydrofuran under an argon atmosphere. The reaction mixture is treated dropwise with 2.5 mL of triethyl aluminum (1.0 M in hexane) over 1.5 hours. The reaction is then slowly treated with ice and partitioned between diethyl ether and dilute aqueous hydrogen chloride. The aqueous phase is extracted with three additional portions of diethyl ether. The combined ether extracts are washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. Flash column chromatography of the residue on silica gel using 20% to 40% ethyl acetate in hexane affords 0.44 g of the title compound as a tan foam.

Physical characteristics are as follows:

^1H NMR δ 0.0, 0.4, 0.6, 1.0, 1.2, 1.7-1.9, 2.0-2.4, 2.6, 4.2, 7.5, 7.8, 8.1, 8.3

PREPARATION 71 3-[1-(3-Amino-phenyl)-propyl]-6,6-bis-(2-cyclopropyl-ethyl)-5,6-dihydro-4-hydroxy-pyran-2-one (Formula Q-4) Refer to Chart Q.

To a solution of 0.44 g of the title compound of Preparation 70 (Formula Q-3) in 6 ml of methanol is added 0.65 g of ammonium formate and 50 mg of 10% palladium on carbon. The black slurry is stirred under argon for 3 hours, then filtered through pad of Celite with methanol washes. The filtrates are combined and the solvent is removed under reduced pressure. The residue is triturated with four portions of dichloromethane. The combined dichloromethane washes are concentrated under reduced pressure to provide 0.37 g of the title compound as a white foam.

Physical characteristics are as follows:

R_f 0.08 (50% diethyl ether in hexane)

EXAMPLE 246 N-[3-(1-[6,6-Bis-(2-cyclopropyl-ethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-yl]propyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula Q-5: R_1 is 1-methylimidazol-4-yl) Refer to Chart Q.

To a flask containing 57 mg of the title compound of Preparation 71 (Formula Q-4) and 24 μl of pyridine in 1.0 ml of dichloromethane is added 27 mg of 1-methylimidazole-4-sulfonyl chloride. After 6 hours the reaction mixture is concentrated under reduced pressure. The pyridine is azeotroped thrice with toluene. The resulting residue is flash column chromatographed on silica gel using 2% to 6% methanol in dichloromethane to provide 51 mg of the title compound as a white foam.

Physical characteristics are as follows:

^1H NMR δ 0.0, 0.4, 0.6, 0.9, 1.1-1.4, 1.7-2.2, 2.5, 3.7, 3.95, 6.9, 7.1, 7.4, 7.5

HRMS: 528.2537 (FAB)

EXAMPLE 247 N-[3-(1-[6,6-Bis-(2-cyclopropyl-ethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-yl]propyl)phenyl]-5-cyano-2-pyridinesulfonamide
(Formula Q-5: R_1 is 5-cyano-2-pyridyl) Refer to Chart Q.

Using the general sulfonylation procedure described in Example 246, 57 mg of the amine of Preparation 71 (Formula Q-4) is reacted with 30 mg of 5-cyanopyridine-2-sulfonyl chloride. Flash column chromatography on silica gel using 1% to 3% methanol in dichloromethane provides 62 mg of the title compound as a tan foam.

Physical characteristics are as follows:

^1H NMR δ 0.0, 0.4, 0.6, 0.9, 1.1-1.4, 1.6-2.2, 2.5, 3.95, 6.9-7.2, 8.0, 8.2, 9.0

HRMS: 550.2370 (FAB)

PREPARATION 72 3-Aminopropiophenone (Formula R-2) Refer to Chart R.

To a solution of 3-nitropropiophenone (Formula R-1) (1.79 g) in diethyl ether is added 5% Pt/C catalyst (0.20 g). The resulting suspension is placed under a hydrogen gas atmosphere and stirred for 6 hours. The reaction mixture is filtered through a pad of Celite and the pad washed with additional portions of diethyl ether. The combined filtrates are concentrated under reduced pressure to provide 1.49 g of the title compound as pale yellow, low melting solid.

Physical characteristics are as follows:

^1H NMR δ 1.2, 3.0, 6.9, 7.2-7.4

R_f 0.45 (33% ethyl acetate in hexane)

PREPARATION 73 1-[3-(Dibenzyl-amino)-phenyl]-propan-1-one (Formula R-3) Refer to Chart R.

To a solution of the title compound of Preparation 72 of Formula R-2 (1.5 g) in dichloromethane (50 mL) is added diisopropylethylamine (6.0 mL) followed by benzyl bromide (3.6 mL). After stirring for 6 hours the reaction mixture is heated to reflux overnight. The reaction mixture is cooled to room temperature, diluted with diethyl ether (50 mL) and washed sequentially with dilute aqueous potassium hydrogen sulfate, water, saturated aqueous sodium bicarbonate, and brine. The organic layer is dried over sodium sulfate and concentrated under reduced pressure. The residue is purified by flash column chromatography on silica gel eluting with 5% to 20% ethyl acetate in hexane to provide 2.38 g of the title compound as pale yellow

solid.

Physical characteristics are as follows:

^1H NMR δ 1.1, 2.9, 4.7, 6.9, 7.2-7.4

Anal. Found: C, 83.88; H, 7.03; N, 4.20

5 MS: 329 (EI)

PREPARATION 74 6-[3-(Dibenzyl-amino)-phenyl]-dihydro-pyran-2,4-dione (Formula R-4) Refer to Chart R.

Using the general procedure described in Preparation 69 for the formation of the dihydropyranone ring, the compound of Formula R-3 of Preparation 73 (1.96 g) is
10 reacted with the dianion of methyl acetoacetate and cyclized to provide 0.76 g of the title compound.

Physical characteristics are as follows:

^1H NMR δ 0.8, 1.9, 2.6-2.9, 3.1-3.2, 4.7, 6.5-6.7, 7.1-7.4

MS: 413 (EI)

15 PREPARATION 75 6-[3-(Dibenzyl-amino)-phenyl]-5,6-dihydro-6-ethyl-4-hydroxy-3-[1-(3-nitro-phenyl)-propyl]-pyran-2-one (Formula R-5) Refer to Chart R.

Using the general procedure described in Preparation 70, aluminum trichloride catalyzed condensation of 3-nitrobenzaldehyde with the compound of
20 Formula R-4 of Preparation 74 (727 mg), followed by copper catalyzed conjugate addition with triethyl aluminum provides 800 mg of the title compound.

Physical characteristics are as follows:

^1H NMR δ 0.6, 1.6-2.1, 2.8, 3.4, 3.8, 4.4, 6.4-6.6, 6.8-7.4, 7.7-8.0

MS: 576 (EI)

25 PREPARATION 76 6-(3-Amino-phenyl)-3-[1-(3-amino-phenyl)-propyl]-6-ethyl-5,6-dihydro-4-hydroxy-pyran-2-one (Formula R-6) Refer to Chart R.

Using the general procedure described in Preparation 71, catalytic hydrogenation of the compound of Formula R-5 of Preparation 75 (114 mg) with ammonium formate and Pd/C affords 61 mg of the title compound. Alternatively, the
30 compound of Formula R-5 of Preparation 75 (114 mg) is reduced with Pd/C and hydrogen gas to give 72 mg of the title compound.

Physical characteristics are as follows:

^1H NMR δ 0.6-0.9, 1.8-2.1, 3.0, 3.8, 6.4-6.6, 6.95, 7.1

R_f 0.40 (10% methanol in dichloromethane)

35 EXAMPLE 248 N-(3-[1-(6-Ethyl-5,6-dihydro-4-hydroxy-6-[3-[(1-methyl-1H-

imidazol-4-yl)sulfonyl]amino)phenyl]-2-2H-pyran-3-
yl)propyl]phenyl)-1-methyl-1H-imidazole-4-sulfonamide
(Formula R-7: R₁ is 1-methylimidazol-4-yl) Refer to Chart R.

Using the general sulfonylation procedure described in Example 246, the
5 compound of Formula R-6 of Preparation 76 (61 mg) is reacted with 1-
methylimidazole-4-sulfonyl chloride to provide 59 mg of the title compound.

Physical characteristics are as follows:

¹H NMR δ 0.3-0.7, 1.6-2.0, 3.0, 3.4-3.7, 6.7-7.5

HRMS: 655.1995 (FAB)

10 EXAMPLE 249 5-Cyano-N-(3-[1-(6-[3-[(5-cyano-2-pyridinyl)sulfonyl]amino)
phenyl]-6-ethyl-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-
yl)propyl]phenyl)-2-pyridinesulfonamide (Formula R-7: R is 5-
cyano-2-pyridyl) Refer to Chart R.

Using the general sulfonylation procedure described in Example 246, the
15 compound of Formula R-6 of Preparation 76 (66 mg) is reacted with 5-cyano-2-
pyridine sulfonyl chloride to provide 40 mg of the title compound.

Physical characteristics are as follows:

¹H NMR δ 0.3-0.9, 1.3, 1.6-1.9, 3.0, 3.7, 6.6-7.2, 7.9-8.2, 8.8-9.0

HRMS: 699.1679 (FAB)

20 PREPARATION 77 N-Methoxy-N-methyl-4-pentenoic amide (Formula S-2) Refer to
Chart S.

To a suspension of 4-pentenoic acid (Formula S-1) (2.00 g) and *N,O*-
dimethylhydroxylamine hydrochloride (2.15 g) in dichloromethane (50 mL) at 0 °C is
added diisopropylethylamine (11.5 mL) followed by bis(2-oxo-3-
25 oxazolidinyl)phosphinic chloride (5.60 g). After stirring overnight, the reaction
mixture is concentrated under reduced pressure. The residue is partitioned between
dilute aqueous potassium hydrogen sulfate and diethyl ether. The aqueous phase is
extracted with two additional portions of diethyl ether. The organic extracts are
combined, washed with brine, dried over sodium sulfate and concentrated under
30 reduced pressure. The residue is purified by flash column chromatography on silica
gel eluting with 50% to 80% diethyl ether in hexane to provide 2.58 g of the title
compound as a tan oil.

Physical characteristics are as follows:

¹H NMR δ 2.3-2.6, 3.20, 3.70, 4.9-5.1, 5.75-5.95

35 R_f 0.17 (25% diethyl ether in hexane)

PREPARATION 78 Nona-1,8-dien-5-one (Formula S-3) Refer to Chart S

To a flame-dried flask under an argon atmosphere containing a solution of the title compound of Preparation 77 (Formula S-2) (1.45 g) in dry tetrahydrofuran (10 mL) at 0 °C is added 3-butenyl-1-magnesium bromide (20 mL, 1 M solution in
 5 tetrahydrofuran. (Preparation of this Grignard reagent from magnesium metal and 4-bromo-1-butene is described in J.Org.Chem. 43:4247 (1978)). After 1 hour at 0 °C, the reaction mixture is warmed to room temperature; after 1 hour at room temperature, the reaction mixture is poured into dilute aqueous potassium hydrogen sulfate and partitioned against diethyl ether. The aqueous phase is extracted with
 10 three additional portions of diethyl ether. The organic extracts are combined, washed with brine, dried over sodium sulfate and carefully concentrated under reduced pressure. The resulting liquid is purified by distillation to provide 1.32 g of the title compound as a tan oil.

Physical characteristics are as follows:

15 ^1H NMR δ 2.3, 2.5, 5.0, 5.7-5.9

R_f 0.66 (25% diethyl ether in hexane)

PREPARATION 79 1,5-Dicyclopropyl-pentan-3-one (Formula S-4) Refer to Chart S.

To a flame-dried flask under an argon atmosphere equipped with a reflux condenser containing zinc metal (8.0 g) and cuprous chloride (1.25 g) is added a
 20 solution of the title compound of Preparation 78 (Formula S-3) (1.32 g) in dry diethyl ether (10 mL). The resulting suspension is charged with diiodomethane (5.0 mL) and the reaction flask placed in 40 °C ultrasound bath (Branson^{*} 2200) and sonicated. After 2 hours heating is ceased and sonication is continued overnight. The reaction mixture is then diluted with diethyl ether (50 mL), cooled to 0 °C, and treated with
 25 excess saturated aqueous ammonium chloride. After 0.25 hours of vigorous stirring, the mixture is filtered and the layers separated. The aqueous phase is extracted with two additional portions of diethyl ether. The organic extracts are combined and washed sequentially with dilute aqueous sodium thiosulfate, saturated aqueous sodium bicarbonate, brine; dried over magnesium sulfate and then carefully
 30 concentrated under reduced pressure. The resulting residue is purified by flash column chromatography on silica gel eluting with 5% to 20% diethyl ether in hexane to provide 0.48 g of the title compound as an oil.

Physical characteristics are as follows:

^1H NMR δ 0.0, 0.4, 0.65, 1.45, 2.50

35 R_f 0.44 (10% diethyl ether in hexane)

*Trade-mark

PREPARATION 80 3-[2,2-Dimethyl-1-(3-nitro-phenyl)-propyl]-5,6-dihydro-4-hydroxy-6-phenethyl-6-propyl-pyran-2-one (Formula T-3) Refer to Chart T.

To a flame-dried flask containing a slurry of 977 mg of activated zinc metal in 1.0 mL of dry tetrahydrofuran under an argon atmosphere is added 40 μ L of 1,2-dibromoethane. The mixture is placed in 45 °C ultrasound bath (Branson 2200) and sonicated with stirring. After 10 minutes the mixture is treated with 0.25 mL of chlorotrimethylsilane (1.0 M in tetrahydrofuran). After 10 minutes, the mixture is diluted with 4 mL of tetrahydrofuran and treated dropwise with 1.50 mL of 2-iodo-2-methylpropane. The mixture is stirred and sonicated at 45 °C for an additional 3 hours, then cooled to room temperature without stirring. In a separate flask 954 mg of anhydrous lithium chloride is heated in an 110 °C oil bath *in vacuo* for 1 hour. The LiCl flask is cooled to room temperature, placed under an argon atmosphere and charged with 1.01 g of copper (I) cyanide followed by 10 mL of tetrahydrofuran. After 15 minutes of stirring at room temperature, the LiCl-CuCN mixture is cooled to -30 °C and treated via cannula with the organozinc mixture prepared as described above in the first flask. The reaction flask is warmed from -30 °C to 0 °C, stirred 10 minutes then cooled to -78 °C. The preparation of this organometallic reagent is analogous to literature procedures (Org. Syn. 70:195-203 (1991)) described for related reagents.

In a separate flask a stirred solution of 1.56 g of 6-phenethyl-6-propyl-dihydro-pyran-2,4-dione of Formula T-2 (prepared from the compound of Formula T-1 as described in Preparation 17 above) and 915 mg of the 3-nitrobenzaldehyde in 22 mL of dry tetrahydrofuran is treated with a solution of 1.60 g of aluminum trichloride in 14 mL of tetrahydrofuran. After 2 hours, the reaction mixture is treated with 3.6 g of sodium carbonate decahydrate, stirred 5 minutes, diluted with diethyl ether and finally charged with magnesium sulfate. The resulting mixture is filtered through a pad of Celite with diethyl ether washes. The filtrates are combined and concentrated under reduced pressure. The resulting residue is charged with 9 mL of dry tetrahydrofuran under an argon atmosphere and is added via cannula to the cooled (-78 °C) organometallic reagent solution prepared as described above. After 0.5 hours the reaction mixture is warmed to 0 °C. After 0.5 hours at 0 °C the reaction is poured into cold dilute ammonium chloride and the aqueous phase is made acidic with dilute aqueous hydrogen chloride. The mixture is treated with ethyl acetate and filtered through a pad of Celite with ethyl acetate washes.

The layers are separated and the aqueous phase is extracted with three additional portions of ethyl acetate. The combined ethyl acetate extracts are washed with aqueous sodium thiosulfate, brine; dried over magnesium sulfate, and concentrated under reduced pressure. Flash column chromatography of the residue on silica gel eluting with 30% to 50% ethyl acetate in hexane affords 1.73 g of the title compound as a tan foam.

Physical characteristics are as follows:

^1H NMR δ 0.9, 1.1, 1.3, 1.6-2.0, 2.5-2.8, 4.3, 6.9-7.3, 7.8, 8.0, 8.5

HRMS: 452.2449 (FAB)

10 PREPARATION 81 3-[1-(3-Amino-phenyl)-2,2-dimethyl-propyl]-5,6-dihydro-4-hydroxy-6-phenethyl-6-propyl-pyran-2-one (Formula T-4) Refer to Chart T.

To a solution of 1.72 g of the title compound of Preparation 80 (Formula T-3) in 25 mL of methanol is added 3.0 g of ammonium formate and 400 mg of 10% palladium on carbon. The black slurry is stirred under nitrogen for 3 hours, then filtered through pad of Celite with methanol washes. The filtrates are combined and the solvent is removed under reduced pressure. The residue is repeatedly triturated with portions of dichloromethane and the combined dichloromethane washes concentrated under reduced pressure. The residue is flash column chromatographed on silica gel eluting with 10% ethyl acetate in dichloromethane to provide 1.48 g of the title compound as a white foam.

Physical characteristics are as follows:

^1H NMR δ 0.7-0.9, 1.1, 1.3-2.6, 4.2, 6.55, 6.9-7.3

HRMS: 422.2686 (FAB)

25 EXAMPLE 250 N-[3-(1-[5,6-Dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula T-5: R_1 is 1-methylimidazol-4-yl) Refer to Chart T.

To a solution of 1.48 g of the title compound of Preparation 81 (Formula T-4) in 25 ml of dichloromethane at 0 °C is added 0.57 mL of pyridine followed by 632 mg of 1-methylimidazole-4-sulfonyl chloride. After 3 hours the reaction mixture is warmed to room temperature and concentrated under reduced pressure. Pyridine is azeotroped thrice with toluene. The resulting residue is flash column chromatographed on silica gel using 2% to 6% methanol in dichloromethane to provide 1.7 g of the title compound as a white solid.

Physical characteristics are as follows:

^1H NMR δ 0.8-1.0, 0.97, 1.35, 1.6-2.0, 2.5-2.7, 3.6, 4.1, 6.9-7.5

HRMS: 566.2684

The individual stereoisomers of this compound are the following:

5 N-[3-(1(S)-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula KK-8 wherein R_4 is 1-methyl-1H-imidazol-4-yl) Refer to Chart KK;

N-[3-(1(R)-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula
10 LL-8 wherein R_4 is 1-methyl-1H-imidazol-4-yl) Refer to Chart LL;

N-[3-(1(S)-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula
MM-8 wherein R_4 is 1-methyl-1H-imidazol-4-yl) Refer to Chart MM; and

N-[3-(1(R)-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula
15 NN-8 wherein R_4 is 1-methyl-1H-imidazol-4-yl) Refer to Chart NN.

EXAMPLE 251 5-Cyano-N-[3-(1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide (Formula T-5: R_1 is 5-cyano-2-pyridyl)

20 Refer to Chart T.

Using the general sulfonylation procedure described in Example 250, 42 mg of the amine of Preparation 81 (Formula T-4) is reacted with 20 mg of 5-cyanopyridine-2-sulfonyl chloride. Flash column chromatography on silica gel using 1% to 3% methanol in dichloromethane provides 56 mg of the title compound as a
25 white foam.

Physical characteristics are as follows:

^1H NMR δ 0.8-1.0, 0.92, 1.35, 1.6-2.0, 2.5-2.7, 4.0, 6.9-7.4, 8.0, 8.9

HRMS: 588.2532

PREPARATION 82 N-Methoxy, N-methyl 3-(4-fluorophenyl)propionamide (Formula
30 U-2) Refer to Chart U.

To a cold (0°), stirred solution of 5.0 g of 3-(4-fluorophenyl)propionic acid of Formula U-1, 3.2 g of (N,O)-dimethylhydroxylamine hydrochloride, and 11.4 ml of diisopropylethylamine in 40 ml of dichloromethane is slowly added a solution of 5.0 ml of diethyl cyanophosphonate in 10 ml of dichloromethane. After 18 hours, the
35 solution is concentrated under reduced pressure. The residue is dissolved in ethyl

acetate, and the solution washed with dilute HCl, water, aqueous sodium bicarbonate, and brine, and dried over magnesium sulfate. Removal of the solvent under reduced pressure provides 6.94 g of the title compound.

Physical characteristics are as follows:

5 ^1H NMR δ 2.7, 2.9, 3.17, 3.61, 7.0, 7.2 ppm

IR 1665, 1511, 1222, 1033, 990 cm^{-1}

TLC R_f 0.34 (5% ethyl acetate in dichloromethane)

PREPARATION 83 1-(4-Fluorophenyl)-3-hexanone (Formula U-3) Refer to Chart U.

A stirred solution of 4.68 g of the title compound of Preparation 82 (Formula U-2) in 25 ml of dry THF under argon is cooled to -15° , and to the solution is added 17 ml of a 1M solution of propylmagnesium chloride in ether. The resulting solid mass is warmed to 0° , kept at that temperature for 90 minutes, then partitioned between ether and cold dilute HCl. The aqueous phase is extracted with one additional portion of ether, and the combined organic phase washed with brine and dried over magnesium sulfate. Following removal of solvent by distillation at atmospheric pressure, the residue is purified by evaporative distillation (*ca* 160° @ 13 mmHg) to provide 3.51 g of the title compound as a colorless liquid.

Physical characteristics are as follows:

^1H NMR δ 0.89, 1.6, 2.36, 2.7, 2.9, 6.9, 7.1 ppm

20 IR 2965, 1714, 1511, 1222 cm^{-1}

PREPARATION 84 5,6-Dihydro-4-hydroxy-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-2-one (Formula U-4) Refer to Chart U.

To a cold (0°), stirred slurry of 950 mg of sodium hydride (60% dispersion in mineral oil) in 30 ml of dry THF, under argon, is added dropwise 2.3 ml of methyl acetoacetate. After 5 minutes, 13.5 ml of butyllithium (1.6M in hexanes) is added, and the mixture stirred another 5 minutes before addition of a solution of 3.51 g of the title compound of Preparation 83 (Formula U-3) in 4 ml of THF. The solution is stirred for 1 hour, then partitioned between ethyl acetate and cold dilute HCl. The aqueous phase is extracted with two additional portions of ethyl acetate, and the combined organic phase washed with brine and dried over magnesium sulfate. Removal of the solvent under reduced pressure provides the intermediate ester with the following physical characteristics: TLC R_f 0.45 (50% ethyl acetate in hexane).

The ester is stirred in 20 ml of 1M sodium hydroxide, 80 ml of water, and 40 ml of methanol for 90 minutes, then the methanol is removed under reduced pressure. The aqueous phase is washed once with ether, the ether phase being

discarded, and then acidified with dilute HCl. The resulting precipitate is extracted with four portions of dichloromethane, and the extract dried over magnesium sulfate and concentrated under reduced pressure. The residue is dissolved in 1:1 ether-hexane and the solution chilled to provide crystals, which are filtered, washed with
5 ether-hexane, and dried under vacuum to afford 3.24 g of the title compound.

Physical characteristics are as follows:

^1H NMR δ 0.96, 1.4, 1.8, 2.0, 2.5, 2.7, 7.0, 7.1 ppm

IR 2962, 1655, 1604, 1510, 1221 cm^{-1}

M.P. 113-114.5°

10 Anal. Found: C, 68.85; H, 6.99

MS: M^+ 278

R_f 0.44 (5% methanol in dichloromethane)

PREPARATION 85 3-(1-(3-Benzoyloxycarbonylaminophenyl)-2,2-dimethylpropyl)-5,6-dihydro-4-hydroxy-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-2-one (Formula U-6: R_1 is *tert*-butyl) Refer to Chart U.
15

To a stirred solution of 3.06 g of the title compound of Preparation 84 (Formula U-4) and 2.81 of the title compound of Preparation 6 above (Formula B-2) in 30 ml of dry THF is added a solution of 2.93 g of AlCl_3 in 20 ml of THF. After two hours, 6.4 g of sodium carbonate decahydrate is added, and after five minutes
20 the mixture is filtered through Celite with ether rinses. Removal of the solvent under reduced pressure provides the intermediate benzylidene compound of Formula U-5.

To this is added, under argon, 1.13 g of copper (I) bromide-dimethyl sulfide complex and 30 ml of THF, and the mixture is cooled to 0° for dropwise addition of
25 18.1 ml of *tert*-butylmagnesium chloride (1.0M in THF). After 10 minutes, the reaction is partitioned between ether and cold dilute HCl. The organic phase is washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. Flash chromatography of the residue on silica gel using 30-35% ethyl acetate in hexane affords 1.83 g of the title compound as a foam.

30 Physical characteristics are as follows:

^1H NMR δ 0.87, 1.1, 1.3, 1.6-2.2, 2.5, 5.12, 6.8-7.6 ppm

HRMS: 574.2955

R_f 0.29 (35% ethyl acetate in hexane)

PREPARATION 86 3-(1-(3-Aminophenyl)-2,2-dimethylpropyl)-5,6-dihydro-4-hydroxy-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-2-one
35

(Formula U-7: R_1 is *tert*-butyl) Refer to Chart U.

A mixture of 1.83 g of the title compound of Preparation 85 (Formula U-6), 2.0 g of ammonium formate, and 400 mg of 10% palladium on carbon in 25 ml of methanol is stirred under argon for 90 minutes, then filtered through Celite. The solvent is removed under reduced pressure and the residue flash chromatographed on silica gel using 10% ethyl acetate in dichloromethane to provide 1.24 g of the title compound as a white foam.

Physical characteristics are as follows:

R_f 0.28 (10% ethyl acetate in dichloromethane)

The compound of Formula U-7 wherein R_1 is ethyl is prepared from U-4 by analogous procedures as in the preparation of U-7 wherein R_1 is *tert*-butyl (Preparations 85 and 86).

EXAMPLE 252 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula U-8: R_1 is *tert*-butyl, R_2 is 1-methylimidazole-4-yl) Refer to Chart U.

To a cold (0°), stirred solution of 88 mg of the title compound of Preparation 86 (Formula U-7) and 32 μ l of pyridine in 0.5 ml of dichloromethane is added 36 mg of 1-methylimidazole-4-sulfonyl chloride. After 90 minutes the reaction mixture is flash chromatographed on silica using 3-4% methanol in dichloromethane to provide 112 mg of the title compound as a white foam.

Physical characteristics are as follows:

^1H NMR δ 0.8-1.0, 0.96, 1.3, 1.7, 2.35, 2.5, 3.6, 3.7, 6.8-7.5 ppm

HRMS: 583.2525

R_f 0.31 (5% methanol in dichloromethane)

EXAMPLE 253 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula U-8: R_1 is *tert*-butyl, R_2 is 5-cyanopyridine-2-yl) Refer to Chart U.

Using the general sulfonylation procedure of Example 252, 88 mg of the amine of Preparation 86 (Formula U-7, R_1 is *tert*-butyl) is reacted with 5-cyanopyridine-2-sulfonyl chloride. Flash chromatography on silica using 10-15% ethyl acetate in dichloromethane provides 107 mg of the title compound as an amorphous white solid.

Physical characteristics are as follows:

2187523

¹H NMR δ 0.92, 1.3, 1.7, 2.5, 6.8-7.5, 8.0, 8.9 ppm

HRMS: 606.2423

EXAMPLE 254 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula U-8: R₁ is ethyl, R₂ is 1-methylimidazole-4-yl) Refer to Chart U.

Using the general sulfonylation procedure of Example 252, 82 mg of the amine of Formula U-7, wherein R₁ is ethyl, is reacted with 1-methylimidazole-4-sulfonyl chloride. Flash chromatography on silica using 3% methanol in dichloromethane provides 101 mg of the title compound as an amorphous white solid.

Physical characteristics are as follows:

¹H NMR δ 0.8, 1.3, 1.6-2.2, 2.5, 3.5, 3.6, 3.9, 6.8-7.4 ppm

HRMS: 555.2192

R_f 0.29 (5% methanol in dichloromethane)

EXAMPLE 255 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula U-8: R₁ is ethyl, R₂ is 5-cyanopyridine-2-yl) Refer to Chart U.

Using the general sulfonylation procedure of Example 252, 82 mg of the amine of Formula U-7, wherein R₁ is ethyl, is reacted with 5-cyanopyridine-2-sulfonyl chloride. Flash chromatography on silica using 10-15% ethyl acetate in dichloromethane provides 101 mg of the title compound as an amorphous white solid.

Physical characteristics are as follows:

¹H NMR δ 0.9, 1.3, 1.6-2.2, 2.5, 3.9, 6.9-7.3, 8.0, 8.1, 8.9 ppm

HRMS: 557.2059

R_f 0.44 (20% ethyl acetate in dichloromethane)

PREPARATION 87 1,5-Bis-(4-fluorophenyl)-penta-1,4-dien-3-one (Formula V-2)

Refer to Chart V.

To a rapidly stirred, ambient temperature solution of 10 g of sodium hydroxide in 100 ml of water and 80 ml of ethanol is added a mixture of 12.4 g of 4-fluorobenzaldehyde of Formula V-1 and 2.9 g of acetone. After 45 minutes, the resulting precipitate is filtered off, washed well with water, and dried under vacuum. Recrystallization from ethyl acetate-hexane yields 10.7 g of the title

compound as light yellow platelets.

Physical characteristics are as follows:

^1H NMR δ 6.9-7.2, 7.6-7.7 ppm

IR 1653, 1587, 1508, 984, 835 cm^{-1}

5 MS: M^+ 270

Anal. Found: C, 75.40; H, 4.41

R_f 0.35 (dichloromethane)

M.P. 152-154°

10 PREPARATION 88 1,5-Bis-(4-fluorophenyl)-pentane-3-one (Formula V-3) Refer to Chart V.

To a solution of 5.41 g of dienone of Preparation 87 (Formula V-2) in 10 ml of THF and 50 ml of methanol is added 2.0 g of magnesium chips. A water bath is used to maintain the temperature of the reaction near ambient. After the magnesium has been consumed, the reaction mixture is partitioned between
15 dichloromethane and dilute HCl, with two additional dichloromethane extractions of the aqueous phase. The combined organic phase is dried over magnesium sulfate and concentrated under reduced pressure. Flash chromatography of the residue on silica using 50% dichloromethane in hexane affords 3.66 g of the title compound as a yellow oil.

20 Physical characteristics are as follows:

^1H NMR δ 2.67, 2.85, 6.9, 7.1 ppm

IR 2932, 1716, 1603, 1511, 1223, 1159, 828 cm^{-1}

MS: M^+ 274

R_f 0.28 (50% dichloromethane in hexane)

25 PREPARATION 89 4-Hydroxy-5,6-dihydro-6,6-bis(2-(4-fluorophenyl)ethyl)-2H-pyran-2-one (Formula V-4) Refer to Chart V.

Using the general acetoacetate condensation and ring closure procedure of Preparation 84 (Formula U-4), 3.9 g of the ketone of Preparation 88 (Formula V-3) is converted to 2.86 g of the title compound, which may be recrystallized from
30 dichloromethane-hexane.

Physical characteristics are as follows:

^1H NMR δ 2.1, 2.57, 2.7, 7.0, 7.1 ppm

IR 2924, 1659, 1578, 1508, 1241, 1216 cm^{-1}

MS: M^+ 358

35 Anal. Found: C, 70.17; H, 5.50

M.P. 140-141°

PREPARATION 90 3-[1-(3-Benzoyloxycarbonylaminophenyl)-2,2-dimethylpropyl]-6,6-bis[2-(4-fluorophenyl)ethyl]-4-hydroxy-5,6-dihydro-2H-pyran-2-one (Formula V-6: R₁ is *tert*-butyl) Refer to Chart V.

5 Using the general benzylidene condensation and cuprate addition procedure of Preparation 85 (Formula U-6), 1.075 g of the dihydropyrone of Preparation 89 (Formula V-4) is converted to 707 mg of the title compound (via the intermediate compound of Formula V-5), which is purified by flash chromatography on silica gel using 40% ethyl acetate in hexane.

10 Physical characteristics are as follows:

¹H NMR δ 1.07, 2.0, 2.6, 3.9, 5.16, 6.8-7.5 ppm

HRMS: 654.3023

R_f 0.25 (40% ethyl acetate in hexane)

15 PREPARATION 91 3-[1-(3-Aminophenyl)-2,2-dimethylpropyl]-6,6-bis[2-(4-fluorophenyl)ethyl]-4-hydroxy-5,6-dihydro-2H-pyran-2-one (Formula V-7: R₁ is *tert*-butyl) Refer to Chart V.

Using the general transfer hydrogenolysis procedure of Preparation 86 (Formula U-7), 684 mg of the carbamate of Preparation 90 (Formula V-6, R₁ is *tert*-butyl) is converted to 497 mg of the title compound, which is purified by flash
20 chromatography on silica gel using 5-10% ethyl acetate in dichloromethane.

Physical characteristics are as follows:

¹H NMR δ 1.09, 2.0, 2.6, 6.8-7.1 ppm

R_f 0.34 (10% ethyl acetate in dichloromethane)

The compound of Formula V-7 wherein R₁ is ethyl is prepared from V-4 by
25 analogous procedures as in the preparation of V-7 wherein R₁ is *tert*-butyl (Preparations 90 and 91).

EXAMPLE 256 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-(4-fluorophenyl)ethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula V-8: R₁ is *tert*-
30 butyl, R₂ is 1-methylimidazole-4-yl) Refer to Chart V.

Using the general sulfonylation procedure of Example 252, 78 mg of the amine of Preparation 91 (Formula V-7, R₁ is *tert*-butyl) is reacted with 1-methylimidazole-4-sulfonyl chloride. Flash chromatography on silica using 3-4% methanol in dichloromethane provides 92 mg of the title compound as an amorphous
35 white solid.

Physical characteristics are as follows:

^1H NMR δ 0.94, 1.7-2.1, 2.5, 3.50, 6.8-7.4 ppm

HRMS: 664.2647

R_f 0.34 (5% methanol in dichloromethane)

- 5 **EXAMPLE 257** N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-(4-fluorophenyl)ethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula V-8: R_1 is *tert*-butyl, R_2 is 5-cyanopyridine-2-yl) Refer to Chart V.

10 Using the general sulfonylation procedure of Example 252, 78 mg of the amine of Preparation 91 (Formula V-7, R_1 is *tert*-butyl) is reacted with 5-cyanopyridine-2-sulfonyl chloride. Flash chromatography on silica using 10-15% ethyl acetate in dichloromethane provides 91.5 mg of the title compound as an amorphous white solid.

Physical characteristics are as follows:

15 ^1H NMR δ 0.92, 1.9, 2.6, 3.2, 6.8-7.5, 8.0, 8.9 ppm

HRMS: 686.2488

R_f 0.28 (10% ethyl acetate in dichloromethane)

- 20 **EXAMPLE 258** N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-(4-fluorophenyl)ethyl)-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula V-8: R_1 is ethyl, R_2 is 1-methylimidazole-4-yl) Refer to Chart V.

25 Using the general sulfonylation procedure of Example 252, 74 mg of the amine of Formula V-7, wherein R_1 is ethyl, is reacted with 1-methylimidazole-4-sulfonyl chloride. Flash chromatography on silica using 3-4% methanol in dichloromethane provides 77 mg of the title compound as an amorphous white solid.

Physical characteristics are as follows:

^1H NMR δ 0.87, 2.0, 2.6, 3.62, 4.0, 4.05, 6.9-7.5 ppm

HRMS: 636.2350

R_f 0.31 (5% methanol in dichloromethane)

- 30 **EXAMPLE 259** N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-(4-fluorophenyl)ethyl)-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula V-8: R_1 is ethyl, R_2 is 5-cyanopyridine-2-yl) Refer to Chart V.

35 Using the general sulfonylation procedure of Example 252, 74 mg of the amine of Formula V-7, wherein R_1 is ethyl, is reacted with 5-cyanopyridine-2-

sulfonyl chloride. Flash chromatography on silica using 10% ethyl acetate in dichloromethane provides 83 mg of the title compound as an amorphous white solid.

Physical characteristics are as follows:

^1H NMR δ 0.83, 2.0, 2.6, 3.96, 6.8-7.2, 8.0, 8.8 ppm

5 HRMS: 658.2200

R_f 0.49 (10% ethyl acetate in dichloromethane)

EXAMPLE 260 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide
(Formula D-6: R_1 is propyl, R_2 is propyl, R_3 is *tert*-butyl, R_4 is
10 5-cyanopyridine-2-yl) Refer to Chart D.

Using the general sulfonylation procedure of Example 252, 54 mg of the amine of Formula D-5 (R_1 and R_2 are propyl, R_3 is *tert*-butyl), prepared by procedures analogous to those described for the preparation of D-5 (where R_1 is phenethyl, R_2 is propyl, R_3 is ethyl) in Preparation 20, is coupled with 5-
15 cyanopyridine-2-sulfonyl chloride of Formula D-7 (R_4 is 5-cyanopyridine-2-yl) to yield, after flash chromatography on silica gel using 10-15% ethyl acetate in dichloromethane, 62 mg of the title compound as an amorphous solid.

Physical characteristics are as follows:

^1H NMR δ 0.90, 1.2-1.8, 2.5, 7.0-7.4, 8.1, 8.2, 8.9 ppm

20 HRMS: 525.2305

R_f 0.44 (20% ethyl acetate in dichloromethane)

EXAMPLE 261 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-
sulfonamide (Formula D-6: R_1 is propyl, R_2 is propyl, R_3 is *tert*-
25 butyl, R_4 is 1-methylimidazole-4-yl) Refer to Chart D.

Using the general sulfonylation procedure of Example 252, 54 mg of the amine of Formula D-5 (R_1 and R_2 are propyl, R_3 is *tert*-butyl), is coupled with 1-methylimidazole-4-sulfonyl chloride of Formula D-7 (R_4 is 1-methylimidazole-4-yl) to
30 yield, after flash chromatography on silica gel using 3-5% methanol in dichloromethane, 53 mg of the title compound as an amorphous solid.

Physical characteristics are as follows:

^1H NMR δ 0.9, 0.96, 1.2-1.8, 2.5, 3.6, 3.7, 6.9-7.5 ppm

MS: 503.2422

R_f 0.26 (5% methanol in dichloromethane)

35 EXAMPLE 262 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-

yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula D-6:
R₁ is propyl, R₂ is propyl, R₃ is ethyl, R₄ is 5-cyanopyridine-2-
yl) Refer to Chart D.

Using the general sulfonylation procedure of Example 252, 67 mg of the amine of
5 Formula D-5 (R₁ and R₂ are propyl, R₃ is ethyl) of Preparation 20 is coupled with 5-
cyanopyridine-2-sulfonyl chloride of Formula D-7 (R₄ is 5-cyanopyridine-2-yl) to
yield, after flash chromatography on silica gel using 10% ethyl acetate in
dichloromethane, 78 mg of the title compound as an amorphous solid.

Physical characteristics are as follows:

10 ¹H NMR δ 0.6-1.0, 1.2-1.8, 3.4, 3.5, 6.9-7.4, 8.0-8.2, 8.9 ppm

HRMS: 498.2072

R_f 0.38 (15% ethyl acetate in dichloromethane)

EXAMPLE 263 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6-phenethyl-6-propyl-2H-
pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide

15 (Formula D-6: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is
5-cyanopyridine-2-yl) Refer to Chart D.

Using the general sulfonylation procedure of Example 252, 79 mg of the
amine of Formula D-5 (R₁ is phenethyl, R₂ is propyl, R₃ is ethyl) of Preparation 20
is coupled with 5-cyanopyridine-2-sulfonyl chloride of Formula D-7 (R₄ is 5-
20 cyanopyridine-2-yl) to yield, after flash chromatography on silica gel using 10% ethyl
acetate in dichloromethane, 102 mg of the title compound as an amorphous solid.

Physical characteristics are as follows:

¹H NMR δ 0.7-1.0, 1.2-2.6, 3.4, 3.5, 6.9-7.3, 7.9-8.2, 8.9 ppm

HRMS: 560.2231

25 R_f 0.37 (15% ethyl acetate in dichloromethane)

EXAMPLES 264-265

The following compounds are prepared using the general sulfonylation
procedure of Example 246. The requisite amine is prepared analogously from the
compound of Formula Q-1 (Preparation 69) following Preparations 80 and 81.

30 264) N-[3-(1-[6,6-Bis-(2-cyclopropyl-ethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-
pyran-3-yl]-2,2-dimethyl-propyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide

Physical characteristics are as follows:

¹H NMR δ 0.0, 0.4, 0.6, 1.0, 1.2, 1.7, 2.5, 3.7, 4.1, 6.9-7.6

HRMS: 556.2833 (FAB)

35 265) N-[3-(1-[6,6-Bis-(2-cyclopropyl-ethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-

2187523

pyran-3-yl]-2,2-dimethyl-propyl)phenyl]-5-cyano-2-pyridinesulfonamide

Physical characteristics are as follows:

^1H NMR δ 0.0, 0.4, 0.6, 1.0, 1.2, 1.7, 2.5, 4.1, 7.0-7.5, 8.0, 8.1, 9.0

HRMS: 578.2689 (FAB)

5 PREPARATION 92 (3(2E),4S)-3-(2-pentenyl)-4-phenyl-2-oxazolidinone (Formula W-4) Refer to Chart W.

A 1 L round-bottomed flask with nitrogen inlet and addition funnel is charged with 6.92 g of (S)-(+)-4-phenyl-2-oxazolidinone and 250 mL of tetrahydrofuran and then cooled to -78°C . To the aforementioned solution is added
10 25.6 mL of n-butyl lithium during which time a white solid separated from the reaction solution, W-3. To that suspension is added 4.88 g of *trans*-2-pentenoyl chloride of formula W-2 (prepared from the treatment of commercially available *trans*-2-pentenoyl acid of formula W-1 with oxalyl chloride) in a small volume of THF. The resulting pale yellow homogeneous solution is allowed to warm to room
15 temperature and stirred for another 20 min. The reaction mixture is quenched by the addition of saturated ammonium chloride solution and is extracted with ethyl acetate. The organic layer is separated, washed with brine and water, dried over magnesium sulfate, filtered and concentrated to give a white solid. Recrystallization from hot hexane gives 9.13 g of the title compound.

20 Physical characteristics are as follows:

MP $86-88^\circ\text{C}$.

^1H NMR (CDCl_3) δ 7.42-7.23, 7.18-7.09, 5.49, 4.70, 4.28, 2.28, 1.08 ppm.

$[\alpha]_D(\text{CHCl}_3) = +109$

Anal. found: C, 68.59; H, 6.25; N, 5.70

25 PREPARATION 93 (3(3R),4S)-3-[3-(3-Aminophenyl)-1-oxopentyl]-4-phenyl-2-oxazolidinone (Formula W-5) Refer to Chart W.

A 1 L three-necked, round-bottomed flask with nitrogen inlet and addition funnel is charged with 8.90 g of copper(I) bromide-dimethyl sulfide complex and 125 mL of THF and then cooled to -40°C . To that suspension is added 43 mL of a 1 M
30 solution (in THF) of 3-[bis(trimethylsilyl)amino] phenylmagnesium chloride dropwise over 15 minutes. The reaction mixture is warmed 0°C for 30 minutes and then a 25 mL THF solution containing 8.85 g of (3(2E),4S)-3-(2-pentenoyl)-4-phenyl-2-oxazolidinone of Preparation 92 (formula W-4) is added. The reaction mixture is stirred for 30 minutes at 0°C and quenched by the addition of 1N HCl and then the
35 pH readjusted with 1N NaOH to pH 8. The reaction is washed with water, brine

and the organic is dried (Na_2SO_4). The organic solvent is evaporated in vacuo and the resulting oil chromatographed over 600 g of silica gel, eluting with ethyl acetate/hexane to afford 7.91 g of the title product.

Physical characteristics are as follows:

5 MP 94-95 °C.

^1H NMR (CDCl_3) δ 7.28-7.25, 7.07-6.99, 6.60-6.51, 5.38, 4.63, 4.16, 3.52-3.44, 3.10-2.92, 1.65-1.53, 0.76 ppm.

IR (mineral oil) 3437, 3355, 1773, 1696, 1605, 1337, 1322, 1299, 1263, 1212, 1096, 1070, 791, 762, 704 cm^{-1} .

10 Anal. found: C, 71.00; H, 6.67; N, 8.17

EI-MS: $[\text{M}^+] = 338$.

$[\alpha]_D$ (19.87 mg/2 mL CHCl_3) = +60°

PREPARATION 94 3-[3-(3-[Bis(phenylmethyl)amino]phenyl)-1-oxopentyl]-4-phenyl-2-oxazolidinone, (3R)(4S) (Formula W-6) Refer to Chart W.

15 To a mixture of 25 mL of Na_2CO_3 and 80 mL of methylene chloride was added 7.90 g of (3(3R),4S)-3-[3-(3-aminophenyl)-1-oxopentyl]-4-phenyl-2-oxazolidinone of Preparation 93 (formula W-5) followed by 15.94 g of benzyl bromide. That mixture is heated at 65°C for 18 hours, the methylene chloride layer separated, dried (Na_2SO_4) and solvent evaporated to yield the crude product as a dark viscous
20 oil. That oil is chromatographed over 700 g of silica gel eluting with 25% ethyl acetate/hexane to yield 8.55 g of the title compound.

Physical characteristics are as follows:

MP 92-3°C

25 ^1H NMR (CDCl_3) δ 7.24, 7.02, 6.53, 5.34, 4.59, 4.14, 3.44, 3.07, 2.89, 1.50, 0.64 ppm

Anal. found: C, 78.47; H, 6.68; N, 5.26

$[\alpha]_D$ (19.602 mg/2 mL CHCl_3) = +32°

PREPARATION 95 (3R)(4S) 3-[3-(3-[bis(phenylmethyl)amino]phenyl)-2-(2-methyl-1,3-dioxolan-2-yl)-1-oxopentyl]-4-phenyl-2-oxazolidinone
30 (Formula W-8) Refer to Chart W.

To 25 mL of methylene chloride is added 2.1 g of the amide of formula W-6 of Preparation 94 and the resulting solution cooled to -78°C under an atmosphere of nitrogen. To that solution is added 872 μL of neat TiCl_4 followed by the addition of 732 μL of diisopropylethylamine. The resulting mixture is warmed to 0°C for 30
35 minutes and then cooled back to -78°C and 1.3 g of 2-methoxy-2-methyl-1,3-

dioxolane of formula W-7 and the resulting reaction is warmed to 0°C and stirred for 1 hour, then quenched with saturated ammonium chloride and extracted with methylene chloride. The organic extract is dried (Na₂SO₄) and solvent removed in vacuo to afford the crude material. Silica gel chromatography using 100g of support and eluting with 10% hexane/methylene chloride afforded 1.76 g of the title product.

Physical characteristics are as follows:

¹H NMR (CDCl₃) δ 7.36, 7.08, 5.99, 5.42, 4.80, 4.68, 4.60, 4.25, 3.68, 3.57, 3.48, 3.07, 2.90, 1.5, 0.86, 0.54 ppm

Anal. found: C, 75.34; H, 6.99; N, 4.87

[α]_D (18.086 mg/2 mL CHCl₃) = +25°

PREPARATION 96 (3R)(4S)-3-[2-acetyl-3-[3-(bis(phenylmethyl)amino)phenyl]-1-oxopentyl]-4-phenyl-2-oxazolidinone (Formula W-9) Refer to Chart W.

To 25 mL of tetrahydrofuran and 10 mL of 30% HClO₄ is added 5.0 g of the title compound of Preparation 95 (formula W-8) and the resulting solution stirred at 40°C for 3 hours. The reaction is neutralized with saturated NaHCO₃ to pH 8 and then extracted with 400 mL of ether. The ether layer is washed with water, brine and then dried (Na₂SO₄) and solvent evaporated in vacuo to afford an oil. Chromatography over 300 g of silica gel eluting with 15% acetone/hexane afforded 4.12 g of the title compound.

Physical characteristics are as follows:

¹H NMR (CDCl₃) δ 7.31, 7.08, 6.59, 6.55, 5.42, 4.67, 4.61, 4.22, 3.09, 1.63, 1.56, 0.61 ppm

Anal. found: C, 77.11; H, 6.76; N, 4.98

[α]_D (20.172 mg/2 mL CHCl₃) = -10°

PREPARATION 97 (3R)(4S) 3-[2-[1-(3-[bis(phenylmethyl)amino]phenyl)propyl]-5-hydroxy-1,3-dioxo-5-propyloctyl]-4-phenyl-2-oxazolidinone (Formula W-10) Refer to Chart W.

To 25 mL of methylene chloride is added 1.32 g of the compound of Preparation 96 (formula W-9) and the resulting solution cooled to -78°C under an atmosphere of nitrogen. To that solution is added 279 μL of TiCl₄ and 450 μL of diisopropylethylamine and stirring continued for 1 hour. To this solution is added 689 μL of heptanone and the reaction temperature raised to 0°C for 1.5 hours. The reaction is then quenched by the addition of a saturated ammonium chloride solution and the mixture extracted with methylene chloride. The organic extract is

washed with saturated NaHCO_3 , dried (Na_2SO_4) and evaporated in vacuo to yield the crude product. Chromatography over 100 g of silica gel eluting with 5% hexane/methylene chloride affords 1.16 g of the title compound as an off white foam.

Physical characteristics are as follows:

5 ^1H NMR (CDCl_3) 7.36, 7.07, 6.58, 6.54, 5.44, 5.24, 4.69, 4.61, 4.27, 3.21, 3.01, 2.48, 1.90, 1.54, 1.15, 0.81, 0.76, 0.58 ppm

Anal. found: C, 76.62; H, 7.63; N, 4.17

$[\alpha]_D$ (15.380 mg/2 mL CHCl_3) = +16°

PREPARATION 98 (3S)-3-[1-(3-(Bis(phenylmethyl)amino)phenyl)propyl]-6,6-
10 dipropyl-5,6-dihydro-4-hydroxy-2H-pyran-2-one (Formula W-11)
Refer to Chart W.

To 10 mL of dry tetrahydrofuran is added 770 mg of the title compound of Preparation 97 (formula W-10) and the resulting solution cooled to 0°C under an atmosphere of nitrogen. To that solution is added 150 mg of a 60% oil dispersion of
15 sodium hydride and the reaction is warmed to 20°C and stirring continued for 16 hours. The reaction is quenched with saturated ammonium chloride and extracted with ethyl acetate. The extract is dried and evaporated in vacuo to yield the crude product. Chromatography over 100 g of silica gel eluting with 15% EtOAc/hexane affords 560 mg of the title product.

20 Physical characteristics are as follows:

^1H NMR (CDCl_3) 7.34, 6.69, 5.87, 4.69, 4.60, 4.09, 2.28, 2.17, 1.89, 1.73, 1.55, 1.32, 0.88 ppm

[Anal. found: C, 79.71; H, 8.07; N, 2.61]

$[\alpha]_D$ (15.998 mg/2 mL CHCl_3) = -56°

25 PREPARATION 99 (3S)-3-[1-(3-aminophenyl)propyl]-6,6-dipropyl-5,6-dihydro-4-hydroxy-2H-pyran-2-one (Formula W-12) Refer to Chart W.

The title compound of Preparation 98 (formula W-11) ((3R)-3-[1-(3-bis-benzylaminophenyl)propyl]-6,6-bispropyl-5,6-dihydro-4-hydroxypyran-2-one) 110 mg is added to 20 mL of ethyl acetate. To that solution is added 50 mg of 10% Pd/C and
30 the resulting mixture is hydrogenated at 50 psi for 6 hours. The reaction is filtered through celite to yield 83 mg of the title product.

Physical characteristics are as follows:

IR 2957, 2922, 2855, 2871, 2854, 1378, 1605, 1459, 1617, 1262, 1319, 1251, 1282, 1107 cm^{-1} .

35 $[\alpha]_D$ (6.526 mg/2 mL CH_3OH) = -34°

PREPARATION 100 (4R)3-(1-oxo-2-pentenyl)-4-phenyl-2-oxazolidinone
(Formula X-4) Refer to Chart X.

A 2-L, three-necked, round-bottomed flask with nitrogen inlet and addition funnel is charged with (*R*)-(-)-4-phenyl-2-oxazolidinone (31.2 g) and tetrahydrofuran (1.2 L) and cooled to -78 °C. The addition funnel is charged with *n*-butyllithium (1.6 M in hexanes, 117 mL), which is added dropwise to the reaction mixture over 20 min. A white precipitate is formed which is X-3. The reaction mixture is stirred for an additional 30 min at -78 °C. The addition funnel is then charged with *trans*-2-pentenoyl chloride of formula X-2, prepared from the acid of formula X-1, (24.4 g) and tetrahydrofuran (50 mL), and this solution is added to the reaction mixture dropwise over 10 min. The resulting pale yellow homogeneous solution is allowed to warm to room temperature and is stirred for another 30 min. The reaction mixture is quenched by the addition of saturated ammonium chloride solution and is extracted with ethyl acetate (2500 mL). The organic layer is separated, washed with brine and water, dried over magnesium sulfate, filtered and concentrated to give 48 g of a white solid. The solid is recrystallized from ethyl acetate (100 mL) and hexane (200 mL) to give 38.0 g the title product as a white solid.

Physical characteristics are as follows:

MP 86-88 °C.

¹H NMR (CDCl₃) δ 7.42-7.23, 7.18-7.09, 5.49, 4.70, 4.28, 2.28, 1.08 ppm.

IR (mineral oil) 1785, 1764, 1686, 1638, 1349, 1336, 1329, 1257, 1234, 1214, 1087, 1076, 756, 716, 699 cm⁻¹

EI-MS: [M⁺] = 245.

PREPARATION 101 (3(3S),4R)-3-[3-(3-aminophenyl)-1-oxopentyl]-4-phenyl-2-oxazolidinone (Formula X-5) Refer to Chart X.

A 2-L, three-necked, round-bottomed flask with nitrogen inlet and addition funnel is charged with copper(I) bromide-dimethyl sulfide complex (25.1 g) and tetrahydrofuran (250 mL) and cooled to -40 °C. The addition funnel is charged with 3-[bis(trimethylsilyl)amino] phenylmagnesium chloride (1.0 M in THF, 122 mL), which is added dropwise to the reaction mixture over 20 min. The reaction mixture is then allowed to warm from -40 °C to -20 °C over 20 min. The addition funnel is charged with 25 g of the title compound of Preparation 100 (formula X-4) and tetrahydrofuran (100 mL), and this solution is added to the reaction mixture dropwise over 30 min at 0 °C. The reaction mixture is then stirred for 15 min at 0 °C and quenched by the addition of saturated ammonium chloride solution (adjusted to

- pH 8 by addition of ammonium hydroxide). The reaction mixture is poured into ether (2 L) and washed with the ammonium chloride solution until the aqueous layer is no longer blue in color. The organic layer is separated, washed with water, dried over magnesium sulfate, filtered and concentrated to give 58 g of a yellow oil.
- 5 The crude reaction mixture is then stirred at room temperature in a slurry of silica gel (75 g) and methylene chloride (100 mL) for 1 h. The mixture is filtered, washed with methanol, and concentrated to give 49 g of an oil. Column chromatography on 300 g silica (eluting with 10-75% ethyl acetate-hexane, 100% ethyl acetate) yields 30.9 g of a yellow oil. The oil is crystallized from ethyl acetate (75 mL) and hexane
- 10 (150 mL) to give 21.4 g of the title compound as a white solid.

Physical characteristics are as follows:

MP 94-97 °C.

¹H NMR (CDCl₃) δ 7.28-7.25, 7.07-6.99, 6.60-6.51, 5.38, 4.63, 4.16, 3.52-3.44, 3.10-2.92, 1.65-1.53, 0.76 ppm.

- 15 IR (mineral oil) 3437, 3355, 1773, 1696, 1605, 1337, 1322, 1299, 1263, 1212, 1096, 1070, 791, 762, 704 cm⁻¹.

EI-MS: [M⁺] = 338.

PREPARATION 102 (3(3S),4R)-3-[3-(3-(phenylmethyl)amino)phenyl]-1-oxopentyl]-4-phenyl-2-oxazolidinone (Formula X-6) Refer to Chart X.

20

- To a mixture of 80 mL of Na₂CO₃ and 280 mL of methylene chloride is added 21.0 g of (3(3S),4R)-3-[3-(3-aminophenyl)-1-oxopentyl]-4-phenyl-2-oxazolidinone (formula X-5) of Preparation 101 followed by 23.4 g of benzyl bromide. That mixture is heated at 65°C for 18 hours, the methylene chloride layer separated, dried
- 25 (Na₂SO₄) and solvent evaporated to yield the crude product as a dark viscous oil. The oil is chromatographed over 700 g of silica gel eluting with 25% ethyl acetate/hexane to yield 31.42 g of the title compound.

Physical characteristics are as follows:

MP 91.8-93.5

- 30 ¹H NMR (CDCl₃) δ 7.32, 7.08, 6.60, 5.34, 4.67, 4.15, 3.43, 3.02, 2.91, 1.56, 0.65 ppm

PREPARATION 103 (3S)(4S)-3-[3-[3-(Bis(phenylmethyl)amino)phenyl]-2-(2-methyl-1,3-dioxolan-2-yl)-1-oxopentyl]-4-phenyl-2-oxazolidinone (Formula X-8) Refer to Chart X.

- 35 To 12 mL of methylene chloride, under nitrogen, is added 1.55 grams of

(3(3S),4R)-3-[3-(3-bisbenzylaminophenyl)pentanoyl]-4-phenyl-2-oxazolidinone (formula X-6) of Preparation 102 and the resulting solution cooled to -78°C. To the aforementioned solution is added 646 µl of TiCl₄ followed by the addition of 525 µl of diisopropylethylamine. After stirring at 0°C for 30 minutes the reaction is cooled
5 back to -78°C and 886 µl of 2-methoxy-2-methyl-1,3-dioxolane (formula X-7) (also W-7) is added. The reaction is stirred for 1 hour and then quenched by the addition of saturated NH₄Cl, then saturated NaHCO₃ (pH 8) and finally extraction of the aqueous with both methylene chloride and ethyl ether. Evaporation of solvent affords a viscous oil which is chromatographed over 150 g of silica gel eluting with
10 7% hexane/methylene chloride to afford 1.14 g of the title compound.

Physical characteristics are as follows:

IR (mineral oil) 2920, 2954, 2854, 2870, 1776, 1376, 1453, 1196, 699 cm⁻¹.

Anal. found: C, 75.27; H, 6.68; N, 4.55

PREPARATION 104 (3S)(4R) 3-[2-Acetyl-3-[3-[bis(phenylmethyl)amino]-
15 phenyl]-1-oxopentyl]-4-phenyl-2-oxazolidinone (Formula X-9) Refer to Chart X.

To 15 mL of THF is added 960 mg of (3(3S),4R)-3-[2-(2-methyl-1,3-dioxan-2-yl)-3-(3-bisbenzylaminophenyl)pentanoyl]-4-phenyl-2-oxazolidinone (formula X-8) of Preparation 103. To that solution is then added 4 mL of 30% perchloric acid and the
20 resulting mixture stirred at 40°C for 2 hours. The reaction is cooled to room temperature and quenched with the addition of excess saturated NaHCO₃. The reaction is extracted with 200 mL of ethyl ether, dried (Na₂SO₄) and solvent removed in vacuo to yield 981 mg of the crude product. Chromatography over 100 g of silica gel eluding with 10% pentane/methylene chloride affords 854 mg of the title
25 compound.

Physical characteristics are as follows:

¹H NMR (CDCl₃) δ 7.40, 7.08, 6.61, 6.56, 5.41, 4.96, 4.66, 4.61, 4.21, 3.09, 1.63, 1.65, 0.61

IR (mineral oil) 1778, 1718, 1600, 1695, 1452, 1335, 1385, 1200 cm⁻¹.

30 EI-MS: [M+] = 560.

Anal. found: C, 76.81; H, 6.59; N, 4.84.

PREPARATION 105 (3S)(4R) 3-[2-[1-[3-[bis(phenylmethyl)amino]phenyl]-propyl]-5-hydroxy-1,3-dioxo-5-propyloctyl]-4-phenyl-2-oxazolidinone (Formula X-10) Refer to Chart X.

35 To 8 mL of methylene chloride under nitrogen is added 440 mg of (3(3S),4R)-

3-[2-(acetyl)-3-(3-bisbenzylaminophenyl)pentanoyl]-4-phenyl-2-oxazolidinone (formula X-9) of Preparation 104 and that solution is cooled to -78°C . To that solution is added 90 μl of TiCl_4 followed by the addition of 143 μl of diisopropylethylamine. That solution is warmed to 0°C for 40 minutes and then cooled back to -78°C at which time 126 μl of 4-heptanone is added and the reaction temperature is elevated to 0°C and stirring continued for 1.5 hours. The reaction is quenched with the addition of saturated NH_4Cl followed by the addition of saturated NaHCO_3 . The reaction is extracted with methylene chloride (3 X 60 mL), dried (Na_2SO_4) and evaporated in vacuo to yield the crude product as an oil. That material is chromatographed over silica gel (100 g) eluting with 10% pentane/methylene chloride to afford 293 mg of the title compound.

Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 7.28, 7.07, 6.56, 5.44, 5.24, 4.68, 4.61, 4.26, 3.21, 3.10, 2.48, 1.90, 1.55, 1.21, 0.81, 0.74, 0.58

IR (mineral oil) 2959, 2931, 1779, 1720, 1690, 1600, 1494, 1452, 1385, 1359, 1334, 1238, 698 cm^{-1} .

PREPARATION 106 (3R) 3-[1-[3-[bis(phenylmethyl)amino]phenyl]propyl]-5,6-dihydro-4-hydroxy-6,6-dipropyl-2H-pyran-2-one (Formula X-11) Refer to Chart X.

To 3 mL of THF was added 28 mg of NaH under nitrogen. To that suspension is added 418 mg of (3(3S),4R)-3-[2-((3-hydroxy-3-propyl)hexanoyl)-3-(3-bisbenzylaminophenyl)pentanoyl]-4-phenyl-2-oxazolidinone (Formula X-10) of Preparation 105 also in 3 mL of THF at 20°C . The reaction is stirred for 16 hours, cooled to 0°C and quenched by addition of 1N HCl. The reaction is then made basic with the addition of saturated NaHCO_3 . The aqueous is extracted several times with ethyl acetate, the organic extracts dried (Na_2SO_4) and solvent is removed in vacuo to yield 518 mg of crude product. Chromatography over silica gel eluting with 15% EtOAc/hexane affords 128 mg of the title compound.

Physical characteristics are as follows:

IR (mineral oil) 2959, 2931, 2873, 1636, 1599, 1451, 1465, 1386, 1363, 1328, 1249, 1260, 696 cm^{-1} .

EI-MS: $[\text{M}^+] = 511$.

PREPARATION 107 (3R) 3-[1-[3-[amino]phenyl]propyl]-5,6-dihydro-4-hydroxy-6,6-dipropyl-2H-pyran-2-one (Formula X-12) Refer to Chart X.

The dihydropyrone of formula X-11 ((3R)-3-[1-(3-bisbenzylaminophenyl)-propyl]-6,6-bispropyl-5,6-dihydro-4-hydroxypyran-2-one) of Preparation 106, 110 mg, is added to 20 mL of ethyl acetate. To that solution is added 50 mg of 10% Pd/C and the resulting mixture is hydrogenated at 50 psi for 6 hours. The reaction is filtered
5 through celite to yield 83 mg of the title product.

Physical characteristics are as follows:

IR (mineral oil) 2961, 2932, 2873, 1682, 1623, 1604, 1458, 1384, 1369, 1319, 1282, 1259, 1150, 1108 cm^{-1} .

EI-MS: $[M^+] = 331$

10 PREPARATION 108 2-Phenethyl-2-propen-1-ol (Formula BB-2) Refer to Chart BB.

To a cooled (-10°C) solution of *N,N,N,N*-tetramethyl-1,2-ethylenediamine (24.1 mL) in hexane (50 mL) is slowly added butyl lithium (100 mL of a 1.6 M solution in hexane). After stirring for 45 minutes at -10°C the mixture is cooled (-78°C) and 2-methyl-2-propen-1-ol (BB-1, 6.41 mL) is added dropwise. The reaction
15 is allowed to warm to room temperature and stirred an additional 72 h. The mixture is cooled to -78°C and a solution of benzyl bromide (8.6 mL) in anhydrous THF (10 mL) is added slowly. The mixture is stirred at -78°C for 1 hour then gradually allowed to warm to room temperature. After stirring an additional 2
20 hours, the reaction is quenched by the addition of saturated aqueous NH_4Cl . The organic layer is diluted with diethyl ether and washed with brine, dried (MgSO_4), filtered and concentrated in vacuo. Purification by flash chromatography using methylene chloride/ethyl acetate/hexane (1:1:6) as eluent affords the title compound (3.5 g) as an oil.

25 Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 7.31-7.16, 5.07, 4.93, 4.09, 2.82-2.76, 2.41-2.36 ppm.

^{13}C NMR (CDCl_3) δ 148.30, 141.69, 128.24, 125.80, 109.76, 65.90, 34.52, 34.16 ppm.

PREPARATION 109 (2S)-2-Phenethyloxiranemethanol (Formula BB-8) Refer
30 to Chart BB.

To a cooled (-20°C) slurry of molecular sieves (4 Å, crushed and freshly activated, 150 mg) in methylene chloride (1.5 mL) is added diethyl L-tartrate (22 mg) and titanium(IV) isopropoxide (25 mg). The mixture is stirred for 30 min at -20°C and *tert*-butyl hydroperoxide (0.84 mL of a 5-6 M solution in nonane) is added. After
35 an additional 25 min at -20°C , a solution of allylic alcohol of formula BB-2 (300 mg)

of Preparation 108 in methylene chloride (0.5 mL) is slowly added. The mixture is stirred overnight at -20°C then warmed to -10°C. After an additional 4 hours the reaction is warmed to 0-5°C and quenched with the addition of water (1 mL). After warming to room temperature, stirring is continued for 1 hour and tartrates
5 hydrolysed by the addition of a 30% aqueous NaOH solution saturated with NaCl (0.1 mL). After 30 minutes, the mixture is filtered through Celite and the aqueous phase extracted with several portions of methylene chloride. The combined organic layers are dried (MgSO₄), filtered and concentrated in vacuo to provide a residue which is purified by flash chromatography using hexane and a gradient of ethyl
10 acetate (10-20%) as eluent to afford the title product of formula BB-3 (223 mg) as an oil. The enantiomeric excess of the reaction is determined to be 86% by analysis of the ¹H NMR (C₆D₆) of the Mosher ester formed by the reaction of BB-3 with (S)-(+)-α-methoxy-α-(trifluoromethyl)phenylacetyl chloride (J. A. Dale, D. L. Dull, H. S. Mosher, J. Org. Chem. (1969) 34: 2543).

15 Physical characteristics are as follows:

¹H NMR (CDCl₃) δ 7.34-7.16, 3.83-3.61, 2.89-2.87, 2.72-2.64, 2.17-2.04, 1.89-1.79 ppm

¹³C NMR (CDCl₃) δ 141.17, 137.58, 128.49, 128.21, 126.11, 63.00 59.57, 49.92, 33.58, 30.82 ppm.

20 PREPARATION 110 (2S)-2-Phenethyl-2-phenylmethoxymethyloxirane
(Formula BB-9) Refer to Chart BB.

To a cooled (0-5 °C) slurry of sodium hydride (124 mg of a 60% suspension in mineral oil) in THF (10 mL) is added alcohol of formula BB-3 (460 mg) of Preparation 109. The mixture is stirred at 0-5 °C for 5 minutes, allowed to warm to
25 room temperature and stirred an additional 30 minutes. Benzyl bromide (441 mg) is added and the mixture stirred at room temperature overnight. The mixture is quenched with brine (10 mL) and diluted with ethyl ether. The organic layer is washed with brine, dried (MgSO₄), filtered and concentrated in vacuo to provide a residue which is purified by flash chromatography using hexane and a gradient of
30 ethyl acetate (2-5%) as eluent to afford the title product (510 mg) as an oil.

Physical characteristics are as follows:

¹H NMR (CDCl₃) δ 7.34-7.13, 4.59-4.49, 3.64-3.45, 2.75-2.59, 2.19-2.09, 1.94-1.84

¹³C NMR (CDCl₃) δ 141.42, 137.93, 128.58, 128.38, 127.88, 125.94, 73.23,
35 71.98, 58.21, 50.37, 33.65, 30.83 ppm.

PREPARATION 110a (3S)-1-Phenyl-3-(phenylmethoxymethyl) hexan-3-ol
(Formula BB-10) Refer to Chart BB.

To a cooled (-45 °C) solution of Li_2CuCl_4 (0.28 mL of a 0.1 M solution in THF) in THF (2 mL) is added ethylmagnesium bromide (0.203 mL of a 3 M solution in ethyl ether). The brown solution is stirred at -45 °C for 45 minutes and the epoxide of formula BB-4 (150 mg) of Preparation 110 is added dropwise over ca. 10 minutes. After one hour the reaction is quenched by the addition of saturated aqueous NH_4Cl and the aqueous layer extracted with ethyl acetate. The combined organic layers are washed with brine, dried (MgSO_4), filtered and concentrated in vacuo to provide a residue which is purified by flash chromatography using hexane/ethyl acetate (5%) as eluent to afford the title product (150 mg) as an oil.

Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 7.38-7.14, 4.54, 3.37, 2.66-2.58, 2.20, 1.88-1.76, 1.58-1.52, 1.39-1.25, 0.92

^{13}C NMR (CDCl_3) δ 142.61, 138.10, 128.41, 128.33, 127.71, 127.63, 125.68, 75.45, 73.79, 73.44, 38.95, 38.52, 29.86, 16.79, 14.37 ppm.

PREPARATION 110b (2R)-2-Phenethyl-2-(p-toluenesulfonyloxymethyl) oxirane
(Formula BB-13) Refer to Chart BB.

To a cooled (ca. -10 °C) solution of the compound of formula BB-8 (245 mg) of Preparation 109 in methylene chloride (4 mL) is added 4-toluenesulfonyl chloride (302 mg), triethylamine (160 mg) and 4-dimethylaminopyridine (8 mg). The mixture is stirred at ca. -10 °C overnight then warmed to 0-5 °C for 1 hour. The mixture is diluted with methylene chloride, washed with saturated aqueous NaHCO_3 and brine, dried (MgSO_4), filtered and concentrated in vacuo to provide a residue which is purified by flash chromatography using hexane/ethyl acetate (5%) as eluent to afford the title compound (448 mg).

Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 7.79, 7.33, 7.28-7.08, 4.13-3.98, 2.64-2.58, 2.44, 2.11-2.00, 1.92-1.82

^{13}C NMR (CDCl_3) δ 145.17, 140.67, 132.53, 129.96, 128.48, 128.19, 127.95, 126.15, 71.98, 56.49, 50.63, 32.89, 30.37, 21.64 ppm.

PREPARATION 110c (2S)-2-Phenethyl-2-propyl oxirane (Formula BB-12) Refer to Chart BB.

To a cooled (-45 °C) solution of Li_2CuCl_4 (0.3 mL of a 0.1 M solution in THF) in THF (2 mL) is added ethylmagnesium bromide (0.22 mL of a 3 M solution in

ethyl ether). The brown solution is stirred at -45 °C for 45 minutes, cooled to -65 °C then tosylate of formula BB-13 (200 mg) of Preparation 110b is added dropwise over ca. 10 minutes. The mixture is stirred for 2.5 hours, warmed to -50 °C for 2 hours and then quenched by the addition of saturated aqueous NH₄Cl. The aqueous layer
5 is extracted with ethyl acetate and the combined organic layers washed with brine, dried (MgSO₄), filtered and concentrated in vacuo to provide a residue which is purified by flash chromatography using hexane/ ethyl acetate (5%) as eluent to afford the title product (60 mg) and hydroxytosylate of formula BB-14 (47 mg).

Hydroxytosylate of formula BB-14 is converted to the epoxide of formula BB-
10 12 as follows: To a cooled (0-5 °C) solution of the compound of formula BB-14 (43 mg) in methanol (2 mL) is added anhydrous K₂CO₃ (20 mg). After 1 hour at 0-5°C the mixture is warmed to room temperature, stirred an additional 90 minutes then quenched by the addition of saturated aqueous NH₄Cl. The aqueous layer is extracted with ethyl acetate and the combined organic layers washed with brine,
15 dried (MgSO₄), filtered and concentrated in vacuo to provide a residue which is purified by flash chromatography using hexane/ ethyl acetate (5%) to afford the epoxide of formula BB-12 (20 mg).

Physical characteristics for BB-12 are as follows:

¹H NMR (CDCl₃) δ 7.31-7.16, 2.68, 2.59, 1.98-1.82, 1.73-1.37, 0.94
20 ¹³C NMR (CDCl₃) δ 141.71, 128.41, 128.24, 125.92, 59.11, 52.57, 36.42, 36.02, 31.03, 18.19, 14.22 ppm.

Physical characteristics for BB-14 are as follows:

¹H NMR (CDCl₃) δ 7.79, 7.34, 7.29-7.11, 3.90, 2.58-2.53, 2.44, 1.87, 1.77-1.72, 1.54-1.48, 1.31-1.21, 0.89
25 ¹³C NMR (CDCl₃) δ 145.10, 141.66, 132.50, 129.97, 128.45, 128.25, 127.97, 125.96, 74.33, 73.07, 38.37, 37.87, 29.37, 21.66, 16.47, 14.48 ppm.

PREPARATION 111 (4S)-3-acetyl-4-phenyl-2-oxazolidinone (Formula FF-3)
Refer to Chart FF

To a solution of (S)-(+)-4-phenyl-2-oxazolidinone of formula FF-2 (20 g) in
30 anhydrous tetrahydrofuran (600 mL), cooled to -78°C is added a solution of 1.6 M n-butyllithium in hexanes (77.8 mL) and the resulting suspension stirred at -78°C for 30 minutes. The suspension is treated with acetyl chloride of formula FF-1 (10.23 mL) and then gradually allowed to warm to room temperature. The reaction mixture is quenched with 1 L of saturated ammonium chloride and then partitioned
35 between water and ethyl acetate. The organic layer is separated and the aqueous

layer reextracted twice with ethyl acetate. The combined organic layers are washed with brine, dried over anhydrous sodium sulfate, and concentrated *in vacuo*. The crude solid is recrystallized from ethyl acetate/hexane affording (21.27 g,) as a white solid.

5 Physical Characteristics are as follows:

Mp 86-87°C

¹H NMR (CDCl₃) δ 7.42-7.26, 5.44-5.40, 4.68, 4.30-4.26, 2.52 ppm

¹³C NMR (CDCl₃) δ 169.50, 153.71, 138.81, 128.97, 128.53, 125.73, 69.73, 57.20, 23.59 ppm

10 PREPARATION 112 (3(2E),4S)-3-[4,4-dimethyl (2-pentenoyl)]-4-phenyl-2-oxazolidinone (Formula FF-4) Refer to Chart FF

To a solution of the compound of formula FF-3 of Preparation 111 (21.27 g) in anhydrous methylene chloride (500 mL), cooled to -78°C, is added titanium tetrachloride (12.0 mL) in a dropwise manner. The suspension is treated with diisopropylethylamine (19.9 mL) and is allowed to stir at -78°C for 30 minutes. The suspension is then treated with trimethylacetaldehyde (11.4 mL) followed by diisopropylethylamine (19.9 mL) and allowed to gradually warm to room temperature. After 1 hour the reaction mixture is quenched with water (200 mL) and stirred vigorously for 15 minutes. The organic layer is separated and the aqueous layer is reextracted with methylene chloride. The combined organic layers are washed with brine, dried over anhydrous sodium sulfate, and concentrated *in vacuo*. The crude solid is recrystallized from ethyl acetate/hexane affording 21.6 grams of the title compound as a off-white solid:

Physical characteristics are as follows:

25 Mp 148-149 °C

¹H NMR (CDCl₃) δ 7.42-7.05, 5.51-5.46, 4.69, 4.30-4.25, 1.09ppm

¹³C NMR (CDCl₃) δ 165.17, 161.61, 153.70, 139.16, 129.12, 128.61, 125.97, 115.71, 69.88, 57.74, 34.31, 28.56 ppm;

30 PREPARATION 113 (3(3S),4S)-3-[3-(3-Aminophenyl)-4,4-dimethylpentanoyl]-4-phenyl-2-oxazolidinone (Formula FF-5) Refer to Chart FF.

To a slurry of copper(I) bromide dimethylsulfide complex (18.76 g) in anhydrous tetrahydrofuran (60 mL), cooled to -78°C, is added a 1.0 M solution of 3-[bis(trimethylsilyl)amino]phenylmagnesium chloride in tetrahydrofuran (182.2 mL) and the resulting slurry stirred at -78°C for 5 minutes. The slurry is allowed to

warm to -15°C for 15 minutes and then cooled to -78°C. The slurry is then treated with the compound of formula FF-4 of Preparation 112 (16.6 g) added via a solid addition funnel and allowed to stir at -78°C for 3 hours. The reaction mixture is poured into saturated ammonium chloride (200 mL) and then partitioned between
5 water and ethyl acetate. The organic layer is separated and the aqueous layer (pH 8) is basified to pH 9.5 with concentrated ammonium hydroxide. The aqueous layer is reextracted three times with ethyl acetate, the combined organic layers are washed with brine, dried over anhydrous sodium sulfate, and concentrated *in vacuo*. The crude residue is slurried in chloroform (400 mL) and 200 g of silica gel (230-400
10 mesh) at room temperature for 2 hours. The slurry is filtered and the solids washed several times with chloroform followed by methanol. The filtrate is concentrated *in vacuo*. Purification by flash chromatography eluting with hexane/ethyl acetate (15-40%) afford 17.52 grams of the title compound as a light yellow solid.

Physical characteristics are as follows:

15 ¹H NMR (CDCl₃) δ 7.26-7.12 , 7.01 , 6.74-6.70, 6.61-6.50 , 5.32-5.28 , 4.56 , 4.11-3.95 , 3.48, 2.97-2.91 , 0.91 ppm

¹³C NMR (CDCl₃) δ 172.34, 153.51, 145.37, 142.24, 138.12, 128.67, 128.23, 127.68, 124.71, 119.80, 116.49, 113.02, 69.39, 57.39, 52.30, 34.75, 33.49, 27.83 ppm

PREPARATION 114

20 (3(3S),4S)-3-[3-(3-Bisbenzylaminophenyl)-4,4-dimethylpentanoyl]-4-phenyl-2-oxazolidinone (Formula FF-6) Refer to Chart FF-6

To a solution of the compound of formula FF-5 of Preparation 113 (15.0 g) in methylene chloride (190 mL) at room temperature is added saturated sodium carbonate (48.7 mL) followed by benzyl bromide (14.3 mL) and the resulting mixture
25 is refluxed for 24 hours. The reaction mixture is allowed to cool to room temperature and partitioned between water (300 mL) and methylene chloride. The organic layer is separated, washed with brine, dried over anhydrous sodium sulfate, and concentrated *in vacuo*. Purification by flash chromatography eluting with hexane/ethyl acetate (10-25%) affords 15.1 grams of the title compound as a white
30 solid.

Physical characteristics are as follows:

¹H NMR (CDCl₃) δ 7.29-6.99, 6.69-6.49 , 5.32-5.26, 4.71-4.50, 4.06-3.94, 2.90-2.81 , 0.73ppm

35 ¹³C NMR (CDCl₃) δ 172.79, 153.78, 148.32, 142.08, 139.04, 138.48, 129.07, 128.59, 128.37, 127.89, 126.83, 124.92, 118.44, 114.64, 110.87, 69.70, 57.68, 54.77,

52.97, 34.84, 33.79, 27.97 ppm

PREPARATION 115 [S,[R*,S*(E)]]-N-(2-hydroxy-1-methyl-2-phenylethyl)-methyl-N-pentenamide (Formula NNN-3) See Chart NNN.

5 A 250-mL, three-necked, round-bottomed flask with nitrogen inlet and addition funnel is charged with the compound of formula NNN-1 (6.6 g) (prepared from the treatment of commercially available trans-2-pentenoic acid with oxalyl chloride) and tetrahydrofuran (80 mL). The addition funnel is charged with a solution of (1R, 2S)-ephedrine of formula NNN-2 (7.2 g) and triethylamine (6.0 mL)
10 in tetrahydrofuran (15 mL), which is added dropwise to the reaction mixture. After stirring an additional hour, the reaction mixture is poured into 200 mL of ethyl acetate, washed with three 25-mL portions of water, and concentrated *in vacuo* to yield 13.5 g of an oil. Column chromatography on 100 g silica (elution with 10-100% ethyl acetate-hexane) yields 10.75 g of the title compound as a colorless oil.

15 Physical characteristics are as follows:

HRMS found: 248.1652.

PREPARATION 116 [1R-[1R*(R*)2S*]]-3-Amino-β-ethyl-N-(2-hydroxy-1-methyl-2-phenylethyl)-N-methyl-benzenepropanamide (Formula NNN-4) See Chart NNN.

20 A 50-mL, three-necked round-bottomed flask with a nitrogen inlet is charged with the title compound of Preparation 115 (0.247 g) and 5 mL of *t*-butyl methyl ether and cooled to 0 °C. Propyl magnesium chloride (0.55 mL of 2.0 M solution in ether) is added dropwise, and the reaction mixture is stirred for an additional 15 min. 3-[Bis(trimethylsilyl)amino]phenylmagnesium chloride (2.0 mL of 1.0 M
25 solution in tetrahydrofuran, 2.0 mmol) is added dropwise, and the resulting mixture is stirred for an additional 2 h at 0 °C and 1 h at room temperature. The reaction mixture is then quenched with saturated aqueous ammonium chloride solution (pH adjusted to 8 with ammonium hydroxide) and partitioned between 100 mL of ethyl acetate and 5 mL of water. The organic layer is separated, washed with additional
30 ammonium chloride solution and water, and concentrated *in vacuo* to give 0.72 g of a yellow oil. The crude oil is then dissolved in chloroform, and silica gel is added to the solution. The resulting mixture is stirred at room temperature for 1.5 h, then filtered through Celite, rinsing with methanol, and concentrated *in vacuo* to give
35 0.38 g of a yellow oil. Column chromatography on 50 g of silica gel (elution with 20-100% ethyl acetate-hexane) yields 0.174 g of the title compound as an oil.

Physical characteristics are as follows:

HRMS found: 340.2162.

PREPARATION 117 [1R-[1R*(R*)2S*]]-3-[bis(phenylmethyl)amino]-b-ethyl-N-(2-hydroxy-1-methyl-2-phenylethyl)-N-methyl-benzenepropanamide (Formula NNN-5) See Chart NNN.

A 50-mL, three-necked, round-bottomed flask with a condenser fitted with a nitrogen inlet is charged with the title product of Preparation 116 (0.548 g) in 8 mL of acetonitrile. Sodium carbonate (0.375 g) and benzyl bromide (0.42 mL) are added, and the reaction mixture is heated to reflux for 4 h. The reaction mixture is then concentrated *in vacuo* and partitioned between 100 mL of ethyl acetate and 10 mL of water. The organic layer is separated, washed with another 10 mL of water, and concentrated *in vacuo* to give 1.0 g of a yellow oil. Column chromatography on 65 g of silica gel (elution with 20-100% ethyl acetate-hexane and 5% methanol-methylene chloride) yields 0.447 g of the title compound as a pale yellow oil.

Physical characteristics are as follows:

HRMS found: 520.3102.

PREPARATION 118 1-phenyl-6,6,6-trifluoro-3-hexanol (Formula PPP-2) Refer to Chart PPP.

To a stirred solution of 4.0 g of ethyl 4,4,4-trifluorobutyrate of formula PPP-1 in 25 mL of tetrahydrofuran at -70°C 24 mL of DiBAL-H (1M in toluene) is added dropwise and the solution stirred for 90 min. In a separate flask containing 680 mg of magnesium turnings and 5 mL of tetrahydrofuran is added 1-phenyl-2-bromoethane in 20 mL of tetrahydrofuran at a rate to maintain reflux. Heating of the mixture is continued for an additional 1 h, then cooled to room temperature and added via cannula to the DiBAL-H reaction above. The resulting white suspension is stirred 30 min at -70°C and then allowed to warm to room temperature. The reaction is quenched with saturated aqueous ammonium chloride, diluted with 1 N hydrochloric acid to dissolve the precipitated salts and extracted with ethyl acetate. The organic layers are combined, washed with water and brine, dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The crude product is flash chromatographed on silica gel eluting with 20% ethyl acetate in hexane to give 2.0 g of the title compound as a colorless oil.

Physical characteristics are as follows:

HRMS: 232.1088

IR (neat liquid): 3385, 2950, 1455, 1255, 1140, 700 cm⁻¹.

PREPARATION 119 1-phenyl-6,6,6-trifluoro-3-hexanone (Formula PPP-3)

Refer to Chart PPP.

To a solution of 0.48 mL of oxalyl chloride in 10 mL of dichloromethane at -60°C is added dropwise 0.81 mL of dimethylsulfoxide. The solution is stirred for 5 min then treated with 860 mg of 1-phenyl-6,6,6-trifluoro-3-hexanol of formula PPP-2 of Preparation 118 in 5 mL of dichloromethane and stirred for 15 min. Triethylamine (1.5 mL) is added, the mixture is allowed to warm to room temperature, diluted with water and the layers separated. The aqueous layer is extracted with dichloromethane, the organic layers combined, dried over anhydrous sodium sulfate, filtered, and concentrated. The resulting oil is flash chromatographed on silica gel to give 600 mg of the title compound as an oil.

Physical characteristics are as follows:

 ^1H NMR (CDCl_3): δ 7.2-7.3, 2.9, 2.7, 2.6, 2.4. ^{13}C NMR (CDCl_3): δ 206, 140, 129, 128, 126, 125, 44, 35, 30, 28.

15 PREPARATION 120 5,6-dihydro-4-hydroxy-6-phenethyl-6-(3',3',3'-trifluoropropyl)-2H-pyran-2-one (Formula PPP-4) Refer to Chart PPP.

A suspension of 350 mg of 50% sodium hydride in 10 mL of tetrahydrofuran at 0°C is treated dropwise with 0.78 mL of methyl acetoacetate. After stirring 30 min, 4.5 mL of 1.6 M n-butyllithium in hexane is added and stirring continued for 15 min. A solution of 840 mg of 1-phenyl-6,6,6-trifluoro-3-hexanone in 5 mL of tetrahydrofuran of formula PPP-3 of Preparation 119, is added, stirred at 0°C for 15 min, then allowed to warm to room temperature and stirred for 1 h. The reaction mixture is quenched with saturated aqueous ammonium chloride and extracted with ethyl acetate. The organic layers are washed with water and brine, concentrated in vacuo, then dissolved in 20 mL of tetrahydrofuran. The solution is diluted with 60 mL of water and treated with 20 mL of 1 N sodium hydroxide, stirred for 3 h at room temperature, concentrated in vacuo to remove the tetrahydrofuran, cooled to 5°C, and acidified with concentrated hydrochloric acid. The mixture is extracted with dichloromethane, dried over anhydrous sodium sulfate, filtered and concentrated. The crude material is flash chromatographed on silica gel eluting with 30% ethyl acetate in hexane to give 870 mg of the title compound.

Physical characteristics are as follows:

ANAL: C, 61.14, H, 5.45.

35 PREPARATION 121 [3-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-

(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl-carbamic acid, phenylmethylester
(Formula QQQ-3 where $R_1=t\text{-Bu}$) Refer to Chart QQQ.

The title compound of Preparation 120 (850 mg) in 25 mL of tetrahydrofuran at 0°C is treated with 750 mg of aluminum trichloride, stirred 15 min, and then 700 mg of 3-benzyloxycarbonylaminobenzaldehyde is added. The mixture is allowed to stir at room temperature for 2 h then treated with 2 g of sodium carbonate monohydrate and 0.1 mL of water, stirred for 30 min, and filtered through celite washing the filter cake with tetrahydrofuran. The filtrate is concentrated in vacuo. The resulting material is dissolved in 25 mL of tetrahydrofuran and 285 mg of cuprous bromide-dimethylsulfide complex is added and the mixture stirred for 15 min before adding 11 mL of 1 M *t*-butylmagnesium bromide in tetrahydrofuran dropwise over 15-20 min. The resulting brown mixture is stirred an additional 15 min then quenched at 0°C with 50 mL of water. The layers are separated and the aqueous layer acidified with concentrated hydrochloric acid to dissolve inorganic salts and then extracted with ethyl acetate. The combined organic layers are washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography on silica gel eluting with ethyl acetate in hexane gives 1.19 g of the title compound as a white to buff colored foam.

Physical characteristics are as follows:

^1H NMR (CDCl_3): δ 7.1-7.7, 6.7, 6.5, 4.4, 1.8-2.8, 1.16.

HRMS: 609.2711.

PREPARATION 122

Preparative resolution of [3-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl-carbamic acid, phenylmethylester (Formula QQQ-3 where $R_1=t\text{-Bu}$) into 4 isomers, 3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(R or S)-(2-phenethyl)-6-(R or S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethyl-propyl]phenyl-carbamic acid, phenylmethylester (Formulas QQQ-4-7 where R_1 is *t*-Bu) Refer to Chart QQQ.

The first phase of the resolution is accomplished with a 5.1x25 cm (R,R)Whelk-O 1 column eluted with 15% (V/V) isopropanol in hexane at 99 mL/min. ((R,R)Whelk-O 1 is a registered trademark of Regis Technologies, Inc., Morton Grove, IL 60053.) The peaks eluting at approximately 54 and 87 min are,

respectively, pure Isomer 3 and Isomer 2 as judged from System A, below. The mixture of unresolved Isomer 1 and Isomer 4 eluted at approximately 64 minutes and is further treated as described below.

In the second phase of the resolution, the mixture from above that elutes near 64 minutes is injected onto a 2.1x25 cm Chiralcel OD column (Chiral Technologies, Inc.) and eluted with 35% isopropanol in hexane (V/V) at 8 mL/min. The peaks that elute near 8.7 and 23.9 minutes are, respectively, Isomer 1 and Isomer 4.

In both phase of the resolution of enantiomers, fractions are pooled after assay with System A and pools are concentrated to dryness on a rotary evaporator at 30mm and a bath set at 50° maximum.

The four constituent enantiomers are (in order of elution from system A) designated (Peak #1), (Peak #2), (Peak #3) and (Peak #4). System A consists of a 0.46x25 cm Chiralcel OD-H column eluted at 1.0 mL/min with 20% isopropanol in hexane (V/V). (Chiralcel OD-H is a registered trademark of Chiral Technologies, Inc., Exton PA 19341.)

PREPARATION 123 3(R or S)-[1-(3-aminophenyl)-2,2-dimethylpropyl]-4-hydroxy-5,6-dihydro-6-(R or S)-phenethyl-6-(3,3,3-trifluoropropyl)-2H-pyran-2-one (Formula QQQ-8, R₁=*t*-Bu) Refer to Chart QQQ.

A solution of 210 mg of the compound identified as peak 1 from Preparation 122 in 10 mL of methanol is treated with 400 mg of ammonium formate and 40 mg of 10% palladium on charcoal, stirred 2 h, filtered through celite washing the filter cake with methanol. The filtrate is diluted with ethyl acetate and washed with water and brine, dried over anhydrous sodium sulfate, filtered and concentrated in vacuo to give 160 mg of the title compound as a white foam.

Physical characteristics are as follows:

¹H NMR (CD₃OD): δ 6.9-7.3, 6.6, 4.1, 2.6-2.7, 1.9-2.4, 1.0.

TLC (silica gel GF): R_f=0.24 (40% ethyl acetate in hexane).

EXAMPLE 266 5-cyano-N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R or S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide (Formula QQQ-13, R₁ is *t*-Bu [R₂ is [5-cyano-2-pyridinyl]]) Refer to Chart QQQ.

A solution of the title product of Preparation 123 (50 mg), pyridine (30 mL), and 5-cyano-pyridine-2-sulfonyl chloride (30 mg) in dichloromethane at 0°C is stirred

for 2 h. The crude reaction mixture is chromatographed on silica gel to give the title compound as a white amorphous solid.

Physical characteristics are as follows:

FAB HRMS: 642.2267.

- 5 **EXAMPLE 267** N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(R or S)-(2-phenethyl)-6-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula QQQ-13, R₁ is *t*-Bu, [R₂ is 1-methyl-4-imidazolyl]) Refer to Chart QQQ.

- 10 Following the procedure described in EXAMPLE 266 and substituting 5-cyano-2-pyridine sulfonyl chloride with 1-methylimidazole-4-sulfonyl chloride the title compound is prepared.

Physical characteristics are as follows:

HRMS: 619.2298

- 15 **EXAMPLE 268** 5-amino-N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R or S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide (Formula QQQ-13, R₁=*t*-Bu, [R₂ is 5-amino-2-pyridinyl]) Refer to Chart QQQ.

- 20 A solution of the title product of Preparation 123 (50 mg), pyridine (30 μ L), and 5-nitro-pyridine-2-sulfonyl chloride (30 mg) in dichloromethane at 0°C is stirred for 2 h. The crude reaction mixture is chromatographed on silica gel to give the sulfonamide as a white amorphous solid. The white solid is dissolved in 4 mL of methanol and treated with 25 mg of ammonium formate and 5 mg of 10% palladium on carbon, stirred for 1 h at room temperature, diluted with water and extracted
25 with dichloromethane, dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to give the title compound as an off-white amorphous solid.

Physical characteristics are as follows:

FAB HRMS: 632.2393.

- 30 **PREPARATION 124** 3(R or S)-[1-(3-aminophenyl)-2,2-dimethyl-propyl]-4-hydroxy-5,6-dihydro-6(S or R)-phenethyl-6-(3',3',3'-trifluoropropyl)-2H-pyran-2-one (Formula QQQ-9, [R₁ is *t*-Bu]) Refer to Chart QQQ.

- 35 Following the procedure described in Preparation 123 beginning with the compound isolated from peak 2 from Preparation 122 and using starting materials and reagents known and available to one of ordinary skill in organic synthesis the

title compound is prepared.

Physical characteristics are as follows:

¹H NMR (CD₃OD): δ 6.9-7.3, 6.6, 4.1, 2.6-2.7, 1.9-2.4, 1.0.

TLC (silica gel GF): R_f=0.24 (40% ethyl acetate in hexane).

- 5 **EXAMPLE 269** 5-cyano-N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R or S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide (Formula QQQ-14, R₁ is *t*-Bu [R₂ is 5-cyano-2-pyridinyl]) Refer to Chart QQQ.

10 A solution of the title product of Preparation 124 (50 mg), pyridine (30 μL), and 5-cyanopyridine-2-sulfonyl chloride (30 mg) in dichloromethane at 0°C is stirred for 2 h. The crude reaction mixture is chromatographed on silica gel to give the title compound as a white amorphous solid.

Physical characteristics are as follows:

FAB HRMS: 642.2260.

- 15 **EXAMPLE 270** N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(R or S)-(2-phenethyl)-6-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula QQQ-14, R₁ is *t*-Bu [R₂ is 1-methyl-4-imidazolyl]) Refer to Chart QQQ.

20 Following the procedure described in Example 266 substituting 5-cyano-2-pyridine sulfonyl chloride with 1-methylimidazole-4-sulfonyl chloride the title compound is prepared.

Physical characteristics are as follows:

HRMS: 619.2362

- 25 **EXAMPLE 271** 5-amino-N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R or S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide (Formula QQQ-14, R₁ is *t*-Bu [R₂ is 5-amino-2-pyridinyl]) Refer to Chart QQQ.

30 Following the procedure described in Example 268 the title compound is prepared.

Physical characteristics are as follows:

FAB HRMS: 632.2387

- PREPARATION 125 3-(S or R)-[1-(3-aminophenyl)-2,2-dimethyl-propyl]-4-hydroxy-5,6-dihydro-6-(R or S)-phenethyl-6-(R or S)-(3',3',3'-trifluoropropyl)-2H-pyran-2-one (Formula QQQ-
- 35

10 [R₁ is *t*-Bu]) Refer to Chart QQQ.

Following the procedure described in Preparation 123 beginning with the compound isolated from peak 3 from Preparation 122 and using starting materials and reagents known and available to one of ordinary skill in organic synthesis the title compound is prepared.

Physical characteristics are as follows:

¹H NMR (CD₃OD): δ 6.9-7.3, 6.6, 4.1, 2.6-2.7, 1.9-2.4, 1.0.

TLC (silica gel GF): R_f=0.24 (40% ethyl acetate in hexane).

EXAMPLE 272 5-cyano-N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R or S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide (Formula QQQ-15, R₁ is *t*-Bu, [R₂ is 5-cyano-2-pyridinyl]) Refer to Chart QQQ.

A solution of the title product of Preparation 125 (50 mg), pyridine (30 mL), and 5-cyanopyridine-2-sulfonyl chloride (30 mg) in dichloromethane at 0°C is stirred for 2 h. The crude reaction mixture is chromatographed on silica gel to give the title compound as a white amorphous solid.

Physical characteristics are as follows:

FAB HRMS: 642.2254

EXAMPLE 273 N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(R or S)-(2-phenethyl)-6-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula QQQ-15, R₁ is *t*-Bu, [R₂ is 1-methyl-4-imidazolyl]) Refer to Chart QQQ.

Following the procedure described in Example 266 substituting 5-cyano-2-pyridine sulfonyl chloride with 1-methylimidazole-4-sulfonyl chloride the title compound is prepared.

Physical characteristics are as follows:

FAB HRMS: 642.2397

EXAMPLE 274 5-amino-N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R or S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide (Formula QQQ-15, R₁ is *t*-Bu [R₂ is 5-amino-2-pyridinyl]) Refer to Chart QQQ.

Following the procedure described in Example 268 the title compound is prepared.

Physical characteristics are as follows:

FAB HRMS: 632.2393

PREPARATION 126 3-(S or R)-[1-(3-aminophenyl)-2,2-dimethyl-propyl]-4-hydroxy-5,6-dihydro-6 (S or R)-phenethyl-6-(3',3',3'-trifluoropropyl)-2H-pyran-2-one (Formula QQQ-11 [R_1 is *t*-Bu]) Refer to Chart QQQ.

Following the procedure described in Preparation 123 beginning with the compound isolated from peak 4 from Preparation 122 and using starting materials and reagents known and available to one of ordinary skill in organic synthesis the title compound is prepared.

Physical characteristics are as follows:

^1H NMR (CD_3OD): δ 6.9-7.3, 6.6, 4.1, 2.6-2.7, 1.9-2.4, 1.0.

TLC (silica gel GF): $R_f=0.24$ (40% ethyl acetate in hexane).

EXAMPLE 275 5-cyano-N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R or S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide (Formula QQQ-16, R_1 is *t*-Bu, [R_2 is 5-cyano-2-pyridinyl]) Refer to Chart QQQ.

A solution of the title product of Preparation 126 (50 mg), pyridine (30 mL), and 5-cyanopyridine-2-sulfonyl chloride (30 mg) in dichloromethane at 0°C is stirred for 2 h. The crude reaction mixture is chromatographed on silica gel to give the title compound as a white amorphous solid.

Physical characteristics are as follows:

FAB HRMS: 642.2248.

EXAMPLE 276 N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(R or S)-(2-phenethyl)-6-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula QQQ-16, R_1 is *t*-Bu, [R_2 is 1-methyl-4-imidazolyl]) Refer to Chart QQQ.

Following the procedure described in Example 266 substituting 5-cyano-2-pyridine sulfonyl chloride with 1-methylimidazole-4-sulfonyl chloride the title compound is prepared.

Physical characteristics are as follows:

FAB HRMS: 620.2403

EXAMPLE 277 5-amino-N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R or S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide (Formula QQQ-

2187523

16, R₁ is *t*-Bu, [R₂ is 5-amino-2-pyridinyl]) Refer to Chart QQQ.

Following the procedure described in Example 268 the title compound is prepared.

Physical characteristics are as follows:

5 FAB HRMS: 632.2406

EXAMPLE 278 N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-[6-(R or S)-propyl]-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide

Following procedures analogous to those described above and using Isomer 2
10 of Preparation 143 the title compound is prepared.

Physical characteristics are as follows:

FAB HRMS: 562.2527.

EXAMPLE 279 N-[3-(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R or S)-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-
15 pyridinesulfonamide

Following procedures analogous to those described above and using Isomer 1
of Preparation 143 the title compound is prepared.

Physical characteristics are as follows:

FAB HRMS: 562.2528.

20 PREPARATION 127 2-Mercapto-4-trifluoromethylpyridine

To 1.0g of 2-chloro-4-trifluoromethylpyridine (Lancaster Chemical Co) is added 10 ml of absolute ethanol and 417mg of thiourea. The reaction mixture is heated at reflux for 4 hours and 1.25 ml of a solution of 7.44g KOH in 20 ml of water is added. The solution is heated at reflux for an additional 1 hour. The
25 reaction solution is cooled and poured into 100 ml of a 0.1N NaOH solution. The resulting solution is extracted three times with 100 ml of methylene chloride and the resulting aqueous solution is acidified to pH 4 by addition of glacial acetic acid. The aqueous solution is extracted three times with 100 ml of methylene chloride and the organic solution is dried over anhydrous sodium sulfate. Filtration followed by
30 evaporation to dryness gives 501mg of a yellow crystalline solid.

Physical characteristics are as follows:

Found C:40.22; H:2.33; N:8.07; S:17.59

HRMS: 179.0019

PREPARATION 128 2-Chlorosulfonyl-4-trifluoromethylpyridine

35 To 425 mg of 2-mercapto-4-trifluoromethylpyridine of Preparation 127 is

added 10ml of 1N aqueous HCl. The reaction mixture is cooled to 0°C and Cl₂ gas is bubbled into the cold reaction mixture for 15 minutes. The reaction mixture is filtered and the resulting solid is washed well with water. The white solid is dissolved in methylene chloride and is washed twice with saturated aqueous NaHCO₃ followed by one wash with water. After drying the organic solution over sodium sulfate (anhydrous), the solution is filtered and evaporated to dryness to give 300mg of 2-chlorosulfonyl-4-trifluoromethylpyridine which is used directly without further purification, and stored at -78°C until ready for use.

PREPARATION 129 2-Chlorosulfonyl-5-trifluoromethylpyridine

Substituting 2-mercapto-5-trifluoromethylpyridine for 2-mercapto-4-trifluoromethylpyridine in the reaction above in Preparation 128 gives 2-chlorosulfonyl-5-trifluoromethylpyridine as a colorless oil which slowly crystallizes. This material is used without further purification and stored at -78°C until ready for use.

PREPARATION 130 3-[1-[5,6-Dihydro-4-hydroxy-2-oxo-6,6-di-n-propyl-2H-pyran-3-yl]-propyl]phenyl-carbamic acid, (Formula SSS-1; R₁ is ethyl; Refer to Chart SSS).

To 7.2g of AlCl₃ at -70°C, under N₂, is added 180 ml of THF. The mixture is allowed to stir at 0°C for 15 minutes and 5.38g of Formula SSS-A; Refer to Chart SSS, prepared by procedures analogous to those described in Preparation 17, is added. The reaction mixture is stirred for 15 minutes and 6.88g of 3-aminoCbZ-benzaldehyde (Formula SSS-B; Refer to Chart SSS) is added. The reaction mixture is stirred for 15 minutes at 0°C followed by 3 hours at room temperature. The reaction is cooled to 0°C and 35g of sodium carbonate monohydrate is added, with vigorous stirring, followed by 1.6 ml of water. After stirring at 0°C for an additional 15 minutes, 120 ml of THF is added and the mixture filtered through celite. The celite is washed well with THF and the THF solution is evaporated to dryness under vacuum to an amber foam. The residue is dissolved in 180 ml of THF, the solution is cooled to -5°C and 3.2g of CuBr.Me₂S added. The mixture is stirred for 15 minutes and 65 ml of a 2M ethylmagnesium chloride in THF solution is added, dropwise, with temperatures not rising above 0°C. The reaction is allowed to stir for an additional 15 minutes and 9 ml of water is slowly added followed by 45 ml of 1N HCl. These additions are done at 0°C. The reaction mixture is poured into 2L of ethyl ether and 200 ml of water is added. The aqueous layer is separated and the organic layer is extracted three times with 10% aqueous ammonium carbonate followed by once with water. The organic solution is dried over anhydrous sodium

2187523

sulfate, filtered and evaporated to dryness to give 10.2g of a crude amorphous foam. This crude material is chromatographed over silica gel using 2% ethyl acetate in methylene chloride as eluent to give 4.74g of 3-[1-[5,6-dihydro-4-hydroxy-2-oxo-6,6-di-n-propyl-2H-pyran-3-yl]-propyl]phenyl-carbamic acid.

- 5 PREPARATION 130A 3-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-phenethyl-6-propyl]-2H-pyran-3-yl]-propyl-phenyl-carbamic acid (Formula RRR-1; R₁ is ethyl; Refer to Chart RRR).

Following the procedure of Preparation 130 beginning with the compound from Preparation 17 the title compound is prepared.

- 10 Physical characteristics are as follows:

¹H NMR (CD₃OD): δ 6.9-7.5, 5.1, 4.0, 1.4-2.7, 0.9.

TLC (silica gel GF): R_f=0.28, 30% ethyl acetate in hexane.

PREPARATION 131

- 15 Preparative chiral resolution of 3-[1-[5,6-dihydro-4-hydroxy-2-oxo-6,6-di-n-propyl-2H-pyran-3-yl]-propyl]phenyl-carbamic acid (Formula SSS-1; R₁ is ethyl; Refer to Chart SSS) to give two isomers of 3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6,6-di-n-propyl-2H-pyran-3-yl]-propyl]phenyl-carbamic acid (Formulas SSS-3 and SSS-4; R₁ is ethyl; Refer to Chart SSS).

- 20 Samples of the title compound of Preparation 130 are injected onto a 2.1x25 cm Chiralcel OD column and eluted with 20% isopropanol (V/V) in hexane at 10 mL/min. The material eluting near 19.1 minutes is 3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6,6-di-n-propyl-2H-pyran-3-yl]-propyl]phenyl-carbamic acid, (α)_D²⁵ +26° (methanol), (Formula SSS-3; Refer to Chart SSS) (peak 1) and that eluting near 37.7 minutes is 3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6,6-di-n-propyl-2H-pyran-3-yl]-propyl]phenyl-carbamic acid ((α)_D²⁵ -27° (methanol), (Formula SSS-4; Refer to Chart SSS) (peak 2). The pools are concentrated separately on a rotary evaporator (ca. 30 mm, bath at 50° maximum) to give white amorphous solids.

- 25 PREPARATION 132 3(R or S)-[1-(3-aminophenyl)-propyl]-4-hydroxy-5,6-dihydro-6,6-dipropyl-2H-pyran-2-one (Formula SSS-5; R₁ is ethyl; Refer to Chart SSS).
- 30

- To 1.04g of 3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6,6-di-n-propyl-2H-pyran-3-yl]-propyl]phenyl-carbamic acid (Formula SSS-3; Refer to Chart SSS) of Preparation 131, the compound identified as peak 1 from the chiral resolution of the product of Preparation 131, is added 20 ml of methanol and 1.29g of ammonium formate. When dissolution is complete, 275mg of 10% Pd/C is added and the
- 35

reaction mixture is stirred at room temperature for 60 minutes. The reaction mixture is filtered (celite) and the methanolic solution is evaporated to dryness to give a crude solid. The crude solid is partitioned between methylene chloride and water. The organic layer is washed twice with water and dried over anhydrous sodium sulfate. The methylene chloride solution is filtered and evaporated to dryness to give 625 mg of 3(R or S)-[1-(3-aminophenyl)-propyl]-4-hydroxy-5,6-dihydro-6,6-dipropyl-2H-pyran-2-one as an amorphous foam.

Physical characteristics are as follows:

MS(EI): 331

10 $(\alpha)^{25}_D +38^\circ$ (c=.3715, methanol).

PREPARATION 133 3(R or S)-[1-(3-aminophenyl)-propyl]-4-hydroxy-5,6-dihydro-6,6-di-n-propyl-2H-pyran-2-one (Formula SSS-6; R_1 is ethyl; Refer to Chart SSS).

To 825mg of 3(R or S)-[1-(3-aminophenyl)-propyl]-4-hydroxy-5,6-dihydro-6,6-dipropyl-2H-pyran-2-one (Formula SSS-4; Refer to Chart SSS), of Preparation 131, the compound identified as peak 2 from the chiral resolution of the product of Preparation 131, is added 20 ml of methanol and 1.02g of ammonium formate. When dissolution is complete, 210mg of 10% Pd/C is added and the reaction mixture is stirred at room temperature for 60 minutes. The reaction mixture is filtered (celite) and the methanolic solution is evaporated to dryness. The crude solid is partitioned between methylene chloride and water. The organic layer is washed twice with water and dried over anhydrous sodium sulfate. The methylene chloride solution is filtered and evaporated to dryness to give 483 mg of title compound as an amorphous foam.

25 Physical characteristics are as follows:

MS(EI): 331

$(\alpha)^{25}_D -39^\circ$ (c=.2680, methanol).

30 EXAMPLE 280 5-Trifluoromethyl-N-[3-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide (Formula SSS-9; R_1 is ethyl; R_2 is 5-trifluoromethylpyridine; Refer to Chart SSS).

The title compound of Preparation 130 is deprotected as in Preparation 132 to give the compound of formula SSS-2. To 132mg of formula SSS-2 is added 15 ml of methylene chloride and 66 microliters of pyridine. The reaction solution is cooled to -5°C and 98mg of 2-chlorosulfonyl-5-trifluoromethylpyridine (product of Preparation

129) is added. After stirring at 0°C for 60 minutes the solution is placed on a silica gel column and eluted with 10% ethyl acetate in methylene chloride until the 5-trifluoromethyl-N-[3-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide is collected. Rf=0.6 in 10% ethyl acetate in methylene chloride. Evaporation of the organic solution to dryness gives 177mg of 5-trifluoromethyl-N-[3-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide.

Physical characteristics are as follows:

MS(EI): 540, 497, 411, 401, 383, 342, 331, 197, 174, 146, 133.

10 HRMS: 540.1938

Rf=0.6 in 10% ethyl acetate in methylene chloride.

¹H NMR(MeOD): 8.91, 8.21-8.19, 7.12, 6.98-6.96, 6.86-6.83, 3.85-3.79, 2.46, 2.10-1.98, 1.84-1.75, 1.58-1.47, 1.27-1.15, .82-.72 ppm

15 EXAMPLE 281 5-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridine-sulfonamide (Formula SSS-7; R₁ is ethyl; R₂ is 5-trifluoromethylpyridine; Refer to Chart SSS).

To 66mg of the title product from Preparation 132 (Formula SSS-5; Chart SSS) is added 8 ml of methylene chloride and 33 microliters of pyridine. The reaction solution is cooled to -5°C and 49mg of 2-chlorosulfonyl-5-trifluoromethylpyridine (product of Preparation 129) is added. After stirring at 0°C for 60 minutes the solution is placed on a silica gel column and eluted with 10% ethyl acetate in methylene chloride until the 5-trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide is collected. Rf=0.6 in 10% ethyl acetate in methylene chloride. Evaporation of the organic solution to dryness gives 69mg of the title compound.

Physical characteristics are as follows:

MS(EI): 540, 497, 411, 401, 383, 342, 331, 197, 174, 146, 133.

Rf=0.6 in 10% ethyl acetate in methylene chloride

30 ¹H NMR(MeOD): 8.91, 8.21-8.19, 7.12, 6.98-6.96, 6.86-6.83, 3.85-3.79, 2.46, 2.10-1.98, 1.84-1.75, 1.58-1.47, 1.27-1.15, .82-.72ppm

EXAMPLE 282 5-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridine-sulfonamide (Formula SSS-8; R₁ is ethyl; R₂ is 5-trifluoromethylpyridine; Refer to Chart SSS).

2187523

Following the procedure of Example 281 but substituting the product of Preparation 133 (formula SSS-6) gives 5-trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide as an amorphous foam.

5 Physical characteristics are as follows:

MS(EI): 540, 497, 411, 401, 383, 342, 331, 197, 174, 146, 133.

R_f=0.6 in 10% ethyl acetate in methylene chloride

¹H NMR(MeOD): 8.91, 8.21-8.19, 7.12, 6.98-6.96, 6.86-6.83, 3.85-3.79, 2.46, 2.10-1.98, 1.84-1.75, 1.58-1.47, 1.27-1.15, .82-.72ppm

10 **EXAMPLE 283** 4-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridine-sulfonamide (Formula SSS-7; R₁ is ethyl; R₂ is 4-trifluoromethylpyridine; Refer to Chart SSS).

Following the procedure of Example 281 but substituting the product of Preparation 128 for the pyridylsulfonylchloride gives 4-trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide as an amorphous foam.

Physical characteristics are as follows:

MS(EI): 146, 145, 139, 133, 71, 57, 55, 43, 41

20 HRMS: 540.1902

EXAMPLE 284 4-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridine-sulfonamide (Formula SSS-8; R₁ is ethyl; R₂ is 4-trifluoromethylpyridine; Refer to Chart SSS).

25 Following the procedure of Example 282 but substituting the product of Preparation 128 for the pyridylsulfonylchloride gives 4-trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide as an amorphous foam.

Physical characteristics are as follows:

30 MS(EI): 146, 145, 139, 133, 71, 57, 55, 43, 41

HRMS: 540.1896

EXAMPLE 285 5-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide (Formula TTT-6; R₁ is *t*-butyl; R₂ is *n*-propyl; R₃ is 5-trifluoromethyl-2-pyridinyl; Refer to Chart TTT).

35

Following the procedure of Example 281 but using Isomer 1 of Preparation 144 (Formula TTT-4; Chart TTT; R₁ is *t*-butyl, R₂ is *n*-propyl) gives 5-trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-*n*-propyl-5,6-dihydro-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide as an amorphous solid.

5 Physical characteristics are as follows:

MS(EI): 163, 162, 147, 146, 69, 57, 56, 43, 41

HRMS: 568.2213

EXAMPLE 286 5-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-*n*-propyl-5,6-dihydro-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide (Formula TTT-7; R₁ is *t*-butyl; R₂ is *n*-propyl; R₃ is 5-trifluoromethylpyridine; Refer to Chart TTT).

Following the procedure of Example 281 but substituting Isomer 2 of Preparation 144 (Formula TTT-5; Chart TTT; R₁ is *t*-butyl; R₂ is *n*-propyl) gives 5-trifluoromethyl-N-[3-[1(R or S)-4-hydroxy-2-oxo-6,6-di-*n*-propyl-5,6-dihydro-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide as an amorphous solid.

Physical characteristics are as follows:

HRMS: 568.2237

EXAMPLE 287 5-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-phenethyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide (Formula TTT-6; R₁ is ethyl; R₂ is phenyl; R₃ is 5-trifluoromethylpyridine; Refer to Chart TTT).

Following the procedure of Example 281 but using Isomer 1 of Preparation 145 (Formula TTT-4; Chart TTT; R₁ is ethyl; R₂ is phenyl) gives 5-trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-*n*-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide as an amorphous solid.

Physical characteristics are as follows:

MS(EI): 665, 647, 456, 455, 333, 134, 133, 117, 105, 91

EXAMPLE 288 5-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-phenethyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide (Formula TTT-7; R₁ is ethyl; R₂ is phenyl; R₃ is 5-trifluoromethylpyridine; Refer to Chart TTT).

Following the procedure of Example 281 but substituting Isomer 2 of Preparation 145 (Formula TTT-5; Chart TTT; R₁ is ethyl; R₂ is phenyl) gives 5-trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-*n*-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide as an amorphous solid.

Physical characteristics are as follows:

HRMS: 665.2300

MS(EI): 665, 647, 456, 455, 333, 134, 133, 117, 105, 91

5 EXAMPLE 289 4-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-phenethyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide (Formula TTT-6; R₁ is ethyl; R₂ is phenyl; R₃ is 4-trifluoromethylpyridine; Refer to Chart TTT).

Following the procedure of Example 281 but substituting Isomer 1 of Preparation 145 (Formula TTT-4; Chart TTT; R₁ is ethyl; R₂ is phenyl) gives 10 trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide as an amorphous solid.

Physical characteristics are as follows:

MS(EI): 666, 665, 647, 134, 133, 117, 105, 91

HRMS: 665.2306

15 EXAMPLE 290 4-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-phenethyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide (Formula TTT-7; R₁ is ethyl; R₂ is phenyl; R₃ is 4-trifluoromethylpyridine; Refer to Chart TTT).

Following the procedure of Example 281 but substituting Isomer 2 of Preparation 145 (Formula TTT-5; Chart TTT; R₁ is ethyl; R₂ is phenyl) gives 20 trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide as an amorphous solid.

Physical characteristics are as follows:

HRMS: 665.2306

25 MS(EI): 666, 665, 647, 134, 133, 117, 105, 91

EXAMPLE 291 4-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide (Formula TTT-6; R₁ is t-butyl; R₂ is methyl; R₃ is 4-trifluoromethylpyridine; Refer to Chart TTT).

30 Following the procedure of Example 283 but substituting Isomer 1 of Preparation 144 (Formula TTT-4; Chart TTT; R₁ is t-butyl; R₂ is methyl) gives 4-trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide as an amorphous solid.

35 Physical characteristics are as follows:

2187523

MS(EI): 525, 512, 428, 411, 302, 284, 258, 146, 57

HRMS: 568.2209

EXAMPLE 292 4-Trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide (Formula TTT-7; R₁ is t-butyl; R₂ is methyl; R₃ is 4-trifluoromethylpyridine; Refer to Chart TTT).

Following the procedure of Example 283 but substituting Isomer 2 of Preparation 144 (Formula TTT-5; Chart TTT; R₁ is t-butyl; R₂ is methyl) gives 4-trifluoromethyl-N-[3-(R or S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide as an amorphous solid.

Physical characteristics are as follows:

MS(EI): 569, 551, 511, 493, 439, 371, 360, 303, 284, 161, 139

HRMS (M⁺+H⁺): 569.2297

PREPARATION 134 N-[3-[1-[5,6-Dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-carbamic acid, phenylmethyl ester (Formula RRR-1; R₁ is t-butyl; Refer to Chart RRR).

To 4.8g of AlCl₃ at -70°C, under N₂, is added 120 ml of THF. The mixture is allowed to stir at 0°C for 15 minutes and 4.68g of (formula RRR-A) of Preparation 17 is added. The reaction mixture is stirred for 15 minutes and 4.59g of 3-aminoCbZ-benzaldehyde (formula RRR-B) is added. The reaction mixture is stirred for 15 minutes at 0°C followed by 3 hours at room temperature. The reaction is cooled to 0°C and 26g of sodium carbonate monohydrate (.21M) is added, with vigorous stirring, followed by 1.08 ml of water. After stirring at 0°C for an additional 15 minutes, the mixture is treated with 120 ml of THF and filtered through celite. The celite is washed well with THF and the THF solution is evaporated to dryness under vacuum to an amber foam. The residue is dissolved in 120 ml of THF, the solution is cooled to -5°C and 2.1g of CuBr.Me₂S added. The mixture is stirred for 15 minutes and 65mL of a 1M t-butylmagnesium chloride in THF solution is added, dropwise, with temperatures not rising above 0°C. The reaction is allowed to stir for an additional 15 minutes at 0°C and 6 ml of water is slowly added followed by 30 ml of 1N HCl. The reaction mixture is poured into 1.3L of ethyl ether. The aqueous layer is separated and the organic layer is extracted three times with 10% aqueous ammonium carbonate followed by once with water.

The organic solution is dried over anhydrous sodium sulfate, filtered and evaporated to dryness to give an amorphous foam. This crude material is chromatographed over silica gel using 30% ethyl acetate in hexane as eluent to give 6.15g the title product.

PREPARATION 135

Preparative resolution of N-[3-[1-[5,6-Dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl-carbamic acid, phenylmethyl ester (Formula RRR-1; R_1 is *t*-butyl) into four isomers, 3-(*R* or *S*)-[1-[5,6-Dihydro-4-hydroxy-2-oxo-6-(*R* or *S*)-(2-phenethyl)-6-(*R* or *S*)-n-propyl-2H-pyran-3-yl]-2,2-dimethyl-propyl]phenyl-carbamic acid, phenylmethyl ester (Formulas RRR-3 to 6; R_1 is *t*-butyl; Refer to Chart RRR).

The four constituent enantiomers are (in order of elution from system A) Isomer 1 (Formula RRR-4; refer to Chart RRR), Isomer 2 (Formula RRR-3; refer to Chart RRR), Isomer 3 (Formula RRR-5; refer to Chart RRR), and Isomer 4 (Formula RRR-6; refer to Chart RRR). System A consists of a 0.46x25 cm Chiralcel OD-H column eluted at 0.5 mL/min with 20% isopropanol and 0.1 % trifluoroacetic acid in hexane (V/V). (Chiralcel OD-H is a registered trademark of Chiral Technologies, Inc., Exton PA 19341.)

The first phase of the resolution is accomplished with a 2.1x25 cm (R,R)Whelk-O 1 column eluted with 20% (V/V) isopropanol in hexane at 12 mL/min. ((R,R)Whelk-O 1 is a registered trademark of Regis Technologies, Inc., Morton Grove, IL 60053.) The peaks eluting at approximately 35 and 41 min are, respectively, a mixture of Isomer 3 and Isomer 4 and a mixture of Isomers 1 and 2 and as judged from System A, above. The two mixtures are further treated as below.

In the second phase of the resolution, the mixture from above that elutes near 41 minutes is injected onto a 2.1x25 cm Chiralcel OD column (Chiral Technologies, Inc.) and elutes with 15% isopropanol and 0.05% trifluoroacetic acid in hexane (V/V) at 9.0 mL/min. The peaks that elute near 11.0 and 22.0 minutes are designated respectively, peaks 1 and 2 and as judged from System A.

In the final phase of the resolution, the mixture that elutes from the Whelk-O column near 35 minutes is injected onto a 2.2x25 cm Chiralcel OD column and elutes with 35% isopropanol and 0.1% trifluoroacetic acid (V/V) in hexane at 9.0 mL/min. The isomer that elutes near 9.7 minutes is designated peak 3 and the one

2187523

that elutes near 16.6 minutes is designated peak 4.

PREPARATION 136 3-[1-(3-Aminophenyl)-2,2-dimethyl-propyl]-5,6-dihydro-4-hydroxy-6-phenethyl-6-n-propyl-pyran-2-one (Formula RRR-2, Refer to Chart RRR).

5 To 590mg of 3-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl-carbamic acid of Preparation 134, is added 10ml of methanol and 660 mg of ammonium formate. When all the reactants are dissolved, 140 mg of 10% Pd/C is added and the reaction is allowed to stir at room temperature for 60 minutes. The reaction is filtered (celite) and the
10 filter pad is washed well with methanol and the methanol solution is evaporated under vacuum to a crude solid. The solid is partitioned between water and methylene chloride, and the methylene chloride layer is washed twice with water, dried over anhydrous sodium sulfate and evaporated to dryness to give 372mg of 3-[1-(3-amino-phenyl)-2,2-dimethyl-propyl]-5,6-dihydro-4-hydroxy-6-phenethyl-6-n-
15 propyl-pyran-2-one. This material is identical to material described earlier (Formula T-4; refer to Chart T).

PREPARATION 137 3(R or S)-[1-(3-amino-phenyl)-2,2-dimethyl-propyl]-5,6-dihydro-4-hydroxy-6(R or S)-phenethyl-6(R or S)-propyl-pyran-2-one (Formula RRR-7; R₁ is t-butyl; Refer to
20 Chart RRR).

Following the procedure of Preparation 136 but substituting the compound in Preparation 135 designated peak 2 for the compound of Preparation 134 gives 3(R or S)-[1-(3-amino-phenyl)-2,2-dimethyl-propyl]-5,6-dihydro-4-hydroxy-6(R or S)-phenethyl-6(R or S)-propyl-pyran-2-one as an amorphous foam.

25 Physical characteristics are as follows:

MS(EI): 421, 365, 164, 163, 147, 146, 118, 107, 91, 57.

HRMS: 421.2617

PREPARATION 138 3(R or S)-[1-(3-Amino-phenyl)-2,2-dimethyl-propyl]-5,6-dihydro-4-hydroxy-6(R or S)-phenethyl-6(R or S)-propyl-pyran-2-one (Formula RRR-8; R₁ is t-butyl; Refer to
30 Chart RRR).

Following the procedure of Preparation 136 but substituting the compound of Preparation 135 designated peak 1 for the compound of Preparation 134 gives 3(R or S)-[1-(3-amino-phenyl)-2,2-dimethyl-propyl]-5,6-dihydro-4-hydroxy-6(R or S)-phenethyl-6(R or S)-propyl-pyran-2-one as an amorphous foam.
35

Physical characteristics are as follows:

MS(EI): 421, 365, 164, 163, 147, 146, 118, 107, 91, 57.

EXAMPLE 293 5-Trifluoromethyl-N-[3-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide (Formula RRR-15; R₁ is t-butyl; R₂ is 5-trifluoromethyl; Refer to Chart RRR).

Following the procedure of Example 281 but substituting the product of Preparation 136 for the product of Preparation 132 gives 5-trifluoromethyl-N-[3-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide as an amorphous foam.

Physical characteristics are as follows:

MS(EI): 497, 411, 401, 383, 343, 331, 197, 174, 146, 133

HRMS: 540.1938

EXAMPLE 294 5-Trifluoromethyl-N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide (Formula RRR-11; R₁ is t-butyl; R₂ is 5-trifluoromethyl; Refer to Chart RRR).

Following the procedure of Example 281 but substituting the product of Preparation 137 for the product of Preparation 132 gives 5-trifluoromethyl-N-[3 (R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide as an amorphous foam.

Physical characteristics are as follows:

MS(EI): 373, 355, 201, 146, 145, 118, 117, 91, 57.

HRMS:630.2394

EXAMPLE 295 5-Trifluoromethyl-N-[3 (R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide (Formula RRR-12; R₁ is t-butyl; R₂ is 5-trifluoromethyl; Refer to Chart RRR).

Following the procedure of Preparation 281 but substituting the product of Preparation 138 for the product of Preparation 132 gives 5-trifluoromethyl-N-[3(R or S)-(-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide as an amorphous foam.

Physical characteristics are as follows:

MS(EI): 373, 355, 201, 146, 145, 118, 117, 91, 57.

HRMS:630.2379

EXAMPLE 296 4-Trifluoromethyl-N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide (Formula RRR-11, R₁ is t-butyl; R₂ is 4-trifluoromethyl; Refer to Chart RRR).

Following the procedure of Example 294 but substituting the product of Preparation 128 for the product of Preparation 129 gives 4-trifluoromethyl-N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide as an amorphous foam.

Physical characteristics are as follows:

MS(EI): 633, 632, 631, 614, 613, 346, 201, 146, 91, 57.

HRMS:631.2444

EXAMPLE 297 4-Trifluoromethyl-N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide (Formula RRR-12; R₁ is t-butyl; R₂ is 4-trifluoromethyl; Refer to Chart RRR).

Following the procedure of Example 295 but substituting the product of Preparation 128 for the product of Preparation 129 gives 4-trifluoromethyl-N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide as an amorphous foam.

Physical characteristics are as follows:

MS(EI): 633, 632, 631, 614, 613, 346, 201, 146, 91, 57.

HRMS:631.2450

EXAMPLE 298 5-Trifluoromethyl-N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide (Formula RRR-11; R₁ is ethyl; R₂ is 5-trifluoromethyl; Refer to Chart RRR).

Following the procedure of Example 294 but substituting the product of Preparation 147A gives 5-trifluoromethyl-N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-propyl]phenyl]-2-pyridinesulfonamide as an amorphous foam.

Physical characteristics are as follows:

MS(EI): 605, 604, 603, 602, 586, 585, 393, 201, 133, 91

HRMS: 603.2153

EXAMPLE 298A 5-Trifluoromethyl-N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-
6(R or S)-(2-phenethyl)-6(R or S)-n-propyl-2H-pyran-3-yl]-
5 propyl]-phenyl]-2-pyridinesulfonamide (Formula RRR-12; R₁ is
ethyl; R₂ is 5-trifluoromethyl; Refer to Chart RRR).

Following the procedure of Example 294 but substituting the amine derived
from Isomer 1 of Preparation 147 (derived following the procedure of Preparation
147A) gives the title compound as an amorphous foam.

10 Physical characteristics are as follows:

¹H NMR (CD₃OD): δ 8.9, 8.2, 8.0, 7.0-7.3, 3.9, 2.4-2.7, 1.2-2.2, 0.8-1.0.

TLC (silica gel GF): R_f=0.19, 40% ethyl acetate in hexane.

PREPARATION 139 (5-Nitro-pyridin-2-yl)-isothioureia hydrochloride (Formula
UUU-2) Refer to Chart UUU.

15 A solution of 3.81 g of thiourea in 75 mL of hot absolute ethanol is treated
with 7.61 g of 2-chloro-5-nitropyridine (Formula UUU-1) and is heated at reflux for 6
hours. The mixture is then cooled to 0°C and the precipitated solid is collected. The
solid is washed sequentially with cold absolute ethanol and chloroform. The solid is
dried *in vacuo* to afford 6.91 g of the title product as a light brown solid.

20 Physical characteristics are as follows:

MP 175 °C (dec.)

¹H NMR (CD₃OD) δ 7.9, 8.6, 9.4 ppm

PREPARATION 140 5-Nitro-2-thiopyridine (Formula UUU-3) Refer to Chart
UUU.

25 A solution of 1.65 g of sodium carbonate in 50 mL of water is treated with
2.35 g of the title compound of Preparation 139. The mixture is charged with a
solution of 2.75 g of sodium hydroxide in 50 mL of water and the resulting mixture
is warmed to room temperature. After stirring for 1 hour, the mixture is heated to
95°C for 1 hour and finally cooled to room temperature. The aqueous mixture is
30 extracted with two portions of diethyl ether and then carefully acidified with 6 N
aqueous hydrochloric acid. The orange precipitated solid is collected and washed
sequentially with cold dilute aqueous hydrochloric acid and water. The solid is dried
in vacuo to afford 1.27 g of the title product as an orange solid.

Physical characteristics are as follows:

35 MP 167-170 °C

¹H NMR (CDCl₃-CD₃OD) δ 7.4, 7.9, 8.5 ppm

PREPARATION 141 5-Nitro-2-pyridinesulfonyl chloride (Formula UUU-4)

Refer to Chart UUU.

To a suspension of 1.27 g of the title compound of Preparation 140 in 25 mL of 1 N aqueous hydrochloric acid and 5 mL of acetic acid at 0°C is vigorously bubbled in chlorine gas. After 15 minutes, the chlorine gas addition is ceased and replaced with nitrogen gas. The resulting solid is collected and washed sequentially with cold dilute aqueous hydrochloric acid and water. The solid is dried *in vacuo* to afford 1.60 g of the title product as a tan solid.

Physical characteristics are as follows:

MP 77-80 °C

¹H NMR (CDCl₃) δ 8.3, 8.8, 9.6 ppm

PREPARATION 142 N-[3-(1-[5,6-Dihydro-4-hydroxy-2-oxo-6-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-5-nitro-2-pyridinesulfonamide (Formula UUU-5: R₁ is 2-phenylethyl, R₂ is propyl, R₃ is *tert*-butyl) Refer to Chart UUU.

To a solution of 210 mg of the title compound of Preparation 81 (Formula T-4) in 2 mL of dichloromethane at 0°C is added 80 µL of pyridine followed by 111 mg of the title compound of Preparation 141 (Formula UUU-4). After warming to room temperature overnight, the reaction mixture is column chromatographed on flash silica gel eluting with 3% to 9% ethyl acetate in dichloromethane to provide 303 mg of the title compound as a yellow foam.

Physical characteristics are as follows:

¹H NMR (CDCl₃-CD₃OD) δ 0.8-1.0, 1.2-1.4, 1.6-1.9, 2.4-2.7, 4.0, 6.9-7.4, 8.0, 8.5, 9.4 ppm

HRMS 608.2412 (EI)

EXAMPLE 299 5-Amino-N-[3-(1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide (Formula UUU-6: R₁ is 2-phenylethyl, R₂ is propyl, R₃ is *tert*-butyl) Refer to Chart UUU.

To a solution of 300 mg of the title compound of Preparation 142 (Formula UUU-5) in 5 mL of methanol under argon is added 500 mg of ammonium formate followed by 100 mg of 10% palladium on carbon. After 1 hour, the reaction mixture is filtered through a pad of Celite with methanol washes. The combined filtrates are

concentrated under reduced pressure and the residue is repeatedly triturated with portions of dichloromethane. The combined dichloromethane washes are concentrated under reduced pressure and the residue is column chromatographed on flash silica gel eluting with 50% ethyl acetate in dichloromethane to provide 246 mg of the title compound as a white solid.

Physical characteristics are as follows:

^1H NMR ($\text{CDCl}_3\text{-CD}_3\text{OD}$) δ 0.8-1.0, 1.2-1.4, 1.5-2.0, 2.4-2.6, 4.0, 6.7, 6.8-7.3, 7.4, 7.9 ppm

HRMS 577.2617 (EI)

10 EXAMPLE 300-327

Following the procedures and preparations described above and using starting materials known and available to one of ordinary skill in organic synthesis, the following additional compounds in Table 3 of the present invention are made from the compounds prepared in the following preparations:

- 15 PREPARATION 143 Preparative separation of N-[3-[1-[5,6-Dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl-carbamic acid phenylmethyl ester, to give 4 isomers.

The title compound of Preparation 134 is separated into four constituent stereoisomers which are (in order of elution from system A) 4 isomers: Isomer 1, Isomer 2, Isomer 3, and Isomer 4 with the following approximate observed retention times 10.5, 14.9, 21.4 and 65.2 minutes respectively. System A consists of a 0.46x25 cm Chiralcel OD-H column eluting at 0.5 mL/min with 20% isopropanol and 0.1 % trifluoroacetic acid in hexane (V/V). (Chiralcel OD-H is a registered trademark of Chiral Technologies, Inc., Exton PA 19341.)

The first phase of the separation is accomplished with a 2.1x25 cm (R,R)Whelk-O 1 column eluting with 20% (V/V) isopropanol in hexane at 12 mL/min. ((R,R)Whelk-O 1 is a registered trademark of Regis Technologies, Inc., Morton Grove, IL 60053.) The peaks eluting at approximately 35 and 41 min are, respectively, a mixture of Isomers 3 and 4 and a mixture of Isomers 1 and 2 as judged from System A, above. The two mixtures are further treated as below.

In the second phase of the separation, the mixture from above that elutes near 41 minutes is injected onto a 2.1x25 cm Chiralcel OD column (Chiral Technologies, Inc.) and eluting with 15% isopropanol and 0.05% trifluoroacetic acid in hexane (V/V) at 9.0 mL/min. The peaks that elute near 11.0 and 22.0 minutes are,

respectively, Isomer 1 and Isomer 2 as judged from System A.

In the final phase of the separation, the mixture that elutes from the (R,R)Whelk-O 1 column near 35 minutes is injected onto a 2.2x25 cm Chiralcel OD column eluting with 35% isopropanol and 0.1% trifluoroacetic acid (V/V) in hexane at 9.0 mL/min. The isomer that elutes near 9.7 minutes is Isomer 3 and the one that elutes near 16.6 minutes is Isomer 4.

In both phases of the separation of stereoisomers, fractions are pooled after assay with System A and pools are concentrated to dryness on a rotary evaporator.

PREPARATION 144 Resolution of N-[3-(1-[5,6-Dihydro-6,6-dipropyl-4-hydroxy-2-oxo-2H-pyran-3-yl]-2,2-dimethylpropyl)-phenyl]carbamic acid, phenylmethyl ester to give 2 isomers.

Samples of the starting compound (up to 1.0 gm each run) are injected onto a 5.1x50 cm Chiralcel OD column (Chiral Technologies, Inc.). The enantiomers elute at about 23 min (This corresponds to the benzyloxycarbonyl protected analogue of amine (Isomer 1) (EI-MS: 359 [M⁺]; ¹H NMR (CDCl₃-CD₃OD): 7.1-6.9, 6.5, 4.2, 2.6-2.3, 1.8-1.2, 1.1, 0.9; TLC: R_f=0.42 (10% ethyl acetate in dichloromethane)), and at about 33 min (This corresponds to the benzyloxycarbonyl protected analogue of amine (Isomer 2) (EI-MS: 359 [M⁺]; ¹H NMR (CDCl₃-CD₃OD): 7.1-6.9, 6.5, 4.2, 2.6-2.3, 1.8-1.2, 1.1, 0.9; TLC: R_f=0.42 (10% ethyl acetate in dichloromethane)). The mobile phase is 20% isopropanol and 0.1% acetic acid in hexane (V/V) pumped at 60 mL/min. The purity is checked on a 0.46x25 cm Chiralcel OD-H column (Chiral Technologies, Inc.). The mobile phase is 20% isopropanol in hexane (V/V) and 0.05% trifluoroacetic acid pumped at 0.5 mL/min. The observed retention times are 8.9 and 16.7 min (monitor set at 238 nm) for Isomer 1 and Isomer 2, respectively.

PREPARATION 145 Resolution of N-[3-(1-[6,6-Bis(2-phenylethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-yl]propyl)phenyl]carbamic acid, phenylmethyl ester to give 2 isomers.

Samples of the starting compound (up to 1.3 gm each run) are injected onto a 5.1x50 cm Chiralcel OD column (Chiral Technologies, Inc.). The enantiomers are eluted with 20% isopropanol and 0.025% acetic acid in hexane (V/V) at 60 mL/min until the first enantiomer elutes. At this point (approximately 120 min into the run) the flow rate is increased to 90 mL/min to expedite elution of the second enantiomer. The enantiomers elute near 91.2 min (This is the corresponding benzyloxycarbonyl analogue of amine Isomer 1 and near 132 min (This is the corresponding

2187523

benzyloxycarbonyl analogue of amine Isomer 2. The purity is checked on a 0.46x25 cm Chiralcel OD-H column. The mobile phase is 30% isopropanol in hexane (V/V) pumped at 0.5 mL/min.

PREPARATION 146 5-Carbamoylpyridine-2-sulfonyl chloride (Formula VVV-2) Refer to Chart VVV.

Into a cold (0°), stirred suspension of 400 mg of 2-mercapto-5-carbamoylpyridine of formula VVV-1 in 7.5 ml of 1N HCl is passed a brisk stream of chlorine gas. After ten minutes, the suspension is filtered, and the solid washed well with water and dried *in vacuo*. Obtained is 517 mg of the title compound as a nearly white solid.

EXAMPLE 328 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-4-cyanobenzenesulfonamide (Formula U-8: R₁ is *tert*-butyl, R₂ is 4-cyanophenyl) Refer to Chart U.

Using the general sulfonation procedure of Example 252, 88 mg of the amine of Preparation 86 (Formula U-7, R₁ is *tert*-butyl) is reacted with 4-cyanobenzenesulfonyl chloride. Flash chromatography on silica gel using 10% ethyl acetate in dichloromethane provides 117 mg of the title compound as an amorphous white solid.

Physical characteristics are as follows:

¹H NMR δ 0.90, 1.3, 1.7, 2.5, 3.6, 6.8-7.4, 7.6, 7.8 ppm

HRMS: 605.2478

R_f 0.36 (10% ethyl acetate in dichloromethane)

EXAMPLE 329 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-8-quinolinesulfonamide (Formula U-8: R₁ is *tert*-butyl, R₂ is 8-quinolyl) Refer to Chart U.

Using the general sulfonylation procedure of Example 252, 88 mg of the amine of Preparation 86 (Formula U-7, R₁ is *tert*-butyl) is reacted with 8-quinolinesulfonyl chloride. Flash chromatography on silica gel using 5-10% ethyl acetate in dichloromethane provides 101 mg of the title compound as an amorphous white solid.

Physical characteristics are as follows:

¹H NMR δ 0.63, 0.9, 1.1, 1.3, 1.6-1.9, 2.4-2.6, 6.7-7.6, 8.0, 8.2, 9.1 ppm

HRMS: 631.2638

R_f 0.30 (5% ethyl acetate in dichloromethane)

EXAMPLE 330 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide
(Formula D-6: R₁ is phenethyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 1-methylimidazole-4-yl) Refer to Chart D.

Using the general sulfonylation procedure of Example 252, 77 mg of the amine of Formula D-5, wherein R₁ and R₂ are phenethyl and R₃ is ethyl, is reacted with 1-methylimidazole-4-sulfonyl chloride. Flash chromatography on silica gel using 3% methanol in dichloromethane provides 97.0 mg of the title compound as a crystalline white solid.

Physical characteristics are as follows:

¹H NMR δ 0.88, 1.9-2.2, 2.6, 3.6-3.8, 3.97, 6.9-7.5 ppm

HRMS: 600.2521

R_f 0.31 (5% methanol in dichloromethane)

EXAMPLE 331 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide
(Formula D-6: R₁ is phenethyl, R₂ is phenethyl, R₃ is ethyl, R₄ is 5-cyanopyridine-2-yl) Refer to Chart D.

Using the general sulfonylation procedure of Example 252, 77 mg of the amine of Formula D-5, wherein R₁ and R₂ phenethyl and R₃ is ethyl, is reacted with 5-cyanopyridine-2-sulfonyl chloride. Flash chromatography on silica gel using 10% ethyl acetate in dichloromethane provides 88.3 mg of the title compound as a crystalline white solid.

Physical characteristics are as follows:

¹H NMR δ 0.85, 1.8-2.2, 2.5-2.7, 3.97, 6.9-7.4, 7.9, 8.8 ppm.

HRMS: 622.2355

R_f 0.28 (10% ethyl acetate in dichloromethane)

EXAMPLE 332 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-carbamoylpyridine-2-sulfonamide (Formula U-8: R₁ is ethyl, R₂ is 5-carbamoylpyridine-2-yl) Refer to Chart U.

Using the general sulfonylation procedure of Example 252, 82 mg of the amine of Formula U-7, wherein R₁ is ethyl, is reacted with 5-carbamoylpyridine-2-sulfonyl chloride of Preparation 146. Flash chromatography on silica gel using 3-6% methanol in dichloromethane provides 55.4 mg of the title compound as an

2187523

amorphous solid.

Physical characteristics are as follows:

^1H NMR δ 0.7-0.9, 1.3, 1.6-2.1, 2.5, 3.9, 6.8-7.3, 7.8, 8.2 ppm.

HRMS: 596.2216

5 R_f 0.16 (5% methanol in dichloromethane)

EXAMPLE 333 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-(4-fluorophenyl)ethyl)-2H-pyran-3-yl)propyl}phenyl]-5-carbamoylpyridine-2-sulfonamide (Formula V-8: R_1 is ethyl, R_2 is 5-carbamoylpyridine-2-yl) Refer to Chart V.

10 Using the general sulfonylation procedure of Example 252, 98 mg of the amine of formula V-7, wherein R_1 is ethyl) is reacted with 5-carbamoylpyridine-2-sulfonyl chloride of Preparation 146. Flash chromatography on silica using 3-6% methanol in dichloromethane provides 58.3 mg of the title compound as an amorphous solid.

15 Physical characteristics are as follows:

^1H NMR δ 0.83, 1.8-2.2, 2.5-2.6, 6.8-7.2, 7.8, 8.1, 9.0 ppm.

HRMS: 676.2297

R_f 0.17 (5% methanol in dichloromethane)

20 EXAMPLE 334 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)propyl}phenyl]-5-carbamoylpyridine-2-sulfonamide (Formula D-6: R_1 is propyl, R_2 is propyl, R_3 is ethyl, R_4 is 5-carbamoylpyridine-2-yl) Refer to Chart D.

25 Using the general sulfonylation procedure of Example 252, 66 mg of the amine of Formula D-5 (R_1 and R_2 are propyl, R_3 is ethyl) is coupled with 5-carbamoylpyridine-2-sulfonyl chloride of Preparation 146 to yield, after flash chromatography on silica gel using 3-6% methanol in dichloromethane, 83.8 mg of the title compound as an amorphous white solid.

Physical characteristics are as follows:

^1H NMR δ 0.7-0.9, 1.2-2.1, 3.87, 7.0-7.3, 7.8, 8.2 ppm.

30 HRMS: 516.2156

R_f 0.22 (5% methanol in dichloromethane)

PREPARATION 147 Resolution of N-[3-[1-(4-Hydroxy-5,6-dihydro-2-oxo-6-phenethyl-6-propyl-2H-pyran-3-yl)propyl]phenyl]-carbamic acid, phenylmethyl ester to give 4 isomers
35 (Formula WWW-2: R_1 is phenethyl, R_2 is propyl, R_3 is

ethyl) Refer to Chart WWW and RRR.

The four isomers of the product of Preparation 130A (Formula RRR-1; R_1 = ethyl) are (in order of increasing retention time on System B): (ca. 16.9 min) (Isomer 1), (ca. 28.0 min) (Isomer 2), (ca. 38.2 min) (Isomer 3) and (ca. 49.8 min) (Isomer 4).

- 5 System B consists of a 0.46x25 cm Chiralcel OD-H column (Chiral Technologies, Inc.) eluting with 25% isopropanol in hexane (V/V) at 0.5 mL/min.

In Phase one of the complete resolution repeatedly inject 55 mg samples of the product of Preparation 130A onto a 2.1x25 cm (R,R)Whelk-O 1 column (Regis Technologies, Inc.). Elute the isomers at 10 mL/min with 35% isopropanol and 0.5% acetic acid in hexane (V/V). The first of the three peaks to elute (near 12 min) is a mixture of Isomers 1 (Formula RRR-4 of Chart RRR) and 2 (Formula RRR-3 of Chart RRR) as shown by injecting aliquots in System B. Resolve this mixture in Phase 2, below.

- The second phase consists of a 2.1x25 cm Chiralcel OD column kept at 30°. Inject 60 mg batches of the mixture obtained in the first phase and elute the enantiomers with 25% isopropanol and 0.05% trifluoroacetic acid (V/V) at 9 mL/min. Separately pooled and concentrated, the fractions eluting near 14.5 and 23.9 min to give Isomers 1 (Formula RRR-4 where R_1 is ethyl of Chart RRR) and 2 (formula RRR-3 where R_1 is ethyl of Chart RRR) respectively.

- 20 PREPARATION 147A 3-(R or S)-[1-(3-aminophenyl)-propyl]-4-hydroxy-5,6-dihydro-6-(R or S)-phenethyl-6-(R or S)-propyl-2H-pyran-2-one (Formula RRR-7; R_1 is ethyl; Refer to Chart RRR).

Following the procedure of Preparation 132 beginning with the peak identified as peak 2 (Formula RRR-3; R_1 is ethyl of Chart RRR) from the chiral resolution of the product of Preparation 147, the title compound is prepared.

Physical characteristics are as follows:

^1H NMR (CD_3OD): δ 6.5-7.3, 3.9-4.0, 2.5-2.7, 1.2-2.3, 0.8-1.0.

TLC (silica gel GF): R_f =0.31, 40% ethyl acetate in hexane.

- 30 EXAMPLE 335 N-[3-{1(R or S)-(4-Hydroxy-5,6-dihydro-2-oxo-6(R or S)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula WWW-4: R_1 is phenethyl, R_2 is propyl, R_3 is ethyl, R_4 is 5-cyanopyridine-2-yl) [Isomer 1] Refer to Chart WWW.

The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R_1 is phenethyl, R_2 is

propyl, and R₃ is ethyl. The amine is derived from the first stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 147. The title compound is obtained as an amorphous solid after flash chromatography on silica using 10% ethyl acetate in dichloromethane.

5 Physical characteristics are as follows:

¹H NMR δ 0.7-1.0, 1.2-2.6, 3.3-3.6, 6.9-7.3, 7.7-8.2, 8.8-9.0 ppm.

HRMS: 560.2210

R_f 0.41 (15% ethyl acetate in dichloromethane)

10 EXAMPLE 336 N-[3-{1(*R or S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S or R*)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula WWW-4: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 5-cyanopyridine-2-yl) [Isomer 2] Refer to Chart WWW.

15 The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine that is the title product of Preparation 147A (Formula WWW-3, where R₁ is phenethyl, R₂ is propyl, and R₃ is ethyl). The amine is derived from the second stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 147A. The title compound is obtained as an amorphous solid after flash chromatography on silica using 10% ethyl acetate in dichloromethane.

Physical characteristics are as follows:

¹H NMR δ 0.6-2.6, 3.3-3.6, 6.9-7.3, 7.7-8.2, 8.8-9.0 ppm.

HRMS: 560.2215

R_f 0.41 (15% ethyl acetate in dichloromethane)

25 EXAMPLE 337 N-[3-{1(*S or R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R or S*)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula WWW-4: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 5-cyanopyridine-2-yl) [Isomer 3] Refer to Chart WWW.

30 The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R₁ is phenethyl, R₂ is propyl, and R₃ is ethyl. The amine is derived from the third stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 147. The title compound is obtained as an amorphous solid after flash chromatography on silica using 10% ethyl acetate in dichloromethane.

2187523

Physical characteristics are as follows:

^1H NMR δ 0.6-2.6, 3.3-3.6, 6.9-7.3, 7.7-8.2, 8.8-9.0 ppm.

HRMS: 560.2210

R_f 0.41 (15% ethyl acetate in dichloromethane)

- 5 **EXAMPLE 338** N-[3-{1(*S* or *R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S* or *R*)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula WWW-4: R_1 is phenethyl, R_2 is propyl, R_3 is ethyl, R_4 is 5-cyanopyridine-2-yl) [Isomer 4]
Refer to Chart WWW.

- 10 The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R_1 is phenethyl, R_2 is propyl, and R_3 is ethyl. The amine is derived from the fourth stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 147. The title compound is obtained as an amorphous solid after flash
15 chromatography on silica using 10% ethyl acetate in dichloromethane.

Physical characteristics are as follows:

^1H NMR δ 0.7-1.0, 1.2-2.6, 3.3-3.6, 6.9-7.3, 7.7-8.2, 8.8-9.0 ppm.

HRMS: 560.2210

R_f 0.41 (15% ethyl acetate in dichloromethane)

- 20 **EXAMPLE 339** N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R* or *S*)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula WWW-4: R_1 is phenethyl, R_2 is propyl, R_3 is ethyl, R_4 is 1-methylimidazol-4-yl) [Isomer 1]
Refer to Chart WWW.

- 25 The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R_1 is phenethyl, R_2 is propyl, and R_3 is ethyl. The amine is derived from the first stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 147. The title compound is obtained as an amorphous solid after flash chromatography on
30 silica using 3% methanol in dichloromethane.

Physical characteristics are as follows:

^1H NMR δ 0.7-2.8, 3.2-3.7, 3.9, 7.0-7.6 ppm.

HRMS: 537.2317

R_f 0.36 (5% methanol in dichloromethane)

- 35 **EXAMPLE 340** N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S* or *R*)-

phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula WWW-4: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 1-methylimidazol-4-yl) [Isomer 2]
Refer to Chart WWW.

- 5 The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R₁ is phenethyl, R₂ is propyl, and R₃ is ethyl. The amine is derived from the second stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 147. The title compound is obtained as an amorphous solid after flash
10 chromatography on silica using 3-4% methanol in dichloromethane.

Physical characteristics are as follows:

¹H NMR δ 0.7-2.7, 3.3-3.7, 4.0, 7.0-7.5 ppm.

HRMS: 537.2275

R_f 0.36 (5% methanol in dichloromethane)

- 15 EXAMPLE 341 N-[3-{1(*S or R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R or S*)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula WWW-4: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 1-methylimidazol-4-yl) [Isomer 3]
Refer to Chart WWW.

- 20 The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R₁ is phenethyl, R₂ is propyl, and R₃ is ethyl. The amine is derived from the third stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 147. The title compound is obtained as an amorphous solid after flash
25 chromatography on silica using 3% methanol in dichloromethane.

Physical characteristics are as follows:

¹H NMR δ 0.7-2.7, 3.3-3.7, 4.0, 7.0-7.5 ppm.

HRMS: 537.2329

R_f 0.36 (5% methanol in dichloromethane)

- 30 EXAMPLE 342 N-[3-{1(*S or R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S or R*)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula WWW-4: R₁ is phenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 1-methylimidazol-4-yl) [Isomer 4]
Refer to Chart WWW.

- 35 The title compound is prepared using the general sulfonylation procedure of

Example 252, using the amine of Formula WWW-3, where R₁ is phenethyl, R₂ is propyl, and R₃ is ethyl. The amine is derived from the fourth stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 147. The title compound is obtained as an amorphous solid after flash

5 chromatography on silica using 3% methanol in dichloromethane.

Physical characteristics are as follows:

¹H NMR δ 0.7-2.8, 3.2-3.7, 3.9, 7.0-7.6 ppm.

HRMS: 537.2312

R_f 0.36 (5% methanol in dichloromethane)

10 PREPARATION 148 3-[(3-Nitrophenyl)methyl]-6,6-diphenethyl-4-hydroxy-5,6-dihydro-2H-pyran-2-one (Formula XXX-3) Refer to Chart XXX.

To a solution of 172 mg of 6,6-Diphenethyl-4-hydroxy-5,6-dihydro-2H-pyran-2-one of formula XXX-1 and 81 mg of *meta*-nitrobenzaldehyde in 2 ml of dry THF, under argon, is added a solution of 142 mg of AlCl₃ in 1 ml of THF. The solution is stirred at room temperature for 2 hours, then quenched with 310 mg of sodium carbonate decahydrate, diluted with ether, and filtered through Celite with ether rinses. Following removal of solvent under reduced pressure, 264 mg of crude benzylidene of Formula XXX-2 is obtained. This material is dissolved in 5 ml of methanol, and the solution cooled to 0° for the addition of 44 mg of sodium cyanoborohydride. After an hour, a further 20 mg aliquot of sodium cyanoborohydride is added. After another 30 minutes, the mixture is acidified with dilute HCl to pH 1 and extracted with three portions of dichloromethane. The extract is dried (MgSO₄) and then concentrated under reduced pressure. Flash chromatography of the residue on silica using 5-20% ethyl acetate in dichloromethane provides 211 mg of the title compound as an amorphous white solid.

Physical characteristics are as follows:

¹H NMR δ 2.0, 2.7, 3.8, 7.0-7.4, 7.6, 8.0, 8.2 ppm.

30 MS: M+ 457

R_f 0.25 (5% ethyl acetate in dichloromethane)

PREPARATION 149 3-[(3-Aminophenyl)methyl]-6,6-diphenethyl-4-hydroxy-5,6-dihydro-2H-pyran-2-one (Formula XXX-4) Refer to Chart XXX.

35 A mixture of 211 mg of the product of Preparation 148 (Formula XXX-3) and

50 mg of 10% palladium on carbon in 5 ml of methanol is stirred at room temperature under 1 atmosphere hydrogen gas. After two hours, the mixture is filtered through Celite and concentrated under reduced pressure. Flash chromatography of the residue on silica using 25% ethyl acetate in dichloromethane affords 133.6 mg of the title compound.

Physical characteristics are as follows:

^1H NMR δ 2.0, 2.6, 3.6, 4.1, 6.5, 6.6, 6.7, 6.9-7.3 ppm.

MS: M^+ 427

R_f 0.33 (25% ethyl acetate in dichloromethane)

10 EXAMPLE 343 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula YYY-5) R_1 and R_2 are phenethyl, R_3 is 1-methylimidazole-4-yl) Refer to Chart YYY.

Using the general sulfonylation procedure of Example 252, 77 mg of the amine of formula YYY-4 (R_1 and R_2 are phenethyl) is reacted with 1-methylimidazole-4-sulfonyl chloride. Flash chromatography on silica using 3% methanol in dichloromethane provides 90.7 mg of the title compound as an amorphous white solid.

Physical characteristics are as follows:

20 ^1H NMR δ 0.96, 1.0, 1.6-2.7, 3.45, 6.8-7.5 ppm.

HRMS: 628.2832

R_f 0.38 (3% methanol in dichloromethane)

EXAMPLE 344 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula YYY-5) R_1 and R_2 are phenethyl, R_3 is 5-cyanopyridine-2-yl) Refer to Chart YYY

Using the general sulfonylation procedure of Example 252, 77 mg of the amine of Formula YYY-4 (R_1 and R_2 are phenethyl) is reacted with 5-cyanopyridine-2-sulfonyl chloride. Flash chromatography on silica using 10% ethyl acetate in dichloromethane provides 86.1 mg of the title compound as an amorphous white solid.

Physical characteristics are as follows:

^1H NMR δ 0.96, 1.8-2.2, 2.5-2.8, 4.1, 4.3, 6.9-7.4, 7.9-8.0, 8.9 ppm.

HRMS: 650.2681

35 R_f 0.27 (10% ethyl acetate in dichloromethane)

PREPARATION 150

Resolution of N-[3-[1-(4-Hydroxy-5,6-dihydro-2-oxo-6-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl]phenyl]carbamic acid, phenylmethyl ester to give 4 isomers (Formula WWW-2: R₁ is 4-fluorophenethyl, R₂ is propyl, R₃ is *t*-butyl) Refer to Chart WWW.

System C is used to track the enantiomers and to monitor the preparative columns. System C consists of a 0.46x25 cm Chiralcel OD-H column (Chiral Technologies, Inc.) with 15% isopropanol in hexane (V/V) at 0.5 mL/min. The peaks eluting near 13.5, 18.8, 37.1 and 79.7 min are, respectively, Isomer 1, Isomer 2, Isomer 3, and Isomer 4.

Separate Isomers 3 and 4 from the mixture on a 2.1x25 cm (R,R)Whelk-O 1 column (Regis Technologies, Inc.). These two isomers elute at about 23.9 and 26.8 min when the column is developed with 20% isopropanol in hexane (V/V) at 10 mL/min at 30°. The desired isomers elute as an unresolved mixture near 28.9 min and are separated in the second stage of the resolution.

For the second stage inject the unresolved mixture onto a 2.1x25 cm Chiralcel OD column (Chiral technologies, Inc.) kept at 30°. With 12% isopropanol in hexane (V/V) at 12 mL/min, Isomer 1 emerges near 14.5 min and Isomer 2 emerges near 20.8 min.

EXAMPLE 345

N-[3-{1(*R or S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R or S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl]phenyl]-5-cyanopyridine-2-sulfonamide (Formula WWW-4: R₁ is 4-fluorophenethyl, R₂ is propyl, R₃ is *tert*-butyl, R₄ is 5-cyanopyridine-2-yl) [Isomer 1] Refer to Chart WWW.

The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R₁ is 4-fluorophenethyl, R₂ is propyl, and R₃ is ethyl. The amine is derived from the first stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 150. The title compound is obtained as an amorphous solid after flash chromatography on silica using 10% ethyl acetate in dichloromethane.

Physical characteristics are as follows:

¹H NMR δ 0.7-2.7, 3.2, 3.5, 3.6, 3.7, 4.1, 6.8-7.4, 7.5, 7.8-8.2, 8.8 ppm.

HRMS: 606.2429

R_f 0.40 (15% ethyl acetate in dichloromethane)

EXAMPLE 346 N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S* or *R*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide
(Formula WWW-4: R_1 is 4-fluorophenethyl, R_2 is propyl, R_3 is *tert*-butyl, R_4 is 5-cyanopyridine-2-yl) [Isomer 2] Refer to Chart WWW.

The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R_1 is 4-fluorophenethyl, R_2 is propyl, and R_3 is ethyl. The amine is derived from the second stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 150. The title compound is obtained as an amorphous solid after flash chromatography on silica using 10% ethyl acetate in dichloromethane.

Physical characteristics are as follows:

^1H NMR δ 0.6, 0.7-2.6, 3.4, 3.5, 3.7, 4.2, 6.8-7.3, 7.5, 7.8-8.2, 8.8-9.0 ppm.

MS: 606.2434

R_f 0.40 (15% ethyl acetate in dichloromethane)

EXAMPLE 347 N-[3-{1(*S* or *R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R* or *S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula WWW-4: R_1 is 4-fluorophenethyl, R_2 is propyl, R_3 is *tert*-butyl, R_4 is 5-cyanopyridine-2-yl) [Isomer 3] Refer to Chart WWW.

The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R_1 is 4-fluorophenethyl, R_2 is propyl, and R_3 is ethyl. The amine is derived from the third stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 150. The title compound is obtained as an amorphous solid after flash chromatography on silica using 10% ethyl acetate in dichloromethane.

Physical characteristics are as follows:

^1H NMR δ 0.6, 0.7-2.6, 3.4, 3.5, 3.7, 4.2, 6.8-7.3, 7.5, 7.8-8.2, 8.8-9.0 ppm.

MS: 606.2423

R_f 0.40 (15% ethyl acetate in dichloromethane)

EXAMPLE 348 N-[3-{1(*S* or *R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S* or *R*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula WWW-4: R_1 is 4-fluorophenethyl, R_2 is propyl, R_3 is *tert*-butyl, R_4 is 5-

cyanopyridine-2-yl) [Isomer 4] Refer to Chart WWW.

The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R₁ is 4-fluorophenethyl, R₂ is propyl, and R₃ is ethyl. The amine is derived from the fourth stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 150. The title compound is obtained as an amorphous solid after flash chromatography on silica using 10% ethyl acetate in dichloromethane.

Physical characteristics are as follows:

¹H NMR δ 0.7-2.7, 3.2, 3.5, 3.6, 3.7, 4.1, 6.8-7.4, 7.5, 7.8-8.2, 8.8 ppm.

HRMS: 606.2429

R_f 0.40 (15% ethyl acetate in dichloromethane)

EXAMPLE 349 N-[3-{1(*R or S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R or S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula WWW-4: R₁ is 4-fluorophenethyl, R₂ is propyl, R₃ is *tert*-butyl, R₄ is 1-methylimidazol-4-yl [Isomer 1] Refer to Chart WWW.

The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R₁ is 4-fluorophenethyl, R₂ is propyl, and R₃ is ethyl. The amine is derived from the first stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 150. The title compound is obtained as an amorphous solid after flash chromatography on silica using 3% methanol in dichloromethane.

Physical characteristics are as follows:

¹H NMR δ 0.8-1.0, 1.4, 1.7, 2.3-2.7, 3.6, 3.9, 4.1, 6.8-7.5 ppm.

HRMS: 584.2585

R_f 0.34 (5% methanol in dichloromethane)

EXAMPLE 350 N-[3-{1(*R or S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S or R*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula WWW-4: R₁ is 4-fluorophenethyl, R₂ is propyl, R₃ is *tert*-butyl, R₄ is 1-methylimidazol-4-yl [Isomer 2] Refer to Chart WWW.

The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R₁ is 4-fluorophenethyl, R₂ is propyl, and R₃ is ethyl. The amine is derived from the second stereoisomer of Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation

150. The title compound is obtained as an amorphous solid after flash chromatography on silica using 3% methanol in dichloromethane.

Physical characteristics are as follows:

^1H NMR δ 0.7-1.1, 1.3, 1.7, 2.3-2.7, 3.6, 3.9, 4.1, 6.8-7.5 ppm.

5 HRMS: 584.2585

R_f 0.34 (5% methanol in dichloromethane)

EXAMPLE 351 N-[3-{1(*S* or *R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R* or *S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethyl-propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula
10 WWW-4: R_1 is 4-fluorophenethyl, R_2 is propyl, R_3 is *tert*-butyl, R_4 is 1-methylimidazol-4-yl [Isomer 3] Refer to Chart WWW.

The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R_1 is 4-fluorophenethyl, R_2 is propyl, and R_3 is ethyl. The amine is derived from the third stereoisomer of
15 Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 150. The title compound is obtained as an amorphous solid after flash chromatography on silica using 3% methanol in dichloromethane.

Physical characteristics are as follows:

^1H NMR δ 0.7-1.1, 1.3, 1.7, 2.3-2.7, 3.6, 3.9, 4.1, 6.8-7.5 ppm.

20 HRMS: 584.2591

R_f 0.34 (5% methanol in dichloromethane)

EXAMPLE 352 N-[3-{1(*S* or *R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S* or *R*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethyl-propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula
25 WWW-4: R_1 is 4-fluorophenethyl, R_2 is propyl, R_3 is *tert*-butyl, R_4 is 1-methylimidazol-4-yl [Isomer 4] Refer to Chart WWW.

The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R_1 is 4-fluorophenethyl, R_2 is propyl, and R_3 is ethyl. The amine is derived from the fourth stereoisomer of
30 Formula WWW-2 to elute from a Chiralcel OD chiral HPLC column of Preparation 150. The title compound is obtained as an amorphous solid after flash chromatography on silica using 3% methanol in dichloromethane.

Physical characteristics are as follows:

^1H NMR δ 0.8-1.0, 1.4, 1.7, 2.3-2.7, 3.6, 3.9, 4.1, 6.8-7.5 ppm.

35 HRMS: 584.2580

2187523

R_f 0.34 (5% methanol in dichloromethane)

5 EXAMPLE 353 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)methyl}phenyl]-5-cyanopyridine-2-sulfonamide
(Formula XXX-5, R₁ is 5-cyanopyridine-2-yl) Refer to Chart XXX.

Using the general sulfonylation procedure of Example 252, 64 mg of the amine of formula XXX-4 is reacted with 5-cyanopyridine-2-sulfonyl chloride. Flash chromatography on silica using 2-3% methanol in dichloromethane provides 73.2 mg of the title compound as an amorphous white solid.

10 Physical characteristics are as follows:
 ¹H NMR δ 1.8-2.1, 2.6, 6.9-7.3, 7.9, 8.8 ppm.
 HRMS: 594.2068

R_f 0.40 (3% methanol in dichloromethane)

15 EXAMPLE 354 N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)methyl}phenyl]-5-aminopyridine-2-sulfonamide
(Formula UUU-6, R₁ and R₂ are phenethyl, R₃ is H). Refer to Chart UUU.

20 Using the general sulfonylation procedure of Example 252, 69 mg of the amine of formula XXX-4 is reacted with 5-nitropyridine-2-sulfonyl chloride. Flash chromatography on silica using 2-3% methanol in dichloromethane provides 107 mg of the intermediate nitro compound of formula UUU-5 (R₁ and R₂ are phenethyl, R₃ is H). Reduction to the amine is accomplished using hydrogen gas and palladium on carbon catalyst. Flash chromatography on silica gel using 4-6% methanol in dichloromethane provides 65.0 mg of the title compound as an amorphous white solid.

25 Physical characteristics are as follows:
 ¹H NMR δ 1.9-2.1, 2.6, 3.5-4.0, 6.7, 6.9-7.3, 7.5, 7.9 ppm.
 HRMS: 584.2215

R_f 0.24 (5% methanol in dichloromethane)

30 PREPARATION 151 Resolution of N-[3-{1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}-phenyl]carbamic acid, phenylmethyl ester to give 2 enantiomers (Formula WWW-2: R₁ and R₂ are phenethyl, R₃ is *t*-butyl) Refer to Chart WWW.

35 Inject 40 mg batches of the starting compound onto a 2.1x25 cm (R,R)Whelk-O 1 column (Regis Technologies, Inc.) that is maintained at 30°. The 2 enantiomers

2187523

elute at about 37 min (Enantiomer 1) and 43 min (Enantiomer 2) using 25% isopropanol and 0.05% acetic acid at 12 mL/min. Fractions are pooled on the basis of results from analysis on a 0.46x25 cm (R,R)Whelk-O 1 column eluted with 30% isopropanol and 0.1% acetic acid (V/V) at 1.0 mL/min. The isomers elute at (Isomer 1) 19.1 and (Isomer 2) 23.0 min respectively.

EXAMPLE 355 N-[3-{1(*R or S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-aminopyridine-2-sulfonamide (Formula WWW-4, R_1 and R_2 are phenethyl, R_3 is *tert*-butyl, R_4 is 5-aminopyridine-2-yl)
[Enantiomer 1] Refer to Chart WWW.

Using the general sulfonylation procedure of Example 252, 73 mg of the amine of Formula WWW-3 (R_1 and R_2 are phenethyl, R_3 is *tert*-butyl) is reacted with 5-nitropyridine-2-sulfonyl chloride. The amine used is derived from the first enantiomer of Formula WWW-2 to elute from an (R,R)Whelk-O chiral HPLC column of Preparation 151. Flash chromatography on silica using 5-10% ethyl acetate in dichloromethane provides 94.0 mg of the intermediate nitro compound of formula UUU-5 (R_1 and R_2 are phenethyl, R_3 is *tert*-butyl). Reduction to the amine is accomplished using hydrogen gas and palladium on carbon catalyst. Flash chromatography on silica gel using 4% methanol in dichloromethane provides 74.8 mg of the title compound as an amorphous white solid.

Physical characteristics are as follows:

^1H NMR δ 0.95, 2.0, 2.6, 6.8, 6.9-7.4, 7.5, 7.9 ppm.

HRMS: 640.2828

R_f 0.27 (5% methanol in dichloromethane)

EXAMPLE 356 N-[3-{1(*S or R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-aminopyridine-2-sulfonamide (Formula WWW-4, R_1 and R_2 are phenethyl, R_3 is *tert*-butyl, R_4 is 5-aminopyridine-2-yl)
[Enantiomer 2] Refer to Chart WWW.

Using the general sulfonylation procedure of Example 252, 73 mg of the amine of Formula WWW-3 (R_1 and R_2 are phenethyl, R_3 is *tert*-butyl) is reacted with 5-nitropyridine-2-sulfonyl chloride. The amine used is derived from the second enantiomer of Formula WWW-2 to elute from an (R,R)Whelk-O chiral HPLC column of Preparation 151. Flash chromatography on silica using 5-10% ethyl acetate in dichloromethane provides 91.3 mg of the intermediate nitro compound of formula

2187523

UUU-5 (R_1 and R_2 are phenethyl, R_3 is *tert*-butyl). Reduction to the amine is accomplished using hydrogen gas and palladium on carbon catalyst. Flash chromatography on silica gel using 4% methanol in dichloromethane provides 54.3 mg of the title compound as an amorphous white solid.

5 Physical characteristics are as follows:

^1H NMR δ 0.95, 2.0, 2.6, 6.8, 6.9-7.4, 7.5, 7.9 ppm.

HRMS: 640.2828

R_f 0.27 (5% methanol in dichloromethane)

10 EXAMPLE 357 N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula WWW-4, R_1 and R_2 are phenethyl, R_3 is *tert*-butyl, R_4 is 1-methylimidazol-4-yl) [Enantiomer 1] Refer to Chart WWW.

15 The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R_1 and R_2 are phenethyl and R_3 is *tert*-butyl. The amine is derived from the first enantiomer of Formula WWW-2 to elute from an (R,R)Whelk-O chiral HPLC column of Preparation 151. The title compound is obtained as an amorphous solid after flash chromatography on silica using 3% methanol in dichloromethane.

20 Physical characteristics are as follows:

^1H NMR δ 0.98, 2.0, 2.6, 3.6, 3.8, 6.9-7.5 ppm.

HRMS: 628.2832

R_f 0.38 (5% methanol in dichloromethane)

25 EXAMPLE 358 N-[3-{1(*S* or *R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula WWW-4, R_1 and R_2 are phenethyl, R_3 is *tert*-butyl, R_4 is 1-methylimidazol-4-yl) [Enantiomer 2] Refer to Chart WWW.

30 The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R_1 and R_2 are phenethyl and R_3 is *tert*-butyl. The amine is derived from the second enantiomer of Formula WWW-2 to elute from an (R,R)Whelk-O chiral HPLC column of Preparation 151. The title compound is obtained as an amorphous solid after flash chromatography on silica using 3% methanol in dichloromethane.

35 Physical characteristics are as follows:

2187523

¹H NMR δ 0.98, 2.0, 2.6, 3.6, 3.8, 6.9-7.5 ppm.

HRMS: 628.2838

R_f 0.38 (5% methanol in dichloromethane)

EXAMPLE 359 N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula WWW-4, R₁ and R₂ are phenethyl, R₃ is *tert*-butyl, R₄ is 5-cyanopyridine-2-yl)
[Enantiomer 1] Refer to Chart WWW.

The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R₁ and R₂ are phenethyl and R₃ is *tert*-butyl. The amine is derived from the first enantiomer of Formula WWW-2 to elute from an (R,R)Whelk-O chiral HPLC column of Preparation 151. The title compound is obtained as an amorphous solid after flash chromatography on silica using 10% ethyl acetate in dichloromethane.

Physical characteristics are as follows:

¹H NMR δ 0.87, 1.9, 2.6, 6.8-7.4, 7.9, 8.8 ppm.

HRMS: 650.2681

R_f 0.46 (15% ethyl acetate in dichloromethane)

EXAMPLE 360 N-[3-{1(*S* or *R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula WWW-4, R₁ and R₂ are phenethyl, R₃ is *tert*-butyl, R₄ is 5-cyanopyridine-2-yl)
[Enantiomer 2] Refer to Chart WWW.

The title compound is prepared using the general sulfonylation procedure of Example 252, using the amine of Formula WWW-3, where R₁ and R₂ are phenethyl and R₃ is *tert*-butyl. The amine is derived from the second enantiomer of Formula WWW-2 to elute from an (R,R)Whelk-O chiral HPLC column of Preparation 151. The title compound is obtained as an amorphous solid after flash chromatography on silica using 10% ethyl acetate in dichloromethane.

Physical characteristics are as follows:

¹H NMR δ 0.87, 1.9, 2.6, 6.8-7.4, 7.9, 8.8 ppm.

HRMS: 650.2681

R_f 0.46 (15% ethyl acetate in dichloromethane)

PREPARATION 152 Resolution of N-[3-[1-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-propyl]phenyl]carbamic acid,

2187523

phenylmethyl ester to give 2 isomers (Formula WWW-2:
R₁ and R₂ are propyl, R₃ is ethyl) Refer to Chart WWW.

Samples of the starting compound are injected onto a 2.1x25 cm Chiralcel OD column and eluted with 20% isopropanol (V/V) in hexane at 10 mL/min. The material eluting near 19.1 minutes is one isomer (Enantiomer 1) and that eluting near 37.7 minutes is another isomer (Enantiomer 2). The pools are concentrated separately on a rotary evaporator (ca. 30 mm, bath at 50° maximum) to give white solids.

EXAMPLE 361 N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-propyl}phenyl]-5-cyanopyridine-2-sulfonamide
(Formula WWW-4, R₁ and R₂ are propyl, R₃ is ethyl, R₄ is 5-cyanopyridine-2-yl) [Enantiomer 1] Refer to Chart WWW.

Following procedures analogous to those described above, but using Enantiomer 1 of Preparation 152, the title compound is obtained.

Physical characteristics are as follows:
¹H NMR δ 0.8-1.0, 1.2-2.2, 3.90, 6.9-7.2, 8.0, 8.15, 8.9 ppm.
HRMS: 497.1984
R_f 0.38 (15% ethyl acetate in dichloromethane)

EXAMPLE 362 N-[3-{1(*S* or *R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-propyl}phenyl]-5-cyanopyridine-2-sulfonamide
(Formula WWW-4, R₁ and R₂ are propyl, R₃ is ethyl, R₄ is 5-cyanopyridine-2-yl) [Enantiomer 2] Refer to Chart WWW.

Following procedures analogous to those described above, but using Enantiomer 2 of Preparation 152, the title compound is obtained.

Physical characteristics are as follows:
¹H NMR δ 0.8-1.0, 1.2-2.2, 3.90, 6.9-7.2, 8.0, 8.15, 8.9 ppm.
HRMS: 497.1980
R_f 0.38 (15% ethyl acetate in dichloromethane)

EXAMPLE 363 N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-propyl}phenyl]-5-aminopyridine-2-sulfonamide
(Formula WWW-4, R₁ and R₂ are propyl, R₃ is ethyl, R₄ is 5-aminopyridine-2-yl) [Enantiomer 1] Refer to Chart WWW.

Following procedures analogous to those described above, but using Enantiomer 1 of Preparation 152, the title compound is obtained.

Physical characteristics are as follows:

^1H NMR δ 0.7-0.9, 1.2-2.2, 3.8, 6.8-7.2, 7.5, 7.9 ppm.

HRMS: 487.2122

R_f 0.28 (5% methanol in dichloromethane)

EXAMPLE 364 N-[3-{1(*S* or *R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-propyl}phenyl]-5-aminopyridine-2-sulfonamide
5 (Formula WWW-4, R_1 and R_2 are propyl, R_3 is ethyl, R_4 is 5-aminopyridine-2-yl) [Enantiomer 2] Refer to Chart WWW.

Following procedures analogous to those described above, but using Enantiomer 2 of Preparation 152, the title compound is obtained.

10 Physical characteristics are as follows:

^1H NMR δ 0.7-0.9, 1.2-2.2, 3.8, 6.8-7.2, 7.5, 7.9 ppm.

HRMS: 487.2140

R_f 0.28 (5% methanol in dichloromethane)

15 PREPARATION 153 Resolution of N-[3-[1-(4-Hydroxy-5,6-dihydro-2-oxo-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl]-phenyl]carbamic acid, phenylmethyl ester to give 4 isomers (Formula WWW-2: R_1 is 4-fluorophenethyl, R_2 is propyl, and R_3 is ethyl) Refer to Chart WWW.

The enantiomers are defined by elution order from System D. HPLC System
20 D consists of a 0.46x25 cm Chiralcel OD-H column (Chiral Technologies, Inc.) with 20% isopropanol and 0.05% trifluoroacetic acid in hexane (V/V) pumped at 0.5 mL/min. The retention times in this system are (Isomer 1) 21.6, (Isomer 2) 34.5, (Isomer 3) 55.2 and (Isomer 4) 66.6 min.

25 Separate the enantiomers on a 2.1x25 cm Chiralcel OD column (Chiral Technologies, Inc.). Aliquots are injected and the enantiomers eluted with 17.5% isopropanol in hexane (V/V) at 10 mL/min. Fractions eluting near 24.6, 42.9, 66.3 and 77.4 min are pooled appropriately after assay with System D. In order of elution, the four isomers are designated Isomers 1-4, respectively.

30 In all cases, whenever solvent is stripped from a pool the following protocol is used: Solvent is removed from pools of fractions on a rotary evaporator with house vacuum (ca. 30 mm Hg) and a water bath set at $45 \pm 5^\circ$. If acetic acid is present in the solvent, add ca. 10 mL of toluene/L of pool before the flask goes dry. Residues are then washed into tared flasks using methylene chloride and the solvent is stripped as above. Final solvent removal is accomplished at ambient temperature, 1
35 mmHg pressure for 2-24 hours before weighing.

EXAMPLE 365 N-[3-{1(*R or S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R or S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula WWW-4: R₁ is 4-fluorophenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 5-cyanopyridine-2-yl) [Isomer 1] Refer to Chart WWW.

Following procedures analogous to those described above, but using Isomer 1 of Preparation 153, the title compound is obtained.

Physical characteristics are as follows:

¹H NMR δ 0.8-1.0, 1.3, 1.6-2.2, 2.5, 3.9, 6.8-7.3, 7.9-8.1, 8.9 ppm.

HRMS: 578.2120

R_f 0.35 (15% ethyl acetate in dichloromethane)

EXAMPLE 366 N-[3-{1(*R or S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S or R*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula WWW-4: R₁ is 4-fluorophenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 5-cyanopyridine-2-yl) [Isomer 2] Refer to Chart WWW

Following procedures analogous to those described above, but using Isomer 2 of Preparation 153, the title compound is obtained.

Physical characteristics are as follows:

¹H NMR δ 0.8-1.0, 1.3, 1.6-2.2, 2.5, 3.9, 6.8-7.3, 7.9-8.1, 8.9 ppm.

HRMS: 578.2120

R_f 0.35 (15% ethyl acetate in dichloromethane)

EXAMPLE 367 N-[3-{1(*S or R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R or S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide (Formula WWW-4: R₁ is 4-fluorophenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 5-cyanopyridine-2-yl) [Isomer 3] Refer to Chart WWW.

Following procedures analogous to those described above, but using Isomer 3 of Preparation 153, the title compound is obtained.

Physical characteristics are as follows:

¹H NMR δ 0.8-1.0, 1.3, 1.6-2.2, 2.5, 3.9, 6.8-7.3, 7.9-8.1, 8.9 ppm.

HRMS: 578.2126

R_f 0.35 (15% ethyl acetate in dichloromethane)

EXAMPLE 367A N-[3-{1(*S or R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S or R*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-

cyanopyridine-2-sulfonamide (Formula WWW-4: R₁ is 4-fluorophenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 5-cyanopyridine-2-yl) [Isomer 4] Refer to Chart WWW.

Following procedures analogous to those described above, but using Isomer 4 of Preparation 153, the title compound is obtained.

Physical characteristics are as follows:

¹H NMR δ 0.8-1.0, 1.3, 1.6-2.2, 2.5, 3.9, 6.8-7.3, 7.9-8.1, 8.9 ppm.

HRMS: 578.2126

R_f 0.35 (15% ethyl acetate in dichloromethane)

10 EXAMPLE 368 N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R* or *S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide (Formula WWW-4: R₁ is 4-fluorophenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 1-methylimidazol-4-yl) [Isomer 1] Refer to Chart WWW.

15 Following procedures analogous to those described above, but using Isomer 1 of Preparation 153, the title compound is obtained.

Physical characteristics are as follows:

¹H NMR δ 0.8-1.0, 1.3, 1.6-2.2, 2.6, 3.63, 4.0, 6.9-7.5 ppm.

HRMS: 556.2265

20 R_f 0.29 (5% methanol in dichloromethane)

EXAMPLE 369 N-[3-{1(*R* or *S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R* or *S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-aminopyridine-2-sulfonamide (Formula WWW-4: R₁ is 4-fluorophenethyl, R₂ is propyl, R₃ is ethyl, R₄ is 5-aminopyridine-2-yl) [Isomer 1] Refer to Chart WWW.

25 Following procedures analogous to those described above, but using Isomer 1 of Preparation 153, the title compound is obtained.

Physical characteristics are as follows:

¹H NMR δ 0.8-1.0, 1.3, 1.6-2.2, 2.5, 3.9, 6.8-7.2, 7.5, 7.9 ppm.

30 HRMS: 568.2271

R_f 0.27 (5% methanol in dichloromethane)

PREPARATION 154 Hexahydro-2H-1-benzopyran-2,4(3H)-dione (Formula DDDD-2, wherein n is 1) Refer to Chart DDDD.

A solution of 0.42 g of platinum oxide and 1.66 g of the compound of formula DDDD-1 wherein n is 1 in 100 mL of acetic acid is placed on a Parr hydrogenation

35

2187523

apparatus under an initial pressure of 50 psi of hydrogen for 1.5 h. The reaction mixture is then filtered through Celite and concentrated *in vacuo* to give a beige solid. The crude material is purified by flash column chromatography on silica gel 60 (230-400 mesh) eluting with 0-5% methanol in chloroform to give 0.94 g of the
5 title product as a white solid.

Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 4.84-4.80, 3.54, 3.40, 2.60-2.53, 2.08-2.02, 1.79-1.65, 1.62-1.54, 1.44-1.40 ppm.

^{13}C NMR (CDCl_3) δ 203.0, 167.4, 74.3, 47.7, 45.6, 29.1, 23.5, 23.2, 19.7 ppm.

10 IR (mineral oil) 3092, 2768, 2714, 2695, 2662, 1657, 1614, 1577, 1444, 1352, 1345, 1340, 1323, 1308, 1295, 1287, 1260, 1244, 1211, 1188, 1057, 1004, 938, 909, 890, 843, 832, 600 cm^{-1} .

EI-MS: $[\text{M}^+]=168$.

Anal. found: C, 64.16; H, 7.16.

15 PREPARATION 155 4a,5,6,7,8,8a-Hexahydro-4-hydroxy-3-[1-(3-nitrophenyl)-propyl]-2H-1-benzopyran-2-one (Formula DDDD-4, wherein n is 1 and R_1 is ethyl) Refer to Chart DDDD.

A solution of 3.17 g of aluminum trichloride in 30 mL of tetrahydrofuran is added to a solution of 2.00 g of the title compound of Preparation 154 and 1.82 g of
20 3-nitrobenzaldehyde in 20 mL of tetrahydrofuran. The resulting mixture is then stirred at room temperature for 2.5 h, at which time, 7.28 g of sodium carbonate decahydrate is added, and the reaction mixture is stirred an additional 20 min. The mixture is then dried over magnesium sulfate, filtered through Celite, and concentrated *in vacuo* to yield 6.05 g of a yellow gum. This crude material is
25 immediately dissolved in 50 mL of tetrahydrofuran containing 0.73 g of cuprous bromide-dimethyl sulfide complex, and 13.1 mL of a 1.0 M solution of triethyl aluminum in hexanes are added to the reaction mixture. After stirring at room temperature for 1 h, the reaction is quenched by the addition of water, and the resulting mixture is partitioned between ether and water. The organic layer is
30 separated, washed with brine, and concentrated *in vacuo* to produce 4.0 g of a yellow oil. The crude material is purified by flash column chromatography eluting with 10-50% ethyl acetate in hexanes to yield 0.63 g of the title product as a yellow foam.

Physical characteristics are as follows:

MP 86-91 $^{\circ}\text{C}$.

35 IR (mineral oil) 3085, 1635, 1569, 1528, 1448, 1394, 1365, 1349, 1325, 1307,

1288, 1270, 1251, 1244 cm^{-1} .

EXAMPLE 370 5-Cyano-N-[3-1-(4a,5,6,7,8,8a-hexahydro-4-hydroxy-2-oxo-2H-1-benzopyran-3-yl)propyl]phenyl]-2-pyridinesulfonamide (Formula DDDD-7, wherein n is 1, R_1 is ethyl, and R_2 is 5-cyano-2-pyridyl) Refer to Chart DDDD.

A solution of 0.63 g of the title compound of Preparation 155 in 50 mL of ethanol with 0.3 g of 10% palladium on carbon is placed on a Parr hydrogenation apparatus at an initial pressure of 50 psi of hydrogen for 3 h. The reaction mixture is then filtered through Celite and concentrated *in vacuo* to give 0.519 g of crude intermediate. 0.25 g of this intermediate is immediately dissolved in 5 mL of methylene chloride, and 0.168 g of 5-cyano-2-pyridylsulfonyl chloride and 0.134 mL of pyridine are added to the solution. The resulting mixture is stirred at room temperature for 18 h. The reaction mixture is then purified by flash column chromatography on silica gel 60 (230-400 mesh) eluting with 0-2.5% methanol in chloroform to give 0.164 g of the title product as a white foam.

Physical characteristics are as follows:

MP 122-125 °C.

HRMS found: 468.1611

EXAMPLE 371 4-Cyano-N-[3-1-(4a,5,6,7,8,8a-hexahydro-4-hydroxy-2-oxo-2H-1-benzopyran-3-yl)propyl]phenyl]-benzenesulfonamide (Formula DDDD-7, wherein n is 1, R_1 is ethyl, and R_2 is 4-cyanophenyl) Refer to Chart DDDD.

Following the general procedure of Example 370, and making non-critical variations, but substituting 4-cyanophenylsulfonyl chloride for 5-cyano-2-pyridylsulfonyl chloride, 0.236 g of the title compound is obtained as white foam.

Physical characteristics are as follows:

MP 127-130 °C.

HRMS found: 466.1583.

PREPARATION 156 4-Hexahydro-cyclohepta[b]pyran-2,4(3H,4aH)-dione (Formula DDDD-2, wherein n is 2) Refer to Chart DDDD.

Following the general procedure of Preparation 154, and making non-critical variations, but substituting the cycloheptylpyranone of Formula DDDD-1 wherein n is 2 for the cyclohexylpyranone of Formula DDDD-1 wherein n is 1, 0.337 g of the title compound is obtained as white solid.

Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 4.97-4.91, 3.52, 3.42, 2.64-2.58, 2.22-2.11, 2.01-1.72, 1.59-1.36 ppm.

5 ^{13}C NMR (CDCl_3) δ 203.0, 167.2, 78.0, 52.1, 46.5, 32.1, 28.6, 27.1, 25.7, 21.3 ppm.

IR (mineral oil) 3074, 2791, 2755, 2736, 2687, 2637, 2608, 2585, 1655, 1625, 1586, 1500, 1480, 1443, 1333, 1324, 1293 (s), 1265, 1254, 1240, 1222, 1196, 1173, 1082, 1053, 1016, 909, 889, 832, 611 cm^{-1} .

EI-MS: $[\text{M}^+]$ = 182.

10 Anal. found: C, 66.16; H, 7.90.

PREPARATION 157 5,6,7,8,9,9a-Hexahydro-4-hydroxy-3-[1-(3-nitrophenyl)-propyl]-cyclohepta[b]pyran-2(4aH)-one (Formula DDDD-4, wherein n is 2 and R_1 is ethyl) Refer to Chart DDDD.

15 Following the general procedure of Preparation 155, and making non-critical variations, but substituting the title compound of Preparation 156 for the title compound of Preparation 154, 2.5 g of the title compound is obtained as a yellow foam.

Physical characteristics are as follows:

MP 75-78 $^{\circ}\text{C}$.

20 IR (mineral oil) 3071, 2667, 1638, 1528, 1395, 1350, 1305, 1276, 1250, 1143, 1130, 1120, 1100, 1066, 782, 764, 741, 697, 685 cm^{-1} .

HRMS found: 345.1590.

Anal. found: C, 58.74; H, 5.63; N, 3.48.

25 EXAMPLE 372 5-Cyano-N-[3-[1-(2,4a,5,6,7,8,9,9a-octahydro-4-hydroxy-2-oxocyclohepta[b]pyran-3-yl)propyl]phenyl]-2-pyridinesulfonamide (Formula DDDD-7, wherein n is 2, R_1 is ethyl, and R_2 is 5-cyano-2-pyridyl) Refer to Chart DDDD.

30 Following the general procedure of Example 370, and making non-critical variations, but substituting the title compound of Preparation 157 for the title compound of Preparation 155, 0.206 g of the title compound is obtained as a white foam.

Physical characteristics are as follows:

MP 163-166 $^{\circ}\text{C}$.

35 IR (mineral oil) 3352, 3128, 3100, 3073, 3029, 1760, 1726, 1641, 1608, 1593, 1584, 1411, 1397, 1355, 1295, 1282, 1242, 1207, 1173, 1125, 1106, 1086, 1074, 1028,

974, 967, 721, 701, 645, 638 cm^{-1} .

HRMS found: 481.1693.

PREPARATION 158 Octahydro-2H-cycloocta[b]pyran-2,4(3H)-dione (Formula DDDD-2, wherein n is 3) Refer to Chart DDDD.

5 Following the general procedure of Preparation 154, and making non-critical variations, but substituting the cyclooctylpyranone of Formula DDDD-1 wherein n = 3 for the cycloheptylpyranone of Formula DDDD-1 wherein n = 2, 1.72 g of the title compound is obtained as a white solid.

Physical characteristics are as follows:

10 ^1H NMR (CDCl_3) δ 4.84-4.78, 3.61, 3.40, 2.75-2.70, 2.14-1.97, 1.90-1.72, 1.68-1.44 ppm.

^{13}C NMR (CDCl_3) δ 204.2, 167.2, 78.2, 49.5, 46.1, 28.5, 27.3, 26.2, 24.7, 23.9, 22.1 ppm.

15 IR (mineral oil) 2659, 2617, 1650, 1612, 1579, 1444, 1356, 1332, 1307, 1287, 1265, 1244, 1227, 1209, 1041, 1035, 1003, 962, 946, 860, 832, 824 cm^{-1} .

HRMS found: 196.1100.

Anal. found: C, 67.06; H, 8.23.

PREPARATION 159 3-[2,2-Dimethyl-1-(3-nitrophenyl)propyl]-
4a,5,6,7,8,9,10,10a-octahydro-4-hydroxy-2H-cyclo-
20 octa[b]pyran-2-one (Formula DDDD-4, wherein n is 3
and R_1 is *t*-butyl) Refer to Chart DDDD.

25 A solution of 1.36 g of aluminum trichloride in 30 mL of tetrahydrofuran is added to a solution of 1.0 g of the title compound of Preparation 158 and 0.77 g of 3-nitrobenzaldehyde in 20 mL of tetrahydrofuran. The resulting mixture is then stirred at room temperature for 2.3 h, at which time, 3.06 g of sodium carbonate decahydrate is added, and the reaction mixture is stirred an additional 15 min. The mixture is then dried over magnesium sulfate, filtered through Celite, and concentrated *in vacuo* to yield a yellow foam. This crude intermediate is immediately dissolved in 5 mL of tetrahydrofuran for use in the second step.

30 A dry flask is charged with 0.82 g of activated zinc, 3 mL of tetrahydrofuran, 0.035 mL of dibromoethane, and 0.21 mL of a 1 M solution of trimethylsilyl chloride in tetrahydrofuran. After the addition of each reagent the mixture is sonicated for 15 m at 45 °C. The mixture is diluted further by the addition of 2 mL tetrahydrofuran and 1.32 mL of *t*-butyl iodide is added dropwise. The resulting mixture is sonicated
35 for 3 h at 45 °C. A separate mixture of 0.85 g of copper(I) cyanide and 0.80 g of

lithium chloride in 4 mL of tetrahydrofuran is stirred at room temperature for 1 h until almost homogeneous and cooled to -30 °C. The organozinc solution is then added via cannula to the copper cyanide solution and the resulting mixture is allowed to warm to 0 °C and to stir for 15 min. The reaction mixture is then cooled to -78 °C, and the solution of crude intermediate prepared above is added. After stirring for 20 min at -78 °C and 30 min at 0 °C, the reaction is quenched with a saturated solution of aqueous ammonium chloride and diluted with an additional 60 mL of tetrahydrofuran. The organic layer is separated, washed with water, and concentrated *in vacuo* to give 2.17 g of an orange foam. The crude material is then purified by flash column chromatography eluting with 10-30% ethyl acetate in hexanes followed by recrystallization in methylene chloride/hexanes to yield 0.60 g of the title product as a yellow solid.

Physical characteristics are as follows:

MP 158-161 °C.

IR (mineral oil) 3077, 2646, 1632, 1599, 1529, 1477, 1450, 1396, 1357, 1349, 1334, 1317, 1283, 1273, 1252, 1232, 1217, 1205, 1181 cm⁻¹.

EXAMPLE 373 5-Cyano-N-[3-[2,2-dimethyl-1-(4a,5,6,7,8,9,10,10a-octahydro-4-hydroxy-2-oxo-2H-cycloocta[b]pyran-3-yl)propyl]phenyl]-2-pyridinesulfonamide (Formula DDDD-7, wherein n is 2, R₁ is *t*-butyl, and R₂ is 5-cyano-2-pyridyl) Refer to Chart DDDD.

Following the general procedure of Example 370, and making non-critical variations, but substituting the title compound of Preparation 159 for the title compound of Preparation 157, 0.034 g of the title compound is obtained as white crystals.

Physical characteristics are as follows:

MP 182-185 °C.

IR (mineral oil) 3246, 3121, 3098, 2615, 1655, 1633, 1607, 1585, 1575, 1491, 1411, 1395, 1354, 1335, 1322, 1311, 1298, 1281, 1275, 1262, 1255, 1233, 1206, 1178, 1121, 1109, 1028, 977, 702, 657, 646, 635, 605 cm⁻¹.

HRMS found: 524.2216.

Anal. found: C, 63.86; H, 6.41; N, 7.82.

PREPARATION 160 (3(3R),4S)-3-[2-[1-[3-[bis(phenylmethyl)amino]phenyl]-propyl]-5-hydroxy-1,3-dioxo-5-(2-phenylethyl)octyl]-4-phenyl-2-oxazolidinone (Formula W-10 wherein R₁ is 2-phenylethyl) Refer to Chart W.

To 100 mL of methylene chloride is added 5.0 g of the title compound of Preparation 95 (W-8) and the resulting solution cooled to -78°C under an atmosphere of nitrogen. To that solution is added 1.0 mL of TiCl_4 and 1.63 mL of diisopropylethylamine, and the resulting solution is stirred for 1 hour. Then, 3.30 g of 1-phenyl-3-hexanone is added, and the reaction temperature raised to 0 °C for 2.5 hours. The reaction is then quenched by the addition of a saturated ammonium chloride solution, and the mixture is extracted with methylene chloride. The organic extract is washed with saturated sodium bicarbonate solution and evaporated *in vacuo* to yield 9.7 g of a yellow oil. Column chromatography on 900 g silica (elution with 10% hexane-methylene chloride, 100% methylene chloride) affords 3.30 g of the title compound as a yellow foam.

Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 7.33-7.23, 7.14, 7.04, 6.61-6.50, 5.45, 5.22, 4.71, 4.60, 4.48, 4.26, 3.33, 3.15-3.03, 2.58, 2.47-2.31, 1.93, 1.40-1.28, 1.24-1.13, 1.11-0.96, 0.88-0.77, 0.62-0.57 ppm.

MP 121-126 °C.

^{13}C NMR (CDCl_3) δ 167.2, 167.1, 153.7, 142.6, 141.0, 138.2, 138.1, 129.6, 129.5, 129.2, 128.9, 128.8, 128.6, 128.4, 128.3, 127.0, 125.8, 125.6, 125.6, 73.1, 70.0, 69.9, 63.9, 57.9, 54.8, 54.7, 51.5, 51.4, 48.3, 41.3, 41.0, 40.8, 40.5, 29.8, 29.6, 27.1, 26.9, 16.8, 16.6, 14.6, 11.7, 11.6 ppm.

IR (mineral oil) 3525, 3061, 3026, 1777, 1720, 1690, 1601, 1495, 1361, 1335, 1238, 1199, 1104, 735, 698 cm^{-1} .

EI-MS: $[\text{M}^+] = 736$.

Anal. found: C, 78.03; H, 7.11; N, 3.79.

25 PREPARATION 161 (3S)-3-[1-[3-Bis(phenylmethyl)amino]phenyl]propyl]-5,6-dihydro-4-hydroxy-6-(2-phenylethyl-6-propyl-pyran-2-one (Formula W-11 wherein R_1 is 2-phenylethyl) Refer to Chart W.

To 5 mL of dry tetrahydrofuran is added 2.7 g of the title compound of Preparation 160 and the resulting solution is cooled to 0°C under an atmosphere of nitrogen. To that solution is added 0.45 mL of a 1 M solution of potassium *t*-butoxide in tetrahydrofuran. The reaction mixture is then warmed to 20°C and stirred for 2 hours. The reaction is quenched with saturated ammonium chloride and extracted with ethyl acetate. The organic layer is washed with water, dried and evaporated *in vacuo* to yield 0.28 g of a yellow oil. Column chromatography on 80 g

of silica gel (elution with 10-30% acetone/hexane) affords 0.195 g of a yellow foam. Crystallization from ethyl acetate/hexane yields 0.146 g of the title compound.

Physical characteristics are as follows:

MP 128-131 °C

5 ^1H NMR (CDCl_3) δ 7.35-7.12, 6.73-6.64, 5.84, 4.73-4.57, 4.12, 2.69-2.61, 2.38-2.20, 1.95-1.65, 1.41-1.32, 0.98-0.87 ppm.

^{13}C NMR (CDCl_3) δ 204.1, 204.0, 171.7, 171.4, 169.6, 140.9, 140.8, 140.6, 140.5, 140.4, 139.9, 139.8, 138.3, 129.7, 129.6, 129.5, 128.9, 128.6, 128.5, 128.4, 128.2, 128.1, 127.1, 126.9, 126.8, 126.7, 126.5, 126.4, 126.3, 126.2, 126.0, 125.9, 10 116.8, 112.6, 112.5, 112.4, 112.3, 112.2, 112.1, 112.0, 82.0, 81.9, 81.8, 80.4, 80.3, 58.6, 58.5, 54.5, 51.4, 50.4, 50.1, 49.9, 47.8, 47.4, 47.0, 46.6, 43.0, 42.9, 42.2, 41.9, 40.2, 40.1, 40.0, 39.2, 29.8, 29.7, 29.6, 29.1, 29.0, 26.8, 26.7, 24.7, 24.6, 24.3, 16.9, 16.5, 14.0, 12.3 ppm.

IR (mineral oil) 3023, 1637, 1599, 1584, 1575, 1494, 1347, 1300, 1257, 1243, 15 1234, 920, 731, 704, 695 cm^{-1} .

EI-MS: $[\text{M}^+] = 573$.

Anal. found: C, 81.53; H, 7.82; N, 2.34.

$[\alpha]_D$ (CHCl_3) = -83°

PREPARATION 162 (3S)-3-[1-(3-Aminophenyl)propyl]-5,6-dihydro-4-hydroxy-
20 6-(2-phenylethyl)-6-propyl-pyran-2-one (Formula W-12
wherein R_1 is 2-phenylethyl) Refer to Chart W.

0.63 g of the title compound of Preparation 161 is dissolved in 45 mL of ethyl acetate and 15 mL of methanol. To that solution is added 0.47 g of 10% Pd/C, and the resulting mixture is hydrogenated at 50 psi for 2.5 hours. The reaction is then
25 filtered through celite and concentrated *in vacuo* to yield 0.466 g of an off-white foam. Column chromatography on 80 g of silica gel (elution with 20-50% ethyl acetate-hexane) affords 0.389 g of the title compound as an off-white solid.

Physical characteristics are as follows:

MP 155-159 °C.

30 ^1H NMR (CD_3OD) δ 7.26-7.20, 7.15-7.04, 6.95, 6.81, 6.74, 6.54-6.51, 3.98-3.91, 2.68-2.54, 2.25-2.17, 2.02-1.67, 1.43-1.28, 0.99-0.87 ppm.

^{13}C NMR (CD_3OD) δ 171.2, 171.0, 148.5, 148.2, 143.9, 130.4, 130.2, 127.8, 120.8, 117.8, 115.3, 107.4, 82.7, 44.6, 44.4, 41.8, 41.7, 41.5, 38.4, 31.8, 26.8, 26.7, 18.8, 15.6, 14.3 ppm.

35 IR (mineral oil) 3085, 3061, 3026, 1617, 1605, 1495, 1314, 1258, 1168, 1119,

1065, 1030, 923, 776, 699 cm^{-1} .

EI-MS: $[M^+] = 393$.

Anal. found: C, 76.13; H, 8.16; N, 3.37.

$[\alpha]_D$ (MeOH) = -41°

5 **EXAMPLE 374** N-[3-[1-(S)-[5,6,-Dihydro-4-hydroxy-2-oxo-6-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]propyl]phenyl]-5-(trifluoromethyl)-2-pyridinesulfonamide (Formula W-13 wherein R_1 is 2-phenylethyl) Refer to Chart W.

To a solution of 0.200 g of the title compound of Preparation 162 in 5 mL of
10 methylene chloride is added 0.12 mL of pyridine. The resulting mixture is cooled to
0 $^\circ\text{C}$ and 0.132 g of 5-trifluoromethylpyridine-2-sulfonyl chloride is added. The
reaction mixture is then stirred at room temperature for 1.5 h, concentrated *in*
vacuo, and partitioned between ethyl acetate and water. The organic layer is
concentrated *in vacuo* to 0.39 g of pink oil. Column chromatography on 50 g silica
15 gel (elution with 20-50% ethyl acetate/hexane) affords 0.252 g of title compound as a
white foam.

MP 170-173 $^\circ\text{C}$.

^1H NMR (CD_3OD) δ 8.95-8.92, 8.23-8.16, 8.04-8.00, 7.25-6.90, 4.86, 3.98-3.90,
3.31, 3.30, 2.69-2.46, 2.18-2.09, 1.96-1.65, 1.41-1.28, 0.99-0.81 ppm.

20 ^{13}C NMR (CD_3OD) δ 167.3, 147.7, 147.5, 142.9, 142.8, 137.7, 137.0, 129.5,
129.2, 126.9, 126.2, 126.1, 124.1, 122.6, 122.5, 120.3, 120.2, 81.8, 81.7, 43.6, 43.2,
40.9, 40.5, 37.5, 30.9, 25.8, 25.6, 17.9, 14.7, 13.3, 13.2 ppm.

IR (mineral oil) 3087, 3027, 1642, 1606, 1595, 1327, 1260, 1173, 1142, 1110,
1074, 1016, 720, 700, 613 cm^{-1} .

25 FAB-MS: $[M+H] = 603$.

Anal. found: C, 61.79; H, 5.86; N, 4.48; S, 5.16.

$[\alpha]_D$ (MeOH) = -31° .

30 **PREPARATION 163** (3S,6R)-3-[1-[3-bis(phenylmethyl)amino]phenyl]propyl]-
5,6-dihydro-4-hydroxy-6-(2-phenylethyl-6-propyl-pyran-2-
one (Formula FFF-2) Refer to Chart FFF.

The title compound of Preparation 161 is chromatographed on a 5.1 x 30 cm
Cyclobond^{*} I 2000 column in an ice bath at 90 mg per injection using an automated
chromatographic system and a mobile phase of acetonitrile containing 0.1% diethyl-
amine and 0.05% glacial acetic acid (v/v). The eluant is monitored at 260 nm, the
35 flow rate is 45 mL/min, and appropriate fractions from multiple injections are

*Trade-mark

2187523

combined and concentrated *in vacuo* to give 0.300 g of a dark oil. The oil is partitioned between ethyl acetate, saturated aqueous sodium bicarbonate solution, and water. The organic layer is separated and concentrated *in vacuo*. Column chromatography on 50 g of silica gel (elution with 10-20% acetone/hexane) affords

5 0.22 g of the title compound of the compound as a colorless oil.

Physical characteristics are as follows:

The retention time of the title compound is 57 min.

PREPARATION 164 (3S,6S)-3-[1-[3-bis(phenylmethyl)amino]phenyl]propyl]-
5,6-dihydro-4-hydroxy-6-(2-phenylethyl)-6-propyl-pyran-2-
10 one (Formula FFF-3) Refer to Chart FFF.

The title compound of Preparation 161 is separated as described in Preparation 163 above. Further purification as described in Preparation 163 affords 0.117 g of the title compound as a colorless oil.

Physical characteristics are as follows:

15 The retention time of the title compound is 66 min.

PREPARATION 165 (3S,6R)-3-[1-(3-aminophenyl)propyl]-5,6-dihydro-4-
hydroxy-6-(2-phenylethyl)-6-propyl-pyran-2-one (Formula
FFF-4) Refer to Chart FFF.

Following the general procedure of Preparation 162, and making non-critical
20 variations, but substituting the title product of Preparation 163 for the title product of Preparation 161, 0.022 g of the title compound is obtained.

Physical characteristics are as follows:

¹H NMR (CD₃OD) δ 7.25-7.18, 7.15-7.12, 7.07-7.05, 6.97, 6.82-, 6.76-6.71,
6.53, 4.00-3.92, 2.67-2.54, 2.29-2.15, 2.06-1.92, 1.90-1.62, 1.46-1.28, 0.97-0.88 ppm.

25 EXAMPLE 375 (3S,6R)-N-[3-[1-[5,6,-dihydro-4-hydroxy-2-oxo-6-(2-phenylethyl)-
6-propyl-2H-pyran-3-yl]propyl]phenyl]-5-(trifluoromethyl)-2-
pyridinesulfonamide (Formula FFF-5) Refer to Chart FFF.

Following the general procedure of Example 374, and making non-critical
variations, but substituting the title product of Preparation 165 for the title product
30 of Preparation 162, 0.024 g of the title compound is obtained as a white foam.

Physical characteristics are as follows:

MP 156-159 °C.

- ¹H NMR (CD₃OD) δ 8.90, 8.20-8.17, 8.02-7.99, 7.28-6.88, 4.00-3.90, 2.71-2.46,
2.20-2.10, 1.98-1.67, 1.41-1.28, 0.98-0.81 ppm.

35 EXAMPLE 376 (3S)-N-[3-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6,6-dipropyl-2H-

pyran-3-yl)propyl]phenyl]-5-(trifluoromethyl)-2-pyridine-sulfonamide (Formula W-13) Refer to Chart W.

The title compound of Preparation 99 (formula W-12) 182 mg is dissolved in 5 mL of methylene chloride and 133 μ L of pyridine added. The reaction is cooled to 0°C and 142 mg of 5-trifluoromethyl-2-pyridinesulfonylchloride added. The reaction is stirred for 30 minutes and the methylene chloride is evaporated and the resulting material diluted with ethyl acetate. The organic solution is washed with water, brine and then dried over sodium sulfate. Evaporation of solvent gives 580 mg of crude product. Silica gel chromatography (50 g) eluting with 50% ethyl acetate/hexane affords 211 mg of the desired product as a white foam.

Physical characteristics are as follows:

Anal. found: C, 57.80; H, 5.95; N, 5.01; S, 5.64

$[\alpha]_D$ (18.094 mg/2mL CHCl_3) = -30°

EXAMPLE 377 (3R)-N-[3-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6,6-dipropyl-2H-pyran-3-yl)propyl]phenyl]-5-(trifluoromethyl)-2-pyridinesulfonamide (Formula X-13) Refer to Chart X.

The title compound of Preparation 107 (formula X-12) 170 mg is dissolved in 5 mL of methylene chloride and 136 μ L of pyridine added. The reaction is cooled to 0°C and 132 mg of 5-trifluoromethyl-2-pyridinesulfonylchloride is added. The reaction is stirred for 30 minutes and the methylene chloride is evaporated and the resulting material diluted with ethyl acetate. The organic solution is washed with water, brine and then dried over sodium sulfate. Evaporation of solvent gives crude product which is chromatographed over 50 g of silica gel eluting with 50% ethyl acetate/hexane affords 225 mg of the desired product as a white foam.

Physical characteristics are as follows:

$[\alpha]_D$ (mg/2mL CHCl_3) = +29°

PREPARATION 166 (3S)(4R) 3-[2-[1-[3-[Bis(phenylmethyl)amino]phenyl]-propyl]-5-hydroxy-1,3-dioxo-5-phenethyloctyl]-4-phenyl-2-oxazolidinone (Formula X-10 where R_1 is phenethyl) Refer to Chart X.

To 1.12 g of the title compound of Preparation 104 is added 20 mL of methylene chloride and the resulting solution cooled to -78°C. To that solution is added 237 μ L of TiCl_4 followed by 400 μ L of diisopropylethylamine and the resulting solution is stirred at -78°C for 1 hour. To the aforementioned solution is added 776 μ L of 1-phenyl-3-hexanone and stirring continued at -40°C for 40 minutes and then

the temperature is raised to -10°C for 1.5 hours. The reaction is quenched with the addition of a saturated ammonium chloride solution, then extraction with methylene chloride and evaporation of the organic extracts. The crude material is chromatographed over 200 g of silica gel eluting with 10% hexane/methylene chloride to afford 870 mg of the title compound.

Physical characteristics are as follows:

IR (mineral oil) 2956, 2926, 2854, 1777, 1600, 1494, 1452, 698 cm^{-1} .

$[\alpha]_{\text{D}}$ (16.578 mg in CHCl_3) = $+4^{\circ}$

Mass Spectrum: molecular ion at 736.

Anal. found. C, 78.00; H, 7.14; N, 3.61.

PREPARATION 167 (3R) 3-[1-[3-[Bis(phenylmethyl)amino]phenyl]propyl]-5,6-dihydro-4-hydroxy-6-phenethyl-6-propyl-2H-pyran-2-one
(Formula X-11 where R_1 is phenethyl) Refer to Chart X.

The compound of Preparation 166 (750 mg) is added to 5 mL of dry THF and potassium tert. butoxide (1.0 M in THF; 1.2 mL) is added. The reaction is stirred at 20°C for 30 minutes and then quenched by the addition of a saturated ammonium chloride solution. The reaction is extracted with ethyl acetate, the organic extracts washed with water and brine and finally evaporated to afford the crude product. Silica gel chromatography over 100 g of silica gel eluting with 15% ethyl acetate/hexane affords 511 mg of the title product.

Physical characteristics are as follows:

IR (mineral oil) 2956, 2855, 1628, 1599, 1577, 1494, 1385, 1364, 697 cm^{-1} .

Anal. found: C, 81.30; H, 7.68; N, 2.30

Mass spectrum: molecular ion at 573.

$[\alpha]_{\text{D}}$ (18.116 mg/2 mL CH_3OH) = $+38^{\circ}$

PREPARATION 168 (3R) 3-[1-[3-aminophenyl]propyl]-5,6-dihydro-4-hydroxy-6-phenethyl-6-propyl-2H-pyran-2-one (Formula X-12 where R_1 is phenethyl) Refer to Chart X.

The compound of Preparation 167 (370 mg) is dissolved in 35 mL of ethyl acetate and 6 mL of methanol. To that solution is added 200 mg of 10% Pd on Carbon catalyst and the reaction is hydrogenated under 50 psi of hydrogen for 2 hours. The reaction is evaporated and chromatographed over 60 g of silica gel to yield 244 mg of the title compound.

Physical characteristics are as follows:

IR (mineral oil) 3025, 2954, 2871, 2854, 1635, 1619, 1604, 1494, 1456, 1383,

1378, 1256 cm^{-1} .

$[\alpha]_D$ (16.764 mg/mL in CH_3OH) = +39°.

Mass spectrum: molecular ion at 393.

Anal. found: C, 75.79; H, 8.05; N, 3.27.

- 5 **EXAMPLE 378** (3R)-N-[3-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6-propyl-6-phenethyl-2H-pyran-3-yl)propyl]phenyl]-5-(trifluoromethyl)-2-pyridinesulfonamide (Formula X-13 where R_1 is phenethyl)
Refer to Chart X.

The product of Preparation 168 (156 mg) is added to 5 mL of methylene
10 chloride. To that solution is added 96 μL of pyridine and then the reaction is cooled to 0°C. To the aforementioned solution is added 102 mg of 5-trifluoromethyl-2-pyridinyl sulfonyl chloride. The reaction is stirred for 1 hour and then poured into ethyl acetate, washed with water, brine and dried with MgSO_4 . The solvent is
15 evaporated in vacuo and the resulting material chromatographed over 100 g of silica gel eluting with 50% ethyl acetate/hexane to yield 200 mg of the title compound.

Physical characteristics are as follows:

Mass spectrum: molecular ion at 602.

IR (mineral oil) 2953, 2922, 2870, 2853, 1642, 1605, 1459, 1457, 1326, 1259, 1180, 1171, 1141 cm^{-1} .

- 20 UV (EtOH) λ_{max} (ϵ) 216 (22300), 264 sh (10700), 270 (11500), 279 (12100)
Anal. found: C, 57.53; H, 5.98; N, 4.84.

- EXAMPLE 379** (3R,6S)-N-[3-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6-propyl-6-phenethyl-2H-pyran-3-yl)propyl]phenyl]-5-(trifluoromethyl)-2-pyridinesulfonamide (Formula X-13 where R_1 is phenethyl)
25 Refer to Chart X.

The product of Example 378 is added to isopropanol and injected onto a 0.46x25 cm Cyclobond I 2000 column (Advanced Separations Technologies, Inc., Whippany, NJ). The column is in an ice-water bath. The sample is eluted at 1.0 mL/min. with acetonitrile containing 0.1% diethylamine and 0.6% glacial acetic acid
30 (V/V). The monitor is set at 250 nm. The earlier eluting diastereomer is identical to the compound of Example 298. The second eluting diastereomer is purified over 60 g of silica gel eluting with 40% ethyl acetate/hexane to afford 13 mg of the title compound.

Physical characteristics are as follows:

- 35 Opposite stereochemistry at C-6 to the compound of Example 298.

^1H -NMR (CD_3OD , δ) 8.91, 8.19, 8.16, 8.02, 7.99, 7.25, 7.18, 7.15, 7.13, 7.11, 7.04, 6.97, 6.89, 6.75, 3.95, 2.69, 2.64, 2.53, 2.48, 2.13, 1.91, 1.71, 1.68, 1.37, 1.19, 1.17, 1.14, 0.94, 0.92, 0.89, 0.85, 0.83, 0.80, 0.93.

PREPARATION 169 (3S)-3-[(3-Bis(phenylmethyl)amino)phenyl]-4,4-dimethylpentanoic acid methyl ester (Formula LLL-9)
Refer to Chart LLL.

To anhydrous methanol (2 mL) at room temperature is added titanium (IV) chloride (0.07 mL). The resulting light green solution is stirred for 2 h, treated with the compound of formula LLL-2 wherein R is phenyl (100 mg), prepared by procedures analogous to those described in Chart FF, and refluxed for 18 h. The reaction mixture is allowed to cool and is partitioned between 1N HCl and diethyl ether. The organic layer is separated washed with brine, dried over anhydrous sodium sulfate, and concentrated *in vacuo*. Purification by flash chromatography eluting with hexane/ethyl acetate (95:5) affords the title compound (58 mg) as a light amber oil.

Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 7.32-7.20, 7.04, 6.61-6.48, 4.61, 3.48, 2.85-2.80, 2.72-2.55, 0.75 ppm

^{13}C NMR (CDCl_3) δ 173.69, 148.45, 142.51, 138.91, 128.53, 128.17, 126.78, 117.98, 114.49, 110.89, 54.54, 52.24, 51.40, 35.56, 33.65, 27.87 ppm

MS (EI) m/z 415.

PREPARATION 170 (3S)-3-[(3-Bis(phenylmethyl)amino)phenyl]-4,4-dimethylpentanoic acid (Formula LLL-10) Refer to Chart LLL.

The compound of formula LLL-9 (406 mg) of Preparation 169 is slurried in glacial acetic acid (2.6 mL) and 6N sulfuric acid. The reaction mixture is refluxed for 5 h, allowed to cool and is partitioned between water and diethyl ether. The aqueous layer is separated and extracted two more times with diethyl ether. The combined organic layers are washed with brine, dried over anhydrous sodium sulfate, and concentrated *in vacuo*. The resulting light brown residue is dissolved in diethyl ether and treated with dicyclohexylamine (0.16 mL) at 0°C . The solids are isolated, washed with diethyl ether and dried *in vacuo*. The light brown solid is suspended in diethyl ether and washed with 0.25N HCl. The organic layer is washed with brine, dried over anhydrous sodium sulfate, and concentrated *in vacuo*, affording the title product (54 mg) as a light brown amorphous solid.

Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 7.31-7.19, 7.04, 6.61-6.48, 4.61, 2.81-2.56, 0.74 ppm

^{13}C NMR (CDCl_3) δ 179.15, 148.56, 142.28, 138.83, 128.55, 128.23, 126.76, 117.90, 114.49, 110.98, 54.51, 51.83, 35.45, 33.67, 27.84 ppm

MS (EI) m/z 401.

5 PREPARATION 171 N-[(S)-4-Benzyl-2-oxazolidinone] 3-aminocinnamate amide (Formula HHH-4) Refer to Chart HHH.

A 1 liter round-bottomed flask with nitrogen inlet and addition funnel is charged with 10.02 g of commercially available (S)-4-benzyl-2-oxazolidinone and 260 mL of tetrahydrofuran and then cooled to -78°C . To the aforementioned solution is added 37 mL of n-butyl lithium during which time a white solid separates from the reaction solution. To that suspension is added 11.46 g of *trans*-3-nitrocinnamic acid chloride (prepared from the treatment of commercially available 3-nitrocinnamic acid with oxalyl chloride) in a small volume of THF. The resulting pale yellow homogeneous solution is allowed to warm to room temperature and quenched with a saturated ammonium chloride solution and is extracted with ethyl acetate. The organic layer is separated, washed with brine and water, dried over magnesium sulfate, filtered and concentrated to give a reddish brown syrup (formula HHH-3 in Chart HHH) which is used without further purification. The aforementioned crude reaction mixture is added to ethanol containing 64.18 grams of $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ and that mixture heated at reflux for 20 minutes. The reaction is cooled to room temperature and poured into ice. The mixture is brought to pH 9-10 with saturated aqueous Na_2CO_3 . The mixture is filtered and the filter cake washed extensively with ethyl acetate. The filtrate is washed with brine and the organic phase is dried (Na_2SO_4), filtered, and concentrated in vacuo to give a yellow solid. Recrystallization from ethanol gives 11.56 g of the title product.

Physical characteristics are as follows:

IR (mineral oil) 3450, 3369, 2924, 1771, 1678, 1620, 1462, 1392, 1357, 1347, 1214 cm^{-1}

$[\alpha]_D$ (14.418 mg/mL in CHCl_3) = $+51^\circ$

30 PREPARATION 172 N-[(S)-4-Benzyl-2-oxazolidinone] 3-(bis(phenylmethyl)-amino) cinnamate amide (Formula HHH-5) Refer to Chart HHH.

The amine of formula HHH-4 from Preparation 172 (10.13 g), 10.48 g of potassium carbonate, 8.3 mL of benzyl bromide and 100 mL of acetonitrile is heated at reflux for 3 hours. The reaction is cooled to room temperature and partitioned

between water and ethyl acetate. The aqueous is extracted several additional times with ethyl acetate. The combined ethyl acetate extracts are dried (Na_2SO_4), filtered and concentrated in vacuo. The residue is purified via silica gel chromatography eluting with 25% ethyl acetate/hexane to yield 8.87 g of the title product.

5 Physical characteristics are as follows:

^{13}C -NMR (CDCl_3 , ppm) 165, 153, 149, 147, 138, 135, 129.6, 129.3, 128.8, 128.6, 127, 126.9, 126.5, 116.54, 116.50, 114, 113, 65, 55, 54, 37

IR (mineral oil) 2954, 2870, 2854, 1776, 1677, 1616, 1595, 1493, 1454, 1353, 1209, 988 cm^{-1}

10 PREPARATION 173 (3S)(4S) 3-[3-(3-(bis(phenylmethyl)aminophenyl)pentanoyl]-4-phenyl-2-oxazolidinone (Formula HHH-6)
Refer to Chart W.

A 100-mL, three-necked flask equipped with a stir-bar, 25-mL pressure-equalizing addition funnel, and a nitrogen inlet is charged with copper(I) bromide
15 dimethyl sulfide complex (1.69 g), 20 mL of tetrahydrofuran and 10 mL of dimethyl sulfide. The addition funnel is charged with the title compound of Preparation 172 (2.747 g) and 10 mL of tetrahydrofuran. The reaction mixture is cooled to -40°C and ethyl magnesium bromide (5.5 mL of a 3.0 M solution in ether) is added dropwise over 5 min. The resulting black mixture is stirred another 10 min at -40°C
20 $^\circ\text{C}$ and then allowed to warm to -10°C . The solution of the title compound of Preparation 172 in tetrahydrofuran is added dropwise to the reaction mixture over 17 min. The addition funnel is then rinsed with another 3 mL of tetrahydrofuran, and the reaction mixture is stirred for 2.5 h at ca. -40 to -60°C . The reaction is quenched by pouring the mixture into 50 mL of saturated aqueous ammonium
25 chloride solution, and the organic solvents are removed by concentration *in vacuo*. The resulting residue is partitioned between 75 mL of ethyl acetate and 50 mL of water and filtered through glass wool. The organic layer is then separated, washed with two 100-mL portions of 10% ammonium hydroxide solution and 50 mL of brine, dried over magnesium sulfate, filtered and concentration *in vacuo* to yield 3.59 g of a
30 yellow oil. Column chromatography on 150 g of silica gel (elution with 5-15% ethyl acetate/hexane) affords two diastereomeric products. 1.602 g of the title compound (the less polar diastereomer) is isolated as a pale yellow oil

Physical characteristics are as follows:

^1H NMR (CDCl_3) δ 7.32-7.17, 7.06, 6.60, 6.55, 4.63, 4.43-4.37, 4.00, 3.85, 3.37,
35 3.20, 3.08, 3.02-2.92, 2.62, 1.71-1.48, 0.73 ppm]

Also isolated from the column is 0.310 g of the more polar diastereomer as a pale yellow oil.

Physical characteristics are as follows:

¹H NMR (CDCl₃) δ 7.32-7.18, 7.12, 7.05, 6.64-6.56, 4.63, 4.60-4.52, 4.08-4.04, 3.48-3.38, 3.07-2.96, 2.48, 1.69-1.48, 0.73 ppm.

In addition, fractions containing 0.708 g of a ca. 1:4 ratio mixture of the less polar to more polar diastereomers are collected from the column.

PREPARATION 174 (3S,6S)-3-[1-(3-aminophenyl)propyl]-5,6-dihydro-4-hydroxy-6-(2-phenylethyl)-6-propyl-2H-pyran-2-one
(Formula FFF-6) Refer to Chart FFF.

Following the general procedure of Preparation 162, and making non-critical variations, but substituting the title product of Preparation [U-141164] for the title product of Preparation 161, 0.040 g of crude title compound is obtained. This compound is used immediately in the next step without further purification.

EXAMPLE 380 (3S,6S)-N-[3-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]propyl]phenyl]-5-(trifluoromethyl)-2-pyridinesulfonamide (Formula FFF-7) Refer to Chart FFF.

Following the general procedure of Example 374, and making non-critical variations, but substituting the title product of Preparation 174 for the title product of Preparation 162, 0.015 g of the title compound is obtained as a white foam.

Physical characteristics are as follows:

¹H NMR (CD₃OD) δ 8.95, 8.25-8.21, 8.07-8.02, 7.25-6.93, 3.94-3.88, 2.70-2.51, 2.20-2.18, 1.97-1.66, 1.40-1.30, 0.92-0.81 ppm.

Thus, for example, the compounds of the present invention include the following individual stereoisomers:

5-cyano-N-[3-(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,

5-cyano-N-[3-(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,

5-cyano-N-[3-(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,

5-cyano-N-[3-(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,

N-[3-(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(R)-(2-phenethyl)-6-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-

sulfonamide,

N-[3-(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(S)-(2-phenethyl)-6-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

5 N-[3-(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(S)-(2-phenethyl)-6-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

10 N-[3-(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(R)-(2-phenethyl)-6-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

5-amino-N-[3-(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,

5-amino-N-[3-(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,

15 5-amino-N-[3-(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,

5-amino-N-[3-(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(S)-(3,3,3-trifluoropropyl)-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,

20 N-[3-(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R)-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,

N-[3-(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(S)-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,

N-[3-(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(R)-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,

25 N-[3-(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenethyl)-6-(S)-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-2-pyridinesulfonamide,

5-Trifluoromethyl-N-[3-(R)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,

30 5-Trifluoromethyl-N-[3-(S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,

4-Trifluoromethyl-N-[3-(R)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,

4-Trifluoromethyl-N-[3-(S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,

35 5-Trifluoromethyl-N-[3-(R)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-

- pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,
5-Trifluoromethyl-N-[3-(S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-
pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,
5-Trifluoromethyl-N-[3-(R)-[1-[4-hydroxy-2-oxo-6,6-di-phenethyl-5,6-dihydro-
5 2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,
5-Trifluoromethyl-N-[3-(S)-[1-[4-hydroxy-2-oxo-6,6-di-phenethyl-5,6-dihydro-
2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,
4-Trifluoromethyl-N-[3-(R)-[1-[4-hydroxy-2-oxo-6,6-di-phenethyl-5,6-dihydro-
2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,
10 4-Trifluoromethyl-N-[3-(S)-[1-[4-hydroxy-2-oxo-6,6-di-phenethyl-5,6-dihydro-
2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,
4-Trifluoromethyl-N-[3-(R)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-
pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,
4-Trifluoromethyl-N-[3-(S)-[1-[4-hydroxy-2-oxo-6,6-di-n-propyl-5,6-dihydro-2H-
15 pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,
5-Trifluoromethyl-N-[3(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenethyl)-
6(R)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,
5-Trifluoromethyl-N-[3(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenethyl)-
6(S)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,
20 5-Trifluoromethyl-N-[3(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenethyl)-
6(R)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,
5-Trifluoromethyl-N-[3(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenethyl)-
6(S)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,
4-Trifluoromethyl-N-[3(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenethyl)-
25 6(R)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,
4-Trifluoromethyl-N-[3(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenethyl)-
6(S)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,
4-Trifluoromethyl-N-[3(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenethyl)-
6(R)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,
30 4-Trifluoromethyl-N-[3(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenethyl)-
6(S)-n-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]-phenyl]-2-pyridinesulfonamide,
5-Trifluoromethyl-N-[3(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenethyl)-
6(R)-n-propyl-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,
5-Trifluoromethyl-N-[3(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenethyl)-
35 6(S)-n-propyl-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,

- 5-Trifluoromethyl-N-[3(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenethyl)-6(R)-n-propyl-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,
5-Trifluoromethyl-N-[3(S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenethyl)-6(S)-n-propyl-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide,
5 5-Cyano-N-[3(R)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,
5-Cyano-N-[3(R)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,
5-Cyano-N-[3(S)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenylethyl)-6-propyl-10 2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,
5-Cyano-N-[3(S)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,
N-[3(R)-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6(R)-[2-phenylethyl]-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
15 N-[3(R)-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6(S)-[2-phenylethyl]-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
N-[3(S)-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6(R)-[2-phenylethyl]-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
N-[3(S)-[1-(5,6-Dihydro-4-hydroxy-2-oxo-6(S)-[2-phenylethyl]-6-propyl-2H-20 pyran-3-yl)-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
5-Amino-N-[3(R)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,
5-Amino-N-[3(S)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,
25 5-Amino-N-[3(S)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,
5-Amino-N-[3(R)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,
5-Amino-N-[3(R)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]propyl)phenyl]-2-pyridinesulfonamide,
30 5-Amino-N-[3(R)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]propyl)phenyl]-2-pyridinesulfonamide,
5-Amino-N-[3(S)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]propyl)phenyl]-2-pyridinesulfonamide,
35 5-Amino-N-[3(S)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(S)-(2-phenylethyl)-6-

propyl-2H-pyran-3-yl)propyl)phenyl]-2-pyridinesulfonamide,

5-Amino-N-[3(R)-(1-[6(R)-(2-[4-fluorophenyl]ethyl)-5,6-dihydro-4-hydroxy-2-oxo-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,

5-Amino-N-[3(R)-(1-[6(S)-(2-[4-fluorophenyl]ethyl)-5,6-dihydro-4-hydroxy-2-oxo-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,

5-Amino-N-[3(S)-(1-[6(R)-(2-[4-fluorophenyl]ethyl)-5,6-dihydro-4-hydroxy-2-oxo-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,

5-Amino-N-[3(S)-(1-[6(S)-(2-[4-fluorophenyl]ethyl)-5,6-dihydro-4-hydroxy-2-oxo-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,

10 N-[3(R)-(1-[5,6-Dihydro-6,6-dipropyl-4-hydroxy-2-oxo-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

N-[3(S)-(1-[5,6-Dihydro-6,6-dipropyl-4-hydroxy-2-oxo-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

15 5-Amino-N-[3(R)-(1-[5,6-dihydro-6,6-dipropyl-4-hydroxy-2-oxo-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,

5-Amino-N-[3(S)-(1-[5,6-dihydro-6,6-dipropyl-4-hydroxy-2-oxo-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,

5-Cyano-N-[3(R)-(1-[5,6-dihydro-6,6-dipropyl-4-hydroxy-2-oxo-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,

20 5-Cyano-N-[3(S)-(1-[5,6-dihydro-6,6-dipropyl-4-hydroxy-2-oxo-2H-pyran-3-yl]-2,2-dimethylpropyl)phenyl]-2-pyridinesulfonamide,

N-[3(R)-(1-[6,6-Bis(2-phenylethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-yl]propyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

25 N-[3(S)-(1-[6,6-Bis(2-phenylethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-yl]propyl)phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

N-[3(R)-(1-[6,6-Bis(2-phenyl-ethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-yl]propyl)phenyl]-5-cyano-2-pyridinesulfonamide,

N-[3(S)-(1-[6,6-Bis(2-phenyl-ethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-yl]propyl)phenyl]-5-cyano-2-pyridinesulfonamide,

30 N-[3-{1(R)-(4-Hydroxy-5,6-dihydro-2-oxo-6(R)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide,

N-[3-{1(R)-(4-Hydroxy-5,6-dihydro-2-oxo-6(S)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide,

35 N-[3-{1(S)-(4-Hydroxy-5,6-dihydro-2-oxo-6(R)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide,

- N-[3-{1(*S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S*)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide,
- N-[3-{1(*R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R*)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
- 5 N-[3-{1(*R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S*)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
- N-[3-{1(*S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R*)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
- N-[3-{1(*S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S*)-phenethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
- 10 N-[3-{1(*R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide,
- N-[3-{1(*R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide,
- 15 N-[3-{1(*S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide,
- N-[3-{1(*S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide,
- N-[3-{1(*R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
- 20 N-[3-{1(*R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
- 25 N-[3-{1(*S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
- N-[3-{1(*S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,
- 30 N-[3-{1(*R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-aminopyridine-2-sulfonamide,
- N-[3-{1(*S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-2,2-dimethylpropyl}phenyl]-5-aminopyridine-2-sulfonamide,
- 35 N-[3-{1(*R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-

2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

N-[3-{1(S)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-

2,2-dimethylpropyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

N-[3-{1(R)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-

5 2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide,

N-[3-{1(S)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-bis(2-phenylethyl)-2H-pyran-3-yl)-

2,2-dimethylpropyl}phenyl]-5-cyanopyridine-2-sulfonamide,

N-[3-{1(R)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-

propyl}phenyl]-5-cyanopyridine-2-sulfonamide,

10 N-[3-{1(S)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-

propyl}phenyl]-5-cyanopyridine-2-sulfonamide,

N-[3-{1(R)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-

propyl}phenyl]-5-aminopyridine-2-sulfonamide,

N-[3-{1(S)-(4-Hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)-

15 propyl}phenyl]-5-aminopyridine-2-sulfonamide,

N-[3-{1(R)-(4-Hydroxy-5,6-dihydro-2-oxo-6(R)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide,

N-[3-{1(S)-(4-Hydroxy-5,6-dihydro-2-oxo-6(R)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide,

20 N-[3-{1(R)-(4-Hydroxy-5,6-dihydro-2-oxo-6(S)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide,

N-[3-{1(S)-(4-Hydroxy-5,6-dihydro-2-oxo-6(S)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide,

25 N-[3-{1(R)-(4-Hydroxy-5,6-dihydro-2-oxo-6(R)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

N-[3-{1(S)-(4-Hydroxy-5,6-dihydro-2-oxo-6(S)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

N-[3-{1(R)-(4-Hydroxy-5,6-dihydro-2-oxo-6(S)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

30 N-[3-{1(S)-(4-Hydroxy-5,6-dihydro-2-oxo-6(R)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-1-methyl-1H-imidazole-4-sulfonamide,

N-[3-{1(R)-(4-Hydroxy-5,6-dihydro-2-oxo-6(R)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-aminopyridine-2-sulfonamide,

35 N-[3-{1(S)-(4-Hydroxy-5,6-dihydro-2-oxo-6(S)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-aminopyridine-2-sulfonamide,


N-[3-{1(*R*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*S*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-aminopyridine-2-sulfonamide,

N-[3-{1(*S*)-(4-Hydroxy-5,6-dihydro-2-oxo-6(*R*)-(2-(4-fluorophenyl)ethyl)-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-aminopyridine-2-sulfonamide,

5


FORMULA CHART

5




I

10




IA

15




IB

20




II

25



IIA

30




IIB

35


FORMULA CHART (Continued)

5




III

10



IV


15



V


20

25



VI


30



VII


FORMULA CHART (Continued)

5




VIII

10




IX

15



X

20



XI

25

CHART A




CHART B




CHART C




CHART D




CHART E




CHART E (Continued)












CHART E (Continued)




E-12




E-13



E-14



E-15



E-16

CHART F




CHART G




CHART H




CHART I




CHART J






CHART K



2187523

CHART L



2187523

CHART M






CHART N




2187523


CHART O



O-1 (N-7)




O-2





O-3

CHART P




P-1 (N-7)




P-2

2187523


CHART Q




Q-1 (S-4)




Q-2



Q-3




Q-4



Q-5

CHART R



2187523

CHART S




CHART T




CHART U




CHART V






CHART W



W-3







CHART W (Cont'd.)









CHART W (Cont'd.)















CHART X



X-3



2187523

CHART X (Cont'd.)














CHART X (Cont'd.)



2187523

CHART Y




CHART Z




CHART AA




CHART AA (Cont'd.)




CHART BB





CHART BB (Cont'd.)



2187523

CHART CC




CHART CC (Cont'd.)





CHART DD



2187523

CHART DD (Cont'd.)




CHART EE




CHART EE (Cont'd.)




CHART FF




CHART FF (Cont'd.)




CHART GG




CHART GG (Cont'd.)







CHART HH




HH-1 (W-6)




HH-2



HH-3




HH-4



HH-5

CHART HH (Cont'd.)



2187523

CHART II







CHART JJ




JJ-1(X-6)




JJ-2



JJ-3




JJ-4



JJ-5

CLAIMS

1. A compound of the formula



wherein R₁ and R₂ are independently selected from

- a) ethyl,
- b) n-propyl,
- c) cyclopropyl-(CH₂)₂-,
- d) F-phenyl-(CH₂)₂-,
- e) het-SO₂NH-phenyl-,
- f) 2-methylpropyl,
- g) phenyl-(CH₂)₂-, and
- h) F₃C-(CH₂)₂-;

R₃ is cyclopropyl, ethyl or t-butyl;

R₄ is het or cyanophenyl; and

het is 4-imidazolyl, 2-pyridinyl or 8-quinolinyl, any of which is optionally substituted by CH₃, CN, CF₃, NH₂ or CONH₂;

or a pharmaceutically acceptable salt thereof.

2. A compound of claim 1,

wherein R₁ is n-propyl or 2-phenylethyl;

R₂ is n-propyl or 2-phenylethyl;

R₃ is ethyl or t-butyl; and

R₄ is 4-imidazolyl or 2-pyridinyl, substituted with CH₃, CN or CF₃.

3. A compound of claim 1, which is 5-amino-N-[3(R or S)-(1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]propyl)phenyl]-2-pyridinesulfonamide, or a pharmaceutically acceptable salt thereof.

4. A compound of claim 1, which is N-[3(R or S)-(1-[6,6-bis(2-phenylethyl)-5,6-dihydro-4-hydroxy-2-oxo-2H-pyran-3-yl]propyl)phenyl]-5-cyano-2-pyridinesulfonamide, or a pharmaceutically acceptable salt thereof.

5. A compound of claim 1, which is N-[3-{1-(4-hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)propyl}phenyl]-5-carbamoylpyridine-2-sulfonamide, or a pharmaceutically acceptable salt thereof.

6. A compound of claim 1, which is N-[3-{1(R or S)-(4-hydroxy-5,6-dihydro-2-oxo-6(R or S)-phenylethyl-6-propyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide, or a pharmaceutically acceptable salt thereof.

7. A compound of claim 1, which is N-[3-{1(R or S)-(4-hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)propyl}phenyl]-5-cyanopyridine-2-sulfonamide, or a pharmaceutically acceptable salt thereof.


8. A compound of claim 1, which is N-[3-{1(R or S)-(4-hydroxy-5,6-dihydro-2-oxo-6,6-dipropyl-2H-pyran-3-yl)propyl}phenyl]-5-aminopyridine-2-sulfonamide, or a pharmaceutically acceptable salt thereof.

9. A compound of claim 1, which is (3R or S)-N-[3-[1-(5,6-dihydro-4-hydroxy-2-oxo-6,6-dipropyl-2H-pyran-3-yl)propyl]phenyl]-5-trifluoromethyl-2-pyridinesulfonamide, or a pharmaceutically acceptable salt thereof.


10. A compound of claim 1, which is N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]propyl]phenyl]-5-trifluoromethyl-2-pyridinesulfonamide, or a pharmaceutically acceptable salt thereof.

11. A compound of claim 1, which is N-[3(R or S)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R or S)-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]-2,2-dimethylpropyl]phenyl]-1-methyl-1H-imidazole-4-sulfonamide, or a pharmaceutically acceptable salt thereof.


12. A compound of Claim 1, which is 5-trifluoromethyl-N-[3(R)-[1-[5,6-dihydro-4-hydroxy-2-oxo-6(R)-(2-phenylethyl)-6(R)-n-propyl-2H-pyran-3-yl]-propyl]-phenyl]-2-pyridinesulfonamide, or a pharmaceutically acceptable salt thereof.




(I)



(II)



(III)



(IV)