

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 June 2006 (29.06.2006)

PCT

(10) International Publication Number
WO 2006/067134 A1

(51) International Patent Classification:

A61K 31/7076 (2006.01) A61P 37/02 (2006.01)
A61K 39/395 (2006.01) C07K 16/28 (2006.01)

(21) International Application Number:

PCT/EP2005/056943

(22) International Filing Date:

20 December 2005 (20.12.2005)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/638,674 22 December 2004 (22.12.2004) US
04106910.5 22 December 2004 (22.12.2004) EP

(71) Applicant (for all designated States except US): APPLIED RESEARCH SYSTEMS ARS HOLDING N.V. [NL/NL]; Pietermaai 15, Curaçao (AN).

(72) Inventor; and

(75) Inventor/Applicant (for US only): DE LUCA, Giampiero [IT/CH]; Chemin De Conches 15b, CH-1231 Conches (CH).

(74) Agent: SERONO INTERNATIONAL S.A.; Intellectual Property, 12, Chemin Des Aulx, CH-1228 Plan-les-Ouates (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMBINATION TREATMENT FOR MULTIPLE SCLEROSIS

(57) Abstract: This invention is directed towards treatments for multiple sclerosis comprising administering to a patient in need thereof a first agent, such as 2-chloro-2'-deoxyadenosine (2-CdA), that reduces the number of lymphocytes in combination with a second agent, such as an anti-alpha-4 integrin antibody, that blocks the adhesion of monocytes and leukocytes to endothelial cells. The use of 2-CdA combined with an anti-alpha-4 integrin antibody may be more effective than either treatment alone for MS. Furthermore, the combination treatment may allow for a lowering or altering the dose of one or more of the agents in order to limit any adverse effects associated with the individual agents, but maintaining the same therapeutic efficacy.



WO 2006/067134 A1

COMBINATION TREATMENT FOR MULTIPLE SCLEROSIS

BACKGROUND OF THE INVENTION

1. Field of the Invention

[0001] This invention relates to a method of treating multiple sclerosis (MS) by administering to a patient in need thereof a combination of agents characterized by a first agent that has the pharmacological activity of reducing the number of monocytes and a second agent that blocks the adhesion of monocytes and leukocytes to endothelial cells.

2. Description of Related Art

[0002] Multiple sclerosis (MS) is a progressive neurological autoimmune disease that affects 250,000 to 350,000 people in the United States alone. It is the result of demyelination in the brain and spinal cord. It is the most widely known chronic inflammatory demyelinating disease of the central nervous system (CNS) in humans. The onset of the disease typically occurs between the ages of twenty to forty, with twice as many women affected as men. MS onset is defined by the occurrence of the first neurological symptoms of CNS dysfunction. Symptoms resulting from this demyelination include weakness, visual impairment, incoordination, parathesis (abnormal tingling), CNS inflammation, brain atrophy and cognitive impairment. Sexual impairment and sphincter dysfunction may also be seen. The course of disease is unpredictable, but often progresses through a cycle of exacerbation of symptoms followed by remission. Remissions vary in length and may last several years but are infrequently permanent. Advances in cerebrospinal fluid (CSF) analysis and magnetic resonance imaging (MRI) have simplified the diagnostic process and facilitated early diagnosis (*Noseworthy et al., NEJM, 2000, 343(13):938-952*).

[0003] MS can be categorized by four courses of disease: relapsing-remitting (RR)-, secondary progressive (SP)-, primary progressive (PP)- and progressive relapsing (PR)- MS. More than 80% of patients will initially display a RR course with the clinical exacerbation of neurological symptoms, followed by a recovery that may or may not be complete (*Lublin and Reingold, Neurology, 1996, 46:907-911*). During RRMS, accumulation of disability

results from incomplete recovery from relapses. Approximately half of the patients with RRMS switch to SPMS within about ten years after disease onset. During the SP phase, worsening of the disability results from the accumulation of residual symptoms after exacerbation but also from insidious progression between exacerbations (*Lublin and Reingold, supra*). Ten per-cent of MS patients present with a PP course of disease, which is characterized by insidious progression of symptoms from disease onset. Less than 5% of patients have PRMS and share the same poor prognosis as PPMS patients.

[0004] It has been suggested that distinct pathogenic mechanisms may be involved in different patient subgroups and have wide-ranging implications for disease classification and treatment. (*Lassmann et al., 2001, Trends Mol. Med., 7, 115-121; Lucchinetti et al. 2001, Curr. Opin. Neurol., 14, 259-269*). While the etiology of MS is not known, numerous studies have implicated activated monocytes and macrophages, in the presence of activated T cells, in the progression and/or exacerbation of MS. (*Fredrickson et al., 1987, Acta Neurol. Scand., 75 :352-355 ; Huber et al., 1984, J. Exp. Med., 160 :310-316*). At the microscopic level, monocytes, microglial cells (macrophages of the CNS), and activated T-cells are found within the demyelinating regions of the nerve cells during MS exacerbations (Cecil, Textbook of Medicine (1979), Beeson et al. (eds.), W.B. Saunders Co., Philadelphia, PA.). Activated lymphocytes and macrophages have been implicated in the pathogenesis of acute inflammatory brain lesions and blood brain barrier (BBB) breakdowns associated with MS (*Coles et al. 1999, Ann. Neurol. 46(3): 296-304*).

[0005] In light of the association of disease progression with the presence of inflammatory cells, current medications for MS that are disease-modifying treatments, i.e., modifying the course of MS, work by modulating or suppressing the immune system. There are four FDA approved immunomodulating agents for RRMS: three beta interferons (Betaseron™, Berlex; Avonex™, Biogen; Rebif®, Serono) and Glatimer acetate (Copaxone™, Amgen). There is also one FDA approved immunosuppressing drug for worsening MS, Mitoxantrone (Novantrone™, Amgen). However, it is unclear whether the drugs actually prevent disease progression or if their benefits can be sustained over time. Also, some patients can not tolerate the side effects, such as flu-like symptoms and, in some cases, hair loss, blood cell

abnormalities, and depression (Fillippini et al. Interferons in relapsing remitting multiple sclerosis: a systematic review. *Lancet*, 361:545-552, 2003).

a. 2-Chloro-2'-Deoxyadenosine

[0006] Various additional agents are in clinical trials for the treatment of MS. Among them, a chlorinated purine analogue 2-chloro-2'-deoxyadenosine (2-CdA) has been suggested to be useful in the treatment of MS (*EP 626853B1 and US 5,506,214*). Its mode of action may be due to its pharmacologic activity toward activated monocytes and macrophages. Several clinical studies have been described wherein 2-CdA is administered intravenously and subcutaneously to MS patients. Effective doses of an oral formulation to treat MS have also been described (*US 5,506,214*).

[0007] Two double-blind, placebo controlled Phase II studies using 2-CdA were conducted respectively in the treatment of progressive forms of MS (*Selby et al., 1998, Can. J. Neurol. Sci., 25: 295-299*) and RRS (*Romine et al., 1999, Proc. Assoc. Amer. Phys., 111(1): 35-44*). In the first trial, the dose was 0.1 mg/kg for 7 days by continuous intravenous injection for four months. In the second trial, the 2-CdA dose was 0.07 mg/kg for 5 days by subcutaneous injection. A placebo controlled Phase III study was conducted in patients with PPMS and SPMS. In this study, patients received 2-CdA by subcutaneous injection at a dose of 0.07 mg/kg for either two months or for six months.

[0008] The above clinical studies demonstrated the efficacy of 2-CdA treatment in MS patients in terms of Kuzke Extended Disability Status Scale (EDSS), Scripps Neurologic Rating Scale (SNRS) scores and MRI findings (*Beutler et al., 1996, PNAS USA, 93:1716-1720; Romine et al., 1999, supra*). The Phase II study showed 2-CdA significantly reduced MRI-measured brain lesions (*Rice et al., 2000, supra*). Some adverse effects such as increased incidence of infections related to compromised immune function or myelosuppression were observed with the highest doses (*Selby et al., 1998, supra; Beutler et al. 1994, Acta hematol., 91: 10-15*). There was a narrow margin of safety between the effective dose and the occurrence of adverse effects, so therefore the oral route of 2-CdA treatment for MS was excluded (*Beutler et al., 1996, Sem. In Hematol., 33 1(S1): 45-52*).

b. Anti-Alpha-4 Integrin Antibody

[0009] An additional agent in clinical trials for the treatment of MS is a humanized monoclonal antibody specific for the alpha-4 chain of the integrin superfamily. Alpha-4-beta-1 integrin (VLA-4) is expressed on the extracellular surface of activated lymphocytes and monocytes, which have been implicated in the pathogenesis of the acute inflammatory brain lesions and blood brain barrier (BBB) breakdown associated with MS (*Coles et al. supra*). Agents against alpha-4 integrin have been tested for their anti-inflammatory potential both *in vitro* and *in vivo* in animal models (*Yednock et al., 1992, Nature, 356: 63-66; US 5,840,229; US 6,001,809*). The *in vitro* experiments demonstrated that anti-alpha-4 integrin antibodies block attachment of lymphocytes to brain endothelium (*Tubridy et al., 1999, Neurology, 53(3), 466-472*). Experiments in an animal model of MS, experimental autoimmune encephalomyelitis (EAE), demonstrated that administration of anti-alpha-4 integrin antibodies prevents inflammation of the brain and subsequent paralysis in the animals. Collectively, these experiments identify anti-alpha-4 agents as potentially useful therapeutic agents for treating MS as they prevent the trafficking of inflammatory cells across the BBB.

[0010] A Phase III trial showed that a monthly single dose of 300mg of anti-alpha-4 antibody significantly reduced MS lesions. Adverse effects seen in this study included a hypersensitivity-type reaction that was immunoglobulin-related and likely due to the fact that the agent is an antibody. Although not reported, a significant issue with protein therapeutics such as an antibody treatment is the eliciting of an immune response and subsequent development of neutralizing antibodies to the treatment making chronic administration difficult.

[0011] Although there have been some recent advances in treatments for MS, further improvements need to be made to facilitate the development of therapeutic agents with improved effectiveness and to reduce side effects. Accordingly, there is a need in the art for improved treatments for MS. This invention fulfills these needs and further provides other related advantages.

SUMMARY OF THE INVENTION

[0012] This invention is directed toward the use of a novel combination of agents to treat multiple sclerosis.

[0013] The method comprises administering to a patient in need thereof a first agent and a second agent, wherein said first agent has the pharmacological property of reducing the number of lymphocytes, and wherein said second agent blocks the adhesion of monocytes and leukocytes to endothelial cells.

[0014] An embodiment of the invention provides that said first agent is 2-CdA and said second agent is an anti-alpha-4 integrin antibody.

[0015] A further embodiment of the invention provides for a reduction in the effective dose of either or both agents needed to provide maximal therapeutic benefit in treating multiple sclerosis.

[0016] A further embodiment of the invention provides for pharmaceutical compositions used to treat multiple sclerosis, comprising a first agent and a second agent, wherein said first agent has the pharmacological property of reducing the number of lymphocytes, and wherein said second agent blocks the adhesion of monocytes and leukocytes to endothelial cells.

DETAILED DESCRIPTION

[0017] This invention is related to a method of treating MS comprising administering to a patient in need thereof 2-CdA and an anti-alpha-4 integrin antibody. Although 2-CdA and anti-alpha-4 integrin antibodies have each been used individually for the treatment of MS, the agents have not been used in combination for the treatment of MS. Combining 2-CdA and an anti-alpha-4 integrin antibody is more effective in the treatment of MS than either agent administered alone. Furthermore, the combination treatment allows for a reduction in the dose of one or both of the agents required to produce a therapeutically beneficial effect or a reduction in the length of treatment required to produce a therapeutically beneficial effect. Reducing the dose of one or both of the agents may reduce adverse effects associated

with administration of the individual agents, such as an immunoglobulin-related hypersensitivity reaction and/or an immune response to the anti-alpha-4 integrin antibody.

[0018] Without being bound by theory, the combination of agents may provide a synergistic effect. A synergistic effect of the agents may be due to different mechanisms of action for each of the individual agents. 2-CdA has the pharmaceutical effect of reducing lymphocyte numbers, whereas an anti-alpha-4 integrin antibody blocks the adhesion of monocytes and leukocytes to endothelial cells. It is specifically contemplated that the methods of this invention include the use of agents other than 2-CdA that reduce the number of lymphocytes, and agents other than anti-alpha-4 integrin antibodies that specifically bind to an integrin comprising an alpha-4 subunit and inhibits activity of the integrin.

[0019] According to one aspect of the invention, is provided a method for treating multiple sclerosis comprising administering to a patient in need thereof a first agent and a second agent, wherein said first agent has the pharmacological property of reducing the number of lymphocytes, and wherein said second agent blocks the adhesion of monocytes and leukocytes to endothelial cells.

[0020] According to one aspect of the invention, is provided a method treating multiple sclerosis comprising administering to a patient in need thereof a first agent and a second agent, wherein said first agent is a pharmaceutical formulation and has the pharmacological property of reducing the number of lymphocytes, and wherein said second agent is a pharmaceutical formulation that blocks the adhesion of monocytes and leukocytes to endothelial cells.

[0021] According to a further aspect, is provided a method according to the invention wherein said first agent is 2-chloro-2'-deoxyadenosine or pharmaceutically acceptable salts thereof.

[0022] According to a further aspect, is provided a method according to the invention wherein said second agent is an anti-alpha-4 integrin antibody.

[0023] According to a further aspect, is provided a method according to the invention wherein said second agent is a monoclonal antibody.

[0024] According to a further aspect, is provided a method according to the invention wherein said second agent is a chimeric antibody.

[0025] According to a further aspect, is provided a method according to the invention wherein said second agent is a human antibody.

[0026] According to a further aspect, is provided a method according to the invention wherein said second agent is a humanized antibody.

[0027] According to a further aspect, is provided a method according to the invention wherein said second agent is a single chain antibody.

[0028] According to a further aspect, is provided a method according to the invention wherein said second agent is an antibody Fab fragment.

[0029] According to a further aspect, is provided a method according to the invention wherein said composition comprises a suboptimal dose of said first agent and an effective dose of said second agent.

[0030] According to a further aspect, is provided a method according to the invention wherein said composition comprises an effective dose of said first agent and a suboptimal dose of said second agent.

[0031] According to a further aspect, is provided a method according to the invention wherein said composition comprises a sub-optimal dose of the first agent and the second agent.

[0032] According to a further aspect, is provided a method according to the invention wherein the first agent and second agent are administered simultaneously.

[0033] According to a further aspect, is provided a method according to the invention wherein the first agent and second agent are administered sequentially.

[0034] According to a further aspect, is provided a kit comprising the agents of claim 1 or 2 and instructions for using said agents to treat multiple sclerosis.

[0035] According to a further aspect, is provided a composition comprising a first agent having the pharmacological property of reducing the number of lymphocytes, and a second agent blocking the adhesion of monocytes and leukocytes to endothelial cells used to treat multiple sclerosis.

[0036] According to a further aspect, is provided a composition comprising as a first agent, a pharmaceutical formulation having the pharmacological property of reducing the number of lymphocytes, and as a second agent, a pharmaceutical formulation that blocks the adhesion of monocytes and leukocytes to endothelial cells used to treat multiple sclerosis.

[0037] According to another aspect, is provided a use of a combination of active agents for the preparation of a medicament for the treatment of an inflammatory or an autoimmune disorder, said combination of active agents comprising:

- (a) An agent which reduces the number of lymphocytes;
- (b) An agent which blocks the adhesion of monocytes and leukocytes to endothelial cells.

[0038] According to another aspect, is provided a use according to the invention wherein the agent (a) is 2-chloro-2'-deoxyadenosine or pharmaceutically acceptable salts thereof.

[0039] According to another aspect, is provided a use according to the invention wherein the agent (b) is an anti-alpha-4 integrin antibody.

[0040] According to another aspect, is provided a use according to the invention wherein the anti-alpha-4 integrin antibody is a monoclonal antibody.

[0041] According to another aspect, is provided a use according to the invention wherein the anti-alpha-4 integrin antibody is a chimeric antibody.

[0042] According to another aspect, is provided a use according to the invention wherein the anti-alpha-4 integrin antibody is a human antibody.

[0043] According to another aspect, is provided a use according to the invention wherein the anti-alpha-4 integrin antibody is a humanized antibody.

[0044] According to another aspect, is provided a use according to the invention wherein the anti-alpha-4 integrin antibody is a single chain antibody.

[0045] According to another aspect, is provided a use according to the invention wherein the anti-alpha-4 integrin antibody is a an antibody Fab fragment.

[0046] According to another aspect, is provided a use according to the invention wherein the said combination comprises a sub-optimal dose of agent (a) and an effective dose of agent (b).

[0047] According to another aspect, is provided a use according to the invention wherein said combination comprises an effective dose of agent (a) and a sub-optimal dose of agent (b).

[0048] According to another aspect, is provided a use according to the invention wherein said combination comprises a sub-optimal dose of agent (a) and agent (b).

[0049] According to another aspect, is provided a use according to the invention wherein agent (a) and agent (b) are used simultaneously.

[0050] According to another aspect, is provided a use according to the invention wherein agent (a) and agent (b) are used sequentially.

[0051] According to another aspect, is provided a use according to the invention wherein the autoimmune disorder is multiple sclerosis.

[0052] According to another aspect, is provided a kit comprising (a) 2-chloro-2'-deoxyadenosine or pharmaceutically acceptable salts thereof, (b) an anti-alpha-4 integrin antibody and instructions for using said agents to treat multiple sclerosis.

[0053] According to another aspect, is provided a pharmaceutical composition comprising (a) 2-chloro-2'-deoxyadenosine or pharmaceutically acceptable salts thereof, (b) an anti-alpha-4 integrin antibody; and a pharmaceutically acceptable carrier, diluent or excipient thereof.

1. Definitions

[0054] As used herein, "duration of treatment" refers to the period of administration of 2-CdA and an anti-alpha-4 integrin antibody according to the invention from the first day of treatment to the end of the treatment.

[0055] As used herein, "effective amount" when used in the context of a dosing regimen of 2-CdA in combination with an anti-alpha-4 integrin antibody, refers to that amount of either 2-CdA or an anti-alpha-4 integrin antibody, which if given in the absence of the other would

result in the maximal treatment of MS that each respective agent could effect if given as a single agent.

[0056] As used herein, “efficacy” when used in the context of a dosing regimen refers to the effectiveness of a particular treatment regimen. Efficacy can be measured based on changes in the course of disease in response to a dosing regimen of this invention. For example, treatment of MS efficacy can be measured by the frequency of relapses in RRMS and the presence or absence of new lesions in the CNS as detected using methods such as MRI.

[0057] As used herein, “elicits an immune response” and “eliciting an immune response” refer to the production of an immunological response to an anti-alpha-4 integrin antibody in a subject upon introduction of the agent. An immune response in the subject can be characterized by serum reactivity with an anti-alpha-4 integrin antibody that is at least twice that of an untreated subject., more preferably three times that of the reactivity of an untreated subject, and even more preferably at least four times the reactivity of an untreated subject.

[0058] As used herein, “end of treatment” refers to the day on which the patient becomes free of 2-CdA and/or an anti-alpha-4 integrin antibody .

[0059] As used herein, “in combination with” where used to describe administration of 2-CdA and an anti-alpha-4 integrin antibody means that the 2-CdA may be administered prior to, together with, or after the anti-alpha-4 integrin antibody.

[0060] As used herein, “pathological inflammation” refers to an inappropriate and chronic inflammation associated with MS (especially to inhibit further demyelination). Such inflammation is characterized by a heightened response of inflammatory cells, including infiltrating leukocytes. Over time, such pathological inflammation often results in damage to tissue in the region of inappropriate inflammation, namely the tissues of the CNS.

[0061] As used herein, “remission” refers to the period during the course of MS progression wherein a patient experiences a reduction or stabilization of symptoms of the disease. This may be a period of partial healing of CNS lesions, and/or a reduction in CNS inflammation.

[0062] As used herein, “specifically binds” or “binds specifically” refer to the situation in which one member of a specific binding pair will not show any significant binding to

molecules other than its specific binding partner (e.g., an affinity of about 1000X or more for its binding partner).

[0063] As used herein, “suboptimal dose” when used in the context of a dosing regimen of 2-CdA in combination with an anti-alpha-4 integrin antibody, refers to doses of either 2-CdA or an anti-alpha-4 integrin antibody that is less than the effective amount as defined supra

[0064] As used herein, “substantially homologous” refers to any polypeptide that has an alteration in the sequence such that a functionally equivalent amino acid is substituted for one or more amino acids in the polypeptide, thus producing a change that has no or relatively little effect on the binding properties of the polypeptide. For example, one or more amino acid residues within the sequence can be substituted by another amino acid of a similar polarity.

[0065] As used herein, “treatment” and “treating” and the like generally mean obtaining a desired pharmacological and physiological effect. The effect may be prophylactic in terms of preventing or partially preventing a disease, symptom or condition thereof and/or may be therapeutic in terms of a partial or complete cure of a disease, condition, symptom or adverse effect attributed to the disease. The term “treatment” as used herein covers any treatment of a disease in a mammal, particularly a human, and includes: (a) preventing the disease from occurring in a subject which may be predisposed to the disease but has not yet been diagnosed as having it; (b) inhibiting the disease, i.e., arresting its development; or relieving the disease, i.e., causing regression of the disease and/or its symptoms or conditions.

2. Combination

[0066] According to this invention 2-CdA is administered in combination with an anti-alpha-4 integrin antibody. The dosage of the agents may each be an effective amount. Preferably, the dosage of at least one of the agents is a suboptimal amount. More preferably, the dosage of each of the agents is as suboptimal amount.

[0067] The two agents may be administered simultaneously or sequentially with respect to each other. When administered simultaneously, the agents may be formulated in the same

composition or in different compositions. When the two agents are not administered as part of the same formulation, the two agents may be administered by the same or different routes of administration. The relative administration of 2-CdA and an anti-alpha-4 integrin antibody may be dependent on the specific route of administration and/or formulation of each component. The relative administration of 2-CdA and an anti-alpha-4 integrin antibody may be different depending on whether the patient is experiencing remission or exacerbation of symptoms.

[0068] The combination of agents may be administered for one or more treatment cycles. Each treatment cycle may be of any duration including, but not limited to, from about two months to six months, wherein 2-CdA is given about 1-2 times per day for a total of about 5-7 days per month and an-anti-alpha-4 integrin antibody is administered about once per monthly cycle, or about 1-4 times per monthly cycle or about once per week. The number of monthly cycles may also be from about 6 to about 12, or may be more than 12. The number of monthly cycles may vary depending on the course of MS (e.g. whether the patient is experiencing RR- or SPMS) and whether the patient is experiencing remission or exacerbation of symptoms.

3. 2-CdA

[0069] 2-CdA and its pharmacologically acceptable salts may be used in the practice of this invention. Processes for preparing 2-CdA are well known in the art. For example, the preparation of 2-CdA is described in *European Patent Application No. 173,059 A2* and *Robins et al., J. Am. Chem. Soc., 1984, 106: 6379, WO 04/028462, US 5,208,327 and WO 00/64918*. Alternatively, pharmaceutical preparations of 2-CdA may be purchased from Bedford Laboratories, Bedford, Ohio.

[0070] In general, the dosage of 2-CdA may vary over a relatively wide range to achieve, and preferably maintain the desired plasma concentration. The dosage of 2-CdA may be sufficient to provide a concentration in the patient's plasma of about 0.5 nM to about 50 nM, preferably about 1 nM to about 10 nM. An effective daily dose of 2-CdA may be about 0.04 to about 1.0 mg/kg body weight, more preferably about 0.04 to about 0.20 mg/kg/day,

and most preferably about 0.1mg/kg body weight. Additional dosage of 2-CdA is described in *US 5,506,214*, the contents of which are incorporated herein by reference.

4. Anti-Alpha-4 Integrin Antibodies

[0071] The anti-alpha-4 integrin antibody may specifically bind to an integrin comprising an alpha-4 subunit and may inhibit activity of the integrin. The antibody may also bind to an integrin dimer comprising the alpha-4 integrin, e.g., alpha-4-beta-1 or alpha-4-beta-7. The anti-alpha-4 integrin antibody may not show significant binding to any polypeptide other than an alpha-4 integrin or a receptor comprising an alpha-4 integrin. The antibodies may bind to an alpha-4 integrin with a binding affinity of 10^7 mole/l or more, preferably 10^8 mole/liters or more.

[0072] In general, the dosage of an anti-alpha-4 integrin antibody may vary over a relatively wide range to achieve, and preferably maintain the desired plasma concentration. An effective dose of an anti-alpha-4 integrin antibody is that which provides maximal reduction and/or suppression of MRI-detected lesions in the brain. The dosage of the antibody may be less than about 300 mg administered per month for a period of at least 6 months, more preferably 12 months and perhaps over the course of several years. Another preferred dosing regimen is 3 mg/kg of patient weight per month. Additional dosages of anti-alpha-4 integrin antibodies are described in *U.S. Patent Publication 2004/0009169*, the contents of which are incorporated herein by reference.

[0073] The antibodies of this invention include antibodies of classes IgG, IgM, IgA, IgD, and IgE, and fragments and derivatives thereof including Fab and F(ab')₂. The antibodies may also be recombinant antibody products including, but not limited to, single chain antibodies, chimeric antibody products, human antibodies, "humanized" antibody products, and "primatized" antibody products, CDR-grafted antibody products. The antibodies of this invention include monoclonal antibodies, polyclonal antibodies, affinity purified antibodies, or mixtures thereof which exhibit sufficient binding specificity to anti-alpha-4 integrin. The antibody may also be an antibody fragment. The antibody may be produced using standard

techniques, including as described in *WO 01/55210*, the contents of which are hereby incorporated by reference

[0074] The antibody may also be attached to a label. Labels can be signal-generating enzymes, antigens, other antibodies, lectins, carbohydrates, biotin, avidin, radioisotopes, toxins, heavy metals, and other compositions known in the art. Attachment techniques are also well known in the art.

[0075] Additional antibodies may be identified using techniques available in the art. For example, monoclonal antibodies of this invention can be produced using phage display technology. Antibody fragments that selectively bind to an alpha-4 integrin or a dimer comprising an alpha-4 integrin are then isolated. Exemplary preferred methods for producing such antibodies via phage display are disclosed in *U.S. Pat. Nos. 6,225,447; 6,180,336; 6,172,197; 6,140,471; 5, 969,108; 5, 885,793; 5,872,215; 5,871,907; 5,858,657; 5,837,242; 5,733, 743; and 5,565, 332*, the contents of which are incorporated herein by reference.

[0076] The antibodies may be provided by administering a polynucleotide encoding a whole or partial antibody (e. g., a single chain Fv) to a patient. The polynucleotide is administered to a subject in an appropriate vehicle to allow the expression of the antibody in the subject in a therapeutically effective amount.

5. Compositions

[0077] Compositions of this invention may further comprise one or more pharmaceutically acceptable additional ingredient(s) such as alum, stabilizers, antimicrobial agents, buffers, coloring agents, flavoring agents, adjuvants, and the like.

[0078] Compositions of this invention may be in the form of tablets or lozenges formulated in a conventional manner. For example, tablets and capsules for oral administration may contain conventional excipients including, but not limited to, binding agents, fillers, lubricants, disintegrants and wetting agents. Binding agents include, but are not limited to, syrup, accacia, gelatin, sorbitol, tragacanth, mucilage of starch and polyvinylpyrrolidone. Fillers include, but are not limited to, lactose, sugar, microcrystalline cellulose, maizestarch,

calcium phosphate, and sorbitol. Lubricants include, but are not limited to, magnesium stearate, stearic acid, talc, polyethylene glycol, and silica. Disintegrants include, but are not limited to, potato starch and sodium starch glycollate. Wetting agents include, but are not limited to, sodium lauryl sulfate). Tablets may be coated according to methods well known in the art.

[0079] Compositions of this invention may also be liquid formulations including, but not limited to, aqueous or oily suspensions, solutions, emulsions, syrups, and elixirs. The compositions may also be formulated as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may contain additives including, but not limited to, suspending agents, emulsifying agents, nonaqueous vehicles and preservatives. Suspending agent include, but are not limited to, sorbitol syrup, methyl cellulose, glucose/sugar syrup, gelatin, hydroxyethylcellulose, carboxymethyl cellulose, aluminum stearate gel, and hydrogenated edible fats. Emulsifying agents include, but are not limited to, lecithin, sorbitan monooleate, and acacia. Nonaqueous vehicles include, but are not limited to, edible oils, almond oil, fractionated coconut oil, oily esters, propylene glycol, and ethyl alcohol. Preservatives include, but are not limited to, methyl or propyl p-hydroxybenzoate and sorbic acid.

[0080] Compositions of this invention may also be formulated as suppositories, which may contain suppository bases including, but not limited to, cocoa butter or glycerides. Compositions of this invention may also be formulated for inhalation, which may be in a form including, but not limited to, a solution, suspension, or emulsion that may be administered as a dry powder or in the form of an aerosol using a propellant, such as dichlorodifluoromethane or trichlorofluoromethane. Compositions of this invention may also be formulated transdermal formulations comprising aqueous or nonaqueous vehicles including, but not limited to, creams, ointments, lotions, pastes, medicated plaster, patch, or membrane.

[0081] Compositions of this invention may also be formulated for parenteral administration including, but not limited to, by injection or continuous infusion. Formulations for injection may be in the form of suspensions, solutions, or emulsions in oily or aqueous vehicles, and

may contain formulation agents including, but not limited to, suspending, stabilizing, and dispersing agents. The composition may also be provided in a powder form for reconstitution with a suitable vehicle including, but not limited to, sterile, pyrogen-free water.

[0082] Compositions of this invention may also be formulated as a depot preparation, which may be administered by implantation or by intramuscular injection. The compositions may be formulated with suitable polymeric or hydrophobic materials (as an emulsion in an acceptable oil, for example), ion exchange resins, or as sparingly soluble derivatives (as a sparingly soluble salt, for example).

[0083] Compositions of this invention may also be formulated as a liposome preparation. The liposome preparation can comprise liposomes which penetrate the cells of interest or the stratum corneum, and fuse with the cell membrane, resulting in delivery of the contents of the liposome into the cell. For example, liposomes such as those described in *U.S. Patent No. 5,077,211 of Yarosh, U.S. Patent No. 4,621,023 of Redziniak et al. or U.S. Patent No. 4,508,703 of Redziniak et al.* can be used. Other suitable formulations can employ niosomes. Niosomes are lipid vesicles similar to liposomes, with membranes consisting largely of non-ionic lipids, some forms of which are effective for transporting compounds across the stratum corneum.

[0084] Representative formulations of 2-CdA are described in *US 6,194,395 and US 5,506,214, WO 96/19230, WO 96/19229, WO 04/087100 and WO 04/087101*, the contents of which are incorporated herein by reference.

6. Administration

[0085] Compositions of this invention may be administered in any manner including, but not limited to, orally, parenterally, sublingually, transdermally, rectally, transmucosally, topically, via inhalation, via buccal administration, or combinations thereof. Parenteral administration includes, but is not limited to, intravenous, intraarterial, intraperitoneal, subcutaneous, intramuscular, intrathecal, and intraarticular. The compositions of this invention may also be

administered in the form of an implant, which allows slow release of the compositions as well as a slow controlled i.v. infusion.

[0086] This invention is further illustrated by the following examples that are not intended to limit the scope of the invention in any way.

Example 1

Treatment of Multiple Sclerosis

A study of sixty patients with chronic multiple sclerosis is undertaken. Each patient is first examined for normal hepatic, renal, and bone marrow functioning to establish baseline values. Patients are randomly assigned to one of the treatment groups listed in Table 1.

Table 1

Group	2-CdA	Natalizumab
1	-	-
2	0.25 mg/kg	-
3	0.5 mg/kg	-
4	1 mg/kg	-
5	-	1 mg/kg
6	-	3 mg/kg
7	-	6 mg/kg
8	0.25 mg/kg	1 mg/kg
9	0.5 mg/kg	3 mg/kg
10	1 mg/kg	6 mg/kg

Each of the patients in Groups 2-4 and 8-10 is treated with 2-CdA dissolved in sterile preservative-free isotonic saline, whereas patients in Groups 1 and 5-7 receive a placebo (saline). The 2-CdA is administered intravenously for a total of seven days, once per month for a total of one year. Each of the patients in Groups 5-10 is treated with natalizumab, whereas Groups 1-4 received placebo (saline). The natalizumab is administered intravenously for a total of seven days, once per month for a total of one year, with the treatment being on the same day as 2-CdA treatment.

Patients are monitored to determine whether there is any progression or improvement of brain lesions associated with progression of MS. All patients have a baseline and MRI study (brain or spinal cord, according to localization of the lesions) at month 12.

Patients in Groups 1-3 and 5 have an increase in brain lesions. Patients in Groups 4 and 6-10 have a decrease in brain lesions. Importantly, patients in Group 10 show the largest decrease in brain lesions. Moreover, patients in Group 8 have a decrease in brain lesions similar to that of those patients in Groups 4 and 6. This data shows that the combination of 2-CdA and natalizumab is more effective than either agent administered alone. In addition, the data show that the combination of 2-CdA and natalizumab is effective at dosages that are ineffective for either 2-CdA or natalizumab administered alone.

CLAIMS

1. Use of a combination of active agents for the preparation of a medicament for the treatment of an inflammatory or an autoimmune disorder, said combination of active agents comprising:
 - (a) An agent which reduces the number of lymphocytes;
 - (b) An agent which blocks the adhesion of monocytes and leukocytes to endothelial cells.
2. Use according to claim 1 wherein the agent (a) is 2-chloro-2'-deoxyadenosine or pharmaceutically acceptable salts thereof.
3. Use according to claims 1 or 2 wherein the agent (b) is an anti-alpha-4 integrin antibody.
4. Use according to claim 3 wherein the anti-alpha-4 integrin antibody is a monoclonal antibody.
5. Use according to claim 3 wherein the anti-alpha-4 integrin antibody is a chimeric antibody.
6. Use according to claim 3 wherein the anti-alpha-4 integrin antibody is a human antibody.
7. Use according to claim 3 wherein the anti-alpha-4 integrin antibody is a humanized antibody.
8. Use according to claim 3 wherein the anti-alpha-4 integrin antibody is a single chain antibody.

9. Use according to claim 3 wherein the anti-alpha-4 integrin antibody is a an antibody Fab fragment.
10. Use according to any of claims 1 to 9 wherein said combination comprises a sub-optimal dose of agent (a) and an effective dose of agent (b).
11. Use according to any of claims 1 to 9 wherein said combination comprises an effective dose of agent (a) and a sub-optimal dose of agent (b).
12. Use according to any of claims 1 to 9 wherein said combination comprises a sub-optimal dose of agent (a) and agent (b).
13. Use according to any of the preceding claims wherein agent (a) and agent (b) are used simultaneously.
14. Use according to any of the preceding claims wherein agent (a) and agent (b) are used sequentially.
15. Use according to any of the preceding claims wherein the autoimmune disorder is multiple sclerosis.
16. A kit comprising (a) 2-chloro-2'-deoxyadenosine or pharmaceutically acceptable salts thereof, (b) an anti-alpha-4 integrin antibody and instructions for using said agents to treat multiple sclerosis.
17. A pharmaceutical composition comprising (a) 2-chloro-2'-deoxyadenosine or pharmaceutically acceptable salts thereof, (b) an anti-alpha-4 integrin antibody; and a pharmaceutically acceptable carrier, diluent or excipient thereof.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2005/056943

A. CLASSIFICATION OF SUBJECT MATTER A61K31/7076 A61K39/395 A61P37/02 //C07K16/28,(A61K39/395,31:7076)		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C07K A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LEARY S M ET AL: "CURRENT MANAGEMENT OF MULTIPLE SCLEROSIS" INTERNATIONAL JOURNAL OF CLINICAL PRACTICE, MEDICON INTERNATIONAL, ESHER, GB, vol. 54, no. 3, April 2000 (2000-04), pages 161-169, XP008018314 ISSN: 1368-5031 page 163, left-hand column, paragraph 4 - paragraph 5	1-17
X	WO 03/072040 A2 (ELAN PHARMACEUTICALS, INC) 4 September 2003 (2003-09-04) cited in the application	1,3-15
Y	claims 38,41; examples 1-3	2,16,17
	----- -/--	
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents :		
* A * document defining the general state of the art which is not considered to be of particular relevance * E * earlier document but published on or after the international filing date * L * document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) * O * document referring to an oral disclosure, use, exhibition or other means * P * document published prior to the international filing date but later than the priority date claimed		* T * later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention * X * document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone * Y * document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. * & * document member of the same patent family
Date of the actual completion of the international search		Date of mailing of the international search report
23 March 2006		30/03/2006
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Luyten, K

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2005/056943

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 506 214 A (BEUTLER ET AL) 9 April 1996 (1996-04-09) cited in the application	2
Y	examples 1,2	1,3-17
X	TUBRIDY N ET AL: "THE EFFECT OF ANTI-ALPHA4 INTEGRIN ANTIBODY ON BRAIN LESION ACTIVITY IN MS" NEUROLOGY, LIPPINCOTT WILLIAMS & WILKINS, PHILADELPHIA, US, vol. 53, no. 3, 11 August 1999 (1999-08-11), pages 466-472, XP008050557 ISSN: 0028-3878	1,3-15
Y	the whole document	2,16,17
X	BEUTLER E ET AL: "The treatment of chronic progressive multiple sclerosis with cladribine." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA. 20 FEB 1996, vol. 93, no. 4, 20 February 1996 (1996-02-20), pages 1716-1720, XP002339785 ISSN: 0027-8424	2
Y	cited in the application the whole document	1,3-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/EP2005/056943

Patent document cited in search report	Publication date	Publication date	Patent family member(s)	Publication date
WO 03072040	A2	04-09-2003	AU 2003213231 A1	09-09-2003
			BR 0307975 A	11-01-2005
			CA 2477178 A1	04-09-2003
			CN 1646163 A	27-07-2005
			EP 1485127 A2	15-12-2004
			JP 2005526045 T	02-09-2005
			MX PA04008267 A	10-11-2004
US 5506214	A	09-04-1996	US 5541164 A	30-07-1996