



US 20050089907A1

(19) **United States**

(12) **Patent Application Publication**
Friddle et al.

(10) **Pub. No.: US 2005/0089907 A1**

(43) **Pub. Date: Apr. 28, 2005**

(54) **NOVEL HUMAN KINASES AND
POLYNUCLEOTIDES ENCODING THE
SAME**

(76) Inventors: **Carl Johan Friddle**, The Woodlands,
TX (US); **Erin Hilbun**, Houston, TX
(US); **Brian Mathur**, The Woodlands,
TX (US); **C. Alexander Turner JR.**,
The Woodlands, TX (US)

Correspondence Address:
Lance K. Ishimoto
LEXICON GENETICS INCORPORATED
8800 Technology Forest Place
The Woodlands, TX 77381 (US)

(21) Appl. No.: **10/948,842**

(22) Filed: **Sep. 23, 2004**

Related U.S. Application Data

(60) Division of application No. 10/434,034, filed on May
8, 2003, now Pat. No. 6,815,188, which is a continu-
ation of application No. 09/992,481, filed on Nov. 19,
2001, now Pat. No. 6,593,125.

(60) Provisional application No. 60/252,011, filed on Nov.
20, 2000.

Publication Classification

(51) **Int. Cl.⁷** **C12Q 1/68**; C07H 21/04;
C12N 9/12

(52) **U.S. Cl.** **435/6**; 435/194; 536/23.2

(57) **ABSTRACT**

Novel human polynucleotide and polypeptide sequences are
disclosed that can be used in therapeutic, diagnostic, and
pharmacogenomic applications.

NOVEL HUMAN KINASES AND POLYNUCLEOTIDES ENCODING THE SAME

[0001] The present application claims the benefit of U.S. Provisional Application No. 60/252,011, which was filed on Nov. 20, 2000 and is herein incorporated by reference in its entirety.

1. INTRODUCTION

[0002] The present invention relates to the discovery, identification, and characterization of novel human polynucleotides encoding proteins sharing sequence similarity with animal kinases. The invention encompasses the described polynucleotides, host cell expression systems, the encoded proteins, fusion proteins, polypeptides and peptides, antibodies to the encoded proteins and peptides, and genetically engineered animals that either lack or overexpress the disclosed genes, antagonists and agonists of the proteins, and other compounds that modulate the expression or activity of the proteins encoded by the disclosed genes, which can be used for diagnosis, drug screening, clinical trial monitoring, the treatment of diseases and disorders, and cosmetic or nutraceutical applications.

2. BACKGROUND OF THE INVENTION

[0003] Kinases mediate the phosphorylation of a wide variety of proteins and compounds in the cell. In conjunction with phosphatases, kinases are involved in a range of regulatory pathways. Given the physiological importance of kinases, they have been subject to intense scrutiny and are proven drug targets.

3. SUMMARY OF THE INVENTION

[0004] The present invention relates to the discovery, identification, and characterization of nucleotides that encode novel human proteins and the corresponding amino acid sequences of these proteins. The novel human proteins (NHPs) described for the first time herein share structural similarity with animal kinases, including, but not limited to, receptor tyrosine kinases (SEQ ID NOS:1-2 show particular similarity to NEK family kinases, and SEQ ID NOS:3-5 are particularly similar to calcium and calmodulin dependent kinases as well as sequences encoding PK 80), and serine-threonine kinases. The described NHPs encode novel kinases having homologues and orthologs across a range of phyla and species.

[0005] The novel human polynucleotides described herein encode open reading frames (ORFs) encoding proteins of 692 and 817 amino acids in length (see respectively SEQ ID NOS:2 and 4).

[0006] The invention also encompasses agonists and antagonists of the described NHPs, including small molecules, large molecules, mutant NHPs, or portions thereof, that compete with native NHP, peptides, and antibodies, as well as nucleotide sequences that can be used to inhibit the expression of the described NHPs (e.g., antisense and ribozyme molecules, and open reading frame or regulatory sequence replacement constructs) or to enhance the expression of the described NHPs (e.g., expression constructs that place the described polynucleotide under the control of a strong promoter system), and transgenic animals that express a NHP sequence, or "knock-outs" (which can be

conditional) that do not express a functional NHP. Knock-out mice can be produced in several ways, one of which involves the use of mouse embryonic stem cells ("ES cells") lines that contain gene trap mutations in a murine homolog of at least one of the described NHPs. When the unique NHP sequences described in SEQ ID NOS:1-5 are "knocked-out" they provide a method of identifying phenotypic expression of the particular gene as well as a method of assigning function to previously unknown genes. In addition, animals in which the unique NHP sequences described in SEQ ID NOS:1-5 are "knocked-out" provide a unique source in which to elicit antibodies to homologous and orthologous proteins that would have been previously viewed by the immune system as "self" and therefore would have failed to elicit significant antibody responses. To these ends, gene trapped knockout ES cells have been generated in murine homologs of the described NHPs.

[0007] Additionally, the unique NHP sequences described in SEQ ID NOS:1-5 are useful for the identification of protein coding sequence and mapping a unique gene to a particular chromosome (the gene encoding SEQ ID NOS:1-2 is apparently encoded on human chromosome 17, see GENBANK accession no. AC010761, and the gene encoding SEQ ID. NOS:3-5 is apparently encoded on human chromosome 3, see GENBANK accession no. AC068979). These sequences identify biologically verified exon splice junctions as opposed to splice junctions that may have been bioinformatically predicted from genomic sequence alone. The sequences of the present invention are also useful as additional DNA markers for restriction fragment length polymorphism (RFLP) analysis, and in forensic biology.

[0008] Further, the present invention also relates to processes for identifying compounds that modulate, i.e., act as agonists or antagonists, of NHP expression and/or NHP activity that utilize purified preparations of the described NHPs and/or NHP products, or cells expressing the same. Such compounds can be used as therapeutic agents for the treatment of any of a wide variety of symptoms associated with biological disorders or imbalances.

4. DESCRIPTION OF THE SEQUENCE LISTING AND FIGURES

[0009] The Sequence Listing provides the sequence of the novel human ORFs encoding the described novel human kinase proteins. SEQ ID NO:5 describes a NHP ORF and flanking sequences.

5. DETAILED DESCRIPTION OF THE INVENTION

[0010] The NHPs described for the first time herein are novel proteins that are expressed in, inter alia, human cell lines and pituitary, thymus, spleen, lymph node, bone marrow, trachea, kidney, prostate, testis, thyroid, adrenal gland, pancreas, salivary gland, stomach, small intestine, skeletal muscle, heart, uterus, placenta, adipose, skin, bladder, rectum, pericardium, ovary, fetal kidney, fetal lung, gall bladder, tongue, aorta, 6-, 9-, and 12-week embryos, adenocarcinoma, osteosarcoma, and embryonic carcinoma cells (SEQ ID NOS:1-2). SEQ ID NOS:3-5 were predominantly expressed in fetal brain, brain, spinal cord, thymus, lymph node, trachea, lung, prostate, testis, thyroid, adrenal gland,

stomach, small intestine, skeletal muscle, uterus, placenta, mammary gland, skin, bladder, pericardium, hypothalamus, fetal kidney, fetal lung, tongue, aorta, 6-, 9-, and 12-week embryos, and embryonic carcinoma cells.

[0011] The described sequences were compiled from sequences available in GENBANK, and cDNAs generated from pituitary, lymph node, mammary gland, brain, adrenal gland, fetus, and testis mRNAs (Edge Biosystems, Gaithersburg, Md.).

[0012] The present invention encompasses the nucleotides presented in the Sequence Listing, host cells expressing such nucleotides, the expression products of such nucleotides, and: (a) nucleotides that encode mammalian homologs of the described genes, including the specifically described NHPs, and the NHP products; (b) nucleotides that encode one or more portions of an NHP that correspond to functional domains, and the polypeptide products specified by such nucleotide sequences, including, but not limited to, the novel regions of any active domain(s); (c) isolated nucleotides that encode mutant versions, engineered or naturally occurring, of the described NHPs in which all or a part of at least one domain is deleted or altered, and the polypeptide products specified by such nucleotide sequences, including, but not limited to, soluble proteins and peptides in which all or a portion of the signal sequence is deleted; (d) nucleotides that encode chimeric fusion proteins containing all or a portion of a coding region of a NHP, or one of its domains (e.g., a receptor/ligand binding domain, accessory protein/self-association domain, etc.) fused to another peptide or polypeptide; or (e) therapeutic or diagnostic derivatives of the described polynucleotides such as oligonucleotides, antisense polynucleotides, ribozymes, dsRNA, or gene therapy constructs comprising a sequence first disclosed in the Sequence Listing.

[0013] As discussed above, the present invention includes the human DNA sequences presented in the Sequence Listing (and vectors comprising the same), and additionally contemplates any nucleotide sequence encoding a contiguous NHP open reading frame (ORF) that hybridizes to a complement of a DNA sequence presented in the Sequence Listing under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65° C., and washing in 0.1×SSC/0.1% SDS at 68° C. (Ausubel et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & Sons, Inc., New York, at p. 2.10.3) and encodes a functionally equivalent expression product. Additionally, contemplated are any nucleotide sequences that hybridize to the complement of a DNA sequence that encodes and expresses an amino acid sequence presented in the Sequence Listing under moderately stringent conditions, e.g., washing in 0.2×SSC/0.1% SDS at 42° C. (Ausubel et al., 1989, supra), yet still encode a functionally equivalent NHP product. Functional equivalents of a NHP include naturally occurring NHPs present in other species and mutant NHPs whether naturally occurring or engineered (by site directed mutagenesis, gene shuffling, and/or directed evolution, as described in, for example, U.S. Pat. No. 5,837,458 or 5,723,323, both of which are herein incorporated by reference). The invention also includes degenerate nucleic acid variants of the disclosed NHP polynucleotide sequences.

[0014] Additionally contemplated are polynucleotides encoding NHP ORFs, or their functional equivalents, encoded by polynucleotide sequences that are about 99, 95, 90, or about 85 percent similar to corresponding regions of a NHP (as measured by BLAST sequence comparison analysis using, for example, the GCG sequence analysis package, as described herein, using default parameters).

[0015] The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore the complements of, the described NHP encoding polynucleotides. Such hybridization conditions can be highly stringent or less highly stringent, as described herein. In instances where the nucleic acid molecules are deoxyoligonucleotides (“DNA oligos”), such molecules are generally about 16 to about 100 bases long, or about 20 to about 80 bases long, or about 34 to about 45' bases long, or any variation or combination of sizes represented therein that incorporate a contiguous region of sequence first disclosed in the Sequence Listing. Such oligonucleotides can be used in conjunction with the polymerase chain reaction (PCR) to screen libraries, isolate clones, and prepare cloning and sequencing templates, etc.

[0016] Alternatively, such NHP oligonucleotides can be used as hybridization probes for screening libraries, and assessing gene expression patterns (particularly using a microarray or high-throughput “chip” format). Additionally, a series of the described NHP oligonucleotide sequences, or the complements thereof, can be used to represent all or a portion of the described NHP sequences. An oligonucleotide or polynucleotide sequence first disclosed in at least a portion of one or more of the sequences of SEQ ID NOS:1-5 can be used as a hybridization probe in conjunction with a solid support matrix/substrate (resins, beads, membranes, plastics, polymers, metal or metallized substrates, crystalline or polycrystalline substrates, etc.). Of particular note are spatially addressable arrays (i.e., gene chips, microtiter plates, etc.) of oligonucleotides and polynucleotides, or corresponding oligopeptides and polypeptides, wherein at least one of the biopolymers present on the spatially addressable array comprises an oligonucleotide or polynucleotide sequence first disclosed in at least one of the sequences of SEQ ID NOS:1-5, or an amino acid sequence encoded thereby. Methods for attaching biopolymers to, or synthesizing biopolymers on, solid support matrices, and conducting binding studies thereon are disclosed in, inter alia, U.S. Pat. Nos. 5,700,637, 5,556,752, 5,744,305, 4,631,211, 5,445,934, 5,252,743, 4,713,326, 5,424,186, and 4,689,405, the disclosures of which are herein incorporated by reference in their entirety.

[0017] Addressable arrays comprising sequences first disclosed in SEQ ID NOS:1-5 can be used to identify and characterize the temporal and tissue specific expression of a gene. These addressable arrays incorporate oligonucleotide sequences of sufficient length to confer the required specificity, yet be within the limitations of the production technology. The length of these probes is within a range of between about 8 to about 2000 nucleotides. Preferably the probes consist of 60 nucleotides and more preferably 25 nucleotides from the sequences first disclosed in SEQ ID NOS:1-5.

[0018] For example, a series of the described oligonucleotide sequences, or the complements thereof, can be used in

chip format to represent all or a portion of the described sequences. The oligonucleotides, typically between about 16 to about 40 (or any whole number within the stated range) nucleotides in length can partially overlap each other and/or the sequence may be represented using oligonucleotides that do not overlap. Accordingly, the described polynucleotide sequences shall typically comprise at least about two or three distinct oligonucleotide sequences of at least about 8 nucleotides in length that are each first disclosed in the described Sequence Listing. Such oligonucleotide sequences can begin at any nucleotide present within a sequence in the Sequence Listing and proceed in either a sense (5'-to-3') orientation vis-a-vis the described sequence or in an antisense orientation.

[0019] Microarray-based analysis allows the discovery of broad patterns of genetic activity, providing new understanding of gene functions and generating novel and unexpected insight into transcriptional processes and biological mechanisms. The use of addressable arrays comprising sequences first disclosed in SEQ ID NOS:1-5 provides detailed information about transcriptional changes involved in a specific pathway, potentially leading to the identification of novel components or gene functions that manifest themselves as novel phenotypes.

[0020] Probes consisting of sequences first disclosed in SEQ ID NOS:1-5 can also be used in the identification, selection and validation of novel molecular targets for drug discovery. The use of these unique sequences permits the direct confirmation of drug targets and recognition of drug dependent changes in gene expression that are modulated through pathways distinct from the intended target of the drug. These unique sequences therefore also have utility in defining and monitoring both drug action and toxicity.

[0021] As an example of utility, the sequences first disclosed in SEQ ID NOS:1-5 can be utilized in microarrays or other assay formats, to screen collections of genetic material from patients who have a particular medical condition. These investigations can also be carried out using the sequences first disclosed in SEQ ID NOS:1-5 in silico and by comparing previously collected genetic databases and the disclosed sequences using computer software known to those in the art.

[0022] Thus the sequences first disclosed in SEQ ID NOS:1-5 can be used to identify mutations associated with a particular disease and also in diagnostic and/or prognostic assays.

[0023] Although the presently described sequences have been specifically described using nucleotide sequence, it should be appreciated that each of the sequences can uniquely be described using any of a wide variety of additional structural attributes, or combinations thereof. For example, a given sequence can be described by the net composition of the nucleotides present within a given region of the sequence in conjunction with the presence of one or more specific oligonucleotide sequence(s) first disclosed in the SEQ ID NOS:1-5. Alternatively, a restriction map specifying the relative positions of restriction endonuclease digestion sites, or various palindromic or other specific oligonucleotide sequences can be used to structurally describe a given sequence. Such restriction maps, which are typically generated by widely available computer programs (e.g., the University of Wisconsin GCG sequence analysis

package, SEQUENCHER 3.0, Gene Codes Corp., Ann Arbor, Mich., etc.), can optionally be used in conjunction with one or more discrete nucleotide sequence(s) present in the sequence that can be described by the relative position of the sequence relative to one or more additional sequence(s) or one or more restriction sites present in the disclosed sequence.

[0024] For oligonucleotide probes, highly stringent conditions may refer, e.g., to washing in 6xSSC/0.05% sodium pyrophosphate at 37° C. (for 14-base oligos), 48° C. (for 17-base oligos), 55° C. (for 20-base oligos), or 60° C. (for 23-base oligos). These nucleic acid molecules may encode or act as NHP gene antisense molecules, useful, for example, in NHP gene regulation (and/or as antisense primers in amplification reactions of NHP gene nucleic acid sequences). With respect to NHP gene regulation, such techniques can be used to regulate biological functions. Further, such sequences can be used as part of ribozyme and/or triple helix sequences that are also useful for NHP gene regulation.

[0025] Inhibitory antisense or double stranded oligonucleotides can additionally comprise at least one modified base moiety that is selected from the group including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

[0026] The antisense oligonucleotide can also comprise at least one modified sugar moiety selected from the group including, but not limited to, arabinose, 2-fluoroarabinose, xylose, and hexose.

[0027] In yet another embodiment, the antisense oligonucleotide will comprise at least one modified phosphate backbone selected from the group including, but not limited to, a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

[0028] In yet another embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al., 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue et al., 1987, FEBS Lett. 215:327-330). Alternatively, double stranded RNA can be used to disrupt the expression and function of a targeted NHP.

[0029] Oligonucleotides of the invention can be synthesized by standard methods known in the art, e.g., by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides can be synthesized (Stein et al., 1988, Nucl. Acids Res. 16:3209), and methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. USA 85:7448-7451), etc.

[0030] Low stringency conditions are well-known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual (and periodic updates thereof), Cold Spring Harbor Press, N.Y.; and Ausubel et al., 1989, supra.

[0031] Alternatively, suitably labeled NHP nucleotide probes can be used to screen a human genomic library using appropriately stringent conditions or by PCR. The identification and characterization of human genomic clones is helpful for identifying polymorphisms (including, but not limited to, nucleotide repeats, microsatellite alleles, single nucleotide polymorphisms, or coding single nucleotide polymorphisms), determining the genomic structure of a given locus/allele, and designing diagnostic tests. For example, sequences derived from regions adjacent to the intron/exon boundaries of the human gene can be used to design primers for use in amplification assays to detect mutations within the exons, introns, splice sites (e.g., splice acceptor and/or donor sites), etc., that can be used in diagnostics and pharmacogenomics.

[0032] For example, the present sequences can be used in restriction fragment length polymorphism (RFLP) analysis to identify specific individuals. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification (as generally described in U.S. Pat. No. 5,272,057, incorporated herein by reference). In addition, the sequences of the present invention can be used to provide polynucleotide reagents, e.g., PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for example, providing another "identification marker" (i.e., another DNA sequence that is unique to a particular individual). Actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments.

[0033] Further, a NHP gene homolog can be isolated from nucleic acid from an organism of interest by performing PCR using two degenerate or "wobble" oligonucleotide primer pools designed on the basis of amino acid sequences within the NHP products disclosed herein. The template for the reaction may be total RNA, mRNA, and/or cDNA obtained by reverse transcription of mRNA prepared from, for example, human or non-human cell lines or tissue known to express or suspected of expressing an allele of a NHP gene.

[0034] The PCR product can be subcloned and sequenced to ensure that the amplified sequences represent the sequence of the desired NHP gene. The PCR fragment can then be used to isolate a full length cDNA clone by a variety

of methods. For example, the amplified fragment can be labeled and used to screen a cDNA library, such as a bacteriophage cDNA library. Alternatively, the labeled fragment can be used to isolate genomic clones via the screening of a genomic library.

[0035] PCR technology can also be used to isolate full length cDNA sequences. For example, RNA can be isolated, following standard procedures, from an appropriate cellular or tissue source (i.e., one known to express, or suspected of expressing, a NHP gene). A reverse transcription (RT) reaction can be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" using a standard terminal transferase reaction, the hybrid may be digested with RNase H, and second strand synthesis may then be primed with a complementary primer. Thus, cDNA sequences upstream of the amplified fragment can be isolated. For a review of cloning strategies that can be used, see e.g., Sambrook et al., 1989, supra.

[0036] A cDNA encoding a mutant NHP sequence can be isolated, for example, by using PCR. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known to express, or suspected of expressing, a mutant NHP allele, in an individual putatively carrying a mutant NHP allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal sequence. Using these two primers, the product is then amplified via PCR, optionally cloned into a suitable vector, and subjected to DNA sequence analysis through methods well-known to those of skill in the art. By comparing the DNA sequence of the mutant NHP allele to that of a corresponding normal NHP allele, the mutation(s) responsible for the loss or alteration of function of the mutant NHP gene product can be ascertained.

[0037] Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of carrying, or known to carry, a mutant NHP allele (e.g., a person manifesting a NHP-associated phenotype such as, for example, immune disorders, obesity, high blood pressure, etc.), or a cDNA library can be constructed using RNA from a tissue known to express, or suspected of expressing, a mutant NHP allele. A normal NHP sequence, or any suitable fragment thereof, can then be labeled and used as a probe to identify the corresponding mutant NHP allele in such libraries. Clones containing mutant NHP sequences can then be purified and subjected to sequence analysis according to methods well-known to those skilled in the art.

[0038] Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known to express, or suspected of expressing, a mutant NHP allele in an individual suspected of carrying, or known to carry, such a mutant allele. In this manner, gene products made by the putatively mutant tissue may be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against a normal NHP product, as described below (for screening techniques, see, for example, Harlow and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor).

[0039] Additionally, screening can be accomplished by screening with labeled NHP fusion proteins, such as, for example, alkaline phosphatase-NHP or NHP-alkaline phosphatase fusion proteins. In cases where a NHP mutation results in an expression product with altered function (e.g., as a result of a missense or a frameshift mutation), polyclonal antibodies to a NHP are likely to cross-react with a corresponding mutant NHP expression product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well-known in the art.

[0040] An additional application of the described novel human polynucleotide sequences is their use in the molecular mutagenesis/evolution of proteins that are at least partially encoded by the described novel sequences using, for example, polynucleotide shuffling or related methodologies. Such approaches are described in U.S. Pat. Nos. 5,830,721, 5,837,458, 6,117,679, and 5,723,323, which are herein incorporated by reference in their entirety.

[0041] The invention also encompasses: (a) DNA vectors that contain any of the foregoing NHP coding sequences and/or their complements (i.e., antisense); (b) DNA expression vectors that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences (for example, baculovirus as described in U.S. Pat. No. 5,869,336 herein incorporated by reference); (c) genetically engineered host cells that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell; and (d) genetically engineered host cells that express an endogenous NHP sequence under the control of an exogenously introduced regulatory element (i.e., gene activation). As used herein, regulatory elements include, but are not limited to, inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include, but are not limited to, the cytomegalovirus (hCMV) immediate early gene, regulatable, viral elements (particularly retroviral LTR promoters), the early or late promoters of SV40 and adenovirus, the lac system, the trp system, the TAC system, the TRC system, the major operator and promoter regions of phage lambda, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase (PGK), the promoters of acid phosphatase, and the promoters of the yeast α -mating factors.

[0042] Where, as in the present instance, some of the described NHP peptides or polypeptides are thought to be cytoplasmic or nuclear proteins (although processed forms or fragments can be secreted or membrane associated), expression systems can be engineered that produce soluble derivatives of a NHP (corresponding to a NHP extracellular and/or intracellular domains, or truncated polypeptides lacking one or more hydrophobic domains) and/or NHP fusion protein products (especially NHP-Ig fusion proteins, i.e., fusions of a NHP domain to an IgFc), NHP antibodies, and anti-idiotypic antibodies (including Fab fragments) that can be used in therapeutic applications. Preferably, the above expression systems are engineered to allow the desired peptide or polypeptide to be recovered from the culture media.

[0043] The present invention also encompasses antibodies and anti-idiotypic antibodies (including Fab fragments),

antagonists and agonists of a NHP, as well as compounds or nucleotide constructs that inhibit expression of a NHP sequence (transcription factor inhibitors, antisense and ribozyme molecules, or open reading frame sequence or regulatory sequence replacement constructs), or promote the expression of a NHP (e.g., expression constructs in which NHP coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, etc.).

[0044] The NHPs or NHP peptides, NHP fusion proteins, NHP nucleotide sequences, antibodies, antagonists and agonists can be useful for the detection of mutant NHPs or inappropriately expressed NHPs for the diagnosis of disease. The NHP proteins or peptides, NHP fusion proteins, NHP nucleotide sequences, host cell expression systems, antibodies, antagonists, agonists and genetically engineered cells and animals can be used for screening for drugs (or high throughput screening of combinatorial libraries) effective in the treatment of the symptomatic or phenotypic manifestations of perturbing the normal function of a NHP in the body. The use of engineered host cells and/or animals can offer an advantage in that such systems allow not only for the identification of compounds that bind to the endogenous receptor/ligand of a NHP, but can also identify compounds that trigger NHP-mediated activities or pathways.

[0045] Finally, the NHP products can be used as therapeutics. For example, soluble derivatives such as NHP peptides/domains corresponding to NHPs, NHP fusion protein products (especially NHP-Ig fusion proteins, i.e., fusions of a NHP, or a domain of a NHP, to an IgFc), NHP antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists (including compounds that modulate or act on downstream targets in a NHP-mediated pathway) can be used to directly treat diseases or disorders. For instance, the administration of an effective amount of soluble NHP, or a NHP-IgFc fusion protein or an anti-idiotypic antibody (or its Fab) that mimics the NHP could activate or effectively antagonize the endogenous NHP or a protein interactive therewith. Nucleotide constructs encoding such NHP products can be used to genetically engineer host cells to express such products *in vivo*; these genetically engineered cells function as "bioreactors" in the body delivering a continuous supply of a NHP, a NHP peptide, or a NHP fusion protein to the body. Nucleotide constructs encoding functional NHPs, mutant NHPs, as well as antisense and ribozyme molecules can also be used in "gene therapy" approaches for the modulation of NHP expression. Thus, the invention also encompasses pharmaceutical formulations and methods for treating biological disorders.

[0046] Various aspects of the invention are described in greater detail in the subsections below.

5.1 The NHP Sequences

[0047] The cDNA sequences and corresponding deduced amino acid sequences of the described NHPs are presented in the Sequence Listing.

[0048] Expression analysis has provided evidence that the described NHPs can be expressed in a range of human tissues, as described in greater detail herein above. In addition to serine-threonine kinases, the described NHPs also share significant similarity to several additional kinase families, including kinases associated with signal transduc-

tion, from a variety of phyla and species. Several polymorphisms were identified in the described NHPs. These include a T/C polymorphism in the sequence region represented by nucleotide position 1170 of SEQ ID NO:1, both of which result in the same amino acid being present at the corresponding amino acid (aa) position of SEQ ID NO:2; a T/C polymorphism in the sequence region represented by nucleotide position 1321 of SEQ ID NO:1, both of which result in the same amino acid being present at the corresponding aa position of SEQ ID NO:2; a C/G polymorphism in the sequence region represented by nucleotide position 94 of SEQ ID NO:3, which can result in either a leu or val being present at corresponding aa position 32 of SEQ ID NO:4; an A/G polymorphism at nucleotide position 112 of SEQ ID NO:3, which can result in either a lys or glu being present at corresponding aa position 38 of SEQ ID NO:4; and an A/T polymorphism at nucleotide position 133 of SEQ ID NO:3, which can result in either a thr or ser being present at corresponding aa position 45 of SEQ ID NO:4. The above polymorphisms can be present either singly, or in any combination or permutation within a given sequence.

[0049] An additional application of the described novel human polynucleotide sequences is their use in the molecular mutagenesis/evolution of proteins that are at least partially encoded by the described novel sequences using, for example, polynucleotide shuffling or related methodologies. Such approaches are described in U.S. Pat. Nos. 5,830,721 and 5,837,458, which are herein incorporated by reference in their entirety.

[0050] NHP gene products can also be expressed in transgenic animals. Animals of any species, including, but not limited to, worms, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, birds, goats, and non-human primates, e.g., baboons, monkeys, and chimpanzees may be used to generate NHP transgenic animals.

[0051] Any technique known in the art may be used to introduce a NHP transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Hoppe and Wagner, 1989, U.S. Pat. No. 4,873,191); retrovirus mediated gene transfer into germ lines (Van der Putten et al., 1985, Proc. Natl. Acad. Sci. USA 82:6148-6152); gene targeting in embryonic stem cells (Thompson et al., 1989, Cell 56:313-321); electroporation of embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814); and sperm-mediated gene transfer (Lavitrano et al., 1989, Cell 57:717-723); etc. For a review of such techniques, see Gordon, 1989, Transgenic Animals, Intl. Rev. Cytol. 115:171-229, which is incorporated by reference herein in its entirety.

[0052] The present invention provides for transgenic animals that carry the NHP transgene in all their cells, as well as animals that carry the transgene in some, but not all their cells, i.e., mosaic animals or somatic cell transgenic animals. The transgene may be integrated as a single transgene or in concatamers, e.g., head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al., 1992, Proc. Natl. Acad. Sci. USA 89:6232-6236. The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

[0053] When it is desired that a NHP transgene be integrated into the chromosomal site of the endogenous NHP gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous NHP gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous NHP gene (i.e., "knockout" animals).

[0054] The transgene can also be selectively introduced into a particular cell type, thus inactivating the endogenous NHP gene in only that cell type, by following, for example, the teaching of Gu et al., 1994, Science, 265:103-106. The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

[0055] Once transgenic animals have been generated, the expression of the recombinant NHP gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques that include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, in situ hybridization analysis, and RT-PCR. Samples of NHP gene-expressing tissue, may also be evaluated immunocytochemically using antibodies specific for the NHP transgene product.

5.2 NHPS and NHP Polypeptides

[0056] NHPs, NHP polypeptides, NHP peptide fragments, mutated, truncated, or deleted forms of the NHPs, and/or NHP fusion proteins can be prepared for a variety of uses. These uses include, but are not limited to, the generation of antibodies, as reagents in diagnostic assays, for the identification of other cellular gene products related to a NHP, as reagents in assays for screening for compounds that can be used as pharmaceutical reagents useful in the therapeutic treatment of mental, biological, or medical disorders and disease. Given the similarity information and expression data, the described NHPs can be targeted (by drugs, oligos, antibodies, etc.) in order to treat disease, or to therapeutically augment the efficacy of therapeutic agents.

[0057] The Sequence Listing discloses the amino acid sequences encoded by the described NHP-encoding polynucleotides. The NHPs display initiator methionines that are present in DNA sequence contexts consistent with eucaryotic translation initiation sites. The NHPs do not display consensus signal sequences, which indicates that they may be cytoplasmic or possibly nuclear proteins, however, the homology data and presence of hydrophobic domains indicates that the NHPs are probably membrane associated, or possibly secreted.

[0058] The NHP amino acid sequences of the invention include the amino acid sequences presented in the Sequence Listing as well as analogues and derivatives thereof. Further, corresponding NHP homologues from other species are encompassed by the invention. In fact, any NHP protein encoded by the NHP nucleotide sequences described above are within the scope of the invention, as are any novel polynucleotide sequences encoding all or any novel portion

of an amino acid sequence presented in the Sequence Listing. The degenerate nature of the genetic code is well-known, and, accordingly, each amino acid presented in the Sequence Listing, is generically representative of the well-known nucleic acid "triplet" codon, or in many cases codons, that can encode the amino acid. As such, as contemplated herein, the amino acid sequences presented in the Sequence Listing, when taken together with the genetic code (see, for example, Table 4-1 at page 109 of "Molecular Cell Biology", 1986, J. Darnell et al., eds., Scientific American Books, New York, N.Y., herein incorporated by reference) are generically representative of all the various permutations and combinations of nucleic acid sequences that can encode such amino acid sequences.

[0059] The invention also encompasses proteins that are functionally equivalent to the NHPs encoded by the presently described nucleotide sequences, as judged by any of a number of criteria, including, but not limited to, the ability to bind and modify a NHP substrate, or the ability to effect an identical or complementary downstream pathway, or a change in cellular metabolism (e.g., proteolytic activity, ion flux, tyrosine phosphorylation, etc.). Such functionally equivalent NHP proteins include, but are not limited to, additions or substitutions of amino acid residues within the amino acid sequence encoded by the NHP nucleotide sequences described above, but that result in a silent change, thus producing a functionally equivalent expression product. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

[0060] A variety of host-expression vector systems can be used to express the NHP nucleotide sequences of the invention. Where the NHP peptide or polypeptide can exist, or has been engineered to exist, as a soluble or secreted molecule, the soluble NHP peptide or polypeptide can be recovered from the culture media. Such expression systems also encompass engineered host cells that express a NHP, or functional equivalent, in situ. Purification or enrichment of a NHP from such expression systems can be accomplished using appropriate detergents and lipid micelles and methods well-known to those skilled in the art. However, such engineered host cells themselves may be used in situations where it is important not only to retain the structural and functional characteristics of the NHP, but to assess biological activity, e.g., in drug screening assays.

[0061] The expression systems that may be used for purposes of the invention include, but are not limited to, microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing NHP nucleotide sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing NHP nucleotide sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing NHP nucleotide sequences; plant

cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing NHP nucleotide sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing NHP nucleotide sequences and promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

[0062] In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the NHP product being expressed. For example, when a large quantity of such a protein is to be produced for the generation of pharmaceutical compositions of or containing NHP, or for raising antibodies to a NHP, vectors that direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited to, the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which a NHP coding sequence may be ligated individually into the vector in frame with the lacZ coding region so that a fusion protein is produced; pIN vectors (Inouye and Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke and Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The PGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target expression product can be released from the GST moiety.

[0063] In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) can be used as a vector to express foreign polynucleotide sequences. The virus grows in *Spodoptera frugiperda* cells. A NHP coding sequence can be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of NHP coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted sequence is expressed (e.g., see Smith et al., 1983, J. Virol. 46:584; Smith, U.S. Pat. No. 4,215,051).

[0064] In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the NHP nucleotide sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric sequence may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing a NHP product in infected hosts (e.g., see Logan and Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals may also be required for efficient transla-

tion of inserted NHP nucleotide sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire NHP gene or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of a NHP coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, may be provided. Furthermore, the initiation codon should be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bitter et al.; 1987, *Methods in Enzymol.* 153:516-544).

[0065] In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the expression product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and expression products. Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. To this end, eukaryotic host cells that possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the expression product may be used. Such mammalian host cells include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in particular, human cell lines.

[0066] For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines that stably express the NHP sequences described above can be engineered. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci, which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines that express the NHP product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the NHP product.

[0067] A number of selection systems may be used, including, but not limited to, the herpes simplex virus thymidine kinase (Wigler et al., 1977, *Cell* 11:223), hypoxanthine-guanine phosphoribosyltransferase. (Szybalska and Szybalski, 1962, *Proc. Natl. Acad. Sci. USA* 48:2026), and adenine phosphoribosyltransferase (Lowy et al., 1980, *Cell* 22:817) genes, which can be employed in tk⁻, hgprt⁻ or aprt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr,

which confers resistance to methotrexate (Wigler et al., 1980, *Proc. Natl. Acad. Sci. USA* 77:3567; O'Hare et al., 1981, *Proc. Natl. Acad. Sci. USA* 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan and Berg, 1981, *Proc. Natl. Acad. Sci. USA* 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., 1981, *J. Mol. Biol.* 150:1); and hygro, which confers resistance to hygromycin (Santerre et al., 1984, *Gene* 30:147).

[0068] Alternatively, any fusion protein can be readily purified by utilizing an antibody specific for the fusion protein being expressed. For example, an exemplary system allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:8972-8976). In this system, the sequence of interest is subcloned into a vaccinia recombination plasmid such that the sequence's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni²⁺ nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

[0069] Also encompassed by the present invention are fusion proteins that direct the NHP to a target organ and/or facilitate transport across the membrane into the cytosol. Conjugation of NHPs to antibody molecules or their Fab fragments could be used to target cells bearing a particular epitope. Attaching the appropriate signal sequence to the NHP would also transport the NHP to the desired location within the cell. Alternatively, targeting of NHP or its nucleic acid sequence might be achieved using liposome or lipid complex based delivery systems. Such technologies are described in "Liposomes: A Practical Approach", New, R.R.C., ed., Oxford University Press, New York, and in U.S. Pat. Nos. 4,594,595, 5,459,127, 5,948,767 and 6,110,490 and their respective disclosures, which are herein incorporated by reference in their entirety. Additionally embodied are novel protein constructs engineered in such a way that they facilitate transport of the NHP to the target site or desired organ, where they cross the cell membrane and/or the nucleus where the NHP can exert its functional activity. This goal may be achieved by coupling of the NHP to a cytokine or other ligand that provides targeting specificity, and/or to a protein transducing domain (see generally U.S. Provisional Patent Application Ser. Nos. 60/111,701 and 60/056,713, both of which are herein incorporated by reference, for examples of such transducing sequences) to facilitate passage across cellular membranes, and can optionally be engineered to include nuclear localization.

5.3 Antibodies to NHP Products

[0070] Antibodies that specifically recognize one or more epitopes of a NHP, or epitopes of conserved variants of a NHP, or peptide fragments of a NHP, are also encompassed by the invention. Such antibodies include, but are not limited to, polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

[0071] The antibodies of the invention can be used, for example, in the detection of NHP in a biological sample and

may, therefore, be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal amounts of NHP. Such antibodies may also be utilized in conjunction with, for example, compound screening schemes for the evaluation of the effect of test compounds on expression and/or activity of a NHP expression product. Additionally, such antibodies can be used in conjunction with gene therapy to, for example, evaluate the normal and/or engineered NHP-expressing cells prior to their introduction into the patient. Such antibodies may additionally be used as a method for the inhibition of abnormal NHP activity. Thus, such antibodies may, therefore, be utilized as part of treatment methods.

[0072] For the production of antibodies, various host animals may be immunized by injection with the NHP, a NHP peptide (e.g., one corresponding to a functional domain of a NHP), truncated NHP polypeptides (NHP in which one or more domains have been deleted), functional equivalents of the NHP, or mutated variants of the NHP. Such host animals may include, but are not limited to, pigs, rabbits, mice, goats, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including, but not limited to, Freund's adjuvant (complete and incomplete), mineral salts such as aluminum hydroxide or aluminum phosphate, chitosan, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, and potentially useful human adjuvants such as BCG (*bacille Calmette-Guerin*) and *Corynebacterium parvum*. Alternatively, the immune response could be enhanced by combination and/or coupling with molecules such as keyhole limpet hemocyanin, tetanus toxoid, diphtheria toxoid, ovalbumin, cholera toxin or fragments thereof. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of the immunized animals.

[0073] Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, can be obtained by any technique that provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique (Kohler and Milstein, 1975, *Nature* 256:495-497; and U.S. Pat. No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, *Immunology Today* 4:72; Cole et al., 1983, *Proc. Natl. Acad. Sci. USA* 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, *Monoclonal Antibodies And Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class, including IgG, IgM, IgE, IgA, and IgD, and any subclass thereof. The hybridoma producing the mAbs of this invention may be cultivated *in vitro* or *in vivo*. Production of high titers of mAbs *in vivo* makes this the presently preferred method of production.

[0074] In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, *Proc. Natl. Acad. Sci. USA*, 81:6851-6855; Neuberger et al., 1984, *Nature*, 312:604-608; Takeda et al., 1985, *Nature*, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region. Such

technologies are described in U.S. Pat. Nos. 6,075,181 and 5,877,397 and their respective disclosures, which are herein incorporated by reference in their entirety. Also encompassed by the present invention is the use of fully humanized monoclonal antibodies, as described in U.S. Pat. No. 6,150,584 and respective disclosures, which are herein incorporated by reference in their entirety.

[0075] Alternatively, techniques described for the production of single chain antibodies (U.S. Pat. No. 4,946,778; Bird, 1988, *Science* 242:423-426; Huston et al., 1988, *Proc. Natl. Acad. Sci. USA* 85:5879-5883; and Ward et al., 1989, *Nature* 341:544-546) can be adapted to produce single chain antibodies against NHP expression products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

[0076] Antibody fragments that recognize specific epitopes may be generated by known techniques. For example, such fragments include, but are not limited to: F(ab')₂ fragments, which can be produced by pepsin digestion of the antibody molecule; and Fab fragments, which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed (Huse et al., 1989, *Science*, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

[0077] Antibodies to a NHP can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" a given NHP, using techniques well-known to those skilled in the art (see, e.g., Greenspan and Bona, 1993, *FASEB J.* 7:437-444; and Nissinoff, 1991, *J. Immunol.* 147:2429-2438). For example, antibodies that bind to a NHP domain and competitively inhibit the binding of NHP to its cognate receptor/ligand can be used to generate anti-idiotypes that "mimic" the NHP and, therefore, bind, activate, or neutralize a NHP, NHP receptor, or NHP ligand. Such anti-idiotypic antibodies, or Fab fragments of such anti-idiotypes, can be used in therapeutic regimens involving a NHP mediated pathway.

[0078] Additionally given the high degree of relatedness of mammalian NHPs, the presently described knock-out mice (having never seen a NHP, and thus never been tolerized to a NHP) have a unique utility, as they can be advantageously applied to the generation of antibodies against the disclosed mammalian NHPs (i.e., a NHP will be immunogenic in NHP knock-out animals).

[0079] The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein, will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims. All cited publications, patents, and patent applications are herein incorporated by reference in their entirety.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 5

<210> SEQ ID NO 1

<211> LENGTH: 2079

<212> TYPE: DNA

<213> ORGANISM: homo sapiens

<400> SEQUENCE: 1

```
atggagaagt acgagcggat ccgagtgggtg gggagaggtg ccttcgggat tgtgcacctg    60
tgctgcgaa aggctgacca gaagctgggtg atcatcaagc agattccagt ggaacagatg    120
accaaggaag agcggcaggc agcccagaat gagtgccagg tcctcaagct gtccaaccac    180
cccaatgtca ttgagtacta cgagaacttc ctggaagaca aagcccttat gatcgccatg    240
gaatatgcac caggcggcac tctggctgag ttcattcaaa agcgtgtaa ttcctgctg    300
gaggaggaga ccatcctgca cttcttcgtg cagatcctgc ttgactgca tcatgtgcac    360
accacactca tcctgcaccg agacctcaag acccagaaca tcctgcttga caaacaccgc    420
atggtcgtca agatcgggtg tttcggcatc tccaagatcc ttagcagcaa gagcaaggcc    480
tacacggtgg tgggtacccc atgctataat tcccctgagc tgtgtgaggg caagccctac    540
aaccagaaga gtgacatctg gcccctgggc tgtgtcctct acgagctggc cagcctcaag    600
agggctttcg aggctgcgaa cttgccagca ctggtgctga agatcatgag tggcaccttt    660
gcacctatct ctgaccggta cagccctgag cttcgccagc tggtcctgag tctactcagc    720
ctggagcctg cccagcggcc accactcagc cacatcatgg cacagcccct ctgcatccgt    780
gccctcctca acctccacac cgacgtgggc agtgtccgca tgcggagggc agagaagtcc    840
gtggccccc gaaacacagg gagcaggacc accagtgtcc gctgcagagg tatccccgg    900
ggactgtgta ggccagccat cccaccacca ctgtcgtcag tgtatgcctg ggtggtggtg    960
ctgggcaccc ccctgcggct gccaatgtct aacacagagg tggccaggt ggcagctggg    1020
cgcacgcaga aagccggcgt cacgcgctct gggcgtctca tcctgtggga ggccccacc    1080
ctaggtgcag gcggaggcag tctccttctt ggggcagtgg agcagccaca gcccagttc    1140
atctcgcgtt tcctggaggg ccagtcgggy gtgacctca agcacgtggc ctgtggggac    1200
ttcttactg cctgctgac tgacagaggc atcatcatga cattcggcag cggcagcaat    1260
gggtgcctag gccatggcag cctcactgac atcagccagc ccaccattgt ggaggctttg    1320
ytgggctatg aaatggtgca ggtggcctgt ggggcctctc acgtgctggc cctgtccact    1380
gagcgagaac tatttgctg gggccgtgga gacagcggca gactggggct aggcaccagg    1440
gagtcccaca gctgccccca gcaggtgccc atgccccag gacaggaagc tcagcgagtt    1500
gtatgtggtg tcgattcctc catgatcctc actgtgctg gccaagccct agcctgtggg    1560
agcaacaggt tcaacaagct gggcctggac cacctctccc tgggggagga gcctgtccc    1620
caccagcaag tggaggaggc cctgagcttc aactactag gctctgcacc cctggaccag    1680
gagcctctgc tgagtataga cctgggcact gctcactcag ctgctgtgac tgctcgggt    1740
gattgttaca cttttggcag caatcagcac ggacagttgg gcaccaatac tcgccgagge    1800
agtcgggcac cctgtaaggt ccaaggcctt gagggcatca agatggcaat ggtagcctgt    1860
ggggatgcct tcactgtagc tattggggca gagagcgaag tgtactcttg gggcaaagg    1920
```

-continued

```

gcgcgaggtc gattgggaag gagggatgag gatgccggac tccctcggcc agtgcagttg 1980
gatgagacac acccttacac ggtgacttcc gtgtcctggt gccatggaaa caccctcctg 2040
gctgttcgat cggtcacaga tgagccggtc cccccctga 2079

```

```

<210> SEQ ID NO 2
<211> LENGTH: 692
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

```

```

<400> SEQUENCE: 2

```

```

Met Glu Lys Tyr Glu Arg Ile Arg Val Val Gly Arg Gly Ala Phe Gly
 1          5          10          15
Ile Val His Leu Cys Leu Arg Lys Ala Asp Gln Lys Leu Val Ile Ile
 20          25          30
Lys Gln Ile Pro Val Glu Gln Met Thr Lys Glu Glu Arg Gln Ala Ala
 35          40          45
Gln Asn Glu Cys Gln Val Leu Lys Leu Leu Asn His Pro Asn Val Ile
 50          55          60
Glu Tyr Tyr Glu Asn Phe Leu Glu Asp Lys Ala Leu Met Ile Ala Met
 65          70          75          80
Glu Tyr Ala Pro Gly Gly Thr Leu Ala Glu Phe Ile Gln Lys Arg Cys
 85          90          95
Asn Ser Leu Leu Glu Glu Glu Thr Ile Leu His Phe Phe Val Gln Ile
100          105          110
Leu Leu Ala Leu His His Val His Thr His Leu Ile Leu His Arg Asp
115          120          125
Leu Lys Thr Gln Asn Ile Leu Leu Asp Lys His Arg Met Val Val Lys
130          135          140
Ile Gly Asp Phe Gly Ile Ser Lys Ile Leu Ser Ser Lys Ser Lys Ala
145          150          155          160
Tyr Thr Val Val Gly Thr Pro Cys Tyr Ile Ser Pro Glu Leu Cys Glu
165          170          175
Gly Lys Pro Tyr Asn Gln Lys Ser Asp Ile Trp Ala Leu Gly Cys Val
180          185          190
Leu Tyr Glu Leu Ala Ser Leu Lys Arg Ala Phe Glu Ala Ala Asn Leu
195          200          205
Pro Ala Leu Val Leu Lys Ile Met Ser Gly Thr Phe Ala Pro Ile Ser
210          215          220
Asp Arg Tyr Ser Pro Glu Leu Arg Gln Leu Val Leu Ser Leu Leu Ser
225          230          235          240
Leu Glu Pro Ala Gln Arg Pro Pro Leu Ser His Ile Met Ala Gln Pro
245          250          255
Leu Cys Ile Arg Ala Leu Leu Asn Leu His Thr Asp Val Gly Ser Val
260          265          270
Arg Met Arg Arg Ala Glu Lys Ser Val Ala Pro Ser Asn Thr Gly Ser
275          280          285
Arg Thr Thr Ser Val Arg Cys Arg Gly Ile Pro Arg Gly Pro Val Arg
290          295          300
Pro Ala Ile Pro Pro Pro Leu Ser Ser Val Tyr Ala Trp Gly Gly Gly
305          310          315          320
Leu Gly Thr Pro Leu Arg Leu Pro Met Leu Asn Thr Glu Val Val Gln
325          330          335

```

-continued

Val Ala Ala Gly Arg Thr Gln Lys Ala Gly Val Thr Arg Ser Gly Arg
 340 345 350
 Leu Ile Leu Trp Glu Ala Pro Pro Leu Gly Ala Gly Gly Ser Leu
 355 360 365
 Leu Pro Gly Ala Val Glu Gln Pro Gln Pro Gln Phe Ile Ser Arg Phe
 370 375 380
 Leu Glu Gly Gln Ser Gly Val Thr Ile Lys His Val Ala Cys Gly Asp
 385 390 395 400
 Phe Phe Thr Ala Cys Leu Thr Asp Arg Gly Ile Ile Met Thr Phe Gly
 405 410 415
 Ser Gly Ser Asn Gly Cys Leu Gly His Gly Ser Leu Thr Asp Ile Ser
 420 425 430
 Gln Pro Thr Ile Val Glu Ala Leu Leu Gly Tyr Glu Met Val Gln Val
 435 440 445
 Ala Cys Gly Ala Ser His Val Leu Ala Leu Ser Thr Glu Arg Glu Leu
 450 455 460
 Phe Ala Trp Gly Arg Gly Asp Ser Gly Arg Leu Gly Leu Gly Thr Arg
 465 470 475 480
 Glu Ser His Ser Cys Pro Gln Gln Val Pro Met Pro Pro Gly Gln Glu
 485 490 495
 Ala Gln Arg Val Val Cys Gly Ile Asp Ser Ser Met Ile Leu Thr Val
 500 505 510
 Pro Gly Gln Ala Leu Ala Cys Gly Ser Asn Arg Phe Asn Lys Leu Gly
 515 520 525
 Leu Asp His Leu Ser Leu Gly Glu Glu Pro Val Pro His Gln Gln Val
 530 535 540
 Glu Glu Ala Leu Ser Phe Thr Leu Leu Gly Ser Ala Pro Leu Asp Gln
 545 550 555 560
 Glu Pro Leu Leu Ser Ile Asp Leu Gly Thr Ala His Ser Ala Ala Val
 565 570 575
 Thr Ala Ser Gly Asp Cys Tyr Thr Phe Gly Ser Asn Gln His Gly Gln
 580 585 590
 Leu Gly Thr Asn Thr Arg Arg Gly Ser Arg Ala Pro Cys Lys Val Gln
 595 600 605
 Gly Leu Glu Gly Ile Lys Met Ala Met Val Ala Cys Gly Asp Ala Phe
 610 615 620
 Thr Val Ala Ile Gly Ala Glu Ser Glu Val Tyr Ser Trp Gly Lys Gly
 625 630 635 640
 Ala Arg Gly Arg Leu Gly Arg Arg Asp Glu Asp Ala Gly Leu Pro Arg
 645 650 655
 Pro Val Gln Leu Asp Glu Thr His Pro Tyr Thr Val Thr Ser Val Ser
 660 665 670
 Cys Cys His Gly Asn Thr Leu Leu Ala Val Arg Ser Val Thr Asp Glu
 675 680 685
 Pro Val Pro Pro
 690

<210> SEQ ID NO 3
 <211> LENGTH: 2454
 <212> TYPE: DNA
 <213> ORGANISM: homo sapiens

-continued

<400> SEQUENCE: 3

```

atgcccgcgc ccaactccagc cccgcagccg ccgcccgcgc cggcccgcgc agccccagcc    60
tgcccggcgc gccctgcccc gggacagcaa ggcctatgtg accattctct aaaatattta    120
agctcgagaa tcacagagcg gaagctgcaa ggctcctggc tgcttgccag ccgaggggat    180
ctggagaaac cattcctggg gccgcgtggc cccgtcgtgc ccttgttctg ccctcgggat    240
ggccttcaact cagcacatcc tgagaacagc cctctgaagc ccagggtcgt gaccgtagtg    300
aagctgggtg ggcagcgcgc ccgaaagatc actctgctcc tcaacaggcg atcagtgcag    360
acgttcgagc agctcttagc tgacatctca gaagccttgg gctctcccag atggaagaat    420
gaccgtgtga gaaactgtt taacctcaag ggcagggaaa tcaggagcgt ctctgatttc    480
ttcagggag gggatgcttt catagctatg ggcaaagaac cactgacact gaagagcatt    540
caggtggctg tagaagaact gtaccccaac aaagcccggg ccctgacact ggcccagcac    600
agccgtgccc ctctccaag gctgaggagc aggtgttcta gcaaggctct gaaaggagac    660
caccgctgtg gggagaccga gacccccaaag agctgcagcg aagtgcagc atgcaaggca    720
gccatgaggc accaggggaa gatccccgag gagctttcac tagatgacag agcagaggacc    780
cagaagaagt gggggagggg gaaatgggag ccagaacca gtagcaagcc ccccagggaa    840
gccactcttg aagagaggca cgcaagggga gagaagcatc ttgggggtga gattgaaaag    900
acctcgggtg aaattatcag atgcgagaag tgcaagagag agagggagct tcagcagagc    960
ctggagcgtg agaggcttc tctggggacc agtgagctgg atatggggaa gggccaatg    1020
tatgatgtgg agaagctggt gaggaccaga agctgcagga ggtctcccga ggcaaactct    1080
gcaagtgggg aggaaggggtg gaaggggtgac agccacagga gcagccccag gaatcccact    1140
caagagctga ggagaccag caagagcatg gacaagaaag aggacagagc cccagaggat    1200
caagaaagcc atgctcaggg agcagccaag gccaaagaag accttgtgga agttcttctc    1260
gtcacagagg aggggctgag ggaggtgaa gaggacacca ggcccatgag caggagcaaa    1320
catggtggct ggctcctgag agagcaccag gcgggctttg agaagctccg caggacccga    1380
ggagaagaga aggagcgaga gaaggagaaa aagccatgta tgtctggagg cagaaggatg    1440
actctcagag atgaccaacc tgcaaagcta gaaaaggagc ccaagacgag gccagaagag    1500
aacaagccag agcggcccag cggctggaag ccacggccca tgggcatcat tgccgccaat    1560
gtggaagagc attatgagac tggccgggtc attggggatg ggaactttgc tgtcgtgaag    1620
gagtgcagac accgcgagac caggcaggcc tatgcgatga agatcattga caagtccaga    1680
ctcaagggca aggaggacat ggtggacagt gagatcttga tcattccagag cctctctcac    1740
cccaacatcg taaaattgca tgaagtctac gaaacagaca tggaaatcta cctgatcctg    1800
gagtacgtgc agggaggaga cctttttgac gccatcatag aaagtgtgaa gttcccggag    1860
cccgatgctg ccctcatgat catggactta tgcaaagccc tcgtccacat gcacgacaag    1920
agcattgtcc accgggacct caagcgggaa aaccttttgg ttcagcgaag tgaggacaaa    1980
tctactacct taaaattggc tgattttgga cttgcaaagc atgtggtgag acctatattt    2040
actgtgtgtg ggacccaac ttacgtagct cccgaaattc tttctgagaa aggttatgga    2100
ctggaggtgg acatgtgggc tgctggcgtg atcctctata tcctgctgtg tggctttccc    2160
ccattccgca gccctgagag ggaccagac gagctcttta acatcatcca gctgggccac    2220

```

-continued

```

tttgagtcc tccccctta ctgggacaat atctctgatg ctgctaaaga tctggtgagc 2280
cggttgctgg tggtagaccc caaaaagcgc tacacagctc atcaggttct tcagcacccc 2340
tggatcgaaa cagctggcaa gaccaataca gtgaaacgac agaagcaggt gtcccccagc 2400
agcgatggtc acttccggag ccagcacaag agggttgtgg agcaggtatc atag      2454

```

```

<210> SEQ ID NO 4
<211> LENGTH: 817
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

```

```

<400> SEQUENCE: 4

```

```

Met Pro Ala Ala Thr Pro Ala Pro Gln Pro Pro Pro Pro Ala Arg
 1           5           10          15
Pro Ala Pro Ala Cys Pro Ala Arg Pro Ala Pro Gly Gln Gln Gly Leu
 20          25          30
Cys Asp His Ser Leu Lys Tyr Leu Ser Ser Arg Ile Thr Glu Arg Lys
 35          40          45
Leu Gln Gly Ser Trp Leu Pro Ala Ser Arg Gly Asn Leu Glu Lys Pro
 50          55          60
Phe Leu Gly Pro Arg Gly Pro Val Val Pro Leu Phe Cys Pro Arg Asn
 65          70          75          80
Gly Leu His Ser Ala His Pro Glu Asn Ser Pro Leu Lys Pro Arg Val
 85          90          95
Val Thr Val Val Lys Leu Gly Gly Gln Arg Pro Arg Lys Ile Thr Leu
100         105         110
Leu Leu Asn Arg Arg Ser Val Gln Thr Phe Glu Gln Leu Leu Ala Asp
115         120         125
Ile Ser Glu Ala Leu Gly Ser Pro Arg Trp Lys Asn Asp Arg Val Arg
130         135         140
Lys Leu Phe Asn Leu Lys Gly Arg Glu Ile Arg Ser Val Ser Asp Phe
145         150         155         160
Phe Arg Glu Gly Asp Ala Phe Ile Ala Met Gly Lys Glu Pro Leu Thr
165         170         175
Leu Lys Ser Ile Gln Val Ala Val Glu Glu Leu Tyr Pro Asn Lys Ala
180         185         190
Arg Ala Leu Thr Leu Ala Gln His Ser Arg Ala Pro Ser Pro Arg Leu
195         200         205
Arg Ser Arg Leu Phe Ser Lys Ala Leu Lys Gly Asp His Arg Cys Gly
210         215         220
Glu Thr Glu Thr Pro Lys Ser Cys Ser Glu Val Ala Gly Cys Lys Ala
225         230         235         240
Ala Met Arg His Gln Gly Lys Ile Pro Glu Glu Leu Ser Leu Asp Asp
245         250         255
Arg Ala Arg Thr Gln Lys Lys Trp Gly Arg Gly Lys Trp Glu Pro Glu
260         265         270
Pro Ser Ser Lys Pro Pro Arg Glu Ala Thr Leu Glu Glu Arg His Ala
275         280         285
Arg Gly Glu Lys His Leu Gly Val Glu Ile Glu Lys Thr Ser Gly Glu
290         295         300
Ile Ile Arg Cys Glu Lys Cys Lys Arg Glu Arg Glu Leu Gln Gln Ser
305         310         315         320

```

-continued

Leu Glu Arg Glu Arg Leu Ser Leu Gly Thr Ser Glu Leu Asp Met Gly
 325 330 335
 Lys Gly Pro Met Tyr Asp Val Glu Lys Leu Val Arg Thr Arg Ser Cys
 340 345 350
 Arg Arg Ser Pro Glu Ala Asn Pro Ala Ser Gly Glu Glu Gly Trp Lys
 355 360 365
 Gly Asp Ser His Arg Ser Ser Pro Arg Asn Pro Thr Gln Glu Leu Arg
 370 375 380
 Arg Pro Ser Lys Ser Met Asp Lys Lys Glu Asp Arg Gly Pro Glu Asp
 385 390 395 400
 Gln Glu Ser His Ala Gln Gly Ala Ala Lys Ala Lys Lys Asp Leu Val
 405 410 415
 Glu Val Leu Pro Val Thr Glu Glu Gly Leu Arg Glu Val Lys Lys Asp
 420 425 430
 Thr Arg Pro Met Ser Arg Ser Lys His Gly Gly Trp Leu Leu Arg Glu
 435 440 445
 His Gln Ala Gly Phe Glu Lys Leu Arg Arg Thr Arg Gly Glu Glu Lys
 450 455 460
 Glu Ala Glu Lys Glu Lys Lys Pro Cys Met Ser Gly Gly Arg Arg Met
 465 470 475 480
 Thr Leu Arg Asp Asp Gln Pro Ala Lys Leu Glu Lys Glu Pro Lys Thr
 485 490 495
 Arg Pro Glu Glu Asn Lys Pro Glu Arg Pro Ser Gly Arg Lys Pro Arg
 500 505 510
 Pro Met Gly Ile Ile Ala Ala Asn Val Glu Lys His Tyr Glu Thr Gly
 515 520 525
 Arg Val Ile Gly Asp Gly Asn Phe Ala Val Val Lys Glu Cys Arg His
 530 535 540
 Arg Glu Thr Arg Gln Ala Tyr Ala Met Lys Ile Ile Asp Lys Ser Arg
 545 550 555 560
 Leu Lys Gly Lys Glu Asp Met Val Asp Ser Glu Ile Leu Ile Ile Gln
 565 570 575
 Ser Leu Ser His Pro Asn Ile Val Lys Leu His Glu Val Tyr Glu Thr
 580 585 590
 Asp Met Glu Ile Tyr Leu Ile Leu Glu Tyr Val Gln Gly Gly Asp Leu
 595 600 605
 Phe Asp Ala Ile Ile Glu Ser Val Lys Phe Pro Glu Pro Asp Ala Ala
 610 615 620
 Leu Met Ile Met Asp Leu Cys Lys Ala Leu Val His Met His Asp Lys
 625 630 635 640
 Ser Ile Val His Arg Asp Leu Lys Pro Glu Asn Leu Leu Val Gln Arg
 645 650 655
 Asn Glu Asp Lys Ser Thr Thr Leu Lys Leu Ala Asp Phe Gly Leu Ala
 660 665 670
 Lys His Val Val Arg Pro Ile Phe Thr Val Cys Gly Thr Pro Thr Tyr
 675 680 685
 Val Ala Pro Glu Ile Leu Ser Glu Lys Gly Tyr Gly Leu Glu Val Asp
 690 695 700
 Met Trp Ala Ala Gly Val Ile Leu Tyr Ile Leu Leu Cys Gly Phe Pro
 705 710 715 720
 Pro Phe Arg Ser Pro Glu Arg Asp Gln Asp Glu Leu Phe Asn Ile Ile

-continued

	725								730										735
Gln	Leu	Gly	His	Phe	Glu	Phe	Leu	Pro	Pro	Tyr	Trp	Asp	Asn	Ile	Ser				
			740					745					750						
Asp	Ala	Ala	Lys	Asp	Leu	Val	Ser	Arg	Leu	Leu	Val	Val	Asp	Pro	Lys				
			755				760					765							
Lys	Arg	Tyr	Thr	Ala	His	Gln	Val	Leu	Gln	His	Pro	Trp	Ile	Glu	Thr				
			770			775					780								
Ala	Gly	Lys	Thr	Asn	Thr	Val	Lys	Arg	Gln	Lys	Gln	Val	Ser	Pro	Ser				
					790					795					800				
Ser	Asp	Gly	His	Phe	Arg	Ser	Gln	His	Lys	Arg	Val	Val	Glu	Gln	Val				
				805					810						815				

Ser

<210> SEQ ID NO 5

<211> LENGTH: 2824

<212> TYPE: DNA

<213> ORGANISM: homo sapiens

<400> SEQUENCE: 5

```

cgggctcgtg gctgctcgtc tcgccccgcc ttccccgcgc tgctcgaccg tcgagccgcg      60
tccccgcgct gccacctctg ctccaggctc tccccgagcc cgccgccgcg ccatgccccg      120
cgccactcca gccccgcagc cgccgccgcc cccggcccgg ccagccccag cctgccccgg      180
gcgccctgcc ccgggacagc aaggcctatg tgaccattct ctaaaatatt taagctcgag      240
aatcacagag cggaagctgc aaggtcctg gctgcctgcc agccgagggg atctggagaa      300
accattcctg gggccgcgct gccccgctgt gcccttgctc tgccctcgga atggccttca      360
ctcagcacat cctgagaaca gccctctgaa gcccagggtc gtgaccgtag tgaagctggg      420
tgggcagcgc ccccgaaga tcaactctgt cctcaacagg cgatcagtgc agacgttcga      480
gcagctctta gctgacatct cagaagcctt gggctctccc agatggaaga atgaccgtgt      540
gaggaaactg tttaacctca agggcagggg aatcaggagc gtctctgatt tcttcagggg      600
aggggatgct ttcatagcta tgggcaaaga accactgaca ctgaagagca ttcagtgggc      660
tgtagaagaa ctgtacccca acaaagcccg ggccctgaca ctggcccagc acagccgtgc      720
cccttctcca aggtgagga gcaggctgtt tagcaaggct ctgaaaggag accaccgctg      780
tggggagacc gagaccccca agagctgcag cgaagttgca ggatgcaagg cagccatgag      840
gcaccagggg aagatccccg aggagcttct actagatgac agagcgagga cccagaagaa      900
gtgggggagg gggaaatggg agccagaacc cagtagcaag cccccaggg aagccactct      960
ggaagagagg cacgcaaggg gagagaagca tcttggggtg gagattgaaa agacctcggg     1020
tgaaattatc agatgcgaga agtgcaagag agagagggag cttcagcaga gcctggagcg     1080
tgagaggctt tctctgggga ccagtgcgct ggatatgggg aagggcccaa tgtatgatgt     1140
ggagaagctg gtgaggacca gaagctgcag gaggtctccc gaggcaaadc ctgcaagtgg     1200
ggaggaaggg tggaagggtg acagccacag gagcagcccc aggaatccca ctcaagagct     1260
gaggagaccc agcaagagca tggacaagaa agaggacaga ggcccagagg atcaagaaag     1320
ccatgctcag ggagcagcca aggccaaaga ggacctgtg gaagttcttc ctgtcacaga     1380
ggaggggctg agggaggtga agaaggacac caggcccatg agcaggagca aacatggtgg     1440
ctggctcctg agagagcacc aggcgggctt tgagaagctc cgcaggaccc gaggagaaga     1500

```

-continued

gaaggaggca gagaaggaga aaaagccatg tatgtctgga ggcagaagga tgactctcag	1560
agatgaccaa cctgcaaagc tagaaaagga gccaagacg aggccagaag agaacaagcc	1620
agagcggccc agcggtcgga agccacggcc catgggcac attgccgcca atgtggaaaa	1680
gcattatgag actggccggg tcattgggga tgggaacttt gctgtcgtga aggagtgcag	1740
acaccgagc accaggcagc cctatgcgat gaagatcatt gacaagtcca gactcaaggg	1800
caaggaggac atggtggaca gtgagatctt gatcatccag agcctctctc accccaacat	1860
cgtgaaattg catgaagtct acgaaacaga catggaaatc tacctgatcc tggagtacgt	1920
gcagggagga gacctttttg acgccatcat agaaagtgtg aagttcccgg agcccgatgc	1980
tgccctcatg atcatggact tatgcaaagc cctcgtccac atgcacgaca agagcattgt	2040
ccaccgggac ctcaagccgg aaaacctttt ggttcagcga aatgaggaca aatctactac	2100
cttgaattg gctgattttg gacttgcaaa gcattgtgtg agacctatat ttactgtgtg	2160
tgggaccca acttacgtag ctcccgaat tctttctgag aaaggttatg gactggaggt	2220
ggacatgtg gctgctggcg tgatcctcta tatcctgctg tgtggctttc cccattccg	2280
cagccctgag agggaccagc acgagctctt taacatcatc cagctgggccc actttgagtt	2340
cctccccct tactgggaca atatctctga tgctgctaaa gatctggtga gccggttgct	2400
ggtggtagac ccaaaaaagc gctacacagc tcatcaggtt cttcagcacc cctggatcga	2460
aacagctggc aagaccaata cagtgaacg acagaagcag gtgtccccca gcagcgatgg	2520
tcacttccgg agccagcaca agagggttgt ggagcaggtg tcatagtcac caccttgga	2580
atctgtccag cccccagttc tgctcaagga cagagaaaag gatagaagtt tgagagaaaa	2640
acaatgaaag aggcttcttc acataattgg tgaatcagag ggagagacac tgagtatatt	2700
ttaaagcata ttaaaaaat taagtcaatg ttaaatgtca caacatattt ttagatttgt	2760
atatttaaag cctttaatac atttttgggg ggtaagcatt gtcacagtg aggaattttg	2820
gtaa	2824

1. (canceled)
2. (canceled)
3. (canceled)
4. (canceled)
5. (canceled)
6. An antibody that specifically recognizes the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4.

7. A method for producing a polypeptide comprising SEQ ID NO:2 or SEQ ID NO:4, comprising culturing a host cell comprising a recombinant expression vector comprising a nucleotide sequence encoding the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4 under conditions sufficient for the production of said polypeptide, and recovering said polypeptide from the host cell culture.

* * * * *