

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2016265929 B2

(54) Title
Methods of agent delivery into eggs of egg-producing aquatic animals

(51) International Patent Classification(s)
A01K 67/027 (2006.01) **C12N 15/87** (2006.01)

(21) Application No: **2016265929** (22) Date of Filing: **2016.05.17**

(87) WIPO No: **WO16/187198**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/163,511	2015.05.19	US
62/274,958	2016.01.05	US

(43) Publication Date: **2016.11.24**

(44) Accepted Journal Date: **2022.03.31**

(71) Applicant(s)
University of Maryland Baltimore County

(72) Inventor(s)
Wong, Ten Tsao;Zohar, Yonathan

(74) Agent / Attorney
IP Gateway Patent and Trade Mark Attorneys Pty Ltd, PO Box 1321, SPRINGWOOD, QLD, 4127, AU

(56) Related Art
EP 2535404 A1

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2016/187198 A3

(43) International Publication Date
24 November 2016 (24.11.2016)

WIPO | PCT

(51) International Patent Classification:
A01K 67/027 (2006.01) *C12N 15/87* (2006.01)

(74) Agent: GRANT, Mary, B.; Hultquist IP, PO Box 14329, Research Triangle Park, NC 27709 (US).

(21) International Application Number:
PCT/US2016/032873

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:
17 May 2016 (17.05.2016)

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,

(25) Filing Language: English

(26) Publication Language: English


(30) Priority Data:
62/163,511 19 May 2015 (19.05.2015) US
62/274,958 5 January 2016 (05.01.2016) US

(71) Applicant: UNIVERSITY OF MARYLAND BALTIMORE COUNTY [US/US]; 1000 Hilltop Circle, Baltimore, MD 21250 (US).

(72) Inventors: WONG, Ten, Tsao; 9 W Main Blvd., Timonium, MD 21093 (US). ZOHAR, Yonathan; 2712 Old Court Road, Baltimore, MD 21208 (US).

[Continued on next page]

(54) Title: METHODS OF AGENT DELIVERY INTO EGGS OF EGG-PRODUCING AQUATIC ANIMALS



WO 2016/187198 A3

FIG. 3



SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, — *with sequence listing part of description (Rule 5.2(a))*
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

(88) Date of publication of the international search report:
12 January 2017

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

METHODS OF AGENT DELIVERY INTO EGGS AND EMBRYOS OF EGG-PRODUCING AQUATIC ANIMALS FOR DRUG SCREENING, AGENT TOXICITY ASSAY AND PRODUCTION OF INFERTILE FISH

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims benefit of priority to U.S. Provisional Application No. 62/163,511 filed May 19, 2015 and U.S. Provisional Application No. 62/274,958 filed January 5, 2016. The disclosures of such related provisional applications are hereby incorporated herein by reference in their respective entireties.

TECHNICAL FIELD

[0002] The present disclosure relates to methods for agent delivery into the eggs and embryos of egg-producing aquatic animals. The subject matter of the invention includes methods used to enhance chorion permeability and administer agents, e.g., large molecules, into the eggs and embryos. The present disclosure also relates to methods for drug and bioactive agent screening in safety and toxicity assays for biotech and pharmaceutical industries, and the production of reproductively sterile fish for aquaculture, the aquarium trade, and control of invasive species.

[0003] The disclosure further includes methods used to administer Morpholino oligomer (MO) into eggs and embryos of egg-producing aquatic animals which lead to the failure of fertile gonad development and to sterile adult fish.

BACKGROUND ART

[0004] The reference to prior art in this specification is not and should not be taken as an acknowledgment or any form of suggestion that the referenced prior art forms part of the common general knowledge in Australia or in any other country.

[0005] Aquaculture is becoming increasingly important to resolve the current and projected global shortfalls in aquatic foods and seafood availability. As the shift in dependence from fisheries' harvests to artificially propagated aquatic species continues, optimization of aquaculture methods is increasingly necessary to maximize food production and minimize ecological impact, thereby achieving long-term environmental sustainability of our seafood supplies.

[0006] Sterilization (induced infertility) of farmed fish and other egg-producing aquatic animals enhances their growth rate by increasing the conversion of food energy to muscle growth,

instead of gonadal development. In addition, if escaped from aquaculture operations to the environment, reproductively sterile farmed fish and other egg-producing aquatic animals, including domesticated, non-native or genetically modified species, will not be able to reproduce or inter-breed with wild population. This will assist biological containment and prevent genetic contamination of wild populations and/or the establishment in the wild of domestic, non-native or genetically modified farmed fish and other egg-producing aquatic animals.

[0007] Additionally, reproductive sterilization of fish and other egg-producing aquatic animals prevents unauthorized breeding and sale of patented, or otherwise protected, genetically selected or modified fish and other egg-producing aquatic animals.

[0008] Furthermore, aquatic animals, such as zebrafish, have provided models for evaluation of drugs or bioactive agents of interest. However, one of the major challenges in the use of eggs and embryos of these aquatic animals is the low permeability of the chorion or egg envelope which prevents such agents from traversing the chorion and reaching the embryo.

[0009] Therefore, methods are needed to enhance the permeability of the chorion of the eggs of aquatic animals and allow candidate drugs, bioactive agents and large molecules to traverse the chorion and to reach the embryo. In addition, methods are needed for reproductive sterilization of egg-producing aquatic animals.

DEFINITION

[0010] In the present description and claims, the term “comprising” shall be understood to have a broad meaning similar to the term “including” and will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps. This definition also applies to variations on the term “comprising” such as “comprise” and “comprises”.

SUMMARY

[0011] It has been discovered that certain compounds, e.g., guanidine, arginine, and their derivatives, dimers, trimers or polymers, are able to effectively enhance permeability of the chorion of eggs of aquatic animals, which allows agents, such as candidate drugs, potentially bioactive agents and/or large molecules, to traverse the chorion of such eggs of egg-producing aquatic animals and reach the embryos.

[0012] One aspect of the disclosure relates to a method of delivering at least one agent into egg(s) from an egg-producing aquatic animal, said method comprising contacting the egg(s)

with the at least one agent in the presence of a guanidine-containing compound that is effective to enhance permeability of the chorion of the egg(s), wherein the egg(s) are contacted with the at least one agent in an immersion medium comprising the at least one agent and the guanidine-containing compound and wherein said guanidine-containing compound is arginine or a salt thereof, guanidine or a salt thereof or a dimer, trimer or polymer of guanidine or arginine and wherein said polymer is not a dendrimeric oligoguanidine. The aquatic animal is suitably a fish.

[0013] Another aspect of the disclosure relates to a method of delivering at least one agent into fertilized or unfertilized egg(s) from an egg-producing aquatic animal, the method comprising contacting the fertilized or unfertilized egg(s) with the at least one agent in the presence of a guanidine-containing compound that is effective to enhance permeability of the chorion of the egg(s). Such methods may be used in screening methods for agents of interest. The egg is suitably a fish egg.

[0014] Thus, another aspect of the disclosure relates to a method of conducting drug screening and compound toxicity assays using aquatic animals, the method comprising contacting egg(s) from an aquatic animal with at least one agent comprising a drug, a drug candidate, a toxic compound or a toxic compound candidate in the presence of a guanidine-containing compound that is effective to enhance permeability of the chorion of the egg(s), the method further comprising a response in the aquatic animal to the agent.

[0015] A further aspect of the disclosure relates to a method of enhancing permeability of the chorion of egg(s) from an egg-producing aquatic animal, such as a fish, comprising contacting fertilized or unfertilized egg(s) from the egg-producing aquatic animal with a guanidine-containing compound capable of enhancing the permeability of the chorion of the egg(s).

[0016] The disclosure further relates to methods of producing populations of sterile egg-producing aquatic animals, wherein the sterilization methods include disruption of primordial germ cell migration and/or development in each treated individual without detrimentally affecting other characteristics of a normal animal.

[0017] A yet further aspect of the disclosure relates to a method of producing reproductively sterile egg-producing aquatic animals, said method comprising contacting fertilized egg(s) with anti-sense Morpholino oligomer selected from oligomer sequences comprising 12 bases that is effective to transfet the egg(s) and render individual(s) produced therefrom reproductively sterile.

[0018] Yet another aspect of the disclosure relates to a method of producing reproductively sterile egg-producing aquatic animals, said method comprising contacting fertilized egg(s)

immediately after fertilization with anti-sense Morpholino oligomer that is effective to transfect the egg(s) and render individual(s) produced therefrom reproductively sterile.

[0019] Other aspects, features and advantages of the invention will be more fully apparent from the ensuing disclosure and appended claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020] FIG. 1 is a flow chart diagram of the drug screening or agent toxicity assays performed according to embodiments of the disclosure that use guanidine-containing compound(s) (GCCs) to enhance chorion permeability.

[0021] FIG. 2 is a flow chart diagram of the production of reproductively sterile fish obtained according to embodiments of the disclosure that use guanidine-containing compound(s) (GCCs) to enhance chorion permeability.

[0022] FIG. 3 is flow chart diagram of the production of reproductively sterile fish obtained according to embodiments of the disclosure.

[0023] FIG. 4 is a fluorescence photomicrograph showing the effects of the administration to fish embryos of fluorescein-dextran using the GCCs arginine, guanidine-HCl and guanidine thiocyanate.

[0024] FIG. 5 is a photomicrograph showing the effects of the administration to fish embryos of poly-arginine/9mer. FIG. 5 shows bright field photographic results that high concentration of poly-arginine/9 mer (100 μ M) generated uncharacterized aggregates within 1 hour of incubation and caused chorion lysis after 18 hours incubation, which were not seen in the 10 μ M poly-arginine/9 mer treated group.

[0025] FIG. 6A-6D are photomicrographs showing the effects of the administration of fluorescein-dextran alone and in combination with poly-arginine to fish embryos. FIG. 6A shows bright field photographic results with poly-arginine and fluorescein-dextran; FIG. 6B shows bright field photographic results with fluorescein-dextran alone. FIG. 6C shows the fluorescently photographic results with poly-arginine and fluorescein-dextran; FIG. 6D shows the fluorescently photographic results with fluorescein-dextran alone.

[0026] FIG. 7A-7G are photomicrographs showing salmonids' *dnd*-MO-Vivo induced sterility in rainbow trout. Examination of gonadal tissue show (FIG. 7A) well-developed testis of untreated male fish; (FIG. 7B) well-developed ovary of untreated female fish; (FIG. 7C) the gonads of salmonids' *dnd*-MO-Vivo treated fish developed into a thin filament-like tissue; (FIG. 7D) a photomicrograph of dissected gonads; (FIG. 7E) the active spermatogenesis of the testis of

an untreated male fish; (FIG. 7F) a well-developed ovary of an untreated control female with oocytes at different developmental stages; (FIG. 7G) histological examinations of gonadal tissue show the gonad of a salmonids' *dnd*-MO-Vivo treated fish appears to be under-developed without advanced gonadal structure or germ cells.

[0027] FIG. 8A-8D are photomicrographs showing that salmonids' *dnd*-MO or *dnd*-MO-Vivo treatment blocked gonadal development in Atlantic salmon. In 10-month-old Atlantic salmon, (FIG. 8A) a developing testis of untreated male; (FIG. 8B) a developing ovary of untreated female; (FIG. 8C) the gonads of *dnd*-MO-Vivo treated or *dnd*-MO treated fish developed into a thin filament-like tissue; (FIG. 8D) a photomicrograph of dissected gonads from (A), (B) and (C).

DETAILED DESCRIPTION OF THE DISCLOSURE

[0028] In one aspect, the disclosure relates to a method of enhancing the permeability of the chorion of egg(s) from an egg-producing aquatic animal comprising contacting fertilized or unfertilized egg(s) from the egg-producing aquatic animal with a guanidine-containing compound capable of enhancing the permeability of the chorion of the egg(s). Such method allows for delivery of agents of interest to the embryo by enhancing the permeability of the chorion sufficiently to allow the agent to access to the interior of the egg.

[0029] Delivery of agents into fish or other egg-producing aquatic animals has traditionally been achieved via the feed, injection, or immersion of embryos or individuals in an agent of interest. Injection of stock, however, is not practical in large-scale commercial aquaculture operations. In addition, use of immersion treatment of fertilized and water-activated eggs has been limited due to low permeability of the chorion of the egg, a thick acellular multi-layer coat, also known as the egg envelope, composed mainly of proteins and glycoproteins. Typically, in immersion treatment of fish or other egg-producing aquatic animal embryos, agents of interest, by way of example, such as large molecular compounds, are not able to traverse the chorion and reach the embryo.

[0030] After ovulation/spawning and prior to fertilization and water-activation, eggs have a permeable and perforated chorion (or outermost coat) that allows for entry of water and substances into the unfertilized eggs through the pores or the micropyle, a small canal in the chorion of the egg allowing for the sperm to penetrate the egg for fertilization. Following fertilization and water-activation, however, the chorion becomes sealed and the egg is rendered

substantially impermeable, preventing further uptake of substances or water from the environment.

[0031] The methods of the disclosure provide for enhancing the permeability of the chorion of egg(s) of an egg-producing aquatic animal, either a fertilized or an unfertilized egg, such that delivery of at least one agent into egg(s) from an egg-producing aquatic animal may be achieved. Such methods comprise contacting fertilized or unfertilized egg(s) from the egg-producing aquatic animal with at least one agent in the presence of a guanidine-containing compound (GCC) capable of enhancing the permeability of the chorion of the egg(s).

[0032] The contacting of fertilized or unfertilized egg(s) comprises chorionic transfection of the egg(s).


[0033] The guanidine-containing compound in aspects of the disclosure may be chosen from guanidine, guanidine derivatives, guanidine dimers, trimers or polymers, e.g. dendrimers, or salts thereof, or arginine, arginine derivatives, arginine dimers, trimers or polymers, e.g. dendrimers, or salts thereof containing a guanidine moiety, and mixtures thereof.

[0034] The guanidine-containing compounds suitable for use in the disclosed methods are guanidine-containing compounds effective to enhance the permeability of the chorion of egg(s) of aquatic animals. As used herein "effective to enhance the permeability of the chorion" means that after an egg of an aquatic animal is contacted with the guanidine-containing compound, the chorion of such egg is more permeable than the chorion of an egg of such aquatic animal not contacted with the guanidine-containing compound. By way of example, the permeability of the chorion of an egg of an aquatic animal after contact with a guanidine-containing compound at time T1 would be more permeable than the chorion of an egg of the same aquatic animal not contacted with the guanidine-containing compound at time T1. The permeability may be determined by methods known to those in the art; see, for example, Hagedorn, M., et al. (1997) "Water distribution and permeability of zebrafish embryos, *Brachydanio rerio*" *J Exp Zool* 278, 356-371; and Kais, B., et al., (2013) "DMSO modifies the permeability of the zebrafish (*Danio rerio*) chorion-implications for the fish embryo test (FET)," *Aquat Toxicol*, 140-141: p. 229-38.

[0035] As used herein, the term "derivatives" is intended to mean derivatives comprising the same functional structure as the compound they are referring to, and that have similar properties, e.g., are effective to enhance the permeability of the chorion of egg(s) of aquatic animals.

[0036] As used herein "guanidine" means any compound comprising in its chemical formula at least one carbon atom doubly bonded to a nitrogen atom and singly bonded to two other nitrogen atoms and includes salts of such compounds.

[0037] The guanidine-containing compound, in one aspect, may be selected from the compounds of general formula (A) below:



(A)


wherein:

R1, R2, R3, R4 and R5 represent, independently:


a hydrogen atom, nitro, amino, cyano, phenyl, cyclohexyl, benzyl, or a linear or branched C₁-C₄ lower alkyl or C₁-C₄ alkenyl radical, optionally substituted with one or two radicals chosen from: hydroxyl, amino, dimethylamino, methoxy, ethoxy, carboxyl, carboxamide, N-methylcarboxamide or SO₃H, or salts thereof.

[0038] Examples of guanidine-containing compounds include arginine, guanidine-HCl, guanidine-thiocyanate, guanidine-acetate, guanidine-carbonate, guanidine-nitrate, guanidine-sulfate, guanidine-bicarbonates and guanidine-hydrobromides.

[0039] In other aspects, the guanidine-containing compound is a guanidine-containing dimer, trimer or polymer. In one aspect, the guanidine-containing compound may be selected from the compounds of general formulas (I) to (IV) below:



R: Variable group : Variable backbone linker $n \geq 2$; $x \geq 1$



R: Variable group : Variable backbone linker $n \geq 2$; $x \geq 1$

[0040] The variable group "R" may be selected from a hydrogen atom, nitro, amino, cyano, phenyl, cyclohexyl, benzyl, or a linear or branched C₁-C₄ lower alkyl or C₁-C₄ alkenyl radical, optionally substituted with one or two radicals chosen from: hydroxyl, amino, dimethylamino, methoxy, ethoxy, carboxyl, carboxamide, N-methylcarboxamide or SO₃H, or salts thereof. The variable backbone linker may be selected from an amino acid, nucleotide, phosphoramidate, glycol, polyethylene glycol, nitro, amino, cyano, phenyl, cyclohexyl, benzyl, or a linear or branched C₁-C₄ lower alkyl. By way of example, polyarginine/9-arginine is a compound of formula I with n=9.

[0041] The guanidine-containing compounds may be used in methods of delivering at least one agent into fertilized or unfertilized egg(s) from an egg-producing aquatic animal, said method

comprising contacting the fertilized or unfertilized egg(s) with the at least one agent in the presence of a guanidine-containing compound that is effective to enhance permeability of the chorion of the egg(s).

[0042] The agent may be any compound, drug or drug candidate, bioactive agent or potentially bioactive agent, pharmaceutical, chemically active substance, therapeutic substance, or other test substance. These include, without limitation, forms such as uncharged molecules, molecular complexes, salts, ethers, esters, amides, etc.

[0043] By way of example, the agent or agents may be placed in an immersion medium containing the fertilized or unfertilized egg(s) and the guanidine-containing compound(s) such that the agent(s) is in the presence of the guanidine-containing compound.

[0044] The guanidine-containing compounds are either covalently bound or not covalently bound to the agent(s) in the methods of the disclosure. It has been discovered that the guanidine-containing compounds are capable of enhancing the permeability of the chorion of egg(s) from an egg-producing aquatic animal and allowing for delivery of agents to the embryo even where the guanidine-containing compounds are not covalently bound to or conjugated with the agent.

[0045] The guanidine-containing compound according to some aspects of the disclosure can be a dendrimer, e.g. an octaguanidine dendrimer comprising a triazine core moiety e.g., of a type as described in U.S. Patent No. 7,935,816, also known in the art as "Vivo". The disclosure of U.S. Patent No. 7,935,816 is hereby incorporated herein by reference in its respective entirety. In certain embodiments of the disclosure where the guanidine-containing compound is an octaguanidine dendrimer comprising a triazine core moiety, the guanidine-containing compound is not covalently bound to or conjugated with the agent(s) of interest.

[0046] The immersion medium in embodiments is an aqueous medium. The immersion medium may be any such medium known to those of skill in the art for use with eggs from aquatic animals. The immersion media, for example, may be an aqueous medium which may further comprise fresh water, brackish water, sea water, fish ovarian fluid or fertilization diluent that contains salt, Tris (pH 7-9), glycine, and/or 0 to 30% of serum and protease inhibitors such as aprotinin or leupeptin.

[0047] The concentration of the guanidine-containing compound in the immersion medium is an amount sufficient for enhancing the permeability of the chorion of the eggs in the immersion medium. By way of example, the concentration of the guanidine-containing compound may be in a range of from about 1 to about 80,000 μ M, preferably in the range of from about 20 to about 40,000 μ M, and more preferably in the range of from about 40 to about 20,000 μ M.

[0048] The discovery that guanidine-containing compounds according to the disclosure are able to effectively enhance permeability of the chorion of eggs of aquatic animals provides the ability to efficiently deliver agents to the eggs and embryos of aquatic animals. The delivery of agents further allows for use of aquatic animals in models for drug or bioactive agent screening or in assays for evaluation of toxicity of a test agent. By way of example, zebrafish have been used in phenotypic models for drug or bioactive agent screening. Accordingly, the methods disclosed herein may be used, for example, in drug or bioactive agent screening methods, assays for assessing drug or bioactive agent safety or toxicity, and methods for evaluating a test agent for biological response.

[0049] FIG. 1 illustrates one example of the application of guanidine-containing compounds in drug screening or agent toxicity assays. As shown, fertilized eggs are placed in an agent bath. One group of eggs is immersed in the agent bath without guanidine-containing compounds therein and one group of eggs is immersed in the agent bath in the presence of guanidine-containing compounds. The therapeutic or toxicity responses of the eggs immersed in compound with GCCs can be monitored and/or evaluated.

[0050] Aspects of the disclosure thus encompass a method of screening a test agent such as a drug or bioactive agent, e.g. antibody, protein, peptide, RNA or DNA, comprising contacting egg(s) from an egg-producing aquatic animal with the at least one agent in the presence of a guanidine-containing compound that is effective to enhance the permeability of the chorion of the egg(s) and identifying a response, e.g., a physiological response. By way of example, the screening method may be used to evaluate the test agent, evaluate the effects of the test agent on the fish embryo, evaluate the toxicity of the test agent, etc.

[0051] The methods of disclosure additionally encompass use of guanidine-containing compounds in compound safety and toxicity assays wherein a test agent is placed in contact with egg(s)/embryo(s) from an egg-producing aquatic animal in the presence of a guanidine-containing compound that is effective to enhance the permeability of the chorion of the egg(s)/embryo(s) and identifying a response, e.g., a disease rescue or a physiological response (abnormality and mortality) and the response to the test agent is monitored and evaluated.

[0052] The disclosure further contemplates methods of producing reproductively sterile egg-producing aquatic animals involving contacting eggs in immersion medium with selected agents resulting in reproductively sterile individuals. The sterilization methods comprise the disruption of gonadal development in the embryo. The present disclosure also relates to methods of preventing interbreeding between domesticated, non-native and genetically modified farmed fish/other egg-producing aquatic animals and their wild stocks, as well as to the establishment of

such aquacultured fish and other egg-producing aquatic animals in the wild. In addition, the disclosed methods may be employed to enable prevention of genetic contamination of a wild population by farmed fish and other egg-producing aquatic animals.

[0053] As defined herein, "sterilizing" egg-producing aquatic animals is understood to mean rendering an individual unable to sexually reproduce. Reproductively sterile egg-producing aquatic animals are defined as individuals that are unable to reach sexual maturity or to reproduce when reaching the age of sexual maturity.

[0054] The disclosure thus provides a method of producing reproductively sterile egg-producing aquatic animals, the method comprising contacting egg(s) with an agent that is effective to transfect the egg(s) and render individual(s) produced therefrom reproductively sterile. In some aspects, the agent is contacted with the eggs in the presence of a guanidine-containing compound. In some aspects, the agent is an anti-sense Morpholino oligomer.

[0055] The methods of producing a reproductively sterile fish and other egg-producing aquatic animals include administration of agents to their eggs to disrupt primordial germ cell (PGC) development, migration and colonization in the gonad of the embryo, which results in failure of gonad development and/or failure of full and proper gonadal functioning at the cellular or tissular level, and ultimately the generation of sterile fish and other egg-producing aquatic animals.

[0056] PGCs are a population of cells in the fish embryo that are precursors of the gametes of the adult fish and other egg-producing aquatic animals. The PGCs are produced during the very early stages of embryonic development. At later stages of embryonic development, the PGCs migrate through the embryo from their original location to the area of the gonadal precursors. At the end of their migration, the PGCs enter the developing gonads, colonize the tissue and start the process of gametogenesis, leading to mature gonads in the adult fish and other egg-producing aquatic animals.

[0057] The methods of the disclosure allow generation of reproductively sterile (infertile) egg-producing aquatic animals. The sterilization strategy will specifically disrupt gonad development in the individuals without detrimentally affecting any other characteristics resulting in the production of completely normal but reproductively sterile egg-producing aquatic animals.

[0058] Thus, in various embodiments, the disclosure provides a method to efficiently administer agents into embryos by contacting eggs with agents suitable to disrupt PGC development, migration and/or survival in large numbers of embryos, resulting in large-scale production of reproductively sterile adult fish and other egg-producing aquatic animals. The methods of the

disclosure are also applicable to single embryos in smaller scale production of reproductively sterile adult fish and other egg-producing aquatic animals.

[0059] In some aspects, the agent administered to render the individual(s) produced therefrom reproductively sterile is delivered into the eggs in the presence of a guanidine-containing compound according to this disclosure. FIG. 2 illustrates an example of the application of guanidine-containing compounds in the production of infertile fish. As shown in FIG. 2, fertilized eggs are either immersed in an agent bath containing guanidine-containing compound(s) or not containing any GCCs. The eggs are immersed in the bath for 24 hours and then washed. The eggs immersed in the bath with the GCC(s) resulted in infertile fish due to mis-migration of the PGCs, whereas the eggs immersed in the bath without the GCCs resulted in fertile fish.

[0060] The agents for use in the methods of the disclosure may include agents known to disrupt PGC development, migration and/or survival which are capable of entering the chorion of a fertilized or unfertilized egg or are provided in the presence of a compound that assists in the entering of the egg. In one aspect, such agent may be an antisense Morpholino oligomer capable of the disruption of PGC development and capable of traversing the chorion of eggs. Thus, in some embodiments, antisense Morpholino oligomer useful in the methods of the disclosure is anti-sense Morpholino oligomer that is effective to transfect the egg(s) and render individual(s) produced therefrom reproductively sterile.

[0061] Antisense Morpholino oligomer is used to transiently silence gene expression by either blocking translation or RNA splicing that is an essential step to generate mRNA. Specific antisense Morpholino oligomers can be identified to transiently block or suppress the expression of genes that are essential for embryonic germ cell development including but not limited to *dead end (dnd)*, *nanos*, *vasa*, *gnrh* or *fsh* receptor which results in the failure of gonadal development and ultimately generates sterile fish and other egg-producing aquatic animals.

[0062] Thus, in one aspect of the disclosure, a method of producing reproductively sterile egg-producing aquatic animals is provided comprising contacting fertilized or unfertilized egg(s) with anti-sense Morpholino oligomer that is effective to transfect the egg(s) and render individual(s) produced therefrom reproductively sterile. The contacting may in some aspects include the presence of a guanidine-containing compounds according to the disclosure herein. The contacting comprises chorionic transfection of the egg(s). The contacting may be at the time of water-activation.

[0063] In such aspect, the disclosure relates to methods of producing reproductively sterile egg-producing aquatic animals by administration of effective Morpholino oligomers to eggs in order

to disrupt primordial germ cell (PGC) development, and migration to, and colonization in, the gonad of the embryo, which results in the failure of gonad development and/or full and proper gonadal functioning at the cellular or tissular level, and ultimately the generation of sterile fish.

[0064] *Dead end (dnd)* is a vertebrate-specific component of the germ plasm and germ-cell granules that is essential for germ cell development. The *dnd* gene is specifically expressed in germ plasm and primordial germ cells. Since *dnd* is considered essential for normal migration and survival of PGCs, embryos devoid of this protein develop to become sterile adults.

[0065] The disclosed methods are useful for the production of reproductively sterile fish and other egg-producing aquatic animals for aquaculture, the aquarium trade, and control of invasive species. In one aspect, the methods include disruption of gonadal development through the administration of antisense Morpholino oligomer against *dead end* mRNA (*dnd*-MO) or other genes that are essential to gonadal development including, but not limited to, *nanos*, *vasa*, *gnrh* or *fsh* receptor, to fertilized egg(s). The action of *dnd*-MO or other antisense Morpholino oligomer against genes that are essential to gonadal development leads to the failure of fertile gonad development and to sterile adult fish.

[0066] In embodiments, the *dnd*-MO is able to transiently suppress expression of Dead end protein that is essential for embryonic germ cell development.

[0067] The present disclosure also relates to specific Morpholino oligomer sequences for use in methods for suppression of expression a *dead end* gene in a fish.

[0068] FIG. 3 is a flow chart for production of reproductively sterile fish by the administration of specific Morpholino oligomers against the *dead end* mRNAs of fish to disrupt primordial germ cell (PGC) development, which results in the failure of gonad development, and ultimately the generation of sterile fish. When eggs are not treated with Morpholino oligomers, they may become fertile broodstock.

[0069] As shown in FIG. 3, eggs of fish (e.g. from salmonids, moronids or cichlids) are contacted with Morpholino oligomers against the *dead end* genes of the relevant fish species. FIG. 3 illustrates the administration of Morpholino oligomers against the *dead end* genes during water activation and water hardening. Alternatively, the eggs may be given no treatment. As shown, where the eggs are contacted with Morpholino oligomers, oligomers effect the suppression of or blocking of Dead end protein translation, resulting in the disruption of PCG development. The adult fish are consequently infertile, since there is no fertile gonadal development. When normal PGC development is permitted, the fish will have normal fertile gonadal development and the fish may be used as broodstock.

[0070] Thus, in embodiments, the antisense Morpholino oligomers are capable of effectively suppressing expression of at least one *dead end* gene, e.g. of at least one *dead end* gene of Salmonidae *dead end* gene, Moronidae *dead end* gene or Cichlidae *dead end* gene.

[0071] In another aspect, the disclosure relates to specific antisense Morpholino oligomers that are able to transiently and effectively suppress the translation of *dead end*, an essential gene for germ cell survival, and specifically disrupt gonadal development resulting in the production of infertile fish, e.g., salmonids (salmons and trouts), moronids (basses) and cichlids (tilapias and ornamental cichlids).

[0072] In another aspect, the disclosure relates to the identification of short sequences of Morpholino oligomer that can be more easily up-taken by eggs to reach target cells. Such short sequences preferably are 12 bases or 18 bases instead of 25 bases, as disclosed in WO 2015/073819, incorporated herein by reference. Such Morpholino oligomer sequences have been found to more efficiently transiently block the expression of specific genes that are essential for germ cell development in cold water species such as Atlantic salmon. The low egg hatching temperature is believed to make short Morpholino oligomers more stable against its target mRNA to transiently block the expression of specific genes.

[0073] The disclosure further relates to a specific Morpholino oligomer, 5'-ACGCTCCTCCAT-3' (SEQ ID NO: 1) and its variants, e.g., 5'-ACTTGAACGCTCCTCCAT-3' (SEQ ID NO: 2), that are able to transiently and effectively suppress the expression of Salmonidae *dead end* gene, which results in the failure of gonad development and/or the failure of full and proper gonadal functioning, and ultimately the generation of sterile salmonids. Accordingly, the methods of the disclosure are applicable to all salmonids, including, but not limited to, Atlantic, coho, chinook, chum, sockeye, pink and masu salmons, rainbow, brook and brown trouts, common and Arctic grayling, and Arctic char, among others.

[0074] The Morpholino oligomers may be variants of the listed sequences. These variations may include but not be limited to other modified nucleic acids and other Morpholino oligomers that cover the whole or partial sequences listed above. Particularly included are antisense oligomers that comprise, consist essentially of, or consist of, one or more of SEQ ID NO:1 and SEQ ID NO: 2. Also included are variants of these antisense oligomers, including variant oligomers having 80%, 85%, 90%, 95%, 97%, 98%, or 99% (including all integers in between) sequence identity or sequence homology to any one of SEQ ID NO:1 and SEQ ID NO: 2., and/or variants that differ from these sequences by about 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides, preferably variants that are effective in reproductively sterilizing fish upon contact with fish egg(s).

[0075] Another aspect of the invention includes antisense Morpholino oligomers which consist of SEQ ID NO: 1 and variants thereof. Yet another aspect of the invention includes transfected eggs of an egg-producing aquatic animal comprising an anti-sense Morpholino oligomer which consists of the nucleotide sequence of SEQ ID NO: 1 or a variant thereof.

[0076] Variants that are effective in reproductively sterilizing fish upon contact with fish egg(s) as used in such context means use of the variants in the methods disclosed herein will result in reproductively sterile fish. In embodiments, the variants will effectively suppress the expression of a *dead end* gene of interest.

[0077] According to embodiments of the methods of the disclosure, fertilized or unfertilized fish egg(s) (one or more eggs) are immersed in an immersion medium comprising an antisense Morpholino oligomer effective to suppress expression of a *dead end* gene in the fish of interest. Such immersion is preferably at the start of water activation of the egg(s). The concentration of the antisense Morpholino oligomer in the immersion medium should be sufficient to allow the antisense Morpholino oligomer to traverse the chorion of the fish egg(s), effectively transfecting the egg(s). In embodiments, such concentration will typically be about 1 to about 200 μ M, more preferably, about 10 to about 100 μ M, and still more preferably, about 20 to about 60 μ M.

[0078] The immersion medium, by way of example, is typically an aqueous medium which may further comprise fresh water, brackish water, sea water, fish ovarian fluid or fertilization diluent that contains salt, Tris (pH 7-9), Glycine, and/or 0 to 30% of serum and protease inhibitors such as aprotinin or leupeptin.

[0079] In accordance with the disclosure, the immersion medium may further comprise a guanidine-containing compound as detailed above capable of enhancing the permeability of the chorion of the eggs(s).


[0080] Although the time required for the immersion of the fertilized eggs to result in satisfactory sterilization of the fish or other egg-producing aquatic animals will depend on the species of fish and other egg-producing aquatic animals, typically the fish or other eggs will be immersed in the immersion medium containing a Morpholino oligomer and, optionally, a guanidine-containing compound for about 2 to about 96 hours, more preferably for about 4 to about 72 hours, and still more preferably from about 5 to about 48 hours.

[0081] In one aspect, the disclosure relates to methods of producing reproductively sterile fish and other egg-producing aquatic animals by administration of guanidine dendrimers to either fish eggs or embryos in order to disrupt primordial germ cell (PGC) development and migration to, and colonization in, the gonad of the embryo, which results in the failure of gonad

development and/or the failure of full and proper gonadal functioning at the cellular or tissular level, and ultimately the generation of sterile fish and other egg-producing aquatic animals.

[0082] In one aspect, the disclosure relates to a guanidine dendrimer comprising a triazine core, e.g. an octaguanidine dendrimer with a triazine core, also known in the art as "Vivo". Such compounds are described in U.S. Patent No. 7,935,816. The disclosure of U.S. Patent No. 7,935,816 is hereby incorporated herein by reference in its respective entirety.

[0083] An illustrative octaguanidine dendrimer transporter compound with morpholino as a representative agent is shown in the following conjugate:



[0084] A particular guanidine dendrimer that is useful with morpholino oligomers of the disclosure is 2-[(4-nitrophenyl)oxycarbonylhexamethylenecarbonylpiperazinyl]-4,6-bis{di[di(trifluoroacetamidohexyl)aminocarbonyloxyethyl]amino}triazine.

[0085] The disclosure further relates to the use of an octaguanidine dendrimer with a triazine core. Such a guanidine dendrimer is effective for chorionic transport of a Morpholino oligomer,

either covalently bound to the Morpholino oligomer or not covalently bound to the Morpholino oligomer.

[0086] In another aspect, the disclosure contemplates a method of producing reproductively sterile egg-producing aquatic animals, said method comprising contacting fertilized egg(s) (or embryos) with anti-sense Morpholino oligomer and a guanidine dendrimer that is effective to transfect the egg(s) and render individual(s) produced therefrom reproductively sterile.

[0087] The selected Morpholino oligomers may disrupt PGC development, migration and/or survival in large numbers of embryos, resulting in large-scale production of reproductively sterile adult fish and other egg-producing aquatic animals. The methods of the disclosure are also applicable to single embryos in smaller scale production of reproductively sterile adult fish and other egg-producing aquatic animals.

[0088] In one embodiment, a method of producing reproductively sterile egg-producing aquatic animals is provided, comprising immersing fertilized or unfertilized fish egg(s) in an immersion medium comprising a guanidine dendrimer, e.g. Vivo, and an antisense Morpholino oligomer capable of effectively suppressing expression of the *dead end* gene or other genes such as *nanos*, *vasa*, *gnrh* or *fsh* receptor that are essential for gonadal development in the fish and other egg-producing aquatic animals. Preferably, the guanidine dendrimer is an octaguanidine dendrimer with a triazine core, e.g. Vivo. The antisense Morpholino oligomer may, by way of example, be SEQ ID NO: 1 or SEQ ID NO: 2.

[0089] According to particular aspects of the methods of the disclosure, fertilized or unfertilized egg(s) (one or more) are immersed in an immersion medium comprising a guanidine dendrimer and an antisense Morpholino oligomer capable of effectively suppressing the expression of a *dead end* gene or other genes that are essential for gonadal development in the fish and other egg-producing aquatic animals. The concentration of the antisense Morpholino oligomer in the immersion medium should be sufficient to allow the antisense Morpholino oligomer to traverse the chorion of the egg(s). In embodiments, such concentration will typically be about 1 to about 200 μ M, more preferably about 10 to about 100 μ M, even more preferably from about 20 to about 60 μ M.

[0090] All aspects of the disclosure may include the administration or use of guanidine-containing compounds effective to enhance the permeability of the chorion of an egg from an aquatic animal. In such aspects, the guanidine-containing compounds are either covalently bound or not covalently bound to any agent intended to transfect the chorion of the egg.

[0091] It will be recognized that the contacting of eggs and/or embryos with the agents and compounds of the present disclosure may be carried out in any suitable manner, e.g., involving

immersion contacting, or alternatively by contacting not involving immersion, although it is to be recognized that immersion contacting provides an efficient and effective contacting technique that is amenable to large-scale operations for the production of reproductively sterile fish and aquatic animals. Non-immersion contacting techniques that may be utilized in the broad practice of the present disclosure include, without limitation, spray or drip methods for contacting, dry transfer techniques, in which eggs are contacted with a carrier or surface coated with or otherwise supporting any of the agents or compounds of the present disclosure that are effective for production of reproductively sterile fish, or other techniques by which the eggs and/or embryos are contacted with the agents or compounds of the present disclosure. The contacting preferably comprises a non-injection contacting.

[0092] The present disclosure relates in one aspect to delivering agents to eggs from fish or other egg-producing aquatic animals by contacting the eggs with the agents during water activation and water hardening of fertilized eggs (FIG. 3). During this period, the eggs actively up-take water (by way of example, up to 14.83 ± 2.05 microliter per Atlantic salmon egg). This active water-uptake force can be used to administer agents, for example, Morpholino oligomers into the eggs.

[0093] The present disclosure in some aspects describes methods of efficiently delivering agents into eggs of egg-producing aquatic animals by contacting the eggs with agents of interest during a window of time starting at water activation and water hardening. In various embodiments, the contacting comprises immersion of the eggs in an immersion medium containing one or more agent(s) of interest. As used herein, water activation and water hardening of fertilized eggs starts at the first contact of newly fertilized eggs with water to the end of water hardening.

[0094] Accordingly, in various aspects, the present disclosure relates to contacting fertilized eggs immediately starting at water-activation with an agent in immersion media. The agent used in the immersion medium may contribute to the sterilization of the egg-producing aquatic animals, and the immersion medium in specific embodiments may include additional agents or other materials that are beneficial to the egg-producing aquatic animals hatched from eggs contacted with the immersion medium, e.g., materials such as DNA/RNA, hormones, growth promoters, protective antigens, nutrients, etc.

[0095] According to particular aspects of the methods of the disclosure, fertilized egg(s) (one or more) are immersed in an aqueous immersion medium, thus starting water activation, comprising an antisense Morpholino oligomer or a guanidine-containing compound and an antisense Morpholino oligomer capable of effectively suppressing the expression of a *dead end* gene or other genes that are essential for gonadal development in the fish or other egg-producing

aquatic animals. The immersion continues until the chorion completes the hardening process. The concentration of the antisense Morpholino oligomer in the bath should be sufficient to allow the antisense Morpholino oligomer to traverse the chorion of the egg(s). In embodiments, such concentration will typically be about 1 to about 200 μ M, more preferably about 10 to about 100 μ M, even more preferably from about 20 to about 60 μ M.

[0096] The methods of the disclosure are applicable to egg-producing aquatic animals. As used herein, egg-producing aquatic animals include egg-bearing species of water-based animals including all fish species and other egg-producing aquatic animals such as crustaceans and/or mollusks.

[0097] Accordingly, egg-producing aquatic animals includes all fish species, including, but not limited to, salmon, Atlantic salmon, coho salmon, chinook salmon, chum salmon, sockeye salmon, pink salmon, masu salmon, trout, rainbow trout, brook trout, brown trout, common grayling, Arctic grayling, Arctic char, bass, hybrid bass, striped bass, white bass, striped-white bass hybrids, yellow bass, perch, white perch, yellow perch, European perch, bass-perch hybrids, Nile tilapia, blue tilapia, blue-Nile tilapia hybrids, Mozambique tilapia, zebrafish, carp species, breams, seabreams, porgies, catfish species, and cod.

[0098] The methods of the disclosure are additionally applicable to egg-producing aquatic animals such as crustaceans and/or mollusks. Such egg-producing aquatic animals may include, but are not limited to, shrimp, prawn, lobster, crayfish, crabs, oysters, squid, octopus, and the like.

[0099] The advantages and features of the disclosure are further illustrated with reference to the following examples, which are not to be construed as in any way limiting the scope of the disclosure but rather as illustrative of particular embodiments of the disclosure in specific applications thereof.

[00100] Examples:

[00101] Zebrafish were selected for initial exemplification of the methods of the disclosure, due to their short generation time and large numbers of embryos produced per mating, which are easily obtained on a daily or weekly basis. Additionally, the embryos of zebrafish are transparent, providing ease of visual observations, and are hardy. The normal development of PGCs and gonads within the embryo is an evolutionarily conserved mechanism that is found in all fish. Accordingly, the methods of the disclosure are applicable to all fish species, including, but not limited to, zebrafish, carp species, trout species, salmon species, breams (including seabreams and porgies), basses (including marine and freshwater seabass and

hybrid basses, etc), perches (yellow perch, white perch, etc), catfish species, cod and other major classes that are candidates for captive culture.

[00102] As described herein, the methods are generally applicable to farmed fish and aquatic egg-producing animals, as production of sterile farmed species is desirable. Accordingly, the methods of the invention are applicable to any farmed species of fish and aquatic egg-producing animals, particularly to commercially important farmed species.

[00103] Example 1

[00104] The disclosure relates to the specific chemicals, e.g., guanidine-containing compounds, which are able to enhance the permeability of chorion, which allows agents such as large molecules to traverse the chorion and reach the egg and embryo.

[00105] Zebrafish eggs were incubated in solutions that contained 1 μ M of 10, 20 or 40 KD fluorescein-dextran with either 0 or 40 mM of arginine, guanidine-HCl or guanidine-thiocyanate for 24 hours.

[00106] Fluorescently photographic results indicated that arginine, guanidine-HCl and guanidine-thiocyanate enhanced the uptake of fluorescein-dextran in the eggs that were treated with 40 mM of arginine, guanidine-HCl or guanidine-thiocyanate (treated eggs) over the up-take in the eggs which were incubated in solutions that did only contain fluorescein-dextran, but no guanidine-containing compounds (untreated eggs). As shown in FIG. 4, the treated eggs up-took more fluorescein-dextran than untreated eggs as indicated by higher green fluorescence found in the treated eggs. Thus, arginine, guanidine-HCl and guanidine-thiocyanate enhanced the permeability of chorion.

[00107] In addition, all three different sizes (molecule weight 10, 20 and 40 KD) of fluorescein-dextran were up-taken by the treated eggs. Accordingly, the methods of the invention are applicable to all egg-producing aquatic animals and are applicable to transfect eggs with various sized molecules, including both smaller and larger molecules.

[00108] Example 2

[00109] 40 zebrafish eggs were transferred into each well of 48-well plates that contained 300 μ l of fresh tank system water. After the allocation of eggs, system water in each well was replaced with 300 μ l of water that contained 10 or 100 μ M of poly-arginine (9-arginine polymer). After 18 hours of incubation the eggs were examined using an Axioplan 2 fluorescence microscope (ZEISS, Thornwood, NY, USA). The microscope is equipped with a DP70 digital camera (Olympus, Center Valley, PA, USA).

[00110] FIG. 5 shows the microphotographs obtained in the study. As shown in bright field photographic results, high concentration of poly-arginine/9 mer (100 μ M) generated

uncharacterized aggregates with in 1 hour of incubation, which was not seen in the 10 μM poly-arginine/9 mer treated group. After 18 hours incubation, high concentration of poly-arginine/9 mer (100 μM) caused chorion lysis that was not seen in the 10 μM poly-arginine/9 mer treated group.

[00111] Example 3

[00112] 40 zebrafish eggs were transferred into each well of 48-well plates that contained 300 μl of fresh tank system water. After the allocation of eggs, system water in each well was replaced with 300 μl of water that contained 0 or 40 μM of poly-arginine (9-arginine polymer) and 1 μM of 10 KD fluorescein-dextran. After 4 hours of incubation the eggs were examined using a MZ12 stereomicroscope (Leica, Buffalo Grove, USA), or an Axioplan 2 fluorescence microscope (ZEISS, Thornwood, NY, USA). Both microscopes were equipped with a DP70 digital camera (Olympus, Center Valley, PA, USA).

[00113] FIG. 6A-6D shows the microphotographs obtained in the study. As shown, poly-arginine enhanced the permeability of chorion. The zebrafish eggs that were incubated in solutions that contained 1 μM of 10 KD fluorescein-dextran with either 0 or 40 μM of poly-arginine are shown in FIG. 6A and 6B. Bright field photographic results indicated that poly-arginine (FIG. 6A) generated uncharacterized aggregates that were not seen in the control (FIG. 6B). FIG. 6C and FIG. 6D show fluorescently photographic results which indicated that poly-arginine enhanced the uptake of fluorescein-dextran as indicated by higher green fluorescence found in the poly-arginine treated eggs (FIG. 6C) than that of control eggs (FIG. 6D).

[00114] Example 4

[00115] A dendrimeric octaguanidine with a triazine core also known as Vivo, was conjugated to zebrafish *dnd*-MO, and used in an immersion containing zebrafish embryos in water. The ability of methods disclosed herein to induce sterility in zebrafish was tested by administering the following:

A: *dnd*-MO-Vivo 60 μM for 0.5 hour, 40 μM for 2 hours, 20 μM for 3 hours, 10 μM for 4.5 hours and 5 μM for 14 hours, administration began at the start of water activation and water hardening (immediately after fertilization);

B: water-only solution as control;

C: *dnd*-MO-Vivo 60 μM for 0.5 hour, 40 μM for 2 hours, 20 μM for 3 hours, 10 μM for 4.5 hours and 5 μM for 14 hours, administration began at one hour post-fertilization.

[00116] In zebrafish, 100% sterility induction can be achieved only when *dnd*-MO-Vivo was administered immediately starting at water activation and water hardening. If *dnd*-MO-Vivo was administered 1 hour after water activation only 44-59 % treated fish were infertile (Table 1).

[00117]

TABLE 1

	A1	A2	B1	B2	C1	C2
embryos survived to 2 dpf (days post-fertilization)	41	40	52	50	45	47
adult fish obtained	28	32	39	43	34	29
adult survival rate	50%	57%	69%	76%	60%	51%
number of males	0	0	16	15	12	9
number of females	0	0	23	28	7	3
number of infertile fish	28	32	0	0	15	17
% of infertile fish	100%	100%	0%	0%	44%	59%
Average % of infertile fish		100%		0%		52%

[00118] **Example 5**

[00119] A dendrimeric octaguanidine with a triazine core also known as Vivo, was conjugated to salmonids' *dnd*-MO, 5'- CTGACTTGAACGCTCCTCCATTATC-3' (SEQ ID NO. 3) and used in an immersion containing rainbow trout embryos in water, or unfertilized eggs in ovarian fluid or fertilization diluent. The ability of methods disclosed herein to induce sterility in rainbow trout was tested by administering the following:

I: Salmonids' *dnd*-MO-Vivo 10 μ M for 48 hours, administration to fertilized eggs at the beginning of water activation (within 1 minute)

II: Salmonids' *dnd*-MO-Vivo 10 μ M for 48 hours, administration to unfertilized eggs with the immersion medium of the ovarian fluid. The fertilization was conducted after 48 hours of incubation.

III: Salmonids' *dnd*-MO-Vivo 10 μ M for 48 hours, administration to unfertilized eggs with the fertilization diluent that contains salt, Tris (pH 8.0), Glycine, and 5% of fish serum. The fertilization was conducted after 48 hours of incubation.

IV: Water or immersion medium only without salmonids' *dnd*-MO-Vivo as controls.

[00120] FIG. 7A-7G shows that salmonids' *dnd*-MO-Vivo treatment induced infertility in rainbow trout. (FIG. 7A) a well-developed testis of untreated control male fish from treatment IV; (FIG. 7B) a well-developed ovary of untreated control female fish from treatment IV; (FIG. 7C) the gonads of salmonids' 10 μ M *dnd*-MO-Vivo treated fish in treatments I, II and III, developed into a thin filament-like tissue; (FIG. 7D) a photomicrograph of dissected gonads;

(FIG. 7E) the active spermatogenesis of the testis of an untreated control (treatment IV) male fish; (FIG. 7F) a well-developed ovary of an untreated control female (treatment IV) with oocytes at different developmental stages; (FIG. 7G) the histological examinations of gonadal tissue show the gonad of a salmonids' *dnd*-MO-Vivo treated fish (treatment I, II and III) that appears to be under-developed without advanced gonadal structure or germ cells.

[00121] Example 6

[00122] A salmonids' *dnd*-MO (see Example 5) or a dendrimeric octaguanidine with a triazine core also known as Vivo, which was conjugated to salmonids' *dnd*-MO, resulting in salmonids' *dnd*-MO-Vivo, was used to induce sterility in Atlantic salmon and was tested by administering the following:

I: Salmonids' *dnd*-MO-Vivo 10 μ M for 48 hours, administration to fertilized eggs at the beginning of water activation (within 1 minute).

II: Salmonids' *dnd*-MO-Vivo 10 μ M for 48 hours, administration to unfertilized eggs with the immersion medium of the ovarian fluid. The eggs were fertilized after 48 hours of incubation.

III: Salmonids' *dnd*-MO-Vivo 10 μ M for 48 hours, administration to unfertilized eggs with the immersion medium that contains salt, Tris (pH 8.0), Glycine, and 5% of fish serum. The fertilization was conducted after 48 hours of incubation.

IV: Salmonids' *dnd*-MO 20 μ M for 48 hours, administration to unfertilized eggs with the immersion medium of the ovarian fluid. The eggs were fertilized after 48 hours of incubation.

V: Salmonids' *dnd*-MO 20 μ M for 48 hours, administration to unfertilized eggs with the immersion medium that contains salt, Tris (pH 8.0), Glycine, and 5% of fish serum. The eggs were fertilized after 48 hours of incubation.

VI: Water or immersion medium only without either salmonids' *dnd*-MO-Vivo nor *dnd*-MO as controls.

[00123] FIG. 8A-8D shows that salmonids' *dnd*-MO or *dnd*-MO-Vivo treatment blocked gonadal development in Atlantic salmon. In 10-month-old Atlantic salmon, (FIG. 8A) A developing testis of untreated (treatment VI) male; (FIG. 8B) a developing ovary of untreated (treatment VI) female; (FIG. 8C) the gonads of *dnd*-MO-Vivo treated (treatments I, II, III) or *dnd*-MO treated (treatments IV, V) fish developed into a thin filament-like tissue; (FIG. 8D) a photomicrograph of dissected gonads from (FIG. 8A), (FIG. 8B) and (FIG. 8C).

[00124] While the disclosure has been set out herein in reference to specific aspects, features and illustrative embodiments, it will be appreciated that the utility of the disclosure is not thus limited, but rather extends to and encompasses numerous other variations, modifications and alternative embodiments, as will suggest themselves to those of ordinary skill

in the field of the present disclosure, based on the description herein. Correspondingly, the invention as hereinafter claimed is intended to be broadly construed and interpreted, as including all such variations, modifications and alternative embodiments, within its spirit and scope.

INDUSTRIAL APPLICABILITY

[00125] The disclosure provides methods for delivering agents to eggs/embryos of aquatic animals using guanidine-containing compounds to enhance the permeability of the chorion of the eggs. Delivery of drug candidates or potentially bioactive agents to the eggs/embryos of aquatic animals, such as zebrafish, offer opportunities for screening and evaluation of the activity and/or toxicity of such test agents.

[00126] The methods and compounds of the disclosure further produce reproductively sterile fish and aquatic egg-producing animals. Sterilization (induced infertility) of farmed fish and aquatic egg-producing animals enhances their growth rate by increasing the conversion of food energy to muscle growth, instead of gonadal development. In addition, if escaped from aquaculture operations to the environment, reproductively sterile farmed fish and egg-producing aquatic animals, including domesticated, non-native or genetically modified species, will not be able to reproduce or inter-breed with wild stock. This will assist biological containment and prevent genetic contamination of wild populations and/or the establishment in the wild of domestic, non-native or genetically modified farmed fish and aquatic egg-producing animals.


THE CLAIMS

What is claimed is:

1. A method of delivering at least one agent into fish egg(s), said method comprising contacting the fish egg(s) with the at least one agent in the presence of a guanidine-containing compound that is effective to enhance permeability of the chorion of the fish egg(s), wherein the fish egg(s) are contacted with the at least one agent in an immersion medium comprising the at least one agent and the guanidine-containing compound and wherein said guanidine-containing compound is arginine or a salt thereof, guanidine or a salt thereof or a dimer, trimer or polymer of guanidine or arginine and wherein said polymer is not a dendrimeric oligoguanidine.
2. The method of claim 1, wherein said contacting comprises chorionic transfection of the fish egg(s).
3. The method of claim 1 or claim 2, wherein concentration of the guanidine-containing compound in the immersion medium is in a range of from 1 to 80,000 μ M.
4. The method of any of claims 1 to 3, wherein the guanidine-containing compound is arginine or a salt thereof, preferably wherein the guanidine-containing compound is arginine.
5. The method of any of claims 1 to 3, wherein the guanidine-containing compound is guanidine or a salt thereof, preferably wherein the guanidine-containing compound is guanidine-HCl, guanidine-thiocyanate, guanidine-acetate, guanidine-carbonate, guanidine-nitrate, guanidine-sulfate, guanidine-bicarbonate or guanidine-hydrobromide.
6. The method of any of claims 1 to 3, wherein the guanidine-containing compound is a dimer, trimer or polymer of guanidine or arginine, wherein said polymer is not a dendrimeric oligoguanidine.
7. The method of any of claims 1 to 6, wherein the at least one agent is selected from the group consisting of antibodies, proteins, peptides, RNAs, and DNAs.
8. The method of any of claims 1 to 7, wherein the at least one agent is an anti-sense Morpholino oligomer that is effective to suppress expression of a gene essential for gonadal development.

9. The method of claim 8, wherein the anti-sense Morpholino oligomer suppresses expression of the *dead end*, *nanos*, *vasa*, *gnrh* or *fsh* receptor gene in the fish.
10. The method of claim 9, wherein the anti-sense Morpholino oligomer suppresses expression of the *dead end* gene in the fish.
11. The method of any of claims 1 to 10 for producing reproductively sterile fish, wherein the agent is an anti-sense Morpholino oligomer that is effective to transfect the fish egg(s) and render individual(s) produced therefrom reproductively sterile.
12. The method of claim 10 or claim 11, wherein the egg(s) are fertilized or unfertilized egg(s) of a fish selected from the group consisting of salmonids, moronids, and cichlids.
13. The method of any of claims 1 to 12, wherein the guanidine-containing compound is not covalently bound to the agent.
14. The method of any of claims 1 to 12, wherein the guanidine-containing compound is covalently bound to the agent.
15. The method of any of claims 8 to 14, wherein the anti-sense Morpholino oligomer comprises the nucleotide sequence of SEQ ID NO: 1 or SEQ ID NO: 2 or a variant thereof.
16. The method of any of claims 1 to 15, wherein the guanidine-containing compound is poly-arginine/9-mer.
17. A transfected fish egg of an egg-producing aquatic animal obtained by a method according to any of claims 2 to 16, said fish egg comprises a chorion-permeated amount of said at least one agent and a chorion-permeated amount of said guanidine-containing compound.
18. The transfected fish egg of claim 17, wherein the at least one agent is an anti-sense Morpholino oligomer that renders an individual produced from such egg reproductively sterile.

19. The transfected fish egg of claim 17 or claim 18, wherein said egg is a salmonids egg and the at least one agent is an anti-sense Morpholino oligomer which comprises the nucleotide sequence of SEQ ID NO: 1 or SEQ ID NO: 2 or a variant thereof.

**FIG. 1**

2/8






FIG. 2

3/8

**FIG. 3**

4/8

**FIG. 4**

5/8




FIG. 5

6/8






FIG. 6A-6D

7/8

**FIG. 7A-7G**

8/8

**FIG. 8A-8D**

268PCT_SeqID_ST25
SEQUENCE LISTING

<110> UNIVERSITY OF MARYLAND BALTIMORE COUNTY
<120> METHODS OF AGENT DELIVERY INTO EGGS AND EMBRYOS OF EGG-PRODUCING AQUATIC ANIMALS FOR DRUG SCREENING, AGENT TOXICITY ASSAY AND PRODUCTION OF INFERTILE FISH
<130> 4451-268-PCT
<140> Not yet assigned
<141> 2016-05-17
<150> US 62/163,511
<151> 2015-05-19
<150> US 62/274,958
<151> 2016-01-05
<160> 3
<170> PatentIn version 3.5
<210> 1
<211> 12
<212> DNA
<213> *Salmo* *salar*
<400> 1
acgctccctcc at 12

<210> 2
<211> 18
<212> DNA
<213> *Salmo* *salar*
<400> 2
acttgAACGc tcctccat 18

<210> 3
<211> 25
<212> DNA
<213> *Salmo* *salar*
<400> 3
ctgacttgaa cgctccctcca ttatc 25