



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2019/01/04
(87) Date publication PCT/PCT Publication Date: 2019/07/11
(85) Entrée phase nationale/National Entry: 2020/05/04
(86) N° demande PCT/PCT Application No.: EP 2019/050186
(87) N° publication PCT/PCT Publication No.: 2019/134982
(30) Priorité/Priority: 2018/01/04 (US62/613,646)

(51) Cl.Int./Int.Cl. *A61K 36/185* (2006.01),
A61K 31/57 (2006.01), *A61K 31/575* (2006.01),
A61K 41/10 (2020.01), *A61K 9/10* (2006.01),
A61K 9/107 (2006.01), *A61K 9/12* (2006.01),
A61P 17/00 (2006.01), *A61P 17/02* (2006.01)

(71) Demandeur/Applicant:
AMRYT RESEARCH LIMITED, DE

(72) Inventeurs/Inventors:
WATSON, JOHN ASHLEIGH, CH;
JAGER, SEBASTIAN, DE;
ZAHN, TOBIAS, DE

(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : EXTRAITS D'ECORCE DE BOULEAU CONTENANT DE LA BETULINE ET LEUR FORMULATION
(54) Title: BETULIN-CONTAINING BIRCH BARK EXTRACTS AND THEIR FORMULATION




FIG. 1

(57) Abrégé/Abstract:

The present disclosure relates to birch bark extracts, methods of producing such extracts, stable pharmaceutical compositions containing such extracts and methods of using of such extracts. The birch bark extracts of the present disclosure contain triterpenes, which are known to improve wound healing.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

11 July 2019 (11.07.2019)



(10) International Publication Number

WO 2019/134982 A1

(51) International Patent Classification:

A61K 45/06 (2006.01) **A61L 15/44** (2006.01)
A61K 9/107 (2006.01) **A61K 31/56** (2006.01)
A61K 9/12 (2006.01) **A61K 36/185** (2006.01)
A61F 13/00 (2006.01) **A61P 17/00** (2006.01)
A61K 9/70 (2006.01) **A61P 17/02** (2006.01)
A61K 31/19 (2006.01)

(21) International Application Number:

PCT/EP2019/050186

(22) International Filing Date:

04 January 2019 (04.01.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/613,646 04 January 2018 (04.01.2018) US

(71) **Applicant: AMRYT RESEARCH LIMITED** [IE/IE]; 90
Harcourt Street, Dublin 2 (IE).

(72) **Inventors: WATSON, John Ashleigh**; Zollweg 11, 8880
Walenstadt (CH). **JÄGER, Sebastian**; Scheffelstrasse 15,

75446 Wiernsheim (DE). **ZAHN, Tobias**; Merkurstr. 8,
75179 Pforzheim (DE).

(74) **Agent: COOLEY (UK) LLP**; Dashwood, 69 Old Broad
Street, London Greater London EC2M 1QS (GB).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) **Title: BETULIN-CONTAINING BIRCH BARK EXTRACTS AND THEIR FORMULATION**




FIG. 1

(57) **Abstract:** The present disclosure relates to birch bark extracts, methods of producing such extracts, stable pharmaceutical compositions containing such extracts and methods of using of such extracts. The birch bark extracts of the present disclosure contain triterpenes, which are known to improve wound healing.

[Continued on next page]

WO 2019/134982 A1

WO 2019/134982 A1

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

BETULIN-CONTAINING BIRCH BARK EXTRACTS AND THEIR FORMULATION

CROSS-REFERENCE TO RELATED APPLICATIONS

[001] This application claims the benefit of U.S. Provisional Application No. 62/613,646, filed January 4, 2018, the entire contents of which are hereby incorporated by reference in its entirety for all purposes.

FIELD OF THE DISCLOSURE

[002] The present disclosure relates to pharmaceutical formulations derived from the extracts of birch bark.

BACKGROUND OF THE DISCLOSURE

[003] The triterpenes found in birch bark extracts are known to have wound healing properties. Methods of extracting these triterpenes from birch bark are reported in U.S. Patent No. 7,482,383. These methods provide solid birch bark extracts that may be used in pharmaceutical formulations. For example, emulsions containing such extracts are described in U.S. Patent No. 7,482,383, and oleogels containing such extracts are described in U.S. Patent Nos. 9,352,041; 8,828,444 and 8,536,380.

[004] For clinical use in wound healing, an oleogel must be applied by touch to the area of the skin where treatment is needed. Touch application is disadvantageous for the treatment of certain skins conditions (e.g., epidermolysis bullosa) because the simple act of applying the oleogel may lead to worsening of the skin condition.

[005] Therefore, there is a need for wound-healing formulations containing solid birch bark extracts that may be applied to the skin without worsening the patient's condition.

[006] The present disclosure provides clinically-advantageous wound-healing formulations (including oleogels, emulsions and foams) with improved rheological properties which overcome the disadvantages of the known solid birch bark-containing emulsions and oleogels.

SUMMARY OF THE DISCLOSURE

[007] The present disclosure provides a solid birch bark extract that contains at least about 70% by weight of betulin and one or more triterpenes selected from the group consisting of betulinic acid, oleanolic acid, erythrodiol and lupeol. The solid birch bark extracts described herein may be formulated in an oleogel that possesses clinically-advantageous rheological properties.

[008] The present disclosure also provides a method of making a solid birch bark extract that may be dispersed in a nonpolar solvent to provide a clinically-advantageous oleogel comprising the steps of (a) contacting birch bark with a suitable solvent to form an extraction solution containing betulin and at least one triterpene; (b) separating the birch bark from the extraction solution; (c) cooling the extraction solution to crystallize a portion of the betulin and triterpene from the solution; (d) separating the crystallized betulin and triterpene; (e) drying the separated, crystallized betulin and triterpene to form a solid birch bark extract. The present disclosure also provides the solid birch bark extracts that are prepared according to these methods.

[009] The present disclosure provides a clinically-advantageous oleogel comprising about 1% to about 20% by weight of particles of the solid birch bark extract dispersed in about 80% to about 99% of one or more nonpolar liquids the solid birch bark extracts are dispersed in a suitable nonpolar liquid to form an oleogel. A sterile wound dressing comprising a pad and a therapeutically effective amount of the oleogel is also provided.

[010] The present disclosure provides an emulsion comprising an oleogel of the present invention and further provides a foam comprising an emulsion of the present disclosure.

[011] The present disclosure also provides methods of treating a wound in a patient by topically administering an effective amount of an oleogel, an emulsion or a foam of the present disclosure to at least a portion of the wound.

[012] The present disclosure also provides methods of treating epidermolysis bullosa in a patient in need thereof comprising topically administering an effective amount of an oleogel, an emulsion or a foam of the present disclosure to an area of epidermolysis bullosa of the patient.

[013] The present disclosure provides a solid birch bark extract, oleogel, sterile wound dressing, process of preparation, emulsion, foam, pressurized container, method of treating a wound, and method of treating epidermolysis bullosa according to the following numbered embodiments:

1. A solid birch bark extract comprising at least about 70 wt. % betulin and one or more triterpenes selected from the group consisting of betulinic acid, oleanolic acid, erythrodiol, and lupeol.
2. The solid birch bark extract of embodiment 1, wherein the solid birch extract has an HPLC chromatogram substantially identical to Figure 1 when chromatographed according to the following HPLC method:

Time (min)	Acetonitrile	Water	H ₃ PO ₄ (v/v)	Flow rate (ml/min)
0	30	70	0.1%	0.6
5	30	30	0.1%	0.6
15	98	2	0.1%	0.6
20	98	2	0.1%	0.6
21	30	70	0.1%	0.6
30	30	70	0.1%	0.6

Column: C18, 2.6 μ m, 150 x 2.1 mm; detector: 210 and 320 nm;

wherein the betulinic acid has a relative retention time of about 0.75 – 0.90, the oleanolic acid has a relative retention time of about 0.84-0.97, the betulin has a relative retention time of 1.00, the erythrodiol has a relative retention time of about 1.25-1.40, and the lupeol has a relative retention time of about 3.50-4.15.

3. The solid birch bark extract of embodiment 1 or 2, prepared by a process comprising:
 - (a) contacting birch bark with a pharmaceutically acceptable solvent, thereby forming an extraction solution comprising betulin and one or more triterpenes;
 - (b) separating the birch bark from the extraction solution;
 - (c) cooling the extraction solution whereby a portion of the betulin and one or more triterpenes crystallizes from the cooled extraction solution;

- (d) separating the crystallized betulin and one or more triterpenes from the cooled extraction solution;
 - (e) drying the separated, crystallized betulin and one or more triterpenes to provide the solid birch bark extract;
- wherein if after drying, about 1 wt.% to about 20 wt.% of the dried solid birch bark extract of step (e) dispersed in refined sunflower oil forms an oleogel.
4. The solid birch bark extract of embodiment 3, wherein one or more of steps (a), (c), and (e) is carried out under the following conditions:
- (i) said contacting of step (a) is carried out at a temperature of about 60° C to about 130° C;
 - (ii) said cooling of step (c) is carried out at a temperature of about -20° C to about 35° C;
 - (iii) said drying of step (e) is carried out at a temperature of about 75° C to about 95° C at a pressure of less than about 70 mbar.
5. The solid birch bark extract of embodiment 3, wherein two or more of steps (a), (c), and (e) are carried out under the following conditions:
- (i) said contacting of step (a) is carried out at a temperature of about 60° C to about 130° C;
 - (ii) said cooling of step (c) is carried out at a temperature of about -20° C to about 35° C;
 - (iii) said drying of step (e) is carried out at a temperature of about 75° C to about 95° C at a pressure of less than about 70 mbar.
6. The solid birch bark extract of embodiment 3, wherein steps (a), (c), and (e) are carried out under the following conditions:
- (i) said contacting of step (a) is carried out at a temperature of about 60° C to about 130° C;
 - (ii) said cooling of step (c) is carried out at a temperature of about -20° C to about 35° C;
 - (iii) said drying of step (e) is carried out at a temperature of about 75° C to about 95° C at a pressure of less than about 70 mbar.

7. The solid birch bark extract of any one of embodiments 3-6, wherein the pharmaceutically acceptable solvent is a hydrocarbon or alcohol, or combinations thereof.
8. The solid birch bark extract of any one of embodiments 3-7, wherein the pharmaceutically acceptable solvent is a hydrocarbon selected from the group consisting of n-pentane, n-hexane, or n-heptane.
9. The solid birch bark extract of embodiment 8, wherein the pharmaceutically acceptable solvent is n-heptane, and said contacting is carried out at a temperature of about 60 °C to about 130 °C for about 8-12 minutes.
10. The solid birch bark extract of any one of embodiments 3-9, wherein in step (c) the cooled extraction solution is supersaturated at least about 2-fold.
11. The solid birch bark extract of embodiment 10, wherein the cooled extraction solution is supersaturated about 5-fold.
12. The solid birch bark extract of any of embodiments 3-11, wherein the amount of residual extraction solvent is no more than about 0.5 wt.%.
13. An oleogel comprising about 1 wt. % to about 20 wt. % of particles of the solid birch bark extract of any one of embodiments 1-12, having an average particle size of less than about 50 µm, dispersed in about 80 wt.% to about 99 wt.% of one or more nonpolar liquids.
14. The oleogel of embodiment 13, wherein the dispersed solid birch bark extract particles are the only oleogel forming agent in the oleogel.
15. The oleogel of embodiment 13 or 14, comprising about 10 wt. % of particles of the solid birch bark extract.
16. The oleogel of any one of embodiments 13 to 15, wherein the nonpolar liquid comprises at least one triglyceride.

17. The oleogel of any one of embodiments 13 to 16, wherein the nonpolar liquid comprises at least one C7 or greater hydrocarbon.
18. The oleogel of embodiment 16, wherein the nonpolar liquid comprises one or more vegetable oils.
19. The oleogel of embodiment 18, wherein the nonpolar liquid comprises sunflower oil.
20. The oleogel of any one of embodiments 13-19, wherein the nonpolar liquid has a peroxide value less than about 10.
21. The oleogel of embodiment 20, wherein the peroxide value is no more than about 3.
22. The oleogel of any one of embodiments 13 - 21, wherein the oleogel is substantially free of solid birch bark extract particles having a size greater than about 50 μm .
23. The oleogel of any one of embodiments 13-22, wherein the oleogel is sterile.
24. The oleogel of embodiment 23, wherein the oleogel is sterilized by ionizing irradiation at doses less than about 20 kGy.
25. The oleogel of embodiment 24, wherein the oleogel is sterilized by ionizing irradiation at doses ranging from about 11 to about 20 kGy.
26. The oleogel of any one of embodiments 13-25, wherein the segregation of the nonpolar liquid from the oleogel is less than about 10% after centrifuging at 25°C for 30 min at 2750 g.
27. The oleogel of any one of embodiments 13-26, wherein the viscosity at 200/s of the oleogel ranges from about 0.5 to about 4.0 Pa.s and the thixotropy value of the oleogel ranges from about 200 to about 1200 Pa.s as measured according to the rotating viscometer method described in Ph. Eur. 2.2.10 using a cone-plate viscometer.

28. The oleogel of any one of embodiments 13-27, wherein the consistency of the oleogel ranges from about 300-3000 mN as measured according using a texture analyser.
29. The oleogel of any one of embodiments 13-28, further comprising a disinfectant.
30. The oleogel of embodiment 29, wherein the disinfectant is selected from the group consisting of ethanol, n-propanol, and isopropanol.
31. The oleogel of any one of embodiments 13-28, further comprising a lipophilic antibiotic.
32. A sterile wound dressing comprising:
 - (a) a pad, and
 - (b) a therapeutically effective layer comprising the oleogel of any of embodiments 13-31 disposed on at least one surface of the pad.
33. The sterile wound dressing of embodiment 32, wherein the pad is an absorbent pad.
34. A process for preparing the solid birch bark extract of embodiment 1 or 2, comprising:
 - (a) contacting birch bark with a pharmaceutically acceptable solvent, thereby forming an extraction solution comprising betulin and one or more triterpenes;
 - (b) separating the birch bark from the extraction solution;
 - (c) cooling the extraction solution whereby a portion of the betulin and one or more triterpenes crystallizes from the cooled extraction solution;
 - (d) separating the crystallized betulin and one or more triterpenes from the cooled extraction solution;
 - (e) drying the separated, crystallized betulin and one or more triterpenes to provide the solid birch bark extract;wherein after drying, about 1 wt.% to about 20 wt.% of the dried solid birch bark extract of step (e) dispersed in refined sunflower oil forms an oleogel.

35. The process of embodiment 34, wherein one or more of steps (a), (c), and (e) is carried out under the following conditions:
- (i) said contacting of step (a) is carried out at a temperature of about 60° C to about 130° C;
 - (ii) said cooling of step (c) is carried out at a temperature of about -20° C to about 35° C;
 - (iii) said drying of step (e) is carried out at a temperature of about 75° C to about 95° C at a pressure of less than about 70 mbar.
36. The process of embodiment 34, wherein two or more of steps (a), (c), and (e) are carried out under the following conditions:
- (i) said contacting of step (a) is carried out at a temperature of about 60° C to about 130° C;
 - (ii) said cooling of step (c) is carried out at a temperature of about -20° C to about 35° C;
 - (iii) said drying of step (e) is carried out at a temperature of about 75° C to about 95° C at a pressure of less than about 70 mbar.
37. The process of embodiment 34, wherein steps (a), (c), and (e) are carried out under the following conditions:
- (i) said contacting of step (a) is carried out at a temperature of about 60° C to about 130° C;
 - (ii) said cooling of step (c) is carried out at a temperature of about -20° C to about 35° C;
 - (iii) said drying of step (e) is carried out at a temperature of about 75° C to about 95° C at a pressure of less than about 70 mbar.
38. The process of any one of embodiments 34-37, wherein the pharmaceutically acceptable solvent is a hydrocarbon selected from the group consisting of n-pentane, n-hexane, or n-heptane.
39. The process of any one of embodiments 34-38, wherein the pharmaceutically acceptable solvent is n-heptane, and said contacting is carried out at a temperature of about 115 °C to about 130 °C for about 8-12 minutes.

40. The process of any one of embodiments 34-39, wherein in step (c) the cooled extraction solution is supersaturated at least about 2-fold.
41. The process of embodiment 40, wherein the cooled extraction solution is supersaturated about 5-fold.
42. The process of any of embodiments 34-41, wherein the amount of residual extraction solvent is no more than about 0.5 wt.%.
43. An emulsion comprising the solid birch bark extract of any of embodiments 1-12.
44. An emulsion comprising the oleogel of any of embodiments 13-31.
45. The emulsion of embodiment 43 or 44, wherein said emulsion is a water-in-oil emulsion.
46. The emulsion of embodiment 45, substantially free of emulsifier.
47. The emulsion of any one of embodiments 43-46, consisting essentially of the solid birch bark extract or oleogel, oil, and water.
48. The emulsion of any one of embodiments 44-46, consisting of the oleogel, oil and water.
49. The emulsion of any one of embodiments 43-46, further comprising a disinfectant.
50. The emulsion of embodiment 49, wherein the disinfectant is selected from the group consisting of ethanol, n-propanol, and isopropanol.
51. The emulsion of any one of embodiments 43-46, further comprising a lipophilic antibiotic.

52. A foam comprising the emulsion of any one of embodiments 43-51.
53. The foam of embodiment 52, further comprising a disinfectant.
54. The foam of embodiment 53, wherein the disinfectant is selected from the group consisting of ethanol, n-propanol, and isopropanol.
55. The foam of embodiment 52, further comprising a lipophilic antibiotic.
56. The foam of any of embodiments 52-55, wherein the interfacial surface tension of the emulsion is greater than about 4 mN/m.
57. The foam of any of embodiments 52-56, wherein the foam index is greater than about 2.
58. The foam of embodiment 52, wherein the oleogel consists of about 5 wt.% to about 10 wt.% solid birch bark extract and the emulsion is a water-in-oil emulsion consisting of the oleogel and about 20 wt.% to about 30 wt.% of water.
59. The foam of embodiment 58, wherein the oleogel consists of about 7 wt.% solid birch bark extract and the amount of water in the emulsion is about 25 wt.%.
60. The foam of any of embodiments 52-56, further comprising an emulsifier selected from the group consisting of phosphatidyl choline, polyglyceryl-3-methyl glucose distearate, and combinations thereof.
61. A pressurized container filled with a mixture comprising the emulsion of any one of embodiments 43-51 and a pharmaceutically acceptable propellant, whereby the emulsion forms a foam upon decanting at least a portion of the mixture from the container.
62. A method of treating a wound in a patient comprising topically administering the oleogel of any one of embodiments 13-31 to at least a portion of the wound.

63. The oleogel of any one of embodiments 13-31, for use in a method of treating a wound.
64. The method or oleogel for use of embodiment 63, wherein the wound is selected from the group consisting of a burns, surgical skin lesions, superficial injuries, chronic wounds, pressure ulcers, diabetic foot ulcers, chronic venous ulcers, artery insufficiency ulcers, wounds from aesthetic skin treatments, wounds from ablative laser skin treatments, wounds from chemical peels, wounds from dermabrasion, wounds resulting from adverse drug reactions, wounds resulting from toxic epidermal necrolysis, wounds resulting from Lyell syndrome, wounds resulting from Stevens-Johnson syndrome, and wounds resulting from radiation dermatitis.
65. A method of treating a wound in a patient comprising topically administering the emulsion of any one of embodiments 43-51 to at least a portion of the wound.
66. The emulsion of any one of embodiments 43-51, for use in a method of treating a wound.
67. The method or emulsion for use of embodiment 66, wherein the wound is selected from the group consisting of burns, surgical skin lesions, superficial injuries, chronic wounds, pressure ulcers, diabetic foot ulcers, chronic venous ulcers, artery insufficiency ulcers, wounds from aesthetic skin treatments, wounds from ablative laser skin treatments, wounds from chemical peels, wounds from dermabrasion, wounds resulting from adverse drug reactions, wounds resulting from toxic epidermal necrolysis, wounds resulting from Lyell syndrome, wounds resulting from Stevens-Johnson syndrome, and wounds resulting from radiation dermatitis.
68. A method of treating a wound in a patient comprising topically administering the foam of any one of embodiments 52-60 to at least a portion of the wound.
69. The foam of any one of embodiments 52-60, for use in a method of treating a wound.
70. The method or foam for use of embodiment 69, wherein the wound is selected from the group consisting of burns, surgical skin lesions, superficial injuries, chronic

wounds, pressure ulcers, diabetic foot ulcers, chronic venous ulcers, artery insufficiency ulcers, wounds from aesthetic skin treatments, wounds from ablative laser skin treatments, wounds from chemical peels, wounds from dermabrasion, wounds resulting from adverse drug reactions, wounds resulting from toxic epidermal necrolysis, wounds resulting from Lyell syndrome, wounds resulting from Stevens-Johnson syndrome, and wounds resulting from radiation dermatitis.

71. A method of treating epidermolysis bullosa in a patient in need thereof comprising topically administering to an area of epidermolysis bullosa of the patient the oleogel of any one of embodiments 13-31.
72. The oleogel of any one of embodiments 13-31, for use in a method of treating epidermolysis bullosa.
73. A method of treating epidermolysis bullosa in a patient in need thereof comprising topically administering to an area of epidermolysis bullosa of the patient the emulsion of any one of embodiments 43-51.
74. The emulsion of any one of embodiments 43-51, for use in a method of treating epidermolysis bullosa.
75. A method of treating epidermolysis bullosa in a patient in need thereof comprising topically administering to an area of epidermolysis bullosa of the patient the foam of any one of embodiments 52-60.
76. The foam of any one of embodiments 52-60, for use in a method of treating epidermolysis bullosa.

DETAILED DESCRIPTION OF THE DRAWINGS

[014] FIG. 1. shows the HPLC chromatograph taken from a solid birch bark extract monitoring 210 nm (top figure) and 320 nm (bottom figure). The x-axis in FIG. 1 is the retention time (min), and the y-axis is the peak intensity (mAU).

[015] FIG 2. illustrates the continuous extraction of birch bark to provide an extraction solution comprising betulin and one or more triterpenes.

DETAILED DESCRIPTION OF THE DISCLOSURE

Definitions

[016] While the following terms are believed to be well understood by one of ordinary skill in the art, the following definitions are set forth to facilitate explanation of the presently disclosed subject matter.

[017] It is to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

[018] Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this disclosure belongs. Preferred methods, devices, and materials are described, although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure. All references cited herein (including U.S. Patent Nos. 9,352,041; 8,828,444; 8,536,380; and 7,482,383) are incorporated for all purposes by reference in their entirety.

[019] Following long-standing patent law convention, the terms "a", "an", and "the" refer to "one or more" when used in this application, including the claims. Thus, for example, reference to "a carrier" includes mixtures of one or more carriers, two or more carriers, and the like.

[020] Unless otherwise indicated, all numbers expressing quantities of ingredients, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term "about". Accordingly, unless indicated to the contrary, the numerical parameters set forth in the present specification and attached claims are approximations that can vary depending upon the desired properties sought to be obtained by the present application. Generally the term "about", as used herein when referring to a measurable value such as an amount of weight, time, dose, etc. is meant to encompass in one example variations of $\pm 15\%$ or $\pm 10\%$, in another example $\pm 5\%$, in another example $\pm 1\%$, and in yet another example $\pm 0.1\%$ from the specified amount, as such variations are appropriate to perform the disclosed method.

[021] “Administering” includes any mode of administration, such as oral, subcutaneous, sublingual, transmucosal, parenteral, intravenous, intra-arterial, buccal, sublingual, topical, vaginal, rectal, ophthalmic, otic, nasal, inhaled, and transdermal. “Administering” can also include prescribing or filling a prescription for a dosage form comprising a particular compound. “Administering” can also include providing directions to carry out a method involving a particular compound or a dosage form comprising the compound.

[022] As used herein, the verb “comprise” as is used in this description and in the claims and its conjugations are used in its non-limiting sense to mean that items following the word are included, but items not specifically mentioned are not excluded.

[023] The term “birch bark” means the cortex of white-barked birch trees. Preferred embodiments include birch bark derived from *Betula pendula* Roth and *Betula pubescens* Ehrh as well as hybrids of both species.

[024] The term “relative retention time” (or “RRT”) means ratio of the retention time of the unknown peak to the retention time the betulin peak and is calculated according to the following equation:

$$RRT = RT_{\text{unknown}} / RT_{\text{betulin}}$$

where RT_{unknown} is the retention time of the unknown peak and RT_{betulin} is the retention time of betulin peak (210 nm) measured by the following HPLC method:

Time (min)	Acetonitrile	Water	H ₃ PO ₄ (v/v)	Flow rate (ml/min)
0	30	70	0.1%	0.6
5	30	30	0.1%	0.6
15	98	2	0.1%	0.6
20	98	2	0.1%	0.6
21	30	70	0.1%	0.6
30	30	70	0.1%	0.6

Column: Phenomenex Kinetix C18, 2.6 μ m, 150 x 2.1 mm.

[025] The term “substantially identical” as used herein means an analytical spectrum, such as HPLC chromatogram, NMR spectrum, and etc., which resembles the reference spectrum to a great degree in both the peak locations and their intensity. For example, a HPLC chromatogram is “substantially identical” to a reference chromatogram if the peak locations (relative retention

times) in the HPLC chromatogram vary by no more than $\pm 5\%$ from the peak locations (relative retention times) in the reference chromatogram. In some embodiments, the relative peak intensities in the HPLC chromatogram may vary by no more than $\pm 10\%$ from the peak intensities in the reference chromatogram.

[026] “Therapeutically effective amount” means the amount of an active substance that, when administered to a subject for treating a disease, disorder, or other undesirable medical condition, is sufficient to have a beneficial effect with respect to that disease, disorder, or condition. The therapeutically effective amount will vary depending on the chemical identity and formulation form of the active substance, the disease or condition and its severity, and the age, weight, and other relevant characteristics of the patient to be treated. Determining the therapeutically effective amount of a given active substance is within the ordinary skill of the art and typically requires no more than routine experimentation.

[027] The term “substantially free” means that the ingredient indicated is not present, or is present in only insignificant amounts. In one embodiment, “substantially free” means less than about 10% (e.g. less than about 10 wt. %). In other embodiments, “substantially free” means less than about 5% (e.g. less than about 5 wt. %), less than about 2% (e.g. less than about 2 wt. %), or less than about 1% (e.g. less than about 1 wt. %), or about 0% (e.g. about 0 wt. %). For example, an emulsion that is substantially free of an emulsifier does not contain any emulsifier in a substantial amount (e.g. contains less than about 10 wt. %, less than about 5 wt. %, less than about 2 wt. %, or less than about 1 wt. % of an emulsifier; or contains 0 wt. % of an emulsifier). Similarly, an oleogel that is substantially free of solid birch bark extract particles having a size greater than about 50 μm does not contain any solid birch bark extract particles having a size greater than about 50 μm in a substantial amount (e.g. contains less than about 10 wt. %, less than about 5 wt. %, less than about 2 wt. %, or less than about 1 wt. % of solid birch bark extract particles having a size greater than about 50 μm ; or contains 0 wt. % of solid birch bark extract particles having a size greater than about 50 μm).

Solid Birch Bark Extracts

[028] The present disclosure provides solid birch bark extracts that may be formulated in clinically-advantageous oleogels. Without being bound by any theory, it is believed that the chemical composition and the particle morphology of the solid birch bark extracts produced according to the methods of the present disclosure are responsible these advantages.

[029] The solid birch bark extracts of the present disclosure may be characterized on the basis of their chemical composition. In some embodiments, the solid birch bark extracts of the present disclosure comprise lupane and oleanane triterpenes. In particular, the birch bark extracts may contain the lupane triterpenes: betulin, lupeol, and betulinic acid, and the oleanane triterpenes: erythrodiol and oleanolic acid.

[030] The presence of a particular triterpene in the birch extract may be determined using relative retention times obtained from HPLC chromatography. The following HPLC method is used to determine the chemical composition of the solid birch bark extracts of the present disclosure:

Time (min)	Acetonitrile	Water	H ₃ PO ₄ (v/v)	Flow rate (ml/min)
0	30	70	0.1%	0.6
5	30	30	0.1%	0.6
15	98	2	0.1%	0.6
20	98	2	0.1%	0.6
21	30	70	0.1%	0.6
30	30	70	0.1%	0.6

Column: Phenomenex Kinetix C18, 2.6 μ m, 150 x 2.1 mm.

[031] A Phenomenex Kinetix C18 column is exemplary of an HPLC column having a C-18 modified silica stationary phase on core-shell silica solid support (2.6 μ m HILIC 100Å) with the above noted column dimensions.

[032] When a solid birch bark extract is subjected to the above-mentioned HPLC chromatography method betulinic acid has a relative retention time of about 0.75 – 0.90, oleanolic acid has a relative retention time of about 0.84-0.97, betulin has a relative retention time of 1.00, erythrodiol has a relative retention time of about 1.25-1.40, and lupeol has a relative retention time of about 3.50-4.15.

[033] In certain embodiments, the solid birch bark extract has an HPLC chromatogram substantially identical to Figure 1.

[034] In some embodiments, the solid birch bark extract comprises at least about 50 wt. %, at least about 55 wt. %, at least about 60 wt. %, at least about 65 wt. %, at least about 70 wt. %, at least about 75 wt. %, at least about 80 wt. %, at least about 85 wt. %, or at least about 90% by weight betulin and one or more triterpenes. In some embodiments, the one or more

triterpenes is selected from the group consisting of betulinic acid, oleanolic acid, erythrodiol and lupeol.

[035] In some embodiments, the solid birch bark extract comprises at least one of the following substances: 3- β -caffeoyl betulin, acetate of the methylester of betulinic acid, acetyloleanolic acid, allobetulin, betulinic aldehyde, betulonic acid, betulonic aldehyde, lupane-3 β ,20,28-triol, lupane-3 β ,20-diol (monogynol), oleanolic aldehyde, sitosterol, ursolic acid, or β -amyrin

[036] The solid birch bark extract of the present disclosure may be characterized by the particle size of the particles of the solid birch bark extract. In some embodiments, the average particle size of the particles of the solid birch bark extract is less than about 100 μm , less than about 90 μm , less than about 80 μm , less than about 70 μm , less than about 60 μm , less than about 50 μm , less than about 40 μm , less than about 30 μm or less than about 25 μm .

[037] In other embodiments, the solid birch bark extract of the present disclosure is substantially free of solid birch bark extract particles having a particle size greater than about 30 μm , greater than about 40 μm , greater than about 50 μm , greater than about 60 μm , greater than about 70 μm , greater than about 80 μm , greater than about 90 μm or greater than about 100 μm .

[038] In preferred embodiments, the solid birch bark extracts are derived from *Betula pendula* Roth and *Betula pubescens* Ehrh as well as hybrids of both species.

Solid Birch Bark Extract Manufacturing Processes

[039] The present disclosure provides methods for preparing solid birch bark extracts that may be formulated in clinically-advantageous oleogels. In general, the methods include the steps of obtaining birch trees, stripping and processing the bark from said birch trees, contacting the processed birch bark with a suitable solvent to provide an extraction solution comprising betulin and one or more triterpenes, and isolating and drying the birch bark extract comprising betulin and one or more triterpenes from the extraction solution. In some embodiments, the isolated the birch bark extract is in the form of a solid.

[040] In some embodiments, the method comprises: (a) contacting birch bark with a pharmaceutically acceptable solvent to form a extraction solution comprising betulin and one

or more triterpenes; (b) separating the birch bark from the extraction solution; (c) cooling the extraction solution whereby a portion of the betulin and one or more triterpenes crystallizes from the cooled extraction solution; (d) separating the crystallized betulin and one or more triterpenes from the cooled extraction solution; and (e) drying the separated, crystallized betulin and one or more triterpenes to provide the solid birch bark extract.

[041] In some embodiments, the birch bark used in step (a) is first processed to improve the extraction efficiency of betulin and one or more triterpenes from the birch bark. For example, experiments have shown that the yield of the dry extract from birch bark is highest when the extracted birch bark particles are small. In certain embodiments, the particle size of the birch bark is reduced prior to step (a) using a mill, or other suitable equipment, until the resulting birch bark passes through a sieve with a mesh size of about 1.25 mm.

[042] In some embodiments, the birch bark is contacted with a pharmaceutically acceptable solvent at a temperature of between about 50°C and about 200°C. In other embodiments, the temperature is between about 60°C and about 100°C, between about 60°C and about 110°C, between about 60°C to about 120°C, between about 60°C to about 130°C, between about 70°C to about 130°C, between about 80°C to about 130°C, between about 90°C to about 130°C, or between about 100°C and 150°C. In certain further embodiments, the temperature is between about 115°C and 130°C.

[043] In some embodiments, the birch bark is contacted with a pharmaceutically acceptable solvent at a pressure between about 2 to 10 bars. In one embodiment, the pressure is about 4.5 bars.

[044] Pharmaceutically acceptable solvents are known to those of skill in the art and include hydrocarbons, alcohols, ketones, ethers esters, sulfoxides, etc. Examples of pharmaceutically acceptable solvents include, without limitation, 1-butanol, 1-pentanol, 1-propanol, 2-butanol, 2-methyl-1-propanol, 2-propanol, 3-methyl-1-butanol, acetic acid, acetone, anisole, butyl acetate, dimethyl sulfoxide, ethanol, ethyl acetate, ethyl ether, ethyl formate, formic acid, heptane, hexane, isobutyl acetate, isopropyl acetate, methyl acetate, methylethyl ketone, methylisobutyl ketone, pentane, propyl acetate, tert-butylmethyl ether, diisopropyl ether, methyl tert-butyl ether, methylisopropyl ketone, and methyltetrahydrofuran.

[045] In some embodiments, two or more pharmaceutically acceptable solvents are mixed to provide the pharmaceutically acceptable solvent.

[046] In some embodiments, the birch bark is contacted with n-heptane at a temperature of between about 60°C and 130°C for between 8 to 12 minutes.

[047] In some embodiments, the (a) contacting birch bark with a pharmaceutically acceptable solvent to form an extraction solution comprising betulin and one or more triterpenes and (b) separating the birch bark from the extraction solution are run using a continuous extraction process (Shown in Figure 2).

[048] In one embodiment of the continuous extraction process, birch bark is mixed with n-heptane in a proportion of about 1:14 to 1:16 (w/v). The mixture is extracted continuously using n-heptane as extraction solvent at a temperature of about 115-130°C and at a pressure of about 4.5 bar (Nitrogen). At the lower end of the sedimentation vessel the extracted cork is rinsed down by cold n-heptane, and at the upper end of the sedimentation vessel the extract (hot solution of triterpenes in n-heptane) is filtrated (10 µm filter) and transported through a second filter (1 µm filter) to the crystallisation step.

[049] In some embodiments, the cooling of step (c) is carried out at a temperature of about -20°C to about 35°C, about -15°C to about 35°C, -10°C to about 35°C, about -5°C to about 35°C, or about 0°C to about 35°C. In other embodiments, in step (c) the cooled extraction solution is supersaturated at least about 1-fold, about 2-fold, about 3-fold, about 4-fold, about 5-fold or about 6-fold. In preferred embodiments, in step (c) the cooled extraction solution is supersaturated about 2-fold or about 5-fold.

[050] In some embodiments, the drying step (e) is carried out under vacuum at a temperature of at least 50°C, at least 60°C, at least 70°C, at least 80°C or at least 85°C. In certain embodiments, the drying step (e) is carried out at a temperature of about 65°C to about 75°C, about 70°C to about 80°C, about 75°C to about 85°C, or about 85°C to about 95°C.

[051] In some embodiments, the drying step (e) is carried out under vacuum at a pressure of less than about 90 mbar, less than about 80 mbar, less than about 70 mbar, less than about 60 mbar or less than about 50 mbar. In certain embodiments, drying step (e) reduces the amount of the pharmaceutically acceptable solvent in the dry birch bark extract to less than about 0.5% (weight/weight).

[052] The present invention also provides the solid birch bark extracts prepared according to the processes set forth above.

Compositions comprising Solid Birch Bark Extracts

[053] The present disclosure provides clinically-advantageous wound-healing formulations comprising solid birch bark extracts. The compositions include oleogels, emulsions, foams, and oleogel-impregnated sterile wound dressings. The compositions are useful as topical wound healing agents.

[054] Gels are finely dispersed systems comprising a liquid phase and a solid phase. The solid phase forms a coherent three-dimensional framework and the two phases permeate one another. Oleogels are hydrophobic gels based on a nonpolar liquid (for example, an oil, a wax, or a paraffin) to which a gel-forming agent is added to achieve the desired physical properties.

[055] The present disclosure provides oleogels comprising a nonpolar liquid and an oleogel-forming agent. Suitable nonpolar liquids for use in oleogels of the present disclosure include, for example, plant, animal, or synthetic oils, waxes, and paraffins. In some embodiments, the nonpolar liquid is a vegetable oil selected from the group consisting of: castor oil, peanut oil, jojoba oil, sunflower oil, olive oil, avocado oil, and almond oil. In a preferred embodiment, the nonpolar liquid is sunflower oil.

[056] In some embodiments, the nonpolar liquid comprises at least one triglyceride. In certain embodiments, the at least one triglyceride is Miglyol. In other embodiments, the nonpolar liquid comprises at least one C7 or greater hydrocarbon. In certain embodiments, the at least one C7 or greater hydrocarbon is a paraffin.

[057] In some embodiments, the nonpolar liquid used in the oleogel has a peroxide value of less than about 15, less than about 10, less than about 5, less than about 4, less than about 3 or less than about 2. In certain embodiments, the nonpolar liquid has a peroxide value of no more than about 3. The term “peroxide value” as used herein means the peroxide value is determined according to Ph. Eur. 2.5.5.

[058] The present disclosure provides methods of making oleogels. In some embodiments, after the solid birch bark extract is dried, about 1 wt% to about 20 wt% of the dried solid birch bark extract is dispersed in nonpolar liquid to form an oleogel. In certain embodiments, the nonpolar liquid is sunflower oil.

[059] In certain embodiments, the oleogel is sterile. The oleogel may be sterilized by suitable methods known to those skilled in the art, for example using ionizing radiation such as electron

beam (EB), X-ray, or gamma . In some embodiments, the oleogel is sterilized by ionizing (e.g., gamma) irradiation at doses less than about 40 kGy, less than about 35 kGy, less than about 30 kGy, less than about 25 kGy, less than about 20 kGy, or less than about 15 kGy. In certain embodiments, the oleogel is sterilized by ionizing (e.g., gamma) irradiation at doses ranging from about 5 to about 25 kGy, about 9 to about 25 kGy, about 10 to about 25 kGy, about 11 to about 25 kGy, about 12 to about 25 kGy, or from about 11 to about 20 kGy.

[060] In some embodiments, the oleogel comprises between about 1 wt. % and about 30 wt. % solid birch bark extract dispersed in about 70 wt. % to about 99 wt. % of one or more nonpolar liquids, wherein the oleogel contains at least one oleogel forming agent in addition to the solid birch bark extract particles. In some embodiments, the oleogel comprises between about 1 wt. % and about 20 wt. % solid birch bark extract dispersed in about 80 wt. % to about 99 wt. % of one or more nonpolar liquids, wherein the oleogel contains at least one oleogel forming agent in addition to the solid birch bark extract particles.

[061] In other embodiments, the oleogel comprises between about 1 wt. % and about 30 wt. % solid birch bark extract particle dispersed in about 70 wt. % to about 99 wt. % of one or more nonpolar liquids, wherein the dispersed solid birch bark extract particles are the only oleogel forming agent in the oleogel. In some embodiments, the oleogel comprises between about 1 wt. % and about 20 wt. % solid birch bark extract dispersed in about 80 wt. % to about 99 wt. % of one or more nonpolar liquids, wherein the oleogel contains at least one oleogel forming agent in addition to the solid birch bark extract particles.

[062] In certain embodiments, the oleogel comprises: about 5 wt. % solid birch bark extract particle dispersed in about 95 wt. % of one or more nonpolar liquids; about 10 wt. % solid birch bark extract particle dispersed in about 90 wt. % of one or more nonpolar liquids; about 15 wt. % solid birch bark extract particle dispersed in about 85 wt. % of one or more nonpolar liquids; or about 20 wt. % solid birch bark extract particle dispersed in about 80 wt. % of one or more nonpolar liquids.

[063] In the foregoing embodiments, the amount of solid birch bark extract particle (for example, about 1 wt. % and about 20 wt. %) includes up to about 0.5 wt % of solid birch bark extract particles that are dissolved in the nonpolar liquid.

[064] Stressing an oleogel by centrifugation provides information on the tendency of the oleogel to separate (or segregate) the nonpolar liquid. In some embodiments of the oleogels of

the present disclosure, the segregation of the nonpolar liquid from the oleogel is less than about 10%, about 9%, about 8%, about 7%, about 6%, about 5%, about 4%, about 3%, about 2%, about 1%, about 0.5%, or about 0.1% after centrifuging at 25°C for about 30 minutes at 2750 g (e.g., 4,400 rpm).

[065] The oleogels of the present disclosure may be characterized by their viscosity. In certain embodiments, the viscosity at 200/s of the oleogel ranges from about 0.5 to about 4.0 Pa.s and the thixotropy value of the oleogel ranges from about 200 to about 1200 Pa.s as measured according to the rotating viscometer method described in Ph. Eur. 2.2.10 using a cone-plate viscometer.

[066] The oleogels of the present disclosure may be also characterized by their consistency values as determined using a texture analyzer. Texture analysis is primarily concerned with measurement of the mechanical properties of a product calculated from the results of a two cycle texture profile analysis test. Texture analysers perform this test by applying controlled forces to the product and recording its response in the form of force, deformation and time.

[067] In present disclosure, consistency is the force [mN] needed to penetrate 1 cm into a sample (e.g., oleogel) as measured using a material testing apparatus (Texture analyser) with a cylindrical penetrating object (0.5 inch = 1.27 cm). The speed of penetration is 0.4 mm/s.

[068] The oleogels of the present disclosure may be distinguished from simple thickened mixtures of the solid birch bark extracts on the basis of consistency values. Specifically, the oleogels of the present disclosure may be characterized by having a consistency value of less than about 3000 mN, and at least about 250 mN as measured by a texture analyzer. In some embodiments, the consistency value of the oleogels is between about 300-2000 mN as measured by a texture analyser.

[069] The present disclosure also provides methods for preparing oleogels. In some embodiments, the method comprises dispersing solid birch bark extract in a suitable nonpolar liquid.

[070] In some embodiments, a sterile wound dressing comprising a pad and a therapeutically effective layer comprising the oleogel of the present disclosure is provided. In certain embodiments, the pad is an adsorbent pad. In other embodiments, the pad is any solid material suitable for covering a wound, including cotton gauze. In some embodiments, the

therapeutically effective layer of oleogel is applied to one or more surfaces of the pad (e.g., the surface of the pad intended to contact the skin or wound of the patient directly). In other embodiments, the pad is immersed in the oleogel such that the oleogel is disposed on at least a portion of the outer surfaces of the pad, and optionally in at least portions of the interior of the pad.

[071] In some embodiments, the pad can comprise a material suitable for dressing a wound that dissolve when used, releasing the oleogel. Such materials can include absorbable materials comprising collagen, alginate, etc.

[072] The term emulsion relates to heterogeneous systems consisting of two liquids that are not miscible with each other or only miscible to a limited extent, which are typically designated as phases. In an emulsion, one of the two liquids is dispersed in the other liquid in the form of minute droplets.

[073] In some embodiments, an emulsion comprising the solid birch bark extract of the present disclosure is provided. Other embodiments provide emulsions comprising the oleogels of the present disclosure.

[074] In some embodiments, the emulsions of the present disclosure include an emulsifier. In certain embodiments, the emulsifier is (hydropropyl)methyl cellulose. In certain other embodiments, the emulsions are substantially free of an emulsifier.

[075] For treating skin wounds, foams may be superior to oleogels because foams can be applied to wounds almost touchless whereas the application of an oleogel requires touch. Foams are generally based on emulsions where a propellant is mixed with the dispersed lipid phase of an emulsion.

[076] The present disclosure provides foams comprising a solid birch bark extract-containing emulsion as described above.

[077] In certain embodiments, the foam comprises an oleogel consisting of about 5 wt. % to about 10 wt. % solid birch bark extract and the emulsion is a water-in-oil emulsion consisting of the oleogel and about 20 wt. % to about 30 wt. % of water.

[078] In certain other embodiments, the foam comprises an oleogel consisting of about 7 wt. % solid birch bark extract and the emulsion is a water-in-oil emulsion consisting of the oleogel and about 25 wt. % of water.

[079] In certain embodiments, the foams of the present disclosure further comprise an emulsifier. Emulsifiers are well known in the art as substances that stabilize emulsions, and include surfactants. Emulsifiers useful in the present invention are those that are acceptable for pharmaceutical use, particularly for skin or wound contact. In some embodiments, suitable emulsifiers include emulsifying waxes, cetearyl alcohol, polysorbate 20, cetareth 20, etc. In certain further embodiments, the emulsifier is selected from the group consisting of phosphatidyl choline, polyglyceryl-3-methyl glucose distearate, and combinations thereof.

[080] In certain embodiments, the oleogels, emulsions, and foams of the present disclosure further comprise a disinfectant. In certain further embodiments, the disinfectant is selected from the group consisting of ethanol, propan-1-ol (n-propanol), and propan-2-ol (isopropanol). In other embodiments, the oleogels, emulsions, and foams of the present disclosure further comprise an antibiotic, particularly lipophilic antibiotics such as fluoroquinolones, macrolides, tigecycline, lincosamides, rifampin, linezolid, tetracyclines, and chloramphenicol.

[081] In certain embodiments, the foams of the present disclosure possess certain physical properties. In some embodiments, the foam index is greater than about 2. In other embodiments, the emulsion used in the foam exhibits an interfacial surface tension of greater than about 4 nM/m using methods known in the art. The term “foam index” as used herein is the ratio of the foamed and unfoamed densities of the specified material. For example, the foam index of a particular foam is the ratio of the density of the foamed material to the density of the material prior to foaming.

[082] The present disclosure also provides for pressurized containers filled with an emulsion of the present invention and a pharmaceutically acceptable propellant whereby the emulsion forms a foam upon decanting at least a portion of the mixture from the container.

Methods of using Solid Birch Bark Extracts

[083] The present disclosure also provides methods of treating a wound in a patient by topically administering an effective amount of an oleogel, an emulsion or a foam of the present disclosure to at least a portion of the wound.

[084] In certain embodiments, the wound treated is selected from the group consisting of burns (such as mild to severe burns), surgical skin lesions, superficial injuries, chronic wounds (such as pressure ulcers, diabetic foot ulcers, chronic venous ulcers, artery insufficiency ulcers, etc.), aesthetic skin treatments (such as ablative laser skin treatments, chemical peels, dermabrasion, etc.), wounds resulting from adverse drug reactions (such as toxic epidermal necrolysis, Lyell syndrome, Stevens-Johnson syndrome, radiation-induced dermatitis, chemotherapy-induced dermatitis, etc.), rare skin diseases (such as epidermolysis bullosa, pemphigus vulgaris, pemphigoid, bullous pemphigoid, pemphigus foliaceus, pyoderma gangrenosa, etc.) and combinations thereof.

[085] The present disclosure also provides methods of treating epidermolysis bullosa in a patient in need thereof comprising topically administering an effective amount of an oleogel, an emulsion or a foam of the present disclosure to an area of epidermolysis bullosa of the patient.

INCORPORATION BY REFERENCE

[086] All references, articles, publications, patents, patent publications, and patent applications cited herein are incorporated by reference in their entireties for all purposes. However, mention of any reference, article, publication, patent, patent publication, and patent application cited herein is not, and should not be taken as acknowledgment or any form of suggestion that they constitute valid prior art or form part of the common general knowledge in any country in the world.

CLAIMS

What is claimed is:

1. A solid birch bark extract comprising at least about 70 wt. % betulin and one or more triterpenes selected from the group consisting of betulinic acid, oleanolic acid, erythrodiol, and lupeol.
2. The solid birch bark extract of claim 1, wherein the solid birch extract has an HPLC chromatogram substantially identical to Figure 1 when chromatographed according to the following HPLC method:

Time (min)	Acetonitrile	Water	H ₃ PO ₄ (v/v)	Flow rate (ml/min)
0	30	70	0.1%	0.6
5	30	30	0.1%	0.6
15	98	2	0.1%	0.6
20	98	2	0.1%	0.6
21	30	70	0.1%	0.6
30	30	70	0.1%	0.6

Column: C18, 2.6 μ m, 150 x 2.1 mm; detector: 210 and 320 nm;

wherein the betulinic acid has a relative retention time of about 0.75 – 0.90, the oleanolic acid has a relative retention time of about 0.84-0.97, the betulin has a relative retention time of 1.00, the erythrodiol has a relative retention time of about 1.25-1.40, and the lupeol has a relative retention time of about 3.50-4.15.

3. The solid birch bark extract of claim 1 or claim 2, prepared by a process comprising:
 - (a) contacting birch bark with a pharmaceutically acceptable solvent, thereby forming an extraction solution comprising betulin and one or more triterpenes;
 - (b) separating the birch bark from the extraction solution;
 - (c) cooling the extraction solution whereby a portion of the betulin and one or more triterpenes crystallizes from the cooled extraction solution;
 - (d) separating the crystallized betulin and one or more triterpenes from the cooled extraction solution;

- (e) drying the separated, crystallized betulin and one or more triterpenes to provide the solid birch bark extract;
wherein if after drying, about 1 wt.% to about 20 wt.% of the dried solid birch bark extract of step (e) dispersed in refined sunflower oil forms an oleogel.
4. The solid birch bark extract of claim 3, wherein one or more of steps (a), (c), and (e) is carried out under the following conditions:
- (i) said contacting of step (a) is carried out at a temperature of about 60° C to about 130° C;
 - (ii) said cooling of step (c) is carried out at a temperature of about -20° C to about 35° C;
 - (iii) said drying of step (e) is carried out at a temperature of about 75° C to about 95° C at a pressure of less than about 70 mbar.
5. The solid birch bark extract of claim 3, wherein two or more of steps (a), (c), and (e) are carried out under the following conditions:
- (i) said contacting of step (a) is carried out at a temperature of about 60° C to about 130° C;
 - (ii) said cooling of step (c) is carried out at a temperature of about -20° C to about 35° C;
 - (iii) said drying of step (e) is carried out at a temperature of about 75° C to about 95° C at a pressure of less than about 70 mbar.
6. The solid birch bark extract of claim 3, wherein steps (a), (c), and (e) are carried out under the following conditions:
- (i) said contacting of step (a) is carried out at a temperature of about 60° C to about 130° C;
 - (ii) said cooling of step (c) is carried out at a temperature of about -20° C to about 35° C;
 - (iii) said drying of step (e) is carried out at a temperature of about 75° C to about 95° C at a pressure of less than about 70 mbar.
7. The solid birch bark extract of any one of claims 3-6, wherein the pharmaceutically acceptable solvent is a hydrocarbon or alcohol, or combinations thereof.

8. The solid birch bark extract of any one of claims 3-7, wherein the pharmaceutically acceptable solvent is a hydrocarbon selected from the group consisting of n-pentane, n-hexane, or n-heptane.
9. The solid birch bark extract of claim 8, wherein the pharmaceutically acceptable solvent is n-heptane, and said contacting is carried out at a temperature of about 60 °C to about 130 °C for about 8-12 minutes.
10. The solid birch bark extract of any one of claims 3-9, wherein in step (c) the cooled extraction solution is supersaturated at least about 2-fold.
11. The solid birch bark extract of claim 10, wherein the cooled extraction solution is supersaturated about 5-fold.
12. The solid birch bark extract of any of claims 3-11, wherein the amount of residual extraction solvent is no more than about 0.5 wt.%.
13. An oleogel comprising about 1 wt. % to about 20 wt. % of particles of the solid birch bark extract of any one of claims 1-12, having an average particle size of less than about 50 µm, dispersed in about 80 wt.% to about 99 wt.% of one or more nonpolar liquids.
14. The oleogel of claim 13, wherein the dispersed solid birch bark extract particles are the only oleogel forming agent in the oleogel.
15. The oleogel of claim 13, comprising about 10 wt. % of particles of the solid birch bark extract.
16. The oleogel of claim 13 or 15, wherein the nonpolar liquid comprises at least one triglyceride.
17. The oleogel of claim 13 or 15, wherein the nonpolar liquid comprises at least one C7 or greater hydrocarbon.

18. The oleogel of claim 16, wherein the nonpolar liquid comprises one or more vegetable oils.
19. The oleogel of claim 18, wherein the nonpolar liquid comprises sunflower oil.
20. The oleogel of any one of claims 13-19, wherein the nonpolar liquid has a peroxide value less than about 10.
21. The oleogel of claim 20, wherein the peroxide value is no more than about 3.
22. The oleogel of any one of claims 13-21, wherein the oleogel is substantially free of solid birch bark extract particles having a size greater than about 50 μm .
23. The oleogel of any one of claims 13-22, wherein the oleogel is sterile.
24. The oleogel of claim 23, wherein the oleogel is sterilized by ionizing irradiation at doses less than about 20 kGy.
25. The oleogel of claim 24, wherein the oleogel is sterilized by ionizing irradiation at doses ranging from about 11 to about 20 kGy.
26. The oleogel of any one of claims 13-25, wherein the segregation of the nonpolar liquid from the oleogel is less than about 10% after centrifuging at 25°C for 30 min at 2750 g.
27. The oleogel of any one of claims 13-26, wherein the viscosity at 200/s of the oleogel ranges from about 0.5 to about 4.0 Pa.s and the thixotropy value of the oleogel ranges from about 200 to about 1200 Pa.s as measured according to the rotating viscometer method described in Ph. Eur. 2.2.10 using a cone-plate viscometer.
28. The oleogel of any one of claims 13-27, wherein the consistency of the oleogel ranges from about 300-3000 mN as measured according using a texture analyser.

29. The oleogel of any one of claims 13-28, further comprising a disinfectant.
30. The oleogel of claim 29, wherein the disinfectant is selected from the group consisting of ethanol, n-propanol, and isopropanol.
31. The oleogel of any one of claims 13-28, further comprising a lipophilic antibiotic.
32. A sterile wound dressing comprising:
(a) a pad, and
(b) a therapeutically effective layer comprising the oleogel of any of claims 13-31 disposed on at least one surface of the pad.
33. The sterile wound dressing of claim 32, wherein the pad is an absorbent pad.
34. A process for preparing the solid birch bark extract of claim 1 or claim 2 comprising:
(a) contacting birch bark with a pharmaceutically acceptable solvent, thereby forming an extraction solution comprising betulin and one or more triterpenes;
(b) separating the birch bark from the extraction solution;
(c) cooling the extraction solution whereby a portion of the betulin and one or more triterpenes crystallizes from the cooled extraction solution;
(d) separating the crystallized betulin and one or more triterpenes from the cooled extraction solution;
(e) drying the separated, crystallized betulin and one or more triterpenes to provide the solid birch bark extract;
wherein after drying, about 1 wt.% to about 20 wt.% of the dried solid birch bark extract of step (e) dispersed in refined sunflower oil forms an oleogel.
35. The process of claim 34, wherein one or more of steps (a), (c), and (e) is carried out under the following conditions:
(i) said contacting of step (a) is carried out at a temperature of about 60° C to about 130° C;

- (ii) said cooling of step (c) is carried out at a temperature of about -20° C to about 35° C;
 - (iii) said drying of step (e) is carried out at a temperature of about 75° C to about 95° C at a pressure of less than about 70 mbar.
- 36. The process of claim 34, wherein two or more of steps (a), (c), and (e) are carried out under the following conditions:
 - (i) said contacting of step (a) is carried out at a temperature of about 60° C to about 130° C;
 - (ii) said cooling of step (c) is carried out at a temperature of about -20° C to about 35° C;
 - (iii) said drying of step (e) is carried out at a temperature of about 75° C to about 95° C at a pressure of less than about 70 mbar.
- 37. The process of claim 34, wherein steps (a), (c), and (e) are carried out under the following conditions:
 - (i) said contacting of step (a) is carried out at a temperature of about 60° C to about 130° C;
 - (ii) said cooling of step (c) is carried out at a temperature of about -20° C to about 35° C;
 - (iii) said drying of step (e) is carried out at a temperature of about 75° C to about 95° C at a pressure of less than about 70 mbar.
- 38. The process of claim 34, wherein the pharmaceutically acceptable solvent is a hydrocarbon selected from the group consisting of n-pentane, n-hexane, or n-heptane.
- 39. The process of any one of claims 34-38, wherein the pharmaceutically acceptable solvent is n-heptane, and said contacting is carried out at a temperature of about 115 °C to about 130 °C for about 8-12 minutes.
- 40. The process of any one of claims 34-39, wherein in step (c) the cooled extraction solution is supersaturated at least about 2-fold.

41. The process of claim 40, wherein the cooled extraction solution is supersaturated about 5-fold.
42. The process of any of claims 34-41, wherein the amount of residual extraction solvent is no more than about 0.5 wt.%.
43. An emulsion comprising the solid birch bark extract of any of claims 1-12.
44. An emulsion comprising the oleogel of any of claims 13-31.
45. The emulsion of claim 43 or 44, wherein said emulsion is a water-in-oil emulsion.
46. The emulsion of claim 45, substantially free of emulsifier.
47. The emulsion of any one of claims 43-46, consisting essentially of the solid birch bark extract or oleogel, oil, and water.
48. The emulsion of any one of claims 44-46, consisting of the oleogel, oil and water.
49. The emulsion of any one of claims 43-46, further comprising a disinfectant.
50. The emulsion of claim 49, wherein the disinfectant is selected from the group consisting of ethanol, n-propanol, and isopropanol.
51. The emulsion of any one of claims 43-46, further comprising a lipophilic antibiotic.
52. A foam comprising the emulsion of any one of claims 43-51.
53. The foam of claim 52, further comprising a disinfectant.
54. The foam of claim 53, wherein the disinfectant is selected from the group consisting of ethanol, n-propanol, and isopropanol.
55. The foam of claim 52, further comprising a lipophilic antibiotic.

56. The foam of any of claims 52-55, wherein the interfacial surface tension of the emulsion is greater than about 4 mN/m.
57. The foam of any of claims 52-56, wherein the foam index is greater than about 2.
58. The foam of claim 52, wherein the oleogel consists of about 5 wt.% to about 10 wt.% solid birch bark extract and the emulsion is a water-in-oil emulsion consisting of the oleogel and about 20 wt.% to about 30 wt.% of water.
59. The foam of claim 58, wherein the oleogel consists of about 7 wt.% solid birch bark extract and the amount of water in the emulsion is about 25 wt.%.
60. The foam of any of claims 52-56, further comprising an emulsifier selected from the group consisting of phosphatidyl choline, polyglyceryl-3-methyl glucose distearate, and combinations thereof.
61. A pressurized container filled with a mixture comprising the emulsion of any one of claims 43-51 and a pharmaceutically acceptable propellant, whereby the emulsion forms a foam upon decanting at least a portion of the mixture from the container.
62. A method of treating a wound in a patient comprising topically administering the oleogel of any one of claims 13-31 to at least a portion of the wound.
63. The oleogel of any one of claims 13-31, for use in a method of treating a wound.
64. The method or oleogel for use of claim 63, wherein the wound is selected from the group consisting of a burns, surgical skin lesions, superficial injuries, chronic wounds, pressure ulcers, diabetic foot ulcers, chronic venous ulcers, artery insufficiency ulcers, wounds from aesthetic skin treatments, wounds from ablative laser skin treatments, wounds from chemical peels, wounds from dermabrasion, wounds resulting from adverse drug reactions, wounds resulting from toxic epidermal necrolysis, wounds resulting from Lyell syndrome, wounds resulting from Stevens-Johnson syndrome, and wounds resulting from radiation dermatitis.

65. A method of treating a wound in a patient comprising topically administering the emulsion of any one of claims 43-51 to at least a portion of the wound.
66. The emulsion of any one of claims 43-51, for use in a method of treating a wound.
67. The method or emulsion for use of claim 66, wherein the wound is selected from the group consisting of burns, surgical skin lesions, superficial injuries, chronic wounds, pressure ulcers, diabetic foot ulcers, chronic venous ulcers, artery insufficiency ulcers, wounds from aesthetic skin treatments, wounds from ablative laser skin treatments, wounds from chemical peels, wounds from dermabrasion, wounds resulting from adverse drug reactions, wounds resulting from toxic epidermal necrolysis, wounds resulting from Lyell syndrome, wounds resulting from Stevens-Johnson syndrome, and wounds resulting from radiation dermatitis.
68. A method of treating a wound in a patient comprising topically administering the foam of any one of claims 52-60 to at least a portion of the wound.
69. The foam of any one of claims 52-60, for use in a method of treating a wound.
70. The method or foam for use of claim 69, wherein the wound is selected from the group consisting of burns, surgical skin lesions, superficial injuries, chronic wounds, pressure ulcers, diabetic foot ulcers, chronic venous ulcers, artery insufficiency ulcers, wounds from aesthetic skin treatments, wounds from ablative laser skin treatments, wounds from chemical peels, wounds from dermabrasion, wounds resulting from adverse drug reactions, wounds resulting from toxic epidermal necrolysis, wounds resulting from Lyell syndrome, wounds resulting from Stevens-Johnson syndrome, and wounds resulting from radiation dermatitis.
71. A method of treating epidermolysis bullosa in a patient in need thereof comprising topically administering to an area of epidermolysis bullosa of the patient the oleogel of any one of claims 13-31.

72. The oleogel of any one of claims 13-31, for use in a method of treating epidermolysis bullosa.
73. A method of treating epidermolysis bullosa in a patient in need thereof comprising topically administering to an area of epidermolysis bullosa of the patient the emulsion of any one of claims 43-51.
74. The emulsion of any one of claims 43-51, for use in a method of treating epidermolysis bullosa.
75. A method of treating epidermolysis bullosa in a patient in need thereof comprising topically administering to an area of epidermolysis bullosa of the patient the foam of any one of claims 52-60.
76. The foam of any one of claims 52-60, for use in a method of treating epidermolysis bullosa.




FIG. 1




FIG. 2




FIG. 1