



Office de la Propriété

Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2658824 A1 2008/01/31

(21) **2 658 824**

(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2007/07/26
(87) Date publication PCT/PCT Publication Date: 2008/01/31
(85) Entrée phase nationale/National Entry: 2009/01/23
(86) N° demande PCT/PCT Application No.: US 2007/016853
(87) N° publication PCT/PCT Publication No.: 2008/013914
(30) Priorités/Priorities: 2006/07/26 (US60/833,757);
2007/03/14 (US60/918,225)


(51) Cl.Int./Int.Cl. *G01N 33/53* (2006.01),
G01N 33/543 (2006.01), *G01N 33/564* (2006.01),
C07K 16/18 (2006.01), *C07K 14/775* (2006.01)

(71) Demandeur/Applicant:
INOVA DIAGNOSTICS, INC., US

(72) Inventeurs/Inventors:
IVERSON, MICHAEL G., US;
BINDER, WALTER L., US;
NORMAN, GARY L., US

(74) Agent: SIM & MCBURNEY

(54) Titre : COMPOSITIONS ET PROCEDES DE DIAGNOSTIQUE DE SYNDROMES DE L'ATHEROSCLEROSE AIGUE
CHEZ DES PATIENTS
(54) Title: COMPOSITIONS AND METHODS FOR DIAGNOSING PATIENTS WITH ACUTE ATHEROSCLEROTIC
SYNDROMES



(57) Abrégé/Abstract:

The present invention is in the field of autoimmunity. More specifically, the present invention relates to the detection of autoantibodies to domain 4 of beta 2-glycoprotein I (β_2 -GPI) as an indicator for atherosclerosis.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
31 January 2008 (31.01.2008)

PCT

(10) International Publication Number
WO 2008/013914 A2(51) International Patent Classification:
G01N 33/53 (2006.01)

(72) Inventors; and

(75) Inventors/Applicants (for US only): IVERSON, G. Michael [US/US]; 13784 Boquita Drive, Del Mar, California 92014 (US). BINDER, Walter, L. [US/US]; 1240 India Street, #2108, San Diego, California 92101 (US). NORMAN, Gary, L. [US/US]; 1920 Colina Salida del Sol, San Clemente, California 92673 (US).

(21) International Application Number:
PCT/US2007/016853

(74) Agents: AXFORD, Laurie, A. et al.; Gordon & Rees LLP, 101 West Broadway, Suite 1600, San Diego, California 92101 (US).

(22) International Filing Date: 26 July 2007 (26.07.2007)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL,

(25) Filing Language: English

(26) Publication Language: English


[Continued on next page]

(30) Priority Data:
60/833,757 26 July 2006 (26.07.2006) US
60/918,225 14 March 2007 (14.03.2007) US

(71) Applicant (for all designated States except US): INOVA DIAGNOSTICS, INC. [US/US]; 9900 Old Grove Road, San Diego, California 92131 (US).

(54) Title: COMPOSITIONS AND METHODS FOR DIAGNOSING PATIENTS WITH ACUTE ATHEROSCLEROTIC SYNDROMES

FIGURE 1

(57) Abstract: The present invention is in the field of autoimmunity. More specifically, the present invention relates to the detection of autoantibodies to domain 4 of beta 2-glycoprotein I (β_2 -GPI) as an indicator for atherosclerosis.

WO 2008/013914 A2

WO 2008/013914 A2



PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *without international search report and to be republished upon receipt of that report*
- *with sequence listing part of description published separately in electronic form and available upon request from the International Bureau*

(48) **Date of publication of this corrected version:**

23 October 2008

(15) **Information about Correction:**

see Notice of 23 October 2008

COMPOSITIONS AND METHODS FOR DIAGNOSING PATIENTS WITH ACUTE ATHEROSCLEROTIC SYNDROMES

TECHNICAL FIELD

[0001] The present invention is in the field of autoimmunity. More specifically, the present invention relates to the detection of autoantibodies to domain 4 of beta 2-glycoprotein I (β_2 -GPI) as an indicator for atherosclerosis.

BACKGROUND OF THE INVENTION

[0002] IgA class autoantibodies to beta 2-glycoprotein I (β_2 -GPI) have recently been reported in patients with acute myocardial infarction and also in patients with ischemic stroke (Ranzolin A, et al., *Arg Bras Cardiol.* 83(2):141-4; 137-40 (2004); Kahles T, et al., *Rheumatology* 44(9):1161-5 (2005); Staub HL, et al., *Arg Neuropsiquiat* 61(3B):757-63 (2003)). A striking observation from two of these studies was that the IgA β_2 -GPI autoantibodies were usually detected in patients that were negative for IgA anti-cardiolipin antibodies (ACA) (Ranzolin A, et al., *Arg Bras Cardiol.* 83(2):141-4; 137-40 (2004); Staub HL, *Arg Neuropsiquiat* 61(3B):757-63 (2003)). This finding is in sharp contrast to that observed in patients with anti-phospholipid syndrome (APS), where both anti- β_2 -GPI and anti-cardiolipin antibodies are usually positive.

[0003] β_2 -GPI is a serum protein composed of five homologous domains numbered 1-5 from the N terminus. Domains 1-4 are composed of ~ 60 amino acids that contain a motif characterized by a framework of four conserved cysteine residues, which form two internal disulfide bridges (Lozier, J. et al., *PNAS* 81:3640-3644 (1984)). The fifth domain differs from domains 1-4 in that it contains 82 amino acid residues with six cysteines. The fifth domain contains the phospholipid binding site (Hunt, J., et al., *PNAS* 90:2141-2145 (1993)).

[0004] Conflicting findings have been published concerning the domain specificity of anti- β_2 -GPI autoantibodies. For example, in one study, it was shown that using recombinant β_2 -GPI and β_2 -GPI domain-deleted mutants (Dms) expressed in insect cells, that anti- β_2 -GPI autoantibodies found in serum samples from patients with APS recognize domain 1 of β_2 -GPI (Iverson, GM, et al., *PNAS* 95:15542-15546 (1998)).

[0005] There are also reports that IgG anti- β_2 -GPI autoantibodies in patients with APS recognize epitopes on domains 3, 4 and 5 of β_2 -GPI (Blank, M, et al., *PNAS* 96:5164-8 (1999); Blank, M, et al., *PNAS* 96:5164-8 (1999); Koike T, et al., *J. Autoimmun* 15: 97-100 (2000); Yang CD, et al., *APLAR J Rheumatol* 1:96-100 (1997); Iverson, GM, et al., *J. of Autoimmunity* 18:289-297 (2002); and McNeeley PA, et al., *Thromb Haemost* 86:590-5 (2001)).

[0006] Accordingly, there is a need for an enhanced understanding of the antibody profiles exhibited by APS patients (cardiolipin IgG positive/ β_2 -GPI IgG/IgA positive), and the differences between these profiles and the profiles exhibited by cardiovascular patients with acute ischemic disease (cardiolipin IgG/IgA negative/ β_2 -GPI IgA positive), which is based on differing domain-specificity of the APS and cardiovascular patient's antibodies. The present invention provides such an enhanced understanding, and relates to the finding that IgA anti- β_2 -GPI autoantibodies that bind to domain 4 are found in a high percentage of patients with APS and various cardiovascular diseases.

SUMMARY OF THE INVENTION

[0007] The present invention relates, in part, to a method for diagnosing a subject suspected of having acute atherosclerotic syndrome (ASS) comprising the steps of: preparing an antigen comprising a polypeptide having an epitope from domain 4 of β_2 -GPI; reacting the antigen with a biological sample from the subject; and detecting IgA antibodies in the sample that bind to the antigen.

[0008] The antigen according to the present invention may consist of domain 4 in its entirety, or an antigenic fragment of domain 4, and/or it may include all or a portion of the sequences of domains 2, 3 and 5. Accordingly, the phrase "an epitope from domain 4" intends that the epitope is recognized by an antibody that is selective for domain 4, in that it binds preferentially to domain 4 when compared to the other 4 domains. Thus, the epitope may be a polypeptide consisting of domain 4 (or a fragment thereof) alone, or a combination of domains 2, 3 and 4; 2, 3, 4 and 5; 3 and 4; 4 and 5; 3, 4 and 5; and/or fragments thereof.

[0009] As shown below, the domain 4 sequence consists of 56 amino acids. Given that the minimum number of contiguous amino acids from domain 4 is in the neighborhood of 6 it is also possible to construct an antigen that is a multimer (such as a dimer, trimer, etc.) of the domain 4 epitope, with the repeating units separated by non-interfering linking regions such as

polyglycine and other small nonpolar amino acids. Such linking regions may or may not include the naturally existing flanking sequences of the epitope.

[0010] According to one embodiment, the invention is a method of diagnosing an acute atherosclerotic syndrome in a subject, comprising determining the presence or absence of IgA domain 4-specific anti-beta 2-glycoprotein I (β_2 -GPI) antibodies in said subject, wherein the presence of said IgA anti- β_2 -GPI antibodies indicates that said subject has an acute atherosclerotic syndrome.

[0011] In an alternative embodiment, the invention is a method of diagnosing an acute atherosclerotic syndrome in a subject, comprising the steps of: a. obtaining a sample from a subject suspected of having an acute atherosclerotic syndrome; b. contacting the sample with a β_2 -GPI antigen comprising a domain 4 epitope; and c. detecting the presence or absence of IgA domain 4-specific anti- β_2 -GPI antibodies that bind to the domain 4 epitope; wherein the presence of said IgA domain 4-specific anti- β_2 -GPI antibodies in said sample indicates that said subject has an acute atherosclerotic syndrome.

[0012] In yet another embodiment, the invention is a method of diagnosing an acute atherosclerotic syndrome in a subject, comprising the steps of: a. contacting a sample from a subject suspected of having an acute atherosclerotic syndrome with an epitope derived from domain 4 of a β_2 -GPI antigen comprising the amino acid sequence of SEQ ID NO:5 without linker sequences under conditions suitable to form a complex of the epitope and IgA domain 4-specific anti- β_2 -GPI antibody; and b. detecting the presence or absence of the IgA domain 4-specific anti- β_2 -GPI antibody in the complex, wherein the presence of said domain 4-specific IgA anti- β_2 -GPI antibodies in said subject indicates that said subject has an acute atherosclerotic syndrome.

[0013] The method for detecting the IgA anti- β_2 -GPI antibodies can be by any known method, such as with a labeled anti-IgA antibody in a variety of well known assay formats such as an enzyme-linked immunosorbent assay.

[0014] In one embodiment, the method further includes determining the presence or absence of anticardiolipin (aCL) antibodies in said subject, wherein the presence of said domain 4-specific IgA anti- β_2 -GPI antibodies in said subject and the absence of anticardiolipin (aCL) antibodies indicates that said subject has an acute atherosclerotic syndrome.

[0015] The domain 4 epitope may exist in the form of a variety of different combinations of domains and fragments thereof. For example, the domain 4 epitope may consist of the known domain 4 fragment antigenic sequence that is adjacent to domain 5 (Kasahara, et al.). It may also be in the form of the full domain 4 plus 5 sequences with all or portion of domain 3, or it may be in the form of the full domain 3, 4 and 5 sequences with all or a portion of domain 2, or it may be in the form of adjacent fragments of domain 4 and 5, and so on. However, in a preferred embodiment, domain 1 is completely absent.

[0016] Other aspects of the invention are described throughout the specification.

BRIEF DESCRIPTION OF THE DRAWINGS

[0017] **Figure 1 depicts:** Competitive inhibition of APS sample 6635 binding to β_2 -GPI by Recombinant β_2 -GPI and Dms. A constant amount of antibody was mixed with varying concentrations of inhibitor in wells coated with β_2 -GPI. Recombinant β_2 -GPI and DMs were used as inhibitors. Upper panel measures inhibition of IgG antibodies. Lower panel measures inhibition of IgA antibodies.

[0018] **Figure 2 depicts:** Competitive inhibition of IgA anti- β_2 -GPI of ACS-71 from binding to β_2 -GPI by recombinant β_2 -GPI and deletion mutants. A constant amount of antibody was mixed with varying concentrations of inhibitors in wells coated with β_2 -GPI. Recombinant β_2 -GPI and deletion mutants were used as inhibitors.

[0019] **Figure 3 depicts:** The entire sequence and tertiary structure of β_2 -GPI.

DETAILED DESCRIPTION

[0020] In the description that follows, a number of terms used in the field of molecular biology, immunology and medicine are extensively utilized. In order to provide a clearer and consistent understanding of the specification and claims, including the scope to be given such terms, the following non-limiting definitions are provided.

[0021] When the terms "one," "a," or "an" are used in this disclosure, they mean "at least one" or "one or more," unless otherwise indicated.

[0022] The term "antibody" refers to a molecule which is capable of binding an epitope or antigenic determinant. The term "antibody" includes whole antibodies and antigen-binding

fragments thereof, including single-chain antibodies. Such antibodies include human antigen binding antibody and antibody fragments, including, but not limited to, Fab, Fab' and F(ab')₂, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and fragments comprising either a V_L or V_H domain. The antibodies may be from any animal origin including birds and mammals, e.g., human, murine, rabbit, goat, guinea pig, camel, horse and the like.

[0023] The term "antigen" refers to a molecule capable of being bound by an antibody or a T cell receptor (TCR) if presented by MHC molecules. An antigen may be additionally capable of being recognized by the immune system and/or being capable of inducing a humoral immune response and/or cellular immune response leading to the activation of B- and/or T-lymphocytes. An antigen may have one or more epitopes (B- and T-epitopes). Antigens as used herein may also be mixtures of several individual antigens.

[0024] The term "antigenic determinant" refers to a portion of an antigen that is specifically recognized by either B- or T-lymphocytes. Antigenic determinants or epitopes are those parts of an antigen that are recognized by antibodies, or in the context of an MHC, by T-cell receptors. An antigenic determinant contains one or more epitopes.

[0025] The term "autoantigen" refers to a constituent of self that binds an autoantibody or that induces a cellular response.

[0026] The term "autoantibody" refers to an immunoglobulin, antigen specific B cell surface receptor (surface immunoglobulin), or antigen specific T cell receptor directed against self-protein, carbohydrate or nucleic acid.

[0027] The term "epitope" refers to a portion of an antigen that is recognized by the immune system, specifically by an antibody (e.g., an autoantibody), B-cell, or T cell, and thus the particular domain, region or molecular structure to which the antibody, B-cell or T-cell binds. An antigen may consist of numerous epitopes while a hapten, typically, may possess few epitopes.

[0028] The term "wild-type" refers to a gene or gene product isolated from a naturally occurring source. A wild-type gene is that which is most frequently observed in a population and is thus arbitrarily designed the "normal" or "wild-type" form of the gene or gene product. In contrast, the term "modified" or "mutant" refers to a gene or gene product that displays modifications in sequence and or functional properties (i.e., altered characteristics, e.g.,

hypomethylation) when compared to the wild-type gene or gene product. It is noted that naturally occurring mutants can be isolated; these are identified by the fact that they have altered characteristics, such as physical and biological properties, when compared to the wild-type gene or gene product.

[0029] The term "native protein" refers to a protein that contains only those amino acids found in the protein as it occurs in nature. A native protein may be produced by recombinant means or may be isolated from a naturally occurring source.

[0030] The term "fragment" means a peptide, polypeptide, or compound containing naturally occurring amino acids, non-naturally occurring amino acids or chemically modified amino acids. The fragments may range in size from two amino acid residues to the entire amino acid sequence minus one amino acid.

[0031] The term "subject" refers to an animal, including, but limited to, an ave, ovine, bovine, ruminant, lagomorph, porcine, equine, canine, feline, rodent or primate, for example a human. Typically, the terms "subject" and "patient" are used interchangeably herein in reference to a mammalian subject, particularly a human subject.

[0032] The term "sample" is used in its broadest sense. In one sense, it is meant to include a specimen or culture obtained from any source, as well as biological and environmental samples. Biological samples may be obtained from animals (including humans) and refers to a biological material or compositions found therein, including, but not limited to, bone marrow, blood, serum, platelet, plasma, interstitial fluid, urine, cerebrospinal fluid, nucleic acid, DNA, tissue, and purified or filtered forms thereof.

[0033] The term "serum sample" refers to a biological sample comprising serum. It is understood that a serum sample for use in the present methods may contain other components, in particular blood components. Thus, whole blood samples, or blood samples which have been only partially fractionated or separated but which still contain serum, are considered "serum samples" for purposes of the present invention. One skilled in the art can readily obtain serum samples, for example by using conventional blood drawing techniques. Furthermore, the presence of preservative, anticoagulants or other chemicals in the serum sample should not interfere the detection of IgA β_2 -GPI antibodies.

[0034] The term "control" or "control sample" refers to one or more sample, such as a serum sample, taken from at least one healthy blood donor. It is understood that when the control comprises multiple samples, the IgA β_2 -GPI -specific antibody level can be expressed as the arithmetic mean, median, mode or other suitable statistical measure of the IgA β_2 -GPI -specific antibody level measured in each sample. Multiple control samples can also be pooled, and IgA β_2 -GPI -specific antibody level of the pooled samples can be determined and compared to the subject's sample.

[0035] Atherosclerosis (also referred to as arteriosclerosis, atheromatous vascular disease, arterial occlusive disease) as used herein, refers to a cardiovascular disease characterized by plaque accumulation on vessel walls and vascular inflammation. The plaque consists of accumulated intracellular and extracellular lipids, smooth muscle cells, connective tissue, inflammatory cells, and glycosaminoglycans. Inflammation occurs in combination with lipid accumulation in the vessel wall, and vascular inflammation is with the hallmark of atherosclerosis disease process.

[0036] The term "acute atherosclerotic syndrome" or "ASS" refers to several types of cardiovascular problems including, but not limited to, acute myocardial infarction, acute coronary syndrome, "carotid artery study," and peripheral artery disease.

[0037] β_2 -GPI is a serum protein composed of five homologous domains numbered 1-5 from the N terminus. The primary and predicted tertiary sequence is shown in Figure 3. Domains 1-4 are composed of ~ 60 amino acids that contain a motif characterized by a framework of four conserved cysteine residues, which form two internal disulfide bridges. The fifth domain differs from domains 1-4 in that it contains 82 amino acid residues with six cysteines. The fifth domain contains the phospholipid binding site.

[0038] The amino acid sequence of domain 1 (SEQ ID NO.:2) as shown in Figure 3, starting with the N-terminal end and ending with the small sequence that links domain 1 to domain 2 (underlined) is as follows:

GRTCPKPDDLPFSTVVPLKTFYEPGEEITYSCKPGYVSRGGMRKFICPLTGLWPIN
TLKCTPRV

[0039] The amino acid sequence of domain 2 (SEQ ID NO.:3) as shown in Figure 3, starting with the N-terminal end of the small sequence that links domain 1 to domain 2 (underlined) and ending with the small sequence that links domain 2 to domain 3 (underlined) is as follows:

TPRVCPFAGILENGAVRYTTFEYPNTISFSCNTGFYLNGADSAKCTEEGKWSP
EPLPVCAPII

[0040] The amino acid sequence of domain 3 (SEQ ID NO.:4) as shown in Figure 3, starting with the N-terminal end of the small sequence that links domain 2 to domain 3 (underlined) and ending with the small sequence that links domain 3 to domain 4 (underlined) is as follows:

APIICPPPSIPTFATLRVYKPSAGNNSLYRDTAVFECLPQHAMFGNDTITCTTHGN
WTKLPECREVK

[0041] The amino acid sequence of domain 4 (SEQ ID NO.:5) as shown in Figure 3, starting with the N-terminal end of the small sequence that links domain 3 to domain 4 (underlined) and ending with the small sequence that links domain 4 to domain 5 (underlined) is as follows:

REVKCPFPSRPDNGFVNYPAKPTLYYKDKATFGCHDGYSLDGPEEIECTKLGNW
SAMPSCKAS

[0042] The amino acid sequence of domain 5 (SEQ ID NO.:6) as shown in Figure 3, starting with the N-terminal end of the small sequence that links domain 4 to domain 5 (underlined) and ending with the C-terminal end is as follows:

KASCKVPVKKATVVYQGERVKIQEKFKNGMLHGDKVSFFCKNKEKKCSYTED
AQCIDGTIEVPKCFKEHSSLAFWKTDASDVKPC

[0043] As used herein, it shall be understood that the term “domain X” alone refers to a polypeptide with the amino acid sequence without the underlined linker sequences identified above, but when terms such as “domain X/Y” or “domain X+Y” are used, it refers to a polypeptide with the amino acid sequence of the two domains linked by the appropriate underlined linker sequence identified above. Likewise, referring to the “domain X amino acid sequence of SEQ. ID NO. A without linker sequences” means the same as saying the

polypeptide that includes all of the SEQ. ID NO. A amino acids, without the underlined linker sequences identified above.

[0044] "Dms" or "domain-deleted mutants" nomenclature for domain deletion mutants uses numbers to indicate the presence of β_2 -GPI domains, while a dash symbolizes the domain is missing. Thus D--345 is the name given to the recombinant protein that contains domains 3, 4 and 5 while lacking domains 1 and 2.

[0045] The β_2 -GPI proteins of the present invention may also be its variants. Unless otherwise indicated, the term " β_2 -GPI" refers both to native β_2 -GPI proteins, as well as variants thereof. As used herein, β_2 -GPI variants are β_2 -GPI proteins which comprises an amino acid sequence having one or more amino acid substitutions, deletions, and/or additions (such as internal additions and/or β_2 -GPI fusion proteins) as compared to the amino acid sequence of a native β_2 -GPI proteins, but which nonetheless retain β_2 -GPI immunologically activity. Such functionally or immunologically equivalent variants may occur as natural biological variations (e.g., polypeptide allelic variants, polypeptide orthologs, and polypeptide splice variants), or they may be prepared using known and standard techniques for example by chemical synthesis or modification, mutagenesis, e.g., site-directed or random mutagenesis, etc. Thus, for example, an amino acid may be replaced by another which preserves the physicochemical character of the β_2 -GPI proteins or its epitope(s), e.g. in terms of charge density, hydrophilicity/hydrophobicity, size and configuration and hence preserve the immunological structure. "Addition" variants may include N- or C-terminal fusions as well as intrasequence insertion of single or multiple amino acids. Deletions may be intrasequence or may be truncations from the N- or C-termini.

[0046] The variants may have from 1 to 3, to 5, to 10, to 15, to 20, to 25, to 50, to 75, or to 100, or more than 100 amino acid substitutions, insertions, additions and/or deletions, wherein the substitutions may be conservative, or non-conservative, or a combination thereof. Additionally, the β_2 -GPI proteins of the present invention may comprise at least 10, at least 12, at least 15, at least 20, at least 25, at least 30, at least 35, at least 40, or at least 50 consecutive amino acid residues of a native β_2 -GPI protein. Such a variant is preferably at least about 50%, at least about 60%, at least about 70%, at least about 80%, as least about 90%, or at least about 95% identical to a native β_2 -GPI proteins. Furthermore, the β_2 -GPI variants may remain immunologically active with an activity of over about 1%, over about 10%, over about 25%,

over about 50%, over about 60%, over about 70%, over about 80%, over about 90%, over about 95%, or over about 100% of the immunological activity of the native protein.

[0047] Conservative modifications to the amino acid sequence of a β_2 -GPI protein generally produce a polypeptide having functional and chemical characteristics similar to those of the original β_2 -GPI proteins. In contrast, substantial modifications in the functional and/or chemical characteristics of a β_2 -GPI protein may be accomplished by selecting substitutions in the amino acid sequence of the β_2 -GPI protein that differ significantly in their effects on maintaining (a) the structure (secondary, tertiary, and/or quaternary) in the area of the substitution or (b) the charge or hydrophobicity of the molecule at the target site. Amino acid sequence modifications can be accomplished by chemical and biological peptide and protein synthetic methods that are well known in the art.

[0048] Desired amino acid substitutions (whether conservative or non-conservative) can be determined by those skilled in the art at the time such substitutions are required. For example, amino acid substitutions can be used to identify important residues, to modulate the biological activity of a β_2 -GPI protein, e.g., the binding interactions with β_2 -GPI-specific antibodies, or to decrease unwanted non-specific binding interactions with other molecules in a sample. Suitable amino acid substitutions include, but are not limited to, substituting Ala with Val, Leu, or Ile; substituting Arg with Lys, Gln, or Asn; substituting Asn with Gln; substituting Asp with Glu; substituting Cys with Ser or Ala; substituting Gln with Asn; substituting Glu with Asp; substituting His with Asn, Gln, Lys, or Arg; substituting Ile with Leu, Val, Met, Ala, Phe, or Norleucine; substituting Leu with Norleucine, Ile, Val, Met, Ala, or Phe; substituting Lys with Arg, 1,4-diamino-butyric acid, Gln, or Asn; substituting Met with Leu, Phe, or Ile; substituting Phe with Leu, Val, Ile, Ala, or Tyr; substituting Pro with Ala; substituting Ser with Thr, Ala, or Cys; substituting Thr with Ser; substituting Trp with Tyr or Phe; substituting Tyr with Trp, Phe, Thr, or Ser; and substituting Val with Ile, Met, Leu, Phe, Ala, or Norleucine. The selection of an amino acid for replacement can also be guided by its hydropathic index and/or hydrophilicity.

[0049] In addition, the polypeptide may be fused to a homologous polypeptide to form a homodimer or to a heterologous polypeptide to form a heterodimer. Heterologous polypeptides include, but are not limited to: an epitope to allow for the detection and/or isolation of a β_2 -GPI fusion polypeptide, such as, polyhistine at either C- or N-terminal to ease the purification; an enzyme or portion thereof which is catalytically active; a polypeptide which promotes

oligomerization, such as a leucine zipper domain; and a polypeptide which increases stability, such as an immunoglobulin constant region.

[0050] Fusions can be made either at the amino-terminus or at the carboxyl-terminus of a β_2 -GPI polypeptide. Fusions may be direct with no linker or adapter molecule or may be through a linker or adapter molecule. A linker or adapter molecule may be one or more amino acid residues, typically from about 20 to about 50 amino acid residues. A linker or adapter molecule may also be designed with a cleavage site for a protease to allow for the separation of the fused moieties. It will be appreciated that once constructed, the fusion polypeptides can further be derivatized according to the methods described herein.

[0051] The β_2 -GPI protein of the present invention may also be β_2 -GPI derivatives, which is a chemically or biologically modified protein, including protein post-translation modification, such as acylation (i.e., acetylation or formylation), biotinylation, carboxylation, deamination, glutathionylation, glycosylation, lipidation (i.e., farnesylation, geranylgeranylation, prenylation, myristoylation, palmitoylation, or stearoylation), methylation, phosphorylation, sulphation, fucosylation, and ubiquitination. Unless otherwise indicated, the term “ β_2 -GPI protein” refers both to native proteins, and variants and derivatives thereof. A protein derivative may be modified in a manner that is different in the type, number, or location of the post-translation modification groups naturally attached to the polypeptide. For example, a derivative may have the number and/or type of glycosylation altered compared to the native protein. The resulting derivative may comprise a greater or a lesser number of N-linked glycosylation sites than the native protein.

[0052] The β_2 -GPI polypeptide may also be modified by the covalent attachment of one or more polymers. Typically, the polymer selected is water-soluble so that the protein to which it is attached does not precipitate in an aqueous environment, such as a physiological environment. The polymer may be of any molecular weight and may be branched or unbranched. The polymer each typically has an average molecular weight of between about 1 kDa to about 100 kDa.

[0053] Suitable water-soluble polymers or mixtures thereof include, but are not limited to, polyalkylene glycol (such as mono-(C₁-C₁₀) alkoxy-, aryloxy-polyethylene glycol, poly-(N-vinyl pyrrolidone) polyethylene glycol, propylene glycol homopolymers, or polypropylene oxide/ethylene oxide co-polymers), carbohydrate-based polymers (such as dextran or cellulose),

polyoxyethylated polyols, and polyvinyl alcohols. Also encompassed by the present invention are bifunctional crosslinking molecules which can be used to prepare covalently attached β_2 -GPI polypeptide multimers.

[0054] In general, chemical derivatization may be performed under a suitable condition by reacting a protein with an activated polymer molecule. Methods for preparing chemical derivatives of polypeptides will generally comprise the steps of: (a) reacting the polypeptide with the activated polymer molecule (such as a reactive ester or aldehyde derivative of the polymer molecule) under conditions whereby the β_2 -GPI protein becomes attached to one or more polymer molecules, and (b) obtaining the reaction products. The optimal reaction conditions may vary depending upon the β_2 -GPI protein selected and chemical reagents used, and are generally determined experimentally. The PEGylation of a polypeptide may be carried out using any of the PEGylation reactions known in the art, including, but not limited to, acylation, alkylation, or Michael addition.

Diagnostic Assay

[0055] There are many different types of immunoassays suitable for use in the present invention. Any of the well known immunoassays may be adapted to detect the level of β_2 -GPI - specific antibodies in a sample which react with the β_2 -GPI antigens, such as, e.g., enzyme linked immunoabsorbent assay (ELISA), fluorescent immunosorbent assay (FIA), chemical linked immunosorbent assay (CLIA), radioimmuno assay (RIA), immunoblotting, gel diffusion precipitation reactions, immunodiffusion assays, in situ immunoassays (e.g., using colloidal gold, enzyme or radioisotope labels, for example), Western blots, precipitation reactions, agglutination assays (e.g., gel agglutination assays, hemagglutination assays, etc.), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc. For a review of the different immunoassays which may be used, see: The Immunoassay Handbook, David Wild, ed., Stockton Press, New York, 1994. A competitive immunoassay with solid phase separation or an immunometric assay for antibody testing is particularly suitable for use in the present invention. See, The Immunoassay Handbook, chapter 2.

[0056] In one exemplary embodiment of the invention, the diagnostic assay is an immunometric assay for detecting the level of β_2 -GPI -specific antibodies in a sample. In the immunometric assay, the β_2 -GPI antigens are immobilized on a solid support directly or

indirectly through a capture agent, such as anti- β_2 -GPI antibodies. An aliquot of a sample, such as a serum sample, from a subject is added to the solid support and allowed to incubate with the β_2 -GPI antigens on the solid phase. A secondary antibody that recognizes a constant region in the autoantibodies present in the sample which have reacted with the β_2 -GPI antigen is added. When the subject is a human, this secondary antibody is an anti-human immunoglobulin. The secondary antibody which is specific for IgA, IgG, or IgM heavy chain constant regions, or combination thereof, may be employed. After separating the solid support from the liquid phase, the support phase is examined for a detectable signal. The presence of the signal on the solid support indicates that autoantibodies to the native β_2 -GPI proteins present in the sample have bound to the β_2 -GPI antigen on the solid support. Increased optical density or radiolabeled signal when compared to the control samples from normal subjects correlates with a diagnosis of APS in a subject.

[0057] Solid supports are known to those skilled in the art and include the walls of wells of a reaction tray (e.g., microtiter plates), test tubes, polystyrene beads, magnetic beads, nitrocellulose strips, membranes, microparticles such as latex particles, glass or silicon chips, sheep (or other animal) red blood cells, duracytes and others. Suitable methods for immobilizing nucleic acids on solid phases include ionic, hydrophobic, covalent interactions and the like. A solid support, as used herein, refers to any material which is insoluble, or can be made insoluble by a subsequent reaction. The solid support can be chosen for its intrinsic ability to attract and immobilize the capture reagent. Alternatively, the solid phase can retain an additional molecule which has the ability to attract and immobilize the capture reagent. The additional molecule can include a charged substance that is oppositely charged with respect to the capture reagent itself or to a charged substance conjugated to the capture reagent. As yet another alternative, the molecule can be any specific binding member which is immobilized upon (attached to) the solid support and which has the ability to immobilize the β_2 -GPI antigen through a specific binding reaction. The molecule enables the indirect binding of the β_2 -GPI antigen to a solid support material before the performance of the assay or during the performance of the assay.

[0058] The signal producing system is made up of one or more components, at least one of which is a label, which generate a detectable signal that relates to the amount of bound and/or unbound label, i.e., the amount of label bound or unbound to the β_2 -GPI antigen. The label is a molecule that produces or which may be induced to produce a signal. Examples of labels

include fluorescers, enzymes, chemiluminescers, photosensitizers or suspendable particles. The signal is detected and may be measured by detecting enzyme activity, luminescence or light absorbance. Radiolabels may also be used and levels of radioactivity detected and measured using a scintillation counter.

[0059] Examples of enzymes which may be used to label the anti-human immunoglobulin include β -D-galactosidase, horseradish peroxidase, alkaline phosphatase, and glucose-6-phosphate dehydrogenase ("G6PDH"). Examples of fluorescers which may be used to label the anti-human immunoglobulin include fluorescein, isothiocyanate, rhodamines, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde, fluorescamine, and Alexa Fluor® dyes (that is, sulfonated courmarin, rhodamine, xanthene, and cyanine dyes). Chemiluminescers include e.g., isoluminol. For example, the anti-human immunoglobulin may be enzyme labeled with either horseradish peroxidase or alkaline phosphatase.

[0060] Enzymes may be covalently linked to β_2 -GPI antigen reactive antibodies for use in the methods of the present invention using well known methods. There are many well known conjugation methods. For example, alkaline phosphatase and horseradish peroxidase may be conjugated to antibodies using glutaraldehyde. Horseradish peroxidase may also be conjugated using the periodate method. Commercial kits for enzyme conjugating antibodies are widely available. Enzyme conjugated anti-human and anti-mouse immunoglobulin specific antibodies are available from multiple commercial sources.

[0061] Biotin labeled antibodies may be used as an alternative to enzyme linked antibodies. In such cases, bound antibody would be detected using commercially available streptavidin-horseradish peroxidase detection systems.

[0062] Enzyme labeled antibodies produce different signal sources, depending on the substrate. Signal generation involves the addition of substrate to the reaction mixture. Common peroxidase substrates include ABTS (2,2'-azinobis(ethylbenzothiazoline-6-sulfonate)), OPD (O-phenylenediamine) and TMB (3,3', 5,5'-tetramethylbenzidine). These substrates require the presence of hydrogen peroxide. p-Nitrophenyl phosphate is a commonly used alkaline phosphatase substrate. During an incubation period, the enzyme gradually converts a proportion of the substrate to its end product. At the end of the incubation period, a stopping reagent is added which stops enzyme activity. Signal strength is determined by measuring optical density, usually via spectrophotometer.

[0063] Alkaline phosphatase labeled antibodies may also be measured by fluorometry. Thus in the immunoassays of the present invention, the substrate 4-methylumbelliferyl phosphate (4-UMP) may be used. Alkaline phosphatase dephosphorylated 4-UMP to form 4-methylumbelliferone (4-MU), the fluorophore. Incident light is at 365 nm and emitted light is at 448 nm.

[0064] The amount of color, fluorescence, luminescence, or radioactivity present in the reaction (depending on the signal producing system used) is proportionate to the amount of autoantibodies in a sample which react with the β_2 -GPI antigens. Quantification of optical density may be performed using spectrophotometric or fluorometric methods, including flow cytometers. Quantification of radiolabel signal may be performed using scintillation counting.

[0065] In another exemplary embodiment, the assay is a competitive immunoassay, which employs one or more β_2 -GPI-specific antibodies that binds to the same epitopes as the β_2 -GPI-specific autoantibodies. In the assay, the β_2 -GPI-specific antibodies and the β_2 -GPI-specific autoantibodies in a sample compete for binding to the β_2 -GPI antigens. Typically, a constant amount of a labeled antibody which is known to bind to β_2 -GPI antigen is incubated with different concentrations of a sample from a subject. The β_2 -GPI-specific antibodies may be monoclonal or polyclonal.

[0066] As described herein above, the β_2 -GPI-specific antibodies may be labeled with a fluorescer, enzyme, chemiluminescer, photosensitizer, suspendable particles, or radioisotope. After incubation, bound labeled antibodies are separated from free antibodies. Depending on the signal producing system used and if necessary, an appropriate substrate with which the labeled antibody reacts is added and allowed to incubate. The signal generated by the sample is then measured. A decrease in optical density or radioactivity from before and after addition of the serum sample or between experimental and control samples, is indicative that autoantibodies in the sample have bound to the β_2 -GPI antigen. Decreased optical density or radiolabeled signal when compared to control samples from normal subjects correlates with a diagnosis of APS in a subject.

[0067] In an alternative exemplary embodiment of the competitive immunoassay, an indirect method using two antibodies is provided. β_2 -GPI antigen specific antibodies are added first as described in the preceding paragraph with the exception that they are not labeled. They are

incubated with different concentrations of a sample from a subject. A constant amount of a second antibody is then added to the mixture of the sample and the first antibody. The second antibody recognizes constant regions of the heavy chains of the first antibody. For example, the second antibody may be an antibody which recognizes constant regions of the heavy chains of mouse immunoglobulin which has reacted with the β_2 -GPI antigens (anti-mouse immunoglobulin). The second antibody may be labeled with a fluorophore, chemilophore or radioisotope, as described above. Free labeled second antibody is separated from bound antibody. If an enzyme-labeled antibody is used, an appropriate substrate with which the enzyme label reacts is added and allowed to incubate. A decrease in optical density or radioactivity from before and after addition of the serum sample in comparison with control samples is indicative that autoantibodies in the serum sample have bound to the β_2 -GPI antigens. Decreased optical density or radioactivity when compared to control samples from normal subject correlates with a diagnosis of a APS in a subject.

[0068] In some embodiments, an automated detection assay is utilized. Methods for the automation of immunoassays include those described in U.S. Pat. Nos. 5,885,530, 4,981,785, 6,159,750, and 5,358,691, each of which is herein incorporated by reference. In some embodiments, the analysis and presentation of results is also automated. For example, in some embodiments, software that generates a prognosis based on the presence or absence of a series of proteins corresponding to autoimmune or chronic inflammatory disease markers is utilized.

[0069] In some embodiments, the β_2 -GPI specific autoantibody level may be used together with other biological markers as a panel for the diagnosis of heart disease. The panel allows for the simultaneous analysis of multiple markers correlating with ASS. For example, a panel may include markers identified as correlating with ASS in a subject that is likely or not to respond to a given treatment. Depending on the subject, panels may be analyzed alone or in combination in order to provide the best possible diagnosis and prognosis. Markers for inclusion on a panel are selected by screening for their predictive value using any suitable method, including but not limited to, those described in the illustrative examples below.

Data Analysis

[0070] In the present invention, a computer-based analysis program may also be used to translate the raw data generated by the detection assay into data of predictive value for a clinician. The clinician can readily access the predictive data using any suitable means. The

clinician is then able to immediately utilize the information in order to optimize the care of the subject.

[0071] The present invention contemplates any method capable of receiving, processing, and transmitting the information to and from laboratories conducting the assays, information provides, medical personal, and subjects. For example, in some embodiments of the present invention, a sample (e.g., a biopsy or a serum or urine sample) is obtained from a subject and submitted to a profiling service (e.g., clinical lab at a medical facility, genomic profiling business, etc.), located in any part of the world (e.g., in a country different than the country where the subject resides or where the information is ultimately used) to generate raw data. Where the sample comprises a tissue or other biological sample, the subject may visit a medical center to have the sample obtained and sent to the profiling center, or subjects may collect the sample themselves (e.g., a urine sample) and directly send it to a profiling center. Where the sample comprises previously determined biological information, the information may be directly sent to the profiling service by the subject (e.g., an information card containing the information may be scanned by a computer and the data transmitted to a computer of the profiling center using an electronic communication system). Once received by the profiling service, the sample is processed and a profile is produced, specific for the diagnostic or prognostic information desired for the subject.

[0072] The profile data is then prepared in a format suitable for interpretation by a treating clinician. For example, rather than providing raw expression data, the prepared format may represent a diagnosis or risk assessment (e.g., likelihood of a liver disease such as HCC to respond to a specific therapy) for the subject, along with recommendations for particular treatment options. The data may be displayed to the clinician by any suitable method. For example, in some embodiments, the profiling service generates a report that can be printed for the clinician (e.g., at the point of care) or displayed to the clinician on a computer monitor.

[0073] In some embodiments, the information is first analyzed at the point of care or at a regional facility. The raw data is then sent to a central processing facility for further analysis and/or to convert the raw data to information useful for a clinician or patient. The central processing facility provides the advantage of privacy (all data is stored in a central facility with uniform security protocols), speed, and uniformity of data analysis. The central processing facility can then control the fate of the data following treatment of the subject. For example,

using an electronic communication system, the central facility can provide data to the clinician, the subject, or researchers.

[0074] In some embodiments, the subject is able to directly access the data using the electronic communication system. The subject may chose further intervention or counseling based on the results. In some embodiments, the data is used for research use. For example, the data may be used to further optimize the inclusion or elimination of markers as useful indicators of a particular condition or severity of disease.

EXAMPLES

[0075] Autoantibodies targeting β_2 -Glycoprotein I (β_2 -GPI), a component of the atherosclerotic plaque, are commonly found in patients with acute ischemic syndromes. Serum samples from APS (Antiphospholipid Syndrome) patients and from cardiovascular patients exhibiting acute atherosclerotic syndromes were analyzed for IgG and IgA antibodies in both anti- β_2 -GPI and anti-cardiolipin (aCL) ELISA assays. All of the APS samples used were positive in both assays. Serum samples from 382 cardiovascular patients were also analyzed for IgG and IgA antibodies in the same assays. In sharp contrast to the APS samples, it was found that only 1% of the samples from cardiovascular patients were positive for IgA aCL, and 1.6% positive for IgG aCL, whereas 35.6% were positive for IgA anti- β_2 -GPI and only 1.6 % for IgG anti- β_2 -GPI. The antigenic specificity of 29 serum samples from cardiovascular patients was evaluated. Six different recombinant domain-deleted mutants (DM) of human β_2 -GPI and full-length human β_2 -GPI (wild-type), were used in competitive inhibition assays to inhibit the autoantibodies from binding in the anti- β_2 -GPI ELISA assays. Domain-deleted mutants D--345 and D---45 inhibited the binding in the IgA anti- β_2 -GPI assay suggesting that these autoantibodies recognize domain 4 of the β_2 -GPI molecule. These results demonstrated that IgA anti- β_2 -GPI autoantibodies from atherosclerotic patients are distinct from IgA autoantibodies found in APS samples.

Example 1

[0076] This example demonstrates that the predominant antibody profile exhibited by APS patients (cardiolipin IgG positive/ β_2 -GPI IgG/IgA positive) differed from the predominant profile exhibited by atherosclerotic patients with acute ischemic disease (cardiolipin IgG/IgA negative/ β_2 -GPI IgA positive) because of differing domain specificity of the APS and

atherosclerosis patient's antibodies. Using a series of full-length β_2 -GPI and β_2 -GPI Dms, a large number of serum samples from patients with APS and various atherosclerotic populations were tested for IgG, IgA, and IgM antibodies to these constructs by using a competitive inhibition ELISA. All specimens were also tested for IgG, IgA, and IgM aCL antibodies. This experiment demonstrated that 29 of 29 IgA anti- β_2 -GPI positive samples from atherosclerosis patients specifically recognized domain 4 of β_2 -GPI.

Materials and Methods

Recombinant β_2 -GPI

[0077] The recombinant β_2 -GPI and β_2 -GPI domain-deleted mutants (Dms) used are as previously described (Igarashi M, et al., *Blood* 87(8):3262-3270 (1996). Briefly, TN5 insect cells were infected with high titer viral stock produced in Sf9 insect cells. Each construct contained a 6his tail that was used for purifying the protein from culture media. The nomenclature for domain deletion mutants uses numbers to indicate the presence of domains while a dash symbolizes the domain is missing. Thus D--345 is the name given to the recombinant protein that contains domains 3, 4 and 5 while lacking domains 1 and 2.

Patient sample selection

[0078] The diagnosis of each syndrome described was done according to clinical presentation, ultrasound, angiography or magnetic angioresonance studies. Patients were enrolled consecutively in a tertiary center (University Hospital). Patients with infective endocarditis, osteonecrosis, neoplasms, cerebral hemorrhage, infection by HIV or treponema pallidum, presence of known heritable causes of thrombosis such as homocystinuria or factor V (Leiden) mutation, previous diagnosis of APS, or other connective tissue disorder (CTD) were excluded.. Control patients were recruited from patients admitted to the Orthopedic clinic for fractures or musculoligamentous disorders and without acute myocardial infarction, stroke, or other cardiac conditions.(Ranzolin A, et al., *Arg Bras Cardiol*, 83(2): 141-4; 137-140 (2004); Staub HL et al., *Arg Bras Neuropsiquiat* 61(3B): 757-63 (2003)).

[0079] A total of 511 archived specimens consisting of 382 sera from individuals with various atherosclerosis conditions and 129 sera from individuals with anti-phospholipid syndrome were studied. The atherosclerosis group included sera from individuals with peripheral arterial disease (117), acute coronary syndrome (117), and acute myocardial

infarction (90). Ten samples were randomly selected from APS patients that were positive for both IgG and IgA in the anti- β_2 -GPI ELISA, and 29 samples from atherosclerosis patients that were positive for IgA in the anti- β_2 -GPI ELISA.

Anti- β_2 -GPI and Anti-Cardiolipin ELISA

[0080] All samples, both from the APS and cardiovascular patients, were tested for the presence of anticardiolipin (aCL) antibodies and anti- β_2 -GPI antibodies by ELISA. Specimens were first tested for the presence of IgG, IgA, or IgM aCL and β_2 -GPI antibodies using polyvalent aCL and anti- β_2 -GPI screening ELISA tests. All ELISA kits used in this study were manufactured by INOVA Diagnostics (INOVA Diagnostics, San Diego, CA) and run according to the manufacturer's instructions.

Competitive inhibition ELISA

[0081] Tests were performed using the appropriate (IgG and/or IgA) anti- β_2 -GPI ELISA kit from INOVA Diagnostics. Each serum was titered to determine the dilution required to give approximately 80% of maximum binding. Test inhibitors were diluted in the sample dilution buffer provided in the kits and 25 μ l of each dilution or sample diluent alone was added to the wells. The serum samples were diluted in sample dilution buffer and 25 μ l of a constant dilution was added to the wells. The contents of the wells were mixed and plates were incubated at room temperature for 30 minutes. Wells were washed 3 times with the wash buffer provided in the kits, 50 μ l of the HRP conjugated anti-IgG or IgA added, incubated for 30 minutes, washed 3 times with wash buffer and 50 μ l of substrate solution was added. Wells were incubated at room temperature for 30 minutes and 50 μ l of stop solution was added. The OD 450 for each well was determined in an Anthos Labtec HT2 microplate reader (Salzburg, Austria). The percent inhibition was determined as follows: [(mean A₄₅₀ obtained from the control wells without inhibitor less A₄₅₀ of background) - (A₄₅₀ obtained in the presence of inhibitor less A₄₅₀ of background)/mean A₄₅₀ obtained from the control wells without inhibitor less A₄₅₀ of background)] X 100.

Results

Anti- β_2 -GPI and Anti-Cardiolipin

[0082] Serum samples from APS and cardiovascular patients were analyzed for IgG and IgA autoantibodies in both the anti- β_2 -GPI and the anti-cardiolipin (aCL) assays. Almost 80% of the APS samples were positive by polyvalent IgG/IgA/IgM aCL and β_2 -GPI screening assays (Table 1).

Table 1: Frequency of aCL and anti- β_2 -GPI antibodies in APS and Cardiovascular Groups by ELISA testing

Total sera =511	APS patients (n= 129)	Cardiovascular patients (n=382)
aCL Screen (IgG/IgA/IgM)	78%	12%
anti- β_2 -GPI Screen (IgG/IgA/IgM)	79%	46%
aCL IgG	64%	1%
aCL IgA	9%	1%
Anti- β_2 -GPI IgG	43%	1%
Anti- β_2 -GPI IgA	48%	33%

[0083] Specific isotype testing of the APS sera revealed that approximately 64% were IgG and 9% were IgA ACA antibody positive, while 43% were IgG and 48% were IgA anti- β_2 -GPI positive. Serum samples from 370 cardiovascular patients were similarly tested for total (IgG/IgA/IgM) and specific IgG and IgA antibodies in both the aCL and anti- β_2 -GPI assays. In sharp contrast to the APS samples, where IgG aCL and IgG anti- β_2 -GPI antibodies were found in 64% and 43% of the specimens, respectively, it was found that IgG aCL and IgG anti- β_2 -GPI antibodies were present in only 1% of the samples from cardiovascular patients. Even more striking was the observation that while the pattern of reactivity for IgA aCL and IgA anti- β_2 -GPI was similar in the APS and cardiovascular patients (both had low levels of IgA aCL and moderate levels of IgA anti- β_2 -GPI), IgA anti- β_2 -GPI was the only major antibody present in the cardiovascular group. In contrast, the APL patients had moderate levels of antibodies to IgG aCL, IgG anti- β_2 -GPI, and IgA anti- β_2 -GPI (Table 1).

Epitope(s) of β_2 -GPI recognized by both IgG and IgA anti- β_2 -GPI I from APS patients.

[0084] Recombinant β_2 -GPI and two deletion mutants were used to determine the antigenic specificity of both the IgG and IgA autoantibodies from 10 different APS patients. Each recombinant form of β_2 -GPI was tested, in a dose-dependent fashion, for its ability to inhibit these autoantibodies from binding to full length β_2 -GPI (Table 2, Figure 1).

Table 2: Competitive inhibition assay using 10 different APS serum samples with indicated recombinant B2GPI and deletion mutants.

IgG						
Antibody Number	D12345 ¹		D12---		D---45	
	Max ²	50% ³	Max	50%	Max	50%
6612	89	9.5	98	24.0	17.0	>125
6626	88	31.8	90	88.0	3.5	>125
6635	95	11.8	96	29.5	3.0	>125
6647	92	25.8	94	66.0	5.0	>125
6656	64	38.4	67	85.0	10.0	>125
6666	79	27.2	85	20.3	2.4	>125
6674	90	20.5	94	52.0	12.0	>125
7002	58	53.7	53	178.0	7.0	>125
7005	77	23.5	71	58.0	1.9	>125
7010	83	15.8	86	41.6	14.0	>125

IgA						
Antibody Number	D12345		D12---		D---45	
	Max	50%	Max	50%	Max	50%
6612	80	13.1	74	36	28	>125
6626	62	42.4	39	>125	61	>125
6635	57	44.7	67	29.6	0	>125
6647	68	30.7	58	66.4	0.7	>125
6656	74	24.3	42	>125	40	>125
6666	48	67.3	42	>125	11	>125
6674	85	39.9	88	39.9	4	>125
7002	29	116.3	21	>125	0	>125
7005	35	50.6	45	>125	40	>125
7010	63	16.7	59	50	12	>125

> = Highest concentration tested.

¹ = Domains included in construct.

² = Maximum Inhibition observed at concentrations tested.

³ = Concentration (micromolar) to give 50% inhibition.

[0085] Only those constructs that contained domain 1 inhibited both the IgG and IgA autoantibodies. As shown in Table 2, both the IgG and IgA anti- β_2 -GPI binding antibodies from all 10 patients were inhibited by both constructs that contain domain 1. None of the samples were effectively inhibited, even at the highest concentration tested, by the construct that lacked domain 1. ID₅₀ values for mutants that contain domain 1 ranged from 1 to 50 μ M for the IgG antibody and 13 to 100 μ M for the IgA antibody. By contrast the mutant that did not contain domain 1 (D---45) did not effectively inhibit either the IgG nor the IgA antibody.

Epitope(s) of β_2 -GPI recognized by IgA anti- β_2 -GPI from patients with acute cardiovascular syndromes.

[0086] The differing β_2 -GPI and aCL profile of the APS and cardiovascular sera (Table 1) suggested to us that the IgA anti- β_2 -GPI antibodies in cardiovascular patients may be distinct from those present in APS patients and might target a different domain on the β_2 -GPI protein.

[0087] Twenty nine samples from the cardiovascular patient cohort were selected which were IgA anti- β_2 -GPI antibody positive and aCL IgG, aCL IgM, and with the exception of one sera, aCL IgA negative. The detailed β_2 -GPI and aCL profiles of these sera are shown in Table 3.

Table 3: Anti-aCL and anti- β_2 -GPI profile of samples from cardiovascular patients¹ for inhibition study.

Sample	β_2 -GPI IgA Units	β_2 -GPI IgG Units ²	β_2 -GPI IgM Units	aCL IgA Units	aCL IgG Units	aCL IgM Units
ACS-52	58.9	0	10.1	n.t.	n.t.	n.t.
ACS-53	20.7	0	4.0	n.t.	n.t.	n.t.
ACS-54	33.1	9.6	0	n.t.	n.t.	n.t.
ACS-58	86.4	0	5.3	n.t.	n.t.	n.t.
ACS-65	234.4	0	4.0	n.t.	n.t.	n.t.
ACS-67	38.7	0	0	n.t.	n.t.	n.t.
ACS-71	77.6	0	0	n.t.	n.t.	n.t.
ACS-74	154.8	0	5.7	n.t.	n.t.	n.t.
ACS-104	23.7	0	58.5	n.t.	n.t.	n.t.
ACS-136	32.0	64.4	0.2	n.t.	n.t.	n.t.
ACS-144	28.3	3.3	16.9	n.t.	n.t.	n.t.
<hr/>						
CAS-5	65.4	0	2.8	n.t.	n.t.	n.t.
CAS-6	55.5	1.2	0	n.t.	n.t.	n.t.
CAS-8	61.2	0	0	n.t.	n.t.	n.t.
CAS-13	39.9	1.7	11.7	n.t.	n.t.	n.t.
CAS-15	130.1	0.1	0	n.t.	n.t.	n.t.
CAS-18	35.5	0	0	n.t.	n.t.	n.t.
CAS-28	60.5	0	0	n.t.	n.t.	n.t.
CAS-29	51.8	4.1	5.7	n.t.	n.t.	n.t.
<hr/>						
MI-5	70.0	0	25.6	n.t.	n.t.	n.t.
MI-7	90.4	0	0	n.t.	n.t.	n.t.
MI-10	183.9	0	25.4	n.t.	n.t.	n.t.
MI-15	58.4	0	77.4	n.t.	n.t.	n.t.
MI-37	32.7	0	41.2	n.t.	n.t.	n.t.
MI-45	25.5	0	11.6	n.t.	n.t.	n.t.
<hr/>						
PAD-30	27.6	0	0	n.t.	n.t.	n.t.
PAD-39	22.1	0	0	n.t.	n.t.	n.t.
PAD-42	45.5	0	0	27.9	n.t.	n.t.
PAD-101	26.4	3.0	25.4	n.t.	n.t.	n.t.

n.t. = Specimens testing negative on aCL screening assay were not tested on isotype-specific assays.

¹ = ACS: Acute Coronary Syndrome. MI: Myocardial Infarction. CAS: Carotid Artery Study. PAD: Peripheral Artery Disease.

² = β_2 -GPI results with negative values (resulting from extrapolation at bottom of the standard curve) were assigned a value of 0.

[0088] Seven different recombinant β_2 -GPI mutant proteins were used to determine the antigenic specificity of the IgA β_2 -GPI binding antibodies from 29 different samples from patients with various cardiovascular conditions, including acute cardiac syndrome (11), acute myocardial infarction (6), carotid artery disease (8), and peripheral artery disease (4). Each mutant recombinant β_2 -GPI protein was tested, in a dose-dependent fashion, for its ability to inhibit the IgA antibody from binding to full-length β_2 -GPI (Table 4).

Table 4: Competitive inhibition assays using 29 different serum samples, from patients with cardiovascular problems¹, with indicated recombinant B2GP1 and deletion-mutants.

Ab#	12345 ²		12---		123--		1234-		---345		---45		----5	
	Max ³	50% ⁴	Max	50%	Max	50%	Max	50%	Max	50%	Max	50%	Max	50%
ACS-104	62	20	0	>125	13	>83	0	>63	79	30	76	24	10	>250
ACS-136	74	12	3	>125	0	>83	0	>63	72	6	76	26	3	>250
ACS-144	52	12	12	>125	12	>83	8	>63	63	6	68	7	9	>250
ACS-52	70	17	0	>125	14	>83	30	>63	86	6	87	2	21	>250
ACS-53	56	23	15	>125	12	>83	24	>83	76	10	80	14	10	>250
ACS-54	71	5	11	>125	0	>83	0	>63	67	3	85	2	3	>250
ACS-58	57	24	0	>125	7	>83	24	>63	84	20	90	16	2	>250
ACS-65	82	12	0	>125	18	>83	88	5	92	4	94	8	0	>250
ACS-67	65	21	8	>125	2	>83	49	>63	75	21	83	23	20	>250
ACS-71	72	12	18	>125	7	>83	13	>63	90	4	93	4	49	227
ACS-74	69	10	3	>125	2	>83	15	>63	90	5	91	1	29	>250
<hr/>														
CAS-13	54	34	7	>125	0	>83	10	>63	74	17	72	33	15	>250
CAS-15	83	2	0	>125	1	>83	0	>63	92	2	91	17	11	>250
CAS-18	72	24	0	>125	13	>83	0	>63	85	31	82	46	0	>250
CAS-28	65	18	7	>125	22	>83	39	>63	84	26	84	29	0	>250
CAS-29	63	26	8	>125	17	>83	33	>63	76	36	75	28	0	>250
CAS-5	70	29	0	>125	19	>83	29	>63	90	6	84	20	0	>250
CAS-6	68	24	14	>125	14	>83	27	>63	78	6	77	17	15	>250
CAS-8	75	23	14	>125	0	>83	6	>63	88	6	87	17	0	>250
<hr/>														
MI-10	74	14	8	>125	12	>83	64	43	90	13	89	4	0	>250
MI-15	71	16	6	>125	19	>83	25	>63	71	34	80	12	19	>250
MI-37	37	52	0	>125	0	>83	0	>63	50	7	58	16	2	>250
MI-45	60	27	5	>125	16	>83	20	>63	67	38	66	8	0	>250
MI-5	73	18	30	>125	0	>83	0	>63	90	2	58	5	0	>250
MI-7	85	2	0	>125	35	>83	48	55	76	21	90	8	56	80
<hr/>														
PAD-101	27	>50	3	>125	0	>83	14	>63	67	51	72	55	6	>250
PAD-30	57	24	0	>125	19	>83	58	41	78	30	79	24	0	>250
PAD-39	59	19	14	>125	10	>83	54	46	75	30	77	24	0	>250
PAD-42	40	>50	9	>125	6	>83	9	>63	79	24	83	29	14	>250

> = Greater than highest concentration tested

¹ = ACS: Acute Coronary Syndrome. MI: Myocardial Infarction. CAS: Carotid Artery Study. PAD: Peripheral Artery Disease.

² = Domains included in construct.

³ = Maximum Inhibition observed at concentrations tested.

⁴ = Concentration (μM) to give 50% inhibition.

[0089] An example of the results is shown graphically in Figure 2. With the exception of the full-length construct, only the D--345 and D---45 constructs inhibited these IgA antibodies.

Four of the 29 samples were also inhibited, albeit to a much lesser extent, by the D1234- construct. Only one of the samples was also inhibited by the D----5 construct. ID₅₀ values for the D--345 and D---45 mutants ranged from 1 to 55 μ M. By contrast, the D12--- and D123-- mutants did not effectively inhibit the binding of any of the 29 samples tested.

Discussion

[0090] It has been previously shown that the antigenic specificity of the IgG autoantibodies found in APS patients recognize domain 1 of the β_2 -GPI molecule (Iverson, GM et al., *PNAS* 95: 15542-15546 (1998); Iverson, GM, et al., *J. Immunol.* 169:7097-7103 (2002)). The antigenic specificity of the IgA autoantibodies from APS patients however, was not known. The inhibition studies reported here (Fig. 1, Table 2) clearly show that the antigenic specificity of the battery of 10 APS samples studied in this report are directed toward an epitope that is contained within domain 1 of the β_2 -GPI molecule. Thus, the antigenic specificity of both the IgG and IgA autoantibodies found in APS patients is domain 1.

[0091] This experiment also shows (Fig. 2 and Table 3) that IgA β_2 -GPI binding antibodies from patients with several types of ASS (acute myocardial infarction, acute coronary syndrome, “carotid artery study”, and peripheral artery disease) recognize an epitope on domain 4 of the β_2 -GPI molecule. This should not be confused with earlier studies that purported to show autoantibodies from APS patients recognize domain 4 of β_2 -GPI (Igarashi M., et al, *Blood*, 87 (8): 3262-70 (1996); George J, et al., *J Immunol.* 160(8):3917-3923 (1998)). These studies were designed to study the antigenic specificity of IgG, not IgA, autoantibodies.

[0092] The Dms that contained domain 4 inhibited in a similar, but not identical pattern, among the various samples tested. For example, only 4 were inhibited by the D1234- construct. This suggests that these antibodies recognize comparable, but distinguishable, epitopes present on domain 4. A recent molecular simulation derived from β_2 -GPI crystal structure supports this possibility. This study suggested 2 discontinuous antigenic sequences in Domain 4 β_2 -GPI (Kasahara H, et al., *Int. Immunol.* 17:1533-1542 (2005)). Domain 4 may have different conformational states when present in constructs containing different domains. For example, a few samples recognized domain 4 when domain 5 was absent, while the majority only recognized domain 4 when domain 5 was present. Thus, these antibodies may recognize an epitope on domain 4 that is affected by the presence of additional domains. This interpretation was also supported by the simulation experiments.

[0093] Previously it has been shown that the orientation of β_2 -GPI on the ELISA plate is important for the binding of anti- β_2 -GPI autoantibodies when measured by ELISA. (George J, et al., *J. Immunol.* 160(8):3917-3923 (1998)) This could explain why these samples recognize β_2 -GPI when adsorbed onto plastic plates, but do not bind β_2 -GPI when adsorbed onto cardiolipin. The binding of β_2 -GPI to cardiolipin via domain 5 may give a different orientation than when bound to plastic. Binding via domain 5 may alter either the availability of domain 4 or the configuration of domain 4, or both. However, the orientation of the β_2 -GPI molecule when adsorbed onto plastic is not completely understood. It is conceivable that sufficient numbers of molecules adsorb to the plastic in such an orientation that domain 4 is neither hindered, that is available for antibody, nor has its conformation altered enough to negate the binding of these antibodies.

Example 2

[0094] A clinical study was performed using an ELISA assay to detect domain 4-specific antibodies using a domain 4/5 combined antigen using 218 clinically characterized sera. The cohort included healthy control individuals (n=30), patients with cardiac stents (without symptoms, n=28; symptomatic, n=23), and patients with acute stroke (ischemic stroke and transient Ischemic attack; without intracerebral hemorrhage, n= 137). The mean IgA antibody value for the healthy control population was 14.1 units, compared to 26.5 units for the patients with stents and 29.5 for patients having an acute stroke (Table 5).

Table 5: ELISA assays using 218 different serum samples

Number of sera tested	30	51	137
Patient diagnosis	Control	Stent	Acute stroke
Mean value of IgA Domain 4 and 5 of β_2 -GPI antibodies	14.09	26.46	29.54

* * * * *

[0095] The examples set forth above are provided to give those of ordinary skill in the art with a complete disclosure and description of how to make and use the preferred embodiments

of the compositions, and are not intended to limit the scope of what the inventors regard as their invention. Modifications of the above-described modes (for carrying out the invention that are obvious to persons of skill in the art) are intended to be within the scope of the following claims. All publications, patents, and patent applications cited in this specification are incorporated herein by reference.


CLAIMS

What is claimed is:


1. A method of diagnosing an acute atherosclerotic syndrome in a subject, comprising determining the presence or absence of IgA domain 4-specific anti-beta 2-glycoprotein I (β_2 -GPI) antibodies in said subject, wherein the presence of said IgA anti- β_2 -GPI antibodies indicates that said subject has an acute atherosclerotic syndrome.
2. A method of diagnosing an acute atherosclerotic syndrome in a subject, comprising the steps of:
 - a. obtaining a sample from a subject suspected of having an acute atherosclerotic syndrome;
 - b. contacting the sample with a β_2 -GPI antigen comprising a domain 4 epitope; and
 - c. detecting the presence or absence of IgA domain 4-specific anti- β_2 -GPI antibodies that bind to the domain 4 epitope;
wherein the presence of said IgA domain 4-specific anti- β_2 -GPI antibodies in said sample indicates that said subject has an acute atherosclerotic syndrome.
3. A method of diagnosing an acute atherosclerotic syndrome in a subject, comprising the steps of:
 - a. contacting a sample from a subject suspected of having an acute atherosclerotic syndrome with an epitope derived from domain 4 of a β_2 -GPI antigen comprising the amino acid sequence of SEQ ID NO:5 without linker sequences under conditions suitable to form a complex of the epitope and IgA domain 4-specific anti- β_2 -GPI antibody; and
 - b. detecting the presence or absence of the IgA domain 4-specific anti- β_2 -GPI antibody in the complex, wherein the presence of said domain 4-specific IgA anti- β_2 -GPI antibodies in said subject indicates that said subject has an acute atherosclerotic syndrome.

4. The method of claim 2, wherein IgA anti- β_2 -GPI antibodies are detected with an enzyme-linked immunosorbent assay.
5. The method of claim 2, further comprising determining the presence or absence of anticardiolipin (aCL) antibodies in said subject, wherein the presence of said domain 4-specific IgA anti- β_2 -GPI antibodies in said subject and the absence of anticardiolipin (aCL) antibodies indicates that said subject has an acute atherosclerotic syndrome.
6. The method of claim 5, wherein IgA domain 4-specific anti- β_2 -GPI antibodies and aCL antibodies are detected with independent enzyme-linked immunosorbent assays.
7. The method of claim 3, wherein the domain 4 epitope further comprises the domain 3 amino acid sequence of SEQ. ID NO. 4, and the domain 2 amino acid sequence of SEQ. ID NO. 3, or a portion thereof adjacent to the domain 3 amino acid sequence of SEQ. ID NO. 4.
8. The method of claim 7, wherein the domain 4 epitope further comprises the domain 5 amino acid sequence of SEQ. ID NO. 6, or a portion thereof adjacent to the domain 4 amino acid sequence of SEQ. ID NO. 5.
9. The method of claim 1, further comprising the step of reacting the IgA domain 4-specific anti-beta 2-glycoprotein I (β_2 -GPI) antibodies if present with a polypeptide having the amino acid sequence of SEQ. ID NO. 5.
10. The method of claim 1, further comprising the step of reacting the IgA domain 4-specific anti-beta 2-glycoprotein I (β_2 -GPI) antibodies if present with a polypeptide having the amino acid sequence of SEQ. ID NO. 5 plus SEQ. ID NO. 6.
11. The method of claim 2, wherein the β_2 -GPI antigen does not include any or a portion of domain 1.

1/3


FIGURE 1



2/3

FIGURE 2

3/3

FIGURE 3





Inhibitor, mg/ml

—◆— D 1 2 3 4 5
—■— D 1 2 ...
—▲— D ... 4 5