



US00PP32448P2

(12) **United States Plant Patent**
Ren et al.

(10) **Patent No.:** **US PP32,448 P2**

(45) **Date of Patent:** **Nov. 10, 2020**

- (54) **PETUNIA PLANT NAMED ‘BALCOBEES’**
- (50) Latin Name: *Petunia x hybrida x (Petunia x Calibrachoa)*
Varietal Denomination: **Balcobees**
- (71) Applicant: **Ball Horticultural Company**, West Chicago, IL (US)
- (72) Inventors: **Jianping Ren**, Geneva, IL (US);
Robert B. Eisenreich, North Aurora, IL (US)
- (73) Assignee: **Ball Horticultural Company**, West Chicago, IL (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **16/873,287**
- (22) Filed: **Mar. 11, 2020**

- (51) **Int. Cl.**
A01H 5/02 (2018.01)
A01H 6/82 (2018.01)
- (52) **U.S. Cl.**
USPC **Plt./356.11**
CPC *A01H 6/824* (2018.05)
- (58) **Field of Classification Search**
USPC Plt./356.11
CPC *A01H 6/824; A01H 6/821; A01H 6/82*
See application file for complete search history.

Primary Examiner — Anne Marie Grunberg
(74) *Attorney, Agent, or Firm* — Audrey Charles

(57) **ABSTRACT**
A new and distinct cultivar of *Petunia* plant named ‘Balcobees’, characterized by its dark yellow and cream-colored flowers, medium green-colored foliage, and moderately vigorous, mounded-trailing growth habit, is disclosed.

1 Drawing Sheet

1

Latin name of genus and species of plant claimed: *Petunia x hybrida x (Petunia x Calibrachoa)*.
Variety denomination: ‘Balcobees’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Petunia* plant botanically known as *Petunia x hybrida x (Petunia x Calibrachoa)* and hereinafter referred to by the cultivar name ‘Balcobees’.

The new cultivar originated in a controlled breeding program in Elburn, Ill. during December 2015. The objective of the breeding program was the development of *Petunia* cultivars with single-type flowers, unique flower coloration and patterns, and moderately vigorous, mounded-trailing growth habit.

The new *Petunia* cultivar is the result of cross-pollination and embryo rescue. The female (seed) parent of the new cultivar is the proprietary *Petunia x hybrida* breeding selection coded Y6265, not patented, characterized by its medium cream-yellow colored flowers, medium green-colored foliage, and moderately vigorous, mounding-trailing growth habit. The male (pollen) parent of the new cultivar is the proprietary *Petunia x Calibrachoa* intergeneric hybrid coded pc15-3-1, not patented, characterized by its medium yellow-colored flowers, medium green-colored foliage, and moderately vigorous, mounded growth habit. The new cultivar was selected as a single flowering plant within the progeny of the above stated cross-pollination during June 2016 in a controlled environment in Elburn, Ill.

Asexual reproduction of the new cultivar by terminal stem cuttings since June 2016 in Elburn, Ill., and Arroyo Grande, Calif. has demonstrated that the new cultivar reproduces true to type with all of the characteristics, as herein described, firmly fixed and retained through successive generations of such asexual propagation.

2

SUMMARY OF THE INVENTION

The following characteristics of the new cultivar have been repeatedly observed and can be used to distinguish ‘Balcobees’ as a new and distinct cultivar of *Petunia* plant:

1. Dark yellow and cream-colored flowers;
2. Medium green-colored foliage; and
3. Moderately vigorous, mounded-trailing growth habit.

Plants of the new cultivar differ from plants of the female parent primarily in having darker yellow-colored flowers and more branches per plant. Plants of the new cultivar differ from plants of the male parent primarily in having a larger flower size, more branches per plant, and a more mounded-trailing growth habit.

Of the many commercially available *Petunia* cultivars, the most similar in comparison to the new cultivar is SUPERCAL Light Yellow ‘SAKPXC017’, U.S. Plant Pat. No. 28,612 and U.S. Utility Pat. No. 10,117,392. However, in comparison, plants of the new cultivar differ from plants of ‘SAKPXC017’ in at least the following characteristics:

1. Plants of the new cultivar have darker yellow-colored flower than plants of ‘SAKPXC017’;
2. Plants of the new cultivar have longer pistils than plants of ‘SAKPXC017’; and
3. Plants of the new cultivar have slightly larger diameter corollas than plants of ‘SAKPXC017’.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs show, as nearly true as it is reasonably possible to make the same in color illustrations of this type, typical flower and foliage characteristics of the new cultivar. Colors in the photographs may differ slightly from the color values cited in the detailed description, which accurately describes the colors of ‘Balcobees’. The approximately 3-month-old plants were grown in 6-inch pots for

approximately 9 weeks in a greenhouse in West Chicago, Ill. Plants were given one pinch one week prior to transplant.

FIG. 1 illustrates a side view of the overall growth and flowering habit of 'Balcobees'.

FIG. 2 illustrates a close-up view of an individual flower of 'Balcobees'.

DETAILED BOTANICAL DESCRIPTION

The new cultivar has not been observed under all possible environmental conditions to date. Accordingly, it is possible that the phenotype may vary somewhat with variations in the environment, such as temperature, light intensity, and day length, without, however, any variance in genotype.

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 2015 edition, except where general color terms of ordinary significance are used. The color values were determined in January 2020 under natural light conditions in West Chicago, Ill.

The following descriptions and measurements describe approximately 3-month-old plants produced from cuttings from stock plants and grown in a glass-covered greenhouse under conditions comparable to those used in commercial practice. The plants were grown in West Chicago, Ill. in 6-inch pots for approximately 9 weeks utilizing a soilless growth medium. Plants were given one pinch one week prior to transplant. Greenhouse temperatures were maintained at approximately 67° F. to 72° F. (19° C. to 22° C.) during the day and approximately 65° F. to 68° F. (18° C. to 20° C.) during the night. Supplemental lighting was used. Measurements and numerical values represent averages of typical plants.

Botanical classification: *Petunia x hybrida x (Petunia x Calibrachoa)* 'Balcobees'.

Parentage:

Female parent.—Proprietary *Petunia x hybrida* breeding selection coded Y6265, not patented.

Male parent.—Proprietary *Petunia x Calibrachoa* intergeneric hybrid coded pc15-3-1, not patented.

Propagation:

Type cutting.—Terminal stem.

Time to initiate roots.—Approximately 6 to 9 days.

Time to produce a rooted cutting.—Approximately 21 to 28 days.

Root description.—Fibrous.

Rooting habit.—Freely branching.

Plant description:

Commercial crop time.—Approximately 5 to 7 weeks from a rooted cutting to finish in a 10 cm pot.

Growth habit and general appearance.—Moderately vigorous, mounded-trailing.

Size.—Height from soil level to top of plant plane: Approximately 18.0 cm. Width: Approximately 39.0 cm.

Branching habit.—Freely branching, pinching improves basal branching. Quantity of main branches per plant: Approximately 6.

Branch.—Strength: Moderate. Length: Approximately 23.0 cm. Diameter: Approximately 4.0 mm. Length of central internode: Approximately 3.0 cm. Texture: Densely glandular pubescent with a mixture of long and short hairs. Gland color: Colorless, transparent. Color of young stems: 144A. Color of mature stems: 146A.

Foliage description:

General description.—Quantity of leaves per main branch: Approximately 14. Fragrance: Slight. Form: Simple. Arrangement on flowering stem: Opposite.

Leaves.—Aspect: Acute angle to stem. Shape: Ovate. Margin: Entire. Apex: Broadly acute. Base: Broadly attenuate. Venation pattern: Pinnate. Length of mature leaf: Approximately 4.6 cm. Width of mature leaf: Approximately 3.2 cm. Texture of upper and lower surfaces: Moderately glandular pubescent. Gland color: Colorless, transparent. Color of upper surface of young and mature foliage: 137A with venation of 146C to indistinguishable. Color of lower surface of young and mature foliage: Closest to 138B with venation of 146D to indistinguishable.

Petiole.—Length: Approximately 2.0 mm. Width: Approximately 2.0 mm. Texture: Densely glandular pubescent with a mixture of long and short hairs. Gland color: Colorless, transparent. Color: 146D.

Flowering description:

Flowering habit.—'Balcobees' is freely flowering under outdoor growing conditions with substantially continuous blooming from spring through autumn and year-round in greenhouse environment.

Lastingness of individual flower on the plant.—Approximately 10 to 12 days.

Flower description:

General description.—Type: Simple, salverform. Quantity per plant: Approximately 12. Fragrance: Slight.

Bud.—Rate of opening: Generally takes 2 to 3 days for bud to progress from first color to fully open flower.

Quantity per plant: Approximately 7.

Bud just before opening.—Shape: Oblong. Length: Approximately 3.5 cm. Diameter at apex: Approximately 9.0 mm. Diameter at base: Approximately 2.0 mm. Texture: Densely glandular pubescent. Gland color: Colorless, transparent. Color of petal portion: 154B with venation of 144A. Color of tube: 145B with venation of 144A.

Corolla.—Diameter: Approximately 5.5 cm.

Petals.—Quantity: 5, fused to form a tube. Shape: Obovate. Appearance: Matte. Margin: Entire, slightly wavy. Apex: Cuspidate. Length from tube: Approximately 2.5 cm. Length of free portion: Approximately 8.0 mm. Width: Approximately 3.0 cm. Texture of upper surface: Glabrous. Texture of lower surface: Sparsely glandular pubescent. Gland color: Colorless, transparent. Color of upper surface when first open: Closest to 9A with 4C at outer edge, midveins of 144A. Color of lower surface when first and fully open: 150D, midveins of 145A. Color of upper surface when fully open: Closest to 9A with 4C at outer edge, midveins of 144B.

Corolla tube.—Length: Approximately 3.0 cm. Diameter at distal end: Approximately 1.0 cm. Diameter at proximal end: Approximately 3.0 mm. Texture of inner surface: Glabrous. Texture of outer surface: Densely glandular pubescent. Gland color: Colorless, transparent. Color of inner surface: Base of 145A transitioning to 12A toward tube opening, venation of 145A. Color of outer surface: Base of 150C transitioning to 150D toward petals, venation of 145A.

Sepals.—Quantity per flower: 5, fused at base. Shape: Linear. Margin: Entire. Apex: Acute. Length: Approximately 2.0 cm. Width: Approximately 5.0 mm. Texture of upper and lower surfaces: Densely glandular pubescent. Gland color: Colorless, transparent. Color of upper surface: 137A. Color of lower surface: 138A with 144A at base.

Peduncle.—Strength: Strong, flexible. Aspect: Acute angle to stem. Length: Approximately 2.5 cm. Diameter: Approximately 2.0 mm. Texture: Densely glandular pubescent with a mixture of long and short hairs. Gland color: Colorless, transparent. Color: 146B.

Reproductive organs.—Androecium: Stamen quantity: 5, basifixed. Stamen length: Approximately 2.3 cm. Filament length of fixed portion: Approximately 9.0 mm. Filament color: 155C. Anther shape: Bilobed.

Anther length: Approximately 1.0 mm. Anther color: 158A. Pollen amount: Abundant. Pollen color: 155A. Gynoecium: Pistil quantity: 1 per flower. Pistil length: Approximately 2.2 cm. Stigma shape: Funnel. Stigma length: Approximately 1.0 mm. Stigma color: 144B. Style length: Approximately 1.8 cm. Style color: 145B. Ovary length: Approximately 3.0 mm. Ovary color: 144A.

Seed and fruit production: Neither seed nor fruit production has been observed.

Disease and pest resistance: Resistance to pathogens and pests common to *Petunia* has not been observed.

What is claimed is:

1. A new and distinct cultivar of *Petunia* plant named 'Balcobees', substantially as herein illustrated and described.

* * * * *



FIG. 1




FIG. 2