



(86) Date de dépôt PCT/PCT Filing Date: 2003/01/23
(87) Date publication PCT/PCT Publication Date: 2003/07/31
(85) Entrée phase nationale/National Entry: 2004/07/21
(86) N° demande PCT/PCT Application No.: NL 2003/000049
(87) N° publication PCT/PCT Publication No.: 2003/062430
(30) Priorité/Priority: 2002/01/23 (02075266.3) EP

(51) Cl.Int.⁷/Int.Cl.⁷ C12N 15/61, C12P 7/56, C12P 7/54,
C12N 15/54, C12P 7/46, C12P 13/04, C12P 35/00,
C12P 7/20, C12N 1/19, C12N 1/14, C12P 7/10

(71) Demandeur/Applicant:
ROYAL NEDALCO B.V., NL

(72) Inventeurs/Inventors:
OP DEN CAMP, HUBERTUS JOHANNES MARIE, NL;
HARHANGI, HARRY RAMANOEDJ, NL;
VAN DER DRIFT, CHRISTIAAN, NL;
PRONK, JACOBUS THOMAS, NL

(74) Agent: MCCARTHY TETRAULT LLP

(54) Titre : FERMENTATION DE SUCRES PENTOSE
(54) Title: FERMENTATION OF PENTOSE SUGARS

(57) **Abrégé/Abstract:**

The present invention relates to host cells transformed with a nucleic acid sequence encoding a eukaryotic xylose isomerase obtainable from an anaerobic fungus. When expressed, the sequence encoding the xylose isomerase confers to the host cell the ability to convert xylose to xylulose which may be further metabolised by the host cell. Thus, the host cell is capable of growth on xylose as carbon source. The host cell preferably is a eukaryotic microorganism such as a yeast or a filamentous fungus. The invention further relates to processes for the production of fermentation products such as ethanol, in which a host cell of the invention uses xylose for growth and for the production of the fermentation product. The invention further relates to nucleic acid sequences encoding eukaryotic xylose isomerases and xylulose kinases as obtainable from anaerobic fungi.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 July 2003 (31.07.2003)

PCT

(10) International Publication Number
WO 03/062430 A1

(51) International Patent Classification⁷: **C12N 15/61**,
15/54, 1/14, 1/19, C12P 7/10, 7/20, 7/46, 7/54, 7/56,
13/04, 35/00

(21) International Application Number: PCT/NL03/00049

(22) International Filing Date: 23 January 2003 (23.01.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
02075266.3 23 January 2002 (23.01.2002) EP

(71) Applicant (for all designated States except US): **ROYAL NEDALCO B.V.** [NL/NL]; P.O. Box 6, NL-4600 AA Bergen Op Zoom (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **OP DEN CAMP, Hubertus, Johannes, Marie** [NL/NL]; Randenrade 60, NL-5807 BM Oostrum (NL). **HARHANGI, Harry, Ramanoedj** [NL/NL]; Mooieweg 84, NL-6836 AJ Arnhem (NL). **VAN DER DRIFT, Christiaan** [NL/NL]; Windvleugel 36, NL-6581 DX Malden (NL).

(74) Agent: **VAN WESTENBRUGGE, Andries**; Nederlandsch Octrooibureau, Scheveningseweg 82, P.O. Box 29720, NL-2502 LS The Hague (NL).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: FERMENTATION OF PENTOSE SUGARS

(57) **Abstract:** The present invention relates to host cells transformed with a nucleic acid sequence encoding a eukaryotic xylose isomerase obtainable from an anaerobic fungus. When expressed, the sequence encoding the xylose isomerase confers to the host cell the ability to convert xylose to xylulose which may be further metabolised by the host cell. Thus, the host cell is capable of growth on xylose as carbon source. The host cell preferably is a eukaryotic microorganism such as a yeast or a filamentous fungus. The invention further relates to processes for the production of fermentation products such as ethanol, in which a host cell of the invention uses xylose for growth and for the production of the fermentation product. The invention further relates to nucleic acid sequences encoding eukaryotic xylose isomerases and xylulose kinases as obtainable from anaerobic fungi.



WO 03/062430 A1

Fermentation of pentose sugarsField of the invention

5 The present invention relates to host cells transformed with a nucleic acid sequence encoding a eukaryotic xylose isomerase. The xylose isomerase is expressed in the host cell to confer the ability of isomerising xylose to xylulose. The host cell is used in a process for the production of ethanol and other fermentation products by fermentation of a pentose-containing medium. The present invention further relates to
10 nucleic acid sequences encoding eukaryotic xylose isomerases.

Background of the invention

Large-scale consumption of the traditional, fossil fuels (petroleum-based fuels) in the last few decades has contributed to high levels of pollution. Moreover, the
15 realisation that the world stock of petroleum is not boundless, combined with the growing environmental awareness, has stimulated new initiatives to investigate the feasibility of alternative fuels such as ethanol, which could realise a 60-90% decrease in CO₂ production. Although biomass-derived ethanol may be produced by fermentation of hexose sugars that are obtained from many different sources, so far,
20 however, the substrates for industrial scale production or fuel alcohol are cane sugar and corn starch. The drawback of these substrates are the high costs.

Expanding fuel ethanol production requires the ability to use lower-cost feedstocks. Presently, only lignocellulosic feedstock from plant biomass would be available in sufficient quantities to substitute the crops used for ethanol production. The
25 major fermentable sugars from lignocellulosic materials are glucose and xylose, constituting respectively about 40% and 25% of lignocellulose. However, most yeasts that are capable of alcoholic fermentation, like *Saccharomyces cerevisiae*, are not capable of using xylose as a carbon source. Additionally, no organisms are known that can ferment xylose to ethanol with both a high ethanol yield and a high ethanol
30 productivity. To enable the commercial production of ethanol from lignocellulose hydrolysate, an organism possessing both these properties would be required. Thus it is an object of the present invention to provide for a yeast that is capable of both alcoholic fermentation and of using xylose as a carbon source.

D-xylose is metabolisable by numerous microorganisms such as enteric bacteria, some yeasts and fungi. In most xylose-utilising bacteria, xylose is directly isomerised to D-xylulose by xylose (glucose) isomerase (XI). Filamentous fungi and yeasts, are however not capable of this one-step isomerisation and first reduce xylose to xylitol by the action of xylose reductase (XR) after which the xylitol is converted to xylulose by xylitol dehydrogenase (XDH). The first step requires NAD(P)H as a co-factor whereas the second step requires NAD⁺. The xylulose that is produced subsequently enters the pentose phosphate pathway (PPP) after it is phosphorylated by xylulose kinase (XK). Anaerobic fermentation of xylose to ethanol is not possible in organisms with a strictly NADPH dependent xylose reductase (XR). This is because xylitol dehydrogenase (XDH) is strictly NAD⁺ dependent resulting in a redox imbalance (*i.e.*, NAD⁺ depletion). To solve the redox imbalance under anaerobic conditions, the organism produces by-products such as glycerol and xylitol. Similarly, aerobic production of β -lactams on xylose is also negatively influenced as compared to β -lactam production on glucose. A likely cause for these low yields again are a relatively high demand of reducing equivalents in the form of NADPH in this route, compared to the use of glucose (W.M. van Gulik et al. Biotechnol. Bioeng. Vol. 68, No. 6, June 20, 2000).

Over the years many attempts have been made to introduce xylose metabolism in *S. cerevisiae* and similar yeasts, as reviewed in Zaldivar et al. (2001, Appl. Microbiol. Biotechnol. 56: 17-34). One approach concerns the expression of at least genes encoding a xylose (aldose) reductase and a xylitol dehydrogenase, e.g. the *XYL1* and *XYL2* of *Pichia stipitis*, in *S. cerevisiae* (US 5,866,382; WO 95/13362; and WO 97/42307). Although this approach enables growth of *S. cerevisiae* on xylose, it generally suffers from a low ethanol productivity and/or yield as well as a high xylitol production, mainly as a result of the redox imbalance between XR and XDH.

The expression of a XI in *S. cerevisiae* or related yeast or in filamentous fungi would circumvent the redox imbalance and consequent xylitol production and excretion. Xylose isomerase genes from several bacteria have been inserted in *S. cerevisiae*, however, expression of mesophilic prokaryotic XIs in *S. cerevisiae* did not lead to active XI (Amore and Hollenberg, 1989, Nucleic Acids Res. 17: 7515; Amore et al., 1989, Appl. Microbiol. Biotechnol. 30: 351-357; Chan et al., 1986, Biotechnol. Lett 8: 231-234; Chan et al., 1989, Appl. Microbiol. Biotechnol. 31: 524-528; Ho et al., 1983, Fed. Proc. Fed. Am. Soc. Exp. Biol. 42: 2167; Hollenberg, 1987, EBC-

Symposium on Brewer's Yeast, Helsinki (Finland), 24-25 Nov 1986; Sarthy et al., 1987, Appl. Environ. Microbiol. 53: 1996-2000; Ueng et al., 1985, Biotechnol. Lett. 7: 153-158). Nevertheless, two XIs from thermophilic bacteria expressed in *S. cerevisiae* showed a specific activity of 1 μmol per minute per mg^{-1} at 85°C (Bao et al., 1999, Weishengwu-Xuebao 39: 49-54; Walfridson et al., 1996, Appl. Environ. Microbiol. 61: 4184-4190). However, at physiological temperature for *S. cerevisiae* (20-35°C) only a few percent of this activity is left, which is not sufficient for efficient alcoholic fermentation from xylose. Thus, there is still a need for nucleic acids encoding an XI that can be expressed in yeasts to provide sufficient XI activity under physiological conditions to allow for the use of xylose as carbon source.

Description of the invention

Definitions

Xylose isomerase

The enzyme "xylose isomerase" (EC 5.3.1.5) is herein defined as an enzyme that catalyses the direct isomerisation of D-xylose into D-xylulose and vice versa. The enzyme is also known as a D-xylose ketoisomerase. Some xylose isomerases are also capable of catalysing the conversion between D-glucose and D-fructose and are therefore sometimes referred to as glucose isomerase. Xylose isomerases require magnesium as cofactor. Xylose isomerases of the invention may be further defined by their amino acid sequence as herein described below. Likewise xylose isomerases may be defined by the nucleotide sequences encoding the enzyme as well as by nucleotide sequences hybridising to a reference nucleotide sequence encoding a xylose isomerase as herein described below.

A unit (U) of xylose isomerase activity is herein defined as the amount of enzyme producing 1 nmol of xylulose per minute, in a reaction mixture containing 50 mM phosphate buffer (pH 7.0), 10 mM xylose and 10 mM MgCl_2 , at 37°C. Xylulose formed was determined by the method of Dische and Borenfreund (1951, J. Biol. Chem. 192: 583-587) or by HPLC as described in the Examples.

Sequence identity and similarity

Sequence identity is herein defined as a relationship between two or more amino acid (polypeptide or protein) sequences or two or more nucleic acid (polynucleotide) sequences, as determined by comparing the sequences. In the art, "identity" also means

the degree of sequence relatedness between amino acid or nucleic acid sequences, as the case may be, as determined by the match between strings of such sequences.

"Similarity" between two amino acid sequences is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one polypeptide to the
5 sequence of a second polypeptide. "Identity" and "similarity" can be readily calculated by known methods, including but not limited to those described in (Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part I, Griffin, A. M., and
10 Griffin, H. G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heine, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., SIAM J. Applied Math., 48:1073 (1988).

Preferred methods to determine identity are designed to give the largest match
15 between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include e.g. the GCG program package (Devereux, J., et al., Nucleic Acids Research 12 (1):387 (1984)), BestFit, BLASTP, BLASTN, and FASTA (Altschul, S. F. et al., J. Mol. Biol. 215:403-410
20 (1990). The BLAST X program is publicly available from NCBI and other sources (BLAST Manual, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., J. Mol. Biol. 215:403-410 (1990). The well-known Smith Waterman algorithm may also be used to determine identity.

Preferred parameters for polypeptide sequence comparison include the following:
25 Algorithm: Needleman and Wunsch, J. Mol. Biol. 48:443-453 (1970); Comparison matrix: BLOSSUM62 from Hentikoff and Hentikoff, Proc. Natl. Acad. Sci. USA. 89:10915-10919 (1992); Gap Penalty: 12; and Gap Length Penalty: 4. A program useful with these parameters is publicly available as the "Ogap" program from Genetics Computer Group, located in Madison, WI. The aforementioned parameters are the
30 default parameters for amino acid comparisons (along with no penalty for end gaps).

Preferred parameters for nucleic acid comparison include the following:
Algorithm: Needleman and Wunsch, J. Mol. Biol. 48:443-453 (1970); Comparison matrix: matches=+10, mismatch=0; Gap Penalty: 50; Gap Length Penalty: 3. Available

as the Gap program from Genetics Computer Group, located in Madison, Wis. Given above are the default parameters for nucleic acid comparisons.

Optionally, in determining the degree of amino acid similarity, the skilled person may also take into account so-called "conservative" amino acid substitutions, as will be clear to the skilled person. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulphur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, and asparagine-glutamine. Substitutional variants of the amino acid sequence disclosed herein are those in which at least one residue in the disclosed sequences has been removed and a different residue inserted in its place. Preferably, the amino acid change is conservative. Preferred conservative substitutions for each of the naturally occurring amino acids are as follows: Ala to ser; Arg to lys; Asn to gln or his; Asp to glu; Cys to ser or ala; Gln to asn; Glu to asp; Gly to pro; His to asn or gln; Ile to leu or val; Leu to ile or val; Lys to arg; gln or glu; Met to leu or ile; Phe to met, leu or tyr; Ser to thr; Thr to ser; Trp to tyr; Tyr to trp or phe; and, Val to ile or leu.

Hybridising nucleic acid sequences

Nucleotide sequences encoding xylose isomerases or xylulose kinases of the invention may also be defined by their capability to hybridise with the nucleotide sequences of SEQ ID NO. 2 or SEQ ID NO. 4, respectively, under moderate, or preferably under stringent hybridisation conditions. Stringent hybridisation conditions are herein defined as conditions that allow a nucleic acid sequence of at least about 25, preferably about 50 nucleotides, 75 or 100 and most preferably of about 200 or more nucleotides, to hybridise at a temperature of about 65°C in a solution comprising about 1 M salt, preferably 6 x SSC or any other solution having a comparable ionic strength, and washing at 65°C in a solution comprising about 0.1 M salt, or less, preferably 0.2 x SSC or any other solution having a comparable ionic strength. Preferably, the

hybridisation is performed overnight, i.e. at least for 10 hours and preferably washing is performed for at least one hour with at least two changes of the washing solution. These conditions will usually allow the specific hybridisation of sequences having about 90% or more sequence identity.

5 Moderate conditions are herein defined as conditions that allow a nucleic acid sequences of at least 50 nucleotides, preferably of about 200 or more nucleotides, to hybridise at a temperature of about 45°C in a solution comprising about 1 M salt, preferably 6 x SSC or any other solution having a comparable ionic strength, and washing at room temperature in a solution comprising about 1 M salt, preferably 6 x
10 SSC or any other solution having a comparable ionic strength. Preferably, the hybridisation is performed overnight, i.e. at least for 10 hours, and preferably washing is performed for at least one hour with at least two changes of the washing solution. These conditions will usually allow the specific hybridisation of sequences having up to 50% sequence identity. The person skilled in the art will be able to modify these
15 hybridisation conditions in order to specifically identify sequences varying in identity between 50% and 90%.

Operably linked

As used herein, the term "operably linked" refers to a linkage of polynucleotide elements in a functional relationship. A nucleic acid is "operably linked" when it is
20 placed into a functional relationship with another nucleic acid sequence. For instance, a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the coding sequence. Operably linked means that the DNA sequences being linked are typically contiguous and, where necessary to join two protein coding regions, contiguous and in reading frame.

25 Promoter

As used herein, the term "promoter" refers to a nucleic acid fragment that functions to control the transcription of one or more genes, located upstream with respect to the direction of transcription of the transcription initiation site of the gene, and is structurally identified by the presence of a binding site for DNA-dependent RNA
30 polymerase, transcription initiation sites and any other DNA sequences, including, but not limited to transcription factor binding sites, repressor and activator protein binding sites, and any other sequences of nucleotides known to one of skill in the art to act directly or indirectly to regulate the amount of transcription from the promoter. A

In a first aspect the present invention relates to a transformed host cell that has the ability of isomerising xylose to xylulose. The ability of isomerising xylose to xylulose is conferred to the host cell by transformation of the host cell with a nucleic acid construct comprising a nucleotide sequence encoding a xylose isomerase. The transformed host cell's ability to isomerise xylose into xylulose is the direct isomerisation of xylose to xylulose. This is understood to mean that xylose isomerised into xylulose in a single reaction catalysed by a xylose isomerase, as opposed to the two step conversion of xylose into xylulose via a xylitol intermediate as catalysed by xylose reductase and xylitol dehydrogenase, respectively.

Preferably, expression of the nucleotide sequence in the host cell produces a xylose isomerase with a K_m for xylose that is less than 50, 40, 30 or 25 mM, more preferably, the K_m for xylose is about 20 mM or less.

A nucleotide sequence encoding the xylose isomerase may be selected from the group consisting of:

- (a) nucleotide sequences encoding a polypeptide comprising an amino acid sequence that has at least 40, 45, 49, 50, 53, 55, 60, 70, 80, 90, 95, 97, 98, or 99% sequence identity with the amino acid sequence of SEQ ID NO. 1;
- (b) nucleotide sequences comprising a nucleotide sequence that has at least 40, 50, 55, 56, 57, 60, 70, 80, 90, 95, 97, 98, or 99% sequence identity with the nucleotide sequence of SEQ ID NO. 2;
- (c) nucleotide sequences the complementary strand of which hybridises to a nucleic acid molecule sequence of (a) or (b);
- (d) nucleotide sequences the sequence of which differs from the sequence of a nucleic acid molecule of (c) due to the degeneracy of the genetic code.

The nucleotide sequence preferably encodes a eukaryotic xylose isomerase, i.e. a xylose isomerase with an amino acid sequence that is identical to that of a xylose isomerase that naturally occurs in an eukaryotic organism. Expression of a eukaryotic xylose isomerase increases the likelihood that the xylose isomerase is expressed in active form in a eukaryotic host cell such as yeast, as opposed to the mesophilic prokaryotic xylose isomerases. More preferably the nucleotide sequence encodes a plant xylose isomerase (e.g. from *Hordeum vulgare*) or a fungal xylose isomerase (e.g. from a *Basidiomycete*). Most preferably, however, the nucleotide sequence encodes a xylose isomerase from an anaerobic fungus, to further increase the likelihood of expression in enzymatically active form in a eukaryotic host cell, particularly in yeast. Most preferred are nucleotide sequences encoding a xylose isomerase from an anaerobic fungus that belongs to the families *Neocallimastix*, *Caecomyces*, *Piromyces*, *Orpinomyces*, or *Ruminomyces*.

A host cell for transformation with a nucleotide sequence encoding a xylose isomerase preferably is a host capable of active or passive xylose transport into the cell. The host cell preferably contains active glycolysis, the pentose phosphate pathway and preferably contains xylulose kinase activity so that the xylulose isomerised from xylose may be metabolised to pyruvate. The host further preferably contains enzymes for conversion of pyruvate to a desired fermentation product such as ethanol, ethylene or lactic acid. A preferred host cell is a host cell that is naturally capable of alcoholic fermentation, preferably, anaerobic alcoholic fermentation. The host cell further preferably has a high tolerance to ethanol and organic acids like lactic acid, acetic acid or formic acid and sugar degradation products such as furfural and hydroxy-

methyolfurfural. Any of these characteristics or activities of the host cell may be naturally present in the host cell or may be introduced or modified by genetic modification. A suitable host cell is a microorganism like a bacterium or a fungus, however, most suitable as host cell are yeasts or filamentous fungi.

5 Yeasts are herein defined as eukaryotic microorganisms and include all species of the subdivision Eumycotina (Alexopoulos, C. J., 1962, In: Introductory Mycology, John Wiley & Sons, Inc., New York) that predominantly grow in unicellular form. Yeasts may either grow by budding of a unicellular thallus or may grow by fission of the organism. Preferred yeasts as host cells belong to the genera *Saccharomyces*,
10 *Kluyveromyces*, *Candida*, *Pichia*, *Schizosaccharomyces*, *Hansenula*, *Kloeckera*, *Schwanniomyces*, and *Yarrowia*. Preferably the yeast is capable of anaerobic fermentation, more preferably anaerobic alcoholic fermentation.

Filamentous fungi are herein defined as eukaryotic microorganisms that include all filamentous forms of the subdivision Eumycotina. These fungi are characterized by a
15 vegetative mycelium composed of chitin, cellulose, and other complex polysaccharides. The filamentous fungi of the present invention are morphologically, physiologically, and genetically distinct from yeasts. Vegetative growth by filamentous fungi is by hyphal elongation and carbon catabolism of most filamentous fungi is obligately aerobic. Preferred filamentous fungi as host cells belong to the genera *Aspergillus*,
20 *Trichoderma*, *Humicola*, *Acremonium*, *Fusarium*, and *Penicillium*.

Over the years suggestions have been made for the introduction of various organisms for the production of bio-ethanol from crop sugars. In practice, however, all major bio-ethanol production processes have continued to use the yeasts of the genus *Saccharomyces* as ethanol producer. This is due to the many attractive features of
25 *Saccharomyces* species for industrial processes, i.e., a high acid-, ethanol- and osmo-tolerance, capability of anaerobic growth, and of course its high alcoholic fermentative capacity. Preferred yeast species as host cells include *S. cerevisiae*, *S. bulderi*, *S. barnetti*, *S. exiguus*, *S. uvarum*, *S. diastaticus*, *K. lactis*, *K. marxianus*, *K. fragilis*.

The host cell is transformed with a nucleic acid construct as further defined
30 below and may comprise a single but preferably comprises multiple copies of the nucleic acid construct. The nucleic acid construct may be maintained episomally and thus comprise a sequence for autonomous replication, such as an ARS sequence. Suitable episomal nucleic acid constructs may e.g. be based on the yeast 2 μ or pKD1

(Fleer et al., 1991, Biotechnology 9:968-975) plasmids. Preferably, however, the nucleic acid construct is integrated in one or more copies into the genome of the host cell. Integration into the host cell's genome may occur at random by illegitimate recombination but preferably nucleic acid construct is integrated into the host cell's genome by homologous recombination as is well known in the art of fungal molecular genetics (see e.g. WO 90/14423, EP-A-0 481 008, EP-A-0 635 574 and US 6,265,186).

In a preferred transformed host cell according to the invention, the nucleic acid construct confers to the host cell the ability to grow on xylose as carbon source, preferably as sole carbon source, and preferably under anaerobic conditions, whereby preferably the transformed host produce essentially no xylitol, e.g. the xylitol produced is below the detection limit or e.g. less than 5, 2, 1% of the carbon consumed on a molar basis. The transformed host cell has the ability to grow on xylose as sole carbon source at a rate of at least 0.01, 0.02, 0.05, 0.1 or 0.2 h⁻¹. The transformed host cell of the invention thus expresses a xylose isomerase at a specific activity level defined above.

A host cell may comprises further genetic modifications that result in one or more of the characteristics selected from the group consisting of (a) increase transport of xylose into the host cell; (b) increased xylulose kinase activity; (c) increased flux of the pentose phosphate pathway; (d) decreased sensitivity to catabolite repression; (e) increased tolerance to ethanol, osmolarity or organic acids; and, (f) reduced production of by-products. By-products are understood to mean carbon-containing molecules other than the desired fermentation product and include e.g. xylitol, glycerol and/or acetic acid. Such genetic modifications may be introduced by classical mutagenesis and screening and/or selection for the desired mutant. Alternatively, the genetic modifications may consist of overexpression of endogenous genes and/or expression of a heterologous genes and/or the inactivation of endogenous genes. The genes are preferably chosen from genes encoding a hexose or pentose transporter; a xylulose kinase such as the xylulose kinase genes from *S. cerevisiae* (*XKS1* Deng and Ho, 1990, Appl. Biochem. Biotechnol. 24-25: 193-199) or *Piromyces* (*xylB*, i.e. SEQ ID NO. 4); an enzyme from the pentose phosphate pathway such as a transaldolase (*TAL1*) or a transketolase (*TKL1*) (see e.g. Meinander et al., 1995, Pharmacol.Toxicol. Suppl.2: 45) glycolytic enzymes, ethanologenic enzymes such as alcohol dehydrogenases. Preferred endogenous genes for inactivation include a hexose kinase gene e.g. the *S. cerevisiae*

HXK2 gene (see Diderich et al., 2001, Appl. Environ. Microbiol. 67: 1587-1593); the *S. cerevisiae* *MIG1* or *MIG2* genes; (unspecific) aldose reductase genes such as the *S. cerevisiae* *GRE3* gene (Träff et al., 2001, Appl. Environm. Microbiol. 67: 5668-5674); genes for enzymes involved in glycerol metabolism such as the *S. cerevisiae* glycerol-phosphate dehydrogenase 1 and/or 2 genes; or (hybridising) homologues of the genes in other host species. Further preferred modifications of host cells for xylose fermentation are reviewed in Zaldivar et al. (2001, *supra*).

In another aspect the invention relates to a transformed host cell for the production of fermentation products other than ethanol. Such non-ethanolic fermentation products include in principle any bulk or fine chemical that is producible by eukaryotic microorganism such as a yeast or a filamentous fungus. Such fermentation products include e.g. lactic acid, acetic acid, succinic acid, amino acids, 1,3-propane-diol, ethylene, glycerol, β -lactam antibiotics and cephalosporins.

Transformation of host cells with the nucleic acid constructs of the invention and additional genetic modification of host cells, preferably yeasts, as described above may be carried out by methods well known in the art. Such methods are e.g. known from standard handbooks, such as Sambrook and Russel (2001) "Molecular Cloning: A Laboratory Manual (3rd edition), Cold Spring Harbor Laboratory, *Cold Spring Harbor Laboratory Press*, or F. Ausubel et al, eds., "Current protocols in molecular biology", Green Publishing and Wiley Interscience, New York (1987). Methods for transformation and genetic modification of fungal host cells are known from e.g. EP-A-0 635 574, WO 98/46772, WO 99/60102 and WO 00/37671.

In another aspect the invention relates to a nucleic acid construct comprising a nucleotide sequence encoding a xylose isomerase as defined above and used for transformation of a host cell as defined above. In the nucleic acid construct, the nucleotide sequence encoding the xylose isomerase preferably is operably linked to a promoter for control and initiation of transcription of the nucleotide sequence in a host cell as defined below. The promoter preferably is capable of causing sufficient expression of the xylose isomerase in the host cell, to confer to the host cell the ability to isomerise xylose into xylulose. Preferably, the promoter causes a specific xylose isomerase activity in the host cell as defined above. Promoters useful in the nucleic acid constructs of the invention include both constitutive and inducible natural promoters as well as engineered promoters. A preferred promoter for use in the present invention

will in addition be insensitive to catabolite (glucose) repression and/or will preferably not require xylose for induction. Promoters having these characteristics are widely available and known to the skilled person. Suitable examples of such promoters include e.g. yeast promoters from glycolytic genes, such as the yeast phosphofructokinase (PPK), triose phosphate isomerase (TPI), glyceraldehyde-3-phosphate dehydrogenase (GPD, TDH3 or GAPDH), pyruvate kinase (PYK), phosphoglycerate kinase (PGK) promoters; more details about such promoters may be found in (WO 93/03159). Other useful promoters are ribosomal protein encoding gene promoters, the lactase gene promoter (LAC4), alcohol dehydrogenase promoters (ADH1, ADH4, and the like), and the enolase promoter (ENO). Other promoters, both constitutive and inducible and enhancers or upstream activating sequences will be known to those of skill in the art. The promoters used in the nucleic acid constructs of the present invention may be modified, if desired, to affect their control characteristics. Preferably, the promoter used in the nucleic acid construct for expression of the xylose isomerase is homologous to the host cell in which the xylose isomerase is expressed.

In the nucleic acid construct, the 3'-end of the nucleotide acid sequence encoding the xylose isomerase preferably is operably linked to a transcription terminator sequence. Preferably the terminator sequence is operable in a host cell of choice, such as e.g. the yeast species of choice. In any case the choice of the terminator is not critical, it may e.g. be from any yeast gene, although terminators may sometimes work if from a non-yeast, eukaryotic, gene. The transcription termination sequence further preferably comprises a polyadenylation signal.

Optionally, a selectable marker may be present in the nucleic acid construct. As used herein, the term "marker" refers to a gene encoding a trait or a phenotype which permits the selection of, or the screening for, a host cell containing the marker. The marker gene may be an antibiotic resistance gene whereby the appropriate antibiotic can be used to select for transformed cells from among cells that are not transformed. Examples of suitable antibiotic resistance markers include e.g. dihydrofolate reductase, hygromycin-B-phosphotransferase, 3'-O-phosphotransferase II (kanamycin, neomycin and G418 resistance). Although the of antibiotic resistance markers may be most convenient for the transformation of polyploid host cells, preferably however, non-antibiotic resistance markers are used, such as auxotrophic markers (URA3, TRP1, LEU2) or the *S. pombe* TPI gene (described by Russell P R, 1985, Gene 40: 125-130).

In a preferred embodiment the host cells transformed with the nucleic acid constructs are marker gene free. Methods for constructing recombinant marker gene free microbial host cells are disclosed in EP-A-0 635 574 and are based on the use of bidirectional markers such as the *A. nidulans* amdS (acetamidase) gene or the yeast URA3 and LYS2 genes. Alternatively, a screenable marker such as Green Fluorescent Protein, *lacZ*, luciferase, chloramphenicol acetyltransferase, beta-glucuronidase may be incorporated into the nucleic acid constructs of the invention allowing to screen for transformed cells.

Optional further elements that may be present in the nucleic acid constructs of the invention include, but are not limited to, one or more leader sequences, enhancers, integration factors, and/or reporter genes, intron sequences, centromeres, telomeres and/or matrix attachment (MAR) sequences. The nucleic acid constructs of the invention may further comprise a sequence for autonomous replication, such as an ARS sequence. Suitable episomal nucleic acid constructs may e.g. be based on the yeast 2 μ or pKD1 (Fleer et al., 1991, *Biotechnology* 9:968-975) plasmids. Alternatively the nucleic acid construct may comprise sequences for integration, preferably by homologous recombination. Such sequences may thus be sequences homologous to the target site for integration in the host cell's genome. The nucleic acid constructs of the invention can be provided in a manner known per se, which generally involves techniques such as restricting and linking nucleic acids/nucleic acid sequences, for which reference is made to the standard handbooks, such as Sambrook and Russel (2001) "Molecular Cloning: A Laboratory Manual (3rd edition), Cold Spring Harbor Laboratory, *Cold Spring Harbor Laboratory Press*, or F. Ausubel et al, eds., "Current protocols in molecular biology", Green Publishing and Wiley Interscience, New York (1987).

In another aspect the invention relates to a nucleic acid molecule comprising a nucleotide sequence that encodes a xylose isomerase. The nucleic acid molecule is preferably selected from the group consisting of:

- (a) nucleic acid molecules encoding a polypeptide comprising an amino acid sequence that has at least 50, 53, 54, 55, 60, 70, 80, 90, 95, 97, 98, or 99% sequence identity with the amino acid sequence of SEQ ID NO. 1;
- (b) nucleic acid molecules comprising a nucleotide sequence that has at least 50, 56, 57, 58, 60, 70, 80, 90, 95, 97, 98, or 99% sequence identity with the nucleotide sequence of SEQ ID NO. 2;

- (c) nucleic acid molecules the complementary strand of which hybridises to a nucleic acid molecule sequence of (a) or (b);
- (d) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (c) due to the degeneracy of the genetic code.

5 Alternatively, a nucleic acid molecule of (a) may encode a polypeptide comprising an amino acid sequence that has at least 67, 68, 69, 70, 80, 90, 95, 97, 98, or 99% sequence similarity with the amino acid sequence of SEQ ID NO. 1. A nucleic acid molecule of (c) preferably hybridises under moderate conditions, more preferably under stringent conditions as herein defined above. Preferably the nucleic acid

10 molecule is from a eukaryote, more preferably from a eukaryotic microorganism such as a fungus, most preferably from an anaerobic fungus, such as e.g. that anaerobic fungi described above.

Yet another aspect of the invention relates to a nucleic acid molecule comprising a nucleotide sequence that encodes a xylulose kinase, preferably a D-xylulose kinase. A

15 D-xylulose kinase (EC 2.7.1.17; also referred to as a D-xylulokinase) is herein defined as an enzyme that catalyses the conversion of D-xylulose into xylulose-5-phosphate.

The nucleic acid molecule is preferably selected from the group consisting of:

- (a) nucleic acid molecules encoding a polypeptide comprising an amino acid sequence that has at least 45, 47, 48, 49, 50, 55, 60, 70, 80, 90, 95, 97, 98, or 99% sequence
- 20 identity with the amino acid sequence of SEQ ID NO. 3;
- (b) nucleic acid molecules comprising a nucleotide sequence that has at least 30, 37, 38, 39, 40, 50, 60, 70, 80, 90, 95, 97, 98, or 99% sequence identity with the nucleotide sequence of SEQ ID NO. 4;
- (c) nucleic acid molecules the complementary strand of which hybridises to a nucleic
- 25 acid molecule sequence of (a) or (b); and,
- (d) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (c) due to the degeneracy of the genetic code.

Alternatively, a nucleic acid molecule of (a) may encode a polypeptide comprising an amino acid sequence that has at least 64, 65, 66, 70, 80, 90, 95, 97, 98,

30 or 99% sequence similarity with the amino acid sequence of SEQ ID NO. 3. A nucleic acid molecule of (c) preferably hybridises under moderate conditions, more preferably under stringent conditions as herein defined above. Preferably the nucleic acid molecule is from a eukaryote, more preferably from a eukaryotic microorganism such

as a fungus, most preferably from an anaerobic fungus, such as e.g. that anaerobic fungi described above.

In a further aspect the invention relates to fermentation processes in which the transformed host cells of the invention are used for the fermentation of carbon source comprising a source of xylose, such as xylose. In addition to a source of xylose the carbon source in the fermentation medium may also comprise a source of glucose. The source of xylose or glucose may be xylose or glucose as such or may be any carbohydrate oligo- or polymer comprising xylose or glucose units, such as e.g. lignocellulose, xylans, cellulose, starch and the like. For release of xylose or glucose units from such carbohydrates, appropriate carbohydrases (such as xylanases, glucanases, amylases and the like) may be added to the fermentation medium or may be produced by the transformed host cell. In the latter case the transformed host cell may be genetically engineered to produce and excrete such carbohydrases. In a preferred process the transformed host cell ferments both the xylose and glucose, preferably simultaneously in which case preferably a transformed host cell is used which is insensitive to glucose repression to prevent diauxic growth. In addition to a source of xylose (and glucose) as carbon source, the fermentation medium will further comprise the appropriate ingredient required for growth of the transformed host cell. Compositions of fermentation media for growth of microorganisms such as yeasts are well known in the art.

The fermentation process is a process for the production of a fermentation product such as ethanol, lactic acid, acetic acid, succinic acid, amino acids, 1,3-propane-diol, ethylene, glycerol, β -lactam antibiotics such as Penicillin G or Penicillin V and fermentative derivatives thereof and cephalosporins. The fermentation process may be an aerobic or an anaerobic fermentation process. An anaerobic fermentation process is herein defined as a fermentation process run in the absence of oxygen or in which substantially no oxygen is consumed, e.g. less than 5 mmol/L/h, and wherein organic molecules serve as both electron donor and electron acceptors. In the absence of oxygen, NADH produced in glycolysis and biomass formation, cannot be oxidised by oxidative phosphorylation. To solve this problem many microorganisms use pyruvate or one of its derivatives as an electron and hydrogen acceptor thereby regenerating NAD^+ . Thus, in a preferred anaerobic fermentation process pyruvate is

used as an electron (and hydrogen acceptor) and is reduced to fermentation products such as ethanol, lactic acid, 1,3-propanediol, ethylene, acetic acid or succinic acid.

The fermentation process is preferably run at a temperature that is optimal for the transformed host cell. Thus, for most yeasts or fungal host cells, the fermentation
5 process is performed at a temperature which is less than 38°C. For yeast or filamentous fungal host cells, the fermentation process is preferably performed at a temperature which is lower than 35, 33, 30 or 28°C and at a temperature which is higher than 20, 22, or 25°C.

A preferred process is a process for the production of ethanol, whereby the
10 process comprises the steps of: (a) fermenting a medium containing a source of xylose with a transformed host cell as defined above, whereby the host cell ferments xylose to ethanol; and optionally, (b) recovery of the ethanol. The fermentation medium may also comprise a source of glucose that is also fermented to ethanol. In the process the volumetric ethanol productivity is preferably at least 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 5.0 or
15 10.0 g ethanol per litre per hour. The ethanol yield on xylose and/or glucose in the process preferably is at least 50, 60, 70, 90, 95 or 98%. The ethanol yield is herein defined as a percentage of the theoretical yield, which, for glucose and xylose is 0.51 g. ethanol per g. glucose or xylose.

In a further aspect the invention relates to a process for producing a fermentation
20 product selected from the group consisting of lactic acid, acetic acid, succinic acid, amino acids, 1,3-propane-diol, ethylene, glycerol, β -lactam antibiotics and cephalosporins. The process preferably comprises the steps of (a) fermenting a medium containing a source of xylose with a transformed host cell as defined herein above, whereby the host cell ferments xylose to the fermentation product, and optionally, (b)
25 recovery of the fermentation product. In a preferred process, the medium also contains a source of glucose.

Description of the figures

30 **Figure 1.** Growth curves of *S. cerevisiae* transformant grown on medium containing 25 mM galactose and 100 mM xylose as carbon source. Transformant pYes contains a yeast expression vector without insertion. Transformants 14.3, 16.2.1 and 16.2.2 are transformed with the pYES vector containing the *Piromyces* sp. E2 xylose isomerase

coding sequence.

Examples

Example 1

Cloning of *Piromyces* xylanase isomerase and xylulose kinase cDNAs

5 Organism and growth conditions

The anaerobic fungus *Piromyces* sp. E2 (ATCC 76762), isolated from faeces of an Indian elephant, was grown anaerobically under N₂/CO₂ (80 %/20 %) at 39°C in medium M2 supplemented with various carbon sources (24). Carbon sources used were Avicel (microcrystalline cellulose type PH 105, Serva, Germany), fructose or xylose (all 10 0.5 %, w/v). After growth ceased, as judged by hydrogen production, the cells were harvested by centrifugation (15,000 x g, 4°C, 15 min;) or by filtration over nylon gauze (30 µm pore size).

Preparation of cell-free extract

The fungal cells were washed with deionized water to remove medium 15 components. Cell-free extracts were prepared by freezing the cells in liquid nitrogen and subsequent grinding with glass beads (0.10-0.11 mm diameter) in a mortar. Tris/HCl buffer (100 mM, pH 7.0) was added to the powder (1:1, w/v) and after thawing for 15 min the suspension was centrifuged (18,000 x g, 4°C, 15 min). The clear supernatant was used as a source of intracellular enzymes.

20 Enzyme assays

Xylose isomerase activity was assayed at 37°C in a reaction mixture containing 50 mM phosphate buffer (pH 7.0), 10 mM xylose, 10 mM MgCl₂ and a suitable amount of cell-free extract. The amount of xylulose formed was determined by the cysteine-carbazole method (9). Xylulose kinase and xylose reductase activities were assayed as 25 described by Witteveen et al. (28). One unit of activity is defined as the amount of enzyme producing 1 nmol of xylulose per min under the assay conditions. Xylulose formed was determined by the method of Dische and Borenfreund (Dische and Borenfreund, 1951, J. Biol. Chem. 192: 583-587) or by HPLC using a Biorad HPX-87N Column operated at 80°C and eluted at 0.6 ml/min using 0.01 M Na₂HPO₄ as the 30 eluens. Xylose and xylulose were detected by a Refractive Index detector at an internal temperature of 60°C.

Specific activity is expressed as units per mg protein. Protein was determined with the Bio-Rad protein reagent (Bio-Rad Laboratories, Richmond, CA, USA) with bovine γ -globulin as a standard.

5 Random sequencing of a *Piromyces* sp. E2 cDNA library

The cDNA library constructed in the vector lambda ZAPII as described previously (2) was used. An aliquot of this library was converted to pBluescript SK-clones by mass excision with the ExAssist helper phage (Stratagene, La Jolla, CA, USA). Randomly selected clones were sequenced with the M13 reverse primer to
10 obtain 5' part sequences. Uncomplete cDNAs were used to synthesize probes which were used to rescreen the library. To obtain full length sequences subclones were generated in pUC18. Sequencing was performed with the ABI prism 310 automated sequencer with the dRhodamine terminator cycle sequencing ready reaction DNA sequencing kit (Perkin-Elmer Applied Biosystems).

15 Results

Randomly selected clones from a cDNA library of the anaerobic fungus *Piromyces* sp. E2 were sequenced and this resulted in two clones (pH97 and pAK44) which sequences showed high homology to xylose isomerase and D-xylulokinase genes, respectively. The clones were analysed in detail.

20 Clone pH97 did not contain a complete ORF and therefore the cDNA library was rescreened with a probe designed on the basis of sequence data from clone pH97. This resulted in a clone pR3 with an insert of 1669 bp. An ORF encoding a protein of 437 amino acids with high similarity to xylose isomerases could be identified. Although the 5' untranslated region comprises only 4 bp, the presumed starting methionine residue
25 fitted well into an alignment of known xylose isomerase sequences. The 3' untranslated region was 351 bp long and had a high AT content, which is typical for anaerobic fungi. The ORF contained the amino acids shown to be important for interaction with the substrate (catalytic triad His 102, Asp 105, Asp 340 and Lys 235) and binding of magnesium (Glu 232) (14, 26). Further, the two signature patterns (residues 185-194
30 and 230-237) developed for xylose isomerases (20) were present. The *Piromyces* sp. E2 xylose isomerase (XylA) shows the highest homology to the enzymes of *Haemophilus influenza* (52 % identity, 68 % similarity) and *Hordeum vulgare* (49 % identity, 67 %

similarity). The polypeptide deduced from the cDNA sequence corresponds to a molecular mass of 49,395 Da and has a calculated pI of 5.2.

The second clone, pAK44, had an insert of 2041 bp and contained a complete ORF encoding a protein of 494 amino acids with a molecular weight of 53,158 Da and a pI of 5.0. The first methionine is preceded by a 111 bp 5' untranslated region, while the 3' untranslated region comprised 445 bp. Both regions are AT-rich. BLAST and FASTA searches revealed high similarity to xylulokinases. The two phosphate consensus regions defined by Rodriguez-Peña et al. (22) were found at positions 6-23 and 254-270 as shown in a partial alignment. Moreover the signatures for this family of carbohydrate kinase as described in the Prosite database were identified (131-145 and 351-372). The *Piromyces* sp. E2 xylulokinase (XylB) showed highest homology with the XylB protein of *Haemophilus influenza* (46 % identity, 64 % similarity).

Example 2

15 Construction of yeast expression vectors

Expression of xylose isomerase from *Piromyces* sp. E2 in *Saccharomyces cerevisiae*

cDNA from *Piromyces* sp. E2 was used in a PCR reaction with *pfu* polymerase (Stratagene). The primers were designed using the sequences from the 5' and 3' ends of the xylose isomerase gene and also contained a Sfi I and a XbaI restriction site. The PCR product was cloned in the pPICZ α vector (Invitrogen, Carlsbad, CA, USA). To obtain the xylose isomerase gene, the pPICZ α vector was digested with EcoRI and XbaI. The digestion product was ligated into the pYes2 vector (Invitrogen). The pYes2 plasmid with the xylose isomerase gene was transformed into *Saccharomyces cerevisiae* (stam BJ1991, gift from Beth Jones, UvA). The genotype of this strain is: $\text{mat}\alpha$, *leu2*, *trp1*, *ura 3-251*, *prb1-1122* and *pep4-3*.

Transformants were plated on SC plates (0.67% YNB medium + 0.05% L-Leu + 0.05% L-Trp + 2% glucose + 2% agarose). Untransformed cells can not grow on these plates.

Induction

30 Transformed *Saccharomyces cerevisiae* cells were grown on glucose medium at 25°C for 72 h (raffinose can be used as an alternative for glucose). Cells were harvested and resuspended in SC medium with galactose instead of glucose. After 8 h of induction cells were harvested and lysed using glass beads (0.10-0.11 mm diameter) and

“breaking buffer” (50mM phosphate buffer + 5% glycerol + protease inhibitor). After lysis the mixture was centrifuged (18,000 x g, 4°C, 15 min). The clear supernatant was used to determine xylose isomerase activity using the method described above (Example 1). An activity of 10 U per mg protein was measured at 37°C.

5

Example 3

Growth of transformed yeast strains on xylose

Medium composition

Saccharomyces cerevisiae strains were grown on SC-medium with the following composition: 0.67% (w/v) yeast nitrogen base; 0.01% (w/v) L-tryptophan; 0.01% (w/v) L-leucine and either glucose, galactose or xylose, or a combination of these substrates (see below). For agar plates the medium was supplemented with 2% (w/v) bacteriological agar.

Growth experiment

Saccharomyces cerevisiae strain BJ1991 (genotype: $\text{mat}\alpha$, *leu2*, *trp1*, *ura 3-251*, *prb1-1122*, *pep4-3*) transformed with pYes2 without insertion and three selected transformants (16.2.1; 16.2.2 and 14.3) containing pYes2 with the *Piromyces* sp. E2 xylose isomerase gene were grown on SC-agar plates with 10 mM glucose as carbon source. When colonies were visible, single colonies were used to inoculate liquid SVC-medium with 100 mM xylose and 25 mM galactose as carbon sources. Growth was monitored by measuring the increase in optical density at 600 nm on a LKB Ultrospec K spectrophotometer.

Results

The results of the growth experiments are compiled in Figure 1. The culture with the BJ1991 strain transformed with pYes2 without insertion shows an increase in OD₆₀₀ up to 80 h. After this time a gradual decrease is observed. This is caused by aggregation of the yeast cells which is often observed at the end of growth. The cultures with the three transformants do not stop growing after 80 h and show a further increase up to at least 150 h.

30

Example 4Construction of a new, improved, yeast expression vector for constitutive expression of the *Piromyces* sp.E2 xylose isomerase in *Saccharomyces cerevisiae*.

The pPICZ α vector, containing the *Piromyces* sp. E2 gene coding for xylose isomerase, was used as a template for PCR with Vent_R DNA polymerase (New England Biolabs). The primers were designed using the 5' and 3' sequences of the gene coding for xylose isomerase and included an EcoRI and an SpeI site. Additionally the primers were designed to remove the XbaI site found in the pPICZ α construct, replacing it with a stopcodon (TAA). The final product was designed to restore the original open reading frame, without the added aminoacids (his and c-Myc tags) found in the pPICZ α construct. The PCR product was cut with EcoRI and SpeI. The final product was cloned into a vector derived from pYES2 (Invitrogen). In this vector the *GAL1* promoter found in pYES2 was replaced by the *TPII* promoter in order to ensure constitutive expression of the xylose isomerase, thereby eliminating the need for galactose in the medium. The *TPII* promoter was cloned from a modified form of plasmid pYX012 (R&D systems). The promoter was cut out as a NheI-EcoRI fragment. Both the *TPII* promoter and the PCR product of the gene coding for the xylose isomerase were ligated into pYES2 cut with SpeI and XbaI. This plasmid was used to transform *Saccharomyces cerevisiae* strain CEN.PK113-5D (gift from Peter Kötter, Frankfurt). The genotype of the strain is: *MatA ura3-52*. Transformants were selected on mineral medium plates (Verduyn et al.: Effect of benzoic acid on metabolic fluxes in yeasts: a continuous-culture study on the regulation of respiration and alcoholic fermentation.(1992) Yeast 8(7):501-17) with 2% glucose as the carbon source. Untransformed cells cannot grow on these plates.

Transformants were grown on glucose/xylose mixtures in carbon-limited chemostat cultures. Transformants grown under these conditions exhibit high xylose isomerase activities (800 units per mg at 30°C) according to a specific enzyme assay as developed by Kersters-Hildersson et al. (Kinetic characterization of D-xylose isomerases by enzymatic assays using D-sorbitol dehydrogenase. Enz. Microb. Technol. 9 (1987) 145-148). The in vitro activity of xylose isomerase in the cell-free extracts of the transformed *S.cerevisiae* strain was dependent on bivalent cations (Mg²⁺ or Co²⁺) and a relatively low K_m value for xylose of approximately 20 mM was measured.

23

SEQUENCE LISTING

<110> Royal Nedalco B.V.

5 <120> Fermentation of pentose sugars

<130> piromyces xylose isomerase

<140> BO 44829

10 <141> 2001-12-31

<160> 4

<170> PatentIn Ver. 2.1

15

<210> 1

<211> 437

<212> PRT

<213> Piromyces sp.

20

<400> 1

Met	Ala	Lys	Glu	Tyr	Phe	Pro	Gln	Ile	Gln	Lys	Ile	Lys	Phe	Glu	Gly
1				5					10					15	

Lys	Asp	Ser	Lys	Asn	Pro	Leu	Ala	Phe	His	Tyr	Tyr	Asp	Ala	Glu	Lys
			20					25					30		

Glu	Val	Met	Gly	Lys	Lys	Met	Lys	Asp	Trp	Leu	Arg	Phe	Ala	Met	Ala
		35					40					45			

Trp	Trp	His	Thr	Leu	Cys	Ala	Glu	Gly	Ala	Asp	Gln	Phe	Gly	Gly	Gly
	50					55					60				

Thr	Lys	Ser	Phe	Pro	Trp	Asn	Glu	Gly	Thr	Asp	Ala	Ile	Glu	Ile	Ala
65					70					75					80

Lys	Gln	Lys	Val	Asp	Ala	Gly	Phe	Glu	Ile	Met	Gln	Lys	Leu	Gly	Ile
				85					90					95	

Pro	Tyr	Tyr	Cys	Phe	His	Asp	Val	Asp	Leu	Val	Ser	Glu	Gly	Asn	Ser
			100					105					110		

Ile	Glu	Glu	Tyr	Glu	Ser	Asn	Leu	Lys	Ala	Val	Val	Ala	Tyr	Leu	Lys
		115					120					125			

Glu	Lys	Gln	Lys	Glu	Thr	Gly	Ile	Lys	Leu	Leu	Trp	Ser	Thr	Ala	Asn
	130					135					140				

Val	Phe	Gly	His	Lys	Arg	Tyr	Met	Asn	Gly	Ala	Ser	Thr	Asn	Pro	Asp
145					150					155					160

Phe	Asp	Val	Val	Ala	Arg	Ala	Ile	Val	Gln	Ile	Lys	Asn	Ala	Ile	Asp
				165					170					175	

Ala	Gly	Ile	Glu	Leu	Gly	Ala	Glu	Asn	Tyr	Val	Phe	Trp	Gly	Gly	Arg
			180					185					190		

Glu	Gly	Tyr	Met	Ser	Leu	Leu	Asn	Thr	Asp	Gln	Lys	Arg	Glu	Lys	Glu
		195					200					205			

60

24

	His	Met	Ala	Thr	Met	Leu	Thr	Met	Ala	Arg	Asp	Tyr	Ala	Arg	Ser	Lys	
	210						215					220					
5	Gly	Phe	Lys	Gly	Thr	Phe	Leu	Ile	Glu	Pro	Lys	Pro	Met	Glu	Pro	Thr	
	225					230					235					240	
	Lys	His	Gln	Tyr	Asp	Val	Asp	Thr	Glu	Thr	Ala	Ile	Gly	Phe	Leu	Lys	
					245					250					255		
10	Ala	His	Asn	Leu	Asp	Lys	Asp	Phe	Lys	Val	Asn	Ile	Glu	Val	Asn	His	
				260					265					270			
	Ala	Thr	Leu	Ala	Gly	His	Thr	Phe	Glu	His	Glu	Leu	Ala	Cys	Ala	Val	
				275				280					285				
15	Asp	Ala	Gly	Met	Leu	Gly	Ser	Ile	Asp	Ala	Asn	Arg	Gly	Asp	Tyr	Gln	
	290						295					300					
	Asn	Gly	Trp	Asp	Thr	Asp	Gln	Phe	Pro	Ile	Asp	Gln	Tyr	Glu	Leu	Val	
20	305					310					315					320	
	Gln	Ala	Trp	Met	Glu	Ile	Ile	Arg	Gly	Gly	Gly	Phe	Val	Thr	Gly	Gly	
					325					330					335		
25	Thr	Asn	Phe	Asp	Ala	Lys	Thr	Arg	Arg	Asn	Ser	Thr	Asp	Leu	Glu	Asp	
				340					345					350			
	Ile	Ile	Ile	Ala	His	Val	Ser	Gly	Met	Asp	Ala	Met	Ala	Arg	Ala	Leu	
			355					360				365					
30	Glu	Asn	Ala	Ala	Lys	Leu	Leu	Gln	Glu	Ser	Pro	Tyr	Thr	Lys	Met	Lys	
	370						375					380					
	Lys	Glu	Arg	Tyr	Ala	Ser	Phe	Asp	Ser	Gly	Ile	Gly	Lys	Asp	Phe	Glu	
35	385					390					395					400	
	Asp	Gly	Lys	Leu	Thr	Leu	Glu	Gln	Val	Tyr	Glu	Tyr	Gly	Lys	Lys	Asn	
					405					410					415		
40	Gly	Glu	Pro	Lys	Gln	Thr	Ser	Gly	Lys	Gln	Glu	Leu	Tyr	Glu	Ala	Ile	
				420					425					430			
	Val	Ala	Met	Tyr	Gln												
				435													
45																	
	<210>	2															
	<211>	1669															
	<212>	DNA															
50	<213>	Piromyces sp.															
	<400>	2															
	gtaa	aatggct	aagga	aatatt	tccc	acaaat	tcaaa	agatt	aagtt	cgaag	gtaag	gattc	60				
	taaga	atcca	ttagc	cttcc	actac	tacga	tgctg	aaaag	gaagt	catgg	gtaag	aaaat	120				
55	gaagg	attgg	ttacg	tttcc	ccatg	gcctg	gtggc	acact	ctttg	cgccg	aaggt	gctga	180				
	ccaat	tcggt	ggagg	tacaa	agtct	tttccc	atgga	acgaa	ggtac	tgatg	ctatt	gaaat	240				
	tgcca	agcaa	aaggt	tgatg	ctgg	tttcga	aatca	tgcaa	aagct	tggta	ttcca	tacta	300				
	ctgtt	tccac	gatgt	tgatc	ttgtt	tccga	aggta	actct	attga	agaat	acga	atccaa	360				
	cctta	aggct	gtcgt	tgtct	acct	caagga	aaagc	aaaag	gaaac	cggtg	ttaag	cttct	420				
60	ctgg	agtact	gcta	acgtct	tcggt	cacaa	gcgtt	acatg	aacgg	tgcct	ccact	aacc	480				
	agact	ttgat	gttg	tcgccc	gtgct	attgt	tcaa	attaag	aacgc	catag	acgc	cggtat	540				

25

tgaacttggt gctgaaaact acgtcttctg ggggtggcgt gaagggttaca tgagtctcct 600
 taacactgac caaaagcgtg aaaaggaaca catggccact atgcttacca tggctcgtga 660
 ctacgctcgt tccaagggat tcaagggtac tttcctcatt gaaccaaagc caatggaacc 720
 5 aaccaagcac caatacgatg ttgacactga aaccgctatt ggtttcctta aggcccacaa 780
 cttagacaag gacttcaagg tcaacattga agttaaccac gctactcttg ctgggtcacac 840
 tttcgaacac gaacttgcct gtgctgttga tgctgggtatg ctcgggttcca ttgatgctaa 900
 ccgtgggtgac taccaaaacg gttgggatac tgatcaattc ccaattgatc aatacgaact 960
 cgtccaagct tggatggaaa tcatccgtgg tgggtggtttc gttactgggtg gtaccaactt 1020
 cgatgccaaag actcgtcgtg actctactga cctcgaagac atcatcattg cccacgtttc 1080
 10 tgggtatggat gctatggctc gtgctcttga aaacgctgcc aagctcctcc aagaatctcc 1140
 atacaccaag atgaagaagg aacgttacgc ttccttcgac agtgggtattg gtaaggactt 1200
 tgaagatggt aagctcacc cgaacaagt ttacgaatac ggtaagaaga acggtgaacc 1260
 aaagcaaact tctggtaagc aagaactcta cgaagctatt gttgccatgt accaataagt 1320
 taatcgtagt taaattggta aaataattgt aaaatcaata aacttgtcaa tcctccaatc 1380
 15 aagtttataaa gatcctatct ctgtactaat taaatatagt acaaaaaaaaa atgtataaac 1440
 aaaaaaaaaagt ctaaaagacg gaagaattta atttagggaa aaaataaaaa taataataaa 1500
 caatagataa atcctttata ttaggaaaat gtcccattgt attattttca tttctactaa 1560
 aaaagaaagt aaataaaaca caagaggaaa ttttcccttt tttttttttt tgtaataaat 1620
 tttatgcaaa tataaatata aataaaataa taaaaaaaaa aaaaaaaaaa 1669

20

<210> 3

<211> 494

<212> PRT

25 <213> *Piromyces* sp.

<400> 3

Met Lys Thr Val Ala Gly Ile Asp Leu Gly Thr Gln Ser Met Lys Val
 1 5 10 15

30

Val Ile Tyr Asp Tyr Glu Lys Lys Glu Ile Ile Glu Ser Ala Ser Cys
 20 25 30

35

Pro Met Glu Leu Ile Ser Glu Ser Asp Gly Thr Arg Glu Gln Thr Thr
 35 40 45

Glu Trp Phe Asp Lys Gly Leu Glu Val Cys Phe Gly Lys Leu Ser Ala
 50 55 60

40

Asp Asn Lys Lys Thr Ile Glu Ala Ile Gly Ile Ser Gly Gln Leu His
 65 70 75 80

Gly Phe Val Pro Leu Asp Ala Asn Gly Lys Ala Leu Tyr Asn Ile Lys
 85 90 95

45

Leu Trp Cys Asp Thr Ala Thr Val Glu Glu Cys Lys Ile Ile Thr Asp
 100 105 110

50

Ala Ala Gly Gly Asp Lys Ala Val Ile Asp Ala Leu Gly Asn Leu Met
 115 120 125

Leu Thr Gly Phe Thr Ala Pro Lys Ile Leu Trp Leu Lys Arg Asn Lys
 130 135 140

55

Pro Glu Ala Phe Ala Asn Leu Lys Tyr Ile Met Leu Pro His Asp Tyr
 145 150 155 160

Leu Asn Trp Lys Leu Thr Gly Asp Tyr Val Met Glu Tyr Gly Asp Ala
 165 170 175

60

Ser Gly Thr Ala Leu Phe Asp Ser Lys Asn Arg Cys Trp Ser Lys Lys

26

	180					185					190					
	Ile	Cys	Asp	Ile	Ile	Asp	Pro	Lys	Leu	Leu	Asp	Leu	Leu	Pro	Lys	Leu
			195					200					205			
5	Ile	Glu	Pro	Ser	Ala	Pro	Ala	Gly	Lys	Val	Asn	Asp	Glu	Ala	Ala	Lys
		210					215					220				
10	Ala	Tyr	Gly	Ile	Pro	Ala	Gly	Ile	Pro	Val	Ser	Ala	Gly	Gly	Gly	Asp
	225					230					235					240
	Asn	Met	Met	Gly	Ala	Val	Gly	Thr	Gly	Thr	Val	Ala	Asp	Gly	Phe	Leu
					245					250					255	
15	Thr	Met	Ser	Met	Gly	Thr	Ser	Gly	Thr	Leu	Tyr	Gly	Tyr	Ser	Asp	Lys
				260					265					270		
	Pro	Ile	Ser	Asp	Pro	Ala	Asn	Gly	Leu	Ser	Gly	Phe	Cys	Ser	Ser	Thr
			275					280					285			
20	Gly	Gly	Trp	Leu	Pro	Leu	Leu	Cys	Thr	Met	Asn	Cys	Thr	Val	Ala	Thr
		290					295					300				
25	Glu	Phe	Val	Arg	Asn	Leu	Phe	Gln	Met	Asp	Ile	Lys	Glu	Leu	Asn	Val
	305					310					315					320
	Glu	Ala	Ala	Lys	Ser	Pro	Cys	Gly	Ser	Glu	Gly	Val	Leu	Val	Ile	Pro
					325					330					335	
30	Phe	Phe	Asn	Gly	Glu	Arg	Thr	Pro	Asn	Leu	Pro	Asn	Gly	Arg	Ala	Ser
				340					345					350		
	Ile	Thr	Gly	Leu	Thr	Ser	Ala	Asn	Thr	Ser	Arg	Ala	Asn	Ile	Ala	Arg
			355					360					365			
35	Ala	Ser	Phe	Glu	Ser	Ala	Val	Phe	Ala	Met	Arg	Gly	Gly	Leu	Asp	Ala
		370					375					380				
40	Phe	Arg	Lys	Leu	Gly	Phe	Gln	Pro	Lys	Glu	Ile	Arg	Leu	Ile	Gly	Gly
	385					390					395					400
	Gly	Ser	Lys	Ser	Asp	Leu	Trp	Arg	Gln	Ile	Ala	Ala	Asp	Ile	Met	Asn
					405					410					415	
45	Leu	Pro	Ile	Arg	Val	Pro	Leu	Leu	Glu	Glu	Ala	Ala	Ala	Leu	Gly	Gly
				420					425					430		
	Ala	Val	Gln	Ala	Leu	Trp	Cys	Leu	Lys	Asn	Gln	Ser	Gly	Lys	Cys	Asp
			435					440					445			
50	Ile	Val	Glu	Leu	Cys	Lys	Glu	His	Ile	Lys	Ile	Asp	Glu	Ser	Lys	Asn
		450					455					460				
55	Ala	Asn	Pro	Ile	Ala	Glu	Asn	Val	Ala	Val	Tyr	Asp	Lys	Ala	Tyr	Asp
	465					470					475					480
	Glu	Tyr	Cys	Lys	Val	Val	Asn	Thr	Leu	Ser	Pro	Leu	Tyr	Ala		
					485					490						

60

<210> 4

<211> 2041

<212> DNA

<213> *Piromyces* sp.

5 <400> 4
attatataaa ataactttaa ataaaacaat ttttatttgt ttattttaatt attcaaaaaa 60
aattaaagta aaagaaaaat aatacagtag aacaatagta ataatatcaa aatgaagact 120
gttgctggta ttgatcttgg aactcaaagt atgaaagtcg ttattttacga ctatgaaaag 180
aaagaaatta ttgaaagtgc tagctgtcca atggaattga tttccgaaag tgacgggtacc 240
10 cgtgaacaaa ccactgaatg gtttgacaag ggtcttgaag ttgttttgg taagcttagt 300
gctgataaca aaaagactat tgaagctatt ggtatttctg gtcaattaca cggttttggt 360
cctcttgatg ctaacggtaa ggctttatac aacatcaaac ttgggtgtga tactgctacc 420
gttgaagaat gtaagattat cactgatgct gccgggtggg acaaggctgt tattgatgcc 480
cttggttaacc ttatgctcac cggtttccacc gctccaaaga tcctctggct caagcgcaac 540
15 aagccagaag ctttcgctaa cttaaagtac attatgcttc cacacgatta cttaaactgg 600
aagcttactg gtgattacgt tatggaatac ggtgatgcct ctgggtaccgc tctcttcgat 660
tctaagaacc gttgctgggc taagaagatt tgcgatatca ttgacccaaa acttttagat 720
ttacttccaa agttaattga accaagcgct ccagctggta aggttaatga tgaagccgct 780
aaggcttacg gtattccagc cgggtattcca gtttccgctg gtgggtggta taacatgatg 840
20 ggtgctggtg gtactgggtac tgttgctgat ggtttcctta ccatgtctat ggggtacttct 900
ggtactcttt acggttacag tgacaagcca attagtacc cagctaatgg tttaagtgg 960
ttctgttctt ctactgggtg atggcttcca ttactttgta ctatgaactg tactgttgcc 1020
actgaattcg ttcgtaacct cttccaaatg gatattaagg aacttaatgt tgaagctgcc 1080
aagtctccat gtggtagtga aggtgtttta gttattccat tcttcaatgg tgaagaact 1140
25 ccaaacttac caaacggctg tgctagtatt actggtctta cttctgctaa caccagccgt 1200
gctaacattg ctgctgctag tttcgaatcc gccgttttcg ctatgcgtgg tggtttagat 1260
gctttccgta agttagggtt ccaaccaaag gaaattcgtc ttattgggtg tggttctaag 1320
tctgatctct ggagacaaat tgccgctgat atcatgaacc ttccaatcag agttccactt 1380
ttagaagaag ctgctgctct tgggtgggtg gttcaagctt tatgggtgtc taagaaccaa 1440
30 tctggtaagt gtgatattgt tgaactttgc aaagaacaca ttaagattga tgaatctaag 1500
aatgctaacc caattgccga aaatgttgct gtttacgaca aggcttacga tgaatactgc 1560
aaggttgtaa atactctttc tccattatat gcttaaattg ccaatgtaaa aaaaaatata 1620
atgccatata attgccttgt caatacactg ttcattgttca tataatcata ggacattgaa 1680
tttacaaggt ttatacaatt aatatctatt atcatattat tatacagcat ttcatttttct 1740
35 aagattagac gaaacaattc ttgggttcctt gcaatataca aaatttacat gaatttttag 1800
aatagtctcg tatttatgcc caataatcag gaaaattacc taatgctgga ttcttggtta 1860
taaaaaacaaa ataaataaat taaataaaca aataaaaatt ataagtaa ataaatatat 1920
aagtaatatata aaaaaaaagt aaataaataa ataaataaat aaaaattttt tgcaaatata 1980
taaataaata aataaaatat aaaaataatt tagcaataa attaaaaaaa aaaaaaaaaa 2040
40 a 2041

13-04-2004

NL0300049

EPO - DG 1

PCT/NL03/00049

1

13.04.2004

Amended Claims

94

1. A eukaryotic host cell transformed with a nucleic acid construct comprising a nucleotide sequence encoding a xylose isomerase comprising an amino acid sequence
5 that has at least 70 % sequence identity with the amino acid sequence of SEQ ID NO: 1, whereby the nucleic acid construct, upon transformation of the host cell, confers to the host cell the ability to grow on xylose as carbon source.
2. A transformed host cell according to claim 1, wherein the host cell is a yeast,
10 preferably a yeast that belongs to one of the genera: *Saccharomyces*, *Kluyveromyces*, *Candida*, *Pichia*, *Schizosaccharomyces*, *Hansenula*, *Kloeckera*, *Schwanniomyces*, and *Yarrowia*.
3. A transformed host cell according to claim 2, wherein the yeast belongs to one of
15 the species: *S. cerevisiae*, *S. bulderi*, *S. barnetti*, *S. exiguus*, *S. uvarum*, *S. diastaticus*, *K. lactis*, *K. marxianus*, and *K. fragilis*.
4. A transformed host cell according to claim 1, wherein the host cell is a filamentous fungus, preferably a filamentous fungus that belongs to one of the genera:
20 *Aspergillus*, *Trichoderma*, *Hemicola*, *Acremonium*, *Fusarium*, and *Penicillium*.
5. A transformed host cell according to any one of the preceding claims, whereby the nucleotide sequence encoding a xylose isomerase is operably linked to a promoter that causes sufficient expression of the xylose isomerase in the host cell, to confer to
25 the host cell the ability to isomerise xylose into xylulose.
6. A transformed host cell according to claim 6, whereby the promoter is insensitive to catabolite repression in the host cell.
- 30 7. A transformed host cell according to any one of the preceding claims, whereby the host cell comprises a genetic modification that result in a characteristic selected from the group consisting of:
(a) increase transport of xylose into the host cell;

13-04-2004

NL0300049

2

- (b) increased xylulose kinase activity;
(c) increased flux of the pentose phosphate pathway;
(d) decreased sensitivity to catabolite repression;
(e) increased tolerance to ethanol, osmolarity or organic acids; and,
5 (f) reduced production of by-products.

8. A transformed host cell according to claim 7, wherein the genetic modification consist of overexpression of endogenous genes, expression of a heterologous genes, or a combination thereof, and whereby the gene is selected from the group consisting of a
10 gene encoding: a hexose or pentose transporter, an xylulose kinase; an enzyme from the pentose phosphate pathway, a glycolytic enzyme, and an ethanologenic enzymes.

9. A transformed host cell according to claim 7, wherein the genetic modification consist of the inactivation of an endogenous genes, whereby the gene is selected from
15 the group consisting of a gene encoding a hexose kinase gene, the *Saccharomyces MIG1* and *MIG2* genes and hybridising homologues thereof.

10. A transformed host cell according to any one of the preceding claims, whereby the host cell expresses one or more enzymes that confer to the host cell the ability to
20 produce lactic acid, acetic acid, succinic acid, amino acids, 1,3-propane-diol, ethylene, glycerol, β -lactam antibiotics and cephalosporins.

11. A transformed host cell according to claim 10, whereby the host cell contains a genetic modification that results in decreased alcohol dehydrogenase activity.
25

12. A process for producing ethanol, whereby the process comprises the steps of:
(a) fermenting a medium containing a source of xylose with a transformed host cell as defined in any one of claims 1 - 9, whereby the host cell ferments xylose to ethanol,
30 and optionally,
(b) recovery of the ethanol.

13-04-2004

NL0300049

3

13. A process according to claim 12, whereby the medium also contains a source of glucose.

14. A process according to claims 12 or 13, whereby the volumetric ethanol
5 productivity is at least 0.5 g ethanol per litre per hour.

15. A process according to claims any one of claims 12 - 14, whereby the ethanol yield is at least 50 %.

10 16. A process for producing a fermentation product selected from the group consisting of lactic acid, acetic acid, succinic acid, amino acids, 1,3-propane-diol, ethylene, glycerol, β -lactam antibiotics and cephalosporins, whereby the process comprises the steps of:

- 15 (a) fermenting a medium containing a source of xylose with a transformed host cell as defined in claims 10 or 11, whereby the host cell ferments xylose to the fermentation product, and optionally,
(b) recovery of the fermentation product.

20 17. A process according to claim 16, whereby the medium also contains a source of glucose.

1/1

Fig 1