



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2128215 C 2005/05/03

(11)(21) **2 128 215**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1993/01/15
(87) Date publication PCT/PCT Publication Date: 1993/07/22
(45) Date de délivrance/Issue Date: 2005/05/03
(85) Entrée phase nationale/National Entry: 1994/07/15
(86) N° demande PCT/PCT Application No.: US 1993/000181
(87) N° publication PCT/PCT Publication No.: 1993/014202
(30) Priorité/Priority: 1992/01/17 (07/822,043) US

(51) Cl.Int.⁵/Int.Cl.⁵ C12N 15/12, A61K 39/395, G01N 33/574,
C07K 14/47, C07K 17/00, C07K 1/36, C07K 7/08,
C07K 7/06

(72) Inventeurs/Inventors:
STRACKE, MARY, US;
LIOTTA, LANCE A., US;
SCHIFFMANN, ELLIOTT, US;
KRUTZSCH, HENRY, US

(73) Propriétaire/Owner:
THE UNITED STATES OF AMERICA REPRESENTED
BY THE SECRETARY, DEPARTMENT OF HEALTH
AND HUMAN SERVICES, US

(74) Agent: SMART & BIGGAR

(54) Titre : L'AUTOTAXINE, PROTEINE STIMULANT LA MOTILITE QUI EST UTILE DANS LE DIAGNOSTIC ET LE
TRAITEMENT DU CANCER

(54) Title: AUTOTAXIN: MOTILITY STIMULATING PROTEIN USEFUL IN CANCER DIAGNOSIS AND THERAPY

(57) **Abrégé/Abstract:**

The present invention relates, in general, to autotaxin. In particular, the present invention relates to a DNA segment encoding autotaxin; recombinant DNA molecules containing the DNA segment; cells containing the recombinant DNA molecule; a method of producing autotaxin; and antibodies to autotaxin.






INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/12, C07K 7/00, 15/00 C07K 15/28, A61K 39/395 G01N 33/574, C12N 1/21	A1	(11) International Publication Number: WO 93/14202 (43) International Publication Date: 22 July 1993 (22.07.93) 2128215
(21) International Application Number: PCT/US93/00181 (22) International Filing Date: 15 January 1993 (15.01.93) (30) Priority data: 07/822,043 17 January 1992 (17.01.92) US (71) Applicant: THE UNITED STATES OF AMERICA, represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; National Institutes of Health, Office of Technology Transfer, Box OTT, Bethesda, MD 20892 (US). (72) Inventors: STRACKE, Mary ; 14414 Barkwood Drive, Rockville, MD 20853 (US). LIOTTA, Lance, A. ; 9027 Mitswood Drive, Potomac, MD 20854 (US). SCHIFFMANN, Elliott ; 3207 Pickwick Lane, Chevy Chase, MD 20815 (US). KRUTZSCH, Henry ; 9704 DePaul Drive, Bethesda, MD 20817 (US).		(74) Agents: SCOTT, Watson, T. et al.; Cushman, Darby & Cushman, 1100 New York Avenue, N.W., Washington, DC 20005 (US). (81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: AUTOTAXIN: MOTILITY STIMULATING PROTEIN USEFUL IN CANCER DIAGNOSIS AND THERAPY

MOTILITY (density units)



(57) Abstract

The present invention relates, in general, to autotaxin. In particular, the present invention relates to a DNA segment encoding autotaxin; recombinant DNA molecules containing the DNA segment; cells containing the recombinant DNA molecule; a method of producing autotaxin; and antibodies to autotaxin.

AUTOTAXIN: MOTILITY STIMULATING PROTEIN USEFUL
IN CANCER DIAGNOSIS AND THERAPY

BACKGROUND OF THE INVENTION

Field of the Invention

5 The present invention relates, in
general, to a motility stimulating peptide and
compositions comprising the same. In particular, the
present invention relates to a purified form of the
peptide, for example, autotaxin (ATX); a DNA segment
10 encoding autotaxin; recombinant DNA molecules
containing the DNA segment; cells containing the
recombinant DNA molecule; a method of producing
autotaxin; antibodies to autotaxin; and methods of
cancer diagnosis and therapy using the above
15 referenced peptides and DNA segments.

Background Information

Cell motility plays an important role in
embryonic events, adult tissue remodeling, wound
healing, angiogenesis, immune defense, and
20 metastasis of tumor cells (Singer, S.J. and Kupfer,
A. (1986) Ann. Rev. Cell Biol. 2, 337-365). In
normal physiologic processes, motility is tightly
regulated. On the other hand, tumor cell motility
may be aberrantly regulated or autoregulated. Tumor
25 cells can respond in a motile fashion to a variety
of agents. These include host-derived factors such
as scatter factor (Rosen, E.M. et al., (1989) In
Vitro Cell Devel. Biol. 25, 163-173) and growth
factors (Kahan, B.W. et al., (1987) Cancer Res. 47,
30 6324-6328; Stracke, M.L. et al., Biochem. Biophys.
Res. Comm. 153, 1076-1083; Tamm, I., et al., (1989)

2128215

2

J. Exp. Med. 170, 1649-1669; Wang, J.M., et al. (1990) Biochem. Biophys. Res. Comm. 169, 165-170; and Jouanneau, J., et al. (1991) Proc. Natl. Acad. Sci. USA 88, 2893-2897), components of the

5 extracellular matrix (McCarthy, J.B., et al. (1984) J. Cell Biol. 98, 1474-1480), and tumor-secreted or autocrine factors (Liotta, L.A., et al. (1988) Cancer Surveys 7, 631-652; Ruff, M., et al. (1985) Clin. Immunol. Immunopath. 37, 387-396; Atnip, K.D.,

10 et al. (1987) Biochem. Biophys. Res. Comm. 146, 996-1002; Ohnishi, T., et al. (1990) J. Neurosurg. 73, 881-888; Silletti, S., et al. (1991) Cancer Res. 51, 3507-3511.; and Watanabe, H., et al. (1991) J. Biol. Chem. 266, 13442-13448).

15 Many types of host-derived soluble factors act in a paracrine fashion to stimulate cell locomotion. Motility-stimulating proteins called "scatter factors" have been identified which are produced by embryonic fibroblasts and by smooth

20 muscle cells (Stoker, M., et al. (1987) Nature 327, 239-242). Scatter factors stimulate random and directed motility by epithelial cells, keratinocytes, vascular endothelial cells and carcinoma cells (Stoker, M., et al. (1987) Nature

25 327, 239-242; Rosen, E.M., et al. (1990) Proc. Soc. Exp. Biol. Med. 195, 34-43; and Weidner, K.M., et al. (1990) J. Cell. Biol. 111, 2097-2108), but not fibroblasts. In addition, a number of host-secreted growth factors have been demonstrated to stimulate

30 motility in tumor cells, including nerve growth factor (Kahan, B.W., et al. (1987) Cancer Res. 47, 6324-6328) insulin-like growth factor-I (Stracke, M.L., et al. (1988) Biochem. Biophys. Res. Comm. 153, 1076-1083), interleukin-6 (Tamm, I., et al.

35 (1989) J. Exp. Med. 170, 1649-1669), interleukin-8

2128215

3

(Wang, J.M., et al. (1990) Biochem. Biophys. Res. Comm. 169, 165-170), and acidic fibroblast growth factor (Jouanneau, J., et al. (1991) Proc. Natl. Acad. Sci. USA 88, 2893-2897). These paracrine
5 factors may influence "homing" or the directionality of tumor cell motility.

In contrast to these host-derived factors, many types of tumor cells have been found to produce proteins termed "autocrine motility factors" which
10 stimulate motility by the same tumor cells which make the factor (Liotta, L.A., et al. (1986) Proc. Natl. Acad. Sci. USA 83, 3302-3306). Autocrine motility factors are not specific for a given type of cancer cell but have a wide spectrum of activity
15 on many types of cancer cells (Kohn, E.C., et al. (1990) Int. J. Cancer 46, 287-292), with little effect on normal fibroblasts or leukocytes.

Autocrine motility factors identified to date act through cell-surface receptors (Stracke, M.L., et al. (1987) Biochem. Biophys. Res. Comm.
20 147, 339-345; Nabi, I.R., et al. (1990) Cancer Res. 50, 409-414; Watanabe, H., et al. (1991) J. Biol. Chem. 266, 13442-13448) resulting in pseudopodial protrusion (Guirguis, R., et al. (1987) Nature 329,
25 261-263) leading to both random and directed migration (Liotta, L.A., et al. (1986) Proc. Natl. Acad. Sci. USA 83, 3302-3306; Atnip, K.D., et al. (1987) Biochem. Biophys. Res. Comm. 146, 996-1002; Ohnishi, T., et al. (1990) J. Neurosurg. 73, 881-
30 888).

Prior studies of human A2058 melanoma cells have demonstrated that these cells are a particularly rich source of autocrine motility factors. An autocrine motility factor with a
35 molecular mass of approximately 60 kDa has been

2128215

4

previously isolated from the conditioned media of these cells. (Liotta, L.A., et al. (1986) Proc. Natl. Acad. Sci. USA 83, 3302-3306). Similar tumor cell derived or induced factors with the same

5 molecular weight have subsequently been reported and purified by several investigators (Atnip, K.D., et al. (1987) Biochem. Biophys. Res. Comm. 146, 996-1002; Schnor, S.L., et al. (1988) J. Cell Sci. 90, 391-399; Ohnishi, T., et al. (1990) J. Neurosurg.

10 73, 881-888; Siletti, S., et al. (1991) Cancer Res. 51, 3507-3511; Watanabe, H., et al. (1990) J. Cell Biol. 111, 2097-2108). Such factors are thought to play a key role in tumor cell invasion.

Most of the motility factors identified to

15 date have not been purified to homogeneity and have not been sequenced. The novel tumor motility factor of the present invention, named herein as autotaxin (ATX), has been purified and verified to be a

20 homogeneous sample by 2-D gel electrophoresis. The protein of the present invention is unique from any previously identified or purified motility factor. The molecular size of ATX is about 125kDa and it has an isoelectric point of approximately 7.7. ATX stimulates both random and directed migration of

25 human A2058 melanoma cells at pM concentrations. The activity of this factor (ATX) is completely sensitive to inhibition by pertussis toxin. No significant homology has been found to exist between the protein of the invention and any mammalian

30 protein including previous factors known to stimulate cell motility.

There is a great clinical need to predict the aggressiveness of a patient's individual tumor, to predict the local recurrence of treated tumors

35 and to identify patients at high risk for

66597-197

5

development of invasive tumors. The present invention provides a functional marker which is functionally related to the invasive potential of human cancer. The invention further provides an assay for this secreted marker in body fluids, or in tissues. The assay of the invention can be used in the detection, diagnosis, and treatment of human malignancies and other inflammatory, fibrotic, infectious or healing disorders.

SUMMARY OF THE INVENTION

10 The present invention relates, generally, to a motility stimulating peptide and to a DNA segment encoding same.

 It is a specific object of the present invention to provide the motility stimulating peptide, hereinafter
15 referred to as autotaxin.

 Thus, according to one aspect of the present invention, there is provided an isolated polypeptide selected from human autotaxin protein in homogeneous form or a fragment of said protein, wherein (A) said protein has an
20 isoelectric point of about 7.7, a molecular weight of about 125 kDa, as determined SDS-polyacrylamide gel electrophoresis under reducing conditions, and induces a motility in human A2508 melanoma cells which is inhibited by pertussis toxin; (B) said fragment comprises an amino acid
25 sequence selected from the group consisting of SEQ ID Nos. 1-11 and 26-33; and (C) antibody suitable for immunohistochemical staining of cancerous human tissue can be raised to each of said protein and said fragment.

 According to another aspect of the present
30 invention, there is provided an isolated polynucleotide coding for the polypeptide described above.

66597-197

5a

According to still another aspect of the present invention, there is provided a recombinant DNA molecule comprising the polynucleotide and a rector.

According to yet another aspect of the present invention, there is provided a cell comprising the recombinant DNA molecule.

According to a further aspect of the present invention, there is provided a method of producing a polypeptide as described above, comprising the steps of:

10 (i) culturing a cell comprising a polynucleotide coding for said polypeptide and a vector under conditions such that said polynucleotide is expressed, and said polypeptide is thereby produced, and (ii) isolating said polypeptide.

The method of the present invention further comprises the steps of (i) collecting and concentrating supernatant from said cultured cell, whereby a first preparation of polypeptide is produced; (ii) salt fractionating said first preparation to produce a second preparation of polypeptide; (iii) isolating said peptide

20 from said second polypeptide preparation, so that said peptide is obtained in substantially pure form.

Further objects and advantages of the present invention will be clear from the description that follows.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Fractionation of ATX by hydrophobic interaction. A 200 ml sample of A2058 conditioned media was chromatographed on a 200 mL column of phenyl Sepharose-4B*. Buffer A was 50 mM Tris (pH 7.5), 5% methanol, and 1.2 M ammonium sulfate. Buffer B was 50 mM Tris (pH 7.5), 5% methanol and 50% ethylene glycol. The gradient (----) represents a double linear gradient with decreasing ammonium sulfate (1.2 to 0.0 M) and increasing ethylene glycol (0 to 50%). Absorbance was monitored at 280 nm (——) and indicated that most of the protein did not bind to the column. Ten ml fractions were assayed for motility stimulating capacity using the Boyden Chamber assay (o). The peak of motility activity occurred between 900 and 1050 minutes, ~ 12% of the gradient.

Figure 2. Isolation of ATX by lectin affinity chromatography. 20 ml portions of the phenyl Sepharose activity peak were affinity purified on a 40 ml Concanavalin A Affi-Gel* column. The bound components were eluted with a step gradient (----) of methyl α -D-mannopyranoside (0.0 mM, 10 mM, and 500 mM) in a buffer consisting of 0.05 M Tris (pH 7.5), 0.1 M NaCl, 0.01 M CaCl₂ and 20% ethylene glycol. Absorbance was monitored at 280 nm (_____) and indicated that the majority of the protein components did not bind to the column. Motility was assayed in 10 mL fractions (...o...) and was found predominantly in the 500 mM elution concentration. One of seven chromatographic runs is shown.

*Trade-mark

Figure 3. Purification of ATX by weak anionic exchange chromatography. Approximately 30% of the activity peak eluted from the Con A affinity column was applied to a ZORBAX BioSeries-WAX column. The bound components were eluted with an NaCl gradient (----) in a buffer consisting of 10 mM Tris (pH 7.5) and 30% ethylene glycol. Motility (o) was assayed in 1.0 ml fractions. The peak of activity eluted in a discrete but broad region in the shallow portion of the gradient. Absorbance was monitored at 230 nm (_____). The majority of the protein components not associated with activity bound strongly to the column were eluted at 1.0 M NaCl. One of two chromatographic runs is shown.

Figure 4. Purification of ATX by molecular sieve exclusion chromatography. The entire activity peak eluted from the weak anion exchange column was applied to a series of TSK* columns (4000SW, 4000SW, 3000SW, and 2000SW, in this order). Proteins were eluted in a buffer consisting of 0.1M NaPO₄ (pH 7.2) with 10% methanol and 10% ethylene glycol. Two major protein peaks were evident by monitoring the absorbance at 235 nm (_____). Motility (...o...) was assayed in 0.4 ml samples and found predominantly in the first, smaller, protein peak.

Figure 5. Final purification of ATX by strong anionic exchange chromatography. Approximately 15% of the activity peak from the molecular sieve exclusion series was applied to a Pro-Pac* PA1 column. Protein which bound to the column was eluted with a NaCl gradient (----) in a buffer consisting of 10 mM Tris (pH 7.5), 5% methanol and 20% ethylene glycol. Absorbance was

*Trade-mark

66597-197

8

monitored at 215 nM (____). Motility activity was assayed in 1.0 ml fractions at two different dilutions: 1/5 (...o....) or 1/15 (.__o.__). Activity was found to correspond to a double protein peak in the central region of the gradient.

Figures 6A, 6B and 6C. Protein components associated with the activity peaks from various stages of purification. The activity peak from each chromatographic fractionation was pooled, concentrated and analyzed by SDS-polyacrylamide gel electrophoresis. Molecular weight standards are in Lane 1 for each panel. FIG. 6A) 8-16% gradient gel of the first three purification steps, run under non-reducing conditions. Lane 2 is an aliquot of the pooled activity peak eluted from the phenyl sepharose fractionation. Lane 3 is an aliquot of the pooled activity peak eluted from the Con A affinity purification. Lanes 4 and 5 show the "peak" and "shoulder" of activity fractionated by weak anion exchange chromatography (FIG. 3). FIG. 6B) 7% gel of the activity peak fractionated by molecular sieve exclusion chromatography. Lanes 2 and 3 show the protein separation pattern of the total pooled activity peak when the gel was run under non-reducing and reducing conditions, respectively. FIG. 6C) 8-16% gradient gel of the final strong anionic exchange chromatographic separation, run under non-reducing conditions. Lane 2 comprises ~1% of the total pooled activity peak eluted from the column.

Figure 7. 2-dimensional gel electrophoresis of ATX. Purified ATX (FIG. 6, Panel C) was subjected to non-equilibrium isoelectric

focusing (5 hr. at 500v), then applied to a 7.5% SDS-polyacrylamide gel for the second dimension. The pH separation which resulted was measured in 0.5 cm samples of concurrently run tube gels and is shown at the top. Molecular weight standards for the second dimension are shown on the right. This analysis reveals a single component with $pI = 7.7 \pm 0.2$ and $M_r = 120,000$.

Figure 8. Dilution curve of ATX. Purified ATX (FIG. 6, Panel C) was serially diluted and tested for motility-stimulating activity. The result, with unstimulated background motility subtracted out, shows that activity is half-maximal at ~ 500 pM ATX.

Figure 9. Pertussis toxin (PT) sensitivity of ATX. A2058 cells were pre-treated for 1 hr. prior to the start of the motility assay with 0.5 μ g/ml PT in 0.1% BSA-DMEM or with 0.1% BSA-DMEM alone (for untreated control). The motility activity stimulated by purified ATX (FIG. 6, Panel C) was then assessed for the two treatment groups. The result, expressed as cells/high power fields (HPF) $\pm$ S.E.M. with unstimulated background motility subtracted out, reveals profound inhibition of PT-treated cells (hatched) compared to untreated cells (solid). PT had no effect on cell viability. S.E.M.'s were < 10%.

Figure 10. Checkerboard analysis of ATX-stimulated motility using a 48-well microchemotaxis chamber. Varying dilutions of autotaxin were added to the upper chamber with the cells and/or to the lower chamber, as shown. Motility

66597-197

10

response, expressed as cells/HPF $\pm$ S.E.M., was assessed for each point in the checkerboard.

Figure 11. Purification of ATX peptides on HPLC. ATX, purified to homogeneity by strong anionic exchange chromatography, was sequentially digested by cyanogen bromide, subjected to reduction and pyridylethylation, and digested by trypsin. The resulting peptides were purified on an Aquapore* RP300 C-8 reverse phase column using a (0-70)% acetonitrile gradient in 0.1% trifluoroacetic acid (----). The absorbance was monitored at 215 nm (____) and peaks were collected. Seven peaks, chosen at random for N-terminal amino acid sequence analysis, are shown with appropriate numbers.

15 DETAILED DESCRIPTION OF THE INVENTION

Autotaxin is secreted by A2058 human melanoma cells cultured in low abundance in serum-free conditioned medium. ATX is believed to be a glycosylated protein due to its high affinity for con A and amino acid sequence analysis of the ATX peptides. The molecular size of ATX is about 125 kDa as determined by SDS polyacrylamide gel electrophoresis under reducing conditions and it has an isoelectric point of 7.7 ± 0.2 . These characteristics distinguish ATX from several small growth factors and interleukins which are implicated in tumor cell motility (Stracke, M.L., et al. (1988) Biochem. Biophys. Res. Comm. 153, 1076-1083; Ruff, M., et al. (1985) Clin. Immunol. Immunopath. 37, 387-396; Maciag, T., et al. (1984) Sci. 225, 932-935; Gospodarowicz, D. (1984) Proc. Natl. Acad. Sci. USA 81, 6963-6967; Van Snick, J. (1990) Ann. Rev.

*Trade-mark

Immunol. 8, 253-278; Yoshimura, T. (1987) Proc. Natl. Acad. Sci. USA 84, 9233-9237).

The protein of the present invention, which in one embodiment is derived from A2058 human melanoma cells, can be prepared substantially free from proteins with which it is normally associated using, for example, the purification protocol disclosed herein. That protocol can be generally described as follows.

10 A large volume of serum-free conditioned medium from appropriate producer cells (e.g., tumor cells) is collected and concentrated approximately 500-fold. This concentrated conditioned medium is then separated from other contaminating proteins by
15 techniques that rely on the chemical and physical characteristics of the protein. These include the molecular weight, relative hydrophobicity, net charge, isoelectric focusing point, and the presence of lectin-binding sugar residues on the protein.

20 Alternatively, the protein, or functional portion thereof, can be synthesized using chemical or recombinant means.

The protein of the present invention has a potent biological activity. The purified ATX is
25 active in the picomolar range and 1 unit of activity corresponds to a concentration of approximately 500 pM as assessed by the cell motility assay described herein and elsewhere (Stracke, M.L., et al. (1989) J. Biol. Chem. 264, 21544-49).

30 The protein of the present invention has a molecular size, as determined by two dimensional gel electrophoresis, of from 120 to 130 kDa, or more specifically, about 125 kDa. Further the protein of the present invention can have a pI in the range of
35 7.5 to 7.9, preferably, approximately 7.7.

2128215

12

The present invention also relates to a DNA segment coding for a polypeptide comprising an amino acid sequence corresponding to ATX, or a unique portion of such a sequence (unique portion being defined herein as at least 5, 10, 25, or 50 amino acids). In one embodiment, the DNA segment encodes any one of the amino acid sequences shown in SEQ ID NO:1 to SEQ ID NO:11 and SEQ ID NO:26 to SEQ ID NO:33.

10 In another embodiment, the present invention relates to a recombinant DNA molecule comprising a vector (for example plasmid or viral vector) and a DNA segment coding for a polypeptide corresponding to ATX, as can be prepared by one skilled in the art. Preferably, the encoding segment is present in the vector operably linked to a promoter.

In a further embodiment, the present invention relates to a cell containing the above described recombinant DNA molecule. Suitable hosts cells include procaryotic cells (such as bacteria, including E. coli) and both lower eucaryotic cells (for example, yeast) and higher eucaryotic cells (for example, mammalian cells). Introduction of the recombinant molecule into the host cells can be effected using methods known in the art.

In another embodiment, the present invention relates to a method of producing a peptide having an amino acid sequence corresponding to ATX. The method comprises culturing the above described cell under conditions such that the DNA segment is expressed, and isolating ATX thereby produced.

In a further embodiment, the present invention relates to an antibody having affinity for autotaxin or peptide fragments thereof. The

2128215

13

invention also relates to binding fragments of such antibodies. In one preferred embodiment, the antibodies are specific for autotaxin peptides having an amino acid sequence set forth in one of
5 SEQ ID NO:1 through SEQ ID NO:11 and SEQ ID NO:26 through SEQ ID NO:33.

Antibodies can be raised to autotaxin or its fragment peptides, either naturally occurring or recombinantly produced, using methods known in the
10 art.

ATX peptide fragments described above can be joined to other materials, particularly polypeptides, as fused or covalently joined polypeptides to be used as carrier proteins. ATX
15 and its fragments can be fused or covalently linked to a variety of carrier proteins, such as keyhole limpet hemocyanin, bovine serum albumin, tetanus toxoid, etc. See for example, Microbiology, Hoeber Medical Division (Harper and Row, 1969),
20 Landsteiner, Specificity of Serological Reactions (Dover Publications, New York, 1962) and Williams et al., Methods in Immunology and Immunochemistry, Vol. 1 (Academic Press, New York, 1967), for descriptions of methods of preparing polyclonal antisera. A
25 typical method involves hyperimmunization of an animal with an antigen. The blood of the animal is then collected shortly after the repeated immunizations and the gamma globulin is isolated.

In some instances, it is desirable to
30 prepare monoclonal antibodies from various mammalian hosts. Description of techniques for preparing such monoclonal antibodies may be found in Stites et al., editors, Basic and Clinical Immunology, (Lange Medical Publications, Los Altos, CA, Fourth edition)
35 and references cited therein, and in particular in

Kohler and Milstein in Nature 256:495-497 (1975), which discusses one method of generating monoclonal antibodies.

5 In another embodiment, the present invention relates to an oligonucleotide probe synthesized according to the sense or antisense degenerative sequence set forth in one of SEQ ID NO:1 through SEQ ID NO:11 and SEQ ID NO:26 through SEQ ID NO:33.

10 No significant homologies to known proteins found upon comparing the sequences set forth in SEQ ID NO:1 through SEQ ID NO:11 and SEQ ID NO:26 through SEQ ID NO:33 with those in the following protein databases: GenBank* (68.0),
15 EMBL (27.0), SWISS-PROT (18.0), and GenPept* (64.3). Thus, autotaxin is a unique, previously undescribed protein.

In a further embodiment, the present invention relates to a method of diagnosing cancer
20 metastasis and to a kit suitable for use in such a method. Preferably, antibodies to ATX can be used in, but not limited to, ELISA, RIA or immunoblots configurations to detect the presence of ATX in body fluids of patients (e.g. serum, urine, pleural
25 effusions, etc.). These antibodies can also be used in immunostains of patient samples to detect the presence of ATX.

In yet another embodiment, the present invention relates to in vivo and in vitro
30 diagnostics. ATX may be radiolabelled, by means known to one skilled in the art, and injected in cancer patients with appropriate ancillary substances also known to one skilled in the art, in order to ultimately detect distant metastatic sites
35 by appropriate imagery. The level of ATX in tissue

*Trade-mark

66597-197

15

or body fluids can be used to predict disease outcomes and/or choice of therapy which may also include ATX inhibitors.

5 In a further embodiment, the present invention relates to a treatment of cancer. ATX antibodies can be cross-linked to toxins (e.g., Ricin A), by means known to one skilled in the art, wherein the cross-linked complex is administered to cancer patients with appropriate ancillary agents by
10 means known to one skilled in the art, so that when the antibody complex binds to the cancer cell, the cell is killed by the cross-linked toxin.

The present invention is described in further detail in the following non-limiting
15 examples.

EXAMPLES

The following protocols and experimental details are referenced in the Examples that follow:

Materials. The polycarbonate Nuclepore* membranes
20 and the 48-well microchemotaxis chambers were obtained from Neuro Probe, Inc. Pertussis toxin (PT), ethylene glycol (biotechnology grade), methyl α -D-mannopyranoside were obtained from commercial vendors. The ampholyte, pH 3-10 Bio-Lyte* and pH 8-
25 10 Bio-Lyte, were obtained from Bio-Rad. Phenyl Sepharose CL-4B; affi-Gel concanavalin A; ZORBAX BioSeries-WAX (weak anion exchange) column (9.4mm x 24cm); Spherogel-TSK* 4000SW, 3000SW and 2000SW columns (each 7.5mm x 30cm); the Pro-Pac PA1 (4 x
30 50mm) strong anion exchange column; the Aquapore RP300 C-8 reverse phase column (220 x 2.1mm); and

*Trade-mark

the AminoQuant* C-18 reverse phase column (200 x 2.1mm) were also obtained from commercial sources.

Cell Culture. The human melanoma cell line A2058, originally isolated by Todaro (Todaro, G.J., et al. (1980) Proc. Natl. Acad. Sci. USA 77, 5258-5262), was maintained as previously described by Liotta (Liotta, L.A., et al. (1986) Proc. Natl. Acad. Sci. USA 83, 3302-3306).

Production of Autotaxin. A2058 cells were grown up in T-150 flasks, trypsinized, and seeded into 24,000 cm² cell factories at a cell density of 1×10^{10} cells/factory. After 5-6 days, the serum-containing medium was removed and the cells were washed with DPBS. The factories were maintained in DMEM without phenol red, supplemented with 4 mM glutamine, 100 units/ml penicillin, 100 µg/ml streptomycin, 5 µg/ml crystallized bovine serum albumin, 10 µg/ml bovine insulin, and 1 µM aprotinin. Culture supernatants were harvested every 3 days, frozen at -40°C and replaced with fresh serum-free medium. Each cycle of supernatant was tested for ATX production with a cell motility assay detailed below. Typically, a cell factory continued to be productive for 9-11 of these cycles.

After accumulation of approximately 45-60 L of supernatant, the culture supernatants were thawed and concentrated down to 2-2.5 L using an Amicon S10Y30 spiral membrane ultrafiltration cartridge. This supernatant was further concentrated in an Amicon high performance ultrafiltration cell using Diaflo* membranes. The final volume achieved from 100-200 L of conditioned

*Trade-mark

66597-197

17

medium was typically 250-400 ml. All ultrafiltrations were performed at 4°C.

Cell Motility Assays. Purification of autotaxin was monitored by testing the motility-stimulating capacity of the fractions collected from the columns. These fractions were in buffers unsuitable for a chemotaxis assay so each fraction had to be washed into an appropriate buffer, i.e., 0.1% (w/v) BSA in DPBS containing calcium and magnesium. This dialysis was performed by adding aliquots of each fraction to be tested into Centricon-30* ultrafiltration tubes, which retain molecular species larger than 30,000 daltons.

The assay to determine motility was performed in triplicate using a 48-well microchemotaxis chamber as described elsewhere in detail (Stracke, M.L., et al. (1987) Biochem. Biophys. Res. Comm. 146, 339-345; Stracke, et al. (1989) J. Biol. Chem. 264, 21544-21549). The Nuclepore membranes used in these modified Boyden chambers were fixed and stained with Diff-Quik*. Chemotaxis was quantitated either by reading the stained membranes with a 2202 Ultrosan* laser densitometer or by counting 5 randomly chosen high power fields (HPF) under light microscopy (400 x) for each replicate. Densitometer units (wavelength - 633 nm) have been shown to be linearly related to the number of cells per HPF (Taraboletti, G., (1987) J. Cell Biol. 105, 2409-2415; Stracke, M.L., et al. (1989) J. Biol. Chem. 264, 21544-21549). Typically, unstimulated motility (background) corresponded to 5-10 cells/HPF and highly responding cells to 70-100 cells/HPF above

*Trade-mark

2128215

18

unstimulated background (i.e., 75-110 total cells/HPF).

For experiments using PT, the toxin was pre-incubated with the cells for 1-2 hr. at room temperature prior to the assay and maintained with the cells throughout the assay (Stracke, M.L., et al. (1987) Biochem. Biophys. Res. Comm. 146, 339-345). The treated cells were tested for their motility response to the chemoattractant as well as for unstimulated random motility.

Purification of Autotaxin. Ammonium sulfate, to a final concentration of 1.2 M, was added to the concentrated A2058 conditioned medium for 1 hr. at 4°C. The solution was spun in a RC2-B Ultraspeed Sorvall centrifuge at 10,000 x g for 15 min. Only the supernatant had the capacity to stimulate motility.

In the first step, the sample was fractionated by hydrophobic interaction chromatography using 200 ml phenyl Sepharose CL-4B column equilibrated into 50 mM Tris (pH 7.5), 5% (v/v) methanol and 1.2 M ammonium sulfate. The supernatant from the ammonium sulfate fractionation was added to this column and eluted using linear gradients of 50 mM Tris (pH 7.5), 5% (v/v) methanol, with decreasing (1.2 - 0.0) M ammonium sulfate and increasing (0-50) % (v/v) ethylene glycol at 1 ml/min.

The active peak was pooled, dialyzed into 50 mM Tris, 0.1 M NaCl, 0.01 M CaCl₂, 20% (v/v) ethylene glycol, and subjected to a second fractionation by lectin affinity chromatography using a 40 ml Affi-Gel concanavalin A column run at 1 ml/min. The sample was eluted in a stepwise

fashion in the same buffer with 0, 10, and 500 mM methyl α -mannopyranoside added successively. Fractions from each step of the gradient were pooled and tested for their capacity to stimulate motility.

5 In the third purification step, the sample that eluted at 500 mM α -methyl-mannopyranoside was dialyzed into 10 mM Tris (pH 7.5) with 30% (v/v) ethylene glycol and fractionated by weak anion exchange chromatography. Chromatography was carried
10 out on a ZORBAX BioSeries-WAX column using a Shimadzu BioLiquid chromatograph and eluted with a linear gradient of (0.0 - 0.4 M) sodium chloride at 3 ml/min.

 The active peak was pooled, dialyzed
15 against 0.1 M sodium phosphate (pH 7.2), 10% (v/v) methanol, and 10% (v/v) ethylene glycol, and subjected to a fourth fractionation step on a series of Spherigel TSK columns (4000SW, 4000SW, 3000SW, 2000SW, in that order). This molecular sieve step
20 was run using the Shimadzu BioLiquid* chromatograph at 0.4 ml/min.

 The active peak was pooled and dialyzed into 10 mM Tris (pH 7.5), 5% (v/v) methanol, 20% (v/v) ethylene glycol and subjected to a fifth
25 (strong anion exchange) chromatography step, a Pro-Pac PA1 column run at 1 ml/min using a Dionex BioLC* with AI450 software. The sample was eluted with a linear gradient of (0.0-0.4M) NaCl.

 In order to calculate activity yields
30 after each step of purification, a unit of activity had to be derived. The dilution curve of autotaxin was biphasic with a broad peak and a linear range at sub-optimal concentrations. One unit of activity/well (i.e., 40 units/ml) was defined as 50%
35 of the maximal activity in a full dilution curve.

*Trade-mark

This allowed calculation of the activity contained in any volume from the dilution needed to achieve 1 unit/well. Therefore, if a 1:10 dilution were needed in order to produce 1 unit of activity/well, the material contained $10 \times 40 = 400$ units/ml.

Gel Electrophoresis. Protein samples were analyzed by SDS-polyacrylamide gel electrophoresis using the conditions of Laemmli (Laemmli U.K. (1970) Nature 227, 680-685). In brief, 7 or 8% SDS-containing polyacrylamide gels were prepared or pre-poured (8-16%) gradient gels were obtained commercially. Samples were prepared with or without reducing conditions (5% β -mercaptoethanol). After electrophoretic separation, the gels were stained using Coomassie Blue G-250 as previously described (Neuhoff, V., et al. (1988) Electrophoresis 9, 255-262). In this staining protocol, which ordinarily requires no destaining step, the Coomassie stain appears to be able to stain as little as 10 ng of protein.

For two-dimensional electrophoresis, the protein, in 20% ethylene glycol, was dried in a Speed-vac and redissolved in loading solution: 9M urea, 1% (v/v) pH 3-10 Bio-Lyte, and 2.5% (v/v) Nonidet* P40. This sample was then subjected to isoelectric focusing (O'Farrell, P.H. (1975) J. Biol. Chem. 250, 4007-4021) using a Bio-Rad tube cell in 120 x 3 mm polyacrylamide tube gels containing 9M urea, 2% (v/v) pH 3-10 Bio-Lyte, 0.25% (v/v) pH 8-10 Bio-Lyte and 2.5% (v/v) Nonidet-P40. Reservoir solutions were 0.01 M phosphoric acid and 0.02 M NaOH. Non-equilibrium isoelectric focusing (O'Farrell, P.H., et al. (1977) Cell 12, 1133-1142) was run initially with constant voltage (500 v) for

*Trade-mark

5 hr. Since the protein was basic, the procedure was repeated under equilibrium conditions (500 v for 17 hr.). Electrophoresis in the second dimension was performed on a 7.5% polyacrylamide using the conditions of Laemmli (1970). The gel was stained with Coomassie Blue G-250 as above.

Preparation of peptides for internal sequence of autotaxin. Homogeneous ATX was sequentially digested with cyanogen bromide and, following reduction and pyridylethylation, with trypsin (Stone, M, et al. (1989) in A Practical Guide to Protein and Peptide Purification for Microsequencing Matsudaira, P.T., ed.) pgs. 33-47, Academic Press, N.Y.). The resulting fragments were then separated by gradient elution on an Aquapore RP300 C-8 reverse phase column: 0.1% (v/v) trifluoroacetic acid and (0-70)% acetonitrile over 85 min. at a flow rate of 0.2 ml/min. A Dionex AI450 BioLC system was utilized and fractions were collected by hand while monitoring the absorbance at 215 nm.

Sequence analysis of peptides. The amino acid sequences of peptides resulting from digestion and purification of ATX peptides #1-7 and 12-18, corresponding to SEQ ID NO:1 through SEQ ID NO:7 and SEQ ID NO:26 through SEQ ID NO:32, respectively, were determined on a Porton Instruments 2020 off-line sequenator using standard program #1. Phenylthiohydantoin amino acid analysis of sequenator runs were performed on a Beckman System Gold HPLC using a modified sodium acetate gradient program and a Hewlett-Packard C-18 column. ATX-100 (SEQ ID NO:8), ATX-101 (SEQ ID NO:9), ATX-102 (SEQ

2128215

22

ID NO:10), ATX-103 (SEQ ID NO:11) and ATX 104 (SEQ ID NO:33) were sequenced from gel-purified ATX.

Protein databases (Pearson, W.R., et al. (1988) Proc. Natl. Acad. Sci. USA 85, 2444-2448) that were searched for homologies in amino acid sequence with the ATX peptides include: GenBank (68.0), EMBL (27.0), SWISS-PROT (18.0), and GenPept (64.3).

EXAMPLE 1

10 Purification of Autotaxin

The A2058 cells had been previously shown to produce protein factors which stimulate motility in an autocrine fashion (Liotta, L.A., et al. (1986) Proc. Natl. Acad. Sci. USA 83, 3302-3306).

15 Conditioned medium from these cells was therefore used to identify and purify a new motility-stimulating factor, which is here named autotaxin. Since the purification was monitored with a biological assay, motility-stimulating activity had to be maintained throughout. The activity proved to be labile to freezing, acidic buffers, proteases (but not DNase or RNase), reduction, strong chaotropic agents (e.g. > 4 M urea), and a variety of organic solvents (isopropanol, ethanol,

20 acetonitrile). An organic solvent, ethylene glycol, which did not decrease bioactivity, was added for both storage and chromatographic separation.

25 100-200 L of serum-free conditioned medium were required in order to produce enough autotaxin for amino acid sequence analysis. The medium contained low concentrations of both BSA (5 µg/ml) which was needed as a carrier protein and insulin (10 µg/ml) which was required to support cell growth

30 100-200 L of serum-free conditioned medium were required in order to produce enough autotaxin for amino acid sequence analysis. The medium contained low concentrations of both BSA (5 µg/ml) which was needed as a carrier protein and insulin (10 µg/ml) which was required to support cell growth

2128215

23

in low protein medium. Ultrafiltration to concentrate this large volume was performed with low protein-binding YM30 membranes which retain molecular species with $M_r > 30,000$. As seen in Table 1, 200 L of conditioned medium prepared in this manner resulted in 10×10^6 units of activity. However, the initial unfractionated conditioned medium contained additional substances known to stimulate activity, particularly insulin, which does not completely wash out in the ultrafiltration step and which is additive to the motility stimulating activity in a complex manner (Stracke, M.L., et al. (1989) J. Biol. Chem. 264, 21544-21549). This had to be taken into account in order to determine yields for subsequent steps in which insulin had been removed.

TABLE 1. PURIFICATION OF AUTOTAXIN

Purification Step	Protein (mg)	Activity ^a (total units)	Specific Activity (units/mg)	Recovery (%) ^b
200 L Conditioned Medium	33,000	10,000,000 ^c	300	
Phenyl Sepharose	1,235	460,000	370	100
Concanavalin A	58	660,000	11,400	100
Weak Anion Exchange	4.5	490,000	110,000	100
25 TSK Molecular Sieves	-0.4 ^d	220,000	550,000	48
Strong Anion Exchange	-0.04 ^d	24,000 ^e	600,000	5.2

^a Activity calculated from Boyden chamber assay. The dilution which resulted in 50% of maximal activity (generally approximately 20 laser density units or ~40 cells/HPF) was chosen to have 1 unit of activity per well (equivalent to 40 units/ml).

30 ^b Recovery was estimated from activity, after the first purification column (i.e., phenyl sepharose).

^c Initial activity in the unfractionated conditioned medium reflected the fact that insulin was used in the medium as a necessary growth factor under low protein conditions.

35 ^d Estimated protein is based on quantification by amino acid analysis.

^e This specific activity for purified protein corresponds to ~10 fmol ATX/unit of motility activity (in a Boyden chamber well).

2128215

24

The first step in the purification involved fractionation by hydrophobic interaction chromatography using a phenyl Sepharose CL-4B column. The results are shown in FIG. 1. Most proteins, including insulin, eluted from the column in early fractions or in the void. However, the peak of activity eluted relatively late. The activity which was purified was estimated as 460,000 units $\pm$ 20% (Table 1). As the pooled peak of activity from the phenyl Sepharose fractionation is considered to be the first sample without significant insulin contamination, subsequent yields are measured against its total activity. Gel electrophoresis of a small portion of the pooled peak of activity (FIG. 6A, column 2) revealed a large number of protein bands with BSA predominant from the original conditioned medium.

In the second step of purification, the active peak was applied to the lectin affinity column, Affi-Gel concanavalin A. As shown in FIG. 2, most protein (estimated to be 90% of the total absorbance at 280 nm) failed to bind to the column at all. The non-binding fraction contained essentially no motility-stimulating activity (see dotted line in FIG. 2). When a linear gradient of methyl α -D-mannopyranoside was applied to the column, chemotactic activity eluted off in a prolonged zone, beginning at a concentration of approximately 20 mM sugar. Consequently, a step gradient was used to elute. Pure BSA failed to bind to con A.

Activity was found primarily in the 500 mM step of methyl α -D-mannopyranoside. There appeared to be no significant loss of activity as seen in

Table 1; however, specific activity (activity/mg total protein) increased thirty-fold. Gel electrophoresis of the pooled and concentrated peak (FIG. 6A, column 3) revealed that the BSA overload was no longer apparent and the number of bands were much reduced. When the unbound protein was concentrated and applied to a gel, it appeared identical to the active peak from phenyl Sepharose-4B with a large BSA band.

The third purification step involved fractionating the previous active peak by weak anion exchange chromatography as shown in FIG. 3. Under the running conditions, activity eluted in a broad peak-shoulder or double peak in the middle of the shallow portion (0.0-0.4 M) of the NaCl gradient. The largest proportion of protein, lacking in motility-stimulating capacity, bound strongly to the column and eluted off in high salt (1 M NaCl). There appeared to be no significant loss of activity, though specific activity increased by twenty-fold (Table 1). Analysis by gel electrophoresis of both the peak (28-34 min. in FIG. 3) and the shoulder (35-45 min. in FIG. 3) is shown in FIG. 6A (columns 4 and 5, respectively). Two predominant protein bands resulted: a broad doublet around 25-35 kDa and a second doublet around 110-130 kDa.

In the fourth purification step, the active peak was applied to a series of molecular sieves. Spectrophotometric monitoring of the eluant revealed two large peaks of protein (FIG. 4). Activity corresponded to the first, higher molecular weight peak. Recovery of activity was ~48% with a five-fold increase in specific activity. Analysis by gel electrophoresis was performed under non-

2128215

26

reducing and reducing conditions as shown in FIG. 6B (columns 2 and 3, respectively). This fractionation step had essentially removed all contaminating protein of molecular weight < 55 kDa. The
5 predominant band remaining has a molecular weight of 120 kDa unreduced and 125 kDa reduced; there are two minor bands with molecular weights 85 kDa and 60 kDa. The fact that the 120 kDa protein changes so little in electrophoretic mobility after reduction
10 would tend to indicate a paucity of disulfide bonds. However, the existing disulfide bonds have functional significance because motility-stimulating activity is labile to reduction.

The fifth purification step involved
15 fractionation of the active peak by strong anion exchange chromatography. As shown in FIG. 5, activity corresponds to two broad optical absorbance peaks in the middle of the gradient with contaminating proteins eluting earlier. These two
20 peaks were identical by amino acid analysis and by polyacrylamide gel electrophoretic separation. They presumably represent different glycosylation states of the same parent protein. Activity is shown in FIG. 5 at two different sample dilutions. Several
25 dilutions of the fractionated samples were often necessary in order to resolve the true "peak" of activity as the shape of the ATX dilution curve was not sharp due to saturation and down regulation at high concentrations. Recovery from this
30 chromatographic step is lower (5% compared to phenyl Sepharose), as might be expected when a minute quantity of protein is applied to a column; however, specific activity again increased (Table 1). Analysis by gel electrophoresis revealed a single

2128215

27

protein band at molecular weight 120 kDa, unreduced (FIG. 6C, column 2).

EXAMPLE 2

Characterization of Autotaxin

5 Two dimensional gel electrophoresis of the purified protein (FIG. 7) revealed a single predominant band. The band slopes downward slightly toward the basic side of the gel in a manner that is characteristic of glycosylated proteins. A basic pI
10 of 7.7 ± 0.2 was essentially the same whether the isoelectric focusing was run under non-equilibrium conditions (5 hr.) or was allowed to go to equilibrium (17 hr.).

 A dilution curve of the purified protein
15 is shown in FIG. 8. The protein is active in the pM range and 1 unit of activity appears to correspond to a concentration of 400-600 pMolar (or approximately 10 fmol of autotaxin/Boyden chamber well). When dilutions were begun at higher
20 concentrations of ATX, the resultant curve showed a broad plateau with down regulation at the highest concentrations. The motility response to purified autotaxin is highly sensitive to pertussis toxin (Table 2 and FIG. 9) with approximately 95%
25 inhibition of activity at 0.5 μ g/ml PT.

2128215

28

TABLE 2. Effect of Pertussis Toxin (PT)
on Autotaxin-stimulated motility

5	A2058 Motility Response (density units ¹)		
	control cells ²		Pertussis toxin- treated cells ³
	Condition medium ⁴	60.3	0.4
10	Purified Autotaxin	38.5	0.0

¹ Chemotaxis quantitated by motility assay (Stracke, M.L., et al. (1978) Biochem. Biophys. Res. Comm. 146, 339-345)

15 ² A2058 cell suspended at 2×10^6 cells/ml in DMEM supplemented with 1 mg/ml bovine serum and rocked at room temperature for 1 hr.

³ As control with 0.5 μ g/ml pertussis toxin.

20 ⁴ Prepared by adding DMEM without phenol red supplemented with 0.1 mg/ml bovine serum albumin to subconfluent flasks of A2058 cells. The medium was harvested after 2 days incubation at 37°C in a humidified atmosphere and concentrated 25-30 fold using an Amicon ultrafiltration assembly with a YM-30 membrane.

25

Checkerboard analysis was performed to assess the random (chemokinetic) versus the directed (chemotactic) nature of the motility response to autotaxin. Chambers were assembled with different concentrations of ATX above and below the filter, using ATX purified through the weak anion exchange fractionation step. Squares below the diagonal reflect response to a positive gradient, squares above reflect response to a negative gradient, and squares on the diagonal reflect random motility in the absence of a gradient. ATX stimulates both chemotactic and chemokinetic responses (FIG. 10),

30

35

2128215

29

with chemotactic responses as high as fifteen-fold above background and chemokinesis as high as eight-fold above background.

5 Amino acid analysis after complete acid hydrolysis was used to quantitate purified protein. This hydrolysis was carried out on protein excised from a polyacrylamide gel and presumed to be pure. The analysis indicated that 2.7 nmol of protein was present after fractionation on the molecular sieve.
10 After fractionation by strong anion exchange chromatography, approximately 300 pmol remained. The results of the analysis are shown in Table 3.

15 TABLE 3. AMINO ACID COMPOSITION OF AUTOTAXIN
(CYS and TRP were not determined in this analysis)

	<u>Amino Acid</u>	<u>Residues/100</u>
	ASX	12.5
	THR	6.0
	SER	5.7
20	GLX	9.4
	PRO	7.4
	GLY	7.0
	ALA	3.9
	VAL	6.7
25	MET	1.2
	ILE	4.3
	LEU	9.0
	TYR	5.2
	PHE	5.2
30	HIS	3.8
	LYS	7.4
	ARG	5.4

EXAMPLE 3ATX Degradation and Determination of
Amino Acid Sequence

Attempts to obtain N-terminal sequence
5 information from purified ATX repeatedly proved
futile. The purified protein was therefore,
sequentially digested and the resulting peptides
fractionated by reverse phase chromatography. The
results are shown in FIG. 11. Multiple sharp peaks
10 including clusters at both the hydrophilic and
hydrophobic ends of the gradient are seen.

Several of these peptide peaks were chosen
randomly for Edman degradation and N-terminal amino
acid sequence analysis. Seven of the eight peaks
15 (shown in FIG. 11) chosen gave clear single sequence
information as seen in Table 4. Using material from
a separate digestion and purification, the remaining
four sequences were also obtained.

Separate sense and antisense
20 oligonucleotide probes were synthesized according to
the fragment sequences of Table 4 by methods known
to one skilled in the art. Representative probes
are shown in Table 5.

TABLE 4. Peptide sequences for Autotaxin.

	PEPTIDE NO.	AMINO ACID SEQUENCE	SEQ ID: NO:	NAME
5	1.	WHVAAN	SEQ ID NO:1	ATX 18
	2.	PXLDVYK	SEQ ID NO:2	ATX 19
	3.	YPAFK	SEQ ID NO:3	ATX 20
	4.	QAEVS	SEQ ID NO:4	ATX 24
	5.	PEEVTPNYL	SEQ ID NO:5	ATX 29
10	6.	YDVPWNETI	SEQ ID NO:6	ATX 47
	7.	SPPFENINLY	SEQ ID NO:7	ATX 48
	8.	GGQPLWITATK	SEQ ID NO:8	ATX 100
	9.	VNSMQTVFVGY- GPTFK	SEQ ID NO:9	ATX 101
15	10.	DIEHLTSLDFFR	SEQ ID NO:10	ATX 102
	11.	TEFLSNYLTNVDD- ITLVPGTLGR	SEQ ID NO:11	ATX 103
	12.	QYLHQYGSS	SEQ ID NO:26	ATX 37
20	13.	VLNYF	SEQ ID NO:27	ATX 39
	14.	YLNAT	SEQ ID NO:28	ATX 40
	15.	HLLYGRPAVLY	SEQ ID NO:29	ATX 41
	16.	SYPEILTPADN	SEQ ID NO:30	ATX 44
	17.	XYGFLFPPYLSSSP	SEQ ID NO:31	ATX 53
	18.	TFPNLYVF/LAQGLYWS	SEQ ID NO:32	ATX 59
	19.	VNVISGPIFDYDYGDLH/A- DTEDK	SEQ ID NO:33	ATX 104

Peptide numbers 1-7 refer to peaks numbered in FIG. 11. Peptide numbers 12-18 refer to peptides purified from the preparation which yielded peptide numbers 1-7. Peptides 8-11 and 19, are from a separate purification, not shown in FIG. 11.

X refers to potentially glycosylated residues.

66597-197

32

TABLE 5.

5 Oligonucleotides synthesized from peptide sequences of autotaxin (ATX). The number of the oligonucleotide corresponds to the ATX peptide number as per Table 4. The final letter suffix distinguishes whether the oligonucleotide is a sense (S) or antisense (A) sequence.

	<u>Oligo</u>	<u>Sequence</u>	<u>SEQ ID NO:</u>
	A-18A	GTT-GGC-AGC-NAC-RTG-CCA	SEQ ID NO:12
10	A-18S	TGG-CAY-GTN-GCT-GCC-AAC	SEQ ID NO:13
	A-20A	CTT-GAA-GGC-AGG-GTA	SEQ ID NO:14
	A-20S	TAY-CCT-GCN-TTY-AAG	SEQ ID NO:15
	A-29A	GGT-NAC-YTC-YTC-AGG	SEQ ID NO:16
	A-29S	CCT-GAR-GAR-GTN-ACC	SEQ ID NO:17
15	A-47A	NGT-NGC-RTC-RAA-TGG-CAC-RTC	SEQ ID NO:18
	A-47S	GAY-GTG-CCA-TTY-GAY-GCN-ACN	SEQ ID NO:19
	A-48A	GTT-DAT-RTT-STC-RAA-TGG-GGG	SEQ ID NO:20
	A-48S	CCC-CCA-TTT-GAG-AAC-ATC-AAC	SEQ ID NO:21
	A-100A	CTT-NGT-NGC-NGT-DAT-CCA-NAR-	SEQ ID NO:22
20		GGG-YTG-GCC-GCC	
	A-100S	GGC-GGC-CAR-CCC-YTN-TGG-ATH-	SEQ ID NO:23
		ACN-GCN-ACN-AAG	
	A-101A	CTT-RAA-GGT-GGG-GCC-RTA-GCC-	SEQ ID NO:24
		CAC-RAA-GAC-TGT-YTG-CAT	
25	A-101S	ATG-CAR-ACA-GTC-TTY-GTG-GGC-	SEQ ID NO:25
		TAY-GGC-CCC-ACC-TTY-AAR	

EXAMPLE 4Molecular cloning of autotaxin cDNA

30 A cDNA library was prepared in phage (λ_{zap})* from the same A2058 tumor cells known to synthesize ATX. This library has been screened for the presence of the ATX sequence by the following methods. First, a 200 base pair (bp) sequence of DNA was prepared using PCR (polymerase chain

*Trade-mark

2128215

33

5 reaction) technology with the same λ_{2p} library as template and the derived oligonucleotides A-48S (SEQ ID NO:21) and A-100A (SEQ ID NO:22) as priming sequences. This 200 bp sequence of the ATX mRNA has been used to perform initial screens of the full library. Alternatively, a mixture of derived oligonucleotides was also used to screen the full cDNA library. Both screening methods have produced positive cDNA clones.

10

EXAMPLE 5Antipeptide Antibodies

15 Rabbits were injected with ATX-102 (SEQ ID NO:10) which had been cross-linked to bovine serum albumin. Antisera from these rabbits was subjected to salt precipitation followed by purification using affinity chromatography with Affi-Gel 10 beads covalently linked to the peptide, ATX-102 (SEQ ID NO:10). This affinity purified antibody reacted with the partially
20 purified protein on immunoblots. This same antibody has been used to perform immunohistochemical stains on human tissue.

66597-197

34

* * * * *

5 While the foregoing invention has been
described in some detail for purposes of clarity
and understanding, it will be appreciated by one
skilled in the art from a reading of this
disclosure that various changes in form and detail
10 can be made without departing from the true scope
of the present invention and appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: STRACKE, MARY
LIOTTA, LANCE
SCHIFFMANN, ELLIOTT
KRUTZSCH, HENRY

(ii) TITLE OF INVENTION: MOTILITY STIMULATING PROTEIN USEFUL IN
CANCER DIAGNOSIS AND THERAPY

(iii) NUMBER OF SEQUENCES: 33

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: CUSHMAN DARBY AND CUSHMAN
(B) STREET: 1615 L STREET, N.W.
(C) CITY: WASHINGTON
(D) STATE: D.C.
(E) COUNTRY: U.S.A.
(F) ZIP: 20036

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Tape
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: SCOTT, WATSON T
(B) REGISTRATION NUMBER: 26581

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (202)-861-3000
(B) TELEFAX: (202) 822-0944
(C) TELEX: 6714627CUSH

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

2128215

36

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Trp His Val Ala Ala Asn
1 5

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Pro Xaa Leu Asp Val Tyr Lys
1 5

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Tyr Pro Ala Phe Lys
1 5

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Gln Ala Glu Val Ser
1 5

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 amino acids

2128215

37

(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Pro Glu Glu Val Thr Xaa Pro Asn Tyr Leu
1 5 10

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Tyr Asp Val Pro Trp Asn Glu Thr Ile
1 5

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Ser Pro Pro Phe Glu Asn Ile Asn Leu Tyr
1 5 10

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Gly Gly Gln Pro Leu Trp Ile Thr Ala Thr Lys
1 5 10

2128215

38

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Val	Asn	Ser	Met	Gln	Thr	Val	Phe	Val	Gly	Tyr	Gly	Pro	Thr	Phe	Lys
1				5					10					15	

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Asp	Ile	Glu	His	Leu	Thr	Ser	Leu	Asp	Phe	Phe	Arg
1				5					10		

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Thr	Glu	Phe	Leu	Ser	Asn	Tyr	Leu	Thr	Asn	Val	Asp	Asp	Ile	Thr	Leu
1				5					10					15	

Val	Pro	Gly	Thr	Leu	Gly	Arg
				20		

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

2128215

39

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GTTGGCAGCN ACRTGCCA

18

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TGGCAYGTNG CTGCCAAC

18

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CTTGAAGGCA GGGTA

15

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

TAYCCTGCNT TYAAG

15

2128215

40

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GGTNACYTCY TCAGG

15

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CCTGARGARG TNACC

15

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

NGTNGCRTCR AATGGCACRT C

21

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

41

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GAYGTGCCAT TYGAYGCNAC N

21

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GTTDATRTTS TCRAATGGGG G

21

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

CCCCCATTTG AGAACATCAA C

21

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

CTTNGTNGCN GTDATCCANA RGGGYTGGCC GCC

33

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid

2128215

42

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GGCGGCCARC CCYTNTGGAT HACNGCNACN AAG

33

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

CTTRAAGGTG GGGCCRTAGC CCACRAAGAC TGTYTGCAT

39

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

ATGCARACAG TCTTYGTGGG CTAYGGCCCC ACCTTYAAR

39

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

43

Gln Tyr Leu His Gln Tyr Gly Ser Ser
1 5

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Val Leu Asn Tyr Phe
1 5

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Tyr Leu Asn Ala Thr
1 5

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

His Leu Leu Tyr Gly Arg Pro Ala Val Leu Tyr
1 5 10

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids

2128215

44

- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Ser	Tyr	Pro	Glu	Ile	Leu	Thr	Pro	Ala	Asp	Asn
1				5					10	

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Xaa	Tyr	Gly	Phe	Leu	Phe	Pro	Pro	Tyr	Leu	Ser	Ser	Ser	Pro
1				5					10				

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Thr	Phe	Pro	Asn	Leu	Tyr	Val	Xaa	Ala	Gln	Gly	Leu	Tyr	Trp	Ser
1				5					10					15

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

45

Val Asn Val Ile Ser Gly Pro Ile Phe Asp Tyr Asp Tyr Asp Gly Leu
1 5 10 15

Xaa Asp Thr Glu Asp Lys
20

66597-197

46

CLAIMS:

1. An isolated polypeptide selected from human autotaxin protein in homogeneous form or a fragment of said protein, wherein

5 (A) said protein has an isoelectric point of about 7.7, a molecular weight of about 125 kDa as determined by SDS-polyacrylamide gel electrophoresis under reducing conditions, and induces a motility in human A2508 melanoma cells which is inhibited by pertussis toxin;

10 (B) said fragment comprises an amino acid sequence selected from the group consisting of SEQ ID Nos. 1-11 and 26-33; and

(C) antibody suitable for immunohistochemical staining of cancerous human tissue can be raised to each of
15 said protein and said fragment.

2. The polypeptide of claim 1, wherein said human autotaxin protein has the amino acid sequence of an autotaxin protein isolated from human A2508 melanoma cells.

3. The polypeptide of claim 1 or claim 2 bound to a
20 solid support.

4. An isolated polynucleotide coding for a polypeptide as defined in any one of claims 1 to 3.

5. The polynucleotide of claim 4, wherein said DNA codes for an amino acid that comprises a sequence selected
25 from the group consisting of SEQ ID Nos. 1-11 and 26-33.

6. The polynucleotide of claim 4, comprising a nucleotide sequence selected from the group consisting of SEQ ID Nos. 12-25.

66597-197

47

7. A recombinant DNA molecule comprising the polynucleotide of any one of claims 4 to 6 and a vector.

8. A cell comprising the recombinant DNA molecule of claim 7.

5 9. An isolated cell comprising the recombinant DNA molecule of claim 7.

10. A method of producing a polypeptide as defined in any one of claims 1 to 3, comprising the steps of:

(i) culturing a cell comprising a polynucleotide
10 coding for said polypeptide and a vector under conditions such that said polynucleotide is expressed, and said polypeptide is thereby produced, and

(ii) isolating said polypeptide.

11. The method of claim 10, further comprising the
15 steps of:

(i) collecting and concentrating supernatant from said cultured cell, whereby a first preparation of polypeptide is produced;

(ii) salt fractionating said first preparation to
20 produce a second preparation of polypeptide;


(iii) isolating said peptide from said second polypeptide preparation, so that said peptide is obtained in substantially pure form.

12. The method of claim 11, wherein said step of
25 isolating said peptide is effected by column chromatography.

13. An isolated antibody having binding affinity for a polypeptide as defined in any one of claims 1 to 3.

1/11


FIG. 1



SUBSTITUTE SHEET


2/11

FIG. 2




3/11

FIG. 3




4/11

FIG. 4



5/11

FIG. 5



6/11

FIG. 6A




FIG. 6B






FIG. 6C




7/11

FIG. 7




8/11

FIG. 8



9/11

FIG. 9



10/11

FIG. 10


		Upper Wells		
		0	0.01%	0.1%
Lower Wells	0	4.8 ± 0.3	13.7 ± 0.8	33.8 ± 1.6
	0.01%	45.4 ± 4.0	39.3 ± 2.6	36.9 ± 1.4
	0.1%	75.6 ± 1.9	58.3 ± 3.1	41.0 ± 3.4

SUBSTITUTE SHEET

2128215

11/11

FIG. II



SUBSTITUTE SHEET