



(86) **Date de dépôt PCT/PCT Filing Date:** 2013/01/09
(87) **Date publication PCT/PCT Publication Date:** 2013/07/18
(85) **Entrée phase nationale/National Entry:** 2014/07/08
(86) **N° demande PCT/PCT Application No.:** US 2013/020910
(87) **N° publication PCT/PCT Publication No.:** 2013/106489
(30) **Priorité/Priority:** 2012/01/09 (US61/584,680)

(51) **Cl.Int./Int.Cl.** **C07K 16/00** (2006.01),
C07K 14/54 (2006.01), **C07K 16/10** (2006.01),
C07K 16/46 (2006.01), **C07K 19/00** (2006.01)
(71) **Demandeur/Applicant:**
THE SCRIPPS RESEARCH INSTITUTE, US
(72) **Inventeurs/Inventors:**
SMIDER, VAUGHN, US;
BAZIRGAN, OMAR A., US;
MAO, HONGYUAN HELEN, US
(74) **Agent:** MBM INTELLECTUAL PROPERTY LAW LLP

(54) **Titre :** ANTICORPS HUMANISES A CDR3 ULTRALONGUES
(54) **Title:** HUMANIZED ANTIBODIES WITH ULTRALONG CDR3S

Figure 1

BLV1H12	CTSVHQ	ETKKYQ	SCPDGYRERSDC	SNRPAC	CGTSD	CCRVSVFGNCLTTLFVSYSYTYNYEW	HVDVWGQGLLTVVSS (SEQ ID NO : 360)
BLV5B8	CTTVHQ	ETRKT	CSDGYIAVDS	CGRGQSDGCVNDCNSCYYGWRNCRQPAIRSYEF		HVDAWGRGLLTVVSS (SEQ ID NO : 361)	
BLV5D3	CSSVTQ	RTHVSR	SCPDGCSGDG	CVDGCC	CSAYRC	CYTPGVRDLSTSYSTITYYEW	NVDAWGRGLLTVVSS (SEQ ID NO : 362)
BLV8C11	CTTVHQ	KTTRKT	CCSDAYRYDS	CGSGCD	CCGAD	CYVFGACTFGLDSSYSYIYIQW	YVDAWGQGLLTVVSS (SEQ ID NO : 363)
BF4E9	CTTVHQ	IF	CPDGYSYGYG	CGYGYG	CSGYD	CYGYGGYGYGGYGGYSSYSYSYEF	YGDWGWQGLLTVVSS (SEQ ID NO : 364)
BF1H1	CTTVHP		SPDGYSYGYG	CGYGYG	CSGYD	CYGYGGYGYGGYGGYSSYSYSYS	(SEQ ID NO : 365)
F18	CTTVHQ	IR	CPDGYSYGYG	CGYGSYGYSGYDCYGYGGYGGYGGYSSYS		(SEQ ID NO : 366)	

(57) **Abrégé/Abstract:**
The present disclosure provides humanized antibodies, including antibodies comprising an ultralong CDR3 (e.g. bovine) and uses thereof.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2013/106489 A1

(43) International Publication Date
18 July 2013 (18.07.2013)

(51) International Patent Classification:

C07K 16/00 (2006.01) *C07K 14/54* (2006.01)
C07K 16/10 (2006.01) *C07K 19/00* (2006.01)
C07K 16/46 (2006.01)

(21) International Application Number:

PCT/US2013/020910

(22) International Filing Date:

9 January 2013 (09.01.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/584,680 9 January 2012 (09.01.2012) US

(71) Applicant: **THE SCRIPPS RESEARCH INSTITUTE**
[US/US]; 10550 North Torrey Pines Road, La Jolla, Cali-
fornia 92037 (US).

(72) Inventors: **SMIDER, Vaughn**; 7483 Healis Place, San
Diego, California 92129 (US). **BAZIRGAN, Omar A.**;
12111 Waverly Downs Lane, San Diego, California 92128
(US). **MAO, Hongyuan Helen**; 3571 Overpark Road, San
Diego, California 92130 (US).

(74) Agents: **MCNICHOLAS, Janet M.** et al.; K&L GATES
LLP, P.O. Box 1135, Chicago, Illinois 60690-1135 (US).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

(54) Title: HUMANIZED ANTIBODIES WITH ULTRALONG CDR3S

Figure 1

BLV1H12	CTEVHQ	ETKKYQ	SCPDGYRTRSDCSNRDACSISDCCRVSVFQNCITLIPVSYSYTNVYEW	HVDVWGQGLLVTVSS (SEQ ID NO : 360)
BLV5B8	CTEVHQ	ETRRK	CSDEYIAYDGCERGGSDGCVNDGNSCYGWRNCRQFALHSYEF	HVDARGRLLVTVSS (SEQ ID NO : 361)
BLV5D3	CSEVTQ	RTHVSK	CCPDGSDGDCGYDCCCSAYRCYTPGVRLSCTSYSTITYEW	NVDARGRLLVTVSS (SEQ ID NO : 362)
BLV6C11	CTEVHQ	ETTRKT	CCSDAYRYDSGCESGDCGAGCYVFGACTFGLSSYSYIYICW	YVDARWGQGLLVTVSS (SEQ ID NO : 363)
BF4E9	CTEVHQ	ET	CPDGYSYGYGCGYGYGCGGYDGYGGYGYGGYSSYSYSEY	YGDARWGQGLLVTVSS (SEQ ID NO : 364)
BF1H1	CTEVH2	ET	SPDGYSYGYGCGYGYGCGGYDGYGGYGYGGYSSYSYS	(SEQ ID NO : 365)
F19	CTEVHQ	ET	CPDGYSYGYGCGYGYGCGGYDGYGGYGYGGYSSYS	(SEQ ID NO : 366)

(57) Abstract: The present disclosure provides humanized antibodies, including antibodies comprising an ultralong CDR3 (e.g. bovine) and uses thereof.

WO 2013/106489 A1

HUMANIZED ANTIBODIES WITH ULTRALONG CDR3S

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No.
5 61/584,680 filed January 9, 2012, which is incorporated by reference herein in its entirety.

FIELD

[0002] The present disclosure relates to humanized antibodies, including
antibodies comprising an ultralong CDR3.

10

BACKGROUND

[0003] Antibodies are natural proteins that the vertebrate immune system forms in
response to foreign substances (antigens), primarily for defense against infection. For over
a century, antibodies have been induced in animals under artificial conditions and
15 harvested for use in therapy or diagnosis of disease conditions, or for biological research.
Each individual antibody producing cell produces a single type of antibody with a
chemically defined composition, however, antibodies obtained directly from animal serum
in response to antigen inoculation actually comprise an ensemble of non-identical
molecules (e.g., polyclonal antibodies) made from an ensemble of individual antibody
20 producing cells.

[0004] Some bovine antibodies have unusually long VH CDR3 sequences
compared to other vertebrates. For example, about 10% of IgM contains "ultralong" CDR3
sequences, which can be up to 61 amino acids long. These unusual CDR3s often have
multiple cysteines. Functional VH genes form through a process called V(D)J
25 recombination, wherein the D-region encodes a significant proportion of CDR3. A unique
D-region encoding an ultralong sequence has been identified in cattle. Ultralong CDR3s
are partially encoded in the cattle genome, and provide a unique characteristic of their
antibody repertoire in comparison to humans. Kaushik *et al.* (U.S. Patent Nos. 6,740,747
and 7,196,185) disclose several bovine germline D-gene sequences unique to cattle stated
30 to be useful as probes and a bovine VDJ cassette stated to be useful as a vaccine vector.

SUMMARY

[0005] The present disclosure provides humanized antibodies, including antibodies
comprising an ultralong CDR3, methods of making same, and uses thereof.

[0006] The present disclosure provides a humanized antibody or binding fragment thereof comprising an ultralong CDR3.

[0007] In some embodiments, the humanized antibody or binding fragment thereof comprises one or more human variable region framework sequences.

5 [0008] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 is 35 amino acids in length or longer, 40 amino acids in length or longer, 45 amino acids in length or longer, 50 amino acids in length or longer, 55 amino acids in length or longer, or 60 amino acids in length or longer.

10 [0009] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 is 35 amino acids in length or longer.

[0010] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises 3 or more cysteine residues, 4 or more cysteine residues, 5 or more cysteine residues, 6 or more cysteine residues, 7 or more cysteine residues, 8 or more cysteine residues, 9 or more cysteine residues, 10 or more
15 cysteine residues, 11 or more cysteine residues, or 12 or more cysteine residues.

[0011] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises 3 or more cysteine residues.

[0012] In some embodiments of each or any of the above or below mentioned embodiments, the antibodies or binding fragments thereof comprise a cysteine motif.

20 [0013] In some embodiments of each or any of the above or below mentioned embodiments, the cysteine motif is selected from the group consisting of:
CX₁₀CX₅CX₅CXCX₇C (SEQ ID NO: 41), CX₁₀CX₆CX₅CXCX₁₅C (SEQ ID NO: 42),
CX₁₁CXCX₅C (SEQ ID NO: 43), CX₁₁CX₅CX₅CXCX₇C (SEQ ID NO: 44),
CX₁₀CX₆CX₅CXCX₁₃C (SEQ ID NO: 45), CX₁₀CX₅CXCX₄CX₈C (SEQ ID NO: 46),
25 CX₁₀CX₆CX₆CXCX₇C (SEQ ID NO: 47), CX₁₀CX₄CX₇CXCX₈C (SEQ ID NO: 48),
CX₁₀CX₄CX₇CXCX₇C (SEQ ID NO: 49), CX₁₃CX₈CX₈C (SEQ ID NO: 50),
CX₁₀CX₆CX₅CXCX₇C (SEQ ID NO: 51), CX₁₀CX₅CX₅C (SEQ ID NO: 52),
CX₁₀CX₅CX₆CXCX₇C (SEQ ID NO: 53), CX₁₀CX₆CX₅CX₇CX₉C (SEQ ID NO: 54),
CX₉CX₇CX₅CXCX₇C (SEQ ID NO: 55), CX₁₀CX₆CX₅CXCX₉C (SEQ ID NO: 56),
30 CX₁₀CXCX₄CX₅CX₁₁C (SEQ ID NO: 57), CX₇CX₃CX₆CX₅CXCX₅CX₁₀C (SEQ ID NO: 58),
CX₁₀CXCX₄CX₅CXCX₂CX₃C (SEQ ID NO: 59), CX₁₆CX₅CXC (SEQ ID NO: 60),
CX₆CX₄CXCX₄CX₅C (SEQ ID NO: 61), CX₁₁CX₄CX₅CX₆CX₃C (SEQ ID NO: 62),
CX₈CX₂CX₆CX₅C (SEQ ID NO: 63), CX₁₀CX₅CX₅CXCX₁₀C (SEQ ID NO: 64),
CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65), CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66),
35 CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67), CX₁₅CX₅CXC (SEQ ID NO: 68),
CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69), CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70),

CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72),
 CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74), CX₁₀CXCX₆C
 (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C
 (SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C
 5 (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCX₁₄C (SEQ ID
 NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83),
 CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85),
 CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87),
 CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89),
 10 CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91),
 CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93),
 CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ
 ID NO: 95).

[0014] In some embodiments of each or any of the above or below mentioned
 15 embodiments, the cysteine motif is selected from the group consisting of:
 CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC
 (SEQ ID NO: 97), CX₇CXCX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 98),
 CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID
 NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101),
 20 CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102), CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID
 NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104),
 CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC
 (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107),
 CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ
 25 ID NO: 109), CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C (SEQ ID NO: 110),
 CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ
 ID NO: 112), CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113),
 CX₉CCX₄CX₂CXCCXCX₄CX₃C (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ
 ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116),
 30 CX₆CCXCX₅CX₄CCXCX₅CX₂C (SEQ ID NO: 117), CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID
 NO: 118), CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119),
 CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO:
 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO:
 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO:
 35 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID
 NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID

NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137),
 5 CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152).

[0015] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises 2 to 6 disulfide bonds.

[0016] In some embodiments of each or any of the above or below mentioned
 15 embodiments, the ultralong CDR3 comprises SEQ ID NO: 40 or a derivative thereof.

[0017] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises amino acid residues 3-6 of any of one SEQ ID NO: 1-4.

[0018] In some embodiments of each or any of the above or below mentioned
 20 embodiments, the ultralong CDR3 comprises a non-human DH or a derivative thereof.

[0019] In some embodiments of each or any of the above or below mentioned embodiments, the non-human DH is SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, or SEQ ID NO: 12.

[0020] In some embodiments of each or any of the above or below mentioned
 25 embodiments, the ultralong CDR3 comprises a JH sequence or a derivative thereof.

[0021] In some embodiments of each or any of the above or below mentioned embodiments, the JH sequence is SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, or SEQ ID NO: 17.

[0022] In some embodiments of each or any of the above or below mentioned
 30 embodiments, the ultralong CDR3 comprises: a non-human or human VH sequence (e.g., a germline VH) or a derivative thereof; a non-human DH sequence or a derivative thereof; and/or JH sequence or derivative thereof.

[0023] In some embodiments of each or any of the above or below mentioned
 35 embodiments, the ultralong CDR3 comprises an additional amino acid sequence comprising two to six amino acid residues or more positioned between the VH sequence and the DH sequence.

[0024] In some embodiments of each or any of the above or below mentioned embodiments, the additional amino acid sequence is selected from the group consisting of: IR, IF, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20 or SEQ ID NO: 21.

5 [0025] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a sequence derived from or based on SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, or SEQ ID NO: 28.

10 [0026] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a bovine sequence, a non-bovine sequence, an antibody sequence, or a non-antibody sequence.

[0027] In some embodiments of each or any of the above or below mentioned embodiments, the non-antibody sequence is a synthetic sequence.

15 [0028] In some embodiments of each or any of the above or below mentioned embodiments, the non-antibody sequence is a cytokine sequence, a lymphokine sequence, a chemokine sequence, a growth factor sequence, a hormone sequence, or a toxin sequence.

20 [0029] In some embodiments of each or any of the above or below mentioned embodiments, the non-antibody sequence is an IL-8 sequence, an IL-21 sequence, an SDF-1 (alpha) sequence, a somatostatin sequence, a chlorotoxin sequence, a Pro-TxII sequence, or a ziconotide sequence.

[0030] In some embodiments of each or any of the above or below mentioned embodiments, the non-antibody sequence is any one of SEQ ID NOS: 475-481.

25 [0031] In some embodiments of each or any of the above or below mentioned embodiments, the non-antibody sequence replaces at least a portion of the ultralong CDR3.

30 [0032] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q).

35 [0033] In some embodiments of each or any of the above or below mentioned embodiments, the $X^1X^2X^3X^4X^5$ motif is TTVHQ (SEQ ID NO: 153), TSVHQ (SEQ ID NO: 154), SSVTQ (SEQ ID NO: 155), STVHQ (SEQ ID NO: 156), ATVRQ (SEQ ID NO: 157), TTVYQ (SEQ ID NO: 158), SPVHQ (SEQ ID NO: 159), ATVYQ (SEQ ID NO: 160), TAVYQ

(SEQ ID NO: 161), TNVHQ (SEQ ID NO: 162), ATVHQ (SEQ ID NO: 163), STVYQ (SEQ ID NO: 164), TIVHQ (SEQ ID NO: 165), AIVYQ (SEQ ID NO: 166), TTVFQ (SEQ ID NO: 167), AAVFQ (SEQ ID NO: 168), GTVHQ (SEQ ID NO: 169), ASVHQ (SEQ ID NO: 170), TAVFQ (SEQ ID NO: 171), ATVFQ (SEQ ID NO: 172), AAAHQ (SEQ ID NO: 173), VVVYQ (SEQ ID NO: 174), GTVFQ (SEQ ID NO: 175), TAVHQ (SEQ ID NO: 176), ITVHQ (SEQ ID NO: 177), ITAHQ (SEQ ID NO: 178), VTVHQ (SEQ ID NO: 179); AAVHQ (SEQ ID NO: 180), GTVYQ (SEQ ID NO: 181), TTVLQ (SEQ ID NO: 182), TTTHQ (SEQ ID NO: 183), or TTDYQ (SEQ ID NO: 184).

[0034] In some embodiments of each or any of the above or below mentioned
10 embodiments, the ultralong CDR3 comprises a $CX^1X^2X^3X^4X^5$ motif.

[0035] In some embodiments of each or any of the above or below mentioned
embodiments, the $CX^1X^2X^3X^4X^5$ motif is CTTVHQ (SEQ ID NO: 185), CTSVHQ (SEQ ID NO: 186), CSSVTQ (SEQ ID NO: 187), CSTVHQ (SEQ ID NO: 188), CATVRQ (SEQ ID NO: 189), CTTVYQ (SEQ ID NO: 190), CSPVHQ (SEQ ID NO: 191), CATVYQ (SEQ ID NO: 192), CTAVYQ (SEQ ID NO: 193), CTNVHQ (SEQ ID NO: 194), CATVHQ (SEQ ID NO: 195), CSTVYQ (SEQ ID NO: 196), CTIVHQ (SEQ ID NO: 197), CAIVYQ (SEQ ID NO: 198), CTTVFQ (SEQ ID NO: 199), CAAVFQ (SEQ ID NO: 200), CGTVHQ (SEQ ID NO: 201), CASVHQ (SEQ ID NO: 202), CTAVFQ (SEQ ID NO: 203), CATVFQ (SEQ ID NO: 204), CAAAHQ (SEQ ID NO: 205), CVVVYQ (SEQ ID NO: 206), CGTVFQ (SEQ ID NO: 207), CTAVHQ (SEQ ID NO: 208), CITVHQ (SEQ ID NO: 209), CITAHQ (SEQ ID NO: 210), CVTVHQ (SEQ ID NO: 211); CAAVHQ (SEQ ID NO: 212), CGTVYQ (SEQ ID NO: 213), CTTVLQ (SEQ ID NO: 214), CTTTHQ (SEQ ID NO: 215), or CTTDYQ (SEQ ID NO: 216).

[0036] In some embodiments of each or any of the above or below mentioned
25 embodiments, the ultralong CDR3 comprises a $(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[0037] In some embodiments of each or any of the above or below mentioned
embodiments, the $(X^aX^b)_z$ motif is CYTYNYEF (SEQ ID NO: 217), HYTYTYDF (SEQ ID NO: 218), HYTYTYEW (SEQ ID NO: 219), KHRITYEW (SEQ ID NO: 220), NYIYKYSF (SEQ ID NO: 221), PYIYTYQF (SEQ ID NO: 222), SFTYTYEW (SEQ ID NO: 223), SYIYIYQW (SEQ ID NO: 224), SYNYTYSW (SEQ ID NO: 225), SYSYSYEF (SEQ ID NO: 226), SYTYNYDF (SEQ ID NO: 227), SYTYNYEW (SEQ ID NO: 228), SYTYNYQF (SEQ ID NO: 229), SYVWTHNF (SEQ ID NO: 230), TYKYVYEW (SEQ ID NO: 231), TYTYTYEF (SEQ ID NO: 232), TYTYTYEW (SEQ ID NO: 233), VFTYTYEF (SEQ ID NO: 234), AYTIEW (SEQ ID NO: 235), DYIYTY (SEQ ID NO: 236), IHSYEF (SEQ ID NO: 237),

SFTYEF (SEQ ID NO: 238), SHSYEF (SEQ ID NO: 239), THTYEF (SEQ ID NO: 240), TWTYEF (SEQ ID NO: 241), TYNTEW (SEQ ID NO: 242), TYSYEF (SEQ ID NO: 243), TYSYEH (SEQ ID NO: 244), TYTYDF (SEQ ID NO: 245), TYTYEF (SEQ ID NO: 246), TYTYEW (SEQ ID NO: 247), ATEF (SEQ ID NO: 248), ATSF (SEQ ID NO: 249), ATSY (SEQ ID NO: 250), CYSF (SEQ ID NO: 251), DTTY (SEQ ID NO: 252), KYEH (SEQ ID NO: 253), KYEW (SEQ ID NO: 254), MYEF (SEQ ID NO: 255), NWIY (SEQ ID NO: 256), NYDY (SEQ ID NO: 257), NYQW (SEQ ID NO: 258), NYSF (SEQ ID NO: 259), PYEW (SEQ ID NO: 260), RYNW (SEQ ID NO: 261), RTTY (SEQ ID NO: 262), SYEF (SEQ ID NO: 263), SYEH (SEQ ID NO: 264), SYEW (SEQ ID NO: 265), SYKW (SEQ ID NO: 266), SYTY (SEQ ID NO: 267), TYDF (SEQ ID NO: 268), TYEF (SEQ ID NO: 269), TYEW (SEQ ID NO: 270), TYQW (SEQ ID NO: 271), TTTY (SEQ ID NO: 272), or VTEW (SEQ ID NO: 273).

[0038] In some embodiments of each or any of the above or below mentioned embodiments, the $(X^aX^b)_z$ motif is YXYXYX.

[0039] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5X_n$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), wherein X^5 is glutamine (Q), and wherein n is 27 – 54.

[0040] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises $X_n(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

[0041] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5X_n(X^aX^b)_z$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q), wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

[0042] In some embodiments of each or any of the above or below mentioned embodiments, the $X^1X^2X^3X^4X^5$ motif is TTVHQ (SEQ ID NO: 153) or TSVHQ (SEQ ID NO: 154), and wherein the $(X^aX^b)_z$ motif is YXYXYX.

[0043] In some embodiments of each or any of the above or below mentioned
 5 embodiments, the ultralong CDR3 comprises: a $CX^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is
 10 glutamine (Q); a cysteine motif selected from the group consisting of: $CX_{10}CX_5CX_5CXCX_7C$ (SEQ ID NO: 41), $CX_{10}CX_6CX_5CXCX_{15}C$ (SEQ ID NO: 42), $CX_{11}CXCX_5C$ (SEQ ID NO: 43), $CX_{11}CX_5CX_5CXCX_7C$ (SEQ ID NO: 44), $CX_{10}CX_6CX_5CXCX_{13}C$ (SEQ ID NO: 45), $CX_{10}CX_5CXCX_4CX_8C$ (SEQ ID NO: 46), $CX_{10}CX_6CX_6CXCX_7C$ (SEQ ID NO: 47), $CX_{10}CX_4CX_7CXCX_8C$ (SEQ ID NO: 48), $CX_{10}CX_4CX_7CXCX_7C$ (SEQ ID NO: 49),
 15 $CX_{13}CX_8CX_8C$ (SEQ ID NO: 50), $CX_{10}CX_6CX_5CXCX_7C$ (SEQ ID NO: 51), $CX_{10}CX_5CX_5C$ (SEQ ID NO: 52), $CX_{10}CX_5CX_6CXCX_7C$ (SEQ ID NO: 53), $CX_{10}CX_6CX_5CX_7CX_9C$ (SEQ ID NO: 54), $CX_9CX_7CX_5CXCX_7C$ (SEQ ID NO: 55), $CX_{10}CX_6CX_5CXCX_9C$ (SEQ ID NO: 56), $CX_{10}CXCX_4CX_5CX_{11}C$ (SEQ ID NO: 57), $CX_7CX_3CX_6CX_5CXCX_5CX_{10}C$ (SEQ ID NO: 58), $CX_{10}CXCX_4CX_5CXCX_2CX_3C$ (SEQ ID NO: 59), $CX_{16}CX_5CXC$ (SEQ ID NO: 60),
 20 $CX_6CX_4CXCX_4CX_5C$ (SEQ ID NO: 61), $CX_{11}CX_4CX_5CX_6CX_3C$ (SEQ ID NO: 62), $CX_8CX_2CX_6CX_5C$ (SEQ ID NO: 63), $CX_{10}CX_5CX_5CXCX_{10}C$ (SEQ ID NO: 64), $CX_{10}CXCX_6CX_4CXC$ (SEQ ID NO: 65), $CX_{10}CX_5CX_5CXCX_2C$ (SEQ ID NO: 66), $CX_{14}CX_2CX_3CXCXC$ (SEQ ID NO: 67), $CX_{15}CX_5CXC$ (SEQ ID NO: 68), $CX_4CX_6CX_9CX_2CX_{11}C$ (SEQ ID NO: 69), $CX_6CX_4CX_5CX_5CX_{12}C$ (SEQ ID NO: 70),
 25 $CX_7CX_3CXCXCX_4CX_5CX_9C$ (SEQ ID NO: 71), $CX_{10}CX_6CX_5C$ (SEQ ID NO: 72), $CX_7CX_3CX_5CX_5CX_9C$ (SEQ ID NO: 73), $CX_7CX_5CXCX_2C$ (SEQ ID NO: 74), $CX_{10}CXCX_6C$ (SEQ ID NO: 75), $CX_{10}CX_3CX_3CX_5CX_7CXCX_6C$ (SEQ ID NO: 76), $CX_{10}CX_4CX_5CX_{12}CX_2C$ (SEQ ID NO: 77), $CX_{12}CX_4CX_5CXCXCX_9CX_3C$ (SEQ ID NO: 78), $CX_{12}CX_4CX_5CX_{12}CX_2C$ (SEQ ID NO: 79), $CX_{10}CX_6CX_5CXCX_{11}C$ (SEQ ID NO: 80), $CX_{16}CX_5CXCXCX_{14}C$ (SEQ ID
 30 NO: 81), $CX_{10}CX_5CXCX_8CX_6C$ (SEQ ID NO: 82), $CX_{12}CX_4CX_5CX_8CX_2C$ (SEQ ID NO: 83), $CX_{12}CX_5CX_5CXCX_8C$ (SEQ ID NO: 84), $CX_{10}CX_6CX_5CXCX_4CXCX_9C$ (SEQ ID NO: 85), $CX_{11}CX_4CX_5CX_8CX_2C$ (SEQ ID NO: 86), $CX_{10}CX_6CX_5CX_8CX_2C$ (SEQ ID NO: 87), $CX_{10}CX_6CX_5CXCX_8C$ (SEQ ID NO: 88), $CX_{10}CX_6CX_5CXCX_3CX_8CX_2C$ (SEQ ID NO: 89), $CX_{10}CX_6CX_5CX_3CX_8C$ (SEQ ID NO: 90), $CX_{10}CX_6CX_5CXCX_2CX_6CX_5C$ (SEQ ID NO: 91),
 35 $CX_7CX_6CX_3CX_3CX_9C$ (SEQ ID NO: 92), $CX_9CX_8CX_5CX_6CX_5C$ (SEQ ID NO: 93), $CX_{10}CX_2CX_2CX_7CXCX_{11}CX_5C$ (SEQ ID NO: 94), and $CX_{10}CX_6CX_5CXCX_2CX_8CX_4C$ (SEQ

ID NO: 95), and a $(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[0044] In some embodiments of each or any of the above or below mentioned
 5 embodiments, the ultralong CDR3 comprises: a $CX^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is
 10 glutamine (Q); a cysteine motif selected from the group consisting of: wherein the cysteine motif is selected from the group consisting of: $CCX_3CXCX_3CX_2CCXCX_5CX_9CX_5CXC$ (SEQ ID NO: 96), $CX_6CX_2CX_5CX_4CCXCX_4CX_6CXC$ (SEQ ID NO: 97), $CX_7CXCX_5CX_4CCCX_4CX_6CXC$ (SEQ ID NO: 98), $CX_9CX_3CXCX_2CXCCCX_6CX_4C$ (SEQ ID NO: 99), $CX_5CX_3CXCX_4CX_4CCX_{10}CX_2CC$ (SEQ ID NO: 100),
 15 $CX_5CXCX_1CXCX_3CCX_3CX_4CX_{10}C$ (SEQ ID NO: 101), $CX_9CCCX_3CX_4CCCX_5CX_6C$ (SEQ ID NO: 102), $CCX_8CX_5CX_4CX_3CX_4CCXCX_1C$ (SEQ ID NO: 103), $CCX_6CCX_5CCCX_4CX_4CX_{12}C$ (SEQ ID NO: 104), $CX_6CX_2CX_3CCCX_4CX_5CX_3CX_3C$ (SEQ ID NO: 105), $CX_3CX_5CX_6CX_4CCXCX_5CX_4CXC$ (SEQ ID NO: 106), $CX_4CX_4CCX_4CX_4CXCX_{11}CX_2CXC$ (SEQ ID NO: 107), $CX_5CX_2CCX_5CX_4CCX_3CCX_7C$ (SEQ ID NO: 108),
 20 $CX_5CX_5CX_3CX_2CXCCX_4CX_7CXC$ (SEQ ID NO: 109), $CX_3CX_7CX_3CX_4CCXCX_2CX_5CX_2C$ (SEQ ID NO: 110), $CX_9CX_3CXCX_4CCX_5CCCX_6C$ (SEQ ID NO: 111), $CX_9CX_3CXCX_2CXCCX_6CX_3CX_3C$ (SEQ ID NO: 112), $CX_8CCXCX_3CCX_3CXCX_3CX_4C$ (SEQ ID NO: 113), $CX_9CCX_4CX_2CXCCXCX_4CX_3C$ (SEQ ID NO: 114), $CX_{10}CXCX_3CX_2CXCCX_4CX_5CXC$ (SEQ ID NO: 115), $CX_9CXCX_3CX_2CXCCX_4CX_5CXC$ (SEQ ID NO: 116), $CX_6CCXCX_5CX_4CCXCX_5CX_2C$ (SEQ ID NO: 117),
 25 $CX_6CCXCX_3CXCCX_3CX_4CC$ (SEQ ID NO: 118), $CX_6CCXCX_3CXCX_2CXCX_4CX_8C$ (SEQ ID NO: 119), $CX_4CX_2CCX_3CXCX_4CCX_2CX_3C$ (SEQ ID NO: 120), $CX_3CX_5CX_3CCCX_4CX_9C$ (SEQ ID NO: 121), $CCX_9CX_3CXCCX_3CX_5C$ (SEQ ID NO: 122), $CX_9CX_2CX_3CX_4CCX_4CX_5C$ (SEQ ID NO: 123), $CX_9CX_7CX_4CCXCX_7CX_3C$ (SEQ ID NO: 124), $CX_9CX_3CCCX_{10}CX_2CX_3C$ (SEQ ID NO: 125), $CX_3CX_5CX_5CX_4CCX_{10}CX_6C$ (SEQ ID NO: 126), $CX_9CX_5CX_4CCXCX_5CX_4C$ (SEQ ID NO: 127), $CX_7CXCX_6CX_4CCCX_{10}C$ (SEQ ID NO: 128), $CX_8CX_2CX_4CCX_4CX_3CX_3C$ (SEQ ID NO: 129), $CX_7CX_5CXCX_4CCX_7CX_4C$ (SEQ ID NO: 130), $CX_{11}CX_3CX_4CCCX_8CX_2C$ (SEQ ID NO: 131), $CX_2CX_3CX_4CCX_4CX_5CX_{15}C$ (SEQ ID NO: 132), $CX_9CX_5CX_4CCX_7C$ (SEQ ID NO: 133), $CX_9CX_7CX_3CX_2CX_6C$ (SEQ ID NO: 134),
 30 $CX_9CX_5CX_4CCX_{14}C$ (SEQ ID NO: 135), $CX_9CX_5CX_4CCX_8C$ (SEQ ID NO: 136), $CX_9CX_6CX_4CCXC$ (SEQ ID NO: 137), $CX_5CCX_7CX_4CX_{12}$ (SEQ ID NO: 138),

CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140),
 CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142),
 CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144),
 CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C
 5 (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149),
 CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152); and a
 (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected
 from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine
 (H), and wherein z is 1 - 4.

10 [0045] In some embodiments of each or any of the above or below mentioned
 embodiments, the ultralong CDR3 comprises an additional sequence that is a linker.

[0046] In some embodiments of each or any of the above or below mentioned
 embodiments, the linker is linked to a C-terminus, a N-terminus, or both C-terminus and N-
 terminus of the non-antibody sequence.

15 [0047] In some embodiments of each or any of the above or below mentioned
 embodiments, the ultralong CDR3 is a ruminant CDR3.

[0048] In some embodiments of each or any of the above or below mentioned
 embodiments, the ruminant is a cow.

20 [0049] In some embodiments of each or any of the above or below mentioned
 embodiments, the humanized antibody or binding fragment thereof comprises a human
 heavy chain variable region framework sequence.

[0050] In some embodiments of each or any of the above or below mentioned
 embodiments, the humanized antibody or binding fragment thereof comprises a human
 heavy chain germline sequence or is derived from a human heavy chain germline
 25 sequence.

[0051] In some embodiments of each or any of the above or below mentioned
 embodiments, the humanized antibody or binding fragment thereof comprises a human
 heavy chain germline sequence that is a VH4 family member germline sequence.

30 [0052] In some embodiments of each or any of the above or below mentioned
 embodiments, the humanized antibody or binding fragment thereof comprises a VH4-34
 germline sequence.

[0053] In some embodiments of each or any of the above or below mentioned
 embodiments, the humanized antibody or binding fragment thereof comprises a human
 heavy chain germline sequence that is derived from the VH4-34 germline sequence.

[0054] In some embodiments of each or any of the above or below mentioned embodiments, the VH4-34 germline sequence comprises a CDR1 comprising a Gly31Asp and Tyr32Lys substitution based on Kabat numbering.

5 [0055] In some embodiments of each or any of the above or below mentioned embodiments, the VH4-34 germline sequence comprises a CDR2 comprising a Glu50Ser substitution based on Kabat numbering.

[0056] In some embodiments of each or any of the above or below mentioned embodiments, the VH4-34 germline sequence comprises a CDR1 comprising Ala23Thr, Val24Ala, Tyr25Ser, Gly27Phe, Phe29Leu, Gly31Asp, Tyr32Lys, Tyr33Ala, Trp34Val, and
10 Ser35Gly substitutions, and a CDR2 comprising Ile48Leu, Glu50Ser, Asn52Asp, His53Thr, Ser54Gly, Ser56Asn, and Asn58Gly substitutions, based on Kabat numbering.

[0057] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469.

15 [0058] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 470.

[0059] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469; and
20 the amino acid sequence set forth in SEQ ID NO: 470.

[0060] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469; the
25 amino acid sequence set forth in SEQ ID NO: 470; and a peptide sequence, wherein the peptide sequence is positioned between SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469, and SEQ ID NO: 470.

[0061] In some embodiments of each or any of the above or below mentioned embodiments, the peptide sequence is a bovine sequence, a non-bovine sequence, an
30 antibody sequence, or a non-antibody sequence.

[0062] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibody or binding fragment thereof comprises a human light chain variable region framework sequence.

[0063] In some embodiments of each or any of the above or below mentioned
35 embodiments, the humanized antibody or binding fragment thereof comprises a light chain variable region sequence that is a lambda light chain variable region sequence.

[0064] In some embodiments of each or any of the above or below mentioned embodiments, the lambda light chain variable region sequence is a bovine lambda light chain variable region sequence or derived from a bovine lambda light chain variable region sequence.

5 [0065] In some embodiments of each or any of the above or below mentioned embodiments, the lambda light chain variable region sequence is a human lambda light chain variable region sequence or derived from a human lambda light chain variable region sequence.

10 [0066] In some embodiments of each or any of the above or below mentioned embodiments, the human lambda light chain variable region sequence is a VL1-51 germline sequence.

[0067] In some embodiments of each or any of the above or below mentioned embodiments, the human lambda light chain variable region sequence is derived from a VL1-51 germline sequence.

15 [0068] In some embodiments of each or any of the above or below mentioned embodiments, the VL1-51 germline sequence comprises a CDR1 comprising Ile29Val and Asn32Gly substitution based on Kabat numbering.

[0069] In some embodiments of each or any of the above or below mentioned embodiments, the VL1-51 germline sequence comprises a CDR2 comprising a substitution
20 of DNN to GDT.

[0070] In some embodiments of each or any of the above or below mentioned embodiments, the VL1-51 germline sequence comprises a CDR2 comprising a substitution of DNNKRP (SEQ ID NO: 471) to GDTSRA (SEQ ID NO: 472).

25 [0071] In some embodiments of each or any of the above or below mentioned embodiments, the VL1-51 germline sequence comprises a S2A, T5N, P8S, A12G, A13S, and P14L substitution based on Kabat numbering.

[0072] In some embodiments of each or any of the above or below mentioned embodiments, the VL1-51 germline sequence comprises a S2A, T5N, P8S, A12G, A13S, and P14L substitution based on Kabat numbering, and a CDR2 comprising a substitution of
30 DNN to GDT.

[0073] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibody or binding fragment thereof comprising the amino acid sequence set forth in SEQ ID NO: 440, 441, 442, 443, or 444.

35 [0074] The present disclosure also provides polynucleotides encoding the humanized antibody or binding fragment thereof disclosed herein.

[0075] The present disclosure also provides polynucleotides encoding a humanized antibody or binding fragment thereof that comprises an ultralong CDR3.

[0076] The present disclosure also provides polynucleotides encoding a humanized variable region that comprises an ultralong CDR3.

5 [0077] The present disclosure also provides vectors that comprise the polynucleotides disclosed herein.

[0078] The present disclosure also provides host cells comprising the vectors disclosed herein.

10 [0079] The present disclosure also provides a nucleic acid library comprising a plurality of polynucleotides comprising sequences coding for humanized antibodies or binding fragments thereof, wherein the antibodies or binding fragments thereof comprise an ultralong CDR3.

[0080] The present disclosure also provides a library of humanized antibodies or binding fragments thereof, wherein the antibodies or binding fragments thereof comprise
15 an ultralong CDR3.

[0081] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 is 35 amino acids in length or longer, 40 amino acids in length or longer, 45 amino acids in length or longer, 50 amino acids in length or longer, 55 amino acids in length or longer, or 60 amino acids in length or longer.

20 [0082] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 is 35 amino acids in length or longer.

[0083] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises 3 or more cysteine residues, 4 or more cysteine residues, 5 or more cysteine residues, 6 or more cysteine residues, 7 or more
25 cysteine residues, 8 or more cysteine residues, 9 or more cysteine residues, 10 or more cysteine residues, 11 or more cysteine residues, or 12 or more cysteine residues.

[0084] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises 3 or more cysteine residues.

[0085] In some embodiments of each or any of the above or below mentioned
30 embodiments, the antibodies or binding fragments thereof comprise a cysteine motif.

[0086] In some embodiments of each or any of the above or below mentioned embodiments, the cysteine motif is selected from the group consisting of:
CX₁₀CX₅CX₅CXCX₇C (SEQ ID NO: 41), CX₁₀CX₆CX₅CXCX₁₅C (SEQ ID NO: 42),
CX₁₁CXCX₅C (SEQ ID NO: 43), CX₁₁CX₅CX₅CXCX₇C (SEQ ID NO: 44),
35 CX₁₀CX₆CX₅CXCX₁₃C (SEQ ID NO: 45), CX₁₀CX₅CXCX₄CX₈C (SEQ ID NO: 46),
CX₁₀CX₆CX₆CXCX₇C (SEQ ID NO: 47), CX₁₀CX₄CX₇CXCX₈C (SEQ ID NO: 48),

CX₁₀CX₄CX₇CXCX₇C (SEQ ID NO: 49), CX₁₃CX₈CX₈C (SEQ ID NO: 50),
 CX₁₀CX₆CX₅CXCX₇C (SEQ ID NO: 51), CX₁₀CX₅CX₅C (SEQ ID NO: 52),
 CX₁₀CX₅CX₆CXCX₇C (SEQ ID NO: 53), CX₁₀CX₆CX₅CX₇CX₉C (SEQ ID NO: 54),
 CX₉CX₇CX₅CXCX₇C (SEQ ID NO: 55), CX₁₀CX₆CX₅CXCX₉C (SEQ ID NO: 56),
 5 CX₁₀CXCX₄CX₅CX₁₁C (SEQ ID NO: 57), CX₇CX₃CX₆CX₅CXCX₅CX₁₀C (SEQ ID NO: 58),
 CX₁₀CXCX₄CX₅CXCX₂CX₃C (SEQ ID NO: 59), CX₁₆CX₅CXC (SEQ ID NO: 60),
 CX₆CX₄CXCX₄CX₅C (SEQ ID NO: 61), CX₁₁CX₄CX₅CX₆CX₃C (SEQ ID NO: 62),
 CX₈CX₂CX₆CX₅C (SEQ ID NO: 63), CX₁₀CX₅CX₅CXCX₁₀C (SEQ ID NO: 64),
 CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65), CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66),
 10 CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67), CX₁₅CX₅CXC (SEQ ID NO: 68),
 CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69), CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70),
 CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72),
 CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74), CX₁₀CXCX₆C
 (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C
 15 (SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C
 (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCX₁₄C (SEQ ID
 NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83),
 CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85),
 CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87),
 20 CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89),
 CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91),
 CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93),
 CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ
 ID NO: 95).

25 [0087] In some embodiments of each or any of the above or below mentioned
 embodiments, the cysteine motif is selected from the group consisting of:
 CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC
 (SEQ ID NO: 97), CX₇CXCX₅CX₄CCCX₄CX₆CXC (SEQ ID NO: 98),
 CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID
 30 NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101),
 CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102), CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID
 NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104),
 CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC
 (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107),
 35 CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCCX₄CX₇CXC (SEQ
 ID NO: 109), CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C (SEQ ID NO: 110),

CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113), CX₉CCX₄CX₂CXCCXCX₄CX₃C (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116),
 5 CX₆CCXCX₅CX₄CCXCX₅CX₂C (SEQ ID NO: 117), CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118), CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO: 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135),
 15 CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145),
 20 CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152).

[0088] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises 2 to 6 disulfide bonds.

25 [0089] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises SEQ ID NO: 40 or a derivative thereof.

[0090] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises amino acid residues 3-6 of any of one SEQ ID NO: 1-4.

30 [0091] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a non-human DH or a derivative thereof.

[0092] In some embodiments of each or any of the above or below mentioned embodiments, the non-human DH is SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, or SEQ ID NO: 12.

35 [0093] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a JH sequence or a derivative thereof.

[0094] In some embodiments of each or any of the above or below mentioned embodiments, the JH sequence is SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, or SEQ ID NO: 17.

5 [0095] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises: a non-human VH sequence or a derivative thereof; a non-human DH sequence or a derivative thereof; and/or a JH sequence or derivative thereof.

10 [0096] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises an additional amino acid sequence comprising two to six amino acid residues or more positioned between the VH sequence and the DH sequence.

[0097] In some embodiments of each or any of the above or below mentioned embodiments, the additional amino acid sequence is selected from the group consisting of: IR, IF, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20 or SEQ ID NO: 21.

15 [0098] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a sequence derived from or based on SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, or SEQ ID NO: 28.

20 [0099] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a non-bovine sequence or a non-antibody sequence.

[00100] In some embodiments of each or any of the above or below mentioned embodiments, the non-antibody sequence is a synthetic sequence.

25 [00101] In some embodiments of each or any of the above or below mentioned embodiments, the non-antibody sequence is a cytokine sequence, a lymphokine sequence, a chemokine sequence, a growth factor sequence, a hormone sequence, or a toxin sequence.

30 [00102] In some embodiments of each or any of the above or below mentioned embodiments, the non-antibody sequence is an IL-8 sequence, an IL-21 sequence, an SDF-1 (alpha) sequence, a somatostatin sequence, a chlorotoxin sequence, a Pro-TxII sequence, or a ziconotide sequence.

[00103] In some embodiments of each or any of the above or below mentioned embodiments, the non-antibody sequence is any one of SEQ ID NOS: 475-481.

35 [00104] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S),

threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q).

5 [00105] In some embodiments of each or any of the above or below mentioned embodiments, the $X^1X^2X^3X^4X^5$ motif is TTVHQ (SEQ ID NO: 153), TSVHQ (SEQ ID NO: 154), SSVTQ (SEQ ID NO: 155), STVHQ (SEQ ID NO: 156), ATVRQ (SEQ ID NO: 157), TTVYQ (SEQ ID NO: 158), SPVHQ (SEQ ID NO: 159), ATVYQ (SEQ ID NO: 160), TAVYQ (SEQ ID NO: 161), TNVHQ (SEQ ID NO: 162), ATVHQ (SEQ ID NO: 163),
 10 STVYQ (SEQ ID NO: 164), TIVHQ (SEQ ID NO: 165), AIVYQ (SEQ ID NO: 166), TTVFQ (SEQ ID NO: 167), AAVFQ (SEQ ID NO: 168), GTVHQ (SEQ ID NO: 169), ASVHQ (SEQ ID NO: 170), TAVFQ (SEQ ID NO: 171), ATVFQ (SEQ ID NO: 172), AAAHQ (SEQ ID NO: 173), VVVYQ (SEQ ID NO: 174), GTVFQ (SEQ ID NO: 175), TAVHQ (SEQ ID NO: 176), ITVHQ (SEQ ID NO: 177), ITAHQ (SEQ ID NO: 178), VTVHQ (SEQ ID NO: 179); AAVHQ
 15 (SEQ ID NO: 180), GTVYQ (SEQ ID NO: 181), TTVLQ (SEQ ID NO: 182), TTTHQ (SEQ ID NO: 183), or TTDYQ (SEQ ID NO: 184).

[00106] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a $CX^1X^2X^3X^4X^5$ motif.

[00107] In some embodiments of each or any of the above or below
 20 mentioned embodiments, the $CX^1X^2X^3X^4X^5$ motif is CTTVHQ (SEQ ID NO: 185), CTSVHQ (SEQ ID NO: 186), CSSVTQ (SEQ ID NO: 187), CSTVHQ (SEQ ID NO: 188), CATVRQ (SEQ ID NO: 189), CTTVYQ (SEQ ID NO: 190), CSPVHQ (SEQ ID NO: 191), CATVYQ (SEQ ID NO: 192), CTAVYQ (SEQ ID NO: 193), CTNVHQ (SEQ ID NO: 194), CATVHQ (SEQ ID NO: 195), CSTVYQ (SEQ ID NO: 196), CTIVHQ (SEQ ID NO: 197), CAIVYQ
 25 (SEQ ID NO: 198), CTTVFQ (SEQ ID NO: 199), CAAVFQ (SEQ ID NO: 200), CGTVHQ (SEQ ID NO: 201), CASVHQ (SEQ ID NO: 202), CTAVFQ (SEQ ID NO: 203), CATVFQ (SEQ ID NO: 204), CAAAHQ (SEQ ID NO: 205), CVVVYQ (SEQ ID NO: 206), CGTVFQ (SEQ ID NO: 207), CTAVHQ (SEQ ID NO: 208), CITVHQ (SEQ ID NO: 209), CITAHQ (SEQ ID NO: 210), CVTVHQ (SEQ ID NO: 211); CAAVHQ (SEQ ID NO: 212), CGTVYQ
 30 (SEQ ID NO: 213), CTTVLQ (SEQ ID NO: 214), CTTTHQ (SEQ ID NO: 215), or CTTDYQ (SEQ ID NO: 216).

[00108] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a $(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of:
 35 tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[00109] In some embodiments of each or any of the above or below mentioned embodiments, the $(X^aX^b)_z$ motif is CYTYNYEF (SEQ ID NO: 217), HYTYTYDF (SEQ ID NO: 218), HYTYTYEW (SEQ ID NO: 219), KHRYTYEW (SEQ ID NO: 220), NYIYKYSF (SEQ ID NO: 221), PYIYTYQF (SEQ ID NO: 222), SFTYTYEW (SEQ ID NO: 223), SYIYIYQW (SEQ ID NO: 224), SYNYTYSW (SEQ ID NO: 225), SYSYSYEF (SEQ ID NO: 226), SYTYNYDF (SEQ ID NO: 227), SYTYNYEW (SEQ ID NO: 228), SYTYNYQF (SEQ ID NO: 229), SYVWTHNF (SEQ ID NO: 230), TYKYVYEW (SEQ ID NO: 231), TYTYTYEF (SEQ ID NO: 232), TYTYTYEW (SEQ ID NO: 233), VFTYTYEF (SEQ ID NO: 234), AYTIEW (SEQ ID NO: 235), DYIYTY (SEQ ID NO: 236), IHSYEF (SEQ ID NO: 237), SFTYEF (SEQ ID NO: 238), SHSYEF (SEQ ID NO: 239), THTYEF (SEQ ID NO: 240), TWTYEF (SEQ ID NO: 241), TYNIEW (SEQ ID NO: 242), TYSYEF (SEQ ID NO: 243), TYSYEH (SEQ ID NO: 244), TYTYDF (SEQ ID NO: 245), TYTYEF (SEQ ID NO: 246), TYTYEW (SEQ ID NO: 247), AYEF (SEQ ID NO: 248), AYSF (SEQ ID NO: 249), AYSY (SEQ ID NO: 250), CYSF (SEQ ID NO: 251), DYTY (SEQ ID NO: 252), KYEH (SEQ ID NO: 253), KYEW (SEQ ID NO: 254), MYEF (SEQ ID NO: 255), NWIY (SEQ ID NO: 256), NYDY (SEQ ID NO: 257), NYQW (SEQ ID NO: 258), NYSF (SEQ ID NO: 259), PYEW (SEQ ID NO: 260), RYNW (SEQ ID NO: 261), RYTY (SEQ ID NO: 262), SYEF (SEQ ID NO: 263), SYEH (SEQ ID NO: 264), SYEW (SEQ ID NO: 265), SYKW (SEQ ID NO: 266), SYTY (SEQ ID NO: 267), TYDF (SEQ ID NO: 268), TYEF (SEQ ID NO: 269), TYEW (SEQ ID NO: 270), TYQW (SEQ ID NO: 271), TYTY (SEQ ID NO: 272), or VYEW (SEQ ID NO: 273).

[00110] In some embodiments of each or any of the above or below mentioned embodiments, the $(X^aX^b)_z$ motif is YXYXYX.

[00111] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5X_n$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q), and wherein n is 27 – 54.

[00112] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises $X_n(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

[00113] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5X_n(X^aX^b)_z$ motif, wherein X_1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X_2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X_3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X_4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), wherein X_5 is glutamine (Q), X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

[00114] In some embodiments of each or any of the above or below mentioned embodiments, the $X^1X^2X^3X^4X^5$ motif is TTVHQ (SEQ ID NO: 153) or TSVHQ (SEQ ID NO: 154), and wherein the $(X^aX^b)_z$ motif is YXYXYX.

[00115] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises: a $CX^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q), a cysteine motif selected from the group consisting of: $CX_{10}CX_5CX_5CXCX_7C$ (SEQ ID NO: 41), $CX_{10}CX_6CX_5CXCX_{15}C$ (SEQ ID NO: 42), $CX_{11}CXCX_5C$ (SEQ ID NO: 43), $CX_{11}CX_5CX_5CXCX_7C$ (SEQ ID NO: 44), $CX_{10}CX_6CX_5CXCX_{13}C$ (SEQ ID NO: 45), $CX_{10}CX_5CXCX_4CX_8C$ (SEQ ID NO: 46), $CX_{10}CX_6CX_6CXCX_7C$ (SEQ ID NO: 47), $CX_{10}CX_4CX_7CXCX_8C$ (SEQ ID NO: 48), $CX_{10}CX_4CX_7CXCX_7C$ (SEQ ID NO: 49), $CX_{13}CX_8CX_8C$ (SEQ ID NO: 50), $CX_{10}CX_6CX_5CXCX_7C$ (SEQ ID NO: 51), $CX_{10}CX_5CX_5C$ (SEQ ID NO: 52), $CX_{10}CX_5CX_6CXCX_7C$ (SEQ ID NO: 53), $CX_{10}CX_6CX_5CX_7CX_9C$ (SEQ ID NO: 54), $CX_9CX_7CX_5CXCX_7C$ (SEQ ID NO: 55), $CX_{10}CX_6CX_5CXCX_9C$ (SEQ ID NO: 56), $CX_{10}CXCX_4CX_5CX_{11}C$ (SEQ ID NO: 57), $CX_7CX_3CX_6CX_5CXCX_5CX_{10}C$ (SEQ ID NO: 58), $CX_{10}CXCX_4CX_5CXCX_2CX_3C$ (SEQ ID NO: 59), $CX_{16}CX_5CXC$ (SEQ ID NO: 60), $CX_6CX_4CXCX_4CX_5C$ (SEQ ID NO: 61), $CX_{11}CX_4CX_5CX_6CX_3C$ (SEQ ID NO: 62), $CX_8CX_2CX_6CX_5C$ (SEQ ID NO: 63), $CX_{10}CX_5CX_5CXCX_{10}C$ (SEQ ID NO: 64), $CX_{10}CXCX_6CX_4CXC$ (SEQ ID NO: 65), $CX_{10}CX_5CX_5CXCX_2C$ (SEQ ID NO: 66), $CX_{14}CX_2CX_3CXCXC$ (SEQ ID NO: 67), $CX_{15}CX_5CXC$ (SEQ ID NO: 68), $CX_4CX_6CX_9CX_2CX_{11}C$ (SEQ ID NO: 69), $CX_6CX_4CX_5CX_5CX_{12}C$ (SEQ ID NO: 70), $CX_7CX_3CXCXCX_4CX_5CX_9C$ (SEQ ID NO: 71), $CX_{10}CX_6CX_5C$ (SEQ ID NO: 72), $CX_7CX_3CX_5CX_5CX_9C$ (SEQ ID NO: 73), $CX_7CX_5CXCX_2C$ (SEQ ID NO: 74), $CX_{10}CXCX_6C$ (SEQ ID NO: 75), $CX_{10}CX_3CX_3CX_5CX_7CXCX_6C$ (SEQ ID NO: 76), $CX_{10}CX_4CX_5CX_{12}CX_2C$

(SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCX₁₄C (SEQ ID NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83), CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85),
 5 CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87), CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89), CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91), CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93), CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ
 10 ID NO: 95); and a (X^aX^b)_z motif, X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[00116] In some embodiments of each or any of the above or below mentioned embodiments, the ultralong CDR3 comprises: a CX¹X²X³X⁴X⁵ motif, wherein X¹ is threonine
 15 (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X² is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X³ is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X⁴ is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X⁵ is glutamine (Q); a cysteine motif selected from the group consisting of: wherein the cysteine
 20 motif is selected from the group consisting of: CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCCX₄CX₆CXC (SEQ ID NO: 98), CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID
 25 NO: 102), CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109), CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C
 30 (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113), CX₉CCX₄CX₂CXCCXCX₄CX₃C (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116), CX₆CCXCX₅CX₄CCXCX₅CX₂C (SEQ ID NO: 117),
 35 CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118), CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C

(SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO: 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128),
 5 CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138),
 10 CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149),
 15 CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152); andba (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[00117] In some embodiments of each or any of the above or below mentioned
 20 embodiments, the ultralong CDR3 comprises an additional sequence that is a linker.

[00118] In some embodiments of each or any of the above or below mentioned embodiments, the linker is linked to a C-terminus, a N-terminus, or both C-terminus and N-terminus of the non-antibody sequence.

[00119] In some embodiments of each or any of the above or below mentioned
 25 embodiments, the ultralong CDR3 is a ruminant CDR3.

[00120] In some embodiments of each or any of the above or below mentioned embodiments, the ruminant is a cow.

[00121] In some embodiments of each or any of the above or below mentioned
 30 embodiments, the humanized antibody or binding fragment thereof comprises a human heavy chain variable region framework sequence.

[00122] In some embodiments of each or any of the above or below mentioned
 35 embodiments, the humanized antibody or binding fragment thereof comprises a human heavy chain germline sequence or is derived from a human heavy chain germline sequence.

[00123] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibody or binding fragment thereof comprises a human heavy chain germline sequence that is a VH4 family member germline sequence.

5 [00124] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibody or binding fragment thereof comprises a VH4-34 germline sequence.

[00125] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibody or binding fragment thereof comprises a human heavy chain germline sequence that is derived from VH4-34.

10 [00126] In some embodiments of each or any of the above or below mentioned embodiments, the VH4-34 germline sequence comprises a CDR1 comprising a Gly31Asp and Tyr32Lys substitution based on Kabat numbering.

[00127] In some embodiments of each or any of the above or below mentioned embodiments, the VH4-34 germline sequence comprises a CDR2 comprising a Glu50Ser
15 substitution based on Kabat numbering.

[00128] In some embodiments of each or any of the above or below mentioned embodiments, the VH4-34 germline sequence comprises a CDR1 comprising Ala23Thr, Val24Ala, Tyr25Ser, Gly27Phe, Phe29Leu, Gly31Asp, Tyr32Lys, Tyr33Ala, Trp34Val, and Ser35Gly substitutions, and a CDR2 comprising Ile48Leu, Glu50Ser, Asn52Asp, His53Thr,
20 Ser54Gly, Ser56Asn, and Asn58Gly substitutions, based on Kabat numbering.

[00129] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 470.

[00130] In some embodiments of each or any of the above or below mentioned
25 embodiments, the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469; and the amino acid sequence set forth in SEQ ID NO: 470.

[00131] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibody or binding fragment thereof comprises the amino
30 acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469; the amino acid sequence set forth in SEQ ID NO: 470; and a peptide sequence, wherein the peptide sequence is positioned between SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469, and SEQ ID NO: 470.

[00132] In some embodiments of each or any of the above or below mentioned
35 embodiments, the peptide sequence is a bovine sequence, a non-bovine sequence, an antibody sequence, or a non-antibody sequence.

[00133] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibody or binding fragment thereof comprises a lambda light chain variable region sequence.

5 [00134] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibody or binding fragment thereof comprises a light chain variable region sequence that is a lambda light chain variable region sequence.

[00135] In some embodiments of each or any of the above or below mentioned embodiments, the lambda light chain variable region sequence is a bovine lambda light chain variable region sequence.

10 [00136] In some embodiments of each or any of the above or below mentioned embodiments, the lambda light chain variable region sequence is a human lambda light chain variable region sequence.

[00137] In some embodiments of each or any of the above or below mentioned embodiments, the human lambda light chain variable region sequence is VL1-51.

15 [00138] In some embodiments of each or any of the above or below mentioned embodiments, the human lambda light chain variable region sequence is derived from VL1-51.

[00139] In some embodiments of each or any of the above or below mentioned embodiments, the VL1-51 germline sequence comprises a CDR1 comprising Ile29Val and
20 Asn32Gly substitution based on Kabat numbering.

[00140] In some embodiments of each or any of the above or below mentioned embodiments, the VL1-51 germline sequence comprises a CDR2 comprising a substitution of DNN to GDT.

[00141] In some embodiments of each or any of the above or below mentioned
25 embodiments, the VL1-51 germline sequence comprises a CDR2 comprising a substitution of DNNKRP (SEQ ID NO: 471) to GDTSRA (SEQ ID NO: 472).

[00142] In some embodiments of each or any of the above or below mentioned embodiments, the VL1-51 germline sequence comprises a S2A, T5N, P8S, A12G, A13S, and P14L substitution based on Kabat numbering.

30 [00143] In some embodiments of each or any of the above or below mentioned embodiments, the VL1-51 germline sequence comprises a S2A, T5N, P8S, A12G, A13S, and P14L substitution based on Kabat numbering, and a CDR2 comprising a substitution of DNN to GDT.

[00144] In some embodiments of each or any of the above or below mentioned
35 embodiments, the humanized antibody or binding fragment thereof comprising the amino acid sequence set forth in SEQ ID NO: 440, 441, 442, 443, or 444.

[00145] In some embodiments of each or any of the above or below mentioned embodiments, the humanized antibodies or binding fragments thereof are present in a spatially addressed format.

5 [00146] The present disclosure also provides a method of humanizing an antibody variable region comprising the step of genetically combining a nucleic acid sequence encoding a non-human ultralong CDR3 with a nucleic acid sequence encoding a human variable region framework (FR) sequence.

10 [00147] The present disclosure also provides a method of generating a library of humanized antibodies that comprises an ultralong CDR3, the method comprising: combining a nucleic acid sequence encoding an ultralong CDR3 with a nucleic acid sequence encoding a human variable region framework (FR) sequence to produce nucleic acids encoding for humanized antibodies that comprises an ultralong CDR3; and expressing the nucleic acids encoding for humanized antibodies that comprises an ultralong CDR3 to generate a library of humanized antibodies that comprises an ultralong
15 CDR3.

[00148] The present disclosure also provides a method of generating a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and which comprises a non-antibody sequence, the method comprising: combining a nucleic acid sequence encoding an ultralong CDR3, a nucleic acid sequence encoding a human
20 variable region framework (FR) sequence, and a nucleic acid sequence encoding a non-antibody sequence to produce nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a non-antibody sequence, and expressing the nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a non-antibody sequence to generate a library of
25 humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a non-antibody sequence. In some embodiments, the ultralong CDR3 comprises a bovine, a non-bovine, an antibody, or a non-antibody sequence.

[00149] The present disclosure also provides a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 which comprises a non-bovine or
30 a non-antibody sequence.

[00150] The present disclosure also provides a method of generating a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 which comprises a cysteine motif, the method comprising: combining a human variable region framework (FR) sequence, and a nucleic acid sequence encoding an ultralong CDR3 and a
35 cysteine motif; introducing one or more nucleotide changes to the nucleic acid sequence encoding one or more amino acid residues that are positioned between one or more

cysteine residues in the cysteine motif for nucleotides encoding different amino acid residues to produce nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a cysteine motif with one or more nucleotide changes introduced between one or more cysteine residues in the cysteine domain; and
 5 expressing the nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a cysteine motif with one or more nucleotide changes introduced between one or more cysteine residues in the cysteine domain to generate a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a cysteine motif with one or more amino acid changes introduced between one or
 10 more cysteine residues in the cysteine domain.

[00151] The present disclosure also provides a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 which comprises a cysteine motif, wherein the antibodies or binding fragments comprise one or more substitutions of amino acid residues that are positioned between cysteine residues in the cysteine motif.

15 [00152] The present disclosure also provides a method of generating a library of humanized antibodies or binding fragments thereof comprising a bovine ultralong CDR3, the method comprising: combining a nucleic acid sequence encoding a human variable region framework (FR) sequence and a nucleic acid encoding a bovine ultralong CDR3, and expressing the nucleic acids encoding a human variable region framework (FR)
 20 sequence and a nucleic acid encoding a bovine ultralong CDR3 to generate a library of humanized antibodies or binding fragments thereof comprising a bovine ultralong CDR3.

[00153] The present disclosure provides humanized antibodies, including antibodies comprising a CDR3 scaffold, methods of making same, and uses thereof.

25 [00154] The present disclosure provides a humanized antibody or binding fragment thereof comprising a CDR3 scaffold.

[00155] In some embodiments, the CDR3 scaffold comprises one or more human variable region framework sequences.

[00156] The present disclosure also provides a library of humanized antibodies or binding fragments thereof comprising a bovine ultralong CDR3.

30 [00157] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises a $X^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H),
 35 threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q).

[00158] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold is 35 amino acids in length or longer, 40 amino acids in length or longer, 45 amino acids in length or longer, 50 amino acids in length or longer, 55 amino acids in length or longer, or 60 amino acids in length or longer.

5 [00159] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold is 35 amino acids in length or longer.

[00160] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises 3 or more cysteine residues, 4 or more cysteine residues, 5 or more cysteine residues, 6 or more cysteine residues, 7 or more
10 cysteine residues, 8 or more cysteine residues, 9 or more cysteine residues, 10 or more cysteine residues, 11 or more cysteine residues, or 12 or more cysteine residues.

[00161] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises 3 or more cysteine residues.

[00162] In some embodiments of each or any of the above or below mentioned
15 embodiments, the CDR3 scaffold comprises a cysteine motif.

[00163] In some embodiments of each or any of the above or below mentioned embodiments, the cysteine motif is selected from the group consisting of:
CX₁₀CX₅CX₅CXCX₇C (SEQ ID NO: 41), CX₁₀CX₆CX₅CXCX₁₅C (SEQ ID NO: 42),
CX₁₁CXCX₅C (SEQ ID NO: 43), CX₁₁CX₅CX₅CXCX₇C (SEQ ID NO: 44),
20 CX₁₀CX₆CX₅CXCX₁₃C (SEQ ID NO: 45), CX₁₀CX₅CXCX₄CX₈C (SEQ ID NO: 46),
CX₁₀CX₆CX₆CXCX₇C (SEQ ID NO: 47), CX₁₀CX₄CX₇CXCX₈C (SEQ ID NO: 48),
CX₁₀CX₄CX₇CXCX₇C (SEQ ID NO: 49), CX₁₃CX₈CX₈C (SEQ ID NO: 50),
CX₁₀CX₆CX₅CXCX₇C (SEQ ID NO: 51), CX₁₀CX₅CX₅C (SEQ ID NO: 52),
CX₁₀CX₅CX₆CXCX₇C (SEQ ID NO: 53), CX₁₀CX₆CX₅CX₇CX₉C (SEQ ID NO: 54),
25 CX₉CX₇CX₅CXCX₇C (SEQ ID NO: 55), CX₁₀CX₆CX₅CXCX₉C (SEQ ID NO: 56),
CX₁₀CXCX₄CX₅CX₁₁C (SEQ ID NO: 57), CX₇CX₃CX₆CX₅CXCX₅CX₁₀C (SEQ ID NO: 58),
CX₁₀CXCX₄CX₅CXCX₂CX₃C (SEQ ID NO: 59), CX₁₆CX₅CXC (SEQ ID NO: 60),
CX₆CX₄CXCX₄CX₅C (SEQ ID NO: 61), CX₁₁CX₄CX₅CX₆CX₃C (SEQ ID NO: 62),
CX₈CX₂CX₆CX₅C (SEQ ID NO: 63), CX₁₀CX₅CX₅CXCX₁₀C (SEQ ID NO: 64),
30 CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65), CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66),
CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67), CX₁₅CX₅CXC (SEQ ID NO: 68),
CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69), CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70),
CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72),
CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74), CX₁₀CXCX₆C
35 (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C
(SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C

(SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCX₁₄C (SEQ ID NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83), CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85), CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87),
 5 CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89), CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91), CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93), CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ ID NO: 95).

10 [00164] In some embodiments of each or any of the above or below mentioned embodiments, the cysteine motif is selected from the group consisting of: CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 98), CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID
 15 NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102), CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107),
 20 CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109), CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113), CX₉CCX₄CX₂CXCCXCX₄CX₃C (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ
 25 ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116), CX₆CCXCX₅CX₄CCXCX₅CX₂C (SEQ ID NO: 117), CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118), CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO: 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137),
 35

CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145),
 5 CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152).

[00165] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises 2 to 6 disulfide bonds.

10 [00166] In some embodiments of each or any of the above or below mentioned embodiments, the X¹X²X³X⁴X⁵ motif in the CDR3 scaffold is TTVHQ (SEQ ID NO: 153), TSVHQ (SEQ ID NO: 154), SSVTQ (SEQ ID NO: 155), STVHQ (SEQ ID NO: 156), ATVRQ (SEQ ID NO: 157), TTVYQ (SEQ ID NO: 158), SPVHQ (SEQ ID NO: 159), ATVYQ (SEQ ID NO: 160), TAVYQ (SEQ ID NO: 161), TNVHQ (SEQ ID NO: 162), ATVHQ (SEQ ID NO: 163), STVYQ (SEQ ID NO: 164), TIVHQ (SEQ ID NO: 165), AIVYQ (SEQ ID NO: 166), TTVFQ (SEQ ID NO: 167), AAVFQ (SEQ ID NO: 168), GTVHQ (SEQ ID NO: 169), ASVHQ (SEQ ID NO: 170), TAVFQ (SEQ ID NO: 171), ATVFQ (SEQ ID NO: 172), AAAHQ (SEQ ID NO: 173), VVVYQ (SEQ ID NO: 174), GTVYQ (SEQ ID NO: 175), TAVHQ (SEQ ID NO: 176), ITVHQ (SEQ ID NO: 177), ITAHQ (SEQ ID NO: 178), VTVHQ (SEQ ID NO: 179); AAVHQ (SEQ ID NO: 180), GTVYQ (SEQ ID NO: 181), TTVLQ (SEQ ID NO: 182), TTTHQ (SEQ ID NO: 183), or TTDYQ (SEQ ID NO: 184).

[00167] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises a CX¹X²X³X⁴X⁵ motif.

25 [00168] In some embodiments of each or any of the above or below mentioned embodiments, the CX¹X²X³X⁴X⁵ motif is CTTVHQ (SEQ ID NO: 185), CTSVHQ (SEQ ID NO: 186), CSSVTQ (SEQ ID NO: 187), CSTVHQ (SEQ ID NO: 188), CATVRQ (SEQ ID NO: 189), CTTVYQ (SEQ ID NO: 190), CSPVHQ (SEQ ID NO: 191), CATVYQ (SEQ ID NO: 192), CTAVYQ (SEQ ID NO: 193), CTNVHQ (SEQ ID NO: 194), CATVHQ (SEQ ID NO: 195), CSTVYQ (SEQ ID NO: 196), CTIVHQ (SEQ ID NO: 197), CAIVYQ (SEQ ID NO: 198), CTTVFQ (SEQ ID NO: 199), CAAVFQ (SEQ ID NO: 200), CGTVHQ (SEQ ID NO: 201), CASVHQ (SEQ ID NO: 202), CTAVFQ (SEQ ID NO: 203), CATVFQ (SEQ ID NO: 204), CAAAHQ (SEQ ID NO: 205), CVVVYQ (SEQ ID NO: 206), CGTVFQ (SEQ ID NO: 207), CTAVHQ (SEQ ID NO: 208), CITVHQ (SEQ ID NO: 209), CITAHQ (SEQ ID NO: 210), CVTVHQ (SEQ ID NO: 211); CAAVHQ (SEQ ID NO: 212), CGTVYQ (SEQ ID NO: 213), CTTVLQ (SEQ ID NO: 214), CTTTHQ (SEQ ID NO: 215), or CTTDYQ (SEQ ID NO: 216).

[00169] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises a $(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

5 [00170] In some embodiments of each or any of the above or below mentioned embodiments, the $(X^aX^b)_z$ motif is CYTYNYEF (SEQ ID NO: 217), HYTYTYDF (SEQ ID NO: 218), HYTYTYEW (SEQ ID NO: 219), KHRITYEW (SEQ ID NO: 220), NYIYKYSF (SEQ ID NO: 221), PYIYTYQF (SEQ ID NO: 222), SFTYTYEW (SEQ ID NO: 223), SYIYIYQW (SEQ ID NO: 224), SYNYTYSW (SEQ ID NO: 225), SYSYSYEF (SEQ ID NO: 226), SYTYNYDF (SEQ ID NO: 227), SYTYNYEW (SEQ ID NO: 228), SYTYNYQF (SEQ ID NO: 229), SYVWTHNF (SEQ ID NO: 230), TYKYVYEW (SEQ ID NO: 231), TYTYTYEF (SEQ ID NO: 232), TYTYTYEW (SEQ ID NO: 233), VFTYTYEF (SEQ ID NO: 234), AYTIEW (SEQ ID NO: 235), DYIYTY (SEQ ID NO: 236), IHSYEF (SEQ ID NO: 237), SFTYEF (SEQ ID NO: 238), SHSYEF (SEQ ID NO: 239), THTYEF (SEQ ID NO: 240), TWTYEF (SEQ ID NO: 241), TYNIEW (SEQ ID NO: 242), TYSYEF (SEQ ID NO: 243), TYSYEH (SEQ ID NO: 244), TYTYDF (SEQ ID NO: 245), TYTYEF (SEQ ID NO: 246), TYTIEW (SEQ ID NO: 247), AYEF (SEQ ID NO: 248), AYSF (SEQ ID NO: 249), AYSY (SEQ ID NO: 250), CYSF (SEQ ID NO: 251), DYT Y (SEQ ID NO: 252), KYEH (SEQ ID NO: 253), KYEW (SEQ ID NO: 254), MYEF (SEQ ID NO: 255), NWIY (SEQ ID NO: 256), NYDY (SEQ ID NO: 257), NYQW (SEQ ID NO: 258), NYSF (SEQ ID NO: 259), PYEW (SEQ ID NO: 260), RYNW (SEQ ID NO: 261), RYT Y (SEQ ID NO: 262), SYEF (SEQ ID NO: 263), SYEH (SEQ ID NO: 264), SYEW (SEQ ID NO: 265), SYKW (SEQ ID NO: 266), SYTY (SEQ ID NO: 267), TYDF (SEQ ID NO: 268), TYEF (SEQ ID NO: 269), TIEW (SEQ ID NO: 270), TYQW (SEQ ID NO: 271), TYTY (SEQ ID NO: 272), or VYEW (SEQ ID NO: 273).

[00171] In some embodiments of each or any of the above or below mentioned embodiments, the $(X^aX^b)_z$ motif is YXYXYX.

[00172] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises a $X^1X^2X^3X^4X^5X_n$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q), and wherein n is 27 - 54.

35 [00173] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises $X_n(X^aX^b)_z$ motif, wherein X^a is any amino acid

residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

[00174] In some embodiments of each or any of the above or below mentioned
 5 embodiments, the CDR3 scaffold comprises a $X^1X^2X^3X^4X^5X_n(X^aX^b)_z$ motif, herein X_1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X_2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X_3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X_4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), wherein X_5 is
 10 glutamine (Q), X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

[00175] In some embodiments of each or any of the above or below mentioned
 15 embodiments, the $X^1X^2X^3X^4X^5$ motif in the CDR3 scaffold is TTVHQ (SEQ ID NO: 153) or TSVHQ (SEQ ID NO: 154), and wherein the $(X^aX^b)_z$ motif is YXYXYX.

[00176] In some embodiments of each or any of the above or below mentioned
 20 embodiments, the CDR3 scaffold comprises: a $CX^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is
 25 valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q), a cysteine motif selected from the group consisting of: $CX_{10}CX_5CX_5CXCX_7C$ (SEQ ID NO: 41), $CX_{10}CX_6CX_5CXCX_{15}C$ (SEQ ID NO: 42), $CX_{11}CXCX_5C$ (SEQ ID NO: 43), $CX_{11}CX_5CX_5CXCX_7C$ (SEQ ID NO: 44), $CX_{10}CX_6CX_5CXCX_{13}C$ (SEQ ID NO: 45),
 30 $CX_{10}CX_5CXCX_4CX_8C$ (SEQ ID NO: 46), $CX_{10}CX_6CX_6CXCX_7C$ (SEQ ID NO: 47), $CX_{10}CX_4CX_7CXCX_8C$ (SEQ ID NO: 48), $CX_{10}CX_4CX_7CXCX_7C$ (SEQ ID NO: 49), $CX_{13}CX_8CX_8C$ (SEQ ID NO: 50), $CX_{10}CX_6CX_5CXCX_7C$ (SEQ ID NO: 51), $CX_{10}CX_5CX_5C$ (SEQ ID NO: 52), $CX_{10}CX_5CX_6CXCX_7C$ (SEQ ID NO: 53), $CX_{10}CX_6CX_5CX_7CX_9C$ (SEQ ID NO: 54), $CX_9CX_7CX_5CXCX_7C$ (SEQ ID NO: 55), $CX_{10}CX_6CX_5CXCX_9C$ (SEQ ID NO: 56),
 35 $CX_{10}CXCX_4CX_5CX_{11}C$ (SEQ ID NO: 57), $CX_7CX_3CX_6CX_5CXCX_5CX_{10}C$ (SEQ ID NO: 58), $CX_{10}CXCX_4CX_5CXCX_2CX_3C$ (SEQ ID NO: 59), $CX_{16}CX_5CXC$ (SEQ ID NO: 60), $CX_6CX_4CXCX_4CX_5C$ (SEQ ID NO: 61), $CX_{11}CX_4CX_5CX_6CX_3C$ (SEQ ID NO: 62), $CX_8CX_2CX_6CX_5C$ (SEQ ID NO: 63), $CX_{10}CX_5CX_5CXCX_{10}C$ (SEQ ID NO: 64), $CX_{10}CXCX_6CX_4CXC$ (SEQ ID NO: 65), $CX_{10}CX_5CX_5CXCX_2C$ (SEQ ID NO: 66),
 $CX_{14}CX_2CX_3CXCXC$ (SEQ ID NO: 67), $CX_{15}CX_5CXC$ (SEQ ID NO: 68), $CX_4CX_6CX_9CX_2CX_{11}C$ (SEQ ID NO: 69), $CX_6CX_4CX_5CX_5CX_{12}C$ (SEQ ID NO: 70),

CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72),
 CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74), CX₁₀CXCX₆C
 (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C
 (SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C
 5 (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCX₁₄C (SEQ ID
 NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83),
 CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85),
 CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87),
 CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89),
 10 CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91),
 CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93),
 CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ
 ID NO: 95); and (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic
 amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F),
 15 tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[00177] In some embodiments of each or any of the above or below mentioned
 embodiments, the CDR3 scaffold comprises: a CX¹X²X³X⁴X⁵ motif, wherein X¹ is threonine
 (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X² is serine (S), threonine
 (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X³ is
 20 valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X⁴ is histidine (H),
 threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X⁵ is
 glutamine (Q); a cysteine motif selected from the group consisting of: wherein the cysteine
 motif is selected from the group consisting of: CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ
 ID NO: 96), CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97),
 25 CX₇CXCX₅CX₄CCCX₄CX₆CXC (SEQ ID NO: 98), CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID
 NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100),
 CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID
 NO: 102), CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C
 (SEQ ID NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105),
 30 CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC
 (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108),
 CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109), CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C
 (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111),
 CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ
 ID NO: 113), CX₉CCX₄CX₂CXCCXCX₄CX₃C (SEQ ID NO: 114),
 35 CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC

(SEQ ID NO: 116), CX₆CCXCX₅CX₄CCXCX₅CX₂C (SEQ ID NO: 117), CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118), CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO: 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152); and a (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[00178] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises SEQ ID NO: 40 or a derivative thereof.

[00179] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises amino acid residues 3-6 of any of one SEQ ID NO: 1-4.

[00180] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises a non-human DH or a derivative thereof.

[00181] In some embodiments of each or any of the above or below mentioned embodiments, the non-human DH is SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, or SEQ ID NO: 12

[00182] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises a JH sequence or a derivative thereof.

[00183] In some embodiments of each or any of the above or below mentioned embodiments, the JH sequence is SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, or SEQ ID NO: 17.

[00184] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises: a non-human VH sequence or a derivative thereof; a non-human DH sequence or a derivative thereof; and/or a JH sequence or derivative thereof.

5 [00185] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises an additional amino acid sequence comprising two to six amino acid residues or more positioned between the VH sequence and the DH sequence.

10 [00186] In some embodiments of each or any of the above or below mentioned embodiments, the additional amino acid sequence is selected from the group consisting of: IR, IF, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20 or SEQ ID NO: 21.

[00187] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold comprises a sequence derived from or based on SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 15 27, or SEQ ID NO: 28.

[00188] In some embodiments of each or any of the above or below mentioned embodiments, the CDR3 scaffold is a ruminant CDR3 scaffold.

[00189] In some embodiments of each or any of the above or below mentioned embodiments, the ruminant is a cow.

20 [00190] The present disclosure also provides a library comprising the CDR3 scaffold disclosed herein.

[00191] The present disclosure also provides a polynucleotide encoding the CDR3 scaffold disclosed herein.

25 [00192] The present disclosure also provides a vector comprising any of the polynucleotides disclosed herein.

[00193] The present disclosure also provides a host cell comprising any of the vectors disclosed herein.

[00194] The present disclosure also provides

30 [00195] In some embodiments of each or any of the above or below mentioned embodiments, the cell is a bacteria, virus, or bacteriophage.

BRIEF DESCRIPTION OF THE DRAWINGS

[00196] The foregoing summary, as well as the following detailed description of the disclosure, will be better understood when read in conjunction with the appended figures.
35 For the purpose of illustrating the disclosure, shown in the figures are embodiments which

are presently preferred. It should be understood, however, that the disclosure is not limited to the precise arrangements, examples and instrumentalities shown.

[00197] Figure 1 shows a sequence alignment of exemplary bovine-derived antibody variable region sequences designated BLV1H12, BLV5B8, BLV5D3, BLV8C11, BF4E9, BF1H1, or F18 that comprise an ultralong CDR3 sequence.

[00198] Figure 2A-C depicts ultralong CDR3 sequences. (Top) Translation from the germline V_H BUL, D_H2 , and J_H . The 5 full length ultralong CDR H3s reported in the literature contain between four and eight cysteines and are not highly homologous to one another; however, some conservation of cysteine residues with D_H2 could be found when the first cysteine of these CDR H3s was “fixed” prior to alignment. Four of the seven sequences (BLV1H12, BLV5D3, BLV8C11, and BF4E9) contain four cysteines in the same positions as D_H2 , but also have additional cysteines. BLV5B8 has two cysteines in common with the germline D_H2 . This limited homology with some cysteine conservation suggests that mutation of D_H2 could generate these sequences. B-L1 and B-L2 are from initial sequences from bovine spleen, and the remaining are selected ultralong CDR H3 sequences from deep sequencing data. The first group contains the longest CDR H3s identified, and appear clonally related. The * indicates a sequence represented 167 times, suggesting it was strongly selected for function. Several of the eight-cysteine sequences appear selected for function as they were represented multiple times, indicated in parentheses. Other representative sequences of various lengths are indicated in the last group. The framework cysteine and tryptophan residues that define the CDR H3 boundaries are double-underlined. The sequences BLV1H12 through UL-77 (left-most column) presented in Tables 2A-C are depicted broken apart into four segments to identify the segments of amino acid residues that are derived from certain germline sequences and V/D/J joining sequences. Moving from left to right, the first segment is derived from the V_H germline and is represented in the disclosure as a $X^1X^2X^3X^4X^5$ motif. The second segment represents sequences from V-D joining and is represented in the disclosure as X_n . The third segment is a string of amino acid residues derived from D_H2 germline, and the fourth segment is a string of amino acid residues derived from J_H1 germline region.

[00199] Figure 3 depicts a sequence alignment of exemplary bovine-derived ultralong CDR3 sequences designated BLV1H12, BLV5B8, BLV5D3, BLV8C11, BF4E9, BF1H1, or F18.

[00200] Figure 4 shows an exemplary bovine germline heavy chain variable region (V_H) sequence designated VH-UL suitable for modification or use with an ultralong CDR3 sequence.

[00201] Figure 5A-B shows exemplary human germline heavy chain variable region sequences designated 4-39, 4-59*03, 4-34*09, and 4-34*02 that are suitable for modification or use with an ultralong CDR3 sequence (A) and an alignment of these sequences (B).

5 [00202] Figure 6 shows an exemplary bovine light chain variable region sequence designated BLV1H12 suitable for modification or use with an ultralong CDR 3 sequence (e.g., a heavy chain variable region sequence comprising an ultralong CDR3 sequence).

[00203] Figure 7A-B shows exemplary light chain variable region sequences designated VI1-47, VI1-40*1, VI1-51*01, and VI2-18*02 that are suitable for modification or
10 use with an ultralong CDR 3 sequence (A) and an alignment of these sequences (B).

DETAILED DESCRIPTION

[00204] The present disclosure provides humanized antibodies comprising ultralong CDR3 sequences, along with materials for (e.g., protein sequences, genetic
15 sequences, cells, libraries) and methods of making the antibodies (e.g., humanizing methods, library methods). Such humanized antibodies may be useful for the treatment or prevention of a variety of disease states or disorders.

[00205] The present disclosure also provides humanized antibodies comprising ultralong CDR3 sequences, wherein the CDR3 sequences are 35 amino acids in length or
20 longer (e.g., 40 or longer, 45 or longer, 50 or longer, 55 or longer, 60 or longer) and/or wherein the CDR3 sequences have at least 3 cysteine residues or more (e.g., 3 or more cysteine residues, 4 or more cysteine residues, 5 or more cysteine residues, 6 or more cysteine residues, 7 or more cysteine residues, 8 or more cysteine residues, 9 or more cysteine residues, 10 or more cysteine residues, 11 or more cysteine residues, or 12 or
25 more cysteine residues). Such antibodies, as described herein, bind (e.g., specifically or selectively bind) a variety of targets, including, for example protein targets such as transmembrane proteins (e.g., GPCRs, ion channels, transporter, cell surface receptors).

[00206] The present disclosure also provides methods and materials for the preparation or making of humanized antibodies comprising ultralong CDR3 sequences.
30 Such materials include proteins, genetic sequences, cells and libraries. Such methods include methods of humanization and method of making and screening libraries.

[00207] The present disclosure provides a humanized antibody or binding fragment thereof comprising an ultralong CDR3. In some embodiments, the ultralong CDR3 may be 35 amino acids in length or longer, 40 amino acids in length or longer, 45
35 amino acids in length or longer, 50 amino acids in length or longer, 55 amino acids in length or longer, or 60 amino acids in length or longer. In some embodiments, the

ultralong CDR3 may comprise 3 or more cysteine residues, 4 or more cysteine residues, 5 or more cysteine residues, 6 or more cysteine residues, 7 or more cysteine residues, 8 or more cysteine residues, 9 or more cysteine residues, 10 or more cysteine residues, 11 or more cysteine residues, or 12 or more cysteine residues. The ultralong CDR3 may

5 comprise a cysteine motif including, for example, where the cysteine motif is selected from the group consisting of: CX₁₀CX₅CX₅CXCX₇C (SEQ ID NO: 41), CX₁₀CX₆CX₅CXCX₁₅C (SEQ ID NO: 42), CX₁₁CXCX₅C (SEQ ID NO: 43), CX₁₁CX₅CX₅CXCX₇C (SEQ ID NO: 44), CX₁₀CX₆CX₅CXCX₁₃C (SEQ ID NO: 45), CX₁₀CX₅CXCX₄CX₈C (SEQ ID NO: 46), CX₁₀CX₆CX₆CXCX₇C (SEQ ID NO: 47), CX₁₀CX₄CX₇CXCX₈C (SEQ ID NO: 48),

10 CX₁₀CX₄CX₇CXCX₇C (SEQ ID NO: 49), CX₁₃CX₈CX₈C (SEQ ID NO: 50), CX₁₀CX₆CX₅CXCX₇C (SEQ ID NO: 51), CX₁₀CX₅CX₅C (SEQ ID NO: 52), CX₁₀CX₅CX₆CXCX₇C (SEQ ID NO: 53), CX₁₀CX₆CX₅CX₇CX₉C (SEQ ID NO: 54), CX₉CX₇CX₅CXCX₇C (SEQ ID NO: 55), CX₁₀CX₆CX₅CXCX₉C (SEQ ID NO: 56), CX₁₀CXCX₄CX₅CX₁₁C (SEQ ID NO: 57), CX₇CX₃CX₆CX₅CXCX₅CX₁₀C (SEQ ID NO: 58),

15 CX₁₀CXCX₄CX₅CXCX₂CX₃C (SEQ ID NO: 59), CX₁₆CX₅CXC (SEQ ID NO: 60), CX₆CX₄CXCX₄CX₅C (SEQ ID NO: 61), CX₁₁CX₄CX₅CX₆CX₃C (SEQ ID NO: 62), CX₈CX₂CX₆CX₅C (SEQ ID NO: 63), CX₁₀CX₅CX₅CXCX₁₀C (SEQ ID NO: 64), CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65), CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66), CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67), CX₁₅CX₅CXC (SEQ ID NO: 68),

20 CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69), CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70), CX₇CX₃CXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72), CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74), CX₁₀CXCX₆C (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 77), CX₁₂CX₄CX₅CXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C

25 (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCX₁₄C (SEQ ID NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83), CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85), CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87), CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89),

30 CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91), CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93), CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ ID NO: 95). Alternatively, the ultralong CDR3 may comprise a cysteine motif including, for example, where the cysteine motif is selected from the group consisting of:

35 CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCCX₄CX₆CXC (SEQ ID NO: 98),

CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID
 NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101),
 CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102), CCX₈CX₅CX₄CX₃CX₄CCX₁C (SEQ ID
 NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104),
 5 CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCX₅CX₄CXC
 (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107),
 CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ
 ID NO: 109), CX₃CX₇CX₃CX₄CCX₂CX₅CX₂C (SEQ ID NO: 110),
 CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ
 10 ID NO: 112), CX₈CCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113),
 CX₉CCX₄CX₂CXCCX₄CX₃C (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ
 ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116),
 CX₆CCX₅CX₄CCX₅CX₂C (SEQ ID NO: 117), CX₆CCX₃CXCCX₃CX₄CC (SEQ ID
 NO: 118), CX₆CCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119),
 15 CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO:
 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO:
 123), CX₉CX₇CX₄CCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO:
 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCX₅CX₄C (SEQ ID
 NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID
 20 NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID
 NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO:
 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135),
 CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137),
 CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139),
 25 CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141),
 CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143),
 CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145),
 CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID
 NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO:
 30 151), and CX₉C (SEQ ID NO: 152).

[00208] The present disclosure provides a humanized antibody or binding
 fragment thereof comprising an ultralong CDR3, wherein the ultralong CDR3 comprises a
 X¹X²X³X⁴X⁵ motif, wherein X¹ is threonine (T), glycine (G), alanine (A), serine (S), or valine
 (V), wherein X² is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine
 35 (V), or asparagine (N), wherein X³ is valine (V), alanine (A), threonine (T), or aspartic acid
 (D), wherein X⁴ is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F),

or leucine (L), and wherein X^5 is glutamine (Q). In some embodiments, the $X^1X^2X^3X^4X^5$ motif may be TTVHQ (SEQ ID NO: 153), TSVHQ (SEQ ID NO: 154), SSVTQ (SEQ ID NO: 155), STVHQ (SEQ ID NO: 156), ATVRQ (SEQ ID NO: 157), TTVYQ (SEQ ID NO: 158), SPVHQ (SEQ ID NO: 159), ATVYQ (SEQ ID NO: 160), TAVYQ (SEQ ID NO: 161),
 5 TNVHQ (SEQ ID NO: 162), ATVHQ (SEQ ID NO: 163), STVYQ (SEQ ID NO: 164), TIVHQ (SEQ ID NO: 165), AIVYQ (SEQ ID NO: 166), TTVFQ (SEQ ID NO: 167), AAVFQ (SEQ ID NO: 168), GTVHQ (SEQ ID NO: 169), ASVHQ (SEQ ID NO: 170), TAVFQ (SEQ ID NO: 171), ATVFQ (SEQ ID NO: 172), AAAHQ (SEQ ID NO: 173), VVYQ (SEQ ID NO: 174), GTVFQ (SEQ ID NO: 175), TAVHQ (SEQ ID NO: 176), ITVHQ (SEQ ID NO: 177), ITAHQ (SEQ ID NO: 178), VTVHQ (SEQ ID NO: 179); AAVHQ (SEQ ID NO: 180), GTVYQ (SEQ ID NO: 181), TTVLQ (SEQ ID NO: 182), TTTHQ (SEQ ID NO: 183), or TTDYQ (SEQ ID NO: 184).

[00209] The present disclosure provides a humanized antibody or binding fragment thereof comprising an ultralong CDR3, wherein the ultralong CDR3 comprises a
 15 $(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4. In some embodiments, the $(X^aX^b)_z$ motif may be CYTYNYEF (SEQ ID NO: 217), HYTYTYDF (SEQ ID NO: 218), HYTYTYEW (SEQ ID NO: 219), KHRITYEW (SEQ ID NO: 220), NYIYKYSF (SEQ ID NO: 221), PYIYTYQF (SEQ ID NO: 222), SFTYTYEW (SEQ ID NO: 223), SYIYIYQW (SEQ ID NO: 224), SYNITYSW (SEQ ID NO: 225), SYSYSYEF (SEQ ID NO: 226), SYTYNYDF (SEQ ID NO: 227), SYTYNYEW (SEQ ID NO: 228), SYTYNYQF (SEQ ID NO: 229), SYVWTHNF (SEQ ID NO: 230), TYKYVYEW (SEQ ID NO: 231), TYTYTYEF (SEQ ID NO: 232), TYTYTYEW (SEQ ID NO: 233), VFTYTYEF (SEQ ID NO: 234), AYTYEW (SEQ ID NO: 235), DYIYTY (SEQ ID NO: 236), IHSYEF (SEQ ID NO: 237), SFTYEF (SEQ ID NO: 238), SHSYEF (SEQ ID NO: 239), THTYEF (SEQ ID NO: 240), TWTYEF (SEQ ID NO: 241), TYNTEW (SEQ ID NO: 242), TYSYEF (SEQ ID NO: 243), TYSYEH (SEQ ID NO: 244), TYTYDF (SEQ ID NO: 245), TYTYEF (SEQ ID NO: 246), TYTYEW (SEQ ID NO: 247), ATEF (SEQ ID NO: 248), AYSF (SEQ ID NO: 249), AYSY (SEQ ID NO: 250), CYSF (SEQ ID NO: 251), DYTE (SEQ ID NO: 252), KYEH (SEQ ID NO: 253), KYEW (SEQ ID NO: 254), MYEF (SEQ ID NO: 255), NWIY (SEQ ID NO: 256), NYDY (SEQ ID NO: 257), NYQW (SEQ ID NO: 258), NYSF (SEQ ID NO: 259), PYEW (SEQ ID NO: 260), RYNW (SEQ ID NO: 261), RYTE (SEQ ID NO: 262), SYEF (SEQ ID NO: 263), SYEH (SEQ ID NO: 264), SYEW (SEQ ID NO: 265), SYKW (SEQ ID NO: 266), SYTY (SEQ ID NO: 267), TYDF (SEQ ID NO: 268),
 35 TYEF (SEQ ID NO: 269), TYEW (SEQ ID NO: 270), TYQW (SEQ ID NO: 271), TYTY (SEQ ID NO: 272), or VTEW (SEQ ID NO: 273).

[00210] The present disclosure provides a humanized antibody or binding fragment thereof comprising an ultralong CDR3, wherein the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5X_n(X^aX^b)_z$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q), wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

[00211] The present disclosure provides a humanized antibody or binding fragment thereof comprising an ultralong CDR3, wherein the ultralong CDR3 comprises: a $CX^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q), a cysteine motif selected from the group consisting of: $CX_{10}CX_5CX_5CXCX_7C$ (SEQ ID NO: 41), $CX_{10}CX_6CX_5CXCX_{15}C$ (SEQ ID NO: 42), $CX_{11}CXCX_5C$ (SEQ ID NO: 43), $CX_{11}CX_5CX_5CXCX_7C$ (SEQ ID NO: 44), $CX_{10}CX_6CX_5CXCX_{13}C$ (SEQ ID NO: 45), $CX_{10}CX_5CXCX_4CX_8C$ (SEQ ID NO: 46), $CX_{10}CX_6CX_6CXCX_7C$ (SEQ ID NO: 47), $CX_{10}CX_4CX_7CXCX_8C$ (SEQ ID NO: 48), $CX_{10}CX_4CX_7CXCX_7C$ (SEQ ID NO: 49), $CX_{13}CX_8CX_8C$ (SEQ ID NO: 50), $CX_{10}CX_6CX_5CXCX_7C$ (SEQ ID NO: 51), $CX_{10}CX_5CX_5C$ (SEQ ID NO: 52), $CX_{10}CX_5CX_6CXCX_7C$ (SEQ ID NO: 53), $CX_{10}CX_6CX_5CX_7CX_9C$ (SEQ ID NO: 54), $CX_9CX_7CX_5CXCX_7C$ (SEQ ID NO: 55), $CX_{10}CX_6CX_5CXCX_9C$ (SEQ ID NO: 56), $CX_{10}CXCX_4CX_5CX_{11}C$ (SEQ ID NO: 57), $CX_7CX_3CX_6CX_5CXCX_5CX_{10}C$ (SEQ ID NO: 58), $CX_{10}CXCX_4CX_5CXCX_2CX_3C$ (SEQ ID NO: 59), $CX_{16}CX_5CXC$ (SEQ ID NO: 60), $CX_6CX_4CXCX_4CX_5C$ (SEQ ID NO: 61), $CX_{11}CX_4CX_5CX_6CX_3C$ (SEQ ID NO: 62), $CX_8CX_2CX_6CX_5C$ (SEQ ID NO: 63), $CX_{10}CX_5CX_5CXCX_{10}C$ (SEQ ID NO: 64), $CX_{10}CXCX_6CX_4CXC$ (SEQ ID NO: 65), $CX_{10}CX_5CX_5CXCX_2C$ (SEQ ID NO: 66), $CX_{14}CX_2CX_3CXCXC$ (SEQ ID NO: 67), $CX_{15}CX_5CXC$ (SEQ ID NO: 68), $CX_4CX_6CX_9CX_2CX_{11}C$ (SEQ ID NO: 69), $CX_6CX_4CX_5CX_5CX_{12}C$ (SEQ ID NO: 70), $CX_7CX_3CXCXCX_4CX_5CX_9C$ (SEQ ID NO: 71), $CX_{10}CX_6CX_5C$ (SEQ ID NO: 72), $CX_7CX_3CX_5CX_5CX_9C$ (SEQ ID NO: 73), $CX_7CX_5CXCX_2C$ (SEQ ID NO: 74), $CX_{10}CXCX_6C$ (SEQ ID NO: 75), $CX_{10}CX_3CX_3CX_5CX_7CXCX_6C$ (SEQ ID NO: 76), $CX_{10}CX_4CX_5CX_{12}CX_2C$ (SEQ ID NO: 77), $CX_{12}CX_4CX_5CXCXCX_9CX_3C$ (SEQ ID NO: 78), $CX_{12}CX_4CX_5CX_{12}CX_2C$

(SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCX₁₄C (SEQ ID NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83), CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85), CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87),
 5 CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89), CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91), CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93), CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ ID NO: 95), and a (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[00212] The present disclosure provides a humanized antibody or binding fragment thereof comprising an ultralong CDR3, wherein the ultralong CDR3 comprises: a CX¹X²X³X⁴X⁵ motif, wherein X¹ is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X² is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X³ is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X⁴ is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X⁵ is glutamine (Q); a cysteine motif selected from the group consisting of: wherein the cysteine motif is selected from the group consisting of:
 15 CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCCX₄CX₆CXC (SEQ ID NO: 98), CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102), CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID NO: 103),
 25 CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109), CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C (SEQ ID NO: 110),
 30 CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113), CX₉CCX₄CX₂CXCCXCX₄CX₃C (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116), CX₆CCXCX₅CX₄CCXCX₅CX₂C (SEQ ID NO: 117), CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118),
 35 CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO:

121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO: 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152); and a (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[00213] The present disclosure also provides methods of generating a library of humanized antibodies that comprises an ultralong CDR3, comprising: combining a nucleic acid sequence encoding an ultralong CDR3 with a nucleic acid sequence encoding a human variable region framework (FR) sequence to produce nucleic acids encoding for humanized antibodies that comprises an ultralong CDR3; and expressing the nucleic acids encoding for humanized antibodies that comprises an ultralong CDR3 to generate a library of humanized antibodies that comprises an ultralong CDR3.

[00214] The present disclosure also provides methods of generating a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 that comprises a non-antibody sequence, comprising: combining a nucleic acid sequence encoding an ultralong CDR3, a nucleic acid sequence encoding a human variable region framework (FR) sequence, and a nucleic acid sequence encoding a non-antibody sequence to produce nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a non-antibody sequence, and expressing the nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a non-antibody sequence to generate a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a non-antibody sequence.

[00215] The present disclosure also provides libraries of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 that comprises a non-antibody sequence.

[00216] The present disclosure also provides methods of generating a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 that comprises a cysteine motif, comprising: combining a human variable region framework (FR) sequence, and a nucleic acid sequence encoding an ultralong CDR3 and a cysteine motif; introducing one or more nucleotide changes to the nucleic acid sequence encoding one or more amino acid residues that are positioned between one or more cysteine residues in the cysteine motif for nucleotides encoding different amino acid residues to produce nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a cysteine motif with one or more nucleotide changes introduced between one or more cysteine residues in the cysteine domain; and expressing the nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a cysteine motif with one or more nucleotide changes introduced between one or more cysteine residues in the cysteine domain to generate a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a cysteine motif with one or more amino acid changes introduced between one or more cysteine residues in the cysteine domain.

[00217] The present disclosure also provides libraries of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 that comprises a cysteine motif, wherein the antibodies or binding fragments comprise one or more substitutions of amino acid residues that are positioned between cysteine residues in the cysteine motif.

[00218] The present disclosure also provides methods of generating a library of humanized antibodies or binding fragments thereof comprising a bovine ultralong CDR3, comprising: combining a nucleic acid sequence encoding a human variable region framework (FR) sequence and a nucleic acid encoding a bovine ultralong CDR3, and expressing the nucleic acids encoding a human variable region framework (FR) sequence and a nucleic acid encoding a bovine ultralong CDR3 to generate a library of humanized antibodies or binding fragments thereof comprising a bovine ultralong CDR3.

[00219] The present disclosure also provides a CDR3 scaffold comprising a $X^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q).

Proteins

[00220] The present disclosure provides humanized antibodies comprising ultralong CDR3 sequences.

5 [00221] In an embodiment, the present disclosure provides a humanized antibody comprising an ultralong CDR3, wherein the CDR3 is 35 amino acids in length or more (e.g., 40 or more, 45 or more, 50 or more, 55 or more, 60 or more). Such a humanized antibody may comprise at least 3 cysteine residues or more (e.g., 4 or more, 6 or more, 8 or more) within the ultralong CDR3.

10 [00222] In another embodiment, the present disclosure provides a humanized antibody comprising an ultralong CDR3, wherein the CDR3 is 35 amino acids in length or more and is derived from or based on a non-human sequence. The ultralong CDR3 sequence may be derived from any species that naturally produces ultralong CDR3 antibodies, including ruminants such as cattle (*Bos taurus*).

15 [00223] In another embodiment, the present disclosure provides a humanized antibody comprising an ultralong CDR3, wherein the CDR3 is 35 amino acids in length or more and is derived from a non-antibody sequence. The non-antibody sequence may be derived from any protein family including, but not limited to, chemokines, growth factors, peptides, cytokines, cell surface proteins, serum proteins, toxins, extracellular matrix
20 proteins, clotting factors, secreted proteins, etc. The non-antibody sequence may be of human or non-human origin and may comprise a portion of a non-antibody protein such as a peptide or domain. The non-antibody sequence of an ultralong CDR3 may contain mutations from its natural sequence, including amino acid changes (e.g., substitutions), insertions or deletions. Engineering additional amino acids at the junction between the
25 non-antibody sequence may be done to facilitate or enhance proper folding of the non-antibody sequence within the humanized antibody.

[00224] In another embodiment, the present disclosure provides a humanized antibody comprising an ultralong CDR3, wherein the CDR3 is 35 amino acids in length or more and comprises at least 3 cysteine residues or more, including, for example, 4 or
30 more, 6 or more, and 8 or more.

[00225] In another embodiment, the present disclosure provides for a humanized antibody comprising an ultralong CDR3 wherein the CDR3 is 35 amino acids in length or more and comprises at least 3 cysteine residues or more and wherein the ultralong CDR3 is a component of a multispecific antibody. The multispecific antibody may be bispecific or
35 comprise greater valencies.

[00226] In another embodiment, the present disclosure provides a humanized antibody comprising an ultralong CDR3, wherein the CDR3 is 35 amino acids in length or more and comprises at least 3 cysteine residues or more, wherein the partially human ultralong CDR3 is a component of an immunoconjugate.

5 [00227] In another embodiment, the present disclosure provides a humanized antibody comprising an ultralong CDR3, wherein the CDR3 is 35 amino acids in length or more and comprises at least 3 cysteine residues or more, wherein the humanized antibody comprising an ultralong CDR3 binds to a transmembrane protein target. Such transmembrane targets may include, but are not limited to, GPCRs, ion channels,
10 transporters, and cell surface receptors.

Genetic Sequences

[00228] The present disclosure provides genetic sequences (e.g., genes, nucleic acids, polynucleotides) encoding humanized antibodies comprising ultralong CDR
15 sequences.

[00229] The present disclosure also provides genetic sequences (e.g., genes, nucleic acids, polynucleotides) encoding an ultralong CDR3.

[00230] In an embodiment, the present disclosure provides genetic sequences encoding a humanized antibody comprising an ultralong CDR3, wherein the CDR3 is 35
20 amino acids in length or more (e.g., 40 or more, 45 or more, 50 or more, 55 or more, 60 or more). Such a humanized antibody may comprise at least 3 cysteine residues or more (e.g., 4 or more, 6 or more, 8 or more) within the ultralong CDR3.

[00231] In another embodiment, the present disclosure provides genetic sequences encoding a humanized antibody comprising an ultralong CDR3, wherein the
25 CDR3 is 35 amino acids in length or more and is derived from or based on a non-human sequence. The genetic sequences encoding the ultralong CDR3 may be derived from any species that naturally produces ultralong CDR3 antibodies, including ruminants such as cattle (*Bos taurus*).

[00232] In another embodiment, the present disclosure provides genetic
30 sequences encoding a humanized antibody comprising an ultralong CDR3, wherein the CDR3 is 35 amino acids in length or more and is derived from a non-antibody protein sequence. The genetic sequences encoding the non-antibody protein sequences may be derived from any protein family including, but not limited to, chemokines, growth factors, peptides, cytokines, cell surface proteins, serum proteins, toxins, extracellular matrix
35 proteins, clotting factors, secreted proteins, etc. The non-antibody protein sequence may be of human or non-human origin and may comprise a portion of a non-antibody protein

such as a peptide or domain. The non-antibody protein sequence of an ultralong CDR3 may contain mutations from its natural sequence, including amino acid changes (e.g., substitutions), insertions or deletions. Engineering additional amino acids at the junction between the non-antibody sequence may be done to facilitate or enhance proper folding of the non-antibody sequence within the humanized antibody.

[00233] In another embodiment, the present disclosure provides genetic sequences encoding a humanized antibody comprising an ultralong CDR3, wherein the CDR3 is 35 amino acids in length or more and comprises at least 3 cysteine residues or more, including, for example, 4 or more, 6 or more, and 8 or more.

[00234] In another embodiment, the present disclosure provides genetic sequences encoding a humanized antibody comprising an ultralong CDR3 wherein the CDR3 is 35 amino acids in length or more and comprises at least 3 cysteine residues or more and wherein the ultralong CDR3 is a component of a multispecific antibody. The multispecific antibody may be bispecific or comprise greater valencies.

[00235] In another embodiment, the present disclosure provides genetic sequences encoding a humanized antibody comprising an ultralong CDR3, wherein the CDR3 is 35 amino acids in length or more and comprises at least 3 cysteine residues or more, wherein the ultralong CDR3 is a component of an immunoconjugate.

[00236] In another embodiment, the present disclosure provides genetic sequences encoding a humanized antibody comprising an ultralong CDR3 wherein the CDR3 is 35 amino acids in length or more and comprises at least 3 cysteine residues or more and wherein the humanized antibody comprising an ultralong CDR3 binds to a transmembrane protein target. Such transmembrane targets may include, but are not limited to, GPCRs, ion channels, transporters, and cell surface receptors.

Libraries and Arrays

[00237] The present disclosure provides collections, libraries, and arrays of humanized antibodies comprising ultralong CDR3 sequences.

[00238] In an embodiment, the present disclosure provides a library or an array of humanized antibodies comprising ultralong CDR3 sequences wherein at least two members of the library or array differ in the positions of at least one of the cysteines in the ultralong CDR3 sequence. Structural diversity may be enhanced through different numbers of cysteines in the ultralong CDR3 sequence (e.g., at least 3 or more cysteine residues such as 4 or more, 6 or more and 8 or more) and/or through different disulfide bond formation, and hence different loop structures.

[00239] In another embodiment, the present disclosure provides for a library or an array of humanized antibodies comprising ultralong CDR3 sequences wherein at least two members of the library or the array differ in at least one amino acid located between cysteines in the ultralong CDR3. In this regard, members of the library or the array can
5 contain cysteines in the same positions of CDR3, resulting in similar overall structural folds, but with fine differences brought about through different amino acid side chains. Such libraries or arrays may be useful for affinity maturation.

[00240] In another embodiment, the present disclosure provides libraries or arrays of humanized antibodies comprising ultralong CDR3 sequences wherein at least two of the
10 ultralong CDR3 sequences differ in length (e.g., 35 amino acids in length or more such as 40 or more, 45 or more, 50 or more, 55 or more and 60 or more). The amino acid and cysteine content may or may not be altered between the members of the library or the array. Different lengths of ultralong CDR3 sequences may provide for unique binding sites, including, for example, due to steric differences, as a result of altered length.

[00241] In another embodiment, the present disclosure provides libraries or arrays of humanized antibodies comprising ultralong CDR3 sequences wherein at least two
15 members of the library differ in the human framework used to construct the humanized antibody comprising an ultralong CDR3.

[00242] In another embodiment, the present disclosure provides libraries or arrays
20 of humanized antibodies comprising ultralong CDR3 sequences wherein at least two members of the library or the array differ in having a non-antibody protein sequence that comprises a portion of the ultralong CDR3. Such libraries or arrays may contain multiple non-antibody protein sequences, including for chemokines, growth factors, peptides, cytokines, cell surface proteins, serum proteins, toxins, extracellular matrix proteins,
25 clotting factors, secreted proteins, viral or bacterial proteins, etc. The non-antibody protein sequence may be of human or non-human origin and may be comprised of a portion of a non-antibody protein such as a peptide or domain. The non-antibody protein sequence of the ultralong CDR3 may contain mutations from its natural sequence, including amino acid changes (e.g., substitutions), or insertions or deletions. Engineering additional amino acids
30 at the junction between the non-antibody sequence within the ultralong CDR3 may be done to facilitate or enhance proper folding of the non-antibody sequence within the humanized antibody.

[00243] The libraries or the arrays of the present disclosure may be in several formats well known in the art. The library or the array may be an addressable library or an
35 addressable array. The library or array may be in display format, for example, the antibody sequences may be expressed on phage, ribosomes, mRNA, yeast, or mammalian cells.

Cells

[00244] The present disclosure provides cells comprising genetic sequences encoding humanized antibodies comprising ultralong CDR3 sequences.

5 [00245] In an embodiment, the present disclosure provides cells expressing a humanized antibody comprising an ultralong CDR3. The cells may be prokaryotic or eukaryotic, and a humanized antibody comprising an ultralong CDR3 may be expressed on the cell surface or secreted into the media. When displayed on the cell surface a humanized antibody preferentially contains a motif for insertion into the plasmid membrane
10 such as a membrane spanning domain at the C-terminus or a lipid attachment site. For bacterial cells, a humanized antibody comprising an ultralong CDR3 may be secreted into the periplasm. When the cells are eukaryotic, they may be transiently transfected with genetic sequences encoding a humanized antibody comprising an ultralong CDR3. Alternatively, a stable cell line or stable pools may be created by transfecting or
15 transducing genetic sequences encoding a humanized antibody comprising an ultralong CDR3 by methods well known to those of skill in the art. Cells can be selected by fluorescence activated cell sorting (FACS) or through selection for a gene encoding drug resistance. Cells useful for producing humanized antibodies comprising ultralong CDR3 sequences include prokaryotic cells like *E. coli*, eukaryotic cells like the yeasts
20 *Saccharomyces cerevisiae* and *Pichia pastoris*, chinese hamster ovary (CHO) cells, monkey cells like COS-1, or human cells like HEK-293, HeLa, SP-1.

Humanization Methods

[00246] The present disclosure provides methods for making humanized
25 antibodies comprising ultralong CDR3 sequences, comprising the steps of engineering an ultralong CDR3 sequence derived from a non-human CDR3 into a human framework. The human framework may be of germline origin, or may be derived from non-germline (e.g. mutated or affinity matured) sequences. Genetic engineering techniques well known to those in the art, including as disclosed herein, may be used to generate a hybrid DNA
30 sequence containing a human framework and a non-human ultralong CDR3. Unlike human antibodies which may be encoded by V region genes derived from one of seven families, bovine antibodies which produce ultralong CDR3 sequences appear to utilize a single V region family which may be considered to be most homologous to the human VH4 family. In a preferred embodiment where ultralong CDR3 sequences derived from cattle
35 are to be humanized to produce an antibody comprising an ultralong CDR3, human V region sequences derived from the VH4 family may be genetically fused to a bovine-

derived ultralong CDR3 sequence. Exemplary VH4 germline gene sequences in the human antibody locus are shown in Figure 5A (e.g., SEQ ID NOS: 31-34; and 368-371).

[00247] The present disclosure also provides methods of humanizing an antibody variable region comprising the step of genetically combining a nucleic acid sequence encoding a non-human ultralong CDR3 (ULCDR3) with a nucleic acid sequence encoding a human variable region framework (FR) sequence. Also provided are methods of making a humanized antibody variable region comprising selecting a human framework sequence comprising FR1, FR2, and FR3; selecting a CDR1 sequence; selecting a CDR2 sequence; selecting an ultralong CDR3 sequence; and combining the sequences as FR1-CDR1-FR2-CDR2-FR3-ULCDR3. Also provided are methods of making a humanized antibody variable region sequence comprising selecting a human antibody variable region sequence comprising a sequence encoding FR1-CDR1-FR2-CDR2-FR3; selecting a sequence encoding a non-human ultralong CDR3 (ULCDR3); and genetically fusing the human sequence of step (a) in frame with the non-human sequence of step (b) to generate a sequence encoding FR1-CDR1-FR2-CDR2-FR3-ULCDR3.

[00248] In an embodiment, the present disclosure provides a fusion of a human VH4 framework sequence to a bovine-derived ultralong CDR3, for example, as may be accomplished through the following steps. First, the second cysteine of a V region genetic sequence is identified along with the nucleotide sequence encoding the second cysteine. Generally, the second cysteine marks the boundary of the framework and CDR3 two residues upstream (N-terminal) of the CDR3. Second, the second cysteine in a bovine-derived V region sequence is identified which similarly marks 2 residues upstream (N-terminal) of the CDR3. Third, the genetic material encoding the human V region is combined with the genetic sequence encoding the ultralong CDR3. Thus, a genetic fusion may be made, wherein the ultralong CDR3 sequence is placed in frame of the human V region sequence. Preferably a humanized antibody comprising an ultralong CDR3 is as near to human in amino acid composition as possible. Optionally, a J region sequence may be mutated from bovine-derived sequence to a human sequence. Also optionally, a humanized heavy chain may be paired with a human light chain.

[00249] In another embodiment, the present disclosure provides pairing of a human ultralong CDR3 heavy chain with a non-human light chain.

[00250] In another embodiment, the present disclosure provides pairing of a humanized heavy chain comprising an ultralong CDR3 with a human light chain. Preferably the light chain is homologous to a bovine light chain known to pair with a bovine ultralong CDR3 heavy chain. An exemplary bovine light chain is shown in Figure 7A (e.g., SEQ ID NO: 36-39; and 373-376).

Library Methods

[00251] The present disclosure provides methods for making libraries comprising humanized antibodies comprising ultralong CDR3 sequences. Methods for making libraries of spatially addressed libraries are described in WO 2010/054007. Methods of making libraries in yeast, phage, *E. coli*, or mammalian cells are well known in the art.

[00252] The present disclosure also provides methods of screening libraries of humanized antibodies comprising ultralong CDR3 sequences.

Definitions

[00253] An “ultralong CDR3” or an “ultralong CDR3 sequence”, used interchangeably herein, comprises a CDR3 or CDR3 sequence that is not derived from a human antibody sequence. An ultralong CDR3 may be 35 amino acids in length or longer, for example, 40 amino acids in length or longer, 45 amino acids in length or longer, 50 amino acids in length or longer, 55 amino acids in length or longer, or 60 amino acids in length or longer. The length of the ultralong CDR3 may include a non-antibody sequence. An ultralong CDR3 may comprise a non-antibody sequence, including, for example, an interleukin sequence, a hormone sequence, a cytokine sequence, a toxin sequence, a lymphokine sequence, a growth factor sequence, a chemokine sequence, a toxin sequence, or combinations thereof. Preferably, the ultralong CDR3 is a heavy chain CDR3 (CDR-H3 or CDRH3). Preferably, the ultralong CDR3 is a sequence derived from or based on a ruminant (e.g., bovine) sequence. An ultralong CDR3 may comprise at least 3 or more cysteine residues, for example, 4 or more cysteine residues, 6 or more cysteine residues, 8 or more cysteine residues, 10 or more cysteine residues, or 12 or more cysteine residues (e.g., 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 or more). An ultralong CDR3 may comprise one or more of the following motifs: a cysteine motif, a $X^1X^2X^3X^4X^5$ motif, a $CX^1X^2X^3X^4X^5$ motif, or a $(X^aX^b)_z$ motif. A “cysteine motif” is a segment of amino acid residues in an ultralong CDR3 that comprises 3 or more cysteine residues including, 4 or more cysteine residues, 5 or more cysteine residues, 6 or more cysteine residues, 7 or more cysteine residues, 8 or more cysteine residues, 9 or more cysteine residues, 10 or more cysteine residues, 11 or more cysteine residues, or 12 or more cysteine residues. A cysteine motif may comprise an amino acid sequence selected from the group consisting of: $CX_{10}CX_5CX_5CXCX_7C$ (SEQ ID NO: 41), $CX_{10}CX_6CX_5CXCX_{15}C$ (SEQ ID NO: 42), $CX_{11}CXCX_5C$ (SEQ ID NO: 43), $CX_{11}CX_5CX_5CXCX_7C$ (SEQ ID NO: 44), $CX_{10}CX_6CX_5CXCX_{13}C$ (SEQ ID NO: 45), $CX_{10}CX_5CXCX_4CX_8C$ (SEQ ID NO: 46), $CX_{10}CX_6CX_6CXCX_7C$ (SEQ ID NO: 47), $CX_{10}CX_4CX_7CXCX_8C$ (SEQ ID NO: 48),

CX₁₀CX₄CX₇CXCX₇C (SEQ ID NO: 49), CX₁₃CX₈CX₈C (SEQ ID NO: 50),
 CX₁₀CX₆CX₅CXCX₇C (SEQ ID NO: 51), CX₁₀CX₅CX₅C (SEQ ID NO: 52),
 CX₁₀CX₅CX₆CXCX₇C (SEQ ID NO: 53), CX₁₀CX₆CX₅CX₇CX₉C (SEQ ID NO: 54),
 CX₉CX₇CX₅CXCX₇C (SEQ ID NO: 55), CX₁₀CX₆CX₅CXCX₉C (SEQ ID NO: 56),
 5 CX₁₀CXCX₄CX₅CX₁₁C (SEQ ID NO: 57), CX₇CX₃CX₆CX₅CXCX₅CX₁₀C (SEQ ID NO: 58),
 CX₁₀CXCX₄CX₅CXCX₂CX₃C (SEQ ID NO: 59), CX₁₆CX₅CXC (SEQ ID NO: 60),
 CX₆CX₄CXCX₄CX₅C (SEQ ID NO: 61), CX₁₁CX₄CX₅CX₆CX₃C (SEQ ID NO: 62),
 CX₈CX₂CX₆CX₅C (SEQ ID NO: 63), CX₁₀CX₅CX₅CXCX₁₀C (SEQ ID NO: 64),
 CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65), CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66),
 10 CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67), CX₁₅CX₅CXC (SEQ ID NO: 68),
 CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69), CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70),
 CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72),
 CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74), CX₁₀CXCX₆C
 (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C
 15 (SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C
 (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCX₁₄C (SEQ ID
 NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83),
 CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85),
 CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87),
 20 CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89),
 CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91),
 CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93),
 CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ
 ID NO: 95). Alternatively, a cysteine motif may comprise an amino acid sequence selected
 25 from the group consisting of: CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96),
 CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCCX₄CX₆CXC (SEQ
 ID NO: 98), CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99),
 CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C
 (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102),
 30 CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID
 NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105),
 CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC
 (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108),
 CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109), CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C
 35 (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111),
 CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ

ID NO: 113), CX₉CCX₄CX₂CXCCXCX₄CX₃C (SEQ ID NO: 114),
 CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC
 (SEQ ID NO: 116), CX₆CCXCX₅CX₄CCXCX₅CX₂C (SEQ ID NO: 117),
 CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118), CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID
 5 NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCX₄CX₉C
 (SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C
 (SEQ ID NO: 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCX₁₀CX₂CX₃C
 (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126),
 CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCX₁₀C (SEQ ID NO: 128),
 10 CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO:
 130), CX₁₁CX₃CX₄CCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID
 NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134),
 CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136),
 CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138),
 15 CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140),
 CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142),
 CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144),
 CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C
 (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149),
 20 CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152). A
 cysteine motif is preferably positioned within an ultralong CDR3 between a X¹X²X³X⁴X⁵
 motif and a (X^aX^b)_z motif. A "X¹X²X³X⁴X⁵ motif" is a series of five consecutive amino acid
 residues in an ultralong CDR3, wherein X¹ is threonine (T), glycine (G), alanine (A), serine
 (S), or valine (V), wherein X² is serine (S), threonine (T), proline (P), isoleucine (I), alanine
 25 (A), valine (V), or asparagine (N), wherein X³ is valine (V), alanine (A), threonine (T), or
 aspartic acid (D), wherein X⁴ is histidine (H), threonine (T), arginine (R), tyrosine (Y),
 phenylalanine (F), or leucine (L), and wherein X⁵ is glutamine (Q). In some embodiments,
 the X¹X²X³X⁴X⁵ motif may be TTVHQ (SEQ ID NO: 153), TSVHQ (SEQ ID NO: 154),
 SSVTQ (SEQ ID NO: 155), STVHQ (SEQ ID NO: 156), ATVRQ (SEQ ID NO: 157), TTVYQ
 30 (SEQ ID NO: 158), SPVHQ (SEQ ID NO: 159), ATVYQ (SEQ ID NO: 160), TAVYQ (SEQ
 ID NO: 161), TNVHQ (SEQ ID NO: 162), ATVHQ (SEQ ID NO: 163), STVYQ (SEQ ID NO:
 164), TIVHQ (SEQ ID NO: 165), AIVYQ (SEQ ID NO: 166), TTVFQ (SEQ ID NO: 167),
 AAVFQ (SEQ ID NO: 168), GTVHQ (SEQ ID NO: 169), ASVHQ (SEQ ID NO: 170),
 TAVFQ (SEQ ID NO: 171), ATVFQ (SEQ ID NO: 172), AAAHQ (SEQ ID NO: 173), VVVYQ
 35 (SEQ ID NO: 174), GTVFQ (SEQ ID NO: 175), TAVHQ (SEQ ID NO: 176), ITVHQ (SEQ ID
 NO: 177), ITAHQ (SEQ ID NO: 178), VTVHQ (SEQ ID NO: 179); AAVHQ (SEQ ID NO:

180), GTVYQ (SEQ ID NO: 181), TTVLQ (SEQ ID NO: 182), TTTHQ (SEQ ID NO: 183), or TTDYQ (SEQ ID NO: 184). A “CX¹X²X³X⁴X⁵ motif” is a series of six consecutive amino acid residues in an ultralong CDR3, wherein the first amino acid residue is cysteine, wherein X¹ is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X² is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X³ is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X⁴ is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X⁵ is glutamine (Q). In some embodiments, the CX¹X²X³X⁴X⁵ motif is CTTVHQ (SEQ ID NO: 185), CTSVHQ (SEQ ID NO: 186), CSSVTQ (SEQ ID NO: 187), CSTVHQ (SEQ ID NO: 188), CATVRQ (SEQ ID NO: 189), CTTVYQ (SEQ ID NO: 190), CSPVHQ (SEQ ID NO: 191), CATVYQ (SEQ ID NO: 192), CTAVYQ (SEQ ID NO: 193), CTNVHQ (SEQ ID NO: 194), CATVHQ (SEQ ID NO: 195), CSTVYQ (SEQ ID NO: 196), CTIVHQ (SEQ ID NO: 197), CAIVYQ (SEQ ID NO: 198), CTTVFQ (SEQ ID NO: 199), CAAVFQ (SEQ ID NO: 200), CGTVHQ (SEQ ID NO: 201), CASVHQ (SEQ ID NO: 202), CTAVFQ (SEQ ID NO: 203), CATVFQ (SEQ ID NO: 204), CAAAHQ (SEQ ID NO: 205), CVVVYQ (SEQ ID NO: 206), CGTVFQ (SEQ ID NO: 207), CTAVHQ (SEQ ID NO: 208), CITVHQ (SEQ ID NO: 209), CITAHQ (SEQ ID NO: 210), CVTVHQ (SEQ ID NO: 211); CAAVHQ (SEQ ID NO: 212), CGTVYQ (SEQ ID NO: 213), CTTVLQ (SEQ ID NO: 214), CTTTHQ (SEQ ID NO: 215), or CTTDYQ (SEQ ID NO: 216). A “(X^aX^b)_z” motif is a repeating series of two amino acid residues in an ultralong CDR3, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4. In some embodiments, the (X^aX^b)_z motif may comprise CYTYNYEF (SEQ ID NO: 217), HYTYTYDF (SEQ ID NO: 218), HYTYTYEW (SEQ ID NO: 219), KHRYTYEW (SEQ ID NO: 220), NYIYKYSF (SEQ ID NO: 221), PYIYTYQF (SEQ ID NO: 222), SFTYTYEW (SEQ ID NO: 223), SYIYIYQW (SEQ ID NO: 224), SYNYTYSW (SEQ ID NO: 225), SYSYSYEF (SEQ ID NO: 226), SYTYNYDF (SEQ ID NO: 227), SYTYNYEW (SEQ ID NO: 228), SYTYNYQF (SEQ ID NO: 229), SYVWTHNF (SEQ ID NO: 230), TYKYVYEW (SEQ ID NO: 231), TYTYTYEF (SEQ ID NO: 232), TYTYTYEW (SEQ ID NO: 233), VFTYTYEF (SEQ ID NO: 234), AYTYEW (SEQ ID NO: 235), DYIYTY (SEQ ID NO: 236), IHSYEF (SEQ ID NO: 237), SFTYEF (SEQ ID NO: 238), SHSYEF (SEQ ID NO: 239), THTYEF (SEQ ID NO: 240), TWTYEF (SEQ ID NO: 241), TYNYEW (SEQ ID NO: 242), TYSYEF (SEQ ID NO: 243), TYSYEH (SEQ ID NO: 244), TYTYDF (SEQ ID NO: 245), TYTYEF (SEQ ID NO: 246), TYTYEW (SEQ ID NO: 247), AYEY (SEQ ID NO: 248), AYSF (SEQ ID NO: 249), AYSY (SEQ ID NO: 250), CYSF (SEQ ID NO: 251), DYT Y (SEQ ID NO: 252), KYEH (SEQ ID NO: 253), KYEW (SEQ ID NO: 254), MYEF (SEQ ID NO: 255), NWIY (SEQ ID NO: 256), NYDY (SEQ ID NO: 257),

NYQW (SEQ ID NO: 258), NYSF (SEQ ID NO: 259), PYEW (SEQ ID NO: 260), RYNW
 (SEQ ID NO: 261), RYTY (SEQ ID NO: 262), SYEF (SEQ ID NO: 263), SYEH (SEQ ID
 NO: 264), SYEW (SEQ ID NO: 265), SYKW (SEQ ID NO: 266), SYTY (SEQ ID NO: 267),
 TYDF (SEQ ID NO: 268), TYEF (SEQ ID NO: 269), TYEW (SEQ ID NO: 270), TYQW
 5 (SEQ ID NO: 271), TYTY (SEQ ID NO: 272), or VYEW (SEQ ID NO: 273). In some
 embodiments, the $(X^aX^b)_z$ motif is YXYXYX. An ultralong CDR3 may comprise an amino
 acid sequence that is derived from or based on SEQ ID NO: 40 (see, e.g., amino acid
 residues 3-6 of SEQ ID NO: 1-4; see also, e.g., VH germline sequences in Figures 2A-C).
 A variable region that comprises an ultralong CDR3 may include an amino acid sequence
 10 that is SEQ ID NO: 1 (CTTVHQ), SEQ ID NO:2 (CTSVHQ), SEQ ID NO:3 (CSSVTQ) or
 SEQ ID NO: 4 (CTTVHP). Such a sequence may be derived from or based on a bovine
 germline VH gene sequence (e.g., SEQ ID NO: 1). An ultralong CDR3 may comprise a
 sequence derived from or based on a non-human DH gene sequence, for example, SEQ
 ID NO: 5 (see also, e.g., Koti, *et al.* (2010) *Mol. Immunol.* 47: 2119-2128), or alternative
 15 sequences such as SEQ ID NO: 6, 7, 8, 9, 10, 11 or 12 (see also, e.g., DH2 germline
 sequences in Figures 2A-C). An ultralong CDR3 may comprise a sequence derived from
 or based on a JH sequence, for example, SEQ ID NO: 13 (see also, e.g., Hosseini, *et al.*
 (2004) *Int. Immunol.* 16: 843-852), or alternative sequences such as SEQ ID NO: 14, 15,
 16 or 17 (see also, e.g., JH1 germline sequences in Figures 2A-C). In an embodiment, an
 20 ultralong CDR3 may comprise a sequence derived from or based on a non-human VH
 sequence (e.g., SEQ ID NO: 1, 2, 3 or 4; alternatively VH sequences in Figures 2A-C)
 and/or a sequence derived from or based on a non-human DH sequence (e.g., SEQ ID
 NO: 5, 6, 7, 8, 9, 10, 11 or 12; alternatively DH sequences in Figures 2A-C) and/or a
 sequence derived from or based on a JH sequence (e.g., SEQ ID NO: 13, 14, 15, 16, or
 25 17; alternatively JH sequences in Figures 2A-C), and optionally an additional sequence
 comprising two to six amino acids or more (e.g., IR, IF, SEQ ID NO: 18, 19, 20 or 21) such
 as, for example, between the VH derived sequence and the DH derived sequence. In
 another embodiment, an ultralong CDR3 may comprise a sequence derived from or based
 on SEQ ID NO: 22, 23, 24, 25, 26, 27, or 28 (see also, e.g., SEQ ID NOs: 276-359 in
 30 Figures 2A-C).

[00254] An "isolated" biological molecule, such as the various polypeptides,
 polynucleotides, and antibodies disclosed herein, refers to a biological molecule that has
 been identified and separated and/or recovered from at least one component of its natural
 environment.

35 [00255] "Antagonist" refers to any molecule that partially or fully blocks, inhibits, or
 neutralizes an activity (e.g., biological activity) of a polypeptide. Also encompassed by

"antagonist" are molecules that fully or partially inhibit the transcription or translation of mRNA encoding the polypeptide. Suitable antagonist molecules include, e.g., antagonist antibodies or antibody fragments; fragments or amino acid sequence variants of a native polypeptide; peptides; antisense oligonucleotides; small organic molecules; and nucleic acids that encode polypeptide antagonists or antagonist antibodies. Reference to "an" antagonist encompasses a single antagonist or a combination of two or more different antagonists.

[00256] "Agonist" refers to any molecule that partially or fully mimics a biological activity of a polypeptide. Also encompassed by "agonist" are molecules that stimulate the transcription or translation of mRNA encoding the polypeptide. Suitable agonist molecules include, e.g., agonist antibodies or antibody fragments; a native polypeptide; fragments or amino acid sequence variants of a native polypeptide; peptides; antisense oligonucleotides; small organic molecules; and nucleic acids that encode polypeptides agonists or antibodies. Reference to "an" agonist encompasses a single agonist or a combination of two or more different agonists.

[00257] An "isolated" antibody refers to one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody (e.g., as determined by the Lowry method), and preferably to more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence (e.g., by use of a spinning cup sequenator), or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions (e.g., using Coomassie™ blue or, preferably, silver stain). Isolated antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody's natural environment will not be present. Similarly, isolated antibody includes the antibody in medium around recombinant cells. An isolated antibody may be prepared by at least one purification step.

[00258] An "isolated" nucleic acid molecule refers to a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the antibody nucleic acid. An isolated nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated nucleic acid molecules therefore are distinguished from the nucleic acid molecule as it exists in natural cells. However, an isolated nucleic acid molecule includes a nucleic

acid molecule contained in cells that express an antibody where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

[00259] Variable domain residue numbering as in Kabat or amino acid position numbering as in Kabat, and variations thereof, refers to the numbering system used for heavy chain variable domains or light chain variable domains of the compilation of antibodies in Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991). Using this numbering system, the actual linear amino acid sequence may contain fewer or additional amino acids corresponding to a shortening of, or insertion into, a FR or CDR of the variable domain. For example, a heavy chain variable domain may include a single amino acid insert (e.g., residue 52a according to Kabat) after residue 52 of H2 and inserted residues (e.g., residues 82a, 82b, and 82c, etc according to Kabat) after heavy chain FR residue 82. The Kabat numbering of residues may be determined for a given antibody by alignment at regions of homology of the sequence of the antibody with a "standard" Kabat numbered sequence.

[00260] "Substantially similar," or "substantially the same", refers to a sufficiently high degree of similarity between two numeric values (generally one associated with an antibody disclosed herein and the other associated with a reference/comparator antibody) such that one of skill in the art would consider the difference between the two values to be of little or no biological and/or statistical significance within the context of the biological characteristic measured by said values (e.g., Kd values). The difference between said two values is preferably less than about 50%, preferably less than about 40%, preferably less than about 30%, preferably less than about 20%, preferably less than about 10% as a function of the value for the reference/comparator antibody.

[00261] "Binding affinity" generally refers to the strength of the sum total of noncovalent interactions between a single binding site of a molecule (e.g., an antibody) and its binding partner (e.g., an antigen). Unless indicated otherwise, "binding affinity" refers to intrinsic binding affinity which reflects a 1:1 interaction between members of a binding pair (e.g., antibody and antigen). The affinity of a molecule X for its partner Y can generally be represented by the dissociation constant. Affinity can be measured by common methods known in the art, including those described herein. Low-affinity antibodies generally bind antigen slowly and tend to dissociate readily, whereas high-affinity antibodies generally bind antigen faster and tend to remain bound longer. A variety of methods of measuring binding affinity are known in the art, any of which can be used for purposes of the present disclosure.

[00262] An "on-rate" or "rate of association" or "association rate" or " k_{on} " can be determined with a surface plasmon resonance technique such as Biacore (e.g., Biacore A100, Biacore™-2000, Biacore™-3000, Biacore, Inc., Piscataway, N.J.) carboxymethylated dextran biosensor chips (CM5, Biacore Inc.) and according to the
5 supplier's instructions.

[00263] "Vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a phage vector. Another type of vector is a viral vector, wherein
10 additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the
15 host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "recombinant expression vectors" (or simply, "recombinant vectors"). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. Accordingly, "plasmid" and "vector" may, at times, be used interchangeably as the plasmid is a
20 commonly used form of vector.

[00264] "Gene" refers to a nucleic acid (e.g., DNA) sequence that comprises coding sequences necessary for the production of a polypeptide, precursor, or RNA (e.g., rRNA, tRNA). The polypeptide can be encoded by a full length coding sequence or by any portion of the coding sequence so long as the desired activity or functional properties (e.g.,
25 enzymatic activity, ligand binding, signal transduction, immunogenicity, etc.) of the full-length or fragment are retained. The term also encompasses the coding region of a structural gene and the sequences located adjacent to the coding region on both the 5' and 3' ends for a distance of about 1 kb or more on either end such that the gene corresponds to the length of the full-length mRNA. Sequences located 5' of the coding region and present on the mRNA are referred to as 5' non-translated sequences. Sequences located
30 3' or downstream of the coding region and present on the mRNA are referred to as 3' non-translated sequences. The term "gene" encompasses both cDNA and genomic forms of a gene. A genomic form or clone of a gene contains the coding region interrupted with non-coding sequences termed "introns" or "intervening regions" or "intervening sequences."
35 Introns are segments of a gene that are transcribed into nuclear RNA (hnRNA); introns can contain regulatory elements such as enhancers. Introns are removed or "spliced out" from

the nuclear or primary transcript; introns therefore are absent in the messenger RNA (mRNA) transcript. The mRNA functions during translation to specify the sequence or order of amino acids in a nascent polypeptide. In addition to containing introns, genomic forms of a gene can also include sequences located on both the 5' and 3' end of the sequences that are present on the RNA transcript. These sequences are referred to as "flanking" sequences or regions (these flanking sequences are located 5' or 3' to the non-translated sequences present on the mRNA transcript). The 5' flanking region can contain regulatory sequences such as promoters and enhancers that control or influence the transcription of the gene. The 3' flanking region can contain sequences that direct the termination of transcription, post transcriptional cleavage and polyadenylation.

[00265] "Polynucleotide," or "nucleic acid," as used interchangeably herein, refers to polymers of nucleotides of any length, and include DNA and RNA. The nucleotides can be deoxyribonucleotides, ribonucleotides, modified nucleotides or bases, and/or their analogs, or any substrate that can be incorporated into a polymer by DNA or RNA polymerase, or by a synthetic reaction. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and their analogs. If present, modification to the nucleotide structure may be imparted before or after assembly of the polymer. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynucleotide may be further modified after synthesis, such as by conjugation with a label. Other types of modifications include, for example, "caps", substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as, for example, those with uncharged linkages (*e.g.*, methyl phosphonates, phosphotriesters, phosphoamidates, carbamates, *etc.*) and with charged linkages (*e.g.*, phosphorothioates, phosphorodithioates, *etc.*), those containing pendant moieties, such as, for example, proteins (*e.g.*, nucleases, toxins, antibodies, signal peptides, poly-L-lysine, *etc.*), those with intercalators (*e.g.*, acridine, psoralen, *etc.*), those containing chelators (*e.g.*, metals, radioactive metals, boron, oxidative metals, *etc.*), those containing alkylators, those with modified linkages (*e.g.*, alpha anomeric nucleic acids, *etc.*), as well as unmodified forms of the polynucleotide(s). Further, any of the hydroxyl groups ordinarily present in the sugars may be replaced, for example, by phosphonate groups, phosphate groups, protected by standard protecting groups, or activated to prepare additional linkages to additional nucleotides, or may be conjugated to solid or semi-solid supports. The 5' and 3' terminal OH can be phosphorylated or substituted with amines or organic capping group moieties of from 1 to 20 carbon atoms. Other hydroxyls may also be derivatized to standard protecting groups. Polynucleotides can also contain analogous forms of ribose or deoxyribose sugars that are generally known in the art, including, for example, 2'-O-methyl-, 2'-O-allyl, 2'-fluoro-

or 2'-azido-ribose, carbocyclic sugar analogs, alpha-anomeric sugars, epimeric sugars such as arabinose, xyloses or lyxoses, pyranose sugars, furanose sugars, sedoheptuloses, acyclic analogs and a basic nucleoside analogs such as methyl riboside. One or more phosphodiester linkages may be replaced by alternative linking groups. These alternative linking groups include, but are not limited to, embodiments wherein phosphate is replaced by P(O)S("thioate"), P(S)S ("dithioate"), "(O)NR₂ ("amidate"), P(O)R, P(O)OR', CO or CH₂ ("formacetal"), in which each R or R' is independently H or substituted or unsubstituted alkyl (1-20 C) optionally containing an ether (-O-) linkage, aryl, alkenyl, cycloalkyl, cycloalkenyl or araldyl. Not all linkages in a polynucleotide need be identical. The preceding description applies to all polynucleotides referred to herein, including RNA and DNA.

[00266] "Oligonucleotide" refers to short, generally single stranded, generally synthetic polynucleotides that are generally, but not necessarily, less than about 200 nucleotides in length. The terms "oligonucleotide" and "polynucleotide" are not mutually exclusive. The description above for polynucleotides is equally and fully applicable to oligonucleotides.

[00267] "Stringent hybridization conditions" refer to conditions under which a probe will hybridize to its target subsequence, typically in a complex mixture of nucleic acids, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Probes, "Overview of principles of hybridization and the strategy of nucleic acid assays" (1993). Generally, stringent conditions are selected to be about 5-10°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength pH. The T_m is the temperature (under defined ionic strength, pH, and nucleic concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at T_m, 50% of the probes are occupied at equilibrium). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. For selective or specific hybridization, a positive signal is at least two times background, preferably 10 times background hybridization. Exemplary stringent hybridization conditions can be as following: 50% formamide, 5xSSC, and 1% SDS, incubating at 42° C, or, 5xSSC, 1% SDS, incubating at 65° C, with wash in 0.2xSSC, and 0.1% SDS at 65° C.

[00268] "Recombinant" when used with reference to a cell, nucleic acid, protein or vector indicates that the cell, nucleic acid, protein or vector has been modified by the

introduction of a heterologous nucleic acid or protein, the alteration of a native nucleic acid or protein, or that the cell is derived from a cell so modified. For example, recombinant cells express genes that are not found within the native (non-recombinant) form of the cell or express native genes that are overexpressed or otherwise abnormally expressed such as, for example, expressed as non-naturally occurring fragments or splice variants. By the term "recombinant nucleic acid" herein is meant nucleic acid, originally formed *in vitro*, in general, by the manipulation of nucleic acid, e.g., using polymerases and endonucleases, in a form not normally found in nature. In this manner, operably linkage of different sequences is achieved. Thus an isolated nucleic acid, in a linear form, or an expression vector formed *in vitro* by ligating DNA molecules that are not normally joined, are both considered recombinant for the purposes of this disclosure. It is understood that once a recombinant nucleic acid is made and introduced into a host cell or organism, it will replicate non-recombinantly, e.g., using the *in vivo* cellular machinery of the host cell rather than *in vitro* manipulations; however, such nucleic acids, once produced recombinantly, although subsequently replicated non-recombinantly, are still considered recombinant for the purposes disclosed herein. Similarly, a "recombinant protein" is a protein made using recombinant techniques, e.g., through the expression of a recombinant nucleic acid as depicted above.

[00269] "Percent (%) amino acid sequence identity" with respect to a peptide or polypeptide sequence refers to the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific peptide or polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or MegAlign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared.

[00270] "Polypeptide," "peptide," "protein," and "protein fragment" may be used interchangeably to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymers.

[00271] "Amino acid" refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function similarly to the naturally

occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, e.g., hydroxyproline, gamma-carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to compounds that have the same basic chemical structure as a naturally occurring amino acid, e.g., an alpha carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, e.g., homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs can have modified R groups (e.g., norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. Amino acid mimetics refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions similarly to a naturally occurring amino acid.

[00272] "Conservatively modified variants" applies to both amino acid and nucleic acid sequences. "Amino acid variants" refers to amino acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical or associated (e.g., naturally contiguous) sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode most proteins. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to another of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations," which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes silent variations of the nucleic acid. One of skill will recognize that in certain contexts each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, silent variations of a nucleic acid which encodes a polypeptide is implicit in a described sequence with respect to the expression product, but not with respect to actual probe sequences. As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a "conservatively modified variant" including where the alteration results in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing functionally similar amino acids are well known in the art. Such conservatively modified variants are in addition to and do not exclude polymorphic variants,

interspecies homologs, and alleles disclosed herein. Typically conservative substitutions include: 1) Alanine (A), Glycine (G); 2) Aspartic acid (D), Glutamic acid (E); 3) Asparagine (N), Glutamine (Q); 4) Arginine (R), Lysine (K); 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W); 7) Serine (S),
 5 Threonine (T); and 8) Cysteine (C), Methionine (M) (see, e.g., Creighton, Proteins (1984)).

[00273] "Antibodies" (Abs) and "immunoglobulins" (Igs) are glycoproteins having similar structural characteristics. While antibodies may exhibit binding specificity to a specific antigen, immunoglobulins may include both antibodies and other antibody-like molecules which generally lack antigen specificity. Polypeptides of the latter kind are, for
 10 example, produced at low levels by the lymph system and at increased levels by myelomas.

[00274] "Antibody" and "immunoglobulin" are used interchangeably in the broadest sense and include monoclonal antibodies (e.g., full length or intact monoclonal antibodies), polyclonal antibodies, multivalent antibodies, multispecific antibodies (e.g.,
 15 bispecific antibodies so long as they exhibit the desired biological activity) and may also include certain antibody fragments (as described in greater detail herein). An antibody can be human, humanized and/or affinity matured. An antibody may refer to immunoglobulins and immunoglobulin portions, whether natural or partially or wholly synthetic, such as recombinantly produced, including any portion thereof containing at least a portion of the
 20 variable region of the immunoglobulin molecule that is sufficient to form an antigen binding site. Hence, an antibody or portion thereof includes any protein having a binding domain that is homologous or substantially homologous to an immunoglobulin antigen binding site. For example, an antibody may refer to an antibody that contains two heavy chains (which can be denoted H and H') and two light chains (which can be denoted L and L'), where
 25 each heavy chain can be a full-length immunoglobulin heavy chain or a portion thereof sufficient to form an antigen binding site (e.g. heavy chains include, but are not limited to, VH, chains VH-CH1 chains and VH-CH1-CH2-CH3 chains), and each light chain can be a full-length light chain or a thereof sufficient to form an antigen binding site (e.g. light chains include, but are not limited to, VL chains and VL-CL chains). Each heavy chain (H and H')
 30 pairs with one light chain (L and L', respectively). Typically, antibodies minimally include all or at least a portion of the variable heavy (VH) chain and/or the variable light (VL) chain. The antibody also can include all or a portion of the constant region. For example, a full-length antibody is an antibody having two full-length heavy chains (e.g. VH-CH1-CH2-CH3 or VH-CH1-CH2-CH3- CH4) and two full-length light chains (VL-CL) and hinge regions,
 35 such as antibodies produced by antibody secreting B cells and antibodies with the same domains that are produced synthetically. Additionally, an "antibody" refers to a protein of

the immunoglobulin family or a polypeptide comprising fragments of an immunoglobulin that is capable of noncovalently, reversibly, and in a specific manner binding a corresponding antigen. An exemplary antibody structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one
 5 "light" (about 25 kD) and one "heavy" chain (about 50-70 kD), connected through a disulfide bond. The recognized immunoglobulin genes include the κ , λ , α , γ , δ , ϵ , and μ constant region genes, as well as the myriad immunoglobulin variable region genes. Light chains are classified as either κ or λ . Heavy chains are classified as γ , μ , α , δ , or ϵ , which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD, and IgE, respectively. The
 10 N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain (VL) and variable heavy chain (VH) refer to these regions of light and heavy chains respectively.

[00275] "Variable" refers to the fact that certain portions of the variable domains (also referred to as variable regions) differ extensively in sequence among antibodies and
 15 are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called complementarity-determining regions (CDRs) or hypervariable regions (HVRs) both in the light-chain and the heavy-chain variable domains. CDRs include those specified as Kabat, Chothia, and IMGT as
 20 shown herein within the variable region sequences. The more highly conserved portions of variable domains are called the framework (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a β -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the β -sheet structure. The CDRs in each chain are held together in close proximity by
 25 the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat *et al.*, Sequences of Proteins of Immunological Interest, Fifth Edition, National Institute of Health, Bethesda, Md. (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-
 30 dependent cellular toxicity.

[00276] Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an $F(ab')_2$ fragment that has two antigen-combining sites and is still capable of cross-
 35 linking antigen.

[00277] "Fv" refers to an antibody fragment which contains an antigen-recognition and antigen-binding site. In a two-chain Fv species, this region consists of a dimer of one heavy and one light chain variable domain in non-covalent association. In a single chain Fv (scFv) species, one heavy chain and one light chain variable domain can be covalently
 5 linked by a flexible peptide linker such that the light and heavy chains can associate in a "dimeric" structure analogous to that in a two-chain Fv (scFv) species. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the VH-VL dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv
 10 comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

[00278] The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain
 15 CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

20 [00279] The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

[00280] Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major
 25 classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these can be further divided into subclasses (isotypes), e.g., IgG₁, IgG₂, IgG₃, IgG₄, IgA₁, and IgA₂. The heavy-chain constant domains that correspond to the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

30 [00281] "Antibody fragments" comprise only a portion of an intact antibody, wherein the portion preferably retains at least one, preferably most or all, of the functions normally associated with that portion when present in an intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂, single-chain Fvs (scFv), Fv, dsFv, diabody, Fd and Fd' fragments Fab fragments, Fd fragments, scFv fragments, linear antibodies,
 35 single-chain antibody molecules, and multispecific antibodies formed from antibody fragments (see, for example, Methods in Molecular Biology, Vol 207: Recombinant

Antibodies for Cancer Therapy Methods and Protocols (2003); Chapter 1; p 3-25, Kipriyanov). Other known fragments include, but are not limited to, scFab fragments (Hust *et al.*, BMC Biotechnology (2007), 7:14). In one embodiment, an antibody fragment comprises an antigen binding site of the intact antibody and thus retains the ability to bind antigen. In another embodiment, an antibody fragment, for example one that comprises the Fc region, retains at least one of the biological functions normally associated with the Fc region when present in an intact antibody, such as FcRn binding, antibody half life modulation, ADCC function and complement binding. In one embodiment, an antibody fragment is a monovalent antibody that has an *in vivo* half life substantially similar to an intact antibody. For example, such an antibody fragment may comprise an antigen binding arm linked to an Fc sequence capable of conferring *in vivo* stability to the fragment. For another example, an antibody fragment or antibody portion refers to any portion of a full-length antibody that is less than full length but contains at least a portion of the variable region of the antibody sufficient to form an antigen binding site (e.g. one or more CDRs) and thus retains the a binding specificity and/or an activity of the full-length antibody; antibody fragments include antibody derivatives produced by enzymatic treatment of full-length antibodies, as well as synthetically, e.g. recombinantly produced derivatives.

[00282] A “dsFv” refers to an Fv with an engineered intermolecular disulfide bond, which stabilizes the VH-VL pair.

[00283] A “Fd fragment” refers to a fragment of an antibody containing a variable domain (VH) and one constant region domain (CH1) of an antibody heavy chain.

[00284] A “Fab fragment” refers to an antibody fragment that contains the portion of the full-length antibody that would result from digestion of a full-length immunoglobulin with papain, or a fragment having the same structure that is produced synthetically, e.g. recombinantly. A Fab fragment contains a light chain (containing a VL and CL portion) and another chain containing a variable domain of a heavy chain (VH) and one constant region domain portion of the heavy chain (CH1); it can be recombinantly produced.

[00285] A “F(ab')₂ fragment” refers to an antibody fragment that results from digestion of an immunoglobulin with pepsin at pH 4.0-4.5, or a synthetically, e.g. recombinantly, produced antibody having the same structure. The F(ab')₂ fragment contains two Fab fragments but where each heavy chain portion contains an additional few amino acids, including cysteine residues that form disulfide linkages joining the two fragments; it can be recombinantly produced.

[00286] A “Fab' fragment” refers to a fragment containing one half (one heavy chain and one light chain) of the F(ab')₂ fragment.

[00287] A “Fd’ fragment refers to a fragment of an antibody containing one heavy chain portion of a F(ab’)₂ fragment.

[00288] A “Fv’ fragment” refers to a fragment containing only the VH and VL domains of an antibody molecule.

5 [00289] A “scFv fragment” refers to an antibody fragment that contains a variable light chain (VL) and variable heavy chain (VH), covalently connected by a polypeptide linker in any order. The linker is of a length such that the two variable domains are bridged without substantial interference. Exemplary linkers are (Gly-Ser)_n residues with some Glu or Lys residues dispersed throughout to increase solubility.

10 [00290] Diabodies are dimeric scFv; diabodies typically have shorter peptide linkers than scFvs, and they preferentially dimerize.

[00291] “HsFv” refers to antibody fragments in which the constant domains normally present in a Fab fragment have been substituted with a heterodimeric coiled-coil domain (see, *e.g.*, Arndt *et al.* (2001) J Mol Biol. 7:312:221-228).

15 [00292] “Hypervariable region”, “HVR”, or “HV”, as well as “complementary determining region” or “CDR”, may refer to the regions of an antibody variable domain which are hypervariable in sequence and/or form structurally defined loops. Generally, antibodies comprise six hypervariable or CDR regions; three in the VH (H1, H2, H3), and three in the VL (L1, L2, L3). A number of hypervariable region or CDR delineations are in use and are
20 encompassed herein. The Kabat Complementarity Determining Regions (Kabat CDRs) are based on sequence variability and are the most commonly used (Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991)). Chothia refers instead to the location of the structural loops (Chothia and Lesk, J. Mol. Biol. 196:901-917 (1987)). The AbM hypervariable regions
25 represent a compromise between the Kabat CDRs and Chothia structural loops, (Chothia “CDRs”) and are used by Oxford Molecular's AbM antibody modeling software. The “contact” hypervariable regions are based on an analysis of the available complex crystal structures. The residues from each of these hypervariable regions are noted below.

Loop	Kabat	AbM	Chothia	Contact
L1	L24-L34	L24-L34	L26-L32	L30-L36
L2	L50-L56	L50-L56	L50-L52	L46-L55
L3	L89-L97	L89-L97	L91-L96	L89-L96
H1	H31-H35B	H26-H35B	H26-H32	H30-H35B

(Kabat Numbering)				
H1	H31-H35	H26-H35	H26-H32	H30-H35
(Chothia Numbering)				
H2	H50-H65	H50-H58	H53-H55	H47-H58
H3	H95-H102	H95-H102	H96-H101	H93-H101

IMGT refers to the international ImMunoGeneTics Information System, as described by Lefranc *et al.*, Nucl. Acids, Res. 37; D1006-D1012 (2009), including for example, IMGT designated CDRs for antibodies.

5 [00293] Hypervariable regions may comprise "extended hypervariable regions" as follows: 24-36 or 24-34 (L1), 46-56 or 50-56 (L2) and 89-97 (L3) in the VL and 26-35 (H1), 50-65 or 49-65 (H2) and 93-102, 94-102 or 95-102 (H3) in the VH. The variable domain residues are numbered according to Kabat *et al.*, *Supra* for each of these definitions.

10 [00294] "Framework" or "FR" residues are those variable domain residues other than the hypervariable region residues as herein defined. "Framework regions" (FRs) are the domains within the antibody variable region domains comprising framework residues that are located within the beta sheets; the FR regions are comparatively more conserved, in terms of their amino acid sequences, than the hypervariable regions.

15 [00295] "Monoclonal antibody" refers to an antibody from a population of substantially homogeneous antibodies, that is, for example, the individual antibodies comprising the population are identical and/or bind the same epitope(s), except for possible variants that may arise during production of the monoclonal antibody, such variants generally being present in minor amounts. Such monoclonal antibody typically includes an antibody comprising a polypeptide sequence that binds a target, wherein the target-binding

20 polypeptide sequence was obtained by a process that includes the selection of a single target binding polypeptide sequence from a plurality of polypeptide sequences. For example, the selection process can be the selection of a unique clone from a plurality of clones, such as a pool of hybridoma clones, phage clones or recombinant DNA clones. It should be understood that the selected target binding sequence can be further altered, for

25 example, to improve affinity for the target, to humanize the target binding sequence, to improve its production in cell culture, to reduce its immunogenicity *in vivo*, to create a multispecific antibody, *etc.*, and that an antibody comprising the altered target binding sequence is also a monoclonal antibody of this disclosure. In contrast to polyclonal

antibody preparations which typically include different antibodies directed against different determinants (e.g., epitopes), each monoclonal antibody of a monoclonal antibody preparation is directed against a single determinant on an antigen. In addition to their specificity, the monoclonal antibody preparations are advantageous in that they are typically uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present disclosure may be made by a variety of techniques, including, for example, the hybridoma method (e.g., Kohler *et al.*, Nature, 256:495 (1975); Harlow *et al.*, Antibodies: A Laboratory Manual, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling *et al.*, in: Monoclonal Antibodies and T-Cell Hybridomas 563-681, (Elsevier, N.Y., 1981)), recombinant DNA methods (see, e.g., U.S. Patent No. 4,816,567), phage display technologies (see, e.g., Clackson *et al.*, Nature, 352:624-628 (1991); Marks *et al.*, J. Mol. Biol., 222:581-597 (1991); Sidhu *et al.*, J. Mol. Biol. 338(2):299-310 (2004); Lee *et al.*, J. Mol. Biol. 340(5):1073-1093 (2004); Fellouse, Proc. Nat. Acad. Sci. USA 101(34):12467-12472 (2004); and Lee *et al.* J. Immunol. Methods 284(1-2):119-132 (2004), and technologies for producing human or human-like antibodies in animals that have parts or all of the human immunoglobulin loci or genes encoding human immunoglobulin sequences (see, e.g., WO 1998/24893; WO 1996/34096; WO 1996/33735; WO 1991/10741; Jakobovits *et al.*, Proc. Natl. Acad. Sci. USA, 90:2551 (1993); Jakobovits *et al.*, Nature, 362:255-258 (1993); Bruggemann *et al.*, Year in Immuno., 7:33 (1993); U.S. Patent Nos. 5,545,806; 5,569,825; 5,591,669; 5,545,807; WO 1997/17852; U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; and 5,661,016; Marks *et al.*, Bio/Technology, 10: 779-783 (1992); Lonberg *et al.*, Nature, 368: 856-859 (1994); Morrison, Nature, 368: 812-813 (1994); Fishwild *et al.*, Nature Biotechnology, 14: 845-851 (1996); Neuberger, Nature Biotechnology, 14: 826 (1996); and Lonberg and Huszar, Intern. Rev. Immunol., 13: 65-93 (1995)).

[00296] "Humanized" or "Human engineered" forms of non-human (e.g., murine) antibodies are chimeric antibodies that contain amino acids represented in human immunoglobulin sequences, including, for example, wherein minimal sequence is derived from non-human immunoglobulin. For example, humanized antibodies may be human antibodies in which some hypervariable region residues and possibly some FR residues are substituted by residues from analogous sites in non-human (e.g., rodent) antibodies. Alternatively, humanized or human engineered antibodies may be non-human (e.g., rodent) antibodies in which some residues are substituted by residues from analogous

sites in human antibodies (see, e.g., U.S. Patent No. 5,766,886). Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody, including, for example non-antibody sequences such as a chemokine, growth factor, peptide, cytokine, cell surface protein, serum protein, toxin, extracellular matrix protein, clotting factor, or secreted protein sequence. These modifications may be made to further refine antibody performance. Humanized antibodies include human engineered antibodies, for example, as described by U.S. Patent No. 5,766,886, including methods for preparing modified antibody variable domains. A humanized antibody may comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence. A humanized antibody optionally may also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, Nature 321:522-525 (1986); Riechmann *et al.*, Nature 332:323-329 (1988); and Presta, Curr. Op. Struct. Biol. 2:593-596 (1992). See also the following review articles and references cited therein: Vaswani and Hamilton, Ann. Allergy, Asthma & Immunol. 1: 105-115 (1998); Harris, Biochem. Soc. Transactions 23:1035-1038 (1995); Hurle and Gross, Curr. Op. Biotech. 5:428-433 (1994).

[00297] "Hybrid antibodies" refer to immunoglobulin molecules in which pairs of heavy and light chains from antibodies with different antigenic determinant regions are assembled together so that two different epitopes or two different antigens can be recognized and bound by the resulting tetramer.

[00298] "Chimeric" antibodies (immunoglobulins) have a portion of the heavy and/or light chain identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (see e.g., Morrison *et al.*, Proc. Natl. Acad. Sci. USA 81:6851-6855 (1984)). Humanized antibody refers to a subset of chimeric antibodies.

[00299] "Single-chain Fv" or "scFv" antibody fragments may comprise the VH and VL domains of antibody, wherein these domains are present in a single polypeptide chain. Generally, the scFv polypeptide further comprises a polypeptide linker between the VH and VL domains which enables the scFv to form the desired structure for antigen binding. For a review of scFv, see *e.g.*, Pluckthun, in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

[00300] An "antigen" refers to a predetermined antigen to which an antibody can selectively bind. The target antigen may be polypeptide, carbohydrate, nucleic acid, lipid, hapten or other naturally occurring or synthetic compound. Preferably, the target antigen is a polypeptide.

[00301] "Epitope" or "antigenic determinant", used interchangeably herein, refer to that portion of an antigen capable of being recognized and specifically bound by a particular antibody. When the antigen is a polypeptide, epitopes can be formed both from contiguous amino acids and noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained upon protein denaturing, whereas epitopes formed by tertiary folding are typically lost upon protein denaturing. An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation. Antibodies may bind to the same or a different epitope on an antigen. Antibodies may be characterized in different epitope bins. Whether an antibody binds to the same or different epitope as another antibody (*e.g.*, a reference antibody or benchmark antibody) may be determined by competition between antibodies in assays (*e.g.*, competitive binding assays).

[00302] Competition between antibodies may be determined by an assay in which the immunoglobulin under test inhibits specific binding of a reference antibody to a common antigen. Numerous types of competitive binding assays are known, for example: solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay or enzyme-linked immunosorbent assay (EIA or ELISA), sandwich competition assay including an ELISA assay (see *Stahli et al.*, *Methods in Enzymology* 9:242-253 (1983)); solid phase direct biotin-avidin EIA (see *Kirkland et al.*, *J. Immunol.* 137:3614-3619 (1986)); solid phase direct labeled assay, solid phase direct labeled sandwich assay (see *Harlow and Lane*, "Antibodies, A Laboratory Manual," Cold Spring Harbor Press (1988)); solid phase direct label RIA using I-125 label (see *Morel et al.*, *Molec. Immunol.* 25(1):7-15 (1988)); solid phase direct biotin-avidin EIA (*Cheung et al.*, *Virology* 176:546-552 (1990)); and direct labeled RIA (*Moldenhauer et al.*, *Scand. J. Immunol.*, 32:77-82 (1990)). Competition binding assays may be performed using Surface Plasmon Resonance (SPR), for example, with a Biacore[®] instrument for kinetic analysis of

binding interactions. In such an assay, a humanized antibody comprising an ultralong CDR3 of unknown epitope specificity may be evaluated for its ability to compete for binding against a comparator antibody (e.g., a BA1 or BA2 antibody as described herein). An assay may involve the use of purified antigen bound to a solid surface or cells bearing
 5 either of these, an unlabeled test immunoglobulin and a labeled reference immunoglobulin. Competitive inhibition may be measured by determining the amount of label bound to the solid surface or cells in the presence of the test immunoglobulin. Usually the test immunoglobulin is present in excess. An assay (competing antibodies) may include antibodies binding to the same epitope as the reference antibody and antibodies binding to
 10 an adjacent epitope sufficiently proximal to the epitope bound by the reference antibody for steric hindrance to occur. Usually, when a competing antibody is present in excess, it will inhibit specific binding of a reference antibody to a common antigen by at least 50%, or at least about 70%, or at least about 80%, or at least about 90%, or at least about 95%, or at least about 99% or about 100% for a competitor antibody.

15 [00303] That an antibody "selectively binds" or "specifically binds" means that the antibody reacts or associates more frequently, more rapidly, with greater duration, with greater affinity, or with some combination of the above to an antigen or an epitope than with alternative substances, including unrelated proteins. "Selectively binds" or "specifically binds" may mean, for example, that an antibody binds to a protein with a K_D of at least
 20 about 0.1 mM, or at least about 1 μ M or at least about 0.1 μ M or better, or at least about 0.01 μ M or better. Because of the sequence identity between homologous proteins in different species, specific binding can include an antibody that recognizes a given antigen in more than one species.

[00304] "Non-specific binding" and "background binding" when used in reference
 25 to the interaction of an antibody and a protein or peptide refer to an interaction that is not dependent on the presence of a particular structure (e.g., the antibody is binding to proteins in general rather than a particular structure such as an epitope).

[00305] "Diabodies" refer to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (VH) connected to a light-chain variable domain (VL) in the same polypeptide chain (VH-VL). By using a linker that is
 30 too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger et. al., Proc. Natl. Acad. Sci. USA, 90:6444-6448 (1993).

35 [00306] A "human antibody" refers to one which possesses an amino acid sequence which corresponds to that of an antibody produced by a human and/or has been

made using any of the techniques for making human antibodies as disclosed herein. This definition of a human antibody specifically excludes a humanized antibody comprising non-human antigen-binding residues.

[00307] An "affinity matured" antibody refers to one with one or more alterations in one or more CDRs thereof which result in an improvement in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). Preferred affinity matured antibodies will have nanomolar or even picomolar affinities for the target antigen. Affinity matured antibodies are produced by procedures known in the art. Marks *et al.*, *Bio/Technology* 10:779-783 (1992) describes affinity maturation by VH and VL domain shuffling. Random mutagenesis of CDR and/or framework residues is described by: Barbas *et al.*, *Proc Nat. Acad. Sci. USA* 91:3809-3813 (1994); Schier *et al.*, *Gene* 169:147-155 (1995); Yelton *et al.*, *J. Immunol.* 155:1994-2004 (1995); Jackson *et al.*, *J. Immunol.* 154(7):3310-9 (1995); and Hawkins *et al.*, *J. Mol. Biol.* 226:889-896 (1992).

[00308] Antibody "effector functions" refer to those biological activities attributable to the Fc region (a native sequence Fc region or amino acid sequence variant Fc region) of an antibody, and vary with the antibody isotype. Examples of antibody effector functions include: Clq binding and complement dependent cytotoxicity; Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down regulation of cell surface receptors (e.g. B cell receptor); and B cell activation.

[00309] "Antibody-dependent cell-mediated cytotoxicity" or "ADCC" refers to a form of cytotoxicity in which secreted Ig bound onto Fc receptors (FcRs) present on certain cytotoxic cells (e.g., Natural Killer (NK) cells, neutrophils, and macrophages) enable these cytotoxic effector cells to bind specifically to an antigen-bearing target cell and subsequently kill the target cell with cytotoxins. The antibodies "arm" the cytotoxic cells and are absolutely required for such killing. The primary cells for mediating ADCC, NK cells, express FcγRIII only, whereas monocytes express FcγRI, FcγRII and FcγRIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991). To assess ADCC activity of a molecule of interest, an *in vitro* ADCC assay, may be performed. Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, e.g., in a animal model such as that disclosed in Clynes *et al.* *Proc. Natl. Acad. Sci. USA* 95:652-656 (1998).

[00310] "Human effector cells" are leukocytes which express one or more FcRs and perform effector functions. Preferably, the cells express at least FcγRIII and perform ADCC effector function. Examples of human leukocytes which mediate ADCC include

peripheral blood mononuclear cells (PBMC), natural killer (NK) cells, monocytes, cytotoxic T cells and neutrophils; with PBMCs and NK cells being preferred. The effector cells may be isolated from a native source, *e.g.*, from blood.

[00311] "Fc receptor" or "FcR" describes a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the FcγRI, FcγRII, and FcγRIII subclasses, including allelic variants and alternatively spliced forms of these receptors. FcγRII receptors include FcγRIIA (an "activating receptor") and FcγRIIB (an "inhibiting receptor"), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor FcγRIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor FcγRIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain. (see review M. in Daeron, *Annu. Rev. Immunol.* 15:203-234 (1997)). FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-92 (1991); Capel *et al.*, *Immunomethods* 4:25-34 (1994); and de Haas *et al.*, *J. Lab. Clin. Med.* 126:330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term "FcR" herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (Guyer *et al.*, *J. Immunol.* 117:587 (1976) and Kim *et al.*, *J. Immunol.* 24:249 (1994)) and regulates homeostasis of immunoglobulins. For example, antibody variants with improved or diminished binding to FcRs have been described (see, *e.g.*, Shields *et al.* *J. Biol. Chem.* 9(2): 6591-6604 (2001)).

[00312] Methods of measuring binding to FcRn are known (see, *e.g.*, Ghetie 1997, Hinton 2004). Binding to human FcRn *in vivo* and serum half life of human FcRn high affinity binding polypeptides can be assayed, *e.g.*, in transgenic mice or transfected human cell lines expressing human FcRn, or in primates administered with the Fc variant polypeptides.

[00313] "Complement dependent cytotoxicity" or "CDC" refers to the lysis of a target cell in the presence of complement. Activation of the classical complement pathway is initiated by the binding of the first component of the complement system (C1q) to antibodies (of the appropriate subclass) which are bound to their cognate antigen. To assess complement activation, a CDC assay, for example, as described in Gazzano-Santoro *et al.*, *J. Immunol. Methods* 202:163 (1996), may be performed.

[00314] Polypeptide variants with altered Fc region amino acid sequences and increased or decreased C1q binding capability have been described (*e.g.*, see, also, Idusogie *et al.* *J. Immunol.* 164: 4178-4184 (2000)).

[00315] "Fc region-comprising polypeptide" refers to a polypeptide, such as an antibody or immunoadhesin (see definitions below), which comprises an Fc region. The C-terminal lysine (residue 447 according to the EU numbering system) of the Fc region may be removed, for example, during purification of the polypeptide or by recombinant engineering the nucleic acid encoding the polypeptide.

[00316] "Blocking" antibody or an "antagonist" antibody refers to one which inhibits or reduces biological activity of the antigen it binds. Preferred blocking antibodies or antagonist antibodies substantially or completely inhibit the biological activity of the antigen.

[00317] "Agonist" antibody refers to an antibody which mimics (*e.g.*, partially or fully) at least one of the functional activities of a polypeptide of interest.

[00318] "Acceptor human framework" refers to a framework comprising the amino acid sequence of a VL or VH framework derived from a human immunoglobulin framework, or from a human consensus framework. An acceptor human framework "derived from" a human immunoglobulin framework or human consensus framework may comprise the same amino acid sequence thereof, or may contain pre-existing amino acid sequence changes. Where pre-existing amino acid changes are present, preferably no more than 5 and preferably 4 or less, or 3 or less, pre-existing amino acid changes are present.

[00319] A "human consensus framework" refers to a framework which represents the most commonly occurring amino acid residues in a selection of human immunoglobulin VL or VH framework sequences. Generally, the selection of human immunoglobulin VL or VH sequences is from a subgroup of variable domain sequences. Generally, the subgroup of sequences is a subgroup as in Kabat *et al.*, Sequences of Proteins of Immunological Interest, Fifth Edition, NIH Publication 91-3242, Bethesda Md. (1991), vols. 1-3. In one embodiment, for the VL, the subgroup is subgroup kappa I as in Kabat *et al.*, supra. In one embodiment, for the VH, the subgroup is subgroup III as in Kabat *et al.*, supra.

[00320] "Disorder" or "disease" refers to any condition that would benefit from treatment with a substance/molecule (*e.g.*, a humanized antibody comprising an ultralong CDR3 as disclosed herein) or method disclosed herein. This includes chronic and acute disorders or diseases including those pathological conditions which predispose the mammal to the disorder in question.

[00321] "Treatment" refers to clinical intervention in an attempt to alter the natural course of the individual or cell being treated, and can be performed either for prophylaxis or during the course of clinical pathology. Desirable effects of treatment include preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease, preventing metastasis, decreasing the

rate of disease progression, amelioration or palliation of the disease state, and remission or improved prognosis. In some embodiments, antibodies disclosed herein are used to delay development of a disease or disorder.

[00322] "Individual" (e.g., a "subject") refers to a vertebrate, preferably a mammal, more preferably a human. Mammals include, but are not limited to, farm animals (such as cows), sport animals, pets (such as cats, dogs and horses), primates, mice and rats.

[00323] "Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, rodents (e.g., mice and rats), and monkeys; domestic and farm animals; and zoo, sports, laboratory, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. In some embodiments, the mammal is selected from a human, rodent, or monkey.

[00324] "Pharmaceutically acceptable" refers to approved or approvable by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, including humans.

[00325] "Pharmaceutically acceptable salt" refers to a salt of a compound that is pharmaceutically acceptable and that possesses the desired pharmacological activity of the parent compound.

[00326] "Pharmaceutically acceptable excipient, carrier or adjuvant" refers to an excipient, carrier or adjuvant that can be administered to a subject, together with at least one antibody of the present disclosure, and which does not destroy the pharmacological activity thereof and is nontoxic when administered in doses sufficient to deliver a therapeutic amount of the compound.

[00327] "Pharmaceutically acceptable vehicle" refers to a diluent, adjuvant, excipient, or carrier with which at least one antibody of the present disclosure is administered.

[00328] "Providing a prognosis", "prognostic information", or "predictive information" refer to providing information, including for example the presence of cancer cells in a subject's tumor, regarding the impact of the presence of cancer (e.g., as determined by the diagnostic methods of the present disclosure) on a subject's future health (e.g., expected morbidity or mortality, the likelihood of getting cancer, and the risk of metastasis).

[00329] Terms such as "treating" or "treatment" or "to treat" or "alleviating" or "to alleviate" refer to both 1) therapeutic measures that cure, slow down, lessen symptoms of, and/or halt progression of a diagnosed pathologic condition or disorder and 2) prophylactic or preventative measures that prevent and/or slow the development of a targeted pathologic condition or disorder. Thus those in need of treatment include those already with

the disorder; those prone to have the disorder; and those in whom the disorder is to be prevented.

[00330] "Providing a diagnosis" or "diagnostic information" refers to any information, including for example the presence of cancer cells, that is useful in
 5 determining whether a patient has a disease or condition and/or in classifying the disease or condition into a phenotypic category or any category having significance with regards to the prognosis of or likely response to treatment (either treatment in general or any particular treatment) of the disease or condition. Similarly, diagnosis refers to providing any type of diagnostic information, including, but not limited to, whether a subject is likely to
 10 have a condition (such as a tumor), whether a subject's tumor comprises cancer stem cells, information related to the nature or classification of a tumor as for example a high risk tumor or a low risk tumor, information related to prognosis and/or information useful in selecting an appropriate treatment. Selection of treatment can include the choice of a particular chemotherapeutic agent or other treatment modality such as surgery or radiation
 15 or a choice about whether to withhold or deliver therapy.

[00331] A "human consensus framework" refers to a framework which represents the most commonly occurring amino acid residues in a selection of human immunoglobulin VL or VH framework sequences. Generally, the selection of human immunoglobulin VL or VH sequences is from a subgroup of variable domain sequences. Generally, the subgroup
 20 of sequences is a subgroup as in Kabat *et al.*, Sequences of Proteins of Immunological Interest, Fifth Edition, NIH Publication 91-3242, Bethesda Md. (1991), vols. 1-3. In one embodiment, for the VL, the subgroup is subgroup kappa I as in Kabat *et al.*, supra. In one embodiment, for the VH, the subgroup is subgroup III as in Kabat *et al.*, supra.

[00332] An "acceptor human framework" for the purposes herein refers to a
 25 framework comprising the amino acid sequence of a light chain variable domain (VL) framework or a heavy chain variable domain (VH) framework derived from a human immunoglobulin framework or a human consensus framework, as defined below. An acceptor human framework "derived from" a human immunoglobulin framework or a human consensus framework may comprise the same amino acid sequence thereof, or it may
 30 contain amino acid sequence changes. In some embodiments, the number of amino acid changes are 10 or less, 9 or less, 8 or less, 7 or less, 6 or less, 5 or less, 4 or less, 3 or less, or 2 or less. In some embodiments, the VL acceptor human framework is identical in sequence to the VL human immunoglobulin framework sequence or human consensus framework sequence.

[00333] "Antigen-binding site" refers to the interface formed by one or more
 35 complementary determining regions. An antibody molecule has two antigen combining

sites, each containing portions of a heavy chain variable region and portions of a light chain variable region. The antigen combining sites can contain other portions of the variable region domains in addition to the CDRs.

[00334] An “antibody light chain” or an “antibody heavy chain” refers to a polypeptide comprising the VL or VH, respectively. The VL is encoded by the minigenes V (variable) and J (junctional), and the VH by minigenes V, D (diversity), and J. Each of VL or VH includes the CDRs as well as the framework regions. In this application, antibody light chains and/or antibody heavy chains may, from time to time, be collectively referred to as “antibody chains.” These terms encompass antibody chains containing mutations that do not disrupt the basic structure of VL or VH, as one skilled in the art will readily recognize.

[00335] “Native antibodies” refer to naturally occurring immunoglobulin molecules with varying structures. For example, native IgG antibodies are heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light chains and two identical heavy chains that are disulfide bonded. From N- to C-terminus, each heavy chain has a variable region (V H), also called a variable heavy domain or a heavy chain variable domain, followed by three constant domains (CH1, CH2, and CH3). Similarly, from N- to C-terminus, each light chain has a variable region (V L), also called a variable light domain or a light chain variable domain, followed by a constant light (CL) domain. The light chain of an antibody may be assigned to one of two types, called kappa (K) and lambda (K), based on the amino acid sequence of its constant domain.

[00336] “Combinatorial library” refers to collections of compounds formed by reacting different combinations of interchangeable chemical “building blocks” to produce a collection of compounds based on permutations of the building blocks. For an antibody combinatorial library, the building blocks are the component V, D and J regions (or modified forms thereof) from which antibodies are formed. For purposes herein, the terms “library” or “collection” are used interchangeably.

[00337] A “combinatorial antibody library” refers to a collection of antibodies (or portions thereof, such as Fabs), where the antibodies are encoded by nucleic acid molecules produced by the combination of V, D and J gene segments, particularly human V, D and J germline segments. The combinatorial libraries herein typically contain at least 50 different antibody (or antibody portions or fragment) members, typically at or about 50, 100, 500, 103, 1 x 10³, 2 x 10³, 3 x 10³, 4x 10³, 5 x 10³, 6 x 10³, 7 x 10³, 8 x 10³, 9 x 10³, 1 x 10⁴, 2 x 10⁴, 3 x 10⁴, 4x 10⁴, 5 x 10⁴, 6 x 10⁴, 7 x 10⁴, 8 x 10⁴, 9 x 10⁴, 1 x 10⁵, 2 x 10⁵, 3 x 10⁵, 4x 10⁵, 5 x 10⁵, 6 x 10⁵, 7 x 10⁵, 8 x 10⁵, 9 x 10⁵, 10⁶, 10⁷, 10⁸, 10⁹, 10¹⁰, or more different members. The resulting libraries or collections of antibodies or

portions thereof, can be screened for binding to a target protein or modulation of a functional activity.

[00338] A “human combinatorial antibody library” refers to a collection of antibodies or portions thereof, whereby each member contains a VL and VH chains or a
5 sufficient portion thereof to form an antigen binding site encoded by nucleic acid containing human germline segments produced as described herein.

[00339] A “variable germline segment” refers to V, D and J groups, subgroups, genes or alleles thereof. Gene segment sequences are accessible from known database (e.g., National Center for Biotechnology Information (NCBI), the international
10 ImMunoGeneTics information system® (IMGT), the Kabat database and the Tomlinson’s VBase database (Lefranc (2003) Nucleic Acids Res., 31:307-310; Martin *et al.*, Bioinformatics Tools for Antibody Engineering in Handbook of Therapeutic Antibodies, Wiley-VCH (2007), pp. 104-107). Tables 3-5 list exemplary human variable germline segments. Sequences of exemplary VH, DH, JH, V κ , J κ , V λ and or J λ , germline segments
15 are set forth in SEQ ID NOS: 10-451 and 868. For purposes herein, a germline segment includes modified sequences thereof, that are modified in accord with the rules of sequence compilation provided herein to permit practice of the method. For example, germline gene segments include those that contain one amino acid deletion or insertion at the 5’ or 3’ end compared to any of the sequences of nucleotides set forth in SEQ ID
20 NOS:10-451, 868.

[00340] “Compilation,” “compile,” “combine,” “combination,” “rearrange,” “rearrangement,” or other similar terms or grammatical variations thereof refers to the process by which germline segments are ordered or assembled into nucleic acid sequences representing genes. For example, variable heavy chain germline segments
25 are assembled such that the VH segment is 5’ to the DH segment which is 5’ to the JH segment, thereby resulting in a nucleic acid sequence encoding a VH chain. Variable light chain germline segments are assembled such that the VL segment is 5’ to the JL segment, thereby resulting in a nucleic acid sequence encoding a VL chain. A constant gene segment or segments also can be assembled onto the 3’ end of a nucleic acid encoding a
30 VH or VL chain.

[00341] “Linked,” or “linkage” or other grammatical variations thereof with reference to germline segments refers to the joining of germline segments. Linkage can be direct or indirect. Germline segments can be linked directly without additional nucleotides between segments, or additional nucleotides can be added to render the entire segment in-
35 frame, or nucleotides can be deleted to render the resulting segment in-frame. It is

understood that the choice of linker nucleotides is made such that the resulting nucleic acid molecule is in-frame and encodes a functional and productive antibody.

[00342] “In-frame” or “linked in-frame” with reference to linkage of human germline segments means that there are insertions and/or deletions in the nucleotide germline segments at the joined junctions to render the resulting nucleic acid molecule in-frame with the 5’ start codon (ATG), thereby producing a “productive” or functional full-length polypeptide. The choice of nucleotides inserted or deleted from germline segments, particularly at joints joining various VD, DJ and VJ segments, is in accord with the rules provided in the method herein for V(D)J joint generation. For example, germline segments are assembled such that the VH segment is 5’ to the DH segment which is 5’ to the JH segment. At the junction joining the VH and the DH and at the junction joining the DH and JH segments, nucleotides can be inserted or deleted from the individual VH, DH or JH segments, such that the resulting nucleic acid molecule containing the joined VDJ segments are in-frame with the 5’ start codon (ATG).

[00343] A portion of an antibody includes sufficient amino acids to form an antigen binding site.

[00344] A “reading frame” refers to a contiguous and non-overlapping set of three-nucleotide codons in DNA or RNA. Because three codons encode one amino acid, there exist three possible reading frames for given nucleotide sequence, reading frames 1, 2 or 3. For example, the sequence ACTGGTCA will be ACT GGT CA for reading frame 1, A CTG GTC A for reading frame 2 and AC TGG TCA for reading frame 3. Generally for practice of the method described herein, nucleic acid sequences are combined so that the V sequence has reading frame 1.

[00345] A “stop codon” refers to a three-nucleotide sequence that signals a halt in protein synthesis during translation, or any sequence encoding that sequence (e.g. a DNA sequence encoding an RNA stop codon sequence), including the amber stop codon (UAG or TAG)), the ochre stop codon (UAA or TAA)) and the opal stop codon (UGA or TGA)). It is not necessary that the stop codon signal termination of translation in every cell or in every organism. For example, in suppressor strain host cells, such as amber suppressor strains and partial amber suppressor strains, translation proceeds through one or more stop codon (e.g. the amber stop codon for an amber suppressor strain), at least some of the time.

[00346] A “variable heavy” (VH) chain or a “variable light” (VL) chain (also termed VH domain or VL domain) refers to the polypeptide chains that make up the variable domain of an antibody. For purposes herein, heavy chain germline segments are designated as VH, DH and JH, and compilation thereof results in a nucleic acid encoding a

VH chain. Light chain germline segments are designated as VL or JL, and include kappa and lambda light chains (V_{κ} and J_{κ} ; V_{λ} and J_{λ} .) and compilation thereof results in a nucleic acid encoding a VL chain. It is understood that a light chain is either a kappa or lambda light chain, but does not include a kappa/lambda combination by virtue of
 5 compilation of a V_{κ} and J_{λ} .

[00347] A “degenerate codon” refers to three-nucleotide codon that specifies the same amino acid as a codon in a parent nucleotide sequence. One of skill in the art is familiar with degeneracy of the genetic code and can identify degenerate codons.

[00348] “Diversity” with respect to members in a collection refers to the number of
 10 unique members in a collection. Hence, diversity refers to the number of different amino acid sequences or nucleic acid sequences, respectively, among the analogous polypeptide members of that collection. For example, a collection of polynucleotides having a diversity of 104 contains 104 different nucleic acid sequences among the analogous polynucleotide members. In one example, the provided collections of polynucleotides and/or polypeptides
 15 have diversities of at least at or about 102, 103, 104, 105, 106, 107, 108, 109, 1010 or more.

[00349] “Sequence diversity” refers to a representation of nucleic acid sequence similarity and is determined using sequence alignments, diversity scores, and/or sequence clustering. Any two sequences can be aligned by laying the sequences side-by-side and
 20 analyzing differences within nucleotides at every position along the length of the sequences. Sequence alignment can be assessed in silico using Basic Local Alignment Search Tool (BLAST), an NCBI tool for comparing nucleic acid and/or protein sequences. The use of BLAST for sequence alignment is well known to one of skill in the art. The Blast search algorithm compares two sequences and calculates the statistical significance of
 25 each match (a Blast score). Sequences that are most similar to each other will have a high Blast score, whereas sequences that are most varied will have a low Blast score.

[00350] A “polypeptide domain” refers to a part of a polypeptide (a sequence of three or more, generally 5 or 7 or more amino acids) that is a structurally and/or functionally distinguishable or definable. Exemplary of a polypeptide domain is a part of
 30 the polypeptide that can form an independently folded structure within a polypeptide made up of one or more structural motifs (e.g. combinations of alpha helices and/or beta strands connected by loop regions) and/or that is recognized by a particular functional activity, such as enzymatic activity or antigen binding. A polypeptide can have one, typically more than one, distinct domains. For example, the polypeptide can have one or more structural
 35 domains and one or more functional domains. A single polypeptide domain can be distinguished based on structure and function. A domain can encompass a contiguous

linear sequence of amino acids. Alternatively, a domain can encompass a plurality of non-contiguous amino acid portions, which are non-contiguous along the linear sequence of amino acids of the polypeptide. Typically, a polypeptide contains a plurality of domains. For example, each heavy chain and each light chain of an antibody molecule contains a plurality of immunoglobulin (Ig) domains, each about 110 amino acids in length.

[00351] An “Ig domain” refers to a domain, recognized as such by those in the art, that is distinguished by a structure, called the Immunoglobulin (Ig) fold, which contains two beta-pleated sheets, each containing anti-parallel beta strands of amino acids connected by loops. The two beta sheets in the Ig fold are sandwiched together by hydrophobic interactions and a conserved intra-chain disulfide bond. Individual immunoglobulin domains within an antibody chain further can be distinguished based on function. For example, a light chain contains one variable region domain (VL) and one constant region domain (CL), while a heavy chain contains one variable region domain (VH) and three or four constant region domains (CH). Each VL, CL, VH, and CH domain is an example of an immunoglobulin domain.

[00352] A “variable domain” with reference to an antibody refers to a specific Ig domain of an antibody heavy or light chain that contains a sequence of amino acids that varies among different antibodies. Each light chain and each heavy chain has one variable region domain (VL, and, VH). The variable domains provide antigen specificity, and thus are responsible for antigen recognition. Each variable region contains CDRs that are part of the antigen binding site domain and framework regions (FRs).

[00353] A “constant region domain” refers to a domain in an antibody heavy or light chain that contains a sequence of amino acids that is comparatively more conserved among antibodies than the variable region domain. Each light chain has a single light chain constant region (CL) domain and each heavy chain contains one or more heavy chain constant region (CH) domains, which include, CH1, CH2, CH3 and CH4. Full-length IgA, IgD and IgG isotypes contain CH1, CH2 CH3 and a hinge region, while IgE and IgM contain CH1, CH2 CH3 and CH4. CH1 and CL domains extend the Fab arm of the antibody molecule, thus contributing to the interaction with antigen and rotation of the antibody arms. Antibody constant regions can serve effector functions, such as, but not limited to, clearance of antigens, pathogens and toxins to which the antibody specifically binds, e.g. through interactions with various cells, biomolecules and tissues.

[00354] An “antibody or portion thereof that is sufficient to form an antigen binding site” means that the antibody or portion thereof contains at least 1 or 2, typically 3, 4, 5 or all 6 CDRs of the VH and VL sufficient to retain at least a portion of the binding specificity of the corresponding full-length antibody containing all 6 CDRs. Generally, a sufficient

antigen binding site at least requires CDR3 of the heavy chain (CDRH3). It typically further requires the CDR3 of the light chain (CDRL3). As described herein, one of skill in the art knows and can identify the CDRs based on Kabat or Chothia numbering (see, e.g., Kabat, E.A. *et al.* (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242, and Chothia, C. *et al.* (1987) J. Mol. Biol. 196:901-917). For example, based on Kabat numbering, CDR-L1 corresponds to residues L24-L34; CDR-L2 corresponds to residues L50-L56; CDR-L3 corresponds to residues L89-L97; CDR-H1 corresponds to residues H31 – H35, 35a or 35b depending on the length; CDR-H2 corresponds to residues H50-H65; and CDR-H3 corresponds to residues H95-H102.

[00355] A “peptide mimetic” refers to a peptide that mimics the activity of a polypeptide. For example, an erythropoietin (EPO) peptide mimetic is a peptide that mimics the activity of Epo, such as for binding and activation of the EPO receptor.

[00356] An “address” refers to a unique identifier for each locus in a collection whereby an addressed member (e.g. an antibody) can be identified. An addressed moiety is one that can be identified by virtue of its locus or location. Addressing can be effected by position on a surface, such as a well of a microplate. For example, an address for a protein in a microwell plate that is F9 means that the protein is located in row F, column 9 of the microwell plate. Addressing also can be effected by other identifiers, such as a tag encoded with a bar code or other symbology, a chemical tag, an electronic, such RF tag, a color-coded tag or other such identifier.

[00357] An “array” refers to a collection of elements, such as antibodies, containing three or more members.

[00358] A “spatial array” refers to an array where members are separated or occupy a distinct space in an array. Hence, spatial arrays are a type of addressable array. Examples of spatial arrays include microtiter plates where each well of a plate is an address in the array. Spatial arrays include any arrangement wherein a plurality of different molecules, e.g., polypeptides, are held, presented, positioned, situated, or supported. Arrays can include microtiter plates, such as 48-well, 96-well, 144-well, 192-well, 240-well, 288-well, 336-well, 384-well, 432-well, 480-well, 576-well, 672-well, 768-well, 864-well, 960-well, 1056-well, 1152-well, 1248-well, 1344-well, 1440-well, or 1536-well plates, tubes, slides, chips, flasks, or any other suitable laboratory apparatus. Furthermore, arrays can also include a plurality of sub-arrays. A plurality of sub-arrays encompasses an array where more than one arrangement is used to position the polypeptides. For example, multiple 96-well plates could constitute a plurality of sub-arrays and a single array.

[00359] An “addressable library” or “spatially addressed library” refers to a collection of molecules such as nucleic acid molecules or protein agents, such as antibodies, in which each member of the collection is identifiable by virtue of its address.

5 [00360] An “addressable array” refers to one in which the members of the array are identifiable by their address, the position in a spatial array, such as a well of a microtiter plate, or on a solid phase support, or by virtue of an identifiable or detectable label, such as by color, fluorescence, electronic signal (*i.e.* RF, microwave or other frequency that does not substantially alter the interaction of the molecules of interest), bar code or other symbology, chemical or other such label. Hence, in general the members of the array are
10 located at identifiable loci on the surface of a solid phase or directly or indirectly linked to or otherwise associated with the identifiable label, such as affixed to a microsphere or other particulate support (herein referred to as beads) and suspended in solution or spread out on a surface.

[00361] “An addressable combinatorial antibody library” refers to a collection of
15 antibodies in which member antibodies are identifiable and all antibodies with the same identifier, such as position in a spatial array or on a solid support, or a chemical or RF tag, bind to the same antigen, and generally are substantially the same in amino acid sequence. For purposes herein, reference to an “addressable arrayed combinatorial antibody library” means that the antibody members are addressed in an array.

20 [00362] “In silico” refers to research and experiments performed using a computer. In silico methods include, but are not limited to, molecular modeling studies, biomolecular docking experiments, and virtual representations of molecular structures and/or processes, such as molecular interactions. For purposes herein, the antibody members of a library can be designed using a computer program that selects component V, D and J germline
25 segments from among those input into the computer and joins them in-frame to output a list of nucleic acid molecules for synthesis. Thus, the recombination of the components of the antibodies in the collections or libraries provided herein, can be performed in silico by combining the nucleotide sequences of each building block in accord with software that contains rules for doing so. The process could be performed manually without a computer,
30 but the computer provides the convenience of speed.

[00363] A “database” refers to a collection of data items. For purposes herein, reference to a database is typically with reference to antibody databases, which provide a collection of sequence and structure information for antibody genes and sequences. Exemplary antibody databases include, but are not limited to, IMGT®, the international
35 ImMunoGeneTics information system (imgt.cines.fr; see *e.g.*, Lefranc *et al.* (2008) *Briefings in Bioinformatics*, 9:263-275), National Center for Biotechnology Information (NCBI), the

Kabat database and the Tomlinson's VBase database (Lefranc (2003) Nucleic Acids Res., 31:307-310; Martin *et al.*, Bioinformatics Tools for Antibody Engineering in Handbook of Therapeutic Antibodies, Wiley-VCH (2007), pp. 104-107). A database also can be created by a user to include any desired sequences. The database can be created such that the
5 sequences are inputted in a desired format (e.g., in a particular reading frame; lacking stop codons; lacking signal sequences). The database also can be created to include sequences in addition to antibody sequences.

[00364] "Screening" refers to identification or selection of an antibody or portion thereof from a collection or library of antibodies and/or portions thereof, based on
10 determination of the activity or property of an antibody or portion thereof. Screening can be performed in any of a variety of ways, including, for example, by assays assessing direct binding (e.g. binding affinity) of the antibody to a target protein or by functional assays assessing modulation of an activity of a target protein.

[00365] "Activity towards a target protein" refers to binding specificity and/or
15 modulation of a functional activity of a target protein, or other measurements that reflects the activity of an antibody or portion thereof towards a target protein.

[00366] A "target protein" refers to candidate proteins or peptides that are specifically recognized by an antibody or portion thereof and/or whose activity is modulated by an antibody or portion thereof. A target protein includes any peptide or protein that
20 contains an epitope for antibody recognition. Target proteins include proteins involved in the etiology of a disease or disorder by virtue of expression or activity. Exemplary target proteins are described herein.

[00367] "Hit" refers to an antibody or portion thereof identified, recognized or selected as having an activity in a screening assay.

[00368] "Iterative" with respect to screening means that the screening is repeated
25 a plurality of times, such as 2, 3, 4, 5 or more times, until a "Hit" is identified whose activity is optimized or improved compared to prior iterations.

[00369] "High-throughput" refers to a large-scale method or process that permits manipulation of large numbers of molecules or compounds, generally tens to hundred to
30 thousands of compounds. For example, methods of purification and screening can be rendered high-throughput. High-throughput methods can be performed manually. Generally, however, high-throughput methods involve automation, robotics or software.

[00370] Basic Local Alignment Search Tool (BLAST) is a search algorithm developed by Altschul *et al.* (1990) to separately search protein or DNA databases, for
35 example, based on sequence identity. For example, blastn is a program that compares a nucleotide query sequence against a nucleotide sequence database (e.g. GenBank).

BlastP is a program that compares an amino acid query sequence against a protein sequence database.

5 [00371] A BLAST bit score is a value calculated from the number of gaps and substitutions associated with each aligned sequence. The higher the score, the more significant the alignment.

[00372] A “human protein” refers to a protein encoded by a nucleic acid molecule, such as DNA, present in the genome of a human, including all allelic variants and conservative variations thereof. A variant or modification of a protein is a human protein if the modification is based on the wildtype or prominent sequence of a human protein.

10 [00373] “Naturally occurring amino acids” refer to the 20 L-amino acids that occur in polypeptides. The residues are those 20 α -amino acids found in nature which are incorporated into protein by the specific recognition of the charged tRNA molecule with its cognate mRNA codon in humans.

15 [00374] “Non-naturally occurring amino acids” refer to amino acids that are not genetically encoded. For example, a non-natural amino acid is an organic compound that has a structure similar to a natural amino acid but has been modified structurally to mimic the structure and reactivity of a natural amino acid. Non-naturally occurring amino acids thus include, for example, amino acids or analogs of amino acids other than the 20 naturally-occurring amino acids and include, but are not limited to, the D-isostereomers of amino acids. Exemplary non-natural amino acids are known to those of skill in the art.

20 [00375] “Nucleic acids” include DNA, RNA and analogs thereof, including peptide nucleic acids (PNA) and mixtures thereof. Nucleic acids can be single or double-stranded. When referring to probes or primers, which are optionally labeled, such as with a detectable label, such as a fluorescent or radiolabel, single-stranded molecules are contemplated. Such molecules are typically of a length such that their target is statistically unique or of low copy number (typically less than 5, generally less than 3) for probing or priming a library. Generally a probe or primer contains at least 14, 16 or 30 contiguous nucleotides of sequence complementary to or identical to a gene of interest. Probes and primers can be 10, 20, 30, 50, 100 or more nucleic acids long.

30 [00376] A “peptide” refers to a polypeptide that is from 2 to 40 amino acids in length.

[00377] The amino acids which occur in the various sequences of amino acids provided herein are identified according to their known, three-letter or one-letter abbreviations (Table 1). The nucleotides which occur in the various nucleic acid fragments are designated with the standard single-letter designations used routinely in the art.

35

[00378] An “amino acid” is an organic compound containing an amino group and a carboxylic acid group. A polypeptide contains two or more amino acids. For purposes herein, amino acids include the twenty naturally-occurring amino acids, non-natural amino acids and amino acid analogs (*i.e.*, amino acids wherein the α -carbon has a side chain).

5 [00379] “Amino acid residue” refers to an amino acid formed upon chemical digestion (hydrolysis) of a polypeptide at its peptide linkages. The amino acid residues described herein are presumed to be in the “L” isomeric form. Residues in the “D” isomeric form, which are so designated, can be substituted for any L-amino acid residue as long as the desired functional property is retained by the polypeptide. NH₂ refers to the free amino
 10 group present at the amino terminus of a polypeptide. COOH refers to the free carboxy group present at the carboxyl terminus of a polypeptide. In keeping with standard polypeptide nomenclature described in J. Biol. Chem., 243: 3552-3559 (1969), and adopted 37 C.F.R. §§ 1.821-1.822, abbreviations for amino acid residues are shown below:

SYMBOL		
1-Letter	3-Letter	AMINO ACID
Y	Tyr	Tyrosine
G	Gly	Glycine
F	Phe	Phenylalanine
M	Met	Methionine
A	Ala	Alanine
S	Ser	Serine
I	Ile	Isoleucine
L	Leu	Leucine
T	Thr	Threonine
V	Val	Valine
P	Pro	Proline
K	Lys	Lysine
H	His	Histidine
Q	Gln	Glutamine
E	Glu	Glutamic acid
Z	Glx	Glu and/or Gln
W	Trp	Tryptophan
R	Arg	Arginine
D	Asp	Aspartic acid
N	Asn	Asparagine
B	Asx	Asn and/or Asp
C	Cys	Cysteine
X	Xaa	Unknown or other

15

[00380] It should be noted that all amino acid residue sequences represented herein by formulae have a left to right orientation in the conventional direction of amino-terminus to carboxyl-terminus. In addition, the phrase “amino acid residue” is broadly

defined to include the amino acids listed in the Table of Correspondence (Table 1) and modified and unusual amino acids, such as those referred to in 37 C.F.R. §§ 1.821-1.822, and incorporated herein by reference. Furthermore, it should be noted that a dash at the beginning or end of an amino acid residue sequence indicates a peptide bond to a further sequence of one or more amino acid residues, to an amino-terminal group such as NH₂ or to a carboxyl-terminal group such as COOH. The abbreviations for any protective groups, amino acids and other compounds, are, unless indicated otherwise, in accord with their common usage, recognized abbreviations, or the IUPAC-IUB Commission on Biochemical Nomenclature (see, (1972) *Biochem.* 11:1726). Each naturally occurring L-amino acid is identified by the standard three letter code (or single letter code) or the standard three letter code (or single letter code) with the prefix "L-"; the prefix "D-" indicates that the stereoisomeric form of the amino acid is D.

[00381] An "immunoconjugate" refers to an antibody conjugated to one or more heterologous molecule(s), including but not limited to a cytotoxic agent. An immunoconjugate may include non-antibody sequences.

General techniques

[00382] The present disclosure relies on routine techniques in the field of recombinant genetics. Basic texts disclosing the general methods of use in this present disclosure include Sambrook and Russell, *Molecular Cloning: A Laboratory Manual* 3d ed. (2001); Kriegler, *Gene Transfer and Expression: A Laboratory Manual* (1990); and Ausubel *et al.*, *Current Protocols in Molecular Biology* (1994).

[00383] For nucleic acids, sizes are given in either kilobases (Kb) or base pairs (bp). These are estimates derived from agarose or polyacrylamide gel electrophoresis, from sequenced nucleic acids, or from published DNA sequences. For proteins, sizes are given in kilo-Daltons (kD) or amino acid residue numbers. Proteins sizes are estimated from gel electrophoresis, from sequenced proteins, from derived amino acid sequences, or from published protein sequences.

[00384] Oligonucleotides that are not commercially available can be chemically synthesized according to the solid phase phosphoramidite triester method first described by Beaucage and Caruthers, *Tetrahedron Letters*, 22:1859-1862 (1981), using an automated synthesizer, as described in Van Devanter *et al.*, *Nucleic Acids Res.*, 12:6159-6168 (1984). Purification of oligonucleotides is by either native polyacrylamide gel electrophoresis or by anion-exchange chromatography as described in Pearson & Reanier, *J. Chrom.*, 255:137-149 (1983). The sequence of the cloned genes and synthetic

oligonucleotides can be verified after cloning using, e.g., the chain termination method for sequencing double-stranded templates of Wallace *et al.*, *Gene*, 16:21-26 (1981).

[00385] The nucleic acids encoding recombinant polypeptides of the present disclosure may be cloned into an intermediate vector before transformation into prokaryotic or eukaryotic cells for replication and/or expression. The intermediate vector may be a prokaryote vector such as a plasmid or shuttle vector.

Humanized Antibodies with Ultralong CDR3 Sequences

[00386] To date, cattle are the only species where ultralong CDR3 sequences have been identified. However, other species, for example other ruminants, may also possess antibodies with ultralong CDR3 sequences.

[00387] Exemplary antibody variable region sequences comprising an ultralong CDR3 sequence identified in cattle include those designated as: BLV1H12 (see, SEQ ID NO: 22), BLV5B8 (see, SEQ ID NO: 23), BLV5D3 (see, SEQ ID NO: 24) and BLV8C11 (see, SEQ ID NO: 25) (see, e.g., Saini, *et al.* (1999) *Eur.J.Immunol.* 29: 2420-2426; and Saini and Kaushik (2002) *Scand. J. Immunol.* 55: 140-148); BF4E9 (see, SEQ ID NO: 26) and BF1H1 (see, SEQ ID NO: 27) (see, e.g., Saini and Kaushik (2002) *Scand. J. Immunol.* 55: 140-148); and F18 (see, SEQ ID NO: 28) (see, e.g., Berens, *et al.* (1997) *Int. Immunol.* 9: 189-199).

[00388] In an embodiment, bovine antibodies are identified and humanized. Multiple techniques exist to identify antibodies.

[00389] Antibodies of the present disclosure may be isolated by screening combinatorial libraries for antibodies with the desired activity or activities. For example, a variety of methods are known in the art for generating phage display libraries and screening such libraries for antibodies possessing the desired binding characteristics. Such methods are reviewed, e.g., in Hoogenboom *et al.* in *Methods in Molecular Biology* 178:1-37 (O'Brien *et al.*, ed., Human Press, Totowa, N.J., 2001) and further described, e.g., in the McCafferty *et al.*, *Nature* 348:552-554; Clackson *et al.*, *Nature* 352: 624-628 (1991); Marks *et al.*, *J. Mol. Biol.* 222: 581-597 (1992); Marks and Bradbury, in *Methods in Molecular Biology* 248:161-175 (Lo, ed., Human Press, Totowa, N.J., 2003); Sidhu *et al.*, *J. Mol. Biol.* 338(2): 299-310 (2004); Lee *et al.*, *J. Mol. Biol.* 340(5): 1073-1093 (2004); Fellouse, *Proc. Natl. Acad. Sci. USA* 101(34): 12467-12472 (2004); and Lee *et al.*, *J. Immunol. Methods* 284(1-2): 119-132 (2004).

[00390] In certain phage display methods, repertoires of VH and VL genes are separately cloned by polymerase chain reaction (PCR) and recombined randomly in phage libraries, which can then be screened for antigen-binding phage as described in Winter *et*

al., Ann. Rev. Immunol., 12: 433-455 (1994). Phage typically display antibody fragments, either as single-chain Fv (scFv) fragments or as Fab fragments. Libraries from immunized sources provide high-affinity antibodies to the immunogen without the requirement of constructing hybridomas. Phage display libraries of bovine antibodies may be a source of
 5 bovine antibody gene sequences, including ultralong CDR3 sequences.

[00391] Typically, a non-human antibody is humanized to reduce immunogenicity to humans, while retaining the specificity and affinity of the parental non-human antibody. Generally, a humanized antibody comprises one or more variable domains in which CDRs (or portions thereof) are derived from a non-human antibody, and FRs (or portions thereof)
 10 are derived from human antibody sequences. A humanized antibody optionally will also comprise at least a portion of a human constant region. In some embodiments, some FR residues in a humanized antibody are substituted with corresponding residues from a non-human antibody (e.g., the antibody from which the CDR residues are derived), e.g., to restore or improve antibody specificity or affinity.

[00392] Humanized antibodies and methods of making them are reviewed, e.g., in Almagro and Fransson, Front. Biosci. 13:1619-1633 (2008), and are further described, e.g., in Riechmann *et al.*, Nature 332:323-329 (1988); Queen *et al.*, Proc. Nat'l Acad. Sci. USA 86:10029-10033 (1989); U.S. Pat. Nos. 5,821,337, 7,527,791, 6,982,321, and 7,087,409; Kashmiri *et al.*, Methods 36:25-34 (2005) (describing SDR (a-CDR) grafting); Padlan, Mol.
 20 Immunol. 28:489-498 (1991) (describing "resurfacing"); Dall'Acqua *et al.*, Methods 36:43-60 (2005) (describing "FR shuffling"); and Osbourn *et al.*, Methods 36:61-68 (2005); Klimka *et al.*, Br. J. Cancer, 83:252-260 (2000) (describing the "guided selection" approach to FR shuffling); and Studnicka *et al.*, U.S. Patent No. 5,766,886.

[00393] Human variable region framework sequences that may be used for
 25 humanization include but are not limited to: framework sequences selected using the "best-fit" method (see, e.g., Sims *et al.* J. Immunol. 151:2296 (1993)); framework sequences derived from the consensus sequence of human antibodies of a particular subgroup of light or heavy chain variable regions (see, e.g., Carter *et al.* Proc. Natl. Acad. Sci. USA, 89:4285 (1992); and Presta *et al.* J. Immunol., 151:2623 (1993)); human mature (somatically
 30 mutated) framework sequences or human germline framework sequences (see, e.g., Almagro and Fransson, Front. Biosci. 13:1619-1633 (2008)); and framework sequences derived from screening FR libraries (see, e.g., Baca *et al.*, Biol. Chem. 272:10678-10684 (1997) and Rosok *et al.*, J. Biol. Chem. 271:22611-22618 (1996)).

[00394] Humanized antibodies with ultralong CDR3 sequences may also include
 35 engineered non-antibody sequences, such as cytokines or growth factors, into the CDR3 region, such that the resultant humanized antibody is effective, for example, in inhibiting

tumor metastasis. Non-antibody sequences may include an interleukin sequence, a hormone sequence, a cytokine sequence, a toxin sequence, a lymphokine sequence, a growth factor sequence, a chemokine sequence, a toxin sequence, or combinations thereof. Non-antibody sequences may be human, non-human, or synthetic. In some

5 embodiments, the cytokine or growth factor may be shown to have an antiproliferative effect on at least one cell population. Such cytokines, lymphokines, growth factors, or other hematopoietic factors include M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IFN, TNF α 1, TNF1, TNF2, G-CSF, Meg-CSF, GM-CSF, thrombopoietin, stem cell factor, and

10 erythropoietin. Additional growth factors for use in humanized antibodies and/or pharmaceutical compositions of the present disclosure include: angiogenin, bone morphogenic protein-1, bone morphogenic protein-2, bone morphogenic protein-3, bone morphogenic protein-4, bone morphogenic protein-5, bone morphogenic protein-6, bone morphogenic protein-7, bone morphogenic protein-8, bone morphogenic protein-9, bone

15 morphogenic protein-10, bone morphogenic protein-11, bone morphogenic protein-12, bone morphogenic protein-13, bone morphogenic protein-14, bone morphogenic protein-15, bone morphogenic protein receptor IA, bone morphogenic protein receptor IB, brain derived neurotrophic factor, ciliary neurotrophic factor, ciliary neurotrophic factor receptor, cytokine-induced neutrophil chemotactic factor 1, cytokine-induced neutrophil, chemotactic

20 factor 2, cytokine-induced neutrophil chemotactic factor 2, endothelial cell growth factor, endothelin 1, epidermal growth factor, epithelial-derived neutrophil attractant, fibroblast growth factor 4, fibroblast growth factor 5, fibroblast growth factor 6, fibroblast growth factor 7, fibroblast growth factor 8, fibroblast growth factor 8b, fibroblast growth factor 8c, fibroblast growth factor 9, fibroblast growth factor 10, fibroblast growth factor acidic,

25 fibroblast growth factor basic, glial cell line-derived neurotrophic factor receptor-1, glial cell line-derived neurotrophic factor receptor-2, growth related protein, growth related protein-1, growth related protein-2, growth related protein-3, heparin binding epidermal growth factor, hepatocyte growth factor, hepatocyte growth factor receptor, insulin-like growth factor I, insulin-like growth factor receptor, insulin-like growth factor II, insulin-like growth factor

30 binding protein, keratinocyte growth factor, leukemia inhibitory factor, leukemia inhibitory factor receptor-1, nerve growth factor nerve growth factor receptor, neurotrophin-3, neurotrophin-4, placenta growth factor, placenta growth factor 2, platelet-derived endothelial cell growth factor, platelet derived growth factor, platelet derived growth factor A chain, platelet derived growth factor AA, platelet derived growth factor AB, platelet

35 derived growth factor B chain, platelet derived growth factor BB, platelet derived growth factor receptor-1, platelet derived growth factor receptor-2, pre-B cell growth stimulating

factor, stem cell factor, stem cell factor receptor, transforming growth factor-1, transforming growth factor-2, transforming growth factor-1, transforming growth factor-1.2, transforming growth factor-2, transforming growth factor-3, transforming growth factor-S, latent transforming growth factor-1, transforming growth factor-1 binding protein I, transforming growth factor-1 binding protein II, transforming growth factor-1 binding protein III, tumor necrosis factor receptor type I, tumor necrosis factor receptor type II, urokinase-type plasminogen activator receptor, vascular endothelial growth factor, and chimeric proteins and biologically or immunologically active fragments thereof. Exemplary non-antibody sequences include interleukin 8 (IL-8, SEQ ID NO: 475), interleukin 21 (IL-21, SEQ ID NO: 480), CXCL12/SDF-1alpha (SEQ ID NO: 479), somatostatin (SEQ ID NO: 477), ProTx-II (SEQ ID NO: 481), chlorotoxin (SEQ ID NO: 478), and ziconotide (SEQ ID NO: 476).

[00395] A non-human antibody may be humanized to reduce immunogenicity to humans, while retaining the specificity and affinity of the parental non-human antibody. Generally, a humanized antibody comprises one or more variable domains in which HVRs, e.g., CDRs, (or portions thereof) are derived from a non-human antibody, and FRs (or portions thereof) are derived from human antibody sequences. A humanized antibody optionally will also comprise at least a portion of a human constant region. In some embodiments, some FR residues in a humanized antibody are substituted with corresponding residues from a non-human antibody (e.g., the antibody from which the HVR residues are derived), e.g., to restore or improve antibody specificity or affinity.

[00396] The humanized antibodies comprising an ultralong CDR3 as disclosed herein are preferably monoclonal. Also encompassed within the scope of the disclosure are Fab, Fab', Fab'-SH and F(ab')² fragments of the humanized antibodies comprising an ultralong CDR3 as provided herein. These antibody fragments can be created by traditional means, such as enzymatic digestion, or may be generated by recombinant techniques. Such antibody fragments may be chimeric or humanized. These fragments are useful for the diagnostic and therapeutic purposes set forth below.

[00397] Monoclonal antibodies are obtained from a population of substantially homogeneous antibodies, e.g., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Thus, the modifier "monoclonal" indicates the character of the antibody as not being a mixture of discrete antibodies.

[00398] The humanized antibodies comprising an ultralong CDR3 as disclosed herein can be made using a hybridoma cell-based method first described by Kohler *et al.*, Nature, 256:495 (1975), or may be made by recombinant DNA methods.

[00399] Hybridoma cells can be generated by fusing B cells producing a desired antibody with an immortalized cell line, usually a myeloma cell line, so that the resulting fusion cells will be an immortalized cell line that secretes a particular antibody. By the same principle, myeloma cells can be first transfected with a nucleic acid encoding a germline antibody V region and can be screened for the expression of the germline V region. Those myeloma cells with highest level of proteolytic light chain expression can be subsequently fused with B cells that produce an antibody with desired target protein specificity. The fusion cells will produce two types of antibodies: one is a heterologous antibody containing an endogenous antibody chain (either heavy or light) operably joined to the recombinant germline V region (either heavy or light), and the other is the same antibody that the parental B cells would secrete (e.g. both endogenous heavy and light chains). The operably joined heterologous heavy and light chains can be isolated by conventional methods such as chromatography and identification can be confirmed by target protein binding assays, assays identifying a unique tag of the germline polypeptide, or endopeptidase activity assays described in other sections of this disclosure. In some cases, where the heterologous antibody is the predominant type in quantity among the two types of antibodies, such isolation may not be needed.

[00400] The hybridoma cells may be seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

[00401] Preferred myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these, myeloma cell lines may be murine myeloma lines, such as those derived from MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, Calif. USA, and SP-2 or X63-Ag8-653 cells available from the American Type Culture Collection, Rockville, Md. USA. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, J. Immunol., 133:3001 (1984); Brodeur *et al.*, Monoclonal Antibody Production Techniques and Applications, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)).

[00402] Culture medium in which hybridoma cells are growing is assayed for production of humanized antibodies comprising an ultralong CDR3. For example, the binding specificity of monoclonal antibodies produced by hybridoma cells may be

determined by immunoprecipitation or by an *in vitro* binding assay, such as an enzyme-linked immunoadsorbent assay (ELISA).

[00403] The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson *et al.*, Anal. Biochem., 107:220 (1980).

5 [00404] After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, Monoclonal Antibodies: Principles and Practice, pp. 59-103 (Academic Press, 1986)). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may
10 be grown *in vivo* as ascites tumors in an animal.

[00405] The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

15 [00406] The humanized antibodies comprising an ultralong CDR3 as disclosed herein may be made by using combinatorial libraries to screen for synthetic antibody clones with the desired activity or activities. For example, synthetic antibody clones are selected by screening phage libraries containing phage that display various fragments of antibody variable regions (e.g., scFv or Fab) fused to phage coat protein. Such phage
20 libraries may be panned, for example, by affinity chromatography against the desired antigen. Clones expressing antibody fragments capable of binding to the desired antigen may be adsorbed to the antigen and thus separated from the non-binding clones in the library. The binding clones may then be eluted from the antigen, and can be further enriched by additional cycles of antigen adsorption/elution. Any of the humanized
25 antibodies comprising an ultralong CDR3 as disclosed herein may be obtained by designing a suitable antigen screening procedure to select for the phage clone of interest followed by construction of a full length antibody comprising an ultralong CDR3 clone using the VH and VL (e.g., from scFv or Fab) sequences from the phage clone of interest and suitable constant region (Fc) sequences described in Kabat *et al.*, Sequences of Proteins
30 of Immunological Interest, Fifth Edition, NIH Publication 91-3242, Bethesda Md. (1991), vols. 1-3.

[00407] The antigen-binding domain of an antibody is formed from two variable (V) regions, one each from the light (VL) and heavy (VH) chains, that both present three hypervariable loops or complementarity-determining regions (CDRs). Variable domains
35 may be displayed functionally on phage, either as single-chain Fv (scFv, also referred to as single-chain antibody (SCA)) fragments, in which VH and VL are covalently linked through

a short, flexible peptide, or as Fab fragments, in which they are each fused to a constant domain and interact non-covalently, as described in Winter *et al.*, Ann. Rev. Immunol., 12: 433-455 (1994). scFv or SCA encoding phage clones and Fab encoding phage clones may be separately or collectively referred to as "Fv phage clones" or "Fv clones".

5 [00408] Repertoires of VH and VL genes may be separately cloned by polymerase chain reaction (PCR) and recombined randomly in phage libraries, which can then be searched for antigen-binding clones as described in Winter *et al.*, Ann. Rev. Immunol., 12: 433-455 (1994). Libraries from immunized sources provide high-affinity antibodies to the immunogen without the requirement of constructing hybridomas. Alternatively, the naive
10 repertoire may be cloned to provide a single source of human antibodies to a wide range of non-self and also self antigens without any immunization as described by Griffiths *et al.*, EMBO J. 12: 725-734 (1993). Finally, naive libraries can also be made synthetically by cloning the unarranged V-gene segments from stem cells, and using PCR primers containing random sequence to encode the highly variable CDR3 regions and to
15 accomplish rearrangement *in vitro* as described by Hoogenboom and Winter, J. Mol. Biol., 227: 381-388 (1992).

[00409] Filamentous phage is used to display antibody fragments by fusion to the minor coat protein pIII. Protein pIII may include truncated forms of pIII. The antibody fragments can be displayed as single chain Fv fragments, in which VH and VL domains are
20 connected on the same polypeptide chain by a flexible polypeptide spacer, (e.g., as described by Marks *et al.*, J. Mol. Biol., 222: 581-597 (1991)), or as Fab fragments, in which one chain is fused to pIII (e.g., a truncated pIII) and the other is secreted into the bacterial host cell periplasm where assembly of a Fab-coat protein structure which becomes displayed on the phage surface by displacing some of the wild type coat proteins,
25 (e.g., as described in Hoogenboom *et al.*, Nucl. Acids Res., 19: 4133-4137 (1991)).

[00410] Nucleic acid encoding antibody variable gene segments (including VH and VL segments) are recovered from the cells of interest and may be amplified or copies made by recombinant DNA techniques (e.g., Kunkel mutagenesis). For example, in the case of rearranged VH and VL gene libraries, the desired DNA may be obtained by
30 isolating genomic DNA or mRNA from lymphocytes followed by polymerase chain reaction (PCR) with primers matching the 5' and 3' ends of rearranged VH and VL genes as described in Orlandi *et al.*, Proc. Natl. Acad. Sci. (USA), 86: 3833-3837 (1989), thereby making diverse V gene repertoires for expression. The V genes may be amplified from cDNA and genomic DNA, with back primers at the 5' end of the exon encoding the mature
35 V-domain and forward primers based within the J-segment as described in Orlandi *et al.* (1989) and in Ward *et al.*, Nature, 341: 544-546 (1989). For amplifying from cDNA, back

primers can also be based in the leader exon as described in Jones *et al.*, Biotechnol., 9: 88-89 (1991), and forward primers within the constant region as described in Sastry *et al.*, Proc. Natl. Acad. Sci. (USA), 86: 5728-5732 (1989). To enhance or maximize complementarity, degeneracy may be incorporated in the primers as described in Orlandi *et al.* (1989) or Sastry *et al.* (1989). Library diversity may be enhanced or maximized by using PCR primers targeted to each V-gene family in order to amplify available VH and VL arrangements present in the immune cell nucleic acid sample, for example, as described in the method of Marks *et al.*, J. Mol. Biol., 222: 581-597 (1991) or as described in the method of Orum *et al.*, Nucleic Acids Res., 21: 4491-4498 (1993). For cloning of the amplified DNA into expression vectors, rare restriction may can be introduced within the PCR primer as a tag at one end as described in Orlandi *et al.* (1989), or by further PCR amplification with a tagged primer as described in Clackson *et al.*, Nature, 352: 624-628 (1991).

[00411] Repertoires of synthetically rearranged V genes may be derived *in vitro* from V gene segments. Most of the human VH-gene segments have been cloned and sequenced (e.g., reported in Tomlinson *et al.*, J. Mol. Biol., 227: 776-798 (1992)), and mapped (e.g., reported in Matsuda *et al.*, Nature Genet., 3: 88-94 (1993); these cloned segments (including all the major conformations of the H1 and H2 loop) may be used to generate diverse VH gene repertoires with PCR primers encoding H3 loops of diverse sequence and length as described in Hoogenboom and Winter, J. Mol. Biol., 227: 381-388 (1992). VH repertoires may also be made with all the sequence diversity focused in a long H3 loop of a single length as described in Barbas *et al.*, Proc. Natl. Acad. Sci. USA, 89: 4457-4461 (1992). Human Vk and VL segments have been cloned and sequenced (reported in Williams and Winter, Eur. J. Immunol., 23: 1456-1461 (1993)) and can be used to make synthetic light chain repertoires. Synthetic V gene repertoires, based on a range of VH and VL folds, and L3 and H3 lengths, will encode antibodies of considerable structural diversity. Following amplification of V-gene encoding DNAs, germline V-gene segments can be rearranged *in vitro* according to the methods of Hoogenboom and Winter, J. Mol. Biol., 227: 381-388 (1992).

[00412] Repertoires of antibody fragments may be constructed by combining VH and VL gene repertoires together in several ways. Each repertoire may be created in different vectors, and the vectors recombined *in vitro*, for example, as described in Hogrefe *et al.*, Gene, 128: 119-126 (1993), or *in vivo* by combinatorial infection, for example, the loxP system described in Waterhouse *et al.*, Nucl. Acids Res., 21: 2265-2266 (1993). The *in vivo* recombination approach exploits the two-chain nature of Fab fragments to overcome the limit on library size imposed by *E. coli* transformation efficiency. Naive VH and VL repertoires are cloned separately, one into a phagemid and the other into a phage

vector. The two libraries are then combined by phage infection of phagemid-containing bacteria so that each cell contains a different combination and the library size is limited only by the number of cells present (about 10^{12} clones). Both vectors contain *in vivo* recombination signals so that the VH and VL genes are recombined onto a single replicon
 5 and are co-packaged into phage virions. These large libraries may provide large numbers of diverse antibodies of good affinity (K_d^{-1} of about 10^{-8} M).

[00413] Alternatively, the repertoires may be cloned sequentially into the same vector, for example, as described in Barbas *et al.*, Proc. Natl. Acad. Sci. USA, 88: 7978-7982 (1991), or assembled together by PCR and then cloned, for example, as described in
 10 Clackson *et al.*, Nature, 352: 624-628 (1991). PCR assembly may also be used to join VH and VL DNAs with DNA encoding a flexible peptide spacer to form single chain Fv (scFv) repertoires. In yet another technique, "in cell PCR assembly" may be used to combine VH and VL genes within lymphocytes by PCR and then clone repertoires of linked genes as described in Embleton *et al.*, Nucl. Acids Res., 20: 3831-3837 (1992).

[00414] The antibodies produced by naive libraries (either natural or synthetic) can be of moderate affinity (K_d^{-1} of about 10^6 to $10^7 M^{-1}$), but affinity maturation may also be mimicked *in vitro* by constructing and reselecting from secondary libraries as described in Winter *et al.* (1994), *supra*. For example, mutation can be introduced at random *in vitro* by using error-prone polymerase (reported in Leung *et al.*, Technique, 1: 11-15 (1989)) in the
 20 method of Hawkins *et al.*, J. Mol. Biol., 226: 889-896 (1992) or in the method of Gram *et al.*, Proc. Natl. Acad. Sci. USA, 89: 3576-3580 (1992). Additionally, affinity maturation may be performed by randomly mutating one or more CDRs, for example, using PCR with primers carrying random sequence spanning the CDR of interest, in selected individual Fv clones and screening for higher affinity clones. WO 9607754 described a method for
 25 inducing mutagenesis in a complementarity determining region of an immunoglobulin light chain to create a library of light chain genes. Another effective approach is to recombine the VH or VL domains selected by phage display with repertoires of naturally occurring V domain variants obtained from unimmunized donors and screen for higher affinity in several rounds of chain reshuffling as described in Marks *et al.*, Biotechnol., 10: 779-783
 30 (1992). This technique allows the production of antibodies and antibody fragments with affinities in the 10^{-9} M range.

[00415] The phage library samples are contacted with an immobilized protein under conditions suitable for binding of at least a portion of the phage particles with the adsorbent. Normally, the conditions, including pH, ionic strength, temperature and the like
 35 are selected to mimic physiological conditions. The phages bound to the solid phase are washed and then eluted by acid, *e.g.*, as described in Barbas *et al.*, Proc. Natl. Acad. Sci.

USA, 88: 7978-7982 (1991), or by alkali, (e.g., as described in Marks *et al.*, J. Mol. Biol., 222: 581-597 (1991)), or by antigen competition, (e.g., in a procedure similar to the antigen competition method of Clackson *et al.*, Nature, 352: 624-628 (1991)). Phages may be enriched 20-1,000-fold in a single round of selection. Moreover, the enriched phages may be grown in bacterial culture and subjected to further rounds of selection.

[00416] The efficiency of selection depends on many factors, including the kinetics of dissociation during washing, and whether multiple antibody fragments on a single phage can simultaneously engage with antigen. Antibodies with fast dissociation kinetics (and weak binding affinities) may be retained by use of short washes, multivalent phage display and high coating density of antigen in solid phase. The high density not only stabilizes the phage through multivalent interactions, but favors rebinding of phage that has dissociated. The selection of antibodies with slow dissociation kinetics (and good binding affinities) may be promoted by use of long washes and monovalent phage display as described in Bass *et al.*, Proteins, 8: 309-314 (1990) and in WO 92/09690, and a low coating density of antigen as described in Marks *et al.*, Biotechnol., 10: 779-783 (1992).

[00417] DNA encoding the hybridoma-derived monoclonal antibodies or phage display Fv clones disclosed herein is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide primers designed to specifically amplify the heavy and light chain coding regions of interest from hybridoma or phage DNA template). Once isolated, the DNA can be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of the desired monoclonal antibodies in the recombinant host cells. Recombinant expression in bacteria of antibody-encoding DNA has been described by Better *et al.*, U.S. Patent No. 6,204,023 (see also, e.g., Skerra *et al.*, Curr. Opinion in Immunol., 5: 256 (1993) and Pluckthun, Immunol. Revs, 130: 151 (1992)).

[00418] DNA encoding Fv clones as disclosed herein may be combined with known DNA sequences encoding heavy chain and/or light chain constant regions (e.g., the appropriate DNA sequences can be obtained from Kabat *et al.*, *supra*) to form clones encoding full or partial length heavy and/or light chains. It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions may be obtained from any human or animal species. A Fv clone derived from the variable domain DNA of one animal (such as human) species and then fused to constant region DNA of another animal species to form coding sequence(s) for "hybrid", full length heavy chain and/or light chain is included in the definition of "chimeric" and "hybrid" antibody as used herein. In a preferred Fv clone

embodiment, aFv clone derived from human variable DNA is fused to human constant region DNA to form coding sequence(s) for all human, full or partial length heavy and/or light chains.

[00419] DNA encoding a humanized antibody comprising an ultralong CDR3
 5 derived from a hybridoma disclosed herein may also be modified, for example, by substituting the coding sequence for human heavy- and light-chain constant domains in place of homologous murine sequences derived from the hybridoma clone (e.g., as in the method of Morrison *et al.*, Proc. Natl. Acad. Sci. USA, 81: 6851-6855 (1984)). DNA encoding a hybridoma or Fv clone-derived antibody or fragment can be further modified by
 10 covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. In this manner, "chimeric" or "hybrid" antibodies are prepared that have the binding specificity of the Fv clone or hybridoma clone-derived antibodies disclosed herein.

15 **Antibody Genes and Proteins**

[00420] The present disclosure provides antibody genes and proteins including, for example, humanized antibody genes or proteins that comprise an ultralong CDR3 sequence and/or a CDR3 scaffold. The present disclosure additionally provides VH, DH, and JH sequences useful in the preparation of ultralong CDR3 sequences. Such
 20 sequences may comprise motifs (e.g., cysteine motifs) as described herein including those as described in the many embodiments disclosed herein. In some embodiments, the antibodies disclosed herein may selectively or specifically bind to an epitope of a target protein. In some embodiments, the antibody may be an antagonist (e.g., blocking) antibody or an agonist antibody.

[00421] The variable region of the heavy and light chains are encoded by multiple
 25 germline gene segments separated by non-coding regions, or introns, and often are present on different chromosomes. For example, the genes for the human immunoglobulin heavy chain region contains approximately 65 variable (VH) genes, 27 Diversity (DH) genes, and 6 Joining (JH) genes. The human kappa (κ) and lambda (λ) light chains are
 30 also each encoded by a similar number of VL and JL gene segments, but do not include any D gene segments. Exemplary VH, DH, JH and VL (V_κ or V_λ) and JL (J_κ or J_λ) germline gene segments are set forth in WO 2010/054007.

[00422] During B cell differentiation germline DNA is rearranged whereby one DH
 and one JH gene segment of the heavy chain locus are recombined, which is followed by
 35 the joining of one VH gene segment forming a rearranged VDJ gene that encodes a VH chain. The rearrangement occurs only on a single heavy chain allele by the process of

allelic exclusion. Allelic exclusion is regulated by in-frame or “productive” recombination of the VDJ segments, which occurs in only about one-third of VDJ recombinations of the variable heavy chain. When such productive recombination events first occur in a cell, this result in production of a μ heavy chain that gets expressed on the surface of a pre- B cell and transmits a signal to shut off further heavy chain recombination, thereby preventing expression of the allelic heavy chain locus. The surface-expressed μ heavy chain also acts to activate the kappa (κ) locus for rearrangement. The lambda (λ) locus is only activated for rearrangement if the κ recombination is unproductive on both loci. The light chain rearrangement events are similar to the heavy chain, except that only the VL and JL segments are recombined. Before primary transcription of each, the corresponding constant chain gene is added. Subsequent transcription and RNA splicing leads to mRNA that is translated into an intact light chain or heavy chain.

[00423] The variable regions of antibodies confer antigen binding and specificity due to recombination events of individual germline V, D and J segments, whereby the resulting recombined nucleic acid sequences encoding the variable region domains differ among antibodies and confer antigen-specificity to a particular antibody. The variation, however, is limited to three complementarity determining regions (CDR1, CDR2, and CDR3) found within the N-terminal domain of the heavy (H) and (L) chain variable regions. The CDRs are interspersed with regions that are more conserved, termed “framework regions” (FR). The extent of the framework region and CDRs has been precisely defined (see e.g., Kabat, E.A. *et al.* (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242, and Chothia, C. *et al.* (1987) J. Mol. Biol. 196:901-917). Each VH and VL is typically composed of three CDRs and four FRs arranged from the amino terminus to carboxy terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. Sequence variability among VL and VH domains is generally limited to the CDRs, which are the regions that form the antigen binding site. For example, for the heavy chain, generally, VH genes encode the N-terminal three framework regions, the first two complete CDRs and the first part of the third CDR, the DH gene encodes the central portion of the third CDR, and the JH gene encodes the last part of the third CDR and the fourth framework region. For the light chain, the VL genes encode the first CDR and second CDR. The third CDR (CDRL3) is formed by the joining of the VL and JL gene segments. Hence, CDRs 1 and 2 are exclusively encoded by germline V gene segment sequences. The VH and VL chain CDR3s form the center of the Ag-binding site, with CDRs 1 and 2 form the outside boundaries; the FRs support the scaffold by orienting the H and L CDRs. On average, an antigen binding site typically requires at least four of the CDRs make contact with the

antigen's epitope, with CDR3 of both the heavy and light chain being the most variable and contributing the most specificity to antigen binding (see, e.g., Janis Kuby, Immunology, Third Edition, New York, W.H. Freeman and Company, 1998, pp. 115-118). CDRH3, which includes all of the D gene segment, is the most diverse component of the Ab-binding site, and typically plays a critical role in defining the specificity of the Ab. In addition to sequence variation, there is variation in the length of the CDRs between the heavy and light chains.

[00424] The constant regions, on the other hand, are encoded by sequences that are more conserved among antibodies. These domains confer functional properties to antibodies, for example, the ability to interact with cells of the immune system and serum proteins in order to cause clearance of infectious agents. Different classes of antibodies, for example IgM, IgD, IgG, IgE and IgA, have different constant regions, allowing them to serve distinct effector functions.

[00425] These natural recombination events of V, D, and J, can provide nearly 2×10^7 different antibodies with both high affinity and specificity. Additional diversity is introduced by nucleotide insertions and deletions in the joining segments and also by somatic hypermutation of V regions. The result is that there are approximately 10^{10} antibodies present in an individual with differing antigen specificities.

[00426] Antibodies include bovine antibody BLVH12 (e.g., heavy chain variable region set forth in SEQ ID NO: 482, and light chain variable region set forth in SEQ ID NO: 483); and bovine antibody BLV5B8 (e.g., heavy chain variable region set forth in SEQ ID NO: 484, and light chain variable region set forth in SEQ ID NO: 485)

Antibody Fragments

[00427] The present disclosure encompasses antibody fragments. In certain circumstances there are advantages of using antibody fragments, rather than whole antibodies. The smaller size of the fragments allows for rapid clearance, and may lead to improved access to solid tumors. Antibody fragments include, but are not limited to, Fab, Fab', Fab'-SH, F(ab')₂, Fv, and scFv fragments, and other fragments described below. For a review of certain antibody fragments, see Hudson *et al.* Nat. Med. 9:129-134 (2003). For a review of scFv fragments, see, e.g., Pluckthun, in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds., (Springer-Verlag, New York), pp. 269-315 (1994); see also WO 93/16185; and U.S. Pat. Nos. 5,571,894 and 5,587,458. For discussion of Fab and F(ab')₂ fragments comprising salvage receptor binding epitope residues and having increased in vivo half-life, see U.S. Pat. No. 5,869,046.

[00428] Diabodies are antibody fragments with two antigen binding sites that may be bivalent or bispecific. See, for example, EP 404,097; WO 1993/01161; Hudson *et al.*, Nat. Med. 9:129-134 (2003); and Hollinger *et al.*, Proc. Natl. Acad. Sci. USA 90: 6444-6448 (1993). Triabodies and tetrabodies are also described in Hudson *et al.*, Nat. Med. 9: 129134 (2003).

[00429] Single-domain antibodies are antibody fragments comprising all or a portion of the heavy chain variable domain or all or a portion of the light chain variable domain of an antibody. In certain embodiments, a single-domain antibody is a human single-domain antibody (Domantis, Inc., Waltham, Mass.; see, e.g., U.S. Pat. No. 6,248,516). Antibody fragments can be made by various techniques, including but not limited to proteolytic digestion of an intact antibody as well as production by recombinant host cells (e.g. *E. coli* or phage), as described herein.

[00430] Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, e.g., Morimoto *et al.*, Journal of Biochemical and Biophysical Methods 24:107-117 (1992); and Brennan *et al.*, Science, 229:81 (1985)). However, these fragments can now be produced directly by recombinant host cells. Fab, Fv and ScFv antibody fragments can all be expressed in and secreted from *E. coli*, thus allowing the facile production of large amounts of these fragments (see, e.g., U.S. Patent No. 6,204,023). Antibody fragments can be isolated from antibody phage libraries as discussed above. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (see, e.g., Carter *et al.*, Bio/Technology 10: 163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Fab and F(ab')₂ fragment with increased *in vivo* half-life comprising a salvage receptor binding epitope residues (see, e.g., in U.S. Patent No. 5,869,046). Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In other embodiments, the antibody of choice is a single chain Fv fragment (scFv or single chain antibody (SCA)). See WO 93/16185; U.S. Patent Nos. 5,571,894; and 5,587,458. Fv and sFv are the only species with intact combining sites that are devoid of constant regions; thus, they are suitable for reduced nonspecific binding during *in vivo* use. sFv fusion proteins may be constructed to yield fusion of an effector protein at either the amino or the carboxy terminus of an sFv. See Antibody Engineering, ed. Borrebaeck, *Supra*. The antibody fragment may also be a "linear antibody", for example, as described in U.S. Patent No. 5,641,870. Such linear antibody fragments may be monospecific or bispecific.

Humanized Antibodies

[00431] The present disclosure provides humanized antibodies comprising an ultralong CDR3. Humanized antibodies may include human engineered antibodies (see, e.g., Studnicka *et al.* (1994) Protein Eng.. 7(6) 805-814; and U.S. Patent No. 5,766,886). Various methods for humanizing non-human antibodies are known in the art. For example, a humanized antibody can have one or more amino acid residues introduced into it from a source which is human or non-human. Humanization may be performed following the method of Studnicka (see, e.g., Studnicka *et al.* (1994) Protein Eng.. 7(6) 805-814; and U.S. Patent No. 5,766,886), including the preparation of modified antibody variable domains. Humanization may alternatively be performed following the method of Winter and co-workers (Jones *et al.* (1986) Nature 321:522-525; Riechmann *et al.* (1988) Nature 332:323-327; Verhoeyen *et al.* (1988) Science 239:1534-1536), by substituting hypervariable region sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" or "human engineered" antibodies are chimeric antibodies, including wherein substantially less than an intact human variable domain has been substituted by or incorporated into the corresponding sequence from a non-human species. For example, humanized antibodies may be human antibodies in which some hypervariable region residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies. Alternatively, humanized or human engineered antibodies may be non-human (e.g., rodent) antibodies in which some residues are substituted by residues from analogous sites in human antibodies (see, e.g., Studnicka *et al.* (1994) Protein Eng.. 7(6) 805-814; and U.S. Patent No. 5,766,886).

[00432] The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is important to reduce antigenicity. For example, to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the rodent is then accepted as the human framework for the humanized antibody (Sims *et al.* (1993) J. Immunol. 151:2296; Chothia *et al.* (1987) J. Mol. Biol. 196:901). Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter *et al.* (1992) Proc. Natl. Acad. Sci. USA, 89:4285; Presta *et al.* (1993) J. Immunol., 151:2623).

[00433] It is further important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to one method, humanized antibodies are prepared by a process of analysis of

the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *e.g.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the hypervariable region residues are directly and most substantially involved in influencing antigen binding.

[00434] In some embodiments, the humanized antibodies comprising an ultralong CDR3 may be deimmunized. Methods of deimmunizing an antibody or protein are well known in the art. The immunogenicity of therapeutic proteins such as antibodies is thought to result from the presence of T-cell epitopes which can bind MHC class II molecules and generate a proliferative and cytokine response in CD4⁺ helper T-cells. These CD4⁺ helper cells then collaborate with B-cells to generate an antibody response against the therapeutic protein. Removal of the T-cell epitopes are thought to be key steps in deimmunizing a recombinant protein. T-cell epitopes can be predicted by *in silico* algorithms that identify residues required for binding MHC. Alternatively, epitopes can be identified directly by utilizing peripheral blood mononuclear cells from panels of human donors and measuring their response against the therapeutic protein when incubated with antigen presenting cells. Such *in silico* and *in vitro* systems are well known in the art [Jones TD, Crompton LJ, Carr FJ, Baker MP. *Methods Mol Biol.* 2009;525:405-23, Deimmunization of monoclonal antibodies; and Baker M, and Jones TD. The identification and removal of immunogenicity in therapeutic proteins. *Curr. Opin. Drug Discovery Dev.* 2007; (2007); 10(2): 219-227]. When peptides are identified that bind MHC II or otherwise stimulate CD4⁺ cell activation, the residues of the peptide can be mutated one by one and tested for T-cell activation until a mutation is found which disrupts MHC II binding and T-cell activation. Such mutations, when found in an individual peptide, can be encoded directly in the recombinant therapeutic protein. Incubation of the whole protein with antigen presenting cells will not induce a significant CD4⁺ response, indicating successful deimmunization.

Bispecific Antibodies

[00435] Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. For example, one of the binding specificities may be for a first antigen and the other may be for any other antigen. Exemplary bispecific antibodies may bind to two different epitopes of the same protein. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express a particular protein. These antibodies possess a binding arm specific for the particular protein and an arm which binds the cytotoxic agent (e.g., saporin, anti-interferon- α , vinca alkaloid, ricin A chain, methotrexate or radioactive isotope hapten). Bispecific antibodies may be prepared as full length antibodies or antibody fragments (e.g., F(ab')₂ bispecific antibodies).

[00436] Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy chain-light chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, Nature, 305: 537 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829, and in Traunecker *et al.*, EMBO J., 10: 3655 (1991).

[00437] According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1), containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are not of particular significance.

[00438] In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one

arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. This asymmetric structure may facilitate the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO 94/04690. For further details of generating bispecific antibodies see, for example, Suresh *et al.*, Methods in Enzymology, 121:210 (1986).

[00439] According to another approach, the interface between a pair of antibody molecules may can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the C_{H3} domain of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g., tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g., alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

[00440] Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate may be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/00373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in U.S. Patent No. 4,676,980, along with a number of cross-linking techniques.

[00441] Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies may be prepared using chemical linkage. Brennan *et al.*, Science, 229: 81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced may be used as agents for the selective immobilization of enzymes.

[00442] Recent progress has facilitated the direct recovery of Fab'-SH fragments from *E. coli*, which can be chemically coupled to form bispecific antibodies. Shalaby *et al.*, J. Exp. Med., 175: 217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and
5 subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the HER2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

[00443] Various techniques for making and isolating bispecific antibody fragments
10 directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. See, *e.g.*, Kostelny *et al.*, J. Immunol., 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-
15 oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger *et al.*, Proc. Natl. Acad. Sci. USA, 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (VH) connected to a light-chain variable domain (VL) by a linker
20 which is too short to allow pairing between the two domains on the same chain. Accordingly, the VH and VL domains of one fragment are forced to pair with the complementary VL and VH domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber *et al.*, J. Immunol.,
25 152:5368 (1994).

[00444] Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. See, *e.g.*, Tutt *et al.* J. Immunol. 147: 60 (1991).

Multivalent Antibodies

[00445] A multivalent antibody may be internalized (and/or catabolized) faster than
30 a bivalent antibody by a cell expressing an antigen to which the antibodies bind. The antibodies of the present disclosure may be multivalent antibodies (which are other than of the IgM class) with three or more antigen binding sites (*e.g.*, tetravalent antibodies), which may be produced by recombinant expression of nucleic acid encoding the polypeptide
35 chains of the antibody. The multivalent antibody may comprise a dimerization domain and three or more antigen binding sites. A preferred dimerization domain may comprise (or

consist of) an Fc region or a hinge region. In this scenario, the antibody will comprise an Fc region and three or more antigen binding sites amino-terminal to the Fe region. A preferred multivalent antibody may comprise (or consist of) three to about eight, but preferably four, antigen binding sites. The multivalent antibody comprises at least one polypeptide chain
 5 (and preferably two polypeptide chains), wherein the polypeptide chain(s) comprise two or more variable domains. For instance, the polypeptide chain(s) may comprise VD1-(X1)_n-VD2-(X2)_n-Fc, wherein VD1 is a first variable domain, VD2 is a second variable domain, Fc is one polypeptide chain of an Fc region, X1 and X2 represent an amino acid or polypeptide, and n is 0 or 1. For instance, the polypeptide chain(s) may comprise: VH-CH1-
 10 flexible linker-VH-CH1-Fc region chain; or VH-CH1-VH-CH1-Fc region chain. A multivalent antibody may preferably further comprises at least two (and preferably four) light chain variable domain polypeptides. A multivalent antibody may, for instance, comprise from about two to about eight light chain variable domain polypeptides. The light chain variable domain polypeptides may comprise a light chain variable domain and, optionally, further
 15 comprise a CL domain.

Antibody Variants

[00446] In some embodiments, amino acid sequence modification(s) of the humanized antibodies comprising an ultralong CDR3 as described herein are
 20 contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. Amino acid sequence variants of the antibody are prepared by introducing appropriate nucleotide changes into the antibody nucleic acid, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of, residues within the amino acid sequences of the
 25 antibody. Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid alterations may be introduced in the subject antibody amino acid sequence at the time that sequence is made.

[00447] A useful method for identification of certain residues or regions of the
 30 antibody that are preferred locations for mutagenesis is called "alanine scanning mutagenesis" as described by Cunningham and Wells (1989) Science, 244:1081-1085. Here, a residue or group of target residues are identified (e.g., charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with
 35 antigen. Those amino acid locations demonstrating functional sensitivity to the substitutions then are refined by introducing further or other variants at, or for, the sites of substitution.

Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation per se need not be predetermined. For example, to analyze the performance of a mutation at a given site, ala scanning or random mutagenesis is conducted at the target codon or region and the expressed immunoglobulins are screened
5 for the desired activity.

[00448] Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue
10 or the antibody fused to a cytotoxic polypeptide. Other insertional variants of the antibody molecule include the fusion to the N- or C-terminus of the antibody to an enzyme (e.g., for ADEPT) or a polypeptide which increases the serum half-life of the antibody.

[00449] Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-
15 acetylglucosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

[00450] Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may
25 also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antibody (for O-linked glycosylation sites).

[00451] Where the antibody comprises an Fc region, the carbohydrate attached thereto may be altered. For example, antibodies with a mature carbohydrate structure that lacks fucose attached to an Fc region of the antibody have been described (see, e.g., US
30 2003/0157108, US 2004/0093621. Antibodies with a bisecting N-acetylglucosamine (GlcNAc) in the carbohydrate attached to an Fc region of the antibody have been described (see, e.g., WO 2003/011878, and U.S. Patent No. 6,602,684). Antibodies with at least one galactose residue in the oligosaccharide attached to an Fc region of the antibody WO 1997/30087; see, also, WO 1998/58964 and WO 1999/22764 concerning antibodies with
35 altered carbohydrate attached to the Fc region thereof). Antigen-binding molecules with modified glycosylation have been described (see, e.g., WO 99/54342, U.S. Patent Nos.

6,602,684 and 7,517,670, and US 2004/0072290; see also, e.g., U.S. Patent Nos. 7,214,775 and 7,682,610).

[00452] The preferred glycosylation variant herein comprises an Fc region, wherein a carbohydrate structure attached to the Fc region lacks fucose. Such variants have improved ADCC function. Optionally, the Fc region further comprises one or more amino acid substitutions therein which further improve ADCC, for example, substitutions at positions 298, 333, and/or 334 of the Fc region (Eu numbering of residues). Examples of publications related to "defucosylated" or "fucose-deficient" antibodies include: US 2003/0157108; WO 2000/61739; WO 2001/29246; US 2003/0115614 (now U.S. Patent No. 6,946,292) US 2002/0164328 (now U.S. Patent No. 7,064,191); US 2004/0093621; US 2004/0132140; US 2004/0110704; US 2004/0110282 (now U.S. Patent No 7,749,753); US 2004/0109865; WO 2003/085119; WO 2003/084570; WO 2005/035586; WO 2005/035778; WO2005/053742; Okazaki *et al.* J. Mol. Biol. 336:1239-1249 (2004); Yamane-Ohnuki *et al.* Biotech. Bioeng. 87: 614 (2004). Examples of cell lines producing defucosylated antibodies include Lec13 CHO cells deficient in protein fucosylation (Ripka *et al.* Arch. Biochem. Biophys. 249:533-545 (1986); US Pat Appl No US 2003/0157108 A1, Presta, L; and WO 2004/056312 A1, Adams *et al.*, especially at Example 11), and knockout cell lines, such as alpha-1,6-fucosyltransferase gene, FUT8, knockout CHO cells (Yamane-Ohnuki *et al.* Biotech. Bioeng. 87: 614 (2004)).

[00453] Another type of variant is an amino acid substitution variant. These variants have at least one amino acid residue in the antibody molecule replaced by a different residue. The sites of greatest interest for substitutional mutagenesis include the hypervariable regions, but FR alterations are also contemplated. Conservative substitutions are shown in Table 2 under the heading of "preferred substitutions". If such substitutions result in a change in biological activity, then more substantial changes, denominated "exemplary substitutions", or as further described below in reference to amino acid classes, may be introduced and the products screened.

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	Val; Leu; Ile	Val
Arg (R)	Lys; Gln; Asn	Lys
Asn (N)	Gln; His; Asp, Lys; Arg	Gln
Asp (D)	Glu; Asn	Glu
Cys (C)	Ser; Ala	Ser
Gln (Q)	Asn; Glu	Asn
Glu (E)	Asp; Gln	Asp
Gly (G)	Ala	Ala

His (H)	Asn; Gln; Lys; Arg	Arg
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Leu
Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe	Ile
Lys (K)	Arg; Gln; Asn	Arg
Met (M)	Leu; Phe; Ile	Leu
Phe (F)	Trp; Leu; Val; Ile; Ala; Tyr	Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Val; Ser	Ser
Trp (W)	Tyr; Phe	Tyr
Tyr (Y)	Trp; Phe; Thr; Ser	Phe
Val (V)	Ile; Leu; Met; Phe; Ala; Norleucine	Leu

[00454] Substantial modifications in the biological properties of the antibody are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties: (1) hydrophobic: norleucine, met, ala, val, leu, ile; (2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln; (3) acidic: asp, glu; (4) basic: his, lys, arg; (5) residues that influence chain orientation: gly, pro; and (6) aromatic: trp, tyr, phe.

[00455] Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

[00456] One type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody (*e.g.*, a humanized or human antibody). Generally, the resulting variant(s) selected for further development will have improved biological properties relative to the parent antibody from which they are generated. A convenient way for generating such substitutional variants involves affinity maturation using phage display. Briefly, several hypervariable region sites (*e.g.*, 6-7 sites) are mutated to generate all possible amino acid substitutions at each site. The antibodies thus generated are displayed from filamentous phage particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed variants are then screened for their biological activity (*e.g.*, binding affinity) as herein disclosed. In order to identify candidate hypervariable region sites for modification, alanine scanning mutagenesis can be performed to identify hypervariable region residues contributing significantly to antigen binding. Alternatively, or additionally, it may be beneficial to analyze a crystal structure of

the antigen-antibody complex to identify contact points between the antibody and antigen. Such contact residues and neighboring residues are candidates for substitution according to the techniques elaborated herein. Once such variants are generated, the panel of variants is subjected to screening as described herein and antibodies with superior
5 properties in one or more relevant assays may be selected for further development.

[00457] Nucleic acid molecules encoding amino acid sequence variants of the antibody are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-directed)
10 mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of the antibody.

[00458] It may be desirable to introduce one or more amino acid modifications in an Fc region of the immunoglobulin polypeptides disclosed herein, thereby generating a Fc region variant. The Fc region variant may comprise a human Fc region sequence (e.g., a human IgG1, IgG2, IgG3 or IgG4 Fc region) comprising an amino acid modification (e.g., a
15 substitution) at one or more amino acid positions including that of a hinge cysteine.

[00459] In accordance with this description and the teachings of the art, it is contemplated that in some embodiments, an antibody used in methods disclosed herein may comprise one or more alterations as compared to the wild type counterpart antibody, e.g., in the Fc region. These antibodies would nonetheless retain substantially the same
20 characteristics required for therapeutic utility as compared to their wild type counterpart. For example, it is thought that certain alterations can be made in the Fc region that would result in altered (e.g., either improved or diminished) Clq binding and/or Complement Dependent Cytotoxicity (CDC), e.g., as described in WO99/51642. See also Duncan & Winter Nature 322:738-40 (1988); U.S. Patent No. 5,648,260; U.S. Patent No. 5,624,821; and WO94/29351 concerning other examples of Fc region variants. WO00/42072 and WO
25 2004/056312 describe antibody variants with improved or diminished binding to FcRs. See, also, Shields *et al.* J. Biol. Chem. 9(2): 6591-6604 (2001). Antibodies with increased half lives and improved binding to the neonatal Fc receptor (FcRn), which is responsible for the transfer of maternal IgGs to the fetus (Guyer *et al.*, J. Immunol. 117:587 (1976) and Kim *et al.*, J. Immunol. 24:249 (1994)), are described in US2005/0014934 (Hinton *et al.*). These antibodies comprise an Fc region with one or more substitutions therein which improve binding of the Fc region to FcRn. Polypeptide variants with altered Fc region amino acid sequences and increased or decreased Clq binding capability are described in U.S. Patent
30 No. 6,194,551, WO99/51642. See, also, Idusogie *et al.* J. Immunol. 164:4178-4184 (2000).

[00460] In certain embodiments, the present disclosure contemplates an antibody variant that possesses some but not all effector functions, which make it a desirable candidate for applications in which the half life of the antibody in vivo is important yet certain effector functions (such as complement and ADCC) are unnecessary or deleterious.

5 In vitro and/or in vivo cytotoxicity assays can be conducted to confirm the reduction/depletion of CDC and/ or ADCC activities. For example, Fc receptor (FcR) binding assays can be conducted to ensure that the antibody lacks FcγR binding (hence likely lacking ADCC activity), but retains FcRn binding ability. The primary cells for mediating ADCC, NK cells, express FcγRIII only, whereas monocytes express FcγRI, FcγRII and FcγRIII. FcR
10 expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-492 (1991). Non-limiting examples of in vitro assays to assess ADCC activity of a molecule of interest is described in U.S. Pat. No. 5,500,362 (see, e.g. Hellstrom, I. *et al.* *Proc. Nat'l Acad. Sci. USA* 83:7059-7063 (1986)) and Hellstrom, I *et al.*, *Proc. Nat'l Acad. Sci. USA* 82:1499-1502 (1985); 5,821,337 (see,
15 Bruggemann, M. *et al.*, *Exp. Med.* 166:1351-1361 (1987)). Alternatively, non-radioactive assays methods may be employed (see, for example, ACTITM non-radioactive cytotoxicity assay for flow cytometry (CellTect1r1ology, Inc. Mountain View, Calif.; and CytoTox 96® non-radioactive cytotoxicity assay (Promega, Madison, Wis.). Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells.
20 Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed in vivo, e.g., in a animal model such as that disclosed in Clynes *et al.* *Proc. Nat'l Acad. Sci. USA* 95:652-656 (1998). C1q binding assays may also be carried out to confirm that the antibody is unable to bind C1q and hence lacks CDC activity. See, e.g., C1q and C3c binding ELISA in WO 2006/029879 and WO 2005/ 100402. To assess complement
25 activation, a CDC assay may be performed (see, for example, Gazzano-Santoro *et al.*, *Immunol. Methods* 202:163 (1996); Cragg, M. S. *et al.*, *Blood* 101:1045-1052 (2003); and Cragg, M. S, and M. J. Glennie, *Blood* 103:2738-2743 (2004)). FcRn binding and in vivo clearance/half life determinations can also be performed using methods known in the art (see, e.g., Petkova, S. B. *et al.*, *Int'l Immunol.* 18(12):1759-1769 (2006)).

30 [00461] Antibodies with reduced effector function include those with substitution of one or more of Fc region residues 238, 265, 269, 270, 297, 327 and 329 (U.S. Pat. No. 6,737, 056). Such Fc mutants include Fc mutants with substitutions at two or more of amino acid positions 265, 269, 270, 297 and 327, including the so-called "DANA" Fc mutant with substitution of residues 265 and 297 to alanine (U.S. Pat. No. 7,332,581).

[00462] Certain antibody variants with improved or diminished binding to FcRs are described. (See, e.g., U.S. Pat. No. 6,737,056; WO 2004/056312, and Shields *et al.*, Biol. Chem. 9(2): 6591-6604 (2001).)

[00463] In certain embodiments, an antibody variant comprises an Fc region with one or more amino acid substitutions which improve ADCC, e.g., substitutions at positions 298, 333, and/or 334 of the Fc region (EU numbering of residues).

[00464] In some embodiments, alterations are made in the Fc region that result in altered (*i.e.*, either improved or diminished) C1q binding and/or Complement Dependent Cytotoxicity (CDC), e.g., as described in U.S. Pat. No. 6,194,551, WO 99/51642, and Idusogie *et al.* Immunol. 164: 41784184 (2000).

[00465] Antibodies with increased half lives and improved binding to the neonatal Fc receptor (FcRn), which is responsible for the transfer of maternal IgGs to the fetus (Guyer *et al.*, Immunol. 117:587 (1976) and Kim *et al.*, Immunol. 24:249 (1994)), are described in US2005/0014934A1 (Hinton *et al.*). Those antibodies comprise an Fc region with one or more substitutions therein which improve binding of the Fc region to FcRn. Such Fc variants include those with substitutions at one or more of Fc region residues: 238, 256, 265, 272, 286,303,305, 307, 311,312, 317, 340,356, 360, 362, 376, 378, 380, 382, 413, 424 or 434, e.g., substitution of Fc region residue 434 (U.S. Pat. No. 7,371,826).

[00466] See also Duncan & Winter, Nature 322:738-40 (1988); U.S. Pat. No. 5,648,260; U.S. Pat. No. 5,624,821; and WO 94/29351 concerning other examples of Fc region variants.

Antibody Derivatives

[00467] The humanized antibodies comprising an ultralong CDR3 as disclosed herein may be further modified to contain additional nonproteinaceous moieties that are known in the art and readily available. Preferably, the moieties suitable for derivatization of the antibody are water soluble polymers. Non-limiting examples of water soluble polymers include, but are not limited to, polyethylene glycol (PEG), copolymers of ethylene glycol/propylene glycol, carboxymethylcellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone, poly-1,3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids (either homopolymers or random copolymers), and dextran or poly(n-vinyl pyrrolidone)polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols (*e.g.*, glycerol), polyvinyl alcohol, and mixtures thereof. Polyethylene glycol propionaldehyde may have advantages in manufacturing due to its stability in water. The polymer may be of any molecular weight, and may be branched or unbranched. The number of polymers attached to the antibody

may vary, and if more than one polymers are attached, they can be the same or different molecules. In general, the number and/or type of polymers used for derivatization can be determined based on considerations including, but not limited to, the particular properties or functions of the antibody to be improved, whether the antibody derivative will be used in
 5 a therapy under defined conditions, *etc.*

Vectors, Host Cells and Recombinant Methods

[00468] For recombinant production of an antibody or fragment thereof as disclosed herein, the nucleic acid encoding it is isolated and inserted into a replicable
 10 vector for further cloning (amplification of the DNA) or for expression. DNA encoding the antibody is readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the antibody). In an exemplary embodiment, nucleic acid encoding a humanized antibody comprising an ultralong CDR3, a variable region comprising an
 15 ultralong CDR3, or an ultralong CDR3, is isolated and inserted into a replicable vector for further cloning (amplification of the DNA) or for expression. Many vectors are available. The choice of vector depends in part on the host cell to be used. Generally, preferred host cells are of either prokaryotic or eukaryotic (generally mammalian) origin. It will be appreciated that constant regions of any isotype can be used for this purpose, including
 20 IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species.

[00469] Expression vectors containing regulatory elements from eukaryotic viruses are typically used in eukaryotic expression vectors, *e.g.*, SV40 vectors, papilloma virus vectors, and vectors derived from Epstein-Barr virus. Other exemplary eukaryotic vectors
 25 include pMSG, pAV009/A+, pMTO10/A+, pMAMneo-5, baculovirus pDSVE, and any other vector allowing expression of proteins under the direction of the CMV promoter, SV40 early promoter, SV40 later promoter, metallothionein promoter, murine mammary tumor virus promoter, Rous sarcoma virus promoter, polyhedrin promoter, or other promoters shown effective for expression in eukaryotic cells.

[00470] Some expression systems have markers that provide gene amplification such as thymidine kinase and dihydrofolate reductase. Alternatively, high yield expression systems not involving gene amplification are also suitable, such as using a baculovirus vector in insect cells, with a nucleic acid sequence encoding a partially human ultralong CDR3 antibody chain under the direction of the polyhedrin promoter or other strong
 35 baculovirus promoters.

a. Generating Antibodies Using Prokaryotic or Eukaryotic Host Cells:

i. Vector Construction

[00471] Polynucleotide sequences encoding polypeptide components of the antibodies disclosed herein can be obtained using standard recombinant techniques. Desired polynucleotide sequences may be isolated and sequenced from antibody
5 producing cells such as hybridoma cells. Alternatively, polynucleotides can be synthesized using nucleotide synthesizer or PCR techniques. Once obtained, sequences encoding the polypeptides are inserted into a recombinant vector capable of replicating and expressing heterologous polynucleotides in prokaryotic hosts. Many vectors that are available and known in the art can be used for the purpose of the present disclosure. Selection of an
10 appropriate vector will depend mainly on the size of the nucleic acids to be inserted into the vector and the particular host cell to be transformed with the vector. Each vector contains various components, depending on its function (amplification or expression of heterologous polynucleotide, or both) and its compatibility with the particular host cell in which it resides. The vector components generally include, but are not limited to: an origin of replication, a
15 selection marker gene, a promoter, a ribosome binding site (RBS), a signal sequence, the heterologous nucleic acid insert and a transcription termination sequence. Additionally, V regions comprising an ultralong CDR3 may optionally be fused to a C-region to produce an antibody comprising constant regions.

[00472] In general, plasmid vectors containing replicon and control sequences
20 which are derived from species compatible with the host cell are used in connection with these hosts. The vector ordinarily carries a replication site, as well as marking sequences which are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is typically transformed using pBR322, a plasmid derived from an *E. coli* species. pBR322 contains genes encoding ampicillin (Amp) and tetracycline (Tet) resistance and
25 thus provides easy means for identifying transformed cells. pBR322, its derivatives, or other microbial plasmids or bacteriophage may also contain, or be modified to contain, promoters which can be used by the microbial organism for expression of endogenous proteins. Examples of pBR322 derivatives used for expression of particular antibodies have been described (see, e.g., U.S. Patent No. 5,648,237).

[00473] In addition, phage vectors containing replicon and control sequences that
30 are compatible with the host microorganism can be used as transforming vectors in connection with these hosts. For example, bacteriophage such as λ GEM™-11 may be utilized in making a recombinant vector which can be used to transform susceptible host cells such as *E. coli* LE392.

[00474] The expression vectors disclosed herein may comprise two or more
35 promoter-cistron pairs, encoding each of the polypeptide components. A promoter is an

untranslated regulatory sequence located upstream (5') to a cistron that modulates its expression. Prokaryotic promoters typically fall into two classes, inducible and constitutive. Inducible promoter is a promoter that initiates increased levels of transcription of the cistron under its control in response to changes in the culture condition, *e.g.*, the presence or
5 absence of a nutrient or a change in temperature.

[00475] A large number of promoters recognized by a variety of potential host cells are well known. The selected promoter can be operably linked to cistron DNA encoding the light or heavy chain by removing the promoter from the source DNA via restriction enzyme digestion and inserting the isolated promoter sequence into the vector disclosed herein.

10 Both the native promoter sequence and many heterologous promoters may be used to direct amplification and/or expression of the target genes. In some embodiments, heterologous promoters are utilized, as they generally permit greater transcription and higher yields of expressed target gene as compared to the native target polypeptide promoter.

15 [00476] Promoters suitable for use with prokaryotic hosts include: an ara B promoter, a PhoA promoter, β -galactamase and lactose promoter systems, a tryptophan (*trp*) promoter system and hybrid promoters such as the *tac* or the *trc* promoter. However, other promoters that are functional in bacteria (such as other known bacterial or phage promoters) are suitable as well. Their nucleotide sequences have been published, thereby
20 enabling a skilled worker operably to ligate them to cistrons encoding the target light and heavy chains (*e.g.*, Siebenlist *et al.* (1980) *Cell* 20: 269) using linkers or adaptors to supply any required restriction sites.

[00477] Suitable bacterial promoters are well known in the art and fully described in scientific literature such as Sambrook and Russell, *supra*, and Ausubel *et al.*, *supra*.
25 Bacterial expression systems for expressing antibody chains of the recombinant catalytic polypeptide are available in, *e.g.*, *E. coli*, *Bacillus sp.*, and *Salmonella* (Palva *et al.*, *Gene*, 22:229-235 (1983); Mosbach *et al.*, *Nature*, 302:543-545 (1983)).

[00478] In one aspect disclosed herein, each cistron within the recombinant vector comprises a secretion signal sequence component that directs translocation of the
30 expressed polypeptides across a membrane. In general, the signal sequence may be a component of the vector, or it may be a part of the target polypeptide DNA that is inserted into the vector. The signal sequence should be one that is recognized and processed (*e.g.*, cleaved by a signal peptidase) by the host cell. For prokaryotic host cells that do not recognize and process the signal sequences native to the heterologous polypeptides, the
35 signal sequence is substituted by a prokaryotic signal sequence selected, for example *PeIB*, *OmpA*, alkaline phosphatase, penicillinase, *Ipp*, or heat-stable enterotoxin II (*STII*)

leaders, LamB, PhoE, and MBP. In one embodiment disclosed herein, the signal sequences used in both cistrons of the expression system are STII signal sequences or variants thereof.

[00479] In another aspect, the production of the immunoglobulins according to the disclosure can occur in the cytoplasm of the host cell, and therefore does not require the presence of secretion signal sequences within each cistron. In that regard, immunoglobulin light and heavy chains are expressed, folded and assembled to form functional immunoglobulins within the cytoplasm. Certain host strains (*e.g.*, the *E. coli* trxB-strains) provide cytoplasm conditions that are favorable for disulfide bond formation, thereby permitting proper folding and assembly of expressed protein subunits (see *e.g.*, Proba and Pluckthun Gene, 159:203 (1995)).

[00480] Suitable host cells for cloning or expression of antibody-encoding vectors include prokaryotic or eukaryotic cells described herein. In one embodiment, the host cell is eukaryotic, *e.g.* a Chinese Hamster Ovary (CHO) cell, Human Embryonic Kidney (HEK) cell or lymphoid cell (*e.g.*, YO, NSO, Sp20 cell). For example, antibodies may be produced in bacteria, in particular when glycosylation and Fc effector function are not needed. For expression of antibody fragments and polypeptides in bacteria, see, *e.g.*, U.S. Pat. Nos. 5,648,237, 5,789,199, and 5,840,523. (See also Charlton, Methods in Molecular Biology, Vol. 248 (B.K.C. Lo, ed., Humana Press, Totowa, N.J., 2003), pp. 245-254, describing expression of antibody fragments in *E. coli*.) After expression, the antibody may be isolated from the bacterial cell paste in a soluble fraction and can be further purified. In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for antibody-encoding vectors, including fungi and yeast strains whose glycosylation pathways have been "humanized," resulting in the production of an antibody with a partially or fully human glycosylation pattern. See Gemgross, Nat. Biotech. 22: 1409-1414 (2004), and Li *et al.*, Nat. Biotech. 24:210-215 (2006). Suitable host cells for the expression of glycosylated antibody are also derived from multicellular organisms (invertebrates and vertebrates). Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains have been identified which may be used in conjunction with insect cells, particularly for transfection of *Spodoptera frugiperda* cells. These examples are illustrative rather than limiting. Methods for constructing derivatives of any of the above-mentioned bacteria having defined genotypes are known in the art and described in, for example, Bass *et al.*, Proteins, 8:309-314 (1990). It is generally necessary to select the appropriate bacteria taking into consideration replicability of the replicon in the cells of a bacterium. For example, *E. coli*, *Serratia*, or *Salmonella* species can be suitably used as the host when well known plasmids such as pBR322, pBR325, pACYC177, or

pKN410 are used to supply the replicon. Typically the host cell should secrete minimal amounts of proteolytic enzymes, and additional protease inhibitors may desirably be incorporated in the cell culture.

[00481] Plant cell cultures can also be utilized as hosts. See, e.g. U.S. Pat. Nos. 5,959,177, 6,040,498, 6,420,548, 7,125, 978, and 6,417,429 (describing PLANTIBODIESTM technology for producing antibodies in transgenic plants). Vertebrate cells may also be used as hosts. For example, mammalian cell lines that are adapted to grow in suspension may be useful. Other examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7); human embryonic kidney line (293 or 293 cells as described, e.g., in Graham *et al.*, Gen VII'01. 36:59 (1977)); baby hamster kidney cells (BHK); mouse sertoli cells (TM4 cells as described, e.g., in Mather, Biol. Reprod. 23:243-251 (1980)); monkey kidney cells (CV1); African green monkey kidney cells (V ERO-76); human cervical carcinoma cells (HELA); canine kidney cells (MDCK; buffalo rat liver cells (BRL 3A); human lung cells (W138); human liver cells (Hep G2); mouse mammary tumor (MMT 060562); TR1 cells, as described, e.g., in Mather *et al.*, Annals NI'.Acad. Sci. 383:44-68 (1982); MRC 5 cells; and FS4 cells. Other useful mammalian host cell lines include Chinese hamster ovary (CHO) cells, including DHFR⁺ CHO cells (Urlaub *et al.*, Proc. Natl. Acad. Sci. USA 77:4216 (1980)); and myeloma cell lines such as YO, NSO and Sp2/0. For a review of certain mammalian host cell lines suitable for antibody production, see, e.g., Yazaki and Wu, Methods in Molecular Biology, Vol. 248 (B.K.C. Lo, ed., Humana Press, Totowa, N.J.), pp. 255-268 (2003).

[00482] In one such embodiment, a host cell comprises (e.g., has been transformed with): (1) a vector comprising a nucleic acid that encodes an amino acid sequence comprising the VL of the antibody and an amino acid sequence comprising the VH of the antibody, or (2) a first vector comprising a nucleic acid that encodes an amino acid sequence comprising the VL of the antibody and a second vector comprising a nucleic acid that encodes an amino acid sequence comprising the VH of the antibody.

ii. Antibody Production

[00483] For recombinant production of a partially human ultralong CDR3 antibody, nucleic acid encoding a humanized antibody comprising an ultralong CDR3 is inserted into one or more expression vectors for further cloning and/or expression in a host cell. Such nucleic acid may be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the antibody). Host cells are transformed with such expression vectors and cultured in conventional nutrient media modified as appropriate for

inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

[00484] Transformation means introducing DNA into the prokaryotic host so that the DNA is replicable, either as an extrachromosomal element or by chromosomal
5 integrant. Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride is generally used for bacterial cells that contain substantial cell-wall barriers. Another method for transformation employs polyethylene glycol/DMSO. Yet another technique used is electroporation.

10 [00485] Prokaryotic cells used to produce the polypeptides disclosed herein are grown in media known in the art and suitable for culture of the selected host cells. Examples of suitable media include luria broth (LB) plus necessary nutrient supplements. In some embodiments, the media also contains a selection agent, chosen based on the construction of the expression vector, to selectively permit growth of prokaryotic cells
15 containing the expression vector. For example, ampicillin is added to media for growth of cells expressing ampicillin resistant gene.

[00486] Any necessary supplements besides carbon, nitrogen, and inorganic phosphate sources may also be included at appropriate concentrations introduced alone or as a mixture with another supplement or medium such as a complex nitrogen source.
20 Optionally the culture medium may contain one or more reducing agents selected from the group consisting of glutathione, cysteine, cystamine, thioglycollate, dithioerythritol and dithiothreitol.

[00487] The prokaryotic host cells are cultured at suitable temperatures. For *E. coli* growth, for example, the preferred temperature ranges from about 20° C. to about 39° C.,
25 more preferably from about 25° C. to about 37° C., even more preferably at about 30° C. The pH of the medium may be any pH ranging from about 5 to about 9, depending mainly on the host organism. For *E. coli*, the pH is preferably from about 6.8 to about 7.4, and more preferably about 7.0.

[00488] If an inducible promoter is used in the expression vector disclosed herein,
30 protein expression is induced under conditions suitable for the activation of the promoter. For example, an ara B or phoA promoter may be used for controlling transcription of the polypeptides. A variety of inducers may be used, according to the vector construct employed, as is known in the art.

[00489] The expressed polypeptides of the present disclosure are secreted into
35 and recovered from the periplasm of the host cells or transported into the culture media. Protein recovery from the periplasm typically involves disrupting the microorganism,

generally by such means as osmotic shock, sonication or lysis. Once cells are disrupted, cell debris or whole cells may be removed by centrifugation or filtration. The proteins may be further purified, for example, by affinity resin chromatography. Alternatively, proteins that are transported into the culture media may be isolated therein. Cells may be removed
5 from the culture and the culture supernatant being filtered and concentrated for further purification of the proteins produced. The expressed polypeptides can be further isolated and identified using commonly known methods such as polyacrylamide gel electrophoresis (PAGE) and Western blot assay.

[00490] Antibody production may be conducted in large quantity by a fermentation
10 process. Various large-scale fed-batch fermentation procedures are available for production of recombinant proteins. Large-scale fermentations have at least 1000 liters of capacity, preferably about 1,000 to 100,000 liters of capacity. These fermentors use agitator impellers to distribute oxygen and nutrients, especially glucose (a preferred carbon/energy source). Small scale fermentation refers generally to fermentation in a
15 fermentor that is no more than approximately 100 liters in volumetric capacity, and can range from about 1 liter to about 100 liters.

[00491] In a fermentation process, induction of protein expression is typically initiated after the cells have been grown under suitable conditions to a desired density, *e.g.*, an OD550 of about 180-220, at which stage the cells are in the early stationary phase.
20 A variety of inducers may be used, according to the vector construct employed, as is known in the art and described above. Cells may be grown for shorter periods prior to induction. Cells are usually induced for about 12-50 hours, although longer or shorter induction time may be used.

[00492] To improve the production yield and quality of the polypeptides disclosed
25 herein, various fermentation conditions can be modified. For example, to improve the proper assembly and folding of the secreted antibody polypeptides, additional vectors overexpressing chaperone proteins, such as Dsb proteins (DsbA, DsbB, DsbC, DsbD and or DsbG) or FkpA (a peptidylprolyl *cis*,*trans*-isomerase with chaperone activity) may be used to co-transform the host prokaryotic cells. The chaperone proteins have been
30 demonstrated to facilitate the proper folding and solubility of heterologous proteins produced in bacterial host cells. (see *e.g.*, Chen *et al.* (1999) J Bio Chem 274:19601-19605; U.S. Patent No. 6,083,715; U.S. Patent No. 6,027,888; Bothmann and Pluckthun (2000) J. Biol. Chem. 275:17100-17105; Ramm and Pluckthun (2000) J. Biol. Chem. 275:17106-17113; Arie *et al.* (2001) Mol. Microbiol. 39:199-210).

[00493] To minimize proteolysis of expressed heterologous proteins (especially
35 those that are proteolytically sensitive), certain host strains deficient for proteolytic

enzymes can be used for the present disclosure. For example, host cell strains may be modified to effect genetic mutation(s) in the genes encoding known bacterial proteases such as Protease III, OmpT, DegP, Tsp, Protease I, Protease Mi, Protease V, Protease VI and combinations thereof. Some *E. coli* protease-deficient strains are available (see, e.g.,
5 Joly *et al.* (1998), *supra*; U.S. Patent No. 5,264,365; U.S. Patent No. 5,508,192; Hara *et al.*, Microbial Drug Resistance, 2:63-72 (1996)).

[00494] *E. coli* strains deficient for proteolytic enzymes and transformed with plasmids overexpressing one or more chaperone proteins may be used as host cells in the expression systems disclosed herein.

10 *iii. Antibody Purification*

[00495] Standard protein purification methods known in the art can be employed. The following procedures are exemplary of suitable purification procedures: fractionation on immunoaffinity or ion-exchange columns, ethanol precipitation, reverse phase HPLC, chromatography on silica or on a cation-exchange resin such as DEAE, chromatofocusing,
15 SDS-PAGE, ammonium sulfate precipitation, and gel filtration using, for example, Sephadex G-75.

[00496] In one aspect, Protein A immobilized on a solid phase is used for immunoaffinity purification of the full length antibody products disclosed herein. Protein A is a 41 kD cell wall protein from *Staphylococcus aureus* which binds with a high affinity to the
20 Fc region of antibodies (see, e.g., Lindmark *et al.* (1983) J. Immunol. Meth. 62:1-13). The solid phase to which Protein A is immobilized is preferably a column comprising a glass or silica surface, more preferably a controlled pore glass column or a silicic acid column. In some applications, the column has been coated with a reagent, such as glycerol, in an attempt to prevent nonspecific adherence of contaminants.

25 [00497] As the first step of purification, the preparation derived from the cell culture as described above is applied onto the Protein A immobilized solid phase to allow specific binding of the antibody of interest to Protein A. The solid phase is then washed to remove contaminants non-specifically bound to the solid phase. Finally the antibody of interest is recovered from the solid phase by elution.

30 *b. Generating Antibodies Using Eukaryotic Host Cells:*

[00498] The vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence.

(i) Signal Sequence Component

35 [00499] A vector for use in a eukaryotic host cell may also contain a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the

mature protein or polypeptide of interest. The heterologous signal sequence selected preferably is one that is recognized and processed (e.g., cleaved by a signal peptidase) by the host cell. In mammalian cell expression, mammalian signal sequences as well as viral secretory leaders, for example, the herpes simplex gD signal, are available.

5 [00500] The DNA for such precursor region is ligated in reading frame to DNA encoding the antibody.

(ii) Origin of Replication

[00501] Generally, an origin of replication component is not needed for mammalian expression vectors. For example, the SV40 origin may be used only because it contains
10 the early promoter.

(iii) Selection Gene Component

[00502] Expression and cloning vectors may contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b)
15 complement auxotrophic deficiencies, where relevant, or (c) supply critical nutrients not available from complex media.

[00503] One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous gene produce a protein conferring drug resistance and thus survive the selection regimen. Examples of
20 such dominant selection use the drugs neomycin, mycophenolic acid and hygromycin.

[00504] Another example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the antibody nucleic acid, such as DHFR, thymidine kinase, metallothionein-I and -II, preferably primate metallothionein genes, adenosine deaminase, ornithine decarboxylase, etc.

25 [00505] For example, cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium that contains methotrexate (Mtx), a competitive antagonist of DHFR. An appropriate host cell when wild-type DHFR is employed is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity (e.g., ATCC CRL-9096).

30 [00506] Alternatively, host cells (particularly wild-type hosts that contain endogenous DHFR) transformed or co-transformed with DNA sequences encoding an antibody, wild-type DHFR protein, and another selectable marker such as aminoglycoside 3'-phosphotransferase (APH) can be selected by cell growth in medium containing a selection agent for the selectable marker such as an aminoglycosidic antibiotic, e.g.,
35 kanamycin, neomycin, or G418. See U.S. Patent No. 4,965,199.

(iv) Promoter Component

[00507] Expression and cloning vectors usually contain a promoter that is recognized by the host organism and is operably linked to the antibody polypeptide nucleic acid. Promoter sequences are known for eukaryotes. Virtually alleukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CNCAAT region where N may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into eukaryotic expression vectors.

[00508] Antibody polypeptide transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, from heat-shock promoters, provided such promoters are compatible with the host cell systems.

[00509] The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment that also contains the SV40 viral origin of replication. The immediate early promoter of the human cytomegalovirus is conveniently obtained as a HindIII E restriction fragment. A system for expressing DNA in mammalian hosts using the bovine papilloma virus as a vector is disclosed in U.S. Patent No. 4,419,446. A modification of this system is described in U.S. Patent No. 4,601,978. Alternatively, the Rous Sarcoma Virus long terminal repeat can be used as the promoter.

(v) Enhancer Element Component

[00510] Transcription of DNA encoding the antibody polypeptide of this disclosure by higher eukaryotes is often increased by inserting an enhancer sequence into the vector. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). An enhancer from a eukaryotic cell virus may also be used. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. See also Yaniv, Nature 297:17-18 (1982) on enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a position 5' or 3' to the antibody polypeptide-encoding sequence, but is preferably located at a site 5' from the promoter.

(vi) Transcription Termination Component

[00511] Expression vectors used in eukaryotic host cells will typically also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding an antibody. One useful transcription termination component is the bovine growth hormone polyadenylation region. See WO94/11026 and the expression vector disclosed therein.

(vii) Selection and Transformation of Host Cells

[00512] Suitable host cells for cloning or expressing the DNA in the vectors herein include higher eukaryote cells described herein, including vertebrate host cells. Propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, J. Gen Virol. 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub *et al.*, Proc. Natl. Acad. Sci. USA 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23:243-251 (1980); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather *et al.*, Annals N.Y. Acad. Sci. 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

[00513] Any of the well-known procedures for introducing foreign nucleotide sequences into host cells may be used. These include the use of calcium phosphate transfection, polybrene, protoplast fusion, electroporation, biolistics, liposomes, microinjection, plasma vectors, viral vectors and any of the other well known methods for introducing cloned genomic DNA, cDNA, synthetic DNA, or other foreign genetic material into a host cell (see, *e.g.*, Sambrook and Russell, *supra*). It is only necessary that the particular genetic engineering procedure used be capable of successfully introducing at least both genes into the host cell capable of expressing germline antibody polypeptide.

[00514] Host cells are transformed with the above-described expression or cloning vectors for antibody production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

(viii) Culturing the Host Cells

[00515] The host cells used to produce an antibody of this disclosure may be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma), Minimal Essential Medium ((MEM), (Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ((DMEM), Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham *et al.*, Meth. Enz. 58:44 (1979), Barnes *et al.*, Anal. Biochem. 102:255 (1980), U.S. Patent Nos. 4,767,704; 4,657,866; 4,927,762; 4,560,655; or 5,122,469; WO 90/03430; WO 87/00195; or U.S. Patent Reissue 30,985 may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleotides (such as adenosine and thymidine), antibiotics (such as GENTAMYCIN™ drug), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

(ix) Purification of Antibody

[00516] When using recombinant techniques, the antibody can be produced intracellularly, or directly secreted into the medium. If the antibody is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, are removed, for example, by centrifugation or ultrafiltration. Where the antibody is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon® ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious contaminants.

[00517] The antibody composition prepared from the cells can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography being the preferred purification technique. The suitability of protein A as an affinity ligand depends on the species and isotype of any immunoglobulin Fc domain that is present in the antibody. Protein A can be used to purify antibodies that are based on human $\gamma 1$, $\gamma 2$, or $\gamma 4$ heavy chains (Lindmark *et al.*, J. Immunol. Meth. 62:1-13 (1983)). Protein G is recommended for all mouse isotypes and for human $\gamma 3$ (Guss *et al.*, EMBO J. 5:15671575 (1986)). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically

stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the antibody comprises a CH3 domain, the Bakerbond ABX™ resin (J. T. Baker, Phillipsburg, N.J.) is useful for purification. Other techniques for protein purification such as

5 fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE™ chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered.

10 [00518] Soluble forms of antibody or fragment present either in the cytoplasm or released from the periplasmic space may be further purified using methods known in the art, for example Fab fragments are released from the bacterial periplasmic space by osmotic shock techniques.

[00519] If inclusion bodies comprising an antibody or fragment have formed, they

15 can often bind to the inner and/or outer cellular membranes and thus will be found primarily in the pellet material after centrifugation. The pellet material can then be treated at pH extremes or with chaotropic agent such as a detergent, guanidine, guanidine derivatives, urea, or urea derivatives in the presence of a reducing agent such as dithiothreitol at alkaline pH or tris carboxyethyl phosphine at acid pH to release, break apart, and solubilize

20 the inclusion bodies. The soluble antibody or fragment can then be analyzed using gel electrophoresis, immunoprecipitation or the like. If it is desired to isolate a solubilized antibody or antigen binding fragment isolation may be accomplished using standard methods such as those set forth below and in Marston *et al.* (Meth. Enz., 182:264-275 (1990)).

25 [00520] Following any preliminary purification step(s), the mixture comprising the antibody of interest and contaminants may be subjected to low pH hydrophobic interaction chromatography using an elution buffer at a pH between about 2.5-4.5, preferably performed at low salt concentrations (e.g., from about 0-0.25 M salt).

[00521] In some cases, an antibody or fragment may not be biologically active

30 upon isolation. Various methods for "refolding" or converting a polypeptide to its tertiary structure and generating disulfide linkages, can be used to restore biological activity. Such methods include exposing the solubilized polypeptide to a pH usually above 7 and in the presence of a particular concentration of a chaotrope. The selection of chaotrope is very similar to the choices used for inclusion body solubilization, but usually the chaotrope is

35 used at a lower concentration and is not necessarily the same as chaotropes used for the solubilization. In most cases the refolding/oxidation solution will also contain a reducing

agent or the reducing agent plus its oxidized form in a specific ratio to generate a particular redox potential allowing for disulfide shuffling to occur in the formation of the protein's cysteine bridge(s). Some of the commonly used redox couples include cysteine/cystamine, glutathione (GSH)/dithiobis GSH, cupric chloride, dithiothreitol(DTT)/dithiane DTT, and 2-mercaptoethanol(bME)/di-thio-b(ME). In many instances, a cosolvent may be used to increase the efficiency of the refolding, and common reagents used for this purpose include glycerol, polyethylene glycol of various molecular weights, arginine and the like.

Immunoconjugates

[00522] The disclosure also provides immunoconjugates (interchangeably termed "antibody-drug conjugates" or "ADC"), comprising any of the humanized antibodies comprising an ultralong CDR3 as described herein conjugated to a cytotoxic agent such as a chemotherapeutic agent, a drug, a growth inhibitory agent, a toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (e.g., a radioconjugate).

[00523] The use of antibody-drug conjugates for the local delivery of cytotoxic or cytostatic agents. For example, drugs to kill or inhibit tumor cells in the treatment of cancer (Syrigos and Epenetos (1999) *Anticancer Research* 19:605-614; Niculescu-Duvaz and Springer (1997) *Adv. Drg Del. Rev.* 26:151-172; U.S. Patent No. 4,975,278) allows targeted delivery of the drug moiety to tumors, and intracellular accumulation therein, where systemic administration of these unconjugated drug agents may result in unacceptable levels of toxicity to normal cells as well as the tumor cells sought to be eliminated (Baldwin *et al.*, (1986) *Lancet* pp. (Mar. 15, 1986):603-05; Thorpe, (1985) "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review," in *Monoclonal Antibodies '84: Biological And Clinical Applications*, A. Pinchera *et al.* (ed.s), pp. 475-506). Maximal efficacy with minimal toxicity is sought thereby. Both polyclonal antibodies and monoclonal antibodies have been reported as useful in these strategies (Rowland *et al.*, (1986) *Cancer Immunol. Immunother.*, 21:183-87). Drugs used in these methods include daunomycin, doxorubicin, methotrexate, and vindesine (Rowland *et al.*, (1986) *Supra*). Toxins used in antibody-toxin conjugates include bacterial toxins such as diphtheria toxin, plant toxins such as ricin, small molecule toxins such as geldanamycin (Mandler *et al* (2000) *Jour. of the Nat. Cancer Inst.* 92(19):1573-1581; Mandler *et al* (2000) *Bioorganic & Med. Chem. Letters* 10: 1025-1028; Mandler *et al* (2002) *Bioconjugate Chem.* 13:786-791), maytansinoids (EP 1391213; Liu *et al.*, (1996) *Proc. Natl. Acad. Sci. USA* 93:8618-8623), and calicheamicin (Lode *et al* (1998) *Cancer Res.* 58:2928; Hinman *et al* (1993) *Cancer Res.* 53:3336-3342). The toxins may effect their cytotoxic and cytostatic effects by

mechanisms including tubulin binding, DNA binding, or topoisomerase inhibition. Some cytotoxic drugs tend to be inactive or less active when conjugated to large antibodies or protein receptor ligands.

[00524] ZEVALIN[®] (ibritumomab tiuxetan, Biogen/Idec) is an antibody-radioisotope
 5 conjugate composed of a murine IgG1 kappa monoclonal antibody directed against the
 CD20 antigen found on the surface of normal and malignant B lymphocytes and ¹¹¹In or ⁹⁰Y
 radioisotope bound by a thiourea linker-chelator (Wiseman et al (2000) Eur. Jour. Nucl.
 Med. 27(7):766-77; Wiseman et al (2002) Blood 99(12):4336-42; Witzig et al (2002) J. Clin.
 Oncol. 20(10):2453-63; Witzig et al (2002) J. Clin. Oncol. 20(15):3262-69). Although
 10 ZEVALIN has activity against B-cell non-Hodgkin's Lymphoma (NHL), administration
 results in severe and prolonged cytopenias in most patients. MYLOTARG[™] (gemtuzumab
 ozogamicin, Wyeth Pharmaceuticals), an antibody drug conjugate composed of a hu CD33
 antibody linked to calicheamicin, was approved in 2000 for the treatment of acute myeloid
 leukemia by injection (Drugs of the Future (2000) 25(7):686; U.S. Patent Nos. 4,970,198;
 15 5,079,233; 5,585,089; 5,606,040; 5,693,762; 5,739,116; 5,767,285; 5,773,001).
 Cantuzumab mertansine (Immunogen, Inc.), an antibody drug conjugate composed of the
 huC242 antibody linked via the disulfide linker SPP to the maytansinoid drug moiety, DM1,
 is advancing into Phase II trials for the treatment of cancers that express CanAg, such as
 colon, pancreatic, gastric, and others. MLN-2704 (Millennium Pharm., BZL Biologics,
 20 Immunogen Inc.), an antibody drug conjugate composed of the anti-prostate specific
 membrane antigen (PSMA) monoclonal antibody linked to the maytansinoid drug moiety,
 DM1, is under development for the potential treatment of prostate tumors. The auristatin
 peptides, auristatin E (AE) and monomethylauristatin (MMAE), synthetic analogs of
 dolastatin, were conjugated to chimeric monoclonal antibodies cBR96 (specific to Lewis Y
 25 on carcinomas) and cAC10 (specific to CD30 on hematological malignancies) (Doronina et
 al (2003) Nature Biotechnology 21(7):778-784) and are under therapeutic development.

[00525] Chemotherapeutic agents useful in the generation of immunoconjugates
 are described herein. Enzymatically active toxins and fragments thereof that can be used
 include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A
 30 chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain,
 alpha-sarcin, Aleurites fordii proteins, dianthin proteins, *Phytolacca americana* proteins
 (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria
 officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the
 tricothecenes. See, e.g., WO 93/21232 published Oct. 28, 1993. A variety of radionuclides
 35 are available for the production of radioconjugated antibodies. Examples include ²¹²Bi, ¹³¹I,
¹³¹In, ⁹⁰Y, and ¹⁸⁶Re. Conjugates of the antibody and cytotoxic agent are made using a

variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin may be prepared as described in Vitetta *et al.*, Science, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

[00526] Conjugates of an antibody and one or more small molecule toxins, such as a calicheamicin, maytansinoids, dolastatins, aurostatins, a trichothecene, and CC1065, and the derivatives of these toxins that have toxin activity, are also contemplated herein.

15 *a. Maytansine and Maytansinoids*

[00527] In some embodiments, the immunoconjugate comprises a humanized antibody (full length or fragments) comprising an ultralong CDR3 as disclosed herein conjugated to one or more maytansinoid molecules.

[00528] Maytansinoids are mitototic inhibitors which act by inhibiting tubulin polymerization. Maytansine was first isolated from the east African shrub *Maytenus serrata* (U.S. Patent No. 3,896,111). Subsequently, it was discovered that certain microbes also produce maytansinoids, such as maytansinol and C-3 maytansinol esters (U.S. Patent No. 4,151,042). Synthetic maytansinol and derivatives and analogues thereof are disclosed, for example, in U.S. Patent Nos. 4,137,230; 4,248,870; 4,256,746; 4,260,608; 4,265,814; 4,294,757; 4,307,016; 4,308,268; 4,308,269; 4,309,428; 4,313,946; 4,315,929; 4,317,821; 4,322,348; 4,331,598; 4,361,650; 4,364,866; 4,424,219; 4,450,254; 4,362,663; and 4,371,533.

[00529] Maytansinoid drug moieties are attractive drug moieties in antibody drug conjugates because they are: (i) relatively accessible to prepare by fermentation or chemical modification, derivatization of fermentation products, (ii) amenable to derivatization with functional groups suitable for conjugation through the non-disulfide linkers to antibodies, (iii) stable in plasma, and (iv) effective against a variety of tumor cell lines.

[00530] Immunoconjugates containing maytansinoids, methods of making same, and their therapeutic use are disclosed, for example, in U.S. Patent Nos. 5,208,020, 5,416,064 and EP 0 425 235. Liu *et al.*, Proc. Natl. Acad. Sci. USA 93:8618-8623 (1996)

described immunoconjugates comprising a maytansinoid designated DM1 linked to the monoclonal antibody C242 directed against human colorectal cancer. The conjugate was found to be highly cytotoxic towards cultured colon cancer cells, and showed antitumor activity in an *in vivo* tumor growth assay. Chari *et al.*, Cancer Research 52:127-131 (1992)

5 describe immunoconjugates in which a maytansinoid was conjugated via a disulfide linker to the murine antibody A7 binding to an antigen on human colon cancer cell lines, or to another murine monoclonal antibody TA.1 that binds the HER-2/neu oncogene. The cytotoxicity of the TA.1-maytansinoid conjugate was tested *in vitro* on the human breast cancer cell line SK-BR-3, which expresses 3×10^5 HER-2 surface antigens per cell. The
10 drug conjugate achieved a degree of cytotoxicity similar to the free maytansinoid drug, which could be increased by increasing the number of maytansinoid molecules per antibody molecule. The A7-maytansinoid conjugate showed low systemic cytotoxicity in mice.

[00531] Antibody-maytansinoid conjugates are prepared by chemically linking an
15 antibody to a maytansinoid molecule without significantly diminishing the biological activity of either the antibody or the maytansinoid molecule. See, e.g., U.S. Patent No. 5,208,020. An average of 3-4 maytansinoid molecules conjugated per antibody molecule has shown efficacy in enhancing cytotoxicity of target cells without negatively affecting the function or solubility of the antibody, although even one molecule of toxin/antibody would be expected
20 to enhance cytotoxicity over the use of naked antibody. Maytansinoids are well known in the art and can be synthesized by known techniques or isolated from natural sources. Suitable maytansinoids are disclosed, for example, in U.S. Patent No. 5,208,020 and in the other patents and nonpatent publications referred to hereinabove. Preferred maytansinoids are maytansinol and maytansinol analogues modified in the aromatic ring or at other
25 positions of the maytansinol molecule, such as various maytansinol esters.

[00532] There are many linking groups known in the art for making antibody-maytansinoid conjugates, including, for example, those disclosed in U.S. Patent Nos. 5,208,020, 6,441,163, or EP Patent 0 425 235, Chari *et al.*, Cancer Research 52:127-131 (1992). Antibody-maytansinoid conjugates comprising the linker component SMCC may be
30 prepared. The linking groups include disulfide groups, thioether groups, acid labile groups, photolabile groups, peptidase labile groups, or esterase labile groups, as disclosed in the above-identified patents, disulfide and thioether groups being preferred. Additional linking groups are described and exemplified herein.

[00533] Conjugates of the antibody and maytansinoid may be made using a
35 variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP), succinimidyl-4-(N-maleimidomethyl)cyclohexane-1-carboxylate

(SMCC), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates
 5 (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). Particularly preferred coupling agents include N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP) (Carlsson *et al.*, Biochem. J. 173:723-737 (1978)) and N-succinimidyl-4-(2-pyridylthio)pentanoate (SPP) to provide for a disulfide linkage.

[00534] The linker may be attached to the maytansinoid molecule at various
 10 positions, depending on the type of the link. For example, an ester linkage may be formed by reaction with a hydroxyl group using conventional coupling techniques. The reaction may occur at the C-3 position having a hydroxyl group, the C-14 position modified with hydroxymethyl, the C-15 position modified with a hydroxyl group, and the C-20 position having a hydroxyl group. In a preferred embodiment, the linkage is formed at the C-3
 15 position of maytansinol or a maytansinol analogue.

b. Auristatins and Dolastatins

[00535] In some embodiments, the immunoconjugate comprises an antibody disclosed herein conjugated to dolastatins or dolostatin peptidic analogs and derivatives, the auristatins (U.S. Patent Nos. 5,635,483; 5,780,588). Dolastatins and auristatins have
 20 been shown to interfere with microtubule dynamics, GTP hydrolysis, and nuclear and cellular division (Woyke *et al* (2001) Antimicrob. Agents and Chemother. 45(12):3580-3584) and have anticancer (U.S. Patent No. 5,663,149) and antifungal activity (Pettit *et al* (1998) Antimicrob. Agents Chemother. 42:2961-2965). The dolastatin or auristatin drug moiety may be attached to the antibody through the N (amino) terminus or the C (carboxyl)
 25 terminus of the peptidic drug moiety (WO 02/088172).

[00536] Exemplary auristatin embodiments include the N-terminus linked monomethylauristatin drug moieties DE and DF, (see, *e.g.*, U.S. Patent No. 7,498,298).

[00537] Typically, peptide-based drug moieties can be prepared by forming a peptide bond between two or more amino acids and/or peptide fragments. Such peptide
 30 bonds can be prepared, for example, according to the liquid phase synthesis method (see, *e.g.*, E. Schroder and K. Lubke, "The Peptides", volume 1, pp 76-136, 1965, Academic Press) that is well known in the field of peptide chemistry. The auristatin/dolastatin drug moieties may be prepared according to the methods of: U.S. Patent No. 5,635,483; U.S. Patent No. 5,780,588; Pettit *et al* (1989) J. Am. Chem. Soc. 111:5463-5465; Pettit *et al.*
 35 (1998) Anti-Cancer Drug Design 13:243-277; Pettit, G. R., *et al.* Synthesis, 1996, 719-725; and Pettit *et al.* (1996) J. Chem. Soc. Perkin Trans. 1 5:859-863. See also Doronina (2003)

Nat Biotechnol 21(7):778-784; U.S. Patent No. 7,498,289, (disclosing, linkers and methods of preparing monomethylvaline compounds such as MMAE and MMAF conjugated to linkers).

c. Calicheamicin

5 [00538] In other embodiments, the immunoconjugate comprises an antibody disclosed herein conjugated to one or more calicheamicin molecules. The calicheamicin family of antibiotics are capable of producing double-stranded DNA breaks at sub-picomolar concentrations. For the preparation of conjugates of the calicheamicin family, see U.S. Patent Nos. 5,712,374, 5,714,586, 5,739,116, 5,767,285, 5,770,701, 5,770,710,
10 5,773,001, 5,877,296. Structural analogues of calicheamicin which may be used include, but are not limited to, γ_1^I , α_2^I , α_3^I , N-acetyl- γ_1^I , PSAG and θ_1^I (see, e.g., Hinman *et al.*, Cancer Research 53:3336-3342 (1993), Lode *et al.*, Cancer Research 58:2925-2928 (1998) and the aforementioned U.S. patents). Another anti-tumor drug that the antibody can be conjugated is QFA which is an antifolate. Both calicheamicin and QFA have
15 intracellular sites of action and do not readily cross the plasma membrane. Therefore, cellular uptake of these agents through antibody mediated internalization greatly enhances their cytotoxic effects.

d. Other Cytotoxic Agents

[00539] Other antitumor agents that can be conjugated to the antibodies disclosed
20 herein include BCNU, streptozocin, vincristine and 5-fluorouracil, the family of agents known collectively LL-E33288 complex described in U.S. Patent Nos. 5,053,394, 5,770,710, as well as esperamicins (U.S. Patent No. 5,877,296).

[00540] Enzymatically active toxins and fragments thereof which can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A
25 chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolaca americana proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. See, for example, WO 93/21232 published Oct. 28, 1993.

30 [00541] The present disclosure further contemplates an immunoconjugate formed between an antibody and a compound with nucleolytic activity (e.g., a ribonuclease or a DNA endonuclease such as a deoxyribonuclease; DNase).

[00542] For selective destruction of the tumor, the antibody may comprise a highly radioactive atom. A variety of radioactive isotopes are available for the production of
35 radioconjugated antibodies. Examples include At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³², Pb²¹² and radioactive isotopes of Lu. When the conjugate is used for detection, it may

comprise a radioactive atom for scintigraphic studies, for example $\text{tc}^{99\text{m}}$ or I^{123} , or a spin label for nuclear magnetic resonance (NMR) imaging (also known as magnetic resonance imaging, mri), such as iodine-123 again, iodine-131, indium-111, fluorine-19, carbon-13, nitrogen-15, oxygen-17, gadolinium, manganese or iron.

5 [00543] The radiolabels or other labels may be incorporated in the conjugate in known ways. For example, the peptide may be biosynthesized or may be synthesized by chemical amino acid synthesis using suitable amino acid precursors involving, for example, fluorine-19 in place of hydrogen. Labels such as $\text{tc}^{99\text{m}}$ or I^{123} , Re^{186} , Re^{188} and In^{111} can be attached via a cysteine residue in the peptide. Yttrium-90 can be attached via a lysine
10 residue. The IODOGEN method (Fraker et al (1978) Biochem. Biophys. Res. Commun. 80: 49-57) can be used to incorporate iodine-123. "Monoclonal Antibodies in Immunoscintigraphy" (Chatal, CRC Press 1989) describes other methods.

[00544] Conjugates of the antibody and cytotoxic agent may be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithio)
15 propionate (SPDP), succinimidyl-4-(N-maleimidomethyl)cyclohexane-1-carboxylate (SMCC), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates
20 (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., Science 238:1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See
25 WO94/11026. The linker may be a "cleavable linker" facilitating release of the cytotoxic drug in the cell. For example, an acid-labile linker, peptidase-sensitive linker, photolabile linker, dimethyl linker or disulfide-containing linker (Chari et al., Cancer Research 52:127-131 (1992); U.S. Patent No. 5,208,020) may be used.

[00545] The compounds disclosed herein expressly contemplate, but are not
30 limited to, ADC prepared with cross-linker reagents: BMPS, EMCS, GMBS, HBVS, LC-SMCC, MBS, MPBH, SBAP, SIA, SIAB, SMCC, SMPB, SMPH, sulfo-EMCS, sulfo-GMBS, sulfo-KMUS, sulfo-MBS, sulfo-SIAB, sulfo-SMCC, and sulfo-SMPB, and SVSB (succinimidyl-(4-vinylsulfone)benzoate) which are commercially available (e.g., from Pierce Biotechnology, Inc., Rockford, Ill., U.S.A). See pages 467-498, 2003-2004 Applications
35 Handbook and Catalog.

e. Preparation of Antibody Drug Conjugates

[00546] In the antibody drug conjugates (ADC) disclosed herein, an antibody (Ab) is conjugated to one or more drug moieties (D), *e.g.*, about 1 to about 20 drug moieties per antibody, through a linker (L). An ADC of Formula I $[Ab-(L-D)_p]$ may be prepared by several routes, employing organic chemistry reactions, conditions, and reagents known to those skilled in the art, including: (1) reaction of a nucleophilic group of an antibody with a bivalent linker reagent, to form Ab-L, via a covalent bond, followed by reaction with a drug moiety D; and (2) reaction of a nucleophilic group of a drug moiety with a bivalent linker reagent, to form D-L, via a covalent bond, followed by reaction with the nucleophilic group of an antibody. Additional methods for preparing ADC are described herein.

[00547] The linker may be composed of one or more linker components. Exemplary linker components include 6-maleimidocaproyl ("MC"), maleimidopropanoyl ("MP"), valine-citrulline ("val-cit"), alanine-phenylalanine ("ala-phe"), *p*-aminobenzyloxycarbonyl ("PAB"), N-Succinimidyl 4-(2-pyridylthio) pentanoate ("SPP"), N-Succinimidyl 4-(N-maleimidomethyl)cyclohexane-1 carboxylate ("SMCC"), and N-Succinimidyl (4-iodo-acetyl)aminobenzoate ("SIAB"). Additional linker components are known in the art and some are disclosed herein (see, *e.g.*, U.S. Patent No. 7,498,298).

[00548] In some embodiments, the linker may comprise amino acid residues. Exemplary amino acid linker components include a dipeptide, a tripeptide, a tetrapeptide or a pentapeptide. Exemplary dipeptides include: valine-citrulline (vc or val-cit), alanine-phenylalanine (af or ala-phe). Exemplary tripeptides include: glycine-valine-citrulline (gly-val-cit) and glycine-glycine-glycine (gly-gly-gly). Amino acid residues which comprise an amino acid linker component include those occurring naturally, as well as minor amino acids and non-naturally occurring amino acid analogs, such as citrulline. Amino acid linker components can be designed and optimized in their selectivity for enzymatic cleavage by a particular enzymes, for example, a tumor-associated protease, cathepsin B, C and D, or a plasmin protease.

[00549] Nucleophilic groups on antibodies include, but are not limited to: (i) N-terminal amine groups, (ii) side chain amine groups, *e.g.*, lysine, (iii) side chain thiol groups, *e.g.*, cysteine, and (iv) sugar hydroxyl or amino groups where the antibody is glycosylated. Amine, thiol, and hydroxyl groups are nucleophilic and capable of reacting to form covalent bonds with electrophilic groups on linker moieties and linker reagents including: (i) active esters such as NHS esters, HOBt esters, haloformates, and acid halides; (ii) alkyl and benzyl halides such as haloacetamides; (iii) aldehydes, ketones, carboxyl, and maleimide groups. Certain antibodies have reducible interchain disulfides, *e.g.*, cysteine bridges. Antibodies may be made reactive for conjugation with linker reagents by treatment with a reducing agent such as DTT (dithiothreitol). Each cysteine

bridge will thus form, theoretically, two reactive thiol nucleophiles. Additional nucleophilic groups can be introduced into antibodies through the reaction of lysines with 2-iminothiolane (Traut's reagent) resulting in conversion of an amine into a thiol. Reactive thiol groups may be introduced into the antibody (or fragment thereof) by introducing one, two, three, four, or more cysteine residues (e.g., preparing mutant antibodies comprising one or more non-native cysteine amino acid residues).

[00550] Antibody drug conjugates disclosed herein may also be produced by modification of the antibody to introduce electrophilic moieties, which can react with nucleophilic substituents on the linker reagent or drug. The sugars of glycosylated antibodies may be oxidized, e.g., with periodate oxidizing reagents, to form aldehyde or ketone groups which may react with the amine group of linker reagents or drug moieties. The resulting imine Schiff base groups may form a stable linkage, or may be reduced, e.g., by borohydride reagents to form stable amine linkages. In one embodiment, reaction of the carbohydrate portion of a glycosylated antibody with either glactose oxidase or sodium meta-periodate may yield carbonyl (aldehyde and ketone) groups in the protein that can react with appropriate groups on the drug (Hermanson, Bioconjugate Techniques). In another embodiment, proteins containing N-terminal serine or threonine residues can react with sodium meta-periodate, resulting in production of an aldehyde in place of the first amino acid (Geoghegan & Stroh, (1992) Bioconjugate Chem. 3:138-146; U.S. Patent No. 5,362,852). Such aldehyde can be reacted with a drug moiety or linker nucleophile.

[00551] Likewise, nucleophilic groups on a drug moiety include, but are not limited to: amine, thiol, hydroxyl, hydrazide, oxime, hydrazine, thiosemicarbazone, hydrazine carboxylate, and arylhydrazide groups capable of reacting to form covalent bonds with electrophilic groups on linker moieties and linker reagents including: (i) active esters such as NHS esters, HOBt esters, haloformates, and acid halides; (ii) alkyl and benzyl halides such as haloacetamides; (iii) aldehydes, ketones, carboxyl, and maleimide groups.

[00552] Alternatively, a fusion protein comprising the antibody and cytotoxic agent may be made, e.g., by recombinant techniques or peptide synthesis. The length of DNA may comprise respective regions encoding the two portions of the conjugate either adjacent one another or separated by a region encoding a linker peptide which does not destroy the desired properties of the conjugate.

[00553] In yet another embodiment, the antibody may be conjugated to a "receptor" (such streptavidin) for utilization in tumor pre-targeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) which is conjugated to a cytotoxic agent (e.g., a radionucleotide).

Engineered Hybridomas

[00554] Hybridoma cells can be generated by fusing B cells producing a desired antibody with an immortalized cell line, usually a myeloma cell line, so that the resulting fusion cells will be an immortalized cell line that secretes a particular antibody. By the same principle, myeloma cells can be first transfected with a nucleic acid encoding a germline antibody V region and can be screened for the expression of the germline V region. Those myeloma cells with highest level of proteolytic light chain expression can be subsequently fused with B cells that produce an antibody with desired target protein specificity. The fusion cells will produce two types of antibodies: one is a heterologous antibody containing an endogenous antibody chain (either heavy or light) operably joined to the recombinant germline V region (either heavy or light), and the other is the same antibody that the parental B cells would secrete (e.g. both endogenous heavy and light chains). The operably joined heterologous heavy and light chains can be isolated by conventional methods such as chromatography and identification can be confirmed by target protein binding assays, assays identifying a unique tag of the germline polypeptide, or endopeptidase activity assays described in other sections of this disclosure. In some cases, where the heterologous antibody is the predominant type in quantity among the two types of antibodies, such isolation may not be needed. Hybridomas. Including bovine hybridomas, may be a source of bovine antibody gene sequences, including ultralong CDR3 sequences.

Transgenic Mammals

[00555] A nucleic acid sequence encoding a germline antibody polypeptide of the present disclosure can be introduced into a non-human mammal to generate a transgenic animal that expresses the germline antibody polypeptide. Unlike the transgenic animal models more commonly seen, the transgene expressed by the transgenic mammals of the present disclosure need not replace at least one allele of the endogenous coding sequence responsible for the variable regions of antibody chains following somatic recombination. Due to allelic exclusion, the presence of an exogenous, post-somatic rearrangement version of the germline V region DNA will inhibit the endogenous alleles of pre-somatic rearrangement V minigenes from undergoing somatic rearrangement and contributing to the makeup of antibody chains this mammal may produce. Thus, when exposed to a particular antigen, the mammal will generate heterologous antibodies comprising one endogenously rearranged antibody chain, and one transgenic gene which was rearranged a priori. Such heterologous antibodies are invaluable in research and in treating certain

conditions in live subjects. On the other hand, a method that directs the integration of the transgene to the locus of an endogenous allele will fully serve the purpose of practicing the present disclosure as well.

[00556] The general methods of generating transgenic animals have been well
 5 established and frequently practiced. For reviews and protocols for generating transgenic animals and related methods for genetic manipulations, see, *e.g.*, Mansour *et al.*, Nature 336:348-352 (1988); Capecchi *et al.*, Trends Genet. 5:70-76 (1989); Capecchi, Science 244:1288-1292 (1989); Capecchi *et al.*, Current Communications in Molecular Biology, pp45-52, Capecchi, M.R. (ed.), Cold Spring Harbor Press, Cold Spring Harbor, N.Y.
 10 (1989); Frohman *et al.*, Cell 56: 145-147 (1989); Brinster *et al.*, Proc. Natl. Acad. Sci. USA 82:4438-4442 (1985); Evans *et al.*, Nature 292:154-156 (1981); Bradley *et al.*, Nature 309:255-258 (1984); Gossler *et al.*, Proc. Natl. Acad. Sci. USA 83:9065-9069 (1986); Robertson *et al.*, Nature 322:445-448 (1986); Jaenisch Science 240:1468-1474 (1988); and Siedel, G. E., Jr., "Critical review of embryo transfer procedures with cattle" in
 15 Fertilization and Embryonic Development in Vitro, page 323, L. Mastroianni, Jr. and J. D. Biggers, ed., Plenum Press, New York, N.Y. (1981).

[00557] An exemplary transgenic animal of the present disclosure is mouse, whereas a number of other transgenic animals can also be produced using the same general method. These animals include, but are not limited to: rabbits, sheep, cattle, and
 20 pigs (Jaenisch Science 240:1468-1474 (1988); Hammer *et al.*, J. Animal. Sci. 63:269 (1986); Hammer *et al.* Nature 315:680 (1985); Wagner *et al.*, Theriogenology 21:29 (1984)).

Pharmaceutical Compositions

[00558] Humanized antibodies comprising an ultralong CDR3, antibody fragments, nucleic acids, or vectors disclosed herein can be formulated in compositions, especially pharmaceutical compositions. Such compositions with humanized antibodies comprising an ultralong CDR3 comprise a therapeutically or prophylactically effective amount of a humanized antibodies comprising an ultralong CDR3, antibody fragment, nucleic acid, or
 30 vector disclosed herein in admixture with a suitable carrier, *e.g.*, a pharmaceutically acceptable agent. Typically, humanized antibodies comprising an ultralong CDR3, antibody fragments, nucleic acids, or vectors disclosed herein are sufficiently purified for administration before formulation in a pharmaceutical composition.

[00559] Pharmaceutically acceptable agents for use in the present pharmaceutical
 35 compositions include carriers, excipients, diluents, antioxidants, preservatives, coloring, flavoring and diluting agents, emulsifying agents, suspending agents, solvents, fillers,

bulking agents, buffers, delivery vehicles, tonicity agents, cosolvents, wetting agents, complexing agents, buffering agents, antimicrobials, and surfactants.

[00560] Neutral buffered saline or saline mixed with serum albumin are exemplary appropriate carriers. The pharmaceutical compositions may include antioxidants such as ascorbic acid; low molecular weight polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween, pluronics, or polyethylene glycol (PEG). Also by way of example, suitable tonicity enhancing agents include alkali metal halides (preferably sodium or potassium chloride), mannitol, sorbitol, and the like. Suitable preservatives include benzalkonium chloride, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid and the like. Hydrogen peroxide also may be used as preservative. Suitable cosolvents include glycerin, propylene glycol, and PEG. Suitable complexing agents include caffeine, polyvinylpyrrolidone, beta-cyclodextrin or hydroxy-propyl-beta-cyclodextrin. Suitable surfactants or wetting agents include sorbitan esters, polysorbates such as polysorbate 80, tromethamine, lecithin, cholesterol, tyloxapal, and the like. The buffers may be conventional buffers such as acetate, borate, citrate, phosphate, bicarbonate, or Tris-HCl. Acetate buffer may be about pH 4-5.5, and Tris buffer can be about pH 7-8.5. Additional pharmaceutical agents are set forth in Remington's Pharmaceutical Sciences, 18th Edition, A. R. Gennaro, ed., Mack Publishing Company, 1990.

[00561] The composition may be in liquid form or in a lyophilized or freeze-dried form and may include one or more lyoprotectants, excipients, surfactants, high molecular weight structural additives and/or bulking agents (see, for example, U.S. Patent Nos. 6,685,940, 6,566,329, and 6,372,716). In one embodiment, a lyoprotectant is included, which is a non-reducing sugar such as sucrose, lactose or trehalose. The amount of lyoprotectant generally included is such that, upon reconstitution, the resulting formulation will be isotonic, although hypertonic or slightly hypotonic formulations also may be suitable. In addition, the amount of lyoprotectant should be sufficient to prevent an unacceptable amount of degradation and/or aggregation of the protein upon lyophilization. Exemplary lyoprotectant concentrations for sugars (e.g., sucrose, lactose, trehalose) in the pre-lyophilized formulation are from about 10 mM to about 400 mM. In another embodiment, a surfactant is included, such as for example, nonionic surfactants and ionic surfactants such as polysorbates (e.g., polysorbate 20, polysorbate 80); poloxamers (e.g., poloxamer 188);

poly(ethylene glycol) phenyl ethers (e.g., Triton); sodium dodecyl sulfate (SDS); sodium laurel sulfate; sodium octyl glycoside; lauryl-, myristyl-, linoleyl-, or stearyl-sulfobetaine; lauryl-, myristyl-, linoleyl- or stearyl-sarcosine; linoleyl, myristyl-, or cetyl-betaine; lauroamidopropyl-, cocamidopropyl-, linoleamidopropyl-, myristamidopropyl-,
 5 palmidopropyl-, or isostearamidopropyl-betaine (e.g., lauroamidopropyl); myristamidopropyl-, palmidopropyl-, or isostearamidopropyl-dimethylamine; sodium methyl cocoyl-, or disodium methyl ofeyle-aurate; and the MONAQUAT™. series (Mona Industries, Inc., Paterson, N.J.), polyethyl glycol, polypropyl glycol, and copolymers of ethylene and propylene glycol (e.g., Pluronic, PF68 etc). Exemplary amounts of surfactant that may be
 10 present in the pre-lyophilized formulation are from about 0.001-0.5%. High molecular weight structural additives (e.g., fillers, binders) may include for example, acacia, albumin, alginic acid, calcium phosphate (dibasic), cellulose, carboxymethylcellulose, carboxymethylcellulose sodium, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, microcrystalline cellulose, dextran, dextrin, dextrates,
 15 sucrose, tylose, pregelatinized starch, calcium sulfate, amylose, glycine, bentonite, maltose, sorbitol, ethylcellulose, disodium hydrogen phosphate, disodium phosphate, disodium pyrosulfite, polyvinyl alcohol, gelatin, glucose, guar gum, liquid glucose, compressible sugar, magnesium aluminum silicate, maltodextrin, polyethylene oxide, polymethacrylates, povidone, sodium alginate, tragacanth microcrystalline cellulose,
 20 starch, and zein. Exemplary concentrations of high molecular weight structural additives are from 0.1% to 10% by weight. In other embodiments, a bulking agent (e.g., mannitol, glycine) may be included.

[00562] Compositions may be suitable for parenteral administration. Exemplary compositions are suitable for injection or infusion into an animal by any route available to
 25 the skilled worker, such as intraarticular, subcutaneous, intravenous, intramuscular, intraperitoneal, intracerebral (intraparenchymal), intracerebroventricular, intramuscular, intraocular, intraarterial, or intralesional routes. A parenteral formulation typically will be a sterile, pyrogen-free, isotonic aqueous solution, optionally containing pharmaceutically acceptable preservatives.

[00563] Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringers' dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils.
 35 Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers, such as those based on Ringer's dextrose, and the like. Preservatives and other additives may

also be present, such as, for example, anti-microbials, anti-oxidants, chelating agents, inert gases and the like. See generally, Remington's Pharmaceutical Science, 16th Ed., Mack Eds., 1980.

[00564] Pharmaceutical compositions described herein may be formulated for controlled or sustained delivery in a manner that provides local concentration of the product (e.g., bolus, depot effect) and/or increased stability or half-life in a particular local environment. The compositions can include the formulation of humanized antibodies comprising an ultralong CDR3, antibody fragments, nucleic acids, or vectors disclosed herein with particulate preparations of polymeric compounds such as polylactic acid, polyglycolic acid, *etc.*, as well as agents such as a biodegradable matrix, injectable microspheres, microcapsular particles, microcapsules, bioerodible particles beads, liposomes, and implantable delivery devices that provide for the controlled or sustained release of the active agent which then can be delivered as a depot injection. Techniques for formulating such sustained- or controlled-delivery means are known and a variety of polymers have been developed and used for the controlled release and delivery of drugs. Such polymers are typically biodegradable and biocompatible. Polymer hydrogels, including those formed by complexation of enantiomeric polymer or polypeptide segments, and hydrogels with temperature or pH sensitive properties, may be desirable for providing drug depot effect because of the mild and aqueous conditions involved in trapping bioactive protein agents (e.g., humanized antibodies comprising an ultralong CDR3). See, for example, the description of controlled release porous polymeric microparticles for the delivery of pharmaceutical compositions in WO 93/15722.

[00565] Suitable materials for this purpose include polylactides (see, e.g., U.S. Patent No. 3,773,919), polymers of poly-(α -hydroxycarboxylic acids), such as poly-D-(-)-3-hydroxybutyric acid (EP 133,988A), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman *et al.*, Biopolymers, 22: 547-556 (1983)), poly(2-hydroxyethyl-methacrylate) (Langer *et al.*, J. Biomed. Mater. Res., 15: 167-277 (1981), and Langer, Chem. Tech., 12: 98-105 (1982)), ethylene vinyl acetate, or poly-D-(-)-3-hydroxybutyric acid. Other biodegradable polymers include poly(lactones), poly(acetals), poly(orthoesters), and poly(orthocarbonates). Sustained-release compositions also may include liposomes, which can be prepared by any of several methods known in the art (see, e.g., Eppstein *et al.*, Proc. Natl. Acad. Sci. USA, 82: 3688-92 (1985)). The carrier itself, or its degradation products, should be nontoxic in the target tissue and should not further aggravate the condition. This can be determined by routine screening in animal models of the target disorder or, if such models are unavailable, in normal animals.

[00566] Microencapsulation of recombinant proteins for sustained release has been performed successfully with human growth hormone (rhGH), interferon-(rhIFN-), interleukin-2, and MN rgp120. Johnson *et al.*, Nat. Med., 2:795-799 (1996); Yasuda, Biomed. Ther., 27:1221-1223 (1993); Hora *et al.*, Bio/Technology. 8:755-758 (1990);

5 Cleland, "Design and Production of Single Immunization Vaccines Using Polylactide Polyglycolide Microsphere Systems," in Vaccine Design: The Subunit and Adjuvant Approach, Powell and Newman, eds, (Plenum Press: New York, 1995), pp. 439-462; WO 97/03692, WO 96/40072, WO 96/07399; and U.S. Patent No. 5,654,010. The sustained-release formulations of these proteins were developed using poly-lactic-coglycolic acid

10 (PLGA) polymer due to its biocompatibility and wide range of biodegradable properties. The degradation products of PLGA, lactic and glycolic acids can be cleared quickly within the human body. Moreover, the degradability of this polymer can be depending on its molecular weight and composition. Lewis, "Controlled release of bioactive agents from lactide/glycolide polymer," in: M. Chasin and R. Langer (Eds.), Biodegradable Polymers as

15 Drug Delivery Systems (Marcel Dekker: New York, 1990), pp. 1-41. Additional examples of sustained release compositions include, for example, EP 58,481A, U.S. Patent No. 3,887,699, EP 158,277A, Canadian Patent No. 1176565, U. Sidman *et al.*, Biopolymers 22, 547 [1983], R. Langer *et al.*, Chem. Tech. 12, 98 [1982], Sinha *et al.*, J. Control. Release 90, 261 [2003], Zhu *et al.*, Nat. Biotechnol. 18, 24 [2000], and Dai *et al.*, Colloids Surf B

20 Biointerfaces 41, 117 [2005].

[00567] Bioadhesive polymers are also contemplated for use in or with compositions of the present disclosure. Bioadhesives are synthetic and naturally occurring materials able to adhere to biological substrates for extended time periods. For example, Carbopol and polycarbophil are both synthetic cross-linked derivatives of poly(acrylic acid).

25 Bioadhesive delivery systems based on naturally occurring substances include for example hyaluronic acid, also known as hyaluronan. Hyaluronic acid is a naturally occurring mucopolysaccharide consisting of residues of D-glucuronic and N-acetyl-D-glucosamine. Hyaluronic acid is found in the extracellular tissue matrix of vertebrates, including in connective tissues, as well as in synovial fluid and in the vitreous and aqueous humor of

30 the eye. Esterified derivatives of hyaluronic acid have been used to produce microspheres for use in delivery that are biocompatible and biodegradable (see, for example, Cortivo *et al.*, Biomaterials (1991) 12:727-730; EP 517,565; WO 96/29998; Illum *et al.*, J. Controlled Rel. (1994) 29:133-141). Exemplary hyaluronic acid containing compositions of the present disclosure comprise a hyaluronic acid ester polymer in an amount of approximately 0.1% to

35 about 40% (w/w) of a humanized antibody comprising an ultralong CDR3 to hyaluronic acid polymer.

[00568] Both biodegradable and non-biodegradable polymeric matrices may be used to deliver compositions of the present disclosure, and such polymeric matrices may comprise natural or synthetic polymers. Biodegradable matrices are preferred. The period of time over which release occurs is based on selection of the polymer. Typically, release over a period ranging from between a few hours and three to twelve months is most desirable. Exemplary synthetic polymers which may be used to form the biodegradable delivery system include: polymers of lactic acid and glycolic acid, polyamides, polycarbonates, polyalkylenes, polyalkylene glycols, polyalkylene oxides, polyalkylene terephthalates, polyvinyl alcohols, polyvinyl ethers, polyvinyl esters, poly-vinyl halides, polyvinylpyrrolidone, polyglycolides, polysiloxanes, polyanhydrides, polyurethanes and copolymers thereof, poly(butic acid), poly(valeric acid), alkyl cellulose, hydroxyalkyl celluloses, cellulose ethers, cellulose esters, nitro celluloses, polymers of acrylic and methacrylic esters, methyl cellulose, ethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, hydroxybutyl methyl cellulose, cellulose acetate, cellulose propionate, cellulose acetate butyrate, cellulose acetate phthalate, carboxylethyl cellulose, cellulose triacetate, cellulose sulphate sodium salt, poly(methyl methacrylate), poly(ethyl methacrylate), poly(butylmethacrylate), poly(isobutyl methacrylate), poly(hexylmethacrylate), poly(isodecyl methacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate), polyethylene, polypropylene, poly(ethylene glycol), poly(ethylene oxide), poly(ethylene terephthalate), poly(vinyl alcohols), polyvinyl acetate, poly vinyl chloride, polystyrene and polyvinylpyrrolidone. Exemplary natural polymers include alginate and other polysaccharides including dextran and cellulose, collagen, chemical derivatives thereof (substitutions, additions of chemical groups, for example, alkyl, alkylene, hydroxylations, oxidations, and other modifications routinely made by those skilled in the art), albumin and other hydrophilic proteins, zein and other prolamines and hydrophobic proteins, copolymers and mixtures thereof. In general, these materials degrade either by enzymatic hydrolysis or exposure to water *in vivo*, by surface or bulk erosion. The polymer optionally is in the form of a hydrogel (see, for example, WO 04/009664, WO 05/087201, Sawhney, *et al.*, *Macromolecules*, 1993, 26, 581-587) that can absorb up to about 90% of its weight in water and further, optionally is cross-linked with multi-valent ions or other polymers.

[00569] Delivery systems also include non-polymer systems that are lipids including sterols such as cholesterol, cholesterol esters and fatty acids or neutral fats such as mono-di- and tri-glycerides; hydrogel release systems; silastic systems; peptide based systems; wax coatings; compressed tablets using conventional binders and excipients;

partially fused implants; and the like. Specific examples include, but are not limited to: (a) erosional systems in which the product is contained in a form within a matrix such as those described in U.S. Patent Nos. 4,452,775, 4,675,189 and 5,736,152 and (b) diffusional systems in which a product permeates at a controlled rate from a polymer such as described in U.S. Patent Nos. 3,854,480, 5,133,974 and 5,407,686. Liposomes containing the product may be prepared by methods known methods, such as for example (DE 3,218,121; Epstein *et al.*, Proc. Natl. Acad. Sci. USA, 82: 3688-3692 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA, 77: 4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; JP 83-118008; U.S. Patent Nos. 4,485,045 and 4,544,545; and EP 102,324).

[00570] Alternatively or additionally, the compositions may be administered locally via implantation into the affected area of a membrane, sponge, or other appropriate material on to which a humanized antibody comprising an ultralong CDR3, antibody fragment, nucleic acid, or vector disclosed herein has been absorbed or encapsulated. Where an implantation device is used, the device may be implanted into any suitable tissue or organ, and delivery of a humanized antibody comprising an ultralong CDR3 antibody fragment, nucleic acid, or vector disclosed herein can be directly through the device via bolus, or via continuous administration, or via catheter using continuous infusion.

[00571] A pharmaceutical composition comprising a humanized antibody comprising an ultralong CDR3, antibody fragment, nucleic acid, or vector disclosed herein may be formulated for inhalation, such as for example, as a dry powder. Inhalation solutions also may be formulated in a liquefied propellant for aerosol delivery. In yet another formulation, solutions may be nebulized. Additional pharmaceutical composition for pulmonary administration include, those described, for example, in WO 94/20069, which discloses pulmonary delivery of chemically modified proteins. For pulmonary delivery, the particle size should be suitable for delivery to the distal lung. For example, the particle size may be from 1 μm to 5 μm ; however, larger particles may be used, for example, if each particle is fairly porous.

[00572] Certain formulations containing humanized antibodies comprising an ultralong CDR3, antibody fragments, nucleic acids, or vectors disclosed herein may be administered orally. Formulations administered in this fashion may be formulated with or without those carriers customarily used in the compounding of solid dosage forms such as tablets and capsules. For example, a capsule can be designed to release the active portion of the formulation at the point in the gastrointestinal tract when bioavailability is maximized and pre-systemic degradation is minimized. Additional agents may be included to facilitate absorption of a selective binding agent. Diluents, flavorings, low melting point waxes,

vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders also can be employed.

[00573] Another preparation may involve an effective quantity of a humanized antibody comprising an ultralong CDR3, antibody fragment, nucleic acid, or vector disclosed herein in a mixture with non-toxic excipients which are suitable for the manufacture of tablets. By dissolving the tablets in sterile water, or another appropriate vehicle, solutions may be prepared in unit dose form. Suitable excipients include, but are not limited to, inert diluents, such as calcium carbonate, sodium carbonate or bicarbonate, lactose, or calcium phosphate; or binding agents, such as starch, gelatin, or acacia; or lubricating agents such as magnesium stearate, stearic acid, or talc.

[00574] Suitable and/or preferred pharmaceutical formulations may be determined in view of the present disclosure and general knowledge of formulation technology, depending upon the intended route of administration, delivery format, and desired dosage. Regardless of the manner of administration, an effective dose may be calculated according to patient body weight, body surface area, or organ size. Further refinement of the calculations for determining the appropriate dosage for treatment involving each of the formulations described herein are routinely made in the art and is within the ambit of tasks routinely performed in the art. Appropriate dosages may be ascertained through use of appropriate dose-response data.

[00575] In some embodiments, humanized antibodies comprising an ultralong CDR3 or fragments thereof are provided with a modified Fc region where a naturally-occurring Fc region is modified to increase the half-life of the antibody or fragment in a biological environment, for example, the serum half-life or a half-life measured by an *in vitro* assay. Methods for altering the original form of a Fc region of an IgG also are described in U.S. Patent No. 6,998,253.

[00576] In certain embodiments, it may be desirable to modify the antibody or fragment in order to increase its serum half-life, for example, adding molecules such as PEG or other water soluble polymers, including polysaccharide polymers, to antibody fragments to increase the half-life. This may also be achieved, for example, by incorporation of a salvage receptor binding epitope into the antibody fragment (*e.g.*, by mutation of the appropriate region in the antibody fragment or by incorporating the epitope into a peptide tag that is then fused to the antibody fragment at either end or in the middle, *e.g.*, by DNA or peptide synthesis) (see, International Publication No. WO96/32478). Salvage receptor binding epitope refers to an epitope of the Fc region of an IgG molecule (*e.g.*, IgG1, IgG2, IgG3, or IgG4) that is responsible for increasing the *in vivo* serum half-life of the IgG molecule.

[00577] A salvage receptor binding epitope may include a region wherein any one or more amino acid residues from one or two loops of a Fc domain are transferred to an analogous position of the antibody fragment. Even more preferably, three or more residues from one or two loops of the Fc domain are transferred. Still more preferred, the epitope is taken from the CH2 domain of the Fc region (e.g., of an IgG) and transferred to the CH1, CH3, or VH region, or more than one such region, of the antibody. Alternatively, the epitope is taken from the CH2 domain of the Fc region and transferred to the C_L region or V_L region, or both, of the antibody fragment. See also WO 97/34631 and WO 96/32478 which describe Fc variants and their interaction with the salvage receptor.

[00578] Mutation of residues within Fc receptor binding sites may result in altered effector function, such as altered ADCC or CDC activity, or altered half-life. Potential mutations include insertion, deletion or substitution of one or more residues, including substitution with alanine, a conservative substitution, a non-conservative substitution, or replacement with a corresponding amino acid residue at the same position from a different IgG subclass (e.g., replacing an IgG1 residue with a corresponding IgG2 residue at that position). For example, it has been reported that mutating the serine at amino acid position 241 in IgG4 to proline (found at that position in IgG1 and IgG2) led to the production of a homogeneous antibody, as well as extending serum half-life and improving tissue distribution compared to the original chimeric IgG4. (Angal *et al.*, Mol. Immunol. 30:105-8, 1993).

Kits/Articles of Manufacture

[00579] As an additional aspect, the present disclosure includes kits which comprise one or more compounds or compositions packaged in a manner which facilitates their use to practice methods of the present disclosure. In one embodiment, such a kit includes a compound or composition described herein (e.g., a composition comprising a humanized antibody comprising an ultralong CDR3 alone or in combination with a second agent), packaged in a container with a label affixed to the container or a package insert that describes use of the compound or composition in practicing the method. Suitable containers include, for example, bottles, vials, syringes, *etc.* The containers may be formed from a variety of materials such as glass or plastic. The container may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The article of manufacture may comprise (a) a first container with a composition contained therein, wherein the composition comprises a humanized antibody comprising an ultralong CDR3 as disclosed herein; and (b) a second container with a composition contained therein, wherein the

composition comprises a further therapeutic agent. The article of manufacture in this embodiment disclosed herein may further comprise a package insert indicating that the first and second compositions can be used to treat a particular condition. Alternatively, or additionally, the article of manufacture may further comprise a second (or third) container comprising a pharmaceutically-acceptable buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes. Preferably, the compound or composition is packaged in a unit dosage form. The kit may further include a device suitable for administering the composition according to a specific route of administration or for practicing a screening assay. Preferably, the kit contains a label that describes use of the humanized antibody comprising an ultralong CDR3 composition.

[00580] The following are examples of the methods and compositions of the disclosure. It is understood that various other embodiments may be practiced, given the general description provided above.

EXAMPLES

Example 1. Generation of a humanized antibody with an ultralong CDR3 sequence

[00581] At least 7 antibody heavy chain variable region sequences are publicly available that comprise an ultralong CDR3 of bovine origin. These sequences are shown in the alignment in Figure 1. As shown in Figure 1, the ultralong CDR3s present within these variable region sequences range in length from 56 to 61 amino acids. Strikingly, the sequence alignment indicates that the amino acid residues positioned in the N-terminal ends of the CDR3s are particularly conserved. Following the first cysteine residue in each of these variable region sequences, an amino acid sequence pattern of "TTVHQ" and variants thereof are found. This is unusual in that most heavy chain variable regions of most species end with the amino acid sequence "CAK" or "CAR". This unusual sequence motif (e.g., "TTVHQ") may be characteristic of the structure of an ultralong CDR3.

[00582] Additionally, the *Bos taurus* genome was searched and an undescribed heavy chain variable region DNA sequence in the antibody locus was found that encodes a "CTTVHQ" motif (e.g., SEQ ID NO: 1). This variable region sequence is designated herein as VH-UL (SEQ ID NO: 29). This sequence motif, discovered for the first time in searching the cow genome, is important in antibodies that comprise an ultralong CDR3.

[00583] To produce a humanized antibody comprising an ultralong CDR3, human variable regions for acceptor human frameworks were identified that are homologous to the bovine-derived VH-UL sequence (SEQ ID NO: 29). Several members of the human VH4

family were identified as being homologous to the VH-UL sequence, however none of VH4 family members contained a “CTTVHQ” motif. The sequences VH4-34*02 (SEQ ID NO: 33), VH4-39 (SEQ ID NO: 31), and VH4-59*03 (SEQ ID NO: 32) are the most homologous to the VH-UL sequence (see, e.g., Figure 4B).

5 [00584] In a first exemplary method to produce a humanized antibody comprising an ultralong CDR3, a VH4-39 human acceptor framework (SEQ ID NO: 31) is used. The nucleotides encoding the last two amino acid residues of VH4-39 (e.g., “AR”) are removed from the VH4-39 human acceptor framework, and nucleotides encoding a “TTVHQ” motif are added to the 3’ end of the DNA sequence encoding the VH4-39 human acceptor
10 framework after the nucleotides that encode the second cysteine residue of the VH4-39 human acceptor framework (e.g., after the nucleotides that code for amino acid position 97 of SEQ ID NO: 31). Next, the DNA encoding a portion of the ultralong CDR3 from BLV1H12 (e.g., the portion of the ultralong CDR3 beginning at the first glutamic acid residue
15 ETKKYQSCPDGYRERSDCSNRPACGTSDCCRVSVFGNCLTTLPVSSYSYTYNYEWHVD) is fused to the 3’ end of the polynucleotide encoding the “TTVHQ” motif. Finally, DNA encoding a partial human JH4 region, beginning with the conserved tryptophan residue common to J region sequences, is added to the 3’ end of the portion of the ultralong CDR3 sequence derived from BLV1H12. The following antibody gene encoding a partially human
20 antibody comprising an ultralong CDR3 is derived:

25 cagctgcagctgcaggagtcgggcccaggactggtgaagccttcggagaccctgtccctcacctgca
ctgtctctggtggctccatcagcagtagtagttactactggggctggatccgccagccccaggga
ggggctggagtggtattgggagtagtattattatagtagggagcacctactacaacccgtccctcaagagt
cgagtcaccatatccgtagacacgtccaagaaccagttctccctgaagctgagctctgtgaccgccg
cagacacggctgtgtattactgt**actactgtgcaccagGAAACAAAAAATACCAAAGTTGTCCTGA**
TGGGTATAGAGAACGTTCTGGATTGTAGTAACAGACCTGCTTGTGGTACTAGTGATTGTTGTCGTGTT
AGTGTTTTTGGTAATTGTCTTACTACTCTTCCTGTGAGTTATAGTTATACTTACAATTACGAATGGC
ACGTCGATGTC*TGGGGCCAGGGAACCTGGTCACCGTCTCCTCAG* (SEQ ID NO: 377)
30

- Regular Font – derived from human VH4-39
- **Underlined** – derived from VH-UL
- **BOLD BLACK** – ultralong CDR3, derived from cow BLVH12
- *ITALICS* – derived from human JH4

35

[00585] The amino acid sequence as translated from this partially human variable region gene is:

40 cagctgcagctgcaggagtcgggcccaggactggtgaagccttcggagaccctgtccctc
Q L Q L Q E S G P G L V K P S E T L S L
acctgcactgtctctggtggctccatcagcagtagtagttactactggggctggatccgc
T C T V S G G S I S S S S Y Y W G W I R
cagccccagggaaggggctggagtggtattgggagtagtattattatagtagggagcacctac
Q P P G K G L E W I G S I Y Y S G S T Y
tacaacccgtccctcaagagtcgagtcaccatatccgtagacacgtccaagaaccagttc

Y N P S L K S R V T I S V D T S K N Q F
 tcctgaagctgagctctgtgaccgccgcagacacggctgtgtattactgtactactgtg
 S L K L S S V T A A D T A V Y Y **C** T T V
 caccaggaaacaaaaaataccaaagttgtcctgatgggtatagagaacgttcggattgt
 5 H Q E T K K Y Q S C P D G Y R E R S D C
 agtaacagacctgcttgtggtactagtgttgttcgtgttagtggttttggttaattgt
 S N R P A C G T S D C C R V S V F G N C
 cttactactcttcctgtgagttatagttatactttacaattacgaatggcacgtcgcgatgtc
 L T T L P V S Y S Y T Y N Y E W H V D V
 10 tggggccagggaaccctggtcaccgtctcctcag (SEQ ID NO: 377)
W G Q G T L V T V S S (SEQ ID NO: 378)

[00586] In an embodiment, this humanized VH sequence is recombinantly fused
 in-frame with a human heavy chain constant region and paired with a light chain for
 15 recombinant antibody production.

[00587] In another exemplary method to produce a humanized antibody
 comprising an ultralong CDR3, an ultralong CDR3 derived from BLV5B8 (SEQ ID NO: 7) is
 incorporated into the VH4-34 human acceptor framework (*e.g.*, 4-34*02 (SEQ ID NO: 33);
 or 4-34*09 (SEQ ID NO: 34), along with a portion of the human JH2 region to produce a
 20 humanized antibody comprising an ultralong CDR3. Next, the nucleotides encoding the
 last two amino acid residues of the VH4-34 human acceptor framework (*e.g.*, "AR") are
 removed from the VH4-34 human acceptor framework, and nucleotides encoding a
 "TTVHQ" motif are added to the 3' end of the VH4-34 human acceptor framework after the
 nucleotides that code for the second conserved cysteine residue in the VH4-34 human
 25 acceptor framework (*e.g.*, after the nucleotides that code for amino acid position 95 of SEQ
 ID NOS: 33 or 34). Finally, DNA encoding a partial human JH2 region, beginning with the
 conserved tryptophan residue common to J region sequences, is added to the 3' end of the
 portion of the ultralong CDR3 sequence derived from BLV5B8. Thus, the following
 sequence encoding a humanized antibody variable region comprising an ultralong CDR3 is
 30 produced:

cagggtgcagctacaacagtgggggcgcaggactgttgaagccttcggagaccctgtccctcacctgcg
 ctgtctatggtgggtccttcagtgggttactactggagctggatccgccagccccaggggaaggggct
 ggagtggattggggaaatcaatcatagtgggaagcaccaactacaaccgtccctcaagagtcgagtc
 35 accatatcagtagacacgtccaagaaccagttctccctgaagctgagctctgtgaccgccgcggaca
 cggctgtgtattactgt**Actactgtgcaccag**GAAACCAGAAAAACCTGTTCTGATGGTTATATGGC
 TGTAGATAGTTGTGGTCGTGGTCAGAGTGATGGTTGTGTCAATGATTGCAATTGTTGTTATTATGGT
 TGGCGGAAGTGTGCGCAGGCAGCCTGCAATTCAAAGTTACGAATTTACGTCGATGCC***TGGGGCCGTG***
 GCACCCTGGTCACTGTCTCCTCAG (SEQ ID NO: 379)

40
 - Regular Font – derived from human VH4-39
 - **Underlined** – derived from VH-UL
 - **BLACK** – ultralong CDR3, derived from cow BLVH12
 - *ITALICS* – derived from human JH2
 45

[00588] The amino acid sequence as translated from this partially human variable region gene is:

```

5  caggtgcagctacaacagtggggcgcaggactggtgaagccttcggagaccctgtccctc
   Q V Q L Q Q W G A G L L K P S E T L S L
acctgcgctgtctatggtgggtccttcagtgggttactactggagctggatccgccagccc
10 T C A V Y G G S F S G Y Y W S W I R Q P
   ccaggggaaggggctggagtggattggggaaatcaatcatagtggaagcaccaactacaac
   P G K G L E W I G E I N H S G S T N Y N
   ccgtccctcaagagtcgagtcaccatatcagtagacacgtccaagaaccagttctccctg
15 P S L K S R V T I S V D T S K N Q F S L
   aagctgagctctgtgaccgccgcggacacggctgtgtattactgtactactgtgcaccag
   K L S S V T A A D T A V Y Y C T T V H Q
   gaaaccagaaaaacctgttctgatgggttatatggctgtagatagttgtggtcgtggtcag
   E T R K T C S D G Y M A V D S C G R G Q
15 agtgatggttgtgtcaatgattgcaattgttgttattatggttggcgggaactgtcgcagg
   S D G C V N D C N C C Y Y G W R N C R R
   cagcctgcaattcaaagttacgaatttcacgtcgatgcctggggccgtggcaccctggtc
   Q P A I Q S Y E F H V D A W G R G T L V
Actgtctcctcag (SEQ ID NO: 379)
20 T V S S (SEQ ID NO: 380)

```

[00589] In an embodiment, this humanized VH sequence is then recombinantly fused in-frame with a human heavy chain constant region and paired with a light chain for recombinant antibody production.

25 [00590] Thus, any heavy chain variable region can be paired with a light chain variable region to produce a recombinant antibody. Bovine antibodies containing ultralong VH CDR3s typically pair with a restricted set of lambda light chains. Several human VL sequences can be used to pair with the sequences above, including VL1-47, VL1-40, VL1-51, VL2-18, which are homologous to the lambda region derived from *Bos taurus*.

30

Example 2: Generation of Libraries of Polynucleotides Encoding Antibodies Comprising an Ultralong CDR3

[00591] Libraries of polynucleotides encoding antibodies that comprise an ultralong CDR3 may be constructed by any method known in the art. Such polynucleotide libraries may be present within a plurality of vectors (e.g., a library of vectors) including, for example, vectors present within a plurality of host cells (e.g., a library of host cells). The libraries may present in any known format including, in a spatially addressed format (see, e.g., WO 11/056997; and Mao *et al.* (2010) *Nat Biotech* 28:1195-1202).

40 [00592] In an exemplary method, bovine spleen and lymph nodes were obtained from Animal Technologies (Tyler, TX), or from Texas A&M University. Total RNA was isolated from bovine tissues from three different cows (MID1, MID10, and MID 11) using TRIzol reagent (Invitrogen, Carlsbad, CA, USA) followed by on column digestion of DNA using the RNeasy Mini Kit (Qiagen, Valencia, CA, USA). Alternatively, cDNA may be

obtained from the lymph nodes of a bovine immunized with an antigen (e.g., BVDV). Next, RNA quantity and quality were assessed with Nanodrop (Thermal Scientific), Qubit RNA and Agilent 2100 Bioanalyzer (Agilent, Santa Clara, CA, USA), following the manufacturer's protocols. Total RNA was used as a template for cDNA synthesis catalyzed by Superscript II (Invitrogen).

[00593] The library of amplified antibody variable regions were then subjected to deep sequencing. Briefly, bar-coded primers (Table 1) for each of the three cows (MID1, MID10, and MID11) were used to amplify V_H from bovine spleen cDNA.

10 Table 1. Bar-coded primers for deep sequencing

Primer #	Isotype	Primers
MID1 FW	IgG	CCTATCCCCTGTGTGCCTTGGCAGTCTCAGACGAGTGCGTTTGAGCGACA AGGCTGTAGGCTG (SEQ ID NO: 381)
MID1 RV	IgG	CCATCTCATCCCTGCGTGTCTCCGACTCAGACGAGTGCGTCTTTTCGGGGC TGTGGTGGAGGC (SEQ ID NO: 382)
MID10 FW	IgM	CCTATCCCCTGTGTGCCTTGGCAGTCTCAGTCTCTATGCGTTGAGCGACA AGGCTGTAGGCTG (SEQ ID NO: 383)
MID10 RV	IgM	CCATCTCATCCCTGCGTGTCTCCGACTCAGTCTCTATGCGAGTGAAGACT CTCGGGTGTGATTCAC (SEQ ID NO: 384)
MID11 FW	IgM	CCTATCCCCTGTGTGCCTTGGCAGTCTCAGTGATACGTCTTTGAGCGACA AGGCTGTAGGCTG (SEQ ID NO: 385)
MID11 RV	IgM	CCATCTCATCCCTGCGTGTCTCCGACTCAGTGATACGTCTAGTGAAGACTC TCGGGTGTGATTCAC (SEQ ID NO: 386)

[00594] Next, the amplicons of V_H were purified from 2% agarose gels and deep sequenced according to Roche 454 GS FLX instructions. Multiple alignments were performed with the MUSCLE algorithm (Edgar (2004) Nucleic Acids Research 32:1792-1797). MUSCLE was executed to generate multiple long CDR H3 nucleotide alignments with relatively high gap open (-20.0) and gap extend (-10.0) penalties due to the large amount of heterogeneity observed in the sequences. Local alignment was executed using the Smith-Waterman algorithm with the following settings, match score = 2.0, mismatch penalty = -1.0, gap opening penalty = -2.0, and gap extension penalty = -0.5. CDR H3s were defined by the third residue following the conserved cysteine in framework 3 to the residue immediately preceding the conserved tryptophan in framework 4. V_HBUL was identified by BLAST searching the bovine genome (assembly Btau_4.6.1) with multiple ultralong V_H sequences identified by deep sequencing. The deep sequencing identified a

total of 11,728 ultralong CDR3 sequences with having a length between 44 and 69 amino acid residues. The results of the deep sequencing are summarized in Table 2 below.

Table 2. Summary of deep sequencing results from bovine spleen

Source (Bar code)	Cow#1 (MID1)	Cow#1 (MID10)	Cow#2 (MID11)
Ig Class	IgG	IgM	IgM
CDR H3 length range	44-66	44-68	44-69
Number of unique cysteine patterns	655	449	847
Total number of unique long CDR H3 sequences	5633	1639	4456

5

[00595] The results of the deep sequencing also revealed that ultralong CDR3 comprise a cysteine motif (e.g., a pattern of cysteine residues) that comprises between 3 and 12 cysteine residues. Representative examples of cysteine patterns are shown for the deep sequencing run for three different cows (MID1, MID10, and MID11) as well as their abundance in the run (Tables 3-5). The cysteines in the ultralong CDR3 regions are symbolized as "C". The amino acids between two cysteines are symbolized as "X_n". Exemplary sequences comprising cysteine motifs identified from the deep sequencing are presented in Figures 2A-C.

10

15

Table 3. Cysteine patterns identified in ultralong CDR3s from MID1

Cysteine pattern (MID1)	Abundance (%)
CX₁₀CX₅CX₅CXCX₇C	10.44%
CX₁₀CX₆CX₅CXCX₁₅C	8.11%
CX₁₁CXCX₅C	5.22%
CX₁₁CX₅CX₅CXCX₇C	2.56%
CX₁₀CX₆CX₅CXCX₁₃C	1.47%
CX₁₀CX₅CXCX₄CX₈C	1.19%
CX₁₀CX₆CX₆CXCX₇C	1.08%
CX₁₀CX₄CX₇CXCX₈C	1.05%
CX₁₀CX₄CX₇CXCX₇C	0.91%

Cysteine pattern (MID1)	Abundance (%)
CX ₁₃ CX ₈ CX ₈ C	0.91%
CX ₁₀ CX ₆ CX ₅ CXCX ₇ C	0.59%
CX ₁₀ CX ₅ CX ₅ C	0.57%
CX ₁₀ CX ₅ CX ₆ CXCX ₇ C	0.50%
CX ₁₀ CX ₆ CX ₅ CX ₇ CX ₉ C	0.43%
CX ₉ CX ₇ CX ₅ CXCX ₇ C	0.41%
CX ₁₀ CX ₆ CX ₅ CXCX ₉ C	0.36%
CX ₁₀ CXCX ₄ CX ₅ CX ₁₁ C	0.32%
CX ₇ CX ₃ CX ₆ CX ₅ CXCX ₅ CX ₁₀ C	0.32%
CX ₁₀ CXCX ₄ CX ₅ CXCX ₂ CX ₃ C	0.30%
CX ₁₆ CX ₅ CXC	0.23%

Table 4. Cysteine patterns identified in ultralong CDR3s from MID10

Cysteine pattern (MID10)	Abundance (%)
CX ₁₀ CXCX ₄ CX ₅ CXCX ₂ CX ₃ C	2.87%
CX ₁₀ CX ₅ CX ₅ C	0.73%
CX ₁₀ CXCX ₄ CX ₅ CX ₁₁ C	0.67%
CX ₆ CX ₄ CXCX ₄ CX ₅ C	0.61%
CX ₁₁ CX ₄ CX ₅ CX ₆ CX ₃ C	0.55%
CX ₈ CX ₂ CX ₆ CX ₅ C	0.43%
CX ₁₀ CX ₅ CX ₅ CXCX ₁₀ C	0.37%
CX ₁₀ CXCX ₆ CX ₄ CXC	0.31%
CX ₁₀ CX ₅ CX ₅ CXCX ₂ C	0.31%
CX ₁₄ CX ₂ CX ₃ CXCXC	0.31%
CX ₁₅ CX ₅ CXC	0.31%
CX ₄ CX ₆ CX ₉ CX ₂ CX ₁₁ C	0.31%

CX ₆ CX ₄ CX ₅ CX ₅ CX ₁₂ C	0.31%
CX ₇ CX ₃ CXCXCX ₄ CX ₅ CX ₉ C	0.31%
CX ₁₀ CX ₆ CX ₅ C	0.24%
CX ₇ CX ₃ CX ₅ CX ₅ CX ₉ C	0.24%
CX ₇ CX ₅ CXCX ₂ C	0.24%
CX ₁₀ CXCX ₆ C	0.18%
CX ₁₀ CX ₃ CX ₃ CX ₅ CX ₇ CXCX ₆ C	0.18%
CX ₁₀ CX ₄ CX ₅ CX ₁₂ CX ₂ C	0.18%

Table 5. Cysteine patterns identified in ultralong CDR3s from MID11

Cysteine pattern (MID11)	Abundance (%)
CX ₁₂ CX ₄ CX ₅ CXCXCX ₉ CX ₃ C	1.19%
CX ₁₂ CX ₄ CX ₅ CX ₁₂ CX ₂ C	0.96%
CX ₁₀ CX ₆ CX ₅ CXCX ₁₁ C	0.92%
CX ₁₆ CX ₅ CXCXCX ₁₄ C	0.70%
CX ₁₀ CX ₅ CXCX ₈ CX ₆ C	0.52%
CX ₁₂ CX ₄ CX ₅ CX ₈ CX ₂ C	0.49%
CX ₁₂ CX ₅ CX ₅ CXCX ₈ C	0.47%
CX ₁₀ CX ₆ CX ₅ CXCX ₄ CXCX ₉ C	0.45%
CX ₁₁ CX ₄ CX ₅ CX ₈ CX ₂ C	0.45%
CX ₁₀ CX ₆ CX ₅ CX ₈ CX ₂ C	0.43%
CX ₁₀ CX ₆ CX ₅ CXCX ₈ C	0.36%
CX ₁₀ CX ₆ CX ₅ C	0.31%
CX ₁₀ CX ₆ CX ₅ CXCX ₃ CX ₈ CX ₂ C	0.29%
CX ₁₀ CX ₆ CX ₅ CX ₃ CX ₈ C	0.29%
CX ₁₀ CX ₆ CX ₅ CXCX ₂ CX ₆ CX ₅ C	0.25%
CX ₇ CX ₆ CX ₃ CX ₃ CX ₉ C	0.25%

CX₉CX₈CX₅CX₆CX₅C	0.22%
CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C	0.20%
CX₁₀CX₆CX₅CXCX₁₃C	0.20%
CX₁₀CX₆CX₅CXCX₂CX₈CX₄C	0.20%

[00596] Bovine V_H regions were amplified from cDNA prepared in example 9 using primers 5'-TTGAGCGACAAGGCTGTAGGCTG-3' (SEQ ID NO: 387) and 5'-CTTTCGGGGCTGTGGTGG-AGGC-3' (SEQ ID NO: 388) producing a library of antibody
5 variable region cDNA biased for ultralong CDRs. Next, the mixture of V_H regions was assembled by overlap PCR with bovine C_H1 and human IgG Fc. Briefly, EcoRI and NheI sites were incorporated for ligation into pFUSE expression vector, to afford a full-length heavy chain library ready for expression in mammalian cells. The ligation product was transformed into *E. coli* and 500 single *E. coli* transformants were picked. Each
10 transformant was then grown overnight in a separate vessel and DNA from each colony was extracted using Qiagen minprep kits (Qiagen, Inc.) and sequenced by BATJ, Inc. (San Diego, CA) using the oligo 5'-AGATCCAAGCTGTGACCGGC-3' (SEQ ID NO: 389). Sequences were analyzed using VectorNTI (Invitrogen, Inc. Carlsbad, CA). Duplicative sequences, sequences with no insert, and sequences encoding a CDR shorter than 35
15 residues were excluded. 132 clones containing unique long CDR heavy chain sequences were selected. Each heavy chain in the 132 member library was then co-transfected in parallel with pFUSE expression vector encoding the invariant bovine light chain BLV1H12 (SEQ ID NO: 412) into 293T cells, to generate a small spatially addressed library (Mao *et al.* (2010) *Nat Biotech* 28:1195-1202). 130,000 293T cells per well were plated in 24 well
20 plates and grown overnight in 500ul DMEM media (Invitrogen) with 10% FBS (Invitrogen), and Penicillin/streptomycin/ glutamine (Invitrogen) at 37°C and 5% CO₂. 0.5μg of Hc-encoding pFUSE vector and 0.5μg of Lc-encoding pFUSE vector were added to 25μl of optimem (Invitrogen). 1μl of Lipofectamine 2000 or 293Fectin transfection reagent (Invitrogen) was added to 25μl of optimem, and incubated 5 minutes. Next, the DNA-
25 optimem mix and transfection reagent-optimem mix were combined and incubated 15 minutes, added to 293T cells, and allowed to incubate on cells 4-6 hours. Then media was aspirated from wells and replaced with fresh media, and cells were allowed to grow and secrete IgG into the media for 4 days. Cell-culture supernatants containing IgG were harvested in 96 well format for further testing. The chimeric antibodies were quantified by
30 sandwich ELISA detecting human F_c and screened for binding to BVDV by ELISA.

[00597] Antibodies were then secreted into culture media and harvested in a 96 well format to generate a small spatially addressed library for further testing including, screening for binding to BVDV by ELISA. For example, an ELISA was conducted to screen the antibody library for binding to BVDV. Briefly, killed BVDV (0.2 µg) in 100 µL DPBS was coated on 96-well MaxiSorp ELISA plates (Nunc) for 1 hour at 37°C. Next, the plates were blocked with 200 µL 3% BSA solution in DPBST, Dulbecco's phosphate buffered saline, 0.25% Tween 20) for 1 hour at 37°C. Samples were then incubated with 3% BSA in DPBST for 1 hour at 37°C. Subsequently, wells were washed 5 times with 200 µL DPBST. Next, Goat Anti-Human IgG (Fc) – HRP conjugated antibody (KPL Inc.) was added at a 1:1,000 dilution in blocking solution and incubated for 1 hour at 37°C. Wells were then washed 10 times with 200 µL DPBST. A 100 µL working solution of QuantaBlu (Pierce) was added to each well and incubated for 5 minutes at room temperature before plates were read in a SpectraMax M5 plate reader at ex325/em420 nm. Several candidate binders were identified. Clone H12 has a 63-residue CDR3 with 6 cysteine residues and was able to strongly bind BVDV in a dose dependent fashion.

[00598] Additionally, binding of the chimeric recombinant antibodies to BVDV antigens was evaluated by immunocytometric analysis of transfected human embryonic kidney (HEK) 293A cells (Invitrogen), as previously described (see, e.g., Njongmeta *et al.* (2012) *Vaccine* 30:1624-1635). Briefly, HEK 293A monolayers grown in 6-well tissue culture plates were transfected with 2 µg/well of plasmid (pCDNA3.3, Invitrogen) encoding BVDV antigens (N^{pro}, E2, or non-structural proteins NS2-3) using Lipofectamine 2000 reagent (Invitrogen), and incubated for 48 hr at 37 °C with 5 % CO₂. The monolayers were fixed with ice-cold 100% methanol for 10 minutes, rinsed with PBS, and after blocking for 1 hour with PBS containing 5% fetal bovine serum (blocking buffer), the monolayers were incubated at room temperature for 1 hr with 10 µg/ml of a mouse anti-FLAG M2-alkaline phosphatase (AP)-conjugate (Sigma) in blocking buffer or 10 µg/ml of the chimeric recombinant antibodies (H12 or B8). Monolayers transfected with empty vector were similarly reacted to serve as negative controls and, following washes in blocking buffer, the monolayers probed with the chimeric recombinant antibodies were incubated with a 1/200 dilution of AP-conjugated goat anti-Human IgG (Fc specific) mAb (Sigma) in blocking buffer for 1 hr. Following washes in blocking buffer, the AP activity in all the wells was detected using Fast Red AS-MX substrate (Sigma). Stained cells were visualized and photographed using an IS70 inverted optical microscope (Olympus, Japan) equipped with a camera. H12 strongly binds HEK293A cells transfected with the NS2-3 non-structural proteins of BVDV but weakly bound to untransfected cells while B8 had weak binding to both HEK293A cells transfected with the NS2-3 non-structural proteins of BVDV and untransfected cells.

Example 3: Construction of Vector Comprising an Antibody with an Ultralong CDR3

[00599] Any vector known in the art may be used, or may be modified to be used, for cloning and/or expression of a nucleotide sequence encoding a heavy chain variable region that comprises an ultralong CDR3. Such vectors may optionally comprise, or be modified to comprise, a nucleotide sequence encoding the Fc portion of a human immunoglobulin (*e.g.*, IgG) linked to the nucleotide sequence encoding the heavy chain variable region comprising an ultralong CDR3. Additionally, the nucleotide sequence encoding the heavy chain variable region comprising the ultralong CDR3 may be modified according to known methods such that the ultralong CDR3 can accept a nucleotide sequence encoding a non-bovine (*e.g.*, non-antibody or human) sequence including, for example, by unidirectional cloning with restriction enzymes. Any vector known in the art may also be used, or may be modified to be used, for cloning and/or expression of a nucleotide sequence encoding a light chain variable region.

[00600] In an exemplary method, a vector may be modified by recombinant techniques to comprise a nucleotide sequence encoding a heavy chain variable region having an ultralong CDR3 linked to a nucleotide sequence encoding an IgG Fc. Briefly, a nucleotide sequence encoding CH1-CH2-CH3 of human IgG1 (SEQ ID NO: 390) was amplified by PCR and assembled via strand overlap elongation. Next, a compatible cohesive-end strategy was employed to replace the human Fc sequence encoded in vector pFUSE-hIgG2-Fc2 (SEQ ID NO: 457, InVivogen, San Diego CA) with CH1-CH2-CH3 of human IgG1, which destroyed the existing 3' NheI site in the pFUSE-hIgG2-Fc2 vector and generated an NheI site at the 5' end of SEQ ID NO: 390. The modified pFUSE-hIgG2-Fc2 vector ("HC pFuse", SEQ ID NO: 458) allows insertion of VH fragments between the signal sequence and the nucleotide sequence encoding CH1-CH2-CH3 of human IgG1 using the existing EcoRI site and the newly introduced NheI site. For example, the VH region from bovine antibody BLV1H12 (SEQ ID NO: 392) was amplified by PCR and subcloned in-frame between the signal sequence and nucleotide sequence encoding CH1-CH2-CH3 of human IgG1 in the HC pFuse vector using EcoRI and NheI restriction enzymes generating SEQ ID NO: 393.

[00601] Alternatively, nucleotide sequences encoding non-bovine sequences were inserted into the nucleotide sequence encoding the CDR3 of BLV1H12 heavy chain. Briefly, a pair of BsaI sites were introduced by PCR strand overlap extension into the nucleotide sequence encoding the CDR3 of BLV1H12 (SEQ ID NO: 395). The modified nucleotide sequence incorporating the BsaI sites within the CDR3 encoding sequence of BLV1H12 was then subcloned in-frame between the signal sequence and the nucleotide

sequence encoding CH1-CH2-CH3 of human IgG1 in the HC pFuse vector described above using EcoRI and NheI restriction enzymes. Next, non-antibody sequences including cDNA encoding for interleukin 8 (IL-8, SEQ ID NO: 475), interleukin 21 (IL-21, SEQ ID NO: 480) and CXCL12/SDF-1alpha (SEQ ID NO: 479) (Origene) and oligoprimers encoding the peptide hormone somatostatin (SEQ ID NO: 477), the venom peptides ProTx-II (SEQ ID NO: 481) and chlorotoxin (SEQ ID NO: 478), and the synthetic conotoxin peptide ziconotide (SEQ ID NO: 476) (IDTDNA) were modified by PCR amplification using oligoprimers with BsaI flanks to comprise BsaI flanks to produce non-bovine sequences with BsaI flanks for insertion into the CDR3 of BLV1H12. Next, the BsaI flanks in the modified non-antibody sequences were cut with BsaI and ligated, in frame, with the BLV1H12 BsaI digested vector, thereby inserting the nucleotide sequence encoding the non-antibody sequence into the BLV1H12 heavy chain variable region and replacing a portion (*e.g.*, cysteine-rich portion) of the ultralong CDR3, and in-frame with the sequence encoding BLV1H12 variable region and CH1-CH2-CH3.

[00602] Similarly, a vector was modified by recombinant techniques to comprise a nucleotide sequence encoding a light chain. Briefly, the BLV1H12 light chain sequence (SEQ ID NO: 412) was amplified by PCR to introduce EcoRI and NheI ends. The BLV1H12 light chain variable region sequence was then digested with EcoRI and NheI and subcloned into pFUSE-hIgG2-Fc2, deleting the portion of pFUSE-hIgG2-Fc2 encoding Fc, and replacing it with only light chain sequence (SEQ ID NO: 459).

[00603] In addition, overlap PCR was used to amplify and join sequence from BLV1H12 light chain variable region and human lambda light chain constant region (SEQ ID NO: 474). This was amplified by PCR to introduce EcoRI and NheI ends. The hybrid BLV1H12 with human lambda sequence was then digested with EcoRI and NheI and subcloned into pFUSE-hIgG2-Fc2, deleting the portion of pFUSE-hIgG2-Fc2 encoding Fc, and replacing it with only light chain sequence.

Example 4: Expression of Polynucleotides Coding for Antibodies that Comprise an Ultralong CDR3

[00604] Polynucleotides coding for antibodies comprising an ultralong CDR3 may be expressed including, transiently expressed, in a host cell by any method known in the art.

[00605] In an exemplary method, the vectors comprising the heavy and light chains generated in Example 2 may be transfected into either 293T cells or Freestyle™ 293-F cells.

[00606] For example, 130,000 293T cells per well were plated in 24 well plates and grown overnight in 500 µl DMEM media (Invitrogen) with 10% FBS (Invitrogen), and penicillin/streptomycin/glutamine (Invitrogen) at 37°C and 5% CO₂. Next, 0.5 µg of Heavy chain-encoding pFuse vector (*e.g.*, generated in Example 2) and 0.5 µg of Light chain-encoding pFuse vector (*e.g.*, generated in Example 2) were added to 25 µl of Opti-MEM (Invitrogen). Subsequently, 1 µl of Lipofectamine 2000 or 293Fectin transfection reagent (Invitrogen) was added to 25 µl of Opti-MEM, and incubated 5 minutes. Then the dna-Opti-MEM solution and transfection reagent-Opti-MEM solution were combined and incubated for 15 minutes, added to 293T cells, and allowed to incubate on cells for 4-6 hours. Then media was aspirated from wells and replaced with fresh media, and the cells were allowed to grow and secrete IgG into the media for 2-6 days.

[00607] Additionally, for example, 1x10⁶ 293Freestyle suspension cells/ml were grown overnight in Freestyle™293 Expression Medium (Invitrogen), and penicillin/streptomycin/ glutamine (Invitrogen) at 37°C and 5% CO₂. Next, for each milliliter of cells, a 30µl solution was made in buffer PBS comprising 0.5 µg Heavy chain-encoding pFuse DNA and 0.5 µg of Lc-encoding pFuse. Additionally, for each milliliter of cells, a solution comprising 1 µl of 293Fectin transfection reagent was added to 30µl PBS and incubated 5 minutes. Subsequently, the DNA and lipofectamine solutions were combined and incubated 15 minutes before adding the mixture to the cells. The cells were then allowed to grow and secrete IgG into media for 2-6 days.

[00608] After the growth period for the 293T cells or Freestyle™ 293-F suspension cells, media was harvested, and IgG secreted into the supernatant was evaluated by sandwich ELISA. Briefly, Fc-specific anti Human IgG (Cat I2136, Sigma-Aldrich) was diluted 1:1000 in PBS and coated onto maxisorp plates (Nunc). Next, plates were blocked with 2% BSA in TBST, washed, and IgG secreted supernatants were incubated for 1 hour. After washing with TBST, the appropriate HRP conjugated anti-Light chain antibody is diluted 1:1000 in TBST and incubated 1 hour (anti kappa-HRP, Cat#A-7164, Sigma-Aldrich; anti-lambda-HRP, Cat# 2070-05, Southern Biotech). After washing with TBST, HRP was detected with TMB (Cat#TMBS-1000-01, BioFX), and neutralized with 0.6M H₂SO₄. Subsequently, absorbance (A450) was measured, and compared to a standard curve of known IgG concentration to determine the concentration of human antibody in cell culture supernatant.

[00609] The yield of antibody from supernatants from cells transfected with heavy chain constructs encoding BLV1H12 with human CH1-CH2-CH3 and a replacement of at least a portion of CDR3 (generated in Example 3) and a light chain construct encoding BLV1H12 LV-human lambda LC was determined by ELISA, and normalized to the yield of

BLV1H12-IL8 (Table 6). The ELISA indicated that some sequences (e.g., zinconotide) that were used to replace a portion of CDR3 have a minor impact on expression of the antibody, while some sequences (e.g., somatostatin, chlorotoxin, SDF1(alpha), IL21) that were used to replace a portion of CDR3 had a modest impact on expression of the antibody, and other sequences (ProTxII) had an even greater impact on expression of the antibody, as compared to BLV1H12-IL8 (the highest expressor) (Table 6).

Table 6. Expression yield of BLV1H12 with replacements to Long CDR3

Heavy Chain V region	Light Chain	Normalized Yield (% of highest expressor)
BLV1H12-IL8	BLV1H12-HuLambdaC	100
BLV1H12-Zinconotide	BLV1H12-HuLambdaC	94.7
BLV1H12-Somatostatin	BLV1H12-HuLambdaC	46.5
BLV1H12-Chlorotoxin	BLV1H12-HuLambdaC	39.7
BLV1H12-SDF1(alpha)	BLV1H12-HuLambdaC	38.9
BLV1H12-IL21	BLV1H12-HuLambdaC	32.3
BLV1H12-ProTxII	BLV1H12-HuLambdaC	2.1

10 **Example 5: Evaluation of Human Germline Sequences Comprising an Ultralong CDR3 with a Non-Bovine Sequence**

[00610] Human germline sequences that comprise an ultralong CDR3 may be evaluated by any method known in the art to identify those human germline sequences that permit expression of an ultralong CDR3 that comprises a non-bovine sequence.

15 [00611] In an exemplary method, overlap PCR was used to insert Bsal sites into human germline variable regions VH1-24, VH1-46, VH1-69, VH3-23, VH4-4, and VH4-34. Next, PCR was used to introduce EcoRI and NheI sites for sub-cloning in between the signal sequence and CH1-CH2-CH3 region of the HC pFuse vector comprising BLV1H12 generated in Example 2. Subsequently, the IL-8-GSG cassette (SEQ ID NO: 399) was introduced into the Bsal sites of each human germline construct, giving VH1-24+CDR3-IL8 (SEQ ID NO: 425), VH1-46+CDR3-IL8 (SEQ ID NO: 426), VH1-69+CDR3-IL8 (SEQ ID NO: 427), VH3-23+CDR3-IL8 (SEQ ID NO: 428), VH4-4+CDR3-IL8 (SEQ ID NO: 429), and VH4-34+CDR3-IL8 (SEQ ID NO: 430). Next, each pFuse vector encoding a human germline heavy chain and CDR3-IL8 construct was co-transfected with pFuse vector encoding BLV1H12 LV (SEQ ID NO: 474), and allowed to express and secrete these IgGs into the media as described in Example 4. ELISA was then used to determine IgG yield as described in Example 4. Yields of the constructs were determined, and normalized to the highest expressing construct. Surprisingly, the CDR3 from cow is not readily transposable

to any human heavy chain variable region. VH4-4 had the highest yield, despite VH4-34 having the closest sequence homology to the BLV1H12 sequence (Table 7).

5 Table 7. Expression Yield of Humanized Heavy Chains with Ultralong CDR3s (removed last column of values, with IgG nM, only showing normalized values.

Heavy Chain Variable Region:	Normalized Yield (% of highest expressor)
BLV1H12+CDR3IL8	100
BLV1H12	58.4
VH1-24 BLVCDR3IL8	4.4
VH1-46 BLVCDR3IL8	2.4
VH1-69 BLVCDR3IL8	1.4
VH3-23 BLVCDR3IL8	14.7
VH4-4 BLVCDR3IL8	22.5
VH4-34 BLVCDR3IL8	5.4

Example 6: Evaluation of Antibodies that Comprise an Ultralong CDR3 with a Non-Bovine Sequence

10 [00612] Antibodies comprising an ultralong CDR3 with a non-bovine sequence (e.g., a replacement of at least a portion of the CDR3 with a non-bovine sequence) including, humanized antibodies, may be evaluated by any method known in the art for binding of the non-bovine sequence to its binding partner including, for example, flow cytometry.

15 [00613] In an exemplary method, BLV1H12 IgG comprising a CDR3 with an IL-8 insert (e.g., the IL-8 sequence replaced a portion of the CDR3) was evaluated for binding to CXCR1 expressing cells by flow cytometry. Briefly, a cell line expressing functionally validated CXCR1 derived from U2OS cells was obtained from DiscoverX and cultured per manufacturer's instructions (Cat# 93-0226C3, DiscoverX Corporation, Freemont, CA). The parental cell line U2OS was obtained from ATCC and cultured under the same conditions
20 as the CXCR1 cells. Cell culture supernatants were then tested for binding to cells by flow cytometry. The adherent U2OS or CXCR1-U2OS cells were dissociated with Accutase (Innovative Cell Technologies, Inc., San Diego, CA), neutralized with an equal volume of media containing 10% serum, centrifuged at 1000 g, and resuspended in PBS with 2%BSA. Next, cells were dispensed into microtiter plates to achieve between 30,000 to
25 300,000 cells per well, centrifuged again, and resuspended in cell culture supernatant containing expressed IgG, or a dilution of IgG-containing cell culture supernatant. A fluorescent-conjugated anti-Human Fc antibody was used to detect binding of the expressed antibody to cells. Subsequently, cell fluorescence was measured by flow

cytometry (e.g., FACS), and median Arbitrary Fluorescence Units (AFU) were calculated for each combination of antibody and cell type tested, revealing the extent of IgG binding to those cells. The ratio of median fluorescence (IgG binding) of CXCR1-U2OS cells versus U2OS parental cells shows that the BLV1H12 IgG comprising a CDR3 with an IL-8 insert
 5 has specificity for CXCR1 (Table 8).

Table 8. Binding of BLV1H12 frameworks to CXCR1 U2OS cells

	ULcowV (Median Arbitrary Fluorescence Units (AFU))	ULcowV+IL8 (Median Arbitrary Fluorescence Units (AFU))
Parental U2OS	4	76
CXCR1-U2OS	4	707

[00614] Additionally, the human germline CDR3-IL8 IgGs described in Example 5
 10 were evaluated for binding to CXCR1 expressing cells by flow cytometry as described above. IgG binding to CXCR1-U2OS cells is indicated by higher fluorescence values, but non-specific binding to U2OS parental cells is also detectable. For a given IgG with CDR3-IL-8, specific binding to CXCR1 is revealed by the ratio of fluorescence on CXCR1-U2OS cells to the fluorescence on parental U2OS cells (Table 9). While the BLV1H12 supported
 15 strong and specific binding to CXCR1 (Table 8), the IL-8 in human germline IgGs showed either weaker binding to CXCR1, or non-specific interaction as seen in strong binding to both CXCR1-U2OS cells as well as the parental control cells. The most specific interaction to CXCR1 of IL-8 within a human germline IgG was seen with VH4-34.

20 **Table 9.** Binding of IgGs with CDR3-IL8 to CXCR1 U2OS cells

IgG with HC V region:	Parental U2OS	CXCR1-U2OS	Ratio of CXCR1 / Parental
BLV1H12	5	9	1.9
VH1-24 +CDR3-IL8	5	23	4.3
VH1-46 +CDR3-IL8	5	6	1.1
VH1-69 +CDR3-IL8	20	78	4.0
VH3-23 +CDR3-IL8	534	1443	2.7
VH4-4 +CDR3-IL8	1007	3079	3.1
VH4-34 +CDR3-IL8	70	648	9.3

[00615] In another exemplary method, IL-8 activation of the CXCR1 receptor was tested using the CXCR1-U2OS DiscoverX cells described above (Cat# 93-0226C3, DiscoverX). The DiscoverX cells are engineered such that upon activation by ligand, the
 25 resultant beta-arrestin recruitment to the GPCR also causes activation of a luminescent

reporter enzyme present cell line. Lysis reagents and luminescence substrate are included in the PathHunter Detection Kit (Cat# 93-001, DiscoverX) which was used following manufacturer's instructions. U2OS-CXCR1 cells were plated at 15,000 cells per well and serum starved overnight in EMEM media without serum. Next, media was removed and cells were incubated 1 hour at 37°C with 80 µl of 1:1 EMEM and PBS, containing dilutions of either IL-8 or IgG with IL-8 insertion in CDR3, or control antibody. After 1 hour, 40 µl of PathHunter Detection reagent mix was added. Subsequently, after an hour at room temperature the luminescence in each well was measured using a luminescence plate reader. Higher luminescence signal reveals more activation of the CXCR1 receptor at the tested concentrations of IL-8 or IgG. IgGs were purified by scaling up the transfection method of Example 4, and purification of the IgG from media using protein A sepharose following manufacturer's instructions (Cat# 17-1279-03 GE Healthcare) and dialysis post-elution into PBS. Protein yield was determined by A280 and calculated molar extinction coefficient. Increasing concentrations of BLV1H12-IL8 IgG activated CXCR1 somewhat similarly to the activation observed with IL-8, whereas increasing concentration of BLV1H12 IgG had no effect on CXCR1 activation (Table 10).

Table 10. Activation of CXCR1 by IL-8 as soluble cytokine or as CDR3 replacement

nM	IL-8 (average RLU)	BLV1H12-IL8 IgG (average RLU)	IgG_BVL1H12 IgG (average RLU)
500.00	5720	6641	484
166.67	6770	6207	525
55.56	7365	5509	511
18.52	7295	5071	529
6.17	5869	4428	517
2.06	4110	3645	495
0.69	1939	2651	516
0.23	940	1515	455
0.08	504	822	461
0.01	440	435	460

[00616] Additionally, antibodies in which IL-8 was grafted into the VH4-4 human germline variable region were tested for activation of CXCR1. IgG with VH4-4+CDR3-IL8 (SEQ ID NO: 429) was compared to BLV1H12+CDR3-IL8, and IgGVH4-4 CDR3 BsaI (lacking the IL-8 insert) (SEQ ID NO: 431). A single concentration of soluble IL-8 was tested as a positive control for assay function at 31.5nM and gave activation of 55525 average RLU. (Table 11). Although the IgG with insertion of IL-8 into the human VH4-4

variable region did activate CXCR1, it was not as potent in equal dose to the IgG with BLV1H12 sequence.

Table 11. Activation of CXCR1 by IgGs with CDR3 IL-8

IgG (nM)	BLV1H12 +CDR3-IL8 (average RLU)	VH4-4 +CDR3-IL8 (average RLU)	VH4-4 +CDR3-BsaI (average RLU)
261.500	56349	36481	7545
87.167	52128	29623	7139
29.056	47621	24889	7104
9.685	39280	15545	7174
3.228	34544	10233	7034
1.076	23700	8012	7184
0.359	13278	7338	6687
0.120	8867	7513	7331
0.040	7736	7323	6973
0.013	7329	6956	7048
0.004	7240	7067	7387
0.001	7437	7142	7500

5

Example 7: Modification of humanized antibodies comprising an ultralong CDR3

[00617] Humanized antibodies generated in Example 5 may be modified to comprise one or more amino acid substitutions in heavy chain CDR1 and/or CDR2. Such amino acid substitutions may be introduced into a human germline CDR1 and CDR2 at positions that are hypothesized to interact with CDR3.

10

[00618] In an exemplary method, certain residues in VH4-34 CDR1 and/or CDR2 were substituted with corresponding CDR1 and/or CDR2 residues from BLV1H12. Briefly, the pFuse vector encoding heavy chain VH4-34 CDR3-IL8 was modified by overlap PCR to replace all of CDR1 with sequence from BLV1H12 (VH4-34+CDR3-IL8_CDR1 Cow, SEQ ID NO: 432) or all of CDR2 with sequence from BLV1H12 (VH4-34+CDR3-IL8_CDR2 Cow, SEQ ID NO: 433). Additionally, overlap PCR was used to introduce point mutants G31D and Y32K into CDR1 (VH4-34+CDR3-IL8_CDR1 G31D, Y32K, SEQ ID NO: 434) or point mutant E50S into CDR2 (VH4-34+CDR3-IL8_CDR2 E50S, SEQ ID NO: 435). These heavy chain constructs were paired with BLV1H12 light chain and were expressed as described in Example 4. Yield and CXCR1 binding were determined as described in Example 6 (Table 12).

15

20

Table 12. CDR1 and 2 modification of VH4-34 CDR3-IL8 IgG

Heavy chain	IgG Yield normalized	FACS CXCR1/Parental Ratio
BLV1H12	100	1.9
VH4-34+CDR3-IL8	20	9.3
VH4-34+CDR3-IL8 + Cow CDR1	29	2.1
VH4-34+CDR3-IL8 + Cow CDR2	9	8.0
VH4-34+CDR3-IL8 + E31D,Y32K	19	8.1
VH4-34+CDR3-IL8 + E50S	27	4.0

Example 8: Pairing of heavy chain comprising an ultralong CDR3 with a human light chain

[00619] Antibodies that comprise an ultralong CDR3 including, antibodies that
 5 comprise an ultralong CDR3 where at least a portion of the CDR3 has been replaced by a non-bovine sequence, may be paired with a human light chain.

[00620] In an exemplary method, BLV1H12+CDR3-IL8 flexibility to pair
 productively with human germline light chains was explored by co-transfecting pFuse
 vector encoding heavy chain BLV1H12+CDR3-IL8 with pFuse vectors encoding each of
 10 several human germline light chains (e.g., SEQ ID NOs: 445-456), or BLV1H12 light chain (SEQ ID NO: 412). These human germline light chain sequences were synthesized (Genscript, Inc) and amplified by PCR as in Example 3 for subcloning via EcoRI and NheI into the pFUSE LC vector. BLV1H12+CDR3-IL8 encoding pFUSE vector was cotransfected
 15 with pFUSE vectors encoding light chains as in Example 3. Secreted IgGs were then tested by FACS as in Example 5 for specific binding to CXCR1 cells (Table 13). Human germ line light chains did not readily support the function of IL-8 when expressed as BLV1H12+CDR3.

Table 13. Evaluation of BLV1H12+CDR3-IL8 when paired with germline human light
 20 chains

Light chain	FACS CXCR1/Parental Ratio
A20J1	0.8
A27J3	0.9
L6J1	0.8
L25J1	1.3
V1-2J7	0.8
V1-7J1	0.6
V1-11J2	0.8
V1-13J5	2.9
V1-16J6	0.6

V2-13J2	0.9
V2-14J4	1.0
V2-15J7	1.1
V2-17J2	2.0
V3-4J1	1.2
V5-4J2	0.8
BLV1H12	5.0

[00621] Alternatively, using the BLV1H12 light chain variable region as a guide, the human germline sequences identified with closest homology were V1-47 and V1-51. These light chain sequences were synthesized (SEQ ID NO: 455, 456) (IDTDNA, Inc.) with the desired EcoRI and NheI restriction sites for subcloning into the pFuse vector. Subsequently, pFuse vector encoding V1-47 or V1-51 were paired with pFuse vectors encoding VH4-34+CDR3-IL8, the VH4-34+CDR3-IL8 with the CDR1 or CDR2 modifications described in Example 6 (SEQ ID NOs: 432-435), as well as vectors encoding the combinations of both CDR1 and CDR2 modifications (e.g., VH4-34+CDR3-IL8_CDR1 Cow_CDR2 Cow, SEQ ID NO: 436; VH4-34+CDR3-IL8_CDR1 Cow_CDR2 E50S, SEQ ID NO: 437; VH4-34+CDR3-IL8_CDR1 G31D,Y32K_CDR2 Cow, SEQ ID NO: 438; VH4-34+CDR3-IL8_CDR1 G31D,Y32K_CDR2 E50S, SEQ ID NO: 439). These IgGs were expressed, yield was determined by ELISA, and CXCR1 binding specificity was measured by flow cytometry as described above (Table 14).

Table 14. Expression yield and CXCR1 specificity for VH4-34 CDR3-IL8 IgGs having CDR1 and CDR2 modifications and paired with Lc V1-47 or V1-51

HC VH4-34 CDR3-IL8+		nM IgG		CXCR1/Parental Ratio	
Hc-CDR1	Hc-CDR2	V1-47	V1-51	V1-47	V1-51
G31D Y32K	VH4-34	1	2.5	3.6	3
Cow	VH4-34	3	8.5	5.1	2.4
VH4-34	VH4-34	1.5	-0.4	2.6	2.9
G31D Y32K	E50S	2	4.6	1.4	3
Cow	E50S	---	22.8	2.6	3.1
VH4-34	E50S	3.3	3.3	3.3	1.8
G31D Y32K	Cow	-0.1	2.3	2	2.6

Cow	Cow	0.7	4.9	4.6	3.1
VH4-34	Cow	0.6	0.6	2.5	3.9

Example 9: Modification of human light chain and pairing with heavy chain comprising an ultralong CDR3

[00622] Human germline light chains including, light chains that may be paired with a heavy chain comprising an ultralong CDR3, may be modified by any method known in the art. Such modifications may include the substitution of certain amino acid residues in the human light chain to those residues at corresponding positions in a bovine light chain sequence. The modified light chains may improve the yield of the antibody comprising the ultralong CDR3 and/or increase its binding specificity.

[00623] In an exemplary method, variants of V1-51 were made by overlap PCR and subcloned into pFuse vector for expression as described in Example 4. The engineered variants of V1-51 had: i) substitutions I29V and N32G introduced in CDR1 (SEQ ID NO: 440), ii) residues DNN (amino acids 51-53) in CDR2 changed to GDT (SEQ ID NO: 441), iii) residues DNNKRP (SEQ ID NO: 471) in and near CDR2 changed to GDTSRA (SEQ ID NO: 472), or iv) the 14 residues at the N-terminus were made identical to the first 14 residues of BLV1H12 light chain with point mutations S2A, T5N, P8S, A12G, A13S, and P14L (SEQ ID NO: 443) or v) combining the changes set forth in ii) and iv) (SEQ ID NO: 444). These V1-51 variants were paired with heavy chains encoding VH4-34+CDR3-IL8 (SEQ ID NO: 430), VH4-34+CDR3-IL8_CDR1 Cow (SEQ ID NO: 432), VH4-34+CDR3-IL8_CDR2 Cow (SEQ ID NO: 433), or VH4-34+CDR3-IL8_CDR1 Cow_CDR2 Cow, (SEQ ID NO: 436). Transfections and ELISA were carried out as described above, and flow cytometry was used to evaluate CXCR1 binding specificity as previously described, with the modification that cells were resuspended in 2% BSA in PBS plus 2ug/ml Heparin sulfate, and IgGs were normalized to 10nM during incubation on cells. Only certain combinations of heavy and light chains supported detectable expression (Table 15). Those combinations that expressed heavy and light chains were tested for CXCR1 binding specificity (Table 16).

Table 15. Expression yield (nM IgG) for VH4-34+CDR3-IL8 variants and V1-51 variants

		VH4-34 CDR3-IL8	VH4-34 CDR3-IL8 CDR1-Cow	VH4-34 CDR3-IL8 CDR2-Cow	VH4-34 CDR3-IL8 CDR1-Cow &CDR2-Cow
	Hc				

	CDR1:	Human	Cow	Human	Cow
	CDR2:	Human	Human	Cow	Cow
V1-51		1	5	0	13
V1-51 CDR1 I29V, N32G		-1	5	0	3
V1-51 CDR2 DNN to GDT		-1	23	-1	5
V1-51 CDR2 DNNKRP (SEQ ID NO: 471) to GDTsRA (SEQ ID NO: 472)		-1	4	1	23
V1-51 S2A,T5N,P8S,A12G,A13S,P14L		0	38	-1	22
V1-51 S2A,T5N,P8S,A12G,A13S,P14L & DNN to GDT		0	37	0	1

Table 16. CXCR1 specificity for VH4-34+CDR3-IL8 variants and V1-51 variants

		VH4-34 CDR3-IL8	VH4-34 CDR3-IL8 CDR1-Cow	VH4-34 CDR3-IL8 CDR2-Cow	VH4-34 CDR3-IL8 CDR1-Cow &CDR2-Cow
	Hc				
	CDR1:	Human	Cow	Human	Cow
	CDR2:	Human	Human	Cow	Cow
V1-51		nt	7.2	nt	11.1
V1-51 CDR1 I29V, N32G		nt	8.9	nt	11.8
V1-51 CDR2 DNN to GDT		nt	5.6	nt	16.7
V1-51 CDR2 DNNKRP (SEQ ID NO: 471) to GDTsRA (SEQ ID NO: 472)		nt	8.0	nt	6.9
V1-51 S2A,T5N,P8S,A12G,A13S,P14L		nt	4.7	nt	12.2
V1-51 S2A,T5N,P8S,A12G,A13S,P14L & DNN to GDT		nt	6.8	nt	

Example 10: Preparation of Antibody Libraries with Ultralong CDR3 Sequences

- 5 [00624] Libraries may be generated that comprise an ultralong CDR3 including, for example libraries that comprise at least a portion of an ultralong CDR3, within an antibody framework (e.g., a heavy chain framework). Such libraries may comprise a diversity of ultralong CDR3 sequences, a diversity of one or more residues that are positioned between one or more cysteine residues in the cysteine domain of the ultralong CDR3, or a
- 10 diversity of non-bovine peptides that may be inserted (e.g., replace a portion of) in the ultralong CDR3. The antibody framework may be derived from a bovine sequence such as VH-UL, a human germline sequence, or a modified human germline sequence such as

described in Example 7. Heavy chains with diverse ultralong CDR3 may be paired with light chains of bovine, human, or modified composition (see, e.g., Example 9) for expression of a library of antibody or antibody fragments comprising a diverse ultralong CDR3.

5 [00625] In an exemplary method, a library is generated in which the library comprises a diversity of ultralong CDR3 sequences (e.g., the diversity in the library resides in that the library contains a plurality of diverse ultralong CDR3 sequences). For example, a plurality of diverse ultralong CDR3 sequences is obtained from cDNA extracted from the spleen and/or lymph nodes from either immunized or non-immunized cows. Alternatively, a
10 plurality of diverse ultralong CDR3 sequences is captured as information from cDNA by sequencing technology, such as described in Example 2. cDNA-derived cow sequences are then amplified from the isolated cDNA or synthesized from sequencing of cDNA, and diverse CDR3 sequences are inserted into an antibody framework for expression of IgGs with ultralong CDR3 sequences, producing a library of IgGs with CDR3s derived from
15 cows. The IgG library can exist in any format, including as a spatially addressed array see, e.g., WO 11/056997; and Mao *et al.* (2010) *Nat Biotech* 28:1195-1202).

[00626] In another exemplary method, a library is generated in which members of the library comprise an ultralong CDR3 with a certain cysteine domain, or multiple cysteine domains, wherein the members are diverse in one or more residues that are positioned
20 between one or more cysteine residues in the cysteine domain. For example, in the sequence CX₁₀CX₅CX₅CXCX₇C (SEQ ID NO: 41), a library is generated whereby any X residue is changed, or combination of X residues are changed, to produce diversity within CDR3. Such diversity is introduced by degenerate nucleotide synthesis, error prone PCR, gene synthesis, or any other method known in the art to make changes in nucleotide
25 sequences. These ultralong CDR3 loops are then integrated with sequence encoding the antibody framework. Heavy chain sequences are then paired with human or modified human light chain sequences for co-transfection and expression to comprise an antibody library engineered from bovine CDR3s.

[00627] In another exemplary method, a library is generated in which members of
30 the library comprise an ultralong CDR3 with diversity in a non-bovine sequence inserted into (e.g., replace a portion of) the ultralong CDR3. Ultralong CDR3s tolerate large non-bovine sequence insertions (e.g., cytokines, peptide hormones, signaling domains, and constituent proteins of arthropod toxins or reptile venoms) including, for example, cysteine rich insertions, within an antibody framework, and support the independent function of the
35 peptide encoded by the non-bovine sequence, as demonstrated in Example 4 and Example 6. Sequences encoding non-bovine sequences are amplified by PCR from cDNA

or synthesized, and incorporated with or without linker sequences, into an antibody sequence such that they are expressed within, in place of, or replace at least a portion of, an ultralong CDR3. Exemplary methods to insert a non-bovine sequence into a vector for expression are set forth in Example 3 above. Heavy chain sequences are then paired with human or modified human light chain for co-transfection and expression. For example, a non-bovine sequence is inserted into HC pFUSE, encoding VH4-34*02 with CDR1 and CDR3 modifications from BLV1H12. Briefly, the portion of the HC pFUSE vector comprising a BsaI cassette (see, Example 3; bolded in the sequence below) which is flanked on both sides by a portion of an ultralong CDR3 (underlined sequence below) is replaced by a non-bovine sequence.

caggtgcagctacagcagtggggcgcaggactgtgaagccttcggagacgctgtccctcacctgcac
agcaagcgggttttactgagcgacaaggcagtgggatggattcgccagccccaggggaaggggctg
gagtggattggggaaatcaatcatagtgaagcaccaactacaacccgtccctcaagagtcgagtcac
catatcagtagacacgtccaagaaccagttctccctgaagctgagctctgtgaccgccgaggacacggc
tgtgtattactgtacctctgtgcaccaggaaactaagaaataccagagcgagacctactatggttcggg
tctctcttatacctacaattatgaatggcatgtggatgtctggggacagggcctgctggtgacagtctctagt
gctagc
(SEQ ID NO: 472)

20

[00628] After replacement of the BsaI cassette with the non-bovine sequence the non-bovine sequence (referred to as insert in bold text in the sequence below) is positioned such that it is flanked on both sides by a portion of the ultralong CDR3.

qvqlqqwgagllkpsetlsltctasgfslsdkavgwirpppgkglewigeinhsgstnynpslksrvtisvdt
sknqfslklssvtaadtavyyctsvhgetkkyqs- **insert** -sytnyewhvdvwgqgllvtvssas
(SEQ ID NO: 473)

[00629] Heavy chain sequences are then paired with human or modified human light chain for co-transfection and expression.

[00630] Embodiments of the present subject matter described above, some of which are set forth below, may be beneficial alone or in combination, with one or more other embodiments. Such embodiments include:

[00631] Embodiment 1. A humanized antibody or binding fragment thereof comprising an ultralong CDR3. In some embodiments, the humanized antibody or binding fragment thereof comprises one or more human variable region framework sequences.

[00632] Embodiment 2. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 is 35 amino acids in length or longer, 40 amino acids in length or longer, 45 amino acids in length or longer, 50 amino acids in length or longer, 55 amino acids in length or longer, or 60 amino acids in length or longer.

5 [00633] Embodiment 3. The humanized antibody or binding fragment thereof of Embodiment 2, wherein the ultralong CDR3 is 35 amino acids in length or longer.

[00634] Embodiment 4. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises 3 or more cysteine residues, 4 or more cysteine residues, 5 or more cysteine residues, 6 or more cysteine residues, 7 or more cysteine residues, 8 or more cysteine residues, 9 or more cysteine residues, 10 or more cysteine residues, 11 or more cysteine residues, or 12 or more cysteine residues.

[00635] Embodiment 5. The humanized antibody or binding fragment thereof of Embodiment 4, wherein the ultralong CDR3 comprises 3 or more cysteine residues.

15 [00636] Embodiment 6. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the antibodies or binding fragments thereof comprise a cysteine motif.

[00637] Embodiment 7. The humanized antibody or binding fragment thereof of Embodiment 6, wherein the cysteine motif is selected from the group consisting of:

20 CX₁₀CX₅CX₅CXCX₇C (SEQ ID NO: 41), CX₁₀CX₆CX₅CXCX₁₅C (SEQ ID NO: 42), CX₁₁CXCX₅C (SEQ ID NO: 43), CX₁₁CX₅CX₅CXCX₇C (SEQ ID NO: 44), CX₁₀CX₆CX₅CXCX₁₃C (SEQ ID NO: 45), CX₁₀CX₅CXCX₄CX₈C (SEQ ID NO: 46), CX₁₀CX₆CX₆CXCX₇C (SEQ ID NO: 47), CX₁₀CX₄CX₇CXCX₈C (SEQ ID NO: 48), CX₁₀CX₄CX₇CXCX₇C (SEQ ID NO: 49), CX₁₃CX₈CX₈C (SEQ ID NO: 50), CX₁₀CX₆CX₅CXCX₇C (SEQ ID NO: 51), CX₁₀CX₅CX₅C (SEQ ID NO: 52),

25 CX₁₀CX₅CX₆CXCX₇C (SEQ ID NO: 53), CX₁₀CX₆CX₅CX₇CX₉C (SEQ ID NO: 54), CX₉CX₇CX₅CXCX₇C (SEQ ID NO: 55), CX₁₀CX₆CX₅CXCX₉C (SEQ ID NO: 56), CX₁₀CXCX₄CX₅CX₁₁C (SEQ ID NO: 57), CX₇CX₃CX₆CX₅CXCX₅CX₁₀C (SEQ ID NO: 58), CX₁₀CXCX₄CX₅CXCX₂CX₃C (SEQ ID NO: 59), CX₁₆CX₅CXC (SEQ ID NO: 60), CX₆CX₄CXCX₄CX₅C (SEQ ID NO: 61), CX₁₁CX₄CX₅CX₆CX₃C (SEQ ID NO: 62),

30 CX₈CX₂CX₆CX₅C (SEQ ID NO: 63), CX₁₀CX₅CX₅CXCX₁₀C (SEQ ID NO: 64), CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65), CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66), CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67), CX₁₅CX₅CXC (SEQ ID NO: 68), CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69), CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70), CX₇CX₃CXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72),

35 CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74), CX₁₀CXCX₆C (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C

(SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCX₁₄C (SEQ ID NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83), CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85),
 5 CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87), CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89), CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91), CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93), CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ
 10 ID NO: 95).

[00638] Embodiment 8. The humanized antibody or binding fragment thereof of Embodiment 6, wherein the cysteine motif is selected from the group consisting of: CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCCX₄CX₆CXC (SEQ ID NO: 98),
 15 CX₉CX₃CXCX₂CXCCX₆CX₄C (SEQ ID NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102), CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109), CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113),
 25 CX₉CCX₄CX₂CXCCXCX₄CX₃C (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116), CX₆CCXCX₅CX₄CCXCX₅CX₂C (SEQ ID NO: 117), CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118), CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO: 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135),
 35

CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143),
 5 CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152).

[00639] Embodiment 9. The humanized antibody or binding fragment thereof of
 10 Embodiment 1, wherein the ultralong CDR3 comprises 2 to 6 disulfide bonds.

[00640] Embodiment 10. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises SEQ ID NO: 40 or a derivative thereof.

[00641] Embodiment 11. The humanized antibody or binding fragment thereof of
 15 Embodiment 1, wherein the ultralong CDR3 comprises amino acid residues 3-6 of any of one SEQ ID NO: 1-4.

[00642] Embodiment 12. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises a non-human DH or a derivative thereof.

20 [00643] Embodiment 13. The humanized antibody or binding fragment thereof of Embodiment 12, wherein the non-human DH is SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, or SEQ ID NO: 12.

[00644] Embodiment 14. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises a JH sequence or a derivative
 25 thereof.

[00645] Embodiment 15. The humanized antibody or binding fragment thereof of Embodiment 14, wherein the JH sequence is SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, or SEQ ID NO: 17.

[00646] Embodiment 16. The humanized antibody or binding fragment thereof of
 30 Embodiment 1, wherein the ultralong CDR3 comprises: a non-human VH sequence or a derivative thereof; a non-human DH sequence or a derivative thereof; and/or a JH sequence or derivative thereof.

[00647] Embodiment 17. The humanized antibody or binding fragment thereof of Embodiment 16, wherein the ultralong CDR3 comprises an additional amino acid sequence
 35 comprising two to six amino acid residues or more positioned between the VH sequence and the DH sequence.

[00648] Embodiment 18. The humanized antibody or binding fragment thereof of Embodiment 17, wherein the additional amino acid sequence is selected from the group consisting of: IR, IF, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20 or SEQ ID NO: 21.

[00649] Embodiment 19. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises a sequence derived from or based on SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, or SEQ ID NO: 28.

[00650] Embodiment 20. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises a non-bovine sequence or a non-antibody sequence.

[00651] Embodiment 21. The humanized antibody or binding fragment thereof of Embodiment 20, wherein the non-antibody sequence is a synthetic sequence.

[00652] Embodiment 22. The humanized antibody or binding fragment thereof of Embodiment 20, wherein the non-antibody sequence is a cytokine sequence, a lymphokine sequence, a chemokine sequence, a growth factor sequence, a hormone sequence, or a toxin sequence.

[00653] Embodiment 23. The humanized antibody or binding fragment thereof of Embodiment 20, wherein the non-antibody sequence is an IL-8 sequence, an IL-21 sequence, an SDF-1 (alpha) sequence, a somatostatin sequence, a chlorotoxin sequence, a Pro-TxII sequence, or a ziconotide sequence.

[00654] Embodiment 24. The humanized antibody or binding fragment thereof of Embodiment 20, wherein the non-antibody sequence is any one of SEQ ID NOS: 475-481.

[00655] Embodiment 25. The humanized antibody or binding fragment thereof of Embodiment 20, wherein the non-antibody sequence replaces at least a portion of the ultralong CDR3.

[00656] Embodiment 26. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q).

[00657] Embodiment 27. The humanized antibody or binding fragment thereof of Embodiment 26, wherein the $X^1X^2X^3X^4X^5$ motif is TTVHQ (SEQ ID NO: 153), TSVHQ (SEQ ID NO: 154), SSVTQ (SEQ ID NO: 155), STVHQ (SEQ ID NO: 156), ATVRQ (SEQ ID NO:

157), TTVYQ (SEQ ID NO: 158), SPVHQ (SEQ ID NO: 159), ATVYQ (SEQ ID NO: 160), TAVYQ (SEQ ID NO: 161), TNVHQ (SEQ ID NO: 162), ATVHQ (SEQ ID NO: 163), STVYQ (SEQ ID NO: 164), TIVHQ (SEQ ID NO: 165), AIVYQ (SEQ ID NO: 166), TTVFQ (SEQ ID NO: 167), AAVFQ (SEQ ID NO: 168), GTVHQ (SEQ ID NO: 169), ASVHQ (SEQ ID NO: 170), TAVFQ (SEQ ID NO: 171), ATVFQ (SEQ ID NO: 172), AAAHQ (SEQ ID NO: 173), VVVYQ (SEQ ID NO: 174), GTVFQ (SEQ ID NO: 175), TAVHQ (SEQ ID NO: 176), ITVHQ (SEQ ID NO: 177), ITAHQ (SEQ ID NO: 178), VTVHQ (SEQ ID NO: 179); AAVHQ (SEQ ID NO: 180), GTVYQ (SEQ ID NO: 181), TTVLQ (SEQ ID NO: 182), TTTHQ (SEQ ID NO: 183), or TTDYQ (SEQ ID NO: 184).

10 [00658] Embodiment 28. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises a $CX^1X^2X^3X^4X^5$ motif.

[00659] Embodiment 29. The humanized antibody or binding fragment thereof of Embodiment 28, wherein the $CX^1X^2X^3X^4X^5$ motif is CTTVHQ (SEQ ID NO: 185), CTSVHQ (SEQ ID NO: 186), CSSVTQ (SEQ ID NO: 187), CSTVHQ (SEQ ID NO: 188), CATVRQ (SEQ ID NO: 189), CTTVYQ (SEQ ID NO: 190), CSPVHQ (SEQ ID NO: 191), CATVYQ (SEQ ID NO: 192), CTAVYQ (SEQ ID NO: 193), CTNVHQ (SEQ ID NO: 194), CATVHQ (SEQ ID NO: 195), CSTVYQ (SEQ ID NO: 196), CTIVHQ (SEQ ID NO: 197), CAIVYQ (SEQ ID NO: 198), CTTVFQ (SEQ ID NO: 199), CAAVFQ (SEQ ID NO: 200), CGTVHQ (SEQ ID NO: 201), CASVHQ (SEQ ID NO: 202), CTAVFQ (SEQ ID NO: 203), CATVFQ (SEQ ID NO: 204), CAAAHQ (SEQ ID NO: 205), CVVVYQ (SEQ ID NO: 206), CGTVFQ (SEQ ID NO: 207), CTAVHQ (SEQ ID NO: 208), CITVHQ (SEQ ID NO: 209), CITAHQ (SEQ ID NO: 210), CVTVHQ (SEQ ID NO: 211); CAAVHQ (SEQ ID NO: 212), CGTVYQ (SEQ ID NO: 213), CTTVLQ (SEQ ID NO: 214), CTTTHQ (SEQ ID NO: 215), or CTTDYQ (SEQ ID NO: 216).

25 [00660] Embodiment 30. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises a $(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[00661] Embodiment 31. The humanized antibody or binding fragment thereof of Embodiment 30, wherein the $(X^aX^b)_z$ motif is CYTYNYEF (SEQ ID NO: 217), HYTYTYDF (SEQ ID NO: 218), HYTYTYEW (SEQ ID NO: 219), KHRITYEW (SEQ ID NO: 220), NYIYKYSF (SEQ ID NO: 221), PYIYTYQF (SEQ ID NO: 222), SFTYTYEW (SEQ ID NO: 223), SYIYIYQW (SEQ ID NO: 224), SYNYTYSW (SEQ ID NO: 225), SYSYSYEF (SEQ ID NO: 226), SYTYNYDF (SEQ ID NO: 227), SYTYNYEW (SEQ ID NO: 228), SYTYNYQF (SEQ ID NO: 229), SYVWTHNF (SEQ ID NO: 230), TYKYVYEW (SEQ ID NO: 231), TYTYTYEF (SEQ ID NO: 232), TYTYTYEW (SEQ ID NO: 233), VFTYTYEF (SEQ ID NO: 234), or VFTYTYEW (SEQ ID NO: 235).

234), AYTIEW (SEQ ID NO: 235), DYIYTY (SEQ ID NO: 236), IHSYEF (SEQ ID NO: 237), SFTYEF (SEQ ID NO: 238), SHSYEF (SEQ ID NO: 239), THTYEF (SEQ ID NO: 240), TWTYEF (SEQ ID NO: 241), TYNIEW (SEQ ID NO: 242), TYSYEF (SEQ ID NO: 243), TYSYEH (SEQ ID NO: 244), TYTYDF (SEQ ID NO: 245), TYTYEF (SEQ ID NO: 246), TYTIEW (SEQ ID NO: 247), AYES (SEQ ID NO: 248), AYSF (SEQ ID NO: 249), AYSY (SEQ ID NO: 250), CYSF (SEQ ID NO: 251), DITY (SEQ ID NO: 252), KYEH (SEQ ID NO: 253), KIEW (SEQ ID NO: 254), MYEF (SEQ ID NO: 255), NWIY (SEQ ID NO: 256), NYDY (SEQ ID NO: 257), NYQW (SEQ ID NO: 258), NYSF (SEQ ID NO: 259), PYEW (SEQ ID NO: 260), RYNW (SEQ ID NO: 261), RITY (SEQ ID NO: 262), SYEF (SEQ ID NO: 263), SYEH (SEQ ID NO: 264), SYEW (SEQ ID NO: 265), SYKW (SEQ ID NO: 266), SYTY (SEQ ID NO: 267), TYDF (SEQ ID NO: 268), TYEF (SEQ ID NO: 269), TIEW (SEQ ID NO: 270), TYQW (SEQ ID NO: 271), TITY (SEQ ID NO: 272), or VIEW (SEQ ID NO: 273).

[00662] Embodiment 32. The humanized antibody or binding fragment thereof of Embodiment 30, wherein the $(X^aX^b)_z$ motif is YXYXYX.

[00663] Embodiment 33. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5X_n$ motif, wherein X_1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X_2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X_3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X_4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), wherein X_5 is glutamine (Q), and wherein n is 27 – 54.

[00664] Embodiment 34. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises $X_n(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

[00665] Embodiment 35. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5X_n(X^aX^b)_z$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q), wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

[00666] Embodiment 36. The humanized antibody or binding fragment thereof of Embodiment 35, wherein the $X^1X^2X^3X^4X^5$ motif is TTVHQ (SEQ ID NO: 153) or TSVHQ (SEQ ID NO: 154), and wherein the $(X^aX^b)_z$ motif is YXYXYX.

[00667] Embodiment 37. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises: a $CX^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q), a cysteine motif selected from the group consisting of: $CX_{10}CX_5CX_5CXCX_7C$ (SEQ ID NO: 41), $CX_{10}CX_6CX_5CXCX_{15}C$ (SEQ ID NO: 42), $CX_{11}CXCX_5C$ (SEQ ID NO: 43), $CX_{11}CX_5CX_5CXCX_7C$ (SEQ ID NO: 44), $CX_{10}CX_6CX_5CXCX_{13}C$ (SEQ ID NO: 45), $CX_{10}CX_5CXCX_4CX_8C$ (SEQ ID NO: 46), $CX_{10}CX_6CX_6CXCX_7C$ (SEQ ID NO: 47), $CX_{10}CX_4CX_7CXCX_8C$ (SEQ ID NO: 48), $CX_{10}CX_4CX_7CXCX_7C$ (SEQ ID NO: 49), $CX_{13}CX_8CX_8C$ (SEQ ID NO: 50), $CX_{10}CX_6CX_5CXCX_7C$ (SEQ ID NO: 51), $CX_{10}CX_5CX_5C$ (SEQ ID NO: 52), $CX_{10}CX_5CX_6CXCX_7C$ (SEQ ID NO: 53), $CX_{10}CX_6CX_5CX_7CX_9C$ (SEQ ID NO: 54), $CX_9CX_7CX_5CXCX_7C$ (SEQ ID NO: 55), $CX_{10}CX_6CX_5CXCX_9C$ (SEQ ID NO: 56), $CX_{10}CXCX_4CX_5CX_{11}C$ (SEQ ID NO: 57), $CX_7CX_3CX_6CX_5CXCX_5CX_{10}C$ (SEQ ID NO: 58), $CX_{10}CXCX_4CX_5CXCX_2CX_3C$ (SEQ ID NO: 59), $CX_{16}CX_5CXC$ (SEQ ID NO: 60), $CX_6CX_4CXCX_4CX_5C$ (SEQ ID NO: 61), $CX_{11}CX_4CX_5CX_6CX_3C$ (SEQ ID NO: 62), $CX_8CX_2CX_6CX_5C$ (SEQ ID NO: 63), $CX_{10}CX_5CX_5CXCX_{10}C$ (SEQ ID NO: 64), $CX_{10}CXCX_6CX_4CXC$ (SEQ ID NO: 65), $CX_{10}CX_5CX_5CXCX_2C$ (SEQ ID NO: 66), $CX_{14}CX_2CX_3CXCXC$ (SEQ ID NO: 67), $CX_{15}CX_5CXC$ (SEQ ID NO: 68), $CX_4CX_6CX_9CX_2CX_{11}C$ (SEQ ID NO: 69), $CX_6CX_4CX_5CX_5CX_{12}C$ (SEQ ID NO: 70), $CX_7CX_3CXCXCX_4CX_5CX_9C$ (SEQ ID NO: 71), $CX_{10}CX_6CX_5C$ (SEQ ID NO: 72), $CX_7CX_3CX_5CX_5CX_9C$ (SEQ ID NO: 73), $CX_7CX_5CXCX_2C$ (SEQ ID NO: 74), $CX_{10}CXCX_6C$ (SEQ ID NO: 75), $CX_{10}CX_3CX_3CX_5CX_7CXCX_6C$ (SEQ ID NO: 76), $CX_{10}CX_4CX_5CX_{12}CX_2C$ (SEQ ID NO: 77), $CX_{12}CX_4CX_5CXCXCX_9CX_3C$ (SEQ ID NO: 78), $CX_{12}CX_4CX_5CX_{12}CX_2C$ (SEQ ID NO: 79), $CX_{10}CX_6CX_5CXCX_{11}C$ (SEQ ID NO: 80), $CX_{16}CX_5CXCXCX_{14}C$ (SEQ ID NO: 81), $CX_{10}CX_5CXCX_8CX_6C$ (SEQ ID NO: 82), $CX_{12}CX_4CX_5CX_8CX_2C$ (SEQ ID NO: 83), $CX_{12}CX_5CX_5CXCX_8C$ (SEQ ID NO: 84), $CX_{10}CX_6CX_5CXCX_4CXCX_9C$ (SEQ ID NO: 85), $CX_{11}CX_4CX_5CX_8CX_2C$ (SEQ ID NO: 86), $CX_{10}CX_6CX_5CX_8CX_2C$ (SEQ ID NO: 87), $CX_{10}CX_6CX_5CXCX_8C$ (SEQ ID NO: 88), $CX_{10}CX_6CX_5CXCX_3CX_8CX_2C$ (SEQ ID NO: 89), $CX_{10}CX_6CX_5CX_3CX_8C$ (SEQ ID NO: 90), $CX_{10}CX_6CX_5CXCX_2CX_6CX_5C$ (SEQ ID NO: 91), $CX_7CX_6CX_3CX_3CX_9C$ (SEQ ID NO: 92), $CX_9CX_8CX_5CX_6CX_5C$ (SEQ ID NO: 93), $CX_{10}CX_2CX_2CX_7CXCX_{11}CX_5C$ (SEQ ID NO: 94), and $CX_{10}CX_6CX_5CXCX_2CX_8CX_4C$ (SEQ

ID NO: 95); and a $(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[00668] Embodiment 38. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 comprises: a $CX^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q); a cysteine motif selected from the group consisting of: wherein the cysteine motif is selected from the group consisting of: $CCX_3CX_3CX_2CCX_5CX_9CX_5CXC$ (SEQ ID NO: 96), $CX_6CX_2CX_5CX_4CCX_4CX_6CXC$ (SEQ ID NO: 97), $CX_7CX_5CX_4CCCX_4CX_6CXC$ (SEQ ID NO: 98), $CX_9CX_3CX_2CX_3CCX_6CX_4C$ (SEQ ID NO: 99), $CX_5CX_3CX_4CX_4CCX_{10}CX_2CC$ (SEQ ID NO: 100), $CX_5CX_1CX_3CCX_3CX_4CX_{10}C$ (SEQ ID NO: 101), $CX_9CCCX_3CX_4CCCX_5CX_6C$ (SEQ ID NO: 102), $CCX_8CX_5CX_4CX_3CX_4CCX_1C$ (SEQ ID NO: 103), $CCX_6CCX_5CCCX_4CX_4CX_{12}C$ (SEQ ID NO: 104), $CX_6CX_2CX_3CCCX_4CX_5CX_3CX_3C$ (SEQ ID NO: 105), $CX_3CX_5CX_6CX_4CCX_5CX_4CXC$ (SEQ ID NO: 106), $CX_4CX_4CCX_4CX_4CXCX_{11}CX_2CXC$ (SEQ ID NO: 107), $CX_5CX_2CCX_5CX_4CCX_3CCX_7C$ (SEQ ID NO: 108), $CX_5CX_5CX_3CX_2CX_4CCX_7CXC$ (SEQ ID NO: 109), $CX_3CX_7CX_3CX_4CCX_2CX_5CX_2C$ (SEQ ID NO: 110), $CX_9CX_3CX_4CCX_5CCCX_6C$ (SEQ ID NO: 111), $CX_9CX_3CX_2CX_3CCX_6CX_3CX_3C$ (SEQ ID NO: 112), $CX_8CCX_3CX_3CX_3CX_4C$ (SEQ ID NO: 113), $CX_9CCX_4CX_2CX_3CCX_4CX_3C$ (SEQ ID NO: 114), $CX_{10}CX_3CX_2CX_4CCX_5CXC$ (SEQ ID NO: 115), $CX_9CX_3CX_2CX_4CCX_5CXC$ (SEQ ID NO: 116), $CX_6CCX_5CX_4CCX_5CX_2C$ (SEQ ID NO: 117), $CX_6CCX_3CX_3CCX_3CX_4CC$ (SEQ ID NO: 118), $CX_6CCX_3CX_2CX_4CX_8C$ (SEQ ID NO: 119), $CX_4CX_2CCX_3CX_4CCX_2CX_3C$ (SEQ ID NO: 120), $CX_3CX_5CX_3CCCX_4CX_9C$ (SEQ ID NO: 121), $CCX_9CX_3CX_3CX_5C$ (SEQ ID NO: 122), $CX_9CX_2CX_3CX_4CCX_4CX_5C$ (SEQ ID NO: 123), $CX_9CX_7CX_4CCX_7CX_3C$ (SEQ ID NO: 124), $CX_9CX_3CCCX_{10}CX_2CX_3C$ (SEQ ID NO: 125), $CX_3CX_5CX_5CX_4CCX_{10}CX_6C$ (SEQ ID NO: 126), $CX_9CX_5CX_4CCX_5CX_4C$ (SEQ ID NO: 127), $CX_7CX_6CX_4CCCX_{10}C$ (SEQ ID NO: 128), $CX_8CX_2CX_4CCX_4CX_3CX_3C$ (SEQ ID NO: 129), $CX_7CX_5CX_4CCX_7CX_4C$ (SEQ ID NO: 130), $CX_{11}CX_3CX_4CCCX_8CX_2C$ (SEQ ID NO: 131), $CX_2CX_3CX_4CCX_4CX_5CX_{15}C$ (SEQ ID NO: 132), $CX_9CX_5CX_4CCX_7C$ (SEQ ID NO: 133), $CX_9CX_7CX_3CX_2CX_6C$ (SEQ ID NO: 134), $CX_9CX_5CX_4CCX_{14}C$ (SEQ ID NO: 135), $CX_9CX_5CX_4CCX_8C$ (SEQ ID NO: 136), $CX_9CX_6CX_4CCXC$ (SEQ ID NO: 137), $CX_5CCX_7CX_4CX_{12}$ (SEQ ID NO: 138),

CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152); and a (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

10 [00669] Embodiment 39. The humanized antibody or binding fragment thereof of Embodiment 20, wherein the ultralong CDR3 comprises an additional sequence that is a linker.

[00670] Embodiment 40. The humanized antibody or binding fragment thereof of Embodiment 39, wherein the linker is linked to a C-terminus, a N-terminus, or both C-terminus and N-terminus of the non-antibody sequence.

[00671] Embodiment 41. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the ultralong CDR3 is a ruminant CDR3.

[00672] Embodiment 42. The humanized antibody or binding fragment thereof of Embodiment 41, wherein the ruminant is a cow.

20 [00673] Embodiment 43. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the humanized antibody or binding fragment thereof comprises a human heavy chain germline sequence that is a VH4 family member germline sequence.

[00674] Embodiment 44. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the humanized antibody or binding fragment thereof comprises a VH4-34 germline sequence.

[00675] Embodiment 45. The humanized antibody or binding fragment thereof of Embodiment 44, wherein the humanized antibody or binding fragment thereof comprises a human heavy chain germline sequence that is derived from the VH4-34 germline sequence.

30 [00676] Embodiment 46. The humanized antibody or binding fragment thereof of Embodiment 45, wherein the VH4-34 germline sequence comprises a CDR1 comprising a Gly31Asp and Tyr32Lys substitution based on Kabat numbering.

[00677] Embodiment 47. The humanized antibody or binding fragment thereof of Embodiment 45, wherein the VH4-34 germline sequence comprises a CDR2 comprising a Glu50Ser substitution based on Kabat numbering.

[00678] Embodiment 48. The humanized antibody or binding fragment thereof of Embodiment 45, wherein the VH4-34 germline sequence comprises a CDR1 comprising Ala23Thr, Val24Ala, Tyr25Ser, Gly27Phe, Phe29Leu, Gly31Asp, Tyr32Lys, Tyr33Ala, Trp34Val, and Ser35Gly substitutions, and a CDR2 comprising Ile48Leu, Glu50Ser, Asn52Asp, His53Thr, Ser54Gly, Ser56Asn, and Asn58Gly substitutions, based on Kabat numbering.

[00679] Embodiment 49. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469.

[00680] Embodiment 50. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 470.

[00681] Embodiment 51. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469; and the amino acid sequence set forth in SEQ ID NO: 470.

[00682] Embodiment 52. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469; the amino acid sequence set forth in SEQ ID NO: 470; and a peptide sequence, wherein the peptide sequence is positioned between SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469, and SEQ ID NO: 470.

[00683] Embodiment 53. The humanized antibody or binding fragment thereof of Embodiment 52, wherein the peptide sequence is a bovine sequence, a non-bovine sequence, an antibody sequence, or a non-antibody sequence.

[00684] Embodiment 54. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the humanized antibody or binding fragment thereof comprises a light chain variable region sequence that is a lambda light chain variable region sequence.

[00685] Embodiment 55. The humanized antibody or binding fragment thereof of Embodiment 54, wherein the lambda light chain variable region sequence is a bovine lambda light chain variable region sequence.

[00686] Embodiment 56. The humanized antibody or binding fragment thereof of Embodiment 54, wherein the lambda light chain variable region sequence is a human lambda light chain variable region sequence.

[00687] Embodiment 57. The humanized antibody or binding fragment thereof of Embodiment 56, wherein the human lambda light chain variable region sequence is a VL1-51 germline sequence.

5 [00688] Embodiment 58. The humanized antibody or binding fragment thereof of Embodiment 56, wherein the human lambda light chain variable region sequence is derived from a VL1-51 germline sequence.

[00689] Embodiment 59. The humanized antibody or binding fragment thereof of Embodiment 58, wherein the VL1-51 germline sequence comprises a CDR1 comprising Ile29Val and Asn32Gly substitution based on Kabat numbering.

10 [00690] Embodiment 60. The humanized antibody or binding fragment thereof of Embodiment 58, wherein the VL1-51 germline sequence comprises a CDR2 comprising a substitution of DNN to GDT.

[00691] Embodiment 61. The humanized antibody or binding fragment thereof of Embodiment 58, wherein the VL1-51 germline sequence comprises a CDR2 comprising
15 a substitution of DNNKRP (SEQ ID NO: 471) to GDTSRA (SEQ ID NO: 472).

[00692] Embodiment 62. The humanized antibody or binding fragment thereof of Embodiment 58, wherein the VL1-51 germline sequence comprises a S2A, T5N, P8S, A12G, A13S, and P14L substitution based on Kabat numbering.

20 [00693] Embodiment 63. The humanized antibody or binding fragment thereof of Embodiment 58, wherein the VL1-51 germline sequence comprises a S2A, T5N, P8S, A12G, A13S, and P14L substitution based on Kabat numbering, and a CDR2 comprising a substitution of DNN to GDT.

[00694] Embodiment 64. The humanized antibody or binding fragment thereof of Embodiment 1, wherein the humanized antibody or binding fragment thereof comprising
25 the amino acid sequence set forth in SEQ ID NO: 440, 441, 442, 443, or 444.

[00695] Embodiment 65. A polynucleotide encoding the humanized antibody or binding fragment thereof of any one of Embodiments 1-64.

[00696] Embodiment 66. A polynucleotide encoding a humanized antibody or binding fragment thereof that comprises an ultralong CDR3.

30 [00697] Embodiment 67. A polynucleotide encoding a humanized variable region that comprises an ultralong CDR3.

[00698] Embodiment 68. A vector comprising the polynucleotide of any one of Embodiments 65-67.

[00699] Embodiment 69. A host cell comprising the vector of Embodiment 68.

35 [00700] Embodiment 70. A nucleic acid library comprising a plurality of polynucleotides comprising sequences coding for humanized antibodies or binding

fragments thereof, wherein the antibodies or binding fragments thereof comprise an ultralong CDR3.

[00701] Embodiment 71. A library of humanized antibodies or binding fragments thereof, wherein the antibodies or binding fragments thereof comprise an ultralong CDR3. In some embodiments, the humanized antibodies or binding fragments thereof in the library comprise one or more human variable region framework sequences.

[00702] Embodiment 72. The library of Embodiment 71, wherein the ultralong CDR3 is 35 amino acids in length or longer, 40 amino acids in length or longer, 45 amino acids in length or longer, 50 amino acids in length or longer, 55 amino acids in length or longer, or 60 amino acids in length or longer.

[00703] Embodiment 73. The library of Embodiment 72, wherein the ultralong CDR3 is 35 amino acids in length or longer.

[00704] Embodiment 74. The library of Embodiment 71, wherein the ultralong CDR3 comprises 3 or more cysteine residues, 4 or more cysteine residues, 5 or more cysteine residues, 6 or more cysteine residues, 7 or more cysteine residues, 8 or more cysteine residues, 9 or more cysteine residues, 10 or more cysteine residues, 11 or more cysteine residues, or 12 or more cysteine residues.

[00705] Embodiment 75. The library of Embodiment 74, wherein the ultralong CDR3 comprises 3 or more cysteine residues.

[00706] Embodiment 76. The library of Embodiment 71, wherein the antibodies or binding fragments thereof comprise a cysteine motif.

[00707] Embodiment 77. The library of Embodiment 76, wherein the cysteine motif is selected from the group consisting of: CX₁₀CX₅CX₅CXCX₇C (SEQ ID NO: 41), CX₁₀CX₆CX₅CXCX₁₅C (SEQ ID NO: 42), CX₁₁CXCX₅C (SEQ ID NO: 43), CX₁₁CX₅CX₅CXCX₇C (SEQ ID NO: 44), CX₁₀CX₆CX₅CXCX₁₃C (SEQ ID NO: 45), CX₁₀CX₅CXCX₄CX₈C (SEQ ID NO: 46), CX₁₀CX₆CX₆CXCX₇C (SEQ ID NO: 47), CX₁₀CX₄CX₇CXCX₈C (SEQ ID NO: 48), CX₁₀CX₄CX₇CXCX₇C (SEQ ID NO: 49), CX₁₃CX₈CX₈C (SEQ ID NO: 50), CX₁₀CX₆CX₅CXCX₇C (SEQ ID NO: 51), CX₁₀CX₅CX₅C (SEQ ID NO: 52), CX₁₀CX₅CX₆CXCX₇C (SEQ ID NO: 53), CX₁₀CX₆CX₅CX₇CX₉C (SEQ ID NO: 54), CX₉CX₇CX₅CXCX₇C (SEQ ID NO: 55), CX₁₀CX₆CX₅CXCX₉C (SEQ ID NO: 56), CX₁₀CXCX₄CX₅CX₁₁C (SEQ ID NO: 57), CX₇CX₃CX₆CX₅CXCX₅CX₁₀C (SEQ ID NO: 58), CX₁₀CXCX₄CX₅CXCX₂CX₃C (SEQ ID NO: 59), CX₁₆CX₅CXC (SEQ ID NO: 60), CX₆CX₄CXCX₄CX₅C (SEQ ID NO: 61), CX₁₁CX₄CX₅CX₆CX₃C (SEQ ID NO: 62), CX₈CX₂CX₆CX₅C (SEQ ID NO: 63), CX₁₀CX₅CX₅CXCX₁₀C (SEQ ID NO: 64), CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65), CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66), CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67), CX₁₅CX₅CXC (SEQ ID NO: 68),

CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69), CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70), CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72), CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCXCX₂C (SEQ ID NO: 74), CX₁₀CXCXCX₆C (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCXCX₁₄C (SEQ ID NO: 81), CX₁₀CX₅CXCXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83), CX₁₂CX₅CX₅CXCXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCXCX₄CXCXCX₉C (SEQ ID NO: 85), CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87), CX₁₀CX₆CX₅CXCXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCXCX₃CX₈CX₂C (SEQ ID NO: 89), CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCXCX₂CX₆CX₅C (SEQ ID NO: 91), CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93), CX₁₀CX₂CX₂CX₇CXCXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCXCX₂CX₈CX₄C (SEQ ID NO: 95).

[00708] Embodiment 78. The library of Embodiment 76, wherein the cysteine motif is selected from the group consisting of: CCX₃CXCXCX₃CX₂CCXCXCX₅CX₉CX₅CXC (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCXCXCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCXCX₅CX₄CCCX₄CX₆CXC (SEQ ID NO: 98), CX₉CX₃CXCXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99), CX₅CX₃CXCXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100), CX₅CXCXCX₁CXCXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102), CCX₈CX₅CX₄CX₃CX₄CCXCXCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCXCXCX₅CX₄CXC (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCXCX₁₁CX₂CXC (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109), CX₃CX₇CX₃CX₄CCXCXCX₂CX₅CX₂C (SEQ ID NO: 110), CX₉CX₃CXCXCX₄CCX₅CCCX₆C (SEQ ID NO: 111), CX₉CX₃CXCXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCXCXCX₃CCX₃CXCXCX₃CX₄C (SEQ ID NO: 113), CX₉CCX₄CX₂CXCCXCXCX₄CX₃C (SEQ ID NO: 114), CX₁₀CXCXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115), CX₉CXCXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116), CX₆CCXCXCX₅CX₄CCXCXCX₅CX₂C (SEQ ID NO: 117), CX₆CCXCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118), CX₆CCXCXCX₃CXCXCX₂CXCXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO: 123), CX₉CX₇CX₄CCXCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128),

CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136),
 5 CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C
 10 (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152).

[00709] Embodiment 79. The library of Embodiment 71, wherein the ultralong CDR3 comprises 2 to 6 disulfide bonds.

[00710] Embodiment 80. The library of Embodiment 71, wherein the ultralong
 15 CDR3 comprises SEQ ID NO: 40 or a derivative thereof.

[00711] Embodiment 81. The library of Embodiment 71, wherein the ultralong CDR3 comprises amino acid residues 3-6 of any of one SEQ ID NO: 1-4.

[00712] Embodiment 82. The library of Embodiment 71, wherein the ultralong CDR3 comprises a non-human DH or a derivative thereof.

20 [00713] Embodiment 83. The library of Embodiment 82, wherein the non-human DH is SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, or SEQ ID NO: 12

[00714] Embodiment 84. The library of Embodiment 71, wherein the ultralong CDR3 comprises a JH sequence or a derivative thereof.

25 [00715] Embodiment 85. The library of Embodiment 84, wherein the JH sequence is SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, or SEQ ID NO: 17.

[00716] Embodiment 86. The library of Embodiment 71, wherein the ultralong CDR3 comprises: a non-human VH sequence or a derivative thereof; a non-human DH
 30 sequence or a derivative thereof; and/or a JH sequence or derivative thereof.

[00717] Embodiment 87. The library of Embodiment 86, wherein the ultralong CDR3 comprises an additional amino acid sequence comprising two to six amino acid residues or more positioned between the VH sequence and the DH sequence.

35 [00718] Embodiment 88. The library of Embodiment 87, wherein the additional amino acid sequence is selected from the group consisting of: IR, IF, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20 or SEQ ID NO: 21.

[00719] Embodiment 89. The library of Embodiment 88, wherein the ultralong CDR3 comprises a sequence derived from or based on SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, or SEQ ID NO: 28.

[00720] Embodiment 90. The library of Embodiment 71, wherein the ultralong CDR3 comprises a non-bovine sequence or a non-antibody sequence.

[00721] Embodiment 91. The library of Embodiment 90, wherein the non-antibody sequence is a synthetic sequence.

[00722] Embodiment 92. The library of Embodiment 90, wherein the non-antibody sequence is a cytokine sequence, a lymphokine sequence, a chemokine sequence, a growth factor sequence, a hormone sequence, or a toxin sequence.

[00723] Embodiment 93. The library of Embodiment 90, wherein the non-antibody sequence is an IL-8 sequence, an IL-21 sequence, an SDF-1 (alpha) sequence, a somatostatin sequence, a chlorotoxin sequence, a Pro-TxII sequence, or a ziconotide sequence.

[00724] Embodiment 94. The library of Embodiment 90, wherein the non-antibody sequence is any one of SEQ ID NOS: 475-481.

[00725] Embodiment 95. The library of Embodiment 71, wherein the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q).

[00726] Embodiment 96. The library of Embodiment 95, wherein the $X^1X^2X^3X^4X^5$ motif is TTVHQ (SEQ ID NO: 153), TSVHQ (SEQ ID NO: 154), SSVTQ (SEQ ID NO: 155), STVHQ (SEQ ID NO: 156), ATVRQ (SEQ ID NO: 157), TTVYQ (SEQ ID NO: 158), SPVHQ (SEQ ID NO: 159), ATVYQ (SEQ ID NO: 160), TAVYQ (SEQ ID NO: 161), TNVHQ (SEQ ID NO: 162), ATVHQ (SEQ ID NO: 163), STVYQ (SEQ ID NO: 164), TIVHQ (SEQ ID NO: 165), AIVYQ (SEQ ID NO: 166), TTVFQ (SEQ ID NO: 167), AAVFQ (SEQ ID NO: 168), GTVHQ (SEQ ID NO: 169), ASVHQ (SEQ ID NO: 170), TAVFQ (SEQ ID NO: 171), ATVFQ (SEQ ID NO: 172), AAAHQ (SEQ ID NO: 173), VVVYQ (SEQ ID NO: 174), GTVFQ (SEQ ID NO: 175), TAVHQ (SEQ ID NO: 176), ITVHQ (SEQ ID NO: 177), ITAHQ (SEQ ID NO: 178), VTVHQ (SEQ ID NO: 179), AAVHQ (SEQ ID NO: 180), GTVYQ (SEQ ID NO: 181), TTVLQ (SEQ ID NO: 182), TTTHQ (SEQ ID NO: 183), or TTDYQ (SEQ ID NO: 184).

[00727] Embodiment 97. The library of Embodiment 95, wherein the ultralong CDR3 comprises a $CX^1X^2X^3X^4X^5$ motif.

[00728] Embodiment 98. The library of Embodiment 97, wherein the CX¹X²X³X⁴X⁵ motif is CTTVHQ (SEQ ID NO: 185), CTSVHQ (SEQ ID NO: 186), CSSVTQ (SEQ ID NO: 187), CSTVHQ (SEQ ID NO: 188), CATVRQ (SEQ ID NO: 189), CTTVYQ (SEQ ID NO: 190), CSPVHQ (SEQ ID NO: 191), CATVYQ (SEQ ID NO: 192), CTAVYQ (SEQ ID NO: 193), CTNVHQ (SEQ ID NO: 194), CATVHQ (SEQ ID NO: 195), CSTVYQ (SEQ ID NO: 196), CTIVHQ (SEQ ID NO: 197), CAIVYQ (SEQ ID NO: 198), CTTVFQ (SEQ ID NO: 199), CAAVFQ (SEQ ID NO: 200), CGTVHQ (SEQ ID NO: 201), CASVHQ (SEQ ID NO: 202), CTAVFQ (SEQ ID NO: 203), CATVFQ (SEQ ID NO: 204), CAAAHQ (SEQ ID NO: 205), CVVVYQ (SEQ ID NO: 206), CGTVFQ (SEQ ID NO: 207), CTAVHQ (SEQ ID NO: 208), CITVHQ (SEQ ID NO: 209), CITAHQ (SEQ ID NO: 210), CVTVHQ (SEQ ID NO: 211); CAAVHQ (SEQ ID NO: 212), CGTVYQ (SEQ ID NO: 213), CTTVLQ (SEQ ID NO: 214), CTTTHQ (SEQ ID NO: 215), or CTTDYQ (SEQ ID NO: 216).

[00729] Embodiment 99. The humanized antibody or binding fragment thereof of Embodiment 71, wherein the ultralong CDR3 comprises a (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[00730] Embodiment 100. The humanized antibody or binding fragment thereof of Embodiment 99, wherein the (X^aX^b)_z motif is CYTYNYEF (SEQ ID NO: 217), HYTYTYDF (SEQ ID NO: 218), HYTYTYEW (SEQ ID NO: 219), KHRITYEW (SEQ ID NO: 220), NYIYKYSF (SEQ ID NO: 221), PYIYTYQF (SEQ ID NO: 222), SFTYTYEW (SEQ ID NO: 223), SYIYIYQW (SEQ ID NO: 224), SYNITYSW (SEQ ID NO: 225), SYSYSYEF (SEQ ID NO: 226), SYTYNYDF (SEQ ID NO: 227), SYTYNYEW (SEQ ID NO: 228), SYTYNYQF (SEQ ID NO: 229), SYVWTHNF (SEQ ID NO: 230), TYKYVYEW (SEQ ID NO: 231), TYTYTYEF (SEQ ID NO: 232), TYTYTYEW (SEQ ID NO: 233), VFTYTYEF (SEQ ID NO: 234), AYTIEW (SEQ ID NO: 235), DYIYTY (SEQ ID NO: 236), IHSYEF (SEQ ID NO: 237), SFTYEF (SEQ ID NO: 238), SHSYEF (SEQ ID NO: 239), THTYEF (SEQ ID NO: 240), TWTYEF (SEQ ID NO: 241), TYNIEW (SEQ ID NO: 242), TYSYEF (SEQ ID NO: 243), TYSYEH (SEQ ID NO: 244), TYTYDF (SEQ ID NO: 245), TYTYEF (SEQ ID NO: 246), TYTYEW (SEQ ID NO: 247), AYES (SEQ ID NO: 248), AYSF (SEQ ID NO: 249), AYSY (SEQ ID NO: 250), CYSF (SEQ ID NO: 251), DYTY (SEQ ID NO: 252), KYEH (SEQ ID NO: 253), KYEW (SEQ ID NO: 254), MYEF (SEQ ID NO: 255), NWIY (SEQ ID NO: 256), NYDY (SEQ ID NO: 257), NYQW (SEQ ID NO: 258), NYSF (SEQ ID NO: 259), PYEW (SEQ ID NO: 260), RYNW (SEQ ID NO: 261), RYTY (SEQ ID NO: 262), SYEF (SEQ ID NO: 263), SYEH (SEQ ID NO: 264), SYEW (SEQ ID NO: 265), SYKW (SEQ ID NO: 266), SYTY (SEQ ID NO: 267), TYDF (SEQ ID NO: 268), TYEF (SEQ ID NO: 269),

TYEW (SEQ ID NO: 270), TYQW (SEQ ID NO: 271), TYTY (SEQ ID NO: 272), or VYEW (SEQ ID NO: 273).

[00731] Embodiment 101. The humanized antibody or binding fragment thereof of Embodiment 99, wherein the $(X^aX^b)_z$ motif is YXYXYX.

5 [00732] Embodiment 102. The library of Embodiment 71, wherein the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5X_n$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R),
10 tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q), and wherein n is 27 – 54.

[00733] Embodiment 103. The library of Embodiment 71, wherein the ultralong CDR3 comprises $X_n(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F),
15 tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

[00734] Embodiment 104. The library of Embodiment 71, wherein the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5X_n(X^aX^b)_z$ motif, wherein X_1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X_2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X_3 is valine (V), alanine
20 (A), threonine (T), or aspartic acid (D), wherein X_4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), wherein X_5 is glutamine (Q), X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

25 [00735] Embodiment 105. The library of Embodiment 104, wherein the $X^1X^2X^3X^4X^5$ motif is TTVHQ (SEQ ID NO: 153) or TSVHQ (SEQ ID NO: 154), and wherein the $(X^aX^b)_z$ motif is YXYXYX.

[00736] Embodiment 106. The library of Embodiment 71, wherein the ultralong CDR3 comprises: a $CX^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine
30 (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q); a cysteine motif selected from the group consisting of: $CX_{10}CX_5CX_5CXCX_7C$ (SEQ ID NO: 41),
35 $CX_{10}CX_6CX_5CXCX_{15}C$ (SEQ ID NO: 42), $CX_{11}CXCX_5C$ (SEQ ID NO: 43), $CX_{11}CX_5CX_5CXCX_7C$ (SEQ ID NO: 44), $CX_{10}CX_6CX_5CXCX_{13}C$ (SEQ ID NO: 45),

CX₁₀CX₅CXCX₄CX₈C (SEQ ID NO: 46), CX₁₀CX₆CX₆CXCX₇C (SEQ ID NO: 47),
 CX₁₀CX₄CX₇CXCX₈C (SEQ ID NO: 48), CX₁₀CX₄CX₇CXCX₇C (SEQ ID NO: 49),
 CX₁₃CX₈CX₈C (SEQ ID NO: 50), CX₁₀CX₆CX₅CXCX₇C (SEQ ID NO: 51), CX₁₀CX₅CX₅C
 (SEQ ID NO: 52), CX₁₀CX₅CX₆CXCX₇C (SEQ ID NO: 53), CX₁₀CX₆CX₅CX₇CX₉C (SEQ ID
 5 NO: 54), CX₉CX₇CX₅CXCX₇C (SEQ ID NO: 55), CX₁₀CX₆CX₅CXCX₉C (SEQ ID NO: 56),
 CX₁₀CXCX₄CX₅CX₁₁C (SEQ ID NO: 57), CX₇CX₃CX₆CX₅CXCX₅CX₁₀C (SEQ ID NO: 58),
 CX₁₀CXCX₄CX₅CXCX₂CX₃C (SEQ ID NO: 59), CX₁₆CX₅CXC (SEQ ID NO: 60),
 CX₆CX₄CXCX₄CX₅C (SEQ ID NO: 61), CX₁₁CX₄CX₅CX₆CX₃C (SEQ ID NO: 62),
 CX₈CX₂CX₆CX₅C (SEQ ID NO: 63), CX₁₀CX₅CX₅CXCX₁₀C (SEQ ID NO: 64),
 10 CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65), CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66),
 CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67), CX₁₅CX₅CXC (SEQ ID NO: 68),
 CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69), CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70),
 CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72),
 CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74), CX₁₀CXCX₆C
 15 (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C
 (SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C
 (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCX₁₄C (SEQ ID
 NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83),
 CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85),
 20 CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87),
 CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89),
 CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91),
 CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93),
 CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ
 25 ID NO: 95); and a (X^aX^b)_z motif, X^a is any amino acid residue, X^b is an aromatic amino acid
 selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and
 histidine (H), and wherein z is 1 - 4.

[00737] Embodiment 107. The library of Embodiment 71, wherein the ultralong
 CDR3 comprises: a CX¹X²X³X⁴X⁵ motif, wherein X¹ is threonine (T), glycine (G), alanine
 30 (A), serine (S), or valine (V), wherein X² is serine (S), threonine (T), proline (P), isoleucine
 (I), alanine (A), valine (V), or asparagine (N), wherein X³ is valine (V), alanine (A),
 threonine (T), or aspartic acid (D), wherein X⁴ is histidine (H), threonine (T), arginine (R),
 tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X⁵ is glutamine (Q); a cysteine
 motif selected from the group consisting of: wherein the cysteine motif is selected from the
 35 group consisting of: CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96),
 CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCCX₄CX₆CXC (SEQ

ID NO: 98), CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99),
 CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C
 (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102),
 CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID
 5 NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105),
 CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC
 (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108),
 CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109), CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C
 (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111),
 10 CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ
 ID NO: 113), CX₉CCX₄CX₂CXCCXCX₄CX₃C (SEQ ID NO: 114),
 CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC
 (SEQ ID NO: 116), CX₆CCXCX₅CX₄CCXCX₅CX₂C (SEQ ID NO: 117),
 CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118), CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID
 15 NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C
 (SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C
 (SEQ ID NO: 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C
 (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126),
 CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128),
 20 CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO:
 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID
 NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134),
 CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136),
 CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138),
 25 CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140),
 CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142),
 CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144),
 CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C
 (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149),
 30 CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152); and a
 (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected
 from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine
 (H), and wherein z is 1 - 4.

[00738] Embodiment 108. The library of Embodiment 90, wherein the ultralong
 35 CDR3 comprises an additional sequence that is a linker.

[00739] Embodiment 109. The library of Embodiment 108, wherein the linker is linked to a C-terminus, a N-terminus, or both C-terminus and N-terminus of the non-antibody sequence.

5 [00740] Embodiment 110. The library of Embodiment 71, wherein the ultralong CDR3 is a ruminant CDR3.

[00741] Embodiment 111. The library of Embodiment 110, wherein the ruminant is a cow.

10 [00742] Embodiment 112. The library of Embodiment 71, wherein the humanized antibody or binding fragment thereof comprises a human heavy chain germline sequence that is a VH4 family member germline sequence.

[00743] Embodiment 113. The library of Embodiment 71, wherein the humanized antibody or binding fragment thereof comprises a VH4-34 germline sequence.

15 [00744] Embodiment 114. The library of Embodiment 71, wherein the humanized antibody or binding fragment thereof comprises a human heavy chain germline sequence that is derived from VH4-34.

[00745] Embodiment 115. The library of Embodiment 114, wherein the VH4-34 germline sequence comprises a CDR1 comprising a Gly31Asp and Tyr32Lys substitution based on Kabat numbering.

20 [00746] Embodiment 116. The library of Embodiment 114, wherein the VH4-34 germline sequence comprises a CDR2 comprising a Glu50Ser substitution based on Kabat numbering.

25 [00747] Embodiment 117. The library of Embodiment 114, wherein the VH4-34 germline sequence comprises a CDR1 comprising Ala23Thr, Val24Ala, Tyr25Ser, Gly27Phe, Phe29Leu, Gly31Asp, Tyr32Lys, Tyr33Ala, Trp34Val, and Ser35Gly substitutions, and a CDR2 comprising Ile48Leu, Glu50Ser, Asn52Asp, His53Thr, Ser54Gly, Ser56Asn, and Asn58Gly substitutions, based on Kabat numbering.

[00748] Embodiment 118. The library of Embodiment 71, wherein the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 470.

30 [00749] Embodiment 119. The library of Embodiment 71, wherein the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469; and the amino acid sequence set forth in SEQ ID NO: 470.

35 [00750] Embodiment 120. The library of Embodiment 71, wherein the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469; the amino acid sequence set forth

in SEQ ID NO: 470; and a peptide sequence, wherein the peptide sequence is positioned between SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469, and SEQ ID NO: 470.

5 [00751] Embodiment 121. The library of Embodiment 120, wherein the peptide sequence is a bovine sequence, a non-bovine sequence, an antibody sequence, or a non-antibody sequence.

[00752] Embodiment 122. The library of Embodiment 71, wherein the humanized antibody or binding fragment thereof comprises a light chain variable region sequence that is a lambda light chain variable region sequence.

10 [00753] Embodiment 123. The library of Embodiment 122, wherein the lambda light chain variable region sequence is a bovine lambda light chain variable region sequence.

[00754] Embodiment 124. The library of Embodiment 122, wherein the lambda light chain variable region sequence is a human lambda light chain variable region
15 sequence.

[00755] Embodiment 125. The library of Embodiment 124, wherein the human lambda light chain variable region sequence is VL1-51.

[00756] Embodiment 126. The library of Embodiment 124, wherein the human lambda light chain variable region sequence is derived from VL1-51.

20 [00757] Embodiment 127. The library of Embodiment 126, wherein the VL1-51 germline sequence comprises a CDR1 comprising Ile29Val and Asn32Gly substitution based on Kabat numbering.

[00758] Embodiment 128. The library of Embodiment 126, wherein the VL1-51 germline sequence comprises a CDR2 comprising a substitution of DNN to GDT.

25 [00759] Embodiment 129. The library of Embodiment 126, wherein the VL1-51 germline sequence comprises a CDR2 comprising a substitution of DNNKRP (SEQ ID NO: 471) to GDTSRA (SEQ ID NO: 472).

[00760] Embodiment 130. The library of Embodiment 126, wherein the VL1-51 germline sequence comprises a S2A, T5N, P8S, A12G, A13S, and P14L substitution
30 based on Kabat numbering.

[00761] Embodiment 131. The library of Embodiment 126, wherein the VL1-51 germline sequence comprises a S2A, T5N, P8S, A12G, A13S, and P14L substitution based on Kabat numbering, and a CDR2 comprising a substitution of DNN to GDT.

35 [00762] Embodiment 132. The library of Embodiment 71, wherein the humanized antibody or binding fragment thereof comprising the amino acid sequence set forth in SEQ ID NO: 440, 441, 442, 443, or 444.

[00763] Embodiment 133. The library of any one of Embodiments 71-132, wherein the humanized antibodies or binding fragments thereof are present in a spatially addressed format.

5 [00764] Embodiment 134. A method of humanizing an antibody variable region comprising the step of genetically combining a nucleic acid sequence encoding a non-human ultralong CDR3 with a nucleic acid sequence encoding a human variable region framework (FR) sequence.

10 [00765] Embodiment 135. A method of generating a library of humanized antibodies that comprises an ultralong CDR3, the method comprising: combining a nucleic acid sequence encoding an ultralong CDR3 with a nucleic acid sequence encoding a human variable region framework (FR) sequence to produce nucleic acids encoding for humanized antibodies that comprises an ultralong CDR3; and expressing the nucleic acids encoding for humanized antibodies that comprises an ultralong CDR3 to generate a library of humanized antibodies that comprises an ultralong CDR3.

15 [00766] Embodiment 136. A method of generating a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 that comprises a non-antibody sequence, the method comprising: combining a nucleic acid sequence encoding an ultralong CDR3, a nucleic acid sequence encoding a human variable region framework (FR) sequence, and a nucleic acid sequence encoding a non-antibody sequence to produce nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a non-antibody sequence, and expressing the nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a non-antibody sequence to generate a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a non-antibody sequence.

[00767] Embodiment 137. A library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 that comprises a non-antibody sequence.

30 [00768] Embodiment 138. A method of generating a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 that comprises a cysteine motif, the method comprising: combining a human variable region framework (FR) sequence, and a nucleic acid sequence encoding an ultralong CDR3 and a cysteine motif; introducing one or more nucleotide changes to the nucleic acid sequence encoding one or more amino acid residues that are positioned between one or more cysteine residues in the cysteine motif for nucleotides encoding different amino acid residues to produce nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a cysteine motif with one or more nucleotide changes introduced between one

or more cysteine residues in the cysteine domain; and expressing the nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a cysteine motif with one or more nucleotide changes introduced between one or more cysteine residues in the cysteine domain to generate a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a cysteine motif with one or more amino acid changes introduced between one or more cysteine residues in the cysteine domain.

[00769] Embodiment 139. A library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 that comprises a cysteine motif, wherein the antibodies or binding fragments comprise one or more substitutions of amino acid residues that are positioned between cysteine residues in the cysteine motif.

[00770] Embodiment 140. A method of generating a library of humanized antibodies or binding fragments thereof comprising a bovine ultralong CDR3, the method comprising: combining a nucleic acid sequence encoding a human variable region framework (FR) sequence and a nucleic acid encoding a bovine ultralong CDR3, and expressing the nucleic acids encoding a human variable region framework (FR) sequence and a nucleic acid encoding a bovine ultralong CDR3 to generate a library of humanized antibodies or binding fragments thereof comprising a bovine ultralong CDR3.

[00771] Embodiment 141. A library of humanized antibodies or binding fragments thereof comprising a bovine ultralong CDR3.

[00772] Embodiment 142. A CDR3 scaffold comprising a $X^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q).

[00773] Embodiment 143. The CDR3 scaffold of Embodiment 142, wherein the CDR3 scaffold is 35 amino acids in length or longer, 40 amino acids in length or longer, 45 amino acids in length or longer, 50 amino acids in length or longer, 55 amino acids in length or longer, or 60 amino acids in length or longer.

[00774] Embodiment 144. The CDR3 scaffold of Embodiment 143, wherein the CDR3 scaffold is 35 amino acids in length or longer.

[00775] Embodiment 145. The CDR3 scaffold of Embodiment 143, wherein the CDR3 scaffold comprises 3 or more cysteine residues, 4 or more cysteine residues, 5 or more cysteine residues, 6 or more cysteine residues, 7 or more cysteine residues, 8 or

more cysteine residues, 9 or more cysteine residues, 10 or more cysteine residues, 11 or more cysteine residues, or 12 or more cysteine residues.

[00776] Embodiment 146. The CDR3 scaffold of Embodiment 145, wherein the CDR3 scaffold comprises 3 or more cysteine residues.

5 [00777] Embodiment 147. The CDR3 scaffold of Embodiment 142, wherein the CDR3 scaffold comprises a cysteine motif.

[00778] Embodiment 148. The CDR3 scaffold of Embodiment 147, wherein the cysteine motif is selected from the group consisting of: CX₁₀CX₅CX₅CXCX₇C (SEQ ID NO: 41), CX₁₀CX₆CX₅CXCX₁₅C (SEQ ID NO: 42), CX₁₁CXCX₅C (SEQ ID NO: 43),
 10 CX₁₁CX₅CX₅CXCX₇C (SEQ ID NO: 44), CX₁₀CX₆CX₅CXCX₁₃C (SEQ ID NO: 45), CX₁₀CX₅CXCX₄CX₈C (SEQ ID NO: 46), CX₁₀CX₆CX₆CXCX₇C (SEQ ID NO: 47), CX₁₀CX₄CX₇CXCX₈C (SEQ ID NO: 48), CX₁₀CX₄CX₇CXCX₇C (SEQ ID NO: 49), CX₁₃CX₈CX₈C (SEQ ID NO: 50), CX₁₀CX₆CX₅CXCX₇C (SEQ ID NO: 51), CX₁₀CX₅CX₅C (SEQ ID NO: 52), CX₁₀CX₅CX₆CXCX₇C (SEQ ID NO: 53), CX₁₀CX₆CX₅CX₇CX₉C (SEQ ID
 15 NO: 54), CX₉CX₇CX₅CXCX₇C (SEQ ID NO: 55), CX₁₀CX₆CX₅CXCX₉C (SEQ ID NO: 56), CX₁₀CXCX₄CX₅CX₁₁C (SEQ ID NO: 57), CX₇CX₃CX₆CX₅CXCX₅CX₁₀C (SEQ ID NO: 58), CX₁₀CXCX₄CX₅CXCX₂CX₃C (SEQ ID NO: 59), CX₁₆CX₅CXC (SEQ ID NO: 60), CX₆CX₄CXCX₄CX₅C (SEQ ID NO: 61), CX₁₁CX₄CX₅CX₆CX₃C (SEQ ID NO: 62), CX₈CX₂CX₆CX₅C (SEQ ID NO: 63), CX₁₀CX₅CX₅CXCX₁₀C (SEQ ID NO: 64),
 20 CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65), CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66), CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67), CX₁₅CX₅CXC (SEQ ID NO: 68), CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69), CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70), CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72), CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74), CX₁₀CXCX₆C (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCX₁₄C (SEQ ID NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83), CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85),
 30 CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87), CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89), CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91), CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93), CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ
 35 ID NO: 95).

[00779] Embodiment 149. The CDR3 scaffold of Embodiment 147, wherein the cysteine motif is selected from the group consisting of: CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCCX₄CX₆CXC (SEQ ID NO: 98),
 5 CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102), CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC
 10 (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109), CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113),
 15 CX₉CCX₄CX₂CXCCXCX₄CX₃C (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116), CX₆CCXCX₅CX₄CCXCX₅CX₂C (SEQ ID NO: 117), CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118), CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO: 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141),
 30 CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152).

35 [00780] Embodiment 150. The CDR3 scaffold of Embodiment 142, wherein the CDR3 scaffold comprises 2 to 6 disulfide bonds.

[00781] Embodiment 151. The CDR3 scaffold of Embodiment 142, wherein the $X^1X^2X^3X^4X^5$ motif in the CDR3 scaffold is TTVHQ (SEQ ID NO: 153), TSVHQ (SEQ ID NO: 154), SSVTQ (SEQ ID NO: 155), STVHQ (SEQ ID NO: 156), ATVRQ (SEQ ID NO: 157), TTVYQ (SEQ ID NO: 158), SPVHQ (SEQ ID NO: 159), ATVYQ (SEQ ID NO: 160), TAVYQ (SEQ ID NO: 161), TNVHQ (SEQ ID NO: 162), ATVHQ (SEQ ID NO: 163), STVYQ (SEQ ID NO: 164), TIVHQ (SEQ ID NO: 165), AIVYQ (SEQ ID NO: 166), TTVFQ (SEQ ID NO: 167), AAVFQ (SEQ ID NO: 168), GTVHQ (SEQ ID NO: 169), ASVHQ (SEQ ID NO: 170), TAVFQ (SEQ ID NO: 171), ATVFQ (SEQ ID NO: 172), AAAHQ (SEQ ID NO: 173), VVVYQ (SEQ ID NO: 174), GTVFQ (SEQ ID NO: 175), TAVHQ (SEQ ID NO: 176), ITVHQ (SEQ ID NO: 177), ITAHQ (SEQ ID NO: 178), VTVHQ (SEQ ID NO: 179); AAVHQ (SEQ ID NO: 180), GTVYQ (SEQ ID NO: 181), TTVLQ (SEQ ID NO: 182), TTTHQ (SEQ ID NO: 183), or TTDYQ (SEQ ID NO: 184).

[00782] Embodiment 152. The CDR3 scaffold of Embodiment 142, wherein the CDR3 scaffold comprises a $CX^1X^2X^3X^4X^5$ motif.

[00783] Embodiment 153. The CDR3 scaffold of Embodiment 152, wherein the $CX^1X^2X^3X^4X^5$ motif in the CDR3 scaffold is CTTVHQ (SEQ ID NO: 185), CTSVHQ (SEQ ID NO: 186), CSSVTQ (SEQ ID NO: 187), CSTVHQ (SEQ ID NO: 188), CATVRQ (SEQ ID NO: 189), CTTVYQ (SEQ ID NO: 190), CSPVHQ (SEQ ID NO: 191), CATVYQ (SEQ ID NO: 192), CTAVYQ (SEQ ID NO: 193), CTNVHQ (SEQ ID NO: 194), CATVHQ (SEQ ID NO: 195), CSTVYQ (SEQ ID NO: 196), CTIVHQ (SEQ ID NO: 197), CAIVYQ (SEQ ID NO: 198), CTTVFQ (SEQ ID NO: 199), CAAVFQ (SEQ ID NO: 200), CGTVHQ (SEQ ID NO: 201), CASVHQ (SEQ ID NO: 202), CTAVFQ (SEQ ID NO: 203), CATVFQ (SEQ ID NO: 204), CAAAHQ (SEQ ID NO: 205), CVVVYQ (SEQ ID NO: 206), CGTVFQ (SEQ ID NO: 207), CTAVHQ (SEQ ID NO: 208), CITVHQ (SEQ ID NO: 209), CITAHQ (SEQ ID NO: 210), CVTVHQ (SEQ ID NO: 211); CAAVHQ (SEQ ID NO: 212), CGTVYQ (SEQ ID NO: 213), CTTVLQ (SEQ ID NO: 214), CTTTHQ (SEQ ID NO: 215), or CTTDYQ (SEQ ID NO: 216).

[00784] Embodiment 154. The CDR3 scaffold of Embodiment 142, wherein the CDR3 scaffold comprises a $(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[00785] Embodiment 155. The CDR3 scaffold of Embodiment 154, wherein the $(X^aX^b)_z$ motif in the CDR3 scaffold is CYTYNYEF (SEQ ID NO: 217), HYTYTYDF (SEQ ID NO: 218), HYTYTYEW (SEQ ID NO: 219), KHRYTYEW (SEQ ID NO: 220), NYIYKYSF (SEQ ID NO: 221), PYIYTYQF (SEQ ID NO: 222), SFTYTYEW (SEQ ID NO: 223), SYIYIYQW (SEQ ID NO: 224), SYNYTYSW (SEQ ID NO: 225), SYSYSYEF (SEQ ID NO: 226), or SYIYIYQW (SEQ ID NO: 227).

226), SYTYNYDF (SEQ ID NO: 227), SYTYNYEW (SEQ ID NO: 228), SYTYNYQF (SEQ ID NO: 229), SYVWTHNF (SEQ ID NO: 230), TYKYVYEW (SEQ ID NO: 231), TYTYTYEF (SEQ ID NO: 232), TYTYTYEW (SEQ ID NO: 233), VFTYTYEF (SEQ ID NO: 234), AYTIEW (SEQ ID NO: 235), DYIYTY (SEQ ID NO: 236), IHSYEF (SEQ ID NO: 237),
 5 SFTYEF (SEQ ID NO: 238), SHSYEF (SEQ ID NO: 239), THTYEF (SEQ ID NO: 240), TWTYEF (SEQ ID NO: 241), TYNIEW (SEQ ID NO: 242), TYSYEF (SEQ ID NO: 243), TYSYEH (SEQ ID NO: 244), TYTYDF (SEQ ID NO: 245), TYTYEF (SEQ ID NO: 246), TYTYEW (SEQ ID NO: 247), AYEF (SEQ ID NO: 248), AYSF (SEQ ID NO: 249), AYSY (SEQ ID NO: 250), CYSF (SEQ ID NO: 251), DYT Y (SEQ ID NO: 252), KYEH (SEQ ID NO: 253), KYEW (SEQ ID NO: 254), MYEF (SEQ ID NO: 255), NWIY (SEQ ID NO: 256), NYDY (SEQ ID NO: 257), NYQW (SEQ ID NO: 258), NYSF (SEQ ID NO: 259), PYEW (SEQ ID NO: 260), RYNW (SEQ ID NO: 261), RYTY (SEQ ID NO: 262), SYEF (SEQ ID NO: 263), SYEH (SEQ ID NO: 264), SYEW (SEQ ID NO: 265), SYKW (SEQ ID NO: 266), SYTY (SEQ ID NO: 267), TYDF (SEQ ID NO: 268), TYEF (SEQ ID NO: 269), TYEW (SEQ ID NO: 270), TYQW (SEQ ID NO: 271), TYTY (SEQ ID NO: 272), or VYEW (SEQ ID NO: 273).

[00786] Embodiment 156. The CDR3 scaffold of Embodiment 154, wherein the $(X^aX^b)_z$ motif in the CDR3 scaffold is YXYXYX.

[00787] Embodiment 157. The CDR3 scaffold of Embodiment 142, wherein the
 20 CDR3 scaffold comprises a $X^1X^2X^3X^4X^5X_n$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q), and
 25 wherein n is 27 – 54.

[00788] Embodiment 158. The CDR3 scaffold of Embodiment 142, wherein the CDR3 scaffold comprises a $X_n(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

[00789] Embodiment 159. The CDR3 scaffold of Embodiment 142, wherein the
 30 CDR3 scaffold comprises a $X^1X^2X^3X^4X^5X_n(X^aX^b)_z$ motif, wherein X_1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X_2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X_3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X_4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), wherein X_5 is glutamine (Q), X^a
 35 is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting

of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

[00790] Embodiment 160. The CDR3 scaffold of Embodiment 159, wherein the $X^1X^2X^3X^4X^5$ motif in the CDR3 scaffold is TTVHQ (SEQ ID NO: 153) or TSVHQ (SEQ ID NO: 154), and wherein the $(X^aX^b)_z$ motif is YXYXYX.

[00791] Embodiment 161. The CDR3 scaffold of Embodiment 142, wherein the CDR3 scaffold comprises: a $CX^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q); a cysteine motif selected from the group consisting of: $CX_{10}CX_5CX_5CXCX_7C$ (SEQ ID NO: 41), $CX_{10}CX_6CX_5CXCX_{15}C$ (SEQ ID NO: 42), $CX_{11}CXCX_5C$ (SEQ ID NO: 43), $CX_{11}CX_5CX_5CXCX_7C$ (SEQ ID NO: 44), $CX_{10}CX_6CX_5CXCX_{13}C$ (SEQ ID NO: 45), $CX_{10}CX_5CXCX_4CX_8C$ (SEQ ID NO: 46), $CX_{10}CX_6CX_6CXCX_7C$ (SEQ ID NO: 47), $CX_{10}CX_4CX_7CXCX_8C$ (SEQ ID NO: 48), $CX_{10}CX_4CX_7CXCX_7C$ (SEQ ID NO: 49), $CX_{13}CX_8CX_8C$ (SEQ ID NO: 50), $CX_{10}CX_6CX_5CXCX_7C$ (SEQ ID NO: 51), $CX_{10}CX_5CX_5C$ (SEQ ID NO: 52), $CX_{10}CX_5CX_6CXCX_7C$ (SEQ ID NO: 53), $CX_{10}CX_6CX_5CX_7CX_9C$ (SEQ ID NO: 54), $CX_9CX_7CX_5CXCX_7C$ (SEQ ID NO: 55), $CX_{10}CX_6CX_5CXCX_9C$ (SEQ ID NO: 56), $CX_{10}CXCX_4CX_5CX_{11}C$ (SEQ ID NO: 57), $CX_7CX_3CX_6CX_5CXCX_5CX_{10}C$ (SEQ ID NO: 58), $CX_{10}CXCX_4CX_5CXCX_2CX_3C$ (SEQ ID NO: 59), $CX_{16}CX_5CXC$ (SEQ ID NO: 60), $CX_6CX_4CXCX_4CX_5C$ (SEQ ID NO: 61), $CX_{11}CX_4CX_5CX_6CX_3C$ (SEQ ID NO: 62), $CX_8CX_2CX_6CX_5C$ (SEQ ID NO: 63), $CX_{10}CX_5CX_5CXCX_{10}C$ (SEQ ID NO: 64), $CX_{10}CXCX_6CX_4CXC$ (SEQ ID NO: 65), $CX_{10}CX_5CX_5CXCX_2C$ (SEQ ID NO: 66), $CX_{14}CX_2CX_3CXCXC$ (SEQ ID NO: 67), $CX_{15}CX_5CXC$ (SEQ ID NO: 68), $CX_4CX_6CX_9CX_2CX_{11}C$ (SEQ ID NO: 69), $CX_6CX_4CX_5CX_5CX_{12}C$ (SEQ ID NO: 70), $CX_7CX_3CXCXCX_4CX_5CX_9C$ (SEQ ID NO: 71), $CX_{10}CX_6CX_5C$ (SEQ ID NO: 72), $CX_7CX_3CX_5CX_5CX_9C$ (SEQ ID NO: 73), $CX_7CX_5CXCX_2C$ (SEQ ID NO: 74), $CX_{10}CXCX_6C$ (SEQ ID NO: 75), $CX_{10}CX_3CX_3CX_5CX_7CXCX_6C$ (SEQ ID NO: 76), $CX_{10}CX_4CX_5CX_{12}CX_2C$ (SEQ ID NO: 77), $CX_{12}CX_4CX_5CXCXCX_9CX_3C$ (SEQ ID NO: 78), $CX_{12}CX_4CX_5CX_{12}CX_2C$ (SEQ ID NO: 79), $CX_{10}CX_6CX_5CXCX_{11}C$ (SEQ ID NO: 80), $CX_{16}CX_5CXCXCX_{14}C$ (SEQ ID NO: 81), $CX_{10}CX_5CXCX_8CX_6C$ (SEQ ID NO: 82), $CX_{12}CX_4CX_5CX_8CX_2C$ (SEQ ID NO: 83), $CX_{12}CX_5CX_5CXCX_8C$ (SEQ ID NO: 84), $CX_{10}CX_6CX_5CXCX_4CXCX_9C$ (SEQ ID NO: 85), $CX_{11}CX_4CX_5CX_8CX_2C$ (SEQ ID NO: 86), $CX_{10}CX_6CX_5CX_8CX_2C$ (SEQ ID NO: 87), $CX_{10}CX_6CX_5CXCX_8C$ (SEQ ID NO: 88), $CX_{10}CX_6CX_5CXCX_3CX_8CX_2C$ (SEQ ID NO: 89), $CX_{10}CX_6CX_5CX_3CX_8C$ (SEQ ID NO: 90), $CX_{10}CX_6CX_5CXCX_2CX_6CX_5C$ (SEQ ID NO: 91),

CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93), CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ ID NO: 95); and a (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

[00792] Embodiment 162. The CDR3 scaffold of Embodiment 142, wherein the CDR3 scaffold comprises: a CX¹X²X³X⁴X⁵ motif, wherein X¹ is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X² is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X³ is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X⁴ is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X⁵ is glutamine (Q); a cysteine motif selected from the group consisting of: wherein the cysteine motif is selected from the group consisting of: CCX₃CXCX₃CX₂CCX₅CX₉CX₅CXC (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCCX₄CX₆CXC (SEQ ID NO: 98), CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102), CCX₈CX₅CX₄CX₃CX₄CCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCX₅CX₄CXC (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109), CX₃CX₇CX₃CX₄CCX₂CX₅CX₂C (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113), CX₉CCX₄CX₂CXCCX₄CX₃C (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116), CX₆CCX₅CX₄CCX₅CX₂C (SEQ ID NO: 117), CX₆CCX₃CXCCX₃CX₄CC (SEQ ID NO: 118), CX₆CCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO: 123), CX₉CX₇CX₄CCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134),

CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136),
 CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138),
 CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140),
 CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142),
 5 CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144),
 CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C
 (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149),
 CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152); and a
 (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected
 10 from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine
 (H), and wherein z is 1 - 4.

[00793] Embodiment 163. The CDR3 scaffold of Embodiment 142, wherein the CDR3 scaffold comprises SEQ ID NO: 40 or a derivative thereof.

[00794] Embodiment 164. The CDR3 scaffold of Embodiment 142, wherein the
 15 CDR3 scaffold comprises amino acid residues 3-6 of any of one SEQ ID NO: 1-4.

[00795] Embodiment 165. The CDR3 scaffold of Embodiment 142, wherein the CDR3 scaffold comprises a non-human DH or a derivative thereof.

[00796] Embodiment 166. The CDR3 scaffold of Embodiment 165, wherein the
 non-human DH is SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID
 20 NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, or SEQ ID NO: 12

[00797] Embodiment 167. The CDR3 scaffold of Embodiment 142, wherein the CDR3 scaffold comprises a JH sequence or a derivative thereof.

[00798] Embodiment 168. The CDR3 scaffold of Embodiment 167, wherein the
 JH sequence is SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, or SEQ
 25 ID NO: 17.

[00799] Embodiment 169. The CDR3 scaffold of Embodiment 142, wherein the CDR3 scaffold comprises: a non-human VH sequence or a derivative thereof; a non-human DH sequence or a derivative thereof; and/or a JH sequence or derivative thereof.

[00800] Embodiment 170. The CDR3 scaffold of Embodiment 169, wherein the
 30 CDR3 scaffold comprises an additional amino acid sequence comprising two to six amino
 acid residues or more positioned between the VH sequence and the DH sequence.

[00801] Embodiment 171. The CDR3 scaffold of Embodiment 170, wherein the additional amino acid sequence is selected from the group consisting of: IR, IF, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20 or SEQ ID NO: 21.

35 [00802] Embodiment 172. The CDR3 scaffold of Embodiment 142, wherein the
 CDR3 scaffold comprises a sequence derived from or based on SEQ ID NO: 22, SEQ ID

NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, or SEQ ID NO: 28.

[00803] Embodiment 173. The CDR3 scaffold of Embodiment 142, wherein the CDR3 scaffold is a ruminant CDR3 scaffold.

5 [00804] Embodiment 174. The CDR3 scaffold of Embodiment 173, wherein the ruminant is a cow.

[00805] Embodiment 175. A library comprising the CDR3 scaffold of any one of Embodiments 142-174.

10 [00806] Embodiment 176. A polynucleotide encoding the CDR3 scaffold of any one of Embodiments 142-174.

[00807] Embodiment 177. A vector comprising the polynucleotide of Embodiment 176.

[00808] Embodiment 178. A host cell comprising the vector of Embodiment 177.

15 [00809] Embodiment 179. The host cell of Embodiment 178, wherein the cell is a bacteria, virus, or bacteriophage.

[00810] For the disclosure herein, the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical parameter should at least be construed in light of the number of reported significant digits and by applying ordinary rounding techniques. Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the disclosure are approximations, the numerical values set forth in the specific examples are reported as precisely as possible. Any numerical value, however, inherently contains certain errors necessarily resulting from the standard deviation found in their respective testing measurements.

20

[00811] The terms “a,” “an,” “the” and similar referents used in the context of describing the exemplary embodiments (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. Recitation of ranges of values herein is merely intended to serve as a shorthand method of referring individually to each separate value falling within the range. Unless otherwise indicated herein, each individual value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided herein is intended merely to better illuminate the exemplary embodiments and does not pose a limitation on the scope of the exemplary

30

35

embodiments otherwise claimed. No language in the specification should be construed as

indicating any non-claimed element essential to the practice of the exemplary embodiments.

[00812] Groupings of alternative elements or embodiments disclosed herein are not to be construed as limitations. Each group member can be referred to and claimed individually or in any combination with other members of the group or other elements found
5 herein. It is anticipated that one or more members of a group can be included in, or deleted from, a group for reasons of convenience and/or patentability. When any such inclusion or deletion occurs, the specification is deemed to contain the group as modified thus fulfilling the written description of all Markush groups used in the appended claims.

[00813] Certain embodiments are described herein, including the best mode known to the inventors for carrying out the exemplary embodiments. Of course, variations on these described embodiments will become apparent to those of ordinary skill in the art upon reading the foregoing description. The inventor expects skilled artisans to employ such variations as appropriate, and the inventors intend for the embodiments to be
15 practiced otherwise than specifically described herein. Accordingly, this disclosure includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the disclosure unless otherwise indicated herein or otherwise clearly contradicted by context.

[00814] Furthermore, numerous references have been made to patents and printed publications. Each of the above-cited references is individually incorporated herein by reference in their entirety.

[00815] Specific embodiments disclosed herein can be further limited in the claims using consisting of or and consisting essentially of language. When used in the claims,
25 whether as filed or added per amendment, the transition term “consisting of” excludes any element, step, or ingredient not specified in the claims. The transition term “consisting essentially of” limits the scope of a claim to the specified materials or steps and those that do not materially affect the basic and novel characteristic(s). Exemplary embodiments so claimed are inherently or expressly described and enabled herein.

[00816] In closing, it is to be understood that the exemplary embodiments disclosed herein are illustrative of the principles of the present disclosure. Other modifications that can be employed are within the scope of the disclosure. Thus, by way of example, but not of limitation, alternative configurations of the present exemplary
35 embodiments can be utilized in accordance with the teachings herein. Accordingly, the present exemplary embodiments are not limited to that precisely as shown and described.

CLAIMS

- 1.) A humanized antibody or binding fragment thereof comprising an ultralong CDR3.
- 2.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong
5 CDR3 is 35 amino acids in length or longer, 40 amino acids in length or longer, 45 amino acids in length or longer, 50 amino acids in length or longer, 55 amino acids in length or longer, or 60 amino acids in length or longer.
- 3.) The humanized antibody or binding fragment thereof of claim 2, wherein the ultralong
10 CDR3 is 35 amino acids in length or longer.
- 4.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises 3 or more cysteine residues, 4 or more cysteine residues, 5 or more cysteine residues, 6 or more cysteine residues, 7 or more cysteine residues, 8 or
15 more cysteine residues, 9 or more cysteine residues, 10 or more cysteine residues, 11 or more cysteine residues, or 12 or more cysteine residues.
- 5.) The humanized antibody or binding fragment thereof of claim 4, wherein the ultralong CDR3 comprises 3 or more cysteine residues.
- 6.) The humanized antibody or binding fragment thereof of claim 1, wherein the
20 antibodies or binding fragments thereof comprise a cysteine motif.
- 7.) The humanized antibody or binding fragment thereof of claim 6, wherein the cysteine
25 motif is selected from the group consisting of: CX₁₀CX₅CX₅CXCX₇C (SEQ ID NO: 41), CX₁₀CX₆CX₅CXCX₁₅C (SEQ ID NO: 42), CX₁₁CXCX₅C (SEQ ID NO: 43), CX₁₁CX₅CX₅CXCX₇C (SEQ ID NO: 44), CX₁₀CX₆CX₅CXCX₁₃C (SEQ ID NO: 45), CX₁₀CX₅CXCX₄CX₈C (SEQ ID NO: 46), CX₁₀CX₆CX₆CXCX₇C (SEQ ID NO: 47), CX₁₀CX₄CX₇CXCX₈C (SEQ ID NO: 48), CX₁₀CX₄CX₇CXCX₇C (SEQ ID NO: 49), CX₁₃CX₈CX₈C (SEQ ID NO: 50), CX₁₀CX₆CX₅CXCX₇C (SEQ ID NO: 51), CX₁₀CX₅CX₅C (SEQ ID NO: 52), CX₁₀CX₅CX₆CXCX₇C (SEQ ID NO: 53), CX₁₀CX₆CX₅CX₇CX₉C (SEQ ID NO: 54), CX₉CX₇CX₅CXCX₇C (SEQ ID NO: 55), CX₁₀CX₆CX₅CXCX₉C (SEQ ID NO: 56), CX₁₀CXCX₄CX₅CX₁₁C (SEQ ID NO: 57), CX₇CX₃CX₆CX₅CXCX₅CX₁₀C (SEQ ID NO: 58), CX₁₀CXCX₄CX₅CXCX₂CX₃C (SEQ ID
30 NO: 59), CX₁₆CX₅CXC (SEQ ID NO: 60), CX₆CX₄CXCX₄CX₅C (SEQ ID NO: 61), CX₁₁CX₄CX₅CX₆CX₃C (SEQ ID NO: 62), CX₈CX₂CX₆CX₅C (SEQ ID NO: 63),

CX₁₀CX₅CX₅CXCX₁₀C (SEQ ID NO: 64), CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65),
 CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66), CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67),
 CX₁₅CX₅CXC (SEQ ID NO: 68), CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69),
 CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70), CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO:
 5 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72), CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73),
 CX₇CX₅CXCX₂C (SEQ ID NO: 74), CX₁₀CXCX₆C (SEQ ID NO: 75),
 CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C (SEQ ID NO:
 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C (SEQ ID
 NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCX₁₄C (SEQ ID NO:
 10 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83),
 CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO:
 85), CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO:
 87), CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID
 NO: 89), CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ
 15 ID NO: 91), CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID
 NO: 93), CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and
 CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ ID NO: 95).

8. The humanized antibody or binding fragment thereof of claim 6, wherein the cysteine
 motif is selected from the group consisting of: CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC
 (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97),
 CX₇CXCX₅CX₄CCX₄CX₆CXC (SEQ ID NO: 98), CX₉CX₃CXCX₂CXCCCX₆CX₄C
 (SEQ ID NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100),
 CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C
 (SEQ ID NO: 102), CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID NO: 103),
 CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C
 (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC (SEQ ID NO: 106),
 CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C
 (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109),
 CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCCX₆C
 (SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112),
 CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113), CX₉CCX₄CX₂CXCCXCX₄CX₃C
 (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115),
 CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116), CX₆CCXCX₅CX₄CCXCX₅CX₂C
 (SEQ ID NO: 117), CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118),
 35 CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C

(SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO: 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152).

- 9.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises 2 to 6 disulfide bonds.
- 10.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises SEQ ID NO: 40 or a derivative thereof.
- 11.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises amino acid residues 3-6 of any of one SEQ ID NO: 1-4.
- 12.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises a non-human DH or a derivative thereof.
- 13.) The humanized antibody or binding fragment thereof of claim 12, wherein the non-human DH is SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, or SEQ ID NO: 12.
- 14.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises a JH sequence or a derivative thereof.

15.) The humanized antibody or binding fragment thereof of claim 14, wherein the JH sequence is SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, or SEQ ID NO: 17.

5

16.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises:

- a non-human VH sequence or a derivative thereof;
- a non-human DH sequence or a derivative thereof; and/or
- a JH sequence or derivative thereof.

10

17.) The humanized antibody or binding fragment thereof of claim 16, wherein the ultralong CDR3 comprises an additional amino acid sequence comprising two to six amino acid residues or more positioned between the VH sequence and the DH sequence.

15

18.) The humanized antibody or binding fragment thereof of claim 17, wherein the additional amino acid sequence is selected from the group consisting of: IR, IF, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20 or SEQ ID NO: 21.

20

19.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises a sequence derived from or based on SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, or SEQ ID NO: 28.

25

20.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises a non-bovine sequence or a non-antibody sequence.

21.) The humanized antibody or binding fragment thereof of claim 20, wherein the non-antibody sequence is a synthetic sequence.

30

22.) The humanized antibody or binding fragment thereof of claim 20, wherein the non-antibody sequence is a cytokine sequence, a lymphokine sequence, a chemokine sequence, a growth factor sequence, a hormone sequence, or a toxin sequence.

35

23.) The humanized antibody or binding fragment thereof of claim 20, wherein the non-antibody sequence is an IL-8 sequence, an IL-21 sequence, an SDF-1 (alpha) sequence, a somatostatin sequence, a chlorotoxin sequence, a Pro-TxII sequence, or a ziconotide sequence.

5

24.) The humanized antibody or binding fragment thereof of claim 20, wherein the non-antibody sequence is any one of SEQ ID NOS: 475-481.

10

25.) The humanized antibody or binding fragment thereof of claim 20, wherein the non-antibody sequence replaces at least a portion of the ultralong CDR3.

26.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5$ motif,

15

wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q).

20

27.) The humanized antibody or binding fragment thereof of claim 26, wherein the $X^1X^2X^3X^4X^5$ motif is TTVHQ (SEQ ID NO: 153), TSVHQ (SEQ ID NO: 154), SSVTQ (SEQ ID NO: 155), STVHQ (SEQ ID NO: 156), ATVRQ (SEQ ID NO: 157), TTVYQ (SEQ ID NO: 158), SPVHQ (SEQ ID NO: 159), ATVYQ (SEQ ID NO: 160), TAVYQ (SEQ ID NO: 161), TNVHQ (SEQ ID NO: 162), ATVHQ (SEQ ID NO: 163), STVYQ (SEQ ID NO: 164), TIVHQ (SEQ ID NO: 165), AIVYQ (SEQ ID NO: 166), TTVFQ (SEQ ID NO: 167), AAVFQ (SEQ ID NO: 168), GTVHQ (SEQ ID NO: 169), ASVHQ (SEQ ID NO: 170), TAVFQ (SEQ ID NO: 171), ATVFQ (SEQ ID NO: 172), AAAHQ (SEQ ID NO: 173), VVVYQ (SEQ ID NO: 174), GTVYQ (SEQ ID NO: 175), TAVHQ (SEQ ID NO: 176), ITVHQ (SEQ ID NO: 177), ITAHQ (SEQ ID NO: 178), VTVHQ (SEQ ID NO: 179); AAVHQ (SEQ ID NO: 180), GTVYQ (SEQ ID NO: 181), TTVLQ (SEQ ID NO: 182), TTTHQ (SEQ ID NO: 183), or TTDYQ (SEQ ID NO: 184).

25

30

28.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises a $CX^1X^2X^3X^4X^5$ motif.

35

- 29.) The humanized antibody or binding fragment thereof of claim 28, wherein the $CX^1X^2X^3X^4X^5$ motif is CTTVHQ (SEQ ID NO: 185), CTSVHQ (SEQ ID NO: 186), CSSVTQ (SEQ ID NO: 187), CSTVHQ (SEQ ID NO: 188), CATVRQ (SEQ ID NO: 189), CTTVYQ (SEQ ID NO: 190), CSPVHQ (SEQ ID NO: 191), CATVYQ (SEQ ID NO: 192), CTAVYQ (SEQ ID NO: 193), CTNVHQ (SEQ ID NO: 194), CATVHQ (SEQ ID NO: 195), CSTVYQ (SEQ ID NO: 196), CTIVHQ (SEQ ID NO: 197), CAIVYQ (SEQ ID NO: 198), CTTVFQ (SEQ ID NO: 199), CAAVFQ (SEQ ID NO: 200), CGTVHQ (SEQ ID NO: 201), CASVHQ (SEQ ID NO: 202), CTAVFQ (SEQ ID NO: 203), CATVFQ (SEQ ID NO: 204), CAAAHQ (SEQ ID NO: 205), CVVVYQ (SEQ ID NO: 206), CGTVFQ (SEQ ID NO: 207), CTAVHQ (SEQ ID NO: 208), CITVHQ (SEQ ID NO: 209), CITAHQ (SEQ ID NO: 210), CVTVHQ (SEQ ID NO: 211), CAAVHQ (SEQ ID NO: 212), CGTVYQ (SEQ ID NO: 213), CTTVLQ (SEQ ID NO: 214), CTTTHQ (SEQ ID NO: 215), or CTTDYQ (SEQ ID NO: 216).
- 30.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises a $(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.
- 31.) The humanized antibody or binding fragment thereof of claim 30, wherein the $(X^aX^b)_z$ motif is CYTYNYEF (SEQ ID NO: 217), HYTYTYDF (SEQ ID NO: 218), HYTYTYEW (SEQ ID NO: 219), KHRITYEW (SEQ ID NO: 220), NYIYKYSF (SEQ ID NO: 221), PYIYTYQF (SEQ ID NO: 222), SFTYTYEW (SEQ ID NO: 223), SYIYIYQW (SEQ ID NO: 224), SYNYTYSW (SEQ ID NO: 225), SYSYSYEF (SEQ ID NO: 226), SYTYNYDF (SEQ ID NO: 227), SYTYNYEW (SEQ ID NO: 228), SYTYNYQF (SEQ ID NO: 229), SYVWTHNF (SEQ ID NO: 230), TYKYVYEW (SEQ ID NO: 231), TYTYTYEF (SEQ ID NO: 232), TYTYTYEW (SEQ ID NO: 233), VFTYTYEF (SEQ ID NO: 234), AYTYEW (SEQ ID NO: 235), DYIYTY (SEQ ID NO: 236), IHSYEF (SEQ ID NO: 237), SFTYEF (SEQ ID NO: 238), SHSYEF (SEQ ID NO: 239), THTYEF (SEQ ID NO: 240), TWTYEF (SEQ ID NO: 241), TYNYEW (SEQ ID NO: 242), TYSYEF (SEQ ID NO: 243), TYSYEH (SEQ ID NO: 244), TYTYDF (SEQ ID NO: 245), TYTYEF (SEQ ID NO: 246), TYTYEW (SEQ ID NO: 247), AYES (SEQ ID NO: 248), AYSF (SEQ ID NO: 249), AYSY (SEQ ID NO: 250), CYSF (SEQ ID NO: 251), DITY (SEQ ID NO: 252), KYEH (SEQ ID NO: 253), KYEW (SEQ ID NO: 254), MYEF (SEQ ID NO: 255), NWIY (SEQ ID NO: 256), NYDY (SEQ ID NO: 257), NYQW (SEQ ID NO: 258), NYSF

(SEQ ID NO: 259), PYEW (SEQ ID NO: 260), RYNW (SEQ ID NO: 261), RYTY (SEQ ID NO: 262), SYEF (SEQ ID NO: 263), SYEH (SEQ ID NO: 264), SYEW (SEQ ID NO: 265), SYKW (SEQ ID NO: 266), SYTY (SEQ ID NO: 267), TYDF (SEQ ID NO: 268), TYEF (SEQ ID NO: 269), TYEW (SEQ ID NO: 270), TYQW (SEQ ID NO: 271), TYTY (SEQ ID NO: 272), or VYEW (SEQ ID NO: 273).

32.) The humanized antibody or binding fragment thereof of claim 30, wherein the $(X^aX^b)_z$ motif is YXYXYX.

33.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5X_n$ motif, wherein X_1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X_2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X_3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X_4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), wherein X_5 is glutamine (Q), and wherein n is 27 – 54.

34.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises $X_n(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

35.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5X_n(X^aX^b)_z$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q), wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

36.) The humanized antibody or binding fragment thereof of claim 35, wherein the $X^1X^2X^3X^4X^5$ motif is TTVHQ (SEQ ID NO: 153) or TSVHQ (SEQ ID NO: 154), and wherein the $(X^aX^b)_z$ motif is YXYXYX.

5 37.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 comprises:

- a $CX^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A),
10 threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q),

- a cysteine motif selected from the group consisting of:

CX₁₀CX₅CX₅CXCX₇C (SEQ ID NO: 41), CX₁₀CX₆CX₅CXCX₁₅C (SEQ ID NO: 42),
CX₁₁CXCX₅C (SEQ ID NO: 43), CX₁₁CX₅CX₅CXCX₇C (SEQ ID NO: 44),
15 CX₁₀CX₆CX₅CXCX₁₃C (SEQ ID NO: 45), CX₁₀CX₅CXCX₄CX₈C (SEQ ID NO: 46),
CX₁₀CX₆CX₆CXCX₇C (SEQ ID NO: 47), CX₁₀CX₄CX₇CXCX₈C (SEQ ID NO: 48),
CX₁₀CX₄CX₇CXCX₇C (SEQ ID NO: 49), CX₁₃CX₈CX₈C (SEQ ID NO: 50),
CX₁₀CX₆CX₅CXCX₇C (SEQ ID NO: 51), CX₁₀CX₅CX₅C (SEQ ID NO: 52),
CX₁₀CX₅CX₆CXCX₇C (SEQ ID NO: 53), CX₁₀CX₆CX₅CX₇CX₉C (SEQ ID NO: 54),
20 CX₉CX₇CX₅CXCX₇C (SEQ ID NO: 55), CX₁₀CX₆CX₅CXCX₉C (SEQ ID NO: 56),
CX₁₀CXCX₄CX₅CX₁₁C (SEQ ID NO: 57), CX₇CX₃CX₆CX₅CXCX₅CX₁₀C (SEQ ID NO:
58), CX₁₀CXCX₄CX₅CXCX₂CX₃C (SEQ ID NO: 59), CX₁₆CX₅CXC (SEQ ID NO: 60),
CX₆CX₄CXCX₄CX₅C (SEQ ID NO: 61), CX₁₁CX₄CX₅CX₆CX₃C (SEQ ID NO: 62),
CX₈CX₂CX₆CX₅C (SEQ ID NO: 63), CX₁₀CX₅CX₅CXCX₁₀C (SEQ ID NO: 64),
25 CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65), CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66),
CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67), CX₁₅CX₅CXC (SEQ ID NO: 68),
CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69), CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70),
CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72),
CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74),
30 CX₁₀CXCX₆C (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76),
CX₁₀CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO:
78), CX₁₂CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80),
CX₁₆CX₅CXCXCX₁₄C (SEQ ID NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82),
CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83), CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84),
35 CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85), CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO:
86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87), CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88),

CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89), CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91), CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93), CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ ID NO: 95),

- 5 - a (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

38.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong
10 CDR3 comprises:

 - a CX¹X²X³X⁴X⁵ motif, wherein X¹ is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X² is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X³ is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X⁴ is histidine (H), threonine (T), arginine
15 (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X⁵ is glutamine (Q);

 - a cysteine motif selected from the group consisting of: wherein the cysteine motif is selected from the group consisting of:
CCX₃CXCX₃CX₂CCX₅CX₉CX₅CXC (SEQ ID NO: 96),
CX₆CX₂CX₅CX₄CCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCCX₄CX₆CXC
20 (SEQ ID NO: 98), CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99),
CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C
(SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102),
CCX₈CX₅CX₄CX₃CX₄CCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ
ID NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105),
25 CX₃CX₅CX₆CX₄CCX₅CX₄CXC (SEQ ID NO: 106),
CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C
(SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109),
CX₃CX₇CX₃CX₄CCX₂CX₅CX₂C (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCCX₆C
(SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112),
30 CX₈CCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113), CX₉CCX₄CX₂CXCCX₄CX₃C
(SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115),
CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116), CX₆CCX₅CX₄CCX₅CX₂C
(SEQ ID NO: 117), CX₆CCX₃CXCCX₃CX₄CC (SEQ ID NO: 118),
CX₆CCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C
35 (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO: 121),
CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO:

123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152); and

- a (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

39.) The humanized antibody or binding fragment thereof of claim 20, wherein the ultralong CDR3 comprises an additional sequence that is a linker.

40.) The humanized antibody or binding fragment thereof of claim 39, wherein the linker is linked to a C-terminus, a N-terminus, or both C-terminus and N-terminus of the non-antibody sequence.

41.) The humanized antibody or binding fragment thereof of claim 1, wherein the ultralong CDR3 is a ruminant CDR3.

42.) The humanized antibody or binding fragment thereof of claim 41, wherein the ruminant is a cow.

43.) The humanized antibody or binding fragment thereof of claim 1, wherein the humanized antibody or binding fragment thereof comprises a human heavy chain germline sequence that is a VH4 family member germline sequence.

44.) The humanized antibody or binding fragment thereof of claim 1, wherein the humanized antibody or binding fragment thereof comprises a VH4-34 germline sequence.

5 45.) The humanized antibody or binding fragment thereof of claim 44, wherein the humanized antibody or binding fragment thereof comprises a human heavy chain germline sequence that is derived from the VH4-34 germline sequence.

10 46.) The humanized antibody or binding fragment thereof of claim 45, wherein the VH4-34 germline sequence comprises a CDR1 comprising a Gly31Asp and Tyr32Lys substitution based on Kabat numbering.

15 47.) The humanized antibody or binding fragment thereof of claim 45, wherein the VH4-34 germline sequence comprises a CDR2 comprising a Glu50Ser substitution based on Kabat numbering.

20 48.) The humanized antibody or binding fragment thereof of claim 45, wherein the VH4-34 germline sequence comprises a CDR1 comprising Ala23Thr, Val24Ala, Tyr25Ser, Gly27Phe, Phe29Leu, Gly31Asp, Tyr32Lys, Tyr33Ala, Trp34Val, and Ser35Gly substitutions, and a CDR2 comprising Ile48Leu, Glu50Ser, Asn52Asp, His53Thr, Ser54Gly, Ser56Asn, and Asn58Gly substitutions, based on Kabat numbering.

25 49.) The humanized antibody or binding fragment thereof of claim 1, wherein the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469.

30 50.) The humanized antibody or binding fragment thereof of claim 1, wherein the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 470.

35 51.) The humanized antibody or binding fragment thereof of claim 1, wherein the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469; and the amino acid sequence set forth in SEQ ID NO: 470.

52.) The humanized antibody or binding fragment thereof of claim 1, wherein the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469; the amino acid sequence set forth in SEQ ID NO: 470; and a peptide sequence, wherein the peptide sequence is positioned between SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469, and SEQ ID NO: 470.

53.) The humanized antibody or binding fragment thereof of claim 52, wherein the peptide sequence is a bovine sequence, a non-bovine sequence, an antibody sequence, or a non-antibody sequence.

54.) The humanized antibody or binding fragment thereof of claim 1, wherein the humanized antibody or binding fragment thereof comprises a light chain variable region sequence that is a lambda light chain variable region sequence.

55.) The humanized antibody or binding fragment thereof of claim 54, wherein the lambda light chain variable region sequence is a bovine lambda light chain variable region sequence.

56.) The humanized antibody or binding fragment thereof of claim 54, wherein the lambda light chain variable region sequence is a human lambda light chain variable region sequence.

57.) The humanized antibody or binding fragment thereof of claim 56, wherein the human lambda light chain variable region sequence is a VL1-51 germline sequence.

58.) The humanized antibody or binding fragment thereof of claim 56, wherein the human lambda light chain variable region sequence is derived from a VL1-51 germline sequence.

59.) The humanized antibody or binding fragment thereof of claim 58, wherein the VL1-51 germline sequence comprises a CDR1 comprising Ile29Val and Asn32Gly substitution based on Kabat numbering.

60.) The humanized antibody or binding fragment thereof of claim 58, wherein the VL1-51 germline sequence comprises a CDR2 comprising a substitution of DNN to GDT.

- 61.) The humanized antibody or binding fragment thereof of claim 58, wherein the VL1-51 germline sequence comprises a CDR2 comprising a substitution of DNNKRP (SEQ ID NO: 471) to GDTsRA (SEQ ID NO: 472).
- 5
- 62.) The humanized antibody or binding fragment thereof of claim 58, wherein the VL1-51 germline sequence comprises a S2A, T5N, P8S, A12G, A13S, and P14L substitution based on Kabat numbering.
- 10
- 63.) The humanized antibody or binding fragment thereof of claim 58, wherein the VL1-51 germline sequence comprises a S2A, T5N, P8S, A12G, A13S, and P14L substitution based on Kabat numbering, and a CDR2 comprising a substitution of DNN to GDT.
- 15
- 64.) The humanized antibody or binding fragment thereof of claim 1, wherein the humanized antibody or binding fragment thereof comprising the amino acid sequence set forth in SEQ ID NO: 440, 441, 442, 443, or 444.
- 20
- 65.) A polynucleotide encoding the humanized antibody or binding fragment thereof of any one of claims 1-64.
- 25
- 66.) A polynucleotide encoding a humanized antibody or binding fragment thereof that comprises an ultralong CDR3.
- 67.) A polynucleotide encoding a humanized variable region that comprises an ultralong CDR3.
- 30
- 68.) A vector comprising the polynucleotide of any one of claims 65-67.
- 69.) A host cell comprising the vector of claim 68.
- 35
- 70.) A nucleic acid library comprising a plurality of polynucleotides comprising sequences coding for humanized antibodies or binding fragments thereof, wherein the antibodies or binding fragments thereof comprise an ultralong CDR3.
- 71.) A library of humanized antibodies or binding fragments thereof, wherein the antibodies or binding fragments thereof comprise an ultralong CDR3.

- 72.) The library of claim 71, wherein the ultralong CDR3 is 35 amino acids in length or longer, 40 amino acids in length or longer, 45 amino acids in length or longer, 50 amino acids in length or longer, 55 amino acids in length or longer, or 60 amino acids in length or longer.
- 73.) The library of claim 72, wherein the ultralong CDR3 is 35 amino acids in length or longer.
- 74.) The library of claim 71, wherein the ultralong CDR3 comprises 3 or more cysteine residues, 4 or more cysteine residues, 5 or more cysteine residues, 6 or more cysteine residues, 7 or more cysteine residues, 8 or more cysteine residues, 9 or more cysteine residues, 10 or more cysteine residues, 11 or more cysteine residues, or 12 or more cysteine residues.
- 75.) The library of claim 74, wherein the ultralong CDR3 comprises 3 or more cysteine residues.
- 76.) The library of claim 71, wherein the antibodies or binding fragments thereof comprise a cysteine motif.
- 77.) The library of claim 76, wherein the cysteine motif is selected from the group consisting of: CX₁₀CX₅CX₅CXCX₇C (SEQ ID NO: 41), CX₁₀CX₆CX₅CXCX₁₅C (SEQ ID NO: 42), CX₁₁CXCX₅C (SEQ ID NO: 43), CX₁₁CX₅CX₅CXCX₇C (SEQ ID NO: 44), CX₁₀CX₆CX₅CXCX₁₃C (SEQ ID NO: 45), CX₁₀CX₅CXCX₄CX₈C (SEQ ID NO: 46), CX₁₀CX₆CX₆CXCX₇C (SEQ ID NO: 47), CX₁₀CX₄CX₇CXCX₈C (SEQ ID NO: 48), CX₁₀CX₄CX₇CXCX₇C (SEQ ID NO: 49), CX₁₃CX₈CX₈C (SEQ ID NO: 50), CX₁₀CX₆CX₅CXCX₇C (SEQ ID NO: 51), CX₁₀CX₅CX₅C (SEQ ID NO: 52), CX₁₀CX₅CX₆CXCX₇C (SEQ ID NO: 53), CX₁₀CX₆CX₅CX₇CX₉C (SEQ ID NO: 54), CX₉CX₇CX₅CXCX₇C (SEQ ID NO: 55), CX₁₀CX₆CX₅CXCX₉C (SEQ ID NO: 56), CX₁₀CXCX₄CX₅CX₁₁C (SEQ ID NO: 57), CX₇CX₃CX₆CX₅CXCX₅CX₁₀C (SEQ ID NO: 58), CX₁₀CXCX₄CX₅CXCX₂CX₃C (SEQ ID NO: 59), CX₁₆CX₅CXC (SEQ ID NO: 60), CX₆CX₄CXCX₄CX₅C (SEQ ID NO: 61), CX₁₁CX₄CX₅CX₆CX₃C (SEQ ID NO: 62), CX₈CX₂CX₆CX₅C (SEQ ID NO: 63), CX₁₀CX₅CX₅CXCX₁₀C (SEQ ID NO: 64), CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65), CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66), CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67), CX₁₅CX₅CXC (SEQ ID NO: 68),

CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69), CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70), CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72), CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74), CX₁₀CXCX₆C (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCX₁₄C (SEQ ID NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83), CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85), CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87), CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89), CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91), CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93), CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ ID NO: 95).

78. The library of claim 76, wherein the cysteine motif is selected from the group consisting of: CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCCX₄CX₆CXC (SEQ ID NO: 98), CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102), CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109), CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113), CX₉CCX₄CX₂CXCCXCX₄CX₃C (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116), CX₆CCXCX₅CX₄CCXCX₅CX₂C (SEQ ID NO: 117), CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118), CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO: 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ

ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152).

79.) The library of claim 71, wherein the ultralong CDR3 comprises 2 to 6 disulfide bonds.

80.) The library of claim 71, wherein the ultralong CDR3 comprises SEQ ID NO: 40 or a derivative thereof.

81.) The library of claim 71, wherein the ultralong CDR3 comprises amino acid residues 3-6 of any of one SEQ ID NO: 1-4.

82.) The library of claim 71, wherein the ultralong CDR3 comprises a non-human DH or a derivative thereof.

83.) The library of claim 82, wherein the non-human DH is SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, or SEQ ID NO: 12

84.) The library of claim 71, wherein the ultralong CDR3 comprises a JH sequence or a derivative thereof.

85.) The library of claim 84, wherein the JH sequence is SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, or SEQ ID NO: 17.

86.) The library of claim 71, wherein the ultralong CDR3 comprises:

- a non-human VH sequence or a derivative thereof;
- a non-human DH sequence or a derivative thereof; and/or
- a JH sequence or derivative thereof.

5

87.) The library of claim 86, wherein the ultralong CDR3 comprises an additional amino acid sequence comprising two to six amino acid residues or more positioned between the VH sequence and the DH sequence.

10 88.) The library of claim 87, wherein the additional amino acid sequence is selected from the group consisting of: IR, IF, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20 or SEQ ID NO: 21.

15 89.) The library of claim 71, wherein the ultralong CDR3 comprises a sequence derived from or based on SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, or SEQ ID NO: 28.

90.) The library of claim 71, wherein the ultralong CDR3 comprises a non-bovine sequence or a non-antibody sequence.

20

91.) The library of claim 90, wherein the non-antibody sequence is a synthetic sequence.

92.) The library of claim 90, wherein the non-antibody sequence is a cytokine sequence, a lymphokine sequence, a chemokine sequence, a growth factor sequence, a hormone
25 sequence, or a toxin sequence.

93.) The library of claim 90, wherein the non-antibody sequence is an IL-8 sequence, an IL-21 sequence, an SDF-1 (alpha) sequence, a somatostatin sequence, a chlorotoxin sequence, a Pro-TxII sequence, or a ziconotide sequence.

30

94.) The library of claim 90, wherein the non-antibody sequence is any one of SEQ ID NOS: 475-481.

95.) The library of claim 71, wherein the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5$ motif,
35 wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or

asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q).

5 96.) The library of claim 95, wherein the $X^1X^2X^3X^4X^5$ motif is TTVHQ (SEQ ID NO: 153), TSVHQ (SEQ ID NO: 154), SSVTQ (SEQ ID NO: 155), STVHQ (SEQ ID NO: 156), ATVRQ (SEQ ID NO: 157), TTVYQ (SEQ ID NO: 158), SPVHQ (SEQ ID NO: 159), ATVYQ (SEQ ID NO: 160), TAVYQ (SEQ ID NO: 161), TNVHQ (SEQ ID NO: 162), ATVHQ (SEQ ID NO: 163), STVYQ (SEQ ID NO: 164), TIVHQ (SEQ ID NO: 165),
 10 AIVYQ (SEQ ID NO: 166), TTVFQ (SEQ ID NO: 167), AAVFQ (SEQ ID NO: 168), GTVHQ (SEQ ID NO: 169), ASVHQ (SEQ ID NO: 170), TAVFQ (SEQ ID NO: 171), ATVFQ (SEQ ID NO: 172), AAAHQ (SEQ ID NO: 173), VVVYQ (SEQ ID NO: 174), GTVFQ (SEQ ID NO: 175), TAVHQ (SEQ ID NO: 176), ITVHQ (SEQ ID NO: 177), ITAHQ (SEQ ID NO: 178), VTVHQ (SEQ ID NO: 179); AAVHQ (SEQ ID NO: 180),
 15 GTVYQ (SEQ ID NO: 181), TTVLQ (SEQ ID NO: 182), TTTHQ (SEQ ID NO: 183), or TTDYQ (SEQ ID NO: 184).

97.) The library of claim 71, wherein the ultralong CDR3 comprises a $CX^1X^2X^3X^4X^5$ motif.

20 98.) The library of claim 97, wherein the $CX^1X^2X^3X^4X^5$ motif is CTTVHQ (SEQ ID NO: 185), CTSVHQ (SEQ ID NO: 186), CSSVTQ (SEQ ID NO: 187), CSTVHQ (SEQ ID NO: 188), CATVRQ (SEQ ID NO: 189), CTTVYQ (SEQ ID NO: 190), CSPVHQ (SEQ ID NO: 191), CATVYQ (SEQ ID NO: 192), CTAVYQ (SEQ ID NO: 193), CTNVHQ (SEQ ID NO: 194), CATVHQ (SEQ ID NO: 195), CSTVYQ (SEQ ID NO: 196), CTIVHQ
 25 (SEQ ID NO: 197), CAIVYQ (SEQ ID NO: 198), CTTVFQ (SEQ ID NO: 199), CAAVFQ (SEQ ID NO: 200), CGTVHQ (SEQ ID NO: 201), CASVHQ (SEQ ID NO: 202), CTAVFQ (SEQ ID NO: 203), CATVFQ (SEQ ID NO: 204), CAAAHQ (SEQ ID NO: 205), CVVVYQ (SEQ ID NO: 206), CGTVFQ (SEQ ID NO: 207), CTAVHQ (SEQ ID NO: 208), CITVHQ (SEQ ID NO: 209), CITAHQ (SEQ ID NO: 210), CVTVHQ (SEQ ID NO: 211); CAAVHQ (SEQ ID NO: 212), CGTVYQ (SEQ ID NO: 213), CTTVLQ (SEQ ID NO: 214), CTTTHQ (SEQ ID NO: 215), or CTTDYQ (SEQ ID NO: 216).

99.) The library of claim 71, wherein the ultralong CDR3 comprises a $(X^aX^b)_z$ motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the
 35 group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

100.) The library of claim 99, wherein the $(X^aX^b)_z$ motif is CYTYNYEF (SEQ ID NO: 217),
 HYTYTYDF (SEQ ID NO: 218), HYTYTYEW (SEQ ID NO: 219), KHRITYEW (SEQ ID
 NO: 220), NYIYKYSF (SEQ ID NO: 221), PYIYTYQF (SEQ ID NO: 222), SFTYTYEW
 5 (SEQ ID NO: 223), SYIYIYQW (SEQ ID NO: 224), SYNITYSW (SEQ ID NO: 225),
 SYSYSYEF (SEQ ID NO: 226), SYTYNYDF (SEQ ID NO: 227), SYTYNYEW (SEQ ID
 NO: 228), SYTYNYQF (SEQ ID NO: 229), SYVWTHNF (SEQ ID NO: 230),
 TYKYVYEW (SEQ ID NO: 231), TYTYTYEF (SEQ ID NO: 232), TYTYTYEW (SEQ ID
 NO: 233), VFTYTYEF (SEQ ID NO: 234), AITYEW (SEQ ID NO: 235), DIIYTY (SEQ
 10 ID NO: 236), IHSYEF (SEQ ID NO: 237), SFTYEF (SEQ ID NO: 238), SHSYEF (SEQ
 ID NO: 239), THTYEF (SEQ ID NO: 240), TWTYEF (SEQ ID NO: 241), TYNIEW
 (SEQ ID NO: 242), TYSYEF (SEQ ID NO: 243), TYSYEH (SEQ ID NO: 244), TYTYDF
 (SEQ ID NO: 245), TYTYEF (SEQ ID NO: 246), TYTYEW (SEQ ID NO: 247), AIEF
 (SEQ ID NO: 248), AYSF (SEQ ID NO: 249), AYSY (SEQ ID NO: 250), CYSF (SEQ
 15 ID NO: 251), DITY (SEQ ID NO: 252), KYEH (SEQ ID NO: 253), KYEW (SEQ ID NO:
 254), MYEF (SEQ ID NO: 255), NWIY (SEQ ID NO: 256), NYDY (SEQ ID NO: 257),
 NYQW (SEQ ID NO: 258), NYSF (SEQ ID NO: 259), PYEW (SEQ ID NO: 260),
 RYNW (SEQ ID NO: 261), RITY (SEQ ID NO: 262), SYEF (SEQ ID NO: 263), SYEH
 (SEQ ID NO: 264), SYEW (SEQ ID NO: 265), SYKW (SEQ ID NO: 266), SYTY (SEQ
 20 ID NO: 267), TYDF (SEQ ID NO: 268), TYEF (SEQ ID NO: 269), TYEW (SEQ ID NO:
 270), TYQW (SEQ ID NO: 271), TITY (SEQ ID NO: 272), or VYEW (SEQ ID NO:
 273).

101.) The library of claim 99, wherein the $(X^aX^b)_z$ motif is YXYXYX.
 25

102.) The library of claim 71, wherein the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5X_n$
 motif,
 wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein
 X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or
 30 asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid
 (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine
 (F), or leucine (L), and wherein X^5 is glutamine (Q), and wherein n is 27 – 54.

103.) The library of claim 71, wherein the ultralong CDR3 comprises $X_n(X^aX^b)_z$ motif,

wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

5 104.) The library of claim 71, wherein the ultralong CDR3 comprises a $X^1X^2X^3X^4X^5X_n(X^aX^b)_z$ motif,

wherein X_1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X_2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X_3 is valine (V), alanine (A), threonine (T), or aspartic acid
10 (D), wherein X_4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), wherein X_5 is glutamine (Q), X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

15 105.) The library of claim 104, wherein the $X^1X^2X^3X^4X^5$ motif is TTVHQ (SEQ ID NO: 153) or TSVHQ (SEQ ID NO: 154), and wherein the $(X^aX^b)_z$ motif is YXYXYX.

106.) The library of claim 71, wherein the ultralong CDR3 comprises:

- a $CX^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A),
20 serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q),

- a cysteine motif selected from the group consisting of:
25 $CX_{10}CX_5CX_5CXCX_7C$ (SEQ ID NO: 41), $CX_{10}CX_6CX_5CXCX_{15}C$ (SEQ ID NO: 42),
 $CX_{11}CXCX_5C$ (SEQ ID NO: 43), $CX_{11}CX_5CX_5CXCX_7C$ (SEQ ID NO: 44),
 $CX_{10}CX_6CX_5CXCX_{13}C$ (SEQ ID NO: 45), $CX_{10}CX_5CXCX_4CX_8C$ (SEQ ID NO: 46),
 $CX_{10}CX_6CX_6CXCX_7C$ (SEQ ID NO: 47), $CX_{10}CX_4CX_7CXCX_8C$ (SEQ ID NO: 48),
 $CX_{10}CX_4CX_7CXCX_7C$ (SEQ ID NO: 49), $CX_{13}CX_8CX_8C$ (SEQ ID NO: 50),
30 $CX_{10}CX_6CX_5CXCX_7C$ (SEQ ID NO: 51), $CX_{10}CX_5CX_5C$ (SEQ ID NO: 52),
 $CX_{10}CX_5CX_6CXCX_7C$ (SEQ ID NO: 53), $CX_{10}CX_6CX_5CX_7CX_9C$ (SEQ ID NO: 54),
 $CX_9CX_7CX_5CXCX_7C$ (SEQ ID NO: 55), $CX_{10}CX_6CX_5CXCX_9C$ (SEQ ID NO: 56),
 $CX_{10}CXCX_4CX_5CX_{11}C$ (SEQ ID NO: 57), $CX_7CX_3CX_6CX_5CXCX_5CX_{10}C$ (SEQ ID NO:
58), $CX_{10}CXCX_4CX_5CXCX_2CX_3C$ (SEQ ID NO: 59), $CX_{16}CX_5CXC$ (SEQ ID NO: 60),
35 $CX_6CX_4CXCX_4CX_5C$ (SEQ ID NO: 61), $CX_{11}CX_4CX_5CX_6CX_3C$ (SEQ ID NO: 62),
 $CX_8CX_2CX_6CX_5C$ (SEQ ID NO: 63), $CX_{10}CX_5CX_5CXCX_{10}C$ (SEQ ID NO: 64),

CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65), CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66),
 CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67), CX₁₅CX₅CXC (SEQ ID NO: 68),
 CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69), CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70),
 CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72),
 5 CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74),
 CX₁₀CXCX₆C (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76),
 CX₁₀CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO:
 78), CX₁₂CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80),
 CX₁₆CX₅CXCXCX₁₄C (SEQ ID NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82),
 10 CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83), CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84),
 CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85), CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO:
 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87), CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88),
 CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89), CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO:
 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91), CX₇CX₆CX₃CX₃CX₉C (SEQ ID
 15 NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93), CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ
 ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ ID NO: 95), and

- a (X^aX^b)_z motif, X^a is any amino acid residue, X^b is an aromatic amino acid
 selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W),
 and histidine (H), and wherein z is 1 - 4.

20

107.) The library of claim 71, wherein the ultralong CDR3 comprises:

- a CX¹X²X³X⁴X⁵ motif, wherein X¹ is threonine (T), glycine (G), alanine (A),
 serine (S), or valine (V), wherein X² is serine (S), threonine (T), proline (P), isoleucine
 (I), alanine (A), valine (V), or asparagine (N), wherein X³ is valine (V), alanine (A),
 25 threonine (T), or aspartic acid (D), wherein X⁴ is histidine (H), threonine (T), arginine
 (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X⁵ is glutamine (Q);

- a cysteine motif selected from the group consisting of: wherein the
 cysteine motif is selected from the group consisting of:
 CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96),
 30 CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCX₄CX₆CXC
 (SEQ ID NO: 98), CX₉CX₃CXCX₂CXCCX₆CX₄C (SEQ ID NO: 99),
 CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C
 (SEQ ID NO: 101), CX₉CCX₃CX₄CCX₅CX₆C (SEQ ID NO: 102),
 CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCX₄CX₄CX₁₂C (SEQ
 35 ID NO: 104), CX₆CX₂CX₃CCX₄CX₅CX₃CX₃C (SEQ ID NO: 105),
 CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC (SEQ ID NO: 106),

CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C
 (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109),
 CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCX₆C
 (SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112),
 5 CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113), CX₉CCX₄CX₂CXCCXCX₄CX₃C
 (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115),
 CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116), CX₆CCXCX₅CX₄CCXCX₅CX₂C
 (SEQ ID NO: 117), CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118),
 CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C
 10 (SEQ ID NO: 120), CX₃CX₅CX₃CCX₄CX₉C (SEQ ID NO: 121),
 CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO:
 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCX₁₀CX₂CX₃C (SEQ ID
 NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C
 (SEQ ID NO: 127), CX₇CXCX₆CX₄CCX₁₀C (SEQ ID NO: 128),
 15 CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID
 NO: 130), CX₁₁CX₃CX₄CCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C
 (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ
 ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO:
 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138),
 20 CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140),
 CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142),
 CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144),
 CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146),
 CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ
 25 ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID
 NO: 152); and

- a (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic
 amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F),
 tryptophan (W), and histidine (H), and wherein z is 1 - 4.

30

108.) The library of claim 90, wherein the ultralong CDR3 comprises an additional sequence that is a linker.

109.) The library of claim 108, wherein the linker is linked to a C-terminus, a N-terminus, or
 35 both C-terminus and N-terminus of the non-antibody sequence.

110.) The library of claim 71, wherein the ultralong CDR3 is a ruminant CDR3.

111.) The library of claim 110, wherein the ruminant is a cow.

5 112.) The library of claim 71, wherein the humanized antibody or binding fragment thereof comprises a human heavy chain germline sequence that is a VH4 family member germline sequence.

10 113.) The library of claim 71, wherein the humanized antibody or binding fragment thereof comprises a VH4-34 germline sequence.

114.) The library of claim 71, wherein the humanized antibody or binding fragment thereof comprises a human heavy chain germline sequence that is derived from VH4-34.

15 115.) The library of claim 114, wherein the VH4-34 germline sequence comprises a CDR1 comprising a Gly31Asp and Tyr32Lys substitution based on Kabat numbering.

116.) The library of claim 114, wherein the VH4-34 germline sequence comprises a CDR2 comprising a Glu50Ser substitution based on Kabat numbering.

20 117.) The library of claim 114, wherein the VH4-34 germline sequence comprises a CDR1 comprising Ala23Thr, Val24Ala, Tyr25Ser, Gly27Phe, Phe29Leu, Gly31Asp, Tyr32Lys, Tyr33Ala, Trp34Val, and Ser35Gly substitutions, and a CDR2 comprising Ile48Leu, Glu50Ser, Asn52Asp, His53Thr, Ser54Gly, Ser56Asn, and Asn58Gly substitutions,
25 based on Kabat numbering.

118.) The library of claim 71, wherein the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 470.

30 119.) The library of claim 71, wherein the humanized antibody or binding fragment thereof comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469; and the amino acid sequence set forth in SEQ ID NO: 470.

120.) The library of claim 71, wherein the humanized antibody or binding fragment thereof
35 comprises the amino acid sequence set forth in SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 469; the amino acid sequence set forth in SEQ ID NO: 470; and a

peptide sequence, wherein the peptide sequence is positioned between SEQ ID NO: 461, 462, 463, 464, 465, 466, 467, 468, or 4697, and SEQ ID NO: 470.

121.) The library of claim 120, wherein the peptide sequence is a bovine sequence, a non-bovine sequence, an antibody sequence, or a non-antibody sequence.

122.) The library of claim 71, wherein the humanized antibody or binding fragment thereof comprises a light chain variable region sequence that is a lambda light chain variable region sequence.

123.) The library of claim 122, wherein the lambda light chain variable region sequence is a bovine lambda light chain variable region sequence.

124.) The library of claim 122, wherein the lambda light chain variable region sequence is a human lambda light chain variable region sequence.

125.) The library of claim 124, wherein the human lambda light chain variable region sequence is VL1-51.

126.) The library of claim 124, wherein the human lambda light chain variable region sequence is derived from VL1-51.

127.) The library of claim 126, wherein the VL1-51 germline sequence comprises a CDR1 comprising Ile29Val and Asn32Gly substitution based on Kabat numbering.

128.) The library of claim 126, wherein the VL1-51 germline sequence comprises a CDR2 comprising a substitution of DNN to GDT.

129.) The library of claim 126, wherein the VL1-51 germline sequence comprises a CDR2 comprising a substitution of DNNKRP (SEQ ID NO: 471) to GDTSRA (SEQ ID NO: 472).

130.) The library of claim 126, wherein the VL1-51 germline sequence comprises a S2A, T5N, P8S, A12G, A13S, and P14L substitution based on Kabat numbering.

131.) The library of claim 126, wherein the VL1-51 germline sequence comprises a S2A, T5N, P8S, A12G, A13S, and P14L substitution based on Kabat numbering, and a CDR2 comprising a substitution of DNN to GDT.

5 132.) The library of claim 71, wherein the humanized antibody or binding fragment thereof comprising the amino acid sequence set forth in SEQ ID NO: 440, 441, 442, 443, or 444.

10 133.) The library of any one of claims 71-132, wherein the humanized antibodies or binding fragments thereof are present in a spatially addressed format.

134.) A method of humanizing an antibody variable region comprising the step of genetically combining a nucleic acid sequence encoding a non-human ultralong CDR3 with a nucleic acid sequence encoding a human variable region framework (FR) sequence.
15

135.) A method of generating a library of humanized antibodies that comprises an ultralong CDR3, the method comprising:

20 a.) combining a nucleic acid sequence encoding an ultralong CDR3 with a nucleic acid sequence encoding a human variable region framework (FR) sequence to produce nucleic acids encoding for humanized antibodies that comprises an ultralong CDR3; and

25 b.) expressing the nucleic acids encoding for humanized antibodies that comprises an ultralong CDR3 to generate a library of humanized antibodies that comprises an ultralong CDR3.

136.) A method of generating a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 that comprises a non-antibody sequence, the method comprising:

30 a.) combining a nucleic acid sequence encoding an ultralong CDR3, a nucleic acid sequence encoding a human variable region framework (FR) sequence, and a nucleic acid sequence encoding a non-antibody sequence to produce nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a non-antibody sequence, and

35 b.) expressing the nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a non-antibody sequence to

generate a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a non-antibody sequence.

137.) A library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 that comprises a non-antibody sequence.

138.) A method of generating a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 that comprises a cysteine motif, the method comprising:

a.) combining a human variable region framework (FR) sequence, and a nucleic acid sequence encoding an ultralong CDR3 and a cysteine motif;

b.) introducing one or more nucleotide changes to the nucleic acid sequence encoding one or more amino acid residues that are positioned between one or more cysteine residues in the cysteine motif for nucleotides encoding different amino acid residues to produce nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a cysteine motif with one or more nucleotide changes introduced between one or more cysteine residues in the cysteine domain ; and

c.) expressing the nucleic acids encoding humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a cysteine motif with one or more nucleotide changes introduced between one or more cysteine residues in the cysteine domain to generate a library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 and a cysteine motif with one or more amino acid changes introduced between one or more cysteine residues in the cysteine domain.

139.) A library of humanized antibodies or binding fragments thereof comprising an ultralong CDR3 that comprises a cysteine motif, wherein the antibodies or binding fragments comprise one or more substitutions of amino acid residues that are positioned between cysteine residues in the cysteine motif.

140.) A method of generating a library of humanized antibodies or binding fragments thereof comprising a bovine ultralong CDR3, the method comprising:

a.) combining a nucleic acid sequence encoding a human variable region framework (FR) sequence and a nucleic acid encoding a bovine ultralong CDR3, and

b.) expressing the nucleic acids encoding a human variable region framework (FR) sequence and a nucleic acid encoding a bovine ultralong CDR3 to generate a library of humanized antibodies or binding fragments thereof comprising a bovine ultralong CDR3.

5

141.) A library of humanized antibodies or binding fragments thereof comprising a bovine ultralong CDR3.

142.) A CDR3 scaffold comprising a $X^1X^2X^3X^4X^5$ motif,

10

wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q).

15

143.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold is 35 amino acids in length or longer, 40 amino acids in length or longer, 45 amino acids in length or longer, 50 amino acids in length or longer, 55 amino acids in length or longer, or 60 amino acids in length or longer.

20

144.) The CDR3 scaffold of claim 143, wherein the CDR3 scaffold is 35 amino acids in length or longer.

145.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises 3 or more cysteine residues, 4 or more cysteine residues, 5 or more cysteine residues, 6 or more cysteine residues, 7 or more cysteine residues, 8 or more cysteine residues, 9 or more cysteine residues, 10 or more cysteine residues, 11 or more cysteine residues, or 12 or more cysteine residues.

25

146.) The CDR3 scaffold of claim 145, wherein the CDR3 scaffold comprises 3 or more cysteine residues.

30

147.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises a cysteine motif.

35

148.) The CDR3 scaffold of claim 147, wherein the cysteine motif is selected from the group consisting of: CX₁₀CX₅CX₅CXCX₇C (SEQ ID NO: 41), CX₁₀CX₆CX₅CXCX₁₅C (SEQ ID NO: 42), CX₁₁CXCX₅C (SEQ ID NO: 43), CX₁₁CX₅CX₅CXCX₇C (SEQ ID NO: 44), CX₁₀CX₆CX₅CXCX₁₃C (SEQ ID NO: 45), CX₁₀CX₅CXCX₄CX₈C (SEQ ID NO: 46), CX₁₀CX₆CX₆CXCX₇C (SEQ ID NO: 47), CX₁₀CX₄CX₇CXCX₈C (SEQ ID NO: 48), CX₁₀CX₄CX₇CXCX₇C (SEQ ID NO: 49), CX₁₃CX₈CX₈C (SEQ ID NO: 50), CX₁₀CX₆CX₅CXCX₇C (SEQ ID NO: 51), CX₁₀CX₅CX₅C (SEQ ID NO: 52), CX₁₀CX₅CX₆CXCX₇C (SEQ ID NO: 53), CX₁₀CX₆CX₅CX₇CX₉C (SEQ ID NO: 54), CX₉CX₇CX₅CXCX₇C (SEQ ID NO: 55), CX₁₀CX₆CX₅CXCX₉C (SEQ ID NO: 56), CX₁₀CXCX₄CX₅CX₁₁C (SEQ ID NO: 57), CX₇CX₃CX₆CX₅CXCX₅CX₁₀C (SEQ ID NO: 58), CX₁₀CXCX₄CX₅CXCX₂CX₃C (SEQ ID NO: 59), CX₁₆CX₅CXC (SEQ ID NO: 60), CX₆CX₄CXCX₄CX₅C (SEQ ID NO: 61), CX₁₁CX₄CX₅CX₆CX₃C (SEQ ID NO: 62), CX₈CX₂CX₆CX₅C (SEQ ID NO: 63), CX₁₀CX₅CX₅CXCX₁₀C (SEQ ID NO: 64), CX₁₀CXCX₆CX₄CXC (SEQ ID NO: 65), CX₁₀CX₅CX₅CXCX₂C (SEQ ID NO: 66), CX₁₄CX₂CX₃CXCXC (SEQ ID NO: 67), CX₁₅CX₅CXC (SEQ ID NO: 68), CX₄CX₆CX₉CX₂CX₁₁C (SEQ ID NO: 69), CX₆CX₄CX₅CX₅CX₁₂C (SEQ ID NO: 70), CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72), CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74), CX₁₀CXCX₆C (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCX₁₄C (SEQ ID NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83), CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85), CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87), CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89), CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91), CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93), CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ ID NO: 95).

149. The CDR3 scaffold of claim 147, wherein the cysteine motif is selected from the group consisting of: CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCCX₄CX₆CXC (SEQ ID NO: 98), CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C

(SEQ ID NO: 102), CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109), CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCCX₆C (SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113), CX₉CCX₄CX₂CXCCXCX₄CX₃C (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116), CX₆CCXCX₅CX₄CCXCX₅CX₂C (SEQ ID NO: 117), CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118), CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCCX₄CX₉C (SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO: 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCCX₁₀CX₂CX₃C (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152).

30

150.) The CDR3 scaffold of claim 142, wherein CDR3 scaffold comprises 2 to 6 disulfide bonds.

151.) The CDR3 scaffold of claim 142, wherein the X¹X²X³X⁴X⁵ motif in the CDR3 scaffold is TTVHQ (SEQ ID NO: 153), TSVHQ (SEQ ID NO: 154), SSVTQ (SEQ ID NO: 155), STVHQ (SEQ ID NO: 156), ATVRQ (SEQ ID NO: 157), TTVYQ (SEQ ID

35

NO: 158), SPVHQ (SEQ ID NO: 159), ATVYQ (SEQ ID NO: 160), TAVYQ (SEQ ID NO: 161), TNVHQ (SEQ ID NO: 162), ATVHQ (SEQ ID NO: 163), STVYQ (SEQ ID NO: 164), TIVHQ (SEQ ID NO: 165), AIVYQ (SEQ ID NO: 166), TTVFQ (SEQ ID NO: 167), AAVFQ (SEQ ID NO: 168), GTVHQ (SEQ ID NO: 169), ASVHQ (SEQ ID NO: 170), TAVFQ (SEQ ID NO: 171), ATVFQ (SEQ ID NO: 172), AAAHQ (SEQ ID NO: 173), VVYQ (SEQ ID NO: 174), GTVFQ (SEQ ID NO: 175), TAVHQ (SEQ ID NO: 176), ITVHQ (SEQ ID NO: 177), ITAHQ (SEQ ID NO: 178), VTVHQ (SEQ ID NO: 179); AAVHQ (SEQ ID NO: 180), GTVYQ (SEQ ID NO: 181), TTVLQ (SEQ ID NO: 182), TTTHQ (SEQ ID NO: 183), or TTDYQ (SEQ ID NO: 184).

10

152.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises a $CX^1X^2X^3X^4X^5$ motif.

15

153.) The CDR3 scaffold of claim 152, wherein the $CX^1X^2X^3X^4X^5$ motif in the CDR3 scaffold is CTTVHQ (SEQ ID NO: 185), CTSVHQ (SEQ ID NO: 186), CSSVTQ (SEQ ID NO: 187), CSTVHQ (SEQ ID NO: 188), CATVRQ (SEQ ID NO: 189), CTTVYQ (SEQ ID NO: 190), CSPVHQ (SEQ ID NO: 191), CATVYQ (SEQ ID NO: 192), CTAVYQ (SEQ ID NO: 193), CTNVHQ (SEQ ID NO: 194), CATVHQ (SEQ ID NO: 195), CSTVYQ (SEQ ID NO: 196), CTIVHQ (SEQ ID NO: 197), CAIVYQ (SEQ ID NO: 198), CTTVFQ (SEQ ID NO: 199), CAAVFQ (SEQ ID NO: 200), CGTVHQ (SEQ ID NO: 201), CASVHQ (SEQ ID NO: 202), CTAVFQ (SEQ ID NO: 203), CATVFQ (SEQ ID NO: 204), CAAAHQ (SEQ ID NO: 205), CVVYQ (SEQ ID NO: 206), CGTVFQ (SEQ ID NO: 207), CTAVHQ (SEQ ID NO: 208), CITVHQ (SEQ ID NO: 209), CITAHQ (SEQ ID NO: 210), CVTVHQ (SEQ ID NO: 211); CAAVHQ (SEQ ID NO: 212), CGTVYQ (SEQ ID NO: 213), CTTVLQ (SEQ ID NO: 214), CTTTHQ (SEQ ID NO: 215), or CTTDYQ (SEQ ID NO: 216).

20

25

154.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises a $(X^aX^b)_z$ motif,

30

wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

35

155.) The CDR3 scaffold of claim 154, wherein the $(X^aX^b)_z$ motif in the CDR3 scaffold is CYTYNYEF (SEQ ID NO: 217), HYTYTYDF (SEQ ID NO: 218), HYTYTYEW (SEQ ID NO: 219), KHRYTYEW (SEQ ID NO: 220), NYIYKYSF (SEQ ID NO: 221), PYIYTYQF

(SEQ ID NO: 222), SFTYTYEW (SEQ ID NO: 223), SYIYIQW (SEQ ID NO: 224),
 SYNYTYSW (SEQ ID NO: 225), SYSYSYEW (SEQ ID NO: 226), SYTYNYDF (SEQ ID
 NO: 227), SYTYNYEW (SEQ ID NO: 228), SYTYNYQF (SEQ ID NO: 229),
 SYVWTHNF (SEQ ID NO: 230), TYKYVYEW (SEQ ID NO: 231), TYTYTYEF (SEQ ID
 NO: 232), TYTYTYEW (SEQ ID NO: 233), VFTYTYEF (SEQ ID NO: 234), AYTIEW
 (SEQ ID NO: 235), DIYITY (SEQ ID NO: 236), IHSYEF (SEQ ID NO: 237), SFTYEF
 (SEQ ID NO: 238), SHSYEF (SEQ ID NO: 239), THTYEF (SEQ ID NO: 240),
 TWTYEF (SEQ ID NO: 241), TYNIEW (SEQ ID NO: 242), TYSYEF (SEQ ID NO:
 243), TYSYEH (SEQ ID NO: 244), TYTYDF (SEQ ID NO: 245), TYTYEF (SEQ ID NO:
 246), TYTYEW (SEQ ID NO: 247), AYES (SEQ ID NO: 248), AYSF (SEQ ID NO:
 249), AYSY (SEQ ID NO: 250), CYSF (SEQ ID NO: 251), DITY (SEQ ID NO: 252),
 KYEH (SEQ ID NO: 253), KYEW (SEQ ID NO: 254), MYEF (SEQ ID NO: 255), NWIY
 (SEQ ID NO: 256), NYDY (SEQ ID NO: 257), NYQW (SEQ ID NO: 258), NYSF (SEQ
 ID NO: 259), PYEW (SEQ ID NO: 260), RYNW (SEQ ID NO: 261), RITY (SEQ ID
 NO: 262), SYEF (SEQ ID NO: 263), SYEH (SEQ ID NO: 264), SYEW (SEQ ID NO:
 265), SYKW (SEQ ID NO: 266), SYTY (SEQ ID NO: 267), TYDF (SEQ ID NO: 268),
 TYEF (SEQ ID NO: 269), TYEW (SEQ ID NO: 270), TYQW (SEQ ID NO: 271), TITY
 (SEQ ID NO: 272), or VYEW (SEQ ID NO: 273).

156.) The CDR3 scaffold of claim 154, wherein the $(X^aX^b)_z$ motif in the CDR3 scaffold is
 YXYXYX.

157.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises a
 $X^1X^2X^3X^4X^5X_n$ motif,
 wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein
 X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or
 asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid
 (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine
 (F), or leucine (L), and wherein X^5 is glutamine (Q), and wherein n is 27 – 54.

158.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises $X_n(X^aX^b)_z$
 motif,
 wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the
 group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H),
 wherein n is 27 – 54, and wherein z is 1 - 4.

159.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises a $X^1X^2X^3X^4X^5X_n(X^aX^b)_z$ motif,

wherein X_1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X_2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X_3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X_4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), wherein X_5 is glutamine (Q), X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), wherein n is 27 – 54, and wherein z is 1 - 4.

160.) The CDR3 scaffold of claim 159, wherein the $X^1X^2X^3X^4X^5$ motif in the CDR3 scaffold is TTVHQ (SEQ ID NO: 153) or TSVHQ (SEQ ID NO: 154), and wherein the $(X^aX^b)_z$ motif is YXYXYX.

161.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises:

- a $CX^1X^2X^3X^4X^5$ motif, wherein X^1 is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X^2 is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X^3 is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X^4 is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X^5 is glutamine (Q),

- a cysteine motif selected from the group consisting of:
 $CX_{10}CX_5CX_5CXCX_7C$ (SEQ ID NO: 41), $CX_{10}CX_6CX_5CXCX_{15}C$ (SEQ ID NO: 42),
 $CX_{11}CXCX_5C$ (SEQ ID NO: 43), $CX_{11}CX_5CX_5CXCX_7C$ (SEQ ID NO: 44),
 $CX_{10}CX_6CX_5CXCX_{13}C$ (SEQ ID NO: 45), $CX_{10}CX_5CXCX_4CX_8C$ (SEQ ID NO: 46),
 $CX_{10}CX_6CX_6CXCX_7C$ (SEQ ID NO: 47), $CX_{10}CX_4CX_7CXCX_8C$ (SEQ ID NO: 48),
 $CX_{10}CX_4CX_7CXCX_7C$ (SEQ ID NO: 49), $CX_{13}CX_8CX_8C$ (SEQ ID NO: 50),
 $CX_{10}CX_6CX_5CXCX_7C$ (SEQ ID NO: 51), $CX_{10}CX_5CX_5C$ (SEQ ID NO: 52),
 $CX_{10}CX_5CX_6CXCX_7C$ (SEQ ID NO: 53), $CX_{10}CX_6CX_5CX_7CX_9C$ (SEQ ID NO: 54),
 $CX_9CX_7CX_5CXCX_7C$ (SEQ ID NO: 55), $CX_{10}CX_6CX_5CXCX_9C$ (SEQ ID NO: 56),
 $CX_{10}CXCX_4CX_5CX_{11}C$ (SEQ ID NO: 57), $CX_7CX_3CX_6CX_5CXCX_5CX_{10}C$ (SEQ ID NO: 58),
 $CX_{10}CXCX_4CX_5CXCX_2CX_3C$ (SEQ ID NO: 59), $CX_{16}CX_5CXC$ (SEQ ID NO: 60),
 $CX_6CX_4CXCX_4CX_5C$ (SEQ ID NO: 61), $CX_{11}CX_4CX_5CX_6CX_3C$ (SEQ ID NO: 62),
 $CX_8CX_2CX_6CX_5C$ (SEQ ID NO: 63), $CX_{10}CX_5CX_5CXCX_{10}C$ (SEQ ID NO: 64),
 $CX_{10}CXCX_6CX_4CXC$ (SEQ ID NO: 65), $CX_{10}CX_5CX_5CXCX_2C$ (SEQ ID NO: 66),
 $CX_{14}CX_2CX_3CXCXC$ (SEQ ID NO: 67), $CX_{15}CX_5CXC$ (SEQ ID NO: 68),
 $CX_4CX_6CX_9CX_2CX_{11}C$ (SEQ ID NO: 69), $CX_6CX_4CX_5CX_5CX_{12}C$ (SEQ ID NO: 70),

CX₇CX₃CXCXCX₄CX₅CX₉C (SEQ ID NO: 71), CX₁₀CX₆CX₅C (SEQ ID NO: 72), CX₇CX₃CX₅CX₅CX₉C (SEQ ID NO: 73), CX₇CX₅CXCX₂C (SEQ ID NO: 74), CX₁₀CXCX₆C (SEQ ID NO: 75), CX₁₀CX₃CX₃CX₅CX₇CXCX₆C (SEQ ID NO: 76), CX₁₀CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 77), CX₁₂CX₄CX₅CXCXCX₉CX₃C (SEQ ID NO: 78), CX₁₂CX₄CX₅CX₁₂CX₂C (SEQ ID NO: 79), CX₁₀CX₆CX₅CXCX₁₁C (SEQ ID NO: 80), CX₁₆CX₅CXCXCX₁₄C (SEQ ID NO: 81), CX₁₀CX₅CXCX₈CX₆C (SEQ ID NO: 82), CX₁₂CX₄CX₅CX₈CX₂C (SEQ ID NO: 83), CX₁₂CX₅CX₅CXCX₈C (SEQ ID NO: 84), CX₁₀CX₆CX₅CXCX₄CXCX₉C (SEQ ID NO: 85), CX₁₁CX₄CX₅CX₈CX₂C (SEQ ID NO: 86), CX₁₀CX₆CX₅CX₈CX₂C (SEQ ID NO: 87), CX₁₀CX₆CX₅CXCX₈C (SEQ ID NO: 88), CX₁₀CX₆CX₅CXCX₃CX₈CX₂C (SEQ ID NO: 89), CX₁₀CX₆CX₅CX₃CX₈C (SEQ ID NO: 90), CX₁₀CX₆CX₅CXCX₂CX₆CX₅C (SEQ ID NO: 91), CX₇CX₆CX₃CX₃CX₉C (SEQ ID NO: 92), CX₉CX₈CX₅CX₆CX₅C (SEQ ID NO: 93), CX₁₀CX₂CX₂CX₇CXCX₁₁CX₅C (SEQ ID NO: 94), and CX₁₀CX₆CX₅CXCX₂CX₈CX₄C (SEQ ID NO: 95), and

- a (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

162.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises:

- a CX¹X²X³X⁴X⁵ motif, wherein X¹ is threonine (T), glycine (G), alanine (A), serine (S), or valine (V), wherein X² is serine (S), threonine (T), proline (P), isoleucine (I), alanine (A), valine (V), or asparagine (N), wherein X³ is valine (V), alanine (A), threonine (T), or aspartic acid (D), wherein X⁴ is histidine (H), threonine (T), arginine (R), tyrosine (Y), phenylalanine (F), or leucine (L), and wherein X⁵ is glutamine (Q);

- a cysteine motif selected from the group consisting of: wherein the cysteine motif is selected from the group consisting of: CCX₃CXCX₃CX₂CCXCX₅CX₉CX₅CXC (SEQ ID NO: 96), CX₆CX₂CX₅CX₄CCXCX₄CX₆CXC (SEQ ID NO: 97), CX₇CXCX₅CX₄CCCX₄CX₆CXC (SEQ ID NO: 98), CX₉CX₃CXCX₂CXCCCX₆CX₄C (SEQ ID NO: 99), CX₅CX₃CXCX₄CX₄CCX₁₀CX₂CC (SEQ ID NO: 100), CX₅CXCX₁CXCX₃CCX₃CX₄CX₁₀C (SEQ ID NO: 101), CX₉CCCX₃CX₄CCCX₅CX₆C (SEQ ID NO: 102), CCX₈CX₅CX₄CX₃CX₄CCXCX₁C (SEQ ID NO: 103), CCX₆CCX₅CCCX₄CX₄CX₁₂C (SEQ ID NO: 104), CX₆CX₂CX₃CCCX₄CX₅CX₃CX₃C (SEQ ID NO: 105), CX₃CX₅CX₆CX₄CCXCX₅CX₄CXC (SEQ ID NO: 106), CX₄CX₄CCX₄CX₄CXCX₁₁CX₂CXC (SEQ ID NO: 107), CX₅CX₂CCX₅CX₄CCX₃CCX₇C (SEQ ID NO: 108), CX₅CX₅CX₃CX₂CXCCX₄CX₇CXC (SEQ ID NO: 109), CX₃CX₇CX₃CX₄CCXCX₂CX₅CX₂C (SEQ ID NO: 110), CX₉CX₃CXCX₄CCX₅CCCX₆C

(SEQ ID NO: 111), CX₉CX₃CXCX₂CXCCX₆CX₃CX₃C (SEQ ID NO: 112), CX₈CCXCX₃CCX₃CXCX₃CX₄C (SEQ ID NO: 113), CX₉CCX₄CX₂CXCCXCX₄CX₃C (SEQ ID NO: 114), CX₁₀CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 115), CX₉CXCX₃CX₂CXCCX₄CX₅CXC (SEQ ID NO: 116), CX₆CCXCX₅CX₄CCXCX₅CX₂C (SEQ ID NO: 117), CX₆CCXCX₃CXCCX₃CX₄CC (SEQ ID NO: 118), CX₆CCXCX₃CXCX₂CXCX₄CX₈C (SEQ ID NO: 119), CX₄CX₂CCX₃CXCX₄CCX₂CX₃C (SEQ ID NO: 120), CX₃CX₅CX₃CCX₄CX₉C (SEQ ID NO: 121), CCX₉CX₃CXCCX₃CX₅C (SEQ ID NO: 122), CX₉CX₂CX₃CX₄CCX₄CX₅C (SEQ ID NO: 123), CX₉CX₇CX₄CCXCX₇CX₃C (SEQ ID NO: 124), CX₉CX₃CCX₁₀CX₂CX₃C (SEQ ID NO: 125), CX₃CX₅CX₅CX₄CCX₁₀CX₆C (SEQ ID NO: 126), CX₉CX₅CX₄CCXCX₅CX₄C (SEQ ID NO: 127), CX₇CXCX₆CX₄CCX₁₀C (SEQ ID NO: 128), CX₈CX₂CX₄CCX₄CX₃CX₃C (SEQ ID NO: 129), CX₇CX₅CXCX₄CCX₇CX₄C (SEQ ID NO: 130), CX₁₁CX₃CX₄CCX₈CX₂C (SEQ ID NO: 131), CX₂CX₃CX₄CCX₄CX₅CX₁₅C (SEQ ID NO: 132), CX₉CX₅CX₄CCX₇C (SEQ ID NO: 133), CX₉CX₇CX₃CX₂CX₆C (SEQ ID NO: 134), CX₉CX₅CX₄CCX₁₄C (SEQ ID NO: 135), CX₉CX₅CX₄CCX₈C (SEQ ID NO: 136), CX₉CX₆CX₄CCXC (SEQ ID NO: 137), CX₅CCX₇CX₄CX₁₂ (SEQ ID NO: 138), CX₁₀CX₃CX₄CCX₄C (SEQ ID NO: 139), CX₉CX₄CCX₅CX₄C (SEQ ID NO: 140), CX₁₀CX₃CX₄CX₇CXC (SEQ ID NO: 141), CX₇CX₇CX₂CX₂CX₃C (SEQ ID NO: 142), CX₉CX₄CX₄CCX₆C (SEQ ID NO: 143), CX₇CXCX₃CXCX₆C (SEQ ID NO: 144), CX₇CXCX₄CXCX₄C (SEQ ID NO: 145), CX₉CX₅CX₄C (SEQ ID NO: 146), CX₃CX₆CX₈C (SEQ ID NO: 147), CX₁₀CXCX₄C (SEQ ID NO: 148), CX₁₀CCX₄C (SEQ ID NO: 149), CX₁₅C (SEQ ID NO: 150), CX₁₀C (SEQ ID NO: 151), and CX₉C (SEQ ID NO: 152); and

- a (X^aX^b)_z motif, wherein X^a is any amino acid residue, X^b is an aromatic amino acid selected from the group consisting of: tyrosine (Y), phenylalanine (F), tryptophan (W), and histidine (H), and wherein z is 1 - 4.

163.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises SEQ ID NO: 40 or a derivative thereof.

164.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises amino acid residues 3-6 of any of one SEQ ID NO: 1-4.

165.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises a non-human DH or a derivative thereof.

166.) The CDR3 scaffold of claim 165, wherein the non-human DH is SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, or SEQ ID NO: 12

5 167.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises a JH sequence or a derivative thereof.

168.) The CDR3 scaffold of claim 167, wherein the JH sequence is SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, or SEQ ID NO: 17.

10

169.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises:

- a non-human VH sequence or a derivative thereof;
- a non-human DH sequence or a derivative thereof; and/or
- a JH sequence or derivative thereof.

15

170.) The CDR3 scaffold of claim 169, wherein the CDR3 scaffold comprises an additional amino acid sequence comprising two to six amino acid residues or more positioned between the VH sequence and the DH sequence.

20 171.) The CDR3 scaffold of claim 170, wherein the additional amino acid sequence is selected from the group consisting of: IR, IF, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20 or SEQ ID NO: 21.

25 172.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold comprises a sequence derived from or based on SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, or SEQ ID NO: 28.

173.) The CDR3 scaffold of claim 142, wherein the CDR3 scaffold is a ruminant CDR3 scaffold.

30

174.) The CDR3 scaffold of claim 173, wherein the ruminant is a cow.

175.) A library comprising the CDR3 scaffold of any one of claims 142-174.

35 176.) A polynucleotide encoding the CDR3 scaffold of any one of claims 142-174.

- 177.) A vector comprising the polynucleotide of claim 176.
- 178.) A host cell comprising the vector of claim 177.
- 5 179.) The host cell of claim 178, wherein the cell is a bacteria, virus, or bacteriophage.

Figure 1

BLV1H12	CTSVHQ	ETKYYQ	S	CPDGYRERSD	CCSNRPA	CGTSD	CCRVSVFGNCL	TTLPVSYSYTYN	VEW	HVDVWGQGGLLVTVSS	(SEQ ID NO : 360)
BLV5B8	CTTVHQ	ETRKT	C	SDGYIAVDS	CCGRGQSD	GCVNDCN	SCYYGWRNCRRQ	PAIHSYEF		HVDWGRGGLLVTVSS	(SEQ ID NO : 361)
BLV5D3	CSSVTQ	RTHVSR	S	CPDGCSDGDG	CCVDGCC	CSAYR	CYTFGVRDL	SCTSYSITYTYEW		NVDWGRGGLLVTVSS	(SEQ ID NO : 362)
BLV8C11	CTTVHQ	KTRKT	C	SDAYRYDSG	CCSGGCD	CCGAD	CYVFGACT	FGLDSSYSYIYIQW		YVDWGGQGLLVTVSS	(SEQ ID NO : 363)
BF4E9	CTTVHQ	IF	C	PDGYSYGYG	CCGYGYG	CCSGYD	CYGYGGYGYGGYSSYSYSY	SEY		YGDWGGQGLLVTVSS	(SEQ ID NO : 364)
BF1H1	CTTVHP		S	PDGYSYGYG	CCGYGYG	CCSGYD	CYGYGGYGYGGYSSYSYS	SYS			(SEQ ID NO : 365)
F18	CTTVHQ	IR	C	PDGYGYGYG	CCGYGSYGYSGYDCYGYGGYGGYGGYSSYS						(SEQ ID NO : 366)

	V_H Germ <u>CTTVHQ</u>				(SEQ ID NO : 185)
	D_{H2} Germ	<u>SCPDGYSYGYGCGGYGYGCGSGYDCYGYGGYGGYGGYSSYSYTYEY</u>			(SEQ ID NO : 274)
	J_{H1} Germ	<u>YVDAMGGQLLVTVSS</u>			(SEQ ID NO : 275)
		Published sequences:		Length	
BLV1H12	<u>CTSVHQ</u> <u>ETKKYQ</u>	<u>SCPDGYRERSDCSNRPACGTSDCCRVSVFGNCLTTLPLVSYSYTYNIEW</u>	<u>HVDVW</u>	61	(SEQ ID NO : 276)
BLV5B8	<u>CTTVHQ</u> <u>ETRKT</u>	<u>CSDGYIAVDS CGRGQSDGCVNDCNSCYGWRNCRQPAHSYEF</u>	<u>HVDAW</u>	56	(SEQ ID NO : 277)
BLV5D3	<u>CSSVTQ</u> <u>RTHVSR</u>	<u>SCPDGCCSDGDCVDCGCCSAYRCYTPGVRLDSCYSITYTYEW</u>	<u>NVDAW</u>	57	(SEQ ID NO : 278)
BLV8C11	<u>CTTVHQ</u> <u>KTRTKT</u>	<u>CCSDAYRDSGCGSGCDCCGADCYVFGACTFGLDSSYSYIYIQW</u>	<u>YVDAW</u>	58	(SEQ ID NO : 279)
BF4E9	<u>CTTVHQ</u> <u>IF</u>	<u>CPDGYSYGYGCGGYGYGCGSYDCYGYGGYGGYSSYSYSYEF</u>	<u>YGDAW</u>	56	(SEQ ID NO : 280)
B-I1	<u>CTSVHQ</u> <u>KTRTTQGN</u>	<u>CPDGYTLKDDCPRCRGGCDGYDCCWGDACRSSGLCWGHNPLVTETTYTEF</u>	<u>YIDAW</u>	66	(SEQ ID NO : 281)
B-L2	<u>CATVRQ</u> <u>TTLRD</u>	<u>CPGGYTEDRS CVNTYS CGADDCCGRGDVGYPALYGYRCAAHIQRYNW</u>	<u>HADAW</u>	59	(SEQ ID NO : 282)
	Longest CDR H3s:				
UL-1	<u>CTSVHQ</u> <u>KTRTTQGEY</u>	<u>LSLMVTLKDDCPRCRGGCDGYDCCWGDACRSSGLCWGHNPLVTETTYTEF</u>	<u>YIDAW</u>	67	(SEQ ID NO : 283)
UL-2	<u>CTSVHQ</u> <u>KTRTTQGN</u>	<u>LSLMVTLKDDCPRCRGGCDGYDCCWGDACRSSGLCWGHNPLVTETTYTEF</u>	<u>YIDAW</u>	67	(SEQ ID NO : 284)
UL-3	<u>CTSVHQ</u> <u>KTRTTQGN</u>	<u>TCPDGYTLKDDCPRCRGGCDGYDCCWGDACRSSGLCWGHNPLVTETTYTEF</u>	<u>YIDAW</u>	66*	(SEQ ID NO : 285)
UL-4	<u>CTSVHQ</u> <u>KTRTTQGN</u>	<u>TCPDGYTFKDDCPRCRGGCDGYDCCWGDACRSSGLCWGHNPLVTETTYTEF</u>	<u>YIDAW</u>	66	(SEQ ID NO : 286)
UL-5	<u>CTTVHQ</u> <u>KTRTTQGN</u>	<u>TCPDGYTLKDDCPRCRGGCDGYDCCWGDACRSSGLCWGHNPLVTETTYTEF</u>	<u>YIDAW</u>	66	(SEQ ID NO : 287)
UL-6	<u>CTSVHQ</u> <u>KTRTTQGN</u>	<u>TCPDGYTLKNDCPRCRGGCDGYDCCWGDACRSSGLCWGHNPLVTETTYTEF</u>	<u>YIDAW</u>	66	(SEQ ID NO : 288)
UK-7	<u>CTTVYQ</u> <u>KTRTTQGN</u>	<u>TCPDGYTLKDDCPRCRGGCDGYDCCWGDACRSSGLCWGHNPLVTETTYTEF</u>	<u>YIDAW</u>	66	(SEQ ID NO : 289)
UL-8	<u>CTSVHQ</u> <u>KPGQHKGI</u>	<u>LVLMTLLKDDCPRCRGGCDGYDCCWGDACRSSGLCWGHNPLVTETTYTEF</u>	<u>YIDAW</u>	66	(SEQ ID NO : 290)
UL-9	<u>CTSVHQ</u> <u>KTRTTQGI</u>	<u>LVLMTLLKDDCPRCRGGCDGYDCCWGDACRSSGLCWGHNPLVTETTYTEF</u>	<u>YIDAW</u>	66	(SEQ ID NO : 291)
	12 Cysteines:				
UL-10	<u>CSPVHQ</u> <u>EIRK</u>	<u>CCPAGCQCGRS CGACCGCAGDEFECGINVYGYVTCGGYRTCSCIDTYDF</u>	<u>YVDAW</u>	59	(SEQ ID NO : 292)
UL-11	<u>CSPVHQ</u> <u>QTRK</u>	<u>CCPAGCQCGRS CGACCGCAGDEFECGINVGYITCGGYRTCSCIDTYDF</u>	<u>YVEAW</u>	59	(SEQ ID NO : 293)
	10 Cysteines:				
UL-12	<u>CATVYQ</u> <u>KTNQSK</u>	<u>NCPEGSAWCRSCDGGAGCADYECCRCGWSGCSWRNGACECSSLSSSYTYEL</u>	<u>HVDAW</u>	64	(SEQ ID NO : 294)
UL-13	<u>CTSVHQ</u> <u>TTHQIH</u>	<u>TCPNGWTGGCVCSSRFNCRGNCCCRTAYCSVDRYVCACPTVITYTEF</u>	<u>NVDSW</u>	61	(SEQ ID NO : 295)
UL-14	<u>CTAVYQ</u> <u>KTTSIR</u>	<u>SCPGGTTLRNGCRSACGCNDCDCCGSSWDICYMSKCTSAPEITYTEL</u>	<u>HIDAW</u>	61	(SEQ ID NO : 296)
UL-15	<u>CTNVHQ</u> <u>KTKK</u>	<u>TCPDDYTCGVSCSSSSGCADYGCCSYITYGVPGDCGGCCSKKHYTYEW</u>	<u>NVDAW</u>	61	(SEQ ID NO : 297)
UL-16	<u>CTTVHQ</u> <u>KTKK</u>	<u>LCPNGRTCGCGDCGSGCCTSYCDSEFGCWGGRDTEGSSCTSATYTYEW</u>	<u>GVDAW</u>	59	(SEQ ID NO : 298)
UL-17	<u>CATVHQ</u> <u>HTNKK</u>	<u>RCPDGYEFSAACCCGEGCSGSDCCCNRLRCSWYEIYCSVSPSDTYEF</u>	<u>HVDAW</u>	60	(SEQ ID NO : 299)
UL-18	<u>CTTVHQ</u> <u>HTNKK</u>	<u>RCPDGYRFSAAACCCGEGCSGNECCCNTRLRCSWYEIYCSVSPSDTYEF</u>	<u>HVDAW</u>	60	(SEQ ID NO : 300)
UL-19	<u>CTTVHQ</u> <u>HTNQN</u>	<u>RCPTGYKHSAGCCCGVGCSSGNDCCCNRLRCSWYETYCSLSPTDMYEF</u>	<u>YVDAW</u>	60	(SEQ ID NO : 301)
UL-20	<u>CTSVHQ</u> <u>HTNQN</u>	<u>RCPAGYKHSAGCCCGVGCSSGNDCCCNRLRCSWYETYCSLSPTDMYEF</u>	<u>YVDAW</u>	60	(SEQ ID NO : 302)

Figure 2B

UL-21	<u>C</u> T <u>T</u> V <u>H</u> Q K <u>T</u> N <u>E</u>	R <u>C</u> R <u>V</u> V <u>S</u> D <u>D</u> G <u>E</u> C <u>G</u> D <u>G</u> N <u>S</u> C <u>H</u> R <u>W</u> L <u>C</u> S <u>D</u> Y <u>C</u> S <u>G</u> D <u>C</u> C <u>A</u> C <u>G</u> C <u>R</u> A <u>Y</u> H <u>Y</u> T <u>Y</u> T <u>Y</u> E <u>W</u>	N <u>I</u> D <u>A</u> W	59	(<u>S</u> E <u>Q</u> ID NO : 303)
UL-22	<u>C</u> T <u>T</u> V <u>H</u> Q K <u>T</u> N <u>E</u> R	C <u>R</u> R <u>V</u> V <u>S</u> D <u>D</u> G <u>E</u> C <u>G</u> D <u>G</u> N <u>S</u> C <u>H</u> R <u>W</u> L <u>C</u> S <u>D</u> Y <u>C</u> S <u>G</u> D <u>C</u> C <u>A</u> C <u>G</u> C <u>R</u> A <u>Y</u> H <u>Y</u> T <u>Y</u> T <u>Y</u> D <u>F</u>	R <u>I</u> D <u>V</u> W	59	(<u>S</u> E <u>Q</u> ID NO : 304)
UL-23	<u>C</u> T <u>T</u> V <u>H</u> Q K <u>T</u> N <u>R</u> E <u>R</u>	C <u>P</u> D <u>G</u> Y <u>Y</u> C <u>C</u> R <u>S</u> V <u>S</u> D <u>C</u> C <u>S</u> T <u>R</u> A <u>C</u> V <u>G</u> D <u>S</u> C <u>G</u> W <u>T</u> D <u>F</u> G <u>S</u> T <u>H</u> N <u>V</u> D <u>C</u> S <u>F</u> T <u>Y</u> E <u>F</u>	H <u>V</u> D <u>A</u> W	60	(<u>S</u> E <u>Q</u> ID NO : 305)
UL-24	<u>C</u> T <u>T</u> V <u>H</u> Q Q <u>T</u> R <u>K</u>	S <u>C</u> P <u>D</u> G <u>Y</u> T <u>Y</u> C <u>H</u> D <u>C</u> G <u>Y</u> C <u>C</u> C <u>G</u> A <u>S</u> E <u>F</u> C <u>R</u> D <u>Y</u> G <u>G</u> C <u>S</u> L <u>C</u> G <u>R</u> Y <u>C</u> T <u>S</u> F <u>D</u> I <u>Y</u> T <u>Y</u> E <u>N</u>	Y <u>V</u> E <u>T</u> W	59	(<u>S</u> E <u>Q</u> ID NO : 306)
UL-25	<u>C</u> T <u>T</u> V <u>H</u> Q E <u>T</u> K <u>K</u>	N <u>C</u> P <u>D</u> N <u>C</u> Y <u>E</u> N <u>S</u> C <u>G</u> D <u>Y</u> G <u>S</u> G <u>C</u> N <u>G</u> G <u>D</u> C <u>C</u> R <u>C</u> G <u>T</u> W <u>L</u> F <u>C</u> S <u>V</u> S <u>G</u> C <u>T</u> C <u>I</u> R <u>A</u> T <u>N</u> T <u>Y</u> Q <u>W</u>	Y <u>V</u> N <u>A</u> W	60	(<u>S</u> E <u>Q</u> ID NO : 307)
UL-26	<u>C</u> T <u>T</u> V <u>H</u> Q S <u>T</u> N <u>K</u> K	S <u>C</u> P <u>D</u> R <u>V</u> C <u>W</u> A <u>V</u> G <u>C</u> F <u>G</u> E <u>D</u> C <u>T</u> S <u>S</u> D <u>C</u> T <u>C</u> Y <u>A</u> S <u>P</u> G <u>N</u> P <u>Y</u> R <u>H</u> D <u>C</u> G <u>N</u> C <u>D</u> C <u>R</u> S <u>S</u> Y <u>E</u> H	H <u>V</u> D <u>A</u> W	60	(<u>S</u> E <u>Q</u> ID NO : 308)
UL-27	<u>C</u> T <u>T</u> V <u>R</u> Q E <u>T</u> L <u>I</u>	R <u>C</u> R <u>D</u> G <u>P</u> S <u>C</u> A <u>A</u> C <u>C</u> R <u>S</u> G <u>R</u> R <u>C</u> S <u>G</u> Y <u>G</u> C <u>T</u> D <u>G</u> C <u>C</u> S <u>D</u> N <u>D</u> Y <u>A</u> D <u>C</u> I <u>R</u> G <u>E</u> F <u>V</u> D <u>V</u> Y <u>E</u> W	N <u>V</u> D <u>A</u> W	59	(<u>S</u> E <u>Q</u> ID NO : 309)
UL-28	<u>C</u> S <u>T</u> V <u>Y</u> Q K <u>T</u> R <u>T</u>	T <u>C</u> P <u>D</u> G <u>Y</u> T <u>C</u> G <u>D</u> G <u>A</u> R <u>C</u> E <u>K</u> A <u>C</u> R <u>G</u> C <u>D</u> C <u>C</u> R <u>T</u> T <u>V</u> C <u>D</u> T <u>V</u> W <u>S</u> S <u>Y</u> C <u>S</u> C <u>S</u> F <u>T</u> D <u>S</u> Y <u>E</u> F	Y <u>V</u> D <u>A</u> W	59	(<u>S</u> E <u>Q</u> ID NO : 310)
UL-29	<u>C</u> A <u>T</u> V <u>Y</u> Q K <u>T</u> N <u>R</u> E <u>M</u>	S <u>C</u> P <u>D</u> G <u>C</u> R <u>I</u> H <u>N</u> A <u>R</u> L <u>C</u> L <u>S</u> G <u>C</u> S <u>G</u> S <u>D</u> C <u>C</u> S <u>C</u> G <u>D</u> C <u>V</u> S <u>D</u> A <u>R</u> C <u>Y</u> N <u>C</u> R <u>S</u> A <u>V</u> T <u>Y</u> T <u>Y</u> E <u>F</u>	H <u>V</u> D <u>A</u> W	62	(<u>S</u> E <u>Q</u> ID NO : 311)
UL-30	<u>C</u> T <u>I</u> V <u>H</u> Q E <u>T</u> K <u>R</u>	S <u>C</u> P <u>D</u> G <u>Y</u> N <u>T</u> G <u>T</u> R <u>C</u> F <u>G</u> S <u>C</u> G <u>C</u> I <u>G</u> S <u>N</u> C <u>C</u> R <u>S</u> T <u>T</u> S <u>C</u> C <u>A</u> G <u>I</u> Y <u>S</u> Q <u>C</u> T <u>T</u> S <u>T</u> L <u>T</u> Y <u>E</u> W	H <u>A</u> D <u>V</u> W	59	(<u>S</u> E <u>Q</u> ID NO : 312)
UL-31	<u>C</u> A <u>I</u> V <u>Y</u> Q R <u>T</u> R <u>Q</u>	R <u>C</u> P <u>D</u> G <u>Y</u> N <u>T</u> G <u>T</u> R <u>C</u> F <u>G</u> T <u>C</u> G <u>C</u> N <u>G</u> S <u>N</u> C <u>C</u> R <u>F</u> T <u>T</u> S <u>C</u> C <u>A</u> G <u>Y</u> S <u>Q</u> C <u>T</u> T <u>S</u> T <u>L</u> Y <u>E</u> W	H <u>A</u> D <u>V</u> W	59	(<u>S</u> E <u>Q</u> ID NO : 313)
UL-32	<u>C</u> T <u>T</u> V <u>H</u> Q K <u>T</u> E <u>T</u>	R <u>C</u> P <u>D</u> G <u>Y</u> S <u>S</u> T <u>N</u> G <u>D</u> A <u>R</u> C <u>G</u> C <u>S</u> D <u>C</u> D <u>C</u> C <u>N</u> V <u>G</u> R <u>W</u> G <u>C</u> P <u>L</u> I <u>C</u> S <u>R</u> N <u>C</u> R <u>S</u> F <u>T</u> Y <u>E</u> W	Y <u>A</u> D <u>A</u> W	59	(<u>S</u> E <u>Q</u> ID NO : 314)
UL-33	<u>C</u> T <u>T</u> V <u>H</u> Q K <u>T</u> N <u>K</u> K <u>E</u>	S <u>C</u> P <u>D</u> G <u>Y</u> T <u>M</u> N <u>E</u> C <u>C</u> G <u>C</u> G <u>Y</u> G <u>C</u> C <u>R</u> G <u>G</u> C <u>V</u> C <u>S</u> A <u>Y</u> C <u>S</u> R <u>P</u> N <u>C</u> W <u>R</u> E <u>L</u> T <u>Y</u> T <u>Y</u> E <u>F</u>	Y <u>V</u> D <u>T</u> W	59	(<u>S</u> E <u>Q</u> ID NO : 315)
UL-34	<u>C</u> T <u>T</u> V <u>Y</u> Q K <u>S</u> R <u>K</u> E <u>S</u>	S <u>C</u> P <u>N</u> G <u>W</u> I <u>Y</u> G <u>K</u> D <u>C</u> S <u>W</u> S <u>Y</u> C <u>T</u> D <u>C</u> D <u>C</u> L <u>C</u> G <u>D</u> L <u>H</u> C <u>Y</u> D <u>G</u> C <u>S</u> S <u>F</u> G <u>V</u> T <u>W</u> T <u>Y</u> E <u>F</u>	H <u>V</u> D <u>A</u> W	59	(<u>S</u> E <u>Q</u> ID NO : 316)
UL-35	<u>C</u> T <u>T</u> V <u>F</u> Q E <u>T</u> R <u>K</u>	S <u>C</u> P <u>T</u> G <u>F</u> Y <u>V</u> D <u>G</u> S <u>T</u> C <u>G</u> C <u>A</u> T <u>Y</u> C <u>R</u> T <u>C</u> D <u>C</u> C <u>G</u> G <u>Y</u> C <u>S</u> G <u>G</u> A <u>D</u> C <u>H</u> C <u>V</u> S <u>Y</u> N <u>T</u> Y <u>S</u> W	H <u>V</u> D <u>A</u> W	58	(<u>S</u> E <u>Q</u> ID NO : 317)
UL-36	<u>C</u> A <u>A</u> V <u>F</u> Q E <u>T</u> R <u>T</u>	N <u>C</u> P <u>S</u> G <u>Y</u> G <u>N</u> A <u>F</u> S <u>C</u> G <u>C</u> P <u>I</u> A <u>C</u> R <u>D</u> C <u>D</u> C <u>C</u> G <u>G</u> Y <u>W</u> C <u>S</u> G <u>G</u> A <u>D</u> C <u>H</u> C <u>V</u> S <u>Y</u> N <u>T</u> Y <u>S</u> W	H <u>V</u> D <u>A</u> W	57	(<u>S</u> E <u>Q</u> ID NO : 318)
UL-37	<u>C</u> A <u>T</u> V <u>Y</u> Q K <u>T</u> E <u>K</u>	H <u>C</u> P <u>L</u> F <u>H</u> S <u>I</u> C <u>H</u> C <u>G</u> E <u>V</u> G <u>S</u> G <u>G</u> D <u>C</u> C <u>G</u> C <u>E</u> R <u>R</u> S <u>G</u> C <u>V</u> C <u>T</u> M <u>R</u> N <u>S</u> Y <u>T</u> N <u>Y</u> Q <u>F</u>	H <u>V</u> D <u>A</u> W	58	(<u>S</u> E <u>Q</u> ID NO : 319)
UL-38	<u>C</u> G <u>T</u> V <u>H</u> Q K <u>T</u> I <u>K</u> E	L <u>C</u> P <u>D</u> D <u>S</u> T <u>Y</u> C <u>C</u> G <u>C</u> V <u>S</u> G <u>A</u> C <u>C</u> T <u>Y</u> G <u>C</u> D <u>V</u> G <u>C</u> C <u>R</u> V <u>S</u> L <u>W</u> T <u>T</u> Y <u>I</u> K <u>D</u> I <u>V</u> G <u>V</u> S <u>Y</u> E <u>W</u>	H <u>V</u> D <u>A</u> W	59	(<u>S</u> E <u>Q</u> ID NO : 320)
UL-39	<u>C</u> A <u>S</u> V <u>H</u> Q H <u>T</u> E <u>P</u>	T <u>C</u> P <u>A</u> G <u>Y</u> T <u>Y</u> C <u>C</u> G <u>C</u> L <u>Y</u> K <u>C</u> N <u>C</u> G <u>D</u> C <u>G</u> C <u>Y</u> N <u>V</u> G <u>C</u> S <u>G</u> W <u>L</u> G <u>K</u> A <u>C</u> D <u>Y</u> R <u>E</u> T <u>Y</u> E <u>W</u>	Y <u>V</u> D <u>A</u> W	57	(<u>S</u> E <u>Q</u> ID NO : 321)
UL-40	<u>C</u> A <u>S</u> V <u>H</u> Q H <u>T</u> E <u>P</u>	T <u>C</u> P <u>A</u> G <u>Y</u> T <u>Y</u> C <u>C</u> G <u>C</u> L <u>Y</u> K <u>C</u> N <u>C</u> G <u>D</u> C <u>G</u> C <u>Y</u> N <u>A</u> G <u>C</u> S <u>G</u> W <u>L</u> G <u>K</u> A <u>C</u> D <u>Y</u> R <u>E</u> T <u>Y</u> E <u>W</u>	Y <u>V</u> D <u>A</u> W	57	(<u>S</u> E <u>Q</u> ID NO : 322)
UL-41	<u>C</u> T <u>T</u> V <u>F</u> Q E <u>T</u> R <u>K</u>	S <u>C</u> P <u>S</u> G <u>F</u> R <u>D</u> R <u>D</u> A <u>C</u> G <u>C</u> A <u>V</u> T <u>C</u> R <u>N</u> C <u>D</u> C <u>C</u> G <u>G</u> G <u>P</u> C <u>N</u> G <u>G</u> S <u>C</u> R <u>C</u> N <u>N</u> Y <u>I</u> Y <u>K</u> Y <u>S</u> F	H <u>V</u> D <u>A</u> W	57	(<u>S</u> E <u>Q</u> ID NO : 323)
UL-42	<u>C</u> T <u>A</u> V <u>F</u> Q E <u>T</u> R <u>K</u>	D <u>C</u> P <u>S</u> G <u>Y</u> G <u>S</u> A <u>F</u> T <u>C</u> G <u>C</u> L <u>A</u> A <u>C</u> H <u>G</u> C <u>D</u> C <u>C</u> G <u>G</u> W <u>C</u> S <u>G</u> G <u>D</u> C <u>R</u> C <u>R</u> S <u>Y</u> S <u>T</u> A <u>Y</u> S <u>F</u>	H <u>I</u> D <u>A</u> W	57	(<u>S</u> E <u>Q</u> ID NO : 324)
UL-43	<u>C</u> A <u>T</u> V <u>F</u> Q E <u>T</u> R <u>K</u>	S <u>C</u> P <u>S</u> G <u>Y</u> A <u>D</u> R <u>F</u> T <u>C</u> D <u>C</u> V <u>Y</u> Y <u>Q</u> T <u>C</u> D <u>C</u> C <u>G</u> G <u>N</u> R <u>C</u> S <u>G</u> G <u>G</u> P <u>C</u> R <u>C</u> S <u>S</u> I <u>N</u> Y <u>S</u> F	H <u>V</u> D <u>T</u> W	57	(<u>S</u> E <u>Q</u> ID NO : 325)
UL-44	<u>C</u> A <u>A</u> A <u>H</u> Q E <u>T</u> K <u>K</u>	S <u>C</u> P <u>D</u> G <u>T</u> C <u>R</u> Q <u>C</u> C <u>G</u> G <u>V</u> C <u>R</u> C <u>H</u> A <u>S</u> G <u>C</u> C <u>Y</u> W <u>C</u> T <u>T</u> G <u>C</u> V <u>G</u> R <u>A</u> L <u>S</u> E <u>S</u> H <u>S</u> Y <u>E</u> F	H <u>V</u> D <u>T</u> W	54	(<u>S</u> E <u>Q</u> ID NO : 326)
8 Cysteines:					
UL-45	<u>C</u> S <u>T</u> V <u>H</u> Q K <u>T</u> R <u>T</u> Q <u>G</u> N	T <u>C</u> P <u>D</u> G <u>Y</u> T <u>L</u> K <u>D</u> D <u>C</u> P <u>R</u> C <u>R</u> G <u>G</u> C <u>D</u> G <u>Y</u> D <u>C</u> C <u>W</u> G <u>D</u> A <u>C</u> R <u>S</u> S <u>G</u> L <u>C</u> W <u>G</u> H <u>N</u> P <u>L</u> V <u>T</u> E <u>T</u> Y <u>T</u> Y <u>E</u> F	Y <u>I</u> D <u>A</u> W	66 (167)	(<u>S</u> E <u>Q</u> ID NO : 327)
UL-46	<u>C</u> V <u>V</u> V <u>Y</u> Q K <u>T</u> I <u>N</u> S <u>Q</u> K	S <u>C</u> P <u>R</u> G <u>Y</u> T <u>E</u> R <u>E</u> T <u>C</u> N <u>R</u> R <u>Y</u> G <u>W</u> C <u>G</u> R <u>Y</u> D <u>C</u> C <u>D</u> C <u>D</u> R <u>W</u> V <u>S</u> G <u>N</u> C <u>A</u> N <u>I</u> C <u>T</u> D <u>Y</u> T <u>D</u> T <u>H</u> T <u>Y</u> E <u>F</u>	H <u>A</u> D <u>A</u> W	64 (23)	(<u>S</u> E <u>Q</u> ID NO : 328)
UL-47	<u>C</u> G <u>T</u> V <u>F</u> Q Q <u>T</u> H <u>K</u> V <u>R</u>	D <u>C</u> P <u>D</u> G <u>F</u> T <u>A</u> A <u>P</u> R <u>C</u> G <u>G</u> E <u>C</u> C <u>C</u> S <u>N</u> V <u>N</u> S <u>R</u> S <u>G</u> W <u>C</u> R <u>Y</u> C <u>G</u> R <u>D</u> C <u>T</u> A <u>P</u> E <u>T</u> S <u>T</u> Y <u>E</u> F	H <u>V</u> D <u>A</u> W	60 (29)	(<u>S</u> E <u>Q</u> ID NO : 329)
UL-48	<u>C</u> T <u>A</u> V <u>Y</u> Q R <u>T</u> G <u>Q</u>	K <u>C</u> P <u>E</u> G <u>C</u> E <u>S</u> R <u>N</u> T <u>C</u> L <u>Y</u> S <u>R</u> N <u>C</u> G <u>D</u> Y <u>T</u> C <u>C</u> G <u>S</u> R <u>A</u> S <u>G</u> S <u>G</u> A <u>C</u> G <u>W</u> N <u>S</u> V <u>D</u> C <u>K</u> N <u>K</u> Y <u>E</u> H	H <u>V</u> D <u>A</u> W	59 (30)	(<u>S</u> E <u>Q</u> ID NO : 330)
UL-49	<u>C</u> T <u>T</u> V <u>Y</u> Q K <u>T</u> K <u>Q</u>	N <u>C</u> P <u>D</u> G <u>Y</u> D <u>F</u> R <u>D</u> T <u>C</u> G <u>S</u> Q <u>S</u> Y <u>C</u> S <u>G</u> Y <u>D</u> C <u>C</u> R <u>C</u> S <u>R</u> F <u>G</u> G <u>C</u> S <u>I</u> G <u>T</u> C <u>I</u> S <u>Y</u> S <u>D</u> A <u>Y</u> T <u>Y</u> E <u>W</u>	Y <u>V</u> D <u>A</u> W	59 (16)	(<u>S</u> E <u>Q</u> ID NO : 331)
UL-50	<u>C</u> T <u>T</u> V <u>H</u> Q Q <u>T</u> H <u>E</u> K <u>R</u>	S <u>C</u> P <u>E</u> S <u>Y</u> S <u>Y</u> S <u>C</u> S <u>C</u> A <u>S</u> G <u>V</u> V <u>G</u> C <u>G</u> P <u>D</u> D <u>C</u> C <u>T</u> Y <u>R</u> I <u>S</u> I <u>R</u> G <u>Y</u> T <u>C</u> S <u>S</u> L <u>S</u> N <u>S</u> Y <u>E</u> W	Y <u>V</u> D <u>A</u> W	59 (15)	(<u>S</u> E <u>Q</u> ID NO : 332)
UL-51	<u>C</u> T <u>A</u> V <u>H</u> Q Q <u>T</u> R <u>K</u> S	G <u>C</u> P <u>D</u> G <u>Y</u> S <u>D</u> E <u>S</u> C <u>S</u> Y <u>C</u> G <u>S</u> W <u>C</u> P <u>V</u> Y <u>W</u> C <u>G</u> S <u>P</u> C <u>S</u> Y <u>R</u> C <u>L</u> R <u>H</u> T <u>D</u> T <u>Y</u> S <u>Y</u> E <u>H</u>	H <u>V</u> D <u>A</u> W	57 (12)	(<u>S</u> E <u>Q</u> ID NO : 333)
UL-52	<u>C</u> A <u>T</u> V <u>Y</u> Q E <u>T</u> K <u>R</u>	T <u>C</u> A <u>G</u> H <u>S</u> V <u>E</u> C <u>D</u> S <u>P</u> Y <u>D</u> C <u>N</u> C <u>R</u> G <u>G</u> D <u>C</u> C <u>R</u> S <u>P</u> I <u>F</u> N <u>D</u> C <u>W</u> A <u>A</u> S <u>C</u> S <u>A</u> T <u>K</u> T <u>Y</u> E <u>W</u>	H <u>V</u> E <u>S</u> W	56 (16)	(<u>S</u> E <u>Q</u> ID NO : 334)
UL-53	<u>C</u> I <u>T</u> V <u>H</u> Q E <u>T</u> Q <u>K</u>	S <u>C</u> P <u>D</u> D <u>Y</u> T <u>Y</u> G <u>D</u> G <u>T</u> C <u>A</u> Y <u>V</u> C <u>S</u> I <u>D</u> K <u>C</u> C <u>C</u> G <u>R</u> T <u>W</u> L <u>S</u> S <u>G</u> C <u>L</u> P <u>C</u> R <u>Y</u> T <u>Y</u> N <u>L</u>	H <u>V</u> D <u>A</u> W	54 (155)	(<u>S</u> E <u>Q</u> ID NO : 335)
UL-54	<u>C</u> I <u>T</u> V <u>H</u> Q E <u>T</u> Q <u>K</u>	S <u>C</u> P <u>D</u> D <u>Y</u> T <u>Y</u> G <u>D</u> A <u>T</u> C <u>A</u> Y <u>V</u> C <u>S</u> T <u>D</u> E <u>C</u> C <u>C</u> G <u>R</u> T <u>W</u> L <u>S</u> A <u>G</u> C <u>R</u> P <u>C</u> R <u>Y</u> T <u>Y</u> N <u>L</u>	H <u>V</u> D <u>A</u> W	54 (61)	(<u>S</u> E <u>Q</u> ID NO : 336)
UL-55	<u>C</u> I <u>T</u> V <u>H</u> Q E <u>T</u> Q <u>K</u>	S <u>C</u> F <u>D</u> D <u>Y</u> T <u>Y</u> G <u>D</u> A <u>S</u> C <u>A</u> Y <u>V</u> C <u>S</u> T <u>D</u> E <u>C</u> C <u>C</u> G <u>R</u> T <u>W</u> L <u>S</u> A <u>G</u> C <u>R</u> P <u>C</u> R <u>Y</u> T <u>Y</u> N <u>L</u>	H <u>V</u> D <u>A</u> W	54 (20)	(<u>S</u> E <u>Q</u> ID NO : 337)
UL-56	<u>C</u> I <u>T</u> A <u>H</u> Q E <u>T</u> Q <u>K</u>	S <u>C</u> S <u>D</u> D <u>Y</u> T <u>Y</u> G <u>D</u> A <u>T</u> C <u>A</u> Y <u>V</u> C <u>S</u> T <u>D</u> E <u>C</u> C <u>C</u> G <u>R</u> T <u>W</u> L <u>S</u> A <u>G</u> C <u>R</u> P <u>C</u> R <u>Y</u> T <u>Y</u> N <u>L</u>	H <u>V</u> D <u>A</u> W	54 (20)	(<u>S</u> E <u>Q</u> ID NO : 338)
UL-57	<u>C</u> I <u>T</u> V <u>H</u> Q E <u>T</u> Q <u>K</u>	S <u>C</u> P <u>D</u> D <u>Y</u> T <u>Y</u> G <u>D</u> G <u>T</u> C <u>A</u> Y <u>V</u> C <u>S</u> I <u>D</u> N <u>C</u> C <u>C</u> G <u>R</u> T <u>W</u> L <u>S</u> S <u>G</u> C <u>L</u> P <u>C</u> R <u>Y</u> T <u>Y</u> N <u>L</u>	H <u>V</u> D <u>A</u> W	54 (10)	(<u>S</u> E <u>Q</u> ID NO : 339)

Figure 2C

Other Representative sequences:				
UL-58	<u>C</u> <u>V</u> <u>T</u> <u>V</u> <u>H</u> <u>Q</u> <u>Q</u> <u>T</u> <u>H</u> <u>A</u> <u>T</u> <u>R</u>	<u>R</u> <u>C</u> <u>P</u> <u>D</u> <u>G</u> <u>Y</u> <u>G</u> <u>D</u> <u>S</u> <u>Y</u> <u>A</u> <u>C</u> <u>K</u> <u>S</u> <u>N</u> <u>Y</u> <u>G</u> <u>S</u> <u>A</u> <u>E</u> <u>G</u> <u>C</u> <u>C</u> <u>R</u> <u>W</u> <u>G</u> <u>P</u> <u>G</u> <u>S</u> <u>G</u> <u>A</u> <u>C</u> <u>T</u> <u>G</u> <u>A</u> <u>I</u> <u>T</u> <u>S</u> <u>P</u> <u>Y</u> <u>E</u> <u>W</u>	YVDAM	(SEQ ID NO : 340)
UL-59	<u>C</u> <u>A</u> <u>A</u> <u>V</u> <u>H</u> <u>Q</u> <u>R</u> <u>T</u> <u>E</u> <u>G</u> <u>Q</u> <u>Q</u>	<u>S</u> <u>C</u> <u>P</u> <u>D</u> <u>G</u> <u>Y</u> <u>L</u> <u>E</u> <u>T</u> <u>R</u> <u>V</u> <u>C</u> <u>P</u> <u>Y</u> <u>R</u> <u>M</u> <u>Y</u> <u>R</u> <u>C</u> <u>I</u> <u>G</u> <u>W</u> <u>D</u> <u>C</u> <u>C</u> <u>R</u> <u>C</u> <u>S</u> <u>D</u> <u>G</u> <u>S</u> <u>R</u> <u>D</u> <u>N</u> <u>Y</u> <u>I</u> <u>M</u> <u>T</u> <u>Y</u> <u>S</u> <u>Y</u> <u>E</u> <u>F</u>	HVDVW	(SEQ ID NO : 341)
UL-60	<u>C</u> <u>T</u> <u>T</u> <u>V</u> <u>Y</u> <u>Q</u> <u>E</u> <u>T</u> <u>K</u> <u>T</u> <u>K</u> <u>S</u>	<u>G</u> <u>C</u> <u>P</u> <u>D</u> <u>G</u> <u>Y</u> <u>S</u> <u>C</u> <u>C</u> <u>Y</u> <u>N</u> <u>G</u> <u>R</u> <u>S</u> <u>R</u> <u>S</u> <u>C</u> <u>R</u> <u>P</u> <u>N</u> <u>D</u> <u>C</u> <u>S</u> <u>T</u> <u>Y</u> <u>G</u> <u>E</u> <u>V</u> <u>R</u> <u>S</u> <u>L</u> <u>S</u> <u>R</u> <u>S</u> <u>C</u> <u>T</u> <u>Y</u> <u>N</u> <u>Y</u> <u>E</u> <u>F</u>	YVDAM	(SEQ ID NO : 342)
UL-61	<u>C</u> <u>G</u> <u>T</u> <u>V</u> <u>Y</u> <u>Q</u> <u>H</u> <u>T</u> <u>K</u> <u>E</u> <u>I</u> <u>K</u>	<u>T</u> <u>C</u> <u>P</u> <u>D</u> <u>G</u> <u>Y</u> <u>S</u> <u>D</u> <u>V</u> <u>F</u> <u>T</u> <u>Y</u> <u>C</u> <u>P</u> <u>V</u> <u>T</u> <u>C</u> <u>P</u> <u>G</u> <u>W</u> <u>D</u> <u>C</u> <u>C</u> <u>R</u> <u>R</u> <u>N</u> <u>D</u> <u>C</u> <u>G</u> <u>R</u> <u>T</u> <u>R</u> <u>Y</u> <u>T</u> <u>V</u> <u>A</u> <u>Y</u> <u>S</u> <u>Y</u> <u>A</u> <u>L</u>	HVDVW	(SEQ ID NO : 343)
UL-62	<u>C</u> <u>T</u> <u>T</u> <u>V</u> <u>L</u> <u>Q</u> <u>E</u> <u>T</u> <u>H</u> <u>Q</u> <u>Q</u> <u>R</u>	<u>G</u> <u>C</u> <u>P</u> <u>A</u> <u>G</u> <u>Y</u> <u>Q</u> <u>V</u> <u>V</u> <u>D</u> <u>G</u> <u>C</u> <u>C</u> <u>R</u> <u>T</u> <u>S</u> <u>Y</u> <u>V</u> <u>C</u> <u>G</u> <u>P</u> <u>L</u> <u>T</u> <u>C</u> <u>T</u> <u>S</u> <u>N</u> <u>T</u> <u>A</u> <u>T</u> <u>R</u> <u>N</u> <u>Y</u> <u>Q</u> <u>W</u>	YVDAM	(SEQ ID NO : 344)
UL-63	<u>C</u> <u>S</u> <u>T</u> <u>V</u> <u>Y</u> <u>Q</u> <u>K</u> <u>T</u> <u>E</u> <u>K</u>	<u>K</u> <u>C</u> <u>P</u> <u>D</u> <u>G</u> <u>Y</u> <u>T</u> <u>D</u> <u>R</u> <u>R</u> <u>D</u> <u>E</u> <u>C</u> <u>P</u> <u>N</u> <u>T</u> <u>C</u> <u>K</u> <u>N</u> <u>F</u> <u>D</u> <u>C</u> <u>E</u> <u>N</u> <u>E</u> <u>G</u> <u>G</u> <u>L</u> <u>R</u> <u>C</u> <u>L</u> <u>C</u> <u>S</u> <u>A</u> <u>I</u> <u>S</u> <u>A</u> <u>Y</u> <u>E</u> <u>F</u>	HVDAW	(SEQ ID NO : 345)
UL-64	<u>C</u> <u>T</u> <u>T</u> <u>H</u> <u>Q</u> <u>R</u> <u>T</u> <u>Q</u> <u>K</u>	<u>S</u> <u>C</u> <u>P</u> <u>D</u> <u>Y</u> <u>A</u> <u>S</u> <u>I</u> <u>D</u> <u>C</u> <u>G</u> <u>S</u> <u>P</u> <u>D</u> <u>D</u> <u>E</u> <u>E</u> <u>C</u> <u>S</u> <u>S</u> <u>C</u> <u>R</u> <u>S</u> <u>C</u> <u>T</u> <u>R</u> <u>W</u> <u>C</u> <u>A</u> <u>P</u> <u>T</u> <u>A</u> <u>P</u> <u>I</u> <u>Y</u> <u>T</u> <u>Y</u> <u>Q</u> <u>F</u>	YIDAW	(SEQ ID NO : 346)
UL-65	<u>C</u> <u>T</u> <u>T</u> <u>V</u> <u>H</u> <u>Q</u> <u>Q</u> <u>T</u> <u>N</u> <u>K</u>	<u>R</u> <u>C</u> <u>P</u> <u>T</u> <u>G</u> <u>Y</u> <u>N</u> <u>S</u> <u>G</u> <u>T</u> <u>L</u> <u>C</u> <u>N</u> <u>M</u> <u>I</u> <u>G</u> <u>C</u> <u>S</u> <u>G</u> <u>D</u> <u>E</u> <u>C</u> <u>C</u> <u>N</u> <u>Y</u> <u>G</u> <u>R</u> <u>V</u> <u>E</u> <u>C</u> <u>T</u> <u>S</u> <u>Y</u> <u>V</u> <u>W</u> <u>T</u> <u>H</u> <u>N</u> <u>F</u>	YVDAM	(SEQ ID NO : 347)
UL-66	<u>C</u> <u>T</u> <u>T</u> <u>V</u> <u>H</u> <u>Q</u> <u>E</u> <u>T</u> <u>Q</u> <u>R</u> <u>T</u>	<u>S</u> <u>C</u> <u>P</u> <u>S</u> <u>G</u> <u>W</u> <u>T</u> <u>Y</u> <u>T</u> <u>C</u> <u>N</u> <u>C</u> <u>R</u> <u>N</u> <u>G</u> <u>C</u> <u>G</u> <u>C</u> <u>Y</u> <u>R</u> <u>P</u> <u>S</u> <u>Q</u> <u>L</u> <u>C</u> <u>G</u> <u>A</u> <u>Y</u> <u>V</u> <u>A</u> <u>V</u> <u>T</u> <u>H</u> <u>T</u> <u>Y</u> <u>E</u> <u>F</u>	HVDAW	(SEQ ID NO : 348)
UL-67	<u>C</u> <u>A</u> <u>T</u> <u>V</u> <u>H</u> <u>Q</u> <u>K</u> <u>D</u> <u>K</u>	<u>H</u> <u>C</u> <u>P</u> <u>A</u> <u>G</u> <u>Y</u> <u>R</u> <u>S</u> <u>G</u> <u>T</u> <u>L</u> <u>C</u> <u>R</u> <u>M</u> <u>I</u> <u>G</u> <u>C</u> <u>T</u> <u>G</u> <u>D</u> <u>D</u> <u>C</u> <u>C</u> <u>N</u> <u>Y</u> <u>D</u> <u>R</u> <u>V</u> <u>E</u> <u>C</u> <u>T</u> <u>N</u> <u>D</u> <u>Y</u> <u>T</u> <u>N</u> <u>N</u> <u>F</u>	YVDAM	(SEQ ID NO : 349)
UL-68	<u>C</u> <u>T</u> <u>A</u> <u>V</u> <u>H</u> <u>Q</u> <u>Q</u> <u>T</u> <u>T</u> <u>E</u> <u>K</u> <u>G</u> <u>K</u>	<u>T</u> <u>C</u> <u>P</u> <u>P</u> <u>R</u> <u>S</u> <u>R</u> <u>D</u> <u>M</u> <u>G</u> <u>T</u> <u>R</u> <u>C</u> <u>R</u> <u>D</u> <u>D</u> <u>R</u> <u>Y</u> <u>Y</u> <u>P</u> <u>W</u> <u>R</u> <u>Y</u> <u>S</u> <u>D</u> <u>Y</u> <u>T</u> <u>Y</u> <u>T</u> <u>Y</u> <u>E</u> <u>W</u>	HVDAW	(SEQ ID NO : 350)
UL-69	<u>C</u> <u>T</u> <u>S</u> <u>V</u> <u>H</u> <u>Q</u> <u>K</u> <u>T</u> <u>D</u> <u>V</u>	<u>T</u> <u>C</u> <u>P</u> <u>S</u> <u>G</u> <u>A</u> <u>T</u> <u>Y</u> <u>R</u> <u>C</u> <u>D</u> <u>C</u> <u>G</u> <u>G</u> <u>R</u> <u>G</u> <u>C</u> <u>G</u> <u>C</u> <u>Y</u> <u>D</u> <u>P</u> <u>W</u> <u>C</u> <u>S</u> <u>T</u> <u>T</u> <u>Y</u> <u>R</u> <u>G</u> <u>T</u> <u>Y</u> <u>T</u> <u>Y</u> <u>D</u> <u>F</u>	HVETW	(SEQ ID NO : 351)
UL-70	<u>C</u> <u>G</u> <u>T</u> <u>V</u> <u>H</u> <u>Q</u> <u>E</u> <u>T</u> <u>H</u> <u>T</u> <u>Q</u> <u>R</u>	<u>T</u> <u>C</u> <u>P</u> <u>D</u> <u>A</u> <u>C</u> <u>D</u> <u>V</u> <u>T</u> <u>G</u> <u>D</u> <u>N</u> <u>C</u> <u>K</u> <u>V</u> <u>R</u> <u>R</u> <u>N</u> <u>G</u> <u>D</u> <u>W</u> <u>C</u> <u>G</u> <u>R</u> <u>A</u> <u>S</u> <u>K</u> <u>T</u> <u>D</u> <u>T</u> <u>Y</u> <u>D</u> <u>F</u>	YVDAM	(SEQ ID NO : 352)
UL-71	<u>C</u> <u>T</u> <u>T</u> <u>D</u> <u>Y</u> <u>Q</u> <u>K</u> <u>T</u> <u>E</u> <u>K</u>	<u>S</u> <u>C</u> <u>P</u> <u>E</u> <u>N</u> <u>Y</u> <u>A</u> <u>E</u> <u>T</u> <u>G</u> <u>Y</u> <u>C</u> <u>M</u> <u>C</u> <u>G</u> <u>S</u> <u>W</u> <u>R</u> <u>C</u> <u>G</u> <u>Y</u> <u>G</u> <u>S</u> <u>T</u> <u>S</u> <u>L</u> <u>I</u> <u>V</u> <u>S</u> <u>Y</u> <u>K</u> <u>W</u>	YVDAM	(SEQ ID NO : 353)
UL-72	<u>C</u> <u>T</u> <u>T</u> <u>V</u> <u>H</u> <u>Q</u> <u>K</u> <u>T</u> <u>N</u> <u>Q</u> <u>K</u> <u>W</u>	<u>G</u> <u>C</u> <u>P</u> <u>D</u> <u>G</u> <u>Y</u> <u>V</u> <u>H</u> <u>M</u> <u>S</u> <u>G</u> <u>S</u> <u>C</u> <u>C</u> <u>R</u> <u>G</u> <u>S</u> <u>I</u> <u>C</u> <u>T</u> <u>N</u> <u>G</u> <u>L</u> <u>F</u> <u>R</u> <u>N</u> <u>T</u> <u>Y</u> <u>T</u> <u>Y</u> <u>E</u> <u>F</u>	NVEAW	(SEQ ID NO : 354)
UL-73	<u>C</u> <u>T</u> <u>T</u> <u>V</u> <u>Y</u> <u>Q</u> <u>E</u> <u>T</u> <u>R</u> <u>T</u>	<u>N</u> <u>C</u> <u>P</u> <u>D</u> <u>G</u> <u>Y</u> <u>N</u> <u>R</u> <u>S</u> <u>G</u> <u>D</u> <u>C</u> <u>R</u> <u>R</u> <u>N</u> <u>H</u> <u>W</u> <u>L</u> <u>G</u> <u>E</u> <u>Q</u> <u>R</u> <u>V</u> <u>S</u> <u>P</u> <u>T</u> <u>Y</u> <u>N</u> <u>Y</u> <u>E</u> <u>W</u>	YVDSW	(SEQ ID NO : 355)
UL-74	<u>C</u> <u>T</u> <u>T</u> <u>V</u> <u>Y</u> <u>Q</u> <u>K</u> <u>T</u> <u>T</u> <u>T</u> <u>K</u>	<u>S</u> <u>C</u> <u>P</u> <u>G</u> <u>G</u> <u>F</u> <u>D</u> <u>N</u> <u>G</u> <u>R</u> <u>R</u> <u>C</u> <u>I</u> <u>M</u> <u>G</u> <u>L</u> <u>G</u> <u>D</u> <u>L</u> <u>R</u> <u>D</u> <u>Y</u> <u>T</u> <u>F</u> <u>N</u> <u>K</u> <u>Y</u> <u>E</u> <u>W</u>	YVETW	(SEQ ID NO : 356)
UL-75	<u>C</u> <u>S</u> <u>T</u> <u>V</u> <u>H</u> <u>Q</u> <u>K</u> <u>T</u> <u>E</u> <u>Q</u>	<u>R</u> <u>C</u> <u>L</u> <u>D</u> <u>G</u> <u>Y</u> <u>D</u> <u>D</u> <u>R</u> <u>G</u> <u>A</u> <u>Y</u> <u>C</u> <u>Y</u> <u>D</u> <u>S</u> <u>V</u> <u>R</u> <u>G</u> <u>L</u> <u>M</u> <u>S</u> <u>W</u> <u>T</u> <u>Y</u> <u>K</u> <u>Y</u> <u>V</u> <u>Y</u> <u>E</u> <u>W</u>	RVDTW	(SEQ ID NO : 357)
UL-76	<u>C</u> <u>T</u> <u>N</u> <u>V</u> <u>H</u> <u>Q</u> <u>M</u> <u>T</u> <u>I</u> <u>K</u>	<u>T</u> <u>C</u> <u>P</u> <u>D</u> <u>G</u> <u>G</u> <u>S</u> <u>Y</u> <u>G</u> <u>W</u> <u>Y</u> <u>P</u> <u>Y</u> <u>G</u> <u>Y</u> <u>G</u> <u>C</u> <u>N</u> <u>G</u> <u>G</u> <u>V</u> <u>S</u> <u>A</u> <u>T</u> <u>Y</u> <u>T</u> <u>Y</u> <u>E</u> <u>F</u>	YVDAM	(SEQ ID NO : 358)
UL-77	<u>C</u> <u>T</u> <u>T</u> <u>V</u> <u>Y</u> <u>Q</u> <u>K</u> <u>T</u> <u>E</u> <u>S</u> <u>V</u> <u>R</u>	<u>S</u> <u>C</u> <u>P</u> <u>D</u> <u>G</u> <u>S</u> <u>M</u> <u>D</u> <u>G</u> <u>W</u> <u>R</u> <u>C</u> <u>R</u> <u>L</u> <u>G</u> <u>T</u> <u>M</u> <u>N</u> <u>W</u> <u>I</u> <u>Y</u> <u>S</u> <u>N</u> <u>T</u> <u>Y</u> <u>E</u> <u>F</u>	YVDAM	(SEQ ID NO : 359)

Figure 3BLV1H12

SVHQETKKYQSCPDGYRERSDCSNRPACGTSDCCRVSVFGNCLTTLPVSSYSYTYNYEWHVD
(SEQ ID NO : 22)

BLV5B8

TVHQETRKTCSGDYIAVDSCGRGQSDGCVNDCNSCYYGWRNCRRQPAIHSYEFHVD
(SEQ ID NO : 23)

BLV5D3

SVTQRTHVSRSCPDGCSGDGCVDGCCCSAYRCYTPGVRDLSCTSYSITYTYEWNVD
(SEQ ID NO : 24)

BLV8C11

TVHQKTTTRKTCCSDAYRYDSGCGSGCDCCGADCYVFGACTFGLDSSYSYIYIYQWYVD
(SEQ ID NO : 25)

BF4E9

TVHQIFCPDGYSYGYGCGYGYGCSGYDCYGYGGYGYGGYGGYSSYSYSYSYEEYGD
(SEQ ID NO : 26)

BF1H1

TVHPSPDGYSYGYGCGYGYGCSGYDCYGYGGYGYGGYGGYSSYSYSYS
(SEQ ID NO : 27)

F18

TVHQIRCPDGYSYGYGCGYGSYGSYGYDCYGYGGYGGYGGYGGYSSYS
(SEQ ID NO : 28)

Figure 4**Bovine VH-UL:**

```
caggtgcagctgcgggagtcggggccccagcctggtgaagccctcacagaccctctcgctc
Q V Q L R E S G P S L V K P S Q T L S L
acctgcacggcctctggattctcattgagcgcacaaggctgtaggctgggtccgccaggct
T C T A S G F S L S D K A V G W V R Q A
ccaggggaaggcgctgggagtggtcgggtggtatagacactggtggaagcacaggctataac
P G K A L E W L G G I D T G G S T G Y N
ccaggcctgaaatcccggtcagcatcaccaaggacaactccaagagccaagtctctctg
P G L K S R L S I T K D N S K S Q V S L
tcagtgagcagcgtgacaactgaggactcggccacatactactgtactactgtgcaccag (SEQ ID NO: 367)
S V S S V T T E D S A T Y Y C T T V H Q (SEQ ID NO: 29)
```

```
QVQLRESGPSLVKPSQTLSTCTASGFSLSDKAVGWVRQAPGKALEWLGGIDTGGSTGYN
PGLKSRLSITKDNSKSQVSLSVSSVTTEDSATYYCTTVHQ (SEQ ID NO: 29)
```

Figure 5A

4-39:

Cagctgcagctgcaggagtcggggcccaggactggtgaagccttcggagaccctgtccctcacctgcactgtctctgg
 tggctccatcagcagtagtagttactactggggctggatccgccagccccaggggaaggggctggagtggttggga
 gtatctattatagtgaggacactactacaaccctgtccctcaagagtcgagtcaccatatccgtagacacgtccaag
 aaccagttctccctgaagctgagctctgtgaccgccgcagacacggctgtgtattactgtgcgagacacacagtgag
 ggg

(SEQ ID NO: 368)

QLQLQESGPGLVKPSETLSLTCTVSGGSISSSYWGWIRQPPGKGLEWIGSIYYSGSTYYNPSLKSRVTISVDTSK
 NQFSLKLSSVTAADTAVYYCAR

(SEQ ID NO: 31)

4-59*03:

Caggtgcagctgcaggagtcggggcccaggactggtgaagccttcggagaccctgtccctcacctgcactgtctctgg
 tggctccatcagtagttactactggagctggatccggcagccccaggggaagggactggagtggttgggtatatct
 attacagtgggagcaccaactacaaccctccctcaagagtcgagtcaccatatcagtagacacgtccaagaaccaa
 ttctccctgaagctgagctctgtgaccgctgcggaacacggccgtgtattactgtgcg

(SEQ ID NO: 369)

QVQLQESGPGLVKPSETLSLTCTVSGGSISSYYWSWIRQPPGKGLEWIGYIYYSGSTNYPNPSLKSRVTISVDTSKNQ
 FSLKLSSVTAADTAVYYCA

(SEQ ID NO: 32)

4-34*09:

Caggtgcagctgcaggagtcggggcccaggactggtgaagccttcacagaccctgtccctcacctgcgctgtctatgg
 tgggtccttcagtggttactactggagctggatccgccagccccaggggaagggactggagtggttggggaaatca
 atcatagtgggaagcaccaactacaaccctgtccctcaagagtcgagttaccatatcagtagacacgtctaagaaccag
 ttctccctgaagctgagctctgtgactgccgcggacacggccgtgtattactgtgcgaga

(SEQ ID NO: 370)

QVQLQESGPGLVKPSQTLSTCAVYGGSFSGYYWSWIRQPPGKGLEWIGEINHSGSTNYPNPSLKSRVTISVDTSKNQ
 FSLKLSSVTAADTAVYYCAR

(SEQ ID NO: 33)

4-34*02:

Caggtgcagctacaacagtggggcgcaggactggtgaagccttcggagaccctgtccctcacctgcgctgtctatgg
 tgggtccttcagtggttactactggagctggatccgccagccccaggggaaggggctggagtggttggggaaatca
 atcatagtgggaagcaccaactacaaccctgtccctcaagagtcgagtcaccatatcagtagacacgtccaagaaccag
 ttctccctgaagctgagctctgtgaccgccgcggacacggctgtgtattactgtgcgagagg

(SEQ ID NO: 371)

QVQLQQWGAGLLKPSETLSLTCAVYGGSFSGYYWSWIRQPPGKGLEWIGEINHSGSTNYPNPSLKSRVTISVDTSKNQ
 FSLKLSSVTAADTAVYYCAR

(SEQ ID NO: 34)

Figure 5B

4-39	QLQLQESGPGLVKPSETLSLTCTVSGGSISSSSYYWGWIRQPPGKGLEWIGSIYYSGSTY	60
4-59*03	QVQLQESGPGLVKPSETLSLTCTVSGGSISS--YYWSWIRQPPGKGLEWIGYIYYSGSTN	58
4-34*02	QVQLQQWGAGLLKPSETLSLTCAVYGGSFSG--YYWSWIRQPPGKGLEWIGEINHSGSTN	58
4-34*09	QVQLQESGPGLVKPSQTLTLTCAVYGGSFSG--YYWSWIRQPPGKGLEWIGEINHSGSTN	58
VH-UL	QVQLRESGPSLVKPSQTLTLTCTASGFSLS <u>D</u> -- <u>K</u> AVGWVRQAPGKALEWLGGIDTGGSTG	58
	*: **:: *..*:***:*****:. * *:*. .*:**.***.***:* * .***	
4-39	YNPSLKSRTVISVDTSKNQFSLKLSSVTAADTAVYYCAR-----	99 (SEQ ID NO: 31)
4-59*03	YNPSLKSRTVISVDTSKNQFSLKLSSVTAADTAVYYCA-----	96 (SEQ ID NO: 32)
4-34*02	YNPSLKSRTVISVDTSKNQFSLKLSSVTAADTAVYYCAR-----	97 (SEQ ID NO: 33)
4-34*09	YNPSLKSRTVISVDTSKNQFSLKLSSVTAADTAVYYCAR-----	97 (SEQ ID NO: 34)
VH-UL	YNPGLKSRLSITKDNSKSQVSLSVSSVTTEDSATYYCTTVHQ	102 (SEQ ID NO: 29)
	.*:.*: *.**.*.*.*:****:	

Figure 6

BLV1H12 VL:

Caggctgtgctgaatcagccatcatccgtgtccgggtccctggggccagaggggtctccatcacctgctctggaagcag
cagcaatggttggaatggatatgtgagctggtagcaactgatcccaggatcggccccccagaaccctcatctatggtg
acaccagtcgagcctcgggggtccccgaccgattctccgggtccagggtctgggaacacagccaccctgaccatcagc
tcgctccaggctgaggacgaggcagattatttctgtgcatctgctgaggatagtagcagtaatgctgttttcggcag
cgggaccacactgaccgtcctg
(SEQ ID NO: 372)

QAVLNQPSVSGSLGQRVSITCSGSSSNVGNGYVSWYQLIPGSAPRTLIIYGDTSRASGVPDRFSGSRSGNTATLTIS
SLQAEDEADYFCASAEDSSSNAVFGSGTTLTVL
(SEQ ID NO: 35)

Figure 7A

V11-47:

Cagtctgtgctgactcagccaccctcagcgtctgggacccccgggcagaggggtcaccatctcttgttctggaagcag
 ctccaacatcggaagtaattatgtatactgggtaccagcagctcccaggaacggcccccaaactcctcatctatagga
 ataatcagcggccctcaggggtccctgaccgattctctggctccaagtctggcacctcagcctccctggccatcagt
 gggctccggtccgaggatgaggctgattattactgtgcagcatgggatgacagcctgagtggtec
 (SEQ ID NO: 373)

QSVLTQPPSASGTPGQRVITISCSGSSSNIGSNYVYWYQQLPGTAPKLLIYRNNQRPSGVPDRFSGSKSGTSASLAIS
 GLRSEDEADYYCAAWDDSLSG
 (SEQ ID NO: 36)

V11-40*1:

Cagtctgtgctgacgcagccgccctcagtgtctggggccccagggcagaggggtcaccatctcctgcactgggagcag
 ctccaacatcggggcaggttatgatgtacactgggtaccagcagcttcaggaacagcccccaaactcctcatctatg
 gtaacagcaatcgccctcaggggtccctgaccgattctctggctccaagtctggcacctcagcctccctggccatc
 actgggctccaggtgaggatgaggctgattattactgccagtcctatgacagcagcctgagtggttc
 (SEQ ID NO: 374)

QSVLTQPPSVSGAPGQRVITISCTGSSSNIGAGYDVHWYQQLPGTAPKLLIYGNSNRPSGVPDRFSGSKSGTSASLAI
 TGLQAEDEADYYCQSYDSSLG
 (SEQ ID NO: 37)

V11-51 *01:

Cagtctgtgttgacgcagccgccctcagtgtctgcggccccaggacagaagggtcaccatctcctgctctggaagcag
 ctccaacattgggaataattatgtatcctgggtaccagcagctcccaggaacagcccccaaactcctcatttatgaca
 ataataagcgaccctcagggattcctgaccgattctctggctccaagtctggcacgtcagccaccctgggcatcacc
 ggactccagactggggacgaggccgattattactgcggaacatgggatagcagcctgagtgctgg
 (SEQ ID NO: 375)

QSVLTQPPSVSAA PGQKVTISCSGSSSNIGNNYVSWYQQLPGTAPKLLIYDNNKRPSGIPDRFSGSKSGTSATLGIT
 GLQTGDEADYYCGTWDSSLSA
 (SEQ ID NO: 38)

V12-18*02:

Cagtctgccctgactcagcctccctcctgtgtccgggtctcctggacagtcagtcaccatctcctgcactgggaaccagcagtgacgt
 tggtagttataaccgtgtctcctgggtaccagcagccccaggcacagcccccaaactcatgatttatgaggtcagtaatcgccct
 caggggtccctgatcgcttctctgggtccaagtctggcaacacggcctccctgaccatctctgggctccagggtgaggacgaggct
 gattattactgcagctcatatacaagcagcagcactttc
 (SEQ ID NO: 376)

QSALTQPPSVSGSPGQSVTISCTGTSSDVGSYNRVSWYQPPGTAPKLMIYEVSNRPSGVPDRFSGSKSGNTASLTI
 SGLQAEDEADYYCSSYTSSSTF
 (SEQ ID NO: 39)

Figure 7B

Vl1-47	QSVLTQPPSASGTPGQRTISCSGSSSNIGSNY-VYWYQQLPGTAPKLLIYRNNQRPSGV	59	
Vl1-51	QSVLTQPPSVSAAPGQKVTISCSGSSSNIGNNY-VSWYQQLPGTAPKLLIYDNNKRPSGI	59	
Vl1-40*1	QSVLTQPPSVSGAPGQRTISCTGSSSNIGAGYDVHWYQQLPGTAPKLLIYGNSNRPSGV	60	
Vl2-18*02	QSALTQPPSVSGSPGQSVTISCTGTSSDVGSYNRVSWYQQLPGTAPKLMIEVSNRPSGV	60	
Vl1x (cow)	QAVLTQPSSVSGSLGQRVSITCSGSSSNVNGY-VSWYQLIPGSAPRTLIIYGDTSRASGV	59	
	*:.*****.*.*.: ** *:.*:~*:~*:~*:~*:~*:~* * *** **:~*:~*:~*:~*:~*:~* ..*.*.*.:		
Vl1-47	PDRFSGSKSGTSASLAISGLRSEDEADYYCAAWDDSLSG	98	(SEQ ID NO: 36)
Vl1-51	PDRFSGSKSGTSATLGITGLQTGDEADYYCGTWDSSLSA	98	(SEQ ID NO: 37)
Vl1-40*1	PDRFSGSKSGTSASLAITGLQAEDEADYYCQSYDSSLSG	99	(SEQ ID NO: 38)
Vl2-18*02	PDRFSGSKSGNTASLTISGLQAEDEADYYCSSYTSSSTF	99	(SEQ ID NO: 39)
Vl1x	PDRFSGSRSGNTATLTISSLQAEDEADYYFCASAEDSSSN	98	(SEQ ID NO: 35)
	*****:~*.*.*.* *:.*.*.: *****:* : .* :		

Figure 1

BLV1H12	CTSVHQ	ETKKYQ	S <u>C</u> PDGYRERSD <u>C</u> SNRPAC <u>C</u> GTSD <u>C</u> CRVSVFGNCLTTLPVSSYSYTYNYEW	HVDVWGQGGLLVTVSS	(SEQ ID NO : 360)
BLV5B8	CTTVHQ	ETRKT	<u>C</u> SDGYIAVDS <u>C</u> GRGQSDGCVNDCNSCYYGWRNCRRQPAIHSYEF	HVDAWGRGLLVTVSS	(SEQ ID NO : 361)
BLV5D3	CSSVTQ	RTHVSR	S <u>C</u> PDGCSDGDG <u>C</u> VDGCC <u>C</u> SAYR <u>C</u> YTPGVRDLSCSYSITYTYEW	NVDAWGRGLLVTVSS	(SEQ ID NO : 362)
BLV8C11	CTTVHQ	KTRKT	C <u>C</u> SDAYRYDSG <u>C</u> GSGCD <u>C</u> CGAD <u>C</u> YVFGACTFGLDSSYSYIYIYQW	YVDAWQGGLLVTVSS	(SEQ ID NO : 363)
BF4E9	CTTVHQ	IF	<u>C</u> PDGYSYGYG <u>C</u> GYGYG <u>C</u> SGYD <u>C</u> YGYGGYGYGGYGGYSSYSYSYSY	YGDAWQGGLLVTVSS	(SEQ ID NO : 364)
BF1H1	CTTVHP		SPDGYSYGYG <u>C</u> GYGYG <u>C</u> SGYD <u>C</u> YGYGGYGYGGYGGYSSYSYSYS		(SEQ ID NO : 365)
F18	CTTVHQ	IR	<u>C</u> PDGYGYGYG <u>C</u> GYGSYGYSGYDCYGYGGYGGYGGYGGYSSYS		(SEQ ID NO : 366)