



(51) International Patent Classification:

C07K 16/30 (2006.01)

(21) International Application Number:

PCT/US2021/033088

(22) International Filing Date:

19 May 2021 (19.05.2021)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/027,465 20 May 2020 (20.05.2020) US

(71) Applicant: REMD BIOTHERAPEUTICS, INC.

[US/US]; 4010 Adolfo Road, Camarillo, CA 93012 (US).

(72) Inventors: YAN, Hai; 198 Via Inez, Thousand Oaks, CA

91320 (US). ZHANG, John; 2125 Via Arandana, Camarillo, CA 93012 (US).

(74) Agent: CRANDALL, Craig, A.; 3034 Deer Valley Avenue, Newbury Park, CA 91320 (US).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD,


ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: HUMAN ENDOTHELIN RECEPTOR A (ET_AR) ANTAGONIST ANTIBODIES

(57) **Abstract:** This application provides isolated antibodies, and antigen-binding fragments thereof, that specifically bind endothelin A receptor (ET_AR). These ET_AR antibodies, or antigen-binding fragments thereof, have a high affinity for ET_AR, function to effectively block ET-1 binding to ET_AR, are less immunogenic compared to their unmodified parent antibodies in a given species (e.g., in a human), and can be used to treat ET_AR - associated disorders. Importantly, compared to small molecule ET_AR antagonists, the antibodies of the present invention have advantages of longer serum half-life, higher target specificity, thereby limiting the risk of off-target toxicity and improving therapeutic window.

#17: 312A10E5 (A3)

#21: 336D8G9 (A2)

FIG. 2

HUMAN ENDOTHELIN RECEPTOR A (ET_AR) ANTAGONIST ANTIBODIES

RELATED PATENT APPLICATIONS

[001] This application claims benefit of U.S. Provisional Application No. 63/027,465, filed on May 20, 2020, incorporated in its entirety by reference herein.

TECHNICAL FIELD

[002] Endothelin (ET) is a 21 amino acid vasoconstriction peptide hormone, important to the homeostasis and regulation of the biological functions of the cardiovascular system. ET is found not only in the endothelium but also in many other tissues and cell types (Barton et al., 2008, Can. J. Physiol. Pharmacol. 86:485-498). ET has three isoforms: ET-1, ET-2 and ET-3 which differ by a few amino acid residues and are all expressed in the human kidney. ET-1 and ET-2 bind to the two basic types of ET receptors: ET_AR and ET_BR. Both of these receptors are coupled to a Gq-protein and the formation of IP₃. Increased IP₃ causes calcium release by the sarcoplasmic reticulum, which causes smooth muscle contraction. In blood vessels, the ET_AR is dominant under normal conditions in terms of ET-1 effects on contraction. ET_AR has 63% sequence similarity with the human ET_BR. The endothelin axis and its receptors are implied in several physiopathological functions and dysfunctions. By way of non-limiting examples, arterial hypertension, atherosclerosis, coronary artery diseases, liver dysfunctions, cerebrovascular diseases, Crohn's disease, pulmonary fibrosis, asthma, etc. can be mentioned (R. Shah, Eur. J. Int. Med., 18:272-282, 2007). The two receptor subtypes exhibit contrasting cardiovascular actions under normal physiologic conditions. Moreover, endothelin receptors also turned out to be associated with the development of many cancers, by promoting proliferation, survival and dissemination of cancer cells as well as angiogenesis (Bagnato & Rosano, Int. J. of Biochem. & Cell Biology, 40:1443-1451, 2008).

[003] ET-1 is the predominant isoform and ET-1 formation and release are stimulated by angiotensin II (Ang), antidiuretic hormone (ADH), thrombin, cytokines, reactive oxygen

species, and shearing forces acting on the vascular endothelium. ET-1 release is inhibited by prostacyclin and atrial natriuretic peptide as well as by nitric oxide. Once ET-1 is released by the endothelial cell, it binds to receptors on the target tissue (e.g., adjacent vascular smooth muscle). On release from the endothelium, the peptide causes vasoconstriction, mainly via ET_AR. Because of its powerful vasoconstrictor properties, and its effects on intracellular calcium, ET-1 has been implicated in the pathogenesis of hypertension, coronary vasospasm, and heart failure. In the latter condition, ET-1 is released by the failing myocardium where it can contribute to calcium overload and hypertrophy. Endothelin receptor antagonists have been shown to decrease mortality and improve hemodynamics in experimental models of heart failure.

[004] Overproduction of ET-1 in pathophysiological conditions from the endothelium and other cellular sources can lead to constriction of vessels in conditions such as pulmonary hypertension (PAH), leading to vascular remodeling. This is exacerbated particularly where there is reduced production of opposing vasodilators such as nitric oxide, prostacyclin, and endothelium derived hyperpolarizing factor. Activation of the ET_AR can drive proliferation in multiple cell types and contributes to fibrosis and inflammation. Thus, ET_AR antagonists have been explored in the treatment of a range of cardio-pulmonary disorders, including portopulmonary hypertension, hypoplastic left heart syndrome, inoperable chronic thromboembolic pulmonary hypertension, PAH associated with systemic sclerosis, and exercise induced PAH (clinical trials.gov). A non-selective ET-1 receptor antagonist (bosentan) is currently used in the treatment of pulmonary hypertension. PAH is an often fatal disease with a dismal prognosis due to the lack of effective curative therapies.

[005] ET-1 and ET_AR are also involved in other pathophysiological conditions. The experimental evidence collected to date indicates that the ET-1 system is activated in the presence of obesity, type 2 diabetes (T2DM) and metabolic syndrome (METs) defined as elevated blood pressure, abnormal fasting blood glucose and atherogenic dyslipidemia. This phenomenon results predominantly in an increased ET_AR-dependent vasoconstrictor tone and is associated with impaired NO bioactivity, hence leading to abnormal vascular function in patients with these conditions (Campia et al., Br J Pharmacol, 165:561–73, 2012; Cardillo et al., Circulation, 106:1783–7, 2002). Considerable evidence supports a mechanistic role of ET-1 in the pathophysiology of adiposity-related vascular dysfunction. Targeting the ET_AR, therefore,

might have the potential for effective cardiovascular prevention in obesity, the MetS, and T2DM. ET_AR antagonists in clinical studies has shown promise in patients with type 2 diabetic nephropathy as Atrasentan reduced albuminuria and improved BP and lipid spectrum with manageable fluid overload–related adverse events in patients with type 2 diabetic nephropathy receiving renin-angiotensin system (RAS) inhibitors (de Zeeuw et al., J Am Soc Nephrol, 25: 1083–1093, 2014).

DISCLOSURE OF THE INVENTION

[006] In accordance with the present invention, there are provided isolated antibodies, and antigen-binding fragments thereof, that specifically bind endothelin A receptor (ET_AR). These ET_AR antibodies, or antigen-binding fragments thereof, have a high affinity for ET_AR, function to effectively block ET-1 binding to ET_AR, are less immunogenic compared to their unmodified parent antibodies in a given species (e.g., in a human), and can be used to treat ET_AR -associated disorders. Importantly, compared to small molecule ET_AR antagonists, the antibodies of the present invention have advantages of longer serum half-life, higher target specificity, thereby limiting the risk of off-target toxicity and improving therapeutic window.

[007] In various embodiments, the antibody or antigen-binding fragment is selected from a human antibody, a humanized antibody, chimeric antibody, a monoclonal antibody, a polyclonal antibody, a recombinant antibody, a single chain antibody, a diabody, a triabody, a tetrabody, a Fab fragment, a Fab' fragment, a Fab₂ fragment, a F(ab)₂ fragment, a domain antibody, an IgD antibody, an IgE antibody, an IgM antibody, an IgG1 antibody, an IgG2 antibody, an IgG3 antibody, an IgG4 antibody, or an IgG4 antibody having at least one mutation in the hinge region that alleviates a tendency to form intra H-chain disulfide bonds. In various embodiments, the antibody is a chimeric antibody. In various embodiments, the antibody is a humanized antibody. In various embodiments, the antibody is a fully human antibody. In various embodiments, isolated antibodies, and antigen-binding fragments thereof, that have a high affinity for the human ET_AR of SEQ ID NO: 1 are provided.

[008] In various embodiments, the antibody or antigen-binding fragment binds to ET_AR protein with a dissociation constant (K_D) of at least about 1×10^{-6} M, at least about 1×10^{-7} M, at

least about 1×10^{-8} M, at least about 1×10^{-9} M, at least about 1×10^{-10} M, at least about 1×10^{-11} M, or at least about 1×10^{-12} M.

[009] In various embodiments, an isolated antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises either: (a) a light chain CDR3 sequence identical, substantially identical or substantially similar to a CDR3 sequence selected from SEQ ID NOs: 16-18; (b) a heavy chain CDR3 sequence identical, substantially identical or substantially similar to a CDR3 sequence selected from SEQ ID NOs: 8-10; or (c) the light chain CDR3 sequence of (a) and the heavy chain CDR3 sequence of (b).

[010] In various embodiments, the isolated antibody or antigen-binding fragment further comprises an amino acid sequence selected from: (d) a light chain CDR1 sequence identical, substantially identical or substantially similar to a CDR1 sequence selected from SEQ ID NOs: 11-13; (e) a light chain CDR2 sequence identical, substantially identical or substantially similar to a CDR2 sequence selected from SEQ ID NOs: 14-15; (f) a heavy chain CDR1 sequence identical, substantially identical or substantially similar to a CDR1 sequence selected from SEQ ID NOS: 2-4; (g) a heavy chain CDR2 sequence identical, substantially identical or substantially similar to a CDR2 sequence selected from SEQ ID NOs: 5-7; (h) the light chain CDR1 sequence of (d) and the heavy chain CDR1 sequence of (f); or (i) the light chain CDR2 sequence of (e) and the heavy chain CDR2 sequence of (g).

[011] In various embodiments, the isolated human monoclonal antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises: (a) a light chain CDR1 sequence identical, substantially identical or substantially similar to a CDR1 sequence selected from SEQ ID NOs: 11-13; (b) a light chain CDR2 sequence identical, substantially identical or substantially similar to a CDR2 sequence selected from SEQ ID NOs: 14-15; (c) a light chain CDR3 sequence identical, substantially identical or substantially similar to a CDR3 sequence selected from SEQ ID NOs: 16-18; (d) a heavy chain CDR1 sequence identical, substantially identical or substantially similar to a CDR1 sequence selected from SEQ ID NOS: 2-4; (e) a heavy chain CDR2 sequence identical, substantially identical or substantially similar to a CDR2 sequence selected from SEQ ID NOs: 5-7; and (f) a heavy chain CDR3 sequence identical, substantially identical or substantially similar to a CDR3 sequence selected from SEQ ID NOs: 8-10.

[012] In various embodiments, the isolated human monoclonal antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises: (a) a light chain CDR1 sequence identical, substantially identical or substantially similar to SEQ ID NO: 11; (b) a light chain CDR2 sequence identical, substantially identical or substantially similar to SEQ ID NO: 14; (c) a light chain CDR3 sequence identical, substantially identical or substantially similar to SEQ ID NO: 16; (d) a heavy chain CDR1 sequence identical, substantially identical or substantially similar to SEQ ID NO: 2; (e) a heavy chain CDR2 sequence identical, substantially identical or substantially similar to SEQ ID NO: 5; and (f) a heavy chain CDR3 sequence identical, substantially identical or substantially similar to SEQ ID NO: 8.

[013] In various embodiments, the isolated human monoclonal antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises: (a) a light chain CDR1 sequence identical, substantially identical or substantially similar to SEQ ID NO: 12; (b) a light chain CDR2 sequence identical, substantially identical or substantially similar to SEQ ID NO: 14; (c) a light chain CDR3 sequence identical, substantially identical or substantially similar to SEQ ID NO: 17; (d) a heavy chain CDR1 sequence identical, substantially identical or substantially similar to SEQ ID NO: 3; (e) a heavy chain CDR2 sequence identical, substantially identical or substantially similar to SEQ ID NO: 6; and (f) a heavy chain CDR3 sequence identical, substantially identical or substantially similar to SEQ ID NO: 9.

[014] In various embodiments, the isolated human monoclonal antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises: (a) a light chain CDR1 sequence identical, substantially identical or substantially similar to SEQ ID NO: 13; (b) a light chain CDR2 sequence identical, substantially identical or substantially similar to SEQ ID NO: 15; (c) a light chain CDR3 sequence identical, substantially identical or substantially similar to SEQ ID NO: 18; (d) a heavy chain CDR1 sequence identical, substantially identical or substantially similar to SEQ ID NO: 4; (e) a heavy chain CDR2 sequence identical, substantially identical or substantially similar to SEQ ID NO: 7; and (f) a heavy chain CDR3 sequence identical, substantially identical or substantially similar to SEQ ID NO: 10.

[015] In various embodiments, an isolated antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises either: (a) a heavy and/or light chain variable domain(s), the variable domain(s) having a set of three light chain CDR1, CDR2, and CDR3 identical, substantially identical or substantially similar to SEQ ID NOS: 11-13, 14-15, and 16-18, and/or a set of three heavy chain CDR1, CDR2, and CDR3 identical, substantially identical or substantially similar to SEQ ID NOS: 2-4, 5-7, and 8-10; and (b) a set of four variable region framework regions from a human immunoglobulin (IgG). In various embodiments, the antibody can optionally include a hinge region. In various embodiments, the framework regions are chosen from human germline exon X_H, J_H, V_k and J_k sequences. In various embodiments, the antibody is a fully humanized antibody. In various embodiments, the antibody is a fully human antibody.

[016] In various embodiments, an isolated antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises the heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 19 and the light chain variable region having the amino acid sequence set forth in SEQ ID NO: 21. In various embodiments, an isolated antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises the heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 23 and the light chain variable region having the amino acid sequence set forth in SEQ ID NO: 25. In various embodiments, an isolated antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises the heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 27 and the light chain variable region having the amino acid sequence set forth in SEQ ID NO: 29.

[017] In various embodiments, the isolated antibody or antigen-binding fragment, when bound to human ET_AR: (a) binds to human ET_AR with substantially the same or greater K_d as a reference antibody; (b) competes for binding to human ET_AR with said reference antibody; or (c) is less immunogenic in a human subject than said reference antibody, wherein said reference antibody comprises a combination of heavy chain variable domain and light chain variable domain sequences selected from SEQ ID NOS: 19/23/27 and 21/25/29, respectively.

[018] In various embodiments, an isolated humanized antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises a heavy chain

variable region having a sequence identical, substantially identical or substantially similar to SEQ ID NOs: 33-35, and a light chain variable region having the sequence identical, substantially identical or substantially similar to SEQ ID NOs: 36-37.

[019] In various embodiments, the isolated antibody or antigen-binding fragment, when bound to human ET_AR: (a) binds to human ET_AR with substantially the same or greater K_d as a reference antibody; (b) competes for binding to human ET_AR with said reference antibody; or (c) is less immunogenic in a human subject than said reference antibody, wherein said reference antibody comprises the combination of heavy chain variable domain and light chain variable domain sequences set forth in SEQ ID NOs: 27 and 29.

[020] In various embodiments, an isolated humanized antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises a heavy chain having a sequence identical, substantially identical or substantially similar to SEQ ID NOs: 38, 42 and 44, and a light chain variable region having the sequence identical, substantially identical or substantially similar to SEQ ID NOs: 40 and 46. In various embodiments, an isolated humanized antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises the heavy chain sequence set forth in SEQ ID NO: 38, and the light chain sequence set forth in SEQ ID NO: 40. In various embodiments, an isolated humanized antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises the heavy chain sequence set forth in SEQ ID NO: 42, and the light chain sequence set forth in SEQ ID NO: 40. In various embodiments, an isolated humanized antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises the heavy chain sequence set forth in SEQ ID NO: 44, and the light chain sequence set forth in SEQ ID NO: 40.

[021] In another aspect, the present invention relates to a pharmaceutical composition comprising an isolated antibody or antigen-binding fragment of the present invention in admixture with a pharmaceutically acceptable carrier. In various embodiments, the pharmaceutical composition comprises an isolated human antibody in admixture with a pharmaceutically acceptable carrier. In various embodiments, the pharmaceutical composition is formulated for administration via a route selected from the group consisting of subcutaneous injection, intraperitoneal injection, intramuscular injection, intrasternal injection, intravenous

injection, intraarterial injection, intrathecal injection, intraventricular injection, intraurethral injection, intracranial injection, intrasynovial injection or via infusions.

[022] In another aspect, the present invention relates to methods of treating a subject suffering from a ET_AR-associated disorder, comprising administering to said subject a therapeutically effective amount (either as monotherapy or in a combination therapy regimen) of an antibody or antigen-binding fragment thereof of the present invention. In various embodiments, the subject is a human subject.

[023] In various embodiments, the ET_AR-associated disorder is a metabolic disorder selected from the group consisting of diabetes, diabetic myopathy, diabetic nephropathy, diabetic neuropathy, diabetic retinopathy, obesity, dyslipidemia, sarcopenic obesity, fatty liver disease (e.g., nonalcoholic fatty liver disease (NAFLD); nonalcoholic steatohepatitis (NASH)), insulin resistance, and hemochromatosis.

[024] In various embodiments, the ET_AR-associated disorder is a cardiovascular disease selected from the group consisting of angina, heart attack (myocardial infarction), atherosclerosis, heart failure, cardiovascular disease, rheumatic heart disease, cardiac arrhythmias (abnormal heart rhythms), cerebrovascular disease, congenital heart defects, cardiomyopathy, left ventricular hypertrophy, right ventricular hypertrophy, post-infarction heart rupture, infections of the heart, coronary artery disease, peripheral arterial disease, renal artery stenosis, aortic aneurysm, myocardial diseases, heart valve disorders, myocarditis, pericarditis, and pulmonary arterial hypertension (PAH).

[025] In another aspect, the present invention provides combination therapies designed to treat a metabolic disorder in a subject, comprising (a) administering to said subject a therapeutically effective amount of an antibody or antigen-binding fragment thereof of the present invention, and (b) administering a glucose-lowering agent. In various embodiments, the glucose-lowering agent is selected from anti-diabetic agents, anti-hyperglycemic agents, lipid lowering agents, anti-hypertensive agents, and anti-obesity agents. In various embodiments, the glucose-lowering agent is selected from biguanides, sulfonylureas, meglitinides, thiazolidinediones (TZDs), α -glucosidase inhibitors, DPP-4 inhibitors, bile acid sequestrants, dopamine-2 agonists, SGLT2 inhibitors, GLP-1R agonists, amylin mimetics, and insulins.

[026] In another aspect, the present invention provides combination therapies designed to treat a cardiovascular disease in a subject, comprising (a) administering to said subject a therapeutically effective amount of an isolated antibody or antigen-binding fragment of the present invention, and (b) administering a second agent selected from the group consisting of antihypertensive agents, agents for treating chronic heart failure, atherosclerosis or related diseases.

[027] In another aspect, an isolated immunoconjugate or fusion protein comprising an antibody or antigen-binding fragment conjugated to, linked to (or otherwise stably associated with) an effector molecule is provided. In various embodiments, the effector molecule is an immunotoxin, cytokine, chemokine, therapeutic agent, or chemotherapeutic agent.

[028] In another aspect, the antibodies or antigen-binding fragments disclosed herein may be covalently linked to (or otherwise stably associated with) an additional functional moiety, such as a label or a moiety that confers desirable pharmacokinetic properties. In various embodiments, the label is selected from the group consisting of: a fluorescent label, a radioactive label, and a label having a distinctive nuclear magnetic resonance signature.

[029] In another aspect, the present invention provides a method for detecting in vitro or in vivo the presence of human ET_AR peptide in a sample, e.g., for diagnosing a human ET_AR-associated disorder.

[030] In another aspect, isolated nucleic acids comprising the polynucleotide sequence that encodes the antibodies or antigen-binding fragments disclosed herein are provided. In various embodiments, the polynucleotide comprises a heavy chain sequences as set forth in SEQ ID NOs: 38, 42 and 44 a light chain sequence as set forth in SEQ ID NOs: 40 and 46, or both. Also provided are expression vectors comprising the nucleic acid of the present invention. Also provided are isolated cells comprising the expression vectors of the invention. In various embodiments, the cell is a host cell comprising an expression vector of the present invention. In various embodiments, the cell is a hybridoma, wherein the chromosome of the cell comprises a nucleic acid of the present invention. Further provided is a method of making the antibody or antigen-binding fragment of the present invention comprising culturing or incubating the cell under conditions that allow the cell to express the antibody or antigen-binding fragment of the present invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[031] FIG. 1 is a bar graph depicting the ET_AR functional assay data for murine mAbs #1 – #21.

[032] FIG. 2 is a line graph depicting the ET_AR functional assay data (IC₅₀) for murine mAbs #17 (312A10E5 (mAb A3)) and #21 (336D8G9 (mAb A2)).

[033] FIG. 3 depict the FACS binding assay data for various ET_AR humanized antibodies.

MODE(S) OF CARRYING OUT THE INVENTION

[034] The present invention relates to antigen binding proteins such as antibodies, or antigen-binding fragments thereof that specifically bind to human ET_AR. In one aspect, there are provided isolated antibodies, and antigen-binding fragments thereof, that specifically bind ET_AR, have a high affinity for ET_AR, function to inhibit ET_AR, are less immunogenic compared to their unmodified parent antibodies in a given species (e.g., a human), and can be used to treat human disorders mediated by ET_AR. Also provided are nucleic acid molecules, and derivatives and fragments thereof, comprising a sequence of polynucleotides that encode all or a portion of a polypeptide that binds to ET_AR, such as a nucleic acid encoding all or part of an anti-ET_AR antibody, antibody fragment, or antibody derivative. Also provided are vectors and plasmids comprising such nucleic acids, and cells or cell lines comprising such nucleic acids and/or vectors and plasmids. Also provided are methods of making, identifying, or isolating antigen binding proteins that bind to human ET_AR, such as anti-ET_AR antibodies, methods of determining whether an antigen binding protein binds to ET_AR, methods of making compositions, such as pharmaceutical compositions, comprising an antigen binding protein that binds to human ET_AR, and methods for administering an antibody, or antigen-binding fragment thereof that binds ET_AR to a subject, for example, methods for treating a condition mediated by ET_AR.

Definitions

[035] Unless otherwise defined herein, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those commonly used and well known in the art. The methods and techniques of the present invention are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, e.g., Green and Sambrook, *Molecular Cloning: A Laboratory Manual*, 4th ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2012), incorporated herein by reference. Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The nomenclature used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those commonly used and well known in the art. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of subjects.

[036] Polynucleotide and polypeptide sequences are indicated using standard one- or three-letter abbreviations. Unless otherwise indicated, polypeptide sequences have their amino termini at the left and their carboxy termini at the right, and single-stranded nucleic acid sequences, and the top strand of double-stranded nucleic acid sequences, have their 5' termini at the left and their 3' termini at the right. A particular section of a polypeptide can be designated by amino acid residue number such as amino acids 80 to 119, or by the actual residue at that site such as Ser80 to Ser119. A particular polypeptide or polynucleotide sequence also can be described based upon how it differs from a reference sequence. Polynucleotide and polypeptide sequences of particular light and heavy chain variable domains are designated L1 ("light chain

variable domain 1") and H1 ("heavy chain variable domain 1"). Antibodies comprising a light chain and heavy chain are indicated by combining the name of the light chain and the name of the heavy chain variable domains. For example, "L4H7," indicates, for example, an antibody comprising the light chain variable domain of L4 and the heavy chain variable domain of H7.

[037] The term "antibody" is used herein to refer to a protein comprising one or more polypeptides substantially or partially encoded by immunoglobulin genes or fragments of immunoglobulin genes and having specificity to a tumor antigen or specificity to a molecule overexpressed in a pathological state. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon and mu constant region genes, as well as subtypes of these genes and myriad of immunoglobulin variable region genes. Light chains (LC) are classified as either kappa or lambda. Heavy chains (HC) are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively. A typical immunoglobulin (e.g., antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kD) and one "heavy" chain (about 50-70 kD). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition.

[038] In a full-length antibody, each heavy chain is comprised of a heavy chain variable region (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3 (and in some instances, CH4). Each light chain is comprised of a light chain variable region (abbreviated herein as LCVR or VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The extent of the framework region and CDRs has been defined. The sequences of the framework regions of different light or heavy chains are relatively conserved within a species, such as humans. The framework region of an antibody, that is the combined framework regions of the constituent light and heavy chains,

serves to position and align the CDRs in three-dimensional space. Immunoglobulin molecules can be of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG 3, IgG4, IgA1 and IgA2) or subclass.

[039] The CDRs are primarily responsible for binding to an epitope of an antigen. The CDRs of each chain are typically referred to as CDR1, CDR2, CDR3, numbered sequentially starting from the N-terminus, and are also typically identified by the chain in which the particular CDR is located. Thus, a VH CDR3 is located in the variable domain of the heavy chain of the antibody in which it is found, whereas a VL CDR1 is the CDR1 from the variable domain of the light chain of the antibody in which it is found. Antibodies with different specificities (*i.e.* different combining sites for different antigens) have different CDRs. Although it is the CDRs that vary from antibody to antibody, only a limited number of amino acid positions within the CDRs are directly involved in antigen binding. These positions within the CDRs are called specificity determining residues (SDRs).

[040] The Kabat definition is a standard for numbering the residues in an antibody and is typically used to identify CDR regions. The Kabat database is now maintained online and CDR sequences can be determined, for example, see IMGT/V-QUEST programme version: 3.2.18 March 29, 2011, available on the internet and Brochet, X. et al., Nucl. Acids Res. 36, W503-508, 2008). The Chothia definition is similar to the Kabat definition, but the Chothia definition takes into account positions of certain structural loop regions. See, e.g., Chothia et al., J. Mol. Biol., 196: 901-17, 1986; Chothia et al., Nature, 342: 877-83, 1989. The AbM definition uses an integrated suite of computer programs produced by Oxford Molecular Group that model antibody structure. See, e.g., Martin et al., Proc. Natl. Acad. Sci. USA, 86:9268-9272, 1989; "AbM™, A Computer Program for Modeling Variable Regions of Antibodies," Oxford, UK; Oxford Molecular, Ltd. The AbM definition models the tertiary structure of an antibody from primary sequence using a combination of knowledge databases and ab initio methods, such as those described by Samudrala et al., "Ab Initio Protein Structure Prediction Using a Combined Hierarchical Approach," in PROTEINS, Structure, Function and Genetics Suppl., 3:194-198, 1999. The contact definition is based on an analysis of the available complex crystal structures. See, e.g., MacCallum et al., J. Mol. Biol., 5:732-45, 1996.

[041] The term "Fc region" is used to define the C-terminal region of an immunoglobulin heavy chain, which may be generated by papain digestion of an intact antibody. The Fc region may be a native sequence Fc region or a variant Fc region. The Fc region of an immunoglobulin generally comprises two constant domains, a CH2 domain and a CH3 domain, and optionally comprises a CH4 domain. The Fc portion of an antibody mediates several important effector functions e.g. cytokine induction, ADCC, phagocytosis, complement dependent cytotoxicity (CDC) and half-life/clearance rate of antibody and antigen-antibody complexes (e.g., the neonatal FcR (FcRn) binds to the Fc region of IgG at acidic pH in the endosome and protects IgG from degradation, thereby contributing to the long serum half-life of IgG). Replacements of amino acid residues in the Fc portion to alter antibody effector function are known in the art (see, e.g., Winter et al., U.S. Patent No. 5,648,260 and 5,624,821).

[042] Antibodies exist as intact immunoglobulins or as a number of well characterized fragments. Such fragments include Fab fragments, Fab' fragments, Fab₂, F(ab')₂ fragments, single chain Fv proteins ("scFv") and disulfide stabilized Fv proteins ("dsFv"), that bind to the target antigen. A scFv protein is a fusion protein in which a light chain variable region of an immunoglobulin and a heavy chain variable region of an immunoglobulin are bound by a linker, while in dsFvs, the chains have been mutated to introduce a disulfide bond to stabilize the association of the chains. While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such fragments may be synthesized de novo either chemically or by utilizing recombinant DNA methodology. Thus, as used herein, the term antibody encompasses e.g., monoclonal antibodies (including full-length monoclonal antibodies), polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies) formed from at least two intact antibodies, human antibodies, humanized antibodies, camelised antibodies, chimeric antibodies, single-chain Fvs (scFv), single-chain antibodies, single domain antibodies, domain antibodies, Fab fragments, F(ab')₂ fragments, antibody fragments that exhibit the desired biological activity, disulfide-linked Fvs (sdFv), intrabodies, and epitope-binding fragments or antigen binding fragments of any of the above.

[043] Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site. A "Fab fragment" comprises one light chain and the CH1 and variable regions of one heavy chain. The heavy chain of a Fab

molecule cannot form a disulfide bond with another heavy chain molecule. A "Fab' fragment" comprises one light chain and a portion of one heavy chain that contains the VH domain and the CH1 domain and also the region between the CH1 and CH2 domains, such that an interchain disulfide bond can be formed between the two heavy chains of two Fab' fragments to form an F(ab')₂ molecule.

[044] Pepsin treatment of an antibody yields an F(ab')₂ fragment that has two antigen-combining sites and is still capable of cross-linking antigen. A "F(ab')₂ fragment" contains two light chains and two heavy chains containing a portion of the constant region between the CH1 and CH2 domains, such that an interchain disulfide bond is formed between the two heavy chains. A F(ab')₂ fragment thus is composed of two Fab' fragments that are held together by a disulfide bond between the two heavy chains.

[045] The "Fv region" comprises the variable regions from both the heavy and light chains but lacks the constant regions.

[046] "Single-chain antibodies" are Fv molecules in which the heavy and light chain variable regions have been connected by a flexible linker to form a single polypeptide chain, which forms an antigen binding region. Single chain antibodies are discussed in detail in International Patent Application Publication No. WO 88/01649, U.S. Patent No. 4,946,778 and 5,260,203, the disclosures of which are incorporated by reference.

[047] The terms "an antigen-binding fragment" and "antigen-binding protein" as used herein means any protein that binds a specified target antigen. "Antigen-binding fragment" includes but is not limited to antibodies and binding parts thereof, such as immunologically functional fragments. An exemplary antigen-binding fragment of an antibody is the heavy chain and/or light chain CDR(s), or the heavy and/or light chain variable region.

[048] The term "immunologically functional fragment" (or simply "fragment") of an antibody or immunoglobulin chain (heavy or light chain) antigen binding protein, as used herein, is a species of antigen binding protein comprising a portion (regardless of how that portion is obtained or synthesized) of an antibody that lacks at least some of the amino acids present in a full-length chain but which is still capable of specifically binding to an antigen. Such fragments are biologically active in that they bind to the target antigen and can compete with other antigen binding proteins, including intact antibodies, for binding to a given epitope. In some

embodiments, the fragments are neutralizing fragments. In one aspect, such a fragment will retain at least one CDR present in the full-length light or heavy chain, and in some embodiments will comprise a single heavy chain and/or light chain or portion thereof. These biologically active fragments can be produced by recombinant DNA techniques or can be produced by enzymatic or chemical cleavage of antigen binding proteins, including intact antibodies. Immunologically functional immunoglobulin fragments include, but are not limited to, Fab, a diabody, Fab', F(ab')₂, Fv, domain antibodies and single-chain antibodies, and can be derived from any mammalian source, including but not limited to human, mouse, rat, camelid or rabbit. It is further contemplated that a functional portion of the antigen binding proteins disclosed herein, for example, one or more CDRs, could be covalently bound to a second protein or to a small molecule to create a therapeutic agent directed to a particular target in the body, possessing bifunctional therapeutic properties, or having a prolonged serum half-life.

[049] Diabodies are bivalent antibodies comprising two polypeptide chains, wherein each polypeptide chain comprises VH and VL regions joined by a linker that is too short to allow for pairing between two regions on the same chain, thus allowing each region to pair with a complementary region on another polypeptide chain (see, e.g., Holliger et al., Proc. Natl. Acad. Sci. USA, 90:6444-48, 1993; and Poljak et al., Structure, 2:1121-23, 1994). If the two polypeptide chains of a diabody are identical, then a diabody resulting from their pairing will have two identical antigen binding sites. Polypeptide chains having different sequences can be used to make a diabody with two different antigen binding sites. Similarly, tribodies and tetrabodies are antibodies comprising three and four polypeptide chains, respectively, and forming three and four antigen binding sites, respectively, which can be the same or different.

[050] Bispecific antibodies or fragments can be of several configurations. For example, bispecific antibodies may resemble single antibodies (or antibody fragments) but have two different antigen binding sites (variable regions). In various embodiments bispecific antibodies can be produced by chemical techniques (Kranz et al., Proc. Natl. Acad. Sci. USA, 78:5807, 1981; by "polydoma" techniques (see, e.g., U.S. Patent No. 4,474,893); or by recombinant DNA techniques. In various embodiments bispecific antibodies of the present disclosure can have binding specificities for at least two different epitopes at least one of which is a tumor associate antigen. In various embodiments the antibodies and fragments can also be heteroantibodies.

Heteroantibodies are two or more antibodies, or antibody binding fragments (e.g., Fab) linked together, each antibody or fragment having a different specificity.

[051] The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigen. Furthermore, in contrast to polyclonal antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. The modifier "monoclonal" is not to be construed as requiring production of the antibody by any particular method.

[052] The term "chimeric antibody" as used herein refers to an antibody which has framework residues from one species, such as human, and CDRs (which generally confer antigen binding) from another species, such as a murine antibody that specifically binds targeted antigen.

[053] The term "human antibody", as used herein, is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies of the disclosure may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*), for example in the CDRs and in particular CDR3. However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

[054] The term "humanized antibody" as used herein refers to an antibody comprising a humanized light chain and a humanized heavy chain immunoglobulin. A humanized antibody binds to the same antigen as the donor antibody that provides the CDRs. The acceptor framework of a humanized immunoglobulin or antibody may have a limited number of substitutions by amino acids taken from the donor framework. Humanized or other monoclonal antibodies can have additional conservative amino acid substitutions which have substantially no effect on antigen binding or other immunoglobulin functions. In various embodiments, the

framework regions are chosen from human germline exon X_H, J_H, V_K and J_K sequences. For example, acceptor sequences for humanization of FR of a V_H domain can be chosen from genuine V_H exons V_H 1-18 (Matsuda et al., Nature Genetics 3:88-94, 1993) or V_H1-2 (Shin et al., EMBO J. 10:3641-3645, 1991) and for the hinge region (J_H), exon J_H-6 (Mattila et al., Eur. J. Immunol. 25:2578-2582, 1995). In other examples, germline V_K exon B3 (Cox et al., Eur. J. Immunol. 24:827-836, 1994) and J_K exon J_K-1 (Hieter et al., J. Biol. Chem. 257:1516-1522, 1982) can be chosen as acceptor sequences for V_L domain humanization.

[055] The term "recombinant human antibody", as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies expressed using a recombinant expression vector transfected into a host cell; antibodies isolated from a recombinant, combinatorial human antibody library; antibodies isolated from an animal (e.g., a mouse) that is transgenic for human immunoglobulin genes; or antibodies prepared, expressed, created or isolated by any other means that involves splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable and constant regions derived from human germline immunoglobulin sequences. In various embodiments, however, such recombinant human antibodies are subjected to *in vitro* mutagenesis (or, when an animal transgenic for human Ig sequences is used, *in vivo* somatic mutagenesis) and thus the amino acid sequences of the V_H and V_L regions of the recombinant antibodies are sequences that, while derived from and related to human germline V_H and V_L sequences, may not naturally exist within the human antibody germline repertoire *in vivo*. All such recombinant means are well known to those of ordinary skill in the art.

[056] The term "anti-ETR antagonist antibody" (interchangeably termed "anti-ETR antibody") refers to an antibody that is able to bind to ETR and inhibit ETR biological activity and/or downstream pathway(s) mediated by ETR signaling. An anti-ETR antagonist antibody encompasses antibodies that block, antagonize, suppress or reduce (including significantly) ETR biological activity, including downstream pathways mediated by ETR signaling, such as receptor binding and/or elicitation of a cellular response to ETR. For purpose of the present invention, it will be explicitly understood that the term "anti-ETR antagonist antibody" encompasses all the previously identified terms, titles, and functional states and characteristics

whereby the ETR itself, an ETR biological activity (including but not limited to its ability to mediate any aspect of headache), or the consequences of the biological activity, are substantially nullified, decreased, or neutralized in any meaningful degree. In some embodiment, an anti-ETR antagonist antibody binds ETR and prevents ETR binding to a ETR receptor. In other embodiments, an anti-ETR antibody binds ETR and prevents activation of a ETR receptor. Examples of anti-ETR antagonist antibodies are provided herein.

[057] The term "epitope" as used herein includes any protein determinant capable of specific binding to an immunoglobulin or T-cell receptor or otherwise interacting with a molecule. Epitopic determinants generally consist of chemically active surface groupings of molecules such as amino acids or carbohydrate or sugar side chains and generally have specific three dimensional structural characteristics, as well as specific charge characteristics. An epitope may be "linear" or "conformational." In a linear epitope, all of the points of interaction between the protein and the interacting molecule (such as an antibody) occur linearly along the primary amino acid sequence of the protein. In a conformational epitope, the points of interaction occur across amino acid residues on the protein that are separated from one another. Once a desired epitope on an antigen is determined, it is possible to generate antibodies to that epitope, e.g., using the techniques described in the present disclosure. Alternatively, during the discovery process, the generation and characterization of antibodies may elucidate information about desirable epitopes. From this information, it is then possible to competitively screen antibodies for binding to the same epitope. An approach to achieve this is to conduct cross-competition studies to find antibodies that competitively bind with one another, e.g., the antibodies compete for binding to the antigen.

[058] An antigen binding protein, including an antibody, "specifically binds" to an antigen if it binds to the antigen with a high binding affinity as determined by a dissociation constant (K_D , or corresponding K_b , as defined below) value of at least 1×10^{-6} M, or at least 1×10^{-7} M, or at least 1×10^{-8} M, or at least 1×10^{-9} M, or at least 1×10^{-10} M, or at least 1×10^{-11} M. An antigen binding protein that specifically binds to the human antigen of interest may be able to bind to the same antigen of interest from other species as well, with the same or different affinities. The term " K_D " as used herein refers to the equilibrium dissociation constant of a particular antibody-antigen interaction.

[059] The term "surface plasmon resonance" as used herein refers to an optical phenomenon that allows for the analysis of real-time biospecific interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIACORE™ system (Pharmacia Biosensor AB, Uppsala, Sweden and Piscataway, N.J.). For further descriptions, see Jonsson U. et al., *Ann. Biol. Clin.*, 51:19-26, 1993; Jonsson U. et al., *Biotechniques*, 11:620-627, 1991; Jonsson B. et al., *J. Mol. Recognit.*, 8:125-131, 1995; and Johnsson B. et al., *Anal. Biochem*, 198:268-277, 1991.

[060] The term "immunogenicity" as used herein refers to the ability of an antibody or antigen binding fragment to elicit an immune response (humoral or cellular) when administered to a recipient and includes, for example, the human anti-mouse antibody (HAMA) response. A HAMA response is initiated when T-cells from a subject make an immune response to the administered antibody. The T-cells then recruit B-cells to generate specific "anti-antibody" antibodies.

[061] The term "immune cell" as used herein means any cell of hematopoietic lineage involved in regulating an immune response against an antigen (e.g., an autoantigen). In various embodiments, an immune cell is, e.g., a T cell, a B cell, a dendritic cell, a monocyte, a natural killer cell, a macrophage, Langerhan's cells, or Kuffer cells.

[062] The terms "polypeptide", "peptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. In various embodiments, "peptides", "polypeptides", and "proteins" are chains of amino acids whose alpha carbons are linked through peptide bonds. The terminal amino acid at one end of the chain (amino terminal) therefore has a free amino group, while the terminal amino acid at the other end of the chain (carboxy terminal) has a free carboxyl group. As used herein, the term "amino terminus" (abbreviated N-terminus) refers to the free α -amino group on an amino acid at the amino terminal of a peptide or to the α -amino group (imino group when participating in a peptide bond) of an amino acid at any other location within the peptide. Similarly, the term "carboxy terminus" refers to the free carboxyl group on the carboxy terminus of a peptide or the carboxyl group of an amino acid at any other location within the peptide. Peptides also include essentially any polyamino acid including, but not limited to, peptide mimetics such as amino acids joined by an ether as opposed to an amide bond.

[063] The term "recombinant polypeptide", as used herein, is intended to include all polypeptides, including fusion molecules that are prepared, expressed, created, derived from, or isolated by recombinant means, such as polypeptides expressed using a recombinant expression vector transfected into a host cell.

[064] Polypeptides of the disclosure include polypeptides that have been modified in any way and for any reason, for example, to: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter binding affinities, and (5) confer or modify other physicochemical or functional properties. For example, single or multiple amino acid substitutions (e.g., conservative amino acid substitutions) may be made in the naturally occurring sequence (e.g., in the portion of the polypeptide outside the domain(s) forming intermolecular contacts). A "conservative amino acid substitution" refers to the substitution in a polypeptide of an amino acid with a functionally similar amino acid. The following six groups each contain amino acids that are conservative substitutions for one another:

Alanine (A), Serine (S), and Threonine (T)

Aspartic acid (D) and Glutamic acid (E)

Asparagine (N) and Glutamine (Q)

Arginine (R) and Lysine (K)

Isoleucine (I), Leucine (L), Methionine (M), and Valine (V)

Phenylalanine (F), Tyrosine (Y), and Tryptophan (W)

[065] A "non-conservative amino acid substitution" refers to the substitution of a member of one of these classes for a member from another class. In making such changes, according to various embodiments, the hydropathic index of amino acids may be considered. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics. They are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

[066] The importance of the hydropathic amino acid index in conferring interactive biological function on a protein is understood in the art (see, for example, Kyte et al., 1982, J. Mol. Biol. 157:105-131). It is known that certain amino acids may be substituted for other amino acids having a similar hydropathic index or score and still retain a similar biological activity. In making changes based upon the hydropathic index, in various embodiments, the substitution of amino acids whose hydropathic indices are within ± 2 is included. In various embodiments, those that are within ± 1 are included, and in various embodiments, those within ± 0.5 are included.

[067] It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity, particularly where the biologically functional protein or peptide thereby created is intended for use in immunological embodiments, as disclosed herein. In various embodiments, the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with its immunogenicity and antigenicity, i.e., with a biological property of the protein.

[068] The following hydrophilicity values have been assigned to these amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0+-.1); glutamate (+3.0+-.1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5+-.1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5) and tryptophan (-3.4). In making changes based upon similar hydrophilicity values, in various embodiments, the substitution of amino acids whose hydrophilicity values are within ± 2 is included, in various embodiments, those that are within ± 1 are included, and in various embodiments, those within ± 0.5 are included. Exemplary amino acid substitutions are set forth in Table 1.

Table 1

<u>Original Residues</u>	<u>Exemplary Substitutions</u>	<u>Preferred Substitutions</u>
Ala	Val, Leu, Ile	Val
Arg	Lys, Gln, Asn	Lys
Asn	Gln	

Asp	Glu	
Cys	Ser, Ala	Ser
Gln	Asn	
Glu	Asp	
Gly	Pro, Ala	Ala
His	Asn, Gln, Lys, Arg	Arg
Ile	Leu, Val, Met, Ala, Phe, Norleucine	Leu
Leu	Norleucine, Ile, Val, Met, Ala, Phe	Ile
Lys	Arg, 1,4 Diamino-butyric Acid, Gln, Asn	Arg
Met	Leu, Phe, Ile	Leu
Phe	Leu, Val, Ile, Ala, Tyr	Leu
Pro	Ala	Gly
Ser	Thr, Ala, Cys	Thr
Thr	Ser	
Trp	Tyr, Phe	Tyr
Tyr	Trp, Phe, Thr, Ser	Phe
Val	Ile, Met, Leu, Phe, Ala, Norleucine	Leu

[069] The term "polypeptide fragment" and "truncated polypeptide" as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion as compared to a corresponding full-length protein. In various embodiments, fragments can be, *e.g.*, at least 5, at least 10, at least 25, at least 50, at least 100, at least 150, at least 200, at least 250, at least 300, at least 350, at least 400, at least 450, at least 500, at least 600, at least 700, at least 800, at least 900 or at least 1000 amino acids in length. In various embodiments, fragments can also be, *e.g.*, at most 1000, at most 900, at most 800, at most 700, at most 600, at most 500, at most 450, at most 400, at most 350, at most 300, at most 250, at most 200, at

most 150, at most 100, at most 50, at most 25, at most 10, or at most 5 amino acids in length.

A fragment can further comprise, at either or both of its ends, one or more additional amino acids, for example, a sequence of amino acids from a different naturally-occurring protein (*e.g.*, an Fc or leucine zipper domain) or an artificial amino acid sequence (*e.g.*, an artificial linker sequence).

[070] The terms "polypeptide variant" and "polypeptide mutant" as used herein refers to a polypeptide that comprises an amino acid sequence wherein one or more amino acid residues are inserted into, deleted from and/or substituted into the amino acid sequence relative to another polypeptide sequence. In various embodiments, the number of amino acid residues to be inserted, deleted, or substituted can be, *e.g.*, at least 1, at least 2, at least 3, at least 4, at least 5, at least 10, at least 25, at least 50, at least 75, at least 100, at least 125, at least 150, at least 175, at least 200, at least 225, at least 250, at least 275, at least 300, at least 350, at least 400, at least 450 or at least 500 amino acids in length. Variants of the present disclosure include fusion proteins.

[071] A "derivative" of a polypeptide is a polypeptide that has been chemically modified, *e.g.*, conjugation to another chemical moiety such as, for example, polyethylene glycol, albumin (*e.g.*, human serum albumin), phosphorylation, and glycosylation.

[072] The term "% sequence identity" is used interchangeably herein with the term "% identity" and refers to the level of amino acid sequence identity between two or more peptide sequences or the level of nucleotide sequence identity between two or more nucleotide sequences, when aligned using a sequence alignment program. For example, as used herein, 80% identity means the same thing as 80% sequence identity determined by a defined algorithm and means that a given sequence is at least 80% identical to another length of another sequence. In various embodiments, the % identity is selected from, *e.g.*, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% or more sequence identity to a given sequence. In various embodiments, the % identity is in the range of, *e.g.*, about 60% to about 70%, about 70% to about 80%, about 80% to about 85%, about 85% to about 90%, about 90% to about 95%, or about 95% to about 99%.

[073] The term "% sequence homology" is used interchangeably herein with the term "% homology" and refers to the level of amino acid sequence homology between two or more

peptide sequences or the level of nucleotide sequence homology between two or more nucleotide sequences, when aligned using a sequence alignment program. For example, as used herein, 80% homology means the same thing as 80% sequence homology determined by a defined algorithm, and accordingly a homologue of a given sequence has greater than 80% sequence homology over a length of the given sequence. In various embodiments, the % homology is selected from, *e.g.*, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% or more sequence homology to a given sequence. In various embodiments, the % homology is in the range of, *e.g.*, about 60% to about 70%, about 70% to about 80%, about 80% to about 85%, about 85% to about 90%, about 90% to about 95%, or about 95% to about 99%.

[074] Exemplary computer programs which can be used to determine identity between two sequences include, but are not limited to, the suite of BLAST programs, *e.g.*, BLASTN, BLASTX, and TBLASTX, BLASTP and TBLASTN, publicly available on the Internet at the NCBI website. See also Altschul et al., J. Mol. Biol. 215:403-10, 1990 (with special reference to the published default setting, *i.e.*, parameters $w=4$, $t=17$) and Altschul et al., Nucleic Acids Res., 25:3389-3402, 1997. Sequence searches are typically carried out using the BLASTP program when evaluating a given amino acid sequence relative to amino acid sequences in the GenBank Protein Sequences and other public databases. The BLASTX program is preferred for searching nucleic acid sequences that have been translated in all reading frames against amino acid sequences in the GenBank Protein Sequences and other public databases. Both BLASTP and BLASTX are run using default parameters of an open gap penalty of 11.0, and an extended gap penalty of 1.0, and utilize the BLOSUM-62 matrix. See *Id.*

[075] In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, *e.g.*, Karlin & Altschul, Proc. Nat'l. Acad. Sci. USA, 90:5873-5787, 1993). One measure of similarity provided by the BLAST algorithm is the smallest sum probability ($P(N)$), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is, *e.g.*, less than about 0.1, less than about 0.01, or less than about 0.001.

[076] The terms "substantial similarity" or "substantially similar," in the context of polypeptide sequences, indicate that a polypeptide region has a sequence with at least 70%, typically at least 80%, more typically at least 85%, or at least 90% or at least 95% sequence similarity to a reference sequence. For example, a polypeptide is substantially similar to a second polypeptide, for example, where the two peptides differ by one or more conservative substitution(s).

[077] "Polynucleotide" refers to a polymer composed of nucleotide units. Polynucleotides include naturally occurring nucleic acids, such as deoxyribonucleic acid ("DNA") and ribonucleic acid ("RNA") as well as nucleic acid analogs. Nucleic acid analogs include those which include non-naturally occurring bases, nucleotides that engage in linkages with other nucleotides other than the naturally occurring phosphodiester bond or which include bases attached through linkages other than phosphodiester bonds. Thus, nucleotide analogs include, for example and without limitation, phosphorothioates, phosphorodithioates, phosphorotriesters, phosphoramidates, boranophosphates, methylphosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides, peptide-nucleic acids (PNAs), and the like. Such polynucleotides can be synthesized, for example, using an automated DNA synthesizer. The term "nucleic acid" typically refers to large polynucleotides. The term "oligonucleotide" typically refers to short polynucleotides, generally no greater than about 50 nucleotides. It will be understood that when a nucleotide sequence is represented by a DNA sequence (i.e., A, T, G, C), this also includes an RNA sequence (i.e., A, U, G, C) in which "U" replaces "T."

[078] Conventional notation is used herein to describe polynucleotide sequences: the left-hand end of a single-stranded polynucleotide sequence is the 5'-end; the left-hand direction of a double-stranded polynucleotide sequence is referred to as the 5'-direction. The direction of 5' to 3' addition of nucleotides to nascent RNA transcripts is referred to as the transcription direction. The DNA strand having the same sequence as an mRNA is referred to as the "coding strand"; sequences on the DNA strand having the same sequence as an mRNA transcribed from that DNA and which are located 5' to the 5'-end of the RNA transcript are referred to as "upstream sequences"; sequences on the DNA strand having the same sequence as the RNA and which are 3' to the 3' end of the coding RNA transcript are referred to as "downstream sequences."

[079] "Complementary" refers to the topological compatibility or matching together of interacting surfaces of two polynucleotides. Thus, the two molecules can be described as complementary, and furthermore, the contact surface characteristics are complementary to each other. A first polynucleotide is complementary to a second polynucleotide if the nucleotide sequence of the first polynucleotide is substantially identical to the nucleotide sequence of the polynucleotide binding partner of the second polynucleotide, or if the first polynucleotide can hybridize to the second polynucleotide under stringent hybridization conditions.

[080] "Hybridizing specifically to" or "specific hybridization" or "selectively hybridize to", refers to the binding, duplexing, or hybridizing of a nucleic acid molecule preferentially to a particular nucleotide sequence under stringent conditions when that sequence is present in a complex mixture (e.g., total cellular) DNA or RNA. The term "stringent conditions" refers to conditions under which a probe will hybridize preferentially to its target subsequence, and to a lesser extent to, or not at all to, other sequences. "Stringent hybridization" and "stringent hybridization wash conditions" in the context of nucleic acid hybridization experiments such as Southern and northern hybridizations are sequence-dependent and are different under different environmental parameters. An extensive guide to the hybridization of nucleic acids can be found in Tijssen, 1993, *Laboratory Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Acid Probes*, part I, chapter 2, "Overview of principles of hybridization and the strategy of nucleic acid probe assays", Elsevier, N.Y.; Sambrook et al., 2001, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, 3rd ed., NY; and Ausubel et al., eds., *Current Edition, Current Protocols in Molecular Biology*, Greene Publishing Associates and Wiley Interscience, NY.

[081] Generally, highly stringent hybridization and wash conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Very stringent conditions are selected to be equal to the T_m for a particular probe. An example of stringent hybridization conditions for hybridization of complementary nucleic acids which have more than about 100 complementary residues on a filter in a Southern or northern blot is 50% formamide with 1 mg of heparin at 42°C, with the hybridization being carried out overnight. An example of highly

stringent wash conditions is 0.15 M NaCl at 72°C for about 15 minutes. An example of stringent wash conditions is a 0.2 x SSC wash at 65°C for 15 minutes. See Sambrook et al. for a description of SSC buffer. A high stringency wash can be preceded by a low stringency wash to remove background probe signal. An exemplary medium stringency wash for a duplex of, e.g., more than about 100 nucleotides, is 1 x SSC at 45°C for 15 minutes. An exemplary low stringency wash for a duplex of, e.g., more than about 100 nucleotides, is 4-6 x SSC at 40°C for 15 minutes. In general, a signal to noise ratio of 2 x (or higher) than that observed for an unrelated probe in the particular hybridization assay indicates detection of a specific hybridization.

[082] "Primer" refers to a polynucleotide that is capable of specifically hybridizing to a designated polynucleotide template and providing a point of initiation for synthesis of a complementary polynucleotide. Such synthesis occurs when the polynucleotide primer is placed under conditions in which synthesis is induced, i.e., in the presence of nucleotides, a complementary polynucleotide template, and an agent for polymerization such as DNA polymerase. A primer is typically single-stranded but may be double-stranded. Primers are typically deoxyribonucleic acids, but a wide variety of synthetic and naturally occurring primers are useful for many applications. A primer is complementary to the template to which it is designed to hybridize to serve as a site for the initiation of synthesis but need not reflect the exact sequence of the template. In such a case, specific hybridization of the primer to the template depends on the stringency of the hybridization conditions. Primers can be labeled with, e.g., chromogenic, radioactive, or fluorescent moieties and used as detectable moieties.

[083] "Probe," when used in reference to a polynucleotide, refers to a polynucleotide that is capable of specifically hybridizing to a designated sequence of another polynucleotide. A probe specifically hybridizes to a target complementary polynucleotide but need not reflect the exact complementary sequence of the template. In such a case, specific hybridization of the probe to the target depends on the stringency of the hybridization conditions. Probes can be labeled with, e.g., chromogenic, radioactive, or fluorescent moieties and used as detectable moieties. In instances where a probe provides a point of initiation for synthesis of a complementary polynucleotide, a probe can also be a primer.

[084] A "vector" is a polynucleotide that can be used to introduce another nucleic acid linked to it into a cell. One type of vector is a "plasmid," which refers to a linear or circular double stranded DNA molecule into which additional nucleic acid segments can be ligated. Another type of vector is a viral vector (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), wherein additional DNA segments can be introduced into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors comprising a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. An "expression vector" is a type of vector that can direct the expression of a chosen polynucleotide.

[085] A "regulatory sequence" is a nucleic acid that affects the expression (e.g., the level, timing, or location of expression) of a nucleic acid to which it is operably linked. The regulatory sequence can, for example, exert its effects directly on the regulated nucleic acid, or through the action of one or more other molecules (e.g., polypeptides that bind to the regulatory sequence and/or the nucleic acid). Examples of regulatory sequences include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Further examples of regulatory sequences are described in, for example, Goeddel, 1990, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, Calif. and Baron et al., 1995, Nucleic Acids Res. 23:3605-06. A nucleotide sequence is "operably linked" to a regulatory sequence if the regulatory sequence affects the expression (e.g., the level, timing, or location of expression) of the nucleotide sequence.

[086] A "host cell" is a cell that can be used to express a polynucleotide of the disclosure. A host cell can be a prokaryote, for example, *E. coli*, or it can be a eukaryote, for example, a single-celled eukaryote (e.g., a yeast or other fungus), a plant cell (e.g., a tobacco or tomato plant cell), an animal cell (e.g., a human cell, a monkey cell, a hamster cell, a rat cell, a mouse cell, or an insect cell) or a hybridoma. Typically, a host cell is a cultured cell that can be transformed or transfected with a polypeptide-encoding nucleic acid, which can then be expressed in the host cell. The phrase "recombinant host cell" can be used to denote a host cell that has been transformed or transfected with a nucleic acid to be expressed. A host cell

also can be a cell that comprises the nucleic acid but does not express it at a desired level unless a regulatory sequence is introduced into the host cell such that it becomes operably linked with the nucleic acid. It is understood that the term host cell refers not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to, e.g., mutation or environmental influence, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

[087] The term "isolated molecule" (where the molecule is, for example, a polypeptide or a polynucleotide) is a molecule that by virtue of its origin or source of derivation (1) is not associated with naturally associated components that accompany it in its native state, (2) is substantially free of other molecules from the same species (3) is expressed by a cell from a different species, or (4) does not occur in nature. Thus, a molecule that is chemically synthesized, or expressed in a cellular system different from the cell from which it naturally originates, will be "isolated" from its naturally associated components. A molecule also may be rendered substantially free of naturally associated components by isolation, using purification techniques well known in the art. Molecule purity or homogeneity may be assayed by a number of means well known in the art. For example, the purity of a polypeptide sample may be assayed using polyacrylamide gel electrophoresis and staining of the gel to visualize the polypeptide using techniques well known in the art. For certain purposes, higher resolution may be provided by using HPLC or other means well known in the art for purification.

[088] A protein or polypeptide is "substantially pure," "substantially homogeneous," or "substantially purified" when at least about 60% to 75% of a sample exhibits a single species of polypeptide. The polypeptide or protein may be monomeric or multimeric. A substantially pure polypeptide or protein will typically comprise about 50%, 60%, 70%, 80% or 90% W/W of a protein sample, more usually about 95%, and preferably will be over 99% pure. Protein purity or homogeneity may be indicated by a number of means well known in the art, such as polyacrylamide gel electrophoresis of a protein sample, followed by visualizing a single polypeptide band upon staining the gel with a stain well known in the art. For certain purposes, higher resolution may be provided by using HPLC or other means well known in the art for purification.

[089] "Linker" refers to a molecule that joins two other molecules, either covalently, or through ionic, van der Waals or hydrogen bonds, e.g., a nucleic acid molecule that hybridizes to one complementary sequence at the 5' end and to another complementary sequence at the 3' end, thus joining two non-complementary sequences. A "cleavable linker" refers to a linker that can be degraded or otherwise severed to separate the two components connected by the cleavable linker. Cleavable linkers are generally cleaved by enzymes, typically peptidases, proteases, nucleases, lipases, and the like. Cleavable linkers may also be cleaved by environmental cues, such as, for example, changes in temperature, pH, salt concentration, etc.

[090] The terms "label" or "labeled" as used herein refers to incorporation of another molecule in the antibody. In one embodiment, the label is a detectable marker, e.g., incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotinyl moieties that can be detected by marked avidin (e.g., streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or calorimetric methods). In another embodiment, the label or marker can be therapeutic, e.g., a drug conjugate or toxin. Various methods of labeling polypeptides and glycoproteins are known in the art and may be used. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I), fluorescent labels (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic labels (e.g., horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase), chemiluminescent markers, biotinyl groups, predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags), magnetic agents, such as gadolinium chelates, toxins such as pertussis toxin, taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. In some embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance.

[091] The term "immunoconjugate" or "fusion protein" as used herein refers to a molecule comprising an antibody or antigen-binding fragment thereof conjugated (or linked)

directly or indirectly to an effector molecule. The effector molecule can be a detectable label, an immunotoxin, cytokine, chemokine, therapeutic agent, or chemotherapeutic agent. The antibody or antigen-binding fragment thereof may be conjugated to an effector molecule via a peptide linker. An immunoconjugate and/or fusion protein retains the immunoreactivity of the antibody or antigen-binding fragment, e.g., the antibody or antigen-binding fragment has approximately the same, or only slightly reduced, ability to bind the antigen after conjugation as before conjugation. As used herein, an immunoconjugate may also be referred to as an antibody drug conjugate (ADC). Because immunoconjugates and/or fusion proteins are originally prepared from two molecules with separate functionalities, such as an antibody and an effector molecule, they are also sometimes referred to as "chimeric molecules."

[092] "Pharmaceutical composition" refers to a composition suitable for pharmaceutical use in an animal. A pharmaceutical composition comprises a pharmacologically effective amount of an active agent and a pharmaceutically acceptable carrier. "Pharmacologically effective amount" refers to that amount of an agent effective to produce the intended pharmacological result. "Pharmaceutically acceptable carrier" refers to any of the standard pharmaceutical carriers, vehicles, buffers, and excipients, such as a phosphate buffered saline solution, 5% aqueous solution of dextrose, and emulsions, such as an oil/water or water/oil emulsion, and various types of wetting agents and/or adjuvants. Suitable pharmaceutical carriers and formulations are described in Remington's Pharmaceutical Sciences, 21st Ed. 2005, Mack Publishing Co, Easton. A "pharmaceutically acceptable salt" is a salt that can be formulated into a compound for pharmaceutical use including, e.g., metal salts (sodium, potassium, magnesium, calcium, etc.) and salts of ammonia or organic amines.

[093] The terms "treat", "treating" and "treatment" refer to a method of alleviating or abrogating a biological disorder and/or at least one of its attendant symptoms. As used herein, to "alleviate" a disease, disorder or condition means reducing the severity and/or occurrence frequency of the symptoms of the disease, disorder, or condition. As used herein, "treatment" is an approach for obtaining beneficial or desired clinical results. For purposes of this invention, beneficial or desired clinical results include, but are not limited to, any one or more of: alleviation of one or more symptoms, diminishment of extent of disease, preventing or delaying spread (e.g., metastasis, for example metastasis to the lung or to the lymph node) of disease,

preventing or delaying recurrence of disease, delay or slowing of disease progression, amelioration of the disease state, and remission (whether partial or total). Also encompassed by "treatment" is a reduction of pathological consequence of a proliferative disease. The methods of the invention contemplate any one or more of these aspects of treatment.

[094] The term "effective amount" or "therapeutically effective amount" as used herein refers to an amount of a compound or composition sufficient to treat a specified disorder, condition or disease such as ameliorate, palliate, lessen, and/or delay one or more of its symptoms.

[095] It is understood that aspects and embodiments of the invention described herein include "consisting" and/or "consisting essentially of" aspects and embodiments.

[096] Reference to "about" a value or parameter herein includes (and describes) variations that are directed to that value or parameter per se. For example, description referring to "about X" includes description of "X".

[097] As used herein and in the appended claims, the singular forms "a," "or," and "the" include plural referents unless the context clearly dictates otherwise. It is understood that aspects and variations of the invention described herein include "consisting" and/or "consisting essentially of" aspects and variations.

ET_AR

[098] ET_AR is a member of class A of the GPCR family, comprises seven transmembrane domains. Its extracellular domain is very small, only 1/7 of the complete protein sequence. Human ET_AR has 63% sequence similarity with the human ET_BR. The two receptor subtypes exhibit contrasting cardiovascular actions under normal physiologic conditions. On release from the endothelium, the peptide causes vasoconstriction, mainly via ET_AR. The expression of ET_AR on endothelial and smooth muscle results in broad receptor distributions in tissues or organs receiving a blood supply. Highest relative expression of ET_AR mRNA is associated with the heart and lungs with relatively low expression in the CNS. ET_AR mRNA was the fifth most abundant GPCR in heart atria and 11th in ventricle, emphasizing the importance of the ET pathway in the cardiovascular system (Regard et al., 2008 Anatomical profiling of G protein coupled receptor expression. Cell 135:561–571).

[0099] The term "ET_AR" as used herein includes human ET_AR (hET_AR), variants, isoforms, and species homologs of hET_AR, and analogs having at least one common epitope with hET_AR. In various embodiments, a hET_AR as used herein may comprise the amino acid sequence set forth in SEQ ID NO: 1:

METLCLRASFWLALVGCVISDNPERYSTNLSNHVDDFTTFRGTELSFLVTTHQPTNLVL
 PSNGSMHNYCPQQTKITSAPKYINTVISCTIFIVGMVGNATLLRIIYQNKCMRNGPNALIA
 SLALGDLIYVVIDLPINVFKLLAGRWPFHDNDFGVFLCKLFPFLQKSSVGITVLNLCALSV
 DRYRAVASWSRVQGGIGIPLVTAIEIVSIWILSFILAIPEAIGFVMVPFEYRGEQHKTCMLNA
 TSKFMEFYQDVKDWLFGFYFCMPLVCTAIFYTLMTCEMLNRRNGSLRIALSEHLKQR
 REVAKTVFCLVVIFALCWFLHLSRILKKTVYNEMDKNRCELLSFLLLMDYIGINLATMNS
 CINPIALYFVSKKFKNCFQSCLCCCCYQSKSLMTSVPMNGTSIQWKNHDQNNHNTDRS
 SHKDSMN (SEQ ID NO: 1)

[0100] In various embodiments, a ET_AR comprises an amino acid sequence that shares an observed homology of, *e.g.*, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% with the human ET_AR sequence of SEQ ID NO: 1. In some embodiments, the has at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, at least 1x, at least 1.5x, at least 2x, at least 2.5x, or at least 3x activity of the human ET_AR of SEQ ID NO: 1. Variants of ET_AR may be described herein by reference to the addition, deletion, or substitution of amino acid residue present at a given position in the 360 amino acid sequence of SEQ ID NO: 1. Thus, for example, the term "P29W" indicates that the "P" (proline, in standard single letter code) residue at position 29 in SEQ ID NO: 1 has been substituted with a "W" (tryptophan, in standard single letter code).

Antibodies

[0101] Methods of generating novel antibodies that bind to human ET_AR are known to those skilled in the art. For example, a method for generating a monoclonal antibody that binds specifically to an ET_AR may comprise administering to a mouse an amount of an immunogenic composition comprising the ET_AR effective to stimulate a detectable immune response, obtaining antibody-producing cells (*e.g.*, cells from the spleen) from the mouse and fusing the antibody-producing cells with myeloma cells to obtain antibody-producing hybridomas, and

testing the antibody-producing hybridomas to identify a hybridoma that produces a monoclonal antibody that binds specifically to the ET_AR. Once obtained, a hybridoma can be propagated in a cell culture, optionally in culture conditions where the hybridoma-derived cells produce the monoclonal antibody that binds specifically to ET_AR. The monoclonal antibody may be purified from the cell culture. A variety of different techniques are then available for testing antibody:antigen interactions to identify particularly desirable antibodies.

[0102] Other suitable methods of producing or isolating antibodies of the requisite specificity can be used, including, for example, methods which select recombinant antibody from a library, or which rely upon immunization of transgenic animals (e.g., mice) capable of producing a full repertoire of human antibodies. See e.g., Jakobovits et al., Proc. Natl. Acad. Sci. USA, 90: 2551-2555, 1993; Jakobovits et al., Nature, 362:255-258, 1993; Lonberg et al., U.S. Pat. No. 5,545,806; Surani et al., U.S. Patent No. 5,545,807.

[0103] Antibodies can be engineered in numerous ways. They can be made as single-chain antibodies (including small modular immunopharmaceuticals or SMIPsTM), Fab and F(ab')₂ fragments, etc. Antibodies can be humanized, chimerized, deimmunized, or fully human. Numerous publications set forth the many types of antibodies and the methods of engineering such antibodies. For example, see U.S. Pat. Nos. 6,355,245; 6,180,370; 5,693,762; 6,407,213; 6,548,640; 5,565,332; 5,225,539; 6,103,889; and 5,260,203.

[0104] Chimeric antibodies can be produced by recombinant DNA techniques known in the art. For example, a gene encoding the Fc constant region of a murine (or other species) monoclonal antibody molecule is digested with restriction enzymes to remove the region encoding the murine Fc, and the equivalent portion of a gene encoding a human Fc constant region is substituted (see Robinson et al., International Patent Publication PCT/US86/02269; Akira, et al., European Patent Application 184,187; Taniguchi, M., European Patent Application 171,496; Morrison et al., European Patent Application 173,494; Neuberger et al., International Application WO 86/01533; Cabilly et al. U.S. Pat. No. 4,816,567; Cabilly et al., European Patent Application 125,023; Better et al., Science, 240:1041-1043, 1988; Liu et al., PNAS USA, 84:3439-3443, 1987; Liu et al., J. Immunol. 139:3521-3526, 1987; Sun et al., PNAS USA, 84:214-218, 1987; Nishimura et al., Canc. Res. 47:999-1005, 1987; Wood et al., Nature 314:446-449, 1985; and Shaw et al., J. Natl Cancer Inst., 80:1553-1559, 1988).

[0105] Methods for humanizing antibodies have been described in the art. In practice, humanized antibodies are typically human antibodies in which some hypervariable region residues and possibly some framework region residues are substituted by residues from analogous sites in rodent antibodies. Accordingly, such "humanized" antibodies are chimeric antibodies wherein substantially less than an intact human variable region has been substituted by the corresponding sequence from a nonhuman species. To a degree, this can be accomplished in connection with techniques of humanization and display techniques using appropriate libraries. It will be appreciated that murine antibodies or antibodies from other species can be humanized or primatized using techniques well known in the art (see e.g., Winter et al., *Immunol Today*, 14:43-46, 1993; and Wright et al., *Crit. Reviews in Immunol.*, 12:125-168, 1992). The antibody of interest may be engineered by recombinant DNA techniques to substitute the CH1, CH2, CH3, hinge domains, and/or the framework domain with the corresponding human sequence (see WO 92/02190 and U.S. Pat. Nos. 5,530,101, 5,585,089, 5,693,761, 5,693,792, 5,714,350, and 5,777,085). Also, the use of Ig cDNA for construction of chimeric immunoglobulin genes is known in the art (Liu et al., *P.N.A.S.* 84:3439, 1987; J. *Immunol.* 139:3521, 1987). mRNA is isolated from a hybridoma or other cell producing the antibody and used to produce cDNA. The cDNA of interest may be amplified by the polymerase chain reaction using specific primers (U.S. Pat. Nos. 4,683,195 and 4,683,202). Alternatively, a library is made and screened to isolate the sequence of interest. The DNA sequence encoding the variable region of the antibody is then fused to human constant region sequences. The sequences of human constant regions to genes may be found in Kabat et al. (1991) *Sequences of Proteins of Immunological Interest*, N.I.H. publication no. 91-3242. Human C region genes are readily available from known clones. The choice of isotype will be guided by the desired effector functions, such as complement fixation, or activity in antibody-dependent cellular cytotoxicity. In various embodiments, the isotype is selected from the group consisting of IgG1, IgG2, IgG3 and IgG4. Either of the human light chain constant regions, kappa or lambda, may be used. The chimeric, humanized antibody is then expressed by conventional methods.

[0106] U.S. Patent No. 5,693,761 to Queen et al, discloses a refinement on Winter et al. for humanizing antibodies, and is based on the premise that ascribes avidity loss to problems in the structural motifs in the humanized framework which, because of steric or other chemical

incompatibility, interfere with the folding of the CDRs into the binding-capable conformation found in the mouse antibody. To address this problem, Queen teaches using human framework sequences closely homologous in linear peptide sequence to framework sequences of the mouse antibody to be humanized. Accordingly, the methods of Queen focus on comparing framework sequences between species. Typically, all available human variable region sequences are compared to a particular mouse sequence and the percentage identity between correspondent framework residues is calculated. The human variable region with the highest percentage is selected to provide the framework sequences for the humanizing project. Queen also teaches that it is important to retain in the humanized framework, certain amino acid residues from the mouse framework critical for supporting the CDRs in a binding-capable conformation. Potential criticality is assessed from molecular models. Candidate residues for retention are typically those adjacent in linear sequence to a CDR or physically within 6Å of any CDR residue.

[0107] In other approaches, the importance of particular framework amino acid residues is determined experimentally once a low-avidity humanized construct is obtained, by reversion of single residues to the mouse sequence and assaying antigen-binding as described by Riechmann et al, 1988. Another example approach for identifying important amino acids in framework sequences is disclosed by U.S. Patent No. 5,821,337 to Carter et al, and by U.S. Patent No. 5,859,205 to Adair et al. These references disclose specific Kabat residue positions in the framework, which, in a humanized antibody may require substitution with the correspondent mouse amino acid to preserve avidity.

[0108] Another method of humanizing antibodies, referred to as "framework shuffling", relies on generating a combinatorial library with nonhuman CDR variable regions fused in frame into a pool of individual human germline frameworks (Dall'Acqua et al., Methods, 36:43, 2005). The libraries are then screened to identify clones that encode humanized antibodies which retain good binding.

[0109] The choice of human variable regions, both light and heavy, to be used in making the desired humanized antibodies is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable region of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence

that is closest to that of the rodent is then accepted as the human framework region (framework region) for the humanized antibody (Sims et al., J. Immunol., 151:2296, 1993; Chothia et al., J. Mol. Biol., 196:901, 1987). Another method uses a particular framework region derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chain variable regions. The same framework may be used for several different humanized antibodies (Carter et al., Proc. Natl. Acad. Sci. USA, 89:4285, 1992; Presta et al., J. Immunol., 151:2623, 1993).

[0110] The choice of nonhuman residues to substitute into the human variable region can be influenced by a variety of factors. These factors include, for example, the rarity of the amino acid in a particular position, the probability of interaction with either the CDRs or the antigen, and the probability of participating in the interface between the light and heavy chain variable domain interface. (See, for example, U.S. Patent Nos. 5,693,761, 6,632,927, and 6,639,055). One method to analyze these factors is through the use of three-dimensional models of the nonhuman and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available that illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, e.g., the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, nonhuman residues can be selected and substituted for human variable region residues in order to achieve the desired antibody characteristic, such as increased affinity for the target antigen(s).

[0111] Methods for making fully human antibodies have been described in the art. By way of example, a method for producing an anti-ET_AR antibody or antigen-binding fragment thereof comprises the steps of synthesizing a library of human antibodies on phage, screening the library with ET_AR or an antibody-binding portion thereof, isolating phage that bind ET_AR, and obtaining the antibody from the phage. By way of another example, one method for preparing the library of antibodies for use in phage display techniques comprises the steps of immunizing a non-human animal comprising human immunoglobulin loci with ET_AR or an antigenic portion thereof to create an immune response, extracting antibody-producing cells from the immunized

animal; isolating RNA encoding heavy and light chains of antibodies of the invention from the extracted cells, reverse transcribing the RNA to produce cDNA, amplifying the cDNA using primers, and inserting the cDNA into a phage display vector such that antibodies are expressed on the phage. Recombinant anti-ET_AR antibodies of the invention may be obtained in this way.

[0112] Recombinant human anti-ETR antibodies of the invention can also be isolated by screening a recombinant combinatorial antibody library. Preferably the library is a scFv phage display library, generated using human VL and VH cDNAs prepared from mRNA isolated from B cells. Methods for preparing and screening such libraries are known in the art. Kits for generating phage display libraries are commercially available (e.g., the Pharmacia Recombinant Phage Antibody System, catalog no. 27-9400-01; and the Stratagene SurfZAPTM phage display kit, catalog no. 240612). There also are other methods and reagents that can be used in generating and screening antibody display libraries (see, e.g., U.S. Patent No. 5,223,409; PCT Publication Nos. WO 92/18619, WO 91/17271, WO 92/20791, WO 92/15679, WO 93/01288, WO 92/01047, WO 92/09690; Fuchs et al., *Bio/Technology* 9:1370-1372 (1991); Hay et al., *Hum. Antibod. Hybridomas* 3:81-85, 1992; Huse et al., *Science* 246:1275-1281, 1989; McCafferty et al., *Nature* 348:552-554, 1990; Griffiths et al., *EMBO J.* 12:725-734, 1993; Hawkins et al., *J. Mol. Biol.* 226:889-896, 1992; Clackson et al., *Nature* 352:624-628, 1991; Gram et al., *Proc. Natl. Acad. Sci. USA* 89:3576-3580, 1992; Garrad et al., *Bio/Technology* 9:1373-1377, 1991; Hoogenboom et al., *Nuc. Acid Res.* 19:4133-4137, 1991; and Barbas et al., *Proc. Natl. Acad. Sci. USA* 88:7978-7982, 1991, each incorporated herein by reference for purposes of teaching preparation and screening of phase display libraries.

[0113] Human antibodies are also produced by immunizing a non-human, transgenic animal comprising within its genome some or all of human immunoglobulin heavy chain and light chain loci with a human IgE antigen, e.g., a XenoMouseTM animal (Abgenix, Inc./Amgen, Inc.--Fremont, Calif.). XenoMouseTM mice are engineered mouse strains that comprise large fragments of human immunoglobulin heavy chain and light chain loci and are deficient in mouse antibody production. See, e.g., Green et al., *Nature Genetics* 7:13-21, 1994; and U.S. Patent Nos. 5,916,771, 5,939,598, 5,985,615, 5,998,209, 6,075,181, 6,091,001, 6,114,598, 6,130,364, 6,162,963 and 6,150,584. See also WO 91/10741, WO 94/02602, WO 96/34096, WO 96/33735, WO 98/16654, WO 98/24893, WO 98/50433, WO 99/45031, WO 99/53049, WO 00/09560, and

WO 00/037504. XenoMouseTM mice produce an adult-like human repertoire of fully human antibodies and generate antigen-specific human antibodies. In some embodiments, the XenoMouseTM mice contain approximately 80% of the human antibody V gene repertoire through introduction of megabase sized, germline configuration fragments of the human heavy chain loci and kappa light chain loci in yeast artificial chromosome (YAC). In other embodiments, XenoMouseTM mice further contain approximately all of the human lambda light chain locus. See Mendez et al., Nature Genetics 15:146-156, 1997, Green and Jakobovits, J. Exp. Med. 188:483-495 (1998), and WO 98/24893 (each incorporated by reference in its entirety for purposes of teaching the preparation of fully human antibodies). In another aspect, the present invention provides a method for making anti-ET_AR antibodies from non-human, non-mouse animals by immunizing non-human transgenic animals that comprise human immunoglobulin loci with a ET_AR antigen. One can produce such animals using the methods described in the above-cited documents.

Characterization of Antibody Binding to Antigen

[0114] Antibodies of the present invention can be tested for binding to human ET_AR by, for example, standard ELISA. As an example, microtiter plates are coated with purified ET_AR in PBS, and then blocked with 5% bovine serum albumin in PBS. Dilutions of antibody (e.g., dilutions of plasma from ET_AR-immunized mice) are added to each well and incubated for 1-2 hours at 37°C. The plates are washed with PBS/Tween and then incubated with secondary reagent (e.g., for human antibodies, a goat-anti-human IgG Fc-specific polyclonal reagent) conjugated to alkaline phosphatase for 1 hour at 37°C. After washing, the plates are developed with pNPP substrate (1 mg/ml) and analyzed at OD of 405-650. Preferably, mice which develop the highest titers will be used for fusions. An ELISA assay can also be used to screen for hybridomas that show positive reactivity with ET_AR immunogen. Hybridomas that bind with high avidity to ET_AR are subcloned and further characterized. One clone from each hybridoma, which retains the reactivity of the parent cells (by ELISA), can be chosen for making a 5-10 vial cell bank stored at -140°C., and for antibody purification.

[0115] To determine if the selected anti-ET_AR monoclonal antibodies bind to unique epitopes, each antibody can be biotinylated using commercially available reagents (Pierce,

Rockford, Ill.). Competition studies using unlabeled monoclonal antibodies and biotinylated monoclonal antibodies can be performed using ET_AR coated-ELISA plates as described above. Biotinylated mAb binding can be detected with a strep-avidin-alkaline phosphatase probe. To determine the isotype of purified antibodies, isotype ELISAs can be performed using reagents specific for antibodies of a particular isotype. For example, to determine the isotype of a human monoclonal antibody, wells of microtiter plates can be coated with 1 µg/ml of anti-human immunoglobulin overnight at 4°C. After blocking with 1% BSA, the plates are reacted with 1 µg/ml or less of test monoclonal antibodies or purified isotype controls, at ambient temperature for one to two hours. The wells can then be reacted with either human IgG1 or human IgM-specific alkaline phosphatase-conjugated probes. Plates are developed and analyzed as described above.

[0116] Anti-ET_AR human IgGs can be further tested for reactivity with ETR antigen by Western blotting. Briefly, ET_AR can be prepared and subjected to sodium dodecyl sulfate polyacrylamide gel electrophoresis. After electrophoresis, the separated antigens are transferred to nitrocellulose membranes, blocked with 10% fetal calf serum, and probed with the monoclonal antibodies to be tested. Human IgG binding can be detected using anti-human IgG alkaline phosphatase and developed with BCIP/NBT substrate tablets (Sigma Chem. Co., St. Louis, Mo.).

Identification of Anti-ET_AR Antibodies

[0117] The present invention provides monoclonal antibodies, and antigen-binding fragments thereof, that specifically bind to ET_AR antigen.

[0118] Further included in the present invention are antibodies that bind to the same epitope as the anti-ET_AR antibodies of the present invention. To determine if an antibody can compete for binding to the same epitope as the epitope bound by the anti-ET_AR antibodies of the present invention, a cross-blocking assay, e.g., a competitive ELISA assay, can be performed. In an exemplary competitive ELISA assay, ET_AR coated on the wells of a microtiter plate is pre-incubated with or without candidate competing antibody and then the biotin labeled anti-ET_AR antibody of the invention is added. The amount of labeled anti-ET_AR antibody bound to the ET_AR antigen in the wells is measured using avidin-peroxidase conjugate and appropriate

substrate. The antibody can be labeled with a radioactive or fluorescent label or some other detectable and measurable label. The amount of labeled anti-ET_AR antibody that bound to the antigen will have an indirect correlation to the ability of the candidate competing antibody (test antibody) to compete for binding to the same epitope, i.e., the greater the affinity of the test antibody for the same epitope, the less labeled antibody will be bound to the antigen-coated wells. A candidate competing antibody is considered an antibody that binds substantially to the same epitope or that competes for binding to the same epitope as an anti-ET_AR antibody of the invention if the candidate antibody can block binding of the ET_AR antibody by at least 20%, preferably by at least 20-50%, even more preferably, by at least 50% as compared to the control performed in parallel in the absence of the candidate competing antibody. It will be understood that variations of this assay can be performed to arrive at the same quantitative value.

[0119] The amino acid sequences of the heavy chain variable region CDRs and the light chain variable region CDRs of three murine antibodies, 178D1F6C1G6 (also referred to hereinafter as “A1”), 336D8G9 (also referred to hereinafter as “A2”) and 312A10E5 (also referred to hereinafter as “A3”), generated as described herein, are shown below in Table 2.

Table 2
Heavy Chain CDRs

<u>Ab</u>	<u>HCDR1</u>	<u>HCDR2</u>	<u>HCDR3</u>
A1	SYWMN (SEQ ID NO: 2)	QIYPGNGDTNYNGKFRG (SEQ ID NO: 5)	RGSLDY (SEQ ID NO: 8)
A2	DYYIH (SEQ ID NO: 3)	RIDPEDGDTEYAPKFQG (SEQ ID NO: 6)	NWYSNSY (SEQ ID NO: 9)
A3	GYTFTNYGMS (SEQ ID NO: 4)	WINTYSGESTYADDFKG (SEQ ID NO: 7)	SLNRDTWFTY (SEQ ID NO: 10)

Light Chain CDRs

<u>Ab</u>	<u>LCDR1</u>	<u>LCDR2</u>	<u>LCDR3</u>
------------------	---------------------	---------------------	---------------------

A1	KSSHSLVHSNGDTYLH (SEQ ID NO: 11)	KVSNRFS (SEQ ID NO: 14)	SQTTYVPYT (SEQ ID NO: 16)
A2	RSSQSLVHSNGNTYLH (SEQ ID NO: 12)	KVSNRFS (SEQ ID NO: 14)	SQSTHVPYT (SEQ ID NO: 17)
A3	RSITGAVTTSNYAN (SEQ ID NO: 13)	GTNNRAP (SEQ ID NO: 15)	ALWYSNHWV (SEQ ID NO: 18)

[0120] In various embodiments of the present invention, the antibody or antigen-binding fragment is a murine antibody, 178D1F6C1G6 ("A1"), comprising the heavy chain variable region sequence of SEQ ID NO: 19:

QVQLQQSGAELVRPGSSVRISCKAFDYAFSSYWMNWVKQRPGQDLEWIGQIYPNG
DTNYNGKFRGKASLTADKSSSTAYMHLSSLTSEDSAVYFCARRGSLDYWGQGTSVTV
SS (SEQ ID NO: 19)

and the light chain variable region sequence of SEQ ID NO: 21:

DVVMTQIPLSLPVSLGDQSSISCKSSHSLVHSNGDTYLHWFLQKPGQSPKLLIYKVSNR
FSGVPDRFTGSGSGTEFTLKISRVEAEDLGVYFCSQTTYVPYTFGGGTKLEIK
(SEQ ID NO: 21)

[0121] In certain alternative embodiments, the antibody is an antibody comprising a heavy chain and a light chain, wherein heavy chain comprises a heavy chain variable region, and wherein the heavy chain variable region comprises a sequence that has at least about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least about 99% identity to the amino acid sequence as set forth in SEQ ID NO: 19, or its corresponding polynucleotide sequence SEQ ID NO: 20:

caggttcagctgcagcagctctggggctgagttggtgaggcctgggtcctcagtgaggatttcctgcaaggctttgactatgc
attcagtagctactggatgaactgggtgaagcagaggcctggacaggatcttgagtgattggacagattatcctggaaat
ggtgataactaactacaatggaaagtcaggggttaaagcctcactgactgcagacaaatcctccagcacagcctacatgca
cctcagcagcctaacatctgaggactctgcggtctatttctgtgcaagacgggggtctttgactactgggggtcaaggaacct
cagtcaccgtctcctca (SEQ ID NO: 20)

and wherein the light chain comprises a light chain variable region, and wherein the light chain variable region comprises a sequence that has at least about 75%, at least about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least about 99% identity to the amino acid sequence as set forth in SEQ ID NO: 21, or its corresponding polynucleotide sequence SEQ ID NO: 22:

gatgtggtgatgacccaaattccactctccctgctgtcagtttgagatcaatcctccatctctgcaaattagtcatagcct
tgtacatagtaatggagacacctaatttaccattggttcctgcagaagccaggccagttcctcctgatctacaaagtct
ccaaccgattttctgggtcccagacaggttcactggcagtggtcagggacagagttcacactcaagatcagcagagtg
aggctgaggatctgggagtttattctgctctcaaactacatatgttcggtacacgttcggaggggggaccaagctggaaata
aaa (SEQ ID NO: 22)

[0122] In various embodiments of the present invention, the antibody or antigen-binding fragment is a murine antibody, 336D8G9 ("A2"), comprising the heavy chain variable region sequence of SEQ ID NO: 23:

EVQLQQSGAELVRPGASVKLSCPASGFTIKDYYIHVVVKQRPEQGLEWIGRIDPEDGDT
EYAPKFQGKATMTADTSSNTAYLHLSSLTSEDVAVYYCTSNWYSNSYWGGGTTTLTVS
S (SEQ ID NO: 23)

and the light chain variable region sequence of SEQ ID NO: 25:

DVVMQTPLSLPVSLGDQASISCRSSQSLVHSNGNTYLHWYLQKPGQSPKVLIIYKVS
NRFSGVPDRFSGSGSGTDFTLKISRVEAEDLGVYFCSQSTHVPYTFGGGTKLEIK
(SEQ ID NO: 25)

[0123] In certain alternative embodiments, the antibody is an antibody comprising a heavy chain and a light chain, wherein heavy chain comprises a heavy chain variable region, and wherein the heavy chain variable region comprises a sequence that has at least about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least about 99% identity to the amino acid sequence as set forth in SEQ ID NO: 23, or its corresponding polynucleotide sequence SEQ ID NO: 24:

gaggttcagctgcagcagctctggggcagagcttgtagggccaggggcctcagtcagttgtcctgccagcttctggcttc
accattaaagactactatatacactgggtgaagcagagggcctgagcagggcctggagtggtggaaggattgatcctga
ggatgggtgatactgaatatgccccgaagttccagggcaaggccactatgactgcagacacatcctccaacacagcctacc
tgcacctcagcagcctgacatctgaggacactccgtctattactgtacttcaaactggatatagtaactcctactggggccaa
ggcaccactctcacagtctcctca (SEQ ID NO: 24)

and wherein the light chain comprises a light chain variable region, and wherein the light chain variable region comprises a sequence that has at least about 75%, at least about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least about 99% identity to the amino acid sequence as set forth in SEQ ID NO: 25, or its corresponding polynucleotide sequence SEQ ID NO: 26:

gatgttgatgacccaaactccactctccctgcctgtcagttctggagatcaagcctccatctcttcagatctagtcagagc
ctgtacacagtaatggaaacacctaattacattggtacctgcaaaagccaggccagttctcaaaggtcctgatctacaaag
ttccaaccgattttctgggtcccagacaggttcagtggtgagtcagggacagatttcacactcaagatcagcagagt
ggaggctgaggatctgggagtttatttctgtctcaaagtacacatgttcgtatcgttcggaggggggaccaagctggaa
ataaaa (SEQ ID NO: 26)

[0124] In various embodiments of the present invention, the antibody or antigen-binding fragment is a murine antibody, 312A10E5 ("A3"), comprising the heavy chain variable region sequence of SEQ ID NO: 27:

QIQLVQSGPDLKKPGETVKISCKASGYTFTNYGMSWVKQAPGKGLKWMGWINTYSGE
STYADDFKGRFAFSLETSASTAYLQINNLKNEDTATYFCARSLNRDWTFTYWGQGTLV
TVSA (SEQ ID NO: 27)

and the light chain variable region sequence of SEQ ID NO: 29:

QAVVTQESALSTSPGETVTLTCRSITGAVTTSNYANWVQEKPDHLFTGLIGGTNNRAP
GVPARFSGSLIGDKAALTITGAQTEDEAIYFCALWYSNHWVFGGGTKLTVL
(SEQ ID NO: 29)

[0125] In certain alternative embodiments, the antibody is an antibody comprising a heavy chain and a light chain, wherein heavy chain comprises a heavy chain variable region, and wherein the heavy chain variable region comprises a sequence that has at least about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least

about 99% identity to the amino acid sequence as set forth in SEQ ID NO: 27, or its corresponding polynucleotide sequence SEQ ID NO: 28:

```
cagatccagttggtacagtctggacctgacctgaagaagcctggagagacagtcaagatctcctgcaaggcttctgggtat
accttcacaaactatggaatgagctgggtgaaacaggctccaggaaagggtttaagtggatgggctggataaacaccta
ttctggagagtcaacatatgctgatgactcaagggacggttgccttctcttggaaacctctgccagcactgcctatttcag
atcaacaacctcaaaaatgaggacacggctacatattctgtgcaagatccctcaaccgggacacctggtttacttactggg
gccaaagggactctggtcactgtctctgca (SEQ ID NO: 28)
```

and wherein the light chain comprises a light chain variable region, and wherein the light chain variable region comprises a sequence that has at least about 75%, at least about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least about 99% identity to the amino acid sequence as set forth in SEQ ID NO: 29, or its corresponding polynucleotide sequence SEQ ID NO: 30:

```
caggctgttgtagctcaggaatctgcactctccacatcacctgggtgaaacagtcacactcactgtcgctcaattactggggct
gttacaactagtaactatgccaactgggtccaagaaaaaccagatcattgttactgggtctaataagggtgtaccaacaacc
gagctccaggtgttctgccagattctcaggctccctgattggagacaaggctgccctcaccatcacaggggcacagactg
aggatgaggcaatatatttctgtgctctatggtacagcaaccactgggtgttcggtggaggaaactgactgtccta
(SEQ ID NO: 30)
```

[0126] In various embodiments, the heavy chain variable domain comprises a sequence of amino acids that is encoded by a polynucleotide that hybridizes under moderately stringent conditions to the complement of a polynucleotide that encodes a heavy chain variable domain having the sequence of SEQ ID NOs: 19, 23 and 27. In another embodiment, the heavy chain variable domain comprises a sequence of amino acids that is encoded by a polynucleotide that hybridizes under stringent conditions to the complement of a polynucleotide that encodes a heavy chain variable domain having the sequence of SEQ ID NOs: 19, 23 and 27. In another embodiment, the light chain variable domain comprises a sequence of amino acids that is encoded by a polynucleotide that hybridizes under moderately stringent conditions to the complement of a polynucleotide that encodes a light chain variable domain having the sequence of SEQ ID NOs: 21, 25 and 29. In another embodiment, the light chain variable domain comprises a sequence of amino acids that is encoded by a polynucleotide that hybridizes under

stringent conditions to the complement of a polynucleotide that encodes a light chain variable domain having the sequence of SEQ ID NOs: 21, 25 and 29.

[0127] In various embodiments, antibodies of the present invention include antibodies that bind to the same epitope as murine antibody A1. In various embodiments, antibodies of the present invention include antibodies that bind to the same epitope as murine antibody A2. In various embodiments, antibodies of the present invention include antibodies that bind to the same epitope as murine antibody A3.

[0128] In various embodiments of the present invention, the antibody or antigen-binding fragment is a murine-human chimeric antibody (derived from murine antibody A3) comprising the heavy chain variable region sequence of SEQ ID NO: 31:

QVQLVQSGAEVKKPGASVKVSCKASGYTFTNYGMSWVRQAPGGGLEWMGWINTY
SGESTYADDFKGRVTMTTDTSTSTAYMELRSLRSDDTAVYYCARSLNRDTWFTYWG
QGTLVTVSS (SEQ ID NO: 31)

and the light chain variable region sequence of SEQ ID NO: 32:

QTVVTQEPSFSVSPGGTVTLTCRSITGAVTTSNYANWYQQTPGQAPRTLTYGTNNRAP
GVPDRFSGSILGNKAALTITGAQADDESDYYCALWYSNHWVFEGGGTKLTVL
(SEQ ID NO: 32)

[0129] In various embodiments of the present invention, the antibody or antigen-binding fragment is an isolated humanized antibody or antigen-binding fragment comprising a heavy chain variable region having a sequence identical, substantially identical or substantially similar to SEQ ID NOs: 33-35, and a light chain variable region having the sequence identical, substantially identical or substantially similar to SEQ ID NOs: 36-37. In various embodiments the antibody is a humanized antibody or antigen-binding fragment thereof which comprises the heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 33 and the light chain variable region having the amino acid sequence set forth in SEQ ID NO: 37. In various embodiments the antibody is a humanized antibody or antigen-binding fragment thereof which comprises the heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 34 and the light chain variable region having the amino acid sequence set forth in

SEQ ID NO: 37. In various embodiments the antibody is a humanized antibody or antigen-binding fragment thereof which comprises the heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 35 and the light chain variable region having the amino acid sequence set forth in SEQ ID NO: 37.

[0130] In various embodiments, the antibodies or antigen-binding fragments thereof comprise a heavy chain variable domain comprising a sequence of amino acids that differs from the sequence of a heavy chain variable domain having the amino acid sequence set forth in SEQ ID NOs: 33-35 only at 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1 or 0 residues, wherein each such sequence difference is independently either a deletion, insertion, or substitution of one amino acid residue. In various embodiments, the antibodies or antigen-binding fragments thereof comprise a heavy chain variable domain comprising a sequence that has at least about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least about 99% identity to the amino acid sequence as set forth in SEQ ID NOs: 33-35.

[0131] In various embodiments, an isolated humanized antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises sequence of amino acids that differs from the sequence of a light chain variable domain having the amino acid sequence set forth in SEQ ID NOs: 36-37 only at 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1 or 0 residues, wherein each such sequence difference is independently either a deletion, insertion, or substitution of one amino acid residue. In various embodiments, the antibodies or antigen-binding fragments thereof comprise a light chain variable domain comprising a sequence that has at least about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least about 99% identity to the amino acid sequence as set forth in SEQ ID NOs: 36-37.

[0132] In various embodiments, an isolated humanized antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises a heavy chain having a sequence identical, substantially identical or substantially similar to SEQ ID NOs: 38, 42 and 44, and a light chain variable region having the sequence identical, substantially identical or substantially similar to SEQ ID NOs: 40 and 46.

[0133] In various embodiments of the present invention, the antibody is a humanized IgG comprising the heavy chain sequence ("H1") of SEQ ID NO: 38:

QVQLVQSGAEVKKPGASVKVSCKASGYTFTNYGMSWVRQAPGQGLEWMGWINTYS
 GESTYADDFKGRVTMTTDTSTSTAYMELRSLRSDDTAVYYCARSLNRDWTFTYWGQG
 TLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH
 TTPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCP
 PCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVH
 NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQP
 REPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD
 GSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK
 (SEQ ID NO: 38)

and the light chain sequence ("L3") of SEQ ID NO: 40:

QTVVTQEPSFSVSPGGTVTLTCRSITGAVTTSNYANWVQETPGQAPRGLIGGTNNRAP
 GVPDRFSGSILGNKAALTITGAQADDEADYFCALWYSNHWVFGGGTKLTVLGQPKAAP
 SVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTTPSKQSNNKY
 AASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS (SEQ ID NO: 40)

[0134] In certain alternative embodiments, the antibody is an antibody comprising a heavy chain and a light chain, wherein the heavy chain comprises a sequence that has at least about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least about 99% identity to the amino acid sequence as set forth in SEQ ID NO: 38, or its corresponding polynucleotide sequence SEQ ID NO: 39:

cagggtgcagctggtgcagctctggagcagaggtgaagaagccaggagcctctgtgaaggtagctgcaaggcctccggc
 tacacctcacaaactatggcatgagctgggtgcggcaggcacctggacagggactggagtgatgggctggatcaata
 catactccggcgagctctacctatgccgacgacttcaagggcaggggtgacaatgaccacagacacctctacaagcaccgc
 ctacatggagctgcggagcctgagatccgacgataccgccgtgtactattgtgccagggtccctgaaccgcgatacatggttc
 acctattggggccagggcacactggtgaccgtgagctccgccagcaccaagggcccttccgtgttccactggccccctcc
 tctaaatccacatctggcggcaccgccgcccctgggctgtctggtgaaggactacttcccagagcctgtgacagtgctctgga
 actctggcgccctgacatccggcgtgcacacatttccagccgtgtgcagagctccggcctgtacagcctgtctagcgtggt
 gacagtgccctcctctagcctgggacacagacctatctgcaacgtgaatcacaagccaagcaataccaaggtggac
 aagaaggtggagcccaagtcctgtgataagacacacacctgccccctgtcctgctcccagagctgctgggcggccctag
 cgtgttccgtgttccaccaagcctaaggacacctgatgatctcccgacacccgaggtgacctgctggtggtggacgtg
 tctcacgaggatcctgaggtgaagttcaactggtatgtggatggcgtggaggtgcacaatgccaagaccaagcccagag
 aggagcagtacaactctacatataggggtggtgagcgtgctgaccgtgctgcaccaggactggctgaacggcaaggagta
 taagtgaaggtgtccaataaggccctgccccgcccccatcgagaagacaatcagcaaggccaagggccagcctcggg
 agccacaggtgtacacctgcctccatccagagacgagctgacaaagaaccaggtgtctctgacatgtctggtgaaggg
 cttctatcctagcgatacgcctggagtgaggagtgccatggccagccagagagaacaattacaagaccacccccctgtgct

ggactccgatggctccttcttctgtattccaagctgaccgtggataagtctcggtggcagcagggcaacgtgttcagctgttc
cgatgatgcacgaagccctgcataatcactatactcagaaatccctgtccctgtcacctggaaag
(SEQ ID NO: 39)

and wherein the light chain comprises a sequence that has at least about 75%, at least about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least about 99% identity to the amino acid sequence as set forth in SEQ ID NO: 40, or its corresponding polynucleotide sequence SEQ ID NO: 41:

cagacagtggtagaccaggagcctagcttctccgtgtctccaggaggaaccgtgacactgacctgcaggtctatcacagg
cgccgtgaccacaagcaactacgccaattgggtgcaggagacaccaggacaggcaccaaggggactgatcgaggga
accaacaatagggcaccaggagtgctgaccggttcagcggatccatcctgggaacaaggccgcctgacaatcacc
ggagcacaggcagacgatgaggcagattacttctgtgcctgtggtattccaatcactgggtgttggcggcggcacaag
ctgaccgtgtgtgggcagcccaaggccgctcctagcgtgacactgttcccccttctccgaggagctgcaggccaaca
ggccaccctggtgtgctgatctccgacttctatcctggcgccgtgacagtggcctggaaggctgatttagccagtgaa
gtggtgggtggagaccacaacccctccaagcagtctaacaataagtatgccgttctcttacctgagcctgacaccaga
gcagtggaaagtcaccggtcttacagctgccaggtcactcacgaaggctctaccgtggaaaagacagtcgcaccacc
gaatgctcatgataagcttaagggttcgatccct (SEQ ID NO: 41)

[0135] In various embodiments of the present invention, the antibody is a humanized IgG comprising the heavy chain sequence ("H2") of SEQ ID NO: 42:

QVQLVQSGAEVKKPGASVKVSCKASGYTFTNYGMSWVRQAPGQGLEWMGWINTYS
GESTYADDFKGRFTMTLDTSTSTAYMELRSLRSDDTAVYFCARSLNRDWTFTYWGQG
TLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH
TFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCP
PCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVH
NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQP
REPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSD
GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO: 42)

and the light chain sequence ("L3") of SEQ ID NO: 40.

[0136] In certain alternative embodiments, the antibody is an antibody comprising a heavy chain and a light chain, wherein the heavy chain comprises a sequence that has at least about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least about 99% identity to the amino acid sequence as set forth in SEQ ID NO: 42, or its corresponding polynucleotide sequence SEQ ID NO: 43:

cagggtgcagctgggtgcagctctggagcagaggtgaagaagccaggagcctctgtgaagggtgagctgcaaggcctccggc
 tacaccttcacaaactatggcatgagctgggtgcggcaggcacctggacagggactggagtggatgggctggatcaata
 catactccggcgagcttacctatgccgacgattcaagggcaggtttaccatgacactggacacctctacaagcaccgcct
 acatggagctgcggagcctgagatccgacgatacagccgtgtattctgtgccaggctccctgaaccgcgatacatggtttac
 ctactggggccagggcacactggtgaccgtgagctccgccagcaccaagggcccttccgtgttccactggccccctcctc
 taaatccacatctggcggcaccgcccgcctgggctgtctggtgaaggactacttcccagagcctgtgacagtgtcctggaa
 ctctggcgccttgacatccggcgtgcacacatttccagccgtgctgcagagctccggcctgtacagcctgtctagcgtggg
 acagtgccctcctctagcctgggcacacagacctatactgcaacgtgaatcaccaagccaagcaataccaaggtggaca
 agaaggtggagcccaagtcctgtgataagacacacacctgcccccttctcctgtcctccgagctgctgggcggccctagc
 gtgtcctgtttccaccaagcctaaggacaccctgatgatctcccgacacccgaggtgacctgctggtgggtggacgtgt
 ctacagaggatcctgaggtgaagtcaactggtatgtggatggcgtggaggtgcacaatgccaaagccaagcccagaga
 ggagcagtacaactctacatataggggtgggtgagcgtgctgaccgtgctgcaccaggactggctgaacggcaaggagtat
 aagtgaagggtgtccaataaggccctgcccgcctccatcgagaagacaatcagcaaggccaagggccagcctcggga
 gccacaggtgtacacctgctccatccagagacgagctgacaaagaaccaggtgtctctgacatgtctggtgaagggt
 tctatcctagcgatatcgccgtggagtgaggagccaatggccagccagagaacaattacaagaccacccccctgtgctg
 gactccgatggctccttcttctgtattccaagctgaccgtggataagctcgggtggcagcagggcaacgtgttcagctgtcc
 gtgatgcacgaagccctgcataatcactatactcagaaatccctgtcctgtcacctggaaag (SEQ ID NO: 43)

and wherein the light chain comprises a sequence that has at least about 75%, at least about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least about 99% identity to the amino acid sequence as set forth in SEQ ID NO: 40, or its corresponding polynucleotide sequence SEQ ID NO: 41.

[0137] In various embodiments of the present invention, the antibody is a humanized IgG comprising the heavy chain sequence (“H3”) of SEQ ID NO: 44:

QVQLVQSGAEVKKPGASVKVSCASGYTFTNYGMSWVRQAPGQGLEWMGWINTYS
 GESTYADDFKGRFTFTLDTSTSTAYMELRSLRSDDTATYFCARSLNRDWTFTYWGQG
 TLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH
 TFPVAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCP
 PCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVH
 NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQP
 REPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD
 GSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO: 44)

and the light chain sequence (“L3”) of SEQ ID NO: 40.

[0138] In certain alternative embodiments, the antibody is an antibody comprising a heavy chain and a light chain, wherein the heavy chain comprises a sequence that has at least

about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least about 99% identity to the amino acid sequence as set forth in SEQ ID NO: 44, or its corresponding polynucleotide sequence SEQ ID NO: 45:

```
cagggtgcagctggtgcagtctggagcagaggtgaagaagccaggagcctctgtgaaggtagctgcaaggcctccggc
tacacctcacaaactatggcatgagctgggtgcggcaggcacctggacagggactggagtggatgggctggatcaata
cactactccggcgagctacatgacgactcaagggcaggttcacctttacactggacacctctacaagcaccgccta
catggagctgcggagcctgagatccgacgatacagccacctatttctgtgccagggtccctgaaccgcgatacatggttacc
tactggggccagggcacactgggtgaccgtgagctccgccagcaccaagggcccttccgtgttccactggccccctctct
aaatccacatctggcggcaccgcccctgggtgtctggtgaaggactacttcccagagcctgtgacagtgtcctggaac
tctggcgccctgacatccggcgtgcacacatttccagccgtgtgcagagctccggcctgtacagcctgtctagcgtggtga
cagtgcctctctagcctgggcacacagacctatatctgcaacgtgaatcacaagccaagcaataccaaggtggacaa
gaagggtggagcccaagtctgtgataagacacacacctgccccctgtctgtctcccgagctgtctggcgcccttagcgt
gttctgtttccaccaagcctaaggacaccctgatgatctcccgacacccgaggtgacctgctggtggtggacgtgtct
cacgaggatcctgaggtgaagttcaactggtatgtggtggcgtggaggtgcacaatgccaagaccaagcccagagag
gagcagtagaactctacatataggggtggtgagcgtgtgaccgtgtgcaccaggactggctgaacggcaaggagtata
agtgaagggtgtccaataaggccctgcccgcctcatcgagaagacaatcagcaaggccaagggccagcctcgggag
ccacaggtgtacacctgcctccatccagagacgagctgacaaagaaccaggtgtctctgacatgtctggtgaagggctc
tatcctagcgatatcgccgtggagtgaggagtcgaatggccagccagagaacaattacaagaccacacccctgtgtgg
actccgatggctccttcttctgtattccaagctgaccgtggataagtctcggtggcagcagggaacgtgttcagctgtccgt
gatgcacgaagccctgcataatactatactcagaaatccctgtccctgtcacctggaaag (SEQ ID NO: 45)
```

and wherein the light chain comprises a sequence that has at least about 75%, at least about 80%, at least about 85%, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least about 99% identity to the amino acid sequence as set forth in SEQ ID NO: 40, or its corresponding polynucleotide sequence SEQ ID NO: 41.

[0139] In various embodiments of the present disclosure, the antibody may be an anti-ET_AR antibody that has the same or higher antigen-binding affinity as that of the antibody comprising the heavy chain sequence as set forth in any of SEQ ID NOs: 38, 42 and 44. In various embodiments, the antibody may be an anti-ET_AR antibody which binds to the same epitope as the antibody comprising the heavy chain sequence as set forth in any of SEQ ID NOs: 38, 42 and 44. In various embodiments, the antibody is an anti-ET_AR antibody which competes with the antibody comprising the heavy chain sequence as set forth in any of SEQ ID NOs: 38, 42 and 44. In various embodiments, the antibody may be an anti-ET_AR antibody which

comprises at least one (such as two or three) CDRs of the heavy chain sequence as set forth in any of SEQ ID NOs: 38, 42 and 44.

[0140] In various embodiments, the antibody contains an heavy chain amino acid sequence that shares an observed homology of, *e.g.*, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% with any of SEQ ID NOs: 38, 42 and 44. In various embodiments, the antibody contains a nucleic acid sequence that shares an observed homology of, *e.g.*, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% with any of SEQ ID NOs: 39, 43 and 45.

[0141] In various embodiments of the present disclosure the antibody may be an anti-ET_AR antibody that has the same or higher antigen-binding affinity as that of the antibody comprising the light chain sequence as set forth in any of SEQ ID NOs: 40 and 46. In various embodiments, the antibody may be an anti-ET_AR antibody which binds to the same epitope as the antibody comprising the light chain sequence as set forth in any of SEQ ID NOs: 40 and 46. In various embodiments, the antibody is an anti-ET_AR antibody which competes with the antibody comprising the light chain sequence as set forth in any of SEQ ID NOs: 40 and 46. In various embodiments, the antibody may be an anti-ET_AR antibody which comprises at least one (such as two or three) CDRs of the light chain sequence as set forth in any of SEQ ID NOs: 40 and 46.

[0142] In various embodiments, the antibody contains an light chain amino acid sequence that shares an observed homology of, *e.g.*, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% with any of SEQ ID NOs: 40 and 46. In various embodiments, the antibody contains a nucleic acid sequence that shares an observed homology of, *e.g.*, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% with any of SEQ ID NOs: 41 and 47.

[0143] In various embodiments, an isolated humanized antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises a heavy chain variable region having a sequence identical, substantially identical or substantially similar to SEQ ID NOs: 38, 42 and 44, and a light chain variable region having the sequence identical,

substantially identical or substantially similar to SEQ ID NO: 40. In various embodiments, an isolated humanized antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises the heavy chain sequence set forth in SEQ ID NO: 38, and the light chain sequence set forth in SEQ ID NO: 40. In various embodiments, an isolated humanized antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises the heavy chain sequence set forth in SEQ ID NO: 42, and the light chain sequence set forth in SEQ ID NO: 40. In various embodiments, an isolated humanized antibody or antigen-binding fragment thereof of the present invention binds to human ET_AR and comprises the heavy chain sequence set forth in SEQ ID NO: 44, and the light chain sequence set forth in SEQ ID NO: 40.

[0144] Antibodies or antigen-binding fragments thereof of the invention can comprise any constant region known in the art. The light chain constant region can be, for example, a kappa- or lambda-type light chain constant region, e.g., a human kappa- or lambda-type light chain constant region. The heavy chain constant region can be, for example, an alpha-, delta-, epsilon-, gamma-, or mu-type heavy chain constant regions, e.g., a IgA-, IgD-, IgE-, IgG- and IgM-type heavy chain constant region. In various embodiments, the light or heavy chain constant region is a fragment, derivative, variant, or mutein of a naturally occurring constant region.

[0145] Techniques are known for deriving an antibody of a different subclass or isotype from an antibody of interest, i.e., subclass switching. Thus, IgG antibodies may be derived from an IgM antibody, for example, and vice versa. Such techniques allow the preparation of new antibodies that possess the antigen-binding properties of a given antibody (the parent antibody), but also exhibit biological properties associated with an antibody isotype or subclass different from that of the parent antibody. Recombinant DNA techniques may be employed. Cloned DNA encoding particular antibody polypeptides may be employed in such procedures, e.g., DNA encoding the constant domain of an antibody of the desired isotype. See also Lanitto et al., *Methods Mol. Biol.* 178:303-16, 2002.

[0146] In various embodiments, an antibody of the invention further comprises a light chain kappa or lambda constant domain, or a fragment thereof, and further comprises a heavy chain constant domain, or a fragment thereof. Sequences of the light chain constant region and

heavy chain constant region used in the exemplified antibodies, and polynucleotides encoding them, are provided below.

Light Chain (Kappa) Constant Region

RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKD
STYLSSTLTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 48)

Light Chain (Lambda) Constant Region

GQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSN
NKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS (SEQ ID NO: 49)

Heavy Chain Constant Region

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY
SLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFP
PKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT
VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKG
FYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFSCSVMHEALH
NHYTQKSLSLSPGK (SEQ ID NO: 50)

[0147] Antibodies of the present invention may also be described or specified in terms of their cross-reactivity. Antibodies that bind ET_ARs, which have at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55%, and at least 50% identity (as calculated using methods known in the art and described herein) to human ET_AR are also included in the present invention.

[0148] Further included in the present invention are antibodies that bind to the same epitope as the anti-ET_AR antibodies of the present invention. To determine if an antibody can compete for binding to the same epitope as the epitope bound by the anti-ET_AR antibodies of the present invention, a cross-blocking assay, e.g., a competitive ELISA assay, can be performed. In an exemplary competitive ELISA assay, ET_AR coated on the wells of a microtiter plate is pre-incubated with or without candidate competing antibody and then the biotin labeled anti-ET_AR antibody of the invention is added. The amount of labeled anti-ET_AR antibody bound to the ET_AR antigen in the wells is measured using avidin-peroxidase conjugate and appropriate substrate. The antibody can be labeled with a radioactive or fluorescent label or some other

detectable and measurable label. The amount of labeled anti-ET_AR antibody that bound to the antigen will have an indirect correlation to the ability of the candidate competing antibody (test antibody) to compete for binding to the same epitope, i.e., the greater the affinity of the test antibody for the same epitope, the less labeled antibody will be bound to the antigen-coated wells. A candidate competing antibody is considered an antibody that binds substantially to the same epitope or that competes for binding to the same epitope as an anti-ET_AR antibody of the invention if the candidate antibody can block binding of the ET_AR antibody by at least 20%, by at least 30%, by at least 40%, or by at least 50% as compared to the control performed in parallel in the absence of the candidate competing antibody. It will be understood that variations of this assay can be performed to arrive at the same quantitative value.

[0149] In certain alternative embodiments, the antibodies of the present invention can be engineered by modifying one or more residues within one or both variable regions (i.e., V_H and/or V_L), or by modifying residues within the constant region(s), e.g., to alter the effector function(s) of the antibody. In various embodiments, the variable region of the antibody will be modified by performing CDR grafting using framework sequences can be obtained from public DNA databases or published references that include germline antibody gene sequences (e.g., Tomlinson, I. M., et al., J. Mol. Biol. 227:776-798, 1992; and Cox, J. P. L. et al., Eur. J. Immunol. 24:827-836, 1994; the contents of each of which are expressly incorporated herein by reference). In various embodiments, the antibodies may be modified using site-directed mutagenesis or PCR-mediated mutagenesis to introduce a mutation(s) in the V_H and/or V_L which improves binding affinity and/or decreases immunogenicity. In various embodiments, the antibodies may be modified in the Fc region for purposes of altering the serum half-life, complement fixation, Fc receptor binding, and/or antigen-dependent cellular cytotoxicity of the antibody. In various embodiments, the antibodies may be modified for purposes of modifying the glycosylation of the antibody. Methods for performing each of the modifications described herein, and others, are well known to the skilled artisan.

Pharmaceutical Compositions

[0150] In another aspect, the present invention provides a pharmaceutical composition comprising an antibody or antigen-binding fragment thereof as described above. The

pharmaceutical compositions, methods and uses of the invention thus also encompass embodiments of combinations (co-administration) with other active agents, as detailed below.

[0151] Generally, the antibodies, or antigen-binding fragments thereof antibodies of the present invention are suitable to be administered as a formulation in association with one or more pharmaceutically acceptable excipient(s). The term 'excipient' is used herein to describe any ingredient other than the compound(s) of the invention. The choice of excipient(s) will to a large extent depend on factors such as the particular mode of administration, the effect of the excipient on solubility and stability, and the nature of the dosage form. As used herein, "pharmaceutically acceptable excipient" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Some examples of pharmaceutically acceptable excipients are water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Additional examples of pharmaceutically acceptable substances are wetting agents or minor amounts of auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or effectiveness of the antibody. Pharmaceutical compositions of the present invention and methods for their preparation will be readily apparent to those skilled in the art. Such compositions and methods for their preparation may be found, for example, in Remington's Pharmaceutical Sciences, 19th Edition (Mack Publishing Company, 1995). Pharmaceutical compositions are preferably manufactured under GMP conditions.

[0152] A pharmaceutical composition of the invention may be prepared, packaged, or sold in bulk, as a single unit dose, or as a plurality of single unit doses. As used herein, a "unit dose" is discrete amount of the pharmaceutical composition comprising a predetermined amount of the active ingredient. The amount of the active ingredient is generally equal to the dosage of the active ingredient which would be administered to a subject or a convenient fraction of such a dosage such as, for example, one-half or one-third of such a dosage.

[0153] Any method for administering peptides, proteins or antibodies accepted in the art may suitably be employed for the antibodies and portions of the invention.

[0154] The pharmaceutical compositions of the invention are typically suitable for parenteral administration. As used herein, "parenteral administration" of a pharmaceutical composition includes any route of administration characterized by physical breaching of a tissue of a subject and administration of the pharmaceutical composition through the breach in the tissue, thus generally resulting in the direct administration into the blood stream, into muscle, or into an internal organ. Parenteral administration thus includes, but is not limited to, administration of a pharmaceutical composition by injection of the composition, by application of the composition through a surgical incision, by application of the composition through a tissue-penetrating non-surgical wound, and the like. In particular, parenteral administration is contemplated to include, but is not limited to, subcutaneous, intraperitoneal, intramuscular, intrasternal, intravenous, intraarterial, intrathecal, intraventricular, intraurethral, intracranial, intrasynovial injection or infusions; and kidney dialytic infusion techniques. Various embodiments include the intravenous and the subcutaneous routes.

[0155] Formulations of a pharmaceutical composition suitable for parenteral administration typically generally comprise the active ingredient combined with a pharmaceutically acceptable carrier, such as sterile water or sterile isotonic saline. Such formulations may be prepared, packaged, or sold in a form suitable for bolus administration or for continuous administration. Injectable formulations may be prepared, packaged, or sold in unit dosage form, such as in ampoules or in multi-dose containers containing a preservative. Formulations for parenteral administration include, but are not limited to, suspensions, solutions, emulsions in oily or aqueous vehicles, pastes, and the like. Such formulations may further comprise one or more additional ingredients including, but not limited to, suspending, stabilizing, or dispersing agents. In one embodiment of a formulation for parenteral administration, the active ingredient is provided in dry (i.e. powder or granular) form for reconstitution with a suitable vehicle (e.g. sterile pyrogen-free water) prior to parenteral administration of the reconstituted composition. Parenteral formulations also include aqueous solutions which may contain excipients such as salts, carbohydrates and buffering agents (preferably to a pH of from 3 to 9), but, for some applications, they may be more suitably formulated as a sterile non-aqueous solution or as a dried form to be used in conjunction with a suitable vehicle such as sterile, pyrogen-free water. Exemplary parenteral administration forms include solutions or

suspensions in sterile aqueous solutions, for example, aqueous propylene glycol or dextrose solutions. Such dosage forms can be suitably buffered, if desired. Other parentally administrable formulations which are useful include those which comprise the active ingredient in microcrystalline form, or in a liposomal preparation. Formulations for parenteral administration may be formulated to be immediate and/or modified release. Modified release formulations include delayed-, sustained-, pulsed-, controlled-, targeted and programmed release.

[0156] For example, in one aspect, sterile injectable solutions can be prepared by incorporating the anti-ETR antibody in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The proper fluidity of a solution can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prolonged absorption of injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

[0157] The antibodies of the invention can also be administered intranasally or by inhalation, typically in the form of a dry powder (either alone, as a mixture, or as a mixed component particle, for example, mixed with a suitable pharmaceutically acceptable excipient) from a dry powder inhaler, as an aerosol spray from a pressurized container, pump, spray, atomizer (preferably an atomizer using electrohydrodynamics to produce a fine mist), or nebulizer, with or without the use of a suitable propellant, or as nasal drops.

[0158] The pressurized container, pump, spray, atomizer, or nebulizer generally contains a solution or suspension of an antibody of the invention comprising, for example, a suitable agent for dispersing, solubilizing, or extending release of the active, a propellant(s) as solvent.

[0159] Prior to use in a dry powder or suspension formulation, the drug product is generally micronized to a size suitable for delivery by inhalation (typically less than 5 microns). This may be achieved by any appropriate comminuting method, such as spiral jet milling, fluid bed jet milling, supercritical fluid processing to form nanoparticles, high pressure homogenization, or spray drying.

[0160] Capsules, blisters and cartridges for use in an inhaler or insufflator may be formulated to contain a powder mix of the compound of the invention, a suitable powder base and a performance modifier.

[0161] Suitable flavours, such as menthol and levomenthol, or sweeteners, such as saccharin or saccharin sodium, may be added to those formulations of the invention intended for inhaled/intranasal administration.

[0162] Formulations for inhaled/intranasal administration may be formulated to be immediate- and/or modified release. Modified release formulations include delayed-, sustained-, pulsed-, controlled-, targeted and programmed release.

[0163] In the case of dry powder inhalers and aerosols, the dosage unit is determined by means of a valve which delivers a metered amount. Units in accordance with the invention are typically arranged to administer a metered dose or "puff" of an antibody of the invention. The overall daily dose will typically be administered in a single dose or, more usually, as divided doses throughout the day.

[0164] The antibodies and antibody portions of the invention may also be formulated for an oral route administration. Oral administration may involve swallowing, so that the compound enters the gastrointestinal tract, and/or buccal, lingual, or sublingual administration by which the compound enters the blood stream directly from the mouth.

[0165] Formulations suitable for oral administration include solid, semi-solid and liquid systems such as tablets; soft or hard capsules containing multi- or nano-particulates, liquids, or powders; lozenges (including liquid-filled); chews; gels; fast dispersing dosage forms; films; ovules; sprays; and buccal/mucoadhesive patches.

[0166] Pharmaceutical compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening

agents in order to provide a pharmaceutically elegant and palatable preparation. For example, to prepare orally deliverable tablets, the antibody or antigen-binding fragment thereof is mixed with at least one pharmaceutical excipient, and the solid formulation is compressed to form a tablet according to known methods, for delivery to the gastrointestinal tract. The tablet composition is typically formulated with additives, e.g. a saccharide or cellulose carrier, a binder such as starch paste or methyl cellulose, a filler, a disintegrator, or other additives typically usually used in the manufacture of medical preparations. To prepare orally deliverable capsules, DHEA is mixed with at least one pharmaceutical excipient, and the solid formulation is placed in a capsular container suitable for delivery to the gastrointestinal tract. Compositions comprising antibodies or antigen-binding fragments thereof may be prepared as described generally in Remington's Pharmaceutical Sciences, 18th Ed. 1990 (Mack Publishing Co. Easton Pa. 18042) at Chapter 89, which is herein incorporated by reference.

[0167] In various embodiments, the pharmaceutical compositions are formulated as orally deliverable tablets containing antibodies or antigen-binding fragments thereof in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for manufacture of tablets. These excipients may be inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, maize starch, gelatin or acacia, and lubricating agents, for example, magnesium stearate, stearic acid, or talc. The tablets may be uncoated or they may be coated with known techniques to delay disintegration and absorption in the gastrointestinal track and thereby provide a sustained action over a longer period of time. For example, a time delay material such as glyceryl monostearate or glyceryl distearate alone or with a wax may be employed.

[0168] In various embodiments, the pharmaceutical compositions are formulated as hard gelatin capsules wherein the antibody or antigen-binding fragment thereof is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate, or kaolin or as soft gelatin capsules wherein the antibody or antigen-binding fragment thereof is mixed with an aqueous or an oil medium, for example, arachis oil, peanut oil, liquid paraffin or olive oil.

[0169] Liquid formulations include suspensions, solutions, syrups and elixirs. Such formulations may be employed as fillers in soft or hard capsules (made, for example, from gelatin or hydroxypropylmethylcellulose) and typically comprise a carrier, for example, water,

ethanol, polyethylene glycol, propylene glycol, methylcellulose, or a suitable oil, and one or more emulsifying agents and/or suspending agents. Liquid formulations may also be prepared by the reconstitution of a solid, for example, from a sachet.

Therapeutic Uses

[0170] In another aspect, the present invention relates to methods of treating a subject suffering from a ET_AR-associated disorder, comprising administering to said subject a therapeutically effective amount (either as monotherapy or in a combination therapy regimen) of an antibody or antigen-binding fragment thereof of the present invention. In various embodiments, the subject is a human subject.

[0171] In various embodiments, the ET_AR-associated disorder is a metabolic disorder selected from the group consisting of diabetes, diabetic myopathy, diabetic nephropathy, diabetic neuropathy, diabetic retinopathy, obesity, dyslipidemia, sarcopenic obesity, fatty liver disease (e.g., nonalcoholic fatty liver disease (NAFLD); nonalcoholic steatohepatitis (NASH)), insulin resistance, and hemochromatosis.

[0172] In various embodiments, the ET_AR-associated disorder is a cardiovascular disease selected from the group consisting of angina, heart attack (myocardial infarction), atherosclerosis, heart failure, cardiovascular disease, rheumatic heart disease, cardiac arrhythmias (abnormal heart rhythms), cerebrovascular disease, congenital heart defects, cardiomyopathy, left ventricular hypertrophy, right ventricular hypertrophy, post-infarction heart rupture, infections of the heart, coronary artery disease, peripheral arterial disease, renal artery stenosis, aortic aneurysm, myocardial diseases, heart valve disorders, myocarditis, pericarditis, and pulmonary arterial hypertension (PAH).

[0173] "Therapeutically effective amount" or "therapeutically effective dose" refers to that amount of the therapeutic agent being administered which will relieve to some extent one or more of the symptoms of the disorder being treated.

[0174] A therapeutically effective dose can be estimated initially from cell culture assays by determining an IC₅₀. A dose can then be formulated in animal models to achieve a circulating plasma concentration range that includes the IC₅₀ as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in

plasma may be measured, for example, by HPLC. The exact composition, route of administration and dosage can be chosen by the individual physician in view of the subject's condition.

[0175] Dosage regimens can be adjusted to provide the optimum desired response (e.g., a therapeutic or prophylactic response). For example, a single bolus can be administered, several divided doses (multiple or repeat or maintenance) can be administered over time and the dose can be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the present disclosure will be dictated primarily by the unique characteristics of the antibody and the particular therapeutic or prophylactic effect to be achieved.

[0176] Thus, the skilled artisan would appreciate, based upon the disclosure provided herein, that the dose and dosing regimen is adjusted in accordance with methods well-known in the therapeutic arts. That is, the maximum tolerable dose can be readily established, and the effective amount providing a detectable therapeutic benefit to a subject may also be determined, as can the temporal requirements for administering each agent to provide a detectable therapeutic benefit to the subject. Accordingly, while certain dose and administration regimens are exemplified herein, these examples in no way limit the dose and administration regimen that may be provided to a subject in practicing the present disclosure.

[0177] It is to be noted that dosage values may vary with the type and severity of the condition to be alleviated and may include single or multiple doses. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that dosage ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition. Further, the dosage regimen with the compositions of this disclosure may be

based on a variety of factors, including the type of disease, the age, weight, sex, medical condition of the subject, the severity of the condition, the route of administration, and the particular antibody employed. Thus, the dosage regimen can vary widely, but can be determined routinely using standard methods. For example, doses may be adjusted based on pharmacokinetic or pharmacodynamic parameters, which may include clinical effects such as toxic effects and/or laboratory values. Thus, the present disclosure encompasses intra-subject dose-escalation as determined by the skilled artisan. Determining appropriate dosages and regimens are well-known in the relevant art and would be understood to be encompassed by the skilled artisan once provided the teachings disclosed herein.

[0178] For administration to human subjects, the total monthly dose of the antibodies or antigen-binding fragments thereof of the disclosure can be in the range of 0.5-1200 mg per subject, 0.5-1100 mg per subject, 0.5-1000 mg per subject, 0.5-900 mg per subject, 0.5-800 mg per subject, 0.5-700 mg per subject, 0.5-600 mg per subject, 0.5-500 mg per subject, 0.5-400 mg per subject, 0.5-300 mg per subject, 0.5-200 mg per subject, 0.5-100 mg per subject, 0.5-50 mg per subject, 1-1200 mg per subject, 1-1100 mg per subject, 1-1000 mg per subject, 1-900 mg per subject, 1-800 mg per subject, 1-700 mg per subject, 1-600 mg per subject, 1-500 mg per subject, 1-400 mg per subject, 1-300 mg per subject, 1-200 mg per subject, 1-100 mg per subject, or 1-50 mg per subject depending, of course, on the mode of administration. For example, an intravenous monthly dose can require about 1-1000 mg/subject. In various embodiments, the antibodies or antigen-binding fragments thereof of the disclosure can be administered at about 1-200 mg per subject, 1-150 mg per subject or 1-100 mg/subject. The total monthly dose can be administered in single or divided doses and can, at the physician's discretion, fall outside of the typical ranges given herein.

[0179] In various embodiments, a non-limiting daily dosing range for a therapeutically or prophylactically effective amount of an antibody or antigen-binding fragment thereof of the disclosure can be 0.001 to 100 mg/kg, 0.001 to 90 mg/kg, 0.001 to 80 mg/kg, 0.001 to 70 mg/kg, 0.001 to 60 mg/kg, 0.001 to 50 mg/kg, 0.001 to 40 mg/kg, 0.001 to 30 mg/kg, 0.001 to 20 mg/kg, 0.001 to 10 mg/kg, 0.001 to 5 mg/kg, 0.001 to 4 mg/kg, 0.001 to 3 mg/kg, 0.001 to 2 mg/kg, 0.001 to 1 mg/kg, 0.010 to 50 mg/kg, 0.010 to 40 mg/kg, 0.010 to 30 mg/kg, 0.010 to 20 mg/kg, 0.010 to 10 mg/kg, 0.010 to 5 mg/kg, 0.010 to 4 mg/kg, 0.010 to 3 mg/kg, 0.010 to 2

mg/kg, 0.010 to 1 mg/kg, 0.1 to 50 mg/kg, 0.1 to 40 mg/kg, 0.1 to 30 mg/kg, 0.1 to 20 mg/kg, 0.1 to 10 mg/kg, 0.1 to 5 mg/kg, 0.1 to 4 mg/kg, 0.1 to 3 mg/kg, 0.1 to 2 mg/kg, 0.1 to 1 mg/kg, 1 to 50 mg/kg, 1 to 40 mg/kg, 1 to 30 mg/kg, 1 to 20 mg/kg, 1 to 10 mg/kg, 1 to 5 mg/kg, 1 to 4 mg/kg, 1 to 3 mg/kg, 1 to 2 mg/kg, or 1 to 1 mg/kg body weight.

[0180] For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of symptoms occurs or until sufficient therapeutic levels are achieved, for example, to reduce pain. An exemplary dosing regimen comprises administering an initial dose of about 2 mg/kg, followed by a weekly maintenance dose of about 1 mg/kg of the anti-ET_AR antibody, or followed by a maintenance dose of about 1 mg/kg every other week. However, other dosage regimens may be useful, depending on the pattern of pharmacokinetic decay that the practitioner wishes to achieve. For example, in some embodiments, dosing from one-four times a week is contemplated. The progress of this therapy is easily monitored by conventional techniques and assays. The dosing regimen (including the ET_AR antagonist(s) used) can vary over time. In various embodiments, the appropriate dosage of an anti-ET_AR antagonist antibody will depend on the anti-ET_AR antagonist antibody (or compositions thereof) employed, the type and severity of headache (e.g., migraine) to be treated, whether the agent is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the agent, and the discretion of the attending physician. Typically, the clinician will administer an anti-ET_AR antagonist antibody, until a dosage is reached that achieves the desired result. Dose and/or frequency can vary over course of treatment.

[0181] It is to be noted that dosage values may vary with the type and severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that dosage ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition.

[0182] In various embodiments, the total dose administered will achieve a plasma antibody concentration in the range of, e.g., about 1 to 1000 µg/ml, about 1 to 750 µg/ml, about 1 to 500 µg/ml, about 1 to 250 µg/ml, about 10 to 1000 µg/ml, about 10 to 750 µg/ml, about 10

to 500 µg/ml, about 10 to 250 µg/ml, about 20 to 1000 µg/ml, about 20 to 750 µg/ml, about 20 to 500 µg/ml, about 20 to 250 µg/ml, about 30 to 1000 µg/ml, about 30 to 750 µg/ml, about 30 to 500 µg/ml, about 30 to 250 µg/ml.

[0183] Toxicity and therapeutic index of the pharmaceutical compositions of the invention can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effective dose is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. Compositions that exhibit large therapeutic indices are generally preferred.

[0184] In various embodiments, single or multiple administrations of the pharmaceutical compositions are administered depending on the dosage and frequency as required and tolerated by the subject. In any event, the composition should provide a sufficient quantity of at least one of the antibodies or antigen-binding fragments thereof disclosed herein to effectively treat the subject. The dosage can be administered once but may be applied periodically until either a therapeutic result is achieved or until side effects warrant discontinuation of therapy.

[0185] The dosing frequency of the administration of the antibody or antigen-binding fragment thereof pharmaceutical composition depends on the nature of the therapy and the particular disease being treated. The subject can be treated at regular intervals, such as weekly or monthly, until a desired therapeutic result is achieved. Exemplary dosing frequencies include, but are not limited to: once weekly without break; once weekly, every other week; once every 2 weeks; once every 3 weeks; weakly without break for 2 weeks, then monthly; weakly without break for 3 weeks, then monthly; monthly; once every other month; once every three months; once every four months; once every five months; or once every six months, or yearly.

Combination Therapy

[0186] As used herein, the terms "co-administration", "co-administered" and "in combination with", referring to the antibodies or antigen-binding fragments thereof of the disclosure and one or more other therapeutic agents, is intended to mean, and does refer to and include the following: simultaneous administration of such combination of antibodies or antigen-binding fragments thereof of the disclosure and therapeutic agent(s) to a subject in need of

treatment, when such components are formulated together into a single dosage form which releases said components at substantially the same time to said subject; substantially simultaneous administration of such combination of antibodies or antigen-binding fragments thereof of the disclosure and therapeutic agent(s) to a subject in need of treatment, when such components are formulated apart from each other into separate dosage forms which are taken at substantially the same time by said subject, whereupon said components are released at substantially the same time to said subject; sequential administration of such combination of antibodies or antigen-binding fragments thereof of the disclosure and therapeutic agent(s) to a subject in need of treatment, when such components are formulated apart from each other into separate dosage forms which are taken at consecutive times by said subject with a significant time interval between each administration, whereupon said components are released at substantially different times to said subject; and sequential administration of such combination of antibodies or antigen-binding fragments thereof of the disclosure and therapeutic agent(s) to a subject in need of treatment, when such components are formulated together into a single dosage form which releases said components in a controlled manner whereupon they are concurrently, consecutively, and/or overlappingly released at the same and/or different times to said subject, where each part may be administered by either the same or a different route.

[0187] In another aspect, the present invention relates to combination therapies designed to treat a metabolic disorder in a subject, comprising administering to the subject a therapeutically effective amount of an isolated antibody or antigen-binding fragment of the present invention, and (b) administering a glucose-lowering agent. In various embodiments, the glucose-lowering agent is selected from anti-diabetic agents, anti-hyperglycemic agents, lipid lowering agents, anti-hypertensive agents, and anti-obesity agents. In various embodiments, the glucose-lowering agent is selected from biguanides, sulfonylureas, meglitinides, thiazolidinediones (TZDs), α -glucosidase inhibitors, DPP-4 inhibitors, bile acid sequestrants, dopamine-2 agonists, SGLT2 inhibitors, GLP-1R agonists, amylin mimetics, and insulins.

[0188] In another aspect, the present invention provides combination therapies designed to treat a cardiovascular disease in a subject, comprising (a) administering to said subject a therapeutically effective amount of an isolated antibody or antigen-binding fragment of the present invention, and (b) administering a second agent selected from the group consisting

of antihypertensive agents, agents for treating chronic heart failure, atherosclerosis or related diseases. In various embodiments such agents contemplated for use include, but are not limited to, bimecromol, angiotensin-converting enzyme inhibitors (such as captopril, enalapril, fosinopril, lisinopril, perindopril, quinapril, ramipril and the like), neutral endopeptidase inhibitors (such as thiorphan, omapatrilat, MDL-100240, fasidotril, sampatrilat, GW-660511, mixanpril, SA-7060, E-4030, SLV-306, ecadotril and the like), angiotensin II receptor antagonists (such as candesartan cilexetil, eprosartan, irbesartan, losartan, olmesartan medoxomil, telmisartan, valsartan, tasosartan, enoltasartan and the like), endothelin-converting enzyme inhibitors (such as CGS 35066, CGS 26303, CGS-31447, SM-19712 and the like), endothelin receptor antagonists (such as traclear, sitaxsentan, ambrisentan, L-749805, TBC-3214, BMS-182874, BQ-610, TA-0201, SB-215355, PD-180988, BMS-193884, darusentan, TBC-3711, bosentan, tezosentan, J-104132, YM-598, S-0139, SB-234551, RPR-118031A, ATZ-1993, RO-61-1790, ABT-546, enlasentan, BMS-207940 and the like), diuretic agents (such as hydrochlorothiazide, bendroflumethiazide, trichlormethiazide, indapamide, metolazone, furosemide, bumetamide, torsemide, chlorthalidone, metolazone, cyclopenthiazide, hydroflumethiazide, triamide, mefruside, benzylhydrochlorothiazide, penflutizide, methyclothiazide, azosemide, etacrynic acid, torasemide, piretamide, meticrane, potassium canrenoate, spironolactone, triamterene, aminophylline, cicletanine, LLU-alpha, PNU-80873A, isosorbide, D-mannitol, D-sorbitol, fructose, glycerin, acetazolamide, methazolamide, FR-179544, OPC-31260, lixivaptan, conivaptan and the like), calcium channel antagonists (such as amlodipine, bepridil, diltiazem, felodipine, isradipine, nicardipen, nimodipine, verapamil, S-verapamil, aranidipine, efonidipine, barnidipine, benidipine, manidipine, cilnidipine, nisoldipine, nitrendipine, nifedipine, nilvadipine, felodipine, pranidipine, lercanidipine, isradipine, elgodipine, azelnidipine, lacidipine, vatanidipine, lemdipine, diltiazem, clentiazem, fasudil, bepridil, gallopamil and the like), vasodilating antihypertensive agents (such as indapamide, todralazine, hydralazine, cadralazine, budralazine and the like), beta blockers (such as acebutolol, bisoprolol, esmolol, propranolol, atenolol, labetalol, carvedilol, metoprolol and the like), sympathetic blocking agents (such as amosulalol, terazosin, bunazosin, prazosin, doxazosin, propranolol, atenolol, metoprolol, carvedilol, nipradilol, celiprolol, nebivolol, betaxolol, pindolol, tertatolol, bevantolol, timolol, carteolol, bisoprolol, bopindolol, nipradilol, penbutolol, acebutolol, tilisolol, nadolol,

urapidil, indoramin and the like), alpha-2-adrenoceptor agonists (such as clonidine, methyldopa, CHF-1035, guanabenz acetate, guanfacine, moxonidine, lofexidine, talipexole and the like), centrally acting antihypertensive agents (such as reserpine and the like), thrombocyte aggregation inhibitors (such as warfarin, dicumarol, phenprocoumon, acenocoumarol, anisindione, phenindione, ximelagatran and the like), antiplatelets agents (such as aspirin, clopidogrel, ticlopidine, dipyridamole, cilostazol, ethyl icosapentate, sarpogrelate, dilazep, trapidil, beraprost and the like), and neuregulins (NRG-1, NRG-2, NRG-3 and NRG-4) and isoforms thereof.

[0189] In various embodiments, the combination therapy comprises administering the antibody or antigen-binding fragment thereof and the one or more additional therapies simultaneously. In various embodiments, antibody or antigen-binding fragment thereof composition and the one or more additional therapies are administered sequentially, i.e., the antibody or antigen-binding fragment thereof composition is administered either prior to or after the administration of the one or more additional therapies.

[0190] In various embodiments, the administrations of the antibody or antigen-binding fragment thereof composition and the one or more additional therapies are concurrent, i.e., the administration period of the antibody or antigen-binding fragment thereof composition and the one or more additional therapies overlap with each other.

[0191] In various embodiments, the administrations of the antibody or antigen-binding fragment thereof composition and the one or more additional therapies are non-concurrent. For example, in various embodiments, the administration of the antibody or antigen-binding fragment thereof composition is terminated before the one or more additional therapies is administered. In various embodiments, the administration of the one or more additional therapies is terminated before the antibody or antigen-binding fragment thereof composition is administered.

[0192] When the antibody or antigen-binding fragment thereof disclosed herein is administered in combination with one or more additional therapies, either concomitantly or sequentially, such antibody or antigen-binding fragment thereof may enhance the therapeutic effect of the one or more additional therapies or overcome cellular resistance to the one or more additional therapies. This allows for decreased dosage or duration of the one or more additional

therapies, thereby reducing the undesirable side effects, or restores the effectiveness of the one or more additional therapies.

Diagnostic Uses

[0193] In another aspect, the present invention provides a method for detecting in vitro or in vivo the presence of human ETR peptide in a sample, e.g., for diagnosing a human ETR-related disorder. In some methods, this is achieved by contacting a sample to be tested, along with a control sample, with a human sequence antibody or a human monoclonal antibody of the invention, or an antigen-binding portion thereof (or a bispecific or multispecific molecule), under conditions that allow for formation of a complex between the antibody and human ETR. Complex formation is then detected (e.g., using an ELISA) in both samples, and any statistically significant difference in the formation of complexes between the samples is indicative the presence of human ETR antigen in the test sample.

[0194] In various embodiments, methods are provided for detecting a ETR-related disorder or confirming the diagnosis of a ETR-related disorder in a subject. The method includes contacting a biological sample from the subject with an isolated antibody or antigen-binding fragment thereof of the invention and detecting binding of the isolated human monoclonal antibody or antigen-binding fragment thereof to the sample. An increase in binding of the isolated human monoclonal antibody or antigen-binding fragment thereof to the sample as compared to binding of the isolated human monoclonal antibody or antigen-binding fragment thereof to a control sample detects a ETR-related disorder in the subject or confirms the diagnosis of a ETR-related disorder in the subject. The control can be a sample from a subject known not to have a ETR-related disorder, or a standard value. The sample can be any sample, including, but not limited to, tissue from biopsies, autopsies and pathology specimens. Biological samples also include sections of tissues, for example, frozen sections taken for histological purposes. Biological samples further include body fluids, such as blood, serum, plasma, sputum, and spinal fluid.

[0195] In one embodiment, a kit is provided for detecting ETR in a biological sample, such as a blood sample. Kits for detecting a polypeptide will typically comprise a human antibody that specifically binds ETR, such as any of the antibodies disclosed herein. In some

embodiments, an antibody fragment, such as an Fv fragment is included in the kit. For *in vivo* uses, the antibody can be a scFv fragment. In a further embodiment, the antibody is labeled (for example, with a fluorescent, radioactive, or an enzymatic label).

[0196] In one embodiment, a kit includes instructional materials disclosing means of use of an antibody that specifically binds ETR. The instructional materials may be written, in an electronic form (such as a computer diskette or compact disk) or may be visual (such as video files). The kits may also include additional components to facilitate the particular application for which the kit is designed. Thus, for example, the kit may additionally contain means of detecting a label (such as enzyme substrates for enzymatic labels, filter sets to detect fluorescent labels, appropriate secondary labels such as a secondary antibody, or the like). The kits may additionally include buffers and other reagents routinely used for the practice of a particular method. Such kits and appropriate contents are well known to those of skill in the art.

[0197] In one embodiment, the diagnostic kit comprises an immunoassay. Although the details of the immunoassays may vary with the particular format employed, the method of detecting ETR in a biological sample generally includes the steps of contacting the biological sample with an antibody which specifically reacts, under immunologically reactive conditions, to ETR. The antibody is allowed to specifically bind under immunologically reactive conditions to form an immune complex, and the presence of the immune complex (bound antibody) is detected directly or indirectly.

[0198] In various embodiments, the antibodies or antigen-binding fragments can be labeled or unlabeled for diagnostic purposes. Typically, diagnostic assays entail detecting the formation of a complex resulting from the binding of an antibody to ETR. The antibodies can be directly labeled. A variety of labels can be employed, including, but not limited to, radionuclides, fluorescers, enzymes, enzyme substrates, enzyme cofactors, enzyme inhibitors and ligands (e.g., biotin, haptens). Numerous appropriate immunoassays are known to the skilled artisan (see, for example, U.S. Patent Nos. 3,817,827; 3,850,752; 3,901,654; and 4,098,876). When unlabeled, the antibodies can be used in assays, such as agglutination assays. Unlabeled antibodies can also be used in combination with another (one or more) suitable reagent which can be used to detect antibody, such as a labeled antibody (e.g., a second antibody) reactive

with the first antibody (e.g., anti-idiotypic antibodies or other antibodies that are specific for the unlabeled immunoglobulin) or other suitable reagent (e.g., labeled protein A).

[0199] The antibody or antigen-binding fragment provided herein may also be used in a method of detecting the susceptibility of a mammal to certain diseases. To illustrate, the method can be used to detect the susceptibility of a mammal to diseases which progress based on the amount of ETR present on cells and/or the number of ETR-positive cells in a mammal. In one embodiment, the application provides a method of detecting susceptibility of a mammal to a tumor. In this embodiment, a sample to be tested is contacted with an antibody which binds to ETR or portion thereof under conditions appropriate for binding of said antibody thereto, wherein the sample comprises cells which express ETR in normal individuals. The binding of antibody and/or amount of binding is detected, which indicates the susceptibility of the individual to a tumor, wherein higher levels of receptor correlate with increased susceptibility of the individual to a tumor.

[0200] In various embodiments, the antibodies or antigen-binding fragments are attached to a label that is able to be detected (e.g., the label can be a radioisotope, fluorescent compound, enzyme or enzyme co-factor). The active moiety may be a radioactive agent, such as: radioactive heavy metals such as iron chelates, radioactive chelates of gadolinium or manganese, positron emitters of oxygen, nitrogen, iron, carbon, or gallium, ^{43}K , ^{52}Fe , ^{57}Co , ^{67}Cu , ^{67}Ga , ^{68}Ga , ^{123}I , ^{125}I , ^{131}I , ^{132}I , or ^{99}Tc . A binding agent affixed to such a moiety may be used as an imaging agent and is administered in an amount effective for diagnostic use in a mammal such as a human and the localization and accumulation of the imaging agent is then detected. The localization and accumulation of the imaging agent may be detected by radioscintigraphy, nuclear magnetic resonance imaging, computed tomography or positron emission tomography.

[0201] Immunoscintigraphy using antibodies or antigen-binding fragments directed at ETR may be used to detect and/or diagnose cancers and vasculature. For example, monoclonal antibodies against the ETR marker labeled with ^{99}Tc Technetium, ^{111}In Indium, or ^{125}I Iodine may be effectively used for such imaging. As will be evident to the skilled artisan, the amount of radioisotope to be administered is dependent upon the radioisotope. Those having ordinary skill in the art can readily formulate the amount of the imaging agent to be administered based upon the specific activity and energy of a given radionuclide used as the active moiety.

Typically, 0.1-100 millicuries per dose of imaging agent, or 1-10 millicuries, or 2-5 millicuries are administered. Thus, the compositions disclosed are useful as imaging agents comprising a targeting moiety conjugated to a radioactive moiety comprise 0.1-100 millicuries, in some embodiments 1-10 millicuries, in some embodiments 2-5 millicuries, in some embodiments 1-5 millicuries.

Immunoconjugates

[0202] The application further provides immunoconjugates comprising an antibody or antigen-binding fragment thereof of the present invention conjugated (or linked) directly or indirectly to an effector molecule. In this regard, the term "conjugated" or "linked" refers to making two polypeptides into one contiguous polypeptide molecule. The linkage can be either by chemical or recombinant means. In one embodiment, the linkage is chemical, wherein a reaction between the antibody moiety and the effector molecule has produced a covalent bond formed between the two molecules to form one molecule. A peptide linker (short peptide sequence) can optionally be included between the antibody and the effector molecule. In various embodiments, an antibody or antigen-binding fragment is joined to an effector molecule. In other embodiments, an antibody or antigen-binding fragment joined to an effector molecule is further joined to a lipid, a protein or peptide to increase its half-life in the body. Accordingly in various embodiments, the antibodies of the present disclosure may be used to deliver a variety of effector molecules.

[0203] The effector molecule can be a detectable label, an immunotoxin, cytokine, chemokine, therapeutic agent, or chemotherapeutic agent.

[0204] Specific, non-limiting examples of immunotoxins include, but are not limited to, abrin, ricin, Pseudomonas exotoxin (PE, such as PE35, PE37, PE38, and PE40), diphtheria toxin (DT), botulinum toxin, cholera toxin, or modified toxins thereof, or other toxic agents that directly or indirectly inhibit cell growth or kill cells.

[0205] A "cytokine" is class of proteins or peptides released by one cell population which act on another cell as intercellular mediators. Cytokines can act as an immune-modulating agent. Examples of cytokines include lymphokines, monokines, growth factors and traditional polypeptide hormones. Thus, embodiments may utilize an interferon (e.g., IFN- α , IFN- β , and

IFN- γ); tumor necrosis factor super family (TNFSF) member; human growth hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; follicle stimulating hormone (FSH); thyroid stimulating hormone (TSH); luteinizing hormone (LH); hepatic growth factor; prostaglandin, fibroblast growth factor; prolactin; placental lactogen, OB protein; TNF- α ; TNF- β ; integrin; thrombopoietin (TPO); a nerve growth factor such as NGF- β ; platelet-growth factor; TGF- α ; TGF- β ; insulin-like growth factor-I and -II; erythropoietin (EPO); colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); an interleukin (IL-1 to IL-21), kit-ligand or FLT-3, angiostatin, thrombospondin, or endostatin. These cytokine include proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence cytokines.

[0206] Chemokines can also be conjugated to the antibodies disclosed herein. Chemokines are a superfamily of small (approximately about 4 to about 14 kDa), inducible and secreted pro-inflammatory cytokines that act primarily as chemoattractants and activators of specific leukocyte cell subtypes. Chemokine production is induced by inflammatory cytokines, growth factors and pathogenic stimuli. The chemokine proteins are divided into subfamilies (alpha, beta, and delta) based on conserved amino acid sequence motifs and are classified into four highly conserved groups--CXC, CC, C and CX3C, based on the position of the first two cysteines that are adjacent to the amino terminus. To date, more than 50 chemokines have been discovered and there are at least 18 human seven-transmembrane-domain (7TM) chemokine receptors. Chemokines of use include, but are not limited to, RANTES, MCAF, MCP-1, and fractalkine.

[0207] The therapeutic agent can be a chemotherapeutic agent. One of skill in the art can readily identify a chemotherapeutic agent of use (e.g. see Slapak and Kufe, Principles of Cancer Therapy, Chapter 86 in Harrison's Principles of Internal Medicine, 14th edition; Perry et al., Chemotherapy, Ch. 17 in Abeloff, Clinical Oncology 2^{sup}.nd ed., 2000 Churchill Livingstone, Inc; Baltzer L., Berkery R. (eds): Oncology Pocket Guide to Chemotherapy, 2nd ed. St. Louis, Mosby-Year Book, 1995; Fischer D S, Knobf M F, Durivage H J (eds): The Cancer Chemotherapy Handbook, 4th ed. St. Louis, Mosby-Year Book, 1993). Useful chemotherapeutic agents for the preparation of immunoconjugates include auristatin, dolastatin, MMAE, MMAF, AFP, DM1, AEB, doxorubicin, daunorubicin, methotrexate, melphalan, chlorambucil, vinca

alkaloids, 5-fluorouridine, mitomycin-C, taxol, L-asparaginase, mercaptopurine, thioguanine, hydroxyurea, cytarabine, cyclophosphamide, ifosfamide, nitrosoureas, cisplatin, carboplatin, mitomycin, dacarbazine, procarbazine, topotecan, nitrogen mustards, cytoxan, etoposide, BCNU, irinotecan, camptothecins, bleomycin, idarubicin, dactinomycin, plicamycin, mitoxantrone, asparaginase, vinblastine, vincristine, vinorelbine, paclitaxel, and docetaxel and salts, solvents and derivatives thereof. In various embodiments, the chemotherapeutic agent is auristatin E (also known in the art as dolastatin-10) or a derivative thereof as well as pharmaceutically salts or solvates thereof. Typical auristatin derivatives include DM1, AEB, AEVB, AFP, MMAF, and MMAE. The synthesis and structure of auristatin E and its derivatives, as well as linkers, are described in, e.g., U.S. Patent Application Publication No. 20030083263; U.S. Patent Application Publication No. 20050238629; and U.S. Patent No. 6,884,869 (each of which is incorporated by reference herein in its entirety). In various embodiments, the therapeutic agent is an auristatin or an auristatin derivative. In various embodiments, the auristatin derivative is dovaline-valine-dolaisoleunine-dolaproine-phenylalanine (MMAF) or monomethauristatin E (MMAE). In various embodiments, the therapeutic agent is a maytansinoid or a maytansinol analogue. In various embodiments, the maytansinoid is DM1.

[0208] The effector molecules can be linked to an antibody or antigen-binding fragment of the present invention using any number of means known to those of skill in the art. Both covalent and noncovalent attachment means may be used. The procedure for attaching an effector molecule to an antibody varies according to the chemical structure of the effector molecule. Polypeptides typically contain a variety of functional groups; such as carboxylic acid (COOH), free amine ($--NH_2$) or sulfhydryl ($--SH$) groups, which are available for reaction with a suitable functional group on an antibody to result in the binding of the effector molecule. Alternatively, the antibody is derivatized to expose or attach additional reactive functional groups. The derivatization may involve attachment of any of a number of linker molecules such as those available from Pierce Chemical Company, Rockford, Ill. The linker can be any molecule used to join the antibody to the effector molecule. The linker is capable of forming covalent bonds to both the antibody and to the effector molecule. Suitable linkers are well known to those of skill in the art and include, but are not limited to, straight or branched-chain carbon linkers, heterocyclic carbon linkers, or peptide linkers. Where the antibody and the

effector molecule are polypeptides, the linkers may be joined to the constituent amino acids through their side groups (such as through a disulfide linkage to cysteine) or to the alpha carbon amino and carboxyl groups of the terminal amino acids.

[0209] In some circumstances, it is desirable to free the effector molecule from the antibody when the immunoconjugate has reached its target site. Therefore, in these circumstances, immunoconjugates will comprise linkages that are cleavable in the vicinity of the target site. Cleavage of the linker to release the effector molecule from the antibody may be prompted by enzymatic activity or conditions to which the immunoconjugate is subjected either inside the target cell or in the vicinity of the target site.

[0210] Procedures for conjugating the antibodies with the effector molecules have been previously described and are within the purview of one skilled in the art. For example, procedures for preparing enzymatically active polypeptides of the immunotoxins are described in WO84/03508 and WO85/03508, which are hereby incorporated by reference for purposes of their specific teachings thereof. Other techniques are described in Shih et al., Int. J. Cancer 41:832-839 (1988); Shih et al., Int. J. Cancer 46:1101-1106 (1990); Shih et al., U.S. Pat. No. 5,057,313; Shih Cancer Res. 51:4192, International Publication WO 02/088172; U.S. Pat. No. 6,884,869; International Patent Publication WO 2005/081711; U.S. Published Application 2003-0130189 A; and US Patent Application No. 20080305044, each of which is incorporated by reference herein for the purpose of teaching such techniques.

[0211] An immunoconjugate of the present invention retains the immunoreactivity of the antibody or antigen-binding fragment, e.g., the antibody or antigen-binding fragment has approximately the same, or only slightly reduced, ability to bind the antigen after conjugation as before conjugation. As used herein, an immunoconjugate is also referred to as an antibody drug conjugate (ADC).

Bispecific Molecules

[0212] In another aspect, the present invention features bispecific molecules comprising an anti-ETR antibody, or antigen-binding fragment thereof, of the invention. An antibody of the invention, or antigen-binding fragment thereof, can be derivatized or linked to another functional molecule, e.g., another peptide or protein (e.g., another antibody or ligand for a receptor) to

generate a bispecific molecule that binds to at least two different binding sites or target molecules. The antibody of the invention may in fact be derivatized or linked to more than one other functional molecule to generate multispecific molecules that bind to more than two different binding sites and/or target molecules; such multispecific molecules are also intended to be encompassed by the term "bispecific molecule" as used herein. To create a bispecific molecule of the invention, an antibody of the invention can be functionally linked (e.g., by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other binding molecules, such as another antibody, antibody fragment, peptide or binding mimetic, such that a bispecific molecule results. In various embodiments, the invention includes bispecific molecules capable of binding both to FcγR or FcαR expressing effector cells (e.g., monocytes, macrophages or polymorphonuclear cells (PMNs)), and to target cells expressing PD. In such embodiments, the bispecific molecules target ETR expressing cells to effector cell and trigger Fc receptor-mediated effector cell activities, e.g., phagocytosis of an ETR expressing cells, antibody dependent cell-mediated cytotoxicity (ADCC), cytokine release, or generation of superoxide anion. Methods of preparing the bispecific molecules of the present invention are well known in the art.

Polynucleotides and Antibody Expression

[0213] The application further provides polynucleotides comprising a nucleotide sequence encoding an anti-ETR antibody or antigen-binding fragment thereof. Because of the degeneracy of the genetic code, a variety of nucleic acid sequences encode each antibody amino acid sequence. The application further provides polynucleotides that hybridize under stringent or lower stringency hybridization conditions, e.g., as defined herein, to polynucleotides that encode an antibody that binds to human ETR.

[0214] Stringent hybridization conditions include, but are not limited to, hybridization to filter-bound DNA in 6xSSC at about 45°C followed by one or more washes in 0.2xSSC/0.1% SDS at about 50-65°C, highly stringent conditions such as hybridization to filter-bound DNA in 6xSSC at about 45°C followed by one or more washes in 0.1xSSC/0.2% SDS at about 60°C, or any other stringent hybridization conditions known to those skilled in the art (see, for example,

Ausubel, F. M. et al., eds. 1989 Current Protocols in Molecular Biology, vol. 1, Green Publishing Associates, Inc. and John Wiley and Sons, Inc., NY at pages 6.3.1 to 6.3.6 and 2.10.3).

[0215] The polynucleotides may be obtained, and the nucleotide sequence of the polynucleotides determined, by any method known in the art. For example, if the nucleotide sequence of the antibody is known, a polynucleotide encoding the antibody may be assembled from chemically synthesized oligonucleotides (e.g., as described in Kutmeier et al., *BioTechniques* 17:242 (1994)), which, briefly, involves the synthesis of overlapping oligonucleotides containing portions of the sequence encoding the antibody, annealing and ligating of those oligonucleotides, and then amplification of the ligated oligonucleotides by PCR. In one embodiment, the codons that are used comprise those that are typical for human or mouse (see, e.g., Nakamura, Y., *Nucleic Acids Res.* 28: 292 (2000)).

[0216] A polynucleotide encoding an antibody may also be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid encoding a particular antibody is not available, but the sequence of the antibody molecule is known, a nucleic acid encoding the immunoglobulin may be chemically synthesized or obtained from a suitable source (e.g., an antibody cDNA library, or a cDNA library generated from, or nucleic acid, preferably polyA+RNA, isolated from, any tissue or cells expressing the antibody, such as hybridoma cells selected to express an antibody) by PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the sequence or by cloning using an oligonucleotide probe specific for the particular gene sequence to identify, e.g., a cDNA clone from a cDNA library that encodes the antibody. Amplified nucleic acids generated by PCR may then be cloned into replicable cloning vectors using any method well known in the art.

[0217] The present invention is also directed to host cells that express a ETR and/or the anti-ETR antibodies of the invention. A wide variety of host expression systems known in the art can be used to express an antibody of the present invention including prokaryotic (bacterial) and eukaryotic expression systems (such as yeast, baculovirus, plant, mammalian and other animal cells, transgenic animals, and hybridoma cells), as well as phage display expression systems.

[0218] An antibody of the invention can be prepared by recombinant expression of immunoglobulin light and heavy chain genes in a host cell. To express an antibody

recombinantly, a host cell is transformed, transduced, infected or the like with one or more recombinant expression vectors carrying DNA fragments encoding the immunoglobulin light and/or heavy chains of the antibody such that the light and/or heavy chains are expressed in the host cell. The heavy chain and the light chain may be expressed independently from different promoters to which they are operably-linked in one vector or, alternatively, the heavy chain and the light chain may be expressed independently from different promoters to which they are operably-linked in two vectors one expressing the heavy chain and one expressing the light chain. Optionally, the heavy chain and light chain may be expressed in different host cells.

[0219] Additionally, the recombinant expression vector can encode a signal peptide that facilitates secretion of the antibody light and/or heavy chain from a host cell. The antibody light and/or heavy chain gene can be cloned into the vector such that the signal peptide is operably-linked in-frame to the amino terminus of the antibody chain gene. The signal peptide can be an immunoglobulin signal peptide or a heterologous signal peptide. Preferably, the recombinant antibodies are secreted into the medium in which the host cells are cultured, from which the antibodies can be recovered or purified.

[0220] An isolated DNA encoding a HCVR can be converted to a full-length heavy chain gene by operably linking the HCVR-encoding DNA to another DNA molecule encoding heavy chain constant regions. The sequences of human, as well as other mammalian, heavy chain constant region genes are known in the art. DNA fragments encompassing these regions can be obtained e.g., by standard PCR amplification. The heavy chain constant region can be of any type, (e.g., IgG, IgA, IgE, IgM or IgD), class (e.g., IgG₁, IgG₂, IgG₃ and IgG₄) or subclass constant region and any allotypic variant thereof as described in Kabat (supra).

[0221] An isolated DNA encoding a LCVR region may be converted to a full-length light chain gene (as well as to a Fab light chain gene) by operably linking the LCVR-encoding DNA to another DNA molecule encoding a light chain constant region. The sequences of human, as well as other mammalian, light chain constant region genes are known in the art. DNA fragments encompassing these regions can be obtained by standard PCR amplification. The light chain constant region can be a kappa or lambda constant region.

[0222] In addition to the antibody heavy and/or light chain gene(s), a recombinant expression vector of the invention carries regulatory sequences that control the expression of

the antibody chain gene(s) in a host cell. The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals), as needed, that control the transcription or translation of the antibody chain gene(s). The design of the expression vector, including the selection of regulatory sequences may depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired. Preferred regulatory sequences for mammalian host cell expression include viral elements that direct high levels of protein expression in mammalian cells, such as promoters and/or enhancers derived from cytomegalovirus (CMV), Simian Virus 40 (SV40), adenovirus, (e.g., the adenovirus major late promoter (AdMLP)) and/or polyoma virus.

[0223] Additionally, the recombinant expression vectors of the invention may carry additional sequences, such as sequences that regulate replication of the vector in host cells (e.g., origins of replication) and one or more selectable marker genes. The selectable marker gene facilitates selection of host cells into which the vector has been introduced. For example, typically the selectable marker gene confers resistance to drugs, such as G418, hygromycin, or methotrexate, on a host cell into which the vector has been introduced. Preferred selectable marker genes include the dihydrofolate reductase (dhfr) gene (for use in dhfr-minus host cells with methotrexate selection/amplification), the neo gene (for G418 selection), and glutamine synthetase (GS) in a GS-negative cell line (such as NSO) for selection/amplification.

[0224] For expression of the light and/or heavy chains, the expression vector(s) encoding the heavy and/or light chains is introduced into a host cell by standard techniques e.g. electroporation, calcium phosphate precipitation, DEAE-dextran transfection, transduction, infection and the like. Although it is theoretically possible to express the antibodies of the invention in either prokaryotic or eukaryotic host cells, eukaryotic cells are preferred, and most preferably mammalian host cells, because such cells are more likely to assemble and secrete a properly folded and immunologically active antibody. Preferred mammalian host cells for expressing the recombinant antibodies of the invention include Chinese Hamster Ovary (CHO cells) [including dhfr minus CHO cells, as described in Urlaub and Chasin, Proc. Natl. Acad. Sci. USA 77:4216-20, 1980, used with a DHFR selectable marker, e.g. as described in Kaufman and Sharp, J. Mol. Biol. 159:601-21, 1982], NSO myeloma cells, COS cells, and SP2/0 cells. When recombinant expression vectors encoding antibody genes are introduced into mammalian host

cells, the antibodies are produced by culturing the host cells for a period of time sufficient to allow for expression of the antibody in the host cells or, more preferably, secretion of the antibody into the culture medium in which the host cells are grown under appropriate conditions known in the art. Antibodies can be recovered from the host cell and/or the culture medium using standard purification methods.

[0225] The invention provides a host cell comprising a nucleic acid molecule of the present invention. Preferably a host cell of the invention comprises one or more vectors or constructs comprising a nucleic acid molecule of the present invention. For example, a host cell of the invention is a cell into which a vector of the invention has been introduced, said vector comprising a polynucleotide encoding a LCVR of an antibody of the invention and/or a polynucleotide encoding a HCVR of the invention. The invention also provides a host cell into which two vectors of the invention have been introduced; one comprising a polynucleotide encoding a LCVR of an antibody of the invention and one comprising a polynucleotide encoding a HCVR present in an antibody of the invention and each operably-linked to enhancer/promoter regulatory elements (e.g., derived from SV40, CMV, adenovirus and the like, such as a CMV enhancer/AdMLP promoter regulatory element or an SV40 enhancer/AdMLP promoter regulatory element) to drive high levels of transcription of the genes.

[0226] Once expressed, the intact antibodies, individual light and heavy chains, or other immunoglobulin forms of the present invention can be purified according to standard procedures of the art, including ammonium sulfate precipitation, ion exchange, affinity (e.g., Protein A), reverse phase, hydrophobic interaction column chromatography, hydroxyapatite chromatography, gel electrophoresis, and the like. Standard procedures for purification of therapeutic antibodies are described, for example, by Feng L1, Joe X. Zhou, Xiaoming Yang, Tim Tressel, and Brian Lee in an article entitled "Current Therapeutic Antibody Production and Process Optimization" (BioProcessing Journal, September/October 2005)(incorporated by reference in its entirety for purposes of teaching purification of therapeutic antibodies). Additionally, standard techniques for removing viruses from recombinantly expressed antibody preparations are also known in the art (see, for example, Gerd Kern and Mani Krishnan, "Viral Removal by Filtration: Points to Consider" (Biopharm International, October 2006)). The effectiveness of filtration to remove viruses from preparations of therapeutic antibodies is known

to be at least in part dependent on the concentration of protein and/or the antibody in the solution to be filtered. The purification process for antibodies of the present invention may include a step of filtering to remove viruses from the mainstream of one or more chromatography operations. Preferably, prior to filtering through a pharmaceutical grade nanofilter to remove viruses, a chromatography mainstream containing an antibody of the present invention is diluted or concentrated to give total protein and/or total antibody concentration of about 1 g/L to about 3 g/L. Even more preferably, the nanofilter is a DV20 nanofilter (e.g., Pall Corporation; East Hills, N.Y.). Substantially pure immunoglobulins of at least about 90%, about 92%, about 94% or about 96% homogeneity are preferred, and about 98 to about 99% or more homogeneity most preferred, for pharmaceutical uses. Once purified, partially or to homogeneity as desired, the sterile antibodies may then be used therapeutically, as directed herein.

[0227] In view of the aforementioned discussion, the present invention is further directed to an antibody obtainable by a process comprising the steps of culturing a host cell including, but not limited to a mammalian, plant, bacterial, transgenic animal, or transgenic plant cell which has been transformed by a polynucleotide or a vector comprising nucleic acid molecules encoding antibodies of the invention so that the nucleic acid is expressed and, optionally, recovering the antibody from the host cell culture medium.

[0228] In certain aspects, the present application provides hybridoma cell lines, as well as to the monoclonal antibodies produced by these hybridoma cell lines. The cell lines disclosed have uses other than for the production of the monoclonal antibodies. For example, the cell lines can be fused with other cells (such as suitably drug-marked human myeloma, mouse myeloma, human-mouse heteromyeloma or human lymphoblastoid cells) to produce additional hybridomas, and thus provide for the transfer of the genes encoding the monoclonal antibodies. In addition, the cell lines can be used as a source of nucleic acids encoding the anti-ETR immunoglobulin chains, which can be isolated and expressed (e.g., upon transfer to other cells using any suitable technique (see e.g., Cabilly et al., U.S. Pat. No. 4,816,567; Winter, U.S. Pat. No. 5,225,539)). For instance, clones comprising a rearranged anti-ETR light or heavy chain can be isolated (e.g., by PCR) or cDNA libraries can be prepared from mRNA isolated from the cell lines, and cDNA clones encoding an anti-ETR immunoglobulin chain can be isolated. Thus,

nucleic acids encoding the heavy and/or light chains of the antibodies or portions thereof can be obtained and used in accordance with recombinant DNA techniques for the production of the specific immunoglobulin, immunoglobulin chain, or variants thereof (e.g., humanized immunoglobulins) in a variety of host T-cells or in an in vitro translation system. For example, the nucleic acids, including cDNAs, or derivatives thereof encoding variants such as a humanized immunoglobulin or immunoglobulin chain, can be placed into suitable prokaryotic or eukaryotic vectors (e.g., expression vectors) and introduced into a suitable host T-cell by an appropriate method (e.g., transformation, transfection, electroporation, infection), such that the nucleic acid is operably linked to one or more expression control elements (e.g., in the vector or integrated into the host T-cell genome). For production, host T-cells can be maintained under conditions suitable for expression (e.g., in the presence of inducer, suitable media supplemented with appropriate salts, growth factors, antibiotic, nutritional supplements, etc.), whereby the encoded polypeptide is produced. If desired, the encoded protein can be recovered and/or isolated (e.g., from the host T-cells or medium). It will be appreciated that the method of production encompasses expression in a host T-cell of a transgenic animal (see e.g., WO 92/03918, GenPharm International, published Mar. 19, 1992)(incorporated by reference in its entirety).

[0229] Host cells can also be used to produce portions, or fragments, of intact antibodies, e.g., Fab fragments or scFv molecules by techniques that are conventional. For example, it may be desirable to transfect a host cell with DNA encoding either the light chain or the heavy chain of an antibody of this invention. Recombinant DNA technology may also be used to remove some or all the DNA encoding either or both of the light and heavy chains that is not necessary for binding to human ETR. The molecules expressed from such truncated DNA molecules are also encompassed by the antibodies of the invention.

[0230] Methods for expression of single chain antibodies and/or refolding to an appropriate active form, including single chain antibodies, from bacteria such as *E. coli* have been described and are well-known and are applicable to the antibodies disclosed herein (see, e.g., Buchner et al., Anal. Biochem. 205:263-270, 1992; Pluckthun, Biotechnology 9:545, 1991; Huse et al., Science 246:1275, 1989 and Ward et al., Nature 341:544, 1989, all incorporated by reference herein).

[0231] Often, functional heterologous proteins from *E. coli* or other bacteria are isolated from inclusion bodies and require solubilization using strong denaturants, and subsequent refolding. During the solubilization step, as is well known in the art, a reducing agent must be present to separate disulfide bonds. An exemplary buffer with a reducing agent is: 0.1 M Tris pH 8, 6 M guanidine, 2 mM EDTA, 0.3 M DTE (dithioerythritol). Reoxidation of the disulfide bonds can occur in the presence of low molecular weight thiol reagents in reduced and oxidized form, as described in Saxena et al., Biochemistry 9: 5015-5021, 1970, incorporated by reference herein, and especially as described by Buchner et al., supra.

[0232] Renaturation is typically accomplished by dilution (for example, 100-fold) of the denatured and reduced protein into refolding buffer. An exemplary buffer is 0.1 M Tris, pH 8.0, 0.5 M L-arginine, 8 mM oxidized glutathione (GSSG), and 2 mM EDTA.

[0233] As a modification to the two-chain antibody purification protocol, the heavy and light chain regions are separately solubilized and reduced and then combined in the refolding solution. An exemplary yield is obtained when these two proteins are mixed in a molar ratio such that a 5-fold molar excess of one protein over the other is not exceeded. Excess oxidized glutathione or other oxidizing low molecular weight compounds can be added to the refolding solution after redox shuffling is completed.

[0234] In addition to recombinant methods, the antibodies, labeled antibodies and antigen-binding fragments thereof that are disclosed herein can also be constructed in whole or in part using standard peptide synthesis. Solid phase synthesis of the polypeptides of less than about 50 amino acids in length can be accomplished by attaching the C-terminal amino acid of the sequence to an insoluble support followed by sequential addition of the remaining amino acids in the sequence. Techniques for solid phase synthesis are described by Barany & Merrifield, The Peptides: Analysis, Synthesis, Biology. Vol. 2: Special Methods in Peptide Synthesis, Part A. pp. 3-284; Merrifield et al., J. Am. Chem. Soc. 85:2149-2156, 1963, and Stewart et al., Solid Phase Peptide Synthesis, 2nd ed., Pierce Chem. Co., Rockford, Ill., 1984. Proteins of greater length may be synthesized by condensation of the amino and carboxyl termini of shorter fragments. Methods of forming peptide bonds by activation of a carboxyl

terminal end (such as by the use of the coupling reagent N,N'-dicyclohexylcarbodiimide) are well known in the art.

[0235] The following examples are offered to more fully illustrate the invention but are not construed as limiting the scope thereof.

Example 1

Generation of Monoclonal Antibodies Targeting Specifically to Human ET_AR

[0236] Balb/c and C57Bl/6 mouse (22 gms, 6-8 weeks old) were each immunized bi-weekly (5 total immunizations) with 100 µg or 50 µg hET_AR plasmid DNA with Gene Gun, and splenocytes were harvested 4 days after the final immunization for fusion with myeloma cell line NS0 from ATCC (Allendale, NJ). Electric fusion methods are used to obtain hybridoma cells and then the hybridoma supernatants are screened for antigen binding, ligand blocking, IgG binning, reference antibody binding, and FACS binding. 21 murine mAbs were ultimately selected from the initial screens for subcloning (limited dilution method). BD Cell MAb Medium was used to grow hybridomas in roller bottles for the collection of supernatants for antibody production. mAbs were purified with Protein A affinity chromatography. Estimated purity of mAbs was higher than 90% based on SDS-PAGE Coomassie staining. Secondary screening of the 21 mAbs comprised: human ET_AR binding assays (FACS) and evaluation of functional effects in an IP1 assay (IP-one-Gq kit, Cisbio).

[0237] To evaluate the functional effects of ET_AR, the 21 purified mAbs were assessed in recombinant CHO cells stably expressing ET_AR using an IP1 assay (IP-one-Gq kit, Cisbio). To assess antagonist activity, recombinant CHO cells expressing tagged ET_AR were added to 96-well half-area white plates (20,000 cells per well) and were treated with EDNRA mAb at 0.5 pM to 1 µM for 30 min at room temperature. 25.0 nM of ET-1 corresponding to the EC80, was then added and cells were further incubated for 30 min at room temperature. The reaction was stopped by adding detection mix to all wells followed by a 60 min incubation at room temperature. The assay plates were read on an EnVision® Multilabel Plate Reader (PerkinElmer; Waltham, MA, USA) at an emission wavelength of 665 nm. The ratio of HTRF

665/620 is used to indicate the blocking of ligand binding. The lower the ratio, the lower degree of the blockage. All results were analyzed by nonlinear regression curve fit using GraphPad Prism version 6 (GraphPad Inc., La Jolla, CA).

[0238] The functional assay data for 21 mAbs are summarized in FIG. 1. The IC₅₀ of the top two Abs are summarized in FIG. 2 and Table 3:

Table 3

Murine mAb	mAb ID	HuET _A R Binding IC ₅₀ (nM)
A2	336D8G9	0.06072
A3	312A10E5	0.08906

[0239] Based on the cumulative results of the binding and secondary assays, purified murine mAb 178D1F6C1G6 ("A1"), mAb 336D8G9 ("A2") and mAb 312A10E5 ("A3") were selected for sequencing and further analysis. Total RNA was extracted from frozen hybridoma cells following the technical manual of TRIzol® Reagent. The total RNA was analyzed by agarose gel electrophoresis. Total RNA was reverse transcribed into cDNA using isotype-specific anti-sense primers or universal primers following the technical manual of PrimeScript™ 1st Strand cDNA Synthesis Kit. PCR was then performed to amplify the variable regions (heavy and light chains) of the antibodies, which were then cloned into a standard cloning vector separately and sequenced. Murine mAb 178D1F6C1G6 ("A1"), mAb 336D8G9 ("A2") and mAb 312A10E5 ("A3") comprise the heavy chain variable region sequences set forth in SEQ ID NOs: 19, 23 and 27, respectively, and the light chain variable region sequences set forth in SEQ ID NOs: 21, 25 and 29, respectively. The heavy chain variable regions of mAb 178D1F6C1G6 ("A1"), mAb 336D8G9 ("A2") and mAb 312A10E5 ("A3") are encoded by the nucleic acid sequences set forth in SEQ ID NOs: 20, 24 and 28, respectively, and the light chain variable regions are encoded by the nucleic acid sequences set forth in SEQ ID NOs: 22, 26 and 30, respectively.

Example 2

Generation of a Chimeric IgG Targeting Human ET_AR

[0240] Using the HCVR and LCVR sequences of mAb 312A10E5 ("A3"), a murine-human IgG chimeric Fab (hereinafter "chimeric IgG") was prepared and which comprised the heavy chain variable region sequence set forth in SEQ ID NO: 31 and the light chain variable region sequence set forth in SEQ ID NO: 32.

[0241] The DNA sequences encoding the chimeric IgG heavy and light chains were synthesized and inserted into pTT5 vector to construct the expression plasmids of full-length IgGs. Expression of chimeric IgG was conducted in HEK293 cell culture and the supernatants were purified with protein A affinity column. The purified antibody was buffer-exchanged into PBS using PD-10 desalting column. The concentration and purity of the purified protein were determined by OD₂₈₀ and SDS-PAGE, respectively. The purified chimeric IgG migrated as ~170 kDa band in SDS-PAGE under non-reducing conditions. Evaluating by the SDS-PAGE result, the purity of the IgGs was > 95%.

Example 3

Generation of Humanized Abs Specifically Targeting Human ETR

[0242] A CDR grafting and back mutation method was used to prepare humanized anti-ETR mAbs derived from murine mAb 312A10E5 ("A3"). The structure of parental antibody was modelled by computer-aided homology modelling program. Humanized antibodies were designed using CDR grafting and subsequent replaced putative back mutation sites of grafted antibody. Briefly, the CDRs of parental mAb A3 were grafted into the human acceptors to obtain humanized light chains and humanized heavy chains for the parental antibody. The sequences of three humanized heavy chain variable regions (referred to as VH1, VH2, VH3) are set forth in SEQ ID NOs: 33-35, respectively and the sequences of three humanized light chain variable

regions (referred to as VL2 and VL3) are set forth in SEQ ID NOs: 36-37, respectively were paired with each other for affinity ranking experiments.

[0243] The DNA sequences encoding the chimeric antibody and humanized IgG heavy and light chains were synthesized and inserted into pCDNA3.4 vector to construct the expression plasmids of full-length IgGs. 16 humanized Abs were expressed in Expi293F cell culture, and then the cells were spun down. For affinity ranking, the supernatants were assayed by FACS against cell lines for the assessment of binding activity. Positive cells lines (CCR5/CHO-K1) and negative cells were cultured following the protocol and harvested by centrifugation. About 1×10^5 cells per well were washed with 1*PBS and incubated in a serial dilution of antibodies for 30 minutes at 4°C. After washing with PBS, secondary antibody [1 µg/ml Alexa Fluor® 647 AffiniPure Goat Anti-Human IgG (H+L)] was added to the cells and incubated in PBS for 30 minutes at 4°C. After washing with PBS, cells were analyzed for binding by using FACS Calibur (BD Bioscience, San Jose, CA) and Flowjo software. The antibodies were ranked by EC50. The binders that interact with ET_AR with similar affinity to parental antibody were selected.

[0244] Top three binders were selected to express in the Expi293F cell culture. The recombinant plasmids encoding target IgG were transiently co-transfected and the supernatants were purified with protein A affinity column. The purified antibody was buffer exchanged into PBS using dialysis bag. The concentration and purity of the purified protein were determined by OD280 and SDS-PAGE, respectively. Binding confirmation were tested by FACS Calibur (BD Bioscience, San Jose, CA).

[0245] Three IgGs (VH1/VL3), (VH2/VL3) and (VH3/VL3) were selected for expression in the Expi293F cell culture. The recombinant plasmids encoding target IgG were transiently co-transfected and the supernatants were purified with protein A affinity column. The purified antibody was buffer exchanged into PBS using dialysis bag. The concentration and purity of the purified protein were determined by OD280 and SDS-PAGE, respectively. Evaluating from the SDS-PAGE, the purity of humanized IgGs were >95%. Binding confirmation were tested by FACS Calibur (BD Bioscience, San Jose, CA). The HC and LC amino acid sequences for the 3 selected humanized ETR antibodies are summarized in Table 4:

Table 4

Humanized IgG	HC	LC
1 (VH1/VL3)	SEQ ID NO: 38	SEQ ID NO: 40
2 (VH2/VL3)	SEQ ID NO: 42	SEQ ID NO: 40
3 (VH3/VL3)	SEQ ID NO: 44	SEQ ID NO: 40

[0246] The FACS binding data is summarized in FIG. 3. All three humanized antibodies demonstrated similar binding affinity as the chimeric antibody.

[0247] All of the articles and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the articles and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the articles and methods without departing from the spirit and scope of the invention. All such variations and equivalents apparent to those skilled in the art, whether now existing or later developed, are deemed to be within the spirit and scope of the invention as defined by the appended claims. All patents, patent applications, and publications mentioned in the specification are indicative of the levels of those of ordinary skill in the art to which the invention pertains. All patents, patent applications, and publications are herein incorporated by reference in their entirety for all purposes and to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference in its entirety for any and all purposes. The invention illustratively described herein suitably may be practiced in the absence of any element(s) not specifically disclosed herein. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the appended claims.

Sequence Listings

The nucleic and amino acid sequences listed in the accompanying sequence listing are shown using standard letter abbreviations for nucleotide bases and three letter code for amino acids, as defined in 37 C.F.R. 1.822.

SEQ ID NO: 1 is an amino acid sequence comprising a human ET_AR.

SEQ ID NOs: 2-4 are the amino acid sequence of a heavy chain CDR1 in a monoclonal antibody which specifically binds ET_AR.

SEQ ID NOs: 5-7 are the amino acid sequences of a heavy chain CDR2 in a monoclonal antibody which specifically binds ET_AR.

SEQ ID NOs: 8-10 are the amino acid sequences of a heavy chain CDR3 in a monoclonal antibody which specifically binds ET_AR.

SEQ ID NOs: 11-13 are the amino acid sequences of a light chain CDR1 in a monoclonal antibody which specifically binds ET_AR.

SEQ ID NO: 14-15 are the amino acid sequences of a light chain CDR2 in a monoclonal antibody which specifically binds ET_AR.

SEQ ID NOs: 16-18 are the amino acid sequences of a light chain CDR3 in a monoclonal antibody which specifically binds ET_AR.

SEQ ID NOs: 19, 23 and 27 are amino acid sequences of a heavy chain variable region of murine monoclonal antibodies which specifically bind ET_AR.

SEQ ID NOs: 20, 24 and 28 are nucleic acid sequences encoding a heavy chain variable region of murine monoclonal antibodies which specifically bind ET_AR.

SEQ ID NOs: 21, 25 and 29 are amino acid sequences of a light chain variable region of murine monoclonal antibodies which specifically bind ET_AR.

SEQ ID NOs: 22, 26 and 30 are nucleic acid sequences encoding a light chain variable region of murine monoclonal antibodies which specifically bind ET_AR.

SEQ ID NO: 31 is the amino acid sequence of a heavy chain variable region of a murine-human chimeric antibody which specifically binds ET_AR.

SEQ ID NO: 32 is the amino acid sequence of a light chain variable region of a murine-human chimeric antibody which specifically binds ET_AR.

SEQ ID NOs: 33-35 are amino acid sequences of a heavy chain variable regions of humanized monoclonal antibodies which specifically binds ET_AR.

SEQ ID NOs: 36-37 are amino acid sequences of a light chain variable regions of humanized monoclonal antibodies which specifically binds ET_AR.

SEQ ID NO: 38, 42 and 44 are amino acid sequences of a heavy chain of humanized monoclonal antibodies which specifically binds ET_AR.

SEQ ID NOs: 39, 43 and 45 are nucleic acid sequences of a heavy chain of humanized monoclonal antibodies which specifically binds ET_AR.

SEQ ID NOs: 40 and 46 are amino acid sequences of a light chain of humanized monoclonal antibodies which specifically binds ET_AR.

SEQ ID NOs: 41 and 47 are nucleic acid sequences of a light chain of humanized monoclonal antibodies which specifically binds ET_AR.

SEQ ID NOs: 48 and 49 are the amino acid sequences of a light chain constant region of a monoclonal antibody which specifically binds ET_AR.

SEQ ID NO: 50 is the amino acid sequence of a heavy chain constant region of a monoclonal antibody which specifically binds ET_AR.

SEQUENCE LISTINGS

SEQ ID NO: 1 - ET_AR amino acid sequence

METLCLRASFWLALVGCVISDNPERYSTNLSNHVDDFTTFRGTELSFLVTTHQPTNLVLPSNGS
MHNYCPQQTKITSAFKYINTVISCTIFIVGMVGNATLLRIIYQNKCMRNGPNALIASLALGDLIYVVI
DLPINVFKLLAGRWPFHDNDFGVFLCKLFPFLQKSSVGITVLNLCALSVDYRAVASWSRVQGI
GIPLVTAIEIVSIWILSFILAIPEAIGFVMVPFEYRGEQHKTCMLNATSKFMEFYQDVKDWWLFGF
YFCMPLVCTAIFYTLMTCEMLNRRNGSLRIALSEHLKQRREVAKTVFCLVVIFALCWFPPLHLSRI
LKKT VYNEMDKNRCELLSFLLLMDYIGINLATMNSCINPIALYFVSKKFKNCFQSCLCCLCCCYQSK
SLMTSVPMNGTSIQWKNHDQNNHNTDRSSHKDSMN

SEQ ID NO: 2 – Murine monoclonal antibody heavy chain CDR1 amino acid sequence

SYWMN

SEQ ID NO: 3 – Murine monoclonal antibody heavy chain CDR1 amino acid sequence

DYYIH

SEQ ID NO: 4 – Murine monoclonal antibody heavy chain CDR1 amino acid sequence

NYGMS

SEQ ID NO: 5 - Murine monoclonal antibody heavy chain CDR2 amino acid sequence

QIYPGNGDTNYNGKFRG

SEQ ID NO: 6 - Murine monoclonal antibody heavy chain CDR2 amino acid sequence

RIDPEDGDTEYAPKFQG

SEQ ID NO: 7 - Murine monoclonal antibody heavy chain CDR2 amino acid sequence

WINTYSGESTYADDFKG

SEQ ID NO: 8 - Murine monoclonal antibody heavy chain CDR3 amino acid sequence

RGSLDY

SEQ ID NO: 9 - Murine monoclonal antibody heavy chain CDR3 amino acid sequence

GYTFTNWYSNSY

SEQ ID NO: 10 - Murine monoclonal antibody heavy chain CDR3 amino acid sequence

SLNRDTWFTY

SEQ ID NO: 11 - Murine monoclonal antibody light chain CDR1 amino acid sequence

KSSHSLVHSNGDTYLH

SEQ ID NO: 12 - Murine monoclonal antibody light chain CDR1 amino acid sequence

RSSQSLVHSNGNTYLH

SEQ ID NO: 13 - Murine monoclonal antibody light chain CDR1 amino acid sequence

RSITGAVTTSNYAN

SEQ ID NO: 14 - Murine monoclonal antibody light chain CDR2 amino acid sequence

KVSNRFS

SEQ ID NO: 15 - Murine monoclonal antibody light chain CDR2 amino acid sequence

GTNNRAP

SEQ ID NO: 16 - Murine monoclonal antibody light chain CDR3 amino acid sequence

SQTTYVPYT

SEQ ID NO: 17 - Murine monoclonal antibody light chain CDR3 amino acid sequence

SQSTHVPYT

SEQ ID NO: 18 - Murine monoclonal antibody light chain CDR3 amino acid sequence

ALWYSNHVV

SEQ ID NO: 19 – Murine monoclonal antibody heavy chain variable region amino acid sequence

QVQLQQSGAELVRPGSSVRISCKAFDYAFSSYWMNWVKQRPGQDLEWIGQIYPGNGDTNYN
GKFRGKASLTADKSSSTAYMHLSSLTSEDSAVYFCARRGSLDYWGQGTSTVSS

SEQ ID NO: 20 – Murine monoclonal antibody heavy chain variable region nucleic acid sequence

cagggtcagctgcagcagctctggggctgagttggtgaggcctgggtcctcagtgaggatttcctgcaaggctttgactatgcattcagta
gctactggatgaactgggtgaagcagaggcctggacaggatcttgagtggtgacagattatcctggaaatgggtgataacta
caatggaaagttcaggggtaaagcctcactgactgcagacaaatcctccagcacagcctacatgcacctcagcagcctaactctg
aggactctgcggtctattctgtgcaagacgggggtcttggactactgggtcaaggaacctcagtcaccgtctcctca

SEQ ID NO: 21 – Murine monoclonal antibody light chain variable region amino acid sequence

DVVMQTQIPLSLPVSLGDQSSISCKSSSHSLVHSNGDTYLHWFLQKPGQSPKLLIYKVSNRFSGVP
DRFTGSGSGTEFTLKISRVEAEDLGVYFCSQTTYVPYTFGGGKLEIK

SEQ ID NO: 22 – Murine monoclonal antibody light chain variable region nucleic acid sequence

gatgtggtgatgacccaaattccactctccctgcctgtcagttctggagatcaatcctccatctctgcaaactctagtcatagcctgtacat
agtaatggagacacattattacattggttcctgcagaagccaggccagtcctcaaagctcctgatctacaaagtcctcaaccgatttctg
gggtcccagacaggttcactggcagtggttcagggacagagttcacactcaagatcagcagagtgaggctgaggatctgggagttt
atttctgctctcaaactacatatgtccgtacacgttcggaggggggaccaagctggaaataaaa

SEQ ID NO: 23 – Murine monoclonal antibody heavy chain variable region amino acid sequence

EVQLQQSGAELVRPGASVKLSCPASGFTIKDYYIHVVVKQRPEQGLEWIGRIDPEDGDTEYAPK
FQGKATMTADTSSNTAYLHLSSLTSEDTAVYYCTSNWYSNSYWGGGTTTLTVSS

SEQ ID NO: 24 – Murine monoclonal antibody heavy chain variable region nucleic acid sequence

gagggtcagctgcagcagctctggggcagagcttgtagggccaggggcctcagtcagttgtcctgccagcttctggcttcaccattaa
agactactatatacactgggtgaagcagaggcctgagcagggcctggagtggattggaaggattgatcctgaggatgggtgatactga
atatgccccgaagtccagggaaggccactatgactgcagacacatcctccaacacagcctacctgcacctcagcagcctgacat
ctgaggacactgccgtctattactgtacttcaaactggtatagtaactcctactggggccaaggcaccactctcacagtctcctca

SEQ ID NO: 25 – Murine monoclonal antibody light chain variable region amino acid sequence

DVVMQTPLSLPVS LGDQASISCRSSQSLVHSNGNTYLHWYLQKPGQSPKVL IYKVS NRFSGV
PDRFSGSGSGTDFTLKISRVEAEDLGVYFCSQSTHVPYTFGGGTKLEIK

SEQ ID NO: 26 – Murine monoclonal antibody light chain variable region nucleic acid sequence

gatgttgatgacccaaactccactctccctgcctgtcagttctggagatcaagcctccatctcttcgagatctagtcagagcctgtaca
cagtaatggaaacacctatttacattgggtacctgcaaaagccaggccagctctcaaaggctctgatctacaaagttccaaccgatttct
tggggctccagacaggttcagtgccagtggtatcagggacagatttcacactcaagatcagcagagtgaggctgaggatctgggag
ttatttctgctctcaaagtacacatgttcgtatcgttcggaggggggaccaagctggaaataaaa

SEQ ID NO: 27 – Murine monoclonal antibody heavy chain variable region amino acid sequence

QIQLVQSGPDLK KPGETVKISCKASGYTFTNYGMSWVKQAPGKGLKWMGWINTYSGESTYAD
DFKGRFAFSLETSASTAYLQINNLKNEDTATYFCARSLNRDWTWFTYWGQGLTVTVSA

SEQ ID NO: 28 – Murine monoclonal antibody heavy chain variable region nucleic acid sequence

cagatccagttggtacagtctggacctgacctgaagaagcctggagagacagtcagatctcctgcaaggcttctgggtataccttca
caaactatggaatgagctgggtgaaacaggctccaggaaagggttaaagtggatgggctggataaacacctattctggagagtca
acatatgctgatgacttcaaggacgggttgccttctcttggaaacctctgccagcactgcctatttgcagatcaacaacctcaaaaatg
aggacacggctacatatttctgtgcaagatccctcaaccgggacacctggttactactggggccaagggactctggtcactgtctctg
ca

SEQ ID NO: 29 – Murine monoclonal antibody light chain variable region amino acid sequence

QAVVTQESALSTSPGETVTLTCSRITGAVTTSNYANWVQE KPDHLFTGLIGGTNNRAPGV PAR
FSGSLIGDKAALTITGAQTEDEAIYFCALWYSNHWVFGGGTKLTVL

SEQ ID NO: 30 – Murine monoclonal antibody light chain variable region nucleic acid sequence

caggctgttgtagctcaggaatctgcactctccacatcacctgggtgaaacagtcacactcactgtcgctcaattactggggctgttaca
ctagtaactatgccaactgggtccaagaaaaaccagatcatttgttactggtctaataaggtgtaccaacaaccgagctccaggtgtt

cctgccagattctcaggctccctgattggagacaaggctgccctcaccatcacaggggcacagactgaggatgaggcaatatattct
gtgctctatggtacagcaaccactgggtgttcggtggaggaaccaaactgactgtccta

SEQ ID NO: 31 – Heavy chain variable region amino acid sequence of a murine-human chimeric antibody

QVQLVQSGAEVKKPGASVKVSCASGYTFTNYGMSWVRQAPGQGLEWMGWINTYSGESTY
ADDFKGRVTMTTDTSTSTAYMELRSLRSDDTAVYYCARSLNRDTWFTYWGQGTLLTVSS

SEQ ID NO: 32 – Light chain variable region amino acid sequence of a murine-human chimeric antibody

QTVVTQEPSFSVSPGGTVTLTCRSITGAVTTSNYANWYQQTPGQAPRTLIYGTNNRAPGVPDR
FSGSILGNKAALTITGAQADDESDYYCALWYSNHWVFGGGTKLTVL

SEQ ID NO: 33 – Humanized heavy chain variable region amino acid sequence

QVQLVQSGAEVKKPGASVKVSCASGYTFTNYGMSWVRQAPGQGLEWMGWINTYSGESTY
ADDFKGRVTMTTDTSTSTAYMELRSLRSDDTAVYYCARSLNRDTWFTYWGQGTLLTVSS

SEQ ID NO: 34 – Humanized heavy chain variable region amino acid sequence

QVQLVQSGAEVKKPGASVKVSCASGYTFTNYGMSWVRQAPGQGLEWMGWINTYSGESTY
ADDFKGRFTMTLDTSTSTAYMELRSLRSDDTAVYFCARSLNRDTWFTYWGQGTLLTVSS

SEQ ID NO: 35 – Humanized heavy chain variable region amino acid sequence

QVQLVQSGAEVKKPGASVKVSCASGYTFTNYGMSWVRQAPGQGLEWMGWINTYSGESTY
ADDFKGRFTFTLDTSTSTAYMELRSLRSDDTATYFCARSLNRDTWFTYWGQGTLLTVSS

SEQ ID NO: 36 – Humanized light chain variable region amino acid sequence

QTVVTQEPSFSVSPGGTVTLTCRSITGAVTTSNYANWYQQTPGQAPRGLIGGTNNRAPGVPD
RFSGSILGNKAALTITGAQADDESDYFCALWYSNHWVFGGGTKLTVL

SEQ ID NO: 37 – Humanized light chain variable region amino acid sequence

QTVVTQEPSFSVSPGGTVTLTCRSITGAVTTSNYANWVQETPGQAPRGLIGGTNNRAPGVPD
RFSGSILGNKAALTITGAQADDEADYFCALWYSNHWVFGGGTKLTVL

SEQ ID NO: 38 – Humanized heavy chain amino acid sequence

QVQLVQSGAEVKKPGASVKVSCASGYTFTNYGMSWVRQAPGQGLEWMGWINTYSGESTY
ADDFKGRVTMTTDTSTSTAYMELRSLRSDDTAVYYCARSLNRDTWFTYWGQGTLLTVSSAST
KGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLS

SVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKP
KDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLH
QDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYP
SDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHY
TQKSLSLSPGK

SEQ ID NO: 39 – Humanized heavy chain nucleic acid sequence

cagggtgcagctgggtgcagctcggagcagaggtgaagaagccaggagcctctgtgaaggtgagctgcaaggcctccggctacacctt
cacaaactatggcatgagctgggtgcggcaggcacctggacagggactggagtgatgggctggatcaatacactccggcgag
tctacctatgccgacgacttcaaggcagggtgacaatgaccacagacacctctacaagcaccgcctacatggagctgcggagcct
gagatccgacgataccgccgtgtactattgtgccaggtccctgaaccgcgatacatggttcacctattggggccagggcacactgggtg
accgtgagctccgccagcaccaagggcccttccgtgttccactggccccctctctaaatccacatctggcggcaccgccgcccctgg
gctgtctggtgaaggactacttcccagagcctgtgacagtgctcctggaactctggcgccctgacatccggcggtgcacacatttccagcc
gtgtgcagagctccggcctgtacagcctgtctagcgtggtgacagtgccctcctctagcctgggcacacagacctatactgcaacgt
gaatcacaagccaagcaataccaaggtggacaagaaggtggagcccaagtcctgtgataagacacacacctgccccctgtcct
gctcccagctgtctggcgccctagcgtgttctgttccaccaagcctaaggacacctgatgatctccggacacccgagggtga
cctgcgtggtgggtgacgtgtctcacgaggatctgaggtgaagtcaactggtatgtggatggcgtggaggtgcacaaatgccaagac
caagcccagagaggagcagtacaactctacatataggggtggtagcgtgtgaccgtgtgcaccaggactggctgaacgggaag
gagtataagtgaaggtgtccaataaggccctgccgcccccatcgagaagacaatcagcaaggccaagggccagcctcgggag
ccacaggtgtacacctgacctcatccagagacgagctgacaaagaaccaggtgtctctgacatgtctggtgaagggcttctatccta
gcatatcgccgtggagtgaggagtgccaatggccagccagagaacaattacaagaccacacccctgtgctggactccgatggctcc
ttctttctgtattccaagctgaccgtggataagtctcgttggcagcaggggaacgtgttcagctgttccgtgatgcacgaagccctgcata
atcactatactcagaaatccctgtccctgtcaccttgaaag

SEQ ID NO: 40 – Humanized light chain amino acid sequence

QTVVTQEPSFSVSPGGTVTLTCRSITGAVTTSNYANWVQETPGQAPRGLIGGTNNRAPGVDP
RFSGSILGNKAALTITGAQADDEADYFCALWYSNHWVFGGGTKLTVLGQPKAAPSVTLFPPSS
EELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNKYAASSYLSLTPEQW
KSHRSYSCQVTHEGSTVEKTVAPTECS

SEQ ID NO: 41 – Humanized light chain nucleic acid sequence

cagacagtggtgacctcaggagcctagcttctccgtgtctccaggaggaaccgtgacactgacctgcaggtctatcacaggcgccgtg
accacaagcaactacgccaattgggtgcaggagacaccaggacaggcaccaaggggactgatcggaggaaccaacaataggg
caccaggagtgacctgaccggttcagcgatccatcctgggaacaaggccgcccctgacaatcaccggagcagaggcagacgatg
aggcagattactctgtgccctgtggtattccaatcactgggtgttggcgccggcacaagctgacctgtctggggcagcccaaggc
cgctcctagcgtgacactgttcccccttctccgaggagctgcaggccaacaaggccaccctggtgtgctgatctccgacttctatcct
ggcgccgtgacagtgccctggaaggctgattctagcccagtgaggctggcgtggagaccacaacccccctcaagcagcttaaca
ataagtatgccgttctcttacctgagcctgacaccagagcagtggaagtcccaccggtcttacagtgccaggtcactcacgaagg
ctctaccgtggaaaagacagctgcacccaccgaatgctcatgataagcttaaggggttcgatccct

SEQ ID NO: 42 – Humanized heavy chain amino acid sequence

QVQLVQSGAEVKKPGASVKVSCKASGYTFTNYGMSWVRQAPGQGLEWMGWINTYSGESTY
ADDFKGRFTMTLDTSTSTAYMELRSLRSDDTAVYFCARSLNRDTWFTYWGQGLTVTVSSAST
KGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLS
SVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKP
KDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLH
QDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYP
SDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHY
TQKSLSLSPGK

SEQ ID NO: 43 – Humanized heavy chain nucleic acid sequence

cagggtgcagctgggtgcagctctggagcagaggtgaagaagccaggagcctctgtgaaggtgagctgaaggcctccggctacacctt
cacaaactatggcatgagctgggtgcggcaggcacctggacagggactggagtgatgggctggatcaatacactccggcgag
tctacctatgccgacgatttcaagggcagggttaccatgacactggacacctctacaagcaccgcctacatggagctgcggagcctga
gatccgacgatacagccgtgtatttctgtgccaggtccctgaaccgcgatacatggtttacctactggggccagggcacactgggtgacc
gtgagctccggcagcaccaagggcccttccgtgtttccactggccccctcctctaaatccacatctggcggcaccgcggcctgggt
gtctgtgaaggactacttccagagcctgtgacagtgctctggaactctggcgcctgacatccggcgtgcacacatttccagccgtg
ctgcagagctccggcctgtacagcctgtctagcgtgggtgacagtgccctcctctagcctgggcacacagacctatatctgaacgtga
atcacaagccaagcaataccaaggtggacaagaaggtggagccaagtctgtgataagacacacacctgccccctgtctgtct
cccgagctgtggcgccctagcgtgttctgtttccaccaagcctaaggacacctgatgatctccgggacacccgaggtgacct
gctgtgtgtggacgtgtctcacgaggatcctgaggtgaagttcaactggatgtggatggcgtggaggtgcacaatgccaaagacca
agccagagaggagcagtacaactctacatataggggtggtagcgtgtgaccgtgtgcaccaggactggctgaacggcaagga
gtataagtcaaggtgtccaataagggccctgcccgcctccatcgagaagacaatcagcaaggccaagggccagcctcgggagcc
acaggtgtacacctgctccatccagagacgagctgacaaagaaccaggtgtctctgacatgtctgtgaagggcttctatcctagc
gatatcgccgtggagtgaggagtgccaatggccagccagagaacaattacaagaccacacccctgtgtgactccgatggctccttc
tttctgtattccaagctgacctgggataagtctcggtggcagcagggcaacgtgttcagctgttccgtgatgcacgaagccctgcataat
cactatactcagaatccctgtccctgtcacctggaaag

SEQ ID NO: 44 – Humanized heavy chain amino acid sequence

QVQLVQSGAEVKKPGASVKVSCKASGYTFTNYGMSWVRQAPGQGLEWMGWINTYSGESTY
ADDFKGRFTFTLDTSTSTAYMELRSLRSDDTATYFCARSLNRDTWFTYWGQGLTVTVSSASTK
GPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLS
VVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKP
DTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQ
DWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPS
DIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT
QKSLSLSPGK

SEQ ID NO: 45 – Humanized heavy chain nucleic acid sequence

cagggtgcagctgggtgagctctggagcagaggtgaagaagccaggagcctctgtgaaggtagctgcaaggcctccggctacacctt
 cacaactatggcatgagctgggtgcggcaggcacctggacagggactggagtggatgggtggatcaatacactccggcgag
 tctacctatgccgacgacttcaagggcaggttcacctttacactggacacctctacaagcaccgctacatggagctgcggagcctga
 gatccgacgatacagccacctatttctgtgccagggtccctgaaccgcgatacatgggttacctactggggccagggcacactgggtgac
 cgtgagctccgccagcaccaagggccctccgtgtttccactgccccctctctaaatccacatctggcggcaccgcccgtgggc
 tgtctgggtgaaggactacttcccagagcctgtgacagtgctcctggaactctggcgcctgacatccggcggtgcacacatttccagccgt
 gctgcagagctccggcctgtacagcctgtctagcgtgggtgacagtgcctcctctagcctgggcacacagacctatctgcaacgtg
 aatcacaagccaagcaataccaaggtggacaagaaggtggagcccaagtcctgtgataagacacacacctgccccctgtcctg
 ctcccgagctgctgggcccctagcgtgttctgtttccaccaagcctaaggacaccctgatgatctccggacaccggaggtgac
 ctgctgtgggtggagctgtctcacgaggatcctgaggtgaagttcaactggtatgtggatggcgtggaggtgcacaatgccaagacc
 aagcccagagaggagcagtagaactctacatataggggtggtagcgtgctgaccgtgctgcaccaggactggctgaacggcaagg
 agtataagtgaagggttccaataaggccctgccgcccccatcgagaagacaatcagcaaggccaagggccagcctcgggagc
 cacagggtgacacctgcctccatccagagacgagctgacaaagaaccagggtgtctctgacatgtctggtagaagggttctatcctag
 cgatatcgccgtggagtgaggatccaatggccagccagagaacaattacaagaccacccccctgtgctggactccgatggctcctt
 ctttctgtattccaagctgaccgtggataagctcgggtggcagcagggcaacgtgttcagctgttccgtgatgcacgaagccctgcataa
 tcactatactcagaaatccctgtccctgtcacctggaaag

SEQ ID NO: 46 – Humanized light chain amino acid sequence

QTVVTQEPSFSVSPGGTVTLTCRSITGAVTTSNYANWYQQTPGQAPRGLIGGTNNRAPGVDP
 RFGSILGNKAALTITGAQADDESDYFCALWYSNHWVFGGGTKLTVLGQPKAAPSVTLFPPSS
 EELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNKYAASSYLSLTPEQW
 KSHRSYSCQVTHEGSTVEKTVAPTECS

SEQ ID NO: 47 – Humanized light chain nucleic acid sequence

cagacagtggtgacctcaggagcctagcttctccgtgtctccaggaggaaccgtgacactgacctgcagggtctatcacaggcgccgtg
 accacaagcaactacgccaattggtatcagcagacaccaggacaggcaccaaggggactgatcggaggaaccaacaataggg
 caccaggagtgcctgaccggttcagcggatccatcctgggaacaaggccgcccgtgacaatcaccggagcacaggcagacgatg
 agtccgattacttctgtccctgtggtattctaactactgggtgtttggcggcggcacaaagctgaccgtgctggggcagcccaaggcc
 gtcctagcgtgacactgtttcccccttctccgaggagctgcaggccaacaaggccaccctggtgtcctgatctccgacttctatcct
 ggcgccgtgacagtggcctggaaggctgattctagcccagtgaggctggcgtggagaccacaacccccctcaagcagcttaaca
 ataagtatgccgttctcttacctgagcctgacaccagagcagtggaagtcaccgggtcttacagctgccagggtcactcacgaagg
 ctctaccgtggaaaagacagctgcacccaccgaatgctcatgataagcttaagggttcgatccct

SEQ ID NO: 48 – Light chain constant region amino acid sequence

RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKD
 STYLSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

SEQ ID NO: 49 – Light chain constant region amino acid sequence

GQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTTPSKQSN
NKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS

SEQ ID NO: 50 – Heavy chain constant region amino acid sequence

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY
SLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFP
PKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT
VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKG
FYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFSCSVMHEALH
NHYTQKSLSLSPGK

What is claimed is:

1. An isolated antibody or antigen-binding fragment thereof which specifically binds human endothelin A receptor (ET_AR) and comprises either: (a) a light chain CDR3 sequence identical, substantially identical or substantially similar to a CDR3 sequence selected from SEQ ID NOs: 16-18; (b) a heavy chain CDR3 sequence identical, substantially identical or substantially similar to a CDR3 sequence selected from SEQ ID NOs: 8-10; or (c) the light chain CDR3 sequence of (a) and the heavy chain CDR3 sequence of (b).
2. An isolated antibody or antigen-binding fragment thereof according to claim 1, further comprising an amino acid sequence selected from: (d) a light chain CDR1 sequence identical, substantially identical or substantially similar to a CDR1 sequence selected from SEQ ID NOs: 11-13; (e) a light chain CDR2 sequence identical, substantially identical or substantially similar to a CDR2 sequence selected from SEQ ID NOs: 14-15; (f) a heavy chain CDR1 sequence identical, substantially identical or substantially similar to a CDR1 sequence selected from SEQ ID NOS: 2-4; (g) a heavy chain CDR2 sequence identical, substantially identical or substantially similar to a CDR2 sequence selected from SEQ ID NOs: 5-7; (h) the light chain CDR1 sequence of (d) and the heavy chain CDR1 sequence of (f); and (i) the light chain CDR2 sequence of (e) and the heavy chain CDR2 sequence of (g).
3. An isolated antibody, or antigen-binding fragment thereof, which specifically binds human ET_AR and comprises: (a) a light chain CDR1 sequence identical, substantially identical or substantially similar to a CDR1 sequence selected from SEQ ID NOs: 11-13; (b) a light chain CDR2 sequence identical, substantially identical or substantially similar to a CDR2 sequence selected from SEQ ID NOs: 14-15; (c) a light chain CDR3 sequence identical, substantially identical or substantially similar to a CDR3 sequence selected from SEQ ID NOs: 16-18; (d) a heavy chain CDR1 sequence identical, substantially identical or substantially similar to SEQ ID NOS: 2-4; (e) a heavy chain CDR2 sequence identical, substantially identical or substantially similar to a CDR2 sequence selected from SEQ ID NOs: 5-7; and (f) a heavy chain CDR3

sequence identical, substantially identical or substantially similar to a CDR3 sequence selected from SEQ ID NOs: 8-10.

4. An isolated antibody or antigen-binding fragment thereof according to claim 3, which comprises: (a) a light chain CDR1 sequence identical, substantially identical or substantially similar to SEQ ID NO: 11; (b) a light chain CDR2 sequence identical, substantially identical or substantially similar to SEQ ID NO: 14; (c) a light chain CDR3 sequence identical, substantially identical or substantially similar to SEQ ID NO: 16 (d) a heavy chain CDR1 sequence identical, substantially identical or substantially similar to SEQ ID NO: 2; (e) a heavy chain CDR2 sequence identical, substantially identical or substantially similar to SEQ ID NO: 5; and (f) a heavy chain CDR3 sequence identical, substantially identical or substantially similar to SEQ ID NO: 8.

5. An isolated antibody or antigen-binding fragment thereof according to claim 3, which comprises: (a) a light chain CDR1 sequence identical, substantially identical or substantially similar to SEQ ID NO: 12; (b) a light chain CDR2 sequence identical, substantially identical or substantially similar to SEQ ID NO: 14; (c) a light chain CDR3 sequence identical, substantially identical or substantially similar to SEQ ID NO: 17; (d) a heavy chain CDR1 sequence identical, substantially identical or substantially similar to SEQ ID NO: 3; (e) a heavy chain CDR2 sequence identical, substantially identical or substantially similar to SEQ ID NO: 6; and (f) a heavy chain CDR3 sequence identical, substantially identical or substantially similar to SEQ ID NO: 9.

6. An isolated antibody or antigen-binding fragment thereof according to claim 3, which comprises: (a) a light chain CDR1 sequence identical, substantially identical or substantially similar to SEQ ID NO: 13; (b) a light chain CDR2 sequence identical, substantially identical or substantially similar to SEQ ID NO: 15; (c) a light chain CDR3 sequence identical, substantially identical or substantially similar to SEQ ID NO: 18; (d) a heavy chain CDR1 sequence identical, substantially identical or substantially similar to SEQ ID NO: 4; (e) a heavy chain CDR2 sequence identical, substantially identical or substantially similar to SEQ ID NO: 7; and (f) a

heavy chain CDR3 sequence identical, substantially identical or substantially similar to SEQ ID NO: 10.

7. An isolated antibody or antigen-binding fragment thereof which specifically binds human ET_AR and comprises either: (a) a heavy and/or light chain variable domain(s), the variable domain(s) having a set of three light chain CDR1, CDR2, and CDR3 identical, substantially identical or substantially similar to SEQ ID NOS: 11-13, 14-15, and 16-18, and/or a set of three heavy chain CDR1, CDR2, and CDR3 identical, substantially identical or substantially similar to SEQ ID NOS: 2-4, 5-7, and 8-10; and (b) a set of four variable region framework regions from a human immunoglobulin (IgG).

8. An isolated antibody or antigen-binding fragment thereof which binds to human ET_AR with substantially the same or greater K_d as a reference antibody; (b) competes for binding to human ET_AR with said reference antibody; or (c) is less immunogenic in a human subject than said reference antibody, wherein said reference antibody comprises the heavy chain variable domain sequence of SEQ ID NO: 27 and the light chain variable domain sequence of SEQ ID NO: 29.

9. The isolated antibody or antigen-binding fragment thereof according to any one of claims 1-8 that binds to ET_AR protein with a dissociation constant (K_D) of at least about 1x10⁻⁶ M, at least about 1x10⁻⁷ M, at least about 1x10⁻⁸ M, at least about 1x10⁻⁹ M, at least about 1x10⁻¹⁰ M, at least about 1x10⁻¹¹ M, or at least about 1x10⁻¹² M.

10. An isolated antibody or antigen-binding fragment thereof according to any one of claims 1-9 wherein the antibody or antigen-binding fragment is selected from a human antibody, a humanized antibody, chimeric antibody, a monoclonal antibody, a polyclonal antibody, a recombinant antibody, an antigen-binding antibody fragment, a single chain antibody, a diabody, a triabody, a tetrabody, a Fab fragment, a Fab' fragment, a Fab₂ fragment, a F(ab')₂ fragment, a domain antibody, an IgD antibody, an IgE antibody, an IgM antibody, an IgG1 antibody, an IgG2

antibody, an IgG3 antibody, an IgG4 antibody, or an IgG4 antibody having at least one mutation in the hinge region that alleviates a tendency to form intra H-chain disulfide bonds.

11. An isolated humanized antibody or antigen-binding fragment thereof that specifically binds human ET_AR and comprises a heavy chain variable region sequence identical, substantially identical or substantially similar to SEQ ID NOs: 33-35, and a light chain variable region sequence identical, substantially identical or substantially similar to SEQ ID NOs: 36-37.

12. An isolated humanized antibody or antigen-binding fragment thereof that specifically binds to human ET_AR and comprises the heavy chain sequence set forth in SEQ ID NO: 38, and the light chain sequence set forth in SEQ ID NO: 40.

13. An isolated humanized antibody or antigen-binding fragment thereof that specifically binds to human ET_AR and comprises the heavy chain sequence set forth in SEQ ID NO: 42, and the light chain sequence set forth in SEQ ID NO: 40.

14. An isolated humanized antibody or antigen-binding fragment thereof that specifically binds to human ET_AR and comprises the heavy chain sequence set forth in SEQ ID NO: 44, and the light chain sequence set forth in SEQ ID NO: 40.

15. A pharmaceutical composition comprising an isolated antibody or antigen-binding fragment thereof according to any one of claims 1-14 in admixture with a pharmaceutically acceptable carrier.

16. A method of treating a subject suffering from a ET_AR-associated disorder, comprising administering to said subject a therapeutically effective amount of an antibody or antigen-binding fragment thereof according to any one of claims 1-14;

17. A method according to claim 16, wherein the ET_AR-associated disorder is a metabolic disorder selected from the group consisting of diabetes, diabetic myopathy, diabetic

nephropathy, diabetic neuropathy, diabetic retinopathy, obesity, dyslipidemia, sarcopenic obesity, fatty liver disease (e.g., nonalcoholic fatty liver disease (NAFLD); nonalcoholic steatohepatitis (NASH)), insulin resistance, and hemochromatosis.

18. A method according to claim 17, further comprising administration of a glucose-lowering agent selected from the group consisting of anti-diabetic agents, anti-hyperglycemic agents, lipid lowering agents, anti-hypertensive agents, and anti-obesity agents.

19. A method according to claim 16, wherein the ET_AR-associated disorder is a cardiovascular disease selected from the group consisting of angina, heart attack (myocardial infarction), atherosclerosis, heart failure, cardiovascular disease, rheumatic heart disease, cardiac arrhythmias (abnormal heart rhythms), cerebrovascular disease, congenital heart defects, cardiomyopathy, left ventricular hypertrophy, right ventricular hypertrophy, post-infarction heart rupture, infections of the heart, coronary artery disease, peripheral arterial disease, renal artery stenosis, aortic aneurysm, myocardial diseases, heart valve disorders, myocarditis, pericarditis, and pulmonary arterial hypertension (PAH).

20. A method according to claim 19, further comprising administration of a second agent selected from the group consisting of antihypertensive agents, agents for treating chronic heart failure, atherosclerosis or related diseases.


21. An isolated immunoconjugate or fusion protein comprising an antibody or antigen-binding fragment thereof according to any one of claims 1-14 coupled to an effector molecule.

22. An isolated nucleic acid comprising a polynucleotide sequence encoding an antibody or antigen-binding fragment thereof according to any one of claims 1-14.

23. A recombinant expression vector comprising the isolated nucleic acid of claim 22.

24. A host cell comprising the vector of claim 23.

1/3




#17: 312A10E5 (A3)

#21: 336D8G9 (A2)

FIG.1

2/3



#17: 312A10E5 (A3)
#21: 336D8G9 (A2)

FIG. 2




FIG.3