



- (51) **International Patent Classification:**
C12N 1/20 (2006.01) *A61K 35/66* (2006.01)
- (21) **International Application Number:**
PCT/US2016/014524
- (22) **International Filing Date:**
22 January 2016 (22.01.2016)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
62/106,892 23 January 2015 (23.01.2015) US
62/260,882 30 November 2015 (30.11.2015) US
- (71) **Applicant:** NOVOZYMES A/S [DK/DK]; Krogshoejvej 36, DK-2880 Bagsvaerd (DK).
- (72) **Inventor; and**
- (71) **Applicant (for BW only):** LEWIS, Derrick [US/US]; 108 TW Alexander Drive Bldg 1A, Durham, North Carolina 27709 (US).
- (72) **Inventors:** COHN, Marianne Thorup; Krogshoejvej 36, DK-2880 Bagsvaerd (DK). NELSON, Adam; 5400 Corporate Circle, Salem, Virginia 24153 (US). NIELSEN, Preben; Krogshoejvej 36, DK-2880 Bagsvaerd (DK).
- (74) **Agents:** LAMBIRIS, Elias et al.; Novozymes North America, Inc., 60 East 42nd Street, Suite 700, New York, New York 10165 (US).

- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description (Rules 13bis.4(d)(i) and 48.2(a)(viii))
- with sequence listing part of description (Rule 5.2(a))

(54) **Title:** BACILLUS STRAINS IMPROVING HEALTH AND PERFORMANCE OF PRODUCTION ANIMALS

(57) **Abstract:** The present invention relates to Bacillus strains which improves health and performance of production animals. The invention further relates to compositions comprising the Bacillus strains.



WO 2016/118864 A1

Bacillus strains improving health and performance of production animals**Reference to a Deposit of Biological Material**

This application contains a reference to a deposit of biological material, which deposit
5 is incorporated herein by reference.

Reference to sequence listing

This application contains a Sequence Listing in computer readable form. The
computer readable form is incorporated herein by reference.

10 Index to sequence listing:

SEQ ID NO: 1 is 16S rDNA of DSM 29869.

SEQ ID NO: 2 is 16S rDNA of DSM 29870.

SEQ ID NO: 3 is 16S rDNA of DSM 29871.

SEQ ID NO: 4 is 16S rDNA of DSM 29872.

15 SEQ ID NO: 5 is 16S rDNA of *Bacillus vallismortis* from AB021198.

SEQ ID NO: 6 is 16S rDNA of *Bacillus subtilis* from AJ276351.

SEQ ID NO: 7 is 16S rDNA of *Bacillus amyloliquefaciens* from AB255669.

SEQ ID NO: 8 to SEQ ID NO: 13: PCR and sequencing primers.

BACKGROUND OF THE INVENTION**Field of the Invention**

The present invention relates to *Bacillus* strains which improves the health and
performance of production animals. The invention further relates to compositions comprising
25 the *Bacillus* strains.

Background of the Invention

Clostridium perfringens (*C. perfringens*) is a Gram-positive, rod-shaped, anaerobic,
spore-forming bacterium of the genus *Clostridium*. *C. perfringens* is widely present in nature
30 and can be found as a component of decaying vegetation, marine sediment, the intestinal
tract of humans and other vertebrates, insects, and soil.

Infections due to *C. perfringens* show evidence of tissue necrosis, bacteremia,
emphysematous cholecystitis, and gas gangrene, which is also known as clostridial
myonecrosis. *C. perfringens* can also result in polymicrobial anaerobic infections.

35 The incidence of *Clostridium perfringens*-associated necrotic enteritis in poultry has
increased in countries that stopped using antibiotic growth promoters. Necrotic enteritis is an
enterotoxemic disease that results in significant economic losses in the poultry industry.

There is a need to develop tools and strategies for the prevention and control of *C. perfringens* in mono-gastric animals such as poultry. Whilst the vaccination of animals has been suggested, there are challenges associated with vaccinating thousands of animals. Thus discovering a solution which could be administered as an additive in an animal feed would be advantageous.

It is thus an object of the invention to provide solutions which prevents and/or controls *C. perfringens* in mono-gastric animals such as poultry by use of an animal feed comprising one or more bacteria with activity against *Clostridium perfringens*.

A further object of the invention is to provide a solution that does not present a risk for the health of the animal. The solution to this problem is use of non-hemolytic *Bacillus* strains with activity against *Clostridium perfringens*.

A challenge of delivering *Bacillus* spp. in feed is the common use of antibiotics as growth promoters in feed. Therefore it is necessary to determine the compatibility of strains with commonly-used feed antibiotics such as monensin in order to identify any potential conflicts with use as a direct fed microbial. The present invention relates in one embodiment to a *Bacillus* strains with high compatibility with monensin.

Description of the Related Art

Knap et al. (2010) describes that *Bacillus licheniformis* has an effect on necrotic enteritis in broiler chickens (Knap, Lund, Kehlet, Hofacre, and Mathis; *Bacillus licheniformis* Prevents Necrotic Enteritis in Broiler Chickens; Avian Diseases 54(2):931-935. 2010). This *Bacillus licheniformis* strain has no effect on *Clostridium perfringens* *in vitro* as demonstrated in Example 2.

Clostat (alias *Bacillus* PB6) is described in WO2007/064741. *Bacillus* PB6 has antagonistic effect against *C. perfringens*. *Bacillus* PB6 is hemolytic as described in Example 1.

SUMMARY OF THE INVENTION

It has been surprisingly found that the addition of direct fed microbes (DFM) from *Bacillus* species to animal feed can be used to prevent and/or control *C. perfringens* infections and/or necrotic enteritis in mono-gastric animal such as pigs and/or poultry. The *Bacillus* species can also improve the body weight gain and/or feed conversion rate (e.g., in both *Clostridium perfringens* challenged and unchallenged chickens).

In a first aspect the invention relates to a *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
 - ii) the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens*
- and

iii) the *Bacillus* strain improves body weight gain and/or feed conversion rate in chickens fed with the *Bacillus* strain.

In a second aspect the invention relates to a *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
- 5 ii) the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens* and
- iii) the *Bacillus* strain is sensitive to at least seven such as eight of the antibiotics selected from the group consisting of Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline.

10 In a third aspect the invention relates to a *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
- ii) the *Bacillus* strain is sensitive to at least seven such as eight of the antibiotics selected from the group consisting of Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline and

15 iii) the *Bacillus* strain improves body weight gain and/or feed conversion rate in chickens fed with the *Bacillus* strain.

In a fourth aspect the invention relates to a *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
- ii) the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens*
- 20 and optionally
- iii) the *Bacillus* strain has enzymatic activity under aerobic and/or anaerobic conditions that hydrolyzes one or more of the substrates selected from the group consisting of Amylose, Arabinan, Arabinoxylan, Casein and Xylan.

The invention further relates to the strain having the deposit accession number DSM 25 29869; a strain having all of the identifying characteristics of *Bacillus* DSM 29869 or a mutant thereof.

The invention further relates to the strain having the deposit accession number DSM 29870; a strain having all of the identifying characteristics of *Bacillus* DSM 29870 or a mutant thereof.

30 The invention further relates to the strain having the deposit accession number DSM 29871; a strain having all of the identifying characteristics of *Bacillus* DSM 29871 or a mutant thereof.

The invention further relates to the strain having the deposit accession number DSM 29872; and a strain having all of the identifying characteristics of *Bacillus* DSM 29872 or a 35 mutant thereof.

The present invention also relates to compositions comprising one or more of the *Bacillus* strains according to the invention which improves health and performance of production animals.

In a preferred embodiment the invention relates to a composition, e.g., comprising a carrier and a *Bacillus* strain wherein:

i. the *Bacillus* strain is selected from the group consisting of:

the strain having the deposit accession number DSM 29869; a strain having all of the identifying characteristics of *Bacillus* DSM 29869 or a mutant thereof,

the strain having the deposit accession number DSM 29870; a strain having all of the identifying characteristics of *Bacillus* DSM 29870 or a mutant thereof,

the strain having the deposit accession number DSM 29871; a strain having all of the identifying characteristics of *Bacillus* DSM 29871 or a mutant thereof,

the strain having the deposit accession number DSM 29872; and a strain having all of the identifying characteristics of *Bacillus* DSM 29872 or a mutant thereof,

and

ii. the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens* (such as *Clostridium perfringens* strains 23 or 48 [Gholamiandekhordi AR, Ducatelle R, Heyndrickx M, Haesebrouck F, Van Immerseel F. 2006. Molecular and phenotypical characterization of *Clostridium perfringens* isolates from poultry flocks with different disease status. Vet. Microbiol. 113:143–152] and/or *E. coli* (such as ATCC10536 or ATCC25922).

In a preferred embodiment the *Bacillus* strain has a high compatibility with monensin such as being compatible with at least 2.4 µg/ml monensin as determined in Example 12.

BRIEF DESCRIPTION OF DRAWINGS

Fig. 1 Alignment of 16 S rDNA sequences. SEQ ID NO: 1 (DSM 29869), SEQ ID NO: 2 (DSM 29870), SEQ ID NO: 3 (DSM 29871), SEQ ID NO: 4 (DSM 29872), SEQ ID NO: 5 (*Bacillus vallismortis* AB021198), SEQ ID NO: 6 (*Bacillus subtilis* from AJ276351), and SEQ ID NO: 7 (*Bacillus amyloliquefaciens* AB255669) have been aligned. The region covering position 481-1200 is not shown in Fig. 1 (in this region all sequences are identical).

DEFINITIONS

In general, the terms and phrases used herein have their art-recognized meaning, which can be found by reference to standard texts, journal references, and context known to those skilled in the art. The following definitions are provided to clarify their specific use in context of the disclosure.

As used herein, the singular forms "a", "an" and "the" are intended to include the plural forms as well, unless the context clearly indicates otherwise.

Antimicrobial activity against *Clostridium perfringens*: The term "Antimicrobial activity against *Clostridium perfringens*" means that the growth of *Clostridium perfringens* is inhibited and/or that some or all of the *Clostridium perfringens* are killed. This can be determined by the assay described in Example 2.

Body Weight Gain: The Body Weight Gain of an animal is the increase of body weight of the animal over a specified time period. An example of Body Weight Gain determination is given in Example 8.

Composition: The term "composition" refers to a composition comprising a carrier and at least one bacterial strain as described herein. The compositions described herein may be mixed with an animal feed(s) and referred to as a "mash feed."

Control *C. perfringens* infections and/or necrotic enteritis: The term "control *C. perfringens* infections and/or necrotic enteritis" means a method and/or composition that partly or completely inhibits *C. perfringens* infections and/or necrotic enteritis in an animal. Accordingly, the term "control *C. perfringens* infections and/or necrotic enteritis" means the *C. perfringens* infections and/or the necrotic enteritis is reduced or completely eliminated or prevented.

Direct Fed Microbial: The term "direct fed microbial" means live micro-organisms including spores which, when administered in adequate amounts, confer a benefit, such as improved digestion or health, on the host.

Enzymatic activity under aerobic conditions: The term "enzymatic activity under aerobic conditions" means activity of enzymes produced by a *Bacillus* strain during growth under aerobic conditions as described in Example 6.

Enzymatic activity under anaerobic conditions: The term "enzymatic activity under anaerobic conditions" means activity of enzymes produced by a *Bacillus* strain during growth under anaerobic conditions as described in Example 6.

European Production Efficacy Factor (EPEF): The European Production Efficacy Factor is a way of comparing the live-bird performance of flocks. This single-figure facilitates comparison of performance within and among farms and can be used to assess environmental, climatic and managerial variables. The EPEF is calculated as $[(\text{liveness} (\%) \times \text{Liveweight (kg)}) / (\text{Age at depletion (days)} \times \text{FCR})] \times 100$, wherein liveness is the percentage of birds alive at slaughter, Liveweight is the average weight of the birds at slaughter, age of depletion is the age of the birds at slaughter and FCR is the feed conversion ratio at slaughter.

Fed: The term "fed" means any type of oral administration such as administration via an animal feed or via drinking water.

FCR (Feed Conversion Rate): FCR is a measure of an animal's efficiency in converting feed mass into increases of the desired output. Animals raised for meat – such as swine, poultry and fish – the output is the mass gained by the animal. Specifically FCR is the mass of the food eaten divided by the output, all over a specified period. FCR can be determined as described in Example 8. Improvement in FCR means reduction of the FCR value. A FCR improvement of 2% means that the FCR was reduced by 2%.

Isolated: The term “isolated” means that the one or more bacterial strains described herein are in a form or environment which does not occur in nature, that is, the one or more bacterial strains are at least partially removed from one or more or all of the naturally occurring constituents with which it is associated in nature.

Non-hemolytic: Hemolysis is the breakdown of red blood cells. The ability of bacterial colonies to induce hemolysis when grown on blood agar is used to classify bacterial strains into hemolytic and non-hemolytic strains. In this context hemolysis is defined as described in EFSA Journal 2011;9(11):2445, using *Bacillus subtilis* 168 (BGSC-1A1, Bacillus Genetic Stock Center) as a negative control. A *Bacillus* strain can be classified as non-hemolytic using the assay described in Example 1.

Poultry: The term “poultry” means domesticated birds kept by humans for the eggs they produce and/or their meat and/or their feathers. Poultry includes broilers and layers. Poultry include members of the superorder Galloanserae (fowl), especially the order Galliformes (which includes chickens, Guineafowls, quails and turkeys) and the family Anatidae, in order Anseriformes, commonly known as “waterfowl” and including domestic ducks and domestic geese. Poultry also includes other birds that are killed for their meat, such as the young of pigeons. Examples of poultry include chickens (including layers, broilers and chicks), ducks, geese, pigeons, turkeys and quail.

Prevent *C. perfringens* infections and/or necrotic enteritis: The term “prevent *C. perfringens* infections and/or necrotic enteritis” means a method and/or composition that prevents development of a *C. perfringens* infection and/or necrotic enteritis in an animal.

Sensitive to antibiotics: The term “sensitive to antibiotics” means the phenotypic property of a bacterial strain, that growth of said bacterial strain is inhibited under conditions where the bacterial strain would otherwise grow. In this context sensitivity to antibiotics is tested after the CLSI guidelines (M07-A9 Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; 2012). The *S. aureus* ATCC 29213 is used as reference strain, which means that it should be included in the test, and that the results are only valid if *S. aureus* ATCC 29213 show results in compliance with the breakpoints of the CLSI guideline (see Example 5 Table 5.5) (M100-S24 Performance Standards for Antimicrobial Susceptibility Testing; informational Supplement, 2014). A strain of *Bacillus* is

considered sensitive if the growth is detected at or below the breakpoint concentration of the appropriate antibiotic specified in EFSA journal 2012; 10(6):2740.

Spore: The terms "spore" and "endospore" are interchangeable and have their normal meaning which is well known and understood by those of skill in the art. As used
5 herein, the term spore refers to a microorganism in its dormant, protected state.

Stable: The term "stable" is a term that is known in the art, and in a preferred aspect, stable is intended to mean the ability of the microorganism to remain in a spore form until it is administered to an animal to improve the health of the animal.

Swine: The term "swine" or "pigs" means domesticated pigs kept by humans for food,
10 such as their meat. Swine includes members of the genus *Sus*, such as *Sus scrofa domesticus* or *Sus domesticus* and include piglets, growing pigs, and sows.

DETAILED DESCRIPTION OF THE INVENTION

It has been surprisingly found that the addition of direct fed microbes (DFM) from
15 *Bacillus* species to animal feed can be used to prevent and/or control *C. perfringens* infections and/or necrotic enteritis in mono-gastric animal such as pigs and/or poultry and at the same time improve the body weight gain and/or feed conversion rate (in both *Clostridium perfringens* challenged and unchallenged chickens).

Bacillus strains of the invention

The invention relates to the following aspects with respect to *Bacillus* strains:

Aspect 1: A *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
- 25 ii) the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens* and
- iii) the *Bacillus* strain improves body weight gain and/or feed conversion rate in chickens fed with the *Bacillus* strain.

Aspect 2: A *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
- ii) the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens* and
- iii) the *Bacillus* strain is sensitive to at least seven such as eight of the antibiotics
35 selected from the group consisting of Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline.

Aspect 3: A *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
- ii) the *Bacillus* strain is sensitive to at least seven such as eight of the antibiotics selected from the group consisting of Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline and
- iii) the *Bacillus* strain improves body weight gain and/or feed conversion rate in chickens fed with the *Bacillus* strain.

Aspect 4: A *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
- ii) the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens* and optionally
- iii) the *Bacillus* strain has enzymatic activity under aerobic and/or anaerobic conditions that hydrolyzes one or more of the substrates selected from the group consisting of Amylose, Arabinan, Arabinoxylan, Casein and Xylan.

In a preferred embodiment the *Bacillus* according to Aspect 1, 2 3 or 4 has a high compatibility with monensin such as being compatible with at least 2.3 µg/ml monensin as determined in Example 12. It is even more preferred that the *Bacillus* strain is compatible with at least 2.4 µg/ml monensin as determined in Example 12 (such as at least 2.5 µg/ml monensin as determined in Example 12, such as at least 2.6 µg/ml monensin as determined in Example 12 or such as at least 2.7 µg/ml monensin as determined in Example 12).

Sensitivity to Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline can, e.g., be determined as described in Example 5.

Enzymatic activity under aerobic and/or anaerobic conditions that hydrolyzes one or more of the substrates selected from the group consisting of Amylose, Arabinan, Arabinoxylan, Casein and Xylan can, e.g., be determined as described in Example 6.

In one embodiment the *Bacillus* strain according to Aspect 1, 2 or 3 has enzymatic activity under aerobic and/or anaerobic conditions that hydrolyzes one or more of the substrates selected from the group consisting of Amylose, Arabinan, Arabinoxylan, Casein and Xylan.

In one embodiment the *Bacillus* strain according to Aspect 2, 3 or 4 improves one or more performance parameters in poultry selected from the list consisting of body weight gain, European Production Efficacy Factor and feed conversion rate in chickens fed with the *Bacillus* strain.

In one embodiment, the *Bacillus* strain according to Aspect 3 has antimicrobial activity against *Clostridium perfringens*. Antimicrobial activity against *Clostridium perfringens* can, e.g., be determined as described in Example 2.

In one embodiment the *Bacillus* strain according to Aspect 1 or 4 is sensitive to at least seven such as eight of the antibiotics selected from the group consisting of Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline.

In one embodiment the improvement of feed conversion rate (FCR) for Aspect 1, 2, 3 or 4 results in a FCR of -2.5% or less than -2.5%, such as less than -2.6%, such as less than -2.7%, such as less than -2.8%, such as less than -2.9%, such as less than -3.0%. In a preferred embodiment the improvement of FCR results in a FCR of from -5% to -2% such as a FCR from -4% to -2%, such as a FCR of from -3.5% to -2.5%. In a specific embodiment the improvement of FCR for Aspect 1, 2, 3 or 4 results in a FCR within an interval selected from the group consisting of from -5% to -4.5%, from -4.5% to -4%, from -4% to -3.8%, from -3.8% to -3.6%, from -3.6% to -3.4%, from -3.4% to -3.2%, from -3.2% to -3.0%, from -3.0% to -2.8% and from -2.8% to -2.5%, or any combination of these intervals. The FCR can be determined as described in Example 8.

In one embodiment the improvement in body weight gain for Aspect 1, 2, 3 or 4 results in a body weight gain of at least 0.5%, such as at least 0.8%, such as at least 1.5%, such as at least 1.8%, such as at least 2.0%, such as at least 2.3%, such as at least 3.5%, such as at least 4.2%, such as at least 5.2%, such as at least 6.5%, such as at least 7.3%. In a preferred embodiment the improvement in body weight gain for Aspect 1, 2, 3 or 4 results in a body weight gain selected from the group consisting of from 1.8% to 2.0%, from 2.0% to 2.2%, from 2.2% to 2.4%, from 2.4% to 2.6%, from 2.6% to 2.8%, from 2.8% to 3.0%, from 3.0% to 3.2%, from 3.2% to 3.4%, from 3.4% to 3.6%, from 3.6% to 3.8%, from 3.8% to 4.0%, from 4% to 5%, from 5% to 7%, from 7% to 10%, or any combination thereof. The body weight gain can be determined as described in Example 8.

In one embodiment the *Bacillus* strain according to Aspect 1, 2, 3 or 4 wherein the *Bacillus* strain comprises 16S rDNA that is more than 98% (such as more than 98.5%, such as more than 99%, such as more than 99.5%, such as more than 99.5%) sequence identity to SEQ ID NO: 1.

In one embodiment the *Bacillus* strain according to Aspect 1, 2, 3 or 4 wherein the *Bacillus* strain comprises 16S rDNA that is more than 98% (such as more than 98.5%, such as more than 99%, such as more than 99.5%) sequence identity to SEQ ID NO: 2.

In one embodiment the *Bacillus* strain according to Aspect 1, 2, 3 or 4 wherein the *Bacillus* strain comprises 16S rDNA that is more than 98% (such as more than 98.5%, such as more than 99%, such as more than 99.5%) sequence identity to SEQ ID NO: 3.

In one embodiment the *Bacillus* strain according to Aspect 1, 2, 3 or 4 wherein the *Bacillus* strain comprises 16S rDNA that is more than 98% (such as more than 98.5%, such as more than 99%, such as more than 99.5%) sequence identity to SEQ ID NO: 4.

5 In one embodiment the *Bacillus* strain according to Aspect 1, 2, 3 or 4 is a *Bacillus subtilis* strain, a *Bacillus amyloliquefaciens* strain or a *Bacillus licheniformis* strain.

In one embodiment the *Bacillus* strain according to Aspect 1, 2, 3 or 4 has antimicrobial effect against *E. coli*. The effect against *E. coli* can, e.g., be determined as described in Example 4.

10 In one embodiment the *Bacillus* strain according to Aspect 1, 2, 3 or 4 is the *Bacillus* strain having deposit accession number DSM 29869, or a strain having all of the identifying characteristics of *Bacillus* DSM 29869 or a mutant thereof that has antimicrobial activity against *Clostridium perfringens*. In another embodiment the *Bacillus* strain according to Aspect 1, 2, 3 or 4 is the *Bacillus* strain having deposit accession number DSM 29870, or a strain having all of the identifying characteristics of *Bacillus* DSM 29870 or a mutant thereof
15 that has antimicrobial activity against *Clostridium perfringens*. In yet another embodiment the *Bacillus* strain according to Aspect 1, 2, 3 or 4 is the *Bacillus* strain having deposit accession number DSM 29871, or a strain having all of the identifying characteristics of *Bacillus* DSM 29871 or a mutant thereof that has antimicrobial activity against *Clostridium perfringens*. In a further embodiment the *Bacillus* strain according to Aspect 1, 2, 3 or 4 is the *Bacillus* strain
20 having deposit accession number DSM 29872, or a strain having all of the identifying characteristics of *Bacillus* DSM 29872 or a mutant thereof that has antimicrobial activity against *Clostridium perfringens*.

The invention relates in one embodiment to a *Bacillus* strain having deposit accession number DSM 29869 or a strain having all of the identifying characteristics of *Bacillus* DSM
25 29869 or a mutant thereof. The invention relates in another embodiment to a *Bacillus* strain having deposit accession number DSM 29870 or a strain having all of the identifying characteristics of *Bacillus* DSM 29870 or a mutant thereof. The invention relates in another embodiment to a *Bacillus* strain having deposit accession number DSM 29871 or a strain having all of the identifying characteristics of *Bacillus* DSM 29871 or a mutant thereof. The
30 invention relates in a further embodiment to a *Bacillus* strain having deposit accession number DSM 29872 or a strain having all of the identifying characteristics of *Bacillus* DSM 29872 or a mutant thereof.

The invention also relates to a biologically pure culture of the *Bacillus* strain according to aspect 1, 2, 3 or 4. In a further embodiment the invention relates to a biologically pure
35 culture of the *Bacillus* strain having deposit accession number DSM 29869 or a strain having all of the identifying characteristics of *Bacillus* DSM 29869 or a mutant thereof. In a further embodiment the invention relates to a biologically pure culture of the *Bacillus* strain having

deposit accession number DSM 29870 or a strain having all of the identifying characteristics of *Bacillus* DSM 29870 or a mutant thereof. In a further embodiment the invention relates to a biologically pure culture of the *Bacillus* strain having deposit accession number DSM 29871 or a strain having all of the identifying characteristics of *Bacillus* DSM 29871 or a mutant thereof. In a further embodiment the invention relates to a biologically pure culture of the *Bacillus* strain having deposit accession number DSM 29872 or a strain having all of the identifying characteristics of *Bacillus* DSM 29872 or a mutant thereof.

The invention also relates to an isolated *Bacillus* strain according to aspect 1, 2, 3 or 4. In a further embodiment the invention also relates to an isolated *Bacillus* strain having deposit accession number DSM 29869 or an isolated strain having all of the identifying characteristics of *Bacillus* DSM 29869 or a mutant thereof. In a further embodiment the invention also relates to an isolated *Bacillus* strain having deposit accession number DSM 29870 or an isolated strain having all of the identifying characteristics of *Bacillus* DSM 29870 or a mutant thereof. In a further embodiment the invention also relates to an isolated *Bacillus* strain having deposit accession number DSM 29871 or an isolated strain having all of the identifying characteristics of *Bacillus* DSM 29871 or a mutant thereof. In a further embodiment the invention also relates to an isolated *Bacillus* strain having deposit accession number DSM 29872 or an isolated strain having all of the identifying characteristics of *Bacillus* DSM 29872 or a mutant thereof.

Compositions of the invention

The invention relates to a composition comprising spores of one or more *Bacillus* strains according to invention.

More specifically the invention relates to the following aspects with respect to compositions comprising *Bacillus* strains:

Aspect 5: A composition comprising spores of a *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
- ii) the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens* and
- iii) the *Bacillus* strain improves body weight gain and/or feed conversion rate in chickens fed with the *Bacillus* strain.

Aspect 6: A composition comprising spores of a *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
- ii) the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens* and

iii) the *Bacillus* strain is sensitive to at least seven such as eight of the antibiotics selected from the group consisting of Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline.

5 Aspect 7: A composition comprising spores of a *Bacillus* strain characterized in that:

i) the *Bacillus* strain is non-hemolytic,

ii) the *Bacillus* strain is sensitive to at least seven such as eight of the antibiotics selected from the group consisting of Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline and

10 iii) the *Bacillus* strain improves body weight gain and/or feed conversion rate in chickens fed with the *Bacillus* strain.

Aspect 8: A composition comprising spores of a *Bacillus* strain characterized in that:

i) the *Bacillus* strain is non-hemolytic,

15 ii) the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens* and optionally

iii) the *Bacillus* strain has enzymatic activity under aerobic and/or anaerobic conditions that hydrolyzes one or more of the substrates selected from the group consisting of Amylose, Arabinan, Arabinoxylan, Casein and Xylan.

20 In a preferred embodiment the *Bacillus* strain(s) of the composition according to Aspect 5, 6, 7 or 8 has a high compatibility with monensin such as being compatible with at least 2.3 µg/ml monensin as determined in Example 12. It is even more preferred that the *Bacillus* strain is compatible with at least 2.4 µg/ml monensin as determined in Example 12 (such as at least 2.5 µg/ml monensin as determined in Example 12, such as at least 2.6
25 µg/ml monensin as determined in Example 12 or such as at least 2.7 µg/ml monensin as determined in Example 12).

Sensitivity to Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline can, e.g., be determined as described in Example 5.

30 Enzymatic activity under aerobic and/or anaerobic conditions that hydrolyzes one or more of the substrates selected from the group consisting of Amylose, Arabinan, Arabinoxylan, Casein and Xylan can, e.g., be determined as described in Example 6.

In one embodiment the *Bacillus* strain of the composition according to Aspect 5, 6 or 7 has enzymatic activity under aerobic and/or anaerobic conditions that hydrolyzes one or
35 more of the substrates selected from the group consisting of Amylose, Arabinan, Arabinoxylan, Casein and Xylan.

In one embodiment the *Bacillus* strain of the composition according to Aspect 6, 7, or 8 improves one or more performance parameters in poultry selected from the list consisting of body weight gain, European Production Efficacy Factor and feed conversion rate in chickens fed with the *Bacillus* strain.

5 In one embodiment the *Bacillus* strain of the composition according to Aspect 8 has antimicrobial activity against *Clostridium perfringens*. Antimicrobial activity against *Clostridium perfringens* can, e.g., be determined as described in Example 2.

10 In one embodiment the *Bacillus* strain of the composition according to Aspect 5 or 8 is sensitive to at least seven such as eight of the antibiotics selected from the group consisting of Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline.

15 In one embodiment the improvement of feed conversion rate (FCR) for Aspect 5, 6, 7 or 8 results in a FCR of -2.5% or less than -2.5%, such as less than -2.6%, such as less than -2.7%, such as less than -2.8%, such less than -2.9%, such as less than -3.0%. In a preferred embodiment the improvement of FCR results in a FCR of from -5% to -2% such as a FCR from -4% to -2%, such as a FCR of from -3.5% to -2.5%. In a specific embodiment the improvement of FCR for Aspect 1, 2, 3 or 4 results in a FCR within an interval selected from the group consisting of from -5% to -4.5%, from -4.5% to -4%, from -4% to -3.8%, from -3.8% to -3.6%, from -3.6% to -3.4%, from -3.4% to -3.2%, from -3.2 to -3.0%, from -3.0% to -2.8% and from -2.8 to -2.5%, or any combination of these intervals. The FCR can be determined as described in Example 8.

25 In one embodiment the improvement in body weight gain for Aspect 5, 6, 7 or 8 results in a body weight gain of at least 0.5%, such as at least 0.8%, such as at least 1.5%, such as at least 1.8%, such as at least 2.0%, such as at least 2.3%, such as at least 3.5%, such as at least 4.2%, such as at least 5.2%, such as at least 6.5%, such as at least 7.3%. In a preferred embodiment the improvement in body weight gain for Aspect 1, 2, 3 or 4 results in a body weight gain selected from the group consisting of from 1.8% to 2.0%, from 2.0% to 2.2%, from 2.2% to 2.4%, from 2.4% to 2.6%, from 2.6% to 2.8%, from 2.8% to 3.0%, from 3.0% to 3.2%, from 3.2% to 3.4%, from 3.4% to 3.6%, from 3.6% to 3.8%, from 3.8% to 4.0%, from 4% to 5%, from 5% to 7%, from 7% to 10%, or any combination thereof. The body weight gain can be determined as described in Example 8.

35 In one embodiment the *Bacillus* strain of the composition according to Aspect 5, 6, 7 or 8 comprises 16S rDNA that has more than 98% (such as more than 98.5%, such as more than 99%, such as more than 99.5%) sequence identity to SEQ ID NO: 1 and/or more than 98% (such as more than 98.5%, such as more than 99%, such as more than 99.5%) sequence identity to SEQ ID NO: 2 and/or more than 98% (such as more than 98.5%, such as more than 99%, such as more than 99.5%) sequence identity to SEQ ID NO: 3 and/or

more than 98% (such as more than 98.5%, such as more than 99%, such as more than 99.5%) sequence identity to SEQ ID NO: 4.

In one embodiment the *Bacillus* strain of the composition according to Aspect 5, 6, 7 or 8 is a *Bacillus subtilis* strain, or a *Bacillus amyloliquefaciens* strain.

5 In one embodiment the *Bacillus* strain of the composition according to Aspect 5, 6, 7 or 8 has antimicrobial effect against *E. coli*. The effect against *E. coli* can, e.g., be determined as described in Example 4.

In one embodiment the *Bacillus* strain of the composition according to Aspect 5, 6, 7 or 8 is the *Bacillus* strain having deposit accession number DSM 29869, or a strain having all
10 of the identifying characteristics of *Bacillus* DSM 29869 or a mutant thereof that has antimicrobial activity against *Clostridium perfringens*. In another embodiment the *Bacillus* strain of the composition according to Aspect 5, 6, 7 or 8 is the *Bacillus* strain having deposit accession number DSM 29870, or a strain having all of the identifying characteristics of
15 *Bacillus* DSM 29870 or a mutant thereof that has antimicrobial activity against *Clostridium perfringens*. In yet another embodiment the *Bacillus* strain of the composition according to Aspect 5, 6, 7 or 8 is the *Bacillus* strain having deposit accession number DSM 29871, or a strain having all of the identifying characteristics of *Bacillus* DSM 29871 or a mutant thereof that has antimicrobial activity against *Clostridium perfringens*. In a further embodiment the
20 *Bacillus* strain of the composition according to Aspect 5, 6, 7 or 8 is the *Bacillus* strain having deposit accession number DSM 29872, or a strain having all of the identifying characteristics of *Bacillus* DSM 29872 or a mutant thereof that has antimicrobial activity against *Clostridium perfringens*.

The invention relates in one embodiment to a composition comprising a *Bacillus* having deposit accession number DSM 29869 or a strain having all of the identifying
25 characteristics of *Bacillus* DSM 29869 or a mutant thereof. The invention relates in another embodiment to a composition comprising a *Bacillus* having deposit accession number DSM 29870 or a strain having all of the identifying characteristics of *Bacillus* DSM 29870 or a mutant thereof. The invention relates in another embodiment to a composition comprising a
30 *Bacillus* having deposit accession number DSM 29871 or a strain having all of the identifying characteristics of *Bacillus* DSM 29871 or a mutant thereof. The invention relates in a further embodiment to a composition comprising a *Bacillus* having deposit accession number DSM 29872 or a strain having all of the identifying characteristics of *Bacillus* DSM 29872 or a mutant thereof.

In one embodiment of Aspect 5, 6, 7, or 8 the *bacillus* spores of the composition are
35 present as dried spores such as spray-dried spores. In one embodiment of Aspect 5, 6, 7, or 8 the *bacillus* spores of the composition are present as stable spores. The composition

according to Aspect 5, 6, 7, or 8 can also be a liquid composition and/or comprise culture supernatant comprising one or more *Bacillus* strain(s) of the invention.

In one embodiment of Aspect 5, 6, 7, or 8 the composition further comprises a carrier.

The carrier can comprise one or more of the following compounds: water, glycerol, ethylene glycol, 1, 2-propylene glycol or 1, 3-propylene glycol, sodium chloride, sodium benzoate, potassium sorbate, sodium sulfate, potassium sulfate, magnesium sulfate, sodium thiosulfate, calcium carbonate, sodium citrate, dextrin, maltodextrin, glucose, sucrose, sorbitol, lactose, wheat flour, wheat bran, corn gluten meal, starch, cellulose farigel, cassava cores, sodium aluminium silicate, colloidal amorphous silica, Sipernat 50S, polyethylene glycol 200, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 1000, polyethylene glycol 1500, polyethylene glycol 4000 and carbopol.

In a preferred embodiment of Aspect 5, 6, 7, or 8 the composition further comprises calcium carbonate and sodium aluminium silicate.

In a preferred embodiment of Aspect 5, 6, 7, or 8 the composition further comprises calcium carbonate, sodium aluminium silicate and sucrose.

In another preferred embodiment of Aspect 5, 6, 7, or 8 the composition further comprises one or more carriers such as one or more carriers selected from the group consisting of Calcium carbonate, sodium sulfate, starch, farigel and cassava cores.

In another preferred embodiment of Aspect 5, 6, 7, or 8 the composition further comprises one or more flowability agents such as sodium aluminium silicate and/or colloidal amorphous silica (e.g., Sipernat 50S).

In another preferred embodiment of Aspect 5, 6, 7, or 8 the composition further comprises one or more binder such as one or more binders selected from the group consisting of sucrose, sorbitol, glycerol, polyethylene glycol 200, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 1000, polyethylene glycol 1500, polyethylene glycol 4000, dextrin, maltodextrin and carbopol.

In a preferred embodiment the composition comprises *Bacillus* DSM 29869, calcium carbonate and sodium aluminium silicate.

In a preferred embodiment the composition comprises *Bacillus* DSM 29869, calcium carbonate, sodium aluminium silicate and sucrose.

In a preferred embodiment the composition comprises *Bacillus* DSM 29869 and one or more carriers such as one or more carriers selected from the group consisting of calcium carbonate, sodium sulphate, starch, farigel and cassava cores.

In a preferred embodiment the composition comprises *Bacillus* DSM 29869 and one or more flowability agents such as sodium aluminium silicate and/or colloidal amorphous silica (e.g., Sipernat 50S).

In a preferred embodiment the composition comprises *Bacillus* DSM 29869 and one or more binder such as one or more binders selected from the group consisting of sucrose, sorbitol, glycerol, polyethylene glycol 200, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 1000, polyethylene glycol 1500, polyethylene glycol 4000, dextrin, maltodextrin and carbopol.

In a preferred embodiment the composition comprises *Bacillus* DSM 29870, calcium carbonate and sodium aluminium silicate.

In a preferred embodiment the composition comprises *Bacillus* DSM 29870, calcium carbonate, sodium aluminium silicate and sucrose.

In a preferred embodiment the composition comprises *Bacillus* DSM 29870 and one or more carriers such as one or more carriers selected from the group consisting of calcium carbonate, sodium sulphate, starch, farigel and cassava cores.

In a preferred embodiment the composition comprises *Bacillus* DSM 29870 and one or more flowability agents such as sodium aluminium silicate and/or colloidal amorphous silica (e.g., Sipernat 50S).

In a preferred embodiment the composition comprises *Bacillus* DSM 29870 and one or more binder such as one or more binders selected from the group consisting of sucrose, sorbitol, glycerol, polyethylene glycol 200, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 1000, polyethylene glycol 1500, polyethylene glycol 4000, dextrin, maltodextrin and carbopol.

In a preferred embodiment the composition comprises *Bacillus* DSM 29871, calcium carbonate and sodium aluminium silicate.

In a preferred embodiment the composition comprises *Bacillus* DSM 29871, calcium carbonate, sodium aluminium silicate and sucrose.

In a preferred embodiment the composition comprises *Bacillus* DSM 29871 and one or more carriers such as one or more carriers selected from the group consisting of calcium carbonate, sodium sulphate, starch, farigel and cassava cores.

In a preferred embodiment the composition comprises *Bacillus* DSM 29871 and one or more flowability agents such as sodium aluminium silicate and/or colloidal amorphous silica (e.g., Sipernat 50S).

In a preferred embodiment the composition comprises *Bacillus* DSM 29871 and one or more binder such as one or more binders selected from the group consisting of sucrose, sorbitol, glycerol, polyethylene glycol 200, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 1000, polyethylene glycol 1500, polyethylene glycol 4000, dextrin, maltodextrin and carbopol.

In a preferred embodiment the composition comprises *Bacillus* DSM 29872, calcium carbonate and sodium aluminium silicate.

In a preferred embodiment the composition comprises *Bacillus* DSM 29872, calcium carbonate, sodium aluminium silicate and sucrose.

In a preferred embodiment the composition comprises *Bacillus* DSM 29872 and one or more carriers such as one or more carriers selected from the group consisting of calcium carbonate, sodium sulphate, starch, farigel and cassava cores.

In a preferred embodiment the composition comprises *Bacillus* DSM 29872 and one or more flowability agents such as sodium aluminium silicate and/or colloidal amorphous silica (e.g., Sipernat 50S).

In a preferred embodiment the composition comprises *Bacillus* DSM 29872 and one or more binder such as one or more binders selected from the group consisting of sucrose, sorbitol, glycerol, polyethylene glycol 200, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 1000, polyethylene glycol 1500, polyethylene glycol 4000, dextrin, maltodextrin and carbopol.

In one embodiment the composition comprises one or more coccidiostats wherein the composition is, e.g., a premix.

In a preferred embodiment the composition according to Aspect 5, 6, 7, or 8 the composition comprises from 10^5 to 10^{18} CFU/g of isolated *Bacillus* spores.

In a further embodiment, the composition according to the invention comprises one or more bacterial strains such as at least two of the above strains up to and including all of the strains in the group consisting of DSM 29869, DSM 29870, DSM 29871 and DSM 29872.

In an embodiment to any of the aforementioned embodiments the *Bacillus* spore kills/inhibits at least 40% (such as at least 45%, at least 50%, at least 60%, at least 70% or at least 80%) of *Clostridium perfringens* after, e.g., 24 hours, e.g., determined as described in Example 2.

In another embodiment of the invention the composition further comprises one or more additional microbes. In another embodiment of the invention the composition further comprises one or more enzymes. In another embodiment of the invention the composition further comprises one or more vitamins. In another embodiment of the invention the composition further comprises one or more minerals. In another embodiment of the invention the composition further comprises one or more amino acids. In another embodiment of the invention the composition further comprises one or more other feed ingredients.

In an embodiment to any of the aforementioned embodiments, the composition also improves the health of the mono-gastric animal when fed to said animal. In another embodiment to any of the aforementioned embodiments, the composition also increases the egg yield of poultry when fed said poultry. In an embodiment to any of the aforementioned embodiments, the composition increases the meat yield of the mono-gastric animal when fed to said animal.

In a preferred embodiment, the composition comprises one or more bacterial strains described herein, wherein the bacterial count of each of the bacterial strains is between 1×10^4 and 1×10^{18} CFU/kg of composition, preferably between 1×10^7 and 1×10^{16} CFU/kg of composition, more preferably between 1×10^{10} and 1×10^{15} CFU/kg of composition and most preferably between 1×10^{11} and 1×10^{14} CFU/kg of composition.

In a preferred embodiment, the bacterial count of each of the bacterial strains in the composition is between 1×10^4 and 1×10^{18} CFU/kg of dry matter, preferably between 1×10^7 and 1×10^{16} CFU/kg of dry matter, more preferably between 1×10^{10} and 1×10^{15} CFU/kg of dry matter and most preferably between 1×10^{11} and 1×10^{14} CFU/kg of dry matter. In a more preferred embodiment the bacterial count of each of the bacterial strains in the animal feed is between 1×10^8 and 1×10^{10} CFU/kg of dry matter.

In a preferred embodiment, the composition has a bacterial count of each *Bacillus* spore between 1×10^3 and 1×10^{13} CFU/animal/day, preferably between 1×10^5 and 1×10^{11} CFU/animal/day, more preferably between 1×10^6 and 1×10^{10} CFU/animal/day and most preferably between 1×10^7 and 1×10^9 CFU/animal/day.

In still yet another embodiment of the invention, the one or more bacterial strains are present in the composition in form of a spore such as a stable spore. In still a further embodiment of the invention, the stable spore will germinate in the intestine and/or stomach of the mono-gastric animal.

In one embodiment, the one or more bacterial strains are stable when subjected to pressures applied/achieved during an extrusion process for pelleting. In a particular embodiment, the one or more bacterial strains are stable at pressures ranging from 1 bar to 40 bar, particularly 10 bar to 40 bar, more particularly 15 bar to 40 bar, even more particularly 20 bar to 40 bar, still even more particularly 35 bar to 37 bar, even still more particularly 36 bar.

In a particular embodiment, the one or more bacterial strains are stable at high temperatures. In particular, the bacterial strains are stable when they are subjected to temperatures achieved during an extrusion process for pelleting. In an even more particular embodiment, the one or more bacterial strains are stable at temperatures ranging from 80°C to 120°C, particularly temperatures ranging from 90°C to 120°C, even more particularly temperatures ranging from 95°C to 120°C.

In another aspect, the invention relates to a composition comprising a carrier and one or more of the bacteria cultures having characteristics substantially identical to that of a strain selected from the group consisting of:

- the strain having the deposit accession number DSM 29869;
- the strain having the deposit accession number DSM 29870;
- the strain having the deposit accession number DSM 29871; and

the strain having the deposit accession number DSM 29872; or
any combination thereof.

In an embodiment, the composition comprises a carrier and the strain having the deposit accession number DSM 29869, or a strain having all of the identifying characteristics
5 of *Bacillus* DSM 29869 or a mutant thereof.

In an embodiment, the composition comprises a carrier and the strain having the deposit accession number DSM 29870, or a strain having all of the identifying characteristics of *Bacillus* DSM 29870 or a mutant thereof.

In an embodiment, the composition comprises a carrier and the strain having the
10 deposit accession number DSM 29871, or a strain having all of the identifying characteristics of *Bacillus* DSM 29871 or a mutant thereof.

In an embodiment, the composition comprises a carrier and the strain having the deposit accession number DSM 29872, or a strain having all of the identifying characteristics of *Bacillus* DSM 29872 or a mutant thereof.

15 In an embodiment, the composition further comprises one or more additional microbes. In a particular embodiment, the composition further comprises a bacterium from one or more of the following genera: *Lactobacillus*, *Lactococcus*, *Streptococcus*, *Bacillus*, *Pediococcus*, *Enterococcus*, *Leuconostoc*, *Carnobacterium*, *Propionibacterium*, *Bifidobacterium*, *Clostridium* and *Megasphaera* or any combination thereof.

20 In a particular embodiment, the composition further comprises a bacterium from one or more of the following strains of *Bacillus amyloliquefaciens*, *Bacillus subtilis*, *Bacillus pumilus*, *Bacillus polymyxa*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus coagulans*, *Bacillus circulans*, or any combination thereof.

In a particular embodiment, the composition further comprises one or more types of
25 yeast. The one or more types of yeast can be selected from the group consisting of *Saccharomycetaceae*, *Saccharomyces* (such as *S. cerevisiae* and/or *S. boulardii*), *Kluyveromyces* (such as *K. marxianus* and *K. lactis*), *Candida* (such as *C. utilis*, also called *Torula* yeast), *Pichia* (such as *P. pastoris*), *Torulaspora* (such as *T. delbrueckii*), *Phaffia* yeasts and *Basidiomycota*.

30 In an embodiment to any of the aforementioned embodiments the composition further comprises a carrier. The carrier can comprise one or more of the following compounds: water, glycerol, ethylene glycol, 1, 2-propylene glycol or 1, 3-propylene glycol, sodium chloride, sodium benzoate, potassium sorbate, sodium sulfate, potassium sulfate, magnesium sulfate, sodium thiosulfate, calcium carbonate, sodium citrate, dextrin,
35 maltodextrin, glucose, sucrose, sorbitol, lactose, wheat flour, wheat bran, corn gluten meal, starch, cellulose, fargel, cassava cores, sodium aluminium silicate, colloidal amorphous silica, Sipernat 50S, polyethylene glycol 200, polyethylene glycol 400, polyethylene glycol

600, polyethylene glycol 1000, polyethylene glycol 1500, polyethylene glycol 4000 and carbopol.

In another embodiment, the composition described herein can optionally include one or more enzymes. Enzymes can be classified on the basis of the handbook Enzyme
5 Nomenclature from NC-IUBMB, 1992), see also the ENZYME site at the internet: <http://www.expasy.ch/enzyme/>. ENZYME is a repository of information relative to the nomenclature of enzymes. It is primarily based on the recommendations of the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology (IUB-MB), Academic Press, Inc., 1992, and it describes each type of characterized enzyme for which an
10 EC (Enzyme Commission) number has been provided (Bairoch, The ENZYME database, 2000, *Nucleic Acids Res.* 28:304-305). This IUB-MB Enzyme nomenclature is based on their substrate specificity and occasionally on their molecular mechanism; such a classification does not reflect the structural features of these enzymes.

Another classification of certain glycoside hydrolase enzymes, such as
15 endoglucanase, xylanase, galactanase, mannanase, dextranase and alpha-galactosidase, in families based on amino acid sequence similarities has been proposed a few years ago. They currently fall into 90 different families: See the CAZy(ModO) internet site (Coutinho, P.M. & Henrissat, B. (1999) Carbohydrate-Active Enzymes server at URL: <http://afmb.cnrs-mrs.fr/~cazy/CAZY/index.html> (corresponding papers: Coutinho, P.M. & Henrissat, B. (1999)
20 Carbohydrate-active enzymes: an integrated database approach. In "Recent Advances in Carbohydrate Bioengineering", H.J. Gilbert, G. Davies, B. Henrissat and B. Svensson eds., The Royal Society of Chemistry, Cambridge, pp. 3-12; Coutinho, P.M. & Henrissat, B. (1999) The modular structure of cellulases and other carbohydrate-active enzymes: an integrated database approach. In "Genetics, Biochemistry and Ecology of Cellulose Degradation", K.
25 Ohmiya, K. Hayashi, K. Sakka, Y. Kobayashi, S. Karita and T. Kimura eds., Uni Publishers Co., Tokyo, pp. 15-23).

Thus the composition of the invention may also comprise at least one other enzyme selected from the group comprising of phytase (EC 3.1.3.8 or 3.1.3.26); xylanase (EC 3.2.1.8); galactanase (EC 3.2.1.89); alpha-galactosidase (EC 3.2.1.22); protease (EC 3.4);
30 phospholipase A1 (EC 3.1.1.32); phospholipase A2 (EC 3.1.1.4); lysophospholipase (EC 3.1.1.5); phospholipase C (3.1.4.3); phospholipase D (EC 3.1.4.4); amylase such as, for example, alpha-amylase (EC 3.2.1.1); lysozyme (EC 3.2.1.17); and beta-glucanase (EC 3.2.1.4 or EC 3.2.1.6), or any mixture thereof.

In a particular embodiment, the composition of the invention comprises a phytase (EC
35 3.1.3.8 or 3.1.3.26). Examples of commercially available phytases include Bio-FeedTM Phytase (Novozymes), Ronozyme® P and HiPhosTM (DSM Nutritional Products), NatuphosTM (BASF), Finase® and Quantum® Blue (AB Enzymes), the Phyzyme® XP (Verenium/DuPont)

and Axtra® PHY (DuPont). Other preferred phytases include those described in, e.g., WO 98/28408, WO 00/43503, and WO 03/066847.

In a particular embodiment, the composition of the invention comprises a xylanase (EC 3.2.1.8). Examples of commercially available xylanases include Ronozyme® WX and G2 (DSM Nutritional Products), Econase® XT and Barley (AB Vista), Xylathin® (Verenium) and Axtra® XB (Xylanase/beta-glucanase, DuPont)

In a particular embodiment, the composition of the invention comprises a protease (EC 3.4). Examples of commercially available proteases include Ronozyme® ProAct (DSM Nutritional Products).

Manufacturing

The composition of the invention can, e.g., be manufactured as mash composition (non-pelleted) or pelleted composition. The bacteria cultures and optionally enzymes can be added as solid or liquid formulations. For example, for mash composition a solid or liquid culture formulation may be added before or during the ingredient mixing step. Typically a liquid culture preparation comprises the culture of the invention optionally with a polyol, such as glycerol, ethylene glycol or propylene glycol, and is added after the pelleting step, such as by spraying the liquid formulation onto the pellets.

The enzyme may be added to the composition as a granule, which is optionally pelleted or extruded. The granule typically comprises a core particle and one or more coatings, which typically are salt and/or wax coatings. The core particle can either be a homogeneous blend of an active compound optionally together with salts (e.g., organic or inorganic zinc or calcium salt) or an inert particle with an active compound applied onto it. The active compound is the culture of the invention optionally combined with one or more enzymes. The inert particle may be water soluble or water insoluble, e.g., starch, a sugar (such as sucrose or lactose), or a salt (such as NaCl, Na₂SO₄). The salt coating is typically at least 1 µm thick and can either be one particular salt or a mixture of salts, such as Na₂SO₄, K₂SO₄, MgSO₄ and/or sodium citrate. Other examples are those described in, e.g., WO 2008/017659, WO 2006/034710, WO 97/05245, WO 98/54980, WO 98/55599, WO 00/70034 or polymer coating such as described in WO 01/00042.

Preferred embodiments

Preferred embodiments of the invention are described in the two set of items herein below.

ITEM SET I:

1. A *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
- ii) the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens*

5 and

- iii) the *Bacillus* strain improves body weight gain and/or feed conversion rate in chickens fed with the *Bacillus* strain.

2. A *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
- ii) the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens*

10 and

- iii) the *Bacillus* strain is sensitive to at least seven such as eight of the antibiotics selected from the group consisting of Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline.

15

3. A *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
- ii) the *Bacillus* strain is sensitive to at least seven such as eight of the antibiotics

20 selected from the group consisting of Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline and

- iii) the *Bacillus* strain improves body weight gain and/or feed conversion rate in chickens fed with the *Bacillus* strain.

25 4. A *Bacillus* strain characterized in that:

- i) the *Bacillus* strain is non-hemolytic,
- ii) the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens*

and optionally

- iii) the *Bacillus* strain has enzymatic activity under aerobic and/or anaerobic conditions that hydrolyzes one or more of the substrates selected from the group consisting of Amylose, Arabinan, Arabinoxylan, Casein and Xylan.

30

5. The *Bacillus* strain according to any of items 1 to 3, wherein the *Bacillus* strain has enzymatic activity under aerobic and/or anaerobic conditions that hydrolyzes one or more of the substrates selected from the group consisting of Amylose, Arabinan, Arabinoxylan, Casein and Xylan (e.g., determined as described in Example 6).

35

6. The *Bacillus* strain according to any of items 2 to 4, wherein the *Bacillus* strain improves one or more performance parameters in poultry selected from the list consisting of body weight gain, European Production Efficacy Factor and feed conversion rate in poultry fed with the *Bacillus* strain.

5

7. The *Bacillus* strain according to item 3, wherein the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens* (e.g., determined as described in Example 2).

8. The *Bacillus* strain according to any of items 1 and 4, wherein the *Bacillus* strain is sensitive to at least seven such as eight of the antibiotics selected from the group consisting of Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline (e.g., determined as described in Example 5).

9. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain comprises 16S rDNA that is more than 98% sequence identity to SEQ ID NO: 1 and/or more than 98% sequence identity to SEQ ID NO: 2 and/or more than 98% sequence identity to SEQ ID NO: 3 and/or more than 98% sequence identity to SEQ ID NO: 4.

10. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain is a *Bacillus subtilis* strain or a *Bacillus amyloliquefaciens* strain.

11. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain has antimicrobial effect against *E. coli* (e.g., determined as described in Example 4).

12. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain is the *Bacillus* strain having deposit accession number DSM 29869, or a strain having all of the identifying characteristics of *Bacillus* DSM 29869 or a mutant thereof that has antimicrobial activity against *Clostridium perfringens*.

13. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain is the *Bacillus* strain having deposit accession number DSM 29870, or a strain having all of the identifying characteristics of *Bacillus* DSM 29870 or a mutant thereof that has antimicrobial activity against *Clostridium perfringens*.

14. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain is the *Bacillus* strain having deposit accession number DSM 29871, or a strain having all of the

identifying characteristics of *Bacillus* DSM 29871 or a mutant thereof that has antimicrobial activity against *Clostridium perfringens*.

15. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain is the *Bacillus* strain having deposit accession number DSM 29872, or a strain having all of the identifying characteristics of *Bacillus* DSM 29872 or a mutant thereof that has antimicrobial activity against *Clostridium perfringens*.

16. A *Bacillus* having deposit accession number DSM 29869 or a strain having all of the identifying characteristics of *Bacillus* DSM 29869 or a mutant thereof.

17. A *Bacillus* having deposit accession number DSM 29870 or a strain having all of the identifying characteristics of *Bacillus* DSM 29870 or a mutant thereof.

18. A *Bacillus* having deposit accession number DSM 29871 or a strain having all of the identifying characteristics of *Bacillus* DSM 29871 or a mutant thereof.

19. A *Bacillus* having deposit accession number DSM 29872 or a strain having all of the identifying characteristics of *Bacillus* DSM 29872 or a mutant thereof.

20. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain has a high compatibility with monensin such as being compatible with at least 2.3 µg/ml monensin as determined in Example 12 (such as at least 2.4 µg/ml monensin, such as at least 2.5 µg/ml monensin, such as at least 2.6 µg/ml monensin or such as at least 2.7 µg/ml monensin as determined in Example 12).

21. A composition comprising spores of the *Bacillus* strain according to any of items 1-20.

22. The composition according to item 21, wherein the *Bacillus* spores of the composition are present as dried spores.

23. The composition according to any of items 21 to 22 which further comprises a carrier.

24. The composition according to item 23, wherein the carrier comprises one or more of the following compounds: water, glycerol, ethylene glycol, 1, 2-propylene glycol or 1, 3-propylene glycol, sodium aluminium silicate, sodium chloride, sodium benzoate, potassium sorbate, sodium sulfate, potassium sulfate, magnesium sulfate, sodium thiosulfate, calcium

carbonate, sodium citrate, dextrin, maltodextrin, glucose, sucrose, sorbitol, lactose, wheat flour, wheat bran, corn gluten meal, starch, farigel, cassava cores, sodium aluminium silicate, colloidal amorphous silica, Sipernat 50S, polyethylene glycol 200, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 1000, polyethylene glycol 1500, polyethylene glycol 4000, carbopol and cellulose.

25. The composition according to any of items 21 to 24, wherein the composition comprises from 10^5 to 10^{18} CFU/g of isolated *Bacillus* spores.

26. The composition according to any of items 21 to 25 which further comprises one or more components selected from the list consisting of:

- one or more enzymes;
- one or more additional microbes;
- one or more vitamins;
- one or more minerals;
- one or more amino acids; and
- one or more other feed ingredients.

27. The composition according to any of items 21 to 26, wherein the bacterial count of each *Bacillus* spore is between 1×10^4 and 1×10^{18} CFU/kg of composition, preferably between 1×10^7 and 1×10^{16} CFU/kg of composition, more preferably between 1×10^{10} and 1×10^{15} CFU/kg of composition and most preferably between 1×10^{11} and 1×10^{14} CFU/kg of composition.

28. The composition according to any of items 21 to 26, wherein the bacterial count of each *Bacillus* spore is between 1×10^4 and 1×10^{18} CFU/kg of composition, preferably between 1×10^6 and 1×10^{15} CFU/kg of composition, and more preferably between 1×10^7 and 1×10^{11} CFU/kg of composition.

29. A biologically pure culture of the *Bacillus* strain according to any of items 1 to 20.

30. An isolated *Bacillus* strain according to any of items 1 to 20.

31. An isolated *Bacillus* strain selected from the group consisting of *Bacillus* strain DSM 29869, DSM 29870, DSM 29871 and DSM 29872, or a strain having all of the identifying characteristics of *Bacillus* DSM 29869, DSM 29870, DSM 29871 and DSM 29872 or a mutant thereof.

32. The isolated *Bacillus* strain according to item 31, wherein the identifying characteristics can one or more (such as all) of the characteristics selected from the group consisting of

i) non-hemolytic, *e.g.*, as determined in Example 1,

ii) antimicrobial activity against *Clostridium perfringens*, *e.g.*, as determined in Example 2,

iii) antimicrobial activity against *E. coli*, *e.g.*, as determined in Example 4,

iv) sensitive to Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracyclin, and

v) high compatibility with monensin such as being compatible with at least 2.3 µg/ml monensin as determined in Example 12 (such as at least 2.4 µg/ml monensin, such as at least 2.5 µg/ml monensin, such as at least 2.6 µg/ml monensin or such as at least 2.7 µg/ml monensin as determined in Example 12).

ITEM SET II:

1. A *Bacillus* strain characterized in:

i) having deposit accession number DSM 29870 or a strain having all of the identifying characteristics of *Bacillus* DSM 29870 or a mutant thereof;

ii) having deposit accession number DSM 29869 or a strain having all of the identifying characteristics of *Bacillus* DSM 29869 or a mutant thereof;

iii) having deposit accession number DSM 29871 or a strain having all of the identifying characteristics of *Bacillus* DSM 29871 or a mutant thereof; or

iv) having deposit accession number DSM 29872 or a strain having all of the identifying characteristics of *Bacillus* DSM 29872 or a mutant thereof.

2. The *Bacillus* strain according to item 1, wherein the *Bacillus* strain is non-hemolytic.

3. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens* (*e.g.*, determined as described in Example 2).

4. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain improves body weight gain and/or feed conversion rate in chickens fed with the *Bacillus* strain.

5. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain comprises 16S rDNA that is more than 98% sequence identity to SEQ ID NO: 1 and/or more

than 98% sequence identity to SEQ ID NO: 2 and/or more than 98% sequence identity to SEQ ID NO: 3 and/or more than 98% sequence identity to SEQ ID NO: 4.

5 6. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain comprises 16S rDNA that is more than 99% sequence identity to SEQ ID NO: 1 and/or more than 99% sequence identity to SEQ ID NO: 2 and/or more than 99% sequence identity to SEQ ID NO: 3 and/or more than 99% sequence identity to SEQ ID NO: 4.

10 7. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain is a *Bacillus subtilis* strain or a *Bacillus amyloliquefaciens* strain.

8. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain has antimicrobial effect against *E. coli* (e.g., determined as described in Example 4).

15 9. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain is an isolated *Bacillus* strain.

10. A biologically pure culture of the *Bacillus* strain according to any of items 1 to 9.

20 11. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain is sensitive to at least seven such as eight of the antibiotics selected from the group consisting of Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin, Streptomycin, Erythromycin and Tetracycline (e.g., determined as described in Example 5).

25 12. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain has enzymatic activity under aerobic and/or anaerobic conditions that hydrolyzes one or more of the substrates selected from the group consisting of Amylose, Arabinan, Arabinoxylan, Casein and Xylan (e.g., determined as described in Example 6).

30 13. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain improves one or more performance parameters in poultry selected from the list consisting of body weight gain, European Production Efficacy Factor and feed conversion rate in poultry fed with the *Bacillus* strain.

35 14. The *Bacillus* strain according to any of the previous items, wherein the *Bacillus* strain has a high compatibility with monensin such as being compatible with at least 2.3 µg/ml monensin as determined in Example 12 (such as at least 2.4 µg/ml monensin, such as at

least 2.5 µg/ml monensin, such as at least 2.6 µg/ml monensin or such as at least 2.7 µg/ml monensin as determined in Example 12).

15. A composition comprising spores of the *Bacillus* strain according to any of items 1-14.

5

16. The composition according to item 15, wherein the *Bacillus* spores of the composition are present as spores such as dried spores such as spray dried spores.

17. The composition according to item 16, wherein the bacterial count of each *Bacillus* spore is between 1×10^4 and 1×10^{18} CFU/kg of composition, preferably between 1×10^6 and 1×10^{15} CFU/kg of composition, and more preferably between 1×10^7 and 1×10^{11} CFU/kg of composition.

10

18. The composition according to any of items 15 to 17 which further comprises a carrier.

15

19. The composition according to item 18, wherein the carrier comprises one or more of the following compounds: water, glycerol, ethylene glycol, 1, 2-propylene glycol or 1, 3-propylene glycol, sodium aluminium silicate, sodium chloride, sodium benzoate, potassium sorbate, sodium sulfate, potassium sulfate, magnesium sulfate, sodium thiosulfate, calcium carbonate, sodium citrate, dextrin, maltodextrin, glucose, sucrose, sorbitol, lactose, wheat flour, wheat bran, corn gluten meal, starch, farigel, cassava cores, sodium aluminium silicate, colloidal amorphous silica, Sipernat 50S, polyethylene glycol 200, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 1000, polyethylene glycol 1500, polyethylene glycol 4000, carbopol. and cellulose.

20

25

20. The composition according to item 18 or 19, wherein the carrier comprises calcium carbonate.

21. The composition according to any of items 15 to 20, wherein the composition comprises from 10^5 to 10^{18} CFU/g of isolated *Bacillus* spores.

30

22. The composition according to any of items 15 to 21 which further comprises one or more components selected from the list consisting of:

one or more enzymes;

35

one or more additional microbes;

one or more vitamins;

one or more minerals;

one or more amino acids; and
one or more other feed ingredients.

23. The composition according to any of items 15 to 22, wherein the bacterial count of each
5 *Bacillus* spore is between 1×10^4 and 1×10^{18} CFU/kg of composition, preferably between
 1×10^7 and 1×10^{16} CFU/kg of composition, more preferably between 1×10^{10} and 1×10^{15}
CFU/kg of composition and most preferably between 1×10^{11} and 1×10^{14} CFU/kg of
composition.

10 24. The composition according to any of items 15 to 23, wherein the bacterial count of each
Bacillus spore is between 1×10^4 and 1×10^{18} CFU/kg of composition, preferably between
 1×10^6 and 1×10^{15} CFU/kg of composition, and more preferably between 1×10^7 and 1×10^{11}
CFU/kg of composition.

15 25. An isolated *Bacillus* strain selected from the group consisting of *Bacillus* strain DSM
29869, DSM 29870, DSM 29871 and DSM 29872, or a strain having all of the identifying
characteristics of *Bacillus* DSM 29869, DSM 29870, DSM 29871 and DSM 29872 or a
mutant thereof.

20 26. The isolated *Bacillus* strain according to item 25, wherein the identifying characteristics
can one or more (such as all) of the characteristics selected from the group consisting of

- i) non-hemolytic, e.g., as determined in Example 1,
- ii) antimicrobial activity against *Clostridium perfringens*, e.g., as determined in
Example 2,
- 25 iii) antimicrobial activity against *E. coli*, e.g., as determined in Example 4,
- iv) sensitive to Vancomycin, Clindamycin, Chloramphenicol, Gentamicin, Kanamycin,
Streptomycin, Erythromycin and Tetracyclin, and
- v) high compatibility with monensin such as being compatible with at least 2.3 µg/ml
monensin as determined in Example 12 (such as at least 2.4 µg/ml monensin, such as at
30 least 2.5 µg/ml monensin, such as at least 2.6 µg/ml monensin or such as at least 2.7 µg/ml
monensin as determined in Example 12).

27. The isolated *Bacillus* strain according to item 25 or 26 for use of treatment of necrotic
enteritis and/or a *clostridium perfringens* infection.

35 28. An isolated *Bacillus* strain selected from the group consisting of *Bacillus* strain DSM
29869, DSM 29871 and DSM 29872, or a strain having all of the identifying characteristics of

Bacillus DSM 29869, DSM 29871 and DSM 29872 or a mutant thereof for treatment of necrotic enteritis and/or a *clostridium perfringens* infection.

29. Use of the *Bacillus* strain or the composition according to any of the previous items to improve one or more performance parameters of an animal selected from the group consisting of: improving the feed conversion ratio, improving the body weight gain, improving the European Production Efficacy Factor, improving the feed efficiency and improving the health, e.g., of poultry such as chickens.

30. The use according to item 29, wherein the *Bacillus* strain is selected from the group consisting of *Bacillus* strain DSM 29869, DSM 29871 and DSM 29872, or a strain having all of the identifying characteristics of *Bacillus* DSM 29869, DSM 29871 and DSM 29872 or a mutant thereof.

31. A method of improving one or more performance parameters of an animal selected from the group consisting of: improving the feed conversion ratio, improving the body weight gain, improving the European Production Efficacy Factor, improving the feed efficiency and improving the health, comprising administering the *Bacillus* strain or the composition according to any of the previous items to the animal.

32. The method according to item 31, wherein the *Bacillus* strain is selected from the group consisting of *Bacillus* strain DSM 29869, DSM 29871 and DSM 29872, or a strain having all of the identifying characteristics of *Bacillus* DSM 29869, DSM 29871 and DSM 29872 or a mutant thereof.

33. A method for feeding an animal comprising administering the *Bacillus* strain or the composition according to any of the previous items to said animal optionally in conjunction with other animal feed ingredients.

34. The method for feeding an animal according to item 33 wherein the animal is selected from the group consisting of poultry and swine.

EXAMPLES

Example 1: Screening for Hemolysis negative *Bacillus* strains

Hemolysis was tested according to the Technical Guidance on the assessment of the toxigenic potential of *Bacillus* species used in animal nutrition, EFSA Journal 2011; 9(11):2445.

Sheep blood agar plates were purchased as ready to use (Becton Dickinson art 254053 or 254087). Alternatively, the agar plates can be prepared by adding 5% defibrinated sheep blood (obtained from Statens Serum Institute, Denmark) to TS-agar (Oxoid CM 131). Agar should be autoclaved at 121°C for 20 minutes and cooled down to about 40°C before adding the blood immediately before pouring the plates.

The *Bacillus* strains were taken from the preservation at -80°C and streaked on TS-agar plate, which was incubated at 30°C overnight or until growth appeared. From a single colony as little as possible of the material was used to streak a line on ¼ of an agar plate. The plate was incubated at 30°C for 72 hours. Hemolysis/clearing zones of the *Bacillus* strains to be tested were compared with the positive and negative control. As positive control *Bacillus subtilis* ATCC21332 was used. As negative control *Bacillus subtilis* 168, (BGSC-1A1, Bacillus Genetic Stock Center) was used.

In a screening of 599 independent isolates of *Bacillus* 223 strains (37%) were hemolysis negative. In another screening 21 of 65 independent isolates of *Bacillus* (32%) were hemolysis negative. Both screenings exclusively comprised strains that based on identification by 16S rDNA sequencing belong to *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus pumilus* or *Bacillus amyloliquefaciens* or close relatives to these four species.

The non-Hemolytic *Bacillus* strains therefore seem to be common and fairly abundant in nature, but only comprising a minority of the natural strains. The non-hemolytic strains seem to be more abundant in *Bacillus licheniformis*, while most *Bacillus amyloliquefaciens* strains appear hemolytic. All the non-hemolytic strains were tested for inhibition of growth of *Clostridium perfringens*. The following strains from screening were all hemolysis negative and among those selected for further studies: DSM 29869, DSM 29870, DSM 29871 and DSM 29872. Clostat (*Bacillus* PB6) was determined to be hemolytic.

Example 2: Determination of inhibition of growth of *Clostridium perfringens*

All the non-hemolytic strains were tested for inhibition of growth of *Clostridium perfringens* strains 23 and 48 (both are netB positive) [Gholamiandekhordi *et al.*, 2006, Molecular and phenotypical characterization of *Clostridium perfringens* isolates from poultry flocks with different diseasestatus, *Vet. Microbiol.* 113:143-152], were grown overnight in tryptic soy broth (BD part 211822) supplemented with 0.6% yeast extract (BD part 212750) at 35°C under static anaerobic conditions. 250 µL of the overnight culture of *Clostridium perfringens* was added to 250 mL of tryptic soy agar supplemented with 0.6% yeast extract at 40°C and poured into rectangular petri plates (Nunc part 267060). The inoculated agar was allowed to cool at room temperature after which an 8 mm diameter well was made in the agar. Plates were stored in absence of oxygen until use.

The *Bacillus* strain DSM 29869, DSM 29870, DSM 29871, or DSM 29872 was grown overnight in tryptic soy broth at 35°C under aerobic conditions. 1000 µL of *Bacillus* culture was collected and fractionated into cell-free supernatant and cells by centrifugation. 20 µL of cell-free supernatant or 100x diluted cells in phosphate buffered saline were added directly to the wells in the *Clostridium perfringens* inoculated agar plates. A control well contained 20 µL of phosphate buffer saline. The plates were incubated for 18 hours at 35°C under anaerobic conditions.

Inhibition of the *Clostridium perfringens* strain was noted by a circular clearing zone around the well of interest. The phosphate buffer saline well was considered a negative control based on lack of clearing zone around the well.

Cell-free supernatant and 100x diluted cells of *Bacillus* strains DSM 29869, DSM 29870, DSM 29871, and DSM 29872 were able to consistently inhibit growth of *C. perfringens* strains 23 and 48 *in vitro*. Inhibition was also seen by competitor strain “CloSTAT”, for both supernatant and cells, but was not seen with competitor strain GalliproTect. “CloSTAT” is a strain of *Bacillus amyloliquefaciens* that was isolated from the commercial DFM product CloSTAT, Kemin. “GalliproTect” is a strain of *Bacillus licheniformis* that was isolated from the commercial product Gallipro Tect, Chr Hansen.

Example 3: Identification, characterization and deposit of the biological material

The following biological materials were deposited under the terms of the Budapest Treaty at Leibniz-Institut DSMZ-Deutsche Sammlung von Mikro-organismen und Zellkulturen GmbH, Inhoffenstraße 7 B, 38124 Braunschweig Germany, and given the following accession numbers:

Table 3.1: Deposit of Biological Material

Identification	Accession Number	Date of Deposit
<i>Bacillus amyloliquefaciens</i>	DSM 29869	Jan 12 2015
<i>Bacillus subtilis</i>	DSM 29870	Jan 12 2015
<i>Bacillus subtilis</i>	DSM 29871	Jan 12 2015
<i>Bacillus amyloliquefaciens</i>	DSM 29872	Jan 12 2015

The strains have been deposited under conditions that assure that access to the culture will be available during the pendency of this patent application to one determined by foreign patent laws to be entitled thereto. The deposits represent a substantially pure culture of the deposited strain. The deposits are available as required by foreign patent laws in

countries wherein counterparts of the subject application or its progeny are filed. However, it should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by governmental action.

5 Sequencing of 16S rDNA genes

DNA was extracted from a culture of DSM 29869, DSM 29870, DSM 29871 and DSM 29872 using QiaAmp DNA Blood Mini Kit (Qiagen art 51106). The kit was used as recommended for extraction of DNA from gram positive bacteria.

16S rDNA was amplified in a total volume of 50 µl by mixing: 10 pmol of each of
 10 Primer 16S F and 16S R, 0.2 mM of each nucleotide, 2.5 units Ampli taq, 1 x Ampli taq buffer, 5 µl DNA template and by using the following PCR program: 94°C 2 min (94°C 30 s, 52°C 30 S, 72°C 1 min)x35, 72°C 10 min on a Perkin Elmer PCR machine. The PCR product was sequenced by Novozymes DNA sequencing facility using primer 530R, 357F, 1390R and 1100F.

15 Table 3.2: Primers:

Primer	Sequence	SEQ ID NO
16S-F	5'-GAGTTTGATCCTGGCTCAG-3'	SEQ ID NO: 8
16S-R	5'-AGAAAGGAGGTGATCCAGCC-3'	SEQ ID NO: 9
794-R	5'-ATCTAATCCTGTTTGCTCCCC-3'	SEQ ID NO: 10
357-F	5'-TACGGGAGGCAGCAG-3'	SEQ ID NO: 11
1390-R	5'-CGGTGTGTRCAAGGCC-3'	SEQ ID NO: 12
1000-F	5'-CAACGAGCGCAACCCT',	SEQ ID NO: 13

Degeneration of primer 1390-R: R is A or G. The 16 S rDNA sequences from DSM 29869, DSM 29870, DSM 29871 and DSM 29872 are shown as SEQ ID NO: 1-4 in the sequence listing respectively. The 16 S rDNA sequences from DSM 29869, DSM 29870,
 20 DSM 29871 and DSM 29872 were analyzed by BLAST against EMBL database and showed identity to 16 S rDNA sequences of *Bacillus subtilis* (SEQ ID NO: 2 and SEQ ID NO: 3) and to *Bacillus amyloliquefaciens* (SEQ ID NO: 1 and SEQ ID NO: 4).

In order to study the phylogenetic affiliation of SEQ ID NO: 1 to SEQ ID NO: 4 the sequences were analyzed by a ClustalW alignment in MegAlign (DNASTAR) using SEQ ID
 25 NO: 5 to SEQ ID NO: 7 as benchmark. These sequences are reference 16S rDNA sequences of the type strains of *Bacillus vallismortis* taken from AB021198 (SEQ ID NO: 5), *Bacillus subtilis* taken from AJ276351 (SEQ ID NO: 6) and *Bacillus amyloliquefaciens* taken from AB255669 (SEQ ID NO: 7).

The ClustalW alignment of SEQ ID NO: 1 to SEQ ID NO: 7 (Fig. 1) shows 7
 30 nucleotide positions where 2 or more sequences have a nucleotide that deviates from the

other. For numbering the positions in SEQ ID NO: 6 are used. At position 152 SEQ ID NO: 2 is identical to *Bacillus amyloliquefaciens* and *Bacillus vallismortis*, but different from the rest that are identical to each other. At position 174 SEQ ID NO: 1 and SEQ ID NO: 4 are identical to *Bacillus amyloliquefaciens*, but different from the rest that are identical to each other. At position 257 SEQ ID NO: 4 is identical to *Bacillus amyloliquefaciens*, but different from the rest that are identical to each other. At position 437, 444 and 455 SEQ ID NO: 2 and SEQ ID NO: 3 are identical to *Bacillus subtilis*, but different from the rest that are identical to each other. At position 1223 SEQ ID NO: 1 is identical to *Bacillus amyloliquefaciens*, but different from the rest that are identical to each other. The variation in the 16S rDNA genes support the species affiliation seen in the BLAST report.

Description of the biological material

Bacillus amyloliquefaciens DSM 29869 was isolated by Novozymes (Novo Nordisk) from an environmental sample collected at Jamaica in 1990. The strain was identified as *Bacillus amyloliquefaciens* based on 16S rDNA sequencing.

Bacillus subtilis DSM 29870 was isolated by Novozymes (Novo Nordisk) from an environmental sample collected at Jamaica in 1990. The strain was identified as *Bacillus subtilis* based on 16S rDNA sequencing.

Bacillus subtilis DSM 29871 was isolated for Novozymes by a Danish high-school student at Brønderslev Gymnasium from a soil sample from private property in Denmark in 2008. The strain was identified as *Bacillus subtilis* based on 16S rDNA sequencing.

Bacillus amyloliquefaciens DSM 29872 was isolated from an environmental sample from the USA in 2008. The strain was identified as *Bacillus amyloliquefaciens* based on 16S rDNA sequencing.

Example 4: Determination of inhibition of growth of *Escherichia coli*

Escherichia coli strains, ATCC10536 or ATCC25922, were grown overnight in tryptic soy broth (BD part 211822) supplemented with 0.6% yeast extract (BD part 212750) at 35°C under static anaerobic conditions. 100 µL of the overnight culture of *Escherichia coli* was added to 250 mL of tryptic soy agar supplemented with 0.6% yeast extract at 40°C and poured into rectangular petri plates (Nunc part 267060). The inoculated agar was then allowed to cool at room temperature after which an 8mm diameter well was made in the agar.

The *Bacillus* strain DSM 29869, DSM 29870, DSM 29871, or DSM 29872 was grown overnight in tryptic soy broth at 35°C under aerobic conditions. 1000 µL of *Bacillus subtilis* culture was collected and fractionated into cell-free supernatant and cells by centrifugation. 20 µL of cell-free supernatant or 100x diluted cells in phosphate buffer saline were added

directly to the wells in the *Escherichia coli* inoculated agar plates. A control well contained 20 µL of phosphate buffer saline. The plates were incubated for 18 hours at 30°C under aerobic conditions.

Inhibition of the *Escherichia coli* strain was noted by a circular clearing zone around the well of interest. The phosphate buffer saline well was considered a negative control based on lack of clearing zone around the well.

Cell-free supernatant and 100x diluted cells of *Bacillus* strains DSM 29869, DSM 29870, DSM 29871, and DSM 29872 were able to consistently inhibit growth of the *E. coli* strains ATCC10535 and ATCC25922 *in vitro*. Inhibition was also seen by competitor strain CloSTAT, for both supernatant and cells.

Example 5: Sensitivity to antibiotics

The minimal inhibitory concentrations (MIC) of eight antibiotics against *Bacillus* strains DSM 29869, DSM 29870, DSM 29871, and DSM 29872 were determined using broth micro dilution essentially as described in the CLSI guidelines (M07-A9 Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; 2012). Only modification was that volumes were changed; 90 µl Mueller Hinton Broth 2 with bacteria was added to 10 µl antibiotics dilutions, cfu's of bacteria and concentration of antibiotics were changed so final concentrations and final cfu's matched the guideline. The plates were incubated for 20-24 h instead of 16-20 h. The control strain recommended in the CLSI standard *Staphylococcus aureus* ATCC 29213 was used as control strain.

Strains:

Bacillus amyloliquefaciens DSM 29869

Bacillus subtilis DSM 29870

Bacillus subtilis DSM 29871

Bacillus amyloliquefaciens DSM 29872

S. aureus ATCC 29213

Antibiotics:

Chloramphenicol (Sigma C1919, 10 mg/ml, solubilized in 96% ethanol)

Clindamycin (Sigma PHR1159, 10 mg/ml, solubilized in water)

Erythromycin (ABBOTICIN from Amdipharm 40501, 50 mg/ml, solubilized in 96% ethanol)

Gentamycin (Biomedicals inc., 190057, 10 mg/ml, solubilized in water)

Kanamycin (Sigma, CAS no. 25389-94-0, 50 mg/ml, solubilized in water)

Streptomycin (Sigma, S1277, 10 mg/ml, solubilized in water)

Tetracycline (Sigma, T3383, 10 mg/ml, solubilized in water)

Vancomycin (Sigma, V-8138, 10 mg/ml, solubilized in water)

Mueller Hinton Broth 2 (Sigma/Fluka 90922)

0.9% NaCl (Sigma/RdH 31434/Merck 106404)

5 Tryptone soya agar plates (Oxoid CM 131)

Microtiter plates: Costar plate, polypropylene, round bottom, Corning 3879

Preparation of bacteria:

A few colonies of *Bacillus spp.* (<1 day old) were inoculated into Mueller Hinton Broth
10 2 (MHB) and incubated for around 4 hours at 37°C. OD₆₀₀ (BioPhotometer plus, Eppendorf)
was measured and adjusted to 0.25 (equivalent to McFarland 0.5) in MHB. For the control
strain direct colony suspension was used. A few colonies of *S. aureus* ATCC 29213 (<1 day
old) were suspended in MHB and OD₆₀₀ (BioPhotometer plus, Eppendorf) was measured and
adjusted to 0.10-0.12 (equivalent to McFarland 0.5) in MHB. The bacterial suspensions were
15 diluted 200x in MHB.

Preparation of assay plates:

Antibiotics were diluted to the concentration of 640 µg/mL in MHB. A two fold dilution
series was prepared in MHB down to the concentration 0.625 µg/ml. 10 µl of each dilution
20 and of each antibiotic was pipetted into a micro-titre plate. Later, when the antibiotics were
mixed with the suspension of bacteria, the samples were diluted 10x (10 µL sample in a total
volume of 100 µl). This resulted in the final test range of 0.06-64 µg/ml.

If the plates were not used right away the plates were stored in the freezer at -20°C
until usage.

25 90 µl of the bacterial suspensions were added to the assay plates. The assay plates
were incubated in a plastic bag with a wet cloth at 37°C for 20-24 h. The MIC was
determined as the lowest concentration of antibiotic that completely inhibited growth of
bacteria as detected by the unaided eye.

30 CFU estimation:

A 10-fold dilution series in 0.9% NaCl was made to the 10⁻³ of the cultures inoculated
into the micro-titre plate. 2x 100 µl from the 10⁻³ dilution were plated onto two TSA plates.
The plates were incubated overnight at 37°C. Number of CFU/ml was counted.

Three biological replicates of the assay were performed for *Bacillus subtilis* DSM
35 29871, *Bacillus amyloliquefaciens* DSM 29869 and *Bacillus subtilis* DSM 29870 (MIC 1-3),
while four biological replicates were performed for *Bacillus amyloliquefaciens* DSM 29872
(MIC 2-5).

Results:

The MIC values obtained for *B. subtilis* DSM 29870 showed that the breakpoint values were equal to or below the breakpoint values given in the EFSA guideline (EFSA journal 2012; 10(6):2740). For *B. subtilis* DSM 29869 the MIC values obtained in two out of the three biological replicates showed MIC values equal to or below the breakpoint. In MIC 1 the value for Tetracycline was 16. However, due to analytical variance of the method MIC results of one dilution above the breakpoint is in general accepted and the strain may be regarded as being sensitive. Thus, according to EFSA these two strains are regarded as sensitive to all the 8 antibiotics included in the test (Tables 5.2 and 5.3).

For the strains *B. subtilis* DSM 29871 and *B. amyloliquefaciens* DSM 29872 the MIC values showed that the two strains were sensitive to seven out of the eight antibiotics (Tables 5.1 and 5.4). An increased tolerance was observed towards Streptomycin and according to EFSA the strains are classified as resistant to Streptomycin.

As a control *S. aureus* ATCC 29213 was tested in parallel and had MIC values within the ranges given by the CLSI standard (M100-S24 Performance Standards for Antimicrobial Susceptibility Testing; informational Supplement, 2014) (Table 5.5).

The amount of bacteria inoculated into the assay plates was measured (CFU/ml). In general the CFU/ml was very close to the target value of 5×10^5 CFU/ml. However, the CFU/ml for *Bacillus* strains may be associated with some uncertainty, since the bacteria tend to aggregate and once aggregated will only result in one colony forming unit (Tables 5.1 and 5.2).

Table 5.1: MIC results for *B. subtilis* DSM 29871

Antibiotic	MIC 1 µg/ml	MIC 2 µg/ml	MIC 3 µg/ml	EFSA* breakpoints µg/ml
Chloramphenicol	8	4	8	8
Clindamycin	0.5	0.25	0.5	4
Erythromycin	0.125	0.125	0.125	4
Gentamycin	0.25	0.125	0.25	4
Kanamycin	2	1	2	8
Streptomycin	32	16	32	8
Tetracycline	2	4	4	8
Vancomycin	0.25	0.25	0.25	4
CFU/ml	3.6×10^5	n.r.	n.r.	

* EFSA Journal 2012; 10(6):2740

n.r. not registered

Table 5.2: MIC results for *B. amyloliquefaciens* DSM 29869

Antibiotic	MIC 1 µg/ml	MIC 2 µg/ml	MIC 3 µg/ml	EFSA* breakpoints µg/ml
Chloramphenicol	4	4	4	8
Clindamycin	1	0.5	1	4
Erythromycin	0.5	0.06	0.06	4
Gentamycin	0.5	0.125	0.06	4
Kanamycin	2	0.5	0.5	8
Streptomycin	2	4	4	8
Tetracycline	16	8	8	8
Vancomycin	0.25	0.25	0.25	4
CFU/ml	3.9*10 ⁵	2.7*10 ⁵	1.2*10 ⁵	

* EFSA Journal 2012; 10(6):2740

Table 5.3: MIC results for *B. subtilis* DSM 29870

Antibiotic	MIC 1 µg/ml	MIC 2 µg/ml	MIC 3 µg/ml	EFSA* breakpoints µg/ml
Chloramphenicol	4	8	4	8
Clindamycin	0.25	0.25	0.25	4
Erythromycin	0.125	0.125	0.125	4
Gentamycin	0.25	0.125	0.25	4
Kanamycin	2	1	2	8
Streptomycin	4	4	8	8
Tetracycline	0.25	0.25	0.25	8
Vancomycin	0.25	0.25	0.25	4
CFU/ml	4.2*10 ⁵	2.1*10 ⁵	1.4*10 ⁵	

* EFSA Journal 2012; 10(6):2740

Table 5.4: MIC results for *B. amyloliquefaciens* DSM 29872

Antibiotic	MIC 2 µg/ml	MIC 3 µg/ml	MIC 4 µg/ml	MIC 5 µg/ml	EFSA* breakpoints µg/ml
Chloramphenicol	4	4	4	4	8
Clindamycin	0.5	0.5	0.5	0.5	4
Erythromycin	0.06	0.125	0.06	0.06	4
Gentamycin	0.06	0.06	0.125	0.125	4
Kanamycin	1	0.5	2	1	8
Streptomycin	8 (32)	8	32	>64	8
Tetracycline	0.25	0.25	0.125	0.125	8
Vancomycin	0.25	0.25	0.125	0.125	4
CFU/ml	2.7*10 ⁵	1.5*10 ⁵	4.7*10 ⁵	6.3*10 ⁵	

* EFSA Journal 2012; 10(6):2740

Table 5.5: MIC results for *S. aureus* ATCC 29213

Antibiotic	MIC 1 µg/ml	MIC 2 µg/ml	MIC 3 µg/ml	MIC 4 µg/ml	MIC 5 µg/ml	CLSI* breakpoints µg/ml
Chloramphenicol	8-16	16	8	16	16	2-16
Clindamycin	0.125	0.25	0.125	0.06	0.06	0.06-0.25
Erythromycin	0.5	0.5	0.5	0.5	0.5	0.25-1
Gentamycin	0.5	0.5	0.5	1	1	0.12-1
Kanamycin	4	4	4	4	4	1-4
Streptomycin	8	8	8	16	8	No information
Tetracycline	0.5	1	1	0.5	0.5	0.12-1
Vancomycin	0.5	1	1	1	1	0.5-2
CFU/ml	5.2*10 ⁵	7.0*10 ⁵	7.0*10 ⁵	7.3*10 ⁵	4.3*10 ⁵	

* M100-S24 Performance Standards for Antimicrobial Susceptibility Testing; informational Supplement, 2014

Example 6: Enzyme activityPreparation of overnight pre-cultures

Overnight cultivation was made in a 96-deep well plate with 1.25 mL of LB broth (Difco; BD #244610) per well. Each strain was tested in duplicate and inoculated with a single colony from a pure Standard Methods Agar plate (SMA plates, Smith River Biologicals #11-00450). Before incubation, a single sterile Aeraseal breathable seal (Fisher #50-212-463) was placed on top of the plate, and the plate was incubated with shaking at 35°C overnight under aerobic conditions.

Determination of anaerobic growth

SMA plates were streaked from the overnight cultures using a 10 µL inoculating loop. Up to four strains were streaked per plate, into separate quadrants.

- 5 Plates were incubated at 39°C in an AnaeroJar (Thermo Scientific Oxoid, Fisher# OXAG0025A) along with an AnaeroGen sachet (Thermo Scientific Oxoid, Fisher #OXAN0025A) to maintain anaerobic conditions (~0.1% oxygen). Work was performed under aseptic conditions in biological hood, using sterile materials.

10 Enzyme activity determination using AZCL-substrate agar plates

Preparation of media with AZCL substrates which can be autoclaved AZCL-cellulose (AZCL-HE-Cellulose, I-AZCEL, Megazyme), arabinoxylan (AZCL-Arabinoxylan, wheat, I-AZWAX, Megazyme), arabinan (AZCL-Arabinan, debranched, I-AZDAR, Megazyme):

- 15 Into two (2) 500 mL beakers, 300 mL of 0.1x and 1x LB agar were prepared for each beaker, with a magnetic stir bars added to each. While stirring, 100 mL of LB condition from the beakers were added to three (3) 250 mL Wheaton bottles, thus giving 3 bottles with 0.1x LB and 3 bottles with 1x LB. AZCL-cellulose, arabinan, and arabinoxylan were then added separately to each of a 1x and a 0.1x bottle with stirring. A total of 0.05 grams of substrate is added per 100 mL of media. If more than 100 mL of media is being used, the amount of
- 20 substrate added is adjusted by using the 0.05g/100mL ratio. The media were autoclaved to sterilize. Note: Only these three AZCL substrates can be autoclaved without obvious dye release indicating substrate instability. After autoclaving, media is kept at ≥50°C until ready to pour plates. The pH was checked, and if necessary adjusted to pH 7 using 10% HCl or 2N NaOH, maintaining sterility.

25

Preparation of media with AZCL substrates which can't be autoclaved

These substrates were: AZCL-amylose (I-AZAMY, Megazyme), AZCL-casein (I-AZCAS, Megazyme), AZCL-xylan (Xylan, birchwood (I-AZXBW, Megazyme)

- 30 Into two (2) 1000 mL Wheaton bottles, 400 mL of 0.1x LB and 1x LB was prepared and a magnetic stir bar was added to each. The media was autoclaved to sterilize, prior to adding AZCL reagent. After autoclaving, media was kept at 50-55°C until ready to add AZCL substrate and pour plates. The pH was checked, and if necessary adjusted to pH 7 using 10% HCl or 2N NaOH, maintaining sterility.

Pouring AZCL substrate media agar plates

Work was performed under aseptic conditions in biological hood, using sterile materials. These steps were done relatively quickly, so that the agar did not solidify before transferring into the 96-well plate.

5 For substrates which could be autoclaved (AZCL-cellulose, AZCL-arabinoxylan, AZCL-arabinan): The warm (~50°C), autoclaved liquid agar media was magnetically stirred in order to suspend the insoluble AZCL substrate. An aliquot was transferred to a sterile solution basin (Periodic mixing of the material in the solution basin was necessary to keep the substrate suspended.) A repeating, multi-channel pipette (e.g., Matrix 1250 µL pipet
10 along with the corresponding 1250 µL sterile pipet tips) was used to dispense 180 µL into each well of a sterile 96-well plate. A set of tips would typically last for up to six columns per plate. Once the plate was filled, a sterile lid was added, and it was allowed to solidify. One plate was made for each substrate (AZCL-cellulose, AZCL-arabinoxylan, AZCL-arabinan) and condition (0.1x and 1x LB).

15 For the substrates that could not be autoclaved (AZCL-amylose, AZCL-casein, AZCL-xylan): Into a sterile 250 mL beaker with magnetic stir-bar was added 100 mL of warm, sterile 0.1x or 1X LB media (no AZCL substrate added). With stirring, add 0.05 mg of AZCL substrate, and continue stirring until re-suspended. It may help to add the substrate slowly at first. Once substrate was re-suspended, it was poured into solution basin and added to
20 plates as described above. One plate was made for each substrate (AZCL-cellulose, AZCL-arabinoxylan, AZCL-arabinan) and condition (0.1x and 1x LB).

Inoculating the 96-well plates, incubation, and evaluation

25 Work was performed under aseptic conditions in biological hood, using sterile materials. The AZCL substrate agar plates were inoculated by applying 2 µL of overnight culture that was dispensed on top of the agar. The added liquid should be visible on top of the agar. When moving to the next plate, be sure that there are no drops left at the end of the pipet tips to ensure that each well has been inoculated. Repeat for all AZCL substrate plates. Control enzyme dilutions were prepared, each at 1/100, diluting in sterile phosphate dilution
30 buffer: 10 µL enzyme preparation + 990 µL buffer. Control enzyme wells were then inoculated with 10 µL of the appropriate enzyme dilution for each substrate

Table 6.1: Enzymes and respective substrates

Enzyme	Substrate
Amylase	Amylose
Arabinase	Arabinan
Cellulase	Cellulose
Protease	Casein
Xylanase	Arabinoxylan, Xylan

Table 6.2: Enzyme production under aerobic conditions

	30°C Aerobic, 24 h					
organism ID	Amylose	Arabinan	Arabinoxylan	Casein	Cellulose	Xylan
DSM 29869	+	+	+	+	-	+
DSM 29870	+	+	+	+	-	+
DSM 29871	+	+	+	+	+	+
DSM 29872	+	+	+	+	-	+

5 Table 6.3: Enzyme production under anaerobic conditions, 24 hours

	30°C Anaerobic, 24 h					
organism ID	Amylose	Arabinan	Arabinoxylan	Casein	Cellulose	Xylan
DSM 29869	+	+	+	-	-	+
DSM 29870	+	+	+/-	+	-	+
DSM 29871	-	+	+	-	+	+
DSM 29872	+	+	+	-	-	+

Table 6.4: Enzyme production under anaerobic conditions, 48 hours

	30°C Anaerobic, 48 h					
organism ID	Amylose	Arabinan	Arabinoxylan	Casein	Cellulose	Xylan
DSM 29869	+	+	+	+	-	+
DSM 29870	+	+	+/-	+	-	+
DSM 29871	-	+	+	-	+	+
DSM 29872	+	+	+	-	-	+

Example 7: Efficacy of the DSM 29870 strain in *Clostridium perfringens* challenge conditions

Material and Methods

Three independent battery studies have been conducted to evaluate the influence of the strain DSM 29870 on the development of an induced necrotic enteritis.

In each experiment, a total of 256 one day-old male broiler chickens Cobb x Cobb were allocated to Petersime battery cages (8 birds/cage). Cages were used in factorial and completely randomized design with 8 cages per treatment (*i.e.*, 64 animals/treatment).

An unmedicated corn/soybean meal-based commercial broiler starter ration was formulated (Table 7.1). Feed and water were available *ad libitum* throughout the trials.

Table 7.1: Composition of the basal experimental diet⁴ (starter d1-d28)

	Starter (0-28)
<i>Ingredients</i>	
Corn, yellow, ground	56.12
Soybean meal (48)	37.50
Fat, poultry	3.00
Dicalcium phosphate	1.75
Limestone	0.80
Salt	0.30
Vitamin premix ¹	0.25
DL-Methionine	0.20
Trace mineral premix ²	0.08
<i>Calculated Nutritional Content</i>	
ME, kcal/kg	3.096
Protein, %	22.30
Lysine, %	1.180
Methionine, %	0.530
Met + Cys, % ³	0.890

¹Vitamin mix provided the following (per kg of diet): thiamin•mononitrate, 2.4 mg; nicotinic acid, 44 mg; riboflavin, 4.4 mg; D-Ca pantothenate, 12 mg; vitamin B₁₂ (cobalamin), 12.0 µg; pyridoxine•HCL, 4.7 mg; D-biotin, 0.11 mg; folic acid, 5.5 mg; menadione sodium bisulfite complex, 3.34 mg; choline chloride, 220 mg; cholecalciferol, 27.5 µg; trans-retinyl acetate, 1,892 µg; all-rac α tocopheryl acetate, 11 mg; ethoxyquin, 125 mg.

²Trace mineral mix provided the following (per kg of diet): manganese ($\text{MnSO}_4 \cdot \text{H}_2\text{O}$), 60 mg; iron ($\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$), 30 mg; zinc (ZnO), 50 mg; copper ($\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$), 5 mg; iodine (ethylene diamine dihydroiodide), 0.15 mg; selenium (NaSeO_3), 0.3 mg.

³Met= methionine; Cys= cysteine.

- 5 ⁴The basal feed will not contain any probiotic/ prebiotic feed additives, NSPases, coccidiostats or antibiotic growth promoter.

The 4 experimental groups consisted in (Table 7.2): T1, non-infected and non-treated animals; T2, *C. perfringens* infected and non-treated animals; T3, *C. perfringens* infected animals fed with the basal diet containing bacitracin (included at 50 ppm, from d1 to d28); T4, *C. perfringens* infected animals fed with the basal diet containing the DSM 29870 strain (included at 1.10^9 CFU/kg of feed in trial 1 and 5.10^8 CFU/kg of feed in trials 2 and 3, from d1 to d28).

15 Table 7.2: Experimental treatments

Treatments	
T1	Non-infected, non-treated
T2	Infected, non-treated
T3	Infected, bacitracin
T4	Infected, DSM 29870

On day 14, all birds were orally inoculated with a coccidial inoculum containing approximately 5,000 oocysts of *Eimeria maxima* per bird.

20 Once daily on d19, d20 and d21, all birds, except T1, were administered (by oral gavage) 1 mL of a fresh broth culture of *C. perfringens*.

On d21, three birds from each cage were selected, sacrificed, weighed, and examined for the degree of presence of necrotic enteritis lesions. The scoring was based on a 0 to 3 score, with 0 being normal and 3 being the most severe.

25 The live body weight and the feed were recorded at the start (d1) and the end of the experiment (d28), on a cage basis, to calculate performance:

- Body Weight Gain (BWG) = $(\text{Live Body Weight})_{\text{d28}} - (\text{Live Body Weight})_{\text{d1}}$
 - Feed Conversion Ratio (FCR) = ratio between feed consumed and weight gained
- Mortality was recorded daily and the FCR was adjusted accordingly.

30 Necrotic enteritis lesion scores, total mortality and % of dead birds with necrotic enteritis lesions were also calculated

Data (n = 32) were subjected to an ANOVA, with complete randomized block design using the ANOVA procedure of XLSTAT (Addinsoft 1995-2014) to establish differences

between diets. Pen was considered as the experimental unit. The model included diets (n=4) and block as fixed effect. Results are reported as least square means. LS means were assumed to be different at $P \leq 0.05$.

5 Experimental results

The results obtained on performance, necrotic enteritis lesion scores, mortality and % of dead birds with necrotic enteritis lesions parameters are shown in Tables 7.3.

Tables 7.3: Effect of dietary supplementation of the strain DSM 29870 on the negative impact on production parameters due to induced necrotic enteritis in three independent trials.

Trial 1	BWG ⁵ (g/bird)	FCR ⁵	Lesion score ⁶	Total mortality (%)	Dead birds with necrotic enteritis lesions (%)
	(d1-d28)	(d1-d28)	(d21)	(d1-d28)	(d1-d28)
T1 ¹	720a	1.666c	0.0a	7.5b	0.0b
T2 ²	549b	2.071a	0.2a	27.5a	17.2a
T3 ³	633ab	1.813bc	0.2a	5.0b	1.6b
T4 ⁴	628ab	1.843bc	0.3a	5.0b	3.1b

¹ Non-infected, non-treated group fed with the basal diet.

² Infected, non-treated group fed with the basal diet.

³ Infected group fed with the basal diet supplemented with bacitracin (50 ppm).

⁴ Infected group fed with the basal diet supplemented with DSM 29870 (1.10^9 CFU/kg of feed).

⁵ Means from 8 replicates of 8 birds per cage for each group. Raw data (n=32) were analyzed using variance analysis with block (n=8) and treatments (n=4) as fixed effect.

⁶ Means from 8 replicates of 3 birds per cage for each group. Raw data (n=32) were analyzed using variance analysis with block (n=8) and treatments (n=4) as fixed effect.

^{a,b,c} Values with different superscripts are significantly ($P < 0.05$) different from each other.

Trial 2	BWG ⁵ (g/bird)	FCR ⁵	Lesion score ⁶	Total mortality (%)	Dead birds with necrotic enteritis lesions (%)
	(d1-d28)	(d1-d28)	(d21)	(d1-d28)	(d1-d28)
T1	685a	1.692b	0.0d	2.5c	0.0c
T2	435c	2.404a	1.2a	32.5a	20.3a
T3	665ab	1.759b	0.6b	12.5bc	6.3c
T4	654ab	1.768b	0.4bc	25.0ab	9.4bc

¹ Non-infected, non-treated group fed with the basal diet.

² Infected, non-treated group fed with the basal diet.

³ Infected group fed with the basal diet supplemented with bacitracin (50 ppm).

⁴ Infected group fed with the basal diet supplemented with DSM 29870 (5.10^8 CFU/kg of feed).

⁵ Means from 8 replicates of 8 birds per cage for each group. Raw data (n=32) were analyzed using variance analysis with block (n=8) and treatments (n=4) as fixed effect.

⁶ Means from 8 replicates of 3 birds per cage for each group. Raw data (n=32) were analyzed using variance analysis with block (n=8) and treatments (n=4) as fixed effect.

^{a,b,c} Values with different superscripts are significantly ($P<0.05$) different from each other.

Trial 3	BWG ⁵ (g/bird)	FCR ⁵	Lesion score ⁶	Total mortality (%)	Dead birds with necrotic enteritis lesions (%)
	(d1-d28)	(d1-d28)	(d21)	(d1-d28)	(d1-d28)
T1	667a	1.752c	0.0b	10.0a	0.0c
T2	521b	1.988a	0.5a	25.0a	15.6a
T3	645a	1.786bc	0.6a	15.0a	4.7bc
T4	638a	1.851b	0.4a	10.0a	4.7bc

¹ Non-infected, non-treated group fed with the basal diet.

² Infected, non-treated group fed with the basal diet.

³ Infected group fed with the basal diet supplemented with bacitracin (50 ppm).

⁴ Infected group fed with the basal diet supplemented with DSM 29870 (5.10^8 CFU/kg of feed).

⁵ Means from 8 replicates of 8 birds per cage for each group. Raw data (n=32) were analyzed using variance analysis with block (n=8) and treatments (n=4) as fixed effect.

⁶ Means from 8 replicates of 3 birds per cage for each group. Raw data (n=32) were analyzed using variance analysis with block (n=8) and treatments (n=4) as fixed effect.

5 ^{a,b,c} Values with different superscripts are significantly ($P<0.05$) different from each other.

The data obtained from the three challenge studies showed that the strain DSM 29870 can counteract the negative impact of an induced necrotic enteritis on the production parameters with a significant effect in all challenge trials on the FCR and the mortality due to necrotic enteritis, relative to the positive control group (non-infected, non-treated animals).

Furthermore, there was no significant difference between the DSM 29870 fed group and the bacitracin fed group with regards to all parameters measured in these trials.

The challenge studies described above showed that the administration of the DSM 29870 strain could prevent necrotic enteritis in broiler chickens.

Example 8: First Performance Trial with Broiler Chickens (521)

Material and methods

A total of 2000 one day-old male broiler chickens Cobb 500 were individually weighed (body weight 42 g $\pm$ 1 g) and allocated in floor pens (50 birds/pen). Pens were used in factorial and completely randomized design with 8 pens per treatment (i.e., 400 animals/treatment). The 5 experimental treatments consisted in (Table 8.1): T1, negative control basal diets; T2, basal diets containing the strain DSM 29870 (included at 5.10^8 CFU/kg of feed, from d1 to d35); T3, basal diets containing the strain DSM 29871 (included at 5.10^8 CFU/kg of feed, from d1 to d35); T4 basal diets containing the strain DSM 29872 (included at 5.10^8 CFU/kg of feed, from d1 to d35); T5, basal diet containing a marketed probiotic, GalliPro Max (included at the commercial dose, 8.10^8 CFU/kg of feed, from d1 to d35).

Basal diets consisted in 3 phases feeding program: starter phase (from 1 to day 12), grower phase (from 13 to day 28) and finisher phase (from 29 to day 35). Every phases were formulated to meet or exceed animal requirement and agree with standard commercial US corn-soybean meal-based broiler diets. The dietary and raw material composition of these diets is given in Table 8.2. All experimental diets include phytase (Ronozyme P, 250 FYT/kg) and did not contain any coccidiostat, NSPase or growth promoting substance. Feed and water were available *ad libitum* throughout the trial.

Table 8.1: Experimental treatments

Treatments	
T1	Control
T2	T1 + DSM 29870
T3	T1 + DSM 29871
T4	T1 + DSM 29872
T5	T1 + GalliPro Max

Table 8.2: Composition of the basal experimental diets

	Starter (0-12d)	Grower (13-28d)	Finisher (29-35d)
Ingredients	%		
Corn, yellow, grain	64.673	66.460	68.491
Soybean meal, dehulled, solvent	29.020	26.662	24.677
Ampro 55 (animal by-product 55% protein)	2.500	3.000	3.000
Calcium carbonate	0.886	0.735	0.684
Fat, vegetable	0.883	1.485	1.702
Dicalcium phosphate	0.706	0.612	0.500
Salt, plain (NaCl)	0.439	0.435	0.436
Methionine MHA	0.358	0.259	0.221
L-Lysine	0.273	0.208	0.145
L-Threonine 98.5	0.103	0.000	0.000
Trace Mineral ¹	0.075	0.075	0.075
Vitamin premix ²	0.065	0.050	0.050
Ronozyme P-(ct)	0.019	0.019	0.019
Calculated Nutritional Content			
ME (kcal/kg)	3,067	3,130	3,165
Crude protein (%)	20.96	20.03	19.16
Dig. Lysine (%)	1.20	1.10	1.00
Dig. Methionine (%)	0.61	0.52	0.48
Dig. TSAA (%)	0.90	0.80	0.75
Dig. Threonine (%)	0.81	0.68	0.65
Calcium (%)	0.90	0.85	0.8
Avail. phosphorus (%)	0.42	0.42	0.4

¹Vitamin mix will provide the following (per kg of diet): thiamin•mononitrate, 2.4 mg; nicotinic acid, 44 mg; riboflavin, 4.4 mg; D-Ca pantothenate, 12 mg; vitamin B₁₂ (cobalamin), 12.0 µg; pyridoxine•HCL, 4.7 mg; D-biotin, 0.11 mg; folic acid, 5.5 mg; menadione sodium bisulfite complex, 3.34 mg; choline chloride, 220 mg; cholecalciferol, 27.5 ug; trans-retinyl acetate, 1,892 ug; all-rac α tocopheryl acetate, 11 mg; ethoxyquin, 125 mg.

²Trace mineral mix provided the following (per kg of diet): manganese (MnSO₄•H₂O), 60 mg; iron (FeSO₄•7H₂O), 30 mg; zinc (ZnO), 50 mg; copper (CuSO₄•5H₂O), 5 mg; iodine (ethylene diamine dihydroiodide), 0.15 mg; selenium (NaSeO₃), 0.3 mg.

The live body weight and the feed were recorded at the start (d1) and the end of the experiment (d35), on a pen basis, to calculate performance:

- Feed Intake (FI) = (Remaining Feed)_{d35} – (Issued Feed)_{d1}
- Body Weight Gain (BWG) = (Live Body Weight)_{d35} – (Live Body Weight)_{d1}
- Feed Conversion Ratio (FCR) = ratio between feed consumed and weight gained

Mortality was recorded daily and the FCR was adjusted accordingly.

Data (n = 40) were subjected to an ANOVA, with complete randomized block design using the ANOVA procedure of XLSTAT (Addinsoft 1995-2014) to establish differences between diets. Pen was considered as the experimental unit. The model included diets (n=5) and block as fixed effect. Results are reported as least square means. LS means were assumed to be different at $P \leq 0.05$.

Similar calculations were realized in the example 9 and 10.

Similar calculations were realized in the example 3 and 4. Experimental results

The results obtained on performance parameters for the whole experimental period are shown in Table 8.3.

Table 8.3: Effect of dietary supplementation of the strains DSM 29870, DSM 29871 and DSM 29872 on broiler performance

Parameters ⁴	Control ¹	DSM 29870 ² supplemented group	DSM 29871 ² supplemented group	DSM 29872 ² supplemented group	GalliPro Max ³ supplemented group
FI (g/bird)	3245a	3256a	3221a	3263a	3203a
Relative to the control	-	+0.3%	-0.7%	+0.6%	-1.3%
BWG (g/bird)	1987ab	2026ab	2002ab	2038a	1973b
Relative to the control	-	+2.0%	+0.8%	+2.6%	-0.7%
FCR	1.633a	1.607b	1.609b	1.602b	1.624ab
Relative to the control	-	-1.6%	-1.5%	-2.0%	-0.6%
Mortality (%)	3.25a	3.50a	5.25a	4.5a	6.25a

¹ Group fed with the basal diet.

² DSM 29870/29871/29872 supplemented group fed with the basal diet including strain DSM 29870/29871/29872 at $5 \cdot 10^8$ CFU/kg of feed.

³ GalliPro Max supplemented group fed with the basal diet including GalliPro Max at 8.10^8 CFU/kg of feed.

⁴ Means from 8 replicates of 50 birds per pen for each group. Raw data (n=24) were analyzed using variance analysis with block (n=8) and treatments (n=3) as fixed effect.

^{a,b,c} Values with different superscripts are significantly ($P<0.05$) different from each other.

Feed additives did not significantly affect animal feed intake relative to the control.

Body weight was slightly increased using the NZB strains whereas GalliPro Max tended to decrease this parameter.

The DSM 29870, DSM 29871 and DSM 29872 strains also improved significantly the feed conversion ratio by 1.6%, 1.5% and 1.9%, respectively, with no significant differences between the groups fed with NZB strains. GalliPro Max had only a slight and non-significant effect on the feed conversion ratio.

None of the probiotic tested here had a significant effect on the mortality level.

Example 9: Second Performance Trial with Broiler Chickens (522)

Material and methods

A total of 900 one day-old male broiler chickens Ross 708 were individually weighed (body weight 47.1 ± 1.1 g) and allocated in floor pens (15 birds/pens). Pens were used in factorial and completely randomized design with 12 pens per treatment (*i.e.*, 180 animals/treatment). The 5 experimental treatments consisted in (Table 9.1): T1, negative control basal diets; T2, basal diets containing the strain DSM 29870 (included at 5.10^8 CFU/kg of feed, from d1 to d35); T3, basal diets containing the strain DSM 29871 (included at 5.10^8 CFU/kg of feed, from d1 to d35); T4 basal diets containing the strain DSM 29872 (included at 5.10^8 CFU/kg of feed, from d1 to d35); T5, basal diet containing a marketed probiotic, GalliPro Max (included at the commercial dose, 8.10^8 CFU/kg of feed, from d1 to d35).

Basal diets consisted in 3 phases feeding program: starter phase (from 1 to day 14), grower phase (from 15 to day 21) and finisher phase (from 22 to day 35). Every phases were formulated to meet or exceed animal requirement and agree with standard commercial US corn-soybean meal-based broiler diets. The dietary and raw material composition of these diets is given in Table 9.2. All experimental diets included vitamin and mineral premixes, but did not contain any coccidiostat, NSPase, phytase or growth promoting substance. Feed and water were available *ad libitum* throughout the trial.

Table 9.1: Experimental treatments

Treatments	
T1	Control
T2	T1 + DSM 29870
T3	T1 + DSM 29871
T4	T1 + DSM 29872
T5	T1 + GalliPro Max

Table 9.2: Composition of the basal experimental diets

	Starter (0-14d)	Grower (15-21d)	Finisher (22-35d)
<i>Ingredients</i>	%		
Corn	57.59	59.20	62.30
Wheat Flour	1.00	1.00	1.00
Soybean Meal (46% CP)	30.00	28.00	24.50
Poultry Meal	5.00	5.00	5.00
Poultry Fat	2.10	2.86	3.52
Limestone (36% Ca)	1.05	1.02	0.98
Dicalcium Phosphate (18% P)	1.76	1.52	1.42
Salt	0.35	0.32	0.29
Sodium Bicarbonate	0.16	0.14	0.11
L-Lysine HCL (78%)	0.23	0.22	0.21
DL-Methionine (99%)	0.36	0.33	0.29
L-Threonine (98%)	0.05	0.04	0.03
Selenium premix	0.05	0.05	0.05
Vitamin Mix	0.15	0.15	0.15
Mineral Mix	0.15	0.15	0.15
<i>Calculated Nutritional Content</i>			
ME (kcal/kg)	2968	3032	3096
Crude Protein (%)	23.3	22.4	20.78
Lysine (%)	1.35	1.29	1.18
Methionine (%)	0.69	0.65	0.59
Met+Cys (%)	1.05	1.00	0.92
Threonine (%)	0.86	0.81	0.75
Calcium (%)	1.03	0.96	0.91
Phosphorus Avail. (%)	0.46	0.41	0.39

¹Vitamin mix will provide the following (per kg of diet): vitamin A, 13 227 513 IU; vitamin D₃, 3 968 254 IU; vitamin E, 66 138 IU; vitamin B₁₂, 40 mg; biotin, 254 mg; menadione, 3 968 mg; thiamine, 3 968 mg; riboflavin, 13 228 mg; d-Pantothenic Acid, 22 046 mg; pyridoxine, 7 937 mg; niacin, 110 229 mg; folic acid, 2 205 mg; Selenium Premix: selenium, 600 ppm + calcium, 36 %

²Trace mineral mix provided the following (per kg of diet): calcium, Min: 15.7% and Max: 18.7%; manganese (Mn), 6.0%; zinc (Zn), 6.0%; iron (Fe), 4.0%; copper (Cu), 5000 ppm; iodine (I), 1250 ppm; cobalt (Co), 500 ppm

5 Experimental results

The results obtained on performance parameters for the whole experimental period are shown in Table 9.3.

Table 9.3: Effect of dietary supplementation of the strains DSM 29870, DSM 29871 and DSM 29872 on broiler performance

Parameters ⁴	Control ¹	DSM 29870 ² supplemented group	DSM 29871 ² supplemented group	DSM 29872 ² supplemented group	GalliPro Max ³ supplemented group
FI (g/bird)	3573a	3542a	3606a	3515a	3541a
<i>Relative to the control</i>	-	-0.9%	+0.9%	-1.0%	-0.9%
BWG (g/bird)	2361a	2416b	2401ab	2424b	2398ab
<i>Relative to the control</i>	-	+2.3%	+1.7%	+2.6%	+1.5%
FCR	1.513a	1.467b	1.501bc	1.451a	1.477ab
<i>Relative to the control</i>	-	-3.6%	-0.8%	-4.2%	-2.4%
Mortality (%)	15.6ab	7.8a	11.7ab	19.4b	10.0a

10 ¹ Group fed with the basal diet

² DSM 29870/29871/29872 supplemented group fed with the basal diet including strain DSM 29870/29871/29872 at 5.10^8 CFU/kg of feed.

³ GalliPro Max supplemented group fed with the basal diet including GalliPro Max at 8.10^8 CFU/kg of feed.

15 ⁴ Means from 12 replicates of 15 birds per pen for each group. Raw data (n=36) were analyzed using variance analysis with block (n=12) and treatments (n=3) as fixed effect.

^{a,b,c} Values with different superscripts are significantly ($P<0.05$) different from each other

Also in this experiment, the administration of the strains DSM 29870, DSM 29871, DSM 29872 and GalliPro Max did not significantly reduce the feed intake, but improved the body weight gain by 2.3%, 1.7%, 2.6% and 1.5%, respectively. Therefore, difference relatively to control was significant for DSM 29870 and DSM 29872.

The strains DSM 29870 and DSM 29872 improved significantly the feed conversion ratio by 3.6% and 4.2%, respectively. DSM 29871 and GalliPro Max also improved (slightly for DSM 29871) the feed conversion ratio in this experiment, but not significantly.

The strain DSM 29870 allowed a decrease of the mortality level in a numerically higher manner than GalliPro Max.

Example 10: Third Performance Trial with Broiler Chickens (051)

Material and methods

A total of 1080 one day-old male broiler chickens Ross PM3 were individually weighed (body weight 42 ± 3.5 g) and allocated in floor pens (18 birds/pens). Pens were used in factorial and completely randomized design with 12 pens per treatment (*i.e.*, 216 animals/treatment). The 3 experimental treatments consisted in (Table 10.1): T1, negative control basal diets; T2, basal diets containing the strain DSM 29870 (included at 5.10^8 CFU/kg of feed, from d1 to d35); T3, basal diets containing the strain DSM 29871 (included at 5.10^8 CFU/kg of feed, from d1 to d35); T4 basal diets containing the strain DSM 29872 (included at 5.10^8 CFU/kg of feed, from d1 to d35); T5, basal diet containing a marketed probiotic, GalliPro Max (included at the commercial dose, 8.10^8 CFU/kg of feed, from d1 to d35).

Basal diets consisted in 3 phases feeding program: starter phase (from 1 to day 21) and grower phase (from 22 to day 35). Every phases were formulated to meet or exceed animal requirement and agree with standard commercial EU corn-soybean meal-based broiler diets. The dietary and raw material composition of these diets is given in Table 10.2. All experimental diets included vitamin and mineral premixes, but did not contain any coccidiostat, NSPase, phytase or growth promoting substance. Feed and water were available *ad libitum* throughout the trial.

Table 10.1: Experimental treatments

Treatments	
T1	Control
T2	T1 + DSM 29870
T3	T1 + DSM 29871
T4	T1 + DSM 29872
T5	T1 + GalliPro Max

Table 10.2: Composition of the basal experimental diets

	Starter (0-21d)	Grower (22-35d)
<i>Ingredients</i>	%	
Maize	51.68	55.58
Soybean Meal (48% CP)	39.03	34.55
Soybean oil	4.27	5.18
DL-Methionine	0.22	0.17
Calcium carbonate	1.05	1.05
Dicalcium Phosphate	1.78	1.5
Salt	0.37	0.37
Wheat middlings	1.00	1.00
Premix ¹	0.6	0.6
<i>Calculated Nutritional Content</i>		
ME (kcal/kg)	3000	3100
Crude Protein (%)	22.25	20.5
Fat (%)	6.76	7.7
Cellulose (%)	3.15	3
Minerals (%)	5.74	5.3
Lysine (%)	1.25	1.13
Methionine (%)	0.55	0.48
Met+Cys (%)	0.92	0.83
Threonine (%)	0.87	0.8
Calcium (%)	0.93	0.85
Total phosphorus	0.7	0.63
Phosphorus Avail. (%)	0.38	0.33

¹The premix will provide the following (per kg of diet): vitamin A, 12 000 IU; vitamin D3, 3 000 IU; vitamin E = 300 IU; vitamin K3, 3 mg; vitamin B, 2 mg; vitamin B2, 8 mg; vitamin B6, 3 mg; vitamin B12, 0.02 mg; folic acid, 1 mg; biotin, 0.2 mg; calcium pantothenate, 15 mg; nicotinic acid, 40 mg; manganese (Mn), 80 mg; zinc (Zn), 60 mg; iodine (I), 1 mg; iron (Fe), 80 mg; copper (Cu), 15 mg; cobalt (Co), 0.4 mg; selenium (Se), 0.2 mg; magnesium (Mg), 5 mg; Etoxyquin, 0.5 mg; BHA, 0.5 mg; citric acid, 5 mg; phosphoric acid, 5 mg.

Experimental results

The results obtained on performance parameters for the whole experimental period are shown in Table 10.3.

5 Table 10.3: Effect of dietary supplementation of the strains DSM 29870, DSM 29871 and DSM 29872 on broiler performance

Parameters ⁴	Control ¹	DSM 29870 ² supplemented group	DSM 29871 ² supplemented group	DSM 29872 ² supplemented group	GalliPro Max ³ supplemented group
FI (g/bird)	3217a	3295a	3281a	3230a	3246a
<i>Relative to the control</i>	-	+2.4%	+2.0%	+0.4%	+0.9%
BWG (g/bird)	2019b	2149a	2167a	2124a	2103ab
<i>Relative to the control</i>	-	+6.5%	+7.3%	+5.2%	+4.2%
FCR	1.598a	1.534b	1.513b	1.521b	1.544b
<i>Relative to the control</i>	-	-4.0%	-5.3%	-4.8%	-3.4%
Mortality (%)	11.1a	9.7a	8.3a	6.5a	7.4a

¹ Group fed with the basal diet.

² DSM 29870/29871/29872 supplemented group fed with the basal diet including strain DSM 29870/29871/29872 at 5.10⁸ CFU/kg of feed.

10 ³ GalliPro Max supplemented group fed with the basal diet including GalliPro Max at 8.10⁸ CFU/kg of feed.

⁴ Means from 12 replicates of 18 birds per pen for each group. Raw data (n=36) were analyzed using variance analysis with block (n=12) and treatments (n=3) as fixed effect.

^{a,b,c} Values with different superscripts are significantly (P<0.05) different from each other.

15

In this experiment, the administration of the strains DSM 29870, DSM 29871, DSM 29872 (as GalliPro Max) had not significant effect on the feed intake, but led to the significant improvement of the body weight gain by 6.5%, 7.3% and 5.2%, respectively. GalliPro Max improved also the body weight gain (+4.2%), but not significantly.

20 All NZB strains improved significantly the feed conversion ratio by 4.0%, 5.3% and 4.8% (DSM 29870, DSM 29871 and DSM 29872, respectively) as GalliPro Max which reached 3.4% of improvement.

Regarding the mortality level, there were no significant differences between the treatments.

EXAMPLE 11: Synthesis and meta-analysis

5 The results of the three performance experiments described above are summarized in Table 11.1.

Table 11.1: Results of the individual studies

Trial	Treatments	FI (g/bird)	Relative to the control	BWG (g/bird)	Relative to the control	FCR	Relative to the control
	- Control	3245a	-	1987ab	-	1.633a	-
	- DSM 29870	3256a	+0.3%	2026ab	+2.0%	1.607b	-1.6%
	- DSM 29871	3221a	-0.7%	2002ab	+0.8%	1.609b	-1.5%
	- DSM 29872	3263a	+0.6%	2038a	+2.6%	1.602b	-2.0%
	- GalliPro Max	3203a	-1.3%	1973b	-0.7%	1.624ab	-0.6%
	- Control	3573a	-	2361a	-	1.513a	-
	- DSM 29870	3542a	-0.9%	2416b	+2.3%	1.467b	-3.6%
	- DSM 29871	3606a	+0.9%	2401ab	+1.7%	1.501bc	-0.8%
	- DSM 29872	3515a	-1.0%	2424b	+2.6%	1.451a	-4.2%
	- GalliPro Max	3541a	-0.9%	2398ab	+1.5%	1.477ab	-2.4%
	- Control	3217a	-	2019b	-	1.598a	-
	- DSM 29870	3295a	+2.4%	2149a	+6.5%	1.534b	-4.0%
	- DSM 29871	3281a	+2.0%	2167a	+7.3%	1.513b	-5.3%
	- DSM 29872	3230a	+0.4%	2124a	+5.2%	1.521b	-4.8%
	- GalliPro Max	3246a	+0.9%	2103ab	+4.2%	1.544b	-3.4%

^{a,b,c} Values with different superscripts are significantly ($P < 0.05$) different from each other.

10

These trials having a similar experimental design, all the data obtained were pooled and combined in a meta-analysis (after having been tested for homogeneity). This meta-analysis involved 3 980 broilers in 160 replicates (32 replicates and 796 animals for each experimental group). The results are shown in Table 11.

15 Data (n = 160) were subjected to a mixed model ANOVA, using the GLIMMIX procedure of SAS (SAS Institute, 2002-2012) to establish differences between diets. Pen

was considered as the experimental unit. The model included diets (n=5) as fixed effect and experiment (n=3) as random effect. Results are reported as least square means. LS means were assumed to be different at $P \leq 0.05$.

5 Table 11.2: meta-analysis of three experiments showing the effect of dietary supplementation of the strain DSM 29870, DSM 29871 and DSM 29872 on broiler performance

Parameters ⁴	Control ¹	DSM 29870 ² supplemented group	DSM 29871 ² supplemented group	DSM 29872 ² supplemented group	GalliPro Max ³ supplemented group
FI (g/bird)	3343a	3360a	3361a	3356a	3347a
<i>Relative to the control</i>	-	+0.5%	+0.5%	+0.4%	+0.1%
BWG (g/bird)	2145b	2208a	2202ab	2211a	2171ab
<i>Relative to the control</i>	-	+2.9%	+2.6%	+3.0%	+1.2%
FCR	1.560a	1.530b	1.53bc	1.52c	1.550ab
<i>Relative to the control</i>	-	-2.4%	-2.2%	-2.7%	-1.1%

¹ Group fed with the basal diet

² DSM 29870/29871/29872 supplemented group fed with the basal diet including strain DSM 29870/29871/29872 at 5.10^8 CFU/kg of feed.

10 ³ GalliPro Max supplemented group fed with the basal diet including GalliPro Max at 8.10^8 CFU/kg of feed.

⁴ Means from 32 observations for each experimental group (796 animals/group).

^{a,b,c} Values with different superscripts are significantly ($P < 0.05$) different from each other.

15 This meta-analysis showed that the use of the strain DSM 29870, DSM 29871 or DSM 29872 in a corn/soybean meal-based diet significantly improved in average the body weight gain (+2.9%, +2.6% and +3.0%, respectively) and the feed conversion ratio (-2.4%, -2.2% and -2.7%, respectively) with no effect on the feed intake.

20 Conclusion

In the working examples above, positive effects of strain DSM 29870 have been demonstrated on broiler performances (*i.e.*, body weight gain and feed conversion ratio) fed a corn/soybean meal-based diet.

All the data obtained from the experiments described in the above examples also showed that the effect of strain DSM 29870 on the feed conversion ratio is due to its effect on the body weight. These effects might be associated with either health effect or metabolism improvement.

5 Strain DSM 29870 showed positive and significant effects on body weight gain and feed conversion ratio in 66% and 100% of the working examples above, respectively.

GalliPro Max showed no significant effect on the body weight gain in any of the working examples above and showed significant and positive effect on feed conversion ratio in one third of the experiments.

10 In the experiments described in the above examples, the better effect of strain DSM 29870 compared to GalliPro Max on broilers performance has been demonstrated. The effect observed using DSM 29870 were in average twice those observed for GalliPro Max.

Example 12: Determination of Monensin Compatibility

15 Monensin compatibility of *Bacillus* strains DSM 29869, DSM 29870, DSM 29871, and DSM 29872 was determined using a modified broth micro dilution similar to the method described in the Example 5. Briefly, a single colony of *Bacillus spp.* (from overnight tryptic soy agar plates) was inoculated into Mueller Hinton Broth (MHB) and cultured overnight. Sterile media was the inoculated with the overnight culture and allowed to grow for 4 hours to
20 test bacteria in log growth phase. Cultures were then diluted once more 1:200 into fresh MHB and 90 µL of this inoculated broth was added to the diluted monensin at the indicated concentrations. Prior art strains were also tested for comparison: NN019785, NN062266 (NRRL B-50013), NN062267 (NRRL B-50104), NN062278 (PTA-6507), NN062319 (FERM BP-1096), NN062440, NN062441 (DSM 17236), NN062439.

25

Strains:

Bacillus amyloliquefaciens DSM 29869

Bacillus subtilis DSM 29870

Bacillus subtilis DSM 29871

30 *Bacillus amyloliquefaciens* DSM 29872

Bacillus licheniformis NN019785

Bacillus amyloliquefaciens NN062266 (NRRL B-50013)

Bacillus subtilis NN062267 (NRRL B-50104)

Bacillus subtilis NN062278 (PTA-6507)

35 *Bacillus amyloliquefaciens* NN062319 (FERM BP-1096)

Bacillus subtilis NN062440

Bacillus licheniformis NN062441 (DSM 17236)

Bacillus amyloliquefaciens NN062439

Materials:

Monensin sodium salt (Sigma, CAS no. 22373-78-0, solubilized in 96% ethanol)

5 Mueller Hinton Broth (Becton, Dickinson and Company, 275730)

Tryptic soy agar (Becton, Dickinson and Company, 236920)

Micro titer plates: Costar plate, polypropylene, flat bottom, Corning, 3628

Borosilicate glass tubes: Kimbale, 16 x 125 mm, 73500-16125

Adhesive gas permeable seals: Thermo Scientific, AB-0718

10

Preparation of bacteria:

Bacillus spp. were grown overnight on tryptic soy agar plates (40 g/L) at 37°C. Mueller Hinton broth (21 g/L) was dissolved in water and autoclaved in glass tubes containing 5 mL of broth each. A single colony of *Bacillus spp.* (from overnight plates) was inoculated into Mueller Hinton Broth (MHB) and incubated overnight at 37°C shaking at 200 rpm. A 5 mL glass tube of fresh, sterile media was then inoculated with 25 mL of overnight culture and allowed to grow for 4 hours at 37°C. Cultures were then diluted once more 1:200 into fresh MHB. 90 µL of this inoculated broth was then added to the diluted antibiotic at the indicated concentrations.

20

Preparation of assay plates:

Monensin was diluted into 96% ethanol to a concentration of 800 µg/mL. This solution was then diluted 10-fold into sterile phosphate buffer to a concentration of 80 µg/mL. A two fold dilution series was prepared in MHB down to the concentration 2.5 µg/mL. 10 µL of each dilution and of each antibiotic was pipetted into a micro titer plate. Later, when the antibiotics were mixed with the suspension of bacteria, the samples were diluted 10x (10µL sample in a total volume of 100 µl). This resulted in the final test range of 0.25-8 µg/ml.

25

90 µl of the bacterial suspensions were added to the assay plates. The assay plates were then covered with an adhesive glass permeable seal and incubated overnight at 37°C shaking at 200 rpm. The maximum compatible concentration was determined similar to a MIC analysis as the concentration above that which inhibited 80% of bacteria as detected by the unaided eye.

30

Results:

A potential challenge of delivering *Bacillus spp.* in feed is the common use of antibiotics as growth promoters in feed. Therefore it is necessary to determine the compatibility of strains with commonly-used feed antibiotics in order to identify any potential

35

conflicts with use as a direct fed microbial. Therefore, the monensin compatibility of *Bacillus* strains DSM 29869, DSM 29870, DSM 29871, and DSM 29872 were determined along with prior art strains. *Bacillus* strains DSM 29869, DSM 29870, DSM 29871, and DSM 29872 indicated a higher level of compatibility with monensin than most of the prior art strains included herein: NN019785, NN062266 (NRRL B-50013), NN062267 (NRRL B-50104), NN062278 (PTA-6507), NN062319 (FERM BP-1096), NN062440, NN062441 (DSM 17236), NN062439.

Table 12.1: Monensin compatibility results

		Species	Product Name	Monensin (µg/mL)
NN062677	DSM29870	<i>Bacillus subtilis</i>		2.7
NN062673	DSM29871	<i>Bacillus subtilis</i>		2.0
NN062683	DSM29872	<i>Bacillus amyloliquefaciens</i>		4.0
NN062676	DSM29869	<i>Bacillus amyloliquefaciens</i>		2.0
NN019785		<i>Bacillus licheniformis</i>	BioPlus 2B (Chr. Hansen)	0.8
NN062266	NRRL B-50013	<i>Bacillus amyloliquefaciens</i>	Eviva Pro (Dupont)	1.1
NN062267	NRRL B-50104	<i>Bacillus subtilis</i>	Eviva Pro (Dupont)	1.1
NN062278	PTA-6507	<i>Bacillus subtilis</i>	Eviva Pro (Dupont)	1.4
NN062319	FERM BP-1096	<i>Bacillus amyloliquefaciens</i>	Calsporin (Calpis)	2.2
NN062440		<i>Bacillus subtilis</i>	GalliPro Max (Chr. Hansen)	0.4
NN062441	DSM 17236	<i>Bacillus licheniformis</i>	GalliPro Tect (Chr. Hansen)	0.9
NN062439		<i>Bacillus amyloliquefaciens</i>	Clostat (Kemin)	2.1

Claims**What is claimed is:**

- 5 1. A *Bacillus* strain characterized in:
- i) having deposit accession number DSM 29870 or a strain having all of the identifying characteristics of *Bacillus* DSM 29870 or a mutant thereof;
 - ii) having deposit accession number DSM 29869 or a strain having all of the identifying characteristics of *Bacillus* DSM 29869 or a mutant thereof;
 - 10 iii) having deposit accession number DSM 29871 or a strain having all of the identifying characteristics of *Bacillus* DSM 29871 or a mutant thereof; or
 - iv) having deposit accession number DSM 29872 or a strain having all of the identifying characteristics of *Bacillus* DSM 29872 or a mutant thereof.
- 15 2. The *Bacillus* strain according to claim 1, wherein the *Bacillus* strain is non-hemolytic.
3. The *Bacillus* strain according to claim 1 or 2, wherein the *Bacillus* strain has antimicrobial activity against *Clostridium perfringens*.
- 20 4. The *Bacillus* strain according to any of the previous claims, wherein the *Bacillus* strain improves body weight gain and/or feed conversion rate in chickens fed with the *Bacillus* strain.
5. The *Bacillus* strain according to any of the previous claims, wherein the *Bacillus* strain comprises 16S rDNA that is more than 98% sequence identity to SEQ ID NO: 1 and/or more than 98% sequence identity to SEQ ID NO: 2 and/or more than 98% sequence identity to SEQ ID NO: 3 and/or more than 98% sequence identity to SEQ ID NO: 4.
- 25 6. The *Bacillus* strain according to any of the previous claims, wherein the *Bacillus* strain is a *Bacillus subtilis* strain or a *Bacillus amyloliquefaciens* strain.
- 30 7. The *Bacillus* strain according to any of the previous claims, wherein the *Bacillus* strain has antimicrobial effect against *E. coli*.
- 35 8. The *Bacillus* strain according to any of the previous claims, wherein the *Bacillus* strain is an isolated *Bacillus* strain.

9. A biologically pure culture of the *Bacillus* strain according to any of claims 1 to 8.
10. A composition comprising spores of the *Bacillus* strain according to any of claims 1-8.
- 5 11. The composition according to claim 10, wherein the *Bacillus* spores of the composition are present as spores such as dried spores.
12. The composition according to claim 11, wherein the bacterial count of each *Bacillus* spore is between 1×10^4 and 1×10^{18} CFU/kg of composition, preferably between 1×10^6 and 1×10^{15} CFU/kg of composition, and more preferably between 1×10^7 and 1×10^{11} CFU/kg of composition.
- 10 13. The composition according to any of claims 10 to 12 which further comprises a carrier.
- 15 14. The composition according to claim 13, wherein the carrier comprises one or more of the following compounds: water, glycerol, ethylene glycol, 1, 2-propylene glycol or 1, 3-propylene glycol, sodium aluminium silicate, sodium chloride, sodium benzoate, potassium sorbate, sodium sulfate, potassium sulfate, magnesium sulfate, sodium thiosulfate, calcium carbonate, sodium citrate, dextrin, maltodextrin, glucose, sucrose, sorbitol, lactose, wheat
- 20 flour, wheat bran, corn gluten meal, starch, farigel, cassava cores, sodium aluminium silicate, colloidal amorphous silica, Sipernat 50S, polyethylene glycol 200, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 1000, polyethylene glycol 1500, polyethylene glycol 4000, carbopol. and cellulose.
- 25 15. The composition according to claim 13 or 14, wherein the carrier comprises calcium carbonate.

Figure 1: Alignment of 16 S rDNA sequences

SEQ ID NO: 1 (DSM 29869), SEQ ID NO: 2 (DSM 29870), SEQ ID NO: 3 (DSM 29871), SEQ ID NO: 4 (DSM 29872), SEQ ID NO: 5 (*Bacillus vallismortis* AB021198), SEQ ID NO: 6 (*Bacillus subtilis* from AJ276351), and SEQ ID NO: 7 (*Bacillus amyloliquefaciens* AB255669)

	90	100	110	120	130	140	150	160	
AJ276351 <i>B. subtilis</i>	GGT	GAG	TAA	CAC	GTG	GGT	AAC	CTG	160
DSM29871	GGT	GAG	TAA	CAC	GTG	GGT	AAC	CTG	160
DSM29870	GGT	GAG	TAA	CAC	GTG	GGT	AAC	CTG	160
AB021198 <i>B. vallismortis</i>	GGT	GAG	TAA	CAC	GTG	GGT	AAC	CTG	160
AB255669 <i>B. amyloliquefaciens</i>	GGT	GAG	TAA	CAC	GTG	GGT	AAC	CTG	160
DSM29869	GGT	GAG	TAA	CAC	GTG	GGT	AAC	CTG	160
DSM29872	GGT	GAG	TAA	CAC	GTG	GGT	AAC	CTG	160
	170	180	190	200	210	220	230	240	
AJ276351 <i>B. subtilis</i>	ACC	GAT	GTT	CAA	ACA	TAA	AGG	TGG	240
DSM29871	ACC	GAT	GTT	CAA	ACA	TAA	AGG	TGG	240
DSM29870	ACC	GAT	GTT	CAA	ACA	TAA	AGG	TGG	240
AB021198 <i>B. vallismortis</i>	ACC	GAT	GTT	CAA	ACA	TAA	AGG	TGG	240
AB255669 <i>B. amyloliquefaciens</i>	ACC	GAT	GTT	CAA	ACA	TAA	AGG	TGG	240
DSM29869	ACC	GAT	GTT	CAA	ACA	TAA	AGG	TGG	240
DSM29872	ACC	GAT	GTT	CAA	ACA	TAA	AGG	TGG	240
	250	260	270	280	290	300	310	320	
AJ276351 <i>B. subtilis</i>	AA	CGG	CT	CA	CAA	GG	CA	AG	320
DSM29871	AA	CGG	CT	CA	CAA	GG	CA	AG	320
DSM29870	AA	CGG	CT	CA	CAA	GG	CA	AG	320
AB021198 <i>B. vallismortis</i>	AA	CGG	CT	CA	CAA	GG	CA	AG	320
AB255669 <i>B. amyloliquefaciens</i>	AA	CGG	CT	CA	CAA	GG	CA	AG	320
DSM29869	AA	CGG	CT	CA	CAA	GG	CA	AG	320
DSM29872	AA	CGG	CT	CA	CAA	GG	CA	AG	320
	330	340	350	360	370	380	390	400	
AJ276351 <i>B. subtilis</i>	CT	A	CGG	A	GG	CA	GCA	GT	400
DSM29871	CT	A	CGG	A	GG	CA	GCA	GT	400
DSM29870	CT	A	CGG	A	GG	CA	GCA	GT	400
AB021198 <i>B. vallismortis</i>	CT	A	CGG	A	GG	CA	GCA	GT	400
AB255669 <i>B. amyloliquefaciens</i>	CT	A	CGG	A	GG	CA	GCA	GT	400
DSM29869	CT	A	CGG	A	GG	CA	GCA	GT	400
DSM29872	CT	A	CGG	A	GG	CA	GCA	GT	400
	410	420	430	440	450	460	470	480	
AJ276351 <i>B. subtilis</i>	CG	GAT	CGT	AAA	GCT	CT	GTT	TTA	480
DSM29871	CG	GAT	CGT	AAA	GCT	CT	GTT	TTA	480
DSM29870	CG	GAT	CGT	AAA	GCT	CT	GTT	TTA	480
AB021198 <i>B. vallismortis</i>	CG	GAT	CGT	AAA	GCT	CT	GTT	TTA	480
AB255669 <i>B. amyloliquefaciens</i>	CG	GAT	CGT	AAA	GCT	CT	GTT	TTA	480
DSM29869	CG	GAT	CGT	AAA	GCT	CT	GTT	TTA	480
DSM29872	CG	GAT	CGT	AAA	GCT	CT	GTT	TTA	480
	1210	1220	1230	1240	1250	1260	1270	1280	
AJ276351 <i>B. subtilis</i>	GG	CTA	CAC	AGT	GCT	ACA	AT	GGA	1280
DSM29871	GG	CTA	CAC	AGT	GCT	ACA	AT	GGA	1280
DSM29870	GG	CTA	CAC	AGT	GCT	ACA	AT	GGA	1280
AB021198 <i>B. vallismortis</i>	GG	CTA	CAC	AGT	GCT	ACA	AT	GGA	1280
AB255669 <i>B. amyloliquefaciens</i>	GG	CTA	CAC	AGT	GCT	ACA	AT	GGA	1280
DSM29869	GG	CTA	CAC	AGT	GCT	ACA	AT	GGA	1280
DSM29872	GG	CTA	CAC	AGT	GCT	ACA	AT	GGA	1280
	1290	1300	1310	1320	1330	1340	1350	1360	
AJ276351 <i>B. subtilis</i>	TT	CGG	AT	CG	AGT	CT	GCA	ACT	1360
DSM29871	TT	CGG	AT	CG	AGT	CT	GCA	ACT	1360
DSM29870	TT	CGG	AT	CG	AGT	CT	GCA	ACT	1360
AB021198 <i>B. vallismortis</i>	TT	CGG	AT	CG	AGT	CT	GCA	ACT	1360
AB255669 <i>B. amyloliquefaciens</i>	TT	CGG	AT	CG	AGT	CT	GCA	ACT	1360
DSM29869	TT	CGG	AT	CG	AGT	CT	GCA	ACT	1360
DSM29872	TT	CGG	AT	CG	AGT	CT	GCA	ACT	1360
	1370	1380	1390	1400	1410	1420	1430	1440	
AJ276351 <i>B. subtilis</i>	T	CCC	GGG	CC	T	G	T	A	1440
DSM29871	T	CCC	GGG	CC	T	G	T	A	1440
DSM29870	T	CCC	GGG	CC	T	G	T	A	1440
AB021198 <i>B. vallismortis</i>	T	CCC	GGG	CC	T	G	T	A	1440
AB255669 <i>B. amyloliquefaciens</i>	T	CCC	GGG	CC	T	G	T	A	1440
DSM29869	T	CCC	GGG	CC	T	G	T	A	1440
DSM29872	T	CCC	GGG	CC	T	G	T	A	1440

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/014524

A. CLASSIFICATION OF SUBJECT MATTER
INV. C12N1/20 A61K35/66
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2007/064741 A2 (KEMIN IND INC [US]; VARGHESE JERRY [IN]; SAS BENEDIKT [BE]; VAN HEMEL) 7 June 2007 (2007-06-07) cited in the application page 8, line 26 - page 9, line 25; figure 1; example 2 ----- -/--	1-4, 6-10,13, 14



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

25 April 2016

Date of mailing of the international search report

04/05/2016

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Stoyanov, Borislav

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/014524

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	S. JAYARAMAN ET AL: "Bacillus subtilis PB6 improves intestinal health of broiler chickens challenged with Clostridium perfringens-induced necrotic enteritis", POULTRY SCIENCE, vol. 92, no. 2, 1 February 2013 (2013-02-01), pages 370-374, XP055237622, US ISSN: 0032-5791, DOI: 10.3382/ps.2012-02528 page 371, left-hand column, paragraph 3 - page 371, right-hand column, paragraph 1 -----	1-4,6-10
X	A. Y.- L. TEO ET AL: "Effect of Bacillus subtilis PB6 (CloSTAT) on Broilers Infected with a Pathogenic Strain of Escherichia coli", JOURNAL OF APPLIED POULTRY RESEARCH, vol. 15, no. 2, 1 June 2006 (2006-06-01), pages 229-235, XP055268047, US ISSN: 1056-6171, DOI: 10.1093/japr/15.2.229 tables 3-4 -----	1-4,6-10
X	B. ESSGHAIER ET AL: "Biological control of grey mould in strawberry fruits by halophilic bacteria", JOURNAL OF APPLIED MICROBIOLOGY., vol. 106, no. 3, 1 March 2009 (2009-03-01), pages 833-846, XP055225876, GB ISSN: 1364-5072, DOI: 10.1111/j.1365-2672.2008.04053.x B. subtilis strain B1-33 16S RNA-gene 100% identical to present SEQ ID NO.:2; figure 1 -----	1-9
X	CN 103 882 097 A (QINGDAO ZHONGREN PHARMACEUTICAL CO LTD) 25 June 2014 (2014-06-25) page 1 - page 4; claims 1-2 -----	1-9

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2016/014524

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-15

A Bacillus strain characterized in:

i) having deposit accession number DSM 29870 and compositions comprising it.

1.1. claims: 1-15(partially)

A Bacillus strain characterized in:

i) having deposit accession number DSM 29869 and compositions comprising it.

1.2. claims: 1-15(partially)

A Bacillus strain characterized in:

i) having deposit accession number DSM 29871 and compositions comprising it.

1.3. claims: 1-15(partially)

A Bacillus strain characterized in:

i) having deposit accession number DSM 29872 and compositions comprising it.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2016/014524

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2007064741	A2	07-06-2007	
		AU 2006320613 A1	07-06-2007
		BR PI0619364 A2	27-09-2011
		CA 2631289 A1	07-06-2007
		CN 101448398 A	03-06-2009
		EP 1956914 A2	20-08-2008
		IL 191795 A	31-07-2013
		JP 5280854 B2	04-09-2013
		JP 2009519238 A	14-05-2009
		JP 2013173750 A	05-09-2013
		KR 20080073771 A	11-08-2008
		NZ 569244 A	27-07-2012
		US 2008057047 A1	06-03-2008
		WO 2007064741 A2	07-06-2007

CN 103882097	A	25-06-2014	NONE
