

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 October 2006 (26.10.2006)

PCT

(10) International Publication Number
WO 2006/111759 A1

(51) International Patent Classification:
C07D 519/00 (2006.01) A61K 31/5517 (2006.01)
A61P 35/00 (2006.01)

(74) Agents: WATSON, Robert et al.; Mewburn Ellis LLP,
York House, 23 Kingsway, London, Greater London
WC2B 6HP (GB).

(21) International Application Number:
PCT/GB2006/001456

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV,
LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI,
NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG,
SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US,
UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 21 April 2006 (21.04.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0508084.1 21 April 2005 (21.04.2005) GB
0522746.7 7 November 2005 (07.11.2005) GB

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US):
SPIROGEN LIMITED [GB/GB]; 79 George Street,
Ryde, Hampshire PO33 2JF (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GREGSON,
Stephen, John [GB/GB]; School of Pharmacy, University
of London, 29/39 Brunswick Square, London, Greater
London WC1N 1AX (GB). HOWARD, Philip, Wilson
[GB/GB]; School of Pharmacy, University of London,
29/39 Brunswick Square, London, Greater London WC1N
1AX (GB). CHEN, Zhizhi [CN/GB]; School of Pharmacy,
University of London, 29/39 Brunswick Square, London,
Greater London WC1N 1AX (GB).

Declaration under Rule 4.17:


— of inventorship (Rule 4.17(iv))

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PYRROLOBENZODIAZEPINES



(57) Abstract: Compounds of the formula: (I) or solvate thereof, wherein: R² is an optionally substituted C₅₋₂₀ aryl group; R⁶ and R⁹ are independently selected from H, R, OH, OR, SH, SR, NH₂, NHR, NRR', nitro, Me₃Sn and halo; where R and R' are independently selected from optionally substituted C₁₋₁₂ alkyl, C₃₋₂₀ heterocyclyl and C₅₋₂₀ aryl groups; R⁷ is selected from H, R, OH, OR, SH, SR, NH₂, NHR, NHR', nitro, Me₃Sn and halo; R'' is a C₃₋₁₂ alkylene group, which chain may be interrupted by one or more heteroatoms and/or aromatic rings; X is selected from O, S, or NH; z is 2 or 3; M is a monovalent pharmaceutically acceptable cation; R², R⁶, R⁷, R⁹, X' and M' are selected from the same groups as R², R⁶, R⁷, R⁹, X and M respectively, or M and M' may together represent a divalent pharmaceutically acceptable cation.

WO 2006/111759 A1


PYRROLOBENZODIAZEPINES

The present invention relates to pyrrolobenzodiazepines (PBDs), and in particular pyrrolobenzodiazepine dimers bearing C2 aryl substitutions.

5

Background to the invention

Some pyrrolobenzodiazepines (PBDs) have the ability to recognise and bond to specific sequences of DNA; the preferred sequence is PuGpu. The first PBD antitumour antibiotic, anthramycin, was discovered in 1965 (Leimgruber, *et al.*, *J. Am. Chem. Soc.*, **87**, 5793-5795 (1965); Leimgruber, *et al.*, *J. Am. Chem. Soc.*, **87**, 5791-5793 (1965)). Since then, a number of naturally occurring PBDs have been reported, and over 10 synthetic routes have been developed to a variety of analogues (Thurston, *et al.*, *Chem. Rev.* **1994**, 433-465 (1994)). Family members include abbeymycin (Hochlowski, *et al.*, *J. Antibiotics*, **40**, 145-148 (1987)), chicamycin (Konishi, *et al.*, *J. Antibiotics*, **37**, 200-206 (1984)), DC-81 (Japanese Patent 58-180 487; Thurston, *et al.*, *Chem. Brit.*, **26**, 767-772 (1990); Bose, *et al.*, *Tetrahedron*, **48**, 751-758 (1992)), mazethramycin (Kuminoto, *et al.*, *J. Antibiotics*, **33**, 665-667 (1980)), neothramycins A and B (Takeuchi, *et al.*, *J. Antibiotics*, **29**, 93-96 (1976)), porothramycin (Tsunakawa, *et al.*, *J. Antibiotics*, **41**, 1366-1373 (1988)), prothracarcin (Shimizu, *et al.*, *J. Antibiotics*, **29**, 2492-2503 (1982); Langley and Thurston, *J. Org. Chem.*, **52**, 91-97 (1987)), sibanomicin (DC-102)(Hara, *et al.*, *J. Antibiotics*, **41**, 702-704 (1988); Itoh, *et al.*, *J. Antibiotics*, **41**, 1281-1284 (1988)), sibiromycin (Leber, *et al.*, *J. Am. Chem. Soc.*, **110**, 2992-2993 (1988)) and tomamycin (Arima, *et al.*, *J. Antibiotics*, **25**, 437-444 (1972)). PBDs are of the general structure:




They differ in the number, type and position of substituents, in both their aromatic A rings and pyrrolo C rings, and in the degree of saturation of the C ring. In the B-ring there is either an imine (N=C), a carbinolamine (NH-CH(OH)), or a carbinolamine methyl ether (NH-CH(OMe)) at the N10-C11 position which is the electrophilic centre responsible for alkylating DNA. All of the known natural products have an (S)-configuration at the chiral C11a position which provides them with a right-handed twist when viewed from the C ring towards the A ring. This gives them the appropriate three-dimensional shape for isohelicity


30

with the minor groove of B-form DNA, leading to a snug fit at the binding site (Kohn, In *Antibiotics III*. Springer-Verlag, New York, pp. 3-11 (1975); Hurley and Needham-VanDevanter, *Acc. Chem. Res.*, **19**, 230-237 (1986)). Their ability to form an adduct in the minor groove, enables them to interfere with DNA processing, hence their use as antitumour agents.

The present inventors have previously disclosed, in WO 2004/043963, cytotoxic compounds having an aryl group at the C2 position, for example:




The present inventors have also previously disclosed, in co-pending PCT application PCT/GB2005/000768 (published as WO 2005/085251), dimeric PBD compounds bearing C2 aryl substituents, such as:



Disclosure of the invention

The present inventors have encountered some issues with the solubility of compounds such as ZC-207, which they have resolved by the use of a different form of these compounds.

The present invention comprises a compound with the formula I:



or solvate thereof, wherein:

R² is an optionally substituted C₅₋₂₀ aryl group;

R⁶ and R⁹ are independently selected from H, R, OH, OR, SH, SR, NH₂, NHR, NRR', nitro, Me₃Sn and halo;

where R and R' are independently selected from optionally substituted C₁₋₁₂ alkyl, C₃₋₂₀ heterocyclyl and C₅₋₂₀ aryl groups;

5 R⁷ is selected from H, R, OH, OR, SH, SR, NH₂, NHR, NHRR', nitro, Me₃Sn and halo;
R'' is a C₃₋₁₂ alkylene group, which chain may be interrupted by one or more heteroatoms, e.g. O, S, NH, and/or aromatic rings, e.g. benzene or pyridine;

X is selected from O, S, or NH;

z is 2 or 3;

10 M is a monovalent pharmaceutically acceptable cation;

R², R⁶, R⁷, R⁹, X' and M' are selected from the same groups as R², R⁶, R⁷, R⁹, X and M respectively, or M and M' may together represent a divalent pharmaceutically acceptable cation.

15 Pyrrolobenzodiazepines having an imine bond are known to convert to the di-carbinolamine form in water, and isolated pyrrolobenzodiazepines often exist as a mixture of the imine, mono-carbinolamine and di-carbinolamine forms. Furthermore, if the compound is isolated as a solid with a mixture of these three forms, the balance between them may change over time. Although this does not pose a problem for administration of
20 the compound, it can provide difficulties in accurately assessing the amount of active substance in a given amount of powder. Compounds of the present invention, at least to some extent, overcome this difficulty whilst remaining active, and are thus suited to formulation as pharmaceuticals.

25 Dimeric pyrrolobenzodiazepines offer advantages over monomeric pyrrolobenzodiazepines in that they possess the ability to cross-link DNA in the minor groove, which can lead to an increase in cytotoxicity.

30 Further aspects of the invention relate to the compounds use in methods of therapy (particularly in treating proliferative diseases), pharmaceutical compositions comprising the compounds, and their use in the manufacture of a medicament for the treatment of a proliferative disease.

Definitions

Substituents

The phrase "optionally substituted" as used herein, pertains to a parent group which may be unsubstituted or which may be substituted.

5

Unless otherwise specified, the term "substituted" as used herein, pertains to a parent group which bears one or more substituents. The term "substituent" is used herein in the conventional sense and refers to a chemical moiety which is covalently attached to, or if appropriate, fused to, a parent group. A wide variety of substituents are well known, and methods for their formation and introduction into a variety of parent groups are also well known.

10

Examples of substituents are described in more detail below.

15

C_{1-12} alkyl: The term " C_{1-12} alkyl" as used herein, pertains to a monovalent moiety obtained by removing a hydrogen atom from a carbon atom of a hydrocarbon compound having from 1 to 12 carbon atoms, which may be aliphatic or alicyclic, and which may be saturated or unsaturated (e.g. partially unsaturated, fully unsaturated). Thus, the term "alkyl" includes the sub-classes alkenyl, alkynyl, cycloalkyl, etc., discussed below.

20

Examples of saturated alkyl groups include, but are not limited to, methyl (C_1), ethyl (C_2), propyl (C_3), butyl (C_4), pentyl (C_5), hexyl (C_6) and heptyl (C_7).

25

Examples of saturated linear alkyl groups include, but are not limited to, methyl (C_1), ethyl (C_2), n-propyl (C_3), n-butyl (C_4), n-pentyl (amyl) (C_5), n-hexyl (C_6) and n-heptyl (C_7).

30

C_{2-12} Alkenyl: The term " C_{2-12} alkenyl" as used herein, pertains to an alkyl group having one or more carbon-carbon double bonds.

35

Examples of unsaturated alkenyl groups include, but are not limited to, ethenyl (vinyl, $-CH=CH_2$), 1-propenyl ($-CH=CH-CH_3$), 2-propenyl (allyl, $-CH-CH=CH_2$), isopropenyl (1-methylvinyl, $-C(CH_3)=CH_2$), butenyl (C_4), pentenyl (C_5), and hexenyl (C_6).

C₂₋₁₂ alkynyl: The term "C₂₋₁₂ alkynyl" as used herein, pertains to an alkyl group having one or more carbon-carbon triple bonds.

5 Examples of unsaturated alkynyl groups include, but are not limited to, ethynyl (ethynyl, -C≡CH) and 2-propynyl (propargyl, -CH₂-C≡CH).

C₃₋₁₂ cycloalkyl: The term "C₃₋₁₂ cycloalkyl" as used herein, pertains to an alkyl group which is also a cyclyl group; that is, a monovalent moiety obtained by removing a hydrogen atom
10 from an alicyclic ring atom of a cyclic hydrocarbon (carbocyclic) compound, which moiety has from 3 to 7 carbon atoms, including from 3 to 7 ring atoms.

Examples of cycloalkyl groups include, but are not limited to, those derived from:

saturated monocyclic hydrocarbon compounds:

15 cyclopropane (C₃), cyclobutane (C₄), cyclopentane (C₅), cyclohexane (C₆), cycloheptane (C₇), methylcyclopropane (C₄), dimethylcyclopropane (C₅), methylcyclobutane (C₅), dimethylcyclobutane (C₆), methylcyclopentane (C₆), dimethylcyclopentane (C₇) and methylcyclohexane (C₇);

unsaturated monocyclic hydrocarbon compounds:

20 cyclopropene (C₃), cyclobutene (C₄), cyclopentene (C₅), cyclohexene (C₆), methylcyclopropene (C₄), dimethylcyclopropene (C₅), methylcyclobutene (C₅), dimethylcyclobutene (C₆), methylcyclopentene (C₆), dimethylcyclopentene (C₇) and methylcyclohexene (C₇); and

saturated polycyclic hydrocarbon compounds:

25 norcarane (C₇), norpinane (C₇), norbornane (C₇).

C₃₋₂₀ heterocyclyl: The term "C₃₋₂₀ heterocyclyl" as used herein, pertains to a monovalent moiety obtained by removing a hydrogen atom from a ring atom of a heterocyclic compound, which moiety has from 3 to 20 ring atoms, of which from 1 to 10 are ring
30 heteroatoms. Preferably, each ring has from 3 to 7 ring atoms, of which from 1 to 4 are ring heteroatoms.

In this context, the prefixes (e.g. C₃₋₂₀, C₃₋₇, C₅₋₆, etc.) denote the number of ring atoms, or range of number of ring atoms, whether carbon atoms or heteroatoms. For example, the

term "C₅₋₆heterocyclyl", as used herein, pertains to a heterocyclyl group having 5 or 6 ring atoms.

Examples of monocyclic heterocyclyl groups include, but are not limited to, those derived from:

N₁: aziridine (C₃), azetidine (C₄), pyrrolidine (tetrahydropyrrole) (C₅), pyrroline (e.g., 3-pyrroline, 2,5-dihydropyrrole) (C₅), 2H-pyrrole or 3H-pyrrole (isopyrrole, isoazole) (C₅), piperidine (C₆), dihydropyridine (C₆), tetrahydropyridine (C₆), azepine (C₇);

O₁: oxirane (C₃), oxetane (C₄), oxolane (tetrahydrofuran) (C₅), oxole (dihydrofuran) (C₅), oxane (tetrahydropyran) (C₆), dihydropyran (C₆), pyran (C₆), oxepin (C₇);

S₁: thiirane (C₃), thietane (C₄), thiolane (tetrahydrothiophene) (C₅), thiane (tetrahydrothiopyran) (C₆), thiepane (C₇);

O₂: dioxolane (C₅), dioxane (C₆), and dioxepane (C₇);

O₃: trioxane (C₆);

N₂: imidazolidine (C₅), pyrazolidine (diazolidine) (C₅), imidazoline (C₅), pyrazoline (dihydropyrazole) (C₅), piperazine (C₆);

N₁O₁: tetrahydrooxazole (C₅), dihydrooxazole (C₅), tetrahydroisoxazole (C₅), dihydroisoxazole (C₅), morpholine (C₆), tetrahydrooxazine (C₆), dihydrooxazine (C₆), oxazine (C₆);

N₁S₁: thiazoline (C₅), thiazolidine (C₅), thiomorpholine (C₆);

N₂O₁: oxadiazine (C₆);

O₁S₁: oxathiole (C₅) and oxathiane (thioxane) (C₆); and,

N₁O₁S₁: oxathiazine (C₆).

Examples of substituted monocyclic heterocyclyl groups include those derived from saccharides, in cyclic form, for example, furanoses (C₅), such as arabinofuranose, lyxofuranose, ribofuranose, and xylofuranse, and pyranoses (C₆), such as allopyranose, altropyranose, glucopyranose, mannopyranose, gulopyranose, idopyranose, galactopyranose, and talopyranose.

C₅₋₂₀ aryl: The term "C₅₋₂₀ aryl", as used herein, pertains to a monovalent moiety obtained by removing a hydrogen atom from an aromatic ring atom of an aromatic compound, which moiety has from 3 to 20 ring atoms. Preferably, each ring has from 5 to 7 ring atoms.

In this context, the prefixes (e.g. C₃₋₂₀, C₅₋₇, C₅₋₆, etc.) denote the number of ring atoms, or range of number of ring atoms, whether carbon atoms or heteroatoms. For example, the term "C₅₋₆ aryl" as used herein, pertains to an aryl group having 5 or 6 ring atoms.

5 The ring atoms may be all carbon atoms, as in "carboaryl groups".

Examples of carboaryl groups include, but are not limited to, those derived from benzene (i.e. phenyl) (C₆), naphthalene (C₁₀), azulene (C₁₀), anthracene (C₁₄), phenanthrene (C₁₄), naphthacene (C₁₈), and pyrene (C₁₆).

10 Examples of aryl groups which comprise fused rings, at least one of which is an aromatic ring, include, but are not limited to, groups derived from indane (e.g. 2,3-dihydro-1H-indene) (C₉), indene (C₉), isoindene (C₉), tetraline (1,2,3,4-tetrahydronaphthalene (C₁₀), acenaphthene (C₁₂), fluorene (C₁₃), phenalene (C₁₃), acephenanthrene (C₁₅), and aceanthrene (C₁₆).

15 Alternatively, the ring atoms may include one or more heteroatoms, as in "heteroaryl groups". Examples of monocyclic heteroaryl groups include, but are not limited to, those derived from:

N₁: pyrrole (azole) (C₅), pyridine (azine) (C₆);

20 O₁: furan (oxole) (C₅);

S₁: thiophene (thiole) (C₅);

N₁O₁: oxazole (C₅), isoxazole (C₅), isoxazine (C₆);

N₂O₁: oxadiazole (furazan) (C₅);

N₃O₁: oxatriazole (C₅);

25 N₁S₁: thiazole (C₅), isothiazole (C₅);

N₂: imidazole (1,3-diazole) (C₅), pyrazole (1,2-diazole) (C₅), pyridazine (1,2-diazine) (C₆), pyrimidine (1,3-diazine) (C₆) (e.g., cytosine, thymine, uracil), pyrazine (1,4-diazine) (C₆);

N₃: triazole (C₅), triazine (C₆); and,

N₄: tetrazole (C₅).

30

Examples of heteroaryl which comprise fused rings, include, but are not limited to:

C₉ (with 2 fused rings) derived from benzofuran (O₁), isobenzofuran (O₁), indole (N₁), isoindole (N₁), indolizine (N₁), indoline (N₁), isoindoline (N₁), purine (N₄) (e.g., adenine, guanine), benzimidazole (N₂), indazole (N₂), benzoxazole (N₁O₁), benzisoxazole (N₁O₁),

benzodioxole (O₂), benzofurazan (N₂O₁), benzotriazole (N₃), benzothiofuran (S₁),
benzothiazole (N₁S₁), benzothiadiazole (N₂S);

C₁₀ (with 2 fused rings) derived from chromene (O₁), isochromene (O₁), chroman
(O₁), isochroman (O₁), benzodioxan (O₂), quinoline (N₁), isoquinoline (N₁), quinolizine (N₁),
5 benzoxazine (N₁O₁), benzodiazine (N₂), pyridopyridine (N₂), quinoxaline (N₂), quinazoline
(N₂), cinnoline (N₂), phthalazine (N₂), naphthyridine (N₂), pteridine (N₄);

C₁₁ (with 2 fused rings) derived from benzodiazepine (N₂);

C₁₃ (with 3 fused rings) derived from carbazole (N₁), dibenzofuran (O₁),
dibenzothiophene (S₁), carboline (N₂), perimidine (N₂), pyridoindole (N₂); and,

10 C₁₄ (with 3 fused rings) derived from acridine (N₁), xanthene (O₁), thioxanthene (S₁),
oxanthrene (O₂), phenoxathiin (O₁S₁), phenazine (N₂), phenoxazine (N₁O₁), phenothiazine
(N₁S₁), thianthrene (S₂), phenanthridine (N₁), phenanthroline (N₂), phenazine (N₂).

The above groups, whether alone or part of another substituent, may themselves optionally
15 be substituted with one or more groups selected from themselves and the additional
substituents listed below.

Halo: -F, -Cl, -Br, and -I.

20 Hydroxy: -OH.

Ether: -OR, wherein R is an ether substituent, for example, a C₁₋₇ alkyl group (also referred
to as a C₁₋₇ alkoxy group, discussed below), a C₃₋₂₀ heterocyclyl group (also referred to as a
C₃₋₂₀ heterocycloxy group), or a C₅₋₂₀ aryl group (also referred to as a C₅₋₂₀ aryloxy group),
25 preferably a C₁₋₇alkyl group.

Alkoxy: -OR, wherein R is an alkyl group, for example, a C₁₋₇ alkyl group. Examples of C₁₋₇
alkoxy groups include, but are not limited to, -OMe (methoxy), -OEt (ethoxy), -O(nPr) (n-
propoxy), -O(iPr) (isopropoxy), -O(nBu) (n-butoxy), -O(sBu) (sec-butoxy), -O(iBu)
30 (isobutoxy), and -O(tBu) (tert-butoxy).

Acetal: -CH(OR¹)(OR²), wherein R¹ and R² are independently acetal substituents, for
example, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably a
C₁₋₇ alkyl group, or, in the case of a "cyclic" acetal group, R¹ and R², taken together with the
35 two oxygen atoms to which they are attached, and the carbon atoms to which they are

attached, form a heterocyclic ring having from 4 to 8 ring atoms. Examples of acetal groups include, but are not limited to, $-\text{CH}(\text{OMe})_2$, $-\text{CH}(\text{OEt})_2$, and $-\text{CH}(\text{OMe})(\text{OEt})$.

Hemiacetal: $-\text{CH}(\text{OH})(\text{OR}^1)$, wherein R^1 is a hemiacetal substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of hemiacetal groups include, but are not limited to, $-\text{CH}(\text{OH})(\text{OMe})$ and $-\text{CH}(\text{OH})(\text{OEt})$.

Ketal: $-\text{CR}(\text{OR}^1)(\text{OR}^2)$, where R^1 and R^2 are as defined for acetals, and R is a ketal substituent other than hydrogen, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples ketal groups include, but are not limited to, $-\text{C}(\text{Me})(\text{OMe})_2$, $-\text{C}(\text{Me})(\text{OEt})_2$, $-\text{C}(\text{Me})(\text{OMe})(\text{OEt})$, $-\text{C}(\text{Et})(\text{OMe})_2$, $-\text{C}(\text{Et})(\text{OEt})_2$, and $-\text{C}(\text{Et})(\text{OMe})(\text{OEt})$.

Hemiketal: $-\text{CR}(\text{OH})(\text{OR}^1)$, where R^1 is as defined for hemiacetals, and R is a hemiketal substituent other than hydrogen, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of hemiacetal groups include, but are not limited to, $-\text{C}(\text{Me})(\text{OH})(\text{OMe})$, $-\text{C}(\text{Et})(\text{OH})(\text{OMe})$, $-\text{C}(\text{Me})(\text{OH})(\text{OEt})$, and $-\text{C}(\text{Et})(\text{OH})(\text{OEt})$.

Oxo (keto, -one): $=\text{O}$.

Thione (thioketone): $=\text{S}$.

Imino (imine): $=\text{NR}$, wherein R is an imino substituent, for example, hydrogen, C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably hydrogen or a C_{1-7} alkyl group. Examples of ester groups include, but are not limited to, $=\text{NH}$, $=\text{NMe}$, $=\text{NEt}$, and $=\text{NPh}$.

Formyl (carbaldehyde, carboxaldehyde): $-\text{C}(=\text{O})\text{H}$.

Acyl (keto): $-\text{C}(=\text{O})\text{R}$, wherein R is an acyl substituent, for example, a C_{1-7} alkyl group (also referred to as C_{1-7} alkylacyl or C_{1-7} alkanoyl), a C_{3-20} heterocyclyl group (also referred to as C_{3-20} heterocyclylacyl), or a C_{5-20} aryl group (also referred to as C_{5-20} arylacyl), preferably a C_{1-7} alkyl group. Examples of acyl groups include, but are not limited to, $-\text{C}(=\text{O})\text{CH}_3$

(acetyl), $-\text{C}(=\text{O})\text{CH}_2\text{CH}_3$ (propionyl), $-\text{C}(=\text{O})\text{C}(\text{CH}_3)_3$ (t-butyryl), and $-\text{C}(=\text{O})\text{Ph}$ (benzoyl, phenone).

Carboxy (carboxylic acid): $-\text{C}(=\text{O})\text{OH}$.

5

Thiocarboxy (thiocarboxylic acid): $-\text{C}(=\text{S})\text{SH}$.

Thiolocarboxy (thiolocarboxylic acid): $-\text{C}(=\text{O})\text{SH}$.

10 Thionocarboxy (thionocarboxylic acid): $-\text{C}(=\text{S})\text{OH}$.

Imidic acid: $-\text{C}(=\text{NH})\text{OH}$.

Hydroxamic acid: $-\text{C}(=\text{NOH})\text{OH}$.

15

Ester (carboxylate, carboxylic acid ester, oxycarbonyl): $-\text{C}(=\text{O})\text{OR}$, wherein R is an ester substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of ester groups include, but are not limited to, $-\text{C}(=\text{O})\text{OCH}_3$, $-\text{C}(=\text{O})\text{OCH}_2\text{CH}_3$, $-\text{C}(=\text{O})\text{OC}(\text{CH}_3)_3$, and $-\text{C}(=\text{O})\text{OPh}$.

20

Acyloxy (reverse ester): $-\text{OC}(=\text{O})\text{R}$, wherein R is an acyloxy substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of acyloxy groups include, but are not limited to, $-\text{OC}(=\text{O})\text{CH}_3$ (acetoxo), $-\text{OC}(=\text{O})\text{CH}_2\text{CH}_3$, $-\text{OC}(=\text{O})\text{C}(\text{CH}_3)_3$, $-\text{OC}(=\text{O})\text{Ph}$, and $-\text{OC}(=\text{O})\text{CH}_2\text{Ph}$.

25

Oxycarboxyloxy: $-\text{OC}(=\text{O})\text{OR}$, wherein R is an ester substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of ester groups include, but are not limited to, $-\text{OC}(=\text{O})\text{OCH}_3$, $-\text{OC}(=\text{O})\text{OCH}_2\text{CH}_3$, $-\text{OC}(=\text{O})\text{OC}(\text{CH}_3)_3$, and $-\text{OC}(=\text{O})\text{OPh}$.

30

Amino: $-\text{NR}^1\text{R}^2$, wherein R^1 and R^2 are independently amino substituents, for example, hydrogen, a C_{1-7} alkyl group (also referred to as C_{1-7} alkylamino or di- C_{1-7} alkylamino), a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably H or a C_{1-7} alkyl group, or, in the case of a "cyclic" amino group, R^1 and R^2 , taken together with the nitrogen atom to which they are attached, form a heterocyclic ring having from 4 to 8 ring atoms. Amino groups


35

may be primary (-NH₂), secondary (-NHR¹), or tertiary (-NHR¹R²), and in cationic form, may be quaternary (-⁺NR¹R²R³). Examples of amino groups include, but are not limited to, -NH₂, -NHCH₃, -NHC(CH₃)₂, -N(CH₃)₂, -N(CH₂CH₃)₂, and -NHPh. Examples of cyclic amino groups include, but are not limited to, aziridino, azetidino, pyrrolidino, piperidino,
 5 piperazino, morpholino, and thiomorpholino.


Amido (carbamoyl, carbamyl, aminocarbonyl, carboxamide): -C(=O)NR¹R², wherein R¹ and R² are independently amino substituents, as defined for amino groups. Examples of amido groups include, but are not limited to, -C(=O)NH₂, -C(=O)NHCH₃, -C(=O)N(CH₃)₂,
 10 -C(=O)NHCH₂CH₃, and -C(=O)N(CH₂CH₃)₂, as well as amido groups in which R¹ and R², together with the nitrogen atom to which they are attached, form a heterocyclic structure as in, for example, piperidinocarbonyl, morpholinocarbonyl, thiomorpholinocarbonyl, and piperazinocarbonyl.

Thioamido (thiocarbamyl): -C(=S)NR¹R², wherein R¹ and R² are independently amino substituents, as defined for amino groups. Examples of amido groups include, but are not limited to, -C(=S)NH₂, -C(=S)NHCH₃, -C(=S)N(CH₃)₂, and -C(=S)NHCH₂CH₃.


Acylamido (acylamino): -NR¹C(=O)R², wherein R¹ is an amide substituent, for example, hydrogen, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably hydrogen or a C₁₋₇ alkyl group, and R² is an acyl substituent, for example, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably hydrogen or a C₁₋₇ alkyl group. Examples of acylamide groups include, but are not limited to, -NHC(=O)CH₃,
 20 -NHC(=O)CH₂CH₃, and -NHC(=O)Ph. R¹ and R² may together form a cyclic structure, as in, for example, succinimidyl, maleimidyl, and phthalimidyl:
 25



succinimidyl



maleimidyl



phthalimidyl


Aminocarbonyloxy: -OC(=O)NR¹R², wherein R¹ and R² are independently amino substituents, as defined for amino groups. Examples of aminocarbonyloxy groups include,
 30 but are not limited to, -OC(=O)NH₂, -OC(=O)NHMe, -OC(=O)NMe₂, and -OC(=O)NEt₂.

Ureido: $-N(R^1)CONR^2R^3$ wherein R^2 and R^3 are independently amino substituents, as defined for amino groups, and R^1 is a ureido substituent, for example, hydrogen, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably hydrogen or a C_{1-7} alkyl group. Examples of ureido groups include, but are not limited to, $-NHCONH_2$, -

5 $NHCONHMe$, $-NHCONHEt$, $-NHCONMe_2$, $-NHCONEt_2$, $-NMeCONH_2$, $-NMeCONHMe$, $-NMeCONHEt$, $-NMeCONMe_2$, and $-NMeCONEt_2$.

Guanidino: $-NH-C(=NH)NH_2$.

10 Tetrazolyl: a five membered aromatic ring having four nitrogen atoms and one carbon atom,



15 Imino: $=NR$, wherein R is an imino substituent, for example, hydrogen, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably H or a C_{1-7} alkyl group. Examples of imino groups include, but are not limited to, $=NH$, $=NMe$, and $=NEt$.

Amidine (amidino): $-C(=NR)NR_2$, wherein each R is an amidine substituent, for example, hydrogen, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably H or a C_{1-7} alkyl group. Examples of amidine groups include, but are not limited to,

20 $-C(=NH)NH_2$, $-C(=NH)NMe_2$, and $-C(=NMe)NMe_2$.

Nitro: $-NO_2$.

25 Nitroso: $-NO$.

Azido: $-N_3$.

Cyano (nitrile, carbonitrile): $-CN$.

30

Isocyano: $-NC$.

Cyanato: $-OCN$.

Isocyanato: -NCO.

Thiocyano (thiocyanato): -SCN.

5

Isothiocyano (isothiocyanato): -NCS.

Sulfhydryl (thiol, mercapto): -SH.

10 Thioether (sulfide): -SR, wherein R is a thioether substituent, for example, a C₁₋₇ alkyl group (also referred to as a C₁₋₇alkylthio group), a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably a C₁₋₇ alkyl group. Examples of C₁₋₇ alkylthio groups include, but are not limited to, -SCH₃ and -SCH₂CH₃.

15 Disulfide: -SS-R, wherein R is a disulfide substituent, for example, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably a C₁₋₇ alkyl group (also referred to herein as C₁₋₇ alkyl disulfide). Examples of C₁₋₇ alkyl disulfide groups include, but are not limited to, -SSCH₃ and -SSCH₂CH₃.

20 Sulfine (sulfinyl, sulfoxide): -S(=O)R, wherein R is a sulfine substituent, for example, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably a C₁₋₇ alkyl group. Examples of sulfine groups include, but are not limited to, -S(=O)CH₃ and -S(=O)CH₂CH₃.

25 Sulfone (sulfonyl): -S(=O)₂R, wherein R is a sulfone substituent, for example, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably a C₁₋₇ alkyl group, including, for example, a fluorinated or perfluorinated C₁₋₇ alkyl group. Examples of sulfone groups include, but are not limited to, -S(=O)₂CH₃ (methanesulfonyl, mesyl), -S(=O)₂CF₃ (triflyl), -S(=O)₂CH₂CH₃ (esyl), -S(=O)₂C₄F₉ (nonaflyl), -S(=O)₂CH₂CF₃ (tresyl), -S(=O)₂CH₂CH₂NH₂ (tauryl), -S(=O)₂Ph (phenylsulfonyl, besyl), 4-methylphenylsulfonyl (tosyl), 4-chlorophenylsulfonyl (closyl), 4-bromophenylsulfonyl (brosyl), 4-nitrophenyl (nosyl), 2-naphthalenesulfonate (napsyl), and 5-dimethylamino-naphthalen-1-ylsulfonate (dansyl).

30 Sulfonic acid (sulfino): -S(=O)OH, -SO₂H.

35

Sulfonic acid (sulfo): $-\text{S}(=\text{O})_2\text{OH}$, $-\text{SO}_3\text{H}$.

Sulfinate (sulfinic acid ester): $-\text{S}(=\text{O})\text{OR}$; wherein R is a sulfinate substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfinate groups include, but are not limited to, $-\text{S}(=\text{O})\text{OCH}_3$ (methoxysulfinyl; methyl sulfinate) and $-\text{S}(=\text{O})\text{OCH}_2\text{CH}_3$ (ethoxysulfinyl; ethyl sulfinate).

Sulfonate (sulfonic acid ester): $-\text{S}(=\text{O})_2\text{OR}$, wherein R is a sulfonate substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfonate groups include, but are not limited to, $-\text{S}(=\text{O})_2\text{OCH}_3$ (methoxysulfonyl; methyl sulfonate) and $-\text{S}(=\text{O})_2\text{OCH}_2\text{CH}_3$ (ethoxysulfonyl; ethyl sulfonate).

Sulfinyloxy: $-\text{OS}(=\text{O})\text{R}$, wherein R is a sulfinyloxy substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfinyloxy groups include, but are not limited to, $-\text{OS}(=\text{O})\text{CH}_3$ and $-\text{OS}(=\text{O})\text{CH}_2\text{CH}_3$.

Sulfonyloxy: $-\text{OS}(=\text{O})_2\text{R}$, wherein R is a sulfonyloxy substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfonyloxy groups include, but are not limited to, $-\text{OS}(=\text{O})_2\text{CH}_3$ (mesylate) and $-\text{OS}(=\text{O})_2\text{CH}_2\text{CH}_3$ (esylate).

Sulfate: $-\text{OS}(=\text{O})_2\text{OR}$; wherein R is a sulfate substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfate groups include, but are not limited to, $-\text{OS}(=\text{O})_2\text{OCH}_3$ and $-\text{SO}(=\text{O})_2\text{OCH}_2\text{CH}_3$.

Sulfamyl (sulfamoyl; sulfinic acid amide; sulfonamide): $-\text{S}(=\text{O})\text{NR}^1\text{R}^2$, wherein R^1 and R^2 are independently amino substituents, as defined for amino groups. Examples of sulfamyl groups include, but are not limited to, $-\text{S}(=\text{O})\text{NH}_2$, $-\text{S}(=\text{O})\text{NH}(\text{CH}_3)$, $-\text{S}(=\text{O})\text{N}(\text{CH}_3)_2$, $-\text{S}(=\text{O})\text{NH}(\text{CH}_2\text{CH}_3)$, $-\text{S}(=\text{O})\text{N}(\text{CH}_2\text{CH}_3)_2$, and $-\text{S}(=\text{O})\text{NHPH}$.

Sulfonamido (sulfinamoyl; sulfonic acid amide; sulfonamide): $-\text{S}(=\text{O})_2\text{NR}^1\text{R}^2$, wherein R^1 and R^2 are independently amino substituents, as defined for amino groups. Examples of sulfonamido groups include, but are not limited to, $-\text{S}(=\text{O})_2\text{NH}_2$, $-\text{S}(=\text{O})_2\text{NH}(\text{CH}_3)$, $-\text{S}(=\text{O})_2\text{N}(\text{CH}_3)_2$, $-\text{S}(=\text{O})_2\text{NH}(\text{CH}_2\text{CH}_3)$, $-\text{S}(=\text{O})_2\text{N}(\text{CH}_2\text{CH}_3)_2$, and $-\text{S}(=\text{O})_2\text{NHPH}$.

Sulfamino: $-\text{NR}^1\text{S}(=\text{O})_2\text{OH}$, wherein R^1 is an amino substituent, as defined for amino groups. Examples of sulfamino groups include, but are not limited to, $-\text{NHS}(=\text{O})_2\text{OH}$ and $-\text{N}(\text{CH}_3)\text{S}(=\text{O})_2\text{OH}$.

5

Sulfonamino: $-\text{NR}^1\text{S}(=\text{O})_2\text{R}$, wherein R^1 is an amino substituent, as defined for amino groups, and R is a sulfonamino substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfonamino groups include, but are not limited to, $-\text{NHS}(=\text{O})_2\text{CH}_3$ and $-\text{N}(\text{CH}_3)\text{S}(=\text{O})_2\text{C}_6\text{H}_5$.

10

Sulfinamino: $-\text{NR}^1\text{S}(=\text{O})\text{R}$, wherein R^1 is an amino substituent, as defined for amino groups, and R is a sulfinamino substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfinamino groups include, but are not limited to, $-\text{NHS}(=\text{O})\text{CH}_3$ and $-\text{N}(\text{CH}_3)\text{S}(=\text{O})\text{C}_6\text{H}_5$.

15

Phosphino (phosphine): $-\text{PR}_2$, wherein R is a phosphino substituent, for example, $-\text{H}$, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably $-\text{H}$, a C_{1-7} alkyl group, or a C_{5-20} aryl group. Examples of phosphino groups include, but are not limited to, $-\text{PH}_2$, $-\text{P}(\text{CH}_3)_2$, $-\text{P}(\text{CH}_2\text{CH}_3)_2$, $-\text{P}(\text{t-Bu})_2$, and $-\text{P}(\text{Ph})_2$.

20

Phospho: $-\text{P}(=\text{O})_2$.

Phosphinyl (phosphine oxide): $-\text{P}(=\text{O})\text{R}_2$, wherein R is a phosphinyl substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group or a C_{5-20} aryl group. Examples of phosphinyl groups include, but are not limited to, $-\text{P}(=\text{O})(\text{CH}_3)_2$, $-\text{P}(=\text{O})(\text{CH}_2\text{CH}_3)_2$, $-\text{P}(=\text{O})(\text{t-Bu})_2$, and $-\text{P}(=\text{O})(\text{Ph})_2$.

25

Phosphonic acid (phosphono): $-\text{P}(=\text{O})(\text{OH})_2$.

Phosphonate (phosphono ester): $-\text{P}(=\text{O})(\text{OR})_2$, where R is a phosphonate substituent, for example, $-\text{H}$, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably $-\text{H}$, a C_{1-7} alkyl group, or a C_{5-20} aryl group. Examples of phosphonate groups include, but are not limited to, $-\text{P}(=\text{O})(\text{OCH}_3)_2$, $-\text{P}(=\text{O})(\text{OCH}_2\text{CH}_3)_2$, $-\text{P}(=\text{O})(\text{O-t-Bu})_2$, and $-\text{P}(=\text{O})(\text{OPh})_2$.

30

Phosphoric acid (phosphonoxy): $-\text{OP}(=\text{O})(\text{OH})_2$.

35

Phosphate (phosphonoxy ester): $-OP(=O)(OR)_2$, where R is a phosphate substituent, for example, -H, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably -H, a C₁₋₇ alkyl group, or a C₅₋₂₀ aryl group. Examples of phosphate groups include, but are not limited to, $-OP(=O)(OCH_3)_2$, $-OP(=O)(OCH_2CH_3)_2$, $-OP(=O)(O-t-Bu)_2$, and $-OP(=O)(OPh)_2$.

Phosphorous acid: $-OP(OH)_2$.

10 Phosphite: $-OP(OR)_2$, where R is a phosphite substituent, for example, -H, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably -H, a C₁₋₇ alkyl group, or a C₅₋₂₀ aryl group. Examples of phosphite groups include, but are not limited to, $-OP(OCH_3)_2$, $-OP(OCH_2CH_3)_2$, $-OP(O-t-Bu)_2$, and $-OP(OPh)_2$.

15 Phosphoramidite: $-OP(OR^1)-NR^2_2$, where R¹ and R² are phosphoramidite substituents, for example, -H, a (optionally substituted) C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably -H, a C₁₋₇ alkyl group, or a C₅₋₂₀ aryl group. Examples of phosphoramidite groups include, but are not limited to, $-OP(OCH_2CH_3)-N(CH_3)_2$, $-OP(OCH_2CH_3)-N(i-Pr)_2$, and $-OP(OCH_2CH_2CN)-N(i-Pr)_2$.

20 Phosphoramidate: $-OP(=O)(OR^1)-NR^2_2$, where R¹ and R² are phosphoramidate substituents, for example, -H, a (optionally substituted) C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably -H, a C₁₋₇ alkyl group, or a C₅₋₂₀ aryl group. Examples of phosphoramidate groups include, but are not limited to, $-OP(=O)(OCH_2CH_3)-N(CH_3)_2$, $-OP(=O)(OCH_2CH_3)-N(i-Pr)_2$, and $-OP(=O)(OCH_2CH_2CN)-N(i-Pr)_2$.

Bisoxylalkylene: $-O-(CH_2)_n-O-$, where n=1-3, which bonds to adjacent atoms. Examples of bisoxylalkylene groups include, but are not limited to, $-O-CH_2-O-$.

30 *Alkylene*

C₃₋₁₂ alkylene: The term "C₃₋₁₂ alkylene", as used herein, pertains to a bidentate moiety obtained by removing two hydrogen atoms, either both from the same carbon atom, or one from each of two different carbon atoms, of a hydrocarbon compound having from 3 to 12 carbon atoms (unless otherwise specified), which may be aliphatic or alicyclic, and which

may be saturated, partially unsaturated, or fully unsaturated. Thus, the term "alkylene" includes the sub-classes alkenylene, alkynylene, cycloalkylene, etc., discussed below.

5 Examples of linear saturated C₃₋₁₂ alkylene groups include, but are not limited to, -(CH₂)_n- where n is an integer from 3 to 12, for example, -CH₂CH₂CH₂- (propylene), -CH₂CH₂CH₂CH₂- (butylene), -CH₂CH₂CH₂CH₂CH₂- (pentylene) and -CH₂CH₂CH₂CH₂CH₂CH₂CH₂- (heptylene).

10 Examples of branched saturated C₃₋₁₂ alkylene groups include, but are not limited to, -CH(CH₃)CH₂-, -CH(CH₃)CH₂CH₂-, -CH(CH₃)CH₂CH₂CH₂-, -CH₂CH(CH₃)CH₂-, -CH₂CH(CH₃)CH₂CH₂-, -CH(CH₂CH₃)-, -CH(CH₂CH₃)CH₂-, and -CH₂CH(CH₂CH₃)CH₂-.

15 Examples of linear partially unsaturated C₃₋₁₂ alkylene groups (C₃₋₁₂ alkenylene, and alkynylene groups) include, but are not limited to, -CH=CH-CH₂-, -CH₂-CH=CH₂-, -CH=CH-CH₂-CH₂-, -CH=CH-CH₂-CH₂-CH₂-, -CH=CH-CH=CH-, -CH=CH-CH=CH-CH₂-, -CH=CH-CH=CH-CH₂-CH₂-, -CH=CH-CH₂-CH=CH-, -CH=CH-CH₂-CH₂-CH=CH-, and -CH₂-C≡C-CH₂-.

20 Examples of branched partially unsaturated C₃₋₁₂ alkylene groups (C₃₋₁₂ alkenylene and alkynylene groups) include, but are not limited to, -C(CH₃)=CH-, -C(CH₃)=CH-CH₂-, -CH=CH-CH(CH₃)- and -C≡C-CH(CH₃)-.

25 Examples of alicyclic saturated C₃₋₁₂ alkylene groups (C₃₋₁₂ cycloalkylenes) include, but are not limited to, cyclopentylene (e.g. cyclopent-1,3-ylene), and cyclohexylene (e.g. cyclohex-1,4-ylene).

30 Examples of alicyclic partially unsaturated C₃₋₁₂ alkylene groups (C₃₋₁₂ cycloalkylenes) include, but are not limited to, cyclopentenylene (e.g. 4-cyclopenten-1,3-ylene), cyclohexenylene (e.g. 2-cyclohexen-1,4-ylene; 3-cyclohexen-1,2-ylene; 2,5-cyclohexadien-1,4-ylene).

Pharmaceutically acceptable cations

Examples of pharmaceutically acceptable monovalent and divalent cations are discussed in Berge, *et al.*, *J. Pharm. Sci.*, **66**, 1-19 (1977), which is incorporated herein by reference.

The pharmaceutically acceptable cation may be inorganic or organic.

5 Examples of pharmaceutically acceptable monovalent inorganic cations include, but are not limited to, alkali metal ions such as Na^+ and K^+ . Examples of pharmaceutically acceptable divalent inorganic cations include, but are not limited to, alkaline earth cations such as Ca^{2+} and Mg^{2+} . Examples of pharmaceutically acceptable organic cations include, but are not limited to, ammonium ion (i.e. NH_4^+) and substituted ammonium ions (e.g. NH_3R^+ , NH_2R_2^+ , NHR_3^+ , NR_4^+). Examples of some suitable substituted ammonium ions are those derived from: ethylamine, diethylamine, dicyclohexylamine, triethylamine, 10 butylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine, benzylamine, phenylbenzylamine, choline, meglumine, and tromethamine, as well as amino acids, such as lysine and arginine. An example of a common quaternary ammonium ion is $\text{N}(\text{CH}_3)_4^+$.

Proliferative Diseases

15 One of ordinary skill in the art is readily able to determine whether or not a candidate compound treats a proliferative condition for any particular cell type. For example, assays which may conveniently be used to assess the activity offered by a particular compound are described in the examples below.

20 The term "proliferative disease" pertains to an unwanted or uncontrolled cellular proliferation of excessive or abnormal cells which is undesired, such as, neoplastic or hyperplastic growth, whether *in vitro* or *in vivo*.

25 Examples of proliferative conditions include, but are not limited to, benign, pre-malignant, and malignant cellular proliferation, including but not limited to, neoplasms and tumours (e.g. histiocytoma, glioma, astrocytoma, osteoma), cancers (e.g. lung cancer, small cell lung cancer, gastrointestinal cancer, bowel cancer, colon cancer, breast carcinoma, ovarian carcinoma, prostate cancer, testicular cancer, liver cancer, kidney cancer, bladder cancer, pancreas cancer, brain cancer, sarcoma, osteosarcoma, Kaposi's sarcoma, melanoma), 30 leukemias, psoriasis, bone diseases, fibroproliferative disorders (e.g. of connective tissues), and atherosclerosis.

35 Any type of cell may be treated, including but not limited to, lung, gastrointestinal (including, e.g. bowel, colon), breast (mammary), ovarian, prostate, liver (hepatic), kidney (renal), bladder, pancreas, brain, and skin.

Methods of Treatment

As described above, the present invention provide the use of a compound of formula I in a method of therapy. Also provided is a method of treatment, comprising administering to a
5 subject in need of treatment a therapeutically-effective amount of a compound of formula I, preferably in the form of a pharmaceutical composition. The term "therapeutically effective amount" is an amount sufficient to show benefit to a patient. Such benefit may be at least amelioration of at least one symptom. The actual amount administered, and rate and time-

10 Prescription of treatment, e.g. decisions on dosage, is within the responsibility of general practitioners and other medical doctors.

A compound may be administered alone or in combination with other treatments, either
15 simultaneously or sequentially dependent upon the condition to be treated. Examples of treatments and therapies include, but are not limited to, chemotherapy (the administration of active agents, including, e.g. drugs; surgery; and radiation therapy.

Pharmaceutical compositions according to the present invention, and for use in accordance
20 with the present invention, may comprise, in addition to the active ingredient, i.e. a compound of formula I, a pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material will depend on the route of administration, which may be oral, or by injection, e.g. cutaneous, subcutaneous, or intravenous.

25 Pharmaceutical compositions for oral administration may be in tablet, capsule, powder or liquid form. A tablet may comprise a solid carrier or an adjuvant. Liquid pharmaceutical compositions generally comprise a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil. Physiological saline solution, dextrose or other
30 saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may be included. A capsule may comprise a solid carrier such a gelatin.

For intravenous, cutaneous or subcutaneous injection, or injection at the site of affliction,
35 the active ingredient will be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the

art are well able to prepare suitable solutions using, for example, isotonic vehicles such as Sodium Chloride Injection, Ringer's Injection, Lactated Ringer's Injection. Preservatives, stabilisers, buffers, antioxidants and/or other additives may be included, as required.

5 Includes Other Forms


Unless otherwise specified, included in the above are the well known ionic, salt, solvate, and protected forms of these substituents. For example, a reference to carboxylic acid (-COOH) also includes the anionic (carboxylate) form (-COO⁻), a salt or solvate thereof, as well as conventional protected forms. Similarly, a reference to an amino group includes the protonated form (-N⁺HR¹R²), a salt or solvate of the amino group, for example, a hydrochloride salt, as well as conventional protected forms of an amino group. Similarly, a reference to a hydroxyl group also includes the anionic form (-O⁻), a salt or solvate thereof, as well as conventional protected forms.

15 In particular, a reference to group (-SO₂M) also includes the anionic form (-SO₂⁻), or solvate thereof, as well as conventional protected forms.

Isomers and Solvates


Certain compounds may exist in one or more particular geometric, optical, enantiomeric, diastereomeric, epimeric, atropic, stereoisomeric, tautomeric, conformational, or anomeric forms, including but not limited to, cis- and trans-forms; E- and Z-forms; c-, t-, and r- forms; endo- and exo-forms; R-, S-, and meso-forms; D- and L-forms; d- and l-forms; (+) and (-) forms; keto-, enol-, and enolate-forms; syn- and anti-forms; synclinal- and anticlinal-forms; α- and β-forms; axial and equatorial forms; boat-, chair-, twist-, envelope-, and halfchair-forms; and combinations thereof, hereinafter collectively referred to as "isomers" (or "isomeric forms").

In some embodiments, compounds of the present invention have the following stereochemistry at the C11 position:



Note that, except as discussed below for tautomeric forms, specifically excluded from the term "isomers", as used herein, are structural (or constitutional) isomers (i.e. isomers which differ in the connections between atoms rather than merely by the position of atoms in space). For example, a reference to a methoxy group, -OCH₃, is not to be construed as a reference to its structural isomer, a hydroxymethyl group, -CH₂OH. Similarly, a reference to ortho-chlorophenyl is not to be construed as a reference to its structural isomer, meta-chlorophenyl. However, a reference to a class of structures may well include structurally isomeric forms falling within that class (e.g. C₁₋₇ alkyl includes n-propyl and iso-propyl; butyl includes n-, iso-, sec-, and tert-butyl; methoxyphenyl includes ortho-, meta-, and para-methoxyphenyl).

The above exclusion does not pertain to tautomeric forms, for example, keto-, enol-, and enolate-forms, as in, for example, the following tautomeric pairs: keto/enol (illustrated below), imine/enamine, amide/imino alcohol, amidine/amidine, nitroso/oxime, thioketone/enethiol, N-nitroso/hydroxyazo, and nitro/aci-nitro.



Note that specifically included in the term "isomer" are compounds with one or more isotopic substitutions. For example, H may be in any isotopic form, including ¹H, ²H (D), and ³H (T); C may be in any isotopic form, including ¹²C, ¹³C, and ¹⁴C; O may be in any isotopic form, including ¹⁶O and ¹⁸O; and the like.


Unless otherwise specified, a reference to a particular compound includes all such isomeric forms, including (wholly or partially) racemic and other mixtures thereof. Methods for the preparation (e.g. asymmetric synthesis) and separation (e.g. fractional crystallisation and chromatographic means) of such isomeric forms are either known in the art or are readily obtained by adapting the methods taught herein, or known methods, in a known manner.

It may be convenient or desirable to prepare, purify, and/or handle a corresponding solvate of the active compound. The term "solvate" is used herein in the conventional sense to refer to a complex of solute (e.g. active compound, salt of active compound) and solvent. If the solvent is water, the solvate may be conveniently referred to as a hydrate, for example, a mono-hydrate, a di-hydrate, a tri-hydrate, etc.

General synthetic routes


The synthesis of PBD compounds is extensively discussed in WO 00/12508, which discussion is incorporated herein by reference.

- 5 As discussed in that patent application, a key step in a preferred route to PBDs is a cyclisation to produce the B-ring, involving generation of an aldehyde (or functional equivalent thereof) at what will be the 11-position, and attack thereon by the Pro-N10-nitrogen:




- 10 wherein the substituents are as defined above, R^8 represents the link (-X-R"-X-) to the other PBD moiety, R^{10} is a nitrogen protecting group and R^{12} is R^2 or a precursor thereof. The "masked aldehyde" -CPQ may be an acetal or thioacetal, in which case the cyclisation involves unmasking. Alternatively, it may be an alcohol -CHOH, in which case the reaction involves oxidation, e.g. by means of TPAP, TEMPO or DMSO (Swern oxidation).

- 15 The masked aldehyde compound can be produced by condensing a corresponding 2,4-substituted pyrrolidine with a 2-nitrobenzoic acid:



- 20 The nitro group can then be reduced to $-NH_2$ and protected by reaction with a suitable agent, e.g. a chloroformate, which provides the removable nitrogen protecting group in the compound of formula IV.

A process involving the oxidation-cyclization procedure is illustrated in scheme 1 (an alternative type of cyclisation will be described later with reference to scheme 2).



Scheme 1

5 Exposure of the alcohol (B) (in which the Pro-N10-nitrogen is generally protected as carbamate) to tetrapropylammonium perruthenate (TPAP)/N-methylmorpholine N-oxide (NMO) over A4 sieves results in oxidation accompanied by spontaneous B-ring closure to afford the desired product IV. The TPAP/NMO oxidation procedure is found to be particularly convenient for small scale reactions while the use of DMSO-based oxidation methods, particularly Swern oxidation, proves superior for larger scale work (e.g. > 1 g). A particularly preferred oxidising agent is (diacetoxyiodo)benzene (1.1 eq) and TEMPO (0.1 eq) dissolved in CH₂Cl₂.

15 The uncyclized alcohol (B) may be prepared by the reaction of a nitrogen protection reagent of formula D, which is preferably a chloroformate or acid chloride, to a solution of the amino alcohol C, generally in solution, generally in the presence of a base such as pyridine (preferably 2 equivalents) at a moderate temperature (e.g. at 0°C). Under these conditions little or no O-acylation is usually observed.

The key amino alcohol C may be prepared by reduction of the corresponding nitro compound E, by choosing a method which will leave the rest of the molecule intact. Treatment of E with tin (II) chloride in a suitable solvent, e.g. refluxing methanol, generally affords, after the removal of the tin salts, the desired product in high yield.

5

Exposure of E to hydrazine/Raney nickel avoids the production of tin salts and may result in a higher yield of C, although this method is less compatible with the range of possible C and A-ring substituents. For instance, if there is C-ring unsaturation (either in the ring itself, or in R₂ or R₃), this technique may be unsuitable. Another suitable means of reduction would be catalytic hydrogenation using palladium on carbon as a catalyst.

10


The nitro compound of formula E may be prepared by coupling the appropriate *o*-nitrobenzoyl chloride to a compound of formula F, e.g. in the presence of K₂CO₃ at -25°C under a N₂ atmosphere. Compounds of formula F can be readily prepared, for example by olefination of the ketone derived from L-trans-hydroxy proline. The ketone intermediate can also be exploited by conversion to the enol triflate for use in palladium mediated coupling reactions.

15

The *o*-nitrobenzoyl chloride is synthesised from the *o*-nitrobenzoic acid (or alkyl ester after hydrolysis) of formula G, which itself is prepared from the vanillic acid (or alkyl ester) derivative H. Many of these are commercially available and some are disclosed in Althuis, T.H. and Hess, H.J., *J. Medicinal Chem.*, **20(1)**, 146-266 (1977).

20

Alternative Cyclisation (Scheme 2)



Scheme 2


- 5 In scheme 1, the final or penultimate step was an oxidative cyclisation. An alternative, using thioacetal coupling, is shown in scheme 2. Mercury-mediated unmasking causes cyclisation to the protected PBD compound IV.

10 The thioacetal compound may be prepared as shown in scheme 2: the thioacetal protected C-ring [prepared via a literature method: Langley, D.R. & Thurston, D.E., *J. Organic Chemistry*, 52, 91-97 (1987)] is coupled to the o-nitrobenzoic acid (or alkyl ester after hydrolysis) (G) using a literature procedure. The resulting nitro compound cannot be reduced by hydrogenation, because of the thioacetal group, so the tin(II) chloride method is used to afford the amine. This is then N-protected, e.g., by reaction with a chloroformate or acid chloride, such as 2,2,2-trichloroethylchloroformate.

15

Acetal-containing C-rings can be used as an alternative in this type of route with deprotection involving other methods, including the use of acidic conditions.

Dimer Synthesis (Scheme 3)



Scheme 3

PBD dimers, as in the present invention, may be synthesized using the strategy developed for the synthesis of the protected PBD monomers. The synthesis routes illustrated in scheme 3 show compounds when the dimer linkage is of the formula $-O-(CH_2)_n-O-$, and may be readily modified for other dimer linkages. The step of dimer formation is normally carried out to form a bis(nitro acid) G'. This compound can then be treated as compound G in either scheme 1 or scheme 2 above.

The bis(nitro acid) G' may be obtained by nitrating (e.g. using 70% nitric acid) the bis(carboxylic acid). This can be synthesised by alkylation of two equivalents of the relevant benzoic acid with the appropriate diiodoalkane under basic conditions. Many

benzoic acids are commercially available and others can be synthesised by conventional methods. Alternatively, the relevant benzoic acid esters can be joined together by a Mitsunobu etherification with an appropriate alkanediol, followed by nitration, and then hydrolysis (not illustrated).


5 An alternative synthesis of the bis(nitro acid) involves oxidation of the bis(nitro aldehyde), e.g. with potassium permanganate. This can be obtained in turn by direct nitration of the bis(aldehyde), e.g. with 70% HNO₃. Finally, the bis(aldehyde) can be obtained via the Mitsunobu etherification of two equivalents of the benzoic aldehyde with the appropriate
10 alkanediol.

Alternative routes to PBDs

Alternative methods of synthesising N10 protected PBDs are disclosed WO 2005/023814, which is incorporated herein, and which describes the use of isocyanate intermediates.

15

Introduction of C2 aryl substituent: introducing leaving group at C2



The C2 aryl substituent may be in place in compounds of formula **IV**, by starting with the appropriate material, in which case the N10 may be deprotected to yield the desired
20 compound (see below). Alternatively, the following method can be used where R¹² is a protected hydroxy group. Following cyclisation to form the B-ring, the C11-alcohol **IV** is then preferably re-protected, by conventional means to provide **IIIb**. For example, if R¹¹ is TBS, the protection can take place by reacting **IV** with TBSOTf and 2,6-lutidine. Cleavage of the C2-protecting group from **IIIb** then provides the C2 alcohol. For example, where the

C2 protecting group (R^{14}) is acyl, this deprotection may be performed by addition of an aqueous solution of K_2CO_3 .

This reprotection at the C11 position and deprotection of the C2 alcohol allows subsequent reaction of selectively the C2 alcohol position leaving the C11 position unaffected.


The C2-alcohol may then be oxidized to the ketone **IIIb**. Preferably this oxidation is performed under Swern conditions, in good yield. However, other oxidation methods involving TPAP or the Dess Martin reagent also provide the ketone in good yield.

R^{12} in the compound of formula **II** may be $-OSO_2CH_3$, $-OSO_2(C_nF_{2n+1})$ where $n=0,1$ or 4 , or $-OSO_2R^S$, in which case the conversion from **IIIb** may be achieved by treatment with the appropriate anhydride. For example, if R^{12} is triflate that reaction with trifluoromethanesulfonic anhydride in DCM in the presence of pyridine.

R^{12} in the compound of formula **II** may also be $-I$ or $-Br$, in which case the conversion from **IIIb** may be achieved by reaction with hydrazine and iodine or bromine respectively.

R^{12} in the compound of formula **II** may also be $-Cl$, where the conversion from **IIIb** may be achieved by reaction with a phosphorous oxychloride (e.g. $POCl_3$).

Introduction of C2 aryl substituent: Replacement of leaving group



This compound of formula **II** may be reacted under a variety of conditions to yield PBD precursor molecules with pendant groups coupled at the C2 position **Ic**.

In particular, the use of palladium catalysed coupling is preferred, such as Suzuki, Stille and Heck coupling. The palladium catalyst may be any suitable catalyst, for example $Pd(PPh_3)_4$, $Pd(OCOCH_3)_2$, $PdCl_2$, $Pd(dba)_3$. The compounds which are coupled may be any suitable reactant, e.g. for Heck, alkenes with an sp^2 H; for Stille, organostannanes; and for Suzuki, organoboron derivatives.

In a preferred aspect of the invention, the coupling may be performed under microwave conditions. Typically, the palladium catalyst, such as Pd(PPh₃)₄, is solid supported, for example on polystyrene, to facilitate work-up and allow potential recycling of catalyst. Unreacted boronic acid can be sequestered following complete consumption of triflate using PS-DEAM, with a phase separator cartridge being used to isolate the coupling product. Such a method allows for the parallel synthesis of more than one (e.g. up to 10, 20 or 30) compound at the same time.

The imine bond in the compound of formula **1c** can be unprotected by standard methods to yield the unprotected compound **1b** (which may be in its carbinolamine or carboinolamine ether form, depending on the solvents used). For example if R¹⁰ is Alloc, then the deprotection is carried using palladium to remove the N10 protecting group, followed by the elimination of water. If R¹⁰ is Troc, then the deprotection is carried out using a Cd/Pb couple to yield the compound of formula **1b**.

Reference is also made to the synthesis discussion and examples in WO 2004/043963 and WO 2005/085251, which are herein incorporated by reference.

Conversion to sulphur containing form

The conversion of compounds of formula **1b** to those of the present invention may be carried out by the addition of the appropriate bisulphite salt or sulphinate salt, followed by an appropriate purification step. Further methods are described in GB 2 053 894, which is herein incorporated by reference.

Further Preferences

It is preferred that X is O.

It is preferred that Rⁿ represents a linear saturated C₃₋₁₂ alkylene group, and more preferably an alkylene group having 3, 5, 7, 8, 9, 10, 11 or 12 carbon atoms. Of these C₃ and C₅ linear saturated alkylene groups are preferred.

R⁹ is preferably H.

R⁶ is preferably selected from H, OH, OR, SH, NH₂, nitro and halo, and is more preferably H or halo, and most preferably is H.

R⁷ is preferably selected from H, OH, OR, SH, SR, NH₂, NHR, NRR', and halo, and more preferably independently selected from H, OH and OR, where R is preferably selected from optionally substituted C₁₋₇ alkyl, C₃₋₁₀ heterocyclyl and C₅₋₁₀ aryl groups. Particularly preferred substituents at the 7- position are OMe and OCH₂Ph.

R² is preferably an optionally substituted C₅₋₇ aryl group, and most preferably an optionally substituted phenyl group.

In some embodiments, R² is a C₉₋₁₂ aryl group, for example naphth-1-yl or naphth-2-yl. Further examples of C₉₋₁₂ aryl groups include quinolinyl, for example, quinolin-2-yl, quinolin-3-yl and quinolin-6-yl.

In other embodiments, R² is a C₅₋₇ heteroaryl group, for example furanyl, thiophenyl and pyridyl. Of these thiophenyl is preferred, for example, thiophen-2-yl and thiophen-3-yl.

The C₅₋₂₀ aryl group may bear any substituent group. It preferably bears from 1 to 3 substituent groups, with 1 and 2 being more preferred, and singly substituted groups being most preferred.


Preferred C₅₋₂₀ aryl substituents, particularly for phenyl, include: halo (e.g. F, Cl, Br); C₁₋₇ alkoxy (e.g. methoxy, ethoxy); C₁₋₇ alkyl (e.g. methyl, trifluoromethyl, ethyl, propyl, t-butyl); bis-oxy-alkylene (e.g. bis-oxy-methylene, -O-CH₂-O-).

Particularly preferred substituted C₅₋₂₀ aryl groups include, but are not limited to, 4-methylphenyl, 4-methoxyphenyl, 3-methoxyphenyl, 4-fluorophenyl, 3,4-bisoxymethylenephenyl, 4-trifluoromethylphenyl, 4-methylthiophenyl, 4-cyanophenyl and 4-phenoxyphenyl.


Particularly preferred unsubstituted C₅₋₂₀ aryl groups include, but are not limited to thiophen-2-yl, naphth-2-yl, quinolin-3-yl and quinolin-6-yl.

If R is optionally substituted C₁₋₁₂ alkyl, it is preferred that it is optionally substituted C₁₋₇ alkyl.



It is preferred that both PBD monomers are identically substituted.




5




More preferred compounds include:



5



A most preferred compound is:




10

The above preferred compounds may also have a C₅ alkylene linking chain.

Examples**Example 1**


(a) (2*S*,4*R*)-*N*-(Benzyloxycarbonyl)-2-*t*-butyldimethylsilyloxymethyl-4-hydroxypyrrolidine (**1**)



5

Compound **1** is formed in high yield in a four step process known in the art starting from *trans*-4-hydroxy-L-proline (S.J. Gregson *et al.*, *J. Med. Chem.*, 2004, 1161-1174).

(b) (2*S*,4*R*)-*N*-(Benzyloxycarbonyl)-2-*t*-butyldimethylsilyloxymethyl-4-oxyacetylpyrrolidine (**2**)




10

Pyridine (18.3 g, 18.7 mL, 232 mmol, 1.1 eq), acetic anhydride (23.6 g, 21.8 mL, 232 mmol, 1.1 eq) and DMAP (5.14 g, 42.1 mmol, 0.2 eq) were added to a stirred solution of **1** (76.9 g, 211 mmol) in dry THF (1 L). The reaction mixture was stirred for 16 hours after which time TLC (95:5 v/v CHCl₃/MeOH) showed the complete consumption of starting material. Excess solvent was removed by rotary evaporation and the residue was dissolved in EtOAc (1 L), washed with 1N HCl (2 x 1 L), H₂O (1 L), brine (1 L) and dried (MgSO₄). Filtration and evaporation of the solvent afforded acetate **2** as a colourless oil (80.7 g, 94%): ¹H NMR (400 MHz, CDCl₃) (rotamers) δ 7.36-7.12 (m, 5H), 5.30-5.10 (m, 3H), 4.09-3.97 (m, 2H), 3.74-3.55 (m, 3H), 2.36-2.29 (m, 1H), 2.11-2.06 (m, 1H), 2.02 (s, 3H), 0.87 (s, 6H), 0.86 (s, 3H), 0.03 and 0.00 (s x 2, 6H); MS (ES), *m/z* (relative intensity) 430 ([M + Na]⁺, 95), 408 ([M + H]⁺, 100).


20

(c) (2*S*,4*R*)-2-*t*-Butyldimethylsilyloxymethyl-4-oxyacetylpyrrolidine (**3**)



A slurry of silyl ether **2** (1.95 g, 4.80 mmol) and 10% Pd/C (0.17 g) in absolute ethanol (10 mL) was subjected to Parr hydrogenation at 45 Psi for 16 h after which time TLC (95:5 v/v CHCl₃/MeOH) showed the complete consumption of starting material. The reaction mixture was filtered through celite to remove the Pd/C, and the filter pad was washed repeatedly with ethanol. Excess solvent was removed by rotary evaporation under reduced pressure to afford the amine **3** as a pale orange waxy oil (1.28 g, 98%): IR (CHCl₃) 3315, 2930, 2858, 1739, 1652, 1472, 1435, 1375, 1251, 1088, 838, 779, 667 cm⁻¹.

(d) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(2*S*,4*R*)-(5-methoxy-2-nitro-1,4-phenylene)carbonyl]]bis[2-(*tert*-butyldimethylsilyloxymethyl)-4-oxyacetylpyrrolidine] (**5**)




A catalytic amount of DMF (2 drops) was added to a stirred solution of the nitro-acid **4** (8.12 g, 17.4 mmol)¹ and oxalyl chloride (3.80 mL, 5.52 g, 43.5 mmol, 2.5 eq) in dry THF (250 mL). The initial precipitate dissolved gradually and the reaction mixture was allowed to stir for 16 h at room temperature. The resulting acid chloride solution was added dropwise to a stirred mixture of the amine **3** (11.9 g, 43.5 mmol, 2.5 eq), TEA (9.71 mL, 7.05 g, 69.7 mmol, 4.0 eq) and H₂O (2.26 mL) in THF (100 mL) at 0°C (ice/acetone) under a nitrogen atmosphere. The reaction mixture was allowed to warm to room temperature and stirred for a further 2.5 h. Excess THF was removed by rotary evaporation and the resulting residue was partitioned between H₂O (400 mL) and EtOAc (400 mL). The layers were allowed to separate and the aqueous layer was extracted with EtOAc (3 x 200 mL). The combined organic layers were then washed with saturated NH₄Cl (200 mL), saturated NaHCO₃ (200 mL), brine (200 mL) and dried (MgSO₄). Filtration and evaporation of the solvent gave the crude product as a dark oil. Purification by flash chromatography (99.7:0.3 v/v CHCl₃/MeOH) isolated the pure amide **5** as a light yellow glass (13.3 g, 78%): ¹H NMR (400 MHz, CDCl₃) δ 7.60 (s, 2H), 6.60 (s, 2H), 5.06 (br s, 2H), 4.44 (br s, 2H), 4.25-4.20 (m, 4H), 4.10-4.08 (m, 2H), 3.80 (s, 6H), 3.64-3.62 (m, 2H), 3.36-3.32 (m, 2H),

3.11-3.08 (m, 2H), 2.36-2.26 (m, 4H), 2.13-2.08 (m, 2H), 1.92 (s, 6H), 0.80 (s, 18H), 0.00 (s x 2, 12H); ^{13}C NMR (100.6 MHz, CDCl_3) δ 171.0, 166.3, 154.5, 148.2, 137.4, 128.0, 127.2, 109.2, 108.5, 72.9, 65.6, 62.6, 57.4, 56.5, 54.8, 33.0, 28.6, 25.8, 21.0, 18.1; MS (ES), m/z (relative intensity) 1000 ($[\text{M} + \text{Na}]^+$, 39), 978 ($[\text{M} + \text{H}]^+$, 63), 977 (M^+ , 100), 812 (13).


5

(e) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(2S,4R)-(5-methoxy-2-amino-1,4-phenylene)carbonyl]]bis[2-(tert-butyldimethylsilyloxymethyl)-4-oxyacetylpyrrolidine] (6)




Sodium dithionite (16.59 g, 95.27 mmol, 5 eq) was added to a stirred solution of amide **5** (18.6 g, 19.1 mmol) in H_2O (200 mL) and THF (400 mL). The reaction mixture was allowed to stir for 36 h after which time excess THF was removed by rotary evaporation and the resulting residue was extracted with EtOAc (3 x 250 mL). The combined organic layers were then washed with H_2O (300 mL), brine (300 mL) and dried (MgSO_4). Filtration and evaporation of the solvent yielded the crude product which was purified by flash chromatography (80:20 v/v hexane/EtOAc then gradient to neat EtOAc) to afford the product **6** as a yellow foam (9.53 g, 55%): ^1H NMR (400 MHz, CDCl_3) (rotamers) δ 6.70 and 6.67 (s x 2, 2H), 6.25 and 6.23 (s x 2, 2H), 5.20 (br s, 2H), 4.49 (br s, 4H), 4.16-4.05 (m, 6H), 3.70 (s, 6H), 3.68-3.57 (m, 4H), 2.36-2.27 (m, 4H), 2.12-2.04 (m, 2H), 1.96 (s, 6H), 0.85 (s, 18H), 0.01 and 0.00 (s x 2, 12H); ^{13}C NMR (100.6 MHz, CDCl_3) δ 170.6, 170.0, 141.1, 116.3, 113.1, 102.3, 102.1, 102.0, 66.2, 65.3, 65.2, 57.0, 28.9, 18.2; MS (ES), m/z (relative intensity) 946 ($\text{M}^+ + 29$, 43), 933 ($[\text{M} + 16]^+$, 61), 932 ($[\text{M} + 15]^+$, 100), 918 ($[\text{M} + \text{H}]^+$, 72).

(f) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(2S,4R)-[5-methoxy-1,4-phenylene-2-(2,2,2-trichloroethoxycarbonylamino)]carbonyl]]bis[2-(tert-butylidimethylsilyloxymethyl)-4-oxycetylpyrrolidine] (7)



5 A solution of 2,2,2-trichloroethyl chloroformate (3.58 mL, 5.50 g, 26.0 mmol, 2.2 eq) in dry DCM (60 mL) was added dropwise to a solution of anhydrous pyridine (3.82 mL, 3.80 g, 47.2 mmol, 4.0 eq) and bis-aniline 6 (10.8 g, 11.8 mmol) in dry DCM (150 mL) at -10 °C (liq. N₂/ethanediol). After 16 h at room temperature, the reaction mixture was washed with saturated NH₄Cl (2 x 150 mL), saturated CuSO₄ (150 mL), H₂O (150 mL), brine (150 mL) and dried (MgSO₄). Filtration and evaporation of the solvent yielded a yellow viscous oil which was purified by flash chromatography (70:30 v/v hexane/EtOAc) to afford the product 7 as a white glass (13.8 g, 92%): ¹H NMR (400 MHz, CDCl₃) δ 9.42 (br s, 1H), 7.83 (s, 2H), 6.76 and 6.74 (s x 2, 2H), 5.21 (br s, 2H), 4.79 and 4.73 (d x 2, 4H, J = 12.0 Hz), 4.56 (br s, 2H), 4.26-4.23 (m, 4H), 4.09-4.04 (m, 2H), 3.74 (s, 6H), 3.72-3.68 (m, 2H), 3.60 (br s, 4H), 2.40-2.32 (m, 4H), 2.23-2.08 (m, 2H), 1.95 (s, 6H), 0.85 (s, 18H), 0.01 and 0.00 (s x 2, 12H); ¹³C NMR (100.6 MHz, CDCl₃) δ 170.4, 169.2, 151.9, 151.5, 150.8, 143.4, 132.6, 114.4, 111.7, 95.3, 74.4, 65.5, 65.4, 57.3, 56.4, 32.5, 28.8, 25.8, 21.1, 18.1, 14.9; MS (ES), m/z (relative intensity) 1306 ([M + 38]⁺, 92), 1304 ([M + 36]⁺, 100), 1282 ([M + 14]⁺, 97), 1280 ([M + 12]⁺, 55).


20 (g) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(2S,4R)-[5-methoxy-1,4-phenylene-2-(2,2,2-trichloroethoxycarbonylamino)]carbonyl]]bis(2-hydroxymethyl-4-oxycetylpyrrolidine) (8)



25 A mixture of glacial acetic acid (310 mL) and H₂O (100 mL) was added to a solution of 7 (13.8 g, 10.9 mmol) in THF (250 mL) and was stirred for 16 h at room temperature. The reaction mixture was diluted with DCM (750 mL) and neutralised with saturated NaHCO₃ (5

L). The aqueous layer was extracted with DCM (3 x 500 mL) and the organic layers were combined, washed with brine (1 L) and dried (MgSO₄). TLC (60:40 v/v hexane/EtOAc) revealed the complete disappearance of the starting material. Filtration and evaporation of the solvent afforded the crude product which was purified by flash column chromatography (99.7:0.3 v/v CHCl₃/MeOH then gradient to 96:4 v/v CHCl₃/MeOH) to provide the product **8** as a white glass (11.6 g, > 100%): ¹H NMR (500 MHz, CDCl₃) δ 8.92 (br s, 2H), 7.55 (s, 1H), 6.71 (s, 1H), 5.18 (br s, 2H), 4.78 (d, 2H, *J* = 12.0 Hz), 4.72 (d, 2H, *J* = 12.0 Hz), 4.50 (br s, 2H), 4.22-4.19 (m, 4H), 4.00 (br s, 2H), 3.78 (s, 6H), 3.76-3.52 (m, 6H), 2.32-2.30 (m, 2H), 2.21-2.17 (m, 2H), 2.09-2.04 (m, 2H) 1.94 (s, 6H); ¹³C NMR (125.8 MHz, CDCl₃) δ 170.4, 152.2, 149.8, 145.0, 111.3, 106.5, 95.6, 74.4, 72.5, 65.4, 64.1, 58.7, 56.5, 56.3, 33.6, 29.1, 21.1.


(h) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11*S*,11*aS*,2*R*)-10-(2,2,2-trichloroethoxycarbonyl)-11-hydroxy-7-methoxy-2-oxyacetyl-1,2,3,10,11,11*a*-hexahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]]] (**9**)



TEMPO (0.69 g, 4.42 mmol, 0.4 eq) and BAIB (15.7 g, 48.7 mmol, 4.4 eq) were added to a stirred solution of diol **8** (11.5 g, 11.1 mmol) in DCM (150 mL). The reaction mixture was allowed to stir for 2 h and diluted with DCM (400 mL), washed with saturated NaHSO₃ (500 mL), saturated NaHCO₃ (500 mL), brine (200 mL) and dried (MgSO₄). Filtration and evaporation of the solvent afforded the crude product which was purified by flash column chromatography (99.9:0.1 v/v CHCl₃/MeOH then gradient to 99.7:0.3 v/v CHCl₃/MeOH) to provide the product **9** as a light yellow glass (4.43 g, 39%): ¹H NMR (400 MHz, CDCl₃) δ 7.28 (s, 2H, **H6**), 6.84 (s, 2H, **H9**), 5.68 (d, 2H, *J* = 9.1 Hz, **H11**), 5.37-5.35 (m, 2H, **H2**), 5.18 (d, 2H, *J* = 12.0 Hz, **Troc CH₂**), 4.32-4.21 (m, 6H, **OCH₂CH₂CH₂O**, **Troc CH₂**), 4.03 (dd, 2H, *J* = 13.2, 2.6 Hz, **H3**), 3.92 (s, 6H, **OCH₃ x 2**), 3.39-3.69 (m, 4H, **H3** and **H11a**), 2.39-2.35 (m, 6H, **OCH₂CH₂CH₂O** and **H1**), 2.03 (s, 6H, **CH₃CO₂ x 2**); ¹³C NMR (100.6 MHz, CDCl₃) δ 170.4 (**CH₃CO₂**), 167.4 (**C_{quat}**), 154.3 (**C_{quat}**), 150.5 (**C_{quat}**), 149.1 (**C_{quat}**), 127.4 (**C_{quat}**), 124.9 (**C_{quat}**), 114.1 (**C9**), 110.9 (**C6**), 95.0 (**Troc CCl₃**), 87.5 (**C11**), 75.0 (**Troc CH₂**), 71.4 (**C2**), 65.5 (**OCH₂CH₂CH₂O**), 58.4 (**C11a**), 56.1 (**OCH₃**), 51.1 (**C3**), 35.8


(C1), 29.1 (OCH₂CH₂CH₂O), 21.0 (CH₃CO₂); MS (ES), *m/z* (relative intensity) 1058 ([M + Na]⁺, 100).

5 (i) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11S,11aS,2R)-10-(2,2,2-trichloroethoxycarbonyl)-11-(tert-butyl dimethylsilyloxy)-7-methoxy-2-oxyacetyl-1,2,3,10,11,11a-hexahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one]]] (**10**)



TBSOTf (2.70 mL, 3.10 g, 11.7 mmol, 3.0 eq) was added to a stirred solution of bis-alcohol
 9 (4.05 g, 3.91 mmol) and 2,6-lutidine (1.82 mL, 1.68 g, 15.6 mmol, 4.0 eq) in DCM (50
 10 mL). The reaction mixture was allowed to stir for 2.5 h and diluted with DCM (150 mL),
 washed with saturated CuSO₄ (2 X 100 mL), saturated NaHCO₃ (100 mL), brine (200 mL)
 and dried (MgSO₄). Filtration and evaporation of the solvent afforded the crude product
 which was purified by flash column chromatography (99.9:0.1 v/v CHCl₃/MeOH) to provide
 the product **10** as a white glass (5.05 g, >100%): ¹H NMR (400 MHz, CDCl₃) δ 7.05 (s, 2H,
 15 **H6**), 6.52 (s, 2H, **H9**), 5.53 (d, 2H, *J* = 9.0 Hz, **H11**), 5.14 (br s, 2H, **H2**), 4.99 (d, 2H, *J* =
 12.0 Hz, **Troc CH₂**), 4.06-3.87 (m, 8H, **OCH₂CH₂CH₂O**, **Troc CH₂** and **H11a**), 3.71 (s, 6H,
OCH₃ x 2), 3.48-3.43 (m, 4H, **H3**), 2.21-2.11 (m, 4H, **OCH₂CH₂CH₂O** and **H1**), 2.03-1.96
 (m, 2H, **H1**), 1.81 (s, 6H, **CH₃CO₂** x 2), 0.63 (s, 18H, **TBS CH₃** x 6), 0.00 (s x 2, 12H, **TBS**
 20 **CH₃** x 4); ¹³C NMR (100.6 MHz, CDCl₃) δ 170.3 (**CH₃CO₂**), 167.9 (**C_{quat}**), 153.6 (**C_{quat}**),
 150.4 (**C_{quat}**), 149.2 (**C_{quat}**), 127.9 (**C_{quat}**), 125.5 (**C_{quat}**), 113.9 (**C9**), 110.7 (**C6**), 95.2 (**Troc**
CCl₃), 88.2 (**C11**), 74.7 (**Troc CH₂**), 71.7 (**C2**), 65.0 (**OCH₂CH₂CH₂O**), 60.5 (**C11a**), 56.1
 (**OCH₃**), 51.2 (**C3**), 36.2 (**C1**), 28.8 (**OCH₂CH₂CH₂O**), 25.6 (**TBS CH₃**), 21.0 (**CH₃CO₂**),
 17.8 (**TBS C_{quat}**), 14.2 and 14.1 (**TBS CH₃**); MS (ES), *m/z* (relative intensity) 1285 ([M +
 25 21]⁺, 100), 1265 ([M + H]⁺, 75).

(j) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11S,11aS,2R)-10-(2,2,2-trichloroethoxycarbonyl)-11-(tert-butyl dimethylsilyloxy)-7-methoxy-2-hydroxy-1,2,3,10,11,11a-hexahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one]]] (**11**)




5 A solution of K_2CO_3 (93 mg, 0.67 mmol, 5.0 eq) in H_2O (2 mL) was added dropwise to a stirred solution of acetate **10** (170 mg, 0.13 mmol) in MeOH (3 mL). The initial colorless solution eventually turned yellow and the formation of a white precipitate was observed. The reaction mixture was allowed to stir for 16 h when TLC (95:5 v/v $CHCl_3/MeOH$) showed the complete consumption of the starting material. Excess solvent was removed by rotary evaporation and the mixture was carefully neutralized with 1N HCl to pH 7. The resulting mixture was extracted with EtOAc (3 x 25 mL) and the combined organic layers were then washed with brine (40 mL) and dried ($MgSO_4$). Filtration and removal of the solvent afforded the product **11** as a white glass (151 mg, 95%): 1H NMR (400 MHz, $CDCl_3$) δ 6.94 (s, 2H, **H6**), 6.52 (s, 2H, **H9**), 5.53 (d, 2H, $J = 9.0$ Hz, **H11**), 5.00 (d, 2H, $J = 12.0$ Hz, **Troc CH₂**), 4.36-4.35 (m, 2H, **H2**), 4.06-3.82 (m, 8H, **OCH₂CH₂CH₂O**, **Troc CH₂** and **H3**), 3.61 (s, 6H, **OCH₃ x 2**), 3.54-3.48 (m, 2H, **H11a**), 3.39-3.34 (m, 2H, **H3**), 2.96 and 2.95 (br s x 2, 2H, **OH x 2**), 2.21-2.20 (m, 2H, **OCH₂CH₂CH₂O**), 2.19-2.08 (m, 2H, **H1**), 1.90-1.74 (m, 2H, **H1**), 0.64 (s, 18H, **TBS CH₃ x 6**), 0.00 (s, 12H, **TBS CH₃ x 4**); ^{13}C NMR (100.6 MHz, $CDCl_3$) δ 168.5 (**C_{quat}**), 153.6 (**C_{quat}**), 150.3 (**C_{quat}**), 149.1 (**C_{quat}**), 127.9 (**C_{quat}**), 125.4 (**C_{quat}**), 113.9 (**C9**), 110.7 (**C6**), 95.2 (**Troc CCl₃**), 88.3 (**C11**), 74.7 (**Troc CH₂**), 69.4 (**C2**), 65.0 (**OCH₂CH₂CH₂O**), 60.9 (**C11a**), 55.9 (**OCH₃**), 54.1 (**C3**), 38.8 (**C1**), 28.9 (**OCH₂CH₂CH₂O**), 25.6 (**TBS CH₃**), 17.8 (**TBS C_{quat}**); MS (ES), m/z (relative intensity) 1196 ($[M + 16]^+$, 100), 1181 ($[M + H]^+$, 82).

10

15

20


(k) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11S,11aS)-10-(2,2,2-trichloroethoxycarbonyl)-11-(tert-butyl)dimethylsilyloxy)-7-methoxy-2-oxo-1,2,3,10,11,11a-hexahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one]]] (**12**)



12

- 5 A solution of anhydrous DMSO (0.82 mL, 0.90 g, 11.5 mmol, 6.0 eq) in dry DCM (20 mL) was added dropwise to a stirred solution of oxalyl chloride (2.88 mL of a 2 M solution in DCM, 5.76 mmol, 3.0 eq) under a nitrogen atmosphere at -60°C (liq N_2/CHCl_3). After stirring at -55°C for 1.5 h, a solution of the substrate **11** (2.26 g, 1.92 mmol) in dry DCM (30 mL) was added dropwise to the reaction mixture, which was then stirred for a further 2 h at
- 10 -45°C . A solution of TEA (10.8 mL, 7.82 g; 71.7 mmol, 4.2 eq) in dry DCM (90 mL) was added dropwise to the mixture and stirred for a further 30 min. The reaction mixture was left to warm to 0°C , washed with cold 1 N HCl (2 x 50 mL), H_2O (50 mL), brine (50 mL) and dried (MgSO_4). Filtration and evaporation of the solvent *in vacuo* afforded the crude product which was purified by flash column chromatography (70:30 v/v hexane/EtOAc then
- 15 gradient to 40:60 v/v hexane/EtOAc) to afford carbinolamine **12** as a white glass (1.62 g, 72%): ^1H NMR (400 MHz, CDCl_3) δ 7.02 (s, 2H, **H6**), 6.54 (s, 2H, **H9**), 5.59 (d, 2H, $J = 9.2$ Hz, **H11**), 4.98 (d, 2H, $J = 12.0$ Hz, **Troc CH₂**), 4.09-3.86 (m, 8H, **OCH₂CH₂CH₂O**, **Troc CH₂** and **H3**), 3.75-3.66 (m, 10H, **OCH₃** x 2, **H11a**, and **H3**), 2.72 (dd, 2H, $J = 10.2, 19.6$ Hz, **H1**), 2.82 (dd, 2H, $J = 2.6, 19.6$ Hz, **H1**), 2.22-2.19 (m, 2H, **OCH₂CH₂CH₂O**), 0.63 (s,
- 20 18H, **TBS CH₃** x 6), 0.00 (s x 2, 12H, **TBS CH₃** x 4); ^{13}C NMR (100.6 MHz, CDCl_3) δ 207.7 (**C2**), 168.0 (**C_{quat}**), 153.7 (**C_{quat}**), 150.7 (**C_{quat}**), 149.4 (**C_{quat}**), 127.8 (**C_{quat}**), 124.6 (**C_{quat}**), 114.0 (**C9**), 110.6 (**C6**), 95.1 (**Troc CCl₃**), 87.4 (**C11**), 74.8 (**Troc CH₂**), 65.0 (**OCH₂CH₂CH₂O**), 58.9 (**C11a**), 56.1 (**OCH₃**), 53.0 (**C3**), 40.3 (**C1**), 28.8 (**OCH₂CH₂CH₂O**), 25.6 (**TBS CH₃**), 17.8 (**TBS C_{quat}**); MS (ES), m/z (relative intensity) 1224 ($[\text{M} + 48]^+$, 100),
- 25 1210 ($[\text{M} + 34]^+$, 60), 1199 ($[\text{M} + \text{Na}]^+$, 35), 1192 ($[\text{M} + 16]^+$, 40), 1176 (M^+ , 18).

(l) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11S,11aS)-10-(2,2,2-trichloroethoxycarbonyl)-11-(tert-butyl)dimethylsilyloxy)-7-methoxy-2-[[[(trifluoromethyl)sulfonyl]oxy]-1,10,11,11a-tetrahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one]]] (13)



5 Anhydrous triflic anhydride (3.09 mL, 5.19 g, 18.4 mmol, 22 eq) taken from a freshly opened ampule was added rapidly in one portion to a vigorously stirred solution of ketone **12** (0.98 g, 0.84 mmol) and anhydrous pyridine (1.49 mL, 1.46 g, 18.4 mmol, 22 eq) in dry DCM (50 mL) at room temperature under a nitrogen atmosphere. The initial precipitate dissolved gradually and the solution eventually turned a dark red colour. The reaction


10 mixture was allowed to stir for 4.5 h when TLC (80:20 v/v EtOAc/hexane) revealed the complete consumption of the starting material. The mixture was poured into cold saturated NaHCO₃ (60 mL) and extracted with DCM (3 x 80 mL). The combined organic layers were then washed with saturated CuSO₄ (2 x 125 mL), brine (125 mL) and dried (MgSO₄). Filtration and evaporation of the solvent afforded the crude product which was purified by

15 flash column chromatography (80:20 v/v hexane/EtOAc) to afford triflate **13** as a light yellow glass (0.74 mg, 61%): $[\alpha]_D^{25} = +46.0^\circ$ ($c = 0.33$, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.23 (s, 2H, **H6**), 7.19 (s, 2H, **H3**), 6.77 (s, 2H, **H9**), 5.94 (d, 2H, $J = 8.9$ Hz, **H11**), 5.23 (d, 2H, $J = 12.0$ Hz, **Troc CH₂**), 4.31-4.28 (m, 2H, **OCH₂CH₂CH₂O**), 4.18 (d, 2H, $J = 12.2$ Hz, **Troc CH₂**), 4.15-4.13 (m, 2H, **OCH₂CH₂CH₂O**), 3.95-3.91 (m, 8H, **OCH₃ x 2, H11a**), 3.35

20 (dd, 2H, $J = 11.0, 16.6$ Hz, **H1**), 2.84 (d, 2H, $J = 16.6$ Hz, **H1**), 2.46-2.44 (m, 2H, **OCH₂CH₂CH₂O**), 0.89 (s, 18H, **TBS CH₃ x 6**), 0.29 and 0.26 (s x 2, 12H, **TBS CH₃ x 4**); ¹³C NMR (100.6 MHz, CDCl₃) δ 164.9 (**C_{quat}**), 153.6 (**C_{quat}**), 151.0 (**C_{quat}**), 149.5 (**C_{quat}**), 136.0 (**C_{quat}**), 127.7 (**C_{quat}**), 123.9 (**C_{quat}**), 121.0 (**C3**), 114.0 (**C9**), 110.9 (**C6**), 95.1 (**Troc CCl₃**), 86.3 (**C11**), 74.8 (**Troc CH₂**), 65.0 (**OCH₂CH₂CH₂O**), 60.6 (**C11a**), 56.2 (**OCH₃**), 34.4 (**C1**),


25 28.8 (**OCH₂CH₂CH₂O**), 25.6 (**TBS CH₃**), 17.8 (**TBS C_{quat}**); IR (CHCl₃) 3020, 2957, 2860, 1725, 1674, 1651, 1604, 1516, 1466, 1454, 1431, 1409, 1329, 1312, 1274, 1216, 1138, 1113, 1083, 1042, 1006, 900, 840, 757, 668, 646, 610 cm⁻¹; MS (ES), m/z (relative intensity) 1461 ($[M + 21]^+$, 100), 1440 (M^+ , 55).

(m) 1,1'-[[*(Propane-1,3-diyl)dioxy*]]bis[(11*S*,11*aS*)-10-(2,2,2-trichloroethoxycarbonyl)-11-(*tert*-butyldimethylsilyloxy)-7-methoxy-2-(*p*-methoxybenzene)-1,10,11,11*a*-tetrahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]] (**14**)




5 A solution of TEA (0.20 mL, 148 mg, 1.46 mmol, 6.0 eq) in H₂O (1.5 mL) and EtOH (10 mL) was added to a solution of triflate **13** (350 mg, 0.24 mmol) in toluene (10 mL) at room temperature. To this mixture 4-methoxybenzeneboronic acid (96 mg, 0.63 mmol, 2.6 eq) and Pd(PPh₃)₄ (11 mg, 9 μmol, 0.04 eq) were added. The reaction mixture was allowed to stir for 15 min when TLC (80:20 v/v EtOAc/hexane) revealed the complete consumption of the starting material. Excess solvent was removed and the residue was dissolved in EtOAc (25 mL), washed with H₂O (15 mL), brine (15 mL) and dried (MgSO₄). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (80:20 v/v hexane/EtOAc then gradient to 50:50 v/v hexane/EtOAc) to afford **14** as a yellow glass (286 mg, 87%): ¹H NMR (400 MHz, CDCl₃) δ 7.38 (s, 2H, **H3**), 7.32-7.28 (m, 6H, **H6** and **H13**), 6.92 (d, 4H, *J* = 8.7 Hz, **H14**), 6.81 (s, 2H, **H9**), 5.93 (d, 2H, *J* = 8.8 Hz, **H11**), 5.24 (d, 2H, *J* = 12.0 Hz, **Troc CH₂**), 4.34-4.29 (m, 2H, **OCH₂CH₂CH₂O**), 4.20-4.11 (m, 4H, **Troc CH₂** and **OCH₂CH₂CH₂O**), 4.00-3.96 (m, 8H, **H11a** and **OCH₃ x 2**), 3.84 (s, 6H, **OCH₃ x 2**), 3.36 (dd, 2H, *J* = 10.8, 16.6 Hz, **H1**), 2.85 (d, 2H, *J* = 16.5 Hz, **H1**), 2.48-2.45 (m, 2H, **OCH₂CH₂CH₂O**), 0.93 (s, 18H, **TBS CH₃ x 6**), 0.30 and 0.27 (s x 2, 12H, **TBS CH₃ x 4**); ¹³C NMR (100.6 MHz, CDCl₃) δ 162.5 (**C_{quat}**), 161.3 (**C_{quat}**), 159.2 (**C_{quat}**), 151.1 (**C_{quat}**), 148.1 (**C_{quat}**), 140.3 (**C_{quat}**), 126.2 (**C13**), 126.0 (**C_{quat}**), 123.2 (**C_{quat}**), 121.9 (**C3**), 119.3 (**C_{quat}**), 114.3 (**C6**), 111.9 (**C14**), 111.2 (**C9**), 95.2 (**Troc CCl₃**), 87.3 (**C11**), 74.8 (**Troc CH₂**), 65.0 (**OCH₂CH₂CH₂O**), 61.5 (**C11a**), 56.1 and 55.3 (**OCH₃**), 35.3 (**C1**), 28.8 (**OCH₂CH₂CH₂O**), 25.7 (**TBS CH₃**), 17.9 (**TBS C_{quat}**); MS (ES), *m/z* (relative intensity) 1357 (**M⁺**, 63), 1114 (48), 955 (59), 919 (78).

(n) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11a*S*)-7-methoxy-2-(*p*-methoxybenzene)-1,11a-dihydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepine-5-one]]] (**ZC-207**)



10% Cd/Pd couple (461 mg, 3.73 mmol, 20 eq) was added to a rapidly stirring mixture of
 5 **14** (253 mg, 0.19 mmol), THF (5 mL) and 1 N NH₄OAc (5 mL). The reaction mixture was
 allowed to stir for 1.5 h when TLC showed the complete consumption of the starting
 material. The solids were filtered and rinsed with H₂O and DCM. The aqueous layer was
 extracted with DCM (3 x 30 mL) and the organic extracts were combined, washed with
 brine (50 mL) and dried (MgSO₄). Filtration and evaporation of solvent afforded the crude
 10 product which was purified by flash column chromatography (99.9:0.1 v/v CHCl₃/MeOH
 then gradient to 95:5 v/v CHCl₃/MeOH) to afford **ZC-207** as a yellow glass (132 mg, 96%):
 [α]_D²⁰ = +880.0 ° (c = 0.22, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.79 (d, 2H, *J* = 3.9 Hz,
H11), 7.44 (s, 2H, **H6**), 7.30 (s, 2H, **H3**), 7.24 (d, 4H, *J* = 8.7 Hz, **H13**), 6.81 (d, 4H, *J* = 8.7
 Hz, **H14**), 6.79 (s, 2H, **H9**), 4.30-4.18 (m, 6H, **OCH₂CH₂CH₂O** and **H11a**), 3.86 (s, 6H,
 15 **OCH₃** x 2), 3.74 (s, 6H, **OCH₃** x 2), 3.48 (dd, 2H, *J* = 11.8, 16.2 Hz, **H1**), 2.85 (d, 2H, *J* =
 16.2 Hz, **H1**), 2.38-2.32 (m, 2H, **OCH₂CH₂CH₂O**); ¹³C NMR (62.9 MHz, CDCl₃) δ 162.5
 (**C11**), 161.3(**C_{quat}**), 159.2 (**C_{quat}**), 151.1 (**C_{quat}**), 148.1 (**C_{quat}**), 140.3 (**C_{quat}**), 126.2 (**C13**),
 126.0 (**C_{quat}**), 123.2 (**C_{quat}**), 121.9 (**C3**), 114.3 (**C14**), 111.9 (**C9**), 111.2 (**C6**), 65.4
 (**OCH₂CH₂CH₂O**), 56.2 and 55.3 (**OCH₃**), 53.8 (**C11a**), 35.6 (**C1**), 28.9 (**OCH₂CH₂CH₂O**);
 20 MS (ES), *m/z* (relative intensity) 741 (M⁺, 43), 660 (71).

(o) 1,1'[[[(Propane-1,5-diyl)dioxy]bis[(11a*S*)-11-sulpho-7-methoxy-2-(4-methoxyphenyl)-1,10,11,11a-tetrahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]]] sodium salt (**ZC-423**)




25 A solution of sodium bisulphite (6.74 mg, 65 μmol) in water (1.5 mL) was added to a stirred
 solution of **ZC-207** (24.01 mg, 32 μmol) in dichloromethane (1.5 mL). The reaction mixture
 was allowed to stir vigorously for 2 h, after which time the organic and aqueous layers were
 separated. TLC analysis (eluent-EtOAc) of the aqueous phase revealed absence of **ZC-**

207 and presence of baseline material with strong uv absorption. The aqueous layer was lyophilised to provide the bisulphite adduct **ZC-423** as a white solid (17 mg, 55%): ¹H NMR (400 MHz, *d*₆-DMSO) δ 7.42 (s, 2H, **H-3**), 7.38 (d, 4H, *J* = 8.72 Hz, **2'-H**), 7.05 (s, 2H, **H-6**), 6.92 (d, 4H, *J* = 8.92 Hz, **3'-H**), 6.52 (s, 2H, **H-9**), 5.27 (s, 2H, **NH**), 4.35-4.25 (m, 2H, **H11a**), 4.15-4.05 (m, 4H, OCH₂CH₂CH₂O), 3.95 (d, 2H, *J* = 10.4 Hz, **H11**), 3.77 (s, 6H, OMe), 3.72 (s, 6H, OMe), 3.55-3.45 (m, 2H, **H1**), 3.30-3.15 (m, 2H, **H1**), 2.25-2.15 (m, 2H, OCH₂CH₂CH₂O).


Example 2

10 (a) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11*S*,11*aS*)-10-(2,2,2-trichloroethoxycarbonyl)-11-(*tert*-butyldimethylsilyloxy)-7-methoxy-2-(naphthalen-2-yl)-1,10,11,11*a*-tetrahydro-5H-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]]] (**17**)




A solution of TEA (0.15 mL, 1.05 mmol, 6.0 eq) in H₂O (1 mL) and EtOH (10 mL) was added to a solution of **13** (251 mg, 0.17 mmol) in toluene (6 mL) at room temperature. To this mixture 2-naphthaleneboronic acid (77.9 mg, 0.45 mmol, 2.6 eq) and Pd(PPh₃)₄ (8.0 mg, 7 μmol, 0.04 eq) were added. The reaction mixture was heated at 100°C under microwave irradiation for 20 minutes when TLC (80:20 v/v EtOAc/hexane) revealed complete consumption of the starting material. Excess solvent was removed and the residue was dissolved in EtOAc (20 mL), washed with H₂O (15 mL), brine (15 mL) and dried (MgSO₄). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (80:20 v/v hexane/EtOAc then gradient to 50:50 v/v hexane/EtOAc) to afford **17** as a yellow glass (191 mg, 81%): ¹H NMR (400 MHz, CDCl₃) δ 7.32-7.29 (m, 6H, **H_{arom}**), 7.61-7.58 (m, 6H, **H_{arom}**), 7.52-7.42 (m, 4H, **H3** and **H_{arom}**), 7.30 (s, 2H, **H6**), 6.81 (s, 2H, **H9**), 5.98 (d, 2H, *J* = 8.8 Hz, **H11**), 5.24 (d, 2H, *J* = 12.0 Hz, **Troc CH₂**), 4.35-4.30 (m, 2H, **OCH₂CH₂CH₂O**), 4.20-4.13 (m, 4H, **Troc CH₂** and **OCH₂CH₂CH₂O**), 4.08-4.00 (m, 2H, **H11a**), 3.97 (s, 6H, **OCH₃ x 2**), 3.50 (dd, 2H, *J* = 10.8, 16.6 Hz, **H1**), 2.98 (d, 2H, *J* = 16.5 Hz, **H1**), 2.50-2.48 (m, 2H, **OCH₂CH₂CH₂O**), 0.94 (s, 18H, **TBS CH₃ x 6**), 0.30 and 0.28 (s x 2, 12H, **TBS CH₃ x 4**).

(b) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11aS)-7-methoxy-2-(naphthalen-2-yl)-1,11a-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-5-one]]] (**18**)



5 10% Cd/Pb couple (455 mg, 3.71 mmol, 26 eq) was added to a rapidly stirring mixture of
17 (190 mg, 0.14 mmol), THF (3.2 mL) and 1 N NH₄OAc (3.2 mL). The reaction mixture
 was allowed to stir for 6.5 hours when TLC showed the incomplete consumption of the
 starting material and formation of side products. The solids were filtered and rinsed with
 H₂O and MeOH. Excess solvent was removed and the residue was diluted with H₂O (25
 10 mL) and DCM (25 mL). The aqueous layer was extracted with DCM (3 x 25 mL) and the
 organic extracts were combined, washed with brine (50 mL) and dried (MgSO₄). Filtration
 and evaporation of solvent afforded the crude product which was purified by flash column
 chromatography (99:1 v/v CHCl₃/MeOH then gradient to 98:2 v/v CHCl₃/MeOH) to afford
18 as a yellow glass (38 mg, 36%): ¹H NMR (400 MHz, CDCl₃) δ 7.85 (d, 2H, *J* = 3.9 Hz,
 15 **H11**), 7.77-7.65 (m, 6H, **H_{arom}**), 7.59-7.50 (m, 6H, **H3** and **H_{arom}**), 7.47 (s, 2H, **H6**), 7.44-
 7.33 (m, 4H, **H_{arom}**), 6.82 (s, 2H, **H9**), 4.42-4.32 (m, 2H, **H11a**), 4.31-4.14 (m, 4H,
OCH₂CH₂CH₂O), 3.89 (s, 6H, **OCH₃** x 2), 3.69-3.56 (m, 2H, **H1**), 3.50-3.37 (m, 2H, **H1**),
 2.45-2.29 (m, 2H, **OCH₂CH₂CH₂O**).

20 (c) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11aS)-11-sulpho-7-methoxy-2-(naphthalen-2-yl)-1,
 10,11,11a-tetrahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one]]] sodium salt (**19**)




A solution of sodium bisulphite (6.67 mg, 0.064 mmol, 2.0 equiv.) in H₂O (1.5 mL) was
 added to a solution of compound **18** (25 mg, 0.032 mmol) in dry DCM (1.5 mL). The
 25 reaction mixture was allowed to stir for 5 hours. The reaction mixture was diluted with
 DCM (5 mL) and H₂O (5 mL). The aqueous layer was separated (without shaking the
 separating funnel) and dried under the freeze dryer to afford **19** as a lightweight white solid
 (7 mg, 22 %): ¹H NMR (400 MHz, DMSO) δ 7.78-7.72 (m, 8H, **H_{arom}**), 7.68 (s, 2H, **H3**),
 7.59-7.27 (m, 6H, **H_{arom}**), 6.92 (s, 2H, **H6**), 6.41 (s, 2H, **H9**), 5.21 (d, 2H, *J* = 5.2 Hz, **NH**),

4.38-4.31 (m, 2H, **H11a**), 4.12-3.96 (m, 4H, **OCH₂CH₂CH₂O**), 3.87 (d, 2H, $J = 10.4$ Hz, **H11**), 3.70 (s, 3H, **OCH₃**), 3.61 (s, 3H, **OCH₃**), 3.40-3.12 (m, 4H, **H1**), 2.28-2.22 (m, 2H, **OCH₂CH₂CH₂O**).

5 Example 3

(a) 1,1'-[[*(Propane-1,3-diyl)dioxy*]bis[(11*S*,11*aS*)-10-(2,2,2-trichloroethoxycarbonyl)-11-(*tert*-butyldimethylsilyloxy)-7-methoxy-2-(thiophen-2-yl)-1,10,11,11*a*-tetrahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]] (**20**)




10 A solution of TEA (0.11 mL, 0.82 mmol, 6.0 eq) in H₂O (0.8 mL) and EtOH (5 mL) was added to a solution of **13** (197 mg, 0.14 mmol) in toluene (5 mL). To this mixture thiophene-2-boronic acid (45.5 mg, 0.36 mmol, 2.6 eq) and Pd(PPh₃)₄ (6.3 mg, 5 μmol, 0.04 eq) were added. The reaction mixture was allowed to stir at room temperature for 3

15 hours when TLC (80:20 v/v EtOAc/hexane) revealed complete consumption of the starting material. Excess solvent was removed and the residue was dissolved in EtOAc (20 mL), washed with H₂O (15 mL), brine (15 mL) and dried (MgSO₄). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (80:20 v/v hexane/EtOAc) to afford **20** as a yellow glass (168 mg, 94%): ¹H NMR (400

20 MHz, CDCl₃) δ 7.32 (s, 2H, **H3**), 7.26 (s, 2H, **H6**), 7.24-7.19 (m, 2H, **H_{arom}**), 7.04-6.79 (m, 2H, **H_{arom}**), 6.94-6.88 (m, 2H, **H_{arom}**), 6.78 (s, 2H, **H9**), 5.92 (d, 2H, $J = 8.9$ Hz, **H11**), 5.24 (d, 2H, $J = 12.0$ Hz, **Troc CH₂**), 4.36-4.25 (m, 2H, **OCH₂CH₂CH₂O**), 4.19-4.08 (m, 4H, **Troc CH₂** and **OCH₂CH₂CH₂O**), 4.02-3.87 (m, 8H, **H11a** and **OCH₃ x 2**), 3.37 (dd, 2H, $J = 10.8, 16.6$ Hz, **H1**), 2.85 (d, 2H, $J = 16.5$ Hz, **H1**), 2.52-2.37 (m, 2H, **OCH₂CH₂CH₂O**), 0.91 (s, 18H, **TBS CH₃ x 6**), 0.28 and 0.25 (s x 2, 12H, **TBS CH₃ x 4**).


25

(b) 1,1'-[[*(Propane-1,3-diyl)dioxy*]bis[(11*aS*)-7-methoxy-2-(thiophen-2-yl)-1,11*a*-dihydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepine-5-one]] (**21**)



10% Cd/Pb couple (288 mg, 2.35 mmol, 20 eq) was added to a rapidly stirring mixture of **20** (154 mg, 0.12 mmol), THF (3 mL) and 1 N NH₄OAc (3 mL). The reaction mixture was allowed to stir for 3 hours when TLC showed complete consumption of the starting material. The solids were filtered and rinsed with H₂O and DCM. The aqueous layer was extracted with DCM (3 x 15 mL) and the organic extracts were combined, washed with brine (40 mL) and dried (MgSO₄). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (99:1 v/v CHCl₃/MeOH then gradient to 98:2 v/v CHCl₃/MeOH) to afford **21** as a yellow glass (59 mg, 72%): ¹H NMR (400 MHz, CDCl₃) δ 7.90 (d, 2H, *J* = 4.0 Hz, **H11**), 7.51 (s, 2H, **H6**), 7.36 (s, 2H, **H3**), 7.22 (d, 2H, *J* = 5.2 Hz, **H_{arom}**), 7.02 (dd, 2H, *J* = 3.6, 5.0 Hz, **H_{arom}**), 6.98 (d, 2H, *J* = 3.4 Hz, **H_{arom}**), 6.88 (s, 2H, **H9**), 4.43-4.23 (m, 6H, **H11a** and **OCH₂CH₂CH₂O**), 3.94 (s, 6H, **OCH₃** x 2), 3.59 (ddd, 2H, *J* = 2.0, 11.5, 16.0 Hz, **H1**), 2.85 (ddd, 2H, *J* = 1.5, 5.2, 16.0 Hz, **H1**), 2.49-2.40 (m, 2H, **OCH₂CH₂CH₂O**).


(c) 1,1'-[[*(Propane-1,3-diyl)dioxy*]bis[(11a*S*)-11-sulpho-7-methoxy-2-(thiophen-2-yl)-1,10,11,11a-tetrahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]] sodium salt (**22**)



A solution of sodium bisulphite (16.5 mg, 0.159 mmol, 2.0 equiv.) in H₂O (3.5 mL) was added to a solution of compound **21** (55.0 mg, 0.079 mmol) in DCM (3.5 mL). The reaction mixture was allowed to stir for 4 hours. TLC (EtOAc) revealed complete consumption of the starting material and a product spot at the base line. The reaction mixture was diluted with DCM and H₂O. The aqueous layer was separated (without shaking the separating funnel) and dried under the freeze dryer to afford **22** as a lightweight yellow solid (53.9 mg, 75 %): ¹H NMR (400 MHz, DMSO) δ 7.42 (dd, 2H, *J* = 1.5, 4.7 Hz, **H_{arom}**), 7.33-7.30 (m, 6H, **H_{arom}**), 6.53 (s, 2H, **H9**), 5.30 (d, 2H, *J* = 3.8 Hz, **NH**), 4.35-4.29 (m, 2H, **H11a**), 4.12-4.06 (m, 4H, **OCH₂CH₂CH₂O**), 3.95 (d, 2H, *J* = 10.3 Hz, **H11**), 3.72 (s, 6H, **OCH₃** x 2), 3.53-3.19 (m, 4H, **H1**), 2.23-2.21 (m, 2H, **OCH₂CH₂CH₂O**).

Example 4

(a) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11*S*,11*aS*)-10-(2,2,2-trichloroethoxycarbonyl)-11-(*tert*-butyldimethylsilyloxy)-7-methoxy-2-(quinolin-6-yl)-1,10,11,11*a*-tetrahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]]] (**23**)




5

A solution of TEA (0.12 mL, 0.84 mmol, 6.0 eq) in H₂O (0.8 mL) and EtOH (5 mL) was added to a solution of triflate **13** (202 mg, 0.14 mmol) in toluene (5 mL) at room temperature. To this mixture 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl) quinoline (93.0 mg, 0.36 mmol, 2.6 eq) and Pd(PPh₃)₄ (6.5 mg, 6 μmol, 0.04 eq) were added. The reaction mixture was heated at 100°C under microwave irradiation for 15 minutes when TLC (80:20 v/v EtOAc/hexane) revealed complete consumption of the starting material. Excess solvent was removed and the residue was dissolved in EtOAc (20 mL), washed with H₂O (15 mL), brine (15 mL) and dried (MgSO₄). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (80:20 v/v hexane/EtOAc then gradient to 50:50 v/v hexane/EtOAc) to afford **23** as a yellow glass (126 mg, 64%): ¹H NMR (400 MHz, CDCl₃) δ 8.88 (dd, 2H, *J* = 1.6, 4.2 Hz, H_{arom}), 8.14 (d, 2H, *J* = 7.6 Hz, H_{arom}), 8.07 (d, 2H, *J* = 9.0 Hz, H_{arom}), 7.85 (dd, 2H, *J* = 1.8, 8.9 Hz, H_{arom}), 7.66 (s, 2H, H₃), 7.55 (d, 2H, *J* = 1.8 Hz, H_{arom}), 7.41 (dd, 2H, *J* = 4.3, 8.3 Hz, H_{arom}), 7.31 (s, 2H, H₆), 6.82 (s, 2H, H₉), 5.98 (d, 2H, *J* = 8.8 Hz, H₁₁), 5.25 (d, 2H, *J* = 12.1 Hz, Troc CH₂), 4.34-4.31 (m, 2H, OCH₂CH₂CH₂O), 4.21-4.15 (m, 4H, Troc CH₂ and OCH₂CH₂CH₂O), 4.09-4.03 (m, 2H, H_{11a}), 3.97 (s, 6H, OCH₃ x 2), 3.48 (ddd, 2H, *J* = 1.6, 10.4, 16.1 Hz, H₁), 2.98 (dd, 2H, *J* = 2.4, 16.1 Hz, H₁), 2.50-2.47 (m, 2H, OCH₂CH₂CH₂O), 0.95 (s, 18H, TBS CH₃ x 6), 0.32 and 0.28 (s x 2, 12H, TBS CH₃ x 4).

15

20


(b) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11aS)-7-methoxy-2-(quinolin-6-yl)-1,11a-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-5-one]]] (**24**)



5
10% Cd/Pb couple (516 mg, 4.21 mmol, 25 eq) was added to a rapidly stirring mixture of **23** (234 mg, 0.17 mmol), THF (5 mL) and 1 N NH₄OAc (5 mL). The reaction mixture was allowed to stir for 5 hours when TLC showed complete consumption of the starting material and formation of a small amount of side products. The solids were filtered and rinsed with H₂O and a large amount of MeOH. Excess solvent was removed and the residue was diluted with H₂O (35 mL) and CHCl₃ (35 mL). The aqueous layer was extracted with CHCl₃ (3 x 30 mL) and the organic extracts were combined, washed with brine (100 mL) and dried (MgSO₄). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (99.5:0.5 v/v CHCl₃/MeOH then gradient to 96:4 v/v CHCl₃/MeOH) to afford **24** as a yellow glass (44 mg, 33%): ¹H NMR (400 MHz, CDCl₃) δ 8.85 (dd, 2H, *J* = 1.6, 4.2 Hz, H_{arom}), 8.09 (d, 2H, *J* = 7.6 Hz, H_{arom}), 8.03 (d, 2H, *J* = 9.0 Hz, H_{arom}), 7.88 (d, 2H, *J* = 3.9 Hz, H₁₁), 7.82 (dd, 2H, *J* = 1.8, 8.9 Hz, H_{arom}), 7.24 (s, 2H, H₃), 7.59 (s, 2H, H_{arom}), 7.52 (s, 2H, H₆), 7.48 (dd, 2H, *J* = 4.3, 8.3 Hz, H_{arom}), 6.88 (s, 2H, H₉), 4.48-4.42 (m, 2H, H_{11a}), 4.39-4.28 (m, 4H, OCH₂CH₂CH₂O), 3.94 (s, 6H, OCH₃ x 2), 3.76-3.63 (m, 2H, H₁), 3.09-2.98 (m, 2H, H₁), 2.46-2.43 (m, 2H, OCH₂CH₂CH₂O).

Example 5


(a) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11S,11aS)-10-(2,2,2-trichloroethoxycarbonyl)-11-(tert-butylidimethylsilyloxy)-7-methoxy-2-(3-methoxybenzene)-1,10,11,11a-tetrahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one]]] (**25**)



A solution of TEA (0.071 mL, 0.51 mmol, 6.0 eq) in H₂O (0.6 mL) and EtOH (3.5 mL) was added to a solution of triflate **13** (122 mg, 0.085 mmol) in toluene (3.5 mL) at room


temperature. To this mixture 3-methoxyphenolboronic acid (33.5 mg, 0.22 mmol, 2.6 eq) and Pd(PPh₃)₄ (3.9 mg, 3.4 μmol, 0.04 eq) were added. The reaction mixture was heated at 100°C under microwave irradiation for 5 minutes when TLC (80:20 v/v EtOAc/hexane) revealed complete consumption of the starting material. Excess solvent was removed and the residue was dissolved in EtOAc (10 mL), washed with H₂O (10 mL), brine (10 mL) and dried (MgSO₄). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (80:20 v/v hexane/EtOAc then gradient to 60:40 v/v hexane/EtOAc) to afford **25** as a yellow glass (66 mg, 57%): ¹H NMR (400 MHz, CDCl₃) δ 7.23 (s, 2H, **H3**), 7.05-7.00 (m, 4H, **H6** and **H_{arom}**), 6.70 (d, 2H, *J* = 8.7 Hz, **H_{arom}**), 6.64 (s, 2H, **H_{arom}**), 6.67-6.64 (m, 4H, **H_{arom}** and **H9**), 5.67 (d, 2H, *J* = 8.8 Hz, **H11**), 4.98 (d, 2H, *J* = 12.0 Hz, **Troc CH₂**), 4.08-4.02 (m, 2H, **OCH₂CH₂CH₂O**), 3.92-3.84 (m, 4H, **Troc CH₂** and **OCH₂CH₂CH₂O**), 3.74-3.66 (m, 8H, **H11a** and **OCH₃ x 2**), 3.68 (s, 6H, **OCH₃ x 2**), 3.10 (dd, 2H, *J* = 10.8, 16.6 Hz, **H1**), 2.60 (d, 2H, *J* = 16.5 Hz, **H1**), 2.22-2.19 (m, 2H, **OCH₂CH₂CH₂O**), 0.68 (s, 18H, **TBS CH₃ x 6**), 0.02 and 0.00 (s x 2, 12H, **TBS CH₃ x 4**).

(b) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11a*S*)-7-methoxy-2-(3-methoxybenzene)-1,11a-dihydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepine-5-one]]] (**26**)



10% Cd/Pb couple (269 mg, 2.20 mmol, 25 eq) was added to a rapidly stirring mixture of **25** (119 mg, 0.088 mmol), THF (3 mL) and 1 N NH₄OAc (3 mL). The reaction mixture was allowed to stir for 3.5 hours when TLC showed complete consumption of the starting material. The solids were filtered and rinsed with H₂O and MeOH. Excess solvent was removed and the residue was diluted with H₂O (20 mL) and DCM (20 mL). The aqueous layer was extracted with DCM (3 x 20 mL) and the organic extracts were combined, washed with H₂O (50 mL) and brine (50 mL) and dried (MgSO₄). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (99.9:0.1 v/v CHCl₃/MeOH then gradient to 98:2 v/v CHCl₃/MeOH) to afford **26** as a yellow glass (35 mg, 54%): ¹H NMR (400 MHz, CDCl₃) δ 7.79 (d, 2H, *J* = 3.9 Hz, **H11**), 7.44 (s, 2H, **H3**), 7.42 (s, 2H, **H6**), 7.21-7.16 (m, 2H, **H_{arom}**), 6.90 (d, 2H, *J* = 8.7 Hz, **H_{arom}**), 6.84-6.83 (m, 2H, **H_{arom}**), 6.80 (s, 2H, **H9**), 4.33-4.17 (m, 6H, **OCH₂CH₂CH₂O** and **H11a**), 3.86 (s, 6H, **OCH₃ x 2**), 3.75 (s, 6H, **OCH₃ x 2**), 3.44 (dd, 2H, *J* = 11.8, 16.2 Hz, **H1**), 3.30 (d, 2H, *J* = 16.2 Hz, **H1**), 2.39-2.33 (m, 2H, **OCH₂CH₂CH₂O**).

(c) 1,1'-[[*(Propane-1,3-diyl)dioxy*]]bis[(11*aS*)-11-sulpho-7-methoxy-2-(3-methoxybenzene)-1, 10,11,11*a*-tetrahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]] sodium salt (**27**)



5

A solution of sodium bisulphite (7.6 mg, 0.073 mmol, 2.0 equiv.) in H₂O (1.7 mL) was added to a solution of compound **26** (27.0 mg, 0.036 mmol) in DCM (1.7 mL). The reaction mixture was allowed to stir for 2 hours. TLC (EtOAc) revealed complete consumption of the starting material and a product spot at the base line. The reaction mixture was diluted with DCM (5 mL) and H₂O (5 mL). The aqueous layer was separated and lyophilized to afford **27** as a lightweight white solid (12.1 mg, 35 %): ¹H NMR (400 MHz, DMSO) δ 7.61 (s, 2H, **H3**), 7.27-7.23 (m, 2H, **H_{arom}**), 7.08-6.99 (m, 6H, **H_{arom}** and **H6**), 6.81-6.78 (m, 2H, **H_{arom}**), 6.54 (s, 2H, **H9**), 5.30 (s, 2H, **NH**), 4.37-4.30 (m, 2H, **H11*a***), 4.17-4.08 (m, 4H, **OCH₂CH₂CH₂O**), 3.84 (d, 2H, *J* = 10.4 Hz, **H11**), 3.78 (s, 6H, **OCH₃ x 2**), 3.62 (s, 6H, **OCH₃ x 2**), 3.52-3.20 (m, 4H, **H1**), 2.26-2.20 (m, 2H, **OCH₂CH₂CH₂O**).


10

15

Example 6

(a) 1,1'-[[*(Propane-1,3-diyl)dioxy*]]bis[(11*S*,11*aS*)-10-(2,2,2-trichloroethoxycarbonyl)-11-(*tert*-butyldimethylsilyloxy)-7-methoxy-2-(3,4-methylenedioxyphenyl)-1,10,11,11*a*-tetrahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]] (**28**)

20




A solution of TEA (0.066 mL, 0.47 mmol, 6.0 eq) in H₂O (0.6 mL) and EtOH (3.5 mL) was added to a solution of triflate **13** (122 mg, 0.085 mmol) in toluene (3.5 mL) at room temperature. To this mixture 3,4-methylenedioxyphenylboronic acid (34.2 mg, 0.21 mmol, 2.6 eq) and Pd(PPh₃)₄ (3.6 mg, 3.2 μmol, 0.04 eq) were added. The reaction mixture was heated at 100°C under microwave irradiation for 5 minutes when TLC (80:20 v/v EtOAc/hexane) revealed complete consumption of the starting material. Excess solvent was removed and the residue was dissolved in EtOAc (10 mL), washed with H₂O (10 mL),

25


brine (10 mL) and dried (MgSO_4). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (80:20 v/v hexane/EtOAc then gradient to 65:35 v/v hexane/EtOAc) to afford **28** as a yellow glass (84 mg, 77%): $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.34 (s, 2H, **H6**), 7.26 (s, 2H, **H3**), 6.89 (s, 2H, **H9**), 6.78-6.76 (m, 4H, **H_{arom}**), 5.98 (s, 2H, **OCH₂O**), 5.89 (d, 2H, $J = 8.8$ Hz, **H11**), 5.23 (d, 2H, $J = 12.0$ Hz, **Troc CH₂**), 4.32-4.27 (m, 2H, **OCH₂CH₂CH₂O**), 4.18-4.09 (m, 4H, **Troc CH₂** and **OCH₂CH₂CH₂O**), 3.97-3.91 (m, 8H, **H11a** and **OCH₃ x 2**), 3.30 (dd, 2H, $J = 10.8, 16.6$ Hz, **H1**), 2.76 (d, 2H, $J = 16.6$ Hz, **H1**), 2.46-2.44 (m, 2H, **OCH₂CH₂CH₂O**), 0.88 (s, 18H, **TBS CH₃ x 6**), 0.28 and 0.25 (s x 2, 12H, **TBS CH₃ x 4**).

(b) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11a*S*)-7-methoxy-2-(3,4-methylenedioxyphenyl)-1,11a-dihydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepine-5-one]] (**29**)



10% Cd/Pb couple (307 mg, 2.51 mmol, 25 eq) was added to a rapidly stirring mixture of **28** (139 mg, 0.101 mmol), THF (3.5 mL) and 1 N NH_4OAc (3.5 mL). The reaction mixture was allowed to stir for 5 hours when TLC showed complete consumption of the starting material and formation of a small amount of side products. The solids were filtered and rinsed with H_2O and MeOH. Excess solvent was removed and the residue was diluted with H_2O (25 mL) and DCM (25 mL). The aqueous layer was extracted with DCM (3 x 40 mL) and the organic extracts were combined, washed with H_2O (80 mL) and brine (80 mL) and dried (MgSO_4). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (99.9:0.1 v/v $\text{CHCl}_3/\text{MeOH}$ then gradient to 98:2 v/v $\text{CHCl}_3/\text{MeOH}$) to afford **26** as a yellow glass (30 mg, 39%): $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.86 (d, 2H, $J = 3.9$ Hz, **H11**), 7.60 (s, 2H, **H6**), 7.36 (s, 2H, **H3**), 6.91 (d, 2H, $J = 0.9$ Hz, **H_{arom}**), 6.86 (s, 2H,), 6.82-6.78 (m, 4H, **H9** and **H_{arom}**), 5.97 (s, 2H, **OCH₂O**), 4.36-4.25 (m, 6H, **H11a** and **OCH₂CH₂CH₂O**), 3.93 (s, 6H, **OCH₃ x 2**), 3.52 (dd, 2H, $J = 10.8, 16.6$ Hz, **H1**), 3.33 (d, 2H, $J = 16.6$ Hz, **H1**), 2.46-2.40 (m, 2H, **OCH₂CH₂CH₂O**).

(c) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11aS)-11-sulpho-7-methoxy-2-(3,4-methylenedioxyphenyl)-1,10,11,11a-tetrahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one]] sodium salt (**30**)




5 A solution of sodium bisulphite (2.2 mg, 0.021 mmol, 2.0 equiv.) in H₂O (0.5 mL) was added to a solution of compound **29** (8.1 mg, 0.0105 mmol) in DCM (0.5 mL). The reaction mixture was allowed to stir for 5 hours. TLC (EtOAc) revealed complete consumption of the starting material and a product spot at the base line. The reaction mixture was diluted with DCM (1.5 mL) and H₂O (1.5 mL). The aqueous layer was separated and lyophilized to afford **30** as a lightweight white solid (3.5 mg, 34 %): ¹H NMR (400 MHz, DMSO) δ 7.47 (s, 2H, **H3**), 6.98 (s x 2, 2H, **H6**), 7.06-6.78 (m, 6H, **H_{arom}**), 6.51 (s x 2, 2H, **H9**), 5.27 (s, 2H, **NH**), 4.32-4.26 (m, 2H, **H11a**), 4.17-4.07 (m, 4H, **OCH₂CH₂CH₂O**), 3.93 (d, 2H, *J* = 10.4 Hz, **H11**), 3.72 (s, 6H, **OCH₃** x 2), 3.50-3.20 (m, 4H, **H1**), 2.27-2.20 (m, 2H, **OCH₂CH₂CH₂O**).

15

Example 7

(a) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11S,11aS)-10-(2,2,2-trichloroethoxycarbonyl)-11-(tert-butylidimethylsilyloxy)-7-methoxy-2-(4-fluorobenzene)-1,10,11,11a-tetrahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one]] (**31**)




20 A solution of TEA (0.072 mL, 0.52 mmol, 6.0 eq) in H₂O (0.6 mL) and EtOH (3.5 mL) was added to a solution of triflate **13** (125 mg, 0.086 mmol) in toluene (3.5 mL) at room temperature. To this mixture 4-fluorobenzeneboronic acid (31.6 mg, 0.22 mmol, 2.6 eq) and Pd(PPh₃)₄ (4.0 mg, 3.4 μmol, 0.04 eq) were added. The reaction mixture was heated at 100 °C under microwave irradiation for 5 minutes when TLC (80:20 v/v EtOAc/hexane) revealed complete consumption of the starting material. Excess solvent was removed and the residue was dissolved in EtOAc (10 mL), washed with H₂O (10 mL), brine (10 mL) and dried (MgSO₄). Filtration and evaporation of solvent afforded the crude product which was

25


purified by flash column chromatography (80:20 v/v hexane/EtOAc then gradient to 65:35 v/v hexane/EtOAc) to afford **31** as a yellow glass (82 mg, 71%): $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.40 (s, 2H, **H3**), 7.42 (dd, 4H, $J = 5.2, 8.7$ Hz, H_{arom}), 7.28 (s, 2H, **H6**), 7.06 (dd, 4H, $J = 8.7$ Hz, H_{arom}), 6.70 (s, 2H, **H9**), 5.92 (d, 2H, $J = 8.8$ Hz, **H11**), 5.22 (d, 2H, $J = 12.0$ Hz, **Troc CH₂**), 4.32-4.29 (m, 2H, **OCH₂CH₂CH₂O**), 4.19-4.10 (m, 4H, **Troc CH₂** and **OCH₂CH₂CH₂O**), 4.00-3.92 (m, 8H, **H11a** and **OCH₃ x 2**), 3.33 (dd, 2H, $J = 10.8, 16.5$ Hz, **H1**), 2.84 (d, 2H, $J = 16.5$ Hz, **H1**), 2.49-2.44 (m, 2H, **OCH₂CH₂CH₂O**), 0.92 (s, 18H, **TBS CH₃ x 6**), 0.30 and 0.27 (s x 2, 12H, **TBS CH₃ x 4**).

10 (b) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11a*S*)-7-methoxy-2-(4-fluorobenzene)-1,11a-dihydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepine-5-one]]] (**32**)



10% Cd/Pb couple (342 mg, 2.80 mmol, 25 eq) was added to a rapidly stirring mixture of **31** (149 mg, 0.112 mmol), THF (4.5 mL) and 1 N NH_4OAc (4.5 mL). The reaction mixture was allowed to stir for 5 hours when TLC showed complete consumption of the starting material. The solids were filtered and rinsed with H_2O and MeOH. Excess solvent was removed and the residue was diluted with H_2O (30 mL) and DCM (30 mL). The aqueous layer was extracted with DCM (3 x 30 mL) and the organic extracts were combined, washed with H_2O (80 mL) and brine (80 mL) and dried (MgSO_4). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (99.9:0.1 v/v $\text{CHCl}_3/\text{MeOH}$ then gradient to 98:2 v/v $\text{CHCl}_3/\text{MeOH}$) to afford **32** as a yellow glass (60 mg, 75%): $^1\text{H NMR}$ (400 MHz, CDCl_3) 7.88 (d, 2H, $J = 3.9$ Hz, **H11**), δ 7.51 (s, 2H, **H6**), 7.43 (s, 2H, **H3**), 7.34 (dd, 4H, $J = 5.3, 8.6$ Hz, H_{arom}), 7.06 (dd, 4H, $J = 8.7$ Hz, H_{arom}), 6.87 (s, 2H, **H9**), 4.41-4.25 (m, 6H, **H11a** and **OCH₂CH₂CH₂O**), 3.93 (s, 3H, **OCH₃ x 2**), 3.55 (dd, 2H, $J = 10.8, 16.5$ Hz, **H1**), 3.37 (d, 2H, $J = 16.5$ Hz, **H1**), 2.46-2.42 (m, 2H, **OCH₂CH₂CH₂O**).


(c) 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11aS)-11-sulpho-7-methoxy-2-(4-fluorobenzene)-1,10,11,11a-tetrahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one]] sodium salt (**33**)



A solution of sodium bisulphite (13.9 mg, 0.134 mmol, 2.0 equiv.) in H₂O (3 mL) was added to a solution of compound **32** (48.0 mg, 0.067 mmol) in DCM (3 mL). The reaction mixture was allowed to stir for 2 hours. TLC (EtOAc) revealed complete consumption of the starting material and a product spot at the base line. The reaction mixture was diluted with DCM (8 mL) and H₂O (8 mL). The aqueous layer was separated and lyophilized to afford **33** as a lightweight white solid (35.5 mg, 58 %): ¹H NMR (400 MHz, DMSO) δ 7.56 (s, 2H, **H3**), 7.49 (dd, 4H, *J* = 5.3, 8.6 Hz, **H_{arom}**), 7.17 (dd, 4H, *J* = 8.7 Hz, **H_{arom}**), 7.07 (s, 2H, **H6**), 6.53 (s, 2H, **H9**), 5.28 (s, 2H, **NH**), 4.36-4.30 (m, 2H, **H11a**), 4.17-4.09 (m, 4H, **OCH₂CH₂CH₂O**), 3.95 (d, 2H, *J* = 10.4 Hz, **H11**), 3.73 (s, 6H, **OCH₃ x 2**), 3.53-3.22 (m, 4H, **H1**), 2.27-2.19 (m, 2H, **OCH₂CH₂CH₂O**).

15 Example 8


(a) 1,1'-[[[(Pentane-1,5-diyl)dioxy]bis[(2S,4R)-(5-methoxy-2-nitro-1,4-phenylene)carbonyl]]bis[2-(tert-butyldimethylsilyloxymethyl)-4-hydroxyacetylpyrrolidine] (**36**)



A catalytic amount of DMF (2 drops) was added to a stirred solution of the nitro-acid **34** (15.0 g, 30.4 mmol) and oxalyl chloride (13.5 mL, 19.3 g, 152 mmol) in dry THF (300 mL). The reaction mixture was allowed to stir for 16 hours at room temperature and the solvent was removed by evaporation *in vacuo*. The resulting residue was re-dissolved in dry THF

(225 mL) and the acid chloride solution was added dropwise to a stirred mixture of the amine **35** (20.7 g, 76.0 mmol) and TEA (84 mL, 61.52 g, 608 mmol) in THF (150 mL) at 0°C (ice/acetone) under a nitrogen atmosphere. The reaction mixture was allowed to warm to room temperature and stirred for a further 16 hours. Excess THF was removed by rotary evaporation and the resulting residue was partitioned between H₂O (400 mL) and EtOAc (400 mL). The layers were allowed to separate and the aqueous layer was extracted with EtOAc (3 x 130 mL). The combined organic layers were then washed with saturated NH₄Cl (200 mL), saturated NaHCO₃ (200 mL), brine (200 mL) and dried (MgSO₄). Filtration and evaporation of the solvent gave the crude product as a dark coloured oil. Purification by flash chromatography (gradient elution: 60:40 v/v Hexane/EtOAc to 40:60 v/v Hexane/EtOAc) isolated the pure amide **36** as a light yellow coloured glass (23.56 g, 77%): ¹H NMR (400 MHz, CDCl₃) (rotamers) δ 7.66 (s, 2H), 6.71 (s, 2H), 5.21-5.17 (m, 2H), 4.57-4.53 (m, 2H), 4.21-4.08 (m, 6H), 3.91 (s, 6H), 3.76-3.72 (m, 2H), 3.46 (dd, 2H, *J* = 11.8, 4.6 Hz), 3.21 (d, 2H, *J* = 11.9 Hz), 2.44-2.37 (m, 2H), 2.24-2.18 (m, 2H), 2.03-1.94 (m, 10H), 1.75-1.67 (m, 2H), 0.91-0.84 (m, 18H), 0.11- -0.05 (m, 12H); ¹³C NMR (100 MHz, CDCl₃) δ 171.0, 166.4, 154.4, 148.5, 137.5, 127.7, 109.2, 108.3, 72.9, 69.3, 62.6, 57.4, 56.5, 54.8, 33.0, 28.5, 25.8, 22.5, 21.0, 18.2, -5.4 and -5.5; LC/MS 2.93 min (ES+) *m/z* (relative intensity) 1006 ([*M* + H]⁺, 100).


(b) 1,1'-[[[(Pentane-1,5-diyloxy)bis[(2*S*,4*R*)-(5-methoxy-2-amino-1,4-phenylene)carbonyl]]bis[2-(*tert*-butyldimethylsilyloxymethyl)-4-hydroxyacetylpyrrolidine]] (37)



A slurry of 10% Pd-C (550 mg) in EtOAc (20 mL) was added to a solution of the nitro compound **36** (5.5 g, 5.47 mmol) in EtOAc (73 mL). The mixture was subjected to hydrogenation using Parr apparatus at 30 psi for a total of 24 hours. The mixture was degassed and analysed by TLC (EtOAc) where complete consumption of starting material was observed. The catalyst was removed by vacuum filtration through Whatman GF/F filter paper and the filtrate evaporated to provide the aniline **37** as a grey coloured foam (5.17 g, 100%): ¹H NMR (400 MHz, CDCl₃) δ 6.71 (s, 2H), 6.22 (s, 2H), 5.25-5.23 (m, 2H), 4.58-4.30 (m, 6H), 4.14-4.05 (m, 2H), 3.99 (t, *J* = 6.6 Hz, 4H), 3.79-3.73 (m, 8H), 3.69-3.54 (m, 4H), 2.40-2.33 (m, 2H), 2.16-2.08 (m, 2H), 2.00 (s, 6H), 1.94-1.87 (m, 4H), 1.68-1.60


(m, 2H), 0.89 (s, 18H), 0.05 and 0.04 (s x 2, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ 170.6, 170.0, 151.5, 141.7, 141.2, 113.1, 111.0, 102.0, 73.5, 68.5, 62.6, 57.0, 56.2, 32.9, 28.7, 25.6, 22.6, 21.1, 18.1, -5.1 and -5.4; LC/MS 2.78 min (ES+) m/z (relative intensity) 946 ($[M + H]^+$, 52), 672 (30), 399 (20), 274 (65), 166 (20).

5
(c) 1,1'-[[[(Pentane-1,5-diyl)dioxy]bis[(2S,4R)-[5-methoxy-1,4-phenylene-2-(2,2,2-trichloroethoxycarbonylamino)]carbonyl]]bis[2-(tert-butylidimethylsilyloxymethyl)-4-hydroxyacetylpyrrolidine] (38)



10 A solution of 2,2,2-trichloroethyl chloroformate (2.63 mL, 4.05 g, 19.1 mmol) in dry DCM (45 mL) was added dropwise to a stirred solution of amine **37** (8.21 g, 8.70 mmol) and anhydrous pyridine (2.81 mL, 2.75 g, 34.8 mmol) in dry DCM (120 mL) at $-10\text{ }^\circ\text{C}$ (liquid N_2 /ethanediol). After 16 hours stirring at room temperature TLC (50:50 v/v Hexane/EtOAc) revealed complete consumption of starting material. The reaction mixture was washed
15 with saturated NH_4Cl (2 x 70 mL), saturated CuSO_4 (70 mL), H_2O (70 mL), brine (70 mL) and dried (MgSO_4). Filtration and evaporation of the solvent yielded the Troc-carbamate **38** as a yellow foam (11.25 g, 99%): ^1H NMR (400 MHz, CDCl_3) δ 9.48 (br s, 1H), 7.85 (s, 2H), 6.79 (s, 2H), 5.28-5.24 (m, 2H), 4.86-4.76 (m, 4H), 4.70-4.53 (m, 2H), 4.12 (t, $J = 6.4$ Hz, 4H), 4.14-4.10 (m, 2H), 3.80 (s, 6H), 3.76 (dd, 2H, $J = 12.2, 3.8$ Hz), 3.69-3.62 (m, 4H),
20 2.44-2.37 (m, 2H), 2.16 (dd, 2H, $J = 14.1, 8.1$ Hz), 2.10-1.92 (m, 10H), 1.72-1.65 (m, 2H), 0.90 (s, 18H), 0.06 and 0.04 (s x 2, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ 170.5, 169.3, 152.0, 151.1, 144.2, 132.4, 114.4, 111.8, 105.2, 95.3, 74.4, 73.4, 68.7, 62.3, 57.3, 57.2, 56.5, 32.6, 28.7, 25.8, 22.6, 21.1, 18.1, -5.4 and -5.5; LC/MS 3.23 min (ES+/ES-) m/z (M^+ not observed), 472 (30), 416 (15), 302 (85), 274 (60), 198 (20), 170 (100), 142 (50), 110
25 (80).

(d) 1,1'-[[[(Pentane-1,5-diyl)dioxy]bis[(2S,4R)-[5-methoxy-1,4-phenylene-2-(2,2,2-trichloroethoxycarbonylamino)]carbonyl]]bis(2-hydroxymethyl-4-hydroxyacetylpyrrolidine)] (39)




39

5 Glacial acetic acid (90 mL) was added to a stirred solution of **38** (5.06 g, 3.91 mmol) in H₂O (30 mL) and THF (30 mL). The reaction mixture was allowed to stir at room temperature for 16 h at which point TLC (95:5 v/v CHCl₃/MeOH) revealed completion of the reaction. The acidic solution was added dropwise to a stirred solution of NaHCO₃ (132 g) in H₂O (1.3 L) at 0°C (ice/acetone). The aqueous layer was extracted with EtOAc (3 x 200 mL) and the
10 organic layers were combined, washed with H₂O (200 mL), brine (200 mL) and dried (MgSO₄). Filtration and evaporation of the solvent *in vacuo* afforded the crude product which was purified by flash column chromatography (98:2 v/v CHCl₃/MeOH) to provide the bis-alcohol **39** as a white glass (4.04 g, 97%): ¹H NMR (400 MHz, CDCl₃) δ 9.01 (br s, 2H), 7.66 (s, 2H), 6.79 (s, 2H), 5.23-5.20 (m, 2H), 4.84 (d, 2H, *J* = 12.0 Hz), 4.77 (d, 2H, *J* =
15 12.0 Hz), 4.63-4.55 (m, 2H), 4.10 (t, *J* = 6.4 Hz, 4H), 4.08-4.01 (m, 2H), 3.82 (s, 6H), 3.75 (dd, 2H, *J* = 12.6, 3.7 Hz), 3.70-3.58 (m, 4H), 2.26 (dd, 2H, *J* = 14.1, 7.6 Hz), 2.15-2.06 (m, 2H), 2.01 (s, 6H), 1.98-1.92 (m, 4H), 1.74-1.65 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 170.6, 170.4, 152.1, 150.9, 144.8, 131.0, 115.6, 111.3, 105.7, 95.3, 74.3, 72.5, 68.7, 64.3, 58.8, 56.6, 56.5, 33.6, 28.5, 22.6, 21.1.

20

(e) 1,1'-[[[(Pentane-1,5-diyl)dioxy]bis[(11S,11aS,2R)-10-(2,2,2-trichloroethoxycarbonyl)-11-hydroxy-7-methoxy-2-hydroxyacetyl-1,2,3,10,11,11a-hexahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one]]] (40)




40

25 TEMPO (234 mg, 1.50 mmol) and BAIB (5.31 g, 16.5 mmol) were added to a stirred solution of bis-alcohol **39** (4.0 g, 3.75 mmol) in DCM (140 mL). The reaction mixture was allowed to stir under a nitrogen atmosphere for 24 hours at which point TLC (95:5 v/v


CHCl₃/MeOH) revealed conversion of starting material to product. The mixture was diluted with DCM (30 mL), washed with saturated aqueous NaHSO₃ (2 x 30 mL), saturated aqueous NaHCO₃ (2 x 30 mL), brine (30 mL) and dried (MgSO₄). Filtration and evaporation of the solvent afforded the crude product which was purified by flash column chromatography (gradient elution: CHCl₃ to 99.5:0.5 v/v CHCl₃/MeOH) to provide the cyclised product **40** as a light coloured yellow glass (1.99 g, 50%): ¹H NMR (400 MHz, CDCl₃) δ 7.27 (s, 2H), 6.78 (s, 2H), 5.68 (d, 2H, *J* = 9.7 Hz), 5.41-5.35 (m, 2H), 5.24 (d, 2H, *J* = 12.0 Hz), 4.23 (d, 2H, *J* = 12.0 Hz), 4.14-3.97 (m, 6H), 3.93 (s, 6H), 3.77-3.69 (m, 4H), 2.46-2.33 (m, 4H), 2.05 (s, 6H), 2.03-1.87 (m, 4H), 1.70-1.62 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 170.4, 167.5, 154.4, 150.6, 149.0, 127.3, 124.5, 113.7, 110.8, 95.0, 87.5, 75.0, 71.4, 68.8, 58.3, 56.2, 51.1, 35.8, 28.5, 22.4, 21.0.

(f) 1,1'-[[[(Pentane-1,5-diyloxy)bis[(11*S*,11*aS*,2*R*)-10-(2,2,2-trichloroethoxycarbonyl)-11-(*tert*-butyldimethylsilyloxy)-7-methoxy-2-hydroxyacetyl-1,2,3,10,11,11*a*-hexahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]]] (**41**)



TBSOTf (1.27 mL, 1.47 g, 5.54 mmol) was added to a stirred solution of bis-alcohol **40** (1.96 g, 1.85 mmol) and 2,6-lutidine (0.86 mL, 0.79 g, 7.39 mmol) in dry DCM (40 mL). The reaction mixture was allowed to stir under a N₂ atmosphere for 5 hours after which time TLC (EtOAc) revealed consumption of starting material. Following dilution with DCM (50 mL), the organic mixture was washed with saturated CuSO₄ (30 mL), saturated NaHCO₃ (30 mL), brine (50 mL) and dried (MgSO₄). Filtration and evaporation of the solvent *in vacuo* afforded the crude product which was purified by flash column chromatography (40:60 v/v Hexane/EtOAc) to provide the product **41** as a white glass (1.31 g, 55%): ¹H NMR (400 MHz, CDCl₃) δ 7.27 (s, 2H), 6.72 (s, 2H), 5.76 (d, 2H, *J* = 9.0 Hz), 5.37 (br s, 2H), 5.24 (d, 2H, *J* = 12.1 Hz), 4.17 (d, 2H, *J* = 12.1 Hz), 4.14-3.96 (m, 6H), 3.93 (s, 6H), 3.75-3.66 (m, 4H), 2.40-2.33 (m, 2H), 2.28-2.18 (m, 2H), 2.04 (s, 6H), 1.98-1.90 (m, 4H), 1.69-1.62 (m, 2H), 0.86 (s, 18H), 0.23 and 0.22 (s x 2, 12H); ¹³C NMR (100 MHz, CDCl₃) δ 170.4, 168.0, 153.6, 150.5, 149.1, 127.9, 125.3, 113.9, 110.7, 95.2, 88.2, 74.7, 71.7, 68.7, 60.5, 56.2, 51.2, 36.3, 28.7, 25.6, 22.7, 21.1, 17.8, -4.2 and -5.2.

(g) 1,1'-[[*(Pentane-1,5-diyl)dioxy*]]bis[*(11S,11aS,2R)*-10-(2,2,2-trichloroethoxycarbonyl)-11-(*tert*-butyldimethylsilyloxy)-7-methoxy-2-hydroxy-1,2,3,10,11,11a-hexahydro-5H-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]] (**42**)




5 A solution of K_2CO_3 (685 mg, 4.96 mmol) in H_2O (15 mL) was added to a stirred solution of acetate compound **41** (1.28 g, 0.99 mmol) in MeOH (15 mL). The initial colourless solution eventually turned yellow and the formation of a white precipitate was observed. The reaction mixture was heated at reflux for 5 hours at which point TLC (EtOAc) revealed complete conversion of starting material to product. Excess solvent was removed by rotary

10 evaporation and the mixture was carefully neutralized with 1N HCl to pH 7. The resulting mixture was extracted with DCM (5 x 20 mL) and the combined organic layers were then washed with brine (50 mL) and dried ($MgSO_4$). Filtration followed by removal of the solvent by evaporation *in vacuo* afforded the product **42** as a white solid (982 mg, 82%): 1H NMR (400 MHz, $CDCl_3$) δ 7.19 (s, 2H), 6.72 (s, 2H), 5.75 (d, 2H, $J = 9.0$ Hz), 5.22 (d, 2H, $J = 12.0$ Hz), 4.58-4.56 (m, 2H), 4.18 (d, 2H, $J = 12.0$ Hz), 4.11-3.95 (m, 6H), 3.85 (s, 6H), 3.75-3.69 (m, 2H), 3.59 (dd, 2H, $J = 12.7, 4.2$ Hz), 2.55 (br s, 2H), 2.38-2.25 (m, 2H), 2.14-2.03 (m, 2H), 1.97-1.85 (m, 4H), 1.74-1.62 (m, 2H), 0.86 (s, 18H), 0.22 and 0.21 (s x 2, 12H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 168.5, 153.6, 150.6, 149.1, 127.9, 125.3, 114.0, 110.8, 95.3, 88.3, 74.7, 69.5, 68.8, 60.8, 56.0, 54.0, 38.8, 28.8, 25.6, 22.7, 17.8, -4.2 and -5.2.

20

(h) 1,1'-[[*(Pentane-1,5-diyl)dioxy*]]bis[*(11S,11aS)*-10-(2,2,2-trichloroethoxycarbonyl)-11-(*tert*-butyldimethylsilyloxy)-7-methoxy-2-oxo-1,2,3,10,11,11a-hexahydro-5H-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]] (**43**)



25


A solution of anhydrous DMSO (0.34 mL, 0.37 g, 4.78 mmol) in dry DCM (4 mL) was added dropwise over a period of 5 minutes to a stirred solution of oxalyl chloride (1.20 mL of a 2M solution in DCM, 2.39 mmol) under a nitrogen atmosphere at -60 °C (liq $N_2/CHCl_3$).

After stirring at -55°C for 15 minutes, a slurry of the substrate **42** (962 mg, 0.80 mmol) in dry DCM (8 mL) was added dropwise over a period of 10 minutes to the reaction mixture. After stirring for a further 1 h at -55 °C, a solution of TEA (1.56 mL, 1.13 g; 11.2 mmol) in dry DCM (4 mL) was added dropwise over a period of 5 minutes to the reaction mixture.

5 The reaction mixture was allowed to warm to 0°C and then diluted with DCM (50 mL). The organic solution was washed with cold 1 N HCl (20 mL), H₂O (20 mL), brine (30 mL) and dried (MgSO₄). Filtration and evaporation of the solvent *in vacuo* afforded the crude product which was purified by flash column chromatography (50:50 v/v Hexane/EtOAc) to afford bis-ketone **43** as a foam (550 mg, 57%): ¹H NMR (400 MHz, CDCl₃) δ 7.25 (s, 2H), 6.76 (s, 2H), 5.82 (d, 2H, *J* = 9.3 Hz), 5.22 (d, 2H, *J* = 12.0 Hz), 4.32 (d, 2H, *J* = 20.4 Hz), 4.22 (d, 2H, *J* = 12.0 Hz), 4.17-4.08 (m, 4H), 4.03-3.89 (m, 10H), 2.96 (dd, 2H, *J* = 19.6, 10.2 Hz), 2.56 (dd, 2H, *J* = 19.6, 2.8 Hz), 1.99-1.92 (m, 4H), 1.72-1.64 (m, 2H), 0.86 (s, 18H), 0.24 and 0.23 (s x 2, 12H); ¹³C NMR (100 MHz, CDCl₃) δ 207.7, 168.0, 153.8, 151.0, 149.5, 127.9, 124.4, 114.1, 110.8, 95.2, 87.5, 74.8, 68.8, 58.9, 56.2, 52.9, 40.4, 28.7, 25.6, 22.8, 17.8, -4.12 and -5.31.

10
15


(i) 1,1'-[[[(Pentane-1,5-diyl)dioxy]bis[(11*S*,11*aS*)-10-(2,2,2-trichloroethoxycarbonyl)-11-(*tert*-butyldimethylsilyloxy)-7-methoxy-2-[[[(trifluoromethyl)sulfonyl]oxy]-1,10,11,11*a*-tetrahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]]] (**44**)



Anhydrous triflic anhydride taken from a freshly opened ampoule (1.61 mL, 2.71 g, 9.60 mmol) was added rapidly in one portion to a vigorously stirred solution of ketone **43** (525 mg, 0.44 mmol) and anhydrous pyridine (0.78 mL, 759 mg, 9.60 mmol) in dry DCM (25 mL) at 0-5°C (ice) under a nitrogen atmosphere. The reaction mixture was immediately allowed to warm to room temperature and eventually turned a dark red colour. The reaction mixture was allowed to stir for a total of 28 hours at which point TLC (80:20 v/v EtOAc/hexane) revealed the complete consumption of starting material. The mixture was poured into cold saturated NaHCO₃ (50 mL) and extracted with DCM (3 x 20 mL). The combined organic layers were then washed with saturated CuSO₄ (30 mL), brine (30 mL) and dried (MgSO₄). Filtration and evaporation of the solvent *in vacuo* afforded the crude product which was purified by flash column chromatography (80:20 v/v Hexane/EtOAc) to


afford triflate **44** as a yellow foam (249 mg, 39%): $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.23 (s, 2H), 7.17 (s, 2H), 6.73 (s, 2H), 5.93 (d, 2H, $J = 8.9$ Hz), 5.22 (d, 2H, $J = 12.1$ Hz), 4.21 (d, 2H, $J = 12.0$ Hz), 4.15-4.07 (m, 2H), 4.02-3.86 (m, 8H), 3.33 (ddd, 2H, $J = 16.6, 10.7, 2.3$ Hz), 2.82 (dd, 2H, $J = 16.7, 2.6$ Hz), 1.98-1.91 (m, 4H), 1.71-1.63 (m, 2H), 0.88 (s, 18H), 0.28 and 0.25 (s x 2, 12H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 164.9, 153.6, 151.3, 149.6, 136.0, 127.8, 123.7, 121.0, 118.6 (q, $J = 321.5$ Hz), 114.2, 111.0, 95.1, 86.4, 74.9, 68.8, 60.6, 56.2, 34.4, 28.7, 25.6, 22.8, 17.8, -4.2 and -5.4.

(j) 1,1'-[[[(Pentane-1,5-diyl)dioxy]bis[(11S,11aS)-10-(2,2,2-trichloroethoxycarbonyl)-11-(tert-butyl)dimethylsilyloxy)-7-methoxy-2-(p-methoxyphenyl)-1,10,11,11a-tetrahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one]]] (**45**)




$\text{Pd}(\text{PPh}_3)_4$ (7.43 mg, 6.43 μmol) was added to a solution of the triflate **44** (236 mg, 0.16 mmol), 4-methoxyphenylboronic acid (63 mg, 0.42 mmol), TEA (0.13 mL, 98 mg, 0.97 mmol) in H_2O (0.8 mL), EtOH (5 mL) and toluene (5 mL) at room temperature. The reaction mixture was heated in the microwave at 100 $^\circ\text{C}$ for 30 minutes after which time TLC (80:20 v/v EtOAc/hexane) revealed the complete consumption of the starting material. Excess solvent was removed by evaporation *in vacuo* and the resulting residue was dissolved in EtOAc (50 mL), washed with H_2O (20 mL), brine (20 mL) and dried (MgSO_4). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (gradient elution: 80:20 v/v Hexane/EtOAc to 50:50 v/v Hexane/EtOAc) to afford C2-aryl product **45** as a yellow glass (117 mg, 52%): $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.35 (s, 2H), 7.31 (s, 2H), 7.31-7.27 (m, 6H), 6.90 (d, 4H, $J = 8.8$ Hz), 6.76 (s, 2H), 5.92 (d, 2H, $J = 8.8$ Hz), 5.23 (d, 2H, $J = 12.0$ Hz), 4.20 (d, 2H, $J = 12.1$ Hz), 4.14-4.05 (m, 4H), 4.04-3.95 (m, 2H), 3.93 (s, 6H), 3.83 (s, 6H), 3.34 (ddd, 2H, $J = 16.3, 10.4, 2.0$ Hz), 2.86-2.75 (m, 2H), 2.02-1.94 (m, 4H), 1.72-1.66 (m, 2H), 0.93 and 0.86 (s x 2, 18H), 0.28 and 0.25 (s x 2, 12H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 163.7, 159.1, 153.7, 150.8, 149.3, 127.8, 126.3, 126.1, 125.4, 122.3, 122.2, 114.3, 114.2, 110.9, 95.3, 87.3, 74.8, 68.8, 61.5, 56.2, 55.4, 35.3, 28.8, 25.7, 22.8, 17.9, -4.0 and -5.1.

(k) 1,1'-[[*(Pentane-1,5-diyloxy)bis*[(11*aS*)-7-methoxy-2-(*p*-methoxyphenyl)-1,11*a*-dihydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepine-5-one]] (46)



10% Cd/Pb couple (192 mg, 1.55 mmol) was added to a rapidly stirring mixture of **45** (107 mg, 77.3 μmol), THF (2.5 mL) and 1N NH_4OAc (2.5 mL). The reaction mixture was allowed to stir for 3 hours at which point TLC (95:5 v/v $\text{CHCl}_3/\text{MeOH}$) showed the formation of desired PBD accompanied by side-products. The solids were collected by filtration and rinsed with H_2O and DCM. The aqueous layer was extracted with DCM (3 x 10 mL) and the organic extracts were combined, washed with brine (50 mL) and dried (MgSO_4). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (gradient elution: CHCl_3 to 99:1 v/v $\text{CHCl}_3/\text{MeOH}$) to afford the imine **46** as a yellow glass (32.5 mg, 55%): $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.89 (d, 2H, $J = 3.9$ Hz), 7.53 (s, 2H), 7.39 (s, 2H), 7.34 (d, 4H, $J = 8.8$ Hz), 6.90 (d, 4H, $J = 8.8$ Hz), 6.82 (s, 2H), 4.43-4.38 (m, 2H), 4.18-4.05 (m, 4H), 3.95 (s, 6H), 3.83 (s, 6H), 3.58 (ddd, 2H, $J = 16.2, 11.5, 1.9$ Hz), 3.38 (ddd, 2H, $J = 16.3, 5.1, 1.6$ Hz), 2.01-1.94 (m, 4H), 1.73-1.66 (m, 2H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 162.6, 161.3, 159.2, 151.3, 148.1, 140.3, 126.2, 126.0 (x2), 123.2, 122.0, 119.1, 114.3, 111.9, 110.9, 68.8, 56.2, 55.4, 53.9, 35.6, 28.7, 22.6.

(j) 1,1'-[[*(Pentane-1,5-diyloxy)bis*[(11*aS*)-11-sulpho-7-methoxy-2-(*p*-methoxyphenyl)-1,10,11,11*a*-tetrahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]] sodium salt (**47**)



A solution of sodium bisulphite (4.90 mg, 47.2 μmol) in water (3 mL) was added to a stirred solution of **46** (18.11 mg, 23.6 μmol) in dichloromethane (1.5 mL). The reaction mixture was allowed to stir vigorously for 24 hours, after which time the organic and aqueous layers were separated. TLC analysis (95:5 v/v $\text{CHCl}_3/\text{MeOH}$) of the aqueous phase revealed absence of **46** ($R_f \sim 0.3$) and the presence of baseline material with strong UV-absorption. The aqueous layer was lyophilised to provide the bisulphite adduct **47** as a lightweight solid (10 mg, 43%): $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ 7.41 (s, 2H), 7.37 (d, 4H, $J = 8.7$ Hz), 7.04

(s, 2H), 6.91 (d, 4H, $J = 8.9$ Hz), 6.47 (s, 2H), 5.24 (s, 2H), 4.31-4.21 (m, 2H), 4.06-3.86 (m, 6H), 3.76 (s, 6H), 3.71 (s, 6H), 3.51-3.21 (m, 4H), 1.90-1.70 (m, 4H), 1.62-1.51 (m, 2H).

Example 9: Determination of *In Vitro* Cytotoxicity

5 *K562 cell line (MTT assay)*

1 hour exposure

K562 human chronic myeloid leukaemia cells were maintained in RPMI 1640 medium supplemented with 10% fetal calf serum and 2 mM glutamine at 37°C in a humidified atmosphere containing 5% CO₂ and were incubated with a specified dose of drug for 1
10 hour at 37°C in the dark. The incubation was terminated by centrifugation (5 min, 300 g) and the cells were washed once with drug-free medium. Following the appropriate drug treatment, the cells were transferred to 96-well microtiter plates (10⁴ cells per well, 8 wells per sample). Plates were then kept in the dark at 37°C in a humidified atmosphere containing 5% CO₂. The assay is based on the ability of viable cells to reduce a yellow
15 soluble tetrazolium salt, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide (MTT, Aldrich-Sigma), to an insoluble purple formazan precipitate. Following incubation of the plates for 4 days (to allow control cells to increase in number by approximately 10 fold), 20 µL of MTT solution (5 mg/mL in phosphate-buffered saline) was added to each well and the plates further incubated for 5 h. The plates were then centrifuged for 5 min at 300 g and
20 the bulk of the medium pipetted from the cell pellet leaving 10–20 µL per well. DMSO (200 µL) was added to each well and the samples agitated to ensure complete mixing. The optical density was then read at a wavelength of 550 nm on a Titertek Multiscan ELISA plate reader, and a dose-response curve was constructed. For each curve, an IC₅₀ value was read as the dose required to reduce the final optical density to 50% of the control
25 value.

96 hour exposure

In a modification of the above method, 5x10⁴ K562 cells (as above) were maintained in RPMI 1640 medium (Sigma) supplemented with 10% foetal calf serum (Sigma) and 20mM
30 L-glutamine (Sigma) and 190 µL of this solution incubated with specific doses of the drug (10µL) for 96 hours at 37°C in 5% CO₂. 250µg/ml (final) MTT was added to each well and the plates incubated for a further 4 hours. Absorbance was read at 450nm using an Envision plate reader (Applied Biosystems). Data was then analysed by Graphpad PRISM, and an IC₅₀ obtained (defined as the compound concentration needed to reduce the cell
35 population by half).

Results

Compound	IC ₅₀ (μM)
ZC-207	0.0053 ^a
ZC-423	0.0015 ^b

^a 1 hour incubation

^b 96 hour incubation

5

K562 cell line (Alamar Blue assay)

The compounds to be tested were dissolved in 100% Biotech grade DMSO (Sigma) and subsequently diluted to stock concentrations of either 2 μM or 200 nM in 2% DMSO. 100 μl of the stock concentrations were serially diluted 1 in 3 in 2% DMSO in intermediate polystyrene 96 well cell culture plates (Nunc). 2% DMSO was placed in the outer columns for use as blanks, and in the 2nd or 11th column (depending on whether the top or bottom of an assay plate was to be used) for use as a control. The entire row of the intermediate plate was then transferred in triplicate to either rows B to D or E to G of a fluorescence compatible polystyrene 96 well cell culture plate (Greiner BioOne); 10 μl to each well. A cell solution containing 5 x 10⁴ cells/ml was made up in phenol red free RPMI 1640 (Sigma) supplemented with 10% foetal calf serum (Sigma) and 20mM L-glutamine (Sigma). 190 μl cell solution was added to each well of the assay plate, from columns 2 to 11. 190 μl culture media was added to columns 1 and 12. Plates were incubated for 96 hours at 37°C in 5% CO₂. 1 μM (final) rezasurin (converted to fluorescent resofurin by viable cells) was added to each well and the plates incubated for a further 4 hours. Fluorescence was read at 530-570nm excitation, 580-620nm emission using an Envision plate reader (Applied Biosystems). Data was then analysed by Graphpad PRISM, and an IC₅₀ obtained (defined as the compound concentration needed to reduce the cell population by half).

10

15

20

Results

Compound	IC ₅₀ (nM)
19	<1
22	<1
26	<1
27	<1
29	<1
30	<1
32	<1
33	<1

LOXIMVI and OVCAR-5 cell lines

5 A cell solution containing 5×10^4 cells/ml of either LOXIMVI human melanoma cells or OVCAR-5 human ovarian tumour cells was made up in phenol red free RPMI 1640 (Sigma) supplemented with 10% foetal calf serum (Sigma) and 20mM L-glutamine (Sigma). 190 μ l cell solution was added to each well of a fluorescence compatible polystyrene 96 well cell culture plate (Greiner BioOne), from columns 2 to 11. 190 μ l culture media was added to
10 columns 1 and 12. Plates were incubated overnight at 37°C in 5% CO₂. Compounds were dissolved in 100% Biotech grade DMSO (Sigma) and subsequently diluted to stock concentrations in 2% DMSO. 100 μ l of the stock concentrations were serially diluted 1 in 3 in 2% DMSO in intermediate polystyrene 96 well cell culture plates (Nunc). 2% DMSO was placed in the outer columns for use as blanks, and in the 2nd or 11th column (depending on
15 whether the top or bottom of an assay plate was to be used) for use as a control. The entire row of the intermediate plate was then transferred in triplicate, 10 μ l per well, to either rows B to D or E to G of the cell plate. Plates were incubated for 96 hours at 37°C in 5% CO₂. 1 μ M (final) rezasurin (converted to fluorescent resofurin by viable cells) was added to each well and the plates incubated for a further 4 hours. Fluorescence was read at 530-
20 570nm excitation, 580-620nm emission using an Envision plate reader (Applied Biosystems). Data was then analysed by Graphpad PRISM, and an IC₅₀ obtained (defined as the compound concentration needed to reduce the cell population by half).

Results

Cell line	ZC-423 IC ₅₀ (nM)
LOXIMVI	4.83
OVCAR-5	5.2

Example 10: Anti tumour activity

5 The following experiments was carried out under a project licence approved by the Home Office, London, UK, and UK CCCR guidelines (Workman, P., *et al.*, *British Journal of Cancer*, 77, 1-10 (1998)) were followed throughout.

10 LOX IMVI (human melanomas) were grown subcutaneously in nude mice (B&K Universal, Hull, UK). Tumours were transplanted as single tumour fragments in the flank by trocar. Groups of 8 tumour-bearing mice were treated with ZC423 at the previously established single intravenous (iv) maximum tolerated dose (MTD) of 3mgkg⁻¹ using 5% DMA/saline as the vehicle. Control mice (n=8) were treated with vehicle alone.

15 Treatment commenced when tumours could be reliably measured by calipers (mean dimensions 4 x 4mm) and therapeutic effects were assessed by daily caliper measurements of the tumour and mouse weights. Tumour volumes were determined by the formula $a^2 \times b/2$ where a is the smaller and b is the larger diameter of the tumour. The results are shown in Fig. 1 as a graphs of relative tumour volume against time (■ ZC423; ♦ Control – solvent alone). The activity of ZC423 was shown to be statistically significant when assessed by Mann-Whitney analysis (as described, for example, in Essential Statistics 2nd Edition (1989), D.G.Rees, ISBN 0 412 32030 4, Chapman and Hall, London).

20

25 Fig. 2 shows the results of the same test (● ZC423; ♦ Control – solvent alone) but extended over a longer period, and with a larger scale for the Mean RTV. The control group was ceased at 16 days, at which point there were no tumour free mice, compared to the treated group where all animals were tumour free at 68 days.

30 A study was carried out using OVCAR-5 (human ovarian tumours) under the same conditions as described above. The results are shown in Fig. 3 (● ZC423; ♦ Control – solvent alone). In this study the control group was ceased at 32 days.

Example 11: Cross-linking and Solubility*DNA Cross-linking*

5 Closed-circular puc18 DNA was linearized with HindIII, then dephosphorylated, and finally
5' end labeled with [γ 32P]-ATP using polynucleotide kinase. Reactions containing 10 ng of
DNA and drug were carried out in aqueous 1 x TEOA (25 mM triethanolamine, 1 mM
EDTA, pH 7.2) buffer at a final volume of 50 μ L at 37°C. Reactions were terminated by
addition of an equal volume of stop solution (0.6 M NaOAc, 20 mM EDTA, 100 μ g/mL
tRNA) followed by precipitation with ethanol. Following centrifugation of the samples, the
10 supernatants were discarded and the pellets were dried by lyophilization. Samples were
resuspended in 10 μ L of alkaline denaturing buffer (4mg bromophenol blue, 600 mg
sucrose and 40 mg NaOH) and vortexed for three minutes at room temperature. The non-
denatured controls were re-suspended in 10 μ L of standard sucrose loading dye (2.5 mg
bromophenol blue, 2.5 mg xylene cyanol blue and 4 g sucrose). Both samples and
15 controls were loaded directly onto an agarose gel.

Electrophoresis was performed on a 0.8% submerged horizontal agarose gel, 20 cm in
length for 16 hours at 38-40 V in 1 x TAE running buffer (2.42 g Tris Base, 0.372 g EDTA,
0.571ml glacial acetic acid). Gels were dried under vacuum for 80 minutes at 80°C on a
20 Savant SG210D SpeedGel gel dryer onto one layer of Whatman 3MM with a layer of DE81
filter paper underneath. An autoradiograph was obtained, after overnight exposure onto
FujiRX x-ray film. The film bands were quantitated using a BioRad GS-670 imaging laser
densitometer. The percentage of cross-linking was calculated by measuring the total DNA
in each lane (the sum of the densities for the double-stranded and single-stranded bands)
25 relative to the amount of density of double-stranded band alone. A dose response curve
was derived by plotting drug concentration against the determined percentage level of
cross-linked DNA, from which was derived the amount required to cross-link 50% of the
DNA (XL₅₀).

30 *Solubility*

Solubility was determined by dissolving the test compound in the minimum amount of water
to produce a saturated solution.

Results*DNA Cross-linking*


Compound	IC ₅₀ (μM)
18	0.2
19	2
21	0.5
22	0.3

Solubility

Compound	Amount of compound (g) in 1L water
ZC-207	Insoluble
ZC-423	11
27	9
30	4
33	3

Claims

1. A compound of the formula I:



- 5 or solvate thereof, wherein:
 R^2 is an optionally substituted C_{5-20} aryl group;
 R^6 and R^9 are independently selected from H, R, OH, OR, SH, SR, NH_2 , NHR, NRR' , nitro, Me_3Sn and halo;
 where R and R' are independently selected from optionally substituted C_{1-12} alkyl, C_{3-20}
 10 heterocyclyl and C_{5-20} aryl groups;
 R^7 is selected from H, R, OH, OR, SH, SR, NH_2 , NHR, $NHRR'$, nitro, Me_3Sn and halo;
 R'' is a C_{3-12} alkylene group, which chain may be interrupted by one or more heteroatoms
 and/or aromatic rings;
 X is selected from O, S, or NH;
 15 z is 2 or 3;
 M is a monovalent pharmaceutically acceptable cation;
 R^2 , R^6 , R^7 , R^9 , X' and M' are selected from the same groups as R^2 , R^6 , R^7 , R^9 , X and M
 respectively, or M and M' may together represent a divalent pharmaceutically acceptable
 cation.
 20
2. A compound according to claim 1, wherein X is O.
3. A compound according to either claim 1 or claim 2, wherein R'' represents a linear
 saturated C_{3-12} alkylene group.
 25
4. A compound according to any one of claims 1 to 3, wherein R^9 is H.
5. A compound according to any one of claims 1 to 4, wherein R^6 is H.
- 30 6. A compound according to any one of claims 1 to 5, wherein R^7 is selected from H,
 OH and OR, where R is selected from optionally substituted C_{1-7} alkyl, C_{3-10} heterocyclyl
 and C_{5-10} aryl groups.

7. A compound according to claim 6, wherein R⁷ is OMe or OCH₂Ph.
8. A compound according to any one of claims 1 to 7, wherein R² is an optionally substituted C₅₋₇ aryl group.
9. A compound according to claim 8, wherein R² is an optionally substituted phenyl group.
10. A compound according to any one of claims 1 to 9, wherein both PBD monomers are identically substituted.
11. A compound according to any one of claims 1 to 10, wherein M and M' are monovalent pharmaceutically acceptable cations.
12. A compound according to claim 11, wherein M and M' are Na⁺.
13. A compound according to any one of claims 1 to 12 wherein z is 3.
14. A compound according to any one of claims 1 to 13 for use in a method of therapy.
15. A pharmaceutical composition comprising a compound according to any one of claims 1 to 13, and a pharmaceutical excipient.
16. The use of a compound according to any one of claims 1 to 13 in the manufacture of a medicament for the treatment of a proliferative disease.
17. A compound according to any one of claims 1 to 13 for use in treating proliferative diseases.




Fig. 1




Fig. 2




Fig. 3

INTERNATIONAL SEARCH REPORT

International application No
PCT/GB2006/001456

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D519/00 A61P35/00 A61K31/5517

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
C07D A61P A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, CHEM ABS Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 1 193 270 A (SPIROGEN LIMITED) 3 April 2002 (2002-04-03) claim 1 compounds 56, 80, 90, 218	1-17
Y	US 4 309 437 A (UEDA ET AL) 5 January 1982 (1982-01-05) the whole document	1-17
P,Y	WO 2005/110423 A (SPIROGEN LIMITED; PEPPER, CHRISTOPHER JOHN; THURSTON, DAVID EDWIN) 24 November 2005 (2005-11-24) the whole document	1-17
P,Y	WO 2005/042535 A (SPIROGEN LIMITED; HOWARD, PHILIP, WILSON; THURSTON, DAVID, EDWIN; GREG) 12 May 2005 (2005-05-12) the whole document	1-17

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier document but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 4 July 2006	Date of mailing of the international search report 14/07/2006
---	---

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer MATES VALDIVIELSO, J
---	---

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/GB2006/001456

Patent document cited in search report	Publication date	Publication date	Patent family member(s)	Publication date
EP 1193270	A	03-04-2002	DK 1109812 T3 EP 1413582 A1	05-09-2005 28-04-2004
<hr style="border-top: 1px dashed black;"/>				
US 4309437	A	05-01-1982	BE 882305 A1 CA 1152985 A1 CH 648848 A5 DE 3010544 A1 FR 2461711 A1 GB 2053894 A JP 1428917 C JP 56015289 A JP 62037631 B LU 82272 A1 NL 8001531 A SE 436882 B SE 8001458 A	16-07-1980 30-08-1983 15-04-1985 29-01-1981 06-02-1981 11-02-1981 09-03-1988 14-02-1981 13-08-1987 01-07-1980 20-01-1981 28-01-1985 18-01-1981
<hr style="border-top: 1px dashed black;"/>				
WO 2005110423	A	24-11-2005	NONE	
<hr style="border-top: 1px dashed black;"/>				
WO 2005042535	A	12-05-2005	NONE	
<hr style="border-top: 1px dashed black;"/>				