



US 20030064474A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2003/0064474 A1**

**Baker et al.** (43) **Pub. Date: Apr. 3, 2003**

(54) **SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC ACIDS ENCODING THE SAME**

(22) Filed: **Sep. 16, 2002**

**Related U.S. Application Data**

(75) Inventors: **Kevin P. Baker**, Darnestown, MD (US); **Dan L. Eaton**, San Rafael, CA (US); **Ellen Filvaroff**, San Francisco, CA (US); **Audrey Goddard**, San Francisco, CA (US); **J. Christopher Grimaldi**, San Francisco, CA (US); **Austin L. Gurney**, Belmont, CA (US); **Victoria Smith**, Burlingame, CA (US); **Jean Philippe Stephan**, Millbrae, CA (US); **Colin K. Watanabe**, Moraga, CA (US); **William I. Wood**, Hillsborough, CA (US); **Zemin Zhang**, Foster City, CA (US); **Sherman Fong**, Alameda, CA (US)

(63) Continuation of application No. 10/197,942, filed on Jul. 18, 2002, which is a continuation of application No. PCT/US01/27099, filed on Aug. 29, 2001.

**Publication Classification**

(51) **Int. Cl.<sup>7</sup>** ..... **C12P 21/02**; C12N 5/06; C07K 14/435; C07H 21/04; C12N 9/00  
(52) **U.S. Cl.** ..... **435/69.1**; 435/183; 435/320.1; 435/325; 530/350; 536/23.2

(57) **ABSTRACT**

Correspondence Address:  
**KNOBBE, MARTENS, OLSON & BEAR, LLP**  
**2040 MAIN STREET**  
**FOURTEENTH FLOOR**  
**IRVINE, CA 92614 (US)**

The present invention is directed to novel polypeptides and to nucleic acid molecules encoding those polypeptides. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention.

(73) Assignee: **Genentech, Inc.**

(21) Appl. No.: **10/245,859**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA102880
><subunit 1 of 1, 184 aa, 1 stop
><MW: 21052, pI: 5.01, NX(S/T): 3
MDNRFATAFVIACVLSLISTIYMAASIGTDFWYEYRSPVQENSSDLNKS IWDEFISDEAD
EKTYNDALFRYNGTVGLWRRCTIPKNNHWYSPPERTESFDVVTKVSTLLEQFMEK FV
DPGNHNSGIDLLR TYLWRCQFLLPFVSLGLMCFGALIGLCACICRSLYPTIATGILHLLA
DTML
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-20

**Transmembrane domain:**

Amino acids 142-163

**N-glycosylation sites:**

Amino acids 42-46; 47-51; 72-76;

**N-myristoylation sites:**

Amino acids 123-129; 154-160; 158-164

**Prokaryotic membrane lipoprotein lipid attachment site:**

Amino acids 152-163

**FIGURE 1**

CGGACGCGTGGGTGCGAGGCGAAGGTGACCGGGGACCGAGCATTTTCAGATCTGCTCGGTAGA  
CCTGGTGCACCACCACC**ATG**TTGGCTGCAAGGCTGGTGTGTCTCCGGACACTACCTTCTAGG  
GTTTTCCACCCAGCTTTTACCAAGGCCTCCCCTGTTGTGAAGAATTCATCACGAAGAATCA  
ATGGCTGTTAACACCTAGCAGGGAATATGCCACCAAAACAAGAATTGGGATCCGGCGTGGGA  
GAACTGGCCAAGAACTCAAAGAGGCAGCATTGGAACCATCGATGGAAAAAATATTTAAAATT  
GATCAGATGGGAAGATGGTTTGTGCTGGAGGGGCTGCTGTTGGTCTTGGAGCATTGTGCTA  
CTATGGCTTGGGACTGTCTAATGAGATTGGAGCTATTGAAAAGGCTGTAATTTGGCCTCAGT  
ATGTCAAGGATAGAATTCATTCACCTATATGTACTTAGCAGGGAGTATTGGTTTAAACAGCT  
TTGTCTGCCATAGCAATCAGCAGAACGCCTGTTCTCATGAACTTCATGATGAGAGGCTCTTG  
GGTGACAATTGGTGTGACCTTTGCAGCCATGGTTGGAGCTGGAATGCTGGTACGATCAATAC  
CATATGACCAGAGCCCAGGCCCAAAGCATCTTGCTTGGTTGCTACATTCTGGTGTGATGGGT  
GCAGTGGTGGCTCCTCTGACAATATTAGGGGGTCTCTTCTCATCAGAGCTGCATGGTACAC  
AGCTGGCATTGTGGGAGGCCTCTCCACTGTGGCCATGTGTGCGCCCAGTGAAAAGTTTCTGA  
ACATGGGTGCACCCCTGGGAGTGGGCCTGGGTCTCGTCTTTGTGTCCTCATTGGGATCTATG  
TTTCTTCCACCTACCACCGTGGCTGGTGCCACTCTTTACTCAGTGGCAATGTACGGTGGATT  
AGTTCTTTTTCAGCATGTTCCCTTCTGTATGATACCCAGAAAGTAATCAAGCGTGCAGAAGTAT  
CACCAATGTATGGAGTTCAAAAATATGATCCCATTAACCTCGATGCTGAGTATCTACATGGAT  
ACATTAATATATTTATGCGAGTTGCAACTATGCTGGCAACTGGAGGCAACAGAAAGAA**TG**  
**A**AGTGACTCAGCTTCTGGCTTCTCTGCTACATCAAATATCTTGTTTAAATGGGGCAGATATGC  
ATTAATAGTTTGTACAAGCAGCTTTCGTTGAAGTTTAGAAGATAAGAAACATGTCATCATA  
TTTAAATGTTCCGGTAATGTGATGCCTCAGGTCTGCCTTTTTTTCTGGAGAATAAATGCAGT  
AATCCTCTCCCAAATAAGCACACACATTTTCAATTCTCATGTTTGAGTGATTTTAAAATGTT  
TTGGTGAATGTGAAAACATAAGTTTGTGTCATGAGAATGTAAGTCTTTTTTCTACTTTAAAA  
TTTAGTAGGTTCACTGAGTAACTAAAATTTAGCAAACCTGTGTTTGCATATTTTTTTGGAGT  
GCAGAATATTGTAATTAATGTCATAAGTGATTTGGAGCTTTGGTAAAGGGACCAGAGAGAAG  
GAGTCACCTGCAGTCTTTTGTTTTTTTAAATACTTAGAACTTAGCACTTGTGTTATTGATTA  
GTGAGGAGCCAGTAAGAAACATCTGGGTATTTGGAAACAAGTGGTCATTGTTACATTCATTT  
GCTGAACTTAACAAAACCTGTTTCATCCTGAAACAGGCACAGGTGATGCATTTCTCCTGCTGTTG  
CTTCTCAGTGCTCTCTTTCCAATATAGATGTGGTCATGTTTGACTTGTACAGAATGTTAATC  
ATACAGAGAATCCTTGATGGAATTATATATGTGTGTTTTACTTTTTGAATGTTACAAAAGGAA  
ATAACTTTAAAACATTTCTCAAGAGAAAATATTCAAAGCATGAAATATGTTGCTTTTTCCAG  
AATACAAACAGTATACTCATG

**FIGURE 2**

MLAARLVCLRTLPSRVFHPAFTKASPVVKN SITKNQWLLTPSREYATKTRIGIRRGRTGQEL  
KEAALEPSMEKIFKIDQMGRWVAVAGGAAVGLGALCYGLGLSNEIGAIEKAVIWPQYVKDRI  
HSTYMYLAGSIGLTALSAIAISRTPVLMNFMMRGSWVTIGVTFAAMVGAGMLVRSIPYDQSP  
GPKHLAWLLHSGVMGAVVAPLTI LGGPLLIRAAWYTAGIVGGLSTVAMCAPSEKFLNMGAPL  
GVGLGLVVFVSSLGSMFLPPTTVAGATLYSVAMYGGLVLF SMFLLYDTQKVIKRAEVSPMYGV  
QKYDPINSMLS IYMDTLNIFMRVATMLATGGNRKK

**FIGURE 3**

CCAATCGCCCGGTGCGGTGGTGCAGGGTCTCGGGCTAGTCATGGCGTCCCCGTCTCGGAGACTGCAGACTAAAC  
CAGTCATTACTTGTTC AAGAGCGTTC TGCTAATCTACACTTTTATTTCTGGATCACTGGCGTTATCCTTCTT  
GCAGTTGGCATTG GGGCAAGGTGAGCCTGGAGAATFACTTTTCTCTTTAAATGAGAAGGCCACCAATGTCCC  
CTTCGTGCTCATTGCTACTGGTACCGTCATTATCTTTTGGGCACCTTTGGTTGTTTTGCTACCTGCCGAGCTT  
CTGCATGGATGCTAAAAC TGTATGCAATGTTCTGACTCTCGTTTTTTGGT CGAACTGGTCGCTGCCATCGTA  
GGATTTGTTTT CAGACATGAGATTAAGAACAGCTTTAAGAATAATTATGAGAAGGCTTTGAAGCAGTATAACTC  
TACAGGAGATTATAGAAGCCATGCAGTAGACAAGATCCAAAATACGTTGCATTGTTGTGGTGT CACCGATTATA  
GAGATTGGACAGATACTAATTATTACTCAGAAAAAGGATTTCC TAAGAGTTGCTGTAAACTTGAAGATTGACT  
CCACAGAGAGATGCAGACAAAGTAAACAATGAAGGTTGTTTTATAAAGGTGATGACCATTATAGAGTCAGAAAT  
GGGAGTCGTTGCAGGAATTTCTTTGGAGTTGCTTGCTTCCAAC TGAATCTTTCTCGCCTACTGCCWCT  
CTCGTGCCATAACAAATAACCAGTATGAGATAGTGTAACCCAATGTATCTGTGGGCCTATTCCTCTCTACCTTT  
AAGGACATTTAGGGTCCCCCTGTGAATTAGAAAGTTGCTTGGCTGGAGA ACTGACAACACTACTFACTGATAG  
ACCAAAAACTACACCAGTAGGTTGATTCAATCAAGATGTATGTAGACCTAAA ACTACACCAATAGGCTGATTC  
AATCAAGATCCGTGCTCGCAGTGGGCTGATTC AATCAAGATGTATGTTTGCTATGTTCTAAGTCCACCTTCTAT  
CCCATT CATGTTAGATCGTTGAAACCCGTATCCCTCTGAAACACTGGAAGAGCTAGTAAATTGTAATGAAGT



## **FIGURE 4**

MASPSRRLQTKPVITCFKSVLLIYTFIFWITGVILLAVGIWGKVSLENYFSLLEKATNVPF  
VLIATGTVIILLGTFGCFATCRASAWMLKLYAMFLTLVFLVELVAAIVGFVFRHEIKNSFKN  
NYEKALKQYNSTGDIRSHAVDKIQNTLHCCGVTDYRDWTDTNYYSEKGFPSCKLEDCTPQ  
RDADKVNNEGCFIKVMTIIIESEMGVVAGISFGVACFQLIGIFLAYCXSRITNNQYEIV

**Important features of the protein:**

**Signal peptide:**

amino acids 1-42

**Transmembrane domains:**

amino acids 19-42, 61-83, 92-114, 209-230,

**N-glycosylation site.**

amino acids 134-138

**Tyrosine kinase phosphorylation site.**

amino acids 160-168, 160-169

**N-myristoylation site.**

amino acids 75-81, 78-84, 210-216, 214-220, 226-232

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 69-80, 211-222

**FIGURE 5**

GGGGCCGCGGTCTAGGGCGGCTACGTGTGTTGCCATAGCGACCATTTTGCATTAACTGGTTG  
GTAGCTTCTATCCTGGGGGCTGAGCGACTGCGGGCCAGCTCTTCCCCTACTCCCTCTCGGCT  
CCTTGTGGCCCAAAGGCCTAACCAGGGTCCGGCGGTCTGGCCTAGGGATCTTCCCCGTTGCC  
CCTTTGGGGCGGG**ATG**GCTGCGGAAGAAGAAGACGAGGTGGAGTGGGTAGTGGAGAGCATCG  
CGGGGTTCCCTGCGAGGCCAGACTGGTCCATCCCCATCTTGGACTTTGTGGAACAGAAATGT  
GAAGTAACTGCAAAGGAGGGCATGTGATAACTCCAGGAAGCCCAGAGCCGGTGATTTTGGT  
GGCCTGTGTTCCCCTTGT'TTTT'GATGATGAAGAAGAAAGCAAATTGACCTATACAGAGATTC  
ATCAGGAATACAAAGAAGACTAGTTGAAAAGCTGTTAGAAGGTTACCTCAAAGAAATTGGAATT  
AATGAAGATCAATTTCAAGAAGCATGCACTTCTCCTCTTGCAAAGACCCATACATCACAGGC  
CATTTTGCAACCTGTGTTGGCAGCAGAAGATTTACTATCTTTAAAGCAATGATGGTCCAGA  
AAAACATTGAAATGCAGCTGCAAGCCATTCGAATAATTCAAGAGAGAAATGGTGTATTACCT  
GACTGCTTAACCGATGGCTCTGATGTGGTCAGTGACCTTGAACACGAAGAGATGAAAATCCT  
GAGGGAAGTTCTTAGAAAAATCAAAGAGGAATATGACCAGGAAGAAGAAAGGAAGAGGAAAA  
AACAGTTATCAGAGGCTAAAACAGAAGAGCCACAGTGCATTCCAGTGAAGCTGCAATAATG  
AATAATTCCCAAGGGGATGGTGAACATTTTGCACACCCACCCTCAGAAGTTAAAATGCATTT  
TGCTAATCAGTCAATAGAACC'TTTGGGAAGAAAAGTGGAAAGGTCTGAAACTCCTCCCTCC  
CACAAAAGGCCCTGAAGATTCCTGGCTTAGAGCATGCGAGCATTGAAGGACCAATAGCAAAC  
TTATCAGTACTTGGAACAGAAGA'ACTTCGGCAACGAGAACACTATCTCAAGCAGAAGAGAGA  
TAAGTTGATGTCCATGAGAAAGGATATGAGGACTAAACAGATACAAAATATGGAGCAGAAAG  
GAAAACCCACTGGGGAGGTAGAGGAAATGACAGAGAAACCAGAAATGACAGCAGAGGAGAAG  
CAAACATTACTAAAGAGGAGATTGCTTGCAGAGAAACTCAAAGAAGAAGTTATTAATAAG**TA**  
**A**ATAATTAAGAACAATTTAACAAAATGGAAGTCAAATTGTCTTAAAAATAAATTATTTAGTC  
CTTACACTG

**FIGURE 6**

MAAEEEEDEVEWVVESIAGFLRGPDWSIPILDFVEQKCEVNCKGGHVITPGSPEPVILVACVP  
LVFDDEEESKLYTEIHQEYKELVEKLLLEGYLKEIGINEDQFQEACTIONSPPLAKTHTSQAILQP  
VLAAEDFTIFKAMMVQKNIEMLQAIRI IQERNGVLPDCLTDGSDVVS DLEHEEMKILREVL  
RKSKEEYDQEEERKRKKQLSEAKTEEPTVHSSEAAIMNNSQGDGEHFHPPSEVKMHFANQS  
IEPLGRKVERSETSSLPQKGLKIPGLEHASIEGPIANLSVLGTEELRQREHYLKQKRDKLMS  
MRKDMRTKQIQNMEQKGPTEGEVEEMTEKPEMTAEEKQTLLKRRLLAEKLKEEVINK

**N-glycosylation sites.**

amino acids 224-228, 246-250, 285-289

**N-myristoylation site.**

amino acids 273-279

**Amidation site.**

amino acids 252-256

**Cytosolic fatty-acid binding proteins.**

amino acids 78-108

**FIGURE 7**

GGGAACGGAAAATGGCGCCTCACGGCCCGGGTAGTCTTACGACCCTGGTGCCCTGGGCTGCCGCCCTGCTCCTC  
GCTCTGGGCGTGGAAAGGGCTCTGGCGCTACCCGAGATATGCACCCAATGTCCAGGGAGCGTGCAAAATTTGTC  
AAAAGTGGCCTTTTATTGTAAAACGACAGGAGAGCTAATGCTGCATGCCCGTTGCTGCCTGAATCAGAAGGGCA  
CCATCTTGGGGCTGGATCTCCAGAACTGTTCTCTGGAGGACCCTGGTCCAAACTTTCATCAGGCACATAACCACT  
GTCATCATAGACCTGCAAGCAAACCCCTCAAAGGTGACTTGGCCAACACCTTCCGTGGCTTTACTCAGCTCCA  
GACTCTGATACTGCCACAACATGTCAACTGTCTGGAGGAATTAATGCCTGGAATACTATCACCTTTATATAG  
ACAACCAATCTGTCAAGGGCAAAGAACCCTTTGCAATAACACTGGGGACCCAGAAATGTGCTCTGAGAATGGA  
TCTTGTGTACCTGATGGTCCAGGTCTTTTGCAGTGTGTTTGTGCTGATGGTTTCCATGGATACAAGTGTATGCG  
CCAGGGCTCGTTCTCACTGCTTATGTTCTTCGGGATTCTGGGAGCCACCACTCTATCCGTCTCCATTCTGCTTT  
GGGCGACCCAGCGCCGAAAAGCCAAGACTTCATGAACTACATAGGTCTTACCATTGACCTAAGATCAATCTGAA  
CTATCTTAGCCAGTCAGGGAGCTCTGCTTCCTAGAAAGGCATCTTTGCCAGTGGATTCCGCTCAAGGTTGAG  
GCCGCCATTGGAAGATGAAAAATTGCACTCCCTTGGTGTAGACAAATACCAGTCCCATTTGGTGTGTTGCCTA  
TAATAAACACTTTTTCTTTTTTNAAAAAAAAAAAAAAAAAAAAAA

## FIGURE 8

**Signal Peptide:**

Amino acids 1-30

**Transmembrane:**

Amino acids 198-212

MAPHGPGSLTTLVPWAAALLLALGVERALALPEICTQCPGSVQNLSKVAFYCKTTREMLHA  
RCCLNQGKGTILGLDLQNCSEDPGPNFHQAHTTVIIDLQANPLKGDLANFRGFTQLQTLIL  
PQHVNCPPGINAWNTITSYIDNQICQGQKNLCNNTGDPEMCPENGSCVPDGPGLLQVCADG  
FHGYKCMRQGSFSLLMFFGILGATTLSVSILLWATQRRKAKTS

**FIGURE 9**

GGGGGAGAAGGCGGCCGAGCCCCAGCTCTCCGAGCACCGGGTCGGAAGCCGCGACCCGAGCC  
GCGCAGGAAGCTGGGACCGGAACCTCGGCGGACCCGCCCCACCCAACCTCACCTGCGCAGGT  
CACCAGCACCCCTCGGAACCCAGAGGCCCGCGCTCTGAAGGTGACCCCCCTGGGGAGGAAGGC  
**GATG**GGCCCCTGCGAGGACGATGGCCCCGCGCCCGCCTCGCCCCGGCCGGCATCCCTGCCGTCCG  
CCTTGTGGCTTCTGTGCACGCTCGGCCTCCAGGGCACCCAGGCCGGGCCACCGCCCCGCGCCC  
CCTGGGCTGCCCGCGGGAGCCGACTGCCTGAACAGCTTTACCGCCGGGGTGCTGGCTTCGT  
GCTGGACACCAACGCCTCGGTGAGCAACGGAGCTACCTTCCTGGAGTCCCCACCCGTGCGCC  
GGGGCTGGGACTGCGTGCGCGCTGCTGCACCACCCAGAACTGCAACTTGGCGCTAGTGGAG  
CTGCAGCCCGACCGCGGGGAGGACGCCATCGCCGCTGCTTCCTCATCAACTGCCTCTACGA  
GCAGAACTTCGTGTGCAAGTTCGCGCCCAGGGAGGGCTTCATCAACTACCTCACGAGGGAAG  
TGTACCGCTCCTACCGCCAGCTGCGGACCCAGGGCTTTGGAGGGTCTGGGATCCCCAAGGCC  
TGGGCAGGCATAGACTTGAAGGTACAACCCAGGAACCCCTGGTGTGAAGGATGTGGAAAA  
CACAGATTGGCGCCTACTGCGGGGTGACACGGATGTCAGGGTAGAGAGGAAAGACCCAAACC  
AGGTGGAACCTGTGGGGACTCAAGGAAGGCACCTACCTGTTCCAGCTGACAGTGACTAGCTCA  
GACCACCCAGAGGACACGGCCAACGTACAGTCACTGTGCTGTCCACCAAGCAGACAGAAGA  
CTACTGCCTCGCATCCAACAAGGTGGGTGCGTGCCTGGGGCTCTTTCCACGCTGGTACTATG  
ACCCACCGGAGCAGATCTGCAAGAGTTTCGTTTATGGAGGCTGCTTGGGCAACAAGAACAAC  
TACCTTCGGGAAGAAGAGTGCATTTCTAGCCTGTGCGGGTGTGCAAGGTGGGCCTTTGAGAGG  
CAGCTCTGGGGCTCAGGCGACTTTCCCCAGGGCCCCCTCCATGGAAAGGCGCCATCCAGTGT  
GCTCTGGCACCTGTCAGCCCACCCAGTTCGCTGCAGCAATGGCTGCTGCATCGACAGTTTC  
CTGGAGTGTGACGACACCCCAACTGCCCCGACGCTCCGACGAGGCTGCCTGTGAAAAATA  
CACGAGTGGCTTTGACGAGCTCCAGCGCATCCATTTCCCCAGTGACAAAGGGCACTGCGTGG  
ACCTGCCAGACACAGGACTCTGCAAGGAGAGCATCCCGCGCTGGTACTACAACCCCTTCAGC  
GAACACTGCGCCCCGCTTTACCTATGGTGGTGTGTTATGGCAACAAGAACAACCTTTGAGGAAGA  
GCAGCAGTGCCTCGAGTCTTGTGCGGCGCATCTCCAAGAAGGATGTGTTTGGCCTGAGGCGGG  
AAATCCCCATTCCCAGCACAGGCTCTGTGGAGATGGCTGTCACAGTGTTCCTGGTCATCTGC  
ATTGTGGTGGTGGTAGCCATCTTGGGTTACTGCTTCTTCAAGAACCAGAGAAAGGACTTCCA  
CGGACACCACCACCACCACCACCACCCTGCCAGTCCACTGTCTCCACTACCGAGGACA  
CGGAGCACCTGGTCTATAACACACCACCCCGCCCCCTCT**TGA**GCCTGGGTCTCACCGGCTCTC  
ACCTGGCCCTGCTTCCCTGCTTGCCAAGGCAGAGGCTGGGCTGGGAAAAAATTTGGAACCAG  
ACTCTTGCCTGTTTCCCAGGCCCACTGTGCCTCAGAGACCAGGGCTCCAGCCCCTCTTGGAG  
AAGTCTCAGCTAAGCTCACGTCTGAGAAAGCTCAAAGGTTTGGAAAGGAGCAGAAAACCCTT  
GGGCCAGAAGTACCAGACTAGATGGACCTGCCTGCATAGGAGTTTGGAGGAAGTTGGAGTTT  
TGTTTCTCTGTTCAAAGCTGCCTGTCCCTACCCCATGGTGTAGGAAGAGGAGTGGGGTGG  
TGTCAGACCCTGGAGGCCCAACCCTGTCCCTCCCGAGCTCCTCTTCCATGCTGTGCGCCCAG  
GGCTGGGAGGAAGGACTTCCCTGTGTAGTTTGTGCTGTAAAGAGTTGCTTTTTGTTTATTTA  
ATGCTGTGGCATGGGTGAAGAGGAGGGGAAGAGGCCTGTTTGGCCTCTCTGTCCTCTCTTCC  
TCTTCCCCAAGATTGAGCTCTCTGCCCTTGATCAGCCCCACCCTGGCCTAGACCAGCAGAC  
AGAGCCAGGAGAGGCTCAGCTGCATTCGCGAGCCCCCACCCTTCTCCAACATCACA  
GCCAGCCCACCCTGGGTAATAAAAGTGGTTTGTGGAAAAAAAAAAAAAAAAAAAAAAAAAAAA

## **FIGURE 10**

MAPARTMARARLAPAGIPAVALWLLCTLGLQGTQAGPPPAPPGLPAGADCLNSFTAGVPGFV  
LDTNASVSNNGATFLESPTVRRGWDCVRACCTTQNCNLALVELQPDRGEDAIAACFLINCLYE  
QNFVCKFAPREGFINYLTVREYRSYRQLRTQGFQGGSGIPKAWAGIDLKVQPQEPLVLKDVEN  
TDWRLLRGDTDVRVERKDPNQVELWGLKEGYL.FQLTVTSSDHPEDTANVTVTVLSTKQTED  
YCLASNKVGRCRGSFPRWYYPTEQICKSFVYGGCLGNKNNYLREEECILACRGVQGGPLRG  
SSGAQATFPQGSMERRHPVCSGTCQPTQFRCSNGCCIDSFLECDTPNCPDASDEAAACEKY  
TSGFDELQRIHFPSDKGHCVDLPDTGLCKESI PRWYYPFSEHCAREFTYGGCYGNKNNFEEE  
QQCLESCRGISKKDVFLRREIPI PSTGSEMAVTVFLVICIVVVVAILGYCFFKNQRKDFH  
GHHHPPTPASSTVSTTEDTEHLVYNHTTRPL

**signal sequence:**

Amino acids 1-35

**transmembrane domain:**

Amino acids 466-483

**N-glycosylation sites:**

Amino acids 66-70;235-239;523-527

**N-myristoylation sites:**

Amino acids 29-35;43-49;161-167;212-218;281-287;282-288;285-291;  
310-316;313-319;422-428;423-429;426-432

**Cell attachment sequence:**

Amino acids 193-199

**Pancreatic trypsin inhibitor (Kunitz) family signatures:**

Amino acids 278-298;419-438





**FIGURE 12**

MRAPGCGRLVLPLLLLAAAAALAEGLKGLKEGETPGNFMEDEQWLSSISQYSGKIKHWNFRDEVEDDYIKSWE  
DNQQGDEALDTTKDFCQKVKCSRHKVCIAQGYQRAMCISRKKLEHRIKQPTVKLHGKDSICKPCHMAQLASVC  
GSDGHTYSSVCKLEQQACLSSKQLAVRCEGPCPCPTEQAATSTADGKPETCTGQDLADLGDRLRDWFQLLHENS  
KQNGSASSVAGPASGLDKSLGASCKDSIGWMFSKLDTSADLFLDQTELAAINLDKYEVCIRPFFNSCDTYKDGR  
VSTAEWCFWREKPPCLAELERIQIQEAAKKPGIFIPSCDEDEGYRKMQCDQSSGDCWRVDQLGLELTGTRT  
HGSPDCDDIVGFSGDFGSGVGVWEDEEEKETEAGEEAEAEAGEAGEADDGGYIW

### FIGURE 13

TGCGGCGACCGTTCGTACACCATGGGCCTCCACCTCCGCCCTACCGTGTGGGGCTGCTCCCG  
GATGGCCTCCTGTTCCCTCTTGCTGCTGCTAATGCTGCTCGCGGACCCAGCGCTCCCGGCCGG  
ACGTCACCCCCCAGTGGTGGTCCCTGGTGATTGGGTAACCAACTGGAAGCCAAGCTGG  
ACAAGCCGACAGTGGTGCACCTACCTCTGCTCCAAGAAGACCGAAAGCTACTTCACAATCTGG  
CTGAACCTGGAAGTGTGCTGCCTGTCATCATTGACTGCTGGATTGACAATATCAGGCTGGT  
TTACAACAAAACATCCAGGGCCACCCAGTTTCCTGATGGTGTGGATGTACGTGTCCCTGGCT  
TTGGGAAGACCTTCTCACTGGAGTTCCTGGACCCCAGCAAAGCAGCGTGGGTTCCTATTTTC  
CACACCATGGTGGAGAGCCTTGTGGGCTGGGGCTACACACGGGGTGGAGATGTCCGAGGGGC  
TCCCTATGACTGGCGCCGAGCCCCAAATGAAAACGGGGCCCTACTTCCTGGCCCTCCGCGAGA  
TGATCGAGGAGATGTACCAGCTGTATGGGGGCCCGTGGTGTGGTTGCCACAGTATGGGC  
AACATGTACACGCTCTACTTTCTGCAGCGGCAGCCGAGGCCCTGGAAGGACAAGTATATCCG  
GGCCTTCGTGTCACTGGGTGCGCCCTGGGGGGGCGTGGCCAAGACCCTGCGCGTCTGGCTT  
CAGGAGACAACAACCGGATCCAGTCATCGGGCCCCGTAAGATCCGGGAGCAGCAGCGGTCA  
GCTGTCTCCACCAGCTGGCTGCTGCCCTACAACCTACACATGGTCACCTGAGAAGGTGTTCTGT  
GCAGACACCACAATCAACTACACACTGCGGGACTACCGCAAGTTCCTCCAGGACATCGGCT  
TTGAAGATGGCTGGCTCATGCGGCAGGACACAGAAGGGCTGGTGGAAAGCCACGATGCCACCT  
GGCGTGCAGCTGCACTGCCTCTATGGTACTGGCGTCCCCACACCAGACTCCTTCTACTATGA  
GAGCTTCCCTGACCGTGACCCATAAATCTGCTTTGGTGACGGCGATGGTACTGTGAACCTGA  
AGAGTGCCCTGCAGTGCCAGGCCCTGGCAGAGCCGCCAGGAGCACCAAGTGTGCTGCAGGAG  
CTGCCAGGCAGCGAGCACATCGAGATGCTGGCCAACGCCACCACCCTGGCCTATCTGAAACG  
TGTGCTCCTTGGGCCCTTGAACTCCTGTGCCACAGGACTCCTGTGGCTCGGGCCGTGGACCTGCT  
GTTGGCCTCTGGGGCTGTCATGGCCCACGCGTTTTTGCAAAGTTTTGTGACTCACCATTCAAGG  
CCCCGAGTCTTGGACTGTGAAGCATCTGCCATGGGGAAGTGTGTTTTGTTATCCTTTCTCTG  
TGGCAGTGAAGAAGGAAGAAATGAGAGTCTAGACTCAAGGGACTGAGTGGCAAGAATGCT  
GCTGATGGTGGAACTGCTGTGACCTTAGGACTGGCTCCACAGGGTGGACTGGCTGGGCCCTG  
GTCCCAGTCCCTGCCTGGGGCCATGTGTCCCCCTATTCCCTGTGGGCTTTTCATACTGTCCTA  
CTGGGGCCCTGGCCCCGCAGCCTTCCTATGAGGGATGTTACTGGGCTGTGGTTCCTGTACCCAG  
AGGTCCCAGGGATCGGCTCCTGGCCCCCTCGGGTGACCCTTCCCACACACCAGCCACAGATAG  
GCCTGCCACTGGTTCATGGGTAGCTAGAGCTGCTGGCTTCCCTGTGGCTTAGCTGGTGGCCAG  
CCTGACTGGCTTCTGGGCGAGCCTAGTAGCTCCTGCAGGCAGGGGAGTTTGTGCTTCT  
TCGTGGTTCCAGGCCCTGGGACATCTCACTCCACTCCTACCTCCCTTACCACCAGGAGCAT  
TCAAGCTCTGGATTGGGCAGCAGATGTGCCCCAGTCCCGCAGGCTGTGTTCCAGGGGCCCT  
GATTTCCCTCGGATGTGCTATTGGCCCCAGGACTGAAGCTGCCTCCCTTACCCTGGGACTGT  
GGTTCCAAGGATGAGAGCAGGGGTTGGAGCCATGGCCTTCTGGGAACCTATGGAGAAAGGGA  
ATCCAAGGAAGCAGCCAAGGCTGCTCGCAGCTTCCCTGAGCTGCACCTCTTGCTAACCCAC  
CATCACACTGCCACCCTGCCCTAGGGTCTCACTAGTACCAAGTGGGTGAGCAGCAGGGCTGAG  
GATGGGGCTCCTATCCACCCTGGCCAGCACCCAGCTTAGTGTGCTGGGACTAGCCCAGAAACTT  
GAATGGGACCCTGAGAGAGCCAGGGGTCCCCTGAGGCCCCCTAGGGGCTTTCTGTCTGCC  
CAGGGTGTCCATGGATCTCCCTGTGGCAGCAGGCATGGAGAGTCAGGGCTGCCTTCATGGC  
AGTAGGCTCTAAGTGGGTGACTGGCCACAGGCCGAGAAAAGGGTACAGCCTCTAGGTGGGGT  
TCCCAAAGACGCCTTCAGGCTGGACTGAGCTGCTCTCCACAGGGTTTCTGTGCAGCTGGAT  
TTTCTCTGTTGCATACATGCCTGGCATCTGTCTCCCTTGTTCCTGAGTGGCCCCACATGGG  
GCTCTGAGCAGGCTGTATCTGGATTCTGGCAATAAAAAGTACTCTGGATGCTGTAAAAA  
AAAAAAAAAAAAA

## FIGURE 14

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA44189
><subunit 1 of 1, 412 aa, 1 stop
><MW: 46658, pI: 6.65, NX(S/T): 4
MGLHLRPYRVGLLPDGLLFLLLLMLLADPALPAGRHPVVLVPGDLGNQLEAKLDKPTV
VHYLCSKKTESYFTIWLNLLELLLPVIIDCWIDNIRLVYNKTSRATQFPDGVDRVPGFGK
TFSLEFLDPSKSSVGSYFHTMVESLVGWGYTRGEDVRGAPYDWRRAPNENGPYFLALREM
IEEMYQLYGGPVVLVAHSMGNMYTLYFLQRQPQAWKDKYIRAFVSLGAPWGGVAKTLRVL
ASGDNNRIPVIGPLKIREQORSVSTSWLLPYNYTWSPEKVFVQTPPTINYTLRDYRKFFQ
DIGFEDGWLMRQDTEGLVEATMPPGVQLHCLYGTGVPTPDSFYYESFPDRDPKICFGDGD
GTVNLKSALQCQAWQSRQEHQVLLQELPGSEHIEMLANATTLAYLKRVLVLLGP
```

**Signal peptide:**

Amino acids 1-28

**Potential lipid substrate binding site:**

Amino acids 147-164

**N-glycosylation sites:**

Amino acids 99-103;273-277;289-293;398-402

**Lipases, serine proteins family:**

Amino acids 189-202

**Beta-transducin family Trp-Asp repeat:**

Amino acids 353-366

**Tyrosine kinase phosphorylation site:**

Amino acids 165-174;178-186

**N-myristoylation sites:**

Amino acids 200-206;227-233;232-238;316-322

**FIGURE 15**

CAGAGCAGATA**ATG**GCAAGCATGGCTGCCGTGCTCACCTGGGCTCTGGCTCTTCTTTCAGCG  
TTTTCGGCCACCCAGGCACGGAAAGGCTTCTGGGACTACTTCAGCCAGACCAGCGGGGACAA  
AGGCAGGGTGGAGCAGATCCATCAGCAGAAGATGGCTCGCGAGCCC GCGACCCTGAAAGACA  
GCCTTGAGCAAGACCTCAACAATATGAACAAGTTCCTGGAAAAGCTGAGGCCTCTGAGTGGG  
AGCGAGGCTCCTCGGCTCCACAGGACCCGGTGGGCATGCGGCGGCAGCTGCAGGAGGAGTTG  
GAGGAGGTGAAGGCTCGCCTCCAGCCCTACATGGCAGAGGCGCACGAGCTGGTGGGCTGGAA  
TTTGGAGGGCTTGCGGCAGCAACTGAAGCCCTACACGATGGATCTGATGGAGCAGGTGGCCC  
TGCGCGTGCAGGAGCTGCAGGAGCAGTTGCGCGTGGTGGGGGAAGACACCAAGGCCCAGTTG  
CTGGGGGGCGTGGACGAGGCTTGGGCTTTGCTGCAGGGACTGCAGAGCCGCGTGGTGCACCA  
CACCGGCCGCTTCAAAGAGCTCTTCCACCCATACGCCGAGAGCCTGGTGAGCGGCATCGGGC  
GCCACGTGCAGGAGCTGCACCGCAGTGTGGCTCCGCACGCCCCCGCCAGCCCCGCGCGCCTC  
AGTCGCTGCGTGCAGGTGCTCTCCCGGAAGCTCACGCTCAAGGCCAAGGCCCTGCACGCACG  
CATCCAGCAGAACCTGGACCAGCTGCGCGAAGAGCTCAGCAGAGCCTTTGCAGGCACTGGGA  
CTGAGGAAGGGGCCGGCCCCGGACCCCT**TAGAT**GCTCTCCGAGGAGGTGCGCCAGCGACTTCAG  
GCTTTCCGCCAGGACACCTACCTGCAGATAGCTGCCTTCACTCGCGCCATCGACCAGGAGAC  
TGAGGAGGTCCAGCAGCAGCTGGCGCCACCTCCACCAGGCCACAGTGCCTTCGCCCCAGAGT  
TTCAACAAACAGACAGTGGCAAGGTTCTGAGCAAGCTGCAGGCCCGTCTGGATGACCTGTGG  
GAAGACATCACTCACAGCCTTCATGACCAGGGCCACAGCCATCTGGGGGACCCCTGAGGATC  
TACCTGCCCAGGCCCATTCCCAGCTTCTTGTCTGGGGAGCCTTGGCTCTGAGCCTCTAGCAT  
GGTTCAGTCCCTGAAAGTGGCCTGTTGGGTGGAGGGTGGAAAGTCCCTGTGCAGGACAGGGAG  
GCCACCAAAGGGGCTGCTGTCTCCTGCATATCCAGCCTCCTGCGACTCCCCAATCTGGATGC  
ATTACATTACCAGGCTTTGCAAA  
AAAAAA

## **FIGURE 16**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA48303
><subunit 1 of 1, 274 aa, 1 stop
><MW: 30754, pI: 7.77, NX(S/T): 0
MASMAAVLTWALALLSAFSATQARKGFWDYFSQTS GDKGRVEQIHQQMAREPATLKDSL
EQDLNMMNKFLEKLRPLSGSEAPRLPQDPVGMRRQLQEELEEVKARLQPYMAEAHELVGW
NLEGLRQQLKPYTMDLMEQVALRVQELQEQLRVVGEDTKAQLLGGVDEAWALLQGLQSRV
VHHTGRFKELFHPYAESLVSGIGRHSVQELHRSVAPHAPASPARLSRCVQVLSRKLTLKAK
ALHARIQQNLDQLREELSRAFAGTGTEEGAGPDP
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-23

**Glycosaminoglycan attachment site:**

Amino acids 200-204

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 233-237

**N-myristoylation sites:**

Amino acids 165-171;265-271

**FIGURE 17**

CTAAGAGGACAAG**ATG**AGGCCCGGCTCTCATTCTCCTAGCCCTTCTGTTCTTCCTTGGCC  
AAGCTGCAGGGGATTTGGGGGATGTGGGACCTCCAATTCAGCCCCGGCTTCAGCTCTTTC  
CCAGGTGTTGACTCCAGCTCCAGCTTCAGCTCCAGCTCCAGGTGGGCTCCAGCTCCAGCCG  
CAGCTTAGGCAGCGGAGGTTCTGTGTCCCAGTTGTTTTCCAATTCACCGGCTCCGTGGATG  
ACCGTGGGACCTGCCAGTGCTCTGTTCCCTGCCAGACACCACCTTTCCCGTGGACAGAGTG  
GAACGCTTGAATTCACAGCTCATGTTCTTCTCAGAAGTTTGAGAAAGAACTTTCTAAAGTG  
AGGGAATATGTCCAATTAATTAGTGTGTATGAAAAGAACTGTAAACCTAACTGTCCGAAT  
TGACATCATGGAGAAGGATACCATTTCTTACACTGAACTGGACTTCGAGCTGATCAAGGTAG  
AAGTGAAGGAGATGGAAAACCTGGTCATACAGCTGAAGGAGAGTTTTGGTGGAAAGCTCAGAA  
ATTGTTGACCAGCTGGAGGTGGAGATAAGAAATATGACTCTCTTGGTAGAGAAGCTTGAGAC  
ACTAGACAAAACAATGTCCTTGCCATTCGCCGAGAAATCGTGGCTCTGAAGACCAAGCTGA  
AAGAGTGTGAGGCCTCTAAAGATCAAACACCCCTGTCGTCCACCCTCCTCCCACTCCAGGG  
AGCTGTGGTCATGGTGGTGTGGTGAACATCAGCAAACCGTCTGTGGTTCAGCTCAACTGGAG  
AGGGTTTTCTTATCTATATGGTGTCTGGGGTAGGGATTACTCTCCCAGCATCCAAACAAAG  
GACTGTATTGGGTGGCGCCATTGAATACAGATGGGAGACTGTTGGAGTATTATAGACTGTAC  
AACACACTGGATGATTTGCTATTGTATATAAATGCTCGAGAGTTGCCGATCACCTATGGCCA  
AGGTAGTGGTACAGCAGTTTACAACAACAACATGTACGTCAACATGTACAACACCGGGAATA  
TTGCCAGAGTTAACCTGACCACCAACACGATTGCTGTGACTCAAACCTCTCCCTAATGCTGCC  
TATAATAACCGCTTTTCATATGCTAATGTTGCTTGGCAAGATATTGACTTTGCTGTGGATGA  
GAATGGATTGTGGGTTATTTATTCAACTGAAGCCAGCACTGGTAACATGGTATTAGTAAAC  
TCAATGACACCACACTTCAGGTGCTAAACACTTGGTATACCAAGCAGTATAAACCATCTGCT  
TCTAACGCCTTCATGGTATGTGGGGTCTGTATGCCACCCGTAATGAAACCCAGAACAGA  
AGAGATTTTTTACTATTATGACACAAACACAGGGAAAGAGGGCAAACCTAGACATTGTAATGC  
ATAAGATGCAGGAAAAGTGCAGAGCATTAACTATAACCCTTTTGACCAGAACTTTATGTC  
TATAACGATGGTTACCTTCTGAATTATGATCTTTCTGTCTTGCAGAAGCCCCAG**TAA**GCTGT  
TTAGGAGTTAGGGTGAAAGAGAAAATGTTTGGTGA AAAAATAGTCTTCTCCACTTACTTAGA  
TATCTGCAGGGGTGTCTAAAAGTGTGTTTCAATTTTGCAGCAATGTTTAGGTGCATAGTTCTAC  
CACACTAGAGATCTAGGACATTTGTCTTGATTTGGTGGAGTTCTCTTGGGAATCATCTGCCTC  
TTCAGGCGCATTTTGAATAAAGTCTGTCTAGGGTGGGATTGTCAGAGGTCTAGGGGCACTG  
TGGGCCTAGTGAAGCCTACTGTGAGGAGGCTTCACTAGAAGCCTTAAATTAGGAATTAAGGA  
ACTTAAACTCAGTATGGCGTCTAGGGATTCTTTGTACAGGAAATATTGCCCAATGACTAGT  
CCTCATCCATGTAGCACCCTAATCTTCCATGCCTGGAAGAACTGGGGACTTAGTTAGG  
TAGATTAATATCTGGAGCTCCTCGAGGGACCAATCTCCAACCTTTTTTTTCCCCTCACTAGC  
ACCTGGAATGATGCTTTGTATGTGGCAGATAAGTAAATTTGGCATGCTTATATATTTCTACAT  
CTGTAAAGTGTGCTGAGTTTTATGGAGAGAGGCCTTTTTATGCATTAAATTTGTACATGGCAAATAA  
ATCCCAGAAGGATCTGTAGATGAGGCACCTGCTTTTTCTTTCTCTCATTTGTCACCTTACT  
AAAAGTCAGTAGAATCTTCTACCTCATAACTTCCTTCCAAGGCAGCTCAGAAGATTAGAAC  
CAGACTTACTAACCAATTCACCCCCACCAACCCCTTCTACTGCCTACTTTAAAAAAAT  
AATAGTTTTCTATGGAACCTGATCTAAGATTAGAAAAATTAATTTCTTTAATTTTATTATGG  
ACTTTTATTTACATGACTCTAAGACTATAAGAAAATCTGATGGCAGTGACAAAGTGCTAGCA  
TTTATTGTTATCTAATAAAGACCTTGGAGCATATGTGCAACTTATGAGTGTATCAGTTGTTG  
CATGTAATTTTGCCTTTGTTTAAAGCCTGGAACCTGTAAGAAAATGAAAATTTAATTTTTTT  
TTCTAGGACGAGCTATAGAAAAGCTATTGAGAGTATCTAGTTAATCAGTGCAGTAGTTGGAA  
ACCTTGCTGGTGTATGTGATGTGCTTCTGTGCTTTTGAATGACTTTATCATCTAGTCTTTGT  
CTATTTTTCCCTTGATGTTCAAGTCCTAGTCTATAGGATTGGCAGTTTAAATGCTTTACTCC  
CCCTTTTAAAAATAAATGATTA AAAATGTGCTTTGAAAAAAAAAAAAAAAAAAAAAAAAAAAA

## FIGURE 18

```
</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA48320
<subunit 1 of 1, 510 aa, 1 stop
<MW: 57280, pI: 5.61, NX(S/T): 6
MRPGLSFLLALLFFLQQAAGDLGDVGPPIPSPGFSSFPGVDSSTSSSRSGSSSSRSL
GSGGSVSQLF'SNFTGSVDDRGTCCQCSVSLPDTTFPVDRVERLEFTAHVLSQKFEKELSKV
REYVQLISVYEKLLNLTVRIDIMEKDTISYTELDFELIKVEVKEMEKLVIQLKESFGGS
SEIVDQLEVEIRNMTLLVEKLETLDKNNVLAIRREIVALKTKLKECEASKDQNTPVVHPP
PTPGSCGHGGVNVNISKPSVVQLNWRGFSYLYGAWGRDYSPQHHPNKGLYWVAPLNTDGRLL
EYYRLYNLTDLLLYINARELRITYGQGSQTAVYNNMYVNMNTGNIARVNLTNTIIV
TQTLPNAAYNRRFSYANVAWQDIDFAVDENGLWVIYSTEASTGNMVISKLNDDTLQVLNT
WYTKQYKPSASNAFMVCGVLYATRMTNTRTEEIIFYYYDTNTGKEGKLDIVMHKMQEKVQS
INYNPFDQKLYVYNDGYLLNYDLSVLQKPQ
```

**Important features:**

**Signal peptide:**

Amino acids 1-20

**N-glycosylation sites:**

Amino acids 72-76;136-140;193-197;253-257;352-356;  
411-415

**Tyrosine kinase phosphorylation site:**

Amino acids 449-457

**N-myristoylation sites:**

Amino acids 16-22;39-45;53-59;61-67;63-69;81-87;  
249-255;326-332;328-334;438-444

**Legume lectins beta-chain proteins:**

Amino acids 20-40

**HBGF/FGF family proteins:**

Amino acids 338-366

**FIGURE 19**

GCACCGCAGACGGCGCGGATCGCAGGGAGCCGGTCCGCCGCCGGAACGGGAGCCTGGGTGTG  
CGTGTGGAGTCCGGACTCGTGGGAGACGATCGCG**ATG**AACACGGTGCTGTCCGGGGCGAACT  
CACTGTTCGCCTTCTCGCTGAGCGTGATGGCGGGCGCTCACCTTCGGCTGCTTCATCACCACC  
GCCTTCAAAGACAGGAGCGTCCCGGTGCGGCTGCACGTCTCGCGGATCATGCTAAAAAATGT  
AGAAGATTTCACTGGACCTAGAGAAAGAAGTGATCTGGGATTTATCACATTTGATATAACTG  
CTGATCTAGAGAATATATTTGATTGGAATGTTAAGCAGTTGTTTCTTTATTTATCAGCAGAA  
TATTCAACAAAAAATAATGCTCTGAACCAAGTTGTCCTATGGGACAAGATTGTTTTGAGAGG  
TGATAATCCGAAGCTGCTGCTGAAAGATATGAAAACAAAATATTTTTCTTTGACGATGGAA  
ATGGTCTCAAGGGAAACAGGAATGTCACCTTTGACCCTGTCTTGGAACGTCGTACCAAATGCT  
GGAATTCTACCTCTTGTGACAGGATCAGGACACGTATCTGTCCCATTTCCAGATACATATGA  
AATAACGAAGAGTTAT**TAA**ATTATTCTGAATTTGAAACAAAA



## **FIGURE 20**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA56049
><subunit 1 of 1, 180 aa, 1 stop
><MW: 20313, pI: 8.91, NX(S/T): 1
MNTVLSRANSLFAFSLSVMAALTFGCFITTAFFKDRSVPVRLHVSRIMLKKNVEDFTGPRER
SDLGFITFDITADLENI FDWNVKQLFLYLSAEYSTKNNALNQVVLWDKIVLRGDNPKLLL
KDMKTKYFFEDDGNGLGKGNRVTLTLSWNVVVPNAGILPLVTGSGHVSVPFPDTYEITKSY
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-25

**Transmembrane domain:**

Amino acids 149-164

**N-glycosylation site:**

Amino acids 141-145

**N-myristoylation sites:**

Amino acids 25-31;135-141

**Cell attachment sequence:**

Amino acids 112-115

**TonB-dependent receptor proteins signature 1:**

Amino acids 1-21

## **FIGURE 21**

AAACTTGACGCC**ATG**AAGATCCCGGTCCTTCCTGCCGTGGTGCTCCTCTCCCTCCTGGTGCT  
CCTCTGCCAGGGAGCCACCCTGGGTGGTCCTGAGGAAGAAAGCACCATTGAGAATTATG  
CGTCACGACCCGAGGCCTTTAACACCCCGTTCCTGAACATCGACAAATTGCGATCTGCGTTT  
AAGGCTGATGAGTTCCTGAACTGGCACGCCCTTTTGAGTCTATCAAAGGAAACTTCCTTT  
CCTCAACTGGGATGCCTTTCCTAAGCTGAAAGGACTGAGGAGCGCAACTCCTGATGCCCAG**T**  
**GA**CCATGACCTCCACTGGAAGAGGGGGCTAGCGTGAGCGCTGATTCTCAACCTACCATAACT  
CTTTCCTGCCTCAGGAACTCCAATAAAACATTTTCCATCCAAA

## FIGURE 22

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA57694
><subunit 1 of 1, 99 aa, 1 stop
><MW: 11050, pI: 7.47, NX(S/T): 0
MKIPVLPVAVLLSLLVLHSAQGATLGGPEEESTIENYASRPEAFNTPFLNIDKLRSFAFKA
DEFLNWHALFESIKRKLPFLNWDAFPCLKGLRSATPDAQ
```

**Important features:**

**Signal peptide:**

Amino acids 1-22

**N-myristoylation sites:**

Amino acids 22-28;90-96

**Homologous region to Peroxidase:**

Amino acids 16-48

**FIGURE 23**

TCTCAGACTCTTGAAGGGGCTATACTAGACACACAAAGACAGCCCCAAGAAGGACGGTGGAGTAGTGTCCCTCGCTAAAAGACAGTAGAT**ATG**CAACGCCTCTTGCTCCTGCCCTTTCTCCTGCTGGGAACAGTTTCTGCTTTCATCTGGAGAATGATGCCCCCATCTGGAGAGCCTAGAGACACAGGCAGACCTAGGCCAGGATCTGGATAGTTCAAAGGAGCAGGAGAGAGACTTGGCTCTGACGGAGGAGGTGATTCAGGCAGAGGGAGAGGAGGTCAAGGCTTCTGCCTGTCAAGACAACCTTGAGGATGAGGAAGCCATGGAGTCGGACCCAGCTGCCTTAGACAAGGACTTCCAGTGCCCCAGGGAAGAAGACATTGTTGAAGTGCAGGGAAGTCCAAGGTGCAAGACCTGCCGCTACCTATTGGTGCGGACTCCTAAAACCTTTGCAGAAGCTCAGAATGTCTGCAGCAGATGCTACGGAGGCAACCTTGTCTCTATCCATGACTTCAACTTCAACTATCGCATTCAAGTGCTGCACTAGCACAGTCAACAAGCCCAGGTCTGGATTGGAGGCAACCTCAGGGGCTGGTTCCTGTGGAAGCGGTTTTGCTGGACTGATGGGAGCCACTGGAATTTGCTTACTGGTCCCCAGGGCAACCTGGGAATGGGCAAGCTCCTGTGTGGCCCTATGCACCAAAGGAGGTTATTGGCGACGAGCTCAATGCGACAAGCAACTGCCCTTCGTCTGCTCCTT**TAA**GCCAGCGGCACGGAGACCCTGCCAGCAGCTCCCTCCCGTCCCCAACCTCTCCTGCTCATAAATCCAGACTTCCCACAGCAAAAAAAAAAAAAAAAAAAAA

## FIGURE 24

```
</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA59208
<subunit 1 of 1, 225 aa, 1 stop
<MW: 25447, pI: 4.79, NX(S/T): 0
MQRLLLLPFLLLGTVSALHLENDAPHLESLETQADLGQDLDSKEQERDLALTEEVIAE
GEEVKASACQDNFEDEEAMESDPAALDKDFQCPREEDIVEVQGS PRCKTCRYLLV RTPKT
FAEAQNVCSR CYGGNLVSIHDFNFYRIQCCTSTVNQAQVWIGGNLRGWFLWKRF CWT DG
SHWNFAYWSPGQPGNGQGSCVALCTKGGYWRRAQCDKQLPFVCSF
```

**Important features:**

**Signal peptide:**

Amino acids 1-17

**N-myristoylation sites:**

Amino acids 13-19;103-109;134-140;164-170;  
180-186;191-197;194-200;196-202;  
198-204

**C-type lectin domain signature:**

Amino acids 200-224

**FIGURE 25**

CAACAGAAGCCAAGAAGGAAGCCGTCTATCTTGTGGCGATC**ATG**TATAAGCTGGCCTCCTGC  
TGTTTGCTTTTCACAGGATTCTTAAATCCTCTCTTATCTCTCCTCCTCCTTGACTCCAGGGA  
AATATCCTTTCAACTCTCAGCACCTCATGAAGACGCGCGCTTAACTCCGGAGGAGCTAGAAA  
GAGCTTCCCTTCTACAGATATTGCCAGAGATGCTGGGTGCAGAAAGAGGGGATATTCTCAGG  
AAAGCAGACTCAAGTACCAACATTTTTAACCCAAGAGGAAATTTGAGAAAGTTTCAGGATTT  
CTCTGGACAAGATCCTAACATTTTACTGAGTCATCTTTTGGCCAGAATCTGGAAACCATA  
AGAAACGTGAGACTCCTGATTGCTTCTGGAAATACTGTGTCT**TGA**AGTGAAATAAGCATCTGT  
TAGTCAGCTCAGAAACACCCATCTTAGAATATGAAAAATAACACAATGCTTGATTTGAAAAC  
AGTGTGGAGAAAAACTAGGCAAACCTACACCCTGTTTCATTGTTACCTGGAAAATAAATCCTCT  
ATGTTTTGCACAAAAAAAAAAAAAAAAA

## **FIGURE 26**

</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA59214

<subunit i of 1, 124 aa, 1 stop

<MW: 14284, pI: 8.14, NX(S/T): 0

MYKLASCCLLEFTGFLNPLLSLPLLLDSREISFQLSAPHEDARLTPEELERASLLQILPEML  
GAERGDILRKADSSTNIFNPRGNLRKFQDFSGQDPNILLSHLLARIWKPYKKRETPDCFW  
KYCV

**Important features:**

**Signal peptide:**

Amino acids 1-20

**Urotensin II signature:**

Amino acids 118-124

**Cell attachment sequence:**

Amino acids 64-67

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 112-116

**N-myristoylation sites:**

Amino acids 61-67;92-98

**FIGURE 27**

CAAGTAAATGCAGCACTAGTGGGTGGGATTGAGGTATGCCCTGGTGCATAAATAGAGACTCA  
GCTGTGCTGGCACACTCAGAAGCTTGGACCGCATCCTAGCCGCCGACTCACACAAGGCAGGT  
GGGTGAGGAAATCCAGAGTTGCC**ATG**GAGAAAATTCCAGTGTGAGCATTCTTGCTCCTTGTG  
GCCCTCTCTTACACTCTGGCCAGAGATACCACAGTCAAACCTGGAGCCAAAAAGGACACAAA  
GGACTCTCGACCCAAACTGCCCCAGACCTCTCCAGAGGTTGGGGTGACCAACTCATCTGGA  
CTCAGACATATGAAGAAGCTCTATATAAAATCCAAGACAAGCAACAAACCTTGGATGATTATT  
CATCACTTGGATGAGTGCCACACAGTCAAGCTTTAAAGAAAGTGTTTGCTGAAAATAAAGA  
AATCCAGAAATTGGCAGAGCAGTTTGTCTCCTCAATCTGGTTTATGAAACAACCTGACAAAC  
ACCTTTCTCCTGATGGCCAGTATGTCCCAGGATTATGTTTGTGACCCATCTCTGACAGTT  
AGAGCCGATATCACTGGAAGATATTCAAATCGTCTCTATGCTTACGAACCTGCAGATACAGC  
TCTGTTGCTTGACAACATGAAGAAAGCTCTCAAGTTGCTGAAGACTGAATTG**TAA**AGAAAAA  
AAATCTCCAAGCCCTTCTGTCTGTCAGGCCTTGAGACTTGAAACCAGAAGAAGTGTGAGAAG  
ACTGGCTAGTGTGGAAGCATAGTGAACACACTGATTAGGTTATGGTTTAATGTTACAACAAC  
TATTTTTTAAGAAAAACAAGTTTTAGAAAATTTGGTTTCAAGTGTACATGTGTGAAAACAATA  
TTGTATACTACCATAGTGAGCCATGATTTTCTAAAAAATAAATGTTA



## **FIGURE 28**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA59485
><subunit 1 of 1, 175 aa, 1 stop
><MW: 19979, pI: 9.26, NX(S/T): 0
MEKIPVSAFLLLVALSYTLARDTTVKPGAKKDTKDSRPKLPQTLSRGWGDQLIWTQTYEE
ALYKSKTSNKPLMIIHHLDECPHSQALKKVFAENKEIQKLAEQFVLLNLVYETTDKHLSP
DGQYVPRIMFVDPSLTVRADITGRYSNRLYAYEPADTALLLDNMKKALKLLKTEL
```

**Important features:**

**Signal peptide:**

Amino acids 1-20

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 30-34

**FIGURE 29**

AAGACCTCTCTTTTCGCTGTTTGAGAGTCTCTCGGCTCAAGGACCGGGAGGTAAGAGGTT  
TGGGACTGCCCCGGCAACTCCAGGGTGTCTGGTCCACGACCTATCCTAGGCGCC**ATGGGT**  
GTGATAGGTATACAGCTGGTTGTTACCATGGTGATGGCCAGTGTGCATGCAGAAGATTATA  
CCTCACTATTCTCTTGCTCGATGGCTACTCTGTAATGGCAGTTTGAGGTGGTATCAACAT  
CCTACAGAAGAAGAATTAAGAATTCTTGCAGGGAAACAACAAAAAGGGAAAACCAAAAA  
GATAGGAAATATAATGGTCACATTGAAAGTAAGCCATTAACCATTCCAAAGGATATTGAC  
CTTCATCTAGAAACAAAGTCAGTTACAGAAGTGGATACTTTAGCATTGCATTACTTTCCA  
GAATACCAGTGGCTGGTGGATTTACAGTGGCTGCTACAGTTGTGTATCTAGTAACTGAA  
GTCTACTACAATTTTATGAAGCCTACACAGGAAATGAATATCAGCTTAGTCTGGTGCCTA  
CTTGTTTTGTCTTTTGCAATCAAAGTTCTATTTTCATTAACACACTATTTTAAAGTA  
GAAGATGGTGGTCAAAGATCTGTTTGTGTCACCTTTGGATTTTTTTTCTTTGTCAAAGCA  
ATGGCAGTGTTGATTGTAACAGAAAATTATCTGGAATTTGGACTTGAAACAGGGTTTACA  
AATTTTTTCAGACAGTGCATGTCAGTTTCTTGAAAAGCAAGGTTTAGAATCTCAGAGTCTT  
GTTTCAAACCTTACTTTCAAATTTTTCCTGGCTATTTTCTGTTTCATTTCATTGGGGCTTTT  
TTGACATTTCTGGATTACGACTGGCTCAAATGCATCTGGATGCCCTGAATTTGGCAACA  
GAAAAAATTACACAACTTTACTTCATATCAACTTCTTGGCACCTTATTTATGGTTTTG  
CTCTGGGTAAAACCAATCACCAAAGACTACATTATGAACCCACCCTGGGCAAAGAAATT  
TCCCATCTGGAAGAT**TGA**AGATAATAGTATCTAACTCACAAGGTTATCATTGGAATAAAT  
GAAAGAACACATGTAATGCAACCAGCTGGAATTAAGTGCTTAATAAATGTTCTTTTCACT  
GCTTTCCTCATCAGAATTAATAAGAAATACTTGACTAGT

## **FIGURE 30**

</usr/seqdb2/sst/DNA/Dnaseqs.full/ss.DNA64966

<subunit 1 of 1, 307 aa, 1 stop

<MW: 35098, pI: 8.11, NX(S/T): 3

MGVIGIQLVVTMVMASVMQKII PHYSLARWLLCNGSLRWYQHPTTEEELRILAGKQKQKGT  
KKDRKYNGHIESKPLTI PKDIDLHLETKSVTEVDTLALHYFPEYQWLVDFTVAATVVYLV  
TEVYYNFMKPTQEMNISLVWCLLVLSFAIKVLFSLTTHYFKVEDGGERSVCVTFGFFFFV  
KAMAVLIVTENYLEFGLETGFTNFSDSAMQFLEKQGLSQSPVSKLTFKFFLAI FCSFIG  
AFLTFPGLRLAQMHLDALNLATEKITQTLLHINFLAPLFMVLLWVKPITKDYIMNPPLGK  
EISPSGR

**Important features:**

**Signal peptide:**

Amino acids 1-15

**Transmembrane domains:**

Amino acids 134-157;169-189;230-248;272-285

**N-glycosylation sites:**

Amino acids 34-38;135-139;203-207

**ATP/GTP-binding site motif A (P-loop):**

Amino acids 53-61

**Tyrosine kinase phosphorylation site:**

Amino acids 59-67

**N-myristoylation sites:**

Amino acids 165-171;196-202;240-246;247-253

**FIGURE 31**

GTAGCATAGTGTGCAGTTCACTGGACCAAAAGCTTTGGCTGCACCTCTTCTGGAAAGCTGGCC  
**ATG**GGGCTCTTCATGATCATTGCAATTCTGCTGTTCCAGAAACCCACAGTAACCGAACAACT  
TAAGAAGTGCTGGAATAACTATGTACAAGGACATTGCAGGAAAATCTGCAGAGTAAATGAAG  
TGCCTGAGGCACTATGTGAAAATGGGAGATACTGTTGCCTCAATATCAAGGAACTGGAAGCA  
TGTAATAAAATTACAAAGCCACCTCGTCCAAAGCCAGCAACACTTGCACTGACTCTTCAAGA  
CTATGTTACAATAATAGAAAATTTCCCAAGCCTGAAGACACAGTCTACAT**TAA**ATCAAATACA  
ATTTTCGTTTTCACTTGCTTCTCAACCTAGTCTAATAAACTAAGGTGATGAGATATACATCTT  
CTTCCTTCTGGTTTCTTGATCCTTAAAATGACCTTCGAGCATATTCTAATAAAGTGCATTGC  
CAGTTAAAAAAAAAAAA

## FIGURE 32

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA82403
><subunit 1 of 1, 99 aa, 1 stop
><MW: 11343, pI: 9.17, NX(S/T): 0
MGLFMIIAILLFQKPTVTEQLKKCWNNYVQGHCRKICRVNEVPEALCENGRYCCLNKEL
EACKKITKPPRPKPATLALTLQDYVTIIENFPSLKTQST
```

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**  
Amino acids 64-68

**FIGURE 33**

CGGACGCGTGGGCGCTGAGCCCCGGAGGCCAGGGCGTCCGGGGCTGCGCCACTTCCGAGGGC  
CGAGCGCTGCCGGTCCC GGCGGTGCGACACGGCCGGGAGGAGGAACAACGCAAGGGGCTC  
AACCGTCGGTCGCTGGAGCCCCCCCCGGGGCGTGGCCTCCCGCCCCCTCAGCTGGGGAGGGC  
GGGGCTCGCTGCCCCCTGCTGCCGACTGCGACCCTTACAGGGGAGGGAGGGCGCAGGCCGCG  
CGGAGATGAGGAGGAGGCTGCGCCTACGCAGGGACGCATTGCTCACGCTGCTCCTTGGCGCC  
TCCCTGGGCTCTTACTCTATGCGCAGCGCGACGGCGCGGCCCCGACGGCGAGCGCGCCGCG  
AGGGCGAGGGAGGGCGGCACCGAGGCCACCCCCGACCCCGCGCGTTCAGTTACCCGACG  
CGGGTGCAGCCCCGCCGGCCTACGAAGGGGACACACCGGGCGCCGCCACGCCTACGGGACCC  
TTTGACTTCGCCCGCTATTTGCGCGCAAGGACCAGCGGGCGGTTTCCACTGCTCATTAAACCA  
GCCGCACAAGTGCCGCGGCGACGGCGCACCCGGTGGCCGCCCGGACCTGCTTATTGCTGTCA  
AGTCGGTGGCAGAGGACTTCGAGCGGCGCCAAGCCGTGCGCCAGACGTGGGGCGCGGAGGGT  
CGCGTGCAGGGGGCGCTGGTGCGCCGCGTGTCTTGGTGGGCGTGCCAGGGGGCGCAGGCTC  
GGGCGGGGCCGACGAAGTTGGGGAGGGCGCGCGAACCCTGCGCGCCCTGCTGCGGGCCG  
AGAGCCTTGCGTATGCGGACATCTGCTCTGGGCCTTCGACGACACCTTTTTTAACTAACG  
CTCAAGGAGATCCACTTTCTAGCCTGGGCCTCAGCTTTCTGCCCCGACGTGCGCTTCGTTTT  
TAAGGGCGACGCAGATGTGTTGTAACGTGGGAAATCTCCTGGAGTTCCTGGCGCCGCGGGAC  
CCGGCGCAAGACCTGCTTGGTGGTACGTAATTGTGCATGCGCGGCCCATCCGCACGCGGGC  
TAGCAAGTACTACATCCCCGAGGCCGTGTACGGCCTGCCCGCCTATCCGGCCTACGCGGGCG  
GCGGTGGCTTTGTGCTTTCCGGGGCCACGCTGCACCCGCTGGCTGGCGCCTGTGCGCAGGTC  
GAGCTCTTCCCCATCGACGACGTCTTTCTGGGCATGTGTCTGCAGCGCCTGCGGCTCACGCC  
CGAGCCTCACCTGCCTTCGACACCTTTGGCATCCCCAGCCTTCAGCCGCGCCGCATTTGA  
GCACCTTCGACCCCTGCTTTTACCGTGAGCTGGTTGTAGTGCACGGGCTCTCGGCCGCTGAC  
ATCTGGCTTATGTGGCGCCTGCTGCACGGGCCGCATGGGCCAGCCTGTGCGCATCCACAGCC  
TGTGCTGCAGGCCCTTCCAATGGGACTCCTAGCTCCCCACTACAGCCCCAAGCTCCTAAC  
TCAGACCCAGAATGGAGCCGTTTCCAGATTATTGCCGTGTATGTGGTTCTTCCCTGATCA  
CCAGGTGCCTGTCTCCACAGGATCCCAGGGGATGGGGGTTAAGCTTGGCTCCTGGCGGTCCA  
CCCTGCTGGAACCAAGTTGAAACCCGTGTAATGGTGAACCTTTGAGCGAGCCAAGGCTGGGTG  
GTAGATGACCATCTCTTGTCCAACAGGTCCCAGAGCAGTGGATATGTCTGGTCTCTAGTA  
GCACAGAGGTGTGTTCTGGTGTGGTGGCAGGGACTTAGGGAATCCTACCCTCTGCTGGATT  
TGGAACCCCTAGGCTGACGCGGACGTATGCAGAGGCTCTCAAGGCCAGGCCCCACAGGGAG  
GTGGAGGGGCTCCGGCCGCCACAGCTGAATTCATGAACCTGGCAGGCACTTTGCCATAGCT  
CATCTGAAAACAGATATTATGCTTCCACAACCTCTCCTGGGCCAGGTGTGGCTGAGCACC  
AGGGATGGAGCCACACATAAGGGACAAATGAGTGCACGGTCTTACCTAGTCTTTCCCTCACCT  
CTGAACTCACACAACAATGCCAGTCTCCCACTGGAGGCTGTATCCCCTCAGAGGAGCCAAG  
GAATGTCTTCCCCTGAGATGCCACCACTATTAATTTCCCATATGCTTCAACCACCCCTTG  
CTCAAAAACCAATACCCACACTTACCTTAATACAAACATCCCAGCAACAGCACATGGCAGG  
CCATTGCTGAGGGCACAGGTGCTTTATTGGAGAGGGGATGTGGGCAGGGGATAAGGAAGGTTCC  
CCATTCCAGGAGGATGGGAACAGTCTGGCTGCCCTGACAGTGGGGATATGCAAGGGGCT  
CTGGCCAGGCCACAGTCCAAATGGGAAGACACCAGTCAGTCAAAAAGTCGGGAGCGCCACA  
CAAACCTGGCTATAAGGCCCAGGAACCATATAGGAGCCTGAGACAGGTCCCCTGCACATTCA  
TCATTAACCTATACAGGATGAGGCTGTACATGAGTTAATTACAAAAGAGTCATATTTACAAA  
AATCTGTACACACATTTGAAAACTCACAAAATGTGCATCTATGTATCACAAGTTGCTAGAC  
CCAAAATATTA AAAATGGGATAAAAATTNNTTTAAAAA AAAAAAAAAAAAAAAAAAAAAA  
AAAAAAAAAAAAA

## **FIGURE 34**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA83505
><subunit 1 of 1, 402 aa, 1 stop
><MW: 43751, pI: 9.42, NX(S/T): 1
MRRRLRLRRDALLTLLLGASLGLLLYAQRDGAAPTASAPRGRGRAAPRPTPGPRAFQLPD
AGAAPPAYEGDTPAPPTPTGPFDFARYLRAKDQRRFPLLINQPHKCRGDGAPGGRPDLLI
AVKSVAEDFERRQAVRQTWGAEGRVQGALVRRVFLLGVPRGAGSGGADEVGEGARTHWR
LLRAESLAYADILLWAFDDTFFNLTLKEIHFLAWASAFCPDVRVFKGDADVFVNVGNLL
EFLAPRDPAQDLLAGDVIVHARPIRTRASKYYIPEAVYGLPAYPAYAGGGGFVLSGATLH
RLAGACAQVELFPIDDVFLGMCLQRLRLTPEPHPAFRFTFGIPQPSAAPHLSTFDPCFYRE
LVVVHGLSAADIWLMWRLLHGPHGPACAHQPVAAGPFQWDS
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-27

**N-glycosylation site:**

Amino acids 203-207

**N-myristoylation sites:**

Amino acids 18-24;31-37;110-116;157-163;161-167  
163-169;366-372

**Cell attachment sequence:**

Amino acids 107-110

## FIGURE 35

AGCAGCCTCTGCCCCACCCGGCTCGTGCGGACCCCAGGACCGGGCGCGGGACGCGTGCGTCC  
AGCCTCCGGCGCTGCGGAGACCCGCGGCTGGGTCCGGGGAGGCCCCAAACCCGCCCCCGCCA  
GAACCCCGCCCCAAATTCCCACCTCCTCCAGAAGCCCCGCCACTCCCAGCCCCGAGAGCT  
CCGCGCACCTGGGCGCCATCCGCCCTGGCTCCGCTGCACGAGCTCCACGCCCGTACCCCGGC  
GTCACGCTCAGCCCCGCGGTGCTCGCACACCTGAGACTCATCTCGCTTCGACCCCGCCGCGC  
CGCCGCCCGGCATCCTGAGCACGGAGACAGTCTCCAGCTGCCGTTCATGCCTTCCCCAGC  
CTTCCGCAGCCCACCAGGGAAGGGGCGGTAGGAGTGGCCTTTTACCAAAGGGACCGGCGATG  
CTCTGCAGGCTGTGCTGGCTGGTCTCGTACAGCTTGGCTGTGCTGTTGCTCGGCTGCCTGCT  
CTTCTGAGGAAGGCGGCCAAGCCCGCAGGAGACCCACGGCCCACCAGCCTTTCTGGGCTCCC  
CAAACACCCCGTCACAGCCGGTGTCCACCCAACCACACAGTGTCTAGCGCCTCTCTGTCCCT  
GCCTAGCCGTCACCGTCTCTTCTTGACCTATCGTCACTGCCGAAATTTCTCTATCTTGCTGG  
AGCCTTCAGGCTGTTCCAAGGATACCTTCTTGCTCCTGGCCATCAAGTACAGCCTGGTCAC  
GTGGAGCGACGTGCGGCTATCCGCAGCACGTGGGGCAGGGTGGGGGGATGGGCTAGGGGCCG  
GCAGCTGAAGCTGGTGTTCCTCCTAGGGGTGGCAGGATCCGCTCCCCAGCCCAGCTGCTGG  
CCTATGAGAGTAGGGAGTTTGATGACATCCTCCAGTGGGACTTCACTGAGGACTTCTTCAAC  
CTGACGCTCAAGGAGCTGCACCTGCAGCGCTGGGTGGTGGCTGCCTGCCCCAGGCCCATTT  
CATGCTAAAGGGAGATGACGATGTCTTTGTCCACGTCCCCAACGTGTTAGAGTTCCTGGATG  
GCTGGGACCCAGCCCAGGACCTCCTGGTGGGAGATGTCATCCGCCAAGCCCTGCCCAACAGG  
AACACTAAGGTCAAATACTTCATCCCACCCCTCAATGTACAGGGCCACCCACTACCCACCCTA  
TGCTGGTGGGGGAGGATATGTCATGTCCAGAGCCACAGTGCGGCGCCTCCAGGCTATCATGG  
AAGATGCTGAACTCTTCCCATTGATGATGTCTTTGTGGGTATGTGCCTGAGGAGGCTGGGG  
CTGAGCCCTATGCACCATGCTGGCTTCAAGACATTTGGAATCCGGCGGCCCCCTGGACCCCTT  
AGACCCCTGCCTGTATAGGGGGCTCCTGCTGGTTCACCGCCTCAGCCCCCTCGAGATGTGGA  
CCATGTGGGCACTGGTGACAGATGAGGGGCTCAAGTGTGCAGCTGGCCCCATACCCAGCGC  
TGAAGGGTGGGTTGGGCAACAGCCTGAGAGTGGACTCAGTGTGATTCTCTATCGTGATGCG  
AAATTGATGCCTGCTGCTCTACAGAAAATGCCAACTTGGTTTTTTAACTCCTCTCACCCCTGT  
TAGCTCTGATTA AAAACACTGCAACCCAA



## **FIGURE 36**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA84927
><subunit 1 of 1, 378 aa, 1 stop
><MW: 42310, pI: 9.58, NX(S/T): 3
MLPPQPSAAHQGRGGRSGLLPKGPAMLCRLCWLVSYSLAVLLLGCLLFLRKA AKPAGDPT
AHQPFWAPPTPRHSRCPNHTVSSASLSLPSRHRLFLTYRHCNFSILLEPSGCSKDTFL
LLAIKSQPGHVERRAAIRSTWGRVGGWARGRQLKLVFLLGVAGSAPPAQLLAYESREFDD
ILQWDFTEDEFFNLTLKELHLQRWVVAACPQAHFMLKGDDDDVFVHVPNVLEFLDGWDPAQD
LLVGDVIRQALPNRNTKVKYFIPPSMYRATHYPPYAGGGGYVMSRATVRRLLQAIMEDAEL
FPIDDFVFGMCLRRLGLSPMHHAGFKTFGIRRPLDPLDPCLYRGLLLVHRLSPLEMWTMW
ALVTDEGLKCAAGPIPQR
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-39

**Transmembrane domain:**

Amino acids 146-171

**N-glycosylation sites:**

Amino acids 79-83;104-108;192-196

**N-myristoylation sites:**

Amino acids 14-20;160-166;367-373

**Prokaryotic membrane lipoprotein lipid attachment site:**

Amino acids 35-46

**FIGURE 37**

ATGAAAGTGATAATCAGGCAGCCCAAATGATTGTTAATAAGGATCAAATGAGATCGTGTATG  
TGGGTCCAATCAATTGATTCTACACAAAGGAGCCTGGGGAGGGGCC**ATGGT**GCCAAATGCACT  
TACTGGGGAGACTGGAGAAGCCGCTTCTCCTCCTGTGCTGCGCCTCCTTCCTACTGGGGCTG  
GCTTTGCTGGGCATAAAGACGGACATCACCCCGTTGCTTATTTCTTTCTCACATTGGGTGG  
CTTCTTCTTGTGTTTGCCTATCTCCTGGTCCGGTTTCTGGAATGGGGGCTTCGGTCCCAGCTCC  
AATCAATGCAGACTGAGAGCCCAGGGCCCTCAGGCAATGCACGGGACAATGAAGCCTTTGAA  
GTGCCAGTCTATGAAGAGGCCGTGGTGGGACTAGAATCCCAGTGCCGCCCCAAGAGTTGGA  
CCAACCACCCCTACAGCACTGTTGTGATACCCCGAGCACCTGAGGAGGAACAACCTAGCC  
ATCCAGAGGGGTCCAGGAGAGCCAAACTGGAACAGAGGCGAATGGCCTCAGAGGGGTCCATG  
GCCCAGGAAGGAAGCCCTGGAAGAGCTCCAATCAACCTTCGGCTTCGGGGACCACGGGCTGT  
GTCCACTGCTCCTGATCTGCAGAGCTTGGCGGCAGTCCCCACATTAGAGCCTCTGACTCCAC  
CCCCTGCCTATGATGTCTGCTTTGGTCACCCTGATGATGATAGTGTTTTTTATGAGGACAAC  
TGGGCACCCCT**TAA**ATGACTCTCCAAGATTTCTTCTTCTCCACACCAGACCTCGTTCAT  
TTGACTAACATTTTCCAGCGCCTACTATGTGTCAGAAACAAGTGTTCCTGCCTGGACATCAT  
AAATGGGGACTTGGACCCTGAGGAGAGTCAGGCCACGGTAAGCCCTTCCCAGCTGAGATATG  
GGTGGCATAAATTTGAGTCTTCTGGCAACATTTGGTGACCTACCCCATATCCAATATTTCCAG  
CGTTAGATTGAGGATGAGGTAGGGAGGTGATCCAGAGAAGGCGGAGAAGGAAGAAGTAACCT  
CTGAGTGGCGGCTATTGCTTCTGTTCCAGGTGCTGTTCCAGCTGTTAGAACCCTTAGGCTTGAC  
AGCTTTGTGAGTTATTATTGAAAATGAGGATTCCAAGAGTCAGAGGAGTTTGATAATGTGC  
ACGAGGGCACACTGCTAGTAAATAACATTAAAATAACTGGAATGAA

## FIGURE 38

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA92264
><subunit 1 of 1, 216 aa, 1 stop
><MW: 23729, pI: 4.73, NX(S/T): 0
MVPMHLLGRLEKPLLLLCCASFLLGLALLGIKTDITPVAYFFLTLGGFFLFAYLLVRFLE
WGLRSQLQSMQTESPGPSGNARDNEAFEVVPVYEEAVVGLESQCRPQELDQPPPYSTVVIP
PAPEEEQPSHPEGSRRAKLEQRRMASEGSM AQEGSPGRAPINLRLRGPRAVSTAPDLQSL
AAVPTLEPLTPPPAYDVCFGHPDDDSV FYEDNWAPP
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-25

**Transmembrane domain:**

Amino acids 41-59

**N-myristoylation site:**

Amino acids 133-139



## FIGURE 40

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA94713
><subunit 1 of 1, 547 aa, 1 stop
><MW: 61005, pI: 6.34, NX(S/T): 2
MPSEVARGKRAALFFAAVAIVLGLPLWWTETETYRASLPYSQISGLNALQLRLMVPVTVV
FTRESVPLDDQEKLPFTVVHEREIPLKYKMKIKCRFQKAYRRALDHEEEALSSGSVQEAE
AMLDEPQEQAEGLTVYVISEHSSLLPQDMMSYIGPKRTAVVVRGIMHREAFNIIGRRIVQ
VAQAMSLTEDVLAALADHLPEDKWSAEKRRPLKSSLGYEITFSLNPDPKSHDVYWDIE
GAVRRYVQPFLNALGAAGNFSVDSQILYYAMLGVNPRFDSASSSYLDMHSLPHVINPVE
SRLGSSAASLYPVLNFLLYPELAHSPLYIQDKDGAPVATNAFHSPRWGGIMVYNVDSKT
YNASVLPVRVEVDMVRVMEVFLAQLRLLFGIAQPQLPPKCLLSGPTSEGLMTWELDRLLW
ARSVENLATATTTTSLAQLLGKISNIVIKDDVASEVYKAVAAVQKSAEELASGHLASAF
VASQEAVTSSELAFFDPSLLHLLYFPDDQKFAIYIPLFLPMAVPILLSLVKIFLETRKSW
RKPEKTD
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-23

**Transmembrane domain:**

Amino acids 511-530

**N-glycosylation sites:**

Amino acids 259-263;362-366

**N-myristoylation sites:**

Amino acids 255-261;304-310;335-341

**Amidation sites:**

Amino acids 7-11;174-178

## FIGURE 41

CCAGCTGCAGAGAGGAGGAGGTGAGCTGCAGAGAAGAGGAGGTTGGTGTGGAGCACAGGCAG  
CACCGAGCCTGCCCCGTGAGCTGAGGGCCTGCAGTCTGCGGCTGGAATCAGGATAGACACCA  
AGGCAGGACCCCCAGAGATGCTGAAGCCTCTTTGGAAAGCAGCAGTGGCCCCACATGGCCA  
TGCTCCATGCCGCCCGCCGCGCCGTGGGACAGAGAGGCTGGCACGTTGCAGGTCTCTGGGAGC  
GCTGGCTGTGCTGTGGCTGGGCTCCGTGGCTCTTATCTGCCTCCTGTGGCAAGTGCCCCGTCT  
CCCACCTGGGGCCAGGTGCAGCCCAAGGACGTGCCCAGGTCTGGGAGCATGGCTCCAGCCC  
AGCTTGGGAGCCCCCTGGAAGCAGAGGCCAGGCAGCAGAGGGACTCCTGCCAGCTTGTCTTGG  
TGGAAAGCATCCCCAGGACCTGCCATCTGCAGCCGGCAGCCCCCTCTGCCAGCCTCTGGGC  
CAGGCCTGGCTGCAGCTGCTGGACACTGCCCAGGAGAGCGTCCACGTGGCTTCATACTACTG  
GTCCCTCACAGGGCCTGACATCGGGGTCAACGACTCGTCTTCCCAGCTGGGAGAGGCTCTTC  
TGCAGAAGCTGCAGCAGCTGCTGGGCAGGAACATTTCCCTGGCTGTGGCCACCAGCAGCCCG  
AACTGGCCAGGACATCCACCGACCTGCAGGTTCTGGCTGCCCGAGGTGCCCATGTACGACA  
GGTGCCCATGGGGCGGCTCACCGGGGTGTTTTGCACTCCAAATTCTGGGTTGTGGATGGAC  
GGCACATATACATGGGCAGTGCCAACATGGACTGGCGGTCTCTGACGCAGGTGAAGGAGCTT  
GGCGCTGTCATCTATAACTGCAGCCACCTGGCCCAAGACCTGGAGAAGACCTTCCAGACCTA  
CTGGGTACTGGGGGTGCCCAAGGCTGTCTCCCCAAAACCTGGCCTCAGAATTCTCATCTC  
ACTTCAACCGTTTTCCAGCCCTTCCACGGCCTCTTTGATGGGGTGCCACCCTGCCTACTTC  
TCAGCGTCGCCACCAGCACTCTGTCCCCAGGGCCGCACCCGGGACCTGGAGGCGCTGCTGGC  
GGTGATGGGGAGCGCCAGGAGTTCATCTATGCCTCCGTGATGGAGTATTTCCCCACCACGC  
GCTTCAGCCACCCCCGAGGTACTGGCCGGTGTGGACAACGCGCTGCGGGCGGCAGCCTTC  
GGCAAGGGCGTGC GCGTGC GCTGGTGGCTGCGGACTCAACACGGACCCACCATGTT  
CCCCTACCTGCGGTCCCTGCAGGCGCTCAGCAACCCCGCGGCCAACGTCTCTGTGGACGTGA  
AAGTCTTCATCGTGCCGGTGGGGAACCATTCCAACATCCCATTCAGCAGGGTGAACCACAGC  
AAGTTCATGGTCACGGAGAAGGCAGCCTACATAGGCACCTCCAACTGGTCGGAGGATTACTT  
CAGCAGCACGGCGGGGGTGGGCTTGGTGGTCAACCAGAGCCCTGGCGCGCAGCCCCGCGGGG  
CCACGGTGCAGGAGCAGCTGCGGCAGCTCTTTGAGCGGGACTGGAGTTCGCGCTACGCCGTC  
GGCCTGGACGGACAGGCTCCGGGCCAGGACTGCGTGTGGCAGGGCTTGAGGGGGGCCTCTTTT  
TCTCTCGGCGACCCCGCCCCGCACGCGCCCTCCCCTCTGACCCCGGCCTGGGCTTCAGCCGC  
TTCCTCCCGCAAGCAGCCGGGTCCGCACTGCGCCAGGAGCCGCTGCGACCGCCCGGGCGT  
CGAAACCGCCCGCTGCTCTCTGATTTCCGAGTCCAGCCCCCCTGAGCCCCACCTCCTCC  
AGGGAGCCCTCCAGGAAGCCCC'TCCCTGACTCCTGGCCCACAGGCCAGGCCTAAAAAAAC  
TCGTGGCTTCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

## FIGURE 42

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA96869
><subunit 1 of 1, 489 aa, 1 stop
><MW: 53745, pI: 8.36, NX(S/T): 8
MPPRRPWDREAGTLQVLGALAVLWLG SVALICLLWQVPRPPTWGQVQPKDVPRSWEHGSS
PAWEPLAEARQQRDSCQLVLVESIPQDLPSAAGSPSAQPLGQAWLQLLDTAQESVHVAS
YYWSLTGPDIGVNDSSSQLGEALLQKLQQLGRNISLAVATSSPTLARTSTD LQVLAARG
AHVRQVPMGRLTRGVLHSKFWVVDGRHIYMG SANMDWRSLTQVKELGAVIYNCSHLAQDL
EKTFFQTYWVLGVPKAVLPKTWPQNFSSHFNRFQPFHGLFDGVP TTAYFSASPPALCPQGR
TRDLEALLAVMGSQAQEFIYASVMEYFPTTRESHPPRYWPVLDNALRAAAF GKGVRVRLLV
GCGLNTDPTMFPYLRSLQALS NPAANVSVDVKVFI VPGNHSNIPFSRVNHSKFMVTEKA
AYIGTSNWSEDYFSSTAGVGLVVTQSPGAQPAGATVQEQLRQLFERDWSSRYAVGLDGQA
PGQDCVWQG
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-29

**N-glycosylation sites:**

Amino acids 133-137;154-158;232-236;264-268;  
386-390;400-404;410-414;427-431

**N-myristoylation sites:**

Amino acids 58-64;94-100;131-137;194-200;251-257;  
277-283;281-287;361-367;399-405;  
440-446;448-454;478-484

## **FIGURE 43**

GGGCCTGGCGATCCGGATCCCGCAGGCGCGCTGGCTGCGCTGCCCGGCTGTCTGTCGTC**ATG**  
GTGGGGCCCTGGGTGTATCTGGTGGCGGCAGTTTTGCTCATCGGCCTGATCCTCTTCCTGAC  
TCGCAGCCGGGGTCGGGCGGCAGCAGCTGACGGAGAACCCTGCACAATGAGGAAGAGAGGG  
CAGGAGCAGGCCAGGTAGGCCGCTCTTTGCCCCAGGAGTCTGAAGAACAGAGAACTGGAAGC  
AGACCCCGGCGTCCGAGGGACTTGGGCAGCCGTCTACAGGCCCAGCGTCGAGCCCAGCGAGT  
GGCCTGGGAAGACGGGGATGAGAATGTGGGTCAAACGTATTATCCAGCCCAGGAGGAAGAAG  
GCATTGAGAAGCCAGCAGAAGTTCACCCAACAGGGAAAATTGGAGCCAAGAACTACGGAAG  
CTAGAGGAAAAACAGGCTCGAAAGGCTCAGCGAGAGGCAGAGGAGGCTGAACGTGAAGAACG  
GAAACGCCTAGAGTCCCAACGTGAGGCCGAATGGAAGAAGGAAGAGGAACGGCTTCGCCTGA  
AGGAAGAACAGAAGGAGGAGGAAGAGAGGAAGGCTCAGGAGGAGCAGGCCCGGCGGGATCAC  
GAGGAGTACCTGAAACTGAAGGAGGCCTTCGTGGTAGAAGAAGAAGGTGTTAGCGAAACCAT  
GACTGAGGAGCAGTCTCACAGCTTCCTGACAGAATTCATCAATTACATCAAGAAGTCCAAGG  
TTGTGCTTTTGGAAAGATCTGGCTTTCAGATGGGCCTAAGGACTCAGGACGCCATAAACCGC  
ATCCAGGACCTGCTGACGGAGGGGACTCTAACAGGTGTGATTGACGACCGGGGCAAGTTTAT  
CTACATAACCCCAGAGGAACTGGCTGCCGTGGCCAATTCATCCGACAGCGGGGCCGGGTGT  
CCATCACAGAGCTTGCCCAGGCCAGCAACTCCCTCATCTCCTGGGGCCAGGACCTCCCTGCC  
CAGGCTTCAGCCT**TGA**CTCCAGTCCTTCCTTGAGTGTATCCTGTGGCCTACATGTGTCTTCAT  
CCTTCCCTAATGCCGTCTTGGGGCAGGGATGGAATATGACCAGAAAGTTGTGGATTAAGGC  
CTGTGAATACTGAA



## **FIGURE 44**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA96881
><subunit 1 of 1, 315 aa, 1 stop
><MW: 35963, pI: 5.38, NX(S/T): 0
MVG PWVYLVA AVLLIGLILFLTRSRG RAAAADG EPLHNEE ERAGAGQVGRSLPQESEEQR
TGSRRRRRDLG SRLQAQRRAQRV AWEDGDENVGQTVIPAQEEEGIEKPAEVHPTGKIGA
KKLRKLEEKQARKAQREAEAAEREERKRLESQREAEWKKEEERLRLKEEQKEEERKAQE
EQARRDHEEYLKLKEAFVVEEEGVSETMTEEQSHSFLTEFINYIKKSKVVLEDLAFQMG
LRTQDAINRIQDLLTEGTLTGVIDDRGKFIYITPEELAAVANFIRQRGRVSI TELAQASN
SLISWGQDLPAQASA
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-26

**N-myristoylation sites:**

Amino acids 203-209;257-263

**FIGURE 45**

ACGGGCCGCAGCGGCAGTGACGTAGGGTTGGCGCACGGATCCGTTGCGGCTGCAGCTCTGCA  
GTCGGGCCGTTTCCTTCGCCGCCGCCAGGGGTAGCGGTGTAGCTGCGCAGCGTCGCGCGCGCT  
ACCGCACCCAGGTTCCGGCCCGTAGGCGTCTGGCAGCCC GGCGCCATCTTCATCGAGCGCCAT  
GGCCGCAGCCTGCGGGCCGGGAGCGGCCGGGTACTGCTTGCTCCTCGGCTTGCATTTGTTTC  
TGCTGACCGCGGGCCCTGCCCTGGGCTGGAACGACCCTGACAGAATGTTGCTGCGGGATGTA  
AAAGCTCTTACCCTCCACTATGACCGCTATAACCACCTCCCGCAGGCTGGATCCCATCCCACA  
GTTGAAATGTGTTGGAGGCACAGCTGGTTGTGATTCTTATACCCCAAAGTCATACAGTGTC  
AGAACAAAGGCTGGGATGGGTATGATGTACAGTGGGAATGTAAGACGGACTTAGATATTGCA  
TACAAATTTGGAAAACTGTGGTGAGCTGTGAAGGCTATGAGTCCTCTGAAGACCAGTATGT  
ACTAAGAGGTTCTTGTGGCTTGGAGTATAATTTAGATTATACAGAACTTGGCCTGCAGAAAC  
TGAAGGAGTCTGGAAGCAGCACGGCTTTGCCTCTTCTCTGATTATTATTATAAGTGGTCC  
TCGGCGGATTCTGTAAACATGAGTGGATTGATTACCATCGTGGTACTCCTTGGGATCGCCTT  
TGTAGTCTATAAGCTGTTCTTGAGTGACGGGCAGTATTCTCCTCCACCGTACTCTGAGTATC  
CTCCATTTTCCCACCGTTACCAGAGATTACCAACTCAGCAGGACCTCCTCCCCCAGGCTTT  
AAGTCTGAGTTCACAGGACCACAGAATACTGGCCATGGTGCAACTTCTGGTTTTGGCAGTGC  
TTTTACAGGACAACAAGGATATGAAAATTCAGGACCAGGGTTCTGGACAGGCTTGGGAACTG  
GTGGAATACTAGGATATTTGTTTGGCAGCAATAGAGCGGCAACACCCTTCTCAGACTCGTGG  
TACTACCCGTCTATCCTCCCTCCTACCCTGGCACGTGGAATAGGGCTTACTCACCCCTTCA  
TGGAGGCTCGGGCAGCTATTCGGTATGTTCAAACCTCAGACACGAAAACCAGAACTGCATCAG  
GATATGGTGGTACCAGGAGACGTAAAGTAGAAAAGTTGGAGTCAAACACTGGATGCAGAAAT  
TTTGGATTTTTTCATCACTTTCTCTTTAGAAAAAAAGTACTACCTGTTAACAATTGGGAAAAG  
GGGATATTCAAAGTTCTGTGGTGTATGTCCAGTGTAGCTTTTTGTTATTCTATTATTTGAG  
GCTAAAAGTTGATGTGTGACAAAATACTTATGTGTTGTATGTCAGTGTAAACATGCAGATGTA  
TATTGCAGTTTTTTGAAAGTGATCATTACTGTGGAATGCTAAAAATACATTAATTTCTAAAAC  
CTGTGATGCCCTAAGAAGCATTAAGAATGAAGGTGTTGTAATAAGAACTAAGTACAGAA  
AATTTTCAGTTTTTAGGTGGTGTAGCTGATGAGTTATTACCTCATAGAGACTATAATATTCTA  
TTTGGTATTATATTATTTGATGTTTGGCTGTTCTTCAAACATTTAAATCAAGCTTTGGACTAA  
TTATGCTAATTTGTGAGTTCTGATCACTTTTGGCTCTGAAGCTTTGAATCATTGAGTGGTG  
GAGATGGCCTTCTGGTAACTGAATATTACCTTCTGTAGGAAAAGGTGGAAAATAAGCATCTA  
GAAGGTTGTTGTGAATGACTCTGTGCTGGCAAAAATGCTTGAAACCTCTATATTTCTTTTCGT  
TCATAAGAGGTAAAGGTCAAATTTTTCAACAAAAGTCTTTTAATAACAAAAGCATGCAGTTCTC  
TGTGAAATCTCAAATATTGTTGTAATAGTCTGTTTCAATCTTAAAAAGAATCA

## FIGURE 46

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA96889
><subunit 1 of 1, 339 aa, 1 stop
><MW: 36975, pI: 7.85, NX(S/T): 1
MAAACGPGAAGYCLLLGLHLFLLTAGPALGWNDPDRMLLRDVKALTLHYDRYTTSRRLDP
IPQLKCVGGTAGCDSYTPKVIQCQNKGWGDYDVQWECKTDLDIAYKFGKTVVSCEGYESS
EDQYVLRGSCGLEYNLDYTELGLQKLKESGKQHGFAFSFDYKSSADSCNMSGLITIV
VLLGIAFVVYKLFSLDQYSPPPYSEYPPFSHRYQRFTNSAGPPPPGFKSEFTGPQNTGH
GATSGFGSAFTGQQGYENSGPGFWTGLGTGGILGYLFGSNRAATPFSDSWYYPSYPPSY
GTWNRAYSPLHGSGSYSVCSNSDTKTRTASGYGGTRRR
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-30

**Transmembrane domain:**

Amino acids 171-190

**N-glycosylation site:**

Amino acids 172-176

**Glycosaminoglycan attachment sites:**

Amino acids 244-248;259-263;331-335

**Tyrosine kinase phosphorylation site:**

Amino acids 98-106

**N-myristoylation sites:**

Amino acids 68-74;69-75;131-137;241-247;  
247-253;266-272;270-276;278-284;  
312-318

## FIGURE 47

CCCGGAGCCGGGGAGGGAGGGAGCGAGGTTCCGGACACCGGCGGGCGGCTGCCTGGCCTTTCCA  
**TG**AGCCCCGCGGCGGACCCCTCCC CGCCCCCTCTCGCTCTGCCTCTCCCTCTGCCTCTGCCTC  
TGCTTGCCCGCGGCTCTGGGAAGTGC GCAGTCCGGGTCGTGTAGGGATAAAAAGAACTGTAA  
GGTGGTCTTTTCCCAGCAGGAACTGAGGAAGCGGCTAACACCCCTGCAGTACCATGTCACTC  
AGGAGAAAGGGACCGAAAAGTGCCTTTGAAGGAGAATACACACATCACAAAGATCCTGGAATA  
TATAAATGTGTTGTTTGTGGAACCTCCATTGTTTAAAGTCAGAAACCAAATTTGACTCCGGTTC  
AGGTTGGCCTTCATTCCACGATGTGATCAATTTCTGAGGCAATCACATTCACAGATGACTTTT  
CCTATGGGATGCACAGGGTGGAAACAAGCTGCTCTCAGTGTGGTGTCTCACCTTGGGCACATT  
TTTGATGATGGGCCTCGTCCAACCTGGGAAAAGATACTGCATAAAATTCGGCTGCCTTGTCTTT  
TACACCTGCGGATAGCAGTGGCACCGCCGAGGGAGGCAGTGGGGTGCAGCCCGGCCAGG  
CAGACAAAGCGGAGCTC**TAG**AGTAATGGAGAGTGATGGAAACAAGTGTACTTAATGCACAG  
CTTATTAATAAAATCAAAATTTGTTATCTTAATAGATATATTTTTTCAA AAACTATAAGGGCA  
GTTTTGTGCTATTGATATTTTTTCTTCTTTTGCTTAAACAGAAGCCCTGGCCATCCATGTAT  
TTTGAATTGACTAGATCAAGAACTGTTTTATAGCTTTAGCAAATGGAGACAGCTTTGTGAAA  
CTTCTTCACAAGCCACTTATACCCCTTGGCATTCTTTTCTTTGAGCACATGGCTTCTTTTGC  
AGTTTTTCCCCCTTTGATTCAGAAGCAGAGGGTTCATGGTCTTCAAACATGAAAATAGAGAT  
CTCCTCTGCAGTGTAGAGACCAGAGCTGGGCAGTGCAGGGCATGGAGACCTGCAAGACACAT  
GGCCTTGAGGCCTTTGCACAGACCCACCTAAGATAAGGTTGGAGTGATGTTTTAATGAGACT  
GTTTCACTTTGTGGAAAGTTTGGAGCTAAGGTCATTTTTTTTTTTCTCACTGAAAGGGTGTGA  
AGGTCTAAAGTCTTTCCTTATGTTAAATTGTTGCCAGATCCAAAGGGGCATACTGAGTGTTG  
TGGCAGAGAAGTAAACATTACCACACTGTTAGGCCTTTATTTTTATTTTTATTTTCCATCGAAA  
GCATTGGAGGCCAGTGCAATGGCTCACGCCTGTGATCCCAGCACTTTGGGAGGCCAAGGCG  
GGTGGATCACGAGGTCAGGAGATGGAGACCATCCTGGCTAACATGGTGAAACCCCGTCTCTA  
CTAAAAATACGAAAAATTAGCCAGGCGTGGTGGTGGGCACCTGTAGTCCCAGCTACTCAGGAGG  
CTGAGGCAGGAGAATGGCGTGAACCCGGAAGGCGGAGCTTGCACTTAGCCGAGATCATGCCA  
CTGCACTCCAGCCTACATGACAATGTGACACTCCATCTCAAAAAATAATAATAATAACAATA  
TAAGA ACTAGCTGGGCATGGTGGCGCATGCATGTAGTCCCAGCTACTCCTGAGGCTCAGTCA  
GGAGAATCGCTTGAACCTGGGAGGCGGAGGTTGCAGTGAGCTGAGCTCATACCACTGCACTC  
CAGCCTGAACAGAGTGAGATCCTGTCAA

## FIGURE 48

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA96898
><subunit 1 of 1, 192 aa, 1 stop
><MW: 20702, pI: 7.50, NX(S/T): 0
MSPRRTLPRPLSLCLSLCLCLCLAAALGSAQSGSCRDKKNCKVVFSQQELRKRLTPLQYH
VTQEKGTESAFEGEYTHHKDPGIYKCVVCGTPLFKSETKFDSGSGWPSFHDVINSEAITF
TDDFSYGMHRVETSCSQCGAHLGHIFDDGPRPTGKRYCINSAALSFTPADSSGTAEGGSG
VASPAQADKAEL
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-24

**Glycosaminoglycan attachment site:**

Amino acids 102-106

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 52-56

**N-myristoylation sites:**

Amino acids 28-34;66-72;82-88;139-145;  
173-179;178-184

**Amidation site:**

Amino acids 153-157

## FIGURE 49

CCCAAAGAGGTGAGGAGCCGGCAGCGGGGGCGGCTGTAAGTGTGAGGAAGGCTGCAGAGTGG  
CGACGTCTACGCCGTAGGTTGGAGGCTGTGGGGGGTGGCCGGGCGCCAGCTCCCAGGCCGCA  
GAAGTGACCTGCGGTGGAGTTCCTCCTCGCTGCTGGAGAACGGAGGGAGAAGGTTGCTGGC  
CGGGTCAAAGTGCCCTCCCTCTGCTTGACGGGGCTGAGGGGCCCCAAGTCTAGGGCGTCCGTA  
GTCGCCCCGGCCTCCGTGAAGCCCCAGGTCTAGAGAT**ATG**ACCCGAGAGTGCCCATCTCCGG  
CCCCGGGGCTGGGGCTCCGCTGAGTGGATCGGTGCTGGCAGAGGCCGGCAGTAGTGTTTGCA  
GTGGTGTGAGCATCCACGCAACCGTATGGGACCGATACTCGTGGTGCGCCGTGGCCCTCGC  
AGTGCAGGCCCTTCTACGTCCAATACAAGTGGGACCGGCTGCTACAGCAGGGAAGCGCCGTCT  
TCCAGTTCCGAATGTCCGCAAACAGTGGCCTATTGCCCGCCTCCATGGTCATGCCTTTGCTT  
GGACTAGTCATGAAGGAGCGGTGCCAGACTGCTGGGAACCCGTTCTTTGAGCGTTTTGGCAT  
TGTGGTGGCAGCCACTGGCATGGCAGTGGCCCTCTTCTCATCAGTGTGGCGCTCGGCATCA  
CTCGCCAGTGCCAAACCAACTTGTGTCATCTTGGGCTTGGCTGGAGGTGTTATCATTAT  
ATCATGAAGCACTCGTTGAGCGTGGGGGAGGTGATCGAAGTCCTGGAAGTCCTTCTGATCTT  
CGTTTATCTCAACATGATCCTGCTGTACCTGCTGCCCCGCTGCTTACCCCTGGTGAGGCAC  
TGCTGGTATTGGGTGGCATTAGCTTTGTCTCAACCAGCTCATCAAGCGCTCTCTGACACTG  
GTGAAAGTCAGGGGGACCCAGTGGACTTCTTCTGCTGGTGGTGGTAGTAGGGATGGTACT  
CATGGGCATTTTCTTACACTCTGTTTGTCTTTCATGGACTCAGGCACCTGGGCCTCCTCCA  
TCTTCTCCACCTCATGACCTGTGTGCTGAGCCTGGTGTGGTCTTACCCTGGCTGCACCGG  
CTCATCCGCAGGAATCCCCTGCTCTGGCTTCTTACAGTTTCTTCTCCAGACAGACACCCGCAT  
CTACCTCCTAGCCTATTGGTCTCTGCTGGCCACCTTGGCCTGCCTGGTGGTGTGTACCAGA  
ATGCCAAGCGGTCATCTTCCGAGTCCAAGAAGCACCAGGCCCCACCATCGCCCGAAAGTAT  
TTCCACCTCATTGTGGTAGCCACCTACATCCCAGGTATCATCTTTGACCGGCCACTGCTCTAT  
GTAGCCGCCACTGTATGCCTGGCGGTCTTCATCTTCTTGGAGTATGTGCGCTACTTCCGCAT  
CAAGCCTTTGGGTACACTCTACGGAGCTTCTGTCCCTTTTTCTGGATGAACGAGACAGTG  
GACCACTCATTCTGACACACATCTACCTGCTCCTGGGCATGTCTCTTCCCATCTGGCTGATC  
CCCAGACCCTGCACACAGAAGGGTAGCCTGGGAGGAGCCAGGGCCCTCGTCCCCTATGCCGG  
TGTCCTGGCTGTGGGTGTGGGTGATACTGTGGCCTCCATCTTCGGTAGCACCATGGGGGAGA  
TCCGCTGGCCTGGAACCAAAAAGACTTTTGAGGGGACCATGACATCTATATTTGCGCAGATC  
ATTTCTGTAGCTCTGATCTTAATCTTTGACAGTGGAGTGGACCTAAACTACAGTTATGCTTG  
GATTTTGGGGTCCATCAGCACTGTGTCCCTCCTGGAAGCATACTACACAGATAGACAATC  
TCCTTCTGCCTCTCTACCTCCTGATATTGCTGATGGCCT**TAG**CTGTTACAGTGCAGCAGCAGT  
GACGGAGGAAACAGACATGGGGAGGGTGAACAGTCCCCACAGCAGACAGCTACTTGGGCATG  
AAGAGCCAAGGTGTGAAAAGCAGATTTGATTTTTAGTTGATTTCAGATTTAAAATAAAAAGC  
AAAGCTCTCCTAGTTCTA

## **FIGURE 50**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA97003
><subunit 1 of 1, 538 aa, 1 stop
><MW: 59268, pI: 8.94, NX(S/T): 1
MTRECPSPAPGPGAPLSGSQLAEAAVVFVAVLSIHAT'VWDRYSWCAVALAVQAFYVQYKW
DRLLQQGSVAVFQFRMSANSGLLPASVMPLLLGI.VMKERCQTAGNPFERFGIVVAATGMA
VALFSSVLALGITRVPPTNTCVILGLAGGVIIYIMKHSLSVGEVIEVLEVLLIFVYLNMI
LLYLLPRCFTPGEALLVLGGISFVLNQLIKRSLTLVESQGDVDFLLVVVVGMVLMGIF
FSTLFVFMDSGTWASSIFFHLMTCVLSLGVVLPWLHRLIRRNPLLWLLQFLEFQTDTRIYL
LAYWLLATLACLVVLYQNAKRSSSESKKHQAPTIAKYFHLIVVATYIPGIIFDRPLLY
VAATVCLAVFIFLEYVRYFRIKPLGHTLRSFSLFLDERDSGPLILTHIYLLLGMSLPIW
LIPRPTQKGSLLGARALVPYAGVLAVGVGDTVASIFGSTMGEIRWPGTKKTFEGTMTSI
FAQIISVALILIFDSGVDLNYSAWILGSISTVSLLEAYTTQIDNLLLPLYLLILLMA
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-36

**Transmembrane domains:**

Amino acids 77-95;111-133;161-184;225-248;  
255-273;299-314;348-373;406-421;  
435-456;480-497

**N-glycosylation sites:**

Amino acids 500-504

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 321-325

**N-myristoylation sites:**

Amino acids 13-19;18-24;80-86;111-117;  
118-124;145-151;238-244;251-257;  
430-436;433-439;448-454;458-464;  
468-474;475-481;496-502;508-514

**Prokaryotic membrane lipoprotein lipid attachment site:**

Amino acids 302-313

**FIGURE 51**

GCTCTATGCCGCTACCTTGCTCTCGCCGCTGCTGCCGGAGCCGAAGCAGAGAAGGCAGCGGGTCCCGTGACCG  
TCCCGAGAGCCCGCGCTCCCGACCAGGGGGCGGGGGCGGCCCCGGGGAGGGCGGGGCAGGGGCGGGGGAAGA  
AAGGGGGTTTTGTGCTGCGCCGGGAGGGCCGGCGCCCTCTTCCGAATGTCTGCGGCCCCAGCCTCTCCTCAGC  
CTCGCGCAGTCTCGCCCGCAGTCTCAGCTGCAGCTGCAGGACTGAGCCGTGCACCCGGAGGAGACCCCGGAGG  
AGGCGACAACTTCGCGAGTGCCGCGACCCAAACCCAGCCCTGGGTAGCCTGCAGCATGGCCAGCTGTTCCCTGC  
CCCTGCTGGCAGCCCTGGTCTGGCCCAGGCTCCTGCAGCTTTAGCAGATGTTCTGGAAGGAGACAGCTCAGAG  
GACCGCGCTTTTCGCGTGCGCATCGCGGGCGACGCGCCACTGCAGGGCCTGCTCGCGGGCGCCCTCACCATCCC  
TTGCCACGTCCACTACCTGCGGCCACCGCCGAGCCCGCGGCTGTGCTGGGCTCTCCGCGGGTCAAGTGGACTT  
TCCTGTCCCAGGGCCGGAGGCAGAGTGTGCTGGTGGCGCGGGGAGTGCAGCTCAAGGTGAACGAGGCCTACCGG  
TTCCGCGTGGCACTGCCTGCGTACCCAGCGTGCCTCACCGACGCTCCCTGGCGCTGAGCGAGCTGCGCCCCAA  
CGACTCAGGTATCTATCGCTGTGAGGTCCAGCACGGCATCGATGACAGCAGCGACGCTGTGGAGGTCAAGGTCA  
AAGGGTCTCTTTCTCTACCGAGAGGGCTCTGCCGCTATGCTTTCTCCTTTTCTGGGGCCAGGAGGCCTGT  
GCCCGCATGGAGCCCACATCGCCACCCCGGAGCAGCTCTATGCCGCTACCTTGGGGGCTATGAGCAATGTGA  
TGCTGGCTGGCTGTCCGATCAGACCGTGAAGTATCCCATCCAGACCCACGAGAGGCCTGTTACGGAGACATGG  
ATGGCTTCCCAGGGTCCGGAACATGGTGTGGTGGACCCGGATGACCTCTATGATGTGTACTGTTATGCTGAA  
GACCTAAATGGAGAAGTGTTCCTGGGTGACCCTCAGAGAAGCTGACATTGGAGGAAGCACGGGCGTACTGCCA  
GGAGCGGGGTGCAGAGATTGCCACCACGGGCCAACTGTATGCAGCCTGGGATGGTGGCCTGGACCCTGCAGCC  
CAGGGTGGCTAGCTGATGGCAGTGTGCGCTACCCCATCGTACACCCAGCCAGCGCTGTGGTGGGGGCTTGCCT  
GGTGTCAAGACTCTTTCCTCTTCCCCAACAGACTGGCTTCCCCAATAAGCACAGCCGCTTCAACGCTCTACTG  
CTTCCGAGACTCGGCCAGCCTTCTGCCATCCCTGAGGCCTCCAACCCAGCCTCCAACCCAGCCTGATGATGAG  
TAGAGCTATCCTCACAGTGACAGAGACCCTTGGAGGAAGTGCAGCTGCCTCAGGAAGCCACAGAGAGTGAATCC  
CGTGGGGCCATCTACTCCATCCCCATCATGGAGGACGGAGGAGGTGGAAGCTCCACTCCAGAAGACCCAGCAGA  
GGCCCCTAGGACGCTCCTAGAATTTGAAACACAATCCATGGTACCGCCCACGGGGTCTCAGAGAGGAAGGTA  
AGGCATTGGAGGAAGAAGAGAAATATGAAGATGAAGAAGAGAAAGAGGAGGAAGAAGAAGAGGAGGAGGTGGAG  
GATGAGGCTCTGTGGGCATGGCCCAGCGAGCTCAGCAGCCCGGGCCCTGAGGCCTCTCTCCCCTGAGCCAGC  
AGCCCAGGAGAAGTCACTCTCCAGGCGCCAGCAAGGGCAGTCTGCGAGCCTGGTGCATCACCCTTCCCTGATG  
GAGAGTCAGAAGCTTCCAGGCTCCAAGGGTCCATGGACCCTACTGAGACTCTGCCACTCCCAGGGAGAGG  
AACCTAGCATCCCCATCACCTTCCACTCTGGTTGAGGCAAGAGAGGTGGGGGAGGCAACTGGTGGTCTGAGCT  
ATCTGGGGTCCCTCGAGGAGAGAGCGAGGAGACAGGAAGCTCCGAGGGTGCCCTTCCCTGCTTCCAGCCACAC  
GGGCCCTGAGGGTACCAGGGAGCTGGAGGCCCTCTGAAGATAATTCTGGAGAAGTGCACCAGCAGGGACC  
TCAGTGCAGGCCAGCCAGTGTGCCCCTGACAGCGCCAGCCGAGGTGGAGTGGCCGTGGTCCCCGCATCAGG  
TGACTGTGTCCCAGCCCCCTGCCACAATGGTGGGACATGCTGGAGGAGGAGGAAGGGTCCGCTGCCATATGTC  
TGCCCTGGCTATGGGGGGGACCTGTGCGATGTTGGCCTCCGCTTCTGCAACCCCGGCTGGGACGCCTTCCAGGGC  
GCCTGCTACAAGCACTTTTCCACACGAAGGAGCTGGGAGGAGGCAGAGACCCAGTGCAGGATGTACGGCGCGCA  
TCTGGCCAGCATCAGCACACCCGAGGAACAGGACTTCATCAACAACCGGTACCGGGAGTACCAGTGGATCGGAC  
TCAACGACAGGACCATCGAAGGCGACTTCTTGTGGTCCGATGGCGTCCCCCTGCTCTATGAGAAGTGAACCCCT  
GGGAGCCTGACAGCTACTTCTGTCTGGAGAGAAGTGCCTGGTGCATGGTGTGGCATGATCAGGGACAATGGAG  
TGACGTGCCCTGCAACTACCCTGTCTACACCTGCAAGATGGGGCTGGTGTCTGTGGCCCGCCACCGGAGC  
TGCCCCCTGGCTCAAGTGTTCGGCCGCCACGGCTGCGCTATGAGGTGGACTGTGCTTCCGCTACCGGTGCCGG  
GAAGGACTGGCCCAGCGCAATCTGCCGCTGATCCGATGCCAAGAGAACGGTCTGTGGGAGGCCCCAGATCTC  
CTGTGTGCCAGAAGACCTGCCCGAGCTCTGCACCCAGAGGAGGCCAGAAGGACGTGAGGGGAGGCTACTGG  
GACGCTGGAAGGCGCTGTTGATCCCCCTTCCAGCCCCATGCCAGGTCCCTTAGGGGGCAAGGCCTTGAACACTGCCG  
GCCACAGCACTGCCCTGTACCCAAATTTTCCCTCACACCTTGCCTCCCGCCACCACAGGAAGTGACAACATG  
ACGAGGGGTGGTGTGGAGTCCAGTGCAGTCTCTGAAGGGGCTTCTGGGAAATACCTAGGAGGCTCCAGCC  
AGCCCAGGCCCTTCCCCCTACCCTGGGCACCAGATCTCCATCAGGGCCGGAGTAAATCCCTAAGTGCCTCAA  
CTGCCCTCTCCCTGGCAGCCATCTTGTCCCCTCTATTCTCTAGGGAGCACTGTGCCACTCTTTCTGGGTTTT  
CCAAGGGAATGGGCTTGCAGGATGGAGTGTCTGAAAATCAACAGGAAATAAACTGTGTATGAGCCCA



## **FIGURE 52**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA98565
><subunit 1 of 1, 911 aa, 1 stop
><MW: 99117, pI: 4.62, NX(S/T): 2
MAQLFLPLLAALVLAQAPAALADVLEGDSSSEDRAFRVRIAGDAPLQGVLLGGALTI PCHVH
YLRPPPSRRRAVLGSPRVKWTFLSRGREAEVLVARGVVRVKVNEAYRFRVALPAYPASLTDV
SLALSELRPNDSGIYRCEVQHGI DDSSDAVEVKVKGVVFLYREGSARYAFS FSGAQEACA
RIGAHIAITPEQLYAAYLGGYEQC DAGWLS DQTVRYPIQT PREACYGDMDGFP GVRNYGVV
DPDDLVDVYCYAEDLNGELFLGDPPEKLTLEEARAYCQERGAELATTGQLYAAWDGGLDH
CSPGWLADGSVRYPIVTPSQRCGGGLPGVKTLFLFPNQTGFPNKHSRFRNVYCFRDSAQPS
AIPASNPASNPDGLEAIVTVTETLEELQLPQEATESESRGAIYSIPIMEDGGGGSSST
PEDPAEAPRTLLEFETQSMVPPTGFSEEEGKALEEEEEKYEDEEEKEEEEEVEDEALW
AWPSELSSPGPEASLPTEPAAQEKSLSQAPARAVLQPGASPLPDGESEASRPPRVHGPPT
ETLPTPRERNLASPSPSTLVEAREVGEATGGPELSGVPRGESEETGSSEGAPSLLPATRA
PEGTRELEAPSEDNSGRAPAGTSVQAQPVLPDTSASRGGVAVVPASGDCVPSPCHNNGT
CLEEEEGVRCLCLPGYGGDLCDVGLRFCNPGWDAFQGACYKHFSTRRSWEEAETQCRMYG
AHLASISTPEEQDFINRYREYQWIGLNDRTIEGDFLWSDGVPLLYENWNPGQPD SYFLS
GENCVVMVWHQDQGWSDVPCNYHLSYTKMGLVSCGPPPELPLAQVFGFRPRLRYEVDTVL
RYRCREGLAQRNLP LIRCQENGRWEAPQISCVPRRPARALHPEEDPEGRQGRLLGRWKAL
LIPSSPMPGP
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-15

**N-glycosylation sites:**

Amino acids 130-134;337-341

**Tyrosine kinase phosphorylation sites:**

Amino acids 128-136;451-460

**N-myristoylation sites:**

Amino acids 47-53;50-56;133-139;142-148;  
174-180;183-189;281-287;288-294;  
297-303;324-330;403-409;414-420;  
415-421;576-582;586-592;677-683;  
684-690;720-726;772-778;811-817

**EGF-like domain cysteine pattern signature:**

Amino acids 670-682

**C-type lectin domain signature:**

Amino acids 784-809

**Immunoglobulins and major histocompatibility complex proteins signature:**

Amino acids 135-142

**Link domain proteins:**

Amino acids 166-216;264-314

**Calcium-binding EGF-like domain proteins pattern proteins.**

Amino acids 655-676

**C-type lectin domain proteins:**

Amino acids 791-800

### FIGURE 53

CTGCCAGGTGACAGCCGCAAGATGGGGTCTTGGGCCCTGCTGTGGCCCTCCCCTGCTGTTACCGGGCTGCTCG  
TCCGACCCCCGGGACCATGGCCCAGGCCAGTACTGCTCTGTGAACAAGGACATCTTTGAAGTAGAGGAGAAC  
ACAAATGTCACCGAGCCGCTGGTGACATCCACGTCCCGGAGGGCCAGGAGGTGACCCTCGGAGCCTTGTCCAC  
CCCCTTTCGATTTTCGGATCCAGGGAAACAGCTGTTTCTCAACGTGACTCCTGATTACGAGGAGAAGTCACTGC  
TTGAGGCTCAGTGTGTGTGAGAGCGGAGGCACATTGGTGAACCCAGCTAAGGGTGTTCGTGTGAGTGTGGAC  
GTCAATGACAATGCCCCGAATCCCCTTTAAGACCAAGGAGATAAGGGTGGAGGAGGACACGAAAGTGAACTC  
CACCGTATCCCTGAGACGCAACTGCAGGCTGAGGACCGGACAAGGACGACATTTCTGTTTACACCCCTCCAGG  
AAATGACAGCAGGTGCCAGTACTACTTCTCCCTGGTGAAGTGTAAACCGTCCCGCCCTGAGGCTGGACCGGCC  
CTGGACTTCTACGAGCGGCCGAACATGACCTTCTGGCTGTGGTGTGGGACACTCCAGGGGAGAATGTGGAACC  
CAGCCACACTGCCACCGCCACACTAGTGCTGAACGTGGTGCCTGCGGACCTGCGGCCCCCGTGGTTTCTGCCCT  
GCACCTTCTCAGATGGCTACGTCTGCATTCAAGCTCAGTACCACGGGGCTGTCCCACGGGGCACATACTGCCA  
TCTCCCCTCGTCTGCGTCCCGGACCCATCTACGCTGAGGACGGAGACCGCGGCATCAACCAGCCCATCATCTA  
CAGCATCTTTAGGGGAAACGTGAATGGTACATTCATCATCCACCCAGACTCGGGCAACCTCACCGTGGCCAGGA  
GTGTCCCCAGCCCCATGACCTTCTTCTGCTGGTGAAGGGCAACAGGCCGACCTTGCCCGCTACTCAGTGACC  
CAGGTACCGTGGAGGCTGTGGCTGCGGCCGGGAGCCCGCCCGCTTCCCCAGAGCCTGTATCGTGGCACCGT  
GGCGCTGGCGTGGAGCGGGCGTTGTGGTCAAGGATGCAGTGTGCCCTTCTCAGCCTCTGAGGATCCAGGCTC  
AGGACCCGGAGTCTCGGACCTCAACTCGGCCATCACATATCGAATTACCAACCACTCACACTTCCGGATGGAG  
GGAGAGGTTGTGTGACCACCACCACTGGCACAGGCGGGAGCCTTCTACGCAGAGGTTGAGGCCACACAACAC  
GGTGAACCTCTGGCACCGCAACCACAGTCAATTGAGATACAAGTTTCCGAACAGGAGCCCCCTCCACAGAGGCTG  
GAGGAACAACGGGCCCTGGACCAGCACCCTTCCGAGGTCCCAGACCCCTGAGCCCTCCCAGGGACCCCTCC  
ACGACCAGCTCTGGGGGAGGCACAGGCCCTCATCCACCCCTGGCACAACTCTGAGGGCCACCAACCTCGTCCAC  
ACCCGGGGGGCCCCGGGTGCAGAAAACAGCACCTCCACCAACCAGCCACTCCCGGTGGGGACACAGCACAGA  
CCCCAAAGCCAGGAACCTCTCAGCCGATGCCCCCGGTGTGGGAACCAGCACCTCCACCAACCAGCCACACCC  
AGTGGGGGCACAGCACAGACCCAGAGCCAGGAACCTCTCAGCCGATGCCCCCAGTATGGGAACCAGCACCTC  
CCACCAACCAGCCACACCCGGTGGGGGCACAGCACAGACCCAGAGGCAGGAACCTCTCAGCCGATGCCCCCG  
GTATGGGAACCAGCACCTCCACCAACCAACCACACCCGGTGGGGGCACAGCACAGACCCAGAGCCAGGAACC  
TCTCAGCCGATGCCCTCAGCAAGAGCACCCCATCTCAGGTGGCGGCCCTCGGAGGACAAGCGCTTCTCGGT  
GGTGGATATGGCGGCCCTGGGCGGGTGTGGGTGCGCTGCTGCTGCTGGCTCTCCTTGGCCTCGCCGTCTTG  
TCCACAAGCACTATGGCCCCCGGCTCAAGTGTGCTCTGGCAAAGCTCCGGAGCCCCAGCCCCAAGGCTTTGAC  
AACCAGGCTTCTCCTCCCTGACCACAAGGCCAACTGGGCGCCCGTCCCAGCCCCACGCACGACCCCAAGCCGC  
GGAGGCACCGATGCCCGCAGAGCCCGCACCCCCGGCCCTGCCCTCCCAGGCGGTGCCCTGAGCCCCCGCAG  
CGCCCCGAGCTGGCGGAAGCCCCACGGCGGTGAGGTCCATCCTGACCAAGGAGCGGCGGCCGGAGGGCGGGTAC  
AAGGCCGTCTGGTTTGGCGAGGACATCGGGACGGAGGCAGACGTGGTCTCTCAACCGCCACCCCTGGACGT  
GGATGGCGCCAGTGACTCCGGCAGCGGCGACGAGGGCGAGGGCGCGGGGAGGGGTGGGGGTCCCTACGATGCAC  
CCGTGGTGATGACTCCTACATCTAAAGTGGCCCTCCACCCCTCCCCAGCCGACGGGCACTGGAGGTCTCG  
CTCCCCAGCCTCCGACCCGAGGCAGAATAAAGCAAGGCTCCCGAAACCCAGGCCATGGCGTGGGGCAGGCGCG  
TGGGTCCCTGGGGGCCCATTCACCTCAGTCCCCTGTCTGCTATTAGCGCTTGAGCCAGGTGTGCAGATGAGGCG  
GTGGGTCTGGCCACGCTGTCCCACCCCAAGGCTGCAGCACTTCCCGTAAACCACCTGCAGTGCCCGCCGCTT  
CCCAGGCTCTGTGCCAGCTAGTCTGGGAAGTTTCTCTCCCGCTTAACCACAGCCGAGGGGGGCTCCCCTCC  
CCCGACCTGCACCAGAGATCTCAGGCACCCGGCTCAACTCAGACCTCCCGCTCCCGACCCCTACACAGAGATTGC  
CTGGGGAGGCTGAGGAGCCGATGCAAACCCCAAGGCGACGCACTTGGGAGCCGGTGGTCTCAAACACCTGCCG  
GGGTCCCTAGTCCCCTCTGAAATCTACATGCTTGGTGGAGCGCAGCAGTAAACACCCCTGCCAGTGACCTG  
GACTGAGGCGCGTGGGGTGGGTGCGCCGTGTGGCTGAGCAGGAGCCAGACCAGGAGCCATAGGGGTGAGAG  
ACACATCCCCTCGCTGCTCCCAAAGCCAGAGCCAGGCTGGGCGCCCATGCCAGAACCATCAAGGGATCCCT  
TGCGGCTTGTGACACTTTCCCTAATGGAATAACCCATTAATTCCTTTCAAATGTTTT

## **FIGURE 54**

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA102846

><subunit 1 of 1, 839 aa, 1 stop

><MW: 87546, pI: 4.84, NX(S/T): 8

MGSWALLWPPLLFTGLLVRRPPGTMAQAQYCSVNKDI FEVEENTNVTEPLVDIHVPEGQEV  
TLGALSTPFAFRIQGNQLFLNVTDPYEEKSLLEAQLLCQSGGTLVTQLRVFVSVLDVNDN  
APEFPFKTKKEIRVEEDTKVNSTVIPETQLQAEDRDKDDILFYTLQEMTAGASDYFSLVSV  
NRPALRLDRPLDFYERPNMTFWLLVRDTPGENVEPSHTATATLVLNVVPADLRPPWFLPC  
TFSDGYVCIQAQYHGAVPTGHILPSPLVLRPGPIYAEDGDRGINQPIIYSIFRGNVNGTF  
IIHPDSGNLTVARSVPSPMTFLLLVKGGQADLARYSVTQVTVEAVAAAGSPPRFPQSLYR  
GTVARGAGAGVVVKDAAAPSQPLRIQAQDPEFSDLNSAITYRITNHSFRMEGEVVLTTT  
TLAQAGAFYAEVEAHNTVTSQTATTVIEIQVSEQEPSTEAGGTTGPWTSTTSEVPRPPE  
PSQGPSTTSSGGGTGPHPPSGTTLRPPTSSTPGGPPGAENSTSHQPATPGGDTAQT PKPG  
TSQPMPPGVGTSTSHQPATPSGGTAQTPEPGTSQPMPPSMGTSTSHQPATPGGGTAQTPE  
AGTSQPMPPGMGTSTSHQPSTPGGGTAQTPEPGTSQPMPLSKSTPSSGGGPPSEDKRFVSV  
DMAALGGVLGALLLLALLGLAVLVHKKHYGPRKCCSGKAPEPQPQGFNDQAFLPDHKANW  
APVPSPTHDPKPAEAPMPAEPAPPGPASPGGAPEPPAAARAGGSPTAVRSILTKERRPEG  
GYKAVWFGEDIGTEADVVLNAPTLDVDGASDSGSGDEGEGAGRGGGPPYDAPGGDDSYI

### **Important features of the protein:**

#### **Signal peptide:**

Amino acids 1-25

#### **Transmembrane domain:**

Amino acids 662-684

#### **N-glycosylation sites:**

Amino acids 44-48;140-144;198-202;297-301;  
308-312;405-409;520-524

#### **Glycosaminoglycan attachment sites:**

Amino acids 490-494;647-651;813-817

#### **cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 655-659

#### **Tyrosine kinase phosphorylation sites:**

Amino acids 154-163;776-783

#### **N-myristoylation sites:**

Amino acids 57-63;102-108;255-261;294-300;  
366-372;426-432;441-447;513-519;  
517-523;530-536;548-554;550-556;  
581-587;592-598;610-616;612-618;  
623-629;648-654;666-672;667-673;  
762-768;763-769;780-786;809-815;  
821-827;833-839

#### **Cadherins extracellular repeated domain signature:**

Amino acids 112-123



**FIGURE 56**

MVGFGANRRAGRLPSLVLVVLLVIVVLAFFNYWSISSRHVLLQEEVAELQGQVQRTEVAR  
GRLEKRNSDLLLLVDTHKKQIDQKEADYGRSSRLQAREGLGKRCEDDKVKLQNNISYQM  
ADIIHHLKEQLAELRQEFRLRQEDQLQDYRKNNTYLVKRLEYESFQCGQQMKELRAQHEENI  
KKLADQFLEEQKQETQKIQSNDGKELDINNQVVPKNI PKVAENVADKNEEPSSNHI PHGK  
EQIKRGGDAGMPGIEENDLAKVDDLPPALRKPPISVSQHESHQAI SHLPTGQPLSPNMPP  
DSHINHNGNPGTSKQNPSSPLQRLIPGSNLDSEPRIQT DILKQATKDRVSDFHKLKQNDE  
ERELQMDPADYGKQHFNDVL

**Important features of the protein:**

**Signal peptide:**

1-29

**Transmembrane domain.**

None

**N-glycosylation site.**

115-119

150-154

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

65-69

**N-myristoylation site.**

246-252

253-259

308-314

**Amidation site.**

101-105

**FIGURE 57**

GGATGGGCGAGCAGTCTGAATGCCAGAAATGGATAACCGTTTTGCTACAGCATTGTAAATTGC  
TTGTGTGCTTAGCCTCATTTCACCATCTACATGGCAGCCTCCATTGGCACAGACTTCTGGT  
ATGAATATCGAAGTCCAGTTCAAGAAAATTCCAGTGATTTGAATAAAAGCATCTGGGATGAA  
TTCATTAGTGATGAGGCAGATGAAAAGACTTATAATGATGCACTTTTTCGATAACAATGGCAC  
AGTGGGATTGTGGAGACGGTGTATCACCATAACCCAAAAACATGCATTGGTATAGCCCACCAG  
AAAGGACAGAGTCATTTGATGTGGTCACAAAATGTGTGAGTTTCACACTAACTGAGCAGTTC  
ATGGAGAAATTTGTTGATCCCGGAAACCACAATAGCGGGATTGATCTCCTTAGGACCTATCT  
TTGGCGTTGCCAGTTCCTTTTACCTTTTGTGAGTTTAGGTTTGATGTGCTTTGGGGCTTTGA  
TCGGACTTTGTGCTTGCATTTGCCGAAGCTTATATCCCACCATTGCCACGGGCATTCTCCAT  
CTCCTTGCAGATACCATGCTGTGAAGTCCAGGCCACATGGAGGTGCCTGTGTAGATGCTCC  
AGCTGAAATCCCAAGCTAAGCTCCCAACTGACAGCCAACATCATTCCAGCCATGTGTGGGA  
GCCATCCTGGATGTCCAGCCTAACAAGCCTTCAGAGGACTTCAGCCACAGCTATTATCTTA  
CTACATCCTTGTGAGACTCTAATAAAGAACCAACTAGCTGAGCCCAATCAACCTATGGAAGT  
ATAGAAATAAAATGAATTGTTGTTTTGTGCCGTT

## **FIGURE 58**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA102880
><subunit 1 of 1, 184 aa, 1 stop
><MW: 21052, pI: 5.01, NX(S/T): 3
MDNRFATAFVIACVLSLISTIYMAASIGTDFWYEYRSPVQENSSDLNKS IWDEFISDEAD
EKTYNDALFRYNGTVGLWRRGITIPKNMHWYSPPERTESFDVVTKCVSFTLTEQFMK FV
DPGNHNSGIDLLR TYLWRCQFLLPFVSLGLMCFGALIGLCACICRSLYPTIATGILHLLA
DTML
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-20

**Transmembrane domain:**

Amino acids 142-163

**N-glycosylation sites:**

Amino acids 42-46;47-51;72-76;

**N-myristoylation sites:**

Amino acids 123-129;154-160;158-164

**Prokaryotic membrane lipoprotein lipid attachment site:**

Amino acids 152-163





## **FIGURE 60**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA105782
><subunit 1 of 1, 156 aa, 1 stop
><MW: 17472, pI: 10.01, NX(S/T): 1
MAPARAGFCPLLLLLLLGLWVAEIPVSAKPKGMTSSQWFKIQHMQPSPQACNSAMKNINK
HTKRCKDLNTFLHEPFSSVAATCQTPKIACKNGDKNCHQSHGPVSLTMCKLTSGKYPNCR
YKEKRQNKSYVVACKPPQKKDSQQFHLVPVHLDRVL
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-22

**N-glycosylation site:**

Amino acids 127-131

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 139-143

**N-myristoylation sites:**

Amino acids 18-24;32-38

**Pancreatic ribonuclease family signature:**

Amino acids 65-72

**Pancreatic ribonuclease family proteins:**

Amino acids 49-93

**FIGURE 61**

CGGGTCATGCGCCGCCGCCTGTGGCTGGGCCTGGCCTGGCTGCTGCTGGCGCGGGCGCCGGA  
CGCCGCGGGAACCCCGAGCGCGTCGCGGGGACCGCGCAGCTACCCGCACCTGGAGGGCGACGTG  
CGCTGGCGGGCGCTCTTCTCCTCCACTCACTTCTTCTGCGCGTGGATCCCGGCGGCCGCGT  
GCAGGGCACCCGCTGGCGCCACGGCCAGGACAGCATCCTGGAGATCCGCTCTGTACACGTGG  
GCGTCGTGGTCATCAAAGCAGTGTCTCAGGCTTCTACGTGGCCATGAACCGCCGGGGCCGC  
CTCTACGGGTCGCGACTCTACACCGTGGACTGCAGGTTCCGGGAGCGCATCGAAGAGAACGG  
CCACAACACCTACGCCTCACAGCGCTGGCGCCGCCGCGGCCAGCCATGTTCTGGCGCTGG  
ACAGGAGGGGGGGGCCCCGGCCAGGCGGCCGGACGCGGGCGGTACCACCTGTCCGCCCACTTC  
CTGCCCCTCCTGGTCTCCTTGAG

## FIGURE 62

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA108912
><subunit 1 of 1, 170 aa, 1 stop
><MW: 19663, pI: 11.81, NX(S/T): 0
MRRRLWLGLAWLLLARAPDAAGTPSASRGPRSYPHLEGDVRWRRLFSSTHFFLRVDPGGR
VQGTRWRHGGQDSILEIRSVHVGVVVIKAVSSGFYVAMNRRGRLYGSRLYTVDCRFERIE
ENGHNTYASQRWRRRGQPMFLALDRRGGPRPGGRTRRYHLSAHFLPVLVS
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-17

**N-myristoylation site:**

Amino acids 22-28

**HGF/FGF family proteins:**

Amino acids 74-125;139-166

**FIGURE 63**

ATCCCTCGACCTCGACCCACGCGTCCGCTGGAAGGTGGCGTGCCCTCCTCTGGCTGGTACCA  
**TG**CAGCTCCCCTGGCCCTGTGTCTCGTCTGCCTGCTGGTACACACAGCCTTCCGTGTAGTG  
GAGGGCCAGGGGTGGCAGGCGTTCAAGAATGATGCCACGGAAATCATCCCCGAGCTCGGAGA  
GTACCCCGAGCCTCCACCGGAGCTGGAGAACAACAAGACCATGAACCGGGCGGAGAACGGAG  
GGCGGCCTCCCCACCACCCCTTTGAGACCAAAGACGTGTCCGAGTACAGCTGCCGCGAGCTG  
CACTTCACCCGCTACGTGACCGATGGGCGGTGCCGCAGCGCCAAGCCGGTCACCGAGCTGGT  
GTGCTCCGGCCAGTGCGGCCCGGCGCGCTGCTGCCCAACGCCATCGGCCGCGGCAAGTGGT  
GGCGACCTAGTGGGCCCGACTTCCGCTGCATCCCCGACCGCTACCGCGCGCAGCGCGTGCAG  
CTGCTGTGTCCCGGTGGTGAAGCGCCGCGCGCGCAAGGTGCGCCTGGTGGCCTCGTGCAA  
GTGCAAGCGCCTCACCCGCTTCCACAACCAGTCGGAGCTCAAGGACTTCGGGACCGAGGCCG  
CTCGGCCGAGAAGGGCCGGAAGCCGCGGCCCGCGCCCGGAGCGCCAAAGCCAACCAGGCC  
GAGCTGGAGAACGCCTACT**TAG**AGCCCGCCCGCGCCCCCTCCCCACCGGCGGGCGCCCCGGCCC  
TGAACCCGCGCCCCACATTTCTGTCTCTGCGCGTGGT'TTGATTGTTTATATTTTATTGTAA  
ATGCCTGCAACCCAGGGCAGGGGGCTGAGACCTTCCAGGCCCTGAGGAATCCCGGGCGCCGG  
CAAGGCCCCCTCAGCCCGCCAGCTGAGGGGTCCACAGGGGCAGGGGAGGGAATTGAGAGTC  
ACAGACACTGAGCCACGCAGCCCCGCTCTGGGGCCGCTACCTTTGCTGGTCCCCTTACAG  
AGGAGGCAGAAATGGAAGCATTTTCACCGCCCTGGGGT'TTTAAGGGAGCGGTGTGGGAGTGG  
GAAAGTCCAGGGACTGGTTAAGAAAGTTGGATAAGATTCCCCCTTGACCTCGCTGCCCATC  
AGAAAGCCTGAGGCGTGCCAGAGCACAAGACTGGGGGCAACTGTAGATGTGGTTTCTAGTCC  
TGGCTCTGCCACTAACTTCTGTGTAACCTTGAACACTACACAATTCTCCTTCGGGACCTCAAT  
TTCCACTTTGTAATAATGAGGGTGGAGGTGGGAATAGGATCTCGAGGAGACTATTGGCATATG  
ATTCCAAGGACTCCAGTGCCTTTTGAATGGGCAGAGGTGAGAGAGAGAGAGAAAGAGAGA  
GAATGAATGCAGTTGCATTGATTCAGTGCCAAGGTCACTTCCAGAATTCAGAGTTGTGATGC  
TCTCTTCTGACAGCCAAAGATGAAAAACAACAGAAAAAAAAAAGTAAAGAGTCTATTTATG  
GCTGACATATTTACGGCTGACAAACTCCTGGAAGAAGCTATGCTGCTTCCCAGCCTGGCTTC  
CCCGGATGTTTGGCTACCTCCACCCCTCCATCTCAAAGAAATAACATCATCCATTGGGGTAG  
AAAAGGAGAGGGTCCGAGGGTGGTGGGAGGGATAGAAATCACATCCGCCCAACTTCCCAAA  
GAGCAGCATCCCTCCCCGACCCATAGCCATGTTTTAAAGTCACCTTCCGAAGAGAAGTGAA  
AGGTTCAAGGACACTGGCCTTGCAAGGCCGAGGGAGCAGCCATCACAACTCACAGACCAGC  
ACATCCCTTTTGAGACACCGCCTTCTGCCACCCTCACGGACACATTTCTGCCTAGAAAAC  
AGCTTCTTACTGCTCTTACATGTGATGGCATATCTTACACTAAAAGAATATTATTGGGGGAA  
AACTACAAGTGCTGTACATATGCTGAGAACTGCAGAGCATAATAGCTGCCACCCAAAAAT  
CTTTTTGAAAATCATTTCCAGACAACCTTACTTTCTGTGTAGTTTTTAAATTGTTAAAAAA  
AAAAAGTTTTAAACAGAAGCACATGACATATGAAAGCCTGCAGGACTGGTCGTTTTTTTTGCG  
AATCTTCCACGTGGGACTTGTCCACAAGAATGAAAGTAGTGGTTTTTAAAGAGTTAAGTTA  
CATATTTATTTTCTCACTTAAGTTATTTATGCAAAGTTTTTCTGTAGAGAATGACAATGT  
TAATATTGCTTTATGAATTAACAGTCTGTTCTTCCAGAGTCCAGAGACATTGTTAATAAAGA  
CAATGAATCATGAAAAAAAAAAAAAAAAAAAAA

## **FIGURE 64**

```
</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA115253
<subunit 1 of 1, 213 aa, 1 stop
<MW: 24031, pI: 9.59, NX(S/T): 2
MQLPLALCLVCLLVHTAFRVVEGQGWQAFKNDATTEIIPELGEYPEPPPELENNKTMNRAE
NGGRPPHHPFETKDVSEYSCRELHFTRYVTDGPCRSAPVTELVCSGQCGPARLLPNAIG
RGKWWRPSPDFRCIPDRYRAQRVQLLCPGGEAPRARKVRLVASCKCKRLTRFHNQSELK
DFGTEAARPQKGRKPRPRARSAKANQAELENAY
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-16

**N-glycosylation sites:**

Amino acids 53-57;175-179

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 168-172

**N-myristoylation site:**

Amino acids 183-189

**Amidation site:**

Amino acids 191-195

### FIGURE 65

CCCCTCGGCGGTTTGGCGGGAGGGAGGGGCTTTGCGCAGGCCCCCGCTCCCGCCCCGCTCC  
**ATG**CGGCCCCGCCCCGATTGCGCTGTGGCTGCGCCTGGTCTTGGCCCTGGCCCTTGTCCGCC  
CCGGGCTGTGGGGTGGGCCCCGGTCCGAGCCCCATCTATGTCAGCAGCTGGGCCGTCCAGG  
TGTCCAGGGTAACCGGGAGGTCGAGCGCCTGGCACGCAAATTCGGCTTCGTCAACCTGGGG  
CCGATCTTCTCTGACGGGCAGTACTTTCACCTGCGGCACCGGGGCGTGGTCCAGCAGTCCCT  
GACCCCGCACTGGGGCCACCGCCTGCACCTGAAGAAAACCCCAAGGTGCAGTGGTTCCAGC  
AGCAGACGCTGCAGCGGCGGGTGAACGCTCTGTCTGGTGGCCACGGACCCCTGGTTCTCC  
AAGCAGTGGTACATGAACAGCGAGGCCCAACCAGACCTGAGCATCCTGCAGGCCCTGGAGTCA  
GGGGCTGTGAGGCCAGGGCATCGTGGTCTCTGTGCTGGACGATGGCATCGAGAAGGACCACC  
CGGACCTCTGGGCCAACTACGACCCCTGGCCAGCTATGACTTCAATGACTACGACCCGGAC  
CCCCAGCCCCGCTACACCCCGAGCAAAGAGAACCGGCACGGGACCCGCTGTGCTGGGGAGGT  
GGCCGCGATGGCCAACAATGGCTTCTGTGGTGTGGGGGTGCTTTCAACGCCCGAATCGGAG  
GCGTACGGATGCTGGACGGTACCATCACCGATGTCATCGAGGCCAGTCGCTGAGCCTGCAG  
CCGCAGCACATCCACATTTACAGCGCCAGCTGGGGTCCCGAGGACGACGGCCGCACGGTGA  
CGGCCCCGGCATCCTCACCCGCGAGGCCCTCCGGCGTGGTGTGACCAAGGGCCGCGGGCGGGC  
TGGGCACGCTCTTCATCTGGGCCTCGGGCAACGGCGGCCTGCACTACGACAACCTGCAACTGC  
GACGGCTACACCAACAGCATCCACACGCTTTCGCTGGGCAGCACCCAGCAGGGCCGCGT  
GCCCTGGTACAGCGAAGCCTGCGCCTCCACCCTCACACCACCTACAGCAGCGGCCTGGCCA  
CCGACCCCGAGATCGTCACCACGGACCTGCATCACGGGTGCACAGACCAGCACACGGGCACC  
TCGGCCTCAGCCCCACTGGCGGCCGGCATGATCGCCCTAGCGCTGGAGGCCAACCCGTTCTCT  
GACGTGGAGAGACATGCAGCACCTGGTGGTCCGCGCGTCCAAGCCGGCGCACCTGCAGGCCG  
AGGACTGGAGGACCAACGGCGTGGGGCGCCAAGTGAGCCATCACTACGGATACGGGCTGCTG  
GACGCCGGGCTGCTGGTGGACACCGCCCGCACCTGGCTGCCACCCAGCCGCAGAGGAAGTG  
CGCCGTCCGGGTCCAGAGCCGCCCCACCCCATCCTGCCGCTGATCTACATCAGGGAAAACG  
TATCGGCCTGCGCCGGCCTCCACAACCTCCATCCGCTCGCTGGAGCACGTGCAGGGCGCAGCTG  
ACGCTGTCCTACAGCCGGCGCGGAGACCTGGAGATCTCGCTCACCCAGCCCCATGGGCACGG  
CTCCACACTCGTGGCCATACGACCCTTGGACGTCAGCACTGAAGGCTACAACAACCTGGGTCT  
TCATGTCCACCCACTTCTGGGATGAGAACCACAGGGCGTGTGGACCCTGGGCCCTAGAGAAC  
AAGGGCTACTATTTCAACACGGGGACGTTGTACCGCTACACGCTGCTGCTCTATGGGACGGC  
CGAGGACATGACAGCGCGGCCTACAGGCCCCAGGTGACCAGCAGCGCGTGTGTGCAGCGGGAC  
ACAGAGGGGCTGTGCCAGGCGTGTGACGGCCCCGCCTACATCCTGGGACAGCTCTGCCTGGC  
CTACTGCCCCCGCGGTTCTTCAACCACACAAGGCTGGTGACCGCTGGGCCTGGGCACACGG  
CGGCGCCCCGCGCTGAGGGTCTGCTCCAGCTGCCATGCCTCCTGCTACACCTGCCGCGGGC  
TCCCCGAGGGACTGCACCTCCTGTCCCCATCCTCCACGCTGGACCAGCAGCAGGGCTCCTG  
CATGGGACCCACCACCCCGACAGCCGCCCGGGCTTAGAGCTGCCGCCTGTCCCCACCACCG  
CTGCCCAGCCTCGGCCATGGTGTGAGCCTCCTGGCCGTGACCCTCGGAGGCCCGCTCCTCT  
GCGGCATGTCCATGGACCTCCACTATACGCCTGGCTCTCCCGTGCCAGGGCCACCCACC  
AAACCCAGGTCTGGCTGCCAGCTGGAACCT**TGA**AGTTGTCAGCTCAGAAAGCGACCTTGCCC  
CCGCTGGGTCCCTGACAGGCACTGCTGCCATGCTGCCTCCCAGGCTGGCCCCAGAGGAGC  
GAGACCAGCACCCGACGCTGGCCTGCCAGGGATGGGCCCCGTGGAACCCCGAAGCCTGGC  
GGGAGAGAGAGAGAGAAGTCTCCTCTGCATTTTGGGTTTGGGCAGGAGTGGGCTGGGGGG  
AGAGGCTGGAGCACCCCAAAGCCAGGGAAAGTGGAGGGAGAGAAACCTGACACTGTCCGT  
CTCGGGCACCGCGTCCAACCTCAGAGTTTGCAAATAAAGGTTGCTTAGAAGGTGAA

## FIGURE 66

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA119302
><subunit 1 of 1, 755 aa, 1 stop
><MW: 82785, pI: 8.71, NX(S/T): 2
MRPAPIALWLRLVLALALVRPRAVGWAPVRAPIYVSSWAVQVSQGNREVERLARKFGFVN
LGPIFSDGQYFHLRHRGVVQQSLTPHWGHRHLKKNPKVQWFQQOTLQRRVKRSVVVPTD
PWFSKQWYMNSEAQPDLNILQAWSQGLSQGIVVSVLDDGIEKDHPDLWANYDPLASYDF
NDYDPPDPPRYTPSKENRHGTRCAGEVAAMANNFGCGVGVAFNARIGGVRMLDGTITDVI
EAQSLSLQPQHIHIYSASWGPEDDGRTVDGPGILTREAFRRGVTKGRGGLTLFIWASGN
GGLHYDNCNCDGYTNSIHTLSVSGSTTQQGRVPWYSEACASTLTTTYSSGVATDPQIVTTD
LHHGCTDQHTGTSASAPLAAGMIALALEANPFLTWRDMQHLVVRASKPAHLQAEDWRTNG
VGRQVSHHYGYGLLDAGLLVDTARTWLPTQPQRKCAVRVQSRPTPILPLIYIRENVSACA
GLHNSIRSLEHVQAQLTLSYSRRGDLEISLTPMGRSTLVAIRPLDVSTEGYNNWVEMS
THFDENPQGVWTLGLENKGYFNTGTLYRYTLLLYGTAEDMTARPTGPQVTSSACVQRD
TEGLCQACDGPAYILGQLCLAYCPPRFNHNTRLVTAGPGHTAAPALRVCSSCHASCYTCR
GGSPRDCTSCPPSSTLDQQQGSCMGPTTPDSRPRLRAAACPHHRCPASAMVLSLLAVTLG
GPVLCGMSMDLPLYAWLSRARATPTKPQVWLPAGT
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-21

**Transmembrane domain:**

Amino acids 706-730

**N-glycosylation sites:**

Amino acids 475-479;629-633

**Glycosaminoglycan attachment sites:**

Amino acids 148-152;298-302

**N-myristoylation sites:**

Amino acids 151-157;200-206;217-223;219-225;  
282-288;288-294;371-377;432-438;  
481-487;515-521;603-609

**Prokaryotic membrane lipoprotein lipid attachment site:**

Amino acids 586-597

**Cell attachment sequence:**

Amino acids 503-506

**Serine proteases, subtilase family, aspartic acid active site:**

Amino acids 154-166

**Serine proteases, subtilase family, histidine active site:**

Amino acids 199-210

**Serine proteases, subtilase family, serine active site:**

Amino acids 371-382

**Cytochrome c family heme-binding site signature:**

Amino acids 649-655

**FIGURE 67**

ATGAGGAAGCTCCAGGGCAGGATGGTTTACCTGCCTGGACAGCAAG**ATG**ATGGCTTACACTAG  
CCCCATTCTCTGGGCGCCTGGATTTGCCACCAGATCTCCTCACCTCTTGCCCTTCACCTC  
CTGCTGTACCTACAAGGTCTCCCCGATTCTCATCTGCCATAATCATGGACACAGCCCCAGG  
ATGTGCAGGACTCTCAGGGACCATCTGGAGTTCCAGCTGGAATCTGGGCCTGGTGGAGTGGG  
AGTGGGGCAGGGGCCTGCATTGGGCTGACTTAGAGAGCACAGTTATTCCATCCATATGGAAA  
**TAA**ACATTTTGGATTCCTGATC



## FIGURE 68

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA119536
><subunit 1 of 1, 88 aa, 1 stop
><MW: 9645, pI: 5.45, NX(S/T): 0
MMATLAPILWAPGFAHQISSPLALHLLLYLQGLPDSHLPIIMDTAPGCAGLSGTIWSSSW
NLGLVEWEWGRGLHWADLESTVIPSIWK
```

**Signal sequence:**

Amino acids 1-15

**N-myristoylation sites:**

Amino acids 32-38;50-56;53-59;72-78

**FIGURE 69**

TTTGCAGTGGGGTCCTCCTCTGGCCTCCTGCCCTCCTGCTGCTGCTGCTGCTTCCATTGCT  
GGCAGCCCAGGGTGGGGGTGGCCTGCAGGCAGCGCTGCTGGCCCTTGAGGTGGGGCTGGTGG  
GTCTGGGGGCCTCCTACCTGCTCCTTTGTACAGCCCTGCACCTGCCCTCCAGTCTTTTCCTA  
CTCCTGGCCCAGGGTACCGCACTGGGGGCCGTCTGGGCCCTGAGCTGGCGCCGAGGCCTC**AT**  
**G**GGTGTTCCTTGGCCCTTGGAGCTGCCTGGCTCTTAGCTTGGCCAGGCCTAGCTCTACCTC  
TGGTGGCTATGGCAGCGGGGGGAGATGGGTGCGGCAGCAGGGCCCCGGGTGCGCCGGGGC  
ATATCTCGACTCTGGTTGCGGGTCTGCTGCGCCTGTCACCCATGGCCTTCCGGGCCCTGCA  
GGGCTGTGGGGCTGTGGGGGACCGGGTCTGTTTGCAGTGTACCCAAAACCAACAAGGATG  
GCTTCCGCAGCCGCTGCCCGTCCCTGGGGCCCCGGCGGCGTAATCCCCGCACCACCAACAC  
CCATTAGCTCTGTTGGCAAGGGTCTGGGTCTGTGCAAGGGCTGGAACGGCGTCTGGCAGC  
GGCCAGCCAGGGTTTAGCATCCCACTTGCCCCCGTGGGCCATCCACACACTGGCCAGCTGGG  
GCCTGCTTCGGGGTGAACGGCCACCCGAATCCCCGGCTACTACCACGCAGCCAGCGCCAG  
CTAGGGCCCCCTGCCTCCCGCCAGCCACTGCCAGGGACTCTAGCCGGGCGGAGGTCACGCAC  
CCGCCAGTCCCGGGCCCTGCCCCCTGGAGG**TAG**CTGACTCCAGCCCTTCCAGCCCAAATCT  
AGAGCATTGAGCACTTTATCTCCCACGACTCAGTGAAGTTTCTCCAGTCCCTAGTCCTCTCT  
TTTCACCCACCTTCCTCAGTTTGCTCACTTACCCAGGCCAGCCCTTCGGACCTCTAGACA  
GGCAGCCTCCTCAGCTGTGGAGTCCAGCAGTCACTCTGTGTTCTCCTGGCGCTCCTCCCCTA  
AGTTATTGCTGTTCGCCCCTGTGTGTGCTCATCCTCACCTCATTGACTCAGGCCTGGGGC  
CAGGGGTGGTGGAGGGTGGGAAGAGTCATGTTTTTTTTCTCCTCTTTGATTTTGTTTTTCTG  
TCTCCCTTCCAACCTGTCCCTTCCCCCACCAAAAAAAAXXXXXXXXXXXXXXXXXXXXXXXXXX  
XXXXXXXXXXXXXXXXAAA

## **FIGURE 70**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA119542
><subunit 1 of 1, 197 aa, 1 stop
><MW: 21992, pI: 12.18, NX(S/T): 0
MGVPLGLGAAWLLAWPGLALPLVAMAAGGRWVRQQGPRVRRGISRLWLRVLLRLSPMAFR
ALQCGAVGDRGLFALYPKTNKDGFRSRLPVPGPRRRNPRTTQHPLALLARVWVLCKGWN
WRLARASQGLASHLPPWAIHTLASWGLLRGERPTRIPRLLPRSQRQLGPPASRQPLPGTL
AGRRSRTRQSRALPPWR
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-21

**N-myristoylation sites:**

Amino acids 2-8;6-12;146-152;178-184

**Amidation site:**

Amino acids 181-185

**FIGURE 71**

GTTTGGGGGTTGTTTGGGATTAGTGAAGCTACTGCCTTTGCCGCCAGCGCAGCCTCAGAGTT  
TGATTATTTGCA**ATG**TCAGGCTTTGAAAACCTAAACACGGATTTCTACCAGACAAGTTACAG  
CATCGATGATCAGTCACAGCAGTCCATGATTATGGAGGAAGTGGAGGACCCATAGCAAAC  
AGTATGCTGGCTATGACTATTCGCAGCAAGGCAGATTTGTCCCTCCAGACATGATGCAGCCA  
CAACAGCCATACACCGGGCAGATTTACCAGCCAACCTCAGGCATATACTCCAGCTTCACCTCA  
GCCTTCTATGGAAACAACTTTGAGGATGAGCCACCTTTATTAGAAGAGTTAGGTATCAATTTT  
GACCACATCTGGCAAAAAACACTAACAGTATTACATCCGTTAAAAGTAGCAGATGGCAGCAT  
CATGAATGAACTGATTTGGCAGGTCCAATGGTTTTTTGCCTTGCTTTTGGAGCCACATTGC  
TACTGGCTGGCAAAATCCAGTTTGGCTATGTATACGGGATCAGTGCAATTGGATGTCTAGGA  
ATGTTTTGTTTATTAACCTTAATGAGTATGACAGGTGTTTCATTTGGTTGTGTGGCAAGTGT  
CCTTGGATATTGTCTTCTGCCCATGATCCTACTTTCCAGCTTTGCAGTGATATTTTCTTTGC  
AAGGAATGGTAGGAATCATTTCTACTGCTGGGATTATTGGATGGTGTAGTTTTTCTGCTTCC  
AAAATATTTATTTCTGCATTAGCCATGGAAGGACAGCAACTTTTAGTAGCATATCCTTGCGC  
TTTGTATATGGAGTCTTTGCCCTGATTTCCGTCTTT**TG**AAAATTTATCTGGGATGTGGACA  
TCAGTGGGCCAGATGTACAAAAGGACCTTGAACCTTTAAATTGGACCAGCAAACCTGCTGCA  
GCGCAACTCTCATGCAGATTTACATTTGACTGTTGGAGCAATGAAAGTAAACGTGTATCTCT  
TGTTTCATTTTTATAGAACTTTTGCATACTATATTGGATTTACCTGCGGTGTGACTAGCTTTA  
AATGTTTGTGTTTATACAGATAAGAAATGCTATTTCTTTCTGGTTCCTGCAGCCATTGAAA  
ACCTTTTTCTTGCAAATTATAATGTTTTTGGATAGATTTTTATCAACTGTGGGAAACCAAAC  
ACAAAGCTGATAACCTTTCTTAAAAACGACCCAGTCACAGTAAAGAAGACACAAGACGGCCG  
GGCGTGGTAGCTCACGCCTGTAATCCAGCACTTTGGGAGGCCGAGGCGGGCGGATCACAAAG  
GGCAGGAGATCGAGACCATCCTGGTTAACACGGTGAAACCCCGACTCTACTAAAACCTACAAA  
AAAAATTAGCTGGGCGTGGTGGCGGGCGCCTGTAGTCCAGCTACTCAGGAGGCTGAGGCAG  
GAGAAGTGTGAACCCAGGAGGCGGAGCTTGCAGTGAGCCGAGATCACACCACTGCACTCCAT  
CCAGCCTGGGTGACAGGGTGAGACTCTGTCTCAAAAAAAAAAAAAAAAAAAGGAGACACAAGACT  
TACTGCAAAAATATTTTTCCAAGGATTTAGGAAAGAAAAATTGCCTTGTATTCTCAAGTCAG  
GTAACCTCAAAGCAAAAAGTGATCCAATGTAGAGTATGAGTTTGCCTCCAAAATTTGAC  
ATTACTGTAAATTATCTCATGGAATTTTTGCTAAAATTCAGAGATACGGGAAGTTCACAATC  
TACCTCATTGTAGACATGAAATGCGAACACTTACTTACATATTAATGTTAACTCAACCTTAG  
GGACCTGGAATGGTTGCATTAATGCTATAATCGTTGGATCGCCACATTTCCCAAAAATAATA  
AAAAATCACTAACCTTTTTTAAGGAAAATATTTAAAGTTTTACAAAATTCATATTGCAAT  
TATCAATGTAAAGTACATTTGAATGCTTATTTAAACCTTTCCAATTAATTTT

## FIGURE 72

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA143498
><subunit 1 of 1, 257 aa, 1 stop
><MW: 27989, pI: 4.16, NX(S/T): 1
MSGFENLNTDFYQTSYSIDDSQQSYDYGGSGGPYSKQYAGYDYSQQGRFVPPDMMQPQQ
PYTGQIYQPTQAYTPASPQPFYGNFEDEPPLLEELGINFDHIWQKTLTVLHPLKVADGS
IMNETDLAGPMVFCLAFGATLLLAGKIQFGYVYGISAIGCLGMFCLLNLMSTGVSFGCV
ASVLGYCLLPMILLSSFAVIFSLQGMVGIILTAGIIGWCSFSASKIFISALAMEGQQLLV
AYPCALLYGVFALISVF
```

**Transmembrane domain:**

Amino acids 129-145;184-203

**N-glycosylation sites:**

Amino acids 123-127

**N-myristoylation sites:**

Amino acids 32-38;119-125;174-180;178-184;208-214

**Prokaryotic membrane lipoprotein lipid attachment site:**

Amino acids 150-161;169-180

## FIGURE 73

ACACTGGCCAAAACGCGGCTCGCCCTCGGCTGCGCTCGGCTCCCGCGGGCGCTCGGCCCCGA  
GCCCTCCTCCCCCTACCCGCCGGCCGGACAGGGAGGAGCCAATGGCTGGGCCTGCCATCCA  
CACCGCTCCCATGCTGTTCCCTCGTCCTCCTGCTGCCCCAGCTGAGCCTGGCAGGCGCCCTTG  
CACCTGGGACCCCTGCCCGGAACCTCCCTGAGAATCACATTGACCTCCCAGGCCCAGCGCTG  
TGGACGCCTCAGGCCAGCCACCACCGCCGGCGGGCCCGGGCAAGAAGGAGTGGGGCCAGG  
CCTGCCCAGCCAGGCCCAGGATGGGGCTGTGGTCACCGCCACCAGGCAGGCCTCCAGGCTGC  
CAGAGGCTGAGGGGCTGCTGCCTGAGCAGAGTCCTGCAGGCCTGCTGCAGGACAAGGACCTG  
CTCCTGGGACTGGCATTGCCCTACCCCGAGAAGGAGAACAGACCTCCAGGTTGGGAGAGGAC  
CAGGAAACGCAGCAGGGAGCACAAAGAGACGCAGGGACAGGTTGAGGCTGCACCAAGGCCGAG  
CCTTGGTCCGAGGTCCCAGCTCCCTGATGAAGAAGGCAGAGCTCTCCGAAGCCCAGGTGCTG  
GATGCAGCCATGGAGGAATCCTCCACCAGCCTGGCGCCACCATGTTCTTTCTACCACCTT  
TGAGGCAGCACCTGCCACAGAAGAGTCCCTGATCCTGCCCGTCACCTCCCTGCGGCCCCAGC  
AGGCACAGCCCAGGTCTGACGGGGAGGTGATGCCACGCTGGACATGGCCTTGTTCGACTGG  
ACCGATTATGAAGACTTAAAACCTGATGGTTGGCCCTCTGCAAAGAAGAAAGAGAAACACCG  
CGGTAAACTCTCCAGTGATGGTAACGAAACATCACCAGCCGAAGGGGAACCATGCGACCATC  
ACCAAGACTGCCTGCCAGGGACTTGCTGCGACCTGCGGGAGCATCTCTGCACACCCCACAAC  
CGAGGCCTCAACAACAAATGCTTCGATGACTGCATGTGTGTGGAAGGGCTGCGCTGCTATGC  
CAAATTCACCGGAACCGCAGGGTTACACGGAGGAAAGGGCGCTGTGTGGAGCCCAGACGG  
CCAACGGCGACCAGGGATCCTTCATCAACGTCTAGCGGCCCCGCGGGACTGGGGACTGAGCC  
CAGGAGGTTTGCACAAGCCGGGCGATTTGTTTGTAACTAGCAGTGGGAGATCAAGTTGGGA  
ACAGATGGCTGAGGCTGCAGACTCAGGCCCAGGACACTCAACCC

## **FIGURE 74**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA145583
><subunit 1 of 1, 348 aa, 1 stop
><MW: 38536, pI: 8.24, NX(S/T): 1
MAGPAIHTAPMLFLVLLLPQLSLAGALAPGTPARNLPENHIDLPGPALWTPQASHHRRRG
PGKKEWGPGLPSQAQDGAVVTATRQASRLPEAEGLLPEQSPAGLLQDKDLLLGLALPYPE
KENRPPGWERTRKRSREHKRRRDRLRLHQGRALVRGPPSSLMKKAELSEAQVLDAAMEESS
TSLAPTMMFFLTTFEAAPATEESLILPVTSLRPQQAQPRSDGEVMPTLDMALFDWTDYEDL
KPDGWPSAKKKEKHRGKLS SDGNETSPAEGEPCDHHQDCLPGTCCDLREHLCTPHNRGLN
NKCFDDCMCVEGLRCYAKFHRNRRVTRRKGRCEPETANGDQGSFINV
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-24

**N-glycosylation site:**

Amino acids 263-267

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 132-136;323-327

**N-myristoylation sites:**

Amino acids 77-83;343-349

**Amidation site:**

Amino acids 61-65

## FIGURE 75

CAGAAGGGCAAAAACATTGACTGCCTCAAGGTCTCAAGCACCAGTCTTCACCGCGGAAAGCA  
**TGTT**TGTGGCTGTTCCAATCGCTCCTGTTTGTCTTCTGCTTTGGCCCAGGGAATGTAGTTTCA  
CAAAGCAGCTTAACCCCATTTGATGGTGAACGGGATTCTGGGGGAGTCAGTAACTCTTCCCCT  
GGAGTTTCTCTGCAGGAGAGAAGGTCAACTTCATCACTTGGCTTTTCAATGAAACATCTCTTG  
CCTTCATAGTACCCCATGAAACCAAAAGTCCAGAAATCCACGTGACTAATCCGAAACAGGGA  
AAGCGACTGAACTTCACCCAGTCTTACTCCCTGCAACTCAGCAACCTGAAGATGGAAGACAC  
AGGCTCTTACAGAGCCCAGATATCCACAAAGACCTCTGCAAAGCTGTCCAGTTACACTCTGA  
GGATATTAAGACAACCTGAGGAACATACAAGTTACCAATCACAGTCAGCTATTTTCAGAATATG  
ACCTGTGAGCTCCATCTGACTTGCTCTGTGGAGGATGCAGATGACAATGTCTCATTCAGATG  
GGAGGCCCTTGGGAAACACACTTTCAAGTCAGCCAAACCTCACTGTCTCTCTGGGACCCCAGGA  
TTTCCAGTGAACAGGACTACACCTGCATAGCAGAGAATGCTGTGAGTAATTTATCCTTCTCT  
GTCTCTGCCCAGAAGCTTTGCGAAGATGTTAAAATTCATATACAGATACCAAAATGATTCT  
GTTTATGGTTTCTGGGATATGCATAGTCTTCGGTTTTCATCATACTGCTGTTACTTGTTTTGA  
GGAAAAGAAGAGATTCCCTATCTTTGTCTACTCAGCGAACACAGGGCCCCGCAGAGTCCGCA  
AGGAACCTAGAGTATGTTTTCAGTGTCTCCAACGAACAACACTGTGTATGCTTCAGTCACTCA  
TTCAAACAGGGAAACAGAAATCTGGACACCTAGAGAAAATGATACTATCACAATTTACTCCA  
CAATTAATCATTCCAAAGAGAGTAAACCCACTTTTTTCCAGGGCAACTGCCCTTGACAATGTC  
GT**GTAA**GTTGCTGAAAGGCCTCAGAGGAATTCGGGAATGACACGTCTTCTGATCCCATGAGA  
CAGAACAAGAACAGGAAGCTTGGTTCCTGTTGTTCCCTGGCAACAGAATTTGAATATCTAGG  
ATAGGATGATCACCTCCAGTCCCTTCGGACTTAAACCTGCCTACCTGAGTCAAACACCTAAGG  
ATAACATCATTTCCAGCATGTGGTTCAAATAATATTTTCCAATCCACTTCAGGCCAAAACAT  
GCTAAAGATAACACACCAGCACATTGACTCTCTCTTTGATAACTAAGCAAATGGAATTATGG  
TTGACAGAGAGTTTATGATCCAGAAGACAACCACTTCTCTCCTTTTAGAAAGCAGCAGGATT  
GACTTATTGAGAAATAATGCAGTGTGTTGGTTACATGTGTAGTCTCTGGAGTTGGATGGGCC  
CATCCTGATACAAGTTGAGCATCCCTTGTCTGAAATGCTTGGGATTAGAAATGTTTCAGATT  
TCAATTTTTTTTTCAGATTTTGGAAATATTTGCATTATATTTAGCGGTTGAGTATCCAAATCCA  
AAAATCCAAAATTCAAAATGCTCCAATAAGCATTTCCTTGGAGTTTTCATTGATGTCGATGCA  
GTGCTCAAAAATCTCAGATTTTGGAGCAATTTGGATATTGGATTTTTGGATTTGGGATGCTCA  
ACTTGTACAATGTTTATTAGACACATCTCCTGGGACATACTGCCTAACCTTTTGGAGCCTTA  
GTCTCCAGACTGAAAAGGAAGAGGATGGTATTACATCAGCTCCATTGTTTGGCCAAGAA  
TCTAAGTC



## FIGURE 76

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA161000
><subunit 1 of 1, 332 aa, 1 stop
><MW: 37345, pI: 6.72, NX(S/T): 10
MLWLFQSLLEFVFCFGPGNVVSQSSLTPLMVNGILGESVTLPLEFPAGEKVNFFITWLFNET
SLAFIVPHETKSPEIHVTNPKQGKRLNFTQSYSLQLSNLKMEDTGSYRAQISTKTSAKLS
SYTLRILRQLRNIQVTNHSQLFQNMTCELHLTCSVEDADDNVVFRWEALGNTLSSQPNTL
VSWDPRISSEQDYTCIAENAVSNLSFSVSAQKLCEDVKIQYTDTKMILFMVSGICIVFGF
IILLLLVLRKRRDLSLSLSTQRTQGPASARNLEYVSVSPTNNTVYASVTHSNRETEIWTP
RENDTITIYSTINHSKESKPTFSRATALDNVV
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-13

**Transmembrane domain:**

Amino acids 228-247

**N-glycosylation sites:**

Amino acids 58-62;87-91;137-141;144-148;161-165;  
178-182;203-207;281-285;303-307;  
313-317

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 251-255

**Tyrosine kinase phosphorylation sites:**

Amino acids 100-108;186-194

**N-myristoylation sites:**

Amino acids 17-23;105-111;170-176

**Amidation site:**

Amino acids 82-86

**Immunoglobulin domain:**

Amino acids 35-111

**FIGURE 77**

GATCCCTCGACCTCGACCCACGCGTCCGCTCTTTAATGCTTTCTTTTTAAGAGATCACCTTC  
 TGACTTCTCACAGAAGAGGTTAACTATTACCTGTGGGAAGTCAGAAGGTGATCTCTTTAATG  
 CTTTCTTTTTAAGAATTTTTCAAATTGAGACTAATTGCAGAGGTTCCAGTTGACCAGCATTTC  
 ATAGGAATGAAGACAAACACAGAGATGGTGTGTCTAAGAACTTCAAAGGTGTAGACCTCC  
 TGACTGAAGCATATTGGATTTATTTAATTTTTTCTACTGTATTTCTGTCTCTTACAAGGGA  
 AAGTC**CATG**ATTACACTAACTGAGCTAAAATGCTTAGCAGATGCCCAGTCATCTTATCACATC  
 TTA AAAACCATGGTGGGACGTCTTCTGGTATTACATCACACTGATCATGCTGCTGGTGGCCGTG  
 CTGGCCGGAGCTCTCCAGCTGACGCAGAGCAGGGTCTGTGCTGTCTTCCATGCAAAGTGGAA  
 ATTTGACAATCACTGTGCCGTGCCCTGGGACATCCTGAAAGCCAGCATGAACACATCCTCTA  
 ATCCTGGGACACCGCTTCCGCTCCCCCTCCGAATTCAGAATGACCTCCACCGACAGCAGTAC  
 TCCTATATTGATGCCGTCTGTTACGAGAAACAGCTCCATTGGTTTTGCAAAGTTTTTCCCCTA  
 TCTGGTGCTCTTGACACGCTCATCTTTGCAGCCTGCAGCAACTTTGGCTTCACTACCCCA  
 GTACCAGTTCAGGCTCGAGCATTTTGTGGCCATCCTTACAAGTGCTTCGATTCTCCATGG  
 ACCACCCGCGCCCTTTCAGAAACAGTGCTGAGCAGTCAGTGAGGCCTCTGAAACTCTCCAA  
 GTCCAAGATTTTGTCTTTCGCTCCTCAGGGTGTTCAGCTGACATAGATTCCGGCAAACAGTCAT  
 TGCCCTACCCACAGCCAGGTTTGGAGTCAGCTGGTATAGAAAGCCCAACTTCCAGTGGCCTG  
 GACAAGAAGGAGGGTGAACAGGCCAAAGCCATCTTTGAAAAAGTGAAAAGATTCCGCATGCA  
 TGTGGAGCAGAAGGACATCATTTATAGAGTATATCTGAAACAGATAATAGTCAAAGTCATTT  
 TGTTTTGTGCTCATCATAACTTATGTTCCATATTTTTTAACCCACATCACTCTTCAAATCGAC  
 TGTTCACTTGTGTCAGGCTTTTACAGGATATAAGCGCTACCAGTGTGTCTATTCTTGGC  
 AGAAATCTTTAAGGTCTGGCTTCATTTTATGTCATTTTGGTTATACTTTATGGTCTGACCT  
 CTTCCTACAGCTGTGGTGGATGCTGAGGAGTCCCTGAAGCAATATTCCTTTGAGGCGTTA  
 AGAGAAAAAAGCAACTACAGTGACATCCCTGATGTCAAGAATGACTTTGCCTTCATCCTTCA  
 TCTGGCTGATCAGTATGATCCTCTTTATTCCAAACGCTTCTCCATATTCCTATCAGAGGTCA  
 GTGAGAACAAACTGAAACAGATCAACCTCAATAATGAATGGACAGTTGAGAACTGAAAAGT  
 AAGCTTGTGAAAAATGCCCAGGACAAGATAGAAGTGCATCTTTTTATGCTCAACGGTCTTCC  
 AGACAATGTCTTTFAGTTAACTGAAATGGAAGTGCTAAGCCTGGAGCTTATCCAGAGGTGA  
 AGCTGCCCTCTGCAGTCTCACAGCTGGTCAACCTCAAGGAGCTTCGTGTGTACCATTCACTCT  
 CTGGTCTGATACCATCCTGCACTGGCCTTCTAGAGGAGAATTTAAAAATCCTCCGCTGAA  
 ATTTACTGAAATGGGAAAAATCCCACGCTGGGTATTTACCTCAAGAATCTCAAGGAACTTT  
 ATCTTTCGGGCTGTGTTCTCCCTGAACAGTTGAGTACTATGCAGTTGGAGGGCTTTCAGGAC  
 TTA AAAAATCTAAGGACCCTGTA CTTGAAGAGCAGCCTCTCCCGGATCCCACAAGTTGTTACA  
 GACCTCTGCCTTCATTGCAGAACTGTCCCTTGATAATGAGGGAAGCAAACCTGGTTGTGTT  
 GAACAACCTGAAAAAGATGGTCAATCTGAAAAGCCTAGA ACTGATCAGCTGTGACCTGGAAC  
 GCATCCCACATTCATTTTCAGCCTGAATAATTTGCATGAGTTAGACCTAAGGGAAAATAAC  
 CTTAAAACCTGTGGAAGAGATTAGCTTTCAGCATCTTCAGAATCTTTCCTGCTTAAAGTTGTG  
 GCACAATAACATTGCTTATATTCCTGCACAGATTGGGGCATTATCTAACCTAGAGCAGCTCT  
 CTTTGGACCATAATAATATTGAGAATCTGCCCTTGCAGCTTTTCCTATGCACTAACTACAT  
 TATTTGGATCTAAGCTATAACCACTTGACCTTCATTCCAGAAGAAATCCAGTATCTGAGTAA  
 TTTGCAGTACTTTGCTGTGACCAACAACAATATTGAGATGCTACCAGATGGGCTGTTTCAGT  
 GCAAAAAGCTGCAGTGTTTACTTTTGGGAAAAATAGCTTGATGAATTTGTCCCCTCATGTG  
 GGTGAGCTGTCAAACCTTACTCATCTGGAGCTCATTTGTAATTACCTGGAAACACTTCCTCC  
 TGA ACTAGAAGGATGTCAGTCCCTAAAACGGA ACTGTCTGATTGTTGAGGAGAACTTGCTCA  
 ATACTCTTCTCCTCTCCCTGTAACAGAACGTTTACAGACGTGCTTAGACAAATGT**TGA**CTTAAA  
 GAAAAGAGACCCGTGTTTCAAATCATTTTTAAAAGTATGCTCGGCCGGGCGTGGTGGCTCA  
 TGCCTATAATCCCAGCACTTTGGGAGGCCAAGATGGGCCGATTGCTTGGAGTCAAGGAGTTCG  
 AGACCAGTCTGGCCAACTGGTGAACCCCATCTCTGCTAAAAC TACAAAAAATTAGCCAG  
 GCGTGGTGGCGTGC GCCTGTAATCCCAGCTACTTGGGAGGCTGACGCAGGGGAATGCTTGA  
 ACCAGGGAGGTGGAGGTTGCAGTGAGCCGAGATTGTGCCACTGTACACCAGCCTGGGTGACA  
 GAGCAAGACTCTTATCTCAAAAAAAAAAAAAA

## FIGURE 78

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA161005
><subunit 1 of 1, 802 aa, 1 stop
><MW: 92235, pI: 6.80, NX(S/T): 5
MITLTELKCLADAQSSYHILKPWWDVFWYYITLIMLLVAVLAGALQLTQSRVLCCLPCKV
EFDNHC AVPWDILKASMN TSSNPGTPLPLPLRIQNDLHRQQYSYIDAVCYEKQLHWFAKF
FPYLVLLHTLIFAACSNFWLHYPSTSSRLEHFVAILHKCFDSPWTTTRALSETVAEQSVRP
LKLSKSKILLSSSGCSADIDSGKQSLPYPQPGLESAGIESPTSSGLDKKEGEQAKAIFEK
VKRFRMHVEQKDIYRVYLKQIIIVKVILFVLIITYVPYFLTHITLEIDCSVDVQAFTGYK
RYQCVYSLAEIFKVLASFYVILVILYGLTSSYSYLWWM LRS SLKQYSFEALREKSNYS DIP
DVKNDFAFILHLADQYDPLYSKRFSIFLSEVSENK LKQINLNNEWTVEK LKSKLVKNAQD
KIELHLFMLNGLPDNVFELTEMEVLSLELIPEVKLPSAVS QLVNLKELRVYHSSLVVDHP
ALAFLEENL KILRLKFT EMGKI PRWVFHLK NLKELYLSGCVLPEQLSTMQLEGFQDLK NL
RTLYLKSSLSRI PQVVTDL LPSLQKLSLDNEGSKLVV LNNLKKMVNLKSLELISCDLERI
PHSIFSLN NLHEL DLRENNLKTVEEISFOHLQNL SCLKLWHNNIAYIPAQIGALS NLEQL
SLDHNNIENLPLQLFLCTKLHYLDLSYNHLTFIPEEIQYLSNLQYFAVTN NNIEMLPDGL
FQCKKLQCLLLGKNSLMNLS PHVGELS NLTHLELIGNYLETL PPELEG CQSLKRNCLIVE
ENLLNTLPLPVTERLQTCLDKC
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-46

**Transmembrane domains:**

Amino acids 118-138;261-281;311-332

**N-glycosylation sites:**

Amino acids 78-82;355-359;633-637;748-752

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 382-386

**Tyrosine kinase phosphorylation site:**

Amino acids 21-30

**N-myristoylation sites:**

Amino acids 212-218;327-333;431-437;652-658;  
719-725

**Prokaryotic membrane lipoprotein lipid attachment site:**

Amino acids 125-136

**Leucine zipper pattern:**

Amino acids 468-490

**Leucine Rich Repeat:**

Amino acids 609-632; 748-770

**FIGURE 79**

CGGACGCGTGGGCGCGCTCCCTCACGGCCCCTCGGCGGCGCCCGTCCGATCCGGCCTCTCT  
CTGCGCCCCGGGGCGCGCCACCTCCCCGCCGGAGGTGTCCACGCGTCCGGCCGTCCATCCGT  
CCGTCCCTCCTGGGGCCGGCGCTGACC**ATG**CCCAGCGGCTGCCGCTGCCTGCATCTCGTGTG  
CCTGTTGTGCATTCTGGGGGCTCCCGGTCAGCCTGTCCGAGCCGATGACTGCAGCTCCCCT  
GTGACCTGGCCCACGGCTGCTGTGCACCTGACGGCTCCTGCAGGTGTGACCCGGGCTGGGAG  
GGGCTGCACTGTGAGCGCTGTGTGAGGATGCCTGGCTGCCAGCACGGTACCTGCCACCAGCC  
ATGGCAGTGCATCTGCCACAGTGGCTGGGCAGGCAAGTTCTGTGACAAAGATGAACATATCT  
GTACCACGCAGTCCCCCTGCCAGAATGGAGGCCAGTGCATGTATGACGGGGGCGGTGAGTAC  
CATTGTGTGTGCTTACCAGGCTTCCATGGGCGTGACTGCGAGCGCAAGGCTGGACCCTGTGA  
ACAGGCAGGCTCCCCATGCCGCAATGGCGGGCAGTGCCAGGACGACCAGGGCTTTGCTCTCA  
ACTTCACGTGCCGCTGCTTGGTGGGCTTTGTGGGTGCCCGCTGTGAGGTAATGTGGATGAC  
TGCCTGATGCCGCCCTTGTGCTAACGGTGCCACCTGCCTTGACGGCATAAACCGCTTCTCCTG  
CCTCTGTCCTGAGGGCTTTGCTGGACGCTTCTGCACCATCAACCTGGATGACTGTGCCAGCC  
GCCATGCCAGAGAGGGGGCCCGCTGTCGGGACCCTGTCCACGACTTCGACTGCCTCTGCCCC  
AGTGGCTATGGTGGCAAGACCTGTGAGCTTGTCTTACCTGTCCAGACCCCCCAACCACAGTG  
GACACCCCTCTAGGGCCACCTCAGCTGTAGTGGTACCTGCTACGGGGCCAGCCCCCACAG  
CGCAGGGGCTGGTCTGCTGCGGATCTCAGTGAAGGAGGTGGTGCAGGCAAGAGGCTGGGC  
TAGGTGAGCCTAGCTTGGTGGCCCTGGTGGTGTTTGGGGCCCTCACTGCTGCCCTGGTTCTG  
GCTACTGTGTTGCTGACCCCTGAGGGCCTGGCGCCGGGGTGTCTGCCCCCTGGACCCTGTTG  
CTACCCTGCCCCACACTATGCTCCAGCGTGCCAGGACCAGGAGTGTGAGGTTAGCATGCTGC  
CAGCAGGGCTCCCCCTGCCACGTGACTTGCCCCCTGAGCCTGGAAAGACCACAGCACT**GTGA**  
TGGAGGTGGGGGCTTTCTGGCCCCCTTCTCACCTCTTCCACCCCTCAGACTGGAGTGGTCC  
GTTCTCACCACCCCTCAGCTTGGGTACACACACAGAGGAGACCTCAGCCTCACACCAGAAAT  
ATTATTTTTTTAATACACAGAATGTAAGATGGAATTTTATCAAATAAACTATGAAAATGCA  
AAAAAAAAAAAAAAAA

## FIGURE 80

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA170245
><subunit 1 of 1, 383 aa, 1 stop
><MW: 40548, pI: 6.48, NX(S/T): 1
MPSGCRCLHLVCLLCILGAPGQPVRADDCSSSHCDLAHGCCAPDGSCRCDPGWEGLHCERC
VRMPGCQHGTCHQPWQCICHSGWAGKFCDEHICTTQSPCQNGGQCMYDGGGEYHCVCL
PGFHGRDCERKAGPCEQAGSPCRNGGQCQDDQGFALNFTCRCLVGFVGARCEVNVDDCLM
RPCANGATCLDGINRFSLCPEGFAGRECTINLDDCASRPCQRGARCRDRVHDFDCLCPS
GYGGKTCELVLPVPDPPTTVDTPLGPTSAVVVPATGPAPHSAGAGLLRISVKEVRRQEA
GLGEPSSLVALVVFALTAALVLATVLLTLRAWRRGVCPPGPCCYPAPHYAPACQDQECQV
SMLPAGLPLPRDLPEPGKTAL
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-21

**Transmembrane domain:**

Amino acids 306-331

**N-glycosylation site:**

Amino acids 157-160

**Glycosaminoglycan attachment site:**

Amino acids 240-243

**N-myristoylation sites:**

Amino acids 44-49;65-70;243-248;314-319

**Aspartic acid and asparagine hydroxylation sites:**

Amino acids 189-200;227-238

**EGF-like domain cysteine pattern signature:**

Amino acids 46-57;77-88;117-128;160-171;198-209;  
236-247

**Zinc finger, C3HC4 type, signature:**

Amino acids 7-16

**EGF-like domain proteins:**

Amino acids 46-58;77-89;117-129;160-172;198-210;  
216-228;236-248

## FIGURE 81

GTTTGTGCTCAAACCGAGTTCTGGAGAACGCCATCAGCTCGCTGCTTAAAATTAACCACA  
GGTTCCATT**ATG**GGTCGACTTGATGGGAAAGTCATCATCCTGACGGCCGCTGCTCAGGGGAT  
TGGCCAAGCAGCTGCCTTAGCTTTTGCAAGAGAAGGTGCCAAAGTCATAGCCACAGACATTA  
ATGAGTCCAAACTTCAGGAACTGGAAAAGTACCCGGGTATTCAAACCTCGTGTCTTATGATGTC  
ACAAAGAAGAAACAAATTGATCAGTTTGCCAGTGAAGTTGAGAGACTTGATGTTCTCTTTAAT  
GTTGCTGGTTTTGTCCATCATGGAACGTCTCTGGATTGTGAGGAGAAAGACTGGGACTTCTC  
GATGAATCTCAATGTGCGCAGCATGTACCTGATGATCAAGGCATTCCTTCCTAAAATGCTTG  
CTCAGAAATCTGGCAATATTATCAACATGTCTTCTGTGGCTTCCAGCGTCAAAGGAGTTGTG  
AACAGATGTGTGTACAGCACAAACCAAGGCAGCCGTGATTGGCCTCACAAAATCTCTGGCTGC  
AGATTTTCATCCAGCAGGGCATCAGGTGCAACTGTGTGTGCCAGGAACAGTTGATACGCCAT  
CTCTACAAGAAAGAATAACAAGCCAGAGGAAATCCTGAAGAGGCACGGAATGATTTCTGAAG  
AGACAAAAGACGGGAAGATTCGCAACTGCAGAAGAAATAGCCATGCTCTGCGTGTATTTGGC  
TTCTGATGAATCTGCTTATGTAAGTGGTAACCCTGTCATCATTGATGGAGGCTGGAGCTT**GT**  
**GA**TTTTTAGGATCTCCATGGTGGGAAGGAAGGCAGGCCCTTCTATCCACAGTGAACCTGGTT  
ACGAAGAAAACCTACCAATCATCTCCTTCCTGTTAATCACATGTTAATGAAAATAAGCTCTT  
TTAATGATGTCAGTCTTTGCAAGAGTCTGATCTTTAAGTATATTAATCTCTTTGTAATCT  
CTTCTGAAATCA'TTGTAAAGAAATAAAAATATTGAACTCAT

## FIGURE 82

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA171771
><subunit 1 of 1, 245 aa, 1 stop
><MW: 26711, pI: 8.00, NX(S/T): 2
MGRLDGKVIILTAAAQGIGQAAALAFAREGAKVIATDINESKLELEKYPGIQTRVLDVT
KKKQIDQFASEVERLDVLFNVAGFVHHGTVLDCEEKDWDFSMNLNVRSMYLMIKAFLPKM
LAQKSGNIINMSSVASSVKGVNRCVYSTTKAAVIGLTKSLAADFIQQGIRCNCVCPGTV
DTPSLQERIQARGNPEEARNDLKRQKTGRFATAEEIAMLCVYLASDESAYVTGNPVIID
GGWSL
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-20

**N-glycosylation sites:**

Amino acids 39-43;130-134

**Tyrosine kinase phosphorylation site:**

Amino acids 42-50

**N-myristoylation sites:**

Amino acids 17-23;19-25;126-132;156-162;169-175

**Short-chain dehydrogenases/reductases family proteins:**

Amino acids 7-19;73-83;127-164; 169-178

**Short chain dehydrogenase:**

Amino acids 7-183

## FIGURE 83

GGGCGGCGGCGGCAGCGGTTGGAGGTTGTAGGACCGGCGAGGAATAGGAATCATGGCGGCTG  
CGCTGTTTCGTGCTGCTGGGATTCGCGCTGCTGGGCACCCACGGAGCCTCCGGGGCTGCCGGC  
TTCGTCCAGGCGCCGCTGTCCCAGCAGAGGTGGGTGGGGGGCAGTGTGGAGCTGCACTGCGA  
GGCCGTGGGCAGCCCGGTGCCCGAGATCCAGTGGTGGTTTGAAGGGCAGGGTCCCAACGACA  
CCTGCTCCCAGCTCTGGGACGGCGCCCGGCTGGACCGCGTCCACATCCACGCCACCTACCAC  
CAGCACGCGGCCAGCACCATCTCCATCGACACGCTCGTGGAGGAGGACACGGGCACTTACGA  
GTGCCGGGCCAGCAACGACCCGGATCGCAACCACCTGACCCGGGCGCCCAGGGTCAAGTGGG  
TCCGCGCCAGGCAGTTCGTGCTAGTCCCTGGAACCCGGCACAGTCTTCACTACCCTAGAAGAC  
CTTGGCTCCAAGATACTCCTCACCTGCTCCTTGAATGACAGCGCCACAGAGGTCACAGGGCA  
CCGCTGGCTGAAGGGGGGCGTGGTGTGAAGGAGGACGCGCTGCCCGGCCAGAAAACGGAGT  
TCAAGGTGGACTCCGACGACCAGTGGGGAGAGTACTCCTGCGTCTTCCCTCCCGAGCCCATG  
GGCACGGCCAACATCCAGCTCCACGGGCCCTCCAGAGTGAAGGCTGTGAAGTCGTCAGAACA  
CATCAACGAGGGGGAGACGGCCATGCTGGTCTGCAAGTCAGAGTCCGTGCCACCTGTCACTG  
ACTGGGCCTGGTACAAGATCACTGACTCTGAGGACAAGGCCCTCATGAACGGCTCCGAGAGC  
AGGTTCTTCGTGAGTTCCCTCGCAGGGCCGGTTCAGAGCTACACATTGAGAACCTGAACATGGA  
GGCCGACCCCGGCCAGTACCGGTGCAACGGCACCCAGCTCCAAGGGCTCCGACCAGGCCATCA  
TCACGCTCCGCGTGCGCAGCCACCTGGCCGCCCTCTGGCCCTTCCCTGGGCATCGTGGCTGAG  
GTGCTGGTGTGTCACCATCATCTTCATCTACGAGAAGCGCCGGAAGCCCGAGGACGTCCT  
GGATGATGACGACGCCGGCTCTGCACCCCTGAAGAGCAGCGGGCAGCACCAGAATGACAAAG  
GCAAGAACGTCCGCCAGAGGAACCTCTCCCTGAGGCAGGTGGCCCGAGGACGCTCCCTGCTCC  
ACGTCTGCGCCCGCCCGGAGTCCACTCCCAGTGCTTGCAAGATTCCAAGTTCTCACCTCTT  
AAAGAAAACCCACCCCGTAGATTCCCATCATACTTCCCTCTTTTTTAAAAAAGTTGGGTT  
TTCTCCATTCAGGATTCTGTTCCCTTAGGTTTTTTTTCCCTTCTGAAGTGTTCACGAGAGCCCG  
GGAGCTGCTGCCCTGCGGCCCGCTGTGTGGCTTTCAGCCTCTGGGTCTGAGTCATGGCCGGG  
TGGGCGGCACAGCCTTCTCCACTGGCCGGAGTCAGTGCCAGGTCCCTTGCCCTTTGTGGAAAGTC  
ACAGGTCACACGAGGGGGCCCGTGTCCCTGCCTGTCTGAAGCCAATGCTGTCTGGTTGCGCCA  
TTTTTGTGCTTTTATGTTTAAATTTTATGAGGGCCACGGGTCTGTGTTCGACTCAGCCTCAGG  
GACACTCTGACCTCTTGGCCACAGAGGACTCACTTGCCACACCGAGGGGCGACCCCGTCAC  
AGCCTCAAGTCACTCCAAGCCCCCTCCTTGTCTGTGCATCCGGGGGCAGCTCTGGAGGGGG  
TTTGCTGGGGAACCTGGCGCCATCGCCGGGACTCCAGAACCGCAGAAGCCTCCCAGCTCACC  
CCTGGAGGACGGCCGGCTCTCTATAGCACCAGGGCTCACGTGGGAACCCCCCTCCACCCAC  
CGCCACAATAAAGATCGCCCCACCTCCACCCAAAAA



## FIGURE 84

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA173157
><subunit 1 of 1, 385 aa, 1 stop
><MW: 42200, pI: 5.57, NX(S/T): 5
MAAALFVLLGFALLGTHGASGAAGFVQAPLSQQRWVGGSVELHCEAVGSPVPEIQWWFEG
QGPNDTCSQLWDGARLDRVHIHATYHQHAASTISIDTLVEEDTGTYECRASNDPDRNHLT
RAPRVKVVRAQAVVLVLEPGTVFTTVEEDLGSKILLTCSLNDSETEVTGHRWLKGGVVLKE
DALPGQKTEFKVDSDDQWGEYSCVFLPEPMGTANIQLHGPPRVKAVKSSEHINEGETAML
VCKSESVPPVTDWAWYKITDSEDKALMNGSESRRFFVSSSQGRSELHIENLNMEADPGQYR
CNGTSSKGSQAIITLVRSHLAALWFLGIVAEVLVLTIIIFIYEKRRKPEDVLDDDDA
GSAPLKSSGQHQNDRKGNVRQRNSS
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-18

**Transmembrane domain:**

Amino acids 320-343

**N-glycosylation sites:**

Amino acids 64-68;160-164;268-272;302-306

**N-myristoylation sites:**

Amino acids 15-21;18-24;60-66;104-110;140-146;  
297-303;308-314;369-375

**Immunoglobulin domain:**

Amino acids 37-110;150-205;235-303

**FIGURE 85**

GGCTCGAGCAAAGACATACGAACAGGGAGGAAGGCCGACTGAAAGAAAGACGGAGAAGAGGA  
GAGAGAAGCCAGGGCCGAGCGTGCCAGCAGGCCGATGGAGGGCGGCCTGGTGGAGGAGGAGA  
CGTAGTGGCCTGGGCTGAGCTGGGTGGGCCGGGAGAAGCGGGTGCCTCAGAGTGGGGGTGGG  
GGC**ATG**GGAGGGGCAGGCATTCTGCTGCTGCTGCTGGCTGGGGCGGGGGTGGTGGTGGCCTGG  
AGACCCCAAAGGGAAAGTGTCCCCTGCGCTGCTCCTGCTCTAAAGACAGCGCCCTGTGTGA  
GGGCTCCCCGGACCTGCCCGTCAGCTTCTCTCCGACCCTGCTGTCACTCTCACTCGTCAGGA  
CGGGAGTCACCCAGCTGAAGGCCGGCAGCTTCTGAGAATTCCGTCTCTGCACCTGCTCCTC  
TTCACCTCCAACCTCTTCTCCGTGATTGAGGACGATGCATTTGCGGGCCTGTCCCACCTGCA  
GTACCTCTTCATCGAGGACAATGAGATTGGCTCCATCTCTAAGAATGCCCTCAGAGGACTTC  
GCTCGCTTACACACCTAAGCCTGGCCAATAACCATCTGGAGACCCTCCCCAGATTCTGTTT  
CGAGGCCTGGACACCCTTACTCACGTGGACCTCCGCGGGAACCCGTTCCAGTGTGACTGCCG  
CGTCTCTGGCTCCTGCAGTGGATGCCACCGTGAATGCCAGCGTGGGGACCGGCCTGTG  
CGGGCCCCGCCTCCCTGAGCCACATGCAGCTCCACCACCTCGACCCCAAGACTTCAAGTGC  
AGAGCCATAGGTGGGGGGCTTCCCGATGGGGTGGGAGGCGGGAGATCTGGGGGAAAGGCTG  
CCAGGGCCAAGAGGCTCGTCTCACTCCCTGCCCTGCCATTTCCCGAGTGGGAAGACCCTGA  
GCAAGCAGCACTGCCTTCTGAGCCCCAGTTTTCTCATCTG**TAA**AGTGGGGTAATAAACAG  
TGATATAGG

## FIGURE 86

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA175734
><subunit 1 of 1, 261 aa, 1 stop
><MW: 28231, pI: 9.28, NX(S/T): 1
MGGAGILLLLL LAGAGVVAVWRPPK GKCP LRCSCSKDSALCEGSPDLPVFSPTLLSLSLV
RTGVTQLKAGSFLRIPSLHLLLF TSNFSVIEDDAFAGLSHLQYLFIEDNEIGSISKNAL
RGLRSLTHLSLANNHLETLP RFLFRGLDTLTHVDLRGNPFQCDCRVLWLLQWMPTVNASV
GTGACAGPASLSHMQ LHHLDPKTFK CRAIGGGLSRWGGREI WGKGCQGEARLT PCPAI
SRSGKTL SKQHCLPEPQF SHL
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-19

**N-glycosylation site:**

Amino acids 177-181

**N-myristoylation sites:**

Amino acids 15-21;181-186;210-215

**Amidation site:**

Amino acids 217-220

**Microbodies C-terminal targeting signal:**

Amino acids 259-262

**ATP/GTP-binding site motif A (P-loop):**

Amino acids 239-246

**Leucine zipper pattern:**

Amino acids 129-150

**Leucine Rich Repeat:**

Amino acids 53-76; 149-171

**Leucine rich repeat C-terminal domain:**

Amino acids 158-207

**FIGURE 87**

CGGACGCGTGGGGCGGCGAGAGCAGCTGCAGTTCGCATCTCAGGCAGTACCTAGAGGAGCTG  
 CCGGTGCCTCCTCAGAACATCTCCTGATCGCTACCCAGGACCAGGCACCAAGGACAGGGAGT  
 CCCAGGCGCACACCCCCATTCTGGGTCCCCAGGCCAGACCCCCACTCTGCCACAGGTTG  
 CATCTTGACCTGGTCCTCCTGCAGAAGTGGCCCCGTGGTCTGCTCTGAGACTCGTCCCTG  
 GCGCCCCCTGCAGCCCCCTTTCTATGACTCCATCTGGATTTGGCTGGCTGTGGGGACGCGGT  
 CGAGGGGCGGCTGGCTCTCAGCGTGGTGGCAGCCAGCTCTCTGGCCACCATGGCAAATGCT  
 GAGATCTGAGGGGACAAGGCTCTACAGCCTCAGCCAGGGGCACTCAGCTGTTGCAGGGTGTG  
**ATG**GAGAACAAGCTATGTACCTACACACCGTCAGCGACTGTGACACCAGCTCCATCTGTGA  
 GGATTCCTTTGATGGCAGGAGCCTGTCCAAGCTGAACCTGTGTGAGGATGGTCCATGTCACA  
 AACGGCGGGCAAGCATCTGCTGTACCCAGCTGGGGTCCCTGTGCGCCCTGAAGCATGCTGTC  
 CTGGGGCTCTACCTGCTGGTCTTCCCTGATTCTTGTGGGCATCTTCATCTTAGCAGGGCCACC  
 GGGACCCAAAGGTGATCAGGGGGATGAAGGAAAGGAAGGCAGGCCTGGCATCCCTGGATTGC  
 CTGGACTTCGAGGTCTGCCCGGGGAGAGAGGTACCCAGGATTGCCCGGGCCCAAGGGCGAT  
 GATGGGAAGCTGGGGCCACAGGACCAATGGGCATGCGTGGGTTCAAAGGTGACCGAGGCC  
 AAAAGGAGAGAAAGGAGAGAAAGGAGACAGAGCTGGGGATGCCAGTGGCGTGGAGGCCCCGA  
 TGATGATCCGCTGGTGAATGGCTCAGGTCCGCACGAGGGCCGCGTGGAAGTGTACCACGAC  
 CGGCGCTGGGGCACCGTGTGTGACGACGGCTGGGACAAGAAGGACGGAGACGTGGTGTGCCG  
 CATGCTCGGCTTCCGCGGTGTGGAGGAGGTGTACCGCACAGCTCGATTCCGGCAAGGCACTG  
 GGAGGATCTGGATGGATGACGTTGCCCTGCAAGGGCACAGAGGAAACCATCTTCCGCTGCAGC  
 TTCTCAAATGGGGGGTGACAAACTGTGGACATGCCGAAGATGCCAGCGTGACATGCAACAG  
 ACAC**TGA**AAGTGGGCAGAGCCCAAGTTCGGGGTCTGCACAGAGCACCCCTTGCTGCATCCCT  
 GGGGTGGGGCACAGCTCGGGGCCACCCTGACCATGCCTCGACCACACCCCGTCCAGCATTCT  
 CAGTCTCACACCTGCATCCCAGGACCGTGGGGGCCGGTCTGTCATTTCCCTCTTGAACATGT  
 GCTCCGAAGTATAACTCTGGGACCTACTGCCCGTCTCTCTTCCACCAGGTTCCCTGCATGA  
 GGAGCCCTGATCAACTGGATCACCACTTTGCCAGCCTCTGAACACCATGCACCAGGCCTCA  
 ATATCCAGTTCCCTTTGGCCTTTTAGTTACAGGTGAATGCTGAGAATGTGTCAGAGACAAG  
 TGCAGCAGCAGCGATGGTTGGTAGTATAGATCATTTACTCTTCAGACAATCCCAAACCTCC  
 ATTAGTCCAAGAGTTTCTACATCTTCCCTCCCCAGCAAGAGGCAACGTCAAGTGATGAATTC  
 CCCCCTTACTCTGCCTCTGCTCCCCATTTGCTAGTTTGGAGGAAGTACATAGAGGAGAAGC  
 CAGCTGTAGGGGCAAGAGGGAAATGCAAGTCACCTGCAGGAATCCAGCTAGATTTGGAGAAG  
 GGAATGAACTAACATGAATGACTACCATGGCACGCTAAATAGTATCTTGGGTGCCAAATTC  
 TGTATCCACTTAGCTGCATTGGTCCAGGGCATGTGAGTCTGGATAACGCTTACCTTCAGGT  
 AGCACTTAAGTGGTCCATTACCTAGACTGCAAGTAAGAAGACAAAATGACTGAGACCGTGT  
 GCCCACCTGAACTTATTGTCTTTACTTGGCCTGAGCTAAAAGCTTGGGTGCAGGACCTGTGT  
 AACTAGAAAGTTGCCTACTTCAGAACCTCCAGGGCGTGAGTGCAAGGTCAAACATGACTGGC  
 TTCCAGGCCGACCATCAATGTAGGAGGAGAGCTGATGTGGAGGGTGACATGGGGGCTGCCCA  
 TGTTAAACCTGAGTCCAGTGCTCTGGCATTGGGCAGTCACGGTTAAAGCCAAGTCATGTGTG  
 TCTCAGCTGTTTGGAGGTGATGATTTTGCATCTTCCAAGCCTCTTCAGGTGTGAATCTGTGG  
 TCAGGAAAACACAAGTCCTAATGGAACCCTTAGGGGGGAAGGAAATGAAGATCCCTATAAC  
 CTCTGGGGGTGGGGAGTAGGAATAAGGGGCCCTTGGGCCCTCATAAATCTGCAATCTGCACC  
 TCCTCCTAGAGACAGGGAGATCGTGTCTGCTTTTTACATGAGGAGCAGAACTGGGCCATAC  
 ACGTGTTCAGAAGTACGGGAGCTACCTGGTAGCAAGTGAGTGCAGACCCACCTCACCTTGG  
 GGAATCTCAAACCTCATAGGCCTCAGATACACGATCACCTGTCATATCAGGTGAGCACTGGC  
 CTGCTTGGGGAGAGACCTGGGCCCTCCAGGTGTAGGAACAGCAAACTCCTGGCTGACAAC  
 TAAGCCAATATGGCCCTAGGTCACTTCTTGGCTTCCAATATGCTTGGCACTCCTTAAATGTCT  
 AATGATGAGAACTCTCTTTCTGACCAATTGCTATGTTTACATAAACCGCATGTACTCATGC  
 ATCCCTTGGCAGAGCCCATATATGTATGCATATATAAACATAGCACTTTTTACTACATAGCT  
 CAGCACATTGCAAGGTTTGCATTTAAGTT

## FIGURE 88

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA176108
><subunit 1 of 1, 270 aa, 1 stop
><MW: 28871, pI: 7.09, NX(S/T): 1
MENKAMYLHTVSDCDTSSICEDSFDGRSLSKLNLCEDGPCHKRRASICCTQLGSLK
AVLGLYLLVFLILVGFILAGPPGPKGDQDQDEGKEGRPGIPGLPGLRGLPGERGT
PKGDDGKLGATGPMGMRGFKGDRGPKGEKGEKGDAGDASGVEAPMMIRLVNGSGP
VEVYHRRRWGTVCDDGWDKKGDVVCRMLGFRGVEEVYRTARFGQGTGRIWDDVACK
EETIFRCSFSKWGVTNCGHAEDASVTCNRH
```

**Transmembrane domain:**

Amino acids 55-80

**N-glycosylation site:**

Amino acids 172-175

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 43-46

**Tyrosine kinase phosphorylation site:**

Amino acids 212-218

**N-myristoylation sites:**

Amino acids 53-58;224-229;239-244;253-258

**Speract receptor repeated domain signature:**

Amino acids 173-211

**Scavenger receptor cysteine-rich domain:**

Amino acids 171-268

**Collagen Collagen triple helix repeat:**

Amino acids 90-149

**FIGURE 89**

GTCGCCGCGAGGGACGCAGAGAGCACCCCTCCACGCCAGATGCCTGCGTAGTTTTTGTGACC  
 AGTCCGCTCCTGCCTCCCCCTGGGGCAGTAGAGGGGGAGCG**ATG**GAGAAGTGGACTGGCAGG  
 CCCTGGCTGTATCTGCTGCTGCTTCTGTCCCTCCCTCAGCTCTGCTTGGATCAGGAGGTGTT  
 GTCCGGACACTCTCTTCAGACACCTACAGAGGAGGGCCAGGGCCCCGAAGGTGTCTGGGGAC  
 CTGGGTCCAGTGGGCCTCTTGCTCCAGCCCTGCGGGGTGGGGGTGCAGCGCAGGAGCCGG  
 ACATGTCAGCTCCCTACAGTGCAGCTCCACCCGAGTCTGCCCTCCCTCCCCGGCCCCCAAG  
 ACATCCAGAAGCCCTCCTCCCCGGGGCCAGGGTCCAGACCCAGACTTCTCCAGAAACCC  
 TCCCCTTGTACAGGACACAGTCTCGGGGAAGGGGTGGCCCACTTCGAGGTCCCCTTCCCAC  
 CTAGGGAGAGAGGAGACCCAGGAGATTCGAGCGGCCAGGAGGTCCCCTTCGAGACCCCAT  
 CAAGCCAGGAATGTTTCGGTTATGGGAGAGTGCCCTTTGCATTGCCACTGCACCCGGAACCGCA  
 GGCACCCCTCGGAGCCCACCCAGATCTGAGCTGTCCCTGATCTCTTCTAGAGGGGAAGAGGCT  
 ATTCGGTCCCCTACTCCAAGAGCAGAGCCATTCTCCGCAAACGGCAGCCCCCAAAGTGGAGCT  
 CCCTCCCACAGAAGTGTCTGTCCACACCCCATCCCCCAAGCAGAACCTCTAAGCCCTGAAA  
 CTGCTCAGACAGAGGTGGCCCCCAGAACCAGGCCTGCCCCCTACGGCATCACCCAGAGCC  
 CAGGCCTCTGGCACAGAGCCCCCTCACCCACGCACTCCTTAGGAGAAGGTGGCTTCTTCGG  
 TGCATCCCCTCAGCCACGAAGGCCAAGTTCCAGGGTTGGGCCAGTCCCAGGTAGCAGGGA  
 GACGCCCTGATCCTTTTTCTTCGGTCCCTCGGGGCCGAGGCCAGCAGGGCCAAGGGCCTTGG  
 GAAACGGGGGGGACTCCTCACGGGCCCGCCTGGAGCCTGACCCTCAGCACCCGGGCGCCTG  
 GCTGCCCTGCTGAGCAACGGCCCCCATGCCAGCTCCCTCTGGAGCCTCTTTGCTCCCAGTA  
 GCCCTATTCCAAGATGTTCTGGGGAGAGTGAACAGCTAAGAGCCTGCAGCCAAGCGCCCTGC  
 CCCCCTGAGCAGCCAGACCCCCGGGCCCTGCAGTGCAGCAGCTTTAACTCCAGGAATTCATG  
 GGCCAGCTGTATCAGTGGGAGCCCTTCACTGAAGTCCAGGGCTCCCAGCGCTGTGAACTGAA  
 CTGCCGGCCCCGTGGCTTCCGCTTCTATGTCCGTCACTGAAAAGGTCCAGGATGGGACCC  
 TGTGTCAGCCTGGAGCCCTGACATCTGTGTGGCTGGACGCTGTCTGAGCCCCGGCTGTGAT  
 GGGATCCTTGGCTCTGGCAGGCGTCTGATGGCTGTGGAGTCTGTGGGGGTGATGATTCTAC  
 CTGTGCGCTTGTTCGGGGAACCTCACTGACCGAGGGGGCCCCCTGGGCTATCAGAAGATCT  
 TGTGGATTCAGCGGGAGCCTTGC GGCTCCAGATTGCCAGCTCCGGCCTAGCTCCAAGTAC  
 CTGGCACTTCGTGGCCCTGGGGGCCGGTCCATCATCAATGGGAACTGGGCTGTGGATCCCC  
 TGGGTCTACAGGGCCGGCGGGACCGTCTTCGATATAACCGTCTCCCAGGGAGGAGGGCA  
 AAGGGGAGAGTCTGTCGGCTGAAGGCCCCACCACCCAGCCTGTGGATGTCTATATGATCTTT  
 CAGGAGGAAAACCCAGGCGTTTTTTATCAGTATGTCATCTCTTCACCTCCTCCAATCCTTGA  
 GAACCCACCCAGAGCCCCCTGTCCCCAGCTTCAGCCGGAGATTCTGAGGGTGGAGCCCC  
 CACTTGCTCCGGCACCCCGCCAGCCGGACCCAGGCACCCCTCCAGCGTCAGGTGCGGATC  
 CCCAGATGCCCGCCCCGCCCATCCAGGACACCCCTGGGGTCTCCAGCTGCGTACTGGAA  
 ACGAGTGGGACACTCTGCATGCTCAGCGTCTGCGGGAAAGGTGTCTGGCGCCCCATTTTCC  
 TCTGCATCTCCCGTGAGTCGGGAGAGGAAGTGGATGAACGCAGCTGTGCCGCGGGTGCCAGG  
 CCCCAGCCTCCCCTGAACCTGCCACGGCACCCCATGCCCCCATACTGGGAGGCTGGCGA  
 GTGGACATCCTGCAGCCGCTCCTGTGGCCCCGGCACCCAGCACCCGAGCTGCAGTGCCGGC  
 AGGAATTTGGGGGGGGTGGCTCCTCGGTGCCCCGGAGCGCTGTGGACATCTCCCCGGCCC  
 AACATCACCCAGTCTTGCCAGCTGCGCCTCTGTGGCCATTGGGAAGTTGGCTCTCCTTGGAG  
 CCAGTGCTCCGTGCGGTGCGGCCGGGGCCAGAGAAGCCGGCAGGTTCGCTGTGTTGGGAACA  
 ACGGTGATGAAGTGAAGCAGCAGGAGTGTGCGTCAAGCCCCCACAGCCCCCAGCAGAGAG  
 GCCTGTGACATGGGGCCCTGTACTACTGCCTGGTTCCACAGCGACTGGAGCTCCAAGGTGAG  
 CCCGGAACCCAGCCATATCCTGCATCCTGGGTAAACCATGCCAGGACACCTCAGCCTTTC  
 CAGCA**TAG**CTCAATAAACTTGTATTGATC

## FIGURE 90

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA190710
><subunit 1 of 1, 877 aa, 1 stop
><MW: 95132, pI: 8.77, NX(S/T): 5
MENWTGRPWL YLLLLLSLPQLCLDQEVLSGHS LQTPTEEGQGPEGVWGPVWQWASCSQPC
GVGVQRRSRTCQLPTVQLHPSLPLPPRPPRHPEALLPRGQGPRPQTS PETLPLYRTQSRG
RGGPLRGPASHLGREETQEIRAARRSRLRDP I KPGMFGYGRVPFALPLHRNRRHPRSPPR
SELSLISSRGEEAIPSPTPRAE PFSANGSPQTELPPELSVHTPSPQAEPLSPETAQTEV
APRTRPAPLRHH PRAQASGTEPPSPTHSLGEGGFFRAS PQPRRPSSQGWAS PQVAGRRPD
PFPSVPRGRGQQGQGPWGTGGTPHGPRLEPDPQH PGAWLPLLSNGPHASSLWSLFA PSSP
IPRCSGESEQLRACSQAPCPPEQPDPRALQCAAFNSQEFMGQLYQWEPFTEVQGSQRCEL
NCRPRGFRFRFYVRHTEKVDGTL CQPGAPDICVAGRCLSPGCDGILGSGRRPDGCGVCGGD
DSTCRLVSGNLTDRGGPLGYQKILWI PAGALRLQIAQLRPSSNYLALRGPGRS IINGNW
AVDPPGSYRAGGTVFRYNRP PREEGKGESLSAEGPTTQPVDVYMI FQEENPGVFYQYVIS
SPPPILENPTPEPPVPQLQPEILRVEPPLAPAPR PARTPGTLQRQVRI PQMPAPPHPRT P
LGSPAAYWKRVGHSACSASC GKGVWRPIFLCISRESGEELDERSCAAGARPPASPEPCHG
TPCPPYWEAGEWTSCSRSCGPGTQHRQLQCRQEF GGGSSVPPERCGHLPRPNITQSCQL
RLCGHWEVGS PWSQCSVRCGRGQRSRQVRCVGNNGDEVSEQECASGPPQPPSREACDMGP
CTTAWFHSDWSSKVSPEPPAISCILGNHAQDTS AFPA
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-24

**N-glycosylation sites:**

Amino acids 3-6; 490-493; 773-776

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 282-285

**N-myristoylation sites:**

Amino acids 208-213; 414-419; 463-468; 473-478; 475-480;  
478-483; 495-500; 546-551; 662-667; 755-760;  
756-761; 789-794

**Amidation sites:**

Amino acids 295-298; 467-470

**Leucine zipper pattern:**

Amino acids 504-526

**VWFC domain proteins:**

Amino acids 53-67; 732-746; 792-806

**Thrombospondin type 1 domain:**

Amino acids 48-87; 727-783; 787-841

**FIGURE 91**

CGAGTATTTTCCCACCATCTCCAGCCGAAACTGACCAAGAACTCTGAGGCGGATGGC**ATGT**  
TCGCGTACGTCTCCATGATGAGTTCGTGGCCTCGATGATTAAGATCCCTTCGGACACCTTC  
ACCATCATCCCTGACTTTGATATCTACTATGTCTATGGTTTTAGCAGTGGCAACTTTGTCTA  
CTTTTGGACCCTCCAACCTGAGATGGTGTCTCCACCAGGCTCCACCACCAAGGAGCAGGTGT  
ATACATCCAAGCTCGTGAGGCTTTGCAAGGAGGACACAGCCTTCAACTCCTATGTAGAGGTG  
CCCATTGGCTGTGAGCGCAGTGGGGTGGAGTACCGCCTGCTGCAGGCTGCCTACCTGTCCAA  
AGCGGGGGCCGTGCTTGGCAGGACCCTTGGAGTCCATCCAGATGATGACCTGCTCTTCACCG  
TCTTCTCCAAGGGCCAGAAGCGGAAAATGAAATCCCTGGATGAGTCCGGCCCTGTGCATCTTC  
ATCTTGAAGCAGATAAATGACCCGATTAAGGAGCGGCTGCAGTCTTGTACCAGGGGCGAGGG  
CACGCTGGACCTGGCCTGGCTCAAGGTGAAGGACATCCCCTGCAGCAGTGCCTCTTAACCA  
TTGACGATAACTTCTGTGGCCTGGACATGAATGCTCCCCTGGGAGTGTCCGACATGGTGCCT  
GGAATCCCCTCTTACGGAGGACAGGGACCGCATGACGTCTGTGCATCGCATATGTCTACAA  
GAACCACTCTCTGGCCTTTGTGGGCACCAAAAGTGGCAAGCTGAAGAAGGTGCCTGGTACCA  
GCCTCTGCCCTACCCTTGAGCTACAGACGGGACCCCGATCCCACAGAGCAACAGTGACTCTG  
GAACTCCTGTTCTCCAGCTGTTCAATCAAACT**TGAG**AAAAAATTCAGAGCTGTGTAGGCTTATT  
TAGTGTGTTGTCAGCCTTGGATATTGAAAAATGGAAACAGATGAGACACATCTACCTCCCTG  
TGACCCAGCCATACATCATAGCTCATGTCTGCCACCCCAAGTCTTAGGGAAAAAAGACT  
TTGGAGAATGTGTCTCTGCTTAGCTTGGCTAGGTAGTTGGTCTCTTTTCTCTGCCCAAGCG  
TCCCCTGGGTAATTTTGGACAATGGAGTGTAGGCATGTTTGACTCTTGTGGTGTATCACTT  
GTATATGTCAGTGAACTAACTGATTCTCCCATCGGAATATAGTTATCTCTTGGGCCTGATA  
TATGGTAGGATAACCTTATGCTCATCTGTCCACTTCTGCAGCCAAGTGCCTGGCCAGTGTG  
TG  
TGCATACACAGGGCAGAGAGGATGGAGCCACCGTACTGCAGCATCATGTAATTAACCTCAGT  
GCTCAGAACCATCCCAGCCTCTGCGGGAAAGAGAAAAAGTAAGCCAACAGTGCCTGATGAGCT  
GATCATATGTGCAAAGCTCTGTTGGCATCTGGTCCAGGAGAGCACCCAAAAAAGTTAATT  
GGTGTGTGTCAGTCTCCTTTCCCTTAAGACTATGGTTACAACAAAGCGTGAGCAGTGTCTCCT  
GCATGGCCACTATCCAGCACAAATCCATAATTCCCCCATAGAGCCGGTGGGGAGGAGGAGGT  
GAGTGGCGAAGGAAGTGGAAACACTTGGTGTGTCATGTGCTCCTATCATTTCTACTAGCTTACT  
GGGAAATAAAGTGTAGTCAAGAGTGTATGAAGGCAAGATGTAAAATTAGCGACTGGTGCTAA  
TCTGGTTACTTGAAAACAAGTGAAGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGT  
CTAAACCTCGTATAGTTCCCTGGAGGATATACAACAGTGTAAATTCCTTTAGGGTGTGCCACA  
GGTTCCCTGGCCTGTGGGAGGGAATGAATCAGGAGGGCTCTTGAGAACCTTCATCTGTGTGCT  
TGCAGTGAAGTGAAGTCCCAAAGCTGGAGATTTAGTGAGAGCAGGCAACCCCTCTGTGTCTC  
ACTGTCCATATTCTGGAGGCAGAGGTTTGTAAACAGGCCATGTGCACCTGCATAGGGATGGGT  
AAAGCAAGGACTTTGAAAGAGTTGAAAAGCATTATAAACAGTTGTTTCAAGAAATACGTCCCAG  
GAGTTCATGTGAACTGGCTCTGTGTGCATTGAAGCATGGCTGTTGGGAATTCTAACTGGT  
CCAACACTCCTGCAAAACAATGTGTAATAATTTAGGAAGAACTTGAAAATAGTCAAATCCT  
TTGAACTGGTGACAATTTTTTAAAGAATCAATTCATTTGTTTCAAGGGTAATAATCACCA  
AGATACACATTTTCAAGATTTATTTAGTCTATCAAAAATTGGAATTGATATATACACTCATTT  
ATAGGAGAATGGTTAGGTAGATTTGGTATATTTATGTAGTCATTGAAAACCTTAGTTTATAAA  
GGCCAATCTTGTAACCTGATCCTTGTGTGATAACATTCAGTGAAAAAGCATGAGACAATTAGA  
AAGCATGATACAATGAATAAAATAAAAATGGAAAGAGAACCATCAAATGCTAA



## **FIGURE 92**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA190803
><subunit 1 of 1, 280 aa, 1 stop
><MW: 31222, pI: 7.40, NX(S/T): 1
MFAYVFHDEFVASMIIKIPSDTFTIIPDFDIYYVYGFSSGNFVYFLTLQPEMVSPPGSTTK
EQVYTSKLVRLCKEDTAFNSYVEVPIGCERSGVEYRLLQAAYLSKAGAVLGRTLGVHPDD
DLLFTVFSKQGQRKMKSLDESALCIFILKQINDRIKERLQSCYRGEGTLDLAWLKVKDIP
CSSALLTIDDNFCGLDMNAPLGVSDMVRGIPVFTEDRDRMTSVIAYVYKNHSLAFVGTKS
GKLKKVPGTSLCPTLELQTGPRSHRATVTLELLFSSCSSN
```

### **Important features of the protein:**

#### **N-glycosylation site:**

Amino acids 230-233

#### **N-myristoylation sites:**

Amino acids 87-92;107-112;194-199;237-242

**FIGURE 93**

CCTTATCAGACAAAGGACGAGATGGAAAATACAAGATAATTTACAGTGGAGAAGAATTAGAA  
TGTAACCTGAAAGATCTTAGACCAGCAACAGATTATCATGTGAGGGTGTATGCC**ATGT**TACAA  
TTCCGTAAAGGGATCCTGCTCCGAGCCTGTTAGCTTCACCACCCACAGCTGTGCACCCGAGT  
GTCCTTTCCCCCTAAGCTGGCACATAGGAGCAAAAGTTCACTAACCCTGCAGTGGAAGGCA  
CCAATTGACAACGGTTCAAAAATCACCACCTACCTTTTAGAGTGGGATGAGGGAAAAAGAAA  
TAGTGGTTTCAGACAGTGTCTTTCGGGAGCCAGAAGCACTGCAAGTTGACAAAGCTTTGTC  
CGGCAATGGGGTACACATTCAGGCTGGCCGCTCGAAACGACATTGGCACCAGTGGTTATAGC  
CAAGAGGTGGTGTGCTACACATTAGGAAATATCCCTCAGATGCCTTCTGCACCTAAGGCTGGT  
TCGAGCTGGCATCACATGGGTCACGTTGCAGTGGAGTAAGCCAGAAGGCTGTTACCCCGAGG  
AAGTGATCACCTACACCTTGGAATTCAGGAGGATGAAAATGATAACCTTTTCCACCCAAAA  
TACACTGGAGAGGATTTAACCTGTACTGTGAAAAATCTCAAAGAAGCACACAGTATAAATT  
CAGGCTGACTGCTTCTAATACGGAAGGAAAAAGCTGTCCAAGCGAAGTTCTTGTGGTGTACGA  
CGAGTCCCTGACAGGCCTGGACCTCCTACCAGACCGCTTGTCAAAGGCCAGTTACATCTCAT  
GGCTTTAGTGTCAAATGGGATCCCCCTAAGGACAATGGTGGTTCAGAAATCCTCAAGTACTT  
GCTAGAGATTACTGATGGAAATCTGAAGCGAATCAGTGGGAAGTGGCCTACAGTGGGTCCG  
CTACCGAATACACCTTCACCCACTTGAAACCAGGCCTTTGTACAAACTCCGAGCATGCTGC  
ATCAGTACCGGCGGACACAGCCAGTGTCTGAAAGTCTCCCTGTTTCGCACACTAAGCATTGC  
ACCAGGTCAATGTGACCACCGAGGGTTTTGGGTAGACCAAAGCACAAAGAAGTCCACTTAG  
AGTGGGATGTTCTGCATCGGAAAGTGGCTGTGAGGTCTCAGAGTACAGCGTGGAGATGACG  
GAGCCCGAAGACGTAGCCTCGGAAGTGTACCATGGCCAGAGCTGGAGTGCACCGTCCGGCAA  
CCTGCTTCCCTGGAACCGTGTATCGCTTCCGGGTGAGGGCTCTGAATGATGGAGGGTATGGTC  
CCTATTCTGATGTCTCAGAAATTAACACTGCTGCAGGGCCTCCTGGACAATGCAAAGCACCT  
TGTATTTCTTGTACACCTGATGGATGTGTCTTAGTGGGTTGGGAGAGTCCCTGATAGTTCTGG  
TGCTGACATCTCAGAGTACAGGTTGGAATGGGGAGAAGATGAAGAATCCTTAGAACTCATTT  
ATCATGGGACAGACACCCGTTTTGAAATAAGAGACCTGTTGCCTGCTGCACAGTATTGCTGT  
AGACTACAGGCCTTCAATCAAGCAGGGGCAGGGCCGTACAGTGAAGTGTCTTGTCCCTTCCAGAC  
GCCAGCGTCTGCCCTGACCCCGTCTCCACTCTCTGTGTCTGGAGGAGGAGCCCTTGTATGCC  
TACCCTGATTCACCTTCTGCGTGCCTTGTACTGAAGTGGGAAGAGCCGTGCAATAACGGATC  
TGAAATCCTTGCTTACACCATGATCTAGGAGACACTAGCATTACCGTGGGCAACACCACCA  
TGCATGTTATGAAAGATCTCCTTCCAGAAACCACCTACCGGATCAGAATTCAGGCTATAAAT  
GAAATTGGAGCTGGACCATTTAGTCAGTTCATTAAGCAAAAACCTCGGCCATTACCACCTT  
GCCTCCTAGGCTAGAATGTGCTGCTGCTGGTCTCAGAGCCTGAAGCTAAAATGGGGAGACA  
GTAACCTCAAGACACATGCTGCTGAGGACATTGTGTACACACTACAGCTGGAGGACAGAAAC  
AAGAGGTTTATTTCAATCTACAGAGGACCCAGCCACACCTACAAGGTCAGAGACTGACGGA  
ATTCACATGCTACTCCTTCAGAATCCAGGCAGCAAGCGAGGCTGGAGAAGGGCCCTTCTCAG  
AAACCTATACCTTCAGCACAAACAAAAGTGTCCCCCACCATCAAAGCACCTCGAGTAACA  
CAGTTAGAAGTAAATTCATGTGAAATTTTATGGGAGACGGTACCATCAATGAAAGGTGACCC  
TGTTAACTACATCTGCAGGTATTGGTTGGAAGAGAATCTGAGTACAAACAGGTGTACAAGG  
GAGAAGAAGCCACATTCCAAATCTCAGGCCTCCAGACCAACACAGACTACAGGTTCCGCGTA  
TGTGCGTGTGCTGCTGTTTAGACACCTCTCAGGAGCTAAGCGGAGCCTTCCAGCCCTCTGC  
GGCTTTTGTATTACAACGAAGTGAGGTCATGCTTACAGGGGACATGGGGAGCTTAGATGATC  
CCAAAATGAAGAGCATGATGCCTACTGATGAACAGTTTGCAGCCATCATTTGTGCTTGGCTTT  
GCAACTTTGTCCATTTTATTTGCCTTTATATTACAGTACTTCTTAATGAAG**TAA**ACCCAACA  
AAACTAGAGGTATGAATTAATGCTACACATTTTAAATACACACATTTATTCAGATACTCCCCT  
TTTTAAAGCCCTTTTGTTTTTTGTATTTATATACTCTGTTTTACAGATTTAGCTAGAAAAAA  
ATGTCAGTGTTTGGTGCACCTTTTTGAAATGCAAACCTAGGAAAAGGTTAACTGGATTTT  
TTTTTAAAAA

## FIGURE 94

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA191064
><subunit 1 of 1, 847 aa, 1 stop
><MW: 93607, pI: 5.33, NX(S/T): 3
MYNSVKGSCSEPVSFTHSCAPECPFPKLAHRKSSSLTLQWKAPIDNGSKITNYLLEWD
EGKRNSGFRQCFFGSQKHCKLTKLCPAMGYTFRLAARNDIGTSGYSQEVVVCYTLGNIPQM
PSALRLVRAGITWVTLQWSKPEGCSPEEVITYTLEIQEDENDNLFHPKYTGEDLTCTVKN
LKRSTQYKFRLTASNTEGKSCPSEVLVCTTSPDRPGPPTPLVKGPVTSHGFSVKWDPK
DNGGSEILKYLLEITDGNSEANQWEVAYSGSATEYTFTHLKPGLYKLRACCISTGGHSQ
CSESLPVRTLSIAPGQCRPPRVLGRPKHKEVHLEWDVPASESGCEVSEYSVEMTEPEDVA
SEVYHGPELECTVGNLLPGTVYRFRVRALNDGGYGPYSDVSEITTAAGPPGQCKAPCISC
TPDGCVLVWESPSSGADISEYRLEWGEDEESLELIYHGTDRFEIRDLLPAAQYCCRL
QAFNQAGAGPYSELVLCQTPASAPDPVSTLCVLEEEPLDAYPDSPSACLVLNWEEPCNNG
SEILAYTIDLGDTSITVGNTTMHVMKDLLPETTYRIRIQAINEIGAGPFSQFIKAKTRPL
PPLPPRLECAAAGPQSLKCLKWGSNSKTHAAEDIVYTLQLEDNRKRFISYRGPSTYK
QRLTEFTCYSFRIQAASEAGEGPFSEYTFSTTKSVPPTIKAPRVTQLEVNSCEILWETV
PSMKGDPVNYILQVLVGRESEYKQVYKGEATFQISGLQTNTDYRFRVCACRRCLDTSQE
LSGAFSPSAAFVLRSEVMLTGMGSLDDPKMKSMPTDEQFAAIVLGFATLSILFAFI
LQYFLMK
```

**Important features of the protein:**

**Transmembrane domain:**

Amino acids 823-843

**N-glycosylation sites:**

Amino acids 48-51;539-542;559-562

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 63-66;182-185

**Tyrosine kinase phosphorylation sites:**

Amino acids 387-394;662-669

**N-myristoylation sites:**

Amino acids 49-54;257-262;343-348;437-442;757-762

**Amidation site:**

Amino acids 61-64

**ATP/GTP-binding site motif A (P-loop):**

Amino acids 193-200

**Fibronectin type III domain:**

Amino acids 22-106;118-203;215-302;314-398;  
410-492;504-590;601-685;697-778

**FIGURE 95A**

CAATTCGGCCTCGCTCCTTGTGATTGCGCTAAACCTTCCGTCCTCAGCTGAGAACGCTCCACCACCTCCCCGGA  
TCGCTCATCTCTTGGCTGCCCTCCCCTGTTCTCTGATGTTATTTTACTCCCCGATCCCCACTCGTTCTTCC  
AATTCGTAGGTGAGTGTTCCAGCTGGCTGCCCTGTGTCTCTTGGATGCCCTGTGGCTTCAGTCCGTCTC  
CTGTTGCCACCACCTCGTCCCTGGGCCGCTGATACCCAGCCCAACAGCTAAGGTGTGGATGGACAGTAGGG  
GGCTGGCTTCTCTACTGGTCAGGGGTCTTCTCCCCTGTCTGCCCTCCCGAGCTAGGACTGCAGAGGGGCTTAT  
CATGGTGTGTCAGGCCCTGGCTGTCTCGCTGTTGCTGCCAGCCTCACACTGCTGGTGTCCACCTCTCCA  
GCTCCCAGGATGTCTCCAGTGAGCCAGCAGTGAGCAGCAGCTGTGCGCCCTTAGCAAGCACCCACCGTGGCC  
TTTGAAGACCTGCAGCCGTGGGTCTCTAACTTCACTACCTGGAGCCCGGGATTTCTCCCAGCTGGCTTGGGA  
CCCCCGGGAACCTGACTCATCGTGGGAGCCAGAACTACTCTTCCAGACTCAGCCTTCCCAATGTCTCTCTC  
TTCAGGCCACAGAGTGGGCCCTCCAGTGAGGACACGCGCCGCTCCTGCCAAAGCAAAGGAAAGACTGAGGAGGAG  
TGTGAGAACTACGTGCGAGTCTGATCGTCCGCGCCGGAAGGTGTTTATGTGTGGAACCAATGCCTTTTCCCC  
CATGTGCACCAGCAGACAGGTGGGGAACCTCAGCCGGACTATTGAGAAGATCAATGGTGTGGCCCGCTGCCCT  
ATGACCCACGCCACAACCTCCACAGCTGTATCTCTCCCAGGGGAGCTCTATGCAGCCACGGTTCATCGACTTC  
TCAGGTCCGGACCTGCCATCTACCGCAGCCTGGGCAGTGGGCCACCGCTTCCGACTGCCCAATATAACTCCAAG  
TGGCTTAATGAGCCAACTTCGTGGCAGCCTATGATATTGGGCTGTTGCATACTTCTTCTGCGGGAGAACGC  
AGTGGAGCAGCAGTGGACCCGACCCGAGCCGTGTACTCTCGCTGGCCCGCTGTGCAAGAAATGACGTGGGGCCGAT  
TCCTGTGAGGACACATGGACCACATTCATGAAGGCCGGCTCAACTGCTCCCGCCGGGCGAGGTCCCCCTC  
TACTATAACGAGCTGCAGAGTGCCTTCCACTTCCCGGAGCAGGACCTCATCTATGGAGTTTTTACAACCAACGT  
AAACAGCATCGCGGCTTCTGCTGTCTGCCCTCAACCTCAGTGCTATCTCCCAGCTTTCAATGGCCCATTTTC  
GCTACCAGGAGAACCCAGGGCTGCCTGGCTCCCCATAGCCAACCCCATCCCCAATTTCCAGTGTGGCACCCCTG  
CCTGAGACCCGTTCCCAACGAGAACCTGACGGAGCGCAGCCTGCAGGACGCGCAGCGCCTTCTCTGATGAGCGA  
GGCCGTGACCCGCTGGACCCGAGCCCTGTGTCACTACCCAGGACCGCTGCGCTTCTCACACCTCGTGGTGGACC  
TGGTGCAGGCTAAAGACACGCTCTACCATGTACTCTACATTGGCACCGAGTCCGGCACCATCCTGAAGGCGCTG  
TCCACGGCGAGCCGAGCCTCCACGGCTGTACTTGGAGGAGTGCACGTGCTGCCCCCGGGCGCCGCGAGCC  
CCTGCGCAGCCTGCGCATCCTGCACAGCGCCGCGCGCTTCTCGTGGGGCTGAGAGACGGCGTCTGCGGGTCC  
CACTGGAGAGGTGCGCCGCTACCGCAGCCAGGGGGCATGCCTGGGGGCCCGGGACCCGTAAGTGTGGCTGGGAC  
GGGAAGCAGCAACGTTGCAGCACACTCGAGGACAGCTCCAACATGAGCCTCTGGACCCAGAACATCACCCGCTG  
TCTGTGCGGAATGTGACACGGGATGGGGCTTCCGCCCCGCTACCATGGCAACCATGCTGAGCATTGGATG  
GGGACAACCTCAGGCTCTTGCCTGTGTGAGCTCGATCCTGTGATTCCCCTCGACCCCGCTGTGGGGCTTGGAC  
TGCTGGGGCCAGCCATCCACATCGCCAACCTGCTCCAGGAATGGGGCTGGACCCCGTGGTTCATCGTGGGCGCT  
GTGCAGCACGTCCTGTGGCATCGGCTTCCAGGTCCGCCAGCGAAGTTGCAGCAACCCGTGCTCCCCGCCACGGGGC  
CGCATCTTCTGTGGGCAAGAGCCGGGAGGAACGGTCTGTAATGAGAACACGCCTTCCCCGGTGGCCATCTTCTG  
GGCTTCTGGGGCTCCTGGAGCAAGTGCAGCAGCAACTGTGGAGGGGGCATGCAGTCCGCGGCTCGGGCCCTGCG  
AGAACGGCAACTCCTGCCTGGGCTGCGGCGAGTTCAGACGCTGCAACCCCGAGGGCTGCCCGAAGTGGGCGC  
AACACCCCTGGACGCGTGGCTGCCCGTGAACGTGACGCGAGGGCGGGGCACGGCAGGAGCAGCGGTTCCGCTT  
CACCTGCCGCGGCCCTTGCAGACCCGACGGCTGCAGTTCGGCAGGAGAAGGACCGAGACGAGGACCTGTC  
CCGCGGACGGCTCCGGCTCCTGGACACCCGACGCCCTGGTGGAGTCTCCTGCGCAGCGGGAGCACCTCCCCG  
CACACGGTGTGAGCGGGGGCTGGGCCGCTGGGGCCCGTGGTGTCTGCTCCCAGGACTGCGAGCTGGGCTTCCG  
CGTCCGCAAGAGAACGTGACTAACCCTGGAGCCCGCAACGGGGGCTGCCCTGCGTGGGCGATGCTGCCGAGT  
ACCAGACTGCAACCCCGAGCTTGGCCAGTTCGGGGTGGTGGTCTGCTGGACTCATGGTCTCATGCTCA  
GCTTCTGTGGTGGGGTCACTATCAACGCAACCGTCTCCTGCACACCCCGCACCTCCCCAGTGGAGACAT  
CTGTCTCGGGCTGCACACGGAGGAGGACTATGTGCCACACAGGCTGCCAGGCTGGTCCGCTTGGTCTGAGT  
GGAGTAAGTGCAGTGCAGACGGAGCCAGAGCCGAAGCCGGCAGTGTGAGGAGCTCCTCCCAGGGTCCAGCGC  
TGTGCTGGAAACAGCAGCCAGAGCCGCCCTGCCCTACAGCGAGATTCCCGTATCCTGCCAGCCTCCAGCAT  
GGAGGAGGCCACCGACTGTGCAGGTAAGAAACCGGACCTACCTCATGCTGCGGTCTCCCAGCCCTCCAGCA  
CCCCACTCCAAAGTCTGGACTTCTCCACATCCTGCTCCAGACAGCAAGCTTTGTTGGGGTCCCCACTGCTTT  
GAGATGGGTTCAATCTCATCCACTTGGTGGCCACGGGCATCTCCTGCTTCTTGGGCTCTGGGCTCCTGACCCTA  
GCAGTGTACTGTCTTGCCAGCACTGCCAGCTCAGTCCCAGGAGTCCACACTGGTCCATCCTGCCACCCCAACC  
ATTTGCACTACAAGGGCGGAGGCACCCGAAGAATGAAAAGTACACACCCATGGAATTCAGACCCCTGAACAAG  
AATAACTTCATCCCTGATGACAGAGCCAACTTCTACCCATTGCAGCAGACCAATGTGTACACGACTACTTACTA  
CCCAAGCCCCCTGAACAAACACAGCTTCCGGCCCGAGGCTCACCTGGACAACGGTGTCTCCCCAACAGCTGAT  
ACCGCCCTTCTGGGACTTGGGCTTCTTGCCTTATAAGGCACAGAGCAGATGGAGATGGGACAGTGGAGCCAG  
TTTGGTTTTCTCCCTGCACTAGGCCAAGAATTGCTGCCTTGCCTGTGGGGGCTCCCCTCCGGTTCAGAGA  
GCTCTGGCTGGCATTGACCATGGGGGAAAGGCTGGTTTTAGGCTGACATATGGCCGAGGTCCAGTTCAGCCC  
AGGTCTCTCATGGTTATCTTCCAACCCACTGTCAGCTGACACTATGCTGCCATGCCTGGGCTGTGGACCTACT  
GGGCATTTGAGGAATTGGAGAATGGAGATGGCAAGAGGGCAGGCTTTTAAAGTTTGGGTTGGAGACAACCTCCTG  
TGGCCCCACAAGCTGAGTCTGGCTTCTCCAGCTGGCCCCAAAAGGCTTTTGTACATCTGATATATCTCT  
GAAAGTAATCAATCAAGTGGCTCCAGTAGCTCTGGATTTTCTGCCAGGGCTGGGCCATTGTGGTGTGCCCCAG  
TATGACATGGGACCAAGGCCAGCGCAGGTATCCACCTCTGCCTGGAAGTCTATACTTACCAGGGCATCCCT  
CTGGTCAGAGCAGTGTACTGGGAAGTGGAGGCTGACCTGTGCTTAGAAGTCTTTAATCTGGGCTGTGTACA  
GGCCTCAGCTTGGCTCAATGCACGAAAGGTGGCCAGGAGAGAGGATCAATGCCATAGGAGGCAGAAGTCTG  
GCCTCTGTGCCTCTATGGAGACTATCTTCCAGTTGCTGCTCAACAGAGTTGTTGGCTGAGACCTGCTTGGGAGT

**FIGURE 95B**

CTCTGCTGGCCCTTCATCTGTTTCAGGAACACACACACACACACTCACACACGCACACACAATCACAATTTGC  
TACAGCAACAAAAAAGACATTGGGCTGTGGCATTATTAATTAAAGATGATATCCAGTC

## **FIGURE 96**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA194909
><subunit 1 of 1, 1092 aa, 1 stop
><MW: 119324, pI: 8.13, NX(S/T): 14
MPCGFSPSPVAHHLVPGPPDTPAQQLRCGWTVGGWLLSLVRGLLPCLPPGARTAE GPIMV
LAGPLAVSLLLPSLTLVSHLSSSQDVSSEPSSEQQLCALSKHPTVAFEDLQPWVSNFTY
PGARDFSQLALDPSGNQLIVGARNYLFRLSLANVSLLOATEWASSEDTRRSCQSKGKTEE
ECQNYVRVLIVAGRKVFMCGTNAFSPMCTSRQVGNLSRTIEKINGVARCPYDPRHNSTAV
ISSQGELYAATVIDFSGRDPAIYRSLGSGPPLRTAQYNSKWLNEPNFVAAYDIGLFAYFF
LRENAVEHDCGRTVYSRVARVCKNDVGGFLEDTWTFMFKARLNCSRPGEVVPFYNELQ
SAFHLPEQDLIYGVFTTNVNSIAASAVCAFNLSAISQAFNGPFRYQENPRAAWLP IANPI
PNFQCGLTPE TGPENLTERSLQDAQRLFLMSEAVQPVTPEPCVTDQSVRFSHLVVDLVQ
AKDTLYHVLYIGTESGITL KALSTASRSLHGCYLEELHVLPPGRREPLRSLRILHSARAL
FVGLRDGVLRVPLERCAAYRSQGA CLGARDPYCGWDGKQQRCTLEDSSNMSLWTQNITA
CPVRNVTRDGGFGPWPSPWQ PCEHLGDNSGSCLCRARSCDSPRPRCGGLDCLGPAIHIAN
CSRNGAWTPWSSWALCSTSCGIGFQVRQRSCSNPAPRHGGRI FVGKSREERFCNENTPCP
VPIFWASWGSWSKCSSNCGGMQSRRRACENGSCLGCGEFKTCNPEGCEVRRNTPWTP
WLPVNVTOGGARQEQRFRFTCRAPLADPHGLQFGRRRRETETRTCPADGSGSCD TDALVEVL
LRSGSTSPHTVSGGWAAWGPWSSCSRDCELGFRVRKRTCTNPEPRNGGLPCVGDAAEYQD
CNPQACPVRGAWSCWTSWSPCSASC GGGHYQRTRSC TSPAPSPGEDICLGLHTEEALCAT
QACPWSPWSEWSKCTDDGAQSR SRHCEELLPGSSACAGNSSQSRPCPYSEIPVILPASS
MEEATDCAGKRNRTYLM LRS SQPSSTPLQSLDSFHILLQTAKLCWGP HCFEMGSI SSTWW
PRASPASWALGS
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-42

**Transmembrane domain:**

Amino acids 56-79;373-395

**N-glycosylation sites:**

Amino acids 117-120;153-156;215-218;236-239;345-348;391-394;  
436-439;590-593;597-600;605-608;660-663;785-788;  
1000-1003;1032-1035

**cAMP- and cGMP-dependent protein kinase phosphorylation sites:**

Amino acids 773-776;815-818;875-878

**Tyrosine kinase phosphorylation site:**

Amino acids 177-185;348-355

**N-myristoylation sites:**

Amino acids 42-47;50-55;373-378;492-497;543-548;563-568;  
630-635;647-652;740-745;810-815;827-832;829-834;  
853-858;887-892;910-915;993-998;1073-1078

**Amidation sites:**

Amino acids 192-195;522-525;813-816;1028-1031

**ATP/GTP-binding site motif A (P-loop):**

Amino acids 700-707

**Cytochrome c oxidase subunit II, copper A binding region signature:**

Amino acids 921-929

**Growth factor and cytokines receptors family signature 2:**

Amino acids 967-973

**Sema domain:**

Amino acids 126-537

**Plexin repeat:**

Amino acids 555-602

**Thrombospondin type 1 domain:**

Amino acids 613-661;668-719;726-769;856-906;913-963;967-1007

**FIGURE 97**

CAAGCCCTCCCAGCATCCCCTCTCCTGTGTTCCCTCCCCAGTTCTCTACTCAGAGTTGACTGACCAGAGATTTAT  
CAGCTTGGAGGGCTGGAGGTGTGGATCCATGGGGTAGCCTCAACGCATCTGCCCTCCACCCAGCCAGCTCAT  
GGCCACGTGGCCTGGCCAGCCTCAGCACCAGGGCCAGTGAACAGAGCCCTGGCTGGAGTCCAAAC**ATG**TGG  
GGCTGGTGGGCTCCTGCTGGCCTGGCTGGGTGGCTGGGGCTGCATGGGGCGTCTGGCAGCCCCAGCCGGGC  
CTGGGCAGGGTCCCAGGAACACCCAGGGCCTGCTCTGCTGCGGACTCGAAGGAGCTGGGTCTGGAACCAAGTTCT  
TTGTCAATGAGGAATATGCTGGTCCAGAGCCTGTTCTCATTGGCAAGCTGCACCTCGGATGTTGACCCGGGAGAG  
GGCCGCACCAAGTACCTGTTGACCGGGGAGGGGGCAGGCACCGTATTTGTGATGATGAGGCCACAGGCAATAT  
TCATGTTACCAAGAGCCTTGACCGGGAGGAAAAGGCGCAATATGTGCTACTGGCCAAAGCCGTGGACCGAGCCT  
CCAACCGGGCCCTGGAGCCCCCATCAGAGTTCATCATCAAAGTGAAGACATCAACGACAATCCACCCATTTTT  
CCCCTTGGGCCCTACCATGCCACCGTGCCCGAGATGTCCAATGTCCGGACATCAGTGATCCAGGTGACTGCTCA  
CGATGCTGATGACCCAGCTATGGGAACAGTGCCAAGCTGGTGTACACTGTTCTGGATGGACTGCCTTTCTTCT  
CTGTGGACCCAGACTGGAGTGGTGGTACAGCCATCCCCAACATGGACCGGGAGACACAGGAGGAGTTCTTG  
TGGTGATCCAGGCCAAGGACATGGGCGGCCACATGGGGGGCTGTCAGGCAGCACTACCGGAGCGTGCAGCT  
CAGCGATGTCAACGACAACCCCCCAAGTTCACACAGAGCCTATACCAGTTCTCCGTGGTGGAGACAGCTGGAC  
CTGGCACACTGGTGGGCCGGCTCCGGGCCAGGACCCAGACCTGGGGGACAACGCCCTGATGGCATAACAGCATC  
CTGGATGGGGAGGGTCTGAGGCCCTCAGCATCAGCACAGACTTGCAGGGTCGAGACGGGCTCCTCACTGTCCG  
CAAGCCCTAGACTTTGAGAGCCAGCGCTCCTACTCCTTCCGTGTCGAGGCCACCAACACGCTCATTGACCCAGCC  
TATCTGCGGCGAGGGCCCTTCAAGGATGTGGCTCTGTGCGTGTGGCAGTGAAGATGCCCCAGAGCCACCTGC  
CTTACCCAGGCTGCCTACCACCTGACAGTGCCTGAGAACAAGGCCCCGGGGACCTGGTAGGCCAGATCTCCG  
CGGCTGACCTGGACTCCCCTGCCAGCCCAATCAGATACTCCATCCTTCCCCACTCAGATCCGGAGCGTTGCTTC  
TCTATCCAGCCCAGGAAGGCACCATCCATACAGCAGCACCCCTGGATCGCGAGGCTCGCGCCTGGCACAACCT  
CACTGTGCTGGCTACAGAGCTCGACAGTTCTGCACAGGCCCTCGCGCGTGAAGTGGCCATCCAGACCCTGGATG  
AGAATGACAATGCTCCCAGCTGGCTGAGCCCTACGATACTTTTGTGTGTGACTCTGCAGCTCCTGGCCAGCTG  
ATTAGGTTCATCCGGGCCCTGGACAGAGATGAAGTGGCAACAGTAGCCATGTCTCCTTTCAAGGTCCCTTGGG  
CCCTGATGCCAACTTTACTGTCCAGGACAACCGAGATGGCTCCGCCAGCCTGCTGCTGCCCTCCCGCCCTGCTC  
CACCCCGCCACTGCCCTACTTGGTTCCTATAGAATGTGGGACTGGGGCAGCCGGCGCTGAGCAGCACTGCC  
ACAGTACTGTTAGTGTGTGCCGCTGCCAGCCTGACGGCTCTGTGGCATCCTGCTGGCCTCAGCTCACCTCTC  
AGCTGCTGGGCTCAGCACCGGCGCCCTGCTTGGCATCATCACCTGTGTGGGTGCCCTGCTTGGCCCTGGTGGTGC  
TCTTCGTGGCCCTGCGGCGGCAGAAGCAAGAAGCACTGATGGTACTGGAGGAGGAGGACGTCCGAGAGAATC  
ATCACCTACGACGACGAGGGCGGCGGCGAGGAGACACCGAGGCTTCGACATCACGGCCTTGAGAAACCCGGA  
CGGGGCGGCCCCCGGCGCCCGGCCCTCCCGCGCGCCGAGACGTGTGCCCGGGCCCCGGGTGTCCGCGCCAGC  
CCAGACCCCGGCCCCCGGCGACGTGGCGCAGCTCCTGGCGCTGCGGCTCCGCGAGGCGGACGAGGACCCCGGC  
GTACCCCGTACGACTCGGTGCAGGTGTACGGCTACGAGGGCCCGGCTCCTCTTGGCGCTCCCTCAGCTCCCT  
GGGCTCCGGCAGCGAAGCCGGCGGCGCCCCGGCCCCGGAGCCGCTGGACGACTGGGGTCCGCTCTTCCGCACC  
CTGGCCGAGCTGTATGGGGCCAAGGAGCCCCGGCCCC**TG**AGCGCCCGGGCTGGCCCCGGCCACCGCGGGGG  
GGGGCAGCGGGCACAGGCCCTCTGAGTGAGCCCCACGGGTCCAGGCGGGCGGCAGCAGCCAGGGGCCCCAGG  
CCTCCTCCCTGTCTTGTGTCCCTCCTTGCCTCCCCGGGGCACCCCTCGCTCTCACCTCCCTCCTCCTGAGTCGG  
TGTGTGTGTCTCTCTCCAGGAATCTTGTCTCTATCTGTGACACGCTCCTCTGTCCGGGCTGGGTTTCTGCC  
CTGGCCCTGGCCCTGCGATCTCTACTGTGATTCTCTCCTTCCCTCCGTGGCGTTTTTGTCTCTGCAGTTCTGAA  
GCTCACACATAGTCTCCCTGCGTCTTCTTGCCATACACATGCTCTGTGTCTGTCTCCTGCCACATCTCCCT  
TCCTTCTCTGGGTCCCTGTGACTGGCTTTTTTGTTTTTTTCTGTTGTCCATCCCCAAATCAAGAGAAACTTCC  
AGCCACTGCTGCCACCCTCCTGCAGGGGATGTTGTGCCCCAGACCTGCCTGCATGGTTCCATCCATTACTCAT  
GGCCTCAGCCTCATCTGGCTCCACTGGCCTCCAGCTGAGAGAGGGAACCAGCCTGCCTCCAGGGCAAGAGCT  
CCAGCCTCCCGTGTGGCCGCTCCCTGGAGCTCTGCCAGCTGCCAGCTTCCCTGGGCATCCAGCCCTGGGC  
ATTGTCTTGTGTGCTTCTGAGGGAGTAGGGAAAGGAAAGGGGGAGCGGCTGGGGAAGGGGAAAGAGGGAGGA  
AGGGGAGGGGCTCCATCTCTAATTTCCATAATAAACAACACTTTATTTTGTAAAAAC

## FIGURE 98

MWGLVRLLLLAWLGGWGCGRRLAAPARAWAGSREHPGPALLRTRRSWVWNQFFVIEEYAGP  
EPVLIGKHLHSDVDRGEGRTKYLLTGEGAGTVFVIDEATGNIHVTKSLDREEKAQYVLLAQ  
AVDRASNRPLEPPSEFI IKVQDINDNPPIFPLGPYHATVPEMSNVGTSVIQVTAHDADDP  
SYGNSAKLVYTVLDGLPFFSVDPQTGVVRTAIPNMDRETQEEFLVVIQAKDMGGHMGGLS  
GSTTIVTTLSDVNDNPPKFPQSLYQFSVVEVETAGPGTLVGRRLRAQDPDLGDNALMAYSILD  
GEGSEAFSISTDLQGRDGLLTVRKPLDFESQRSYSFRVEATNTLIDPAYLRRGPFKDVAS  
VRVAVQDAPEPPAFTQAAYHLTVPENKAPGTLVGQISAADLSPASPIRYSILPHSDPER  
CFSIQPEEGTIHTAAPLDREARAWHNLTVLATELDSSAQASRVQVAIQTLDENDNAPQLA  
EPYDTFVCDSAAPGQLIQVIRALDRDEVGNSSHVSFQGPLGPDANFTVQDNRDGSASLLL  
PSRPAPPRHAPYLVPIELWDWGQPALSSTATVTVSVCRCPDGSVASCWPEAHLAAGLS  
TGALLAIITCVGALLALVVLVVALRRQKQEALMVLEEDVRENIITYDDEGGGEEDTEAF  
DITALQNPDGAAPPAPGPPARRDVLPRARVSRQPRPPGPADVAQLLALRLREADEDPGVP  
PYDSVQVYGYEGRGSSCGSLSSILGSGSEAGGAPGPAEPLDDWGPLFRFLAELYGAKEPPA  
P

**Signal peptide:**

Amino acids 1-16

**Transmembrane domain:**

Amino acids 597-624

**N-glycosylation sites:**

Amino acids 446-449;510-513;525-528

**N-myristoylation sites:**

Amino acids 13-18;206-211;233-238;237-242;238-243;275-280;390-395;  
394-399;429-434;583-588;598-603;602-607;612-617;  
734-739;738-743;746-751

**ATP synthase c subunit signature:**

Amino acids 691-712

**Cadherins extracellular repeated domain signature:**

Amino acids 138-148;247-257

**Cadherin domain:**

Amino acids 50-141;155-250;264-366;379-470;483-577

**Cadherin cytoplasmic region:**

Amino acids 625-776





## **FIGURE 100**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA213858
><subunit 1 of 1, 627 aa, 1 stop
><MW: 66189, pI: 7.31, NX(S/T): 5
MAILPLLLCLLPLAPASSPPQSATPSPCPRRCRCQTQSLPLSVLCPGAGLLFVPPSLDRR
AAELRLADNFIASVRRRDLANMTGLLHLSLSRNTIRHVAAGAFADLRALRALHLDGNRLT
SLGEGQLRGLVNLRLHLILSNNQLAALAAGALDDCAETLEDLDLSYNNLEQLPWEALGRIG
NVNTLGLDHNLLASVPGAFSRLHKLARLDMTSNRLTTIPDPPLFSRLPLLARPRGSPASA
LVLAFIGGNPLHCNCELVWLRRLAREDDLEACASPPALGGRYFWAVGEEEFVCEPPVVTHR
SPPLAVPAGRPAALRCRAVGDPPEPRVRWVSPQGRLGNSRRARAFPNGTLELLLVTEPGDG
GIFTCIAANAAGEATAAVELTVGPPPPQLANSTSCDPPRDGDPDALTPPSAASASAKVA
DTGPPTDRGVQVTEHGATAALVQWPDQRPIPGIRMYQIQYNSSADDILVYRMI PAESRSF
LLTDLASGRTYDLCVLAVYEDSATGLTATRPVGCAREFSTEPALRPCGAPHAPFLGGTMI I
ALGGVIVASVLVFI FVLLMRYKVHGGQPPGKAKIPAPVSSVCSQTNGALGPTPTPAPPAP
EPAALRAHTVVQLDCEPWGPGHEPVGP
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-16

**Transmembrane domain:**

Amino acids 35-55; 536-556

**N-glycosylation sites:**

Amino acids 81-84;338-341;347-350;392-395;461-464

**N-myristoylation sites:**

Amino acids 116-121;125-130;180-185;186-191;235-240;  
360-365;361-366;429-434;436-441;505-510;  
544-549;566-571

**Leucine Rich Repeat:**

Amino acids 60-83;84-107;108-131;132-155;157-180;  
181-203;204-227

**Leucine rich repeat C-terminal domain:**

Amino acids 248-293

**Immunoglobulin domain:**

Amino acids 309-367

**Fibronectin type III domain:**

Amino acids 424-504

**FIGURE 101**

CGACTCCATAACCGTGGCCTTGGCCCCAGTCCCCCTGACTTCCGGACTTCAGACCAGATACTGCCCATATCCCC  
TTATGAAGTCTTGGCCAGGCAACCCCTAGGGGTACGTTTTCTAAAGATTAAAGAGGCGGTGCTAAGCTGCAGA  
CGGACTTGC GACTCAGCCACTGGTGTAAGTCAGGCGGGAGGTGGCGCCCAATAAGCTCAAGAGAGGAGGCGGGT  
5 TCTGGAAAAAGGCCAATAGCCTGTGAAGGCGAGTCTAGCAGCAACCAATAGCTATGAGCGAGAGGCGGGACTCT  
GAGGGAAGTCAATCGCTGCCGCAGGTACCGCCAATGGCTTTTGGCGGGGGCGTTCCCAACCCCTGCCCTCTCTC  
ATGACCCCGTCCCGGATTATGCGCCGGGACTGGGCTGCTGGCGCTGCGGACGCTGCCAGGGCCCAGCTGGGTGC  
GAGGCTCGGGCCCTTCCGTGCTGAGCCGCTGCAGGACGCGGCCGTGGTGC GGCCCTGGCTTCCTGAGCAGGCA  
GAGGAGGAGACGCTGAGCCGAGA AACTGGAGCCGAGCTGCGCCGCGCCGCTACGAATACGATCACTGGGACGC  
10 GGCCATCCACGGCTTCCGAGAGACAGAGAAGTCGCGCTGGT CAGAAGCCAGCCGGGCCATCCTGCAGCGCGTGC  
AGGCGGCCGCCTTTGGCCCCGGCCAGACCCTGCTCTCCTCCGTGCACGTGCTGGACCTGGAAGCCCGCGGCTAC  
ATCAAGCCCACGTGGACAGCATCAAGTTC TCGGGGGCCACCATCGCCGGCCTGTCTCTCTGTCTCCAGCGT  
TATGCGGCTGGTGCACACCAGGAGCCGGGGAGTGGCTGGA AACTCTTGCTGGAGCCGGGCTCCCTCTACATCC  
TTAGGGGCTCAGCCCGTTATGACTTCTCCCATGAGATCCTTCGGGATGAAGAGTCTTCTTTGGGGAACGCCGG  
15 ATTCCCCGGGGCCGGCGCATCTCCGTGATCTGCCGCTCCCTCCCTGAGGGCATGGGGCCAGGGGAGTCTGGACA  
GCCGCCCCAGCCTGCTGACCCCCAGCTTCTACAGACACCAGATTTGTGAATAAAGTTGGGGAATGGACAGCCT

## FIGURE 102

MAGTGLLLALRTLPGPSWVRGSGPSVLSRLQDAAVVRPGFLSTABEEETLSRELEPELRRRRYEYDHWDAAIHGFR  
ETEKSRWSEASRAILQRVQAAAFGPGQTLSSVHVLDLLEARGYIKPHVDSIKFCGATIAGLSLLSPSVMRLVHT  
QEPGEWLELLLLLEPGSLYILRGSARYDFSHEILRDEESFFGERRIPRGRRISVICRSLPEGMGPGESGQPPPAC

Important features of the protein:

Signal peptide:

1-18

Transmembrane domain:

None

cAMP- and cGMP-dependent protein kinase phosphorylation site.

196-199

N-myristoylation site.

20-25

129-134

208-213

Amidation site.

194-197

**FIGURE 103**

CTCCCCGGCGCCGAGGCAGCGTCTCCTCCGAAGCAGCTGCACCTGCAACTGGGCAGCCTGGACCCTCGTGCC  
CTGTTCCCGGGACCTCGCGCAGGGGGCGCCCGGGACACCCCTGCGGGCCGGGTGGAGGAGGAAGAGGAGGAG  
GAGGAAGAAGACGTGGACAAGGACCCCATCCTACCCAGAACACCTGCCTGCGCTGCCGCACTTCTCTTTAAG  
GGAGAGGAAAAGAGAGCCTAGGAGAACCATGGGGGGCTGCGAAGTCCGGGAATTTCTTTTGAATTTGGTTTTCT  
TCTTGCCCTGCTGACAGCGTGGCCAGGCGACTGCAGTCACTGCTCCAACAACCAAGTTGTGTTGCTTGATACA  
ACAACGTACTGGGAGAGCTAGGATGGAAAACATATCCATTAATGGGTGGGATGCCATCACTGAAATGGATGA  
ACATAATAGGCCATTACACATACCAGGTATGTAATGTAATGGAACCAAACCAAACAACTGGCTTCGTACAA  
ACTGGATCTCCCGTGATGCAGCTCAGAAAATTTATGTGGAATGAAATTCACACTAAGGGATTGTAACAGCATC  
CCATGGGTCTTGGGACTTGCAAAGAAACATTTAATCTGTTTTATATGGAATCAGATGAGTCCACGGAATTA  
ATTCAGCCAAACCAGTATACAAAGATCGACACAATTGCTGCTGATGAGAGTTTTACCCAGATGGATTTGGGTG  
ATCGCATCCTCAAACCTCAACACTGAAATTCGTGAGGTGGGGCCTATAGAAAAGGAAAGGATTTATCTGGCTTTT  
CAAGACATGGGGCGTGCATTGCCCTGGTTTTAGTCCGTGTTTTCTACAAGAAATGCCCTTCACTGTTTCGTA  
CTTGGCCATGTTTCTGATACCATTCCAAGGTTGATTCCCTCCTCTTTGGTTGAAGTACGGGGTCTTGTTGTA  
AGAGTGTGAAGAGCGTGACACTCCTAAACTGTATTGTGGAGCTGATGGAGATTGGCTGGTTCCTCTTGAAGG  
TGCATCTGCAGTACAGGATATGAAGAAATGAGGGTCTTGCCATGCTTGACAGCAGGATTCTATAAAGCTTT  
TGCTGGGAACACAAAATGTTCTAAATGTCCTCCACACAGTTTAACATACATGGAAGCAACTTCTGTCTGTCAGT  
GTGAAAAGGGTTATTTCCGAGCTGAAAAGAGCCACCTTCTATGGCATGTACCAGGCCACCTTCACTCCTAGG  
AATGTGGTTTTTAACATCAATGAAACAGCCCTTATTTGGAAATGGAGCCCACCAAGTACACAGGAGGGAGAAA  
AGATCTCACATACAGTGTAATCTGTAAGAAATGTGGCTTAGACACCAGCCAGTGTGAGGACTGTGGTGGAGGAC  
TCCGCTTCATCCCAAGACATACAGGCTGATCAACAATTCGCTGATAGTACTTGACTTTGTGTCTCACGTGAAT  
TACACCTTTGAAATAGAAGCAATGAATGGAGTTCTGAGTTGAGTTTTCTCCCAAGCCATTCACAGCTATTAC  
AGTGACCACGGATCAAGATGCACCTTCCCTGATAGGTGTGGTAAGGAAAGGACTGGGCATCCCAAATAGCATGGC  
CTATCATGGCAAGCACCTGCTTTTTCCAATGGAGCCATTCGGACTACGAGATCAAGTACTATGAGAAAAGAAC  
TGAGAGAGTACACTACTCTTCCACAAGGTCCAAAGCCCCAGTGTGTCATCACAGGCTTAAGCCAGCCACCA  
AATATGTATTTACATCCGAGTGAGAACTGCGACAGGATACAGTGGCTACAGTCAAGAAATTTGAATTTGAAACA  
GGAGATGAACTTCTGACATGGCAGCAGAACAAGGACAGATTTCTCGTGATAGCCACCGCCGCTGTTGGCGGAT  
CACCTCCTCGTCACTCCTCACTTTATTCTTCTTGATCACTGGGAGATGTCAGTGGTACATAAAAGCCAAGATGA  
AGTCAGAAGAGAAGAGAAGAAACCCTTACAGAATGGGCATTTGCGCTTCCCGGGAATTAAGAACTTACATTTGAT  
CCAGATACATATGAAGACCCATCCCTAGCAGTCCATGAATTTGCAAAGGAGATTGATCCCTCAAGAATTCGTAT  
TGAGAGAGTACATGGGGCAGGTGAATTTGGAGAAGTCTGTAGTGGCGCTTTGAAGACACCAGGGAAAAGAGAGA  
TCCCAGTTGCCATTAAGAACTTTGAAAGGTGGCCACATGGATCGGCLLGLLGLLGLLGLLGLLGLLGLLGLLGLL  
ATCATGGGCCAGTTTGACCATCAAACATCATTGCGCTAGAAGGGGTTGTACCAAAGATCCTTCCCGGCCAT  
TGGGGTGGAGGCGTTTTGCCCCAGCTTCTGAGGGCAGGGTTTTTAAATAGCATCCAGGCCCCGCATCCAGTGC  
CAGGGGGAGGATCTTTGCCCCCCAGGATTCCTGCTGGCAGACCAGTAATGATTGTGGTGGAAATATATGGAGAA  
GGATCCCTAGACTCCTTTTTCGGAAGCATGATGGCCACTTCACAGTCACTCCAGTTGGTCGGAATGCTCCGAG  
CATTGCCATCAGGCATGAAGTATCTTCTGATATGGGTTATGTTTCATCGAGACCTAGCGGCTCGGAATATACGG  
TCAATAGCAACTTAGTATGCAAAGTTTCTGATTTTGGTCTCTCCAGAGTGTGGAAAGATGATCCAGAAGCTGCT  
TATACAACAACCTGGTGGAAAATCCCATAAGGTGGACAGCCCAGAAGCCATCGCTACAGAAAATTTCTCCTC  
AGCAAGCGATGCATGGAGCTATGGCATTGTCATGTGGGAGGTCATGTCCTATGGAGAGAGACCTTATGGGAAATG  
TCTAACCAAGATGTCTTCTGTCCATGAAGAAGGGTACAGACTCCAGCTCCCATGGGCTGTCCAGCATCTCT  
ACACCAGTGTGCTCCACTGCTGGCAGAAGGAGAAAATCACAGACCAAATTTACTGACATTTGTCAGCTTCC  
TTGACAAAATGATCCGAAATCCAGTGCCTTACACCCTGGTGGAGGACATCCTTGTAATGCCAGAGTCCCT  
GGTGAAGTTCCGGAATATCCTTTGTTTGTACAGTTGGTACTGGCTAGATTTCTATAAAGATGGGGCAATACAA  
GAATAACTTCGTGGCAGCAGGGTTTACAACATTTGACCTGATTTCAAGAATGAGCATTGATGACATTAGAAGAA  
TTGGAGTCATACTTATTTGGACACCAGAGACGAATAGTCAGCAGCATAACAGACTTTACGTTTACACATGATGCAC  
ATACAGGAGAAGGGATTTGATGTATGAAAGTACCACAAGCACCTGTGTTTTGTGCCTCAGCATTTCTAAAATGA  
ACGATACCTCTCTACTACTCTCTCTTCTGATTTCCAAACATCACTTCAAACTGCAGTCTTCTGTTTCAGAC  
TATAGGCACACACTTATGTTTTATGCTTCCAACAGGATTTTAAATCATGCTACATAAATCCGTTCTGAATAA  
CCTGCAACTAAAAAAAAAAAAAAAAAAAA

## FIGURE 104

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA222653

><subunit 1 of 1, 1036 aa, 1 stop

><MW: 116379, pI: 6.94, NX(S/T): 5

MGGCEVREFLLQFGFFLPLLTAWPGDCSHVSNNQVLLDTTTVLGGELGWKTYPLNGWDAI  
TEMDEHNRPIHTYQVCNVMENPNQNNWLRTNWISRDAAQKIYVEMKFTLRDCNSIPWVLGT  
CKETFNLFYMESDESHGIKFKPNQYTKIDTIAADESFTQMDLGDRILKINTEIREVGP  
RKGFYLAFAQDIGACIALVSVRVFYKCKPFTVRNLAMFPDTPRVDSSSLVEVRGSCVKSA  
EERDTPKLYCGADGDWLVPLGRICSTGYEEIEGSCHACRPGFYKAFAGNTKCSKCP  
LTYMEATSVCQCEKGYFRAEKDPPSMACRPPSAPRNVVFNINETALILEWSPPSDTGGR  
KDLTYSVICKKCGLDTSQCEDCGGGLRFIPRHTGLINNSVIVLDFVSHVNYTFEIEAMNG  
VSELSFSPKPFMTAITVTTDQDAPSLIGVVRKDWASQNSIALSWQAPAFSNGAILDYEIKY  
YEKEHEQLTYSSTRSKAPSVIITGLKPKTYVFHIRVRTATGYSQKFEFETGDETS  
MAAEQOQILVIATAAVGGFTLLVILTLFFLITGRQWYIKAKMKSEEKRRNHLQNGHLRF  
PGIKTYIDPDTYEDPSLAVHEFAKEIDPSRIRIERVIGAGEFGEVCSGRLKTPGKREIPV  
AIKTLKGGHMDRQRDFLEASIMGQFDHPNIIRLEGVVTKRSFPAIGVEAFCPSFLRAG  
FLNSIQAPHPVPPGGSLPPRIAGRPVMIVVEYMENGLDSFLRKHDGHFTVIQLVGLMR  
GIASGMKYLSDMGYVHRDLAARNILVNSNLVCKVSDFGLSRVLEDDPEAAAYTTTGGKIP  
RWTAPEAIAYRKFSSASDAWSYGIVMWEVMSYGERPYWEMSNQDVILSIEEGYRLPAPMG  
CPASLHQLMLHCWQKERNHRPKFTDIVSFLDKLIRNPSALHTLVEDILVMPESPGEVPEY  
PLEFVTVDGLWDSIKMGQYKNNFVAAGFTTFDLISRMSIDDIRRIGVILIGHQRRIVSSIQ  
TLRLHMMHIQEKGFHV

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-22

**Transmembrane domain:**

Amino acids 551-571

**N-glycosylation sites:**

Amino acids 343-346;397-400;410-413;756-759

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 851-854

**Tyrosine kinase phosphorylation sites:**

Amino acids 483-490;604-612;787-794

**N-myristoylation sites:**

Amino acids 192-197;274-279;289-294;373-378;394-399;504-509;  
757-762;777-782;781-786;900-905;976-981

**Amidation site:**

Amino acids 358-361;653-656

**Tyrosine protein kinases specific active-site signature:**

Amino acids 794-806

**Receptor tyrosine kinase class V signature 1:**

Amino acids 192-208

**Ephrin receptor ligand binding domain:**

Amino acids 34-207

**pkine Protein kinase domain:**

Amino acids 631-927

**Fibronectin type III domain:**

Amino acids 332-425;440-527

**SAM domain (Sterile alpha motif):**

Amino acids 959-1023

**FIGURE 105**

GGCGGGGGCTGCGCGGAGCGGCGTCCCCTGCAGCCGCGGACCGAGGCAGCGGGCGCACCTGCCGGCCGAGCAA  
TGCCAAGTGAGTACACCTATGTGAAACTGAGAAGTGATTGCTCGAGGCCCTTCCCTGCAATGGTACACCCGAGCT  
CAAAGCAAGATGAGAAAGGCCAGCTTGTATTATAAAGACATCCTCAAATGTACATTGCTTGTGTTTGGAGTGTG  
GATCCCTTTATATCCTCAAGTTAAATFATACTACTGAAGAATGTGACATGAAAAAATGCATTATGTGGACCCTG  
ACCATGTAAGAGAGACTCAGAAATATGCTCAGCAAGTCTTGCAGAAGGAATGTCGTCCCAAGTTTCCAAGACA  
TCAATGGCGCTGTTATTTGAGCACAGGTATAGCGTGGACTTACTCCCTTTTGTGCAGAAGGCCCCCAAAGACAG  
TGAAGCTGAGTCCAAGTACGATCCTCCTTTTGGGTTCGGAAGTTCTCCAGTAAAGTCCAGACCCTCTTGGAAC  
TCTTGCCAGAGCACGACCTCCCTGAACACTTGAAGCCAAGACCTGTGCGGCGTGTGTGGTTATTGGAAGCGGA  
GGAATACTGCACGGATTAGAACTGGGCCACACCCTGAACCAGTTCGATGTTGTGATAAGGTTAAACAGTGCACC  
AGTTGAGGGATATTCAGAACATGTTGGAATAAAACTACTATAAGGATGACTTATCCAGAGGGCGCACCACCTGT  
CTGACCTTGAATATTTTCCAATGACTTATTTGTTGCTGTTTTATTTAAGAGTGTGATTTCAACTGGCTTCAA  
GCAATGGTAAAAAAGGAAACCCTGCCATTCTGGGTACGACTCTTCTTTTGAAGCAGGTGGCAGAAAAAATCCC  
ACTGCAGCCAAAACATTTCAGGATTTTGAATCCAGTTATCATCAAAGAGACTGCCTTTGACATCCTTCAGTACT  
CAGAGCCTCAGTCAAGTTCTGGGGCCGAGATAAGAACGTCCCACAATCGGTGTCATTGCCGTTGTCTTAGCC  
ACACATCTGTGCGATGAAGTCAGTTTGGCGGGTTTTGGATATGACCTCAATCAACCAGAACACCTTTGCACTA  
CTTCGACAGTCAATGCATGGCTGCTATGAACTTTCAGACCATGCATAATGTGACAACGGAACCAAGTTCCTCT  
TAAAGCTGGTCAAAGAGGGAGTGGTGAAGATCTCAGTGGAGGCATTGATCGTGAATTTTGAACACAGAAAACC  
TCAGTTGAAAATGCAACTCTAACTCTGAGAGCTGTTTTTGAACAGCCTTCTTGATGATTTTCTCCATCCTGCAGA  
TACTTTGAAGTGCAGCTCATGTTTTAACTTTTAATTTAAAAACACAAAAAATTTTAGCTCTTCCCACTTTT  
TTTTTCTATTTATTTGAGGTCACTGTTTGTGTTTTGACACCATTTTGTAAATGAACTTAAGAATTGAATTGG  
AAAGACTTCTCAAAGAGAATTGTATGTAACGATGTTGTATTGATTTTTAAGAAAAGTAATTTAATTTGTAAAAC  
TCTGCTCGTTTACTGCACATTGAATACAGGTAACATAATGGAAGGAGAGGGGAGGTCACTCTTTTGTGTTG  
GCCCTGAACCTCATTCTGGTTCCTGCTGCGCTGCTTGGTGTGACCCACGGAGGATCCACTCCCAGGATGACGT  
GCTCCGTAGCTCTGCTGCTGATACTGGGTCTGCGATGCAGCGCGTGAAGCCTGGGCTGGTTGGAGAAGGTCAC  
AACCTTCTCTGTTGGTCTGCCTTCTGCTGAAAGACTCGAGAACCAACCAGGGAAGCTGTCTGAGGTCCTG  
GTCGGAGAGGGACATAGAATCTGTGACCTCTGACAAC'GTGAAGCCACCCTGGGCTACAGAAACCACAGTCTTC  
CCAGCAATTATTACAATTCTTGAATTCCTTGGGGATTTTTTACTGCCCTTTCAAAGCACTTAAGTGTTAGATCT  
AACGTGTTCCAGTGTCTGCTGAGGTGACTAAAAAATCAGAACAAAACCTTCTAT'ATCCAGAGTCATGGGAGA  
GTACACCCTTTCAGGAATAATGTTTTGGGAAACACTGAAATGAAATCTTCCAGTATTATAAATTGTGTATTTAA

## **FIGURE 106**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA96897
><subunit 1 of 1, 362 aa, 1 stop
><MW: 41736, pI: 8.80, NX(S/T): 3
MRRPSLLLKDILKCTLLVFGVWILYILKLNYYTTEECMDKMHYVDPDHVKRAQKYAQQVLQK
ECRPKFAKTSMAALLFEHRYSVDLLPFVQKAPKDSEAESKYDPPFGFRKFSSKVQTLLELLPE
HDLPEHLKAKTCRRCVVGSGGILHGLELGHNTLNQFDVVIRLNSAPVEGYSEHVGNKTTIRM
TYPEGAPLSDLEYYSNDLFAVLFKSVDFNWLQAMVKKETLPFWVRLFFWKQVAEKIPLQPK
HFRILNPVIIKETAFDILQYSEPQSRFWGRDKNVPTIGVIAVVLATHLCDEVSLAGFGYDLN
QPRTPHLYFDSQCMAAMNFQTMHNVTTETKFLKLVKEGVVKDLSGGIDREF
```

### **Important features of the protein:**

#### **Transmembrane domain:**

Amino acids 11-27;281-297

#### **N-glycosylation sites:**

Amino acids 30-34;180-184;334-338

#### **cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 2-6;109-113;223-227

#### **N-myristoylation sites:**

Amino acids 146-152;150-156;179-185;191-197



## **FIGURE 107**

TGACGCGGGGCGCCAGCTGCCAACTTCGCGCGGGAGCTCCCCGGCGGTGCAGTCCCGTCCCGGCGGGCGCGG  
GCGGC**ATGA**AAGACTAGCCGCGCGGGCCGAGCGCTCCTGGCCGTGGCCCTGAACCTGCTGGCGCTGCTGTTTCG  
CCACCACCGCTTTCCTCACCACGCACTGGTGCCAGGGCACGCAGCGGGTCCCCAAGCCGGGCTGCGGCCAGG  
GCGGGCGCGCAACTGCCCAACTCGGGCGCCAAACGCCACGGCCAACGGCACCGGCCCGCCCGCCGCGCCG  
CCGCGCGCGCCACCGCTCGGGGAACGGCCCCCTGGCGCGCGCTCTACAGCTGGGAGACCGGGCGACGACC  
GCTTCCTCTCAGGAATTTCCACACCGGCATCTGGTACTCGTGCGAGGAGGAGCTCAGCGGGCTTGGTGAAA  
AATGTCGCAGCTTCATTGACCTGGCCCCGGCGTCGGAGAAAGGCCTCCTGGGAATGGTCGCCACATGATGT  
ACACGCAGGTGTTCCAGGTACCGTGAGCCTCGGTCTGAGGACTGGAGACCCCATTCCTGGGACTACGGGT  
GGTCTTCTGCCTGGCGTGGGGCTCCTTTACCTGCTGCATGGCAGCCTCTGTCAACACGCTCAACTCCTACA  
CCAAGACGGTCATTGAGTTCGGGCACAAGCGCAAGGTCTTTGAGCAGGGCTACCGGGAAGAGCCGACCTTCA  
TAGACCTGAGGCCATCAAGTACTTCCGGGAGAGGATGGAGAAGAGGGACGGGAGCGAGGAGGACTTTCACT  
TAGACTGCCGCCACGAGAGATACCCTGCCCGACACCAGCCACACATGGCGGATTCCTGGCCCCGGAGCTCCG  
CACAGGAAGCACCAGAGCTGAACCGACAGTGTGGGTCTTGGGGCACTGGGT**TGA**CCAAGACCTCAACCTG  
GCCCCGGACCTCAGGCCATCGCTGGCACCAGCCCCCTGCTGCAAGACCACCAGAGTGGTGCCCCAGAACCC  
TGGCCTGTGTGCCGTGAACTCAGTCAGCCTGCGTGGGAGATGCCAGGCCTGTCTGCCATCGCTGCCTGGG  
TCCCATGGCCTTGAAATGGGGCCAGGGCAGGCCAAGGGAATGCACAGGGCTGCACAGAGTGACTTTGGGA  
CAGCAGCCCCGGACTCTTGCCATCATCACATGAGCCCTGCTGGGCACAGCTGCGATGCCAGGAGACACATGG  
CCACTGGCCACTGAATGGCTGGCACCCACAAGCCAGTCAGGTGCCAGAGGGGCAGAGCCCTTTGGGGGCA  
GAGAGTGGCTTCTGAAGGAGGGGGCAGTGGCGCAGGCACTGCAGGGGTGCACACAGCAGGCACACAGCAG  
GGGCTCAATAAATGCTTGTGAACTTGT

## **FIGURE 108**

MKTSRRGRALLAVALNLLALLFATTAFLTTHWCQGTQRVPKPGCGQGGRANCPNSGANATANGTAAPAAAA  
AAATASGNGPPGGALYSWETGDDRFLFRNFHTGIWYSCEEELSGLGEKCRSFIDLAPASEKGLLGMVAHMM  
YTQVFQVTVSLGPEDWRPHSWDYGWSFCLAWGSFTCCMAASVTTLNSYTKTVIEFRHKRKVFEQGYREEPT  
FIDPEAIKYFRERMEKRDGSEEDFHLDRCRHERYPARHQPHMADSWPRSSAQEAPELNRQCWVLGHWV

Important features of the protein:

Signal peptide:

1-26

Transmembrane domain:

169-189

N-glycosylation site.

58-61

62-65

Glycosaminoglycan attachment site.

77-80

114-117

Tyrosine kinase phosphorylation site.

202-208

N-myristoylation site.

43-48

47-52

56-61

84-89

104-109

174-179



## FIGURE 110

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA142930
><subunit 1 of 1, 512 aa, 1 stop
><MW: 54535, pI: 4.89, NX(S/T): 7
MKAI IHL TLLALLSVNTATNQGNSADAVTTTETATSGPTVAAAADTTETNFPETASTTANT
PSFPTATSPAPPI IISTHSSSTIPTPAPPI IISTHSSSTIPIPTAADSESTTNVNSLATSDI
ITASSPNDGLITMVPSETQSNNEMSPTTEDNQSSGPPTGTALLETSTLNSTGSPNPCQDD
PCADNSLCVKLHNTSFCLCLEGYYNSSTCKKKGKVFPGKISVTVSETFDPEEKHSMAYQD
LHSEITSLFKDVFGTSVYGQTVILTVSTLSLSPRSEMRADDKFVNVTIVTILAETTS DNEK
TVTEKINKAIRSSSNFLNYDLTLRCDYYGCNQTADDCLNGLACDCKSDLQRPNPQSPFC
VASSLKCPDACNAQHKQCLIKKSGGAPECACVPGYQEDANGNCQKCAFYSGLDCKDKFQ
LILTIVGTIAGIVILSMIALIVTARSNNKTKHIEEENLIDEDFQNLKLRSTGFTNLGAE
GSVFPKVRITASRDSQMNPYSSHSSMPRPDY
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-17

**Transmembrane domain:**

Amino acids 421-442

**N-glycosylation sites:**

Amino acids 151-155;169-173;193-197;206-210;284-288;  
332-336;449-453

**N-myristoylation sites:**

Amino acids 330-336;385-391;427-433;478-484

**SEA domain:**

Amino acids 212-328

**FIGURE 111**

CTGGGACTTGGCTTTCTCCGGATAAGCGGGCGGCACCGGCGTCAGCGATGACCGTGCAGAGAC  
TCGTGGCCGCGGCCGTGCTGGTGGCCCTGGTCTCACTCATCCTCAACAACGTGGCGGCCTTC  
ACCTCCAACCTGGGTGTGCCAGACGCTGGAGGATGGGCGCAGGCGCAGCGTGGGGCTGTGGAG  
GTCCTGCTGGCTGGTGGACAGGACCCGGGGAGGGCCGAGCCCTGGGGCCAGAGCCGGCCAGG  
TGGACGCACATGACTGTGAGGCGCTGGGCTGGGGCTCCGAGGCAGCCGGCTTCCAGGAGTCC  
CGAGGCACCGTCAAACCTGCAGTTCGACATGATGCGCGCCTGCAACCTGGTGGCCACGGCCGC  
GCTCACCGCAGGCCAGCTCACCTTCCTCCTGGGGCTGGTGGGCCTGCCCTGCTGTCACCCG  
ACGCCCCGTGCTGGGAGGAGGCCATGGCCGCTGCATTCCAACCTGGCGAGTTTTGTCCTGGTC  
ATCGGGCTCGTGACTTTCTACAGAATTGGCCCATACACCAACCTGTCCTGGTCCTGCTACCT  
GAACATTGGCGCCTGCCTTCTGGCCACGCTGGCGGCAGCCATGCTCATCTGGAACATTCTCC  
ACAAGAGGGAGGACTGCATGGCCCCCGGGTGATTGTCATCAGCCGCTCCCTGACAGCGCGC  
TTTCGCCGTGGGCTGGACAATGACTACGTGGAGTCACCATGCTGAGTGCCCTTCTCAGCGC  
TCCATCAACGCACACCTGCTATCGTGGAACAGCCTAGAAACCAAGGGACTCCACCACCAAGT  
CACTTCCCCTGCTCGTGCAGAGGCACGGGATGAGTCTGGGTGACCTCTGCGCCATGCGTGCG  
AGACACGTGTGCGTTTACTGTTATGTCGGTCATATGTCTGTACGTGTCGTGGGCCAACCTCG  
TTCTGCCTCCAGC

## FIGURE 112

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA147253
><subunit 1 of 1, 226 aa, 1 stop
><MW: 24540, pI: 8.27, NX(S/T): 1
MTVQRLVAAAVALVALVSLILNNVAAFTSNWVCQTLEDGRRRSVGLWRSCWLVDTRGGPS
PGARAGQVDAHDCEALGWGSEAAGFQESRGTVKLQFDMMRACNLVATAALTAGQLTFLLG
LVGLPLLSPDAPCWEEAMAAAFQLASFVLVIGLVTFYRIGPYTNLSWSCYLNIGACLLAT
LAAAMLIWNILHKREDCMAPRVIVISRSLTARFRRGLDNDYVESPC
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-25

**Transmembrane domains:**

Amino acids 105-125;139-157;169-188

**N-glycosylation site:**

Amino acids 164-168

**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

Amino acids 39-43

**Tyrosine kinase phosphorylation site:**

Amino acids 214-222

**N-myristoylation sites:**

Amino acids 44-50;62-68;66-72;79-85

**Amidation site:**

Amino acids 37-41

**FIGURE 113**

GACTTTACCACTACTCGCTATAGAGCCCTGGTCAAGTTCTCTCCACCTCTCTATCTATGTCT  
CAGTTTCTTCATCTGTAACATCAAATGAATAATAATACCAATCTCCTAGACTTCATAAGAGG  
ATTAACAAAGACAAAATATGGGAAAAACATAACATGGCGTCCCATAATTATTAGATCTTATT  
ATTGACACTAAAATGGCATTAAAATTACCAAAGGAAGACAGCATCTGTTTCCTCTTTGGTC  
CTGAGCTGGTTAAAAGGAACACTGGTTGCCTGAACAGTCACACTTGCAACC**ATG**ATGCCTAA  
ACATTGCTTTCTAGGCTTCCTCATCAGTTTCTTCCTTACTGGTGTAGCAGGAACTCAGTCAA  
CGCATGAGTCTCTGAAGCCTCAGAGGGTACAATTTT**CAGT**CCCGAAATTTT**CACA**ACATTTTG  
CAATGGCAGCCTGGGAGGGCACTTACTGGCAACAGCAGTGTCTATTTTGTGCAGTACAAAAT  
ATATGGACAGAGACAATGGAAAAATAAAGAAGACTGTTGGGGTACTCAAGAACTCTCTTGTG  
ACCTTACCAGTGAAACCTCAGACATACAGGAACCTTATTACGGGAGGGTGAGGGCGGCCTCG  
GCTGGGAGCTACTCAGAATGGAGCATGACGCCGCGGTTCACTCCCTGGTGGGAAACAAAAT  
AGATCCTCCAGTCATGAATATAACCCAAGTCAATGGCTCTTTGTTGGTAATTCTCCATGCTC  
CAAATTTACCATATAGATACCAAAGGAAAAAATGTATCTATAGAAGATTACTATGAACTA  
CTATACCGAGTTTTTATAATTAACAATTCACTAGAAAAGGAGCAAAGGTTTATGAAGGGGC  
TCACAGAGCGGTTGAAATTGAAGCTCTAACACCACACTCCAGCTACTGTGTAGTGGCTGAAA  
TATATCAGCCCATGTTAGACAGAAGAAGTCAGAGAAGTGAAGAGAGATGTGTGGAAATCCA  
**TGA**CTTGTGGAATTTGGCATT**CAGCA**ATGTGGAAATCTAAAGCTCCCTGAGAACAGGATGA  
CTCGTGTTTGAAGGATCTTATTTAAAATTGTTTTTGTATTTTCTTAAAGCAATATTC**ACTGT**  
TACACCTTGGGGACTTCTTTGTTTACCATTCTTTTATCCTTTATATTT**CATTTGTAA**ACTA  
TATTTGAACGACATTCACCCCGAAAATTGAAATGTAAAGATGAGGCAGAGAATAAAGTGT  
CTATGAAATTCAGAACTTTATTTCTGAATGTAACATCCCTAATAACAACCTT**CATTCTT**CTA  
ATACAGCAAATAAAAATTTAACAACCAAGGAATAGTATTTAAGAAAATGTTGAAATAATTT  
TTTTAAAATAGCATTACAGACTGAG

## **FIGURE 114**

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA149927
><subunit 1 of 1, 231 aa, 1 stop
><MW: 26980, pI: 7.06, NX(S/T): 5
MMPKHCFLGLISFFLTGVAGTQSTHESLKPQRVQFQSRNFNHILQWQPGRALTGNSSVY
FVQYKIYGQRQWKNKEDCWGTQELSCDLTSETSDIQEPYYGRVRAASAGSYSEWSMTPRF
TPWWETKIDPPVMNITQVNGSLLVILHAPNLPYRYQKEKNVSIEDYYELLYRVFIINNSL
EKEQKVYEGAHRAVEIEALTPHSSYCVVAEIQPMLDRRSQRSEERCVEIP
```

**Important features of the protein:**

**Signal peptide:**

Amino acids 1-21

**N-glycosylation sites:**

Amino acids 56-60;134-138;139-143;160-164;177-181

**N-myristoylation sites:**

Amino acids 18-24;21-27;189-195



**SECRETED AND TRANSMEMBRANE  
POLYPEPTIDES AND NUCLEIC ACIDS  
ENCODING THE SAME**

**FIELD OF THE INVENTION**

[0001] The present invention relates generally to the identification and isolation of novel DNA and to the recombinant production of novel polypeptides.

**BACKGROUND OF THE INVENTION**

[0002] Extracellular proteins play important roles in, among other things, the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. These secreted polypeptides or signaling molecules normally pass through the cellular secretory pathway to reach their site of action in the extracellular environment.

[0003] Secreted proteins have various industrial applications, including as pharmaceuticals, diagnostics, biosensors and bioreactors. Most protein drugs available at present, such as thrombolytic agents, interferons, interleukins, erythropoietins, colony stimulating factors, and various other cytokines, are secretory proteins. Their receptors, which are membrane proteins, also have potential as therapeutic or diagnostic agents. Efforts are being undertaken by both industry and academia to identify new, native secreted proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.* 93:7108-7113 (1996); U.S. Pat. No. 5,536,637].

[0004] Membrane-bound proteins and receptors can play important roles in, among other things, the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. Such membrane-bound proteins and cell receptors include, but are not limited to, cytokine receptors, receptor kinases, receptor phosphatases, receptors involved in cell-cell interactions, and cellular adhesion molecules like selectins and integrins. For instance, transduction of signals that regulate cell growth and differentiation is regulated in part by phosphorylation of various cellular proteins. Protein tyrosine kinases, enzymes that catalyze that process, can also act as growth factor receptors. Examples include fibroblast growth factor receptor and nerve growth factor receptor.

[0005] Membrane-bound proteins and receptor molecules have various industrial applications, including as pharma-

ceutical and diagnostic agents. Receptor immunoadhesins, for instance, can be employed as therapeutic agents to block receptor-ligand interactions. The membrane-bound proteins can also be employed for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction.

[0006] Efforts are being undertaken by both industry and academia to identify new, native receptor or membrane-bound proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel receptor or membrane-bound proteins.

**SUMMARY OF THE INVENTION**

[0007] In one embodiment, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a PRO polypeptide.

[0008] In one aspect, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule encoding a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

[0009] In other aspects, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about

90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule comprising the coding sequence of a full-length PRO polypeptide cDNA as disclosed herein, the coding sequence of a PRO polypeptide lacking the signal peptide as disclosed herein, the coding sequence of an extracellular domain of a transmembrane PRO polypeptide, with or without the signal peptide, as disclosed herein or the coding sequence of any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

[0010] In a further aspect, the invention concerns an isolated nucleic acid molecule comprising a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule that encodes the same mature polypeptide encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein, or (b) the complement of the DNA molecule of (a).

[0011] Another aspect the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding a PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated, or is complementary to such encoding nucleotide sequence, wherein the transmembrane domain(s) of such polypeptide are disclosed herein. Therefore, soluble extracellular domains of the herein described PRO polypeptides are contemplated.

[0012] Another embodiment is directed to fragments of a PRO polypeptide coding sequence, or the complement thereof, that may find use as, for example, hybridization probes, for encoding fragments of a PRO polypeptide that may optionally encode a polypeptide comprising a binding site for an anti-PRO antibody or as antisense oligonucleotide

probes. Such nucleic acid fragments are usually at least about 10 nucleotides in length, alternatively at least about 15 nucleotides in length, alternatively at least about 20 nucleotides in length, alternatively at least about 30 nucleotides in length, alternatively at least about 40 nucleotides in length, alternatively at least about 50 nucleotides in length, alternatively at least about 60 nucleotides in length, alternatively at least about 70 nucleotides in length, alternatively at least about 80 nucleotides in length, alternatively at least about 90 nucleotides in length, alternatively at least about 100 nucleotides in length, alternatively at least about 110 nucleotides in length, alternatively at least about 120 nucleotides in length, alternatively at least about 130 nucleotides in length, alternatively at least about 140 nucleotides in length, alternatively at least about 150 nucleotides in length, alternatively at least about 160 nucleotides in length, alternatively at least about 170 nucleotides in length, alternatively at least about 180 nucleotides in length, alternatively at least about 190 nucleotides in length, alternatively at least about 200 nucleotides in length, alternatively at least about 250 nucleotides in length, alternatively at least about 300 nucleotides in length, alternatively at least about 350 nucleotides in length, alternatively at least about 400 nucleotides in length, alternatively at least about 450 nucleotides in length, alternatively at least about 500 nucleotides in length, alternatively at least about 600 nucleotides in length, alternatively at least about 700 nucleotides in length, alternatively at least about 800 nucleotides in length, alternatively at least about 900 nucleotides in length and alternatively at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. It is noted that novel fragments of a PRO polypeptide-encoding nucleotide sequence may be determined in a routine manner by aligning the PRO polypeptide-encoding nucleotide sequence with other known nucleotide sequences using any of a number of well known sequence alignment programs and determining which PRO polypeptide-encoding nucleotide sequence fragment(s) are novel. All of such PRO polypeptide-encoding nucleotide sequences are contemplated herein. Also contemplated are the PRO polypeptide fragments encoded by these nucleotide molecule fragments, preferably those PRO polypeptide fragments that comprise a binding site for an anti-PRO antibody.

[0013] In another embodiment, the invention provides isolated PRO polypeptide encoded by any of the isolated nucleic acid sequences hereinabove identified.

[0014] In a certain aspect, the invention concerns an isolated PRO polypeptide, comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93%

amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein.

[0015] In a further aspect, the invention concerns an isolated PRO polypeptide comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to an amino acid sequence encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein.

[0016] In a specific aspect, the invention provides an isolated PRO polypeptide without the N-terminal signal sequence and/or the initiating methionine and is encoded by a nucleotide sequence that encodes such an amino acid sequence as hereinbefore described. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

[0017] Another aspect the invention provides an isolated PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

[0018] In yet another embodiment, the invention concerns agonists and antagonists of a native PRO polypeptide as

defined herein. In a particular embodiment, the agonist or antagonist is an anti-PRO antibody or a small molecule.

[0019] In a further embodiment, the invention concerns a method of identifying agonists or antagonists to a PRO polypeptide which comprise contacting the PRO polypeptide with a candidate molecule and monitoring a biological activity mediated by said PRO polypeptide. Preferably, the PRO polypeptide is a native PRO polypeptide.

[0020] In a still further embodiment, the invention concerns a composition of matter comprising a PRO polypeptide, or an agonist or antagonist of a PRO polypeptide as herein described, or an anti-PRO antibody, in combination with a carrier. Optionally, the carrier is a pharmaceutically acceptable carrier.

[0021] Another embodiment of the present invention is directed to the use of a PRO polypeptide, or an agonist or antagonist thereof as hereinbefore described, or an anti-PRO antibody, for the preparation of a medicament useful in the treatment of a condition which is responsive to the PRO polypeptide, an agonist or antagonist thereof or an anti-PRO antibody.

[0022] In other embodiments of the present invention, the invention provides vectors comprising DNA encoding any of the herein described polypeptides. Host cell comprising any such vector are also provided. By way of example, the host cells may be CHO cells, *E. coli*, or yeast. A process for producing any of the herein described polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of the desired polypeptide and recovering the desired polypeptide from the cell culture.

[0023] In other embodiments, the invention provides chimeric molecules comprising any of the herein described polypeptides fused to a heterologous polypeptide or amino acid sequence. Example of such chimeric molecules comprise any of the herein described polypeptides fused to an epitope tag sequence or a Fc region of an immunoglobulin.

[0024] In another embodiment, the invention provides an antibody which binds, preferably specifically, to any of the above or below described polypeptides. Optionally, the antibody is a monoclonal antibody, humanized antibody, antibody fragment or single-chain antibody.

[0025] In yet other embodiments, the invention provides oligonucleotide probes which may be useful for isolating genomic and cDNA nucleotide sequences, measuring or detecting expression of an associated gene or as antisense probes, wherein those probes may be derived from any of the above or below described nucleotide sequences. Preferred probe lengths are described above.

[0026] In yet other embodiments, the present invention is directed to methods of using the PRO polypeptides of the present invention for a variety of uses based upon the functional biological assay data presented in the Examples below.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0027] FIG. 1 shows a nucleotide sequence (SEQ ID NO:1) of a native sequence PRO281 cDNA, wherein SEQ ID NO:1 is a clone designated herein as "DNA16422-1209".

[0028] FIG. 2 shows the amino acid sequence (SEQ ID NO:2) derived from the coding sequence of SEQ ID NO:1 shown in FIG. 1.

- [0029] FIG. 3 shows a nucleotide sequence (SEQ ID NO:3) of a native sequence PRO1560 cDNA, wherein SEQ ID NO:3 is a clone designated herein as "DNA19902-1669".
- [0030] FIG. 4 shows the amino acid sequence (SEQ ID NO:4) derived from the coding sequence of SEQ ID NO:3 shown in FIG. 3.
- [0031] FIG. 5 shows a nucleotide sequence (SEQ ID NO:5) of a native sequence PRO189 cDNA, wherein SEQ ID NO:5 is a clone designated herein as "DNA21624-1391".
- [0032] FIG. 6 shows the amino acid sequence (SEQ ID NO:6) derived from the coding sequence of SEQ ID NO:5 shown in FIG. 5.
- [0033] FIG. 7 shows a nucleotide sequence (SEQ ID NO:7) of a native sequence PRO240 cDNA, wherein SEQ ID NO:7 is a clone designated herein as "DNA34387-1138".
- [0034] FIG. 8 shows the amino acid sequence (SEQ ID NO:8) derived from the coding sequence of SEQ ID NO:7 shown in FIG. 7.
- [0035] FIG. 9 shows a nucleotide sequence (SEQ ID NO:9) of a native sequence PRO256 cDNA, wherein SEQ ID NO:9 is a clone designated herein as "DNA35880-1160".
- [0036] FIG. 10 shows the amino acid sequence (SEQ ID NO:10) derived from the coding sequence of SEQ ID NO:9 shown in FIG. 9.
- [0037] FIG. 11 shows a nucleotide sequence (SEQ ID NO:11) of a native sequence PRO306 cDNA, wherein SEQ ID NO:11 is a clone designated herein as "DNA39984-1221".
- [0038] FIG. 12 shows the amino acid sequence (SEQ ID NO:12) derived from the coding sequence of SEQ ID NO:11 shown in FIG. 11.
- [0039] FIG. 13 shows a nucleotide sequence (SEQ ID NO:13) of a native sequence PRO540 cDNA, wherein SEQ ID NO:13 is a clone designated herein as "DNA44189-1322".
- [0040] FIG. 14 shows the amino acid sequence (SEQ ID NO:14) derived from the coding sequence of SEQ ID NO:13 shown in FIG. 13.
- [0041] FIG. 15 shows a nucleotide sequence (SEQ ID NO:15) of a native sequence PRO773 cDNA, wherein SEQ ID NO:15 is a clone designated herein as "DNA48303-2829".
- [0042] FIG. 16 shows the amino acid sequence (SEQ ID NO:16) derived from the coding sequence of SEQ ID NO:15 shown in FIG. 15.
- [0043] FIG. 17 shows a nucleotide sequence (SEQ ID NO:17) of a native sequence PRO698 cDNA, wherein SEQ ID NO:17 is a clone designated herein as "DNA48320-1433".
- [0044] FIG. 18 shows the amino acid sequence (SEQ ID NO:18) derived from the coding sequence of SEQ ID NO:17 shown in FIG. 17.
- [0045] FIG. 19 shows a nucleotide sequence (SEQ ID NO:19) of a native sequence PRO3567 cDNA, wherein SEQ ID NO:19 is a clone designated herein as "DNA56049-2543".
- [0046] FIG. 20 shows the amino acid sequence (SEQ ID NO:20) derived from the coding sequence of SEQ ID NO:19 shown in FIG. 19.
- [0047] FIG. 21 shows a nucleotide sequence (SEQ ID NO:21) of a native sequence PRO826 cDNA, wherein SEQ ID NO:21 is a clone designated herein as "DNA57694-1341".
- [0048] FIG. 22 shows the amino acid sequence (SEQ ID NO:22) derived from the coding sequence of SEQ ID NO:21 shown in FIG. 21.
- [0049] FIG. 23 shows a nucleotide sequence (SEQ ID NO:23) of a native sequence PRO1002 cDNA, wherein SEQ ID NO:23 is a clone designated herein as "DNA59208-1373".
- [0050] FIG. 24 shows the amino acid sequence (SEQ ID NO:24) derived from the coding sequence of SEQ ID NO:23 shown in FIG. 23.
- [0051] FIG. 25 shows a nucleotide sequence (SEQ ID NO:25) of a native sequence PRO1068 cDNA, wherein SEQ ID NO:25 is a clone designated herein as "DNA59214-1449".
- [0052] FIG. 26 shows the amino acid sequence (SEQ ID NO:26) derived from the coding sequence of SEQ ID NO:25 shown in FIG. 25.
- [0053] FIG. 27 shows a nucleotide sequence (SEQ ID NO:27) of a native sequence PRO1030 cDNA, wherein SEQ ID NO:27 is a clone designated herein as "DNA59485-1336".
- [0054] FIG. 28 shows the amino acid sequence (SEQ ID NO:28) derived from the coding sequence of SEQ ID NO:27 shown in FIG. 27.
- [0055] FIG. 29 shows a nucleotide sequence (SEQ ID NO:29) of a native sequence PRO1313 cDNA, wherein SEQ ID NO:29 is a clone designated herein as "DNA64966-1575".
- [0056] FIG. 30 shows the amino acid sequence (SEQ ID NO:30) derived from the coding sequence of SEQ ID NO:29 shown in FIG. 29.
- [0057] FIG. 31 shows a nucleotide sequence (SEQ ID NO:31) of a native sequence PRO6071 cDNA, wherein SEQ ID NO:31 is a clone designated herein as "DNA82403-2959".
- [0058] FIG. 32 shows the amino acid sequence (SEQ ID NO:32) derived from the coding sequence of SEQ ID NO:31 shown in FIG. 31.
- [0059] FIG. 33 shows a nucleotide sequence (SEQ ID NO:33) of a native sequence PRO4397 cDNA, wherein SEQ ID NO:33 is a clone designated herein as "DNA83505-2606".
- [0060] FIG. 34 shows the amino acid sequence (SEQ ID NO:34) derived from the coding sequence of SEQ ID NO:33 shown in FIG. 33.
- [0061] FIG. 35 shows a nucleotide sequence (SEQ ID NO:35) of a native sequence PRO4344 cDNA, wherein SEQ ID NO:35 is a clone designated herein as "DNA84927-2585".

[0062] FIG. 36 shows the amino acid sequence (SEQ ID NO:36) derived from the coding sequence of SEQ ID NO:35 shown in FIG. 35.

[0063] FIG. 37 shows a nucleotide sequence (SEQ ID NO:37) of a native sequence PRO4407 cDNA, wherein SEQ ID NO:37 is a clone designated herein as "DNA92264-2616".

[0064] FIG. 38 shows the amino acid sequence (SEQ ID NO:38) derived from the coding sequence of SEQ ID NO:37 shown in FIG. 37.

[0065] FIG. 39 shows a nucleotide sequence (SEQ ID NO:39) of a native sequence PRO4316 cDNA, wherein SEQ ID NO:39 is a clone designated herein as "DNA94713-2561".

[0066] FIG. 40 shows the amino acid sequence (SEQ ID NO:40) derived from the coding sequence of SEQ ID NO:39 shown in FIG. 39.

[0067] FIG. 41 shows a nucleotide sequence (SEQ ID NO:41) of a native sequence PRO5775 cDNA, wherein SEQ ID NO:41 is a clone designated herein as "DNA96869-2673".

[0068] FIG. 42 shows the amino acid sequence (SEQ ID NO:42) derived from the coding sequence of SEQ ID NO:41 shown in FIG. 41.

[0069] FIG. 43 shows a nucleotide sequence (SEQ ID NO:43) of a native sequence PRO6016 cDNA, wherein SEQ ID NO:43 is a clone designated herein as "DNA96881-2699".

[0070] FIG. 44 shows the amino acid sequence (SEQ ID NO:44) derived from the coding sequence of SEQ ID NO:43 shown in FIG. 43.

[0071] FIG. 45 shows a nucleotide sequence (SEQ ID NO:45) of a native sequence PRO4499 cDNA, wherein SEQ ID NO:45 is a clone designated herein as "DNA96889-2641".

[0072] FIG. 46 shows the amino acid sequence (SEQ ID NO:46) derived from the coding sequence of SEQ ID NO:45 shown in FIG. 45.

[0073] FIG. 47 shows a nucleotide sequence (SEQ ID NO:47) of a native sequence PRO4487 cDNA, wherein SEQ ID NO:47 is a clone designated herein as "DNA96898-2640".

[0074] FIG. 48 shows the amino acid sequence (SEQ ID NO:48) derived from the coding sequence of SEQ ID NO:47 shown in FIG. 47.

[0075] FIG. 49 shows a nucleotide sequence (SEQ ID NO:49) of a native sequence PRO4980 cDNA, wherein SEQ ID NO:49 is a clone designated herein as "DNA97003-2649".

[0076] FIG. 50 shows the amino acid sequence (SEQ ID NO:50) derived from the coding sequence of SEQ ID NO:49 shown in FIG. 49.

[0077] FIG. 51 shows a nucleotide sequence (SEQ ID NO:51) of a native sequence PRO6018 cDNA, wherein SEQ ID NO:51 is a clone designated herein as "DNA98565-2701".

[0078] FIG. 52 shows the amino acid sequence (SEQ ID NO:52) derived from the coding sequence of SEQ ID NO:51 shown in FIG. 51.

[0079] FIG. 53 shows a nucleotide sequence (SEQ ID NO:53) of a native sequence PRO7168 cDNA, wherein SEQ ID NO:53 is a clone designated herein as "DNA102846-2742".

[0080] FIG. 54 shows the amino acid sequence (SEQ ID NO:54) derived from the coding sequence of SEQ ID NO:53 shown in FIG. 53.

[0081] FIG. 55 shows a nucleotide sequence (SEQ ID NO:55) of a native sequence PRO6308 cDNA, wherein SEQ ID NO:55 is a clone designated herein as "DNA102847-2726".

[0082] FIG. 56 shows the amino acid sequence (SEQ ID NO:56) derived from the coding sequence of SEQ ID NO:55 shown in FIG. 55.

[0083] FIG. 57 shows a nucleotide sequence (SEQ ID NO:57) of a native sequence PRO6000 cDNA, wherein SEQ ID NO:57 is a clone designated herein as "DNA102880-2689".

[0084] FIG. 58 shows the amino acid sequence (SEQ ID NO:58) derived from the coding sequence of SEQ ID NO:57 shown in FIG. 57.

[0085] FIG. 59 shows a nucleotide sequence (SEQ ID NO:59) of a native sequence PRO6006 cDNA, wherein SEQ ID NO:59 is a clone designated herein as "DNA105782-2693".

[0086] FIG. 60 shows the amino acid sequence (SEQ ID NO:60) derived from the coding sequence of SEQ ID NO:59 shown in FIG. 59.

[0087] FIG. 61 shows a nucleotide sequence (SEQ ID NO:61) of a native sequence PRO5800 cDNA, wherein SEQ ID NO:61 is a clone designated herein as "DNA108912-2680".

[0088] FIG. 62 shows the amino acid sequence (SEQ ID NO:62) derived from the coding sequence of SEQ ID NO:61 shown in FIG. 61.

[0089] FIG. 63 shows a nucleotide sequence (SEQ ID NO:63) of a native sequence PRO7476 cDNA, wherein SEQ ID NO:63 is a clone designated herein as "DNA115253-2757".

[0090] FIG. 64 shows the amino acid sequence (SEQ ID NO:64) derived from the coding sequence of SEQ ID NO:63 shown in FIG. 63.

[0091] FIG. 65 shows a nucleotide sequence (SEQ ID NO:65) of a native sequence PRO6496 cDNA, wherein SEQ ID NO:65 is a clone designated herein as "DNA119302-2737".

[0092] FIG. 66 shows the amino acid sequence (SEQ ID NO:66) derived from the coding sequence of SEQ ID NO:65 shown in FIG. 65.

[0093] FIG. 67 shows a nucleotide sequence (SEQ ID NO:67) of a native sequence PRO7422 cDNA, wherein SEQ ID NO:67 is a clone designated herein as "DNA119536-2752".

- [0094] FIG. 68 shows the amino acid sequence (SEQ ID NO:68) derived from the coding sequence of SEQ ID NO:67 shown in FIG. 67.
- [0095] FIG. 69 shows a nucleotide sequence (SEQ ID NO:69) of a native sequence PRO7431cDNA, wherein SEQ ID NO:69 is a clone designated herein as "DNA119542-2754".
- [0096] FIG. 70 shows the amino acid sequence (SEQ ID NO:70) derived from the coding sequence of SEQ ID NO:69 shown in FIG. 69.
- [0097] FIG. 71 shows a nucleotide sequence (SEQ ID NO:71) of a native sequence PRO10275 cDNA, wherein SEQ ID NO:71 is a clone designated herein as "DNA143498-2824".
- [0098] FIG. 72 shows the amino acid sequence (SEQ ID NO:72) derived from the coding sequence of SEQ ID NO:71 shown in FIG. 71.
- [0099] FIG. 73 shows a nucleotide sequence (SEQ ID NO:73) of a native sequence PRO10268 cDNA, wherein SEQ ID NO:73 is a clone designated herein as "DNA145583-2820".
- [0100] FIG. 74 shows the amino acid sequence (SEQ ID NO:74) derived from the coding sequence of SEQ ID NO:73 shown in FIG. 73.
- [0101] FIG. 75 shows a nucleotide sequence (SEQ ID NO:75) of a native sequence PRO20080 cDNA, wherein SEQ ID NO:75 is a clone designated herein as "DNA161000-2896".
- [0102] FIG. 76 shows the amino acid sequence (SEQ ID NO:76) derived from the coding sequence of SEQ ID NO:75 shown in FIG. 75.
- [0103] FIG. 77 shows a nucleotide sequence (SEQ ID NO:77) of a native sequence PRO21207 cDNA, wherein SEQ ID NO:77 is a clone designated herein as "DNA161005-2943".
- [0104] FIG. 78 shows the amino acid sequence (SEQ ID NO:78) derived from the coding sequence of SEQ ID NO:77 shown in FIG. 77.
- [0105] FIG. 79 shows a nucleotide sequence (SEQ ID NO:79) of a native sequence PRO28633 cDNA, wherein SEQ ID NO:79 is a clone designated herein as "DNA170245-3053".
- [0106] FIG. 80 shows the amino acid sequence (SEQ ID NO:80) derived from the coding sequence of SEQ ID NO:79 shown in FIG. 79.
- [0107] FIG. 81 shows a nucleotide sequence (SEQ ID NO:81) of a native sequence PRO20933 cDNA, wherein SEQ ID NO:81 is a clone designated herein as "DNA171771-2919".
- [0108] FIG. 82 shows the amino acid sequence (SEQ ID NO:82) derived from the coding sequence of SEQ ID NO:81 shown in FIG. 81.
- [0109] FIG. 83 shows a nucleotide sequence (SEQ ID NO:83) of a native sequence PRO21383 cDNA, wherein SEQ ID NO:83 is a clone designated herein as "DNA173157-2981".
- [0110] FIG. 84 shows the amino acid sequence (SEQ ID NO:84) derived from the coding sequence of SEQ ID NO:83 shown in FIG. 83.
- [0111] FIG. 85 shows a nucleotide sequence (SEQ ID NO:85) of a native sequence PRO21485 cDNA, wherein SEQ ID NO:85 is a clone designated herein as "DNA175734-2985".
- [0112] FIG. 86 shows the amino acid sequence (SEQ ID NO:86) derived from the coding sequence of SEQ ID NO:85 shown in FIG. 85.
- [0113] FIG. 87 shows a nucleotide sequence (SEQ ID NO:87) of a native sequence PRO28700 cDNA, wherein SEQ ID NO:87 is a clone designated herein as "DNA176108-3040".
- [0114] FIG. 88 shows the amino acid sequence (SEQ ID NO:88) derived from the coding sequence of SEQ ID NO:87 shown in FIG. 87.
- [0115] FIG. 89 shows a nucleotide sequence (SEQ ID NO:89) of a native sequence PRO34012 cDNA, wherein SEQ ID NO:89 is a clone designated herein as "DNA190710-3028".
- [0116] FIG. 90 shows the amino acid sequence (SEQ ID NO:90) derived from the coding sequence of SEQ ID NO:89 shown in FIG. 89.
- [0117] FIG. 91 shows a nucleotide sequence (SEQ ID NO:91) of a native sequence PRO34003 cDNA, wherein SEQ ID NO:91 is a clone designated herein as "DNA190803-3019".
- [0118] FIG. 92 shows the amino acid sequence (SEQ ID NO:92) derived from the coding sequence of SEQ ID NO:91 shown in FIG. 91.
- [0119] FIG. 93 shows a nucleotide sequence (SEQ ID NO:93) of a native sequence PRO34274 cDNA, wherein SEQ ID NO:93 is a clone designated herein as "DNA191064-3069".
- [0120] FIG. 94 shows the amino acid sequence (SEQ ID NO:94) derived from the coding sequence of SEQ ID NO:93 shown in FIG. 93.
- [0121] FIGS. 95A-95B shows a nucleotide sequence (SEQ ID NO:95) of a native sequence PRO34001 cDNA, wherein SEQ ID NO:95 is a clone designated herein as "DNA194909-3013".
- [0122] FIG. 96 shows the amino acid sequence (SEQ ID NO:96) derived from the coding sequence of SEQ ID NO:95 shown in FIGS. 95A-95B.
- [0123] FIG. 97 shows a nucleotide sequence (SEQ ID NO:97) of a native sequence PRO34009 cDNA, wherein SEQ ID NO:97 is a clone designated herein as "DNA203532-3029".
- [0124] FIG. 98 shows the amino acid sequence (SEQ ID NO:98) derived from the coding sequence of SEQ ID NO:97 shown in FIG. 97.
- [0125] FIG. 99 shows a nucleotide sequence (SEQ ID NO:99) of a native sequence PRO34192 cDNA, wherein SEQ ID NO:99 is a clone designated herein as "DNA213858-3060".

[0126] FIG. 100 shows the amino acid sequence (SEQ ID NO:100) derived from the coding sequence of SEQ ID NO:99 shown in FIG. 99.

[0127] FIG. 101 shows a nucleotide sequence (SEQ ID NO:101) of a native sequence PRO34564 cDNA, wherein SEQ ID NO:101 is a clone designated herein as "DNA216676-3083".

[0128] FIG. 102 shows the amino acid sequence (SEQ ID NO:102) derived from the coding sequence of SEQ ID NO:101 shown in FIG. 101.

[0129] FIG. 103 shows a nucleotide sequence (SEQ ID NO:103) of a native sequence PRO35444 cDNA, wherein SEQ ID NO:103 is a clone designated herein as "DNA222653-3104".

[0130] FIG. 104 shows the amino acid sequence (SEQ ID NO:104) derived from the coding sequence of SEQ ID NO:103 shown in FIG. 103.

[0131] FIG. 105 shows a nucleotide sequence (SEQ ID NO:105) of a native sequence PRO5998 cDNA, wherein SEQ ID NO:105 is a clone designated herein as "DNA96897-2688".

[0132] FIG. 106 shows the amino acid sequence (SEQ ID NO:106) derived from the coding sequence of SEQ ID NO:105 shown in FIG. 105.

[0133] FIG. 107 shows a nucleotide sequence (SEQ ID NO:107) of a native sequence PRO19651 cDNA, wherein SEQ ID NO:107 is a clone designated herein as "DNA142917-3081".

[0134] FIG. 108 shows the amino acid sequence (SEQ ID NO:108) derived from the coding sequence of SEQ ID NO:107 shown in FIG. 107.

[0135] FIG. 109 shows a nucleotide sequence (SEQ ID NO:109) of a native sequence PRO20221 cDNA, wherein SEQ ID NO:109 is a clone designated herein as "DNA142930-2914".

[0136] FIG. 110 shows the amino acid sequence (SEQ ID NO:110) derived from the coding sequence of SEQ ID NO:109 shown in FIG. 109.

[0137] FIG. 111 shows a nucleotide sequence (SEQ ID NO:111) of a native sequence PRO21434 cDNA, wherein SEQ ID NO:111 is a clone designated herein as "DNA147253-2983".

[0138] FIG. 112 shows the amino acid sequence (SEQ ID NO:112) derived from the coding sequence of SEQ ID NO:111 shown in FIG. 111.

[0139] FIG. 113 shows a nucleotide sequence (SEQ ID NO:113) of a native sequence PRO19822 cDNA, wherein SEQ ID NO:113 is a clone designated herein as "DNA149927-2887".

[0140] FIG. 114 shows the amino acid sequence (SEQ ID NO:114) derived from the coding sequence of SEQ ID NO:113 shown in FIG. 113.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

### [0141] I. Definitions

[0142] The terms "PRO polypeptide" and "PRO" as used herein and when immediately followed by a numerical designation refer to various polypeptides, wherein the complete designation (i.e., PRO/number) refers to specific polypeptide sequences as described herein. The terms "PRO/number polypeptide" and "PRO/number" wherein the term "number" is provided as an actual numerical designation as used herein encompass native sequence polypeptides and polypeptide variants (which are further defined herein). The PRO polypeptides described herein may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods. The term "PRO polypeptide" refers to each individual PRO/number polypeptide disclosed herein. All disclosures in this specification which refer to the "PRO polypeptide" refer to each of the polypeptides individually as well as jointly. For example, descriptions of the preparation of, purification of, derivation of, formation of antibodies to or against, administration of, compositions containing, treatment of a disease with, etc., pertain to each polypeptide of the invention individually. The term "PRO polypeptide" also includes variants of the PRO/number polypeptides disclosed herein.

[0143] A "native sequence PRO polypeptide" comprises a polypeptide having the same amino acid sequence as the corresponding PRO polypeptide derived from nature. Such native sequence PRO polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. The term "native sequence PRO polypeptide" specifically encompasses naturally-occurring truncated or secreted forms of the specific PRO polypeptide (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. In various embodiments of the invention, the native sequence PRO polypeptides disclosed herein are mature or full-length native sequence polypeptides comprising the full-length amino acids sequences shown in the accompanying figures. Start and stop codons are shown in bold font and underlined in the figures. However, while the PRO polypeptide disclosed in the accompanying figures are shown to begin with methionine residues designated herein as amino acid position 1 in the figures, it is conceivable and possible that other methionine residues located either upstream or downstream from the amino acid position 1 in the figures may be employed as the starting amino acid residue for the PRO polypeptides.

[0144] The PRO polypeptide "extracellular domain" or "ECD" refers to a form of the PRO polypeptide which is essentially free of the transmembrane and cytoplasmic domains. Ordinarily, a PRO polypeptide ECD will have less than 1% of such transmembrane and/or cytoplasmic domains and preferably, will have less than 0.5% of such domains. It will be understood that any transmembrane domains identified for the PRO polypeptides of the present invention are identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than about 5 amino

acids at either end of the domain as initially identified herein. Optionally, therefore, an extracellular domain of a PRO polypeptide may contain from about 5 or fewer amino acids on either side of the transmembrane domain/extracellular domain boundary as identified in the Examples or specification and such polypeptides, with or without the associated signal peptide, and nucleic acid encoding them, are contemplated by the present invention.

[0145] The approximate location of the “signal peptides” of the various PRO polypeptides disclosed herein are shown in the present specification and/or the accompanying figures. It is noted, however, that the C-terminal boundary of a signal peptide may vary, but most likely by no more than about 5 amino acids on either side of the signal peptide C-terminal boundary as initially identified herein, wherein the C-terminal boundary of the signal peptide may be identified pursuant to criteria routinely employed in the art for identifying that type of amino acid sequence element (e.g., Nielsen et al., *Prot. Eng.* 10:1-6 (1997) and von Heinje et al., *Nucl. Acids. Res.* 14:4683-4690 (1986)). Moreover, it is also recognized that, in some cases, cleavage of a signal sequence from a secreted polypeptide is not entirely uniform, resulting in more than one secreted species. These mature polypeptides, where the signal peptide is cleaved within no more than about 5 amino acids on either side of the C-terminal boundary of the signal peptide as identified herein, and the polynucleotides encoding them, are contemplated by the present invention.

[0146] “PRO polypeptide variant” means an active PRO polypeptide as defined above or below having at least about 80% amino acid sequence identity with a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Such PRO polypeptide variants include, for instance, PRO polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of the full-length native amino acid sequence. Ordinarily, a PRO polypeptide variant will have at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a full-length native sequence PRO polypeptide sequence as disclosed herein, a

PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, PRO variant polypeptides are at least about 10 amino acids in length, alternatively at least about 20 amino acids in length, alternatively at least about 30 amino acids in length, alternatively at least about 40 amino acids in length, alternatively at least about 50 amino acids in length, alternatively at least about 60 amino acids in length, alternatively at least about 70 amino acids in length, alternatively at least about 80 amino acids in length, alternatively at least about 90 amino acids in length, alternatively at least about 100 amino acids in length, alternatively at least about 150 amino acids in length, alternatively at least about 200 amino acids in length, alternatively at least about 300 amino acids in length, or more.

[0147] “Percent (%) amino acid sequence identity” with respect to the PRO polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific PRO polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, Calif. or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

[0148] In situations where ALIGN-2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

[0149] where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B,



and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations using this method, Tables 2 and 3 demonstrate how to calculate the % amino acid sequence identity of the amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO", wherein "PRO" represents the amino acid sequence of a hypothetical PRO polypeptide of interest, "Comparison Protein" represents the amino acid sequence of a polypeptide against which the "PRO" polypeptide of interest is being compared, and "X", "Y" and "Z" each represent different hypothetical amino acid residues.

**[0150]** Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % amino acid sequence identity values may also be obtained as described below by using the WU-BLAST-2 computer program (Altschul et al., *Methods in Enzymology* 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=11, and scoring matrix=BLOSUM62. When WU-BLAST-2 is employed, a % amino acid sequence identity value is determined by dividing (a) the number of matching identical amino acid residues between the amino acid sequence of the PRO polypeptide of interest having a sequence derived from the native PRO polypeptide and the comparison amino acid sequence of interest (i.e., the sequence against which the PRO polypeptide of interest is being compared which may be a PRO variant polypeptide) as determined by WU-BLAST-2 by (b) the total number of amino acid residues of the PRO polypeptide of interest. For example, in the statement "a polypeptide comprising an amino acid sequence A which has or having at least 80% amino acid sequence identity to the amino acid sequence B", the amino acid sequence A is the comparison amino acid sequence of interest and the amino acid sequence B is the amino acid sequence of the PRO polypeptide of interest.

**[0151]** Percent amino acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov> or otherwise obtained from the National Institute of Health, Bethesda, Md. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask=yes, strand=all, expected occurrences=10, minimum low complexity length=15/5, multi-pass e-value=0.01, constant for multi-pass=25, dropoff for final gapped alignment=25 and scoring matrix=BLOSUM62.

**[0152]** In situations where NCBI-BLAST2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A

that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

**[0153]** where X is the number of amino acid residues scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A.

**[0154]** "PRO variant polynucleotide" or "PRO variant nucleic acid sequence" means a nucleic acid molecule which encodes an active PRO polypeptide as defined below and which has at least about 80% nucleic acid sequence identity with a nucleotide acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, a PRO variant polynucleotide will have at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity with a nucleic acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal sequence, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Variants do not encompass the native nucleotide sequence.

**[0155]** Ordinarily, PRO variant polynucleotides are at least about 30 nucleotides in length, alternatively at least about 60 nucleotides in length, alternatively at least about 90 nucleotides in length, alternatively at least about 120 nucleotides in length, alternatively at least about 150 nucleotides in length, alternatively at least about 180 nucleotides in length, alternatively at least about 210 nucleotides in length, alternatively at least about 240 nucleotides in length, alter-

natively at least about 270 nucleotides in length, alternatively at least about 300 nucleotides in length, alternatively at least about 450 nucleotides in length, alternatively at least about 600 nucleotides in length, alternatively at least about 900 nucleotides in length, or more.

[0156] “Percent (%) nucleic acid sequence identity” with respect to PRO-encoding nucleic acid sequences identified herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in the PRO nucleic acid sequence of interest, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. For purposes herein, however, % nucleic acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, Calif. or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

[0157] In situations where ALIGN-2 is employed for nucleic acid sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

100 times the fraction  $W/Z$

[0158] where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program’s alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of % nucleic acid sequence identity calculations, Tables 4 and 5, demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated “Comparison DNA” to the nucleic acid sequence designated “PRO-DNA”, wherein “PRO-DNA” represents a hypothetical PRO-encoding nucleic acid sequence of interest, “Comparison DNA” represents the nucleotide sequence of a nucleic acid molecule against which the “PRO-DNA” nucleic acid molecule of interest is being compared, and “N”, “L” and “V” each represent different hypothetical nucleotides.

[0159] Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as

described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % nucleic acid sequence identity values may also be obtained as described below by using the WU-BLAST-2 computer program (Altschul et al., *Methods in Enzymology* 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=11, and scoring matrix=BLOSUM62. When WU-BLAST-2 is employed, a % nucleic acid sequence identity value is determined by dividing (a) the number of matching identical nucleotides between the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest having a sequence derived from the native sequence PRO polypeptide-encoding nucleic acid and the comparison nucleic acid molecule of interest (i.e., the sequence against which the PRO polypeptide-encoding nucleic acid molecule of interest is being compared which may be a variant PRO polynucleotide) as determined by WU-BLAST-2 by (b) the total number of nucleotides of the PRO polypeptide-encoding nucleic acid molecule of interest. For example, in the statement “an isolated nucleic acid molecule comprising a nucleic acid sequence A which has or having at least 80% nucleic acid sequence identity to the nucleic acid sequence B”, the nucleic acid sequence A is the comparison nucleic acid molecule of interest and the nucleic acid sequence B is the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest.

[0160] Percent nucleic acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov> or otherwise obtained from the National Institute of Health, Bethesda, Md. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask=yes, strand=all, expected occurrences=10, minimum low complexity length=15/5, multi-pass e-value=0.01, constant for multi-pass=25, dropoff for final gapped alignment=25 and scoring matrix=BLOSUM62.

[0161] In situations where NCBI-BLAST2 is employed for sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

100 times the fraction  $W/Z$

[0162] where W is the number of nucleotides scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program’s alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C.

[0163] In other embodiments, PRO variant polynucleotides are nucleic acid molecules that encode an active PRO polypeptide and which are capable of hybridizing, prefer-

ably under stringent hybridization and wash conditions, to nucleotide sequences encoding a full-length PRO polypeptide as disclosed herein. PRO variant polypeptides may be those that are encoded by a PRO variant polynucleotide.

[0164] “Isolated,” when used to describe the various polypeptides disclosed herein, means polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide in situ within recombinant cells, since at least one component of the PRO polypeptide natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

[0165] An “isolated” PRO polypeptide-encoding nucleic acid or other polypeptide-encoding nucleic acid is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the polypeptide-encoding nucleic acid. An isolated polypeptide-encoding nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated polypeptide-encoding nucleic acid molecules therefore are distinguished from the specific polypeptide-encoding nucleic acid molecule as it exists in natural cells. However, an isolated polypeptide-encoding nucleic acid molecule includes polypeptide-encoding nucleic acid molecules contained in cells that ordinarily express the polypeptide where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

[0166] The term “control sequences” refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

[0167] Nucleic acid is “operably linked” when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, “operably linked” means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

[0168] The term “antibody” is used in the broadest sense and specifically covers, for example, single anti-PRO mono-

clonal antibodies (including agonist, antagonist, and neutralizing antibodies), anti-PRO antibody compositions with polypeptopic specificity, single chain anti-PRO antibodies, and fragments of anti-PRO antibodies (see below). The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts.

[0169] “Stringency” of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature which can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel et al., *Current Protocols in Molecular Biology*, Wiley Interscience Publishers, (1995).

[0170] “Stringent conditions” or “high stringency conditions”, as defined herein, may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50° C.; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42° C.; or (3) employ 50% formamide, 5×SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5×Denhardt’s solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42° C., with washes at 42° C. in 0.2×SSC (sodium chloride/sodium citrate) and 50% formamide at 55° C., followed by a high-stringency wash consisting of 0.1×SSC containing EDTA at 55° C.

[0171] “Moderately stringent conditions” may be identified as described by Sambrook et al., *Molecular Cloning: A Laboratory Manual*, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and % SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37° C. in a solution comprising: 20% formamide, 5×SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5×Denhardt’s solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1×SSC at about 37-50° C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

[0172] The term “epitope tagged” when used herein refers to a chimeric polypeptide comprising a PRO polypeptide

fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

[0173] As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM.

[0174] "Active" or "activity" for the purposes herein refers to form(s) of a PRO polypeptide which retain a biological and/or an immunological activity of native or naturally-occurring PRO, wherein "biological" activity refers to a biological function (either inhibitory or stimulatory) caused by a native or naturally-occurring PRO other than the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO and an "immunological" activity refers to the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO.

[0175] The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of a native PRO polypeptide disclosed herein. In a similar manner, the term "agonist" is used in the broadest sense and includes any molecule that mimics a biological activity of a native PRO polypeptide disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments or amino acid sequence variants of native PRO polypeptides, peptides, antisense oligonucleotides, small organic molecules, etc. Methods for identifying agonists or antagonists of a PRO polypeptide may comprise contacting a PRO polypeptide with a candidate agonist or antagonist molecule and measuring a detectable change in one or more biological activities normally associated with the PRO polypeptide.

[0176] "Treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented.

[0177] "Chronic" administration refers to administration of the agent(s) in a continuous mode as opposed to an acute

mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time. "Intermittent" administration is treatment that is not consecutively done without interruption, but rather is cyclic in nature.

[0178] "Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. Preferably, the mammal is human.

[0179] Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

[0180] "Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are non-toxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, polyethylene glycol (PEG), and PLURONICS™.

[0181] "Antibody fragments" comprise a portion of an intact antibody, preferably the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')<sub>2</sub>, and Fv fragments; diabodies; linear antibodies (Zapata et al., *Protein Eng.* 8(10): 1057-1062 [1995]); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

[0182] Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, a designation reflecting the ability to crystallize readily. Pepsin treatment yields an F(ab')<sub>2</sub> fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

[0183] "Fv" is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the V<sub>H</sub>-V<sub>L</sub> dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

[0184] The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab fragments differ from Fab' fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or

more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')<sub>2</sub> antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

[0185] The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and lambda, based on the amino acid sequences of their constant domains.

[0186] Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2.

[0187] "Single-chain Fv" or "sFv" antibody fragments comprise the V<sub>H</sub> and V<sub>L</sub> domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V<sub>H</sub> and V<sub>L</sub> domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv, see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

[0188] The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (V<sub>H</sub>) connected to a light-chain variable domain (V<sub>L</sub>) in the same polypeptide chain (V<sub>H</sub>-V<sub>L</sub>). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993).

[0189] An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or

internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody in situ within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

[0190] An antibody that "specifically binds to" or is "specific for" a particular polypeptide or an epitope on a particular polypeptide is one that binds to that particular polypeptide or epitope on a particular polypeptide without substantially binding to any other polypeptide or polypeptide epitope.

[0191] The word "label" when used herein refers to a detectable compound or composition which is conjugated directly or indirectly to the antibody so as to generate a "labeled" antibody. The label may be detectable by itself (e.g. radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable.

[0192] By "solid phase" is meant a non-aqueous matrix to which the antibody of the present invention can adhere. Examples of solid phases encompassed herein include those formed partially or entirely of glass (e.g., controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g., an affinity chromatography column). This term also includes a discontinuous solid phase of discrete particles, such as those described in U.S. Pat. No. 4,275,149.

[0193] A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as a PRO polypeptide or antibody thereto) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

[0194] A "small molecule" is defined herein to have a molecular weight below about 500 Daltons.

[0195] An "effective amount" of a polypeptide disclosed herein or an agonist or antagonist thereof is an amount sufficient to carry out a specifically stated purpose. An "effective amount" may be determined empirically and in a routine manner, in relation to the stated purpose.

**Table 1**

```

/*
 *
 * C-C increased from 12 to 15
 * Z is average of EQ
 5 * B is average of ND
 * match with stop is _M; stop-stop = 0; J (joker) match = 0
 */
#define _M -8 /* value of a match with a stop */

) int _day[26][26] = {
/* A B C D E F G H I J K L M N O P Q R S T U V W X Y Z */
/* A */ { 2, 0, -2, 0, 0, -4, 1, -1, -1, 0, -1, -2, -1, 0, _M, 1, 0, -2, 1, 1, 0, 0, -6, 0, -3, 0},
/* B */ { 0, 3, -4, 3, 2, -5, 0, 1, -2, 0, 0, -3, -2, 2, _M, -1, 1, 0, 0, 0, 0, -2, -5, 0, -3, 1},
/* C */ {-2, -4, 15, -5, -5, -4, -3, -3, -2, 0, -5, -6, -5, -4, _M, -3, -5, -4, 0, -2, 0, -2, -8, 0, 0, -5},
5 /* D */ { 0, 3, -5, 4, 3, -6, 1, 1, -2, 0, 0, -4, -3, 2, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 2},
/* E */ { 0, 2, -5, 3, 4, -5, 0, 1, -2, 0, 0, -3, -2, 1, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 3},
/* F */ {-4, -5, -4, -6, -5, 9, -5, -2, 1, 0, -5, 2, 0, -4, _M, -5, -5, -4, -3, -3, 0, -1, 0, 0, 7, -5},
/* G */ { 1, 0, -3, 1, 0, -5, 5, -2, -3, 0, -2, -4, -3, 0, _M, -1, -1, -3, 1, 0, 0, -1, -7, 0, -5, 0},
/* H */ {-1, 1, -3, 1, 1, -2, -2, 6, -2, 0, 0, -2, -2, 2, _M, 0, 3, 2, -1, -1, 0, -2, -3, 0, 0, 2},
) /* I */ {-1, -2, -2, -2, -2, 1, -3, -2, 5, 0, -2, 2, 2, -2, _M, -2, -2, -2, -1, 0, 0, 4, -5, 0, -1, -2},
/* J */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* K */ {-1, 0, -5, 0, 0, -5, -2, 0, -2, 0, 5, -3, 0, 1, _M, -1, 1, 3, 0, 0, 0, -2, -3, 0, -4, 0},
/* L */ {-2, -3, -6, -4, -3, 2, -4, -2, 2, 0, -3, 6, 4, -3, _M, -3, -2, -3, -3, -1, 0, 2, -2, 0, -1, -2},
/* M */ {-1, -2, -5, -3, -2, 0, -3, -2, 2, 0, 0, 4, 6, -2, _M, -2, -1, 0, -2, -1, 0, 2, -4, 0, -2, -1},
5 /* N */ { 0, 2, -4, 2, 1, -4, 0, 2, -2, 0, 1, -3, -2, 2, _M, -1, 1, 0, 1, 0, 0, -2, -4, 0, -2, 1},
/* O */ { _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M},
/* P */ { 1, -1, -3, -1, -1, -5, -1, 0, -2, 0, -1, -3, -2, -1, _M, 6, 0, 0, 1, 0, 0, -1, -6, 0, -5, 0},
/* Q */ { 0, 1, -5, 2, 2, -5, -1, 3, -2, 0, 1, -2, -1, 1, _M, 0, 4, 1, -1, -1, 0, -2, -5, 0, -4, 3},
/* R */ {-2, 0, -4, -1, -1, -4, -3, 2, -2, 0, 3, -3, 0, 0, _M, 0, 1, 6, 0, -1, 0, -2, 2, 0, -4, 0},
) /* S */ { 1, 0, 0, 0, 0, -3, 1, -1, -1, 0, 0, -3, -2, 1, _M, 1, -1, 0, 2, 1, 0, -1, -2, 0, -3, 0},
/* T */ { 1, 0, -2, 0, 0, -3, 0, -1, 0, 0, 0, -1, -1, 0, _M, 0, -1, -1, 1, 3, 0, 0, -5, 0, -3, 0},
/* U */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* V */ { 0, -2, -2, -2, -2, -1, -1, -2, 4, 0, -2, 2, 2, -2, _M, -1, -2, -2, -1, 0, 0, 4, -6, 0, -2, -2},
/* W */ {-6, -5, -8, -7, -7, 0, -7, -3, -5, 0, -3, -2, -4, -4, _M, -6, -5, 2, -2, -5, 0, -6, 17, 0, 0, -6},
5 /* X */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* Y */ {-3, -3, 0, -4, -4, 7, -5, 0, -1, 0, -4, -1, -2, -2, _M, -5, -4, -4, -3, -3, 0, -2, 0, 0, 10, -4},
/* Z */ { 0, 1, -5, 2, 3, -5, 0, 2, -2, 0, 0, -2, -1, 1, _M, 0, 3, 0, 0, 0, 0, -2, -6, 0, -4, 4}
};

```

**Table 1 (cont')**

```

/*
*/
#include <stdio.h>
#include <ctype.h>
5
#define MAXJMP      16      /* max jumps in a diag */
#define MAXGAP      24      /* don't continue to penalize gaps larger than this */
#define JMPS        1024    /* max jmps in an path */
0
#define MX          4       /* save if there's at least MX-1 bases since last jmp */

#define DMAT        3       /* value of matching bases */
#define DMIS        0       /* penalty for mismatched bases */
#define DINS0       8       /* penalty for a gap */
5
#define DINS1       1       /* penalty per base */
#define PINS0       8       /* penalty for a gap */
#define PINS1       4       /* penalty per residue */

struct jmp {
0
    short           n[MAXJMP]; /* size of jmp (neg for dely) */
    unsigned short x[MAXJMP]; /* base no. of jmp in seq x */
}; /* limits seq to 2^16 -1 */

struct diag {
5
    int             score;     /* score at last jmp */
    long            offset;    /* offset of prev block */
    short           ijmp;     /* current jmp index */
    struct jmp      jp;       /* list of jmps */
};

0
struct path {
    int             spc;       /* number of leading spaces */
    short           n[JMPS]; /* size of jmp (gap) */
    int             x[JMPS]; /* loc of jmp (last elem before gap) */
5
};

char             *ofile;     /* output file name */
char             *namex[2];  /* seq names: getseqs() */
char             *prog;     /* prog name for err msgs */
char             *seqx[2];   /* seqs: getseqs() */
0
int              dmax;       /* best diag: nw() */
int              dmax0;     /* final diag */
int              dna;       /* set if dna: main() */
int              endgaps;   /* set if penalizing end gaps */
int              gapx, gapy; /* total gaps in seqs */
5
int              len0, len1; /* seq lens */
int              ngapx, ngapy; /* total size of gaps */
int              smax;     /* max score: nw() */
int              *xbm;     /* bitmap for matching */
long             offset;    /* current offset in jmp file */
0
struct           diag      *dx; /* holds diagonals */
struct           path      pp[2]; /* holds path for seqs */

char             *calloc(), *malloc(), *index(), *strcpy();
5
char             *getseq(), *g_calloc();

```

**Table 1 (cont')**

```

/* Needleman-Wunsch alignment program
*
* usage: prog file1 file2
* where file1 and file2 are two dna or two protein sequences.
* The sequences can be in upper- or lower-case and may contain ambiguity
* Any lines beginning with ';', '>' or '<' are ignored
* Max file length is 65535 (limited by unsigned short x in the jmp struct)
* A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
* Output is in the file "align.out"
*
* The program may create a tmp file in /tmp to hold info about traceback.
* Original version developed under BSD 4.3 on a vax 8650
*/
#include "nw.h"
#include "day.h"

static _dbval[26] = {
    1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};

static _pbval[26] = {
    1, 2|(1<<('D'-'A'))|(1<<('N'-'A')), 4, 8, 16, 32, 64,
    128, 256, 0xFFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
    1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
    1<<23, 1<<24, 1<<25|(1<<('E'-'A'))|(1<<('Q'-'A'))
};

main(ac, av)
int ac;
char *av[];
{
    prog = av[0];
    if (ac != 3) {
        fprintf(stderr, "usage: %s file1 file2\n", prog);
        fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
        fprintf(stderr, "The sequences can be in upper- or lower-case\n");
        fprintf(stderr, "Any lines beginning with ';', '>' or '<' are ignored\n");
        fprintf(stderr, "Output is in the file \"align.out\"\n");
        exit(1);
    }
    namex[0] = av[1];
    namex[1] = av[2];
    seqx[0] = getseq(namex[0], &len0);
    seqx[1] = getseq(namex[1], &len1);
    xbm = (dna)? _dbval : _pbval;

    endgaps = 0; /* 1 to penalize endgaps */
    ofile = "align.out"; /* output file */

    nw(); /* fill in the matrix, get the possible jmps */
    readjmps(); /* get the actual jmps */
    print(); /* print stats, alignment */

    cleanup(0); /* unlink any tmp files */
}

```



**Table 1 (cont')**

```

/* do the alignment, return best score: main()
* dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
* pro: PAM 250 values
* When scores are equal, we prefer mismatches to any gap, prefer
* a new gap to extending an ongoing gap, and prefer a gap in seqx
* to a gap in seq y.
*/
nw()
{
    char          *px, *py;          /* seqs and ptrs */
    int           *ndely, *dely;     /* keep track of dely */
    int           ndelx, delx;       /* keep track of delx */
    int           *tmp;              /* for swapping row0, row1 */
    int           mis;               /* score for each type */
    int           ins0, ins1;        /* insertion penalties */
    register      id;                /* diagonal index */
    register      ij;                /* jmp index */
    register      *col0, *col1;      /* score for curr, last row */
    register      xx, yy;            /* index into seqs */

    dx = (struct diag *)g_calloc("to get diags", len0+len1+1, sizeof(struct diag));

    ndely = (int *)g_calloc("to get ndely", len1+1, sizeof(int));
    dely = (int *)g_calloc("to get dely", len1+1, sizeof(int));
    col0 = (int *)g_calloc("to get col0", len1+1, sizeof(int));
    col1 = (int *)g_calloc("to get col1", len1+1, sizeof(int));
    ins0 = (dna)? DINS0 : PINS0;
    ins1 = (dna)? DINS1 : PINS1;

    smax = -10000;
    if (endgaps) {
        for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
            col0[yy] = dely[yy] = col0[yy-1] - ins1;
            ndely[yy] = yy;
        }
        col0[0] = 0;          /* Waterman Bull Math Biol 84 */
    }
    else
        for (yy = 1; yy <= len1; yy++)
            dely[yy] = -ins0;

    /* fill in match matrix
    */
    for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
        /* initialize first entry in col
        */
        if (endgaps) {
            if (xx == 1)
                col1[0] = delx = -(ins0+ins1);
            else
                col1[0] = delx = col0[0] - ins1;
            ndelx = xx;
        }
        else {
            col1[0] = 0;
            delx = -ins0;
            ndelx = 0;
        }
    }
}

```

**Table 1 (cont')****...NW**

```

for (py = seqx[1], yy = 1; yy <= len1; py++, yy++) {
    mis = col0[yy-1];
    if (dna)
        mis += (xbrm[*px-'A']&xbrm[*py-'A'])? DMAT : DMIS;
    else
        mis += _day[*px-'A'][*py-'A'];

    /* update penalty for del in x seq;
     * favor new del over ongong del
     * ignore MAXGAP if weighting endgaps
     */
    if (endgaps || ndely[yy] < MAXGAP) {
        if (col0[yy] - ins0 >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0+ins1);
            ndely[yy] = 1;
        } else {
            dely[yy] -= ins1;
            ndely[yy]++;
        }
    } else {
        if (col0[yy] - (ins0+ins1) >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0+ins1);
            ndely[yy] = 1;
        } else
            ndely[yy]++;
    }

    /* update penalty for del in y seq;
     * favor new del over ongong del
     */
    if (endgaps || ndelx < MAXGAP) {
        if (col1[yy-1] - ins0 >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else {
            delx -= ins1;
            ndelx++;
        }
    } else {
        if (col1[yy-1] - (ins0+ins1) >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else
            ndelx++;
    }

    /* pick the maximum score; we're favoring
     * mis over any del and delx over dely
     */
}

```

**Table 1 (cont')**

...nw

```

id = xx - yy + len1 - 1;
if (mis >= delx && mis >= dely[yy])
    coll[yy] = mis;
else if (delx >= dely[yy]) {
    coll[yy] = delx;
    ij = dx[id].ijmp;
    if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writejmps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = ndelx;
    dx[id].jp.x[ij] = xx;
    dx[id].score = delx;
}
else {
    coll[yy] = dely[yy];
    ij = dx[id].ijmp;
if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writejmps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = -ndely[yy];
    dx[id].jp.x[ij] = xx;
    dx[id].score = dely[yy];
}
if (xx == len0 && yy < len1) {
    /* last col
    */
    if (endgaps)
        coll[yy] -= ins0+ins1*(len1-yy);
    if (coll[yy] > smax) {
        smax = coll[yy];
        dmax = id;
    }
}
}
if (endgaps && xx < len0)
    coll[yy-1] -= ins0+ins1*(len0-xx);
if (coll[yy-1] > smax) {
    smax = coll[yy-1];
    dmax = id;
}
}
tmp = col0; col0 = coll; coll = tmp;
}
(void) free((char *)ndely);
(void) free((char *)dely);
(void) free((char *)col0);
(void) free((char *)coll);
}

```

**Table 1 (cont')**

```

/*
 *
 * print() -- only routine visible outside this module
 *
5  * static:
 * getmat() -- trace back best path, count matches: print()
 * pr_align() -- print alignment of described in array p[]: print()
 * dumpblock() -- dump a block of lines with numbers, stars: pr_align()
 * nums() -- put out a number line: dumpblock()
0  * putline() -- put out a line (name, [num], seq, [num]): dumpblock()
 * stars() - -put a line of stars: dumpblock()
 * stripname() -- strip any path and prefix from a seqname
 */

5  #include "nw.h"

#define SPC      3
#define P_LINE  256 /* maximum output line */
#define P_SPC   3   /* space between name or num and seq */

0  extern _day[26][26];
int olen; /* set output line length */
FILE *fx; /* output file */

5  print() print
{
    int lx, ly, firstgap, lastgap; /* overlap */

    if ((fx = fopen(ofile, "w")) == 0) {
0      fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
    }
    fprintf(fx, "<first sequence: %s (length = %d)\n", namex[0], len0);
    fprintf(fx, "<second sequence: %s (length = %d)\n", namex[1], len1);
5    olen = 60;
    lx = len0;
    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) { /* leading gap in x */
0        pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
    }
    else if (dmax > len1 - 1) { /* leading gap in y */
5        pp[1].spc = firstgap = dmax - (len1 - 1);
        lx -= pp[1].spc;
    }
    if (dmax0 < len0 - 1) { /* trailing gap in x */
0        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
    }
    else if (dmax0 > len0 - 1) { /* trailing gap in y */
        lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
5    getmat(lx, ly, firstgap, lastgap);
    pr_align();
}

```

**Table 1 (cont')**

```

/*
 * trace back the best path, count matches
 */
static
5 getmat(lx, ly, firstgap, lastgap)
    int    lx, ly;          /* "core" (minus endgaps) */
    int    firstgap, lastgap; /* leading trailing overlap */
{
    int    nm, i0, i1, siz0, siz1;
0    char    outx[32];
    double    pct;
    register    n0, n1;
    register char    *p0, *p1;

5    /* get total matches, score
    */
    i0 = i1 = siz0 = siz1 = 0;
    p0 = seqx[0] + pp[1].spc;
    p1 = seqx[1] + pp[0].spc;
0    n0 = pp[1].spc + 1;
    n1 = pp[0].spc + 1;

    nm = 0;
5    while ( *p0 && *p1 ) {
        if (siz0) {
            p1++;
            n1++;
            siz0--;
        }
0        else if (siz1) {
            p0++;
            n0++;
            siz1--;
        }
5        else {
            if (xbm[*p0-'A']&xbm[*p1-'A'])
                nm++;
            if (n0++ == pp[0].x[i0])
                siz0 = pp[0].n[i0++];
0            if (n1++ == pp[1].x[i1])
                siz1 = pp[1].n[i1++];
            p0++;
            p1++;
        }
5    }

    /* pct homology:
    * if penalizing endgaps, base is the shorter seq
    * else, knock off overhangs and take shorter core
    */
    if (endgaps)
        lx = (len0 < len1)? len0 : len1;
    else
        lx = (lx < ly)? lx : ly;
5    pct = 100.*((double)nm)/((double)lx);
    fprintf(fx, "\n");
    fprintf(fx, "< %d match%s in an overlap of %d: %.2f percent similarity\n",
        nm, (nm == 1)? "" : "es", lx, pct);
)

```

**getmat**

**Table 1 (cont')**

```

fprintf(fx, "< gaps in first sequence: %d", gapx);
if (gapx) {
    (void) sprintf(outh, "(%d %s%s)",
        ngapx, (dna)? "base":"residue", (ngapx == 1)? "" : "s");
5     fprintf(fx, "%s", outh);

    fprintf(fx, ", gaps in second sequence: %d", gapy);
    if (gapy) {
        (void) sprintf(outh, "(%d %s%s)",
0         ngapy, (dna)? "base":"residue", (ngapy == 1)? "" : "s");
        fprintf(fx, "%s", outh);
    }
    if (dna)
5     fprintf(fx,
        "\n< score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
        smax, DMAT, DMIS, DINSO, DINS1);
    else
        fprintf(fx,
0         "\n< score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
        smax, PINSO, PINS1);
    if (endgaps)
        fprintf(fx,
5         "< endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
        firstgap, (dna)? "base" : "residue", (firstgap == 1)? "" : "s",
        lastgap, (dna)? "base" : "residue", (lastgap == 1)? "" : "s");
    else
        fprintf(fx, "< endgaps not penalized\n");
}

0     static      nm;          /* matches in core -- for checking */
    static      lmax;        /* lengths of stripped file names */
    static      ij[2];       /* jmp index for a path */
    static      nc[2];       /* number at start of current line */
    static      ni[2];       /* current elem number -- for gapping */
5     static      siz[2];
    static char *ps[2];      /* ptr to current element */
    static char *po[2];      /* ptr to next output char slot */
    static char out[2][P_LINE]; /* output line */
    static char star[P_LINE]; /* set by stars() */
0
    /*
    * print alignment of described in struct path pp[]
    */
    static
5     pr_align()
    {
        int      nn;         /* char count */
        int      more;
        register i;

0         for (i = 0, lmax = 0; i < 2; i++) {
            nn = stripname(namex[i]);
            if (nn > lmax)
5                 lmax = nn;

            nc[i] = 1;
            ni[i] = 1;
            siz[i] = ij[i] = 0;
            ps[i] = seqx[i];
            po[i] = out[i];
        }
}

```

...getmat

pr\_align

**Table 1 (cont')**

```

for (nn = nm = 0, more = 1; more;) {
    for (i = more = 0; i < 2; i++) {
        /*
         * do we have more of this sequence?
         */
        if (!*ps[i])
            continue;

        more++;

        if (pp[i].spc) { /* leading space */
            *po[i]++ = ' ';
            pp[i].spc--;
        }
        else if (siz[i]) { /* in a gap */
            *po[i]++ = '-';
            siz[i]--;
        }
        else {
            /* we're putting a seq element
             */
            *po[i] = *ps[i];
            if (islower(*ps[i]))
                *ps[i] = toupper(*ps[i]);
            po[i]++;
            ps[i]++;

            /*
             * are we at next gap for this seq?
             */
            if (ni[i] == pp[i].x[ij[i]]) {
                /*
                 * we need to merge all gaps
                 * at this location
                 */
                siz[i] = pp[i].n[ij[i] + +];
                while (ni[i] == pp[i].x[ij[i]])
                    siz[i] += pp[i].n[ij[i] + +];
            }
            ni[i]++;
        }
    }
    if (++nn == olen || !more && nn) {
        dumpblock();
        for (i = 0; i < 2; i++)
            po[i] = out[i];
        nn = 0;
    }
}

/*
 * dump a block of lines, including numbers, stars: pr_align()
 */
static
dumpblock()
{
    register i;

    for (i = 0; i < 2; i++)
        *po[i]-- = '\0';
}

```

...pr\_align

dumpblock





**Table 1 (cont')**

```

5      int          i;
      register char *px;

      for (px = namex[ix], i = 0; *px && *px != ':'; px++, i++)
          (void) putc(*px, fx);
0      for (; i < lmax+P_SPC; i++)
          (void) putc(' ', fx);

      /* these count from 1:
      * ni[] is current element (from 1)
      * nc[] is number at start of current line
      */
5      for (px = out[ix]; *px; px++)
          (void) putc(*px&0x7F, fx);
      (void) putc('\n', fx);
  }

  /*
  * put a line of stars (seqs always in out[0], out[1]): dumpblock()
  */
5  static
  stars()
  {
      int          i;
      register char *p0, *p1, cx, *px;

0      if (!*out[0] || (*out[0] == ' ' && *(po[0]) == ' ') ||
          !*out[1] || (*out[1] == ' ' && *(po[1]) == ' '))
          return;
      px = star;
5      for (i = lmax+P_SPC; i; i--)
          *px++ = ' ';

      for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
          if (isalpha(*p0) && isalpha(*p1)) {
0              if (xbm[*p0-'A']&xbm[*p1-'A']) {
                  cx = '*';
                  nm++;
              }
5              else if (!dna && _day[*p0-'A'][*p1-'A'] > 0)
                  cx = '.';
              else
                  cx = ' ';
          }
0          else
              cx = ' ';
          *px++ = cx;
      }
5      *px++ = '\n';
      *px = '\0';
  }

```

...putline

stars

**Table 1 (cont')**

```
/*
 * strip path or prefix from pn, return len: pr_align()
 */
static
5 stripname(pn)
   char *pn; /* file name (may be path) */
   {
   register char *px, *py;
0   py = 0;
   for (px = pn; *px; px++)
       if (*px == '/')
           py = px + 1;
5   if (py)
       (void) strcpy(pn, py);
   return(strlen(pn));
   }
0
```

**stripname**

**Table 1 (cont')**

```

/*
 * cleanup() -- cleanup any tmp file
 * getseq() -- read in seq, set dna, len, maxlen
 * g_malloc() -- calloc() with error checkin
5  * readjumps() -- get the good jumps, from tmp file if necessary
 * writejumps() -- write a filled array of jumps to a tmp file: nw()
 */
#include "nw.h"
#include <sys/file.h>
10 char *jname = "/tmp/homgXXXXXX"; /* tmp file for jumps */
FILE *fj;

int cleanup(); /* cleanup tmp file */
15 long lseek();

/*
 * remove any tmp file if we blow
 */
20 cleanup(i)
    int i;
    {
        if (fj)
            (void) unlink(jname);
25         exit(i);
    }

/*
 * read, return ptr to seq, set dna, len, maxlen
 * skip lines starting with ';', '<', or '>'
 * seq in upper or lower case
 */
30 char *
getseq(file, len)
35 char *file; /* file name */
    int *len; /* seq len */
    {
        char line[1024], *pseq;
        register char *px, *py;
        int natgc, tlen;
        FILE *fp;

        if ((fp = fopen(file, "r")) == 0) {
45             fprintf(stderr, "%s: can't read %s\n", prog, file);
             exit(1);
        }
        tlen = natgc = 0;
        while (fgets(line, 1024, fp)) {
50             if (*line == ';' || *line == '<' || *line == '>')
                 continue;
             for (px = line; *px != '\n'; px++)
                 if (isupper(*px) || islower(*px))
                     tlen++;
        }
55         if ((pseq = malloc((unsigned)(tlen+6))) == 0) {
             fprintf(stderr, "%s: malloc() failed to get %d bytes for %s\n", prog, tlen+6, file);
             exit(1);
        }
        pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0';
60

```

**cleanup****getseq**

**Table 1 (cont')**

```

py = pseq + 4;
*len = tlen;
rewind(fp);

while (fgets(line, 1024, fp)) {
    if (*line == ';' || *line == '<' || *line == '>')
        continue;
    for (px = line; *px != '\n'; px++) {
        if (isupper(*px))
            *py++ = *px;
        else if (islower(*px))
            *py++ = toupper(*px);
        if (index("ATGCU", *(py-1)))
            natgc++;
    }
    *py++ = '\0';
    *py = '\0';
    (void) fclose(fp);
    dna = natgc > (tlen/3);
    return(pseq+4);
}

char *
g_alloc(msg, nx, sz)
char *msg; /* program, calling routine */
int nx, sz; /* number and size of elements */
{
    char *px, *alloc();

    if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
        if (*msg) {
            fprintf(stderr, "%s: g_alloc() failed %s (n=%d, sz=%d)\n", prog, msg, nx, sz);
            exit(1);
        }
    }
    return(px);
}

/*
 * get final jmps from dx[] or tmp file, set pp[], reset dmax: main()
 */
readjmps()
{
    int fd = -1;
    int siz, i0, i1;
    register i, j, xx;

    if (fj) {
        (void) fclose(fj);
        if ((fd = open(jname, O_RDONLY, 0)) < 0) {
            fprintf(stderr, "%s: can't open() %s\n", prog, jname);
            cleanup(1);
        }
    }
    for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; ; i++) {
        while (1) {
            for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--);

```

...getseq

g\_alloc

readjmps

**Table 1 (cont')****...readjumps**

```

    if (j < 0 && dx[dmax].offset && fj) {
        (void) lseek(fd, dx[dmax].offset, 0);
        (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
        (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
        dx[dmax].ijmp = MAXJMP-1;
    }
    else
        break;
}
if (i >= JMPS) {
    fprintf(stderr, "%s: too many gaps in alignment\n", prog);
    cleanup(1);
}
if (j >= 0) {
    siz = dx[dmax].jp.n[j];
    xx = dx[dmax].jp.x[j];
    dmax += siz;
    if (siz < 0) { /* gap in second seq */
        pp[1].n[i1] = -siz;
        xx += siz;
        /* id = xx - yy + len1 - 1
        */
        pp[1].x[i1] = xx - dmax + len1 - 1;
        gapy++;
        ngapy -= siz;
        /* ignore MAXGAP when doing endgaps */
        siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
        i1++;
    }
    else if (siz > 0) { /* gap in first seq */
        pp[0].n[i0] = siz;
        pp[0].x[i0] = xx;
        gapx++;
        ngapx += siz;
        /* ignore MAXGAP when doing endgaps */
        siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
        i0++;
    }
}
else
    break;
}

/* reverse the order of jumps
*/
for (j = 0, i0--, j < i0; j++, i0--) {
    i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;
    i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
}
for (j = 0, i1--, j < i1; j++, i1--) {
    i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
    i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
}
if (fd >= 0)
    (void) close(fd);
if (fj) {
    (void) unlink(jname);
    fj = 0;
    offset = 0;
}
}
}

```

**Table 1 (cont')**

```
/*
 * write a filled jmp struct offset of the prev one (if any): nw()
 */
writejumps(ix)
{
    int ix;
    char *mktemp();

    if (!fj) {
        if (mktemp(jname) < 0) {
            fprintf(stderr, "%s: can't mktemp() %s\n", prog, jname);
            cleanup(1);
        }
        if ((fj = fopen(jname, "w")) == 0) {
            fprintf(stderr, "%s: can't write %s\n", prog, jname);
            exit(1);
        }
    }
    (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
    (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
}
```

**writejumps**

TABLE 2

PRO	XXXXXXXXXXXXXXXXXX	(Length = 15 amino acids)
Comparison Protein	XXXXXXXXXXXXXXX	(Length = 12 amino acids)

% amino acid sequence identity = (the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) = 5 divided by 15 = 33.3%

[0196]

TABLE 3

PRO	XXXXXXXXXX	(Length = 10 amino acids)
Comparison Protein	XXXXXXXXXXXXZZZ	(Length = 15 amino acids)

% amino acid sequence identity = (the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) = 5 divided by 10 = 50%

[0197]

TABLE 4

PRO-DNA	NNNNNNNNNNNN	(Length = 14 nucleotides)
Comparison DNA	NNNNNNLLLLLLLL	(Length = 16 nucleotides)

% nucleic acid sequence identity = (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) = 6 divided by 14 = 42.9%

[0198]

TABLE 5

PRO-DNA	NNNNNNNNNNNN	(Length = 12 nucleotides)
Comparison DNA	NNNNLLLVV	(Length = 9 nucleotides)

% nucleic acid sequence identity = (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) = 4 divided by 12 = 33.3%

[0199] II. Compositions and Methods of the Invention

[0200] A. Full-Length PRO Polypeptides

[0201] The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO polypeptides. In particular, cDNAs encoding various PRO polypeptides have been identified and isolated, as disclosed in further detail in the Examples below. It is noted that proteins produced in separate expression rounds may be given different PRO numbers but the UNQ number is unique for any given DNA and the encoded protein, and will not be changed. However, for sake of simplicity, in the present specification the protein encoded by the full length native nucleic acid molecules disclosed herein as well as all further native homologues and variants included in the foregoing definition of PRO, will be referred to as "PRO/number", regardless of their origin or mode of preparation.

[0202] As disclosed in the Examples below, various cDNA clones have been deposited with the ATCC. The actual nucleotide sequences of those clones can readily be deter-

mined by the skilled artisan by sequencing of the deposited clone using routine methods in the art. The predicted amino acid sequence can be determined from the nucleotide sequence using routine skill. For the PRO polypeptides and encoding nucleic acids described herein, Applicants have identified what is believed to be the reading frame best identifiable with the sequence information available at the time.

[0203] B. PRO Polypeptide Variants

[0204] In addition to the full-length native sequence PRO polypeptides described herein, it is contemplated that PRO variants can be prepared. PRO variants can be prepared by introducing appropriate nucleotide changes into the PRO DNA, and/or by synthesis of the desired PRO polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the PRO, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

[0205] Variations in the native full-length sequence PRO or in various domains of the PRO described herein, can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Pat. No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the PRO that results in a change in the amino acid sequence of the PRO as compared with the native sequence PRO. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the PRO. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the PRO with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

[0206] PRO polypeptide fragments are provided herein. Such fragments may be truncated at the N-terminus or C-terminus, or may lack internal residues, for example, when compared with a full length native protein. Certain fragments lack amino acid residues that are not essential for a desired biological activity of the PRO polypeptide.

[0207] PRO fragments may be prepared by any of a number of conventional techniques. Desired peptide fragments may be chemically synthesized. An alternative approach involves generating PRO fragments by enzymatic digestion, e.g., by treating the protein with an enzyme known to cleave proteins at sites defined by particular amino acid residues, or by digesting the DNA with suitable restriction enzymes and isolating the desired fragment. Yet another suitable technique involves isolating and amplifying a DNA fragment encoding a desired polypeptide fragment, by polymerase chain reaction (PCR). Oligonucleotides that define

the desired termini of the DNA fragment are employed at the 5' and 3' primers in the PCR. Preferably, PRO polypeptide fragments share at least one biological and/or immunological activity with the native PRO polypeptide disclosed herein.

[0208] In particular embodiments, conservative substitutions of interest are shown in Table 6 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in Table 6, or as further described below in reference to amino acid classes, are introduced and the products screened.

TABLE 6

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; lys; arg	gln
Asp (D)	glu	glu
Cys (C)	ser	ser
Gln (Q)	asn	asn
Glu (E)	asp	asp
Gly (G)	pro; ala	ala
His (H)	asn; gln; lys; arg	arg
Ile (I)	leu; val; met; ala; phe; norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala; tyr	leu
Pro (P)	ala	ala
Ser (S)	thr	thr
Thr (T)	ser	ser
Trp (W)	tyr; phe	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

[0209] Substantial modifications in function or immunological identity of the PRO polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

[0210] (1) hydrophobic: norleucine, met, ala, val, leu, ile;

[0211] (2) neutral hydrophilic: cys, ser, thr;

[0212] (3) acidic: asp, glu;

[0213] (4) basic: asn, gln, his, lys, arg;

[0214] (5) residues that influence chain orientation: gly, pro; and

[0215] (6) aromatic: trp, tyr, phe.

[0216] Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

[0217] The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter et al., *Nucl. Acids Res.*, 13:4331 (1986); Zoller et al., *Nucl. Acids Res.*, 10:6487 (1987)], cassette mutagenesis [Wells et al., *Gene*, 34:315 (1985)], restriction selection mutagenesis [Wells et al., *Philos. Trans. R. Soc. London SerA*, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the PRO variant DNA.

[0218] Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, *Science*, 244: 1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, *The Proteins*, (W. H. Freeman & Co., N.Y.); Chothia, *J. Mol. Biol.*, 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

[0219] C. Modifications of PRO

[0220] Covalent modifications of PRO are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a PRO polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues of the PRO. Derivatization with bifunctional agents is useful, for instance, for crosslinking PRO to a water-insoluble support matrix or surface for use in the method for purifying anti-PRO antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis-(diazocetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propioimidate.

[0221] Other modifications include deamidation of glutamyl and asparagyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the  $\alpha$ -amino groups of lysine, arginine, and histidine side chains [T. E. Creighton, *Proteins: Structure and Molecular Properties*, W. H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

[0222] Another type of covalent modification of the PRO polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence PRO (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means),



and/or adding one or more glycosylation sites that are not present in the native sequence PRO. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

[0223] Addition of glycosylation sites to the PRO polypeptide may be accomplished by altering the amino acid sequence. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence PRO (for O-linked glycosylation sites). The PRO amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the PRO polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

[0224] Another means of increasing the number of carbohydrate moieties on the PRO polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published Sep. 11, 1987, and in Aplin and Wriston, *CRC Crit. Rev. Biochem.*, pp. 259-306 (1981).

[0225] Removal of carbohydrate moieties present on the PRO polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, et al., *Arch. Biochem. Biophys.*, 259:52 (1987) and by Edge et al., *Anal. Biochem.*, 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., *Meth. Enzymol.*, 138:350 (1987).

[0226] Another type of covalent modification of PRO comprises linking the PRO polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Pat. Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

[0227] The PRO of the present invention may also be modified in a way to form a chimeric molecule comprising PRO fused to another, heterologous polypeptide or amino acid sequence.

[0228] In one embodiment, such a chimeric molecule comprises a fusion of the PRO with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl-terminus of the PRO. The presence of such epitope-tagged forms of the PRO can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the PRO to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., *Mol. Cell. Biol.*, 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., *Molecular and Cellular Biology*, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., *Protein Engineering*, 3(6):547-

553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., *BioTechnology*, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., *Science*, 255:192-194 (1992)]; an  $\alpha$ -tubulin epitope peptide [Skinner et al., *J. Biol. Chem.*, 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., *Proc. Natl. Acad. Sci. USA*, 87:6393-6397 (1990)].

[0229] In an alternative embodiment, the chimeric molecule may comprise a fusion of the PRO with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of a PRO polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions see also U.S. Pat. No. 5,428,130 issued Jun. 27, 1995.

[0230] D. Preparation of PRO

[0231] The description below relates primarily to production of PRO by culturing cells transformed or transfected with a vector containing PRO nucleic acid. It is, of course, contemplated that alternative methods, which are well known in the art, may be employed to prepare PRO. For instance, the PRO sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., *Solid-Phase Peptide Synthesis*, W. H. Freeman Co., San Francisco, Calif. (1969); Merrifield, *J. Am. Chem. Soc.*, 85:2149-2154 (1963)]. In vitro protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, Calif.) using manufacturer's instructions. Various portions of the PRO may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length PRO.

[0232] 1. Isolation of DNA Encoding PRO

[0233] DNA encoding PRO may be obtained from a cDNA library prepared from tissue believed to possess the PRO mRNA and to express it at a detectable level. Accordingly, human PRO DNA can be conveniently obtained from a cDNA library prepared from human tissue, such as described in the Examples. The PRO-encoding gene may also be obtained from a genomic library or by known synthetic procedures (e.g., automated nucleic acid synthesis).

[0234] Libraries can be screened with probes (such as antibodies to the PRO or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual* (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding PRO is to use PCR methodology [Sambrook et al., supra; Dieffenbach et al., *PCR Primer: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1995)].

[0235] The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like  $^{32}\text{P}$ -labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., supra.

[0236] Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined using methods known in the art and as described herein.

[0237] Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook et al., supra, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

#### [0238] 2. Selection and Transformation of Host Cells

[0239] Host cells are transfected or transformed with expression or cloning vectors described herein for PRO production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in *Mammalian Cell Biotechnology: a Practical Approach*, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., supra.

[0240] Methods of eukaryotic cell transfection and prokaryotic cell transformation are known to the ordinarily skilled artisan, for example,  $\text{CaCl}_2$ ,  $\text{CaPO}_4$ , liposome-mediated and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., supra, or electroporation is generally used for prokaryotes. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., *Gene*, 23:315 (1983) and WO 89/05859 published Jun. 29, 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, *Virology*, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transfections have been described in U.S. Pat. No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., *J. Bact.*, 130:946 (1977) and Hsiao et al., *Proc. Natl. Acad. Sci. (USA)*, 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene,

polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown et al., *Methods in Enzymology*, 185:527-537 (1990) and Mansour et al., *Nature*, 336:348-352 (1988).

[0241] Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published Apr. 12, 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype tonA; *E. coli* W3110 strain 9E4, which has the complete genotype tonA ptr3; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype tonA ptr3 phoA E15 (argF-lac)169 degP ompT kan<sup>r</sup>; *E. coli* W3110 strain 37D6, which has the complete genotype tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7 ilvG kan<sup>r</sup>; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant degP deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Pat. No. 4,946,783 issued Aug. 7, 1990. Alternatively, in vitro methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

[0242] In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for PRO-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, *Nature*, 290: 140 [1981]; EP 139,383 published May 2, 1985); Kluyveromyces hosts (U.S. Pat. No. 4,943,529; Fleer et al., *Bio/Technology*, 9:968-975 (1991)) such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt et al., *J. Bacteriol.*, 154(2):737-742 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickeramii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilum* (ATCC 36,906; Van den Berg et al., *Bio/Technology*, 8:135 (1990)), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna et al., *J. Basic Microbiol.*, 28:265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case et al., *Proc. Natl. Acad. Sci. USA*, 76:5259-5263 [1979]); Schwanniomycetes such as *Schwanniomycetes occidentalis* (EP 394,538 published Oct. 31, 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published Jan. 10, 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance et al., *Biochem. Biophys. Res. Commun.*, 112:284-289

[1983]; Tilburn et al., *Gene*, 26:205-221 [1983]; Yelton et al., *Proc. Natl. Acad. Sci. USA*, 81: 1470-1474 [1984] and *A. niger* (Kelly and Hynes, *EMBO J.*, 4:475-479 [1985]). Methylophilic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, *The Biochemistry of Methylophilic Yeasts*, 269 (1982).

[0243] Suitable host cells for the expression of glycosylated PRO are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful mammalian host cell lines include Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., *J. Gen. Virol.*, 36:59 (1977)); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.*, 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

### [0244] 3. Selection and Use of a Replicable Vector

[0245] The nucleic acid (e.g., cDNA or genomic DNA) encoding PRO may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

[0246] The PRO may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the PRO-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces*  $\alpha$ -factor leaders, the latter described in U.S. Pat. No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published Apr. 4, 1990), or the signal described in WO 90/13646 published Nov. 15, 1990. In mammalian cell expression, mammalian signal sequences

may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

[0247] Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the  $2\mu$  plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

[0248] Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

[0249] An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the PRO-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb et al., *Nature*, 282:39 (1979); Kingsman et al., *Gene*, 7:141 (1979); Tschemper et al., *Gene*, 10:157 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, *Genetics*, 85:12 (1977)].

[0250] Expression and cloning vectors usually contain a promoter operably linked to the PRO-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the  $\beta$ -lactamase and lactose promoter systems [Chang et al., *Nature*, 275:615 (1978); Goeddel et al., *Nature*, 281:544 (1979)], alkaline phosphatase, a tryptophan (*trp*) promoter system [Goeddel, *Nucleic Acids Res.*, 8:4057 (1980); EP 36,776], and hybrid promoters such as the *tac* promoter [deBoer et al., *Proc. Natl. Acad. Sci. USA*, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding PRO.

[0251] Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., *J. Biol. Chem.*, 255:2073 (1980)] or other glycolytic enzymes [Hess et al., *J. Adv. Enzyme Reg.*, 7:149 (1968); Holland, *Biochemistry*, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

[0252] Other yeast promoters, which are inducible promoters having the additional advantage of transcription

controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

**[0253]** PRO transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published Jul. 5, 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

**[0254]** Transcription of a DNA encoding the PRO by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin,  $\alpha$ -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the PRO coding sequence, but is preferably located at a site 5' from the promoter.

**[0255]** Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding PRO.

**[0256]** Still other methods, vectors, and host cells suitable for adaptation to the synthesis of PRO in recombinant vertebrate cell culture are described in Gething et al., *Nature*, 293:620-625 (1981); Mantei et al., *Nature*, 281:40-46 (1979); EP 117,060; and EP 117,058.

#### **[0257]** 4. Detecting Gene Amplification/Expression

**[0258]** Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, *Proc. Natl. Acad. Sci. USA*, 77:5201-5205 (1980)], dot blotting (DNA analysis), or in situ hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

**[0259]** Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence PRO polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to PRO DNA and encoding a specific antibody epitope.

#### **[0260]** 5. Purification of Polypeptide

**[0261]** Forms of PRO may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of PRO can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

**[0262]** It may be desired to purify PRO from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the PRO. Various methods of protein purification may be employed and such methods are known in the art and described for example in Deutscher, *Methods in Enzymology*, 182 (1990); Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular PRO produced.

#### **[0263]** E. Uses for PRO

**[0264]** Nucleotide sequences (or their complement) encoding PRO have various applications in the art of molecular biology, including uses as hybridization probes, in chromosome and gene mapping and in the generation of anti-sense RNA and DNA. PRO nucleic acid will also be useful for the preparation of PRO polypeptides by the recombinant techniques described herein.

**[0265]** The full-length native sequence PRO gene, or portions thereof, may be used as hybridization probes for a cDNA library to isolate the full-length PRO cDNA or to isolate still other cDNAs (for instance, those encoding naturally-occurring variants of PRO or PRO from other species) which have a desired sequence identity to the native PRO sequence disclosed herein. Optionally, the length of the probes will be about 20 to about 50 bases. The hybridization probes may be derived from at least partially novel regions of the full length native nucleotide sequence wherein those regions may be determined without undue experimentation or from genomic sequences including promoters, enhancer elements and introns of native sequence PRO. By way of example, a screening method will comprise isolating the coding region of the PRO gene using the known DNA sequence to synthesize a selected probe of about 40 bases.

Hybridization probes may be labeled by a variety of labels, including radionucleotides such as  $^{32}\text{P}$  or  $^{35}\text{S}$ , or enzymatic labels such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems. Labeled probes having a sequence complementary to that of the PRO gene of the present invention can be used to screen libraries of human cDNA, genomic DNA or mRNA to determine which members of such libraries the probe hybridizes to. Hybridization techniques are described in further detail in the Examples below.

[0266] Any EST sequences disclosed in the present application may similarly be employed as probes, using the methods disclosed herein.

[0267] Other useful fragments of the PRO nucleic acids include antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence (either RNA or DNA) capable of binding to target PRO mRNA (sense) or PRO DNA (antisense) sequences. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment of the coding region of PRO DNA. Such a fragment generally comprises at least about 14 nucleotides, preferably from about 14 to 30 nucleotides. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence encoding a given protein is described in, for example, Stein and Cohen (*Cancer Res.* 48:2659, 1988) and van der Krol et al. (*BioTechniques* 6:958, 1988).

[0268] Binding of antisense or sense oligonucleotides to target nucleic acid sequences results in the formation of duplexes that block transcription or translation of the target sequence by one of several means, including enhanced degradation of the duplexes, premature termination of transcription or translation, or by other means. The antisense oligonucleotides thus may be used to block expression of PRO proteins. Antisense or sense oligonucleotides further comprise oligonucleotides having modified sugar-phosphodiester backbones (or other sugar linkages, such as those described in WO 91/06629) and wherein such sugar linkages are resistant to endogenous nucleases. Such oligonucleotides with resistant sugar linkages are stable in vivo (i.e., capable of resisting enzymatic degradation) but retain sequence specificity to be able to bind to target nucleotide sequences.

[0269] Other examples of sense or antisense oligonucleotides include those oligonucleotides which are covalently linked to organic moieties, such as those described in WO 90/10048, and other moieties that increases affinity of the oligonucleotide for a target nucleic acid sequence, such as poly-(L-lysine). Further still, intercalating agents, such as ellipticine, and alkylating agents or metal complexes may be attached to sense or antisense oligonucleotides to modify binding specificities of the antisense or sense oligonucleotide for the target nucleotide sequence.

[0270] Antisense or sense oligonucleotides may be introduced into a cell containing the target nucleic acid sequence by any gene transfer method, including, for example,  $\text{CaPO}_4$ -mediated DNA transfection, electroporation, or by using gene transfer vectors such as Epstein-Barr virus. In a preferred procedure, an antisense or sense oligonucleotide is inserted into a suitable retroviral vector. A cell containing the target nucleic acid sequence is contacted with the recombinant retroviral vector, either in vivo or ex vivo. Suitable retroviral vectors include, but are not limited to, those derived from the murine retrovirus M-MuLV, N2 (a retro-

virus derived from M-MuLV), or the double copy vectors designated DCT5A, DCT5B and DCT5C (see WO 90/13641).

[0271] Sense or antisense oligonucleotides also may be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell.

[0272] Alternatively, a sense or an antisense oligonucleotide may be introduced into a cell containing the target nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. The sense or antisense oligonucleotide-lipid complex is preferably dissociated within the cell by an endogenous lipase.

[0273] Antisense or sense RNA or DNA molecules are generally at least about 5 bases in length, about 10 bases in length, about 15 bases in length, about 20 bases in length, about 25 bases in length, about 30 bases in length, about 35 bases in length, about 40 bases in length, about 45 bases in length, about 50 bases in length, about 55 bases in length, about 60 bases in length, about 65 bases in length, about 70 bases in length, about 75 bases in length, about 80 bases in length, about 85 bases in length, about 90 bases in length, about 95 bases in length, about 100 bases in length, or more.

[0274] The probes may also be employed in PCR techniques to generate a pool of sequences for identification of closely related PRO coding sequences.

[0275] Nucleotide sequences encoding a PRO can also be used to construct hybridization probes for mapping the gene which encodes that PRO and for the genetic analysis of individuals with genetic disorders. The nucleotide sequences provided herein may be mapped to a chromosome and specific regions of a chromosome using known techniques, such as in situ hybridization, linkage analysis against known chromosomal markers, and hybridization screening with libraries.

[0276] When the coding sequences for PRO encode a protein which binds to another protein (example, where the PRO is a receptor), the PRO can be used in assays to identify the other proteins or molecules involved in the binding interaction. By such methods, inhibitors of the receptor/ligand binding interaction can be identified. Proteins involved in such binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction. Also, the receptor PRO can be used to isolate correlative ligand(s). Screening assays can be designed to find lead compounds that mimic the biological activity of a native PRO or a receptor for PRO. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates. Small molecules contemplated include synthetic organic or inorganic compounds. The assays can be performed in a variety of formats, including protein-protein

binding assays, biochemical screening assays, immunoassays and cell based assays, which are well characterized in the art.

[0277] Nucleic acids which encode PRO or its modified forms can also be used to generate either transgenic animals or “knock out” animals which, in turn, are useful in the development and screening of therapeutically useful reagents. A transgenic animal (e.g., a mouse or rat) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an embryonic stage. A transgene is a DNA which is integrated into the genome of a cell from which a transgenic animal develops. In one embodiment, cDNA encoding PRO can be used to clone genomic DNA encoding PRO in accordance with established techniques and the genomic sequences used to generate transgenic animals that contain cells which express DNA encoding PRO. Methods for generating transgenic animals, particularly animals such as mice or rats, have become conventional in the art and are described, for example, in U.S. Pat. Nos. 4,736,866 and 4,870,009. Typically, particular cells would be targeted for PRO transgene incorporation with tissue-specific enhancers. Transgenic animals that include a copy of a transgene encoding PRO introduced into the germ line of the animal at an embryonic stage can be used to examine the effect of increased expression of DNA encoding PRO. Such animals can be used as tester animals for reagents thought to confer protection from, for example, pathological conditions associated with its overexpression. In accordance with this facet of the invention, an animal is treated with the reagent and a reduced incidence of the pathological condition, compared to untreated animals bearing the transgene, would indicate a potential therapeutic intervention for the pathological condition.

[0278] Alternatively, non-human homologues of PRO can be used to construct a PRO “knock out” animal which has a defective or altered gene encoding PRO as a result of homologous recombination between the endogenous gene encoding PRO and altered genomic DNA encoding PRO introduced into an embryonic stem cell of the animal. For example, cDNA encoding PRO can be used to clone genomic DNA encoding PRO in accordance with established techniques. A portion of the genomic DNA encoding PRO can be deleted or replaced with another gene, such as a gene encoding a selectable marker which can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector [see e.g., Thomas and Capecchi, *Cell*, 51:503 (1987) for a description of homologous recombination vectors]. The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected [see e.g., Li et al., *Cell*, 69:915 (1992)]. The selected cells are then injected into a blastocyst of an animal (e.g., a mouse or rat) to form aggregation chimeras [see e.g., Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113-152]. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a “knock out” animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knock-

out animals can be characterized for instance, for their ability to defend against certain pathological conditions and for their development of pathological conditions due to absence of the PRO polypeptide.

[0279] Nucleic acid encoding the PRO polypeptides may also be used in gene therapy. In gene therapy applications, genes are introduced into cells in order to achieve in vivo synthesis of a therapeutically effective genetic product, for example for replacement of a defective gene. “Gene therapy” includes both conventional gene therapy where a lasting effect is achieved by a single treatment, and the administration of gene therapeutic agents, which involves the one time or repeated administration of a therapeutically effective DNA or mRNA. Antisense RNAs and DNAs can be used as therapeutic agents for blocking the expression of certain genes in vivo. It has already been shown that short antisense oligonucleotides can be imported into cells where they act as inhibitors, despite their low intracellular concentrations caused by their restricted uptake by the cell membrane. (Zamecnik et al., *Proc. Natl. Acad. Sci. USA* 83:4143-4146 [1986]). The oligonucleotides can be modified to enhance their uptake, e.g. by substituting their negatively charged phosphodiester groups by uncharged groups.

[0280] There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells in vitro, or in vivo in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells in vitro include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. The currently preferred in vivo gene transfer techniques include transfection with viral (typically retroviral) vectors and viral coat protein-liposome mediated transfection (Dzau et al., *Trends in Biotechnology* 11, 205-210 [1993]). In some situations it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins which bind to a cell surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, e.g. capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu et al., *J. Biol. Chem.* 262, 4429-4432 (1987); and Wagner et al., *Proc. Natl. Acad. Sci. USA* 87, 3410-3414 (1990). For review of gene marking and gene therapy protocols see Anderson et al., *Science* 256, 808-813 (1992).

[0281] The PRO polypeptides described herein may also be employed as molecular weight markers for protein electrophoresis purposes and the isolated nucleic acid sequences may be used for recombinantly expressing those markers.

[0282] The nucleic acid molecules encoding the PRO polypeptides or fragments thereof described herein are useful for chromosome identification. In this regard, there exists an ongoing need to identify new chromosome markers, since relatively few chromosome marking reagents, based upon actual sequence data are presently available. Each PRO nucleic acid molecule of the present invention can be used as a chromosome marker.

[0283] The PRO polypeptides and nucleic acid molecules of the present invention may also be used diagnostically for tissue typing, wherein the PRO polypeptides of the present invention may be differentially expressed in one tissue as compared to another, preferably in a diseased tissue as compared to a normal tissue of the same tissue type. PRO nucleic acid molecules will find use for generating probes for PCR, Northern analysis, Southern analysis and Western analysis.

[0284] The PRO polypeptides described herein may also be employed as therapeutic agents. The PRO polypeptides of the present invention can be formulated according to known methods to prepare pharmaceutically useful compositions, whereby the PRO product hereof is combined in admixture with a pharmaceutically acceptable carrier vehicle. Therapeutic formulations are prepared for storage by mixing the active ingredient having the desired degree of purity with optional physiologically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate and other organic acids; antioxidants including ascorbic acid; Low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone, amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, PLURONICS™ or PEG.

[0285] The formulations to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution.

[0286] Therapeutic compositions herein generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[0287] The route of administration is in accord with known methods, e.g. injection or infusion by intravenous, intraperitoneal, intracerebral, intramuscular, intraocular, intraarterial or intralesional routes, topical administration, or by sustained release systems.

[0288] Dosages and desired drug concentrations of pharmaceutical compositions of the present invention may vary depending on the particular use envisioned. The determination of the appropriate dosage or route of administration is well within the skill of an ordinary physician. Animal experiments provide reliable guidance for the determination of effective doses for human therapy. Interspecies scaling of effective doses can be performed following the principles laid down by Mordenti, J. and Chappell, W. "The use of interspecies scaling in toxicokinetics" In *Toxicokinetics and New Drug Development*, Yacobi et al., Eds., Pergamon Press, New York 1989, pp. 42-96.

[0289] When in vivo administration of a PRO polypeptide or agonist or antagonist thereof is employed, normal dosage

amounts may vary from about 10 ng/kg to up to 100 mg/kg of mammal body weight or more per day, preferably about 1 µg/kg/day to 10 mg/kg/day, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see, for example, U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. It is anticipated that different formulations will be effective for different treatment compounds and different disorders, that administration targeting one organ or tissue, for example, may necessitate delivery in a manner different from that to another organ or tissue.

[0290] Where sustained-release administration of a PRO polypeptide is desired in a formulation with release characteristics suitable for the treatment of any disease or disorder requiring administration of the PRO polypeptide, microencapsulation of the PRO polypeptide is contemplated. Microencapsulation of recombinant proteins for sustained release has been successfully performed with human growth hormone (rhGH), interferon-(rhIFN-), interleukin-2, and MN rgp120. Johnson et al., *Nat. Med.*, 2:795-799 (1996); Yasuda, *Biomed. Ther.*, 27:1221-1223 (1993); Hora et al., *Bio/Technology*, 8:755-758 (1990); Cleland, "Design and Production of Single Immunization Vaccines Using Poly-lactide Polyglycolide Microsphere Systems," in *Vaccine Design: The Subunit and Adjuvant Approach*, Powell and Newman, eds, (Plenum Press: New York, 1995), pp. 439-462; WO 97/03692, WO 96/40072, WO 96/07399; and U.S. Pat. No. 5,654,010.

[0291] The sustained-release formulations of these proteins were developed using poly-lactic-coglycolic acid (PLGA) polymer due to its biocompatibility and wide range of biodegradable properties. The degradation products of PLGA, lactic and glycolic acids, can be cleared quickly within the human body. Moreover, the degradability of this polymer can be adjusted from months to years depending on its molecular weight and composition. Lewis, "Controlled release of bioactive agents from lactide/glycolide polymer," in: M. Chasin and R. Langer (Eds.), *Biodegradable Polymers as Drug Delivery Systems* (Marcel Dekker: New York, 1990), pp. 1-41.

[0292] This invention encompasses methods of screening compounds to identify those that mimic the PRO polypeptide (agonists) or prevent the effect of the PRO polypeptide (antagonists). Screening assays for antagonist drug candidates are designed to identify compounds that bind or complex with the PRO polypeptides encoded by the genes identified herein, or otherwise interfere with the interaction of the encoded polypeptides with other cellular proteins. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates.

[0293] The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays, and cell-based assays, which are well characterized in the art.

[0294] All assays for antagonists are common in that they call for contacting the drug candidate with a PRO polypeptide encoded by a nucleic acid identified herein under conditions and for a time sufficient to allow these two components to interact.

[0295] In binding assays, the interaction is binding and the complex formed can be isolated or detected in the reaction

mixture. In a particular embodiment, the PRO polypeptide encoded by the gene identified herein or the drug candidate is immobilized on a solid phase, e.g., on a microtiter plate, by covalent or non-covalent attachments. Non-covalent attachment generally is accomplished by coating the solid surface with a solution of the PRO polypeptide and drying. Alternatively, an immobilized antibody, e.g., a monoclonal antibody, specific for the PRO polypeptide to be immobilized can be used to anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which may be labeled by a detectable label, to the immobilized component, e.g., the coated surface containing the anchored component. When the reaction is complete, the non-reacted components are removed, e.g., by washing, and complexes anchored on the solid surface are detected. When the originally non-immobilized component carries a detectable label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally non-immobilized component does not carry a label, complexing can be detected, for example, by using a labeled antibody specifically binding the immobilized complex.

**[0296]** If the candidate compound interacts with but does not bind to a particular PRO polypeptide encoded by a gene identified herein, its interaction with that polypeptide can be assayed by methods well known for detecting protein-protein interactions. Such assays include traditional approaches, such as, e.g., cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields and co-workers (Fields and Song, *Nature (London)*, 340:245-246 (1989); Chien et al., *Proc. Natl. Acad. Sci. USA*, 88:9578-9582 (1991)) as disclosed by Chevray and Nathans, *Proc. Natl. Acad. Sci. USA*, 89: 5789-5793 (1991). Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, the other one functioning as the transcription-activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-lacZ reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for  $\beta$ -galactosidase. A complete kit (MATCHMAKER™) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

**[0297]** Compounds that interfere with the interaction of a gene encoding a PRO polypeptide identified herein and other intra- or extracellular components can be tested as follows: usually a reaction mixture is prepared containing the product of the gene and the intra- or extracellular component under conditions and for a time allowing for the interaction and binding of the two products. To test the ability of a candidate compound to inhibit binding, the

reaction is run in the absence and in the presence of the test compound. In addition, a placebo may be added to a third reaction mixture, to serve as positive control. The binding (complex formation) between the test compound and the intra- or extracellular component present in the mixture is monitored as described hereinabove. The formation of a complex in the control reaction(s) but not in the reaction mixture containing the test compound indicates that the test compound interferes with the interaction of the test compound and its reaction partner.

**[0298]** To assay for antagonists, the PRO polypeptide may be added to a cell along with the compound to be screened for a particular activity and the ability of the compound to inhibit the activity of interest in the presence of the PRO polypeptide indicates that the compound is an antagonist to the PRO polypeptide. Alternatively, antagonists may be detected by combining the PRO polypeptide and a potential antagonist with membrane-bound PRO polypeptide receptors or recombinant receptors under appropriate conditions for a competitive inhibition assay. The PRO polypeptide can be labeled, such as by radioactivity, such that the number of PRO polypeptide molecules bound to the receptor can be used to determine the effectiveness of the potential antagonist. The gene encoding the receptor can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting. Coligan et al., *Current Protocols in Immun.*, 1(2): Chapter 5 (1991). Preferably, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the PRO polypeptide and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the PRO polypeptide. Transfected cells that are grown on glass slides are exposed to labeled PRO polypeptide. The PRO polypeptide can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase. Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an interactive sub-pooling and re-screening process, eventually yielding a single clone that encodes the putative receptor.

**[0299]** As an alternative approach for receptor identification, labeled PRO polypeptide can be photoaffinity-linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE and exposed to X-ray film. The labeled complex containing the receptor can be excised, resolved into peptide fragments, and subjected to protein micro-sequencing. The amino acid sequence obtained from micro-sequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the gene encoding the putative receptor.

**[0300]** In another assay for antagonists, mammalian cells or a membrane preparation expressing the receptor would be incubated with labeled PRO polypeptide in the presence of the candidate compound. The ability of the compound to enhance or block this interaction could then be measured.

**[0301]** More specific examples of potential antagonists include an oligonucleotide that binds to the fusions of immunoglobulin with PRO polypeptide, and, in particular, antibodies including, without limitation, poly- and monoclonal antibodies and antibody fragments, single-chain anti-



bodies, anti-idiotypic antibodies, and chimeric or humanized versions of such antibodies or fragments, as well as human antibodies and antibody fragments. Alternatively, a potential antagonist may be a closely related protein, for example, a mutated form of the PRO polypeptide that recognizes the receptor but imparts no effect, thereby competitively inhibiting the action of the PRO polypeptide.

[0302] Another potential PRO polypeptide antagonist is an antisense RNA or DNA construct prepared using antisense technology, where, e.g., an antisense RNA or DNA molecule acts to block directly the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes the mature PRO polypeptides herein, is used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix—see Lee et al., *Nucl. Acids Res.*, 6:3073 (1979); Cooney et al., *Science*, 241: 456 (1988); Dervan et al., *Science*, 251:1360 (1991)), thereby preventing transcription and the production of the PRO polypeptide. The antisense RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of the mRNA molecule into the PRO polypeptide (antisense—Okano, *Neurochem.*, 56:560 (1991); *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression* (CRC Press: Boca Raton, Fla., 1988). The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed in vivo to inhibit production of the PRO polypeptide. When antisense DNA is used, oligodeoxyribonucleotides derived from the translation-initiation site, e.g., between about -10 and +10 positions of the target gene nucleotide sequence, are preferred.

[0303] Potential antagonists include small molecules that bind to the active site, the receptor binding site, or growth factor or other relevant binding site of the PRO polypeptide, thereby blocking the normal biological activity of the PRO polypeptide. Examples of small molecules include, but are not limited to, small peptides or peptide-like molecules, preferably soluble peptides, and synthetic non-peptidyl organic or inorganic compounds.

[0304] Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. Ribozymes act by sequence-specific hybridization to the complementary target RNA, followed by endonucleolytic cleavage. Specific ribozyme cleavage sites within a potential RNA target can be identified by known techniques. For further details see, e.g., Rossi, *Current Biology*, 4:469-471 (1994), and PCT publication No. WO 97/33551 (published Sep. 18, 1997).

[0305] Nucleic acid molecules in triple-helix formation used to inhibit transcription should be single-stranded and composed of deoxynucleotides. The base composition of these oligonucleotides is designed such that it promotes triple-helix formation via Hoogsteen base-pairing rules, which generally require sizeable stretches of purines or pyrimidines on one strand of a duplex. For further details see, e.g., PCT publication No. WO 97/33551, supra.

[0306] These small molecules can be identified by any one or more of the screening assays discussed hereinabove and/or by any other screening techniques well known for those skilled in the art.

[0307] Diagnostic and therapeutic uses of the herein disclosed molecules may also be based upon the positive functional assay hits disclosed and described below.

[0308] F. Anti-PRO Antibodies

[0309] The present invention further provides anti-PRO antibodies. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies.

[0310] 1. Polyclonal Antibodies

[0311] The anti-PRO antibodies may comprise polyclonal antibodies. Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the PRO polypeptide or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

[0312] 2. Monoclonal Antibodies

[0313] The anti-PRO antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, *Nature*, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized in vitro.

[0314] The immunizing agent will typically include the PRO polypeptide or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, *Monoclonal Antibodies: Principles and Practice*, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine

phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

[0315] Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, Calif. and the American Type Culture Collection, Manassas, Va. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur et al., *Monoclonal Antibody Production Techniques and Applications*, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

[0316] The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against PRO. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, *Anal. Biochem.*, 107:220 (1980).

[0317] After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods [Goding, *supra*]. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown in vivo as ascites in a mammal.

[0318] The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

[0319] The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Pat. No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences [U.S. Pat. No. 4,816,567; Morrison et al., *supra*] or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide.

Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

[0320] The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

[0321] In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly, Fab fragments, can be accomplished using routine techniques known in the art.

[0322] 3. Human and Humanized Antibodies

[0323] The anti-PRO antibodies of the invention may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub> or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.*, 2:593-596 (1992)].

[0324] Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones et al., *Nature*, 321:522-525 (1986); Riechman et al., *Nature*, 332:323-327 (1988); Verhoeven et al., *Science*, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567), wherein substantially less than an intact human

variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

[0325] Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985) and Boerner et al., *J. Immunol.*, 147(1):86-95 (1991)]. Similarly, human antibodies can be made by introducing of human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Pat. Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks et al., *Bio/Technology* 10, 779-783 (1992); Lonberg et al., *Nature* 368 856-859 (1994); Morrison, *Nature* 368, 812-13 (1994); Fishwild et al., *Nature Biotechnology* 14, 845-51 (1996); Neuberger, *Nature Biotechnology* 14, 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol.* 13 65-93 (1995).

[0326] The antibodies may also be affinity matured using known selection and/or mutagenesis methods as described above. Preferred affinity matured antibodies have an affinity which is five times, more preferably 10 times, even more preferably 20 or 30 times greater than the starting antibody (generally murine, humanized or human) from which the matured antibody is prepared.

#### [0327] 4. Bispecific Antibodies

[0328] Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the PRO, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

[0329] Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities [Milstein and Cuello, *Nature*, 305:537-539 (1983)]. Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published May 13, 1993, and in Traunecker et al., *EMBO J.*, 10:3655-3659 (1991).

[0330] Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain

constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., *Methods in Enzymology*, 121:210 (1986).

[0331] According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

[0332] Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')<sub>2</sub> bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared can be prepared using chemical linkage. Brennan et al., *Science* 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')<sub>2</sub> fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

[0333] Fab' fragments may be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby et al., *J. Exp. Med.* 175:217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')<sub>2</sub> molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling in vitro to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

[0334] Various technique for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., *J. Immunol.* 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the

hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain ( $V_H$ ) connected to a light-chain variable domain ( $V_L$ ) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the  $V_H$  and  $V_L$  domains of one fragment are forced to pair with the complementary  $V_L$  and  $V_H$  domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber et al., *J. Immunol.* 152:5368 (1994). Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al., *J. Immunol.* 147:60 (1991).

[0335] Exemplary bispecific antibodies may bind to two different epitopes on a given PRO polypeptide herein. Alternatively, an anti-PRO polypeptide arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (Fc $\gamma$ R), such as Fc $\gamma$ RI (CD64), Fc $\gamma$ RII (CD32) and Fc $\gamma$ RIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular PRO polypeptide. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express a particular PRO polypeptide. These antibodies possess a PRO-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the PRO polypeptide and further binds tissue factor (TF).

#### [0336] 5. Heteroconjugate Antibodies

[0337] Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells [U.S. Pat. No. 4,676,980], and for treatment of HIV infection [WO 91/00360; WO 92/200373; EP 03089]. It is contemplated that the antibodies may be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Pat. No. 4,676,980.

#### [0338] 6. Effector Function Engineering

[0339] It may be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) may be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., *J. Exp. Med.*, 176: 1191-1195 (1992) and Shopes, *J. Immunol.*, 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using

heterobifunctional cross-linkers as described in Wolff et al. *Cancer Research*, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson et al., *Anti-Cancer Drug Design*. 3: 219-230 (1989).

#### [0340] 7. Immunoconjugates

[0341] The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

[0342] Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcun, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include  $^{212}\text{Bi}$ ,  $^{131}\text{I}$ ,  $^{131}\text{In}$ ,  $^{90}\text{Y}$ , and  $^{186}\text{Re}$ . Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimide HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., *Science*, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyl-diethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionuclide to the antibody. See WO94/11026.

[0343] In another embodiment, the antibody may be conjugated to a "receptor" (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is conjugated to a cytotoxic agent (e.g., a radionuclide).

#### [0344] 8. Immunoliposomes

[0345] The antibodies disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein et al., *Proc. Natl. Acad. Sci. USA*, 82: 3688 (1985); Hwang et al., *Proc. Natl. Acad. Sci. USA*, 77: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Pat. No. 5,013,556.

[0346] Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid compo-

sition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin et al., *J. Biol. Chem.*, 257: 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See Gabizon et al., *J. National Cancer Inst.*, 81(19): 1484 (1989).

#### [0347] 9. Pharmaceutical Compositions of Antibodies

[0348] Antibodies specifically binding a PRO polypeptide identified herein, as well as other molecules identified by the screening assays disclosed hereinbefore, can be administered for the treatment of various disorders in the form of pharmaceutical compositions.

[0349] If the PRO polypeptide is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, lipofections or liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, e.g., Marasco et al., *Proc. Natl. Acad. Sci. USA*, 90: 7889-7893 (1993). The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition may comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

[0350] The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's *Pharmaceutical Sciences*, supra.

[0351] The formulations to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

[0352] Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and  $\gamma$  ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™

(injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37° C., resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S—S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

#### [0353] G. Uses for Anti-PRO Antibodies

[0354] The anti-PRO antibodies of the invention have various utilities. For example, anti-PRO antibodies may be used in diagnostic assays for PRO, e.g., detecting its expression (and in some cases, differential expression) in specific cells, tissues, or serum. Various diagnostic assay techniques known in the art may be used, such as competitive binding assays, direct or indirect sandwich assays and immunoprecipitation assays conducted in either heterogeneous or homogeneous phases [Zola, *Monoclonal Antibodies: A Manual of Techniques*, CRC Press, Inc. (1987) pp. 147-158]. The antibodies used in the diagnostic assays can be labeled with a detectable moiety. The detectable moiety should be capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ , or  $^{125}\text{I}$ , a fluorescent or chemiluminescence compound, such as fluorescein isothiocyanate, rhodamine, or luciferin, or an enzyme, such as alkaline phosphatase, beta-galactosidase or horseradish peroxidase. Any method known in the art for conjugating the antibody to the detectable moiety may be employed, including those methods described by Hunter et al., *Nature*, 144:945 (1962); David et al., *Biochemistry*, 13:1014 (1974); Pain et al., *J. Immunol. Meth.*, 40:219 (1981); and Nygren, *J. Histochem. and Cytochem.*, 30:407 (1982).

[0355] Anti-PRO antibodies also are useful for the affinity purification of PRO from recombinant cell culture or natural sources. In this process, the antibodies against PRO are immobilized on a suitable support, such as a Sephadex resin or filter paper, using methods well known in the art. The immobilized antibody then is contacted with a sample containing the PRO to be purified, and thereafter the support is washed with a suitable solvent that will remove substantially all the material in the sample except the PRO, which is bound to the immobilized antibody. Finally, the support is washed with another suitable solvent that will release the PRO from the antibody.

[0356] The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

[0357] All patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

## EXAMPLES

[0358] Commercially available reagents referred to in the examples were used according to manufacturer's instructions unless otherwise indicated. The source of those cells identified in the following examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, Va.

## Example 1

[0359] Extracellular Domain Homology Screening to Identify Novel Polypeptides and cDNA Encoding therefor

[0360] The extracellular domain (ECD) sequences (including the secretion signal sequence, if any) from about 950 known secreted proteins from the Swiss-Prot public database were used to search EST databases. The EST databases included public databases (e.g., Dayhoff, GenBank), and proprietary databases (e.g., LIFESEQ™, Incyte Pharmaceuticals, Palo Alto, Calif.). The search was performed using the computer program BLAST or BLAST-2 (Altschul et al., *Methods in Enzymology*, 266:460-480 (1996)) as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequences. Those comparisons with a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Wash.).

[0361] Using this extracellular domain homology screen, consensus DNA sequences were assembled relative to the other identified EST sequences using phrap. In addition, the consensus DNA sequences obtained were often (but not always) extended using repeated cycles of BLAST or BLAST-2 and phrap to extend the consensus sequence as far as possible using the sources of EST sequences discussed above.

[0362] Based upon the consensus sequences obtained as described above, oligonucleotides were then synthesized and used to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for a PRO polypeptide. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than about 1-1.5 kbp. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel et al., *Current Protocols in Molecular Biology*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

[0363] The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, Calif. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as PRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

## Example 2

[0364] Isolation of cDNA Clones by Amylase Screening

[0365] 1. Preparation of Oligo dT Primed cDNA Library

[0366] mRNA was isolated from a human tissue of interest using reagents and protocols from Invitrogen, San Diego, Calif. (Fast Track 2). This RNA was used to generate an oligo dT primed cDNA library in the vector pRK5D using reagents and protocols from Life Technologies, Gaithersburg, Md. (Super Script Plasmid System). In this procedure, the double stranded cDNA was sized to greater than 1000 bp and the SalI/NotI linker cDNA was cloned into XhoI/NotI cleaved vector. pRK5D is a cloning vector that has an sp6 transcription initiation site followed by an SfiI restriction enzyme site preceding the XhoI/NotI cDNA cloning sites.

[0367] 2. Preparation of Random Primed cDNA Library

[0368] A secondary cDNA library was generated in order to preferentially represent the 5' ends of the primary cDNA clones. Sp6 RNA was generated from the primary library (described above), and this RNA was used to generate a random primed cDNA library in the vector pSST-AMY.0 using reagents and protocols from Life Technologies (Super Script Plasmid System, referenced above). In this procedure the double stranded cDNA was sized to 500-1000 bp, linker with blunt to NotI adaptors, cleaved with SfiI, and cloned into SfiI/NotI cleaved vector. pSST-AMY.0 is a cloning vector that has a yeast alcohol dehydrogenase promoter preceding the cDNA cloning sites and the mouse amylase sequence (the mature sequence without the secretion signal) followed by the yeast alcohol dehydrogenase terminator, after the cloning sites. Thus, cDNAs cloned into this vector that are fused in frame with amylase sequence will lead to the secretion of amylase from appropriately transfected yeast colonies.

[0369] 3. Transformation and Detection

[0370] DNA from the library described in paragraph 2 above was chilled on ice to which was added electrocompetent DH10B bacteria (Life Technologies, 20 ml). The bacteria and vector mixture was then electroporated as recommended by the manufacturer. Subsequently, SOC media (Life Technologies, 1 ml) was added and the mixture was incubated at 37° C. for 30 minutes. The transformants were then plated onto 20 standard 150 mm LB plates containing ampicillin and incubated for 16 hours (37° C.). Positive colonies were scraped off the plates and the DNA was isolated from the bacterial pellet using standard protocols, e.g. CsCl-gradient. The purified DNA was then carried on to the yeast protocols below.

[0371] The yeast methods were divided into three categories: (1) Transformation of yeast with the plasmid/cDNA combined vector; (2) Detection and isolation of yeast clones secreting amylase; and (3) PCR amplification of the insert directly from the yeast colony and purification of the DNA for sequencing and further analysis.

[0372] The yeast strain used was HD56-5A (ATCC-90785). This strain has the following genotype: MAT alpha, ura3-52, leu2-3, leu2-112, his3-11, his3-15, MAL<sup>+</sup>, SUC<sup>+</sup>, GAL<sup>+</sup>. Preferably, yeast mutants can be employed that have deficient post-translational pathways. Such mutants may have translocation deficient alleles in sec71, sec72, sec62, with truncated sec71 being most preferred. Alternatively,

antagonists (including antisense nucleotides and/or ligands) which interfere with the normal operation of these genes, other proteins implicated in this post translation pathway (e.g., SEC61p, SEC72p, SEC62p, SEC63p, TDJ1p or SSA1p-4p) or the complex formation of these proteins may also be preferably employed in combination with the amylase-expressing yeast.

[0373] Transformation was performed based on the protocol outlined by Gietz et al., *Nucl. Acid. Res.*, 20:1425 (1992). Transformed cells were then inoculated from agar into YEPD complex media broth (100 ml) and grown overnight at 30° C. The YEPD broth was prepared as described in Kaiser et al., *Methods in Yeast Genetics*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y., p. 207 (1994). The overnight culture was then diluted to about 2×10<sup>7</sup> cells/ml (approx. OD<sub>600</sub>=0.1) into fresh YEPD broth (500 ml) and regrown to 1×10<sup>7</sup> cells/ml (approx. OD<sub>600</sub>=0.4-0.5).

[0374] The cells were then harvested and prepared for transformation by transfer into GS3 rotor bottles in a Sorval GS3 rotor at 5,000 rpm for 5 minutes, the supernatant discarded, and then resuspended into sterile water, and centrifuged again in 50 ml falcon tubes at 3,500 rpm in a Beckman GS-6KR centrifuge. The supernatant was discarded and the cells were subsequently washed with LiAc/TE (10 ml, 10 mM Tris-HCl, 1 mM EDTA pH 7.5, 100 mM Li<sub>2</sub>OOCCH<sub>3</sub>), and resuspended into LiAc/TE (2.5 ml).

[0375] Transformation took place by mixing the prepared cells (100 μl) with freshly denatured single stranded salmon testes DNA (Lofstrand Labs, Gaithersburg, Md.) and transforming DNA (1 μg, vol. <10 μl) in microfuge tubes. The mixture was mixed briefly by vortexing, then 40% PEG/TE (600 μl, 40% polyethylene glycol-4000, 10 mM Tris-HCl, 1 mM EDTA, 100 mM Li<sub>2</sub>OOCCH<sub>3</sub>, pH 7.5) was added. This mixture was gently mixed and incubated at 30° C. while agitating for 30 minutes. The cells were then heat shocked at 42° C. for 15 minutes, and the reaction vessel centrifuged in a microfuge at 12,000 rpm for 5-10 seconds, decanted and resuspended into TE (500 μl, 10 mM Tris-HCl, 1 mM EDTA pH 7.5) followed by recentrifugation. The cells were then diluted into TE (1 ml) and aliquots (200 μl) were spread onto the selective media previously prepared in 150 mm growth plates (VWR).

[0376] Alternatively, instead of multiple small reactions, the transformation was performed using a single, large scale reaction, wherein reagent amounts were scaled up accordingly.

[0377] The selective media used was a synthetic complete dextrose agar lacking uracil (SCD-Ura) prepared as described in Kaiser et al., *Methods in Yeast Genetics*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y., p. 208-210 (1994). Transformants were grown at 30° C. for 2-3 days.

[0378] The detection of colonies secreting amylase was performed by including red starch in the selective growth media. Starch was coupled to the red dye (Reactive Red-120, Sigma) as per the procedure described by Biely et al., *Anal. Biochem.*, 172:176-179 (1988). The coupled starch was incorporated into the SCD-Ura agar plates at a final concentration of 0.15% (w/v), and was buffered with potassium phosphate to a pH of 7.0 (50-100 mM final concentration).

[0379] The positive colonies were picked and streaked across fresh selective media (onto 150 mm plates) in order to obtain well isolated and identifiable single colonies. Well isolated single colonies positive for amylase secretion were detected by direct incorporation of red starch into buffered SCD-Ura agar. Positive colonies were determined by their ability to break down starch resulting in a clear halo around the positive colony visualized directly.

[0380] 4. Isolation of DNA by PCR Amplification

[0381] When a positive colony was isolated, a portion of it was picked by a toothpick and diluted into sterile water (30 μl) in a 96 well plate. At this time, the positive colonies were either frozen and stored for subsequent analysis or immediately amplified. An aliquot of cells (5 μl) was used as a template for the PCR reaction in a 25 μl volume containing: 0.5 μl KlenTaq (Clontech, Palo Alto, Calif.); 4.0 μl 10 mM dNTP's (Perkin Elmer-Cetus); 2.5 μl Kentaq buffer (Clontech); 0.25 μl forward oligo 1; 0.25 μl reverse oligo 2; 12.5 μl distilled water. The sequence of the forward oligonucleotide 1 was:

[0382] 5'-TGTA<sup>u</sup>AAACGACGGCCAGTTAAATA-GACCTGCAATTATTAATCT-3' (SEQ ID NO:115)

[0383] The sequence of reverse oligonucleotide 2 was:

[0384] 5'-CAGGAAACAGCTATGACCACCTGCA-CACCTGCAAATCCATT-3' (SEQ ID NO:116)

[0385] PCR was then performed as follows:

a.	Denature	92° C., 5 minutes
b. 3 cycles of:	Denature	92° C., 30 seconds
	Anneal	59° C., 30 seconds
	Extend	72° C., 60 seconds
c. 3 cycles of:	Denature	92° C., 30 seconds
	Anneal	57° C., 30 seconds
	Extend	72° C., 60 seconds
d. 25 cycles of:	Denature	92° C., 30 seconds
	Anneal	55° C., 30 seconds
	Extend	72° C., 60 seconds
e.	Hold	4° C.

[0386] The underlined regions of the oligonucleotides annealed to the ADH promoter region and the amylase region, respectively, and amplified a 307 bp region from vector pSST-AMY.0 when no insert was present. Typically, the first 18 nucleotides of the 5' end of these oligonucleotides contained annealing sites for the sequencing primers. Thus, the total product of the PCR reaction from an empty vector was 343 bp. However, signal sequence-fused cDNA resulted in considerably longer nucleotide sequences.

[0387] Following the PCR, an aliquot of the reaction (5 μl) was examined by agarose gel electrophoresis in a 1% agarose gel using a Tris-Borate-EDTA (TBE) buffering system as described by Sambrook et al., supra. Clones resulting in a single strong PCR product larger than 400 bp were further analyzed by DNA sequencing after purification with a 96 Qiaquick PCR clean-up column (Qiagen Inc., Chatsworth, Calif.).

## Example 3

**[0388]** Isolation of cDNA Clones Using Signal Algorithm Analysis

**[0389]** Various polypeptide-encoding nucleic acid sequences were identified by applying a proprietary signal sequence finding algorithm developed by Genentech, Inc. (South San Francisco, Calif.) upon ESTs as well as clustered and assembled EST fragments from public (e.g., GenBank) and/or private (LIFESEQ®, Incyte Pharmaceuticals, Inc., Palo Alto, Calif.) databases. The signal sequence algorithm computes a secretion signal score based on the character of the DNA nucleotides surrounding the first and optionally the second methionine codon(s) (ATG) at the 5'-end of the sequence or sequence fragment under consideration. The nucleotides following the first ATG must code for at least 35 unambiguous amino acids without any stop codons. If the first ATG has the required amino acids, the second is not examined. If neither meets the requirement, the candidate sequence is not scored. In order to determine whether the EST sequence contains an authentic signal sequence, the DNA and corresponding amino acid sequences surrounding the ATG codon are scored using a set of seven sensors (evaluation parameters) known to be associated with secretion signals. Use of this algorithm resulted in the identification of numerous polypeptide-encoding nucleic acid sequences.

## Example 4

**[0390]** Isolation of cDNA Clones Encoding Human PRO Polypeptides

**[0391]** Using the techniques described in Examples 1 to 3 above, numerous full-length cDNA clones were identified as encoding PRO polypeptides as disclosed herein. These cDNAs were then deposited under the terms of the Budapest Treaty with the American Type Culture Collection, 10801 University Blvd., Manassas, Va. 20110-2209, USA (ATCC) as shown in Table 7 below.

TABLE 7

Material	ATCC Dep. No.	Deposit Date
DNA16422-1209	209929	Jun. 2, 1998
DNA19902-1669	203454	Nov. 3, 1998
DNA21624-1391	209917	Jun. 2, 1998
DNA34387-1138	209260	Sep. 16, 1997
DNA35880-1160	209379	Oct. 16, 1997
DNA39984-1221	209435	Nov. 7, 1997
DNA44189-1322	209699	Mar. 26, 1998
DNA48303-2829	PTA-1342	Feb. 8, 2000
DNA48320-1433	209904	May 27, 1998
DNA56049-2543	203662	Feb. 9, 1999
DNA57694-1341	203017	Jun. 23, 1998
DNA59208-1373	209881	May 20, 1998
DNA59214-1449	203046	Jul. 1, 1998
DNA59485-1336	203015	Jun. 23, 1998
DNA64966-1575	203575	Jan. 12, 1999
DNA82403-2959	PTA-2317	Aug. 1, 2000
DNA83505-2606	PTA-132	May 25, 1999
DNA84927-2585	203865	Mar. 23, 1999
DNA92264-2616	203969	Apr. 27, 1999
DNA94713-2561	203835	Mar. 9, 1999
DNA96869-2673	PTA-255	Jun. 22, 1999
DNA96881-2699	PTA-553	Aug. 17, 1999
DNA96889-2641	PTA-119	May 25, 1999
DNA96898-2640	PTA-122	May 25, 1999
DNA97003-2649	PTA-43	May 11, 1999

TABLE 7-continued

Material	ATCC Dep. No.	Deposit Date
DNA98565-2701	PTA-481	Aug. 3, 1999
DNA102846-2742	PTA-545	Aug. 17, 1999
DNA102847-2726	PTA-517	Aug. 10, 1999
DNA102880-2689	PTA-383	Jul. 20, 1999
DNA105782-2683	PTA-387	Jul. 20, 1999
DNA108912-2680	PTA-124	May 25, 1999
DNA115253-2757	PTA-612	Aug. 31, 1999
DNA119302-2737	PTA-520	Aug. 10, 1999
DNA119536-2752	PTA-551	Aug. 17, 1999
DNA119542-2754	PTA-619	Aug. 31, 1999
DNA143498-2824	PTA-1263	Feb. 2, 2000
DNA145583-2820	PTA-1179	Jan. 11, 2000
DNA161000-2896	PTA-1731	Apr. 18, 2000
DNA161005-2943	PTA-2243	Jun. 27, 2000
DNA170245-3053	PTA-2952	Jan. 23, 2001
DNA171771-2919	PTA-1902	May 23, 2000
DNA173157-2981	PTA-2388	Aug. 8, 2000
DNA175734-2985	PTA-2455	Sep. 12, 2000
DNA176108-3040	PTA-2824	Dec. 19, 2000
DNA190710-3028	PTA-2822	Dec. 19, 2000
DNA190803-3019	PTA-2785	Dec. 12, 2000
DNA191064-3069	PTA-3016	Feb. 6, 2001
DNA194909-3013	PTA-2779	Dec. 12, 2000
DNA203532-3029	PTA-2823	Dec. 19, 2000
DNA213858-3060	PTA-2958	Jan. 23, 2001
DNA216676-3083	PTA-3157	Mar. 6, 2001
DNA222653-3104	PTA-3330	Apr. 24, 2001
DNA96897-2688	PTA-379	Jul. 20, 1999
DNA142917-3081	PTA-3155	Mar. 6, 2001
DNA142930-2914	PTA-1901	May 23, 2000
DNA147253-2983	PTA-2405	Aug. 22, 2000
DNA149927-2887	PTA-1782	Apr. 25, 2000

**[0392]** These deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposits will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC §122 and the Commissioner's rules pursuant thereto (including 37 CFR §1.14 with particular reference to 886 OG 638).

**[0393]** The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

## Example 5

**[0394]** Use of PRO as a Hybridization Probe

**[0395]** The following method describes use of a nucleotide sequence encoding PRO as a hybridization probe.



[0396] DNA comprising the coding sequence of full-length or mature PRO as disclosed herein is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of PRO) in human tissue cDNA libraries or human tissue genomic libraries.

[0397] Hybridization and washing of filters containing either library DNAs is performed under the following high stringency conditions. Hybridization of radiolabeled PRO-derived probe to the filters is performed in a solution of 50% formamide, 5×SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2×Denhardt's solution, and 10% dextran sulfate at 42° C. for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1×SSC and 0.1% SDS at 42° C.

[0398] DNAs having a desired sequence identity with the DNA encoding full-length native sequence PRO can then be identified using standard techniques known in the art.

#### Example 6

[0399] Expression of PRO in *E. coli*

[0400] This example illustrates preparation of an unglycosylated form of PRO by recombinant expression in *E. coli*.

[0401] The DNA sequence encoding PRO is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector. A variety of expression vectors may be employed. An example of a suitable vector is pBR322 (derived from *E. coli*; see Bolivar et al., *Gene*, 2:95 (1977)) which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme and dephosphorylated. The PCR amplified sequences are then ligated into the vector. The vector will preferably include sequences which encode for an antibiotic resistance gene, a trp promoter, a polyhis leader (including the first six STII codons, polyhis sequence, and enterokinase cleavage site), the PRO coding region, lambda transcriptional terminator, and an argU gene.

[0402] The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook et al., supra. Transformants are identified by their ability to grow on LB plates and antibiotic resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA sequencing.

[0403] Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture may subsequently be used to inoculate a larger scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

[0404] After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the solubilized PRO protein can then be purified using a metal chelating column under conditions that allow tight binding of the protein.

[0405] PRO may be expressed in *E. coli* in a poly-His tagged form, using the following procedure. The DNA encoding PRO is initially amplified using selected PCR primers. The primers will contain restriction enzyme sites which correspond to the restriction enzyme sites on the

selected expression vector, and other useful sequences providing for efficient and reliable translation initiation, rapid purification on a metal chelation column, and proteolytic removal with enterokinase. The PCR-amplified, poly-His tagged sequences are then ligated into an expression vector, which is used to transform an *E. coli* host based on strain 52 (W3110 fuhA(tonA) lon galE rpoHts(htpRts) clpP(lacIq)). Transformants are first grown in LB containing 50 mg/ml carbenicillin at 30° C. with shaking until an O.D.600 of 3-5 is reached. Cultures are then diluted 50-100 fold into CRAP media (prepared by mixing 3.57 g (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.71 g sodium citrate.2H<sub>2</sub>O, 1.07 g KCl, 5.36 g Difco yeast extract, 5.36 g Sheffield hycase SF in 500 mL water, as well as 110 mM MPOS, pH 7.3, 0.55% (w/v) glucose and 7 mM MgSO<sub>4</sub>) and grown for approximately 20-30 hours at 30° C. with shaking. Samples are removed to verify expression by SDS-PAGE analysis, and the bulk culture is centrifuged to pellet the cells. Cell pellets are frozen until purification and refolding.

[0406] *E. coli* paste from 0.5 to 1 L fermentations (6-10 g pellets) is resuspended in 10 volumes (w/v) in 7 M guanidine, 20 mM Tris, pH 8 buffer. Solid sodium sulfite and sodium tetrathionate is added to make final concentrations of 0.1M and 0.02 M, respectively, and the solution is stirred overnight at 4° C. This step results in a denatured protein with all cysteine residues blocked by sulfitolization. The solution is centrifuged at 40,000 rpm in a Beckman Ultracentrifuge for 30 min. The supernatant is diluted with 3-5 volumes of metal chelate column buffer (6 M guanidine, 20 mM Tris, pH 7.4) and filtered through 0.22 micron filters to clarify. The clarified extract is loaded onto a 5 ml Qiagen Ni-NTA metal chelate column equilibrated in the metal chelate column buffer. The column is washed with additional buffer containing 50 mM imidazole (Calbiochem, Utrol grade), pH 7.4. The protein is eluted with buffer containing 250 mM imidazole. Fractions containing the desired protein are pooled and stored at 4° C. Protein concentration is estimated by its absorbance at 280 nm using the calculated extinction coefficient based on its amino acid sequence.

[0407] The proteins are refolded by diluting the sample slowly into freshly prepared refolding buffer consisting of: 20 mM Tris, pH 8.6, 0.3 M NaCl, 2.5 M urea, 5 mM cysteine, 20 mM glycine and 1 mM EDTA. Refolding volumes are chosen so that the final protein concentration is between 50 to 100 micrograms/ml. The refolding solution is stirred gently at 4° C. for 12-36 hours. The refolding reaction is quenched by the addition of TFA to a final concentration of 0.4% (pH of approximately 3). Before further purification of the protein, the solution is filtered through a 0.22 micron filter and acetonitrile is added to 2-10% final concentration. The refolded protein is chromatographed on a Poros R1/H reversed phase column using a mobile buffer of 0.1% TFA with elution with a gradient of acetonitrile from 10 to 80%. Aliquots of fractions with A280 absorbance are analyzed on SDS polyacrylamide gels and fractions containing homogeneous refolded protein are pooled. Generally, the properly refolded species of most proteins are eluted at the lowest concentrations of acetonitrile since those species are the most compact with their hydrophobic interiors shielded from interaction with the reversed phase resin. Aggregated species are usually eluted at higher acetonitrile concentrations. In addition to resolving misfolded forms of proteins from the desired form, the reversed phase step also removes endotoxin from the samples.

[0408] Fractions containing the desired folded PRO polypeptide are pooled and the acetonitrile removed using a gentle stream of nitrogen directed at the solution. Proteins are formulated into 20 mM Hepes, pH 6.8 with 0.14 M sodium chloride and 4% mannitol by dialysis or by gel filtration using G25 Superfine (Pharmacia) resins equilibrated in the formulation buffer and sterile filtered.

[0409] Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

#### Example 7

[0410] Expression of PRO in Mammalian Cells

[0411] This example illustrates preparation of a potentially glycosylated form of PRO by recombinant expression in mammalian cells.

[0412] The vector, pRK5 (see EP 307,247, published Mar. 15, 1989), is employed as the expression vector. Optionally, the PRO DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the PRO DNA using ligation methods such as described in Sambrook et al., supra. The resulting vector is called pRK5-PRO.

[0413] In one embodiment, the selected host cells may be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10  $\mu$ g pRK5-PRO DNA is mixed with about 1  $\mu$ g DNA encoding the VA RNA gene [Thimmappaya et al., *Cell*, 31:543 (1982)] and dissolved in 500  $\mu$ l of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl<sub>2</sub>. To this mixture is added, dropwise, 500  $\mu$ l of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO<sub>4</sub>, and a precipitate is allowed to form for 10 minutes at 25° C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37° C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, fresh medium is added and the cells are incubated for about 5 days.

[0414] Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200  $\mu$ Ci/ml <sup>35</sup>S-cysteine and 200  $\mu$ Ci/ml <sup>35</sup>S-methionine. After a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of PRO polypeptide. The cultures containing transfected cells may undergo further incubation (in serum free medium) and the medium is tested in selected bioassays.

[0415] In an alternative technique, PRO may be introduced into 293 cells transiently using the dextran sulfate method described by Somparyrac et al., *Proc. Natl. Acad. Sci.*, 12:7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700  $\mu$ g pRK5-PRO DNA is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5  $\mu$ g/ml bovine insulin and 0.1  $\mu$ g/ml bovine transferrin. After about

four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed PRO can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography.

[0416] In another embodiment, PRO can be expressed in CHO cells. The pRK5-PRO can be transfected into CHO cells using known reagents such as CaPO<sub>4</sub> or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as <sup>35</sup>S-methionine. After determining the presence of PRO polypeptide, the culture medium may be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. The medium containing the expressed PRO can then be concentrated and purified by any selected method.

[0417] Epitope-tagged PRO may also be expressed in host CHO cells. The PRO may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-his tag into a Baculovirus expression vector. The poly-his tagged PRO insert can then be subcloned into a SV40 driven vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 driven vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the expressed poly-His tagged PRO can then be concentrated and purified by any selected method, such as by Ni<sup>2+</sup>-chelate affinity chromatography.

[0418] PRO may also be expressed in CHO and/or COS cells by a transient expression procedure or in CHO cells by another stable expression procedure.

[0419] Stable expression in CHO cells is performed using the following procedure. The proteins are expressed as an IgG construct (immunoadhesin), in which the coding sequences for the soluble forms (e.g. extracellular domains) of the respective proteins are fused to an IgG1 constant region sequence containing the hinge, CH2 and CH2 domains and/or is a poly-His tagged form.

[0420] Following PCR amplification, the respective DNAs are subcloned in a CHO expression vector using standard techniques as described in Ausubel et al., *Current Protocols of Molecular Biology*, Unit3.16, John Wiley and Sons (1997). CHO expression vectors are constructed to have compatible restriction sites 5' and 3' of the DNA of interest to allow the convenient shuttling of cDNA's. The vector used expression in CHO cells is as described in Lucas et al., *Nucl. Acids Res.* 24:9 (1774-1779 (1996), and uses the SV40 early promoter/enhancer to drive expression of the cDNA of interest and dihydrofolate reductase (DHFR). DHFR expression permits selection for stable maintenance of the plasmid following transfection.

[0421] Twelve micrograms of the desired plasmid DNA is introduced into approximately 10 million CHO cells using commercially available transfection reagents Superfect® (Qiagen), Dosper® or Fugene® (Boehringer Mannheim). The cells are grown as described in Lucas et al., supra. Approximately 3 $\times$ 10<sup>7</sup> cells are frozen in an ampule for further growth and production as described below.

[0422] The ampules containing the plasmid DNA are thawed by placement into water bath and mixed by vortex-

ing. The contents are pipetted into a centrifuge tube containing 10 mLs of media and centrifuged at 1000 rpm for 5 minutes. The supernatant is aspirated and the cells are resuspended in 10 mL of selective media (0.2  $\mu$ m filtered PS20 with 5% 0.2  $\mu$ m diafiltered fetal bovine serum). The cells are then aliquoted into a 100 mL spinner containing 90 mL of selective media. After 1-2 days, the cells are transferred into a 250 mL spinner filled with 150 mL selective growth medium and incubated at 37° C. After another 2-3 days, 250 mL, 500 mL and 2000 mL spinners are seeded with  $3 \times 10^5$  cells/mL. The cell media is exchanged with fresh media by centrifugation and resuspension in production medium. Although any suitable CHO media may be employed, a production medium described in U.S. Pat. No. 5,122,469, issued Jun. 16, 1992 may actually be used. A 3L production spinner is seeded at  $1.2 \times 10^6$  cells/mL. On day 0, the cell number pH is determined. On day 1, the spinner is sampled and sparging with filtered air is commenced. On day 2, the spinner is sampled, the temperature shifted to 33° C., and 30 mL of 500 g/L glucose and 0.6 mL of 10% antifoam (e.g., 35% polydimethylsiloxane emulsion, Dow Corning 365 Medical Grade Emulsion) taken. Throughout the production, the pH is adjusted as necessary to keep it at around 7.2. After 10 days, or until the viability dropped below 70%, the cell culture is harvested by centrifugation and filtering through a 0.22  $\mu$ m filter. The filtrate was either stored at 4° C. or immediately loaded onto columns for purification.

[0423] For the poly-His tagged constructs, the proteins are purified using a Ni-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 ml Ni-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4° C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein is subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80° C.

[0424] Immunoadhesin (Fc-containing) constructs are purified from the conditioned media as follows. The conditioned medium is pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 ml fractions into tubes containing 275  $\mu$ L of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity is assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation.

[0425] Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

#### Example 8

[0426] Expression of PRO in Yeast

[0427] The following method describes recombinant expression of PRO in yeast.

[0428] First, yeast expression vectors are constructed for intracellular production or secretion of PRO from the

ADH2/GAPDH promoter. DNA encoding PRO and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of PRO. For secretion, DNA encoding PRO can be cloned into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, a native PRO signal peptide or other mammalian signal peptide, or, for example, a yeast alpha-factor or invertase secretory signal/leader sequence, and linker sequences (if needed) for expression of PRO.

[0429] Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

[0430] Recombinant PRO can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected cartridge filters. The concentrate containing PRO may further be purified using selected column chromatography resins.

[0431] Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

#### Example 9

[0432] Expression of PRO in Baculovirus-Infected Insect Cells

[0433] The following method describes recombinant expression of PRO in Baculovirus-infected insect cells.

[0434] The sequence coding for PRO is fused upstream of an epitope tag contained within a baculovirus expression vector. Such epitope tags include poly-his tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the sequence encoding PRO or the desired portion of the coding sequence of PRO such as the sequence encoding the extracellular domain of a transmembrane protein or the sequence encoding the mature protein if the protein is extracellular is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

[0435] Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (PharMingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4-5 days of incubation at 28° C., the released viruses are harvested and used for further amplifications. Viral infection and protein expression are performed as described by O'Reilley et al., *Baculovirus expression vectors: A Laboratory Manual*, Oxford: Oxford University Press (1994).

[0436] Expressed poly-his tagged PRO can then be purified, for example, by Ni<sup>2+</sup>-chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert et al., *Nature*,

362:175-179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 mL Hepes, pH 7.9; 12.5 mM MgCl<sub>2</sub>; 0.1 mM EDTA; 10% glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% glycerol, pH 7.8) and filtered through a 0.45  $\mu$ m filter. A Ni<sup>2+</sup>-NTA agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 mL, washed with 25 mL of water and equilibrated with 25 mL of loading buffer. The filtered cell extract is loaded onto the column at 0.5 mL per minute. The column is washed to baseline A<sub>280</sub> with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% glycerol, pH 6.0), which elutes nonspecifically bound protein. After reaching A<sub>280</sub> baseline again, the column is developed with a 0 to 500 mM Imidazole gradient in the secondary wash buffer. One mL fractions are collected and analyzed by SDS-PAGE and silver staining or Western blot with Ni<sup>2+</sup>-NTA-conjugated to alkaline phosphatase (Qiagen). Fractions containing the eluted His<sub>10</sub>-tagged PRO are pooled and dialyzed against loading buffer.

[0437] Alternatively, purification of the IgG tagged (or Fc tagged) PRO can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

[0438] Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

#### Example 10

[0439] Preparation of Antibodies that Bind PRO

[0440] This example illustrates preparation of monoclonal antibodies which can specifically bind PRO.

[0441] Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, *supra*. Immunogens that may be employed include purified PRO, fusion proteins containing PRO, and cells expressing recombinant PRO on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

[0442] Mice, such as Balb/c, are immunized with the PRO immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1-100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, Mont.) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice may also be boosted with additional immunization injections. Serum samples may be periodically obtained from the mice by retro-orbital bleeding for testing in ELISA assays to detect anti-PRO antibodies.

[0443] After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of PRO. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The

fusions generate hybridoma cells which can then be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

[0444] The hybridoma cells will be screened in an ELISA for reactivity against PRO. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against PRO is within the skill in the art.

[0445] The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-PRO monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

#### Example 11

[0446] Purification of PRO Polypeptides Using Specific Antibodies

[0447] Native or recombinant PRO polypeptides may be purified by a variety of standard techniques in the art of protein purification. For example, pro-PRO polypeptide, mature PRO polypeptide, or pre-PRO polypeptide is purified by immunoaffinity chromatography using antibodies specific for the PRO polypeptide of interest. In general, an immunoaffinity column is constructed by covalently coupling the anti-PRO polypeptide antibody to an activated chromatographic resin.

[0448] Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway, N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated SEPHAROSE™ (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

[0449] Such an immunoaffinity column is utilized in the purification of PRO polypeptide by preparing a fraction from cells containing PRO polypeptide in a soluble form. This preparation is derived by solubilization of the whole cell or of a subcellular fraction obtained via differential centrifugation by the addition of detergent or by other methods well known in the art. Alternatively, soluble PRO polypeptide containing a signal sequence may be secreted in useful quantity into the medium in which the cells are grown.

[0450] A soluble PRO polypeptide-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PRO polypeptide (e.g., high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/PRO polypeptide binding (e.g., a low pH buffer such as approximately pH 2-3,

or a high concentration of a chaotrope such as urea or thiocyanate ion), and PRO polypeptide is collected.

#### Example 12

##### [0451] Drug Screening

[0452] This invention is particularly useful for screening compounds by using PRO polypeptides or binding fragment thereof in any of a variety of drug screening techniques. The PRO polypeptide or fragment employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the PRO polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, the formation of complexes between PRO polypeptide or a fragment and the agent being tested. Alternatively, one can examine the diminution in complex formation between the PRO polypeptide and its target cell or target receptors caused by the agent being tested.

[0453] Thus, the present invention provides methods of screening for drugs or any other agents which can affect a PRO polypeptide-associated disease or disorder. These methods comprise contacting such an agent with an PRO polypeptide or fragment thereof and assaying (i) for the presence of a complex between the agent and the PRO polypeptide or fragment, or (ii) for the presence of a complex between the PRO polypeptide or fragment and the cell, by methods well known in the art. In such competitive binding assays, the PRO polypeptide or fragment is typically labeled. After suitable incubation, free PRO polypeptide or fragment is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular agent to bind to PRO polypeptide or to interfere with the PRO polypeptide/cell complex.

[0454] Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to a polypeptide and is described in detail in WO 84/03564, published on Sep. 13, 1984. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. As applied to a PRO polypeptide, the peptide test compounds are reacted with PRO polypeptide and washed. Bound PRO polypeptide is detected by methods well known in the art. Purified PRO polypeptide can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies can be used to capture the peptide and immobilize it on the solid support.

[0455] This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding PRO polypeptide specifically compete with a test compound for binding to PRO polypeptide or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PRO polypeptide.

#### Example 13

##### [0456] Rational Drug Design

[0457] The goal of rational drug design is to produce structural analogs of biologically active polypeptide of inter-

est (i.e., a PRO polypeptide) or of small molecules with which they interact, e.g., agonists, antagonists, or inhibitors. Any of these examples can be used to fashion drugs which are more active or stable forms of the PRO polypeptide or which enhance or interfere with the function of the PRO polypeptide in vivo (c.f., Hodgson, *Bio/Technology*, 9: 19-21 (1991)).

[0458] In one approach, the three-dimensional structure of the PRO polypeptide, or of an PRO polypeptide-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the PRO polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of the PRO polypeptide may be gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design analogous PRO polypeptide-like molecules or to identify efficient inhibitors. Useful examples of rational drug design may include molecules which have improved activity or stability as shown by Braxton and Wells, *Biochemistry*, 31:7796-7801 (1992) or which act as inhibitors, agonists, or antagonists of native peptides as shown by Athauda et al., *J. Biochem.*, 113:742-746 (1993).

[0459] It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original receptor. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides would then act as the pharmacore.

[0460] By virtue of the present invention, sufficient amounts of the PRO polypeptide may be made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the PRO polypeptide amino acid sequence provided herein will provide guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

#### Example 14

[0461] Ability of PRO Polypeptides to Stimulate the Release of Proteoglycans from Cartilage (Assay 97)

[0462] The ability of various PRO polypeptides to stimulate the release of proteoglycans from cartilage tissue was tested as follows.

[0463] The metacarpophalangeal joint of 4-6 month old pigs was aseptically dissected, and articular cartilage was removed by free hand slicing being careful to avoid the underlying bone. The cartilage was minced and cultured in bulk for 24 hours in a humidified atmosphere of 95% air, 5% CO<sub>2</sub> in serum free (SF) media (DME/F12 1:1) with 0.1% BSA and 100 U/ml penicillin and 100 µg/ml streptomycin.

After washing three times, approximately 100 mg of articular cartilage was aliquoted into microtubes and incubated for an additional 24 hours in the above SF media. PRO polypeptides were then added at 1% either alone or in combination with 18 ng/ml interleukin-1 $\alpha$ , a known stimulator of proteoglycan release from cartilage tissue. The supernatant was then harvested and assayed for the amount of proteoglycans using the 1,9-dimethyl-methylene blue (DMB) colorimetric assay (Farndale and Buttle, *Biochem. Biophys. Acta* 883:173-177 (1985)). A positive result in this assay indicates that the test polypeptide will find use, for example, in the treatment of sports-related joint problems, articular cartilage defects, osteoarthritis or rheumatoid arthritis.

[0464] When various PRO polypeptides were tested in the above assay, the polypeptides demonstrated a marked ability to stimulate release of proteoglycans from cartilage tissue both basally and after stimulation with interleukin-1 $\alpha$  and at 24 and 72 hours after treatment, thereby indicating that these PRO polypeptides are useful for stimulating proteoglycan release from cartilage tissue. As such, these PRO polypeptides are useful for the treatment of sports-related joint problems, articular cartilage defects, osteoarthritis or rheumatoid arthritis. PRO6018 polypeptide testing positive in this assay.

#### Example 15

[0465] Human Microvascular Endothelial Cell Proliferation (Assay 146)

[0466] This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce proliferation of human microvascular endothelial cells in culture and, therefore, function as useful growth factors.

[0467] On day 0, human microvascular endothelial cells were plated in 96-well plates at 1000 cells/well per 100 microliter and incubated overnight in complete media [EBM-2 growth media, plus supplements: IGF-1; ascorbic acid; VEGF; hEGF; hFGF; hydrocortisone, gentamicin (GA-1000), and fetal bovine serum (FBS, Clonetics)]. On day 1, complete media was replaced by basal media [EBM-2 plus 1% FBS] and addition of PRO polypeptides at 1%, 0.1% and 0.01%. On day 7, an assessment of cell proliferation was performed using the ViaLight HS kit [ATP/luciferase Lumitech]. Results are expressed as % of the cell growth observed with control buffer.

[0468] The following PRO polypeptides stimulated human microvascular endothelial cell proliferation in this assay: PRO1313, PRO20080, and PRO21383.

[0469] The following PRO polypeptides inhibited human microvascular endothelial cell proliferation in this assay: PRO6071, PRO4487, and PRO6006.

#### Example 16

[0470] Microarray Analysis to Detect Overexpression of PRO Polypeptides in Cancerous Tumors

[0471] Nucleic acid microarrays, often containing thousands of gene sequences, are useful for identifying differentially expressed genes in diseased tissues as compared to their normal counterparts. Using nucleic acid microarrays,

test and control mRNA samples from test and control tissue samples are reverse transcribed and labeled to generate cDNA probes. The cDNA probes are then hybridized to an array of nucleic acids immobilized on a solid support. The array is configured such that the sequence and position of each member of the array is known. For example, a selection of genes known to be expressed in certain disease states may be arrayed on a solid support. Hybridization of a labeled probe with a particular array member indicates that the sample from which the probe was derived expresses that gene. If the hybridization signal of a probe from a test (disease tissue) sample is greater than hybridization signal of a probe from a control (normal tissue) sample, the gene or genes overexpressed in the disease tissue are identified. The implication of this result is that an overexpressed protein in a diseased tissue is useful not only as a diagnostic marker for the presence of the disease condition, but also as a therapeutic target for treatment of the disease condition.

[0472] The methodology of hybridization of nucleic acids and microarray technology is well known in the art. In the present example, the specific preparation of nucleic acids for hybridization and probes, slides, and hybridization conditions are all detailed in U.S. Provisional Patent Application Serial No. 60/193,767, filed on Mar. 31, 2000 and which is herein incorporated by reference.

[0473] In the present example, cancerous tumors derived from various human tissues were studied for PRO polypeptide-encoding gene expression relative to non-cancerous human tissue in an attempt to identify those PRO polypeptides which are overexpressed in cancerous tumors. Cancerous human tumor tissue from any of a variety of different human tumors was obtained and compared to a "universal" epithelial control sample which was prepared by pooling non-cancerous human tissues of epithelial origin, including liver, kidney, and lung. mRNA isolated from the pooled tissues represents a mixture of expressed gene products from these different tissues. Microarray hybridization experiments using the pooled control samples generated a linear plot in a 2-color analysis. The slope of the line generated in a 2-color analysis was then used to normalize the ratios of (test:control detection) within each experiment. The normalized ratios from various experiments were then compared and used to identify clustering of gene expression. Thus, the pooled "universal control" sample not only allowed effective relative gene expression determinations in a simple 2-sample comparison, it also allowed multi-sample comparisons across several experiments.

[0474] In the present experiments, nucleic acid probes derived from the herein described PRO polypeptide-encoding nucleic acid sequences were used in the creation of the microarray and RNA from a panel of nine different tumor tissues (listed below) were used for the hybridization thereto. A value based upon the normalized ratio:experimental ratio was designated as a "cutoff ratio". Only values that were above this cutoff ratio were determined to be significant. Table 8 below shows the results of these experiments, demonstrating that various PRO polypeptides of the present invention are significantly overexpressed in various human tumor tissues, as compared to a non-cancerous human tissue control or other human tumor tissues. As described above, these data demonstrate that the PRO polypeptides of the present invention are useful not only as diagnostic markers for the presence of one or more cancerous tumors, but also serve as therapeutic targets for the treatment of those tumors.

TABLE 8

Molecule	is overexpressed in:	as compared to normal control:
PRO240	breast tumor	universal normal control
PRO240	lung tumor	universal normal control
PRO256	colon tumor	universal normal control
PRO256	lung tumor	universal normal control
PRO256	breast tumor	universal normal control
PRO306	colon tumor	universal normal control
PRO306	lung tumor	universal normal control
PRO540	lung tumor	universal normal control
PRO540	colon tumor	universal normal control
PRO773	breast tumor	universal normal control
PRO773	colon tumor	universal normal control
PRO698	colon tumor	universal normal control
PRO698	breast tumor	universal normal control
PRO698	lung tumor	universal normal control
PRO698	prostate tumor	universal normal control
PRO698	rectal tumor	universal normal control
PRO3567	colon tumor	universal normal control
PRO3567	breast tumor	universal normal control
PRO3567	lung tumor	universal normal control
PRO826	colon tumor	universal normal control
PRO826	lung tumor	universal normal control
PRO826	breast tumor	universal normal control
PRO826	rectal tumor	universal normal control
PRO826	liver tumor	universal normal control
PRO1002	colon tumor	universal normal control
PRO1002	lung tumor	universal normal control
PRO1068	colon tumor	universal normal control
PRO1068	breast tumor	universal normal control
PRO1030	colon tumor	universal normal control
PRO1030	breast tumor	universal normal control
PRO1030	lung tumor	universal normal control
PRO1030	prostate tumor	universal normal control
PRO1030	rectal tumor	universal normal control
PRO4397	colon tumor	universal normal control
PRO4397	breast tumor	universal normal control
PRO4344	colon tumor	universal normal control
PRO4344	lung tumor	universal normal control
PRO4344	rectal tumor	universal normal control
PRO4407	colon tumor	universal normal control
PRO4407	breast tumor	universal normal control
PRO4407	lung tumor	universal normal control
PRO4407	liver tumor	universal normal control
PRO4407	rectal tumor	universal normal control
PRO4316	colon tumor	universal normal control
PRO4316	prostate tumor	universal normal control
PRO5775	colon tumor	universal normal control
PRO6016	colon tumor	universal normal control
PRO4980	breast tumor	universal normal control
PRO4980	colon tumor	universal normal control
PRO4980	lung tumor	universal normal control
PRO6018	colon tumor	universal normal control
PRO7168	colon tumor	universal normal control
PRO6000	colon tumor	universal normal control
PRO6006	colon tumor	universal normal control
PRO5800	colon tumor	universal normal control
PRO5800	breast tumor	universal normal control
PRO5800	lung tumor	universal normal control
PRO5800	rectal tumor	universal normal control
PRO7476	colon tumor	universal normal control
PRO10268	colon tumor	universal normal control
PRO6496	colon tumor	universal normal control
PRO6496	breast tumor	universal normal control
PRO6496	lung tumor	universal normal control
PRO7422	colon tumor	universal normal control
PRO7431	colon tumor	universal normal control
PRO28633	colon tumor	universal normal control
PRO28633	lung tumor	universal normal control
PRO28633	liver tumor	universal normal control
PRO21485	colon tumor	universal normal control
PRO28700	breast tumor	universal normal control
PRO28700	lung tumor	universal normal control
PRO28700	colon tumor	universal normal control
PRO34012	colon tumor	universal normal control
PRO34012	lung tumor	universal normal control

TABLE 8-continued

Molecule	is overexpressed in:	as compared to normal control:
PRO34003	colon tumor	universal normal control
PRO34003	lung tumor	universal normal control
PRO34001	colon tumor	universal normal control
PRO34009	colon tumor	universal normal control
PRO34009	breast tumor	universal normal control
PRO34009	lung tumor	universal normal control
PRO34009	rectal tumor	universal normal control
PRO34192	colon tumor	universal normal control
PRO34564	colon tumor	universal normal control
PRO35444	colon tumor	universal normal control
PRO5998	colon tumor	universal normal control
PRO5998	lung tumor	universal normal control
PRO5998	kidney tumor	universal normal control
PRO19651	colon tumor	universal normal control
PRO20221	liver tumor	universal normal control
PRO21434	liver tumor	universal normal control

## Example 17

**[0475]** Fetal Hemoglobin Induction in an Erythroblastic Cell Line (Assay 107)

**[0476]** This assay is useful for screening PRO polypeptides for the ability to induce the switch from adult hemoglobin to fetal hemoglobin in an erythroblastic cell line. Molecules testing positive in this assay are expected to be useful for therapeutically treating various mammalian hemoglobin-associated disorders such as the various thalassemias. The assay is performed as follows. Erythroblastic cells are plated in standard growth medium at 1000 cells/well in a 96 well format. PRO polypeptides are added to the growth medium at a concentration of 0.2% or 2% and the cells are incubated for 5 days at 37° C. As a positive control, cells are treated with 100 μM hemin and as a negative control, the cells are untreated. After 5 days, cell lysates are prepared and analyzed for the expression of gamma globin (a fetal marker). A positive in the assay is a gamma globin level at least 2-fold above the negative control.

**[0477]** PRO20080 polypeptide tested positive in this assay.

## Example 18

**[0478]** Microarray Analysis to Detect Overexpression of PRO Polypeptides in HUVEC Cells Treated with Growth Factors

**[0479]** This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce angiogenesis by stimulating endothelial cell tube formation in HUVEC cells.

**[0480]** Nucleic acid microarrays, often containing thousands of gene sequences, are useful for identifying differentially expressed genes in tissues exposed to various stimuli (e.g., growth factors) as compared to their normal, unexposed counterparts. Using nucleic acid microarrays, test and control mRNA samples from test and control tissue samples are reverse transcribed and labeled to generate cDNA probes. The cDNA probes are then hybridized to an array of nucleic acids immobilized on a solid support. The array is configured such that the sequence and position of

each member of the array is known. Hybridization of a labeled probe with a particular array member indicates that the sample from which the probe was derived expresses that gene. If the hybridization signal of a probe from a test (exposed tissue) sample is greater than hybridization signal of a probe from a control (normal, unexposed tissue) sample, the gene or genes overexpressed in the exposed tissue are identified. The implication of this result is that an overexpressed protein in an exposed tissue may be involved in the functional changes within the tissue following exposure to the stimuli (e.g., tube formation).

**[0481]** The methodology of hybridization of nucleic acids and microarray technology is well known in the art. In the present example, the specific preparation of nucleic acids for hybridization and probes, slides, and hybridization conditions are all detailed in U.S. Provisional Patent Application Serial No. 60/193,767, filed on Mar. 31, 2000 and which is herein incorporated by reference.

**[0482]** In the present example, HUVEC cells grown in either collagen gels or fibrin gels were induced to form tubes by the addition of various growth factors. Specifically, collagen gels were prepared as described previously in Yang et al., *American J. Pathology*, 1999, 155(3):887-895 and Xin et al., *American J. Pathology*, 2001, 158(3):1111-1120. Following gelation of the HUVEC cells, 1×basal medium containing M199 supplemented with 1% FBS, 1×ITS, 2 mM L-glutamine, 50 µg/ml ascorbic acid, 26.5 mM NaHCO<sub>3</sub>, 100 U/ml penicillin and 100 U/ml streptomycin was added. Tube formation was elicited by the inclusion in the culture media of either a mixture of phorbol myristate acetate (50 nM), vascular endothelial cell growth factor (40 ng/ml) and basic fibroblast growth factor (40 ng/ml) ("PMA growth factor mix") or hepatocyte growth factor (40 ng/ml) and vascular endothelial cell growth factor (40 ng/ml) (HGF/VEGF mix) for the indicated period of time. Fibrin Gels were prepared by suspending Huvec (4×10<sup>5</sup> cells/ml) in M199 containing 1% fetal bovine serum (Hyclone) and human fibrinogen (2.5 mg/ml). Thrombin (50 U/ml) was then added to the fibrinogen suspension at a ratio of 1 part thrombin solution:30 parts fibrinogen suspension. The solution was then layered onto 10 cm tissue culture plates (total volume: 15 ml/plate) and allowed to solidify at 37° C. for 20 min. Tissue culture media (10 ml of BM containing PMA (50 nM), bFGF (40 ng/ml) and VEGF (40 ng/ml)) was then added and the cells incubated at 37° C. in 5% CO<sub>2</sub> in air for the indicated period of time.

**[0483]** Total RNA was extracted from the HUVEC cells incubated for 0, 4, 8, 24, 40 and 50 hours in the different matrix and media combinations using a TRIzol extraction followed by a second purification using RNeasy Mini Kit (Qiagen). The total RNA was used to prepare cRNA which was then hybridized to the microarrays.

**[0484]** In the present experiments, nucleic acid probes derived from the herein described PRO polypeptide-encoding nucleic acid sequences were used in the creation of the microarray and RNA from the HUVEC cells described above were used for the hybridization thereto. Pairwise comparisons were made using time 0 chips as a baseline. Three replicate samples were analyzed for each experimental condition and time. Hence there were 3 time 0 samples for each treatment and 3 replicates of each successive time point. Therefore, a 3 by 3 comparison was performed for

each time point compared against each time 0 point. This resulted in 9 comparisons per time point. Only those genes that had increased expression in all three non-time-0 replicates in each of the different matrix and media combinations as compared to any of the three time zero replicates were considered positive. Although this stringent method of data analysis does allow for false negatives, it minimizes false positives.

**[0485]** PRO281, PRO1560, PRO189, PRO4499, PRO6308, PRO6000, PRO10275, PRO21207, PRO20933, and PRO34274 tested positive in this assay.

#### Example 19

**[0486]** Tumor Versus Normal Differential Tissue Expression Distribution

**[0487]** Oligonucleotide probes were constructed from some of the PRO polypeptide-encoding nucleotide sequences shown in the accompanying figures for use in quantitative PCR amplification reactions. The oligonucleotide probes were chosen so as to give an approximately 200-600 base pair amplified fragment from the 3' end of its associated template in a standard PCR reaction. The oligonucleotide probes were employed in standard quantitative PCR amplification reactions with cDNA libraries isolated from different human tumor and normal human tissue samples and analyzed by agarose gel electrophoresis so as to obtain a quantitative determination of the level of expression of the PRO polypeptide-encoding nucleic acid in the various tumor and normal tissues tested. β-actin was used as a control to assure that equivalent amounts of nucleic acid was used in each reaction. Identification of the differential expression of the PRO polypeptide-encoding nucleic acid in one or more tumor tissues as compared to one or more normal tissues of the same tissue type renders the molecule useful diagnostically for the determination of the presence or absence of tumor in a subject suspected of possessing a tumor as well as therapeutically as a target for the treatment of a tumor in a subject possessing such a tumor. These assays provided the following results:

**[0488]** (1) DNA161005-2943 molecule is very highly expressed in human umbilical vein endothelial cells (HUVEC), substantia niagra, hippocampus and dendrocytes; highly expressed in lymphoblasts; expressed in spleen, prostate, uterus and macrophages; and is weakly expressed in cartilage and heart. Among a panel of normal and tumor tissues examined, it is expressed in esophageal tumor, and is not expressed in normal esophagus, normal stomach, stomach tumor, normal kidney, kidney tumor, normal lung, lung tumor, normal rectum, rectal tumor, normal liver and liver tumor.

**[0489]** (2) DNA170245-3053 molecule is highly expressed in cartilage, testis, adrenal gland, and uterus, and not expressed in HUVEC, colon tumor, heart, placenta, bone marrow, spleen and aortic endothelial cells. In a panel of tumor and normal tissue samples examined, the DNA170245-3053 molecule was found to be expressed in normal esophagus and esophageal tumor, expressed in normal stomach and in stomach tumor, not expressed in normal kidney, but expressed in kidney tumor, not expressed in normal lung, but expressed in lung



- tumor, not expressed in normal rectum nor in rectal tumor, and not expressed in normal liver, but is expressed in liver tumor.
- [0490] (3) DNA173157-2981 molecule is significantly expressed in the following tissues: cartilage, testis, HUVEC, heart, placenta, bone marrow, adrenal gland, prostate, spleen, aortic endothelial cells, and uterus. When these assays were conducted on a tumor tissue panel, it was found that the DNA173157-2981 molecule is significantly expressed in the following tissues: normal esophagus and esophageal tumor, normal stomach and stomach tumor, normal kidney and kidney tumor, normal lung and lung tumor, normal rectum and rectal tumor, normal liver and liver tumor, and colon tumor.
- [0491] (4) DNA175734-2985 molecule is significantly expressed in the adrenal gland and the uterus. The DNA175734-2985 molecule is not significantly expressed in the following tissues: cartilage, testis, HUVEC, colon tumor, heart, placenta, bone marrow, prostate, spleen and aortic endothelial cells. Screening of a tumor panel revealed that DNA175734-2985 is significantly expressed in normal esophagus but not in esophageal tumor. Similarly, while highly expressed in normal rectum, DNA175734-2985 is expressed to a lesser extent in rectal tumor. DNA175734-2985 is expressed equally in normal stomach and stomach tumor as well as normal liver and liver tumor. While not expressed in normal kidney, DNA175734-2985 is highly expressed in kidney tumor.
- [0492] (5) DNA176108-3040 molecule is highly expressed in prostate and uterus, expressed in cartilage, testis, heart, placenta, bone marrow, adrenal gland and spleen, and not significantly expressed in HUVEC, colon tumor, and aortic endothelial cells. In a panel of tumor and normal tissue samples examined, the DNA176108-3040 molecule was found to be highly expressed in normal esophagus, but expressed at lower levels in esophageal tumor, highly expressed in normal stomach, and expressed at a lower level in stomach tumor, expressed in kidney and in kidney tumor, expressed in normal rectum and at a lower level in rectal tumor, and expressed in normal liver and not expressed in liver tumor.
- [0493] (6) DNA191064-3069 molecule is significantly expressed in the following tissues: cartilage, testis, HUVEC, heart, placenta, bone marrow, adrenal gland, prostate, spleen, aortic endothelial cells, and uterus and not significantly expressed in colon tumor. In a panel of tumor and normal tissue samples, the DNA191064-3069 molecule was found to be expressed in normal esophagus and in esophageal tumors, expressed in normal stomach and in stomach tumors, expressed in normal kidney and in kidney tumors, expressed in normal lung and in lung tumors, expressed in normal rectum and in rectal tumors, expressed in normal liver and in liver tumors.
- [0494] (7) DNA194909-3013 molecule is highly expressed in placenta, and expressed in cartilage, testis, HUVEC, colon tumor, heart, bone marrow, adrenal gland, prostate, spleen, aortic endothelial cells and uterus. In a panel of tumor and normal tissue samples examined, the DNA194909-3013 molecule was found to be expressed in normal esophagus and esophageal tumor, not expressed in normal stomach nor stomach tumor, expressed in normal kidney and kidney tumor, expressed in normal lung and lung tumor, expressed in normal rectum and rectal tumor, and not expressed in normal liver, but is expressed in liver tumor.
- [0495] (8) The PRO34009 encoding genes of the invention (DNA203532-3029) were screened in normal tissues and the following primary tumors and the resulting values are reported below.
- [0496] Tumor Panel:
- [0497] PRO34009 encoding genes were expressed 39.3 fold higher in lung tumor than normal lung. It is expressed 9.5 fold higher in esophageal tumors than normal esophagus. It is expressed 6.7 fold higher in kidney tumor than normal kidney. It is expressed 4.0 fold higher in colon tumor than normal colon. It is expressed 2.7 fold higher in stomach tumor than normal stomach. It is expressed at similar levels in normal rectum and rectal tumor, normal liver and liver tumor, normal uterus and uterine tumor.
- [0498] Normal Panel:
- [0499] For the normal tissue values, the normal tissue with the highest expression, in this case normal thymus, was given a value of 1 and all other normal tissues were given a value of less than 1, and described as expressed, weakly expressed or not expressed, based on their expression relative to thymus. PRO34009 encoding genes were expressed in normal thymus. It is weakly expressed in lymphoblast, spleen, heart, fetal limb, fetal lung, placenta, HUVEC, testis, fetal kidney, uterus, prostate, macrophage, substantia nigra, hippocampus, liver, skin, esophagus, stomach, rectum, kidney, thyroid, skeletal muscle, or fetal articular cartilage. It is not expressed in bone marrow, fetal liver, colon, lung or dendrocytes.
- [0500] (9) DNA213858-3060 molecule is not significantly expressed in cartilage, testis, HUVEC, colon tumor, heart, placenta, bone marrow, adrenal gland, prostate, spleen, aortic endothelial cells or uterus. In a panel of tumor and normal tissue samples examined, the DNA213858-3060 molecule was found to be expressed in normal esophagus and esophageal tumor, expressed in normal stomach and in stomach tumor, expressed in normal kidney and kidney tumor, expressed in normal lung and in lung tumor, expressed in normal rectum and in rectal tumor, and expressed in normal liver and in liver tumor.
- [0501] (10) DNA216676-3083 molecule is significantly expressed in the following tissues: testis, heart, bone marrow, and uterus, and not significantly expressed in the following tissues: cartilage, HUVEC, colon tumor, placenta, adrenal gland, prostate, spleen, or aortic endothelial cells. In a panel of tumor and normal tissues samples examined, the DNA216676-3083 molecule was found to be expressed in normal esophagus and esophageal tumor, not expressed in normal stomach, but is ex-

pressed in stomach tumor, not expressed in normal kidney nor in kidney tumor, not expressed in normal lung, but is expressed in lung tumor, not expressed in normal rectum, but is expressed in rectal tumor, and not expressed in normal liver nor in liver tumor.

[0502] (11) DNA222653-3104 molecule is significantly expressed testis, and not significantly expressed in cartilage, HUVEC, colon tumor, heart, placenta, bone marrow, adrenal gland, prostate, spleen, aortic endothelial cells and uterus. In a panel of tumor and normal tissue samples examined, the DNA222653-3104 molecule was not expressed in normal esophagus, esophageal tumor, normal stomach, stomach tumor, normal kidney, kidney tumor, normal lung, lung tumor, normal rectum, rectal tumor, normal liver and liver tumor.

#### Example 20

[0503] Guinea Pig Vascular Leak (Assay 51)

[0504] This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce vascular permeability. Polypeptides testing positive in this assay are expected to be useful for the therapeutic treatment of conditions which would benefit from enhanced vascular permeability including, for example, conditions which may benefit from enhanced local immune system cell infiltration.

[0505] Hairless guinea pigs weighing 350 grams or more were anesthetized with Ketamine (75-80 mg/kg) and 5 mg/kg Xylazine intramuscularly. Test samples containing the PRO polypeptide or a physiological buffer without the test polypeptide are injected into skin on the back of the test animals with 100  $\mu$ l per injection site intradermally. There were approximately 16-24 injection sites per animal. One ml of Evans blue dye (1% in PBS) is then injected intracardially. Skin vascular permeability responses to the compounds (i.e., blemishes at the injection sites of injection) are visually scored by measuring the diameter (in mm) of blue-colored leaks from the site of injection at 1 and 6 hours post administration of the test materials. The mm diameter of blueness at the site of injection is observed and recorded as well as the severity of the vascular leakage. Blemishes of

at least 5 mm in diameter are considered positive for the assay when testing purified proteins, being indicative of the ability to induce vascular leakage or permeability. A response greater than 7 mm diameter is considered positive for conditioned media samples. Human VEGF at 0.1  $\mu$ g/100  $\mu$ l is used as a positive control, inducing a response of 15-23 mm diameter.

[0506] PRO19822 polypeptides tested positive in this assay.

#### Example 21

[0507] Skin Vascular Permeability Assay (Assay 64)

[0508] This assay shows that certain polypeptides of the invention stimulate an immune response and induce inflammation by inducing mononuclear cell, eosinophil and PMN infiltration at the site of injection of the animal. Compounds which stimulate an immune response are useful therapeutically where stimulation of an immune response is beneficial. This skin vascular permeability assay is conducted as follows. Hairless guinea pigs weighing 350 grams or more are anesthetized with ketamine (75-80 mg/Kg) and 5 mg/Kg xylazine intramuscularly (IM). A sample of purified polypeptide of the invention or a conditioned media test sample is injected intradermally onto the backs of the test animals with 100  $\mu$ l per injection site. It is possible to have about 10-30, preferably about 16-24, injection sites per animal. One  $\mu$ l of Evans blue dye (1% in physiologic buffered saline) is injected intracardially. Blemishes at the injection sites are then measured (mm diameter) at 1 hr and 6 hr post injection. Animals were sacrificed at 6 hrs after injection. Each skin injection site is biopsied and fixed in formalin. The skins are then prepared for histopathologic evaluation. Each site is evaluated for inflammatory cell infiltration into the skin. Sites with visible inflammatory cell inflammation are scored as positive. Inflammatory cells may be neutrophilic, eosinophilic, monocytic or lymphocytic. At least a minimal perivascular infiltrate at the injection site is scored as positive, no infiltrate at the site of injection is scored as negative.

[0509] PRO19822 polypeptide tested positive in this assay.

---

#### SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 116

<210> SEQ ID NO 1  
<211> LENGTH: 1943  
<212> TYPE: DNA  
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 1

cgagcgcgtg ggtgcgaggc gaaggtgacc gggaccgag catttcagat	50
ctgctcggta gacctggtgc accaccacca tgttgctgc aaggctggtg	100
tgtctcggga cactaccttc tagggttttc caccagctt tcaccaaggc	150
ctcccctgtt gtgaagaatt ccatcacgaa gaatcaatgg ctgttaacac	200

-continued

ctagcagga atatgccacc aaacaagaa ttgggatccg gcgtgggaga	250
actggccaag aactcaaaga ggcagcattg gaaccatcga tggaaaaaat	300
atntaaaatt gatcagatgg gaagatgggt tggctgctga ggggctgctg	350
ttggtccttg agcattgtgc tactatggct tgggactgtc taatgagatt	400
ggagctattg aaaaggctgt aatttggcct cagtatgtca aggatagaat	450
tcattccacc tatatgtact tagcaggag tattggttta acagctttgt	500
ctgccatagc aatcagcaga acgcctgttc tcatgaactt catgatgaga	550
ggctcctggg tgacaattgg tgtgaccttt gcagccatgg ttggagctgg	600
aatgctggta cgatcaatac catatgacca gagccaggc ccaaagcatc	650
ttgcttggtt gctacattct ggtgtgatgg gtgcagtggg ggctcctctg	700
acaatattag ggggtcctct tctcatcaga gctgcatggt acacagctgg	750
cattgtggga ggcctctcca ctgtggccat gtgtgccc agtgaaaagt	800
ttctgaacat ggggtcaccc ctgggagtg gctgggtct cgtctttgtg	850
tcctcattgg gatctatggt tcttccacct accaccgtgg ctggtgccac	900
tctttactca gtggcaatgt acggtggatt agttcttttc agcatgttc	950
ttctgtatga taccagaaa gtaatcaagc gtgcagaagt atcaccaatg	1000
tatggagttc aaaaatatga tcccattaac togatgctga gtatctacat	1050
ggatacatta aatataatga tgcgagttgc aactatgctg gcaactggag	1100
gcaacagaaa gaaatgaagt gactcagctt ctggcttctc tgctacatca	1150
aatactctgt ttaatggggc agatagcat taaatagttt gtacaagcag	1200
ctttcgttga agtttagaag ataagaaaca tgtcatcata tttaaatggt	1250
ccgtaaatgt gatgcctcag gtctgccttt tttctggag aataaatgca	1300
gtaatcctct cccaataag cacacacatt ttcaattctc atgtttgagt	1350
gattttaaaa tgttttggtg aatgtgaaaa ctaaagtttg tgtcatgaga	1400
atgtaagtct ttttctact ttaaaattha gtaggtcac tgagtaacta	1450
aaatttagca aacctgtgtt tgcataatth tttggagtgc agaattgtt	1500
aattaatgtc ataagtgatt tggagctttg gtaaaggac cagagagaag	1550
gagtcacctg cagtcttttg tttttttaa tacttagaac ttagcacttg	1600
tgttattgat tagtgaggag ccagtaagaa acatctgggt atttgaaac	1650
aagtgtcat tgttacattc atttgctgaa cttacaaaa ctgttcatcc	1700
tgaaacagcg acaggtgatg cattctcctg ctgttgcttc tcagtgetct	1750
ctttccaata tagatgtggt catgtttgac ttgtacagaa tgttaatcat	1800
acagagaatc cttgatggaa ttatatatgt gtgttttact tttgaatggt	1850
acaaaaggaa ataactttaa aactattctc aagagaaaat attcaaagca	1900
tgaaatatgt tgctttttcc agaatacaaa cagtatactc atg	1943

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 345

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

-continued

&lt;400&gt; SEQUENCE: 2

---

```

Met Leu Ala Ala Arg Leu Val Cys Leu Arg Thr Leu Pro Ser Arg
 1           5           10           15
Val Phe His Pro Ala Phe Thr Lys Ala Ser Pro Val Val Lys Asn
          20           25           30
Ser Ile Thr Lys Asn Gln Trp Leu Leu Thr Pro Ser Arg Glu Tyr
          35           40           45
Ala Thr Lys Thr Arg Ile Gly Ile Arg Arg Gly Arg Thr Gly Gln
          50           55           60
Glu Leu Lys Glu Ala Ala Leu Glu Pro Ser Met Glu Lys Ile Phe
          65           70           75
Lys Ile Asp Gln Met Gly Arg Trp Phe Val Ala Gly Gly Ala Ala
          80           85           90
Val Gly Leu Gly Ala Leu Cys Tyr Tyr Gly Leu Gly Leu Ser Asn
          95           100          105
Glu Ile Gly Ala Ile Glu Lys Ala Val Ile Trp Pro Gln Tyr Val
          110          115          120
Lys Asp Arg Ile His Ser Thr Tyr Met Tyr Leu Ala Gly Ser Ile
          125          130          135
Gly Leu Thr Ala Leu Ser Ala Ile Ala Ile Ser Arg Thr Pro Val
          140          145          150
Leu Met Asn Phe Met Met Arg Gly Ser Trp Val Thr Ile Gly Val
          155          160          165
Thr Phe Ala Ala Met Val Gly Ala Gly Met Leu Val Arg Ser Ile
          170          175          180
Pro Tyr Asp Gln Ser Pro Gly Pro Lys His Leu Ala Trp Leu Leu
          185          190          195
His Ser Gly Val Met Gly Ala Val Val Ala Pro Leu Thr Ile Leu
          200          205          210
Gly Gly Pro Leu Leu Ile Arg Ala Ala Trp Tyr Thr Ala Gly Ile
          215          220          225
Val Gly Gly Leu Ser Thr Val Ala Met Cys Ala Pro Ser Glu Lys
          230          235          240
Phe Leu Asn Met Gly Ala Pro Leu Gly Val Gly Leu Gly Leu Val
          245          250          255
Phe Val Ser Ser Leu Gly Ser Met Phe Leu Pro Pro Thr Thr Val
          260          265          270
Ala Gly Ala Thr Leu Tyr Ser Val Ala Met Tyr Gly Gly Leu Val
          275          280          285
Leu Phe Ser Met Phe Leu Leu Tyr Asp Thr Gln Lys Val Ile Lys
          290          295          300
Arg Ala Glu Val Ser Pro Met Tyr Gly Val Gln Lys Tyr Asp Pro
          305          310          315
Ile Asn Ser Met Leu Ser Ile Tyr Met Asp Thr Leu Asn Ile Phe
          320          325          330
Met Arg Val Ala Thr Met Leu Ala Thr Gly Gly Asn Arg Lys Lys
          335          340          345

```

&lt;210&gt; SEQ ID NO 3

&lt;211&gt; LENGTH: 1110

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapien

-continued

&lt;400&gt; SEQUENCE: 3

```

ccaatcgccc ggtgcggtgg tgcagggtct cgggctagtc atggcgctccc      50
cgtctcggag actgcagact aaaccagtca ttacttgttt caagagcgtt      100
ctgctaactc acacttttat tttctggatc actggcgtaa tccttcttgc      150
agttggcatt tggggcaagg tgagcctgga gaattacttt tctcttttaa      200
atgagaaggc caccaatgtc cccttcgtgc tcattgctac tggtagcgtc      250
attattcttt tgggcacctt tggttgtttt gctacctgcc gagcttctgc      300
atggatgcta aaactgtatg caatgtttct gactctcggt tttttggtcg      350
aactggtcgc tgccatcgta ggatttgttt tcagacatga gattaagaac      400
agctttaaga ataattatga gaaggctttg aagcagtata actctacag      450
agattataga agccatgcag tagacaagat ccaaaatagc ttgcattggt      500
gtggtgtcac cgattataga gattggacag atactaatta ttactcagaa      550
aaagatttcc ctaagagttg ctgtaaactt gaagattgta ctccacagag      600
agatgcagac aaagtaaaca atgaaggttg tttataaag gtgatgacca      650
ttatagagtc agaaatggga gtcgttcgag gaatttcctt tggagttgct      700
tgcttccaac tgattggaat ctttctcgcc tactgcccwt ctcgtgccat      750
aacaataaac cagtatgaga tagtgaacc caatgatctc gtgggcctat      800
tcctctctac cttaaggac atttagggtc cccctgtga attagaaagt      850
tgcttggtcg gagaactgac aacactactt actgatagac caaaaaacta      900
caccagtagg ttgattcaat caagatglat gtagacctaa aactacacca      950
ataggctgat tcaatcaaga tccgtgctcg cagtggtgctg attcaatcaa      1000
gatgtatggt tgctatgttc taagtccacc ttctatccca ttcgatgtag      1050
atcgttgaaa ccctgtatcc ctctgaaaca ctggaagagc tagtaaattg      1100
taaatgaagt                                     1110

```

&lt;210&gt; SEQ ID NO 4

&lt;211&gt; LENGTH: 245

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: unsure

&lt;222&gt; LOCATION: 233

&lt;223&gt; OTHER INFORMATION: unknown amino acid

&lt;400&gt; SEQUENCE: 4

```

Met Ala Ser Pro Ser Arg Arg Leu Gln Thr Lys Pro Val Ile Thr
 1                5                10                15
Cys Phe Lys Ser Val Leu Leu Ile Tyr Thr Phe Ile Phe Trp Ile
                20                25                30
Thr Gly Val Ile Leu Leu Ala Val Gly Ile Trp Gly Lys Val Ser
                35                40                45
Leu Glu Asn Tyr Phe Ser Leu Leu Asn Glu Lys Ala Thr Asn Val
                50                55                60
Pro Phe Val Leu Ile Ala Thr Gly Thr Val Ile Ile Leu Leu Gly
                65                70                75

```

-continued

Thr	Phe	Gly	Cys	Phe	Ala	Thr	Cys	Arg	Ala	Ser	Ala	Trp	Met	Leu
				80					85					90
Lys	Leu	Tyr	Ala	Met	Phe	Leu	Thr	Leu	Val	Phe	Leu	Val	Glu	Leu
				95					100					105
Val	Ala	Ala	Ile	Val	Gly	Phe	Val	Phe	Arg	His	Glu	Ile	Lys	Asn
				110					115					120
Ser	Phe	Lys	Asn	Asn	Tyr	Glu	Lys	Ala	Leu	Lys	Gln	Tyr	Asn	Ser
				125					130					135
Thr	Gly	Asp	Tyr	Arg	Ser	His	Ala	Val	Asp	Lys	Ile	Gln	Asn	Thr
				140					145					150
Leu	His	Cys	Cys	Gly	Val	Thr	Asp	Tyr	Arg	Asp	Trp	Thr	Asp	Thr
				155					160					165
Asn	Tyr	Tyr	Ser	Glu	Lys	Gly	Phe	Pro	Lys	Ser	Cys	Cys	Lys	Leu
				170					175					180
Glu	Asp	Cys	Thr	Pro	Gln	Arg	Asp	Ala	Asp	Lys	Val	Asn	Asn	Glu
				185					190					195
Gly	Cys	Phe	Ile	Lys	Val	Met	Thr	Ile	Ile	Glu	Ser	Glu	Met	Gly
				200					205					210
Val	Val	Ala	Gly	Ile	Ser	Phe	Gly	Val	Ala	Cys	Phe	Gln	Leu	Ile
				215					220					225
Gly	Ile	Phe	Leu	Ala	Tyr	Cys	Xaa	Ser	Arg	Ala	Ile	Thr	Asn	Asn
				230					235					240
Gln	Tyr	Glu	Ile	Val										
				245										

<210> SEQ ID NO 5  
 <211> LENGTH: 1373  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 5

ggggccgcgg	tctagggcgg	ctacgtgtgt	tgccatagcg	accattttgc	50
attaactggt	tggtagcttc	tatcctgggg	gctgagcgc	tgccggccag	100
ctcttccct	actccctctc	ggctccttgt	ggcccaaagg	cctaaccggg	150
gtccggcggg	ctggcctagg	gatcttcccc	gttggcccctt	tggggcggga	200
tggtgcgcca	agaagaagac	gaggtggagt	gggtagtgga	gagcatcgcg	250
gggttcctgc	gaggcccaga	ctggctccatc	cccattcttg	actttgtgga	300
acagaaatgt	gaagttaact	gcaaaggagg	gcatgtgata	actccaggaa	350
gcccagagcc	ggtgattttg	gtggcctgtg	ttccccttgt	ttttgatgat	400
gaagaagaaa	gcaaattgac	ctatacagag	attcatcagg	aatacaaaaga	450
actagttaa	aagctgttag	aaggttacct	caaagaaatt	ggaattaatg	500
aagatcaatt	tcaagaagca	tgcaactctc	ctcttgcaaa	gaccataca	550
tcacaggcca	ttttgcaacc	tgtgttgcca	gcagaagatt	ttactatctt	600
taaagcaatg	atggtccaga	aaaacattga	aatgcagctg	caagccattc	650
gaataattca	agagagaaat	ggtgtattac	ctgactgctt	aaccgatggc	700
tctgatgtgg	tcagtgacct	tgaacacgaa	gagatgaaaa	tcctgagggg	750
agttcttaga	aaatcaaaaag	aggaatatga	ccaggaagaa	gaaaggaaga	800

-continued

---

```

ggaaaaaaca gttatcagag gctaaaacag aagagcccac agtgcattcc      850
agtgaagctg caataatgaa taattcccaa ggggatggtg aacattttgc      900
acacccaccc tcagaagtta aaatgcattt tgctaatacag tcaatagaac      950
ctttgggaag aaaagtggaa aggtctgaaa cttcctccct cccacaaaaa     1000
ggcctgaaga ttcttgctt agagcatgag agcattgaag gaccaatagc     1050
aaacttatca gtacttggaa cagaagaact tcggcaacga gaacactatc     1100
tcaagcagaa gagagataag ttgatgtcca tgagaaagga tatgaggact     1150
aaacagatac aaaatatgga gcagaaagga aaacccactg gggaggtaga     1200
ggaaatgaca gagaaccag aaatgacagc agaggagaag caaacattac     1250
taaagaggag attgcttgcg gagaaactca aagaagaagt tattaataag     1300
taataattaa gaacaattta acaaaatgga agttcaaatt gtcttaaaaa     1350
taaattattt agtccttaca ctg                                     1373

```

&lt;210&gt; SEQ ID NO 6

&lt;211&gt; LENGTH: 367

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 6

```

Met Ala Ala Glu Glu Glu Asp Glu Val Glu Trp Val Val Glu Ser
 1          5          10          15
Ile Ala Gly Phe Leu Arg Gly Pro Asp Trp Ser Ile Pro Ile Leu
          20          25          30
Asp Phe Val Glu Gln Lys Cys Glu Val Asn Cys Lys Gly Gly His
          35          40          45
Val Ile Thr Pro Gly Ser Pro Glu Pro Val Ile Leu Val Ala Cys
          50          55          60
Val Pro Leu Val Phe Asp Asp Glu Glu Glu Ser Lys Leu Thr Tyr
          65          70          75
Thr Glu Ile His Gln Glu Tyr Lys Glu Leu Val Glu Lys Leu Leu
          80          85          90
Glu Gly Tyr Leu Lys Glu Ile Gly Ile Asn Glu Asp Gln Phe Gln
          95          100          105
Glu Ala Cys Thr Ser Pro Leu Ala Lys Thr His Thr Ser Gln Ala
          110          115          120
Ile Leu Gln Pro Val Leu Ala Ala Glu Asp Phe Thr Ile Phe Lys
          125          130          135
Ala Met Met Val Gln Lys Asn Ile Glu Met Gln Leu Gln Ala Ile
          140          145          150
Arg Ile Ile Gln Glu Arg Asn Gly Val Leu Pro Asp Cys Leu Thr
          155          160          165
Asp Gly Ser Asp Val Val Ser Asp Leu Glu His Glu Glu Met Lys
          170          175          180
Ile Leu Arg Glu Val Leu Arg Lys Ser Lys Glu Glu Tyr Asp Gln
          185          190          195
Glu Glu Glu Arg Lys Arg Lys Lys Gln Leu Ser Glu Ala Lys Thr
          200          205          210
Glu Glu Pro Thr Val His Ser Ser Glu Ala Ala Ile Met Asn Asn
          215          220          225

```

-continued

---

Ser Gln Gly Asp Gly Glu His Phe Ala His Pro Pro Ser Glu Val  
 230 235 240

Lys Met His Phe Ala Asn Gln Ser Ile Glu Pro Leu Gly Arg Lys  
 245 250 255

Val Glu Arg Ser Glu Thr Ser Ser Leu Pro Gln Lys Gly Leu Lys  
 260 265 270

Ile Pro Gly Leu Glu His Ala Ser Ile Glu Gly Pro Ile Ala Asn  
 275 280 285

Leu Ser Val Leu Gly Thr Glu Glu Leu Arg Gln Arg Glu His Tyr  
 290 295 300

Leu Lys Gln Lys Arg Asp Lys Leu Met Ser Met Arg Lys Asp Met  
 305 310 315

Arg Thr Lys Gln Ile Gln Asn Met Glu Gln Lys Gly Lys Pro Thr  
 320 325 330

Gly Glu Val Glu Glu Met Thr Glu Lys Pro Glu Met Thr Ala Glu  
 335 340 345

Glu Lys Gln Thr Leu Leu Lys Arg Arg Leu Leu Ala Glu Lys Leu  
 350 355 360

Lys Glu Glu Val Ile Asn Lys  
 365

<210> SEQ ID NO 7  
 <211> LENGTH: 932  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien  
 <220> FEATURE:  
 <221> NAME/KEY: unsure  
 <222> LOCATION: 911  
 <223> OTHER INFORMATION: unknown base

<400> SEQUENCE: 7

```

gggaacggaa aatggcgccct cacggcccgg gtagtcttac gaccctggtg      50
ccctgggctg cgcacctgct cctcgctctg ggcgtgaaa ggcctctggc      100
gctacccgag atatgcaccc aatgtccagg gagcgtgcaa aatttgtcaa      150
aagtggcctt ttattgtaaa acgacacgag agctaagtct gcatgcccgt      200
tgctgcctga atcagaaggg caccatcttg gggctggatc tccagaactg      250
ttctctggag gaccctggtc caaactttca tcaggcacat accactgtca      300
tcatagacct gcaagcaaac cccctcaaag gtgacttggc caacaccttc      350
cgtggcttta ctacgtcca gactctgata ctgccacaac atgtcaactg      400
tcctggagga attaatgcct ggaatactat cacctcttat atagacaacc      450
aaatctgtca agggcaaaag aacctttgca ataactctgg ggaccagaa      500
atgtgtcctg agaatggatc ttgtgtacct gatggtccag gtcttttgca      550
gtgtgtttgt gctgatggtt tccatggata caagtgtatg cgccagggtc      600
cgttctcact gcttatgttc ttcgggatcc tgggagccac cactctatcc      650
gtctccattc tgctttgggc gaccagcgc cgaaaagcca agacttcatg      700
aactacatag gtcttaccat tgacctaaaga tcaatctgaa ctatcttagc      750
ccagtcaggg agctctgctt cctagaaaag catctttcgc cagtggatcc      800
gcctcaaggt tgaggccgcc attggaagat gaaaaattgc actoccttgg      850
    
```



-continued

tgtagacaaa taccagttcc cattggtgtt gttgcctata ataaacactt 900

tttctttttt naaaaaaaaa aaaaaaaaaa aa 932

<210> SEQ ID NO 8

<211> LENGTH: 229

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 8

Met Ala Pro His Gly Pro Gly Ser Leu Thr Thr Leu Val Pro Trp  
1 5 10 15

Ala Ala Ala Leu Leu Leu Ala Leu Gly Val Glu Arg Ala Leu Ala  
20 25 30

Leu Pro Glu Ile Cys Thr Gln Cys Pro Gly Ser Val Gln Asn Leu  
35 40 45

Ser Lys Val Ala Phe Tyr Cys Lys Thr Thr Arg Glu Leu Met Leu  
50 55 60

His Ala Arg Cys Cys Leu Asn Gln Lys Gly Thr Ile Leu Gly Leu  
65 70 75

Asp Leu Gln Asn Cys Ser Leu Glu Asp Pro Gly Pro Asn Phe His  
80 85 90

Gln Ala His Thr Thr Val Ile Ile Asp Leu Gln Ala Asn Pro Leu  
95 100 105

Lys Gly Asp Leu Ala Asn Thr Phe Arg Gly Phe Thr Gln Leu Gln  
110 115 120

Thr Leu Ile Leu Pro Gln His Val Asn Cys Pro Gly Gly Ile Asn  
125 130 135

Ala Trp Asn Thr Ile Thr Ser Tyr Ile Asp Asn Gln Ile Cys Gln  
140 145 150

Gly Gln Lys Asn Leu Cys Asn Asn Thr Gly Asp Pro Glu Met Cys  
155 160 165

Pro Glu Asn Gly Ser Cys Val Pro Asp Gly Pro Gly Leu Leu Gln  
170 175 180

Cys Val Cys Ala Asp Gly Phe His Gly Tyr Lys Cys Met Arg Gln  
185 190 195

Gly Ser Phe Ser Leu Leu Met Phe Phe Gly Ile Leu Gly Ala Thr  
200 205 210

Thr Leu Ser Val Ser Ile Leu Leu Trp Ala Thr Gln Arg Arg Lys  
215 220 225

Ala Lys Thr Ser

<210> SEQ ID NO 9

<211> LENGTH: 2482

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 9

gggggagaag gcgccgagc cccagctctc cgagcaccgg gtcggaagcc 50

gcgaccgag ccgcgagga agctgggacc ggaacctcgg cggaccggc 100

cccccaac tcacctgcg agtccaccag caccctcgga acccagaggc 150

ccgctctg aagtgacc cctggggag gaaggcgatg gccctgcga 200

-continued

---

ggacgatggc ccgcgcccgc ctgcgcccg cggcatccc tgcgctgcc	250
ttgtggcttc tgtgcacgct cggcctccag ggcacccagg ccgggcccacc	300
gcccgcgccc cctgggctgc ccgcgggagc cgactgcctg aacagcttta	350
ccgcgggggt gcctggcttc gtgctggaca ccaacgcctc ggtcagcaac	400
ggagctacct tcctggagtc ccccaccgtg cgccggggct gggactgctg	450
gcgcgctcgc tgcaccacc agaactgcaa cttggcgcta gtggagctgc	500
agcccgaccg cggggaggac gccatcgccg cctgcttctt catcaactgc	550
ctctacgagc agaacttcgt gtgcaagttc gcgccaggg agggcttcat	600
caactacctc acgagggaag tgtaccgctc ctaccgcag ctgcggacc	650
agggctttgg agggctctgg atcccgaag cctgggcagg catagacttg	700
aaggtacaac ccaggaacc cctggtgctg aaggatgtgg aaaacacaga	750
ttggcgcta ctgcggggtg acacggatgt cagggtagag aggaaagacc	800
caaaccaggt ggaactgtgg ggactcaagg aaggcaccta cctgttccag	850
ctgacagtga ctagctcaga ccaccagag gacacggcca acgtcacagt	900
cactgtgctg tccaccaagc agacagaaga ctactgcctc gcatccaaca	950
aggtgggtcg ctgcggggc tctttccac gotggtacta tgaccccacg	1000
gagcagatct gcaagagttt cgtttatgga ggctgcttg gcaacaagaa	1050
caactacctt cgggaagaag agtgcattct agcctgtcgg ggtgtgcaag	1100
gtggcccttt gagaggcagc tctggggctc aggcgacttt ccccagggc	1150
ccctccatgg aaaggcgcca tccagtgtgc tctggcacct gtcagcccac	1200
ccagttccgc tgcagcaatg gctgctgcat cgacagtttc ctggagtgtg	1250
acgacacccc caactgcccc gacgcctccg acgaggctgc ctgtgaaaaa	1300
tacacgagtg gctttgacga gctccagcgc atccatttcc ccagtgacaa	1350
agggcaactgc gtggacctgc cagacacagg actctgcaag gagagcatcc	1400
cgcgctggtg ctacaacccc ttcagcgaac actgcgcccg ctttacctat	1450
ggtggttgtt atggcaacaa gaacaacttt gaggaagagc agcagtgctc	1500
cgagtcttgt cgcgcatct ccaagaagga tgtgtttggc ctgaggcggg	1550
aaatcccctt tcccagcaca ggctctgtgg agatggctgt cacagtgttc	1600
ctggctatct gcatttgtgt ggtggtagcc atcttgggtt actgcttctt	1650
caagaaccag agaaagact tccacggaca ccaccaccac ccaccacca	1700
cccctgccag ctccactgtc tccactaccg aggacacgga gcacctggtc	1750
tataaccaca ccacccggcc cctctgagcc tgggtctcac cggctctcac	1800
ctggccctgc ttcctgcttg ccaaggcaga ggctgggct gggaaaaact	1850
ttggaaccag actcttgctt gtttcccagg cccactgtgc ctgagagacc	1900
agggctccag ccctcttgg agaagtctca gctaagctca cgtcctgaga	1950
aagctcaaag gtttgaagg agcagaaaac ccttgggcca gaagtaccag	2000
actagatgga cctgcctgca taggagtttg gaggaagttg gagttttgtt	2050
tcctctgttc aaagctgctt gtocctaccc catggtgcta ggaagaggag	2100

-continued

---

tggggtggtg tcagaccctg gaggcccaaa cctgtcctc ccgagctcct	2150
cttccatgct gtgcgccag ggctgggagg aaggacttcc ctgtgtagtt	2200
tgtgctgtaa agagttgctt tttgtttatt taatgctgtg gcatgggtga	2250
agaggagggg aagaggcctg tttggcctct ctgtcctctc ttctcttcc	2300
cccaagattg agctctctgc ccttgatcag ccccaccctg gcctagacca	2350
gcagacagag ccaggagagg ctacagctgca ttccgcagcc cccaccccca	2400
aggttctcca acatcacagc ccagcccacc cactgggtaa taaaagtgtt	2450
ttgtggaaaa aaaaaaaaaa aaaaaaaaaa aa	2482

&lt;210&gt; SEQ ID NO 10

&lt;211&gt; LENGTH: 529

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 10

Met Ala Pro Ala Arg Thr Met Ala Arg Ala Arg Leu Ala Pro Ala	1	5	10	15
Gly Ile Pro Ala Val Ala Leu Trp Leu Leu Cys Thr Leu Gly Leu	20	25	30	
Gln Gly Thr Gln Ala Gly Pro Pro Pro Ala Pro Pro Gly Leu Pro	35	40	45	
Ala Gly Ala Asp Cys Leu Asn Ser Phe Thr Ala Gly Val Pro Gly	50	55	60	
Phe Val Leu Asp Thr Asn Ala Ser Val Ser Asn Gly Ala Thr Phe	65	70	75	
Leu Glu Ser Pro Thr Val Arg Arg Gly Trp Asp Cys Val Arg Ala	80	85	90	
Cys Cys Thr Thr Gln Asn Cys Asn Leu Ala Leu Val Glu Leu Gln	95	100	105	
Pro Asp Arg Gly Glu Asp Ala Ile Ala Ala Cys Phe Leu Ile Asn	110	115	120	
Cys Leu Tyr Glu Gln Asn Phe Val Cys Lys Phe Ala Pro Arg Glu	125	130	135	
Gly Phe Ile Asn Tyr Leu Thr Arg Glu Val Tyr Arg Ser Tyr Arg	140	145	150	
Gln Leu Arg Thr Gln Gly Phe Gly Gly Ser Gly Ile Pro Lys Ala	155	160	165	
Trp Ala Gly Ile Asp Leu Lys Val Gln Pro Gln Glu Pro Leu Val	170	175	180	
Leu Lys Asp Val Glu Asn Thr Asp Trp Arg Leu Leu Arg Gly Asp	185	190	195	
Thr Asp Val Arg Val Glu Arg Lys Asp Pro Asn Gln Val Glu Leu	200	205	210	
Trp Gly Leu Lys Glu Gly Thr Tyr Leu Phe Gln Leu Thr Val Thr	215	220	225	
Ser Ser Asp His Pro Glu Asp Thr Ala Asn Val Thr Val Thr Val	230	235	240	
Leu Ser Thr Lys Gln Thr Glu Asp Tyr Cys Leu Ala Ser Asn Lys	245	250	255	
Val Gly Arg Cys Arg Gly Ser Phe Pro Arg Trp Tyr Tyr Asp Pro				

-continued

---

	260									265									270
Thr	Glu	Gln	Ile	Cys	Lys	Ser	Phe	Val	Tyr	Gly	Gly	Cys	Leu	Gly					
				275					280					285					
Asn	Lys	Asn	Asn	Tyr	Leu	Arg	Glu	Glu	Glu	Cys	Ile	Leu	Ala	Cys					
				290					295					300					
Arg	Gly	Val	Gln	Gly	Gly	Pro	Leu	Arg	Gly	Ser	Ser	Gly	Ala	Gln					
				305					310					315					
Ala	Thr	Phe	Pro	Gln	Gly	Pro	Ser	Met	Glu	Arg	Arg	His	Pro	Val					
				320					325					330					
Cys	Ser	Gly	Thr	Cys	Gln	Pro	Thr	Gln	Phe	Arg	Cys	Ser	Asn	Gly					
				335					340					345					
Cys	Cys	Ile	Asp	Ser	Phe	Leu	Glu	Cys	Asp	Asp	Thr	Pro	Asn	Cys					
				350					355					360					
Pro	Asp	Ala	Ser	Asp	Glu	Ala	Ala	Cys	Glu	Lys	Tyr	Thr	Ser	Gly					
				365					370					375					
Phe	Asp	Glu	Leu	Gln	Arg	Ile	His	Phe	Pro	Ser	Asp	Lys	Gly	His					
				380					385					390					
Cys	Val	Asp	Leu	Pro	Asp	Thr	Gly	Leu	Cys	Lys	Glu	Ser	Ile	Pro					
				395					400					405					
Arg	Trp	Tyr	Tyr	Asn	Pro	Phe	Ser	Glu	His	Cys	Ala	Arg	Phe	Thr					
				410					415					420					
Tyr	Gly	Gly	Cys	Tyr	Gly	Asn	Lys	Asn	Asn	Phe	Glu	Glu	Glu	Gln					
				425					430					435					
Gln	Cys	Leu	Glu	Ser	Cys	Arg	Gly	Ile	Ser	Lys	Lys	Asp	Val	Phe					
				440					445					450					
Gly	Leu	Arg	Arg	Glu	Ile	Pro	Ile	Pro	Ser	Thr	Gly	Ser	Val	Glu					
				455					460					465					
Met	Ala	Val	Thr	Val	Phe	Leu	Val	Ile	Cys	Ile	Val	Val	Val	Val					
				470					475					480					
Ala	Ile	Leu	Gly	Tyr	Cys	Phe	Phe	Lys	Asn	Gln	Arg	Lys	Asp	Phe					
				485					490					495					
His	Gly	His	His	His	His	Pro	Pro	Pro	Thr	Pro	Ala	Ser	Ser	Thr					
				500					505					510					
Val	Ser	Thr	Thr	Glu	Asp	Thr	Glu	His	Leu	Val	Tyr	Asn	His	Thr					
				515					520					525					

Thr Arg Pro Leu

<210> SEQ ID NO 11  
 <211> LENGTH: 1899  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 11

gtgctgggct ttttcagaca agtgcacatc ctaaccaggt cacatttcag	50
ccgcgaccca ctctccgcca gtcaccggag gcagaccgcg ggaggagagc	100
tgaggacagc cgcgtgcgct tcgccagcag cggggtgggg ggaaggacat	150
taaaatactg cagaagtcaa gaccccccca ggtcgaacc agaccacgat	200
gcgcgccccc ggctgcgggc ggctggtgct gccgctgctg ctctggccc	250
cggcagccct ggccgaaggc gacgccaagg ggctcaagga gggcgagacc	300
cccggcaatt tcatggagga cgagcaatgg ctgtcgtcca tctcgcagta	350

-continued

---

```

cagcggcaag atcaagcact ggaaccgctt ccgagacgaa gtggaggatg      400
actatatcaa gagctgggag gacaatcagc aaggagatga agccctggat      450
accaccaagg acccctgcca gaagggtgaag tgcagccgcc acaagggtgtg      500
cattgccagc ggctaccagc gggccatgtg catcagtcgc aagaagctgg      550
agcacaggat caagcagccg accgtgaaac tccatggaaa caaagactcc      600
atctgcaagc cctgccacat ggcccagctt gcctctgtct gcggctcaga      650
tggccacact tacagctctg tgtgtaagct ggagcaacag gcgtgcctga      700
gcagcaagca gctggcgggt cgatgcgagg gccctgccc ctgcccacg      750
gagcaggctg ccacctccac cgccgatggc aaaccagaga cttgcaccgg      800
tcaggacctg gctgacctgg gagatcggct gcgggactgg ttccagctcc      850
ttcatgagaa ctccaagcag aatggctcag ccagcagtgt agcgggccc      900
gccagcgggc tggacaagag cctggggggc agctgcaagg actccattgg      950
ctggatgttc tccaagctgg acaccagtgc tgacctcttc ctggaccaga     1000
cggagctggc cgccatcaac ctggacaagt acgaggtctg catccgtccc     1050
ttcttcaact cctgtgacac ctacaaggat ggccgggtct ctactgctga     1100
gtggtgcttc tgcttctgga gggagaagcc cccctgcctg gcagagctgg     1150
agcgcatcca gatccaggag gcgcacaaga agaagccagg catcttcatc     1200
ccgagctcgc acgaggatgg ctactaccgg aagatgcagt gtgaccagag     1250
cagcgggtgac tgctggcgtg tggaccagct gggcctggag ctgactggca     1300
cgcgcacgca tgggagcccc gactgcgatg acatcgtggg cttctcgggg     1350
gactttgtaa gcggtgtcgg ctgggaggat gaggaggaga aggagacgga     1400
ggaaagcagg gaggaggccc aggaggagga gggcgaggca ggcgaggctg     1450
acgacggggg ctacatctgg tagacgccct caggagccgg ctgccggggg     1500
ggactcaaca gcagagctct gagcagcagc aggcaacttc gagaacggat     1550
ccagaaatgc agtcagaagg accctgctcc acctgggggg actgggagtg     1600
tgagtgtgca tggcatgtgt gtggcacaga tggctgggac gggtgacagt     1650
gtgagtgcac gtgtgcatgc atgtgtgtat gtgtgtgtgt gtgtggcatg     1700
cgctgacaaa tgtgtccttg atccacactg ctctggcagc agtgagtac     1750
ccaaaggccc ctcggcctc ctgtagctg ttttcttcc ttttgttgtt     1800
ggttttaaaa tacattcaca cacaaataca aaaaaaaaaa aaaaaaaaaa     1850
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     1899

```

&lt;210&gt; SEQ ID NO 12

&lt;211&gt; LENGTH: 424

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 12

```

Met Arg Ala Pro Gly Cys Gly Arg Leu Val Leu Pro Leu Leu Leu
 1             5             10             15

```

```

Leu Ala Ala Ala Ala Leu Ala Glu Gly Asp Ala Lys Gly Leu Lys
 20             25             30

```

-continued

---

Glu Gly Glu Thr Pro Gly Asn Phe Met Glu Asp Glu Gln Trp Leu  
 35 40 45  
 Ser Ser Ile Ser Gln Tyr Ser Gly Lys Ile Lys His Trp Asn Arg  
 50 55 60  
 Phe Arg Asp Glu Val Glu Asp Asp Tyr Ile Lys Ser Trp Glu Asp  
 65 70 75  
 Asn Gln Gln Gly Asp Glu Ala Leu Asp Thr Thr Lys Asp Pro Cys  
 80 85 90  
 Gln Lys Val Lys Cys Ser Arg His Lys Val Cys Ile Ala Gln Gly  
 95 100 105  
 Tyr Gln Arg Ala Met Cys Ile Ser Arg Lys Lys Leu Glu His Arg  
 110 115 120  
 Ile Lys Gln Pro Thr Val Lys Leu His Gly Asn Lys Asp Ser Ile  
 125 130 135  
 Cys Lys Pro Cys His Met Ala Gln Leu Ala Ser Val Cys Gly Ser  
 140 145 150  
 Asp Gly His Thr Tyr Ser Ser Val Cys Lys Leu Glu Gln Gln Ala  
 155 160 165  
 Cys Leu Ser Ser Lys Gln Leu Ala Val Arg Cys Glu Gly Pro Cys  
 170 175 180  
 Pro Cys Pro Thr Glu Gln Ala Ala Thr Ser Thr Ala Asp Gly Lys  
 185 190 195  
 Pro Glu Thr Cys Thr Gly Gln Asp Leu Ala Asp Leu Gly Asp Arg  
 200 205 210  
 Leu Arg Asp Trp Phe Gln Leu Leu His Glu Asn Ser Lys Gln Asn  
 215 220 225  
 Gly Ser Ala Ser Ser Val Ala Gly Pro Ala Ser Gly Leu Asp Lys  
 230 235 240  
 Ser Leu Gly Ala Ser Cys Lys Asp Ser Ile Gly Trp Met Phe Ser  
 245 250 255  
 Lys Leu Asp Thr Ser Ala Asp Leu Phe Leu Asp Gln Thr Glu Leu  
 260 265 270  
 Ala Ala Ile Asn Leu Asp Lys Tyr Glu Val Cys Ile Arg Pro Phe  
 275 280 285  
 Phe Asn Ser Cys Asp Thr Tyr Lys Asp Gly Arg Val Ser Thr Ala  
 290 295 300  
 Glu Trp Cys Phe Cys Phe Trp Arg Glu Lys Pro Pro Cys Leu Ala  
 305 310 315  
 Glu Leu Glu Arg Ile Gln Ile Gln Glu Ala Ala Lys Lys Lys Pro  
 320 325 330  
 Gly Ile Phe Ile Pro Ser Cys Asp Glu Asp Gly Tyr Tyr Arg Lys  
 335 340 345  
 Met Gln Cys Asp Gln Ser Ser Gly Asp Cys Trp Arg Val Asp Gln  
 350 355 360  
 Leu Gly Leu Glu Leu Thr Gly Thr Arg Thr His Gly Ser Pro Asp  
 365 370 375  
 Cys Asp Asp Ile Val Gly Phe Ser Gly Asp Phe Gly Ser Gly Val  
 380 385 390  
 Gly Trp Glu Asp Glu Glu Glu Lys Glu Thr Glu Glu Ala Gly Glu  
 395 400 405

-continued

---

Glu Ala Glu Glu Glu Glu Gly Glu Ala Gly Glu Ala Asp Asp Gly  
410 415 420

Gly Tyr Ile Trp

<210> SEQ ID NO 13  
<211> LENGTH: 2680  
<212> TYPE: DNA  
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 13

tgccggcgacc gtcgtacacc atgggcctcc acctccgccc ctaccgtgtg	50
gggctgctcc cggatggcct cctgttcctc ttgctgctgc taatgctgct	100
cgccggaccga gcgctcccgg ccggacgtca cccccagtg gtgctggctc	150
ctggtgattt gggtaaccaa ctggaagcca agctggacaa gccgacagt	200
gtgcactaac tctgctccaa gaagaccgaa agctacttca caatctggct	250
gaacctggaa ctgctgctgc ctgtcatcat tgactgctgg attgacaata	300
tcaggctggt ttacaacaaa acatccaggg ccaccocagtt tcctgatggt	350
gtggatgtac gtgtccctgg ctttgggaag accttctcac tggagttcct	400
ggaccccagc aaaagcagcg tgggttccta tttccacacc atggtgagga	450
gccttgtggg ctggggctac acacgggggt aggatgtccg aggggctccc	500
tatgactggc gccgagcccc aatatgaaaac gggcctact tcctggcctc	550
ccgcgagatg atcgaggaga tgtaccagct gtatgggggc cccgtgggtc	600
tggttgccca cagtatgggc aacatgtaca cgctctactt tctgcagcgg	650
cagcccgagg cctggaagga caagtatac cgggccttcg tgtcactggg	700
tgcgccctgg gggggcgtgg ccaagaccct gcgcgtcctg gcttcaggag	750
acaacaaccg gatcccagtc atcgggcccc tgaagatccg ggagcagcag	800
cggtcagctg tctccaccag ctggctgctg ccctacaact acacatggtc	850
acctgagaag gtgttcgtgc agacaccac aatcaactac acactgcggg	900
actaccgcaa gttcttccag gacatcggct ttgaagatgg ctggctcatg	950
cgccagcaga cagaagggct ggtggaagcc acgatgccac ctggcgtgca	1000
gctgcactgc ctctatggta ctggcgtccc cacaccagac tccttctact	1050
atgagagctt ccctgaccgt gaccctaaaa tctgctttgg tgacggcgat	1100
ggtactgtga acttgaagag tgcctgcag tgccaggcct ggcagagccg	1150
ccaggagcac caagtgttc tgcaggagct gccaggcagc gagcacatcg	1200
agatgctggc caaccgcc accctggcct atctgaaacg tgtgctcctt	1250
gggcctgac tcctgtgcca caggactcct gtggctcggc cgtggacctg	1300
ctgttggcct ctggggctgt catggcccac gcgttttgca aagtttgtga	1350
ctcaccattc aaggccccga gtcttgact gtgaagcatc tgccatgggg	1400
aagtgctggt tgttatacct tctctgtggc agtgaagaag gaagaaatga	1450
gagtctagac tcaagggaca ctggatggca agaagctgc tgatggtgga	1500
actgctgtga ccttaggact ggctccacag ggtggactgg ctgggcctg	1550
gtcccagctc ctgcctgggg ccattgtgct ccctattcct gtgggctttt	1600

-continued

```

catacttgcc tactgggccc tggccccgca gccttcctat gagggatggt      1650
actgggctgt ggtcctgtac ccagaggtcc cagggatcgg ctccctggccc      1700
ctcgggtgac ccttcccaca caccagccac agataggcct gccactggtc      1750
atgggtagct agagctgctg gcttcctctg ggcttagctg gtggccagcc      1800
tgactggctt cctgggctgag cctagtagct cctgcaggca ggggcagttt      1850
gttgcgttct tcgtggttcc caggccctgg gacatctcac tccactccta      1900
cctcccttac caccaggagc attcaagctc tggattgggc agcagatgtg      1950
ccccagctcc cgcaggctgt gttccagggg ccttgatttc ctgggatgtg      2000
ctattggccc caggactgaa gctgcctccc ttcaccctgg gactgtggtt      2050
ccaaggtaga gagcaggggt tggagccatg gccttctggg aacctatgga      2100
gaaagggaat ccaaggaagc agccaaggct gctcgcagct tccctgagct      2150
gcacctcttg ctaacccccc catcacactg ccaccctgcc ctagggtctc      2200
actagtacca agtgggtcag cacagggctg aggatggggc tcctatccac      2250
cctggccagc acccagctta gtgctgggac tagcccagaa acttgaatgg      2300
gaccctgaga gagccagggg tcccctgagg ccccctagg ggctttctgt      2350
ctgccccagg gtgctccatg gatctccctg tggcagcagg catggagagt      2400
cagggtctgc ttcattggcag taggtctctaa gtgggtgact ggccacaggc      2450
cgagaaaagg gtacagcctc taggtggggt tcccгаагac gccttcaggc      2500
tggactgagc tgctctccca cagggtttct gtgcagctgg attttctctg      2550
ttgcatacat gcctggcatc tgtctcccct tgttctgag tggcccaca      2600
tggggctctg agcaggctgt atctggattc tggcaataaa agtactctgg      2650
atgctgtaaa aaaaaaaaaa aaaaaaaaaa      2680
    
```

```

<210> SEQ ID NO 14
<211> LENGTH: 412
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien
    
```

<400> SEQUENCE: 14

```

Met Gly Leu His Leu Arg Pro Tyr Arg Val Gly Leu Leu Pro Asp
 1          5          10          15
Gly Leu Leu Phe Leu Leu Leu Leu Leu Met Leu Leu Ala Asp Pro
          20          25          30
Ala Leu Pro Ala Gly Arg His Pro Pro Val Val Leu Val Pro Gly
          35          40          45
Asp Leu Gly Asn Gln Leu Glu Ala Lys Leu Asp Lys Pro Thr Val
          50          55          60
Val His Tyr Leu Cys Ser Lys Lys Thr Glu Ser Tyr Phe Thr Ile
          65          70          75
Trp Leu Asn Leu Glu Leu Leu Leu Pro Val Ile Ile Asp Cys Trp
          80          85          90
Ile Asp Asn Ile Arg Leu Val Tyr Asn Lys Thr Ser Arg Ala Thr
          95          100          105
Gln Phe Pro Asp Gly Val Asp Val Arg Val Pro Gly Phe Gly Lys
          110          115          120
    
```



-continued

---

Thr Phe Ser Leu Glu Phe Leu Asp Pro Ser Lys Ser Ser Val Gly  
 125 130 135

Ser Tyr Phe His Thr Met Val Glu Ser Leu Val Gly Trp Gly Tyr  
 140 145 150

Thr Arg Gly Glu Asp Val Arg Gly Ala Pro Tyr Asp Trp Arg Arg  
 155 160 165

Ala Pro Asn Glu Asn Gly Pro Tyr Phe Leu Ala Leu Arg Glu Met  
 170 175 180

Ile Glu Glu Met Tyr Gln Leu Tyr Gly Gly Pro Val Val Leu Val  
 185 190 195

Ala His Ser Met Gly Asn Met Tyr Thr Leu Tyr Phe Leu Gln Arg  
 200 205 210

Gln Pro Gln Ala Trp Lys Asp Lys Tyr Ile Arg Ala Phe Val Ser  
 215 220 225

Leu Gly Ala Pro Trp Gly Gly Val Ala Lys Thr Leu Arg Val Leu  
 230 235 240

Ala Ser Gly Asp Asn Asn Arg Ile Pro Val Ile Gly Pro Leu Lys  
 245 250 255

Ile Arg Glu Gln Gln Arg Ser Ala Val Ser Thr Ser Trp Leu Leu  
 260 265 270

Pro Tyr Asn Tyr Thr Trp Ser Pro Glu Lys Val Phe Val Gln Thr  
 275 280 285

Pro Thr Ile Asn Tyr Thr Leu Arg Asp Tyr Arg Lys Phe Phe Gln  
 290 295 300

Asp Ile Gly Phe Glu Asp Gly Trp Leu Met Arg Gln Asp Thr Glu  
 305 310 315

Gly Leu Val Glu Ala Thr Met Pro Pro Gly Val Gln Leu His Cys  
 320 325 330

Leu Tyr Gly Thr Gly Val Pro Thr Pro Asp Ser Phe Tyr Tyr Glu  
 335 340 345

Ser Phe Pro Asp Arg Asp Pro Lys Ile Cys Phe Gly Asp Gly Asp  
 350 355 360

Gly Thr Val Asn Leu Lys Ser Ala Leu Gln Cys Gln Ala Trp Gln  
 365 370 375

Ser Arg Gln Glu His Gln Val Leu Leu Gln Glu Leu Pro Gly Ser  
 380 385 390

Glu His Ile Glu Met Leu Ala Asn Ala Thr Thr Leu Ala Tyr Leu  
 395 400 405

Lys Arg Val Leu Leu Gly Pro  
 410

<210> SEQ ID NO 15  
 <211> LENGTH: 1371  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 15

cagagcagat aatggcaagc atggctgccc tgctcacctg ggctctggct	50
cttctttcag cgttttcggc caccaggca cggaaaggct tctgggacta	100
cttcagccag accagcgggg acaaaggcag ggtggagcag atccatcagc	150
agaagatggc tcgagagccc gcgacctga aagacagcct tgagcaagac	200

-continued

```

ctcaacaata tgaacaagtt cctgaaaaag ctgaggcctc tgagtgggag      250
cgaggctcct cggctcccac aggaccgggt gggcatgcgg cggcagctgc      300
aggaggagtt ggaggagggt aaggctcgcc tccagcccta catggcagag      350
gcgcacgagc tggtagggctg gaatttggag ggcttgccgc agcaactgaa      400
gccctacacg atggatctga tggagcaggt ggcctgcgcg gtgcaggagc      450
tgcaggagca gttgcgcgtg gtgggggaag acaccaaggc ccagttgctg      500
gggggctggt acgaggcttg ggctttgctg cagggactgc agagccgctg      550
gggtgaccac accggccgct tcaaagagct cttccacca tacgccgaga      600
gcctggtgag cggcatcggg cgcacgtgc aggagctgca ccgcagtgtg      650
gtcccgcaag ccccgccag ccccgccgc ctcagtcgct gcgtgcaggt      700
gctctcccgg aagctcacgc tcaaggccaa ggcctgcac gcaogcatcc      750
agcagaacct ggaccagctg cgcgaagagc tcagcagagc ctttgaggc      800
actgggactg aggaaggggc cggcccggac ccctagatgc tctccgagga      850
gggtgcgccg cgacttcagg ctttccgcca ggacacctac ctgcagatag      900
ctgccttacc tcgcgccatc gaccaggaga ctgaggaggt ccagcagcag      950
ctggcgccac ctccaccagg ccacagtgcc ttcgcccag agtttcaaca     1000
aacagacagt ggcaaggttc tgagcaagct gcaggcccgt ctggatgacc     1050
tgtgggaaga catcactcac agccttcacg accagggcca cagccatctg     1100
ggggaccctt gaggatctac ctgcccaggc ccattcccag cttcttgtct     1150
ggggagcctt ggctctgagc ctctagcatg gttcagtcct tgaagtggc     1200
ctgttgggtg gagggtgaa ggtcctgtgc aggacagga ggccacaaa     1250
ggggctgctg tctctgcat atccagcctc ctgcgactcc ccaatctgga     1300
tgcattacat tcaccaggct ttgcaaaaaa aaaaaaaaaa aaaaaaaaaa     1350
aaaaaaaaaa aaaaaaaaaa a                                     1371
    
```

```

<210> SEQ ID NO 16
<211> LENGTH: 274
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien
    
```

<400> SEQUENCE: 16

```

Met Ala Ser Met Ala Ala Val Leu Thr Trp Ala Leu Ala Leu Leu
 1           5           10           15
Ser Ala Phe Ser Ala Thr Gln Ala Arg Lys Gly Phe Trp Asp Tyr
 20          25          30
Phe Ser Gln Thr Ser Gly Asp Lys Gly Arg Val Glu Gln Ile His
 35          40          45
Gln Gln Lys Met Ala Arg Glu Pro Ala Thr Leu Lys Asp Ser Leu
 50          55          60
Glu Gln Asp Leu Asn Asn Met Asn Lys Phe Leu Glu Lys Leu Arg
 65          70          75
Pro Leu Ser Gly Ser Glu Ala Pro Arg Leu Pro Gln Asp Pro Val
 80          85          90
Gly Met Arg Arg Gln Leu Gln Glu Glu Leu Glu Glu Val Lys Ala
    
```

-continued

													95				100				105		
Arg	Leu	Gln	Pro	Tyr	Met	Ala	Glu	Ala	His	Glu	Leu	Val	Gly	Trp	110				115				120
Asn	Leu	Glu	Gly	Leu	Arg	Gln	Gln	Leu	Lys	Pro	Tyr	Thr	Met	Asp	125				130				135
Leu	Met	Glu	Gln	Val	Ala	Leu	Arg	Val	Gln	Glu	Leu	Gln	Glu	Gln	140				145				150
Leu	Arg	Val	Val	Gly	Glu	Asp	Thr	Lys	Ala	Gln	Leu	Leu	Gly	Gly	155				160				165
Val	Asp	Glu	Ala	Trp	Ala	Leu	Leu	Gln	Gly	Leu	Gln	Ser	Arg	Val	170				175				180
Val	His	His	Thr	Gly	Arg	Phe	Lys	Glu	Leu	Phe	His	Pro	Tyr	Ala	185				190				195
Glu	Ser	Leu	Val	Ser	Gly	Ile	Gly	Arg	His	Val	Gln	Glu	Leu	His	200				205				210
Arg	Ser	Val	Ala	Pro	His	Ala	Pro	Ala	Ser	Pro	Ala	Arg	Leu	Ser	215				220				225
Arg	Cys	Val	Gln	Val	Leu	Ser	Arg	Lys	Leu	Thr	Leu	Lys	Ala	Lys	230				235				240
Ala	Leu	His	Ala	Arg	Ile	Gln	Gln	Asn	Leu	Asp	Gln	Leu	Arg	Glu	245				250				255
Glu	Leu	Ser	Arg	Ala	Phe	Ala	Gly	Thr	Gly	Thr	Glu	Glu	Gly	Ala	260				265				270
Gly Pro Asp Pro																							

<210> SEQ ID NO 17  
 <211> LENGTH: 2854  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 17

ctaagaggac aagatgaggc cgggcctctc atttctocta gccottctgt	50
tcttctctgg ccaagctgca ggggatttgg gggatgtggg acctccaatt	100
cccagccccg gcttcagctc tttcccaggt gttgactcca gctccagctt	150
cagctccagc tccaggtcgg gctccagctc cagccgcagc ttaggcagcg	200
gaggttctgt gtcccagttg ttttccaatt tcaccggctc cgtggatgac	250
cgtgggacct gccagtgtc tgtttccctg ccagacacca cctttcccgt	300
ggacagagtg gaacgcttgg aattcacagc tcatgttctt tctcagaagt	350
ttgagaaaga actttctaaa gtgagggaa atgtccaatt aattagtgtg	400
tatgaaaaga aactgttaaa cctaactgtc cgaattgaca tcatggagaa	450
ggataccatt tcttacctg aactggactt cgagctgac aaggtagaag	500
tgaaggagat ggaaaaactg gtcatacagc tgaaggagag ttttggtgga	550
agctcagaaa ttgttgacca gctggaggty gagataagaa atatgactct	600
cttggtagag aagcttgaga cactagacaa aaacaatgtc cttgccattc	650
gccgagaaat cgtggctctg aagaccaagc tgaagagty tgaggcctct	700
aaagatcaaa acaccctgt cgtccacct cctcccactc cagggagctg	750
tggctcatggt ggtgtgtgta acatcagcaa accgtctgtg gttcagctca	800

-continued

---

actggagagg gttttcttat ctatatggtg cttggggtag ggattactct	850
ccccagcatc caaaciaaagg actgtattgg gtggcgccat tgaatacaga	900
tgggagactg ttggagtatt atagactgta caacacactg gatgatttgc	950
tattgtatat aaatgctcga gagttgcgga tcacctatgg ccaaggtagt	1000
ggtacagcag tttacaacaa caacatgtac gtcaacatgt acaacaccgg	1050
gaatattgcc agagttaacc tgaccaccaa cacgattgct gtgactcaaa	1100
ctctccctaa tgctgctat aataaccgct tttcatatgc taatgttgct	1150
tggcaagata ttgactttgc tgtggatgag aatggattgt gggttattta	1200
ttcaactgaa gccagcactg gtaacatggt gattagtaaa ctcaatgaca	1250
ccacacttca ggtgctaaac acttgggtata ccaagcagta taaacctct	1300
gcttctaacg ccttcatggt atgtgggggt ctgtatgcc cccgtactat	1350
gaacaccaga acagaagaga tttttacta ttatgacaca aacacaggga	1400
aagagggcaa actagacatt gtaatgcata agatgcagga aaaagtgcag	1450
agcattaact ataacccttt tgaccagaaa ctttatgtct ataacgatgg	1500
ttacctctg aattatgac tttctgtctt gcagaagccc cagtaagctg	1550
tttaggagtt agggtgaaag agaaaatggt tgttgaaaa atagtcttct	1600
ccacttactt agatatctgc aggggtgtct aaaagtgtgt tcattttgca	1650
gcaatgttta ggtgcatagt tctaccacac tagagatcta ggacatttgt	1700
cttgatttgg tgagttctct tgggaatcat ctgcctcttc aggcgcattt	1750
tgcaataaag tctgtctagg gtgggattgt cagaggctca ggggcactgt	1800
gggcctatg aagcctactg tgaggaggct tcactagaag ccttaaatta	1850
ggaattaagg aacttaaac tcagtatggc gtctagggat tctttgtaca	1900
ggaaatattg cccaatgact agtcctcatc catgtagcac cactaattct	1950
tccatgcctg gaagaaacct ggggacttag ttaggtagat taatatctgg	2000
agctcctcga gggaccaaat ctccaacttt tttttcccct cactagcacc	2050
tggaatgatg ctttgtatgt ggcagataag taaatttggc atgottatat	2100
attctacatc tgtaaaagtgc tgagttttat ggagagaggc ctttttatgc	2150
attaaattgt acatggcaaa taaatcccag aaggatctgt agatgaggca	2200
cctgcttttt cttttctctc attgtccacc ttactaaaag tcagtagaat	2250
cttctacctc ataacttctc tccaaaggca gctcagaaga ttagaaccag	2300
acttactaac caattccacc ccccaccaac ccccttctac tgcctacttt	2350
aaaaaaatta atagttttct atggaactga tctaagatta gaaaaattaa	2400
ttttctttaa tttcattatg gacttttatt tacatgactc taagactata	2450
agaaaaactg atggcagtga caaagtgcta gcatttattg ttatctaata	2500
aagaccttgg agcatatgtg caacttatga gtgtatcagt tgttgcattg	2550
aatttttgcc tttgtttaag cctggaactt gtaagaaaat gaaaatttaa	2600
tttttttttc taggacgagc tatagaaaag ctattgagag tatctagtta	2650
atcagtgcag tagttggaaa ccttgcctgt gtatgtgatg tgcctctgtg	2700

-continued

---

```

cttttgaatg actttatcat ctagtctttg tctatctttc ctttgatgtt      2750
caagtcctag tctatagatg tggcagttta aatgctttac tccccctttt      2800
aaaataaatg attaaaatgt gctttgaaaa aaaaaaaaaa aaaaaaaaaa      2850
aaaa                                                              2854

```

```

<210> SEQ ID NO 18
<211> LENGTH: 510
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

```

```

<400> SEQUENCE: 18

```

```

Met Arg Pro Gly Leu Ser Phe Leu Leu Ala Leu Leu Phe Phe Leu
 1          5          10          15
Gly Gln Ala Ala Gly Asp Leu Gly Asp Val Gly Pro Pro Ile Pro
 20          25          30
Ser Pro Gly Phe Ser Ser Phe Pro Gly Val Asp Ser Ser Ser Ser
 35          40          45
Phe Ser Ser Ser Ser Arg Ser Gly Ser Ser Ser Ser Arg Ser Leu
 50          55          60
Gly Ser Gly Gly Ser Val Ser Gln Leu Phe Ser Asn Phe Thr Gly
 65          70          75
Ser Val Asp Asp Arg Gly Thr Cys Gln Cys Ser Val Ser Leu Pro
 80          85          90
Asp Thr Thr Phe Pro Val Asp Arg Val Glu Arg Leu Glu Phe Thr
 95          100         105
Ala His Val Leu Ser Gln Lys Phe Glu Lys Glu Leu Ser Lys Val
 110         115         120
Arg Glu Tyr Val Gln Leu Ile Ser Val Tyr Glu Lys Lys Leu Leu
 125         130         135
Asn Leu Thr Val Arg Ile Asp Ile Met Glu Lys Asp Thr Ile Ser
 140         145         150
Tyr Thr Glu Leu Asp Phe Glu Leu Ile Lys Val Glu Val Lys Glu
 155         160         165
Met Glu Lys Leu Val Ile Gln Leu Lys Glu Ser Phe Gly Gly Ser
 170         175         180
Ser Glu Ile Val Asp Gln Leu Glu Val Glu Ile Arg Asn Met Thr
 185         190         195
Leu Leu Val Glu Lys Leu Glu Thr Leu Asp Lys Asn Asn Val Leu
 200         205         210
Ala Ile Arg Arg Glu Ile Val Ala Leu Lys Thr Lys Leu Lys Glu
 215         220         225
Cys Glu Ala Ser Lys Asp Gln Asn Thr Pro Val Val His Pro Pro
 230         235         240
Pro Thr Pro Gly Ser Cys Gly His Gly Gly Val Val Asn Ile Ser
 245         250         255
Lys Pro Ser Val Val Gln Leu Asn Trp Arg Gly Phe Ser Tyr Leu
 260         265         270
Tyr Gly Ala Trp Gly Arg Asp Tyr Ser Pro Gln His Pro Asn Lys
 275         280         285
Gly Leu Tyr Trp Val Ala Pro Leu Asn Thr Asp Gly Arg Leu Leu
 290         295         300

```

-continued

---

Glu Tyr Tyr Arg Leu Tyr Asn Thr Leu Asp Asp Leu Leu Leu Tyr  
 305 310 315

Ile Asn Ala Arg Glu Leu Arg Ile Thr Tyr Gly Gln Gly Ser Gly  
 320 325 330

Thr Ala Val Tyr Asn Asn Asn Met Tyr Val Asn Met Tyr Asn Thr  
 335 340 345

Gly Asn Ile Ala Arg Val Asn Leu Thr Thr Asn Thr Ile Ala Val  
 350 355 360

Thr Gln Thr Leu Pro Asn Ala Ala Tyr Asn Asn Arg Phe Ser Tyr  
 365 370 375

Ala Asn Val Ala Trp Gln Asp Ile Asp Phe Ala Val Asp Glu Asn  
 380 385 390

Gly Leu Trp Val Ile Tyr Ser Thr Glu Ala Ser Thr Gly Asn Met  
 395 400 405

Val Ile Ser Lys Leu Asn Asp Thr Thr Leu Gln Val Leu Asn Thr  
 410 415 420

Trp Tyr Thr Lys Gln Tyr Lys Pro Ser Ala Ser Asn Ala Phe Met  
 425 430 435

Val Cys Gly Val Leu Tyr Ala Thr Arg Thr Met Asn Thr Arg Thr  
 440 445 450

Glu Glu Ile Phe Tyr Tyr Tyr Asp Thr Asn Thr Gly Lys Glu Gly  
 455 460 465

Lys Leu Asp Ile Val Met His Lys Met Gln Glu Lys Val Gln Ser  
 470 475 480

Ile Asn Tyr Asn Pro Phe Asp Gln Lys Leu Tyr Val Tyr Asn Asp  
 485 490 495

Gly Tyr Leu Leu Asn Tyr Asp Leu Ser Val Leu Gln Lys Pro Gln  
 500 505 510

<210> SEQ ID NO 19  
 <211> LENGTH: 663  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 19

```

gcaccgcaga cggcgcggat cgcagggagc cggtcgcccg ccggaacggg           50
agcctgggtg tgctgtgga gtccggactc gtgggagacg atcgcgatga           100
acacggtgct gtcgcgggcg aactcactgt tcgccttctc gctgagcgtg           150
atggcggcgc tcaccttcg ctgcttcac accaccgctt tcaaagacag           200
gagcgtcccg gtgcggctgc acgtctcgcg gatcatgcta aaaaatgtag           250
aagatttcac tggacctaga gaaagaagtg atctgggatt tatcacattt           300
gatataactg ctgatctaga gaatatattt gattggaatg ttaagcagtt           350
gtttctttat ttatcagcag aatattcaac aaaaaataat gctctgaacc           400
aagttgtcct atgggacaag attgttttga gaggtgataa tccgaagctg           450
ctgctgaaa atagaaaac aaaatatttt ttctttgacg atggaaatgg           500
tctcaaggga aacaggaatg tacttttgac cctgtcttgg aacgtcgtac           550
caaatgctgg aattctacct ctgtgacag gatcaggaca cgtatctgtc           600
ccatttccag atacatatga aatacgaag agttattaaa ttattctgaa           650
    
```

-continued

tttgaacaaa aaa

663

<210> SEQ ID NO 20  
 <211> LENGTH: 180  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 20

Met	Asn	Thr	Val	Leu	Ser	Arg	Ala	Asn	Ser	Leu	Phe	Ala	Phe	Ser
1				5					10					15
Leu	Ser	Val	Met	Ala	Ala	Leu	Thr	Phe	Gly	Cys	Phe	Ile	Thr	Thr
			20						25					30
Ala	Phe	Lys	Asp	Arg	Ser	Val	Pro	Val	Arg	Leu	His	Val	Ser	Arg
			35						40					45
Ile	Met	Leu	Lys	Asn	Val	Glu	Asp	Phe	Thr	Gly	Pro	Arg	Glu	Arg
			50						55					60
Ser	Asp	Leu	Gly	Phe	Ile	Thr	Phe	Asp	Ile	Thr	Ala	Asp	Leu	Glu
			65						70					75
Asn	Ile	Phe	Asp	Trp	Asn	Val	Lys	Gln	Leu	Phe	Leu	Tyr	Leu	Ser
			80						85					90
Ala	Glu	Tyr	Ser	Thr	Lys	Asn	Asn	Ala	Leu	Asn	Gln	Val	Val	Leu
			95						100					105
Trp	Asp	Lys	Ile	Val	Leu	Arg	Gly	Asp	Asn	Pro	Lys	Leu	Leu	Leu
			110						115					120
Lys	Asp	Met	Lys	Thr	Lys	Tyr	Phe	Phe	Phe	Asp	Asp	Gly	Asn	Gly
			125						130					135
Leu	Lys	Gly	Asn	Arg	Asn	Val	Thr	Leu	Thr	Leu	Ser	Trp	Asn	Val
			140						145					150
Val	Pro	Asn	Ala	Gly	Ile	Leu	Pro	Leu	Val	Thr	Gly	Ser	Gly	His
			155						160					165
Val	Ser	Val	Pro	Phe	Pro	Asp	Thr	Tyr	Glu	Ile	Thr	Lys	Ser	Tyr
			170						175					180

<210> SEQ ID NO 21  
 <211> LENGTH: 415  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 21

aaacttgaag	ccatgaagat	cccggtcctt	cctgccgtgg	tgctcctctc	50
cctcctgggt	ctccactctg	cccagggagc	cacctgggt	ggtcctgagg	100
aagaaagcac	cattgagaat	tatgctcac	gaccgaggc	ctttaaacc	150
ccgttctga	acatgacaa	attgcgatct	ggtttaagg	ctgatgagtt	200
cctgaactgg	cacgccctct	ttgagtctat	caaaaggaaa	cttcctttcc	250
tcaactggga	tgcttttct	aagctgaaag	gactgaggag	cgcaactcct	300
gatgcccagt	gaccatgacc	tccactggaa	gaggggcta	gcgtgagcgc	350
tgattctcaa	cctaccataa	ctctttctctg	cctcaggaac	tccaataaaa	400
cattttccat	ccaaa				415

<210> SEQ ID NO 22  
 <211> LENGTH: 99

-continued

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 22

```

Met Lys Ile Pro Val Leu Pro Ala Val Val Leu Leu Ser Leu Leu
 1             5             10             15
Val Leu His Ser Ala Gln Gly Ala Thr Leu Gly Gly Pro Glu Glu
                20             25             30
Glu Ser Thr Ile Glu Asn Tyr Ala Ser Arg Pro Glu Ala Phe Asn
                35             40             45
Thr Pro Phe Leu Asn Ile Asp Lys Leu Arg Ser Ala Phe Lys Ala
                50             55             60
Asp Glu Phe Leu Asn Trp His Ala Leu Phe Glu Ser Ile Lys Arg
                65             70             75
Lys Leu Pro Phe Leu Asn Trp Asp Ala Phe Pro Lys Leu Lys Gly
                80             85             90
Leu Arg Ser Ala Thr Pro Asp Ala Gln
                95

```

&lt;210&gt; SEQ ID NO 23

&lt;211&gt; LENGTH: 866

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 23

```

tctcagactc ttggaagggg ctatactaga cacacaaaga cagccccaag           50
aaggacggtg gtagtagtgc ctgcctaaaa gacagtagat atgcaacgcc           100
tcttgctcct gccctttctc ctgctgggaa cagtttctgc tcttcatctg           150
gagaatgatg cccccatctt ggagagccta gagacacagg cagacctagg           200
ccaggatctg gatagttcaa aggagcagga gagagacttg gctotgacgg           250
aggaggtgat tcaggcagag ggagaggagg tcaaggcttc tgctgtcaa           300
gacaactttg aggatgagga agccatggag tcggaccacg ctgccttaga           350
caaggacttc cagtgcacca ggaagaaga cattggtgaa gtgcagggaa           400
gtccaaggtg caagacctgc cgctacctat tgggtcggac tcctaaaact           450
tttgcagaag ctcagaatgt ctgcagcaga tgctacggag gcaaccttgt           500
ctctatccat gacttcaact tcaactatcg cttcagtgct gcactagca           550
cagtcacaac agcccagggtc tggattggag gcaacctcag gggctgggtc           600
ctgtggaagc ggttttgctg gactgatggg agccactgga attttgctta           650
ctggtcccca gggcaacctg ggaatgggca aggtcctgtg gtggccctat           700
gcaccaaagg aggttattgg cgacgagctc aatgcgacaa gcaactgcc           750
ttcgtctgct ctttctaagc cagcggcagc gagaccctgc cagcagctcc           800
ctcccgctcc ccaacctctc ctgctcataa atccagactt cccacagcaa           850
aaaaaaaaaa aaaaaa                                           866

```

&lt;210&gt; SEQ ID NO 24

&lt;211&gt; LENGTH: 225

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien



-continued

&lt;400&gt; SEQUENCE: 24

Met	Gln	Arg	Leu	Leu	Leu	Leu	Pro	Phe	Leu	Leu	Leu	Gly	Thr	Val	
1			5						10					15	
Ser	Ala	Leu	His	Leu	Glu	Asn	Asp	Ala	Pro	His	Leu	Glu	Ser	Leu	
			20						25					30	
Glu	Thr	Gln	Ala	Asp	Leu	Gly	Gln	Asp	Leu	Asp	Ser	Ser	Lys	Glu	
			35						40					45	
Gln	Glu	Arg	Asp	Leu	Ala	Leu	Thr	Glu	Glu	Val	Ile	Gln	Ala	Glu	
			50						55					60	
Gly	Glu	Glu	Val	Lys	Ala	Ser	Ala	Cys	Gln	Asp	Asn	Phe	Glu	Asp	
			65						70					75	
Glu	Glu	Ala	Met	Glu	Ser	Asp	Pro	Ala	Ala	Leu	Asp	Lys	Asp	Phe	
			80						85					90	
Gln	Cys	Pro	Arg	Glu	Glu	Asp	Ile	Val	Glu	Val	Gln	Gly	Ser	Pro	
			95						100					105	
Arg	Cys	Lys	Thr	Cys	Arg	Tyr	Leu	Leu	Val	Arg	Thr	Pro	Lys	Thr	
			110						115					120	
Phe	Ala	Glu	Ala	Gln	Asn	Val	Cys	Ser	Arg	Cys	Tyr	Gly	Gly	Asn	
			125						130					135	
Leu	Val	Ser	Ile	His	Asp	Phe	Asn	Phe	Asn	Tyr	Arg	Ile	Gln	Cys	
			140						145					150	
Cys	Thr	Ser	Thr	Val	Asn	Gln	Ala	Gln	Val	Trp	Ile	Gly	Gly	Asn	
			155						160					165	
Leu	Arg	Gly	Trp	Phe	Leu	Trp	Lys	Arg	Phe	Cys	Trp	Thr	Asp	Gly	
			170						175					180	
Ser	His	Trp	Asn	Phe	Ala	Tyr	Trp	Ser	Pro	Gly	Gln	Pro	Gly	Asn	
			185						190					195	
Gly	Gln	Gly	Ser	Cys	Val	Ala	Leu	Cys	Thr	Lys	Gly	Gly	Tyr	Trp	
			200						205					210	
Arg	Arg	Ala	Gln	Cys	Asp	Lys	Gln	Leu	Pro	Phe	Val	Cys	Ser	Phe	
			215						220					225	

&lt;210&gt; SEQ ID NO 25

&lt;211&gt; LENGTH: 584

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 25

caacagaagc caagaaggaa gccgtctatc ttgtggcgat catgtataag	50
ctggcctcct gctgtttgc tttcacagga ttcttaaadc ctctcttctc	100
tcttctcttc cttgactcca gggaaatadc ctttcaactc tcagcacctc	150
atgaagacgc gcgcttaact ccggaggagc tagaaagagc ttcccttcta	200
cagatattgc cagagatgct gggcgagaa agaggggata ttctcaggaa	250
agcagactca agtaccaca tttttaacc aagaggaaat ttgagaaagt	300
ttcaggattt ctctggacaa gatcctaaca ttttactgag tcactctttg	350
gccagaatct gaaaccata caagaacgt gagactcctg attgcttctg	400
gaaatactgt gtctgaagt aaataagcat ctgttagtca gctcagaaac	450
accatctta gaatatgaaa aataacaca tgcttgattt gaaaacagt	500
tggagaaaa ctaggcaaac tacacctgt tcattgttac ctggaaaata	550

-continued

aatcctctat gttttgcaca aaaaaaaaaa aaaa 584

<210> SEQ ID NO 26  
 <211> LENGTH: 124  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 26

Met Tyr Lys Leu Ala Ser Cys Cys Leu Leu Phe Thr Gly Phe Leu  
 1 5 10 15  
 Asn Pro Leu Leu Ser Leu Pro Leu Leu Asp Ser Arg Glu Ile Ser  
 20 25 30  
 Phe Gln Leu Ser Ala Pro His Glu Asp Ala Arg Leu Thr Pro Glu  
 35 40 45  
 Glu Leu Glu Arg Ala Ser Leu Leu Gln Ile Leu Pro Glu Met Leu  
 50 55 60  
 Gly Ala Glu Arg Gly Asp Ile Leu Arg Lys Ala Asp Ser Ser Thr  
 65 70 75  
 Asn Ile Phe Asn Pro Arg Gly Asn Leu Arg Lys Phe Gln Asp Phe  
 80 85 90  
 Ser Gly Gln Asp Pro Asn Ile Leu Leu Ser His Leu Leu Ala Arg  
 95 100 105  
 Ile Trp Lys Pro Tyr Lys Lys Arg Glu Thr Pro Asp Cys Phe Trp  
 110 115 120

Lys Tyr Cys Val

<210> SEQ ID NO 27  
 <211> LENGTH: 920  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 27

caagtaaag cagcactagt ggggtgggatt gaggtatgcc ctgggtgcata 50  
 aatagagact cagctgtgct ggcacactca gaagcttggga ccgcatccta 100  
 gccgcccact cacacaaggc aggtgggtga ggaaatccag agttgccatg 150  
 gagaaaattc cagtgtcagc attcttgctc cttgtggccc tctcctacac 200  
 tctggccaga gataccacag tcaaacctgg agcctaaaag gacacaaagg 250  
 actctcgacc caaactgccc cagaccctct ccagaggttg gggtgaccaa 300  
 ctcatctgga ctcagacata tgaagaagct ctatataaat ccaagacaag 350  
 caacaaacc ttgatgatta ttcatacactt ggatgagtgc ccacacagtc 400  
 aagctttaa gaaagtgtt gctgaaaata aagaaatcca gaaattggca 450  
 gagcagtttg tcctcctcaa tctggtttat gaaacaactg acaaacacct 500  
 ttctcctgat ggccagtatg tccccaggat tatgtttggt gacctatctc 550  
 tgacagttag agccgatatc actggaagat attcaaactg tctctatgct 600  
 tacgaaactg cagatacagc tctgttgctt gacaacatga agaaagctct 650  
 caagttgctg aagactgaat tgtaaagaaa aaaaatctcc aagcccttct 700  
 gtctgtcagg ccttgagact tgaaaccaga agaagtgatga gaagactggc 750  
 tagtgtggaa gcatagtga cacactgatt aggttatggt ttaatgttac 800

-continued

---

```

aacaactatt ttttaagaaa aacaagtttt agaaatttgg tttcaagtgt      850
acatgtgtga aaacaatatt gtatactacc atagttagcc atgattttct      900
aaaaaaaaa ataatgtta                                           920

```

```

<210> SEQ ID NO 28
<211> LENGTH: 175
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

```

```

<400> SEQUENCE: 28

```

```

Met Glu Lys Ile Pro Val Ser Ala Phe Leu Leu Leu Val Ala Leu
 1             5             10             15
Ser Tyr Thr Leu Ala Arg Asp Thr Thr Val Lys Pro Gly Ala Lys
                20             25             30
Lys Asp Thr Lys Asp Ser Arg Pro Lys Leu Pro Gln Thr Leu Ser
                35             40             45
Arg Gly Trp Gly Asp Gln Leu Ile Trp Thr Gln Thr Tyr Glu Glu
                50             55             60
Ala Leu Tyr Lys Ser Lys Thr Ser Asn Lys Pro Leu Met Ile Ile
                65             70             75
His His Leu Asp Glu Cys Pro His Ser Gln Ala Leu Lys Lys Val
                80             85             90
Phe Ala Glu Asn Lys Glu Ile Gln Lys Leu Ala Glu Gln Phe Val
                95             100            105
Leu Leu Asn Leu Val Tyr Glu Thr Thr Asp Lys His Leu Ser Pro
                110            115            120
Asp Gly Gln Tyr Val Pro Arg Ile Met Phe Val Asp Pro Ser Leu
                125            130            135
Thr Val Arg Ala Asp Ile Thr Gly Arg Tyr Ser Asn Arg Leu Tyr
                140            145            150
Ala Tyr Glu Pro Ala Asp Thr Ala Leu Leu Leu Asp Asn Met Lys
                155            160            165
Lys Ala Leu Lys Leu Leu Lys Thr Glu Leu
                170            175

```

```

<210> SEQ ID NO 29
<211> LENGTH: 1181
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

```

```

<400> SEQUENCE: 29

```

```

aagaccctct ctttcgctgt ttgagagtct ctcggctcaa ggaccgggag      50
gtaagagggt tgggactgcc ccggcaactc caggggtgtct ggtccacgac      100
ctatcctagg cgccatgggt gtgataggta tacagctggt tgttaccatg      150
gtgatggcca gtgtcatgca gaagattata cctcactatt ctcttgctcg      200
atggctactc tgtaatggca gtttgaggty gtatcaacat cctacagaag      250
aagaattaag aattccttga gggaaacaac aaaaagggaa aaccaaaaaa      300
gataggaatg ataatgttca cattgaaagt aagccattaa ccattccaaa      350
ggatattgac cttcatctag aaacaaagtc agttacagaa gtggatactt      400
tagcattgca ttactttcca gaataccagt ggctgggtgga tttcacagtg      450

```

-continued

```

gctgctacag ttgtgtatct agtaactgaa gtctactaca attttatgaa      500
gcctacacag gaaatgaata tcagcttagt ctggcgccta cttgttttgt      550
cttttgcaat caaagttcta ttttcattaa ctacacacta ttttaaagta      600
gaagatggtg gtgaaagatc tgtttgtgtc acctttggat tttttttcct      650
tgtcaaagca atggcagtggt tgattgtaac agaaaattat ctggaatttg      700
gacttgaaac aggggtttaca aatttttcag acagtgcgat gcagtttctt      750
gaaaagcaag gtttagaatc tcagagtctt gtttcaaac ttactttcaa      800
atttttcctg gctattttct gttcattcat tggggctttt ttgacatttc      850
ctggattacg actggctcaa atgcactctgg atgccctgaa tttggcaaca      900
gaaaaaatta cacaaacttt acttcatatc aacttcttgg cacctttatt      950
tatggttttg ctctgggtaa aaccaatcac caaagactac attatgaacc     1000
caccactggg caaagaaatt tcccacatcg gaagatgaag ataatagtat     1050
ctaactcaca aggttatcat tggaataaat gaaagaacac atgtaatgca     1100
accagctgga attaagtgct taataaatgt tcttttctact gctttgcttc     1150
atcagaatta aaatagaaat acttgactag t                          1181
    
```

```

<210> SEQ ID NO 30
<211> LENGTH: 307
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien
    
```

<400> SEQUENCE: 30

```

Met Gly Val Ile Gly Ile Gln Leu Val Val Thr Met Val Met Ala
 1           5           10          15
Ser Val Met Gln Lys Ile Ile Pro His Tyr Ser Leu Ala Arg Trp
          20          25          30
Leu Leu Cys Asn Gly Ser Leu Arg Trp Tyr Gln His Pro Thr Glu
          35          40          45
Glu Glu Leu Arg Ile Leu Ala Gly Lys Gln Gln Lys Gly Lys Thr
          50          55          60
Lys Lys Asp Arg Lys Tyr Asn Gly His Ile Glu Ser Lys Pro Leu
          65          70          75
Thr Ile Pro Lys Asp Ile Asp Leu His Leu Glu Thr Lys Ser Val
          80          85          90
Thr Glu Val Asp Thr Leu Ala Leu His Tyr Phe Pro Glu Tyr Gln
          95          100         105
Trp Leu Val Asp Phe Thr Val Ala Ala Thr Val Val Tyr Leu Val
          110         115         120
Thr Glu Val Tyr Tyr Asn Phe Met Lys Pro Thr Gln Glu Met Asn
          125         130         135
Ile Ser Leu Val Trp Cys Leu Leu Val Leu Ser Phe Ala Ile Lys
          140         145         150
Val Leu Phe Ser Leu Thr Thr His Tyr Phe Lys Val Glu Asp Gly
          155         160         165
Gly Glu Arg Ser Val Cys Val Thr Phe Gly Phe Phe Phe Phe Val
          170         175         180
Lys Ala Met Ala Val Leu Ile Val Thr Glu Asn Tyr Leu Glu Phe
    
```

-continued

	185		190		195
Gly Leu Glu Thr	Gly Phe Thr Asn Phe Ser Asp Ser Ala Met Gln				
	200		205		210
Phe Leu Glu Lys	Gln Gly Leu Glu Ser Gln Ser Pro Val Ser Lys				
	215		220		225
Leu Thr Phe Lys	Phe Phe Leu Ala Ile Phe Cys Ser Phe Ile Gly				
	230		235		240
Ala Phe Leu Thr	Phe Pro Gly Leu Arg Leu Ala Gln Met His Leu				
	245		250		255
Asp Ala Leu Asn	Leu Ala Thr Glu Lys Ile Thr Gln Thr Leu Leu				
	260		265		270
His Ile Asn Phe	Leu Ala Pro Leu Phe Met Val Leu Leu Trp Val				
	275		280		285
Lys Pro Ile Thr	Lys Asp Tyr Ile Met Asn Pro Pro Leu Gly Lys				
	290		295		300
Glu Ile Ser Pro	Ser Gly Arg				
	305				

<210> SEQ ID NO 31  
 <211> LENGTH: 513  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 31

```

gtagcatagt gtgcagttca ctggacccaa agctttggct gcacctcttc      50
tggaagctg gccatggggc tcttcatgat cattgcaatt ctgotgttc      100
agaaacccc agtaaccgaa caacttaaga agtgctggaa taactatgta      150
caaggacatt gcaggaaaat ctgcagagta aatgaagtgc ctgaggcact      200
atgtgaaaaa gggagatact gttgcctcaa tatcaaggaa ctggaagcat      250
gtaaaaaaat tacaaagcca cctcgtccaa agccagcaac acttgactg      300
actcttcaag actatgttac aataatagaa aatttcccaa gcctgaagac      350
acagtctaca taaatcaaat acaatttcgt tttcacttgc ttctcaacct      400
agtctaataa actaaggtga tgagataac atcttcttcc ttctggtttc      450
ttgatcctta aaatgacctt cgagcatatt ctaataaagt gcattgccag      500
ttaaaaaaaaa aaa                                             513
    
```

<210> SEQ ID NO 32  
 <211> LENGTH: 99  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 32

Met Gly Leu Phe Met Ile Ile Ala Ile Leu Leu Phe Gln Lys Pro															
1				5				10							15
Thr Val Thr Glu Gln Leu Lys Lys Cys Trp Asn Asn Tyr Val Gln															
				20				25							30
Gly His Cys Arg Lys Ile Cys Arg Val Asn Glu Val Pro Glu Ala															
				35				40							45
Leu Cys Glu Asn Gly Arg Tyr Cys Cys Leu Asn Ile Lys Glu Leu															
				50				55							60

-continued

Glu Ala Cys Lys Lys Ile Thr Lys Pro Pro Arg Pro Lys Pro Ala  
 65 70 75  
 Thr Leu Ala Leu Thr Leu Gln Asp Tyr Val Thr Ile Ile Glu Asn  
 80 85 90  
 Phe Pro Ser Leu Lys Thr Gln Ser Thr  
 95

<210> SEQ ID NO 33  
 <211> LENGTH: 2684  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien  
 <220> FEATURE:  
 <221> NAME/KEY: unsure  
 <222> LOCATION: 2636-2637  
 <223> OTHER INFORMATION: unknown base

<400> SEQUENCE: 33

cgagacgctg ggcgctgagc cccggaggcc agggcgctccg gggctgcgcc 50  
 acttccgagg gccgagcgtc gccggtcccc gcggtgcgac acggccggga 100  
 ggagagaaac aacgcaaggc gctcaaccgt cggtcgctgg agccccccc 150  
 ggggctggcc ctcccggccc ctccagctggg gagggcgggg ctgctgcgc 200  
 cctgctgcgc actgcgaccc ttacagggga gggagggcgc aggccgcgcg 250  
 gagatgagga ggaggctgcg cctacgcagg gacgcattgc tcacgctgct 300  
 ccttgccgcc tccctgggccc tcttactcta tgcgcagcgc gacggcgcgg 350  
 ccccagcgcg gagcgcgcgc cgaggcgcag ggagggcggc accgagggccc 400  
 acccccggac cccgcgcgct ccagttaccc gacgcgggtg cagccccgcc 450  
 ggcctacgaa ggggacacac cggcgcccgc cacgcctacg ggaccctttg 500  
 acttcgcccc ctatttgcgc gccaaaggacc agcggcggtt tccactgctc 550  
 attaaccagc cgcacaagtg ccgcgggcgc ggcgcaccgc gtggccgccc 600  
 ggacctgttt attgctgtca agtcggtggc agaggacttc gagcggcgcc 650  
 aagccgtgcg ccagacgtgg ggcgcggagg gtcgctgca gggggcgctg 700  
 gtgcgccgcg tgttcttctt gggcgtgccc agggggcgag gctcggggcg 750  
 ggccgacgaa gttggggagg gcgcgcgaac ccaactggcg gccctgctgc 800  
 gggccgagag ccttgcgtat gcggacatcc tgctctgggc cttcgacgac 850  
 acctttttta acctaacgct caaggagatc cactttctag cctgggcctc 900  
 agctttctgc cccgacgtgc gcttcgtttt taagggcgac gcagatgtgt 950  
 tcgtgaacct gggaaatctc ctggagtccc tggcgcgcgc ggaccggcg 1000  
 caagacctgc ttgctggtga cgtaattgtg catgcgcggc ccatccgcac 1050  
 gcggctagc aagtactaca tccccaggc cgtgtacggc ctgcccgcct 1100  
 atccggccta cgcggggcgc ggtggctttg tgctttccgg ggcacgctg 1150  
 caccgcctgg ctggcgcctg tgcgcaggtc gagctcttcc ccatcgacga 1200  
 cgtctttctg ggcattgtgc tgcagcgcct goggetcacg cccgagcctc 1250  
 accctgcctt ccgacacctt ggcaccccc agccttcagc cgcgccgcat 1300  
 ttgagcacct tcgaccctg cttttaccgt gagctggttg tagtgcacgg 1350  
 gctctcgccc gctgacatct ggcttatgtg gcgcctgctg cacgggccc 1400

-continued

---

```

atgggccagc ctgtgcgcat ccacagcctg tcgctgcagg ccccttccaa      1450
tgggactcct agctccccac tacagcccca agctcctaac tcagaccag      1500
aatggagccg gtttcccaga ttattgccgt gtatgtgggtt cttccctgat      1550
caccaggtgc ctgtctccac aggatcccag gggatggggg ttaagcttgg      1600
ctcctggcgg tccaccctgc tggaaaccagt tgaaacccgt gtaatggtga      1650
ccctttgagc gagccaaggc tgggtggtag atgaccatct cttgtccaac      1700
aggtcccaga gcagtggata tgtctgggcc tcctagtagc acagaggtgt      1750
gttctggtgt ggtggcaggg acttagggaa tcctaccact ctgctggatt      1800
tggaaacccc taggctgacg cggacgtatg cagaggctct caaggccagg      1850
ccccacaggg aggtggaggg gtcctggcgg ccacagcctg aattcatgaa      1900
cctggcaggg actttgccat agctcatctg aaaacagata ttatgcttcc      1950
cacaacctct cctgggcccc ggtgtggctg agcaccaggg atggagccac      2000
acataagggg caaatgagtg cacggtccta cctagtcttt cctcacctcc      2050
tgaactcaca caacaatgcc agtctccac tggaggctgt atcccctcag      2100
aggagccaag gaatgtcttc ccctgagatg ccaccactat taatttcccc      2150
atatgcttca accacccctt tgctcaaaaa accaataccc aacttacct      2200
taatacaaac atcccagcaa cagcacatgg caggccattg ctgagggcac      2250
aggtgcttta ttggagaggg gatgtgggca ggggataagg aaggttcccc      2300
cattccagga ggatgggaac agtcctggct gccctgaca gtggggatat      2350
gcaagggggt ctggccaggc cacagtccaa atgggaagac accagtcagt      2400
cacaaaagtc gggagcgcca cacaaacctg gctataaggc ccaggaacca      2450
tataggagcc tgagacaggt ccctgcaca ttcacatta aactatacag      2500
gatgaggctg tacatgagtt aattacaaaa gagtcattt tacaaaaatc      2550
tgtacacaca tttgaaaaac tcacaaaatt gtcattatg tatcacaagt      2600
tgctagaccc aaaatattaa aaatgggata aaatnnnttt aaaaaaaaaa      2650
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa      2684

```

&lt;210&gt; SEQ ID NO 34

&lt;211&gt; LENGTH: 402

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 34

```

Met Arg Arg Arg Leu Arg Leu Arg Arg Asp Ala Leu Leu Thr Leu
 1           5           10           15
Leu Leu Gly Ala Ser Leu Gly Leu Leu Leu Tyr Ala Gln Arg Asp
          20           25           30
Gly Ala Ala Pro Thr Ala Ser Ala Pro Arg Gly Arg Gly Arg Ala
          35           40           45
Ala Pro Arg Pro Thr Pro Gly Pro Arg Ala Phe Gln Leu Pro Asp
          50           55           60
Ala Gly Ala Ala Pro Pro Ala Tyr Glu Gly Asp Thr Pro Ala Pro
          65           70           75

```

-continued

Pro	Thr	Pro	Thr	Gly	Pro	Phe	Asp	Phe	Ala	Arg	Tyr	Leu	Arg	Ala
				80					85					90
Lys	Asp	Gln	Arg	Arg	Phe	Pro	Leu	Leu	Ile	Asn	Gln	Pro	His	Lys
				95					100					105
Cys	Arg	Gly	Asp	Gly	Ala	Pro	Gly	Gly	Arg	Pro	Asp	Leu	Leu	Ile
				110					115					120
Ala	Val	Lys	Ser	Val	Ala	Glu	Asp	Phe	Glu	Arg	Arg	Gln	Ala	Val
				125					130					135
Arg	Gln	Thr	Trp	Gly	Ala	Glu	Gly	Arg	Val	Gln	Gly	Ala	Leu	Val
				140					145					150
Arg	Arg	Val	Phe	Leu	Leu	Gly	Val	Pro	Arg	Gly	Ala	Gly	Ser	Gly
				155					160					165
Gly	Ala	Asp	Glu	Val	Gly	Glu	Gly	Ala	Arg	Thr	His	Trp	Arg	Ala
				170					175					180
Leu	Leu	Arg	Ala	Glu	Ser	Leu	Ala	Tyr	Ala	Asp	Ile	Leu	Leu	Trp
				185					190					195
Ala	Phe	Asp	Asp	Thr	Phe	Phe	Asn	Leu	Thr	Leu	Lys	Glu	Ile	His
				200					205					210
Phe	Leu	Ala	Trp	Ala	Ser	Ala	Phe	Cys	Pro	Asp	Val	Arg	Phe	Val
				215					220					225
Phe	Lys	Gly	Asp	Ala	Asp	Val	Phe	Val	Asn	Val	Gly	Asn	Leu	Leu
				230					235					240
Glu	Phe	Leu	Ala	Pro	Arg	Asp	Pro	Ala	Gln	Asp	Leu	Leu	Ala	Gly
				245					250					255
Asp	Val	Ile	Val	His	Ala	Arg	Pro	Ile	Arg	Thr	Arg	Ala	Ser	Lys
				260					265					270
Tyr	Tyr	Ile	Pro	Glu	Ala	Val	Tyr	Gly	Leu	Pro	Ala	Tyr	Pro	Ala
				275					280					285
Tyr	Ala	Gly	Gly	Gly	Gly	Phe	Val	Leu	Ser	Gly	Ala	Thr	Leu	His
				290					295					300
Arg	Leu	Ala	Gly	Ala	Cys	Ala	Gln	Val	Glu	Leu	Phe	Pro	Ile	Asp
				305					310					315
Asp	Val	Phe	Leu	Gly	Met	Cys	Leu	Gln	Arg	Leu	Arg	Leu	Thr	Pro
				320					325					330
Glu	Pro	His	Pro	Ala	Phe	Arg	Thr	Phe	Gly	Ile	Pro	Gln	Pro	Ser
				335					340					345
Ala	Ala	Pro	His	Leu	Ser	Thr	Phe	Asp	Pro	Cys	Phe	Tyr	Arg	Glu
				350					355					360
Leu	Val	Val	Val	His	Gly	Leu	Ser	Ala	Ala	Asp	Ile	Trp	Leu	Met
				365					370					375
Trp	Arg	Leu	Leu	His	Gly	Pro	His	Gly	Pro	Ala	Cys	Ala	His	Pro
				380					385					390
Gln	Pro	Val	Ala	Ala	Gly	Pro	Phe	Gln	Trp	Asp	Ser			
				395					400					

<210> SEQ ID NO 35  
 <211> LENGTH: 1643  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien  
 <400> SEQUENCE: 35

agcagctctct gcccgaccgg gctcgtgctgg accccaggac cgggcgctgg 50  
 acgctgtcgt ccagcctccg gcgctgcgga gaccgcggc tgggtccggg 100



-continued

---

```

gaggcccaaa acccgcccc gccagaaccc cgccccaaat tcccacctcc      150
tccagaagcc ccgcccactc ccgagccccg agagctccgc gcacctgggc      200
gccatccgcc ctggctccgc tgcacagact ccacgcccgt accccggcgt      250
cacgctcagc ccgcggtgct cgcacacctg agactcatct cgcttcgacc      300
ccgcccggcg cgccgcccgg catcctgagc acggagacag tctccagctg      350
ccgttcatgc ttcctcccca gccttcgcga gccaccacag gaaggggagg      400
taggagtgcc cttttaccaa agggaccggc gatgctctgc aggctgtgct      450
ggctgggtct gtacagcttg gctgtgctgt tgctcggctg cctgctcttc      500
ctgaggaagg cggccaagcc cgcagagac  cccacggccc accagccttt      550
ctgggctccc ccaacacccc gtcacagccg gtgtccaccc aaccacacag      600
tgtctagcgc ctctctgtcc ctgctagacc gtcaccgtct cttcttgacc      650
tatcgtcact gccgaaatth ctctatcttg ctggagcctt caggctgttc      700
caagataacc ttcttgctcc tggccatcaa gtcacagcct ggtcacgtgg      750
agcgacgtgc ggctatccgc agcacgtggg gcagggtggg gggatgggct      800
aggggccggc agctgaagct ggtgttcctc ctaggggagg caggatccgc      850
tccccagcc  cagctgctgg cctatgagag tagggagttt gatgacatcc      900
tccagtgagg cttcactgag gacttcttca acctgacgct caaggagctg      950
cacctgcagc gctgggtggt ggctgcctgc ccccaggccc atttcatgct     1000
aaagggagat gacgatgtct ttgtccacgt cccaacgtg ttagagttcc     1050
tggatggctg ggaccagcc caggacctcc tgggtggaga tgtcatccgc     1100
caagccctgc ccaacaggaa cactaaggtc aaatacttca tcccaccctc     1150
aatgtacagg gccaccact acccacccta tgctgggtggg ggaggatag      1200
tcatgtccag agccacagtg cggcgctccc aggctatcat ggaagatgct     1250
gaaactcttc ccattgatga tgtctttgtg ggatgtgcc tgaggaggct     1300
ggggctgagc cctatgcacc atgctggctt caagacattt ggaatccggc     1350
ggccctgga  ccccttagac cctgctctgt atagggggct cctgctggtt     1400
caccgcctca gcccctcga gatgtggacc atgtgggcac tggtagacaga     1450
tgaggggctc aagtgtgcag ctggcccat  accccagcgc tgaagggagg      1500
gttgggcaac agcctgagag tggactcagt gttgattctc tatcgtgatg     1550
cgaaattgat gcctgctgct ctacagaaaa tgccaacttg gtttttaac     1600
tcctctcacc ctgttagctc tgattaaaaa cactgcaacc  caa           1643

```

&lt;210&gt; SEQ ID NO 36

&lt;211&gt; LENGTH: 378

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 36

```

Met Leu Pro Pro Gln Pro Ser Ala Ala His Gln Gly Arg Gly Gly
 1             5             10            15

```

```

Arg Ser Gly Leu Leu Pro Lys Gly Pro Ala Met Leu Cys Arg Leu
 20            25            30

```

-continued

---

Cys Trp Leu Val Ser Tyr Ser Leu Ala Val Leu Leu Leu Gly Cys  
 35 40 45  
 Leu Leu Phe Leu Arg Lys Ala Ala Lys Pro Ala Gly Asp Pro Thr  
 50 55 60  
 Ala His Gln Pro Phe Trp Ala Pro Pro Thr Pro Arg His Ser Arg  
 65 70 75  
 Cys Pro Pro Asn His Thr Val Ser Ser Ala Ser Leu Ser Leu Pro  
 80 85 90  
 Ser Arg His Arg Leu Phe Leu Thr Tyr Arg His Cys Arg Asn Phe  
 95 100 105  
 Ser Ile Leu Leu Glu Pro Ser Gly Cys Ser Lys Asp Thr Phe Leu  
 110 115 120  
 Leu Leu Ala Ile Lys Ser Gln Pro Gly His Val Glu Arg Arg Ala  
 125 130 135  
 Ala Ile Arg Ser Thr Trp Gly Arg Val Gly Gly Trp Ala Arg Gly  
 140 145 150  
 Arg Gln Leu Lys Leu Val Phe Leu Leu Gly Val Ala Gly Ser Ala  
 155 160 165  
 Pro Pro Ala Gln Leu Leu Ala Tyr Glu Ser Arg Glu Phe Asp Asp  
 170 175 180  
 Ile Leu Gln Trp Asp Phe Thr Glu Asp Phe Phe Asn Leu Thr Leu  
 185 190 195  
 Lys Glu Leu His Leu Gln Arg Trp Val Val Ala Ala Cys Pro Gln  
 200 205 210  
 Ala His Phe Met Leu Lys Gly Asp Asp Asp Val Phe Val His Val  
 215 220 225  
 Pro Asn Val Leu Glu Phe Leu Asp Gly Trp Asp Pro Ala Gln Asp  
 230 235 240  
 Leu Leu Val Gly Asp Val Ile Arg Gln Ala Leu Pro Asn Arg Asn  
 245 250 255  
 Thr Lys Val Lys Tyr Phe Ile Pro Pro Ser Met Tyr Arg Ala Thr  
 260 265 270  
 His Tyr Pro Pro Tyr Ala Gly Gly Gly Gly Tyr Val Met Ser Arg  
 275 280 285  
 Ala Thr Val Arg Arg Leu Gln Ala Ile Met Glu Asp Ala Glu Leu  
 290 295 300  
 Phe Pro Ile Asp Asp Val Phe Val Gly Met Cys Leu Arg Arg Leu  
 305 310 315  
 Gly Leu Ser Pro Met His His Ala Gly Phe Lys Thr Phe Gly Ile  
 320 325 330  
 Arg Arg Pro Leu Asp Pro Leu Asp Pro Cys Leu Tyr Arg Gly Leu  
 335 340 345  
 Leu Leu Val His Arg Leu Ser Pro Leu Glu Met Trp Thr Met Trp  
 350 355 360  
 Ala Leu Val Thr Asp Glu Gly Leu Lys Cys Ala Ala Gly Pro Ile  
 365 370 375  
 Pro Gln Arg

&lt;210&gt; SEQ ID NO 37

&lt;211&gt; LENGTH: 1226

&lt;212&gt; TYPE: DNA

-continued

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 37

```

atgaaagtga taatcaggca gcccaaatga ttgtaataa ggatcaaagt      50
agatcgtgta tgtgggtcca atcaattgat tctacacaaa ggagcctggg      100
gaggggccat ggtgccaatg cacttactgg ggagactgga gaagccgctt      150
ctcctcctgt gctgcgcctc cttcctactg gggctggctt tgctgggcat      200
aaagacggac atcacccccg ttgcttattt ctttctcaca ttgggtggct      250
tcttcttggt tgcctatctc ctggctccgt ttctggaatg ggggcttcgg      300
tcccagctcc aatcaatgca gactgagagc ccagggcctt caggcaatgc      350
acgggacaat gaagcctttg aagtgccagt ctatgaagag gccgtgggtg      400
gactagaatc ccagtgcgcg ccccaagagt tggaccaacc acccccctac      450
agcactgttg tgatacccc agcacctgag gaggaacaac ctagccatcc      500
agaggggtcc aggagagcca aactggaaca gaggcgaatg gcctcagagg      550
ggtccatggc ccaggaagga agccctggaa gagctccaat caaccttcgg      600
cttcggggac cacgggctgt gtccactgct cctgatctgc agagcttggc      650
ggcagtcacc acattagagc ctctgactcc acccctgcc tatgatgtct      700
gctttgtgca ccctgatgat gatagtgttt tttatgagga caactgggca      750
cccccttaaa tgactctccc aagatttctc ttctctccac accagacctc      800
gttcatttga ctaacatttt ccagcgccta ctatgtgtca gaaacaagtg      850
tttctgcctg gacatcataa atggggactt ggaccctgag gagagtcagg      900
ccacggtaag cccttcccag ctgagatatg ggtggcataa tttgagtctt      950
ctggcaacat ttggtgacct accccatctc caatatttcc agcgttagat      1000
tgaggatgag gtagggaggt gatccagaga aggcggagaa ggaagaagta      1050
acctctgagt ggcggctatt gcttctgttc cagggtgctgt tcgagctggt      1100
agaaccctta ggcttgacag ctttgtgagt tattattgaa aaatgaggat      1150
tccaagagtc agaggagtgt gataatgtgc acgagggcac actgctagta      1200
aataacatta aaataactgg aatgaa                                1226

```

&lt;210&gt; SEQ ID NO 38

&lt;211&gt; LENGTH: 216

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 38

```

Met Val Pro Met His Leu Leu Gly Arg Leu Glu Lys Pro Leu Leu
 1                5                10                15

Leu Leu Cys Cys Ala Ser Phe Leu Leu Gly Leu Ala Leu Leu Gly
 20                25                30

Ile Lys Thr Asp Ile Thr Pro Val Ala Tyr Phe Phe Leu Thr Leu
 35                40                45

Gly Gly Phe Phe Leu Phe Ala Tyr Leu Leu Val Arg Phe Leu Glu
 50                55                60

Trp Gly Leu Arg Ser Gln Leu Gln Ser Met Gln Thr Glu Ser Pro
 65                70                75

```

-continued

Gly Pro Ser Gly Asn Ala Arg Asp Asn Glu Ala Phe Glu Val Pro  
 80 85 90  
 Val Tyr Glu Glu Ala Val Val Gly Leu Glu Ser Gln Cys Arg Pro  
 95 100 105  
 Gln Glu Leu Asp Gln Pro Pro Pro Tyr Ser Thr Val Val Ile Pro  
 110 115 120  
 Pro Ala Pro Glu Glu Glu Gln Pro Ser His Pro Glu Gly Ser Arg  
 125 130 135  
 Arg Ala Lys Leu Glu Gln Arg Arg Met Ala Ser Glu Gly Ser Met  
 140 145 150  
 Ala Gln Glu Gly Ser Pro Gly Arg Ala Pro Ile Asn Leu Arg Leu  
 155 160 165  
 Arg Gly Pro Arg Ala Val Ser Thr Ala Pro Asp Leu Gln Ser Leu  
 170 175 180  
 Ala Ala Val Pro Thr Leu Glu Pro Leu Thr Pro Pro Pro Ala Tyr  
 185 190 195  
 Asp Val Cys Phe Gly His Pro Asp Asp Asp Ser Val Phe Tyr Glu  
 200 205 210  
 Asp Asn Trp Ala Pro Pro  
 215

<210> SEQ ID NO 39  
 <211> LENGTH: 2770  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 39

cccacgcgctc cggcggctac acacctaggt gcggtgggct tcgggtgggg 50  
 ggcctgcagc tagctgatgg caagggagga atagcagggg tggggattgt 100  
 ggtgtgcgag aggtcccgcg gacggggggc tcgggggtct cttcagacga 150  
 gattcccttc aggcttgggc cgggtccctt cgcacggaga tcccaatgaa 200  
 cgcgggcccc tggaggcccg tggttggggc ttctcccgct cggggatggg 250  
 gccggtaccc tagcccgttt ccagcgcctc agtcggttcc ccatgcccctc 300  
 agaggtggcc cggggcaagc gcgccccctt cttcttcgct gcggtggcca 350  
 tcgtgctggg gctaccgctc tgggtggaaga ccacggagac ctaccggggc 400  
 tcgttgctt actcccagat cagtggcctg aatgcccttc agctccgcct 450  
 catggtgcct gtcactgtcg tgtttacgcg ggagtcagtg ccctgggacg 500  
 accaggagaa gctgcccttc accgttgtgc atgaaagaga gattcctctg 550  
 aaatacaaaa tgaaaatcaa atgccgttcc cagaagcct atcggaggggc 600  
 tttgaccat gaggaggagg ccctgtcatc gggcagtggt caagaggcag 650  
 aagccatggt agatgagcct caggaacaag cggagggctc cctgactgtg 700  
 tacgtgatat ctgaacactc ctcacttctt cccagagaca tgatgagcta 750  
 cattgggccc aagaggacag cagtgggtgc ggggataatg caccgggagg 800  
 cctttaacat cattggccgc cgcatagtcc aggtggccca ggccatgtct 850  
 ttgactgagg atgtgcttgc tgctgctctg gctgaccacc ttccagagga 900  
 caagtggagc gctgagaaga ggcggcctct caagtccagc ttgggctatg 950

-continued

---

agatcacctt cagtttactc aaccagacc ccaagtcca tgatgtctac	1000
tgggacattg agggggctgt cggcgctat gtgcaacctt tcctgaatgc	1050
cctcggtgcc gctggcaact tctctgtgga ctctcagatt ctttactatg	1100
caatgttggg ggtgaatccc cgctttgact cagcttctc cagctactat	1150
ttggacatgc acagcctccc ccatgtcatc aaccagtggt agtcccggct	1200
gggatccagt gctgcctcct tgtaccctgt gctcaacttt ctactctacg	1250
tgccctgagct tgcacactca ccgctgtaca ttcaggacaa ggatggcgct	1300
ccagtgggcca ccaatgcctt ccatagtccc cgctgggggtg gcattatggt	1350
atataatggt gactccaaaa cctataatgc ctcaagtctg ccagtgagag	1400
tcgaggtgga catgggtcga gtgatggagg tgttctctggc acagttgcgg	1450
ttgctctttg ggattgctca gcccagctg cctccaaaat gcctgctttc	1500
agggcctacg agtgaagggc taatgacctg ggagctagac cggctgctct	1550
gggctcggtc agtggagaac ctggccacag ccaccaccac ccttacctcc	1600
ctggcgacgc ttctgggcaa gatcagcaac attgtcatta aggacgacgt	1650
ggcatctgag gtgtacaagg ctgtagctgc cgtccagaag tcggcagaag	1700
agttggcgtc tgggcacctg gcatctgcct ttgtcgccag ccaggaagct	1750
gtgacatcct ctgagcttgc cttctttgac ccgtcactcc tccacctcct	1800
ttatttccct gatgaccaga agtttgccat ctacatccca ctcttctgctc	1850
ctatggctgt gccatcctc ctgtccctgg tcaagatcct cctggagacc	1900
cgcaagtctt ggagaaagcc tgagaagaca gactgagcag ggcagacct	1950
ccataggaag ccttcctttc tggccaaggt gggcggtggt agattgtgag	2000
gcacgtacat ggggcctgcc ggaatgacct aaatatttgt ctccagtctc	2050
cactgttggc tctccagcaa ccaaagtaca acaactccaag atgggttcat	2100
cttttcttcc tttcccattc acctggctca atcctcctcc accaccaggg	2150
gcctcaaaag gcacatcctc cgggtctcct tatcttgttt gataaggctg	2200
ctgcctgtct ccctctgtgg caaggactgt ttgttctttt gccccatttc	2250
tcaacatagc acacttgtgc actgagagga gggagcatta tgggaaagtc	2300
cctgccttcc acacctctct ctagtccctg tgggacagcc ctagcccctg	2350
ctgtcatgaa ggggccaggc attggtcacc tgtgggacct tctccctcac	2400
tcccctccct cctagttggc tttgtctgtc aggtgcagtc tggcgggagt	2450
ccagaggcoa gcagctcagg acatggtgct gtgtgtgtgt gtgtgtgtgt	2500
gtgtgtgtgt gtgtgtgtca gaggttccag aaagttccag atttggaaac	2550
aaacagtctt gaattcaaat ccttgttttt gcacttattg tctggagagc	2600
tttgataag gtattgaatc tctctgagcc tcagtttttc atttgttcaa	2650
atggcaactga tgatgtctcc cttacaagat ggtgtgaggt agtaaatgtg	2700
atcagcatgt aaagtgtctg gcgtgtagta ggctottaat aaacactggc	2750
tgaatatgaa ttggaatgat	2770

-continued

---

```

<210> SEQ ID NO 40
<211> LENGTH: 547
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 40

Met Pro Ser Glu Val Ala Arg Gly Lys Arg Ala Ala Leu Phe Phe
 1          5          10          15
Ala Ala Val Ala Ile Val Leu Gly Leu Pro Leu Trp Trp Lys Thr
 20          25          30
Thr Glu Thr Tyr Arg Ala Ser Leu Pro Tyr Ser Gln Ile Ser Gly
 35          40          45
Leu Asn Ala Leu Gln Leu Arg Leu Met Val Pro Val Thr Val Val
 50          55          60
Phe Thr Arg Glu Ser Val Pro Leu Asp Asp Gln Glu Lys Leu Pro
 65          70          75
Phe Thr Val Val His Glu Arg Glu Ile Pro Leu Lys Tyr Lys Met
 80          85          90
Lys Ile Lys Cys Arg Phe Gln Lys Ala Tyr Arg Arg Ala Leu Asp
 95          100         105
His Glu Glu Glu Ala Leu Ser Ser Gly Ser Val Gln Glu Ala Glu
 110         115         120
Ala Met Leu Asp Glu Pro Gln Glu Gln Ala Glu Gly Ser Leu Thr
 125         130         135
Val Tyr Val Ile Ser Glu His Ser Ser Leu Leu Pro Gln Asp Met
 140         145         150
Met Ser Tyr Ile Gly Pro Lys Arg Thr Ala Val Val Arg Gly Ile
 155         160         165
Met His Arg Glu Ala Phe Asn Ile Ile Gly Arg Arg Ile Val Gln
 170         175         180
Val Ala Gln Ala Met Ser Leu Thr Glu Asp Val Leu Ala Ala Ala
 185         190         195
Leu Ala Asp His Leu Pro Glu Asp Lys Trp Ser Ala Glu Lys Arg
 200         205         210
Arg Pro Leu Lys Ser Ser Leu Gly Tyr Glu Ile Thr Phe Ser Leu
 215         220         225
Leu Asn Pro Asp Pro Lys Ser His Asp Val Tyr Trp Asp Ile Glu
 230         235         240
Gly Ala Val Arg Arg Tyr Val Gln Pro Phe Leu Asn Ala Leu Gly
 245         250         255
Ala Ala Gly Asn Phe Ser Val Asp Ser Gln Ile Leu Tyr Tyr Ala
 260         265         270
Met Leu Gly Val Asn Pro Arg Phe Asp Ser Ala Ser Ser Ser Tyr
 275         280         285
Tyr Leu Asp Met His Ser Leu Pro His Val Ile Asn Pro Val Glu
 290         295         300
Ser Arg Leu Gly Ser Ser Ala Ala Ser Leu Tyr Pro Val Leu Asn
 305         310         315
Phe Leu Leu Tyr Val Pro Glu Leu Ala His Ser Pro Leu Tyr Ile
 320         325         330
Gln Asp Lys Asp Gly Ala Pro Val Ala Thr Asn Ala Phe His Ser
 335         340         345

```

-continued

---

Pro Arg Trp Gly Gly Ile Met Val Tyr Asn Val Asp Ser Lys Thr  
 350 355 360

Tyr Asn Ala Ser Val Leu Pro Val Arg Val Glu Val Asp Met Val  
 365 370 375

Arg Val Met Glu Val Phe Leu Ala Gln Leu Arg Leu Leu Phe Gly  
 380 385 390

Ile Ala Gln Pro Gln Leu Pro Pro Lys Cys Leu Leu Ser Gly Pro  
 395 400 405

Thr Ser Glu Gly Leu Met Thr Trp Glu Leu Asp Arg Leu Leu Trp  
 410 415 420

Ala Arg Ser Val Glu Asn Leu Ala Thr Ala Thr Thr Thr Leu Thr  
 425 430 435

Ser Leu Ala Gln Leu Leu Gly Lys Ile Ser Asn Ile Val Ile Lys  
 440 445 450

Asp Asp Val Ala Ser Glu Val Tyr Lys Ala Val Ala Ala Val Gln  
 455 460 465

Lys Ser Ala Glu Glu Leu Ala Ser Gly His Leu Ala Ser Ala Phe  
 470 475 480

Val Ala Ser Gln Glu Ala Val Thr Ser Ser Glu Leu Ala Phe Phe  
 485 490 495

Asp Pro Ser Leu Leu His Leu Leu Tyr Phe Pro Asp Asp Gln Lys  
 500 505 510

Phe Ala Ile Tyr Ile Pro Leu Phe Leu Pro Met Ala Val Pro Ile  
 515 520 525

Leu Leu Ser Leu Val Lys Ile Phe Leu Glu Thr Arg Lys Ser Trp  
 530 535 540

Arg Lys Pro Glu Lys Thr Asp  
 545

<210> SEQ ID NO 41  
 <211> LENGTH: 1964  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 41

```

ccagctgcag agaggaggag gtgagctgca gagaagagga ggttggtgtg          50
gagcacaggc agcaccgagc ctgccccgtg agctgagggc ctgcagtctg          100
cggctggaat caggatagac accaaggcag gacccccaga gatgctgaag          150
cctctttgga aagcagcagt ggccccaca tggccatgct ccatgccgcc          200
ccgcccccg tgggacagag aggctggcac gttgcaggtc ctgggagcgc          250
tggctgtgct gtggctgggc tccgtggctc ttatctgcct cctgtggcaa          300
gtgccccctg ctcccacctg gggccagggt cagcccaagg acgtgccccg          350
gtcctgggag catggctcca gccagcttg ggagcccctg gaagcagagg          400
ccaggcagca gagggactcc tgccagcttg tccttgtgga aagcatcccc          450
caggacctgc catctgcagc cggcagcccc tctgcccagc ctctgggccca          500
ggcctggctg cagctgctgg aactgcccga ggagagcgtc cacgtggctt          550
catactactg gtcctcaca gggcctgaca toggggtcaa cgactcgtct          600
tcccagctgg gagaggctct tctgcagaag ctgcagcagc tgctgggcag          650
    
```

-continued

```

gaacatttcc ctggctgtgg ccaccagcag cccgacactg gccaggacat      700
ccaccgacct gcaggttctg gctgcccagc gtgccatgt acgacaggtg      750
cccatggggc ggctcaccag ggggtgtttg cactccaaat tctgggttgt      800
ggatggacgg cacatataca tgggcagtgc caacatggac tggcggttct      850
tgacgcaggt gaaggagcct ggcgctgtca tctataactg cagccacctg      900
gcccagaacc tggagaagac cttccagacc tactgggtac tgggggtgcc      950
caaggctgtc ctccccaaaa cctggcctca gaacttctca tctcaactca     1000
accgtttcca gcccttccac ggctctttg atgggggtgcc caccactgcc     1050
tacttctcag cgtcgccacc agcactctgt ccccagggcc gcacccggga     1100
cctggaggcg ctgctggcgg tgatggggag cgcccaggag ttcattatg      1150
cctccgtgat ggagtatttc cccaccacgc gcttcagcca cccccgagg      1200
tactggccgg tgctggacaa cgcgctgcgg gcggcagcct tcggcaaggg      1250
cgtgcgcgtg cgcctgctgg tcggctgcgg actcaacacg gaccccacca     1300
tgttccccta cctgcggtcc ctgcaggcgc tcagcaaccc cgcggccaac     1350
gtctctgtgg acgtgaaagt ctatcctgtg ccggtgggga accattocaa     1400
catcccattc agcaggggta accacagcaa gttcatggtc acggagaagg     1450
cagcctacat aggcacctcc aactggtcgg aggattactt cagcagcacg     1500
gcgggggtgg gcttgggtgt caccagagc cctggcgcgc agcccgcggg     1550
ggccacggtg caggagcagc tgccgagcct ctttgagcgg gactggagtt     1600
cgcgctacgc cgtcggcctg gacggacagg ctccgggcca ggactgcgtt     1650
tggcagggct gaggggggcc tctttttctc tcggcgacc cgccccgac      1700
gcgccttccc ctctgacccc ggctgggctc tcagccgctt cctcccgcaa     1750
gcagcccggg tccgcaactgc gccaggagcc gcctgcgacc gcccgggcgt     1800
cgcaaacccg ccgctgtctc tctgatttcc gagtccagcc cccctgagc     1850
cccactcctc ccaggagacc ctccaggaag ccccttccct gactcctggc     1900
ccacaggcca ggcctaaaaa aaactcgtgg cttcaaaaaa aaaaaaaaaa     1950
aaaaaaaaaa aaaa                                             1964

```

&lt;210&gt; SEQ ID NO 42

&lt;211&gt; LENGTH: 489

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 42

```

Met Pro Pro Arg Arg Pro Trp Asp Arg Glu Ala Gly Thr Leu Gln
 1           5           10
Val Leu Gly Ala Leu Ala Val Leu Trp Leu Gly Ser Val Ala Leu
          20           25           30
Ile Cys Leu Leu Trp Gln Val Pro Arg Pro Pro Thr Trp Gly Gln
          35           40           45
Val Gln Pro Lys Asp Val Pro Arg Ser Trp Glu His Gly Ser Ser
          50           55           60
Pro Ala Trp Glu Pro Leu Glu Ala Glu Ala Arg Gln Gln Arg Asp
          65           70           75

```



-continued

---

Ser	Cys	Gln	Leu	Val	Leu	Val	Glu	Ser	Ile	Pro	Gln	Asp	Leu	Pro
				80					85					90
Ser	Ala	Ala	Gly	Ser	Pro	Ser	Ala	Gln	Pro	Leu	Gly	Gln	Ala	Trp
				95					100					105
Leu	Gln	Leu	Leu	Asp	Thr	Ala	Gln	Glu	Ser	Val	His	Val	Ala	Ser
				110					115					120
Tyr	Tyr	Trp	Ser	Leu	Thr	Gly	Pro	Asp	Ile	Gly	Val	Asn	Asp	Ser
				125					130					135
Ser	Ser	Gln	Leu	Gly	Glu	Ala	Leu	Leu	Gln	Lys	Leu	Gln	Gln	Leu
				140					145					150
Leu	Gly	Arg	Asn	Ile	Ser	Leu	Ala	Val	Ala	Thr	Ser	Ser	Pro	Thr
				155					160					165
Leu	Ala	Arg	Thr	Ser	Thr	Asp	Leu	Gln	Val	Leu	Ala	Ala	Arg	Gly
				170					175					180
Ala	His	Val	Arg	Gln	Val	Pro	Met	Gly	Arg	Leu	Thr	Arg	Gly	Val
				185					190					195
Leu	His	Ser	Lys	Phe	Trp	Val	Val	Asp	Gly	Arg	His	Ile	Tyr	Met
				200					205					210
Gly	Ser	Ala	Asn	Met	Asp	Trp	Arg	Ser	Leu	Thr	Gln	Val	Lys	Glu
				215					220					225
Leu	Gly	Ala	Val	Ile	Tyr	Asn	Cys	Ser	His	Leu	Ala	Gln	Asp	Leu
				230					235					240
Glu	Lys	Thr	Phe	Gln	Thr	Tyr	Trp	Val	Leu	Gly	Val	Pro	Lys	Ala
				245					250					255
Val	Leu	Pro	Lys	Thr	Trp	Pro	Gln	Asn	Phe	Ser	Ser	His	Phe	Asn
				260					265					270
Arg	Phe	Gln	Pro	Phe	His	Gly	Leu	Phe	Asp	Gly	Val	Pro	Thr	Thr
				275					280					285
Ala	Tyr	Phe	Ser	Ala	Ser	Pro	Pro	Ala	Leu	Cys	Pro	Gln	Gly	Arg
				290					295					300
Thr	Arg	Asp	Leu	Glu	Ala	Leu	Leu	Ala	Val	Met	Gly	Ser	Ala	Gln
				305					310					315
Glu	Phe	Ile	Tyr	Ala	Ser	Val	Met	Glu	Tyr	Phe	Pro	Thr	Thr	Arg
				320					325					330
Phe	Ser	His	Pro	Pro	Arg	Tyr	Trp	Pro	Val	Leu	Asp	Asn	Ala	Leu
				335					340					345
Arg	Ala	Ala	Ala	Phe	Gly	Lys	Gly	Val	Arg	Val	Arg	Leu	Leu	Val
				350					355					360
Gly	Cys	Gly	Leu	Asn	Thr	Asp	Pro	Thr	Met	Phe	Pro	Tyr	Leu	Arg
				365					370					375
Ser	Leu	Gln	Ala	Leu	Ser	Asn	Pro	Ala	Ala	Asn	Val	Ser	Val	Asp
				380					385					390
Val	Lys	Val	Phe	Ile	Val	Pro	Val	Gly	Asn	His	Ser	Asn	Ile	Pro
				395					400					405
Phe	Ser	Arg	Val	Asn	His	Ser	Lys	Phe	Met	Val	Thr	Glu	Lys	Ala
				410					415					420
Ala	Tyr	Ile	Gly	Thr	Ser	Asn	Trp	Ser	Glu	Asp	Tyr	Phe	Ser	Ser
				425					430					435
Thr	Ala	Gly	Val	Gly	Leu	Val	Val	Thr	Gln	Ser	Pro	Gly	Ala	Gln
				440					445					450

-continued

Pro Ala Gly Ala Thr Val Gln Glu Gln Leu Arg Gln Leu Phe Glu  
 455 460 465

Arg Asp Trp Ser Ser Arg Tyr Ala Val Gly Leu Asp Gly Gln Ala  
 470 475 480

Pro Gly Gln Asp Cys Val Trp Gln Gly  
 485

<210> SEQ ID NO 43  
 <211> LENGTH: 1130  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 43

```

gggcctggcg atccgcatcc cgcagcgcg ctggtgctgc tgcccggctg      50
tctgtcgtca tggtagggcc ctgggtgtat ctggtggcgg cagttttgct      100
catcggcctg atcctcttcc tgactcgag cgggggtcgg gcggcagcag      150
ctgacggaga accactgcac aatgaggaag agagggcagg agcaggccag      200
gtagggcctg ctttgcccca ggagtctgaa gaacagagaa ctggaagcag      250
accccgcgct cggagggact tgggcagccg tctacaggcc cagcgtcgag      300
cccagcaggt ggcctgggaa gacggggatg agaatgtggg tcaaactggt      350
attccagccc aggaggaaga aggcattgag aagccagcag aagttcacc      400
aacagggaaa attggagcca agaaactacg gaagctagag gaaaaacagg      450
ctcgaaaagg tcagcgagag gcagaggagg ctgaactga agaacggaaa      500
cgcctagagt cccaactga ggccgaatgg aagaaggaa aggaacggct      550
tcgcctgaag gaagaacaga aggaggagga agagaggaa gctcaggagg      600
agcaggcccc gcgggatcac gaggagtacc tgaaactgaa ggaggccttc      650
gtggtagaag aagaaggtgt tagcgaaacc atgactgagg agcagtctca      700
cagttctctg acagaattca tcaattacat caagaagtcc aagttgtgc      750
ttttggaaga tctggctttc cagatgggcc taaggactca ggacgccata      800
aaccgcatcc aggacctgct gacggagggg actctaacag gtgtgattga      850
cgaccggggg aagtttatct acataacccc agaggaactg gctgccgtgg      900
ccaatttcat ccgacagcgg ggccgggtgt ccatcacaga gcttgcccat      950
gccagcaact ccctcatctc ctggggccag gacctccctg cccaggcttc     1000
agcctgactc cagtccttcc ttgagtgtat cctgtggcct acatgtgtct     1050
tcctccttcc ctaatgccgt cttggggcag ggatggaata tgaccagaaa     1100
gttgtggatt aaagcctgt gaatactgaa                                1130
    
```

<210> SEQ ID NO 44  
 <211> LENGTH: 315  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 44

Met Val Gly Pro Trp Val Tyr Leu Val Ala Ala Val Leu Leu Ile  
 1 5 10 15

Gly Leu Ile Leu Phe Leu Thr Arg Ser Arg Gly Arg Ala Ala Ala  
 20 25 30

-continued

---

Ala Asp Gly Glu Pro Leu His Asn Glu Glu Glu Arg Ala Gly Ala  
 35 40 45

Gly Gln Val Gly Arg Ser Leu Pro Gln Glu Ser Glu Glu Gln Arg  
 50 55 60

Thr Gly Ser Arg Pro Arg Arg Arg Arg Asp Leu Gly Ser Arg Leu  
 65 70 75

Gln Ala Gln Arg Arg Ala Gln Arg Val Ala Trp Glu Asp Gly Asp  
 80 85 90

Glu Asn Val Gly Gln Thr Val Ile Pro Ala Gln Glu Glu Glu Gly  
 95 100 105

Ile Glu Lys Pro Ala Glu Val His Pro Thr Gly Lys Ile Gly Ala  
 110 115 120

Lys Lys Leu Arg Lys Leu Glu Glu Lys Gln Ala Arg Lys Ala Gln  
 125 130 135

Arg Glu Ala Glu Glu Ala Glu Arg Glu Glu Arg Lys Arg Leu Glu  
 140 145 150

Ser Gln Arg Glu Ala Glu Trp Lys Lys Glu Glu Glu Arg Leu Arg  
 155 160 165

Leu Lys Glu Glu Gln Lys Glu Glu Glu Glu Arg Lys Ala Gln Glu  
 170 175 180

Glu Gln Ala Arg Arg Asp His Glu Glu Tyr Leu Lys Leu Lys Glu  
 185 190 195

Ala Phe Val Val Glu Glu Glu Gly Val Ser Glu Thr Met Thr Glu  
 200 205 210

Glu Gln Ser His Ser Phe Leu Thr Glu Phe Ile Asn Tyr Ile Lys  
 215 220 225

Lys Ser Lys Val Val Leu Leu Glu Asp Leu Ala Phe Gln Met Gly  
 230 235 240

Leu Arg Thr Gln Asp Ala Ile Asn Arg Ile Gln Asp Leu Leu Thr  
 245 250 255

Glu Gly Thr Leu Thr Gly Val Ile Asp Asp Arg Gly Lys Phe Ile  
 260 265 270

Tyr Ile Thr Pro Glu Glu Leu Ala Ala Val Ala Asn Phe Ile Arg  
 275 280 285

Gln Arg Gly Arg Val Ser Ile Thr Glu Leu Ala Gln Ala Ser Asn  
 290 295 300

Ser Leu Ile Ser Trp Gly Gln Asp Leu Pro Ala Gln Ala Ser Ala  
 305 310 315

<210> SEQ ID NO 45  
 <211> LENGTH: 1977  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 45

```

acggggcgca gcggcagtga cgtagggttg ggcacggat ccgttgcggc           50
tgca gctctg cagtcgggcc gttccttcgc cgccgccagg ggtagcggtg           100
tagctgcgca gcgtcgcgcg cgctaccgca cccaggttcg gcccgtaggc           150
gtctggcagc ccggcgccat cttcatcgag cgccatggcc gcagcctgag           200
ggccgggagc ggccgggtac tgcttctcc tcggcttgca tttgtttctg           250
    
```

-continued

---

ctgaccgcgg gccctgccct gggctggaac gaccctgaca gaatgttgct	300
gcgggatgta aaagctctta ccctccacta tgaccgctat accacctccc	350
gcaggctgga tcccatccca cagttgaaat gtgttgagg cacagctggt	400
tgtgattctt ataccccaaa agtcatacag tgtcagaaca aaggctggga	450
tgggtatgat gtacagtggg aatgtaagac ggacttagat attgcataca	500
aatttgaaa aactgtggtg agctgtgaag gctatgagtc ctctgaagac	550
cagtatgtac taagaggttc ttgtggcttg gagtataatt tagattatac	600
agaacttggc ctgcagaaac tgaaggagtc tggaaagcag cacggctttg	650
cctctttctc tgattattat tataagtgtt cctcggcggg ttctgtaac	700
atgagtggat tgattaccat cgtggtactc cttgggatcg cctttgtagt	750
ctataagctg ttctgagtg acgggcagta ttctcctcca ccgtactctg	800
agtatcctcc attttcccac cgttaccaga gattcaccaa ctcagcagga	850
cctcctccc caggctttaa gtctgagttc acaggaccac agaatactgg	900
ccatggtgca acttctgggt ttggcagtc ttttacagga caacaaggat	950
atgaaaattc aggaccaggg ttctgacag gcttgggaaac tggtggaata	1000
ctaggatatt tgtttggcag caatagagcg gcaacaccct tctcagactc	1050
gtggtactac ccgtcctatc ctcccctcta ccctggcacg tggaaatagg	1100
cttactcacc ccttcatgga ggctcgggca gctattcggg atgttcaaac	1150
tcagacacga aaaccagaac tgcacagga tatggtggtg ccaggagacg	1200
ataaagtaga aagttggagt caaacactgg atgcagaaat tttggatttt	1250
tcatacactt ctctttagaa aaaaagtact acctgttaac aattgggaaa	1300
aggggatatt caaaagtctt gtggtgttat gtccagtgta gctttttgta	1350
ttctattatt tgaggctaaa agttgatgtg tgacaaaata cttatgtgtt	1400
gtatgtcagt gtaacatgca gatgtatatt gcagtttttg aaagtatca	1450
ttactgtgga atgctaaaa tacattaatt tctaaaacct gtgatgccct	1500
aagaagcatt aagaatgaag gtgtgttact aatagaaact aagtacagaa	1550
aatttcagtt ttaggtggtt gtagctgatg agttattacc tcatagagac	1600
tataatattc tatttggtat tatattatth gatgtttgct gttcttcaaa	1650
catttaaatac aagctttgga ctaattatgc taatttga gttctgatca	1700
cttttgagct ctgaagcttt gaatcattca gtggtggaga tggccttctg	1750
gtaactgaat attaccttct gtaggaaaag gtgaaaata agcatctaga	1800
aggtgtgtgt gaatgactct gtgctggcaa aaatgctga aacctctata	1850
tttctttcgt tcataagagg taaaggctca atttttcaac aaaagtcttt	1900
taataacaaa agcatgcagt tctctgtgaa atctcaata ttgttgaat	1950
agtctgtttc aatcttaaaa agaataca	1977

&lt;210&gt; SEQ ID NO 46

&lt;211&gt; LENGTH: 339

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

-continued

&lt;400&gt; SEQUENCE: 46

---

```

Met Ala Ala Ala Cys Gly Pro Gly Ala Ala Gly Tyr Cys Leu Leu
 1           5           10
Leu Gly Leu His Leu Phe Leu Leu Thr Ala Gly Pro Ala Leu Gly
 20           25           30
Trp Asn Asp Pro Asp Arg Met Leu Leu Arg Asp Val Lys Ala Leu
 35           40           45
Thr Leu His Tyr Asp Arg Tyr Thr Thr Ser Arg Arg Leu Asp Pro
 50           55           60
Ile Pro Gln Leu Lys Cys Val Gly Gly Thr Ala Gly Cys Asp Ser
 65           70           75
Tyr Thr Pro Lys Val Ile Gln Cys Gln Asn Lys Gly Trp Asp Gly
 80           85           90
Tyr Asp Val Gln Trp Glu Cys Lys Thr Asp Leu Asp Ile Ala Tyr
 95           100          105
Lys Phe Gly Lys Thr Val Val Ser Cys Glu Gly Tyr Glu Ser Ser
 110          115          120
Glu Asp Gln Tyr Val Leu Arg Gly Ser Cys Gly Leu Glu Tyr Asn
 125          130          135
Leu Asp Tyr Thr Glu Leu Gly Leu Gln Lys Leu Lys Glu Ser Gly
 140          145          150
Lys Gln His Gly Phe Ala Ser Phe Ser Asp Tyr Tyr Tyr Lys Trp
 155          160          165
Ser Ser Ala Asp Ser Cys Asn Met Ser Gly Leu Ile Thr Ile Val
 170          175          180
Val Leu Leu Gly Ile Ala Phe Val Val Tyr Lys Leu Phe Leu Ser
 185          190          195
Asp Gly Gln Tyr Ser Pro Pro Pro Tyr Ser Glu Tyr Pro Pro Phe
 200          205          210
Ser His Arg Tyr Gln Arg Phe Thr Asn Ser Ala Gly Pro Pro Pro
 215          220          225
Pro Gly Phe Lys Ser Glu Phe Thr Gly Pro Gln Asn Thr Gly His
 230          235          240
Gly Ala Thr Ser Gly Phe Gly Ser Ala Phe Thr Gly Gln Gln Gly
 245          250          255
Tyr Glu Asn Ser Gly Pro Gly Phe Trp Thr Gly Leu Gly Thr Gly
 260          265          270
Gly Ile Leu Gly Tyr Leu Phe Gly Ser Asn Arg Ala Ala Thr Pro
 275          280          285
Phe Ser Asp Ser Trp Tyr Tyr Pro Ser Tyr Pro Pro Ser Tyr Pro
 290          295          300
Gly Thr Trp Asn Arg Ala Tyr Ser Pro Leu His Gly Gly Ser Gly
 305          310          315
Ser Tyr Ser Val Cys Ser Asn Ser Asp Thr Lys Thr Arg Thr Ala
 320          325          330
Ser Gly Tyr Gly Gly Thr Arg Arg Arg
 335

```

&lt;210&gt; SEQ ID NO 47

&lt;211&gt; LENGTH: 1766

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapien

-continued

---

<400> SEQUENCE: 47

cccggagccg gggagggagg gagcgagggt cggacaccgg cggcggctgc	50
ctggcctttc catgagcccg cggcggaccc tcccgcgccc cctctcgctc	100
tgccctctcc tctgcctctg cctctgcctg gcccgggctc tgggaagtgc	150
gcagtcocgg tcgtgtaggg ataaaaagaa ctgtaagggt gtcttttccc	200
agcaggaact gaggaagcgg ctaacacccc tgcagtacca tgtaactcag	250
gagaaagga ccgaaagtgc ctttgaagga gaatacacac atcacaaga	300
tcttgaata tataaatgtg ttgtttgtgg aactccattg ttttaagtca	350
aaaccaaatt tgactccggt tcaggttggc cttcattcca cgatgtgatc	400
aattctgagg caatcacatt cacagatgac ttttctatg ggatgcacag	450
ggtgaaaca agctgctctc agtgtgtgct tcacctggg cacatttttg	500
atgatgggccc tcgtccaact gggaaaagat actgcataaa ttcggctgccc	550
ttgtctttta cacctgcgga tagcagtggc accgccgagg gaggcagtgg	600
ggtcgcacag ccggcccagg cagacaaaag ggagctctag agtaatggag	650
agtgatggaa acaaagtgta cttaatgac agcttattaa aaaaatcaaa	700
attgttatct taatagatat attttttcaa aaactataag ggcagttttg	750
tgctattgat attttttctt cttttgctta aacagaagcc ctggccatcc	800
atgtattttg caattgacta gatcaagaac tgtttatagc tttagcaaat	850
ggagacagct ttgtgaaact tcttcacaag ccacttatac cctttggcat	900
tcttttcttt gagcacatgg cttcttttgc agtttttccc cctttgattc	950
agaagcagag ggttcatggt cttcaaacat gaaaatagag atctcctctg	1000
cagtgtagag accagagctg ggcagtgcag ggcatggaga cctgcaagac	1050
acatggcctt gaggcctttg cacagaccca cctaagataa ggttgagtg	1100
atgttttaat gagactgttc agctttgtgg aaagtttgag ctaaggtcat	1150
tttttttttt ctactgaaa ggggtgtaag gtctaaagtc tttccttatg	1200
ttaaattggt gccagatcca aaggggcata ctgagtgttg tggcagagaa	1250
gtaaacatta ccacactggt aggcctttat tttattttat tttccatcga	1300
aagcattgga ggcccagtgc aatggctcac gcctgtgatc ccagcacttt	1350
gggaggccaa ggcgggtgga tcacgaggtc aggagatgga gaccatcctg	1400
gctaactcgg tgaaccccg tctctactaa aaatacga aattagccag	1450
gcgtggtggt gggcacctgt agtcccagct actcaggagg ctgaggcagg	1500
agaatggcgt gaacccggaa ggcggagctt gcagttagcc gagatcatgc	1550
cactgcactc cagcctacat gacaatgtga cactccatct caaaaaataa	1600
taataataac aatataagaa ctagctgggc atggtggcgc atgcatgtag	1650
tcccagctac tcctgaggct cagtcaggag aatcgcttga acttggagg	1700
cggaggttgc agtgagctga gctcatacca ctgcactcca gcctgaacag	1750
agtgagatcc tgtcaa	1766

-continued

<210> SEQ ID NO 48  
 <211> LENGTH: 192  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 48

Met Ser Pro Arg Arg Thr Leu Pro Arg Pro Leu Ser Leu Cys Leu  
 1 5 10 15  
 Ser Leu Cys Leu Cys Leu Cys Leu Ala Ala Ala Leu Gly Ser Ala  
 20 25 30  
 Gln Ser Gly Ser Cys Arg Asp Lys Lys Asn Cys Lys Val Val Phe  
 35 40 45  
 Ser Gln Gln Glu Leu Arg Lys Arg Leu Thr Pro Leu Gln Tyr His  
 50 55 60  
 Val Thr Gln Glu Lys Gly Thr Glu Ser Ala Phe Glu Gly Glu Tyr  
 65 70 75  
 Thr His His Lys Asp Pro Gly Ile Tyr Lys Cys Val Val Cys Gly  
 80 85 90  
 Thr Pro Leu Phe Lys Ser Glu Thr Lys Phe Asp Ser Gly Ser Gly  
 95 100 105  
 Trp Pro Ser Phe His Asp Val Ile Asn Ser Glu Ala Ile Thr Phe  
 110 115 120  
 Thr Asp Asp Phe Ser Tyr Gly Met His Arg Val Glu Thr Ser Cys  
 125 130 135  
 Ser Gln Cys Gly Ala His Leu Gly His Ile Phe Asp Asp Gly Pro  
 140 145 150  
 Arg Pro Thr Gly Lys Arg Tyr Cys Ile Asn Ser Ala Ala Leu Ser  
 155 160 165  
 Phe Thr Pro Ala Asp Ser Ser Gly Thr Ala Glu Gly Gly Ser Gly  
 170 175 180  
 Val Ala Ser Pro Ala Gln Ala Asp Lys Ala Glu Leu  
 185 190

<210> SEQ ID NO 49  
 <211> LENGTH: 2065  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 49

cccaaagagg tgaggagccg gcagcggggg cggctgtaac tgtgaggaag 50  
 gctgcagagt ggcgacgtct acgccgtagg ttggaggctg tgggggggtg 100  
 ccggggcgcca gctcccaggc cgcagaagtg acctgcggtg gagttccctc 150  
 ctcgctgctg gagaacggag ggagaaggtt gctggccggg tgaagtgcc 200  
 tccctctgct tgacggggct gagggcccg aagtctaggg cgtccgtagt 250  
 cgccccggcc tccgtgaagc cccaggtcta gagatatgac ccgagagtgc 300  
 ccatctccgg ccccggggcc tggggctccg ctgagtggat cgggtctggc 350  
 agaggcggca gtagtgtttg cagtgtgct gagcatccac gcaaccgtat 400  
 gggaccgata ctcggtgtgc gcogtggccc togcagtgca ggcottctac 450  
 gtccaataca agtgggaccg gctgctacag caggaagcg ccgtcttcca 500  
 gttccgaatg tccgcaaaca gtggcctatt gcccgctcc atggtcatgc 550

-continued

---

```

ctttgcttgg actagtcatg aaggagcggg gccagactgc tgggaacccg      600
ttctttgagc gttttggcat tgtggtggca gccactggca tggcagtggc      650
cctctttcoa tcagtgttgg cgctcggcat cactcgccca gtgccaacca      700
acacttgtgt catcttgggc ttggctggag gtgttatcat ttatatcatg      750
aagcactcgt tgagcgtggg ggaggtgatc gaagtcctgg aagtccttct      800
gatcttctgt tatctcaaca tgatcctgct gtacctgctg ccccgctgct      850
tcacccctgg tgaggcactg ctggtattgg gtggcattag ctttgcctc      900
aaccagctca tcaagcgctc tctgacactg gtgaaaagtc agggggaccc      950
agtggacttc ttctctgctg tgggtgtagt agggatgta ctcatgggca     1000
ttttcttcag cactctgttt gtcttcatgg actcaggcac ctgggcctcc     1050
tccatcttct tccacctcat gacctgtgtg ctgagccttg gtgtggtcct     1100
accctggctg caccggtcca tccgcaggaa tcccctgctc tggcttcttc     1150
agtttctctt ccagacagac acccgcatct acctctagc ctattggtct     1200
ctgctggcca ccttggcctg cctggtggtg ctgtaccaga atgccaagcg     1250
gtcatcttcc gagtccaaga agcaccaggc ccccaccatc gcccgaaagt     1300
atctccacct cattgtggta gccacctaca tcccaggtat catctttgac     1350
cggcactcgc tctatgtagc cgccactgta tgcctggcgg tcttcatctt     1400
cctggagtat gtgcgctact tccgcatcaa gcctttgggt cacactctac     1450
ggagcttctc gtcccttttt ctggatgaac gagacagtgg accactcatt     1500
ctgacacaca tctacctgct cctgggcatg tctcttccca tctggctgat     1550
ccccagacct tgcacacaga agggtagcct gggaggagcc agggccctcg     1600
tcccctatgc cgggtgcctg gctgtgggtg tgggtgatac tgtggcctcc     1650
atctctggta gcacatggg ggagatccgc tggcctggaa ccaaaaagac     1700
ttttgagggg accatgacat ctatatattg gcagatcatt tctgtagctc     1750
tgatcttaat ctttgacagt ggagtggacc taaactacag ttatgcttgg     1800
atcttggggg ccatcagcac tgtgtccctc ctggaagcat aactacaca     1850
gatagacaa ctccttctgc ctctctacct cctgatattg ctgatggcct     1900
agctgttaca gtgcagcagc agtgacggag gaaacagaca tggggagggt     1950
gaacagtccc cacagcagac agctacttgg gcatgaagag ccaaggtgtg     2000
aaaagcagat ttgattttc agttgattca gatttaaaat aaaaagcaaa     2050
gctctcctag ttcta      2065

```

```

<210> SEQ ID NO 50
<211> LENGTH: 538
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

```

<400> SEQUENCE: 50

```

Met Thr Arg Glu Cys Pro Ser Pro Ala Pro Gly Pro Gly Ala Pro
 1           5           10          15
Leu Ser Gly Ser Val Leu Ala Glu Ala Ala Val Val Phe Ala Val
          20          25          30

```



-continued

Val	Leu	Ser	Ile	His	Ala	Thr	Val	Trp	Asp	Arg	Tyr	Ser	Trp	Cys	35	40	45
Ala	Val	Ala	Leu	Ala	Val	Gln	Ala	Phe	Tyr	Val	Gln	Tyr	Lys	Trp	50	55	60
Asp	Arg	Leu	Leu	Gln	Gln	Gly	Ser	Ala	Val	Phe	Gln	Phe	Arg	Met	65	70	75
Ser	Ala	Asn	Ser	Gly	Leu	Leu	Pro	Ala	Ser	Met	Val	Met	Pro	Leu	80	85	90
Leu	Gly	Leu	Val	Met	Lys	Glu	Arg	Cys	Gln	Thr	Ala	Gly	Asn	Pro	95	100	105
Phe	Phe	Glu	Arg	Phe	Gly	Ile	Val	Val	Ala	Ala	Thr	Gly	Met	Ala	110	115	120
Val	Ala	Leu	Phe	Ser	Ser	Val	Leu	Ala	Leu	Gly	Ile	Thr	Arg	Pro	125	130	135
Val	Pro	Thr	Asn	Thr	Cys	Val	Ile	Leu	Gly	Leu	Ala	Gly	Gly	Val	140	145	150
Ile	Ile	Tyr	Ile	Met	Lys	His	Ser	Leu	Ser	Val	Gly	Glu	Val	Ile	155	160	165
Glu	Val	Leu	Glu	Val	Leu	Leu	Ile	Phe	Val	Tyr	Leu	Asn	Met	Ile	170	175	180
Leu	Leu	Tyr	Leu	Leu	Pro	Arg	Cys	Phe	Thr	Pro	Gly	Glu	Ala	Leu	185	190	195
Leu	Val	Leu	Gly	Gly	Ile	Ser	Phe	Val	Leu	Asn	Gln	Leu	Ile	Lys	200	205	210
Arg	Ser	Leu	Thr	Leu	Val	Glu	Ser	Gln	Gly	Asp	Pro	Val	Asp	Phe	215	220	225
Phe	Leu	Leu	Val	Val	Val	Val	Gly	Met	Val	Leu	Met	Gly	Ile	Phe	230	235	240
Phe	Ser	Thr	Leu	Phe	Val	Phe	Met	Asp	Ser	Gly	Thr	Trp	Ala	Ser	245	250	255
Ser	Ile	Phe	Phe	His	Leu	Met	Thr	Cys	Val	Leu	Ser	Leu	Gly	Val	260	265	270
Val	Leu	Pro	Trp	Leu	His	Arg	Leu	Ile	Arg	Arg	Asn	Pro	Leu	Leu	275	280	285
Trp	Leu	Leu	Gln	Phe	Leu	Phe	Gln	Thr	Asp	Thr	Arg	Ile	Tyr	Leu	290	295	300
Leu	Ala	Tyr	Trp	Ser	Leu	Leu	Ala	Thr	Leu	Ala	Cys	Leu	Val	Val	305	310	315
Leu	Tyr	Gln	Asn	Ala	Lys	Arg	Ser	Ser	Ser	Glu	Ser	Lys	Lys	His	320	325	330
Gln	Ala	Pro	Thr	Ile	Ala	Arg	Lys	Tyr	Phe	His	Leu	Ile	Val	Val	335	340	345
Ala	Thr	Tyr	Ile	Pro	Gly	Ile	Ile	Phe	Asp	Arg	Pro	Leu	Leu	Tyr	350	355	360
Val	Ala	Ala	Thr	Val	Cys	Leu	Ala	Val	Phe	Ile	Phe	Leu	Glu	Tyr	365	370	375
Val	Arg	Tyr	Phe	Arg	Ile	Lys	Pro	Leu	Gly	His	Thr	Leu	Arg	Ser	380	385	390
Phe	Leu	Ser	Leu	Phe	Leu	Asp	Glu	Arg	Asp	Ser	Gly	Pro	Leu	Ile	395	400	405
Leu	Thr	His	Ile	Tyr	Leu	Leu	Leu	Gly	Met	Ser	Leu	Pro	Ile	Trp			

-continued

	410		415		420
Leu Ile Pro Arg Pro Cys Thr Gln Lys Gly Ser Leu Gly Gly Ala	425		430		435
Arg Ala Leu Val Pro Tyr Ala Gly Val Leu Ala Val Gly Val Gly	440		445		450
Asp Thr Val Ala Ser Ile Phe Gly Ser Thr Met Gly Glu Ile Arg	455		460		465
Trp Pro Gly Thr Lys Lys Thr Phe Glu Gly Thr Met Thr Ser Ile	470		475		480
Phe Ala Gln Ile Ile Ser Val Ala Leu Ile Leu Ile Phe Asp Ser	485		490		495
Gly Val Asp Leu Asn Tyr Ser Tyr Ala Trp Ile Leu Gly Ser Ile	500		505		510
Ser Thr Val Ser Leu Leu Glu Ala Tyr Thr Thr Gln Ile Asp Asn	515		520		525
Leu Leu Leu Pro Leu Tyr Leu Leu Ile Leu Leu Met Ala	530		535		

<210> SEQ ID NO 51  
 <211> LENGTH: 3476  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 51

```

gctctatgcc gcctaccttg ctctcgccgc tgctgcccga gccgaagcag      50
agaaggcagc gggccccgtg accgtcccga gagccccgcg ctcccacca      100
gggggcccgg gcggccccgg ggaggccggg gcaggggcccgg ggggaagaaa    150
gggggttttg tgctgcgccg ggaggcccg cgccctcttc cgaatgtcct    200
gcggccccag cctctcctca cgctcgcgca gtctccgccg cagtctcagc    250
tgacagtgca ggactgagcc gtgcaccggg aggagacccc cggaggaggc    300
gacaaacttc gcagtgccgc gaccacaacc cagccctggg tagcctgcag    350
catggcccag ctgttcctgc cctgctggc agccctggtc ctggcccagg    400
ctctgcagc tttagcagat gttctggaag gagacagctc agaggaccgc    450
gcttttcgag tgcgcatcgc ggggcacggc ccaactgcagg gcgtgctcgg    500
cggcgccctc accatccctt gccacgtcca ctacctgcgg caaccgccga    550
gccgcccggc tgtgctgggc tctccgccc tcaagtggac tttcctgtcc    600
cggggcccgg aggcagaggt gctggtggcg cggggagtgc gcgtcaaggt    650
gaacgaggcc taccggttcc gcgtggcact gcctgcgtac ccagcgtcgc    700
tcaccgacgt ctccctggcg ctgagcgagc tgcgcccaca cgactcaggt    750
atctatcgct gtgaggtcca gcacggcacc gatgacagca gcgacgctgt    800
ggaggtcaag gtcaaagggg tcgtctttct ctaccgagag ggetctgccc    850
gctatgcttt ctctttttct ggggcccagg aggctgtgc ccgcattgga    900
gccacatog ccaccccga gcagctctat gccgcctacc ttgggggcta    950
tgagcaatgt gatgctggct ggctgtcggg tcagaccgtg aggtatocca   1000
tcacagcccc acgagaggcc tgttacggag acatggatgg cttcccggg    1050
    
```

-continued

---

gtccggaact atggtgtggt ggacccggat gacctctatg atgtgtactg	1100
ttatgctgaa gacctaaatg gagaactgtt cctgggtgac cctccagaga	1150
agctgacatt ggaggaagca cgggcgtact gccaggagcg ggggagag	1200
attgccacca cgggccaact gtatgcagcc tgggatggtg gcctggacca	1250
ctgcagccca ggggtggctag ctgatggcag tgtgcgctac cccatcgtca	1300
caccagccca gcgctgtggt gggggcttgc ctggtgtcaa gactctcttc	1350
ctcttcccca accagactgg ctccccaat aagcacagcc gcttcaacgt	1400
ctactgcttc cgagactcgg cccagccttc tgccatcctt gaggcctcca	1450
accagcctc caaccagcc tctgatggac tagaggctat cgtcacagt	1500
acagagacc tggaggaact gcagctgcct caggaagcca cagagagtga	1550
atcccgtggg gccatctact ccatcccat catggaggac ggaggagggtg	1600
gaagctccc tccagaagac ccagcagagg cccctagac gctcctagaa	1650
tttgaaac aatccatggt accgcccacg gggttctcag aagaggaagg	1700
taaggcattg gaggaagaag aaaaatata agatgaagaa gagaaagagg	1750
aggaagaaga agaggaggag gtggaggatg aggctctgtg ggcattggccc	1800
agcgagctca gcagcccggg ccctgaggcc tctctcccca ctgagccagc	1850
agcccaggag aagtcaact cccagcgcgc agcaaggca gtcctgcagc	1900
ctgggtgcat accacttctt gatggagagt cagaagcttc caggcctcca	1950
agggtcctat gaccacctac tgagactctg cccactccca gggagaggaa	2000
cctagcatcc ccatcacctt ccaactctgt tgaggcaaga gagggtggggg	2050
aggaactgg tggctcctgag ctatctgggg tccctcgagg agagagcgag	2100
gagacaggaa gctccgaggg tgccccttc ctgctccag ccacacgggc	2150
ccctgagggt accaggagc tggaggcccc ctctgaagat aattctggaa	2200
gaactgcccc agcagggacc tcagtgcagg cccagccagt gctgcccact	2250
gacagcgcca gccgaggtg agtggccgtg gtccccgat cagggtgactg	2300
tgtccccagc ccctgccaca atggtgggac atgcttgag gaggaggag	2350
gggtccgctg cctatgtctg cctggctatg gggggacct gtgcgatgtt	2400
ggcctccgct tctgcaacc cggctgggac gccttccag gcgcctgcta	2450
caagcacttt tccacacgaa ggagctggga ggaggcagag acccagtgcc	2500
ggatgtacgg cgcgcatctg gccagcatca gcacaccga ggaacaggac	2550
ttcatcaaca accggtaccg ggagtaccag tggatcggac tcaacgacag	2600
gaccatcgaa ggcgacttct tgtggtcggg tggcgtcccc ctgctctatg	2650
agaactggaa ccctgggag cctgacagct acttctgtc tggagagAAC	2700
tgcgtgtgca tgggtgtggca tgatcaggga caatggagt acgtgcctctg	2750
caactaccac ctgtcctaca cctgcaagat ggggctggtg tctgtgggc	2800
cgccaccgga gctgcccctg gctcaagtgt toggccgcc acggctgcgc	2850
tatgaggtgg aactgtgct tcgctaccgg tgccgggag gactggccca	2900
gcgcaactct cgcctgatcc gatgccaaga gaacggctct tgggaggccc	2950

-continued

---

```

cccagatctc ctgtgtgcc agaagacctg cccgagctct gcaccagag      3000
gaggaccag aaggacgtca ggggaggcta ctgggacgct ggaaggcgct      3050
gttgatcccc ccttccagcc ccatgccagg tccctagggg gcaaggcctt      3100
gaacactgcc gcccacagca ctgccctgtc acccaaattt tccctcacac      3150
cttgcgctcc cgccaccaca ggaagtgaca acatgacgag gggtggtgct      3200
ggagtccagg tgacagttcc tgaaggggct tctgggaaat acctaggagg      3250
ctccagccca gccccaggccc tctcccccta cctggggcac cagatcttcc      3300
atcagggccg gagtaaatcc ctaagtgcct caactgcctt ctccctggca      3350
gccatcttgt ccctctatt cctctagga gcaactgtcc cactctttt      3400
gggttttcca agggaatggg cttgcaggat ggagtgtctg taaatcaac      3450
aggaataaaa actgtgtatg agccca      3476

```

&lt;210&gt; SEQ ID NO 52

&lt;211&gt; LENGTH: 911

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 52

```

Met Ala Gln Leu Phe Leu Pro Leu Leu Ala Ala Leu Val Leu Ala
 1           5           10          15
Gln Ala Pro Ala Ala Leu Ala Asp Val Leu Glu Gly Asp Ser Ser
          20          25          30
Glu Asp Arg Ala Phe Arg Val Arg Ile Ala Gly Asp Ala Pro Leu
          35          40          45
Gln Gly Val Leu Gly Gly Ala Leu Thr Ile Pro Cys His Val His
          50          55          60
Tyr Leu Arg Pro Pro Pro Ser Arg Arg Ala Val Leu Gly Ser Pro
          65          70          75
Arg Val Lys Trp Thr Phe Leu Ser Arg Gly Arg Glu Ala Glu Val
          80          85          90
Leu Val Ala Arg Gly Val Arg Val Lys Val Asn Glu Ala Tyr Arg
          95          100         105
Phe Arg Val Ala Leu Pro Ala Tyr Pro Ala Ser Leu Thr Asp Val
          110         115         120
Ser Leu Ala Leu Ser Glu Leu Arg Pro Asn Asp Ser Gly Ile Tyr
          125         130         135
Arg Cys Glu Val Gln His Gly Ile Asp Asp Ser Ser Asp Ala Val
          140         145         150
Glu Val Lys Val Lys Gly Val Val Phe Leu Tyr Arg Glu Gly Ser
          155         160         165
Ala Arg Tyr Ala Phe Ser Phe Ser Gly Ala Gln Glu Ala Cys Ala
          170         175         180
Arg Ile Gly Ala His Ile Ala Thr Pro Glu Gln Leu Tyr Ala Ala
          185         190         195
Tyr Leu Gly Gly Tyr Glu Gln Cys Asp Ala Gly Trp Leu Ser Asp
          200         205         210
Gln Thr Val Arg Tyr Pro Ile Gln Thr Pro Arg Glu Ala Cys Tyr
          215         220         225
Gly Asp Met Asp Gly Phe Pro Gly Val Arg Asn Tyr Gly Val Val

```

-continued

										230											235											240				
Asp	Pro	Asp	Asp	Leu	Tyr	Asp	Val	Tyr	Cys	Tyr	Ala	Glu	Asp	Leu	245	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Asn	Gly	Glu	Leu	Phe	Leu	Gly	Asp	Pro	Pro	Glu	Lys	Leu	Thr	Leu	260	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Glu	Glu	Ala	Arg	Ala	Tyr	Cys	Gln	Glu	Arg	Gly	Ala	Glu	Ile	Ala	275	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Thr	Thr	Gly	Gln	Leu	Tyr	Ala	Ala	Trp	Asp	Gly	Gly	Leu	Asp	His	290	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Cys	Ser	Pro	Gly	Trp	Leu	Ala	Asp	Gly	Ser	Val	Arg	Tyr	Pro	Ile	305	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Val	Thr	Pro	Ser	Gln	Arg	Cys	Gly	Gly	Gly	Leu	Pro	Gly	Val	Lys	320	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Thr	Leu	Phe	Leu	Phe	Pro	Asn	Gln	Thr	Gly	Phe	Pro	Asn	Lys	His	335	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Ser	Arg	Phe	Asn	Val	Tyr	Cys	Phe	Arg	Asp	Ser	Ala	Gln	Pro	Ser	350	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Ala	Ile	Pro	Glu	Ala	Ser	Asn	Pro	Ala	Ser	Asn	Pro	Ala	Ser	Asp	365	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Gly	Leu	Glu	Ala	Ile	Val	Thr	Val	Thr	Glu	Thr	Leu	Glu	Glu	Leu	380	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Gln	Leu	Pro	Gln	Glu	Ala	Thr	Glu	Ser	Glu	Ser	Arg	Gly	Ala	Ile	395	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Tyr	Ser	Ile	Pro	Ile	Met	Glu	Asp	Gly	Gly	Gly	Gly	Ser	Ser	Thr	410	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Pro	Glu	Asp	Pro	Ala	Glu	Ala	Pro	Arg	Thr	Leu	Leu	Glu	Phe	Glu	425	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Thr	Gln	Ser	Met	Val	Pro	Pro	Thr	Gly	Phe	Ser	Glu	Glu	Glu	Gly	440	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Lys	Ala	Leu	Glu	Glu	Glu	Glu	Lys	Tyr	Glu	Asp	Glu	Glu	Glu	Lys	455	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Glu	Glu	Glu	Glu	Glu	Glu	Glu	Glu	Val	Glu	Asp	Glu	Ala	Leu	Trp	470	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Ala	Trp	Pro	Ser	Glu	Leu	Ser	Ser	Pro	Gly	Pro	Glu	Ala	Ser	Leu	485	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Pro	Thr	Glu	Pro	Ala	Ala	Gln	Glu	Lys	Ser	Leu	Ser	Gln	Ala	Pro	500	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Ala	Arg	Ala	Val	Leu	Gln	Pro	Gly	Ala	Ser	Pro	Leu	Pro	Asp	Gly	515	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Glu	Ser	Glu	Ala	Ser	Arg	Pro	Pro	Arg	Val	His	Gly	Pro	Pro	Thr	530	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Glu	Thr	Leu	Pro	Thr	Pro	Arg	Glu	Arg	Asn	Leu	Ala	Ser	Pro	Ser	545	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Pro	Ser	Thr	Leu	Val	Glu	Ala	Arg	Glu	Val	Gly	Glu	Ala	Thr	Gly	560	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Gly	Pro	Glu	Leu	Ser	Gly	Val	Pro	Arg	Gly	Glu	Ser	Glu	Glu	Thr	575	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Gly	Ser	Ser	Glu	Gly	Ala	Pro	Ser	Leu	Leu	Pro	Ala	Thr	Arg	Ala	590	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270
Pro	Glu	Gly	Thr	Arg	Glu	Leu	Glu	Ala	Pro	Ser	Glu	Asp	Asn	Ser	605	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270	Pro	Glu	Lys	Leu	Thr	Leu	270

-continued

---

Gly Arg Thr Ala Pro Ala Gly Thr Ser Val Gln Ala Gln Pro Val  
 620 625 630

Leu Pro Thr Asp Ser Ala Ser Arg Gly Gly Val Ala Val Val Pro  
 635 640 645

Ala Ser Gly Asp Cys Val Pro Ser Pro Cys His Asn Gly Gly Thr  
 650 655 660

Cys Leu Glu Glu Glu Gly Val Arg Cys Leu Cys Leu Pro Gly  
 665 670 675

Tyr Gly Gly Asp Leu Cys Asp Val Gly Leu Arg Phe Cys Asn Pro  
 680 685 690

Gly Trp Asp Ala Phe Gln Gly Ala Cys Tyr Lys His Phe Ser Thr  
 695 700 705

Arg Arg Ser Trp Glu Glu Ala Glu Thr Gln Cys Arg Met Tyr Gly  
 710 715 720

Ala His Leu Ala Ser Ile Ser Thr Pro Glu Glu Gln Asp Phe Ile  
 725 730 735

Asn Asn Arg Tyr Arg Glu Tyr Gln Trp Ile Gly Leu Asn Asp Arg  
 740 745 750

Thr Ile Glu Gly Asp Phe Leu Trp Ser Asp Gly Val Pro Leu Leu  
 755 760 765

Tyr Glu Asn Trp Asn Pro Gly Gln Pro Asp Ser Tyr Phe Leu Ser  
 770 775 780

Gly Glu Asn Cys Val Val Met Val Trp His Asp Gln Gly Gln Trp  
 785 790 795

Ser Asp Val Pro Cys Asn Tyr His Leu Ser Tyr Thr Cys Lys Met  
 800 805 810

Gly Leu Val Ser Cys Gly Pro Pro Pro Glu Leu Pro Leu Ala Gln  
 815 820 825

Val Phe Gly Arg Pro Arg Leu Arg Tyr Glu Val Asp Thr Val Leu  
 830 835 840

Arg Tyr Arg Cys Arg Glu Gly Leu Ala Gln Arg Asn Leu Pro Leu  
 845 850 855

Ile Arg Cys Gln Glu Asn Gly Arg Trp Glu Ala Pro Gln Ile Ser  
 860 865 870

Cys Val Pro Arg Arg Pro Ala Arg Ala Leu His Pro Glu Glu Asp  
 875 880 885

Pro Glu Gly Arg Gln Gly Arg Leu Leu Gly Arg Trp Lys Ala Leu  
 890 895 900

Leu Ile Pro Pro Ser Ser Pro Met Pro Gly Pro  
 905 910

<210> SEQ ID NO 53  
 <211> LENGTH: 3316  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 53

ctgccagggtg acagccgcca agatggggctg ttggggccctg ctgtggcctc	50
cctgtctgtt caccgggctg ctgctccgac ccccggggac catggcccag	100
gcccagtaact gctctgtgaa caaggacatc tttgaagtag aggagaacac	150
aaatgtcacc gagccgctgg tggacatcca cgtcccggag ggcaggagag	200

-continued

---

tgaccctcgg agccttgtcc accccctttg catttcggat ccagggaaac	250
cagctgtttc tcaacgtgac tcctgattac gaggagaagt cactgcttga	300
ggctcagctg ctgtgtcaga gcggaggcac attggtgacc cagctaaggg	350
tgttcgtgtc agtgctggac gtcaatgaca atgccccga attccccttt	400
aagaccaagg agataagggt ggaggaggac acgaaagtga actccaccgt	450
catccctgag acgcaactgc aggctgagga ccgacgacaag gacgacattc	500
tgttctacac cctccaggaa atgacagcag gtgccagtga ctacttctcc	550
ctggtgagtg taaaccgtcc cgcctgagg ctggaccggc ccctggactt	600
ctacgagcgg ccgaacatga cttctggct gctggtgctg gacactccag	650
gggagaatgt ggaaccacgc cacactgcca ccgccacact agtgctgaac	700
gtggtgcccg ccgacctgcg gcccccgtgg ttccctgcct gcaacctctc	750
agatggctac gtctgcattc aagctcagta ccacggggct gtccccacgg	800
ggcacatact gccatctccc ctgctctgct gtcccggacc catctacgct	850
gaggacggag accgcgcat caaccagccc atcatctaca gcatctttag	900
gggaaacgtg aatggtacat tcatcatcca cccagactcg ggcaacctca	950
ccgtggccag gagtgtcccc agcccataga ccttctctct gctggtgaag	1000
ggccaacagc ccgacctgac ccgctactca gtgaccagc tcaccgtgga	1050
ggctgtgctg gcggccggga gcccccggc cttccccag agcctgtatc	1100
gtggcacctg ggccgctggc gctggagcgg gcggtgtggt caaggatgca	1150
gctgcccctt ctacgctctt gaggatccag gctcaggacc cggagtctc	1200
ggacctcaac tcggccatca catatcgaat taccaaccac tcacacttcc	1250
ggatggaggg agaggttggt ctgaccacca ccacactggc acaggcggga	1300
gccttctacg cagaggttga ggcccacaac acggtgacct ctggcaaccg	1350
aaccacagtc attgagatac aagtttccga acaggagccc ccctccacag	1400
aggctggagg aacaactggg ccctggacca gcaccacttc cgaggctccc	1450
agaccccctg agccctcca gggaccctcc acgaccagct ctgggggagg	1500
cacaggccct catccaccct ctggcacaac tctgaggcca ccaacctcgt	1550
ccacaccggg ggggccccg ggtgcagaaa acagcacctc ccaccaacca	1600
gccactcccc gtggggacac agcacagacc ccaagccag gaacctctca	1650
gccgatgcc cccggtgtgg gaaccagcac ctcccacaa ccagccacac	1700
ccagtggggg cacagcacag accccagagc caggaaacct tcagccgatg	1750
ccccccagta tgggaaccag caactccac caaccagcca caccgggtgg	1800
gggcacagca cagaccccag aggcaggaac ctctcagccg atgcccccg	1850
gtatgggaac cagcacctcc caccaaccaa ccacaccggg tgggggcaca	1900
gcacagacc cagagccagg aacctctcag ccgatgcccc tcagcaagag	1950
cacccatct tcaggtggcg gccctcggga ggacaagcgc ttctcgggtg	2000
tggatatggc gccctgggc ggggtgctgg gtgcgctgct gctgctggct	2050
ctccttgccc tcgcccctct tgtccacaag cactatggcc cccggtcaa	2100

-continued

---

```

gtgctgctct ggcaaagctc cggagcccca gccccaaggc ttgacaacc      2150
agcggttctc ccctgaccac aaggccaact gggcgcccgt cccagcccc      2200
acgcacgacc ccaagcccgc ggaggcaccc atgcccgcag agcccgcacc      2250
ccccggccct gcctccccag gcggtgcccc tgagccccc gcagcggccc      2300
gagctggcgg aagccccacg gcggtgaggt ccctcctgac caaggagcgg      2350
cggccggagg gcggttacia gccctctggt ttggcgagg acatcgggac      2400
ggaggcagac gtggtcgttc tcaacgcgcc caccctggac gtggatggcg      2450
ccagtgactc cggcagcggc gacgagggcg agggcgcggg gaggggtggg      2500
ggtccctacg atgcaccggg tggatgatgac tcctacatct aagtggcccc      2550
tccaccctct ccccagccc caccgggact ggaggtctcg ctccccagc      2600
ctccgaccgg aggcagaata aagcaaggct ccgaaaccc aggccatggc      2650
gtggggcagg cgcgtgggtc cctgggggccc ccattcactc agtcccctgt      2700
cgtcattagc gcttgagccc aggtgtgcag atgaggcggg gggctctggc      2750
acgctgtccc caccccaagg ctgcagcact tcccgtaac cacctgcagt      2800
gcccggccc ttcccaggc tctgtgccag ctagtctggg aagtccctct      2850
cccgtctaa ccacagccc aggggggctc ccctcccccg acctgcacca      2900
gagatctcag gcacccggct caactcagac ctcccgtccc cgaccctaca      2950
cagagattgc ctggggaggc tgaggagccg atgcaaaccc ccaaggcgac      3000
gcacttggga gccggtggtc tcaaacacct gccggggggtc ctagtcccct      3050
tctgaaatct acatgcttgg gttggagcgc agcagtaaac accctgccca      3100
gtgacctgga ctgaggcgcg ctgggggtgg gtgcgccgtg tggcctgagc      3150
aggagccaga ccaggaggcc taggggtgag agacacattc ccctcgtgc      3200
tcccaaagcc agagcccagg ctgggcgccc atgccagaa ccatcaaggg      3250
atcccttgcg gcttgtcagc actttcccta atgaaatac accattaatt      3300
cctttccaaa tgtttt      3316

```

&lt;210&gt; SEQ ID NO 54

&lt;211&gt; LENGTH: 839

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 54

```

Met Gly Ser Trp Ala Leu Leu Trp Pro Pro Leu Leu Phe Thr Gly
 1           5           10          15
Leu Leu Val Arg Pro Pro Gly Thr Met Ala Gln Ala Gln Tyr Cys
          20          25          30
Ser Val Asn Lys Asp Ile Phe Glu Val Glu Glu Asn Thr Asn Val
          35          40          45
Thr Glu Pro Leu Val Asp Ile His Val Pro Glu Gly Gln Glu Val
          50          55          60
Thr Leu Gly Ala Leu Ser Thr Pro Phe Ala Phe Arg Ile Gln Gly
          65          70          75
Asn Gln Leu Phe Leu Asn Val Thr Pro Asp Tyr Glu Glu Lys Ser
          80          85          90

```



-continued

---

Leu	Leu	Glu	Ala	Gln	Leu	Leu	Cys	Gln	Ser	Gly	Gly	Thr	Leu	Val
				95					100					105
Thr	Gln	Leu	Arg	Val	Phe	Val	Ser	Val	Leu	Asp	Val	Asn	Asp	Asn
				110					115					120
Ala	Pro	Glu	Phe	Pro	Phe	Lys	Thr	Lys	Glu	Ile	Arg	Val	Glu	Glu
				125					130					135
Asp	Thr	Lys	Val	Asn	Ser	Thr	Val	Ile	Pro	Glu	Thr	Gln	Leu	Gln
				140					145					150
Ala	Glu	Asp	Arg	Asp	Lys	Asp	Asp	Ile	Leu	Phe	Tyr	Thr	Leu	Gln
				155					160					165
Glu	Met	Thr	Ala	Gly	Ala	Ser	Asp	Tyr	Phe	Ser	Leu	Val	Ser	Val
				170					175					180
Asn	Arg	Pro	Ala	Leu	Arg	Leu	Asp	Arg	Pro	Leu	Asp	Phe	Tyr	Glu
				185					190					195
Arg	Pro	Asn	Met	Thr	Phe	Trp	Leu	Leu	Val	Arg	Asp	Thr	Pro	Gly
				200					205					210
Glu	Asn	Val	Glu	Pro	Ser	His	Thr	Ala	Thr	Ala	Thr	Leu	Val	Leu
				215					220					225
Asn	Val	Val	Pro	Ala	Asp	Leu	Arg	Pro	Pro	Trp	Phe	Leu	Pro	Cys
				230					235					240
Thr	Phe	Ser	Asp	Gly	Tyr	Val	Cys	Ile	Gln	Ala	Gln	Tyr	His	Gly
				245					250					255
Ala	Val	Pro	Thr	Gly	His	Ile	Leu	Pro	Ser	Pro	Leu	Val	Leu	Arg
				260					265					270
Pro	Gly	Pro	Ile	Tyr	Ala	Glu	Asp	Gly	Asp	Arg	Gly	Ile	Asn	Gln
				275					280					285
Pro	Ile	Ile	Tyr	Ser	Ile	Phe	Arg	Gly	Asn	Val	Asn	Gly	Thr	Phe
				290					295					300
Ile	Ile	His	Pro	Asp	Ser	Gly	Asn	Leu	Thr	Val	Ala	Arg	Ser	Val
				305					310					315
Pro	Ser	Pro	Met	Thr	Phe	Leu	Leu	Leu	Val	Lys	Gly	Gln	Gln	Ala
				320					325					330
Asp	Leu	Ala	Arg	Tyr	Ser	Val	Thr	Gln	Val	Thr	Val	Glu	Ala	Val
				335					340					345
Ala	Ala	Ala	Gly	Ser	Pro	Pro	Arg	Phe	Pro	Gln	Ser	Leu	Tyr	Arg
				350					355					360
Gly	Thr	Val	Ala	Arg	Gly	Ala	Gly	Ala	Gly	Val	Val	Val	Lys	Asp
				365					370					375
Ala	Ala	Ala	Pro	Ser	Gln	Pro	Leu	Arg	Ile	Gln	Ala	Gln	Asp	Pro
				380					385					390
Glu	Phe	Ser	Asp	Leu	Asn	Ser	Ala	Ile	Thr	Tyr	Arg	Ile	Thr	Asn
				395					400					405
His	Ser	His	Phe	Arg	Met	Glu	Gly	Glu	Val	Val	Leu	Thr	Thr	Thr
				410					415					420
Thr	Leu	Ala	Gln	Ala	Gly	Ala	Phe	Tyr	Ala	Glu	Val	Glu	Ala	His
				425					430					435
Asn	Thr	Val	Thr	Ser	Gly	Thr	Ala	Thr	Thr	Val	Ile	Glu	Ile	Gln
				440					445					450
Val	Ser	Glu	Gln	Glu	Pro	Pro	Ser	Thr	Glu	Ala	Gly	Gly	Thr	Thr
				455					460					465

-continued

---

Gly	Pro	Trp	Thr	Ser	Thr	Thr	Ser	Glu	Val	Pro	Arg	Pro	Pro	Glu
				470					475					480
Pro	Ser	Gln	Gly	Pro	Ser	Thr	Thr	Ser	Ser	Gly	Gly	Gly	Thr	Gly
				485					490					495
Pro	His	Pro	Pro	Ser	Gly	Thr	Thr	Leu	Arg	Pro	Pro	Thr	Ser	Ser
				500					505					510
Thr	Pro	Gly	Gly	Pro	Pro	Gly	Ala	Glu	Asn	Ser	Thr	Ser	His	Gln
				515					520					525
Pro	Ala	Thr	Pro	Gly	Gly	Asp	Thr	Ala	Gln	Thr	Pro	Lys	Pro	Gly
				530					535					540
Thr	Ser	Gln	Pro	Met	Pro	Pro	Gly	Val	Gly	Thr	Ser	Thr	Ser	His
				545					550					555
Gln	Pro	Ala	Thr	Pro	Ser	Gly	Gly	Thr	Ala	Gln	Thr	Pro	Glu	Pro
				560					565					570
Gly	Thr	Ser	Gln	Pro	Met	Pro	Pro	Ser	Met	Gly	Thr	Ser	Thr	Ser
				575					580					585
His	Gln	Pro	Ala	Thr	Pro	Gly	Gly	Gly	Thr	Ala	Gln	Thr	Pro	Glu
				590					595					600
Ala	Gly	Thr	Ser	Gln	Pro	Met	Pro	Pro	Gly	Met	Gly	Thr	Ser	Thr
				605					610					615
Ser	His	Gln	Pro	Thr	Thr	Pro	Gly	Gly	Gly	Thr	Ala	Gln	Thr	Pro
				620					625					630
Glu	Pro	Gly	Thr	Ser	Gln	Pro	Met	Pro	Leu	Ser	Lys	Ser	Thr	Pro
				635					640					645
Ser	Ser	Gly	Gly	Gly	Pro	Ser	Glu	Asp	Lys	Arg	Phe	Ser	Val	Val
				650					655					660
Asp	Met	Ala	Ala	Leu	Gly	Gly	Val	Leu	Gly	Ala	Leu	Leu	Leu	Leu
				665					670					675
Ala	Leu	Leu	Gly	Leu	Ala	Val	Leu	Val	His	Lys	His	Tyr	Gly	Pro
				680					685					690
Arg	Leu	Lys	Cys	Cys	Ser	Gly	Lys	Ala	Pro	Glu	Pro	Gln	Pro	Gln
				695					700					705
Gly	Phe	Asp	Asn	Gln	Ala	Phe	Leu	Pro	Asp	His	Lys	Ala	Asn	Trp
				710					715					720
Ala	Pro	Val	Pro	Ser	Pro	Thr	His	Asp	Pro	Lys	Pro	Ala	Glu	Ala
				725					730					735
Pro	Met	Pro	Ala	Glu	Pro	Ala	Pro	Pro	Gly	Pro	Ala	Ser	Pro	Gly
				740					745					750
Gly	Ala	Pro	Glu	Pro	Pro	Ala	Ala	Ala	Arg	Ala	Gly	Gly	Ser	Pro
				755					760					765
Thr	Ala	Val	Arg	Ser	Ile	Leu	Thr	Lys	Glu	Arg	Arg	Pro	Glu	Gly
				770					775					780
Gly	Tyr	Lys	Ala	Val	Trp	Phe	Gly	Glu	Asp	Ile	Gly	Thr	Glu	Ala
				785					790					795
Asp	Val	Val	Val	Leu	Asn	Ala	Pro	Thr	Leu	Asp	Val	Asp	Gly	Ala
				800					805					810
Ser	Asp	Ser	Gly	Ser	Gly	Asp	Glu	Gly	Glu	Gly	Ala	Gly	Arg	Gly
				815					820					825
Gly	Gly	Pro	Tyr	Asp	Ala	Pro	Gly	Gly	Asp	Asp	Ser	Tyr	Ile	
				830					835					

-continued

&lt;210&gt; SEQ ID NO 55

&lt;211&gt; LENGTH: 3846

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 55

gcagctgggt tctcccggtt cccttgggca ggtgcagggt cgggttcaaa	50
gcctccggaa cgcgttttgg cctgatttga ggaggggggc ggggaggac	100
ctgCGGcttg cggccccgcc cccttctcgg gctcgcagcc gaccgtaag	150
cccgcctcct ccctcggcgg gccttggggc cgtgtccgcc gggcaactcc	200
agccgaggcc tgggttcttg cctgcagggt tctcggcga gggccctagg	250
gtacagcccg atttgcccc atggtggggtt tcggggccea cggcgggct	300
ggccgcctcg cctctctcgt gctgggtgtg ctgctggtgg tgatcgtcgt	350
cctcgccttc aactactgga gcatctcctc cggccacgtc ctgcttcagg	400
aggaggtggc cgagctgcag ggccaggctc agcgcaccga agtggcccgc	450
ggcggtctgg aaaagcgcaa ttccggacctc ttgctgttgg tggacacgca	500
caagaaacag atcgaccaga aggaggccga ctacggccgc ctcagcagcc	550
ggctgcaggc cagagagggc ctcggaaga gatgcgagga tgacaagggt	600
aaactacaga acaacatata gtatcagatg gcagacatac atcatttaaa	650
ggagcaactt gctgagcttc gtcaggaatt tottogacaa gaagaccagc	700
ttcaggacta taggaagaac aatacttacc ttgtgaagag gttagaatat	750
gaaagttttc agtgtggaca gcagatgaag gaattgagag cacagcatga	800
agaaaatatt aaaagttag cagaccagtt tttagaggaa caaaagcaag	850
agacccaaaa gattcaatca aatgatggaa aggaattgga tataaacaat	900
caagtagtac ctaaaaatat tccaaaagta gctgagaatg ttgcagataa	950
gaatgaagaa ccctcaagca atcatattcc acatgggaaa gaacaaatca	1000
aaagaggtgg tgatgcaggg atgcctggaa tagaagagaa tgacctagca	1050
aaagtgtatg atcttcccc tgctttaagg aagcctccta tttcagtttc	1100
tcaaacatgaa agtcatcaag caatctccca tcttccaact ggacaacctc	1150
tctccccaaa tatgcctcca gattcacaca taaaccacaa tggaaacccc	1200
ggtacttcaa aacagaatcc ttccagtcct cttcagcggt taattccagg	1250
ctcaaaacttg gacagtgaac ccagaattca aacagatata ctaaagcagg	1300
ctaccaagga cagagtccgt gatttccata aattgaagca aaatgatgaa	1350
gaacgagagc ttcaaatgga tcctgcagac tatggaaagc aacatttcaa	1400
tgatgtcctt taagtcttaa agaatgctt cagaaaacct aaagtgcgtg	1450
aaaatgaaat cattctactt tgtcctttct gacttttgtt gtaaagacga	1500
attgtatcag ttgtaaagat acattgagat agaattaagg aaaaacttta	1550
atgaaggaat gtacctatgt acatatgtga actttttcat attgtattat	1600
caaggatatg acttttttgg ttatgataca gttaaagcaa aaacagctaa	1650
tctttgcata taaagcaaac taatgtatat ttcacatttt attgagccga	1700
cttattttcca caaatagata aacaggacaa aatagttgta caggttatat	1750

-continued

---

gtggcatagc	ataaccacag	taagaacaga	acagatattc	agcagaaaac	1800
ttttatact	ctaattcttt	ttttttttt	tttgagacag	agtttttagtc	1850
ttgtttccca	ggctggagtg	caatggcaca	atcttggtc	actgcaacct	1900
ccgcctcctg	ggttcaggca	atcttctctg	ctcagcctcc	caagtagctg	1950
ggattacagg	caccaccac	catgccacg	taatttttgt	atttttaata	2000
gagagcta	aattgtatat	ttaataaaga	cgggtttcac	catggtggcc	2050
aggctggtct	tgaactcctg	acctcaggtg	atcctcctgc	attggcctcc	2100
caaagtctg	gaattccagg	catgagccac	tgcccccagt	ctacacacta	2150
attcttgcta	gccaacacg	tgctctgttc	tatctacccc	tcatttcacg	2200
ctcaaggagt	catacctaga	atagttacac	acaagaggga	aactggaagc	2250
caaacactgt	acagtattgt	gtagaaagtc	acctccctac	tccttttatt	2300
ttacatgagt	gctgatgtgt	tttggcagat	gagctttcag	ctgaggcctg	2350
atggaaattg	agataacctg	caaagacata	acagtattta	tgagttatat	2400
cttagttctt	gaaattgtgg	aatgcatgat	tgacaatata	tttttaattt	2450
ttattttttc	aagtaatacc	agtactgttt	aactatagcc	agaactggct	2500
aaaattttta	tattttcaga	gttgaagttg	gtgaagacat	tcattgattta	2550
aacaccagat	cctgaaaggg	gttaaactta	ctttgaaatg	aatctgcaat	2600
cagtatttca	aagcttttct	ggtaatttta	gtgatcttat	ttgattagac	2650
tttttcagaa	gtactaaata	aggaatttta	acaggttttt	attaatgcac	2700
agataaatag	aagtacagtg	aggtctatag	ccatttttatt	aaaatagctt	2750
aaaagtgtgt	aaaaaaaaatga	atctttgtaa	ttacttaata	tgtagtttaa	2800
gaaccctgca	agcttatatt	tgctagacct	acaaattatt	ttaaattgcat	2850
ttatcttttt	tgacactatt	cagtggaaatg	tgtaagctag	ctaattcttg	2900
ttttctgatt	taaagcactt	ttaaacttta	tcctgcccc	taaaaacaaa	2950
aggttttgat	cacaagggga	aatttaagat	tgtaaccct	gtttttcaga	3000
agggctactg	ttaattgcac	ataaacatga	aatgtgtttt	ccctgtgta	3050
ctaacacatt	ctaggcaaaa	ttcaaacctta	tagtggtaaa	gaaacagggt	3100
gttcacttgc	tgaggtgcaa	aaattcttaa	gacttctggt	tgaattgct	3150
caatgactag	gaaaagatgt	agtagtttac	taaaattggt	ttctaccat	3200
atcaaatata	acaattcatg	cctttatagg	gtcaggccta	caatgaatag	3250
gtatggtggt	ttcacagaat	tttaaaatag	agttaaaggg	aagtgatgta	3300
catttcgggg	gcattagggg	agggagatga	atcaaaaaat	accctagta	3350
atgctttata	ttttaatact	gcaaaagctt	tacaaatgga	aacctgcaa	3400
ttacctgcct	tagttctttt	gtcataaaaa	caatcacttg	gttggttgta	3450
ttgtagctat	tacttataca	gcaacatttc	ttcaattagc	agtctagaca	3500
ttttataaac	agaaatcttg	gaccaattga	taattttct	gactgtatta	3550
atattttagt	gctataaaat	actatgtgaa	tccttaaaa	atctgacatt	3600
ttacagtctg	tattagacat	actgttttta	taatgtttta	cttctgctt	3650

-continued

---

```

aagatttagg ttttttaaat gtatttttgc cctgaattaa gtgtaattt      3700
gatggaaact ctgcttttaa aatcatcatt tactgggttc taataaatta      3750
aaaattaaac ttgaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      3800
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa          3846

```

```

<210> SEQ ID NO 56
<211> LENGTH: 380
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

```

```

<400> SEQUENCE: 56

```

```

Met Val Gly Phe Gly Ala Asn Arg Arg Ala Gly Arg Leu Pro Ser
 1                5                10                15
Leu Val Leu Val Val Leu Leu Val Val Ile Val Val Leu Ala Phe
                20                25                30
Asn Tyr Trp Ser Ile Ser Ser Arg His Val Leu Leu Gln Glu Glu
                35                40                45
Val Ala Glu Leu Gln Gly Gln Val Gln Arg Thr Glu Val Ala Arg
                50                55                60
Gly Arg Leu Glu Lys Arg Asn Ser Asp Leu Leu Leu Leu Val Asp
                65                70                75
Thr His Lys Lys Gln Ile Asp Gln Lys Glu Ala Asp Tyr Gly Arg
                80                85                90
Leu Ser Ser Arg Leu Gln Ala Arg Glu Gly Leu Gly Lys Arg Cys
                95                100               105
Glu Asp Asp Lys Val Lys Leu Gln Asn Asn Ile Ser Tyr Gln Met
                110               115               120
Ala Asp Ile His His Leu Lys Glu Gln Leu Ala Glu Leu Arg Gln
                125               130               135
Glu Phe Leu Arg Gln Glu Asp Gln Leu Gln Asp Tyr Arg Lys Asn
                140               145               150
Asn Thr Tyr Leu Val Lys Arg Leu Glu Tyr Glu Ser Phe Gln Cys
                155               160               165
Gly Gln Gln Met Lys Glu Leu Arg Ala Gln His Glu Glu Asn Ile
                170               175               180
Lys Lys Leu Ala Asp Gln Phe Leu Glu Glu Gln Lys Gln Glu Thr
                185               190               195
Gln Lys Ile Gln Ser Asn Asp Gly Lys Glu Leu Asp Ile Asn Asn
                200               205               210
Gln Val Val Pro Lys Asn Ile Pro Lys Val Ala Glu Asn Val Ala
                215               220               225
Asp Lys Asn Glu Glu Pro Ser Ser Asn His Ile Pro His Gly Lys
                230               235               240
Glu Gln Ile Lys Arg Gly Gly Asp Ala Gly Met Pro Gly Ile Glu
                245               250               255
Glu Asn Asp Leu Ala Lys Val Asp Asp Leu Pro Pro Ala Leu Arg
                260               265               270
Lys Pro Pro Ile Ser Val Ser Gln His Glu Ser His Gln Ala Ile
                275               280               285
Ser His Leu Pro Thr Gly Gln Pro Leu Ser Pro Asn Met Pro Pro
                290               295               300

```

-continued

Asp Ser His Ile Asn His Asn Gly Asn Pro Gly Thr Ser Lys Gln  
 305 310 315  
 Asn Pro Ser Ser Pro Leu Gln Arg Leu Ile Pro Gly Ser Asn Leu  
 320 325 330  
 Asp Ser Glu Pro Arg Ile Gln Thr Asp Ile Leu Lys Gln Ala Thr  
 335 340 345  
 Lys Asp Arg Val Ser Asp Phe His Lys Leu Lys Gln Asn Asp Glu  
 350 355 360  
 Glu Arg Glu Leu Gln Met Asp Pro Ala Asp Tyr Gly Lys Gln His  
 365 370 375  
 Phe Asn Asp Val Leu  
 380

<210> SEQ ID NO 57  
 <211> LENGTH: 841  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 57

```

ggatgggcga gcagtctgaa tgccagaatg gataaccggt ttgctacagc          50
at ttgttaatt gcttgtgtgc ttagcctcat ttccaccatc tacatggcag          100
cctccattgg cacagacttc tggatgaat atcgaagtcc agttcaagaa          150
aattccagtg atttgaataa aagcatctgg gatgaattca ttagtgatga          200
ggcagatgaa aagacttata atgatgcact ttttogatac aatggcacag          250
tgggattgtg gagacggtgt atcaccatac ccaaaaacat gcattgggtat          300
agccccaccg aaaggacaga gtcatttgat gtggtcacia aatgtgtgag          350
tttcacacta actgagcagt tcatggagaa atttgttgat cccggaaacc          400
acaatagcgg gattgatctc cttaggacct atctttggcg ttgccagttc          450
cttttaccct ttgtgagttt aggtttgatg tgctttgggg ctttgatcgg          500
actttgtgct tgcatttgcc gaagcttata tcccaccatt gccacgggca          550
ttctccatct ccttgcagat accatgctgt gaagtccagg ccacatggag          600
gtgtcctgtg tagatgctcc agctgaaatc ccaagctaag ctcccaactg          650
acagccaaca tcatttccag ccatgtgtgg gagccatcct ggatgtccag          700
ccttaacaag ccttcagagg acttcagcca cagctattat cttactacat          750
ccttgtgaga ctctaataaa gaaccaacta gctgagccca atcaacctat          800
ggaactgata gaaataaaat gaattgttgt tttgtgocgt t          841
    
```

<210> SEQ ID NO 58  
 <211> LENGTH: 184  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 58

Met Asp Asn Arg Phe Ala Thr Ala Phe Val Ile Ala Cys Val Leu  
 1 5 10 15  
 Ser Leu Ile Ser Thr Ile Tyr Met Ala Ala Ser Ile Gly Thr Asp  
 20 25 30  
 Phe Trp Tyr Glu Tyr Arg Ser Pro Val Gln Glu Asn Ser Ser Asp

-continued

	35		40		45
Leu Asn Lys Ser	Ile Trp Asp Glu Phe	Ile Ser Asp Glu Ala Asp			
	50	55			60
Glu Lys Thr Tyr	Asn Asp Ala Leu Phe	Arg Tyr Asn Gly Thr Val			
	65	70			75
Gly Leu Trp Arg	Arg Cys Ile Thr Ile	Pro Lys Asn Met His Trp			
	80	85			90
Tyr Ser Pro Pro	Glu Arg Thr Glu Ser	Phe Asp Val Val Thr Lys			
	95	100			105
Cys Val Ser Phe	Thr Leu Thr Glu Gln	Phe Met Glu Lys Phe Val			
	110	115			120
Asp Pro Gly Asn	His Asn Ser Gly Ile	Asp Leu Leu Arg Thr Tyr			
	125	130			135
Leu Trp Arg Cys	Gln Phe Leu Leu Pro	Phe Val Ser Leu Gly Leu			
	140	145			150
Met Cys Phe Gly	Ala Leu Ile Gly Leu	Cys Ala Cys Ile Cys Arg			
	155	160			165
Ser Leu Tyr Pro	Thr Ile Ala Thr Gly	Ile Leu His Leu Leu Ala			
	170	175			180
Asp Thr Met Leu					

<210> SEQ ID NO 59  
 <211> LENGTH: 997  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 59

gcgtggacac cacctcagcc cactgagcag gagtcacagc acgaagacca	50
agcgcaaaag gacccctgcc ctccatcctg actgctcctc ctaagagaga	100
tggcaccggc cagagcagga ttctgcccc ttctgctgct tctgctgctg	150
gggctgtggg tggcagagat cccagtcagt gccaaagcca agggcatgac	200
ctcatcacag tggtttaaaa ttcagcacat gcagcccagc cctcaagcat	250
gcaactcagc catgaaaaac attaacaagc acacaaaacg gtgcaaagac	300
ctcaaacctt tcctgcacga gcctttctcc agtgtggccg ccacctgcca	350
gacccccaaa atagcctgca agaatggcga taaaaactgc caccagagcc	400
acgggcccgt gtcctgacc atgtgtaagc tcacctcagg gaagtatccg	450
aactgcaggt acaaagagaa gcgacagaac aagtottacg tagtggcctg	500
taagcctccc cagaaaaagg actctcagca attccacctg gttcctgtac	550
acttgagacag agtcctttag gtttccagac tggcttgctc tttggctgac	600
cttcaattcc ctctccagga ctccgcacca ctcccctaca cccagagcat	650
tctcttcccc tcatctcttg gggctgttcc tggttcagcc tctgctggga	700
ggctgaagct gacactctgg tgagctgagc tctagaggga tggcttttca	750
tctttttggt gctgttttcc cagatgctta toccaagaa acagcaagct	800
caggctctggt ggttccctgg tctatgccat tgcacatgct tcccctgccc	850
cctggcatta gggcagcatg acaaggagag gaaataaatg gaaagggggc	900
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	950

-continued

aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaa 997

<210> SEQ ID NO 60  
 <211> LENGTH: 156  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 60

Met Ala Pro Ala Arg Ala Gly Phe Cys Pro Leu Leu Leu Leu Leu  
 1 5 10 15  
 Leu Leu Gly Leu Trp Val Ala Glu Ile Pro Val Ser Ala Lys Pro  
 20 25 30  
 Lys Gly Met Thr Ser Ser Gln Trp Phe Lys Ile Gln His Met Gln  
 35 40 45  
 Pro Ser Pro Gln Ala Cys Asn Ser Ala Met Lys Asn Ile Asn Lys  
 50 55 60  
 His Thr Lys Arg Cys Lys Asp Leu Asn Thr Phe Leu His Glu Pro  
 65 70 75  
 Phe Ser Ser Val Ala Ala Thr Cys Gln Thr Pro Lys Ile Ala Cys  
 80 85 90  
 Lys Asn Gly Asp Lys Asn Cys His Gln Ser His Gly Pro Val Ser  
 95 100 105  
 Leu Thr Met Cys Lys Leu Thr Ser Gly Lys Tyr Pro Asn Cys Arg  
 110 115 120  
 Tyr Lys Glu Lys Arg Gln Asn Lys Ser Tyr Val Val Ala Cys Lys  
 125 130 135  
 Pro Pro Gln Lys Lys Asp Ser Gln Gln Phe His Leu Val Pro Val  
 140 145 150  
 His Leu Asp Arg Val Leu  
 155

<210> SEQ ID NO 61  
 <211> LENGTH: 520  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 61

cgggatcatgc gccgcccct gtggctgggc ctggcctggc tgctgctggc 50  
 gcgggcccgc gacgccgcgg gaaccccag cgcgtcgcgg ggaccgcgca 100  
 gctaccgcga cctggagggc gacgtgcgct ggccggcgcct cttctcctcc 150  
 actcacttct tcctgcgcgt ggatcccggc ggccgcgtgc agggcaccgc 200  
 ctggcgccac ggccaggaca gcatcctgga gatccgctct gtacacgtgg 250  
 gcgtcgtggc catcaaagca gtgtcctcag gcttctacgt ggccatgaac 300  
 cgccggggcc gcctctacgg gtcgcgactc tacaccgtgg actgcaggtt 350  
 ccgggagcgc atcgaagaga acggccacaa cacctacgcc tcacagcgct 400  
 ggcgccgcgc cggccagccc atgttctctg cgctggacag gagggggggg 450  
 ccccgccag gcggccggac gcggcggtac cacctgtccg cccacttctc 500  
 gcccgctctg gtctcctgag 520

<210> SEQ ID NO 62



-continued

---

<211> LENGTH: 170  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 62

Met Arg Arg Arg Leu Trp Leu Gly Leu Ala Trp Leu Leu Leu Ala  
 1 5 10 15  
 Arg Ala Pro Asp Ala Ala Gly Thr Pro Ser Ala Ser Arg Gly Pro  
 20 25 30  
 Arg Ser Tyr Pro His Leu Glu Gly Asp Val Arg Trp Arg Arg Leu  
 35 40 45  
 Phe Ser Ser Thr His Phe Phe Leu Arg Val Asp Pro Gly Gly Arg  
 50 55 60  
 Val Gln Gly Thr Arg Trp Arg His Gly Gln Asp Ser Ile Leu Glu  
 65 70 75  
 Ile Arg Ser Val His Val Gly Val Val Val Ile Lys Ala Val Ser  
 80 85 90  
 Ser Gly Phe Tyr Val Ala Met Asn Arg Arg Gly Arg Leu Tyr Gly  
 95 100 105  
 Ser Arg Leu Tyr Thr Val Asp Cys Arg Phe Arg Glu Arg Ile Glu  
 110 115 120  
 Glu Asn Gly His Asn Thr Tyr Ala Ser Gln Arg Trp Arg Arg Arg  
 125 130 135  
 Gly Gln Pro Met Phe Leu Ala Leu Asp Arg Arg Gly Gly Pro Arg  
 140 145 150  
 Pro Gly Gly Arg Thr Arg Arg Tyr His Leu Ser Ala His Phe Leu  
 155 160 165  
 Pro Val Leu Val Ser  
 170

<210> SEQ ID NO 63  
 <211> LENGTH: 2329  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 63

atccctcgac ctcgaccac gcgtccgctg gaaggtggcg tgcctcctc 50  
 tggctggtac catgcagctc cactggccc tgtgtctcgt ctgctgctg 100  
 gtacacacag ccttccgtgt agtggagggc caggggtggc aggcgttaa 150  
 gaatgatgcc acggaatca tccccgagct cggagagtac cccgagcctc 200  
 caccggagct ggagaacaac aagaccatga accgggcgga gaacggaggg 250  
 cggcctcccc accaccctt tgagaccaa gacgtgtccg agtacagctg 300  
 ccgcgagctg cacttcacc gctacgtgac cgatgggccc tgccgagcg 350  
 ccaagccggt caccgagctg gtgtgctccg gccagtgcgg cccggcgcgc 400  
 ctgtgtccca acgcatcgg ccgcggcaag tgggtggcgac ctagtggggc 450  
 cgacttccgc tgcatcccc accgctaccg cgcgcagcgc gtgcagctgc 500  
 tgtgtcccgg tggtagggcg ccgcgcgcgc gcaaggtgcg cctggtggcc 550  
 tcgtgcaagt gcaagcgcct caccgccttc cacaaccagt cggagctcaa 600  
 ggacttcggg accgaggccg ctcggccgca gaaggccgg aagccgcggc 650

-continued

cccgcgcccg gagcgccaaa gccaacccagg ccgagctgga gaacgcctac	700
tagagcccgc ccgcgcccct ccccaccggc gggcgccccg gccctgaacc	750
cgcgcccacc atttctgtcc tctgcgctg gtttgattgt ttatatttca	800
ttgtaaatgc ctgcaaccga gggcaggggg ctgagacctt ccaggccctg	850
aggaatcccg ggcgcccggca aggccccct cagcccgcca gctgaggggt	900
cccacggggc aggggaggga attgagagtc acagacctg agccacgcag	950
ccccgcctct ggggcccct acctttgctg gtcccacttc agaggaggca	1000
gaaatggaag cattttcacc gccctggggg ttaaggag cggtgtggga	1050
gtgggaaagt ccagggactg gtaagaaag ttggataaga ttccccctg	1100
cacctcgctg cccatcagaa agcctgaggc gtgccagag cacaagactg	1150
ggggcaactg tagatgtggt ttctagtctt ggctctgcca ctaacttcct	1200
gtgtaacctt gaactacaca attctccttc gggacctca ttocacttt	1250
gtaaaatgag ggtggagggt ggaataggat ctcgaggaga ctattggcat	1300
atgattccaa ggactccagt gcctttttaa tgggcagagg tgagagagag	1350
agagagaaa agagagaatg aatgcagttg cattgattca gtgccaaagt	1400
cacttcagaa attcagagtt gtgatgctct cttctgacag ccaaagatga	1450
aaaacaaaca gaaaaaaaa agtaaagagt ctatttatgg ctgacatatt	1500
tacggctgac aaactcctgg aagaagctat gctgcttccc agcctggctt	1550
ccccggatgt ttggctacct ccaccctcc atctcaaaga aataacatca	1600
tccattgggg tagaaaagga gagggctcga ggggtgggg agggatagaa	1650
atcacatccg cccaacttc caaaagagca gcacccctcc cccgacccat	1700
agccatgttt taaagtcacc ttccgaagag aagtgaaggt ttcaaggaca	1750
ctggccttgc aggcccagag gagcagccat cacaaactca cagaccagca	1800
catccctttt gagacaccgc cttctgccc ccaactcagg acacatttct	1850
gcctagaaaa cagcttctta ctgctcttac atgtgatggc atatcttaca	1900
ctaaaagaat attattgggg gaaaaactac aagtgctgta catatgctga	1950
gaaactgcag agcataatag ctgccacca aaaatctttt tgaaaatcat	2000
ttccagacaa cctcttactt tctgtgtagt ttttaattgt taaaaaaaa	2050
aagttttaa cagaagcaca tgacatatga aagcctgcag gactggtcgt	2100
ttttttgca attcttccac gtgggacttg tccacaagaa tgaaagtagt	2150
ggtttttaa gagttaagtt acatatttat tttctcactt aagttattta	2200
tgcaaaagtt tttctgtag agaatgacaa tgtaaatatt gctttatgaa	2250
ttaacagtct gttcttccag agtccagaga cattgttaat aaagacaatg	2300
aatcatgaaa aaaaaaaaa aaaaaaaaa	2329

&lt;210&gt; SEQ ID NO 64

&lt;211&gt; LENGTH: 213

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 64

-continued

---

Met	Gln	Leu	Pro	Leu	Ala	Leu	Cys	Leu	Val	Cys	Leu	Leu	Val	His
1				5					10					15
Thr	Ala	Phe	Arg	Val	Val	Glu	Gly	Gln	Gly	Trp	Gln	Ala	Phe	Lys
				20					25					30
Asn	Asp	Ala	Thr	Glu	Ile	Ile	Pro	Glu	Leu	Gly	Glu	Tyr	Pro	Glu
				35					40					45
Pro	Pro	Pro	Glu	Leu	Glu	Asn	Asn	Lys	Thr	Met	Asn	Arg	Ala	Glu
				50					55					60
Asn	Gly	Gly	Arg	Pro	Pro	His	His	Pro	Phe	Glu	Thr	Lys	Asp	Val
				65					70					75
Ser	Glu	Tyr	Ser	Cys	Arg	Glu	Leu	His	Phe	Thr	Arg	Tyr	Val	Thr
				80					85					90
Asp	Gly	Pro	Cys	Arg	Ser	Ala	Lys	Pro	Val	Thr	Glu	Leu	Val	Cys
				95					100					105
Ser	Gly	Gln	Cys	Gly	Pro	Ala	Arg	Leu	Leu	Pro	Asn	Ala	Ile	Gly
				110					115					120
Arg	Gly	Lys	Trp	Trp	Arg	Pro	Ser	Gly	Pro	Asp	Phe	Arg	Cys	Ile
				125					130					135
Pro	Asp	Arg	Tyr	Arg	Ala	Gln	Arg	Val	Gln	Leu	Leu	Cys	Pro	Gly
				140					145					150
Gly	Glu	Ala	Pro	Arg	Ala	Arg	Lys	Val	Arg	Leu	Val	Ala	Ser	Cys
				155					160					165
Lys	Cys	Lys	Arg	Leu	Thr	Arg	Phe	His	Asn	Gln	Ser	Glu	Leu	Lys
				170					175					180
Asp	Phe	Gly	Thr	Glu	Ala	Ala	Arg	Pro	Gln	Lys	Gly	Arg	Lys	Pro
				185					190					195
Arg	Pro	Arg	Ala	Arg	Ser	Ala	Lys	Ala	Asn	Gln	Ala	Glu	Leu	Glu
				200					205					210

Asn Ala Tyr

<210> SEQ ID NO 65  
 <211> LENGTH: 2663  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 65

cccactcggc ggtttggcgg gagggagggg ctttgcgag gccccgctcc	50
cgccccgcct ccatgcggcc cgccccgatt gcgctgtggc tgcgcctggt	100
cttgcccctg gcccttgtcc gccccgggc tgtgggtgg gccccggtcc	150
gagcccccat ctatgtcagc agctgggccc tccaggtgtc ccagggtaac	200
cgggaggtgc agcgcctggc acgcaaatcc ggcttcgtca acctggggcc	250
gatcttctct gacgggcagt actttcacct gcggcaccgg ggcgtggtcc	300
agcagtcctc gacccccgac tggggccacc gctgcacct gaagaaaaac	350
cccaaggtgc agtggttcca gcagcagacg ctgcagcggc gggtgaaacg	400
ctctgtcgtg gtgcccacgg acccctggtt ctccaagcag tggatcatga	450
acagcgaggc ccaaccagac ctgagcatcc tgcaggcctg gagtcagggg	500
ctgtcaggcc agggcatcgt ggtctctgtg ctggacgatg gcatcgagaa	550
ggaccaccgg gacctctggg ccaactacga cccctggcc agctatgact	600

-continued

tcaatgacta cgacccggac ccccagcccc gctacacccc cagcaaagag	650
aaccggcacg ggacccgctg tgctggggag gtggccgca tggccaacaa	700
tggtctctgt ggtgtggggg tcgctttcaa cgcccgaatc ggaggcgtac	750
ggatgctgga cggtagcatc accgatgtca tcgaggccca gtcgctgagc	800
ctgcagccgc agcacatcca cattedacgc gccagctggg gtcccagga	850
cgacggccgc acggtggaag gccccggcat cctcaccgc gaggccttc	900
ggcgtggtgt gaccaagggc cgcggcgggc tgggcacgct cttcatctgg	950
gcctcgggca acggcggcct gcactacgac aactgcaact gcgacggcta	1000
caccaacagc atccacacgc tttccgtggg cagcaccacc cagcagggcc	1050
gcgtgccctg gtacagcga gacctgcct ccacctcac caccacctac	1100
agcagcggcg tggccaccga cccccagatc gtcaccacgg acctgcatca	1150
cgggtgcaca gaccagcaca cgggcacctc ggcctcagcc ccaactggcg	1200
ccggcatgat cgcctagcg ctggaggcca acccgttctt gacgtggaga	1250
gacatgcagc acctggtggt ccgcgcgtcc aagccggcgc acctgcaggc	1300
cgaggactgg aggaccaacg gcgtggggcg ccaagtgagc catcactacg	1350
gatacggggt gctggacgcc gggctgctgg tggacaccgc ccgacactgg	1400
ctgccccacc agccgcagag gaagtgcgcc gtccgggtcc agagccgcc	1450
cacccccatc ctgccgctga tctacatcag ggaaaacgta tcggcctgcg	1500
ccggcctcca caactccatc cgtctgctgg agcactgca ggcgcagctg	1550
acgtgttctt acagccggcg cggagacctg gagatctcgc tcaccagccc	1600
catgggcaag cgtccacac tcgtggccat acgacccttg gacgtcagca	1650
ctgaaggcta caacaactgg gtcttcatgt ccacctactt ctgggatgag	1700
aaccacagc gcgtgtggac cctgggccta gagaacaagg gctactatct	1750
caacacgggg acgttgtacc gctacacgct gctgctctat gggacggccg	1800
aggacatgac agcgcggcct acaggcccc aggtgaccag cagcgcgtgt	1850
gtgcagcggg acacagaggg gctgtgccag gcgtgtgacg gccccgccta	1900
catctctgga cagctctgcc tggcctactg cccccgcgg ttcttcaacc	1950
acacaaggct ggtgaccgct gggcctgggc acacggcggc gcccgcgctg	2000
agggctctgt ccagctgcca tgctctctgc tacacctgcc gcggcggtc	2050
cccaggggac tgcacctcct gtccccatc ctccacgctg gaccagcagc	2100
agggctcctg catgggaccc accacccccg acagccgcc cggccttaga	2150
gctgcgcctg gtccccacca ccgctgcccc gcctcggcca tgggtctgag	2200
cctcctggcc gtgacctcg gaggccccgt cctctgcggc atgtccatgg	2250
acctcccact atacgcctgg ctctcccgtg ccagggccac ccccacaaa	2300
ccccaggctt ggtgcccagc tggaaactga agttgtcagc tcagaaagcg	2350
accttgcccc cgcctgggtc cctgacagcg actgctgcca tgctgcctcc	2400
ccaggctggc cccagaggag cagcaccag caccgcagc ctggcctgcc	2450
agggatgggc cccgtggaac cccgaagcct ggcgggagag agagagagag	2500

-continued

---

aagtctcctc tgcattttgg gtttgggcag gagtgggctg gggggagagg	2550
ctggagcacc ccaaaagcca ggggaaagtg gagggagaga aacgtgacac	2600
tgtcctgtctc gggcaccgcg tccaacctca gagtttgcaa ataaaggttg	2650
cttagaaggt gaa	2663

<210> SEQ ID NO 66  
 <211> LENGTH: 755  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 66

Met Arg Pro Ala Pro Ile Ala Leu Trp Leu Arg Leu Val Leu Ala	1	5	10	15
Leu Ala Leu Val Arg Pro Arg Ala Val Gly Trp Ala Pro Val Arg	20	25	30	
Ala Pro Ile Tyr Val Ser Ser Trp Ala Val Gln Val Ser Gln Gly	35	40	45	
Asn Arg Glu Val Glu Arg Leu Ala Arg Lys Phe Gly Phe Val Asn	50	55	60	
Leu Gly Pro Ile Phe Ser Asp Gly Gln Tyr Phe His Leu Arg His	65	70	75	
Arg Gly Val Val Gln Gln Ser Leu Thr Pro His Trp Gly His Arg	80	85	90	
Leu His Leu Lys Lys Asn Pro Lys Val Gln Trp Phe Gln Gln Gln	95	100	105	
Thr Leu Gln Arg Arg Val Lys Arg Ser Val Val Val Pro Thr Asp	110	115	120	
Pro Trp Phe Ser Lys Gln Trp Tyr Met Asn Ser Glu Ala Gln Pro	125	130	135	
Asp Leu Ser Ile Leu Gln Ala Trp Ser Gln Gly Leu Ser Gly Gln	140	145	150	
Gly Ile Val Val Ser Val Leu Asp Asp Gly Ile Glu Lys Asp His	155	160	165	
Pro Asp Leu Trp Ala Asn Tyr Asp Pro Leu Ala Ser Tyr Asp Phe	170	175	180	
Asn Asp Tyr Asp Pro Asp Pro Gln Pro Arg Tyr Thr Pro Ser Lys	185	190	195	
Glu Asn Arg His Gly Thr Arg Cys Ala Gly Glu Val Ala Ala Met	200	205	210	
Ala Asn Asn Gly Phe Cys Gly Val Gly Val Ala Phe Asn Ala Arg	215	220	225	
Ile Gly Gly Val Arg Met Leu Asp Gly Thr Ile Thr Asp Val Ile	230	235	240	
Glu Ala Gln Ser Leu Ser Leu Gln Pro Gln His Ile His Ile Tyr	245	250	255	
Ser Ala Ser Trp Gly Pro Glu Asp Asp Gly Arg Thr Val Asp Gly	260	265	270	
Pro Gly Ile Leu Thr Arg Glu Ala Phe Arg Arg Gly Val Thr Lys	275	280	285	
Gly Arg Gly Gly Leu Gly Thr Leu Phe Ile Trp Ala Ser Gly Asn	290	295	300	

-continued

---

Gly	Gly	Leu	His	Tyr	Asp	Asn	Cys	Asn	Cys	Asp	Gly	Tyr	Thr	Asn
				305					310					315
Ser	Ile	His	Thr	Leu	Ser	Val	Gly	Ser	Thr	Thr	Gln	Gln	Gly	Arg
				320					325					330
Val	Pro	Trp	Tyr	Ser	Glu	Ala	Cys	Ala	Ser	Thr	Leu	Thr	Thr	Thr
				335					340					345
Tyr	Ser	Ser	Gly	Val	Ala	Thr	Asp	Pro	Gln	Ile	Val	Thr	Thr	Asp
				350					355					360
Leu	His	His	Gly	Cys	Thr	Asp	Gln	His	Thr	Gly	Thr	Ser	Ala	Ser
				365					370					375
Ala	Pro	Leu	Ala	Ala	Gly	Met	Ile	Ala	Leu	Ala	Leu	Glu	Ala	Asn
				380					385					390
Pro	Phe	Leu	Thr	Trp	Arg	Asp	Met	Gln	His	Leu	Val	Val	Arg	Ala
				395					400					405
Ser	Lys	Pro	Ala	His	Leu	Gln	Ala	Glu	Asp	Trp	Arg	Thr	Asn	Gly
				410					415					420
Val	Gly	Arg	Gln	Val	Ser	His	His	Tyr	Gly	Tyr	Gly	Leu	Leu	Asp
				425					430					435
Ala	Gly	Leu	Leu	Val	Asp	Thr	Ala	Arg	Thr	Trp	Leu	Pro	Thr	Gln
				440					445					450
Pro	Gln	Arg	Lys	Cys	Ala	Val	Arg	Val	Gln	Ser	Arg	Pro	Thr	Pro
				455					460					465
Ile	Leu	Pro	Leu	Ile	Tyr	Ile	Arg	Glu	Asn	Val	Ser	Ala	Cys	Ala
				470					475					480
Gly	Leu	His	Asn	Ser	Ile	Arg	Ser	Leu	Glu	His	Val	Gln	Ala	Gln
				485					490					495
Leu	Thr	Leu	Ser	Tyr	Ser	Arg	Arg	Gly	Asp	Leu	Glu	Ile	Ser	Leu
				500					505					510
Thr	Ser	Pro	Met	Gly	Thr	Arg	Ser	Thr	Leu	Val	Ala	Ile	Arg	Pro
				515					520					525
Leu	Asp	Val	Ser	Thr	Glu	Gly	Tyr	Asn	Asn	Trp	Val	Phe	Met	Ser
				530					535					540
Thr	His	Phe	Trp	Asp	Glu	Asn	Pro	Gln	Gly	Val	Trp	Thr	Leu	Gly
				545					550					555
Leu	Glu	Asn	Lys	Gly	Tyr	Tyr	Phe	Asn	Thr	Gly	Thr	Leu	Tyr	Arg
				560					565					570
Tyr	Thr	Leu	Leu	Leu	Tyr	Gly	Thr	Ala	Glu	Asp	Met	Thr	Ala	Arg
				575					580					585
Pro	Thr	Gly	Pro	Gln	Val	Thr	Ser	Ser	Ala	Cys	Val	Gln	Arg	Asp
				590					595					600
Thr	Glu	Gly	Leu	Cys	Gln	Ala	Cys	Asp	Gly	Pro	Ala	Tyr	Ile	Leu
				605					610					615
Gly	Gln	Leu	Cys	Leu	Ala	Tyr	Cys	Pro	Pro	Arg	Phe	Phe	Asn	His
				620					625					630
Thr	Arg	Leu	Val	Thr	Ala	Gly	Pro	Gly	His	Thr	Ala	Ala	Pro	Ala
				635					640					645
Leu	Arg	Val	Cys	Ser	Ser	Cys	His	Ala	Ser	Cys	Tyr	Thr	Cys	Arg
				650					655					660
Gly	Gly	Ser	Pro	Arg	Asp	Cys	Thr	Ser	Cys	Pro	Pro	Ser	Ser	Thr
				665					670					675
Leu	Asp	Gln	Gln	Gln	Gly	Ser	Cys	Met	Gly	Pro	Thr	Thr	Pro	Asp

-continued

	680		685		690
Ser Arg Pro Arg Leu Arg Ala Ala Ala Cys Pro His His Arg Cys					
	695		700		705
Pro Ala Ser Ala Met Val Leu Ser Leu Leu Ala Val Thr Leu Gly					
	710		715		720
Gly Pro Val Leu Cys Gly Met Ser Met Asp Leu Pro Leu Tyr Ala					
	725		730		735
Trp Leu Ser Arg Ala Arg Ala Thr Pro Thr Lys Pro Gln Val Trp					
	740		745		750
Leu Pro Ala Gly Thr					
	755				

<210> SEQ ID NO 67  
 <211> LENGTH: 332  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 67

atgaggaagc tccagggcag gatggtttac ctgcctggac agcaagatga	50
tggtctacact agccccatt ctctggcgcg ctggatttgc ccaccagatc	100
tctctcacctc ttgcccttca cctctgtctg tacctacaag gtctccccga	150
ttctcatctg ccataatca tggacacagc cccaggatgt gcaggactct	200
cagggaccat ctggagtcc agctggaatc tgggctggt ggagtgggag	250
tggggcaggg gcctgcattg ggctgactta gagagcacag ttattccatc	300
catatggaaa taaacatttt ggattcctga tc	332

<210> SEQ ID NO 68  
 <211> LENGTH: 88  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 68

Met Met Ala Thr Leu Ala Pro Ile Leu Trp Ala Pro Gly Phe Ala					
1	5		10		15
His Gln Ile Ser Ser Pro Leu Ala Leu His Leu Leu Leu Tyr Leu					
	20		25		30
Gln Gly Leu Pro Asp Ser His Leu Pro Ile Ile Met Asp Thr Ala					
	35		40		45
Pro Gly Cys Ala Gly Leu Ser Gly Thr Ile Trp Ser Ser Ser Trp					
	50		55		60
Asn Leu Gly Leu Val Glu Trp Glu Trp Gly Arg Gly Leu His Trp					
	65		70		75
Ala Asp Leu Glu Ser Thr Val Ile Pro Ser Ile Trp Lys					
	80		85		

<210> SEQ ID NO 69  
 <211> LENGTH: 1302  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien  
 <220> FEATURE:  
 <221> NAME/KEY: unsure  
 <222> LOCATION: 1218-1253  
 <223> OTHER INFORMATION: unknown base

<400> SEQUENCE: 69

-continued

```

tttgcagtgg ggtcctcctc tggcctcctg cccctcctgc tgetgctgct      50
gcttccattg ctggcagccc aggggtggggg tggcctgcag gcagcgctgc      100
tggcccttga ggtggggctg gtgggtcttg gggcctccta cctgctcctt      150
tgtacagccc tgcacctgcc ctccagtctt ttcctactcc tggcccaggg      200
taccgcaactg ggggcccgtcc tgggcctgag ctggcgccga ggcctcatgg      250
gtgttcccct gggccttga gctgcctggc tcttagcttg gccaggccta      300
gctctacctc tgggtggctat ggcagcgggg ggcagatggg tgcggcagca      350
gggccccggg gtgcgcccgg gcatactctg actctggttg cgggttctgc      400
tgcgcctgtc acctatggcc ttcggggccc tgcagggctg tggggctgtg      450
ggggaccggg gtctgtttgc actgtacccc aaaaccaaca aggatggctt      500
ccgcagcccg ctgcccgtcc ctgggccccg gggcgtaaat ccccgacca      550
cccaacaccc attagctctg ttggcaaggg tctgggtcct gtgcaagggc      600
tggaactggc gtctggcacg ggccagccag ggttagcat cccactggc      650
cccgtggggc atccacacac tggccagctg gggcctgctt cggggtgaa      700
ggcccaccgg aatcccccg ctactaccac gcagccagcg ccagctaggg      750
ccccctgctc cccgccagcc actgccaggg actctagccg ggcggaggtc      800
acgcaccggc cagtcccggg cccctgcccc ctggaggtag ctgactccag      850
cccttcacgc ccaaactctag agcattgagc actttatctc ccacgactca      900
gtgaagtttc tccagtccct agtcctctct tttcaccac cttcctcagt      950
ttgtcactt accccaggcc cagcccttcg gacctctaga caggcagcct     1000
cctcagctgt ggagtccagc agtcactctg tgttctcctg gcgctcctcc     1050
cctaagttaa tgctgttcgc ccgctgtgtg tgctcactct caccctcatt     1100
gactcagggc tggggccagg ggtggtggag ggtgggaaga gtcatgtttt     1150
ttttctctc tttgatthttg tttttctgtc tcccttccaa cctgtcccct     1200
tccccccacc aaaaaaannn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn     1250
nnnaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     1300
aa                                                                 1302
    
```

```

<210> SEQ ID NO 70
<211> LENGTH: 197
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien
    
```

<400> SEQUENCE: 70

```

Met Gly Val Pro Leu Gly Leu Gly Ala Ala Trp Leu Leu Ala Trp
 1           5           10          15
Pro Gly Leu Ala Leu Pro Leu Val Ala Met Ala Ala Gly Gly Arg
 20          25          30
Trp Val Arg Gln Gln Gly Pro Arg Val Arg Arg Gly Ile Ser Arg
 35          40          45
Leu Trp Leu Arg Val Leu Leu Arg Leu Ser Pro Met Ala Phe Arg
 50          55          60
Ala Leu Gln Gly Cys Gly Ala Val Gly Asp Arg Gly Leu Phe Ala
    
```



-continued

---

	65		70		75
Leu Tyr Pro Lys Thr Asn Lys Asp Gly Phe Arg Ser Arg Leu Pro	80		85		90
Val Pro Gly Pro Arg Arg Arg Asn Pro Arg Thr Thr Gln His Pro	95		100		105
Leu Ala Leu Leu Ala Arg Val Trp Val Leu Cys Lys Gly Trp Asn	110		115		120
Trp Arg Leu Ala Arg Ala Ser Gln Gly Leu Ala Ser His Leu Pro	125		130		135
Pro Trp Ala Ile His Thr Leu Ala Ser Trp Gly Leu Leu Arg Gly	140		145		150
Glu Arg Pro Thr Arg Ile Pro Arg Leu Leu Pro Arg Ser Gln Arg	155		160		165
Gln Leu Gly Pro Pro Ala Ser Arg Gln Pro Leu Pro Gly Thr Leu	170		175		180
Ala Gly Arg Arg Ser Arg Thr Arg Gln Ser Arg Ala Leu Pro Pro	185		190		195

Trp Arg

<210> SEQ ID NO 71  
 <211> LENGTH: 1976  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 71

```

gtttgggggt tgtttgggat tagtgaagct actgcctttg ccgccagcgc           50
agcctcagag tttgattatt tgcaatgtca ggctttgaaa acttaaacac           100
ggattttctac cagacaagtt acagcatcga tgatcagtca cagcagtcct           150
atgattatgg aggaagtgga ggaccctata gcaaacagta tgctggctat           200
gactattcgc agcaaggcag atttgtccct ccagacatga tgcagccaca           250
acagccatac accgggcaga ttaccagcc aactcaggca tatactccag           300
cttcacctca gcctttctat ggaacaact ttgaggatga gccaccttta           350
ttagaagagt taggtatcaa ttttgaccac atctggcaaa aaacactaac           400
agtattacat ccgttaaaag tagcagatgg cagcatcatg aatgaaactg           450
atttggcagg tccaatgggt ttttgcttg cttttggagc cacattgcta           500
ctggctggca aaatccagtt tggctatgta tacgggatca gtgcaattgg           550
atgtctagga atgttttggt tattaactt aatgagtatg acaggtgttt           600
catttggttg tgtggcaagt gtccctggat attgtcttct gcccatgatc           650
ctactttcca gctttgcagt gatattttct ttgcaaggaa tggtaggaat           700
cattctcact gctgggatta ttggatggtg tagtttttct gttccaaaa           750
tatttttttc tgcattagcc atggaaggac agcaactttt agtagcatat           800
ccttgcgctt tgttatatgg agtctttgcc ctgatttccg tcttttgaaa           850
atztatctgg gatgtggaca tcagtgggcc agatgtacaa aaaggacctt           900
gaactcttaa attggaccag caaactgctg cagcgcaact ctcatgcaga           950
tttacatttg actgttggag caatgaaagt aaactgttat ctcttgttca          1000
    
```

-continued

```

ttttataga acttttgcat actatattgg atttacctgc ggtgtgacta      1050
gctttaaag tttgtgttta tacagataag aaatgctatt tctttctggt      1100
tcttcgagcc attgaaaaac ctttttcctt gcaaattata atgtttttga      1150
tagattttta tcaactgtgg gaaaccaaac acaaagctga taacctttct      1200
taaaaaagc ccagtcacag taaagaagac acaagacggc cgggcgtggt      1250
agctcaagcc tgtaatccca gacttttggg aggccgaggc gggcgatca      1300
caagggcagg agatcgagac catcctggtt aacacggtga aaccccgact      1350
ctactaaaaa tacaaaaaaa attagctggg cgtggtggcg ggcgcctgta      1400
gtcccagcta ctcaggaggc tgaggcagga gaagtgtgaa cccaggaggc      1450
ggagcttgca gtgagccgag atcacaccac tgcactccat ccagcctggg      1500
tgacaggggt agactctgtc tcaaaaaaaaa aaaaaaaaaagg agacacaaga      1550
cttactgcaa aaatattttt ccaaggattt aggaaagaaa aattgcottg      1600
tattctcaag tcaggttaact caagcaaaa aagtgatcca aatgtagagt      1650
atgagtttgc actccaaaaa ttgacatta ctgtaattaa tctcatggaa      1700
tttttgctaa aattcagaga tacgggaagt tcacaatcta cctcattgta      1750
gacatgaaat gcgaacactt acttacatat taatgttaac tcaaccttag      1800
ggacctggaa tggttgcatt aatgctataa tcgttgatc gccacatttc      1850
ccaaaaataa taaaaaaatc actaaccttt ttttaaggaaa atatttaaag      1900
ttttacaaaa ttcaatattg caattatcaa tgtaaagtac atttgaatgc      1950
ttattaaaac tttccaatt aattttt                                     1976
    
```

<210> SEQ ID NO 72  
 <211> LENGTH: 257  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 72

```

Met Ser Gly Phe Glu Asn Leu Asn Thr Asp Phe Tyr Gln Thr Ser
  1           5           10          15
Tyr Ser Ile Asp Asp Gln Ser Gln Gln Ser Tyr Asp Tyr Gly Gly
  20          25          30
Ser Gly Gly Pro Tyr Ser Lys Gln Tyr Ala Gly Tyr Asp Tyr Ser
  35          40          45
Gln Gln Gly Arg Phe Val Pro Pro Asp Met Met Gln Pro Gln Gln
  50          55          60
Pro Tyr Thr Gly Gln Ile Tyr Gln Pro Thr Gln Ala Tyr Thr Pro
  65          70          75
Ala Ser Pro Gln Pro Phe Tyr Gly Asn Asn Phe Glu Asp Glu Pro
  80          85          90
Pro Leu Leu Glu Glu Leu Gly Ile Asn Phe Asp His Ile Trp Gln
  95          100         105
Lys Thr Leu Thr Val Leu His Pro Leu Lys Val Ala Asp Gly Ser
  110         115         120
Ile Met Asn Glu Thr Asp Leu Ala Gly Pro Met Val Phe Cys Leu
  125         130         135
Ala Phe Gly Ala Thr Leu Leu Leu Ala Gly Lys Ile Gln Phe Gly
    
```

-continued

	140		145		150
Tyr Val Tyr Gly Ile Ser Ala Ile Gly Cys Leu Gly Met Phe Cys	155		160		165
Leu Leu Asn Leu Met Ser Met Thr Gly Val Ser Phe Gly Cys Val	170		175		180
Ala Ser Val Leu Gly Tyr Cys Leu Leu Pro Met Ile Leu Leu Ser	185		190		195
Ser Phe Ala Val Ile Phe Ser Leu Gln Gly Met Val Gly Ile Ile	200		205		210
Leu Thr Ala Gly Ile Ile Gly Trp Cys Ser Phe Ser Ala Ser Lys	215		220		225
Ile Phe Ile Ser Ala Leu Ala Met Glu Gly Gln Gln Leu Leu Val	230		235		240
Ala Tyr Pro Cys Ala Leu Leu Tyr Gly Val Phe Ala Leu Ile Ser	245		250		255

Val Phe

<210> SEQ ID NO 73

<211> LENGTH: 1285

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 73

```

acactggcca aaacgcggct cgcctcggc tgcgctcggc tcccgcgggc      50
gctcggcccc gagcccctcc tccccctacc cgccggccgg acagggagga      100
gccaatggct gggcctgcca tccacaccgc tcccatgctg ttctctgtcc      150
tctctgtgcc ccagctgagc ctggcaggcg cccttgacc tgggaccctt      200
gcccgaacc tccctgagaa tcacattgac ctcccaggcc cagcgtgtg      250
gacgcctcag gccagccacc accgcccggc gggcccgggc aagaaggagt      300
ggggcccagg cctgcccagc caggcccagg atggggctgt ggtcaccgcc      350
accaggcagg cctccaggct gccagaggct gaggggctgc tgctgagca      400
gagtcctgca ggctgctgc aggacaagga cctgctcctg ggactggcat      450
tgccctacc cgagaaggag aacagacctc caggttggga gaggaccagg      500
aaacgcagca gggagcacia gagacgcagg gacaggttga ggctgcacca      550
aggccgagcc ttggtccgag gtcccagctc cctgatgaag aaggcagagc      600
tctccgaagc ccagggtgct gatgcagcca tggaggaatc ctccaccagc      650
ctggcgcccc ccatgttctt tctcaccacc tttgaggcag cacctgccac      700
agaagagtcc ctgatcctgc ccgtcacctc cctgcggccc cagcaggcac      750
agcccaggtc tgacggggag gtgatgcccc cgctggacat gccttggtc      800
gactggaccg attatgaaga cttaaacctc gatggttggc cctctgcaaa      850
gaagaaagag aaacaccgcg gtaaactctc cagtgatggt aacgaaacat      900
caccagccga aggggaacca tgcgaccatc accaagactg cctgccaggg      950
acttgctgcg acctgcggga gcatctctgc acaccccaca accgaggcct     1000
caacaacaaa tgcttcgatg actgcatgtg tgtggaaggg ctgogctgct     1050
atgcccgaatt ccaccggaac cgcagggtta cacggaggaa agggcgctgt     1100
    
```

-continued

---

```

gtggagcccg agacggccaa cggcgaccag ggatccttca tcaacgtcta      1150
cgcgccccgc gggactgggg actgagccca ggaggtttgc acaagccggg      1200
cgatttgttt gtaactagca gtgggagatc aagttgggga acagatggct      1250
gaggtgcag actcaggccc aggacactca accccc                        1285

```

```

<210> SEQ ID NO 74
<211> LENGTH: 348
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

```

```

<400> SEQUENCE: 74

```

```

Met Ala Gly Pro Ala Ile His Thr Ala Pro Met Leu Phe Leu Val
 1          5          10          15
Leu Leu Leu Pro Gln Leu Ser Leu Ala Gly Ala Leu Ala Pro Gly
 20          25          30
Thr Pro Ala Arg Asn Leu Pro Glu Asn His Ile Asp Leu Pro Gly
 35          40          45
Pro Ala Leu Trp Thr Pro Gln Ala Ser His His Arg Arg Arg Gly
 50          55          60
Pro Gly Lys Lys Glu Trp Gly Pro Gly Leu Pro Ser Gln Ala Gln
 65          70          75
Asp Gly Ala Val Val Thr Ala Thr Arg Gln Ala Ser Arg Leu Pro
 80          85          90
Glu Ala Glu Gly Leu Leu Pro Glu Gln Ser Pro Ala Gly Leu Leu
 95          100         105
Gln Asp Lys Asp Leu Leu Leu Gly Leu Ala Leu Pro Tyr Pro Glu
 110         115         120
Lys Glu Asn Arg Pro Pro Gly Trp Glu Arg Thr Arg Lys Arg Ser
 125         130         135
Arg Glu His Lys Arg Arg Arg Asp Arg Leu Arg Leu His Gln Gly
 140         145         150
Arg Ala Leu Val Arg Gly Pro Ser Ser Leu Met Lys Lys Ala Glu
 155         160         165
Leu Ser Glu Ala Gln Val Leu Asp Ala Ala Met Glu Glu Ser Ser
 170         175         180
Thr Ser Leu Ala Pro Thr Met Phe Phe Leu Thr Thr Phe Glu Ala
 185         190         195
Ala Pro Ala Thr Glu Glu Ser Leu Ile Leu Pro Val Thr Ser Leu
 200         205         210
Arg Pro Gln Gln Ala Gln Pro Arg Ser Asp Gly Glu Val Met Pro
 215         220         225
Thr Leu Asp Met Ala Leu Phe Asp Trp Thr Asp Tyr Glu Asp Leu
 230         235         240
Lys Pro Asp Gly Trp Pro Ser Ala Lys Lys Lys Glu Lys His Arg
 245         250         255
Gly Lys Leu Ser Ser Asp Gly Asn Glu Thr Ser Pro Ala Glu Gly
 260         265         270
Glu Pro Cys Asp His His Gln Asp Cys Leu Pro Gly Thr Cys Cys
 275         280         285
Asp Leu Arg Glu His Leu Cys Thr Pro His Asn Arg Gly Leu Asn
 290         295         300

```

-continued

Asn Lys Cys Phe Asp Asp Cys Met Cys Val Glu Gly Leu Arg Cys  
 305 310  
 Tyr Ala Lys Phe His Arg Asn Arg Arg Val Thr Arg Arg Lys Gly  
 320 325  
 Arg Cys Val Glu Pro Glu Thr Ala Asn Gly Asp Gln Gly Ser Phe  
 335 340 345

Ile Asn Val

<210> SEQ ID NO 75  
 <211> LENGTH: 1868  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 75

cagaagggca aaaacattga ctgcctcaag gtctcaagca ccagtcttca 50  
 ccgcggaag catgttgtgg ctgttccaat cgctoctggt tgtctttotgc 100  
 tttggcccag ggaatgtagt ttcacaaagc agcttaacct cattgatggt 150  
 gaacgggatt ctgggggagt cagtaactct tcccctggag tttcctgcag 200  
 gagagaaggt caacttcate acttggcttt tcaatgaaac atctcttgcc 250  
 ttcatagtac cccatgaaac caaaagtcca gaaatccacg tgactaatcc 300  
 gaaacaggga aagcgactga acttcaccca gtcctactcc ctgcaactca 350  
 gcaacctgaa gatggaagac acaggctctt acagagccca gatatccaca 400  
 aagacctctg caaagctgtc cagttacact ctgaggatat taagacaact 450  
 gaggaacata caagttacca atcacagtca gctatttcag aatatgacct 500  
 gtgagctcca tctgacttgc tctgtggagg atgcagatga caatgtotca 550  
 ttcagatggg aggccctggg aaacacactt tcaagtcagc caaacctcac 600  
 tgtctcctgg gaccccagga tttccagtga acaggactac acctgcatag 650  
 cagagaatgc tgtcagtaat ttatccttct ctgtctctgc ccagaagctt 700  
 tgcgaaagat ttaaaattca atatacagat accaaaatga ttctgtttat 750  
 ggtttctcgg atatgcatag tcttcggttt catcatactg ctgttacttg 800  
 ttttgaggaa aagaagagat tccctatctt tgtctactca gcgaaacacag 850  
 ggccccgcag agtccgcaag gaacctagag tatgtttcag tgtctccaac 900  
 gaaacaact gtgtatgctt cagtcactca ttcaaacagg gaaacagaaa 950  
 tctggacacc tagagaaaat gatactatca caatttactc cacaattaat 1000  
 cattccaaag agagtaaacc cactttttcc agggcaactg cccttgacaa 1050  
 tgtctgtgtaa gttgctgaaa ggcctcagag gaattcggga atgacacgtc 1100  
 ttctgatccc atgagacaga acaaagaaca ggaagcttgg ttctgttgt 1150  
 tctctggcaac agaatttgaa tatctaggat aggatgatca cctccagtc 1200  
 ttcggactta aacctgccta cctgagtcaa acacctaaag ataacatcat 1250  
 ttccagcatg tggttcaaat aatattttcc aatccacttc aggocaaaac 1300  
 atgctaaaga taacacacca gcacattgac tctctctttg ataactaaagc 1350  
 aaatggaatt atggttgaca gagagtttat gatccagaag acaaccactt 1400

-continued

---

```

ctctcctttt agaaagcagc aggattgact tattgagaaa taatgcagtg      1450
tgttggttac atgtgtagtc tctggagttg gatgggcca tcctgataca      1500
agttgagcat cccttgtctg aaatgcttg gattagaaat gtttcagatt      1550
tcaatttttt ttcagatttt ggaatatttg cattatattt agcggttgag      1600
tatccaaatc caaaaatcca aaattcaaaa tgctccaata agcatttccc      1650
ttgatgttca ttgatgtcga tgcagtgtc aaaatctcag attttgagc      1700
aatttgata ttggattttt ggatttggga tgctcaactt gtacaatgtt      1750
tattagacac atctcctggg acatactgcc taaccttttg gagccttagt      1800
ctcccagact gaaaaaggaa gaggatggtta ttacatcagc tccattgttt      1850
gagccaagaa tctaagtc      1868

```

&lt;210&gt; SEQ ID NO 76

&lt;211&gt; LENGTH: 332

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 76

```

Met Leu Trp Leu Phe Gln Ser Leu Leu Phe Val Phe Cys Phe Gly
 1                    5                    10          15
Pro Gly Asn Val Val Ser Gln Ser Ser Leu Thr Pro Leu Met Val
                20                    25          30
Asn Gly Ile Leu Gly Glu Ser Val Thr Leu Pro Leu Glu Phe Pro
                35                    40          45
Ala Gly Glu Lys Val Asn Phe Ile Thr Trp Leu Phe Asn Glu Thr
                50                    55          60
Ser Leu Ala Phe Ile Val Pro His Glu Thr Lys Ser Pro Glu Ile
                65                    70          75
His Val Thr Asn Pro Lys Gln Gly Lys Arg Leu Asn Phe Thr Gln
                80                    85          90
Ser Tyr Ser Leu Gln Leu Ser Asn Leu Lys Met Glu Asp Thr Gly
                95                    100         105
Ser Tyr Arg Ala Gln Ile Ser Thr Lys Thr Ser Ala Lys Leu Ser
                110                   115         120
Ser Tyr Thr Leu Arg Ile Leu Arg Gln Leu Arg Asn Ile Gln Val
                125                   130         135
Thr Asn His Ser Gln Leu Phe Gln Asn Met Thr Cys Glu Leu His
                140                   145         150
Leu Thr Cys Ser Val Glu Asp Ala Asp Asp Asn Val Ser Phe Arg
                155                   160         165
Trp Glu Ala Leu Gly Asn Thr Leu Ser Ser Gln Pro Asn Leu Thr
                170                   175         180
Val Ser Trp Asp Pro Arg Ile Ser Ser Glu Gln Asp Tyr Thr Cys
                185                   190         195
Ile Ala Glu Asn Ala Val Ser Asn Leu Ser Phe Ser Val Ser Ala
                200                   205         210
Gln Lys Leu Cys Glu Asp Val Lys Ile Gln Tyr Thr Asp Thr Lys
                215                   220         225
Met Ile Leu Phe Met Val Ser Gly Ile Cys Ile Val Phe Gly Phe
                230                   235         240

```

-continued

---

Ile Ile Leu Leu Leu Leu Val Leu Arg Lys Arg Arg Asp Ser Leu  
 245 250 255

Ser Leu Ser Thr Gln Arg Thr Gln Gly Pro Ala Glu Ser Ala Arg  
 260 265 270

Asn Leu Glu Tyr Val Ser Val Ser Pro Thr Asn Asn Thr Val Tyr  
 275 280 285

Ala Ser Val Thr His Ser Asn Arg Glu Thr Glu Ile Trp Thr Pro  
 290 295 300

Arg Glu Asn Asp Thr Ile Thr Ile Tyr Ser Thr Ile Asn His Ser  
 305 310 315

Lys Glu Ser Lys Pro Thr Phe Ser Arg Ala Thr Ala Leu Asp Asn  
 320 325 330

Val Val

<210> SEQ ID NO 77  
 <211> LENGTH: 3073  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 77

```

gatccctcga cctcgacca cgcgtccgct ctttaatgct ttctttttaa           50
gagatcacct tctgacttct cacagaagag gttaactatt acctgtggga           100
agtcagaagg tgatctcttt aatgctttct ttttaagaat ttttcaaatt           150
gagactaatt gcagagggtc cagttgacca gcattcatag gaatgaagac           200
aaacacagag atgggtgtgc taagaaactt caaaagtggt agacctctg           250
actgaagcat attggattta ttttaatttt ttcactgtat ttctgtcctc           300
ctacaaggga aagtcattgat tacactaact gagctaaaat gcttagcaga           350
tgcccagtca tcttatcaca tcttaaaacc atgggtgggac gtcttctggt           400
attacatcac actgatcatg ctgctgggtg ccgctgctggc cggagctctc           450
cagctgacgc agagcagggt tctgtgctgt cttccatgca aagtggaatt           500
tgacaatcac tgtgccgtgc cttgggacat cctgaaagcc agcatgaaca           550
catcctctaa tcctgggaca cgcctccgc tccccctccg aattcagaat           600
gacctccaac gacagcagta ctccatatatt gatgccgtct gttacgagaa           650
acagctccat tggtttgcaa agtttttccc ctatctggtg ctcttgcaaca           700
cgctcatctt tgcagcctgc agcaactttt ggcttcacta ccccagttacc           750
agttccaggc tcgagcattt tgtggccatc cttcacaagt gcttcgattc           800
tccatggacc acccgcgccc tttcagaaac agtggctgag cagtcagtga           850
ggcctctgaa actctccaag tccaagattt tgctttcgtc ctcagggtgt           900
tcagctgaca tagattccgg caaacagtca ttgccttacc cacagccagg           950
tttgagtgca gctggtatag aaagcccaac ttccagtggt cttgacaaga           1000
aggagggtga acaggccaaa gccatctttg aaaaagttaa aagattccgc           1050
atgcatgtgg agcagaagga catcatttat agagtatata tgaacagat           1100
aatagtcaaa gtcattttgt ttgtgctcat cataacttat gttocatatt           1150
ttttaacca catcactctt gaaatcgact gttcagttga tgtgcaggct           1200
    
```

-continued

---

tttacaggat ataagcgcta ccagtgtgtc tattccttgg cagaaatctt	1250
taaggtcctg gcttcatttt atgtcatttt gggtatactt tatggctgta	1300
cctcttccta cagcctgtgg tggatgctga ggagttccct gaagcaatat	1350
tcctttgagg cgtaagaga aaaaagcaac tacagtgaca tccctgatgt	1400
caagaatgac tttgccttca tccttcactt ggctgatcag tatgatcctc	1450
tttattccaa acgcttctcc atattcctat cagaggtcag tgagaacaaa	1500
ctgaaacaga tcaacctcaa taatgaatgg acagttgaga aactgaaaag	1550
taagcttgtg aaaaatgcc aggacaagat agaactgcat ctttttatgc	1600
tcaacggtct tccagacaat gtctttgagt taactgaaat ggaagtgcta	1650
agcctggagc ttatcccaga ggtgaagctg cctctgcag tctcacagct	1700
ggtcaacctc aaggagcttc gtgtgtacca ttcactctctg gtcgtagacc	1750
atcctgcact ggcctttcta gaggagaatt taaaaatcct ccgcctgaaa	1800
tttactgaaa tgggaaaaat cccacgctgg gtatttcacc tcaagaatct	1850
caaggaactt tatctttcgg gctgtgttct cctgaacag ttgagtacta	1900
tgcagttgga gggctttcag gacttaaaaa atctaagac cctgtacttg	1950
aagagcagcc tctcccggat cccacaagtt gttacagacc tctgccttc	2000
attgcagaaa ctgtcccttg ataatgaggg aagcaaaactg gttgtgttga	2050
acaacttgaa aaagatggtc aatctgaaaa gcctagaact gatcagctgt	2100
gacctggaac gcatcccaca ttccattttc agcctgaata attgcatga	2150
gtagaccta agggaaaata accttaaaac tgtggaagag attagctttc	2200
agcatcttca gaatctttcc tgcttaaaagt tgtggcaca taacattgct	2250
tatattcctg cacagattgg ggcattatct aacctagac agctctcttt	2300
ggaccataat aatattgaga atctgccctt gcagcttttc ctatgcacta	2350
aactacatta tttggatcta agctataacc acttgacctt cattccagaa	2400
gaaatccagt atctgagtaa tttgcagtac tttgctgtga ccaacaacaa	2450
tattgagatg ctaccagatg ggctgtttca gtgcaaaaag ctgcagtggt	2500
tacttttggg gaaaaatagc ttgatgaatt tgtcccctca tgtgggtgag	2550
ctgtcaaacc ttactcatct ggagctcatt ggtaattacc tggaaacact	2600
tcctcctgaa ctagaaggat gtcagtcctt aaaacggaac tgtctgattg	2650
ttgaggagaa cttgctcaat actcttcctc tcctgtaac agaacgttta	2700
cagacgtgct tagacaaatg ttgacttaaa gaaaagagac ccgtgtttca	2750
aaatcatttt taaaagtatg ctcgccggg cgtgggtgct catgcctata	2800
atcccagcac tttgggaggc caagatgggc ggattgcttg aggtcaggag	2850
ttcgagacca gtctggccaa cctggtgaaa ccccatctct gctaaaacta	2900
caaaaaaatt agccaggcgt ggtggcgtgc gcctgtaatc ccagctactt	2950
gggaggctga cgcaggggaa ttgcttgaac cagggagggtg gaggttgag	3000
tgagccgaga ttgtgccact gtacaccagc ctgggtgaca gagcaagact	3050
cttatctcaa aaaaaaaaaa aaa	3073



-continued

<210> SEQ ID NO 78  
 <211> LENGTH: 802  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 78

```

Met Ile Thr Leu Thr Glu Leu Lys Cys Leu Ala Asp Ala Gln Ser
 1           5           10           15
Ser Tyr His Ile Leu Lys Pro Trp Trp Asp Val Phe Trp Tyr Tyr
 20           25           30
Ile Thr Leu Ile Met Leu Leu Val Ala Val Leu Ala Gly Ala Leu
 35           40           45
Gln Leu Thr Gln Ser Arg Val Leu Cys Cys Leu Pro Cys Lys Val
 50           55           60
Glu Phe Asp Asn His Cys Ala Val Pro Trp Asp Ile Leu Lys Ala
 65           70           75
Ser Met Asn Thr Ser Ser Asn Pro Gly Thr Pro Leu Pro Leu Pro
 80           85           90
Leu Arg Ile Gln Asn Asp Leu His Arg Gln Gln Tyr Ser Tyr Ile
 95           100          105
Asp Ala Val Cys Tyr Glu Lys Gln Leu His Trp Phe Ala Lys Phe
 110          115          120
Phe Pro Tyr Leu Val Leu Leu His Thr Leu Ile Phe Ala Ala Cys
 125          130          135
Ser Asn Phe Trp Leu His Tyr Pro Ser Thr Ser Ser Arg Leu Glu
 140          145          150
His Phe Val Ala Ile Leu His Lys Cys Phe Asp Ser Pro Trp Thr
 155          160          165
Thr Arg Ala Leu Ser Glu Thr Val Ala Glu Gln Ser Val Arg Pro
 170          175          180
Leu Lys Leu Ser Lys Ser Lys Ile Leu Leu Ser Ser Ser Gly Cys
 185          190          195
Ser Ala Asp Ile Asp Ser Gly Lys Gln Ser Leu Pro Tyr Pro Gln
 200          205          210
Pro Gly Leu Glu Ser Ala Gly Ile Glu Ser Pro Thr Ser Ser Gly
 215          220          225
Leu Asp Lys Lys Glu Gly Glu Gln Ala Lys Ala Ile Phe Glu Lys
 230          235          240
Val Lys Arg Phe Arg Met His Val Glu Gln Lys Asp Ile Ile Tyr
 245          250          255
Arg Val Tyr Leu Lys Gln Ile Ile Val Lys Val Ile Leu Phe Val
 260          265          270
Leu Ile Ile Thr Tyr Val Pro Tyr Phe Leu Thr His Ile Thr Leu
 275          280          285
Glu Ile Asp Cys Ser Val Asp Val Gln Ala Phe Thr Gly Tyr Lys
 290          295          300
Arg Tyr Gln Cys Val Tyr Ser Leu Ala Glu Ile Phe Lys Val Leu
 305          310          315
Ala Ser Phe Tyr Val Ile Leu Val Ile Leu Tyr Gly Leu Thr Ser
 320          325          330
Ser Tyr Ser Leu Trp Trp Met Leu Arg Ser Ser Leu Lys Gln Tyr
 335          340          345

```

-continued

---

Ser	Phe	Glu	Ala	Leu	Arg	Glu	Lys	Ser	Asn	Tyr	Ser	Asp	Ile	Pro
				350					355					360
Asp	Val	Lys	Asn	Asp	Phe	Ala	Phe	Ile	Leu	His	Leu	Ala	Asp	Gln
				365					370					375
Tyr	Asp	Pro	Leu	Tyr	Ser	Lys	Arg	Phe	Ser	Ile	Phe	Leu	Ser	Glu
				380					385					390
Val	Ser	Glu	Asn	Lys	Leu	Lys	Gln	Ile	Asn	Leu	Asn	Asn	Glu	Trp
				395					400					405
Thr	Val	Glu	Lys	Leu	Lys	Ser	Lys	Leu	Val	Lys	Asn	Ala	Gln	Asp
				410					415					420
Lys	Ile	Glu	Leu	His	Leu	Phe	Met	Leu	Asn	Gly	Leu	Pro	Asp	Asn
				425					430					435
Val	Phe	Glu	Leu	Thr	Glu	Met	Glu	Val	Leu	Ser	Leu	Glu	Leu	Ile
				440					445					450
Pro	Glu	Val	Lys	Leu	Pro	Ser	Ala	Val	Ser	Gln	Leu	Val	Asn	Leu
				455					460					465
Lys	Glu	Leu	Arg	Val	Tyr	His	Ser	Ser	Leu	Val	Val	Asp	His	Pro
				470					475					480
Ala	Leu	Ala	Phe	Leu	Glu	Glu	Asn	Leu	Lys	Ile	Leu	Arg	Leu	Lys
				485					490					495
Phe	Thr	Glu	Met	Gly	Lys	Ile	Pro	Arg	Trp	Val	Phe	His	Leu	Lys
				500					505					510
Asn	Leu	Lys	Glu	Leu	Tyr	Leu	Ser	Gly	Cys	Val	Leu	Pro	Glu	Gln
				515					520					525
Leu	Ser	Thr	Met	Gln	Leu	Glu	Gly	Phe	Gln	Asp	Leu	Lys	Asn	Leu
				530					535					540
Arg	Thr	Leu	Tyr	Leu	Lys	Ser	Ser	Leu	Ser	Arg	Ile	Pro	Gln	Val
				545					550					555
Val	Thr	Asp	Leu	Leu	Pro	Ser	Leu	Gln	Lys	Leu	Ser	Leu	Asp	Asn
				560					565					570
Glu	Gly	Ser	Lys	Leu	Val	Val	Leu	Asn	Asn	Leu	Lys	Lys	Met	Val
				575					580					585
Asn	Leu	Lys	Ser	Leu	Glu	Leu	Ile	Ser	Cys	Asp	Leu	Glu	Arg	Ile
				590					595					600
Pro	His	Ser	Ile	Phe	Ser	Leu	Asn	Asn	Leu	His	Glu	Leu	Asp	Leu
				605					610					615
Arg	Glu	Asn	Asn	Leu	Lys	Thr	Val	Glu	Glu	Ile	Ser	Phe	Gln	His
				620					625					630
Leu	Gln	Asn	Leu	Ser	Cys	Leu	Lys	Leu	Trp	His	Asn	Asn	Ile	Ala
				635					640					645
Tyr	Ile	Pro	Ala	Gln	Ile	Gly	Ala	Leu	Ser	Asn	Leu	Glu	Gln	Leu
				650					655					660
Ser	Leu	Asp	His	Asn	Asn	Ile	Glu	Asn	Leu	Pro	Leu	Gln	Leu	Phe
				665					670					675
Leu	Cys	Thr	Lys	Leu	His	Tyr	Leu	Asp	Leu	Ser	Tyr	Asn	His	Leu
				680					685					690
Thr	Phe	Ile	Pro	Glu	Glu	Ile	Gln	Tyr	Leu	Ser	Asn	Leu	Gln	Tyr
				695					700					705
Phe	Ala	Val	Thr	Asn	Asn	Asn	Ile	Glu	Met	Leu	Pro	Asp	Gly	Leu
				710					715					720

-continued

---

Phe Gln Cys Lys Lys Leu Gln Cys Leu Leu Leu Gly Lys Asn Ser  
 725 730 735

Leu Met Asn Leu Ser Pro His Val Gly Glu Leu Ser Asn Leu Thr  
 740 745 750

His Leu Glu Leu Ile Gly Asn Tyr Leu Glu Thr Leu Pro Pro Glu  
 755 760 765

Leu Glu Gly Cys Gln Ser Leu Lys Arg Asn Cys Leu Ile Val Glu  
 770 775 780

Glu Asn Leu Leu Asn Thr Leu Pro Leu Pro Val Thr Glu Arg Leu  
 785 790 795

Gln Thr Cys Leu Asp Lys Cys  
 800

<210> SEQ ID NO 79  
 <211> LENGTH: 1504  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 79

```

cggacgcgtg ggccgcgctc cctcacggcc cctcggcggc gcccgtegga      50
tccggcctct ctctgcgccc cggggcgcgc cacctcccgc cgggaggtgt      100
ccacgcgtcc ggccgtccat ccgtccgtcc ctccctggggc cggcgctgac      150
catgcccagc ggctgcccgt gcttgcattc cgtgtgcctg ttgtgcatc      200
tgggggctcc cggtcagcct gtccgagccg atgactgcag ctcccactgt      250
gacctggccc acggctgctg tgcacctgac ggctcctgca ggtgtgacct      300
gggctgggag gggctgcact gtgagcgtg tgtgaggatg cctggctgcc      350
agcacggtac ctgccaccag ccatggcagt gcatctgcca cagtggctgg      400
gcaggcaagt tctgtgacaa agatgaacat atctgtacca cgcagtcccc      450
ctgccagaat ggaggccagt gcatgtatga cgggggcggg gagtaccatt      500
gtgtgtgctt accaggcttc catgggcgtg actgcgagcg caaggctgga      550
ccctgtgaac aggcaggctc cccatgccgc aatggcgggc agtgccagga      600
cgaccagggc tttgctctca acttcacgtg ccgctgcttg gtgggctttg      650
tgggtgcccg ctgtgaggtg aatgtggatg actgcctgat gcggccttgt      700
gctaacggtg ccacctgcct tgacggcata aaccgcttct cctgcctctg      750
tctctgaggg tttgctggac gcttctgcac catcaacctg gatgactgtg      800
ccagccgccc atgccagaga ggggcccgct gtcgggaccg tgtccacgac      850
ttcgactgcc tctgccccag tggctatggt ggcaagacct gtgagcttgt      900
cttacctgtc ccagaccccc caaccacagt ggacaccctc ctagggccca      950
cctcagctgt agtggctacct gctacggggc cagcccccca cagcgcaggg      1000
gtggtctctc tgcggatctc agtgaaggag gtggtgcgga ggcaagaggc      1050
tgggctaggt gagcctagct tgggtgccct ggtggtgttt ggggccctca      1100
ctgctgccct ggttctggct actgtgttgc tgacctgag gccctggcgc      1150
cggggtgtct gccccctgg accctgttgc taccctgccc cacactatgc      1200
tccagcgtgc caggaccagg agtgtcaggt tagcatgctg ccagcagggc      1250
    
```

-continued

---

```

tccccctgcc acgtgacttg cccccctgagc ctggaaagac cacagcactg      1300
tgatggaggt gggggctttc tggccccctt cctcacctct tccaccctc      1350
agactggagt ggtccgttct caccaccctt cagcttgggt acacacacag      1400
aggagacctc agcctcacac cagaaatatt atttttttaa tacacagaat      1450
gtaagatgga attttatcaa ataaaactat gaaaatgcaa aaaaaaaaaa      1500
aaaa                                                              1504

```

&lt;210&gt; SEQ ID NO 80

&lt;211&gt; LENGTH: 383

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 80

```

Met Pro Ser Gly Cys Arg Cys Leu His Leu Val Cys Leu Leu Cys
 1          5          10          15
Ile Leu Gly Ala Pro Gly Gln Pro Val Arg Ala Asp Asp Cys Ser
          20          25          30
Ser His Cys Asp Leu Ala His Gly Cys Cys Ala Pro Asp Gly Ser
          35          40          45
Cys Arg Cys Asp Pro Gly Trp Glu Gly Leu His Cys Glu Arg Cys
          50          55          60
Val Arg Met Pro Gly Cys Gln His Gly Thr Cys His Gln Pro Trp
          65          70          75
Gln Cys Ile Cys His Ser Gly Trp Ala Gly Lys Phe Cys Asp Lys
          80          85          90
Asp Glu His Ile Cys Thr Thr Gln Ser Pro Cys Gln Asn Gly Gly
          95          100          105
Gln Cys Met Tyr Asp Gly Gly Gly Glu Tyr His Cys Val Cys Leu
          110          115          120
Pro Gly Phe His Gly Arg Asp Cys Glu Arg Lys Ala Gly Pro Cys
          125          130          135
Glu Gln Ala Gly Ser Pro Cys Arg Asn Gly Gly Gln Cys Gln Asp
          140          145          150
Asp Gln Gly Phe Ala Leu Asn Phe Thr Cys Arg Cys Leu Val Gly
          155          160          165
Phe Val Gly Ala Arg Cys Glu Val Asn Val Asp Asp Cys Leu Met
          170          175          180
Arg Pro Cys Ala Asn Gly Ala Thr Cys Leu Asp Gly Ile Asn Arg
          185          190          195
Phe Ser Cys Leu Cys Pro Glu Gly Phe Ala Gly Arg Phe Cys Thr
          200          205          210
Ile Asn Leu Asp Asp Cys Ala Ser Arg Pro Cys Gln Arg Gly Ala
          215          220          225
Arg Cys Arg Asp Arg Val His Asp Phe Asp Cys Leu Cys Pro Ser
          230          235          240
Gly Tyr Gly Gly Lys Thr Cys Glu Leu Val Leu Pro Val Pro Asp
          245          250          255
Pro Pro Thr Thr Val Asp Thr Pro Leu Gly Pro Thr Ser Ala Val
          260          265          270
Val Val Pro Ala Thr Gly Pro Ala Pro His Ser Ala Gly Ala Gly
          275          280          285

```

-continued

Leu Leu Arg Ile Ser Val Lys Glu Val Val Arg Arg Gln Glu Ala  
 290 295 300  
 Gly Leu Gly Glu Pro Ser Leu Val Ala Leu Val Val Phe Gly Ala  
 305 310 315  
 Leu Thr Ala Ala Leu Val Leu Ala Thr Val Leu Leu Thr Leu Arg  
 320 325 330  
 Ala Trp Arg Arg Gly Val Cys Pro Pro Gly Pro Cys Cys Tyr Pro  
 335 340 345  
 Ala Pro His Tyr Ala Pro Ala Cys Gln Asp Gln Glu Cys Gln Val  
 350 355 360  
 Ser Met Leu Pro Ala Gly Leu Pro Leu Pro Arg Asp Leu Pro Pro  
 365 370 375  
 Glu Pro Gly Lys Thr Thr Ala Leu  
 380

<210> SEQ ID NO 81  
 <211> LENGTH: 1034  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 81

```

gtttgttgct caaaccgagt tctggagaac gccatcagct cgctgcttaa      50
aattaaacca caggttccat tatgggtcga cttgatggga aagtcatcat      100
cctgacggcc gctgctcagg ggattggcca agcagctgcc ttagcttttg      150
caagagaagg tgccaaagtc atagccacag acattaatga gtccaaactt      200
caggaactgg aaaagtaccc gggatttcaa actcgtgtcc ttgatgtcac      250
aaagaagaaa caaattgata agtttgccag tgaagttgag agacttgatg      300
ttctctttaa tgttgctggt ttgtgccatc atggaactgt cctggattgt      350
gaggagaaag actgggactt ctcgatgaat ctcaatgtgc gcagcatgta      400
cctgatgata aaggcattcc ttctaaaaat gcttgctcag aaatctggca      450
atattatcaa catgtcttct gtggcttcca gcgtcaaagg agttgtgaac      500
agatgtgtgt acagcacaac caaggcagcc gtgattggcc tcacaaaatc      550
tctggctgca gatttcatcc agcagggcat caggtgcaac tgtgtgtgcc      600
caggaacagt tgatagcca tctctacaag aaagaatata agccagagga      650
aatcctgaag aggcacggaa tgatttcctg aagagacaaa agacgggaag      700
attcgcgaact gcagaagaaa tagccatgct ctgcgtgtat ttggcttctg      750
atgaatctgc ttatgtaact ggtaaccctg tcatcattga tggagctgg      800
agcttgtgat tttaggatct ccatggtggg aaggaaggca ggccttctc      850
atccacagtg aacctggtta cgaagaaaac tcaccaatca tctcttctc      900
gttaatcaca tgtaaatgaa aataagctct ttttaatgat gtcactgttt      950
gcaagagtct gattctttaa gtatattaat ctctttgtaa tctctctga     1000
aatcattgta aagaaataaa aatattgaac tcat                               1034
    
```

<210> SEQ ID NO 82  
 <211> LENGTH: 245  
 <212> TYPE: PRT

-continued

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 82

```

Met Gly Arg Leu Asp Gly Lys Val Ile Ile Leu Thr Ala Ala Ala
 1          5          10          15
Gln Gly Ile Gly Gln Ala Ala Ala Leu Ala Phe Ala Arg Glu Gly
 20          25          30
Ala Lys Val Ile Ala Thr Asp Ile Asn Glu Ser Lys Leu Gln Glu
 35          40          45
Leu Glu Lys Tyr Pro Gly Ile Gln Thr Arg Val Leu Asp Val Thr
 50          55          60
Lys Lys Lys Gln Ile Asp Gln Phe Ala Ser Glu Val Glu Arg Leu
 65          70          75
Asp Val Leu Phe Asn Val Ala Gly Phe Val His His Gly Thr Val
 80          85          90
Leu Asp Cys Glu Glu Lys Asp Trp Asp Phe Ser Met Asn Leu Asn
 95          100         105
Val Arg Ser Met Tyr Leu Met Ile Lys Ala Phe Leu Pro Lys Met
 110         115         120
Leu Ala Gln Lys Ser Gly Asn Ile Ile Asn Met Ser Ser Val Ala
 125         130         135
Ser Ser Val Lys Gly Val Val Asn Arg Cys Val Tyr Ser Thr Thr
 140         145         150
Lys Ala Ala Val Ile Gly Leu Thr Lys Ser Leu Ala Ala Asp Phe
 155         160         165
Ile Gln Gln Gly Ile Arg Cys Asn Cys Val Cys Pro Gly Thr Val
 170         175         180
Asp Thr Pro Ser Leu Gln Glu Arg Ile Gln Ala Arg Gly Asn Pro
 185         190         195
Glu Glu Ala Arg Asn Asp Phe Leu Lys Arg Gln Lys Thr Gly Arg
 200         205         210
Phe Ala Thr Ala Glu Glu Ile Ala Met Leu Cys Val Tyr Leu Ala
 215         220         225
Ser Asp Glu Ser Ala Tyr Val Thr Gly Asn Pro Val Ile Ile Asp
 230         235         240
Gly Gly Trp Ser Leu
 245

```

<210> SEQ ID NO 83

<211> LENGTH: 1961

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 83

```

gggcggcgcc ggcagcgggt ggaggttgta ggaccggcga ggaataggaa           50
tcatggcgcc tgcgctgttc gtgctgctgg gattcgcgct gctgggcacc           100
cacggagcct ccggggctgc cggttcgctc caggcggcgc tgtcccagca           150
gaggtgggtg gggggcagtg tggagctgca ctgagggcc gtgggcagcc           200
cgggtcccga gatccagtgg tggtttgaag ggcagggtcc caacgacacc           250
tgctcccagc tctgggacgg cgcccggctg gaccgcgtcc acatccacgc           300
cacctaccac cagcacgcgg ccagcaccat ctccatcgac acgctcgtgg           350

```

-continued

---

aggaggacac gggcacttac gagtgccggg ccagcaacga cccggatcgc	400
aaccaccta cccgggcgcc cagggtaag tgggtccg cccaggcagt	450
cgtgctagtc ctggaaccg gcacagtctt cactaccgta gaagacctg	500
gtccaagat actectcacc tgctcctga atgacagcgc cacagaggtc	550
acagggcacc gctggctgaa gggggcgctg gtgctgaagg aggacgcct	600
gcccggccag aaaacggagt tcaagggtga ctccgacgac cagtggggag	650
agtactcctg cgtcttctc cccgagccca tgggcacggc caacatccag	700
ctccacgggc ctccagagt gaaggctgtg aagtctcag aacacatcaa	750
cgagggggag acggccatgc tggctctgcaa gtcagagtcc gtgccacctg	800
tcaactgactg ggctctgtac aagatcaactg actctgagga caaggccctc	850
atgaacggct ccgagagcag gttctctctg agttctctgc agggccggtc	900
agagctacac attgagaacc tgaacatgga ggccgacccc ggccagtacc	950
ggtgcaacgg caccagctcc aagggtctcc accaggccat catcaccgtc	1000
cgcgtgcgca gccacctggc cgcctctctg ccttctctgg gcatcgtggc	1050
tgaggtgctg gtgctgtgca ccatcatctt catctacgag aagcgcgga	1100
agcccgagga cgtcctggat gatgacgacg ccggctctgc acccctgaag	1150
agcagcgggg agcaccagaa tgacaaggc aagaacgtcc gccagaggaa	1200
ctcttctgta ggcaggtggc ccgagagcgc tccctgctcc acgtctgcgc	1250
cgccgccgga gtccactccc agtgcttgca agattccaag ttctcaacct	1300
ttaaagaaaa cccaccctgt agattcccat catacacttc cttotTTTT	1350
aaaaaagtgg ggttttctcc attcaggatt ctgttcotta ggtttttttc	1400
cttctgaagt gtttcaagag agcccgggag ctgctgccct gcggccccgt	1450
ctgtggcttt cagcctctgg gtctgagtca tggccgggtg ggcggcacag	1500
ccttctccac tggccggagt cagtgccagg tccttgccct ttgtgaaaag	1550
tcacaggta cacgaggggc cccgtgtcct gcctgtctga agccaatgct	1600
gtctggttgc gccatttttg tgcttttatg ttaatttta tgagggccac	1650
gggtctgtgt tcgactcagc ctccagggag actctgacct cttggccaca	1700
gaggactcac ttgcccacac cgagggcgac cccgtcacag cctcaagtca	1750
ctcccaagcc ccctccttgt ctgtgcatcc gggggcagct ctggaggggg	1800
tttctgggg aactggcgcc atcgccggga ctccagaacc gcagaagcct	1850
ccccagctca cccctggagg acggccggct ctctatagca ccagggctca	1900
cgtgggaacc cccctccac ccaccgccac aataaagatc gccccacct	1950
ccacccaaaa a	1961

&lt;210&gt; SEQ ID NO 84

&lt;211&gt; LENGTH: 385

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 84

Met Ala Ala Ala Leu Phe Val Leu Leu Gly Phe Ala Leu Leu Gly

-continued

1	5	10	15
Thr His Gly Ala Ser	Gly Ala Ala Gly	Phe Val Gln Ala Pro	Leu
	20	25	30
Ser Gln Gln Arg Trp	Val Gly Gly Ser	Val Glu Leu His Cys	Glu
	35	40	45
Ala Val Gly Ser Pro	Val Pro Glu Ile	Gln Trp Trp Phe Glu	Gly
	50	55	60
Gln Gly Pro Asn Asp	Thr Cys Ser Gln	Leu Trp Asp Gly Ala	Arg
	65	70	75
Leu Asp Arg Val His	Ile His Ala Thr	Tyr His Gln His Ala	Ala
	80	85	90
Ser Thr Ile Ser Ile	Asp Thr Leu Val	Glu Glu Asp Thr Gly	Thr
	95	100	105
Tyr Glu Cys Arg Ala	Ser Asn Asp Pro	Asp Arg Asn His Leu	Thr
	110	115	120
Arg Ala Pro Arg Val	Lys Trp Val Arg	Ala Gln Ala Val Val	Leu
	125	130	135
Val Leu Glu Pro Gly	Thr Val Phe Thr	Thr Val Glu Asp Leu	Gly
	140	145	150
Ser Lys Ile Leu Leu	Thr Cys Ser Leu	Asn Asp Ser Ala Thr	Glu
	155	160	165
Val Thr Gly His Arg	Trp Leu Lys Gly	Gly Val Val Leu Lys	Glu
	170	175	180
Asp Ala Leu Pro Gly	Gln Lys Thr Glu	Phe Lys Val Asp Ser	Asp
	185	190	195
Asp Gln Trp Gly Glu	Tyr Ser Cys Val	Phe Leu Pro Glu Pro	Met
	200	205	210
Gly Thr Ala Asn Ile	Gln Leu His Gly	Pro Pro Arg Val Lys	Ala
	215	220	225
Val Lys Ser Ser Glu	His Ile Asn Glu	Gly Glu Thr Ala Met	Leu
	230	235	240
Val Cys Lys Ser Glu	Ser Val Pro Pro	Val Thr Asp Trp Ala	Trp
	245	250	255
Tyr Lys Ile Thr Asp	Ser Glu Asp Lys	Ala Leu Met Asn Gly	Ser
	260	265	270
Glu Ser Arg Phe Phe	Val Ser Ser Ser	Gln Gly Arg Ser Glu	Leu
	275	280	285
His Ile Glu Asn Leu	Asn Met Glu Ala	Asp Pro Gly Gln Tyr	Arg
	290	295	300
Cys Asn Gly Thr Ser	Ser Lys Gly Ser	Asp Gln Ala Ile Ile	Thr
	305	310	315
Leu Arg Val Arg Ser	His Leu Ala Ala	Leu Trp Pro Phe Leu	Gly
	320	325	330
Ile Val Ala Glu Val	Leu Val Leu Val	Thr Ile Ile Phe Ile	Tyr
	335	340	345
Glu Lys Arg Arg Lys	Pro Glu Asp Val	Leu Asp Asp Asp Ala	Ala
	350	355	360
Gly Ser Ala Pro Leu	Lys Ser Ser Gly	Gln His Gln Asn Asp	Lys
	365	370	375
Gly Lys Asn Val Arg	Gln Arg Asn Ser	Ser	
	380	385	



-continued

<210> SEQ ID NO 85  
 <211> LENGTH: 1002  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 85

```

ggctcgagca aagacatacg aacagggagg aaggccgact gaaagaaaga      50
cggagaagag gagagagaag ccagggccga gcgtgccagc aggcggatgg      100
agggcgccct ggtggaggag gagacgtagt ggcctgggct gagctgggtg      150
ggccgggaga agcgggtgcc tcagagtggg ggtgggggca tgggaggggc      200
aggcattctg ctgctgctgc tggctggggc gggggtggtg gtggcctgga      250
gaccccaaaa gggaaagtgt cccctgcgct gctcctgctc taaagacagc      300
gccctgtgtg agggctcccc ggacctgccc gtcagcttct ctccgacct      350
gctgtcactc tcactcgtca ggacgggagt caccagctg aaggccggca      400
gcttctctgag aattccgtct ctgcacctgc tcctottcac ctccaaactc      450
ttctccgtga ttgaggacga tgcatttgcg ggcctgtccc acctgcagta      500
cctcttcatac gaggacaatg agattggctc catctctaag aatgccctca      550
gaggactctg ctgccttaca cacctaagcc tggccaataa ccatctggag      600
accctcccca gattcctgtt ccgaggcctg gacaccctta ctacgtgga      650
cctccgcggg aacccttcc agtgtgactg ccgctcctc tggctcctgc      700
agtgatgacc caccgtgaat gccagcgtgg ggaccggcgc ctgtgcgggc      750
cccgcctccc tgagccacat gcagctccac cacctcgacc ccaagacttt      800
caatgacaga gccatagggtg gggggctttc ccgatggggt gggaggcggg      850
agatctgggg gaaaggctgc cagggccaag aggctcgtct cactccctgc      900
cctgccattt cccggagtgg gaagaccctg agcaagcagc actgccttcc      950
tgagcccagc ttttctcctc tgtaaagtgg gggtaataaa cagtgatata     1000
gg                                                                1002
    
```

<210> SEQ ID NO 86  
 <211> LENGTH: 261  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 86

```

Met Gly Gly Ala Gly Ile Leu Leu Leu Leu Ala Gly Ala Gly
 1           5           10          15
Val Val Val Ala Trp Arg Pro Pro Lys Gly Lys Cys Pro Leu Arg
 20          25          30
Cys Ser Cys Ser Lys Asp Ser Ala Leu Cys Glu Gly Ser Pro Asp
 35          40          45
Leu Pro Val Ser Phe Ser Pro Thr Leu Leu Ser Leu Ser Leu Val
 50          55          60
Arg Thr Gly Val Thr Gln Leu Lys Ala Gly Ser Phe Leu Arg Ile
 65          70          75
Pro Ser Leu His Leu Leu Leu Phe Thr Ser Asn Ser Phe Ser Val
 80          85          90
    
```

-continued

Ile Glu Asp Asp Ala Phe Ala Gly Leu Ser His Leu Gln Tyr Leu  
 95 100 105  
 Phe Ile Glu Asp Asn Glu Ile Gly Ser Ile Ser Lys Asn Ala Leu  
 110 115 120  
 Arg Gly Leu Arg Ser Leu Thr His Leu Ser Leu Ala Asn Asn His  
 125 130 135  
 Leu Glu Thr Leu Pro Arg Phe Leu Phe Arg Gly Leu Asp Thr Leu  
 140 145 150  
 Thr His Val Asp Leu Arg Gly Asn Pro Phe Gln Cys Asp Cys Arg  
 155 160 165  
 Val Leu Trp Leu Leu Gln Trp Met Pro Thr Val Asn Ala Ser Val  
 170 175 180  
 Gly Thr Gly Ala Cys Ala Gly Pro Ala Ser Leu Ser His Met Gln  
 185 190 195  
 Leu His His Leu Asp Pro Lys Thr Phe Lys Cys Arg Ala Ile Gly  
 200 205 210  
 Gly Gly Leu Ser Arg Trp Gly Gly Arg Arg Glu Ile Trp Gly Lys  
 215 220 225  
 Gly Cys Gln Gly Gln Glu Ala Arg Leu Thr Pro Cys Pro Ala Ile  
 230 235 240  
 Ser Arg Ser Gly Lys Thr Leu Ser Lys Gln His Cys Leu Pro Glu  
 245 250 255  
 Pro Gln Phe Ser His Leu  
 260

<210> SEQ ID NO 87  
 <211> LENGTH: 2945  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 87

cggacgctg gggcggcgg agcagctgca gttcgcatct caggcagtac 50  
 cttagaggagc tgccggtgcc tcctcagaac atctcctgat cgctacccag 100  
 gaccaggcac caaggacagg gagtcccagg cgcacacccc ccattctggg 150  
 tccccaggc ccagaccccc actctgccac aggttgcatc ttgacctggt 200  
 cctcctgcaag aagtggcccc tgtggtcctg ctctgagact cgtccctggg 250  
 cgcccctgca gccccctttct atgactccat ctggatttgg ctggctgtgg 300  
 ggacgctgct cgaggggctg cctggtcttc agcgtggtgg cagccagctc 350  
 tctggccacc atggcaaatg ctgagatctg aggggacaag gctctacagc 400  
 cttagccagg ggcactcagc tgttgacagg tgtgatggag aacaaagcta 450  
 tgtacctaca caccgtcagc gactgtgaca ccagctccat ctgtgaggat 500  
 tcctttgatg gcaggagcct gtccaagctg aacctgtgtg aggatggtcc 550  
 atgtcacaaa cggcgggcaa gcatctgctg taccagctg gggccctgt 600  
 cggccctgaa gcatgctgtc ctggggctct acctgctggt cttcctgatt 650  
 cttgtgggca tcttcatctt agcagggcca cggggacca aaggatgatca 700  
 ggggatgtaa gaaaggaag gcaggcctgg catcctgga ttgctggac 750  
 ttcagaggtc gcccggggag agaggtacc caggattgcc cgggcccaag 800

-continued

---

ggcgatgatg ggaagctggg ggccacagga ccaatgggca tgcgtgggtt	850
caaaggtgac cgagcccaa aaggagagaa aggagagaaa ggagacagag	900
ctggggatgc cagtggcgtg gaggccccga tgatgatccg cctgggtgaat	950
ggctcaggtc cgcacgaggg ccgctgggaa gtgtaccacg accggcgctg	1000
gggcaccgtg tgtgacgacg gctgggacaa gaaggacgga gacgtgggtg	1050
gccgcatgct cggcttccgc ggtgtggagg aggtgtaccg cacagctcga	1100
ttcgggcaag gcaactggag gatctggatg gatgacgttg cctgcaaggg	1150
cacagaggaa accatcttcc gctgcagctt ctccaaatgg ggggtgacaa	1200
actgtggaca tgccgaagat gccagcgtga catgcaacag aactgaaaag	1250
tgggcagagc ccaagttcgg ggtcctgcac agagcacctt tgctgcatcc	1300
ctgggggtgg gcacagctcg gggccaccct gaccatgcct cgaccacacc	1350
ccgtccagca ttctcagtcc tcacacctgc atcccaggac cgtggggggc	1400
ggtcgtcatt tccctcttga acatgtgctc cgaagtataa ctctgggacc	1450
tactgccctc ctctctcttc caccaggttc ctgcatgagg agcctgatc	1500
aactggatca ccactttgcc cagcctctga acaccatgca ccaggcctca	1550
atatcccagt tccctttggc cttttagtta caggtgaatg ctgagaatgt	1600
gtcagagaca agtgacgacg cagcgtatgt tggtagtata gatcatttac	1650
tcttcagaca attcccaaac ctccattagt ccaagagttt ctacatcttc	1700
ctccccagca agaggcaacg tcaagtgatg aatttcccc ctttactctg	1750
cctctgctcc ccatttgcta gtttgaggaa gtgacataga ggagaagcca	1800
gctgtagggg caagagggaa atgcaagtca cctgcaggaa tccagctaga	1850
tttgagaga ggaatgaaac taacattgaa tgactaccat ggcacgctaa	1900
atagtatctt ggggtccaaa ttcattgtat cacttagctg cattgggtcca	1950
gggcatgta gtctggatac agccttacct tcaggtagca cttaactggg	2000
ccattcacct agactgcaag taagaagaca aaatgactga gaccgtgtgc	2050
ccacctgaac ttattgtctt taactggcct gagctaaaag cttgggtgca	2100
ggacctgtgt aactagaaag ttgcctactt cagaacctcc agggcgtgag	2150
tgcaaggta aacatgactg gcttccaggc cgaccatcaa tgtaggagga	2200
gagctgatgt ggaggggtgac atgggggctg cccatgttaa acctgagtc	2250
agtgtctggt cattgggcag tcacggttaa agccaagtca tgtgtgtctc	2300
agctgtttgg aggtgatgat tttgcatctt ccaagcctct tcagggtgga	2350
atctgtggtc aggaaaacac aagtcctaata ggaaccctta ggggggaagg	2400
aaatgaagat tccctataac ctctgggggt ggggagtagg aataaggggc	2450
cttgggcctc cataaatctg caatctgcac cctcctccta gagacagggg	2500
gatcgtgttc tgctttttac atgaggagca gaactgggcc atacacgtgt	2550
tcaagaacta ggggagctac ctggtagcaa gtgagtgcag acccacctca	2600
ccttggggga atctcaaaact cataggcctc agatacacga tcaactgtca	2650
tatcaggtga gcaactggcct gcttggggag agacctgggc ccctccaggt	2700

-continued

---

```

gttagaacag caacactcct ggctgacaac taagccaata tggccctag 2750
tcattcttgc ttccaatgat ctggcactc cttaaatgtc ctaatgatga 2800
gaaactctct ttctgaccaa ttgctatggt tacataacac gcatgtactc 2850
atgcatccct tgccagagcc catatatgta tgcataatata aacatagcac 2900
tttttactac atagctcagc acattgcaag gtttgcatth aagtt 2945

```

```

<210> SEQ ID NO 88
<211> LENGTH: 270
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

```

```

<400> SEQUENCE: 88

```

```

Met Glu Asn Lys Ala Met Tyr Leu His Thr Val Ser Asp Cys Asp
 1          5          10          15
Thr Ser Ser Ile Cys Glu Asp Ser Phe Asp Gly Arg Ser Leu Ser
 20          25          30
Lys Leu Asn Leu Cys Glu Asp Gly Pro Cys His Lys Arg Arg Ala
 35          40          45
Ser Ile Cys Cys Thr Gln Leu Gly Ser Leu Ser Ala Leu Lys His
 50          55          60
Ala Val Leu Gly Leu Tyr Leu Leu Val Phe Leu Ile Leu Val Gly
 65          70          75
Ile Phe Ile Leu Ala Gly Pro Pro Gly Pro Lys Gly Asp Gln Gly
 80          85          90
Asp Glu Gly Lys Glu Gly Arg Pro Gly Ile Pro Gly Leu Pro Gly
 95          100         105
Leu Arg Gly Leu Pro Gly Glu Arg Gly Thr Pro Gly Leu Pro Gly
 110         115         120
Pro Lys Gly Asp Asp Gly Lys Leu Gly Ala Thr Gly Pro Met Gly
 125         130         135
Met Arg Gly Phe Lys Gly Asp Arg Gly Pro Lys Gly Glu Lys Gly
 140         145         150
Glu Lys Gly Asp Arg Ala Gly Asp Ala Ser Gly Val Glu Ala Pro
 155         160         165
Met Met Ile Arg Leu Val Asn Gly Ser Gly Pro His Glu Gly Arg
 170         175         180
Val Glu Val Tyr His Asp Arg Arg Trp Gly Thr Val Cys Asp Asp
 185         190         195
Gly Trp Asp Lys Lys Asp Gly Asp Val Val Cys Arg Met Leu Gly
 200         205         210
Phe Arg Gly Val Glu Glu Val Tyr Arg Thr Ala Arg Phe Gly Gln
 215         220         225
Gly Thr Gly Arg Ile Trp Met Asp Asp Val Ala Cys Lys Gly Thr
 230         235         240
Glu Glu Thr Ile Phe Arg Cys Ser Phe Ser Lys Trp Gly Val Thr
 245         250         255
Asn Cys Gly His Ala Glu Asp Ala Ser Val Thr Cys Asn Arg His
 260         265         270

```

```

<210> SEQ ID NO 89
<211> LENGTH: 2758

```

---

-continued

---

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 89

gtcgccgcga gggacgcaga gaggaccctc cacgcccaga tgctgcgta 50  
gtttttgtga ccagtccgct cctgcctccc cctggggcag tagaggggga 100  
gcgatggaga actggactgg caggccctgg ctgtatctgc tgctgcttct 150  
gtccctccct cagctctgct tggatcagga ggtgtgtcc ggacactctc 200  
ttcagacacc tacagaggag ggccagggcc cgaaggtgt ctggggacct 250  
tgggtccagt gggcctcttg ctcccagccc tgcggggtgg gggtcagcg 300  
caggagccgg acatgtcagc tccctacagt gcagctccac ccgagtctgc 350  
ccctccctcc ccggccccc agacatccag aagcctcct cccccgggc 400  
cagggtccca gaccccagac ttctccagaa accctcccct tgtacaggac 450  
acagtctcgg ggaaggggtg gccactctg aggtcccgt tcccacctag 500  
ggagagagga gaccaggag attcgagcgg ccaggaggtc ccggcttcca 550  
gaccccatca agccaggaat gttcggttat gggagagtgc cctttgcatt 600  
gccactgcac cggaaccgca ggcaccctcg gagcccacc agatctgagc 650  
tgtccctgat ctcttctaga ggggaagagg ctattccgtc ccetactcca 700  
agagcagagc ctttctccgc aaacggcagc ccccaaactg agctccctcc 750  
cacagaactg tctgtccaca ccccacccc ccaagcagaa cctctaagcc 800  
ctgaaaactg tcagacagag gtggccccc gaaccaggcc tgcccccta 850  
cggcatcacc ccagagccca ggctctggc acagagcccc cctaccccac 900  
gcactcctta ggagaagggtg gcttcttccg tgcacccct cagccacgaa 950  
ggccaagtcc ccagggttgg gccagtcccc aggtagcagg gagacgccct 1000  
gatccttttc cttcgggtccc tcggggccga ggccagcagg gccaaaggcc 1050  
ttggggaacg ggggggactc ctcaagggcc ccgctggag cctgaccctc 1100  
agcaccgggg cgcctggctg cccctgctga gcaacggccc ccatgccagc 1150  
tccctctgga gcctctttgc tcccagtagc cctattcaa gatgttctgg 1200  
ggagagttaa cagctaagag cctgcagcca agcgcctgc ccccctgagc 1250  
agccagaccc ccgggccctg cagtgcgcag cctttaactc ccaggaattc 1300  
atgggccagc tgtatcagtg ggagcccttc actgaagtcc agggctccca 1350  
gcgctgtgaa ctgaactgcc ggcccgttg cttccgttc tatgtccgtc 1400  
aactgaaaa ggtccaggat gggaccctgt gtcagcctgg agcccctgac 1450  
atctgtgtgg ctggacgctg tctgagcccc ggctgtgatg ggatccttgg 1500  
ctctggcagg cgtcctgatg gctgtggagt ctgtgggggt gatgattcta 1550  
cctgtcgcct tgtttcgggg aaactcactg accgaggggg ccccctgggc 1600  
tatcagaaga tcttgtggat tccagcggga gccttgccgc tccagattgc 1650  
ccagctccgg cctagctcca actacctgc acttcgtggc cctgggggcc 1700  
ggtccatcat caatgggaac tgggctgtgg atccccctgg gtcctacag 1750  
gccggcggga ccgtctttcg atataaccgt cctcccaggg aggagggcaa 1800

-continued

---

```

aggggagagt ctgtcggctg aaggcccccac caccagcct gtggatgtct      1850
atatgatctt tcaggaggaa aaccaggcg tttttatca gtatgtcatc      1900
tcttcacctc ctccaatcct tgagaacccc accccagagc ccctgtccc      1950
ccagctttag ccggagattc tgagggtgga gccccactt gtcocggcac      2000
cccggccagc ccggacccca ggcaccctcc agcgtcaggt gcggatcccc      2050
cagatgcccg ccccgcccca tcccaggaca cccctggggt ctccagctgc      2100
gtactggaaa cgagtgggac actctgcatg ctcagcgtcc tgcgggaaag      2150
gtgtctggcg ccccattttc ctctgcatct cccgtgagtc gggagaggaa      2200
ctggatgaac gcagctgtgc cgcgggtgcc aggccccag cctcccctga      2250
accctgccac ggcaccccat gcccccata ctgggaggct ggcgagtgga      2300
catcctgcag ccgctcctgt ggccccgca cccagcaccg ccagctgcag      2350
tgccggcagg aatttggggg ggggtgctcc tcggtgcccc cggagcgtg      2400
tggacatctc ccccgccca acatcacca gtcttgccag ctgcccctct      2450
gtggccattg ggaagttggc tctccttga gccagtgtc cgtgcggtgc      2500
ggccggggcc agagaagccg gcaggttcgc tgtgttggga acaacggtga      2550
tgaagtgagc gagcaggagt gtgctcagg cccccacag cccccagca      2600
gagagccctg tgacatgggg ccctgtacta ctgcttggtt ccaacagcag      2650
tggagctcca aggtgagccc ggaacccca gccatatect gcatcctggg      2700
taaccatgcc caggacacct cagcctttcc agcatagctc aataaacttg      2750
tattgatc      2758

```

&lt;210&gt; SEQ ID NO 90

&lt;211&gt; LENGTH: 877

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 90

```

Met Glu Asn Trp Thr Gly Arg Pro Trp Leu Tyr Leu Leu Leu Leu
 1          5          10          15
Leu Ser Leu Pro Gln Leu Cys Leu Asp Gln Glu Val Leu Ser Gly
 20          25          30
His Ser Leu Gln Thr Pro Thr Glu Glu Gly Gln Gly Pro Glu Gly
 35          40          45
Val Trp Gly Pro Trp Val Gln Trp Ala Ser Cys Ser Gln Pro Cys
 50          55          60
Gly Val Gly Val Gln Arg Arg Ser Arg Thr Cys Gln Leu Pro Thr
 65          70          75
Val Gln Leu His Pro Ser Leu Pro Leu Pro Pro Arg Pro Pro Arg
 80          85          90
His Pro Glu Ala Leu Leu Pro Arg Gly Gln Gly Pro Arg Pro Gln
 95          100         105
Thr Ser Pro Glu Thr Leu Pro Leu Tyr Arg Thr Gln Ser Arg Gly
 110         115         120
Arg Gly Gly Pro Leu Arg Gly Pro Ala Ser His Leu Gly Arg Glu
 125         130         135

```

-continued

---

Glu Thr Gln Glu Ile Arg Ala Ala Arg Arg Ser Arg Leu Arg Asp	140	145	150
Pro Ile Lys Pro Gly Met Phe Gly Tyr Gly Arg Val Pro Phe Ala	155	160	165
Leu Pro Leu His Arg Asn Arg Arg His Pro Arg Ser Pro Pro Arg	170	175	180
Ser Glu Leu Ser Leu Ile Ser Ser Arg Gly Glu Glu Ala Ile Pro	185	190	195
Ser Pro Thr Pro Arg Ala Glu Pro Phe Ser Ala Asn Gly Ser Pro	200	205	210
Gln Thr Glu Leu Pro Pro Thr Glu Leu Ser Val His Thr Pro Ser	215	220	225
Pro Gln Ala Glu Pro Leu Ser Pro Glu Thr Ala Gln Thr Glu Val	230	235	240
Ala Pro Arg Thr Arg Pro Ala Pro Leu Arg His His Pro Arg Ala	245	250	255
Gln Ala Ser Gly Thr Glu Pro Pro Ser Pro Thr His Ser Leu Gly	260	265	270
Glu Gly Gly Phe Phe Arg Ala Ser Pro Gln Pro Arg Arg Pro Ser	275	280	285
Ser Gln Gly Trp Ala Ser Pro Gln Val Ala Gly Arg Arg Pro Asp	290	295	300
Pro Phe Pro Ser Val Pro Arg Gly Arg Gly Gln Gln Gly Gln Gly	305	310	315
Pro Trp Gly Thr Gly Gly Thr Pro His Gly Pro Arg Leu Glu Pro	320	325	330
Asp Pro Gln His Pro Gly Ala Trp Leu Pro Leu Leu Ser Asn Gly	335	340	345
Pro His Ala Ser Ser Leu Trp Ser Leu Phe Ala Pro Ser Ser Pro	350	355	360
Ile Pro Arg Cys Ser Gly Glu Ser Glu Gln Leu Arg Ala Cys Ser	365	370	375
Gln Ala Pro Cys Pro Pro Glu Gln Pro Asp Pro Arg Ala Leu Gln	380	385	390
Cys Ala Ala Phe Asn Ser Gln Glu Phe Met Gly Gln Leu Tyr Gln	395	400	405
Trp Glu Pro Phe Thr Glu Val Gln Gly Ser Gln Arg Cys Glu Leu	410	415	420
Asn Cys Arg Pro Arg Gly Phe Arg Phe Tyr Val Arg His Thr Glu	425	430	435
Lys Val Gln Asp Gly Thr Leu Cys Gln Pro Gly Ala Pro Asp Ile	440	445	450
Cys Val Ala Gly Arg Cys Leu Ser Pro Gly Cys Asp Gly Ile Leu	455	460	465
Gly Ser Gly Arg Arg Pro Asp Gly Cys Gly Val Cys Gly Gly Asp	470	475	480
Asp Ser Thr Cys Arg Leu Val Ser Gly Asn Leu Thr Asp Arg Gly	485	490	495
Gly Pro Leu Gly Tyr Gln Lys Ile Leu Trp Ile Pro Ala Gly Ala	500	505	510
Leu Arg Leu Gln Ile Ala Gln Leu Arg Pro Ser Ser Asn Tyr Leu			

-continued

---

	515		520		525
Ala Leu Arg Gly Pro	Gly Gly Arg Ser	Ile Ile Asn Gly Asn Trp	530	535	540
Ala Val Asp Pro Pro	Gly Ser Tyr Arg	Ala Gly Gly Thr Val Phe	545	550	555
Arg Tyr Asn Arg Pro	Pro Arg Glu Glu	Gly Lys Gly Glu Ser Leu	560	565	570
Ser Ala Glu Gly Pro	Thr Thr Gln Pro	Val Asp Val Tyr Met Ile	575	580	585
Phe Gln Glu Glu Asn	Pro Gly Val Phe	Tyr Gln Tyr Val Ile Ser	590	595	600
Ser Pro Pro Pro Ile	Leu Glu Asn Pro	Thr Pro Glu Pro Pro Val	605	610	615
Pro Gln Leu Gln Pro	Glu Ile Leu Arg	Val Glu Pro Pro Leu Ala	620	625	630
Pro Ala Pro Arg Pro	Ala Arg Thr Pro	Gly Thr Leu Gln Arg Gln	635	640	645
Val Arg Ile Pro Gln	Met Pro Ala Pro	Pro His Pro Arg Thr Pro	650	655	660
Leu Gly Ser Pro Ala	Ala Tyr Trp Lys	Arg Val Gly His Ser Ala	665	670	675
Cys Ser Ala Ser Cys	Gly Lys Gly Val	Trp Arg Pro Ile Phe Leu	680	685	690
Cys Ile Ser Arg Glu	Ser Gly Glu Glu	Leu Asp Glu Arg Ser Cys	695	700	705
Ala Ala Gly Ala Arg	Pro Pro Ala Ser	Pro Glu Pro Cys His Gly	710	715	720
Thr Pro Cys Pro Pro	Tyr Trp Glu Ala	Gly Glu Trp Thr Ser Cys	725	730	735
Ser Arg Ser Cys Gly	Pro Gly Thr Gln	His Arg Gln Leu Gln Cys	740	745	750
Arg Gln Glu Phe Gly	Gly Gly Gly Ser	Ser Val Pro Pro Glu Arg	755	760	765
Cys Gly His Leu Pro	Arg Pro Asn Ile	Thr Gln Ser Cys Gln Leu	770	775	780
Arg Leu Cys Gly His	Trp Glu Val Gly	Ser Pro Trp Ser Gln Cys	785	790	795
Ser Val Arg Cys Gly	Arg Gly Gln Arg	Ser Arg Gln Val Arg Cys	800	805	810
Val Gly Asn Asn Gly	Asp Glu Val Ser	Glu Gln Glu Cys Ala Ser	815	820	825
Gly Pro Pro Gln Pro	Pro Ser Arg Glu	Ala Cys Asp Met Gly Pro	830	835	840
Cys Thr Thr Ala Trp	Phe His Ser Asp	Trp Ser Ser Lys Val Ser	845	850	855
Pro Glu Pro Pro Ala	Ile Ser Cys Ile	Leu Gly Asn His Ala Gln	860	865	870
Asp Thr Ser Ala Phe	Pro Ala		875		



-continued

&lt;211&gt; LENGTH: 2597

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 91

cgagtatttt cccaccatct ccagccgaa actgaccaag aactctgagg	50
cggatggcat gttcgcgtac gtcttccatg atgagttcgt ggccctcgatg	100
attaagatcc cttcggacac cttcaccatc atccctgact ttgatatcta	150
ctatgtctat ggttttagca gtggcaactt tgtctacttt ttgaccctcc	200
aacctgagat ggtgtctcca ccaggtcca ccaccaagga gcagggtgat	250
acatccaagc tcgtgaggct ttgcaaggag gacacagcct tcaactccta	300
tgtagaggtg ccattggct gtgagcgag tggggggag tacgcctgc	350
tgcaggctgc ctacctgtcc aaagcgggg cgtgcttg caggaccctt	400
ggagtccatc cagatgatga cctgtcttc accgtcttct ccaaggcca	450
gaagcggaaa atgaaatccc tggatgagtc ggccctgtgc atcttcatct	500
tgaagcagat aaatgaccgc attaaggagc ggctgcagtc ttgttaccgg	550
ggcgaggcca cgctggacct ggccctggctc aaggtgaagg acatcccctg	600
cagcagtggc ctcttaacca ttgacgataa cttctgtggc ctggacatga	650
atgtctccct gggagtgtcc gacatggctc gtggaattcc cgtcttcacg	700
gaggacaggc accgcatgac gtctgtcatc gcataatgtct acaagaacca	750
ctctctggcc tttgtgggca ccaaaagtgg caagctgaag aaggtgctg	800
gtaccagcct ctgccctacc cttgagctac agacgggacc ccgatcccac	850
agagcaacag tgactctgga actcctgttc tccagctgtt catcaactg	900
agaaaaactt cagagctgtg taggcttatt tagtgtgttg tcagccttg	950
atatttgaaa atggaacag atgagacaca tctacctccc tgtgacccca	1000
gccatacatc atagctcatg tcctgccacc ccaagtcctt agggaaaaaa	1050
gactttggag aatgtgtctc tgccttagctt ggctaggtag ttggtctctt	1100
ttctctgccc caagcgtccc ctgggtaatt ttgacaatg gagtgtaggc	1150
atgtttgact cttgtggtgt tatcacttgt atatgtcagt gaaactaact	1200
gattctccca tcggaatata gttatctctt gggcctgata tatggttagga	1250
taaccttatg ctcatctgtc cacttctgca gccaaagtcgc ctggccagtg	1300
tgtgtgtgtg tgtgtgtgtg tgtgtgtgtg tgtgtgtatg cttatctgtg	1350
tttaagggtg tgtgtgcata cacagggcag agaggatgga gccaccgta	1400
ctgcagcatc atgtaattaa ctcagtgctc agaaccatcc cagcctctgc	1450
gggaaagaga aaagtaagcc aacagtcct gatgagctga tcatatgtgc	1500
aaaagctctg ttggcatctg gtccaggaga gcacccaaaa aaagttaatt	1550
ggtgtgtgoc agtctccttt ccttaagact atggttacaa caaagcgtga	1600
gcagtgtctc ctgcatggcc actatccagc acaattccat aattccccca	1650
tagagccggt ggggaggagg aggtgagtg cgaaggaagt ggaaacactt	1700
ggtgtcatgt gctcctatca tttctactag cttactggga aataaagtgt	1750

-continued

---

```

agtcaagagt gtatgaaggc aagatgtaaa attagcgact ggtgctaadc      1800
tggttacttg aaaacaagtg aaagtgctgt agattgttc tgttgctaag      1850
aaccaccaca ctaaacctcg tatagttcct ggaggatata caacagtgta      1900
attctcttta ggggtgcca caggttcctg gctgtggga gggaatgaat      1950
caggagggct cttgagaacc ttcactctgt tgcttgact gaaagtgagt      2000
cccaaagctg gagatttagt gagagcaggc aaccctctg tgtctcactg      2050
tccatattct ggaggcagag gtttgaaca ggccatgtgc acctgcatag      2100
ggatgggtaa agcaaggact ttgaaagagt tgaaaagcat tataaacagt      2150
tgttcagaaa tacgtccag gagttccatg tgaactggc tctgtgtgca      2200
ttgaaagcat gctgttggga attctaactg gtccaacact cctgcaaac      2250
aatgtgtaa tatttaggaa gaaacttgaa aatagtcaa tcctttgaa      2300
tggtgacaat ttttaaga atcaattcta atttgttca agggtaataa      2350
tcaccaagat acacatttca gcatttattt agtctatcaa aaattggaat      2400
tgatatatac actcatttat aggagaatgg ttaggtagat ttggtatatt      2450
tatgtagtca ttgaaaactt agtttataaa ggccaatctt gtaactgatt      2500
cttgtgtgat aacattcagt gaaaagcat gagacaatta gaaagcatga      2550
tacaatgaat aaaataaaaa ctggaagag aaccatcaa atgctaa      2597

```

&lt;210&gt; SEQ ID NO 92

&lt;211&gt; LENGTH: 280

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 92

```

Met Phe Ala Tyr Val Phe His Asp Glu Phe Val Ala Ser Met Ile
  1           5           10          15
Lys Ile Pro Ser Asp Thr Phe Thr Ile Ile Pro Asp Phe Asp Ile
          20          25          30
Tyr Tyr Val Tyr Gly Phe Ser Ser Gly Asn Phe Val Tyr Phe Leu
          35          40          45
Thr Leu Gln Pro Glu Met Val Ser Pro Pro Gly Ser Thr Thr Lys
          50          55          60
Glu Gln Val Tyr Thr Ser Lys Leu Val Arg Leu Cys Lys Glu Asp
          65          70          75
Thr Ala Phe Asn Ser Tyr Val Glu Val Pro Ile Gly Cys Glu Arg
          80          85          90
Ser Gly Val Glu Tyr Arg Leu Leu Gln Ala Ala Tyr Leu Ser Lys
          95          100         105
Ala Gly Ala Val Leu Gly Arg Thr Leu Gly Val His Pro Asp Asp
          110         115         120
Asp Leu Leu Phe Thr Val Phe Ser Lys Gly Gln Lys Arg Lys Met
          125         130         135
Lys Ser Leu Asp Glu Ser Ala Leu Cys Ile Phe Ile Leu Lys Gln
          140         145         150
Ile Asn Asp Arg Ile Lys Glu Arg Leu Gln Ser Cys Tyr Arg Gly
          155         160         165
Glu Gly Thr Leu Asp Leu Ala Trp Leu Lys Val Lys Asp Ile Pro

```

-continued

	170		175		180
Cys Ser Ser Ala	Leu Leu Thr Ile Asp	Asp Asn Phe Cys Gly	Leu		
	185		190		195
Asp Met Asn Ala	Pro Leu Gly Val Ser	Asp Met Val Arg Gly	Ile		
	200		205		210
Pro Val Phe Thr	Glu Asp Arg Asp Arg	Met Thr Ser Val Ile	Ala		
	215		220		225
Tyr Val Tyr Lys	Asn His Ser Leu Ala	Phe Val Gly Thr Lys	Ser		
	230		235		240
Gly Lys Leu Lys	Lys Val Pro Gly Thr	Ser Leu Cys Pro Thr	Leu		
	245		250		255
Glu Leu Gln Thr	Gly Pro Arg Ser His	Arg Ala Thr Val Thr	Leu		
	260		265		270
Glu Leu Leu Phe	Ser Ser Cys Ser Ser	Asn			
	275		280		

<210> SEQ ID NO 93  
 <211> LENGTH: 2883  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 93

```

ccttatcaga caaaggacga gatggaaaat acaagataat ttacagtgga          50
gaagaattag aatgtaacct gaaagatctt agaccagcaa cagattatca          100
tgtgaggggtg tatgccatgt acaattccgt aaagggatcc tgctccgagc          150
ctgttagctt caccacccac agctgtgcac ccgagtgtcc tttccccctt          200
aagctggcac ataggagcaa aagttcacta accctgcagt ggaaggcacc          250
aattgacaac ggttcaaaaa tcaccaacta ccttttagag tgggatgagg          300
gaaaagaaa tagtggtttc agacagtgtc tcttcgggag ccagaagcac          350
tgcaagttag caaagctttg tccggcaatg ggggtacacat tcaggctggc          400
cgctcgaaac gacattggca ccagtgggta tagccaagag gtgggtgtgct          450
acacattagg aaatatccct cagatgcctt ctgcactaag gctggttcga          500
gctggcatca catgggtcac gttgcagtgg agtaagccag aaggctgttc          550
acccgaggaa gtgatcacct acaccttgga aattcaggag gatgaaaatg          600
ataacctttt ccacccaaaa tacactggag aggatttaac ctgtactgtg          650
aaaaatctca aaagaagcac acagtataaa ttcaggctga ctgcttctaa          700
tacggaagga aaaagctgtc caagcgaagt tcttgtttgt acgacgagtc          750
ctgacaggcc tggacctcct accagaccgc ttgtcaaagg cccagttaca          800
tctcatggct ttagtgtcaa atgggatccc cctaaggaca atgggtggtc          850
agaaatcctc aagtacttgc tagagattac tgatggaaat tctgaagcga          900
atcagtggga agtggcctac agtgggtcgg ctaccgaata caccttcacc          950
cacttgaaac caggcacttt gtacaaaactc cgagcatgct gcatcagtac          1000
cggcggacac agccagtgtt ctgaaagtct cctgttctgc aactaagca          1050
ttgaccaggg tcaatgtcga ccaccgaggg ttttgggtag accaaagcac          1100
aaagaagtcc acttagagtg ggatgttctc gcatcgaaa gtggctgtga          1150
    
```

-continued

---

ggtctcagag tacagcgtgg agatgacgga gccccaagac gtagcctcgg	1200
aagtgtacca tggcccagag ctggagtgca cgcgcggcaa cctgcttcct	1250
ggaaccgtgt atcgcttccg ggtgagggct ctgaatgatg gagggatgg	1300
tccctattct gatgtctcag aaattaccac tgctgcaggg cctcctggac	1350
aatgcaaac accttgatatt tcttgctcac ctgatggatg tgtcttagtg	1400
ggttgggaga gtcctgatag ttctgggtct gacatctcag agtacaggtt	1450
ggaatgggga gaagatgaag aatccttaga actcatttat catgggacag	1500
acacccgttt tgaataaaga gacctgttgc ctgctgcaca gtattgctgt	1550
agactacagg ccttcaatca agcaggggca gggccgtaca gtgaacttgt	1600
cctttgccag acgccagcgt ctgccctga ccccgctccc actctctgtg	1650
tcttgaggga ggagcccctt gatgcctacc ctgattcacc ttctgcgtgc	1700
cttgactgta actgggaaga gccgtgcaat aacggatctg aaatccttgc	1750
ttacaccatt gatctaggag aactagcat tacctggggc aacaccacca	1800
tgcatgttat gaaagatctc cttccagaaa ccacctaccg gatcagaatt	1850
caggctataa atgaaattgg agctggacca tttagtcagt tcattaagc	1900
aaaaactcgg ccattaccac ccttgctccc taggctagaa tgtgctgctg	1950
ctggctcctca gagcctgaag ctaaaatggg gagacagtaa ctccaagaca	2000
catgctgctg aggacattgt gtacacacta cagctggagg acagaaacaa	2050
gaggtttatt tcaatctaca gaggaccag ccacacctac aaggctccaga	2100
gactgacgga attcacatgc tactccttca gaatccaggc agcaagcgag	2150
gctggagaag ggcccttctc agaaacctat accttcagca caacaaaag	2200
tgccccccc accatcaaag cacctcagat aacacagtta gaagtaaatt	2250
catgtgaaat tttatgggag acggtagcat caatgaaag tgacctggt	2300
aactacattc tgcaggatatt ggttgaaga gaatctgagt acaaacaggt	2350
gtacaaggga gaagaagcca cattccaat ctcaggcctc cagaccaaca	2400
cagactacag gttccgcgta tgtgcgtgct gtcgctgttt agacacctct	2450
caggagctaa gcggagcctt cagcccctct gcggcttttg tattacaacg	2500
aagtgaggtc atgcttacag gggacatggg gagcttagat gatcccaaaa	2550
tgaagagcat gatgcctact gatgaacagt ttgcagccat cattgtgctt	2600
ggctttgcaa ctttgtccat tttatttggc tttatattac agtacttctt	2650
aatgaagtaa acccaacaaa actagaggta tgaattaatg ctacacattt	2700
taatacacac atttattcag atactcccct ttttaaagcc cttttgtttt	2750
ttgatttata tactctgttt tacagattta gctagaaaaa aaatgtcagt	2800
gttttggtgc acctttttga aatgcaaac taggaaaagg ttaaactgga	2850
ttttttttta aaaaaaaaaa aaaaaaaaaa aaa	2883

&lt;210&gt; SEQ ID NO 94

&lt;211&gt; LENGTH: 847

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

-continued

&lt;400&gt; SEQUENCE: 94

Met	Tyr	Asn	Ser	Val	Lys	Gly	Ser	Cys	Ser	Glu	Pro	Val	Ser	Phe
1				5					10					15
Thr	Thr	His	Ser	Cys	Ala	Pro	Glu	Cys	Pro	Phe	Pro	Pro	Lys	Leu
				20					25					30
Ala	His	Arg	Ser	Lys	Ser	Ser	Leu	Thr	Leu	Gln	Trp	Lys	Ala	Pro
				35					40					45
Ile	Asp	Asn	Gly	Ser	Lys	Ile	Thr	Asn	Tyr	Leu	Leu	Glu	Trp	Asp
				50					55					60
Glu	Gly	Lys	Arg	Asn	Ser	Gly	Phe	Arg	Gln	Cys	Phe	Phe	Gly	Ser
				65					70					75
Gln	Lys	His	Cys	Lys	Leu	Thr	Lys	Leu	Cys	Pro	Ala	Met	Gly	Tyr
				80					85					90
Thr	Phe	Arg	Leu	Ala	Ala	Arg	Asn	Asp	Ile	Gly	Thr	Ser	Gly	Tyr
				95					100					105
Ser	Gln	Glu	Val	Val	Cys	Tyr	Thr	Leu	Gly	Asn	Ile	Pro	Gln	Met
				110					115					120
Pro	Ser	Ala	Leu	Arg	Leu	Val	Arg	Ala	Gly	Ile	Thr	Trp	Val	Thr
				125					130					135
Leu	Gln	Trp	Ser	Lys	Pro	Glu	Gly	Cys	Ser	Pro	Glu	Glu	Val	Ile
				140					145					150
Thr	Tyr	Thr	Leu	Glu	Ile	Gln	Glu	Asp	Glu	Asn	Asp	Asn	Leu	Phe
				155					160					165
His	Pro	Lys	Tyr	Thr	Gly	Glu	Asp	Leu	Thr	Cys	Thr	Val	Lys	Asn
				170					175					180
Leu	Lys	Arg	Ser	Thr	Gln	Tyr	Lys	Phe	Arg	Leu	Thr	Ala	Ser	Asn
				185					190					195
Thr	Glu	Gly	Lys	Ser	Cys	Pro	Ser	Glu	Val	Leu	Val	Cys	Thr	Thr
				200					205					210
Ser	Pro	Asp	Arg	Pro	Gly	Pro	Pro	Thr	Arg	Pro	Leu	Val	Lys	Gly
				215					220					225
Pro	Val	Thr	Ser	His	Gly	Phe	Ser	Val	Lys	Trp	Asp	Pro	Pro	Lys
				230					235					240
Asp	Asn	Gly	Gly	Ser	Glu	Ile	Leu	Lys	Tyr	Leu	Leu	Glu	Ile	Thr
				245					250					255
Asp	Gly	Asn	Ser	Glu	Ala	Asn	Gln	Trp	Glu	Val	Ala	Tyr	Ser	Gly
				260					265					270
Ser	Ala	Thr	Glu	Tyr	Thr	Phe	Thr	His	Leu	Lys	Pro	Gly	Thr	Leu
				275					280					285
Tyr	Lys	Leu	Arg	Ala	Cys	Cys	Ile	Ser	Thr	Gly	Gly	His	Ser	Gln
				290					295					300
Cys	Ser	Glu	Ser	Leu	Pro	Val	Arg	Thr	Leu	Ser	Ile	Ala	Pro	Gly
				305					310					315
Gln	Cys	Arg	Pro	Pro	Arg	Val	Leu	Gly	Arg	Pro	Lys	His	Lys	Glu
				320					325					330
Val	His	Leu	Glu	Trp	Asp	Val	Pro	Ala	Ser	Glu	Ser	Gly	Cys	Glu
				335					340					345
Val	Ser	Glu	Tyr	Ser	Val	Glu	Met	Thr	Glu	Pro	Glu	Asp	Val	Ala
				350					355					360
Ser	Glu	Val	Tyr	His	Gly	Pro	Glu	Leu	Glu	Cys	Thr	Val	Gly	Asn

-continued

365										370					375				
Leu	Leu	Pro	Gly	Thr	Val	Tyr	Arg	Phe	Arg	Val	Arg	Ala	Leu	Asn					
				380					385					390					
Asp	Gly	Gly	Tyr	Gly	Pro	Tyr	Ser	Asp	Val	Ser	Glu	Ile	Thr	Thr					
				395					400					405					
Ala	Ala	Gly	Pro	Pro	Gly	Gln	Cys	Lys	Ala	Pro	Cys	Ile	Ser	Cys					
				410					415					420					
Thr	Pro	Asp	Gly	Cys	Val	Leu	Val	Gly	Trp	Glu	Ser	Pro	Asp	Ser					
				425					430					435					
Ser	Gly	Ala	Asp	Ile	Ser	Glu	Tyr	Arg	Leu	Glu	Trp	Gly	Glu	Asp					
				440					445					450					
Glu	Glu	Ser	Leu	Glu	Leu	Ile	Tyr	His	Gly	Thr	Asp	Thr	Arg	Phe					
				455					460					465					
Glu	Ile	Arg	Asp	Leu	Leu	Pro	Ala	Ala	Gln	Tyr	Cys	Cys	Arg	Leu					
				470					475					480					
Gln	Ala	Phe	Asn	Gln	Ala	Gly	Ala	Gly	Pro	Tyr	Ser	Glu	Leu	Val					
				485					490					495					
Leu	Cys	Gln	Thr	Pro	Ala	Ser	Ala	Pro	Asp	Pro	Val	Ser	Thr	Leu					
				500					505					510					
Cys	Val	Leu	Glu	Glu	Glu	Pro	Leu	Asp	Ala	Tyr	Pro	Asp	Ser	Pro					
				515					520					525					
Ser	Ala	Cys	Leu	Val	Leu	Asn	Trp	Glu	Glu	Pro	Cys	Asn	Asn	Gly					
				530					535					540					
Ser	Glu	Ile	Leu	Ala	Tyr	Thr	Ile	Asp	Leu	Gly	Asp	Thr	Ser	Ile					
				545					550					555					
Thr	Val	Gly	Asn	Thr	Thr	Met	His	Val	Met	Lys	Asp	Leu	Leu	Pro					
				560					565					570					
Glu	Thr	Thr	Tyr	Arg	Ile	Arg	Ile	Gln	Ala	Ile	Asn	Glu	Ile	Gly					
				575					580					585					
Ala	Gly	Pro	Phe	Ser	Gln	Phe	Ile	Lys	Ala	Lys	Thr	Arg	Pro	Leu					
				590					595					600					
Pro	Pro	Leu	Pro	Pro	Arg	Leu	Glu	Cys	Ala	Ala	Ala	Gly	Pro	Gln					
				605					610					615					
Ser	Leu	Lys	Leu	Lys	Trp	Gly	Asp	Ser	Asn	Ser	Lys	Thr	His	Ala					
				620					625					630					
Ala	Glu	Asp	Ile	Val	Tyr	Thr	Leu	Gln	Leu	Glu	Asp	Arg	Asn	Lys					
				635					640					645					
Arg	Phe	Ile	Ser	Ile	Tyr	Arg	Gly	Pro	Ser	His	Thr	Tyr	Lys	Val					
				650					655					660					
Gln	Arg	Leu	Thr	Glu	Phe	Thr	Cys	Tyr	Ser	Phe	Arg	Ile	Gln	Ala					
				665					670					675					
Ala	Ser	Glu	Ala	Gly	Glu	Gly	Pro	Phe	Ser	Glu	Thr	Tyr	Thr	Phe					
				680					685					690					
Ser	Thr	Thr	Lys	Ser	Val	Pro	Pro	Thr	Ile	Lys	Ala	Pro	Arg	Val					
				695					700					705					
Thr	Gln	Leu	Glu	Val	Asn	Ser	Cys	Glu	Ile	Leu	Trp	Glu	Thr	Val					
				710					715					720					
Pro	Ser	Met	Lys	Gly	Asp	Pro	Val	Asn	Tyr	Ile	Leu	Gln	Val	Leu					
				725					730					735					
Val	Gly	Arg	Glu	Ser	Glu	Tyr	Lys	Gln	Val	Tyr	Lys	Gly	Glu	Glu					
				740					745					750					

-continued

Ala Thr Phe Gln Ile Ser Gly Leu Gln Thr Asn Thr Asp Tyr Arg  
 755 760 765  
 Phe Arg Val Cys Ala Cys Arg Arg Cys Leu Asp Thr Ser Gln Glu  
 770 775 780  
 Leu Ser Gly Ala Phe Ser Pro Ser Ala Ala Phe Val Leu Gln Arg  
 785 790 795  
 Ser Glu Val Met Leu Thr Gly Asp Met Gly Ser Leu Asp Asp Pro  
 800 805 810  
 Lys Met Lys Ser Met Met Pro Thr Asp Glu Gln Phe Ala Ala Ile  
 815 820 825  
 Ile Val Leu Gly Phe Ala Thr Leu Ser Ile Leu Phe Ala Phe Ile  
 830 835 840  
 Leu Gln Tyr Phe Leu Met Lys  
 845

<210> SEQ ID NO 95  
 <211> LENGTH: 4725  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 95

caattcggcc tcgctccttg tgattcgcgt aaaccttccg tcctcagctg 50  
 agaacgctcc accacctccc cggatcgcgc atctcttggc tgccctccca 100  
 ctgttctcga tgttatttta ctcccgtat cccctactcg ttcttcacaa 150  
 ttctgtaggt gagtggttcc agctggtgcc tggcctgtgt ctcttggatg 200  
 ccctgtggct tcagtccgct tcctgttgcc caccacctcg tccctgggcc 250  
 gcctgatacc ccagcccaac agctaagggt tggatggaca gtagggggct 300  
 ggcttctctc actggtcagg ggtcttctcc cctgtctgcc tcccgagct 350  
 aggactgcag aggggcctat catggtgctt gcaggcccc tggctgtctc 400  
 gctgttgctg cccagcctca cactgctggt gtcccacctc tccagctccc 450  
 aggatgtctc cagtgcagcc agcagtgcgc agcagctgtg cgcccttagc 500  
 aagcacccca ccgtggcctt tgaagacctg cagccgtggg tctctaactt 550  
 cacctaccct ggagcccggg atttctccca gctggctttg gaccctccg 600  
 ggaaccagct catcgtggga gccaggaact acctcttcag actcagcctt 650  
 gccaatgtct ctcttcttca ggccacagag tgggcctcca gtgaggacac 700  
 gcgccgctcc tgccaaagca aaggaagac tgaggaggag tgcagaact 750  
 acgtgcgagt cctgatcgtc gccggccgga agtggttcat gttggaacc 800  
 aatgcctttt ccccatgtg caccagcaga caggtgggga acctcagccg 850  
 gactattgag aagatcaatg gtgtggcccg ctgcccctat gaccacgcc 900  
 acaactccac agctgtcatc tcctcccagg gggagcteta tgcagccacg 950  
 gtcatcgact tctcaggtcg ggacctgcc atctaccgca gcctgggcag 1000  
 tgggccacog cttcgcactg cccaatataa ctccaagtgg cttaatgagc 1050  
 caaacttcgt ggcagcctat gatattgggc tgtttgcata cttcttctg 1100  
 cgggagaacg cagtggagca cgactgtgga cgcacctgt actctcgcgt 1150

-continued

---

ggccccgctg tgcaagaatg acgtgggggg cggattcctg ctggaggaca	1200
catggaccac attcatgaag gcccgctca actgctccc cccggcgag	1250
gtccccctct actataacga gctgcagagt gccttccact tgccggagca	1300
ggacctcatc tatggagttt tcacaaccaa cgtaaacagc atcggcgctt	1350
ctgctgtctg cgccttcaac ctcaagtcta tctcccaggc tttcaatggc	1400
ccatttgcct accaggagaa ccccagggtc gctgggtcc ccatagccaa	1450
ccccatcccc aatttccagt gtggaccctc gctgagacc ggtcccaacg	1500
agaacctgac ggagcgagc ctgcaggagc cgcagcgctt ctctctgatg	1550
agcgaggcgc tgcagccggt gacacccgag cctgtgtca cccaggacag	1600
cgtgctctc tcacacctc tgggtggacct ggtgcaggct aaagacagc	1650
tctaccatgt actctacatt ggcaccgagt cgggcacat cctgaaggcg	1700
ctgtccacgg cgagccgag cctccacggc tgctacctg aggagctgca	1750
cgtgctgccc cccgggccc gcgagcccct ggcagcctg cgcctctgc	1800
acagcgccc cgcgctctc gtggggctga gagacggcgt cctgcgggtc	1850
ccactggaga ggtgcccgc ctaccgagc cagggggcat gcctgggggc	1900
ccgggaccgc tactgtggct gggacgggaa gcagcaactg tgcagcacac	1950
tcgaggacag ctccaacatg agcctctgga cccagaacat caccgcctgt	2000
cctgtgcgga atgtgacagc ggatgggggc ttcggccat ggtcaccatg	2050
gcaaccatgt gacacttg atggggacaa ctacggctct tgcctgtgtc	2100
gagctcgtc ctgtgattcc cctcgacccc gctgtgggg ccttgactgc	2150
ctggggccag ccatccacat cgcacaactg tccaggaatg gggcgtggac	2200
cccgtggtoa tcgtgggccc tgtgcagcac gtcctgtggc atcggcttc	2250
aggctcccca gcgaagtgc agcaaccctg ctcccccca cgggggccc	2300
atcttcgtgg gcaagagccc ggaggaacgg ttctgtaatg agaacacgcc	2350
ttgcccgggt cccatctctt gggcttctg gggctcctg agcaagtga	2400
gcagcaactg tggagggggc atgcagctgc ggcgtcggc ctgcgagAAC	2450
ggcaactcct gcctgggctg cggcaggtc aagacgtga accccgagg	2500
ctgccccgaa gtgcggcga acaccccctg gacccctg ctgcccgtga	2550
acgtgacgca gggcggggca cggcaggagc agcggttccg cttoacctgc	2600
cgcgcgccc ttgcagacc gcacggcctg cagttcgga ggagaaggac	2650
cgagacgag acctgtccc cggacggctc cggctcctg gacaccgag	2700
ccctggtgga ggtcctctg cgcagcggga gcacctccc gcacacggtg	2750
agcgggggct gggcccctg gggcccgtg togtcctgct cccgggactg	2800
cgagctgggc ttccgctcc gcaagagaac gtgactaac cggagcccc	2850
gcaacgggg cctgcctgc gtggcgatg ctgcccagta ccaggactgc	2900
aacccccagg cttgccagt tcgggggtg tggctcctg gaacctcatg	2950
gtctccatgc tcagcttct gtggggggg tcaactataa cgcaccgctt	3000
cctgcaccag ccccgcccc tcccagggtg aggacatctg tctcgggctg	3050



-continued

---

cacacggagg aggcactatg tgccacacag gcctgccag gctggtcgcc	3100
ctggtctgag tggagtaagt gcaactgacga cggagcccag agccgaagcc	3150
ggcactgtga ggagctcctc ccagggtcca gcgcctgtgc tggaaacagc	3200
agccagagcc gccctgccc ctacagcgag attcccgtca tcctgccagc	3250
ctccagcatg gaggaggcca ccgactgtgc aggtaaaaga aaccggacct	3300
acctcatgct gcggtcctcc cagccctcca gcaccccact ccaaagtctg	3350
gactctttcc acatcctgct ccagacagcc aagctttggt ggggtcccca	3400
ctgctttgag atggggtcaa tctcatccac ttggtggcca cgggcatctc	3450
ctgcttctg ggctctgggc tcctgacct agcagtgtac ctgtcttgcc	3500
agcactgcca gcgtcagtcc caggagtcca cactgggtcca tcctgccacc	3550
cccaaccatt tgcactacaa gggcggaggc acccgaaga atgaaaagta	3600
cacaccatg gaattcaaga ccctgaacaa gaataacttg atccctgatg	3650
acagagccaa cttctaccca ttgcagcaga ccaatgtgta cacgactact	3700
tactaccxaa gcccctgaa caaacacagc ttccggcccg aggcctcacc	3750
tggacaacgg tgcttcccca acagctgata ccgccgtcct ggggacttgg	3800
gcttcttgcc tcataaggc acagagcaga tggagatggg acagtggagc	3850
cagtttggtt ttctccctct gcaactggcc aagaactgc tgcttgcct	3900
gtggggggtc ccatccggct tcagagagct ctggctggca ttgacctgg	3950
gggaaagggc tggtttcagg ctgacatatg gccgcaggtc cagttcagcc	4000
caggctcttc atggttatct tccaaccac tgtaacgctg aactatgct	4050
gccatgcctg ggctgtggac ctactgggca tttgaggaat tggagaatgg	4100
agatggcaag agggcaggct ttaagtctg ggtggagac aacttcctgt	4150
ggccccaca agctgagtct ggcttctcc agctggcccc aaaaaaggcc	4200
tttgctacat cctgattatc tctgaaagta atcaatcaag tggctccagt	4250
agctctggat tttctgccag ggctgggcca ttgtgggtct gcccagtat	4300
gacatgggac caaggccagc gcaggttatc cacctctgcc tggaaagtcta	4350
tactctaccc agggcatccc tctggtcaga ggcagtgagt actgggaact	4400
ggaggctgac ctgtgcttag aagtccctta atctgggctg gtacaggcct	4450
cagccttgcc ctcaatgcac gaaaggtggc ccaggagaga ggatcaatgc	4500
cataggaggc agaagtctgg cctctgtgcc tctatggaga ctatcttcca	4550
gttgctgctc aacagagttg ttggctgaga cctgcttggg agtctctgct	4600
ggcccttcat ctgttcagga acacacacac acacacactc acacacgcac	4650
acacaatcac aatttgctac agcaacaaaa aagacattgg gctgtggcat	4700
tattaattaa agatgatatc cagtc	4725

&lt;210&gt; SEQ ID NO 96

&lt;211&gt; LENGTH: 1092

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 96

-continued

Met	Pro	Cys	Gly	Phe	Ser	Pro	Ser	Pro	Val	Ala	His	His	Leu	Val
1				5					10					15
Pro	Gly	Pro	Pro	Asp	Thr	Pro	Ala	Gln	Gln	Leu	Arg	Cys	Gly	Trp
				20					25					30
Thr	Val	Gly	Gly	Trp	Leu	Leu	Ser	Leu	Val	Arg	Gly	Leu	Leu	Pro
				35					40					45
Cys	Leu	Pro	Pro	Gly	Ala	Arg	Thr	Ala	Glu	Gly	Pro	Ile	Met	Val
				50					55					60
Leu	Ala	Gly	Pro	Leu	Ala	Val	Ser	Leu	Leu	Leu	Pro	Ser	Leu	Thr
				65					70					75
Leu	Leu	Val	Ser	His	Leu	Ser	Ser	Ser	Gln	Asp	Val	Ser	Ser	Glu
				80					85					90
Pro	Ser	Ser	Glu	Gln	Gln	Leu	Cys	Ala	Leu	Ser	Lys	His	Pro	Thr
				95					100					105
Val	Ala	Phe	Glu	Asp	Leu	Gln	Pro	Trp	Val	Ser	Asn	Phe	Thr	Tyr
				110					115					120
Pro	Gly	Ala	Arg	Asp	Phe	Ser	Gln	Leu	Ala	Leu	Asp	Pro	Ser	Gly
				125					130					135
Asn	Gln	Leu	Ile	Val	Gly	Ala	Arg	Asn	Tyr	Leu	Phe	Arg	Leu	Ser
				140					145					150
Leu	Ala	Asn	Val	Ser	Leu	Leu	Gln	Ala	Thr	Glu	Trp	Ala	Ser	Ser
				155					160					165
Glu	Asp	Thr	Arg	Arg	Ser	Cys	Gln	Ser	Lys	Gly	Lys	Thr	Glu	Glu
				170					175					180
Glu	Cys	Gln	Asn	Tyr	Val	Arg	Val	Leu	Ile	Val	Ala	Gly	Arg	Lys
				185					190					195
Val	Phe	Met	Cys	Gly	Thr	Asn	Ala	Phe	Ser	Pro	Met	Cys	Thr	Ser
				200					205					210
Arg	Gln	Val	Gly	Asn	Leu	Ser	Arg	Thr	Ile	Glu	Lys	Ile	Asn	Gly
				215					220					225
Val	Ala	Arg	Cys	Pro	Tyr	Asp	Pro	Arg	His	Asn	Ser	Thr	Ala	Val
				230					235					240
Ile	Ser	Ser	Gln	Gly	Glu	Leu	Tyr	Ala	Ala	Thr	Val	Ile	Asp	Phe
				245					250					255
Ser	Gly	Arg	Asp	Pro	Ala	Ile	Tyr	Arg	Ser	Leu	Gly	Ser	Gly	Pro
				260					265					270
Pro	Leu	Arg	Thr	Ala	Gln	Tyr	Asn	Ser	Lys	Trp	Leu	Asn	Glu	Pro
				275					280					285
Asn	Phe	Val	Ala	Ala	Tyr	Asp	Ile	Gly	Leu	Phe	Ala	Tyr	Phe	Phe
				290					295					300
Leu	Arg	Glu	Asn	Ala	Val	Glu	His	Asp	Cys	Gly	Arg	Thr	Val	Tyr
				305					310					315
Ser	Arg	Val	Ala	Arg	Val	Cys	Lys	Asn	Asp	Val	Gly	Gly	Arg	Phe
				320					325					330
Leu	Leu	Glu	Asp	Thr	Trp	Thr	Thr	Phe	Met	Lys	Ala	Arg	Leu	Asn
				335					340					345
Cys	Ser	Arg	Pro	Gly	Glu	Val	Pro	Phe	Tyr	Tyr	Asn	Glu	Leu	Gln
				350					355					360
Ser	Ala	Phe	His	Leu	Pro	Glu	Gln	Asp	Leu	Ile	Tyr	Gly	Val	Phe
				365					370					375
Thr	Thr	Asn	Val	Asn	Ser	Ile	Ala	Ala	Ser	Ala	Val	Cys	Ala	Phe

-continued

380					385					390				
Asn	Leu	Ser	Ala	Ile	Ser	Gln	Ala	Phe	Asn	Gly	Pro	Phe	Arg	Tyr
				395					400					405
Gln	Glu	Asn	Pro	Arg	Ala	Ala	Trp	Leu	Pro	Ile	Ala	Asn	Pro	Ile
				410					415					420
Pro	Asn	Phe	Gln	Cys	Gly	Thr	Leu	Pro	Glu	Thr	Gly	Pro	Asn	Glu
				425					430					435
Asn	Leu	Thr	Glu	Arg	Ser	Leu	Gln	Asp	Ala	Gln	Arg	Leu	Phe	Leu
				440					445					450
Met	Ser	Glu	Ala	Val	Gln	Pro	Val	Thr	Pro	Glu	Pro	Cys	Val	Thr
				455					460					465
Gln	Asp	Ser	Val	Arg	Phe	Ser	His	Leu	Val	Val	Asp	Leu	Val	Gln
				470					475					480
Ala	Lys	Asp	Thr	Leu	Tyr	His	Val	Leu	Tyr	Ile	Gly	Thr	Glu	Ser
				485					490					495
Gly	Thr	Ile	Leu	Lys	Ala	Leu	Ser	Thr	Ala	Ser	Arg	Ser	Leu	His
				500					505					510
Gly	Cys	Tyr	Leu	Glu	Glu	Leu	His	Val	Leu	Pro	Pro	Gly	Arg	Arg
				515					520					525
Glu	Pro	Leu	Arg	Ser	Leu	Arg	Ile	Leu	His	Ser	Ala	Arg	Ala	Leu
				530					535					540
Phe	Val	Gly	Leu	Arg	Asp	Gly	Val	Leu	Arg	Val	Pro	Leu	Glu	Arg
				545					550					555
Cys	Ala	Ala	Tyr	Arg	Ser	Gln	Gly	Ala	Cys	Leu	Gly	Ala	Arg	Asp
				560					565					570
Pro	Tyr	Cys	Gly	Trp	Asp	Gly	Lys	Gln	Gln	Arg	Cys	Ser	Thr	Leu
				575					580					585
Glu	Asp	Ser	Ser	Asn	Met	Ser	Leu	Trp	Thr	Gln	Asn	Ile	Thr	Ala
				590					595					600
Cys	Pro	Val	Arg	Asn	Val	Thr	Arg	Asp	Gly	Gly	Phe	Gly	Pro	Trp
				605					610					615
Ser	Pro	Trp	Gln	Pro	Cys	Glu	His	Leu	Asp	Gly	Asp	Asn	Ser	Gly
				620					625					630
Ser	Cys	Leu	Cys	Arg	Ala	Arg	Ser	Cys	Asp	Ser	Pro	Arg	Pro	Arg
				635					640					645
Cys	Gly	Gly	Leu	Asp	Cys	Leu	Gly	Pro	Ala	Ile	His	Ile	Ala	Asn
				650					655					660
Cys	Ser	Arg	Asn	Gly	Ala	Trp	Thr	Pro	Trp	Ser	Ser	Trp	Ala	Leu
				665					670					675
Cys	Ser	Thr	Ser	Cys	Gly	Ile	Gly	Phe	Gln	Val	Arg	Gln	Arg	Ser
				680					685					690
Cys	Ser	Asn	Pro	Ala	Pro	Arg	His	Gly	Gly	Arg	Ile	Phe	Val	Gly
				695					700					705
Lys	Ser	Arg	Glu	Glu	Arg	Phe	Cys	Asn	Glu	Asn	Thr	Pro	Cys	Pro
				710					715					720
Val	Pro	Ile	Phe	Trp	Ala	Ser	Trp	Gly	Ser	Trp	Ser	Lys	Cys	Ser
				725					730					735
Ser	Asn	Cys	Gly	Gly	Gly	Met	Gln	Ser	Arg	Arg	Arg	Ala	Cys	Glu
				740					745					750
Asn	Gly	Asn	Ser	Cys	Leu	Gly	Cys	Gly	Glu	Phe	Lys	Thr	Cys	Asn
				755					760					765

-continued

---

Pro Glu Gly Cys Pro Glu Val Arg Arg Asn Thr Pro Trp Thr Pro  
 770 775 780

Trp Leu Pro Val Asn Val Thr Gln Gly Gly Ala Arg Gln Glu Gln  
 785 790 795

Arg Phe Arg Phe Thr Cys Arg Ala Pro Leu Ala Asp Pro His Gly  
 800 805 810

Leu Gln Phe Gly Arg Arg Arg Thr Glu Thr Arg Thr Cys Pro Ala  
 815 820 825

Asp Gly Ser Gly Ser Cys Asp Thr Asp Ala Leu Val Glu Val Leu  
 830 835 840

Leu Arg Ser Gly Ser Thr Ser Pro His Thr Val Ser Gly Gly Trp  
 845 850 855

Ala Ala Trp Gly Pro Trp Ser Ser Cys Ser Arg Asp Cys Glu Leu  
 860 865 870

Gly Phe Arg Val Arg Lys Arg Thr Cys Thr Asn Pro Glu Pro Arg  
 875 880 885

Asn Gly Gly Leu Pro Cys Val Gly Asp Ala Ala Glu Tyr Gln Asp  
 890 895 900

Cys Asn Pro Gln Ala Cys Pro Val Arg Gly Ala Trp Ser Cys Trp  
 905 910 915

Thr Ser Trp Ser Pro Cys Ser Ala Ser Cys Gly Gly Gly His Tyr  
 920 925 930

Gln Arg Thr Arg Ser Cys Thr Ser Pro Ala Pro Ser Pro Gly Glu  
 935 940 945

Asp Ile Cys Leu Gly Leu His Thr Glu Glu Ala Leu Cys Ala Thr  
 950 955 960

Gln Ala Cys Pro Gly Trp Ser Pro Trp Ser Glu Trp Ser Lys Cys  
 965 970 975

Thr Asp Asp Gly Ala Gln Ser Arg Ser Arg His Cys Glu Glu Leu  
 980 985 990

Leu Pro Gly Ser Ser Ala Cys Ala Gly Asn Ser Ser Gln Ser Arg  
 995 1000 1005

Pro Cys Pro Tyr Ser Glu Ile Pro Val Ile Leu Pro Ala Ser Ser  
 1010 1015 1020

Met Glu Glu Ala Thr Asp Cys Ala Gly Lys Arg Asn Arg Thr Tyr  
 1025 1030 1035

Leu Met Leu Arg Ser Ser Gln Pro Ser Ser Thr Pro Leu Gln Ser  
 1040 1045 1050

Leu Asp Ser Phe His Ile Leu Leu Gln Thr Ala Lys Leu Cys Trp  
 1055 1060 1065

Gly Pro His Cys Phe Glu Met Gly Ser Ile Ser Ser Thr Trp Trp  
 1070 1075 1080

Pro Arg Ala Ser Pro Ala Ser Trp Ala Leu Gly Ser  
 1085 1090

<210> SEQ ID NO 97  
 <211> LENGTH: 3391  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 97

caagccctcc cagcatcccc tctcctgtgt tctcctccag ttctctactc

-continued

---

agagttgact gaccagagat ttatcagctt ggagggctgg aggtgtggat	100
ccatggggta gcctcaacgc atctgcccct ccaccccagc cagctcatgg	150
gccacgtggc ctggcccagc ctcagcaccc agggccagtg aacagagccc	200
tggctggagt ccaaacatgt ggggcctggt gaggtcctg ctggcctggc	250
tgggtggctg gggctgcatg gggcgtctgg cagccccagc ccgggcctgg	300
gcaggtccc gggaacacc agggcctgct ctgctgcgga ctcgaaggag	350
ctgggtctg aaccagttct ttgtcattga ggaatatgct ggtccagagc	400
ctgttctcat tggcaagctg cactcggatg ttgaccgggg agagggccgc	450
accaagtacc tgttgaccgg ggagggggca ggcaccgtat ttgtgattga	500
tgagccaca ggcaatattc atgttaccaa gagccttgac cgggaggaaa	550
aggcgcaata tgtgctactg gcccaagccg tggaccgagc ctccaaccgg	600
cccctggagc ccccatcaga gttcatcato aaagtgaag acatcaacga	650
caatccaccc atttttcccc ttgggcccta ccatgccacc gtgcccgaga	700
tgtccaatg cgggacatca gtgatccagg tgactgctca cgatgctgat	750
gaccccgct atgggaacag tgccaagctg gtgtacactg ttctggatgg	800
actgcctttc ttctctgtgg acccccagac tggagtggty cgtacagcca	850
tcccacaat ggaccgggag acacaggagg agttcttggg ggtgatccag	900
gccaaagaca tgggcggcca catggggggg ctgtcaggca gcactacggt	950
gactgtcaog ctcagcgatg tcaacgacaa ccccccaag ttcccacaga	1000
gcctatacca gttctccgtg gtggagacag ctggacctgg cacactggtg	1050
ggccggctcc gggcccagga ccagacctg ggggacaacg ccctgatggc	1100
atacagcadc ctggatgggg aggggtctga ggccttcagc atcagcacag	1150
acttgacagg tcgagacggg ctctcactg tccgcaagcc cctagacttt	1200
gagagccagc gctcctactc ctccgtgtc gaggccacca acacgctcat	1250
tgacccagcc tatctgcggc gagggccctt caaggatgtg gcctctgtgc	1300
gtgtggcagt gcaagatgcc ccagagccac ctgccttcac ccaggctgcc	1350
taccacctga cagtgcctga gaacaaggcc ccggggaccc tggtaggcca	1400
gatctccog gctgacctgg actcccctgc cagcccaatc agatactcca	1450
tctccccca ctcagatccg gagcgttgct tctctatcca gcccgaggaa	1500
ggcaccatcc atacagcagc acccctggat cgcgaggctc gcgcctggca	1550
caacctcact gtgctggcta cagagctcga cagttctgca caggcctcgc	1600
gcgtgcaagt ggccatccag accctggatg agaatgacaa tgctccccag	1650
ctggctgagc cctacgatac ttttgtgtgt gactctgag ctctggcca	1700
gctgattcag gtcacccgg ccctggacag agatgaagtt ggcaacagta	1750
gccatgtctc ctttcaaggt cctctgggccc ctgatgcaa ctttactgtc	1800
caggacaacc gagatggctc cgcagcctg ctgctgcct cccgccctgc	1850
tccacccgc catgcccct acttggttcc catagaactg tgggactggg	1900
ggcagccggc gctgagcagc actgccacag tgactgttag tgtgtgccgc	1950

-continued

---

```

tgccagcctg acggctctgt ggcacccctgc tggcctgagg ctacacctc 2000
agctgctggg ctacagcaccg gcgccctgct tgccatcatc acctgtgtgg 2050
gtgccctgct tgccctgggt gtgctcttcg tggccctgcg gcggcagaag 2100
caagaagcac tgatggtact ggaggaggag gacgtccgag agaacatcat 2150
cacctacgac gacgagggcg gcggcgagga ggacaccgag gccttcgaca 2200
tcacggcctt gcagaaccg gacggggcg ccccccgcc gcccgccct 2250
cccgcgcgcc gagacgtgtt gccccggcc cgggtgtcgc gccagcccag 2300
accccccgcc ccgcccagc tggcgagct cctggcgctg cggtcccg 2350
aggcggacga ggacccccgc gtacccccgt acgactcggg gcaggtgtac 2400
ggctacgagg gcccgggctc ctcttgccgc tccctcagct cctgggctc 2450
cggcagcgaa gccggcgcg cccccggcc cggcgagccg ctggacgact 2500
gggtcccgct ctccgcacc ctggccgagc tgtatggggc caaggagccc 2550
ccggccccct gagcgcccgg gctggcccgg cccaccgcyg ggggggggca 2600
gcgggcacag gccctctgag tgagccccc ggggtccagg cgggcggcag 2650
cagcccaggg gcccaggcc tcctccctgt ccttgtgtcc ctcttgctt 2700
ccccggggca ccctcgctct cacctccctc ctctgagtc ggtgtgtgtg 2750
tctctctcca ggaatctttg tctctatctg tgacacgctc ctctgtccgg 2800
gcctgggttt cctgccctgg ccctggccct gcgatctctc actgtgattc 2850
ctctccttcc tccgtggcgt tttgtctctg cagttctgaa gctcacacat 2900
agtctccctg cgtcttcctt gccatacac atgctctgtg tctgtctcct 2950
gcccacatct cccttcttc tctctgggtc cctgtgactg gctttttgtt 3000
ttttctgtt gtccatccca aaatcaagag aaactccag ccaactgctg 3050
ccacctctct gcaggggatg ttgtgcccc gacctgcctg catggttcca 3100
tccattactc atggcctcag cctcctcctg gctccactgg cctccagctg 3150
agagagggaa ccagcctgcc tcccagggca agagctccag cctcccgtgt 3200
ggccgcctcc ctggagctct gccagctgc cagcttccc tgggcatccc 3250
agccctggg attgtctgt gtgcttcctg agggagttag gaaaggaaag 3300
ggggagggcg ctggggaagg gaaaagagg aggaagggga ggggcctcca 3350
tctctaattt cataataaac aacacttta tttgtaaaa c 3391

```

&lt;210&gt; SEQ ID NO 98

&lt;211&gt; LENGTH: 781

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 98

```

Met Trp Gly Leu Val Arg Leu Leu Leu Ala Trp Leu Gly Gly Trp
 1           5           10          15

```

```

Gly Cys Met Gly Arg Leu Ala Ala Pro Ala Arg Ala Trp Ala Gly
          20          25          30

```

```

Ser Arg Glu His Pro Gly Pro Ala Leu Leu Arg Thr Arg Arg Ser
          35          40          45

```

-continued

Trp	Val	Trp	Asn	Gln	Phe	Phe	Val	Ile	Glu	Glu	Tyr	Ala	Gly	Pro	50	55	60
Glu	Pro	Val	Leu	Ile	Gly	Lys	Leu	His	Ser	Asp	Val	Asp	Arg	Gly	65	70	75
Glu	Gly	Arg	Thr	Lys	Tyr	Leu	Leu	Thr	Gly	Glu	Gly	Ala	Gly	Thr	80	85	90
Val	Phe	Val	Ile	Asp	Glu	Ala	Thr	Gly	Asn	Ile	His	Val	Thr	Lys	95	100	105
Ser	Leu	Asp	Arg	Glu	Glu	Lys	Ala	Gln	Tyr	Val	Leu	Leu	Ala	Gln	110	115	120
Ala	Val	Asp	Arg	Ala	Ser	Asn	Arg	Pro	Leu	Glu	Pro	Pro	Ser	Glu	125	130	135
Phe	Ile	Ile	Lys	Val	Gln	Asp	Ile	Asn	Asp	Asn	Pro	Pro	Ile	Phe	140	145	150
Pro	Leu	Gly	Pro	Tyr	His	Ala	Thr	Val	Pro	Glu	Met	Ser	Asn	Val	155	160	165
Gly	Thr	Ser	Val	Ile	Gln	Val	Thr	Ala	His	Asp	Ala	Asp	Asp	Pro	170	175	180
Ser	Tyr	Gly	Asn	Ser	Ala	Lys	Leu	Val	Tyr	Thr	Val	Leu	Asp	Gly	185	190	195
Leu	Pro	Phe	Phe	Ser	Val	Asp	Pro	Gln	Thr	Gly	Val	Val	Arg	Thr	200	205	210
Ala	Ile	Pro	Asn	Met	Asp	Arg	Glu	Thr	Gln	Glu	Glu	Phe	Leu	Val	215	220	225
Val	Ile	Gln	Ala	Lys	Asp	Met	Gly	Gly	His	Met	Gly	Gly	Leu	Ser	230	235	240
Gly	Ser	Thr	Thr	Val	Thr	Val	Thr	Leu	Ser	Asp	Val	Asn	Asp	Asn	245	250	255
Pro	Pro	Lys	Phe	Pro	Gln	Ser	Leu	Tyr	Gln	Phe	Ser	Val	Val	Glu	260	265	270
Thr	Ala	Gly	Pro	Gly	Thr	Leu	Val	Gly	Arg	Leu	Arg	Ala	Gln	Asp	275	280	285
Pro	Asp	Leu	Gly	Asp	Asn	Ala	Leu	Met	Ala	Tyr	Ser	Ile	Leu	Asp	290	295	300
Gly	Glu	Gly	Ser	Glu	Ala	Phe	Ser	Ile	Ser	Thr	Asp	Leu	Gln	Gly	305	310	315
Arg	Asp	Gly	Leu	Leu	Thr	Val	Arg	Lys	Pro	Leu	Asp	Phe	Glu	Ser	320	325	330
Gln	Arg	Ser	Tyr	Ser	Phe	Arg	Val	Glu	Ala	Thr	Asn	Thr	Leu	Ile	335	340	345
Asp	Pro	Ala	Tyr	Leu	Arg	Arg	Gly	Pro	Phe	Lys	Asp	Val	Ala	Ser	350	355	360
Val	Arg	Val	Ala	Val	Gln	Asp	Ala	Pro	Glu	Pro	Pro	Ala	Phe	Thr	365	370	375
Gln	Ala	Ala	Tyr	His	Leu	Thr	Val	Pro	Glu	Asn	Lys	Ala	Pro	Gly	380	385	390
Thr	Leu	Val	Gly	Gln	Ile	Ser	Ala	Ala	Asp	Leu	Asp	Ser	Pro	Ala	395	400	405
Ser	Pro	Ile	Arg	Tyr	Ser	Ile	Leu	Pro	His	Ser	Asp	Pro	Glu	Arg	410	415	420
Cys	Phe	Ser	Ile	Gln	Pro	Glu	Glu	Gly	Thr	Ile	His	Thr	Ala	Ala			

-continued

---

	425		430		435
Pro Leu Asp Arg Glu Ala Arg Ala Trp His Asn Leu Thr Val Leu	440		445		450
Ala Thr Glu Leu Asp Ser Ser Ala Gln Ala Ser Arg Val Gln Val	455		460		465
Ala Ile Gln Thr Leu Asp Glu Asn Asp Asn Ala Pro Gln Leu Ala	470		475		480
Glu Pro Tyr Asp Thr Phe Val Cys Asp Ser Ala Ala Pro Gly Gln	485		490		495
Leu Ile Gln Val Ile Arg Ala Leu Asp Arg Asp Glu Val Gly Asn	500		505		510
Ser Ser His Val Ser Phe Gln Gly Pro Leu Gly Pro Asp Ala Asn	515		520		525
Phe Thr Val Gln Asp Asn Arg Asp Gly Ser Ala Ser Leu Leu Leu	530		535		540
Pro Ser Arg Pro Ala Pro Pro Arg His Ala Pro Tyr Leu Val Pro	545		550		555
Ile Glu Leu Trp Asp Trp Gly Gln Pro Ala Leu Ser Ser Thr Ala	560		565		570
Thr Val Thr Val Ser Val Cys Arg Cys Gln Pro Asp Gly Ser Val	575		580		585
Ala Ser Cys Trp Pro Glu Ala His Leu Ser Ala Ala Gly Leu Ser	590		595		600
Thr Gly Ala Leu Leu Ala Ile Ile Thr Cys Val Gly Ala Leu Leu	605		610		615
Ala Leu Val Val Leu Phe Val Ala Leu Arg Arg Gln Lys Gln Glu	620		625		630
Ala Leu Met Val Leu Glu Glu Glu Asp Val Arg Glu Asn Ile Ile	635		640		645
Thr Tyr Asp Asp Glu Gly Gly Gly Glu Glu Asp Thr Glu Ala Phe	650		655		660
Asp Ile Thr Ala Leu Gln Asn Pro Asp Gly Ala Ala Pro Pro Ala	665		670		675
Pro Gly Pro Pro Ala Arg Arg Asp Val Leu Pro Arg Ala Arg Val	680		685		690
Ser Arg Gln Pro Arg Pro Pro Gly Pro Ala Asp Val Ala Gln Leu	695		700		705
Leu Ala Leu Arg Leu Arg Glu Ala Asp Glu Asp Pro Gly Val Pro	710		715		720
Pro Tyr Asp Ser Val Gln Val Tyr Gly Tyr Glu Gly Arg Gly Ser	725		730		735
Ser Cys Gly Ser Leu Ser Ser Leu Gly Ser Gly Ser Glu Ala Gly	740		745		750
Gly Ala Pro Gly Pro Ala Glu Pro Leu Asp Asp Trp Gly Pro Leu	755		760		765
Phe Arg Thr Leu Ala Glu Leu Tyr Gly Ala Lys Glu Pro Pro Ala	770		775		780

Pro

<210> SEQ ID NO 99  
 <211> LENGTH: 2855



---

-continued

---

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 99

gccaaactg gccaaacata tggggctgga atctcaacat cggtcactgg 50  
gacctcaata tttggagcgg gaaccccaca atttggaaca cagacccc aa 100  
tattttgagc agaacccc aa gatttgacat ctaaacctc aagcctggag 150  
ctgaactctg aattctgggc ctgggacctt gaaatctggg actggatttc 200  
cagtactgta ccctggaacc cactcttggg gacctgaacc ctgggattca 250  
ggcctcaaat tccaagatct ggactgtggg attccaaggg gcctgaacct 300  
gagtttgggc ctgaagtcct tgctgcagac ctgagtgctt aaatctgggg 350  
cttgagacct cccaatcttg actcagcacc ccaatatctg aatgcagaac 400  
cccgggatgg gatctcagac tctaaacccc accgtttggc tgcttagcat 450  
cccaagactg gacctgggag accctgacct tgaacaacct aaactggacc 500  
cgtaaaactg gaccctagag gcccaatatt taggggtctg gaaccccag 550  
tattaaggctc tggagactcc gttgccacag atttgagccg agtcaggaca 600  
cagtcctctc acagaagcct tggggacagc aaaagcatga ccagatgctc 650  
cctccagagc cctgacctct gactccccctg gagctaggac tetgctccct 700  
ggggctgctt ctagctcagc acaccctgc cgcgatggc catcctcccg 750  
ttgctcctgt gcctgctgcc gctggccctt gcctcatccc caccocagtc 800  
agccacaccc agcccatgtc cccgcccgtg ccgctgccag acacagtcgc 850  
tgcccctaag cgtgctgtgc ccaggggcag gcctcctgct cgtgccacct 900  
tcgctggacc gccgggcagc cgagctgcgg ctggcagaca acttcacgc 950  
ctccgtgctc cgcgcgac tggccaacat gacaggcctg ctgcatctga 1000  
gcctgtcgcg gaacaccatc cgcacgtgg ctgcccgcgc cttgcgcgac 1050  
ctgcccggcc tgctgcccct gcacctggat ggcaaccggc tgacctcaact 1100  
gggcgagggc cagctgcgcg gcctggctca cttgcgccac ctcatcctca 1150  
gcaacaacca gctggcagcg ctggcggccg gcgcctgga tgattgtgcc 1200  
gagacactgg aggacctcga cctctctac aacaacctcg agcagctgcc 1250  
ctgggagggc ctgggccgcc tgggcaacgt caacacgttg ggcctcgacc 1300  
acaacctgct ggcttctgtg ccggcgctt tttcccgcct gcacaagctg 1350  
gcccggctgg acatgacctc caaccgctg accacaatcc caccggacc 1400  
actcttctcc cgcctgcccc tgctcggcag gcccccgggc tcgcccgcct 1450  
ctgcctggtg gctggccttt ggccggaaac ccctgcaact caactgcgag 1500  
ctggtgtggc tgctgcccct ggcgcccggg gacgacctc aggcctgccc 1550  
gtccccacct gctctggcg gcgcctactt ctggggggtg ggcgaggagg 1600  
agtttctctc cgagccgccc gtggtgactc accgctcacc acctctggct 1650  
gtgcccgcag gtcggccggc tgcccctgctc tgcccggcag tgggggacct 1700  
agagccccgt gtgcgttggg tgtaaccca gggccggctg ctaggcaact 1750  
caagccgtgc ccgcccctc cccaatggga cgctggagct gctggtcacc 1800

-continued

---

```

gagccgggtg atggtggcat cttcacctgc attgcgcca atgcagctgg      1850
cgagcccaca gctgctgtgg agctgactgt gggccccca ccacctctc      1900
agctagccaa cagcaccagc tgtgaccccc cgcgggacgg ggatcctgat      1950
gctctcacc caccctccgc tgctctgtct tctgccaagg tggccgacac      2000
tgggccccct accgaccgtg gcgtccaggt gactgagcac ggggccacag      2050
ctgctcttgt ccagtggccg gatcagcggc ctatcccggg catccgcatg      2100
taccagatcc agtacaacag ctcggtgat gacatcctcg tctacaggat      2150
gatccccggg gagagccgct cgttctgtct gacggacctg gcgtcaggcc      2200
ggacctacga tctgtgctgt ctgcgctgtg atgaggacag cgccacgggg      2250
ctcacggcca cgcggcctgt gggctcggcc cgcttctcca ccgaacctgc      2300
gctgcggcca tgcggggcgc cgcacgctcc cttctgggc ggcacgatga      2350
tcacgcgct gggcggcgtc atcgtagcct cggtagtgg tttcatcttc      2400
gtgctgctaa tgcgctacaa ggtgcacggc ggcagcccc ccggcaaggc      2450
caagattccc gcgctgtta gcagcgtttg ctcccagacc aacggcggcc      2500
tgggcccac gccacgccc gcccccccg cccggagcc cgcggcgtc      2550
agggcccaca ccgtgtcca gctggactgc gagccctggg ggcccggcca      2600
cgaacctgtg ggacctagc caggcggccc cccctctaag ggtcctctgg      2650
ccccacggac agcaggacct ggacacctg tgggacctgg cctcaaactc      2700
accaaatcgc tcatggtttt taaaactctg atggggaggg tgcggggac      2750
accggggcaa aacaagaaag tcctattttt caaaaaaaaa aaaaaaaaaa      2800
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      2850
aaaaaa                                           2855

```

&lt;210&gt; SEQ ID NO 100

&lt;211&gt; LENGTH: 627

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 100

```

Met Ala Ile Leu Pro Leu Leu Leu Cys Leu Leu Pro Leu Ala Pro
 1           5           10          15
Ala Ser Ser Pro Pro Gln Ser Ala Thr Pro Ser Pro Cys Pro Arg
          20          25          30
Arg Cys Arg Cys Gln Thr Gln Ser Leu Pro Leu Ser Val Leu Cys
          35          40          45
Pro Gly Ala Gly Leu Leu Phe Val Pro Pro Ser Leu Asp Arg Arg
          50          55          60
Ala Ala Glu Leu Arg Leu Ala Asp Asn Phe Ile Ala Ser Val Arg
          65          70          75
Arg Arg Asp Leu Ala Asn Met Thr Gly Leu Leu His Leu Ser Leu
          80          85          90
Ser Arg Asn Thr Ile Arg His Val Ala Ala Gly Ala Phe Ala Asp
          95          100         105
Leu Arg Ala Leu Arg Ala Leu His Leu Asp Gly Asn Arg Leu Thr
          110          115          120

```

-continued

---

Ser	Leu	Gly	Glu	Gly	Gln	Leu	Arg	Gly	Leu	Val	Asn	Leu	Arg	His
				125					130					135
Leu	Ile	Leu	Ser	Asn	Asn	Gln	Leu	Ala	Ala	Leu	Ala	Ala	Gly	Ala
				140					145					150
Leu	Asp	Asp	Cys	Ala	Glu	Thr	Leu	Glu	Asp	Leu	Asp	Leu	Ser	Tyr
				155					160					165
Asn	Asn	Leu	Glu	Gln	Leu	Pro	Trp	Glu	Ala	Leu	Gly	Arg	Leu	Gly
				170					175					180
Asn	Val	Asn	Thr	Leu	Gly	Leu	Asp	His	Asn	Leu	Leu	Ala	Ser	Val
				185					190					195
Pro	Gly	Ala	Phe	Ser	Arg	Leu	His	Lys	Leu	Ala	Arg	Leu	Asp	Met
				200					205					210
Thr	Ser	Asn	Arg	Leu	Thr	Thr	Ile	Pro	Pro	Asp	Pro	Leu	Phe	Ser
				215					220					225
Arg	Leu	Pro	Leu	Leu	Ala	Arg	Pro	Arg	Gly	Ser	Pro	Ala	Ser	Ala
				230					235					240
Leu	Val	Leu	Ala	Phe	Gly	Gly	Asn	Pro	Leu	His	Cys	Asn	Cys	Glu
				245					250					255
Leu	Val	Trp	Leu	Arg	Arg	Leu	Ala	Arg	Glu	Asp	Asp	Leu	Glu	Ala
				260					265					270
Cys	Ala	Ser	Pro	Pro	Ala	Leu	Gly	Gly	Arg	Tyr	Phe	Trp	Ala	Val
				275					280					285
Gly	Glu	Glu	Glu	Phe	Val	Cys	Glu	Pro	Pro	Val	Val	Thr	His	Arg
				290					295					300
Ser	Pro	Pro	Leu	Ala	Val	Pro	Ala	Gly	Arg	Pro	Ala	Ala	Leu	Arg
				305					310					315
Cys	Arg	Ala	Val	Gly	Asp	Pro	Glu	Pro	Arg	Val	Arg	Trp	Val	Ser
				320					325					330
Pro	Gln	Gly	Arg	Leu	Leu	Gly	Asn	Ser	Ser	Arg	Ala	Arg	Ala	Phe
				335					340					345
Pro	Asn	Gly	Thr	Leu	Glu	Leu	Leu	Val	Thr	Glu	Pro	Gly	Asp	Gly
				350					355					360
Gly	Ile	Phe	Thr	Cys	Ile	Ala	Ala	Asn	Ala	Ala	Gly	Glu	Ala	Thr
				365					370					375
Ala	Ala	Val	Glu	Leu	Thr	Val	Gly	Pro	Pro	Pro	Pro	Pro	Gln	Leu
				380					385					390
Ala	Asn	Ser	Thr	Ser	Cys	Asp	Pro	Pro	Arg	Asp	Gly	Asp	Pro	Asp
				395					400					405
Ala	Leu	Thr	Pro	Pro	Ser	Ala	Ala	Ser	Ala	Ser	Ala	Lys	Val	Ala
				410					415					420
Asp	Thr	Gly	Pro	Pro	Thr	Asp	Arg	Gly	Val	Gln	Val	Thr	Glu	His
				425					430					435
Gly	Ala	Thr	Ala	Ala	Leu	Val	Gln	Trp	Pro	Asp	Gln	Arg	Pro	Ile
				440					445					450
Pro	Gly	Ile	Arg	Met	Tyr	Gln	Ile	Gln	Tyr	Asn	Ser	Ser	Ala	Asp
				455					460					465
Asp	Ile	Leu	Val	Tyr	Arg	Met	Ile	Pro	Ala	Glu	Ser	Arg	Ser	Phe
				470					475					480
Leu	Leu	Thr	Asp	Leu	Ala	Ser	Gly	Arg	Thr	Tyr	Asp	Leu	Cys	Val
				485					490					495

-continued

---

Leu Ala Val Tyr Glu Asp Ser Ala Thr Gly Leu Thr Ala Thr Arg  
 500 505 510

Pro Val Gly Cys Ala Arg Phe Ser Thr Glu Pro Ala Leu Arg Pro  
 515 520 525

Cys Gly Ala Pro His Ala Pro Phe Leu Gly Gly Thr Met Ile Ile  
 530 535 540

Ala Leu Gly Gly Val Ile Val Ala Ser Val Leu Val Phe Ile Phe  
 545 550 555

Val Leu Leu Met Arg Tyr Lys Val His Gly Gly Gln Pro Pro Gly  
 560 565 570

Lys Ala Lys Ile Pro Ala Pro Val Ser Ser Val Cys Ser Gln Thr  
 575 580 585

Asn Gly Ala Leu Gly Pro Thr Pro Thr Pro Ala Pro Pro Ala Pro  
 590 595 600

Glu Pro Ala Ala Leu Arg Ala His Thr Val Val Gln Leu Asp Cys  
 605 610 615

Glu Pro Trp Gly Pro Gly His Glu Pro Val Gly Pro  
 620 625

<210> SEQ ID NO 101  
 <211> LENGTH: 1111  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien  
 <400> SEQUENCE: 101

```

cgactccata accgtggcct tggcccccagt ccccttgact tccggacttc           50
agaccagata ctgcccataat ccccttatga agtcttgcc aggcaacccc           100
taggggtgtc gttttctaaa gattaagag gcggtgctaa gctgcagacg           150
gacttgcgac tcagccactg gtgtaagtca ggcgggaggt ggcgccaat           200
aagctcaaga gaggaggcgg gttctggaaa aaggccaata gcctgtgaag           250
gcgagtctag cagcaaccaa tagctatgag cgagaggcgg gactctgagg           300
gaagtcaatc gctgccgcag gtaccgcaa tggcttttgg cggggcgctt           350
ccccaacctt gccctctctc atgaccccgc tccgggatta tggccgggac           400
tggggtgctg gcgctgcgga cgctgccagg gccagctgg gtgcgaggct           450
cggggccctc cgtgctgagc cgcctgcagg acgcggccgt ggtgcggcct           500
ggcttcctga gcacggcaga ggaggagacg ctgagccgag aactggagcc           550
cgagctgcgc cgccgcgctt acgaatacga tcaactgggac gcggccatcc           600
acggcttcgc agagacagag aagtccgctt ggtcagaagc cagccggggc           650
atcctgcagc gcgtgcaggc ggccgccttt ggccccggcc agaccctgct           700
ctcctccgtg cactgtctgg acctggaagc ccgcggctac atcaagcccc           750
acgtggacag catcaagttc tgcggggcca ccctgcgagg cctgtctctc           800
ctgtctccca gcgttatgcg gctggtgcac acccaggagc cgggggagtg           850
gtggaactc ttgctggagc cgggctccct ctacatcett aggggctcag           900
cccgttatga cttctcccat gagatccttc gggatgaaga gtccttcttt           950
ggggaacgoc ggattccccg gggccggcgc atctccgtga tctgccgctc          1000
cctccctgag ggcatggggc caggggagtc tggacagccg cccccagcct          1050
    
```

-continued

---

 gctgaccccc agctttctac agacaccaga tttgtgaata aagttgggga 1100

atggacagcc t 1111

&lt;210&gt; SEQ ID NO 102

&lt;211&gt; LENGTH: 221

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 102

Met Ala Gly Thr Gly Leu Leu Ala Leu Arg Thr Leu Pro Gly Pro  
1 5 10 15Ser Trp Val Arg Gly Ser Gly Pro Ser Val Leu Ser Arg Leu Gln  
20 25 30Asp Ala Ala Val Val Arg Pro Gly Phe Leu Ser Thr Ala Glu Glu  
35 40 45Glu Thr Leu Ser Arg Glu Leu Glu Pro Glu Leu Arg Arg Arg Arg  
50 55 60Tyr Glu Tyr Asp His Trp Asp Ala Ala Ile His Gly Phe Arg Glu  
65 70 75Thr Glu Lys Ser Arg Trp Ser Glu Ala Ser Arg Ala Ile Leu Gln  
80 85 90Arg Val Gln Ala Ala Ala Phe Gly Pro Gly Gln Thr Leu Leu Ser  
95 100 105Ser Val His Val Leu Asp Leu Glu Ala Arg Gly Tyr Ile Lys Pro  
110 115 120His Val Asp Ser Ile Lys Phe Cys Gly Ala Thr Ile Ala Gly Leu  
125 130 135Ser Leu Leu Ser Pro Ser Val Met Arg Leu Val His Thr Gln Glu  
140 145 150Pro Gly Glu Trp Leu Glu Leu Leu Leu Glu Pro Gly Ser Leu Tyr  
155 160 165Ile Leu Arg Gly Ser Ala Arg Tyr Asp Phe Ser His Glu Ile Leu  
170 175 180Arg Asp Glu Glu Ser Phe Phe Gly Glu Arg Arg Ile Pro Arg Gly  
185 190 195Arg Arg Ile Ser Val Ile Cys Arg Ser Leu Pro Glu Gly Met Gly  
200 205 210Pro Gly Glu Ser Gly Gln Pro Pro Pro Ala Cys  
215 220

&lt;210&gt; SEQ ID NO 103

&lt;211&gt; LENGTH: 3583

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 103

ctccccggcg ccgcaggcag cgtcctccto cgaagcagct gcacctgcaa 50

ctgggcagcc tggaccctcg tgccctgttc cggggacctc ggcaggggg 100

cgccccggga caccctctgc gggccgggtg gaggaggaag aggaggagga 150

ggaagaagac gtggacaag accccatcc taccagaac acctgcctgc 200

gctgcccca cttctcttta agggagagga aaagagagcc taggagaacc 250

-continued

---

atggggggct gcgaagtccg ggaatttctt ttgcaatttg gtttcttctt	300
gcctctgctg acagcgtggc caggcgactg cagtcacgtc tccaacaacc	350
aagttgtggt gcttgataca acaactgtac tgggagagct aggatggaaa	400
acatatccat taaatgggtg ggatgccatc actgaaatgg atgaacataa	450
taggcccatt cacacatacc aggtatgtaa tgtaatggaa ccaaaccaaa	500
acaactggct tcgtacaaac tggatctccc gtgatgcagc tcagaaaatt	550
tatgtgtaaa tgaattcac actaagggat tgtaacagca tcccatgggt	600
cttggggact tgcaaagaaa catttaactc gttttatatg gaatcagatg	650
agtcccacgg aattaaattc aagccaaacc agtatacaaa gatcgacaca	700
attgctgctg atgagagttt taaccagatg gatttgggtg atcgcaccc	750
caaactcaac actgaaattc gtgaggtggg gcctatagaa aggaaaggat	800
tttatctggc tttcaagac attggggcgt gcattgcctt ggtttcagtc	850
cgtgttttct acaagaaatg ccccttcaact gttcgttaact tggccatggt	900
tcctgatacc attccaaggg ttgattcctc ctctttggtt gaagtacggg	950
gttcttgtgt gaagagtgct gaagagcgtg acaactcctaa actgtattgt	1000
ggagctgatg gagattggct ggttcctctt ggaaggtgca tctgcagtac	1050
aggatatgaa gaaattgagg gttcttgcca tgcttgacaga ccaggattct	1100
ataaagcttt tgctgggaac acaaaatggt ctaaatgtcc tccacacagt	1150
ttaacataca tggagcaaac ttctgtctgt cagtgtgaaa agggttattt	1200
ccgagctgaa aaagaccac cttctatggc atgtaccagg ccaccttcag	1250
ctcctaggaa tgtggttttt aacatcaatg aaacagccct tattttggaa	1300
tggagcccac caagtgcac agggaggaga aaagatctca catacagtgt	1350
aatctgtaag aaatgtggct tagacaccag ccagtgtgag gactgtgggtg	1400
gaggactccg cttcatocca agacatacag gcctgatcaa caattccgtg	1450
atagtacttg actttgtgtc tcacgtgaat tacacctttg aaatagaagc	1500
aatgaatgga gtttctgagt tgagtttttc tccaagcca tcacagcta	1550
ttacagtgac cacggatcaa gatgcacctt ccctgatagg tgtggttaagg	1600
aaggactggg catcccaaaa tagcattgcc ctatcatggc aagcacctgc	1650
tttttccaat ggagccattc tggactacga gatcaagtac tatgagaaa	1700
aacatgagca gctgacctac tcttccacaa ggtccaaagc ccccagtgtc	1750
atcatcacag gtcttaagcc agccacaaa tatgtatttc acatccgagt	1800
gagaactgcg acaggataca gtggctacag tcagaaaattt gaatttgaaa	1850
caggagatga aacttctgac atggcagcag aacaaggaca gatttctgtg	1900
atagccacgg ccgctgttgg cggattcact ctccctcgta tcctcacttt	1950
attcttcttg atcactggga gatgtcagtg gtacataaaa gccaaagatga	2000
agtcagaaga gaagagaaga aaccacttac agaatgggca tttgcgcttc	2050
ccgggaatta aaacttacat tgatccagat acatatgaag acccatccct	2100
agcagtcatt gaatttgcaa aggagattga tccctcaaga attcgtattg	2150

-continued

---

```

agagagtcat tggggcaggt gaatttgag aagtctgtag tgggcgtttg      2200
aagacaccag gaaaagaga gatcccagtt gccattaaaa cttgaaagg      2250
tggccacatg gatcggcaaa gaagagattt tctaagagaa gctagtatca      2300
tgggccagtt tgaccatcca aacatcattc gcctagaagg gttgtgcacc      2350
aaaagatcct tcccgcccat tggggtggag gcgttttgcc ccagcttcct      2400
gagggcaggg tttttaaata gcatccaggc cccgcatcca gtgccagggg      2450
gaggatcttt gccccccagg attcctgctg gcagaccagt aatgattgtg      2500
gtggaatata tggagaatgg atccctagac tcctttttgc ggaagcatga      2550
tggccacttc acagtcatcc agttggtcgg aatgctccga ggcattgcat      2600
caggcatgaa gtatctttct gatatgggtt atgttcatcg agaactagcg      2650
gctcggaaata tactggtcaa tagcaactta gtatgcaaag tttctgattt      2700
tggctctctc agagtgctgg aagatgatcc agaagctgct tatacaacaa      2750
ctggtgaaa aatccccata aggtggacag cccagaagc catgcctac      2800
agaaaattct cctcagcaag cgatgcatgg agctatggca ttgtcatgtg      2850
ggaggtcatg tcctatggag agagacctta ttgggaaatg tctaaccaag      2900
atgtcattct gtccattgaa gaagggtaca gacttccagc tcccatgggc      2950
tgtccagcat ctctacacca gctgatgctc cactgctggc agaaggagag      3000
aaatcacaga ccaaaattta ctgacattgt cagcttcctt gacaaaactga      3050
tccgaaatcc cagtgcctct cacacctgg tggaggacat cttgtaatg      3100
ccagagtccc ctggtgaagt tccggaatat cctttgtttg tcacagtgg      3150
tgactggcta gattctataa agatggggca atacaagaat aacttcgtgg      3200
cagcaggggt tacaacattt gacctgattt caagaatgag cattgatgac      3250
attagaagaa ttggagtcat acttattgga caccagagac gaatagtcag      3300
cagcatacag actttacggt tacacatgat gcacatacag gagaagggat      3350
ttcatgtatg aaagtaccac aagcacctgt gttttgtgcc tcagcatttc      3400
taaaatgaac gatatcctct ctactactct ctcttctgat tctccaaaca      3450
tcacttcaca aactgcagtc ttctgttcag actataggca cacaccttat      3500
gtttatgctt ccaaccagga ttttaaaatc atgctacata aatccgttct      3550
gaataacctg caactaaaaa aaaaaaaaaa aaa      3583

```

```

<210> SEQ ID NO 104
<211> LENGTH: 1036
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

```

<400> SEQUENCE: 104

```

Met Gly Gly Cys Glu Val Arg Glu Phe Leu Leu Gln Phe Gly Phe
 1           5           10          15
Phe Leu Pro Leu Leu Thr Ala Trp Pro Gly Asp Cys Ser His Val
 20          25          30
Ser Asn Asn Gln Val Val Leu Leu Asp Thr Thr Thr Val Leu Gly
 35          40          45
Glu Leu Gly Trp Lys Thr Tyr Pro Leu Asn Gly Trp Asp Ala Ile

```

-continued

															50	55	60
Thr	Glu	Met	Asp	Glu	His	Asn	Arg	Pro	Ile	His	Thr	Tyr	Gln	Val	65	70	75
Cys	Asn	Val	Met	Glu	Pro	Asn	Gln	Asn	Asn	Trp	Leu	Arg	Thr	Asn	80	85	90
Trp	Ile	Ser	Arg	Asp	Ala	Ala	Gln	Lys	Ile	Tyr	Val	Glu	Met	Lys	95	100	105
Phe	Thr	Leu	Arg	Asp	Cys	Asn	Ser	Ile	Pro	Trp	Val	Leu	Gly	Thr	110	115	120
Cys	Lys	Glu	Thr	Phe	Asn	Leu	Phe	Tyr	Met	Glu	Ser	Asp	Glu	Ser	125	130	135
His	Gly	Ile	Lys	Phe	Lys	Pro	Asn	Gln	Tyr	Thr	Lys	Ile	Asp	Thr	140	145	150
Ile	Ala	Ala	Asp	Glu	Ser	Phe	Thr	Gln	Met	Asp	Leu	Gly	Asp	Arg	155	160	165
Ile	Leu	Lys	Leu	Asn	Thr	Glu	Ile	Arg	Glu	Val	Gly	Pro	Ile	Glu	170	175	180
Arg	Lys	Gly	Phe	Tyr	Leu	Ala	Phe	Gln	Asp	Ile	Gly	Ala	Cys	Ile	185	190	195
Ala	Leu	Val	Ser	Val	Arg	Val	Phe	Tyr	Lys	Lys	Cys	Pro	Phe	Thr	200	205	210
Val	Arg	Asn	Leu	Ala	Met	Phe	Pro	Asp	Thr	Ile	Pro	Arg	Val	Asp	215	220	225
Ser	Ser	Ser	Leu	Val	Glu	Val	Arg	Gly	Ser	Cys	Val	Lys	Ser	Ala	230	235	240
Glu	Glu	Arg	Asp	Thr	Pro	Lys	Leu	Tyr	Cys	Gly	Ala	Asp	Gly	Asp	245	250	255
Trp	Leu	Val	Pro	Leu	Gly	Arg	Cys	Ile	Cys	Ser	Thr	Gly	Tyr	Glu	260	265	270
Glu	Ile	Glu	Gly	Ser	Cys	His	Ala	Cys	Arg	Pro	Gly	Phe	Tyr	Lys	275	280	285
Ala	Phe	Ala	Gly	Asn	Thr	Lys	Cys	Ser	Lys	Cys	Pro	Pro	His	Ser	290	295	300
Leu	Thr	Tyr	Met	Glu	Ala	Thr	Ser	Val	Cys	Gln	Cys	Glu	Lys	Gly	305	310	315
Tyr	Phe	Arg	Ala	Glu	Lys	Asp	Pro	Pro	Ser	Met	Ala	Cys	Thr	Arg	320	325	330
Pro	Pro	Ser	Ala	Pro	Arg	Asn	Val	Val	Phe	Asn	Ile	Asn	Glu	Thr	335	340	345
Ala	Leu	Ile	Leu	Glu	Trp	Ser	Pro	Pro	Ser	Asp	Thr	Gly	Gly	Arg	350	355	360
Lys	Asp	Leu	Thr	Tyr	Ser	Val	Ile	Cys	Lys	Lys	Cys	Gly	Leu	Asp	365	370	375
Thr	Ser	Gln	Cys	Glu	Asp	Cys	Gly	Gly	Gly	Leu	Arg	Phe	Ile	Pro	380	385	390
Arg	His	Thr	Gly	Leu	Ile	Asn	Asn	Ser	Val	Ile	Val	Leu	Asp	Phe	395	400	405
Val	Ser	His	Val	Asn	Tyr	Thr	Phe	Glu	Ile	Glu	Ala	Met	Asn	Gly	410	415	420
Val	Ser	Glu	Leu	Ser	Phe	Ser	Pro	Lys	Pro	Phe	Thr	Ala	Ile	Thr	425	430	435



-continued

---

Val Thr Thr Asp	Gln Asp Ala Pro Ser	Leu Ile Gly Val Val	Arg
440		445	450
Lys Asp Trp Ala	Ser Gln Asn Ser Ile	Ala Leu Ser Trp Gln	Ala
455		460	465
Pro Ala Phe Ser	Asn Gly Ala Ile Leu	Asp Tyr Glu Ile Lys	Tyr
470		475	480
Tyr Glu Lys Glu	His Glu Gln Leu Thr	Tyr Ser Ser Thr Arg	Ser
485		490	495
Lys Ala Pro Ser	Val Ile Ile Thr Gly	Leu Lys Pro Ala Thr	Lys
500		505	510
Tyr Val Phe His	Ile Arg Val Arg Thr	Ala Thr Gly Tyr Ser	Gly
515		520	525
Tyr Ser Gln Lys	Phe Glu Phe Glu Thr	Gly Asp Glu Thr Ser	Asp
530		535	540
Met Ala Ala Glu	Gln Gly Gln Ile Leu	Val Ile Ala Thr Ala	Ala
545		550	555
Val Gly Gly Phe	Thr Leu Leu Val Ile	Leu Thr Leu Phe Phe	Leu
560		565	570
Ile Thr Gly Arg	Cys Gln Trp Tyr Ile	Lys Ala Lys Met Lys	Ser
575		580	585
Glu Glu Lys Arg	Arg Asn His Leu Gln	Asn Gly His Leu Arg	Phe
590		595	600
Pro Gly Ile Lys	Thr Tyr Ile Asp Pro	Asp Thr Tyr Glu Asp	Pro
605		610	615
Ser Leu Ala Val	His Glu Phe Ala Lys	Glu Ile Asp Pro Ser	Arg
620		625	630
Ile Arg Ile Glu	Arg Val Ile Gly Ala	Gly Glu Phe Gly Glu	Val
635		640	645
Cys Ser Gly Arg	Leu Lys Thr Pro Gly	Lys Arg Glu Ile Pro	Val
650		655	660
Ala Ile Lys Thr	Leu Lys Gly Gly His	Met Asp Arg Gln Arg	Arg
665		670	675
Asp Phe Leu Arg	Glu Ala Ser Ile Met	Gly Gln Phe Asp His	Pro
680		685	690
Asn Ile Ile Arg	Leu Glu Gly Val Val	Thr Lys Arg Ser Phe	Pro
695		700	705
Ala Ile Gly Val	Glu Ala Phe Cys Pro	Ser Phe Leu Arg Ala	Gly
710		715	720
Phe Leu Asn Ser	Ile Gln Ala Pro His	Pro Val Pro Gly Gly	Gly
725		730	735
Ser Leu Pro Pro	Arg Ile Pro Ala Gly	Arg Pro Val Met Ile	Val
740		745	750
Val Glu Tyr Met	Glu Asn Gly Ser Leu	Asp Ser Phe Leu Arg	Lys
755		760	765
His Asp Gly His	Phe Thr Val Ile Gln	Leu Val Gly Met Leu	Arg
770		775	780
Gly Ile Ala Ser	Gly Met Lys Tyr Leu	Ser Asp Met Gly Tyr	Val
785		790	795
His Arg Asp Leu	Ala Ala Arg Asn Ile	Leu Val Asn Ser Asn	Leu
800		805	810

-continued

---

Val	Cys	Lys	Val	Ser	Asp	Phe	Gly	Leu	Ser	Arg	Val	Leu	Glu	Asp
				815					820				825	
Asp	Pro	Glu	Ala	Ala	Tyr	Thr	Thr	Thr	Gly	Gly	Lys	Ile	Pro	Ile
				830					835				840	
Arg	Trp	Thr	Ala	Pro	Glu	Ala	Ile	Ala	Tyr	Arg	Lys	Phe	Ser	Ser
				845					850				855	
Ala	Ser	Asp	Ala	Trp	Ser	Tyr	Gly	Ile	Val	Met	Trp	Glu	Val	Met
				860					865				870	
Ser	Tyr	Gly	Glu	Arg	Pro	Tyr	Trp	Glu	Met	Ser	Asn	Gln	Asp	Val
				875					880				885	
Ile	Leu	Ser	Ile	Glu	Glu	Gly	Tyr	Arg	Leu	Pro	Ala	Pro	Met	Gly
				890					895				900	
Cys	Pro	Ala	Ser	Leu	His	Gln	Leu	Met	Leu	His	Cys	Trp	Gln	Lys
				905					910				915	
Glu	Arg	Asn	His	Arg	Pro	Lys	Phe	Thr	Asp	Ile	Val	Ser	Phe	Leu
				920					925				930	
Asp	Lys	Leu	Ile	Arg	Asn	Pro	Ser	Ala	Leu	His	Thr	Leu	Val	Glu
				935					940				945	
Asp	Ile	Leu	Val	Met	Pro	Glu	Ser	Pro	Gly	Glu	Val	Pro	Glu	Tyr
				950					955				960	
Pro	Leu	Phe	Val	Thr	Val	Gly	Asp	Trp	Leu	Asp	Ser	Ile	Lys	Met
				965					970				975	
Gly	Gln	Tyr	Lys	Asn	Asn	Phe	Val	Ala	Ala	Gly	Phe	Thr	Thr	Phe
				980					985				990	
Asp	Leu	Ile	Ser	Arg	Met	Ser	Ile	Asp	Asp	Ile	Arg	Arg	Ile	Gly
				995					1000				1005	
Val	Ile	Leu	Ile	Gly	His	Gln	Arg	Arg	Ile	Val	Ser	Ser	Ile	Gln
				1010					1015				1020	
Thr	Leu	Arg	Leu	His	Met	Met	His	Ile	Gln	Glu	Lys	Gly	Phe	His
				1025					1030				1035	

Val

<210> SEQ ID NO 105  
 <211> LENGTH: 2148  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 105

ggcggcgggc tgcgcgagc ggcgtcccct gcagccgcyg accgagcag	50
cggcgccacc tgccggccga gcaatgcaa gtgagtacac ctatgtgaaa	100
ctgagaagtg attgctcgag gccttccctg caatggtaca cccgagctca	150
aagcaagatg agaaggccca gcttgttatt aaaagacatc ctcaaagtga	200
cattgcttgt gtttgagtg tggatccttt atatcctcaa gttaaattat	250
actactgaa aatgtgacat gaaaaaatg cattatgtgg accctgacca	300
tgtaaagaga gctcagaaat atgctcagca agtcttgca gaggaaatgc	350
gtcccaagtt tgccaagaca tcaatggcgc tgttatttga gcacaggtat	400
agcgtggact tactcccttt tgtgcagaag gccccaaag acagtgaagc	450
tgagtccaag tacgatcctc cttttgggtt coggaagttc tccagtaaag	500
tccagacctt cttggaactc ttgccagagc acgacctccc tgaacacttg	550

-continued

---

aaagccaaga cctgtcggcg ctgtgtggtt attggaagcg gaggaatact	600
gcacggatta gaactgggcc acaccctgaa ccagttcgat gttgtgataa	650
ggttaaacag tgcaccagtt gagggatatt cagaacatgt tggaaataaa	700
actactataa ggatgactta tccagagggc gcaccactgt ctgacctga	750
atattattcc aatgacttat ttgttctgtt tttatttaag agtgttgatt	800
tcaactggct tcaagcaatg gtaaaaaagg aaaccctgcc attctgggta	850
cgactctctt tttggaagca ggtggcagaa aaaatccac tgacagccaaa	900
acatttcagg attttgaatc cagttatcat caaagagact gcctttgaca	950
tccttcagta ctacagacct cagtcaaggt tctggggccg agataagaac	1000
gtccccacaa tcggtgtcat tgccgttctc ttagccacac atctgtgcga	1050
tgaagtcaag ttggcggggt ttggatata cctcaatcaa cccagaacac	1100
ctttgcacta cttcgacagt caatgcattg ctgctatgaa ctttcagacc	1150
atgcataatg tgacaacgga aaccaagttc ctcttaaagc tggtaaaga	1200
gggagtggtg aaagatctca gtggaggcat tgatcgtgaa tttgaaacac	1250
agaaaaacct agttgaaaat gcaactctaa ctctgagagc tgtttttgac	1300
agccttcttg atgtatttct ccatcctgca gatactttga agtgcagctc	1350
atgtttttta cttttaattt aaaaacacaa aaaaaatttt agctcttccc	1400
actttttttt tcctatttat ttgaggtcag tgtttgtttt tgcacaccat	1450
tttgtaaag aaacttaaga attgaattgg aaagacttct caaagagaat	1500
tgatgtaac gatgttgat tgatttttaa gaaagtaatt taatttgtaa	1550
aactctgct cgtttacact gcacattgaa tacaggtaac taattggaag	1600
gagaggggag gtcactcttt tgatgggtgc cctgaacctc attctgggtc	1650
cctgtctgcg tgcttggtgt gaccacgga ggatccactc ccaggatgac	1700
gtgctccgta gctctgctgc tgatactggg tctgcatgc agcggcgtga	1750
ggcctgggct ggttgagaa ggtcacaacc cttctctggt ggtctgcctt	1800
ctgctgaaa actcagaaac caaccagga agctgtcctg gagtccctg	1850
gtcggagag gacatagaat ctgtgacctc tgacaactgt gaagccaccc	1900
tgggctacag aaaccacagt cttcccagca attattaca ttcttgaatt	1950
ccttgggat ttttactgc ctttcaaag cacttaagtg ttagatctaa	2000
cggttccag tgtctgtctg aggtgactta aaaaatcaga acaaaacttc	2050
tattatccag agtcatggga gagtacacc tttccaggaa taatgttttg	2100
ggaaacactg aatgaaatc ttcccagtat tataaattgt gtatttaa	2148

&lt;210&gt; SEQ ID NO 106

&lt;211&gt; LENGTH: 362

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 106

Met	Arg	Arg	Pro	Ser	Leu	Leu	Leu	Lys	Asp	Ile	Leu	Lys	Cys	Thr
1				5				10						15

-continued

Leu	Leu	Val	Phe	Gly	Val	Trp	Ile	Leu	Tyr	Ile	Leu	Lys	Leu	Asn
				20					25					30
Tyr	Thr	Thr	Glu	Glu	Cys	Asp	Met	Lys	Lys	Met	His	Tyr	Val	Asp
				35					40					45
Pro	Asp	His	Val	Lys	Arg	Ala	Gln	Lys	Tyr	Ala	Gln	Gln	Val	Leu
				50					55					60
Gln	Lys	Glu	Cys	Arg	Pro	Lys	Phe	Ala	Lys	Thr	Ser	Met	Ala	Leu
				65					70					75
Leu	Phe	Glu	His	Arg	Tyr	Ser	Val	Asp	Leu	Leu	Pro	Phe	Val	Gln
				80					85					90
Lys	Ala	Pro	Lys	Asp	Ser	Glu	Ala	Glu	Ser	Lys	Tyr	Asp	Pro	Pro
				95					100					105
Phe	Gly	Phe	Arg	Lys	Phe	Ser	Ser	Lys	Val	Gln	Thr	Leu	Leu	Glu
				110					115					120
Leu	Leu	Pro	Glu	His	Asp	Leu	Pro	Glu	His	Leu	Lys	Ala	Lys	Thr
				125					130					135
Cys	Arg	Arg	Cys	Val	Val	Ile	Gly	Ser	Gly	Gly	Ile	Leu	His	Gly
				140					145					150
Leu	Glu	Leu	Gly	His	Thr	Leu	Asn	Gln	Phe	Asp	Val	Val	Ile	Arg
				155					160					165
Leu	Asn	Ser	Ala	Pro	Val	Glu	Gly	Tyr	Ser	Glu	His	Val	Gly	Asn
				170					175					180
Lys	Thr	Thr	Ile	Arg	Met	Thr	Tyr	Pro	Glu	Gly	Ala	Pro	Leu	Ser
				185					190					195
Asp	Leu	Glu	Tyr	Tyr	Ser	Asn	Asp	Leu	Phe	Val	Ala	Val	Leu	Phe
				200					205					210
Lys	Ser	Val	Asp	Phe	Asn	Trp	Leu	Gln	Ala	Met	Val	Lys	Lys	Glu
				215					220					225
Thr	Leu	Pro	Phe	Trp	Val	Arg	Leu	Phe	Phe	Trp	Lys	Gln	Val	Ala
				230					235					240
Glu	Lys	Ile	Pro	Leu	Gln	Pro	Lys	His	Phe	Arg	Ile	Leu	Asn	Pro
				245					250					255
Val	Ile	Ile	Lys	Glu	Thr	Ala	Phe	Asp	Ile	Leu	Gln	Tyr	Ser	Glu
				260					265					270
Pro	Gln	Ser	Arg	Phe	Trp	Gly	Arg	Asp	Lys	Asn	Val	Pro	Thr	Ile
				275					280					285
Gly	Val	Ile	Ala	Val	Val	Leu	Ala	Thr	His	Leu	Cys	Asp	Glu	Val
				290					295					300
Ser	Leu	Ala	Gly	Phe	Gly	Tyr	Asp	Leu	Asn	Gln	Pro	Arg	Thr	Pro
				305					310					315
Leu	His	Tyr	Phe	Asp	Ser	Gln	Cys	Met	Ala	Ala	Met	Asn	Phe	Gln
				320					325					330
Thr	Met	His	Asn	Val	Thr	Thr	Glu	Thr	Lys	Phe	Leu	Leu	Lys	Leu
				335					340					345
Val	Lys	Glu	Gly	Val	Val	Lys	Asp	Leu	Ser	Gly	Gly	Ile	Asp	Arg
				350					355					360

Glu Phe

&lt;210&gt; SEQ ID NO 107

&lt;211&gt; LENGTH: 1399

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapien

-continued

<400> SEQUENCE: 107

tgacgcgggg cgccagctgc caacttcgcg cgcggagctc cccggcgggtg	50
cagtcccgtc ccggcggcgc gggcggcatg aagactagcc gccgcggccg	100
agcgctcctg gccgtggccc tgaacctgct ggcgctgctg ttcgccacca	150
ccgctttcct caccacgcac tggtgccagg gcacgcagcg ggtccccaag	200
ccgggctgcg gccagggcgg gcgcgccaac tgccccaact cgggcgccaa	250
cgccacggcc aacggcaccg ccgccccgcg cgcgcgcccc gccgccgcca	300
ccgcctcggg gaacggcccc cctggcggcg cgctctacag ctgggagacc	350
ggcgacgacc gcttcctcct caggaatttc cacaccgca tctggtactc	400
gtgcgaggag gagctcagcg ggcttggtga aaaatgtcgc agcttcattg	450
acctggcccc ggcgtcggag aaaggcctcc tgggaatggt cgcccacatg	500
atgtacacgc aggtgttcca ggtcaccgtg agcctcggtc ctgaggactg	550
gagaccccat tcctgggact acgggtggtc cttctgcctg gcgtggggct	600
cctttacctg ctgcatggca gcctctgtca ccacgctcaa ctctacacc	650
aagacggta ttgagttccg gcacaagcgc aaggtctttg agcagggcta	700
ccgggaagag ccgacctca tagacctga ggccatcaag tacttccggg	750
agaggatgga gaagagggac gggagcagag aggactttca cttagactgc	800
cgccacgaga gataccctgc ccgacaccag ccacacatgg cggattcctg	850
gccccggagc tccgcacagg aagcaccaga gctgaaccga cagtgtgagg	900
tcttggggca ctgggtgtga ccaagacctc aacctggccc gcggacctca	950
ggccatcgct ggcaccagcc cctgctgcaa gaccaccaga gtggtgcccc	1000
cagaacctg gcctgtgtgc cgtgaactca gtcagcctgc gtgggagatg	1050
ccaggcctgt cctgcccatc gctgcctggg tcccatggcc ttggaaatgg	1100
ggccagggca ggcccaaggg aatgcacagg gctgcacaga gtgactttgg	1150
gacagcagcc ccggactcct gccatcatca catgagccct gctgggcaca	1200
gctgcgatgc caggagacac atggccactg gccactgaat ggctggcacc	1250
cacaagccag tcaggtgccc agaggggcag agccctttgg ggggcagaga	1300
gtggcttcct gaaggagggg gcagtggcgc aggcactgca ggggtgtcac	1350
acagcaggca cacagcaggg gctcaataaa tgcttgttga acttgtttt	1399

<210> SEQ ID NO 108

<211> LENGTH: 280

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 108

Met	Lys	Thr	Ser	Arg	Arg	Gly	Arg	Ala	Leu	Leu	Ala	Val	Ala	Leu
1				5					10					15
Asn	Leu	Leu	Ala	Leu	Leu	Phe	Ala	Thr	Thr	Ala	Phe	Leu	Thr	Thr
				20					25					30
His	Trp	Cys	Gln	Gly	Thr	Gln	Arg	Val	Pro	Lys	Pro	Gly	Cys	Gly
				35					40					45

-continued

Gln	Gly	Gly	Arg	Ala	Asn	Cys	Pro	Asn	Ser	Gly	Ala	Asn	Ala	Thr	
				50					55					60	
Ala	Asn	Gly	Thr	Ala	Ala	Pro	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Thr	
				65					70					75	
Ala	Ser	Gly	Asn	Gly	Pro	Pro	Gly	Gly	Ala	Leu	Tyr	Ser	Trp	Glu	
				80					85					90	
Thr	Gly	Asp	Asp	Arg	Phe	Leu	Phe	Arg	Asn	Phe	His	Thr	Gly	Ile	
				95					100					105	
Trp	Tyr	Ser	Cys	Glu	Glu	Glu	Leu	Ser	Gly	Leu	Gly	Glu	Lys	Cys	
				110					115					120	
Arg	Ser	Phe	Ile	Asp	Leu	Ala	Pro	Ala	Ser	Glu	Lys	Gly	Leu	Leu	
				125					130					135	
Gly	Met	Val	Ala	His	Met	Met	Tyr	Thr	Gln	Val	Phe	Gln	Val	Thr	
				140					145					150	
Val	Ser	Leu	Gly	Pro	Glu	Asp	Trp	Arg	Pro	His	Ser	Trp	Asp	Tyr	
				155					160					165	
Gly	Trp	Ser	Phe	Cys	Leu	Ala	Trp	Gly	Ser	Phe	Thr	Cys	Cys	Met	
				170					175					180	
Ala	Ala	Ser	Val	Thr	Thr	Leu	Asn	Ser	Tyr	Thr	Lys	Thr	Val	Ile	
				185					190					195	
Glu	Phe	Arg	His	Lys	Arg	Lys	Val	Phe	Glu	Gln	Gly	Tyr	Arg	Glu	
				200					205					210	
Glu	Pro	Thr	Phe	Ile	Asp	Pro	Glu	Ala	Ile	Lys	Tyr	Phe	Arg	Glu	
				215					220					225	
Arg	Met	Glu	Lys	Arg	Asp	Gly	Ser	Glu	Glu	Asp	Phe	His	Leu	Asp	
				230					235					240	
Cys	Arg	His	Glu	Arg	Tyr	Pro	Ala	Arg	His	Gln	Pro	His	Met	Ala	
				245					250					255	
Asp	Ser	Trp	Pro	Arg	Ser	Ser	Ala	Gln	Glu	Ala	Pro	Glu	Leu	Asn	
				260					265					270	
Arg	Gln	Cys	Trp	Val	Leu	Gly	His	Trp	Val						
				275					280						

<210> SEQ ID NO 109  
 <211> LENGTH: 2964  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 109

gattaccaag caagaacagc taaaatgaaa gccatcattc atcttactct	50
tcttgctctc ctttctgtaa acacagccac caaccaaggc aactcagctg	100
atgctgtaac aaccacagaa actgcgacta gtggctctac agtagctgca	150
gctgatacca ctgaaactaa tttcctgaa actgctagca ccacagcaaa	200
tacaccttot ttccaacag ctacttcacc tgctccccc ataattagta	250
cacatagttc ctccacaatt cctacacctg ctcccccat aattagtaca	300
catagttctc ccacaattcc tatactact gctgcagaca gtgagtcaac	350
cacaaatgta aattcattag ctacctctga cataatcacc gcttcatctc	400
caaatgatgg attaatacaca atgggtcctt ctgaacaca aagtaacaat	450
gaaatgtccc ccaccacaga agacaatcaa tcatcagggc ctcccactgg	500

-continued

---

caccgcttta ttggagacca gcaccctaaa cagcacaggt cccagcaatc	550
cttgccaaga tgatccctgt gcagataatt cgttatgtgt taagctgcat	600
aatacaagtt tttgcctgtg tttagaaggg tattactaca actcttctac	650
atgtaagaaa ggaaggtat tccttgggaa gatttcagtg acagtatcag	700
aaacatttga cccagaagag aaacattcca tggcctatca agacttgcat	750
agtgaatta ctagcttggt taaagatgta tttggccat ctgtttatgg	800
acagactgta attcttactg taagcacatc tctgtcacca agatctgaaa	850
tgcgtgctga tgacaagttt gttaatgtaa caatagtaac aattttggca	900
gaaaccacaa gtgacaatga gaagactgtg actgagaaaa ttaataaagc	950
aattagaagt agctcaagca actttctaaa ctatgatttg acccttgggt	1000
gtgattatta tggctgtaac cagactgctg atgactgcct caatggttta	1050
gcatgctgatt gcaaactctga cctgcaaagg cctaaccac agagcccttt	1100
ctgctgtgct tccagtctca agtgccttga tgcctgcaac gcacagcaca	1150
agcaatgctt aataaagaag agtgggtggg cccctgagtg tgcgtgctg	1200
cccggctacc aggaagatgc taatgggaac tgccaaaagt gtgcatttgg	1250
ctacagtgga ctgactgta aggacaaatt tcagctgatc ctactattg	1300
tgggaccat cgctggcatt gtcattctca gcatgataat tgcattgatt	1350
gtcacagcaa gatcaaataa caaaacgaag catattgaag aagagaactt	1400
gattgacgaa gactttcaaa atctaaaact goggtcgaca ggcttcacca	1450
atcttggagc agaagggagc gtctttccta aggtcaggat aacggcctcc	1500
agagacagcc agatgcaaaa tccttattca agccacagca gcatgccccg	1550
ccctgactat tagaatcata agaattgtga acccgccatg gccccaac	1600
aatgtacaag ctattattta gagtgtttag aaagactgat ggagaagtga	1650
gcaccagtaa agatctggcc tccggggttt ttcttccatc tgacatctgc	1700
cagcctctct gaatggaagt tgtgaatgtt tgcaacgaat ccagctcact	1750
tgctaaataa gaatctatga cattaatgt agtagatgct attagcgctt	1800
gtcagagagg tggttttctt caatcagtac aaagtactga gacaatggtt	1850
agggttgttt tcttaattct tttcctggta gggcaacaag aaccatttcc	1900
aatctagagg aaagctcccc agcattgctt gctcctgggc aaacattgct	1950
cttgagttaa gtgacctaat tcccctggga gacatacgca tcaactgtgg	2000
aggtccgagg ggatgagaag ggataccac catctttcaa gggtcacaag	2050
ctcactctct gacaagtcag aatagggaca ctgcttctat ccctccaatg	2100
gagagattct ggcaaccttt gaacagccca gagcttgcaa cctagcctca	2150
cccaagaaga ctggaaagag acatatctct cagctttttc aggaggcgtg	2200
cctgggaatc caggaacttt ttgatgctaa ttagaaggcc tggactaaaa	2250
atgtccacta tggggtgcac tctacagttt ttgaaatgct aggaggcaga	2300
aggggcagag agtaaaaaac atgacctggt agaaggaaga gaggcaag	2350
aaactgggtg gggaggatca attagagagg aggcacctgg gatccacctt	2400

-continued

---

```

cttccttagg tcccctcctc catcagcaaa ggagcacttc tctaatacatg      2450
cctcccgaa gactggctgg gagaagggtt aaaaacaaaa atccaggag      2500
taagagcctt aggtcagttt gaaattggag acaaaactgtc tggcaaagg      2550
tgcgagaggg agcttgtgct caggagtcca gccgccagc ctcggggtgt      2600
aggtttctga ggtgtgccat tggggcctca gccttctctg gtgacagagg      2650
ctcagctgtg gccaccaaca cacaaccaca cacacacaac cacacacaca      2700
aatgggggga accacatcca gtacaagctt ttacaaatgt tattagtgtc      2750
cttttttatt tctaatagcct tgcctctta aaagttattt tatttgttat      2800
tattatttgt tcttgactgt taattgtgaa tggaatgca ataaagtgc      2850
tttgtagat ggtgaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      2900
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      2950
aaaaaaaaaa aaaa      2964
    
```

```

<210> SEQ ID NO 110
<211> LENGTH: 512
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien
    
```

<400> SEQUENCE: 110

```

Met Lys Ala Ile Ile His Leu Thr Leu Leu Ala Leu Leu Ser Val
 1             5             10             15
Asn Thr Ala Thr Asn Gln Gly Asn Ser Ala Asp Ala Val Thr Thr
          20             25             30
Thr Glu Thr Ala Thr Ser Gly Pro Thr Val Ala Ala Ala Asp Thr
          35             40             45
Thr Glu Thr Asn Phe Pro Glu Thr Ala Ser Thr Thr Ala Asn Thr
          50             55             60
Pro Ser Phe Pro Thr Ala Thr Ser Pro Ala Pro Pro Ile Ile Ser
          65             70             75
Thr His Ser Ser Ser Thr Ile Pro Thr Pro Ala Pro Pro Ile Ile
          80             85             90
Ser Thr His Ser Ser Ser Thr Ile Pro Ile Pro Thr Ala Ala Asp
          95             100            105
Ser Glu Ser Thr Thr Asn Val Asn Ser Leu Ala Thr Ser Asp Ile
          110            115            120
Ile Thr Ala Ser Ser Pro Asn Asp Gly Leu Ile Thr Met Val Pro
          125            130            135
Ser Glu Thr Gln Ser Asn Asn Glu Met Ser Pro Thr Thr Glu Asp
          140            145            150
Asn Gln Ser Ser Gly Pro Pro Thr Gly Thr Ala Leu Leu Glu Thr
          155            160            165
Ser Thr Leu Asn Ser Thr Gly Pro Ser Asn Pro Cys Gln Asp Asp
          170            175            180
Pro Cys Ala Asp Asn Ser Leu Cys Val Lys Leu His Asn Thr Ser
          185            190            195
Phe Cys Leu Cys Leu Glu Gly Tyr Tyr Tyr Asn Ser Ser Thr Cys
          200            205            210
Lys Lys Gly Lys Val Phe Pro Gly Lys Ile Ser Val Thr Val Ser
          215            220            225
    
```



-continued

Glu Thr Phe Asp Pro Glu Glu Lys His Ser Met Ala Tyr Gln Asp  
 230 235 240  
 Leu His Ser Glu Ile Thr Ser Leu Phe Lys Asp Val Phe Gly Thr  
 245 250 255  
 Ser Val Tyr Gly Gln Thr Val Ile Leu Thr Val Ser Thr Ser Leu  
 260 265 270  
 Ser Pro Arg Ser Glu Met Arg Ala Asp Asp Lys Phe Val Asn Val  
 275 280 285  
 Thr Ile Val Thr Ile Leu Ala Glu Thr Thr Ser Asp Asn Glu Lys  
 290 295 300  
 Thr Val Thr Glu Lys Ile Asn Lys Ala Ile Arg Ser Ser Ser Ser  
 305 310 315  
 Asn Phe Leu Asn Tyr Asp Leu Thr Leu Arg Cys Asp Tyr Tyr Gly  
 320 325 330  
 Cys Asn Gln Thr Ala Asp Asp Cys Leu Asn Gly Leu Ala Cys Asp  
 335 340 345  
 Cys Lys Ser Asp Leu Gln Arg Pro Asn Pro Gln Ser Pro Phe Cys  
 350 355 360  
 Val Ala Ser Ser Leu Lys Cys Pro Asp Ala Cys Asn Ala Gln His  
 365 370 375  
 Lys Gln Cys Leu Ile Lys Lys Ser Gly Gly Ala Pro Glu Cys Ala  
 380 385 390  
 Cys Val Pro Gly Tyr Gln Glu Asp Ala Asn Gly Asn Cys Gln Lys  
 395 400 405  
 Cys Ala Phe Gly Tyr Ser Gly Leu Asp Cys Lys Asp Lys Phe Gln  
 410 415 420  
 Leu Ile Leu Thr Ile Val Gly Thr Ile Ala Gly Ile Val Ile Leu  
 425 430 435  
 Ser Met Ile Ile Ala Leu Ile Val Thr Ala Arg Ser Asn Asn Lys  
 440 445 450  
 Thr Lys His Ile Glu Glu Glu Asn Leu Ile Asp Glu Asp Phe Gln  
 455 460 465  
 Asn Leu Lys Leu Arg Ser Thr Gly Phe Thr Asn Leu Gly Ala Glu  
 470 475 480  
 Gly Ser Val Phe Pro Lys Val Arg Ile Thr Ala Ser Arg Asp Ser  
 485 490 495  
 Gln Met Gln Asn Pro Tyr Ser Ser His Ser Ser Met Pro Arg Pro  
 500 505 510

Asp Tyr

<210> SEQ ID NO 111  
 <211> LENGTH: 943  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien  
 <400> SEQUENCE: 111

ctgggacttg gctttctccg gataagcggc ggcaccggcg tcagcgatga 50  
 ccgtgcagag actcgtggcc gcggccgtgc tggtagccct ggtctcactc 100  
 atcctcaaca acgtggcggc cttcacctcc aactgggtgt gccagacgct 150  
 ggaggatggg cgcagggcga gcgtggggct gtggaggtcc tgctggctgg 200

-continued

---

```

tggacaggac cgggggaggg ccgagccctg gggccagagc cggccagggtg      250
gacgcacatg actgtgaggc gctgggctgg ggctccgagg cagccggctt      300
ccaggagtcc cgaggcaccg tcaaactgca gttcgacatg atgcgcgcct      350
gcaacctggt gggcacggcc gcgctcaccg caggccagct caccttctctc      400
ctggggctgg tgggcctgcc cctgctgtca cccgacgccc cgtgctggga      450
ggaggccatg gccgctgcat tccaactggc gagttttgtc ctggtcacatg      500
ggctcgtgac tttctacaga attggcccat acaccaacct gtcctgggtcc      550
tgctacctga acattggcgc ctgccttctg gccacgctgg cggcagccat      600
gtcatctggg aacattctcc acaagaggga ggactgcatg gcccccggg      650
tgattgtcat cagccgctcc ctgacagcgc gtttcgccg tgggctggac      700
aatgactacg tggagtcacc atgctgagtc gcccttctca gcgctccatc      750
aacgcacacc tgctatcgtg gaacagccta gaaaccaagg gactccacca      800
ccaagtcact tcccctgctc gtgcagaggc acgggatgag tctgggtgac      850
ctctgcgcga tgctgctgag acacgtgtgc gtttactggt atgtcgggtca      900
tatgtctgta cgtgtcgtgg gccaacctcg ttctgcctcc agc      943

```

&lt;210&gt; SEQ ID NO 112

&lt;211&gt; LENGTH: 226

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 112

```

Met Thr Val Gln Arg Leu Val Ala Ala Ala Val Leu Val Ala Leu
 1             5             10             15
Val Ser Leu Ile Leu Asn Asn Val Ala Ala Phe Thr Ser Asn Trp
                20             25             30
Val Cys Gln Thr Leu Glu Asp Gly Arg Arg Arg Ser Val Gly Leu
                35             40             45
Trp Arg Ser Cys Trp Leu Val Asp Arg Thr Arg Gly Gly Pro Ser
                50             55             60
Pro Gly Ala Arg Ala Gly Gln Val Asp Ala His Asp Cys Glu Ala
                65             70             75
Leu Gly Trp Gly Ser Glu Ala Ala Gly Phe Gln Glu Ser Arg Gly
                80             85             90
Thr Val Lys Leu Gln Phe Asp Met Met Arg Ala Cys Asn Leu Val
                95             100            105
Ala Thr Ala Ala Leu Thr Ala Gly Gln Leu Thr Phe Leu Leu Gly
                110            115            120
Leu Val Gly Leu Pro Leu Leu Ser Pro Asp Ala Pro Cys Trp Glu
                125            130            135
Glu Ala Met Ala Ala Ala Phe Gln Leu Ala Ser Phe Val Leu Val
                140            145            150
Ile Gly Leu Val Thr Phe Tyr Arg Ile Gly Pro Tyr Thr Asn Leu
                155            160            165
Ser Trp Ser Cys Tyr Leu Asn Ile Gly Ala Cys Leu Leu Ala Thr
                170            175            180
Leu Ala Ala Ala Met Leu Ile Trp Asn Ile Leu His Lys Arg Glu
                185            190            195

```

-continued

Asp Cys Met Ala Pro Arg Val Ile Val Ile Ser Arg Ser Leu Thr  
 200 205 210  
 Ala Arg Phe Arg Arg Gly Leu Asp Asn Asp Tyr Val Glu Ser Pro  
 215 220 225

Cys

<210> SEQ ID NO 113  
 <211> LENGTH: 1389  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 113

gactttacca ctactcgcta tagagccctg gtcaagttct ctccacctct 50  
 ctatctatgt ctcagtttct tcatctgtaa catcaaatga ataataatac 100  
 caatctccta gacttcataa gaggattaac aaagacaaaa tatgggaaaa 150  
 acataacatg gcgtcccata attattagat cttattattg aactaaaaat 200  
 ggcattaaaa ttacaaaag gaagacagca tctgtttcct ctttggtcct 250  
 gagctggtta aaaggaacac tggttgccctg aacagtcaca cttgcaacca 300  
 tgatgcctaa acattgcttt ctaggcttcc tcatcagttt cttocttact 350  
 ggtgtagcag gaactcagtc aacgcatgag tctctgaagc ctcagagggt 400  
 acaatttcag tcccgaat ttcacaacat tttgcaatgg cagoctggga 450  
 gggcacttac tggcaacagc agtgtctatt ttgtgcagta caaaatata 500  
 ggacagagac aatggaaaaa taaagaagac tgttggggta ctcaagaact 550  
 ctctttgtgac cttaccagtg aaacctcaga catacaggaa cttattacg 600  
 ggagggtagg ggcggcctcg gctgggagct actcagaatg gagcatgacg 650  
 ccgcggttca ctccctggtg ggaaacaaaa atagatcctc cagtcatgaa 700  
 tataaccxaa gtcaatggct ctttgttggg aattctccat gctccaaatt 750  
 taccatatag ataccaaaag gaaaaaatg tatctataga agattactat 800  
 gaactactat accgagtttt tataattaac aattcactag aaaaggagca 850  
 aaaggtttat gaaggggctc acagagcggg tgaattgaa gctctaacac 900  
 cacactccag ctactgtgta gtggctgaaa tataatcagcc catggttagc 950  
 agaagaagtc agagaagtga agagagatgt gtggaaattc catgacttgt 1000  
 ggaattttggc attcagcaat gtggaaattc taaagctccc tgagaacagg 1050  
 atgactcgtg tttgaaggat cttattttaa attgtttttg tattttctta 1100  
 aagcaatatt cactgttaca cttggtggac ttctttgttt acccattctt 1150  
 ttatccttta tatttcattt gtaaactata tttgaacgac attccccccg 1200  
 aaaaattgaa atgtaaagat gaggcagaga ataaagtgtt ctatgaaatt 1250  
 cagaacttta tttctgaatg taacatccct aataacaacc ttcattcttc 1300  
 taatacagca aaataaaaat ttaacaacca aggaatagta tttaagaaaa 1350  
 tgttgaaata atttttttaa aatagcatta cagactgag 1389

<210> SEQ ID NO 114  
 <211> LENGTH: 231

-continued

---

```

<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 114

Met Met Pro Lys His Cys Phe Leu Gly Phe Leu Ile Ser Phe Phe
 1          5          10          15
Leu Thr Gly Val Ala Gly Thr Gln Ser Thr His Glu Ser Leu Lys
 20          25          30
Pro Gln Arg Val Gln Phe Gln Ser Arg Asn Phe His Asn Ile Leu
 35          40          45
Gln Trp Gln Pro Gly Arg Ala Leu Thr Gly Asn Ser Ser Val Tyr
 50          55          60
Phe Val Gln Tyr Lys Ile Tyr Gly Gln Arg Gln Trp Lys Asn Lys
 65          70          75
Glu Asp Cys Trp Gly Thr Gln Glu Leu Ser Cys Asp Leu Thr Ser
 80          85          90
Glu Thr Ser Asp Ile Gln Glu Pro Tyr Tyr Gly Arg Val Arg Ala
 95          100         105
Ala Ser Ala Gly Ser Tyr Ser Glu Trp Ser Met Thr Pro Arg Phe
 110         115         120
Thr Pro Trp Trp Glu Thr Lys Ile Asp Pro Pro Val Met Asn Ile
 125         130         135
Thr Gln Val Asn Gly Ser Leu Leu Val Ile Leu His Ala Pro Asn
 140         145         150
Leu Pro Tyr Arg Tyr Gln Lys Glu Lys Asn Val Ser Ile Glu Asp
 155         160         165
Tyr Tyr Glu Leu Leu Tyr Arg Val Phe Ile Ile Asn Asn Ser Leu
 170         175         180
Glu Lys Glu Gln Lys Val Tyr Glu Gly Ala His Arg Ala Val Glu
 185         190         195
Ile Glu Ala Leu Thr Pro His Ser Ser Tyr Cys Val Val Ala Glu
 200         205         210
Ile Tyr Gln Pro Met Leu Asp Arg Arg Ser Gln Arg Ser Glu Glu
 215         220         225

Arg Cys Val Glu Ile Pro
 230

```

```

<210> SEQ ID NO 115
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

```

```

<400> SEQUENCE: 115

```

```

tgtaaacga cggccagta aatagacctg caattattaa tct

```

43

```

<210> SEQ ID NO 116
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

```

-continued

&lt;223&gt; OTHER INFORMATION: Synthetic Oligonucleotide Probe

&lt;400&gt; SEQUENCE: 116

caggaacag ctatgaccac ctgcacacct gcaaatccat t

41

What is claimed is:

1. Isolated nucleic acid having at least 80% nucleic acid sequence identity to a nucleotide sequence that encodes an amino acid sequence selected from the group consisting of the amino acid sequence shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) and **FIG. 114** (SEQ ID NO:114).

2. Isolated nucleic acid having at least 80% nucleic acid sequence identity to a nucleotide sequence selected from the group consisting of the nucleotide sequence shown in **FIG. 1** (SEQ ID NO:1), **FIG. 3** (SEQ ID NO:3), **FIG. 5** (SEQ ID NO:5), **FIG. 7** (SEQ ID NO:7), **FIG. 9** (SEQ ID NO:9), **FIG. 11** (SEQ ID NO:11), **FIG. 13** (SEQ ID NO:13), **FIG. 15** (SEQ ID NO:15), **FIG. 17** (SEQ ID NO:17), **FIG. 19** (SEQ ID NO:19), **FIG. 21** (SEQ ID NO:21), **FIG. 23** (SEQ ID NO:23), **FIG. 25** (SEQ ID NO:25), **FIG. 27** (SEQ ID NO:27), **FIG. 29** (SEQ ID NO:29), **FIG. 31** (SEQ ID NO:31), **FIG. 33** (SEQ ID NO:33), **FIG. 35** (SEQ ID NO:35), **FIG. 37** (SEQ ID NO:37), **FIG. 39** (SEQ ID NO:39), **FIG. 41** (SEQ ID NO:41), **FIG. 43** (SEQ ID NO:43), **FIG. 45** (SEQ ID NO:45), **FIG. 47** (SEQ ID NO:47), **FIG. 49** (SEQ ID NO:49), **FIG. 51** (SEQ ID NO:51), **FIG. 53** (SEQ ID NO:53), **FIG. 55** (SEQ ID NO:55), **FIG. 57** (SEQ ID NO:57), **FIG. 59** (SEQ ID NO:59), **FIG. 61** (SEQ ID NO:61), **FIG. 63** (SEQ ID NO:63), **FIG. 65** (SEQ ID NO:65), **FIG. 67** (SEQ ID NO:67), **FIG. 69** (SEQ ID NO:69), **FIG. 71** (SEQ ID NO:71), **FIG. 73** (SEQ ID NO:73), **FIG. 75** (SEQ ID NO:75), **FIG. 77** (SEQ ID NO:77), **FIG. 79** (SEQ ID NO:79), **FIG. 81** (SEQ ID NO:81), **FIG. 83** (SEQ ID NO:83), **FIG. 85** (SEQ ID NO:85), **FIG. 87** (SEQ ID NO:87), **FIG. 89** (SEQ ID NO:89), **FIG. 91** (SEQ ID NO:91), **FIG. 93** (SEQ ID NO:93), **FIGS. 95A-95B** (SEQ ID NO:95), **FIG. 97** (SEQ ID NO:97), **FIG. 99** (SEQ ID NO:99), **FIG. 101** (SEQ ID NO:101), **FIG. 103** (SEQ ID NO:103), **FIG. 105** (SEQ ID NO:105), **FIG. 107** (SEQ ID NO:107), **FIG. 109** (SEQ ID NO:109), **FIG. 111** (SEQ ID NO:111) and **FIG. 113** (SEQ ID NO:113).

**FIG. 81** (SEQ ID NO:81), **FIG. 83** (SEQ ID NO:83), **FIG. 85** (SEQ ID NO:85), **FIG. 87** (SEQ ID NO:87), **FIG. 89** (SEQ ID NO:89), **FIG. 91** (SEQ ID NO:91), **FIG. 93** (SEQ ID NO:93), **FIGS. 95A-95B** (SEQ ID NO:95), **FIG. 97** (SEQ ID NO:97), **FIG. 99** (SEQ ID NO:99), **FIG. 101** (SEQ ID NO:101), **FIG. 103** (SEQ ID NO:103), **FIG. 105** (SEQ ID NO:105), **FIG. 107** (SEQ ID NO:107), **FIG. 109** (SEQ ID NO:109), **FIG. 111** (SEQ ID NO:111) and **FIG. 113** (SEQ ID NO:113).

3. Isolated nucleic acid having at least 80% nucleic acid sequence identity to a nucleotide sequence selected from the group consisting of the full-length coding sequence of the nucleotide sequence shown in **FIG. 1** (SEQ ID NO:1), **FIG. 3** (SEQ ID NO:3), **FIG. 5** (SEQ ID NO:5), **FIG. 7** (SEQ ID NO:7), **FIG. 9** (SEQ ID NO:9), **FIG. 11** (SEQ ID NO:11), **FIG. 13** (SEQ ID NO:13), **FIG. 15** (SEQ ID NO:15), **FIG. 17** (SEQ ID NO:17), **FIG. 19** (SEQ ID NO:19), **FIG. 21** (SEQ ID NO:21), **FIG. 23** (SEQ ID NO:23), **FIG. 25** (SEQ ID NO:25), **FIG. 27** (SEQ ID NO:27), **FIG. 29** (SEQ ID NO:29), **FIG. 31** (SEQ ID NO:31), **FIG. 33** (SEQ ID NO:33), **FIG. 35** (SEQ ID NO:35), **FIG. 37** (SEQ ID NO:37), **FIG. 39** (SEQ ID NO:39), **FIG. 41** (SEQ ID NO:41), **FIG. 43** (SEQ ID NO:43), **FIG. 45** (SEQ ID NO:45), **FIG. 47** (SEQ ID NO:47), **FIG. 49** (SEQ ID NO:49), **FIG. 51** (SEQ ID NO:51), **FIG. 53** (SEQ ID NO:53), **FIG. 55** (SEQ ID NO:55), **FIG. 57** (SEQ ID NO:57), **FIG. 59** (SEQ ID NO:59), **FIG. 61** (SEQ ID NO:61), **FIG. 63** (SEQ ID NO:63), **FIG. 65** (SEQ ID NO:65), **FIG. 67** (SEQ ID NO:67), **FIG. 69** (SEQ ID NO:69), **FIG. 71** (SEQ ID NO:71), **FIG. 73** (SEQ ID NO:73), **FIG. 75** (SEQ ID NO:75), **FIG. 77** (SEQ ID NO:77), **FIG. 79** (SEQ ID NO:79), **FIG. 81** (SEQ ID NO:81), **FIG. 83** (SEQ ID NO:83), **FIG. 85** (SEQ ID NO:85), **FIG. 87** (SEQ ID NO:87), **FIG. 89** (SEQ ID NO:89), **FIG. 91** (SEQ ID NO:91), **FIG. 93** (SEQ ID NO:93), **FIGS. 95A-95B** (SEQ ID NO:95), **FIG. 97** (SEQ ID NO:97), **FIG. 99** (SEQ ID NO:99), **FIG. 101** (SEQ ID NO:101), **FIG. 103** (SEQ ID NO:103), **FIG. 105** (SEQ ID NO:105), **FIG. 107** (SEQ ID NO:107), **FIG. 109** (SEQ ID NO:109), **FIG. 111** (SEQ ID NO:111) and **FIG. 113** (SEQ ID NO:113).

4. Isolated nucleic acid having at least 80% nucleic acid sequence identity to the full-length coding sequence of the DNA deposited under any ATCC accession number shown in Table 7.

5. A vector comprising the nucleic acid of claim 1.

6. A host cell comprising the vector of claim 5.

7. The host cell of claim 6, wherein said cell is a CHO cell.

8. The host cell of claim 6, wherein said cell is an *E. coli*.

9. The host cell of claim 6, wherein said cell is a yeast cell.

10. A process for producing a PRO polypeptide comprising culturing the host cell of claim 6 under conditions suitable for expression of said PRO polypeptide and recovering said PRO polypeptide from the cell culture.

11. An isolated polypeptide having at least 80% amino acid sequence identity to an amino acid sequence selected from the group consisting of the amino acid sequence shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) and **FIG. 114** (SEQ ID NO:114).

12. An isolated polypeptide having at least 80% amino acid sequence identity to an amino acid sequence encoded by the full-length coding sequence of the DNA deposited under any ATCC accession number shown in Table 7.

13. A chimeric molecule comprising a polypeptide according to claim 11 fused to a heterologous amino acid sequence.

14. The chimeric molecule of claim 13, wherein said heterologous amino acid sequence is an epitope tag sequence.

15. The chimeric molecule of claim 13, wherein said heterologous amino acid sequence is a Fc region of an immunoglobulin.

16. An antibody which specifically binds to a polypeptide according to claim 11.

17. The antibody of claim 16, wherein said antibody is a monoclonal antibody, a humanized antibody or a single-chain antibody.

18. Isolated nucleic acid having at least 80% nucleic acid sequence identity to:

(a) a nucleotide sequence encoding the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG.**

**46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) or **FIG. 114** (SEQ ID NO:114), lacking its associated signal peptide;

(b) a nucleotide sequence encoding an extracellular domain of the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) or **FIG. 114** (SEQ ID NO:114), with its associated signal peptide; or

(c) a nucleotide sequence encoding an extracellular domain of the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID

NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) or **FIG. 114** (SEQ ID NO:114), lacking its associated signal peptide.

**19.** An isolated polypeptide having at least 80% amino acid sequence identity to:

- (a) an amino acid sequence of the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) or **FIG. 114** (SEQ ID NO:114), lacking its associated signal peptide;
- (b) an amino acid sequence of an extracellular domain of the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40**

(SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) or **FIG. 114** (SEQ ID NO:114), with its associated signal peptide; or

- (c) an amino acid sequence of an extracellular domain of the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) or **FIG. 114** (SEQ ID NO:114), lacking its associated signal peptide.

**20.** A method for stimulating the proliferation or differentiation of chondrocyte cells, said method comprising contacting said cells with a PRO6018 polypeptide, wherein the proliferation or differentiation of said cells is stimulated.

**21.** A method for stimulating the proliferation of human microvascular endothelial cells, said method comprising contacting said cells with a PRO1313, PRO20080 or PRO21383 polypeptide, wherein the proliferation of said cells is stimulated.

**24.** A method for inhibiting the proliferation of human microvascular endothelial cells, said method comprising

contacting said cells with a PRO6071, PRO4487 or PRO6006 polypeptide, wherein the proliferation of said cells is inhibited.

**25.** A method for detecting the presence of tumor in a mammal, said method comprising comparing the level of expression of any PRO polypeptide shown in Table 8 in (a) a test sample of cells taken from said mammal and (b) a control sample of normal cells of the same cell type, wherein a higher level of expression of said PRO polypeptide in the test sample as compared to the control sample is indicative of the presence of tumor in said mammal.

**26.** The method of claim **25**, wherein said tumor is lung tumor, colon tumor, breast tumor, prostate tumor, rectal tumor, kidney tumor or liver tumor.

**27.** A method for inducing endothelial cell tube formation comprising administering to the endothelial cell a PRO281, PRO1560, PRO189, PRO4499, PRO6308, PRO6000, PRO10275, PRO21207, PRO20933 or PRO34274 polypeptide, or agonist thereof, wherein tube formation in said endothelial cell is induced.

**28.** An oligonucleotide probe derived from any of the nucleotide sequences shown in the accompanying figures.

\* \* \* \* \*