



- (51) **International Patent Classification:**
A61K 35/17 (2015.01) C07K 14/705 (2006.01)
- (21) **International Application Number:**
PCT/US2017/025602
- (22) **International Filing Date:**
31 March 2017 (31.03.2017)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
62/316,489 31 March 2016 (31.03.2016) US
- (71) **Applicant:** UNIVERSITY OF SOUTHERN CALIFORNIA [US/US]; 1150 South Olive Street, Suite 2300, Los Angeles, California 90015 (US).
- (72) **Inventor:** CHAUDHARY, Preet, M.; 4606 Arcola Avenue, Toluca Lake, California 91602 (US).
- (74) **Agents:** VAKHARIA-RAO, Hema et al.; Nixon Peabody LLP, 300 South Grand Avenue, Suite 4100, Los Angeles, California 90071-3151 (US).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,


AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) **Title:** A HIGHLY SENSITIVE AND SPECIFIC LUCIFERASE BASED REPORTER ASSAY FOR ANTIGEN DETECTION

Fig. 11

(57) **Abstract:** Provided herein are methods for detecting an antigen or for detecting expression of a chimeric antigen receptor (CAR). The methods include obtaining a sample from a subject, contacting the sample with a fusion protein comprising a reporter fused to a single chain antibody specific to the antigen or fused to an extracellular domain of an antigen targeted by the CAR or fused to Protein L and assaying the activity of the reporter, wherein presence of reporter activity or increase in reporter activity relative to a reference value is indicative of presence of the antigen or presence of the expression of the chimeric antigen receptor in the sample.

A HIGHLY SENSITIVE AND SPECIFIC LUCIFERASE BASED REPORTER ASSAY FOR ANTIGEN DETECTION

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

[0001] This invention was made with government support under Grant No. DE019811 awarded by National Institutes of Health. The government has certain rights in the invention.

CROSS-REFERENCE TO RELATED APPLICATIONS

[0002] This application claims benefit of priority under 35 U.S.C. §119(e) of U.S. Provisional Patent Application No. 62/316,489 filed on March 31, 2016, the contents of which are incorporated by reference in their entirety.

TECHNICAL FIELD

[0003] Provided herein are assays to detect expression of antigens and cells expressing chimeric antigen receptors.

BACKGROUND

[0004] All publications herein are incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference. The following description includes information that may be useful in understanding the present invention. It is not an admission that any of the information provided herein is prior art or relevant to the presently claimed invention, or that any publication specifically or implicitly referenced is prior art.

[0005] There are several methods currently in use for detection of a particular antigen, including flow cytometry, ELISA and western blotting. However, these methods require expensive equipment (such as flow cytometry), suffer from poor sensitivity and specificity (e.g., ELISA and Western blotting) or are time consuming. The poor sensitivity of the current methods of antigen detection is particularly significant in light of development of new targeted therapies using antibody drug conjugates and chimeric antigen receptors. These new therapies are extremely potent at killing cells expressing their target antigen even at very low levels, which are below the detection limits of current methods. For example, Brentuximab

vedotin (Adcetris), an antibody drug conjugate, that targets CD30 antigen has shown impressive clinical activity in diffuse large cell lymphoma in several clinical trials. However, the response to Brentuximab vedotin does not always correlate with CD30 positivity, as determined by conventional immunohistochemistry, and patients with diffuse large cell lymphomas that are CD30–ve frequently respond to this agent. This has led to the hypothesis that Brentuximab vedotin requires very low level expression of CD30 for its activity and its efficacy in CD30-ve diffuse large cell lymphoma is, in part, due to low level expression of CD30 in these samples, which is below the detection of conventional methods.

[0006] Chimeric antigen receptor based T cell therapy represents another example where detection of low level expression of an antigen in a patient sample is critically important to predict response to therapy. It is believed that CAR-T cells can kill a target cell expressing as few as 100 molecules of their target antigen. Consistent with this, CAR-T cells targeting CD19 antigen have shown impressive clinical activity in multiple myeloma, a disease which is believed to be CD19-ve based on conventional detection methods (Hajek, R *et al.*, Br J Haematol 163:551-564,2013). Thus, availability of highly sensitive and specific method for detection of CD30, CD19 and other antigens would be of great benefit in determining the patients who are likely to respond to targeted therapies as well as to predict toxicity to normal tissues. In addition to clinical use, a highly sensitive and specific method for antigen detection, which is inexpensive, quick and does not require special equipment, would also have utility for research applications.

[0007] Another challenge in the field of CAR-T cell therapy is the lack of a fast, economical, sensitive and robust assay for detection of chimeric antigen receptors on the surface of T cells. Current methods for detection of CAR on cell surface have several limitations. For example, CD19-specific CARs have been detected following staining with an Alexa Flour 488-conjugated CD19-Fc fusion protein consisting of human CD19 extracellular domain and Fc region of human IgG1 (De Oliveira, SN *et al.*, Journal of translational medicine 11:23,2013). However, this protocol required the extra steps and cost of purification of the fusion protein followed by its conjugation with Alexa Flour 488. CAR-expressing T cells have been also detected using anti-human IgG1 Fc or biotinylated Protein L. However, these approaches suffer from significant background and limited discrimination (De Oliveira, SN *et al.*, Journal of translational medicine 11:23,2013). Staining using biotinylated Protein L

necessitates secondary staining with labeled streptavidin, with additional protocol steps and potential loss of cells (De Oliveira, SN *et al.*, Journal of translational medicine 11:23,2013).

[0008] In various embodiments, the instant invention provides a fast, economical, sensitive and specific assay for detection of chimeric antigen receptors on the surface of T cells that does not require any expensive equipment. Due to its extreme sensitivity and broad dynamic range, this assay can be used as a quality-control tool during the manufacturing of CAR-T cells and to monitor the persistence of CAR-T cells in the body after infusion.

[0009] There is a need in the art for sensitive methods for detecting antigens that are expressed at low levels and selecting patients for therapy based on detection of antigens of interest. There is also need in the art for sensitive, specific, fast and economical method for detecting chimeric antigen receptor-modified immune cells.

SUMMARY

[0010] The following embodiments and aspects thereof are described and illustrated in conjunction with systems, compositions and methods which are meant to be exemplary and illustrative, not limiting in scope.

[0011] Provided herein is a method for detecting expression of a chimeric antigen receptor (CAR), comprising: obtaining a sample from a subject in need determination of expression of the CAR; contacting the sample with a fusion protein comprising a reporter fused to an extracellular domain of an antigen targeted by the CAR; and assaying the activity of the reporter; wherein presence of reporter activity or increase in reporter activity relative to a reference value is indicative of the expression of the CAR in the sample.

[0012] In one embodiment, the reporter is a non-secretory form of a luciferase. In exemplary embodiments, the non-secretory form of luciferase is obtained from copepods, deep sea shrimp or homologs or orthologs thereof or mutants or derivatives thereof.

[0013] In some embodiments, the copepods are selected from the group consisting of any one or more of *Gaussia princeps*, *Pleuromamma abdominalis*, *Metridia pacifica*, *Metridia curticauda*, *Metridia asymmetrica*, *Metridia okhotensis*, *Metridia longa*, *Lucicutia ovaliformis*, *Heterorhabdus tanneri*, and *Pleuromamma scutellata*.

[0014] In some embodiments, the luciferase is any one or more of GLuc, NanoLuc (NLuc), MLuc7, HtLuc, LoLuc, PaLuc1, PaLuc2, MpLuc1, McLuc1, MaLuc1, MoLuc1, MoLuc2, MLuc39, PsLuc1, LocLuc1-3, HtLuc2 Renilla, TurboLuc16 (TLuc) or homologs or orthologs thereof or mutants or functional derivatives thereof.

[0015] In some embodiments, the reporter activity is assayed by exposing the target cells to a luciferase specific substrate. In one embodiment, the luciferase-specific substrate is coelentrastazine or a derivative thereof. In another embodiment, the luciferase-specific substrate is imidazopyrazinone substrate (furimazine) or a derivative thereof.

[0016] In some embodiments, the CAR is expressed on an immune cell. In one embodiment, the immune cell is a T cell. In another embodiment, the immune cell is a CD4 T cell. In a further embodiment, the immune cell is a CD8 T cell. In an embodiment, the immune cell is a Treg cell. In some embodiments, the immune cell is a naive T cell. In some embodiments, the immune cell is a memory T cell. In some embodiments, the immune cell is central memory T cell. In an embodiment, the immune cell is an effector memory T cell. In an embodiment, immune cell is an NK cell.

[0017] In some embodiments, the CAR is expressed on a stem cell. In an embodiment, the CAR is expressed on a hematopoietic stem cell. In an embodiment, the CAR is expressed on an induced pluripotent stem cell.

[0018] In some embodiments, the target antigen of the CAR is any one or more of but are not limited to CD19, CD20, CS1/SLAMF7, BCMA, CD123, CD33, MPLLym1, Lym2, CD200R, 5T4, adenocarcinoma antigen, alpha-fetoprotein, BAFF, B-lymphoma cell, C242 antigen, CA-125, carbonic anhydrase 9 (CA-IX), C-MET, CCR4, CD152, , CD200, CD22, CD221, CD23 (IgE receptor), CD28, CD30 (TNFRSF8), , CD4, CD40, CD44 v6, CD51, CD52, CD56, CD74, CD80, CEA, CNTO888, CTLA-4, DR5, EGFR, EpCAM, CD3, FAP, fibronectin extra domain-B, folate receptor 1, GD2, GD3 ganglioside, glycoprotein 75, GPNMB, HER2/neu, HGF, human scatter factor receptor kinase, IGF-1 receptor, IGF-I, IgG1, L1-CAM, IL-13, IL-6, insulin-like growth factor I receptor, integrin $\alpha 5\beta 1$, integrin $\alpha v\beta 3$ FLT3, MORAb-009, MS4A1, MUC1, mucin CanAg, N-glycolylneuraminic acid, NPC-1C, PDGF-R α , PDL192, phosphatidylserine, prostatic carcinoma cells, RANKL, RON, ROR1, SCH 900105, SDC1, SLAMF7, (MPL), TAG-72, tenascin C, TGF beta 2, TGF- β ,

TRAIL-R1, TRAIL-R2, tumor antigen CTAA16.88, VEGF-A, VEGFR-1, VEGFR2, or vimentin.

[0019] In an embodiment, the target antigen of CAR is a complex of a HLA molecule with a peptide antigen. In an embodiment, the HLA molecule is HLA-A2.

[0020] In an embodiment, the subject has cancer or an immune disorder. In an embodiment, the subject is being assessed for immune therapy. In an embodiment, the immune therapy comprises CAR T cells

[0021] In some embodiments, the fusion protein further comprises a tag. In exemplary embodiments, the tag is any one or more of chitin binding protein (CBP), glutathione-S-transferase (GST), polyhistidine (His) tag, FLAG tag, HA tag, Myc tag, V5 tag, AcV5 tag, Streptag or a combination thereof.

[0022] In some embodiments, the reference value is the reporter activity in any one or more of (i) cells that do not express the CAR; (ii) cells that express the CAR but are treated with fusion protein which is not targeted by the CAR; (iii) cells that are not treated with the substrate for the reporter; or (iv) combinations thereof.

[0023] In some embodiments, the reporter is fused to the extracellular domain of an antigen targeted by the chimeric antigen receptor through a covalent bond.

[0024] In some embodiments, the reporter is fused to the extracellular domain of an antigen targeted by the chimeric antigen receptor through a non-covalent bond.

[0025] In some embodiments, the reporter is fused to the extracellular domain of an antigen targeted by the chimeric antigen receptor through an intermediate molecule.

[0026] In some embodiments, a single molecule of the reporter is fused to the extracellular domain of an antigen targeted by the chimeric antigen receptor.

[0027] In some embodiments, more than one molecules of the reporter are fused to the extracellular domain of an antigen targeted by the chimeric antigen receptor.

[0028] In one embodiment, the assay is performed in vitro. In another embodiment, the assay is performed in vivo.

[0029] Also provided herein are kits for practicing the invention.

BRIEF DESCRIPTION OF FIGURES

[0030] Exemplary embodiments are illustrated in referenced figures. It is intended that the embodiments and figures disclosed herein are to be considered illustrative rather than restrictive.

[0031] **FIG. 1A-FIG. 1B** depict, in accordance with various embodiments of the invention, that strong binding with 161-GGSG-NLuc-AcV5 was observed on HEL.92.1.7-Gluc-vector cells.

[0032] **FIG. 2** depicts, in accordance with various embodiments of the invention, increased binding of 161-GGSG-NLuc-AcV5 supernatant to cells that had been transfected with MPL as compared to the untransfected cells.

[0033] **FIG. 3** depicts, in accordance with various embodiments of the invention, NLuc assay to measure expression of CAR in T cells. The uninfected T cells (UI) and T cells expressing CD19 (FMC63) and control (4C3) CAR were incubated with CD19-GGSG-NLuc-AcV5 supernatants as described in material and methods section followed by washing with PBS and measurement of NLuc activity by Coeleoentraine (CTZ; Nanolight) diluted in PBS. Luminescence was quantified using a BioTek plate reader. Data represents mean values of triplicate wells +/- standard deviation (SD).

[0034] **FIG. 4A-FIG. 4B** depict, in accordance with various embodiments of the invention, NLuc assay to measure expression of CAR in 293FT cells. The untransfected 293FT cells, and those transfected with CD19 (FMC63-BBZ-PAC) and 161-BBZ-PAC CAR were incubated with CD19-GGSG-NLuc-AcV5 and MPL-GGSG-NLuc-AcV5 supernatants as described in material and methods section followed by washing with PBS and measurement of NLuc activity by Coeleoentraine (CTZ; Nanolight) diluted in PBS. Luminescence was quantified using a BioTek plate reader. Data represents mean values of triplicate wells +/- standard deviation (SD).

[0035] **FIG. 5** depicts, in accordance with various embodiments of the invention, modest binding of MPL-GGSG-NLuc-AcV5 to 293FT cells transfected with hTPO-CD28z-Pac and mTPO-CD28z-Pac constructs and strong binding to 293FT cells transfected with 161-BBz-

Pac construct as compared to untransfected cells or cells that had been transfected with control CAR construct.

[0036] **FIG. 6A-FIG. 6C** depicts, in accordance with various embodiments of the invention, LUC activity of FMC63-GGSG-MLuc7-MM-LL-HA, FMC63-GGS-TurboLuc16-x3Flag, FMC63-GGS-GLuc-x3FLAG, and FMC63-GGS-NLuc-AcV5 on RAJI cells (known to be CD19 positive cell line) as compared to the Luc activity observed with media control.

[0037] **FIG. 7A-FIG. 7B** depicts, in accordance with various embodiments of the invention, fluorescence activity of RAJI cells stained with FMC63-GGS-GLuc-x3FLAG and FMC63-GGS-TurboLuc16-x3Flag.

[0038] **FIG. 8A-FIG. 8B** depicts, in accordance with various embodiments of the invention, relative LUC activity of different fusion proteins on the 3 cell lines.

[0039] **FIG. 9A-FIG. 9B** depicts, in accordance with various embodiments of the invention, show binding of CD30-Ac10(vL-vH)-GGSG-NLuc-x3-Flag to Jurkat and CHTR74 cells compared to Kasumi cells.

[0040] **FIG. 10A-FIG. 10B** depicts, in accordance with various embodiments of the invention, binding of both TCRB1-Jov1(vL-vH)-GGSG-NLuc-x3-Flag and TCRB1-E09(vL-vH)-GGSG-NLuc-x3-Flag fusion proteins to Jurkat and as compared to Kasumi and CHTR74 cells.

[0041] **FIG. 11** depicts, in accordance with various embodiments of the invention, the sensitivity of the FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170] fusion protein to detect 100 NK92MI cells expressing the CD19-specific CAR FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208] in the background of 1 million parental NK92MI cells.

[0042] **FIG. 12** depicts, in accordance with various embodiments of the invention, TurboLuc activity in various fractions of the protocol for purification of FLAG-CD19-ECD-GGS-TurboLuc16-4xFlag-2xStreptag-8xHis-T2A-Pac(111616-J04)[SEQ ID NO:1172] fusion protein from the supernatant of 293FT cells using Strep-Tactin resin.

[0043] FIG. 13 depicts, in accordance with various embodiments of the invention, binding of two Protein L-NLuc fusion proteins to Jurkat cells expressing the CAR HuLuc64-(vL-vH)-Myc-BBz-T2A-Pac(052616-U05)[SEQ ID NO:1228], indicating that Protein L based Luc fusion proteins can be used to detect the expression of CARs containing a kappa light chain,

DETAILED DESCRIPTION

[0044] All references cited herein are incorporated by reference in their entirety as though fully set forth. Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Allen et al., Remington: The Science and Practice of Pharmacy 22nd ed., Pharmaceutical Press (September 15, 2012); Hornyak et al., Introduction to Nanoscience and Nanotechnology, CRC Press (2008); Singleton and Sainsbury, Dictionary of Microbiology and Molecular Biology 3rd ed., revised ed., J. Wiley & Sons (New York, NY 2006); Smith, March's Advanced Organic Chemistry Reactions, Mechanisms and Structure 7th ed., J. Wiley & Sons (New York, NY 2013); Singleton, Dictionary of DNA and Genome Technology 3rd ed., Wiley-Blackwell (November 28, 2012); and Green and Sambrook, Molecular Cloning: A Laboratory Manual 4th ed., Cold Spring Harbor Laboratory Press (Cold Spring Harbor, NY 2012), provide one skilled in the art with a general guide to many of the terms used in the present application. For references on how to prepare antibodies, see Greenfield, Antibodies A Laboratory Manual 2nd ed., Cold Spring Harbor Press (Cold Spring Harbor NY, 2013); Köhler and Milstein, Derivation of specific antibody-producing tissue culture and tumor lines by cell fusion, Eur. J. Immunol. 1976 Jul, 6(7):511-9; Queen and Selick, Humanized immunoglobulins, U. S. Patent No. 5,585,089 (1996 Dec); and Riechmann et al., Reshaping human antibodies for therapy, Nature 1988 Mar 24, 332(6162):323-7.

[0045] One skilled in the art will recognize many methods and materials similar or equivalent to those described herein, which could be used in the practice of the present invention. Other features and advantages of the invention will become apparent from the following detailed description, taken in conjunction with the accompanying drawings, which illustrate, by way of example, various features of embodiments of the invention. Indeed, the present invention

is in no way limited to the methods and materials described. For convenience, certain terms employed herein, in the specification, examples and appended claims are collected here.

[0046] Unless stated otherwise, or implicit from context, the following terms and phrases include the meanings provided below. Unless explicitly stated otherwise, or apparent from context, the terms and phrases below do not exclude the meaning that the term or phrase has acquired in the art to which it pertains. The definitions are provided to aid in describing particular embodiments, and are not intended to limit the claimed invention, because the scope of the invention is limited only by the claims. Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.

[0047] As used herein the term “comprising” or “comprises” is used in reference to compositions, methods, and respective component(s) thereof, that are useful to an embodiment, yet open to the inclusion of unspecified elements, whether useful or not. It will be understood by those within the art that, in general, terms used herein are generally intended as “open” terms (e.g., the term “including” should be interpreted as “including but not limited to,” the term “having” should be interpreted as “having at least,” the term “includes” should be interpreted as “includes but is not limited to,” etc.).

[0048] Unless stated otherwise, the terms “a” and “an” and “the” and similar references used in the context of describing a particular embodiment of the application (especially in the context of claims) can be construed to cover both the singular and the plural. The recitation of ranges of values herein is merely intended to serve as a shorthand method of referring individually to each separate value falling within the range. Unless otherwise indicated herein, each individual value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (for example, “such as”) provided with respect to certain embodiments herein is intended merely to better illuminate the application and does not pose a limitation on the scope of the application otherwise claimed. The abbreviation, “e.g.” is derived from the Latin *exempli gratia*, and is used herein to indicate a non-limiting example. Thus, the abbreviation “e.g.” is synonymous with the term “for example.” No

language in the specification should be construed as indicating any non-claimed element essential to the practice of the application.

[0049] As used herein, the term "about" refers to a measurable value such as an amount, a time duration, and the like, and encompasses variations of $\pm 20\%$, $\pm 10\%$, $\pm 5\%$, $\pm 1\%$, $\pm 0.5\%$ or $\pm 0.1\%$ from the specified value.

[0050] As used herein, "amino acid" is meant to include both natural and synthetic amino acids, and both D and L amino acids. "Standard amino acid" means any of the twenty L-amino acids commonly found in naturally occurring peptides. "Nonstandard amino acid" means any amino acid, other than the standard amino acids, regardless of whether it is prepared synthetically or derived from a natural source.

[0051] As used herein, "synthetic amino acid" also encompasses chemically modified amino acids, including but not limited to salts, amino acid derivatives (such as amides), and substitutions. Amino acids contained within the peptides disclosed herein, and particularly at the carboxy- or amino-terminus, can be modified by methylation, amidation, acetylation or substitution with other chemical groups which can change the peptide's circulating half-life without adversely affecting their biological activity. Additionally, a disulfide linkage may be present or absent in the peptides disclosed herein.

[0052] Herein, "peptide" and "protein" are used interchangeably, and refer to a compound comprised of at least two amino acid residues covalently linked by peptide bonds or modified peptide bonds (*e.g.*, peptide isosteres). No limitation is placed on the maximum number of amino acids which may comprise a protein or peptide. The amino acids comprising the peptides or proteins described herein and in the appended claims are understood to be either D or L amino acids with L amino acids being preferred. The amino acid comprising the peptides or proteins described herein may also be modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Modifications can occur anywhere in a peptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It is understood that the same type of modification may be present in the same or varying degrees at several sites in a given peptide. Also, a given peptide may contain many types of modifications. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent

attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. *See*, for instance, *Proteins--Structure and Molecular Properties*, 2nd Ed., T. E. Creighton, W.H. Freeman and Company, New York, 1993 and Wold F, *Posttranslational Protein Modifications: Perspectives and Prospects*, pgs. 1-12 in *Posttranslational Covalent Modification of Proteins*, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and nonprotein cofactors," *Meth. Enzymol.* (1990) 182: 626-646 and Rattan *et al.* (1992), "Protein Synthesis: Posttranslational Modifications and Aging," *Ann NY Acad Sci* 663: 48-62.

[0053] "Polynucleotide" as used herein includes but is not limited to DNA, RNA, cDNA (complementary DNA), mRNA (messenger RNA), rRNA (ribosomal RNA), shRNA (small hairpin RNA), snRNA (small nuclear RNA), snoRNA (short nucleolar RNA), miRNA (microRNA), genomic DNA, synthetic DNA, synthetic RNA, and/or tRNA.

[0054] As used herein, the term "polypeptide" encompasses amino acid chains of any length, including full length proteins (i.e., antigens), wherein the amino acid residues are linked by covalent peptide bonds.

[0055] It is to be inferred without explicit recitation and unless otherwise intended, that when the present disclosure relates to a polypeptide, protein, polynucleotide, antibody or fragment thereof, an equivalent or a biologically equivalent of such is intended within the scope of this disclosure. As used herein, the term "biological equivalent thereof" is intended to be synonymous with "equivalent thereof" when referring to a reference protein, antibody or fragment thereof, polypeptide or nucleic acid, intends those having minimal homology while still maintaining desired structure or functionality. Unless specifically recited herein, it is contemplated that any of the above also includes equivalents thereof. For example, an

equivalent intends at least about 70% homology or identity, or at least 80% homology or identity and alternatively, or at least about 85%, or alternatively at least about 90%, or alternatively at least about 95%, or alternatively at least 98% percent homology or identity and exhibits substantially equivalent biological activity to the reference protein, polypeptide, antibody or fragment thereof or nucleic acid. Alternatively, when referring to polynucleotides, an equivalent thereof is a polynucleotide that hybridizes under stringent conditions to the reference polynucleotide or its complement. Alternatively, when referring to polypeptides or proteins, an equivalent thereof is a expressed polypeptide or protein from a polynucleotide that hybridizes under stringent conditions to the polynucleotide or its complement that encodes the reference polypeptide or protein.

[0056] A polypeptide "variant," as used herein, is a polypeptide that differs from the recited polypeptide only in conservative substitutions and/or modifications, such that the therapeutic, antigenic and/or immunogenic properties of the polypeptide are retained. Polypeptide variants preferably exhibit at least about 70%, more preferably at least about 90% and most preferably at least about 95% homology to the identified polypeptides. For polypeptides with immunoreactive properties, variants can, alternatively, be identified by modifying the amino acid sequence of one of the above polypeptides, and evaluating the immunoreactivity of the modified polypeptide. Such modified sequences can be prepared and tested using, for example, the representative procedures described herein.

[0057] As used herein, a "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. In general, the following groups of amino acids represent conservative changes: (1) ala, pro, gly, asp, gln, asn, ser, thr; (2) cys, ser, pyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his.

[0058] Variants can also, or alternatively, contain other modifications, including the deletion or addition of amino acids that have minimal influence on the antigenic properties, secondary structure and hydrophobic nature of the polypeptide. For example, a polypeptide can be conjugated to a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the protein. The polypeptide can also

be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (e.g., poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide can be conjugated to an immunoglobulin Fc region.

[0059] A polynucleotide or polynucleotide region (or a polypeptide or polypeptide region) having a certain percentage (for example, 80%, 85%, 90%, or 95%) of “sequence identity” to another sequence means that, when aligned, that percentage of bases (or amino acids) are the same in comparing the two sequences. The alignment and the percent homology or sequence identity can be determined using software programs known in the art, for example those described in Current Protocols in Molecular Biology (Ausubel et al., eds. 1987) Supplement 30, section 7.7.18, Table 7.7.1. Preferably, default parameters are used for alignment. A preferred alignment program is BLAST, using default parameters. In particular, preferred programs are BLASTN and BLASTP, using the following default parameters: Genetic code=standard; filter=none; strand=both; cutoff=60; expect=10; Matrix=BLOSUM62; Descriptions=50 sequences; sort by=HIGH SCORE; Databases=non-redundant, GenBank+EMBL+DDBJ+PDB+GenBank CDS translations+SwissProtein+SPupdate+PIR. Details of these programs can be found at the following Internet address: ncbi.nlm.nih.gov/cgi-bin/BLAST.

[0060] The term “isolated” as used herein refers to molecules or biologicals or cellular materials being substantially free from other materials. In one aspect, the term “isolated” refers to nucleic acid, such as DNA or RNA, or protein or polypeptide (e.g., an antibody or derivative thereof), or cell or cellular organelle, or tissue or organ, separated from other DNAs or RNAs, or proteins or polypeptides, or cells or cellular organelles, or tissues or organs, respectively, that are present in the natural source. The term “isolated” also refers to a nucleic acid or peptide that is substantially free of cellular material, viral material, or culture medium when produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized. Moreover, an “isolated nucleic acid” is meant to include nucleic acid fragments which are not naturally occurring as fragments and would not be found in the natural state. The term “isolated” is also used herein to refer to polypeptides which are isolated from other cellular proteins and is meant to encompass both purified and recombinant polypeptides. The term “isolated” is also used herein to refer to cells or tissues

that are isolated from other cells or tissues and is meant to encompass both, cultured and engineered cells or tissues.

[0061] “Sample” or “biological sample” as used herein refers to tissues or body fluids removed from a mammal, preferably human, and which contain CD8⁺ T cells. Samples can be blood and/or blood fractions, including peripheral blood sample like peripheral blood mononuclear cell (PBMC) sample or blood, bone marrow cell sample. A sample can also include any specific tissues/organ sample of interest, including, without limitation, lymphoid, thymus, pancreas, eye, heart, liver, nerves, intestine, skin, muscle, cartilage, ligament, synovial fluid, and/or joints. The samples can be taken from any individual including a healthy individual or an individual having cells, tissues, and/or an organ afflicted with the unwanted immune response. For example, a sample can be taken from an individual having an allergy, a graft vs. host disease, a cell or organ transplant, or an autoimmune disease or disorder. Methods for obtaining such samples are well known to a person of ordinary skill in the art of immunology and medicine. They include drawing and processing blood and blood components using routine procedures, or obtaining biopsies from the bone marrow or other tissue or organ using standard medical techniques.

[0062] “Chimeric antigen receptor” or “CAR” or “CARs” as used herein refers to engineered receptors, which graft an antigen specificity onto cells (for example T cells such as naïve T cells, central memory T cells, effector memory T cells or combination thereof). CARs are also known as artificial T-cell receptors, chimeric T-cell receptors or chimeric immunoreceptors. In various embodiments, CARs are recombinant polypeptides comprising an antigen-specific domain (ASD), a hinge region (HR), a transmembrane domain (TMD), co-stimulatory domain (CSD) and an intracellular signaling domain (ISD).

[0063] “Endogenous T cell receptor” or “TCR” as used herein refers to a T cell receptors that is expressed in T cells endogenously.

[0064] “Exogenous T cell receptor” or “transgenic T cell receptor” or “transgenic TCR” as used herein refers to T cell receptors that are expressed in T cells from outside.

[0065] As used herein, the term “antibody” refers to an intact immunoglobulin or to a monoclonal or polyclonal antigen-binding fragment with the Fc (crystallizable fragment)

region or FcRn binding fragment of the Fc region, referred to herein as the "Fc fragment" or "Fc domain". Antigen-binding fragments may be produced by recombinant DNA techniques or by enzymatic or chemical cleavage of intact antibodies. Antigen-binding fragments include, inter alia, Fab, Fab', F(ab')₂, Fv, dAb, and complementarity determining region (CDR) fragments, single-chain antibodies (scFv), single domain antibodies, chimeric antibodies, diabodies and polypeptides that contain at least a portion of an immunoglobulin that is sufficient to confer specific antigen binding to the polypeptide. The Fc domain includes portions of two heavy chains contributing to two or three classes of the antibody. The Fc domain may be produced by recombinant DNA techniques or by enzymatic (e.g. papain cleavage) or via chemical cleavage of intact antibodies.

[0066] The term "antibody fragment," as used herein, refer to a protein fragment that comprises only a portion of an intact antibody, generally including an antigen binding site of the intact antibody and thus retaining the ability to bind antigen. Examples of antibody fragments encompassed by the present definition include: (i) the Fab fragment, having VL (or vL), CL, VH (or vH) and CH1 domains; (ii) the Fab' fragment, which is a Fab fragment having one or more cysteine residues at the C-terminus of the CH1 domain; (iii) the Fd fragment having VH and CH1 domains; (iv) the Fd' fragment having VH and CH1 domains and one or more cysteine residues at the C-terminus of the CH1 domain; (v) the Fv fragment having the VL and VH domains of a single arm of an antibody; (vi) the dAb fragment (Ward et al., Nature 341, 544-546 (1989)) which consists of a VH domain; (vii) isolated CDR regions; (viii) F(ab')₂ fragments, a bivalent fragment including two Fab' fragments linked by a disulphide bridge at the hinge region; (ix) single chain antibody molecules (e.g., single chain Fv; scFv) (Bird et al., Science 242:423-426 (1988); and Huston et al., PNAS (USA) 85:5879-5883 (1988)); (x) "diabodies" with two antigen binding sites, comprising a heavy chain variable domain (VH) connected to a light chain variable domain (VL) in the same polypeptide chain (see, e.g., EP 404,097; WO 93/11161; and Hollinger et al., Proc. Natl. Acad. Sci. USA, 90:6444-6448 (1993)); (xi) "linear antibodies" comprising a pair of tandem Fd segments (VH-CH1-VH-CH1) which, together with complementary light chain polypeptides, form a pair of antigen binding regions (Zapata et al. Protein Eng. 8(10):1057-1062 (1995); and U.S. Pat. No. 5,641,870).

[0067] “Single chain variable fragment”, “single-chain antibody variable fragments” or “scFv” antibodies as used herein refer to forms of antibodies comprising the variable regions of only the heavy (V_H) and light (V_L) chains, connected by a linker peptide. The scFvs are capable of being expressed as a single chain polypeptide. The scFvs retain the specificity of the intact antibody from which it is derived. The light and heavy chains may be in any order, for example, V_H -linker- V_L or V_L -linker- V_H , so long as the specificity of the scFv to the target antigen is retained.

[0068] “Complementarity determining region” (CDR) as used herein refers to the amino acid sequences within the variable regions of antibodies which regions confer specificity and binding affinity. In general, there are three CDRs in each of the light chain variable regions (LCDR1, LCDR2 and LCDR3) and three CDRs in each of the heavy chain variable regions (HCD1, HCDr2 and HCDR3). The boundaries of the CDRs may be determined using methods well known in the art including the “Kabat” numbering scheme and/or “Chothia” number scheme (Kabat et al. Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Services, National Institutes of Health, Bethesda, MD; Al-Lazikani *et al.*, (1997) JMB 273,927-948).

[0069] As used herein, the term “specific binding” means the contact between an antibody and an antigen with a binding affinity of at least 10^{-6} M. In certain aspects, antibodies bind with affinities of at least about 10^{-7} M, and preferably 10^{-8} M, 10^{-9} M, 10^{-10} M, 10^{-11} M, or 10^{-12} M.

[0070] “Cancer” and “cancerous” refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. The term “cancer” is meant to include all types of cancerous growths or oncogenic processes, metastatic tissues or malignantly transformed cells, tissues, or organs, irrespective of histopathologic type or stage of invasiveness. Examples of solid tumors include malignancies, e.g., sarcomas, adenocarcinomas, and carcinomas, of the various organ systems, such as those affecting liver, lung, breast, lymphoid, gastrointestinal (e.g., colon), genitourinary tract (e.g., renal, urothelial cells), prostate and pharynx. Adenocarcinomas include malignancies such as most colon cancers, rectal cancer, renal-cell carcinoma, liver cancer, non-small cell carcinoma of the lung, cancer of the small intestine and cancer of the esophagus. In one embodiment, the

cancer is a melanoma, e.g., an advanced stage melanoma. Metastatic lesions of the aforementioned cancers can also be treated or prevented using the methods and compositions of the invention. Examples of other cancers that can be treated include bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular malignant melanoma, uterine cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, testicular cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, Hodgkin Disease, non-Hodgkin lymphoma, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, chronic or acute leukemias including acute myeloid leukemia, chronic myeloid leukemia, acute lymphoblastic leukemia, chronic lymphocytic leukemia, solid tumors of childhood, lymphocytic lymphoma, cancer of the bladder, cancer of the kidney or ureter, carcinoma of the renal pelvis, neoplasm of the central nervous system (CNS), primary CNS lymphoma, tumor angiogenesis, spinal axis tumor, brain stem glioma, pituitary adenoma, Kaposi's sarcoma, epidermoid cancer, squamous cell cancer, T-cell lymphoma, environmentally induced cancers including those induced by asbestos, and combinations of said cancers. Treatment of metastatic cancers, e.g., metastatic cancers that express PD-L1 (Iwai et al. (2005) Int. Immunol. 17:133-144) can be effected using the antibody molecules described herein.

[0071] "Immunogenic," as used herein, refers to the ability to elicit an immune response (e.g., cellular) in a patient, such as a human, and/or in a biological sample.

[0072] "Luciferase" or "Luciferases" as used herein refer class of oxidative enzymes that produce bioluminescence. The non-secretory form of luciferases for use in the compositions and methods described herein lack the N-terminal secretory sequence but produce bioluminescence. In exemplary embodiments, the luciferase is obtained from copepods, deep sea shrimp or homologs or orthologs thereof or mutants or derivatives thereof. In some embodiments, the copepods are any one or more of *Gaussia princeps*, *Pleuromamma abdominalis*, *Metridia pacifica*, *Metridia curticauda*, *Metridia asymmetrica*, *Metridia okhotensis*, *Metridia longa*, *Lucicutia ovaliformis*, *Heterorhabdus tanneri*, and *Pleuromamma scutellata*. In exemplary embodiments, the luciferases are any one or more of GLuc,

NanoLuc (NLuc), MLuc7, HtLuc, LoLuc, PaLuc1, PaLuc2, MpLuc1, McLuc1, MaLuc1, MoLuc1, MoLuc2, MLuc39, PsLuc1, LocLuc1-3, HtLuc2 Renilla, TurboLuc16 (TLuc) or homologs or orthologs thereof or mutants or functional derivatives thereof. In various embodiments, mutants or functional derivatives of the luciferases retain at least 100%, 90%, 80%, 70%, 60% or 50% of the bioluminescence activity from which the mutant or function derivative is derived. As described in Takenaka et al (Evolution of bioluminescence in marine planktonic copepods. 2012 *Mol Biol Evol* 29(6): 1669-1681), the copepod luciferases comprise two domains and each domain includes conserved sequences across the various luciferases. In some embodiments, any luciferase comprising the consensus sequence C-x(3)-C-L-x(2)-L-x(4)-C-x(8)-P-x-R-C (SEQ ID NO: 2437) in each of domains 1 and 2 as described by Takenaka et al., may be used in the compositions and methods described herein. In some embodiments, any luciferase comprising the consensus sequence K-x(7)-E-M-E-A-N-A-x(3)-G-C-x-R-x-C-L-I-x-L-S-x-I-K-C-T-x-K-M-x(4)-P-G-R-C-H-X-Y-x(8)-G (SEQ ID NO: 2438) in domain 1 as described by Takenaka et al. and consensus sequences I-x-G-x(6)-M-x-Q-F-x(2)-Q-V-x(2)-C-x(2)-C-x(3)-C-L-K-G-L-A-N-x(2)-C-S-x(2)-L-x(3)-L-P-x-R-C-x(2)-F-x(3)-I-x(8)-G (SEQ ID NO: 2439) in domain 2 as described by Takenaka et al., may be used in the compositions and methods described herein. In various embodiments, the functional fragments or mutants of luciferases for use with the compositions or methods described herein comprise the consensus sequence C-x(3)-C-L-x(2)-L-x(4)-C-x(8)-P-x-R-C (SEQ ID NO: 2437) in domain 1 as described by Takenaka et al. In various embodiments, the functional fragments or mutants of luciferases for use with the compositions or methods described herein comprise the consensus sequence C-x(3)-C-L-x(2)-L-x(4)-C-x(8)-P-x-R-C (SEQ ID NO: 2437) in domain 2 as described by Takenaka et al. In various embodiments, the functional fragments or mutants of luciferases for use with the compositions or methods described herein comprise the consensus sequence C-x(3)-C-L-x(2)-L-x(4)-C-x(8)-P-x-R-C (SEQ ID NO: 2437) in domain 1 and in domain 2, as described by Takenaka et al.

[0073] “Label” as used herein refers to as used herein refers to any atom or molecule that can be used to provide a detectable (preferably quantifiable) signal, and that can be attached to a nucleic acid or protein. Labels may provide signals detectable by fluorescence, radioactivity, colorimetry, gravimetry, X-ray diffraction or absorption, magnetism, enzymatic activity, and the like. A label may be a charged moiety (positive or negative charge) or alternatively, may

be charge neutral. Labels can include or consist of nucleic acid or protein sequence, so long as the sequence comprising the label is detectable.

[0074] There are several methods currently in use for detection of a particular antigen, including flow cytometry, ELISA and Western blotting. However, these methods require expensive equipment (such as flow cytometry), suffer from poor sensitivity and specificity (e.g. ELISA and Western blotting) or are time consuming. The poor sensitivity of the current methods of antigen detection is particularly significant in light of development of new targeted therapies using antibody drug conjugates and chimeric antigen receptors. These new therapies are extremely potent at killing cells expressing their target antigen even at very low levels, which are below the detection limits of current methods. For example, Brentuximab vedotin (Adcetris), an antibody drug conjugate, that targets CD30 antigen, has shown impressive clinical activity in diffuse large cell lymphoma in several clinical trials. However, the response to Brentuximab vedotin (Adcetris) does not correlate with CD30 positivity, as determined by conventional immunohistochemistry, and patients with lymphomas that are CD30 –ve frequently respond to this agent. This has led to the hypothesis that Brentuximab vedotin (Adcetris) requires very low level expression of CD30 for its activity and its efficacy in CD30-ve diffuse large cell lymphoma is, in part, due to low level expression of CD30 in these samples, which is below the detection of conventional methods.

[0075] Chimeric antigen receptor based T cell therapy represents another example where detection of low level expression of an antigen in a patient sample is critically important to predict response to therapy. It is believed that CAR-T cells can kill a target cell expressing as few as 100 molecules of their target antigen. Consistent with this, CAR-T cells targeting CD19 antigen have shown impressive clinical activity in multiple myeloma (Hajek, R *et al.*, Br J Haematol 163:551-564,2013), a disease which is believed to be CD19-ve based on conventional detection methods. Thus, availability of highly sensitive and specific method for detection of CD30, CD19 and other antigens would be of great benefit in determining the patients who are likely to respond to targeted therapies as well as to predict toxicity to normal tissues.

[0076] A frequent challenge in the field of CAR-T cell therapy is the lack of a fast, economical, sensitive and robust assay for detection of chimeric antigen receptors on the

surface of T cells. A sensitive, specific, economical assay for the detection of CAR would be of great benefit as a quality-control tool during the manufacturing of CAR-T cells and to monitor the persistence of CAR-T cells in the body after infusion.

[0077] The current method also has advantage over molecular assay, such as PCR, that rely on mRNA measurement for detection of low level antigens. These molecular tests, although quite sensitive, measure mRNA levels, which may or may not correlate with the level of expression of the protein due to alternative splicing of mRNA or protein degradation. In contrast, the present invention relates to the measurement of low levels of protein antigens, which are the real target of CAR-T cells and therapeutic antibodies. Additionally, the antigen binding fragment (e.g. scFv or the CDRs contained therein) used for the detection of the target antigen in the current invention could be derived from or identical to the antigen binding fragment of a therapeutic antibody or CAR-T cells and therefore is likely to have similar or identical non-specific binding profile. This has obvious advantage in predicting the off-target side-effects of CAR-T cells and therapeutic antibodies due to their even low-level non-specific binding and cross-reactivity with other protein antigens. In addition to clinical use, a highly sensitive and specific method for antigen detection, which is inexpensive, quick and does not require special equipment, would also have utility for research applications.

[0078] The inventor has taken advantage of the extremely high brightness, sensitivity, specificity, stability and small size of luciferases from the copepods (e.g. GLuc, MLuc7, TurboLuc16 etc.) and deep sea shrimp (e.g. NLuc) to develop a sensitive, specific, cost-effective and rapid assay for detection of an antigen. The assay is extremely useful for detection of CAR and for detection of low level antigens on the surface of cells.

Compositions

[0079] Provided herein are compositions comprising fusion proteins for detecting antigens in a sample obtained from a subject. In various embodiments, the fusion protein comprises a first polypeptide chain comprising an antigen recognition domain joined to a second polypeptide chain comprising a reporter protein. In some embodiments, the first and second polypeptide chains are linked via a linker (**Table 5**).

[0080] In one embodiment, the antigen recognition domain comprises a single chain antibody fragments (scFv) or the CDRs contained therein. In some embodiments, the antigen recognition domain comprises more than one single chain antibody, wherein each copy of the single chain antibody recognizes the same epitope on the target antigen. In another embodiment, the antigen recognition domain comprises more than one single chain antibody, wherein each single chain antibody recognizes a different epitope on the same target antigen. Examples of single chain antibodies are described in **Table 14**. Provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, single chain antibodies as described in **Table 14**.

[0081] In one embodiment, the antigen recognition domain comprises a VL fragment (**Table 4**). In some embodiments, the antigen recognition domain comprises more than one VL fragment, wherein each copy of the VL fragment recognizes the same epitope on the target antigen. In another embodiment, the antigen recognition domain comprises more than one VL fragment, wherein each VL fragment recognizes a different epitope on the same target antigen. Examples of VL fragments are described in **Table 4**. Provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, VL fragments as described in **Table 4**.

[0082] In one embodiment, the antigen recognition domain comprises a VH fragment (**Table 6**). In some embodiments, the antigen recognition domain comprises more than one VH fragment, wherein each copy of the VH fragment recognizes the same epitope on the target antigen. In another embodiment, the antigen recognition domain comprises more than one VH fragment, wherein each VH fragment recognizes a different epitope on the same target antigen. Examples of VH fragments are described in **Table 6**. Provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, VH fragments as described in **Table 6**.

[0083] In one embodiment, the antigen recognition domain comprises a VHH domain (nanobodies) (**Table 7**). In some embodiments, the antigen recognition domain comprises more than one VHH domain, wherein each copy of the VHH domain recognizes the same epitope on the target antigen. In another embodiment, the antigen recognition domain comprises more than one VHH domain, wherein each VHH domain recognizes a different

epitope on the same target antigen. Examples of VHH domains are described in **Table 7**. Provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, VHH domains as described in **Table 7**.

[0084] In one embodiment, the antigen recognition domain comprises a non-immunoglobulin antigen binding domain (affibody) (**Table 8**). In some embodiments, the antigen recognition domain comprises more than one affibody, wherein each copy of the affibody recognizes the same epitope on the target antigen. In another embodiment, the antigen recognition domain comprises more than one affibody, wherein each affibody recognizes a different epitope on the same target antigen. Examples of affibodies are described in **Table 8**. Provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, affibodies as described in **Table 8**.

[0085] In one embodiment, the antigen recognition domain comprises a darpin (**Table 8**). In some embodiments, the antigen recognition domain comprises more than one darpin, wherein each copy of the darpin recognizes the same epitope on the target antigen. In another embodiment, the antigen recognition domain comprises more than one darpin, wherein each darpin recognizes a different epitope on the same target antigen. Examples of darpins are described in **Table 8**. Provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, darpins as described in **Table 8**.

[0086] In one embodiment, the antigen recognition domain comprises an extracellular domain of a receptor (**Table 9**). In some embodiments, the antigen recognition domain comprises more than one extracellular domain of a receptor, wherein each extracellular domain of a receptor recognizes the same epitope on the target antigen. In another embodiment, the antigen recognition domain comprises more than one extracellular domain of a receptor, wherein each extracellular domain of a receptor recognizes a different epitope on the same target antigen. Examples of an extracellular domain of a receptor are described in **Table 9**. Provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, extracellular domains of a receptor as described in **Table 9**.

[0087] In one embodiment, the antigen recognition domain comprises a ligand (**Table 10**). In some embodiments, the antigen recognition domain comprises more than ligand, wherein

the ligand recognizes the same epitope on the target antigen. In another embodiment, the antigen recognition domain comprises more than one ligand, wherein each ligand recognizes a different epitope on the same target antigen. Examples of ligands are described in **Table 10**. Provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ligands as described in **Table 10**.

[0088] In various embodiments, the reporter protein component of the composition comprising a fusion protein is a non-secretory form of a luciferase. In various embodiments, the non-secretory form of luciferase is obtained from copepods, deep sea shrimp or homologs or orthologs thereof or mutants or derivatives thereof.

[0089] In exemplary embodiments, the copepods are selected from the group consisting of any one or more of *Gaussia princeps*, *Pleuromamma abdominalis*, *Metridia pacifica*, *Metridia curticauda*, *Metridia asymmetrica*, *Metridia okhotensis*, *Metridia longa*, *Lucicutia ovaliformis*, *Heterorhabdus tanneri*, and *Pleuromamma scutellata*.

[0090] In exemplary embodiments, the reporter luciferase is any one or more of GLuc, NanoLuc (NLuc), MLuc7, HtLuc, LoLuc, PaLuc1, PaLuc2, MpLuc1, McLuc1, MaLuc1, MoLuc1, MoLuc2, MLuc39, PsLuc1, LocLuc1-3, HtLuc2 Renilla, TurboLuc16 (TLuc) or homologs or orthologs thereof or mutants or derivatives thereof.

[0091] In one embodiment, the reporter luciferase is GLuc (*Gaussia princeps* Luc) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 813 or SEQ ID NO: 2029 (**Table 11**) or a sequence with 70-99% homology to sequence set forth in SEQ ID NO: 813 or SEQ ID NO: 2029 (**Table 11**).

[0092] In another embodiment, the reporter luciferase is NLuc (NanoLuc) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 814 or SEQ ID NO: 2030 (**Table 11**) or a sequence with 70-99% homology to sequence set forth in SEQ ID NO: 814 or SEQ ID NO: 2030 (**Table 11**).

[0093] In a further embodiment, the reporter luciferase is TLuc (TurboLuc16) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 815 or SEQ ID NO: 2031 (**Table 11**) or a sequence with 70-99% homology to sequence set forth in SEQ ID NO: 815 or SEQ ID NO: 2031 (**Table 11**).

[0094] In a further embodiment, the reporter luciferase is MLuc7-(*Metridia longa*) Luc M43L/M110L variant and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 816 or SEQ ID NO: 2032 (**Table 11**) or a sequence with 70-99% homology to sequence set forth in SEQ ID NO: 816 or SEQ ID NO: 2032 (**Table 11**).

[0095] In another embodiment, the reporter luciferase is LoLuc (*Lucicutia ovaliformis* Luc) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 817 or SEQ ID NO: 2033 (**Table 11**) or a sequence with 70-99% homology to sequence set forth in SEQ ID NO: 817 or SEQ ID NO: 2033 (**Table 11**).

[0096] In another embodiment, the reporter luciferase is HtLuc (*Heterorhabdus. tanneri* Luc) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 818 or SEQ ID NO: 2034 (**Table 11**) or a sequence with 70-99% homology to sequence set forth in SEQ ID NO: 818 or SEQ ID NO: 2034 (**Table 11**).

[0097] In another embodiment, the reporter luciferase is PaLuc1 (*Pleuromamma abdominalis* Luc) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 819, or SEQ ID NO: 2035 (**Table 11**) or a sequence with 70-99% homology to sequence set forth in SEQ ID NO: 819 or SEQ ID NO: 2035 (**Table 11**).

[0098] In another embodiment, the reporter luciferase is PaLuc2 (*Pleuromamma abdominalis* Luc2) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 820, or SEQ ID NO: 2036 (**Table 11**) or a sequence with 70-99% homology to sequence set forth in SEQ ID NO: 820 or SEQ ID NO: 2036 (**Table 11**).

[0099] In another embodiment, the reporter luciferase is MpLuc1 (*Metridia pacifica* Luc1) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 821, , or SEQ ID NO: 2037 (**Table 11**) or a sequence with 70-99% homology to sequence set forth in SEQ ID NO: 821 or SEQ ID NO: 2037 (**Table 11**).

[00100] In another embodiment, the reporter luciferase is McLuc1 (*Metridia curticauda* Luc1) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 822, or SEQ ID NO: 2038 (**Table 11**) or a sequence with 70-99% homology to sequence set forth in SEQ ID NO: 822 or SEQ ID NO: 2038 (**Table 11**).

[00101] In another embodiment, the reporter luciferase is MaLuc1 (*Metridia asymmetrica* Luc1) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 823 or SEQ ID NO: 2039 (**Table 11**) or a sequence with 70-99% homology to sequence set forth in SEQ ID NO: 823 or SEQ ID NO: 2039 (**Table 11**).

[00102] In another embodiment, the reporter luciferase is MoLuc1 (*Metridia okhotensis* Luc1) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 824 or SEQ ID NO: 2040 (**Table 11**) or a sequence with 70-99% homology to sequence set forth in SEQ ID NO: 824 or SEQ ID NO: 2040 (**Table 11**).

[00103] In another embodiment, the reporter luciferase is MoLuc2 (*Metridia okhotensis* Luc2) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 825, or 2041 (**Table 11**) or a sequence with 70-99% homology to sequence set forth in SEQ ID NO: 825 or 2041 (**Table 11**).

[00104] In another embodiment, the reporter luciferase is MLuc39 (*Metridia longa* Luc39) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 826 or 2042 (**Table 11**).

[00105] In another embodiment, the reporter luciferase is PsLuc1 (*Pleuromamma scutellata* Luc1) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 827 or 2043 (**Table 11**).

[00106] In another embodiment, the reporter luciferase is LoLuc1-3 (*Lucicutia ovaliformis* Luc1-3) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 828 or 2044 (**Table 11**).

[00107] In another embodiment, the reporter luciferase is HtLuc2 (*Heterorhabdus tanneri* Luc2) and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 829 or 2045 (**Table 11**).

[00108] In another embodiment, the reporter luciferase is Renilla Luc and comprises, consists of or consists essentially of the sequence set forth in SEQ ID NO: 830 or 2046 (**Table 11**).

[00109] In some embodiments, the luciferases are fused to antigen recognition domains that bind to a target antigen. In one embodiment, the luciferases are fused to antigen recognition domains that bind to extracellular antigens. In one embodiment, the luciferases are fused to antigen recognition domains that bind to intracellular antigens. In one embodiment, the luciferases are fused to antigen recognition domains that bind to antigens that are expressed on the surface of cells. In some embodiments, the luciferases are fused to antigen recognition domains that bind to antigens that are naturally expressed on the surface of cells. Exemplary antigens that are naturally expressed on the surface of cells include CD19, CD20, CD33 and an endogenous TCR. In some embodiments, the luciferases are fused to antigen recognition domains that bind to antigens that are expressed on the surface of cells in association with MHC complex. In some embodiments, the luciferases are fused to antigen recognition domains that bind to antigens that are expressed on the surface of cells in association with MHC class I complex. In some embodiments, the luciferases are fused to antigen recognition domains that bind to antigens that are expressed on the surface of cells in association with MHC class II complex. In some embodiments, the luciferases are fused to antigen recognition domains that bind to antigens that are not naturally expressed on the surface of cells. In some embodiments, the luciferases are fused to antigen recognition domains that bind to antigens that are expressed on the surface of cells as a result of an intervention, such as genetic modification. Exemplary antigens that are expressed on the surface of cells as a result of an intervention include artificial or synthetic receptors, such as a chimeric antigen receptor, a transgenic T cell receptor, a synthetic immune receptor or a T-body.

[00110] In some embodiments, the luciferases are fused to antigen recognition domains that bind to any one or more of CD19, CD22, CD23, MPL, CD123, CD32, CD138, CD200R, CD276, CD324, CD30, CD171, CS-1, CLL-1 (CLECL1), CD33, EGFRvIII, GD2, GD3, BCMA, Tn Ag, PSMA, ROR1, FLT3, FAP, TAG72, CD38, CD44v6, CEA, EPCAM, B7H3, KIT, IL-13Ra2, IL11Ra, Mesothelin, PSCA, VEGFR2, Lewis Y, CD24, PDGFR-beta, PRSS21, SSEA-4, CD20, Folate receptor alpha, ERBB2 (Her2/neu), MUC1, EGFR, NCAM, Prostase, PAP, ELF2M, Ephrin B2, IGF-I receptor, CAIX, LMP2, gpl00, bcr-abl, tyrosinase, EphA2, Fucosyl GM1, sLe, GM3, TGS5, HMWMAA, o-acetyl-GD2, Folate receptor beta, TEM1/CD248, TEM7R, CLDN6, TSHR, TCR-beta1 constant chain, TCR

beta2 constant chain, TCR gamma-delta, GPRC5D, CXORF61, CD97, CD179a, ALK, Polysialic acid, PLAC1, GloboH, NY-BR-1, UPK2, HAVCR1, ADRB3, PANX3, GPR20, LY6K, OR51E2, TARP, WTI, NY-ESO-1, LAGE-Ia, legumain, HPV E6, E7, HTLV1-Tax, KSHV K8.1 protein, EBV gp350, HIV1-envelop glycoprotein gp120, MAGE-A1, MAGE A1, ETV6-AML, sperm protein 17, XAGE1, Tie 2, MAD-CT-1, MAD-CT-2, Fos-related antigen 1, p53, p53 mutant, prostein, survivin and telomerase, PCTA-1/Galectin 8, MelanA/MART1, Ras mutant, hTERT, DLL3, TROP2, PTK7, GCC, AFP, sarcoma translocation breakpoints, ML-IAP, ERG (TMPRSS2 ETS fusion gene), NA17, PAX3, Androgen receptor, Cyclin B1, MYCN, RhoC, TRP-2, CYP1B1, BORIS, SART3, PAX5, OY-TES1, LCK, AKAP-4, SSX2, RAGE-1, human telomerase reverse transcriptase, RUI, RU2, intestinal carboxyl esterase, mut hsp70-2, CD79a, CD79b, CD72, LAIR1, FCAR, LILRA2, CD300LF, CLEC12A, BST2, EMR2, LY75, GPC3, FCRL5, IGLL1, FITC, Leutenizing hormone receptor (LHR), Follicle stimulating hormone receptor (FSHR), Chorionic Gonadotropin Hormone receptor (CGHR), CCR4, GD3, SLAMF6, SLAMF4, , FITC, Leutenizing hormone receptor (LHR), Follicle stimulating hormone receptor (FSHR), Chorionic Gonadotropin Hormone receptor (CGHR), CCR4, GD3, SLAMF6, SLAMF4, or combinations thereof.

[00111] In some embodiments, the luciferases are fused to antigen recognition domains that bind to an endogenous T cell receptor (TCR), an exogenous (or transgenic) TCR, or a synthetic (or chimeric) T cell receptors (cTCR) targeting any antigen including but not limited to one or more of CD19, CD22, CD23, MPL, CD99, CD123, CD32, CD138, CD200R, CD276, CD324, CD30, CD70, CD179b CD171, CS-1, CLL-1 (CLECL1), CD33, EGFRvIII, GD2, GD3, BCMA, Tn Ag, FcRH5, PSMA, ROR1, FLT3, FAP, TAG72, CD38, CD44v6, CEA, EPCAM, B7H3, KIT, IL-13Ra2, IL11Ra, Mesothelin, , PSCA, VEGFR2, Lewis Y, CD24, PDGFR-beta, PRSS21, SSEA-4, CD20, Folate receptor alpha, ERBB2 (Her2/neu), MUC1, EGFR, NCAM, Prostase, PAP, ELF2M, Ephrin B2, IGF-I receptor, CAIX, LMP2, gpl00, bcr-abl, tyrosinase, EphA2, Fucosyl GM1, sLea, GM3, TGS5, HMWMAA, o-acetyl-GD2, Folate receptor beta, TEM1/CD248, TEM7R, CLDN6, TSHR, TCR-beta1 constant chain, TCR beta2 constant chain, TCR gamma-delta, GPRC5D, CXORF61, CD97, CD179a, ALK, Polysialic acid, PLAC1, GloboH, NY-BR-1, UPK2, HAVCR1, ADRB3, PANX3, GPR20, LY6K, OR51E2, TARP, WTI, NY-ESO-1, LAGE-Ia, legumain, HPV E6, E7, HTLV1-Tax, KSHV K8.1 protein, EBV gp350, HIV1-envelop

glycoprotein gp120, MAGE-A1, MAGE A1, ETV6-AML, sperm protein 17, XAGE1, Tie 2, MAD-CT-1, MAD-CT-2, Fos-related antigen 1, p53, p53 mutant, prostein, survivin and telomerase, PCTA-1/Galectin 8, MelanA/MART1, Ras mutant, hTERT, DLL3, TROP2, PTK7, GCC, AFP, sarcoma translocation breakpoints, ML-IAP, ERG (TMPRSS2 ETS fusion gene), NA17, PAX3, Androgen receptor, Cyclin B1, MYCN, RhoC, TRP-2, CYP11B1, BORIS, SART3, PAX5, OY-TES1, LCK, AKAP-4, SSX2, RAGE-1, human telomerase reverse transcriptase, RUI, RU2, intestinal carboxyl esterase, mut hsp70-2, CD79a, CD79b, CD72, LAIR1, FCAR, LILRA2, CD300LF, CLEC12A, BST2, EMR2, LY75, GPC3, FCRL5, IGLL1, FITC, Leutenizing hormone receptor (LHR), Follicle stimulating hormone receptor (FSHR), Chorionic Gonadotropin Hormone receptor (CGHR), CCR4, GD3, SLAMF6, SLAMF4, , FITC, Leutenizing hormone receptor (LHR), Follicle stimulating hormone receptor (FSHR), Chorionic Gonadotropin Hormone receptor (CGHR), CCR4, GD3, SLAMF6, SLAMF4, or combinations thereof.

[00112] In some embodiments, the luciferases are fused to antigen recognition domains that are scFv fragments (or the CDRs contained therein) specific to the antigen. In some embodiments, the luciferases are fused to antigen recognition domains that are VL fragments (or the CDRs contained therein) specific to the antigen. In some embodiments, the luciferases are fused to antigen recognition domains that are VH fragments (or the CDRs contained therein) specific to the antigen. In some embodiments, the luciferases are fused to antigen recognition domains that are camelid VHH fragments (or the CDRs contained therein) specific to the antigen. In some embodiments, the antigen recognition domains are non-immunoglobulin binding scaffolds specific to the antigen. In some embodiments, the antigen recognition domains are receptor extracellular domains (ECD) that bind to different ligands or are recognized by the antigen specific domains of various natural or synthetic receptors, such as CARs or T cell receptors (TCRs) or artificial TCRs or synthetic TCRs. In some embodiments, the antigen recognition domains are ligands that bind to different receptor or are recognized by the antigen specific domains of various natural or synthetic receptors, such as CARs or T cell receptors (TCRs) or artificial TCRs or synthetic TCRs.

[00113] In an embodiment, the luciferase is fused to an antigen recognition domain comprised by a protein that binds to an artificial receptor such as a CAR, a TCR or a chimeric TCR. In some embodiments, the luciferase is fused to an antigen recognition domain

comprised by a protein that binds to the antigen recognition domain of a CAR. In some embodiments, the protein that binds to a CAR is a scFv fragment. An exemplary protein containing an antigen recognition domain derived from an scFv fragment that binds to a CAR FMC63-(vL-vH)-Myc-BBz-T2A-Pac(SEQ ID NO: 1208) is CD8SP-FMC136-20-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC. The nucleic acid and protein sequences of this protein are represented by SEQ ID NOs: 1167 and 2367. This protein contains a scFv derived from an anti-idiotypic antibody FMC136-20 that recognizes the FMC63 scFv fragment that constitutes the antigen recognition domain of the corresponding CAR FMC63-(vL-vH)-Myc-BBz-T2A-Pac(SEQ ID NO: 1208). In an alternate embodiment, the luciferase is fused to an antigen recognition domain comprised by an antibody or an antibody fragment directed against the variable region of an antibody or an antibody fragment of another species. Such a luciferase fusion protein can be used to detect a CAR containing an antigen recognition domain comprised by the variable region of the said antibody. In an exemplary embodiment, a scFv fragment derived from a mouse antibody directed against the variable region of human IgG can be fused to NLuc and the resulting fusion protein can be used to detect a CAR whose antigen recognition domain is comprised of a scFv fragment derived from a human or a humanized antibody. In an alternate embodiment, the luciferase is fused to an antigen recognition domain comprised by an antibody or an antibody fragment (e.g., scFv) directed against an epitope tag, such as a Myc Tag, that is present on a CAR. In an alternate embodiment, the luciferase is fused to an antigen recognition domain comprised by an antibody or an antibody fragment (e.g., scFv) directed against a label, such as FITC. In some embodiments, the luciferase is fused to an antigen recognition domain comprised by Protein L (SEQ ID NOS. 2431-2434) that is known to bind to the kappa light chain of a CAR. Exemplary nucleotide sequences encoding luciferase and Protein L fusion proteins are CD8SP-Protein-L-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(112316-Q02)[SEQ ID NO:1206] and CD8SP-Protein-L-2-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(101916-P03)[SEQ ID NO:1207]. The corresponding protein sequences are represented by SEQ ID NO: 2406 and 2407, respectively. In some embodiments, the luciferase is fused to an antigen recognition domain comprised by the extracellular domain of a protein which is the target of a CAR. In an exemplary embodiment, the luciferase NLuc is fused to the extracellular domain of MPL (SEQ ID NO: 1749) and the resulting MPL-ECD-GGSG-NLuc-AcV5(082214-Z01)[SEQ ID NO:1169] fusion protein is used to detect a CAR 161-(vL-vH)-

Myc-BBz-T2A-Pac (SEQ ID NO: 1209) that targets MPL. In one embodiment, one or more luciferases are fused to antigen recognition domains that target specific antigens.

[00114] In exemplary embodiments, the antigen recognition domains that target specific antigens are scFv fragments (or the CDRs contained within) and comprise any one or more of the antigen specific scFv sequences described in **Table 14** or sequences with 70-99% identity to antigen specific scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the antigen specific scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigens as the antigen specific scFv sequences in **Table 14**. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target specific antigens and comprise the sequences described in **Table 15**.

[00115] In exemplary embodiments, one or more luciferases are fused to antigen recognition domains that are receptor extracellular domains (ECD) and comprise any one or more of the antigen-specific ECD sequences described in **Table 9** or sequences with 70-99% identity to antigen-specific ECDs sequences described in **Table 9** or deletion mutants and point mutants of the antigen-specific ECD sequences described in **Table 9** as long as they retain the ability to bind to the target antigen. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target an antigen and comprise the antigen-specific sequences described in **Table 16**.

CD19

[00116] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target CD19. In exemplary embodiments, the antigen recognition domains that targets CD19 are scFv fragments and comprise any one or more of the CD19 scFv sequences described in **Table 14** or sequences with 70-99% identity to CD19 scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the CD19 scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigen as the CD19 scFv sequences in **Table 14**. The nucleotide and proteins sequences of exemplary scFv sequences targeting CD19 are provided

in SEQ ID NOs: 564-577 and 1783-1796, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target CD19 and comprise the sequences described in **Table 15**.

MPL

[00117] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target MPL. In exemplary embodiments, the antigen recognition domains that targets MPL are scFv fragments and comprise any one or more of the MPL scFv sequences described in **Table 14** or sequences with 70-99% identity to MPL scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the MPL scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigen as the MPL scFv sequences in **Table 14**. The nucleotide and proteins sequences of exemplary scFv sequences targeting MPL are provided in SEQ ID NOs: 729-736 and 1948-1955, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target MPL and comprise the sequences described in **Table 15**.

CD20

[00118] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target CD20. In exemplary embodiments, the antigen recognition domains that targets CD20 are scFv fragments and comprise any one or more of the CD20 scFv sequences described in **Table 14** or sequences with 70-99% identity to CD20 scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the CD20 scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigen as the CD20 scFv sequences in **Table 14**. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target CD20 and comprise the sequences described in **Table 15**.

PDL1

[00119] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target PDL1. In exemplary embodiments, the antigen recognition domains that targets PDL1 are scFv fragments and comprise any one or more of the PDL1 scFv sequences described in **Table 14** or sequences with 70-99% identity to PDL1 scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the PDL1 scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigen as the PDL1 scFv sequences in **Table 14**. The nucleotide and proteins sequences of exemplary scFv sequences targeting PDL1 are provided in SEQ ID NOs: 745-747 and 1964-1966, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target PDL1 and comprise the sequences (SEQ ID NO: 1062-1064 and 2062-2064) described in **Table 15**.

BCMA

[00120] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target BCMA. In exemplary embodiments, the antigen recognition domains that targets BCMA are scFv fragments and comprise any one or more of the BCMA scFv sequences described in **Table 14** or sequences with 70-99% identity to BCMA scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the BCMA scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigen as the BCMA scFv sequences in **Table 14**. The nucleotide and proteins sequences of exemplary scFv sequences targeting BCMA are provided in SEQ ID NOs: 586-592 and 1805-1811, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target BCMA and comprise the sequences (SEQ ID NO: 878-884 and 2078-2084) described in **Table 15**.

CD30

[00121] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target CD30. In exemplary embodiments, the antigen recognition

domains that targets CD30 are scFv fragments and comprise any one or more of the CD30 scFv sequences described in **Table 14** or sequences with 70-99% identity to CD30 scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the CD30 scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigen as the CD30 scFv sequences in **Table 14**. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target CD30 and comprise the sequences described in **Table 15**.

CD33

[00122] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target CD33. In exemplary embodiments, the antigen recognition domains that targets CD33 are scFv fragments and comprise any one or more of the CD33 scFv sequences described in **Table 14** or sequences with 70-99% identity to CD33 scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the CD33 scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigen as the CD33 scFv sequences in **Table 14**. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target CD33 and comprise the sequences described in **Table 15**.

CD123

[00123] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target CD123. In exemplary embodiments, the antigen recognition domains that targets CD123 are scFv fragments and comprise any one or more of the CD123 scFv sequences described in **Table 14** or sequences with 70-99% identity to CD123 scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the CD123 scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigen as the CD123 scFv sequences in **Table 14**. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target CD123 and comprise the sequences described in **Table 15**.

CD138

[00124] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target CD138. In exemplary embodiments, the antigen recognition domains that targets CD138 are scFv fragments and comprise any one or more of the CD138 scFv sequences described in **Table 14** or sequences with 70-99% identity to CD138 scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the CD138 scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigen as the CD138 scFv sequences in **Table 14**. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target CD138 and comprise the sequences described in **Table 15**.

Her2

[00125] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target Her2. In exemplary embodiments, the antigen recognition domains that targets Her2 are scFv fragments and comprise any one or more of the Her2 scFv sequences described in **Table 14** or sequences with 70-99% identity to Her2 scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the Her2 scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigen as the Her2 scFv sequences in **Table 14**. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target Her2 and comprise the sequences described in **Table 15**.

HIV1 envelop glycoprotein gp120

[00126] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target HIV1 envelop glycoprotein gp120. In exemplary embodiments, the antigen recognition domains that targets HIV1 envelop glycoprotein gp120 are scFv fragments and comprise any one or more of the HIV1 envelop glycoprotein gp120 scFv sequences described in **Table 14** or sequences with 70-99% identity to HIV1 envelop glycoprotein gp120 scFv sequence in **Table 14** or sequences with 70-99% identity in the six

complementarity determining regions (CDRs) to the HIV1 envelop glycoprotein gp120 scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigens as the HIV1 envelop glycoprotein gp120 scFv sequences in **Table 14**. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target HIV1 envelop glycoprotein gp120 and comprise the sequences described in **Table 15**.

Amyloid

[00127] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target Amyloid. In exemplary embodiments, the antigen recognition domains that targets Amyloid are scFv fragments and comprise any one or more of the Amyloid scFv sequences described in **Table 14** or sequences with 70-99% identity to Amyloid scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the Amyloid scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigen as the Amyloid scFv sequences in **Table 14**. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target Amyloid and comprise the sequences described in **Table 15**.

AFP/MHC class I complex

[00128] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target AFP/MHC class I complex. In exemplary embodiments, the antigen recognition domains that targets AFP/MHC class I complex are scFv fragments and comprise any one or more of the AFP/MHC class I complex scFv sequences described in **Table 14** or sequences with 70-99% identity to AFP/MHC class I complex scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the AFP/MHC class I complex scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigen as the AFP/MHC class I complex scFv sequences in **Table 14**. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target AFP/MHC class I complex and comprise the sequences described in **Table 15**.

CD22

[00129] In one embodiment, the one or more luciferases are fused to antigen recognition domains that target CD22. In exemplary embodiments, the antigen recognition domains that targets CD22 are scFv fragments and comprise any one or more of the CD22 scFv sequences described in **Table 14** or sequences with 70-99% identity to CD22 scFv sequences in **Table 14** or sequences with 70-99% identity in the six complementarity determining regions (CDRs) to the CD22 scFv sequences in **Table 14** or sequences that bind to the same epitopes on the target antigen as the CD22 scFv sequences in **Table 14**. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, scFv-GGS-Luc constructs that target CD22 and comprise the sequences described in **Table 15**.

CD19-specific CAR

[00130] In exemplary embodiments, the antigen recognition domain that targets a CD19-specific CARs is a receptor extracellular domains (ECD) and comprises the CD19 ECD sequences as described in **Table 9** or sequences with 70-99% identity to CD19 ECD sequences described in **Table 9** or deletion mutants and point mutants of the CD19 ECD sequences described in **Table 9** as long as they retain the ability to bind to CD19-specific CARs. The nucleic acid and protein sequences of exemplary receptor ECD sequences targeting CD19-specific CARs are provided in SEQ ID NOs: 529 and 1748, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target CD19 and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting CD19-CARs are provided in SEQ ID NOs: 1170-1176 and 2370-2376, respectively.

[00131] In exemplary embodiments, the antigen recognition domain that targets a CD19-specific FMC63-scFv based CAR is comprised of the scFV fragment of an anti-idiotypic antibody FMC136-20 directed against the CD19-specific antibody FMC63. The nucleotide and proteins sequences of exemplary FMC136-20-GGS-Luc constructs targeting CD19-specific FMC63-scFv based CAR are provided in SEQ ID NOs: 1167 and 2367, respectively.

MPL-CAR

[00132] In exemplary embodiments, the antigen recognition domain that targets an MPL-specific CAR is a receptor extracellular domains (ECD) and comprise the MPL ECD sequence described in **Table 9** or sequences with 70-99% identity to MPL ECDs sequences described in **Table 9** or deletion mutants and point mutants of the MPL ECD sequences described in **Table 9** as long as they retain the ability to bind to MPL-specific CARs. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting MPL-CARs are provided in SEQ ID NOs: 530 and 1749, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target MPL and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting MPL-CAR are provided in SEQ ID NOs: 1169 and 2369, respectively.

CD20-sepcific CAR

In exemplary embodiments, the antigen recognition domain that targets a CD20 -specific CAR is a receptor extracellular domain (ECD) and comprises the CD20 ECD sequences described in **Table 9** or sequences with 70-99% identity to CD20 ECDs sequences described in **Table 9** or deletion mutants and point mutants of the CD20 ECD sequences described in **Table 9** as long as they retain the ability to bind to CD20-specific CARs. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting CD20-specific CARs are provided in SEQ ID NOs: 545-546 and 1764-1765, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target CD20 and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting CD20 -specific CARs are provided in SEQ ID NOs: 1197-1198 and 2397-2398, respectively.

CD33-sepcific CAR

[00133] In exemplary embodiments, the antigen recognition domain that targets a CD33-specific CAR is a receptor extracellular domain (ECD) and comprises the CD33 ECD

sequence described in **Table 9** or sequences with 70-99% identity to CD33 ECDs sequences described in **Table 9** or deletion mutants and point mutants of the CD33 ECD sequences described in **Table 9** as long as they retain the ability to bind to CD33-specific CARs. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting CD33-specific CARs are provided in SEQ ID NOs: 541 and 1761, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target CD33 and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting CD33-specific CARs are provided in SEQ ID NOs: 1177-1182 and 2377-2382, respectively.

CD123-specific CAR

[00134] In exemplary embodiments, the antigen recognition domain that targets a CD123-specific CAR is a receptor extracellular domain (ECD) and comprises the CD123 ECD sequence described in **Table 9** or sequences with 70-99% identity to CD123 ECDs sequences described in **Table 9** or deletion mutants and point mutants of the CD123 ECD sequences described in **Table 9** as long as they retain the ability to bind to CD123-specific CARs. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting CD123-specific CARs are provided in SEQ ID NOs: 536 and 1755, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target CD123 and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting CD123-specific CARs are provided in SEQ ID NOs: 1188 and 2388, respectively.

CD138-specific CAR

[00135] In exemplary embodiments, the antigen recognition domain that targets a CD138-specific CAR is a receptor extracellular domains (ECD) and comprise the CD138 ECD sequence described in **Table 9** or sequences with 70-99% identity to CD138 ECDs sequences described in **Table 9** or deletion mutants and point mutants of the CD138 ECD sequences described in **Table 9** as long as they retain the ability to bind to CD138-specific CARs. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting

CD138-specific CARs are provided in SEQ ID NOs: 535 and 1754, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target CD138 and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting CD138-specific CARs are provided in SEQ ID NOs: 1183-1187 and 2383-2387, respectively.

CD200R-specific CAR

[00136] In exemplary embodiments, the antigen recognition domain that targets a CD200R-specific CAR is a receptor extracellular domain (ECD) and comprises the CD200R ECD sequence described in **Table 9** or sequences with 70-99% identity to CD200R ECDs sequences described in **Table 9** or deletion mutants and point mutants of the CD200R ECD sequences described in **Table 9** as long as they retain the ability to bind to CD200R-specific CARs. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting CD200R-specific CARs are provided in SEQ ID NOs: 538 and 1757, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target CD200R and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting CD200R-specific CARs are provided in SEQ ID NOs: 1190 and 2390, respectively.

CD22-specific CAR

[00137] In exemplary embodiments, the antigen recognition domain that targets a CD22-specific CAR is a receptor extracellular domain (ECD) and comprises the CD22 ECD sequence described in **Table 9** or sequences with 70-99% identity to CD22 ECDs sequences described in **Table 9** or deletion mutants and point mutants of the CD22 ECD sequences described in **Table 9** as long as they retain the ability to bind to CD22-specific CARs. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting CD22-specific CARs are provided in SEQ ID NOs: 547 and 1766, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target CD22 and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of

exemplary ECD-GGS-Luc constructs targeting CD22-specific CARs are provided in SEQ ID NOs: 1198 and 2398, respectively.

BCMA-specific CAR

[00138] In exemplary embodiments, the antigen recognition domain that targets a BCMA-specific CAR is a receptor extracellular domain (ECD) and comprises the BCMA ECD sequence described in **Table 9** or sequences with 70-99% identity to BCMA ECDs sequences described in **Table 9** or deletion mutants and point mutants of the BCMA ECD sequences described in **Table 9** as long as they retain the ability to bind to BCMA-specific CARs. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting BCMA-specific CARs are provided in SEQ ID NOs: 550 and 1769, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target BCMA and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting BCMA-specific CARs are provided in SEQ ID NOs: 1201 and 2401, respectively.

SLAMF7/CS1-specific CAR

[00139] In exemplary embodiments, the antigen recognition domain that targets a SLAMF7/CS1-specific CAR is a receptor extracellular domain (ECD) and comprises the SLAMF7/CS1 ECD sequence described in **Table 9** or sequences with 70-99% identity to SLAMF7/CS1 ECDs sequences described in **Table 9** or deletion mutants and point mutants of the SLAMF7/CS1 ECD sequences described in **Table 9** as long as they retain the ability to bind to SLAMF7/CS1-specific CARs. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting SLAMF7/CS1-specific CARs are provided in SEQ ID NOs: 551 and 1770, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target SLAMF7/CS1 and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting SLAMF7/CS1-specific CARs are provided in SEQ ID NOs: 1202 and 2402, respectively.

PTK7-specific CAR

[00140] In exemplary embodiments, the antigen recognition domain that targets a PTK7-specific CAR is a receptor extracellular domain (ECD) and comprises the PTK7 ECD sequence described in **Table 9** or sequences with 70-99% identity to *PTK7* ECDs sequences described in **Table 9** or deletion mutants and point mutants of the *PTK7* ECD sequences described in **Table 9** as long as they retain the ability to bind to *PTK7*-specific CARs. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting PTK7 - specific CARs are provided in SEQ ID NOs: 540 and 1759, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target PTK7 and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting PTK7-specific CARs are provided in SEQ ID NOs: 1192 and 2392, respectively.

EGFRviii-specific CAR

[00141] In exemplary embodiments, the antigen recognition domain that targets a EGFRviii-specific CAR is a receptor extracellular domain (ECD) and comprises the EGFRviii ECD sequence described in **Table 9** or sequences with 70-99% identity to *EGFRviii* ECDs sequences described in **Table 9** or deletion mutants and point mutants of the *EGFRviii* ECD sequences described in **Table 9** as long as they retain the ability to bind to *EGFRviii*-specific CARs. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting EGFRviii -specific CARs are provided in SEQ ID NOs: 549 and 1768, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target EGFRviii and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting EGFRviii-specific CARs are provided in SEQ ID NOs: 1200 and 2400, respectively.

TSHR-specific CAR

[00142] In exemplary embodiments, the antigen recognition domain that targets a TSHR-specific CAR is a receptor extracellular domain (ECD) and comprises the TSHR ECD

sequence described in **Table 9** or sequences with 70-99% identity to TSHR ECDs sequences described in **Table 9** or deletion mutants and point mutants of the TSHR ECD sequences described in **Table 9** as long as they retain the ability to bind to TSHR-specific CARs. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting TSHR - specific CARs are provided in SEQ ID NOs: 548 and 1767, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target TSHR and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting TSHR-specific CARs are provided in SEQ ID NOs: 1191 and 2391, respectively.

gpNMB-specific CAR

[00143] In exemplary embodiments, the antigen recognition domain that targets a gpNMB-specific CAR is a receptor extracellular domain (ECD) and comprises the gpNMB ECD sequence described in **Table 9** or sequences with 70-99% identity to gpNMB ECDs sequences described in **Table 9** or deletion mutants and point mutants of the gpNMB ECD sequences described in **Table 9** as long as they retain the ability to bind to gpNMB-specific CARs. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting gpNMB -specific CARs are provided in SEQ ID NOs: 539 and 1758, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target gpNMB and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting gpNMB-specific CARs are provided in SEQ ID NOs: 1169 and 2369, respectively.

EpCam-specific CAR

[00144] In exemplary embodiments, the antigen recognition domain that targets an EpCam-specific CAR is a receptor extracellular domain (ECD) and comprises the EpCam ECD sequence described in **Table 9**. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting EpCam-specific CARs are provided in SEQ ID NOs: 543 and 1762, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc

constructs that target Epcam and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting EpCam-specific CARs are provided in SEQ ID NOs: 1194 and 2394, respectively.

CD34 -specific CARs

[00145] In exemplary embodiments, the antigen recognition domain that targets a CD34-specific CAR is a receptor extracellular domain (ECD) and comprises the CD34 ECD sequence described in **Table 9**. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting CD34-specific CARs are provided in SEQ ID NOs: 542 and 1761, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target CD34 and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting CD34-specific CARs are provided in SEQ ID NOs: 1193 and 2393, respectively.

CDH1-specific CAR

[00146] In exemplary embodiments, the antigen recognition domain that targets a CDH1-specific CAR is a receptor extracellular domain (ECD) and comprises the CDH1(CD324) ECD sequence described in **Table 9**. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting CDH1-specific CARs are provided in SEQ ID NOs: 537 and 1756, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target CDH1 and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting CDH1-specific CARs are provided in SEQ ID NOs: 1189 and 2389, respectively.

PD1-specific CAR

[00147] In exemplary embodiments, the antigen recognition domain that targets a PD1-specific CAR is a receptor extracellular domain (ECD) and comprises the PD1 ECD sequence described in **Table 9**. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting PD1-specific CARs are provided in SEQ ID NOs: 531-534 and 1750-1752, respectively. In some embodiments, provided herein are compositions comprising

polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target PD1 and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting PD1-specific CARs are provided in SEQ ID NOs: 1203 and 2403, respectively.

CTLA4-specific CAR

[00148] In exemplary embodiments, the antigen recognition domain that targets a CTLA4-specific CAR is a receptor extracellular domain (ECD) and comprises the CTLA4 ECD sequence described in **Table 9**. The nucleotide and protein sequences of exemplary receptor ECD sequences targeting CTLA4-specific CARs are provided in SEQ ID NOs: 534 and 1753, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target CTLA4 and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting CTLA4-specific CARs are provided in SEQ ID NOs: 1204 and 2404, respectively.

NKG2D-specific CAR

[00149] In exemplary embodiments, the antigen recognition domain that targets a NKG2D-specific CAR is a receptor extracellular domain (ECD) and comprises the NKG2D ECD sequence described in **Table 9**. The nucleotide and protein sequences of an exemplary receptor ECD sequence (lacking the signal peptide) targeting NKG2D-specific CARs is provided in SEQ ID NOs: 552 and 1771, respectively. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target NKG2D and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting NKG2D-specific CARs are provided in SEQ ID NOs: 1205 and 2405, respectively.

Kappa light chain containing CAR

[00150] In exemplary embodiments, the antigen recognition domain that targets any kappa light chain containing CAR comprises, consists or essentially consists of Protein L. The nucleotide and protein sequences of exemplary Protein L sequences targeting any kappa

light chain containing CAR are provided in SEQ ID NOs: 2431-2432 and 2433-2434, respectively or sequences with 70-99% identity to Protein L sequences provided in SEQ ID NOs: 2431-2432 and 2433-2434, respectively, or deletion mutants and point mutants of the Protein L sequences provided in SEQ ID NOs: 2431-2432 and 2433-2434, respectively, as long as they retain the ability to bind to Kappa light chains. In some embodiments, provided herein are compositions comprising polynucleotides encoding, polypeptides encoded by and vectors encoding, ECD-GGS-Luc constructs that target NKG2D and comprise the sequences described in **Table 16**. The nucleotide and proteins sequences of exemplary ECD-GGS-Luc constructs targeting NKG2D-specific CARs are provided in SEQ ID NOs: 1206-1207 and 2406-2407, respectively.

Detection methods

[00151] Broadly, the detection methods include linking an antigen binding domain to a luciferase and using the fusion protein to detect the antigen.

[00152] In the preferred embodiment, the detection methods include fusing the nucleotide sequence encoding a scFV fragment or, another antigen binding fragment (such as nanobodies, affibodies, Darpins etc.), in frame to the nucleotide sequences encoding the luciferases. The expression cassette may also carry nucleotide sequences encoding single or multiple copies of additional epitope tags, such as FLAG, HA and MYC tags, as well tags for purification, such as Streptag and polyhistidine tag. They could also carry nucleotide sequences encoding fluorescent proteins (e.g. EGFP, mcherry) or other reporters (e.g. Horse reddish peroxidase or alkaline phosphatase). The above fusion protein is expressed in appropriate host and then used to detect the expression of the antigen targeted by the scFV (or other antigen binding fragment) by incubation with the target cell (or tissue or biological specimen) followed by extensive washes and then measurement of the bound fusion protein by luciferase reporter assay. In an alternate embodiment, the fusion protein is used to detect an antigen by indirect labeling. In some embodiments, a luciferase fusion protein with an antigen recognition domain directed against another antibody, an epitope tag or a label (e.g., FITC) can be used to identify a target antigen in an indirect labeling protocol. In an exemplary embodiment, a cell expressing a CAR containing a humanized scFv fragment can be first labeled with a FITC-conjugated antibody against human IgG variable region (e.g., a

Goat anti-Human IgG F(ab')₂ fragment specific; Jackson Immuno Research Laboratory; Cat No: 109-096-006) and then labeled with a luciferase fusion protein containing an antigen recognition domain comprised by a scFv directed against FITC (e.g., SEQ ID NO: 2184 or 2185). Such an indirect labeling approach is expected to improve the sensitivity of the assay by amplifying the signal as each molecule of CAR is expected to be bound by many molecules of the FITC-conjugated goat anti-human antibody, which in turn would be bound by many molecules of the anti-FITC-luciferase fusion protein (e.g., SEQ ID NO: 2184 or 2185). In another exemplary embodiment, a CAR containing a humanized scFv fragment is first labeled with an antibody against human IgG variable region (e.g., a Goat anti-Human IgG F(ab')₂ fragment specific; Jackson Immuno Research Laboratory; Cat No: 109-006-006) and then labeled with a luciferase fusion protein containing an antigen recognition domain comprised by a scFv directed against Goat IgG Fc region. The fusion proteins can be also used to detect the presence of very low level target antigens by other methods, including western blotting and microscopy. Finally, the fusion proteins described in this invention can be used to detect the presence of low level antigens in in vivo by bioluminescence imaging for research as well as clinical applications. A potential limitation of the current invention for in vivo application is the immunogenicity of the Luc moiety, which could limit the use to a single or a few applications. However, since the current invention is not limited to a single Luc moiety, it is possible to develop a panel of fusion proteins containing the same antigen binding fragment but different Luc fragments. Thus, a subject who has received FMC63-NLuc fusion protein and has become immunized to it, might receive FMC63-GLuc fusion protein in the future, thereby overcoming the immune response to the NLuc moiety.

[00153] The CAR-T cells are living drugs as the cells once infused can multiply in the patient. They can also kill a target cell that expresses as few as 100 molecules of their target antigen. In addition, CAR cells can kill healthy cells that do not express their target via a bystander effect through production of cytokines. Therefore, toxicity of the CAR and other immunotherapy products (including antibody drug conjugates, and Bispecific T cell engagers or BiTE, and DARTs) on normal tissues is a major concern as even low level expression of the target antigen on a vital organ can be lethal. The assay described herein is sensitive provides a useful platform to monitor low level target antigen expression on normal tissues.

Antigen detection methods

[00154] Provided herein are methods for detecting the presence or absence of an antigen. The methods include contacting a fusion protein comprising an antigen recognition domain fused to a reporter and assaying the activity of the reporter in the presence of a reporter-specific substrate. In one embodiment, the presence of reporter activity is indicative of presence of antigen. In another embodiment, an increase in reporter activity relative to a reference value is indicative of presence of antigen. In one embodiment, the antigen recognition domain of the fusion protein is directed against the antigen to be detected. In another embodiment, the antigen recognition domain of the fusion protein is directed against an antigen (i.e., a protein) that is directed against the antigen to be detected. In some embodiments, the fusion protein comprising an antigen recognition domain fused to a reporter binds directly to the antigen to be detected (i.e., direct labeling). In some embodiments, the fusion protein comprising an antigen recognition domain fused to a reporter binds to the antigen to be detected through one or more intermediates (i.e., indirect labeling). In some embodiments, the reporter is a luciferase as described herein. In one embodiment, the antigen is detected in a subject (i.e., *in vivo*). In another embodiment, the antigen is detected outside a subject. In one embodiment, the antigen is detected in a sample obtained from a subject. In some embodiments, the antigens are expressed on cancer cells. In one embodiment, the subject is diagnosed with cancer. In another embodiment, the subject is suspected of having cancer. The methods further comprise removing the unbound fusion protein prior to assaying using the reporter-specific substrates. In some embodiments, the reporter is a non-secretory form of a luciferase. In some embodiments, the non-secretory form of luciferase is obtained from copepods, deep sea shrimp or homologs or orthologs thereof or mutants or derivatives thereof. In exemplary embodiments, the copepods are selected from the group consisting of any one or more of *Gaussia princeps*, *Pleuromamma abdominalis*, *Metridia pacifica*, *Metridia curticauda*, *Metridia asymmetrica*, *Metridia okhotensis*, *Metridia longa*, *Lucicutia ovaliformis*, *Heterorhabdus tanneri*, and *Pleuromamma scutallata*. In further exemplary embodiments, the luciferase is any one or more of GLuc, NLuc, MLuc7, PaLuc1, PaLuc2, MpLuc1, MoLuc1, MoLuc2, MLuc39, HtLuc, HtLuc2, PsLuc1, H-tanneri, *Lucicutia ovaliformis* (LoLuc), LoLuc1-3, Renilla or TLuc (TurboLuc16) or derivatives thereof. In some embodiments, the reporter activity is assayed

by exposing the target cells to a luciferase-specific substrate. In one embodiment, the luciferase-specific substrate is coelentraine or a derivative thereof. In another embodiment, the luciferase-specific substrate is imidazopyrazinone substrate (furimazine) or a derivative thereof.

[00155] Provided herein are methods for detecting the presence or absence of an antigen in a subject in need thereof. The methods include providing a fusion protein comprising an antigen recognition domain fused to a reporter; providing a sample from the subject; exposing the sample to the fusion protein; and assaying the activity of the reporter in the presence of a reporter-specific substrate. In one embodiment, the presence of reporter activity is indicative of presence of antigen. In another embodiment, an increase in reporter activity relative to a reference value is indicative of presence of antigen. In some embodiments, the antigens are expressed on cancer cells. In one embodiment, the subject has cancer. The methods further comprise removing the unbound fusion protein prior to assaying using reporter-specific substrates. In some embodiments, the reporter is a non-secretory form of a luciferase. In some embodiments, the non-secretory form of luciferase is obtained from copepods, deep sea shrimp or homologs or orthologs thereof or mutants or derivatives thereof. In exemplary embodiments, the copepods are selected from the group consisting of any one or more of *Gaussia princeps*, *Pleuromamma abdominalis*, *Metridia pacifica*, *Metridia curticauda*, *Metridia asymmetrica*, *Metridia okhotensis*, *Metridia longa*, *Lucicutia ovaliformis*, *Heterorhabdus tanneri*, and *Pleuromamma scutellata*. In further exemplary embodiments, the luciferase is any one or more of GLuc, NLuc, MLuc7, PaLuc1, PaLuc2, MpLuc1, MoLuc1, MoLuc2, MLuc39, HtLuc, HtLuc2, PsLuc, LoLuc, LoLuc1-3, Renilla or TLuc or derivative thereof. In some embodiments, the reporter activity is assayed by exposing the target cells to a luciferase-specific substrate. In one embodiment, the luciferase-specific substrate is coelentraine or a derivative thereof. In another embodiment, the luciferase-specific substrate is imidazopyrazinone substrate (furimazine) or a derivative thereof.

[00156] In some embodiments of the methods described herein, the antigen targeting domains comprise an antibody or a functional equivalent thereof or a fragment thereof or a derivative thereof specific to the antigen of interest. The targeting regions may comprise full length heavy chain, Fab fragments, single chain Fv (scFv) fragments (or the CDRs contained

therein), divalent single chain antibodies or diabodies, each of which are specific to the target antigen. In some embodiments, the antigen specific targeting region is linked cytokines (which leads to recognition of cells bearing the cytokine receptor), affibodies, ligand binding domains from naturally occurring receptors, soluble protein/peptide ligand for a receptor (for example on a tumor cell), peptides, and vaccines to prompt an immune response, which may each be used in various embodiments of the invention. In some embodiments, almost any molecule that binds a given antigen with high affinity can be used as an antigen specific targeting region, as will be appreciated by those of skill in the art.

[00157] In some embodiments of the methods described herein, the fusion proteins may be tagged with different tags (e.g. FLAG, HA, MYC, T7, AcV5, V5, StrepTagII, poly His, etc.), either singly or in combination. The fusion proteins comprising the tags may then be detected with fluorescently labeled antibodies against the epitope tags. Alternatively, the fusion proteins may be fused to fluorescent proteins, such as EGFP, RFP and mcherry, for detection directly. Finally, the epitope tags, such as StrepTagII and polyHis tags, can be also used for the purification of the scFv fusion proteins. The tags may be also used to join the antigen specific domain to the luciferase. In an exemplary embodiment, both the antigen specific domain (e.g., a scFv) and the luciferase (e.g., NLuc) carry a StrepTagII. Both the StrepTagII fusion proteins are expressed separately and purified. Subsequently, the scFv-StrepTag fusion protein is incubated with Streptavidin at a molar ratio of 1:1 or 1:2 or 1:5 so that not all StreptTagII binding sites on streptavidin are occupied by the scFv-StrepTagII fusion protein. Next, the Streptavidin-scFv-StrepTagII complex is incubated with NLuc-StrepTagII fusion protein so that the free sites on streptavidin are occupied by the StrepTagII present on the NLuc fusion protein. Thus, in alternate embodiments of the invention, it is possible to link an antigen specific domain to a luciferase by methods other than expressing them as a single polypeptide.

[00158] In various embodiments, examples of disease specific antigens that may be detected by the methods described herein include but are not limited to PDL1, Her2, PD1, 4-1BB, 5T4, adenocarcinoma antigen, alpha-fetoprotein, BAFF, B-lymphoma cell, C242 antigen, CA-125, carbonic anhydrase 9 (CA-IX), C-MET, CCR4, CD152, CD19, CD20, CD200, CD22, CD221, CD23 (IgE receptor), CD28, CD30 (TNFRSF8), CD33, CD4, CD40, CD44 v6, CD51, CD52, CD56, CD74, CD80, CEA, CNT0888, CTLA-4, DR5, EGFR,

EpCAM, CD3, FAP, fibronectin extra domain-B, folate receptor 1, GD2, GD3 ganglioside, glycoprotein 75, GPNMB, HER2/neu, HGF, human scatter factor receptor kinase, IGF-1 receptor, IGF-I, IgG1, L1-CAM, IL-13, IL-6, insulin-like growth factor I receptor, integrin $\alpha 5\beta 1$, integrin $\alpha v\beta 3$, MORAb-009, MS4A1, MUC1, mucin CanAg, N-glycolylneuraminic acid, NPC-1C, PDGF-R α , PDL192, phosphatidylserine, prostatic carcinoma cells, RANKL, RON, ROR1, SCH 900105, SDC1, SLAMF7, thrombopoietin receptor, TAG-72, tenascin C, TGF beta 2, TGF- β , TRAIL-R1, TRAIL-R2, tumor antigen CTAA16.88, VEGF-A, VEGFR-1, VEGFR2 or vimentin. Other antigens specific for disease, including cancer, infectious diseases, allergic diseases and autoimmune diseases, will be apparent to those of skill in the art and may be used in connection with alternate embodiments of the invention.

[00159] In one embodiment of the antigen detection methods described herein, the antigen recognition domain comprises VL fragments of antibodies that recognize and bind specific antigens. In exemplary embodiments, the VL fragments for use with the antigen detection methods are described in **Table 4** and are fused with any luciferase reporters described in **Table 11**. In some embodiments, the VL fragments are linked to the luciferase reporter via the linkers described in **Table 5**. In some embodiments, the fusion protein comprising VL fragments and luciferase reporter further comprise one or more epitope tags described in **Table 13**.

[00160] In one embodiment of the antigen detection methods described herein, the antigen recognition domain comprises VH fragments of antibodies that recognize and bind specific antigens. In exemplary embodiments, the VH fragments for use with the antigen detection methods are described in **Table 6** and are fused with any luciferase reporter described in **Table 11**. In some embodiments, the VH fragments are linked to the luciferase reporter via the linkers described in **Table 5**. In some embodiments the fusion protein comprising VH fragments and luciferase reporter further comprise one or more epitope tags described in **Table 13**.

[00161] In one embodiment of the antigen detection methods described herein, the antigen recognition domain comprises VHH fragments of antibodies that recognize and bind specific antigens. In exemplary embodiments, the VHH fragments for use with the antigen detection methods are described in **Table 7** and are fused with any luciferase described in

Table 11. In some embodiments, the VHH fragments are linked to the luciferase reporter via the linkers described in **Table 5**. In some embodiments the fusion protein comprising VHH fragments and luciferase reporter further comprise one or more epitope tags described in **Table 13**.

[00162] In one embodiment of the antigen detection methods described herein, the antigen recognition domain comprises non-immunoglobulin antigen binding scaffolds, such as affibodies, that recognize and bind specific antigens. In exemplary embodiments, the affibodies for use with the antigen detection methods are described in **Table 8** and are fused with any luciferase described in **Table 11**. In some embodiments, the affibodies are linked to the luciferase reporter via the linkers described in **Table 5**. In some embodiments the fusion protein comprising affibodies and luciferase reporter further comprise one or more epitope tags described in **Table 13**.

[00163] In one embodiment of the antigen detection methods described herein, the antigen recognition domain comprises darpins that recognize and bind specific antigens. In exemplary embodiments, the darpins for use with the antigen detection methods are described in **Table 8** and are fused with any luciferase described in **Table 11**. In some embodiments, the darpins are linked to the luciferase reporter via the linkers described in **Table 5**. In some embodiments the fusion protein comprising darpins and luciferase reporter further comprise one or more epitope tags described in **Table 13**.

[00164] In one embodiment of the antigen detection methods described herein, the antigen recognition domain comprises receptor extracellular domains that recognize and bind specific antigens. In exemplary embodiments, the extracellular domains for use with the antigen detection methods are described in **Table 9** and are fused with any luciferase described in **Table 11**. In some embodiments, the extracellular domains are linked to the luciferase reporter via the linkers described in **Table 5**. In some embodiments the fusion protein comprising receptor extracellular domains and luciferase reporter further comprise one or more epitope tags described in **Table 13**.

[00165] In one embodiment of the antigen detection methods described herein, the antigen recognition domain comprises ligands that recognize and bind specific antigens. In exemplary embodiments, the ligands for use with the antigen detection methods are described

in **Table 10** and are fused with any luciferase described in **Table 11**. In some embodiments, the ligands are linked to the luciferase reporter via the linkers described in **Table 5**. In some embodiments the fusion protein comprising ligands and luciferase reporter further comprise one or more epitope tags described in **Table 13**.

[00166] In one embodiment of the antigen detection methods described herein, the antigen recognition domain comprises scFv fragments (or the CDRs contained therein) of antibodies that recognize and bind specific antigens. In exemplary embodiments, the scFv fragments for use with the antigen detection methods are described in **Table 14** and are fused with any luciferase described in **Table 11**. In some embodiments, the VL fragments are linked to the luciferase reporter via the linkers described in **Table 5**. In some embodiments the fusion protein comprising scFv fragments (or the CDRs contained therein) and luciferase reporter further comprise one or more epitope tags described in **Table 13**.

[00167] In some embodiments, provided herein are methods for detecting presence or absence of CD19 expressing cells comprising contacting a sample from a subject with a fusion protein comprising scFv specific to CD19 for a sufficient amount of time to allow the CD19-specific fusion protein to bind CD19 expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of presence of CD19 expressing cells. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of presence of CD19 expressing cells. In one embodiment, the method further comprises selecting a therapy if presence of CD19 expressing cells in the sample is determined. In exemplary embodiments, the CD19 specific fusion protein comprises GGS linker and NLuc reporter as described in any one or more of SEQ ID Nos: 854-868 or 2054-2068 (**Table 15**).

[00168] In some embodiments, provided herein are methods for detecting presence or absence of CD20 expressing cells comprising contacting a sample from a subject with a fusion protein comprising scFv specific to CD20 for a sufficient amount of time to allow the CD20-specific fusion protein to bind CD20 expressing cells in the sample. In some

embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of presence of CD20 expressing cells. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of presence of CD20 expressing cells. In one embodiment, the method further comprises selecting a therapy if presence of CD20 expressing cells in the sample is determined. In exemplary embodiments, the CD20 specific fusion protein comprises GGS linker and NLuc reporter as described in any one or more of SEQ ID Nos 891-903 or 2091-2103 (**Table 15**).

[00169] In some embodiments, provided herein are methods for detecting presence or absence of CD30 expressing cells comprising contacting a sample from a subject with a fusion protein comprising scFv specific to CD30 for a sufficient amount of time to allow the CD30-specific fusion protein to bind CD30 expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of presence of CD30 expressing cells. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of presence of CD30 expressing cells. In one embodiment, the method further comprises selecting a therapy if presence of CD30 expressing cells in the sample is determined. In exemplary embodiments, the CD30 specific fusion protein comprises GGS linker and NLuc reporter as described in any one or more of SEQ ID Nos: 908-909 or 2108-2109 (**Table 15**).

[00170] In some embodiments, provided herein are methods for detecting presence or absence of Lym1 expressing cells comprising contacting a sample from a subject with a fusion protein comprising scFv specific to Lym1 for a sufficient amount of time to allow the Lym1-specific fusion protein to bind Lym1 expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate

and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of presence of Lym1 expressing cells. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of presence of Lym1 expressing cells. In one embodiment, the method further comprises selecting a therapy if presence of Lym1 expressing cells in the sample is determined. In exemplary embodiments, the Lym1 specific fusion protein comprises GGS linker and NLuc reporter as described in any one or more of SEQ ID NO: 1036 or 2236 (**Table 15**).

[00171] In some embodiments, provided herein are methods for detecting presence or absence of Lym2 expressing cells comprising contacting a sample from a subject with a fusion protein comprising scFv specific to Lym2 for a sufficient amount of time to allow the Lym2-specific fusion protein to bind Lym2 expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of presence of Lym2 expressing cells. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of presence of Lym2 expressing cells. In one embodiment, the method further comprises selecting a therapy if presence of Lym2 expressing cells in the sample is determined. In exemplary embodiments, the Lym2 specific fusion protein comprises GGS linker and NLuc reporter as described in any one or more of SEQ ID NO: 1037 or 2237 (**Table 15**).

[00172] In some embodiments, provided herein are methods for detecting presence or absence of MPL expressing cells comprising contacting a sample from a subject with a fusion protein comprising scFv specific to MPL for a sufficient amount of time to allow the MPL-specific fusion protein to bind MPL expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of presence of MPL expressing cells. In another embodiment, an increase in luciferase activity relative to reference value is indicative of presence of MPL expressing cells. In one

embodiment, the method further comprises selecting a therapy if presence of MPL expressing cells in the sample is determined. In exemplary embodiments, the MPL specific fusion protein comprises GGS linker and NLuc reporter as described in any one or more of SEQ ID NOs: 1044-1051 or 2244-2251 (**Table 15**).

[00173] In some embodiments, provided herein are methods for detecting presence or absence of PDL1 expressing cells comprising contacting a sample from a subject with a fusion protein comprising scFv specific to PDL1 for a sufficient amount of time to allow the PDL1 -specific fusion protein to bind PDL1 expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of presence of PDL1 expressing cells. In another embodiment, an increase in luciferase activity relative to reference value is indicative of presence of PDL1 expressing cells. In one embodiment, the method further comprises selecting a therapy if presence of PDL1 expressing cells in the sample is determined. In exemplary embodiments, the PDL1 specific fusion protein comprises GGS linker and NLuc reporter as described in any one or more of SEQ ID NOs: 1062-1064 or 2262-2264 (**Table 15**).

[00174] In some embodiments, provided herein are methods for detecting presence or absence of cells expressing targeting antigens described in **Table 15**. The methods comprise contacting a sample from a subject with a fusion protein comprising a scFv, a camelid VHH, a non-immunoglobulin antigen binding scaffold, a cytokine or a receptor specific to the target antigens, for a sufficient amount of time to allow the fusion protein to bind the target antigen expressed on the cells in the sample wherein the fusions proteins are any one or more of the fusion proteins described in **Table 15**. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of presence of targeting antigen expressing cells. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of presence of target antigen expressing cells. In one

embodiment, the method further comprises selecting a therapy if presence of the target antigen expressing cells in the sample is determined. In exemplary embodiments, the antigen specific fusion protein comprises GGS linker and Luc reporter as described in **Table 15**.

[00175] In exemplary embodiments, the sample and the fusion protein are incubated for any of 1min, 5mins, 10mins, 20mins, 30mins, 1 hour, 3 hours, 5 hours, 10 hours, 15 hours, 20 hours, 24 hour or combinations thereof. An appropriate length of time will be apparent to a person of skill in the art.

[00176] In exemplary embodiments, repeated washing comprises washing the sample-fusion protein mixture 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 times or more with an appropriate buffer. Appropriate number of washings of the mixture will be apparent to a person of skill in the art.

[00177] In exemplary embodiments, buffers include but are not limited to PBS, PBS with 2% BSA, PBS with 10% BSA, normal saline and RPMI media with 10% FCS. Appropriate buffers for use with the instant invention will be apparent to a person of skill in the art.

Detection of cells expressing chimeric antigen receptors

[00178] Provided herein are methods for detecting the presence or absence of expression of an artificial receptor or synthetic receptor in a sample obtained from a subject. Non-limiting exemplary embodiments of artificial or synthetic receptors include a chimeric antigen receptor, a chimeric T cell receptor, a transgenic (exogenous) TCR, and a synthetic notch receptor. The methods include contacting a fusion protein comprising an antigen recognition domain fused to a reporter and assaying the activity of the reporter in the presence of a reporter-specific substrate. In one embodiment, the presence of reporter activity is indicative of presence of CAR expressing cells. In another embodiment, an increase in reporter activity relative to a reference value is indicative of presence of CAR expressing cells in the sample. In some embodiments, the antigen recognition domain comprises of, consists of, or consists essentially of the antigen that is targeted by the CAR. In some embodiment, the antigen recognition domain comprises of, consists of, or consists essentially of a protein that binds to the antigen specific domain of CAR. In an exemplary embodiment, the antigen recognition domain comprises of, consists of, or consists essentially

of the antigen that is targeted by the CAR. In an exemplary embodiment, the antigen recognition domain comprises of, consists of, or consists essentially of Protein L. In an exemplary embodiment, the antigen recognition domain comprises of, consists of, or consists essentially of an antibody or an antibody fragment (e.g., a scFv) that binds to any one or more of the components of the CAR. In some embodiments, the reporter is a luciferase as described herein. In some embodiments, the CARs are expressed by the immune effector cells (T cells, NK cells) in the sample. The methods further comprise removing the unbound fusion protein prior to assaying for reporter activity using reporter-specific substrates. In some embodiments, the reporter is a non-secretory form of a luciferase. In some embodiments, the non-secretory form of luciferase is obtained from copepods, deep sea shrimp or homologs or orthologs thereof or mutants or derivatives thereof. In exemplary embodiments, the copepods are selected from the group consisting of any one or more of *Gaussia princeps*, *Pleuromamma abdominalis*, *Metridia pacifica*, *Metridia curticauda*, *Metridia asymmetrica*, *Metridia okhotensis*, *Metridia longa*, *Lucicutia ovaliformis*, *Heterorhabdus tanneri*, and *Pleuromamma scutellata*. In further exemplary embodiments, the luciferase is any one or more of GLuc, NLuc, MLuc7, PaLuc1, PaLuc2, MpLuc1, MoLuc1, MoLuc2, MLuc39, HtLuc, HtLuc2, PsLuc1, H-tanneri, *Lucicutia ovaliformis* LoLuc, LoLuc1-3, *Renilla* or TLuc (TurboLuc16) or derivatives thereof. In some embodiments, the reporter activity is assayed by exposing the target cells to a luciferase-specific substrate. In one embodiment, the luciferase-specific substrate is coelenterazine or a derivative thereof. In another embodiment, the luciferase-specific substrate is imidazopyrazinone substrate (furimazine) or a derivative thereof.

[00179] Provided herein are methods for detecting the presence or absence of expression of chimeric antigen receptors in a sample obtained from a subject. The methods include providing a fusion protein comprising an antigen recognition domain fused to a reporter; providing a sample from the subject; exposing the sample to the fusion protein; and assaying the activity of the reporter in the presence of a reporter-specific substrate. In one embodiment, the presence of reporter activity is indicative of presence of CAR expressing cells. In another embodiment, an increase in reporter activity relative to a reference value is indicative of presence of CAR expressing cells in the sample. In some embodiments, the reporter is a luciferase as described herein. In some embodiments, the CARs are expressed

by the immune effector cells (T cells, NK cells) in the sample. The methods further comprise removing the unbound fusion protein prior to assaying for reporter activity using reporter-specific substrates. In some embodiments, the reporter is a non-secretory form of a luciferase. In some embodiments, the non-secretory form of luciferase is obtained from copepods, deep sea shrimp or homologs or orthologs thereof or mutants or derivatives thereof. In exemplary embodiments, the copepods are selected from the group consisting of any one or more of *Gaussia princeps*, *Pleuromamma abdominalis*, *Metridia pacifica*, *Metridia curticauda*, *Metridia asymmetrica*, *Metridia okhotensis*, *Metridia longa*, *Lucicutia ovaliformis*, *Heterorhabdus tanneri*, and *Pleuromamma scutellata*. In further exemplary embodiments, the luciferase is any one or more of GLuc, NLuc, MLuc7, PaLuc1, PaLuc2, MpLuc1, MoLuc1, MoLuc2, MLuc39, HtLuc, HtLuc2, PsLuc1, H-tanneri, *Lucicutia ovaliformis* LoLuc, LoLuc1-3, Renilla or TLuc (TurboLuc16) or derivatives thereof. In some embodiments, the reporter activity is assayed by exposing the target cells to a luciferase-specific substrate. In one embodiment, the luciferase-specific substrate is coelentraxine or a derivative thereof. In another embodiment, the luciferase-specific substrate is imidazopyrazinone substrate (furimazine) or a derivative thereof.

[00180] In various embodiments of the methods for detecting expression of chimeric antigen receptors, the antigen recognition domain comprises the extracellular domain (ECD) of the antigen of interest. For example, a fusion protein comprising an ECD of an antigen binds the scFv of a CAR that is specific to the ECD. In an exemplary embodiment, the fusion protein comprises an ECD of CD19 and the fusion protein binds to a CD19 specific CAR.

[00181] In various embodiments, the extracellular domains in the fusion protein for detection of CAR expressing cells may be the extracellular domains of disease specific antigens including but are not limited to 4-1BB, 5T4, adenocarcinoma antigen, alpha-fetoprotein, BAFF, B-lymphoma cell, C242 antigen, CA-125, carbonic anhydrase 9 (CA-IX), C-MET, CCR4, CD152, CD19, CD20, CD200, CD22, CD221, CD23 (IgE receptor), CD28, CD30 (TNFRSF8), CD33, CD4, CD40, CD44 v6, CD51, CD52, CD56, CD74, CD80, CEA, CNT0888, CTLA-4, DR5, EGFR, EpCAM, CD3, FAP, fibronectin extra domain-B, folate receptor 1, GD2, GD3 ganglioside, glycoprotein 75, GPNMB, HER2/neu, HGF, human scatter factor receptor kinase, IGF-1 receptor, IGF-I, IgG1, L1-CAM, IL-13, IL-6, insulin-like growth factor I receptor, integrin $\alpha 5\beta 1$, integrin $\alpha v\beta 3$, MORAb-009, MS4A1, MUC1,

mucin CanAg, N-glycolylneuraminic acid, NPC-1C, PDGF-R α , PDL192, phosphatidylserine, prostatic carcinoma cells, RANKL, RON, ROR1, SCH 900105, SDC1, SLAMF7, thrombopoietin receptor, TAG-72, tenascin C, TGF beta 2, TGF- β , TRAIL-R1, TRAIL-R2, tumor antigen CTAA16.88, VEGF-A, VEGFR-1, VEGFR2 or vimentin. Other antigens specific for cancer will be apparent to those of skill in the art and may be used in connection with alternate embodiments of the invention.

[00182] In some embodiments, the fusion proteins further comprise a tag. In exemplary embodiments, that tag is tag is any one or more of chitin binding protein (CBP), glutathione-S-transferase (GST), polyhistidine (His) tag, FLAG tag, HA tag, Myc tag, V5 tag, AcV5 tag, Myristoylation (Myr) tag or a combination thereof.

[00183] In some embodiments, the luciferase activity is measured by any one or more of methods for measuring light production such as a luminometer, x-ray films, microscopy, or combinations thereof.

[00184] In exemplary embodiments of the methods for detecting expression of chimeric antigen receptors, extracellular domains of antigens for use in detecting CARs are described in **Table 9**.

[00185] In some embodiments, provided herein are methods for detecting presence or absence of expression of a CD19 CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of CD19 fused to luciferase, for a sufficient amount of time to allow the CD19 ECD fusion protein to bind to CD19 CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of CD19 CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of CD19 CAR in the sample. In exemplary embodiments, the CD19 ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID Nos 1170-1176 or 2370-2376 (**Table 16**).

[00186] In some embodiments, provided herein are methods for detecting presence or absence of expression of a MPL CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of MPL fused to luciferase, for a sufficient amount of time to allow the MPL ECD fusion protein to bind to MPL CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of MPL CAR sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of MPL CAR in the sample. In exemplary embodiments, the MPL ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1169 or 2369 (**Table 16**).

[00187] In some embodiments, provided herein are methods for detecting presence or absence of expression of a CD33 CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of CD33 fused to luciferase, for a sufficient amount of time to allow the CD33 ECD fusion protein to bind to CD33 CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of CD33 CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of CD33 CAR in the sample. In exemplary embodiments, the CD33 ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1177-1182 or 2377-2382 (**Table 16**).

[00188] In some embodiments, provided herein are methods for detecting presence or absence of expression of a CD138 CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of CD138 fused to luciferase, for a sufficient amount of time to allow the CD138 ECD fusion protein to bind to CD138 CAR expressing cells in the sample. In some embodiments, the method

further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of CD138 CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of CD138 CAR in the sample. In exemplary embodiments, the CD138 ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1183-1187 or 2383-2387 (**Table 16**).

[00189] In some embodiments, provided herein are methods for detecting presence or absence of expression of a CD123 CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of CD123 fused to luciferase, for a sufficient amount of time to allow the CD123 ECD fusion protein to bind to CD123 CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of CD123 CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of CD123 CAR in the sample. In exemplary embodiments, the CD123 ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1188 or 2388 (**Table 16**).

[00190] In some embodiments, provided herein are methods for detecting presence or absence of expression of a CDH1 CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of CDH1 fused to luciferase, for a sufficient amount of time to allow the CDH1 ECD fusion protein to bind to CDH1 CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of CDH1 CAR in the sample. In another embodiment, an increase in luciferase activity relative

to a reference value is indicative of expression of CDH1 CAR in the sample. In exemplary embodiments, the CDH1 ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1189 or 2389 (**Table 16**).

[00191] In some embodiments, provided herein are methods for detecting presence or absence of expression of a CD200R CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of CD200R fused to luciferase, for a sufficient amount of time to allow the CD200R ECD fusion protein to bind to CD200R CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of CD200R in the CAR sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of CD200R CAR in the sample. In exemplary embodiments, the CD200R ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1190 or 2390 (**Table 16**).

[00192] In some embodiments, provided herein are methods for detecting presence or absence of expression of a GPNMB CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of GPNMB fused to luciferase, for a sufficient amount of time to allow the GPNMB ECD fusion protein to bind to GPNMB CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of GPNMB in the CAR sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of GPNMB CAR in the sample. In exemplary embodiments, the GPNMB ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1191 or 2391 (**Table 16**).

[00193] In some embodiments, provided herein are methods for detecting presence or absence of expression of a PTK7 CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of PTK7 fused to luciferase, for a sufficient amount of time to allow the PTK7 ECD fusion protein to bind to PTK7 CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of PTK7 CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of PTK7 CAR in the sample. In exemplary embodiments, the PTK7 ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1192 or 2392 (**Table 16**).

[00194] In some embodiments, provided herein are methods for detecting presence or absence of expression of a CD34 CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of CD34 fused to luciferase, for a sufficient amount of time to allow the CD34 ECD fusion protein to bind to CD34 CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of CD34 CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of CD34 CAR in the sample. In exemplary embodiments, the CD34 ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1193 or 2393 (**Table 16**).

[00195] In some embodiments, provided herein are methods for detecting presence or absence of expression of a EpCAM CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of EpCAM fused to luciferase, for a sufficient amount of time to allow the EpCAM ECD fusion protein to bind to EpCAM CAR expressing cells in the sample. In some embodiments, the

method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of EpCAM in the CAR sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of EpCAM CAR in the sample. In exemplary embodiments, the EpCAM ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1194 or 2394 (**Table 16**).

[00196] In some embodiments, provided herein are methods for detecting presence or absence of expression of a CLEC12A CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of CLEC12A fused to luciferase, for a sufficient amount of time to allow the CLEC12A ECD fusion protein to bind to CLEC12A CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of CLEC12A in the CAR sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of CLEC12A CAR in the sample. In exemplary embodiments, the CLEC12A ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1195 or 2395 (**Table 16**).

[00197] In some embodiments, provided herein are methods for detecting presence or absence of expression of a CD20 CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of CD20 fused to luciferase, for a sufficient amount of time to allow the CD20 ECD fusion protein to bind to CD20 CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of

CD20 CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of CD20 CAR in the sample. In exemplary embodiments, the CD20 ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1196-1197 or 2396-2397 (**Table 16**).

[00198] In some embodiments, provided herein are methods for detecting presence or absence of expression of a CD22 CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of CD22 fused to luciferase, for a sufficient amount of time to allow the CD22 ECD fusion protein to bind to CD22 CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of CD22 CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of CD22 CAR in the sample. In exemplary embodiments, the CD22 ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1198 or 2398 (**Table 16**).

[00199] In some embodiments, provided herein are methods for detecting presence or absence of expression of a TSHR CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of TSHR fused to luciferase, for a sufficient amount of time to allow the TSHR ECD fusion protein to bind to TSHR CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of TSHR CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of TSHR CAR in the sample. In exemplary embodiments, the TSHR ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1199 or 2399 (**Table 16**).

[00200] In some embodiments, provided herein are methods for detecting presence or absence of expression of a EGFRviii CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of EGFRviii fused to luciferase, for a sufficient amount of time to allow the EGFRviii ECD fusion protein to bind to EGFRviii CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of EGFRviii CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of EGFRviii CAR in the sample. In exemplary embodiments, the EGFRviii ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1200 or 2400 (**Table 16**).

[00201] In some embodiments, provided herein are methods for detecting presence or absence of expression of a BCMA CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of BCMA fused to luciferase, for a sufficient amount of time to allow the BCMA ECD fusion protein to bind to BCMA CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of BCMA CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of BCMA CAR in the sample. In exemplary embodiments, the BCMA ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1201 or 2401 (**Table 16**).

[00202] In some embodiments, provided herein are methods for detecting presence or absence of expression of a SLAMF7 CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of SLAMF7 fused to luciferase, for a sufficient amount of time to allow the EGFRviii ECD

fusion protein to bind to SLAMF7 CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of SLAMF7 CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of SLAMF7 CAR in the sample. In exemplary embodiments, the SLAMF7 ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1202 or 2402 (**Table 16**).

[00203] In some embodiments, provided herein are methods for detecting presence or absence of expression of a PD1 CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of PD1 fused to luciferase, for a sufficient amount of time to allow the PD1 ECD fusion protein to bind to PD1 CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of PD1 CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of PD1 CAR in the sample. In exemplary embodiments, the PD1 ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1203 or 2403 (**Table 16**).

[00204] In some embodiments, provided herein are methods for detecting presence or absence of expression of a CTLA4 CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of CTLA4 fused to luciferase, for a sufficient amount of time to allow the CTLA4 ECD fusion protein to bind to CTLA4 CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase

activity. In one embodiment, presence of luciferase activity relative to a reference value is indicative of expression of CTLA4 CAR in the sample. In another embodiment, an increase in luciferase activity is indicative of expression of CTLA4 CAR in the sample. In exemplary embodiments, the CTLA4 ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1204 or 2404 (**Table 16**).

[00205] In some embodiments, provided herein are methods for detecting presence or absence of expression of a NKG2D CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising ECD of NKG2D fused to luciferase, for a sufficient amount of time to allow the NKG2D ECD fusion protein to bind to NKG2D CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of NKG2D CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of NKG2D CAR in the sample. In exemplary embodiments, the NKG2D ECD fusion protein comprises GGS linker and Luc reporter as described in any one or more of SEQ ID NO: 1205 or 2405 (**Table 16**).

[00206] In some embodiments, provided herein are methods for detecting presence or absence of expression of a kappa light chain of a CAR in a sample obtained from a subject. The method includes contacting a sample from a subject with a fusion protein comprising Protein L fused to luciferase, for a sufficient amount of time to allow the Protein L fusion protein to bind to the kappa light chain of CAR expressing cells in the sample. In some embodiments, the method further comprises removing unbound fusion protein, for example by repeatedly washing the sample/fusion protein mixture with a buffer. The method further comprises contacting the sample/fusion protein mixture with a luciferase specific substrate and detecting the luciferase activity. In one embodiment, presence of luciferase activity is indicative of expression of a CAR in the sample. In another embodiment, an increase in luciferase activity relative to a reference value is indicative of expression of a CAR in the sample. In exemplary embodiments, the Protein L fusion protein comprises GGS linker and

NLuc reporter as described in any one or more of SEQ ID NO: 1206-1207 or 2406-2407 (Table 16).

[00207] In exemplary embodiments, repeated washing comprises washing the sample-fusion protein mixture 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 times or more with an appropriate buffer. Appropriate number of washings of the mixture will be apparent to a person of skill in the art.

[00208] In exemplary embodiments, buffers include but are not limited to PBS, normal saline, RPMI, PBS with 5% BSA, PBS with 10% BSA. Appropriate buffers for use with the instant invention will be apparent to a person of skill in the art.

Reference Value

[00209] In some embodiments, the reference value is the reporter activity (for example, luciferase activity as described herein) in any one or more of (i) control cells or samples that do not express the target antigen; (ii) control cells or samples that express reporter but are not treated with the test agent(s); (iii) controls cells or samples that are not treated with the substrate for the reporter; (iv) control cells or samples that do express the target antigen but are treated with a fusion protein directed against an antigen that is not expressed on the target cells; (v) a combination thereof.

[00210] In some embodiments, the reference value is the reporter activity (for example, luciferase activity as described herein) in any one or more of (i) cells that do not express the CAR, or an exogenous T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor; (ii) cells that express the CAR, or an exogenous T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor but are treated with fusion protein which is not targeted by the CAR, or an exogenous T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor; (iii) cells that are not treated with the substrate for the reporter.

[00211] In some embodiments, the reporter activity (for example, luciferase activity as described herein) is increased relative to the reference value by at least or about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100%.

[00212] In some embodiments, the reporter activity (for example, luciferase activity as described herein) is increased relative to the reference value by at least or about 1-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold, 15-fold, 20-fold, 25-fold, 30-fold, 35-fold, 40-fold, 45-fold, 50-fold, 55-fold, 60-fold, 65-fold, 70-fold, 75-fold, 80-fold, 85-fold, 90-fold, 95-fold, 100-fold, 1000-fold, 10,000-fold or a combination thereof.

[00213] In some embodiments, the detection methods described herein may be used to screen patients for CAR therapies. For example, in patients who have been diagnosed with cancer and are candidates for CAR therapies, the methods described herein may be used to determine the presence or absence and/or the amount of antigen expressed in a sample from the patient, wherein the antigen is targeted by the CAR. If the antigen is detected in the patient, the patient may be a candidate for CAR therapy.

[00214] In some embodiments, the detection methods described herein may be used to monitor the persistence of CARs in patients who have received CAR therapies. For example, in patients who have received CAR therapies, the methods described herein may be used to determine the presence or absence of CAR T cells in a sample from the patient at different time intervals following the administration of CAR-T cells. If the CAR-T cells are not detected in the patient, the patient may be at higher risk of relapse.

[00215] In some embodiments, the detection methods described herein may be used as a quality control tool to monitor the quality of different lots of CAR-T cells during the manufacturing of CAR-T cells. For example, different lots of CD19-directed CAR-T cells can be compared for their relative binding to CD19-ECD-NLuc fusion protein (SEQ ID NO: 1170) as compared to a reference lot. A key advantage of the detection method of the instant invention is that it not only measures the expression of the CAR protein on the surface of immune cells, but it also measures the ability of the CAR protein to bind to the target antigen.

[00216] There are many variables in the design and manufacturing of CARs. Many different CAR constructs can be generated against a particular antigen. These constructs differ in their antigen specific domains, which are generally derived from different antibodies. There exists a need for a fast, economical, sensitive and specific assay to select the scFv fragments for incorporation into a CAR construct. In some embodiments, the detection methods described herein may be used to select the best candidate scFv for

incorporation into a CAR construct. For example, different scFv fusion proteins targeting a candidate CAR antigen (e.g., CD19) can be generated and tested rapidly for their relative expression and ability to bind to a panel of CD19+ve and CD19-ve cell lines. The scFv fragment(s) showing the highest expression, binding affinity and specificity would be candidate for incorporation into a CAR. In some embodiments, the detection methods described herein may be used to select the best candidate CAR construct for further development. For example, a number of candidate CD19-specific CAR constructs, which differ base on their antigen specific domain, hinge domain, transmembrane domain, costimulatory domain, activation domain etc., can be rapidly compared using the current method for their ability to bind to FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04) [SEQ ID NO:1170] fusion protein, which would provide a relative measure of the ability of the CAR to bind to CD19 on the target cells.

[00217] In some embodiments, the detection methods described herein may be used in antibody engineering. For example, antibody engineering, such as affinity maturation and humanization, rely on expression and testing of several different scFv candidate constructs. These constructs can be constructed as luciferase-fusion proteins, which will allow rapid screening of the different constructs for relative expression and binding affinity to their target antigens, including binding affinity in cell-based assays.

Methods for detecting luciferase activity

[00218] In various embodiments, luciferase activity described herein is measured by any one or more of methods for measuring light production such as a luminometer, X-ray films, microscopy, or combinations thereof. The nucleic acid and amino acid sequences of the fusion proteins described in the above section are presented in the corresponding sequence sections.

Methods for expression of Luc fusion proteins

[00219] The host cell line utilized for expression of the Luc fusion proteins of the present invention can be of mammalian or non-mammalian origin. Those skilled in the art are credited with ability to preferentially determine particular host cell lines which are best suited for the desired gene product to be expressed therein. In one embodiment, the host cell line

utilized for generation of the Luc fusion proteins of the invention is mammalian in origin. Exemplary host cell lines include, but are not limited to, DG44 and DUXBII (Chinese Hamster Ovary lines, DHFR minus), HELA (human cervical carcinoma), CVI (monkey kidney line), COS (a derivative of CVI with SV40 T antigen), R1610 (Chinese hamster fibroblast) BALBC/3T3 (mouse fibroblast), HAK (hamster kidney line), SP2/0 (mouse myeloma), P3×63-Ag3.653 (mouse myeloma), BFA-lclBPT (bovine endothelial cells), RAJI (human lymphocyte), 293FT and 293 (human kidney). Host cell lines are typically available from commercial services, the American Tissue Culture Collection or from published literature. In another embodiment, the host cell line utilized for generation of the Luc fusion proteins of the invention is of insect origin. In another embodiment, the host cells utilized for generation of the Luc fusion proteins of the invention are bacterial in origin. In another embodiment, the host cells utilized for generation of the Luc fusion proteins of the invention are of yeast origin. Those skilled in the art are credited with ability to preferentially determine particular promoters and vectors which are best suited for the expression of a desired gene product in a particular host cell line. Exemplary vector for expression of the Luc fusion protein of the invention is pLenti-EF1a (SEQ ID NO: 842). Vectors for expression of proteins in other host cell lines are typically available from commercial sources, such as Thermofisher, Novagen and Sigma. In various embodiments, the fusion proteins described herein may be expressed in bacterial cells, yeasts or insects. The optimum vectors and promoters to be used for expression of the fusions proteins described herein and for the methods described herein will be apparent to a person of skill in the art.

Therapies

[00220] As described herein, in various embodiments, if an antigen of interest or a CAR of interest is detected, the methods further comprise selecting a therapy. Examples of therapy include, but are not limited to, active surveillance, observation, surgical intervention, chemotherapy, immunotherapy, radiation therapy (such as external beam radiation, stereotactic radiosurgery (gamma knife), and fractionated stereotactic radiotherapy (FSR)), focal therapy, systemic therapy, vaccine therapies, viral therapies, molecular targeted therapies, or combinations thereof.

[00221] In some embodiments, therapies include surgery, chemotherapy, radiation, immunosuppressive agents, such as cyclosporin, azathioprine, methotrexate, mycophenolate, and FK506, antibodies, or other immunoablative agents such as CAMPATH, anti-CD3 antibodies or other antibody therapies, cytoxin, fludarabine, cyclosporin, FK506, rapamycin, mycophenolic acid, steroids, FR901228, cytokines, and irradiation, peptide vaccine, such as that described in Izumoto et al. 2008 J Neurosurg 108:963-971. Exemplary chemotherapeutic agents include an anthracycline (e.g., doxorubicin (e.g., liposomal doxorubicin)), a vinca alkaloid (e.g., vinblastine, vincristine, vindesine, vinorelbine), an alkylating agent (e.g., cyclophosphamide, decarbazine, melphalan, ifosfamide, temozolomide), an immune cell antibody (e.g., alemtuzamab, gemtuzumab, rituximab, ofatumumab, tositumomab, brentuximab), an antimetabolite (including, e.g., folic acid antagonists, pyrimidine analogs, purine analogs and adenosine deaminase inhibitors (e.g., fludarabine)), an mTOR inhibitor, a TNFR glucocorticoid induced TNFR related protein (GITR) agonist, a proteasome inhibitor (e.g., aclacinomycin A, gliotoxin or bortezomib), an immunomodulator such as thalidomide or a thalidomide derivative (e.g., lenalidomide).

[00222] In some embodiments, therapies include bone marrow transplantation, T cell ablative therapy using chemotherapy agents such as, fludarabine, external-beam radiation therapy (XRT), cyclophosphamide, and/or antibodies such as OKT3 or CAMPATH.

[00223] Optimum dosages, frequencies of administration and modes of administration will be apparent to a person of skill in the art.

Kits

[00224] Kits to practice the invention are also provided. For example, kits for expressing one or more luciferase proteins of the invention. The kits may include a nucleic acid molecule or a polypeptide molecule encoding the luciferase fusion constructs along with a method to introduce the nucleic acid into an appropriate host cell to produce the protein. The kit may include a vector comprising a nucleic acid encoding the fusion protein of the invention and chemicals, such as lipofectamine, to introduce the vector into appropriate host cells. The kit may contain components for the expression, purification and detection of the luciferase fusion protein. For example, the kit may contain buffer and reagents for measuring the activity luciferases. More than one of the disclosed vector can be included in the kit. The kit

can include a container and a label or package insert on or associated with the container. The kit may include instructions for storage and shipment of its contents.

[00225] Suitable containers include, for example, bottles, vials, syringes, etc. The containers may be formed from a variety of materials such as glass or plastic. The container typically holds a composition including one or more of the nucleic acid molecules, viruses, vectors and proteins. A label or package insert indicates that the composition is used for diagnosing a particular condition. The label or package insert typically will further include instructions for use of a disclosed nucleic acid molecules, proteins, for example, in a method of detecting the presence of a CAR. The package insert typically includes instructions customarily included in commercial packages of diagnostic products that contain information about the composition and health hazards. The instructional materials may be written, in an electronic form (such as a computer diskette or compact disk) or may be visual (such as video files). The kits may also include additional components to facilitate the particular application for which the kit is designed. Thus, for example, the kit may additionally contain means for measuring the expression of CAR on T cells or of determining the number or percentage of T cells that express the CAR or of determining the functionality of CAR-T cells. The kits may additionally include buffers and other reagents routinely used for the practice of a particular method. Such kits and appropriate contents are well known to those of skill in the art.

[00226] Table 3: Exemplary signal peptides

NAME	SEQ ID (DNA)	SEQ ID (PRT)
CD8_Signal_Peptide	1	1781
CD8_Signal_Peptide	2	1781
CD8_Signal_Peptide	3	1781
CD8-SIGNAL-PEPTIDE	4	1781
IgH_Signal_Peptide	5	1782
IgH_Signal_Peptide	6	1782
IgH_Signal-Peptide	7	1782
IgH_Signal Peptide	8	1782
Amyloid-158-158-Signal-peptide	9	1783
hCD19 Signal Peptide	10	1784

[00227] Table 4: Exemplary VL fragments

Target	SEQ ID (DNA)	SEQ ID (PRT)	NAME
AFP/MHC class I complex	11	1785	AFP-61-vL
AFP/MHC class I complex	12	1786	AFP-76-vL
AFP/MHC class I complex	13	1787	AFP-79-vL
ALK	14	1788	Alk-48-vL
ALK	15	1789	Alk-58-vL
Amyloid	16	1790	Amyloid-158-vL
BCMA	17	1791	BCMA-ET-40-vL
BCMA	18	1792	BCMA-ET-54-vL
BCMA	19	1793	BCMA-huC12A3-vL
BCMA	20	1794	BCMA-J6M0-vL
BCMA	21	1795	BCMA-ET-03-vL
BCMA	22	1796	BCMA-huC11.D5.3L1H3-vL
BCMA	23	1797	BCMA-huC13-F12-vL
CCR4	24	1798	CCR4-humAb1567-vL
CD123	25	1799	CD123-CSL362-vL
CD123	26	1800	CD123-1172-vL
CD123	27	1801	CD123-DART-1-vL
CD123	28	1802	CD123-DART-2-vL
CD123	29	1803	CD123-I3RB18-vL
CD123	30	1804	CD123-hu3E3-vL
CD123	31	1805	CD123-9F6-vL
CD123	32	1806	CD123-I3RB2-vL
CD123	33	1807	CD123-1176-vL
CD123	34	1808	CD123-8B11-vL
CD123	35	1809	CD123-2B8-vL
CD123	36	1810	CD123-9D7-vL
CD123	37	1811	CD123-3B10-vL
CD138	38	1812	CD138-vL
CD179b	39	1813	CD179b-vL
CD19	40	1814	CD19-4G7-vL
CD19	41	1815	CD19Bu12-vL
CD19	42	1816	CD19MM-vL
CD19	43	1817	FMC63-vL
CD19	44	1818	FMC63-[2]-vL
CD19	45	1819	FMC63-[3]-vL
CD19	46	1820	huFMC63-11-vL
CD19	47	1821	CD19-MEDI-3649-vL
CD19	48	1822	CD19-Medrex-24D1-vL
CD19	49	1823	CD19-MOR0028-vL
CD19	50	1824	CD19-HD37-H2L1-vL
CD19	51	1825	CD19-huBly3-vL
CD19	52	1826	CD19-huSJ25C1-vL
CD19	53	1827	CD19-hB4-vL
CD19	54	1828	CD19-hu-mROO5-1-vL
CD19	55	1829	CD19-hA19-vL

Target	SEQ ID (DNA)	SEQ ID (PRT)	NAME
CD20	56	1830	CD20-2F2-vL
CD20	57	1831	CD20-GA101-vL
CD20	58	1832	CD20-Leu16-vL
CD20	59	1833	CD20-11B8-vL
CD20	60	1834	CD20-2C6-vL
CD20	61	1835	CD20-2H7-vL
CD20	62	1836	CD20-hA20-vL
CD20	63	1837	CD20-BM-CA-1925-v4-vL
CD20	64	1838	CD20-Ubli-v4-vL
CD20	65	1839	CD20-2H7-vL
CD20	66	1840	CD20-h1F5-vL
CD20	67	1841	CD20-7D8-vL
CD20	68	1842	CD20-AME-33-vL
CD22	69	1843	CD22-h10F4-vL
CD22	70	1844	CD22-H22Rhov2ACDRKA-vL
CD22	71	1845	CD22m971-vL
CD276	72	1846	CD276-17-vL
CD30	73	1847	CD30-5F11-vL
CD30	74	1848	CD30-Ac10-vL
CD32	75	1849	CD32-Med9-vL
CD324	76	1850	CD324-hSC10-17-vL
CD324	77	1851	CD324-SC10-6-vL
CD33	78	1852	CD33-huMyc9-vL
CD33	79	1853	CD33-AF5-vL
CD33	80	1854	CD33-Boehr2800308-vL
CD33	81	1855	CD33-Him3-4-vL
CD33	82	1856	CD33-SGNh2H12-vL
CD33	83	1857	CD33-15G15-33-vL
CD33	84	1858	CD33-33H4-vL
CD33	85	1859	CD33-9C3-2-vL
CD34	86	1860	CD34-hu4C7-[2]-vL
CD34	87	1861	CD34-hu4C7-vL
CD44v6	88	1862	CD44v6-Biwa8-vL
CD5	89	1863	CD5-18-vL
CD5	90	1864	CD5-9-vL
CD70	91	1865	CD70-h1F6-vL
CD79b	92	1866	CD79b-2F2-vL
CD79b	93	1867	huMA79bv28-vL
CD99	94	1868	CD99-hu12E7-vL
CDH17	95	1869	CDH17-PTA001A4-vL
CDH19	96	1870	CDH19-16A4-vL
CDH6	97	1871	CDH6-NOV710-vL
CDH6	98	1872	CDH6-NOV712-vL
CLEC5A	99	1873	CLEC5A-3E12A2-vL
CLEC5A	100	1874	CLEC5A-8H8F5-vL
CLL1	101	1875	CLL1-M26-vL
CLL1	102	1876	CLL1-M32-vL

Target	SEQ ID (DNA)	SEQ ID (PRT)	NAME
CD123	103	1877	CLL1-21C9-L2H3-vL
CLL1	104	1878	CLL1-6E7L4H1e-vL
CLL1	105	1879	CLL1-hu1075-v1-vL
CLL1	106	1880	CLL1-hu1075-v2-vL
CMVpp65/MHC class I complex	107	1881	CMVpp65-F5-vL
CS1	108	1882	huLuc63-vL
CS1	109	1883	HuLuc64-[2]-vL
CS1	110	1884	HuLuc64-vL
CS1	111	1885	Luc90-vL
CS1	112	1886	CS1-PDL241-vL
CS1	113	1887	CS1-Hu27A-vL
CS1	114	1888	CS1-ScHu34C3-vL
CS1	115	1889	CS1-Hu31-D2-vL
CS1	116	1890	CS1-Luc34-vL
CS1	117	1891	CS1-LucX2-vL
CSF2RA	118	1892	CSF2RA-Ab1-vL
CSF2RA	119	1893	CSF2RA-Ab6-vL
DLL3	120	1894	DLL3-hSC16-13-vL
DLL3	121	1895	DLL3-hSC16-56-vL
EBNA3c//MHC class I complex	122	1896	EBNA3c-315-vL
EGFR	123	1897	Cetuximab-vL
EGFR	124	1898	Nimotuzumab-vL
EGFRviii	125	1899	EGFRviii-139-vL
EGFRviii	126	1900	EGFRviii-2173-vL
EpCam1	127	1901	EpCam1-D5K5-vL
EpCam1	128	1902	Epcam1-MM1-vL
FITC	129	1903	FITC-vL
FITC	130	1904	FITC-4M-53-vL
FITC	131	1905	FITC-E2-vL
FLT3	132	1906	FLT3-NC7-vL
HIV1-envelop glycoprotein	133	1907	HIV1-N6-vL
FR1 (Folate Receptor a)	134	1908	FR1-huMov19-vL
GAD	135	1909	GAD-G3H8-vL
GD2	136	1910	GD2-hu14-18-vL
GD2	137	1911	GD2-hu3F8-vL
GD3	138	1912	GD3-KM-641-vL
GFRa4	139	1913	GFRa4-P4-10-2-vL
GFRa4	140	1914	GFRa4-P4-10-vL
GFRa4	141	1915	GFRAlpha4-P4-6-vL
GM1	142	1916	GM1-5B2-vL
GM1	143	1917	GM1-7E5-vL
gp100//MHC class I complex	144	1918	gp100-G2D12-vL
gp100//MHC class I complex	145	1919	gp100-vL
GPC3	146	1920	GPC3-4E5-vL
gpNMB	147	1921	gpNMB-115-vL
GPRC5D	148	1922	GPRC5D-ET150-18-vL
GPRC5D	149	1923	GPRC5D-ET150-5-vL

Target	SEQ ID (DNA)	SEQ ID (PRT)	NAME
GPRC5D	150	1924	GPRC5D-ET150-1-vL
GPRC5D	151	1925	GPRC5D-ET150-2-vL
Her2	152	1926	Her2-Hu4D5-vL
HIV1-gag (77-85)/MHC class I complex	153	1927	HIV1-3BNC117-vL
HIV1-envelop glycoprotein	154	1928	HIV1-E5-vL
HIV1-envelop glycoprotein	155	1929	HIV1-PGT-128-vL
HIV1-envelop glycoprotein	156	1930	HIV1-VR-C01-vL
HIV1-envelop glycoprotein	157	1931	HIV1-X5-vL
HLA-A2	158	1932	HLA-A2-3PB2-vL
HMW-MAA	159	1933	HMW-MAA-hIND-vL
HPV16-E7/MHC class I complex	160	1934	HPV16-7-8-vL
HPV16-E7/MHC class I complex	161	1935	HPV16-2-vL
HTLV1-TAX/MHC class I complex	162	1936	TAX-T3E3-vL
HTLV1-TAX/MHC class I complex	163	1937	TAX-T3F2-vL
IL11Ra	164	1938	IL11Ra-8E2-vL
IL13Ra2	165	1939	IL13Ra2-hu107-vL
IL13Ra2	166	1940	IL13Ra2-Hu108-vL
IL6R	167	1941	IL6R-M83-vL
Influenza A HA	168	1942	FLU-MEDI-8852-vL
KSHV-gH	169	1943	YC15-vL
KSHV-K8.1	170	1944	4C3-vL
L1CAM	171	1945	L1CAM-9-3-HU3-vL
LAMP1	172	1946	LAMP1-humab1-2-vL
LAMP1	173	1947	LAMP1-Mb4-vL
LewisY	174	1948	LewisY-huS193-vL
Lym1	175	1949	Lym1-vL
Lym2	176	1950	Lym2-vL
MART1/MHC class I complex	177	1951	MART1-CAG10-vL
MART1/MHC class I complex	178	1952	MART1-CLA12-vL
Mesothelin	179	1953	Mesothelin-m912-vL
MPL	180	1954	MPL-111-vL
MPL	181	1955	MPL-161-HL-vL
MPL	182	1956	MPL-161-vL
MPL	183	1957	MPL-175-vL
MPL	184	1958	MPL-178-vL
MPL	185	1959	MPL-huVB22Bw5-vL
MPL	186	1960	MPL-12E10-vL
MPL	187	1961	MPL-AB317-vL
Muc1/MHC class I complex	188	1962	MUC1-D6-M3A1-vL
Muc1/MHC class I complex	189	1963	Muc1-D6-M3B8-vL
Muc16	190	1964	Muc16-4H11-vL
NKG2D	191	1965	NKG2D-MS-vL
NYBR1	192	1966	NYBR1-vL
NY-ESO/MHC class I complex	193	1967	NY-ESO-T1-vL
PD1	194	1968	PD1-4H1-vL
PD1	195	1969	PD1-5C4-vL

Target	SEQ ID (DNA)	SEQ ID (PRT)	NAME
PDL1	196	1970	PDL1-10A5-vL
PDL1	197	1971	PDL1-Atezoli-vL
PDL1	198	1972	PDL1-SP142-vL
PR1/MHC class I complex	199	1973	PR1-vL
PSCA	200	1974	PSCA-Ha14-117-vL
PSCA	201	1975	PSCA-Ha14-121-vL
PSMA	202	1976	PSMA-006-vL
PSMA	203	1977	PSMA-J591-vL
PTK7	204	1978	PTK7-hSC6-23-vL
PTK7	205	1979	PTK7-SC6-10-2-vL
ROR1	206	1980	ROR1-4A5-vL
ROR1	207	1981	ROR1-4C10-vL
SLea	208	1982	SLea-5B1-vL
SLea	209	1983	SLea-7E3-vL
SSEA4	210	1984	SSEA4-vL
TCRB1 (TCR b1 constant chain)	211	1985	TCRB1-E09-vL
TCRB1 (TCR b1 constant chain)	212	1986	TCRB1-Jovi1-vL
TCRB2 (TCR b2 constant chain)	213	1987	TCRB2-CP01-D05-vL
TCRB2 (TCR b2 constant chain)	214	1988	TCRB2-CP01-E05-vL
TCRgamma delta (TCRgd)	215	1989	TCRgd-G5-4-vL
TERT/MHC class I complex	216	1990	TERT-3G3-T865-vL
TERT/MHC class I complex	217	1991	TERT-4A9-T540-vL
TF1 (Tissue Factor)	218	1992	TF1-98-vL
TGFBR2	219	1993	TGFBR2-Ab1-vL
TIM1	220	1994	TIM1-HVCR1-270-2-vL
TIM1	221	1995	Tim1HVCR1-ARD5-vL
TnAg	222	1996	TnAg-vL
Tn-Muc1	223	1997	Tn-Muc1-hu5E5-vL
TROP2	224	1998	TROP2-ARA47-HV3KV3-vL
TROP2	225	1999	TROP2-h7E6-SVG-vL
TSHR	226	2000	TSHR-5C9-vL
TSHR	227	2001	TSHR-K1-70-vL
TSHR	228	2002	TSHR-KB1-vL
TSLRP	229	2003	TSLPR-vL
Tyrosinase/MHC class I complex	230	2004	Tyro-B2-vL
Tyrosinase/MHC class I complex	231	2005	Tyro-Mc1-vL
Tyrosinase/MHC class I complex	232	2006	TA2-vL
VEGFR3	233	2007	VEGFR3-Ab1-vL
WT1/MHC class I complex	234	2008	WT1-Ab13-vL
WT1/MHC class I complex	235	2009	WT1-Ab15-vL
WT1/MHC class I complex	236	2010	WT1-Ab1-vL
WT1/MHC class I complex	237	2011	WT1-Ab5-vL
EBV-gp350	238	2012	EBV-gp350-vL
CDH19	239	2013	CDH19-4B10-vL
Folate Receptor b (FRbeta)	240	2014	FRbeta-m923-vL
LHR	241	2015	LHR-8B7-vL
LHR	242	2016	LHR-5F4-21-vL

Target	SEQ ID (DNA)	SEQ ID (PRT)	NAME
B7H4	243	2017	B7H4-hu22C10-vL
B7H4	244	2018	B7H4-hu1D11-vL
IgE	245	2019	IgE-omalizumab-vL
CD23	246	2020	CD23-p5E8-vL
GCC	247	2021	GCC-5F9-vL
GCC	248	2022	GCC-Ab229-vL
CD200R	249	2023	CD200R-huDx182-vL

[00228] Table 5: Exemplary Linkers

NAME	SEQ ID (DNA)	SEQ ID (PRT)
(GGGGS)x3 LINKER	250	2024
(GGGGS)x3 Linker	251	2024
(Gly4Ser)x3 Linker	252	2024
(Gly4Ser)x3 Linker	253	2024
(GGGGS)x3 Linker	254	2024
(GGGGS)x3 Linker	255	2024
(GGSG)7 Linker	256	2025
(GGSG)7 Linker 2	257	2026
DDAKK linker	258	2027

[00229] Table 6: Exemplary V_H fragments

Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
AFP/MHC class I complex	259	2028	AFP-61-vH
AFP/MHC class I complex	260	2029	AFP-76-vH
AFP/MHC class I complex	261	2030	AFP-79-vH
ALK	262	2031	Alk-48-vH
ALK	263	2032	Alk-58-vH
Amyloid	264	2033	Amyloid-158-vH
BCMA	265	2034	BCMA-ET-40-vH
BCMA	266	2035	BCMA-ET-54-vH
BCMA	267	2036	BCMA-huC12A3-vH
BCMA	268	2037	BCMA-J6M0-vH
BCMA	269	2038	BCMA-ET-03-vH
BCMA	270	2039	BCMA-huC11.D5.3L1H3-vH
BCMA	271	2040	BCMA-huC13-F12-vH
CCR4	272	2041	CCR4-humAb1567-vH
CD123	273	2042	CD123-CSL362-vH
CD123	274	2043	CD123-1172-vH
CD123	275	2044	CD123-DART-1-vH
CD123	276	2045	CD123-DART-2-vH

Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
CD123	277	2046	CD123-13RB18-vH
CD123	278	2047	CD123-hu3E3-vH
CD123	279	2048	CD123-9F6-vH
CD123	280	2049	CD123-I3RB2-vH
CD123	281	2050	CD123-1176-vH
CD123	282	2051	CD123-8B11-vH
CD123	283	2052	CD123-2B8-vH
CD123	284	2053	CD123-9D7-vH
CD123	285	2054	CD123-3B10-vH
CD138	286	2055	CD138-vH
CD179b	287	2056	CD179b-vH
CD19	288	2057	CD19-4G7-vH
CD19	289	2058	CD19Bu12-vH
CD19	290	2059	CD19Bu12-[2]-vH
CD19	291	2060	CD19MM-vH
CD19	292	2061	FMC63-vH
CD19	293	2062	FMC-63-vH
CD19	294	2063	huFMC63-11-vH
CD19	295	2064	CD19-MEDI-3649-vH
CD19	296	2065	CD19-Medrex-24D1-vH
CD19	297	2066	CD19-MOR0028-vH
CD19	298	2067	CD19-HD37-H2L1-vH
CD19	299	2068	CD19-huBly3-vH
CD19	300	2069	CD19-huSJ25C1-vH
CD19	301	2070	CD19-hB4-vH
CD19	302	2071	CD19-hu-mROO5-1-vH
CD19	303	2072	CD19-hA19-vH
CD20	304	2073	CD20-2F2-vH
CD20	305	2074	CD20-GA101-vH
CD20	306	2075	CD20-Leu16-vH
CD20	307	2076	CD20-11B8-vH
CD20	308	2077	CD20-2C6-vH
CD20	309	2078	CD20-2H7-vH
CD20	310	2079	CD20-hA20-vH
CD20	311	2080	CD20-BM-CA-1925-v4-vH
CD20	312	2081	CD20-Ubli-v4-vH
CD20	313	2082	CD20-2H7-vH
CD20	314	2083	CD20-h1F5-vH
CD20	315	2084	CD20-7D8-vH
CD20	316	2085	CD20-AME-33-vH
CD22	317	2086	CD22-h10F4-vH
CD22	318	2087	CD22-H22Rhov2ACDRKA-vH
CD22	319	2088	CD22m971-vH
CD276	320	2089	CD276-17-vH
CD30	321	2090	CD30-5F11-vH
CD30	322	2091	CD30-Ac10-vH
CD32	323	2092	CD32-Med9-vH

Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
CD324	324	2093	CD324-hSC10-17-vH
CD324	325	2094	CD324-SC10-6-vH
CD33	326	2095	CD33-huMyc9-vH
CD33	327	2096	CD33-AF5-vH
CD33	328	2097	CD33-Boehr2800308-vH
CD33	329	2098	CD33-Him3-4-vH
CD33	330	2099	CD33-SGNh2H12-vH
CD33	331	2100	CD33-15G15-33-vH
CD33	332	2101	CD33-33H4-vH
CD33	333	2102	CD33-33H4-vH
CD34	334	2103	CD33-9C3-2-vH
CD34	335	2104	CD34-hu4C7-vH
CD44v6	336	2105	CD44v6-Biwa8-vH
CD5	337	2106	CD5-18-vH
CD5	338	2107	CD5-9-vH
CD70	339	2108	CD70-h1F6-vH
CD79b	340	2109	CD79b-2F2-vH
CD79b	341	2110	huMA79bv28-vH
CD99	342	2111	CD99-hu12E7-vH
CDH17	343	2112	CDH17-PTA001A4-vH
CDH19	344	2113	CDH19-16A4-vH
CDH6	345	2114	CDH6-NOV710-vH
CDH6	346	2115	CDH6-NOV712-vH
CLEC5A	347	2116	CLEC5A-3E12A2-vH
CLEC5A	348	2117	CLEC5A-8H8F5-vH
CLL1	349	2118	CLL1-M26-vH
CLL1	350	2119	CLL1-M32-vH
CD123	351	2120	CLL1-21C9-L2H3-vH
CLL1	352	2121	CLL1-6E7L4H1e-vH
CLL1	353	2122	CLL1-hu1075-v1-vH
CLL1	354	2123	CLL1-hu1075-v2-vH
CMVpp65/MHC class I complex	355	2124	CMVpp65-F5-vH
CS1	356	2125	huLuc63-vH
CS1	357	2126	HuLuc64-vH
CS1	358	2127	Luc90-vH
CS1	359	2128	CS1-PDL241-vH
CS1	360	2129	CS1-Hu27A-vH
CS1	361	2130	CS1-ScHu34C3-vH
CS1	362	2131	CS1-Hu31-D2-vH
CS1	363	2132	CS1-Luc34-vH
CS1	364	2133	CS1-LucX2-vH
CSF2RA	365	2134	CSF2RA-Ab1-vH
CSF2RA	366	2135	CSF2RA-Ab6-vH
DLL3	367	2136	DLL3-hSC16-13-vH
DLL3	368	2137	DLL3-hSC16-56-vH
EBNA3c/MHC class I	369	2138	EBNA3c-315-vH

Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
complex			
EGFR	370	2139	Cetuximab-vH
EGFR	371	2140	Nimotuzumab-vH
EGFRviii	372	2141	EGFRviii-139-vH
EGFRviii	373	2142	EGFRviii-2173-vH
EpCam1	374	2143	EpCam1-D5K5-vH
EpCam1	375	2144	Epcam1-MM1-vH
FITC	376	2145	FITC-vH
FITC	377	2146	FITC-4M-53-vH
FITC	378	2147	FITC-E2-vH
FLT3	379	2148	FLT3-NC7-vH
HIV1-envelop glycoprotein	380	2149	HIV1-N6-vH
FR1 (Folate Receptor a)	381	2150	FR1-huMov19-vH
GAD	382	2151	GAD-G3H8-vH
GD2	383	2152	GD2-hu14-18-vH
GD2	384	2153	GD2-hu3F8-vH
GD3	385	2154	GD3-KM-641-vH
GFRa4	386	2155	GFRa4-P4-10-vH
GFRa4	387	2156	GFRAlpha4-P4-6-vH
GM1	388	2157	GM1-5B2-vH
GM1	389	2158	GM1-7E5-vH
gp100/MHC class I complex	390	2159	gp100-G2D12-vH
gp100/MHC class I complex	391	2160	gp100-vH
GPC3	392	2161	GPC3-4E5-vH
gpNMB	393	2162	gpNMB-115-vH
GPRC5D	394	2163	GPRC5D-ET150-18-vH
GPRC5D	395	2164	GPRC5D-ET150-5-vH
GPRC5D	396	2165	GPRC5D-ET150-1-vH
GPRC5D	397	2166	GPRC5D-ET150-2-vH
Her2	398	2167	Her2-Hu4D5-vH
HIV1-gag (77-85) MHC class I complex	399	2168	HIV1-3BNC117-vH
HIV1-envelop glycoprotein	400	2169	HIV1-E5-vH
HIV1-envelop glycoprotein	401	2170	HIV1-PGT-128-vH
HIV1-envelop glycoprotein	402	2171	HIV1-VR-C01-vH
HIV1-envelop glycoprotein	403	2172	HIV1-X5-vH
HLA-A2	404	2173	HLA-A2-3PB2-vH
HMW-MAA	405	2174	HMW-MAA-hIND-vH
HPV16-E7/MHC class I complex	406	2175	HPV16-7-8-vH
HPV16-E7/MHC class I complex	407	2176	HPV16-2-vH
HTLV1-TAX/MHC class I	408	2177	TAX-T3E3-vH
HTLV1-TAX/ MHC class I	409	2178	TAX-T3F2-vH
IL11Ra	410	2179	IL11Ra-8E2-vH
IL13Ra2	411	2180	IL13Ra2-hu107-vH
IL13Ra2	412	2181	IL13Ra2-Hu108-vH

Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
IL6R	413	2182	IL6R-M83-vH
Influenza A HA	414	2183	FLU-MEDI-8852-vH
KSHV-gH	415	2184	YC15-vH
KSHV-K8.1	416	2185	4C3-vH
L1CAM	417	2186	L1CAM-9-3-HU3-vH
LAMP1	418	2187	LAMP1-humab1-2-vH
LAMP1	419	2188	LAMP1-Mb4-vH
LewisY	420	2189	LewisY-huS193-vH
Lym1	421	2190	Lym1-vH
Lym2	422	2191	Lym2-vH
MART1/MHC class I complex	423	2192	MART1-CAG10-vH
MART1/MHC class I complex	424	2193	MART1-CLA12-vH
Mesothelin	425	2194	Mesothelin-m912-[2]-vH
Mesothelin	426	2195	Mesothelin-m912-vH
MPL	427	2196	MPL-111-vH
MPL	428	2197	MPL-161-HL-vH
MPL	429	2198	MPL-161-vH
MPL	430	2199	MPL-175-vH
MPL	431	2200	MPL-178-vH
MPL	432	2201	MPL-huVB22Bw5-vH
MPL	433	2202	MPL-12E10-vH
MPL	434	2203	MPL-AB317-vH
Muc1/MHC class I complex	435	2204	MUC1-D6-M3A1-vH
Muc1/MHC class I complex	436	2205	Muc1-D6-M3B8-vH
Muc16	437	2206	Muc16-4H11-vH
NKG2D	438	2207	NKG2D-MS-vH
NYBR1	439	2208	NYBR1-vH
NY-ESO/MHC class I complex	440	2209	NY-ESO-T1-vH
NY-ESO/MHC class I complex	441	2210	NY-ESO-T2-vH
PD1	442	2211	PD1-4H1-vH
PD1	443	2212	PD1-5C4-vH
PDL1	444	2213	PDL1-Atezoli-vH
PDL1	445	2214	PDL1-SP142-vH
PDL1	446	2215	PR1-vH
PR1	447	2216	PSCA-Ha14-117-vH
PSCA	448	2217	PSCA-Ha14-121-vH
PSCA	449	2218	PSMA-006-vH
PSMA	450	2219	PSMA-J591-vH
PSMA	451	2220	PTK7-hSC6-23-vH
PTK7	452	2221	PTK7-SC6-10-2-vH
PTK7	453	2222	ROR1-4A5-vH
ROR1	454	2223	ROR1-4C10-vH
ROR1	455	2224	SLea-5B1-vH
SLea	456	2225	SLea-7E3-vH
SLea	457	2226	SSEA4-vH

Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
SSEA4	458	2227	TCRB1-E09-vH
TCRB1	459	2228	TCRB1-Jovi1-vH
TCRB1	460	2229	TCRB2-CP01-D05-vH
TCRB2	461	2230	TCRB2-CP01-E05-vH
TCRB2	462	2231	TCRgd-G5-4-vH
TCRgd	463	2232	TERT-3G3-T865-vH
TERT/MHC class I complex	464	2233	TERT-4A9-T540-vH
TERT/MHC class I complex	465	2234	TF1-98-vH
TF1	466	2235	TGFBR2-Ab1-vH
TGFBR2	467	2236	TIM1-HVCR1-270-2-vH
TIM1	468	2237	Tim1HVCR1-ARD5-vH
TIM1	469	2238	TnAg-vH
TnAg	470	2239	Tn-Muc1-hu5E5-vH
Tn-Muc1	471	2240	TROP2-ARA47-HV3KV3-vH
TROP2	472	2241	TROP2-h7E6-SVG-vH
TROP2	473	2242	TSHR-5C9-vH
TSHR	474	2243	TSHR-K1-70-vH
TSHR	475	2244	TSHR-KB1-vH
TSHR	476	2245	TSLPR-vH
TSLRP	477	2246	Tyro-B2-vH
Tyrosinase/MHC class I complex	478	2247	Tyro-Mc1-vH
Tyrosinase/MHC class I complex	479	2248	TA2-vH
Tyrosinase/MHC class I complex	480	2249	VEGFR3-Ab1-vH
VEGFR3	481	2250	WT1-Ab13-vH
WT1/MHC class I complex	482	2251	WT1-Ab15-vH
WT1/MHC class I complex	483	2252	WT1-Ab1-vH
WT1/MHC class I complex	484	2253	WT1-Ab5-[2]-vH
WT1/MHC class I complex	485	2254	WT1-Ab5-vH
EBV-gp350	486	2255	EBV-gp350-vH
CDH19	487	2256	CDH19-4B10-vH
FRbeta	488	2257	FRbeta-m923-vH
LHR	489	2258	LHR-8B7-vH
LHR	490	2259	LHR-5F4-21-vH
B7H4	491	2260	B7H4-hu22C10-vH
B7H4	492	2261	B7H4-hu1D11-vH
IgE	493	2262	IgE-omalizumab-vH
CD23	494	2263	CD23-p5E8-vH
GCC	495	2264	GCC-5F9-vH
GCC	496	2265	GCC-Ab229-vH
CD200R	497	2266	CD200R-huDx182-vH
PDL1	498	2268	PDL1-10A5-vH

[00230] Table 7: Exemplary V_{HH} fragments (Nanobodies)

Target	SEQ ID (DNA)	SEQ ID (PRT)	NAME
Her2	499	2269	Her2-2D3-vHH
Her2	500	2270	Her2-5F7-vHH
Her2	501	2271	Her2-47D5-vHH
Her3	502	2272	Her3-17B05So-vHH
Her3	503	2273	Her3-21F06-vHH
CEA	504	2274	CEA1-vHH
CEA	505	2275	CEA5-vHH
EGFR	506	2276	EGFR1-vHH
EGFR	507	2277	EGFR33-vHH
cMet	508	2278	cMET-171-vHH
CXCR4	509	2279	CXCR4-2-vHH
CXCR4	510	2280	CXCR4-1-vHH
Mesothelin	511	2281	SD1-vHH
Mesothelin	512	2282	SD2-vHH
Albumin	513	2283	Alb8-vHH
CD123	514	2284	CD123-1-vHH
CD123	515	2285	CD123-2-vHH
IL6R	516	2286	IL6R-304-vHH
EGFR & CEA	517	2287	EGFR1-vHH-Gly-Ser-Linker-CEA1-vHH
EGFR & CEA	518	2288	EGFR33-vHH-Gly-Ser-Linker-CEA5-vHH
Her2	519	2289	Her2-5F7-vHH-Gly-Ser-Linker-Her2-47D5-vHH
Her2	520	2290	Her2-Hu4D5-vL-Gly-Ser-Linker-Her2-Hu4D5-vH
Her3 & Her2	521	2291	Her3-17B05So-vHH-Gly-Ser-Linker-Her2-2D3-vHH
cMet & Her3	522	2292	cMET-171-vHH-Gly-Ser-Linker-Her3-21F06-vHH
Mesothelin	523	2293	SD1-vHH-Gly-Ser-Linker-SD2-vHH

[00231] Table 8: Exemplary non-immunoglobulin antigen binding scaffolds (affibodies, darpins)

Table 8. Non-Immunoglobulin Antigen Binding Scaffolds			
Target	SEQ ID (DNA)	SEQ ID (PRT)	NAME
Her2	524	1743	Her2-DARPIN-1
Her2	525	1744	Her2-DARPIN-2
Her3	526	1745	Her3-affi
Her2	527	1746	Her2-affi
EGFR	528	1747	EGFR-affi

[00232] Table 9: Exemplary receptor extracellular domain

SEQ ID (DNA)	SEQ ID (PRT)	NAME
529	2299	CD19-Extracellular-Domain-minus-signal-peptide(61-867)
530	2300	MPL-Extracellular-Domain-with-signal-peptide
531	2301	CD8-SP-PD1-opt-ECD
532	2302	PD1-opt-ECD-minus-signal-peptide
533	2303	PD1-ECD-with-native-Signal-Peptide
534	2304	CTLA4-opt-ECD with signal peptide
535	2305	CD138-SDC1-ECD
536	2306	Synth-CD123-ECD
537	2307	CDH1-ECD
538	2308	CD200R1L-ECD
539	2309	GPNMB-ECD
540	2310	PTK7-ECD
541	2311	CD33-ECD
542	2312	CD34-ECD
543	2313	EpCAM-ECD
544	2314	CLEC12A-ECD
545	2315	CD20-ECx2-ECD
546	2316	CD20-ECx1-ECD
547	2317	CD22-v5-ECD
548	2318	Thyroid Stimulating Hormone Receptor (TSHR)-ECD
549	2319	EGFRviii-ECD
550	2320	BCMA-ECD
551	2321	SLAMF7-CS1-ECD
552	2322	NKG2D-ECD-minus-signal-peptide

[00233] Table 10: Exemplary ligands

SEQ ID (DNA)	SEQ ID (PRT)	NAME
553	2323	hTPO (1-187)
554	2324	mTPO(1-187)
555	2325	CGH-alpha-minus-Signal-Peptide
556	2326	CGH-beta-with-Signal-Peptide
557	2327	FSH-beta-minus-Signal-Peptide
558	2328	LH-beta-with-Signal-Peptide
559	2329	TSH-beta-with-Signal-Peptide
560	2330	SP-CGHb-Gly-Ser-Linker-CGHa
561	2331	CD8SP-FSHb-Gly-Ser-Linker-CGHa
562	2332	SP-LHb-Gly-Ser-Linker-CGHa
563	2333	SP-TSHb-Gly-Ser-Linker-CGHa

[00234] **Table 11: Exemplary Reporter Fragments**

SEQ ID (DNA)	SEQ ID (Prt)	NAME
813	2029	GLuc (<i>Gussia princeps</i> Luc) Minus Secretory Signal
814	2030	NLuc (NanoLuc)
815	2031	TLuc (TurboLuc16) Minus Secretory Signal
816	2032	MLuc7-(<i>Metridia longa</i> Luc7) M43L/M110L Variant Minus Secretory Signal
817	2033	LoLuc (<i>Lucicutia ovaliformis</i> Luc) Minus Secretory Signal
818	2034	HtLuc (<i>H. tanneri</i> Luc) Minus Secretory Signal
819	2035	PaLuc1 (<i>Pleuromamma abdominalis</i> Luc1) minus Secretory Signal
820	2036	PaLuc2 (<i>Pleuromamma abdominalis</i> Luc2)-minus Secretory Signal
821	2037	MpLuc1 [<i>Metridia pacifica</i> Luc1] minus secretory signal
822	2038	McLuc1 [<i>Metridia curticauda</i> Luc1] minus secretory signal
823	2039	MaLuc1 [<i>Metridia asymmetrica</i> Luc1] minus secretory signal
824	2040	MoLuc1 [<i>Metridia okhotensis</i> Luc1] minus secretory signal
825	2041	MoLuc2 [<i>Metridia okhotensis</i> Luc2] minus secretory signal
826	2042	MLuc39 [<i>Metridia longa</i> Luc39] minus secretory signal
827	2043	PsLuc1 [<i>Pleuromamma scutellata</i> Luc1] minus secretory signal
828	2044	LoLuc1-3 [<i>Lucicutia ovaliformis</i> Luc1-3] minus secretory signal
829	2045	HtLuc2 [<i>Heterorhabdus tanneri</i> Luc2] minus secretory signal
830	2046	Renilla Luc

[00235] **Table 12: Exemplary cleavable linkers and furine cleavage site**

NAME	SEQ ID (DNA)	SEQ ID (PRT)
F2A	831	2598
T2A	832	2599
T2A	833	2599
P2A	834	2600
P2a-variant	835	2601
E2A	836	2602
SGSG	837	2603
SGSG	838	2603
FURINE CLEAVAGE SITE	839	2604
FURINE CLEAVAGE SITE	840	2604
FURINE Cleavage Site	841	2604

[00236] **Table 13: Epitope Tags**

SEQ ID (DNA)	SEQ ID (PRT)	NAME
553	2569	Myc-TAG
554	2570	FLAG-TAG
555	2571	AcV5

SEQ ID (DNA)	SEQ ID (PRT)	NAME
556	2572	V5-TAG
557	2573	HA-TAG
558	2574	HIS-TAG
559	2575	AVI-TAG-delta-GSG
560	2576	StrepTagII
561	2577	4XFLAG-2xSTREP-8xHIS
562	2578	3xFLAG
563	2579	4xHA-Strep-8xHis
3526	3530	RITX-TAG
3527	3531	RITX2-TAG
3528	3532	RITX4-TAG
3529	3533	GA-TAG

[00237] Table 14: Exemplary scFv fragments

Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
CD19	564	2334	FMC63-(vL-vH)
CD19	565	2335	huFMC63-11-(vL-vH)
CD19	566	2336	CD19Bu12-(vL-vH)
CD19	567	2337	CD19MM-(vL-vH)
CD19	568	2338	CD19-4G7-(vL-vH)
CD19	569	2339	CD19-MEDI-3649-(vL-vH)
CD19	570	2340	CD19-Medrex-24D1-(vL-vH)
CD19	571	2341	CD8SP-Ritx-CD19-MOR0028-(vL-vH)
CD19	572	2342	CD19-HD37-H2L1-(vL-vH)
CD19	573	2343	CD19-huBly3-(vL-vH)
CD19	574	2344	CD19-huSJ25C1-(vL-vH)
CD19	575	2345	CD8SP-Ritx-CD19-hB4-(vL-vH)
CD19	576	2346	CD19-hu-mROO5-1-(vL-vH)
CD19	577	2347	CD19-hA19-(vL-vH)
AFP/MHC class I complex	578	2348	AFP-61-(vL-vH)
AFP/MHC class I complex	579	2349	AFP-76-(vL-vH)
AFP/MHC class I complex	580	2350	AFP-79-(vL-vH)
HIV1-env	581	2351	HIV1-N6-(vL-vH)
ALK	582	2352	Alk-48-(vL-vH)
ALK	583	2353	Alk-58-(vL-vH)
Amyloid	584	2354	Amyloid-158-(vL-vH)
CD45	585	2355	BC8-CD45-(vL-vH)
BCMA	586	2356	BCMA-J6M0-(vL-vH)
BCMA	587	2357	BCMA-huC12A3-L3H3-(vL-vH)
BCMA	588	2358	BCMA-ET-40-(vL-vH)
BCMA	589	2359	BCMA-ET-54-(vL-vH)
BCMA	590	2360	BCMA-ET-03-(vL-vH)
BCMA	591	2361	BCMA-huC11.D5.3L1H3-(vL-vH)
BCMA	592	2362	BCMA-huC13-F12-(vL-vH)

Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
CCR4	593	2363	CCR4-humAb1567-(vL-vH)
CD5	594	2364	CD5-9-(vL-vH)
CD5	595	2365	CD5-18-(vL-vH)
CD20	596	2366	CD20-2F2-(vL-vH)
CD20	597	2367	CD20-GA101-(vL-vH)
CD22	598	2368	CD22-h10F4v2-(vL-vH)
CD20	599	2369	CD20-Leu16-(vL-vH)
CD20	600	2370	CD20-11B8-(vL-vH)
CD20	601	2371	CD20-2C6-(vL-vH)
CD20	602	2372	CD20-2H7-(vL-vH)
CD20	603	2373	CD20-hA20-(vL-vH)
CD20	604	2374	CD20-BM-CA-1925-v4-(vL-vH)
CD20	605	2375	CD20-Ubli-v4-(vL-vH)
CD20	606	2376	CD20-2H7-(vL-vH)
CD20	607	2377	CD20-h1F5-(vL-vH)
CD20	608	2378	CD20-7D8-(vL-vH)
CD20	609	2379	CD20-7D8-(vL-GA-tag-vH)
CD20	610	2380	CD20-AME-33-(vL-vH)
CD22	611	2381	CD22-H22Rhov2ACDRKA-(vL-vH)
CD22	612	2382	CD22-m971-(vL-vH)
CD22	613	2383	CD22-m971-HL-(vL-vH)
CD30	614	2384	CD30-5F11-(vL-vH)
CD30	615	2385	CD30-Ac10-(vL-vH)
CD32	616	2386	CD32-Med9-(vL-vH)
CD33	617	2387	CD33-AF5-(vL-vH)
CD33	618	2388	CD33-huMyc9-(vL-vH)
CD33	619	2389	CD8SP-Ritx2-BC33-Boehr2800308-(vL-vH)
CD33	620	2390	CD8SP-Ritx2-CD33-Him3-4-(vL-vH)
CD33	621	2391	CD33-SGNh2H12-(vL-vH)
CD33	622	2392	CD33-15G15-33-(vL-vH)
CD33	623	2393	CD33-33H4-(vL-vH)
CD33	624	2394	CD33-9C3-2-(vL-vH)
CD34	625	2395	CD34-hu4C7-(vL-vH)
CD44v6	626	2396	CD44v6-Biwa8-(vL-vH)
CD70	627	2397	CD70-h1F6-(vL-vH)
CD79b	628	2398	CD79b-2F2-(vL-vH)
CD99	629	2399	CD99-hu12E7-(vL-vH)
CD123	630	2400	CD123-CSL362-(vL-vH)
CD123	631	2401	CD123-1172-(vL-vH)
CD123	632	2402	CD123-DART1-1-(vL-vH)
CD123	633	2403	CD123-DART1-2-(vL-vH)
CD123	634	2404	CD123-I3RB18-(vL-vH)
CD123	635	2405	CD123-hu3E3-(vL-vH)
CD123	636	2406	CD123-9F6-(vL-vH)
CD123	637	2407	CD123-I3RB2-(vL-vH)
CD123	638	2408	CD123-1176-(vL-vH)

Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
CD123	639	2409	CD8SP-Ritx2-CD123-8B11-(vL-vH)
CD123	640	2410	CD123-2B8-(vL-vH)
CD123	641	2411	CD123-9D7-(vL-vH)
CD123	642	2412	CD123-3B10-(vL-vH)
CD138	643	2413	CD138-(vL-vH)
CD179b	644	2414	CD179b-(vL-vH)
CD276	645	2415	CD276-17-(vL-vH)
CD324	646	2416	CD324-SC10-6-(vL-vH)
CD324	647	2417	CD324-hSC10-17-(vL-vH)
CDH6	648	2418	CDH6-NOV710-(vL-vH)
CDH6	649	2419	CDH6-NOV712-(vL-vH)
CDH17	650	2420	CDH17-PTA001A4-(vL-vH)
CDH19	651	2421	CDH19-16A4-(vL-vH)
EGFR	652	2422	Cetuximab-(vL-vH)
CLEC5A	653	2423	CLEC5A-8H8F5-(vL-vH)
CLEC5A	654	2424	CLEC5A-3E12A2-(vL-vH)
CLL1	655	2425	CLL1-M26-(vL-vH)
CLL1	656	2426	CLL1-M32-(vL-vH)
CLL1	657	2427	CLL1-21C9-L2H3-(vL-vH)
CLL1	658	2428	CLL1-6E7L4H1c-(vL-vH)
CLL1	659	2429	CLL1-hu1075-v1-(vL-vH)
CLL1	660	2430	CLL1-hu1075-v2-(vL-vH)
CMVpp65/MHC class I complex	661	2431	CMVpp65-F5-(vL-vH)
CS1	662	2432	CS1-huLuc63-(vL-vH)
CS1	663	2433	CS1-HuLuc64-(vL-vH)
CS1	664	2434	CS1-Luc90-(vL-vH)
CS1	665	2435	CS1-PDL241-(vL-vH)
CS1	666	2436	CS1-Hu27A-(vL-vH)
CS1	667	2437	CS1-ScHu34C3-(vL-vH)
CS1	668	2438	CS1-Hu31-D2-(vL-vH)
CS1	669	2439	CS1-Luc34-(vL-vH)
CS1	670	2440	CS1-LucX2-(vL-vH)
CSF2RA	671	2441	CSF2RA-Ab6-(vL-vH)
CSF2RA	672	2442	CSF2RA-Ab1-(vL-vH)
DLL3	673	2443	DLL3-hSC16-13-(vL-vH)
DLL3	674	2444	DLL3-hSC16-56-(vL-vH)
EBNA3c	675	2445	EBNA3c-315-(vL-vH)
EBV-gp350	676	2446	EBV-gp350-(vL-vH)
EGFRvIII	677	2447	EGFRvIII-139-(vL-vH)
EGFRvIII	678	2448	EGFRvIII-2173-(vH-vL)
EpCam1	679	2449	Epcam1-MM1-(vL-vH)
EpCam1	680	2450	Epcam1-D5K5-(vL-vH)
FLT3	681	2451	FLT3-NC7-(vL-vH)
FITC	682	2452	FITC-(vL-vH)
FITC	683	2453	FITC-4M-53-(vL-vH)
FITC	684	2454	FITC-E2-HL-(vH-vL)

Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
Influenza A HA	685	2455	FLU-MEDI-8852-(vL-vH)
FR1 (Folate Receptor a)	686	2456	FR1-huMov19-(vL-vH)
GAD	687	2457	GAD-G3H8-(vL-vH)
GD2	688	2458	GD2-hu14-18-(vL-vH)
GD2	689	2459	GD2-hu3F8-(vL-vH)
GD3	690	2460	GD3-KM-641-(vL-vH)
GFRa4	691	2461	GFRAlpha4-P4-6-(vL-vH)
GFRa4	692	2462	GFRa4-P4-10-(vL-vH)
GM1	693	2463	GM1-5B2-(vL-vH)
GM1	694	2464	GM1-7E5-(vL-vH)
GPRC5D	695	2465	GPRC5D-ET150-5-(vL-vH)
GPRC5D	696	2466	GPRC5D-ET150-18-(vL-vH)
GPRC5D	697	2467	GPRC5D-ET150-1-(vL-vH)
GPRC5D	698	2468	GPRC5D-ET150-2-(vL-vH)
gp100/MHC class I complex	699	2469	gp100-(vL-vH)
gp100/MHC class I complex	700	2470	gp100-G2D12-(vL-vH)
GPC3	701	2471	GPC3-4E5-(vL-vH)
gpNMB	702	2472	gpNMB-115-(vL-vH)
GRP78	703	2473	GRP78-GC18-(vL-vH)
HIV-gag/MHC class I complex	704	2474	HIV1-E5-(vL-vH)
HIV1-env	705	2475	HIV1-3BNC117-(vL-vH)
HIV1-env	706	2476	HIV1-PGT-128-(vL-vH)
HIV1-env	707	2477	HIV1-VR-C01-(vL-vH)
HIV1-env	708	2478	HIV1-X5-(vL-vH)
HLA-A2	709	2479	HLA-A2-3PB2-(vL-vH)
HMW-MAA	710	2480	HMW-MAA-hIND-(vL-vH)
HPV16	711	2481	HPV16-7-8-(vL-vH)
HPV16	712	2482	HPV16-2-(vL-vH)
HTLV1-TAX/MHC class I complex	713	2483	HTLV-TAX-T3F2-(vL-vH)
HTLV1-TAX/MHC class I complex	714	2484	HTLV-TAX-T3E3-(vL-vH)
IL11Ra	715	2485	IL11Ra-8E2-Ts107-(vL-vH)
IL13Ra2	716	2486	IL13Ra2-hu107-(vL-vH)
IL13Ra2	717	2487	IL13Ra2-Hu108-(vL-vH)
KSHV-K8.1	718	2488	KSHV-4C3-(vL-vH)
LAMP1	719	2489	LAMP1-humab1-2-(vL-vH)
LAMP1	720	2490	LAMP1-Mb4-(vL-vH)
LewisY	721	2491	LewisY-huS193-(vL-vH)
L1CAM	722	2492	L1CAM-9-3-HU3-(vL-vH)
Lym1	723	2493	Lym1-(vL-vH)
Lym2	724	2494	Lym2-(vL-vH)
CD79b	725	2495	huMA79bv28-(vL-vH)
MART/MHC class I	726	2496	MART1-CAG10-(vL-vH)

Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
complex1			
MART/MHC class I complex1	727	2497	MART1-CLA12-(vL-vH)
Mesothelin	728	2498	Mesothelin-m912-(vL-vH)
MPL	729	2499	MPL-175-(vL-vH)
MPL	730	2500	MPL-161-(vL-vH)
MPL	731	2501	MPL-161-HL-(vH-vL)
MPL	732	2502	MPL-111-(vL-vH)
MPL	733	2503	MPL-178-(vL-vH)
MPL	734	2504	MPL-AB317-(vL-vH)
MPL	735	2505	MPL-12E10-(vL-vH)
MPL	736	2506	MPL-huVB22Bw5-(vL-vH)
Muc1/MHC class I complex	737	2507	Muc1-D6-M3B8-(vL-vH)
Muc1/MHC class I complex	738	2508	MUC1-D6-M3A1-(vL-vH)
Muc16	739	2509	Muc16-4H11-(vL-vH)
EGFR	740	2510	Nimotuzumab-(vL-vH)
NKG2D	741	2511	NKG2D-MS-(vL-vH)
NYBR1	742	2512	NYBR1-(vL-vH)
NY-ESO/MHC class I complex	743	2513	NYESO-T1-(vL-vH)
NY-ESO/MHC class I complex	744	2514	NYESO-T2-(vL-vH)
PDL1	745	2515	PDL1-Atezoli-(vL-vH)
PDL1	746	2516	PDL1-SP142-(vL-vH)
PDL1	747	2517	PDL1-10A5-(vL-vH)
PSCA	748	2518	PSCA-Ha14-121-(vL-vH)
PSCA	749	2519	PSCA-Ha14-117-(vL-vH)
PR1/MHC class I complex	750	2520	PR1-(vL-vH)
PSMA	751	2521	PSMA-006-(vL-vH)
PSMA	752	2522	PSMA-J591-(vL-vH)
PTK7	753	2523	PTK7-hSC6-23-(vL-vH)
PTK7	754	2524	PTK7-SC6-10-2-(vL-vH)
ROR1	755	2525	ROR1-4A5-(vL-vH)
ROR1	756	2526	ROR1-4C10-(vL-vH)
Mesothelin	757	2527	SD1-vHH-Gly-Ser-Linker-SD2-vHH
SLea	758	2528	SLea-7E3-(vL-vH)
SLea	759	2529	SLea-5B1-(vL-vH)
SSEA4	760	2530	SSEA4-(vL-vH)
TCRB1	761	2531	TCRB1-CP01-E09-(vL-vH)
TCRB1	762	2532	TCRB1-Jovi1-(vL-vH)
TCRB2	763	2533	TCRB2-CP01-D05-(vL-vH)
TCRB2	764	2534	TCRB2-CP01-E05-(vL-vH)
TCRgd	765	2535	TCRgd-G5-4-(vL-vH)
hTERT	766	2536	TERT-4A9-T540-(vL-vH)
hTERT	767	2537	TERT-3G3-T865-(vL-vH)
Tissue Factor	768	2538	TF1-98-(vL-vH)
TGFBR2	769	2539	TGFBR2-Ab1-(vL-vH)

Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
TIM1	770	2540	TIM1-HVCR1-270-2-(vL-vH)
TIM1	771	2541	TIM1-HVCR1-ARD5-(vL-vH)
TnAg	772	2542	TnAg-(vL-vH)
Tn-Muc1	773	2543	TnMuc1-hu5E5-RHA8-RKA-2-(vL-vH)
TROP2	774	2544	TROP2-ARA47-HV3KV3-(vL-vH)
TROP2	775	2545	TROP2-h7E6-SVG-(vL-vH)
TSHR	776	2546	TSHR-K1-70-(vL-vH)
TSHR	777	2547	TSHR-KB1-(vL-vH)
TSHR	778	2548	TSHR-5C9-(vL-vH)
TSLPR	779	2549	TSLPR-(vL-vH)
Tyrosinase/MHC class I complex	780	2550	Tyros-B2-(vL-vH)
Tyrosinase/MHC class I complex	781	2551	Tyros-MC1-(vL-vH)
Tyrosinase/MHC class I complex	782	2552	Tyros-TA2-(vL-vH)
VEGFR3	783	2553	VEGFR3-Ab1-(vL-vH)
WT1/MHC class I complex	784	2554	WT1-Ab1-(vL-vH)
WT1/MHC class I complex	785	2555	WT1-Ab5-(vL-vH)
WT1/MHC class I complex	786	2556	WT1-Ab13-(vL-vH)
WT1/MHC class I complex	787	2557	WT1-Ab15-(vL-vH)
CDH19	788	2558	CDH19-4B10-(vL-vH)
Folate Receptor beta	789	2559	FRbeta-m923-(vL-vH)
LHR	790	2560	LHR-8B7-(vL-vH)
LHR	791	2561	LHR-5F4-21-(vL-vH)
B7H4	792	2562	B7H4-hu22C10-(vL-vH)
B7H4	793	2563	B7H4-hu1D11-(vL-vH)
IgE	794	2564	IgE-omalizumab-(vL-vH)
CD23	795	2565	CD23-p5E8-(vL-vH)
GCC	796	2566	GCC-5F9-(vL-vH)
GCC	797	2567	GCC-Ab229-(vL-vH)
CD200R	798	2568	CD200R-huDx182-(vL-vH)

[00238] Table 15: Exemplary scFv-GGS-Luc Constructs

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
	CD19	854	2054	CD8SP-FMC63-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
020317-E05	CD19	855	2055	CD8SP-huFMC63-11-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
121516-F01	CD19	856	2056	CD8SP-huFMC63-11-N203Q-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD19	857	2057	CD8SP-CD19Bu12-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
021517	CD19	858	2058	CD8SP-2-CD19MM-(vL-vH)-GGSG-NLuc-

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
-L02				4xFLAG-x2STREP-8xHis-T2A-PAC
062916-C08	CD19	859	2059	CD8SP-CD19-4G7-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
011817-A01	CD19	860	2060	CD8SP-CD19-MEDI-3649-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD19	861	2061	CD8SP-CD19-Medrex-24D1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD19	862	2062	CD8SP-Ritx-CD19-MOR0028-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD19	863	2063	CD8SP-CD19-HD37-H2L1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
020317-I04	CD19	864	2064	CD8SP-CD19-huBly3-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
022817-C04	CD19	865	2065	CD8SP-CD19-huSJ25C1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
022817-F06	CD19	866	2066	CD8SP-Ritx-CD19-hB4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD19	867	2067	CD8SP-CD19-hu-mROO5-1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
022817-H03	CD19	868	2068	CD8SP-CD19-hA19-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
022217-K04	AFP/MHC class I complex	869	2069	CD8SP-AFP-61-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	AFP/MHC class I complex	870	2070	CD8SP-AFP-76-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	AFP/MHC class I complex	871	2071	CD8SP-AFP-79-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
121516-D05	HIV1-envelop glycoprotein	872	2072	CD8SP-HIV1-N6-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
061316-B05	ALK (Anaplastic Lymphoma Kinase)	873	2073	CD8SP-Alk-48-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
061416-C04	ALK (Anaplastic Lymphoma Kinase)	874	2074	CD8SP-Alk-58-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Amyloid	875	2075	SP-Amyloid-158-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
062116-A05	Biotin	876	2076	CD8SP-dc-Avidin-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD45	877	2077	CD8SP-BC8-CD45-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	BCMA	878	2078	CD8SP-BCMA-J6M0-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
	BCMA	879	2079	CD8SP-BCMA-huC12A3-L3H3-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
092016-A02	BCMA	880	2080	CD8SP-BCMA-ET-40-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
092016-B04	BCMA	881	2081	CD8SP-BCMA-ET-54-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	BCMA	882	2082	CD8SP-BCMA-ET-03-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	BCMA	883	2083	CD8SP-BCMA-huC11.D5.3L1H3-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	BCMA	884	2084	CD8SP-BCMA-huC13-F12-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CCR4	885	2085	CD8SP-CCR4-humAb1567-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	HIV1-envelop glycoprotein	886	2086	CD8SP-CD4-ECD-Gly-Ser-Linker-DC-SIGN-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD5	887	2087	CD8SP-CD5-9-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD5	888	2088	CD8SP-CD5-18-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Ig Fc	889	2089	CD8SP-CD16A-V158-ECD-v2-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Ig Fc	890	2090	CD8SP-CD16A-V158-ECD-v1-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD20	891	2091	CD8SP-CD20-2F2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
121516-G07	CD20	892	2092	CD8SP-CD20-GA101-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD20	893	2093	CD8SP-CD20-Leu16-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD20	894	2094	CD8SP-CD20-11B8-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD20	895	2095	CD8SP-CD20-2C6-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD20	896	2096	CD8SP-CD20-2H7-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD20	897	2097	CD8SP-CD20-hA20-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD20	898	2098	CD8SP-CD20-BM-CA-1925-v4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD20	899	2099	CD8SP-CD20-Ubli-v4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
022817-E05	CD20	900	2100	CD8SP-CD20-2H7-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD20	901	2101	CD8SP-CD20-h1F5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD20	902	2102	CD8SP-CD20-7D8-(vL-vH)-GGSG-NLuc-

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
				4xFLAG-x2STREP-8xHis-T2A-PAC
	CD20	903	2103	CD8SP-CD20-AME-33-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD22	904	2104	CD8SP-CD22-h10F4v2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
060716-H05	CD22	905	2105	CD8SP-CD22-H22RhoV2ACDRKA-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD22	906	2106	CD8SP-CD22-m971-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD22	907	2107	CD8SP-CD22-m971-HL-(vH-vL)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD30	908	2108	CD8SP-CD30-5F11-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
031816-D02	CD30	909	2109	CD8SP-CD30-Ac10-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
081516-W03	CD32	910	2110	CD8SP-CD32-Med9-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD33	911	2111	CD8SP-CD33-AF5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD33	912	2112	CD8SP-CD33-huMyc9-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD33	913	2113	CD8SP-CD33-Boehr2800308-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
022217-A02	CD33	914	2114	CD8SP-CD33-Him3-4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD33	915	2115	CD8SP-CD33-SGNh2H12-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD33	916	2116	CD8SP-CD33-15G15-33-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
022217-D03	CD33	917	2117	CD8SP-CD33-33H4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
022217-E04	CD33	918	2118	CD8SP-CD33-9C3-2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
062816-B06	CD34	919	2119	CD8SP-CD34-hu4C7-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
120716-O05	CD44v6	920	2120	CD8SP-CD44v6-Biwa8-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD70	921	2121	CD8SP-CD70-h1F6-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD79b	922	2122	CD8SP-CD79b-2F2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD79b	923	2123	CD8SP-huMA79bv28-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD99	924	2124	CD8SP-CD99-hu12E7-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD123	925	2125	CD8SP-CD123-CSL362-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
112216	CD123	926	2126	CD8SP-CD123-1172-(vL-vH)-GGSG-NLuc-

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
-B03				4xFLAG-x2STREP-8xHis-T2A-PAC
	CD123	927	2127	CD8SP-CD123-DART-1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD123	928	2128	CD8SP-CD123-DART-2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD123	929	2129	CD8SP-CD123-I3RB18-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD123	930	2130	CD8SP-CD123-hu3E3-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD123	931	2131	CD8SP-CD123-9F6-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD123	932	2132	CD8SP-CD123-I3RB2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD123	933	2133	CD8SP-CD123-1176-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD123	934	2134	CD8SP-Ritx2-CD123-8B11-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD123	935	2135	CD8SP-CD123-2B8-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD123	936	2136	CD8SP-CD123-9D7-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD123	937	2137	CD8SP-CD123-3B10-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
041715-D08	CD138	938	2138	CD8SP-CD138-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD179b	939	2139	CD8SP-CD179b-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
060716-C02	CD276	940	2140	CD8SP-CD276-17-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD324	941	2141	CD8SP-CD324-SC10-6-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD324	942	2142	CD8SP-CD324-hSC10-17-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CDH6	943	2143	CD8SP-CDH6-NOV710-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CDH6	944	2144	CD8SP-CDH6-NOV712-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CDH17	945	2145	CD8SP-CDH17-PTA001A4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CDH19	946	2146	CD8SP-CDH19-16A4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
C03-071916	EGFR	947	2147	CD8SP-Cetuximab-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
060716-E05	CLEC5A	948	2148	CD8SP-CLEC5A-8H8F5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
060716-G01	CLEC5A	949	2149	CD8SP-CLEC5A-3E12A2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
082216	GR/LHR	950	2150	SP-CGHb-Gly-Ser-Linker-CGHa-GGSG-NLuc-

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
-X02	(Gonadotropin Receptor)			4xFLAG-x2STREP-8xHis-T2A-PAC
	CLL1	951	2151	CD8SP-CLL1-M26-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CLL1	952	2152	CD8SP-CLL1-M32-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
020117-F06	CLL1	953	2153	CD8SP-CLL1-21C9-L2H3-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CLL1	954	2154	CD8SP-CLL1-6E7L4H1e-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
022217-G06	CLL1	955	2155	CD8SP-CLL1-hu1075-v1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CLL1	956	2156	CD8SP-CLL1-hu1075-v2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CMVpp65/MHC class I complex	957	2157	CD8SP-CMVpp65-F5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CS1 (SLAMF7)	958	2158	CD8SP-CS1-huLuc63-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
101216-A04	CS1 (SLAMF7)	959	2159	CD8SP-CS1-HuLuc64-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CS1 (SLAMF7)	960	2160	CD8SP-CS1-Luc90-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CS1 (SLAMF7)	961	2161	CD8SP-CS1-PDL241-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CS1 (SLAMF7)	962	2162	CD8SP-CS1-Hu27A-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CS1 (SLAMF7)	963	2163	CD8SP-CS1-ScHu34C3-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CS1 (SLAMF7)	964	2164	CD8SP-CS1-Hu31-D2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CS1(SLAMF7)	965	2165	CD8SP-CS1-Luc34-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CS1 (SLAMF7)	966	2166	CD8SP-CS1-LucX2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
061316-H08	CSF2RA	967	2167	CD8SP-CSF2RA-Ab6-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
061316-G03	CSF2RA	968	2168	CD8SP-CSF2RA-Ab1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CXCR4 and CD123	969	2169	CD8SP-CXCR4-1-vHH-Gly-Ser-Linker-CD123-1-vHH-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CXCR4 and CD123	970	2170	CD8SP-CXCR4-2-VHH-Gly-Ser-Linker-CD123-2-VHH-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	DLL3 (Delta Like Ligand 3)	971	2171	CD8SP-DLL3-hSC16-13-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
	DLL3 (Delta Like Ligand 3)	972	2172	CD8SP-DLL3-hSC16-56-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	EBNA3c/MHC class I complex	973	2173	CD8SP-EBNA3c-315-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	EBV-gp350	974	2174	CD8SP-EBV-gp350-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	EGFR	975	2175	CD8SP-EGFR1-vHH-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
041116-E06	EGFR & CEA	976	2176	CD8SP-EGFR1-vHH-Gly-Ser-Linker-CEA1-vHH-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
041116-F02	EGFR & CEA	977	2177	CD8SP-EGFR33-vHH-Gly-Ser-Linker-CEA5-vHH-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
112316-O02	EGFRvIII	978	2178	CD8SP-EGFRvIII-139-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
112316-P02	EGFRvIII	979	2179	CD8SP-EGFRvIII-2173-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	EpCam1	980	2180	CD8SP-Epcam1-MM1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	EpCam1	981	2181	CD8SP-Epcam1-D5K5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	FLT3	982	2182	CD8SP-FLT3-NC7-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
060716-B07	FITC	983	2183	CD8SP-FITC-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	FITC	984	2184	CD8SP-FITC-4M-53-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
030617-H05	FITC	985	2185	CD8SP-FITC-E2-HL-(vH-vL)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
092916-B03	Influenza A HA	986	2186	CD8SP-FLU-MEDI-8852-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
111516-K08	FR1 (Folate Receptor alpha)	987	2187	CD8SP-FR1-huMov19-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
082216-U02	FSHR (Follicle Stimulating Hormone Receptor)	988	2188	CD8SP-FSHb-Gly-Ser-Linker-CGHa-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	GAD (Glutamic Acid Decarboxylase)/MHC class I complex	989	2189	CD8SP-GAD-G3H8-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	GD2	990	2190	CD8SP-GD2-hu14-18-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
	GD2	991	2191	CD8SP-GD2-hu3F8-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
060716-A03	GD3	992	2192	CD8SP-GD3-KM-641-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	GFRa4 (GDNF Family Receptor Alpha 4)	993	2193	CD8SP-GFRAlpha4-P4-6-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	GFRa4 (GDNF Family Receptor Alpha 4)	994	2194	CD8SP-GFRa4-P4-10-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	GM1	995	2195	CD8SP-GM1-5B2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	GM1	996	2196	CD8SP-GM1-7E5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
092016-C06	GPRC5D (G-protein coupled receptor family C group 5 member D)	997	2197	CD8SP-GPRC5D-ET150-5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
100516-A06	GPRC5D	998	2198	CD8SP-GPRC5D-ET150-18-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	GPRC5D	999	2199	CD8SP-GPRC5D-ET150-1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	GPRC5D	1000	2200	CD8SP-GPRC5D-ET150-2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	gp100/MHC class I complex	1001	2201	CD8SP-gp100-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	gp100/MHC class I complex	1002	2202	CD8SP-gp100-G2D12-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
092116-C04	GPC3 (Glypican 3)	1003	2203	CD8SP-GPC3-4E5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	gpNMB (Glycoprotein Nmb)	1004	2204	CD8SP-gpNMB-115-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	GRP78	1005	2205	CD8SP-GRP78-GC18-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Her2	1006	2206	CD8SP-Her2-5F7-vHH-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Her2	1007	2207	IgHSP-Her2-Affi-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Her2	1008	2208	CD8SP-Her2-1-Darpin-GGSG-NLuc-4xFLAG-

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
				x2STREP-8xHis-T2A-PAC
	Her2	1009	2209	IgHSP-Her2-2-Darpin-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
041116-G07	Her2	1010	2210	CD8SP-Her2-5F7-vHH-Gly-Ser-Linker-Her2-47D5-vHH-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
061316-L02	Her2	1011	2211	CD8SP-Her2-Hu4D5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Her3	1012	2212	CD8SP-Her3-17B05So-vHH-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Her3	1013	2213	CD8SP-Her3-Affi-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
041116-D05	Her2 and Her3	1014	2214	CD8SP-Her3-17B05So-vHH-Gly-Ser-Linker-Her2-2D3-vHH-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
090716-B05	HIV1-gag/MHC class I complex	1015	2215	CD8SP-HIV1-E5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
093016-Z11	HIV1-envelop glycoprotein	1016	2216	CD8SP-HIV1-3BNC117-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	HIV1-envelop glycoprotein	1017	2217	CD8SP-HIV1-PGT-128-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
093016-X10	HIV1-envelop glycoprotein	1018	2218	CD8SP-HIV1-VR-C01-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
093016-Y15	HIV1-envelop glycoprotein	1019	2219	CD8SP-HIV1-X5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	HLA-A2	1020	2220	CD8SP-HLA-A2-3PB2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
061316-E01	HMW-MAA	1021	2221	CD8SP-HMW-MAA-hIND-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	HPV16-E7/MHC class I complex	1022	2222	CD8SP-HPV16-7-8-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	HPV16-E7/MHC class I complex	1023	2223	CD8SP-HPV16-2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	HTLV1-TAX/MHC class I complex	1024	2224	CD8SP-HTLV-TAX-T3F2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
081216-O01	HTLV1-TAX/MHC	1025	2225	CD8SP-HTLV-TAX-T3E3-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
	class I complex			
060716-D06	IL11Ra	1026	2226	CD8SP-IL11Ra-8E2-Ts107-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	IL6Ra	1027	2227	IgHSP-IL6R-304-vHH-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	IL13Ra2	1028	2228	CD8SP-IL13Ra2-hu107-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
060716-F03	IL13Ra2	1029	2229	CD8SP-IL13Ra2-Hu108-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	KSHV-K8.1	1030	2230	CD8SP-KSHV-4C3-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	LAMP1 (Lysosomal-associated membrane protein 1)	1031	2231	CD8SP-LAMP1-humab1-2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
061416-B04	LAMP1 (Lysosomal-associated membrane protein 1)	1032	2232	CD8SP-LAMP1-Mb4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
081016-F11	LewisY	1033	2233	CD8SP-LewisY-huS193-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	L1CAM	1034	2234	CD8SP-L1CAM-9-3-HU3-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
082216-W02	LHR	1035	2235	SP-LHb-Gly-Ser-Linker-CGHa-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Lym1	1036	2236	CD8SP-Lym1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Lym2	1037	2237	CD8SP-Lym2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD79b	1038	2238	CD8SP-huMA79bv28-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	MART1/MH C class I complex	1039	2239	CD8SP-MART1-CAG10-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	MART1/MH C class I complex	1040	2240	CD8SP-MART1-CLA12-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Mesothelin	1041	2241	CD8SP-Mesothelin-m912-HL-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	cMet	1042	2242	CD8SP-cMet-171-vHH-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
041116-H01	cMet and Her3	1043	2243	CD8SP-cMET-171-vHH-Gly-Ser-Linker-Her3-21F06-vHH-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	MPL (Thrombopoi	1044	2244	CD8SP-MPL-175-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
	etin receptor)			
	MPL (Thrombopoietin receptor)	1045	2245	CD8SP-MPL-161-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	MPL (Thrombopoietin receptor)	1046	2246	CD8SP-MPL-161-HL-(vH-vL)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
041116-C03	MPL (Thrombopoietin receptor)	1047	2247	CD8SP-2-MPL-111-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	MPL (Thrombopoietin receptor)	1048	2248	CD8SP-MPL-178-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	MPL (Thrombopoietin receptor)	1049	2249	CD8SP-MPL-AB317-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	MPL (Thrombopoietin receptor)	1050	2250	CD8SP-MPL-12E10-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	MPL (Thrombopoietin receptor)	1051	2251	CD8SP-MPL-huVB22Bw5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Muc1/MHC class I complex	1052	2252	CD8SP-Muc1-D6-M3B8-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Muc1/MHC class I complex	1053	2253	CD8SP-MUC1-D6-M3A1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Muc16	1054	2254	CD8SP-Muc16-4H11-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
062916-A02	EGFR	1055	2255	CD8SP-Nimotuzumab-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
060716-J06	NKG2D Ligand	1056	2256	CD8SP-NKG2D-(GGGGS-GGGGD)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
090716-C02	NKG2D	1057	2257	CD8SP-NKG2D-MS-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	NY-BR1	1058	2258	CD8SP-NYBR1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
081216-M04	NY-ESO/MHC class I complex	1059	2259	CD8SP-NYESO-T1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	NY-ESO/MHC class I complex	1060	2260	CD8SP-NYESO-T1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	PD1 ligand (e.g., PDL1)	1061	2261	CD8SP-PD1-ECD-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
100516	PDL1	1062	2262	CD8SP-PDL1-Atezoli-(vL-vH)-GGSG-NLuc-

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
-G03				4xFLAG-x2STREP-8xHis-T2A-PAC
100516-H05	PDL1	1063	2263	CD8SP-PDL1-SP142-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
100516-F03	PDL1	1064	2264	CD8SP-PDL1-10A5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
090716-E02	PSCA (Prostate stem cell antigen)	1065	2265	CD8SP-PSCA-Ha14-121-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	PSCA (Prostate stem cell antigen)	1066	2266	CD8SP-PSCA-Ha14-117-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	PR1/MHC class I complex	1067	2267	CD8SP-PR1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	PSMA (Prostate Specific Membrane Antigen)	1068	2268	CD8SP-PSMA-006-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	PSMA (Prostate Specific Membrane Antigen)	1069	2269	CD8SP-PSMA-J591-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	PTK7 (Tyrosine-protein kinase-like 7)	1070	2270	CD8SP-PTK7-hSC6-23-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	PTK7 (Tyrosine-protein kinase-like 7)	1071	2271	CD8SP-PTK7-SC6-10-2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	ROR1	1072	2272	CD8SP-ROR1-4A5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	ROR1	1073	2273	CD8SP-ROR1-4C10-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Mesothelin	1074	2274	CD8SP-SD1-vHH-Gly-Ser-Linker-SD2-vHH-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
111116-A05	SLea	1075	2275	CD8SP-SLea-7E3-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	SLea	1076	2276	CD8SP-SLea-5B1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
082216-Y02	SSEA4 (stage-specific embryonic antigen 4)	1077	2277	CD8SP-SSEA4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
	TCRB1 (TCR beta 1 constant chain)	1078	2278	CD8SP-TCRB1-CP01-E09-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	TCRB1 (TCR beta 1 constant chain)	1079	2279	CD8SP-TCRB1-Jov1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	TCRB2 (TCR beta 2 constant chain)	1080	2280	CD8SP-TCRB2-CP01-D05-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	TCRB2 (TCR beta 2 constant chain)	1081	2281	CD8SP-TCRB2-CP01-E05-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
110216-B02-PB	TCRgd (TCR gamma/delta)	1082	2282	CD8SP-TCRgd-G5-4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	hTERT/MH C class I complex	1083	2283	CD8SP-TERT-4A9-T540-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
081516-U07	hTERT/MH C class I complex	1084	2284	CD8SP-TERT-3G3-T865-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Tissue Factor	1085	2285	CD8SP-TF1-98-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	TGFBR2	1086	2286	CD8SP-TGFBR2-Ab1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	TIM1/HAVCR	1087	2287	CD8SP-TIM1-HVCR1-270-2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	TIM1/HAVCR	1088	2288	CD8SP-TIM1-HVCR1-ARD5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
061316-F06	TnAg	1089	2289	CD8SP-TnAg-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
080916-C02	Tn-Muc1	1090	2290	CD8SP-TnMuc1-hu5E5-RHA8-RKA-2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	MPL (Thrombopoietin receptor)	1091	2291	CD8SP-hTPO-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	TROP2 (Trophoblast cell-surface antigen-2)	1092	2292	CD8SP-TROP2-ARA47-HV3KV3-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	TROP2 (Trophoblast cell-surface antigen-2)	1093	2293	CD8SP-TROP2-h7E6-SVG-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
082216	TSHR	1094	2294	SP-TSHb-Gly-Ser-Linker-CGHa-GGSG-NLuc-

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
-V02	(Thyrotropin receptor)			4xFLAG-x2STREP-8xHis-T2A-PAC
	TSHR (Thyrotropin receptor)	1095	2295	CD8SP-TSHR-K1-70-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
061316-D01	TSHR (Thyrotropin receptor)	1096	2296	CD8SP-TSHR-KB1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
062916-B07	TSHR (Thyrotropin receptor)	1097	2297	CD8SP-TSHR-5C9-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	TSLPR (thymic stromal lymphopoietin receptor)	1098	2298	CD8SP-TSLPR-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Tyrosinase/MHC class I complex	1099	2299	CD8SP-Tyros-B2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Tyrosinase/MHC class I complex	1100	2300	CD8SP-Tyros-MC1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	Tyrosinase/MHC class I complex	1101	2301	CD8SP-Tyros-TA2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	VEGFR3	1102	2302	CD8SP-VEGFR3-Ab1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	WT1/MHC class I complex	1103	2303	CD8SP-WT1-Ab1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	WT1/MHC class I complex	1104	2304	CD8SP-WT1-Ab5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	WT1/MHC class I complex	1105	2305	CD8SP-MYC3-WT1-Ab13-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	WT1/MHC class I complex	1106	2306	CD8SP-MYC3-WT1-Ab15-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CDH19	1107	2307	CD8SP-CDH19-4B10-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
112216-A02	Folate Receptor beta	1108	2308	CD8SP-FRbeta-m923-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	LHR (Luteinizing hormone Receptor)	1109	2309	CD8SP-LHR-8B7-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	LHR	1110	2310	CD8SP-LHR-5F4-21-(vL-vH)-GGSG-NLuc-

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
	(Luteinizing hormone Receptor)			4xFLAG-x2STREP-8xHis-T2A-PAC
	B7H4	1111	2311	CD8SP-B7H4-hu22C10-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	B7H4	1112	2312	CD8SP-B7H4-hu1D11-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	IgE	1113	2313	CD8SP-IgE-omalizumab-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
121516-A08	CD23	1114	2314	CD8SP-CD23-p5E8-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
121516-C03	GCC (Guanylyl cyclase C)	1115	2315	CD8SP-GCC-5F9-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
121516-B06	GCC (Guanylyl cyclase C)	1116	2316	CD8SP-GCC-Ab229-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	CD200R	1117	2317	CD8SP-CD200R-huDx182-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
021015-T07	MPL	1118	2318	CD8SP-161-(vL-vH)-GGSG-NLuc-AcV5
021015-U09	CD19	1119	2319	CD8SP-FMC63(vL-vH)-GGSG-NLuc-AcV5
040915-J04	Lym1	1120	2320	CD8SP-Lym1(vL-vH)-GGSG-NLuc-AcV5
040915-K05	Lym2	1121	2321	CD8SP-Lym2(vL-vH)-GGSG-NLuc-AcV5
041715-E01	CD33	1122	2322	CD8SP-CD33(vL-vH)-GGSG-NLuc-AcV5
041715-D01	CD138	1123	2323	CD8SP-CD138(vL-vH)-GGSG-NLuc-AcV5
041615-X08	BCMA	1124	2324	CD8SP-BCMA-J6MO-(vL-vH)-NLuc-AcV5
041715-C06	FLT3	1125	2325	CD8SP-FLT3-NC7-(vL-vH)-NLuc-AcV5
041415-V08	CS1	1126	2326	CD8SP-Luc63-(vL-vH)-GGGS-NLuc-AcV5
041515-W03	CS1	1127	2327	CD8SP-Luc90-(vL-vH)-NLuc-AcV5
041515-U04	CD79b	1128	2328	CD8SP-huMA79bv28-(vL-vH)-GGGS-Nluc-AcV5
042115-A09	CD45	1129	2329	CD8SP-CD45-BC8-(vL-vH)-GGSG-NLuc-AcV5
040915-M06	GRP78	1130	2330	CD8SP-GRP78-GC18(vL-vH)-NLuc-AcV5
041715-B03	CD3	1131	2331	CD8SP-CD3-Hum291-(vL-vH)-GGs-NLuc-AcV5
040915	MPL	1132	2332	hTPO(1-187)-GGSG-NLuc-AcV5

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
-N01				
040915-O07	MPL	1133	2333	mTPO(1-187)-GGSG-NLuc-AcV5
021216-A06	CD19	1134	2334	CD8SP-FMC63(vL-vH)-GGSG-NLuc-x3-FLAG
113015-A04	CD19	1135	2335	CD8SP-FMC63(vL-vH)-GGSG-TurboLuc16-x3-FLAG
113015-B02	CD19	1136	2336	CD8SP-FMC63(vL-vH)-GGSG-GLuc-x3-FLAG
111815-K07	CD19	1137	2337	CD8SP-FMC63(vL-vH)-GGSG-MLuc-MM-LL-HA
111815-G04	CD19	1138	2338	CD8SP-FMC63(vL-vH)-GGSG-L-ovaliformis-Luc-AcV5
111815-L05	CD19	1139	2339	CD8SP-FMC63(vL-vH)-GGSG-Htanneri-Luc-x3-FLAG
071416-M08	FMC63	1140	2340	CD8SP-FMC136-20-L-ovalformi-luc-AcV5
030916-B02	CD30	1141	2341	CD8SP-CD30-5F11(vL-vH)-GGSG-NLuc-x3-Flag
	CD30	1142	2342	CD8SP-CD30-Ac10(vL-vH)-NLuc-x3-Flag
030816-E04	TCRB1 (TCR beta 1 constant chain)	1143	2343	CD8SP-TCRB1-Jovi1(vL-vH)-GGSG-NLuc-x3-FLAG
030816-G01	TCRB1 (TCR beta 1 constant chain)	1144	2344	CD8SP-TCRB1-E09(vL-vH)-GGSG-NLuc-x3-Flag
021417-T05	CD19	1145	2345	CD8SP-hCD19-Bu12-(vL-vH)-GGSG-NLuc-AcV5
021417-U01	CD19	1146	2346	CD8SP-hCD19MM-(vL-vH)-GGSG-NLuc-AcV5
021417-V05	CD19	1147	2347	CD8SP-huFMC63-11-(vL-vH)-GGSG-NLuc-AcV5
021417-W05	CD19	1148	2348	CD8SP-huFMC63-11-N203Q-(vL-vH)-GGSG-NLuc-AcV5
021417-X05	CD19	1149	2349	CD8SP-huFMC63-11-N203S-(vL-VH)-GGSG-NLuc-AcV5
021417-Y05	CD19	1150	2350	CD8SP-CD19-Medi-3649-(vL-vH)-GGSG-NLuc-AcV5
021417-Z04	CD19	1151	2351	CD8SP-CD19-4G7-Mlu-GSG-NLuc-AcV5
021417-A05	CD19	1152	2352	CD8SP-RTX-CD19-MOR0028-(vL-vH)-GGSG-NLuc-AcV5
031816-B04	CD19	1153	2353	CD8SP-FMC63-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
071416-F03	CD22	1154	2354	CD8SP-CD22-h10F4v2-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (PRT)	CONSTRUCT NAME
071416-C03	CD276	1155	2355	CD8SP-CD276-17-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
071416-E03	CLEC5A	1156	2356	CD8SP-CLEC5A-3E12A2-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
072616-M04	CSF2RA	1157	2357	CD8SP-CSF2RA-Ab6-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
071416-G03	Her2	1158	2358	CD8SP-Her2-Hu4D5-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
072616-E04	IL13Ra2	1159	2359	CD8SP-IL13Ra2-hu107-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
072616-G04	L1CAM	1160	2360	CD8SP-L1CAM-9-3-HU3-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
072616-F02	Mesothelin	1161	2361	CD8SP-Mesothelin-m912-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
072616-D04	PSMA	1162	2362	CD8SP-PSMA-006-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
071416-I03	PTK7	1163	2363	CD8SP-PTK7-hSC6-23-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
072616-L02	TnAg	1164	2364	CD8SP-TnAg-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
072616-C03	WT1	1165	2365	CD8SP-WT1-Ab5-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
072616-P01	WT1	1166	2366	CD8SP-MYC3-WT1-Ab15-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
071416-L06	FMC63	1167	2367	CD8SP-FMC136-20-(vL-vH)-GGSG-Turboluc16-4xFLAG-x2STREP-8xHis-T2A-PAC
022217-I05	NY-ESO	1168	2368	CD8SP-NY-ESO-IG4-HA-scTCR-Nluc-GGSG-4xFLAG-x2STREP-8xHis-T2A-PAC

[00239] Table 16: Exemplary ECD-GGS-Luc Constructs

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (Prt)	CONSTRUCT NAME
082214-Z01	MPL-CAR	1169	2369	MPL-ECD-GGSG-Nluc-AcV5
062615-C04	CD19-CAR	1170	2370	FLAG-CD19-ECD-GGSG-NLuc-AcV5
	CD19-CAR	1171	2371	CD19-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
111616-J04	CD19-CAR	1172	2372	FLAG-CD19-ECD-GGS-Turboluc16-4xFlag-2xStreptag-8xHis-T2A-Pac
111616-K07	CD19-CAR	1173	2373	FLAG-CD19-ECD-GGS-PaLuc1-HA-Streptag-3xHA-8xHis-T2A-pac
111616-L02	CD19-CAR	1174	2374	FLAG-CD19-ECD-GGS-LoLuc-Luc-AcV5
111616-M07	CD19-CAR	1175	2375	FLAG-CD19-ECD-GGS-MLuc7-MM-LL-HA

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (Prt)	CONSTRUCT NAME
111616-N05	CD19-CAR	1176	2376	CD19-ECD-GGS-HtLuc-x3Flag
060816-A02	CD33-CAR	1177	2377	CD33-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
100616-J04	CD33-CAR	1178	2378	CD33-ECD-GGSG-TurboLuc16-4xFlag-2xStreptag-8xHis-T2A-Pac
100616-K07	CD33-CAR	1179	2379	CD33-ECD-PaLuc1-HA-Streptag-3xHA-8xHis-T2A-pac
100616-L07	CD33-CAR	1180	2380	CD33-ECD-LoLuc-AcV5
100616-M02	CD33-CAR	1181	2381	CD33-ECD-MLuc7-MM-LL-HA
100616-N05	CD33-CAR	1182	2382	CD33-ECD-HtLuc-x3Flag
060816-C02	CD138-CAR	1183	2383	CD138-SDC1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
101216-E04	CD138-CAR	1184	2384	CD138-SDC1-ECD-GGS-TurboLuc16-4xFlag-2xStreptag-8xHis-T2A-Pac
101216-F04	CD138-CAR	1185	2385	CD138-SDC1-ECD-GGS-PaLuc1-HA-Streptag-3xHA-8xHis-T2A-pac
101216-G04	CD138-CAR	1186	2386	CD138-SDC1-ECD-GGS-LoLuc-Luc-AcV5
111616-H04	CD138-CAR	1187	2387	CD138-SDC1-ECD-GGS-MLuc7-MM-LL-HA
060816-D02	CD123-CAR	1188	2388	Synth-CD123-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
062816-G02	CDH1 (CD324) CAR	1189	2389	CDH1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
082616-C07	CD200R-CAR	1190	2390	CD200R-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
081716-R07	GPNMB-CAR	1191	2391	GPNMB-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
082216-S02-F	PTK7-CAR	1192	2392	PTK7-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
062816-B06	CD34-CAR	1193	2393	CD34-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
060816-F08	EpCAM-CAR	1194	2394	EpCAM-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
061716-B07	CLEC12A-CAR	1195	2395	CLEC12A-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
060816-I04	CD20-CAR	1196	2396	CD20-ECx2-ECD-GGSG-TurboLuc16-4xFlag-2xStreptag-8xHis-T2A-Pac
060816-J14	CD20-CAR	1197	2397	CD20-ECx1-ECD-GGSG-TurboLuc16-4xFlag-2xStreptag-8xHis-T2A-Pac
082616-B03	CD22-CAR	1198	2398	CD22v5-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
062816-C06	TSHR-CAR	1199	2399	TSHR-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (Prt)	CONSTRUCT NAME
060816-E01	EGFRviii-CAR	1200	2400	EGFRviii-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
103116-Q07	BCMA-CAR	1201	2401	BCMA-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
062816-A01	SLAMF7-CS1-CAR	1202	2402	SLAMF7-CS1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
	PD1-CAR or PDL1	1203	2403	PD1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac
	CTLA4-CAR	1204	2404	CTLA4-opt-ECD-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
	NKG2D-CAR	1205	2405	CD8SP-NKG2D-ECD-4xFLAG-x2STREP-8xHis-T2A-PAC
112316-Q02	Kappa Light chain CAR	1206	2406	CD8SP-Protein-L-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC
101916-P03	Kappa Light chain CAR	1207	2407	CD8SP-Protein-L-2-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC

[00240] Table 17: Exemplary CAR Constructs

Clone ID #	Target	SEQ ID (DNA)	SEQ ID (Prt)	CONSTRUCT NAME
112014-A13	CD19	1208	2408	FMC63-(vL-vH)-Myc-BBz-T2A-Pac
021015-R07	MPL	1209	2409	161-(vL-vH)-Myc-BBz-T2A-Pac
032415-N06	CD138	1210	2410	CD138-(vL-vH)-Myc-28z-T2A-Pac
032415-O01	CD33	1211	2411	CD33-(vL-vH)-Myc-28z-T2A-Pac
052115-A02	CD138	1212	2412	CD138-(vL-vH)-MYC-BBz-T2A-Pac
031915-G02	BCMA	1213	2413	J6MO-(vL-vH)-MYC-28z-T2A-Pac
052516-I07	CD22	1214	2414	CD22-h10F4v2-(vL-vH)-MYC-BBz-T2A-Pac
081115-X01	CD123	1215	2415	CD123-CSL362-(vL-vH)-MYC-BBz-T2A-Pac
081115-A02	EGFRvii	1216	2416	EGFRvIII-139-(vL-vH)-Myc-BBz-T2A-PAC
081115-E01	EGFRvii	1217	2417	EGFRvIII-2173-(vH-vL)-Myc-BBz-T2A-PAC
	CD200R	1218	2418	CD200R-huDx182-(vL-vH)-Myc-BBz-T2A-PAC
111915-E05	Epcam	1219	2419	Epcam1-D5K5-(vL-vH)-Myc-BBz-T2A-PAC
111815-A03	CD20	1220	2420	CD20-GA10-(vL-vH)-MYC-BBz-T2A-PAC
052516-B07	CD324	1221	2421	CD324-SC10-6-(vL-vH)-MYC-BBz-T2A-PAC
052516-A07	CD324	1222	2422	CD324-hSC10-17-(vL-vH)-MYC-BBz-T2A-PAC
052516-F07	PTK7	1223	2423	PTK7-hSC6-23-(vL-vH)-MYC-BBz-T2A-PAC
052516-E07	PTK7	1224	2424	PTK7-SC6-10-2-(vL-vH)-MYC-BBz-T2A-PAC
052516-G07	gpNMB	1225	2425	gpNMB-115-(vL-vH)-MYC-BBz-T2A-PAC
052616-H06	CD34	1226	2426	CD34-hu4C7-(vL-vH)-MYC-BBz-T2A-PAC
052616-E05	TSHR	1227	2427	TSHR-KB1-(vL-vH)-MYC-BBz-T2A-PAC
052616-U05	CS1	1228	2428	HuLuc64-(vL-vH)-Myc-BBz-T2A-Pac

031915-F03	CS1	1229	2429	Luc90-(vL-vH)-MYC-CD28z-T2A-Pac
042315-N01	KSHV-K8.1	1230	2430	KSHV-4C3(vL-vH)-Myc-BBz-T2A-Pac

[00241] Table 18: Protein L

SEQ ID (DNA)	SEQ ID (PRT)	
2431	2433	Protein-L
2432	2434	Protein-L-2

EXAMPLES

[00242] The following examples are not intended to limit the scope of the claims to the invention, but are rather intended to be exemplary of certain embodiments. Any variations in the exemplified methods which occur to the skilled artisan are intended to fall within the scope of the present invention

[00243] The cell lines used in this invention and their growth media are shown in the **Table 19**. Cells were cultured at 37°C, in a 5% CO₂ humidified incubator. The cell lines were obtained from ATCC, NIH AIDS reagent program or were available in our laboratory or obtained from other laboratories.

[00244] Table 19: Cell lines and growth media

Cell lines	Media	Cell lines	Media
BC-1	RPMI, 20% FCS	THP-1	RPMI, 10% FCS
BC-3	RPMI, 20% FCS	U87MG	DMEM, 10% FCS
BCBL-1	RPMI, 20% FCS	NCI-H540	DMEM, 10% FCS
JSC-1	RPMI, 20% FCS	LoVo	DMEM, 10% FCS
UMPEL-1	RPMI, 20% FCS	SKOV-3	DMEM, 10% FCS
MM1S	RPMI, 10% FCS	NCI-H1993	DMEM, 10% FCS
U266	RPMI, 10% FCS	Kasumi-1	RPMI, 20% FCS
L363	RPMI, 10% FCS	Jeko-1	RPMI, 20% FCS
K562	RPMI, 10% FCS	PC-3	DMEM, 10% FCS
BV173	RPMI, 10% FCS	HeLa	DMEM, 10% FCS
Nalm6	RPMI, 10% FCS	NCI-H2452	DMEM, 10% FCS
HL60	RPMI, 10% FCS	LnCap	DMEM, 10% FCS
U937	RPMI, 10% FCS	SNU-5	RPMI, 20% FCS
RS:411	RPMI, 20% FCS	OVCAR-3	DMEM, 10% FCS
MV:411	RPMI, 10% FCS	MEL-624	DMEM, 10% FCS
Raji	RPMI, 10% FCS	LS174-T	DMEM, 10% FCS
HEL-92.1.7	RPMI, 10% FCS	MEL-526	DMEM, 10% FCS

Cell lines	Media	Cell lines	Media
Meg-01	RPMI, 10% FCS	MDA-MB231	DMEM, 10% FCS
Jurkat	RPMI, 10% FCS	L1236	RPMI, 20% FCS
MM1	DMEM, 10% FCS	L428	RPMI, 20% FCS
Daudi	RPMI, 10% FCS	Molt-16	RPMI, 20% FCS
REC-1	RPMI, 10% FCS	RPMI8402	RPMI, 20% FCS
U2932	RPMI, 10% FCS	KN5-5F2	RPMI, 20% FCS
H929	RPMI, 10% FCS beta Mercaptoethanol (BME)	CCRF-CEM (ATCC)	RPMI, 10% FCS
KMS28	RPMI, 10% FCS	MG-63	DMEM, 10% FCS
EJM	RPMI, 10% FCS	Karpas-299	RPMI, 20% FCS
MRC-5	DMEM, 10% FCS	MCF7	DMEM, 10% FCS
CMK	RPMI, 20% FCS	SUDHL-1	RPMI, 10% FCS
TF-1	RPMI, 10% FCS + GMCSF	AA-2	RPMI, 10% FCS
ML-2	RPMI, 20% FCS	HL2/3	DMEM, 10% FCS
A20	RPMI, 10% FCS + BME	TF228.1.16	DMEM, 10% FCS
KMS28BM	RPMI, 10% FCS	MT-4	RPMI, 10% FCS
KG-1	RPMI, 20% FCS	Sup-T1	RPMI, 10% FCS
CEM	RPMI, 10% FCS	HuT-78	RPMI, 10% FCS
U937	RPMI, 10% FCS	TT	DMEM, 10% FCS
LAMA5	RPMI, 10% FCS	DMS79	RPMI, 10% FCS
A549	DMEM, 10% FCS	LAN-5	DMEM, 10% FCS
HT29	DMEM, 10% FCS	PEER1	RPMI, 10% FCS
Molm-13	RPMI, 20% FCS	SK-MEL-37	DMEM, 10% FCS
A431	DMEM, 10% FCS	Jurkat-NFAT-GFP	RPMI, 10% FCS
P19	DMEM, 10% FCS	F9	DMEM, 10% FCS

[00245] Jurkat cell line (clone E6–1) engineered with a NFAT-dependent EGFP (or GFP) reporter gene was a gift from Dr. Arthur Weiss at University of California San Francisco and have been described to study CAR-signaling ((Wu, CY et al., Science 350:293-302,2015). Jurkat cells were maintained in RPMI-1640 medium supplemented with 10% FBS, penicillin and streptomycin. 293FT cells were obtained from Thermofisher and maintained in DMEM medium supplemented with 10% FBS, penicillin and streptomycin.

Generation of lentiviral vectors encoding chimeric antigen receptors against MPL

[00246] The pLENTI-Blast vector was derived from pLenti6v5gw_lacZ vector (Invitrogen; ThermoFisher Scientific) by removal of the LacZ gene. pLenti-MP2 was a gift from Pantelis Tsoulfas (Addgene plasmid # 36097) and was used to generate pLenti-EF1a or pLenti-EF1 α lentiviral vector by replacement of the CMV promoter with human EF1 α promoter using standard molecular biology techniques. pLenti-EF1a-DWPRE was derived

from the pLENTI-EF1 α vector by deletion of WPRE sequence. The psPAX2 vector was a gift from Didier Trono (Addgene plasmid # 12260). The pLP/VSVG envelope plasmid and 293FT cells were obtained from Invitrogen (ThermoFisher Scientific). The retroviral transfer vector MSCVneo, MSCVhygro, and MSCVpac and the packaging vector pKAT were obtained from Dr. Robert Illaria's laboratory. phRGTK Renilla Luciferase plasmid was from Promega.

Example 1. Assay to detect the expression of CD19 and MPL (Thrombopoietin receptor) antigens

[00247] Both CD19 and MPL (also known as Thrombopoietin receptor or TPO-R) are expressed on hematopoietic cells but show differential expression in cells of different lineages. FMC63 is a well characterized mouse monoclonal antibody that specifically recognizes human CD19. Similarly, 161 (also designated as 1.6.1) is a monoclonal antibody that recognizes human MPL and is described in U.S. patent application US 2012/0269814 A1. We generated a FMC63 single chain Fv (scFv) fragment based on the known sequence of FMC63 vL and vH fragments. The cDNA encoding FMC63 scFv fragment consisted from 5' to 3' ends a nucleotide sequences encoding a signal peptide derived from human CD8 molecule, FMC63 vL fragment, a (Gly₄Ser)₃ linker and FMC63-vH fragment. The cDNA encoding the FMC63 scFv fragment was then fused in-frame at its 3' end to cDNA encoding AcV5-tagged NLuc through a Gly-Gly-Ser-Gly (GGSG) linker to generate FMC63-GGSG-NLuc-AcV5, which was then cloned downstream of the human EF1 α promoter into the lentiviral vector pLenti-EF1 (SEQ ID NO: 842). The DNA and PRT sequences of the insert fragment are provided in SEQ ID NO: 1119 and SEQ ID NO: 2319, respectively. A construct encoding 161-GGSG-NLuc was similarly generated using the vL and vH fragment of 161 (1.6.1) monoclonal antibody against MPL. The DNA and PRT sequences of the insert fragment are provided in SEQ ID NO: 1118 and SEQ ID NO: 2318, respectively.

[00248] The pLenti-EF1-FMC63-GGSG-NLuc-AcV5 and pLenti-EF1-161-GGSG-NLuc-AcV5 plasmids were transfected into 293FT cells by calcium phosphate co-precipitation method. Approximately 20 h post-transfection, the cell culture media was replaced with XIVO medium. The conditioned media containing the secreted FMC63-GGSG-NLuc-AcV5 and 161-GGSG-NLuc-AcV5 proteins was collected 48-72 h later.

[00249] The supernatant containing FMC63-GGSG-NLuc-AcV5 and 161-GGSG-NLuc-AcV5 proteins were used to detect the expression of CD19 and MPL on the surface of Jurkat, K562, RAJI, RS-4-11 (RS411) and HEL.92.1.7 (HEL) cells that had been engineered to express a c-MPL cDNA by transducing these cells with a lentiviral vector expressing human c-MPL cDNA or an empty vector. The cells also expressed a humanized Gluc cDNA lacking its signal peptide.

[00250] The vector- and MPL-expressing Jurkat-Gluc, K562-Gluc, HEL.92.1.7-Gluc, RAJI-Gluc and RS411-Gluc cells were incubated with the FMC63-GGSG-NLuc-AcV5 and 161-GGSG-NLuc-AcV5 supernatants at 4°C for 1h followed by extensive washings with cold PBS supplemented with 0.1%BSA. The cells were re-suspended in cold PBS and 30µl of cell suspension was plated per well in a flat-bottom 384 well plate (Greiner, 384 well white plate cat. # 781075) in triplicate. NLuc assay buffer containing native coelenterazine (CTZ) as NLuc substrate (30µl/well of native coelenterazine diluted in PBS) was added to each well by an automatic dispenser in a well mode using a BioTek synergy H4 plate reader and light emission as a measure of NLuc activity was measured.

[00251] **FIG. 1A-FIG. 1B** show that strong binding with 161-GGSG-NLuc-AcV5 was observed on HEL.92.1.7-Gluc-vector cells suggesting significant expression of MPL endogenously. Ectopic expression of MPL in HEL.92.1.7-Gluc-MPL cells led to a modest increase in 161-GGSG-NLuc-AcV5 binding. In contrast, very weak binding with 161-GGSG-NLuc-AcV5 was observed on vector-expressing Jurkat, RAJI and RS411 cells and was only modestly increased upon ectopic expression of MPL. Finally, weak but stronger binding of 161-GGSG-NLuc-AcV5 was observed on K562-vector cells, and was significantly increased on K562-MPL cells. In contrast to 161-GGSG-NLuc-AcV5, the FMC63-GGSG-NLuc-AcV5 supernatant showed strongest binding on vector- and MPL-expressing RAJI cells, modestly strong binding on RS411 cells and very weak to negligible binding on the other cells.

Example 2. Assay to detect the expression of MPL (Thrombopoietin receptor) using 161-GGSG-NLuc-AcV5 supernatant.

[00252] 293FT cells were transiently transfected with a construct encoding human MPL (pLenti-MPL) or left untransfected. Next day morning, approximately 18 hours post-transfection, cells were collected by pipetting up and down in 1.5 ml tubes. The tubes were

spun down at 1500 RPM for 5 minutes. Then the cells were washed once with wash buffer (1% FBS in PBS), followed by incubation with 100 μ l of 161-GGSG-NLuc-AcV5 supernatant. The cells were incubated at 4°C for 1 hour. After the incubation, cells were washed 5 times with wash buffer (1 ml each wash). Finally the pellet was resuspended in 200 μ l wash buffer. Resuspended cells were placed in a 384 well plate in triplicate (25 μ l each). Luciferase activity was measured using a BioTek synergy H4 plate reader after addition of NLuc assay buffer (Promega) containing native coelenterazine (25 μ l each well) directly to each well (one at a time). **FIG. 2** shows significant increased binding of 161-GGSG-NLuc-AcV5 supernatant to cells that had been transfected with MPL as compared to the untransfected cells.

Example 3. Assay to detect the expression of chimeric antigen receptors targeting CD19 and MPL (Thrombopoietin receptor)

[00253] A frequent problem in the field of chimeric antigen receptors is lack of a sensitive and specific assay that can detect cells that express chimeric antigen receptors. To detect the expression of CAR targeting CD19 and MPL, we fused the extracellular domains (ECD) of human CD19 and human MPL, including their signal peptides, in frame with nucleotide sequence encoding a Gly-Gly-Ser-Gly linker, NLuc (without a secretory signal) and an AcV5 epitope tag. In the case of CD19 construct, a FLAG tag was inserted between the signal peptide and the beginning of the extracellular domain. The whole cassette was cloned downstream of the human EF1 α promoter into the lentiviral vector pLenti-EF1 (SEQ ID NO: 842) to make constructs pLenti-EF1-FLAG-CD19-ECD-GGSG-NLuc-AcV5 (SEQ ID NO: 845) and pLenti-EF1-MPL-ECD-GGSG-NLuc-AcV5, respectively. The nucleic acid sequences of the insert fragments are provided in SEQ ID NO: 1170 and SEQ ID NO: 1169, respectively. The protein sequences of the insert fragments are provided in SEQ ID NO: 2370 and SEQ ID NO: 2369, respectively. The constructs were transfected into 293FT cells by calcium phosphate co-precipitation method. Approximately 20 h post-transfection, the cell culture media was replaced with fresh medium. The conditioned media containing the secreted FLAG-CD19-ECD-GGSG-NLuc-AcV5 (also referred to as CD19-GGSG-NLuc-AcV5) and MPL-ECD-GGSG-NLuc-AcV5 (also referred to as MPL-GGSG-NLuc-AcV5) proteins was collected 48-72 h later.

[00254] T-cells engineered to express a chimeric antigen receptor targeting CD19-specific CAR(FMC63-CAR; or FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208]) or a control CAR (4C3-CAR or KSHV-4C3(vL-vH)-Myc-BBz-T2A-Pac(042315-N01)[SEQ ID NO:1230]) targeting a protein encoded by Kaposi's sarcoma associated herpesvirus were incubated with CD19-GGSG-NLuc-AcV5 supernatant at 4°C for 1h followed by extensive washing with cold PBS supplemented with 0.1% Bovine Serum Albumin (BSA). The cells were re-suspended in cold PBS and 30µl of cell suspension was plated per well in a flat-bottom 384 well plate (Greiner, 384 well white plate cat. # 781075) in triplicate. NLuc assay buffer (Promega) containing native coelenterazine (CTZ) (30 µl/well of native coelenterazine diluted in PBS) was added to each well by an automatic dispenser in a well mode using a BioTek synergy H4 plate reader and light emission as a measure of NLuc activity was measured.

[00255] As shown in **FIG. 3**, T cells expressing CD19 (FMC63)-CAR demonstrated strong binding to CD19-GGSG-NLuc-AcV5 as measured by NLuc assay while very little binding was seen on uninfected T cells (UI) or those expressing control (4C3) CAR. The NLuc value on CD19 (FMC63)-CAR cells was more than 85 fold more than on the uninfected or control CAR expressing cells, thereby demonstrating the extreme sensitivity of the assay.

Example 4. Assay to detect the expression of chimeric antigen receptors targeting CD19 and MPL (Thrombopoietin receptor) on 293FT cells that had been transfected with the corresponding constructs.

[00256] 293FT-cells were transiently transfected (in a 24-well plate, 500 ul volume) with lentiviral constructs expressing chimeric antigen receptors targeting CD19 (FMC63-BBZ-PAC-A13; also referred to as FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208]) and MPL (161-BBZ-PAC-R07; also referred to as 161-(vL-vH)-Myc-BBz-T2A-Pac(021015-R07)[SEQ ID NO:1209]) using calcium phosphate cotransfection method or left untransfected. Next day morning, approximately 18 hours post-transfection, cells were collected by pipetting up and down in 1.5 ml tubes. The tubes were spun down at 1500 RPM for 5 minutes. Then the cells were washed once with wash buffer (1% FBS in PBS), followed by incubation with 100 µl of indicated secretory forms of GGS NLuc supernatant. The cells

were incubated at 4°C for 1 hour. After the incubation, cells were washed 5 times with wash buffer (1 ml each wash). Finally the pellet was resuspended in 200 µl wash buffer. Resuspended cells were placed in a 384 well plate in triplicate (25 µl each). Luciferase activity was measured using a BioTek synergy H4 plate reader after addition of NLuc assay buffer (Promega) containing native coelenterazine (25 µl each well) directly to each well (one at a time).

[00257] As shown in **FIG. 4A - FIG. 4B**, 293FT cells expressing CD19 (FMC63-BBZ-PAC)-CAR demonstrated strong binding to CD19-GGSG-NLuc-AcV5 as measured by NLuc assay while very little binding was seen on uninfected T cells (UI) or those expressing control 161-BBZ-PAC CAR. Similarly, 293FT cells expressing 161-BBZ-PAC CAR showed strong binding with MPL-GGSG-NLuc-AcV5 supernatant as compared to untransfected 293FT cells or those transfected with FMC63-BBZ-PAC CAR.

Example 5. Assay to detect the expression of chimeric antigen receptors targeting MPL (Thrombopoietin receptor) on 293FT cells that had been transfected with the different CAR constructs.

[00258] Three different chimeric antigen receptor constructs were made. The construct hTPO-CD28z-Pac consisted of the extracellular receptor binding domains of human thrombopoietin (TPO; SEQ ID NO: 553) fused to the CD28 hinge, transmembrane and cytosolic domains and CD3z cytosolic domains. The construct mTPO-CD28z-Pac was similar in design except that it consisted of the extracellular receptor binding domains of mouse thrombopoietin (TPO; SEQ ID NO: 554) in place of human TPO. Finally, the construct 161-BBz-Pac (161-(vL-vH)-Myc-BBz-T2A-Pac(021015-R07)[SEQ ID NO:1209]) consisted of a scFv fragment (SEQ ID NO: 730) derived from a monoclonal antibody against human MPL fused to a cassette containing the hinge and transmembrane domain of human CD8, the cytosolic domain of human 41BB (CD137) receptor and the cytosolic domain of human CD3z. A control CAR containing a scFv (SEQ ID NO: 718) derived from an irrelevant antibody against a viral protein was constructed as well. All the constructs were transiently transfected into 293FT cells and incubated with MPL-GGSG-NLuc-AcV5 supernatant essentially as described in the preceding example.

[00259] FIG. 5 shows modest binding of MPL-GGSG-NLuc-AcV5 to 293FT cells transfected with hTPO-CD28z-Pac and mTPO-CD28z-Pac constructs and strong binding to 293FT cells transfected with 161-BBz-Pac construct as compared to untransfected cells or cells that had been transfected with control CAR construct.

Example 6. Assay to detect the expression of CD19 using FMC63 scFv fused to GLuc, MLuc7, TurboLuc16, L. Ovaliformis-Luc and Htanneri-Luc

[00260] In Example 1, we demonstrated the ability of FMC63-GGSG-NLuc-AcV5 fusion protein to detect the expression of CD19 antigen. To demonstrate that this ability is not limited to NLuc fusion proteins, we generated lentiviral expression vectors encoding EF1 α promoter driven fusion constructs containing FMC63 scFv fragment linked via a GGS linker to luciferases from *Gaussia princeps* (GLuc), *Lucicutia ovaliformis* (Lovaliformis-Luc or LoLuc), *Heterorhabdus tanneri* (Htanneri Luc or HtLuc), *Metridia longa* 7 (MLuc7) and TurboLuc16 (or TLuc purchased from Life Technologies). All the luciferases lacked any N-terminal secretory peptide. The GLuc, TurboLuc16 and Htanneri-Luc constructs carried a carboxy terminal x3Flag epitope tag, while the Lovaliformis-Luc carried a carboxy terminal AcV5 tag and the MLuc7 construct carried a carboxy terminal HA epitope tag. The MLuc7 cDNA also carried M43L and M110L substitutions. The corresponding substitutions in GLuc have been previously shown to result in Glow type luminescence. The expression vectors encoding the above fusion proteins were transiently transfected into 293FT cells and supernatant containing the fusion proteins collected essentially as described in Example 1. The FMC63-GGSG-NLuc-AcV5 supernatant was generated as described in Example 1 and stored at -20°C prior to use.

[00261] The supernatant containing the fusion proteins were used to detect the expression of CD19 on the surface of RAJI, K562, and HL60 cells. Approximately, 250,000 cells of each kind were mixed with 100 μ l of supernatant (50 μ l of supernatant in case of GLuc and MLuc7) on ice for approximately 1 h. Cells were washed 4 times with PBS containing 1% FBS. Finally after the 4th wash, the cells were re-suspended in cold PBS and 25 μ l of cell suspension was plated per well in a flat-bottom 384 well plate (Greiner, 384 well white plate cat. # 781075) in duplicate. Luc assay buffer containing native coelenterazine (25 μ l/well of native coelenterazine diluted in PBS) was added to each well by an automatic dispenser in a

well mode using a BioTek synergy H4 plate reader and light emission as a measure of Luc activity was measured.

[00262] As shown in **FIG. 6A**, CD8SP-FMC63(vL-vH)-GGs-MLuc-MM-LL-HA(111815-K07)[SEQ ID NO:1137], CD8SP-FMC63(vL-vH)-GGs-TurboLuc16-x3-FLAG(113015-A04)[SEQ ID NO:1135], CD8SP-FMC63(vL-vH)-GGs-GLuc-x3-FLAG(113015-B02)[SEQ ID NO:1136], and CD8SP-FMC63(vL-vH)-GGSG-NLuc-AcV5(021015-U09)[SEQ ID NO:1119] showed approximately 11676 fold, 13691fold, 14956 fold and 4823 fold higher Luc activity on RAJI cells (known to be CD19 positive cell line) as compared to the Luc activity observed with media control. Weak binding with all the fusion proteins was also observed on K562 cells, while very weak binding was seen on HL60 cells. Even though K562 cells showed weak binding as compared to RAJI cells, they showed approximately 285 fold, 766 fold, 829 fold and 51 fold increase in Luc activity with CD8SP-FMC63(vL-vH)-GGs-MLuc-MM-LL-HA(111815-K07)[SEQ ID NO:1137], CD8SP-FMC63(vL-vH)-GGs-TurboLuc16-x3-FLAG(113015-A04)[SEQ ID NO:1135], CD8SP-FMC63(vL-vH)-GGs-GLuc-x3-FLAG(113015-B02)[SEQ ID NO:1136] and CD8SP-FMC63(vL-vH)-GGSG-NLuc-AcV5(021015-U09)[SEQ ID NO:1119], respectively, over that observed with media alone control, thereby demonstrating the extreme sensitivity of the assay. Similarly, even the very weak Luc activity observed on HL60 cells with CD8SP-FMC63(vL-vH)-GGs-MLuc-MM-LL-HA(111815-K07)[SEQ ID NO:1137], CD8SP-FMC63(vL-vH)-GGs-TurboLuc16-x3-FLAG(113015-A04)[SEQ ID NO:1135], CD8SP-FMC63(vL-vH)-GGs-GLuc-x3-FLAG(113015-B02)[SEQ ID NO:1136] and CD8SP-FMC63(vL-vH)-GGSG-NLuc-AcV5(021015-U09)[SEQ ID NO:1119] fusion proteins was still approximately 65 fold, 262 fold, 178 fold and 15 fold higher than that observed with the media alone. CD8SP-4C3-(vL-vH)-GGSG-NLuc-AcV5(040815-H04)[SEQ ID NO:2435], a fusion protein targeting a lytic protein expressed by Kaposi's sarcoma associated was used as a negative control and showed negligible binding, demonstrating the specificity of the assay. Essentially similar results were obtained upon staining with CD8SP-FMC63(vL-vH)-GGs-L-ovaliformis-Luc-AcV5(111815-G04)[SEQ ID NO:1138] and CD8SP-FMC63(vL-vH)-GGs-Htanneri-Luc-x3-FLAG(111815-L05)[SEQ ID NO:1139] fusion proteins (**FIG. 6B** and **FIG. 6C**), although the light produced by these fusion proteins was of lower intensity.

Example 7. scFV-Luc fusion proteins can be used to detect the expression of their target antigen by immunofluorescence staining and flow cytometry

[00263] We also tested the ability of fusion proteins to detect the expression of CD19 by immunofluorescence staining followed by flow cytometry. For this purpose, RAJI, K562 and HL60 cells (300,000 cells each) were stained with the supernatant (100 μ l) containing the CD8SP-FMC63(vL-vH)-GGG-GLuc-x3-FLAG(113015-B02)[SEQ ID NO:1136], CD8SP-FMC63(vL-vH)-GGG-TurboLuc16-x3-FLAG(113015-A04)[SEQ ID NO:1135] and CD8SP-4C3-(vL-vH)-GGSG-NLuc-AcV5(040815-H04)[SEQ ID NO:2435] (-ve control) fusion proteins (1 hour at 4° C). After extensive washes (5 washes, 1mL each) with PBS containing 1% FBS (Wash buffer), cells were stained (1 hour at 4°C) with FITC-conjugated antibody against FLAG-epitope tag (1 μ g/mL; SIGMA, FITC-M2 FLAG antibody; catalog #F4049). After incubation, cells were further washed 3 times with wash buffer. Finally, the cell pellets were re-suspended with 400 μ l wash buffer and were analyzed by flow cytometry on a BD FACS Verse machine.

[00264] As shown in **FIG. 7A-FIG. 7B**, RAJI cells CD8SP-FMC63(vL-vH)-GGG-GLuc-x3-FLAG(113015-B02)[SEQ ID NO:1136] and CD8SP-FMC63(vL-vH)-GGG-TurboLuc16-x3-FLAG(113015-A04)[SEQ ID NO:1135] showed clear increase (approximately 3-5 fold) in fluorescence [as seen by a significant shift in the FITC peak and by mean fluorescence intensity (MFI)], when compared with those stained with CD8SP-4C3-(vL-vH)-GGSG-NLuc-AcV5(040815-H04)[SEQ ID NO:2435] or medium alone (-ve controls). However, the 3-5 fold increase in mean fluorescence observed over the media alone control using Flow cytometry was still weak as compared to the approximately 13691 fold, 14956 fold increase in Luc activity observed using the Luc assay using the same fusion proteins in Example 6. No significant increase in fluorescence was detected on K562 and HL60 cells when stained with CD8SP-FMC63(vL-vH)-GGG-GLuc-x3-FLAG(113015-B02)[SEQ ID NO:1136] and CD8SP-FMC63(vL-vH)-GGG-TurboLuc16-x3-FLAG(113015-A04)[SEQ ID NO:1135] as compared to the CD8SP-4C3-(vL-vH)-GGSG-NLuc-AcV5(040815-H04)[SEQ ID NO:2435] and media control. In contrast, both K562 and HL60 cells showed easily detectable activity, which was several hundred fold higher over the media control, when measured by Luc assay (see **FIG. 6A-FIG. 6C** of Example 6).

[00265] Taken together, these results demonstrate that the greater sensitivity of the Luc assay as compared to the immunofluorescence/Flow cytometry analysis, which has significance for development of companion diagnostics for cellular therapy products, bispecific antibodies and antibody drug conjugates. These results also demonstrate the versatility of the epitope-tagged scFV-Luc-fusion proteins for detection of their target antigens by immunofluorescence staining. The fusion proteins can be tagged with different tags (e.g. FLAG, HA, MYC, T7, AcV5, V5, StrepTagII, poly His, etc.), either singly or in combination, and can then be detected by flouorochrome labeled antibodies against the epitope tags. Alternatively, they can be fused to fluorescent proteins, such as EGFP, RFP and mcherry, for detection directly. Finally, the epitope tags, such as StrepTagII and polyHis tags, can be also used for the purification of the scFV fusion proteins.

Example 8. scFV-NLuc fusion proteins targeting different tumor antigens.

[00266] To demonstrate that the assay described in Example 1 is not limited to detection of CD19 and MPL, lentiviral expression vectors encoding NLuc fusion proteins containing NH2-terminal scFV fragments derived from antibodies targeting different antigens were generated essentially as described for FMC63-NLuc fusion proteins in Example 1. The names of the different scFVs and their target antigens are shown in **Table 1**. In addition NLuc fusion proteins containing extracellular ligand binding domains (amino acid residues 1-187) of human and mouse thrombopoietin (TPO) were generated. The expression vectors encoding the NLuc fusion proteins were transfected in 293FT cells and supernatant containing the secreted fusion proteins were collected as described for Example 1. The supernatant containing the secreted NLuc fusion proteins was used to stain RAJI, K562 and HL60 cell lines and bound proteins detected by addition of coelenterazine (25µl/well of native coelenterazine diluted in PBS), essentially as described for Example 1.

Table 1	
scFV/Ligand	Target
FMC63	CD19
161	MPL
huMA79b-v28	CD79b
Luc90	CS1
HuLuc63	CS1
J6M0	BCMA
Hum291	CD3
Lym1	Lym1
Lym2	Lym2
CD33	CD33
CD138	CD138
NC7	FLT3
BC8	CD45
GRP78-GC18	GRP78
hTPO (1-187)	MPL
mTPO (1-187)	MPL

[00267] As shown in **FIG. 8A-FIG. 8B**, the different fusion proteins showed differential staining on the 3 cell lines. Thus, greater binding of NLuc fusion proteins containing 161, hTPO (1-187), mTPO (1-87), huMA79b-v28, Luc90, HuLuc63, J6M0, Hum291, CD138, BC8 and GRP78-GC18 was observed on K562 cells as compared to HL60 cells. RAJI cells showed highest binding of Lym1 and BC8 fusion proteins, while HL60 cells showed highest binding of CD33 fusion protein.

Example 9. scFv-NLuc Fusion proteins targeting CD30

[00268] A scFV-GGSG-NLuc-x3-Flag fusion protein targeting CD30 were constructed based on Ac10 scFv fragment, which also carried 3 copies of Flag tag at the carboxy-terminal. Robust expression of fusion protein was observed upon transient transfection of the CD8SP-CD30-Ac10(vL-vH)-NLuc-x3-Flag[SEQ ID NO:1142]construct in 293FT cells. The supernatants containing the CD8SP-CD30-Ac10(vL-vH)-NLuc-x3-Flag[SEQ ID NO:1142]fusion protein was tested for binding to Jurkat (CD30⁺) and Kasumi (CD30⁻) cell lines as well as CHTR74, a lymphoma cell line derived from a patient with lymphoma, using the assay described above. **FIG. 9** shows strong binding of CD30-Ac10(vL-vH)-GGSG-NLuc-x3-Flag to Jurkat and CHTR74 cells as compared to Kasumi cells.

Example 10. scFv-NLuc Fusion proteins targeting the human TCR β 1 isoform constant region.

[00269] Two scFV-GGSG-NLuc-x3-Flag fusion proteins targeting the constant region of the β 1 isoform of human T cell receptor β were constructed based on Jov1 and E09 scFv fragments, which also carried three copies of a Flag tag at the carboxy-terminal. Robust expression of fusion proteins was observed upon transient transfection of the two fusion protein encoding constructs in 293FT cells. The supernatants containing the two fusion proteins were tested for binding to Jurkat (TCRB1+) and Kasumi (TCRB⁻) cell lines as well as CHTR74, a lymphoma cell line derived from a patient with lymphoma using the assay described above. **FIG. 10A-FIG. 10B** show strong binding of both CD8SP-TCRB1-Jov1(vL-vH)-GGSG-NLuc-x3-FLAG(030816-E04)[SEQ ID NO:1143] and CD8SP-TCRB1-E09(vL-vH)-GGSG-NLuc-x3-Flag(030816-G01)[SEQ ID NO:1144] fusion proteins to Jurkat cells as compared to Kasumi and CHTR74 cells.

Example 11. Generation of scFV-NLuc-4xFlag-2x-Streptag-8xHis fusion proteins

[00270] Next expression constructs were constructed in which the FMC63-scFV and CD30-Ac10-scFV fragments were fused in frame at their carboxy-termini to an NLuc-4xFlag-2xStreptag-8xHis cassette via a GGSG linker. The NLuc-4xFlag-2x-Streptag-8xHis cassette was in turn fused in frame to a puromycin resistance gene through a T2A ribosome skip sequence. The 4X Flag-2x-Streptag-8xHis tag can be used for detection of the fusion protein by FACS analysis and immunofluorescence microscopy as well as allows purification of the fusion protein. Similar construct in which NLuc was replaced by H-tanneri, TurboLuc16 and GLuc were also constructed. The pLenti-EF1a-CD30-Ac10(vL-vH)-GGSG-NLuc-4xFlag-2xStrep-8xHis-T2A-Pac construct encoding SEQ ID NO:909 was used to generate scFV-NLuc fusion proteins targeting a number of antigens as described in **Table 15** by replacing the CD30 Ac10 scFV region (i.e. vL-vH) with the corresponding regions of the above scFV fragments targeting the different antigens. In addition, camelid derived vHH fragments targeting EGFR1, Her2, Her3, c-Met, IL6R α are fused to the NLuc-4xFlag-2xStrep-8xHis-T2A-Pac cassettes to generate the corresponding secreted fusion proteins. Affibody targeting Her3 and Her2 are also fused to NLuc-4xFlag-2xStrep-8xHis-T2A-Pac cassette to generate the corresponding secreted fusion proteins. In addition, Darpins targeting

Her2 are fused to NLuc-4xFlag-2xStrep-8xHis-T2A-Pac cassettes to generate the corresponding secreted fusion proteins. A number of intracellular proteins, such as gp100, WT1, NY-ESO, tyrosinase, Muc1, MART1, TERT, HTLV-Tax, EBV-derived EBNA-c3, HIV-1 gag and CMV pp65 etc. are expressed on cell surface in association with HLA antigens. T Cell Receptor (TCR) mimics against such peptide antigens have been generated which recognize the peptide in association with the HLA molecules. The DNA encoding the scFV fragments targeting the peptides derived from the above antigens in complex with HLA-A2 (i.e. TCR mimics) are generated and fused in frame to the NLuc-4xFlag-2xStrep-8xHis-T2A-Pac cassettes to generate the corresponding secreted fusion proteins of the above TCR mimics. The sequence of some of the peptide antigens used to generate the TCR mimics is given in **Table 20**.

[00271] **Table 20. HLA-A2 restricted peptides used for generation of TCR mimics-Luc fusion proteins**

Protein	Fragment Name	Amino Acid Seq
gp100	G9-209	(IMDQVPFSV)
gp100	G9-280	(YLEPGPVTV)
gp100	G9-154	(KTWGQYWQV)
MUC1-A7 (130-138)	A7	(NLTISDVSV)
MUC1-D6 (13-21)	D6	(LLTVLTVV)
TAX (11-19)		(LLFGYPVYV)
hTERT(540-548)	T540	(ILAKFLHWL)
hTERT (865-873)	T865	(RLVDDFLLV)
HIV1 gag (77-85)	SL9	(SLYNTVATL)
CMV-pp65(495-503)		(NLVPMVATV)
MART (26-35)		(EAAGIGILTV)
EBNA-3A (596-604)		(SVRDLRL)
EBNA-3c		(LLDFVRFMGV)
WT1		(RMFPNAPYL)
PR1		(VLQELNVTV)
AFP	AFP-158	(FMNKFIYEI)
HPV16-E7	E7 (11-19)	(YMLDLQPET)
Tyrosinase	Tyr (369-377)	(YMDGTMSQV)

Example 12. Generation of scFV- fusion proteins targeting CD33, CD19 and CD20

[00272] Expression constructs encoding multiple scFv-NLuc fusion proteins targeting CD33, CD19 and CD20 antigens were constructed. These fusion proteins differed in the scFv

fragments. They either carried 4xFLAG-x2STREP-8xHis tag or the AcV5 tag. The expression constructs were transfected in 293FT cells and supernatant containing the fusion proteins collected as described above. The supernatants were tested for binding to RAJI (CD19+, CD20+, CD33-) and HL60 (CD19-, CD20-, CD33+) cell lines. **Table 21** shows stronger binding of several CD33 specific scFv-Luc fusion proteins to HL60 cells as compared to RAJI cells. **Table 21** also shows stronger binding of several CD19- and CD20-specific scFv-Luc fusion proteins to RAJI cells as compared to HL60 cells. However, there was a difference in the binding of the different scFv-fusion proteins to the cell lines expressing their target antigens. In particular, several CD19 specific scFv-Luc fusion proteins, such as CD8SP-CD19-huSJ25C1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022817-C04)[SEQ ID NO:865], CD8SP-Ritx-CD19-hB4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022817-F06)[SEQ ID NO:866] and CD8SP-CD19-hA19-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022817-H03)[SEQ ID NO:868] showed relatively weak to negligible binding to RAJI cells. This difference could be due to the difference in the expression and/or antigen-affinity of the different scFv fusion proteins. The CD8SP-CD19-huSJ25C1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022817-C04)[SEQ ID NO:865] was generated by humanizing the sequence of the mouse monoclonal antibody SJ25C1 directed against CD19. The weak binding of this scFv-Luc fusion protein to RAJI cells suggest that it has lost its binding affinity during the process of humanization. Thus, the luciferase based antigen detection method of the instant invention can be used in the process of protein engineering (e.g. antibody engineering) to rapidly identify the candidate clones with different properties. For example, the assay can be used in the process of humanization of antibodies and antibody fragments (e.g., scFvs) to identify clones that have lost the binding affinity to their target antigens. The method can be also used to rapidly identify scFv fragments and other antigen binding domains for incorporation into CARs based on their relative expression levels and binding affinities.

[00273] Table 21: Binding of CD33, CD19 and CD20 fusion proteins

	TARGET	LUC SUPERNATANT	Raji		HL60	
			Mean	STD	Mean	STD
1	CD33	CD8SP-CD33-SGNh2H12-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-	4702	271	67728	9770

	TARGET	LUC SUPERNATANT	Raji		HL60	
		PAC[SEQ ID NO:915]				
2	CD33	CD8SP-CD33-33H4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022217-D03)[SEQ ID NO:917]	171	48	1137	343
3	CD33	CD8SP-CD33-9C3-2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022217-E04)[SEQ ID NO:918]	167	40	533	179
4	CD33	CD8SP-CD33-33H4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022217-D03)[SEQ ID NO:917]	541	83	20472	3736
5	CD33	CD8SP-CD33-9C3-2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022217-E04)[SEQ ID NO:918]	3489	135	32213	3464
6	CD33	CD8SP-CD33-Him3-4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022217-A02)[SEQ ID NO:914]	4195	1052	2919	91
7	CD19	CD8SP-CD19-huSJ25C1-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022817-C04)[SEQ ID NO:865]	1120	30	705	63
8	CD19	CD8SP-Ritx-CD19-hB4-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022817-F06)[SEQ ID NO:866]	562	170	465	164
9	CD19	CD8SP-CD19-hA19-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022817-H03)[SEQ ID NO:868]	673	398	682	92
10	CD19	CD8SP-hCD19-Bu12-(vL-vH)-GGSG-NLuc-AcV5(021417-T05)[SEQ ID NO:1145]	358453	36785	1606	134
11	CD19	CD8SP-hCD19MM-(vL-vH)-GGSG-NLuc-AcV5(021417-U01)[SEQ ID NO:1146]	213706	32671	223	38
12	CD19	CD8SP-huFMC63-11-(vL-vH)-GGSG-NLuc-AcV5(021417-V05)[SEQ ID NO:1147]	283921	29103	1865	42
13	CD19	CD8SP-huFMC63-11-N203Q-(vL-vH)-GGSG-NLuc-AcV5(021417-W05)[SEQ ID NO:1148]	264816	7404	1218	75
14	CD19	CD8SP-huFMC63-11-N203S-(vL-VH)-GGSG-NLuc-AcV5(021417-X05)[SEQ ID NO:1149]	247192	20801	1462	82
15	CD19	CD8SP-CD19-Medi-3649-(vL-vH)-GGSG-NLuc-AcV5(021417-Y05)[SEQ ID NO:1150]	1428	693	298	97
16	CD19	CD8SP-CD19-4G7-Mlu-GSG-NLuc-AcV5(021417-Z04)[SEQ ID NO:1151]	380049	17232	4320	548
17	CD19	CD8SP-RTX-CD19-MOR0028-(vL-vH)-GGSG-NLuc-AcV5(021417-A05)[SEQ ID NO:1152]	210868	10706	1069	29
18	CD19	CD8SP-huFMC63-11-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(020317-E05)[SEQ ID NO:855]	242476	1841	31121	5049
19	CD20	CD8SP-CD20-2H7-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-	11192	302	3716	372

	TARGET	LUC SUPERNATANT	Raji		HL60	
		PAC(022817-E05)[SEQ ID NO:900]				
20	CD20	CD8SP-CD20-GA101-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(121516-G07)[SEQ ID NO:892]	6015	479	3336	317
21		Control	173	68	151	64

Example 13. Generation of scFV- fusion proteins targeting multiple antigens

[00274] Expression constructs encoding multiple scFv-Luc fusion proteins targeting different antigens were constructed. The fusion proteins were expressed in 293FT cells and supernatants tested for binding to BC1 (primary effusion lymphoma) and RAJI (lymphoma) cell lines. **Table 22** shows differential binding of the different scFv-Luc fusion proteins to the two cell lines. For example, CD8SP-CD138-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(041715-D08)[SEQ ID NO:938] showed stronger binding to BC1 cells as compared to RAJI cells, while CD8SP-FMC63(vL-vH)-GGSG-NLuc-x3-FLAG(113015-B02)[SEQ ID NO:1136] showed stronger binding to RAJI cells as compared to BC1 cells.

[00275] **Table 22:** differential binding of scFv-Luc fusion proteins to two cell lines

	TARGET	LUC SUPERNATANT	BC1		RAJI	
			Mean	SD	Mean	SD
1		Control	61	1	60	561.4
2	CD33	CD8SP-CD33(vL-vH)-GGSG-NLuc-AcV5(041715-E01)[SEQ ID NO:1122]	3259	51	1796	70.7
3	CS1	CD8SP-Luc63-(vL-vH)-GGSG-NLuc-AcV5(041415-V08)[SEQ ID NO:1126]	4310	1913	1458	140.0
4	CD138	CD8SP-CD138-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(041715-D08)[SEQ ID NO:938]	1648679	36109	22100	671.0
5	CD19	CD8SP-FMC63(vL-vH)-GGSG-NLuc-x3-FLAG(021216-A06)[SEQ ID NO:1134]	24611	249	746186	2568.2
6	CD19	CD8SP-FMC63(vL-vH)-GGSG-NLuc-x3-FLAG(113015-B02)[SEQ ID NO:1136]	57529	970	893290	14181.0
7	CD30	CD8SP-CD30-5F11(vL-vH)-GGSG-NLuc-x3-Flag(030916-B02)[SEQ ID NO:1141]	7553	419	2651	1003.4
8	IL11Ra	CD8SP-IL11Ra-8E2-Ts107-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(060716-D06)[SEQ ID NO:1026]	948	4	99	16.3
9	IL13Ra2	CD8SP-IL13Ra2-Hu108-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(060716-F03)[SEQ ID NO:1029]	12049	270	5553	59.4

	TARGET	LUC SUPERNATANT	BC1		RAJI	
10	CLEC5A	CD8SP-CLEC5A-8H8F5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(060716-E05)[SEQ ID NO:948]	124	1	56	9.2
11	CLEC5A	CD8SP-CLEC5A-3E12A2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(060716-G01)[SEQ ID NO:949]	765	0	284	1.4
12	CD22	CD8SP-CD22-H22RhoV2ACDRKA-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(060716-H05)[SEQ ID NO:905]	183	6	97	3.5
13	CD19	CD8SP-CD19-4G7-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(062916-C08)[SEQ ID NO:859]	52301	415	397429	6269.9
14	CD32	CD8SP-CD32-Med9-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(081516-W03)[SEQ ID NO:910]	8615	173	1618	14.1

Example 14. Generation of scFv-fusion proteins targeting HIV1 envelop glycoprotein gp120

[00276] Expression constructs encoding multiple scFv-Luc fusion proteins targeting HIV1 envelop glycoprotein gp120 were constructed. The fusion proteins were generated in 293FT cells and supernatants containing the fusion proteins tested for binding to HL2/3 cells which express HIV1 envelop glycoprotein gp120. **Table 23** shows strong binding of the different scFv-Luc fusion proteins to the HL2/3 cell line as compared to media alone (control).

[00277] **Table 23:** Binding of scFv-Luc fusion proteins to HL2/3 cell line

	LUC SUPERNATANT	HL2/3	
		Mean Luc Activity	STD
1	Control	5700	138
2	CD8SP-HIV1-3BNC117-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(093016-Z11)[SEQ ID NO:1016]	8872	107
3	CD8SP-HIV1-VR-C01-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(093016-X10)[SEQ ID NO:1018]	9330	384
4	CD8SP-HIV1-X5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(093016-Y15)[SEQ ID NO:1019]	12868	464

Example 15. Generation of scFv-fusion proteins targeting BCMA

[00278] Expression constructs encoding multiple scFv-Luc fusion proteins targeting BCMA protein were constructed. The fusion proteins were generated in 293FT cells and supernatants containing the fusion proteins tested for binding to U266 cells which express BCMA. **Table 24** shows strong binding (Mean +/- STD) of the different scFv-Luc fusion proteins to the U266 cell line as compared to media alone (control).

[00279] **Table 24:** Binding of the different scFv-Luc fusion proteins to the U266 cell line

	LUC SUPERNATANT	U266	
		Mean Luc Activity	STD
1	Control	766	20
2	CD8SP-BCMA-ET-40-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(092016-A02)[SEQ ID NO:880]	11973	118
3	CD8SP-BCMA-ET-54-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(092016-B04)[SEQ ID NO:881]	10701	68
4	CD8SP-GPRC5D-ET150-5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(092016-C06)[SEQ ID NO:997]	18545	334

[00280] *Example 16. Generation of scFv-fusion proteins targeting PDL1*

[00281] Expression constructs encoding multiple scFv-Luc fusion proteins targeting PDL1 protein were constructed. The fusion proteins were generated in 293FT cells and supernatants containing the fusion proteins tested for binding to L363 cells which express PDL1. **Table 25** shows strong binding (Mean +/- STD) of the different scFv-Luc fusion proteins to the L363 cell line as compared to media alone (control).

[00282] **Table 25** shows strong binding of the different scFv-Luc fusion proteins to the L363 cell line

		LUC SUPERNATANT	L363	
			MEAN LUC ACTIVITY	STD
1		Control	5467	589
2	PDL1	CD8SP-PDL1-SP142-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(100516-H05)[SEQ ID NO:1063]	52984	145
3	PDL1	CD8SP-PDL1-10A5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(100516-F03)[SEQ ID NO:1064]	45924	498
4	PDL1	CD8SP-PDL1-Atezoli-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(100516-G03)[SEQ ID NO:1062]	35251	2252

Example 17. Generation of scFV-Luc fusion proteins targeting Lym1 and Lym2

[00283] Expression constructs encoding scFv-Luc fusion proteins targeting Lym1 and Lym2 proteins were constructed. The fusion proteins were generated in 293FT cells and supernatant containing the fusion protein tested for binding to a panel of cell lines. **Table 26 and Table 27** show differential binding (Mean +/- STD) of the different scFv-Luc fusion proteins to different cell line.

[00284] **Table 26:** Differential binding of CD8SP-Lym1(vL-vH)-GGSG-NLuc-AcV5(040915-J04)[SEQ ID NO:1120] to different cell lines

CD8SP-Lym1(vL-vH)-GGSG-NLuc-AcV5(040915-J04)[SEQ ID NO:1120]		
CELL LINE	MEAN LUC ACTIVITY	STD
BC-1	7386	165
BC-3	11664	587
BCBL-1	7779	669
JSC-1	6766	498
MM1S	1579	13
U266	258	9
BV173	5561	112
Raji	579907	18122

[00285] **Table 27:** Differential binding of CD8SP-Lym2(vL-vH)-GGSG-NLuc-AcV5(040915-K05)[SEQ ID NO:1121] to different cell lines

CD8SP-Lym2(vL-vH)-GGSG-NLuc-AcV5(040915-K05)[SEQ ID NO:1121]		
CELL LINE	MEAN LUC ACTIVITY	STD
BC-1	86094	747
BC-3	1743	144
BCBL-1	3076	427
JSC-1	13256	419
MM1S	1111	131
U266	621	3
BV173	915	25
Raji	54186	344

Example 18. Generation of a scFv-Luc fusion protein targeting FITC

[00286] An expression construct encoding a scFv-Luc fusion protein targeting FITC was constructed. The fusion protein was generated in the supernatant of 293FT cells and supernatant containing the fusion proteins tested for binding to RAJI cells (CD45+/TCRB1-) that had been pre-labeled with a FITC-conjugated CD45 antibody (clone 2D1, eBioscience Dx) or a FITC-conjugated TCRB1(JOV1) antibody (SantaCruz Biotech). For labeling, approximately 100,000 Raji cells were incubated with 100µl of 1:100 diluted FITC-Abs for an hour at 4°C, washed 3 times with PBS and then stained with 100µl of supernatant containing the CD8SP-FITC-E2-HL-(vH-vL)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(030617-H05)[SEQ ID NO:985] fusion protein. Cells were washed 3 times and then assayed for NLuc activity as described before. **Table 28** shows strong binding (Mean +/- STD) of the CD8SP-FITC-E2-HL-(vH-vL)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(030617-H05)[SEQ ID NO:985] fusion protein to the RAJI cells that had been pre-labelled with a FITC-CD45 antibody while only weak binding was observed on RAJI cells that had been pre-labelled with the negative control FITC-TCRB1(JOV1) antibody or had not been prelabelled.

[00287] **Table 28:** binding (Mean +/- STD) of the CD8SP-FITC-E2-HL-(vH-vL)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(030617-H05)[SEQ ID NO:985] fusion protein.

	FITC-Conjugated Ab	Luc Supernatant	RAJI CELLS	
			Mean Luc	STD
1		MEDIA	44.5	5.1962
2		CD8SP-FITC-E2-HL-(vH-vL)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(030617-H05)[SEQ ID NO:985]	1227	508.62
3	TCRB1 (JOVI)-FITC	CD8SP-FITC-E2-HL-(vH-vL)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(030617-H05)[SEQ ID NO:985]	1435	760.21
4	CD45-FITC	CD8SP-FITC-E2-HL-(vH-vL)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(030617-H05)[SEQ ID NO:985]	57647	11022

Example 19. Generation of scFv-fusion proteins targeting Alpha Fetal Protein (AFP)

[00288] Expression constructs encoding multiple scFv-Luc fusion proteins targeting an AFP-derived peptide in complex with MHC class I (HLA-A2) were constructed. The fusion proteins were generated in 293FT cells and supernatants containing the fusion proteins tested for binding to HepG2 cells which express AFP and HLA-A2. **Table 29** shows strong binding (Mean +/- STD) of the different scFv-Luc fusion proteins to the HepG2 cell line as compared to media alone (control). These results demonstrate that scFv-Luc fusion proteins can be used to detect cell surface expression of intracellular protein antigens when they are presented in association with MHC molecules.

[00289] **Table 29:** Binding (Mean +/- STD) of the different scFv-Luc fusion proteins to the HepG2 cell line

	Luc Supernatant	HepG2 Cells	
		Mean	STD
1	Media	23	1
2	CD8SP-AFP-61-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022217-K04)[SEQ ID NO:869]	2460	95

		HepG2 Cells	
		Mean	STD
3	CD8SP-AFP-76-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022217-L01)[SEQ ID NO:870]	1417	20

Example 20. Generation of scFv- fusion proteins targeting multiple antigens

[00290] Expression constructs encoding multiple scFv-Luc fusion proteins targeting different antigens were constructed. The fusion proteins were expressed in 293FT cells and supernatants tested for binding to a panel of cell lines. **Table 30** shows differential binding of the different scFv-Luc fusion proteins to the different cell lines.

[00291] **Table 30:** Differential binding of scFv-Luc fusion proteins to the different cell lines

TARGET	LUC SUPERNATANT	BC-3	BCBL-1	JSC-1	MM1S	U266	BV173
CD19	CD8SP-FMC63(vL-vH)-GGSG-NLuc-AcV5(021015-U09)[SEQ ID NO:1119]	2517	5687	3209	4040	2358	181552
CD19	CD8SP-FMC63(vL-vH)-GGSG-NLuc-x3-FLAG(021216-A06)[SEQ ID NO:1134]	2770	7817	5698	4601	6206	234481
CD19	CD8SP-FMC63(vL-vH)-GGSG-NLuc-x3-FLAG(113015-B02)[SEQ ID NO:1136]	5656	12685	12371	9053	9897	212871
CD19	CD8SP-CD19-4G7-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(062916-C08)[SEQ ID NO:859]	6182	13985	13285	10443	6540	121316
CD19	CD8SP-CD19-hA19-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(022817-H03)[SEQ ID NO:868]	52	76	106	67	68	125
CD22	CD8SP-CD22-H22RhoV2ACDRKA-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(060716-H05)[SEQ ID NO:905]	305	4410	638	1437	2806	413

TARGET	LUC SUPERNATANT	BC-3	BCBL-1	JSC-1	MM1S	U266	BV173
CD30	CD8SP-CD30-Ac10-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(031816-D02)[SEQ ID NO:909]	10348	25996	2093	2097	1464	1697
CD33	CD8SP-CD33(vL-vH)-GGSG-NLuc-AcV5(041715-E01)[SEQ ID NO:1122]	241	653	328	1995	509	13114
Lym1	CD8SP-Lym1(vL-vH)-GGSG-NLuc-AcV5(040915-J04)[SEQ ID NO:1120]	11664	7779	6766	1579	258	5561
Lym2	CD8SP-Lym2(vL-vH)-GGSG-NLuc-AcV5(040915-K05)[SEQ ID NO:1121]	1743	3076	13256	1111	621	915
GRP78	CD8SP-GRP78-GC18-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC[SEQ ID NO:1005]	106	173	183	304	141	142
GRP78	CD8SP-GRP78-GC18(vL-vH)-NLuc-AcV5(040915-M06)[SEQ ID NO:1130]	1142	2531	1587	2444	1348	1112
CD79b	CD8SP-huMA79bv28-(vL-vH)-GGSG-NLuc-AcV5(041515-U04)[SEQ ID NO:1128]	806	1826	1136	988	635	1173
CS1	CD8SP-Luc90-(vL-vH)-NLuc-AcV5(041515-W03)[SEQ ID NO:1127]	22092	80958	11134 2	53941	1722 6	603
BCMA	CD8SP-BCMA-J6MO-(vL-vH)-NLuc-AcV5(041615-X08)[SEQ ID NO:1124]	23676	11444	24487	58922	2713 2	855
Amyloid	SP-Amyloid-158-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC[SEQ ID NO:875]	3112	5686	8229	8168	3933	6220
FLT3	CD8SP-FLT3-NC7-(vL-vH)-NLuc-AcV5(041715-C06)[SEQ ID NO:1125]	1146	2821	1167	1023	714	13551
CD45	CD8SP-CD45-BC8-(vL-vH)-GGSG-NLuc-AcV5(042115-A09)[SEQ ID NO:1129]	2507	35103	67510	1086	1097 7	1041
CD138	CD8SP-CD138-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(041715-D08)[SEQ ID	72070	250385	13280 8	496546	1232 482	14690

TARGET	LUC SUPERNATANT	BC-3	BCBL-1	JSC-1	MM1S	U266	BV173
	NO:938]						
IL13Ra 2	CD8SP-IL13Ra2-Hu108-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(060716-F03)[SEQ ID NO:1029]	60	86	80	72	67	151
IL13Ra 2	CD8SP-IL13Ra2-Hu108-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(060716-F03)[SEQ ID NO:1029]	3375	4623	4211	3792	3469	2509
CLEC5 A	CD8SP-CLEC5A-8H8F5-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(060716-E05)[SEQ ID NO:948]	31	42	45	45	34	53
CLEC5 A	CD8SP-CLEC5A-3E12A2-(vL-vH)-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(060716-G01)[SEQ ID NO:949]	311	7804	502	526	634	677

Example 21. Generation of a CD20-ECx2-ECD-Luc fusion protein to detect CD20-specific CARs

[00292] CD20 is a type III membrane protein with two extracellular loops. The CD20-ECx2-ECD-GGSG-TurboLuc16-4xFlag-2xStreptag-8xHis-T2A-Pac(060816-I04)[SEQ ID NO:1196] fusion construct was generated which contains a CD20-ECx2-ECD domain as represented by SEQ ID NO: 546. The fusion proteins were expressed in 293FT cells and supernatants tested for binding to parental NK92MI and Jurkat cells and those expressing a CD20-specific CAR CD20-GA10-(vL-vH)-MYC-BBz-T2A-PAC(111815-A03)[SEQ ID NO:1220]. **Table 31** shows strong binding of CD20-ECx2-ECD-GGSG-TurboLuc16-4xFlag-2xStreptag-8xHis-T2A-Pac(060816-I04)[SEQ ID NO:1196] fusion protein to NK92MI and Jurkat cells expressing the CD20-specific CAR CD20-GA10-(vL-vH)-MYC-BBz-T2A-PAC(111815-A03)[SEQ ID NO:1220]. These results demonstrate that Luc fusion proteins can be generated using the extracellular domains of type III membrane proteins and such fusion proteins can be used to detect CARs directed against the said type III membrane protein.

[00293] **Table 31:** Binding of CD20 fusion protein to CD20-specific CARs.

	CAR	Luc Supernatant	NK92MI		Jurkat	
			Mean	STD	Mean	STD
1	None	CD20-ECx2-ECD-GGSG-TurboLuc16-4xFlag-2xStreptag-8xHis-T2A-Pac(060816-I04)[SEQ ID NO:1196]	510	146	484	320
2	CD20-GA10-(vL-vH)-MYC-BBz-T2A-PAC(111815-A03)[SEQ ID NO:1220]	CD20-ECx2-ECD-GGSG-TurboLuc16-4xFlag-2xStreptag-8xHis-T2A-Pac(060816-I04)[SEQ ID NO:1196]	1417	415	7606	5791

Example 22. Generation of a BCMA-ECD-Luc fusion protein to detect BCMA-specific CARs

[00294] A BCMA-ECD-NLuc fusion construct was generated. The fusion protein was expressed in 293FT cells and supernatants tested for binding to parental NK92MI cells and those expressing BCMA- or CD19-specific CAR constructs. **Table 32** shows strong binding of BCMA-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(103116-Q07)[SEQ ID NO:1201] fusion protein to NK92MI cells expressing the BCMA-specific CAR J6MO-(vL-vH)-MYC-28z-T2A-Pac(031915-G02)[SEQ ID NO:1213] as compared to parental NK92MI cells. As a control the binding of FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170] fusion protein, which contains the extracellular domain of CD19, to NK92MI cells expressing the BCMA-specific CAR J6MO-(vL-vH)-MYC-28z-T2A-Pac(031915-G02)[SEQ ID NO:1213] and CD19-specific CAR FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208] was examined. The FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170] fusion protein failed to show significant binding to NK92MI cells expressing the BCMA-specific CAR J6MO-(vL-vH)-MYC-28z-T2A-Pac(031915-G02)[SEQ ID NO:1213] but showed strong binding to NK92MI cells expressing the CD19-specific CAR FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208], thereby demonstrating the specificity of the assay.

[00295] **Table 32:** Binding of BCMA fusion protein to BCMA-specific CARs

Cell Line	CAR-Construct	Luc Supernatant	Mean	STD
NK92MI	J6MO-(vL-vH)-MYC-28z-T2A-Pac(031915-G02)[SEQ ID NO:1213]	BCMA-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(103116-Q07)[SEQ ID NO:1201]	602	102
NK92MI	None	BCMA-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(103116-Q07)[SEQ ID NO:1201]	89	10
NK92MI	J6MO-(vL-vH)-MYC-28z-T2A-Pac(031915-G02)[SEQ ID NO:1213]	FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170]	57	3
NK92MI	FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208]	FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170]	217346	49538

Example 23. Generation of a SLAMF7-CS1-ECD-GGSG-NLuc fusion protein to detect SLAMF7/CS1-specific CARs

[00296] A SLAMF7-CS1-ECD-GGSG-NLuc fusion construct was generated. The fusion protein was expressed in 293FT cells and supernatants tested for binding to parental Jurkat cells and those expressing SLAMF7/CS1-specific CAR constructs. **Table 33** shows strong binding of SLAMF7-CS1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(062816-A01)[SEQ ID NO:1202] fusion protein to Jurkat cells expressing the SLAMF7/CS1-specific CARs HuLuc64-(vL-vH)-Myc-BBz-T2A-Pac(052616-U05)[SEQ ID NO:1228] and Luc90-(vL-vH)-MYC-CD28z-T2A-Pac(031915-F03)[SEQ ID NO:1229] as compared to parental Jurkat cells.

[00297] **Table 33:** Binding of SLAMF7-CS1 fusion protein to SLAMF7-CS1-specific CARs

	Jurkat Cells			
	CAR	Luc supernatant	Mean	STD
1	None	SLAMF7-CS1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(062816-A01)[SEQ ID NO:1202]	354	19

	Jurkat Cells			
	CAR	Luc supernatant	Mean	STD
2	HuLuc64-(vL-vH)-Myc-BBz-T2A-Pac(052616-U05)[SEQ ID NO:1228]	SLAMF7-CS1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(062816-A01)[SEQ ID NO:1202]	2932	103
3	Luc90-(vL-vH)-MYC-CD28z-T2A-Pac(031915-F03)[SEQ ID NO:1229]	SLAMF7-CS1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(062816-A01)[SEQ ID NO:1202]	1143	24

Example 24. Generation of a SLAMF7-CS1-ECD-GGSG-NLuc fusion protein to detect SLAMF7/CS1-specific CARs

[00298] 293FT cells were transiently transfected with a SLAMF7/CS1-specific CAR Luc90-(vL-vH)-MYC-CD28z-T2A-Pac(031915-F03)[SEQ ID NO:1229] and a CD19-specific CAR FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208]. Approximately 24 hours later, cells were stained with supernatants containing SLAMF7-CS1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(062816-A01)[SEQ ID NO:1202] and FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170] fusion proteins. **Table 34** shows strong binding of SLAMF7-CS1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(062816-A01)[SEQ ID NO:1202] to 293FT cells expressing the SLAMF7/CS1-specific CAR Luc90-(vL-vH)-MYC-CD28z-T2A-Pac(031915-F03)[SEQ ID NO:1229] and strong binding of FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170] to cells expressing the CD19-specific CAR FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208]. However, no significant binding of FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170] to 293FT cells transfected with the SLAMF7/CS1-specific CAR was observed, thereby demonstrating the specificity of the assay.

[00299] **Table 34:** Binding of SLAMF7-CS1 and CD19 fusion protein to SLAMF7-CS1-specific and CD19-specific CARs

293FT CELLS			
CAR	Luc Supernatant	Mean Luc Activity	STD
Luc90-(vL-vH)-MYC-CD28z-T2A-Pac(031915-F03)[SEQ ID NO:1229]	SLAMF7-CS1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(062816-A01)[SEQ ID	100008	2740

293FT CELLS			
CAR	Luc Supernatant	Mean Luc Activity	STD
	NO:1202]		
FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208]	SLAMF7-CS1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(062816-A01)[SEQ ID NO:1202]	37801	624
Luc90-(vL-vH)-MYC-CD28z-T2A-Pac(031915-F03)[SEQ ID NO:1229]	FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170]	3827	738
FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208]	FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170]	237068	7993

Example 25. Generation of a CDH1-ECD-GGSG-NLuc fusion protein to detect CDH1/CD324-specific CARs

[00300] The CDH1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(062816-G02)[SEQ ID NO:1189] fusion protein was expressed in the supernatant of 293FT cells as described previously. Another batch of 293FT cells were transiently transfected with *CDH1/CD324*-specific CARs CD324-SC10-6-(vL-vH)-MYC-BBz-T2A-PAC(052516-B07)[SEQ ID NO:1221] and CD324-hSC10-17-(vL-vH)-MYC-BBz-T2A-PAC(052516-A07)[SEQ ID NO:1222] and a CD19-specific CAR Luc90-(vL-vH)-MYC-CD28z-T2A-Pac(031915-F03)[SEQ ID NO:1229]. Approximately 24 hours later, cells were stained with supernatants containing CDH1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(062816-G02)[SEQ ID NO:1189] and FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170] fusion proteins. **Table 35** shows strong binding of CDH1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(062816-G02)[SEQ ID NO:1189] fusion protein to 293FT cells expressing the CDH1/CD324-specific CARs CD324-SC10-6-(vL-vH)-MYC-BBz-T2A-PAC(052516-B07)[SEQ ID NO:1221] and CD324-hSC10-17-(vL-vH)-MYC-BBz-T2A-PAC(052516-A07)[SEQ ID NO:1222] and strong binding of FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170] to cells expressing the CD19-specific CAR FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208]. However, no significant binding of FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-

C04)[SEQ ID NO:1170] to 293FT cells transfected with the CDH1-specific CARs was observed, thereby demonstrating the specificity of the assay.

[00301] **Table 35:** Binding of CDH1-ECD fusion protein to CDH1/CD324-specific CARs

293FT CELLS			
CAR	Luc Supernatant	Mean	STD
CD324-SC10-6-(vL-vH)-MYC-BBz-T2A-PAC(052516-B07)[SEQ ID NO:1221]	CDH1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(062816-G02)[SEQ ID NO:1189]	2904158	44104
CD324-hSC10-17-(vL-vH)-MYC-BBz-T2A-PAC(052516-A07)[SEQ ID NO:1222]	CDH1-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(062816-G02)[SEQ ID NO:1189]	2018952	23267
CD324-SC10-6-(vL-vH)-MYC-BBz-T2A-PAC(052516-B07)[SEQ ID NO:1221]	FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170]	16409	1159
CD324-hSC10-17-(vL-vH)-MYC-BBz-T2A-PAC(052516-A07)[SEQ ID NO:1222]	FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170]	5186	232
FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208]	FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170]	237068	7993

Example 26. Generation of a CD200R-ECD-GGSG-NLuc fusion protein to detect CD200R-specific CARs

[00302] 293FT cells were transiently transfected with a CD200R-specific CAR CD200R-huDx182-(vL-vH)-Myc-BBz-T2A-PAC[SEQ ID NO:1218] and a CD19-specific CAR FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208]. Approximately 24 hours later, cells were stained with supernatants containing CD200R-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(082616-C07)[SEQ ID NO:1190] and FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170] fusion proteins. **Table 36** shows strong binding of CD200R-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(082616-C07)[SEQ ID NO:1190] to 293FT cells expressing the CD200R-specific CAR CD200R-huDx182-(vL-vH)-Myc-BBz-T2A-PAC[SEQ ID NO:1218] and strong binding of FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170] to cells expressing the CD19-specific CAR FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208].

However, no significant binding of FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170] to 293FT cells transfected with the CD200R-specific CAR was observed, thereby demonstrating the specificity of the assay.

[00303] **Table 36:** CD200R-ECD fusion protein to detect CD200R-specific CARs

293FT CELLS			
CAR	Luc Supernatant	Mean	STD
CD200R-huDx182-(vL-vH)-Myc-BBz-T2A-PAC[SEQ ID NO:1218]	CD200R-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(082616-C07)[SEQ ID NO:1190]	18736	3038
CD200R-huDx182-(vL-vH)-Myc-BBz-T2A-PAC[SEQ ID NO:1218]	FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170]	1063	165
FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208]	FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170]	237068	7993

Example 27. Generation of fusion protein containing different Luc fragments to detect CD19-specific CARs

[00304] Multiple constructs containing the extracellular domain of CD19 in fusion with different Luc reporter fragments were generated. The fusion protein was expressed in 293FT cells and supernatants tested for binding to parental NK92MI cells and those expressing a CD19-specific CAR construct. **Table 37** shows strong binding of all fusion protein to NK92MI cells expressing the CD19-specific CAR FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208] as compared to parental NK92MI cells. However, there was a difference in signal intensity observed with the different fusion proteins with the FLAG-CD19-ECD-GGS-LoLuc-AcV5(111616-L02)[SEQ ID NO:1174] and CD19-ECD-GGS-HtLuc-x3Flag(111616-N05)[SEQ ID NO:1176] fusion proteins showing relatively weak signal. This is probably due to the lower brightness of LoLuc and HtLuc as compared to the other luciferases.

[00305] **Table 37:** Binding of fusion proteins to NK92MI cells expressing the CD19-specific CAR

	LUC SUPERNATANT	NK92-MI-Parental		NK92MI-FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208]	
		MEAN	STD	MEAN	STD
1	FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170]	80	23	217346	49538
2	FLAG-CD19-ECD-GGS-PaLuc1-HA-Streptag-3xHA-8xHis-T2A-pac(111616-K07)[SEQ ID NO:1173]	70	9	14064	2569
3	FLAG-CD19-ECD-GGS-Turboluc16-4xFlag-2xStreptag-8xHis-T2A-Pac(111616-J04)[SEQ ID NO:1172]	350	44	14172	2197
4	FLAG-CD19-ECD-GGS-LoLuc-AcV5(111616-L02)[SEQ ID NO:1174]	77	7	185	75
5	FLAG-CD19-ECD-GGS-MLuc7-MM-LL-HA(111616-M07)[SEQ ID NO:1175]	229	93	9045	1637
6	CD19-ECD-GGS-HtLuc-x3Flag(111616-N05)[SEQ ID NO:1176]	88	56	205	155

Example 28. Generation of fusion protein containing different Luc fragments to detect CD33-specific CARs

[00306] Multiple constructs containing the extracellular domain of CD33 in fusion with different Luc reporter fragments were generated. The fusion protein was expressed in 293FT cells and supernatants tested for binding to parental NK92MI cells and those expressing a CD33-specific CAR construct. **Table 38** shows stronger binding of all fusion protein to NK92MI cells expressing the CD33-specific CAR CD33-(vL-vH)-Myc-28z-T2A-Pac(032415-O01)[SEQ ID NO:1211] as compared to the parental NK92MI cells. However, there was a difference in signal intensity observed with the different fusion proteins with the CD33-ECD-LoLuc-AcV5(100616-L07) [SEQ ID NO:1180] and CD33-ECD-HtLuc-x3Flag(100616-N05)[SEQ ID NO:1182] fusion proteins showing relatively weak signal. This is probably due to the lower brightness of LoLuc and HtLuc as compared to the other luciferases.

[00307] **Table 38:** Binding of fusion proteins to NK92MI cells expressing the CD33-specific CAR

	LUC SUPERNATANT	NK92-MI-Parental		NK92MI-CD33-(vL-vH)-Myc-28z-T2A-Pac(032415-O01)[SEQ ID NO:1211]	
		MEAN	STD	MEAN	STD
1	CD33-ECD-GGSG-NLuc-4xFlag-2xStreptag-8xHis-T2A-Pac(060816-A02)[SEQ ID NO:1177]	68	15	2453	315
2	CD33-ECD-GGSG-Turboluc16-4xFlag-2xStreptag-8xHis-T2A-Pac(100616-J04)[SEQ ID NO:1178]	5679	318	2148794	150613
3	CD33-ECD-PaLuc1-HA-Streptag-3xHA-8xHis-T2A-pac(100616-K07)[SEQ ID NO:1179]	404	46	285377	20648
4	CD33-ECD-LoLuc-AcV5(100616-L07)[SEQ ID NO:1180]	55	3	2158	426
5	CD33-ECD-MLuc7-MM-LL-HA(100616-M02)[SEQ ID NO:1181]	1318	140	955672	48712
6	CD33-ECD-HtLuc-x3Flag(100616-N05)[SEQ ID NO:1182]	87	21	1871	1925

Example 29. Generation of fusion protein containing different Luc fragments to detect CD138-specific CARs

[00308] Multiple constructs containing the extracellular domain of CD138 in fusion with different Luc reporter fragments were generated. The fusion protein was expressed in 293FT cells and supernatants tested for binding to parental NK92MI cells and those expressing CD138-specific CARs construct. **Table 39** shows stronger binding of all fusion protein to NK92MI cells expressing the CD138-specific CARs CD138-(vL-vH)-Myc-28z-T2A-Pac(032415-N06)[SEQ ID NO:1210] and CD138-(vL-vH)-MYC-BBz-T2A-Pac(052115-A02)[SEQ ID NO:1212] as compared to the parental NK92MI cells. However, there was a difference in signal intensity observed with the different fusion proteins with the CD138-SDC1-ECD-GGS-LoLuc-AcV5(101216-G04)[SEQ ID NO:1186] fusion protein

showing relatively weak signal. This is probably due to the lower brightness of LoLuc as compared to the other luciferases.

[00309] **Table 39:** Binding of fusion proteins to NK92MI cells expressing the CD138-specific CAR

	LUC SUPERNATANT	NK92-MI-Parental		CD138-(vL-vH)-Myc-28z-T2A-Pac(032415-N06)[SEQ ID NO:1210]		CD138-(vL-vH)-MYC-BBz-T2A-Pac(052115-A02)[SEQ ID NO:1212]	
		MEAN	STD	MEAN	STD	MEAN	STD
1	CD138-SDC1-ECD-GGS-Turboluc16-4xFlag-2xStreptag-8xHis-T2A-Pac(101216-E04) [SEQ ID NO:1184]	49199	5404	136266	29578	186935	9085
2	CD138-SDC1-ECD-GGS-PaLuc1-HA-Streptag-3xHA-8xHis-T2A-pac(101216-F04)[SEQ ID NO:1185]	6988	503	26350	3272	38653	4182
3	CD138-SDC1-ECD-GGS-LoLuc-Luc-AcV5(101216-G04) [SEQ ID NO:1186]	185	50	850	110	782	59
4	CD138-SDC1-ECD-GGS-MLuc7-MM-LL-HA(111616-H04)[SEQ ID NO:1187]	1303	217	47765	4000	72967	1037

Example 30: Sensitivity of Luc fusion protein assay for detecting CAR-expressing cells.

[00310] To determine the sensitivity of the Luc fusion protein assay for detecting CAR-expressing cells, the indicated numbers of NK92MI cells stably expressing a CD19-specific CAR FMC63-(vL-vH)-Myc-BBz-T2A-Pac(112014-A13)[SEQ ID NO:1208] in 25µl volume were added to 1 million parental NK92MI cells (in 25µl volume) in a 96 well U-bottom plate. Next, 100µl of 293FT supernatant containing FLAG-CD19-ECD-GGSG-NLuc-AcV5(062615-C04)[SEQ ID NO:1170] fusion protein was added to each well and cells were mixed by pipetting up and down a couple of times and then incubated for 1 hour on ice. After incubation, cells were spun down at 1400 rpm for 5 minutes at 4°C, followed by 5 washes with ice-cold wash buffer (0.5% FBS in PBS). Finally, the pellets were resuspended in 40µl. 30µl of the resuspended mixture was transferred to one well of a 384 well plate. The experiment was conducted in quadruplicate. The luminescence was read by adding 30µl of

1X CTZ assay buffer. **FIG. 11** shows that 100 CAR-expressing NK92MI cells in the background of a million parental cells can be easily detected by the assay giving a sensitivity of 1 CAR cell in the background of 10,000 non-CAR cells. The sensitivity of the assay can be further improved by using purified and concentrated fusion protein.

Example 31: Single-step Purification of Luc fusion protein containing Streptag using Strep-tactin.

[00311] 293FT cells were transfected with expression vector encoding FLAG-CD19-ECD-GGS-Turboluc16-4xFlag-2xStreptag-8xHis-T2A-Pac(111616-J04)[SEQ ID NO:1172]. The supernatant containing the secreted fusion protein was collected between 48-72h post-transfection and secreted protein purified over a Strep-Tactin resin (IBA Lifesciences) and following the manufacturer's recommendations. The Turboluc activity in the starting supernatant and each step of purification was determined by adding 30µl of 1X CTZ assay buffer to 30µl of 1:500 diluted samples. **FIG. 12** shows that FLAG-CD19-ECD-GGS-Turboluc16-4xFlag-2xStreptag-8xHis-T2A-Pac(111616-J04)[SEQ ID NO:1172] fusion protein can be purified in a single step from crude supernatant with excellent yield and recovery. The purified sample (Elute 2) was tested for binding to RAJI cells and was found to be biologically active in this assay.

Example 32: Use of Protein L-Luc fusion protein for detection of CAR

[00312] Two NLuc fusion constructs containing an N-terminal Protein L coding region downstream of a CD8 signal peptide were constructed. The two constructs are identical except that the construct CD8SP-Protein-L-2-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(101916-P03)[SEQ ID NO:1207] lacks a single amino acid in the Protein L coding region, which is present in the construct CD8SP-Protein-L-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(112316-Q02)[SEQ ID NO:1206]. 293FT cells were transfected with the two constructs and supernatants containing the secreted fusion proteins collected between 48 to 72 hours post-transfection. The secreted proteins were tested for binding to Jurkat cells expressing the CS1-specific CAR HuLuc64-(vL-vH)-Myc-BBz-T2A-Pac(052616-U05)[SEQ ID NO:1228] that comprises a humanized scFv fragment as its antigen recognition domain. **FIG. 13** shows effective binding of the CD8SP-Protein-L-2-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(101916-P03)[SEQ ID NO:1207] and

CD8SP-Protein-L-GGSG-NLuc-4xFLAG-x2STREP-8xHis-T2A-PAC(112316-Q02)[SEQ ID NO:1206] fusion proteins to Jurkat cells expressing the CS1-specific CAR HuLuc64-(vL-vH)-Myc-BBz-T2A-Pac(052616-U05)[SEQ ID NO:1228] as compared to parental cells (Jurkat cells-UI). These results demonstrate that Protein L containing Luc fusion proteins can be used to detect a kappa light chain containing CAR where the kappa light chain binds to Protein L.

[00313] References

1. Kohn DB et al. (2011). "CARs on track in the clinic." Mol. Ther. 19(3): 432.
2. De Oliveira, S. N., J. Wang, C. Ryan, S. L. Morrison, D. B. Kohn and R. P. Hollis (2013). "A CD19/Fc fusion protein for detection of anti-CD19 chimeric antigen receptors." Journal of translational medicine 11: 23.
3. Hajek, R., S. A. Okubote and H. Svachova (2013). "Myeloma stem cell concepts, heterogeneity and plasticity of multiple myeloma." Br J Haematol 163(5): 551-564.

[00314] The various methods and techniques described above provide a number of ways to carry out the application. Of course, it is to be understood that not necessarily all objectives or advantages described can be achieved in accordance with any particular embodiment described herein. Thus, for example, those skilled in the art will recognize that the methods can be performed in a manner that achieves or optimizes one advantage or group of advantages as taught herein without necessarily achieving other objectives or advantages as taught or suggested herein. A variety of alternatives are mentioned herein. It is to be understood that some preferred embodiments specifically include one, another, or several features, while others specifically exclude one, another, or several features, while still others mitigate a particular feature by inclusion of one, another, or several advantageous features.

[00315] Furthermore, the skilled artisan will recognize the applicability of various features from different embodiments. Similarly, the various elements, features and steps discussed above, as well as other known equivalents for each such element, feature or step, can be employed in various combinations by one of ordinary skill in this art to perform methods in accordance with the principles described herein. Among the various elements,

features, and steps some will be specifically included and others specifically excluded in diverse embodiments.

[00316] Although the application has been disclosed in the context of certain embodiments and examples, it will be understood by those skilled in the art that the embodiments of the application extend beyond the specifically disclosed embodiments to other alternative embodiments and/or uses and modifications and equivalents thereof.

[00317] Preferred embodiments of this application are described herein, including the best mode known to the inventors for carrying out the application. Variations on those preferred embodiments will become apparent to those of ordinary skill in the art upon reading the foregoing description. It is contemplated that skilled artisans can employ such variations as appropriate, and the application can be practiced otherwise than specifically described herein. Accordingly, many embodiments of this application include all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the application unless otherwise indicated herein or otherwise clearly contradicted by context.

[00318] All patents, patent applications, publications of patent applications, and other material, such as articles, books, specifications, publications, documents, things, and/or the like, referenced herein are hereby incorporated herein by this reference in their entirety for all purposes, excepting any prosecution file history associated with same, any of same that is inconsistent with or in conflict with the present document, or any of same that may have a limiting affect as to the broadest scope of the claims now or later associated with the present document. By way of example, should there be any inconsistency or conflict between the description, definition, and/or the use of a term associated with any of the incorporated material and that associated with the present document, the description, definition, and/or the use of the term in the present document shall prevail.

[00319] It is to be understood that the embodiments of the application disclosed herein are illustrative of the principles of the embodiments of the application. Other modifications that can be employed can be within the scope of the application. Thus, by way of example, but not of limitation, alternative configurations of the embodiments of the application can be

utilized in accordance with the teachings herein. Accordingly, embodiments of the present application are not limited to that precisely as shown and described.

[00320] Various embodiments of the invention are described above in the Detailed Description. While these descriptions directly describe the above embodiments, it is understood that those skilled in the art may conceive modifications and/or variations to the specific embodiments shown and described herein. Any such modifications or variations that fall within the purview of this description are intended to be included therein as well. Unless specifically noted, it is the intention of the inventors that the words and phrases in the specification and claims be given the ordinary and accustomed meanings to those of ordinary skill in the applicable art(s).

[00321] The foregoing description of various embodiments of the invention known to the applicant at this time of filing the application has been presented and is intended for the purposes of illustration and description. The present description is not intended to be exhaustive nor limit the invention to the precise form disclosed and many modifications and variations are possible in the light of the above teachings. The embodiments described serve to explain the principles of the invention and its practical application and to enable others skilled in the art to utilize the invention in various embodiments and with various modifications as are suited to the particular use contemplated. Therefore, it is intended that the invention not be limited to the particular embodiments disclosed for carrying out the invention.

[00322] While particular embodiments of the present invention have been shown and described, it will be obvious to those skilled in the art that, based upon the teachings herein, changes and modifications may be made without departing from this invention and its broader aspects and, therefore, the appended claims are to encompass within their scope all such changes and modifications as are within the true spirit and scope of this invention.

[00323] The present application and invention further includes the subject matter of the following numbered clauses.

[00324] 1. A method for detecting the presence of an antigen in a subject in need thereof comprising: providing a fusion protein comprising an antigen specific single chain

antibody or a nanobody (camel or LAMA vHH fragment), or an affibody or a DARPIN or a non-immunoglobulin antigen binding domain or the extracellular domain of a receptor or the extracellular domain of a ligand fused to a reporter; providing a sample from the subject; exposing the sample to the fusion protein for sufficient time and under conditions to allow the fusion protein to bind to the sample; washing the sample to remove any unbound fusion protein; and assaying the activity of the reporter; wherein a change in reporter activity relative to a reference value is indicative of the presence of an antigen in the subject.

[00325] 2. A method for detecting the expression of a chimeric antigen receptor or a T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor on the cells of a subject comprising: providing a fusion protein comprising a reporter fused to an extracellular domain of the antigen targeted by the chimeric antigen receptor or a T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor; providing a sample from the subject; exposing the sample to the fusion protein for sufficient time and under conditions to allow the fusion protein to bind to the sample; washing the sample to remove any unbound fusion protein; and assaying the activity of the reporter; wherein a change in reporter activity relative to a reference value is indicative of the expression of the chimeric antigen receptor or an exogenous T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor on the cells of the subject.

[00326] 3. The method of clauses 1 or 2, wherein the reporter is a non-secretory form of a luciferase.

[00327] 4. The method of clause 3, wherein the non-secretory form of luciferase is obtained from copepods, deep sea shrimp or homologs or orthologs thereof or mutants or derivatives thereof.

[00328] 5. The method of clause 4, wherein the copepods are selected from the group consisting of any one or more of *Gaussia princeps*, *Pleuromamma abdominalis*, *Metridia pacifica*, *Metridia curticauda*, *Metridia asymmetrica*, *Metridia okhotensis*, *Metridia longa*, *Lucicutia ovaliformis*, *Heterorhabdus tanneri*, and *Pleuromamma scutullata*.

[00329] 6. The method of clause 3, wherein the luciferase is any one or more of GLuc, NLuc, MLuc7, PaLuc1, PaLuc2, MpLuc1, MoLuc1, MoLuc2, MLuc39, HtLuc, HtLuc2, PsLuc1, Lucifcutia ovaliformis Luc (LoLuc), LoLuc1-3, Renilla or TLuc (TurboLuc16) or derivatives thereof.

[00330] 7. The method of clauses 1 or 2, wherein the reporter activity is assayed by exposing the target cells to a luciferase specific substrate.

[00331] 8. The method of clause 7, wherein the luciferase-specific substrate is coelentraine or a derivative thereof.

[00332] 9. The method of clause 7, wherein the luciferase-specific substrate is imidazopyrazinone substrate (furimazine) or a derivative thereof.

[00333] 10. The method of clause 1, wherein the luciferase activity is measured by any one or more of methods for measuring light production such as a luminometer, Western blotting, ELISA, x-ray films, microscopy, BRET, nanoBRET, or combinations thereof.

[00334] 11. The method of clause 1, wherein the antigen is expressed on a cancer cell.

[00335] 12. The method of clause 11, wherein the antigen is any one or more of but are not limited to 4-1BB, 5T4, adenocarcinoma antigen, alpha-fetoprotein, BAFF, B-lymphoma cell, C242 antigen, CA-125, carbonic anhydrase 9 (CA-IX), C-MET, CCR4, CD152, CD19, CD20, CD200, CD22, CD221, CD23 (IgE receptor), CD28, CD30 (TNFRSF8), CD33, CD4, CD40, CD44 v6, CD51, CD52, CD56, CD74, CD80, CEA, CNTO888, CTLA-4, DR5, EGFR, EpCAM, CD3, FAP, fibronectin extra domain-B, folate receptor 1, GD2, GD3 ganglioside, glycoprotein 75, GPNMB, HER2/neu, HGF, human scatter factor receptor kinase, IGF-1 receptor, IGF-I, IgG1, L1-CAM, IL-13, IL-6, insulin-like growth factor I receptor, integrin $\alpha 5\beta 1$, integrin $\alpha v\beta 3$, MORAb-009, MS4A1, MUC1, mucin CanAg, N-glycolylneuraminic acid, NPC-1C, PDGF-R α , PDL192, phosphatidylserine, prostatic carcinoma cells, RANKL, RON, ROR1, SCH 900105, SDC1, SLAMF7, thrombopoietin receptor, TAG-72, tenascin C, TGF beta 2, TGF- β , TRAIL-R1, TRAIL-R2, tumor antigen CTAA16.88, VEGF-A, VEGFR-1, VEGFR2, MPL, or vimentin.

[00336] 13. The method of clauses 1 or 11, wherein the antigen is the complex of an HLA molecule with a peptide antigen.

- [00337] 14. The method of clause 13, wherein the HLA molecule is HLA-A2.
- [00338] 15. The method of clause 13, wherein the antigen is derived from any one or more of the following proteins but are not limited to NY-ESO, Mart-1, TERT, MAGE, gp100, Tyrosinase, WT1, PR1, HTLV1-Tax, EBNA3c, CMV-pp65, HIV1 gag and HTLV1-Tax.
- [00339] 16. The method of clause 2, wherein the CAR or a T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor is expressed on an immune cell.
- [00340] 17. The method of clause 16, wherein the immune cell is a T cell.
- [00341] 18. The method of clause 17, wherein the immune cell is a CD4 T cell.
- [00342] 19. The method of clause 17, wherein the immune cell is a CD8 T cell.
- [00343] 20. The method of clause 17, wherein the immune cell is a Treg cell.
- [00344] 21. The method of clause 17, wherein the immune cell is a naive T cell.
- [00345] 22. The method of clause 17, wherein the immune cell is a memory T cell.
- [00346] 23. The method of clause 17, wherein the immune cell is central memory T cell.
- [00347] 24. The method of clause 17, wherein the immune cell is an effector memory T cell.
- [00348] 25. The method of clause 16, wherein the immune cell is an NK cell
- [00349] 26. The method of clause 2, wherein the CAR or a T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor is expressed on a stem cell.
- [00350] 27. The method of clause 2, wherein the CAR or a T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor is expressed on a hematopoietic stem cell.

[00351] 28. The method of clause 2, wherein the CAR or a T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor is expressed on an induced pluripotent stem cell.

[00352] 29. The method of any one of clauses 2, 16 or 26-28, wherein the target antigen of CAR or a T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor is any one or more of but are not limited to 4-1BB, 5T4, adenocarcinoma antigen, alpha-fetoprotein, BAFF, B-lymphoma cell, C242 antigen, CA-125, carbonic anhydrase 9 (CA-IX), C-MET, CCR4, CD152, CD19, CD20, CD123, CD200, CD22, CD221, CD23 (IgE receptor), CD28, CD30 (TNFRSF8), CD33, CD4, CD40, CD44 v6, CD51, CD52, CD56, CD74, CD80, CEA, CNTO888, CTLA-4, DR5, EGFR, EpCAM, CD3, FAP, fibronectin extra domain-B, folate receptor 1, GD2, GD3 ganglioside, glycoprotein 75, GPNMB, HER2/neu, HGF, human scatter factor receptor kinase, IGF-1 receptor, IGF-I, IgG1, L1-CAM, IL-13, IL-6, insulin-like growth factor I receptor, integrin $\alpha 5\beta 1$, integrin $\alpha v\beta 3$, Lym1, Lym2, FLT3, MORAb-009, MS4A1, MUC1, mucin CanAg, N-glycolylneuraminic acid, NPC-1C, PDGF-R α , PDL192, phosphatidylserine, prostatic carcinoma cells, RANKL, RON, ROR1, SCH 900105, SDC1, SLAMF7, thrombopoietin receptor (MPL), TAG-72, tenascin C, TGF beta 2, TGF- β , TRAIL-R1, TRAIL-R2, tumor antigen CTAA16.88, VEGF-A, VEGFR-1, VEGFR2, or vimentin.

[00353] 30. The method of any one of clauses 2, 16 or 26-28, wherein the target antigen of CAR or a T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor is a complex of a HLA molecule with a peptide antigen.

[00354] 31. The method of clause 11, wherein the target antigen of CAR or a T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor is the complex of an HLA molecule with a peptide antigen.

[00355] 32. The method of clauses 30 and 31, wherein the HLA molecule is HLA-A2.

[00356] 33. The method of clauses 30 and 31, wherein the antigen is derived from any one or more of but are not limited to NY-ESO, Mart-1, TERT, MAGE, gp100, Tyrosinase, WT1, PR1, HTLV1-Tax, EBNA3c, CMV-pp65, HIV1 gag and HTLV1-Tax.

- [00357] 34. The method of clause 1, wherein the subject has cancer or an immune disorder and is being assessed for immune therapy.
- [00358] 35. The method of clause 34, wherein the immune therapy involves CAR T cells.
- [00359] 36. The method of clause 34, wherein the immune therapy involves exogenous T cell receptor expressing T cells.
- [00360] 37. The method of clause 34, wherein the immune therapy involves synthetic T cell receptor therapy or a chimeric T cell receptor therapy or a hybrid T cell receptor therapy.
- [00361] 38. The method of clause 34, wherein the immune therapy involves antibody therapy.
- [00362] 39. The method of clause 34, wherein the immune therapy involves antibody-drug conjugate therapy.
- [00363] 40. The method of clause 34, wherein the immune therapy involves bispecific antibody.
- [00364] 41. The method of clause 34, wherein the immune therapy involves BiTE.
- [00365] 42. The method of clause 34, wherein the immune therapy involves DART.
- [00366] 43. The method of clause 34, wherein the immune therapy involves NK cells.
- [00367] 44. The method of clause 34, wherein the immune therapy involves stem cells.
- [00368] 45. The method of clause 34, wherein the immune therapy involves pluripotent stem cells.
- [00369] 46. The method of clause 34, wherein the immune therapy involves induced pluripotent stem cells. The method of clause 34, wherein the immune therapy involves stem cell transplantation.
- [00370] 47. The method of clauses 1 or 2, wherein the fusion protein further comprises a tag.

[00371] 48. The method of clause 47, wherein the tag is any one or more of chitin binding protein (CBP), glutathione-S-transferase (GST), polyhistidine (His) tag, FLAG tag, HA tag, Myc tag, V5 tag, AcV5 tag, Streptag or a combination thereof.

[00372] 49. The method of clause 1, wherein the reference value is the reporter activity in any one or more of (i) control cells or samples that do not express the target antigen; (ii) control cells or samples that express reporter but are not treated with the test agent(s); (iii) controls cells or samples that are not treated with the substrate for the reporter; (iv) control cells or samples that do express the target antigen but are treated with a fusion protein directed against an antigen that is not expressed on the target cells; (v) a combination thereof.

[00373] 50. The method of clause 2, wherein the reference value is the reporter activity in any one or more of (i) cells that do not express the CAR, or an exogenous T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor; (ii) cells that express the CAR, or an exogenous T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor but are treated with fusion protein which is not targeted by the CAR, or an exogenous T cell receptor or a synthetic T cell receptor or a chimeric T cell receptor or a hybrid T cell receptor; (iii) cells that are not treated with the substrate for the reporter.

[00374] 51. The method of clauses 1 or 2, wherein the assay is performed on samples outside the patient (in vitro).

[00375] 52. The method of clauses 1 or 2, wherein the assay is performed in vivo.

CLAIMS

1. A method for detecting expression of a chimeric antigen receptor (CAR), comprising:
obtaining a sample from a subject in need determination of expression of the CAR;
contacting the sample with a fusion protein comprising a reporter fused to an extracellular domain of an antigen targeted by the CAR; and
assaying the activity of the reporter; wherein presence of reporter activity or increase in reporter activity relative to a reference value is indicative of the expression of the CAR in the sample.
2. The method of claim 1, wherein the reporter is a non-secretory form of a luciferase.
3. The method of claim 2, wherein the non-secretory form of luciferase is obtained from copepods, deep sea shrimp or homologs or orthologs thereof or mutants or derivatives thereof.
4. The method of claim 3, wherein the copepods are selected from the group consisting of any one or more of *Gaussia princeps*, *Pleuromamma abdominalis*, *Metridia pacifica*, *Metridia curticauda*, *Metridia asymmetrica*, *Metridia okhotensis*, *Metridia longa*, *Lucicutia ovaliformis*, *Heterorhabdus tanneri*, and *Pleuromamma scutullata*.
5. The method of claim 2, wherein the luciferase is any one or more of GLuc, NanoLuc (NLuc), MLuc7, HtLuc, LoLuc, PaLuc1, PaLuc2, MpLuc1, McLuc1, MaLuc1, MoLuc1, MoLuc2, MLuc39, PsLuc1, LocLuc1-3, HtLuc2 Renilla, TurboLuc16 (TLuc) or homologs or orthologs thereof or mutants or functional derivatives thereof.
6. The method of claim 1, wherein the reporter activity is assayed by exposing the target cells to a luciferase specific substrate.
7. The method of claim 6, wherein the luciferase-specific substrate is coelentraine or a derivative thereof.
8. The method of claim 6, wherein the luciferase-specific substrate is imidazopyrazinone substrate (furimazine) or a derivative thereof.
9. The method of claim 1, wherein the CAR is expressed on an immune cell.

10. The method of claim 9, wherein the immune cell is a T cell.
11. The method of claim 9, wherein the immune cell is a CD4 T cell.
12. The method of claim 9, wherein the immune cell is a CD8 T cell.
13. The method of claim 9, wherein the immune cell is a Treg cell.
14. The method of claim 9, wherein the immune cell is a naive T cell.
15. The method of claim 9, wherein the immune cell is a memory T cell.
16. The method of claim 9, wherein the immune cell is central memory T cell.
17. The method of claim 9, wherein the immune cell is an effector memory T cell.
18. The method of claim 9, wherein the immune cell is an NK cell.
19. The method of claim 1, wherein the CAR is expressed on a stem cell.
20. The method of claim 1, wherein the CAR is expressed on a hematopoietic stem cell.
21. The method of claim 1, wherein the CAR is expressed on an induced pluripotent stem cell.
22. The method of any one of claim 1, wherein the target antigen of the CAR is any one or more of but are not limited to CD19, CD20, CS1/SLAMF7, BCMA, CD123, CD33, MPLLym1, Lym2, CD200R, 5T4, adenocarcinoma antigen, alpha-fetoprotein, BAFF, B-lymphoma cell, C242 antigen, CA-125, carbonic anhydrase 9 (CA-IX), C-MET, CCR4, CD152, , CD200, CD22, CD221, CD23 (IgE receptor), CD28, CD30 (TNFRSF8), , CD4, CD40, CD44 v6, CD51, CD52, CD56, CD74, CD80, CEA, CNTO888, CTLA-4, DR5, EGFR, EpCAM, CD3, FAP, fibronectin extra domain-B, folate receptor 1, GD2, GD3 ganglioside, glycoprotein 75, GPNMB, HER2/neu, HGF, human scatter factor receptor kinase, IGF-1 receptor, IGF-I, IgG1, L1-CAM, IL-13, IL-6, insulin-like growth factor I receptor, integrin $\alpha 5\beta 1$, integrin $\alpha v\beta 3$ FLT3, MORAb-009, MS4A1, MUC1, mucin CanAg, N-glycolylneuraminic acid, NPC-1C, PDGF-R α , PDL192, phosphatidylserine, prostatic carcinoma cells, RANKL, RON, ROR1, SCH 900105, SDC1, SLAMF7, (MPL), TAG-72, tenascin C, TGF beta 2,

- TGF- β , TRAIL-R1, TRAIL-R2, tumor antigen CTAA16.88, VEGF-A, VEGFR-1, VEGFR2, or vimentin.
23. The method of claim 1, wherein the target antigen of CAR is a complex of a HLA molecule with a peptide antigen.
 24. The method of claim 23, wherein the HLA molecule is HLA-A2.
 25. The method of claim 1, wherein the subject has cancer or an immune disorder.
 26. The method of claim 25, wherein the subject is being assessed for immune therapy.
 27. The method of claim 26, wherein the immune therapy comprises CAR T cells
 28. The method of claim 1, wherein the fusion protein further comprises a tag.
 29. The method of claim 28, wherein the tag is any one or more of chitin binding protein (CBP), glutathione-S-transferase (GST), polyhistidine (His) tag, FLAG tag, HA tag, Myc tag, V5 tag, AcV5 tag, Streptag or a combination thereof.
 30. The method of claim 1, wherein the reference value is the reporter activity in any one or more of (i) cells that do not express the CAR; (ii) cells that express the CAR but are treated with fusion protein which is not targeted by the CAR; (iii) cells that are not treated with the substrate for the reporter; or (iv) combinations thereof.
 31. The method of claim 1, wherein the reporter is fused to the extracellular domain of an antigen targeted by the chimeric antigen receptor through a covalent bond.
 32. The method of claim 1, wherein the reporter is fused to the extracellular domain of an antigen targeted by the chimeric antigen receptor through a non-covalent bond.
 33. The method of claim 1, wherein the reporter is fused to the extracellular domain of an antigen targeted by the chimeric antigen receptor through an intermediate molecule.
 34. The method of claim 1, wherein a single molecule of the reporter is fused to the extracellular domain of an antigen targeted by the chimeric antigen receptor.
 35. The method of claim 1, wherein more than one molecules of the reporter are fused to the extracellular domain of an antigen targeted by the chimeric antigen receptor.

36. The method of claim 1, wherein the assay is performed in vitro.
37. The method of claim 1, wherein the assay is performed in vivo.
38. A kit for practicing the invention.

Fig. 1A

GLuc-GGSG--161-N-Luc--V-Mpl1 target cells--4-6-15




Fig. 1B

GLuc-GGSG--FMC63-N-Luc--V-Mpl1 target cells--4-6-15




Fig. 2

hPML transient transfection with 161 GGS-NLuc
binding assay:
092915





**Fig. 3****Fig. 4A****Fig. 4B**



Fig. 5**Fig. 6A**





Fig. 6B**Fig. 6C**

Fig. 7A**Fig. 7B**

Mean Fluorescence Intensity			
	Raji	K562	HL60
Medium alone	22	66	8
4C3 GGS NLuc AcV5 H04	22	47	8
FMC63 GGS TurboLuc16 x 3 flag A04	59	49	8
FMC63 GGS GLuc x 3 flag B02	111	49	8

Fig. 8A







Fig. 10A**Fig. 10B**




Fig. 11**Fig. 12**

Fig. 13

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2017/025602

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 35/17; C07K 14/705 (2017.01)

CPC - A61K 35/17; C07K 14/7051; C07K 2317/66; C07K 2319/30; C07K 2319/61 (2017.02)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

See Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

USPC - 424/93.2; 424/133.1; 435/8; 435/328; 435/6.17; 435/7.1 (keyword delimited)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

See Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2015/0376296 A1 (SLOAN-KETTERING INSTITUTE FOR CANCER RESEARCH) 31 December 2015 (31.12.2015) entire document	38
—		-----
Y		1-37
Y	BURBELO et al. "Luciferase Immunoprecipitation Systems for Measuring Antibodies in Autoimmune and Infectious Diseases," Translational Research, 01 September 2014 (01.09.2014), Vol. 165, No. 2, Pgs. 325-335. entire document	1-37
Y	US 2013/0273582 A1 (DALY et al) 17 October 2013 (17.10.2013) entire document	3, 4
Y	US 2014/0194307 A1 (PROMEGA CORP) 10 July 2014 (10.07.2014) entire document	8
Y	WO 2014/127261 A1 (THE REGENTS OF THE UNIVERSITY OF CALIFORNIA) 21 August 2014 (21.08.2014) entire document	20, 21
Y	WO 2012/135854 A2 (SCHEINBERG et al) 04 October 2012 (04.10.2012) entire document	24
Y	ZUBAIR et al. "Microfluidic LIPS for serum antibody detection: Demonstration of a rapid test for HSV-2 infection," Biomed Microdevices, 31 December 2011 (31.12.2011), Vol. 13, No. 6, Pgs. 1053-1062. entire document	28, 29
Y	US 2015/0219647 A1 (US DEPARTMENT OF HEALTH AND HUMAN SERVICES (HHS)) 06 August 2015 (06.08.2015) entire document	31, 33, 35
Y	US 4,959,303 A (MILBURN et al) 25 September 1990 (25.09.1990) entire document	32

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

07 June 2017

Date of mailing of the international search report

28 JUN 2017

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, VA 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Blaine R. Copenheaver

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774