



(72) RICIGLIANO, Joseph W., US


(72) ARANEO, Barbara A., US

(71) Pharmadigm, Inc., US

(51) Int.Cl.<sup>6</sup> C12N 15/85, A61K 31/70, A61K 48/00

(30) 1995/09/19 (08/530,529) US

(54) **ADN DE RECOMBINAISON CONTENANT UN ELEMENT  
REGULATEUR A SPECIFICITE MUSCULAIRE POUR DES  
IMMUNISATIONS OU POUR DES THERAPIES GENIQUES**  
(54) **DNA CONSTRUCT COMPRISING A MUSCLE SPECIFIC  
REGULATORY ELEMENT FOR IMMUNIZATION OR GENE  
THERAPY**



(57) L'invention concerne un ADN de recombinaison qui est utile pour des immunisations ou pour des thérapies géniques. Le produit de recombinaison de l'invention contient des éléments régulateurs à spécificité musculaire, comme par exemple un promoteur ou un promoteur et un ou plusieurs éléments activateurs, et une séquence d'ADN fonctionnant sous le contrôle d'éléments régulateurs à spécificité musculaire. Plusieurs séquences d'ADN peuvent être incorporées dans l'ADN de recombinaison. Dans une forme d'exécution, la séquence d'ADN code pour un antigène, un déterminant antigénique ou un épitope d'un antigène.

(57) The present invention is directed to a DNA construct which is useful for immunization or gene therapy. The construct of the invention comprises muscle specific regulatory elements, such as a promoter or a promoter and one or more enhancer elements, and a DNA sequence under control of the muscle specific regulatory elements. Several DNA sequences may be incorporated into the DNA construct. In one embodiment, the DNA sequence codes for an antigen, antigenic determinant or an epitope of an antigen. In a second embodiment, the DNA sequence is a normal muscle gene which is effected in a muscle disease. In a third embodiment, the



(21) (A1) **2,231,806**  
(86) 1996/09/19  
(87) 1997/03/27

Dans une seconde forme d'exécution, la séquence d'ADN est un gène musculaire normal qui est atteint d'une maladie musculaire. Dans une troisième forme d'exécution, la séquence d'ADN est un ADN anti-sens pour bloquer un gène musculaire anormal. Dans une quatrième forme d'exécution, la séquence d'ADN code pour une protéine qui circule dans le sang du mammifère ou dans son système lymphatique. La présente invention est utile pour combattre les effets de maladies touchant les muscles en assurant une expression d'un gène normal ou en bloquant l'expression d'un gène anormal dans les cellules musculaires, en assurant l'expression hétérologue d'un transgène qui code pour une protéine circulante ou une protéine qui modifie une maladie dans laquelle les muscles ne sont pas impliqués d'une manière fondamentale. L'invention est également utile pour effectuer des vaccinations.

DNA sequence is an antisense for blocking an abnormal muscle gene. In a fourth embodiment, the DNA sequence codes for a protein which circulates in the mammalian blood or lymphatic systems. The present invention is useful for ameliorating the effects of diseases of muscle by expression of the normal gene or blocking abnormal gene expression within muscle cells, for the heterologous expression of a transgene which codes for a circulating protein or a protein which modifies a disease state in which muscle is not primarily involved and for vaccine development.



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau


## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

|                                                                                                                                                                          |  |                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| (51) International Patent Classification <sup>6</sup> :<br><b>C12N 15/85, 15/11, 15/90, A61K 39/00</b>                                                                   |  | (11) International Publication Number: <b>WO 97/11190</b>                                                                                                                                                                                                                                                                                                                                                                                            |
| <b>A3</b>                                                                                                                                                                |  | (43) International Publication Date: 27 March 1997 (27.03.97)                                                                                                                                                                                                                                                                                                                                                                                        |
| (21) International Application Number: PCT/US96/14829                                                                                                                    |  | (81) Designated States: AL, AM, AU, BB, BG, BR, CA, CN, CU, CZ, EE, FI, GE, HU, IL, IS, JP, KG, KP, KR, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). |
| (22) International Filing Date: 19 September 1996 (19.09.96)                                                                                                             |  |                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| (30) Priority Data:<br>08/530,529 19 September 1995 (19.09.95) US                                                                                                        |  |                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| (71) Applicant: PARADIGM BIOSCIENCES, INC. [US/US]; 2401 Foothill Drive, Salt Lake City, UT 84109 (US).                                                                  |  |                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| (72) Inventors: RICIGLIANO, Joseph, W.; 1880 Laurelhurst Drive, Salt Lake City, UT 84108 (US). ARANEO, Barbara, A.; 2434 Kentucky Avenue, Salt Lake City, UT 84117 (US). |  |                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| (74) Agents: IHNEN, Jeffrey, L. et al.; Venable, Baetjer, Howard & Civiletti, Suite 1000, 1201 New York Avenue, N.W., Washington, DC 20005 (US).                         |  | <b>Published</b><br><i>With international search report.<br/>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>                                                                                                                                                                                                                                                    |
|                                                                                                                                                                          |  | (88) Date of publication of the international search report:<br>26 June 1997 (26.06.97)                                                                                                                                                                                                                                                                                                                                                              |

(54) Title: DNA CONSTRUCT COMPRISING A MUSCLE SPECIFIC REGULATORY ELEMENT FOR IMMUNIZATION OR GENE THERAPY

## (57) Abstract

The present invention is directed to a DNA construct which is useful for immunization or gene therapy. The construct of the invention comprises muscle specific regulatory elements, such as a promoter or a promoter and one or more enhancer elements, and a DNA sequence under control of the muscle specific regulatory elements. Several DNA sequences may be incorporated into the DNA construct. In one embodiment, the DNA sequence codes for an antigen, antigenic determinant or an epitope of an antigen. In a second embodiment, the DNA sequence is a normal muscle gene which is effected in a muscle disease. In a third embodiment, the DNA sequence is an antisense for blocking an abnormal muscle gene. In a fourth embodiment, the DNA sequence codes for a protein which circulates in the mammalian blood or lymphatic systems. The present invention is useful for ameliorating the effects of diseases of muscle by expression of the normal gene or blocking abnormal gene expression within muscle cells, for the heterologous expression of a transgene which codes for a circulating protein or a protein which modifies a disease state in which muscle is not primarily involved and for vaccine development.



TITLE OF THE INVENTION**DNA CONSTRUCT COMPRISING A MUSCLE SPECIFIC REGULATORY ELEMENT FOR IMMUNIZATION OR GENE THERAPY**BACKGROUND OF THE INVENTION

The present invention is directed to a DNA construct which can be used for either direct or indirect gene therapy. The DNA construct contains muscle specific regulatory elements and a DNA sequence which encodes an antigen for immunization or a protein for gene therapy or the DNA sequence is an antisense sequence for gene therapy.

The publications and other materials used herein to illuminate the background of the invention, and in particular, cases to provide additional details respecting the practice, are incorporated herein by reference, and for convenience, are referenced by author and date in the following text and respectively grouped in the appended List of References.

Initial efforts toward postnatal gene therapy have relied on indirect means of introducing new genetic information into tissues: target cells are removed from the body, infected with viral vectors carrying the new genetic information and then replanted into the body. (Ledley, 1987; Eglitis and Anderson, 1988; Friedmann, 1989). However, indirect gene transfer is not useful for many applications of gene therapy. In these instances, direct introduction of genes into tissues *in vivo* is desired.

Several gene transfer systems have been developed to directly or indirectly introduce genes into tissues *in vivo*. These systems include viral and nonviral transfer methods. A number of viruses have been used as gene transfer vectors, including papovaviruses, e.g., SV40 (Madzak *et al.*, 1992), adenovirus (Berkner, 1992; Berkner *et al.*, 1988; Gorziglia and Kapikian, 1992; Quantin *et al.*, 1992; Rosenfeld *et al.*, 1992; Wilkinson *et al.*, 1992; Stratford-Perricaudet *et al.*, 1990), vaccinia virus (Moss, 1992), adeno-associated virus (Muzyczka, 1992; Ohi *et al.*, 1990; Srivastava, 1993), herpesviruses including HSV and EBV (Margolskee, 1992; Johnson *et al.*, 1992; Fink *et al.*, 1992; Breakfield and Geller, 1987; Freese *et al.*, 1990), and retroviruses of avian (Brandyopadhyay and Temin, 1984; Petropoulos *et al.*, 1992), murine (Miller, 1992; Miller *et al.*, 1985; Sorge *et al.*, 1984; Mann and Baltimore, 1985; Miller *et al.*, 1988), and human origin (Shimada *et al.*, 1991; Helseth *et al.*, 1990; Page *et al.*, 1990; Buchsacher and Panganiban, 1992). Most human gene therapy protocols have been based on disabled murine retroviruses.

Nonviral gene transfer methods known in the art include chemical techniques such as calcium phosphate coprecipitation (Graham and van der Eb, 1973; Pellicer *et al.*, 1980); mechanical techniques, for example microinjection (Anderson *et al.*, 1980; Gordon *et al.*, 1980; Brinster *et al.*, 1981; Constantini and Lacy, 1981); membrane fusion-mediated transfer via liposomes (Felgner *et al.*, 1987; Wang and Huang, 1989; Kaneda *et al.*, 1989; Stewart *et al.*, 1992; Nabel *et al.*, 1990; Lim *et al.*, 1992); and direct DNA uptake and receptor-mediated DNA transfer (Wolff *et al.*, 1990; Wu *et al.*, 1991; Zenke *et al.*, 1990; Wu *et al.*, 1989b; Wolff *et al.*, 1991; Wagner *et al.*, 1990; Wagner *et al.*, 1991; Cotten *et al.*, 1990; Curiel *et al.*, 1991a; Curiel *et al.*, 1991b). Viral-mediated gene transfer can be combined with direct *in vivo* gene transfer using liposome delivery, allowing one to direct the viral vectors to the tumor cells and not into the surrounding nondividing cells. Alternatively, the retroviral vector producer cell line can be injected into tumors (Culver *et al.*, 1992). Injection of producer cells would then provide a continuous source of vector particles. This technique has been approved for use in humans with inoperable brain tumors.

In an approach which combines biological and physical gene transfer methods, plasmid DNA of any size is combined with a polylysine-conjugated antibody specific to the adenovirus hexon protein, and the resulting complex is bound to an adenovirus vector. The trimolecular complex is then used to infect cells. The adenovirus vector permits efficient binding, internalization, and degradation of the endosome before the coupled DNA is damaged.

Liposome/DNA complexes have been shown to be capable of mediating direct *in vivo* gene transfer. While in standard liposome preparations the gene transfer process is nonspecific, localized *in vivo* uptake and expression have been reported in tumor deposits, for example, following direct *in situ* administration (Nabel, 1992).

Direct gene transfer into mammalian somatic tissues *in vivo* is a developing technology with potential applications in human gene therapy. The principal advantages of such an approach are the simplicity and safety of the techniques. Three types of direct gene transfer methodology have been developed: particle bombardment, liposome-mediated deliver and naked DNA transfer. In particle bombardment methods, first applied to the transformation of plant tissue (Klein *et al.*, 1987), the DNA-coated particles are accelerated to high velocity so that they are able to penetrate target organs, tissues or single cells efficiently. Gene transfer to various mammalian somatic tissue has been effectively achieve *in vivo*, *ex vivo* and *in vitro* with particle

bombardment (Yang *et al.*, 1990). Liposome-mediated gene transfer is also an effective method for *in vivo* gene transfer. For example, DNA-liposome complexes have been used for direct gene transfer to human melanoma cells (Nabel *et al.*, 1993).

A challenge to the development of vaccines against viruses such as influenza A or human immunodeficiency virus (HIV), against which neutralizing antibodies are generated, is the diversity of the viral envelope proteins among different isolates or strains. Because CTLs in both mice and humans are capable of recognizing epitopes derived from conserved internal viral proteins (Townsend *et al.*, 1989) and are thought to be important in the immune response against viruses (Taylor *et al.* 1986), efforts have been directed toward the development of CTL vaccines capable of providing heterologous protection against different viral strains. CD8<sup>+</sup> CTLs kill virally infected cells when their T cell receptors recognize viral peptides associated with major histocompatibility complex (MHC) class I molecules (Germain, 1981). These peptides are derived from endogenously synthesized viral proteins, regardless of the protein's location or function in the virus. Thus, by recognition of epitopes from conserved viral proteins, CTLs may provide cross-strain protection. Peptides capable of associating with MHC class I molecules for CTL recognition originate from proteins that are present in or pass through the cytoplasm or endoplasmic reticulum (Yewdell *et al.* 1989). Therefore, in general, exogenous proteins, which enter the endosomal processing pathway (as in the case of antigens presented by MHC class II molecules), are not effective at generating CD8<sup>+</sup> CTL responses.

Most efforts to generate CTL responses have either used replicating vectors to produce the protein antigen in the cell (Hahn *et al.* 1992) or have focused on the introduction of peptides into the cytosol (Collins *et al.* 1992). Both of these approaches have limitations that may reduce their usefulness as vaccines. Retroviral vectors have restrictions on the site and structure of polypeptides that can be expressed as fusion proteins and still maintain the ability of the recombinant virus to replicate (Miller *et al.* 1992), and the effectiveness of vectors such as vaccinia for subsequent immunizations may be compromised by immune responses against the vectors themselves (Cooney *et al.* 1991). Also, viral vectors and modified pathogens have inherent risks that may hinder their use in humans (Mascola *et al.* 1989). Furthermore, the selection of peptide epitopes to be presented is dependent on the structure of an individual's MHC antigens, and peptide vaccines may therefore have limited effectiveness due to the diversity of MHC haplotypes in outbred populations (Townsend *et al.* 1989; Taylor *et al.* 1986;

Germain, 1981). Hence, immunization with nonreplicating plasmid DNA encoding viral proteins may be advantageous because no infectious agent is involved, no assembly of virus particles is required, and determinant selection is permitted. Because the sequence of nucleoprotein (NP) is conserved among various strains of influenza (Gammelin *et al.* 1989; Gorman *et al.* 1991), protection was achieved here against subsequent challenge by a virulent strain of influenza A that was heterologous to the strain from which the gene for NP was cloned. Vectors used vaccines have also been described by Kieny *et al.* (1992), Hock *et al.* (1993) and Yankaukas *et al.* (1993).

Intramuscular (i.m.) injection of DNA expression vectors in mice has been demonstrated to result in the uptake of DNA by the muscle cells and expression of the protein encoded by the DNA (Ascadi *et al.* 1991; Fazio *et al.*, 1994). Plasmids were shown to be maintained episomally and did not replicate. Subsequently, persistent expression was observed after i.m. injection in skeletal muscle of rats, fish and primates, and in cardiac muscle of rats (Wolff *et al.* 1992).

Muscle creatine kinase (MCK) is expressed at high levels in both skeletal and cardiac muscle of adult animals (Eppenberger *et al.* 1964; Jockers-Wretou *et al.* 1975; Richterich *et al.* 1967; Tanzer *et al.* 1959). Activation of MCK transcription during skeletal myoblast differentiation has been shown (Chamberlain *et al.* 1985; Jaynes *et al.* 1986; Perriard 1979; Perriard *et al.* 1978; Rosenberg *et al.* 1982), and multiple *cis*-acting regulatory sequences have been identified in the 5' flanking sequence and first intron of the MCK gene (Jaynes *et al.* 1988; Sternberg *et al.* 1988). The best-characterized element is a 207-base-pair (bp) muscle-specific enhancer located about 1,100 nucleotides (nt) 5' of the MCK transcription start site. Another enhancer element is located within a 900-nt region in the first intron. The proximal 776-nt 5' MCK sequence also displays muscle cell type specificity in cultured cells, but the absolute level of expression from this element is quite low compared with expression when either enhancer is present (Jaynes *et al.* 1988). A myocyte-specific binding activity, MEF1, that interacts with both enhancers but not the proximal element, has been identified (Buskin *et al.* 1989). Furthermore, the intact MEF1 site is required for the 5' enhancer to function in MCK expression during skeletal myoblast differentiation in culture.

## SUMMARY OF THE INVENTION

The present invention is directed to a DNA construct which is useful for immunization or gene therapy. The construct of the invention comprises muscle specific regulatory elements, such as a promoter or a promoter and one or more enhancer elements, and a DNA sequence under control of the muscle specific regulatory elements. Several DNA sequences may be incorporated into the DNA construct. In one embodiment, the DNA sequence codes for an antigen, antigenic determinant or an epitope of an antigen. In a second embodiment, the DNA sequence is a normal muscle gene which is effected in a muscle disease. In a third embodiment, the DNA sequence is an antisense for blocking an abnormal muscle gene. In a fourth embodiment, the DNA sequence codes for a protein which circulates in the mammalian blood or lymphatic systems. The present invention is useful for ameliorating the effects of diseases of muscle by expression of the normal gene or blocking abnormal gene expression within muscle cells, for the heterologous expression of a transgene which codes for a circulating protein or a protein which modifies a disease state in which muscle is not primarily involved and for vaccine development.

## BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows the DNA sequence of -1354 to +7 of MCK.

Figure 2 shows a map of pMCK.

Figure 3 shows a representation of a Western blot demonstrating production of Herpes simplex virus glycoprotein gD2 in transfected muscle cells.

Figure 4 shows the geometric mean viral titres post-challenge in vaginal washings of mice immunized with the pMCKgDf or control plasmids.

Figure 5 shows the development of severe clinical disease among mice immunized with pMCKgDf or control plasmids.

Figure 6 shows the geometric mean viral titres post-challenge in vaginal washings of mice immunized with varying doses of pMCKgDf.

Figure 7 shows the development of severe clinical disease among mice immunized with varying doses of pMCKgDf.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to a DNA construct which is useful for immunization or gene therapy. The construct of the invention comprises muscle specific regulatory elements, such as a promoter or a promoter and one or more enhancer elements, and a DNA sequence under control of the muscle specific regulatory elements. The DNA sequence is generally a heterologous sequence, i.e., one which is not natively operably linked to the muscle specific regulatory element. Several DNA sequences may be incorporated into the DNA construct. In one embodiment, the DNA sequence codes for an antigen, antigenic determinant or an epitope of an antigen. In a second embodiment, the DNA sequence is a normal muscle gene which is effected in a muscle disease. In a third embodiment, the DNA sequence is an antisense for blocking an abnormal muscle gene. In a fourth embodiment, the DNA sequence codes for a protein which circulates in the mammalian blood or lymphatic systems. The present invention is useful for ameliorating the effects of diseases of muscle by expression of the normal gene or blocking abnormal gene expression within muscle cells, for the heterologous expression of a transgene which codes for a circulating protein or a protein which modifies a disease state in which muscle is not primarily involved and for vaccine development.

The first element of the DNA constructs of the present invention is a muscle specific regulatory element. A muscle specific regulatory element is any regulatory element which affects the transcription or expression of a gene specifically in muscle tissue and not in other body tissues. The muscle specific regulatory element is generally a muscle specific promoter, but it may also include one or more enhancers. Examples of muscle specific regulatory elements include those which are isolated from muscle specific genes, such as the muscle isozyme of creatine kinase (MCK) (Sternberg *et al.*, 1988), myosin light kinase (Merlie 1992a, 1992b), muscle-specific aldolase (Concordet *et al.*, 1993), muscle-specific enolase (Gaillongo *et al.*, 1993), troponin C (Prigozy *et al.*, 1993), myosin (Kitsis *et al.*, 1991; Takeda *et al.*, 1992, von Harsdorf *et al.*, 1993). Many of these promoters are under the control of the MyoD family of transcription factors (Olsen 1990; Hart 1992). These regulatory elements, as well as other muscle specific regulatory elements, may be modified to remove unnecessary sequences as long as they retain the muscle specificity of action.

-7-

The second element of the DNA constructs of the present invention is a DNA sequence which is operably linked to the muscle specific regulatory element. "Operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. For instance, a promoter is operably linked to a coding sequence if the promoter affects its transcription or expression. Any DNA sequence which is useful for immunization or gene therapy can be used as the second element of the DNA constructs of the present invention. The DNA sequence is generally a heterologous sequence, i.e., one which is not natively operably linked to the muscle specific regulatory element. In one embodiment, the DNA sequence codes for an antigen, antigenic determinant or an epitope of an antigen. An "antigen" refers to a molecule containing one or more epitopes that will stimulate a host's immune system to make a secretory, humoral and/or cellular antigen-specific response. The term is also used interchangeably with "immunogen."

The DNA sequence will code for the protein portion of the antigen. The host will appropriately modify the protein to its native state in accordance with the signals provided by the protein produced in accordance with the present invention. Thus, the antigen produced in the host antigen can be a protein or a host modified protein. The antigen can be a fusion peptide of two or more antigens. Particularly, the antigen can include a native protein or protein fragment, or a synthetic protein or protein fragment or peptide. The antigen can include glycoprotein, glycopeptide, lipoprotein, lipopeptide, nucleoprotein, nucleopeptide. It can also include a peptide-peptide conjugate. Examples of antigens include, but are not limited to, those that are capable of eliciting an immune response against viral or bacterial hepatitis, influenza, diphtheria, tetanus, pertussis, measles, mumps, rubella, polio, pneumococcus, herpes, respiratory syncytial virus, hemophilus influenza type b, chlamydia, varicella-zoster virus or rabies.

In a second embodiment, the DNA sequence is a normal muscle gene which is effected in a muscle disease. A kinase gene is known to be effected in myotonic dystrophy. In this embodiment, the normal kinase gene would be operably linked to the muscle specific promoter for introduction into muscle. In a third embodiment, the DNA sequence is an antisense for blocking the expression of an abnormal muscle gene. The antisense may bind to the mRNA produced by the muscle gene to prevent translation or it may bind to a regulatory region of the abnormal gene to prevent transcription of the abnormal gene.

In a fourth embodiment, the DNA sequence codes for a protein which circulates in the mammalian blood or lymphatic systems. Examples of circulating proteins include, but are not limited to, insulin, peptide hormones, hemoglobin, growth factors, liver enzymes, clotting factors and enzymes, complement factors, cytokines, tissue necrosis factor and erythropoietin.

The DNA constructs of the present invention can be constructed by a variety of well known methods, and the order of ligation of the parts can be varied. In one embodiment, the DNA constructs are prepared by separately ligating the muscle specific regulatory element and the DNA sequence into any desired vector for immunization or gene therapy or an intermediate vector used in the construction of the vector used for immunization of gene therapy. In a second embodiment, the muscle specific regulatory element and the DNA sequence are ligated together to provide a cassette which can be inserted into any desired vector for immunization or gene therapy. Vectors which can be utilized for immunization and gene therapy include those vectors described above and those generally known in the art. In addition, the DNA construct can be introduced directly into muscle tissue without the use of a vector by techniques known in the art.

The present invention provides a vaccine (or a vaccine composition) comprising the DNA construct and a pharmaceutically acceptable carrier. The DNA construct may be used by itself or it may be incorporated into a vector which is used to make the vaccine. This vaccine is used to immunize mammals, including humans, against a disease by administering to the mammal (human) an effective immunizing amount of the vaccine. An effective immunizing amount of a vaccine is known or can be readily determined by skilled artisans. The vaccine is administered to a subject by injection into muscle tissue. The vaccine induces a continuing protective level of antibody and/or cellular immune response which is directed against the antigen of the DNA construct.

The present invention further provides a therapeutic for ameliorating the effects of diseases of muscle by expression of the normal gene or by blocking abnormal gene expression within muscle cells, or a therapeutic for the heterologous expression of a transgene which codes for a circulating protein or a protein which modifies a disease state in which muscle is not primarily involved. The treatment of a disease is accomplished by introducing the DNA construct of the present invention into muscle tissue. The DNA construct, alone or in a vector, can be introduced directly by injection into muscle tissue. Alternatively, the DNA construct in a vector can be introduced into muscle tissue by any of the known gene therapy techniques.

discussed above. The DNA construct in a vector can also be introduced into muscle myoblasts *in vitro* by known techniques and then the treated myoblasts can be returned to an *in vivo* environment. The DNA construct of the present invention can be used for permanent gene transfer as known in the art. Alternatively, the DNA construct of the present invention can be used for a reversible gene transfer in which the DNA construct does not become integrated into the host's DNA. In reversible gene transfer, the DNA construct is periodically administered much like a pharmaceutical. In this instance the DNA construct is administered when the transcription and expression of the heterologous DNA sequence decreases below a predetermined level.

There are several advantages to the use of regulatory elements (promoters, enhancers and the like) which are selectively expressed in muscle cells. These regulatory elements in a suitable vector offer safety advantages over comparable vectors employing potent constitutive promoters (such as the cytomegalovirus (CMV) promoter). Promoters which require transcription factors specific to differentiated muscle cells will not be active in other cell types. Therefore, should the nucleic acid construct become incorporated into the genome of a non-muscle cell by homologous or non-homologous recombination, it will not cause the constitutive expression of normally silent or regulated genes. The lack of constitutive expression is a distinct advantage over the use of powerful constitutive promoters which could promote the production of undesirable gene products in an unrepressed manner. While the cloned nucleotide sequence being driven by the muscle specific regulatory elements will be constitutively expressed in muscle cells, recombination leading to insertion of the vector in these cells will be less likely to lead to deleterious events, disrupt normal cell function or homeostasis. This advantage is principally due to the multi-nucleated nature of myotubes. Integration via homologous recombination will only knock out one of multiple copies of the gene from which the muscle specific regulatory elements were derived, and the effects of a nonhomologous recombination could be diluted by alleles present in other unaffected nuclei.

The present invention is described by reference to the following Examples, which are offered by way of illustration and are not intended to limit the invention in any manner. Standard techniques well known in the art or the techniques specifically described below were utilized. For illustration purposes, the examples utilize the regulatory elements of MCK and the glycoprotein D2

gene of Herpes simplex virus type 2. However, any of the regulatory elements mentioned above or other muscle specific regulatory elements could be used in place of the MCK regulatory elements and other DNA sequences can be used in place of the glycoprotein D2 gene.

### EXAMPLE 1

#### Isolation of the MCK Promoter

The mouse MCK gene promoter and enhancer elements isolated in accordance with this example correspond to nucleotides -1354 to +7 of the mouse MCK gene (Sternberg et al., 1988) and is set forth in Figure 1 and SEQ ID NO:1. The MCK gene promoter and enhancer was isolated from mouse genomic DNA by PCR using conventional techniques and manufacturer's recommended procedures. The PWO polymerase was utilized to limit potential errors in the amplification of the MCK gene elements. The following primers were utilized:

forward: 5'-GAAGATCTCAGCTGAGGTGCAAAAGGCTCCTG-3' (SEQ ID NO:2) and

reverse: 5'-CCCAAAGCTTGTGACCCGGGGGCAGCCCCTGTGCC-3' (SEQ ID NO:3).

The nucleotides underlined in the forward primer indicate a Bgl II site, and the nucleotides underlined in the reverse primer indicate a Hind III site.

The plasmid pCDNA3 was obtained from Invitrogen (San Diego, California). This plasmid contains a cytomegalovirus (CMV) promoter with a polylinker downstream of the promoter. The CMV promoter was removed from pCDNA3 by digestions with Hind III and Bgl II. The amplified MCK gene was digested with Hind II and Bgl II and inserted into the digested pCDNA3 to produce pMCK. A map of pMCK is shown in Figure 2.

### EXAMPLE 2

#### Insertion of gD2 Gene of Herpes Simplex Type 2 in pMCK

The plasmid pWW65 (Muggeridge *et al.*, 1990) was obtained from Gary Cohen and Roselyn Rosenberg of the University of Pennsylvania (Philadelphia, PA). This plasmid contains

-11-

the Herpes simplex virus glycoprotein D2 (gD2) gene with Hind III linkers which was inserted into the Hind III site of pRSVnt EPA under control of the Rous sarcoma virus long terminal repeat promoter. The gD2 gene was isolated from pWW65 by digestion with Hind III and inserted into the Hind III site of pMCK downstream of the MCK regulatory elements. Two plasmids were isolated, pMCKgDf, in which the gD gene was inserted in the correct orientation, and pMCKgDr, in which the gD gene was inserted in the reverse orientation.

### EXAMPLE 3

#### In Vitro Production of gD2 in Muscle Cells

The C2C12 muscle cell line was transfected by conventional techniques with either the pMCKgDf or pMCKgDr, the latter acting as the control. In addition, to test for the specificity of the construct, Cos cells were transfected by conventional techniques with either pMCKgDf or pMCKgDr. Cos cells were chosen to be biased against the muscle specificity of the vector, since the vector can be amplified in Cos cells due to the combination of SV40 T antigen expression by the cells and the presence of an SV40 origin of replication in the vector. This amplification would lead to a larger number of vectors per Cos cell than per C2C12 cell and hence a greater chance for gene expression. The transfected cells were grown and the expression of gD was monitored by western blot analysis. The results are shown in Figure 4. This figure shows that C2C12 cells transfected with pMCKgDf produced immunogenic gD, whereas C2C12 cells transfected with pMCKgDr did not produce immunogenic gD. Cos cells transfected with either pMCKgDf or pMCKgDr did not produce detectable amounts of gD.

### EXAMPLE 4

#### In Vivo Production of gD in Muscle Tissue and Generation of an Immune Response

Six to eight week old female BALB/c were used for this example. The mice were divided into three groups and immunized with 100 µg of either pMCKgDf, pMCKgDr or pRSVnt three times at weekly intervals. The pMCKgDr acted as a first control and the pRSVnt acted as a second control. Prior to viral challenge, the mice were pretreated subcutaneously with

2.0 mg of medroxyprogesterone to induce uniform susceptibility to viral challenge (Teepe *et al.*, 1990). An intravaginal live viral challenge with  $5.3 \times 10^4$  to  $1.7 \times 10^5$  PFU/ml HSV-2 MS strain adsorbed to Dacron pledgets was performed two to three weeks after the last vaccination. Vaginal washing viral titres were obtained by plaque assay of vaginal washing specimens. Viral titres were compared using the Mann-Whitney *U* test. Mice were scored daily for development of genital disease. Severe disease was defined as :

- the presence of genital maceration on two consecutive days;
- the development of urinary retention;
- the development of profound cachexia; or
- the development of hindlimb paralysis.

Serology assays for antibody titres and neutralizing antibodies were conducted using conventional techniques. ELISA was conducted by coating ninety-six well plates with 3 µg of baculovirus expressed gD2. Non-specific binding was prevented by incubating ELISA plates with a solution of normal goat serum, Tween and saline. Samples were diluted serially 1:2 starting at 1:1000. Binding of anti[gD2 antibodies was detected by incubation with peroxidase tagged goat-anti-mouse IgG antibodies. ELISA units were obtained by regression analysis of log-transformed optical densities. Serum anti-gD2 antibody titres (by ELISA) and neutralizing activities were compared using the Mann-Whitney *U* test.

Figure 4 shows the geometric mean viral titres post-challenge in vaginal washings from the immunized mice. The viral titres were similar in the three groups of mice two days post-challenge . The viral titres were decreased in the pMCKgDf immunized mice six days post-challenge compared to pMCKgDr or pRSVnt immunized mice (geometric mean titres 126, 288 489 PFU/ml, respectively;  $p < 0.05$ ).

Figure 5 shows the development of severe disease in the immunized mice following challenge. Development of severe disease was markedly delayed among the pMCKgDf immunized mice compared to the pMCKgDr or pRSVnt immunized mice (median 9, 7 , 7 days, respectively;  $p < 0.01$ ).

### EXAMPLE 5

#### In Vivo Production of gD in Muscle Tissue and Generation of an Immune Response

Example 4 was repeated in which 0.1  $\mu\text{g}$  of 1,25-dihydroxy Vitamin D<sub>3</sub> (1,25(OH)<sub>2</sub>D<sub>3</sub>) was used as an adjuvant for the immunizations. The use of 1,25(OH)<sub>2</sub>D<sub>3</sub> as an adjuvant resulted in an enhanced reduction of viral titres as measured in the vaginal washings and further resulted in enhanced serum antibody titres as measured by ELISA and enhanced serum neutralizing antibodies.

### EXAMPLE 6

#### Dose Effects of pMCKgD Immunization

Six to eight week old female BALB/c were divided into five groups and immunized with varying doses (3  $\mu\text{g}$ , 10 $\mu\text{g}$ , 30  $\mu\text{g}$  or 100  $\mu\text{g}$ ) of pMCKgDf or 100  $\mu\text{g}$  pMCKgDr three times at weekly intervals. The immunizations were made using 0.1  $\mu\text{g}$  of (1,25(OH)<sub>2</sub>D<sub>3</sub>) as an adjuvant. The mice were treated and challenged with HSV as described in Example 4. The severity of disease and serology assays were analyzed as described in Example 4.

Figure 6 shows the geometric mean viral titres post-challenge in vaginal washings from the immunized mice. This figure shows dose related effects on the viral titres in vaginal washings. Figure 6 shows that the viral titre for the 100 $\mu\text{g}$  pMCKgDf immunized mice was significantly lower ( $p = 0.05$ ) than the viral titre for the control pMCKgDr immunized mice at two days post challenge. Figure 6 shows that at six days post challenge, the viral titres for the 30  $\mu\text{g}$  and 100  $\mu\text{g}$  pMCKgDf immunized mice were significantly lower than the viral titre for the control pMCKgDr ( $p = 0.05$ ,  $p < 0.01$ , respectively).

Figure 7 shows the development of severe disease in the immunized mice following challenge. This figure shows dose related effects on the development of severe clinical disease. Development of severe disease was markedly delayed among the pMCKgDf immunized mice compared to the pMCKgDr immunized mice ( $p = 0.02$ ).

It will be appreciated that the methods and compositions of the instant invention can be incorporated in the form of a variety of embodiments, only a few of which are disclosed herein. It will be apparent to the artisan that other embodiments exist and do not depart from the spirit of the invention. Thus, the described embodiments are illustrative and should not be construed as restrictive.

## LIST OF REFERENCES

- Anderson, *et al.* (1980). *Proc. Natl. Acad. Sci. USA* **77**:5399-5403.
- Ascadi, G. *et al.* (1991). *Nature* **352**:815.
- Berkner (1992). *Curr. Top. Microbiol. Immunol.* **158**:39-61.
- Berkner, *et al.* (1988). *BioTechniques* **6**:616-629.
- Brandyopadhyay and Temin (1984). *Mol. Cell. Biol.* **4**:749-754.
- Breakfield and Geller (1987). *Mol. Neurobiol.* **1**:337-371.
- Brinster, *et al.* (1981). *Cell* **27**:223-231.
- Buchschacher and Panganiban (1992). *J. Virol.* **66**:2731-2739.
- Buskin, J. *et al.* (1989). *Mol. Cell Biol.* **9**:2627-2640.
- Chamberlain, J.S. *et al.* (1985) *Mol. Cell Biol.* **5**:484-492.
- Collins, D. *et al.* (1992). *J. Immunol.* **148**:3335.
- Concordet, J. *et al.* (1993). *Mol. Cell. Biol.* **13**:9.
- Constantini and Lacy (1981). *Nature* **294**:92-94.
- Cooney, E. *et al.* (1991). *Lancet* **337**:567.
- Cotten, *et al.* (1990). *Proc. Natl. Acad. Sci. USA* **87**:4033-4037.
- Culver, *et al.* (1992). *Science* **256**:1550-1552.
- Curiel, *et al.* (1991a). *Proc. Natl. Acad. Sci. USA* **88**:8850-8854.
- Curiel, *et al.* (1991b). *Hum. Gene Ther.* **3**:147-154.
- Eglitis, M. and Anderson, W.F. (1988). *Biotechniques* **6**:608.
- Eppenberger, H.M. *et al.* (1964). *Dev. Biol.* **10**:1-16.
- Fazio, V.M., *et al.* (1994). *Biochem. Biophys. Res. Comm.* **200**:298.
- Felgner, *et al.* (1987). *Proc. Natl. Acad. Sci. USA* **84**:7413-7417.

- Fink, *et al.* (1992). *Hum. Gene Ther.* **3**:11-19.
- Freese, *et al.* (1990). *Biochem. Pharmacol.* **40**:2189-2199.
- Friedmann, T. (1989). *Science* **244**:1275.
- Gaillongo, A. *et al.* (1993). *Eur. J. Biochem.* **214**:367.
- Gammelin *et al.* (1989). *Virology* **70**:71.
- Germain, R. (1981). *Nature* **353**:605.
- Gordon, *et al.* (1980). *Proc. Natl. Acad. Sci. USA* **77**:7380-7384.
- Gorman *et al.* (1991). *J. Virol.* **65**:3704.
- Gorziglia and Kapikian (1992). *J. Virol.* **66**:4407-4412.
- Graham and van der Eb (1973). *Virology* **52**:456-467.
- Hahn, C. *et al.* (1992). *Proc. Nat. Acad. Sci. USA* **89**:2679.
- Hart, S. (1992). *BioScience* **42**:582.
- Helseth, *et al.* (1990). *J. Virol.* **64**:2416-2420.
- Hock, L.J. *et al.* (1993). U.S. Patent No. 5,273,876.
- Jaynes, J.B. *et al.* (1986). *Mol. Cell Biol.* **6**:2855-2864.
- Jaynes, J.B. *et al.* (1988). *Mol. Cell Biol.* **8**:62-70.
- Jockers-Wretou, E. *et al.* (1975). *Clin. Chem. Acta.* **50**:233-232.
- Johnson, *et al.* (1992). *J. Virol.* **66**:2952-2965.
- Kaneda, *et al.* (1989). *J. Biol. Chem.* **264**:12126-12129.
- Kieny, M-P. *et al.* (1992). U.S. Patent No. 5,169,763.
- Kitsis, R. *et al.* (1991). *Proc. Nat. Acad. Sci. USA* **88**:4138.
- Klein, T.M., *et al.* (1987). *Nature* **327**:70.
- Ledley, F.D. (1987). *J. Pediatrics* **110**:1.

- Lim, *et al.* (1992). *Circulation* **83**:2007-2011.
- Madzak, *et al.* (1992). *J. Gen. Virol.* **73**:1533-1536.
- Mann and Baltimore (1985). *J. Virol.* **54**:401-407.
- Margolskee (1992). *Curr. Top. Microbiol. Immunol.* **158**:67-90.
- Mascola, L. *et al.* (1989). *Arch. Intern. Med.* **149**:1569.
- Merlie, J. (1992a). *Cell* **69**:67.
- Merlie, J. (1992b). *Cell* **69**:79.
- Miller, A. (1992). *Curr. Top. Microbiol. Immunol.* **158**:1.
- Miller, *et al.* (1985). *Mol. Cell. Biol.* **5**:431-437.
- Miller, *et al.* (1988). *J. Virol.* **62**:4337-4345.
- Moss (1992). *Curr. Top. Microbiol. Immunol.* **158**:25-38.
- Muzyczka (1992). *Curr. Top. Microbiol. Immunol.* **158**:97-123.
- Nabel, *et al.* (1990). *Science* **249**:1285-1288.
- Nabel (1992). *Hum. Gene Ther.* **3**:399-410.
- Nabel, G.J., *et al.* (1993). *Proc. Natl. Acad. Sci. USA* **87**:11307.
- Ohi, *et al.* (1990). *Gene* **89**:279-282.
- Olsen, E. (1990). *Genes and Development* **4**:1454.
- Page, *et al.* (1990). *J. Virol.* **64**:5370-5276.
- Pellicer, *et al.* (1980). *Science* **209**:1414-1422.
- Perriard, J.C. (1979). *J. Biol. Chem.* **254**:7036-7041.
- Perriard, J.C. *et al.* (1978). *Arch. Biochem. Biophys.* **191**:90-100.
- Petropoulos, *et al.* (1992). *J. Virol.* **66**:3391-3397.
- Prigozy, T. *et al.* (1993). *Somatic Cell Mol. Genet.* **19**:111.

Quantin, *et al.* (1992). *Proc. Natl. Acad. Sci. USA* **89**:2581-2584.

Richterich, R. *et al.* (1967). Comparative studies on creatine kinase and its isoenzymes, pp. 243-253. In: N van Thoai *et al.* (eds.), *Homologous Enzymes and Biochemical Evolution*, Gordon and Breach Publishers, New York, N.Y.

Rosenberg, U.B. *et al.* (1982). *Proc. Nat. Acad. Sci. USA* **79**:6589-6592.

Rosenfeld, *et al.* (1992). *Cell* **68**:143-155.

Shimada, *et al.* (1991). *J. Clin. Invest.* **88**:1043-1047.

Sorge, *et al.* (1984). *Mol. Cell. Biol.* **4**:1730-1737.

Srivastava, A. (1993). U.S. Patent No. 5,242,479.

Sternberg, E.A., *et al.* (1988). *Mol. Cell. Biol.* **8**:2896.

Stewart, *et al.* (1992). *Hum. Gene Ther.* **3**:267-275.

Stratford-Perricaudet, *et al.* (1990). *Hum. Gene Ther.* **1**:241-256.

Takeda, S. *et al.* (1992). *J. Biol. Chem.* **267**:16957.

Tanzer, M.L. *et al.* (1959). *J. Biol. Chem.* **234**:3201-3204.

Taylor, P. *et al.* (1986). *Immunology* **58**:417.

Teepe, A.G. *et al.* (1990). *Antiviral Res.* **14**:227.

Townsend, A. *et al.* (1989). *Ann. Rev. Immunol.* **7**:601.

von Harsdorf, R. *et al.* (1993). *Circulat. Res.* **72**:689.

Wagner, *et al.* (1990). *Proc. Natl. Acad. Sci. USA* **87**:3410-3414.

Wagner, *et al.* (1991). *Proc. Natl. Acad. Sci. USA* **88**:4255-4259.

Wang and Huang (1989). *Biochemistry* **28**:9508-9514.

Wilkinson, *et al.* (1992). *Nucleic Acids Res.* **20**:2233-2239.

Wolff, J.A., *et al.* (1990). *Science* **247**:1465 (1990).

Wolff, J.A., *et al.* (1991). *BioTechniques* **11**:474-485.

- Wolff, J. *et al.* (1992). *Hum. Mol. Genet.* 1:363.
- Wooster, R., *et al.* (1994). *Science* **265**:2088.
- Wu, *et al.* (1989a). *Genomics* **4**:560-569.
- Wu, *et al.* (1989b). *J. Biol. Chem.* **264**:16985-16987.
- Wu, *et al.* (1991). *J. Biol. Chem.* **266**:14338-14342.
- Yang, N.S., *et al.* (1990). *Proc. Natl. Acad. Sci. USA* **87**:9568.
- Yankauckas, M.A., *et al.* (1993). *DNA Cell Biol.* **12**:771.
- Yewdell, J. *et al.* (1989). *Science* **244**:1072.
- Zenke, *et al.* (1990). *Proc. Natl. Acad. Sci. USA* **87**:3655-3659.

-20-

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Ricigliano, Joseph W.  
Araneo, Barbara A.
- (ii) TITLE OF INVENTION: DNA CONSTRUCT FOR IMMUNIZATION OR GENE  
THERAPY
- (iii) NUMBER OF SEQUENCES: 3
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Venable, Baetjer, Howard & Civiletti, LLP
  - (B) STREET: 1201 New York Avenue, N.W., Suite 1000
  - (C) CITY: Washington
  - (D) STATE: D.C.
  - (E) COUNTRY: U.S.A.
  - (F) ZIP: 20005
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER: US
  - (B) FILING DATE: 19-SEP-1995
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Ihnen, Jeffrey L.
  - (B) REGISTRATION NUMBER: 28,957
  - (C) REFERENCE/DOCKET NUMBER: 23691-115854
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: 202-962-4810
  - (B) TELEFAX: 202-962-8300

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1361 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO

WO 97/11190

PCT/US96/14829

-21-

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mouse

(ix) FEATURE:

(A) NAME/KEY: misc\_feature

(B) LOCATION: 1..1354

(D) OTHER INFORMATION: /note= "5' nontranscribed  
sequence."

(ix) FEATURE:

(A) NAME/KEY: 5'UTR

(B) LOCATION: 1355..1361

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

|                                                                    |      |
|--------------------------------------------------------------------|------|
| CAGCTGAGGT GCAAAAGGCT CCTGTCATAT TGTGTCCTGC TCTGGTCTGC CTTACAGCT   | 60   |
| TGGGGGCCAC CTAGCCCACC TCTCCCTAGG GATGAGAGCA GCCACTATGG GTCTAGGCTG  | 120  |
| CCCATGTAAG GAGGCAAGGC CTGGGGACAC CCGAGATGCC TGGTTATAAT TAACCCAGAC  | 180  |
| ATGTGGCTGC TCCCCCCCCC CAACACCTGC TGCCTGAGCC TCACCCCCAC CCCGGTGCCT  | 240  |
| GGGTCTTAGG CTCTGTACAC CATGGAGGAG AAGCTCGCTC TAAAATAAC CCTGTCCCTG   | 300  |
| GTGGATCCNN TCCGGAGGGG CAGGCTGAGG GCGGCCACTT CCCTCAGCCG CAGTTTGT    | 360  |
| TCCCAAGAAT GGTTTTTCTG CTTCTGTAGC TTTTCCTGTC AATTCTGCCA TGGTGGAGCA  | 420  |
| GCCTGCACTG GGCTTCTGGG AGAAACCAA CCGGGTTCTA ACCTTTCAGC TACAGTCATT   | 480  |
| GCCTTTCCTG TAGATGGGCG ACTACAGCCC CACCCCACC CCCGTCTCCT GTATCCTTCC   | 540  |
| TGGGCCTGGG GATCCTAGGC TTTCACCTGGA AATTTCCCCC CAGGTGCTGT AGGCTAGAGT | 600  |
| CACGGCTCCC AAGAACAGTG CTTGCCTGGC ATGCATGGTT CTGAACCTCC AACTGCAAAA  | 660  |
| AATGACACAT ACCTTGACCC TTGGAAGGCT GAGGCAGGGG GATTGCCATG AGTGCAAAGC  | 720  |
| CAGACTGGGT GGCATAGTTA GACCCTGTCT CAAAAACCA AAAACAATTA AATAACTAAA   | 780  |
| GTCAGGCAAG TAATCCTACT CAGGAGACTG AGGCAGAGGG ATTGTTACAT GTCTGAGGCC  | 840  |
| AGCCTGGACT ACATAGGGTT TCAGGCTAGC CCTGTCTACA GAGTAAGGCC CTATTTCAAA  | 900  |
| AACACAAACA AAATGGTTCT CCCAGCTGCT AATGCTCACC AGGCAATGAA GCCTGGTGAG  | 960  |
| CATTAGCAAT GAAGGCAATG AAGGAGGGTG CTGGCTACAT CAGGCTGTGG GGGACTGAGG  | 1020 |
| GCAGGCTGTA ACAGGCTTGG GGGCCAGGGC TTATACGTGC CTGGGACTCC CAAAGTATTA  | 1080 |
| CTGTTCCATG TTCCCGGCGA AGGGCCAGCT GTCCCCCGCC AGCTAGACTC AGCACTTAGT  | 1140 |
| TTAGGAACCA GTGAGCAAGT CAGCCCTTGG GGCAGCCCAT ACAAGGCCAT GGGGCTGGGC  | 1200 |

WO 97/11190

PCT/US96/14829

-22-

AAGCTGCACG CCTGGGTCCG GGGTGGGCAC GGTGCCCCGG CAACGAGCTG AAAGCTCATC 1260  
TGCTCTCAGG GGCCCCCTCCC TGGGGACAGC CCCTCCTGGC TAGTCACACC CTGTAGGCTC 1320  
CTCTATATAA CCCAGGGGCA CAGGGGCTGC CCCCGGGTCA C 1361

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "Synthetic primer"

## (iii) HYPOTHETICAL: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GAAGATCTCA GCTGAGGTGC AAAAGGCTCC TG 32

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "Synthetic primer"

## (iii) HYPOTHETICAL: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CCCAAGCTTG TGACCCGGGG GCAGCCCCTG TGCC 34

CLAIMS

1. A DNA construct for the expression of a heterologous DNA sequence in muscle tissue consisting of a muscle specific regulatory element of muscle isozyme of creatine kinase (MCK) and a heterologous DNA sequence operably linked to said element, wherein said regulatory element of MCK consists of the nucleic acid set forth in SEQ ID NO:1 and wherein said heterologous DNA sequence is expressed when said DNA construct is introduced into muscle tissue.
2. The DNA construct of claim 1, wherein said heterologous DNA sequence codes for an antigen.
3. The DNA construct of claim 2, wherein said antigen is selected from the group consisting of viral or bacterial hepatitis, influenza, diphtheria, tetanus, pertussis, measles, mumps, rubella, polio, pneumococcus, herpes, respiratory syncytial virus, hemophilus influenza type b, chlamydia, varicella-zoster virus or rabies.
4. The DNA construct of claim 1, wherein said DNA sequence codes for an antisense RNA.
5. The DNA construct of claim 1, wherein said DNA sequence codes for circulating protein.
6. The DNA construct of claim 5, wherein said circulating protein is selected from the group consisting of insulin, peptide hormones, hemoglobin, growth factors, liver enzymes, clotting factors and enzymes, complement factors, cytokines, tissue necrosis factor and erythropoietin.
7. The DNA construct of claim 1, wherein said DNA sequence codes for a biologically functional muscle protein.

8. A vector which comprises the DNA construct of any claims 1 and 4 to 7.
9. The vector of claim 8 further comprising a nucleotide sequence capable of integration into a host chromosome.
10. The vector of claim 8 further comprising a nucleotide sequence capable of replication in a host.
11. A vector which comprises the DNA construct of claim 2 or 3.
12. The vector of claim 11 further comprising a nucleotide sequence capable of integration into a host chromosome.
13. The vector of claim 11 further comprising a nucleotide sequence capable of replication in a host.
14. A vaccine comprising the DNA construct of claim 2 and a physiologically acceptable carrier.
15. A vaccine comprising the DNA construct of claim 3 and a physiologically acceptable carrier.
16. A vaccine comprising the vector of claim 11 and a physiologically acceptable carrier.
17. A vaccine comprising the vector of claim 13 and a physiologically acceptable carrier.

-1354

CAGCTGAGGT GCAAAAGGCT CCTGTCATAT TGTGTCCTGC TCTGGTCTGC CTTACAGCT  
TGGGGGCCAC CTAGCCCACC TCTCCCTAGG GATGAGAGCA GCCACTATGG GTCTAGGCTG  
CCCATGTAAG GAGGCAAGGC CTGGGGACAC CCGAGATGCC TGGTTATAAT TAACCCAGAC  
ATGTGGCTGC TCCCCCCCCC CAACACCTGC TGCCTGAGCC TCACCCCCAC CCCGGTGCCT  
GGGTCTTAGG CTCTGTACAC CATGGAGGAG AAGCTCGCTC TAAAAATAAC CCTGTCCCTG  
GTGGATCCNN TCCGGAGGGG CAGGCTGAGG GCGGCCACTT CCCTCAGCCG CAGTTTGT  
TCCCAAGAAT GGTTTTTCTG CTTCTGTAGC TTTTCCTGTC AATTCTGCCA TGGTGGAGCA  
GCCTGCACTG GGCTTCTGGG AGAAACCAAA CCGGGTTCTA ACCTTTCAGC TACAGTCATT  
GCCTTTCCTG TAGATGGGCG ACTACAGCCC CACCCCCACC CCCGTCTCCT GTATCCTTCC  
TGGGCCTGGG GATCCTAGGC TTTCACTGGA AATTTCCCCC CAGGTGCTGT AGGCTAGAGT  
CACGGCTCCC AAGAACAGTG CTTGCCTGGC ATGCATGGTT CTGAACCTCC AACTGCAAAA  
AATGACACAT ACCTTGACCC TTGGAAGGCT GAGGCAGGGG GATTGCCATG AGTGCAAAGC  
CAGACTGGGT GGCATAGTTA GACCCTGTCT CAAAAACCA AAAACAATTA AATAACTAAA  
GTCAGGCAAG TAATCCTACT CAGGAGACTG AGGCAGAGGG ATTGTTACAT GTCTGAGGCC  
AGCCTGGACT ACATAGGGTT TCAGGCTAGC CCTGTCTACA GAGTAAGGCC CTATTTCAAA  
AACACAAACA AAATGGTTCT CCCAGCTGCT AATGCTCACC AGGCAATGAA GCCTGGTGAG  
CATTAGCAAT GAAGGCAATG AAGGAGGGTG CTGGCTACAT CAGGCTGTGG GGGACTGAGG  
GCAGGCTGTA ACAGGCTTGG GGGCCAGGGC TTATACGTGC CTGGGACTCC CAAAGTATTA  
CTGTTCCATG TTCCCGGCGA AGGGCCAGCT GTCCCCGCC AGCTAGACTC AGCACTTAGT  
TTAGGAACCA GTGAGCAAGT CAGCCCTTGG GGCAGCCCAT ACAAGGCCAT GGGGCTGGGC  
AAGCTGCACG CCTGGGTCCG GGGTGGGCAC GGTGCCCCGG CAACGAGCTG AAAGCTCATC  
TGCTCTCAGG GGCCCCCCC TGGGGACAGC CCCTCCTGGC TAGTCACACC CTGTAGGCTC  
CTCTATATAA CCCAGGGGCA CAGGGGCTGC CCCC GGGTCA C<sup>+7</sup>

FIG. 1

SUBSTITUTE SHEET (RULE 26)




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6

7/7




FIG. 7

