



(86) Date de dépôt PCT/PCT Filing Date: 2000/02/15

(87) Date publication PCT/PCT Publication Date: 2000/08/24

(45) Date de délivrance/Issue Date: 2009/11/17

(85) Entrée phase nationale/National Entry: 2001/08/15

(86) N° demande PCT/PCT Application No.: US 2000/003697

(87) N° publication PCT/PCT Publication No.: 2000/049036

(30) Priorité/Priority: 1999/02/18 (US09/252,072)

(51) Cl.Int./Int.Cl. *C07K 5/072* (2006.01),
A23L 1/22 (2006.01), *A23L 1/236* (2006.01),
A23L 2/60 (2006.01), *C07K 1/04* (2006.01),
C07K 17/00 (2006.01), *C07K 17/02* (2006.01),
C07K 17/10 (2006.01), *C07K 5/06* (2006.01)

(72) Inventeurs/Inventors:

FOTOS, JIM, US;
BISHAY, IHAB, US

(73) Propriétaire/Owner:

NUTRASWEET PROPERTY HOLDINGS, INC., US

(74) Agent: MCFADDEN, FINCHAM

(54) Titre : AGGLOMERAT DE N-[N-(3,3-DIMETHYLBUTYL)-L- α -ASPARTYL]-L-PHENYLALANINE 1-METHYL ESTER

(54) Title: N-[N-(3,3-DIMETHYLBUTYL)-L- α -ASPARTYL]-L-PHENYLALANINE 1-METHYL ESTER AGGLOMERATE

(57) **Abrégé/Abstract:**

This invention provides a process for preparing an agglomerate of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and a carrier comprising the steps of: (a) providing a premix solution comprising N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and a binding agent; (b) fluidizing a carrier; and (c) applying the premix solution of step (a) onto said fluidized carrier to form an agglomerate of said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said carrier. This invention is also directed to the novel agglomerates prepared by the process of this invention.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 5/06, A23L 1/236	A1	(11) International Publication Number: WO 00/49036 (43) International Publication Date: 24 August 2000 (24.08.00)
(21) International Application Number: PCT/US00/03697 (22) International Filing Date: 15 February 2000 (15.02.00) (30) Priority Data: 09/252,072 18 February 1999 (18.02.99) US (71) Applicant: THE NUTRASWEET COMPANY [US/US]; 200 World Trade Center, Suite 900, Merchandise Mart, Chicago, IL 60654-1001 (US). (72) Inventors: FOTOS, Jim; 1370 Longacre Lane, Wheeling, IL 60090 (US). BISHAY, Ihab; 187 Springbrook Court, Mundelein, IL 60060 (US). (74) Agents: MANDRA, Raymond, R. et al.; Fitzpatrick, Cella, Harper & Scinto, 30 Rockefeller Plaza, New York, NY 10112-3801 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: N-[N-(3,3-DIMETHYLBUTYL)-L- α -ASPARTYL]-L-PHENYLALANINE 1-METHYL ESTER AGGLOMERATE (57) Abstract This invention provides a process for preparing an agglomerate of N-[N-(3,3-dimethylbutyl)-L-α-aspartyl]-L-phenylalanine 1-methyl ester and a carrier comprising the steps of: (a) providing a premix solution comprising N-[N-(3,3-dimethylbutyl)-L-α-aspartyl]-L-phenylalanine 1-methyl ester and a binding agent; (b) fluidizing a carrier; and (c) applying the premix solution of step (a) onto said fluidized carrier to form an agglomerate of said N-[N-(3,3-dimethylbutyl)-L-α-aspartyl]-L-phenylalanine 1-methyl ester and said carrier. This invention is also directed to the novel agglomerates prepared by the process of this invention.		

TITLE

5 N-[N-(3,3-DIMETHYLBUTYL)-L- α -ASPARTYL]-L-
 PHENYLALANINE 1-METHYL ESTER AGGLOMERATE

BACKGROUND OF THE INVENTION

10 Field of the Invention


 The present invention relates to a process for
 preparing agglomerates of N-[N-(3,
 3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine
15 1-methyl ester (also known as neotame) and a
 carrier. The invention also relates to the novel
 agglomerates.

- 2 -


Discussion of the Related Art

N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is a derivative of aspartame that has a sweetening potency that is about 40 to 50 times that of aspartame (and about 8,000 times that of sucrose). N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester may be prepared from aspartame as described in U.S. Patent No. 5,480,668, U.S. Patent No. 5,510,508, and U.S. Patent No. 5,728,862.

Structurally, N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and aspartame differ in that, in N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester, a bulky neohexyl substituent is present on the amine nitrogen.



Aspartame



Neotame

This structural difference results in dramatic differences in the physical and chemical properties of these compounds. For example, the melting point of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is about 82.2°C, while that of aspartame is 248°C. In addition, N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester has much higher solubility in organic solvents than aspartame. It is also known that N-

- 3 -

[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester has a higher stability than aspartame under some pH conditions, as described in U.S. Patent No. 5,480,688. The pronounced difference in sweetness between the two compounds is further evidence of their chemical dissimilarity.

The solubility of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in water is about 1%, which is approximately equal to the water solubility of other high intensity sweeteners such as aspartame. However, while the solubility of other high intensity sweeteners in ethanol is very low (e.g., less than 0.1% for aspartame), the solubility of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in ethanol is greater than 100g/100ml.

Thus, unlike less soluble sweeteners such as aspartame which are agglomerated as a powder onto a powder carrier, N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester may be agglomerated as a solution onto a powder carrier. The process of agglomeration using a N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester solution, rather than a powder, provides numerous significant advantages including improved content uniformity and ease of processing.

N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester may be used for sweetening a variety of products, including drinks, foods, confectionery, pastries, chewing gums, hygiene products and toiletries, as well as cosmetic, pharmaceutical and veterinary products. Its superior sweetening potency makes N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-

- 4 -

5 methyl ester an attractive alternative to aspartame because it permits the use of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in substantially smaller quantities than is required for aspartame to achieve an equivalent sweetening effect.

10 Because of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester's superior potency, it is convenient to prepare compositions of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester which deliver sweetness on par with that of aspartame. For liquid products such as beverages, this may be accomplished by preparing
15 dilute solutions of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester. Since N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester's sweetening potency is about 40 times greater than that of aspartame, these
20 solutions can be prepared using about 2.5% as much N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester as would be required to prepare solutions of equal sweetness using aspartame. This amount can be adjusted to take into
25 account the relative potency differences relating to the concentration effect of the respective sweeteners.

30 For solid products, such as tabletop sugar substitutes, this may be accomplished by forming a dry blend of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester with a bulking agent such as, for example, Unidex brand mixture of dextrose (97%) and maltodextrin (3%) available from
35 CPC International. It is common to blend 95% by weight Unidex with 5% by weight aspartame to provide a tabletop sugar substitute. An equivalent N-[N-

- 5 -

(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester sugar substitute would require blending about 99.875% by weight Unidex with about 0.125% by weight N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester.

The use of small amounts of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester powder to prepare dry blends with a bulking agent presents several manufacturing difficulties. Not surprisingly, dry blends prepared using a relatively small amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and a relatively large amount of a bulking agent may exhibit poor content uniformity. Because of the relatively small amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester as compared to bulking agent, these dry blends also may exhibit a high degree of segregation and dusting. Moreover, the content uniformity and segregation problems associated with such dry blends often worsen during shipping and also during periods of storage. Consequently, these dry blends may be somewhat unattractive for manufacturers of consumer products as well as for consumers.

It would be highly desirable to overcome the difficulties described above by preparing agglomerates of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester with a carrier and using the agglomerates to provide N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in a deliverable form. In addition, it would be extremely advantageous to provide a tabletop sweetener with agglomerates of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester. Similarly, it would be extremely

- 6 -

advantageous to provide a powdered soft drink mix with agglomerates of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester. It is also important to deliver a blend of sweetness in an efficient manner.

Fluidized bed agglomeration is well known in the art. The process is described in U.S. Patent Nos. 2,856,290, 3,251,695, and 3,433,644. Typically, in both continuous and batch fluid bed agglomeration processes, finely divided particles are sprayed onto a fluidized bed of particles under moisture and temperature conditions which promote formation of an agglomerate. Often the process involves heating at least one of the components of the agglomerate to a temperature above its melting point.

U.S. Patent No. 4,554,167 discloses a method for preparing agglomerates of aspartame and acid-containing food mixes. The disclosed agglomeration process does not involve heating one or more of the agglomerate components to a temperature above its melting point. Neither does the disclosed process involve dissolving the high intensity sweetener in a solvent prior to formation of the agglomerate.

Agglomerates of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester would facilitate adjustment of the amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester to be delivered. These agglomerates also would provide N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester preparations with excellent content uniformity. In addition, these agglomerates also would exhibit

- 7 -

reduced segregation and dusting as compared to the N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester dry blend preparations.

5

SUMMARY OF THE INVENTION

This invention provides a process for preparing an agglomerate of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and a carrier comprising the steps of: (a) providing a premix solution comprising N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and a binding agent; (b) fluidizing a carrier; and (c) applying the premix solution of step (a) onto said fluidized carrier to form an agglomerate of said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said carrier. This invention is also directed to the novel agglomerates prepared by the process of this invention. This invention is also directed to tabletop sweeteners, powdered soft drink mixes and sweetener blends having the above-described agglomerates.

25

DETAILED DESCRIPTION OF THE INVENTION

In the agglomeration process of the present invention, a premix solution is formed which comprises N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and a binding agent. The premix solution may be formed by adding the N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and the binding agent to the same solvent. Alternatively, the premix solution may be formed by adding the N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and the binding agent to separate solvents and then combining the two solutions. In the latter

- 8 -

approach, the solvent used to dissolve the N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester may be the same as, or different from, the solvent used to dissolve the binding agent.

5 Under either process, the resultant premix is applied onto a fluidized carrier using a fluid bed agglomeration mixer. Preferably, the premix is applied onto the fluidized carrier by spraying the premix onto the fluidized carrier. An agglomerate
10 of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and the carrier is thus formed.

The agglomerates of this invention comprise: (i) an
15 effective sweetener amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester; (ii) a binding agent; and (iii) a carrier. The binders and carriers employed in the agglomerates of this invention are described in
20 greater detail herein.

The agglomeration process of the present invention may be performed using a batch fluid bed agglomerator. Other equipment that may be used in
25 the present invention include a continuous fluid bed agglomerator or a continuous turbulent flow agglomerator (e.g., Schugi Flex-O-Mix and Turbulizer, Hosokawa Bepex Corp., Minneapolis, MN).

30 In order to prepare the premix solution, N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is added to a solvent. Any solvent in which N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester dissolves may be used
35 in the present invention. Preferably, the solvent is a food grade solvent. Exemplary solvents which may be used in this invention include ethanol, water,

- 9 -

isopropanol, methanol, and mixtures thereof.
Preferably, the solvent is ethanol.

5 The binding agent is dissolved in a solvent which
may be the same as, or different from, the solvent
used to dissolve the N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl ester. Any
solvent in which the particular binding agent chosen
dissolves may be used in the present invention.
10 Solvents which may be used to dissolve the binding
agent include water, ethanol, isopropanol, methanol
and mixtures thereof. Preferably, the solvent is a
food grade solvent. Most preferably, the solvent is
water.

15 As mentioned above, it is not necessary to dissolve
the N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-
phenylalanine 1-methyl ester in one solvent, the
binding agent in a separate solvent, and then
20 combine the two solutions to form a premix. As is
readily apparent to those skilled in the art, the
premix solution may be prepared by dissolving the N-
[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine
1-methyl ester and the binding agent in the same
25 solvent. The solvent may be a single compound
(e.g., ethanol), or it may be a mixture of compounds
(e.g., ethanol and water). This invention is not
limited in the manner of making the premix of the
invention.

30 In order to effect complete mixing of the premix,
the premix may be heated up to about 90°C. As used
herein, the term "effect mixing" means blending
sufficiently so as to form a mixture. Preferably,
35 the premix solution is heated to between about 30°C
and about 50°C, more preferably between about 35°C
and about 45°C.

- 10 -

The carrier is fluidized and its temperature is adjusted to between about 20°C and about 50°C. Preferably, the carrier is heated to between about 35°C and about 45°C. More preferably, the carrier is heated to about 40°C.

The carrier may be placed into a removable bowl of a fluid bed agglomerator. After the bowl is secured to the fluid bed agglomerator, the carrier is fluidized and heated as necessary by adjusting the inlet air temperature. Preferably, the temperature of the inlet air is maintained between 50°C and 100°C. For example, to heat the fluidized carrier to about 40°C, the inlet air temperature may be adjusted to between 70°C - 75°C.

Once the fluidized carrier reaches the desired temperature, the premix solution may be applied through the spray nozzle of the fluid bed agglomerator. The premix solution may be sprayed onto the fluidized carrier at any rate which is effective to produce an agglomerate having the desired particle size distribution. Those skilled in the art will recognize that a number of parameters may be adjusted to obtain the desired particle size distribution. After spraying is completed, the agglomerate may be allowed to dry. Preferably, the agglomerate is allowed to dry until the outlet air temperature reaches 35°C to 40°C.

As stated above, the solubility of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in ethanol is greater than 100g/100ml. In contrast, the solubility of other high intensity sweeteners in ethanol is very low (e.g., less than 0.1% for aspartame).

- 11 -

As a result of N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl ester's unusually
high solubility in ethanol and other alcohols,
solutions of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-
5 L-phenylalanine 1-methyl ester in ethanol may be
prepared over a broad range of concentrations. Such
solutions may be used in a fluidized bed
agglomeration process to make N-[N-(3,3-
dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-
10 methyl ester agglomerates with various carriers.
The resultant N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl ester
agglomerates provide a delivery vehicle for N-[N-
(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-
15 methyl ester in a form which is convenient for use
in many food applications.

As a result of the very small amount of N-[N-(3,3-
dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-
20 methyl ester used, the amount of ethanol required
for solubilization is insignificant. In contrast,
even if other high intensity sweeteners could be
solubilized in ethanol, the amount of ethanol
required could be cause for concern.

25 An advantageous feature of the present invention is
that the level of N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl ester that is
present in these agglomerates may be controlled by
30 varying the amount of N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl ester that is
added to the premix solution. This is particularly
important when trying to match the sweetness
delivered by other sweeteners such as aspartame in
35 different products at different concentration
levels. The use of a N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl ester/binding

- 12 -

agent premix also provides advantages in the manufacturing process by facilitating the preparation of especially desirable tabletop products.

5

Another advantageous aspect of the present invention is that the N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerates may be blended with blending agents. As used herein, the term blending agents includes a broad range of ingredients commonly used in foods or beverages including, but not limited to, those used herein as binding agents, carriers, bulking agents, and sweeteners. Exemplary blending agents that may be blended with the N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerates of this invention include aspartame, acesulfame and its salts, sucralose, saccharin, alitame, cyclamates, stevia derivatives, thaumatin, sucrose, fructose, dextrose, polyol sugar alcohols, citric acid, dextrin, maltodextrin, dextrose-maltodextrin blends, lactose, inulin, erythritol, sorbitol, stevioside, hydroxypropylmethyl cellulose, carboxymethyl cellulose, polyvinylpyrrolidone, and mixtures thereof. A preferred blending agent that may be dry blended with the N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerates of this invention is Unidex.

30

Such blends are particularly convenient to manufacture, as they take advantage of procedures used in conventional tabletop manufacturing processes. To illustrate, in conventional manufacturing of tabletop sweeteners with aspartame, the aspartame alone is blended with Unidex. The agglomerate of this invention can be tailored to

35

- 13 -

match the level of sweetness and bulk of aspartame, thereby allowing the agglomerate to be a 1:1 substitute for aspartame. Thus, the agglomerates of this invention provide great flexibility in the use of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in consumer products which use high intensity sweeteners.

The N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerates of this invention may be dry blended with blending agents to prepare highly desirable consumer products. For example, the agglomerates of this invention may be used to prepare particularly desirable tabletop sweeteners and powdered soft drink mixes. The tabletop sweeteners and powdered soft drink mixes of this invention may be prepared by dry blending the agglomerates of this invention with blending agents commonly used to prepare tabletop sweeteners and powdered soft drink mixes using methods well known to those skilled in the art.

The N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerates of this invention may be dry blended with sweeteners to provide highly desirable sweetener blends. Exemplary sweeteners that may be dry blended with the N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerates of this invention include aspartame, acesulfame and its salts, sucralose, saccharin, alitame, cyclamates, stevia derivatives, thaumatin, sucrose, fructose, dextrose, polyol sugar alcohols, and mixtures thereof. It is also possible to prepare agglomerates containing N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and one of the above described other sweeteners.

- 14 -

The amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate that may be present in the sweetener blends of this invention may vary over a broad range. The
5 agglomerates of this invention are extremely compatible for blending with other sweeteners in a wide range of usage levels. This compatibility will allow one skilled in the art to readily formulate the blends depending upon the desired end use
10 functionality.

Another advantageous feature of the present invention is that the concentration of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine
15 1-methyl ester in the solvent may vary widely depending upon the particular solvent used, as well as upon the desired level of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in the agglomerate. Thus, where it
20 is desired to have a high level of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in the agglomerate, a greater amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester would be dissolved in
25 the solvent. Conversely, where it is desired to have a low level of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in the agglomerate, a lesser amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine
30 1-methyl ester may be dissolved in the solvent.

The available range of concentration of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in the solvent will, of course,
35 depend upon the solubility of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in the particular solvent chosen.

- 15 -

A particular advantage of the present invention is that because of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester's solubility properties in ethanol, the broad range of concentrations that can be achieved may be used to produce a broad range of concentrations of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in the agglomerate.

A large number of solvents may be used in the process of this invention, depending upon the particular processing conditions employed. When ethanol is used as the solvent, the available range of concentrations of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is very broad. For example, the concentration of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in ethanol solutions may be between about 0.01% and about 50% by weight. Preferably, the concentration of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in ethanol solutions will be between about 1% and about 15%, more preferably between about 5% and about 10% by weight.

Binding agents facilitate the agglomeration of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester to the carrier. Any binding agent with sufficient binding strength may be used in the present invention. Exemplary binding agents which may be used in the present invention include maltodextrin, sucrose, gellan gum, hydroxypropylmethyl cellulose (HPMC), carboxymethyl cellulose (CMC), polyvinyl pyrrolidone (PVP) and mixtures thereof. Preferably, the binding agent is maltodextrin.

- 16 -

The concentration of binding agent in the solution may vary depending on a variety of factors including the binding strength of the particular binding agent and the particular solvent chosen. When water is the solvent, it is preferable that the concentration of binding agent in the aqueous solution be between about 1% and about 50% by weight. More preferably, the concentration of binding agent in the aqueous solution is between about 5% and about 25% by weight.

The weight ratio of binding agent to N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in the premix solution may vary from as low as about 1:10 to as high as about 10:1. Preferably, the weight ratio of binding agent to N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is from about 0.5:1.0 to about 2:1.

An advantageous aspect of this invention is that the premixes of this invention provide excellent formulation flexibility. These premixes enable the agglomerates of this invention to be used as delivery vehicles for other ingredients, such as, for example, flavors. Consequently, the agglomerates of this invention may be used as flavor modifiers. The use of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester as a flavor modifier is discussed in U.S. Patent Publication No. 2003/0008046, filed on December 18, 1998.

The premixes of this invention may be used in the process of the present invention to provide an agglomerate comprising (i) an effective sweetener amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-

- 17 -

phenylalanine 1-methyl ester; (ii) a binding agent; (iii) a carrier; and (iv) an adjuvant, for example, a sweetener, a flavor or the like. Such an agglomerate may be prepared by adding the sweetener or other soluble adjuvant to the premix before the premix is applied to the carrier. Any soluble adjuvant which enhances the temporal properties of the agglomerate may be used in this invention. Adjuvants which may be added to the premix include, for example, glycerin and flavors.

The carriers which may be used in the present invention include, for example, dextrose, citric acid, maltodextrin, dextrose-maltodextrin blends, lactose, inulin, erythritol, sorbitol, sucrose, aspartame, acesulfame and its salts, sucralose, cyclamate, saccharin, stevioside, alitame and mixtures thereof.

The weight ratio of carrier to N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester may vary considerably depending upon a number of factors including the selection of binding agent and carrier, as well as the desired sweetening potency of the agglomerate. Generally, the weight ratio of carrier to N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester will be between 200:1 and 5:1. Preferably, the weight ratio should be between 100:1 and 40:1.

The amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester present in the agglomerates of the present invention may vary over a broad range, depending upon the particular end use desired. Preferably, the amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester present in the agglomerates of the

- 18 -

present invention is between about 0.001 weight percent and about 50 weight percent based upon the total amount of the agglomerate. More preferably, the amount of N-[N-(3,3-dimethylbutyl)-L- α -
5 aspartyl]-L-phenylalanine 1-methyl ester is between about 0.1 weight percent and about 5 weight percent.

The amount of carrier present in the agglomerates of the present invention also may vary over a broad
10 range, depending upon the particular carrier selected as well as the end use desired. Preferably, the amount of carrier present in the agglomerates of the present invention is between about 50 weight percent and about 99.9 weight
15 percent based upon the total amount of the agglomerate. More preferably, the amount of carrier is between about 75 weight percent and about 99 weight percent.

The amount of binding agent present in the agglomerates of the present invention also may vary over a broad range, depending upon the particular binding agent selected as well as the end use
20 desired. Preferably, the amount of binding agent present in the agglomerates of the present invention is between about 0.1 weight percent and about 15 weight percent based upon the total amount of the
25 agglomerate. More preferably, the amount of binding agent is between about 1 weight percent and about 7 weight percent.
30

The agglomerates of this invention may be dry blended with blending agents to prepare various products including, for example, tabletop sweeteners
35 and powdered soft drink mixes. Exemplary blending agents which may be dry blended with the agglomerates of this invention include aspartame,

- 19 -

acesulfame and its salts, sucralose, saccharin, alitame, cyclamates, stevia derivatives, thaumatin, sucrose, fructose, dextrose, polyol sugar alcohols, citric acid, dextrin, maltodextrin, dextrose-
5 maltodextrin blends, lactose, inulin, erythritol, sorbitol, stevioside, hydroxypropylmethyl cellulose, carboxymethyl cellulose, polyvinylpyrrolidone, and mixtures thereof.

10 The amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate that may be present in the tabletop sweeteners of this invention may vary over a broad range. Preferably, the amount of N-[N-(3,3-dimethylbutyl)-
15 L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate present in the tabletop sweeteners of this invention is between about 0.01 weight percent and about 100 weight percent based upon the total amount of the tabletop sweetener. More preferably,
20 the amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate is between about 0.5 weight percent and about 30 weight percent. Most preferably, the amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine
25 1-methyl ester agglomerate is between about 1 weight percent and about 10 weight percent.

The amount of blending agent that may be present in the tabletop sweeteners of this invention may vary
30 over a broad range. Preferably, the amount of blending agent present in the tabletop sweeteners of this invention is between about 50 weight percent and about 99 weight percent based upon the total amount of the tabletop sweetener. More preferably,
35 the amount of blending agent is between about 90 weight percent and about 99 weight percent.

- 20 -

The amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate that may be present in the powdered soft drink mixes of this invention may vary over a broad range.

5 Preferably, the amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate present in the powdered soft drink mixes of this invention is between about 4 weight percent and about 100 weight percent based upon the total
10 amount of the powdered soft drink mix. More preferably, the amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate is between about 30 weight percent and about 45 weight percent. In particular, preblend
15 products which carry flavors and then are dry blended are preferred. Such preblend products include, for example, citric acid or other dissolving aids.

20 The particle size distribution of the N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate may be determined by sifting the agglomerate through screens of various sizes. For example, the agglomerate may be sifted
25 with screens ranging in size from 14 mesh to 140 mesh or higher. Typically, at least about 65% of the particles of the agglomerate will pass through a 40 mesh screen and less than about 20% of the agglomerate particles will pass through 100 mesh
30 screen. In general, less than about 70% of the particles of the agglomerate will pass through a 60 mesh screen and less than about 5% of the agglomerate particles will pass through a 140 mesh screen. Typically, at least 50% of the particles of
35 the agglomerate are between about 60 mesh and 100 mesh in size.

- 21 -

The product may be screened to produce a narrower particle size distribution, if desired. For example, a 14 mesh screen may be used to remove large particles and produce a product of especially good appearance. Particles smaller than 120 mesh may be removed to obtain an agglomerate with improved flow properties. A narrower particle size distribution may be obtained if desired for particular applications.

The particle size distribution of the N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate may be controlled by a variety of factors including the selection of binding agent, the concentration of the binding agent in solution, the spray rate of the spray solution, the atomization air pressure and the particular carrier used. Those skilled in the art will appreciate that a desired particle size distribution may be obtained by varying one or more of the aforementioned factors. For example, increasing the spray rate is known to increase the average particle size.

The examples which follow are intended to illustrate certain preferred embodiments of the invention, and no limitation of the invention is implied.

EXAMPLE 1

Preparation of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester/Dextrose Agglomerate
(Maltodextrin Binder)

529 grams of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester was dissolved in 3.143 Kg of ethanol. In a separate

- 22 -

container, 635 grams of maltodextrin was dissolved
in 7.333 Kg of water. The two solutions were
combined and heated to 40°C. 20.0 Kg of dextrose
was charged into a removable bowl of a batch fluid
bed agglomeration unit. The dextrose was fluidized
and heated to 40°C by adjusting the inlet air
temperature of the agglomeration unit to between
70°C and 75°C. The N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl
ester/maltodextrin solution was sprayed onto the
fluidized dextrose at a spray rate of 200 ml/min.
The atomization air pressure was maintained at 2.5
bar. The particle size distribution of the
resulting N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-
phenylalanine 1-methyl ester/dextrose agglomerate is
set forth in Table 1.

EXAMPLE 2

Preparation of N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl ester/Dextrose
Agglomerate
(Maltodextrin Binder)

A N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-
phenylalanine 1-methyl ester/dextrose agglomerate
was prepared as described in Example 1, except that
the atomization air pressure was lowered to 1.5 bar.
The particle size distribution of the resultant
agglomerate is set forth in Table 1.

- 23 -

EXAMPLE 3

Preparation of N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl ester/Dextrose
Agglomerate
(Maltodextrin Binder)

A N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-
phenylalanine 1-methyl ester/dextrose agglomerate
was prepared as described in Example 1, except that
the atomization air pressure was lowered to 1.0 bar,
and the spray rate was increased to 250-275 ml/min.
The particle size distribution of the resultant
agglomerate is set forth in Table 1.

EXAMPLE 4

Preparation of N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl ester/Dextrose
Agglomerate
(Hydroxypropylmethyl Cellulose Binder)

539 grams of N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl ester was
dissolved in 3.754 Kg of ethanol. In a separate
container, 204 grams of hydroxypropylmethyl
cellulose (HPMC) was dissolved in 8.759 Kg of water.
The two solutions were combined to form a spray
solution, and heated to 40°C. 20.0 Kg of dextrose
was charged into a removable bowl of a batch fluid
bed agglomeration unit. The dextrose was fluidized
and heated to 38°C to 40°C. The spray solution was
applied to the fluidized dextrose using the spray
nozzle of the agglomeration unit. The atomization
air pressure was set at 1.5 bar. The particle size
distribution of the resultant N-[N-(3,3-
dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-
methyl ester/dextrose agglomerate is set forth in
Table 1.

- 24 -

Table 1. Particle Size Distribution of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester/Dextrose Agglomerates

Screen Size	Ex. 1	Ex. 2	Ex. 3	Ex. 4
20 Mesh	0.00	0.10	0.50	0.10
40 Mesh	2.70	3.80	5.50	9.00
60 Mesh	21.90	35.40	38.50	69.80
100 Mesh	82.10	90.10	89.80	97.40
140 Mesh	94.30	98.90	99.10	99.90
Pan	4.70	1.10	0.90	0.10

EXAMPLE 5

Preparation of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester/Citric Acid Agglomerate
(Hydroxypropylmethyl Cellulose Binder)

212 grams of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester was dissolved in 900 grams of ethanol. In a separate container, 71 grams of HPMC was dissolved in 5.102 Kg of water. The two solutions were combined to form a spray solution, and heated to 40°C. 28.0 Kg of fine granular citric acid was charged into a removable bowl of a batch fluid bed agglomeration unit. The citric acid was fluidized and heated to 40°C. The spray solution was applied to the fluidized citric acid using the spray nozzle of the agglomeration unit. The atomization air pressure was maintained at 2.5 bar. The spray rate was set at 100 ml/min. The particle size distribution of the resultant N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester/citric acid agglomerate is set forth in Table 2.

- 25 -

EXAMPLE 6

Preparation of N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl ester/Citric Acid
Agglomerate

5 (Hydroxypropylmethyl Cellulose Binder)

10 A N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-
phenylalanine 1-methyl ester/citric acid agglomerate
was prepared as described in Example 5, except that
the spray rate was increased to 150 ml/min. The
particle size distribution of the resultant N-[N-
(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-
methyl ester/citric acid agglomerate is set forth in
Table 2.

15

Table 2. Particle Size Distribution of N-[N-(3,3-
dimethylbutyl)-L- α -aspartyl]-L-
phenylalanine 1-methyl ester/Citric Acid
Agglomerates

20

Screen Size	Ex. 5	Ex. 6
20 Mesh	0.00	0.10
40 Mesh	20.30	37.60
60 Mesh	89.40	96.00
100 Mesh	100.00	100.40
Pan	0.00	0.10

25

EXAMPLE 7

30 Preparation of Tabletop Sweetener Product with a N-
[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine
1-methyl ester Agglomerate

35 A tabletop sweetener product was prepared
by blending a N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl ester/dextrose
agglomerate prepared as described in Example 3 with

- 26 -

Unidex using a 4.6 ft³ Littleford Lodige high shear mixer (Littleford Brothers, Inc., Cincinnati, Ohio).

5 Eight 40 Kg batches of the tabletop
sweetener were prepared by blending in each batch,
1.306 Kg of the N-[N-(3,3-dimethylbutyl)-L- α -
aspartyl]-L-phenylalanine 1-methyl ester/dextrose
agglomerate with 38.7 Kg of Unidex. The amount of
10 N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-
phenylalanine 1-methyl ester in each batch was
therefore 0.07956 weight percent based on the total
weight of the tabletop sweetener product.

15 After thorough mixing, six samples were
taken from each batch and analyzed for N-[N-(3,3-
dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-
methyl ester content. The results of these
measurements are set forth in Table 3. These
results show the excellent content uniformity that
20 may obtained in products prepared according to the
present invention. The 48 samples had an average
N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-
phenylalanine 1-methyl ester content of 0.07713%,
with a standard deviation of 0.00361, and a relative
25 standard deviation of only 4.68%.

- 27 -

Table 3. N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester content of Tabletop Sweetener Products

5

Sample	Batch 1	Batch 2	Batch 3	Batch 4	Batch 5	Batch 6	Batch 7	Batch 8
1	0.08334	0.07638	0.08067	0.07794	0.07624	0.07938	0.07808	0.07492
2	0.07532	0.07602	0.07941	0.07603	0.07834	0.08165	0.07231	0.07241
3	0.07782	0.07576	0.07440	0.08125	0.07638	0.09162	0.07708	0.07946
4	0.07532	0.07415	0.07465	0.07426	0.07634	0.07558	0.07932	0.07495
5	0.07333	0.07701	0.07174	0.08110	0.07628	0.08141	0.07752	0.07308
6	0.08052	0.07904	0.07618	0.07607	0.08188	0.07610	0.07246	0.07167

10

15

Other variations and modifications of this invention will be obvious to those skilled in the art. This invention is not limited except as set forth in the claims.

- 28 -

WHAT IS CLAIMED IS:

1. A process for preparing an agglomerate of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and a carrier comprising the steps of:
 - (a) providing a premix solution comprising N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and a binding agent;
 - (b) heating the premix solution of step (a) to a temperature effective to effect mixing of said premix;
 - (c) fluidizing a carrier; and
 - (d) applying the premix solution of step (b) onto said fluidized carrier to form an agglomerate comprising said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said carrier.
2. The process of claim 1, wherein in step (d), said premix solution is applied onto said fluidized carrier by spraying said premix solution onto said fluidized carrier.
3. The process of claim 1, wherein step (a) comprises the steps of:
 - (i) preparing a solution of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester;
 - (ii) preparing a solution of a binding agent; and
 - (iii) combining the solution of step (i) with the solution of step (ii).
4. The process of claim 1, wherein said premix solution further comprises a sweetener selected from the group consisting of aspartame, acesulfame salts, sucralose, saccharin, alitame, cyclamates, stevia

- 29 -

derivatives, thaumatin, sucrose, fructose, dextrose, polyol sugar alcohols, and mixtures thereof.

5. The process of claim 2, wherein preparation of the premix solution comprises combining said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester with a solvent selected from the group consisting of ethanol, isopropanol, methanol, water and mixtures thereof.

6. The process of claim 5, wherein said solvent is ethanol.

7. The process of claim 1, wherein said binding agent is selected from the group consisting of maltodextrin, dextrose-maltodextrin blends, hydroxypropylmethyl cellulose, carboxymethyl cellulose, polyvinylpyrrolidone, sucrose and mixtures thereof.

8. The process of claim 7, wherein said binding agent is maltodextrin.

9. The process of claim 7, wherein said binding agent is hydroxypropylmethyl cellulose.

10. The process of claim 7, wherein said binding agent is combined with a solvent selected from the group consisting of water, ethanol, isopropanol, methanol and mixtures thereof.

11. The process of claim 10, wherein said solvent is water.

12. The process of claim 7, wherein said carrier is selected from the group consisting of dextrose, citric acid, maltodextrin, dextrose-maltodextrin blends, lactose, inulin, erythritol, sorbitol, sucrose,

- 30 -

aspartame, acesulfame salts, sucralose, cyclamate, saccharin, stevioside, alitame and mixtures thereof.

13. The process of claim 12, wherein said carrier is dextrose.

14. The process of claim 12, wherein said carrier is citric acid.

15. The process of claim 12, wherein said carrier is sucrose.

16. The process of claim 12, wherein said carrier is fluidized at a temperature between about 30°C and about 50°C.

17. The process of claim 16, wherein step (b) comprises heating the premix solution of step (a) to between about 35°C and about 50°C.

18. The process of claim 1, wherein steps (c) and (d) are performed using a batch fluid bed agglomerator.

19. The process of claim 1, wherein steps (c) and (d) are performed using a continuous fluid bed agglomerator.

20. The process of claim 1, wherein the weight ratio of said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester to said carrier is from about 1:5 to about 1:200.

21. The process of claim 1, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is present in an amount from about 0.001 weight percent to about 50 weight percent based upon the total amount of said agglomerate.

- 31 -

22. The process of claim 1, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is present in an amount from about 0.1 weight percent to about 5 weight percent based upon the total amount of said agglomerate.

23. The process of claim 1, wherein said carrier is present in an amount from about 50 weight percent to about 99.9 weight percent based upon the total amount of said agglomerate.

24. The process of claim 1, wherein said carrier is present in an amount from about 75 weight percent to about 99 weight percent based upon the total amount of said agglomerate.

25. The process of claim 1, wherein said binding agent is present in an amount from about 0.1 weight percent to about 15 weight percent based upon the total amount of said agglomerate.

26. The process of claim 1, wherein said binding agent is present in an amount from about 1 weight percent to about 7 weight percent based upon the total amount of said agglomerate.

27. A process for preparing a blend comprising a N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate and a blending agent, said process comprising the steps of:

(a) providing a premix solution comprising N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and a binding agent;

(b) heating the premix solution of step (a) to a temperature effective to effect mixing of said premix;

(c) fluidizing a carrier;

- 32 -

(d) applying the premix solution of step (b) onto said fluidized carrier to form an agglomerate comprising said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said carrier; and
(e) dry blending said agglomerate of step (d) with a blending agent.

28. The process of claim 27, wherein said blending agent comprises a substance selected from the group consisting of aspartame, acesulfame salts, sucralose, saccharin, alitame, cyclamates, stevia derivatives, thaumatin, sucrose, fructose, dextrose, polyol sugar alcohols, citric acid, dextrin, maltodextrin, dextrose-maltodextrin blends, lactose, inulin, erythritol, sorbitol, stevioside, hydroxypropylmethyl cellulose, carboxymethyl cellulose, polyvinylpyrrolidone, and mixtures thereof.

29. The process of claim 28, wherein said substance is dextrose.

30. The process of claim 28, wherein said substance is maltodextrin.

31. The process of claim 28, wherein said substance is a mixture of dextrose and maltodextrin.

32. The process of claim 28, wherein said substance is lactose.

33. The product obtained by the process comprising the steps of:

(a) providing a premix solution comprising N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and a binding agent;

- 33 -

(b) heating the premix solution of step (a) to a temperature effective to effect mixing of said premix;

(c) fluidizing a carrier; and

(d) applying the premix solution of step (b) onto said fluidized carrier to form an agglomerate comprising said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said carrier.

34. The product of claim 33, wherein in step (d), said premix solution is applied onto said fluidized carrier by spraying said premix solution onto said fluidized carrier.

35. The product of claim 33, wherein step (a) comprises the steps of:

(i) preparing a solution of N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester;

(ii) preparing a solution of a binding agent; and

(iii) combining the solution of step (i) with the solution of step (ii).

36. The product of claim 33, wherein said premix solution further comprises a sweetener selected from the group consisting of aspartame, acesulfame salts, sucralose, saccharin, alitame, cyclamates, stevia derivatives, thaumatin, sucrose, fructose, dextrose, polyol sugar alcohols, and mixtures thereof.

37. The product of claim 33, wherein preparation of the premix solution comprises combining said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester with a solvent selected from the group consisting

- 34 -

of ethanol, isopropanol, methanol, water and mixtures thereof.

38. The product of claim 37, wherein said solvent is ethanol.

39. The product of claim 33, wherein said binding agent is selected from the group consisting of maltodextrin, dextrose-maltodextrin blends, hydroxypropylmethyl cellulose, carboxymethyl cellulose, polyvinylpyrrolidone, sucrose and mixtures thereof.

40. The product of claim 39, wherein said binding agent is maltodextrin.

41. The product of claim 39, wherein said binding agent is hydroxypropylmethyl cellulose.

42. The product of claim 39, wherein said binding agent is combined with a solvent selected from the group consisting of water, ethanol, isopropanol, methanol and mixtures thereof.

43. The product of claim 42, wherein said solvent is water.

44. The product of claim 39, wherein said carrier is selected from the group consisting of dextrose, citric acid, maltodextrin, dextrose-maltodextrin blends, lactose, inulin, erythritol, sorbitol, sucrose, aspartame, acesulfame salts, sucralose, cyclamate, saccharin, stevioside, alitame and mixtures thereof.

45. The product of claim 44, wherein said carrier is dextrose.

- 35 -

46. The product of claim 44, wherein said carrier is citric acid.

47. The product of claim 44, wherein said carrier is sucrose.

48. The product of claim 44, wherein said carrier is fluidized at a temperature between about 30°C and about 50°C.

49. The product of claim 48, wherein step (b) comprises heating the premix solution of step (a) to between about 35°C and about 50°C.

50. The product of claim 33, wherein steps (c) and (d) are performed using a batch fluid bed agglomerator.

51. The product of claim 33, wherein steps (c) and (d) are performed using a continuous fluid bed agglomerator.

52. The product of claim 33, wherein the weight ratio of said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester to said carrier is from about 1:5 to about 1:200.

53. The product of claim 33, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is present in an amount from about 0.001 weight percent to about 50 weight percent based upon the total amount of said agglomerate.

54. The product of claim 33, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is present in an amount from about 0.1 weight percent to about 5 weight percent based upon the total amount of said agglomerate.

- 36 -

55. The product of claim 33, wherein said carrier is present in an amount from about 50 weight percent to about 99.9 weight percent based upon the total amount of said agglomerate.

56. The product of claim 33, wherein said carrier is present in an amount from about 75 weight percent to about 99 weight percent based upon the total amount of said agglomerate.

57. The product of claim 33, wherein said binding agent is present in an amount from about 0.1 weight percent to about 15 weight percent based upon the total amount of said agglomerate.

58. The product of claim 33, wherein said binding agent is present in an amount from about 1 weight percent to about 7 weight percent based upon the total amount of said agglomerate.

59. The product of claim 33, wherein said agglomerate is comprised of particles, more than about 80 percent of which are greater in size than 100 mesh and more than about 95 percent of which are greater in size than 140 mesh.

60. The product of claim 59, wherein more than about 70 percent of the particles of said agglomerate are smaller in size than 60 mesh.

61. The product of claim 60, wherein at least 50 percent of the particles of said agglomerate are between about 60 mesh and 100 mesh in size.

62. The product of claim 33, wherein said process of preparing said product further comprises the step of

- 37 -

dry blending said agglomerate of step (d) with a blending agent.

63. The product of claim 62, wherein said blending agent comprises a substance selected from the group consisting of aspartame, acesulfame salts, sucralose, saccharin, alitame, cyclamates, stevia derivatives, thaumatin, sucrose, fructose, dextrose, polyol sugar alcohols, dextrin, citric acid, maltodextrin, dextrose-maltodextrin blends, lactose, inulin, erythritol, sorbitol, stevioside, hydroxypropylmethyl cellulose, carboxymethyl cellulose, polyvinylpyrrolidone, and mixtures thereof.

64. The product of claim 63, wherein said substance is dextrose.

65. The product of claim 63, wherein said substance is maltodextrin.

66. The product of claim 63, wherein said substance is a mixture of dextrose and maltodextrin.

67. The product of claim 63, wherein said substance is lactose.

68. A N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate comprising:

(i) an effective sweetener amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester;

(ii) a binding agent; and

(iii) a carrier;

wherein the sweetener and the binding agent are applied onto the carrier.

69. The agglomerate of claim 68, wherein said binding agent is selected from the group consisting of

- 38 -

maltodextrin, dextrose-maltodextrin blends, hydroxypropylmethyl cellulose, carboxymethyl cellulose, polyvinylpyrrolidone, sucrose and mixtures thereof.

70. The agglomerate of claim 69, wherein said binding agent is maltodextrin.

71. The agglomerate of claim 69, wherein said binding agent is hydroxypropylmethyl cellulose.

72. The agglomerate of claim 69, wherein said carrier is selected from the group consisting of dextrose, citric acid, maltodextrin, dextrose-maltodextrin blends, lactose, inulin, erythritol, sorbitol, sucrose, aspartame, acesulfame salts, sucralose, cyclamate, saccharin, stevioside, alitame and mixtures thereof.

73. The agglomerate of claim 72, wherein said carrier is dextrose.

74. The agglomerate of claim 72, wherein said carrier is citric acid.

75. The agglomerate of claim 72, wherein said binding agent and said carrier are the same.

76. The agglomerate of claim 68, wherein the weight ratio of said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester to said carrier is from about 1:5 to about 1:200.

77. The agglomerate of claim 68, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is present in an amount from about 0.001 weight percent to about 50 weight percent based upon the total amount of said agglomerate.

- 39 -

78. The agglomerate of claim 68, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is present in an amount from about 0.1 weight percent to about 5 weight percent based upon the total amount of said agglomerate.

79. The agglomerate of claim 68, wherein said carrier is present in an amount from about 50 weight percent to about 99.9 weight percent based upon the total amount of said agglomerate.

80. The agglomerate of claim 68, wherein said carrier is present in an amount from about 75 weight percent to about 99 weight percent based upon the total amount of said agglomerate.

81. The agglomerate of claim 68, wherein said binding agent is present in an amount from about 0.1 weight percent to about 15 weight percent based upon the total amount of said agglomerate.

82. The agglomerate of claim 68, wherein said binding agent is present in an amount from about 1 weight percent to about 7 weight percent based upon the total amount of said agglomerate.

83. The agglomerate of claim 68, wherein said agglomerate is comprised of particles, more than about 80 percent of which are greater in size than 100 mesh and more than about 95 percent of which are greater in size than 140 mesh.

84. The agglomerate of claim 83, wherein more than about 70 percent of the particles of said agglomerate are smaller in size than 60 mesh.

- 40 -

85. The agglomerate of claim 84, wherein at least 50 percent of the particles of said agglomerate are between about 60 mesh and 100 mesh in size.

86. The agglomerate of claim 68, further comprising a flavor.

87. The agglomerate of claim 68, further comprising a natural or high intensity sweetener.

88. The agglomerate of claim 87, wherein said binding agent is selected from the group consisting of maltodextrin, dextrose-maltodextrin blends, hydroxypropylmethyl cellulose, carboxymethyl cellulose, polyvinylpyrrolidone, sucrose and mixtures thereof.

89. The agglomerate of claim 88, wherein said carrier is selected from the group consisting of dextrose, citric acid, maltodextrin, dextrose-maltodextrin blends, lactose, inulin, erythritol, sorbitol, sucrose, aspartame, acesulfame salts, sucralose, cyclamate, saccharin, stevioside, alitame and mixtures thereof.

90. The agglomerate of claim 89, wherein said natural or high intensity sweetener is selected from the group consisting of aspartame, acesulfame salts, sucralose, saccharin, alitame, cyclamates, stevia derivatives, thaumatin, sucrose, fructose, dextrose, polyol sugar alcohols, and mixtures thereof.

91. A tabletop sweetener comprising a N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate, said agglomerate comprising:

(i) an effective sweetener amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester;

- 41 -

- (ii) a binding agent; and
- (iii) a carrier;

wherein the sweetener and the binding agent are applied onto the carrier.

92. The tabletop sweetener of claim 91, wherein said binding agent is selected from the group consisting of maltodextrin, dextrose-maltodextrin blends, hydroxypropylmethyl cellulose, carboxymethyl cellulose, polyvinylpyrrolidone, sucrose and mixtures thereof.

93. The tabletop sweetener of claim 92, wherein said binding agent is maltodextrin.

94. The tabletop sweetener of claim 92, wherein said binding agent is hydroxypropylmethyl cellulose.

95. The tabletop sweetener of claim 91, wherein said carrier is selected from the group consisting of dextrose, citric acid, maltodextrin, dextrose-maltodextrin blends, lactose, inulin, erythritol, sorbitol, sucrose, aspartame, acesulfame salts, sucralose, cyclamate, saccharin, stevioside, alitame and mixtures thereof.

96. The tabletop sweetener of claim 95, wherein said carrier is dextrose.

97. The tabletop sweetener of claim 95, wherein said carrier is citric acid.

98. The tabletop sweetener of claim 95, wherein said binding agent and said carrier are the same.

99. A tabletop sweetener according to claim 91, wherein said tabletop sweetener comprises:

(a) N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in an amount from about

- 42 -

0.01 weight percent to about 0.5 weight percent based upon the total amount of said tabletop sweetener;

(b) maltodextrin in an amount from about 2 weight percent to about 4 weight percent based upon the total amount of said tabletop sweetener; and

(c) dextrose in an amount from about 96 weight percent to about 98 weight percent based upon the total amount of said tabletop sweetener.

100. A tabletop sweetener according to claim 91, wherein said tabletop sweetener comprises:

(a) N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in an amount from about 0.01 weight percent to about 0.5 weight percent based upon the total amount of said tabletop sweetener;

(b) maltodextrin in an amount from about 2 weight percent to about 4 weight percent based upon the total amount of said tabletop sweetener;

(c) dextrose in an amount from about 92 weight percent to about 97 weight percent based upon the total amount of said tabletop sweetener; and

(d) aspartame in an amount of from about 1% to about 4% based upon the total amount of said tabletop sweetener.

101. A tabletop sweetener according to claim 91, wherein said tabletop sweetener comprises:

(a) N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in an amount from about 0.01 weight percent to about 0.5 weight percent based upon the total amount of said tabletop sweetener; and

(b) sucrose in an amount from about 95 weight percent to about 99.9 weight percent based upon the total amount of said tabletop sweetener.

- 43 -

102. The tabletop sweetener of claim 91, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is present in an amount from about 0.001 weight percent to about 5 weight percent based upon the total amount of said tabletop sweetener.

103. The tabletop sweetener of claim 91, wherein said carrier is present in an amount from about 0.005 weight percent to about 99.9 weight percent based upon the total amount of said tabletop sweetener.

104. The tabletop sweetener of claim 91, wherein said carrier is present in an amount from about 0.0075 weight percent to about 99 weight percent based upon the total amount of said tabletop sweetener.

105. The tabletop sweetener of claim 91, wherein said binding agent is present in an amount from about 0.00001 weight percent to about 15 weight percent based upon the total amount of said tabletop sweetener.

106. The tabletop sweetener of claim 91, wherein said binding agent is present in an amount from about 0.01 weight percent to about 7 weight percent based upon the total amount of said tabletop sweetener.

107. The tabletop sweetener of claim 91, further comprising a blending agent.

108. The tabletop sweetener of claim 107, wherein said blending agent comprises a substance selected from the group consisting of aspartame, acesulfame salts, sucralose, saccharin, alitame, cyclamates, stevia

- 44 -

derivatives, thaumatin, sucrose, fructose, dextrose, polyol sugar alcohols, dextrose, citric acid, dextrin, maltodextrin, dextrose-maltodextrin blends, lactose, inulin, erythritol, sorbitol, stevioside, hydroxypropylmethyl cellulose, carboxymethyl cellulose, polyvinylpyrrolidone, and mixtures thereof.

109. The tabletop sweetener of claim 108, wherein said substance is dextrose.

110. The tabletop sweetener of claim 108, wherein said substance is maltodextrin.

111. The tabletop sweetener of claim 108, wherein said substance is a mixture of dextrose and maltodextrin.

112. The tabletop sweetener of claim 108, wherein said substance is lactose.

113. The tabletop sweetener of claim 107, wherein said blending agent is present in an amount from about 50 weight percent to about 99 weight percent based upon the total amount of said tabletop sweetener.

114. A powdered soft drink mix comprising a N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate, said agglomerate comprising

(i) an effective sweetener amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester;

(ii) a binding agent; and

(iii) a carrier;

wherein the sweetener and the binding agent are applied onto the carrier.

115. The powdered soft drink mix of claim 114, wherein said binding agent is selected from the group consisting of maltodextrin, dextrose-maltodextrin blends, hydroxypropylmethyl cellulose, carboxymethyl

- 45 -

cellulose, polyvinylpyrrolidone, sucrose and mixtures thereof.

116. The powdered soft drink mix of claim 115, wherein said binding agent is maltodextrin.

117. The powdered soft drink mix of claim 115, wherein said binding agent is hydroxypropylmethyl cellulose.

118. The powdered soft drink mix of claim 114, wherein said carrier is selected from the group consisting of dextrose, citric acid, maltodextrin, dextrose-maltodextrin blends, lactose, inulin, erythritol, sorbitol, sucrose, aspartame, acesulfame salts, sucralose, cyclamate, saccharin, stevioside, alitame and mixtures thereof.

119. The powdered soft drink mix of claim 118, wherein said carrier is dextrose.

120. The powdered soft drink mix of claim 118, wherein said carrier is citric acid.

121. The powdered soft drink mix of claim 114, wherein said binding agent and said carrier are the same.

122. The powdered soft drink mix of claim 114, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is present in an amount from about 0.00004 weight percent to about 50 weight percent based upon the total weight of said powdered soft drink mix.

123. The powdered soft drink mix of claim 114, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is present in an amount from about 0.004 weight percent to about 5 weight

- 46 -

percent based upon the total weight of said powdered soft drink mix.

124. The powdered soft drink mix of claim 114, wherein said carrier is present in an amount from about 2 weight percent to about 99.9 weight percent based upon the total amount of said powdered soft drink mix.

125. The powdered soft drink mix of claim 114, wherein said carrier is present in an amount from about 3 weight percent to about 99 weight percent based upon the total amount of said powdered soft drink mix.

126. The powdered soft drink mix of claim 114, wherein said binding agent is present in an amount from about 0.004 weight percent to about 15 weight percent based upon the total amount of said powdered soft drink mix.

127. The powdered soft drink mix of claim 114, wherein said binding agent is present in an amount from about 0.04 weight percent to about 7 weight percent based upon the total amount of said powdered soft drink mix.

128. The powdered soft drink mix of claim 114, further comprising a blending agent.

129. The powdered soft drink mix of claim 128, wherein said blending agent comprises a substance selected from the group consisting of aspartame, acesulfame salts, sucralose, saccharin, alitame, cyclamates, stevia derivatives, thaumatin, sucrose, fructose, dextrose, polyol sugar alcohols, dextrose, citric acid, dextrin, maltodextrin, dextrose-maltodextrin blends, lactose, inulin, erythritol, sorbitol, stevioside, hydroxypropylmethyl cellulose, carboxymethyl cellulose, polyvinylpyrrolidone, and mixtures thereof.

- 47 -

130. The powdered soft drink mix of claim 129, wherein said substance is dextrose.

131. The powdered soft drink mix of claim 129, wherein said substance is maltodextrin.

132. The powdered soft drink mix of claim 129, wherein said substance is a mixture of dextrose and maltodextrin.

133. The powdered soft drink mix of claim 129, wherein said substance is lactose.

134. A sweetener blend comprising a sweetener and a N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester agglomerate, said agglomerate comprising:

- (i) an effective sweetener amount of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester;
- (ii) a binding agent; and
- (iii) a carrier;

wherein the sweetener and the binding agent are applied onto the carrier.

135. The sweetener blend of claim 134, wherein said binding agent is selected from the group consisting of maltodextrin, dextrose-maltodextrin blends, hydroxypropylmethyl cellulose, carboxymethyl cellulose, polyvinylpyrrolidone, sucrose and mixtures thereof.

136. The sweetener blend of claim 135, wherein said binding agent is maltodextrin.

137. The sweetener blend of claim 135, wherein said binding agent is hydroxypropylmethyl cellulose.

138. The sweetener blend of claim 134, wherein said carrier is selected from the group consisting of

- 48 -

dextrose, citric acid, maltodextrin, dextrose-maltodextrin blends, lactose, inulin, erythritol, sorbitol, sucrose, aspartame, acesulfame salts, sucralose, cyclamate, saccharin, stevioside, alitame and mixtures thereof.

139. The sweetener blend of claim 138, wherein said carrier is dextrose.

140. The sweetener blend of claim 138, wherein said carrier is citric acid.

141. The sweetener blend of claim 138, wherein said binding agent and said carrier are the same.

142. The sweetener blend of claim 138, wherein said sweetener is selected from the group consisting of aspartame, acesulfame salts, sucralose, saccharin, alitame, cyclamates, stevia derivatives, thaumatin, sucrose, fructose, dextrose, polyol sugar alcohols, and mixtures thereof.