

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

06 January 2022 (06.01.2022)



(10) International Publication Number

WO 2022/003426 A1

(51) International Patent Classification:

A61P 1/16 (2006.01)

C07D 417/12 (2006.01)

C07B 59/00 (2006.01)

A61K 31/4427 (2006.01)

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with information concerning incorporation by reference of correct element or part (Rule 20.6)

(21) International Application Number:

PCT/IB2021/000661

(22) International Filing Date:

24 June 2021 (24.06.2021)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/046,304

30 June 2020 (30.06.2020)

US

63/046,309

30 June 2020 (30.06.2020)

US

17/314,538

07 May 2021 (07.05.2021)

US

(71) Applicant: POXEL SA [FR/FR]; Immeuble Le Sunway, 259-261 Avenue Jean Jaurès, 69007 Lyon (FR).

(72) Inventors: NAVARRE, Laure Françoise Valérie; 8 Rue Michel Félizat, Bâtiment I, 69007 Lyon (FR). GARDETTE, Emeline; 31 rue Soeur Bouvier, Bâtiment C, 69005 Lyon (FR). BOLZE, Sébastien; 382 Chemin des Dorriers, 01600 Massieux (FR). DEWITT, Sheila; 346 Bunker Hill Rd., Auburn, New Hampshire 03032 (US). JACQUES, Vincent; 118 Roger Avenue, Apt. #2, Somerville, Massachusetts 02144 (US).

(74) Agent: GRAHAM WATT & CO LLP; St Botolph's House, 7-9 St Botolph's Road, Sevenoaks Kent TN13 3AJ (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: CRYSTALLINE FORMS OF DEUTERIUM-ENRICHED PIOGLITAZONE

(57) Abstract: Provided herein are crystalline forms of deuterium-enriched (R)-pioglitazone and compositions thereof. Also provided herein are methods of using the crystalline forms of deuterium-enriched (R)-pioglitazone and compositions thereof.

WO 2022/003426 A1

CRYSTALLINE FORMS OF DEUTERIUM-ENRICHED PIOGLITAZONE**RELATED APPLICATIONS**

[001] This application claims the benefit of and priority to United States Patent
5 Application serial number 17/314,538, filed May 7, 2021, United States Provisional Patent
Application serial number 63/046,304, filed June 30, 2020, and United States Provisional
Patent Application serial number 63/046,309, filed June 30, 2020, the contents of each of
which are hereby incorporated by reference in their entirety.

BACKGROUND

10 **[002]** Thiazolidinediones (TZDs) are antidiabetic drugs that sensitize the body to insulin.
These compounds were empirically discovered to be agonists of peroxisome proliferator-
activated-receptors (PPARs), including PPAR- γ (PPAR γ), a ligand-activated nuclear receptor
that drives broad transcriptional programs associated with adipogenesis, lipid metabolism,
innate immune function, and metabolic homeostasis (see, e.g., J. Clin. Invest. 2000, 106,
15 1305-1307, Trends Endocrinol. Metab. 2012, 23, 205-215). As such, the antidiabetic
mechanism of action of TZDs has up to now been attributed to binding to, and activation of,
PPAR γ (see, e.g., J Biol Chem 1995, 270, 12953-12956, Nat Med 2013, 19, 557-566).

[003] Therapeutics that modulate PPARs have been commercialized for treating medical
disorders, such as metabolic disorders. One such example is the TZD, pioglitazone
20 hydrochloride, which has been approved by the United States Food and Drug
Administration as an adjunct to diet and exercise to improve glycemic control in adults with
type 2 diabetes mellitus in multiple clinical settings. Pioglitazone hydrochloride is marketed
under the registered trademark ACTOS[®] and the prescribing information for ACTOS[®]
explains that pioglitazone is an agonist of PPAR γ . PPAR γ -related adverse side effects of
25 ACTOS[®] have been reported including, for example, weight gain, edema and increased
incidence of bone fracture.

[004] However, emerging genetic and pharmacologic evidence suggests that TZDs exert
many of their beneficial effects independently of PPAR γ activation (see Biochem.
Pharmacol. 2005, 70, 177-188). For instance, liver and skeletal muscle remain responsive to
30 TZDs despite tissue-specific deletion of PPAR γ (see J Clin Invest 2003, 112, 608-618, J Biol
Chem 2012, 287, 23537-23548), and the whole-body insulin sensitizing effect of
rosiglitazone persists upon PPAR γ deletion in mature adipocytes (see Mol Cell Biol 2018, 38,
e00677-17). Pharmacologic evidence also suggests TZDs have PPAR γ -independent effects.
They can acutely alter metabolic homeostasis on a timescale that is likely too rapid to be

driven by broad gene expression changes (see Am. J. Physiol. Endocrinol. Metab. 2006, 291, E175-81), and several *in vitro* and *in vivo* experiments have shown that rank-order affinity for PPAR γ does not always correlate with efficacy (see Biochem. Pharmacol. 2005, 70, 177-188). Pioglitazone and other TZDs have also been shown to have anti-inflammatory activity, which seems to be, at least in part, mediated by a mechanism not involving PPARs (Curr. Drug Targets Inflamm. Allergy 2002, 1, 243-248).

[005] Recent work shows that most of the PPAR γ -independent effects of TZDs could be attributed to inhibition of the mitochondrial pyruvate carrier (MPC), an inner mitochondrial membrane transporter responsible for the uptake of glucose-derived pyruvate from the cytoplasm into the mitochondrial matrix (see Biochim. Biophys. Acta - Mol Cell Res, 2016, 1863, 2436–2442). TZDs are acute, specific inhibitors of MPC activity at clinically relevant concentrations (see Proc. Natl. Acad. Sci. USA, 2013, 110, 5422–5427). TZDs also inhibit mitochondrial complex I (see Diabetes 2004, 53, 1052–1059), though at higher and perhaps supraphysiological concentrations relative to the effect on the MPC. Despite pioglitazone decades of use in humans with T2DM, the repertoire of targets for pioglitazone and its exact mechanism of action is only now beginning to be unraveled.

[006] All TZDs, including pioglitazone, are a mixture of enantiomers and are characterized by the presence of a chiral center. This chiral center is prone to rapid, non-enzymatic inversion of configuration. Thus, the instability of the chiral center has prevented exploitation of the potential for differentiated pharmacology between the enantiomers of TZDs. For example, while the left-handed, or (*S*)- enantiomer, of rosiglitazone was identified as the most potent PPAR γ agonist, (*S*)-rosiglitazone rapidly equilibrated to create a 1:1 mixture of (*R*)- and (*S*)-rosiglitazone, preventing further *in vivo* characterization of the enantiomers (see Bioorg. Med. Chem. Lett., 1998, 8, 3657–8).

Furthermore, the anti-inflammatory effects of pioglitazone were shown to be uniquely associated with the (*R*)-enantiomer in a rat model of chronic obstructive pulmonary disease. This was achieved only after stabilization of the enantiomers in an acidic solution, followed by immediate intranasal dosing, which limited inversion during the time course of the study (see international patent application WO2010015818A1).

[007] Due to the increasing number of patients suffering from disorders such as those mentioned above, and the limitations of existing therapies, such as adverse side effects, there is a need for new therapeutic agents for treating medical disorders in which modulation of PPAR γ , anti-inflammatory, and/or MPC activity are predicted to be beneficial.


[008] The use of deuterium-enriched enantiomers of pioglitazone to treat such medical disorders has gained significant interest, as it is believed that they may provide additional therapeutic efficacy and reduce the rate and severity of side effects compared to pioglitazone. Recently, the characterization of the unique pharmacological and pharmacokinetic properties of the deuterium-enriched (*R*) and/or (*S*)-enantiomers of pioglitazone has been reported (see Hepatol. Commun. 2021, Apr 10 doi.org/10.1002/hep4.1723). Preclinical studies demonstrated that (*R*)-pioglitazone retains the efficacy of pioglitazone in nonalcoholic steatohepatitis (NASH), including reduced hepatic triglycerides, free fatty acids, cholesterol, steatosis, inflammation, hepatocyte enlargement, and fibrosis. Although both enantiomers inhibit MPC, deuterium-enriched (*R*)-pioglitazone (PXL065) shows limited to no PPAR γ activity, whereas (*S*)-pioglitazone appears responsible for the PPAR γ activity and associated weight gain. Both enantiomers reduce plasma glucose and hepatic fibrosis to the same extent as pioglitazone in preclinical mouse models. In a phase 1a clinical study, the safety and tolerability of PXL065 (7.5, 22.5, 30 mg) was demonstrated as well as preferential exposure to the (*R*)-enantiomer in comparison to 45 mg Actos[®]. These results concluded that PXL065 at a dose lower than 22.5 mg is predicted to exhibit efficacy for NASH equal to, or greater than, 45 mg pioglitazone without the potentially detrimental weight gain and edema.

[009] Developing solid forms of these deuterium-enriched pioglitazone enantiomers, however, that provide the physicochemical properties necessary to manufacture a commercial drug product with the required efficacy remains a significant challenge. This is due to the unpredictability in the outcome of solid form screenings for any given compound and the subsequent unpredictability of the physicochemical properties of any solid forms discovered. Thus, there is a need for new solid forms of deuterium-enriched pioglitazone derivatives that achieve the required physicochemical properties necessary to provide superior efficacy over pioglitazone.

SUMMARY


[010] In one aspect, the invention provides salt forms (e.g., crystalline salt forms) of deuterium-enriched (*R*)-pioglitazone (PXL065).

[011] In various embodiments of the invention, a deuterium chloride salt of a deuterium-enriched compound of formula (I):




wherein R^1 is H or D, provided that the total deuterium abundance in the compound of formula (I) is about 1.3 to about 4.

[012] In various embodiments of the invention, a crystalline deuterium chloride salt of a
5 deuterium-enriched compound of formula (I)




wherein R^1 is H or D, provided that the total deuterium abundance in the compound of formula (I) is about 1.3 to about 4, and wherein the crystalline deuterium chloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following
10 diffraction angles (2θ): $15.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, and $26.0^\circ \pm 0.2^\circ$.

[013] In various embodiments of the invention, a crystalline hydrochloride salt of a compound of formula (I-A):




[014] In various embodiments of the invention, a crystalline hydrochloride salt of the
15 compound of formula (I-A)



wherein the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): $15.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, and $26.0^\circ \pm 0.2^\circ$.


[015] In another aspect, the invention provides pharmaceutical materials comprising particles of a crystalline salt (e.g., deuterium chloride salt, hydrochloride salt) of deuterium-enriched (*R*)-pioglitazone.

[016] In various embodiments of the invention, the pharmaceutical material comprises
 5 particles of a crystalline deuterium chloride salt of a deuterium-enriched compound of formula (I)




wherein R¹ is H or D, provided that the total deuterium abundance in the compound of formula (I) is about 1.3 to about 4, and wherein the particles in the composition have a
 10 crystal shape selected from hexagonal, rod, and combinations thereof.

[017] In various embodiments of the invention, the pharmaceutical material comprises particles of a crystalline deuterium chloride salt of a deuterium-enriched compound of formula (I)




15 wherein R¹ is H or D, provided that the total deuterium abundance in the compound of formula (I) is about 1.3 to about 4, and wherein the particles have a particle size distribution which is defined by a d(0.9) of about 10 μm to about 800 μm preferably below 500 μm.

[018] In various embodiments of the invention, the pharmaceutical material comprising
 20 particles of a crystalline hydrochloride salt of the compound of formula (I-A)



wherein the particles in the composition have a needle-like crystal shape.

[019] In various embodiments of the invention, the pharmaceutical material comprising particles of a crystalline hydrochloride salt of the compound of formula (I-A)



wherein the particles have a particle size distribution which is defined by a d(0.9) of about 10 μm to about 800 μm preferably below 500 μm .

[020] In another aspect, the invention provides pharmaceutical compositions comprising a crystalline salt of deuterium-enriched (*R*)-pioglitazone described herein, or a pharmaceutical material described herein, and a pharmaceutically acceptable excipient. In various embodiments, the pharmaceutical compositions comprise a crystalline deuterium chloride salt of deuterium-enriched (*R*)-pioglitazone and a pharmaceutically acceptable excipient. In various embodiments, the pharmaceutical compositions comprise a crystalline hydrochloride salt of deuterium-enriched (*R*)-pioglitazone and a pharmaceutically acceptable excipient.

[021] In another aspect, the invention provides crystalline salt forms of deuterium-enriched (*R*)-pioglitazone, pharmaceutical materials, and pharmaceutical compositions useful for the treatment of the various conditions, diseases, and disorders described herein. In certain embodiments, the condition, disease, or disorder is a metabolic disorder. In certain embodiments, the condition, disease, or disorder is diabetes mellitus type 2 and/or nonalcoholic steatohepatitis. In certain embodiments, the condition, disease, or disorder is a neurological disorder. In some embodiments, the neurological disorder is adrenoleukodystrophy or adrenomyeloneuropathy.

BRIEF DESCRIPTION OF THE DRAWINGS

[022] FIG. 1A is an exemplary X-ray powder diffraction (XRPD) pattern of the crystalline deuterium chloride salt of deuterium-enriched (*R*)-pioglitazone.

[023] FIG. 1B is an exemplary X-ray powder diffraction (XRPD) pattern of the crystalline hydrogen chloride salt of deuterium-enriched (*R*)-pioglitazone overlaid with an X-ray powder diffraction (XRPD) pattern of the crystalline hydrogen chloride salt of pioglitazone.

[024] FIG. 2 is an exemplary optical micrograph of the crystalline deuterium chloride salt of deuterium-enriched (*R*)-pioglitazone at (A) 10x magnification and (B) 20x magnification.

- [025] FIG. 3 is an exemplary differential scanning calorimetry (DSC) curve of the crystalline deuterium chloride salt of deuterium-enriched (*R*)-pioglitazone.
- [026] FIG. 4A is an exemplary thermogravimetric analysis (TGA) curve of the crystalline deuterium chloride salt of deuterium-enriched (*R*)-pioglitazone.
- 5 [027] FIG. 4B is an exemplary thermal gravimetric mass spectrometry (TGMS) plot of the crystalline deuterium chloride salt of deuterium-enriched (*R*)-pioglitazone.
- [028] FIG. 5 is an exemplary proton nuclear magnetic resonance (¹H-NMR) spectrum of the crystalline deuterium chloride salt of deuterium-enriched (*R*)-pioglitazone.
- [029] FIG. 6 an exemplary X-ray powder diffraction (XRPD) pattern of the crystalline
- 10 hydrochloride salt of deuterium-enriched (*R*)-pioglitazone.
- [030] FIG. 7 is an exemplary optical micrograph of the crystalline hydrochloride salt of deuterium-enriched (*R*)-pioglitazone at (A) 10x magnification and (B) 20x magnification.
- [031] FIG. 8 is an exemplary differential scanning calorimetry (DSC) curve of the crystalline hydrochloride salt of deuterium-enriched (*R*)-pioglitazone.
- 15 [032] FIG. 9A is an exemplary thermogravimetric analysis (TGA) curve of the crystalline hydrochloride salt of deuterium-enriched (*R*)-pioglitazone.
- [033] FIG. 9B is an exemplary thermal gravimetric mass spectrometry (TGMS) plot of the crystalline hydrochloride salt of deuterium-enriched (*R*)-pioglitazone.
- [034] FIG. 10 is an exemplary ¹H-NMR spectrum of the crystalline hydrochloride salt of
- 20 deuterium-enriched (*R*)-pioglitazone.
- [035] FIG. 11 an exemplary X-ray powder diffraction (XRPD) pattern of the crystalline free base of deuterium-enriched (*R*)-pioglitazone.
- [036] FIG. 12 is an exemplary differential scanning calorimetry (DSC) curve of the crystalline free-base form of deuterium-enriched (*R*)-pioglitazone.
- 25 [037] FIG. 13A is an exemplary thermogravimetric analysis (TGA) curve of the crystalline free-base form of deuterium-enriched (*R*)-pioglitazone.
- [038] FIG. 13B is an exemplary thermal gravimetric mass spectrometry (TGMS) plot of the crystalline free-base form of deuterium-enriched (*R*)-pioglitazone.
- [039] FIG. 14 is an overlay of pH-dependent solubility profiles of the crystalline DCl and
- 30 HCl salts of deuterium-enriched (*R*)-pioglitazone, of the crystalline DCl salt of deuterium-enriched pioglitazone and of the crystalline HCl salt of pioglitazone.

[040] FIG. 15 is an exemplary plot of tablet hardness (N) vs compression force (kN) for active pharmaceutical ingredient (API) batches with varying particle size distribution (PSD) in μm .

5 [041] FIG. 16 is an exemplary plot of tablet hardness (N) vs compression force (kN) for API batches with varying particle size distribution (PSD) in μm at different tableting speeds.

[042] FIG. 17 is an exemplary plot of plasma levels of total pioglitazone (sum of deuterated and protonated (*R*)- and (*S*)-enantiomers) over time for doses of PXL065 (deuterium-enriched (*R*) pioglitazone) in dogs.

10 [043] FIG. 18 is an exemplary plot of plasma levels of total pioglitazone (sum of deuterated and protonated (*R*)- and (*S*)-enantiomers) over time for doses of PXL061 (deuterium-enriched pioglitazone) in dogs.

[044] FIG. 19 is an exemplary plot of plasma levels over time of total pioglitazone (sum of deuterated and protonated (*R*)- and (*S*)-enantiomers) for doses of PXL064 (deuterium-enriched (*S*)-pioglitazone) in dogs.

15 [045] FIG. 20 is an exemplary plot of plasma levels of total pioglitazone (sum of deuterated and protonated (*R*)- and (*S*)-enantiomers) over time for doses of pioglitazone and PXL065 in dogs.

[046] FIG. 21 is an overlay of total pioglitazone (sum of deuterated and protonated (*R*)- and (*S*)-enantiomers) dose normalized exposure curves for the crystalline hydrochloride salt of deuterium-enriched (*R*)-pioglitazone (22.5 mg dose) and Actos® (pioglitazone HCl, 45 mg dose) in humans, as further described in Example 11.

20 [047] FIG. 22 is an overlay of total pioglitazone (sum of deuterated and protonated (*R*)- and (*S*)-enantiomers) dose normalized exposure curves for the crystalline hydrochloride salt of deuterium-enriched (*R*)-pioglitazone (15 mg and 30 mg doses) and Actos® (pioglitazone HCl, 45 mg dose) in humans, as further described in Example 12.

DETAILED DESCRIPTION

[048] As generally described herein, the invention provides pharmaceutically acceptable salts of deuterium-enriched (*R*)-pioglitazone (e.g., crystalline salts of deuterium-enriched (*R*)-pioglitazone), also referred to herein as a compound of formula I, pharmaceutical compositions containing the same, and methods of using the salt forms of deuterium-enriched (*R*)-pioglitazone to treat medical conditions, diseases, and disorders (e.g., a neurological disorder, a cancer, a respiratory disorder, a metabolic disorder, a hepatitis,

cardiovascular disease, an inflammatory disorder, an immune-mediated disorder, a dermatological disorder, or a skin defect) in a subject in need thereof. In particular, the invention provides crystalline hydrochloride and deuterium chloride salts of deuterium-enriched (*R*)-pioglitazone with unexpectedly improved solubilities at physiologically
5 relevant pHs and in vivo bioavailability compared to the commercially available pioglitazone hydrochloride salt (Actos®).

Definitions

[049] To facilitate an understanding of the present invention, a number of terms and
10 phrases are defined below.

[050] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. The abbreviations used herein have their conventional meaning within the chemical and biological arts. The chemical structures and formulae set forth herein are
15 constructed according to the standard rules of chemical valency known in the chemical arts.

[051] Throughout the description, where compositions and kits are described as having, including, or comprising specific components, or where processes and methods are described as having, including, or comprising specific steps, it is contemplated that, additionally, there are compositions and kits of the present invention that consist
20 essentially of, or consist of, the recited components, and that there are processes and methods according to the present invention that consist essentially of, or consist of, the recited processing steps.

[052] In the application, where an element or component is said to be included in and/or selected from a list of recited elements or components, it should be understood
25 that the element or component can be any one of the recited elements or components, or the element or component can be selected from a group consisting of two or more of the recited elements or components.

[053] Further, it should be understood that elements and/or features of a composition or a method described herein can be combined in a variety of ways without departing from
30 the spirit and scope of the present invention, whether explicit or implicit herein. For example, where reference is made to a particular compound, that compound can be used in various embodiments of compositions of the present invention and/or in methods of the present invention, unless otherwise understood from the context. In other words, within this application, embodiments have been described and depicted in a way that enables a

clear and concise application to be written and drawn, but it is intended and will be appreciated that embodiments may be variously combined or separated without parting from the present teachings and invention(s). For example, it will be appreciated that all features described and depicted herein can be applicable to all aspects of the invention(s) described and depicted herein.

[054] The articles “a” and “an” are used in this disclosure to refer to one or more than one (i.e., to at least one) of the grammatical object of the article, unless the context is inappropriate. By way of example, “an element” means one element or more than one element.

[055] The term “and/or” is used in this disclosure to mean either “and” or “or” unless indicated otherwise.

[056] It should be understood that the expression “at least one of” includes individually each of the recited objects after the expression and the various combinations of two or more of the recited objects unless otherwise understood from the context and use. The expression “and/or” in connection with three or more recited objects should be understood to have the same meaning unless otherwise understood from the context.

[057] The use of the term “include,” “includes,” “including,” “have,” “has,” “having,” “contain,” “contains,” or “containing,” including grammatical equivalents thereof, should be understood generally as open-ended and non-limiting, for example, not excluding additional unrecited elements or steps, unless otherwise specifically stated or understood from the context.

[058] Where the use of the term “about” is before a quantitative value, the present invention also includes the specific quantitative value itself, unless specifically stated otherwise. As used herein, the term “about” refers to a $\pm 10\%$ variation from the nominal value unless otherwise indicated or inferred from the context.

[059] At various places in the present specification, variable or parameters are disclosed in groups or in ranges. It is specifically intended that the description include each and every individual subcombination of the members of such groups and ranges. For example, an integer in the range of 0 to 40 is specifically intended to individually disclose 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, and 40, and an integer in the range of 1 to 20 is specifically intended to individually disclose 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, and 20.

[060] The use of any and all examples, or exemplary language herein, for example, “such as” or “including,” is intended merely to illustrate better the present invention and does not pose a limitation on the scope of the invention unless claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the present invention.

[061] Deuterium, which may also be referred to herein as ^2H and/or D, is a stable, non-radioactive isotope of ^1H hydrogen and has an atomic weight of 2.014. Hydrogen naturally occurs as a mixture of the isotopes ^1H hydrogen (i.e., protium), deuterium (^2H), and tritium (^3H). The natural abundance of deuterium is about 0.015%. One of ordinary skill in the art recognizes that in all chemical compounds with an H atom, the H atom actually represents a mixture of ^1H hydrogen, deuterium (^2H), and tritium (^3H), where about 0.015% is deuterium. In various embodiments, compounds with a level of deuterium that has been enriched to be greater than its natural abundance of about 0.015% are considered unnatural and, as a result, novel over their non-enriched counterparts. In certain embodiments, compounds with a level of deuterium that has been enriched to be greater than its natural abundance of about 0.015% is a deuterium-enriched compound.

[062] As used herein, “total deuterium content” or “total deuterium abundance” refer to number of deuteriums that are present in a molecule (if the molecule is a salt, the counter ion is also counted, e.g., the HCl or DCl salt of deuterium-enriched (*R*)-pioglitazone). The total deuterium content may be determined, for example, by a ^2H -NMR. The deuterium content at the chiral center of a molecule (e.g., deuterium-enriched pioglitazone) may be determined, for an example, using ^1H -NMR.

[063] As a general matter, compositions specifying a percentage are by weight unless otherwise specified. Further, if a variable is not accompanied by a definition, then the previous definition of the variable controls.

[064] As used herein, “pharmaceutical composition” or “pharmaceutical formulation” refers to the combination of an active agent with a carrier, inert or active, making the composition especially suitable for diagnostic or therapeutic use *in vivo* or *ex vivo*.

[065] “Pharmaceutically acceptable” means approved or approvable by a regulatory agency of the federal or a state government or the corresponding agency in countries other than the United States, or that is listed in the U.S. Pharmacopoeia or other generally recognized pharmacopoeia for use in animals, and more particularly, in humans.

[066] As used herein, “pharmaceutically acceptable excipient” refers to a substance that aids the administration of an active agent to and/or absorption by a subject and can be

included in the compositions of the present invention without causing a significant adverse toxicological effect on the patient. Non-limiting examples of pharmaceutically acceptable excipients include water, NaCl, normal saline solutions, such as a phosphate buffered saline solution, emulsions (e.g., such as an oil/water or water/oil emulsions), lactated Ringer's, 5 normal sucrose, normal glucose, binders, fillers, disintegrants, lubricants, coatings, sweeteners, flavors, salt solutions (such as Ringer's solution), alcohols, oils, gelatins, carbohydrates such as lactose, amylose or starch, fatty acid esters, hydroxymethylcellulose, polyvinyl pyrrolidone, and colors, and the like. Such preparations can be sterilized and, if desired, mixed with auxiliary agents such as lubricants, preservatives, stabilizers, wetting 10 agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, and/or aromatic substances and the like that do not deleteriously react with the compounds of the invention. For examples of excipients, see Martin, Remington's Pharmaceutical Sciences, 15th Ed., Mack Publ. Co., Easton, PA (1975).

[067] As used herein, "solid dosage form" means a pharmaceutical dose(s) in solid form, 15 e.g., tablets, capsules, granules, powders, sachets, reconstitutable powders, dry powder inhalers and chewables.

[068] As used herein, "administering" means oral administration, administration as a suppository, topical contact, intravenous administration, parenteral administration, intraperitoneal administration, intramuscular administration, intralesional administration, 20 intrathecal administration, intracranial administration, intranasal administration or subcutaneous administration, or the implantation of a slow-release device, e.g., a mini-osmotic pump, to a subject. Administration is by any route, including parenteral and transmucosal (e.g., buccal, sublingual, palatal, gingival, nasal, vaginal, rectal, or transdermal). Parenteral administration includes, e.g., intravenous, intramuscular, intra- 25 arterial, intradermal, subcutaneous, intraperitoneal, intraventricular, and intracranial.

Other modes of delivery include, but are not limited to, the use of liposomal formulations, intravenous infusion, transdermal patches, etc. By "co-administer" it is meant that a composition described herein is administered at the same time, just prior to, or just after the administration of one or more additional therapies (e.g., anti-cancer agent, 30 chemotherapeutic, or treatment for a neurodegenerative disease). (*R*)-²H-pioglitazone, or a pharmaceutically acceptable salt thereof, can be administered alone or can be co-administered to the patient. Co-administration is meant to include simultaneous or sequential administration of the compound individually or in combination (more than one


compound or agent). Thus, the preparations can also be combined, when desired, with other active substances (e.g., to reduce metabolic degradation).

[069] The terms “disease,” “disorder,” and “condition” are used interchangeably herein.

[070] As used herein, and unless otherwise specified, the terms “treat,” “treating” and “treatment” contemplate an action that occurs while a subject is suffering from the specified disease, disorder or condition, which reduces the severity of the disease, disorder or condition, or retards or slows the progression of the disease, disorder or condition (e.g., “therapeutic treatment”).


[071] In general, an “effective amount” of a compound refers to an amount sufficient to elicit the desired biological response, e.g., to treat a condition, disease, or disorder described herein. As will be appreciated by those of ordinary skill in this art, the effective amount of a compound of the disclosure may vary depending on such factors as the desired biological endpoint, the pharmacokinetics of the compound, the disease being treated, the mode of administration, and the age, weight, health, and condition of the subject.

[072] “Pioglitazone” as used herein refers to compound (A) below which is a racemic (1:1) mixture of (*R*)- and (*S*)-enantiomers and as a bulk material has approximately the natural abundance of deuterium at each hydrogen atom position. Analogs and/or enantiomers of pioglitazone may be specified herein, for example “deuterium-enriched (*R*)-pioglitazone.”



Deuterium-Enriched (*R*)-Pioglitazone

[073] Deuterium-enriched (*R*)-pioglitazone, as depicted in formula (I), is a MPC inhibitor:



wherein R¹ is H or D.

[074] As depicted above, the deuterium-enriched (*R*)-pioglitazone described herein contains deuterium enrichment at the chiral center of pioglitazone and optionally in other

locations in the compound (e.g., the thiazolidinedione nitrogen). Without wishing to be bound by theory, it is thought that deuterium-enrichment at the chiral center reduces or inhibits the rate at which the two enantiomers of pioglitazone may interconvert.

[075] Methods of chemically synthesizing deuterium-enriched (*R*)-pioglitazone and preparing the crystalline salt forms described herein are provided in Examples 1, 2, and 3.

[076] In certain embodiments, the abundance of deuterium at the chiral center is about 30% to about 100%, about 40% to about 100%, about 50% to about 100%, about 60% to about 100%, about 70% to about 100%, about 75% to about 100%, about 80% to about 100%, about 85% to about 100%, about 90% to about 100%, about 95% to about 100%, about 30% to about 95%, about 30% to about 90%, about 30% to about 85%, about 30% to about 80%, about 30% to about 75%, about 30% to about 70%, about 30% to about 60%, about 30% to about 50%, about 30% to about 40%, about 40% to about 95%, about 40% to about 90%, about 40% to about 85%, about 40% to about 80%, about 40% to about 75%, about 40% to about 70%, about 40% to about 60%, about 40% to about 50%, about 50% to about 95%, about 50% to about 90%, about 50% to about 85%, about 50% to about 80%, about 50% to about 75%, about 50% to about 70%, about 50% to about 60%, about 60% to about 95%, about 60% to about 90%, about 60% to about 85%, about 60% to about 80%, about 60% to about 75%, about 60% to about 70%, about 70% to about 95%, about 70% to about 90%, about 70% to about 85%, about 70% to about 80%, about 70% to about 75%, about 75% to about 95%, about 75% to about 90%, about 75% to about 85%, about 75% to about 80%, about 80% to about 95%, about 80% to about 90%, about 80% to about 85%, about 85% to about 95%, about 85% to about 90%, or about 90% to about 95%.

[077] In certain embodiments, the abundance of deuterium at the chiral center is at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%.

[078] In certain embodiments, the abundance of deuterium at the chiral center is about 30%, about 31%, about 32%, about 33%, about 34%, about 35%, about 36%, about 37%, about 38%, about 39%, about 40%, about 41%, about 42%, about 43%, about 44%, about 45%, about 46%, about 47%, about 48%, about 49%, about 50%, about 51%, about 52%, about 53%, about 54%, about 55%, about 56%, about 57%, about 58%, about 59%, about 60%, about 61%, about 62%, about 63%, about 64%, about 65%, about 66%, about 67%, about 68%, about 69%, about 70%, about 71%, about 72%, about 73%, about 74%, about 75%, about 76%, about 77%, about 78%, about 79%, about 80%, about 81%, about 82%,

about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100%.

[079] In certain embodiments, the abundance of deuterium at R^1 is 0.015% to about 100%, about 10% to about 100%, about 20% to about 100%, about 30% to about 100%, about 40% to about 100%, about 50% to about 100%, about 60% to about 100%, about 70% to about 100%, about 75% to about 100%, about 80% to about 100%, about 85% to about 100%, about 90% to about 100%, about 95% to about 100%, 0.015% to about 95%, 0.015% to about 90%, 0.015% to about 85%, 0.015% to about 80%, 0.015% to about 75%, 0.015% to about 70%, 0.015% to about 60%, 0.015% to about 50%, 0.015% to about 40%, 0.015% to about 30%, 0.015% to about 20%, 0.015% to about 10%, about 10% to about 95%, about 10% to about 90%, about 10% to about 85%, about 10% to about 80%, about 10% to about 75%, about 10% to about 70%, about 10% to about 60%, about 10% to about 50%, about 10% to about 40%, about 10% to about 30%, about 10% to about 20%, about 20% to about 95%, about 20% to about 90%, about 20% to about 85%, about 20% to about 80%, about 20% to about 75%, about 20% to about 70%, about 20% to about 60%, about 20% to about 50%, about 20% to about 40%, about 20% to about 30%, about 30% to about 95%, about 30% to about 90%, about 30% to about 85%, about 30% to about 80%, about 30% to about 75%, about 30% to about 70%, about 30% to about 60%, about 30% to about 50%, about 30% to about 40%, about 40% to about 95%, about 40% to about 90%, about 40% to about 85%, about 40% to about 80%, about 40% to about 75%, about 40% to about 70%, about 40% to about 60%, about 40% to about 50%, about 50% to about 95%, about 50% to about 90%, about 50% to about 85%, about 50% to about 80%, about 50% to about 75%, about 50% to about 70%, about 50% to about 60%, about 60% to about 95%, about 60% to about 90%, about 60% to about 85%, about 60% to about 80%, about 60% to about 75%, about 60% to about 70%, about 70% to about 95%, about 70% to about 90%, about 70% to about 85%, about 70% to about 80%, about 70% to about 75%, about 75% to about 95%, about 75% to about 90%, about 75% to about 85%, about 75% to about 80%, about 80% to about 95%, about 80% to about 90%, about 80% to about 85%, about 85% to about 95%, about 85% to about 90%, or about 90% to about 95%.

[080] In certain embodiments, the abundance of deuterium at R^1 is at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 95%, at least 96%, at least 97%, at least 98%, or at least

99%.

[081] In certain embodiments, the abundance of deuterium at R¹ is about 1%, about 2%, about 3%, about 4%, about 5%, about 6%, about 7%, about 8%, about 9%, about 10%, about 11%, about 12%, about 13%, about 14%, about 15%, about 16%, about 17%, about 18%, about 19%, about 20%, about 21%, about 22%, about 23%, about 24%, about 25%, about 26%, about 27%, about 28%, about 29%, about 30%, about 31%, about 32%, about 33%, about 34%, about 35%, about 36%, about 37%, about 38%, about 39%, about 40%, about 41%, about 42%, about 43%, about 44%, about 45%, about 46%, about 47%, about 48%, about 49%, about 50%, about 51%, about 52%, about 53%, about 54%, about 55%, about 56%, about 57%, about 58%, about 59%, about 60%, about 61%, about 62%, about 63%, about 64%, about 65%, about 66%, about 67%, about 68%, about 69%, about 70%, about 71%, about 72%, about 73%, about 74%, about 75%, about 76%, about 77%, about 78%, about 79%, about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100%.

[082] In certain embodiments, the total deuterium abundance in the compound of formula (I) is about 0.3 to about 4, about 0.5 to about 4, about 1 to about 4, about 1.2 to about 4, about 1.3 to about 4, about 1.4 to about 4, about 1.6 to about 4, about 1.8 to about 4, about 2 to about 4, about 2.2 to about 4, about 2.4 to about 4, about 2.6 to about 4, about 2.8 to about 4, about 3 to about 4, about 3.2 to about 4, about 3.4 to about 4, about 3.6 to about 4, about 3.8 to about 4, about 0.3 to about 3.8, about 0.3 to about 3.6, about 0.3 to about 3.4, about 0.3 to about 3.2, about 0.3 to about 3, about 0.3 to about 2.8, about 0.3 to about 2.6, about 0.3 to about 2.4, about 0.3 to about 2.2, about 0.3 to about 2, about 0.3 to about 1.8, about 0.3 to about 1.6, about 0.3 to about 1.4, about 0.3 to about 1.2, about 0.3 to about 1, about 0.3 to about 0.5, about 0.5 to about 3.8, about 0.5 to about 3.6, about 0.5 to about 3.4, about 0.5 to about 3.2, about 0.5 to about 3, about 0.5 to about 2.8, about 0.5 to about 2.6, about 0.5 to about 2.4, about 0.5 to about 2.2, about 0.5 to about 2, about 0.5 to about 1.8, about 0.5 to about 1.6, about 0.5 to about 1.4, about 0.5 to about 1.2, about 0.5 to about 1, about 1 to about 3.8, about 1 to about 3.6, about 1 to about 3.4, about 1 to about 3.2, about 1 to about 3, about 1 to about 2.8, about 1 to about 2.6, about 1 to about 2.4, about 1 to about 2.2, about 1 to about 2, about 1 to about 1.8, about 1 to about 1.6, about 1 to about 1.4, about 1 to about 1.2, about 1.2 to about 3.8, about 1.2 to about 3.6, about 1.2 to about 3.4, about 1.2 to about 3.2, about 1.2 to about 3, about 1.2 to about 2.8, about 1.2 to about 2.6, about 1.2 to about 2.4, about 1.2 to about

2.2, about 1.2 to about 2, about 1.2 to about 1.8, about 1.2 to about 1.6, about 1.2 to about 1.4, about 1.2 to about 1.3, about 1.3 to about 3.8, about 1.3 to about 3.6, about 1.3 to about 3.4, about 1.3 to about 3.2, about 1.3 to about 3, about 1.3 to about 2.8, about 1.3 to about 2.6, about 1.3 to about 2.4, about 1.3 to about 2.2, about 1.3 to about 2, about 1.3 to about 1.8, about 1.3 to about 1.6, about 1.3 to about 1.4, about 1.4 to about 3.8, about 1.4 to about 3.6, about 1.4 to about 3.4, about 1.4 to about 3.2, about 1.4 to about 3, about 1.4 to about 2.8, about 1.4 to about 2.6, about 1.4 to about 2.4, about 1.4 to about 2.2, about 1.4 to about 2, about 1.4 to about 1.8, about 1.4 to about 1.6, about 1.6 to about 3.8, about 1.6 to about 3.6, about 1.6 to about 3.4, about 1.6 to about 3.2, about 1.6 to about 3, about 1.6 to about 2.8, about 1.6 to about 2.6, about 1.6 to about 2.4, about 1.6 to about 2.2, about 1.6 to about 2, about 1.6 to about 1.8, about 1.8 to about 3.8, about 1.8 to about 3.6, about 1.8 to about 3.4, about 1.8 to about 3.2, about 1.8 to about 3, about 1.8 to about 2.8, about 1.8 to about 2.6, about 1.8 to about 2.4, about 1.8 to about 2.2, about 1.8 to about 2, about 2 to about 3.8, about 2 to about 3.6, about 2 to about 3.4, about 2 to about 3.2, about 2 to about 3, about 2 to about 2.8, about 2 to about 2.6, about 2 to about 2.4, about 2 to about 2.2, about 2.2 to about 3.8, about 2.2 to about 3.6, about 2.2 to about 3.4, about 2.2 to about 3.2, about 2.2 to about 3, about 2.2 to about 2.8, about 2.2 to about 2.6, about 2.2 to about 2.4, about 2.4 to about 3.8, about 2.4 to about 3.6, about 2.4 to about 3.4, about 2.4 to about 3.2, about 2.4 to about 3, about 2.4 to about 2.8, about 2.4 to about 2.6, about 2.6 to about 3.8, about 2.6 to about 3.6, about 2.6 to about 3.4, about 2.6 to about 3.2, about 2.6 to about 3, about 2.6 to about 2.8, about 2.8 to about 3.8, about 2.8 to about 3.6, about 2.8 to about 3.4, about 2.8 to about 3.2, about 2.8 to about 3, about 3 to about 3.8, about 3 to about 3.6, about 3 to about 3.4, about 3 to about 3.2, about 3.2 to about 3.8, about 3.2 to about 3.6, about 3.2 to about 3.4, about 3.4 to about 3.8, about 3.4 to about 3.6, or about 3.6 to about 3.8. In certain embodiments, the total deuterium abundance in the compound of formula (I) is about 1.3 to about 4. In certain embodiments, the total deuterium abundance in the compound of formula (I) is about 1.8 to about 3.

[083] In certain embodiments, the deuterium-enriched (*R*)-pioglitazone described herein has an enantiomeric excess of about 70% to about 100%, about 75% to about 100%, about 80% to about 100%, about 85% to about 100%, about 90% to about 100%, about 95% to about 100%, about 70% to about 95%, about 70% to about 90%, about 70% to about 85%, about 70% to about 80%, about 70% to about 75%, about 75% to about 95%, about 75% to about 90%, about 75% to about 85%, about 75% to about 80%, about 80% to about 95%, about 80% to about 90%, about 80% to about 85%, about 85% to about 95%, about 85% to

about 90%, or about 90% to about 95%.

[084] In certain embodiments, the deuterium-enriched (*R*)-pioglitazone described herein has an enantiomeric excess of at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%.


[085] In certain embodiments, the deuterium-enriched (*R*)-pioglitazone described herein has an enantiomeric excess of about 70%, about 71%, about 72%, about 73%, about 74%, about 75%, about 76%, about 77%, about 78%, about 79%, about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100%. In certain embodiments, the deuterium-enriched (*R*)-pioglitazone described herein is enantiopure.

[086] In various embodiments, provided herein is the free-base form of deuterium-enriched (*R*)-pioglitazone.

[087] In various embodiments, provided herein is a pharmaceutically acceptable salt of deuterium-enriched (*R*)-pioglitazone. In certain embodiments, the pharmaceutically acceptable salt of deuterium-enriched (*R*)-pioglitazone is a deuterium chloride salt. In certain embodiments, the pharmaceutically acceptable salt of deuterium-enriched (*R*)-pioglitazone is a hydrochloride salt.

Deuterium Chloride (DCI) Salt of Deuterium-Enriched (*R*)-Pioglitazone


[088] In one aspect, the invention provides a deuterium chloride salt of the deuterium-enriched compound of formula (I):



wherein R^1 is H or D, provided that the total deuterium abundance in the compound of formula (I) is about 1.3 to about 4.

[089] In certain embodiments, the deuterium chloride salt is a crystalline deuterium chloride salt. In certain embodiments, the crystalline deuterium chloride salt is an anhydrous crystalline deuterium chloride salt.

[090] In another aspect, the invention provides a crystalline deuterium chloride salt of the deuterium-enriched compound of formula (I)



wherein R¹ is H or D, provided that the total deuterium abundance in the compound of formula (I) is about 1.3 to about 4, and

wherein the crystalline deuterium chloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): 15.8° ± 0.2°, 22.8° ± 0.2°, and 26.0° ± 0.2°.

[091] In certain embodiments, the crystalline deuterium chloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): 20.0° ± 0.2°, 20.8° ± 0.2°, and 22.8° ± 0.2°. In certain embodiments, the crystalline deuterium chloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): 15.8° ± 0.2°, 20.0° ± 0.2°, 20.8° ± 0.2°, 22.8° ± 0.2°, and 26.0° ± 0.2°. In certain embodiments, the crystalline deuterium chloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): 8.6° ± 0.2°, 8.8° ± 0.2°, 12.8° ± 0.2°, 12.9° ± 0.2°, 15.8° ± 0.2°, 18.8° ± 0.2°, 18.9° ± 0.2°, 19.7° ± 0.2°, 20.0° ± 0.2°, 20.8° ± 0.2°, 22.8° ± 0.2°, 26.0° ± 0.2°, and 31.3° ± 0.2°.

[092] In certain embodiments, the crystalline deuterium chloride salt is characterized by the X-ray powder diffraction pattern expressed in terms of diffraction angle 2θ and optionally relative intensity (expressed as a percentage with respect to the most intense peak) as set forth in Table 1.

Table 1 – X-ray Powder Diffraction Data (XRPD) of the Crystalline DCI Salt

2θ (°)	d value (Å)	Intensity (%)
8.6	10.27	18
8.8	10.07	17
12.8	6.93	22
12.9	6.87	26
15.9	5.59	4
18.7	4.73	18

2θ (°)	d value (Å)	Intensity (%)
18.9	4.70	16
19.7	4.51	18
20.0	4.45	34
20.8	4.26	56
21.2	4.19	13
22.1	4.02	8
22.8	3.89	100
23.7	3.75	3
25.6	3.47	5
26.0	3.43	20
26.7	3.34	9
27.3	3.26	6
28.1	3.18	12
29.8	3.00	12
31.3	2.86	21
32.1	2.79	10
32.3	2.77	5
33.4	2.68	3
33.6	2.66	3
34.4	2.60	4
36.0	2.50	3

[093] In certain embodiments, the crystalline deuterium chloride salt is characterized by an X-ray powder diffraction pattern substantially the same as shown in FIG. 1A.

[094] In certain embodiments, the crystalline deuterium chloride salt exists in a monoclinic crystal system and has a $P2_1/c$ space group. In certain embodiments, the crystalline deuterium chloride salt is characterized by the crystallographic unit cell parameters as set forth in Table 2.

Table 2 – Unit Cell Parameters of the Crystalline Form of Compound of formula (V)

a [Å]	10.1319 (9)
b [Å]	9.4067 (9)
c [Å]	10.3231 (6)

β [°]	94.397 (3)
$V[\text{\AA}^3]$	980.98 (14)
Z	2
$D_c [\text{g}/\text{cm}^3]$	1.330

[095] The crystalline deuterium chloride salt may also be characterized according to the temperature of melting point onset. In certain embodiments, the crystalline deuterium chloride salt has a melting point onset as determined by differential scanning calorimetry at about 190 °C to about 200 °C. In certain embodiments, the crystalline deuterium chloride salt has a melting point onset as determined by differential scanning calorimetry at about 191 °C. In certain embodiments, the crystalline deuterium chloride salt exhibits a melting endotherm with a peak at about 200 °C to about 210 °C. In certain embodiments, the crystalline deuterium chloride salt exhibits a melting endotherm with a peak at about 204 °C. In certain embodiments, the crystalline deuterium chloride salt has a differential scanning calorimetry curve substantially the same as shown in FIG. 3.


[096] In certain embodiments, the crystalline deuterium chloride salt has a chemical purity of about 70% to about 100%, about 75% to about 100%, about 80% to about 100%, about 85% to about 100%, about 90% to about 100%, about 95% to about 100%, about 70% to about 95%, about 70% to about 90%, about 70% to about 85%, about 70% to about 80%, about 70% to about 75%, about 75% to about 95%, about 75% to about 90%, about 75% to about 85%, about 75% to about 80%, about 80% to about 95%, about 80% to about 90%, about 80% to about 85%, about 85% to about 95%, about 85% to about 90%, or about 90% to about 95%.

[097] In certain embodiments, the crystalline deuterium chloride salt has a chemical purity of at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, at least 99.1%, at least 99.2%, at least 99.3%, at least 99.4%, at least 99.5%, at least 99.6%, at least 99.7%, at least 99.8%, or at least 99.9%.

[098] In certain embodiments, the crystalline deuterium chloride salt has a chemical purity of about 70%, about 71%, about 72%, about 73%, about 74%, about 75%, about 76%, about 77%, about 78%, about 79%, about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about

99%, about 99.1%, about 99.2%, about 99.3%, about 99.4%, about 99.5%, about 99.6%, about 99.7%, about 99.8%, about 99.9%, or about 100%.

[099] In various embodiments, the invention provides a pharmaceutical material comprising particles of a crystalline deuterium chloride salt of a deuterium-enriched compound of formula (I)



wherein R^1 is H or D, provided that the total deuterium abundance in the compound of formula (I) is about 1.3 to about 4, and

wherein the particles in the composition have a crystal shape selected from hexagonal, rod, and combinations thereof.

[100] In certain embodiments, the particles have a particle size distribution which is defined by a $d(0.1)$ of about 10 μm to 200 μm , about 20 μm to 200 μm , about 40 μm to 200 μm , about 60 μm to 200 μm , about 80 μm to 200 μm , about 100 μm to 200 μm , about 120 μm to 200 μm , about 140 μm to 200 μm , about 160 μm to 200 μm , about 180 μm to 200 μm , about 10 μm to 180 μm , about 10 μm to 160 μm , about 10 μm to 140 μm , about 10 μm to 120 μm , about 10 μm to 100 μm , about 10 μm to 80 μm , about 10 μm to 60 μm , about 10 μm to 40 μm , about 10 μm to 20 μm , about 20 μm to 180 μm , about 20 μm to 160 μm , about 20 μm to 140 μm , about 20 μm to 120 μm , about 20 μm to 100 μm , about 20 μm to 80 μm , about 20 μm to 60 μm , about 20 μm to 40 μm , about 40 μm to 180 μm , about 40 μm to 160 μm , about 40 μm to 140 μm , about 40 μm to 120 μm , about 40 μm to 100 μm , about 40 μm to 80 μm , about 40 μm to 60 μm , about 60 μm to 180 μm , about 60 μm to 160 μm , about 60 μm to 140 μm , about 60 μm to 120 μm , about 60 μm to 100 μm , about 60 μm to 80 μm , about 80 μm to 180 μm , about 80 μm to 160 μm , about 80 μm to 140 μm , about 80 μm to 120 μm , about 80 μm to 100 μm , about 100 μm to 180 μm , about 100 μm to 160 μm , about 100 μm to 140 μm , about 100 μm to 120 μm , about 120 μm to 180 μm , about 120 μm to 160 μm , about 120 μm to 140 μm , about 140 μm to 180 μm , about 140 μm to 160 μm , or about 160 μm to 180 μm . In certain embodiments, the particles have a particle size distribution which is defined by a $d(0.1)$ of about 10 μm to about 200 μm .

[101] In certain embodiments, the particles have a particle size distribution which is defined by a $d(0.5)$ of about 10 μm to about 400 μm , about 50 μm to about 400 μm , about


[illegible]

about 275 μm , about 200 μm to about 250 μm , about 200 μm to about 225 μm , about 225 μm to about 375 μm , about 225 μm to about 350 μm , about 225 μm to about 325 μm , about 225 μm to about 300 μm , about 225 μm to about 275 μm , about 225 μm to about 250 μm , about 250 μm to about 375 μm , about 250 μm to about 350 μm , about 250 μm to about 325 μm , about 250 μm to about 300 μm , about 250 μm to about 275 μm , about 275 μm to about 375 μm , about 275 μm to about 350 μm , about 275 μm to about 325 μm , about 275 μm to about 300 μm , about 300 μm to about 375 μm , about 300 μm to about 350 μm , about 300 μm to about 325 μm , about 325 μm to about 375 μm , about 325 μm to about 350 μm , or about 350 μm to about 375 μm . In certain embodiments, the particles have a particle size distribution which is defined by a $d(0.5)$ about 10 μm to about 400 μm .

[102] In certain embodiments, the particles have a particle size distribution which is defined by a d(0.9) of about 10 μm to about 800 μm, about 50 μm to about 800 μm, about 100 μm to about 800 μm, about 150 μm to about 800 μm, about 200 μm to about 800 μm, about 250 μm to about 800 μm, about 300 μm to about 800 μm, about 350 μm to about 800 μm, about 400 μm to about 800 μm, about 450 μm to about 800 μm, about 500 μm to about 800 μm, about 550 μm to about 800 μm, about 600 μm to about 800 μm, about 650 μm to about 800 μm, about 700 μm to about 800 μm, about 750 μm to about 800 μm, about 150 μm to about 750 μm, about 150 μm to about 700 μm, about 150 μm to about 650 μm, about 150 μm to about 600 μm, about 150 μm to about 550 μm, about 150 μm to about 500 μm, about 150 μm to about 450 μm, about 150 μm to about 400 μm, about 150 μm to about 350 μm, about 150 μm to about 300 μm, about 150 μm to about 250 μm, about 150 μm to about 200 μm, about 200 μm to about 750 μm, about 200 μm to about 700 μm, about 200 μm to about 650 μm, about 200 μm to about 600 μm, about 200 μm to about 550 μm, about 200 μm to about 500 μm, about 200 μm to about 450 μm, about 200 μm to about 400 μm, about 200 μm to about 350 μm, about 200 μm to about 300 μm, about 200 μm to about 250 μm, about 250 μm to about 750 μm, about 250 μm to about 700 μm, about 250 μm to about 650 μm, about 250 μm to about 600 μm, about 250 μm to about 550 μm, about 250 μm to about 500 μm, about 250 μm to about 450 μm, about 250 μm to about 400 μm, about 250 μm to about 350 μm, about 250 μm to about 300 μm, about 300 μm to about 750 μm, about 300 μm to about 700 μm, about 300 μm to about 650 μm, about 300 μm to about 600 μm, about 300 μm to about 550 μm, about 300 μm to about 500 μm, about 300 μm to about 450 μm, about 300 μm to about 400 μm, about 300 μm to about 350 μm, about 350 μm to about 750 μm, about 350 μm to about 700 μm, about 350 μm to about 650 μm, about 350 μm to about 600 μm, about 350 μm to about

550 μm , about 350 μm to about 500 μm , about 350 μm to about 450 μm , about 350 μm to about 400 μm , about 400 μm to about 750 μm , about 400 μm to about 700 μm , about 400 μm to about 650 μm , about 400 μm to about 600 μm , about 400 μm to about 550 μm , about 400 μm to about 500 μm , about 400 μm to about 450 μm , about 450 μm to about 750 μm , about 450 μm to about 700 μm , about 450 μm to about 650 μm , about 450 μm to about 600 μm , about 450 μm to about 550 μm , about 450 μm to about 500 μm , about 500 μm to about 750 μm , about 500 μm to about 700 μm , about 500 μm to about 650 μm , about 500 μm to about 600 μm , about 500 μm to about 550 μm , about 550 μm to about 750 μm , about 550 μm to about 700 μm , about 550 μm to about 650 μm , about 550 μm to about 600 μm , about 600 μm to about 750 μm , about 600 μm to about 700 μm , about 600 μm to about 650 μm , about 650 μm to about 750 μm , about 650 μm to about 700 μm , or about 700 μm to about 750 μm . In certain embodiments, the particles have a particle size distribution which is defined by a $d(0.9)$ of about 10 μm to about 800 μm .


[103] In various embodiments, the invention provides a pharmaceutical material comprising particles of a crystalline deuterium chloride salt of a deuterium-enriched compound of formula (I)



wherein R^1 is H or D, provided that the total deuterium abundance in the compound of formula (I) is about 1.3 to about 4, and wherein the particles have a particle size distribution which is defined by a $d(0.9)$ of about 10 μm to about 800 μm , preferably below 500 μm .

Hydrochloride (HCl) Salt of Deuterium-Enriched Pioglitazone


[104] In one aspect, the invention provides a crystalline hydrochloride salt of the deuterium-enriched compound of formula (I-A):



[105] In certain embodiments, the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ):

$15.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, and $26.0^\circ \pm 0.2^\circ$. In certain embodiments, the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): $20.0^\circ \pm 0.2^\circ$, $20.8^\circ \pm 0.2^\circ$, and $22.8^\circ \pm 0.2^\circ$.

[106] In various embodiments, the invention provides a crystalline hydrochloride salt of the compound of formula (I-A)



wherein the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): $15.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, and $26.0^\circ \pm 0.2^\circ$.

[107] In certain embodiments, the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ):

$15.8^\circ \pm 0.2^\circ$, $20.0^\circ \pm 0.2^\circ$, $20.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, and $26.0^\circ \pm 0.2^\circ$. In certain embodiments, the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): $8.6^\circ \pm 0.2^\circ$, $8.8^\circ \pm 0.2^\circ$, $12.8^\circ \pm 0.2^\circ$, $12.9^\circ \pm 0.2^\circ$, $15.9^\circ \pm 0.2^\circ$, $18.8^\circ \pm 0.2^\circ$, $19.7^\circ \pm 0.2^\circ$, $20.0^\circ \pm 0.2^\circ$, $20.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, $26.0^\circ \pm 0.2^\circ$, $28.1^\circ \pm 0.2^\circ$, and $31.3^\circ \pm 0.2^\circ$.

[108] In certain embodiments, the crystalline hydrochloride salt is characterized by the X-ray powder diffraction pattern expressed in terms of diffraction angle 2θ and optionally relative intensity (expressed as a percentage with respect to the most intense peak) as set forth in Table 3.

Table 3 – X-ray Powder Diffraction Data of the Crystalline Hydrochloride Salt

2θ (°)	d value (Å)	Intensity (%)
8.6	10.27	32
8.8	10.08	31
12.8	6.93	30
12.9	6.87	27
15.9	5.59	6
18.7	4.73	25
18.9	4.70	12
19.7	4.51	29
20.0	4.45	80
20.8	4.26	50
21.2	4.19	18
22.1	4.02	14
22.8	3.89	100
23.7	3.75	4
25.6	3.47	6
26.0	3.43	36
26.7	3.34	9
27.3	3.26	15
28.1	3.17	30
29.8	3.00	9
31.3	2.86	19
32.1	2.79	13
32.3	2.77	7
33.4	2.68	3
33.6	2.66	3
34.4	2.61	5
36.0	2.50	5

[109] In certain embodiments, the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern substantially the same as shown in FIG. 6.

[110] The crystalline hydrochloride salt may also be characterized according to the temperature of melting point onset. In certain embodiments, the crystalline hydrochloride salt has a melting point onset as determined by differential scanning calorimetry at about 190 °C to about 210 °C. In certain embodiments, the crystalline hydrochloride salt has a melting point onset as determined by differential scanning calorimetry at about 190 °C. In certain embodiments, the crystalline deuterium chloride salt exhibits a melting endotherm with a peak at about 195 °C to about 205 °C. In certain embodiments, the crystalline deuterium chloride salt exhibits a melting endotherm with a peak at about 200 °C. In certain embodiments, the crystalline hydrochloride salt has a differential scanning calorimetry curve substantially the same as shown in FIG. 8.


[111] In certain embodiments, the crystalline hydrochloride salt is an anhydrous crystalline hydrochloride salt.

[112] In certain embodiments, the crystalline hydrochloride salt has a chemical purity of about 70% to about 100%, about 75% to about 100%, about 80% to about 100%, about 85% to about 100%, about 90% to about 100%, about 95% to about 100%, about 70% to about 95%, about 70% to about 90%, about 70% to about 85%, about 70% to about 80%, about 70% to about 75%, about 75% to about 95%, about 75% to about 90%, about 75% to about 85%, about 75% to about 80%, about 80% to about 95%, about 80% to about 90%, about 80% to about 85%, about 85% to about 95%, about 85% to about 90%, or about 90% to about 95%.

[113] In certain embodiments, the crystalline hydrochloride salt has a chemical purity of at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, at least 99.1%, at least 99.2%, at least 99.3%, at least 99.4%, at least 99.5%, at least 99.6%, at least 99.7%, at least 99.8%, or at least 99.9%.

[114] In certain embodiments, the crystalline hydrochloride salt has a chemical purity of about 70%, about 71%, about 72%, about 73%, about 74%, about 75%, about 76%, about 77%, about 78%, about 79%, about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, about 99.1%, about 99.2%, about 99.3%, about 99.4%, about 99.5%, about 99.6%, about 99.7%, about 99.8%, about 99.9%, or about 100%.

[115] In various embodiments, the invention provides a pharmaceutical material comprising particles of a crystalline hydrochloride salt of the compound of formula (I-A)



wherein the particles in the composition have a needle-like crystal shape.

[116] In certain embodiments, the particles have a particle size distribution which is defined by a $d(0.1)$ of about 10 μm to 200 μm , about 20 μm to 200 μm , about 40 μm to 200 μm , about 60 μm to 200 μm , about 80 μm to 200 μm , about 100 μm to 200 μm , about 120 μm to 200 μm , about 140 μm to 200 μm , about 160 μm to 200 μm , about 180 μm to 200 μm , about 10 μm to 180 μm , about 10 μm to 160 μm , about 10 μm to 140 μm , about 10 μm to 120 μm , about 10 μm to 100 μm , about 10 μm to 80 μm , about 10 μm to 60 μm , about 10 μm to 40 μm , about 10 μm to 20 μm , about 20 μm to 180 μm , about 20 μm to 160 μm , about 20 μm to 140 μm , about 20 μm to 120 μm , about 20 μm to 100 μm , about 20 μm to 80 μm , about 20 μm to 60 μm , about 20 μm to 40 μm , about 40 μm to 180 μm , about 40 μm to 160 μm , about 40 μm to 140 μm , about 40 μm to 120 μm , about 40 μm to 100 μm , about 40 μm to 80 μm , about 40 μm to 60 μm , about 60 μm to 180 μm , about 60 μm to 160 μm , about 60 μm to 140 μm , about 60 μm to 120 μm , about 60 μm to 100 μm , about 60 μm to 80 μm , about 80 μm to 180 μm , about 80 μm to 160 μm , about 80 μm to 140 μm , about 80 μm to 120 μm , about 80 μm to 100 μm , about 100 μm to 180 μm , about 100 μm to 160 μm , about 100 μm to 140 μm , about 100 μm to 120 μm , about 120 μm to 180 μm , about 120 μm to 160 μm , about 120 μm to 140 μm , about 140 μm to 180 μm , about 140 μm to 160 μm , or about 160 μm to 180 μm . In certain embodiments, the particles have a particle size distribution which is defined by a $d(0.1)$ of about 10 μm to about 200 μm .

[117] In certain embodiments, the particles have a particle size distribution which is defined by a $d(0.5)$ of about 10 μm to about 400 μm , about 50 μm to about 400 μm , about 75 μm to about 400 μm , about 100 μm to about 400 μm , about 125 μm to about 400 μm , about 150 μm to about 400 μm , about 175 μm to about 400 μm , about 200 μm to about 400 μm , about 225 μm to about 400 μm , about 250 μm to about 400 μm , about 275 μm to about 400 μm , about 300 μm to about 400 μm , about 325 μm to about 400 μm , about 350 μm to about 400 μm , about 375 μm to about 400 μm , about 10 μm to about 375 μm , about 10 μm to about 350 μm , about 10 μm to about 325 μm , about 10 μm to about 300 μm , about 10 μm to about 275 μm , about 10 μm to about 250 μm , about 10 μm to about 225 μm , about 10 μm to about 200 μm , about 50 μm to about 375 μm , about 50 μm to about


350 μm, about 50 μm to about 325 μm, about 50 μm to about 300 μm, about 50 μm to
about 275 μm, about 50 μm to about 250 μm, about 50 μm to about 225 μm, about 50 μm
to about 200 μm, about 50 μm to about 175 μm, about 50 μm to about 150 μm, about 50
μm to about 125 μm, about 50 μm to about 100 μm, about 50 μm to about 75 μm, about
5 75 μm to about 375 μm, about 75 μm to about 350 μm, about 75 μm to about 325 μm,
about 75 μm to about 300 μm, about 75 μm to about 275 μm, about 75 μm to about 250
μm, about 75 μm to about 225 μm, about 75 μm to about 375 μm, about 75 μm to about
350 μm, about 75 μm to about 325 μm, about 75 μm to about 300 μm, about 75 μm to
about 275 μm, about 75 μm to about 250 μm, about 75 μm to about 225 μm, about 75 μm
10 to about 200 μm, about 75 μm to about 175 μm, about 75 μm to about 150 μm, about 75
μm to about 125 μm, about 75 μm to about 100 μm, about 100 μm to about 375 μm, about
100 μm to about 350 μm, about 100 μm to about 325 μm, about 100 μm to about 300 μm,
about 100 μm to about 275 μm, about 100 μm to about 250 μm, about 100 μm to about
225 μm, about 100 μm to about 200 μm, about 100 μm to about 175 μm, about 100 μm to
15 about 150 μm, about 100 μm to about 125 μm, about 125 μm to about 375 μm, about 125
μm to about 350 μm, about 125 μm to about 325 μm, about 125 μm to about 300 μm,
about 125 μm to about 275 μm, about 125 μm to about 250 μm, about 125 μm to about
225 μm, about 125 μm to about 200 μm, about 125 μm to about 175 μm, about 125 μm to
about 150 μm, about 150 μm to about 375 μm, about 150 μm to about 350 μm, about 150
20 μm to about 325 μm, about 150 μm to about 300 μm, about 150 μm to about 275 μm,
about 150 μm to about 250 μm, about 150 μm to about 225 μm, about 150 μm to about
200 μm, about 150 μm to about 175 μm, about 175 μm to about 375 μm, about 175 μm to
about 350 μm, about 175 μm to about 325 μm, about 175 μm to about 300 μm, about 175
μm to about 275 μm, about 175 μm to about 250 μm, about 175 μm to about 225 μm,
25 about 175 μm to about 200 μm, about 200 μm to about 375 μm, about 200 μm to about
350 μm, about 200 μm to about 325 μm, about 200 μm to about 300 μm, about 200 μm to
about 275 μm, about 200 μm to about 250 μm, about 200 μm to about 225 μm, about 225
μm to about 375 μm, about 225 μm to about 350 μm, about 225 μm to about 325 μm,
about 225 μm to about 300 μm, about 225 μm to about 275 μm, about 225 μm to about
30 250 μm, about 250 μm to about 375 μm, about 250 μm to about 350 μm, about 250 μm to
about 325 μm, about 250 μm to about 300 μm, about 250 μm to about 275 μm, about 275
μm to about 375 μm, about 275 μm to about 350 μm, about 275 μm to about 325 μm,
about 275 μm to about 300 μm, about 300 μm to about 375 μm, about 300 μm to about
350 μm, about 300 μm to about 325 μm, about 325 μm to about 375 μm, about 325 μm to

about 350 μm , or about 350 μm to about 375 μm . In certain embodiments, the particles have a particle size distribution which is defined by a d(0.5) of from about 10 μm to about 400 μm .

[118] In certain embodiments, the particles have a particle size distribution which is defined by a d(0.9) of about 10 μm to about 800 μm, about 50 μm to about 800 μm, about 100 μm to about 800 μm, about 150 μm to about 800 μm, about 200 μm to about 800 μm, about 250 μm to about 800 μm, about 300 μm to about 800 μm, about 350 μm to about 800 μm, about 400 μm to about 800 μm, about 450 μm to about 800 μm, about 500 μm to about 800 μm, about 550 μm to about 800 μm, about 600 μm to about 800 μm, about 650 μm to about 800 μm, about 700 μm to about 800 μm, about 750 μm to about 800 μm, about 150 μm to about 750 μm, about 150 μm to about 700 μm, about 150 μm to about 650 μm, about 150 μm to about 600 μm, about 150 μm to about 550 μm, about 150 μm to about 500 μm, about 150 μm to about 450 μm, about 150 μm to about 400 μm, about 150 μm to about 350 μm, about 150 μm to about 300 μm, about 150 μm to about 250 μm, about 150 μm to about 200 μm, about 200 μm to about 750 μm, about 200 μm to about 700 μm, about 200 μm to about 650 μm, about 200 μm to about 600 μm, about 200 μm to about 550 μm, about 200 μm to about 500 μm, about 200 μm to about 450 μm, about 200 μm to about 400 μm, about 200 μm to about 350 μm, about 200 μm to about 300 μm, about 200 μm to about 250 μm, about 250 μm to about 750 μm, about 250 μm to about 700 μm, about 250 μm to about 650 μm, about 250 μm to about 600 μm, about 250 μm to about 550 μm, about 250 μm to about 500 μm, about 250 μm to about 450 μm, about 250 μm to about 400 μm, about 250 μm to about 350 μm, about 250 μm to about 300 μm, about 300 μm to about 750 μm, about 300 μm to about 700 μm, about 300 μm to about 650 μm, about 300 μm to about 600 μm, about 300 μm to about 550 μm, about 300 μm to about 500 μm, about 300 μm to about 450 μm, about 300 μm to about 400 μm, about 300 μm to about 350 μm, about 350 μm to about 750 μm, about 350 μm to about 700 μm, about 350 μm to about 650 μm, about 350 μm to about 600 μm, about 350 μm to about 550 μm, about 350 μm to about 500 μm, about 350 μm to about 450 μm, about 350 μm to about 400 μm, about 400 μm to about 750 μm, about 400 μm to about 700 μm, about 400 μm to about 650 μm, about 400 μm to about 600 μm, about 400 μm to about 550 μm, about 400 μm to about 500 μm, about 400 μm to about 450 μm, about 450 μm to about 750 μm, about 450 μm to about 700 μm, about 450 μm to about 650 μm, about 450 μm to about 600 μm, about 450 μm to about 550 μm, about 450 μm to about 500 μm, about 500 μm to about 750 μm, about 500 μm to about 700 μm, about 500 μm to about 650 μm,

about 500 μm to about 600 μm , about 500 μm to about 550 μm , about 550 μm to about 750 μm , about 550 μm to about 700 μm , about 550 μm to about 650 μm , about 550 μm to about 600 μm , about 600 μm to about 750 μm , about 600 μm to about 700 μm , about 600 μm to about 650 μm , about 650 μm to about 750 μm , about 650 μm to about 700 μm , or
 5 about 700 μm to about 750 μm . In certain embodiments, the particles have a particle size distribution which is defined by a $d(0.9)$ of about 10 μm to about 800 μm , preferably below 500 μm .

[119] In various embodiments, the invention provides a pharmaceutical material comprising particles of a crystalline hydrochloride salt of the compound of formula (I-A)



10

wherein the particles have a particle size distribution which is defined by a $d(0.9)$ of from about 10 μm to about 800 μm preferably below 500 μm .

[120] In certain embodiments, the particles in the composition have a needle-like crystal shape.

15 **[121]** In certain embodiments, the total deuterium abundance in the compound of formula (I-A) is about 0.3 to about 2, about 0.4 to about 2, about 0.6 to about 2, about 0.8 to about 2, about 1 to about 2, about 1.2 to about 2, about 1.4 to about 2, about 1.6 to about 2, about 1.8 to about 2, about 0.3 to about 1.8, about 0.3 to about 1.6, about 0.3 to about 1.4, about 0.3 to about 1.2, about 0.3 to about 1, about 0.3 to about 0.8, about 0.3 to about 0.6, about 0.3 to about 0.4, about 0.4 to about 1.8, about 0.4 to about 1.6, about 0.4 to about 1.4, about 0.4 to about 1.2, about 0.4 to about 1, about 0.4 to about 0.8, about 0.4 to about 0.6, about 0.6 to about 1.8, about 0.6 to about 1.6, about 0.6 to about 1.4, about 0.6 to about 1.2, about 0.6 to about 1, about 0.6 to about 0.8, about 0.8 to about 1.8, about 0.8 to about 1.6, about 0.8 to about 1.4, about 0.8 to about 1.2, about 0.8 to about 1.0,
 20 about 1 to about 1.8, about 1 to about 1.6, about 1 to about 1.4, about 1 to about 1.2, about 1.2 to about 1.8, about 1.2 to about 1.6, about 1.2 to about 1.4, about 1.4 to about 1.8, about 1.4 to about 1.6, or about 1.6 to about 1.8.

30

Pharmaceutical Compositions

[122] In one aspect, the invention provides pharmaceutical compositions comprising a pharmaceutically acceptable salt of the compound of formula I or formula I-A as described herein, including any of the pharmaceutical materials, and a pharmaceutically acceptable
5 excipient, for the treatment of a condition, disease or disorder described herein (e.g., a neurological disorder, a cancer, a respiratory disorder, a metabolic disorder, a hepatitis, a cardiovascular disease, an inflammatory or immune-mediated disorder, a dermatological disorder, a wound, a skin defect, etc.). In certain embodiments, the pharmaceutically acceptable salt form of the compound of formula I or formula I-A is a deuterium chloride
10 salt. In certain embodiments, the pharmaceutically acceptable salt form of the compound of formula I or formula I-A is a hydrochloride salt.

[123] In various embodiments, a pharmaceutical composition comprises a deuterium chloride salt of the compound of formula I and a pharmaceutically acceptable excipient. In certain embodiments, the deuterium chloride salt of the compound of formula I is a
15 crystalline deuterium chloride salt.


[124] In various embodiments, a pharmaceutical composition comprises a crystalline deuterium chloride salt of the compound of formula I and a pharmaceutically acceptable excipient.

[125] In various embodiments, a pharmaceutical composition comprises a hydrochloride salt of the compound of formula I-A and a pharmaceutically acceptable excipient. In certain
20 embodiments, the hydrochloride salt of the compound of formula I-A is a crystalline hydrochloride salt.

[126] In various embodiments, a pharmaceutical composition comprises a crystalline hydrochloride salt of the compound of formula I-A and a pharmaceutically acceptable
25 excipient.

[127] In various embodiments, the invention provides a pharmaceutical composition comprising

(i) particles of a crystalline deuterium chloride salt of a deuterium-enriched compound of formula (I)




(I), and

(ii) a pharmaceutically acceptable excipient,

wherein R¹ is H or D, provided that the abundance of deuterium in R¹ is at least 80%, and wherein the particles in the composition have a crystal shape selected from hexagonal, rod, and combinations thereof.

5 **[128]** In various embodiments, the invention provides a pharmaceutical composition comprising

(i) particles of a crystalline deuterium chloride salt of a deuterium-enriched compound of formula (I)




10 (ii) a pharmaceutically acceptable excipient,

wherein R¹ is H or D, provided that the abundance of deuterium in R¹ is at least 80%, and wherein the particles have a particle size distribution which is defined by a d(0.9) of about 10 μm to about 800 μm preferably below 500 μm.

[129] In various embodiments, the invention provides a pharmaceutical composition
15 comprising

(i) particles of a crystalline hydrochloride salt of the compound of formula (I-A)




(ii) a pharmaceutically acceptable excipient,

wherein the particles in the pharmaceutical composition have a needle-like crystal shape.

20 **[130]** In various embodiments, the invention provides a pharmaceutical composition comprising

(i) particles of a crystalline hydrochloride salt of the compound of formula (I-A)



(ii) a pharmaceutically acceptable excipient,

wherein the particles have a particle size distribution which is defined by a d(0.9) of from about 10 μm to 800 μm preferably below 500 μm .

[131] In certain embodiments, the pharmaceutical compositions described herein
5 comprise a therapeutically effective amount of a pharmaceutically acceptable salt of the compound of formula I or formula I-A. In certain embodiments, the pharmaceutical compositions described herein comprise a therapeutically effective amount of a deuterium chloride salt of the compound of formula I. In certain embodiments, the pharmaceutical compositions described herein comprise a therapeutically effective amount of a crystalline
10 deuterium chloride salt of the compound of formula I. In certain embodiments, the pharmaceutical compositions described herein comprise a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A. In certain embodiments, the pharmaceutical compositions described herein comprise a therapeutically effective amount of a crystalline hydrochloride salt of the compound of formula I-A.

[132] In certain embodiments, the crystalline deuterium chloride salt of the compound of formula I is a crystalline deuterium chloride salt as described herein.

[133] In certain embodiments, the crystalline hydrochloride salt of the compound of formula I-A is a crystalline hydrochloride salt as described herein.

[134] The pharmaceutical compositions provided herein can be administered by a
20 variety of routes including, but not limited to, oral (enteral) administration, parenteral (by injection) administration, rectal administration, transdermal administration, intradermal administration, intrathecal administration, subcutaneous (SC) administration, intravenous (IV) administration, intramuscular (IM) administration, and intranasal administration. In some embodiments, the pharmaceutical compositions disclosed herein are administered
25 orally.

[135] The pharmaceutical compositions provided herein may also be administered chronically ("chronic administration"). Chronic administration refers to administration of a compound or pharmaceutical composition thereof over an extended period of time, *e.g.*, for example, over 3 months, 6 months, 1 year, 2 years, 3 years, 5 years, *etc.*, or may be
30 continued indefinitely, for example, for the rest of the subject's life. In certain embodiments, the chronic administration is intended to provide a constant level of the compound in the blood, *e.g.*, within the therapeutic window over the extended period of time.

[136] The pharmaceutical compositions provided herein may be presented in unit dosage forms to facilitate accurate dosing. The term “unit dosage forms” refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient. Typical unit dosage forms include prefilled, premeasured ampules or syringes of the liquid compositions or pills, tablets, capsules or the like in the case of solid compositions.

[137] Although the descriptions of pharmaceutical compositions provided herein are principally directed to pharmaceutical compositions which are suitable for administration to humans, it will be understood by the skilled artisan that such compositions are generally suitable for administration to animals of all sorts. Modification of pharmaceutical compositions suitable for administration to humans in order to render the compositions suitable for administration to various animals is well understood, and the ordinarily skilled veterinary pharmacologist can design and/or perform such modification with ordinary experimentation. General considerations in the formulation and/or manufacture of pharmaceutical compositions can be found, for example, in *Remington: The Science and Practice of Pharmacy* 21st ed., Lippincott Williams & Wilkins, 2005.

Formulations

[138] Pharmaceutical compositions can be used in the preparation of individual, single unit dosage forms. Pharmaceutical compositions and dosage forms provided herein comprise a compound provided herein, or a pharmaceutically acceptable salt, solvate, or enantiomer thereof. Pharmaceutical compositions and dosage forms can further comprise one or more excipients.

[139] Pharmaceutical compositions and dosage forms provided herein can comprise one or more additional active ingredients. Examples of optional second, or additional, active ingredients are described above.

[140] Single unit dosage forms provided herein are suitable for oral, mucosal (e.g., nasal, sublingual, vaginal, buccal, or rectal), parenteral (e.g., subcutaneous, intravenous, bolus injection, intramuscular, or intraarterial), topical (e.g., eye drops or other ophthalmic preparations), transdermal or transcutaneous administration to a patient. Examples of dosage forms include, but are not limited to: tablets; caplets; capsules, such as soft elastic gelatin capsules; cachets; troches; lozenges; dispersions; suppositories; powders; aerosols (e.g., nasal sprays or inhalers); gels; liquid dosage forms suitable for oral or mucosal

administration to a patient, including suspensions (e.g., aqueous or non-aqueous liquid suspensions, oil-in-water emulsions, or a water-in-oil liquid emulsions), solutions, and elixirs; liquid dosage forms suitable for parenteral administration to a patient; eye drops or other ophthalmic preparations suitable for topical administration; and sterile solids (e.g.,
5 crystalline or amorphous solids) that can be reconstituted to provide liquid dosage forms suitable for parenteral administration to a patient.

[141] The composition, shape, and type of dosage forms will typically vary depending on their use. For example, a dosage form used in the acute treatment of a disease may contain larger amounts of one or more of the active ingredients it comprises than a dosage
10 form used in the chronic treatment of the same disease. Similarly, a parenteral dosage form may contain smaller amounts of one or more of the active ingredients it comprises than an oral dosage form used to treat the same disease. These and other ways in which specific dosage forms are used will vary from one another will be readily apparent to those skilled in the art. See, e.g., Remington's Pharmaceutical Sciences, 18th ed., Mack Publishing,
15 Easton PA (1990).

[142] In another aspect, the invention the pharmaceutical compositions and dosage forms comprise one or more excipients. Suitable excipients are well known to those skilled in the art of pharmacy, and non-limiting examples of suitable excipients are provided herein. Whether a particular excipient is suitable for incorporation into a pharmaceutical
20 composition or dosage form depends on a variety of factors well known in the art including, but not limited to, the way in which the dosage form will be administered to a patient. For example, oral dosage forms such as tablets may contain excipients not suited for use in parenteral dosage forms. The suitability of a particular excipient may also depend on the specific active ingredients in the dosage form. For example, the decomposition of some
25 active ingredients may be accelerated by some excipients, or when exposed to water. Active ingredients that comprise primary or secondary amines are particularly susceptible to such accelerated decomposition. Consequently, provided are pharmaceutical compositions and dosage forms that contain little, if any, lactose other mono- or di-saccharides. As used herein, the term "lactose-free" means that the amount of lactose
30 present, if any, is insufficient to substantially increase the degradation rate of an active ingredient.

[143] Lactose-free compositions can comprise excipients that are well known in the art and are listed, for example, in the U.S. Pharmacopeia (USP) 25-NF20 (2002). In general, lactose-free compositions comprise active ingredients, a binder/filler, and a lubricant in

pharmaceutically compatible and pharmaceutically acceptable amounts. In another aspect, lactose-free dosage forms comprise active ingredients, microcrystalline cellulose, pre-gelatinized starch, and magnesium stearate.

[144] Also provided are anhydrous pharmaceutical compositions and dosage forms comprising active ingredients, since water can facilitate the degradation of some compounds. For example, the addition of water (e.g., 5%) is widely accepted in the pharmaceutical arts as a means of simulating long-term storage in order to determine characteristics such as shelf-life or the stability of formulations over time. See, e.g., Jens T. Carstensen, *Drug Stability: Principles & Practice*, 2d. Ed., Marcel Dekker, NY, NY, 1995, pp. 379-80. In effect, water and heat accelerate the decomposition of some compounds. Thus, the effect of water on a formulation can be of great significance since moisture and/or humidity are commonly encountered during manufacture, handling, packaging, storage, shipment, and use of formulations.

[145] Anhydrous pharmaceutical compositions and dosage forms can be prepared using anhydrous or low moisture containing ingredients and low moisture or low humidity conditions. Pharmaceutical compositions and dosage forms that comprise lactose and at least one active ingredient that comprises a primary or secondary amine are preferably anhydrous if substantial contact with moisture and/or humidity during manufacturing, packaging, and/or storage is expected.

[146] An anhydrous pharmaceutical composition should be prepared and stored such that its anhydrous nature is maintained. Accordingly, anhydrous compositions are, in another aspect, packaged using materials known to prevent exposure to water such that they can be included in suitable formulary kits. Examples of suitable packaging include, but are not limited to, hermetically sealed foils, plastics, dose containers (e.g., vials), blister packs, and strip packs.

[147] Also provided are pharmaceutical compositions and dosage forms that comprise one or more compounds that reduce the rate by which an active ingredient will decompose. Such compounds, which are referred to herein as “stabilizers,” include, but are not limited to, antioxidants such as ascorbic acid, pH buffers, or salt buffers.

[148] Like the amounts and types of excipients, the amounts and specific types of active ingredients in a dosage form may differ depending on factors such as, but not limited to, the route by which it is to be administered to patients. In another aspect, dosage forms comprise a compound provided herein in an amount of from about 0.10 to about 500 mg.

Examples of dosages include, but are not limited to, 0.1, 1, 2, 5, 7.5, 10, 12.5, 15, 17.5, 20, 25, 50, 100, 150, 200, 250, 300, 350, 400, 450, or 500 mg.

[149] In another aspect, dosage forms comprise the second active ingredient in an amount of 1-about 1000 mg, from about 5 to about 500 mg, from about 10 to about 350 mg, or from about 50 to about 200 mg. Of course, the specific amount of the second active agent will depend on the specific agent used, the diseases or disorders being treated or managed, and the amount(s) of a compound provided herein, and any optional additional active agents concurrently administered to the patient.

[150] Pharmaceutical compositions that are suitable for oral administration can be provided as discrete dosage forms, such as, but not limited to, tablets (e.g., chewable tablets), caplets, capsules, and liquids (e.g., flavored syrups). Such dosage forms contain predetermined amounts of active ingredients, and may be prepared by methods of pharmacy well known to those skilled in the art. See generally, Remington's Pharmaceutical Sciences, 18th ed., Mack Publishing, Easton PA (1990).

[151] Oral dosage forms provided herein are prepared by combining the active ingredients in an intimate admixture with at least one excipient according to conventional pharmaceutical compounding techniques. Excipients can take a wide variety of forms depending on the form of preparation desired for administration. For example, excipients suitable for use in oral liquid or aerosol dosage forms include, but are not limited to, water, glycols, oils, alcohols, flavoring agents, preservatives, and coloring agents. Examples of excipients suitable for use in solid oral dosage forms (e.g., powders, tablets, capsules, and caplets) include, but are not limited to, starches, sugars, micro-crystalline cellulose, diluents, granulating agents, lubricants, binders, and disintegrating agents.

[152] In another aspect, the invention provides oral dosage forms that are tablets or capsules, in which case solid excipients are employed. In another aspect, the tablets can be coated by standard aqueous or nonaqueous techniques. Such dosage forms can be prepared by any of the methods of pharmacy. In general, pharmaceutical compositions and dosage forms are prepared by uniformly and intimately admixing the active ingredients with liquid carriers, finely divided solid carriers, or both, and then shaping the product into the desired presentation if necessary.

[153] For example, a tablet can be prepared by compression or molding. Compressed tablets can be prepared by compressing in a suitable machine the active ingredients in a free-flowing form such as powder or granules, optionally mixed with an excipient. Molded

tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

[154] Examples of excipients that can be used in oral dosage forms provided herein include, but are not limited to, binders, fillers, disintegrants, and lubricants. Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (e.g., ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, (e.g., Nos. 2208, 2906, 2910), microcrystalline cellulose, and mixtures thereof.

[155] Suitable forms of microcrystalline cellulose include, but are not limited to, the materials sold as AVICEL-PH-101, AVICEL-PH- 103 AVICEL RC-581, AVICEL-PH- 105 (available from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, PA), and mixtures thereof. A specific binder is a mixture of microcrystalline cellulose and sodium carboxymethyl cellulose sold as AVICEL RC- 581. Suitable anhydrous or low moisture excipients or additives include AVICEL-PH- 103™ and Starch 1500 LM.

[156] Examples of fillers suitable for use in the pharmaceutical compositions and dosage forms provided herein include, but are not limited to, talc, calcium carbonate (e.g., granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The binder or filler in pharmaceutical compositions is, in another aspect, present in from about 50 to about 99 weight percent of the pharmaceutical composition or dosage form.

[157] Disintegrants may be used in the compositions to provide tablets that disintegrate when exposed to an aqueous environment. Tablets that contain too much disintegrant may disintegrate in storage, while those that contain too little may not disintegrate at a desired rate or under the desired conditions. Thus, a sufficient amount of disintegrant that is neither too much nor too little to detrimentally alter the release of the active ingredients may be used to form solid oral dosage forms. The amount of disintegrant used varies based upon the type of formulation, and is readily discernible to those of ordinary skill in the art. In another aspect, pharmaceutical compositions comprise from about 0.5 to about 15 weight percent of disintegrant, or from about 1-about 5 weight percent of disintegrant.

[158] Disintegrants that can be used in pharmaceutical compositions and dosage forms include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline

cellulose, croscarmellose sodium, crospovidone, polacrillin potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other algin, other celluloses, gums, and mixtures thereof.

[159] Lubricants that can be used in pharmaceutical compositions and dosage forms include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (e.g., peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laureate, agar, and mixtures thereof. Additional lubricants include, for example, a
10 syloid® silica gel (AEROSIL200, manufactured by W.R. Grace Co. of Baltimore, MD), a coagulated aerosol of synthetic silica (marketed by Degussa Co. of Plano, TX), CAB-O-SIL (a pyrogenic silicon dioxide product sold by Cabot Co. of Boston, MA), and mixtures thereof. If used at all, lubricants may be used in an amount of less than about 2 weight percent of the pharmaceutical compositions or dosage forms into which they are incorporated.

15 **[160]** In another aspect, the invention provides a solid oral dosage form comprising a compound provided herein, anhydrous lactose, microcrystalline cellulose, polyvinylpyrrolidone, stearic acid, colloidal anhydrous silica, and gelatin.

[161] Active ingredients provided herein can also be administered by controlled release means or by delivery devices that are well known to those of ordinary skill in the art. Examples include, but are not limited to, those described in U.S. Patent Nos.: 3,845,770; 20 3,916,899; 3,536,809; 3,598,123; and 4,008,719, 5,674,533, 5,059,595, 5,591,767, 5,120,548, 5,073,543, 5,639,476, 5,354,556, and 5,733,566, each of which is incorporated in its entirety herein by reference. Such dosage forms can be used to provide slow or controlled-release of one or more active ingredients using, for example, hydropropylmethyl
25 cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the active agents provided herein. In another aspect,
30 the invention provides single unit dosage forms suitable for oral administration such as, but not limited to, tablets, capsules, gelcaps, and caplets that are adapted for controlled-release.

[162] Controlled-release pharmaceutical products improve drug therapy over that achieved by their non-controlled counterparts. In another aspect, the invention provides

the use of a controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include extended activity of the drug, reduced dosage frequency, and increased patient compliance. In addition, controlled-release formulations can be used to affect the time of onset of action or other characteristics, such as blood levels of the drug, and can thus affect the occurrence of side (e.g., adverse) effects.

[163] In another aspect, the controlled-release formulations are designed to initially release an amount of drug (active ingredient) that promptly produces the desired therapeutic or prophylactic effect, and gradually and continually release of other amounts of drug to maintain this level of therapeutic or prophylactic effect over an extended period of time. In another aspect, in order to maintain a constant level of drug in the body, the drug can be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled release of an active ingredient can be stimulated by various conditions including, but not limited to, pH, temperature, enzymes, water, or other physiological conditions or compounds.

[164] Parenteral dosage forms can be administered to patients by various routes including, but not limited to, subcutaneous, intravenous (including bolus injection), intramuscular, and intraarterial. Administration of a parenteral dosage form bypasses patients' natural defenses against contaminants, and thus, in these aspects, parenteral dosage forms are sterile or capable of being sterilized prior to administration to a patient. Examples of parenteral dosage forms include, but are not limited to, solutions ready for injection, dry products ready to be dissolved or suspended in a pharmaceutically acceptable vehicle for injection, suspensions ready for injection, and emulsions.

[165] Suitable vehicles that can be used to provide parenteral dosage forms are well known to those skilled in the art. Examples include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and nonaqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

[166] Compounds that increase the solubility of one or more of the active ingredients disclosed herein can also be incorporated into the parenteral dosage forms. For example,

cyclodextrin and its derivatives can be used to increase the solubility of a compound provided herein. See, e.g., U.S. Patent No. 5,134,127, which is incorporated in its entirety herein by reference.

[167] Topical and mucosal dosage forms provided herein include, but are not limited to, 5 sprays, aerosols, solutions, emulsions, suspensions, eye drops or other ophthalmic preparations, or other forms known to one of skill in the art. See, e.g., Remington's Pharmaceutical Sciences, 16th and 18th eds., Mack Publishing, Easton PA (1980 & 1990); and Introduction to Pharmaceutical Dosage Forms, 4th ed., Lea & Febiger, Philadelphia (1985). Dosage forms suitable for treating mucosal tissues within the oral cavity can be 10 formulated as mouthwashes or as oral gels.

[168] Suitable excipients (e.g., carriers and diluents) and other materials that can be used to provide topical and mucosal dosage forms encompassed herein are well known to those skilled in the pharmaceutical arts, and depend on the particular tissue to which a given pharmaceutical composition or dosage form will be applied. In another aspect, 15 excipients include, but are not limited to, water, acetone, ethanol, ethylene glycol, propylene glycol, butane-1,3-diol, isopropyl myristate, isopropyl palmitate, mineral oil, and mixtures thereof to form solutions, emulsions or gels, which are nontoxic and pharmaceutically acceptable. Moisturizers or humectants can also be added to pharmaceutical compositions and dosage forms. Examples of additional ingredients are 20 well known in the art. See, e.g., Remington's Pharmaceutical Sciences, 16th and 18th eds., Mack Publishing, Easton PA (1980 & 1990).

[169] The pH of a pharmaceutical composition or dosage form may also be adjusted to improve delivery of one or more active ingredients. Also, the polarity of a solvent carrier, its ionic strength, or tonicity can be adjusted to improve delivery. Compounds such as 25 stearates can also be added to pharmaceutical compositions or dosage forms to alter the hydrophilicity or lipophilicity of one or more active ingredients so as to improve delivery. In other aspects, stearates can serve as a lipid vehicle for the formulation, as an emulsifying agent or surfactant, or as a delivery-enhancing or penetration-enhancing agent. In other aspects, salts, solvates, prodrugs, or enantiomers of the active ingredients can be used to 30 further adjust the properties of the resulting composition.

[170] In another aspect, the active ingredients provided herein are not administered to a patient at the same time or by the same route of administration. In another aspect, provided are kits which can simplify the administration of appropriate amounts of active ingredients.

[171] In another aspect, the invention provides a kit comprising a dosage form of a compound provided herein. Kits can further comprise additional active ingredients.

[172] In other aspects, the kits can further comprise devices that are used to administer the active ingredients. Examples of such devices include, but are not limited to, syringes, drip bags, patches, and inhalers.

[173] Kits can further comprise cells or blood for transplantation as well as pharmaceutically acceptable vehicles that can be used to administer one or more active ingredients. For example, if an active ingredient is provided in a solid form that must be reconstituted for parenteral administration, the kit can comprise a sealed container of a suitable vehicle in which the active ingredient can be dissolved to form a particulate-free sterile solution that is suitable for parenteral administration. Examples of pharmaceutically acceptable vehicles include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

[174] In an aspect, provided is a pharmaceutical composition comprising a hydrogen chloride salt or deuterium chloride salt of deuterium-enriched pioglitazone as described herein, lactose, a carmellose, a hypromellose, and a stearate salt.

Methods of Use and Treatment

[175] Provided herein are methods of treating a condition, disease, or disorder (e.g., a neurological disorder, a cancer, a respiratory disorder, an endocrine disorder, a metabolic disorder, a renal disorder, a hepatitis, a cardiovascular disease, an inflammatory or immune-mediated disorder, a dermatological disorder, a wound, a skin defect, etc.), the method comprising administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat the condition, disease, or disorder. In various embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to treat the condition,

disease, or disorder. In various embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to treat condition, disease, or disorder.

5 **[176]** In certain embodiments, the condition, disease, or disorder is a neurological disorder. In certain embodiments, the condition, disease, or disorder is a cancer. In certain
embodiments, the condition, disease, or disorder is a respiratory disorder. In certain
embodiments, the condition, disease, or disorder is an endocrine or metabolic disorder. In
certain embodiments, the condition, disease, or disorder is a metabolic disorder. In certain
10 embodiments, the condition, disease, or disorder is a hepatitis. In certain embodiments,
the condition, disease, or disorder is a cardiovascular disease. In certain embodiments, the
condition, disease, or disorder is a renal disease. In certain embodiments, the condition,
disease, or disorder is an inflammatory or immune-mediated disorder. In certain
embodiments, the condition, disease, or disorder is a dermatological disorder. In certain
15 embodiments, the condition, disease, or disorder is a wound. In certain embodiments, the
condition, disease, or disorder is a skin defect.

[177] Also provided herein are methods of modulating the amount and/or function of an
endogenous biological molecule (e.g., a triglyceride, a fatty acid, a carbohydrate or sugar, a
low-density lipoprotein, a high-density lipoprotein, a cytokine, etc.) for the prevention or
20 treatment of a condition, disease, or disorder described herein, the method comprising
administering to a patient in need thereof a therapeutically effective amount of a
pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the
compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt
described herein, to modulate the endogenous biological molecule. In various
25 embodiments, the method comprises administering to a patient in need thereof a
therapeutically effective amount of a deuterium chloride salt of the compound of formula I,
e.g., a crystalline deuterium chloride salt form as described herein, to modulate the
endogenous biological molecule. In various embodiments, the method comprises
administering to a patient in need thereof a therapeutically effective amount of a
30 hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt
form described herein, to modulate the endogenous biological molecule.

[178] In certain embodiments, the endogenous biological molecule is a triglyceride. In
certain embodiments, the endogenous biological molecule is a fatty acid. In certain
embodiments, the endogenous biological molecule is a carbohydrate or sugar. In certain

embodiments, the endogenous biological molecule is a low-density lipoprotein. In certain embodiments, the endogenous biological molecule is a high-density lipoprotein. In certain embodiments, the endogenous biological molecule is a cytokine.

5 (i) *Treating Metabolic Disorders*

[179] Another aspect of the invention provides a method of treating a metabolic disorder or hepatic disorder selected from the group consisting of nonalcoholic fatty liver disease, nonalcoholic steatohepatitis, viral hepatitis, liver cirrhosis, liver fibrosis, diabetic retinopathy, diabetic neuropathy, diabetic nephropathy, beta cell depletion insulin resistance in a patient with congenital adrenal hyperplasia treated with a glucocorticoid, polycystic ovary syndrome, leukodystrophies including adrenoleukodystrophy and adrenomyeloneuropathy, dysmetabolism in peritoneal dialysis patients, reduced insulin secretion, improper distribution of brown fat cells and white fat cells, obesity, or improper modulation of leptin levels. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat the metabolic disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to treat the metabolic disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to treat the metabolic disorder. In certain embodiments, the metabolic disorder is further selected from a complication of diabetes. In certain embodiments, the metabolic disorder is nonalcoholic fatty liver disease, diabetic retinopathy, diabetic neuropathy, diabetic nephropathy, or beta cell depletion insulin resistance in a patient with congenital adrenal hyperplasia treated with a glucocorticoid. In certain embodiments, the metabolic disorder is nonalcoholic fatty liver disease, diabetic retinopathy, diabetic neuropathy, diabetic nephropathy, beta cell depletion, reduced insulin secretion, improper distribution of brown fat cells and white fat cells, obesity, or improper modulation of leptin levels. In certain other embodiments, the metabolic disorder is non-alcoholic fatty liver disease. In certain other embodiments, the metabolic disorder is non-alcoholic steatohepatitis. In certain other embodiments, the metabolic

disorder is Type II diabetes mellitus. In certain other embodiments, the metabolic disorder is beta cell loss treatable by B-cell regeneration. In certain other embodiments, the metabolic disorder is central obesity, dyslipidemia, or pre-diabetes. In certain other embodiments, the metabolic disorder is polycystic ovary syndrome. In certain other
5 embodiments, the metabolic disorder is leukodystrophy including adrenoleukodystrophy and adrenomyeloneuropathy.

Nonalcoholic Fatty Liver Disease

[180] In certain embodiments, a method is provided for treatment of nonalcoholic fatty
10 liver disease, the method comprising administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat the nonalcoholic fatty liver disease. The therapeutic methods are contemplated to provide particular benefits to
15 patients suffering from nonalcoholic fatty liver disease. Exemplary benefits include little to no occurrence of PPAR gamma side effects (e.g., weight gain, edema, and/or bone loss) while achieving improvement in the patient's nonalcoholic fatty liver disease (which may include reduced amount of hepatic fat due to the therapy).

Nonalcoholic Steatohepatitis

[181] In certain embodiments, a method is provided for treatment of nonalcoholic steatohepatitis, the method comprising administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium
25 chloride salt or a hydrochloride salt described herein, to treat the nonalcoholic steatohepatitis. The therapeutic methods are contemplated to provide particular benefits to patients suffering from nonalcoholic steatohepatitis. Exemplary benefits include little to no occurrence of PPAR gamma side effects (e.g., weight gain, edema, and/or bone loss) while achieving improvement in the patient's nonalcoholic steatohepatitis (which may include
30 reduced amount of hepatic fat due to the therapy).

Type II Diabetes Mellitus

[182] In certain embodiments, a method is provided for treatment of Type II diabetes mellitus, the method comprising administering to a patient in need thereof a

therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat the Type II diabetes mellitus. The therapeutic methods are contemplated to provide particular benefits to patients suffering from Type II diabetes mellitus. Exemplary benefits include little to no occurrence of PPAR gamma side effects (e.g., weight gain, edema, and/or bone loss) while achieving improvement in the patient's Type II diabetes mellitus (which may include improvement in the patient's glycemic control).

10 *(ii) Treatment of Cancer*

[183] Another aspect of the invention provides a method of treating cancer. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat the cancer. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to treat the cancer. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to treat the cancer.

[184] In certain embodiments, the cancer is lung cancer, hepatocellular carcinoma, astrocytoma, glioma, glioblastoma, meningioma, liver cancer, lymphoma, melanoma, multiple myeloma, pancreatic cancer, colorectal cancer, pituitary cancer, thyroid cancer, esophageal cancer, or prostate cancer. In certain embodiments, the cancer is non-small cell lung cancer or hepatocellular carcinoma.

[185] In certain other embodiments, the cancer is lung cancer, hepatocellular carcinoma, astrocytoma, glioma, glioblastoma, meningioma, liver cancer, lymphoma, melanoma, multiple myeloma, pancreatic cancer, colorectal cancer, pituitary cancer, thyroid cancer, esophageal cancer, prostate cancer, nose cancer, throat cancer, kidney cancer, breast cancer, stomach cancer, or uterine cancer. In certain other embodiments, the cancer is brain cancer, bladder cancer, breast cancer, cervical cancer, colon cancer, colorectal cancer, endometrial cancer, esophageal cancer, leukemia, lung cancer, liver cancer, melanoma, ovarian cancer, pancreatic cancer, prostate cancer, rectal cancer, renal

cancer, stomach cancer, testicular cancer, or uterine cancer. In yet other embodiments, the cancer is a vascularized tumor, squamous cell carcinoma, adenocarcinoma, small cell carcinoma, melanoma, glioma, neuroblastoma, sarcoma (e.g., an angiosarcoma or chondrosarcoma), larynx cancer, parotid cancer, biliary tract cancer, thyroid cancer, acral

5 lentiginous melanoma, actinic keratosis, acute lymphocytic leukemia, acute myeloid leukemia, adenoid cystic carcinoma, adenoma, adenosarcoma, adenosquamous carcinoma, anal canal cancer, anal cancer, anorectal cancer, astrocytic tumor, Bartholin's gland carcinoma, basal cell carcinoma, biliary cancer, bone cancer, bone marrow cancer, bronchial cancer, bronchial gland carcinoma, carcinoid, cholangiocarcinoma, chorioid

10 plexus papilloma/carcinoma, chronic lymphocytic leukemia, chronic myeloid leukemia, clear cell carcinoma, connective tissue cancer, cystadenoma, digestive system cancer, duodenum cancer, endocrine system cancer, endodermal sinus tumor, endometrial hyperplasia, endometrial stromal sarcoma, endometrioid adenocarcinoma, endothelial cell cancer, ependymal cancer, epithelial cell cancer, Ewing's sarcoma, eye and orbit cancer,

15 female genital cancer, focal nodular hyperplasia, gallbladder cancer, gastric antrum cancer, gastric fundus cancer, gastrinoma, glioblastoma, glucagonoma, heart cancer, hemangioblastoma, hemangioendothelioma, hemangioma, hepatic adenoma, hepatic adenomatosis, hepatobiliary cancer, hepatocellular carcinoma, Hodgkin's disease, ileum cancer, insulinoma, intraepithelial neoplasia, interepithelial squamous cell neoplasia,

20 intrahepatic bile duct cancer, invasive squamous cell carcinoma, jejunum cancer, joint cancer, Kaposi's sarcoma, pelvic cancer, large cell carcinoma, large intestine cancer, leiomyosarcoma, lentigo maligna melanoma, lymphoma, male genital cancer, malignant melanoma, malignant mesothelial tumor, medulloblastoma, medulloepithelioma, meningeal cancer, mesothelial cancer, metastatic carcinoma, mouth cancer,

25 mucoepidermoid carcinoma, multiple myeloma, muscle cancer, nasal tract cancer, nervous system cancer, neuroepithelial adenocarcinoma, nodular melanoma, non-epithelial skin cancer, non-Hodgkin's lymphoma, oat cell carcinoma, oligodendroglial cancer, oral cavity cancer, osteosarcoma, papillary serous adenocarcinoma, penile cancer, pharynx cancer, pituitary tumor, plasmacytoma, pseudosarcoma, pulmonary blastoma, rectal cancer, renal

30 cell carcinoma, respiratory system cancer, retinoblastoma, rhabdomyosarcoma, sarcoma, serous carcinoma, sinus cancer, skin cancer, small cell carcinoma, small intestine cancer, smooth muscle cancer, soft tissue cancer, somatostatin-secreting tumor, spine cancer, squamous cell carcinoma, striated muscle cancer, submesothelial cancer, superficial spreading melanoma, T-cell leukemia, tongue cancer, undifferentiated carcinoma, ureter

cancer, urethral cancer, urinary bladder cancer, urinary system cancer, uterine cervix cancer, uterine corpus cancer, uveal melanoma, vaginal cancer, verrucous carcinoma, VIPoma, vulva cancer, well differentiated carcinoma, or Wilms tumor.

[186] In certain other embodiments, the cancer is non-Hodgkin's lymphoma, such as a B-cell lymphoma or a T-cell lymphoma. In certain embodiments, the non-Hodgkin's lymphoma is a B-cell lymphoma, such as a diffuse large B-cell lymphoma, primary mediastinal B-cell lymphoma, follicular lymphoma, small lymphocytic lymphoma, mantle cell lymphoma, marginal zone B-cell lymphoma, extranodal marginal zone B-cell lymphoma, nodal marginal zone B-cell lymphoma, splenic marginal zone B-cell lymphoma, Burkitt lymphoma, lymphoplasmacytic lymphoma, hairy cell leukemia, or primary central nervous system (CNS) lymphoma. In certain other embodiments, the non-Hodgkin's lymphoma is a T-cell lymphoma, such as a precursor T-lymphoblastic lymphoma, peripheral T-cell lymphoma, cutaneous T-cell lymphoma, angioimmunoblastic T-cell lymphoma, extranodal natural killer/T-cell lymphoma, enteropathy type T-cell lymphoma, subcutaneous panniculitis-like T-cell lymphoma, anaplastic large cell lymphoma, or peripheral T-cell lymphoma.

(iii) Treating Respiratory Disorders

[187] Another aspect of the invention provides a method of treating a respiratory disorder. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat the respiratory disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to treat the respiratory disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to treat the respiratory disorder.

[188] In certain embodiments, the respiratory disorder is chronic obstructive pulmonary disease, asthma, bronchitis, cystic fibrosis, pulmonary edema, pulmonary embolism, pulmonary arterial hypertension, pneumonia, pulmonary sarcoidosis, silicosis, pulmonary fibrosis, respiratory failure, acute respiratory distress syndrome, emphysema, chronic

bronchitis, tuberculosis, lung cancer, or a chronic respiratory condition. In certain embodiments, the respiratory disorder is chronic obstructive pulmonary disease, asthma, or a chronic respiratory condition. In certain other embodiments, the respiratory disorder is chronic obstructive pulmonary disease. In yet other embodiments, the respiratory disorder is bronchitis, cystic fibrosis, pulmonary edema, pulmonary embolism, pneumonia, pulmonary sarcoidosis, silicosis, pulmonary fibrosis, respiratory failure, acute respiratory distress syndrome, emphysema, chronic bronchitis, tuberculosis, or lung cancer. In certain embodiments, the asthma is mild asthma, moderate asthma, severe asthma, or steroid-resistant asthma.

10

(iv) Treatment of Neurological Disorders

[189] Accordingly, one aspect of the invention provides a method of treating a neurological disorder selected from the group consisting of Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, Friedreich's ataxia, autism spectrum disorder, depression, mild cognitive impairment, Down syndrome, neurodegeneration, adrenoleukodystrophy, adrenomyeloneuropathy, Zellweger's disease, Huntington's disease, stroke, traumatic brain injury, substance abuse, spinal cord injury, neuronal injury, major depression or bipolar disorder comorbid with metabolic syndrome, and a neurological disorder caused by functional mitochondrial impairment. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat the neurological disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to treat the neurological disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to treat the neurological disorder. In certain embodiments, the neurological disorder is selected from the group consisting of Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, Friedreich's ataxia, autism spectrum disorder, depression, mild cognitive impairment, neurodegeneration, adrenoleukodystrophy, adrenomyeloneuropathy, Huntington's disease, stroke, traumatic brain injury, substance abuse, spinal cord injury, neuronal injury,

and major depression or bipolar disorder comorbid with metabolic syndrome. In certain embodiments, the neurological disorder is selected from the group consisting of Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, Friedreich's ataxia, depression, mild cognitive impairment, neurodegeneration, adrenoleukodystrophy, adrenomyeloneuropathy, and Huntington's disease. In certain other embodiments, the neurological disorder is Alzheimer's disease. In certain other embodiments, the neurological disorder is Down syndrome. In certain other embodiments, the neurological disorder is adrenoleukodystrophy. In certain other embodiments, the neurological disorder is adrenomyeloneuropathy.

10 **[190]** In certain other embodiments, the neurological disorder is a cognitive disorder, such as cognitive impairment and/or memory impairment. The cognitive impairment may be, for example, cognitive impairment associated with Alzheimer's disease.

[191] In certain embodiments, the substance abuse is one or more of alcohol craving, heroin dependence, and nicotine dependence.

15

(v) Treating a Symptom of Hepatitis

[192] Another aspect of the invention provides a method of treating a symptom of hepatitis. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat the hepatitis. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to treat the hepatitis. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to treat the hepatitis.

(vi) Treating Cardiovascular Disease

30 **[193]** Another aspect of the invention provides a method of treating a cardiovascular disease. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat the cardiovascular disease. In certain

embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to treat the cardiovascular disease. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to treat the cardiovascular disease. In certain embodiments, the cardiovascular disease is hypertension, hyperlipidemia, atherosclerosis, improper vascular function, dyslipidemia, stenosis, restenosis, myocardial infarction, stroke, intracranial hemorrhage, acute coronary syndrome, stable angina pectoris, or unstable angina pectoris. In certain other embodiments, the cardiovascular disorder is intracranial hemorrhage, acute coronary syndrome, stable angina pectoris, or unstable angina pectoris.

[194] In another aspect, the invention provides a method for preventing stroke in a patient. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to prevent stroke in a patient. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to prevent stroke in a patient. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to prevent stroke in a patient.

[195] The method of treatment or the method of prevention may involve a patient at risk for central nervous system ischemic stroke, or may involve a patient at risk for stroke due to cardiovascular disease.

(vii) Reducing the Amount of a Triglyceride or Low-Density Lipoprotein

[196] Another aspect of the invention provides a method of reducing the amount of a triglyceride or low-density lipoprotein (LDL) in a patient. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt

described herein, to reduce the amount of a triglyceride or low-density lipoprotein (LDL) in a patient. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein,

5 to reduce the amount of a triglyceride or low-density lipoprotein (LDL) in a patient. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to reduce the amount of a triglyceride or low-density lipoprotein (LDL) in a patient.

10 **[197]** In certain embodiments, the method provides a reduction of at least 1%, 5%, 10%, or 25% in the amount of a triglyceride or low-density lipoprotein (LDL) in the patient.

(viii) Increasing the Amount of High-Density Lipoprotein

[198] Another aspect of the invention provides a method of increasing the amount of high-density lipoprotein (HDL) in a patient. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to increase the amount of high-density lipoprotein (HDL) in a patient. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to increase the amount of high-density lipoprotein (HDL) in a patient. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to increase the amount of high-density lipoprotein (HDL) in a patient.

25 **[199]** In certain embodiments, the method provides an increase of at least 1%, 5%, 10%, or 25% in the amount of high-density lipoprotein (HDL) in a patient.

30 *(ix) Treating an Inflammatory or Immune-Mediated Disorder*

[200] Another aspect of the invention provides a method of treating an inflammatory or immune-mediated disorder selected from the group consisting of chronic kidney disease, arthritis, a primary cicatricial alopecia, lung fibrosis, multiple sclerosis, endotoxemia, sepsis, septic shock, laminitis, inflammatory bowel disease, colitis, Crohn's disease, rheumatoid

arthritis, lupus, myasthenia gravis, vasculitis, chronic pancreatitis, a hyperproliferative skin disorder, an inflammatory skin disorder, rhinitis (e.g., allergic rhinitis), and a dermatological condition. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat the inflammatory or immune-mediated disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to treat the inflammatory or immune-mediated disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to treat the inflammatory or immune-mediated disorder. In certain embodiments, the inflammatory or immune-mediated disorder is selected from the group consisting of chronic kidney disease, arthritis, a primary cicatricial alopecia, lung fibrosis, multiple sclerosis, endotoxemia, sepsis, septic shock, laminitis, inflammatory bowel disease, colitis, Crohn's disease, rheumatoid arthritis, lupus, myasthenia gravis, vasculitis, chronic pancreatitis, a hyperproliferative skin disorder, an inflammatory skin disorder, and a dermatological condition. In certain embodiments, the inflammatory or immune-mediated disorder is selected from the group consisting of chronic kidney disease, arthritis, a primary cicatricial alopecia, lung fibrosis, multiple sclerosis, endotoxemia, sepsis, septic shock, laminitis, inflammatory bowel disease, colitis, Crohn's disease, rheumatoid arthritis, lupus, myasthenia gravis, vasculitis, chronic pancreatitis, a hyperproliferative skin disorder, an inflammatory skin disorder, and a dermatological condition, in certain embodiments, the chronic kidney disease may be, for example, polycystic kidney disease (such as autosomal dominant or autosomal recessive).

(x) Treating a Dermatological Disorder

[201] Another aspect of the invention provides a method of treating a dermatological disorder selected from the group consisting of psoriasis, atopic dermatitis, acne, leukoplakia, scleroderma, and a skin malignancy. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat the

dermatological disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to treat the dermatological disorder. In certain embodiments, the method

5 comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to treat the dermatological disorder. In certain embodiments, the administering is by topical administration.

10 *(xi) Modulating Expression of Pro-Inflammatory Cytokines*

[202] Another aspect of the invention provides a method of modulating expression of a pro-inflammatory cytokine (e.g., TNF α , IL-1 β , IL-6, IL-17, IL-23, or MCP-1) in a patient suffering from an inflammatory disorder. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of

15 a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to modulate expression of a pro-inflammatory cytokine (e.g., TNF α , IL-1 β , or IL-6) in a patient suffering from an inflammatory disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of

20 the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to modulate expression of a pro-inflammatory cytokine (e.g., TNF α , IL-1 β , or IL-6) in a patient suffering from an inflammatory disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt

25 form described herein, to modulate expression of a pro-inflammatory cytokine (e.g., TNF α , IL-1 β , or IL-6) in a patient suffering from an inflammatory disorder. In certain embodiments, the pro-inflammatory cytokine is TNF α .

[203] Another aspect of the invention provides a method of modulating expression of an anti-inflammatory cytokine in a patient suffering from an inflammatory disorder. The

30 method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to modulate expression of an anti-inflammatory cytokine in a patient suffering from an inflammatory disorder. In certain embodiments, the

method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to modulate expression of an anti-inflammatory cytokine in a patient suffering from an inflammatory disorder. In certain

5 embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to modulate expression of an anti-inflammatory cytokine in a patient suffering from an inflammatory disorder.

10 *(xii) Modulating Macrophage Function*

[204] Another aspect of the invention provides a method of modulating macrophage function in a patient suffering from an infection, inflammatory disorder, or autoimmune disease. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or

15 formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to modulate macrophage function in a patient suffering from an infection. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described

20 herein, to modulate macrophage function in a patient suffering from an infection, inflammatory disorder, or autoimmune disease. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to modulate macrophage function in a patient suffering from an

25 infection.

(xiii) Method of Promoting Wound Healing

[205] Another aspect of the invention provides a method of promoting wound healing. The method comprises administering to a patient in need thereof a therapeutically

30 effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to promote wound healing. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline

deuterium chloride salt form as described herein, to promote wound healing. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to promote wound healing. In
5 certain embodiments, the administering is by topical administration.

(xiv) Treating Skin Defects

[206] Another aspect of the invention provides a method of treating skin defects caused by exposure to ultraviolet radiation. The method comprises administering to a patient in
10 need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat skin defects caused by exposure to ultraviolet radiation. In certain embodiments, the method
15 comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to treat skin defects caused by exposure to ultraviolet radiation. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to
20 treat skin defects caused by exposure to ultraviolet radiation.

(xv) Method of Modulating Stem Cell Differentiation

[207] Another aspect of the invention provides a method of modulating stem cell differentiation, such as in a patient. The method comprises administering to a patient in
25 need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to modulate stem cell differentiation, such as in a patient. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a
30 deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to modulate stem cell differentiation, such as in a patient. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A,

e.g., a crystalline hydrochloride salt form described herein, to modulate stem cell differentiation, such as in a patient.

(xvi) Additional Medical Disorders

- 5 **[208]** Another aspect of the invention provides a method of treating a disorder selected from the group consisting of transplant rejection, liver functional impairment, Rabson-Mendenhall syndrome, Donohue syndrome, Leber hereditary optic neuropathy, myotonic dystrophy, ototoxicity, Niemann Pick disease, autosomal dominant optic atrophy, spinal bulbar muscular atrophy, Mohr-Tranebjaerg syndrome, hereditary spastic paraplegia,
- 10 MELAS syndrome, monoclonal immunoglobulin deposition disease (MIDD), deafness, insulin resistance in a patient receiving growth hormone, and chronic progressive external ophthalmo-plegia with mitochondrial myopathy. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula
- 15 I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat the disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to treat the disorder. In certain embodiments, the method comprises administering to a
- 20 patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to treat the disorder.

(xvii) Preventing Medical Disorders

- 25 **[209]** Also provided are methods of preventing a medical disorder in a patient. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to prevent the medical disorder. In certain
- 30 embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to prevent the medical disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of

formula I-A, e.g., a crystalline hydrochloride salt form described herein, to prevent the medical disorder. The medical disorder may be one or more of the medical disorders recited above, such as a neurological disorder (e.g., Alzheimer's disease or Parkinson's disease), cancer (e.g., non-small cell lung cancer or hepatocellular carcinoma), a metabolic disorder, a cardiovascular disorder (e.g., in-stent renarrowing in diabetes patients, 5 reinfarction in diabetes patients, or cardiac allograft vasculopathy after heart transplant), or a respiratory disorder (e.g., chronic obstructive pulmonary disease).

(xviii) Additional Medical Uses

10 **[210]** The invention provides methods of using the compounds and solid forms described herein for therapy comprising regenerative medicine. Also provided herein are methods of treating a veterinary disorder, such as laminitis. The method comprises administering to a patient in need thereof a therapeutically effective amount of a pharmaceutically acceptable salt of a compound of formula I or formula I-A, e.g., the 15 compound of formula I or formula I-A as a deuterium chloride salt or a hydrochloride salt described herein, to treat the veterinary disorder. In certain embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a deuterium chloride salt of the compound of formula I, e.g., a crystalline deuterium chloride salt form as described herein, to treat the veterinary disorder. In certain 20 embodiments, the method comprises administering to a patient in need thereof a therapeutically effective amount of a hydrochloride salt of the compound of formula I-A, e.g., a crystalline hydrochloride salt form described herein, to treat the veterinary disorder.

[211] In certain embodiments, the pharmaceutically acceptable salts of the compound of formula I or formula I-A provided herein may be administered as the sole active agent, or 25 they may be administered in combination with other therapeutically active agents (e.g., a combination therapy).

[212] It is contemplated that combination therapies of the invention described herein may work synergistically in the treatment of particular conditions, diseases or disorders described herein, and/or one or more symptoms associated with such conditions, diseases 30 or disorders. It is further contemplated that the pharmaceutically acceptable salts of the compound of formula I or formula I-A provided herein may also work to alleviate adverse effects associated with a second therapeutically active agent, and vice versa.

[213] In various embodiments, one or more second therapeutically active agents can be used in the methods and compositions provided herein. In certain embodiments, the one

or more second therapeutically active agents may be large molecules (e.g., proteins) or small molecules (e.g., synthetic inorganic, organometallic, or organic molecules).

[214] In certain embodiments, the combination therapy comprises a pharmaceutically acceptable salt of the compound of formula I or formula I-A described herein and a second therapeutically active agent for the treatment of a condition, disease or disorder described herein (e.g., a neurological disorder, a cancer, a respiratory disorder, a metabolic disorder, a hepatitis, a cardiovascular disease, an inflammatory or immune-mediated disorder, a dermatological disorder, a wound, a skin defect, etc.).

[215] In certain embodiments, the second therapeutically active agent may be an agent useful for the treatment of a metabolic disorder, such therapeutically active agents may include, but are not limited to, metformin, imeglimin, a dipeptidyl peptidase IV inhibitor (e.g., sitagliptin, vildagliptin, or the like), a statin (e.g., a HMG-CoA reductase inhibitor, such as atorvastatin, cerivastatin, fluvastatin, lovastatin, mevastatin, simvastatin, rosuvastatin, pravastatin, or combinations thereof), an AMPK activator, a thyroid- β agonist, a GLP-1 agonist, a GLP-2 agonist, or an SGLT2 inhibitor.

[216] In certain embodiments, the second therapeutically active agent is a diuretic agent (e.g., hydrochlorothiazide).

[217] In certain embodiments, the second therapeutically active agent may be an agent useful for the treatment of hypertension, diabetes, or an inflammatory disorder. In certain embodiments, the second therapeutically active agent may be an agent that limits the activity of the renin-angiotensin system, such as an angiotensin converting enzyme inhibitor (e.g., an ACE inhibitor, such as ramipril, captopril, enalapril, or the like), an angiotensin receptor blocker (e.g., candesartan, losartan, olmesartan, or the like), or a renin inhibitor. In certain embodiments, the second therapeutic agent may limit hypertension by alternate means, for example, a beta-adrenergic receptor blocker or calcium channel blocker (e.g., amlodipine).

[218] In certain embodiments, second therapeutically active agent is a glucocorticoid agonist. In certain embodiments, a combination therapy comprising a pharmaceutically acceptable salt of the compound of formula I or formula I-A described herein and a glucocorticoid agonist may be useful for the treatment of an inflammatory disorder, such as therapy for suppressing an immune response, preventing transplant rejection, and treating autoimmune disease. Exemplary disorders include, for example, rheumatoid arthritis, lupus, myasthenia gravis, muscular dystrophy vasculitis, multiple sclerosis, chronic

obstructive pulmonary disease (COPD), inflammatory bowel disease, treatment of acute allergic reactions, and transplant rejection.

[219] In certain embodiments, a combination therapy comprising a pharmaceutically acceptable salt of the compound of formula I or formula I-A described herein and a second therapeutically active agent that increases cAMP or a beta-adrenergic agonist may be useful in the treatment of a kidney disease. Exemplary beta-adrenergic agonists include, but are not limited to, a beta-1-adrenergic agonist, a beta-2-adrenergic agonist, a beta-3-adrenergic agonist, or combinations thereof. In certain embodiments, the second therapeutically active agent is noradrenaline, isoprenaline, dobutamine, salbutamol, levosalbutamol, terbutaline, pirbuterol, procaterol, metaproterenol, fenoterol, bitolterol mesylate, salmeterol, formoterol, bambuterol, clenbuterol, indacaterol, L-796568, amibegron, solabegron, isoproterenol, albuterol, metaproterenol, arbutamine, befunolol, bromoacetylalprenololmenthane, broxaterol, cimaterol, cirazoline, denopamine, dopexamine, epinephrine, etilefrine, hexoprenaline, higenamine, isoetharine, isoxsuprine, mabuterol, methoxyphenamine, nylidrin, oxyfedrine, prenalterol, ractopamine, reproterol, rimiterol, ritodrine, tretoquinol, tulobuterol, xamoterol, zilpaterol, zinterol, or a pharmaceutically acceptable salt thereof; or a combination of any of the foregoing.

[220] In certain embodiments, the combination therapy comprises a pharmaceutically acceptable salt of the compound of formula I or formula I-A described herein and a second therapeutically active agent useful in the treatment of a cancer. Exemplary second therapeutically active agents useful for the treatment of cancer include, but are not limited to, an alkylating agent, an anti-metabolite (e.g., a molecule that impedes DNA and/or RNA synthesis), an anti-microtubule agent, a topoisomerase inhibitor, a cytotoxic antibiotic, a tyrosine kinase inhibitor, an inhibitor of tumor necrosis factor alpha, anti-neoplastic radiation therapy, or a Programmed Death protein-1 (PD-1) modulator (e.g., an inhibitor). In certain embodiments, the second therapeutically active agent useful for the treatment of a cancer is azacitidine, azathioprine, bleomycin, carboplatin, capecitabine, carmustine, cisplatin, chlorambucil, cyclophosphamide, cytarabine, dacarbazine, daunorubicin, docetaxel, doxifluridine, doxorubicin, epirubicin, epothilone, etoposide, fluorouracil, fulvestrant, gemcitabine, hydroxyurea, idarubicin, imatinib, lomustine, mechlorethamine, mercaptopurine, methotrexate, mitoxantrone, oxaliplatin, paclitaxel, pemetrexed, procarbazine, raloxifene, teniposide, temozolomide, tamoxifen, toremifene, valrubicin, vinblastine, vincristine, vindesine, vinorelbine, or a pharmaceutically acceptable salt thereof; or a combination of any of the foregoing.

[221] In certain embodiments, the second therapeutically active agent useful for the treatment of a cancer is abraxane; acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; amrubicin; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; 5 azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefmgol; celecoxib; chlorambucil; cirolemycin; cisplatin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; dactinomycin; 10 daunorubicin hydrochloride; decitabine; dexormaplatin; de/.aguanine; dezaguanine mesylate; diaziquone; docetaxel; doxorubicin; doxorubicin hydrochloride; raloxifene; raloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitrucin; enloplatm; enpromate; epipropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; 15 etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; flurocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; herceptin; hydroxyurea; idarubicin hydrochloride; ifosfamide; ilmofofosine; iproplatin; irinotecan; irinotecan hydrochloride; lanreotide acetate; lapatinib; letrozole; leuprolide acetate; 20 liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedopa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; 25 nocodazole; nogalamycin; ormaplatin; oxisuran; paclitaxel; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman; piposulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; portiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; romidepsin; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; 30 sparsomycin; spirogermanium hydrochloride; spiromustine; spiroplatin; a stem cell treatment; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; taxotere; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepa; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate; triciribine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin;

tubulozole hydrochloride; uracil mustard; uredepa; vapreotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinat sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; zorubicin hydrochloride, or combinations thereof.

- 5 **[222]** Administration of a pharmaceutically acceptable salt of the compound of formula I or formula I-A described herein and the second therapeutically active agent(s) to a patient can occur simultaneously or sequentially by the same or different routes of administration. The suitability of a particular route of administration employed for a particular active agent will depend on the active agent itself (e.g., whether it can be administered orally without
10 decomposing prior to entering the blood stream) and the disease being treated. One route of administration for compounds provided herein is oral. Routes of administration for the second active agents or ingredients are known to those of ordinary skill in the art. See, e.g., Physicians' Desk Reference (60th Ed., 2006).

15

EXAMPLES

[223] In order that the disclosure described herein may be more fully understood, the following examples are set forth. The synthetic and biological examples described in this application are offered to illustrate the compounds, pharmaceutical compositions, and methods provided herein and are not to be construed in any way as limiting their scope.

20

Example 1: Synthesis of Deuterium-Enriched (*R*)-Pioglitazone L-Dibenzoyl Tartrate Salt

[224] Reaction temperatures are reported as internal temperatures. Chemical intermediates, reagents, and solvents were obtained from commercial sources.

- [225]** Filtration: Solid products were isolated by filtration over a PTFE Büchner funnel
25 using MN 617 G (fast filtration, phosphate-free; ref: MN 494024) and MN640m (ref: MN 203015) filter paper (unless otherwise stated) and washed with 1:1 MeOD:D₂O (v/v) in a plug-flow method (portions; 2x / 3x) (unless otherwise stated).

[226] Drying: After filtration, first solids were air-dried *in vacuo* for a few hours (3-4 h) and then dried (24-72 h) in a drying oven at 50 °C / 42 °C fitted with a vacuum pump.


- 30 **[227]** NMR: ¹H spectra were recorded on a Bruker Avance 300 MHz or higher spectrometer. Chemical shifts were referenced to residual solvent signals at δ 2.50 (DMSO-d₆) relative to TMS as internal standard wherever applied.

[228] HPLC: For %d.e. / %e.e. measurement, all samples were dissolved in MeOH to make a solution of 1 mg/mL.

[229] HPLC for identity and chemical purity measurement: Samples were run on a 150 mm x 4.6 mm, 5 μ m YMC triart C18 column. The mobile phase was an isocratic elution system with 1:1 0.1 M NH_4OAc : CH_3CN plus 2% HOAc. Flow rate was 0.75 mL/min; run time, 35 min; and detector wavelength, 269 nm.

- 5 [230] In the following examples, when details of an example run are presented, notes in square brackets ([]) are used to denote optional changes that are not part of the example run but could be performed or were performed in another run.

Step 1: Preparation of Racemic Deuterium-Enriched Pioglitazone (Compound B)



- 10 [231] Pioglitazone·HCl (compound A HCl salt) was dissolved in D_2O at elevated temperature. After stirring at 90 °C to 100 °C for at least 5 h, a small amount of DCl was added followed by cooling down to ambient temperature. After aging the suspension for a few hours at 15 °C to 25 °C, the product was filtered off as compound B DCl salt. After this first cycle, approximately 95% of the hydrogen at the chiral center had been exchanged by
15 deuterium.

[232] If desired, the crude product was then subjected to a second cycle of the same treatment to further increase the deuteration grade to 98+ % in compound (B) as determined by ^1H -NMR.


Example run:

- 20 [233] 1st deuteration cycle: 19.96 kg pioglitazone (compound (A)) was charged in a 100 L glass-lined vessel. 79.54 kg (71.86 L; 4.0 equivalents w/w) D_2O was then charged in the same vessel. The reaction mixture was heated and stirred to about 95 °C and stirred at that temperature for 5 h. [Temperatures can be from about 90 °C to about 100 °C (preferably 95 °C) for at least 5 h. A temperature of 95 °C or lower will lead to lower deuteration grade,
25 while temperatures from about 95 °C to about 100 °C worked better. Lower amounts of D_2O were tried, but the reaction mixture was difficult to stir with less than 4 w/w equivalents, and increase in deuteration was minimal above 5 w/w equivalents. Significant increase in deuteration grade was observed up until 3 to 4 hours at elevated temperature.] A thin white suspension was observed and 0.13 kg (0.10 L, 0.005 v/w equivalents) 35% DCl
30 in D_2O was added, and the transfer line was rinsed with approximately 0.25 L D_2O (0.0125 v/w equivalents). The reaction mixture was cooled to about 25 °C over 240 min

approximately [Cooling time is at least 3 h to a final temperature from about 15 °C to about 25 °C]. The reaction mixture was then stirred for 13 h approximately at 15 °C [Holding time is at least 5 h from about 15 °C to about 25 °C. The solid obtained is more stirrable and filterable if cooled over 3 hours or more rather than only 1 hour.] The crude product was
 5 filtered off on a 140 L stainless steel nutsche filter. The product was washed with 5.52 kg (4.99 L; 0.25 v/w equivalents) D₂O. The crude product was blown dry in a stream of nitrogen for 3 h approximately [Drying time should be at least 1 h]. The moist crude product was isolated, and a monitoring sample was used for ¹H-NMR. The sample was then subjected to a second round of deuteration to increase the deuterium content. [Over three
 10 runs, % deuteration on chiral center of 96.5%, 97.2%, and 96.9% were observed at this stage.]

[234] Second deuteration cycle: The entire amount of moist crude product was charged in a vessel. 75.57 kg (68.26 L) D₂O was then charged. The reaction mixture was heated to about 95 °C for 5 h approximately [Heating can be to a temperature from about 90 °C to
 15 about 100 °C (preferably 95 °C) for at least 5 h]. A thin white suspension was observed. 0.12 kg (0.09 L) DCI in D₂O 35% was then added, and the transfer line was rinsed with approximately 0.25 L D₂O. The reaction mixture was cooled to 25 °C over 4 h [Cooling ramp should be at least 3 h to a temperature from about 15 °C to about 25 °C]. The reaction mixture was stirred at about 20 °C for 9 h [Stirring can be done at 15 °C to 25 °C for at least
 20 5 h]. Then the crude product was filtered off on a 140 L stainless steel nutsche filter and was rinsed with 5.52 kg (4.99 L) D₂O. Deuterium NMR showed more than 2 deuteriums per molecule. The compound (B) DCI salt product (or optionally a mixture of HCl and DCI salt) was dried on a 140 L stainless steel nutsche filter in vacuo at about max. 60 °C until LOD (loss on drying) ≤2% (48 h approximately) [Drying temperature should not exceed a
 25 maximum of 60 °C].

Step 2: Preparation of Deuterium-Enriched (R)-Pioglitazone L-dibenzoyl tartrate salt (Compound (C))



[235] The synthesis started with dissolving compound (B) or salt thereof and L-dibenzoyl tartrate in a mixture of MeOD (methanol-d₁) and D₂O at elevated temperature. The mixture was then cooled to approximately 55 °C. Seed crystals (2% or less w/w relative to (compound B)) were optionally added, followed by a rather slow cooling ramp and


extended aging time. Eventually, the product compound (C) was filtered off, washed with a mixture of MeOD and D₂O and dried *in vacuo*.

Example run:

[236] 28.10 kg racemic deuterium-enriched pioglitazone (compound (B)) DCI salt was charged in a 1000 L glass vessel, followed by 25.49 kg L,L-dibenzoyltartaric acid (1 molar equiv.) and 187.3 kg MeOD (230.4 L; 6.67 w/w equivalents) [It is possible to use 0.6 molar equivalents of L,L-dibenzoyltartaric acid resulting in slightly lower yield but comparable % ee]. The reaction mixture was warmed to a temperature from about 60 °C to about 70 °C (65 °C achieved), and 280.9 kg D₂O (253.7 L; 10.0 w/w equivalents) was added. The reaction was warmed to 70 °C to 80 °C (76 °C achieved), and the mixture was stirred at that temperature for at least 30 minutes. An almost clear solution was obtained. The mixture was then cooled to 55 °C over 66 min [Cooling time should be at least one hour to a temperature from about 52 °C to about 58 °C]. The mixture was then stirred for 2 h at 56 °C [Stirring can be done for at least 2 hours at a temperature from about 52 °C to about 58 °C]), during which time seed crystals (0.25 kg) of compound (C) were added [Seeding can be with crystals of compound (C) or (C')]. The suspension was cooled to 25 °C for 7.5 hours [Cooling time to a temperature from about 22 °C to about 28 °C should be at least 5 hours]. Then the suspension was stirred for 37 h [Stirring should take at least 15 hours at 22 °C to 28 °C]).

20 **[237]** The product was filtered off on a stainless-steel centrifuge, and was washed with
a premixed mixture of D₂O (13.38 kg; 12.08 L; 0.48 w/w equiv.) and MeOD (9.82 kg; 12.08 L;
0.35 w/w equiv.). The product was blown dry in a stream of nitrogen for at least 1 h. The
product compound (C) was then dried in vacuo at a maximum temperature of 60 °C on a
Hastelloy vacuum tray dryer. Deuterium NMR showed approximately 2 deuteriums per
25 molecule.

Step 3: Recrystallization of Enantioenriched (R)-Deuterium-enriched Pioglitazone L-Dibenzoyl Tartrate Salt (Compound (C) to Recrystallized Compound (C'))



Example run:

30 **[238]** To prepare HCl salt derivative: 46.88 kg enantioenriched (*R*)-deuterium-enriched pioglitazone dibenzoyl tartrate salt (compound (C)) was charged in a 1000 L glass vessel, followed by 185.6 kg MeOH (234.4 L; 4.0 w/w equivalents), 55.0 kg D₂O (49.7 L; 1.17 w/w

equivalents), and 7.1 kg 35% DCl in D₂O (5.6 L; 0.15 w/w equivalents). The reaction mixture was warmed to 60 °C to 70 °C (63 °C achieved), and 218.0 kg D₂O (196.9 L; 4.65 w/w equivalents) was added. The reaction was warmed to 70 °C to 80 °C, and the mixture was stirred at that temperature for at least 30 minutes. An almost clear solution was obtained.

- 5 The mixture was then cooled to 55 °C during 85 min. [Cooling time should be at least one hour]. The mixture was then stirred for at 55 °C for 120 min and seed crystals of (compound C') (0.14 kg) were added [Stirring can be performed at 52-58 °C for at least 2 hours during which time seed crystals of compound (C) or (C') could be added if desired]. The suspension was cooled to 26 °C over 10 h 50 min [Cooling ramp to 22 °C to 28 °C should be at least 5 hours long]. Then the suspension was stirred at 25 °C for 34 h 20 min [Suspension should be stirred for at least 15 hours at 22 °C to 28 °C].

- [239] The product was filtered off on a stainless-steel centrifuge, and was washed with a premixed mixture of D₂O (17.1 kg; 15.5 L; 0.36 w/w equiv.) and MeOD (12.3 kg; 15.5 L; 0.26 w/w equiv.). The product compound (C') was blown dry in a stream of nitrogen for at least 1 h. The product was then submitted to HPLC for measurement of optical purity. If optical purity was adequate (above 91%), compound (C') was dried in vacuo at a maximum temperature of 60 °C on a Hastelloy vacuum tray dryer. If not, a second recrystallization was performed (actual example: 93% ee; no second recrystallization). Deuterium NMR showed approximately 2 deuteriums per molecule.

- 20 [240] If a second or further recrystallizations are necessary, the above procedure can be repeated and the recrystallized material separated and dried in vacuo at a maximum temperature of 60 °C on a Hastelloy vacuum tray dryer as described above.

[241] It should be noted that the details given below include the procedures of both steps b) and c).

- 25 [242] To prepare DCl salt derivative: only deuterated solvents were required to be used.

Example 2: Preparation and Characterization of the Crystalline DCl Salt of Deuterium-Enriched (*R*)-Pioglitazone

- 30 [243] 278.5 g of deuterium-enriched (*R*)-pioglitazone L-dibenzoyl tartrate salt (as prepared in Example 1), 450 mL of MeOD and about 38% DCl in D₂O (36 g, 2.9 equiv.) was heated to 50 °C. The slightly turbid solution was filtered over a Büchner funnel (100 mL) fitted with porcelain-1 filter and the filtrate was reheated to 50 °C. Ethyl acetate (1500 mL) was then added in seven portions (temp. dropped to 30 °C) and the mixture was allowed to

attain 20 °C. After stirring for 2 hours at this temperature, the crystalline DCI Salt of deuterium-enriched (*R*)-pioglitazone was isolated by filtration over a Büchner funnel (250 mL) fitted with porcelain-4 filter. After washing thoroughly (in portions) with ethyl acetate (5 x 250 mL), isolated wet cake was dried overnight at 50 °C in vacuo to obtain 44.6 g (Y = 89.9%) of the deuterium chloride salt as a white crystalline material. The enantiomeric excess was found to be 97.1% (HPLC) and the D-content on chiral center 99.0% (¹H-NMR). Note: Other samples synthesized using exactly the same process were analyzed by ²H-NMR and the deuterium content by ²H-NMR indicated about 3 deuteriums on the molecule (one on the chiral center, DCI and one deuterium on the exchangeable NH position).

10 **[244]** Chiral HPLC method: Samples were run on a 250 mm x 4.6 mm, 3 µm Chiralpak IC or equivalent. The mobile phase was an isocratic elution system with 70:30 (v/v) Hexane/IPA. Flow rate was 1.0 mL/min; run time, 30 min; and detector wavelength, 225 nm.

15 **[245]** ¹H-NMR method: About 5 to 10 mg sample were dissolved in DMSO D6 for ¹H-NMR analysis. ¹H-NMR analysis was performed using a Bruker Avance 300 MHz or higher spectrometer.

Characterization by X-ray Powder Diffraction

20 **[246]** An X-ray powder diffractogram of the crystalline DCI salt of deuterium-enriched (*R*)-pioglitazone is provided in FIG. 1A. X-ray powder diffraction data were collected using a D8 Advance diffractometer using Cu K_{α1} radiation (1.54056 Å) with a germanium monochromator at room temperature (e.g., about 21 °C to about 23 °C). Detector scans on a solid state LynxEye detector were performed using 0.016° per step, with a scan speed of 5 sec/step. Samples were analyzed in 8 mm long glass capillary tubes with a 0.3 mm outer diameter. Tabulated characteristics of the X-ray powder diffractogram in FIG. 1A are provided below in Table 4, which lists diffraction angle 2θ, inter-planar distances d, and relative intensity (expressed as a percentage with respect to the most intense peak).

25

Table 4 – X-ray Powder Diffractogram Data of the Crystalline Deuterium Chloride Salt of Deuterium-Enriched (*R*)-Pioglitazone

2θ (°)	d value (Å)	Intensity (%)
8.6	10.27	18
8.8	10.07	17
12.8	6.93	22
12.9	6.87	26
15.9	5.59	4
18.7	4.73	18
18.9	4.70	16
19.7	4.51	18
20.0	4.45	34
20.8	4.26	56
21.2	4.19	13
22.1	4.02	8
22.8	3.89	100
23.7	3.75	3
25.6	3.47	5
26.0	3.43	20
26.7	3.34	9
27.3	3.26	6
28.1	3.18	12
29.8	3.00	12
31.3	2.86	21
32.1	2.79	10
32.3	2.77	5
33.4	2.68	3
33.6	2.66	3
34.4	2.60	4
36.0	2.50	3

[247] As shown in FIG. 1B, it can be noted that the X-ray powder diffraction (XRPD) pattern of (*R*)-2H-pioglitazone HCl salt (upper trace) is not identical with the XRPD pattern of pioglitazone HCl salt (lower trace), in particular showing doubling of several peaks.

Characterization by Single Crystal X-ray Diffraction

5 [248] Single crystals of the crystalline DCl salt of deuterium-enriched (*R*)-pioglitazone were analyzed by single crystal X-ray diffraction. Data were collected using a Nonius Kappa-CCD instrument at 296 K.

[249] Data reduction was performed using HKL Scalepack (Otwinowski & Minor 1997) and cell parameters were obtained using Denzo and Scalepak (Otwinowski & Minor, 1997).

10 [250] The crystal structure of the crystalline DCl salt of deuterium-enriched (*R*)-pioglitazone was solved using direct methods by SHELXT-2014/7 (Sheldrick, G. M., 2015a). The structure was refined by least square full matrix refinement using SHELXL-2014/7 (Sheldrick, G. M., 2015b). All H-atoms connected to the C were included from the geometry and kept with fixed thermal parameters. The H atoms involved in the hydrogen bond
15 network were found in the Fourier difference map and were refined isotropically.

[251] The unit cell parameters of the crystalline DCl salt of deuterium-enriched (*R*)-pioglitazone and the data collection and structure refinement methods are shown in Table 5.

Table 5 – Unit Cell Parameters and Data Collection and Structure Refinement Methods for the Crystalline DCl Salt of Deuterium-Enriched (*R*)-Pioglitazone

Empirical formula	C ₁₉ H ₁₈ D ₃ N ₂ O ₃ S ⁺ Cl ⁻
formula weight	395.9
T [K]	296(2) K
λ [Å]	0.71073 Å
Crystal system	Monoclinic
Space group	P2 ₁
Unit cell dimensions	
a [Å]	10.1319(9)
b [Å]	9.4067(9)
c [Å]	10.3231(6)
β [°]	94.397(3)
V[Å ³]	980.98(14)
Z	2

D_c [g/cm³]	1.330
μ [mm⁻¹]	0.322
F(000)	412
Crystal size [mm³]	0.45 x 0.25 x 0.18
θ range for data collection [°]	2.7 → 32.6
Reflections collected	11335
Independent reflections	6943 [R _{int} = 0.0191]
Completeness to θ = 25.242° [%]	98.7
Absorption correction	Integration
Max. and min. transmission	0.958 and 0.877
Data/restraints/parameters	6943 / 1 / 248
Goodness-of-fit on F²	1.030
R₁, wR₂ [I > 2σ(I)]	0.0371, 0.0962
R₁, wR₂ (all data)	0.0444, 0.1034
Flack parameter	-0.013(18)
Absolute configuration	<i>R</i>
Largest diff. peak and hole [e/Å³]	0.191 and -0.252

Characterization by Optical Microscopy

[252] An optical micrograph of the crystalline DCI salt of deuterium-enriched (*R*)-pioglitazone is shown in FIG. 2. Optical micrographs were obtained using a Leica DM
5 2500M optical microscope. The crystals exhibited both hexagonal and rod-like crystal shapes.

Characterization by Differential Scanning Calorimetry

[253] A differential scanning calorimetry (DSC) curve of the crystalline DCI salt of
10 deuterium-enriched (*R*)-pioglitazone is provided in FIG. 3. DSC data were collected using a heat flux DSC3+ STARE system. Samples (~2 mg) were sealed in standard 40 μL aluminum pans, pin-holed, and heated in the DSC from 25 °C to 300 °C, at a heating rate of 10 °C/min. During measurement, dry N₂ gas at a flow rate of 50 mL/min was used to purge the sample chamber. The DSC curve displayed an endothermic event, corresponding to the melting of
15 the crystalline DCI salt form of deuterium-enriched (*R*)-pioglitazone, with onset and peak values at about 191 °C and about 202 °C, respectively.

Characterization by Thermogravimetric Analysis and Thermal Gravimetric Mass

Spectrometry

[254] Thermogravimetric analysis (TGA) and thermal gravimetric mass spectrometry (TGMS) data for the crystalline DCl salt form of deuterium-enriched (*R*)-pioglitazone are provided in FIG. 4A and FIG. 4B, respectively. TGA data were collected using a TGA/DSC 3+ STARe system (Mettler-Toledo GmbH, Switzerland) calibrated for temperature using samples of indium and aluminum. Samples were weighed in to 100 μ L aluminum crucibles and sealed. The seals were pin-holed, and the crucibles heated from 25 °C to 300 °C at a heating rate of 10 °C. During measurement dry N₂ gas was used for purging. The volatiles produced by the TGA samples upon heating were analyzed by an Omnistar GSD 301 T2 (Pfeiffer Vacuum GmbH, Germany) mass spectrometer. The TGA and TGMS data showed that the crystalline DCl salt form of deuterium-enriched (*R*)-pioglitazone is anhydrous, with a mass loss of 0.07% between 40 °C and 170 °C, and that thermal decomposition occurred above 190 °C.

Characterization by NMR Spectroscopy

[255] ¹H-NMR spectra for the crystalline DCl salt form of deuterium-enriched (*R*)-pioglitazone were collected at room temperature on a 500 MHz Bruker instrument using standard pulse sequences. Samples of the crystalline DCl salt form of deuterium-enriched (*R*)-pioglitazone were dissolved in DMSO-*d*₆. The ¹H-NMR chemical shifts (ppm) for the crystalline DCl salt form of deuterium-enriched (*R*)-pioglitazone were as follows: 1.25 (t, 3H), 2.79 (q, 2H), 3.07 (d, 1H), 3.30 - 3.34 (m, 1H), 3.43 (t, 2H), 4.38 (t, 2H), 6.83 - 6.93 (m, 2H), 7.11 - 7.24 (m, 2H), 7.94 (br d, 1H), 8.37 (br d, 1H), 8.66 - 8.80 (m, 1H), 12.04 (s, 1H). A representative ¹H-NMR spectrum of the crystalline DCl salt is provided in FIG. 5.

Example 3: Preparation and Characterization of the Crystalline HCl Salt Form of Deuterium-Enriched (*R*)-Pioglitazone

[256] 2.2 g of deuterium-enriched (*R*)-pioglitazone L-dibenzoyl tartrate salt (as prepared in Example 1), 11 mL of MeOH and 37% HCl in H₂O (0.308 g, 2.9 equiv.) was heated to 40 °C. Subsequently, 40 mL of ethyl acetate was added to the slightly turbid solution and the mixture was allowed to cool to 20 °C. After stirring for 2 hours at this temperature the HCl salt form of deuterium-enriched (*R*)-pioglitazone was isolated by filtration (P-4 Büchner funnel with a filter paper) and washed thoroughly with ethyl acetate

(5 x 20 mL). After drying overnight at 50 °C *in vacuo* 1.1 g (Y = 90.9%) of white crystalline material was obtained. The enantiomeric excess was found to be 97.2% (HPLC) and the D-content >98% (¹H-NMR). Note: Deuterium content by ²H-NMR indicated about 1 deuterium on the molecule which was confirmed to be on the chiral center by ¹H-NMR.

5 **[257]** Chiral HPLC method: Samples were run on a 250 mm x 4.6 mm, 3 μm Chiralpak IC or equivalent. The mobile phase was an isocratic elution system with 70:30 (v/v) Hexane/IPA. Flow rate was 1.0 mL/min; run time, 30 min; and detector wavelength, 225 nm

[258] ¹H-NMR method: About 5 to 10 mg sample were dissolved in DMSO D6 for ¹H-NMR analysis. ¹H-NMR analysis was performed using a Bruker Avance 300 MHz or higher spectrometer.

Characterization by Powder X-ray Diffraction

[259] An X-ray powder diffractogram of the crystalline HCl salt form of deuterium-enriched (*R*)-pioglitazone is provided in FIG. 6. X-ray powder diffraction data were collected as described in Example 2. Tabulated characteristics of the X-ray powder diffractogram in FIG. 6 are provided below in Table 6, which lists diffraction angle 2θ, inter-planar distances d, and relative intensity (expressed as a percentage with respect to the most intense peak).

Table 6 – X-ray Powder Diffractogram Data of the Crystalline HCl Salt Form of Deuterium-Enriched (*R*)-Pioglitazone

2θ (°)	d value (Å)	Intensity (%)
8.6	10.27	32
8.8	10.08	31
12.8	6.93	30
12.9	6.87	27
15.9	5.59	6
18.7	4.73	25
18.9	4.70	12
19.7	4.51	29
20.0	4.45	80
20.8	4.26	50
21.2	4.19	18
22.1	4.02	14
22.8	3.89	100

2 θ (°)	d value (Å)	Intensity (%)
23.7	3.75	4
25.6	3.47	6
26.0	3.43	36
26.7	3.34	9
27.3	3.26	15
28.1	3.17	30
29.8	3.00	9
31.3	2.86	19
32.1	2.79	13
32.3	2.77	7
33.4	2.68	3
33.6	2.66	3
34.4	2.61	5
36.0	2.50	5

Characterization by Optical Microscopy

[260] An optical micrograph of the crystalline HCl salt of deuterium-enriched (*R*)-pioglitazone is shown in FIG. 7. Optical micrographs were obtained using a Leica DM
5 2500M optical microscope. The crystals exhibited both hexagonal and rod-like crystal shapes.

Characterization by Differential Scanning Calorimetry

[261] A differential scanning calorimetry (DSC) curve of the crystalline HCl salt form of
10 deuterium-enriched (*R*)-pioglitazone is provided in FIG. 8. DSC data were collected using the method described in Example 2. The DSC curve displayed an endothermic event, corresponding to the melting of the crystalline HCl salt form of deuterium-enriched (*R*)-pioglitazone, with onset and peak values at about 190 °C and about 200 °C, respectively.

Characterization by Thermogravimetric Analysis/Simultaneous Difference Thermal Analysis and Thermal Gravimetric Mass Spectrometry

[262] Thermogravimetric analysis (TGA) and thermal gravimetric mass spectrometry (TGMS) data for the crystalline HCl salt form of deuterium-enriched (*R*)-pioglitazone are

provided in FIG. 9A and FIG. 9B, respectively. TGA and TGMS data were collected using the methods described in Example 2. The TGA and TGMS data showed that the crystalline HCl salt form of deuterium-enriched (*R*)-pioglitazone is anhydrous, with a mass loss of 0.2% between 40 °C and 180 °C, and that thermal decomposition occurred above 220 °C.

5

Characterization by NMR Spectroscopy

[263] ¹H-NMR spectra for the crystalline HCl salt form of deuterium-enriched (*R*)-pioglitazone were collected as described in Example 2. Samples of the crystalline HCl salt form of deuterium-enriched (*R*)-pioglitazone were dissolved in DMSO-*d*₆. The ¹H-NMR chemical shifts (ppm) for the crystalline HCl salt form of deuterium-enriched (*R*)-pioglitazone were as follows: 1.25 (t, 3H), 2.79 (q, 2H), 3.07 (d, 1H), 3.30 - 3.34 (m, 1H), 3.43 (t, 2H), 4.38 (t, 2H), 6.83 - 6.93 (m, 2H), 7.11 - 7.24 (m, 2H), 7.94 (br d, 1H), 8.37 (br d, 1H), 8.66 - 8.80 (m, 1H), 12.04 (s, 1H). A representative ¹H-NMR spectrum of the crystalline HCl salt is provided in FIG. 10.

15

Example 4: Preparation and Characterization of the Crystalline Free-Base Form of Deuterium-Enriched (*R*)-Pioglitazone

[264] The free-base form of deuterium-enriched *R*-Pioglitazone was prepared by incubating the crystalline DCl salt of deuterium-enriched *R*-Pioglitazone from Example 2 in water at 50 °C for 2 weeks.

[265] The enantiomeric excess, as measured by HPLC, was determined to be 43%. The HPLC method used is described in Examples 2 and 3.

[266] The D-content on the chiral center, as measured by ¹H-NMR, was determined to be about 66%. The ¹H-NMR method used is described in Examples 2 and 3.

25 Characterization by Powder X-ray Diffraction

[267] An X-ray powder diffractogram of the crystalline free-base form of deuterium-enriched *R*-pioglitazone is provided in FIG. 11. X-ray powder diffraction data were collected as described in Example 2. Representative diffraction peaks from the X-ray powder diffractogram in FIG. 11, expressed in terms diffraction angle (2θ), are provided below in Table 7.

30

**Table 7 – X-ray Powder Diffractogram Data of the Crystalline Free-Base Form of
Deuterium-Enriched *R*-Pioglitazone**

2θ (°)
6.2
7.8
8.6
8.8
9.2
10.3
12.4
12.9
15.1
15.6
16.3
18.6
19.6
20.0
20.8
21.4
22.6
22.8
24.0
25.0
25.6
26.4
26.9
27.4
28.1
29.1
29.5
31.1

Characterization by Differential Scanning Calorimetry

[268] A differential scanning calorimetry (DSC) curve of the crystalline free-base form of deuterium-enriched *R*-pioglitazone is provided in FIG. 12. DSC data were collected using the method described in Example 2. The DSC curve displayed an endothermic event,
5 corresponding to the melting of the crystalline free-base form of deuterium-enriched *R*-pioglitazone, with onset and peak values at about 174 °C and about 178 °C, respectively.

Characterization by Thermogravimetric Analysis/Simultaneous Difference Thermal Analysis and Thermal Gravimetric Mass Spectrometry

10 [269] Thermogravimetric analysis (TGA) and thermal gravimetric mass spectrometry (TGMS) data for crystalline free-base form of deuterium-enriched *R*-pioglitazone are provided in FIG. 13A and FIG. 13B, respectively. TGA and TGMS data were collected using the methods described in Example 2. The TGA and TGMS data showed that the crystalline free-base form of deuterium-enriched *R*-pioglitazone is anhydrous, with thermal
15 decomposition occurring above 160 °C.

Characterization by NMR Spectroscopy

[270] ¹H-NMR spectra for the crystalline free-base form of deuterium-enriched (*R*)-pioglitazone were collected as described in Example 2. Samples of the crystalline free-base
20 form of deuterium-enriched (*R*)-pioglitazone were dissolved in DMSO-*d*₆. The ¹H-NMR chemical shifts (ppm) for the crystalline free-base form of deuterium-enriched (*R*)-pioglitazone were as follows: 1.20 (t, 3H), 2.56 - 2.70 (m, 2H), 3.06 (d, 1H), 3.17 (t, 2H), 3.24 - 3.33 (m, 1H), 4.32 (t, 2H), 4.88 (dd, 1H), 6.80 - 6.93 (m, 2H), 7.15 (m, 2H), 7.35 (br d, 1H), 7.66 (br d, 1H), 8.41 (s, 1H), 12.03 (s, 1H).

25

Example 5: Solubility of the Crystalline DCl and HCl Salts of Deuterium-Enriched (*R*)-Pioglitazone in Various Organic Solvents

[271] Crystalline DCl salt of deuterium-enriched (*R*)-pioglitazone: suspensions of the crystalline material in ethanol, methanol, 3-pentanone, 2-propanol, DMSO, *tert*-butyl
30 methyl ether, *N,N*-dimethylformamide, ethyl acetate, acetone, water, tetrahydrofuran, and chloroform were prepared and incubated for 24 hours at room temperature (about 21 °C to about 23 °C). Aliquots of the mother liquor were then taken from each suspension and analyzed by HPLC to determine the concentration of deuterium-enriched (*R*)-pioglitazone in the different solvents. HPLC methods are shown in Table 8.

Table 8: HPLC Method

HPLC system:			
Instrument:	Agilent 1200		
Detector 1:	DAD set at 265 nm		
Detector 2:	HP1100 LC/MSD in positive scan mode		
HPLC conditions:			
Auto sampler temp.:	15 °C		
Column:	Waters Sunfire C18 (100 x 4.6 mm; 3.5 µm)		
Column temp.:	35 °C		
Flow cell:	10 mm path		
Gradient:	Mobile phase A:	10 mM ammonium acetate	
	Mobile phase B:	acetonitrile LC/MS grade	
Flow:	1.0 mL/min		
Gradient:	Time [min]:	Eluent A:	Eluent B:
	0	90%	10%
	1	90%	10%
	6	10%	90%
	9	10%	90%
	12	90%	10%
Run time:	13 min		
Sample:			
Concentration:	ca. 1 mg/mL		
Solvent:	10 mM ammonium acetate:acetonitrile (50:50 v/v)		
Injection volume:	5 µL		
Retention time:	6.2 min		

[272] Crystalline HCl salt of deuterium-enriched (*R*)-pioglitazone: **Qualitative testing:** (i)

- 5 About 5 mg of the crystalline HCl salt was added to 1 mL of each solvent (DMSO, N,N-dimethylformamide, methanol, water, ethanol, tetrahydrofuran, ethyl acetate, toluene, and dichloromethane); (ii) the resulting mixtures were then stirred at ambient temperature (about 21 °C to about 23 °C) for 5-10 minutes; (iii) the mixtures were then visually inspected

to determine if the crystalline HCl salt had completely dissolved; (iv) if the crystalline HCl salt had not completely dissolved more solvent was added until complete dissolution occurred. **Quantitative testing:** (i) 100 mg of crystalline HCl salt was added to 4 vol. of each solvent (0.4 mL) dimethylformamide, methanol and DMSO; (ii) the resulting mixtures were then stirred at ambient temperature (about 20 °C to about 25 °C) for 5-10 minutes; (iii) the mixtures were then visually inspected to determine if the crystalline HCl salt had completely dissolved; (iv) if the crystalline HCl salt had not completely dissolved more solvent was added until complete dissolution occurred.

[273] The solubilities were then calculated using the total quantity of crystalline HCl salt and solvent used for each solvent.

[274] Solubility data for the crystalline DCl and HCl salts of deuterium-enriched (*R*)-pioglitazone in various organic solvents are shown in Table 9 and Table 10, respectively.

Table 9. Solubility of the Crystalline DCl Salt Form of Deuterium-Enriched (*R*)-Pioglitazone in Various Organic Solvents

Solvent	Solubility (mg/mL)
DMSO	225.9
N,N-dimethylformamide	46.4
Methanol	45.0
Water	7.7
Ethanol	4.4
Chloroform	2.0
Tetrahydrofuran	1.8
3-pentanone	0.5
2-propanol	0.5
Acetone	0.5
<i>tert</i> -butyl methyl ether	< 0.1
Ethyl acetate	< 0.1

Table 10. Solubility of the Crystalline HCl Salt Form of Deuterium-Enriched (*R*)-Pioglitazone in Various Organic Solvents

Solvent	Solubility (mg/mL)
DMSO	> 50
N,N-dimethylformamide	~ 50
Methanol	~ 50
Ethyl acetate	< 5 - 10
Ethanol	< 5 – 10
Toluene	< 5 – 10
Tetrahydrofuran	< 5 – 10
Dichloromethane	< 5 – 10
Water	< 5 - 10

Example 6: Effect of pH on the Solubility of the Crystalline DCl and HCl Salts of Deuterium-Enriched (*R*)-Pioglitazone and Crystalline Pioglitazone HCl

[275] The pH-dependent solubility profiles of the crystalline DCl salt form of deuterium-enriched (*R*)-pioglitazone, and crystalline pioglitazone HCl were determined at room temperature (about 21 °C to about 23 °C) in the pH range of 0.3 – 7.6. For the crystalline HCl salt form of deuterium-enriched (*R*)-pioglitazone, the pH-dependent solubility profile was measured at room temperature (about 21 °C to about 23 °C) in the pH range of 0.3 – 3.5.

[276] For the crystalline DCl salt form of deuterium-enriched pioglitazone, the pH-dependent solubility profile was measured at room temperature (about 21 °C to about 23 °C) in the pH range of 0.7 – 3.5.

[277] Suspensions of the three crystalline forms were prepared in water or USP buffer solutions. The pH of the suspensions was adjusted through the addition of 0.1 M HCl or 0.1 M NaOH. For the preparation of suspensions with a pH of 0.5 or lower, the pH was adjusted using 1 M HCl.

[278] The suspensions were then allowed to equilibrate at room temperature (about 21 °C to about 23 °C) for 48 hours. After equilibration the solid phases were separated from the liquid phases by centrifugation, subsequently dried and then analyzed by XRPD. The concentration of deuterium-enriched (*R*)-pioglitazone or pioglitazone in the liquid phases (mother liquors) was determined by HPLC (see HPLC method described in Example 5).

[279] The solubility data collected for the four crystalline forms is shown in Table 11 and an overlay of the pH-dependent solubility profiles of the crystalline DCI and HCl salt forms of deuterium-enriched (*R*)-pioglitazone, crystalline DCI salt form of deuterium-enriched pioglitazone and crystalline pioglitazone HCl is presented in FIG. 14.

5 **[280]** The crystalline DCI and HCl salts of deuterium-enriched (*R*)-pioglitazone exhibited a similar pH-dependent solubility profile: the solubility was higher at very low pH values - 0.4 - 1.4, the maximum solubility was reached at pH 1.4 (4.54 mg/mL and 5.25 mg/mL for the DCI and HCl salts, respectively). A decrease in solubility was observed by increasing the pH thereafter. At pH 1.6, the solubility decreased to 2.58 mg/mL for the crystalline DCI salt of
10 deuterium-enriched (*R*)-pioglitazone, and to 3.47 mg/mL for the crystalline HCl salt of deuterium-enriched (*R*)-pioglitazone. At pH values of 2.4 and 3.0 the observed decrease in solubility was more significant (0.39 mg/mL and 0.20 mg/mL for the DCI and HCl salts, respectively). It was observed that both salts were practically insoluble above pH 4.

[281] In contrast, the maximum solubility for pioglitazone HCl was determined to be
15 4.00 mg/mL at pH 0.7. The solubility of pioglitazone HCl was found to decrease as the pH of the solution was increased: pH 0.6 - 2.88 mg/mL and pH 1.2 - 0.67 mg/mL. Further decreases in solubility were observed at higher pH values. Pioglitazone HCl was observed to be practically insoluble at pH values of greater than 3. The crystalline DCI salt form of deuterium-enriched pioglitazone presents behavior similar to that of pioglitazone HCl.

20 **[282]** For all four solid forms, in the pH range 0.3 - 0.8 there was no observed change in the solid form after equilibration. In the pH range 1.2 - 1.4, all three solid forms were observed to either partially or completely convert to the respective free-base form.

25 **Table 11. pH-Dependent Solubility Data for the Crystalline DCI and HCl Salt Forms of Deuterium-Enriched (*R*)-Pioglitazone and Crystalline Pioglitazone HCl**

Solvent	Initial Suspension pH	pH After Acid/Base Addition	Filtered Mother Liquor pH	Solubility (mg/mL)	Solid Phase After Equilibration
Deuterium-Enriched (<i>R</i>)-Pioglitazone DCI Salt					
Water	-	0.3	0.4	1.60	deuterium-enriched (<i>R</i>)-pioglitazone DCI salt
Water	1.4	0.5	0.6	3.01	deuterium-enriched (<i>R</i>)-pioglitazone DCI salt

Solvent	Initial Suspension pH	pH After Acid/Base Addition	Filtered Mother Liquor pH	Solubility (mg/mL)	Solid Phase After Equilibration
Water	-	0.8	0.8	3.30	deuterium-enriched (<i>R</i>)- pioglitazone DCl salt
Water	1.5	1.2	1.4	4.54	deuterium-enriched (<i>R</i>)- pioglitazone free base
Water	1.4	1.5	1.6	2.58	deuterium-enriched (<i>R</i>)- pioglitazone free base
Water	1.4	2.5	2.4	0.39	deuterium-enriched (<i>R</i>)- pioglitazone free base
Water	1.6	3.1	3.7	0.05	deuterium-enriched (<i>R</i>)- pioglitazone free base
Acetate Buffer	2.0	4.5	4.4	< 0.05	deuterium-enriched (<i>R</i>)- pioglitazone free base
Water	1.5	5.5	6.8	< 0.05	deuterium-enriched (<i>R</i>)- pioglitazone free base
Phosphate Buffer	2.3	6.6	6.5	< 0.05	deuterium-enriched (<i>R</i>)- pioglitazone
Phosphate Buffer	6.5	7.6	7.4	< 0.05	deuterium-enriched (<i>R</i>)- pioglitazone free base
Deuterium-Enriched (<i>R</i>)-Pioglitazone HCl Salt					
Water	1.6	0.3	0.5	2.33 ^a	deuterium-enriched (<i>R</i>)- pioglitazone HCl salt
Water	1.5	0.3	0.5	2.33 ^a	deuterium-enriched (<i>R</i>)- pioglitazone HCl salt
Water	1.5	0.5	0.5	2.33 ^a	deuterium-enriched (<i>R</i>)- pioglitazone HCl salt
Water	1.6	0.5	0.6	2.60	deuterium-enriched (<i>R</i>)- pioglitazone HCl salt
Water	-	0.3	0.7	3.27 ^a	deuterium-enriched (<i>R</i>)- pioglitazone HCl salt

Solvent	Initial Suspension pH	pH After Acid/Base Addition	Filtered Mother Liquor pH	Solubility (mg/mL)	Solid Phase After Equilibration
Water	1.4	0.5	0.7	3.27 ^a	deuterium-enriched (<i>R</i>)- pioglitazone HCl salt
Water	-	0.8	0.8	3.77 ^a	deuterium-enriched (<i>R</i>)- pioglitazone HCl salt
Water	1.4	0.8	0.8	3.77 ^a	deuterium-enriched (<i>R</i>)- pioglitazone HCl salt
Water	1.5	0.8	0.8	3.77 ^a	deuterium-enriched (<i>R</i>)- pioglitazone HCl salt
Water	1.3	1.3	1.4	5.25	deuterium-enriched (<i>R</i>)- pioglitazone HCl salt
Water	1.5	1.5	1.6	3.47	deuterium-enriched (<i>R</i>)- pioglitazone HCl salt
Water	1.6	2.5	3.0	0.20	deuterium-enriched (<i>R</i>)- pioglitazone HCl salt
Water	1.6	3.6	5.7	< 0.05	deuterium-enriched (<i>R</i>)- pioglitazone HCl salt
Pioglitazone HCl Salt					
Water	1.3	0.5	0.6	2.88	pioglitazone HCl salt
Water	-	0.3	0.7	4.00 ^a	pioglitazone HCl salt
Water	-	0.8	0.7	4.00 ^a	pioglitazone HCl salt
Water	1.2	1.2	1.2	0.67	pioglitazone HCl salt + pioglitazone free base
Water	1.3	1.5	1.6	0.44	pioglitazone HCl salt + pioglitazone free base
Water	1.2	2.5	3.0	0.03	pioglitazone free base
Water	1.2	3.5	4.6	< 0.03	pioglitazone free base
Acetate Buffer	1.8	4.5	4.5	< 0.03	pioglitazone free base
Water	1.2	5.5	5.6	< 0.03	pioglitazone free base

Solvent	Initial Suspension pH	pH After Acid/Base Addition	Filtered Mother Liquor pH	Solubility (mg/mL)	Solid Phase After Equilibration
Phosphate Buffer	2.0	6.6	6.6	< 0.03	pioglitazone free base
Phosphate Buffer	6.0	7.6	7.6	< 0.03	pioglitazone free base
Crystalline DCl salt form of deuterium-enriched pioglitazone					
Water	1.4-1.5	0.7	0.7	5.24	deuterium-enriched pioglitazone HCl salt
Water	1.4-1.5	1.0	0.9	2.49	deuterium-enriched pioglitazone HCl salt + deuterium-enriched pioglitazone free base
Water	1.4-1.5	1.2	1.1	1.63	deuterium-enriched pioglitazone HCl salt + deuterium-enriched pioglitazone free base
Water	1.4-1.5	1.5	1.6	0.42	deuterium enriched pioglitazone free base
Water	1.4-1.5-	3.0	3.1	<0.03	deuterium-enriched pioglitazone free base
Water	1.4-1.5	4.5	4.8	<0.03	deuterium-enriched pioglitazone free base

^a Averaged solubility value.

Example 7: Determination of Particle Size Distribution for the Crystalline DCl and HCl Salts of Deuterium-Enriched (*R*)-Pioglitazone

- 5 **[283]** The particle size distribution of the crystalline DCl and HCl salts of deuterium-enriched (*R*)-pioglitazone prepared using the methods described in Examples 2 and 3, respectively, were measured using a laser diffraction method. Samples of the crystalline DCl and HCl salts (100 mg – 200 mg) were wetted and dispersed in 30 mL of heptane. The experimental details are given in Table 12.

[284] Representative particle size distributions for the HCl (before and after milling) and DCl salts are given in Table 13.

[285] The crystalline HCl salt was milled using a fluidized air jet mill system or equivalent.

5

Table 12. Particle Size Distribution Analysis – Laser Diffraction Method

Materials and Equipment Used	
Equipment:	Beckman Coulter LS 13320
Dispersant:	Heptane and dispersant like Span 80 (2 drops) if needed
Operational Parameters:	
Optical model:	Fraunhofer approximation
Refractive index of dispersant:	1.39
Measures:	
Time of measure:	10 – 30 seconds
Speed agitation:	50 ± 2 %
Time measure of the background:	30 seconds
Obscuration range:	1 – 25 %

Table 13. Representative Particle Size Distributions of the Crystalline DCl and HCl Salts of Deuterium-Enriched (*R*)-Pioglitazone

Crystalline DCl salt	D10 = 64 µm D50 = 211 µm D90 = 662 µm	
Crystalline HCl salt	<u>Before Milling</u> D10 = 97 µm D50 = 323 µm D90 = 792 µm	<u>After Milling</u> D10 = 27 µm D50 = 139 µm D90 = 441 µm

10

Example 8: A Study to Assess the impact on Processability of Particle Size Distributions of the Crystalline HCl Salt of Deuterium-Enriched (*R*)-Pioglitazone (API)

[286] Study was conducted on the 15 mg strength PXL065 tablets

Table 14. Particle Size Distribution of 3 Micronized Batches of PXL065 Used in the Study:

	d10	d50	d90
Batch #1 with D90 <10µm	1µm	3µm	7µm
Batch #2 with D90 <20µm	1 µm	6µm	18µm
Batch #3 with D90 <100µm	2µm	17µm	57µm

- 5 **[287]** The tablet includes 15 mg of PXL065, as well as lactose, carmellose calcium, hyprolose, and magnesium stearate. The manufacturing process involves the steps of blending, lubrication, and compression. The particle size distributions of the 3 batches are shown in Table 14.

Flowability

- 10 **[288]** Blend manufactured with API with D90<10µm is stickier.

Blend Uniformity

Table 15: Blend Uniformity

Batch	Before lubrication	After lubrication
LF21026 (D90<10µm)	96.2% (SD=0.6)	96.2% (SD= 4.5)
LF21027 (D90<20µm)	98.4% (SD=0.7)	99.6% (SD=0.5)
LF21028 (D90<100µm)	99.1% (SD=0.8)	100.1% (SD=0.5)

Conclusion :

- 15 **[289]** Blend uniformity (shown in Table 15) is compliant (acceptance criteria = 90-110%) for all three batches. However, a lower uniformity value is obtained for blend manufactured with API with D90<10µm. Some losses during manufacture due to stickiness of the API might have occurred.

- [290]** Mean content uniformity values are <97% for batch LF21026, whereas >97% for
20 batches LF21027 and LF21028, which is consistent with the blend uniformity results.

Extensometric Study

[291] Objective: assess the evolution of tablet hardness (Critical Quality Attributes = CQA) depending on compression force and speed. Tablet hardness is shown in Table 16.

Conclusions

[292] As shown in FIG. 15, The lower the PSD of the API, the lower maximal hardness.

5 Tablet hardness is not specified, and is analyzed for information at release.

[293] As shown in FIG. 16, Tablets produced from blend with API with D90<10µm are more affected by variation of the tableting speed.

Table 16 -Tablet Hardness in Newtons

LF21026 <10µm		LF21027 <20µm		LF21028 <100µm	
Dwell-time (ms) 25rpm	41	Dwell-time (ms) 25rpm	39	Dwell-time (ms) 25rpm	38
Dwell-time (ms) 50rpm	10	Dwell-time (ms) 50rpm	8	Dwell-time (ms) 50rpm	8
Dwell-time (ms) 100rpm	4	Dwell-time (ms) 100rpm	4	Dwell-time (ms) 100rpm	4

10 **Example 9. Dog Pharmacokinetic (PK) Study**

[294] The objective of the study was to evaluate and compare the pharmacokinetic profiles of the deuterium-enriched (*R*)- and (*S*)-enantiomers of pioglitazone (PXL064 and PXL065) to deuterium-enriched pioglitazone (PXL061) after single administration by the oral route (capsule). The 3 Beagle male dogs were fasted before dosing. Then all test drugs were administered by the oral route using API in a capsule. The dosing schedule is shown in Table 17.

Table 17 Administration of Test Drugs for Dog PK Study

Day	Administration	Dose (mg/kg)
1	PXL065	5
4	PXL064	5
8	PXL061	10

[295] Food was given no sooner than one hour after dose administration on the days of treatment. Each administration was separated by a wash-out period of at least 2 days. Blood collection was performed for analysis of plasma concentrations and area under the curve (AUC) of the d-*R*-pioglitazone, d-*S*-pioglitazone, h-*R*-pioglitazone and h-*S*-pioglitazone

20

quantification at: Predose (just before dosing) and then T0.25h, T0.5h, T1h, T2h, T4h, T8h, T12h and T24h after dosing.

Table 18. Mean Plasma PK Parameters of Total Pioglitazone and Pioglitazone Enantiomers
5 Following Single Oral Administration of PXL065, PXL064, PXL061 or Pioglitazone in Dogs

		T_{max} (h)	C_{max} (ng/mL)	AUC₀₋₂₄ (ng.h/mL)	T_{1/2z} (h)
PXL065	Total R-pio	1 [1-12]	1330 ± 676	5770 ± 2500	1.43 ^a (1.42;1.43)
	Total S-pio	2 [2-12]	227 ± 56.7	1240 ± 665	1.62 ^a (1.58 ;1.65)
	Total Pio	1 [1-12]	1520 ± 696	7000 ± 3150	1.46 ^a (1.45 ;1.56)
PXL064	Total R-pio	2 [2-2]	54.0 ± 48.9	203 ± 169	1.91 ^a (1.55 ;2.26)
	Total S-pio	2 [0.5-2]	538 ± 391	1200 ± 852	1.75 ± 1.20
	Total Pio	2 [1-2]	588 ± 441	1410 ± 1020	1.73 ± 0.949
PXL061	Total R-pio	1 [0.5-2]	638 ± 849	2650 ± 3690	2.78 ± 0.669
	Total S-pio	1 [0.25-2]	683 ± 844	2150 ± 3090	3.61 ± 1.32
	Total Pio	1 [0.25-2]	1310 ± 1700	4800 ± 6780	4.03 ± 1.75

AUC0-24=area under the plasma concentration-time curve from 0 to 24 hours;

C_{max}=maximum plasma concentration; pio=pioglitazone; SD=standard deviation;

t_{1/2z}=terminal elimination phase half-life; t_{max}=time to maximum plasma concentration

Note: Table shows exposure to total pioglitazone (sum of deuterated and protonated (R)-
 10 and (S)-enantiomers of pioglitazone),

Note: Unless indicated otherwise, data are presented as mean ± SD values, except for t_{max}, which is presented as median [minimum, maximum] values.

Note: Data are shown for n=3, unless indicated otherwise.

a Data are presented for n=2, instead of n=3. Data are presented as mean (individual animal values).

Mean plasma PK parameters are shown in Table 18, and plasma exposures and relative
5 bioavailability are shown in Table 19.

The plasma PK profiles over time are shown in FIGS. 17-19.

**Table 19. Plasma Exposure (AUC_{0-24}) in Dogs of Total Pioglitazone and Relative
Bioavailability of Deuterium-enriched (*R*)-Pioglitazone (PXL065) and Deuterium-enriched
10 (*S*)-Pioglitazone (PXL064) Compared to Deuterium-Enriched Pioglitazone (PXL061)**

Total Pioglitazone	Animal No.	PXL065 5 mg/kg	PXL064 5 mg/kg	PXL061 10 mg/kg
AUC_{0-24} (ng.h/mL)	1001	3904	1034	12617
	1002	6904	2561	684
	1003	10193	625	1084
	Mean	7000	1410	4800
F_{rel} vs PXL061	1001	0.619	0.164	1
	1002	20.2	7.48	1
	1003	18.8	1.15	1
	Geo Mean	6.17	1.12	1

AUC_{0-24} =area under the plasma concentration-time curve from 0 to 24 hours; F_{rel} =relative oral bioavailability; geo=geometric; total pioglitazone (sum of deuterated and protonated (*R*)- and (*S*)-enantiomers of pioglitazone)

15 **[296]** The results from these studies support the fact that the increase in total pioglitazone (sum of protonated and deuterated (*R*)- and (*S*)-enantiomers) bioavailability seen with PXL065 compared to the deuterium-enriched pioglitazone (PXL061) and not seen with the deuterium-enriched (*S*)-enantiomer (PXL064), indicates that the enhanced bioavailability of PXL065 is not due to the presence of deuterium.

20

Example 10. 13-Week Dog Toxicology Study

[297] The objective of the study was to evaluate the potential toxicity and the toxicokinetics of deuterium-enriched (*R*)-pioglitazone (PXL065) following daily oral administration (capsule) to beagle dogs for 13 weeks in comparison to pioglitazone. Four

groups of beagle dogs each comprising three or five animals/sex, were administered 0 (empty capsule), 1.5, 5 or 15 mg/kg/day of PXL065 orally (API in one capsule/day) for 13 weeks; one other group was given 10 mg/kg/day of pioglitazone (API in one capsule/day). Blood was collected for toxicokinetic evaluations at various time-points (pre-dose, 0.25, 0.5, 1, 2, 4, 8, 12 and 24 hours post-dose) during the dosing period in order to determine the plasma concentrations of d-*R*-pioglitazone, d-*S*-pioglitazone, h-*R*-pioglitazone and h-*S*-pioglitazone.

[298] The full toxicological evaluation is not presented here, but relative bioavailability data based on total pioglitazone plasma concentrations at day 91 is shown in Table 20 and FIG. 20.

[299] Table 20: Mean Total Pioglitazone Exposure in Dogs on Day 91

Total pioglitazone exposure in dogs	AUC _{0-24, ss} (ng.h/mL)	AUC _{0-24, ss} dose normalized (ng.h/mL)	Relative Bioavailability (%)
Pioglitazone 10 mg/kg	2970 (M)	297	
	5620 (F)	562	
PXL065 5mg/kg	10900 (M)	2180	734
	8710 (F)	1762	310

AUC_{0-24,ss} = area under the plasma concentration-time curve from 0 to 24 hours at steady state; total pioglitazone (sum of deuterated and protonated (*R*)- and (*S*)-enantiomers of pioglitazone)

[300] The results from these studies show that there is an increased bioavailability to total pioglitazone (sum of protonated and deuterated (*R*)- and (*S*)-enantiomers) with PXL065 compared to pioglitazone.

Example 11: A Phase Ia Study to Assess the Safety, Tolerability, and Pharmacokinetics of Repeated Doses of the Crystalline HCl Salt of Deuterium-Enriched (*R*)-Pioglitazone (PXL065) in Healthy Human Subjects Compared to Pioglitazone HCl (Actos®)

Objectives

[301] The primary objective of this study was to assess the safety and tolerability, in healthy subjects, of single oral doses of PXL-065 (7.5 mg, 22.5, and 30 mg as capsules).

A secondary objective of this study was to assess the PK of PXL065 in healthy subjects after single doses of PXL065 compared to Actos[®], 45 mg, with a specific focus on assessing the exposures to both deuterated and protonated (*R*)- and (*S*)-pioglitazone to determine the extent, if any, of interconversion of PXL065 to the protonated (*R*)- and (*S*)-enantiomers.

5

Study Methodology

[302] This study was conducted in two parts. Part 1 utilized an open-label, parallel-group design. Within 21 days of screening, eligible subjects were admitted to the clinical study unit. On Study Day 1, subjects were randomly allocated to receive a single dose of PXL065 22.5 mg (6 subjects) or a single dose of Actos[®] 45 mg (6 subjects). Doses of study medication were administered at approximately 8 AM on Day 1 in a fasted state. Subjects remained in the clinical unit for 36 hours post-dose administration. Subjects returned to the clinic as out-patients on Days 4 and 7 for follow-up assessments. Following a review of safety and tolerability of PXL065 in Part 1 by a Data Review Committee (DRC) and determination of comparative PK exposure to enantiomers, 6 healthy subjects were enrolled into Part 2.

[303] Part 2 utilized an open-label design in which a single dose of PXL065 7.5 mg was administered in the morning of Day 1 in a fasted state. Subjects remained in the clinical unit for at least 36 hours post-dose and returned to the clinic as out-patients on Days 4 and 7 for follow-up assessments. Following review of safety, tolerability, and PK data from the PXL065 7.5 mg dose group, an additional treatment group (PXL065 30 mg) of 6 healthy subjects was evaluated. Before dosing the additional group, a review of the safety and tolerability from the preceding group was performed by the DRC.

Diagnosis and Criteria for Inclusion

[304] Subjects were healthy adult male or female, 18-40 years of age (inclusive), with a body mass index (BMI) of ≥ 17 to ≤ 32 kg/m². Female subjects were not pregnant or breastfeeding.

30

Test Product, Dose, and Mode of Administration

[305] PXL065, orally administered. Dose = 1 x 7.5 mg capsule, 1 x 22.5 mg capsule, or 1 x 30 mg capsule. The PXL065 capsule includes 7.5 mg, 22.5 mg, or 30 mg of PXL065, as well as lactose.

Control Product, Dose, and Mode of Administration

[306] Actos[®], orally administered. Dose = 1 x 45 mg, tablet.

Actos[®] was purchased as the branded product. A 45 mg tablet includes 45 mg pioglitazone HCl salt, lactose, carmellose calcium, hypromellose, and magnesium stearate

5 Duration of Treatment

[307] In Part 1, a single dose of PXL065 22.5 mg or a single dose of Actos[®] 45 mg was administered to each subject and they were evaluated for 7 days postdose. In Part 2, a single dose of PXL065 7.5 mg or 30 mg was administered to each subject and they were evaluated for 7 days postdose. The total duration of the study from the time of
10 confinement (Day -1) through the end of study visit is 8 days.

Criteria for Evaluation

[308] Safety: The Investigator evaluated safety using the following assessments: physical examinations, electrocardiograms (ECGs), vital sign measurements, clinical laboratory evaluations, and reported or observed adverse events (AEs). Subjects were
15 monitored for any AEs from the beginning of confinement through the end of the study. Pharmacokinetics: Plasma PK parameters including, but not limited to $t_{1/2}$, t_{max} , C_{max} , AUC_{0-last} and AUC_{0-inf} were calculated for the deuterated and protonated forms of (R)- and (S)-pioglitazone.

Statistical Methods

20 [309] In general, all data were summarized with descriptive statistics (number of subjects, means, standard deviations, minimums, medians, and maximums) for continuous endpoints, and frequency and percentage for categorical endpoints. Within each part of the clinical study, data was presented separately for each cohort and by treatment group within each cohort.

25 [310] Safety and tolerability: All safety and tolerability data were listed. In the case of continuous variables, descriptive statistics was used to summarize the results and changes from baseline by treatment and time point.

[311] The values of categorical assessments were tabulated. Adverse Events (AEs) were coded according to MedDRA.

30 Results

As shown in Table 21 and FIG. 21, PXL065 has improved bioavailability after single dose administration in healthy human subjects compared to Actos[®].

Table 21: Total Pioglitazone Exposure in Phase Ia Study in Humans

	AUC _{inf} (h.ng/mL)	AUC _{inf} dose normalized (h.ng/mL/mg)	Relative bioavailability (%)
Actos [®] 45 mg	15981	355	
PXL065 22.5 mg	13896	618	174

AUC_{inf} = area under the plasma concentration-time curve from time 0 to infinity; total pioglitazone (sum of deuterated and protonated (*R*)- and (*S*)-enantiomers of pioglitazone)

Example 12 A Phase Ib Study to Assess the Safety, Tolerability, and Pharmacokinetics of

5 Repeated Doses of the Crystalline HCl Salt of Deuterium-Enriched (*R*)-Pioglitazone (PXL065) in Healthy Human Subjects Compared to Pioglitazone HCl (Actos[®])

Objectives

[312] The primary objective of this study was to assess the safety and tolerability in healthy subjects of repeated administrations of different doses of PXL065 compared to
10 Pioglitazone HCl (Actos[®]) 45 mg.

[313] A secondary objective of this study was to assess the PK of PXL065 in healthy subjects after single and multiple administrations compared to Actos[®], 45 mg, with a specific focus on assessing the exposures to both deuterated and protonated (*R*)- and (*S*)-enantiomers of pioglitazone to determine the extent, if any, of interconversion of PXL065
15 to the protonated (*R*)- and (*S*)-enantiomers.

Study Methodology

[314] Subjects were screened within 28 days before their first dose of study medication. They stayed in the clinic from 1 day before dosing (Day -1) until at least 48 hours after their final dose of trial medication (Day 11). Subjects returned to the clinic for
20 outpatient assessments and/or PK sampling on Day 12 (approximately 72 hours postdose), Day 13 (approximately 96 hours postdose), Day 14 (approximately 120 hours postdose), Day 15 (approximately 144 hours postdose), and for a follow up visit at 10 days (± 2 days) after their final dose.

[315] Up to 30 healthy subjects were enrolled in this trial:

25

- Three (3) PXL065 dose groups: 8 subjects (6 on active and 2 on placebo) in each dose group. Each dose group was composed of 4 males and 4 females in a 3:1 active/placebo ratio.
- One (1) Actos[®] group: 6 subjects (3 males and 3 females) were given Actos[®]. Placebo was not given to any subjects in this group.

[316] The dose levels in Groups 1-4 were as follows (Table 22):

Table 22. Dose Levels in Groups 1-4

Group	PXL065	Actos®
1	-	45
2	7.5 mg	-
3	15 mg	-
4	30 mg	-

5 [317] Subjects received a single oral dose of study drug (7.5, 15, or 30 mg tablets of PXL065 or matching placebo tablets, or 45 mg Actos® tablets), under fasting conditions on Day 1. They then received repeated administrations of each study drug for 7 days from Day 3 to Day 9. All study drugs were dosed in fasting conditions.

[318] The PXL065 PK parameters of the three single oral doses on Day 1 and after seven
10 daily administrations on Day 9 were compared to the Actos® PK parameters on Day 1 and after seven daily administrations on Day 9.

Diagnosis and Criteria for Inclusion

[319] Subjects were healthy adult male or female, 18-45 years of age (inclusive), with a body mass index (BMI) of ≥ 18.5 to ≤ 32 kg/m² and a body weight of ≥ 60 kg at screening.
15 Female subjects were not pregnant or breastfeeding.

Test Product, Dose, and Mode of Administration

[320] PXL065, orally administered. Dose = 1 x 7.5 mg tablet, 1 x 15 mg tablet, or 1 x 30 mg tablet.

[321] The PXL065 tablet include 7.5 mg, 15 mg, or 30 mg of PXL065, as well as lactose, carmellose calcium, hypolose, and magnesium stearate.
20

Control Product, Dose, and Mode of Administration

[322] Actos®, orally administered. Dose = 1 x 45 mg, tablet.

[323] Actos® was purchased as the branded product. A 45 mg tablet includes 45 mg pioglitazone HCl salt, lactose, carmellose calcium, hypolose, and magnesium stearate.

25 Duration of Treatment

[324] One single oral administration of PXL065 or matching placebo or Actos® was given on Day 1 in fasting conditions followed by once daily multiple oral administrations of PXL065 or matching placebo or Actos® from Day 3 to Day 9 (7 days) in fasting conditions.

Criteria for Evaluation

[325] Pharmacokinetics: Plasma samples were analyzed using a validated assay. The samples from all evaluable subjects were analyzed. Samples from subjects who experienced emesis within 4 hours of dosing were not analyzed.

[326] Plasma PK parameters included, but were not limited to:

- Following single dose on Day 1 and Day 3 (for PXL065 groups): $t_{\max}$, t_{lag} , $C_{\max}$, $AUC_{0-\text{last}}$, AUC_{0-24} , $AUC_{0-\text{inf}}$, λ_z , $t_{1/2}$, $\%AUC_{\text{ext}}$, CL/F , V_z/F , $C_{\max}/\text{Dose}$, $AUC_{0-\text{last}}/\text{Dose}$, $AUC_{0-\text{inf}}/\text{Dose}$ and MRT each calculated using compartmental or non-compartmental methods as appropriate for the deuterated and protonated (*R*)- and (*S*)-enantiomers of pioglitazone.
- Day 4 to Day 9: C_{trough} for the deuterated and protonated (*R*)- and (*S*)-enantiomers of pioglitazone; and
- Following repeated doses on Day 9: $t_{\max}$, $C_{\max}$, C_{avg} , t_{lag} , $AUC_{0-\text{last}}$, AUC_{0-24} , $AUC_{0-\text{inf}}$, λ_z , $t_{1/2}$, $\%AUC_{\text{ext}}$, CL_{ss}/F , $V_{z\text{ss}}/F$, MRT, PTF, $R_{\text{ac}}(AUC_{0-24})$, $R_{\text{ac}}(C_{\max})$, $SR(AUC)$, $C_{\max}/\text{Dose}$, $AUC_{0-\text{last}}/\text{Dose}$, $AUC_{0-\text{inf}}/\text{Dose}$, AUC_{0-24}/Dose .

Safety

[327] The Investigator evaluated safety using the following assessments: physical examinations, vital sign measurements, clinical laboratory evaluations, electrocardiograms, and reported or observed adverse events (AEs). Subjects were monitored for any AEs from the signing of the informed consent through the end of the study.

Statistical Methods

[328] In general, all data was summarized with descriptive statistics (number of subjects, means, standard deviations, minimums, medians, and maximums) for continuous endpoints, and frequency and percentage for categorical endpoints. Within each part of the clinical study, data was presented separately for each cohort and by treatment group within each cohort.

[329] Safety and tolerability: all safety and tolerability data were listed. In the case of continuous variables, descriptive statistics was used to summarize the results and changes from baseline by treatment and time point. The values of categorical assessments were tabulated. Adverse Events (AEs) were coded according to MedDRA.

Results

[330] Pharmacokinetic results: the relative bioavailability in humans of PXL065 increased by 55% - 65% compared to Actos® (see Table 23 and FIG. 18).

Table 23. Pharmacokinetic Parameters for PXL065 and Actos® in Humans

Total Pioglitazone Exposure in Humans	AUC_{0-24,ss} (h*ng/mL)	AUC_{0-24,ss} dose normalized (h*ng/mL/mg)	Relative Bioavailability (%)
Actos® 45 mg	12567	279	-
PXL065 15 mg	6505	434	155
PXL065 30 mg	13802	460	165

As shown in Table 23 and FIG. 22, PXL065 has improved bioavailability after repeated dose
 5 administrations in healthy human subjects compared to Actos®.

[331] Combined, the results from the two dog and two human PK studies described
 above in Examples 9-12 show that bioavailability to total pioglitazone (sum of protonated
 and deuterated (*R*)- and (*S*)-enantiomers) with PXL065 compared to pioglitazone is
 10 independent of the formulation (capsule or tablet).

INCORPORATION BY REFERENCE

[332] This application refers to various issued patents, published patent applications,
 journal articles, and other publications, all of which are incorporated herein by reference. If
 15 there is a conflict between any of the incorporated references and the instant specification,
 the specification shall control. In addition, any particular embodiment of the present
 disclosure that falls within the prior art may be explicitly excluded from any one or more of
 the claims. Because such embodiments are deemed to be known to one of ordinary skill in
 the art, they may be excluded even if the exclusion is not set forth explicitly herein. Any
 20 particular embodiment of the disclosure can be excluded from any claim, for any reason,
 whether or not related to the existence of prior art.

EQUIVALENTS


The various described embodiments of the invention may be used in conjunction
 25 with one or more other embodiments unless technically incompatible.

[333] The invention may be embodied in other specific forms without departing from
 the spirit or essential characteristics thereof. The foregoing embodiments are therefore to

be considered in all respects illustrative rather than limiting the invention described herein. Scope of the invention is thus indicated by the appended claims rather than by the foregoing description, and all changes that come within the meaning and range of equivalency of the claims are intended to be embraced therein.

CLAIMS:

1. A deuterium chloride salt of a deuterium-enriched compound of formula (I):




- 5 wherein R^1 is H or D, provided that the total deuterium abundance in the compound of formula (I) is about 1.3 to about 4.

2. The deuterium chloride salt of claim 1, wherein the deuterium chloride salt is a crystalline deuterium chloride salt.

10

3. The deuterium chloride salt of claim 1 or 2, wherein the crystalline deuterium chloride salt is an anhydrous crystalline deuterium chloride salt.

4. A crystalline deuterium chloride salt of a deuterium-enriched compound of formula (I)



15

- wherein R^1 is H or D, provided that the total deuterium abundance in the compound of formula (I) is about 1.3 to about 4, and

- wherein the crystalline deuterium chloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): $15.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, and $26.0^\circ \pm 0.2^\circ$.

20

5. The crystalline deuterium chloride salt of claim 4, wherein the crystalline deuterium chloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): $8.6^\circ \pm 0.2^\circ$, $8.8^\circ \pm 0.2^\circ$, $12.8^\circ \pm 0.2^\circ$, $12.9^\circ \pm 0.2^\circ$, $15.8^\circ \pm 0.2^\circ$, $18.8^\circ \pm 0.2^\circ$, $18.9^\circ \pm 0.2^\circ$, $19.7^\circ \pm 0.2^\circ$, $20.0^\circ \pm 0.2^\circ$, $20.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, $26.0^\circ \pm 0.2^\circ$, and $31.3^\circ \pm 0.2^\circ$.


25

6. The crystalline deuterium chloride salt of any one of claims 4-5, wherein the crystalline deuterium chloride salt is characterized by an X-ray powder diffraction pattern substantially the same as shown in FIG. 1A.

5 7. The crystalline deuterium chloride salt of any one of claims 4-6, wherein the crystalline deuterium chloride salt has a melting point onset as determined by differential scanning calorimetry at about 190 °C to about 200 °C.

8. The crystalline deuterium chloride salt of any one of claims 4-7, wherein the crystalline
10 deuterium chloride salt is an anhydrous crystalline deuterium chloride salt.

9. A pharmaceutical material comprising particles of a crystalline deuterium chloride salt of a deuterium-enriched compound of formula (I)




15 wherein R¹ is H or D, provided that the total deuterium abundance in the compound of formula (I) is about 1.3 to about 4, and

wherein the particles in the composition have a crystal shape selected from hexagonal, rod, and combinations thereof.

20 10. The pharmaceutical material of claim 9, wherein the particles have a particle size distribution which is defined by a d(0.9) of about 10 μm to about 800 μm preferably below 500 μm.

11. A pharmaceutical material comprising particles of a crystalline deuterium chloride salt
25 of a deuterium-enriched compound of formula (I)



wherein R^1 is H or D, provided that the total deuterium abundance in the compound of formula (I) is about 1.3 to about 4, and

wherein the particles have a particle size distribution which is defined by a $d(0.9)$ of about 10 μm to about 800 μm preferably below 500 μm .

5

12. The pharmaceutical material of any one of claims 9-11, wherein the crystalline deuterium chloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): $15.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, and $26.0^\circ \pm 0.2^\circ$.


10

13. The pharmaceutical material of any one of claims 9-12, wherein the crystalline deuterium chloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): $8.6^\circ \pm 0.2^\circ$, $8.8^\circ \pm 0.2^\circ$, $12.8^\circ \pm 0.2^\circ$, $12.9^\circ \pm 0.2^\circ$, $15.8^\circ \pm 0.2^\circ$, $18.8^\circ \pm 0.2^\circ$, $18.9^\circ \pm 0.2^\circ$, $19.7^\circ \pm 0.2^\circ$, $20.0^\circ \pm 0.2^\circ$, $20.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, $26.0^\circ \pm 0.2^\circ$, and $31.3^\circ \pm 0.2^\circ$.

15

14. The pharmaceutical material of any one of claims 9-13, wherein the crystalline deuterium chloride salt is an anhydrous crystalline deuterium chloride salt.

15. A crystalline hydrochloride salt of a compound of formula (I-A):



20


16. The crystalline hydrochloride salt of claim 15, wherein the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): $15.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, and $26.0^\circ \pm 0.2^\circ$.

25

17. The crystalline hydrochloride salt of claim 15, wherein the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): $20.0^\circ \pm 0.2^\circ$, $20.8^\circ \pm 0.2^\circ$, and $22.8^\circ \pm 0.2^\circ$.

30

18. A crystalline hydrochloride salt of the compound of formula (I-A)



wherein the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): $15.8^\circ \pm 0.2^\circ$, $20.0^\circ \pm 0.2^\circ$, $20.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, and $26.0^\circ \pm 0.2^\circ$.

5

19. The crystalline hydrochloride salt of any one of claims 15-18, wherein the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): $8.6^\circ \pm 0.2^\circ$, $8.8^\circ \pm 0.2^\circ$, $12.8^\circ \pm 0.2^\circ$, $12.9^\circ \pm 0.2^\circ$, $15.8^\circ \pm 0.2^\circ$, $18.8^\circ \pm 0.2^\circ$, $19.7^\circ \pm 0.2^\circ$, $20.0^\circ \pm 0.2^\circ$, $20.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, $26.0^\circ \pm 0.2^\circ$, $28.1^\circ \pm 0.2^\circ$, and $31.3^\circ \pm 0.2^\circ$.

10

20. The crystalline hydrochloride salt of any one of claims 15-19, wherein the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern substantially the same as shown in FIG. 6.


15

21. The crystalline hydrochloride salt of any one of claims 15-20, wherein the crystalline hydrochloride salt has a melting point onset as determined by differential scanning calorimetry at about 190°C to about 210°C .

20

22. The crystalline hydrochloride salt of any one of claims 15-21, wherein the crystalline hydrochloride salt is an anhydrous crystalline hydrochloride salt.

23. A pharmaceutical material comprising particles of a crystalline hydrochloride salt of the compound of formula (I-A)




25

wherein the particles in the composition have a needle-like crystal shape.

24. The pharmaceutical material of claim 23, wherein the particles have a particle size distribution which is defined by a d(0.9) of about 150 μm to about 800 μm , preferably below 500 μm .

- 5 25. A pharmaceutical material comprising particles of a crystalline hydrochloride salt of the compound of formula (I-A)



wherein the particles have a particle size distribution which is defined by a d(0.9) of about 10 μm to about 800 μm , preferably below 500 μm .

10

26. The pharmaceutical material of claim 25, wherein the particles in the composition have a needle-like crystal shape.

- 15 27. The pharmaceutical material of any one of claims 23-26, wherein the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern comprising peaks at the following diffraction angles (2θ): $15.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, and $26.0^\circ \pm 0.2^\circ$.

28. The pharmaceutical material of any one of claims 23-27, wherein the crystalline hydrochloride salt is characterized by an X-ray powder diffraction pattern comprising peaks
20 at the following diffraction angles (2θ): $8.6^\circ \pm 0.2^\circ$, $8.8^\circ \pm 0.2^\circ$, $12.8^\circ \pm 0.2^\circ$, $12.9^\circ \pm 0.2^\circ$, $15.8^\circ \pm 0.2^\circ$, $18.8^\circ \pm 0.2^\circ$, $19.7^\circ \pm 0.2^\circ$, $20.0^\circ \pm 0.2^\circ$, $20.8^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, $26.0^\circ \pm 0.2^\circ$, $28.1^\circ \pm 0.2^\circ$, and $31.3^\circ \pm 0.2^\circ$.

29. The pharmaceutical material of any one of claims 23-28, wherein the crystalline
25 hydrochloride salt is an anhydrous crystalline hydrochloride salt.

30. A pharmaceutical composition comprising a deuterium chloride salt of any one of claims 1-3, a crystalline deuterium chloride salt of any one of claims 4-8, or a pharmaceutical material of any one of claims 9-14, and a pharmaceutically acceptable
30 excipient.

31. A pharmaceutical composition comprising a crystalline hydrochloride salt of any one of claims 15-22, or a pharmaceutical material of any one of claims 23-29, and a pharmaceutically acceptable excipient.

5

32. A method of treating a metabolic disorder in a patient in need thereof, the method comprising administering to the patient an effective amount of a deuterium chloride salt of any one of claims 1-3, a crystalline deuterium chloride salt of any one of claims 4-8, a pharmaceutical material of any one of claims 9-14, or a pharmaceutical composition of claim 30.

10

33. The method of claim 32, wherein the metabolic disorder is polycystic ovary syndrome.

15

34. A method of treating diabetes mellitus type 2 in a patient in need thereof, the method comprising administering to the patient an effective amount of a deuterium chloride salt of any one of claims 1-3, a crystalline deuterium chloride salt of any one of claims 4-8, a pharmaceutical material of any one of claims 9-14, or a pharmaceutical composition of claim 30.

20

35. A method of treating nonalcoholic steatohepatitis in a patient in need thereof, the method comprising administering to the patient an effective amount of a deuterium chloride salt of any one of claims 1-3, a crystalline deuterium chloride salt of any one of claims 4-8, a pharmaceutical material of any one of claims 9-14, or a pharmaceutical composition of claim 30.

25

36. A method of treating nonalcoholic fatty liver disease in a patient in need thereof, the method comprising administering to the patient an effective amount of a deuterium chloride salt of any one of claims 1-3, a crystalline deuterium chloride salt of any one of claims 4-8, a pharmaceutical material of any one of claims 9-14, or a pharmaceutical composition of claim 30.

30

37. A method of treating a neurological disorder in a patient in need thereof, the method comprising administering to the patient an effective amount of a deuterium chloride salt of

any one of claims 1-3, a crystalline deuterium chloride salt of any one of claims 4-8, a pharmaceutical material of any one of claims 9-14, or a pharmaceutical composition of claim 30.

5 38. The method of claim 37, wherein the neurological disorder is adrenoleukodystrophy or adrenomyeloneuropathy.

39. A method of treating a metabolic disorder in a patient in need thereof, the method comprising administering to the patient an effective amount of a crystalline hydrochloride
10 salt of any one of claims 15-22, a pharmaceutical material of any one of claims 23-29, or a pharmaceutical composition of claim 31.

40. The method of claim 39, wherein the metabolic disorder is polycystic ovary syndrome.
15

41. A method of treating diabetes mellitus type 2 in a patient in need thereof, the method comprising administering to the patient an effective amount of a crystalline hydrochloride salt of any one of claims 15-22, a pharmaceutical material of any one of claims 23-29, or a pharmaceutical composition of claim 31.
20

42. A method of treating nonalcoholic steatohepatitis in a patient in need thereof, the method comprising administering to the patient an effective amount of a crystalline hydrochloride salt of any one of claims 15-22, a pharmaceutical material of any one of claims 23-29, or a pharmaceutical composition of claim 31.
25

43. A method of treating nonalcoholic fatty liver disease in a patient in need thereof, the method comprising administering to the patient an effective amount of a crystalline hydrochloride salt of any one of claims 15-22, a pharmaceutical material of any one of claims 23-29, or a pharmaceutical composition of claim 31.
30

44. A method of treating a neurological disorder in a patient in need thereof, the method comprising administering to the patient an effective amount of a crystalline hydrochloride salt of any one of claims 15-22, a pharmaceutical material of any one of claims 23-29, or a pharmaceutical composition of claim 31.

45. The method of claim 44, wherein the neurological disorder is adrenoleukodystrophy or adrenomyeloneuropathy.




FIG. 1A




FIG. 1B




FIG. 2




FIG. 3




FIG. 4A




FIG. 4B




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9A




FIG. 9B




FIG. 10.




FIG. 11




FIG. 12




FIG. 13A




FIG. 13B




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5A





FIG. 5B












FIG. 8

FIG. 7










FIG. 9H






FIG. 10C




FIG. 10D




FIG. 11




FIG. 12





FIG. 13






FIG. 15