(12) PATENT **(11)** Application No. **AU 199656287 B2**
(19) AUSTRALIAN PATENT OFFICE **(10)** Patent No. **707350**

(54) Title
Coccidiosis poultry vaccine

(51)[6] International Patent Classification(s)
C07K 014/455 **C07K 016/20**
A61K 039/012 **C12N 015/30**

(21) Application No: **199656287** (22) Application Date: **1996.07.02**

(30) Priority Data

(31) Number	(32) Date	(33) Country
95201801	**1995.07.03**	**EP**

(43) Publication Date : **1997.01.16**
(43) Publication Journal Date : **1997.01.16**
(44) Accepted Journal Date : **1999.07.08**

(71) Applicant(s)
Akzo Nobel N.V.

(72) Inventor(s)
Jacobus Johannes Kok; Paul van den Boogaart; Arnoldus Nicolaas Vermeulen

(74) Agent/Attorney
SPRUSON and FERGUSON,GPO Box 3898,SYDNEY NSW 2001

AU9656287

(12) PATENT ABSTRACT (11) Document No. AU-A-56287/96
(19) AUSTRALIAN PATENT OFFICE

(54) Title
COCCIDIOSIS POULTRY VACCINE

International Patent Classification(s)
(51)[6] **C07K 014/455** **C07K 016/20** **C12N 015/30** **A61K 039/012**

(21) Application No. : **56287/96** (22) Application Date : **02.07.96**

(30) Priority Data

(31) Number	(32) Date	(33) Country
95201801	**03.07.95**	**EP EUROPEAN PATENT OFFICE (EP)**

(43) Publication Date : **16.01.97**

(71) Applicant(s)
AKZO NOBEL N.V.

(72) Inventor(s)
JACOBUS JOHANNES KOK; PAUL VAN DEN BOOGAART; ARNOLDUS NICOLAAS VERMEULEN

(74) Attorney or Agent
SPRUSON & FERGUSON , GPO Box 3898, SYDNEY NSW 2001

(57)

This invention relates to novel *Eimeria* proteins with immunogenic properties as well as to DNA sequences encoding these proteins. These proteins can be administered to poultry thereby protecting the birds against coccidiosis. In addition the DNA encoding these proteins can be used for the preparation of a vector vaccine against coccidiosis.

Coccidiosis Poultry Vaccine

Abstract

This invention relates to novel *Eimeria* proteins with immunogenic properties as well as to DNA sequences encoding these proteins. These proteins can be administered to poultry thereby protecting the birds against coccidiosis. In addition the DNA encoding these proteins can be used for the preparation of a vector vaccine against coccidiosis.

S & F Ref: 344847

AUSTRALIA

PATENTS ACT 1990

COMPLETE SPECIFICATION

FOR A STANDARD PATENT

ORIGINAL

Name and Address of Applicant: Akzo Nobel N.V.
Velperweg 76
6824 BM Arnhem
THE NETHERLANDS

Actual Inventor(s): Jacobus Johannes Kok, Paul van den Boogaart and Arnoldus Nicolaas Vermeulen

Address for Service: Spruson & Ferguson, Patent Attorneys
Level 33 St Martins Tower, 31 Market Street
Sydney, New South Wales, 2000, Australia

Invention Title: Coccidiosis Poultry Vaccine

The following statement is a full description of this invention, including the best method of performing it known to me/us:-

Coccidiosis Poultry Vaccine

The present invention relates to a protein derived from *Eimeria acervulina*, which is capable of stimulating immune lymphocytes. It also relates to a nucleic acid sequence encoding all or an antigenically significant part of this protein, a recombinant vector comprising such a nucleic acid sequence, a host cell or organism transformed with such a recombinant vector and a vaccine for the protection of poultry against coccidiosis.

Coccidiosis is a disease caused by infection with one or more of the many species of coccidia, intracellular protozoal parasites of the subphylum *Apicomplexa* and the genus *Eimeria*. Poultry is defined herein as domesticated birds that serve as a source of eggs or meat and that include such commercially important kinds as chickens, turkeys, ducks, geese, guinea fowl, pheasants, pigeons and peafowl.

Coccidiosis in chickens is known to be caused by several different species of *Eimeria*, namely *Eimeria acervulina*, *E. maxima*, *E. tenella*, *E. necatrix*, *E. brunetti*, *E. mitis*, *E. praecox*, *E. mivati* and *E. hagani*. Some people, however, doubt the true existence of the last two species. Low level infection with any of these *Eimeria* species results in a protective immunity to reinfection.

The species do differ in their pathogenic effect on chickens, the type of chicken also playing a role; thus, a broiler chicken will be subjected to a great deal of damage by a parasite such as *E. acervulina* or *E. maxima* because these parasitise large portions of the small intestine, where food digestion plays a major role.

E. acevulina is one of the most common species found in the litter of broiler houses in both Europe and the USA. It has a great reproductive potential and is regarded as pathogenic because it produces a marked depression in gain of body weight, higher feed conversion and it produces gross lesions in the upper small intestine.

During the life cycle (see also Table 1), the *Eimeria* parasite passes through a number of stages. The life cycle begins when the chicken ingests the infectious stage, known as the sporulated oocyst, during ground feeding or by inhalation of dust. The wall of the sporulated oocyst is ruptured by a combination of mechanical grinding action and chemical action in the gizzard and intestinal tract, resulting in the release of four sporocysts. The sporocysts pass into the duodenum where they are exposed to bile and digestive enzymes resulting in the release of two sporozoites per sporocyst.

Table 1.

Endogenous stages of *Eimeria acervulina* in stained sections of infected duodenum (after McDonald V. *et al.*, Parasitol. 8, 21-30, 1982).

Time of infection	Histological observations
24h	Immature 1st generation asexual stages
36h	Semi-mature 1st generation schizonts
42h	Mature 1st gen. schizonts. Immature 2nd gen. parasites
48h	Mature 2nd gen. schizonts. A few 3rd gen. schizonts with 8-16 merozoites
60h	Mature 3rd gen. schizonts, immature 4th gen. parasites

The sporozoites are mobile and search for suitable host epithelium cells in order to penetrate and reproduce in them. Following infection of an epithelium cell, the parasite enters the schizont phase of its life cycle, producing from 8 to 16 to >200 merozoites per schizont. Once released from the schizont, the merozoites are free to infect further epithelium cells. After from two to five of these asexual reproduction cycles, the intracellular merozoites grow into sexual forms known as the female or macrogametocyte and the male or microgametocyte. Following fertilisation of the macrogametocyte by the microgametes released from the microgametocyte, a zygote is formed which creates a cyst wall about itself. The newly formed oocyst is passed out of the infected chicken with the droppings.

With the correct environmental conditions of temperature and humidity and sufficient oxygen in the air, the oocyst will sporulate into the infectious stage, ready to infect a new host and thereby spreading the disease. Thus no intermediate host is required for transfer of the parasite from bird to bird.

The result of the *Eimeria* parasite infecting the digestive tract of a chicken may be a reduction in weight gain, increased feed conversion, cessation of egg production and, in some cases, death. The increase in intensive production of poultry has been accompanied by severe losses due to this parasite; indeed, coccidiosis has become the most economically important parasitic disease. In the Netherlands, the losses that poultry farmers suffer every year run into millions of guilders; in 1986 the loss was about 13 million guilders. In the same year, a loss of 300 million dollars was suffered in the United States.

In the past, several methods have been used in attempts to control coccidiosis. Prior to the advent of chemotherapeutic agents, improved sanitation using disinfectants, together with the mechanical removal of litter, was the main method employed; sufficient oocysts, however, usually remained to transmit the disease.

The introduction of coccidiostatic agents in the feed or drinking water, in addition to good management, resulted in some success at disease control. Such agents have been found to suffer from a drop in effectiveness over the years, due partly to the development of drug resistant strains of coccidia. Furthermore, several chemotherapeutic agents have been found to leave residues in the meat, making it unsuitable for consumption.

Attempts have been made to control the disease immunologically by administering to chickens a live vaccine comprising oocysts from all seven species of *Eimeria*, the oocysts administered being from precocious lines. Such precocious lines are obtained by inoculating chickens with a wild population of an *Eimeria* species and collecting the very first parasites that are excreted as a result of the infection. The collected parasites are put back into chickens and the cycle is repeated several times. Eventually a precocious line of parasite is produced which has fewer cycles of asexual reproduction in the gut. Thus such lines retain their immunogenicity, whilst producing fewer parasites in the gut with less consequential damage being caused to the host chicken.

The disadvantage of this type of vaccine is that it is expensive to produce because of the necessity of producing it in live chickens and its lower reproductive potential.

The advent of genetic engineering has provided new methods for producing effective vaccines. Using these methods, the DNA coding for the antigenic proteins of some pathogenic microorganisms has been cloned into such host microorganisms as *Escherichia coli* or *Salmonella* spec., with the result that the protein has been expressed at sufficiently high levels such that it can be incorporated into a vaccine. The advantage of proteins produced in this way is that they are noninfectious and are relatively cheap to produce. In this way, vaccines have been prepared against a number of viruses such as hepatitis, herpes simplex and foot and mouth disease.

Attempts have been made to genetically engineer a coccidiosis vaccine. EP 337 589 describes the isolation of a Group B *Eimeria tenella* protein and its insertion into a novel expression vector which, in turn, has been used to transform appropriate hosts. WO 92/04461 describes the construction of a microorganism that produces an antigenic protein using either the "mRNA route" or the "nuclear DNA route". In this way, certain antigens from *E. tenella* and *E. maxima* were prepared and sequenced. Taking this type of route to prepare antigens for incorporation into a vaccine relies only upon selecting antigens which could induce antibodies in an heterologous species. This approach does not necessarily end up with selecting the most protective antigen.

From H.S. Lillehoj (Vet. Immunol. Immunopath., **13**, 321-330, 1986) it can be conceived that development of protective immunity in chickens infected with coccidia may be due to the development of a species-specific T cell response.

It has now been found that a very immunogenic protein can be isolated from the 42h developmental stage of *Eimeria* schizonts. Surprisingly, this protein is found intracellularly in *Eimeria* and it appears to contain high sequence homology with known heterologous lactate dehydrogenases (LDH).

Thus, the invention provides a protein having one or more immunoreactive and/or antigenic determinants of *Eimeria* lactate dehydrogenase, which has a monomeric molecular weight of about 37kD.

More specifically the lactate hydrogenase is derived from *Eimeria acervulina*.

According to one embodiment of this invention there is provided a purified and isolated protein having one or more immunoreactive and/or antigenic determinants of *Eimeria* lactate dehydrogenase (LDH), said LDH having a relative monomeric molecular mass of about 37,000.

According to a second aspect of the invention, there is provided a nucleic acid sequence encoding all or a substantial part, in particular the immunologically active part, of a purified *Eimeria* lactate dehydrogenase. Such a nucleic acid sequence may be operatively linked to expression control sequences resulting in a recombinant nucleic acid molecule which, when inserted into a suitable vector, results in a recombinant vector capable of expressing the nucleic acid sequence.

Such a recombinant vector, or nucleic acid sequence as defined above, may be used to transform a suitable host cell or organism. Such a transformed host cell or organism may, in turn, be used to produce the stimulatory protein for incorporation into a vaccine

AUSTRALIAN PATENT OFFICE

for the protection of poultry against coccidiosis. Alternatively, the transformed host cell or organism may itself be incorporated into a vaccine.

In general, the term "protein" refers to a molecular chain of amino acids with biological activity. A protein is not of a specific length and can, if required, be modified *in vivo* or *in vitro*, by, for example, glycosylation, amidation, carboxylation or phosphorylation; thus, *inter alia*, peptides, oligopeptides and polypeptides are included within the definition.

More particularly, this invention provides proteins possessing LDH activity, or immunogenically active parts thereof, which have the amino acid sequence shown in SEQ ID NO. 2 and their biologically functional equivalents or variants.

The biologically functional equivalents or variants of the proteins specifically disclosed herein are proteins derived from the above noted amino acid sequences, for example by deletions, insertions and/or substitutions of one or more amino acids, but retain one or more immunogenic determinants of the *Eimeria* antigens, ie. said variants have one or more epitopes capable of eliciting an immune response in a host animal.

It will be understood that, for the particular proteins embraced herein, natural variations can exist between individual *Eimeria* parasites or strains. These variations may be demonstrated by (an) amino acid difference(s) in the overall sequence or by deletions, substitutions, insertions, inversions or additions of (an) amino acid(s) in said sequence. Amino acid substitutions which do not essentially alter biological and immunological activities, have been described, eg. by Neurath *et al* in "The Proteins" Academic Press New York (1979). Amino acid replacements between related amino acids or replacements which have occurred frequently in evolution are, *inter alia*, Ser/Ala, Ser/Gly, Asp/Gly, Asp/Asn, Ile/Val (see Dayhof, M.D., Atlas of protein sequence and structure, Nat. Biomed. Res. Found., Washington D.C., 1978, vol. **5**, suppl. 3). Other amino acid substitutions include Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Thr/Phe, Ala/Pro, Lys/Arg, Leu/Ile, Leu/Val and Ala/Glu. Based on this information, Lipman and Pearson developed a method for rapid and sensitive protein comparison (Science, **227**, 1435-1441, 1985) and determining the functional similarity between homologous proteins. Such amino acid substitutions of the exemplary embodiments of this invention are within the scope of the invention as long as the resulting proteins retain their immunoreactivity.

Furthermore, also immunogenic fragments of the proteins specifically disclosed herein or their functional variants are included in the present invention.

The term "fragment" as used herein means a DNA or amino acid sequence comprising a subsequence of the nucleic acid sequence or protein of the invention. Said fragment is or encodes a polypeptide having one or more immunogenic determinants of an *Eimeria* antigen. Methods for determining useable immunogenic polypeptide fragments are outlined below. Fragments can *inter alia* be produced by enzymatic cleavage of precursor molecules, using restriction endonucleases for the DNA and proteases for the

polypeptides. Other methods include chemical synthesis of the fragments or the expression of polypeptide fragments by DNA fragments.

Suitable immunogenic polypeptide fragments of a protein according to the invention containing (an) epitope(s) can be found by means of the method described in WO 86/06487, Geysen, H.M. *et al.* (Proc. Natl. Acad. Sci. **81**, 3998-4002, 1984), Geysen, H.M. *et al.* (J. Immunol. Meth. 102, 259-274, 1987) based on the so-called pepscan method, wherein a series of partially overlapping peptides corresponding with partial sequences of the complete polypeptide under consideration, are synthesised and their reactivity with antibodies is investigated.

In addition, a number of regions of the polypeptide, with the stated amino acid sequence, can be designated epitopes on the basis of theoretical considerations and structural agreement with epitopes which are now known. The determination of these regions is based on a combination of the hydrophilicity criteria according to Hopp and Woods (Proc. Natl. Acad. Sci. 78, 3824-3828, 1981) and the secondary structure aspects according to Chou and Fasman (Advances in Enzymology **47**, 45-148, 1987).

T-cell epitopes which may be necessary can likewise be derived on theoretical grounds, eg. with the aid of Berzofsky's amphiphilicity criterion (Science **235**, 1059-62, 1987).

The invention further provides isolated and purified nucleic acid sequences encoding the above mentioned proteins of *Eimeria*. One of these nucleic acid sequences is shown in SEQ. ID. NO. 1. It is well known in the art that the degeneracy of the genetic code permits substitution of bases in the codon resulting in another codon but still coding for the same amino acid, eg. the codon for the amino acid glutamic acid is both GAT and GAA. Consequently, it is clear that, for the expression of a protein with the amino acid sequence shown in SEQ. ID. NO. 2, the nucleic acid sequence may have a codon composition different from the nucleic acid sequence shown in SEQ. ID. NO. 1.

A nucleic acid sequence according to the present invention may be isolated from an *Eimeria* strain and multiplied by recombinant DNA techniques including polymerase chain reaction (PCR) technology or may be chemically synthesised *in vitro* by techniques known in the art.

A nucleic acid sequence according to the invention can be ligated to various replication effecting DNA sequences with which it is not associated, or linked in nature, resulting in a so-called recombinant vector which can be used for the transformation of a suitable host. Useful recombinant vectors are preferably derived from plasmids, bacteriophages, cosmids or viruses.

Specific vectors or cloning vehicles which can be used to clone nucleic acid sequences according to the invention are known in the art and include *inter alia* plasmid vectors such as pBR322, the various pUC, pGEM and Bluescript plasmids; bacteriophages, eg. kgt-Wes, Charon 28 and the M13 derived phages or viral vectors such as Sv40, adenovirus or polyoma virus (see also Rodriquez, R.L. and D.T. Denhardt, ed.,

Vectors: A survey of molecular cloning vectors and their uses, Butterworths, 1988; Lenstra, J.A. *et al.*, Arch. Virol., **110**, 1-24, 1990). The methods to be used for the construction of a recombinant vector according to the invention are known to those of ordinary skill in the art and are *inter alia* set forth in Maniatis, T. *et al.* (Molecular Cloning A Laboratory Manual, second edition; Cold Spring Harbor Laboratory, 1989).

For example, the insertion of the nucleic acid sequence according to the invention into a cloning vector can easily be achieved when both the genes and the desired cloning vehicle have been cut with the same restriction enzyme(s) as complementary DNA termini are thereby produced.

Alternatively, it may be necessary to modify the restriction sites that are produced into blunt ends either by digesting the single-stranded DNA or by filling in the single-stranded termini with an appropriate DNA polymerase. Subsequently, blunt end ligation with an enzyme such as T4 DNA ligase may be carried out.

If desired, any restriction site may be produced by ligating linkers onto the DNA termini. Such linkers may comprise specific oligonucleotide sequences that encode restriction site sequences. The restriction enzyme cleaved vector and nucleic acid sequence may also be modified by homopolymeric tailing.

"Transformation", as used herein, refers to the introduction of an heterologous nucleic acid sequence into a host cell, irrespective of the method used, for example direct uptake or transduction. The heterologous nucleic acid sequence may be maintained through autonomous replication or, alternatively, may be integrated into the host genome. If desired, the recombinant vectors are provided with appropriate control sequences compatible with the designated host. These sequences can regulate the expression of the inserted nucleic acid sequence. In addition to microorganisms, cell cultures derived from multicellular organisms may also be used as hosts.

The recombinant vectors according to the invention preferably contain one or more marker activities that may be used to select for desired transformants, such as ampicillin and tetracycline resistance in pBR322, ampicillin resistance and α-peptide of β-galactosidase in pUC8.

A suitable host cell is a microorganism or cell which can be transformed by a nucleic acid sequence encoding a polypeptide or by a recombinant vector comprising such a nucleic acid sequence, and which can, if desired, be used to express said polypeptide encoded by said nucleic acid sequence. The host cell can be of prokaryotic origin, eg. bacteria such as *Escherichia coli*, *Bacillus subtilis* and *Pseudomonas* species; or of eukaryotic origin such as yeasts, eg. *Saccharomyces cerevisiae* or higher eukaryotic cells such as insect, plant or mammalian cells, including HeLa cells and Chinese hamster ovary (CHO) cells. Insect cells include the Sf9 cell line of *Spodoptera frugiperda* (Luckow *et al.*, Biotechnology **6**, 47-55, 1988). Information with respect to the cloning and expression of the nucleic acid sequence of the present invention in eukaryotic cloning

systems can be found in Esser, K. *et al.* (Plasmids of Eukaryotes, Springer-Verlag, 1986).

In general, prokaryotes are preferred for the construction of the recombinant vectors useful in the present invention. *E. coli* K12 strains are particularly useful, especially DH5a or MC1061 strains.

For expression, nucleic acid sequences of the present invention are introduced into an expression vector, ie. said sequences are operably linked to expression control sequences. Such control sequences may comprise promotors, enhancers, operators, inducers, ribosome binding sites etc. Therefore, the present invention provides a recombinant vector comprising a nucleic acid sequence encoding an *Eimeria* protein identified above operably linked to expression control sequences, which is capable of expressing the DNA sequences contained therein in (a) transformed host cell(s).

It should be understood, of course, that the nucleotide sequences inserted at the selected site of the cloning vector may include nucleotides which are not part of the actual structural gene for the desired polypeptide, or may include only a fragment of the complete structural gene for the desired protein as long as the transformed host will produce a polypeptide having at least one or more immunogenic determinants of an *Eimeria* protein antigen.

When the host cells are bacteria, useful expression control sequences which may be used include the Trp promotor and operator (Goeddel, *et al.*, Nucl. Acids Res., **8**, 4057, 1980); the lac promotor and operator (Chang, *et al.*, Nature, **275**, 615, 1978); the outer membrane protein promotor (Nakamura, K. and Inouge, M., EMBO J., **1**, 771-775, 1982); the bacteriophage lambda promotors and operators (Remaut, E. *et al.*, Nucl. Acids Res., **11**, 4677-4688, 1983); the α-amylase (*B. subtilis*) promotor and operator, termination sequences and other expression enhancement and control sequences compatible with the selected host cell. When the host cell is yeast, illustrative useful expression control sequences include, eg., α-mating factor. For insect cells the polyhedrin or p10 promotors of baculoviruses can be used (Smith, G.E. *et al.*, Mol. Cell. Biol. **3**, 2156-65, 1983). When the host cell is of mammalian origin illustrative useful expression control sequences include the SV-40 promotor (Berman, P.W. *et al.*, Science, **222**, 524-527, 1983) or the metallothionein promotor (Brinster, R.L., Nature, **296**, 39-42, 1982) or a heat shock promotor (Voellmy *et al.*, Proc. Natl. Acad. Sci. USA, **82**, 4949-53, 1985). Alternatively, expression control sequences present in *Eimeria* may also be applied. For maximising gene expression, see also Roberts and Lauer (Methods in Enzymology, **68**, 473, 1979).

Therefore, the invention also comprises (a) host cell(s) containing a nucleic acid sequence or a recombinant nucleic acid molecule or a recombinant vector described above, capable of producing the *Eimeria* protein by expression of the nucleic acid sequence.

Immunisation of poultry against *Eimeria* infection can be achieved by administering to the birds a protein according to the invention in an immunologically relevant context as a so-called subunit vaccine. The subunit vaccine according to the invention may comprise a protein in a pure form, optionally in the presence of a pharmaceutically acceptable carrier. The protein can optionally be covalently bonded to a non-related protein, which can be of advantage in the purification of the fusion product. Examples are β-galactosidase, protein A, prochymosine, blood clotting factor Xa, etc.

In some cases the ability to raise protective immunity using these proteins per se may be low. Small fragments are preferably conjugated to carrier molecules in order to raise their immunogenicity.

Suitable carriers for this purpose are macromolecules, such as natural polymers (proteins like key hole limpet hemocyanin, albumin, toxins), synthetic polymers like polyamino acids (polylysine, polyalanine), or micelles of amphiphilic compounds like saponins. Alternatively these fragments may be provided as polymers thereof, preferably linear polymers.

If required, the proteins according to the invention which are to be used in a vaccine can be modified *in vitro* or *in vivo*, for example by glycosylation, acylation, amidation, carboxylation or phosphorylation.

A newly developed vaccine version is a vaccine in which the DNA coding for the protein of the invention is administered in a pharmaceutically acceptable form, for instance in the form of "bullets", which can be shot into the tissue. This naked DNA can be used as vaccine provided it is presented in a plasmid or in combination with suitable eukaryotic promoter sequences such as those from SV40 virus. In this way one can achieve the introduction of this DNA into the genomic DNA, thus ensuring the expression of the antigen *in situ*.

An alternative to subunit vaccines is live vaccines. A nucleic acid sequence according to the invention is introduced by recombinant DNA techniques into a microorganism (eg. a bacterium or virus) in such a way that the recombinant microorganism is still able to replicate, thereby expressing a polypeptide coded by the inserted nucleic acid sequence and eliciting an immune response in the infected host bird.

A preferred embodiment of the present invention is a recombinant vector virus comprising an heterologous nucleic acid sequence described above, capable of expressing the DNA sequence in (a) host cell(s) or host bird infected with the recombinant vector virus. The term "heterologous" indicates that the nucleic acid sequence according to the invention is not normally present in nature in the vector virus.

Furthermore, the invention also comprises (a) host cell(s) or cell culture infected with the recombinant vector virus, capable of producing the *Eimeria* protein by expression of the nucleic acid sequence.

For example the well known technique of *in vivo* homologous recombination can be used to introduce an heterologous nucleic acid sequence according to the invention into the genome of the vector virus.

First, a DNA fragment corresponding with an insertion region of the vector genome, ie. a region which can be used for the incorporation of an heterologous sequence without disrupting essential functions of the vector such as those necessary for infection or replication, is inserted into a cloning vector according to standard recDNA techniques. Insertion-regions have been reported for a large number of microorganisms (eg. EP 80,806, EP 110,385, EP 83,286, EP 314,569, WO 88/02022, WO 88/07088, US 4,769,330 and US 4,722,848).

Second, if desired, a deletion can be introduced into the insertion region present in the recombinant vector molecule obtained from the first step. This can be achieved for example by appropriate exonuclease III digestion or restriction enzyme treatment of the recombinant vector molecule from the first step.

Third, the heterologous nucleic acid sequence is inserted into the insertion-region present in the recombinant vector of the first step or in place of the DNA deleted from said recombinant vector. The insertion region DNA sequence should be of appropriate length as to allow homologous recombination with the vector genome to occur. Thereafter, suitable cells can be infected with wild-type vector virus or transformed with vector genomic DNA in the presence of the recombinant vector containing the insertion flanked by appropriate vector DNA sequences whereby recombination occurs between the corresponding regions in the recombinant vector and the vector genome. Recombinant vector progeny can now be produced in cell culture and can be selected for example genotypically or phenotypically, eg. by hybridisation, detecting enzyme activity encoded by a gene co-integrated along with the heterologous nucleic acid sequence, or detecting the antigenic heterologous polypeptide expressed by the recombinant vector immunologically.

Next, this recombinant microorganisms can be administered to poultry for immunisation whereafter it maintains itself for some time, or even replicates in the body of the inoculated animal, expressing *in vivo* a polypeptide coded for by the inserted nucleic acid sequence according to the invention resulting in the stimulation of the immune system of the inoculated animal. Suitable vectors for the incorporation of a nucleic acid sequence according to the invention can be derived from viruses such as pox viruses, eg. vaccinia virus (EP 110,385, EP 83,286, US 4,769,330 and US 4,722 848) or fowl pox virus (WO 88/02022), herpes viruses such as HVT (WO 88/07088) or Marek's Disease virus, adenovirus or influenza virus, or bacteria such as *E. coli* or specific *Salmonella* species. With recombinant microorganisms of this type, the polypeptide synthesised in the host animal can be exposed as a surface antigen. In this context fusion of the polypeptide with OMP proteins, or pilus proteins of for example *E. coli* or synthetic provision of signal and anchor sequences which are recognised by the organism

are conceivable. It is also possible that the *Eimeria* polypeptide, if desired as part of a larger whole, is released inside the animal to be immunised. In all of these cases it is also possible that one or more immunogenic products will find expression which generate protection against various pathogens and/or against various antigens of a given pathogen.

A vector vaccine according to the invention can be prepared by culturing a recombinant bacterium or a host cell infected with a recombinant vector comprising a nucleic acid sequence according to the invention, whereafter recombinant bacteria or vector containing cells and/or recombinant vector viruses grown in the cells can be collected, optionally in a pure form, and formed into a vaccine optionally in a lyophilised form.

Host cells transformed with a recombinant vector according to the invention can also be cultured under conditions which are favourable for the expression of a polypeptide coded by said nucleic acid sequence. Vaccines may be prepared using samples of the crude culture, host cell lysates or host cell extracts, although in another embodiment more purified polypeptides according to the invention are formed into a vaccine, depending on its intended use. In order to purify the polypeptides produced, host cells transformed with a recombinant vector according to the invention are cultured in an adequate volume and the polypeptides produced are isolated from such cells, or from the medium if the protein is excreted. Polypeptides excreted into the medium can be isolated and purified by standard techniques, eg. salt fractionation, centrifugation, ultrafiltration, chromatography, gel filtration or immunoaffinity chromatography, whereas intra cellular polypeptides can be isolated by first collecting said cells, disrupting the cells, for example by sonication or by other mechanically disruptive means such as French press, followed by separation of the polypeptides from the other intracellular components and forming the polypeptides into a vaccine. Cell disruption could also be achieved by chemical (eg. EDTA or detergents such as Triton X114) or enzymatic means, such as lysozyme digestion.

Antibodies or antiserum directed against a polypeptide according to the invention have a potential use in passive immunotherapy, diagnostic immunoassays and generation of anti-idiotypic antibodies.

The *Eimeria* proteins as characterised above can be used to produce antibodies, both polyclonal, monospecific and monoclonal. If polyclonal antibodies are desired, techniques for producing and processing polyclonal sera are known in the art (eg. Mayer and Walter. eds, Immunochemical Methods in Cell and Molecular Biology, Academic Press, London, 1987). Monospecific antibodies to an immunogen can be affinity purified from polyspecific antisera by a modification of the method of Hall *et al.* (Nature, **311**, 379-387, 1984). Monospecific antibody, as used herein, is defined as a single antibody species or multiple antibody species with homogeneous binding characteristics for the relevant antigen. Homogeneous binding, as used herein, refers to the ability of the antibody species to bind to a specific antigen or epitope.

Monoclonal antibodies, reactive against the *Eimeria* proteins according to the present invention, can be prepared by immunising inbred mice by techniques known in the art (Kohler and Milstein, Nature, **256**, 495-497, 1975). Hybridoma cells are selected by growth in hypoxanthine, thymidine and aminopterin in an appropriate cell culture medium such as Dulbecco's modified Eagle's medium. Antibody producing hybridomas are cloned, preferably using the soft agar technique of MacPherson, (Soft Agar Techniques, Tissue Culture Methods and Applications, Kruse and Paterson, eds., Academic Press, **276**, 1973). Discrete colonies are transferred into individual wells of culture plates for cultivation in an appropriate culture medium. Antibody producing cells are identified by screening with the appropriate immunogen. Immunogen positive hybridoma cells are maintained by techniques known in the art. Specific anti-monoclonal antibodies are produced by cultivating the hybridomas *in vitro* or preparing ascites fluid in mice following hybridoma injection by procedures known in the art.

Anti-idiotypic antibodies are immunoglobulins which carry an "internal image" of the antigen of the pathogen against which protection is desired and can be used as an immunogen in a vaccine (Dreesman *et al.*, J. Infect. Disease, **151**, 761, 1985). Techniques for raising anti-idiotypic antibodies are known in the art (MacNamara *et al.*, Science, **226**, 1325, 1984).

The vaccine according to the invention can be administered in a conventional active immunisation scheme: single or repeated administration in a manner compatible with the dosage formulation, and in such amount as will be prophylactically effective, ie. the amount of immunising antigen or recombinant microorganism capable of expressing said antigen that will induce immunity in poultry against challenge by virulent *Eimeria* parasites. Immunity is defined as the induction of a significant level of protection in a population of chickens after vaccination compared to an unvaccinated group.

Next to an increase in protection a vaccine comprising the polypeptide of the invention will also reduce the number of oocysts shedded by the infected animals. Normally, the shedded oocysts will infect other animals in the flock. A decrease in the number of oocysts shedded will then also give a decrease in the number of animals which is subsequently infected and also a decrease in the number of oocysts shedded will give rise to a lesser infective load.

Furthermore, even without effect on the parasite itself, a vaccine can decrease the incidence of disease. This is especially so when the symptoms of the disease are caused by products released by the parasite. Vaccines directed against such products alleviate the symptoms without attacking the parasite.

For live viral vector vaccines the dose rate per chicken may range from 10^5-10^8pfu. A typical subunit vaccine according to the invention comprises 1μg - 1mg of the protein according to the invention. Such vaccines can be administered intradermally, subcutaneously, intramuscularly, intraperitoneally, intravenously, orally or intranasally.

Additionally the vaccine may also contain an aqueous medium or a water containing suspension, often mixed with other constituents in order to increase the activity and/or the shelf life. These constituents may be salts, pH buffers, stabilisers (such as skimmed milk or casein hydrolysate), emulsifiers, adjuvants to improve the immune response (eg. oils, muramyl dipeptide, aluminium hydroxide, saponin, polyanions and amphipatic substances) and preservatives.

A vaccine comprising the polypeptide of the invention may also comprise other immunogenic proteins of *E. maxima* or immunogenic proteins of other *Eimeria* species. Such a combination vaccine will decrease the parasitic load in a flock of poultry and will increase the level of protection against coccidiosis.

It is clear that a vaccine according to the invention may also contain immunogens related to other pathogens of poultry, or may contain nucleic acid sequences encoding these immunogens, like antigens of Marek's Disease virus (MDV), Newcastle Disease virus (NDV), Infectious Bronchitis virus (IBV), Chicken Anemia Agent (CAA), Reo virus, Avian Retro virus, Fowl Adeno virus, Turkey Rhinotracheitis virus or *E. coli* to produce a multivalent vaccine.

The invention is illustrated by the following examples.

Example 1

Handling of parasites

Eimeria acervulina (Houghton strain) and *Eimeria tenella* (Weybridge strain) parasites were collected after deliberate infection of chickens reared in the absence of coccidia. *E. acervulina* oocysts were isolated from faecal material on days 4 and 5 post-infection (p. i.). *E. tenella* oocysts were harvested from the caeca on day 7 p. i.

The oocysts were sporulated with strong aeration at 30°C for 7 hours, resulting in partially sporulated oocysts. Release of sporocysts and sporozoites of 48h sporulated oocysts was performed as described earlier in A.N. Vermeulen *et al.* FEMS Microbiological Letters **110**, (1993), 223-230.

To obtain *E. acervulina* intra-cellular stages, chickens were infected at 5 weeks with 10^8 sporulated *E. acervulina* oocysts. Intracellular parasites were harvested from the duodenum after 42 hours. Hereto chickens were exsanguinated 42h post inoculation and duodenum was removed from the stomach to Meckel's diverticulum. The tissue was washed and cut into small pieces of approximately $1cm^3$. The pieces were suspended in calcium/magnesium free Hanks BSS containing 10mg/mL glucose (CMF-Hanks). Epithelial cells were released from the matrix by 10min incubation in EDTA (2mM EDTA in CMF Hanks at 35-37°C). Supernatants of four incubations were pooled and centrifuged 10min at 750g, which pelleted the cells. The intracellular parasites (further called "schizonts", although also trophozoites were present) were subsequently released from the host cells by saponin lysis (15min in 0.1% saponin in CMF-Hanks at room temperature) and mechanical shearing.

The schizonts were pelleted and separated from host material after centrifugation through 45% Percoll (Pharmacia Fine Chemicals) (20min. 700g, 4°C). Dry pellets of schizonts were stored at -70°C until further use.

Triton X114 extraction

Triton X114 extractions were carried out to obtain the hydrophilic protein fraction of schizonts. The procedure used was described earlier by C. Bordier (1981) Journal of Biological Chemistry, vol. **256** no. 4 (feb) pp. 1604-1607. 10^8 to 10^9 *E acervulina* schizonts per mL of TBS (10mM Tris-HCl, 150mM NaCl pH7.4) were sonified ±3 x 20sec on ice with the microtip (Branson sonifier, position 7). PMSF (final concentration 1mM) and DNase/RNase (final concentration for both 0.02mg/mL) was added (DNase/RNase stock: 2mg/mL DNase, 2mg/mL RNase in 5mM $MgCl_2$).

Precondensed TritonX114 was added to the sonified schizonts in suspension to a final concentration of 10%(v/v) and mixed well to dissolve the proteins. The non-extractable material was pelleted by centrifugation 20min 12 000g at 4°C. The soluble fraction was layered over a sucrose cushion (6% sucrose, 0.06%(v/v) TX114 in TBS), incubated 10min 40°C and spun 10min 400g at room temperature. The hydrophilic fraction was extracted again by the same procedure.

The hydrophilic fractions were stored at -70°C until further use. Total protein concentration was determined using the BCA (Pierce Chemicals) assay.

Prep-cell fractionation

Hydrophilic proteins were further separated with respect to their relative molecular mass on SDS-PAGE under reducing conditions in the Laemmli buffer system. Hereto we made use of preparative electrophoresis in the so-called Prepcell.

Materials: Prep cell apparatus (Biorad Labs) with Prep cell column (37mm ID) Dialyse membrane for Prep cell (cut off 6kD) Powersupply (EPS 600 Pharmacia) Reducing sample buffer: 62.5mM Tris-HCl pH 6.8; 10% glycerol; 28 SDS; 0.018 bromophenol blue (Merck); 0.13M DTT (dithiothreitol, Merck) Electrophoresis buffer/elution buffer: 25mM Tris, 192mM Glycine, 0.1% SDS pH8.6

Method and results:

All procedures were performed at 4°C. For the fractionation of the hydrophilic proteins a 4% stacking/9% separating gel (polyacrylamide) was used in the 37mm tube (filled to 6cm) of the Prepcell according to the manufacturers protocol, but with the addition of 0.1% SDS.

The hydrophilic phase of TX114 extractions kept at -70°C was thawed and the hydrophilic proteins (about 8mg per run) were diluted in reducing sample buffer (total volume was + 6mL), boiled 3min 100°C, and were loaded on the surface of the 4% stacking gel using a narrow tube affixed to a syringe.

The Prepcell was connected to the powersupply and electrophoresis was started at 40mA, 500V max.

The collection of fractions (fraction volume ±2.5-3mL; flow 0.6mL/min) started after about 6 hours, when the tracking dye eluted from the cell. Fractions were collected overnight (±100 fractions) in plastic 3.5mL tubes (Sarstedt).

Samples of the fractions were taken for analysis by SDS-PAGE and Western Blotting. Fractions were stored at -70°C.

This purification method resulted in fractions containing almost pure proteins as follows from analyses shown below.

Amino acid sequencing

Selected fractions of Prepcellrun COC9314612 containing an almost pure band around Mr=37k (designated as EASC2) were pooled, concentrated by cetoneprecipitation and run on a 12% PAAgel. The gel was shortly stained with a non-denaturing Coomassie Brilliant Blue staining protocol: staining: 20min at ambient temperature in 0.2% CBB in 20% methanol/0.5% acetic acid. Destaining: 60min in 30% methanol.

The staining 37kD band was cut out. Internal amino acid sequencing was performed on a selected HPLC-purified peptide of a trypsin digest of the EASC2, all performed by Eurosequence BV Groningen The Netherlands.

The amino-acid sequence of the tryptic peptide was GWIKQEEVDDIVQK (see SEQ.ID.No:2 amino acids 212-225).

This coding sequence for this peptide was also detected after DNA sequencing of the clone.

Example 2

Preparation of monospecific antibodies in rabbits

Prevaccination sera of SPF rabbits were screened on western blotted *E. acervulina* antigens of different developmental stages and on a blot of *E coli* proteins. 'Negative' rabbits were selected for the raising of antibodies.

Fractions of Prepcell runs containing EASC2 (37kD) were selected by SDS-PAGE, pooled and concentrated (±3x) with an Amiconcell (YM10 filter) to 3.5mL.

The rabbit was twice immunised with concentrated antigen in GNE (8 x 0.25mL i. c.; 1mL i.p.) with an interval of 4 weeks. Two weeks after the second immunisation the rabbit was bled and sera were tested on western blots of *Eimeria acervulina* en tenella sporozoites and schizonts 42h. Figure 1 shows the result of the immunodetection of the monospecific antiserum on sporozoite antigens of both species. It appeared that the antibodies recognised a parasite product of about 37kD in both *E. acervulina* (Lane A1) and *E. tenella* (Lane B1). Control sera of the same rabbit prior to immunisation did not recognise these bands (Lanes A/B2). The protein is also present in schizont stages of the two species (not shown).

Example 3

Vaccination of chickens with E. acervulina TX114 hydrophilic fraction and EASC2

The TX114 hydrophilic phase of schizont material was separated and dialysed extensively against 0.01M PBS pH 7.3 at 4°C.

Selected fractions containing the EASC2 37kD protein were dialysed extensively against 3 x 5 litre 0.01M PBS pH 7.3 at 4°C.

The concentration of protein in the vaccine preparations was estimated by staining different concentrations of sample with CBB after SDS-PAGE and comparing the intensity of the staining with a reference sample of BSA.

The volumes were corrected to obtain ±5μg protein/dose for the purified protein and about 15μg/dose for the total hydrophilic fraction.

These were stored as aliquotted volumes for priming and booster vaccination at -70°C. Frozen vaccine preparations were thawed.

To every 15mL of vaccine 3.2mg Quil A Superfos Biosector was added as adjuvant in a volume of 1mL 0.01M PBS pH 7.3.

Vaccine was mixed well by vortexing and injected in 4-6 week old coccidia-free White Leghorn chickens in 0.75mL given subcutaneously.

The vaccine contained 150μg Quil A/dose.

Figure 2 shows a Coomassie BB stained SDS-PAGE of the EASC2 (Lane 1) and 42h TX114 hydrophilic fraction (Lane 2) injected into the chickens as vaccine.

Four weeks after priming birds were boosted with the same dose *via* the same route. The booster-vaccine was prepared freshly from the frozen antigen stock.

Control chickens were inoculated with 150μg Quil A/dose in PBS. Each group comprised 14 chickens.

Eleven days after the boosting vaccination all chickens were inoculated orally with 240 sporulated oocysts of *Eimeria acervulina* H in 1mL of 15% sucrose in water.

Chickens were placed in cages 2 birds per cage. Oocyst output was assessed in faecal samples taken from days 4 to 8 after challenge.

Table 2 shows the results of this experiment.

Oocyst output is expressed as % oocysts from the output in the control animals. Statistical evaluation of the data was performed on the LOG of the number of oocysts using Student's T-test or Mann-Whitney's test if data distribution was not normal. When $p < 0.05$ the difference was regarded significant.

This table shows that both the TX114 fraction and the EASC2 prepcell purified fraction induce a statistically significant reduction ($p < 0.05$) in oocyst output after challenge.

Prep cell purification seemed to improve the protection induced by the TX114 vaccine.

Table 2.

Oocyst output in percents from control and statistical value of difference

Immunogen	%oocyst output from control+S.D.	p value different from control
EASC2 prep cell pure ±5μg/dose	72 ± 30	p=0.01
Hydrophilic TX114 proteins of Schizonts±15μg/dose	84 + 17	p=0.02

In another experiment in which only total extracts of 42h schizonts were used as vaccine no significant oocyst reduction could be induced (results not shown).

In a second experiment prepsell purified EASC2 was used in two different concentrations of 0.2 and 2μg/dose. Following the same protocol for immunisation and challenge, protection was measured in ten chickens per group as reduction of oocyst output compared to the group inoculated with PBS/QuilA.

Table 3 summarises the average percentual oocyst output of the control for the two EASC2 vaccinated groups. This table demonstrates that the EASC2 protected in a dose dependent manner showing a statistically significant difference at a dose of 2μg/dose.

Table 3.

Oocyst output in percents from control and statistical value of difference

Group	% oocysts + S.D. (control output=100%)	significance of difference from control (p-value)
EASC2/Quil A 2μg/dose	64.0 + 22	0.008
EASC2/Quil A 0.2μg/dose	90.2 + 27	NOT SIGNIFICANT

Example 4

Immunological stimulation after vaccination with EASC2 or TX 114 hydrophilic proteins.

In both protection experiments mentioned above chickens were assayed for stimulation of immunological parameters such as T-lymphocyte proliferation and serum antibodies.

Serum antibodies

Antibodies recognising the vaccine constituents were only detected in sera from the groups vaccinated with the 42h TX114-hydrophilic fraction and not the group vaccinated with the purified EASC2.

Lymphocyte proliferation

Lymphocyte proliferation after antigenic stimulus was tested in a lymphocyte stimulation test (LST).

Method:

Prior to challenge peripheral blood cells were taken from all chickens of each group.

Peripheral blood leucocytes (PBL) were isolated by centrifugation 3mL of the total blood for 7min at 64g at ambient temperature. The buffy coat was collected in RPMI 1640 (Dutch modification) and washed two times. Cell concentration was adjusted to 1 x 10^7 cells per mL in RPMI 1640. The RPMI 1640 (Dutch modification) used was supplemented with sodium pyruvate (1mM), Glutamine (2mM), penicillin 200U/mL and streptomycin 200μg/mL.

96 well round-bottom tissue culture plates were seeded with 0.05mL cell suspension with 3.0% chicken serum (Gibco BRL), 0.05mL "stimulating antigen" suspension and 0.05mL RPMI 1640 , cultured for 64h at 41°C under 5% CO_2 atmosphere. Subsequently 18.5kBq ^{3}H-thymidine (Amersham Beckenham U.K.) was added per well and 8h later the

cells were harvested on a glass-fibre filter (Skatron Norway Bluemat) using a 96 well Cell Harvester (Skatron Norway). The filters were saturated with scintillation fluid (LKB BetaScint) and counted in a Betaplate 1205 (Pharmacia/LKB Sweden).

As "stimulating antigen" *E. acervulina* schizonts were used, which were sonicated using a microtip-equipped Branson sonifier at position 6 for 3 x 20 seconds with intermediate cooling and store at -70°C. The antigens were thawed before use and diluted to meet the concentration used for the stimulation. PBL of all groups were stimulated with 3.10^5 *E. acervulina* schizonts.

Statistical evaluation was performed using Student's T-test on the LOG of the Stimulation Index (SI) (the number of counts per min (cpm) of the stimulated cultures divided by the cpm of the non-stimulated control). When $p<0.05$ the difference was regarded significant.

Results:

Table 4. shows the mean S.I. for the groups from both experiments described above. The first experiment in which EASC2 vaccine was compared with the TX114-hydrophilic fraction, and the second experiment dealing with the two dosages of the EASC2 vaccine.

It appeared that all antigens or dosages induced a significant positive T-cell response detectable in the peripheral blood at the time of challenge.

In both experiments, however, the higher dose of the prepcell pure EASC2 vaccine (2 or 5µg/dose) induced the very highest stimulation of T-cells. The ranking of the T-cell stimulation correlated with the reduction in oocyst output after challenge.

Table 4.

Mean incorporation of 3H-Thymidine after stimulation with *E. acervulina* schizonts by PBL from groups immunised with the different vaccines, expressed as Stimulation Index (S.I) ± Standard Error (SE).

Experiment	Group	^{3}H-thymidine incorporation in S.I. ± SE
I	EASC2 5µg	120 ± 47 @
	TX114 hydrophilic proteins	31 ± 12 @
	Placebo	6 ± 1
II	EASC2 2µg	112 ± 28 @
	EASC2 0.2µg	24 ± 4 @
	Placebo	2.3 ± 0.3

@ Significant from control group p<0 001

Example 5

Cloning Experiments

Sporulation of *E. acervulina* oocysts

A suspension of $5*10^8$ *E. acervulina* oocysts in 60mL 10-4M sodium dithionite was centrifuged, after which the pellet was washed once with 100mL sterile water. The cells were resuspended in 500mL 2% potassium bichromate and then incubated under the

influence of strong aeration for 7 hours at 30°C. The oocysts were then collected by centrifuging and washed three times with 200mL sterile water.

Isolation of RNA

For the isolation of RNA (Pasternak J. *et al.*, Mol. & Bioch. Par. **3**, 133-142, 1981) the cell pellet was taken up into 2.8mL of buffer containing 10mM Tris acetate (pH7.6), 75mM sodium acetate, 1% SDS, 2mM EDTA, 0.2mg/mL proteinase K and 10mM vanadyl ribonucleoside complexes. The oocysts were destroyed by vortexing for 60 seconds (max) in the presence of 13g glass beads (0 0.5mm). 5mL of phenol was added to the total extract and the mixture was vortexed for a further 60 seconds. After centrifuging, the supernatant liquor was pipetted off and again extracted with an equal volume of phenol/chloroform/isoamyl alcohol (25:24:1). RNA was precipitated after adding 2.5 volume ethanol and the resulting precipitate was dissolved in 800mL Tris 10mM, EDTA 0.1mM pH 7.6 ($T_{10}E_{0.1}$), after which the product was extracted a further twice with an equal volume of phenol/chloroform/isoamyl alcohol (25:24:1) and twice with chloroform/isoamyl alcohol (24:1) and then precipitated with ethanol. PolyA$^+$-RNA was isolated by means of oligo (dT) -cellulose chromatography (Maniatis T. *et al.*: Molecular Cloning Cold Spring Harbor Laboratory, 1982). Approximately 100µg polyA$^+$-RNA was isolated from $5*10^8$ oocysts.

cDNA synthesis

PolyA$^+$-RNA was converted to cDNA by means of the enzyme MMLV reverse transcriptase. For this purpose 25µg polyA$^+$-RNA was dissolved in 90mL of water and denatured for 5 minutes at 20°C by adding mercury methyl hydroxide to 10mM, after which β-mercaptoethanol was added to 45mM and the mixture incubated for a further 3 minutes at 20°C. The enzyme reaction was carried out in 190mL buffer containing 4mg oligo (dT) 15, 150 U RNasin(R), 20mM Tris (pH 7.6), 30mM KCl, 4mM dithiothreitol (DTT), 2mM $MgCl_2$, 1mM of each dNTP and 3000 U MMLV reverse transcriptase.

The reaction was stopped after 1 hour incubation at 37°C by adding 10mL 0.5M EDTA. After extraction with an equal volume of phenol/chloroform/isoamyl alcohol (25:24:1), the RNA/DNA hybrid was precipitated by adding ammonium acetate to 2M and 2.5 volumes ethanol. The combined action of the enzymes DNA-polymerase I and RNase H (Gubbler U. *et al.*, Gene **25**, 263-269, 1983) results in the synthesis of the second string. The pellet was dissolved in 960µL of buffer containing 20mM Tris (pH 7.6), 5mM $MgCl_2$, 100mM $(NH_4)_2SO_4$, 0.6mM B-NAD, 16 U RNase H. 200 U DNA-polymerase I and 20 U DNA-ligase (*E.coli*). The incubation time was 1 hour at 12°C and then 1 hour at 22°C, after which the reaction was stopped by adding an equal volume of phenol/chloroform/isoamyl alcohol (25:24:1) and precipitating with ethanol.

Before the cDNA was cloned in a vector suitable for this purpose it was first modified. cDNA (5µg.) was dissolved in 100µL of buffer containing 30mM sodium acetate (pH 5.6), 50mM NaCl, 1mM $ZnSO_4$ and 21 U Mung Bean Nuclease. After incubation for 30 minutes at 37°C the reaction was stopped by adding EDTA to 10mM

and Tris to 25mM. After extraction with phenol/chloroform/isoamylalcohol (25:24:1) the mixture was desalinated over a Sephadex G50 column. The following were added to the eluate (125µL): Tris pH 7.6 to 50mM, EDTA to 2.5mM, DTT to 5mM, S'-adenosylmethionine to 0.5mM and 100 U EcoRI-methylase. After incubation for 30 minutes at 37°C, the reaction was stopped by heating for 15 minutes at 65°C, after which 1/10 volume of a solution containing Tris-HCl 100mM, $MgCl_2$ 100mM and NaCl 500mM (pH7.5) was added, and, at the same time, each dNTP to 1mM and 12.5 U Klenow DNA-polymerase. The reaction was stopped by adding an equal volume of phenol/chloroform/isoamyl alcohol (25:24:1) after incubating for 60 minutes at 22°C. The supernatant liquor was precipitated after adding 350µL H_2O and 50µl 3M sodium acetate (pH 5.6) with 500µL isopropanol. After dissolving in 100mL H_2O, the pellet was desalinated over Sephadex G50 and the eluate precipitated with ethanol. After dissolving the pellet in 24µL H_2O, ligation was carried out in 50µL by adding 2µg EcoRI linker, Tris-HCl(pH 8.0) to 30mM, $MgCl_2$ to 10mM, dithiothreitol to 10mM, ATP to 1mM, gelatin to 0.1mg/mL and 10 U T4DNA-ligase. The reaction was stopped after 16 hours incubation at 4°C by heating (for 15 minutes at 70°C) after which cutting was carried out with restriction endonuclease EcoRI in 210µl buffer containing 100mM Tris-HCl (pH 7.6), 50mM NaCl, 10mM $MgCl_2$, 2.5mM DTT and 500 U EcoRI. After 90 minutes incubation at 37°C, the reaction was stopped by means of extraction with an equal volume of phenol/chloroform/isoamylalcohol (25:24:1). The supernatant liquor was precipitated with 2.5 volume ethanol after adding sodium acetate (pH 5.6) to 300mM cDNA and linkers were separated by means of a Biogel A15M column. The cDNA was precipitated with ethanol, after which the precipitate was dissolved in Tris-HCl 10mM, EDTA 0.1mM(pH 7.6). The cDNA molecules were then cloned in phage lambda ZAPII (Stratagene).

Screening of the cDNA bank ($2*10^5$ pfu) with antibodies directed against the EASC2 protein fraction of *E acervulina* schizonts revealed six positive phage clones. These antibodies were diluted 1:2000 with 1 x Tris salt (Tris-HCl 10mM, NaCl 150mM, pH 8.0) + 0.05% Tween 20 + 10% Foetal Calf Serum (FCS) and incubated for two hours at room temperature (RT) with the filters. The filters were then washed 4 times, for 10 minutes each time, with 50 mL 1 x Tris salt + 0.05% Tween 20, each filter. For the second antibody incubation a conjugate of goat-anti-rabbit antibodies and alkaline phosphatase was used (diluted 1:7500 in 1 x Tris salt + 0.05% Tween 20 + 10% FCS) and incubated for 30 minutes at RT, after which the filters were washed as described after the first antibody incubation. The colour reaction was carried out in Tris-HCl 100mM, NaCl 100mM, $MgCl_2$ 10mM, (pH 9.6), in which 0.33mg/mL Nitrobluetetrazolium and 0.17mg/mL 5-bromo-4-chloro-3-indolyl phosphate were dissolved. The filters were evaluated after 30 minutes incubation at RT. The immunopositive clones were plaque-purified and rescued by means of *in vivo* excision, according to the protocol of the manufacturer (Stratagene). Plasmid DNA was isolated, from the resulting *in vivo*

excision clones, for sequencing purposes according to standard protocols (Maniatis T., *et al. supra*). Partial sequence information showed all clones to be homologous, from the largest clone the nucleotide sequence was determined completely. This clone, designated pBLUE EASC2, contained an insert of 156 bp.

Legend to the figures.

Fig. 1. Western blot of *E. acervulina* (A) and *E. tenella* (B) sporozoite proteins probed with antiserum raised against Prep cell purified EASC2 protein (Lane 1) or pre-immune control serum (Lane 2). Markers indicate molecular weight calibration in kD.

Fig. 2. Coomassie Brilliant Blue stained SDS-PAGE of Prep cell purified EASC2 protein (Lane 1) or TX114 hydrophilic fraction of *E. acervulina* 42h schizonts (lane 2). Lane M contains molecular weight calibration markers in kD.

Sequence Listing

Information for SEQ ID NO: 1:

Sequence Characteristics: (a) length: 1679 base pairs (b) type: nucleic acid (c) strandedness: double (d) topology: linear (ii) molecule type: cDNA to mRNA (iii) hypothetical: no (iv) anti-sense: no (vi) original source: (a) organism: *Eimeria acervulina* (d) developmental stage: schizont (vii) immediate source: (b.) clone: easc2 1 (ix) feature: (a) name/key: cds (b) location:280..1269 (d) other information:/function= "*Eimeria* lactate dehydrogenase" (ix) feature: (a) name/key: misc feature (b) location: 1. . 51 (d) other information:/label= pbluescriptii (ix) feature: (a) name/key: misc feature (b) location:1624..1679 (d) other information: /label= pbluescriptii (ix) feature: (a) name/key: misc feature (b) location: 45 . . 54 (d) other information: /label= ECORI-linker (ix) feature: (a) name/key: misc_feature (b) location: 1621. .1630 (d) other information:/label= ECORI-linker

Sequence Description: SEQ ID NO: 1

```
GCGGTGGCGG CCGCTCTAGA ACTAGTGGAT CCCCCGGGCT GCAGGAATTC GGGTTTTTTT   60
TTTTTTTTCT ACACATTAAT ATTCTTCGTT TACGTTTATT TTGCTACAAA TAAACCCCTT  120
AAACTCTCTA TTTCCTCATA TTCTACCGCT TCATCGGTGG GTGTGTAAGA CGTACGTACG  180
TACAGCTGGG GCTGGCTTAC TGCGCACCGC TTATTTATTA CTTAATTCAT ACACATTTTA  240
TATCTTTCTT CTTCTTTTTT CTTGCTCTTT CTTGTGAAA ATG GCG GTC TTC GAG     294
                                          Met Ala Val Phe Glu
                                           1               5

AAG AAT ACA CGC CCC AAG ATT GCT ATG GTG GGC TCC GGT ATG ATT GGA   342
Lys Asn Thr Arg Pro Lys Ile Ala Met Val Gly Ser Gly Met Ile Gly
                   10                  15                  20

GGC ACC ATG GCT TTC CTG TGC AGC TTG AGG GAA CTC GGA GAT GTT GTC   390
Gly Thr Met Ala Phe Leu Cys Ser Leu Arg Glu Leu Gly Asp Val Val
               25                  30                  35

CTC TTC GAC GTT GTA CCG AAC ATG CCG ATG GGG AAG GCG ATG GAT ATA   438
Leu Phe Asp Val Val Pro Asn Met Pro Met Gly Lys Ala Met Asp Ile
           40                  45                  50

TCG CAC AAT TCG TCG GTG GTT GAC ACG GGT ATA ACA GTA TAC GGC TCA   486
Ser His Asn Ser Ser Val Val Asp Thr Gly Ile Thr Val Tyr Gly Ser
       55                  60                  65
```

AAT TCA TAC GAG TGC TTG AAG GGT GCG GAC GTA GTA ATA A'I'A ACA GCA 534
Asn Ser Tyr Glu Cys Leu Lys Gly Ala Asp Val Val Ile Ile Thr Ala
70 75 80 85

GGG ATA ACA AAG ATA CCC GGA AAG AGC GAT AAA GAA TGG TCT AGA ATG 582
Gly Ile Thr Lys Ile Pro Gly Lys Ser Asp Lys Glu Trp Ser Arg Met
90 95 l00

GAT CTA TTA CCT GTG AAT ATA AAG ATA ATG AGG GAG GTC GGT GCA GCA 630
Asp Leu Leu Pro Val Asn Ile Lys Ile Met Arg Glu Val Gly Ala Ala
105 110 115

ATT AAA TCT TAC TGT CCT AAT GCA TTT GTT ATT AAT ATA ACA AAT CCT 678
Ile Lys Ser Tyr Cys Pro Asn Ala Phe Val Ile Asn Ile Thr Asn Pro
120 125 130

TTA GAT GTG ATG GTA GCT GCT CTT CAA GAG TCA TCA GGA CTA CCT CAT 726
Leu Asp Val Met Val Ala Ala Leu Gln Glu Ser Ser Gly Leu Pro His
135 140 145

CAT AGA ATC TGC GGT ATG GCT GGG ATG CTT GAT AGC TCT CGT TTT AGA 774
His Arg Ile Cys Gly Met ala Gly Met Leu Asp Ser Ser Arg Phe Arg
150 155 160 165

CGT ATG ATA GCT GAT AAA TTA GAA GTC TCT CCT AGA GAT GTA CAG GGG .822
Arg Met Ile Ala Asp Lys Leu Glu Val Ser Pro Arg Asp Val Gln Gly
170 175 180

ATG GTC ATA GGT GTA CAC GGC GAT CAT ATG GTG CCC CTA AGT AGA TAT 870
Met Val Ile Gly Val His Gly Asp His Met Val Pro Leu Ser Arg Tyr
185 190 195

GCA ACA GTT AAC GGC ATC CCG CTT TCT GAG TTT GTT AAG AAG GGC TGG 918
Ala Thr Val Asn Gly Ile Pro Leu Ser Glu Phe Val Lys Lys Gly Trp
200 205 210

ATC AAG CAA GAA GAA GTA GAT GAT ATC GTT CAG AAG ACC AAG GTC GCT 966
Ile Lys Gln Glu Glu Val Asp Asp Ile Val Gln Lys Thr Lys Val Ala
215 220 225

GGA GGA GAG ATC GTA CGC CTA TTA GGA CAA GGC TCT GCT TAC TAT GCT 1014
Gly Gly Glu Ile Val Arg Leu Leu Gly Gln Gly Ser Ala Tyr Tyr Ala
230 235 240 245

CCA GGG GCT TCA GCT ATT CAG ATG GCT GAG AGC TAT CTA AAG GAT AGA 1062
Pro Gly Ala Ser Ala Ile Gln Met ala Glu Ser Tyr Leu Lys Asp Arg
250 255 260

AAG AGA GTG ATG GTT TGC TCT TGC TAC TTG CAA GGA CAA TAT GGT GTA 1110
Lys Arg Val Met Val Cys Ser Cys Tyr Leu Gln Gly Gln Tyr Gly Val
265 270 275

CAG AAT CAC TAC TTA GGA GTA CCT TGT GTT ATC GGT GGG AGA GGT GTT 1158
Gln Asn His Tyr Leu Gly Val Pro Cys Val Ile Gly Gly Arg Gly Val
280 285 290

GAG AAG ATT ATT GAG TTA GAA TTG ACC GCA CAA GAA AGA CAG GAG CTT 1206
Glu Lys Ile Ile Glu Leu Glu Leu Thr Ala Gln Glu Arg Gln Glu Leu
295 300 305

CAG GGA TCT ATC GAT GAG GTT AAG GAG ATG CAG AAG GCT ATT GCT GCT 1254
Gln Gly Ser Ile Asp Glu Val Lys Glu Met Gln Lys Ala Ile Ala Ala
310 315 320 325

CTT GAT GCA TCC AAG TAAGCAGCAG CAAAATCGCA GAAGTTGCAG CGCTAGAACA 1309
Leu Asp Ala Ser Lys
330

ACCAGCAGCA GCAGCAGCAG CAGCCTATAG TTCTTGCTGC TGCTGTTCCT ACTACAGCTG 1369

CGGCTTTCTT CCTCGTGTTA TTATCATGAT AGTAAGCTGC TGTACCAGCA GCAGCAGCAG 1429

CAGCAGATTT TGCTTGCACC GCCGTTTGTT TTGCGTACAC CGGCAGAAAT ATTGACTTGC 1489
AGTTAGGAGA AAGAAAGAAA ACAAACACGA TCCCATCGAT CCCAATAAAC CCCACACTGT 1549
CGATCCCATC GATCCCAGCA ACTCCACGGG GCTCTTAACT GTTAAACCTA TTATTCTTAT 1609
CATTACTGTC TCCCGAATTC GATATCAAGC TTATCGATAC CGTCGACCTC GAGGGGGGGC 1669
CCGGTACCCA 1679

Information for SEQ ID NO: 2

Sequence characteristics: (a) length: 330 amino acids (b) type: amino acid (d) topology: linear (ii) molecule type: protein

Sequence Description: SEQ ID NO: 2

Met 1	Ala	Val	Phe	Glu 5	Lys	Asn	Thr	Arg	Pro 10	Lys	Ile	Ala	Met	Val 15	Gly
Ser	Gly	Met	Ile 20	Gly	Gly	Thr	Met	Ala 25	Phe	Leu	Cys	Ser	Leu 30	Arg	Glu
Leu	Gly	Asp 35	Val	Val	Leu	Phe	Asp 40	Val	Val	Pro	Asn	Met 45	Pro	Met	Gly
Lys	Ala 50	Met	Asp	Ile	Ser	His 55	Asn	Ser	Ser	Val	Val 60	Asp	Thr	Gly	Ile
Thr 65	Val	Tyr	Gly	Ser	Asn 70	Ser	Tyr	Glu	Cys	Leu 75	Lys	Gly	Ala	Asp	Val 80
Val	Ile	Ile	Thr	Ala 85	Gly	Ile	Thr	Lys	Ile 90	Pro	Gly	Lys	Ser	Asp 95	Lys
Glu	Trp	Ser	Arg 100	Met	Asp	Leu	Leu	Pro 105	Val	Asn	Ile	Lys	Ile 110	Met	Arg
Glu	Val	Gly 115	Ala	Ala	Ile	Lys	Ser 120	Tyr	Cys	Pro	Asn	Ala 125	Phe	Val	Ile
Asn	Ile 130	Thr	Asn	Pro	Leu	Asp 135	Val	Met	Val	Ala	Ala 140	Leu	Gln	Glu	Ser
Ser 145	Gly	Leu	Pro	His	His 150	Arg	Ile	Cys	Gly	Met 155	Ala	Gly	Met	Leu	Asp 160
Ser	Ser	Arg	Phe	Arg 165	Arg	Met	Ile	Ala	Asp 170	Lys	Leu	Glu	Val	Ser 175	Pro
Arg	Asp	Val	Gln 180	Gly	Met	Val	Ile	Gly 185	Val	His	Gly	Asp	His 190	Met	Val
Pro	Leu	Ser 195	Arg	Tyr	Ala	Thr	Val 200	Asn	Gly	Ile	Pro	Leu 205	Ser	Glu	Phe
Val	Lys 210	Lys	Gly	Trp	Ile	Lys 215	Gln	Glu	Glu	Val	Asp 220	Asp	Ile	Val	Gln
Lys 225	Thr	Lys	Val	Ala	Gly 230	Gly	Glu	Ile	Val	Arg 235	Leu	Leu	Gly	Gln	Gly 240
Ser	Ala	Tyr	Tyr	Ala 245	Pro	Gly	Ala	Ser	Ala 250	Ile	Gln	Met	Ala	Glu 255	Ser
Tyr	Leu	Lys	Asp 260	Arg	Lys	Arg	Val	Met 265	Val	Cys	Ser	Cys	Tyr 270	Leu	Gln
Gly	Gln	Tyr 275	Gly	Val	Gln	Asn	His 280	Tyr	Leu	Gly	Val	Pro 285	Cys	Val	Ile
Gly	Gly 290	Arg	Gly	Val	Glu	Lys 295	Ile	Ile	Glu	Leu	Glu 300	Leu	Thr	Ala	Gln
Glu 305	Arg	Gln	Glu	Leu	Gln 310	Gly	Ser	Ile	Asp	Glu 315	Val	Lys	Glu	Met	Gln 320
Lys	Ala	Ile	Ala	Ala 325	Leu	Asp	Ala	Ser	Lys 330						

The claims defining the invention are as follows:

1. A purified and isolated protein having one or more immunoreactive and/or antigenic determinants of *Eimeria* lactate dehydrogenase (LDH), said LDH having a relative monomeric molecular mass of about 37,000.

2. A protein according to claim 1 wherein the *Eimeria* species is *Eimeria acervulina*.

3. A protein according to claim 1 characterised in that it comprises at least part of the amino acid sequence shown in SEQ ID NO. 2 or a biologically functional equivalent thereof.

4. A purified and isolated protein having one or more immunoreactive and/or antigenic determinants of *Eimeria* lactate dehydrogenase, substantially as hereinbefore described with reference to any one of the examples.

5. A nucleic acid sequence encoding a protein according to any one of claims 1 to 4.

6. A nucleic acid sequence according to claim 5 characterised in that the nucleic acid sequence contains at least part of the DNA sequence shown in SEQ ID No. 1.

7. A recombinant nucleic acid molecule comprising a nucleic acid sequence according to claim 5 or 6 operatively linked to expression control sequences enabling expression of said nucleic acid sequence.

8. A recombinant vector comprising a nucleic acid sequence according to claim 5 or 6.

9. A recombinant vector according to claim 8 characterised in that the nucleic acid sequence is operatively linked to expression control sequences.

10. A host cell or organism transformed with a nucleic acid sequence according to claim 4 or 5 or a recombinant nucleic acid molecule according to claim 7 or a recombinant vector molecule according to claim 8 or 9.

11. A process for expressing the protein according to any one of claims 1 to 4 comprising culturing a host cell according to claim 10.

12. A vaccine for the protection of poultry against coccidiosis characterised in that it comprises a protein according to any one of claims 1 to 4, a recombinant nucleic acid molecule according to claim 6, a recombinant vector according to claim 7 or claim 8, or a host cell or organism according to claim 9 together with a pharmaceutically acceptable carrier.

13. A vaccine for the protection of poultry against coccidiosis, substantially as hereinbefore described with reference to any one of the examples.

14. A process for the preparation of a coccidiosis vaccine comprising the steps of culturing an infected host cell according to claim 10, collecting the recombinant vector and formulating said recombinant vector into a pharmaceutical preparation with immunising activity.

15. A process for the preparation of a coccidiosis vaccine comprising formulating a protein according to any one of claims 1 to 4 or a protein prepared according to the process of claim 11 into a pharmaceutical preparation with immunising activity.

AUSTRALIAN PATENT OFFICE

16. A process for the preparation of a coccidiosis vaccine, substantially as hereinbefore described with reference to any one of the examples.

17. An antibody or antiserum immunoreactive with a protein according to any one of claims 1 to 4.

18. A method for the protection of poultry against coccidiosis comprising administering a vaccine according to claim 12 to the poultry.

19. Use of a protein according to any one of claims 1 to 4, a recombinant nucleic acid molecule according to claim 6, a recombinant vector according to claim 7 or 8 or a host cell or organism according to claim 9 in the preparation of a vaccine for the protection of poultry against coccidiosis.

20. The protein according to any one of claims 1 to 4, the recombinant nucleic acid molecule according to claim 6, the recombinant vector according to claim 7 or 8 or the host cell or organism according to claim 9 when used in the protection of poultry against coccidiosis.

21. A coccidiosis vaccine when prepared by the process of claim 15 or 16.

22. A method for the protection of poultry against coccidiosis comprising administering a vaccine according to claim 13 or 21 to the poultry.

Dated 7 May 1999
Akzo Nobel N.V.

Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON

AUSTRALIAN PATENT OFFICE



Figure 1



Figure 2