


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 47/48, C07D 305/14	A1	(11) International Publication Number: WO 94/00156 (43) International Publication Date: 6 January 1994 (06.01.94)
(21) International Application Number: PCT/EP93/01433 (22) International Filing Date: 7 June 1993 (07.06.93) (30) Priority data: 9213077.2 19 June 1992 (19.06.92) GB (71) Applicant: FARMITALIA CARLO ERBA S.R.L. [IT/IT]; Via Carlo Imbonati, 24, I-20159 Milan (IT). (72) Inventors: MONGELLI, Nicola ; Via Tertulliano, 38, I-20137 Milan (IT). ANGELUCCI, Francesco ; Via G. Washington, 106, I-20146 Milan (IT). PESENTI, Enrico ; Viale Visconti, 9, I-20093 Cologno Monzese (IT). SUARATO, Antonio ; Via Degli Imbriani, 39, I-20158 Milan (IT). BIASOLI, Giovanni ; Via Scalarini, 31, I-21046 Gavirate (IT).		(74) Agent: FERRARIO, Vittorino; R.S./Documentation and Patent Department, Farmitalia Carlo Erba S.r.l., Via Carlo Imbonati, 24, I-20159 Milan (IT). (81) Designated States: AU, BY, CA, CZ, FI, HU, JP, KR, KZ, NO, NZ, PL, RU, UA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>

(54) Title: POLYMER-BOUND PACLITAXEL DERIVATIVES**(57) Abstract**

A polymer conjugate consisting essentially of: from 90 to 99.9 mol % of units represented by formula (a); from 0.1 to 5 mol % of units represented by formula (b), wherein one of R₁ and R₂ is a copolymer residue of formula (c), and the other one is hydrogen atom; from 0 to 9.9 mol % of units represented by formula (d), wherein R is a phenyl or t-butoxy group, R₃ is H or an acetyl group, A and A₁ which may be the same or different, represent a chemical single bond, an amino acid residue or peptide spacer selected from β Ala, Gly, Phe-Gly, Phe-Phe-, Leu-Gly, Val-Ala, Phe-Ala, Leu-Phe, Leu-Ala, Phe-Leu-Gly, Phe-Phe-Leu, Leu-Leu-Gly, Phe-Tyr-Ala, Phe-Gly-Phe, Phe-Phe-Gly, Phe-Leu-Gly-Phe, Gly-Phe-Leu-Gly-Phe, Gly-βAla, Phe-Gly-βAla, Phe-Phe-βAla, Leu-Gly-βAla, Val-Ala-βAla, Phe-Ala-βAla, Leu-Phe-βAla, Leu-Gly-βAla, Phe-Leu-Gly-βAla, Phe-Phe-Leu-βAla, Leu-Leu-Gly-βAla, Phe-Tyr-Ala-βAla, Phe-Gly-Phe-β, Phe-Phe-GlyβAla, Phe-Leu-Gly-Phe-βAla or Gly-Phe-Leu-Gly-Phe-βAla. The compounds are endowed with antitumor activity and show improved water solubility and decreased toxicity in comparison with paclitaxel or its known analogs. A method for their preparation and pharmaceutical compositions containing them are also described.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				


- 1 -

POLYMER-BOUND PACLITAXEL DERIVATIVES

The present invention is directed to polymer-bound paclitaxel and polymer-bound paclitaxel derivatives endowed with antitumor activity, to a method for their preparation and to pharmaceutical compositions containing them.


Paclitaxel (also named Taxol in several publications) is a member of the taxane family of diterpenes, isolated and characterized from an extract of bark of Taxus brevifolia L.; other analogues of paclitaxel are also known and were prepared by semisynthesis starting from 10-deacetyl baccatin III, extracted from the needles of Taxus baccata L., see Wain et al. in JACS, 93, 2325 (1971) and Lovelle et al., Proc. Am. Assoc. Cancer Res., 31, p. 417, (1990). These compounds have been shown to possess potent antitumor activity, but they are very slightly soluble in water and some toxic side-effects are associated also with their administration by injection or intravenous infusion using as carrier cremophor EL (TM). The present invention provides a polymer conjugate of the formula I consisting essentially of:

from 90 to 99.9 mol % of units of the formula:




from 0.1 to 5 mol % units represented by the formula

- 2 -




wherein one of R₁ and R₂ is a copolymer residue of the formula



5 and the other one is hydrogen atom;

from 0 to 9.9 mol % of units represented by the formula



10 wherein R is a phenyl or t-butoxy group, R₃ is H or an acetyl group, A and A₁ which may be the same or different, represent a chemical single bond, an amino acid residue or peptide spacer selected from β Ala, Gly, Phe-Gly, Phe-Phe-, Leu-Gly, Val-Ala, Phe-Ala, Leu-Phe, Leu-Ala, Phe-Leu-Gly, Phe-Phe-Leu, Leu-Leu-Gly, Phe-Tyr-Ala, Phe-Gly-Phe, Phe-Phe-Gly, Phe-Leu-Gly-Phe, Gly-Phe-Leu-Gly-Phe, Gly-βAla, Phe-Gly-βAla, Phe-Phe-βAla,

15

- 3 -

Leu-Gly- β Ala, Val-Ala- β Ala, Phe-Ala- β Ala, Leu-Phe- β Ala,
Leu-Gly- β Ala, Phe-Leu-Gly- β Ala, Phe-Phe-Leu β Ala, Leu-
Leu-Gly- β Ala, Phe-Tyr-Ala- β Ala, Phe-Gly-Phe- β , Phe-Phe-
Gly- β Ala, Phe-Leu-Gly-Phe- β Ala or Gly-Phe-Leu-Gly-Phe-
5 β Ala.

More particularly the present invention provides polymer
conjugates of paclitaxel and derivatives of paclitaxel
with improved water solubility and decreased toxicity.

Preferably, the mol % of units containing the paclitaxel
10 and paclitaxel derivatives is from 0.5 to 2, more
preferably, the content of paclitaxel in the polymer was
from 2 to 10 % (w/w), most preferred compounds are those
characterized by a content of from 4 to 7 % (w/w). The
wavy line denotes that the oxygen linked at position 7 of
15 the paclitaxel structure may be in both configurations,
i.e. β (natural) or α .

Preferably R represents a phenyl group, R₃ is an acetyl
group and A is a Phe-Leu-Gly or Phe-Leu-Gly- β Ala residue.
All the amino acid residues have the natural L-
20 configuration. According to a preferred embodiment, the
polymer conjugate is a copolymer of 1-methacryloylamino-
2-hydroxypropane, (methacryloylglycyl-phenylalanyl-
leucylglycyl)3-amino-2 hydroxypropane and 2'-
(methacryloylglycylphenylalanylleucylglycyl- β alanyl)
25 paclitaxel.

The % w/w content was determined after enzymatic
hydrolysis by HPLC method analogous to that described in
Cancer Treatment Rep. 71 (1), 1987, p. 53-59.

- 4 -

Enzymatic hydrolysis

To 1 ml of murine or human plasmas various concentrations of the polymer-bound paclitaxel were added and at appropriate time (24, 48, 72, 96 h) 100 μ l were collected and stored at -70°C until further processing.

Extraction procedure


Samples were extracted by adding 75 μ l of Tetra Butyl Ammonium Phosphate (TBAP) 0.5 M, 1250 μ l CH_3CN and 150 μ l of NaCl 5M and vigorously shaken for 20' at 4°C . After that time samples were spun at 15000 x g for 10' and the supernatants were collected and evaporated using a high vacuum centrifuge . Samples were recovered by adding 500 μ l of MeOH:H₂O (75:25 V/V) and injected into HPLC for determining the total paclitaxel percentage content.

HPLC system


Column : Nova Pak C₁₈ (Waters) 3.9 x 300 mm
Flow Rate : 1.5 ml/m
Detector : UV 231 nm
Injection : 20 μ l
Mobile Phase : 57.5 % H₂O brought to pH 2
42.5 % CH₃CN

The present invention also provides a process for preparing a polymer conjugate which process comprises reacting a compound of the formula II


- 5 -



wherein one of A_2 and A_3 is a chemical bond and the other one is A, and A, R and R_3 are as defined above, with an activated polymer consisting essentially of from 90 to 99.9 mol % of units represented by formula



and from 10 to 0.1 mol % of units represented by the formula



wherein A_1 is as defined above and then treating the resultant polymer conjugate with 2-hydroxypropylamine.

15 Preferably the reaction between the compounds of the formula II and the activated polymer is carried out in an anhydrous polar organic solvent such as dimethylsulfoxide or dimethylformamide optionally in presence of an organic or inorganic base such as an alkaline carbonate, dimethylaminopyridine or

- 6 -

triethylamine. The reaction can typically be effected for from 1 to 24 hours.

The reaction is typically carried out at a temperature of from 15 to 40°C, preferably at room temperature. The
5 alkaline carbonate is, for example, an alkali metal carbonate or an alkaline earth metal carbonate.

Some of the starting compounds of the formula II are known compounds, i.e. paclitaxel or paclitaxel analogues or may be prepared starting from known compounds.

10 For example 7-epi derivatives may be prepared by refluxing in toluene paclitaxel or its analogs in the presence of a base (Na_2CO_3 or diazabicycloundecene)

Other compounds of the formula II are new, in particular those in which A_3 is β Ala residue and those in which
15 either A_2 or A_3 represents di, tri, or tetra peptide spacer as defined above for A and are within the scope of the invention.

The compounds of formula II wherein A_2 is not a chemical single bond may be prepared by reacting a paclitaxel or a
20 paclitaxel analog with protected amino acid or peptide in the presence of a condensating reagent, and with or without the additional presence of a catalyst, preferably at room temperature, followed by the removal of the protecting group with known methods.

25 The condensation may be also carried out using activated esters such as paranitrophenyl ester of peptide or amino acid. Suitable condensing reagents include carbodiimides such as dicyclohexyl carbodiimide (DCC).

- 7 -

Suitable catalysts include 4-dimethylamino-pyridine (DMAP), pyridine or triethylamine.

Various known amino protecting groups can be utilized and commercially available protected amino acid or peptides can be utilized as the starting materials. Amino acids or peptides protected with t-BOC, trityl, Fmoc or carbobenzyloxy (CBZ) can be utilized. Amino acid or peptides protected with t-BOC, trityl or Fmoc groups are preferred.

10 The compounds of the formula II wherein A_3 is not a chemical single bond may be prepared by protecting or blocking the 2'-hydroxy group and then esterifying the 7-position hydroxyl and then removing the 2'-protecting or blocking group.

15 More preferably, the compounds of the formula II wherein A_3 is Gly or β Ala residue are prepared by reacting paclitaxel with 2-3 equivalents of N-protected amino acid to produce 2',7-disubstituted paclitaxel, the 2'-position amino acid is cleaved and then the 7-position amino acid is deprotected.

20 Reaction of paclitaxel and the protected amino acid is conducted in the presence of a condensing reagent and a catalyst, like those above defined.

Cleavage of the 2'-amino acid is conducted by adjusting the pH of the 2'-7-(amino acid) paclitaxel solution to pH 25 7-7.4 for example by mixture of the 2',7-di(amino acid) paclitaxel in a phosphate buffer at pH 7-7.4 or with a slight excess of NaHCO_3 .

- 8 -

Deprotection of the amino acid is conducted under a known amino acid deprotection method, such as mild acid treatment with, for example, acetic acid, or by reduction.

5 Thus, for example, paclitaxel is allowed to react with 2-3 mol. equivalent of N-protected amino acid (t.boc, CBZ or FMOC protected) in methylene chloride in the presence of DCC and a catalytic amount of 4-dimethylaminopyridine. In this manner, the protected amino acid is introduced at
10 2' and 7-position. The 2',7-bis amino acid derivative of paclitaxel is allowed to stand in presence of NaHCO₃ in H₂O/MeOH for 2-5 hours, whereby selective deprotection at the 2'-position occurs to yield the 7-substituted derivative of paclitaxel. The protecting groups are
15 removed by appropriate deprotecting agent (e.g., acid, mild base or hydrogenolysis).

The activated polymer is a synthetic, water soluble polymer prepared by the copolymerization of N'-(2-hydroxypropyl) methacrylamide with p-nitrophenylesters
20 of N-methacryloyl oligopeptides, as described in US-A-4062831 and US-A-4097470.

The polymer conjugates of the formula I and the new paclitaxel derivatives of the formula II exhibit good water solubility, biocompatibility and release the
25 paclitaxel or paclitaxel derivative in the plasma or after internalization into cells by cleaving of the oligopeptide spacers.

- 9 -

Biological activityMicrotubule assembly and disassembly assay

5 Calf brain tubulin was prepared by two cycles of assembly-disassembly (Shelanski M.L. Gaskin F. and Cantor C.R., Proc. Natl. Acad.Sci. U.S.A. 70, 765-768, 1973) and stored in liquid nitrogen in MAB (0.1 M MES, 2.5 mM EGTA, 0.5 mM MgSO₄, 0.1 mM EDTA, 0.1 mM DTT pH 6.4).

All the experiments were carried out on protein stored for less than 4 weeks.

10 Before each experiment, the tubulin was kept 30 min at 4°C.

Assembly was monitored by the method of Gaskin et al (Gaskin F., Cantor C.R. Shelanski M.L. J. Molec . Biol 89, 737-758, 1974). The cuvette (1 cm path) containing 15 tubulin (1 mg/ml) and 1 mM GTP was shifted to 37°C and continuous turbidity measurements were made at 340 nm on Perkin-Elmer 557 Double Wavelength Double Beam Spectrophotometer equipped with an automatic recorder and a thermostatically regulated sample chamber.

20 After 30 minutes 4 mM CaCl₂ was added and depolymerisation was measured for 10 minutes as decreased turbidity. At regular intervals of 15 minutes scalar doses of the tested compounds were added and variations in the turbidity were monitored. Data are expressed as 25 percentage of ripolymerisation induced by the tested compounds. The results are shown in Table I.

- 10 -

In vitro drug sensitivity assay

Exponentially growing B16-F10 murine melanoma cells were seeded (2×10^4 /ml) in RPMI 1640 medium were supplemented with 10% heat-inactivated fetal calf serum and 2 mM glutamine in 24 well-plates (Costar). Scalar concentrations of tested compounds were added immediately after seeding. The inhibition of cell growth was evaluated by counting cells with a coulter counter after 72 hrs incubation. For each tested compound concentration triplicate cultures were used. The antiproliferative activity of the tested compounds was calculated from dose-response curves and expressed as IC_{50} (dose causing 50% inhibition cell growth in treated cultures relative to untreated controls). The result are shown in Table I.

15

Table I

20

Compound prepared in	Tubulin Assembly %		IC_{50} (μ M)
	0.5 μ M	5 μ M	
Example 1	0	15	19
Example 9	79	145	47
Example 3	46	86	23
Example 4	0	10	11
Example 6	0	10	51
Reference compound :			
Paclitaxel	41	96	39

- 11 -

The copolymer-paclitaxel prepared in Example 6 was tested in vivo against B16 F10 murine melanoma in comparison with paclitaxel.

Mice

5 C57B16 female mice were obtained from Charles River Italy.

Animals were 8 to 10 weeks old at the beginning of the experiment.

Drugs

10 Because of its limited aqueous solubility, paclitaxel was dissolved in a vehicle consisting of polyoxyethylated castor oil (Cremophor EL) 50% and ethanol 50% and then diluted with glucose 5% solution at the desired concentrations. The solution was slightly hazy and precipitates formation was observed after short time.

15 The compound of example 6 was easily dissolved in glucose 5% and the resulting solution was clear for long time (more than 2 hours). Final concentrations were referred to the paclitaxel content of the compound (4% of total).

20 Tumor

The murine melanoma B16F10 was used. A suspension of 10^5 tumor cells in 0.2 ml was injected subcutaneously in the flank of mice.

Tumor size was measured with caliper and tumor weight was
25 calculated with the formula:

$$\frac{a^2+b^2}{2}$$

- 12 -

Drugs administration

Paclitaxel was given intraperitoneally because of its poor solubility and vehicle toxicity.

The compound of example 6 was injected intravenously.

5 Both compounds were administered at day 1, 5, 9 after tumor implantation.

The data reported in Table 2 show that the compound of the present invention is more active than paclitaxel.

10 The dose of the polymer-conjugate is referred to the paclitaxel content.

Table 2

Compound	Dose (mg/Kg)	Tumor Inhibition (%)	Tox
Control	-	-	-
Paclitaxel	14.6	53	0/10
	22	38	0/10
	33	92	1/7
15 Compound of ex. 6	14.6	77	0/9
	22	79	0/10

TOX: number of mice which died for toxicity.

TOX determination was made when mice died before the control or when significant body weight loss and/or spleen and or/or liver size reduction were observed.

20 From the above data, it can be seen that the polymer

- 13 -

conjugates of the present invention exhibit excellent antitumor activity. These compounds, therefore, are useful antitumor agents due to the lower toxicity and increased water solubility as compared to paclitaxel or its derivative. Examples of tumors that can be treated are for instance, sarcomas, carcinoma, lymphomas, neuroblastoma, melanoma, myeloma, Wilms tumor, leukemias and adenocarcinoma.

The improved solubility and decreased toxicity of the polymer-conjugates of the present invention means that they are suitable for intravenous injection or infusion. The dosage depends upon the age, weight and condition of the patient. The dosage may be from 1 mg/kg body weight to 1 g/kg body weight, preferably from 4 to 800 mg/kg body weight. Typical formulations contain a quantity of polymer-bound paclitaxel/polymerbound paclitaxel derivative equivalent to 0.5, 1.5, 10, 20, 25, or 50 mg of the active paclitaxel/paclitaxel derivative.

The polymer conjugates may be formulated as pharmaceutical compositions with a pharmaceutically acceptable carrier or diluent. Any appropriate carrier or diluent may be used. The solutions for intravenous injection or infusion may contain as carrier or diluent, for example, sterile water or preferably they may be in the form of sterile, aqueous or isotonic saline solutions.

The following examples illustrate the invention.

- 14 -

Example 1

Copolymer of 1-methacryloylamino-2-hydroxypropane,
1 (methacryloyl-glycyl-phenylalanyl-leucyl-glycyl)amino-
2-hydroxypropane and 2'(methacryloyl-glycyl-phenyl -
5 alanyl-leucyl-glycyl)paclitaxel.

To a solution of 1.4 g of copolymer of 1-methacryloyl-
amino-2-hydroxypropane and N-(methylencarbonyl-Phe- Leu-
Gly 4-nitrophenoxy)methacrylamide, prepared according to
10 J. Kopecek et al., Makromol Chem 177, p. 2833 (1976), in
15 ml of anhydrous dimethylformamide were added 100 mg of
paclitaxel and 15 mg of dimethylaminopyridine.

The yellow solution was stirred for 8 hours at room
temperature under anhydrous conditions. Then 2-hydroxy
propylamine (0.2 ml) was dropped into reaction flask, and
15 then the whole was stirred for 30 minutes.

The reaction solution was quenched with 0.3 ml glacial
acetic acid, concentrated under vacuum to a small volume,
and then poured into 200 ml of acetone.

Ater 30' mixing, the precipitate was filtered and washed
20 with acetone to yield 1.25 g of the title compound.

The paclitaxel content was 4.5% (evaluated by enzymatic
hydrolysis and HPLC analysis).

Unreacted paclitaxel was recovered from acetone solution.

Example 2

25 2'(N-trityl-phenylalanyl-leucyl-glycyl)paclitaxel

To a solution of 170 mg of paclitaxel in 16 ml of
acetonitrile were added 24 mg of dimethylamino-pyridine
and 150 mg of N-trityl-phenylanyl-leucyl-glycine 4-

- 15 -

nitrophenyl ester.

The yellow solution was stirred for 20 hours at room temperature, and then evaporated under vacuum to dryness.

The residue was chromatographed on silica gel with ethyl-
5 acetate-hexane 35:25 as eluant, affording 380 mg of the
title compound.

$^1\text{H-NMR}$ (400 MHz, CDCl_3): δ

- 0.82 (d, $J=6.4$ Hz, 3H, δ -Leu)
0.85 (d, $J=6.7$ Hz, 3H, δ -Leu)
10 1.15 (s, 3H, 16)
1.26 (s, 3H, 17)
1.2-1.6 (m, 3H, $\beta+\beta+\gamma$ -Leu)
1.69 (s, 3H, 19)
1.85 (s, 1H, OH-1)
15 1.89 (m, 1H, 6β)
1.96 (d, $J=1.2$ Hz, 3H, 18)
2.14 (dd, $J=5.9$ Hz, $J=13.5$ Hz, 1H, β -Phe)
2.24 (s, 3H, $\text{CH}_3\text{CO-10}$)
2.2-2.7 (m, 5H, $\text{CH}_2\text{-14+OH-7+6}\alpha+\beta\text{Phe+NH-Phe}$)
20 2.47 (s, 3H, $\text{CH}_3\text{CO-4}$)
3.50 (m, 1H, α -Phe)
3.74 (dd, $J=4.7$ Hz, $J=18.2$ Hz, 1H, α -Gly)
3.80 (m, 1H, α -Leu)
3.83 (d, $J=7.0$ Hz, 1H, 3)
25 4.17 (dd, $J=7.0$ Hz, $J=18.2$ Hz, 1H, α' -Gly)
4.22, 4.33 (two-d, $J=8.5$ Hz, 2H, $\text{CH}_2\text{-20}$)
4.46 (m, 1H, 7)
4.97 (dd, $J=2.2$ Hz, $J=9.9$ Hz, 1H, 5)

- 16 -

- 5.44 (d, J=2.3 Hz, 1H, 2')
5.71 (d, J=7.0 Hz, 1H, 2)
5.97 (dd, J=4.7 Hz, J=7.0 Hz, 1H, NH-Gly)
6.07 (dd, J=2.3 Hz, J=9.4 Hz, 1H, 3')
- 5 6.2-6.3 (m, 2H, 13+10)
6.8-9.2 (m, 30 H, 6- Ph)
6.95 (d, J=6.7 Hz, 1H, NH-Leu)
8.00 (d, J=9.4 Hz, 1H, NH-4')

Example 3

10 2' (phenylalanyl-leucyl-glycyl)paclitaxel.

The compound 2' (N-Trit-Phe-Leu-Gly)paclitaxel (250 mg) was dissolved into a mixture of glacial acetic (22 ml) and water (6 ml) and the whole was stirred for 1 hour at room temperature.

15 The solvents were evaporated in vacuum to the dryness, the residue stirred with diethylether-hexane 1:1 for 30 minutes and filtered to obtain 160 mg of the title compound.

FAB-MS: m/z 1171, $M+H^+$; 1112, $M-CH_3COOH + 2H$; 1051,
20 1024, 911, 603, 569, 509.

1H -NMR (400MHz, $CDCl_3$): δ

- 0.88 (d, J=6.4Hz, 3H, δ Leu)
0.92 (d, J=6.4Hz, 3H, δ' Leu)
1.13 (s, 3H, 16)
25 1.16 (s, 3H, 17)
1.4-2.0 (m, 4H, $\beta+\beta'+\gamma$ Leu+6 β)
1.69 (s, 3H, 19)
1.91 (d, J=1.2Hz, 3H, 18)

- 17 -

- 2.16 (dd, J=6.0 Hz, J=13.8Hz, 1H, 14)
2.23 (s, 3H, COCH₃-10)
2.4-2.6 (m, 3H, 6 α +14+ β Phe)
2.53 (s, 3H, COCH₃-4)
5 2.90 (dd, J=4.1Hz, J=13.5Hz, 1H, β' -Phe)
3.49 (dd, J=4.1 Hz, J=9.1Hz, 1H, α Phe)
3.82 (d, J=7.3Hz, 1H, 3)
3.9-4.1 (m, 2H, α + α' Gly)
4.22, 4.33 (two-d, J=8.7 Hz, 2H, CH₂-20)
10 4.27 (m, 1H, α -Leu)
4.44 (dd, J=6.4Hz, J=10.8Hz, 7)
4.98 (dd, J=2.4Hz, J=9.7 Hz, 5)
5.61 (d, J=3.2Hz, 1H, 2')
5.70 (d, J=7.3Hz, 1H, 2)
15 6.12 (dd, J=3.2Hz, J=9.4Hz, 1H, 3')
6.21 (m, 1H, 13)
6.28 (s, 1H, 10)
6.8-8.2 (m, 21H, 4-Ph + NHLeu)
7.87 (d, J=9.4Hz, 1H, NH-4')

20 **Example 4**

Copolymer of 1-methacryloylamino-2-hydroxypropane, 1-(methacryloyl-glycyl)amino-2-hydroxypropane and 2' (methacryloyl-glycyl-phenylalanyl-leucyl-glycyl)paclitaxel.

- 25 To a solution of 1 g of copolymer of 1-methacryloylamino-2-hydroxypropane and N-(methylenecarbonyl-4-nitro - phenoxy)methacrylamide, prepared according to P. Rejmanova et al., Makromol. Chem. 178, p. 2159-2168, in

- 18 -

10 ml of anhydro dimethylformamide were added 100 mg of
2' Phe-Leu-Gly-paclitaxel and 10 mg of
dimethylaminopyridine.

The yellow solution was stirred for 2 hours at room
5 temperature under anhydrous conditions. Then 2-hydroxy -
propylamine (0.15 ml) was added to the reaction solution,
and the whole was stirred for 30 minutes. The reaction
solution was quenched with 0.2 ml of glacial acetic acid,
concentrated under vacuum to a small volume and then
10 poured into 200 ml of acetone.

The mixture was stirred for 1 hours, the precipitate was
filtered and washed with acetone to yield 960 mg of the
title compound.

The paclitaxel content was 6% (evaluated by enzymatic
15 hydrolysis and HPLC analysis).

Example 5

Copolymer of 1-methacryloylamino-2-hydroxy
-propane, 1(methacryloyl-glycyl)amino-2-hydroxypropane
and 2'(methacryloyl-glycyl)paclitaxel.

20 To a solution of 1.6 g of copolymer of 1-methacryloyl -
amino-2-hydroxypropane and N-(methylen-carbonyl-4-nitro-
phenoxy)methacrylamide in 16 ml of anhydrous dimethyl-
formamide were added 100 mg of paclitaxel and 20 mg of
dimethylamino-pyridine. The yellow solution was stirred
25 for 20 hours at room temperature, then 2-
hydroxypropylamine (0.2 ml) was added, and the whole was
stirred for 30 minutes. The reaction solution was
quenched with 0.3 ml of glacial acetic acid, concentrated

- 19 -

under vacuum to a small volume, and then poured into 200 ml of acetone. The mixture was stirred for 1 hour, the precipitate was filtered and washed with acetone to yield 1440 mg of the title compound.

5 The paclitaxel content was 2.75% w/w%

Example 6

Copolymer of 1-methacryloylamino -2-hydroxypropane, 1-(methacryloyl-glycyl-phenylanyl-leucyl-glycyl)amino-2-hydroxypropane and 2'(methacryloyl-glycyl-phenylalanyl-leucyl-glycyl- β alanyl)paclitaxel.

To a solution of 620 mg of copolymer of 1-metacrylamino-2-hydroxypropane and N-(methylenecarbonyl-Phe-Leu-Gly-4-nitro- phenoxy)methacrylamide in 6 ml of anhydrous dimethyl-formamide were added 62 mg of 2'-(β alanyl)paclitaxel, prepared according to N.F. Magri et al, J Nat. Products 51 298-306, 1988, and 10 mg of dimethylamino pyridine. The yellow solution was stirred for 5 hours at room temperature under anhydrous conditions.

20 Then 2-hydroxypropylamine (0.1 ml) was added and the whole was stirred for 30 minutes.

The reaction solution was quenched with 0,15 ml of glacial acetic acid, concentrated under vacuum to a small volume and poured in 150 ml of acetone. The mixture was stirred for 1 hour, the precipitate was filtered and washed with acetone to yield 5.85 mg of the title compound.

The paclitaxel content was 4% w/w%

- 20 -

Example 72',7-di(carbobenzyloxy- β -alanyl)paclitaxel

To a solution of 200 mg of paclitaxel in 15 ml of acetonitrile were added 400 mg of N,N'-
5 dicyclohexylcarbodiimide, 200 mg of carbobenzyloxy- β -alanine and 60 mg of dimethylamino-pyridine. The reaction mixture was stirred for 20 hours, the precipitate was filtered and the solvent evaporated under vacuum to dryness.

10 The residue was chromatographed on silica gel with ethyl acetate-hexane 1:1 as eluant, affording 300 mg of the title compound.

FAB-MS: m/z 1264 $M+H^+$, 1204, 1130, 1070

- 21 -

Example 87-(carbobenzyloxy- β alanyl)paclitaxel

To a solution of 171 mg of 2'7-di(carbobenzyloxy- β -
alanyl)paclitaxel in 60 ml of methanol, were added 30 mg
5 of sodium bicarbonate and 7 ml of water . The reaction
mixture was stirred for 3 hours at room temperature.

The methanol was evaporated and the product extracted
with ethylacetate.

The solvent was evaporated under vacuum to dryness,
10 affording 134 mg of the title compound.

Example 97-(β alanyl)paclitaxel

To a solution of 135 mg of 7-(carbobenzyloxy-
 β alanyl)paclitaxel in 20 ml of methanol and 13 ml of
15 formic acid, was added 200 mg of Pd/C 5%. The reaction
mixture was stirred for 6 hours at room temperature. The
catalyst was filtered, washed with methanol and the
solvents were evaporated to dryness under vacuum. The
residue was dissolved in 8 ml of methanol and
20 precipitated with 150 ml of diethylether, affording 85 mg
of the title compound.

FAB-MS: M/z 925, $M+H^+$; 947, $M+Na^+$ 1H NMR(400MHz, $CDCl_3$): δ 1.14 (s, 3H, CH_3 -16)25 1.20 (s, 3H, CH_3 -17)1.79 (s, 3H, CH_3 -19)1.85 (s, 3H, CH_3 -18)

- 22 -

- 2.17 (s, 3H, CH₃CO-10)
 2.2-2.6 (m, 6H, CH₂-14+CH₂-6+OCOCH₂CH₂NH₂)
 2.42 (s, 3H, CH₃CO-4)
 3.0-3.2 (m, 2H, OCOCH₂CH₂NH₂)
 5 3.90 (d, J=6.8Hz, 1H, 3)
 4.18, 4.31 (two d, J=8.2Hz, 2H, CH₂-20)
 4.80 (d, J=3.2Hz, 1H, 2')
 4.91 (d, J=8.5Hz, 1H, 5)
 5.62 (dd, J=10.2Hz, J=7.0Hz, 1H, 7)
 10 5.66 (d, J=6.8Hz, 1H, 2)
 5.81 (dd, J=2.9Hz, J=9.1Hz, 1H, 3')
 6.17 (m, 1H, 13)
 6.19 (s, 1H, 10)
 7.3-8.2 (m, 16H, NH-4'+3-Ph)

15 **Example 10**

Copolymer of 1-methacryloylamino-2-hydroxypropane, 1-(methacryloyl-glycyl-phenylalanyl-leucyl-glycyl)amino-2-hydroxypropane and 7-(methacryloyl-glycyl-phenylalanyl-leucyl-glycyl-βalanyl)paclitaxel.

- 20 To a solution of 1500 mg of copolymer of 1-methacryloylamino-2-hydroxypropane and N-(methylencarbonyl-Phe-Leu-Gly-4nitrophenoxy)methacrylamide in 13 ml of anhydrous dimethylformamide were added 135 mg of 7-(βalanyl)paclitaxel and 20 mg of dimethylaminopyridine.
- 25 The yellow solution was stirred under anhydrous conditions for 5 hours at room temperature.

Then 2-hydroxypropylamine (0.2 ml) was added, and the

- 23 -

whole was stirred for 30 minutes.


The reaction solution was quenched with 0.3 ml of glacial acetic acid, concentrated under vacuum to a small volume and poured in 250 ml of acetone. The mixture was stirred
5 for 1 hour, the precipitate was filtered and washed with acetone to yield 1520 mg of the title compound.

The paclitaxel content was 7.8% w/w%.


CLAIMS

1. A polymer conjugate consisting essentially of: from 90 to 99.9 mol % of units represented by the formula

5




from 0.1 to 5 mol % of units represented by the formula



10

wherein one of R_1 and R_2 is a copolymer residue of the formula




and the other one is hydrogen atom;

15

from 0 to 9.9 mol % of units represented by the formula

- 25 -




wherein R is a phenyl or t-butoxy group, R₃ is H or an
 acetyl group, A and A₁ which may be the same or
 5 different, represent a chemical single bond, an amino
 acid residue or peptide spacer selected from β Ala,
 Gly, Phe-Gly, Phe-Phe-, Leu-Gly, Val-Ala, Phe-Ala, Leu-
 Phe, Leu-Ala, Phe-Leu-Gly, Phe-Phe-Leu, Leu-Leu-Gly, Phe-
 Tyr-Ala, Phe-Gly-Phe, Phe-Phe-Gly, Phe-Leu-Gly-Phe, Gly-
 10 Phe-Leu-Gly-Phe, Gly-βAla, Phe-Gly-βAla, Phe-Phe-βAla,
 Leu-Gly-βAla, Val-Ala-βAla, Phe-Ala-βAla, Leu-Phe-βAla,
 Leu-Gly-βAla, Phe-Leu-Gly-βAla, Phe-Phe-Leu βAla, Leu-
 Leu-Gly-βAla, Phe-Tyr-Ala-βAla, Phe-Gly-Phe-β, Phe-Phe-
 Gly βAla, Phe-Leu-Gly-Phe-βAla or Gly-Phe-Leu-Gly-Phe-
 15 βAla.


2. A polymer conjugate according to claim 1, in which
 R represents a phenyl group, R₃ is an acetyl group and A
 is a Phe-Leu-Gly or Phe-Leu-Gly-βAla residue.

3. A polymer conjugate according claim 1 or 2, which
 20 is a copolymer of 1-methacryloylamino-2-hydroxypropane,
 (methacryloylglycyl-phenylalanyl-leucylglycyl)3-amino-2-
 hydroxypropane and 2'-(methacryloylglycylphenylalanyl-
 leucylglycyl-βalanyl) paclitaxel.


4. A process for preparing a polymer conjugate as defined in claim 1, which process comprises reacting a compound of the formula II



wherein one of A_2 and A_3 is a chemical bond and the other one is A, and A, R and R_3 are as defined in claim 1, with an activated polymer consisting essentially of from 90 to 99.9 mol % of units represented by formula



and from 10 to 0.1 mol % of units represented by the formula



wherein A_1 is as defined in claim 1 and then treating the resultant polymer conjugate with 2-hydroxypropylamine.

5. A process according to claim 4, in which the reaction is carried out in an anhydrous polar solvent at

- 27 -

a temperature of from 15 to 40°C for from 1 to 24 hours, in the presence of an organic or inorganic base.

6. A compound of the formula II as defined in claim 4 wherein A₂ represents di, tri or tetra peptide spacer as defined in claim 1 for A.

7. A process for preparing a compound of the formula II as defined in claim 6, which process comprises reacting a paclitaxel or a paclitaxel analog with desired protected amino acid or peptide in the presence of a condensating reagent or using activated esters and with or without the additional presence of a catalyst and then removing the protecting groups.

8. A compound of formula II as defined in claim 4 wherein A₃ represents βAla or di, tri or tetra peptide spacer as defined in claim 1 for A.

9. A process for preparing a compound of the formula II as defined in claim 8, which process comprises either (i) protecting or blocking the 2'-hydroxy group of paclitaxel and esterifying the 7-position hydroxyl and then removing the 2'-protecting or blocking group, or (ii) reacting paclitaxel with 2-3 moles of the desired N-protected amino acid to produce 2',7-disubstituted taxol, cleaving the 2'-position amino acid and then deprotecting the 7-position amino acid.

10. A pharmaceutical composition comprising, as active ingredient, a polymer conjugate as defined in claim 1, 6 or 8 and a pharmaceutically acceptable diluent or carrier.

INTERNATIONAL SEARCH REPORT

PCT/EP 93/01433

International Application No

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all)⁶

According to International Patent Classification (IPC) or to both National Classification and IPC
 Int.Cl. 5 A61K47/48; C07D305/14

II. FIELDS SEARCHED

Minimum Documentation Searched⁷

Classification System

Classification Symbols

Int.Cl. 5

A61K ; C07D

Documentation Searched other than Minimum Documentation
 to the Extent that such Documents are Included in the Fields Searched⁸

III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹

Category¹⁰ Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹² Relevant to Claim No.¹³

A	US,A,4 097 470 (J. DROBNIK ET AL.) 27 June 1978 cited in the application see claims; example D	1-10
X,Y	MAKROMOL. CHEM. vol. 178, 1977, pages 2169 - 2183 J. KOPECEK 'REACTIVE COPOLYMERS OF N-(2HYDROXYPROPYL)METHACRYLAMIDE WITH N-METHACRYLOYLATED DERIVATIVES OF L-LEUCINE AND L-PHENYLALANINE, 1' see page 2171; figures	1-10

¹⁰ Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

IV. CERTIFICATION

3

Date of the Actual Completion of the International Search
 11 OCTOBER 1993

Date of Mailing of this International Search Report
 12. 10. 93

International Searching Authority
 EUROPEAN PATENT OFFICE

Signature of Authorized Officer
 BERTE M.J.

III. DOCUMENTS CONSIDERED TO BE RELEVANT

(CONTINUED FROM THE SECOND SHEET)

Category °	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	<p>JOURNAL OF CONTROLLED RELEASE vol. 18, no. 2, February 1992, pages 123 - 132 V. SUBR ET AL. 'POLYMERS CONTAINING ENZYMATICALLY DEGRADABLE BONDS, XII. EFFECT OF SPACER STRUCTURE ON THE RATE OF RELEASE OF DAUNOMYCIN AND ADRIAMYCIN FROM HMPA COPOLYMER DRUG CARRIERS IN VITRO AND ANTITUMOUR ACTIVITY MEASURED IN VIVO.'</p>	1-10
Y	<p>EP,A,0 187 547 (CARLTON MEDICAL PRODUCTS LTD.) 16 July 1986 see page 9, line 14 - line 21; claims</p>	1-10
Y	<p>J. MED. CHEM. vol. 35, 1992, pages 145 - 151 A. E. MATHEW ET AL. 'SYNTHESIS AND EVALUATION OF SOME WATER-SOLUBLE PRODRUGS AND DERIVATIVES OF TAXOL WITH ANTITUMOR ACTIVITY.' cited in the application</p>	1-10
A	<p>EP,A,0 393 575 (SEARLE & CO.) 24 October 1990 see page 4, line 10 - page 5, line 38 see page 7, line 57; claims</p>	1-10
Y	<p>JOURNAL OF CONTROLLED RELEASE vol. 16, 1991, pages 121 - 136 R. DUNCAN ET AL. 'MACROMOLECULAR PRODRUGS FOR USE IN TARGETED CANCER CHEMOTHERAPY: MELPHALAN COVALENTLY COUPLED TO HMPA COPOLYMERS.' see figure 1; table 1</p>	1-10
A,P	<p>EP,A,0 537 905 (VIRGINIA TECH. INTELLECT. PROPERTIES INC.) 21 April 1993</p>	1-10

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

EP 9301433
SA 74848

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 11/10/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4097470	27-06-78	None	
EP-A-0187547	16-07-86	AU-B- 589587 AU-A- 5183386 CA-A- 1305053 JP-A- 61243026 US-A- 5037883	19-10-89 10-07-86 14-07-92 29-10-86 06-08-91
EP-A-0393575	24-10-90	CA-A- 2014732 JP-A- 2292227	17-10-90 03-12-90
EP-A-0537905	21-04-93	None	