



(86) Date de dépôt PCT/PCT Filing Date: 2009/02/27
(87) Date publication PCT/PCT Publication Date: 2009/09/03
(45) Date de délivrance/Issue Date: 2017/05/30
(85) Entrée phase nationale/National Entry: 2010/08/20
(86) N° demande PCT/PCT Application No.: AU 2009/000231
(87) N° publication PCT/PCT Publication No.: 2009/105824
(30) Priorités/Priorities: 2008/02/29 (US61/032,912);
2009/02/24 (US12/391,748)


(51) Cl.Int./Int.Cl. *A61K 31/551* (2006.01),
A61P 17/08 (2006.01), *A61P 17/10* (2006.01),
C07D 243/08 (2006.01), *C07D 401/04* (2006.01),
C07D 401/06 (2006.01), *C07D 401/12* (2006.01),
C07D 403/06 (2006.01), *C07D 403/12* (2006.01),
C07D 405/12 (2006.01), *C07D 409/12* (2006.01),
C07D 413/12 (2006.01), *C07D 413/14* (2006.01),
C07D 417/12 (2006.01)

(72) Inventeurs/Inventors:
BLASKOVICH, MARK ARNOLD THOMAS, AU;
CASSIDY, PETER JOSEPH, AU

(73) Propriétaire/Owner:
MIMETICA PTY LTD, AU

(74) Agent: SIM & MCBURNEY

(54) Titre : PROCEDES DE MODULATION DE L'ACTIVITE DU RECEPTEUR MC5 ET TRAITEMENT DE PATHOLOGIES ASSOCIEES AUDIT RECEPTEUR
(54) Title: METHODS OF MODULATING THE ACTIVITY OF THE MC5 RECEPTOR AND TREATMENT OF CONDITIONS RELATED TO THIS RECEPTOR



Formula (I)

(57) Abrégé/Abstract:

The present invention provides compounds of Formula (I) that are useful for modulating the biological activity of the melanocortin-5 receptor (MC5R). Compounds of this invention can be used to treat diseases and/or conditions in which down-regulation of MC5R is beneficial. Such diseases and/or conditions include, but are not limited to, acne, seborrhea, seborrheic dermatitis, cancer, and inflammatory diseases.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
3 September 2009 (03.09.2009)

PCT

(10) International Publication Number
WO 2009/105824 A1

(51) International Patent Classification:

A61K 31/551 (2006.01) **C07D 403/06** (2006.01)
C07D 401/06 (2006.01) **C07D 413/14** (2006.01)
C07D 409/12 (2006.01) **C07D 243/08** (2006.01)
A61P 17/08 (2006.01) **C07D 403/12** (2006.01)
C07D 401/12 (2006.01) **C07D 417/12** (2006.01)
C07D 413/12 (2006.01) **C07D 401/04** (2006.01)
A61P 17/10 (2006.01) **C07D 405/12** (2006.01)

(21) International Application Number:

PCT/AU2009/000231

(22) International Filing Date:

27 February 2009 (27.02.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/032,912 29 February 2008 (29.02.2008) US
 12/391,748 24 February 2009 (24.02.2009) US

(71) **Applicant** (for all designated States except US):
MIMETICA PTY LTD [AU/AU]; Level 2, 143 Corona-
 tion Drive, Milton, Queensland 4064 (AU).

(72) Inventors; and

(75) **Inventors/Applicants** (for US only): **BLASKOVICH, Mark, Arnold, Thomas** [CA/AU]; 17 Gerler Street, Bar-
 don, Queensland 4065 (AU). **CASSIDY, Peter, Joseph**
 [AU/AU]; 24 Cavell Terrace, Ashgrove, Queensland 4065
 (AU).

(74) **Agent:** **PHILLIPS ORMONDE FITZPATRICK**; Level
 21, 22 & 23, 367 Collins Street, Melbourne, Victoria
 3000 (AU).


(81) **Designated States** (unless otherwise indicated, for every
 kind of national protection available): AE, AG, AL, AM,
 AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,
 CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ,
 EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
 HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR,
 KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
 MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO,
 NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG,
 SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA,
 UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every
 kind of regional protection available): ARIPO (BW, GH,
 GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
 ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,
 TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
 ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
 MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR),
 OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML,
 MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) **Title:** METHODS OF MODULATING THE ACTIVITY OF THE MC5 RECEPTOR AND TREATMENT OF CONDI-
 TIONS RELATED TO THIS RECEPTOR



(57) **Abstract:** The present invention provides compounds of Formula (I) that are useful for modulating the biological activity of the melanocortin-5 receptor (MC5R). Compounds of this invention can be used to treat diseases and/or conditions in which down-regulation of MC5R is beneficial. Such diseases and/or conditions include, but are not limited to, acne, seborrhea, seborrheic dermatitis, cancer, and inflammatory diseases.

METHODS OF MODULATING THE ACTIVITY OF THE MC5 RECEPTOR AND
TREATMENT OF CONDITIONS RELATED TO THIS RECEPTOR

FIELD OF THE INVENTION

5 The present invention relates to methods of modulating the activity of the melanocortin-5 receptor. In particular the present invention relates to the use of a family of 1,4-diazepan-2-ones and derivatives thereof to modulate the activity of the melanocortin-5 receptor. The invention also relates to methods and uses of the compounds in the treatment of conditions in which antagonism of the melanocortin-5 receptor is beneficial.

BACKGROUND OF THE INVENTION

10 The melanocortin-5 receptor (MC5R) is a G-protein coupled receptor (GPCR) belonging to the family of melanocortin receptors. There are five melanocortin receptors that have been isolated and cloned to date: MC1R, MC2R, MC3R, MC4R and MC5R. The melanocortin receptors participate in a variety of physiologic functions, providing a number of opportunities for therapeutic intervention in physiologic processes through alteration (i.e., a statistically significant increase or decrease) or modulation (e.g., up-regulation or down-regulation) of melanocortin receptor signalling activity.

20 Reviews of the melanocortin receptors, and their potential as targets for therapy have been published (Wikberg 2001; Bohm 2006). The melanocortin receptor family members are regulated by natural peptide agonists such as ACTH and the melanocyte-stimulating hormones (α -, β -, γ -MSH) derived from pro-opiomelanocortin (POMC) and by peptide antagonists such as Agouti signal protein (ASP) and Agouti-related peptide (AGRP). The MC1R is widely expressed and is associated with pigmentation in melanocytes and with inflammation responses in many cells involved in the immune system. The MC2R differs from the other melanocortin receptors in that it binds only ACTH but not MSH ligands. It is highly expressed in the adrenal gland and controls corticosteroid synthesis. The MC3R is found in the brain, but also elsewhere in the body, and appears to play a role in the regulation of energy homeostasis, and possibly sexual dysfunction. The MC4R is found almost exclusively in the brain, with some reports of its presence elsewhere. It has been strongly associated with feeding control, and also implicated with sexual desire. The MC5R is widely expressed in peripheral tissues, particularly in the exocrine glands, with some receptor also expressed in the brain. Given the breadth of activity associated with the melanocortin receptors it is desirable when seeking to target one of these receptors to do so selectively in order to avoid side effects associated with antagonism or agonism of another receptor in this family.

35

The MC5R has been cloned and expressed from multiple species, including humans in 1993 (though called MC2 in this paper) (Chhajlani 1993), rat in 1994 (Griffon 1994) mice in 1994 (Gantz 1994; Labbé 1994) and in 1995 (Fathi 1995), canine (Houseknecht 2003),
5 rhesus monkey (Huang 2000), sheep (Barrett 1994), zebrafish (Ringholm 2002), goldfish (Cerdá-Reverter 2003), spiny dogfish (Klovins 2004), rainbow trout (Haitina 2004), and chicken (Ling 2004), with the MC5R gene also identified in pig (Kim 2000). Patents covering the MC5R sequence in humans (Wikberg 2002), mice (Yamada 1997), rhesus monkey (Fong 2003) and dogs (Houseknecht 2003) have been published.

10 The MC5R has been implicated in regulating sebum secretion by a number of studies, as summarized in 2006 (Zhang 2006). Mice lacking MC5R have reduced sebum production, as evidenced by a marked inability to shed water from their fur, and a reduced quantity of sebum isolated from their hair. Significantly, these mice were otherwise generally healthy,
15 with no readily visible abnormalities (appearance, behaviour, growth, muscle mass, adipose mass, reproduction, basal and stress-induced corticosterone, glucose and insulin levels) (Chen 1997). Further studies have identified reductions in pheromones, causing alterations in aggressive behaviours between mice (Caldwell 2002; Morgan 2004a; Morgan 2004b; Morgan 2006). Mice in which the POMC-derived peptide native ligands of MC5R have been
20 knocked out show a similar phenotype (Yaswen 1999). Rats injected with α -MSH had 30-37% increased rates of sebum production, while removal of the neurointermediate lobe (the source of MSH) caused a 35% decrease in sebum secretion, which was restored upon administration of α -MSH (Thody 1973). A synergistic effect between α -MSH and testosterone was observed in rats, with testosterone increasing sebaceous gland and cell
25 volumes (presumably via increased proliferation), α -MSH increasing dermal lipogenesis, and the combination increasing sebum secretion (Thody 1975a; Thody 1975b).

At a cellular level human sebocytes have been shown to express MC5R, via detection of MC5R transcripts in micro-dissected sebaceous glands (Thiboutot 2000), detection of
30 MC5R in human facial sebaceous glands by immunostaining (Hatta 2001), detection of MC5R mRNA and MC5R in human sebaceous glands, cultured human sebocytes and rat preputial cells (Thiboutot 2000) and detection of MC5R as punctate particles within sebaceous glands by staining with polyclonal antibodies, seen in differentiated but not undifferentiated sebocytes (Zhang 2006). MC5R mRNA was also detected in sebaceous glands from the skin
35 of wild-type mice, but not in skin sections of the MC5R-knockout mice (Chen 1997). Treatment of human sebocytes with cholera toxin (ChT), bovine pituitary extract (BPE), α -MSH or NDP-MSH increases lipid droplet formation, squalene synthesis, and MC5R

expression (Zhang 2003; Zhang 2006), While both MC1R and MC5R have been detected in sebaceous cells, treatment of primary human sebocyte cell culture with NDP-MSH or BPE caused a substantial increase in human MC5R expression compared to serum-free conditions, correlating with sebocyte differentiation. Immortalized sebaceous cell lines (SZ-95, TSS-1 and SEB-1) also show MC5R expression (Jeong 2007; Smith 2007a; Phan 2007). These studies suggest that MC5R antagonists could be useful in reducing sebum secretion in mammals and hence in treating conditions associated with excess sebum secretion.

A family of 1,2,4-thiadiazole derivatives with MC5R antagonist activity (138-320 nM) were found to reduce sebum formation both in human sebocyte cell cultures and when applied topically to human skin grafted onto immunodeficient mice (Eisinger 2003a-d; 2006a,b).

Excessive sebum secretion, or seborrhoea, is a common affliction. Sebaceous glands occur over most of the body, with dense concentrations of large glands on the face, scalp and upper trunk (Simpson and Cunliffe p43.1). Sebaceous secretion is dependent in part on androgenic hormones, possibly partly mediated by 5 α -reductase processing of testosterone to 5 α -DHT (dihydrotestosterone). Sebum consists of a species-specific mixture of lipids. In humans this consists of approximately 58% glycerides, 26% wax esters, 12% squalene, and 4% cholesterol/cholesterol esters (Simpson and Cunliffe p43.5). The presence of squalene is almost exclusively characteristic of human sebum. The function of sebum is not well defined, but it is believed to have fungistatic properties, and play a role in moisture loss from, and water repellence of, the epidermis (Simpson and Cunliffe p43.6; Danby 2005; Porter 2001; Shuster 1976; Kligman 1963).

Excessive sebum secretion has been associated with the development of acne vulgaris. Acne vulgaris is a common disease affecting an estimated 80% of the world's population at some stage in their lives. A person is more likely to develop acne than any other disease, although the severity varies greatly (Simpson and Cunliffe p43.16). Acne peaks in prevalence and severity in adolescents aged 14-19 years old, with approximately 35-40% affected, but in a significant number of patients (7-24%) it persists beyond 25 years of age (Simpson and Cunliffe p43.15). Of patients treated for acne, one study found 80% still had symptoms at 30-40 years of age (Simpson and Cunliffe p43.16). While acne is not a life-threatening disease it can have a severe impact on a patient's quality of life (Follador 2006), with one study of severe acne patients showing similar impact as much more serious chronic medical conditions such as asthma, epilepsy, diabetes, back pain or arthritis (Mallon 1999).

Four major factors are believed to be involved in the pathogenesis of acne: (i) increased sebum production (seborrhoea), (ii) hypercornification/blockage of the pilosebaceous duct (comedogenesis), (iii) infection of the duct with *P. acnes*, and (iv) inflammation of the pilosebaceous duct (Simpson and Cunliffe p43.15; Williams 2006). A number of studies have demonstrated a clear link between increased production of sebum, and the presence and severity of acne (Simpson and Cunliffe p43.17; Youn 2005; Piérard 1987; Harris 1983; Cotterill 1981; Thody 1975c; Pochi 1964). A 2007 study found a correlation between sebum excretion and development of acne in preadolescent children (Mourelatos 2007). Sebum is the main nutrient of *P. acnes*, thus reduction of sebum will reduce the subsequent bacterial infection and inflammation response.

Androgenic sex hormones appear to play a role in the development of acne, with strong correlations with sebum production (Makrantonaki 2007). Two oral contraceptive pills are approved by the FDA for the treatment of acne vulgaris (Harper 2005), and these compounds appear to act by reducing androgen mediated sebum formation. Diet (Cordain 2005; Smith 2007b), stress (Zouboulis 2004) and genetic factors (Goulden 1999; Bataille 2006) also may play a role in acne, again potentially via increased sebum production.

Current treatments for acne vulgaris focus predominantly on treating the infection and inflammation stages of the disease, with a large number of different formulations of topical antibiotics (e.g. benzoyl peroxide, tetracycline, erythromycin, clindamycin) and retinoids (e.g. retinoic acid, isotretinoin, adapalene, tazarotene) in use, either alone or in combination; some of these also possess anti-inflammatory action (Simpson and Cunliffe p43.36-43.38). Many of these treatments are of limited efficacy, particularly for severe cases of acne. A growing problem is the development of antibiotic-resistant strains of *P. acnes* (Simpson and Cunliffe p43.37, 43.46; Williams 2006). Both topical retinoids and benzoyl peroxide cause skin irritation, and retinoids can cause photosensitivity (Williams 2006). Oral therapies include isotretinoin, antibiotics, hormones, and steroids. In females, antiandrogens have been shown to reduce sebum production (by approximately 40-80%, though with no placebo control group) and improve acne (Simpson and Cunliffe p43.44; Burke 1984; Goodfellow 1984). Laser and UV-based therapies are gaining acceptance, and are believed to act through heating of the sebaceous gland followed by reduction in sebum formation; with reductions in both sebum formation and acne lesions measured (Jih 2006; Bhardwaj 2005). Of the many therapies available for acne, only oral isotretinoin and hormonal therapies act by regulating the sebaceous gland to reduce sebum secretion (Clarke 2007).

The most effective acne treatment, oral isotretinoin (13-cis-retinoic acid, Roaccutane, Accutane) was introduced in 1983 and still remains the most clinically effective anti-acne therapy. It is the only known treatment with strong sebusuppressive activity, reducing sebum excretion by up to 90% after 8-12 weeks of therapy (60-70% by 2 weeks) (Simpson and Cunliffe p43.47; Jones 1983; Goldstein 1982; King 1982). Topical retinoids, in contrast, have no effect on sebum production. Oral isotretinoin is also anti-inflammatory, reduces comedogenesis, and reduces *P. acnes* infection. The mechanism of action is still unclear, and metabolites of isotretinoin appear to play a significant role. Isotretinoin induces apoptosis and cell cycle arrest in human immortalized SEB-1 sebocyte cell culture (Nelson 2006). Unfortunately, oral isotretinoin has serious side effects; most significantly it is a teratogen and requires a registration program for use in the USA. The FDA has issued a warning against online purchases of isotretinoin. Blood testing for fasting lipids and liver function is also recommended during treatment (Williams 2006). Isotretinoin has been implicated (though not substantively) with adverse psychological effects, including suicide and depression (Marqueling 2005).

Other forms of acne, such as acne conglobata or acne fulminans, may also respond to a sebum-reducing agent. Seborrhoea, or excessive skin oil production, is often associated with severe acne. Seborrheic dermatitis (SD) is a skin disease associated with sebum-rich areas of the scalp, face and trunk with scaly, flaky, itchy red skin affecting 3-5% of the population; dandruff represents a mild form of this dermatitis affecting 15-20% of the population. Seborrhoea and SD appear more common in patients with Parkinson's disease or mood disorders (facial paralysis, supraorbital injury, poliomyelitis, syringomyelia, quadriplegia, unilateral injury to the ganglion Gasser and those with HIV/AIDS) (Plewig 1999). Studies have shown that seborrheic dermatitis is also associated with chronic alcoholic pancreatitis, hepatitis C virus and various cancers. It is also common in patients with genetic disorders, such as Down's syndrome, Hailey-Hailey disease and cardio-facio cutaneous syndrome (Gupta 2004). MC5R antagonists may be useful for treating these indications.

Although rare, a variety of tumours involving sebaceous glands or sebaceous cells have been described (e.g. Ide 1999; Mariappan 2004; Kruse 2003). Muir-Torre syndrome consists of sebaceous gland adenomas associated with an internal adenocarcinoma (usually colon, breast, ovary or prostate). Preventing sebaceous cell differentiation may provide an effective treatment for arresting tumour growth. Oral isotretinoin has been used for this purpose (Graefe 2000). Sebaceous hyperplasia is a benign hyperplasia of the sebaceous glands, generating yellowish small papules on the skin surface, usually the face. The disease is associated with excessive undifferentiated sebocyte proliferation, but not excessive sebum

formation. Ectopic sebaceous glands (Fordyce spots) are similar yellow papules found in the mouth or on the penile shaft. Both respond to oral isotretinoin. A compound that reduced sebocyte proliferation could be an effective treatment.


5 α -MSH shows immunosuppressive effects in humans, suppressing a variety of inflammation responses, and the MC5R has been implicated in these immunomodulating activities. MC5R mRNA was found to be expressed at high levels in human CD4+ T helper (Th) cells and in moderate levels in other human peripheral blood leukocytes (Andersen 2005). In mice, MC5R was detected in the lymphoid organs (Labbé, 1994), and MC5R was
10 found on the surface of mouse pro-B-lymphocyte cells where it appears to mediate α -MSH activation of the JAK2 signalling pathway, enhancing cellular proliferation (Buggy 1998). Induction of CD25+ CD4+ regulatory T-cells by α -MSH also appears to be through MC5R (Taylor 2001).

15 For the reasons described above it would be desirable to provide MC5R antagonists that could be used in a number of therapeutic areas. Therapeutic regulation of biological signal transduction includes modulation of MC5R-mediated cellular events including, *inter alia*, inhibition or potentiation of interactions among MC5R-binding and activating or deactivating molecules, or of other agents that regulate MC5R activities. An increased ability
20 to so regulate MC5R may facilitate the development of methods for modulating sebum secretion or other biological processes, and for treating conditions associated with such pathways such as acne as described above.

 Accordingly there is still the need to develop improved methods of modulating the
25 activity of MC5R which would facilitate the treatment of MC5R related conditions.

SUMMARY OF THE INVENTION

The present invention provides a method of down-regulating the activity of MC5R or a fragment thereof *in vitro* comprising exposing the MC5R or a fragment thereof to a compound of the formula (I):



Formula (I)

wherein

10 Y is a group of formula $-(CR^9R^{10})_n-$;

X is selected from the group consisting $-C(=O)-$, $-OC(=O)-$, $-NHC(=O)-$, $-(CR^{11}R^{12})_s$, and $-S(=O)_2-$;

15 R is a group of the formula $-Z-R^4$ wherein

Z is a group of formula $-(CR^{13}R^{14})_q-$;

20 R^4 is selected from the group consisting of H, C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_6 - C_{18} aryl, optionally substituted C_1 - C_{18} heteroaryl, $NR^{4a}R^{4b}$, $C(=O)R^{15}$, $C(=O)NR^{16}R^{17}$, $-C(=NR^{16})NR^{17}R^{18}$, SR^{20} , $SC(=O)R^{20}$, SO_2R^{20} , OR^{20} , $ONR^{16}R^{17}$, $OCR^{17}R^{18}R^{20}$, $OC(=O)R^{20}$, $OC(=O)OR^{20}$, $OC(=O)NR^{16}R^{17}$, and $ONR^{16}C(=NR^{17})NR^{18}R^{19}$;

25 R^{4a} is selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, optionally substituted C_1 - C_{18} heteroaryl, $C(=O)R^{15a}$, $C(=O)NR^{15a}R^{16a}$, $C(=O)OR^{15a}$, SO_2R^{15a} , $C(=O)H$, $-C(=NR^{15a})-$
 30 $NR^{16a}R^{17a}$, and OR^{15a} ,

R^{4b} is selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, optionally substituted C_1 - C_{18} heteroaryl,
 5 $C(=O)R^{15a}$, $C(=O)NR^{15a}R^{16a}$, $C(=O)OR^{15a}$, or

R^{4a} and R^{4b} when taken together with the nitrogen atom to which they are attached form an optionally substituted heterocyclic moiety, or

10 one of R^{4a} and R^{4b} when taken together with any R^{13} or R^{14} and the atoms to which they are attached forms an optionally substituted heterocyclic moiety;

R^{13} and R^{14} are each independently selected from the group consisting of H, halogen, OH, C_1 - C_{12} alkyl, C_6 - C_{18} aryl, C_1 - C_{12} hydroxyalkyl, C_1 - C_{12} haloalkyl, C_1 - C_{12} alkyloxy and C_1 -
 15 C_{12} haloalkyloxy, or

when taken together with the carbon to which they are attached R^{13} and R^{14} form an optionally substituted C_3 - C_{12} cycloalkyl, or an optionally substituted C_1 - C_{12} heterocycloalkyl group, or

20 one of R^{13} and R^{14} when taken together with one of R^{4a} , and R^{4b} and the atoms to which they are attached form an optionally substituted heterocyclic moiety, or

one of R^{13} and R^{14} when taken together with one of R^{15} , R^{16} , R^{17} , R^{18} , R^{19} or R^{20} and
 25 the atoms to which they are attached form an optionally substituted cyclic moiety;

each R^{15} , R^{15a} , R^{16} , R^{16a} , R^{17} , R^{17a} , R^{18} , R^{19} and R^{20} is independently selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12}
 30 heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl, or

any two of R^{15} , R^{15a} , R^{16} , R^{16a} , R^{17} , R^{17a} , R^{18} , R^{19} and R^{20} when taken together with the atoms to which they are attached form an optionally substituted cyclic group, or

35 one of R^{15} , R^{16} , R^{17} , R^{18} , R^{19} and R^{20} when taken together with one of R^{13} and R^{14} and the atoms to which they are attached form an optionally substituted cyclic moiety;

8a

q is an integer selected from the group consisting of 0, 1, 2, 3, 4, and 5;

R¹ is selected from the group consisting of H, optionally substituted C₁-C₁₂alkyl, optionally substituted C₂-C₁₂alkenyl, optionally substituted C₂-C₁₂alkynyl, optionally substituted C₁-C₁₂heteroalkyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂heterocycloalkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl;

R² and R³ are each independently selected from the group consisting of H, optionally substituted C₁-C₁₂alkyl, optionally substituted C₂-C₁₂alkenyl, optionally substituted C₂-C₁₂alkynyl, optionally substituted C₁-C₁₂heteroalkyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂heterocycloalkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl;

each R^{5a} and R^{5b} are independently selected from the group consisting of H, halogen, C₁-C₁₂alkyl, C₁-C₁₂hydroxyalkyl and C₁-C₁₂haloalkyl, or

one or more of R^{5a} and R^{5b} when taken together with one or more of R⁶, R⁷ and R⁸ and the atoms to which they are attached form a moiety selected from the group consisting of an optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂ heterocycloalkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl;

R⁶, R⁷ and R⁸ are each independently selected from the group consisting of H, halogen, hydroxy, optionally substituted C₁-C₁₂alkyl, optionally substituted C₂-C₁₂alkenyl, optionally substituted C₂-C₁₂alkynyl, optionally substituted C₁-C₁₂ heteroalkyl, optionally substituted C₁-C₁₀ heteroalkenyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂ heterocycloalkyl, optionally substituted C₆-C₁₈aryl, optionally substituted C₁-C₁₈heteroaryl, optionally substituted amino, optionally substituted carboxy, C₁-C₁₂alkoxy, and optionally substituted thio, or

(a) when taken together with the carbon atom to which they are attached two or more of R⁶, R⁷ and R⁸ form a moiety selected from the group consisting of optionally substituted C₂-C₁₂alkenyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂heterocycloalkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl, or

8b

(b) one or more of R^6 , R^7 and R^8 when taken together with one or more of R^{5a} and R^{5b} and the atoms to which they are attached form a moiety selected from the group consisting of an optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl;

each R^9 and R^{10} is independently selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl;

each R^{11} and R^{12} is independently selected from the group consisting of H, and optionally substituted C_1 - C_{12} alkyl;


n is an integer selected from the group consisting of 1, 2, 3 and 4;

r is an integer selected from the group consisting of 1, 2, 3, and 4;

s is an integer selected from the group consisting of 1, 2, 3, and 4;

or a pharmaceutically acceptable salt or prodrug thereof.

The invention also provides use of a compound of the formula (I):



Formula (I)

wherein

Y is a group of formula $-(CR^9R^{10})_n-$;

8c

X is selected from the group consisting $-C(=O)-$, $-OC(=O)-$, $-NHC(=O)-$, $-(CR^{11}R^{12})_s$, and $-S(=O)_2-$;

5 R is a group of the formula $-Z-R^4$ wherein

Z is a group of formula $-(CR^{13}R^{14})_q-$;

10 R^4 is selected from the group consisting of H, C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_6 - C_{18} aryl, optionally substituted C_1 - C_{18} heteroaryl, $NR^{4a}R^{4b}$, $C(=O)R^{15}$, $C(=O)NR^{16}R^{17}$, $-C(=NR^{16})NR^{17}R^{18}$, SR^{20} , $SC(=O)R^{20}$, SO_2R^{20} , OR^{20} , $ONR^{16}R^{17}$, $OCR^{17}R^{18}R^{20}$, $OC(=O)R^{20}$, $OC(=O)OR^{20}$, $OC(=O)NR^{16}R^{17}$, and $ONR^{16}C(=NR^{17})NR^{18}R^{19}$;

15 R^{4a} is selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, optionally substituted C_1 - C_{18} heteroaryl, $C(=O)R^{15a}$, $C(=O)NR^{15a}R^{16a}$, $C(=O)OR^{15a}$, SO_2R^{15a} , $C(=O)H$, $-C(=NR^{15a})-$
 20 $NR^{16a}R^{17a}$, and OR^{15a} ,

R^{4b} is selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, optionally substituted C_1 - C_{18} heteroaryl, $C(=O)R^{15a}$, $C(=O)NR^{15a}R^{16a}$, $C(=O)OR^{15a}$, or
 25 $C(=O)R^{15a}$, $C(=O)NR^{15a}R^{16a}$, $C(=O)OR^{15a}$, or

R^{4a} and R^{4b} when taken together with the nitrogen atom to which they are attached form an optionally substituted heterocyclic moiety, or

30

one of R^{4a} and R^{4b} when taken together with any R^{13} or R^{14} and the atoms to which they are attached forms an optionally substituted heterocyclic moiety;

R^{13} and R^{14} are each independently selected from the group consisting of H, halogen,
 35 OH, C_1 - C_{12} alkyl, C_6 - C_{18} aryl, C_1 - C_{12} hydroxyalkyl, C_1 - C_{12} haloalkyl, C_1 - C_{12} alkyloxy and C_1 - C_{12} haloalkyloxy, or

8d

when taken together with the carbon to which they are attached R^{13} and R^{14} form an optionally substituted C_3 - C_{12} cycloalkyl, or an optionally substituted C_1 - C_{12} heterocycloalkyl group, or

one of R^{13} and R^{14} when taken together with one of R^{4a} , and R^{4b} and the atoms to which they are attached form an optionally substituted heterocyclic moiety, or

one of R^{13} and R^{14} when taken together with one of R^{15} , R^{16} , R^{17} , R^{18} , R^{19} or R^{20} and the atoms to which they are attached form an optionally substituted cyclic moiety;

each R^{15} , R^{15a} , R^{16} , R^{16a} , R^{17} , R^{17a} , R^{18} , R^{19} and R^{20} is independently selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl, or

any two of R^{15} , R^{15a} , R^{16} , R^{16a} , R^{17} , R^{17a} , R^{18} , R^{19} and R^{20} when taken together with the atoms to which they are attached form an optionally substituted cyclic group, or

one of R^{15} , R^{16} , R^{17} , R^{18} , R^{19} and R^{20} when taken together with one of R^{13} and R^{14} and the atoms to which they are attached form an optionally substituted cyclic moiety;

q is an integer selected from the group consisting of 0, 1, 2, 3, 4, and 5;

R^1 is selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl;

R^2 and R^3 are each independently selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl;

each R^{5a} and R^{5b} are independently selected from the group consisting of H, halogen, C_1 - C_{12} alkyl, C_1 - C_{12} hydroxyalkyl and C_1 - C_{12} haloalkyl, or

5 one or more of R^{5a} and R^{5b} when taken together with one or more of R^6 , R^7 and R^8 and the atoms to which they are attached form a moiety selected from the group consisting of an optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl;

10 R^6 , R^7 and R^8 are each independently selected from the group consisting of H, halogen, hydroxy, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_1 - C_{10} heteroalkenyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, optionally substituted
15 C_1 - C_{18} heteroaryl, optionally substituted amino, optionally substituted carboxy, C_1 - C_{12} alkyloxy, and optionally substituted thio, or

(a) when taken together with the carbon atom to which they are attached two or more of R^6 , R^7 and R^8 form a moiety selected from the group consisting of optionally
20 substituted C_2 - C_{12} alkenyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl, or

(b) one or more of R^6 , R^7 and R^8 when taken together with one or more of R^{5a}
25 and R^{5b} and the atoms to which they are attached form a moiety selected from the group consisting of an optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl;

30 each R^9 and R^{10} is independently selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl;

each R^{11} and R^{12} is independently selected from the group consisting of H, and
35 optionally substituted C_1 - C_{12} alkyl;

n is an integer selected from the group consisting of 1, 2, 3 and 4;

r is an integer selected from the group consisting of 1, 2, 3, and 4;

s is an integer selected from the group consisting of 1, 2, 3, and 4;

5

or a pharmaceutically acceptable salt or prodrug thereof, in down-regulating the activity of MC5R or a fragment thereof.

In one embodiment the MC5R or fragment or analogue or functional equivalent thereof is in a cell and the method comprises exposing the cell to a compound of formula (I).
5 In one embodiment the invention provides a method of down-regulating the activity of MC5R or fragment or analogue or functional equivalent thereof in a mammal comprising administering a MC5R-modulating amount of a compound of the invention to the mammal.

10 In yet a further aspect the invention provides the use of a compound of the formula (I) in down-regulating the activity of MC5R or a fragment, analogue or functional equivalent thereof.

In yet a further aspect the invention provides the use of a compound of formula (I) in
15 the preparation of a medicament for down-regulating the activity of MC5R or fragment or analogue or functional equivalent thereof in a mammal.

In yet an even further aspect the invention provides a method of treating, preventing, or controlling a condition associated with the activity of MC5R or a fragment, analogue or
20 functional equivalent thereof in a mammal the method comprising administering a therapeutically effective amount of a compound of formula (I). The compound may be administered in any way known in the art although in one aspect the compound is administered topically. In another aspect the compound is administered orally. In another aspect the compound is administered parenterally. In one embodiment of the method the
25 condition is selected from the group consisting of acne, seborrhoea, and seborrheic dermatitis. In one embodiment the condition is acne vulgaris. In one embodiment the compound of formula (I) is administered in combination with a second active agent. In one embodiment the second active agent is selected from the group consisting of antibiotics, retinoids, anti-androgens, and steroids.

30

In yet an even further aspect the invention provides the use of a compound of formula (I) in the preparation of a medicament for treating, preventing, or controlling a condition associated with the activity of MC5R or a fragment, analogue or functional equivalent thereof
35 in a mammal. In one aspect the medicament is adapted to be administered topically. In another aspect the medicament is adapted to be administered orally. In another aspect the compound is administered parenterally. In one embodiment of the method the condition is

selected from the group consisting of acne, seborrhoea, and seborrheic dermatitis. In a specific embodiment of the invention the condition is acne vulgaris.

In yet an even further aspect the invention provides a method of reducing sebum secretion by a mammal the method comprising administering a therapeutically effective amount of a compound of formula (I) to the mammal. The compound of the invention may be administered in any way known in the art although in one aspect the compound is administered topically. In another aspect the compound is administered orally. In another aspect the compound is administered parenterally.

In yet an even further aspect the invention provides the use of a compound of formula (I) in the preparation of a medicament for reducing sebum secretion in a mammal. In one embodiment the medicament is adapted to be administered topically. In another embodiment the medicament is adapted to be administered orally. In another embodiment the compound is administered parenterally.

DETAILED DESCRIPTION OF THE INVENTION

In this specification a number of terms are used which are well known to a skilled addressee. Nevertheless for the purposes of clarity a number of terms will be defined.

As used herein, the term "unsubstituted" means that there is no substituent or that the only substituents are hydrogen.

The term "optionally substituted" as used throughout the specification denotes that the group may or may not be further substituted or fused (so as to form a condensed polycyclic system), with one or more non-hydrogen substituent groups. In certain embodiments the substituent groups are one or more groups independently selected from the group consisting of halogen, =O, =S, -CN, -NO₂, -CF₃, -OCF₃, alkyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, heteroalkyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, aryl, heteroaryl, cycloalkylalkyl, heterocycloalkylalkyl, heteroarylalkyl, arylalkyl, cycloalkylalkenyl, heterocycloalkylalkenyl, arylalkenyl, heteroarylalkenyl, cycloalkylheteroalkyl, heterocycloalkylheteroalkyl, arylheteroalkyl, heteroarylheteroalkyl, hydroxy, hydroxyalkyl, alkyloxy, alkyloxyalkyl, alkyloxycycloalkyl, alkyloxyheterocycloalkyl, alkyloxyaryl, alkyloxyheteroaryl, alkyloxycarbonyl, alkylaminocarbonyl, alkenyloxy, alkynyloxy, cycloalkyloxy, cycloalkenyloxy, heterocycloalkyloxy, heterocycloalkenyloxy, aryloxy, phenoxy, benzyloxy, heteroaryloxy, arylalkyloxy, amino, alkylamino, acylamino, aminoalkyl, arylamino, sulfonylamino, sulfinylamino, sulfonyl, alkylsulfonyl, arylsulfonyl, aminosulfonyl, sulfinyl,

alkylsulfinyl, arylsulfinyl, aminosulfinylaminoalkyl, $-C(=O)OH$, $-C(=O)R^a$, $-C(=O)OR^a$, $C(=O)NR^aR^b$, $C(=NOH)R^a$, $C(=NR^a)NR^bR^c$, NR^aR^b , $NR^aC(=O)R^b$, $NR^aC(=O)OR^b$, $NR^aC(=O)NR^bR^c$, $NR^aC(=NR^b)NR^cR^d$, $NR^aSO_2R^b$, $-SR^a$, $SO_2NR^aR^b$, $-OR^a$, $OC(=O)NR^aR^b$, $OC(=O)R^a$ and acyl,

wherein R^a , R^b , R^c and R^d are each independently selected from the group consisting of H, C_1 - C_{12} alkyl, C_1 - C_{12} haloalkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} alkynyl, C_1 - C_{10} heteroalkyl, C_3 - C_{12} cycloalkyl, C_3 - C_{12} cycloalkenyl, C_1 - C_{12} heterocycloalkyl, C_1 - C_{12} heterocycloalkenyl, C_6 - C_{18} aryl, C_1 - C_{18} heteroaryl, and acyl, or any two or more of R^a , R^b , R^c and R^d , when taken together with the atoms to which they are attached form a heterocyclic ring system with 3 to 12 ring atoms.

In one embodiment each optional substituent is independently selected from the group consisting of: halogen, $=O$, $=S$, $-CN$, $-NO_2$, $-CF_3$, $-OCF_3$, alkyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, heteroalkyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, aryl, heteroaryl, hydroxy, hydroxyalkyl, alkyloxy, alkyloxyalkyl, alkyloxyaryl, alkyloxyheteroaryl, alkenyloxy, alkynyloxy, cycloalkyloxy, cycloalkenyloxy, heterocycloalkyloxy, heterocycloalkenyloxy, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, arylalkyloxy, amino, alkylamino, acylamino, aminoalkyl, arylamino, sulfonyl, alkylsulfonyl, arylsulfonyl, aminosulfonyl, aminoalkyl, $-COOH$, $-SH$, and acyl.

Examples of particularly suitable optional substituents include F, Cl, Br, I, CH_3 , CH_2CH_3 , OH, OCH_3 , CF_3 , OCF_3 , NO_2 , NH_2 , and CN.

The term "amino acid side chain group" represents a natural or unnatural side chain group present in a protein. The term includes side chain moieties present in naturally occurring proteins including the naturally occurring amino acid side chain moieties identified in table 1 below.

Table 1. Amino Acid Side Chain Moieties

Amino Acid Side Chain Moiety	Amino Acid
H	Glycine
CH_3	Alanine
$CH(CH_3)_2$	Valine
$CH_2CH(CH_3)_2$	Leucine
$CH(CH_3)CH_2CH_3$	Isoleucine
$(CH_2)_4NH_3^+$	Lysine

$(\text{CH}_2)_3\text{NHC}(\text{NH}_2)\text{NH}_2^+$	Arginine
$\text{CH}_2\text{-(imidazol-4-yl)}$	Histidine
CH_2COO^-	Aspartic Acid
$\text{CH}_2\text{CH}_2\text{COO}^-$	Glutamic acid
CH_2CONH_2	Asparagine
$\text{CH}_2\text{CH}_2\text{CONH}_2$	Glutamine
CH_2Ph	Phenylalanine
$\text{CH}_2\text{C}_6\text{H}_4\text{OH}$	Tyrosine
$\text{CH}_2\text{(Indolin-3-yl)}$	Tryptophan
CH_2SH	Cysteine
$\text{CH}_2\text{CH}_2\text{SCH}_3$	Methionine
CH_2OH	Serine
$\text{CH}(\text{OH})\text{CH}_3$	Threonine

In addition to naturally occurring amino acid side chain groups as identified above the term also includes derivatives or analogs thereof. As used herein the term derivative or analogue of an amino acid side chain group includes modifications and variations to naturally occurring side chain groups. With reference to the table above most of the naturally occurring amino acid side chain groups may be classified as alkyl, aryl, arylalkyl or heteroalkyl moieties. As such derivatives of amino acid side chain groups include straight or branched, cyclic or non-cyclic alkyl, aryl, heteroaryl, heteroarylalkyl, arylalkyl or heteroalkyl moieties.

Amino acid side chain groups as discussed above also include optionally substituted derivatives of alkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, or heteroalkyl moieties. The optional substituents may be selected from the group defined above. For example, the optional substituents may be selected from but are not limited to OH, Cl, Br, F, COOH, COOR^Z, CONH₂, NH₂, NHR^Z, NR^ZR^Z, SH, SR^Z, SO₂R^Z, SO₂H and SOR^Z wherein R^Z is an alkyl, aryl or arylalkyl moiety.

In the definitions of a number of substituents below it is stated that “the group may be a terminal group or a bridging group”. This is intended to signify that the use of the term is intended to encompass the situation where the group is a linker between two other portions of the molecule as well as where it is a terminal moiety. Using the term alkyl as an example, some publications would use the term “alkylene” for a bridging group and hence in these other publications there is a distinction between the terms “alkyl” (terminal group) and “alkylene” (bridging group). In the present application no such distinction is made and most groups may be either a bridging group or a terminal group.

Several terms are prefaced by the modifier indicating the number of carbon atoms present in the moiety. For example, the modifier "C₁-C₆" in front of the term "alkyl" indicates that the alkyl moiety has from 1 to 6 carbon atoms. Further, the modifier "C₁-C₁₈" in front of the term "heteroaryl" indicates that the heteroaromatic ring may have from 1 to 18 carbon atoms as part of the total number of atoms in the ring system.

"Acyl" means an R-C(=O)- group in which the R group may be an alkyl, cycloalkyl, heterocycloalkyl, aryl or heteroaryl group as defined herein. Examples of acyl include acetyl and benzoyl. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the carbonyl carbon.

"Acylamino" means an R-C(=O)-NH- group in which the R group may be an alkyl, cycloalkyl, heterocycloalkyl, aryl or heteroaryl group as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the nitrogen atom.

"Alkenyl" as a group or part of a group denotes an aliphatic hydrocarbon group containing at least one carbon-carbon double bond and which may be straight or branched preferably having 2-14 carbon atoms, more preferably 2-12 carbon atoms, most preferably 2-6 carbon atoms, in the normal chain. The group may contain a plurality of double bonds in the normal chain and the orientation about each is independently E or Z. Exemplary alkenyl groups include, but are not limited to, ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl and nonenyl. The group may be a terminal group or a bridging group.

"Alkenyloxy" refers to an alkenyl-O- group in which alkenyl is as defined herein. Preferred alkenyloxy groups are C₁-C₆ alkenyloxy groups. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the oxygen atom.

"Alkyl" as a group or part of a group refers to a straight or branched aliphatic hydrocarbon group, preferably a C₁-C₁₄ alkyl, more preferably a C₁-C₁₀ alkyl, most preferably C₁-C₆ unless otherwise noted. Examples of suitable straight and branched C₁-C₆ alkyl substituents include methyl, ethyl, n-propyl, 2-propyl, n-butyl, sec-butyl, t-butyl, hexyl, and the like. The group may be a terminal group or a bridging group.

"Alkylamino" includes both mono-alkylamino and dialkylamino, unless specified. "Mono-alkylamino" means a Alkyl-NH- group, in which alkyl is as defined herein. "Dialkylamino" means a (alkyl)₂N- group, in which each alkyl may be the same or different and are each as defined herein for alkyl. The alkyl group is preferably a C₁-C₆ alkyl group.
5 The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the nitrogen atom.

"Alkylaminocarbonyl" refers to a group of the formula (Alkyl)_x(H)_yNC(=O)- in which x is 1 or 2, and the sum of x+y =2. The group may be a terminal group or a bridging group. If the
10 group is a terminal group it is bonded to the remainder of the molecule through the carbonyl carbon.

"Alkyloxy" refers to an alkyl-O- group in which alkyl is as defined herein. Preferably the alkyloxy is a C₁-C₆alkyloxy. Examples include, but are not limited to, methoxy and ethoxy.
15 The group may be a terminal group or a bridging group.

"Alkyloxyalkyl" refers to an alkyloxy-alkyl- group in which the alkyloxy and alkyl moieties are as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the alkyl
20 group.

"Alkyloxyaryl" refers to an alkyloxy-aryl- group in which the alkyloxy and aryl moieties are as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the aryl group.
25

"Alkyloxycarbonyl" refers to an alkyl-O-C(=O)- group in which alkyl is as defined herein. The alkyl group is preferably a C₁-C₆ alkyl group. Examples include, but are not limited to, methoxycarbonyl and ethoxycarbonyl. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule
30 through the carbonyl carbon.

"Alkyloxycycloalkyl" refers to an alkyloxy-cycloalkyl- group in which the alkyloxy and cycloalkyl moieties are as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through
35 the cycloalkyl group.

"Alkyloxyheteroaryl" refers to an alkyloxy-heteroaryl- group in which the alkyloxy and heteroaryl moieties are as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the heteroaryl group.

5

"Alkyloxyheterocycloalkyl" refers to an alkyloxy-heterocycloalkyl- group in which the alkyloxy and heterocycloalkyl moieties are as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the heterocycloalkyl group.

10

"Alkylsulfinyl" means an alkyl-S(=O)- group in which alkyl is as defined herein. The alkyl group is preferably a C₁-C₆ alkyl group. Exemplary alkylsulfinyl groups include, but not limited to, methylsulfinyl and ethylsulfinyl. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the sulfur atom.

15

"Alkylsulfonyl" refers to an alkyl-S(=O)₂- group in which alkyl is as defined above. The alkyl group is preferably a C₁-C₆ alkyl group. Examples include, but not limited to methylsulfonyl and ethylsulfonyl. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the sulfur atom.

20

"Alkynyl" as a group or part of a group means an aliphatic hydrocarbon group containing a carbon-carbon triple bond and which may be straight or branched preferably having from 2-14 carbon atoms, more preferably 2-12 carbon atoms, more preferably 2-6 carbon atoms in the normal chain. Exemplary structures include, but are not limited to, ethynyl and propynyl. The group may be a terminal group or a bridging group.

25

"Alkynyloxy" refers to an alkynyl-O- group in which alkynyl is as defined herein. Preferred alkynyloxy groups are C₁-C₆ alkynyloxy groups. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the oxygen atom.

30

"Aminoalkyl" means an NH₂-alkyl- group in which the alkyl group is as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the alkyl group.

35

"Aminosulfonyl" means an $\text{NH}_2\text{-S(=O)}_2\text{-}$ group. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the sulfur atom.

5 "Aryl" as a group or part of a group denotes (i) an optionally substituted monocyclic, or fused polycyclic, aromatic carbocycle (ring structure having ring atoms that are all carbon) preferably having from 5 to 12 atoms per ring. Examples of aryl groups include phenyl, naphthyl, and the like; (ii) an optionally substituted partially saturated bicyclic aromatic carbocyclic moiety in which a phenyl and a C_{5-7} cycloalkyl or C_{5-7} cycloalkenyl group are fused
10 together to form a cyclic structure, such as tetrahydronaphthyl, indenyl or indanyl. The group may be a terminal group or a bridging group. Typically an aryl group is a $\text{C}_6\text{-C}_{18}$ aryl group.

"Arylalkenyl" means an aryl-alkenyl- group in which the aryl and alkenyl are as defined herein. Exemplary arylalkenyl groups include phenylallyl. The group may be a terminal
15 group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the alkenyl group.

"Arylalkyl" means an aryl-alkyl- group in which the aryl and alkyl moieties are as defined herein. Preferred arylalkyl groups contain a C_{1-5} alkyl moiety. Exemplary arylalkyl
20 groups include benzyl, phenethyl, 1-naphthalenemethyl and 2-naphthalenemethyl. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the alkyl group.

"Arylalkyloxy" refers to an aryl-alkyl-O- group in which the alkyl and aryl are as defined
25 herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the oxygen atom.

"Arylamino" includes both mono-arylamino and di-arylamino unless specified. Mono-arylamino means a group of formula arylNH- , in which aryl is as defined herein.
30 di-arylamino means a group of formula $(\text{aryl})_2\text{N-}$ where each aryl may be the same or different and are each as defined herein for aryl. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the nitrogen atom.

35 "Arylheteroalkyl" means an aryl-heteroalkyl- group in which the aryl and heteroalkyl moieties are as defined herein. The group may be a terminal group or a bridging group. If

the group is a terminal group it is bonded to the remainder of the molecule through the heteroalkyl group.

"Aryloxy" refers to an aryl-O- group in which the aryl is as defined herein. Preferably the aryloxy is a C₆-C₁₈aryloxy, more preferably a C₆-C₁₀aryloxy. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the oxygen atom.

"Arylsulfonyl" means an aryl-S(=O)₂- group in which the aryl group is as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the sulfur atom.

A "bond" is a linkage between atoms in a compound or molecule. The bond may be a single bond, a double bond, or a triple bond.

15

"Cyclic group" refers to saturated, partially unsaturated or fully unsaturated monocyclic, bicyclic or polycyclic ring system. Examples of cyclic groups include cycloalkyl, cycloalkenyl and aryl.

20

"Cycloalkenyl" means a non-aromatic monocyclic or multicyclic ring system containing at least one carbon-carbon double bond and preferably having from 5-10 carbon atoms per ring. Exemplary monocyclic cycloalkenyl rings include cyclopentenyl, cyclohexenyl or cycloheptenyl. The cycloalkenyl group may be substituted by one or more substituent groups. The group may be a terminal group or a bridging group.

25

"Cycloalkyl" refers to a saturated monocyclic or fused or spiro polycyclic, carbocycle preferably containing from 3 to 9 carbons per ring, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like, unless otherwise specified. It includes monocyclic systems such as cyclopropyl and cyclohexyl, bicyclic systems such as decalin, and polycyclic systems such as adamantane. The group may be a terminal group or a bridging group.

30

"Cycloalkylalkyl" means a cycloalkyl-alkyl- group in which the cycloalkyl and alkyl moieties are as defined herein. Exemplary monocycloalkylalkyl groups include cyclopropylmethyl, cyclopentylmethyl, cyclohexylmethyl and cycloheptylmethyl. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the alkyl group.

35

"Cycloalkylalkenyl" means a cycloalkyl-alkenyl- group in which the cycloalkyl and alkenyl moieties are as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the alkenyl group.

"Cycloalkylheteroalkyl" means a cycloalkyl-heteroalkyl- group in which the cycloalkyl and heteroalkyl moieties are as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the heteroalkyl group.

"Cycloalkyloxy" refers to a cycloalkyl-O- group in which cycloalkyl is as defined herein. Preferably the cycloalkyloxy is a C₁-C₆cycloalkyloxy. Examples include, but are not limited to, cyclopropanoxy and cyclobutanoxy. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the oxygen atom.

"Cycloalkenyloxy" refers to a cycloalkenyl-O- group in which the cycloalkenyl is as defined herein. Preferably the cycloalkenyloxy is a C₁-C₆cycloalkenyloxy. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the oxygen atom.

"Haloalkyl" refers to an alkyl group as defined herein in which one or more of the hydrogen atoms has been replaced with a halogen atom selected from the group consisting of fluorine, chlorine, bromine and iodine. A haloalkyl group typically has the formula C_nH_(2n+1-m)X_m wherein each X is independently selected from the group consisting of F, Cl, Br and I. In groups of this type n is typically from 1 to 10, more preferably from 1 to 6, most preferably 1 to 3. m is typically 1 to 6, more preferably 1 to 3. Examples of haloalkyl include fluoromethyl, difluoromethyl and trifluoromethyl.

"Haloalkenyl" refers to an alkenyl group as defined herein in which one or more of the hydrogen atoms has been replaced with a halogen atom independently selected from the group consisting of F, Cl, Br and I.

"Haloalkynyl" refers to an alkynyl group as defined herein in which one or more of the hydrogen atoms has been replaced with a halogen atom independently selected from the group consisting of F, Cl, Br and I.

"Halogen" represents chlorine, fluorine, bromine or iodine.

"Heteroalkyl" refers to a straight- or branched-chain alkyl group preferably having from 2 to 14 carbons, more preferably 2 to 10 carbons in the chain, one or more of which has been replaced by a heteroatom selected from S, O, P and N. Exemplary heteroalkyls include alkyl ethers, secondary and tertiary alkyl amines, amides, alkyl sulfides, and the like. The group may be a terminal group or a bridging group.

"Heteroaryl" either alone or part of a group refers to groups containing an aromatic ring (preferably a 5 or 6 membered aromatic ring) having one or more heteroatoms as ring atoms in the aromatic ring with the remainder of the ring atoms being carbon atoms. Suitable heteroatoms include nitrogen, oxygen and sulphur. Examples of heteroaryl include thiophene, benzothiophene, benzofuran, benzimidazole, benzoxazole, benzothiazole, benzisothiazole, naphtho[2,3-b]thiophene, furan, isoindolizine, xantholene, phenoxatine, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, tetrazole, indole, isoindole, 1H-indazole, purine, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, cinnoline, carbazole, phenanthridine, acridine, phenazine, thiazole, isothiazole, phenothiazine, oxazole, isooxazole, furazane, phenoxazine, 2-, 3- or 4- pyridyl, 2-, 3-, 4-, 5-, or 8- quinolyl, 1-, 3-, 4-, or 5- isoquinolinyl 1-, 2-, or 3- indolyl, and 2-, or 3-thienyl. The group may be a terminal group or a bridging group.

"Heteroarylalkyl" means a heteroaryl-alkyl group in which the heteroaryl and alkyl moieties are as defined herein. Preferred heteroarylalkyl groups contain a lower alkyl moiety. Exemplary heteroarylalkyl groups include pyridylmethyl. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the alkyl group.

"Heteroarylalkenyl" means a heteroaryl-alkenyl- group in which the heteroaryl and alkenyl moieties are as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the alkenyl group.

"Heteroarylheteroalkyl" means a heteroaryl-heteroalkyl- group in which the heteroaryl and heteroalkyl moieties are as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the heteroalkyl group.

"Heteroaryloxy" refers to a heteroaryl-O- group in which the heteroaryl is as defined herein. Preferably the heteroaryloxy is a C₁-C₁₂heteroaryloxy. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the oxygen atom.

5

"Heterocyclic" refers to saturated, partially unsaturated or fully unsaturated monocyclic, bicyclic or polycyclic ring system containing at least one heteroatom selected from the group consisting of nitrogen, sulfur and oxygen as a ring atom. Examples of heterocyclic moieties include heterocycloalkyl, heterocycloalkenyl and heteroaryl.

10

"Heterocycloalkenyl" refers to a heterocycloalkyl as defined herein but containing at least one double bond. The group may be a terminal group or a bridging group.

"Heterocycloalkyl" refers to a saturated monocyclic, bicyclic, or polycyclic ring containing at least one heteroatom selected from nitrogen, sulfur, oxygen, preferably from 1 to 3 heteroatoms in at least one ring. Each ring is preferably from 3 to 10 membered, more preferably 4 to 7 membered. Examples of suitable heterocycloalkyl substituents include pyrrolidyl, tetrahydrofuryl, tetrahydrothiofuranyl, piperidyl, piperazyl, tetrahydropyranyl, morpholino, 1,3-diazapane, 1,4-diazapane, 1,4-oxazepane, and 1,4-oxathiapane. The group may be a terminal group or a bridging group.

15

"Heterocycloalkylalkyl" refers to a heterocycloalkyl-alkyl- group in which the heterocycloalkyl and alkyl moieties are as defined herein. Exemplary heterocycloalkylalkyl groups include (2-tetrahydrofuryl)methyl, (2-tetrahydrothiofuranyl) methyl. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the alkyl group.

20

"Heterocycloalkylalkenyl" refers to a heterocycloalkyl-alkenyl- group in which the heterocycloalkyl and alkenyl moieties are as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the alkenyl group.

25

"Heterocycloalkylheteroalkyl" means a heterocycloalkyl-heteroalkyl- group in which the heterocycloalkyl and heteroalkyl moieties are as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the heteroalkyl group.

30

35

"Heterocycloalkyloxy" refers to a heterocycloalkyl-O- group in which the heterocycloalkyl is as defined herein. Preferably the heterocycloalkyloxy is a C₁-C₆heterocycloalkyloxy. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the oxygen atom.

"Heterocycloalkenyloxy" refers to a heterocycloalkenyl-O- group in which heterocycloalkenyl is as defined herein. Preferably the heterocycloalkenyloxy is a C₁-C₆ heterocycloalkenyloxy. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the oxygen atom.

"Hydroxyalkyl" refers to an alkyl group as defined herein in which one or more of the hydrogen atoms has been replaced with an OH group. A hydroxyalkyl group typically has the formula C_nH_(2n+1-x)(OH)_x. In groups of this type n is typically from 1 to 10, more preferably from 1 to 6, most preferably 1 to 3. x is typically 1 to 6, more preferably 1 to 3.

"Lower alkyl" as a group means unless otherwise specified, an aliphatic hydrocarbon group which may be straight or branched having 1 to 6 carbon atoms in the chain, more preferably 1 to 4 carbons such as methyl, ethyl, propyl (n-propyl or isopropyl) or butyl (n-butyl, isobutyl or tertiary-butyl). The group may be a terminal group or a bridging group.

"Sulfinyl" means an R-S(=O)- group in which the R group may be OH, alkyl, cycloalkyl, heterocycloalkyl; aryl or heteroaryl group as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the sulfur atom.

"Sulfinylamino" means an R-S(=O)-NH- group in which the R group may be OH, alkyl, cycloalkyl, heterocycloalkyl; aryl or heteroaryl group as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the nitrogen atom.

"Sulfonyl" means an R-S(=O)₂- group in which the R group may be OH, alkyl, cycloalkyl, heterocycloalkyl; aryl or heteroaryl group as defined herein. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the sulfur atom.

"Sulfonylamino" means an $R-S(=O)_2-NH-$ group. The group may be a terminal group or a bridging group. If the group is a terminal group it is bonded to the remainder of the molecule through the nitrogen atom.

5 It is understood that included in the family of compounds of Formula (I) are isomeric forms including diastereoisomers, enantiomers, tautomers, and geometrical isomers in "E" or "Z" configurational isomer or a mixture of E and Z isomers. It is also understood that some isomeric forms such as diastereomers, enantiomers, and geometrical isomers can be separated by physical and/or chemical methods and by those skilled in the art.

10 Some of the compounds of the disclosed embodiments may exist as single stereoisomers, racemates, and/or mixtures of enantiomers and /or diastereomers. All such single stereoisomers, racemates and mixtures thereof, are intended to be within the scope of the subject matter described and claimed.

15 The present invention includes all pharmaceutically acceptable isotopically-labeled compounds of formula (I) wherein one or more atoms have the same atomic number as, but an atomic mass or mass number different from, the atomic mass or mass number usually found in nature.

20 Examples of isotopes suitable for inclusion in the compounds of the invention include isotopes of hydrogen, such as ^2H and ^3H , carbon, such as ^{11}C , ^{13}C and ^{14}C , chlorine, such as ^{36}Cl , fluorine, such ^{18}F , iodine, such as ^{123}I and ^{125}I , nitrogen, such as ^{13}N and ^{15}N , oxygen, such as ^{15}O , ^{17}O and ^{18}O , phosphorus, such as ^{32}P , and sulphur, such as ^{35}S .

25 Certain isotopically-labeled compounds of formula (I), for example, those incorporating a radioactive isotope, are useful in drug and/or substrate tissue distribution studies. The radioactive isotopes tritium, *i.e.* ^3H , and carbon-14, *i.e.* ^{14}C , are particularly useful for this purpose in view of their ease of incorporation and ready means of detection.

30 Substitution with heavier isotopes such as deuterium, *i.e.* ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life or reduced dosage requirements, and hence may be preferred in some circumstances.

Substitution with positron emitting isotopes, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , can be useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy.

5 Isotopically-labeled compounds of formula (I) can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the accompanying Examples and Preparations using appropriate isotopically-labeled reagents in place of the non-labeled reagent previously employed.

10 Additionally, Formula (I) is intended to cover, where applicable, solvated as well as unsolvated forms of the compounds. Thus, each formula includes compounds having the indicated structure, including the hydrated as well as the non-hydrated forms.

The term "pharmaceutically acceptable salts" refers to salts that retain the desired
15 biological activity of the above-identified compounds, and include pharmaceutically acceptable acid addition salts and base addition salts. Suitable pharmaceutically acceptable acid addition salts of compounds of Formula (I) may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, sulfuric, and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic,
20 aromatic, heterocyclic carboxylic and sulfonic classes of organic acids, examples of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, fumaric, maleic, alkyl sulfonic, arylsulfonic. Additional information on pharmaceutically acceptable salts can be found in Remington's Pharmaceutical Sciences, 19th Edition, Mack Publishing Co., Easton, PA 1995. In the case of agents that are solids, it is understood by those skilled
25 in the art that the inventive compounds, agents and salts may exist in different crystalline or polymorphic forms, all of which are intended to be within the scope of the present invention and specified formulae.

"Prodrug" means a compound that undergoes conversion to a compound of formula
30 (I) within a biological system, usually by metabolic means (e.g. by hydrolysis, reduction or oxidation). For example an ester prodrug of a compound of formula (I) containing a hydroxyl group may be convertible by hydrolysis *in vivo* to the parent molecule. Suitable esters of compounds of formula (I) containing a hydroxyl group, are for example acetates, citrates, lactates, tartrates, malonates, oxalates, salicylates, propionates, succinates, fumarates,
35 maleates, methylene-bis- β -hydroxynaphthoates, gestisates, isethionates, di-*p*-toluoyltartrates, methanesulphonates, ethanesulphonates, benzenesulphonates, *p*-toluenesulphonates, cyclohexylsulphamates and quinate. As another example an ester prodrug of a compound


of formula (I) containing a carboxy group may be convertible by hydrolysis *in vivo* to the parent molecule. (Examples of ester prodrugs are those described by F.J. Leinweber, Drug Metab. Res., 18:379, 1987). Similarly, an acyl prodrug of a compound of formula (I) containing an amino group may be convertible by hydrolysis *in vivo* to the parent molecule
 5 (Many examples of prodrugs for these and other functional groups, including amines, are described in Prodrugs: Challenges and Rewards (Parts 1 and 2); Ed V. Stella, R. Borchardt, M. Hageman, R. Oliyai, H. Maag and J Tilley; Springer, 2007).

The term "therapeutically effective amount" or "effective amount" is an amount
 10 sufficient to effect beneficial or desired clinical results. An effective amount can be administered in one or more administrations. An effective amount is typically sufficient to palliate, ameliorate, stabilize, reverse, slow or delay the progression of the disease state.

The term "functional equivalent" is intended to include variants of the specific receptor
 15 described herein. It will be understood that receptors may have isoforms, such that while the primary, secondary, tertiary or quaternary structure of a given receptor isoform is different to the prototypical receptor; the molecule maintains biological activity as a receptor. Isoforms may arise from normal allelic variation within a population and include mutations such as amino acid substitution, deletion, addition, truncation, or duplication. Also included within the
 20 term "functional equivalent" are variants generated at the level of transcription.

In the methods and uses of the invention it is observed that certain of the compounds of the Formula (I), are more active than others and therefore it is desirable to use these compounds in the methods and uses of the present invention .


In the methods and uses of the invention the preferred stereochemistry at the 3 and 5 positions of the ring of the compounds of formula (I) is the 3S, 5S diastereomer. Accordingly, in one embodiment of the methods and uses of the invention the compound of formula (I) used is a compound of formula (Ia):



Formula (Ia)

wherein R, R¹, R², R³, R^{5a}, R^{5b}, R⁶, R⁷, R⁸, X, Y and r are as defined in formula 1, or a pharmaceutically acceptable salt or prodrug thereof.

5 In the methods and uses of the invention a particularly useful subset of compounds of formula (I) are compounds of formula (Ib) as shown below.



Formula (Ib)

10 wherein

R¹, R², R³, R^{5a}, R^{5b}, R⁶, R⁷, R⁸, X, Y and r are as defined above,

Z is a group of formula $-(CR^{13}R^{14})_q-$;

15

R⁴ is selected from the group consisting of H, C₁-C₁₂alkyl, optionally substituted C₂-C₁₂alkenyl, optionally substituted C₂-C₁₂alkynyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₆-C₁₈aryl, optionally substituted C₁-C₁₈heteroaryl, NR^{4a}R^{4b}, C(=O)R¹⁵, C(=O)NR¹⁶R¹⁷, -C(=NR¹⁶)NR¹⁷R¹⁸, SR²⁰, SC(=O)R²⁰, SO₂R²⁰, OR²⁰, ONR¹⁶R¹⁷, OCR¹⁷R¹⁸R²⁰, OC(=O)R²⁰, OC(=O)OR²⁰, OC(=O)NR¹⁶R¹⁷, and ONR¹⁶C(=NR¹⁷)NR¹⁸R¹⁹;

20

R^{4a} is selected from the group consisting of H, optionally substituted C₁-C₁₂alkyl, optionally substituted C₂-C₁₂alkenyl, optionally substituted C₂-C₁₂alkynyl, optionally substituted C₁-C₁₂heteroalkyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂heterocycloalkyl, optionally substituted C₆-C₁₈aryl, optionally substituted C₁-C₁₈heteroaryl, C(=O)R^{15a}, C(=O)NR^{15a}R^{16a}, C(=O)OR^{15a}, SO₂R^{15a}, C(=O)H, -C(=NR^{15a})-NR^{16a}R^{17a}, and OR^{15a},

25

R^{4b} is selected from the group consisting of H, optionally substituted C₁-C₁₂alkyl, optionally substituted C₂-C₁₂alkenyl, optionally substituted C₂-C₁₂alkynyl, optionally

30

substituted C₁-C₁₂heteroalkyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂heterocycloalkyl, optionally substituted C₆-C₁₈aryl, optionally substituted C₁-C₁₈heteroaryl, C(=O)R^{15a}, C(=O)NR^{15a}R^{16a}, C(=O)OR^{15a}, or

5 R^{4a} and R^{4b} when taken together with the nitrogen atom to which they are attached form an optionally substituted heterocyclic moiety, or

one of R^{4a} and R^{4b} when taken together with any R¹³ or R¹⁴ and the atoms to which they are attached forms an optionally substituted heterocyclic moiety;

10

R¹³ and R¹⁴ are each independently selected from the group consisting of H, halogen, OH, C₁-C₁₂alkyl, C₆-C₁₈aryl, C₁-C₁₂hydroxyalkyl, C₁-C₁₂haloalkyl, C₁-C₁₂alkyloxy and C₁-C₁₂haloalkyloxy, or

15

when taken together with the carbon to which they are attached R¹³ and R¹⁴ form an optionally substituted C₃-C₁₂cycloalkyl, or an optionally substituted C₁-C₁₂heterocycloalkyl group, or

20

one of R¹³ and R¹⁴ when taken together with one of R^{4a}, and R^{4b} and the atoms to which they are attached form an optionally substituted heterocyclic moiety, or

one of R¹³ and R¹⁴ when taken together with one of R¹⁵, R¹⁶, R¹⁷, R¹⁸, R¹⁹ or R²⁰ and the atoms to which they are attached form an optionally substituted cyclic moiety;

25

each R¹⁵, R^{15a}, R¹⁶, R^{16a}, R¹⁷, R^{17a}, R¹⁸, R¹⁹ and R²⁰ is independently selected from the group consisting of H, optionally substituted C₁-C₁₂alkyl, optionally substituted C₁-C₁₂heteroalkyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂heterocycloalkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl, or

30

any two of R¹⁵, R^{15a}, R¹⁶, R^{16a}, R¹⁷, R^{17a}, R¹⁸, R¹⁹ and R²⁰ when taken together with the atoms to which they are attached form an optionally substituted cyclic group, or

35

one of R¹⁵, R¹⁶, R¹⁷, R¹⁸, R¹⁹ and R²⁰ when taken together with one of R¹³ and R¹⁴ and the atoms to which they are attached form an optionally substituted cyclic moiety;

q is an integer selected from the group consisting of 0, 1, 2, 3, 4, and 5;

or a pharmaceutically acceptable salt or prodrug thereof.

In the methods and uses of the invention a particularly useful subset of compounds of
 5 formula (I) are compounds where Y is a group of the formula
 $-(CR^9R^{10})_n-$. In one embodiment of the suitable compounds n is 1 and Y is $-CR^9R^{10}-$. In
 another embodiment of the suitable compounds n is 2 and Y is $-CR^9R^{10}CR^9R^{10}-$.

In one embodiment of the compounds suitable for use in the invention each R^9 and
 10 R^{10} is independently selected from H and CH_3 . In one specific embodiment R^9 and R^{10} are
 both H. Accordingly in one embodiment of the compounds suitable for use in the invention Y
 is $-CH_2-$. In another embodiment of compounds suitable for use in the invention Y is $-$
 CH_2CH_2- . In yet an even further embodiment of compounds suitable for use in the invention
 Y is $-C(CH_3)_2-$.

15 In one embodiment of the compounds suitable for use in the invention R^2 is H or C_1-C_6
 alkyl. In a specific embodiment R^2 is H.

In one embodiment of the compounds suitable for use in the invention R^3 is H or C_1-C_6
 20 alkyl. In a specific embodiment R^3 is H.

In one embodiment of the compounds suitable for use in the invention X is selected
 from the group consisting of $-C(=O)-$ and $-(CR^{11}R^{12})_s-$. In one specific embodiment X is $-$
 $C(=O)-$. In one embodiment of the compounds suitable for use in the invention X is $-$
 25 $(CR^{11}R^{12})_s-$, s is 1. In another embodiment of compounds suitable for use in the invention X is
 $-(CR^{11}R^{12})_s-$, s is 2. In one form of each of these embodiments R^{11} and R^{12} are each
 independently selected from the group consisting of H and C_1-C_6 alkyl. In a specific
 embodiment both R^{11} and R^{12} are H, and s is 1 such that X is $-CH_2-$.


30 In one embodiment of the compounds suitable for use in the present invention $R^2 = H$,
 $R^3 = H$, $X = C(=O)$ and $Y = CH_2$. This provides compounds of formula (Ic).

Chemical structure of a substituted piperazine derivative. The piperazine ring has a carbonyl group at position 2 and a substituent Z at position 4. The nitrogen at position 1 is attached to a group R¹ via a carbonyl group. The nitrogen at position 3 is attached to a group R^{5a}. The carbon at position 5 is attached to a group R^{5b}. The carbon at position 6 is attached to a group R⁶ and a group R⁷. The carbon at position 7 is attached to a group R⁸.

In one embodiment of the compounds suitable for use in the invention R¹⁶ is selected from the group consisting of H, CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂, CH₂CH₂CH₂CH₃, CH(CH₃)CH₂CH₃, CH₂CH(CH₃)₂, C(CH₃)₃, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, benzyl, and phenyl, or a halogenated derivative thereof.

In one embodiment of the compounds suitable for use in the invention R^{17} is selected from the group consisting of H, CH_3 , CH_2CH_3 , $CH_2CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH_2CH_2CH_3$, $CH(CH_3)CH_2CH_3$, $CH_2CH(CH_3)_2$, $C(CH_3)_3$, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, benzyl, and phenyl, or a halogenated derivative thereof.

In one embodiment of the methods and uses of the invention the compound of formula (I) used is one in which $R^4 = NR^{4a}R^{4b}$. Accordingly a useful subset of compounds for use in the methods and uses of the present invention are compounds of formula (Id):




Formula (Id)

wherein R^1 , R^{4a} , R^{4b} , R^{5a} , R^{5b} , R^6 , R^7 , R^8 , Z and r are as defined for formula (I).

In one embodiment of the compounds suitable for use in the invention, r is selected from the group consisting of 1, 2, 3, and 4. In one specific embodiment r is 1. In another specific embodiment r is 2. In yet a further specific embodiment r is 3. In an even further specific embodiment r is 4.

In one embodiment of the compounds suitable for use in the invention, R^{5a} and R^{5b} are independently selected from H and C_1 - C_6 alkyl. In one embodiment R^{5a} and R^{5b} are each independently selected from H and CH_3 . In one specific embodiment R^{5a} and R^{5b} are both H. In yet another embodiment at least one of R^{5a} and R^{5b} when taken together with at least one of R^6 , R^7 and R^8 and the atoms to which they are attached form an optionally substituted cycloalkyl group. In one specific embodiment at least one of R^{5a} and R^{5b} when taken together with at least one of R^6 , R^7 and R^8 and the atoms to which they are attached forms a cyclohexyl group.


In one embodiment of the compounds of the invention, Y is CH₂, R² is H, R³ is H, R^{5a} and R^{5b} are H and X is -C(=O)-, and r is 1. This provides compounds of formula (II).



Formula (II)

wherein R¹, R⁴, R⁶, R⁷, R⁸ and Z are as defined for formula (I).

In one embodiment of the compounds of the invention, Y is CH₂, R² is H, R³ is H, R^{5a} and R^{5b} are H, X is -C(=O)-, R⁴ is NR^{4a}R^{4b}, and r is 1. This provides compounds of formula (IIa).



Formula (IIa)


wherein R¹, R^{4a}, R^{4b}, R⁶, R⁷, R⁸, and Z are as defined for formula (I).

In the compounds of the invention Z is a group of formula -(CR¹³R¹⁴)_q-. In one embodiment of the compounds suitable for use in the invention, and in particular the compounds of formula (I), formula (Ia), formula (Ib), formula (Ic), formula (Id), formula (II) and

Formula (IIa) R^{13} and R^{14} are independently selected from H and C_1-C_6 alkyl. In one embodiment R^{13} and R^{14} are each independently selected from H and CH_3 . In one specific embodiment R^{13} and R^{14} are both H. In yet another embodiment at least one of R^{13} and R^{14} when taken together with at least one of R^{4a} and R^{4b} and the atoms to which they are attached form an optionally substituted heterocycloalkyl group. In one embodiment Z is –
 5 $(CH_2)_q$ –


In one embodiment of the compounds suitable for use in the invention q is an integer selected from the group consisting of 0, 1, 2, 3, 4, and 5. In one specific embodiment q is 1.
 10 In another specific embodiment q is 2, in yet an even further specific embodiment q is 3, and in yet an even further specific embodiment q is 4.

Compounds in which R^7 , R^{13} and R^{14} are H and q is from 1 to 4 lead to compounds (IIIa), (IIIb), (IIIc) and (IIId) respectively.




Formula (IIIa)

wherein R^1 , R^{4a} , R^{4b} , R^6 , and R^8 are as defined for formula (I).




Formula (IIIb)

wherein R¹, R^{4a}, R^{4b}, R⁶, and R⁸ are as defined for formula (I).



Formula (IIIc)

wherein R¹, R^{4a}, R^{4b}, R⁶, and R⁸ are as defined for formula (I).



wherein R^1 , R^{4a} , R^{4b} , R^6 , and R^8 are as defined for formula (I).

5

In one form of the compounds suitable for use in the invention R^{4a} is selected from the group consisting of H, $-C(=N)NH_2$, $-C(=N)N(CH_3)_2$, $-C(=N)NCH(CH_3)_2$, $-C(=O)CH_3$, $-C(=O)cyclohexyl$, CH_3 , CH_2CH_3 , $CH_2CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH_2CH_2CH_3$, $CH(CH_3)CH_2CH_3$, $CH_2CH(CH_3)_2$, $C(CH_3)_3$, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, benzyl, and phenyl, or a halogenated derivative thereof. In one form of the compounds suitable for use in the invention R^{4b} is selected from the group consisting of H, CH_3 , CH_2CH_3 , $CH_2CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH_2CH_2CH_3$, $CH(CH_3)CH_2CH_3$, $CH_2CH(CH_3)_2$, $C(CH_3)_3$, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, benzyl, and phenyl, or a halogenated derivative thereof.

10

15










In another form of the compounds suitable for use in the invention R^{4a} and R^{4b} when taken together with the nitrogen atom to which they are attached form an optionally substituted C_2 - C_{12} heterocycloalkyl group, an optionally substituted C_2 - C_{12} heterocycloalkenyl group or an optionally substituted C_1 - C_{18} heteroaryl group.

20

















In a particular embodiment of the compounds suitable for use in the invention R^{4a} and R^{4b} when taken together with the nitrogen atom to which they are attached form an optionally substituted heterocycloalkyl group selected from the group consisting of piperidin-1-yl, pyrrolidin-1-yl, azetidin-1-yl, piperazin-1-yl, morpholin-4-yl and azepan-1-yl.

In one embodiment of the methods and uses of the invention, the compound of formula (I) is one in which one of R^{4a} and R^{4b} when taken together with the nitrogen atom to which it is attached and one of R^{13} and R^{14} and the carbon atom to which it is attached form an optionally substituted C_2 - C_{12} heterocycloalkyl group. In a particular embodiment one of R^{4a} and R^{4b} when taken together with the nitrogen atom to which it is attached and one of R^{13} and R^{14} and the carbon atom to which it is attached form an optionally substituted heterocycloalkyl group selected from the group consisting of piperidinyl, pyrrolidinyl, azetidiny, morpholinyl, piperazinyl, and azepanyl.

Specific examples of $NR^{4a}R^{4b}$ in the compounds which are useful in the methods and uses of the invention include:



35



5

10

In one embodiment of the compounds suitable for use in the invention, R⁷ is H.

In one embodiment of the compounds suitable for use in the invention, R⁶ and R⁸ are each independently selected from the group consisting of H, optionally substituted C₁-C₁₂ alkyl, optionally substituted C₂-C₁₂ alkenyl, optionally substituted C₆-C₁₈ aryl and optionally substituted C₁-C₁₈heteroaryl.

In one specific embodiment of the compounds suitable for use in the invention R⁶ is selected from the group consisting of H, methyl, trifluoromethyl, ethyl, 2,2,2-trifluoroethyl, isopropyl, isopropenyl, propyl, 2-ethyl-propyl, 3,3-dimethyl-propyl, butyl, 2-methylbutyl, isobutyl, 3,3-dimethyl-butyl, 2-ethyl-butyl, pentyl, 2-methyl-pentyl, optionally substituted phenyl and optionally substituted C₁-C₅ heteroaryl

In one specific embodiment of the compounds suitable for use in the invention R⁶ is optionally substituted phenyl or optionally substituted C₁-C₁₈heteroaryl.

In one specific embodiment of the compounds suitable for use in the invention R⁸ is selected from the group consisting of H, methyl, trifluoromethyl, ethyl, 2,2,2-trifluoroethyl, isopropyl, isopropenyl, propyl, 2-ethyl-propyl, 3,3-dimethyl-propyl, butyl, 2-methyl-butyl, isobutyl, 3,3-dimethyl-butyl, 2-ethyl-butyl, pentyl, 2-methyl-pentyl, pent-4-enyl, hexyl, heptyl, octyl, optionally substituted phenyl and optionally substituted C₁-C₅ heteroaryl.

In one specific embodiment of the compounds suitable for use in the invention R⁸ is methyl, ethyl, phenyl or optionally substituted C₁-C₅ heteroaryl.

In one specific embodiment of the compounds suitable for use in the invention R⁶, R⁷ and R⁸ when taken together with the carbon to which they are attached form a moiety selected from the group consisting of optionally substituted C₂-C₁₂alkenyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂heterocycloalkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl.

In one specific embodiment of the compounds suitable for use in the invention R⁶, R⁷ and R⁸ when taken together with the carbon to which they are attached form an optionally substituted C₆-C₁₈aryl group.

In one specific embodiment of the compounds of the invention and specifically of the compounds of formula (I), (II) (IIa) (IIIa), (IIIb), (IIIc), and (IIId), R⁶, R⁷ and R⁸ when taken together with the carbon atom to which they are attached form a disubstituted phenyl group.

In one embodiment the disubstituted phenyl group is a 2,4-disubstituted phen-1-yl group or a 3,5-disubstituted phen-1-yl group. A wide variety of substituents may be present on the disubstituted phenyl group as defined above. Examples of particularly suitable substituents include, but are not limited to, F, Br, Cl, methyl, trifluoromethyl, ethyl, 2,2,2-trifluoroethyl, isopropyl, propyl, 2-ethyl-propyl, 3,3-dimethyl-propyl, butyl, isobutyl, 3,3-dimethyl-butyl, 2-ethyl-butyl, pentyl, 2-methyl-pentyl, pent-4-enyl, hexyl, heptyl, octyl, phenyl, NH₂, cyano, phenoxy, hydroxy, methoxy, ethoxy, methylenedioxy, pyrrol-1-yl, and 3,5-dimethyl-pyrazol-1-yl. In one specific embodiment the disubstituted phenyl group is a 3,5-dichlorophen-1-yl group.

In one specific embodiment of the compounds suitable for use in the invention R¹ is selected from the group consisting of optionally substituted C₂-C₁₂alkenyl, optionally substituted C₆-C₁₈aryl and optionally substituted C₁-C₁₈heteroaryl.

In one specific embodiment of the compounds suitable for use in the invention R¹ is optionally substituted C₆-C₁₈aryl. The C₆-C₁₈aryl may be a monocyclic, bicyclic or polycyclic moiety. In certain embodiments the C₆-C₁₈aryl is a monocyclic moiety. In certain embodiments the C₆-C₁₈aryl is a bicyclic moiety.

In one specific embodiment R¹ is an optionally substituted C₆-C₁₈aryl selected from the group consisting of optionally substituted phenyl, biphenyl, and optionally substituted naphthyl. The moieties may be unsubstituted or may be substituted with one or more optional substituents. A wide variety of optional substituents may be used as defined above. Examples of particularly suitable optional substituents include, but are not limited to, F, Br, Cl, methyl, trifluoromethyl, ethyl, 2,2,2-trifluoroethyl, isopropyl, propyl, 2-ethyl-propyl, 3,3-dimethyl-propyl, butyl, isobutyl, 3,3-dimethyl-butyl, 2-ethyl-butyl, pentyl, 2-methyl-pentyl, pent-4-enyl, hexyl, heptyl, octyl, phenyl, NH₂, cyano, phenoxy, hydroxy, methoxy, ethoxy, pyrrol-1-yl, and 3,5-dimethyl-pyrazol-1-yl.

The substituents may be located at any substitutable position around the aryl ring available for substitution as would be clear to a skilled addressee. Examples of suitable optionally substituted phenyl compounds include, but are not limited to, 2-methoxy-phenyl, 3-methoxy-phenyl, 4-methoxy-phenyl, 2-trifluoromethyl-phenyl, 3-trifluoromethyl-phenyl, 4-trifluoromethyl-phenyl, 2-chloro-phenyl, 3-chloro-phenyl, 4-chloro-phenyl, 4-bromo-phenyl, 2-fluoro-phenyl, 3-fluoro-phenyl, 4-fluoro-phenyl, 4-hydroxy-phenyl, 4-phenyl-phenyl, 4-methyl-phenyl, 2,4-dichloro-phenyl, 3,4-dichloro-phenyl, 2,5-dichloro-phenyl, 2,6-difluoro-phenyl, 2-chloro-6-fluoro-phenyl, 3-fluoro-4-chloro-phenyl, 3-methyl-4-chloro-phenyl, 3-chloro-4-fluoro-

phenyl, 3-chloro-4-methyl-phenyl, 2-hydroxy-phenyl, 3-hydroxy-phenyl, 4-hydroxy-phenyl, 4-ethoxy-phenyl, 3-phenoxy-phenyl, 4-phenoxy-phenyl, 2-methyl-phenyl, 3-methyl-phenyl, 4-methyl-phenyl, 4-isopropyl-phenyl, 4-cyano-phenyl, 3,4-dimethyl-phenyl, 2,4-dimethyl-phenyl, 4-*t*-butyl-phenyl, 2,4-dimethoxy-phenyl, and 3,4-methylenedioxy-phenyl.

5

When R¹ is optionally substituted biphenyl the point of attachment of R¹ to the remainder of the molecule may be at the 2-, 3- or 4- position relative to the point of attachment of the second phenyl ring. As such the biphenyl may be an optionally substituted biphen-2-yl, or an optionally substituted biphen-3-yl, or an optionally substituted biphen-4-yl. In general the optionally substituted biphenyl is an optionally substituted biphen-4-yl. The optionally substituted biphenyl may be substituted in any suitable position.

10

When R¹ is optionally substituted naphthyl the point of attachment of R¹ to the remainder of the molecule may be at the 1 or 2 position. As such the naphthyl may be an optionally substituted naphth-1-yl, or an optionally substituted naphth-2-yl. In general the optionally substituted naphthyl is an optionally substituted naphth-2-yl. The optionally substituted naphthyl may be substituted in any suitable position. Examples of suitable optionally substituted naphth-2-yls include, but are not limited to, 6-fluoro-naphth-2-yl, 6-bromo-naphth-2-yl, 6-chloro-naphth-2-yl, 1-methoxy-naphth-2-yl, 3-methoxy-naphth-2-yl, 6-methoxy-naphth-2-yl, 1-hydroxy-naphth-2-yl, and 6-amino-naphth-2-yl.

15

20

In one specific embodiment of the compounds suitable for use in the invention R¹ is optionally substituted C₁-C₁₈heteroaryl. The C₁-C₁₈heteroaryl may be a monocyclic, bicyclic or polycyclic moiety. In certain embodiments the C₁-C₁₈heteroaryl is a monocyclic moiety. In certain embodiments the C₁-C₁₈heteroaryl is a bicyclic moiety. Examples of suitable heteroaryl moieties include, but are not limited to, indol-2-yl, indol-3-yl, quinolin-2-yl, quinolin-3-yl, isoquinolin-3-yl, quinoxaline-2-yl, benzo[b]furan-2-yl, benzo[b]thiophen-2-yl, benzo[b]thiophen-5-yl, thiazole-4-yl, benzimidazole-5-yl, benzotriazol-5-yl, furan-2-yl, benzo[d]thiazole-6-yl, pyrazole-1-yl, pyrazole-4-yl and thiophen-2-yl. These may also be optionally substituted as discussed above.


25

30

In one specific embodiment of the compounds suitable for use in the invention R¹ is an optionally substituted C₂-C₁₂alkenyl. The optionally substituted alkenyl may contain one or more double bonds with each of the double bonds being independently in the E or Z configuration. In one embodiment of the invention the alkenyl contains a single double bond which is in the E configuration.

35


In one specific form of this embodiment R^1 is an optionally substituted C_2 - C_{12} alkenyl of the formula:



R^{1a} is selected from the group consisting of H, halogen and optionally substituted C_1 - C_{12} alkyl;

R^{1b} and R^{1c} are each independently selected from the group consisting of H, halogen, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl.

In one form of this embodiment R^{1a} is H. In one form of this embodiment R^{1b} is H. This provides compounds where R^1 is of the formula:



In one embodiment of the compounds of the invention R^{1c} is optionally substituted C_6 - C_{18} aryl. The C_6 - C_{18} aryl may be monocyclic, bicyclic or polycyclic moiety. In certain embodiments the C_6 - C_{18} aryl is a monocyclic moiety. In certain embodiments the C_6 - C_{18} aryl is a bicyclic moiety.







In one specific embodiment R^{1c} is an optionally substituted C_6 - C_{18} aryl selected from the group consisting of optionally substituted phenyl and optionally substituted naphthyl. The moieties may be unsubstituted or may be substituted with one or more optional substituents. A wide variety of optional substituents may be used as defined above. Examples of particularly suitable optional substituents include, but are not limited to, F, Br, Cl, methyl,

trifluoromethyl, ethyl, 2,2,2-trifluoroethyl, isopropyl, propyl, 2-ethyl-propyl, 3,3-dimethyl-propyl, butyl, isobutyl, 3,3-dimethyl-butyl, 2-ethyl-butyl, pentyl, 2-methyl-pentyl, pent-4-enyl, hexyl, heptyl, octyl, phenyl, NH₂, cyano, phenoxy, hydroxy, methoxy, ethoxy, methylenedioxy, pyrrol-1-yl, and 3,5-dimethyl-pyrazol-1-yl.



















5

The substituents may be located at any substitutable position around the aryl ring available for substitution as would be clear to a skilled addressee. Examples of suitable optionally substituted phenyl compounds include, but are not limited to, 2-methoxy-phenyl, 3-methoxy-phenyl, 4-methoxy-phenyl, 2-trifluoromethyl-phenyl, 3-trifluoromethyl-phenyl, 4-trifluoromethyl-phenyl, 2-chloro-phenyl, 3-chloro-phenyl, 4-chloro-phenyl, 4-bromo-phenyl, 2-fluoro-phenyl, 3-fluoro-phenyl, 4-fluoro-phenyl, 4-hydroxy-phenyl, 4-phenyl-phenyl, 4-methyl-phenyl, 2,4-dichloro-phenyl, 3,4-dichloro-phenyl, 2,5-dichloro-phenyl, 2,6-difluoro-phenyl, 2-chloro-6-fluoro-phenyl, 3-fluoro-4-chloro-phenyl, 3-methyl-4-chloro-phenyl, 3-chloro-4-fluoro-phenyl, 3-chloro-4-methyl-phenyl, 2-hydroxy-phenyl, 3-hydroxy-phenyl, 4-hydroxy-phenyl, 4-ethoxy-phenyl, 3-phenoxy-phenyl, 4-phenoxy-phenyl, 2-methyl-phenyl, 3-methyl-phenyl, 4-methyl-phenyl, 4-isopropyl-phenyl, 4-cyano-phenyl, 3,4-dimethyl-phenyl, 2,4-dimethyl-phenyl, 4-t-butyl-phenyl, 2,4-dimethoxy-phenyl, and 3,4-methylenedioxy-phenyl.
















Specific compounds suitable for use in the methods and uses of the invention include the following:





















41


















42


















43


















44


















45


















46


















47


















48


















49





















50





















51


















52


















53





















54





















55





















56


















57





















58


















59


















60


















61


















62


















63


















64


















65



































67


















68
































70



71





or a pharmaceutically acceptable salt or prodrug thereof.

In order to assist the reader the names of compounds suitable for use in the invention as discussed above are as follows:

(14) N-(((3S,5S)-1-(3,5-dichlorobenzyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

(25) N-(((3S,5S)-3-(2-(diethylamino)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide

- (31) N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide
- (33) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 5 (37) N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-(2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (38) N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (39) N-(((3S,5S)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 (49) N-(((3S,5S)-3-(2-aminoethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (50) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)naphthalene-2-sulfonamide
- 15 (54) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (60) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-bromo-N-methyl-2-naphthamide
- (62) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-4-methyl-2-oxo-1,4-diazepan-5-yl)methyl)-6-bromo-2-naphthamide
- 20 (63) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (64) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 (65) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-(isopropylamino)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (67) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-(3-methylguanidino)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (71) (E)-N-(((3S,5S)-3-butyl-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- 30 (79) N-((S)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)acetamide
- (81) (S)-2-acetamido-N-((S)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)-3-(1H-imidazol-5-yl)propanamide
- 35 (83) propyl (S)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethylcarbamate

- (85) N-((R)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)acetamide
- (86) (S)-2-acetamido-N-((R)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)-3-(1H-imidazol-4-yl)propanamide
- 5 (87) propyl (R)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethylcarbamate
- (102) N-(((3R,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (103) N-(((3R,5R)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 (104) N-(((3S,5R)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (105) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 (106) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)biphenyl-4-carboxamide
- (107) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1H-indole-2-carboxamide
- (108) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)biphenyl-4-carboxamide
- 20 (109) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1H-indole-2-carboxamide
- (110) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(naphthalen-2-yl)acetamide
- 25 (111) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,2,3,4-tetrahydronaphthalene-2-carboxamide
- (112) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)quinoline-3-carboxamide
- (113) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)quinoxaline-2-carboxamide
- 30 (114) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)isoquinoline-3-carboxamide
- (115) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- 35 (116) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)quinoline-2-carboxamide

- (117) N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-1-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (118) N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-1-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(naphthalen-2-yl)acetamide
- 5 (119) N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-1-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1-naphthamide
- (120) N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-1-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(1H-indol-3-yl)acetamide
- (121) N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-2-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(biphenyl-4-yl)acetamide
- 10 (122) N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-2-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (123) N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-2-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(naphthalen-2-yl)acetamide
- 15 (124) N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-2-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1-naphthamide
- (125) N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-2-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(naphthalen-1-yl)acetamide
- (126) N-(((3S,5S)-3-(4-aminobutyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-
- 20 naphthamide
- (127) (S)-N-(((3S,5S)-3-(4-aminobutyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-3-carboxamide
- (128) (R)-N-(((3S,5S)-3-(4-aminobutyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-3-carboxamide
- 25 (129) N-(((3S,5S)-3-(4-aminobutyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzofuran-2-carboxamide
- (130) (R)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-(3-methylguanidino)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-3-carboxamide
- (131) (S)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-3-carboxamide
- 30 (132) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzofuran-2-carboxamide
- (133) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2,3-dihydro-1H-indene-2-carboxamide
- 35 (134) (R)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-3-carboxamide

- (135) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzo[b]thiophene-2-carboxamide
- (136) 2,4-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- 5 (137) 2,5-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- (138) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- (139) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)cyclohexanecarboxamide
- 10 (140) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-phenoxybenzamide
- (141) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-phenoxybenzamide
- 15 (142) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1H-indole-2-carboxamide
- (143) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-phenylpropanamide
- (144) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dimethylbenzamide
- 20 (145) 4-tert-butyl-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- (146) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2,4-dimethoxybenzamide
- 25 (147) 2-cyclohexyl-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acetamide
- (148) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzo[d][1,3]dioxole-5-carboxamide
- (149) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide
- 30 (150) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1H-benzo[d][1,2,3]triazole-5-carboxamide
- (151) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)cyclopentanecarboxamide
- 35 (152) 3,4-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide

- (153) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)cinnamamide
- (154) 3,5-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- 5 (155) 2-(2,4-dichlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acetamide
- (156) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1-methoxy-2-naphthamide
- (157) 2-(3,4-dichlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acetamide
- 10 (158) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-methoxy-2-naphthamide
- (159) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(2-(guanidinooxy)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 (160) (E)-3-(2,4-dichlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidino-propyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- (161) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-adamantane-1-carboxamide
- (162) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-phenoxyacetamide
- 20 (163) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-methoxy-2-naphthamide
- (164) 4-bromo-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- 25 (165) (S)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2,3-dihydrobenzo[b][1,4]dioxine-2-carboxamide
- (166) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- (167) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(thiophen-2-yl)acrylamide
- 30 (168) (R)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2,3-dihydrobenzo[b][1,4]dioxine-2-carboxamide
- (169) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-hydroxyphenyl)acrylamide
- 35 (170) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(2-methoxyphenyl)acrylamide

- (171) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-p-tolylacrylamide
- (172) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(2-(trifluoromethyl)phenyl)acrylamide
- 5 (173) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(3-fluorophenyl)acrylamide
- (174) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-methyl-3-phenylacrylamide
- (175) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-phenylcyclopropanecarboxamide
- 10 (176) 2-(2,4-dichlorophenoxy)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acetamide
- (177) (E)-3-(3-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- 15 (178) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzo[d]thiazole-6-carboxamide
- (179) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-5-phenylfuran-2-carboxamide
- (180) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(3-methoxyphenyl)acrylamide
- 20 (181) 6-bromo-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (182) N-(((3S,5S)-3-(3-guanidinopropyl)-2-oxo-1-phenethyl-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 (183) N-(((3S,5S)-1-(3,4-dichlorophenethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (184) N-(((3S,5S)-1-(2,4-dichlorophenethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (185) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzo[b]thiophene-5-carboxamide
- 30 (186) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-5-methyl-1-phenyl-1H-pyrazole-4-carboxamide
- (187) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-methoxyphenyl)acrylamide
- 35 (188) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide

- (189) (E)-3-(2-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- (190) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(2-hydroxyphenyl)acrylamide
- 5 (191) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-m-tolylacrylamide
- (192) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(3-(trifluoromethyl)phenyl)acrylamide
- (193) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(3-hydroxyphenyl)acrylamide
- 10 (194) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(2-fluorophenyl)acrylamide
- (195) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-o-tolylacrylamide
- 15 (196) (Z)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-fluoro-3-phenylacrylamide
- (197) N-((1-(2,2-diphenylethyl)-2-oxo-3-(piperidin-4-yl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (198) N-((1-(2,2-diphenylethyl)-2-oxo-3-(piperidin-4-ylmethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 20 (199) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)acrylamide
- (200) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-(trifluoromethyl)phenyl)acrylamide
- 25 (201) N-(((3S,5S)-1-(2,2-diphenylpropyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (202) N-(((3S,5S)-1-(cyclohexylmethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (203) N-(((3S,5S)-1-(1-adamantylmethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 30 (204) N-(((3S,5S)-1-((S)-1,1-diphenylpropan-2-yl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (205) N-(((3S,5S)-1-((R)-1,1-diphenylpropan-2-yl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 35 (206) N-(((3S,5S)-1-cyclohexyl-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

- (207) N-(((3S,5S)-1-((R)-1-fluoro-1,1-diphenylpropan-2-yl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (208) (E)-3-(2,6-difluorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- 5 (209) (E)-3-(2-chloro-6-fluorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- (210) (E)-3-(4-bromophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- (211) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-ethoxyphenyl)acrylamide
- 10 (212) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-bromo-2-naphthamide
- (213) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)cinnamamide
- 15 (214) (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- (215) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,4-dimethoxy-2-naphthamide
- (216) N-(((3S,5S)-3-(3-(3,3-dimethylguanidino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 20 (217) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-hydroxy-2-naphthamide
- (218) 6-amino-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 (219) (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-p-tolylacrylamide
- (220) (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)acrylamide
- (221) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide
- 30 (222) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-ethylhexanamide
- (223) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dimethylbenzamide
- 35 (224) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide

- (225) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-ethylhexanamide
- (226) N-(((3S,5S)-3-(3-(cyclohexylamino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 (227) N-(((3S,5S)-3-(3-guanidinopropyl)-1-(naphthalen-2-yl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (228) N-(((3S,5S)-1-((9H-fluoren-9-yl)methyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (229) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(pyridin-3-ylmethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 (230) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(pyridin-4-ylmethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (231) (E)-N-(((3S,5S)-3-(3-(cyclohexylamino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)acrylamide
- 15 (232) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(4-(isopropylamino)butyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (233) (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(2,4-difluorophenyl)acrylamide
- (234) (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-cyanophenyl)acrylamide
- 20 (235) (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(naphthalen-2-yl)acrylamide
- (236) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(4-fluorophenoxy)acetamide
- 25 (237) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-5-(4-chlorophenyl)furan-2-carboxamide
- (238) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-(1H-pyrrol-1-yl)benzamide
- (239) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-oxo-1-phenylpyrrolidine-3-carboxamide
- 30 (240) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-5-(4-chlorophenyl)isoxazole-3-carboxamide
- (241) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-5-(furan-2-yl)isoxazole-3-carboxamide
- 35 (242) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-phenylthiazole-4-carboxamide

- (243) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-(3,5-dimethyl-1H-pyrazol-1-yl)benzamide
- (244) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-5-methyl-1-phenyl-1H-pyrazole-4-carboxamide
- 5 (245) N-(((3S,5S)-1-(2-cyclohexylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (246) N-(((3S,5S)-1-(2-(bicyclo[2.2.1]heptan-2-yl)ethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (247) N-(((3S,5S)-1-(2,2-bis(4-methoxyphenyl)ethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 (248) N-(((3S,5S)-3-(3-(benzylamino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (249) N-(((3S,5S)-3-(3-(cyclopentylamino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 (250) N-(((3S,5S)-3-(3-(cyclobutylamino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (251) N-(((3S,5S)-3-(3-(dicyclobutylamino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (252) N-(((3S,5S)-1-benzyl-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 20 (253) N-(((3S,5S)-1-(2,2-bis(4-fluorophenyl)ethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (254) N-(((3S,5S)-3-(3-guanidinopropyl)-1-(naphthalen-2-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 (255) (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(5-methylthiophen-2-yl)acrylamide
- (256) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-phenyl-1H-pyrazole-5-carboxamide
- (257) (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)-N-methylacrylamide
- 30 (258) (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-4-methyl-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)acrylamide
- (259) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-(3-methyl-5-oxo-4,5-dihydro-1H-pyrazol-1-yl)benzamide
- 35 (260) (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-bromophenyl)acrylamide

- (261) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(pyrrolidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (262) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 5 (263) N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (264) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(pyrrolidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (265) N-(((3S,5S)-3-(3-(azetidin-1-yl)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 (266) N-(((3S,5S)-3-(3-guanidinopropyl)-1-(naphthalen-1-yl)methyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (267) N-(((3S,5S)-3-(3-guanidinopropyl)-1-(2-(naphthalen-2-yl)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 (268) N-(((3S,5S)-1-((S)-2-acetamido-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (269) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)cinnamamide
- (270) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-3,4-dimethylbenzamide
- 20 (271) 3,4-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)benzamide
- (272) N-(((3S,5S)-1-((S)-2-(cyclobutanecarboxamido)-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 (273) N-(((3S,5S)-1-((S)-2-(cyclohexanecarboxamido)-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (274) N-(((3S,5S)-3-(aminomethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (275) (E)-N-(((3S,5S)-3-(aminomethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- 30 (276) (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)acrylamide
- (277) (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-p-tolylacrylamide
- 35 (278) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(piperidin-1-yl)methyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

- (279) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(piperidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (280) N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- 5 (281) N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dimethylbenzamide
- (282) N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)cinnamamide
- (283) (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- 10 (284) 3,4-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- (285) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-3,4-dimethylbenzamide
- 15 (286) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)cinnamamide
- (287) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)acrylamide
- (288) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-3-p-tolylacrylamide
- 20 (289) N-(((3S,5S)-1-(3,5-dimethylbenzyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (290) N-(((3S,5S)-1-((S)-2-benzamido-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 (291) N-(((3S,5S)-1-((R)-2-benzamido-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (292) N-(((3S,5S)-3-(3-guanidinopropyl)-1-(2-methoxy-2-phenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (293) N-(((3S,5S)-3-(3-guanidinopropyl)-2-oxo-1-(2-phenyl-2-propoxyethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 30 (294) N-(((3S,5S)-1-(2-(benzyloxy)-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (295) N-(((3S,5S)-1-(2-(allyloxy)-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 35 (296) N-(((3S,5S)-3-butyl-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

- (297) (E)-N-(((3S,5S)-3-butyl-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- (298) (E)-N-(((3S,5S)-3-(3-amino-3-oxopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- 5 (299) N-(((3S,5S)-3-(3-amino-3-oxopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (300) (E)-N-(((3S,5S)-3-(3-acetamidopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-p-tolylacrylamide
- (301) N-(3-((2S,7S)-4-(2,2-diphenylethyl)-3-oxo-7-(((E)-3-p-tolylacrylamido)methyl)-1,4-diazepan-2-yl)propyl)cyclohexanecarboxamide
- 10 (302) N-(((3S,5S)-3-(3-guanidinopropyl)-2-oxo-1-(2-phenoxy-2-phenylethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (303) ethyl 3-(((3S,5S)-5-((2-naphthamido)methyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-1-yl)-2-phenylpropanoate
- 15 (304) N-(((3S,5S)-1-(2-ethylbutyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (305) N-(((3S,5S)-1-((3,5-dimethylcyclohexyl)methyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (306) N-(2-((2S,7S)-7-(((E)-3-(4-chlorophenyl)acrylamido)methyl)-4-(2,2-diphenylethyl)-3-oxo-1,4-diazepan-2-yl)ethyl)cyclohexanecarboxamide
- 20 (307) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-3-(2-(2-cyclohexylacetamido)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- (308) N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-aminoethyl)-1,4-diazepan-5-yl)ethyl)benzamide
- 25 (309) 3,4-dichloro-N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-aminoethyl)-1,4-diazepan-5-yl)ethyl)benzamide
- (310) N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-aminoethyl)-1,4-diazepan-5-yl)ethyl)-2-naphthamide
- (311) N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)ethyl)benzamide
- 30 (312) 3,4-dichloro-N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)ethyl)benzamide
- (313) N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)ethyl)-2-naphthamide
- 35 (314) N-(((3S,5S)-1-(3-(dimethylamino)-3-oxo-2-phenylpropyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

- (315) N-(((3S,5S)-3-(cyclohexylmethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (316) N-(((3S,5S)-3-(2-aminoethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 (317) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-oxo-3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (318) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-oxo-3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (319) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-phenethyl-1,4-diazepan-5-yl)methyl)-2-10 naphthamide
- (320) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-phenethyl-1,4-diazepan-5-yl)methyl)acrylamide
- (321) N-(((3S,5S)-3-(2-cyclohexylethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 (322) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-3-(2-cyclohexylethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- (323) N-(((3S,5S)-3-benzyl-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (324) (E)-N-(((3S,5S)-3-benzyl-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-20 chlorophenyl)acrylamide
- (325) N-(((3S,5S)-3-(2-aminoethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (326) (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- 25 (327) N-(((3S,5S)-1-(3-chloro-5-fluorobenzyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (328) N-(((3S,5S)-1-(3,5-difluorobenzyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (329) N-(((3S,5S)-3-(3-aminopropyl)-1-(3-chloro-5-fluorobenzyl)-2-oxo-1,4-diazepan-5-30 yl)methyl)-2-naphthamide
- (330) N-(((3S,5S)-3-(3-aminopropyl)-1-(3,5-difluorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (331) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 35 (332) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,6-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

- (333) N-(((3S,5S)-3-(3-aminopropyl)-1-(3,5-dimethoxybenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (334) N-(((3S,5S)-3-(3-aminopropyl)-1-(2-chlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 (335) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,3-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (336) N-(((3S,5S)-3-(3-aminopropyl)-1-(2,4-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (337) N-(((3S,5S)-3-(3-aminopropyl)-1-(3,4-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-
10 2-naphthamide
- (338) N-(((3S,5S)-3-(3-aminopropyl)-1-(3-fluoro-5-methylbenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (339) N-(((3S,5S)-3-(3-aminopropyl)-1-(3-fluoro-5-(trifluoromethyl)benzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 (340) N-(((3S,5S)-3-(3-aminopropyl)-1-(4-chlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (341) N-(((3S,5S)-3-(3-aminopropyl)-2-oxo-1-(2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (342) N-(((3S,5S)-3-(3-aminopropyl)-2-oxo-1-((1-phenylcyclohexyl)methyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
20
- (343) 3,4-dichloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- (344) N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 (345) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (346) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (347) (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-
30 (4-chlorophenyl)acrylamide
- (348) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (349) 3,4-dichloro-N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 35 (350) N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

- (351) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-chloro-3-fluorobenzamide
- (352) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-chloro-3-methylbenzamide
- 5 (353) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-chloro-4-fluorobenzamide
- (354) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-chloro-4-methylbenzamide
- (355) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(piperidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 10 (356) N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(pyrrolidin-1-yl)ethyl)-1,4-diazepan-5-yl)ethyl)-2-naphthamide
- (357) N-(((3S,5S)-3-(3-aminopropyl)-1-(3,5-bis(trifluoromethyl)benzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 (358) N-(((3S,5S)-3-(3-aminopropyl)-1-(3-chlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (359) N-(((3S,5S)-2-oxo-1-(2-phenylbutyl)-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (360) N-(((3S,5S)-3-(3-guanidinopropyl)-2-oxo-1-(3-oxo-2-phenyl-3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 20 (361) N-(((3S,5S)-3-(3-guanidinopropyl)-2-oxo-1-(3-oxo-2-phenyl-3-(phenylamino)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (362) 3,4-dichloro-N-(2-((3S,5R)-1-(2,2-diphenylethyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-diazepan-5-yl)ethyl)benzamide
- 25 (363) 3,4-dichloro-N-(2-((3S,5R)-3-(2-(diisopropylamino)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)ethyl)benzamide
- (364) N-(((3S,5S)-3-(aminomethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (365) N-(((3S,5S)-3-(aminomethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 30 (366) (E)-N-(((3S,5S)-3-(aminomethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- (367) 3,4-dichloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(piperidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 35 (368) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(piperidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)acrylamide

- (369) N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-(2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (370) (E)-N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-(2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- 5 (371) 3,4-dichloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(pyrrolidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)benzamide
- (372) N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(pyrrolidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (373) N-(((3S,5S)-3-(3-aminopropyl)-2-oxo-1-((S)-2-phenylpropyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 (374) N-(((3S,5S)-3-(3-aminopropyl)-2-oxo-1-((R)-2-phenylpropyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (375) N-(((3S,5S)-3-(2-(dimethylamino)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 (376) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide
- (377) N-(((3S,5S)-3-(3-aminopropyl)-1-(3,5-diethynylbenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (378) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-3-(2-(diethylamino)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- 20 (379) N-(((3S,5S)-3-(2-(diethylamino)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (380) 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 25 (381) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (382) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (383) N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- 30 (384) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(pyrrolidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (385) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 35 (386) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide

- (387) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- (388) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-bromo-2-naphthamide
- 5 (389) N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide
- (390) 6-chloro-N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (391) 6-bromo-N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 (392) N-(((3S,5S)-3-(2-aminoethyl)-1-((3,5-dimethylcyclohexyl)methyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (393) N-(((3S,5S)-3-(2-aminoethyl)-1-((3,5-dimethylcyclohexyl)methyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 (394) (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-((3,5-dimethylcyclohexyl)methyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- (395) 6-chloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (396) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide
- 20 (397) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(pyrrolidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (398) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- 25 (399) N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (400) 3,4-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- (401) N-(((3S,5S)-3-(3-aminopropyl)-1-((2,6-dimethylcyclohexyl)methyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 30 (402) N-(((3S,5S)-3-(3-aminopropyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (403) 3,4-dichloro-N-(((3S,5S)-1-((3,5-dimethylcyclohexyl)methyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 35 (404) 3,4-dichloro-N-(((3S,5S)-1-((3,5-dimethylcyclohexyl)methyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide

- (405) N-(((3S,5S)-3-(2-aminoethyl)-1-(3-methyl-2-phenylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (406) N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 (407) N-(((3S,5S)-3-(3-aminopropyl)-2-oxo-1-((R)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (408) (E)-N-(((3S,5S)-3-((1H-imidazol-4-yl)methyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- (409) 3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)propanamide
- 10 (410) N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)propanamide
- (411) N-(((2S,7S)-7-(((E)-3-(4-chlorophenyl)acrylamido)methyl)-4-(2,2-diphenylethyl)-3-oxo-1,4-diazepan-2-yl)methyl)picolinamide
- 15 (412) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(pyridin-2-ylmethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (413) N-(((3S,5S)-1-((R)-3-methyl-2-phenylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (414) N-(((3S,5S)-1-((S)-3-methyl-2-phenylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 20 (415) (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-3-(4-isopropylphenyl)acrylamide
- (416) (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-isopropylphenyl)acrylamide
- 25 (417) (E)-3-(2,4-dimethylphenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (418) (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(2,4-dimethylphenyl)acrylamide
- (419) (E)-3-(2,4-difluorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 30 (420) (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(2,4-difluorophenyl)acrylamide
- (421) N-(((2S,7S)-7-(((E)-3-(4-chlorophenyl)acrylamido)methyl)-4-(2,2-diphenylethyl)-3-oxo-1,4-diazepan-2-yl)methyl)cyclohexanecarboxamide
- 35 (422) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(2-morpholinoethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide

- (423) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-3-(2-(2,5-dimethyl-1H-pyrrol-1-yl)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- (424) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-3-(2-(2,5-dimethylpyrrolidin-1-yl)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- 5 (425) 6-chloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (426) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(pyrrolidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (427) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-3-(2-(isopropylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 10 (428) 6-chloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(pyrrolidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (429) 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 15 (430) 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(pyrrolidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- (431) benzyl ((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methylcarbamate
- (432) (E)-3-(4-bromophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 20 (433) 5-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)isoxazole-3-carboxamide
- (434) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-oxo-2-(pyridin-2-ylamino)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 25 (435) (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-oxo-2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (436) 6-chloro-N-(((3S,5S)-2-oxo-3-(3-oxo-3-(piperidin-1-yl)propyl)-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (437) 3,4-dichloro-N-(((3S,5S)-2-oxo-3-(3-oxo-3-(piperidin-1-yl)propyl)-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 30 (438) 6-chloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(piperidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (439) 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(piperidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 35 (440) 6-chloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

- (441) 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)benzamide
- (442) (E)-N-(2-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)propan-2-yl)-3-(4-chlorophenyl)acrylamide
- 5 (443) (E)-3-(4-chlorophenyl)-N-(2-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)propan-2-yl)acrylamide
- (444) N-(((3S,5S)-3-(2-aminoethyl)-1-((R)-2-(4-chlorophenyl)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (445) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-(4-chlorophenyl)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 (446) N-(((3S,5S)-1-((R)-2-(4-chlorophenyl)propyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (447) N-(((3S,5S)-1-((S)-2-(4-chlorophenyl)propyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 (448) 3,4-dichloro-N-(((3S,5S)-3-(2-(methyl(phenyl)amino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- (449) 3,4-dichloro-N-(((3S,5S)-3-(2-(diethylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- (450) 3,4-dichloro-N-(((3S,5S)-3-(2-morpholinoethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 20 (451) 3,4-dichloro-N-(((3S,5S)-2-oxo-3-(2-(phenylamino)ethyl)-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- (452) N-(((3S,5S)-3-(2-(benzylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- 25 (453) (5S,9aS)-5-(2-aminobenzyl)-2-((E)-3-(4-chlorophenyl)acryloyl)-7-(2,2-diphenylethyl)hexahydro-1H-imidazo[1,5-d][1,4]diazepin-6(5H)-one
- (454) N-(((3S,5S)-3-(2-(tert-butylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (455) 3,4-dichloro-N-(((3S,5S)-3-(2-(4-methylpiperazin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 30 (456) N-(((3S,5S)-2-oxo-1-((R)-2-phenylpentyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (457) N-(((3S,5S)-2-oxo-1-((S)-2-phenylpentyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 35 (458) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-4-(trifluoromethyl)benzamide

- (459) N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-(trifluoromethyl)benzamide
- (460) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-3-(trifluoromethyl)benzamide
- 5 (461) N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(trifluoromethyl)benzamide
- (462) 6-chloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (463) 3,4-dichloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-
- 10 diazepan-5-yl)methyl)benzamide
- (464) 6-chloro-N-(((3S,5S)-3-(2-(isopropylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (465) N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- 15 (466) N-(((3S,5S)-3-(2-aminobenzyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (467) N-(((3S,5S)-3-(2-(benzyl(methyl)amino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (468) 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperazin-1-yl)ethyl)-1,4-
- 20 diazepan-5-yl)methyl)benzamide
- (469) 3,4-dichloro-N-(((3S,5S)-3-(2-(methyl(pentyl)amino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- (470) 3,4-dichloro-N-(((3S,5S)-3-(2-(diisopropylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 25 (471) 3,4-dichloro-N-(((3S,5S)-3-(2-(4-methylpiperidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- (472) (S)-6-chloro-N-((2-oxo-1-(2-phenylbutyl)-3-(piperidin-4-yl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (473) (S)-6-chloro-N-(3-(1-isopentylpiperidin-4-yl)-2-oxo-1-(2-phenylbutyl)-1,4-diazepan-5-
- 30 yl)methyl)-2-naphthamide
- (474) N-(((3S,5S)-3-butyl-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- (475) 6-chloro-N-(((3S,5S)-3-isopentyl-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 35 (476) 3,4-dichloro-N-(((3S,5S)-3-(2-(3,5-dimethylpiperidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide

- (477) 3,4-dichloro-N-(((3S,5S)-3-(2-(4-hydroxypiperidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- (478) 1-(2-((2S,7S)-7-((3,4-dichlorobenzamido)methyl)-3-oxo-4-((S)-2-phenylbutyl)-1,4-diazepan-2-yl)ethyl)piperidine-4-carboxylic acid
- 5 (479) N-(((3S,5S)-3-(2-(azepan-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (480) 3,4-dichloro-N-(((3S,5S)-3-(2-((S)-2-methylpiperidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- (481) N-(((3S,5S)-3-(2-(tert-butyl(methyl)amino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-
- 10 diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (482) N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)benzyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (483) N-(((3S,5S)-3-(3-(butyl(methyl)amino)-3-oxopropyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-
- 15 diazepan-5-yl)methyl)-2-naphthamide
- (484) N-(((3S,5S)-3-(3-(cyclohexylamino)-3-oxopropyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-
- 20 diazepan-5-yl)methyl)-2-naphthamide
- (485) 6-chloro-N-((3-(1-ethylpiperidin-4-yl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (486) (3S,5S)-5-((3,4-dichlorobenzylamino)methyl)-1-(2,2-diphenylethyl)-3-(2-(piperidin-1-
- 25 yl)ethyl)-1,4-diazepan-2-one
- (487) 6-chloro-N-(((3S,5S)-3-(2-guanidinoethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (488) 6-chloro-N-(((3S,5S)-3-(2-(3-methylguanidino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-
- 30 diazepan-5-yl)methyl)-2-naphthamide
- (489) N-(((3S,5S)-3-(2-aminoethyl)-1-((R)-2-ethyl-3-methylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (490) N-(((3S,5S)-3-(2-aminoethyl)-1-((S)-2-ethyl-3-methylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (491) 3,4-dichloro-N-(((3S,5S)-1-((R)-2-ethyl-3-methylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-
- 35 1,4-diazepan-5-yl)methyl)benzamide
- (492) 3,4-dichloro-N-(((3S,5S)-1-((S)-2-ethyl-3-methylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- (493) N-(((3S,5S)-3-(2-amino-2-methylpropyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- (494) N-(((3R,5R)-3-(2-amino-2-methylpropyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide

- (495) N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (496) 3,4-dichloro-N-(((3S,5S)-3-(2-(isopropylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 5 (497) N-(((3S,5S)-3-(2-aminoethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- (498) 6-chloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (499) 6-chloro-N-(((3S,5S)-3-(2-methyl-2-(piperidin-1-yl)propyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 (500) 6-chloro-N-(((3R,5R)-3-(2-methyl-2-(piperidin-1-yl)propyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (501) N-(((3S,5S)-3-(2-aminoethyl)-1-((R)-2-ethyl-3-methylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- 15 (502) N-(((3S,5S)-3-(2-aminoethyl)-1-((S)-2-ethyl-3-methylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- (503) 6-chloro-N-(((3S,5S)-1-((R)-2-ethyl-3-methylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (504) 6-chloro-N-(((3S,5S)-1-((S)-2-ethyl-3-methylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 20 (505) 6-chloro-N-(((3S,5S)-3-(2-cyclohexylethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (506) 6-chloro-N-(((3S,5S)-3-hexyl-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 (507) 6-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)carbamoyl)-2-naphthoic acid
- (508) 6-chloro-N-(((3S,5S)-3-(2-(3-isopropylguanidino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (509) 6-chloro-N-(((3S,5S)-3-(4-hydroxybutyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 30 (510) 6-chloro-N-(((3S,5S)-3-(2-methoxyethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (511) N-(((3S,5S)-3-(2-(benzyloxy)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- 35 (512) 6-chloro-N-(((3S,5S)-3-isobutyl-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

- (513) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethyl-3-methylbut-3-enyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (514) 3,4-dichloro-N-(((3S,5S)-1-(2-ethyl-3-methylbut-3-enyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 5 (515) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethyl-3-methylbut-3-enyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- (516) 6-chloro-N-(((3S,5S)-1-((R)-2-ethyl-3-methylbut-3-enyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (517) 6-chloro-N-(((3S,5S)-1-((S)-2-ethyl-3-methylbut-3-enyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 (518) N-(((3S,5S)-1-(cyclohexylmethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)biphenyl-4-carboxamide
- (519) N-(((3S,5S)-1-(cyclohexylmethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(1H-indol-3-yl)acetamide
- 15 (520) N-(((3S,5S)-1-(cyclohexylmethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)quinoline-3-carboxamide
- (521) 3,4-dichloro-N-(((3S,5S)-3-(2-(4,4-difluoropiperidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- (522) 3,4-dichloro-N-(((3S,5S)-3-(2-hydroxyethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 20 (523) 3,4-dichloro-N-(((3S,5S)-3-(2-(3,3-difluoropiperidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- (524) (3S,5S)-5-((3,4-dichlorobenzylamino)methyl)-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-2-one
- 25 (525) 3,4-dichloro-N-(((3S,5S)-1-(2-cyclopropylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- (526) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-cyclopropylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- (527) 6-chloro-N-(((3S,5S)-1-(2-cyclopropylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 30 (528) 3,4-dichloro-N-(((3S,5S)-3-(2-(2,5-dioxopyrrolidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- (529) 6-chloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(3-ureidopropyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 35 (530) 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(1,1,1-trifluoropropan-2-ylamino)ethyl)-1,4-diazepan-5-yl)methyl)benzamide

- (531) 3,4-dichloro-N-(((3S,5S)-3-(2-(3,3-dimethyl-2,5-dioxopyrrolidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- (532) N-(((3S,5S)-3-(2-(azepan-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- 5 (533) 6-chloro-N-(((3S,5S)-3-(2-(3-isopropylureido)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (534) N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)biphenyl-4-carboxamide
- (535) N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-phenylthiazole-4-carboxamide
- 10 (536) 4'-chloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)biphenyl-2-carboxamide
- (537) 6-chloro-N-(((3S,5S)-3-(2-(N-isopropylacetamido)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 (538) 6-chloro-N-(((3S,5S)-3-((isopropylamino)methyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (539) 6-chloro-N-(((3S,5S)-3-(guanidinomethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (540) 2-(2,4-dichlorophenyl)-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acetamide
- 20 (541) N-(((3S,5S)-3-(2-aminoethyl)-1-(2,4-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (542) 3,4-dichloro-N-(((3S,5S)-1-(2,4-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 25 (543) 3,4-dichloro-N-(((3S,5S)-1-(2,4-dichlorobenzyl)-3-(2-(methylsulfonamido)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- (544) 3,4-dichloro-N-(((3S,5S)-1-(2,4-dichlorobenzyl)-3-(2-(4-methylphenylsulfonamido)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- (545) N-(((3S,5S)-3-(2-((S)-2-amino-3-methylbutanamido)ethyl)-1-(2,4-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- 30 (546) N-(((3S,5S)-3-(2-aminoethyl)-1-(2,4-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- (547) 6-chloro-N-(((3S,5S)-1-(2,4-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 35 (548) N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-(2-(thiophen-3-yl)butyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide

- (549) 6-chloro-N-(((3S,5S)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1-((R)-2-(thiophen-3-yl)butyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (550) 6-chloro-N-(((3S,5S)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1-((S)-2-(thiophen-3-yl)butyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 (551) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethyl-2-methylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- (552) 6-chloro-N-(((3S,5S)-1-(2-ethyl-2-methylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (553) 6-chloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(2-morpholinoethyl)-2-oxo-1,4-diazepan-10 5-yl)methyl)-2-naphthamide
- (554) 6-chloro-N-(((3S,5S)-3-(2-morpholinoethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (555) 6-chloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-3-(2-morpholinoethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 (556) 3,4-dichloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-3-(2-morpholinoethyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- (557) N-(3,4-dichlorobenzyl)-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acetamide
- (558) 1-(4-chlorobenzyl)-3-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-20 1,4-diazepan-5-yl)methyl)urea
- (559) N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethyl-2-methylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (560) 3,4-dichloro-N-(((3S,5S)-1-(2-ethyl-2-methylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 25 (561) 6-chloro-N-(((3S,5S)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1-(2,3,5-trichlorobenzyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (562) 6-chloro-N-(((3S,5S)-3-(2-(1-methylethylsulfonamido)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (563) butyl 2-((2S,7S)-7-((6-chloro-2-naphthamido)methyl)-3-oxo-4-((S)-2-phenylbutyl)-1,4-30 diazepan-2-yl)ethylcarbamate
- (564) (S)-6-chloro-N-((3-(1-isopropylpiperidin-4-yl)-2-oxo-1-(2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (565) 6-chloro-N-(((3S,5S)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 35 (566) 5-(4-chlorophenyl)-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)isoxazole-3-carboxamide

- (567) 2,4-dichloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- (568) N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-6-methoxy-2-naphthamide
- 5 (569) 6-chloro-N-([5-¹³C,4-¹⁵N](3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)[¹³C]methyl)-2-naphthamide
- (570) N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-1-methoxy-2-naphthamide
- (571) (E)-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-3-(4-(trifluoromethoxy)phenyl)acrylamide
- 10 (572) 5-(4-chlorophenyl)-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)isoxazole-3-carboxamide
- (573) 2,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 15 (574) 5,6-dichloro-2-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)isoindoline-1,3-dione
- (575) (E)-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)acrylamide
- (576) 6-methoxy-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 20 (577) 1-methoxy-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (578) (E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 25 (579) 6-chloro-N-([5,6,6-²H₃](3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)[²H₂]methyl)-2-naphthamide
- (580) 6-chloro-N-(((3R,5R)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (581) 6-chloro-N-(((3S,5R)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 30 (582) 6-chloro-N-(((3R,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (583) 6-chloro-N-(((3S,5R)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 35 (584) 6-chloro-N-(((3R,5S)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

(585) 6-chloro-N-(((3R,5R)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

5 As stated previously the compounds of formula (I) are antagonists of the MC5R and therefore may be used to modulate the activity of MC5R or a fragment or analogue or functional equivalent thereof by exposing MC5R or a fragment or analogue or functional equivalent thereof to a compound of the invention. This can occur *in vitro* in assays where the downregulation of MC5R activity is desirable, however it is typically more beneficial when
10 utilised in downregulation of MC5R activity in a patient. The amount of downregulation provided by the compounds of the invention will vary from compound to compound and will also be affected by the amount of compound administered. In one embodiment the amount of downregulation is at least 10%. In another embodiment the amount of downregulation is at least 20%. In an even further embodiment the amount of downregulation is at least 50%.

15 Accordingly the methods of the present invention may be used in the treatment of any condition in which modulation of the activity of MC5R or a fragment or analogue or functional equivalent thereof would lead to a beneficial effect on that condition. As such the compounds suitable for use in the present invention may be used in methods of treating, preventing, or
20 controlling a condition associated either directly or indirectly with the activity of MC5R or a fragment or analogue or functional equivalent thereof in a mammal wherein an MC5R modulating amount of the compound of the invention is administered to the mammal. One condition associated with MC5R activity is excess sebum secretion and conditions related thereto. In one embodiment of the method the condition is selected from the group consisting
25 of acne, seborrhoea, and seborrheic dermatitis. In one embodiment the acne is selected from the group consisting of acne vulgaris, acne, acne conglobata and acne fulminans. In one specific embodiment the condition is acne vulgaris.

For example, downregulation of MC5R leads to a reduction of sebum secretion and
30 can thus be used in the treatment or prophylaxis of a number of conditions in which excess sebum secretion is observed such as acne, seborrhoea and seborrheic dermatitis.

The methods of the invention may also be useful in the treatment, prevention or control of a number of conditions that relate to biological processes controlled by MC5R, such
35 as diseases related to inflammation. The compounds of formula (I) may also be useful for the treatment or prevention of cancers, such as Muir-Torre syndrome or other cancers of the sebaceous gland.

Due to their impact on sebum secretion the compounds of formula (I) may also find application in treatments where reduced sebum secretion is desirable such as in cosmetic treatments. The compounds may thus be used in methods of reducing sebum secretion by a mammal, the method comprising administering an effective amount of a compound of formula (I).

The compounds of formula (I) may be used in the treatment of conditions in any species in which MC5R is present, most typically mammals. Examples of species in which MC5R is found and hence species in which the compounds may be used include humans, rats, mice, dogs, rhesus monkey, sheep, zebrafish, goldfish, spiny dogfish, rainbow trout and chickens. In a specific embodiment the mammal is a human.

Administration of compounds within Formula (I) to a patient such as humans can be by topical administration, by any of the accepted modes for enteral administration such as oral or rectal, or by parenteral administration such as subcutaneous, intramuscular, intravenous and intradermal routes. Injection can be bolus or via constant or intermittent infusion. The active compound is typically included in a pharmaceutically acceptable carrier or diluent and in an amount sufficient to deliver to the patient a therapeutically effective dose.

In using the compounds of formula (I) they can be administered in any form or mode which makes the compound bioavailable. One skilled in the art of preparing formulations can readily select the proper form and mode of administration depending upon the particular characteristics of the compound selected, the condition to be treated, the stage of the condition to be treated and other relevant circumstances. We refer the reader to Remingtons Pharmaceutical Sciences, 19th edition, Mack Publishing Co. (1995) for further information.

The compounds of formula (I) can be administered alone or in the form of a pharmaceutical composition in combination with a pharmaceutically acceptable carrier, diluent or excipient. The compounds of formula (I), while effective themselves, are typically formulated and administered in the form of their pharmaceutically acceptable salts as these forms are typically more stable, more easily crystallised and have increased solubility.

The compounds are, however, typically used in the form of pharmaceutical compositions which are formulated depending on the desired mode of administration. The compositions are prepared in manners well known in the art.

A compound of formula (I) is typically combined with the carrier to produce a dosage form suitable for the particular patient being treated and the particular mode of administration. For example, a formulation intended for the oral administration to humans may contain from about 0.5 mg to about 5 g of the compound of the invention, compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 99.95 percent of the total composition. Representative dosage forms will generally contain between from about 1 mg to about 500 mg of a compound of the invention, typically 25 mg, 50 mg, 100 mg, 200 mg, 300 mg, 400 mg, 500 mg, 600 mg, 800 mg, or 1000 mg. Compounds of the present invention may also be formulated for topical delivery in formulations such as solutions, ointments, lotions, gels, creams, microemulsions or transdermal patches. For example, these topical formulations may contain from 0.005 to 5% (wt/wt or wt/vol) of a compound of the invention.

The compounds of formula (I) may be used or administered in combination with one or more additional drug (s). The compounds of the present invention may be used in combination with one or more other pharmaceutically-active compounds, such as other anti-acne treatments. In one embodiment the other pharmaceutically active agent is selected from the group consisting of antibiotics, retinoids, anti-androgens, and steroids. Examples of other pharmaceutically active compounds that may be combined with a compound of formula (I) and administered in concurrent or sequential combination therewith may include, by way of non-limiting example, other anti-acne agents such as oral retinoids (e.g. isotretinoin), topical retinoids (e.g. isotretinoin, adapalene, tazarotene), oral or topical antibiotics (e.g. clindamycin, erythromycin, minocycline, tetracycline, benzoyl peroxide), or hormonal therapies (e.g. drospirenone, norgestimate - ethinyl estradiol, cyproterone acetate). As stated these components can be administered in the same formulation or in separate formulations. If administered in separate formulations the compounds of the invention may be administered sequentially or simultaneously with the other drug(s).

Pharmaceutical compositions suitable for use in the invention for parenteral injection comprise pharmaceutically acceptable sterile aqueous or non aqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and non aqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the

maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preservative, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of micro-organisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents that delay absorption such as aluminium monostearate and gelatin.

If desired, and for more effective distribution, the compounds can be incorporated into slow release or targeted delivery systems such as polymer matrices, liposomes, and microspheres.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions that can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

If desired, and for more effective distribution, the compounds can be incorporated into slow release or targeted delivery systems such as polymer matrices, liposomes, and microspheres.

The active compounds can also be in microencapsulated form, if appropriate, with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminium metahydroxide, bentonite, agar-agar, and tragacanth, and mixtures thereof.

Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which

are solid at room temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

For topical administration, the active agent may be in the form of an ointment, cream,
5 suspension, lotion, powder, solution, paste, gel, spray, aerosol or oil. Alternatively, the composition may be delivered via a liposome, nanosome, rivosome, or nutri-diffuser vehicle. Alternately, a formulation may comprise a transdermal patch or dressing such as a bandage impregnated with an active ingredient and optionally one or more carriers or diluents. To be administered in the form of a transdermal delivery system, the dosage administration will, of
10 course, be continuous rather than intermittent throughout the dosage regimen. Methods for producing formulations for topical administration are known in the art.

The compositions used for topical administration typically contain a pharmaceutically acceptable carrier which may be any vehicle that is toxicologically and pharmaceutically
15 acceptable. Typical pharmaceutically acceptable carriers that can be used in compositions of the present invention include water, ethanol, acetone, isopropyl alcohol, stearyl alcohol, freons, polyvinyl pyrrolidone, propylene glycol, polyethylene glycol, fragrances, gel-producing materials, mineral oil, stearic acid, spermaceti, sorbitan, monoleate, polysorbates, "Tweens," sorbitol, methyl cellulose, petrolatum, a mineral oil (vaseline oil), which may be any petroleum
20 based product; modified or unmodified vegetable oils such as peanut oil, wheatgerm oil, linseed oil, jojoba oil, apricot kernel oil, walnut oil, palm oil, pistachio oil, sesame oil, colza oil, cade oil, corn germ oil, peach kernel oil, poppyseed oil, pine oil, castor oil, soya oil, safflower oil, coconut oil, hazelnut oil, grapeseed oil, avocado oil, soy oil, sweet almond oil, calophyllum oil, castor oil, olive oil, sunflower oil, or animal oils such as whale oil, seal oil, menhaden oil,
25 halibut liver oil, cod liver oil, cod, tuna, turtle tallow, horse's hoof, sheep's foot, mink, otter, marmot oil and the like; synthetic oils such as silicon oil such as dimethylpolysiloxane; alkyl and alkenyl esters of fatty acids, such as isopropyl esters of myristic, palmitic and stearic acids and fatty esters which are solid at room temperature; waxes such as lanolin wax, candelilla wax, spermaceti, cocoa butter, karite butter, silicon waxes, hydrogenated oils which
30 are solid at room temperature, sucro-glycerides, oleates, myristates, linoleates, stearates, paraffin, beeswax, carnauba wax, ozokerite, candelilla wax, microcrystalline wax; fatty alcohols such as lauryl, cetyl, myristyl, stearyl, palmityl and oleyl alcohols; polyoxyethylated fatty alcohols; and wax esters, lanolin and its derivatives, perhydrosqualene and saturated esters, ethyl palmitate, isopropyl palmitate, alkyl myristates such as isopropyl myristate, butyl
35 myristate and decyl myristate, hexyl stearate, triglyceride esters, triglycerides of octanoic and decanoic acid, cetyl ricinoleate, stearyl octanoate (Purcellin oil), fatty acids, polyhydric alcohols, polyether derivatives, fatty acid monoglycerides, polyethylene glycol, propylene

glycol, alkyl ethoxy ether sulfonates, ammonium alkyl sulfates, fatty acid soaps, and hydrogenated polyisobutene, and mixtures of waxes and oils.

The compositions for topical administration may be formulated in numerous forms. However, the composition may often take the form of an aqueous or oily solution or dispersion or emulsion or a gel or a cream. An emulsion may be an oil-in-water emulsion or a water-in-oil emulsion.

The oil phase of water-in-oil or oil-in-water emulsions may comprise for example: a) hydrocarbon oils such as paraffin or mineral oils; b) waxes such as beeswax or paraffin wax; c) natural oils such as sunflower oil, apricot kernel oil, shea butter or jojoba oil; d) silicone oils such as dimethicone, cyclomethicone or cetyldimethicone; e) fatty acid esters such as isopropyl palmitate, isopropyl myristate, dioctylmaleate, glyceryl oleate and cetostearyl isononanoate; f) fatty alcohols such as cetyl alcohol or stearyl alcohol and mixtures thereof (eg cetearyl alcohol); g) polypropylene glycol or polyethylene glycol ethers, eg PPG-14 butyl ether; or h) mixtures thereof.

Emulsifiers used may be any emulsifiers known in the art for use in water-in-oil or oil-in-water emulsions. Known cosmetically acceptable emulsifiers include: a) sesquioleates such as sorbitan sesquioleate, available commercially for example under the trade name Arlacel 83 (ICI), or polyglyceryl-2-sesquioleate; b) ethoxylated esters of derivatives of natural oils such as the polyethoxylated ester of hydrogenated castor oil available commercially for example under the trade name Arlacel 989 (ICI); c) silicone emulsifiers such as silicone polyols available commercially for example under the trade name ABIL WS08 (Th. Goldschmidt AG); d) anionic emulsifiers such as fatty acid soaps e.g. potassium stearate and fatty acid sulphates e.g. sodium cetostearyl sulphate available commercially under the trade name Dehydag (Henkel); e) ethoxylated fatty alcohols, for example the emulsifiers available commercially under the trade name Brij (ICI); f) sorbitan esters, for example the emulsifiers available commercially under the trade name Span (ICI); g) ethoxylated sorbitan esters, for example the emulsifiers available commercially under the trade name Tween (ICI); h) ethoxylated fatty acid esters such as ethoxylated stearates, for example the emulsifiers available commercially under the trade name Myrj (ICI); i) ethoxylated mono-, di-, and tri-glycerides, for example the emulsifiers available commercially under the trade name Labrafil (Alfa Chem.); j) non-ionic self-emulsifying waxes, for example the wax available commercially under the trade name Polawax (Croda); k) ethoxylated fatty acids, for example, the emulsifiers available commercially under the trade name Tefose (Alfa Chem.); l)

methylglucose esters such as polyglycerol-3 methyl glucose distearate available commercially under the name Tegocare 450 (Degussa Goldschmidt); or m) mixtures thereof.

Gels for topical administration may be aqueous or non-aqueous. Aqueous gels are preferred. The gel will contain a thickening agent or gelling agent in order to give sufficient viscosity to the gel. A variety of thickening agents may be used according to the nature of the liquid carrier and the viscosity required and these are recited hereinafter. A particularly suitable thickener is a copolymer of acryloyl dimethyl tauric acid (or a salt thereof), preferably a copolymer of that monomer with another vinylic monomer. For example, the thickening agent is a copolymer of a salt of acryloyl dimethyl tauric acid with another vinylic monomer. The salt may be a salt of a Group I alkali metal, but is more preferably an ammonium salt. Examples of suitable copolymer thickening agents are: i) Ammonium acryloyl dimethyl taurate I vinyl pyrrolidone copolymer, ie a copolymer of ammonium acryloyl dimethyl taurate and vinyl pyrrolidone (1-vinyl-2-pyrrolidone).

15

The composition may additionally comprise other skincare active agents which are well known in the art which may be effective to aid the normal functioning of the skin. One group of preferred compositions comprise hydrolysed milk protein to regulate sebum production.

20

The composition may additionally comprise other components which will be well known to those skilled in the art such as emollients, humectants, emulsion stabilising salts, preservatives, chelating agents or sequestering agents (sequestrants), abrasives, anti-oxidants, stabilisers, pH adjusters, surfactants, thickeners, diluents, perfumes and colourings.

25

The topical formulations may desirably include a compound that enhances absorption or penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethylsulfoxide and related analogues.

30

The amount of compound administered will preferably treat and reduce or alleviate the condition. A therapeutically effective amount can be readily determined by an attending diagnostician by the use of conventional techniques and by observing results obtained under analogous circumstances. In determining the therapeutically effective amount a number of factors are to be considered including but not limited to, the species of animal, its size, age and general health, sex, diet, the specific condition involved, the severity of the condition, the response of the patient to treatment, the particular compound administered, the mode of

35


administration, the bioavailability of the preparation administered, the dose regime selected, the use of other medications and other relevant circumstances.

A preferred dosage will be a range from about 0.01 to 300 mg per kilogram of body weight per day. A more preferred dosage will be in the range from 0.1 to 100 mg per kilogram of body weight per day, more preferably from 0.2 to 80 mg per kilogram of body weight per day, even more preferably 0.2 to 50 mg per kilogram of body weight per day. A suitable dose can be administered in multiple sub-doses per day.

SYNTHESIS OF COMPOUNDS FOR USE IN THE INVENTION

The general synthetic route to the compounds for use in the invention proceeds through the key intermediate A, produced as outlined in Schemes 1 or 2.

In Scheme 1, an amino acid derivative $V-N(R^2)-Y-CO_2H$ ($V = R^1X$ or an amine protecting group P^1) is converted to a Weinreb amide via activation of the carboxyl group and amidation with *N*-methyl methoxyamine. Addition of a vinyl Grignard reagent produces the aminoalkyl vinyl ketone, which undergoes conjugate addition by the $R^6R^7R^8C-(CR^{5a}R^{5b})_rNH_2$ amine component (shown as WNH_2 for simplicity). The resulting secondary amine is acylated under standard peptide coupling conditions with the protected amino acid, $P^2-NHCH(U)-CO_2H$, where U represents either the final R side chain, a protected final side chain $R-P^3$, or a precursor that requires chemical modification to form the final R side chain. Deprotection of the P^2 protecting group is followed by intramolecular reductive amination of the ketone using standard reduction conditions, such as H_2/Pd catalyst, $NaBH_4$, $NaBH_3CN$, or $NaBH(OAc)_3$, forming key intermediate **A**. If $Y = CH_2$ or CH_2CH_2 , **A** is formed as the predominant diastereomer. If $V = R^1X$ and $U = R$, **A** is the final product.


Scheme 1: Synthesis of Intermediate A via Intramolecular Reductive Amination

where U = R, a protected form thereof or a precursor thereof,

V = P¹ or R¹X, and W = R⁶R⁷R⁸C(CR^{5a}R^{5b})_r-

Final product if V = R¹X, U = R



In Scheme 2, an alternate route to the desired intermediate **A** begins with the same Weinreb amide formation, vinyl Grignard addition, and amine conjugate addition. At this point, the secondary amine is protected with an amine protecting group P⁴. The ketone is then reductively aminated with a protected amino ester, H₂NCH(U)-CO₂P⁵, producing a mixture of diastereomers that are carried through the next reaction steps. The ring system is generated by deprotection of the P⁴ and P⁵ protecting groups, followed by amide bond formation using standard peptide coupling reagents. Alternatively, the P⁴ protecting group is removed and cyclization achieved by thermal or base-induced cyclization with the P⁵-protected ester. The cyclization produces a mixture of two diastereomers, A and B, from which the preferred diastereomers **A** can be separated by chromatography.

Scheme 2: Synthesis of Intermediate A via Intermolecular Reductive Amination

where U = R, a protected form thereof or a precursor thereof,
V = P¹ or R¹X, and W = R⁶R⁷R⁸C(CR^{5a}R^{5b})_r-
final product if V = R¹X, U = R


The key intermediate A may be the final product if U = R and V = R¹X, but otherwise is converted into the final product as illustrated in Schemes 3, 4 and 5.

- 5 In Scheme 3, where V = R¹X, the final product is obtained by modification of the U side chain, such as removal of a P³ protecting group, or removal of a P³ protecting group followed by further chemical modification.

Scheme 3: V = R¹X


In Scheme 4, where $V = P^1$, the final product is obtained by removal of the P^1 protecting group followed by introduction of the R^1X substituent. If $U = R$, this produces the final product. Alternatively, the U side chain is then modified to produce the final R group as in Scheme 3.

Scheme 4: $V = P^1$



In Scheme 5, where $V = P^1$, the final product is obtained by first modifying the U side chain to produce the final R group as in Scheme 3. This is followed by removal of the P^1 protecting group followed by introduction of the R^1X substituent.

Scheme 5: $V = P^1$



It is also possible to modify the W substituent, if desired, during these reaction sequences.

Examples

The following examples are intended to illustrate the embodiments disclosed and are not to be construed as being limitations thereto. Additional compounds, other than those described below, may be prepared using the following described reaction schemes as


discussed above or appropriate variations or modifications thereof. All starting materials described in the Examples below are commercially available or readily synthesized by those skilled in the art.

5 Instrumentation

HPLC analyses were carried out on an Agilent 1100 Series Purification System with a Phenomenex Synergi 4 μ Max-RP 80A, 50 x 2.00 mm analytical HPLC column, with peak detection by UV. The standard analysis employed a 1 mL/min flow rate of 0.05% trifluoroacetic acid (TFA) in water (Solvent A) and 0.05% TFA in 90:10 acetonitrile:water (Solvent B), using a gradient of 5% B (initial) to 95% B over 9 min. Mass spectra were run on an Applied Biosystems MDS Sciex API 2000 LC/MS/MS triple quadrupole mass spectrometer and analyzed by ion spray mass spectrometry (ISMS). Preparative scale HPLC was carried out on a Waters Delta Prep 3000 HPLC system with peak detection by UV (Waters model 486 tunable absorbance detector), using Phenomenex Luna 10 μ C5 100A, 250 x 21.20 mm (20 mg scale), Phenomenex Luna 15 μ C8(2) 100A, 250 x 30.00 mm (50 mg scale), or Phenomenex Luna 15 μ C8(2) 100A, 250 x 50.00 mm (100 mg scale) HPLC columns. The solvent system employed various gradients of 0.05% TFA in water (Solvent A) and 0.05% TFA in 90:10 acetonitrile:water (Solvent B).

The following examples 1 to 7 provide general synthetic procedures that may be followed in order to carry out the transformations described in schemes 1 to 5. In order to make different end products using these procedures it is necessary to either vary a variable group on the starting material or to vary a variable group on one of the reagents depending upon the nature of the reaction. It will be apparent to a skilled addressee from a reading of the general procedures how to vary either the starting material or the reagents used in the procedure to produce differing end products. In addition depending upon the starting materials and the reagents it may be necessary and/or desirable to make slight modifications to the described general procedures in order to provide the most facile synthesis of the desired end product.


Example 1 – General Procedure - Weinreb Amide Formation



BOP reagent (100 mmol) and diisopropylethylamine (DIPEA) (100 mmol) is added to a stirred solution of the amino acid (1) (100 mmol) in dichloromethane (DCM) (100 mL). The


solution is then stirred at room temperature for 10 mins, before addition of a premixed solution of *N,O*-dimethylhydroxylamine hydrochloride (100 mmol) and DIPEA (100 mmol) followed by stirring at room temperature overnight. The DCM is then removed by rotary evaporation and the residue taken up in ethyl acetate (EtOAc) (200 mL). The organic phase is then washed with 1N HCl (3 x 100 mL), H₂O (3 x 100 mL), saturated NaHCO₃ aqueous solution (3 x 100 mL) and brine (1 x 10 mL). The organic phase is then dried (MgSO₄) and the EtOAc removed to give the Weinreb amide (2) as a white solid or an oil.

Example 2 – General Procedure - Vinyl Grignard Addition to Weinreb Amide to Form α,β -unsaturated ketones of formula (3)




To the Weinreb amide (2) (15 mmol) in DCM (10 mL) at 0°C is added vinyl magnesium bromide (45 mmol) in THF (45 mL). The reaction is stirred for 2 hrs and monitored by HPLC. The reaction is then quenched by adding it to a mixture of ice and 1M HCl (200 mL). The aqueous mixture is extracted with DCM (3x 100 mL) and the organic layers combined and washed with 1M HCl (2x 200 mL) and H₂O (3x 100 mL). The organic phase is dried (MgSO₄) to provide a solution of the α,β -unsaturated ketone (3). The α,β -unsaturated ketone (3) may be isolated by rotary evaporation or it may be used in solution without further purification. If the intention is to use the α,β -unsaturated ketone (3) in solution the volume is reduced to 100 mL by rotary evaporation and stored for later use.

Example 3 General Procedure – Conjugate Addition of Amine to α,β -unsaturated ketones of formula (3) to produce compounds of formula (4)




To the amine W-NH₂ (7.4 mmol) in DCM (10 mL) is added a solution of the α,β -unsaturated ketone (3) (5.7 mmol) in DCM (50 mL). The solution is stirred at room temperature for 15 mins, or until analysis indicates that all of (3) has been consumed. The solution of compound (4) is immediately used without purification for the subsequent reaction.

Example 4 General Procedure – Acylation of Aminoketone (4)

The amine acid $\text{P}^2\text{-NHCH}(\text{U})\text{-CO}_2\text{H}$ (15 mmol) and DIC (15 mmol) is added to a solution of DCM containing 10 mmol of the conjugate addition adduct **4**. The reaction is stirred at room temperature overnight. The DCM is removed by rotary evaporation and the residue is then subjected to column chromatography on silica gel using petroleum spirit:EtOAc to give **5**.

As an alternative, the DIC may be replaced with HATU (15 mmol) and DIPEA (15 mmol). The reaction is stirred at room temperature overnight. The DCM is removed by rotary evaporation and the residue is taken up in EtOAc (100 mL). The organic layer is washed with saturated sodium bicarbonate solution (2 x 100 mL), saturated ammonium chloride solution (2 x 100 mL) and brine (2 x 100 mL). The organic phase is dried and the solvent removed under reduced pressure. The residue is subjected to column chromatography on silica gel using petroleum ether:EtOAc to give **5**.

Example 5 General Procedure – P^2 Deprotection and Cyclization

The procedure adopted for the removal of the P^2 protecting group will vary depending upon the exact nature of the protecting group. As will be appreciated by a skilled addressee a large number of possible protecting groups may be used and a skilled worker in the art will readily be able to determine an appropriate procedure for the removal of any particular protecting group from procedures known in the art. Nevertheless in order to assist the reader general procedures for the removal of the more common protecting groups are provided.


P^2 = Fmoc: To compound **5** (2 mmol) in DCM (3 mL) is added diethylamine (20 mmol). The reaction is stirred at room temperature for 1 hr. The DCM and diethylamine is then removed

by rotary evaporation. DCM (5 mL) and sodium triacetoxyborohydride (3 mmol) are then added, and the reaction stirred overnight at room temperature. The organic phase is washed with saturated sodium bicarbonate solution (25 mL), dried (MgSO₄) and the DCM removed to give the cyclised product **A**. This may be purified by flash chromatography on silica gel or used without purification.

P² = Boc: To compound **5** (2 mmol) in DCM (3 mL) is added TFA (3 mL) and the reaction stirred at room temperature for 2 hrs. The DCM and TFA are then removed by rotary evaporation. DCM (5 mL) and sodium triacetoxyborohydride (3 mmol) is then added, and the reaction stirred overnight at room temperature. The organic phase is washed with saturated sodium bicarbonate solution (25 mL), dried (MgSO₄) and the DCM removed to give the cyclised product **A**. This may be purified by flash chromatography on silica gel or used without purification.

P² = Cbz: A mixture of crude **5** (1 mmol) and 5% Pd/C (200 mg) in 2-propanol (15 mL) is shaken at room temperature under hydrogen (30 psi) for 24 hrs. The mixture is then filtered through a pad of Celite and the filtrate concentrated under reduced pressure to give a crude product. Purification by flash chromatography on silica gel (100% EtOAc) may be used to give **A**.

Example 6 General Procedure – P¹ Deprotection and Derivatization with R¹X



The procedure adopted for the removal of the P¹ protecting group will vary depending upon the exact nature of the protecting group. As will be appreciated by a skilled addressee a large number of possible protecting groups may be used and a skilled worker in the art will readily be able to determine an appropriate procedure for the removal of any particular protecting group from procedures known in the art. Nevertheless in order to assist the reader general procedures for the removal of the more common protecting groups are provided.

Deprotection, $P^1 = \text{Cbz}$:

To the cyclised product **A** (1 mmol) in methanol (5 mL) is added catalytic Pd/C. The reaction is stirred under a hydrogen atmosphere overnight. The reaction mixture is filtered through Celite and the methanol removed by rotary evaporation to give the free amine. The amine
5 may be used in the next reaction without purification.

Deprotection, $P^1 = \text{Boc}$:

To the cyclised product **A** (1 mmol) in DCM (1 mL) is added TFA (1mL) and the reaction stirred at room temperature for 2 hrs. The solvent is removed by rotary evaporation to give
10 the amine TFA salt, which may be used in the next reaction without purification.


Deprotection, $P^1 = \text{Alloc}$:

To the cyclised product **A** (1 mmol) in DCM (6 mL) is added 1,3-dimethylbarbituric acid (0.2 mmol) and palladium tetrakis triphenylphosphine (10 mg). The reaction is evacuated and
15 stirred at room temperature for 1 hr. The DCM is removed under reduced pressure to give the crude free amine, which may be used in the next reaction without purification.

Derivatisation with R^1X when $X = \text{C}(=\text{O})$:

To the free amine (1 mmol) in DCM (5 mL) is added DIPEA (1 mmol), BOP reagent (1.5 mmol) and acid component $R^1\text{CO}_2\text{H}$ (1.5 mmol). The reaction is stirred at room temperature
20 for 2 hrs. Rotary evaporation and preparative HPLC gives the purified adduct.

Example 7 General Procedure – U Modification via P^3 Deprotection and Dialkylation with Dibromide




The procedure adopted for modification of U via deprotection and derivatization will vary depending on the exact nature of the U group. As will be appreciated by a skilled
30 addressee a large number of modifications are possible, and a skilled worker in the art will readily be able to determine an appropriate procedure for the conversion into a desired R group. Nevertheless in order to assist the reader, one general modification procedure commonly employed for a number of the following examples is provided.

P³ = Boc:

To the protected amine (1 mmol) in DCM (5 mL) is added TFA (5 mL) and the reaction stirred at room temperature for 2 hrs. DCM (20 mL) is added and the solution is washed with saturated sodium bicarbonate solution (20 mL), dried (MgSO₄) and evaporated to give the crude amine. To the crude amine is added DMF (0.5 mL), potassium carbonate (50 mg) and 1,5-dibromopentane (5 mmol). The reaction mixture is stirred at room temperature for 1.5 hrs, after which DCM (20 mL) is added, the organic layer washed with saturated sodium bicarbonate solution (20 mL) and H₂O (20 mL), dried (MgSO₄) and evaporated. The residue may be purified by preparative HPLC to give the piperidinyll product. The purified product is isolated as the TFA salt, but is readily converted into the free base via neutralisation with aqueous NaHCO₃ and extraction into an organic solvent, or further converted into the HCl salt by acidification with 1N HCl.

Example 8 – Synthesis of Compound 8 N-(2-(methoxy(methyl)amino)-2-oxoethyl)-2-naphthamide




8

N-(2-(methoxy(methyl)amino)-2-oxoethyl)-2-naphthamide

To a mixture of 2-naphthoic acid (5.8 g, 33.7 mmol), 2-amino-*N*-methoxy-*N*-methylacetamide (Gly Weinreb amide; prepared from Boc-Gly Weinreb amide **15** as in Example 43) (3.8 g, 32.1 mmol) and DIPEA (12.0 mL, 68.9 mmol) in DCM (70 mL) was added BOP (14.9 g, 33.7 mmol) in one portion at room temperature. The resulting mixture was stirred for 1 hr then saturated NaHCO₃ aqueous solution was added. The organic layer was washed with brine (5 x 60 mL) and 1 N HCl (2 x 30 mL), dried over MgSO₄, filtered and concentrated under reduced pressure to give the crude product, which was used in the next reaction without further purification.

Example 9 – Synthesis of Compound 9 N-(2-(methoxy(methyl)amino)-2-oxoethyl)-2-naphthamide




9

N-(2-oxobut-3-enyl)-2-naphthamide


To a solution of **8** (3.5 g, 12.85 mmol) in dry THF (10 mL) was added a solution of vinylmagnesium bromide in THF (1 M, 31 mL) slowly at 0°C. After addition, the resulting mixture was stirred at room temperature for 1 hr then was poured into an icy 1 N HCl solution (50 mL). The aqueous layer was extracted with DCM (3 x 80 mL) and the combined organic layers were dried over MgSO₄, filtered and concentrated under reduced pressure to give the crude product. **MS (ESI)** 240 (M+1); HPLC *t_R* 5.46 min.

Example 10 – Synthesis of Compound 10 N-(4-(3,5-dichlorobenzylamino)-2-oxobutyl)-2-naphthamide



To a solution of 3,5-dichlorobenzylamine (12 mg, 0.068 mmol) in DCM (0.2 mL) was added a solution of **9** (13 mg, 0.054 mmol) in DCM (0.5 mL) at room temperature. The resulting mixture was stirred until all of the **9** had been consumed (within one hr) and then was used straight in the next reaction. **MS (ESI)** 415 (M+1); HPLC *t_R* 6.00 min.


Example 11 – Synthesis of Compound 11 (S)-N-(4-(5-(3-Pbf-guanidino)-2-(Fmoc-amino)-N-(3,5-dichlorobenzyl)pentanamido)-2-oxobutyl)-2-naphthamide



To a solution of freshly prepared aminoketone **10** in DCM (2 mL) was added Fmoc-L-Arg(Pbf)-OH (53 mg, 0.082 mmol) followed by DIC (12.5 µl, 0.082 mmol) at room temperature. The resulting mixture was stirred for 2 hrs then the solvent was removed under

reduced pressure. The residue was filtered through a short plug of silica gel eluting with DCM followed by EtOAc to give the desired product **11** as a white solid. It was used in the next step without further purification. **MS (ESI)** 1045 (M+1); HPLC t_R 9.99 min.


5 **Example 12 – Synthesis of Compound 12 (S)-N-(4-(5-(3-Pbf-guanidino)-2-amino-N-(3,5-dichlorobenzyl)pentanamido)-2-oxobutyl)-2-naphthamide**

**12**

(S)-N-(4-(5-(3-Pbf-guanidino)-2-amino-N-(3,5-dichlorobenzyl)pentanamido)-2-oxobutyl)-2-naphthamide

Diethylamine (0.5 mL) was added to Fmoc-protected **11** (56 mg, 0.054 mmol) at room temperature and the resulting mixture was stirred for 30min. The excess amount of the diethylamine was removed under reduced pressure to give the desired free amine **12**. It was used in the next step without further purification. **MS (ESI)** 823 (M+1); HPLC t_R 7.49 min.

15 **Example 13 – Synthesis of Compound 13 N-(((3S,5S)-3-(3-(3-Pbf-guanidino)propyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide**

**13**


N-(((3S,5S)-3-(3-(3-Pbf-guanidino)propyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

The amino ketone **12** (44 mg, 0.053 mmol) in DCM (2 mL) was cyclized by addition of NaBH(OAc)₃ (40mg, 0.18 mmol) in one portion at room temperature. The resulting mixture was stirred for 3 hrs, followed by addition of saturated NaHCO₃ aqueous solution (3 mL). The aqueous layer was extracted with DCM (3 x 3 mL) and the combined organic layers were

dried over MgSO_4 , filtered and concentrated under reduced pressure. The residue was filtered through a short plug of silica gel eluting with DCM followed by EtOAc then EtOAc/IPA (9:1) to give the desired product **13** as a white solid. It was used in the next step without further purification. **MS (ESI)** 807 ($\text{M}+1$); HPLC t_R 7.75 min.

5


Example 14 – Synthesis of Compound 14 *N*-(((3*S*,5*S*)-1-(3,5-dichlorobenzyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

**14**

N-(((3*S*,5*S*)-1-(3,5-dichlorobenzyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

A solution of TFA/DCM (2:1) (1 mL) with 5% H_2O was added to **13** at room temperature and the resulting mixture was stirred for 4 hrs. The solvents were removed under reduced pressure and the residue was purified by prep HPLC (100% H_2O to MeCN/ H_2O 9:1, gradient) to give **14** (7.6 mg) as a white solid (TFA salt). The overall yield (from **9**) was ca. 18%. **MS (ESI)** 556.2 ($\text{M}+1$); HPLC t_R 5.74 min.

Example 15 – Synthesis of Compound 15 *tert*-butyl 2-(methoxy(methyl)amino)-2-oxoethylcarbamate (Boc-Gly Weinreb amide)

**15**


tert-butyl 2-(methoxy(methyl)amino)-2-oxoethylcarbamate

To a stirred mixture of Boc-Gly-OH (20 g, 114.1 mmol), DIPEA (19.8 mL, 114.1 mmol) and BOP (50.5 g, 114.1 mmol) in DCM (20 mL) was added a pre-mixed solution of *N*,*O*-dimethylhydroxylamine hydrochloride (11.2 g, 114.1 mmol) and DIPEA (19.8 mL, 114.1 mmol) in DCM (20 mL) at room temperature. The resulting mixture was stirred for 16 h then washed with 1N HCl (3 x 120 mL), H_2O (3 x 120 mL), saturated NaHCO_3 aqueous solution (3 x 120 mL) and brine (40 mL), dried over MgSO_4 , filtered and concentrated under reduced

pressure to give **15** as a white solid (20 g, 80%), which was used in the next step without further purification.

MS(ESI) 219 (M+1); HPLC t_R 4.12 min.

5 **Example 16 – Synthesis of Compound 16 tert-butyl 2-oxobut-3-enylcarbamate**




16

tert-butyl 2-oxobut-3-enylcarbamate

At 0°C a solution of vinylmagnesium bromide in THF (184 mL, 1 M) was added in one portion to Weinreb amide **15** (20 g, 91.6 mmol) under nitrogen with stirring. The resulting mixture was allowed to stir for 2 h, and poured into a 1N HCl/ice mixture (400 mL). The aqueous mixture was extracted with DCM (5 x 100 mL), the combined DCM extract was washed with 1N HCl (2 x 100 mL), saturated NaHCO₃ aqueous solution (100 mL) and brine (100 mL), then dried over MgSO₄. Solvent was removed under reduced pressure gave the ketone **16** (12.9 g, 76%) as a pale yellow oil, which was used in the next step without further purification. **MS(ESI)** 186 (M+1); HPLC t_R 4.19 min.

15

Example 17 – Synthesis of Compound 17 tert-butyl 4-(2,2-diphenylethylamino)-2-oxobutylcarbamate




17

tert-butyl 4-(2,2-diphenylethylamino)-2-oxobutylcarbamate

To a stirred solution of 2,2-diphenylethylamine (0.33 g, 1.66 mmol) in DCM (10 mL) was added α,β -unsaturated ketone **16** (0.31 g, 1.66 mmol) at room temperature. Stirring continued for 2 h; the crude reaction mixture of **17** was used in the next step without purification. **MS(ESI)** 383 (M+1); HPLC t_R 5.98 min

20


Example 18 – Synthesis of Compound 18 (S)-tert-butyl 3-methyl-4,8-dioxo-10-phenyl-2,9-dioxa-3,7-diazadecane-6-carboxylate



(S)-tert-butyl 3-methyl-4,8-dioxo-10-phenyl-2,9-dioxa-3,7-diazadecane-6-carboxylate

To a suspension of Cbz-L-Asp-OtBu-DCHA salt (10.1 g, 20.0 mmol), N,O-dimethylhydroxylamine-HCl (5.9 g, 60.5 mmol) and DIPEA (12.0 mL, 68.9 mmol) in DCM (150 mL) was added BOP (10.6 g, 24.0 mmol) in one portion at room temperature. The resulting suspension was stirred for 3 hrs then H₂O (100 mL) was added. The organic layer was washed with 1 N HCl (2 x 100mL), saturated NaHCO₃ aqueous solution (2 x 100 mL) and brine (3 x 100 mL) and then dried over MgSO₄, filtered and concentrated under reduced pressure to give the crude product. Purification by flash chromatography on silica gel (PET ether/ EtOAc 1:2) gave **18** (6.4 g, 87%) as a colorless oil. **MS (ESI)** 367 (M+1); HPLC *t_R* 6.87 min.


Example 19 – Synthesis of Compound 19 (S)-3-methyl-4,8-dioxo-10-phenyl-2,9-dioxa-3,7-diazadecane-6-carboxylic acid



(S)-3-methyl-4,8-dioxo-10-phenyl-2,9-dioxa-3,7-diazadecane-6-carboxylic acid

Compound **18** (300 mg, 0.82 mmol) was dissolved in a TFA/DCM (1:1) solution (2 mL) and the resulting mixture was stirred at room temperature for 2 hrs. The solvents were removed under reduced pressure and the residue was re-dissolved in DCM (10 mL). This solution was washed with 1 N HCl (1 x 10mL) and the organic layer was dried over MgSO₄, filtered and concentrated under reduced pressure to give the crude product **19** (235 mg, 92%), which was used in the next reaction without further purification. **MS (ESI)** 311 (M+1); HPLC *t_R* 4.96 min.


Example 20 – Synthesis of Compound 20 (S)-benzyl 8-(2,2-diphenylethyl)-3,16,16-trimethyl-4,7,11,14-tetraoxo-2,15-dioxa-3,8,13-triazaheptadecan-6-ylcarbamate



(S)-benzyl 8-(2,2-diphenylethyl)-3,16,16-trimethyl-4,7,11,14-tetraoxo-2,15-dioxa-3,8,13-triazaheptadecan-6-ylcarbamate

Compound 20 was prepared from Compound 17 and 19 following the procedure of
 5 Example 5. **MS (ESI)** 675 (M+1); HPLC t_R 8.31 min.


Example 21 – Synthesis of Compound 21 tert-butyl ((3S,5S)-1-(2,2-diphenylethyl)-3-(2-(methoxy(methyl)amino)-2-oxoethyl)-2-oxo-1,4-diazepan-5-yl)methylcarbamate



tert-butyl ((3S,5S)-1-(2,2-diphenylethyl)-3-(2-(methoxy(methyl)amino)-2-oxoethyl)-2-oxo-1,4-diazepan-5-yl)methylcarbamate

10 A mixture of crude 20 (350 mg) and 5% Pd/C (200 mg) in 2-propanol (15 mL) was shaken at room temperature under hydrogen (30 psi) for 24 hrs. The mixture was then filtered through a pad of Celite and the filtrate was concentrated under reduced pressure to give the crude product. Purification by flash chromatography on silica gel (100% of EtOAc) gave 21 (175 mg, 65% over 3 steps) as a white solid. **MS (ESI)** 525 (M+1); HPLC t_R 6.24
 15 min.


Example 22 – Synthesis of Compound 22 2-((2*S*,7*S*)-7-(aminomethyl)-4-(2,2-diphenylethyl)-3-oxo-1,4-diazepan-2-yl)-*N*-methoxy-*N*-methylacetamide

**22**

2-((2*S*,7*S*)-7-(aminomethyl)-4-(2,2-diphenylethyl)-3-oxo-1,4-diazepan-2-yl)-*N*-methoxy-*N*-methylacetamide

Compound **21** (175 mg, 0.333 mmol) was dissolved in a TFA/DCM (1:1) solution (1 mL) and the resulting mixture was stirred at room temperature for 2 hrs. The solvents were removed under reduced pressure and the residue was re-dissolved in EtOAc (20 mL). Saturated NaHCO₃ aqueous solution (10 mL) and brine (10 mL) were added to the above solution and the aqueous layer was extracted with EtOAc (9 x 20 mL). The combined organic layers were dried over MgSO₄, filtered and concentrated under reduced pressure to give the crude product **22** (120 mg, 85%) as a yellow solid, which was used in the next reaction without further purification. **MS (ESI)** 425 (M+1); HPLC *t_R* 5.20 min.


Example 23 – Synthesis of Compound 23 N-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(2-(methoxy(methyl)amino)-2-oxoethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide

**23**

N-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(2-(methoxy(methyl)amino)-2-oxoethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide

To a solution of **22** (50 mg, 0.118 mmol) and 6-fluoro-2-naphthoic acid (27 mg, 0.142 mmol) in DCM (4 mL) was added DIC (22 µl, 0.142 mmol) at room temperature. The resulting mixture was stirred for 2 hrs then the solvent was removed under reduced pressure to give the crude product. Purification by flash chromatography on silica gel (eluting with Petroleum ether:EtOAc (1:1) then EtOAc) gave **23** (29 mg, 41%) as a white solid. **MS (ESI)** 597 (M+1); HPLC *t_R* 6.75 min.


Example 24 – Synthesis of Compound 24 N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-oxoethyl)-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide

**24**

N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-oxoethyl)-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide

To a solution of **23** (29 mg, 0.049 mmol) in dry THF (1 mL) was added LiAlH(OtBu)₃ (38 mg, 0.145 mmol) in one portion at room temperature and the resulting suspension was stirred overnight. This suspension was then slowly poured into a cold (0°C) 0.4 M KHSO₄ aqueous solution (2 mL, 0.8 mmol) and the resulting mixture was diluted with EtOAc (3 mL). The aqueous layer was extracted with EtOAc (3 x 3 mL) and the combined organic layers were washed with 1 N HCl (3 x 6 mL), saturated NaHCO₃ aqueous solution (1 x 6 mL), and brine (1 x 6 mL). The organic solution was then dried over MgSO₄, filtered and concentrated under reduced pressure to give the crude product **24** (24 mg, 91%), which was used in the next reaction without further purification. **MS (ESI)** 538 (M+1); HPLC *t_R* 6.41 min.

Example 25 – Synthesis of Compound 25 N-(((3S,5S)-3-(2-(diethylamino)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide

**25**


N-(((3S,5S)-3-(2-(diethylamino)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide

To a solution of **24** (24 mg, 0.044 mmol) in DCM (2 mL) was added diethylamine (55 µL, 0.532 mmol) at room temperature. After stirring for 5 mins, NaBH(OAc)₃ (20 mg, 0.090 mmol) was added to the above solution in one portion and the resulting suspension was stirred for another 10 mins. Saturated NaHCO₃ aqueous solution (4 mL) was added to the suspension and the aqueous layer was extracted with DCM (3 x 4 mL). The combined

organic layers were washed with brine (10 mL), dried over MgSO_4 , filtered and concentrated under reduced pressure to give the crude product. This crude product was purified by prep HPLC (100% H_2O to $\text{MeCN}/\text{H}_2\text{O}$ 9:1, gradient) to give the **25** as a white solid (TFA salt). **MS** (ESI) 595.3 ($\text{M}+1$); HPLC t_R 6.22 min.

5


Example 26 – Synthesis of Compound 26 benzyl 2-(methoxy(methyl)amino)-2-oxoethylcarbamate

**26**

benzyl 2-
(methoxy(methyl)amino)-2-
oxoethylcarbamate

To Cbz-glycine (10 g, 47.8 mmol, Aldrich) in DCM (100 mL) was added BOP reagent (21.5 g, 48.6 mmol) and DIPEA (6.5 mL, 46.0 mmol). After stirring at room temperature for 10 mins, N,O-dimethylhydroxylamine hydrochloride (4.9 g, 50.2 mmol) and DIPEA (6.5 mL, 46.0 mmol) were added. The reaction was stirred at room temperature overnight. The DCM was removed by rotary evaporation and the residue taken up in EtOAc (100 mL). The organic phase was washed with H_2O (3x 100 mL), saturated sodium bicarbonate solution (3x 100 mL), H_2O (3x 100 mL), 1M HCl (3x 100 mL), brine (3x 100 mL). The organic phase was dried (MgSO_4) and the EtOAc removed to give the Weinreb amide **26** as a white solid (7.78 g, 64%).

Example 27 – Synthesis of Compound 27 benzyl 2-oxobut-3-enylcarbamate

**27**


benzyl 2-oxobut-3-
enylcarbamate

20

To the Weinreb amide **26** (3.89 g, 15.42 mmol) in DCM (10mL) at 0°C was added vinyl magnesium bromide (45 mmol) in THF (45 mL). The reaction was stirred for 2 hrs and monitored by HPLC. The reaction was added to a mixture of ice and 1M HCl (200 mL). The aqueous mixture was extracted with DCM (3x 100 mL) and washed with 1M HCl (2x 200mL) and H_2O (3x 100mL). The organic phase was dried (MgSO_4) and the volume reduced to 100 mL by rotary evaporation. The α,β -unsaturated ketone **27** was stored and used in solution without purification.

25

Example 28 – Synthesis of Compound 28 (S)-9-fluorenylmethyl 10-(2,2-diphenylethyl)-2,2-dimethyl-18-phenyl-4,9,13,16-tetraoxo-3,17-dioxo-5,10,15-triazaoctadecan-8-ylcarbamate


**28**

(S)-9-fluorenylmethyl 10-(2,2-diphenylethyl)-2,2-dimethyl-18-phenyl-4,9,13,16-tetraoxo-3,17-dioxo-5,10,15-triazaoctadecan-8-ylcarbamate

5 To 2,2-diphenylethylamine (0.95 g, 7.4 mmol) in DCM (10 mL) was added the α,β -unsaturated ketone **27** (5.7 mmol) in DCM (75 mL). After stirring at room temperature for 15 mins, Fmoc-L-2,4-diaminobutyric acid(Boc)-OH (2.4 g, 8.55 mmol) and DIC (0.87 mL, 5.6 mmol) were added. The reaction was stirred at room temperature overnight. The DCM was removed by rotary evaporation and the residue was subjected to column chromatography on
10 silica gel using petroleum spirit:EtOAc (1:1 to 0:1) to give **28** (1.5 g, 31%)

Alternatively, to 2,2-diphenylethylamine (0.97 g, 7.4 mmol) in DCM (20 mL) was added the α,β -unsaturated ketone **27** (5.95 mmol) in DCM (40mL). After stirring at room temperature for 15 mins, Fmoc-L-2,4-diaminobutyric acid(Boc)-OH (2.4 g, 8.55 mmol), DIPEA
15 (2.5 mL) and HATU (2.3 g, 6.0 mmol) were added. The reaction was stirred at room temperature overnight. The DCM was removed by rotary evaporation and the residue was taken up in EtOAc (100 mL). The organic layer was washed with saturated sodium bicarbonate solution (2x 100 mL), saturated ammonium chloride solution (2x 100 mL) and brine (2x 100 mL). The organic phase was dried and the solvent removed under reduced
20 pressure. The residue was subjected to column chromatography on silica gel using petroleum spirit:EtOAc (3:1 to 1:1 to 0:1) to give **28** (0.86 g, 17%).


Example 29 – Synthesis of Compound 29 (3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-(benzyloxycarbonylaminomethyl)-1-(2,2-diphenylethyl)-1,4-diazepan-2-one

**29**

(3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-(benzyloxycarbonylaminomethyl)-1-(2,2-diphenylethyl)-1,4-diazepan-2-one

To Compound **28** (1.5 g, 1.8 mmol) in DCM (3 mL) was added diethylamine (1.5 mL, 14.5 mmol). The reaction was stirred at room temperature for 1hr. The DCM and diethylamine was removed by rotary evaporation. DCM (5 mL), sodium triacetoxyborohydride (0.4 g, 1.9 mmol) was added, and the reaction was stirred overnight at room temperature. The organic phase was washed with saturated sodium bicarbonate solution (25 mL), dried (MgSO₄) and the DCM removed to give the cyclised product **29**, which was used in the next step without purification.

Example 30 – Synthesis of Compound 30 (3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-aminomethyl-1-(2,2-diphenylethyl)-1,4-diazepan-2-one


**30**

(3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-aminomethyl-1-(2,2-diphenylethyl)-1,4-diazepan-2-one

To the cyclised product **29** in methanol (5 mL) was added catalytic Pd/C. The reaction was stirred under a hydrogen atmosphere overnight. The reaction mixture was filtered

through Celite™ and the methanol removed by rotary evaporation to give the amine **30** (0.7 g, 83% from **28**).


Example 31 – Synthesis of Compound 31 *N*-(((3*S*,5*S*)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide

**31**

N-(((3*S*,5*S*)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide

To the amine **30** (0.08 g, 0.17 mmol) in DCM (1 mL) was added DIPEA (0.25 mL), BOP reagent (0.08 g, 0.18 mmol) and 6-fluoro-2-naphthoic acid (0.06 g, 0.32 mmol). The reaction was stirred at room temperature for 2 hrs. TFA (1mL) was added and the reaction stirred at room temperature for 2 hrs. Rotary evaporation and preparative HPLC gave **31** (0.05 g, 54%). **MS (ESI)** 539.3 (*M*+1); HPLC *t_R* min 5.92


Example 32 – Synthesis of Compound 32 6-chloro-2-naphthoic acid

**32**

6-chloro-2-naphthoic acid

A suspension of 6-bromo-2-naphthoic acid (3.0 g, 11.47 mmol), CuCl (11.7 g, 114.64 mmol) and CuI (2.19 g, 11.50 mmol) in degassed DMF (45 mL) was heated to reflux under argon in dark for 4 hrs. After cooling to room temperature, the solution was decanted into H₂O (200 mL) and the resulting mixture was extracted with EtOAc (2 x 500 mL). The combined organic layers were then washed with H₂O (4 x 500 mL) followed by brine (1 x 500 mL), dried over MgSO₄, filtered and concentrated under reduced pressure to dryness. The residue was triturated with CH₃CN and the solid obtained was then re-crystallized from EtOAc to give the pure product **32** (2.2 g, 93%) as a off-white solid. HPLC *t_R* 6.47 min.


Example 33 – Synthesis of Compound 33 (*E*)-3-(4-chlorophenyl)-*N*-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide

**33**

(*E*)-3-(4-chlorophenyl)-*N*-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide

To the amine (*E*)-3-(4-chlorophenyl)-*N*-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-2-oxo-3-(2-aminoethyl)-1,4-diazepan-5-yl)methyl)acrylamide (21 mg, 0.05 mmol) in DMF (0.25 mL) was added K₂CO₃ (5 mg) and 1,5-dibromopropane (0.066 mL, 0.5 mmol). The reaction mixture was left at room temperature for 4 hrs. The solvent was removed under high vacuum, and the residue purified by preparative HPLC to give 8 mg (~30%) of **33** as the TFA salt. **MS** (ESI) 599.4 (M+1); HPLC *t_R* min 6.31

Example 34 – Synthesis of Compound 34 (*S*)-9-fluorenylmethyl 10-(2-phenylbutyl)-2,2-dimethyl-18-phenyl-4,9,13,16-tetraoxo-3,17-dioxa-5,10,15-triazaoctadecan-8-ylcarbamate


**34**

(*S*)-9-fluorenylmethyl 10-(2-phenylbutyl)-2,2-dimethyl-18-phenyl-4,9,13,16-tetraoxo-3,17-dioxa-5,10,15-triazaoctadecan-8-ylcarbamate

To 2-phenylbutylamine hydrochloride (0.26 g, 1.4 mmol) in DCM (10 mL) and DIPEA (0.25 mL, 1.8 mmol) was added the α,β -unsaturated ketone **27** (1.06 mmol) in DCM (20 mL). After stirring at room temperature for 15 mins, Fmoc-diaminobutyric acid(Boc)-OH (0.7 g, 1.56 mmol) and DIC (0.25 mL, 1.61 mmol) were added. The reaction was stirred at room

temperature overnight. The DCM was removed by rotary evaporation and the residue was subjected to column chromatography on silica gel using petroleum spirit:EtOAc (1:1 to 0:1), providing Compound **34** as a mixture of diastereomers (0.17 g, 21%).

5 **Example 35– Synthesis of Compound 35 (3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-(benzyloxycarbonylaminomethyl)-1-(2-phenylbutyl)-1,4-diazepan-2-one**



35


(3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-(benzyloxycarbonylaminomethyl)-1-(2-phenylbutyl)-1,4-diazepan-2-one

To Compound **34** (0.17 g, 0.2 mmol) in DCM (3 mL) was added diethylamine (1.5 mL). The reaction was stirred at room temperature for 1hr. The DCM and diethylamine was removed by rotary evaporation. DCM (5 mL) and sodium triacetoxyborohydride (0.1 g, 0.47 mmol) were added and the reaction was stirred overnight at room temperature. The organic phase was washed with saturated sodium bicarbonate solution (25 mL), dried (MgSO₄) and the DCM removed to give the cyclised product **35** as a mixture of diastereomers (0.11 g, 100%).

15

Example 36– Synthesis of Compound 36 (3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-(aminomethyl)-1-(2-phenylbutyl)-1,4-diazepan-2-one

133




36

(3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-(aminomethyl)-1-(2-phenylbutyl)-1,4-diazepan-2-one

To the cyclised product **35** (0.11 g) in methanol (5 mL) was added catalytic Pd/C. The reaction was stirred under a hydrogen atmosphere overnight. The reaction mixture was filtered through Celite and the methanol removed by rotary evaporation to give the amine **36** as a mixture of diastereomers (0.11 g, 100%).

Example 37– Synthesis of Compound **37** (3*S*,5*S*)-3-(2-aminoethyl)-5-(*N*-2-naphthamidomethyl)-1-(2-phenylbutyl)-1,4-diazepan-2-one



37


(3*S*,5*S*)-3-(2-aminoethyl)-5-(2-naphthamidomethyl)-1-(2-phenylbutyl)-1,4-diazepan-2-one

10

To the amine **36** (0.02 mg, 0.05 mmol) in DCM (1mL) was added DIPEA (0.1 mL, 0.7 mmol), BOP reagent (0.02 mg, 0.045 mmol) and 2-naphthoic acid (0.015 mg, 0.09 mmol). The reaction was stirred at room temperature for 2 hrs. TFA (1 mL) was added and the reaction stirred at room temperature for 2 hrs. Rotary evaporation and preparative HPLC gave **37** as a mixture of diastereomers (13.4 mg, 57%). **MS (ESI)** 473.4 ($M+1$); HPLC t_R 5.59 min

15

Example 38 – Synthesis of Compounds 38-39 *N*-(((3*S*,5*S*)-2-oxo-1-((*S*)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide and *N*-(((3*S*,5*S*)-2-oxo-1-((*R*)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide




5

Prepared from Compound **37** by alkylation as in Example 33. Preparative HPLC purification separated the two diastereomers. The correct configuration was assigned by resynthesis of the compounds using (*S*)-2-phenylbutylamine **43** or (*R*)-2-phenylbutylamine.
38: MS (ESI) 541.3 (M+1); HPLC t_R 5.78 min; 39: MS (ESI) 541.3 (M+1); HPLC t_R 5.67 min

10

Example 39 – Synthesis of Compound 40 (*S*)-2-phenylbutanol



15


20

To a suspension of sodium borohydride (2.36 g, 62.4 mmol) in THF (50 mL) was added a solution of (*S*)-2-phenylbutyric acid (4.27 g, 26.0 mmol) in THF (40 mL) slowly at 0°C. The mixture was stirred until the evolution of gas ceased. A solution of iodine (6.60 g, 26.0 mmol) in THF (40 mL) was then added slowly at 0 °C. After addition, the resulting mixture was allowed to warm to room temperature and stirred for 1 hr. The reaction solution was then slowly poured into a 1 N HCl solution (280 mL) and the resulting mixture was diluted with EtOAc (250 mL). The aqueous layer was extracted with EtOAc (150 mL x 3) and the combined organic layers were then washed with saturated NaHCO₃ (aq), 0.5 M Na₂S₂O₃ (aq) and brine. This organic solution was dried over MgSO₄, filtered and concentrated under reduced pressure to give the crude product. Purification by flash chromatography on silica

gel (Petroleum ether:EtOAc 4:1) gave the desired product **40** as a colorless oil in quantitative yield. HPLC t_R 5.24 min.

Example 40 – Synthesis of Compound 41 (S)-1-mesyloxy-2-phenylbutane


5



To a mixture of **40** (3.9 g, 26.0 mmol) and triethylamine (5.5 mL, 39.5 mmol) in DCM (90 mL) was added a solution of methanesulfonyl chloride (4.47 g, 39.0 mmol) in DCM (30 mL) slowly at 0°C. After addition, the resulting mixture was allowed to warm to room temperature and stirred for 2 hrs. 1 N HCl (70 mL) was then added to the above mixture and the aqueous layer was extracted with DCM (1 x 70 mL). The combined organic layers were washed with brine (150 mL), dried over MgSO₄, filtered and concentrated under reduced pressure to give the crude product **41** as a colorless oil. This crude product was used in the next step without further purification. HPLC t_R 6.48 min.

15


Example 41 – Synthesis of Compound 42 (S)-1-azido-2-phenylbutane



A suspension of **41** (5.93 g, 26.0 mmol) and sodium azide (5.7 g, 78.0 mmol) in DMF (60 mL) was heated at 85°C for 3 hrs. After cooling to room temperature, the mixture was diluted with H₂O (200 mL) and extracted with EtOAc (250 mL). The organic layer was then washed with H₂O (4 x 150 mL) followed by brine (150 mL), dried over MgSO₄, filtered and concentrated under reduced pressure to give the crude product. Purification by flash chromatography on silica gel (100% petroleum ether as the eluent) gave the pure product **42** (4.03 g, 88%) as a colorless oil. HPLC t_R 7.67 min.


25

Example 42– Synthesis of Compound 43 (S)-2-phenylbutylamine



A mixture of **42** (4.0 g, 22.8 mmol) and Lindlar's catalyst (1.5 g) in EtOAc (50 mL) was shaken at room temperature under H₂ (40 psi) over-night. The mixture was then filtered through a pad of Celite[™] and the filtrate was concentrated under reduced pressure to give the crude product **43** (3.4 g, 100%) as a light yellowish oil. This crude product was used for the conjugate addition reactions without further purification. MS (ESI) 150 (M+1); HPLC *t_R* 1.84 min.

Example 43 – Synthesis of Compound 44 allyl 2-(methoxy(methyl)amino)-2-oxoethylcarbamate



44

allyl 2-(methoxy(methyl)-
amino)-2-oxoethylcarbamate


10

To Alloc-glycine (1.45 g, 9.1 mmol) in DCM (20 mL) was added BOP reagent (3.3 g, 7.46 mmol) and DIPEA (1.5 mL, 10.7 mmol). After stirring at room temperature for 10 mins, N,O-dimethylhydroxylamine hydrochloride (0.8 g, 8.2 mmol) and DIPEA (1.5 mL, 10.7 mmol) were added. The reaction was stirred at room temperature overnight. The DCM was removed by rotary evaporation and the residue taken up in EtOAc (100 mL). The organic phase was washed with H₂O (3x 100 mL), saturated sodium bicarbonate solution (3x 50 mL), H₂O (3x 50 mL), 1M HCl (3x 50 mL), brine (3x 50 mL). The organic phase was dried (MgSO₄) and the EtOAc removed to give the Weinreb amide **44** as a white solid (0.43 g, 23%).

20


Alternatively, *tert*-butyl 2-(methoxy(methyl)amino)-2-oxoethylcarbamate **16** (Boc-Gly Weinreb amide, 1.4 g, 6.4 mmol) in DCM (5 mL) and TFA (3 mL) were stirred at room temperature 1 hr. The solvent was removed under reduced pressure, followed by addition of DCM (20 mL) and then DIPEA until basic. The solution was cooled to 0° C and allyl chloroformate added (1.4 mL, 13.2 mmol). The reaction was stirred at room temperature overnight. The reaction mixture was neutralised with 1M HCl and extracted with EtOAc. The EtOAc was removed by rotary evaporation and the residue was subjected to column chromatography on silica gel using petroleum spirit:EtOAc (1:1 to 0:1), providing **44** (0.86 g, 66%).

30

Example 44 – Synthesis of Compound 45 allyl 2-oxobut-3-enylcarbamate**45**

allyl 2-oxobut-3-enylcarbamate


To the Weinreb amide **44** (0.43 g, 2.1 mmol) in DCM (5 mL) at 0° C was added vinyl magnesium bromide (10 mmol) in THF (10 mL). The reaction was stirred for 2 hrs and monitored by HPLC. The reaction was added to a mixture of ice and 1M HCl (100 mL). The aqueous mixture was extracted with DCM (3x 50 mL) and washed with 1M HCl (2x 100 mL) and H₂O (3x 50 mL). The organic phase was dried (MgSO₄) and the volume reduced to 50 mL by rotary evaporation. The α,β -unsaturated ketone **45** was stored and used in solution without further purification.

Example 45 – Synthesis of Compound 46 (S)-9-fluorenylmethyl 10-(3,5-dichlorobenzyl)-2,2-dimethyl-4,9,13,16-tetraoxo-3,17-dioxo-5,10,15-triazaiscos-19-en-8-ylcarbamate**46**

(S)-9-fluorenylmethyl 10-(3,5-dichlorobenzyl)-2,2-dimethyl-4,9,13,16-tetraoxo-3,17-dioxo-5,10,15-triazaiscos-19-en-8-ylcarbamate

To 3,5-dichlorobenzylamine (0.49 g, 2.8 mmol) in DCM (5 mL) was added the α,β -unsaturated ketone **45** (2.12 mmol) in DCM (10 mL). After stirring at room temperature for 15 mins, Fmoc-diaminobutyric acid(Boc)-OH (1.35 g, 3.1 mmol) and DIC (0.5 mL, 3.2 mmol) was added. The reaction was stirred at room temperature overnight. The DCM was removed by rotary evaporation and the residue was subjected to column chromatography on silica gel using petroleum spirit:EtOAc (1:1 to 0:1), providing compound **46** (0.48 g, 22%).


Example 46 – Synthesis of Compound 47 (3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-(allyloxycarbonylaminomethyl)-1-(3,5-dichlorobenzyl)-1,4-diazepan-2-one

**47**

(3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-(allyloxycarbonylaminomethyl)-1-(3,5-dichlorobenzyl)-1,4-diazepan-2-one

To Compound **46** (0.48 g, 0.63 mmol) in DCM (3 mL) was added diethylamine (1.5 mL). The reaction was stirred at room temperature for 1hr. The DCM and diethylamine was removed by rotary evaporation. DCM (5 mL), sodium triacetoxyborohydride (0.2 g, 0.94 mmol) was added, and the reaction was stirred overnight at room temperature. The organic phase was washed with saturated sodium bicarbonate solution (25 mL), dried (MgSO₄) and the DCM removed to give the cyclised product **47** (0.24 g, 72%).

Example 47 – Synthesis of Compound 48 (3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-aminomethyl-1-(3,5-dichlorobenzyl)-1,4-diazepan-2-one


**48**

(3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-aminomethyl-1-(3,5-dichlorobenzyl)-1,4-diazepan-2-one

To the cyclised product **47** (0.24 g, 0.45 mmol) in DCM (3 mL) was added 1,3-dimethylbarbituric acid (13 mg, 0.08 mmol) and palladium tetrakis triphenylphosphine (5 mg). The reaction was evacuated and stirred at room temperature for 1 hr. The DCM was

removed under reduced pressure to give the crude product **48** (0.15 g, 75%) which was used in the next reaction without purification.


Example 48 – Synthesis of Compound 49 (3*S*,5*S*)-3-(2-aminoethyl)-5-(2-naphthoylaminoethyl)-1-(3,5-dichlorobenzyl)-1,4-diazepan-2-one

**49**

(3*S*,5*S*)-3-(2-aminoethyl)-5-(2-naphthoylaminoethyl)-1-(3,5-dichlorobenzyl)-1,4-diazepan-2-one

To the amine **48** (0.05 mg, 0.11 mmol) in DCM (1 mL) was added DIPEA (0.1 mL, 0.7 mmol), BOP reagent (0.05 mg, 0.11 mmol) and 2-naphthoic acid (0.04 mg, 0.23 mmol). The reaction was stirred at room temperature for 2 hrs. TFA (1 mL) was added and the reaction stirred at room temperature for 2 hrs. Rotary evaporation and preparative HPLC gave **49** (48 mg, 90%). **MS (ESI)** 499.3 (M+1); HPLC *t_R* 5.77 min

Example 49 – Synthesis of Compound 50 N-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthalene-2-sulfonamide


**50**

N-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)naphthalene-2-sulfonamide

Prepared from allyl 2-oxobut-3-enylcarbamate **45**, Boc-L-Arg(Fmoc)₂-OH and 2,2-diphenylethylamine according to the procedures of Examples 46-48, with the following

modification: the Boc group was removed with TFA during the deprotection/cyclization procedure of Example 47, rather than using diethylamine for Fmoc removal. Following Alloc deprotection by the procedure of Example 48, the free amine was dissolved in DCM to which was added naphthalene-2-sulfonyl chloride (10 mg) and DIPEA (20 μ L) and the reaction stirred for 2h at room temperature. Diethylamine (1 mL) was added and stirred overnight to remove the Fmoc protection, and the reaction evaporated to dryness. Preparative HPLC gave title compound **50** (13 mg). **MS (ESI)** 613.5 (M+1); HPLC t_R 5.89 min.


Example 50 – Synthesis of Compound 51 (S)-9-fluorenylmethyl 10-(2-ethylbutyl)-2,2-dimethyl-18-phenyl-4,9,13,16-tetraoxo-3,17-dioxo-5,10,15-triazaoctadecan-8-ylcarbamate

**51**

(S)-9-fluorenylmethyl 10-(2-ethylbutyl)-2,2-dimethyl-18-phenyl-4,9,13,16-tetraoxo-3,17-dioxo-5,10,15-triazaoctadecan-8-ylcarbamate

To 2-ethylbutylamine (0.15 g, 1.48 mmol) in DCM (10 mL) was added the α,β -unsaturated ketone **27** (1.47 mmol) in DCM (30 mL). After stirring at room temperature for 15 mins, Fmoc-diaminobutyric acid(Boc)-OH (0.95 g, 2.16 mmol) and DIC (0.34 mL, 2.19 mmol) were added. The reaction was stirred at room temperature overnight. The DCM was removed by rotary evaporation and the residue was subjected to column chromatography on silica gel using petroleum spirit:EtOAc (1:1 to 0:1), providing Compound **51** (0.5 g, 46%).

Example 51– Synthesis of Compound 52 (3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-(benzyloxycarbonylaminomethyl)-1-(2-ethylbutyl)-1,4-diazepan-2-one


**52**

(3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-(benzyloxycarbonylaminomethyl)-1-(2-ethylbutyl)-1,4-diazepan-2-one

To Compound **51** (0.5 g, 0.67 mmol) in DCM (3 mL) was added diethylamine (1.5 mL). The reaction was stirred at room temperature for 1hr. The DCM and diethylamine were removed by rotary evaporation. DCM (5 mL) and sodium triacetoxyborohydride (0.2 g, 0.94 mmol) were added and the reaction was stirred overnight at room temperature. The organic phase was washed with saturated sodium bicarbonate solution (25 mL), dried (MgSO₄) and the DCM removed to give the crude cyclised product **52** (0.4 g).

10

Example 52– Synthesis of Compound 53 (3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-(aminomethyl)-1-(2-ethylbutyl)-1,4-diazepan-2-one


**53**

(3*S*,5*S*)-3-(2-*tert*-butoxycarbonylaminoethyl)-5-(aminomethyl)-1-(2-ethylbutyl)-1,4-diazepan-2-one

To the cyclised product **52** (0.4 g) in methanol (5 mL) was added catalytic Pd/C. The reaction was stirred under a hydrogen atmosphere overnight. The reaction mixture was filtered through Celite and the methanol removed by rotary evaporation to give the amine **53** (0.17 g, 68% from **51**).

15


Example 53– Synthesis of Compound 54 N-(((3*S*,5*S*)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

**54**

N-(((3*S*,5*S*)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

To the amine **53** (0.030 g, 0.08 mmol) in DCM (1 mL) was added DIPEA (0.1 mL, 0.7 mmol), BOP reagent (0.03 g, 0.07 mmol) and 2-naphthoic acid (0.025 g, 0.14 mmol). The reaction was stirred at room temperature for 2 hrs. TFA (1 mL) was added and the reaction stirred at room temperature for 2 hrs. Rotary evaporation and preparative HPLC gave Compound **54** (23 mg, 68%). **MS (ESI)** 425.7 (M+1); HPLC *t_R* 5.27


Example 54 - Synthesis of Compound 55 (3*S*,5*S*)-3-[2-(piperidin-1-yl)ethyl]-5-(benzyloxycarbonylaminomethyl)-1-(2-ethylbutyl)-1,4-diazepan-2-one

**55**

(3*S*,5*S*)-3-[2-(piperidin-1-yl)ethyl]-5-(benzyloxycarbonylaminomethyl)-1-(2-ethylbutyl)-1,4-diazepan-2-one

To Compound **54** (0.25 g, 0.5 mmol) in DCM (3 mL) was added TFA (1.5 mL), with the solution stirred at room temperature for 1 hr. DCM (20 mL) was added and the solution was washed with saturated sodium bicarbonate solution (20 mL), dried over MgSO₄ and evaporated to give the crude amine (0.16 g). To this was added DMF (0.25 mL), potassium carbonate (10 mg) and 1,5-dibromopentane (0.35 mL, 2.5 mmol). The reaction mixture was stirred at room temperature for 1.5 hrs, after which DCM (20 mL) was added, the organic layer washed with saturated sodium bicarbonate solution (20 mL) and H₂O (20 mL), dried (MgSO₄) and evaporated to give crude **55**, which was used in the next reaction without purification.


Example 55– Synthesis of Compound 56 (3*S*,5*S*)-3-[2-(piperidin-1-yl)ethyl]-5-aminomethyl-1-(2-ethylbutyl)-1,4-diazepan-2-one

**56**

(3*S*,5*S*)-3-[2-(piperidin-1-yl)ethyl]-5-aminomethyl-1-(2-ethylbutyl)-1,4-diazepan-2-one

To the cyclised product **55** (0.4 g) in methanol (5 mL) was added catalytic Pd/C. The reaction was stirred under a hydrogen atmosphere overnight. The reaction mixture was filtered through Celite and the methanol removed by rotary evaporation to give the amine **56** (0.12 g).


Example 56 – Synthesis of Boc-L-Glu(piperidine)-OH **57** (*S*)-2-(tert-butoxycarbonylamino)-5-oxo-5-(piperidin-1-yl)pentanoic acid

**57**

(*S*)-2-(*tert*-butoxycarbonylamino)-5-oxo-5-(piperidin-1-yl)pentanoic acid

HATU (2.5 g) and DIPEA (1.5 mL) was added to Boc-L-Glu(OH)-OBn (2.0 g) in DCM (50 mL), stirred for 10 min, then piperidine (0.7 mL) was added and the reaction stirred overnight at room temperature. The reaction was washed with sodium bicarbonate solution (2x), saturated NH₄Cl (2x), brine (2x), dried over MgSO₄, filtered, and evaporated to give 2.9 g of Boc-L-Glu(piperidine)-OBn. The benzyl ester (0.6 g) was dissolved in EtOH (15 mL) with catalytic Pd/C and hydrogenated for 1 h, filtered over Celite, and the EtOH evaporated by rotary evaporation to give 0.51 g of **57**.


Example 57 – Synthesis of Compound 58 (3*S*,5*S*)-3-(2-*tert*-butoxycarbonylamino)propyl)-5-[benzyloxycarbonyl(methylamino)methyl]-1-(2,2-diphenylethyl)-1,4-diazepan-2-one

**58**

(3*S*,5*S*)-3-(2-*tert*-butoxycarbonylamino)propyl)-
5-[benzyloxycarbonyl(methylamino)methyl]-
1-(2,2-diphenylethyl)-1,4-diazepan-2-one

- 5 Prepared from Cbz-Sar, 2,2-diphenylethylamine and Fmoc-L-Orn(Boc) according to the procedures of Examples 27-30.

Example 58 – Synthesis of Compound 59 (3*S*,5*S*)-3-(2-*tert*-butoxycarbonylamino)propyl)-5-(methylamino)methyl-1-(2,2-diphenylethyl)-1,4-diazepan-2-one


**59**

(3*S*,5*S*)-3-(2-*tert*-butoxycarbonylamino)propyl)-
5-(methylamino)methyl-
1-(2,2-diphenylethyl)-1,4-diazepan-2-one

The cyclised product **58** (1.9 g) was dissolved in methanol (10 mL) with catalytic Pd/C and hydrogenated under a hydrogen atmosphere (40 psi) overnight. The reaction mixture was filtered through Celite[™] and the methanol removed by rotary evaporation to give the amine

15 **59** (1.86 g, 97%).


Example 59– Synthesis of Compound 60 N-(((3*S*,5*S*)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-bromo-*N*-methyl-2-naphthamide

**60**

N-(((3*S*,5*S*)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-bromo-*N*-methyl-2-naphthamide

The amine **59** was coupled with 6-bromo-2-naphthoic acid then deprotected with TFA according to Example 31. Rotary evaporation and preparative HPLC gave **60** (7.8 mg). **MS** (ESI) 629.4 (M+1); HPLC *t_R* 6.27 min.

Example 60– Synthesis of Compound 61 (3*S*,5*S*)-3-(*tert*-butoxycarbonylaminopropyl)-5-(6-bromo-2-naphthamidomethyl)-1-(2,2-diphenylethyl)-1,4-diazepan-2-one


**61**

(3*S*,5*S*)-3-(*tert*-butoxycarbonylaminopropyl)-5-(6-bromo-2-naphthamidomethyl)-1-(2,2-diphenylethyl)-1,4-diazepan-2-one

10

Prepared from 2,2-diphenylethylamine, Fmoc-L-Orn(Boc) and 6-bromo-2-naphthoic acid according to the procedures of Examples 28-31, without the TFA deprotection step of Example 31.

Example 61– Synthesis of Compound 62 *N*-(((3*S*,5*S*)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-4-methyl-2-oxo-1,4-diazepan-5-yl)methyl)-6-bromo-2-naphthamide


**62**

N-(((3*S*,5*S*)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-4-methyl-2-oxo-1,4-diazepan-5-yl)methyl)-6-bromo-2-naphthamide

Compound **61** (20.8 mg) was dissolved in DMF (1 mL) and treated with methyl iodide (6 μ L) at room temperature for 1 week. Additional methyl iodide (0.5 mL) and K_2CO_3 were added and the reaction left at room temperature for an additional 2 days. TFA (2 mL) was added and the reaction stirred at room temperature for 2h. Rotary evaporation followed by evaporation under high vacuum then preparative HPLC gave **62** (8.5 mg). **MS (ESI)** 629.3 (M+1); HPLC t_R 6.22 min.

10

Example 62 - Synthesis of Compound 63 *N*-(((3*S*,5*S*)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide


**63**

N-(((3*S*,5*S*)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

Obtained from **9**, 2,2-diphenylethylamine and Fmoc-L-Orn(Boc) according to Examples 10-12. The Boc group was removed under standard conditions to give the free amine. **MS(ESI)** 535 (M+1); HPLC t_R 5.78 min

15

Example 63 - Synthesis of Compound 64 *N*-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide


**64**

N-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

5 The amine **63** (0.79 g, 1.48 mmol), 1,5-dibromopentane (0.2 mL, 1.48 mmol) and K_2CO_3 (0.79 g) in DMF (11 mL) was stirred at room temperature for 4 h. The resulting mixture was diluted with ethylacetate (30 mL), washed with H_2O (5 x 30 mL), brine (10 mL) and dried over $MgSO_4$. Purification by preparative HPLC yielded **64** (0.23 g, 25%) **MS(ESI)** 603.3 ($M+1$); HPLC t_R 6.04 min

10

Example 64- Synthesis of Compound 65 *N*-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(3-(isopropylamino)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide


**65**

N-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(3-(isopropylamino)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

15 To a stirred mixture of the amine **63** (11 mg, 0.02 mmol), acetone (2 mL) and $MgSO_4$ (50 mg) in DCM (5 mL) was added sodium triacetoxyborohydride (8.5 mg, 0.04 mmol) at room temperature. Stirring continued for 2 h, the mixture was concentrated, re-dissolved in EtOAc (5 mL), washed with saturated aqueous $NaHCO_3$ solution (10 mL) and brine (10 mL),

dried over MgSO_4 and concentrated under reduced pressure. Purification by preparative HPLC yielded **65** (9.5 mg, 80%). **MS(ESI)** 577.2 ($\text{M}+1$); HPLC t_R 5.97 min.


Example 65 - Synthesis of Compound 66 tert-butyl (methylamino)(methylthio)methylenecarbamate

**66**

tert-butyl (methylamino)(methylthio)-methylenecarbamate

DIAD (2.7 mL, 13.8 mmol) was added to a stirred mixture of thiopseudourea (2.0 g, 6.9 mmol), PPh_3 (3.6g, 13.8 mmol), and MeOH (0.55 mL, 13.8 mmol) in dry THF (5 mL) at 0°C under nitrogen. Stirring continued at 0°C for 3 h then at room temperature for 16 h. The solvent was removed under reduced pressure and the resulting residue was re-dissolved in EtOAc, washed with saturated aqueous NaHCO_3 solution (20 mL) and brine (20 mL), and dried over MgSO_4 . Purification by silica gel chromatography using 20% EtOAc in petroleum ether as eluent gave **66** (1.63 g, 78%) as a colourless oil. **MS(ESI)** 305 ($\text{M}+1$); HPLC t_R 7.97 min.

Example 66 - Synthesis of Compound 67 N-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(3-(3-methylguanidino)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide


**67**

N-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(3-(3-methylguanidino)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

A mixture of Compound **63** (10 mg, 0.019 mmol), guanylylating agent **66** (56.9 mg, 0.19 mmol) and DIPEA (6.6 μL , 0.038 mmol) in DCM (5 mL) was stirred at room temperature for 16 h. TFA (5 mL) was added, and the resulting mixture was stirred at room temperature for 30

min. Solvent was removed under reduced pressure, and the crude product was purified by preparative HPLC to give **67** (0.53 mg, 4.7%) as a white solid. **MS(ESI)** 591.3 (M+1); HPLC t_R 5.94 min

5 **Example 67** - Synthesis of Compound **68** (S)-9-fluorenylmethyl 1-phenyl-10-(2,2-diphenylethyl)-18,18-dimethyl-4,9,13,16-tetraoxo-2,17-dioxa-4,10,15-triazanonadecan-8-ylcarbamate




68

(S)-9-fluorenylmethyl 1-phenyl-10-(2,2-diphenylethyl)-18,18-dimethyl-4,9,13,16-tetraoxo-2,17-dioxa-4,10,15-triazanonadecan-8-ylcarbamate

To a stirred mixture of Fmoc-L-Orn(Cbz)-OH (1.78 g, 3.65 mmol), DIPEA (0.64 mL, 3.65 mmol) and HATU (1.39 g, 3.65 mmol) in DCM (10 mL) was added a solution of amine **17** at room temperature. Stirring continued for 3 h, the reaction mixture was washed with saturated NaHCO₃ aqueous solution (20 mL) and brine (20 mL), and dried over MgSO₄. The solvent was removed under reduced pressure, with the crude **68** used in the next step without further purification. **MS(ESI)** 853 (M+1); HPLC t_R 9.90 min.

15 **Example 68** - Synthesis of Compound **69** (S)-(9H-fluoren-9-yl)methyl 7-((4-(tert-butoxycarbonylamino)-3-oxobutyl)(2,2-diphenylethyl)carbamoyl)-3-methyl-1,3-diazepane-1-carboxylate

150


**69**

(*S*)-(9*H*-fluoren-9-yl)methyl 7-((4-(*tert*-butoxycarbonylamino)-3-oxobutyl)(2,2-diphenylethyl)carbamoyl)-3-methyl-1,3-diazepane-1-carboxylate

A mixture of **68** (136 mg, 0.159 mmol) and Pd/C (20 mg) in ethanol (5 mL) was shaken under H₂ at 30 psi for 16 h, then filtered, concentrated under reduced pressure to give the crude amine (90 mg, 78%). The amine (90 mg, 0.125 mmol) was treated with excess formaldehyde solution in H₂O (37 mmol) in MeOH (5 mL) followed by sodium triacetoxyborohydride (23.5 mg, 0.375 mmol). After 1 h, the reaction mixture was washed with saturated NaHCO₃ aqueous solution (10 mL) and brine (10 mL), dried over MgSO₄, filtered and concentrated. The crude material was used in the next step without further purification. **MS(ESI)** 745 (M+1); HPLC ; HPLC *t*_R 7.83 min.

10


Example 69 - Synthesis of Compound 70 (*S*)-(9*H*-fluoren-9-yl)methyl 7-((4-(2-naphthamido)-3-oxobutyl)(2,2-diphenylethyl)-carbamoyl)-3-methyl-1,3-diazepane-1-carboxylate

**70**

(*S*)-(9*H*-fluoren-9-yl)methyl 7-((4-(2-naphthamido)-3-oxobutyl)(2,2-diphenylethyl)carbamoyl)-3-methyl-1,3-diazepane-1-carboxylate


Compound **69** (8 mg, 0.011 mmol) was treated with 1:1 v/v trifluoacetic acid/DCM mixture (2 mL) for 30 min at room temperature. The mixture was concentrated under reduced pressure, re-dissolved in DCM (5 mL), washed with saturated NaHCO₃ aqueous solution (5 mL) and brine (5 mL), dried over MgSO₄ and filtered. The filtrate was then treated with a mixture of 2-naphthoic acid (1.8 mg, 0.011 mmol), DIPEA (5.7 µL, 0.033 mmol) and BOP (4.8 mg, 0.011 mmol) in DCM (1 mL) with stirring at room temperature. After 3 h, the reaction mixture was washed with saturated NaHCO₃ aqueous solution (10 mL) and brine (10 mL), dried over MgSO₄, filtered and concentrated. The crude material was used in the next step without further purification. **MS(ESI)** 799 (M+1); HPLC *t_R* 7.90 min.

Example 70 - Synthesis of Compound 71 N-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(3-(methylamino)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

**71**


N-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(3-(methylamino)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

To a stirred solution of **70** (0.011 mmol) in DCM (5 mL) was added diethylamine (5 mL) at room temperature. The reaction was stirred for 1 h, then concentrated under reduced pressure. The residue was re-dissolved in DCM (5 mL) followed by addition sodium triacetoxyborohydride (5 mg, 0.08 mmol) at room temperature. Stirring continued for 1 h, with the resulting mixture washed with saturated NaHCO₃ aqueous solution (10 mL) and brine (10 mL), dried over MgSO₄, filtered and concentrated. Purification by preparative HPLC yielded **71** (0.21 mg) as a white solid. **MS(ESI)** 549.3 (M+1); HPLC *t_R* 5.93 min

Example 71 - Synthesis of Compound 72 ((S)-2-(allyloxycarbonylamino)-3-(naphthalen-2-yl)propanoic acid


((S)-2-(allyloxycarbonylamino)-3-(naphthalen-2-yl)propanoic acid

To a stirred mixture of L-3-(2-naphthyl)alanine hydrochloride (5.0 g, 19.8 mmol),
 5 Na₂CO₃ (7.3 g, 69.3 mmol) and 1,4-dioxane (30 mL) in H₂O (50 mL) was added allylchloroformate (2.1 mL, 19.8 mmol) at 0°C. The resulting mixture was stirred for 16 h then concentrated under reduced pressure. The residue was diluted with ethylacetate (50 mL), and at 0°C acidified to pH 2. The aqueous phase was extracted with ethylacetate (3 x 20 mL), the combined organic phase was washed with H₂O (50 mL) and brine (20 mL), dried over
 10 MgSO₄, filtered and concentrated under reduced pressure to give **72** as a colourless oil (5.8 g, 97%), which was used in the next step without further purification. HPLC *t_R* 6.60 min.

Example 72 - Synthesis of Compound 73 (S)-allyl 1-(methoxy(methyl)amino)-3-(naphthalen-2-yl)-1-oxopropan-2-ylcarbamate


(S)-allyl 1-(methoxy(methyl)amino)-3-(naphthalen-2-yl)-1-oxopropan-2-ylcarbamate

To a stirred mixture of the acid **72** (5.84 g, 19.5 mmol), DIPEA (3.7 mL, 2.09 mmol) and BOP (8.63 g, 19.5 mmol) in DCM (10 mL) was added a pre-mixed solution of *N,O*-dimethylhydroxylamine hydrochloride (1.9 g, 19.5 mmol) and DIPEA (7.3 mL, 41.6 mmol) in DCM (10 mL) at room temperature. Stirring continued for 16 h the reaction mixture was
 20 washed with 1N HCl (3 x 60 mL), H₂O (3 x 60 mL), saturated NaHCO₃ aqueous solution (3 x 60 mL) and brine (60 mL), dried over MgSO₄. Purification by silica gel chromatography using 20% ethylacetate in petroleum ether as eluent gave **73** (4.83 g, 71%) as a colourless oil. **MS (ESI)** 343 (M+1); HPLC *t_R* 7.07 min.

Example 73 - Synthesis of Compound 74 (S)-allyl 1-(naphthalen-2-yl)-3-oxopent-4-en-2-ylcarbamate**74**

(S)-allyl 1-(naphthalen-2-yl)-3-oxopent-4-en-2-ylcarbamate


At 0° C a solution of vinylmagnesium bromide in THF (11.5 mL, 1 M) was added in one portion to Weinreb amide **73** (1.58 g, 4.62 mmol) under nitrogen with stirring. The resulting mixture was allowed to stir for 2 h, and poured into a 1N HCl/ice mixture (50 mL). The aqueous mixture was extracted with DCM (3x 20 mL), the combined DCM extract was washed with 1N HCl (50 mL), saturated NaHCO₃ aqueous solution (50 mL) and brine (20 mL), dried over MgSO₄. Solvent was removed under reduced pressure producing the product **74** (1.14 g, 80%), which was used in the next step without further purification. **MS(ESI)** 310 (M+1); HPLC *t_R* 7.51 min.

Example 74 - Synthesis of Compound 75 (S)-allyl 5-(2,2-diphenylethylamino)-1-(naphthalen-2-yl)-3-oxopentan-2-ylcarbamate**75**

(S)-allyl 5-(2,2-diphenylethylamino)-1-(naphthalen-2-yl)-3-oxopentan-2-ylcarbamate

To a stirred solution of 2,2-diphenylethylamine (0.45 g, 2.3 mmol) in DCM (55 mL) was added the vinyl ketone **74** (0.71 g, 2.3 mmol) in one portion. Stirring continued for 2 h, with the reaction mixture used in the next step without purification. **MS(ESI)** 507 (M+1); HPLC *t_R* 7.22 min.


Example 75 - Synthesis of Compound 76 (S)-allyl 5-(N-(Boc-L-Arg(Cbz)₂) 2,2-diphenylethylamino)-1-(naphthalen-2-yl)-3-oxopentan-2-ylcarbamate

**76**

(S)-allyl 5-(N-(Boc-L-Arg(Cbz)₂) 2,2-diphenylethylamino)-1-(naphthalen-2-yl)-3-oxopentan-2-ylcarbamate

To a stirred solution of the amine adduct **75** (2.3 mmol) was added a mixture of Boc-Arg(Cbz)₂-OH (1.25 g, 2.3 mmol), DIPEA (0.8 mL, 4.6 mmol) and HATU (0.87 g, 2.3 mmol) in DCM (15 mL) at room temperature. Stirring continued for 16 h, after which the reaction mixture was washed with saturated NaHCO₃ aqueous solution (3x 20 mL) and brine (10 mL) then dried over MgSO₄. Purification by silica gel chromatography using 20% ethylacetate in petroleum ether as eluent gave **76** as a colourless oil (708 mg, 30% over 3 steps). **MS(ESI)** 1031 (M+1); HPLC *t_R* 10.80 min.


Example 76 - Synthesis of Compound 77 allyl (S)-1-((3S,5RS)-1-(2,2-diphenylethyl)-3-(bis Cbz 3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethylcarbamate

**77**

allyl (*S*)-1-((3*S*,5*RS*)-1-(2,2-diphenylethyl)-3-(bis Cbz 3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethylcarbamate

To a stirred solution of acyclic intermediate **76** (0.48 g, 0.47 mmol) in DCM (5 mL) was added TFA (5 mL) at room temperature. Stirring continued for 30 min, after which the mixture was diluted with DCM (20 mL) then washed with saturated NaHCO₃ aqueous solution (3 x 20 mL) and brine (10 mL), and dried over MgSO₄. To the resulting solution was added sodium triacetoxyborohydride (0.2 g, 0.94 mmol) with stirring at room temperature, after 30 min the mixture was washed with saturated NaHCO₃ aqueous solution (3 x 20 mL) and brine (10 mL), then dried over MgSO₄. The crude **77**, a mixture of diastereomers at the diazepan-2-one C5, was used in the next step without further purification. **MS(ESI)** 915 (M+1)

Example 77 - Synthesis of Compound 78 bis (Cbz) 1-(3-((2*S*,7*RS*)-7-((*S*)-1-amino-2-(naphthalen-2-yl)ethyl)-4-(2,2-diphenylethyl)-3-oxo-1,4-diazepan-2-yl)propyl)guanidine


**78**

bis (Cbz) 1-(3-((2*S*,7*RS*)-7-((*S*)-1-amino-2-(naphthalen-2-yl)ethyl)-4-(2,2-diphenylethyl)-3-oxo-1,4-diazepan-2-yl)propyl)guanidine


A mixture of compound **77** (36 mg, 0.039 mmol), 1,3-dimethylbarbituric acid (7.4 mg, 0.047 mmol) and Pd(PPh₃)₄ in DCM (5 mL) was stirred at room temperature under vacuum for 4 h. The resulting mixture was used in the next step without further purification. **MS(ESI)** 832 (M+1)

5

Example 78 - Synthesis of Compounds 79 and 80 *N*-((*S*)-1-((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)acetamide and *N*-((*S*)-1-((3*S*,5*R*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)acetamide

**79**

N-((*S*)-1-((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)acetamide

**80**

N-((*S*)-1-((3*S*,5*R*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)acetamide

10


A solution of the amine **78** (0.09 mmol) in DCM (5 mL) was treated with acetic anhydride (8.6 μL, 0.09 mmol) with stirring at room temperature. After 3 h the mixture was concentrated, re-dissolved in EtOAc, washed with saturated NaHCO₃ aqueous solution (10 mL) and brine (10 mL), dried over MgSO₄, then concentrated under reduced pressure. The residue was dissolved in MeOH (10 mL), Pd/C (5 mg) was added, and the solution shaken under H₂ at 20 psi for 16 h. The reaction was filtered, concentrated and purified by preparative HPLC to give the preferred diastereomer **79** (3 mg) and the less preferred diastereomer **80** (6 mg) as white solids.

79: **MS(ESI)** 606.4 (M+1); HPLC *t_R* 6.033 min


80: **MS(ESI)** 606.3 (M+1); HPLC *t_R* 6.046 min

20

Example 79 - Synthesis of Compounds 81 and 82 (S)-2-acetamido-N-((S)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)-3-(1H-imidazol-5-yl)propanamide and (S)-2-acetamido-N-((S)-1-((3S,5R)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)-3-(1H-imidazol-5-yl)propanamide

**81**

(S)-2-acetamido-N-((S)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)-3-(1H-imidazol-5-yl)propanamide

**82**


(S)-2-acetamido-N-((S)-1-((3S,5R)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)-3-(1H-imidazol-5-yl)propanamide

To a stirred mixture of Ac-L-His-OH (33.6 mg, 0.156 mmol), DIPEA (112.5 μ L, 0.312 mmol) and BOP (68.8 mg, 0.156 mmol) in DMF (1 mL) was added the amine **78** (0.039 mmol) at room temperature. Stirring continued for 16 h, then the reaction mixture was diluted with DCM/H₂O mixture (10 mL, 1:1 v/v), and the aqueous phase was extracted with DCM (3 x 5 mL). The combined DCM extracts were washed with saturated NaHCO₃ aqueous solution (3 x 20 mL) and brine (10 mL), dried over MgSO₄, and concentrated under reduced pressure. The residue was re-dissolved in MeOH (5 mL), and Pd/C (20 mg) was added. The resulting mixture was shaken under H₂ at 30 psi for 16 h, then was filtered, concentrated and purified by preparative HPLC to give the preferred diastereomer **81** (1.9 mg) and the less preferred diastereomer **82** (0.9 mg) as white solids.

81: MS(ESI) 743.4 (M+1); HPLC *t_R* 5.489 min

82: MS(ESI) 743.4 (M+1); HPLC *t_R* 5.555 min

Example 80 - Synthesis of Compounds **83** and **84** propyl (*S*)-1-((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethylcarbamate and propyl (*S*)-1-((3*S*,5*R*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethylcarbamate

**83**

propyl (*S*)-1-((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethylcarbamate

84


propyl (*S*)-1-((3*S*,5*R*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethylcarbamate

A mixture of **77** (36 mg, 0.039 mmol) and Pd/C (5 mg) in MeOH (5 mL) was shaken under H₂ at 20 psi for 16 h, then was filtered, concentrated and purified by preparative HPLC to give the preferred diastereomer **83** (0.07 mg) and the less preferred diastereomer **84** (2.7 mg) as white solids.

83: MS(ESI) 650.3 (M+1); HPLC *t_R* 6.52 min

84: MS(ESI) 650.2 (M+1); HPLC *t_R* 6.64 min

Example 81 - Synthesis of Compounds **85-87** 1-(3-((2*S*,7*S*)-7-(*N*-R1 (*R*)-1-amino-2-(naphthalen-2-yl)ethyl)-4-(2,2-diphenylethyl)-3-oxo-1,4-diazepan-2-yl)propyl)guanidine

**85-87**


1-(3-((2*S*,7*S*)-7-(*N*-R1 (*R*)-1-amino-2-(naphthalen-2-yl)ethyl)-4-(2,2-diphenylethyl)-3-oxo-1,4-diazepan-2-yl)propyl)guanidine

Compounds **85-87** were prepared in the same fashion as the preferred diastereomers Compounds **79**, **81** and **83** using the procedures described in Examples 72-81 with D-(2-naphthyl)alanine hydrochloride as the starting material.

Compound	R ₁ group	MS (M+1)	t _R (min)
85	Ac	606.2	6.01
86	Ac-His	743.5	5.41
87	Propyloxycarbonyl	650.4	6.42

5


Examples 82-90: Synthesis via Scheme 2: Preparation of all Four Diastereomers of N-((1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide **88**

**88**

N-((1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

10

Example 82– Synthesis of Compound 89 2,2-dimethyl-10-(2,2-diphenylethyl)-4,7,11-trioxo-3,12-dioxo-5,10-diazapentadec-14-ene

**89**


2,2-dimethyl-10-(2,2-diphenylethyl)-4,7,11-trioxo-3,12-dioxo-5,10-diazapentadec-14-ene

2,2-diphenylethylamine (3g) was added to Boc-vinylketone **16** (2.8g) as in Example 17. To the crude adduct **17** was added Alloc-Cl (1.6 mL) and the reaction stirred

15

until TLC indicated consumption of the secondary amine. THE solvent was evaporated and the residue purified by column chromatography (SiO₂ gel, pet. ether/EtOAc) to give 3.2 g (57%) of **89**.

5 **Example 83– Synthesis of Compound 90 (S)-allyl 2-amino-5-(benzyloxycarbonylamino)pentanoate L-H-Orn(Cbz)-Oallyl**


**90**

(S)-allyl 2-amino-5-(benzyloxycarbonylamino)pentanoate

H-L-Orn(Cbz)-OH (6.66g, 25 mmol), allyl alcohol (17.56 mL, 25 mmol) and p-TsOH (5.7g, 30 mmol) were dissolved in benzene (200 mL) and refluxed under Dean-Stark conditions for 5h. The majority of the solvent was then distilled off, with the remainder removed under vacuum. The resulting solid was recrystallized from DCM, filtered and dried to give 11.19g (94%) of the tosylate salt. To obtain the free amine the solid was dissolved in DCM, washed with sat. NaHCO₃, the aqueous layer washed with DCM (3x), and the organic layers dried over MgSO₄ and evaporated to dryness.

15

Example 84– Synthesis of Compound 91 (R)-allyl 2-amino-5-(benzyloxycarbonylamino)pentanoate D-H-Orn(Cbz)-Oallyl


**91**

(R)-allyl 2-amino-5-(benzyloxycarbonylamino)pentanoate

H-D-Orn(Cbz)-OH (6.66g, 25 mmol) was converted into 10.93 g (91%) of the tosylate salt of **91** as in Example 83, then converted into the free amine.

20

Example 85– Synthesis of Compound 92 (2*R*)-allyl 5-(benzyloxycarbonylamino)-2-(10-(2,2-diphenylethyl)-2,2-dimethyl-4,11-dioxo-3,12-dioxo-5,10-diazapentadec-14-en-7-ylamino)pentanoate




92

(2*R*)-allyl 5-(benzyloxycarbonylamino)-2-(10-(2,2-diphenylethyl)-2,2-dimethyl-4,11-dioxo-3,12-dioxo-5,10-diazapentadec-14-en-7-ylamino)pentanoate

- 5 The protected aminoketone **89** (746 mg, 1.6 mmol), D-Orn(Cbz)-Oallyl **91** (538 mg, 1.76 mmol) and NaBH(OAc)₃ (678 mg, 3.2 mmol) in a minimum volume of DCM were stirred for 24 h. A drop of AcOH was added just before workup, at which point saturated NaHCO₃ was added, extracted with DCM (3x), and the organic extracts combined and washed with saturated NaHCO₃ and H₂O, dried over MgSO₄, and evaporated to dryness. The product was
- 10 purified by column chromatography (SiO₂ gel, pet. ether/EtOAc) to give 890 mg (74%) of **92** as a mixture of diastereoisomers.


Example 86– Synthesis of Compound 93 (2S)-allyl 5-(benzyloxycarbonylamino)-2-(10-(2,2-diphenylethyl)-2,2-dimethyl-4,11-dioxo-3,12-dioxo-5,10-diazapentadec-14-en-7-ylamino)pentanoate

**93**

(2*S*)-allyl 5-(benzyloxycarbonylamino)-2-(10-(2,2-diphenylethyl)-2,2-dimethyl-4,11-dioxo-3,12-dioxo-5,10-diazapentadec-14-en-7-ylamino)pentanoate

- 5 L-Orn(Cbz)-Oallyl **90** (592 mg, 1.93 mmol) was converted into a mixture of the set of diastereomers **93** (925 mg, 76%) following the procedures of Example 86.

Example 87– Synthesis of Compounds 94 and 95 (3*R*,5*S*)-5-(*N*-Boc aminomethyl)-3-(*N*-Cbz 3-aminopropyl)-1-(2,2-diphenylethyl)-1,4-diazepan-2-one and (3*R*,5*R*)-5-(*N*-Boc aminomethyl)-3-(*N*-Cbz 3-aminopropyl)-1-(2,2-diphenylethyl)-1,4-diazepan-2-one

**94**


(3*R*,5*S*)-5-(*N*-Boc aminomethyl)-3-(*N*-Cbz 3-aminopropyl)-1-(2,2-diphenylethyl)-1,4-diazepan-2-one

95

(3*R*,5*R*)-5-(*N*-Boc aminomethyl)-3-(*N*-Cbz 3-aminopropyl)-1-(2,2-diphenylethyl)-1,4-diazepan-2-one

The Alloc/allyl protected derivative **92** (840 mg, 1.11 mmol) was dissolved in a minimum of DCM. 1,3-Dimethylbarbituric acid (346 mg, 2.22 mmol) and catalytic Pd(PPh₃)₄ were added, and the reaction degassed under vacuum, sealed and stirred overnight. The reaction was diluted to 50 mL with DCM, DIPEA (430 mg, 3.33 mmol) and BOP (540 mg, 1.22 mmol) were added, and the reaction stirred for 30 min. The DCM was removed under vacuum and the residue taken up in EtOAc, washed (saturated NaHCO₃, H₂O, saturated NaCl), dried (MgSO₄) and evaporated to dryness (TLC: EtOAc, 2 spots, R_f 0.33 and 0.57). The two diastereomeric products were separated by column chromatography (SiO₂ gel, pet. ether/EtOAc) to give 362 mg of the earlier eluting (3R,5S) isomer **94**, and 342 mg of the later eluting (3R,5R) isomer **95**.

Example 88– Synthesis of Compounds 96 and 97 (3S,5R)-5-(N-Boc aminomethyl)-3-(N-Cbz 3-aminopropyl)-1-(2,2-diphenylethyl)-1,4-diazepan-2-one and (3S,5S)-5-(N-Boc aminomethyl)-3-(N-Cbz 3-aminopropyl)-1-(2,2-diphenylethyl)-1,4-diazepan-2-one

**96**


(3R,5S)-5-(N-Boc aminomethyl)-3-(N-Cbz 3-aminopropyl)-1-(2,2-diphenylethyl)-1,4-diazepan-2-one

97

(3R,5R)-5-(N-Boc aminomethyl)-3-(N-Cbz 3-aminopropyl)-1-(2,2-diphenylethyl)-1,4-diazepan-2-one

The (3S,5R) (312 mg) and (3S,5S) (331 mg) isomers were obtained from the L-Orn-derived acyclic material **93** (870 mg) following the procedure of Example 87.

Example 89– Synthesis of Compounds 98-101 5-(N-Boc aminomethyl)-3-(N,N'-Cbz₂ 3-guanidinopropyl)-1-(2,2-diphenylethyl)-1,4-diazepan-2-one



98-101

The Orn Cbz group of **94** was removed by hydrogenation (H₂, 30 psi) over catalytic Pd/C in methanol overnight. The solution was filtered through Celite and evaporated to give a solid. A solution of the resulting amine (187 mg, 0.39 mmol) in DCM was mixed with a solution of the guanylation reagent CbzNHC(=NCbz)NHTf (196 mg, 0.43 mmol) in DCM. TEA (43 mg, 0.43 mmol) was added, and the reaction stirred overnight. The solution was diluted with DCM, washed (KHSO₄, sat. NaHCO₃, brine), dried (MgSO₄) and evaporated to dryness, then purified by flash chromatography over SiO₂ using hexanes/EtOAc as eluent, to give (3R,5S) **98** (182 mg, 59%). The other isomers **95-97** were converted in a similar manner to give:

99 (3R,5R): 171mg (68%) from 148 mg of amine

100 (3S,5S): 80 mg (65%) from 72 mg of amine

101 (3S,5R): 142 mg (58%) from 144 mg of amine


Example 90– Synthesis of Compounds 102-105

102 N-(((3R,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide


103 N-(((3R,5R)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

104 N-(((3S,5R)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide


105 N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

**102**


N-(((3*R*,5*S*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

**103**

N-(((3*R*,5*R*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

**104**

N-(((3*S*,5*R*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

**105**

N-(((3*S*,5*S*)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

The Boc derivative **99** (180 mg) in DCM (1 mL) was treated with TFA (1 mL) for 20 mL. The solvent was removed by evaporation, a solution of NaHCO₃ was added, and extracted 3x with DCM. The dichloromethane solution was dried over MgSO₄, filtered and evaporated to dryness. A portion (56 mg, 0.086 mmol) of the crude deprotected amine in DCM was stirred with 2-naphthoic acid (16 mg), DIPEA (60 μ L) and BOP (42 mg) for 30 min. MeOH was added and the reaction stirred overnight. The reaction was filtered, then purified by flash chromatography over SiO₂ using petroleum ether/EtOAc as eluent, to give the (3*R*,5*R*) isomer (43 mg, 94%). The other isomers were converted in a similar manner to give: (3*R*,5*S*): 41 mg (85%) from 60 mg **98**, (3*S*,5*R*): 27 mg (70%) from 40 mg **101**, and (3*S*,5*S*): 13 mg (74%) from 20 mg **100**.

Each compound was dissolved in dioxane:MeOH and hydrogenated over catalytic Pd/C under 30 psi H₂ overnight. The solution was filtered through Celite and evaporated to

give a solid. **102** (3R,5S): 27mg (96%) from 41 mg, **103** (3R,5R): 25mg (85%) from 43 mg, **104** (3S,5R): 11 mg (quantitative) from 13 mg, and **105** (3S,5S): 3 mg (73%) from 6 mg.

Compound	stereochemistry	MS (M+1)	t _R (min)
102	(3R,5S)	577.4	5.775
103	(3R,5R)	577.5	5.750
104	(3S,5R)	577.5	5.783
105	(3S,5S)	577.3	5.787

5 **Example 91 – Synthesis of Compounds 425,565, 580-585**

425 6-chloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

565 6-chloro-N-(((3S,5S)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

10 **580** 6-chloro-N-(((3R,5R)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide


581 6-chloro-N-(((3S,5R)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

15 **582** 6-chloro-N-(((3R,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

583 6-chloro-N-(((3S,5R)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide


584 6-chloro-N-(((3R,5S)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

20 **585** 6-chloro-N-(((3R,5R)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide



Compounds **425**, **565**, and **580-585** were prepared following similar procedures as used to prepare Compounds 102-104 (Scheme 2 route). In addition, compounds **425**, **565**, **580** and **585** were prepared according to the Scheme 1 route; a detailed procedure for the preparation of Compound **425** is contained in Examples 92-99.


Compound	stereochemistry	MS (M+1)	<i>t_R</i> (min)
425	(3S,5S,2'S)	575.3	6.269
565	(3S,5S,2'R)	574.8	6.265
580	(3R,5R,2'R)	575.4	6.404
581	(3S,5R,2'S)	575.2	6.262
582	(3R,5S,2'S)	575.2	6.110
583	(3S,5R,2'R)	575.1	6.211
584	(3R,5S,2'R)	575.2	6.253
585	(3R,5R,2'S)	575.4	6.274

Example 92– Synthesis of Compound 586 (*S*)-N-(2-oxo-4-(2-phenylbutylamino)butyl)-3-phenylpropanamide

To a solution of (*S*)-phenylbutylamine (8.5 g, 57.07 mmol) in DCM (100 ml) was added a solution of α,β -unsaturated ketone **27** (12.5 g, 57.1 mmol) in DCM (100 ml) at room temperature in one portion. The resulting mixture was stirred until all of the α,β -unsaturated ketone had been consumed (within one hour), then the conjugate addition adduct **586** was used straight in the next reaction.

HPLC t_R 5.71 min

MS (ESI) 369.3 (M+1)

Example 93– Synthesis of Compound 587 (*S*)-9-fluorenylmethyl 10-[(*S*)-2-phenylbutyl]-2,2-dimethyl-18-phenyl-4,9,13,16-tetraoxo-3,17-dioxa-5,10,15-triazaoctadecan-8-ylcarbamate


To the freshly prepared amine **586** in DCM (200 mL) was added Fmoc-L-Dab(Boc)OH (32.7 g, 74.2 mmol) followed by DIPC (11.5 g, 74.2 mmol) at room temperature. The resulting mixture was stirred for 2 hours, the by-product diisopropylurea was removed by filtration through a pad of Celite[®], and the filtrate was concentrated under reduced pressure to give the crude product, which was purified by silica gel column chromatography using 30-70% EtOAc/Pet. Spirit as eluent to give **587** (19.9 g, 44% yield over two steps).

TLC r_f 0.23 (50% EtOAc/Pet. Spirit)

HPLC t_R 10.03 min

MS (ESI) 791.2 (M+1)

Example 94– Synthesis of Compound 588 (S)-10-[(S)-2-phenylbutyl]-2,2-dimethyl-8-amino-18-phenyl-4,9,13,16-tetraoxo-3,17-dioxa-5,10,15-triazaoctadecane




5

Diethylamine (30 mL) was added to a solution of the acylated amine **587** (19.9 g, 25.19 mmol) in DCM (30 mL) at room temperature and the resulting mixture was stirred for 30 min. The solvent and diethylamine were removed under reduced pressure to give the desired product **588**. It was used in the next step without further purification.

10 HPLC t_R 6.85 min

MS (ESI) 569.3 (M+1)

Example 95– Synthesis of Compound 589 (3S,5S)-3-(2-tert-butoxycarbonylaminoethyl)-5-(benzyloxycarbonylaminomethyl)-1-[(S)-2-phenylbutyl]-1,4-diazepan-2-one



15

To a solution of crude Fmoc deprotected material **588** in DCM (50 ml) was added AcOH (15 mL) followed by NaBH(OAc)₃ (5.34 g, 25.2 mmol) in one portion at room temperature. The resulting mixture was stirred for 30 min, then washed with saturated NaHCO₃(aq) solution (80 mL x 3), brine (80 mL) and dried over MgSO₄. Filtration and concentration of the organic phase under reduced pressure gave the crude product, which was purified by silica gel column chromatography using 50-100% EtOAc/Pet. Spirit followed by 20% MeCN/EtOAc to give the product **589** (12.3 g, 88% over two steps).


20

TLC r_f 0.19 (70% EtOAc/Pet. Spirit)

HPLC t_R 7.06 min

MS (ESI) 553.3 (M+1)

5 **Example 96**– Synthesis of Compound 590 *tert*-butyl 2-((2*S*,7*S*)-7-aminomethyl-3-oxo-4-[(*S*)-2-phenylbutyl]-1,4-diazepan-2-yl)ethylcarbamate




10 A mixture of Cbz-protected product **589** (12.3 g, 22.3 mmol) and 5% Pd/C (2 g) in MeOH (100 mL) was shaken at room temperature under hydrogen at atmospheric pressure for one hour. The mixture was then filtered through a pad of Celite[®], and the filtrate was concentrated under reduced pressure to give the crude amine **590**. The crude material was used in the next step without further purification.

HPLC t_R 5.77 min

15 MS (ESI) 419.3 (M+1)

Example 97– Synthesis of Compound 591 *tert*-butyl 2-((2*S*,7*S*)-7-((6-chloro-2-naphthamido)methyl)-3-oxo-4-((*S*)-2-phenylbutyl)-1,4-diazepan-2-yl)ethylcarbamate



20

To a solution of the free amine **590** and 6-chloro-2-naphthoic acid (4.58 g, 22.3 mmol) in DCM (125 ml) was added diisopropylethylamine (7.74 mL, 44.5 mmol) and BOP (9.84 g, 22.3


mmol) at room temperature. The resulting mixture was stirred for 16 hours then DCM was removed under reduced pressure. The residue was taken up in EtOAc (80 mL), then washed with saturated NaHCO₃(aq) (100 mL x 5), brine (100 mL) and dried over MgSO₄. Filtration and concentration of the organic phase gave the crude material, which was purified by silica gel column chromatography using 80-100% EtOAc/Pet. Spirit as eluent to give the product **591** (10.7 g, 79%).

TLC *r_f* 0.31 (80% EtOAc/Pet. Spirit)

HPLC *t_R* 7.66 min

MS (ESI) 607.2 (M+1)

Example 98– Synthesis of Compound 465 *N*-(((3*S*,5*S*)-3-(2-aminoethyl)-2-oxo-1-((*S*)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide




To the Boc protected material **591** (10.7 g, 17.6 mmol) in DCM (26 mL) was added TFA (26 mL) in one portion, the resulting mixture was stirred at room temperature for one hour. DCM was removed under reduced pressure and the residue was taken up in EtOAc (30 mL), washed with saturated NaHCO₃(aq) (30 mL x 3), brine (30 mL) and dried over MgSO₄. Filtration and concentration of the organic phase under reduced pressure gave the crude amine **465**, which was used in the next step without further purification.

HPLC *t_R* 5.98 min

MS (ESI) 507.0 (M+1)

Example 99– Synthesis of Compound 425 6-chloro-*N*-(((3*S*,5*S*)-2-oxo-1-((*S*)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide



To a mixture of crude amine **465** in CH₃CN (800 mL) was added 1,5-dibromopentane (23.9 mL, 175.7 mmol) followed by K₂CO₃ (48.6 g, 351.4 mmol). The resulting mixture was stirred at room temperature for 44 hours, monitored by HPLC for conversion of sm (6.0 min) to product (6.4 min), avoiding extended reaction times leading to overalkylation to generate a bromopentyl-alkylated byproduct (7.1 min). During isolation, excessive heating/concentration of the crude solution must be avoided before removal of excess dibromopropane, to avoid overalkylation of the piperidine ring. The K₂CO₃ was removed by filtration through a pad of Celite[®], and the filtrate was washed with Pet. Spirit (800 mL x 2). The MeCN phase was concentrated under reduced pressure to 400 mL, and was washed with Pet. Spirit (400 mL x 2). The MeCN was further concentrated under reduced pressure to 200 mL, and was washed with Pet. Spirit (200 mL x 2). Evaporation of the final Pet. Spirit washing showed no more 1,5-dibromopentane was extracted, so the MeCN phase was concentrated under reduced pressure.

Aminopropyl-functionalized TLC rf 0.05-0.47 (80% EtOAc/Pet. Spirit)

Analytical HPLC *t_R* 6.41 min

MS (ESI) 575.2 (M+1).

The crude product was purified by a combination of flash column chromatography over aminopropyl-functionalized silica gel, and/or by recrystallization from acetonitrile.

Flash column: To a column packed with amino propyl-functionalized silica gel (154 g) in 20% ethyl acetate/petroleum spirit was loaded the crude free base oil (7.2 g). The column was eluted with 20% ethyl acetate/petroleum spirit (150 mL), followed by 50% ethyl acetate/petroleum spirit (150 mL), 80% ethyl acetate/petroleum spirit (150 mL x 2), 100% ethyl acetate (150 mL) and finally 100% acetonitrile (150 mL). Fractions containing **425** were combined and evaporated to dryness to yield a white crystalline solid.

Crystallization: The white crystalline solid (2.87 g) obtained by column purification was dissolved in boiling acetonitrile (50 mL) 85 °C. Activated carbon (Darco® G-60, -100 mesh, Sigma-Aldrich) (200 mg) was added to remove colour impurity. A further portion of acetonitrile (50 mL) was added, and the resulting mixture was heated to boiling for 5 min. The charcoal was filtered off while the solution was hot, with the filter paper and the charcoal rinsed with hot acetonitrile (25 mL). The clear acetonitrile solution was reduced down to 50 mL and left to stand to cool to room temperature for 16 h. The white crystals were filtered off and dried by suction to give 2.22 g (99.0% pure by HPLC analysis). An additional 117.2 mg (93.3% purity) was recovered by additional crystallization from the filtrate.

Conversion to bisHCl Salt: The free base (2.4229 g, 42.1 mmol) was suspended in a 1:1 mixture of acetonitrile and milliQ H₂O (10 mL). A solution of 1 M HCl (aq.) was added until all the solids dissolved (approximately 5 mL). An additional quantity of milliQ H₂O was then added (20 mL) and the resulting solution frozen and lyophilised overnight, resulting in a white powder (2.61 g, 95.6% yield).


HPLC t_R 6.27 min

MS (ESI) 575.1 (M+1).

¹H NMR (600 MHz, CDCl₃): δ 0.75 (t, 3 H, J = 7.2 Hz), 1.40 (m, 1 H), 1.56 (m, 1 H), 1.65 (m, 1 H), 1.76~1.90 (m, 4 H), 1.90~2.06 (m, 2 H), 2.13 (m, 1 H), 2.30 (br, 1 H), 2.57 (m, 1 H), 2.64~2.86 (m, 4 H), 2.90~3.10 (m, 2 H), 3.25 (dd, 1 H, J = 15.2, 10.4 Hz), 3.53 (m, 2 H), 3.70~3.85 (m, 3 H), 4.00 (m, 2 H), 4.10 (dd, 1 H, J = 13.6, 5.6 Hz), 4.45 (m, 1 H), 7.10 (d, 2 H, J = 7.2 Hz), 7.18 (t, 1 H, J = 7.2 Hz), 7.26 (t, 1 H, J = 7.2 Hz), 7.37 (dd, 1 H, J = 9.0, 1.8 Hz), 7.71 (d, 1 H, J = 8.4 Hz), 7.75 (s, 1 H), 7.86 (d, 1 H, J = 9.0 Hz), 8.09 (d, 1 H, J = 9.0 Hz), 8.64 (s, 1 H), 8.68 (m, 1 H), 9.85 (br, 1 H).

¹³C NMR (100 MHz, CDCl₃): δ 11.78, 21.86, 23.08 (2), 24.59, 26.14, 28.09, 42.01, 46.24, 47.22, 53.14, 53.53, 54.03, 56.74, 57.79, 61.96, 125.35, 126.24, 126.89, 127.20, 127.33, 127.85, 128.51, 128.72, 130.69, 130.76, 130.91, 133.48, 135.28, 142.29, 167.18, 167.74

Example 100– Synthesis of Compound 579 6-chloro-N-([5,6,6-²H₃](3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)[²H₂]methyl)-2-naphthamide



Compound **579** was synthesized according to the procedures in Examples 92-99, except that the Fmoc deprotection / reductive amination steps of Examples 94 and 95 were replaced by the following procedures in order to introduce the deuterium atoms.

To a solution of (S)-9-fluorenylmethyl 10-[(S)-2-phenylbutyl]-2,2-dimethyl-18-phenyl-4,9,13,16-tetraoxo-3,17-dioxo-5,10,15-triazaoctadecan-8-ylcarbamate **587** (370.5 mg, 0.47 mmol) in dry THF (7.5 ml) was added dry triethylamine (7.5 mL, 54 mmol) in one portion at room temperature followed by D₂O (99.96 atom % deuterium, 3.0 ml, 168 mmol). This mixture was stirred under nitrogen at room temperature for 16 h, with the reaction mixture used in the next step without isolation.

MS (ESI) 573.0 (M+1).

t_R 6.95 min.


To the crude deuterium exchange reaction mixture was added NaBD₃CN (152 mg, 2.31 mmol) in one portion, with the reaction stirred at room temperature for 24 h. A further portion of NaBD₃CN (182.4 mg, 3.28 mmol) was added and stirring continued at room temperature for 24 h. The reaction was quenched by the addition of saturated NaHCO₃(aq) and the aqueous mixture extracted with EtOAc (3 x 10 mL x 3). The combined organic extracts were washed with brine (15 mL), dried over MgSO₄ and concentrated in vacuo. Flash chromatography (60% EtOAc/Pet. Spirit) yielded the product (175.8 mg, 67%).

TLC R_f 0.32 (70% EtOAc/Pet. Spirit)

Analytical HPLC t_R 7.06 min; **MS (ESI)** m/z 558.0 (M+1), 559.0, 557.0, 560.0.

Example 101 – Syntheses of Compounds 106-579.

Compounds **106-579**, with substituents as identified in Table 1, were prepared as in the previous examples according to the routes identified in Schemes 1-5, as summarized in Table 2, with experimental properties summarized in Table 4.



5

Table 1: Identity of Compounds

Cpd	R ¹ X	R ²	R ³	Y	R	W
14	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	3,5-dichlorobenzyl
25	6-fluoro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ N(CH ₂ CH ₃) ₂	2,2-diphenylethyl
31	6-fluoro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
33	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	2,2-diphenylethyl
37	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	(2-phenylbutyl
38	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	(S)-2-phenylbutyl
39	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	(R)-2-phenylbutyl
49	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	3,5-dichlorobenzyl
50	2-naphthylsulfonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
54	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethylbutyl
60	6-bromo-2-naphthoyl	Me	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
62	6-bromo-2-naphthoyl	H	Me	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
63	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
64	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ (1-piperidiny)	2,2-diphenylethyl
65	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHCH(CH ₃) ₂	2,2-diphenylethyl
67	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NHMe	2,2-diphenylethyl
71	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ CH ₂ NHMe	2,2-diphenylethyl
79	acetyl	H	H	(S)-CHCH ₂ -(2-naphthyl)	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
81	Ac-L-His	H	H	(S)-CHCH ₂ -(2-naphthyl)	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
83	propyloxycarbonyl	H	H	(S)-CHCH ₂ -(2-naphthyl)	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
85	acetyl	H	H	(R)-CHCH ₂ -(2-naphthyl)	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
86	Ac-L-His	H	H	(R)-CHCH ₂ -(2-naphthyl)	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
87	propyloxycarbonyl	H	H	(R)-CHCH ₂ -(2-naphthyl)	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
105	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
106	4-biphenylcarbonyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
107	indole-2-carbonyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
108	4-biphenylcarbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
109	indole-2-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
110	2-naphthylacetyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl

Cpd	R ¹ X	R ²	R ³	Y	R	W
111	1,2,3,4-tetrahydro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
112	quinolin-3-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
113	quinoxaline-2-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
114	isoquinoline-3-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
115	benzoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
116	quinaldoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
117	2-naphthoyl	H	H	CH ₂	(CH ₂) ₄ NH ₂	1-naphthylmethyl
118	2-naphthylacetyl	H	H	CH ₂	(CH ₂) ₄ NH ₂	1-naphthylmethyl
119	1-naphthoyl	H	H	CH ₂	(CH ₂) ₄ NH ₂	1-naphthylmethyl
120	indole-3-acetyl	H	H	CH ₂	(CH ₂) ₄ NH ₂	1-naphthylmethyl
121	4-biphenylacetyl	H	H	CH ₂	(CH ₂) ₄ NH ₂	2-naphthylmethyl
122	2-naphthoyl	H	H	CH ₂	(CH ₂) ₄ NH ₂	2-naphthylmethyl
123	2-naphthylacetyl	H	H	CH ₂	(CH ₂) ₄ NH ₂	2-naphthylmethyl
124	1-naphthoyl	H	H	CH ₂	(CH ₂) ₄ NH ₂	2-naphthylmethyl
125	1-naphthylacetyl	H	H	CH ₂	(CH ₂) ₄ NH ₂	2-naphthylmethyl
126	2-naphthoyl	H	H	CH ₂	(CH ₂) ₄ NH ₂	2,2-diphenylethyl
127	S-Tic	H	H	CH ₂	(CH ₂) ₄ NH ₂	2,2-diphenylethyl
128	R-Tic	H	H	CH ₂	(CH ₂) ₄ NH ₂	2,2-diphenylethyl
129	2-benzofurananoyl	H	H	CH ₂	(CH ₂) ₄ NH ₂	2,2-diphenylethyl
130	R-Tic	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NHMe	2,2-diphenylethyl
131	S-Tic	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
132	2-benzofuranoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
133	indane-2-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
134	R-Tic	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
135	benzothiophene-2-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
136	2,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
137	2,5-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
138	benzoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
139	cyclohexanoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
140	3-phenoxybenzoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
141	4-phenoxybenzoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
142	indole-2-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
143	3-phenyl-propanoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
144	3,4-dimethylbenzoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
145	4-tert-butylbenzoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
146	2,4-dimethoxybenzoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
147	cyclohexylacetyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
148	piperonyloyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
149	benzimidazole-5-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
150	benzotriazole-5-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
151	cyclopentanoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
152	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl

Cpd	R ¹ X	R ²	R ³	Y	R	W
153	trans-cinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
154	3,5-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
155	2,4-dichloro-phenylacetyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
156	1-methoxy-2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
157	3,4-dichloro-phenylacetyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
158	6-methoxy-2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
159	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
160	2,4-dichloro-cinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
161	adamantane-1-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
162	phenoxyacetyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
163	3-methoxy-2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
164	4-bromobenzoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
165	S-benzodioxan-2-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
166	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
167	3-(2-thienyl)acryloyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
168	R-benzodioxan-2-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
169	4-hydroxycinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
170	2-methoxycinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
171	4-methylcinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
172	2-trifluoromethyl-cinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
173	3-fluorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
174	alpha-methyl cinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
175	trans- 2-phenylcyclopropane-1-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
176	2,4-dichloro-phenoxyacetyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
177	3-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
178	1,3-benzothiazole-6-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
179	5-phenyl-2-furoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
180	3-methoxycinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
181	6-bromo-2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
182	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	phenethyl
183	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	3,4-dichlorophenethyl
184	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,4-dichlorophenethyl
185	benzothiophene-5-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
186	3-methyl-2-phenyl-pyrazole-4-carbonyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
187	4-methoxycinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
188	6-fluoro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
189	2-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
190	2-hydroxycinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl

Cpd	R ¹ X	R ²	R ³	Y	R	W
191	3-methylcinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
192	3-trifluoromethyl-cinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
193	3-hydroxycinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
194	2-fluorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
195	2-methylcinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
196	alpha-fluorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
197	2-naphthoyl	H	H	CH ₂	4-piperidinyI	2,2-diphenylethyl
198	2-naphthoyl	H	H	CH ₂	CH ₂ (4-piperidinyI)	2,2-diphenylethyl
199	4-fluorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
200	4-trifluoromethyl-cinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
201	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylpropyl
202	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	cyclohexanemethyl
203	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	1-adamantane-methyl
204	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	(S)-1,1-diphenyl-2-propyl
205	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	(R)-1,1-diphenyl-2-propyl
206	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	cyclohexyl
207	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	(R)-1,1-diphenyl-1-fluoro-2-propyl
208	2,6-difluorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
209	2-chloro-6-fluorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
210	4-bromocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
211	4-ethoxycinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
212	6-bromonaphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
213	trans-cinnamoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
214	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
215	1,4-dimethoxy-2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
216	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)N(Me) ₂	2,2-diphenylethyl
217	6-hydroxy-2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
218	6-amino-2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
219	4-Me cinnamoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
220	4-fluoro cinnamoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
221	6-fluoro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
222	2-ethylhexanoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
223	3,4-dimethylbenzoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
224	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
225	2-ethylhexanoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-diphenylethyl
226	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH(cyclohexyl)	2,2-diphenylethyl
227	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2-naphthyl
228	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	(9-fluorenyl)methyl
229	2-naphthoyl	H	H	CH ₂	CH ₂ (3-pyridinyI)	2,2-diphenylethyl
230	2-naphthoyl	H	H	CH ₂	CH ₂ (4-pyridinyI)	2,2-diphenylethyl
231	4-fluorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NH(cyclohexyl)	2,2-diphenylethyl

Cpd	R ¹ X	R ²	R ³	Y	R	W
232	2-naphthoyl	H	H	CH ₂	(CH ₂) ₄ NHCH(CH ₃) ₂	2,2-diphenylethyl
233	2,4-difluorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
234	4-cyanocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
235	3-(2-naphthyl)acryloyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
236	4-fluoro-phenoxyacetyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
237	5-(4-chlorophenyl)-2-furoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
238	4-(pyrrol-1-yl)-benzoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
239	2-oxo-1-phenyl-pyrrolidine-3-carbonyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
240	5-(4-chlorophenyl)-isoxazole-3-carbonyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
241	5-(2-furyl)-isoxazole-3-carbonyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
242	2-phenyl-4-thiazole carbonyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
243	4-(3,5-dimethyl-1H-pyrazol-1-yl) benzoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
244	3-methyl-2-phenyl pyrazole-4-carbonyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
245	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	cyclohexaneethyl
246	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2-norbornaneethyl
247	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-bis(4-methoxyphenyl)ethyl
248	2-naphthoyl	H	H	CH ₂	(CH ₂) ₄ NHCH ₂ Ph	2,2-diphenylethyl
249	2-naphthoyl	H	H	CH ₂	(CH ₂) ₄ NH(cyclopentyl)	2,2-diphenylethyl
250	2-naphthoyl	H	H	CH ₂	(CH ₂) ₄ NH(cyclobutyl)	2,2-diphenylethyl
251	2-naphthoyl	H	H	CH ₂	(CH ₂) ₄ NH(cyclobutyl) ₂	2,2-diphenylethyl
252	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	benzyl
253	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2,2-bis(4-fluorophenyl)ethyl
254	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2-naphthalenemethyl
255	3-(5-methyl-2-thienyl)-acryloyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
256	5-phenyl-pyrazole-3-carbonyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
257	4-fluorocinnamoyl	Me	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
258	4-fluorocinnamoyl	H	Me	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
259	4-(3-methyl-5-oxo-2-pyrazolin-1yl)benzoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
260	4-bromocinnamoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,2-diphenylethyl
261	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ (1-pyrrolidinyl)	2,2-diphenylethyl
262	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ (1-piperidinyl)	2,2-diphenylethyl
263	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
264	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ (1-pyrrolidinyl)	2,2-diphenylethyl
265	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ (1-azetidyl)	2,2-diphenylethyl
266	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	1-naphthalenemethyl
267	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2-(2-naphthyl)ethyl
268	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	(S)-CH ₂ CH(Ph)NHCOMe

Cpd	R ¹ X	R ²	R ³	Y	R	W
269	trans-cinnamoyl	H	H	CH ₂	(CH ₂) ₃ (1-piperidiny)	2,2-diphenylethyl
270	3,4-dimethylbenzoyl	H	H	CH ₂	(CH ₂) ₃ (1-piperidiny)	2,2-diphenylethyl
271	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₃ (1-piperidiny)	2,2-diphenylethyl
272	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	(S)-CH ₂ CH(Ph)-NHCOCu
273	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	(S)-CH ₂ CH(Ph)-NHCOC ₆ H ₁₃
274	2-naphthoyl	H	H	CH ₂	CH ₂ NH ₂	2,2-diphenylethyl
275	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ NH ₂	2,2-diphenylethyl
276	4-fluorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
277	4-methylcinnamoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
278	2-naphthoyl	H	H	CH ₂	CH ₂ (1-piperidiny)	2,2-diphenylethyl
279	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ (1-piperidiny)	2,2-diphenylethyl
280	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
281	3,4-dimethylbenzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
282	trans-cinnamoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
283	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
284	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	2,2-diphenylethyl
285	3,4-dimethylbenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	2,2-diphenylethyl
286	trans-cinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	2,2-diphenylethyl
287	4-fluorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	2,2-diphenylethyl
288	4-methylcinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	2,2-diphenylethyl
289	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	3,5-dimethylbenzyl
290	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	(S)-CH ₂ CH(Ph)NHCOPh
291	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	(R)-CH ₂ CH(Ph)NHCOPh
292	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	CH ₂ CH(Ph)OMe
293	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	CH ₂ CH(Ph)OnPr
294	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	CH ₂ CH(Ph)OBn
295	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	CH ₂ CH(Ph)Oallyl
296	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ CH ₃	2,2-diphenylethyl
297	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₃ CH ₃	2,2-diphenylethyl
298	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ CONH ₂	2,2-diphenylethyl
299	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ CONH ₂	2,2-diphenylethyl
300	4-methylcinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHCOCH ₃	2,2-diphenylethyl
301	4-methylcinnamoyl	H	H	CH ₂	(CH ₂) ₃ NHCO(cyclohexyl)	2,2-diphenylethyl
302	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	CH ₂ CH(Ph)OPh
303	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	CH ₂ CH(Ph)CO ₂ Et
304	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	2-ethylbutyl
305	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	3,5-dimethyl-cyclohexylmethyl
306	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ NHCO(cyclohexyl)	2,2-diphenylethyl
307	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ NHCOCH ₂ (cyclohexyl)	2,2-diphenylethyl
308	benzoyl	H	H	(CH ₂) ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
309	3,4-dichlorobenzoyl	H	H	(CH ₂) ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
310	2-naphthoyl	H	H	(CH ₂) ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl

Cpd	R ¹ X	R ²	R ³	Y	R	W
311	benzoyl	H	H	(CH ₂) ₂	(CH ₂) ₂ (1-piperidiny)	2,2-diphenylethyl
312	3,4-dichlorobenzoyl	H	H	(CH ₂) ₂	(CH ₂) ₂ (1-piperidiny)	2,2-diphenylethyl
313	2-naphthoyl	H	H	(CH ₂) ₂	(CH ₂) ₂ (1-piperidiny)	2,2-diphenylethyl
314	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	CH ₂ CH(Ph)CONMe ₂
315	2-naphthoyl	H	H	CH ₂	CH ₂ cyclohexyl	2,2-diphenylethyl
316	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ cyclohexyl	2,2-diphenylethyl
317	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ CO(1-piperidiny)	2,2-diphenylethyl
318	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ CO(1-piperidiny)	2,2-diphenylethyl
319	2-naphthoyl	H	H	CH ₂	CH ₂ CH ₂ Ph	2,2-diphenylethyl
320	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ CH ₂ Ph	2,2-diphenylethyl
321	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ cyclohexyl	2,2-diphenylethyl
322	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ cyclohexyl	2,2-diphenylethyl
323	2-naphthoyl	H	H	CH ₂	CH ₂ Ph	2,2-diphenylethyl
324	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ Ph	2,2-diphenylethyl
325	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	3,5-dichlorobenzyl
326	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	3,5-dichlorobenzyl
327	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	3-chloro-5-fluorobenzyl
328	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	3,5-difluorobenzyl
329	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	3-chloro-5-fluorobenzyl
330	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	3,5-difluorobenzyl
331	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,5-dichlorobenzyl
332	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,6-dichlorobenzyl
333	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	3,5-dimethoxybenzyl
334	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2-chlorobenzyl
335	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,3-dichlorobenzyl
336	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,4-dichlorobenzyl
337	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	3,4-dichlorobenzyl
338	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	3-fluoro-5-methylbenzyl
339	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	3-fluoro-5-(trifluoromethyl)-benzyl
340	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	4-chlorobenzyl
341	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2-phenylbutyl
342	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	1-(1-phenyl-cyclohexyl)-methyl
343	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny))	3,5-dichlorobenzyl
344	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	3,5-dichlorobenzyl
345	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	3,5-dichlorobenzyl
346	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethylbutyl
347	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethylbutyl
348	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	2-ethylbutyl
349	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	2-ethylbutyl
350	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	2-ethylbutyl
351	4-chloro-3-fluoro-benzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethylbutyl

Cpd	R ¹ X	R ²	R ³	Y	R	W
352	4-chloro-3-methyl-benzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethylbutyl
353	3-chloro-4-fluoro-benzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethylbutyl
354	3-chloro-4-methyl-benzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethylbutyl
355	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ (1-piperidiny)	2-ethylbutyl
356	2-naphthoyl	H	H	(CH ₂) ₂	CH ₂ CH ₂ (1-pyrrolidiny)	2,2-diphenylethyl
357	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	3,5-bis(trifluoromethyl)-benzyl
358	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	3-chlorobenzyl
359	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ (1-piperidiny)	2-phenylbutyl (mixture isomers)
360	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	CH ₂ CH(Ph)CON[-(CH ₂) ₅ -]
361	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	CH ₂ CH(Ph)CONHPh
362	3,4-dichlorobenzoyl	H	H	(CH ₂) ₂	(CH ₂) ₂ NHCH(CH ₃) ₂	2,2-diphenylethyl
363	3,4-dichlorobenzoyl	H	H	(CH ₂) ₂	(CH ₂) ₂ N(CH(CH ₃) ₂) ₂	2,2-diphenylethyl
364	3,4-dichlorobenzoyl	H	H	CH ₂	CH ₂ NH ₂	3,5-dichlorobenzyl
365	2-naphthoyl	H	H	CH ₂	CH ₂ NH ₂	3,5-dichlorobenzyl
366	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ NH ₂	3,5-dichlorobenzyl
367	3,4-dichlorobenzoyl	H	H	CH ₂	CH ₂ (1-piperidiny)	3,5-dichlorobenzyl
368	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ (1-piperidiny)	3,5-dichlorobenzyl
369	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-phenylbutyl
370	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-phenylbutyl
371	3,4-dichlorobenzoyl	H	H	CH ₂	CH ₂ (1-pyrrolidiny)	3,5-dichlorobenzyl
372	2-naphthoyl	H	H	CH ₂	CH ₂ (1-pyrrolidiny)	3,5-dichlorobenzyl
373	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	(S)-β-methylphenethyl
374	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ CH ₂ NH ₂	(R)-β-methylphenethyl
375	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ N(CH ₃) ₂	2,2-diphenylethyl
376	6-fluoro-2--naphthoyl	H	H	CH ₂	(CH ₂) ₃ (1-piperidiny)	2,2-diphenylethyl
377	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	3,5-diethynylbenzyl
378	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ N(CH ₂ CH ₃) ₂	2,2-diphenylethyl
379	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ N(CH ₂ CH ₃) ₂	2,2-diphenylethyl
380	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	(R)-2-phenylbutyl
381	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	(R)-2-phenylbutyl
382	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	(S)-2-phenylbutyl
383	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
384	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-pyrrolidiny)	2,2-diphenylethyl
385	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	2,2-diphenylethyl
386	6-fluoro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethylbutyl
387	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethylbutyl
388	6-bromo-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethylbutyl
389	6-fluoro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	2-ethylbutyl
390	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	2-ethylbutyl
391	6-bromo-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidiny)	2-ethylbutyl

Cpd	R ¹ X	R ²	R ³	Y	R	W
392	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	3,5-dimethyl-cyclohexyl
393	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	3,5-dimethyl-cyclohexyl
394	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	3,5-dimethyl-cyclohexyl
395	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,2-diphenylethyl
396	6-fluoro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,2-diphenylethyl
397	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-pyrrolidinyI)	2,2-diphenylethyl
398	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ NHCH(CH ₃) ₂	2,2-diphenylethyl
399	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NHCH(CH ₃) ₂ r	2,2-diphenylethyl
400	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NHCH(CH ₃) ₂	2,2-diphenylethyl
401	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	2,6-dimethyl-cyclohexylmethyl
402	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	(S)-2-phenylbutyl
403	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	3,5-dimethyl-cyclohexylmethyl
404	3,4-dichlorobenzoyl	H	H	CH ₂	CH ₂ CH ₂ NHCH(CH ₃) ₂	3,5-dimethyl-cyclohexylmethyl
405	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	3-methyl-2-phenylbutyl
406	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	(S)-2-phenylbutyl
407	2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NH ₂	(R)-2-phenylbutyl
408	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ (imidazol-3-yl)	2,2-diphenylethyl
409	3-(4-chlorophenyl)-propanoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,2-diphenylethyl
410	3-(4-chlorophenyl)-propanoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
411	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ NHCO(2-pyridyl)	2,2-diphenylethyl
412	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ (2-pyridinyI)	2,2-diphenylethyl
413	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(R)-3-methyl-2-phenylbutyl
414	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-3-methyl-2-phenylbutyl
415	4-isopropylcinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,2-diphenylethyl
416	4-isopropylcinnamoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
417	2,4-dimethylcinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,2-diphenylethyl
418	2,4-dimethylcinnamoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
419	2,4-difluorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ 1-piperidinyI	2,2-diphenylethyl
420	2,4-difluorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
421	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ NHCO(cyclohexyl)	2,2-diphenylethyl
422	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ (4-morpholinyI)	2,2-diphenylethyl
423	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ CH ₂ N[-C(Me)=CHCH=C(Me)-]	2,2-diphenylethyl
424	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ CH ₂ (2,5-dimethyl-2-pyrrolidin-1-yl)	2,2-diphenylethyl
425	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
426	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-pyrrolidinyI)	(S)-2-phenylbutyl
427	4-chlorocinnamoyl	H	H	CH ₂	(CH ₂) ₂ NHCH(CH ₃) ₂	(S)-2-phenylbutyl
428	6-chloro-2-naphthoyl	H	H	CH ₂	CH ₂ CH ₂ (1-pyrrolidinyI)	(S)-2-phenylbutyl

Cpd	R ¹ X	R ²	R ³	Y	R	W
429	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
430	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ 1-pyrrolidinyI	(S)-2-phenylbutyl
431	Cbz	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,2-diphenylethyl
432	4-bromocinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,2-diphenylethyl
433	5-(4-chlorophenyl)- isoxazole-3-carbonyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,2-diphenylethyl
434	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ CONH(2-pyridyl)	2,2-diphenylethyl
435	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ CO(1-piperidinyI)	2,2-diphenylethyl
436	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ CO(1-piperidinyI)	(S)-2-phenylbutyl
437	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ CO(1-piperidinyI)	(S)-2-phenylbutyl
438	6-chloro-2-naphthoyl	H	H	CH ₂	CH ₂ (1-piperidinyI)	(S)-2-phenylbutyl
439	3,4-dichlorobenzoyl	H	H	CH ₂	CH ₂ (1-piperidinyI)	(S)-2-phenylbutyl
440	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ (1-piperidinyI)	(S)-2-phenylbutyl
441	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₃ (1-piperidinyI)	(S)-2-phenylbutyl
442	4-chlorocinnamoyl	H	H	C(Me) ₂	CH ₂ CH ₂ NH ₂	2,2-diphenylethyl
443	4-chlorocinnamoyl	H	H	C(Me) ₂	(CH ₂) ₂ (1-piperidinyI)	2,2-diphenylethyl
444	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	(R)-2-(4-chloro-phenyl)propyl
445	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-(4-chloro-phenyl)propyl
446	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(R)-2-(4-chloro-phenyl)propyl
447	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-(4-chloro-phenyl)propyl
448	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ N(phenyl)CH ₃	(S)-2-phenylbutyl
449	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ N(CH ₂ CH ₃) ₂	(S)-2-phenylbutyl
450	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (4-morpholinyI)	(S)-2-phenylbutyl
451	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH(phenyl)	(S)-2-phenylbutyl
452	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH(benzyl)	(S)-2-phenylbutyl
453	4-chlorocinnamoyl	H	H	CH ₂	CH ₂ (2-NH ₂ -Ph)	2,2-diphenylethyl
454	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NHC(CH ₃) ₃	(S)-2-phenylbutyl
455	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (4-CH ₃ -piperazin-1-yl)	(S)-2-phenylbutyl
456	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(R)-2-phenylpentyl
457	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylpentyl
458	p-trifluoromethyl-benzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,2-diphenylethyl
459	p-trifluoromethyl-benzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
460	m-trifluoromethyl-benzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,2-diphenylethyl
461	m-trifluoromethyl-benzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,2-diphenylethyl
462	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NHCH(CH ₃) ₂	3,5-dichlorobenzyl
463	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NHCH(CH ₃) ₂	3,5-dichlorobenzyl
464	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NHCH(CH ₃) ₂	(S)-2-phenylbutyl
465	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	(S)-2-phenylbutyl
466	2-naphthoyl	H	H	CH ₂	CH ₂ (2-NH ₂ -Ph)	2,2-diphenylethyl
467	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ N(benzyl)CH ₃	(S)-2-phenylbutyl
468	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (piperazin-1-yl)	(S)-2-phenylbutyl

Cpd	R ¹ X	R ²	R ³	Y	R	W
469	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ N(<i>n</i> -pentyl)CH ₃	(S)-2-phenylbutyl
470	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ N[(CH(CH ₃) ₂) ₂]	(S)-2-phenylbutyl
471	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (4-CH ₃ -piperidin-1-yl)	(S)-2-phenylbutyl
472	6-chloro-2-naphthoyl	H	H	CH ₂	4-piperidinyI	(S)-2-phenylbutyl
473	6-chloro-2-naphthoyl	H	H	CH ₂	1-isopentyl-4-piperidinyI	(S)-2-phenylbutyl
474	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ CH ₃	(S)-2-phenylbutyl
475	6-chloro-2-naphthoyl	H	H	CH ₂	CH ₂ CH ₂ <i>i</i> Pr	(S)-2-phenylbutyl
476	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (3,5-Me ₂ -piperidin-1-yl)	(S)-2-phenylbutyl
477	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (4-OH-piperidin-1-yl)	(S)-2-phenylbutyl
478	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (4-CO ₂ H-piperidin-1-yl)	(S)-2-phenylbutyl
479	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH[-(CH ₂) ₆ -]	(S)-2-phenylbutyl
480	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ [(S)-2-Me-piperidin-1-yl]	(S)-2-phenylbutyl
481	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ N(<i>t</i> Bu)CH ₃	(S)-2-phenylbutyl
482	2-naphthoyl	H	H	CH ₂	CH ₂ (2-(piperidin-1-yl)phenyl)-	2,2-diphenylethyl
483	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ CON(Me) <i>n</i> Bu	(S)-2-phenylbutyl
484	2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ CONHcHex	(S)-2-phenylbutyl
485	6-chloro-2-naphthoyl	H	H	CH ₂	1-ethyl-piperidin-4-yl	(S)-2-phenylbutyl
486	3,4-dichlorobenzyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,2-diphenylethyl
487	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NHC(=NH)NH ₂	(S)-2-phenylbutyl
488	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NHC(=NH)NH Me	(S)-2-phenylbutyl
489	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	(R)-2-isopropylbutyl
490	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	(S)-2-isopropylbutyl
491	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(R)-2-isopropylbutyl
492	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-isopropylbutyl
493	6-chloro-2-naphthoyl	H	H	CH ₂	S-CH ₂ C(Me ₂)NH ₂	(S)-2-phenylbutyl
494	6-chloro-2-naphthoyl	H	H	CH ₂	R-CH ₂ C(Me ₂)NH ₂	(S)-2-phenylbutyl
495	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	(S)-2-phenylbutyl
496	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NHCH(CH ₃) ₂	(S)-2-phenylbutyl
497	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	3,5-dichlorobenzyl
498	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	3,5-dichlorobenzyl
499	6-chloro-2-naphthoyl	H	H	CH ₂	S-CH ₂ C(Me ₂)(1-piperidinyI)	(S)-2-phenylbutyl
500	6-chloro-2-naphthoyl	H	H	CH ₂	R-CH ₂ C(Me ₂)(1-piperidinyI)	(S)-2-phenylbutyl
501	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	(R)-2-isopropylbutyl

Cpd	R ¹ X	R ²	R ³	Y	R	W
502	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	(S)-2-isopropylbutyl
503	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(R)-2-isopropylbutyl
504	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-isopropylbutyl
505	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ cHex	(S)-2-phenylbutyl
506	6-chloro-2-naphthoyl	H	H	CH ₂	nHex	(S)-2-phenylbutyl
507	6-carboxy-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,2-diphenylethyl
508	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NHC(=NH)NH-CH(CH ₃) ₂	(S)-2-phenylbutyl
509	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₄ OH	(S)-2-phenylbutyl
510	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ OMe	(S)-2-phenylbutyl
511	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ OBn	(S)-2-phenylbutyl
512	6-chloro-2-naphthoyl	H	H	CH ₂	iBu	(S)-2-phenylbutyl
513	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethyl-3-methyl-but-3-enyl
514	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2-ethyl-3-methyl-but-3-enyl
515	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethyl-3-methyl-but-3-enyl
516	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(R)-2-ethyl-3-methyl-but-3-enyl
517	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-ethyl-3-methyl-but-3-enyl
518	4-biphenylcarboxylyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	cyclohexanemethyl
519	indole-3-acetyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	cyclohexanemethyl
520	3-quinolinecarboxylyl	H	H	CH ₂	(CH ₂) ₃ NHC(=NH)NH ₂	cyclohexanemethyl
521	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (4,4-difluoro-1-piperidinyI)	(S)-2-phenylbutyl
522	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ OH	(S)-2-phenylbutyl
523	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (3,3-difluoro-1-piperidinyI)	(S)-2-phenylbutyl
524	3,4-dichlorobenzyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
525	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2-cyclopropylbutyl
526	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-cyclopropylbutyl
527	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2-cyclopropylbutyl
528	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ N[-CO(CH ₂) ₂ CO-]	(S)-2-phenylbutyl
529	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₃ NHCONH ₂	(S)-2-phenylbutyl
530	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NHCH(Me)CF ₃	(S)-2-phenylbutyl
531	3,4-dichlorobenzoyl	H	H	CH ₂	CH ₂ CH ₂ N[-COC(Me) ₂ CH ₂ CO-]	(S)-2-phenylbutyl
532	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ N[-(CH ₂) ₆ -]	(S)-2-phenylbutyl

Cpd	R ¹ X	R ²	R ³	Y	R	W
533	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NHCONHiPr	(S)-2-phenylbutyl
534	4-biphenyl carboxylic	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
535	2-phenylthiazole-4-carbonyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
536	4-chloro-biphenyl-2-carbonyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
537	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ N(Ac)iPr	(S)-2-phenylbutyl
538	6-chloro-2-naphthoyl	H	H	CH ₂	CH ₂ NHiPr	(S)-2-phenylbutyl
539	6-chloro-2-naphthoyl	H	H	CH ₂	CH ₂ NC(=NH)NH ₂	(S)-2-phenylbutyl
540	2,4-dichlorophenylacetyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
541	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,4-dichlorobenzyl
542	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,4-dichlorobenzyl
543	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NHSO ₂ Me	2,4-dichlorobenzyl
544	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NHSO ₂ (4-Me-Ph)	2,4-dichlorobenzyl
545	3,4-dichlorobenzoyl	H	H	CH ₂	(S)- (CH ₂) ₂ NHCO-CH(iPr)NH ₂	2,4-dichlorobenzyl
546	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2,4-dichlorobenzyl
547	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,4-dichlorobenzyl
548	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-(3-thienyl)butyl
549	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(R)-2-(3-thienyl)butyl
550	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-(3-thienyl)butyl
551	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethyl-2-methylbutyl
552	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2-ethyl-2-methylbutyl
553	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (4-morpholinyI)	2,2-diphenylethyl
554	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (4-morpholinyI)	(S)-2-phenylbutyl
555	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (4-morpholinyI)	3,5-dichlorobenzyl
556	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (4-morpholinyI)	3,5-dichlorobenzyl
557	(4-chloro-benzyl)NHCO	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
558	3,4-dichlorobenzyl + MeCO	Ac	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
559	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ NH ₂	2-ethyl-2-methylbutyl
560	3,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2-ethyl-2-methylbutyl
561	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	2,3,5-trichlorobenzyl
562	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NHSO ₂ iPr	(S)-2-phenylbutyl
563	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ NHCO ₂ nBu	(S)-2-phenylbutyl

Cpd	R ¹ X	R ²	R ³	Y	R	W
564	6-chloro-2-naphthoyl	H	H	CH ₂	1-iPr-4-piperidinyI	(S)-2-phenylbutyl
565	6-chloro-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(R)-2-phenylbutyl
566	5-(4-chlorophenyl)-isoxazole-3-carbonyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	3,5-dichlorobenzyl
567	2,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	3,5-dichlorobenzyl
568	6-methoxy-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	3,5-dichlorobenzyl
569	6-chloro-2-naphthoyl	H	H	¹³ CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
570	1-methoxy-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	3,5-dichlorobenzyl
571	4-(trifluoromethoxy)cinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	3,5-dichlorobenzyl
572	5-(4-chlorophenyl)-isoxazole-3-carbonyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
573	2,4-dichlorobenzoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
574	4,5-dichlorophthaloyl	R1	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	3,5-dichlorobenzyl
575	3-fluoro-4-(trifluoromethoxy)cinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	3,5-dichlorobenzyl
576	6-methoxy-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
577	1-methoxy-2-naphthoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
578	3-fluoro-4-(trifluoromethoxy)cinnamoyl	H	H	CH ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl
579	6-chloro-2-naphthoyl	H	H	CD ₂	(CH ₂) ₂ (1-piperidinyI)	(S)-2-phenylbutyl

Table 2: Synthesis of Compounds

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
14	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
25	Scheme 1	Boc-Gly-OH	Cbz-L-Asp[(NMe)OMe]-OH	Scheme 4	reduction to aldehyde then reductive amination
31	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
33	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
37	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
38	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
38	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
39	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
49	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
50	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
54	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
60	Scheme 1	Cbz-Sar	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
62	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P1 deprotect, R1 acylate, ring methylate, P3 deprotect
63	Scheme 2	Boc-Gly-OH	H-L-Orn(Cbz)-Oallyl	Scheme 4	P3 deprotection
63	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
64	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
65	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection, reductive alkylation
67	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection, guanylation
79	Scheme 1	Alloc-β-(2-naphthyl)-L-Ala	Boc-L-Arg-(Cbz) ₂ -OH	Scheme 4	P3 deprotection
81	Scheme 1	Alloc-β-(2-naphthyl)-L-Ala	Boc-L-Arg-(Cbz) ₂ -OH	Scheme 4	P3 deprotection
83	Scheme 1	Alloc-β-(2-naphthyl)-L-Ala	Boc-L-Arg-(Cbz) ₂ -OH	Scheme 4	P3 deprotection
85	Scheme 1	Alloc-β-(2-naphthyl)-L-Ala	Boc-L-Arg-(Cbz) ₂ -OH	Scheme 4	P3 deprotection
86	Scheme 1	Alloc-β-(2-naphthyl)-L-Ala	Boc-L-Arg-(Cbz) ₂ -OH	Scheme 4	P3 deprotection
87	Scheme 1	Alloc-β-(2-naphthyl)-L-Ala	Boc-L-Arg-(Cbz) ₂ -OH	Scheme 4	P3 deprotection
105	Scheme 2	Boc-Gly-OH	H-L-Orn(Cbz)-Oallyl	Scheme 5	P3 deprotection, guanidinylation, deprotection
105	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
105	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
105	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
106	Scheme 2	Boc-Gly-OH	H-L-Orn(Cbz)-Oallyl	Scheme 4	P3 deprotection
106	Scheme 2	Boc-Gly-OH	H-L-Orn(Cbz)-Oallyl	Scheme 4	P3 deprotection
107	Scheme 2	Boc-Gly-OH	H-L-Orn(Cbz)-Oallyl	Scheme 4	P3 deprotection
108	Scheme 2	Boc-Gly-OH	H-L-Orn(Cbz)-Oallyl	Scheme 5	P3 deprotection, guanidinylation, deprotection
109	Scheme 2	Boc-Gly-OH	H-L-Orn(Cbz)-Oallyl	Scheme 5	P3 deprotection, guanidinylation, deprotection
110	Scheme 2	Boc-Gly-OH	H-L-Orn(Cbz)-Oallyl	Scheme 5	P3 deprotection, guanidinylation, deprotection
111	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
112	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
113	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
114	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
115	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
116	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
117	Scheme 2	Boc-Gly-OH	Boc-L-Lys(Cbz)-OH	Scheme 4	P3 deprotection
118	Scheme 2	Boc-Gly-OH	Boc-L-Lys(Cbz)-OH	Scheme 4	P3 deprotection
119	Scheme 2	Boc-Gly-OH	Boc-L-Lys(Cbz)-OH	Scheme 4	P3 deprotection

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
120	Scheme 2	Boc-Gly-OH	Boc-L-Lys(Cbz)-OH	Scheme 4	P3 deprotection
121	Scheme 2	Boc-Gly-OH	Boc-L-Lys(Cbz)-OH	Scheme 4	P3 deprotection
122	Scheme 2	Boc-Gly-OH	Boc-L-Lys(Cbz)-OH	Scheme 4	P3 deprotection
123	Scheme 2	Boc-Gly-OH	Boc-L-Lys(Cbz)-OH	Scheme 4	P3 deprotection
124	Scheme 2	Boc-Gly-OH	Boc-L-Lys(Cbz)-OH	Scheme 4	P3 deprotection
125	Scheme 2	Boc-Gly-OH	Boc-L-Lys(Cbz)-OH	Scheme 4	P3 deprotection
126	Scheme 2	Boc-Gly-OH	Boc-L-Lys(Cbz)-OH	Scheme 4	P3 deprotection
127	Scheme 2	Boc-Gly-OH	Boc-L-Lys(Cbz)-OH	Scheme 4	P3 deprotection
128	Scheme 2	Boc-Gly-OH	Boc-L-Lys(Cbz)-OH	Scheme 4	P3 deprotection
129	Scheme 2	Boc-Gly-OH	Boc-L-Lys(Cbz)-OH	Scheme 4	P3 deprotection
130	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
131	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
132	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
133	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
134	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
135	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
136	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
137	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
138	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
139	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
140	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
141	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
142	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
143	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
144	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
145	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
146	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
147	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
148	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
149	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
150	Scheme 2	Boc-Gly-OH	H-L-Arg(Cbz) ₂ -Oallyl	Scheme 4	P3 deprotection
151	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
152	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
153	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
154	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
155	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
156	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
157	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
158	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
159	Scheme 1	Alloc-Gly-OH	Boc-L-Canavanine (Fmoc)-OH	Scheme 4	P3 deprotection
160	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
161	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
162	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
163	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
164	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
165	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
166	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
167	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
168	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
169	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
170	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
171	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
172	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
173	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
174	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
175	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
176	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
177	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
178	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
179	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
180	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
181	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
182	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
183	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
184	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
185	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
186	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
187	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
188	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
189	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
190	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
191	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
192	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
193	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
194	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
195	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
196	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
197	Scheme 1	2-naphthoic-Gly-OH	Fmoc- α -(1-Boc-4-piperidiny)-DL-Gly-OH	Scheme 3	P3 deprotection
198	Scheme 1	2-naphthoic-Gly-OH	Fmoc- β -(1-Boc-4-piperidiny)-DL-Ala-OH	Scheme 3	P3 deprotection
199	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
200	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
201	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
202	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
203	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
204	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
205	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
206	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
207	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
208	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
209	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
210	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
211	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
212	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
213	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
214	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
215	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
216	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(NMe) ₂ Pbf-OH	Scheme 3	P3 deprotection
217	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
218	Scheme 1	Alloc-Gly-OH	Boc-L-Arg(Fmoc) ₂ -OH	Scheme 4	P3 deprotection
219	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
220	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
221	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
222	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
223	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn (Boc)-OH	Scheme 4	P3 deprotection
224	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
225	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection, guanidinylation
226	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection, reductive alkylation
227	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
228	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
229	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-3-PyridylAla-OH	Scheme 3	none
230	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-4-PyridylAla-OH	Scheme 3	none
231	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection, reductive alkylation
232	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Lys(i-Pr)Fmoc-OH	Scheme 3	P3 deprotection
233	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
234	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
235	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
236	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
237	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
238	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
239	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
240	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
241	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
242	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
243	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
244	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
245	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
246	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
247	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
248	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection, reductive alkylation
249	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection, reductive alkylation
250	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection, reductive alkylation
251	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection, reductive alkylation
252	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
253	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
254	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
255	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
256	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
257	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
258	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P1 deprotect, R1 acylate, ring methylate, P3 deprotect
259	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
260	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection
261	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
262	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
262	Scheme 1	Boc-Gly-OH	Fmoc-L-Orn(Cbz)-OH	Scheme 5	P3 deprotection, dialkylation
263	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection
263	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
264	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
265	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
266	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
267	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
268	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
269	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
270	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn (Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
271	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
272	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
273	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
274	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection
275	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection
276	Scheme 1	Cbz-Gly-OH	Boc-L-Dab(Fmoc)-OH	Scheme 4	P3 deprotection
276	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
277	Scheme 1	Cbz-Gly-OH	Boc-L-Dab(Fmoc)-OH	Scheme 4	P3 deprotection
277	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
278	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
279	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
279	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
280	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
281	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
282	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
283	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
284	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
285	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
286	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
287	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
288	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
289	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
290	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
291	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
292	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
293	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
294	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
295	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
296	Scheme 1	Cbz-Gly-OH	Boc-L-Nle-OH	Scheme 4	none
297	Scheme 1	Cbz-Gly-OH	Boc-L-Nle-OH	Scheme 4	none
298	Scheme 1	Cbz-Gly-OH	Boc-L-Gln-OH	Scheme 4	none
299	Scheme 1	Cbz-Gly-OH	Boc-L-Gln-OH	Scheme 4	none
300	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection then acylation
301	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection then acylation
302	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
303	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
304	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
305	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
306	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then acylation
307	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then acylation
308	Scheme 1	Cbz-β-Ala	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
309	Scheme 1	Cbz-β-Ala	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
310	Scheme 1	Cbz-β-Ala	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
311	Scheme 1	Cbz-β-Ala	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
312	Scheme 1	Cbz-β-Ala	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
313	Scheme 1	Cbz-β-Ala	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
314	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	R5 deprotection and amidation then P3 deprotection
315	Scheme 1	Cbz-Gly-OH	Boc-L-Cha-OH	Scheme 4	none
316	Scheme 1	Cbz-Gly-OH	Boc-L-Cha-OH	Scheme 4	none
317	Scheme 1	Cbz-Gly-OH	Boc-L-Glu(1-piperidiny)-OH	Scheme 4	none
318	Scheme 1	Cbz-Gly-OH	Boc-L-Glu(1-piperidiny)-OH	Scheme 4	none
319	Scheme 1	Cbz-Gly-OH	Boc-L-Hfe-OH	Scheme 4	none
320	Scheme 1	Cbz-Gly-OH	Boc-L-Hfe-OH	Scheme 4	none
321	Scheme 1	Cbz-Gly-OH	Boc-L-hCha-OH	Scheme 4	none
322	Scheme 1	Cbz-Gly-OH	Boc-L-hCha-OH	Scheme 4	none
323	Scheme 1	Cbz-Gly-OH	Boc-L-Phe-OH	Scheme 4	none
324	Scheme 1	Cbz-Gly-OH	Boc-L-Phe-OH	Scheme 4	none
325	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
326	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
327	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
328	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	P3 deprotection
329	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
330	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
331	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
332	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
333	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
334	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
335	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
336	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
337	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
338	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
339	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
340	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
341	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
342	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
343	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
344	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
345	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
346	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
347	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
348	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
348	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
349	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
350	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
351	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
352	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
353	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
354	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
355	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
356	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
357	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
358	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
359	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
360	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	R5 deprotection and amidation then P3 deprotection
361	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Arg(Pbf)-OH	Scheme 3	R5 deprotection and amidation then P3 deprotection
362	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection, alkylation
363	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection, dialkylation
364	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection
365	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection
366	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
367	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
367	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
368	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
369	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
370	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
371	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
372	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
373	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
374	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
375	Scheme 1	2-naphthoic-Gly-OH	Cbz-L-Asp[N(Me)OMe]-OH	Scheme 3	P3 conversion to aldehyde then reductive amination
376	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
377	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
378	Scheme 1	Boc-Gly-OH	Cbz-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
379	Scheme 1	Boc-Gly-OH	Cbz-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
380	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
381	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
382	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
383	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
384	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
385	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
386	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
387	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
388	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
389	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
390	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
391	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
392	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
393	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
394	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
395	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
396	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
397	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
398	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, alkylation
399	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, alkylation
400	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, alkylation
401	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
402	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
403	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
404	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, alkylation

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
405	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection
406	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection
407	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 3	P3 deprotection
408	Scheme 1	Cbz-Gly-OH	Fmoc-L-His(Boc)-OH	Scheme 4	P3 deprotection
409	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
410	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
411	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection, acylation
412	Scheme 2	Cbz-Gly-OH	H-β-(2-pyridyl)-L-Ala-Oallyl	Scheme 4	none
413	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
414	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
415	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
416	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
417	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
418	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
419	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
420	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
421	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection, acylation
422	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
423	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, condensation
424	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, condensation, reduction
425	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
426	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
427	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
428	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
429	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
430	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
431	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
432	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
433	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
434	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp(OtBu)-OH	Scheme 4	P3 deprotection, amidation
435	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp(OtBu)-OH	Scheme 4	P3 deprotection, amidation
436	Scheme 1	Cbz-Gly-OH	Boc-L-Gln(piperidyl)-OH	Scheme 4	none
437	Scheme 1	Cbz-Gly-OH	Boc-L-Gln(piperidyl)-OH	Scheme 4	none
438	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
439	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
440	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
441	Scheme 1	Cbz-Gly-OH	Fmoc-L-Orn(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
442	Scheme 1	Boc-Aib	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
443	Scheme 1	Boc-Aib	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
444	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection
445	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection
446	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
447	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
448	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
449	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
450	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
451	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
452	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 conversion to aldehyde then reductive amination

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
453	Scheme 1	Fmoc-Gly-OH	Boc-L-(2-NO ₂)-Phe-OH	Scheme 5	nitro hydrogenation
454	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
455	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
456	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
457	Scheme 1	2-naphthoic-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
458	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
459	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	none
460	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
461	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	none
462	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, reductive alkylation
463	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, reductive alkylation
464	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, reductive alkylation
465	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection
466	Scheme 1	Fmoc-Gly-OH	Boc-L-(2-NO ₂)-Phe-OH	Scheme 4	nitro hydrogenation
467	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
468	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
469	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
470	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
471	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
472	Scheme 1	Cbz-Gly-OH	N-Fmoc-1(1-Boc-piperidin-4yl)-D,L-Gly-OH	Scheme 4	P3 deprotection
473	Scheme 1	Cbz-Gly-OH	N-Fmoc-1(1-Boc-piperidin-4yl)-D,L-Gly-OH	Scheme 4	P3 deprotection, reductive alkylation
474	Scheme 1	Cbz-Gly-OH	Boc-L-Nle-OH	Scheme 4	none
475	Scheme 1	Cbz-Gly-OH	Fmoc-L-HoLeu-OH	Scheme 4	none
476	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
477	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
478	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
479	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
480	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
481	Scheme 1	Cbz-Gly-OH	Fmoc-L-Asp[N(Me)OMe]-OH	Scheme 4	P3 conversion to aldehyde then reductive amination
482	Scheme 1	Fmoc-Gly-OH	Boc-L-(2-NO ₂)-Phe-OH	Scheme 5	nitro hydrogenation then dialkylation with alkyl dibromide
483	Scheme 1	2-naphthoic-Gly-OH	Boc-L-Gln(Me,nBu)-OH	Scheme 4	none
484	Scheme 1	2-naphthoic-Gly-OH	Boc-L-Gln(chex)-OH	Scheme 4	none
485	Scheme 1	Cbz-Gly-OH	N-Fmoc-1-(1-Boc-piperidin-4-yl)-D,L-Gly-OH	Scheme 4	P3 deprotection, reductive alkylation
486	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection, alkylation
487	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection, alkylation
488	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection, alkylation
489	Scheme 1	Boc-Gly-OH	Cbz-DL-γ-nitro-Leu-OH	Scheme 4	P3 reduction to amine
490	Scheme 1	Boc-Gly-OH	Cbz-DL-γ-nitro-Leu-OH	Scheme 4	P3 reduction to amine
491	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
492	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
493	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, dialkylation with alkyl dibromide
494	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, dialkylation with alkyl dibromide
495	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
496	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then reductive alkylation
497	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
498	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation
499	Scheme 1	Boc-Gly-OH	Cbz-DL-γ-nitro-Leu-OH	Scheme 4	P3 reduction to amine then dialkylation with alkyl dibromide
500	Scheme 1	Boc-Gly-OH	Cbz-DL-γ-nitro-Leu-OH	Scheme 4	P3 reduction to amine then dialkylation with alkyl dibromide
501	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
502	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
503	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, dialkylation with alkyl dibromide

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
504	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, dialkylation with alkyl dibromide
505	Scheme 1	Cbz-Gly-OH	Fmoc-L-HoCha-OH	Scheme 4	none
506	Scheme 1	Cbz-Gly-OH	Fmoc-L-2-aminooctanoic acid	Scheme 4	none
507	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection, alkylation
508	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection, alkylation
509	Scheme 1	Cbz-Gly-OH	Boc-L-5-HO-Nle-OH	Scheme 4	none
510	Scheme 1	Cbz-Gly-OH	Fmoc-L-HoSer(Me)-OH	Scheme 4	none
511	Scheme 1	Alloc-Gly-OH	Boc-L-HoSer(Bzl)-OH	Scheme 4	none
512	Scheme 1	Cbz-Gly-OH	Boc-L-Leu-OH	Scheme 4	none
513	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
514	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, dialkylation with alkyl dibromide
515	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
516	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, dialkylation with alkyl dibromide
517	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, dialkylation with alkyl dibromide
518	Scheme 2	Boc-Gly-OH	Boc-L-Arg(Cbz) ₂ -OH	Scheme 4	P3 deprotection
519	Scheme 2	Boc-Gly-OH	Boc-L-Arg(Cbz) ₂ -OH	Scheme 4	P3 deprotection
520	Scheme 2	Boc-Gly-OH	Boc-L-Arg(Cbz) ₂ -OH	Scheme 4	P3 deprotection
521	Scheme 1	Boc-Gly-OH	Cbz-L-Asp[N(Me)OMe]	Scheme 4	P3 conversion to aldehyde then reductive amination
522	Scheme 1	Boc-Gly-OH	Cbz-L-Asp[N(Me)OMe]	Scheme 4	P3 conversion to aldehyde then reduction
523	Scheme 1	Boc-Gly-OH	Cbz-L-Asp[N(Me)OMe]	Scheme 4	P3 conversion to aldehyde then reductive amination
524	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5; reductive alkylation for R1X	P3 deprotection then dialkylation with alkyl dibromide
525	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
526	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
527	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
528	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)	Scheme 4	P3 deprotection, diacylation with anhydride
529	Scheme 1	Cbz-Gly-OH	Boc-L-citrulline-OH	Scheme 4	none

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
530	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)	Scheme 4	P3 deprotection, reductive alkylation
531	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)	Scheme 4	P3 deprotection, diacylation with anhydride
532	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
533	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)	Scheme 4	P3 deprotection, then acylation with isocyanate
534	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
535	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
536	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
537	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then reductive alkylation then acetylation
538	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection then reductive alkylation
539	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dap(Boc)-OH	Scheme 4	P3 deprotection then guanylation
540	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection, guanidinylation, deprotection
541	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
542	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
543	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then sulfonylation
544	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then sulfonylation
545	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection, acylation, deprotection
546	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
547	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
548	Scheme 1	N-(6-Cl-2-napthoic)-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection
549	Scheme 1	N-(6-Cl-2-napthoic)-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
550	Scheme 1	N-(6-Cl-2-napthoic)-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 3	P3 deprotection then dialkylation with alkyl dibromide
551	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
552	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
553	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
554	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
555	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
556	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
557	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5, use isocyanate for R1X	P3 deprotection then dialkylation with alkyl dibromide
558	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5; reductive alkylation then acetylation for R1X and R2	P3 deprotection then dialkylation with alkyl dibromide
559	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection
560	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
561	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
562	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then sulfonylation
563	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then acylation with chloroformate
564	Scheme 1	Cbz-Gly-OH	Fmoc-DL-2-(1-Boc-4-piperidyl)-Gly-OH	Scheme 4	P3 deprotection then reductive alkylation with ketone
565	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
566	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
567	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
568	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
569	Scheme 1	Cbz-[¹⁵ N,1,2- ¹³ C ₂]-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then reductive alkylation then acetylation
570	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
571	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
572	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide

Cpd.	Route to A	Scheme 1: VN(R ²)-Y-CO ₂ H	P ² NH-CH(U)-CO ₂ H	Conversion of A to Product	U modification
573	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
574	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
575	Scheme 1	Alloc-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
576	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
577	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
578	Scheme 1	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 5	P3 deprotection then dialkylation with alkyl dibromide
579	Scheme 1 with D ₂ O exchange during Fmoc deprotection and NaBD ₃ CN reduction	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then reductive alkylation then acetylation
580	Scheme 1	Cbz-Gly-OH	Fmoc-D-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
581	Scheme 2	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
582	Scheme 2	Cbz-Gly-OH	Fmoc-D-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
583	Scheme 2	Cbz-Gly-OH	Fmoc-L-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
584	Scheme 2	Cbz-Gly-OH	Fmoc-D-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide
585	Scheme 1	Cbz-Gly-OH	Fmoc-D-Dab(Boc)-OH	Scheme 4	P3 deprotection then dialkylation with alkyl dibromide

Example 102 – Human MC5R Radioligand Binding Assay

Assessments of compound binding to human MC5R (hMC5R) by displacement of an ¹²⁵I-labeled NDP-MSH receptor ligand peptide were performed essentially as described in the data sheets produced by Perkin Elmer to accompany their frozen hMC5R membranes (Perkin Elmer catalog number RBXMC5M400UA).

[¹²⁵I] NDP-MSH: radiolabeled in house and purified by HPLC:

Na¹²⁵I (0.5 mCi, 17.4 Ci/mg) was added to 50 μL sodium phosphate (50 mM, pH 7.4) in an eppendorf tube precoated with IODOGEN. After incubation for 10 mins the phosphate buffer

containing the iodine was added to NDP-MSH (10 μ L at 1 mg/mL) in a separate eppendorf tube. This was incubated for a further 10 mins. The iodinated NDP-MSH was purified by HPLC on a Zorbax SB 300 column using solvent A: 0.05% TFA and solvent B: 90% acetonitrile 0.045% TFA with a linear gradient, 0-67% B over 60 mins. The ^{125} I NDP-MSH
5 eluted at 52 mins after the unlabeled starting material (48 min) and was counted and stored in the freezer. It was used within 48 hrs, as radioactive decay and ligand decomposition resulted in greatly reduced specific binding observed after 72 hrs.

Reagents:

10 Incubation buffer: 25 mM HEPES-KOH (pH 7.0), 1.5 mM CaCl_2 , 1 mM MgSO_4 , 0.1 M NaCl, 1 mM 1,10-phenanthroline, and 1 CompleteTM protease inhibitor tablet/100 mL (Roche, catalog number 1873580)

Perkin Elmer frozen hMC5 membranes: catalog number RBXMC5M400UA, 0.4 mL/vial; 400 microassays/vial, 0.78 mg/mL protein concentration

15 Vials of frozen membranes were thawed rapidly immediately before use, diluted with binding buffer and vortexed. Resuspended membranes were kept on ice until they were added to the wells of the plate.

Binding Protocol for 400 microassays per vial:

20 Assays were performed in 96 well polypropylene plates. Membranes (0.78 μ g 40 μ L of a 1:40 dilution in incubation buffer) were added to [^{125} I] NDP-MSH (0.84 nM; 2200 Ci/mmol) and test compounds in a total volume of 140 μ L. This was incubated for 1 hr at 37 °C. Non-specific binding was determined with 3 mM NDP-MSH. Plates were filtered using a Tomtec cell harvester with GF/A filters (Wallac) (presoaked in 0.6% polyethylenimine) and washed three
25 times with 1.0 mL ice-cold wash buffer (the above incubation buffer without 1,10-phenanthroline and CompleteTM protease inhibitor tablet). The filters were dried in a 37 °C oven, placed in a sample bag and 5 mL Betaplatescint (Wallac) was added. Prepared filters were counted in cassettes in a Microbeta Trilux (Wallac) for 1 min. Non-specific binding just under 5%. Data analysis was performed using GraphPad Prism 4, employing competition
30 binding with a single site model and a fixed Hill coefficient. The following equation was used: $Y = \text{Bottom} + (\text{Top}-\text{Bottom}) / (1 + 10^{(X - \log EC_{50})})$, where $X = \log(\text{concentration})$ and $Y = \text{binding}$ to fit the data.

Example 103 – Identification of Preferred Diastereomer for Binding to MC5R

35 The four diastereomers of one set of substituents were tested for binding in the hMC5R assay as in Examples 102, as listed in Table 3.

Table 3: Activity of Four Diastereomers

Cpd.	stereochemistry	human MC5R IC ₅₀ (nM)
102	(3R,5S)	3500
103	(3R,5R)	500
104	(3S,5R)	1500
105	(3S,5S)	56

As can be seen the 3S, 5S isomer is almost ten times more active than the next most active isomer and significantly more active than the other two possible isomers. This unexpectedly high level of differential activity and hence specificity of the (S,S) diastereoisomer was unexpected and is not predictable from a knowledge of the hMC5R or its previously known ligands.

Example 104 – Activity of Selected Compounds: hMC5R binding

Representative compounds of the present invention were tested for binding in the hMC5R assay as in Example 102, as listed in Table 4. The compounds were tested as their trifluoroacetate or hydrochloride salts, or as their free base.

Table 4: Properties of Compounds

x = <10 μ M; xx = < 1 μ M, xxx = < 100 nM, xxxx = < 10 nM

Cpd.	MS (M+1)	t _R (min)	MC5R radioligand IC ₅₀
14	556.2	5.74	xxx
25	595.3	6.22	xxx
31	539.3	5.92	xx
33	599.4	6.31	xxxx
37	473.4	5.59	xxx
38	541.3	5.78	xxx
39	541.3	5.67	xxx
49	499.3	5.77	xx
50	613.5	5.89	x
54	425.7	5.27	xx
60	629.4	6.27	x
62	629.3	6.22	xx
63	535.3	5.76	xx
64	603.3	6.04	xxx
65	577.2	5.97	xxx
67	591.3	5.94	xxxx
71	549.3	5.93	xx
79	606.4	6.033	x
81	743.4	5.489	xx

Cpd.	MS (M+1)	t_R (min)	MC5R radioligand IC ₅₀
83	650.3	6.524	XX
85	606.2	6.008	X
86	743.5	5.410	XX
87	650.4	6.424	X
102	577.4	5.775	X
103	577.5	5.750	XX
104	577.5	5.783	X
105	577.3	5.79	XXX
106	561.4	6.05	XX
107	524.3	5.63	XX
108	603.3	6.11	XXX
109	566.2	5.65	XX
110	591.2	5.82	XX
111	581.3	5.95	XXX
112	578.3	5.26	XXX
113	579.3	5.52	XX
114	578.3	5.72	XX
115	527.3	5.41	XX
116	578.3	5.78	XX
117	509.2	5.51	XX
118	523.3	5.56	X
119	523.2	5.51	X
120	512.3	5.10	X
121	549.4	5.96	XX
122	509.2	5.56	XX
123	523.4	5.63	X
124	509.2	5.41	X
125	523.3	5.68	X
126	549.3	5.79	XX
127	554.2	5.87	X
128	554.2	5.87	XX
129	539.1	5.58	X
130	596.5	5.87	X
131	582.4	5.88	X
132	567.4	5.62	X
133	567.4	5.62	X
134	582.4	5.88	XX
135	583.4	5.86	XX
136	595.4	5.31	XXX
137	595.4	5.87	XX
138	527.2	5.33	XX
139	533.3	5.54	X
140	620.2	6.16	XXX
141	620.2	6.21	XX
142	566.3	5.70	XXX
143	555.2	5.55	XX
144	555.2	5.74	XXX

Cpd.	MS (M+1)	t_R (min)	MC5R radioligand IC ₅₀
145	583.4	6.21	XX
146	587.2	4.90	X
147	547.4	5.78	XX
148	571.2	5.34	XX
149	567.1	4.48	X
150	568.1	4.87	X
151	519.5	5.23	X
152	595.4	5.92	XXX
153	553.5	5.58	XXX
154	595.4	5.95	XX
155	609.4	5.88	XX
156	607.5	5.96	XXX
157	609.4	-	X
158	607.4	5.88	XXX
159	579.3	5.83	XX
160	621.3	6.22	XXX
161	585.6	6.00	X
162	557.4	5.50	X
163	607.5	5.94	XX
164	607.2	5.69	XX
165	585.4	5.64	XX
166	557.3	6.06	XXX
167	559.5	5.47	XXX
168	585.5	5.58	XX
169	569.5	5.17	XX
170	583.6	5.70	XX
171	567.6	5.79	XXX
172	621.4	6.01	XX
173	571.5	5.65	XXX
174	567.5	5.50	XX
175	567.5	5.37	XX
176	625.5	5.81	XXX
177	587.4	5.65	XXX
178	584.5	4.84	XX
179	593.4	5.60	XX
180	583.6	5.41	XX
181	655.2	5.97	XXXX
182	501.4	5.20	XX
183	570.2	5.64	X
184	570.2	5.66	XX
185	583.5	5.43	XXX
186	607.3	5.28	XXX
187	583.4	5.37	XXX
188	595.6	5.64	XXX
189	587.4	5.78	XX
190	569.5	5.23	XX
191	567.7	5.92	XXX

Cpd.	MS (M+1)	t_R (min)	MC5R radioligand IC ₅₀
192	621.4	6.19	XX
193	569.6	5.23	XX
194	571.5	5.69	XXX
195	567.5	5.98	XXX
196	571.5	6.00	XX
197	561.3	5.84	XX
198	575.4	5.98	XX
199	571.1	5.69	XXX
200	621.3	6.19	XXX
201	591.2	6.02	XX
202	493.3	5.41	XX
203	545.2	5.91	XX
204	591.3	5.88	XXX
205	591.3	5.90	XX
206	479.4	5.09	XX
207	609.4	6.13	XX
208	589.3	5.69	XXX
209	605.3	5.85	XXX
210	631.4	6.09	XXXX
211	597.4	5.89	XXX
212	615.3	6.20	XXX
213	511.3	5.63	XX
214	545.4	5.92	XXX
215	637.6	6.15	XX
216	605.5	5.94	XXX
217	553.3	5.88	XXX
218	592.4	4.99	X
219	525.3	5.79	XXX
220	529.5	5.59	XXX
221	553.5	5.87	XXX
222	507.2	5.64	X
223	513.5	5.68	XX
224	553.3	5.89	XXX
225	549.7	5.87	XX
226	617.4	6.21	XXX
227	523.3	5.49	X
228	575.5	5.72	X
229	569.2	5.87	XX
230	569.2	5.83	X
231	611.2	6.20	XXXX
232	591.4	6.03	XXX
233	547.5	5.70	XXX
234	536.5	5.47	XX
235	561.7	6.11	XX
236	533.5	5.53	XX
237	585.5	6.23	XXX
238	550.5	5.81	XXX

Cpd.	MS (M+1)	t_R (min)	MC5R radioligand IC ₅₀
239	568.5	5.45	X
240	586.5	6.18	XXX
241	542.5	5.57	XX
242	568.4	5.91	XX
243	579.7	5.60	XX
244	565.5	5.42	X
245	506.4	5.73	XX
246	519.3	5.78	XX
247	637.5	5.84	X
248	624.3	6.28	XXX
249	603.2	6.14	XXX
250	589.4	6.04	XXX
251	543.3	6.30	XXX
252	487.2	5.13	X
253	613.5	6.06	XXX
254	537.4	5.66	X
255	531.6	5.65	XX
256	551.5	5.47	XX
257	543.5	5.77	X
258	543.5	5.66	XX
259	581.5	5.10	XX
260	591.4	5.94	XXX
261	599.4	5.99	XXX
262	613.5	6.08	XXX
263	521.4	5.85	XX
264	589.3	5.81	XXX
265	575.5	5.79	XXX
266	537.2	5.61	X
267	551.4	5.70	X
268	558.4	4.86	X
269	579.6	5.73	XXX
270	581.4	5.84	XXX
271	621.2	6.07	XXX
272	598.6	5.27	XX
273	626.7	5.64	X
274	507.3	6.35	XX
275	517.4	6.51	XXX
276	515.3	5.18	XXX
277	511.4	5.81	XXX
278	575.3	6.71	XXX
279	585.4	6.83	XXXX
280	539.3	5.87	XX
281	499.5	5.62	XX
282	497.6	5.58	XX
283	531.4	5.89	XXX
284	607.3	6.29	XXXX
285	567.3	5.99	XXX

Cpd.	MS (M+1)	t_R (min)	MC5R radioligand IC ₅₀
286	565.4	5.93	XXX
287	583.5	6.02	XXXX
288	579.6	6.18	XXX
289	515.3	5.58	XX
290	620.5	5.44	XX
291	620.5	5.38	X
292	531.5	5.22	XX
293	559.5	5.74	XX
294	607.4	6.06	XX
295	557.4	5.61	XX
296	534.4	7.27	XX
297	544.5	7.42	XX
298	559.4	6.59	XX
299	549.4	6.42	XX
300	567.3	6.52	XX
301	635.5	7.34	XX
302	593.5	6.02	XXX
303	573.4	5.47	XX
304	481.4	5.47	XX
305	521.5	6.10	XXX
306	641.4	7.38	XX
307	655.3	7.59	XX
308	485.3	5.17	X
309	553.3	5.82	XX
310	535.3	5.72	XX
311	553.5	5.39	X
312	621.2	6.06	XX
313	603.4	5.94	XX
314	572.3	4.91	X
315	574.5	7.69	XX
316	584.5	7.83	XX
317	617.7	7.04	XX
318	627.5	7.11	XXX
319	582.3	7.44	XX
320	592.4	7.55	XX
321	588.4	8.00	XX
322	598.4	8.15	XX
323	568.1	7.28	XX
324	578.3	7.45	XX
325	519.2	5.93	XX
326	511.2	5.94	XXX
327	540.3	5.61	XX
328	523.2	5.37	XX
329	498.4	5.49	XX
330	481.5	5.27	X
331	514.3	5.52	X
332	514.2	5.42	X

Cpd.	MS (M+1)	<i>t_R</i> (min)	MC5R radioligand IC₅₀
333	505.4	5.27	X
334	480.3	5.33	X
335	514.4	5.65	X
336	514.4	5.62	XX
337	514.3	5.63	X
338	477.3	5.35	X
339	531.5	5.65	X
340	480.4	5.38	X
341	487.2	5.55	XX
342	527.3	5.96	XX
343	587.2	6.33	XXX
344	567.3	6.12	XXX
345	577.2	6.31	XXXX
346	443.2	5.43	XX
347	435.3	5.46	XX
348	503.1	5.58	XX
349	511.4	5.73	XXX
350	493.7	5.71	XXX
351	427.3	5.04	X
352	423.4	5.28	X
353	427.3	5.13	X
354	423.3	5.32	X
355	489.4	6.42	XX
356	589.4	5.92	XX
357	581.3	6.03	X
358	480.4	5.33	X
359	555.4	5.81	XXX
360	612.6	5.49	X
361	620.5	-	X
362	595.4	6.12	XX
363	637.2	6.30	X
364	505.1	6.46	XX
365	485.2	6.26	XX
366	491.2	5.69	XX
367	573.1	6.07	XX
368	565.2	6.88	XXX
369	491.2	5.69	XX
370	483.4	5.77	XXX
371	559.1	6.12	XX
372	539.2	5.84	XX
373	473.3	5.29	XX
374	473.2	5.21	X
375	549.4	6.03	XX
376	621.4	6.13	XXX
377	493.3	5.32	XX
378	587.2	6.17	XXX
379	577.4	6.04	XXX

Cpd.	MS (M+1)	t_R (min)	MC5R radioligand IC ₅₀
380	559.3	6.01	XXX
381	551.3	5.99	XXX
382	551.4	6.04	XXXX
383	555.3	6.19	XXX
384	575.4	6.11	XXX
385	589.3	6.06	XXX
386	443.6	5.36	XX
387	459.9	5.66	XX
388	503.2	5.74	XX
389	511.4	5.65	XXX
390	527.3	5.95	XXX
391	573.2	6.07	XXX
392	483.3	5.97	XX
393	465.4	5.81	XX
394	475.3	5.98	XX
395	623.2	6.41	XXXX
396	607.4	6.17	XXXX
397	585.4	6.12	XXX
398	573.2	6.12	XXX
399	563.4	5.95	XX
400	581.3	6.12	XX
401	479.1	5.68	X
402	487.4	5.45	XX
403	551.5	6.42	XXX
404	525.3	6.31	XX
405	487.3	5.69	X
406	473.4	5.51	XX
407	487.4	5.40	X
408	568.1	5.91	X
409	601.3	6.08	XXX
410	533.3	5.82	X
411	622.3	6.90	X
412	579.3	6.60	X
413	555.4	6.00	XX
414	555.4	6.10	XXX
415	607.5	6.60	XXX
416	539.4	6.32	XX
417	593.5	6.26	XXX
418	525.3	6.03	XX
419	601.3	6.09	XXX
420	533.2	5.81	XX
421	627.4	7.20	X
422	601.3	6.10	XXX
423	609.4	7.64	XX
424	613.4	6.31	XXX
425	575.3	6.26	XXXX
426	537.4	5.98	XXX

Cpd.	MS (M+1)	t_R (min)	MC5R radioligand IC ₅₀
427	525.1	5.96	XXX
428	561.4	6.26	XXXX
429	559.1	6.11	XXX
430	545.1	6.01	XXX
431	569.3	5.92	XX
432	645.3	6.28	XXXX
433	640.2	6.70	XXX
434	622.3	6.49	XX
435	613.4	7.03	XX
436	603.2	7.23	XXX
437	587.2	7.01	X
438	561.4	6.88	XXX
439	545.1	6.68	XX
440	589.4	6.24	XXX
441	573.1	5.95	XXX
442	559.1	5.90	XX
443	627.4	6.56	XX
444	493.2	5.68	X
445	493.2	5.71	X
446	561.4	5.93	X
447	561.4	6.09	XX
448	581.3	6.93	XX
449	547.3	6.00	XXX
450	561.2	5.92	XXX
451	567.2	6.94	XX
452	581.2	6.45	XXX
453	593.3	6.56	XX
454	547.2	6.12	XXX
455	574.3	5.67	XXX
456	555.3	6.15	XXX
457	555.3	6.25	XXX
458	607.2	6.20	XXX
459	539.2	5.90	X
460	607.3	6.11	XX
461	539.1	5.94	X
462	577.1	6.37	XXXX
463	561.1	6.17	XXX
464	549.2	6.28	XXXX
465	507.2	6.00	XXX
466	583.3	6.38	XX
467	595.2	6.50	XXXX
468	560.3	5.64	XXX
469	575.2	6.62	XXX
470	575.2	6.27	XXXX
471	573.2	6.28	XXXX
472	547.2	5.96	XXX
473	617.2	6.52	XXX

Cpd.	MS (M+1)	t_R (min)	MC5R radioligand IC ₅₀
474	520.2	7.40	XX
475	534.3	7.66	XXX
476	587.4	6.55	XXX
477	575.2	5.82	XXX
478	603.4	5.98	XXX
479	573.1	6.32	XXX
480	573.1	6.18	XXXX
481	561.2	6.21	XXX
482	651.3	6.85	X
483	571.1	7.11	XX
484	583.3	6.98	XX
485	575.2	6.06	XXX
486	593.4	6.01	XXX
487	549.2	5.87	XXX
488	563.3	6.05	XXX
489	457.2	5.64	X
490	457.2	5.78	X
491	525.2	6.05	XXX
492	525.3	6.13	XXX
493	535.2	6.33	XX
494	535.2	6.50	X
495	491.3	5.63	XX
496	533.2	5.94	XXX
497	533.1	6.44	XXX
498	603.3	6.44	XXX
499	603.2	7.15	XXX
500	603.2	6.96	XXX
501	473.1	5.99	XX
502	473.2	6.09	XX
503	541.2	6.34	XXX
504	541.1	6.43	XXX
505	574.2	8.16	X
506	548.3	7.86	XXX
507	633.4	5.66	XX
508	591.1	6.36	XXXX
509	536.2	6.57	XXX
510	522.4	6.72	XXX
511	598.2	7.49	X
512	520.1	7.36	X
513	455.2	5.16	X
514	523.3	6.09	XX
515	471.1	5.53	XXXX
516	539.3	6.11	XXX
517	539.2	6.19	XXXX
518	519.3	5.72	XX
519	496.3	4.91	XXX
520	494.4	4.65	XX

Cpd.	MS (M+1)	t_R (min)	MC5R radioligand IC ₅₀
521	595.3	6.30	XXX
522	492.2	6.16	X
523	595.3	6.39	XX
524	545.2	5.91	XXX
525	523.0	5.88	XX
526	471.1	5.79	XX
527	539.3	6.14	XXX
528	573.1	6.50	X
529	564.5	6.60	XX
530	587.2	6.61	XX
531	601.3	7.03	X
532	589.3	6.59	XXX
533	592.3	7.08	XX
534	567.3	6.34	XXX
535	574.2	6.14	XXX
536	601.3	6.69	XXX
537	591.3	7.24	XX
538	535.3	6.94	XXX
539	535.3	6.37	XXX
540	573.1	6.13	XX
541	519.2	5.98	XX
542	587.1	6.34	XX
543	597.1	6.61	X
544	673.0	7.50	X
545	618.4	-	X
546	533.1	6.20	XXX
547	603.1	6.57	XXX
548	513.3	6.00	XXX
549	581.2	6.22	XXX
550	581.0	6.33	XXXX
551	473.3	6.03	XX
552	541.1	6.36	XXX
553	625.4	6.57	XXX
554	577.2	6.30	XXX
555	603.1	6.45	XXX
556	589.1	6.23	XXX
557	554.3	6.02	XX
558	587.1	6.50	X
559	457.0	5.73	X
560	525.2	6.20	XXX
561	637.0	6.75	XXX
562	613.3	7.03	XXX
563	607.2	-	XX
564	589.5	6.19	XXX
565	574.8	6.27	XXXX
566	618.9	6.74	XXX
567	587.7	6.19	XXXX

Cpd.	MS (M+1)	<i>t_R</i> (min)	MC5R radioligand IC ₅₀
568	596.9	6.20	XXX
569	578.0	6.38	XXXX
570	597.0	6.42	XXX
571	626.9	6.70	XXXX
572	591.9	6.61	XXXX
573	559.0	6.03	XXXX
574	612.8	7.04	XX
575	644.9	6.79	XXX
576	570.9	6.04	XXX
577	570.8	6.21	XXX
578	618.8	6.61	XXXX
579	579.9	6.39	XXXX
580	575.4	6.40	XX
581	575.2	6.26	XXX
582	575.2	6.11	XXX
583	575.1	6.21	XX
584	575.2	6.25	XX
585	575.4	6.27	X

Example 105 – MC5R Radioligand Binding Assay Using MC5 Receptors from Other Species

Radioligand binding and cAMP assays were also conducted using membranes and cells expressing MC5R cloned from other species (mouse MC5R membranes were obtained from Euroscreen; canine, rhesus monkey, cyno monkey and guinea pig MC5 receptors were cloned and expressed from cDNA libraries and transiently transfected as described in Examples 107 and 109. Plasma membranes from the cells were tested in the radioligand assay as in Example 102.

Example 106 – Activity of Selected Compounds: other species MC5R

Representative compounds of the present invention were tested for binding to MC5R from other species, as described in Example 105, the results are listed in Table 5.

Table 5: Binding of Selected Compounds to MC5R from Different Species

Cpd.	human MC5R (membrane) IC ₅₀ (nM)	human MC5R (whole cells) IC ₅₀ (nM)	mouse MC5R (membrane) IC ₅₀ (nM)	canine MC5R (whole cells) IC ₅₀ (nM)	rhesus monkey MC5R (membranes) IC ₅₀ (nM)	rhesus monkey MC5R (whole cells) IC ₅₀ (nM)
105	57	219	4000	6400	6027	3000
64	30	127	-	13000	7307	>5000

These results show the selectivity of the compounds of the invention for human MC5R in comparison to MC5R in other species. Whilst there is activity in other species it is significantly reduced in comparison to human MC5R, which would not be expected given the high receptor homology between species.

Example 107 – Human MC1R, MC3R and MC4R Radioligand Binding Assay

Radioligand binding assays were carried out using commercial or in-house prepared hMC1R, hMC3R and hMC4R membranes and [¹²⁵I] NDP-MSH, as per the hMC5R procedure in Example 102.

In-house plasma membranes were prepared from transfected mammalian cells (prepared as in Example 109, using plasmid DNA containing the human MC1R, MC3R or MC4R gene or other gene of interest in a plasmid vector with a mammalian origin of replication):

Adherent cells were washed with warm Hanks buffered saline solution (HBSS). 1 mL of cold HBSS was added to each flask and the cells were scraped off with a rubber policeman. The scraped cells were added to a 50 mL tube on ice. The plates were then rinsed twice with 5 mL cold HBSS and this was also added to the tube. The cells were centrifuged at 1000 x *g* for 5 mins in a bench top centrifuge and the supernatant was decanted. The remaining cell pellet was resuspended in 0.25 M sucrose. The cell suspension was centrifuged again as previously and the pellet resuspended in 5 mL of 0.25 M sucrose containing protease inhibitors. The cells were homogenised by a 10 second pulse with an Ika disperser followed by 30 seconds on ice. The homogenisation and ice incubation was repeated three times. The mixture was then centrifuged at 1260 x *g* for 5 mins. The supernatant was decanted into another centrifuge tube, to which a buffer containing 50 mM Tris, pH 7.4, 12.5 mM MgCl₂, 5 mM EGTA and protease inhibitors was added to make the volume up to 30 mL. This was centrifuged at 30,000 x *g* for 90 mins at 4°C. The resulting pellet was resuspended in 1 mL of the buffer above also containing 10% glycerol. Membranes were aliquoted into cryovials which were snap-frozen in a dry-ice/ethanol bath before being stored at -80°C until required for use.

Example 108 – Selectivity of Selected Compounds: hMCR binding

Representative compounds of the present invention were tested for binding in the hMC1R, hMC3R, hMC4R and hMC5R assays, as in Examples 102 and 107, as listed in Table 6.

Table 6: hMCR Binding Selectivity of Selected Compounds

Cpd.	human MC5R IC ₅₀ (nM)	human MC1R IC ₅₀ (nM)	human MC3R IC ₅₀ (nM)	human MC4R IC ₅₀ (nM)
105	57	6660	1750	3280
64	31	9220	2240	3490
33	9	2850	1500	6060
348	150	20000	1830	20000

These results demonstrate the selectivity of the compounds of the invention for human MC5R in comparison to other receptor subtypes of the human melanocortin receptor family..

5

Example 109 – Inhibition or stimulation of cAMP signal in cells expressing human MC5R

Transient transfection of mammalian cell lines:

10 The mammalian cell line, human embryonic kidney cells (HEK 293), were maintained in Dulbeccos Modified Eagle's medium (DMEM) with 5% fetal bovine serum, L- glutamine, high glucose and antibiotics/antimycotics. On the day prior to transfection, cells were passaged using trypsin/EDTA and seeded into 75 cm² flasks so that they would be approximately 90% confluent the next day. The next day, the cell media was replaced with
15 fresh antibiotic/antimycotic-containing DMEM. Approximately 100 µl of the transfection lipid Turbofectin 8.0 (Origene Technologies, MD, USA), was diluted in 1.0 mL of serum and antibiotic/antimycotic-free OptiMEM in a sterile 15 mL tube and incubated for 5 mins at room temperature. Following incubation, approximately 10-20 µg of plasmid DNA expressing the gene of interest (for example: pCMV6-XL4:Homo sapiens melanocortin 5 receptor (Origene
20 Technologies, MD, USA)) was diluted into the transfection mix and incubated for a further 30 mins at room temperature. The DNA/lipid solution was then added drop-wise to the media covering the cells while rocking the flask gently. 24 hrs post-transfection, the cells were passaged and seeded directly into two, 75cm² flasks and left to recover. 48 hrs post transfection, cells were harvested for use in assays with cell dissociation solution.

25

Cyclic-Adenosine Monophosphate [cAMP] stimulation assay:

HEK 293 cells transiently expressing the melanocortin MC5 receptor were suspended in stimulation buffer (Hanks buffered saline solution (HBSS), 0.1 % bovine serum albumin, protease inhibitors and 0.5 mM 3-Isobutyl-1-methylxanthine) at 4 x 10⁶ cells/mL. 5 µl of cells,
30 plus the compounds/peptides as described below, were added to wells of a 384-well plate as soon as possible after resuspension.

To detect antagonist activity, test compounds at varying concentrations were diluted in stimulation buffer at four times concentrate and 2.5 µl was added to wells containing cells. 2.5 µl of a four times required concentration of NDP-MSH or alpha-MSH was added to all wells containing compounds. Negative control wells contained two times concentrated NDP-MSH or alpha-MSH alone without compound.

To detect agonist activity, test compounds at varying concentrations were diluted in stimulation buffer at two times concentrate and 5 µl was added to wells containing cells. Positive control wells contained NDP-MSH or alpha-MSH alone (no compound) at two times concentrate

Basal level (of cAMP) control wells contained stimulation buffer only (no agonist or compounds). Known concentrations of cAMP (standards) in stimulation buffer were included on the plate, but no cells were added to these wells. The plate was then incubated for 30 mins at 37 °C with gentle shaking. After incubation, 10 µl of lysis buffer (10 % Tween 20, 1 M HEPES, 0.1% BSA, protease inhibitors, ddH₂O) was added to all wells to be measured. Detection of cAMP was then achieved using the Alphascreen cAMP kit (Perkin Elmer, USA), briefly described as follows. A dilution of 10 µl acceptor beads/mL of lysis buffer was prepared in low light conditions. 5 µl of diluted acceptor beads were added to each well to be measured, then the plate was incubated for 30 mins at room temperature, in the dark, with gentle shaking. In low light conditions, donor beads were diluted at 10 µl/mL of lysis buffer, to which 0.75 µl biotinylated cAMP/mL of lysis buffer was added. This mixture was allowed to incubate for 30 mins at room temperature (in the dark) before proceeding with the assay. Following incubation, 5 µl/mL of biotinylated cAMP/Donor bead mix were added per well in low light conditions and the plate was incubated in the dark, at room temperature, for a further hr. Plates were read on an Envision plate reader (Perkin Elmer) after 1 hr and ~16hrs incubation. cAMP concentration in the cells was determined by the use of a 'standard curve' generated from the output of known cAMP concentrations as described below.

Each assay plate contained a "standard curve" of known concentrations of cAMP, in 10 fold dilutions. This is an essential part of the assay as there is high inter-plate variability. The plates were read on an Envision multilabel plate reader fitted with Alphascreen technology and the raw data was imported into GraphPad Prism 4 software (GraphPad, USA) for analysis. A curve was fitted to the known concentrations using non-linear regression, specifically using a sigmoidal dose-response equation ($Y = \text{Bottom} + (\text{Bottom} + (\text{Top} - \text{Bottom}) / (1 + 10^{\log \text{EC}_{50} - X}))$), where the equation shows the response as a function of the logarithm

of concentration. X is the logarithm of peptide/compound concentration and Y is the response. Also considered in this equation are bottom plateau, top plateau of the curve and EC₅₀ (effective concentration, 50%)

5 **Example 110 – Activity of Selected Compounds: hMC5R**

Representative compounds of the present invention were tested for agonism or antagonism of the hMC5R, as in Example 109, results are listed in Table 7.

Table 7: Agonism and Antagonism of hMC5 by Selected Compounds

Cpd.	human MC5R EC ₅₀ (cAMP, agonism) (nM)	human MC5R IC ₅₀ (cAMP, antagonism of 10 ⁻⁶ M alpha-MSH) (nM)
105	>10000	400
64	>10000	70
33	>10000	190
348	>10000	94

References

- Andersen, G.N.; Hägglund, M.; Nagaeva, O.; Frängsmyr, L.; Petrovska, R.; Mincheva-
Nilsson, L.; Wikberg, J.E.S. *Scand. J. Immunol.* **2005**, *61*, 279-284 "Quantitative
measurement of the levels of melanocortin receptor subtype 1, 2, 3 and 5 and pro-opio-
melanocortin peptide gene expression in subtypes of human peripheral blood leukocytes"
- Barrett, P.; MacDonald, A.; Helliwell, R.; Davidson, G.; Morgan, P. *J. Molec. Endocrin.* **1994**,
12, 203-213 "Cloning and expression of a new member of the melanocyte-stimulating
hormone receptor family"
- Bataille, V.; Snieder, H.; MacGregor, A.J.; Sasieni, P.; Spector, T.D. *J. Invest. Dermatol.*
2002, *119*, 1317-1322 "The Influence of Genetics and Environmental Factors in the
Pathogenesis of Acne: A Twin Study of Acne in Women"
- Bhardwaj, S.S.; Rohrer, T.E.; Arndt, K.A. *Semin. Cutan. Med. Surg.* **2005**, *24*, 107-112
"Lasers and light therapy for acne vulgaris"
- Bohm, M.; Luger, T.A.; Tobin, D.J.; Garcia-Borrón, J.C. *J. Invest. Dermatol.* **2006**, *126*, 1966-
1975 "Melanocortin Receptor Ligands: New Horizons for Skin Biology and Clinical
Dermatology"
- Buggy, J.J. *Biochem J.* **1998**, *331*, 211-216 "Binding of α -melanocyte-stimulating hormone to
its G-protein-coupled receptor on B-lymphocytes activates the Jak/STAT pathway"
- Burke, B.M.; Cunliffe, W.J.; *Br. J. Dermatol.* **1984**, *112* 124-126 "Oral spironolactone therapy
for female patients with acne, hirsutism or androgenic alopecia"

Caldwell, H.K.; Lepri, J.J.. *Chem. Senses* **2002**, 27, 91-94 "Disruption of the fifth melanocortin receptor alters the urinary excretion of aggression-modifying pheromones in male house mice"

Cerdá-Reverter, J.M.; Ling, M.K.; Schiöth, H.B.; Peter, R.E. *J. Neurochem.* **2003**, 1354-1367
 5 "Molecular cloning, characterization and brain mapping of the melanocortin 5 receptor in goldfish"

Chen, W.; Kelly, M.A.; Opitz-Araya, X.; Thomas, R.E.; Low, M.J.; Cone, R.D. *Cell*, **1997**, 91, 789-798 "Exocrine gland dysfunction in MC5-R-deficient mice: evidence for coordinated regulation of exocrine gland function by melanocortin peptides"

10 Chhajlani, V.; Muceniece, R.; Wikberg, J.E.S. *BBRC* **1993**, 195, 866-873 "Molecular Cloning of a Novel Human Melanocortin Receptor"

Clarke, S.B.; Nelson, A.M.; George, R.E.; Thiboutot, D.M. *Dermatol. Clin.* **2007**, 25, 137-146
 "Pharmacologic Modulation of Sebaceous Gland Activity: Mechanisms and Clinical Applications".

15 Cordain, L. *Sem. Cut. Med Surg.* **2005**, 24, 84-91 "Implications for the Role of Diet in Acne"

Cotterill, J.A.; Cunliffe, W.J.; Williamson, B. *Brit. J. Dermatol.* **1971**, 85, 93-94 "Severity of Acne and Sebum Excretion Rate"

Danby, F.W. *J. Am. Acad. Dermatol.* **2005**, 52, 1071-1072 "Why we have sebaceous glands"

Eisinger, M.; Fitzpatrick, L.J.; Lee, D.H.; Pan, K.; Plata-Salaman, C.; Reitz, A.B.; Smith-
 20 Swintosky, V.L.; Zhao, B. WO03/040117 15 May 2003a "Novel 1,2,4-thiadiazole derivatives as melanocortin receptor modulators"

Eisinger, M.; Fitzpatrick, L.J.; Lee, D.H.; Pan, K.; Plata-Salaman, C.; Reitz, A.B.; Smith-
 Swintosky, V.L.; Zhao, B. WO03040118A1 15 May 2003b "Novel 1,2,4-thiadiazolium derivatives as melanocortin receptor modulators"

25 Eisinger, M.; Fitzpatrick, L.J.; Lee, D.H.; Pan, K.; Plata-Salaman, C.; Reitz, A.B.; Smith-
 Swintosky, V.L.; Zhao, B. US2003/0162819A1 Aug 28 2003c "Novel 1,2,4-thiadiazolium derivatives as melanocortin receptor modulators"

Eisinger, M.; Fitzpatrick, L.J.; Lee, D.H.; Pan, K.; Plata-Salaman, C.; Reitz, A.B.; Smith-
 Swintosky, V.L.; Zhao, B. US2003/0176425A1 Sep 18 2003d "Novel 1,2,4-thiadiazole
 30 derivatives as melanocortin receptor modulators"

Eisinger, M.; Fitzpatrick, L.J.; Lee, D.H.; Pan, K.; Plata-Salaman, C.; Reitz, A.B.; Smith-
 Swintosky, V.L.; Zhao, B. US2006/0030604A1 Feb 9 2006a "Novel 1,2,4-thiadiazolium derivatives as melanocortin receptor modulators"

Eisinger, M.; Fitzpatrick, L.J.; Lee, D.H.; Pan, K.; Plata-Salaman, C.; Reitz, A.B.; Smith-
 35 Swintosky, V.L.; Zhao, B. US2006/0128772A1 Jun 15 2006b "Novel 1,2,4-thiadiazole derivatives as melanocortin receptor modulators"

Fathi, Z.; Iben, L.G.; Parker, E.M. *Neurochemical Res.* **1995**, *20*, 107-113 "Cloning, Expression, and Tissue Distribution of a Fifth Melanocortin Receptor Subtype"

Follador, I.; Campelo, L. *Expert Rev. Dermatol.* **2006**, *1* 181-184 "Impact of acne on quality of life"

5 Fong, T.M.; Van der Ploeg, L.H.T.; Huang, R.-R.C. US6645738B1 Nov 11 2003 "DNA molecules encoding the melanocortin 5 receptor protein from rhesus monkey"

Gantz, I.; Shimoto, Y.; Konda, Y.; Miwa, H.; Dickinson, C.J.; Yamada, T. *BBRC* **1994**, *200*, 1214-1220 "Molecular cloning, expression and characterization of a fifth melanocortin receptor"

10 Goldstein, J.A.; Socha-Szott, A.; Thomsen, R.J.; Pochi, P.E.; Shalita, A.R.; Strauss, J.S. *Am. J. Dermatol.* **1982**, *6*, 760-765 "Comparative effect of isotretinoin and etretinate on acne and sebaceous gland secretion"

Goodfellow, A.; Alaghband-Zadeh, J.; Carter, G.; Cream, J.J.; Holland, S.; Scully, J.; Wise, P. *Brit. J. Dermatol.* **1984**, *111*, 209-214 "Oral spironolactone improves acne vulgaris and
15 reduces sebum excretion"

Goulden, V.; Mcgeown, C.H.; Cunliffe, W.J. *Brit. J. Dermatol.* **1999**, *141*, 297-300 "Familial Risk of Adult Acne: A comparison between first-degree relatives of affected and unaffected individuals"

Graefe, T.; Wollina, U.; Schulz, H.-J.; Burgdorf, W. *Dermatology* **2000**, *200*, 331-333 "Muir-Torre Syndrome - Treatment with Isotretinoin and Interferon Alpha-2a Can Prevent
20 Tumour Development"

Griffon, N.; Mignon, V.; Facchinetti, P.; Diaz, J.; Schwartz, J.-C.; Sokoloff, P. *BBRC* **1994**, *200*, 1007-1014 "Molecular cloning and characterization of the rat fifth melanocortin receptor"

25 Gupta, A.K.; Bluhm, R. *Journal of the European Academy of Dermatology and Venereology* **2004** *18*:1 13 "Seborrheic dermatitis"

Haitina, T.; Klovins, J.; Andersson, J.; Fredriksson, R.; Lagerström, M.C.; Larhammar, D.; Larson, E.T.; Schiöth, H.B. *Biochem. J.* **2004**, *380*, 475-486 "Cloning, tissue distribution, pharmacology and three-dimensional modelling of melanocortin receptors 4 and 5 in
30 rainbow trout suggest close evolutionary relationships of these subtypes"

Harper, J.C. *Semin. Cutan. Med. Surg.* **2005**, *24*, 103-106 "Hormonal Therapy for Acne using oral contraceptive pills"

Harris, H.H.; Downing, D.T.; Stewart, M.E.; Strauss, J.S. *J. Am. Acad. Dermatol.* **1983**, *8*, 200-203 "Sustainable rates of sebum secretion in acne patients and matched normal
35 controls"

Hatta, N.; Dixon, C.; Ray, A.J.; Phillips, S.R.; Cunliffe, W.J.; Dale, M.; Todd, C.; Meggit, S.; Birch-Machin, M.A.; Rees, J.L. *J. Invest. Dermatol.* **2001**, *116*, 564-570 "Expression, candidate gene, and population studies of the melanocortin 5 receptor"

Houseknecht, K.L.; Robertson, A.S.; Xiao, X. US2003/0110518A1 Jun 12 2003

"Melanocortin-5 receptor sequences and uses thereof"

Huang, R.-R.C.; Singh, G.; Van der Ploeg, L.H.T.; Fong, T.M. *J. Receptor & Signal Transduction Res.* **2000**, *20*, 47-59 "Species-dependent pharmacological properties of the melanocortin-5 receptor"

Ide, F.; Shimoyama, T.; Horie, N.; Kaneko, T.; Matsumoto, M. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol. Endod.* **1999**, *87*, 721-724 "Benign lymphoepithelial lesion of the parotid gland with sebaceous differentiation"

Jeong, S.K.; Hwang, S.W.; Choi, S.Y.; An, J.M.; Seo, J.T.; Zouboulis, C.C.; Lee, S.H. *J. Investigative Dermatol.* **2007**, *127*, pS72 "Intracellular calcium mobilization is mediated by the melanocortin receptors in SZ95 sebocytes" (Abstract 431, Society for Investigative Dermatology, May 2007, Los Angeles CA)

Jih, M.H.; Friedman, P.M.; Goldberg, L.H.; Robles, M.; Glaich, A.S.; Kimyai-Asadi, A. *J. Am. Acad. Dermatol.* **2006**, *55*, 80-87 "The 1450-nm diode laser for facial inflammatory acne vulgaris: Dose-response and 12-month follow-up study".

Jones, D.H.; King, K.; Miller, A.J.; Cunliffe, W.J. *Brit. J. Dermatol.* **1983**, *108*, 333-343 "A dose-response study of 13-cis-retinoic acid in acne vulgaris"

Kim, K.S.; Marklund, S.; Rothschild, M.F. *Animal Genetics* **2000**, *31*, 230-231. "The porcine melanocortin-5-receptor (MC5R) gene: polymorphisms, linkage and physical mapping"

King, K.; Jones, D.H.; Daltrey, D.C.; Cunliffe, W.J. *Brit. J. Dermatol.* **1982**, *107*, 583-590 "A double-blind study of the effects of 13-cis-retinoic acid on acne, sebum excretion rate and microbial population"

Kligman, A.M. *Brit. J. Dermatol.* **1963**, *75*, 307-319 "The uses of sebum"

Klovins, J.; Haitina, T.; Ringholm, A.; Löwgren, M.; Fridmanis, D.; Slaidina, M.; Stier, S.; Schiöth, H.B. *Eur. J. Biochem.* **2004**, *271*, 4320-4331 "Cloning of two melanocortin (MC) receptors in spiny dogfish"

Kruse, R.; Rütten, A.; Schweiger, N.; Jakob, E.; Mathiak, M.; Propping, P.; Mangold, E.; Bisceglia, M.; Ruzicka, T. *J. Invest. Dermatol.* **2003**, *120*, 858-864 "Frequency of Microsatellite Instability in Unselected Sebaceous Gland Neoplasias and Hyperplasias"

Labbé, O.; Desarnaud, F.; Eggerickx, D.; Vassart, G.; Parmentier, M. *Biochem.* **1994**, *33*, 4543-4549 "Molecular Cloning of a mouse melanocortin 5 receptor gene widely expressed in peripheral tissues"

- Ling, M.K.; Hotta, E.; Kilianova, Z.; Haitina, T.; Ringholm, A.; Johansson, L.; Gallo-Payet, N.; Takeuchi, S.; Schiöth, H.B. *Brit. J. Pharmacol.* **2004**, *143*, 626-637 "The melanocortin receptor subtypes in chicken have high preference to ACTH-derived peptides"
- Makrantonaki, E.; Zouboulis, C.C. *Brit. J. Dermatol.* **2007**, *156*, 428-432 "Testosterone metabolism to 5 α -dihydrotestosterone and synthesis of sebaceous lipids is regulated by the peroxisome proliferator-activated receptor ligand linoleic acid in human sebocytes"
- Mariappan, M.R.; Fadare, O.; Jain, D. *Arch. Pathol. Lab. Med.* **2004**, *128*, 245-246 "Sebaceous Differentiation in Salivary Glands"
- Mallon, E.; Newton, J.N.; Klassen, A.; Stewart-Brown, S.L.; Ryan, T.J.; Finlay, A.Y. *Brit. J. Dermatol.* **1999**, *140*, 672-676 "The quality of life in acne: a comparison with general medical conditions using generic questionnaires"
- Marqueling A.L.; Zane, L.T. *Semin. Cutan. Med. Surg.* **2005**, *24*, 92-102 "Depression and Suicidal Behavior in Acne Patients Treated with Isotretinoin: A Systematic Review"
- Morgan, C.; Thomas, R.E.; Ma, W.; Novotny, M.V.; Cone, R.D. *Chem. Senses* **2004a**, *29*, 111-115 "Melanocortin-5 receptor deficiency reduces a pheromonal signal for aggression in male mice"
- Morgan, C.; Thomas, R.E.; Cone, R.D. *Horm. Behav.* **2004b**, *45*, 58-83 "Melanocortin-5 receptor deficiency promotes defensive behaviour in male mice"
- Morgan, C.; Cone, R.D. *Behaviour Genetics* **2006**, *36*, 291-300 "Melanocortin-5 receptor deficiency in mice blocks a novel pathway influencing pheromone-induced aggression"
- Mourelatos, K.; Eady, E.A.; Cunliffe, W.J.; Clark, S.M.; Cove, J.H. *Brit. J. Dermatol.* **2007**, *156*, 22-31 "Temporal changes in sebum excretion and propionibacterial colonization in preadolescent children with and without acne"
- Nelson, A.M.; Gilliland, K.L.; Cong, Z.; Thiboutot, D.M. *J. Investigative Dermatol.* **2006**, *126*, 2178-2189 "13-*cis*-Retinoic Acid Induces Apoptosis and Cell Cycle Arrest in Human SEB-1 Sebocytes"
- Phan, J.; Kanchanapoomi, M.; Liu, P.; Jalian, H.; Gilliland, K.; Nelson, A.; Thiboutot, D.; Kim, J. *J. Investigative Dermatol.* **2007**, *127*, pS126 "P. acnes induces inflammation via TLR2 and upregulates antimicrobial activity in sebocytes" (Abstract 754, Society for Investigative Dermatology, May 2007, Los Angeles CA)
- Piérard, G.E.; Piérard-Franchimont, T.L. *Dermatologica* **1987**, *175*, 5-9 "Seborrhoea in Acne-Prone and Acne-Free Patients"
- Plewig G, Jansen T. Seborrheic dermatitis. In: Freedberg IM, Eisen AZ, Wolff K, Austen KF, Goldsmith LA, Katz SI, Fitzpatrick TB, (Eds). *Dermatology in General Medicine*, 5th ed. New York: McGraw Hill, 1999: 1482-1489

- Pochi, P.E.; Strauss, J.S. *J. Invest. Dermatol.* **1964**, *43*, 383-388 "Sebum production, casual sebum levels, titratable acidity of sebum and urinary fractional 17-ketosteroid excretion in males with acne"
- Porter, A.M.W. *J. Royal Soc. Med.* **2001**, *94*, 236-237 "Why do we have apocrine and sebaceous glands"
- Ringholm, A.; Fredriksson, R.; Poliakova, N.; Yan, Y.-L.; Postlethwait, J.H.; Larhammar, D.; Schiöth, H.B. *J. Neurochem.* **2002**, *82*, 6-18 "One melanocortin 4 and two melanocortin 5 receptors from zebrafish show remarkable conservation in structure and pharmacology"
- Simpson, N.B. and Cunliffe, W.J. in Rooks' Textbook of Dermatology, 7th Ed **2004** Blackwell Science, Malden Mass, p 43.1-43.75 "Chapter 43. Disorders of the Sebaceous Glands"
- Smith, K.R.; Nelson, A.; Cong, Z.; Thiboutot, D. *J. Investigative Dermatol.* **2007a**, *127*, pS68 "Iron status affects human sebocyte survival" (Abstract 408, Society for Investigative Dermatology, May 2007, Los Angeles CA)
- Smith, R.N.; Mann, N.J.; Braue, A.; Makelainen, H.; Varigos, G.A. *J. Am. Acad. Dermatol.* **2007b**, *57*, 247-256 "The effect of a high-protein, low glycemic-load diet versus a conventional, high glycemic-load diet on biochemical parameters associated with acne vulgaris: A randomized investigator-masked, controlled trial"
- Shuster, S. *Lancet* **1976**, *7973*, 1328-1329 "Biological purpose of acne"
- Taylor, A.; Namba, K. *Immunology Cell Biol.* **2001**, *79*, 358-367 "*In vitro* induction of CD25+CD4+ regulatory T cells by the neuropeptide alpha-melanocyte stimulating hormone (α -MSH)"
- Thiboutot, D.; Sivarajah, A.; Gilliland, K.; Cong, Z.; Clawson, G. *J. Invest. Dermatol.* **2000**, *115*, 614-619 "The melanocortin 5 receptor is expressed in human sebaceous glands and rat preputial cells"
- Thody, A.J.; Shuster, S. *Nature* **1973**, *245*, 207-209 "Possible role of MSH in the mammal"
- Thody, A.J.; Cooper, M.F.; Bowden, P.E.; Shuster, S. *J. Endocrinol.* **1975a**, *67*, 18P-19P "The sebaceous gland response to α -melanocyte-stimulating hormone and testosterone"
- Thody, A.J.; Shuster, S. *J. Endocrinol.* **1975b**, *64*, 503-510 "Control of sebaceous gland function in the rat by α -melanocyte-stimulating hormone"
- Thody, A.J.; Goolamali, S.K.; Burton, J.L.; Plummer, N.A.; Shuster, S. *Brit. J. Dermatol.* **1975c**, *92*, 43-47 "Plasma β -MSH levels in acne vulgaris"
- Wikberg, J.E.S. *Exp. Opin. Ther. Patents* **2001**, *11*, 61-76 "Melanocortin receptors: new opportunities in drug discovery";
- Wikberg, J.; Chhajlani, V. US6448032B1 Sep 10 2002 "Human melanocyte stimulating hormone receptor polypeptide and DNA"


- Williams, C; Layton, A.M. *Exp. Rev. Dermatol.* **2006**, 1, 429-438 "Treatment of Acne: an update"
- Yamada, T.; Gantz, I. US5622860, Apr. 22 1997, "Genes Encoding Melanocortin Receptors"
- Yaswen, L.; Diehl, N.; Brennan, M. B.; Hochgeschwender, U. *Nature Med.* **1999**, 5, 1066-1070 "Obesity on the mouse model of proo-opiomelanocortin deficiency responds to peripheral melanocortin"
- Youn, S.-W.; Park, E.-S.; Lee, D.-H.; Huh, C.-H.; Park, K.-C. *Brit. J. Dermatol.* **2005**, 153, 919-924 "Does facial sebum secretion really affect the development of acne?"
- Zhang, L.; Anthonavage, M.; Huang, Q.; Li, W.-H.; Eisinger, M. *Ann. N. Y. Acad. Sci.* **2003**, 994, 154-161 " Proopiomelanocortin peptides and sebogenesis"
- Zhang, L.; Li, W.-H.; Anthonavage, M.; Eisinger, M. *Peptides* **2006**, 27, 413-420 "Melanocortin-5 receptor: a marker of human sebocyte differentiation"
- Zouboulis, CC; Bohm, M. *Exp. Dermatol.* **2004**, 13, 31-35 "Neurocrine regulation of sebocytes - a pathogenetic link between stress and acne"

The details of specific embodiments described in this invention are not to be construed as limitations. Various equivalents and modifications may be made without departing from the scope of this invention, and it is understood that such equivalent embodiments are part of this invention.

What is claimed is:

1. A method of down-regulating the activity of MC5R or a fragment thereof *in vitro* comprising exposing the MC5R or a fragment thereof to a compound of the formula (I):

5



Formula (I)

wherein

- 10 Y is a group of formula $-(CR^9R^{10})_n-$;

X is selected from the group consisting $-C(=O)-$, $-OC(=O)-$, $-NHC(=O)-$, $-(CR^{11}R^{12})_s$, and $-S(=O)_2-$;

- 15 R is a group of the formula $-Z-R^4$ wherein

Z is a group of formula $-(CR^{13}R^{14})_q-$;

- 20 R^4 is selected from the group consisting of H, C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_6 - C_{18} aryl, optionally substituted C_1 - C_{18} heteroaryl, $NR^{4a}R^{4b}$, $C(=O)R^{15}$, $C(=O)NR^{16}R^{17}$, $-C(=NR^{16})NR^{17}R^{18}$, SR^{20} , $SC(=O)R^{20}$, SO_2R^{20} , OR^{20} , $ONR^{16}R^{17}$, $OCR^{17}R^{18}R^{20}$, $OC(=O)R^{20}$, $OC(=O)OR^{20}$, $OC(=O)NR^{16}R^{17}$, and $ONR^{16}C(=NR^{17})NR^{18}R^{19}$;

- 25 R^{4a} is selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, optionally substituted C_1 -

C_{18} heteroaryl, $C(=O)R^{15a}$, $C(=O)NR^{15a}R^{16a}$, $C(=O)OR^{15a}$, SO_2R^{15a} , $C(=O)H$, $-C(=NR^{15a})-$, $NR^{16a}R^{17a}$, and OR^{15a} ,

R^{4b} is selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, optionally substituted C_1 - C_{18} heteroaryl, $C(=O)R^{15a}$, $C(=O)NR^{15a}R^{16a}$, $C(=O)OR^{15a}$, or

R^{4a} and R^{4b} when taken together with the nitrogen atom to which they are attached form an optionally substituted heterocyclic moiety, or

one of R^{4a} and R^{4b} when taken together with any R^{13} or R^{14} and the atoms to which they are attached forms an optionally substituted heterocyclic moiety;

R^{13} and R^{14} are each independently selected from the group consisting of H, halogen, OH, C_1 - C_{12} alkyl, C_6 - C_{18} aryl, C_1 - C_{12} hydroxyalkyl, C_1 - C_{12} haloalkyl, C_1 - C_{12} alkyloxy and C_1 - C_{12} haloalkyloxy, or

when taken together with the carbon to which they are attached R^{13} and R^{14} form an optionally substituted C_3 - C_{12} cycloalkyl, or an optionally substituted C_1 - C_{12} heterocycloalkyl group, or

one of R^{13} and R^{14} when taken together with one of R^{4a} , and R^{4b} and the atoms to which they are attached form an optionally substituted heterocyclic moiety, or

one of R^{13} and R^{14} when taken together with one of R^{15} , R^{16} , R^{17} , R^{18} , R^{19} or R^{20} and the atoms to which they are attached form an optionally substituted cyclic moiety;

each R^{15} , R^{15a} , R^{16} , R^{16a} , R^{17} , R^{17a} , R^{18} , R^{19} and R^{20} is independently selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl, or

any two of R^{15} , R^{15a} , R^{16} , R^{16a} , R^{17} , R^{17a} , R^{18} , R^{19} and R^{20} when taken together with the atoms to which they are attached form an optionally substituted cyclic group, or

one of R^{15} , R^{16} , R^{17} , R^{18} , R^{19} and R^{20} when taken together with one of R^{13} and R^{14}
 5 and the atoms to which they are attached form an optionally substituted cyclic moiety;

q is an integer selected from the group consisting of 0, 1, 2, 3, 4, and 5;

R^1 is selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl,
 10 optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl;

15 R^2 and R^3 are each independently selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl;

20 each R^{5a} and R^{5b} are independently selected from the group consisting of H, halogen, C_1 - C_{12} alkyl, C_1 - C_{12} hydroxyalkyl and C_1 - C_{12} haloalkyl, or

one or more of R^{5a} and R^{5b} when taken together with one or more of R^6 , R^7 and R^8
 25 and the atoms to which they are attached form a moiety selected from the group consisting of an optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl;

R^6 , R^7 and R^8 are each independently selected from the group consisting of H,
 30 halogen, hydroxy, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_1 - C_{10} heteroalkenyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, optionally substituted C_1 - C_{18} heteroaryl, optionally substituted amino, optionally substituted carboxy, C_1 -
 35 C_{12} alkyloxy, and optionally substituted thio, or

(a) when taken together with the carbon atom to which they are attached two or more of R^6 , R^7 and R^8 form a moiety selected from the group consisting of optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 -
5 C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl, or

(b) one or more of R^6 , R^7 and R^8 when taken together with one or more of R^{5a} and R^{5b} and the atoms to which they are attached form a moiety selected from the group
10 consisting of an optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl;

each R^9 and R^{10} is independently selected from the group consisting of H, optionally
15 substituted C_1 - C_{12} alkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl;

each R^{11} and R^{12} is independently selected from the group consisting of H, and
optionally substituted C_1 - C_{12} alkyl;


20 n is an integer selected from the group consisting of 1, 2, 3 and 4;

r is an integer selected from the group consisting of 1, 2, 3, and 4;

25 s is an integer selected from the group consisting of 1, 2, 3, and 4;

or a pharmaceutically acceptable salt or prodrug thereof.

2. A method according to claim 1 wherein the compound of formula (I) is a compound of
30 the formula (Ia):



wherein R, R¹, R², R³, R^{5a}, R^{5b}, R⁶, R⁷, R⁸, X, Y and r are as defined in claim 1 and the stereochemistry is as shown, or a pharmaceutically acceptable salt or prodrug thereof.

3. A method according to claim 1 wherein in the compound of formula (I) R⁴ = NR^{4a}R^{4b}.

4. A method according to claim 3 wherein in the compound of formula (I) R^{4a} is selected from the group consisting of H, -C(=NH)NH₂, -C(=NH)N(CH₃)₂, -C(=NH)NHCH(CH₃)₂, -C(=O)CH₃, -C(=O)cyclohexyl, CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂, CH₂CH₂CH₂CH₃, -CH(CH₃)CH₂CH₃, CH₂CH(CH₃)₂, C(CH₃)₃, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, benzyl, and phenyl, or a halogenated derivative thereof.

5. A method according to claim 3 wherein in the compound of formula (I) R^{4b} is selected from the group consisting of H, CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂, CH₂CH₂CH₂CH₃, CH(CH₃)CH₂CH₃, CH₂CH(CH₃)₂, C(CH₃)₃, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, benzyl, and phenyl, or a halogenated derivative thereof.

6. A method according to claim 3 wherein in the compound of formula (I) R^{4a} and R^{4b} when taken together with the nitrogen atom to which they are attached form an optionally substituted C₂-C₁₂ heterocycloalkyl group, an optionally substituted C₂-C₁₂ heterocycloalkenyl group or an optionally substituted C₁-C₁₈ heteroaryl group.

7. A method according to claim 6 wherein in the compound of formula (I) R^{4a} and R^{4b} when taken together with the nitrogen atom to which they are attached form an optionally substituted heterocycloalkyl group selected from the group consisting of piperidin-1-yl, pyrrolidin-1-yl, azetidin-1-yl, azepan-1-yl, piperazin-1-yl, and morpholin-4-yl.

8. A method according to claim 3 wherein in the compound of formula (I) one of R^{4a} and R^{4b} when taken together with the nitrogen atom to which it is attached and one of R^{13} and R^{14} and the carbon atom to which it is attached form an optionally substituted C_2 - C_{12} heterocycloalkyl group.

5

9. A method according to claim 8 wherein in the compound of formula (I) one of R^{4a} and R^{4b} when taken together with the nitrogen atom to which it is attached and one of R^{13} and R^{14} and the carbon atom to which it is attached form an optionally substituted heterocycloalkyl group selected from the group consisting of piperidinyl, pyrrolidinyl, azetidiny, morpholinyl, piperazinyl, and azepanyl.

10

10. A method according to claim 1 wherein in the compound of formula (I) Z is $-(CH_2)_q-$.

11. A method according to claim 10 wherein in the compound of formula (I), q is selected from the group consisting of 1, 2, 3 and 4.

15

12. A method according to claim 10 wherein in the compound of formula (I) q is 1.

13. A method according to claim 10 wherein in the compound of formula (I) q is 2.

20

14. A method according to claim 10 wherein in the compound of formula (I) q is 3.

15. A method according to claim 10 wherein in the compound of formula (I) q is 4.

16. A method according to any one of claims 1 to 15 wherein in the compound of formula (I) Y is CH_2 .

25

17. A method according to any one of claims 1 to 16 wherein in the compound of formula (I) R^2 is H.

30

18. A method according to any one of claims 1 to 17 wherein in the compound of formula (I) R^3 is H.

19. A method according to any one of claims 1 to 18 wherein in the compound of formula (I) X is $C(=O)$.

35

20. A method according to any one of claims 1 to 19 wherein in the compound of formula (I), R^1 is selected from the group consisting of optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_6 - C_{18} aryl and optionally substituted C_1 - C_{18} heteroaryl.

5

21. A method according to any one of claims 1 to 20 wherein in the compound of formula (I) R^1 is optionally substituted C_6 - C_{18} aryl.


10

22. A method according to claim 21 wherein in the compound of formula (I) R^1 is an optionally substituted C_6 - C_{18} aryl selected from the group consisting of optionally substituted phenyl and optionally substituted naphthyl.

23. A method according to any one of claims 1 to 20 wherein in the compound of formula (I) R^1 is optionally substituted C_2 - C_{12} alkenyl.

15

24. A method according to claim 23 wherein in the compound of formula (I) R^1 is an optionally substituted C_2 - C_{12} alkenyl of the formula:



20

R^{1a} is selected from the group consisting of H, halogen and optionally substituted C_1 - C_{12} alkyl;

R^{1b} and R^{1c} are each independently selected from the group consisting of H, halogen, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl.

30

25. A method according to claim 24 wherein in the compound of formula (I) R^{1a} is H.

26. A method according to claim 24 or 25 wherein in the compound of formula (I) R^{1b} is H.

27. A method according to any one of claims 24 to 26 wherein in the compound of formula (I) R^{1c} is optionally substituted C_6-C_{18} aryl.

28. A method according to claim 24 wherein in the compound of formula (I) R^{1c} is optionally substituted phenyl.

29. A method according to any one of claims 1 to 28 wherein in the compound of formula (I) R^{5a} and R^{5b} are H.

30. A method according to any one of claims 1 to 29 wherein in the compound of formula (I) r is 1.

31. A method according to any one of claims 1 to 30 wherein in the compound of formula (I) R^7 is H.

32. A method according to any one of claims 1 to 31 wherein in the compound of formula (I), R^6 and R^8 are each independently selected from the group consisting of H, optionally substituted C_1-C_{12} alkyl, optionally substituted C_2-C_{12} alkenyl, optionally substituted C_6-C_{18} aryl and optionally substituted C_1-C_{18} heteroaryl.

33. A method according to any one of claims 1 to 32 wherein in the compound of formula (I) R^6 is selected from the group consisting of H, methyl, trifluoromethyl, ethyl, 2,2,2-trifluoroethyl, isopropyl, isopropenyl, propyl, 2-ethyl-propyl, 3,3-dimethyl-propyl, butyl, 2-methylbutyl, isobutyl, 3,3-dimethyl-butyl, 2-ethyl-butyl, pentyl, 2-methyl-pentyl, optionally substituted phenyl and optionally substituted C_1-C_5 heteroaryl.

34. A method according to any one of claims 1 to 33 wherein in the compound of formula (I) R^6 is optionally substituted phenyl or optionally substituted C_1-C_{18} heteroaryl.

35. A method according to any one of claims 1 to 34 wherein in the compound of formula (I) R^8 is selected from the group consisting of H, methyl, trifluoromethyl, ethyl, 2,2,2-trifluoroethyl, isopropyl, isopropenyl, propyl, 2-ethyl-propyl, 3,3-dimethyl-propyl, butyl, 2-

methyl-butyl, isobutyl, 3,3-dimethyl-butyl, 2-ethyl-butyl, pentyl, 2-methyl-pentyl, pent-4-enyl, hexyl, heptyl, octyl, optionally substituted phenyl and optionally substituted C₁-C₅ heteroaryl.

36. A method according to any one of claims 1 to 35 wherein in the compound of formula
5 (I) R⁸ is methyl, ethyl, phenyl or optionally substituted C₁-C₅ heteroaryl.

37. A method according to any one of claims 1 to 30, wherein in the compound of formula (I) R⁶, R⁷ and R⁸ when taken together with the carbon to which they are attached form a moiety selected from the group consisting of optionally substituted C₂-C₁₂alkenyl,
10 optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₃-C₁₂cycloalkenyl, optionally substituted C₂-C₁₂heterocycloalkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl.

38. A method according to claim 37 wherein in the compound of formula (I) R⁶, R⁷ and R⁸
15 when taken together with the carbon to which they are attached form an optionally substituted C₆-C₁₈aryl group.

39. A method according to claim 37 or 38 wherein in the compound of formula (I) R⁶, R⁷ and R⁸ when taken together with the carbon atom to which they are attached form a
20 disubstituted phenyl group.

40. A method according to claim 39 wherein the disubstituted phenyl group is a 2,4 disubstituted -phen-1-yl group or a 3,5 disubstituted -phen-1-yl group.

25 41. A method according to claim 39 or 40 wherein the disubstituted phenyl group is a 3,5-dichlorophen-1-yl group.


42. A method according to any one of claims 1 to 41 wherein in the compound of formula (I) each optional substituent is independently selected from the group consisting of: halogen,
30 =O, =S, -CN, -NO₂, -CF₃, -OCF₃, alkyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, heteroalkyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, aryl, heteroaryl, cycloalkylalkyl, heterocycloalkylalkyl, heteroarylalkyl, arylalkyl, cycloalkylalkenyl, heterocycloalkylalkenyl, arylalkenyl, heteroarylalkenyl, cycloalkylheteroalkyl, heterocycloalkylheteroalkyl, arylheteroalkyl, heteroarylheteroalkyl, hydroxy, hydroxyalkyl,
35 alkyloxy, alkyloxyalkyl, alkyloxycycloalkyl, alkyloxyheterocycloalkyl, alkyloxyaryl,

alkyloxyheteroaryl, alkyloxycarbonyl, alkylaminocarbonyl, alkenyloxy, alkynyloxy, cycloalkyloxy, cycloalkenyloxy, heterocycloalkyloxy, heterocycloalkenyloxy, aryloxy, phenoxy, benzyloxy, heteroaryloxy, arylalkyloxy, amino, alkylamino, acylamino, aminoalkyl, arylamino, sulfonylamino, sulfinylamino, sulfonyl, alkylsulfonyl, arylsulfonyl, aminosulfonyl, sulfinyl, alkylsulfinyl, arylsulfinyl, aminosulfinylaminoalkyl, -C(=O)OH, -C(=O)R^a, -C(=O)OR^a, C(=O)NR^aR^b, C(=NOH)R^a, C(=NR^a)NR^bR^c, NR^aR^b, NR^aC(=O)R^b, NR^aC(=O)OR^b, NR^aC(=O)NR^bR^c, NR^aC(=NR^b)NR^cR^d, NR^aSO₂R^b, -SR^a, SO₂NR^aR^b, -OR^a, OC(=O)NR^aR^b, OC(=O)R^a and acyl,

wherein R^a, R^b, R^c and R^d are each independently selected from the group consisting of H, C₁-C₁₂ alkyl, C₁-C₁₂ haloalkyl, C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, C₁-C₁₀ heteroalkyl, C₃-C₁₂ cycloalkyl, C₃-C₁₂ cycloalkenyl, C₁-C₁₂ heterocycloalkyl, C₁-C₁₂ heterocycloalkenyl, C₆-C₁₈aryl, C₁-C₁₈heteroaryl, and acyl, or any two or more of R^a, R^b, R^c and R^d, when taken together with the atoms to which they are attached form a heterocyclic ring system with 3 to 12 ring atoms.

43. A method according to any one of claims 1 to 42 wherein in the compound of formula (I) each optional substituent is independently selected from the group consisting of F, Cl, Br, I, CH₃, CH₂CH₃, OH, OCH₃, CF₃, OCF₃, NO₂, NH₂, and CN.

44. Use of a compound of the formula (I):



Formula (I)

wherein

Y is a group of formula $-(CR^9R^{10})_n-$;

X is selected from the group consisting -C(=O)-, -OC(=O)-, -NHC(=O)-, -(CR¹¹R¹²)_s, and -S(=O)₂-;

R is a group of the formula -Z-R⁴ wherein

Z is a group of formula -(CR¹³R¹⁴)_q;

R⁴ is selected from the group consisting of H, C₁-C₁₂alkyl, optionally substituted C₂-C₁₂alkenyl, optionally substituted C₂-C₁₂alkynyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₆-C₁₈aryl, optionally substituted C₁-C₁₈heteroaryl, NR^{4a}R^{4b}, C(=O)R¹⁵, C(=O)NR¹⁶R¹⁷, -C(=NR¹⁶)NR¹⁷R¹⁸, SR²⁰, SC(=O)R²⁰, SO₂R²⁰, OR²⁰, ONR¹⁶R¹⁷, OCR¹⁷R¹⁸R²⁰, OC(=O)R²⁰, OC(=O)OR²⁰, OC(=O)NR¹⁶R¹⁷, and ONR¹⁶C(=NR¹⁷)NR¹⁸R¹⁹;

R^{4a} is selected from the group consisting of H, optionally substituted C₁-C₁₂alkyl, optionally substituted C₂-C₁₂alkenyl, optionally substituted C₂-C₁₂alkynyl, optionally substituted C₁-C₁₂heteroalkyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂heterocycloalkyl, optionally substituted C₆-C₁₈aryl, optionally substituted C₁-C₁₈heteroaryl, C(=O)R^{15a}, C(=O)NR^{15a}R^{16a}, C(=O)OR^{15a}, SO₂R^{15a}, C(=O)H, -C(=NR^{15a})NR^{16a}R^{17a}, and OR^{15a},

R^{4b} is selected from the group consisting of H, optionally substituted C₁-C₁₂alkyl, optionally substituted C₂-C₁₂alkenyl, optionally substituted C₂-C₁₂alkynyl, optionally substituted C₁-C₁₂heteroalkyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂heterocycloalkyl, optionally substituted C₆-C₁₈aryl, optionally substituted C₁-C₁₈heteroaryl, C(=O)R^{15a}, C(=O)NR^{15a}R^{16a}, C(=O)OR^{15a}, or

R^{4a} and R^{4b} when taken together with the nitrogen atom to which they are attached form an optionally substituted heterocyclic moiety, or

one of R^{4a} and R^{4b} when taken together with any R¹³ or R¹⁴ and the atoms to which they are attached forms an optionally substituted heterocyclic moiety;

R¹³ and R¹⁴ are each independently selected from the group consisting of H, halogen, OH, C₁-C₁₂alkyl, C₆-C₁₈aryl, C₁-C₁₂hydroxyalkyl, C₁-C₁₂haloalkyl, C₁-C₁₂alkyloxy and C₁-C₁₂haloalkyloxy, or

when taken together with the carbon to which they are attached R^{13} and R^{14} form an optionally substituted C_3 - C_{12} cycloalkyl, or an optionally substituted C_1 - C_{12} heterocycloalkyl group, or

5

one of R^{13} and R^{14} when taken together with one of R^{4a} , and R^{4b} and the atoms to which they are attached form an optionally substituted heterocyclic moiety, or

10

one of R^{13} and R^{14} when taken together with one of R^{15} , R^{16} , R^{17} , R^{18} , R^{19} or R^{20} and the atoms to which they are attached form an optionally substituted cyclic moiety;

15

each R^{15} , R^{15a} , R^{16} , R^{16a} , R^{17} , R^{17a} , R^{18} , R^{19} and R^{20} is independently selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl, or

20

any two of R^{15} , R^{15a} , R^{16} , R^{16a} , R^{17} , R^{17a} , R^{18} , R^{19} and R^{20} when taken together with the atoms to which they are attached form an optionally substituted cyclic group, or

one of R^{15} , R^{16} , R^{17} , R^{18} , R^{19} and R^{20} when taken together with one of R^{13} and R^{14} and the atoms to which they are attached form an optionally substituted cyclic moiety;

25

q is an integer selected from the group consisting of 0, 1, 2, 3, 4, and 5;

30

R^1 is selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl, optionally substituted C_2 - C_{12} heterocycloalkyl, optionally substituted C_6 - C_{18} aryl, and optionally substituted C_1 - C_{18} heteroaryl;

R^2 and R^3 are each independently selected from the group consisting of H, optionally substituted C_1 - C_{12} alkyl, optionally substituted C_2 - C_{12} alkenyl, optionally substituted C_2 - C_{12} alkynyl, optionally substituted C_1 - C_{12} heteroalkyl, optionally substituted C_3 - C_{12} cycloalkyl,

optionally substituted C₂-C₁₂heterocycloalkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl;

5 each R^{5a} and R^{5b} are independently selected from the group consisting of H, halogen, C₁-C₁₂alkyl, C₁-C₁₂hydroxyalkyl and C₁-C₁₂haloalkyl, or

one or more of R^{5a} and R^{5b} when taken together with one or more of R⁶, R⁷ and R⁸ and the atoms to which they are attached form a moiety selected from the group consisting of an optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂ heterocycloalkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl;

R⁶, R⁷ and R⁸ are each independently selected from the group consisting of H, halogen, hydroxy, optionally substituted C₁-C₁₂alkyl, optionally substituted C₂-C₁₂alkenyl, optionally substituted C₂-C₁₂alkynyl, optionally substituted C₁-C₁₂ heteroalkyl, optionally substituted C₁-C₁₀ heteroalkenyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂ heterocycloalkyl, optionally substituted C₆-C₁₈aryl, optionally substituted C₁-C₁₈heteroaryl, optionally substituted amino, optionally substituted carboxy, C₁-C₁₂alkyloxy, and optionally substituted thio, or

20 (a) when taken together with the carbon atom to which they are attached two or more of R⁶, R⁷ and R⁸ form a moiety selected from the group consisting of optionally substituted C₂-C₁₂alkenyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂heterocycloalkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl, or

25 (b) one or more of R⁶, R⁷ and R⁸ when taken together with one or more of R^{5a} and R^{5b} and the atoms to which they are attached form a moiety selected from the group consisting of an optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂ heterocycloalkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl;

30 each R⁹ and R¹⁰ is independently selected from the group consisting of H, optionally substituted C₁-C₁₂alkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl;

35

each R^{11} and R^{12} is independently selected from the group consisting of H, and optionally substituted C_1 - C_{12} alkyl;

n is an integer selected from the group consisting of 1, 2, 3 and 4;

r is an integer selected from the group consisting of 1, 2, 3, and 4;

s is an integer selected from the group consisting of 1, 2, 3, and 4;

or a pharmaceutically acceptable salt or prodrug thereof, in down-regulating the activity of MC5R or a fragment thereof.

45. Use of a compound of formula (I) as described in claim 44 in the preparation of a medicament for treating, preventing, or controlling a condition associated with the activity of MC5R or a fragment thereof in a mammal.

46. Use of a compound of formula (I) as described in claim 44 in the preparation of a medicament for reducing sebum secretion in a mammal.

47. A use according to claim 45 or 46 wherein the mammal is a human.

48. A use according to claim 45 wherein the condition is selected from the group consisting of acne, seborrhea, and seborrheic dermatitis.

49. A use according to claim 48 wherein the condition is acne vulgaris.

50. A use according to any one of claims 45 to 49 wherein the medicament is adapted to be administered topically.


51. A use according to any one of claims 45 to 49 wherein the medicament is adapted to be administered orally.

52. A use according to any one of claims 45 to 49 wherein the medicament is adapted to be administered parenterally.

53. A use according to any one of claims 45 to 52 wherein the medicament contains a second active agent.

54. A use according to claim 53 wherein the second active agent is selected from the group consisting of antibiotics, retinoids, anti-androgens, and steroids.

55. A use according to any one of claims 44 to 54 wherein the compound of formula (I) is a compound of the formula (Ia):



Formula (Ia)

wherein R , R^1 , R^2 , R^3 , R^{5a} , R^{5b} , R^6 , R^7 , R^8 , X , Y and r are as defined in claim 54 and the stereochemistry is as shown, or a pharmaceutically acceptable salt or prodrug thereof.

56. A use according to claim 44 wherein in the compound of formula (I) $R^4 = NR^{4a}R^{4b}$.

57. A use according to claim 56 wherein in the compound of formula (I) R^{4a} is selected from the group consisting of H , $C(=NH)NH_2$, $C(=NH)N(CH_3)_2$, $C(=NH)NHCH(CH_3)_2$, $C(=O)CH_3$, $C(=O)cyclohexyl$, CH_3 , CH_2CH_3 , $CH_2CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH_2CH_2CH_3$, $CH(CH_3)CH_2CH_3$, $CH_2CH(CH_3)_2$, $C(CH_3)_3$, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, benzyl, and phenyl, or a halogenated derivative thereof.

58. A use according to claim 56 wherein in the compound of formula (I) R^{4b} is selected from the group consisting of H , CH_3 , CH_2CH_3 , $CH_2CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH_2CH_2CH_3$, $CH(CH_3)CH_2CH_3$, $CH_2CH(CH_3)_2$, $C(CH_3)_3$, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, benzyl, and phenyl, or a halogenated derivative thereof.

59. A use according to claim 56 wherein in the compound of formula (I) R^{4a} and R^{4b} when taken together with the nitrogen atom to which they are attached form an optionally substituted C_2 - C_{12} heterocycloalkyl group, an optionally substituted C_2 - C_{12} heterocycloalkenyl group or an optionally substituted C_1 - C_{18} heteroaryl group.

5

60. A use according to claim 59 wherein in the compound of formula (I) R^{4a} and R^{4b} when taken together with the nitrogen atom to which they are attached form an optionally substituted heterocycloalkyl group selected from the group consisting of piperidin-1-yl, pyrrolidin-1-yl, azetidin-1-yl, azepan-1-yl, piperazin-1-yl, and morpholin-4-yl.

10

61. A use according to claim 56 wherein in the compound of formula (I) one of R^{4a} and R^{4b} when taken together with the nitrogen atom to which it is attached and one of R^{13} and R^{14} and the carbon atom to which it is attached form an optionally substituted C_2 - C_{12} heterocycloalkyl group.

15

62. A use according to claim 61 wherein in the compound of formula (I) one of R^{4a} and R^{4b} when taken together with the nitrogen atom to which it is attached and one of R^{13} and R^{14} and the carbon atom to which it is attached form an optionally substituted heterocycloalkyl group selected from the group consisting of piperidinyl, pyrrolidinyl, azetidiny, morpholinyl, piperazinyl, and azepanyl.

20

63. A use according to claim 44 wherein in the compound of formula (I) Z is $-(CH_2)_q-$.

64. A use according to claim 63 wherein in the compound of formula (I) q is an integer selected from the group consisting of 1, 2, 3 and 4.

25

65. A use according to claim 63 wherein in the compound of formula (I) q is 1.

66. A use according to claim 63 wherein in the compound of formula (I) q is 2.

30

67. A use according to claim 63 wherein in the compound of formula (I) q is 3.

68. A use according to claim 63 wherein in the compound of formula (I) q is 4.

246

69. A use according to any one of claims 44 to 68 wherein in the compound of formula (I) Y is CH₂.

70. A use according to any one of claims 44 to 69 wherein in the compound of formula (I)
5 R² is H.

71. A use according to any one of claims 44 to 70 wherein in the compound of formula (I) R³ is H.

10 72. A use according to any one of claims 44 to 71 wherein in the compound of formula (I) X is C(=O).


73. A use according to any one of claims 44 to 72 wherein in the compound of formula (I) R¹ is selected from the group consisting of optionally substituted C₂-C₁₂alkenyl, optionally
15 substituted C₆-C₁₈aryl and optionally substituted C₁-C₁₈heteroaryl.

74. A use according to any one of claims 44 to 73 wherein in the compound of formula (I) R¹ is optionally substituted C₆-C₁₈aryl.

20 75. A use according to claim 74 wherein in the compound of formula (I) R¹ is an optionally substituted C₆-C₁₈aryl selected from the group consisting of optionally substituted phenyl and optionally substituted naphthyl.

76. A use according to any one of claims 44 to 73 wherein in the compound of formula (I)
25 R¹ is optionally substituted C₂-C₁₂alkenyl.

77. A use according to claim 76 wherein in the compound of formula (I) R¹ is an optionally substituted C₂-C₁₂alkenyl of the formula:



R^{1a} is selected from the group consisting of H, halogen and optionally substituted C₁-C₁₂ alkyl;

R^{1b} and R^{1c} are each independently selected from the group consisting of H, halogen, optionally substituted C₁-C₁₂alkyl, optionally substituted C₂-C₁₂alkenyl, optionally substituted C₂-C₁₂alkynyl, optionally substituted C₁-C₁₂heteroalkyl, optionally substituted C₃-C₁₂cycloalkyl, optionally substituted C₂-C₁₂ heterocycloalkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl.

78. A use according to claim 77 wherein in the compound of formula (I) R^{1a} is H.

79. A use according to claim 77 or 78 wherein in the compound of formula (I) R^{1b} is H.

80. A use according to any one of claims 77 to 79 wherein in the compound of formula (I) R^{1c} is optionally substituted C₆-C₁₈aryl.

81. A use according to claim 80 wherein in the compound of formula (I) R^{1c} is optionally substituted phenyl.

82. A use according to any one of claims 44 to 81 wherein in the compound of formula (I) R^{5a} and R^{5b} are H.

83. A use according to any one of claims 44 to 82 wherein in the compound of formula (I) r is 1.

84. A use according to any one of claims 44 to 83 wherein in the compound of formula (I) R⁷ is H.

85. A use according to any one of claims 44 to 84 wherein in the compound of formula (I) R⁶ and R⁸ are each independently selected from the group consisting of H, optionally substituted C₁-C₁₂ alkyl, optionally substituted C₂-C₁₂ alkenyl, optionally substituted C₆-C₁₈ aryl and optionally substituted C₁-C₁₈ heteroaryl.

86. A use according to any one of claims 44 to 85 wherein in the compound of formula (I) R⁶ is selected from the group consisting of H, methyl, trifluoromethyl, ethyl, 2,2,2-

trifluoroethyl, isopropyl, isopropenyl, propyl, 2-ethyl-propyl, 3,3-dimethyl-propyl, butyl, 2-methylbutyl, isobutyl, 3,3-dimethyl-butyl, 2-ethyl-butyl, pentyl, 2-methyl-pentyl, optionally substituted phenyl and optionally substituted C₁-C₅ heteroaryl.

5 87. A use according to any one of claims 44 to 86 wherein in the compound of formula (I) R⁶ is optionally substituted phenyl or optionally substituted C₁-C₁₈heteroaryl.

88. A use according to any one of claims 44 to 87 wherein in the compound of formula (I) R⁸ is selected from the group consisting of H, methyl, trifluoromethyl, ethyl, 2,2,2-
10 trifluoroethyl, isopropyl, isopropenyl, propyl, 2-ethyl-propyl, 3,3-dimethyl-propyl, butyl, 2-methyl-butyl, isobutyl, 3,3-dimethyl-butyl, 2-ethyl-butyl, pentyl, 2-methyl-pentyl, pent-4-enyl, hexyl, heptyl, octyl, optionally substituted phenyl and optionally substituted C₁-C₅ heteroaryl.

89. A use according to any one of claims 44 to 88 wherein in the compound of formula (I)
15 R⁸ is methyl, ethyl, phenyl or optionally substituted C₁-C₅ heteroaryl.

90. A use according to any one of claims 44 to 83 wherein in the compound of formula (I) R⁶, R⁷ and R⁸ when taken together with the carbon to which they are attached form a moiety selected from the group consisting of optionally substituted C₂-C₁₂alkenyl, optionally
20 substituted C₃-C₁₂cycloalkyl, optionally substituted C₃-C₁₂cycloalkenyl, optionally substituted C₂-C₁₂heterocycloalkyl, optionally substituted C₆-C₁₈aryl, and optionally substituted C₁-C₁₈heteroaryl.

91. A use according to claim 90 wherein in the compound of formula (I) R⁶, R⁷ and R⁸
25 when taken together with the carbon to which they are attached form an optionally substituted C₆-C₁₈aryl group.

92. A use according to claim 90 or 91 wherein in the compound of formula (I) wherein R⁶, R⁷ and R⁸ when taken together with the carbon atom to which they are attached form a
30 disubstituted phenyl group.

93. A use according to claim 92 wherein the disubstituted phenyl group is a 2,4 disubstituted -phen-1-yl group or a 3,5 disubstituted -phen-1-yl group.

94. A use according to claim 92 or 93 wherein the disubstituted phenyl group is a 3,5-dichlorophen-1-yl group.

95. A use according to any one of claims 44 to 94 wherein in the compound of formula (I) each optional substituent is independently selected from the group consisting of: halogen, =O, =S, -CN, -NO₂, -CF₃, -OCF₃, alkyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, heteroalkyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, aryl, heteroaryl, cycloalkylalkyl, heterocycloalkylalkyl, heteroarylalkyl, arylalkyl, cycloalkylalkenyl, heterocycloalkylalkenyl, arylalkenyl, heteroarylalkenyl, cycloalkylheteroalkyl, heterocycloalkylheteroalkyl, arylheteroalkyl, heteroarylheteroalkyl, hydroxy, hydroxyalkyl, alkyloxy, alkyloxyalkyl, alkyloxycycloalkyl, alkyloxyheterocycloalkyl, alkyloxyaryl, alkyloxyheteroaryl, alkyloxycarbonyl, alkylaminocarbonyl, alkenyloxy, alkynyloxy, cycloalkyloxy, cycloalkenyloxy, heterocycloalkyloxy, heterocycloalkenyloxy, aryloxy, phenoxy, benzyloxy, heteroaryloxy, arylalkyloxy, amino, alkylamino, acylamino, aminoalkyl, arylamino, sulfonylamino, sulfinylamino, sulfonyl, alkylsulfonyl, arylsulfonyl, aminosulfonyl, sulfinyl, alkylsulfinyl, arylsulfinyl, aminosulfinylaminoalkyl, -C(=O)OH, -C(=O)R^a, -C(=O)OR^a, C(=O)NR^aR^b, C(=NOH)R^a, C(=NR^a)NR^bR^c, NR^aR^b, NR^aC(=O)R^b, NR^aC(=O)OR^b, NR^aC(=O)NR^bR^c, NR^aC(=NR^b)NR^cR^d, NR^aSO₂R^b, -SR^a, SO₂NR^aR^b, -OR^a, OC(=O)NR^aR^b, OC(=O)R^a and acyl,

wherein R^a, R^b, R^c and R^d are each independently selected from the group consisting of H, C₁-C₁₂ alkyl, C₁-C₁₂ haloalkyl, C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, C₁-C₁₀ heteroalkyl, C₃-C₁₂ cycloalkyl, C₃-C₁₂ cycloalkenyl, C₁-C₁₂ heterocycloalkyl, C₁-C₁₂ heterocycloalkenyl, C₆-C₁₈aryl, C₁-C₁₈heteroaryl, and acyl, or any two or more of R^a, R^b, R^c and R^d, when taken together with the atoms to which they are attached form a heterocyclic ring system with 3 to 12 ring atoms.

96. A use according to any one of claims 44 to 95 wherein in the compound of formula (I) each optional substituent is independently selected from the group consisting of F, Cl, Br, I, CH₃, CH₂CH₃, OH, OCH₃, CF₃, OCF₃, NO₂, NH₂, and CN.

97. A method according to any one of claims 1 to 43 or a use according to any one of claims 44 to 96 wherein the compound of formula (I) is selected from the group consisting of:

- N-(((3S,5S)-1-(3,5-dichlorobenzyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

- N-(((3S,5S)-3-(2-(diethylamino)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide
- 5 • (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-(2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 • N-(((3S,5S)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)naphthalene-2-sulfonamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-bromo-N-methyl-2-naphthamide
- 20 • N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-4-methyl-2-oxo-1,4-diazepan-5-yl)methyl)-6-bromo-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 • N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-(isopropylamino)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-(3-methylguanidino)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 30 • (E)-N-(((3S,5S)-3-butyl-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- N-((S)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)acetamide

- (S)-2-acetamido-N-((S)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)-3-(1H-imidazol-5-yl)propanamide
- propyl (S)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethylcarbamate
- 5 • N-((R)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)acetamide
- (S)-2-acetamido-N-((R)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethyl)-3-(1H-imidazol-4-yl)propanamide
- propyl (R)-1-((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)-2-(naphthalen-2-yl)ethylcarbamate
- 10 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)biphenyl-4-carboxamide
- 15 • N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1H-indole-2-carboxamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)biphenyl-4-carboxamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1H-indole-2-carboxamide
- 20 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(naphthalen-2-yl)acetamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,2,3,4-tetrahydronaphthalene-2-carboxamide
- 25 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)quinoline-3-carboxamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)quinoxaline-2-carboxamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)isoquinoline-3-carboxamide
- 30 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)quinoline-2-carboxamide

- N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-1-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-1-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(naphthalen-2-yl)acetamide
- 5 • N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-1-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1-naphthamide
- N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-1-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(1H-indol-3-yl)acetamide
- N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-2-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(biphenyl-4-yl)acetamide
- 10 • N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-2-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-2-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(naphthalen-2-yl)acetamide
- 15 • N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-2-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1-naphthamide
- N-(((3S,5S)-3-(4-aminobutyl)-1-(naphthalen-2-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(naphthalen-1-yl)acetamide
- N-(((3S,5S)-3-(4-aminobutyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 20 • (S)-N-(((3S,5S)-3-(4-aminobutyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-3-carboxamide
- (R)-N-(((3S,5S)-3-(4-aminobutyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-3-carboxamide
- 25 • N-(((3S,5S)-3-(4-aminobutyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzofuran-2-carboxamide
- (R)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-(3-methylguanidino)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-3-carboxamide
- (S)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-3-carboxamide
- 30 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzofuran-2-carboxamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2,3-dihydro-1H-indene-2-carboxamide

- (R)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-3-carboxamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzo[b]thiophene-2-carboxamide
- 5 • 2,4-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- 2,5-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- 10 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)cyclohexanecarboxamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-phenoxybenzamide
- 15 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-phenoxybenzamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1H-indole-2-carboxamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-phenylpropanamide
- 20 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dimethylbenzamide
- 4-tert-butyl-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- 25 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2,4-dimethoxybenzamide
- 2-cyclohexyl-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acetamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzo[d][1,3]dioxole-5-carboxamide
- 30 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1H-benzo[d][1,2,3]triazole-5-carboxamide

- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)cyclopentanecarboxamide
- 3,4-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- 5 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)cinnamamide
- 3,5-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- 10 • 2-(2,4-dichlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acetamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1-methoxy-2-naphthamide
- 2-(3,4-dichlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acetamide
- 15 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-methoxy-2-naphthamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(2-(guanidinooxy)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-3-(2,4-dichlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- 20 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-adamantane-1-carboxamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-phenoxyacetamide
- 25 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-methoxy-2-naphthamide
- 4-bromo-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- (S)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2,3-dihydrobenzo[b][1,4]dioxine-2-carboxamide
- 30 • (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(thiophen-2-yl)acrylamide

- (R)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2,3-dihydrobenzo[b][1,4]dioxine-2-carboxamide
- (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-hydroxyphenyl)acrylamide
- 5 • (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(2-methoxyphenyl)acrylamide
- (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-p-tolylacrylamide
- (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(2-(trifluoromethyl)phenyl)acrylamide
- 10 • (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(3-fluorophenyl)acrylamide
- (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-methyl-3-phenylacrylamide
- 15 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-phenylcyclopropanecarboxamide
- 2-(2,4-dichlorophenoxy)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acetamide
- (E)-3-(3-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- 20 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzo[d]thiazole-6-carboxamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-5-phenylfuran-2-carboxamide
- 25 • (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(3-methoxyphenyl)acrylamide
- 6-bromo-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-guanidinopropyl)-2-oxo-1-phenethyl-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 30 • N-(((3S,5S)-1-(3,4-dichlorophenethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2,4-dichlorophenethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzo[b]thiophene-5-carboxamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-5-methyl-1-phenyl-1H-pyrazole-4-carboxamide
- 5 • (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-methoxyphenyl)acrylamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide
- (E)-3-(2-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- 10 • (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(2-hydroxyphenyl)acrylamide
- (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-m-tolylacrylamide
- 15 • (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(3-(trifluoromethyl)phenyl)acrylamide
- (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(3-hydroxyphenyl)acrylamide
- (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(2-fluorophenyl)acrylamide
- 20 • (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-o-tolylacrylamide
- (Z)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-fluoro-3-phenylacrylamide
- 25 • N-((1-(2,2-diphenylethyl)-2-oxo-3-(piperidin-4-yl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-((1-(2,2-diphenylethyl)-2-oxo-3-(piperidin-4-yl)methyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)acrylamide
- 30 • (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-(trifluoromethyl)phenyl)acrylamide
- N-(((3S,5S)-1-(2,2-diphenylpropyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

- N-(((3S,5S)-1-(cyclohexylmethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(1-adamantylmethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 • N-(((3S,5S)-1-((S)-1,1-diphenylpropan-2-yl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-((R)-1,1-diphenylpropan-2-yl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-cyclohexyl-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-
- 10 naphthamide
- N-(((3S,5S)-1-((R)-1-fluoro-1,1-diphenylpropan-2-yl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-3-(2,6-difluorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- 15 • (E)-3-(2-chloro-6-fluorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- (E)-3-(4-bromophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-ethoxyphenyl)acrylamide
- 20 • N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-bromo-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)cinnamamide
- 25 • (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-1,4-dimethoxy-2-naphthamide
- N-(((3S,5S)-3-(3-(3,3-dimethylguanidino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-
- 30 diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-hydroxy-2-naphthamide
- 6-amino-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

- (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-p-tolylacrylamide
- (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)acrylamide
- 5 • N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-ethylhexanamide
- 10 • N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dimethylbenzamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-ethylhexanamide
- 15 • N-(((3S,5S)-3-(3-(cyclohexylamino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-guanidinopropyl)-1-(naphthalen-2-yl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-((9H-fluoren-9-yl)methyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 20 • N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(pyridin-3-ylmethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(pyridin-4-ylmethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 • (E)-N-(((3S,5S)-3-(3-(cyclohexylamino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)acrylamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(4-(isopropylamino)butyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(2,4-difluorophenyl)acrylamide
- 30 • (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-cyanophenyl)acrylamide
- (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(naphthalen-2-yl)acrylamide

- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(4-fluorophenoxy)acetamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-5-(4-chlorophenyl)furan-2-carboxamide
- 5 • N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-(1H-pyrrol-1-yl)benzamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-oxo-1-phenylpyrrolidine-3-carboxamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-5-
- 10 (4-chlorophenyl)isoxazole-3-carboxamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-5-(furan-2-yl)isoxazole-3-carboxamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-phenylthiazole-4-carboxamide
- 15 • N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-(3,5-dimethyl-1H-pyrazol-1-yl)benzamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-5-methyl-1-phenyl-1H-pyrazole-4-carboxamide
- N-(((3S,5S)-1-(2-cyclohexylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-
- 20 yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2-(bicyclo[2.2.1]heptan-2-yl)ethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2,2-bis(4-methoxyphenyl)ethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 • N-(((3S,5S)-3-(3-(benzylamino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-(cyclopentylamino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-(cyclobutylamino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-
- 30 yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-(dicyclobutylamino)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-benzyl-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

- N-(((3S,5S)-1-(2,2-bis(4-fluorophenyl)ethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-guanidinopropyl)-1-(naphthalen-2-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 • (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(5-methylthiophen-2-yl)acrylamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-phenyl-1H-pyrazole-5-carboxamide
- (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)-N-methylacrylamide
- 10 • (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-4-methyl-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)acrylamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-(3-methyl-5-oxo-4,5-dihydro-1H-pyrazol-1-yl)benzamide
- 15 • (E)-N-(((3S,5S)-3-(3-aminopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-bromophenyl)acrylamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(pyrrolidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 20 • N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(pyrrolidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 • N-(((3S,5S)-3-(3-(azetidin-1-yl)propyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-guanidinopropyl)-1-(naphthalen-1-ylmethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-guanidinopropyl)-1-(2-(naphthalen-2-yl)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 30 • N-(((3S,5S)-1-((S)-2-acetamido-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)cinnamamide

- N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-3,4-dimethylbenzamide
- 3,4-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)benzamide
- 5 • N-(((3S,5S)-1-((S)-2-(cyclobutanecarboxamido)-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-((S)-2-(cyclohexanecarboxamido)-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(aminomethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 • (E)-N-(((3S,5S)-3-(aminomethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)acrylamide
- 15 • (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-p-tolylacrylamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(piperidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(piperidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 20 • N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dimethylbenzamide
- 25 • N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)cinnamamide
- (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- 3,4-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 30 • N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-3,4-dimethylbenzamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)cinnamamide

- (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-3-(4-fluorophenyl)acrylamide
- (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-3-p-tolylacrylamide
- 5 • N-(((3S,5S)-1-(3,5-dimethylbenzyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-((S)-2-benzamido-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-((R)-2-benzamido-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 • N-(((3S,5S)-3-(3-guanidinopropyl)-1-(2-methoxy-2-phenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-guanidinopropyl)-2-oxo-1-(2-phenyl-2-propoxyethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 • N-(((3S,5S)-1-(2-(benzyloxy)-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2-(allyloxy)-2-phenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-butyl-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 20 • (E)-N-(((3S,5S)-3-butyl-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- (E)-N-(((3S,5S)-3-(3-amino-3-oxopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- 25 • N-(((3S,5S)-3-(3-amino-3-oxopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-N-(((3S,5S)-3-(3-acetamidopropyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-p-tolylacrylamide
- N-(3-((2S,7S)-4-(2,2-diphenylethyl)-3-oxo-7-(((E)-3-p-tolylacrylamido)methyl)-1,4-diazepan-2-yl)propyl)cyclohexanecarboxamide
- 30 • N-(((3S,5S)-3-(3-guanidinopropyl)-2-oxo-1-(2-phenoxy-2-phenylethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- ethyl 3-((3S,5S)-5-((2-naphthamido)methyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-1-yl)-2-phenylpropanoate

- N-(((3S,5S)-1-(2-ethylbutyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-((3,5-dimethylcyclohexyl)methyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 • N-(2-((2S,7S)-7-(((E)-3-(4-chlorophenyl)acrylamido)methyl)-4-(2,2-diphenylethyl)-3-oxo-1,4-diazepan-2-yl)ethyl)cyclohexanecarboxamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-3-(2-(2-cyclohexylacetamido)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-aminoethyl)-1,4-diazepan-5-yl)ethyl)benzamide
- 10 • 3,4-dichloro-N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-aminoethyl)-1,4-diazepan-5-yl)ethyl)benzamide
- N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-aminoethyl)-1,4-diazepan-5-yl)ethyl)-2-naphthamide
- 15 • N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)ethyl)benzamide
- 3,4-dichloro-N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)ethyl)benzamide
- N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)ethyl)-2-naphthamide
- 20 • N-(((3S,5S)-1-(3-(dimethylamino)-3-oxo-2-phenylpropyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(cyclohexylmethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 • N-(((3S,5S)-3-(2-aminoethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-oxo-3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-oxo-3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 30 • N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-phenethyl-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-phenethyl-1,4-diazepan-5-yl)methyl)acrylamide

- N-(((3S,5S)-3-(2-cyclohexylethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-3-(2-cyclohexylethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- 5 • N-(((3S,5S)-3-benzyl-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-N-(((3S,5S)-3-benzyl-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-10 3,4-dichlorobenzamide
- (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- N-(((3S,5S)-1-(3-chloro-5-fluorobenzyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 • N-(((3S,5S)-1-(3,5-difluorobenzyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(3-chloro-5-fluorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(3,5-difluorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-20 2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,6-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 • N-(((3S,5S)-3-(3-aminopropyl)-1-(3,5-dimethoxybenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2-chlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,3-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-30 2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(2,4-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(3,4-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

- N-(((3S,5S)-3-(3-aminopropyl)-1-(3-fluoro-5-methylbenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(3-fluoro-5-(trifluoromethyl)benzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 • N-(((3S,5S)-3-(3-aminopropyl)-1-(4-chlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-2-oxo-1-(2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 • N-(((3S,5S)-3-(3-aminopropyl)-2-oxo-1-((1-phenylcyclohexyl)methyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 3,4-dichloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 • (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- 20 • (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 3,4-dichloro-N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 25 • N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-chloro-3-fluorobenzamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-chloro-3-methylbenzamide
- 30 • N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-chloro-4-fluorobenzamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-chloro-4-methylbenzamide

- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(piperidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- N-(2-((3S,5R)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(pyrrolidin-1-yl)ethyl)-1,4-diazepan-5-yl)ethyl)-2-naphthamide
- 5 • N-(((3S,5S)-3-(3-aminopropyl)-1-(3,5-bis(trifluoromethyl)benzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-1-(3-chlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-2-oxo-1-(2-phenylbutyl)-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 • N-(((3S,5S)-3-(3-guanidinopropyl)-2-oxo-1-(3-oxo-2-phenyl-3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-guanidinopropyl)-2-oxo-1-(3-oxo-2-phenyl-3-(phenylamino)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 • 3,4-dichloro-N-(2-((3S,5R)-1-(2,2-diphenylethyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-diazepan-5-yl)ethyl)benzamide
- 3,4-dichloro-N-(2-((3S,5R)-3-(2-(diisopropylamino)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)ethyl)benzamide
- N-(((3S,5S)-3-(aminomethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- 20 • N-(((3S,5S)-3-(aminomethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-N-(((3S,5S)-3-(aminomethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- 25 • 3,4-dichloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(piperidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)benzamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(piperidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-(2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- 30 • (E)-N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-(2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- 3,4-dichloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(pyrrolidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)benzamide

- N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(pyrrolidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-2-oxo-1-((S)-2-phenylpropyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 • N-(((3S,5S)-3-(3-aminopropyl)-2-oxo-1-((R)-2-phenylpropyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(2-(dimethylamino)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide
- 10 • N-(((3S,5S)-3-(3-aminopropyl)-1-(3,5-diethynylbenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-3-(2-(diethylamino)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- 15 • N-(((3S,5S)-3-(2-(diethylamino)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 20 • (E)-3-(4-chlorophenyl)-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- 25 • N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(pyrrolidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide
- 30 • N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-bromo-2-naphthamide

- N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide
- 6-chloro-N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 • 6-bromo-N-(((3S,5S)-1-(2-ethylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-((3,5-dimethylcyclohexyl)methyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-((3,5-dimethylcyclohexyl)methyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 • (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-((3,5-dimethylcyclohexyl)methyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- 6-chloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 • N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-6-fluoro-2-naphthamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(pyrrolidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- 20 • N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 3,4-dichloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- 25 • N-(((3S,5S)-3-(3-aminopropyl)-1-((2,6-dimethylcyclohexyl)methyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 3,4-dichloro-N-(((3S,5S)-1-((3,5-dimethylcyclohexyl)methyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 30 • 3,4-dichloro-N-(((3S,5S)-1-((3,5-dimethylcyclohexyl)methyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(3-methyl-2-phenylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide

- N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-aminopropyl)-2-oxo-1-((R)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 • (E)-N-(((3S,5S)-3-((1H-imidazol-4-yl)methyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-chlorophenyl)acrylamide
- 3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)propanamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-10 (4-chlorophenyl)propanamide
- N-(((2S,7S)-7-(((E)-3-(4-chlorophenyl)acrylamido)methyl)-4-(2,2-diphenylethyl)-3-oxo-1,4-diazepan-2-yl)methyl)picolinamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(pyridin-2-yl)methyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 15 • N-(((3S,5S)-1-((R)-3-methyl-2-phenylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-((S)-3-methyl-2-phenylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-20 yl)methyl)-3-(4-isopropylphenyl)acrylamide
- (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(4-isopropylphenyl)acrylamide
- (E)-3-(2,4-dimethylphenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 25 • (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-(2,4-dimethylphenyl)acrylamide
- (E)-3-(2,4-difluorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (E)-N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-30 yl)methyl)-3-(2,4-difluorophenyl)acrylamide
- N-(((2S,7S)-7-(((E)-3-(4-chlorophenyl)acrylamido)methyl)-4-(2,2-diphenylethyl)-3-oxo-1,4-diazepan-2-yl)methyl)cyclohexanecarboxamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(2-morpholinoethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide

- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-3-(2-(2,5-dimethyl-1H-pyrrol-1-yl)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-3-(2-(2,5-dimethylpyrrolidin-1-yl)ethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)acrylamide
- 5 • 6-chloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(pyrrolidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-3-(2-(isopropylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 10 • 6-chloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(pyrrolidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 15 • 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(pyrrolidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- benzyl ((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methylcarbamate
- (E)-3-(4-bromophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 20 • 5-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)isoxazole-3-carboxamide
- (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-oxo-2-(pyridin-2-ylamino)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 25 • (E)-3-(4-chlorophenyl)-N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-oxo-2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide
- 6-chloro-N-(((3S,5S)-2-oxo-3-(3-oxo-3-(piperidin-1-yl)propyl)-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 3,4-dichloro-N-(((3S,5S)-2-oxo-3-(3-oxo-3-(piperidin-1-yl)propyl)-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 30 • 6-chloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(piperidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(piperidin-1-ylmethyl)-1,4-diazepan-5-yl)methyl)benzamide

- 6-chloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(3-(piperidin-1-yl)propyl)-1,4-diazepan-5-yl)methyl)benzamide
- 5 • (E)-N-(2-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)propan-2-yl)-3-(4-chlorophenyl)acrylamide
- (E)-3-(4-chlorophenyl)-N-(2-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)propan-2-yl)acrylamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-((R)-2-(4-chlorophenyl)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 • N-(((3S,5S)-3-(2-aminoethyl)-1-(2-(4-chlorophenyl)propyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-((R)-2-(4-chlorophenyl)propyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 15 • N-(((3S,5S)-1-((S)-2-(4-chlorophenyl)propyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 3,4-dichloro-N-(((3S,5S)-3-(2-(methyl(phenyl)amino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 3,4-dichloro-N-(((3S,5S)-3-(2-(diethylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 20 • 3,4-dichloro-N-(((3S,5S)-3-(2-morpholinoethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 3,4-dichloro-N-(((3S,5S)-2-oxo-3-(2-(phenylamino)ethyl)-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 25 • N-(((3S,5S)-3-(2-(benzylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- (5S,9aS)-5-(2-aminobenzyl)-2-((E)-3-(4-chlorophenyl)acryloyl)-7-(2,2-diphenylethyl)hexahydro-1H-imidazo[1,5-d][1,4]diazepin-6(5H)-one
- N-(((3S,5S)-3-(2-(tert-butylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- 30 • 3,4-dichloro-N-(((3S,5S)-3-(2-(4-methylpiperazin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- N-(((3S,5S)-2-oxo-1-((R)-2-phenylpentyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

- N-(((3S,5S)-2-oxo-1-((S)-2-phenylpentyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-4-(trifluoromethyl)benzamide
- 5 • N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-4-(trifluoromethyl)benzamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-3-(trifluoromethyl)benzamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3-
- 10 (trifluoromethyl)benzamide
- 6-chloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 3,4-dichloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-3-(2-(isopropylamino)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide
- 15 • 6-chloro-N-(((3S,5S)-3-(2-(isopropylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- N-(((3S,5S)-3-(2-aminobenzyl)-1-(2,2-diphenylethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-
- 20 2-naphthamide
- N-(((3S,5S)-3-(2-(benzyl(methyl)amino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperazin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 25 • 3,4-dichloro-N-(((3S,5S)-3-(2-(methyl(pentyl)amino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 3,4-dichloro-N-(((3S,5S)-3-(2-(diisopropylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 3,4-dichloro-N-(((3S,5S)-3-(2-(4-methylpiperidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-
- 30 1,4-diazepan-5-yl)methyl)benzamide
- (S)-6-chloro-N-((2-oxo-1-(2-phenylbutyl)-3-(piperidin-4-yl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- (S)-6-chloro-N-(3-(1-isopentylpiperidin-4-yl)-2-oxo-1-(2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide

- N-(((3S,5S)-3-butyl-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- 6-chloro-N-(((3S,5S)-3-isopentyl-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 • 3,4-dichloro-N-(((3S,5S)-3-(2-(3,5-dimethylpiperidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 3,4-dichloro-N-(((3S,5S)-3-(2-(4-hydroxypiperidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 10 • 1-(2-((2S,7S)-7-((3,4-dichlorobenzamido)methyl)-3-oxo-4-((S)-2-phenylbutyl)-1,4-diazepan-2-yl)ethyl)piperidine-4-carboxylic acid
- N-(((3S,5S)-3-(2-(azepan-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- 3,4-dichloro-N-(((3S,5S)-3-(2-((S)-2-methylpiperidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- 15 • N-(((3S,5S)-3-(2-(tert-butyl(methyl)amino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- N-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)benzyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(3-(butyl(methyl)amino)-3-oxopropyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 20 • N-(((3S,5S)-3-(3-(cyclohexylamino)-3-oxopropyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 6-chloro-N-((3-(1-ethylpiperidin-4-yl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 25 • (3S,5S)-5-((3,4-dichlorobenzylamino)methyl)-1-(2,2-diphenylethyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-2-one
- 6-chloro-N-(((3S,5S)-3-(2-guanidinoethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 6-chloro-N-(((3S,5S)-3-(2-(3-methylguanidino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 30 • N-(((3S,5S)-3-(2-aminoethyl)-1-((R)-2-ethyl-3-methylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-((S)-2-ethyl-3-methylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide

- 3,4-dichloro-N-(((3S,5S)-1-((R)-2-ethyl-3-methylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 3,4-dichloro-N-(((3S,5S)-1-((S)-2-ethyl-3-methylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 5 • N-(((3S,5S)-3-(2-amino-2-methylpropyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- N-(((3R,5R)-3-(2-amino-2-methylpropyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- 10 • 3,4-dichloro-N-(((3S,5S)-3-(2-(isopropylamino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-(3,5-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- 15 • (498) 6-chloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 6-chloro-N-(((3S,5S)-3-(2-methyl-2-(piperidin-1-yl)propyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 6-chloro-N-(((3R,5R)-3-(2-methyl-2-(piperidin-1-yl)propyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 20 • N-(((3S,5S)-3-(2-aminoethyl)-1-((R)-2-ethyl-3-methylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- N-(((3S,5S)-3-(2-aminoethyl)-1-((S)-2-ethyl-3-methylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- 25 • 6-chloro-N-(((3S,5S)-1-((R)-2-ethyl-3-methylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 6-chloro-N-(((3S,5S)-1-((S)-2-ethyl-3-methylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 6-chloro-N-(((3S,5S)-3-(2-cyclohexylethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 30 • 6-chloro-N-(((3S,5S)-3-hexyl-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 6-(((3S,5S)-1-(2,2-diphenylethyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)carbamoyl)-2-naphthoic acid

- 6-chloro-N-(((3S,5S)-3-(2-(3-isopropylguanidino)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 6-chloro-N-(((3S,5S)-3-(4-hydroxybutyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 5 • 6-chloro-N-(((3S,5S)-3-(2-methoxyethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- N-(((3S,5S)-3-(2-(benzyloxy)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- 6-chloro-N-(((3S,5S)-3-isobutyl-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 10 • N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethyl-3-methylbut-3-enyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide
- 3,4-dichloro-N-(((3S,5S)-1-(2-ethyl-3-methylbut-3-enyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide
- 15 • N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethyl-3-methylbut-3-enyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide
- 6-chloro-N-(((3S,5S)-1-((R)-2-ethyl-3-methylbut-3-enyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 6-chloro-N-(((3S,5S)-1-((S)-2-ethyl-3-methylbut-3-enyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide
- 20 • N-(((3S,5S)-1-(cyclohexylmethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)biphenyl-4-carboxamide
- N-(((3S,5S)-1-(cyclohexylmethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-(1H-indol-3-yl)acetamide
- 25 • N-(((3S,5S)-1-(cyclohexylmethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)quinoline-3-carboxamide;

or a pharmaceutically acceptable salt or prodrug thereof.

98. A method according to any one of claims 1 to 43 or a use according to any one of claims 44 to 96 wherein the compound of formula (I) is selected from the group consisting of:

- 3,4-dichloro-N-(((3S,5S)-3-(2-(4,4-difluoropiperidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide;
- 3,4-dichloro-N-(((3S,5S)-3-(2-hydroxyethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide;


- 3,4-dichloro-N-(((3S,5S)-3-(2-(3,3-difluoropiperidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide;
- (3S,5S)-5-((3,4-dichlorobenzylamino)methyl)-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-2-one;
- 5 • 3,4-dichloro-N-(((3S,5S)-1-(2-cyclopropylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide;
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2-cyclopropylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide;
- 6-chloro-N-(((3S,5S)-1-(2-cyclopropylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 10 • 3,4-dichloro-N-(((3S,5S)-3-(2-(2,5-dioxopyrrolidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide;
- 6-chloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(3-ureidopropyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 15 • 3,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(1,1,1-trifluoropropan-2-ylamino)ethyl)-1,4-diazepan-5-yl)methyl)benzamide;
- 3,4-dichloro-N-(((3S,5S)-3-(2-(3,3-dimethyl-2,5-dioxopyrrolidin-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)benzamide;
- N-(((3S,5S)-3-(2-(azepan-1-yl)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide;
- 20 • 6-chloro-N-(((3S,5S)-3-(2-(3-isopropylureido)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)biphenyl-4-carboxamide;
- 25 • N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-phenylthiazole-4-carboxamide;
- 4'-chloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)biphenyl-2-carboxamide;
- 6-chloro-N-(((3S,5S)-3-(2-(N-isopropylacetamido)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 30 • 6-chloro-N-(((3S,5S)-3-((isopropylamino)methyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 6-chloro-N-(((3S,5S)-3-(guanidinomethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;

- 2-(2,4-dichlorophenyl)-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acetamide;
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2,4-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide;
- 5 • 3,4-dichloro-N-(((3S,5S)-1-(2,4-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide;
- 3,4-dichloro-N-(((3S,5S)-1-(2,4-dichlorobenzyl)-3-(2-(methylsulfonamido)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide;
- 3,4-dichloro-N-(((3S,5S)-1-(2,4-dichlorobenzyl)-3-(2-(4-methylphenylsulfonamido)ethyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide;
- 10 • N-(((3S,5S)-3-(2-((S)-2-amino-3-methylbutanamido)ethyl)-1-(2,4-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide;
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2,4-dichlorobenzyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide;
- 15 • 6-chloro-N-(((3S,5S)-1-(2,4-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- N-(((3S,5S)-3-(2-aminoethyl)-2-oxo-1-(2-(thiophen-3-yl)butyl)-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide;
- 6-chloro-N-(((3S,5S)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1-((R)-2-(thiophen-3-yl)butyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 20 • 6-chloro-N-(((3S,5S)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1-((S)-2-(thiophen-3-yl)butyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethyl-2-methylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-6-chloro-2-naphthamide;
- 25 • 6-chloro-N-(((3S,5S)-1-(2-ethyl-2-methylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 6-chloro-N-(((3S,5S)-1-(2,2-diphenylethyl)-3-(2-morpholinoethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 6-chloro-N-(((3S,5S)-3-(2-morpholinoethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 30 • 6-chloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-3-(2-morpholinoethyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 3,4-dichloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-3-(2-morpholinoethyl)-2-oxo-1,4-diazepan-5-yl)methyl)benzamide;

- N-(3,4-dichlorobenzyl)-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acetamide;
- 1-(4-chlorobenzyl)-3-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)urea;
- 5 • N-(((3S,5S)-3-(2-aminoethyl)-1-(2-ethyl-2-methylbutyl)-2-oxo-1,4-diazepan-5-yl)methyl)-3,4-dichlorobenzamide;
- 3,4-dichloro-N-(((3S,5S)-1-(2-ethyl-2-methylbutyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide;
- 6-chloro-N-(((3S,5S)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1-(2,3,5-trichlorobenzyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 10 • 6-chloro-N-(((3S,5S)-3-(2-(1-methylethylsulfonamido)ethyl)-2-oxo-1-((S)-2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- butyl 2-((2S,7S)-7-((6-chloro-2-naphthamido)methyl)-3-oxo-4-((S)-2-phenylbutyl)-1,4-diazepan-2-yl)ethylcarbamate;
- 15 • (S)-6-chloro-N-((3-(1-isopropylpiperidin-4-yl)-2-oxo-1-(2-phenylbutyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 6-chloro-N-(((3S,5S)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 5-(4-chlorophenyl)-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)isoxazole-3-carboxamide;
- 20 • 2,4-dichloro-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide;
- N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-6-methoxy-2-naphthamide;
- 25 • 6-chloro-N-([5-¹³C,4-¹⁵N](3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)[¹³C]methyl)-2-naphthamide;
- N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-1-methoxy-2-naphthamide;
- (E)-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-3-(4-(trifluoromethoxy)phenyl)acrylamide;
- 30 • 5-(4-chlorophenyl)-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)isoxazole-3-carboxamide;
- 2,4-dichloro-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)benzamide;

- 5,6-dichloro-2-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)isoindoline-1,3-dione;
- (E)-N-(((3S,5S)-1-(3,5-dichlorobenzyl)-2-oxo-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)acrylamide;
- 5 • 6-methoxy-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 1-methoxy-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 10 • (E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)-N-(((3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)acrylamide;
- 6-chloro-N-([5,6,6-²H₃](3S,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)[²H₂]methyl)-2-naphthamide;
- 6-chloro-N-(((3R,5R)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 15 • 6-chloro-N-(((3S,5R)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 6-chloro-N-(((3R,5S)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 20 • 6-chloro-N-(((3S,5R)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 6-chloro-N-(((3R,5S)-2-oxo-1-((R)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 25 • 6-chloro-N-(((3R,5R)-2-oxo-1-((S)-2-phenylbutyl)-3-(2-(piperidin-1-yl)ethyl)-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- N-(((3R,5S)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- N-(((3R,5R)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide;
- 30 • N-(((3S,5R)-1-(2,2-diphenylethyl)-3-(3-guanidinopropyl)-2-oxo-1,4-diazepan-5-yl)methyl)-2-naphthamide;

or a pharmaceutically acceptable salt or prodrug thereof.



Formula (I)