



US 20130122024A1

(19) **United States**(12) **Patent Application Publication**
LERCHEN et al.(10) **Pub. No.: US 2013/0122024 A1**(43) **Pub. Date: May 16, 2013**(54) **NEW BINDER-DRUG CONJUGATES (ADCs)**
AND USE THEREOF(75) Inventors: **Hans-Georg LERCHEN**, Leverkusen (DE); **Beatrix STELTE-LUDWIG**, Wülfrath (DE); **Sven GOLFIER**, Berlin (DE); **Charlotte Christine KOPITZ**, Berlin (DE); **Rudolf BEIER**, Berlin (DE); **Sherif EL SHEIKH**, Essen (DE); **Iring HEISLER**, Wuppertal (DE); **Axel HARRENGA**, Wuppertal (DE); **Karl-Heinz THIERAUCH**, Berlin (DE); **Joachim SCHUHMACHER**, Wuppertal (DE); **Sandra BRUDER**, Leverkusen (DE); **Simone GREVEN**, Dormagen (DE); **Lars LINDEN**, Berlin (DE); **Jörg WILLUDA**, Berlin (DE); **Christoph MAHLERT**, Wuppertal (DE); **Heike PETRUL**, Berlin (DE); **Hannah JÖRIBEN**, Essen (DE); **Sandra Borkowski**, Neuendorf (DE)

Jun. 1, 2011	(EP)	11168556.6
Jun. 1, 2011	(EP)	11168557.4
Jun. 1, 2011	(EP)	11168558.2
Jun. 1, 2011	(EP)	11168559.0
Dec. 14, 2011	(EP)	11193609.2
Dec. 14, 2011	(EP)	11193618.3
Dec. 14, 2011	(EP)	11193621.7
Dec. 14, 2011	(EP)	11193623.3

Publication Classification(51) **Int. Cl.**
C07K 16/18 (2006.01)
A61K 39/395 (2006.01)
A61K 45/06 (2006.01)
C07K 7/06 (2006.01)
(52) **U.S. Cl.**
CPC . **C07K 16/18** (2013.01); **C07K 7/06** (2013.01);
A61K 39/395 (2013.01); **A61K 45/06** (2013.01)
USPC ... **424/179.1**; 530/391.9; 530/330; 530/389.1(73) Assignee: **Bayer Intellectual Property GmbH**,
Monheim (DE)(21) Appl. No.: **13/451,916**(22) Filed: **Apr. 20, 2012**(30) **Foreign Application Priority Data**

Apr. 21, 2011	(EP)	11163467.1
Apr. 21, 2011	(EP)	11163470.5
Apr. 21, 2011	(EP)	11163472.1
Apr. 21, 2011	(EP)	11163474.7

(57) **ABSTRACT**

The present application relates to new binder-drug conjugates (ADCs) of N,N-dialkylauristatins that are directed against the target mesothelin, to active metabolites of these ADCs, to processes for preparing these ADCs, to the use of these ADCs for treating and/or preventing illnesses, and also to the use of these ADCs for producing medicaments for treating and/or preventing illnesses, more particularly hyperproliferative and/or angiogenic diseases such as, for example, cancer diseases. Such treatments may be practised as a monotherapy or else in combination with other medicaments or further therapeutic measures.

NEW BINDER-DRUG CONJUGATES (ADCs) AND USE THEREOF

[0001] The file named SEQUENCE-LISTING.txt submitted herewith that is 327 kilobytes and was created on Apr. 20, 2012 is hereby incorporated by reference herein in its entirety.

[0002] The present application relates to new binder-drug conjugates (ADCs) of N,N-dialkylauristatins that are directed against the target mesothelin, to active metabolites of these ADCs, to processes for preparing these ADCs, to the use of these ADCs for treating and/or preventing illnesses, and also to the use of these ADCs for producing medicaments for treating and/or preventing illnesses, more particularly hyper-proliferative and/or angiogenic diseases such as, for example, cancer diseases. Such treatments may be practised as a monotherapy or else in combination with other medicaments or further therapeutic measures.

[0003] Cancer diseases are the consequence of uncontrolled cell growth in a wide variety of tissues. In many cases the new cells penetrate existing tissue (invasive growth), or they metastasize into remote organs. Cancer diseases occur in a wide variety of organs, and the illnesses often progress in a tissue-specific manner. The designation "cancer disease" as a generic term therefore describes a large group of defined diseases of different organs, tissues and cell types.

[0004] Early-stage tumours may be able to be removed by surgical and radiotherapeutic measures. Metastasized tumours can generally only be given palliative therapy by means of chemotherapeutic agents. The objective in that case is to achieve the optimum combination of improving quality of life and prolonging remaining lifetime.

[0005] The majority of the chemotherapeutic agents which are presently administered parenterally are often not target-directed at the tumour tissue or the tumour cells, but instead, as a result of their systemic administration, are distributed non-specifically within the body, hence including at locations at which exposure to the drug is undesirable, such as in healthy cells, tissues and organs, for example. This may lead to unwanted side-effects and even to serious effects of general toxicity, which then often greatly limit the therapeutically useful dose range of the drug, or necessitate complete cessation of medication.

[0006] The improved and selective availability of these chemotherapeutic agents in the tumour cell or the immediately surrounding tissue, and the associated boost in effect, on the one hand, and minimization of toxic side-effects, on the other hand, have therefore been a focal point for a number of years in the development of new chemotherapeutic agents. Many attempts have been made to date to develop efficient methods of introducing the drug into the target cell. Optimizing the association between drug and intracellular target and minimizing the intercellular distribution of drug, to adjacent cells, for example, nevertheless continue to constitute a difficult problem.

[0007] Monoclonal antibodies, for example, are suitable for the target-directed addressing of tumour tissue and tumour cells. The significance of such antibodies for the clinical treatment of cancer diseases has seen a considerable general increase in recent years, based on the activity of such agents as trastuzumab (Herceptin), rituximab (Rituxan), cetuximab (Erbix) and bevacizumab (Avastin), which have since been approved for the therapy of individual, specific tumour diseases [see e.g. G. P. Adams and L. M. Weiner, *Nat. Biotechnol.* 23, 1147-1157 (2005)]. Consequently there has also been a marked increase in interest in so-called immuno-conjugates such as, for example, the aforementioned ADCs, in which an internalizing antibody directed against a tumour-associated antigen is joined covalently via a linking unit ("linker") to a cytotoxic agent. Following introduction of the ADC into the tumour cell and subsequent cleavage of the conjugate, either the cytotoxic agent itself or another metabolite with cytotoxic activity, formed from the cytotoxic agent, is released within the tumour cell, where it is able to develop its effect directly and selectively. In this way it would be

possible to keep the damage to normal tissue within significantly closer limits in comparison to a conventional chemotherapy of the cancer disease [see e.g. J. M. Lambert, *Curr. Opin. Pharmacol.* 5, 543-549 (2005); A. M. Wu and P. D. Senter, *Nat. Biotechnol.* 23, 1137-1146 (2005); P. D. Senter, *Curr. Opin. Chem. Biol.* 13, 235-244 (2009); L. Ducry and B. Stump, *Bioconjugate Chem.* 21, 5-13 (2010)].

[0008] Instead of antibodies, it is also possible for binders from the small-molecule drug sphere to be used as binders which bind selectively to a specific target location ("target"), such as to a receptor, for example [see e.g. E. Ruoslahti et al., *Science* 279, 377-380 (1998); D. Karkan et al., *PLoS ONE* 3 (6), e2469 (Jun. 25, 2008)]. Also known are conjugates of cytotoxic drug and addressing ligand that exhibit a defined cleavage point between ligand and drug for the release of the drug. A "predetermined break point" of this kind may exist, for example, within a peptide chain which can be cleaved selectively at a particular site by a specific enzyme at the location of action [see e.g. R. A. Firestone and L. A. Telan, US Patent Application US 2002/0147138].


[0009] The activity of monoclonal antibodies is demonstrated in various types of cancer. Thus HERCEPTIN® and Erbitux® are used successfully in the treatment of HER2-positive breast cancer and EGFR-positive colorectal cancer, respectively.

[0010] The coupling of cytotoxic compounds to antibodies forms an extended possibility for additionally improving cancer therapy, since these conjugates allow tumour-specific toxophore accumulation and at the same time reduce the systemic toxicity. With respect to activity and tolerance, clinical studies with brentuximab vedotin in Hodgkin's lymphoma and with trastuzumab-DM1 in breast cancer have yielded highly promising results, which support the development of new antibodies and new ADCs against other tumour antigens.

[0011] The mesothelin precursor polypeptide is a glycosylated glycosylated cell surface protein which is cleaved proteolytically into a 30 kDa N-terminal secreted polypeptide (megakaryocyte potentiating factor, MPF) and a 40 kDa C-terminal polypeptide which is present on the cell membrane predominantly with GPI anchoring (mesothelin) (Chang, K. and I. Pastan, *Proc. Natl. Acad. Sci. USA*, (1996) 93(1):136).

[0012] On account of this surface expression and of the overexpression in a number of types of tumour, more particularly in ovarian carcinoma, in carcinoma of the pancreas and in mesotheliomas, and also on account of the highly limited expression in normal tissue, mesothelin represents a suitable antigen for monoclonal antibody therapy (Argani, P. et al., *Clin. Cancer Res.* (2001) 7(12): 3862; Hassan, R., et al., *Clin. Cancer Res.* (2004) 10(12 Pt 1):3937). The precise function of mesothelin is unknown to date; mesothelin gene-deficient mice do not exhibit any instantaneous anatomical, haematological or reproduction-restricting peculiarities (Bera, T. K. and I. Pastan, *Mol. Cell. Biol.* (2000) 20(8):2902). Mesothelin, however, is a highly variable protein, which is subject to posttranslational modifications such as proteolytic digestion and glycosylations (Hassan, R., et al., *Clin. Cancer Res.* (2004) 10(12 Pt 1): 3937). Therapeutic antibodies ought therefore to recognize an invariant epitope independently of the individual mesothelin modifications. Antibodies can be selected according to their desired properties, such as invariant affinity or internalization, for example, with the use of monoclonal antibodies guaranteeing that these properties on the one hand apply in respect of all antibodies and on the other hand can also be produced with long-term consistency.

[0013] Auristatin E (AE) and monomethylauristatin E (MMAE) are synthetic analogues of the dola-statins, a specific group of linear pseudopeptides which were originally isolated from marine sources and which have in some cases very potent cytotoxic activity with respect to tumour cells [for a review see e.g. G. R. Pettit, *Prog. Chem. org. Nat. Prod.* 70, 1-79 (1997); G. R. Pettit et al., *Anti-Cancer Drug Design* 10, 529-544 (1995); G. R. Pettit et al., *Anti-Cancer Drug Design* 13, 243-277 (1998)].



[0014] MMAE, however, possesses the disadvantage of a comparatively high systemic toxicity. For improving tumour selectivity, MMAE is used more particularly in conjunction with enzymatically cleavable valine-citrulline linkers in the ADC setting for more targeted tumour therapy [WO 2005/081711-A2; S. O. Doronina et al., *Bioconjugate Chem.* 17, 114-124 (2006)]. Following proteolytic cleavage, MMAE is released preferably intracellularly from corresponding ADCs.

[0015] When employed in the form of binder-drug conjugates (ADCs), however, MMAE is not compatible with linking units (linkers) between binder and drug that do not have an enzymatically cleavable predetermined break point [S. O. Doronina et al., *Bioconjugate Chem.* 17, 114-124 (2006)].


[0016] Monomethylauristatin F (MMAF) is an auristatin derivative having a C-terminal phenylalanine unit which exhibits only moderate antiproliferative activity in comparison to MMAE. This fact is very probably attributable to the free carboxyl group, whose polarity and charge adversely affect the capacity of this compound to access cells. In this connection, the methyl ester of MMAF (MMAF-OMe) has been described, as a neutral-charged prodrug derivative with cell access capability, which, in comparison to MMAF, has an in vitro cytotoxicity for various carcinoma cell lines that is increased by a number of orders of magnitude [S. O. Doronina et al., *Bioconjugate Chem.* 17, 114-124 (2006)]. It can be assumed that this effect is brought about by MMAF itself, which, following uptake of the prodrug into the cells, is rapidly released by intracellular ester hydrolysis.

[0017] However, drug compounds based on simple ester derivatives are generally subject to the risk of chemical instability on account of non-specific ester hydrolysis, independent of the intended site of action, by means, for example, of esterases that are present in the blood plasma; this non-specific hydrolysis may significantly restrict the usefulness of such compounds in therapy.


[0018] Monomethylauristatin F (MMAF) and also various ester derivatives and amide derivatives thereof have been disclosed in WO 2005/081711-A2. Further auristatin analogues with a C-terminal, amidically substituted phenylalanine unit are described in WO 01/18032-A2. WO 02/088172-A2 and WO 2007/008603-A1 claim MMAF analogues which relate to side-chain modifications of the phenylalanine, while WO 2007/008848-A2 claims those in which the carboxyl group of the phenylalanine has been modified. Auristatin conjugates linked via the C-terminus have been recently described in WO 2009/117531-A1 [see also S. O. Doronina et al., *Bioconjugate Chem.* 19, 1960-1963 (2008)].

[0019] Furthermore, auristatin derivatives such as MMAE and MMAF are also substrates for transporter proteins which are expressed by many tumour cells, and this may lead to the development of resistance to these drugs.

[0020] The problem addressed with the present invention was that of providing new binder-drug conjugates (ADCs) which, through combination of new N,N-dialkylauristatin derivatives with innovative, suitable linkers and binder, exhibit a very attractive activity profile, such as, for example, in terms of their specific tumour effect and/or the reduced potential of the metabolites formed intracellularly to be a substrate with respect to transporter proteins, and which are therefore suitable for the treatment and/or prophylaxis of hyperproliferative and/or angiogenic diseases, such as cancer diseases, for example.



[0021] The present invention provides binder-drug conjugates of the general formula (Ia)



in which

[0022] n is a number from 1 to 50,

[0023] AK is a binder,


[0024] the group $\text{-G-L}^1\text{-B-L}^2\text{-}\S\S$ is a linker,

[0025] where

[0026] $\S$ marks the linkage site with the group AK and

[0027] $\S\S$ marks the linkage site with the nitrogen atom,

[0028] D is a group of the formula



[0029] where


[0030] #³ marks the linkage site with the nitrogen atom,

[0031] R¹ is hydrogen or methyl,

[0032] R² is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

[0033] or

[0034] R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[0035] in which

[0036] #⁴ marks the linkage site with the adjacent nitrogen atom,

[0037] #⁵ marks the linkage site with the carbonyl group,

[0038] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[0039] in which

[0040] #⁶ marks the linkage site with the carbonyl group,


[0041] R⁶ is hydrogen, hydroxy or benzyloxy,

[0042] R³ is hydrogen or methyl,

[0043] R⁴ is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

[0044] or

[0045] R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[0046] in which

[0047] #⁷ marks the linkage site with the adjacent nitrogen atom,


[0048] #⁸ marks the linkage site with the group T¹,

[0049] T¹ is a group of the formula —C(=O)—OR^7 , $\text{—C(=O)—NR}^8\text{R}^9$, $\text{—C(=O)—NH—NH—R}^{10}$ or $\text{—CH}_2\text{—O—R}^{11}$,

[0050] in which

[0051] R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

- [0052] R^8 is hydrogen or methyl,
 [0053] R^9 is hydrogen, methyl, ethyl, n-propyl or benzyl,
 [0054] or
 [0055] R^8 and R^9 together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,
 [0056] R^{10} is benzoyl,
 [0057] R^{11} is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,
 [0058] R^5 is hydrogen, methyl or a group of the formula



- [0059] in which
 [0060] $\#^9$ marks the linkage site with $-\text{CHC}(R^{26})-\text{T}^2$,
 [0061] R^{12} is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula $-\text{S}(\text{O})_2\text{OH}$,
 [0062] R^{13} is phenyl which may be substituted by methoxycarbonyl or carboxyl,
 [0063] R^{26} is hydrogen or hydroxy,
 [0064] T^2 is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,
 [0065] R^{35} is methyl or hydroxy,
 and also their salts, solvates and solvates of the salts.

[0066] Compounds of the invention are the compounds of the formula (Ia) and (I) and their salts, solvates and solvates of the salts, the compounds of the formulae identified below and encompassed by formula (Ia) and (I), and their salts, solvates and solvates of the salts, and also the compounds identified below as working examples and encompassed by formula (Ia) and (I), and their salts, solvates and solvates of the salts, to the extent that the compounds identified below and encompassed by formula (Ia) and (I) are not already salts, solvates and solvates of the salts.

[0067] Depending on their structure, the compounds of the invention may exist in different stereoisomeric forms, i.e. in the form of configurational isomers or else where appropriate as conformational isomers (enantiomers and/or diastereoisomers, including those in the case of atropisomers). The present invention therefore encompasses the enantiomers and diastereomers and their respective mixtures. The stereoisomerically homogeneous constituents can be isolated from such mixtures of enantiomers and/or diastereomers in a known way; for this purpose it is preferred to use chromatographic processes, more particularly HPLC chromatography on an achiral or chiral phase.

[0068] Where the compounds of the invention can occur in tautomeric forms, the present invention encompasses all of the tautomeric forms.

[0069] The present invention also encompasses all suitable isotopic variants of the compounds of the invention. An isotopic variant of a compound of the invention is understood here to mean a compound in which at least one atom within the compound of the invention has been exchanged for another atom of the same atomic number but with a different atomic mass from the atomic mass which occurs commonly

or predominantly in nature. Examples of isotopes which can be incorporated into an inventive compound are those of hydrogen, carbon, nitrogen, oxygen, phosphorus, sulphur, fluorine, chlorine, bromine and iodine such as ^2H (deuterium), ^3H (tritium), ^{13}C , ^{14}C , ^{15}N , ^{17}O , ^{18}O , ^{32}P , ^{33}P , ^{34}S , ^{36}S , ^{18}F , ^{36}Cl , ^{82}Br , ^{123}I , ^{124}I , ^{129}I and ^{131}I . Particular isotope variants of a compound of the invention, such as more particularly those in which one or more radioactive isotopes are incorporated, may be of benefit, for example, for investigating the mechanism of action or the distribution of drug in the body; owing to the comparative ease of preparation and detectability, compounds labelled with ^3H or ^{14}C isotopes are especially suitable for these purposes. Furthermore, the incorporation of isotopes, such as of deuterium, for example, may lead to certain therapeutic advantages as a consequence of greater metabolic stability of the compound, such as an extension to the half-life in the body or a reduction in the active dose required, for example; such modifications of the compounds of the invention may therefore, where appropriate, also constitute a preferred embodiment of the present invention. Isotopic variants of the compounds of the invention can be prepared by the processes known to the skilled person, as for example in accordance with the methods described later on below and the procedures reproduced in the working examples, by using corresponding isotopic modifications of the respective reagents and/or starting compounds.

[0070] Preferred salts in the context of the present invention are physiologically acceptable salts of the compounds of the invention. Also encompassed are salts which although themselves not suitable for pharmaceutical applications may nevertheless be used, for example, for isolating or purifying the compounds of the invention.

[0071] Physiologically acceptable salts of the compounds of the invention encompass acid addition salts of mineral acids, carboxylic acids and sulphonic acids, examples being salts of hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, methanesulphonic acid, ethanesulphonic acid, benzenesulphonic acid, toluenesulphonic acid, naphthalenedisulphonic acid, acetic acid, trifluoroacetic acid, propionic acid, lactic acid, tartaric acid, malic acid, citric acid, fumaric acid, maleic acid and benzoic acid.

[0072] Physiologically acceptable salts of the compounds of the invention also encompass salts of customary bases, such as, by way of example and preferably, alkali metal salts (e.g. sodium and potassium salts), alkaline earth metal salts (e.g. calcium and magnesium salts) and ammonium salts, derived from ammonia or organic amines having 1 to 16 C atoms, such as, by way of example and preferably, ethylamine, diethylamine, triethylamine, ethyldiisopropylamine, monoethanolamine, diethanolamine, triethanolamine, dicyclohexylamine, dimethylaminoethanol, procaine, dibenzylamine, N-methylpiperidine, N-methylmorpholine, arginine, lysine and 1,2-ethylenediamine.

[0073] Solvates in the context of the invention are those forms of the compounds of the invention that form a complex in the solid or liquid state through coordination with solvent molecules. Hydrates are one specific form of solvates, in which the coordination takes place with water. Preferred solvates in the context of the present invention are hydrates.

[0074] Furthermore, the present invention also encompasses prodrugs of the compounds of the invention. The term "prodrugs" here identifies compounds which may themselves be biologically active or inactive but are converted during

their residence in the body into compounds of the invention (by metabolism or hydrolysis, for example).

[0075] In the context of the present invention the definitions of the substituents, unless otherwise specified, are as follows: (C₁-C₄)-Alkyl in the context of the invention is a linear or branched alkyl radical having 1 to 4 carbon atoms. By way of example and with preference, the following may be mentioned: methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, 1-methylpropyl and tert-butyl.

(C₁-C₄)-Alkylcarbonyl in the context of the invention is a linear or branched alkyl radical having 1 to 4 carbon atoms which is linked via a carbonyl group. By way of example and with preference, the following may be mentioned: methylcarbonyl, ethylcarbonyl, n-propylcarbonyl, isopropylcarbonyl, n-butylcarbonyl, isobutylcarbonyl, 1-methylpropylcarbonyl and tert-butylcarbonyl.

Alkanediyl in the context of the invention is a linear, α,ω -divalent alkyl radical having the particular number of carbon atoms indicated. By way of example and of preference, the following may be mentioned: methylene, ethane-1,2-diyl (1,2-ethylene), propane-1,3-diyl (1,3-propylene), butane-1,4-diyl (1,4-butylene), pentane-1,5-diyl (1,5-pentylene), hexane-1,6-diyl (1,6-hexylene), heptane-1,7-diyl (1,7-hexylene), octane-1,8-diyl (1,8-octylene), nonane-1,9-diyl (1,9-nonylene), decane-1,10-diyl (1,10-decylene).

(C₃-C₇)-Cycloalkyl and 3- to 7-membered carbocycle respectively in the context of the invention is a monocyclic, saturated cycloalkyl group having 3 to 7 carbon atoms. By way of example and of preference, the following may be mentioned: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

[0076] The side group of an α -amino acid in the definition of R¹⁹ encompasses not only the side groups of the naturally occurring α -amino acids but also the side groups of homologues and isomers of these α -amino acids. The α -amino acid here may be in the L or D configuration or else may be present as a mixture of the L and D forms. Examples that may be given of side groups are as follows: methyl (alanine), propan-2-yl (valine), propan-1-yl (norvaline), 2-methylpropan-1-yl (leucine), 1-methylpropan-1-yl (isoleucine), butan-1-yl (norleucine), tert-butyl (2-tert-butylglycine), phenyl (2-phenylglycine), benzyl (phenylalanine), p-hydroxybenzyl (tyrosine), indol-3-ylmethyl (tryptophan), imidazol-4-ylmethyl (histidine), hydroxymethyl (serine), 2-hydroxyethyl (homoserine), 1-hydroxyethyl (threonine), mercaptomethyl (cysteine), methylthiomethyl (S-methylcysteine), 2-mercaptoethyl (homocysteine), 2-methylthio ethyl (methionine), carbamoylmethyl (asparagine), 2-carbamoyl ethyl (glutamine), carboxymethyl (aspartic acid), 2-carboxyethyl (glutamic acid), 4-aminobutan-1-yl (lysine), 4-amino-3-hydroxybutan-1-yl (hydroxylysine), 3-aminopropan-1-yl (ornithine), 2-aminoethyl (2,4-diaminobutyric acid), aminomethyl (2,3-diaminopropionic acid), 3-guanidinopropan-1-yl (arginine), 3-ureidopropan-1-yl (citrulline). Preferred α -amino acid side groups in the definition of R¹⁹ are methyl (alanine), propan-2-yl (valine), 2-methylpropan-1-yl (leucine), benzyl (phenylalanine), imidazol-4-ylmethyl (histidine), hydroxymethyl (serine), 1-hydroxyethyl (threonine), 4-aminobutan-1-yl (lysine), 3-aminopropan-1-yl (ornithine), 2-aminoethyl (2,4-diaminobutyric acid), aminomethyl (2,3-diaminopropionic acid), 3-guanidinopropan-1-yl (arginine). The L configuration is preferred in each case.

[0077] A 4- to 7-membered heterocycle in the context of the invention is a monocyclic, saturated heterocycle having a

total of 4 to 7 ring atoms, which contains one or two ring heteroatoms from the series N, O, S, SO and/or SO₂ and is linked via a ring carbon atom or optionally a ring nitrogen atom. Preference is given to a 5- to 7-membered heterocycle having one or two ring heteroatoms from the series N, O and/or S, more preferably a 5- or 6-membered heterocycle having one or two ring heteroatoms from the series N and/or O. By way of example, the following may be mentioned: azetidiny, oxetany, pyrrolidinyl, pyrazolidinyl, tetrahydrofuranyl, thiolanyl, piperidinyl, piperazinyl, tetrahydropyran-1-yl, tetrahydrothiopyran-1-yl, morpholinyl, thiomorpholinyl, hexahydroazepinyl and hexahydro-1,4-diazepinyl. Preference is given to pyrrolidinyl, tetrahydrofuranyl, piperidinyl, piperazinyl, tetrahydropyran-1-yl and morpholinyl.

[0078] In the formula of the group which may be represented by A, B, B¹, D, G, L¹, L², L⁴, L⁶, R¹, R², R³, R⁴ and R⁵, respectively, the end point of the line at which the symbol #⁶, *, **, #³, #¹, #², ##¹, ##², ##³, ##⁴, ##⁵, ##⁶, ##⁷, ##⁸, ***, ****, #⁴, #⁵, #⁶, #⁷, #⁸ or #⁹ is located is not a carbon atom or a CH₂ group, but instead is part of the bond to the atom designated in each case, to which the A, B, B¹, D, G, L¹, L², L⁴, L⁶, R¹, R², R³, R⁴ or R⁵ is bonded.

[0079] In the context of the present invention, all radicals which occur multiply have their definition independently of one another. If radicals in the compounds of the invention are substituted, the radicals, unless otherwise specified, may be substituted one or more times. Substitution by one or by two identical or different substituent(s) is preferred. Particularly preferred is substitution by one substituent.

[0080] In the context of the present invention the terms used, unless otherwise specified, have the following definitions:

[0081] The term "linker" is understood in the broadest sense as a chemical unit which comprises a covalent bond or a series of atoms that links a binder covalently to a drug. The term "linker" is understood preferably as a series of atoms in the sense of the present invention that links a binder covalently to a drug. Furthermore, linkers may be represented, for example, by divalent chemical units, such as alkylidyls, arylidyls, heteroarylidyls, heterocyclyldyls, dicarbonyl acid esters, dicarbonyl acid amides.

[0082] The term "binder" is understood in the broadest sense as a molecule which binds to a target molecule which is present on a particular target cell population to be addressed with the binder-drug conjugate. The term "binder" should be understood in its broadest interpretation and encompasses, for example, lectins, proteins which are able to bind particular sugar chains, or phospholipid-binding proteins. Such binders comprise, for example, high molecular mass proteins (binding proteins), polypeptides or peptides (binding peptides), non-peptidic (e.g. aptamers (U.S. Pat. No. 5,270,163) review article by Keefe A D., et al., Nat. Rev. Drug Discov. 2010; 9:537-550), or vitamins) and all other cell-binding molecules or substances. Binding proteins are, for example, antibodies and antibody fragments or antibody mimetics such as, for example, affibodies, adnectins, anticalins, DARPins, avimers, nanobodies (review articles by Gebauer M. et al., Curr. Opinion in Chem. Biol. 2009; 13:245-255; Nuttall S. D. et al., Curr. Opinion in Pharmacology 2008; 8:608-617). Binding peptides are, for example, ligands of a ligand-receptor pair, such as VEGF in the ligand-receptor pair VEGF/KDR, such as transferrin of the ligand-receptor pair transfer-

rin/transferrin receptor, or cytokines/cytokine receptor, such as TNFalpha in the ligand receptor pair TNFalpha/TNFalpha receptor.

[0083] Preferred binders in accordance with the invention are (more particularly human, monoclonal) antibodies or antigen-binding antibody fragments which bind to mesothelin. In the case of anti-mesothelin antibodies, *n*, in other words the number of toxophore molecules per antibody molecule, is preferably in the range from 1 to 10, more preferably 2 to 8.

[0084] A “target molecule” is understood in the broadest sense to be a molecule which is present in the target cell population, and may be a protein (e.g. a receptor of a growth factor) or a non-peptidic molecule (e.g. a sugar or phospholipid). Preferably it is a receptor or an antigen.

[0085] The term “extracellular” target molecule describes a target molecule which is attached to the cell and which is located on the outside of a cell or the part of a target molecule which is located on the outside of a cell, i.e. a binder may bind to an intact cell at its extracellular target molecule. An extracellular target molecule may be anchored in the cell membrane or may be part of the cell membrane. The skilled person knows of methods for identifying extracellular target molecules. For proteins this may be done via determination of the transmembrane domain(s) and the orientation of the protein in the membrane. This data is generally recorded in protein databases (e.g. SwissProt).

[0086] The term “cancer target molecule” describes a target molecule which is multiply present on one or more cancer cell types in comparison to non-cancer cells of the same tissue type. The cancer target molecule is preferably present selectively on one or more cancer cell types in comparison to non-cancer cells of the same tissue type, with “selectively” describing an at least twofold accumulation on cancer cells in comparison to non-cancer cells of the same tissue type (a “selective cancer target molecule”). The use of cancer target molecules allows selective therapy of cancer cells with the conjugates of the invention.

[0087] The binder may be linked via a bond to the linker. Known from the literature are various possibilities of covalent coupling (conjugation) of organic molecules to antibody. The linking of the binder may take place by means of a heteroatom of the binder. Inventive heteroatoms of the binder that may be used for linking are sulphur (in one embodiment via a sulphhydryl group of the binder), oxygen (in accordance with the invention by means of a carboxyl or hydroxy group of the binder) and nitrogen (in one embodiment via a primary or secondary amine group or amide group of the binder). Preferred in accordance with the invention is the conjugation of the toxophores to the antibody via one or more sulphur atoms of cysteine residues of the antibody and/or via one or more NH groups of lysine residues of the antibody. These heteroatoms may be present in the natural binder or may be introduced by means of methods of chemistry or molecular biology. In accordance with the invention, the linking of the binder to the toxophore has little influence over the binding activity of the binder to the target molecule. In a preferred embodiment the linking has no influence on the binding activity of the binder to the target molecule.

[0088] The term “antibody” is understood in accordance with the present invention in its broadest sense and encompasses immunoglobulin molecules, examples being intact or modified monoclonal antibodies, polyclonal antibodies or multispecific antibodies (e.g. bispecific antibodies). An

immunoglobulin molecule preferably comprises a molecule having four polypeptide chains, two heavy chains (H chains) and two light chains (L chains), which are linked typically by disulphide bridges. Each heavy chain comprises a variable domain of the heavy chain (abbreviated to VH) and a constant domain of the heavy chain. The constant domain of the heavy chain may encompass, for example, three domains CH1, CH2 and CH3. Each light chain comprises a variable domain (abbreviated to VL) and a constant domain. The constant domain of the light chain comprises one domain (abbreviated to CL). The VH and VL domains may be further subdivided into regions having hypervariability, also called complementarity-determining regions (abbreviated to CDR), and regions having a low sequence variability (“framework region”, abbreviated to FR). Each VH and VL region is typically composed of three CDRs and up to four FRs. For example, in the following order from the amino terminus to the carboxy terminus: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. An antibody may be obtained from any species suitable for the antibody, such as, for example, rabbit, lama, camel, mouse or rat. In one embodiment the antibody is of human or murine origin. An antibody may for example be human, humanized or chimeric.

[0089] The term “monoclonal” antibody identifies antibodies which have been obtained from a population of substantially homogeneous antibodies, i.e. individual antibodies of the population are identical except for naturally occurring mutations which may occur in small numbers. Monoclonal antibodies recognize a single antigenic binding site with a high specificity. The term “monoclonal antibody” does not refer to a particular production method.

[0090] The term “intact” antibody refers to antibodies which comprise not only an antigen-binding domain but also the constant domain of the light and heavy chain. The constant domain may be a naturally occurring domain, or a variant thereof in which a plurality of amino acid positions have been altered.

[0091] The term “modified intact” antibody refers to intact antibodies which have been fused with another polypeptide or protein, not originating from an antibody, via the amino terminus or carboxyl terminus thereof, by means of a covalent bond (e.g. a peptide linkage). Furthermore, antibodies may be modified by introducing reactive cysteines at defined locations, in order to facilitate coupling to a toxophore (see Junutula et al. *Nat. Biotechnol.* 2008 August; 26(8):925-32).

[0092] The term “human” antibody identifies antibodies which can be obtained from a human being or are synthetic human antibodies. A “synthetic” human antibody is an antibody which in parts or as a whole is obtainable from synthetic sequences *in silico* which are based on the analysis of human antibody sequences. A human antibody may be encoded, for example, by a nucleic acid which has been isolated from a library of antibody sequences which are of human origin. One example of such antibodies can be found in Soderlind et al., *Nature Biotech.* 2000, 18:853-856.

[0093] The term “humanized” or “chimeric” antibody describes antibodies which consist of a non-human and of a human sequence component. In these antibodies, part of the sequences of the human immunoglobulin (recipient) is replaced by sequence components of a non-human immunoglobulin (donor). In many cases the donor is a murine immunoglobulin. With humanized antibodies, amino acids of the CDR in the recipient are replaced by amino acids of the donor. In some cases, amino acids of the framework as well are

replaced by corresponding amino acids of the donor. In some cases the humanized antibody contains amino acids which were present neither in the recipient nor in the donor and which were inserted during the optimization of the antibody. In the case of chimeric antibodies, the variable domains of the donor immunoglobulin are fused with the constant regions of a human antibody.

[0094] The term complementarity-determining region (CDR) as used here refers to those amino acids in a variable antibody domain that are necessary for binding to the antigen. Every variable region typically has three CDR regions, identified as CDR1, CDR2 and CDR3. Each CDR region may comprise amino acids according to the definition of Kabat and/or amino acids of a hypervariable loop, defined according to Chotia. The definition according to Kabat encompasses, for example, the region of approximately amino acid position 24-34 (CDR1), 50-56 (CDR2) and 89-97 (CDR3) of the variable light chain and 31-35 (CDR1), 50-65 (CDR2) and 95-102 (CDR3) of the variable heavy chain (Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991)). The definition according to Chotia encompasses, for example, the region of approximately amino acid position 26-32 (CDR1), 50-52 (CDR2) and 91-96 (CDR3) of the variable light chain and 26-32 (CDR1), 53-55 (CDR2) and 96-101 (CDR3) of the variable heavy chain (Chotia and Lesk; J Mol Biol 196: 901-917 (1987)). In some cases a CDR may comprise amino acids from one CDR region as defined by Kabat and Chotia.

[0095] Depending on the amino acid sequence of the constant domain of the heavy chain, antibodies may be divided into different classes. There are five main classes of intact antibodies: IgA, IgD, IgE, IgG and IgM, and a number of them may be broken down into further subclasses (isotypes), e.g. IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2. The constant domains of the heavy chain that correspond to the different classes are identified as $[\alpha/\alpha]$, $[\delta/\delta]$, $[\epsilon/\epsilon]$, $[\gamma/\gamma]$ and $[\mu/\mu]$. Both the three-dimensional structure and the subunit structure of antibodies are known.

[0096] The term "functional fragment" or "antigen-binding antibody fragments" of an antibody/immunoglobulin is defined as a fragment of an antibody/immunoglobulin (e.g. the variable domains of an IgG) which further encompasses the antigen binding domains of the antibody/immunoglobulin. The "antigen binding domain" of an antibody typically encompasses one or more hypervariable regions of an antibody, e.g. the CDR, CDR2 and/or CDR3 region. However, the "framework" or "scaffold" region of an antibody may also play a part with regard to the binding of the antibody to the antigen. The framework region forms the scaffold for the CDRs. The antigen-binding domain preferably encompasses at least amino acids 4 to 103 of the variable light chain and amino acid 5 to 109 of the variable heavy chain, more preferably amino acid 3 to 107 of the variable light chain and 4 to 111 of the variable heavy chain, particular preference being given to the complete variable light and heavy chains, i.e. amino acid 1-109 of the VL and 1 to 113 of the VH (numbering according to WO97/08320).

[0097] "Functional fragments" or "antigen-binding antibody fragments" of the invention encompass, non-conclusively, Fab, Fab', F(ab')₂ and Fv fragments, diabodies, Single Domain Antibodies (DABs), linear antibodies, individual chains of antibodies (single-chain Fv, abbreviated to ScFv); and multispecific antibodies, such as bi and tri-specific anti-

bodies, for example, formed from antibody fragments C. A. K. Borrebaeck, editor (1995) Antibody Engineering (Breakthroughs in Molecular Biology), Oxford University Press; R. Kontermann & S. Duebel, editors (2001) Antibody Engineering (Springer Laboratory Manual), Springer Verlag). Antibodies other than "multispecific" or "multifunctional" antibodies are those having identical binding sites. Multispecific antibodies may be specific for different epitopes of an antigen or may be specific for epitopes of more than one antigen (see, for example WO 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, et al., 1991, J. Immunol. 147:60 69; U.S. Pat. Nos. 4,474,893; 4,714,681; 4,925,648; 5,573,920; 5,601,819; or Kostelny et al., 1992, J. Immunol. 148: 1547 1553). An F(ab')₂ or Fab molecule may be constructed such that the number of intermolecular disulphide interactions occurring between the Ch1 and the CL domains can be reduced or else completely prevented.

[0098] "Functional fragments" or "antigen-binding antibody fragments" may be fused with another polypeptide or protein, not originating from an antibody, via the amino terminus or carboxyl terminus thereof, by means of a covalent bond (e.g. a peptide linkage). Furthermore, antibodies and antigen-binding fragments may be modified by introducing reactive cysteines at defined locations, in order to facilitate coupling to a toxophore (see Junutula et al. Nat. Biotechnol. 2008 August; 26(8):925-32).

[0099] Polyclonal antibodies can be prepared by methods known to a person of ordinary skill in the art. Monoclonal antibodies may be prepared by methods known to a person of ordinary skill in the art (Köhler and Milstein, Nature, 256, 495-497, 1975). Human and humanized monoclonal antibodies may be prepared by methods known to a person of ordinary skill in the art (Olsson et al., Meth Enzymol. 92, 3-16 or Cabilly et al U.S. Pat. No. 4,816,567 or Boss et al U.S. Pat. No. 4,816,397).

[0100] A person of ordinary skill in the art is aware of diverse methods for preparing human antibodies and fragments thereof, such as, for example, by means of transgenic mice (N Lonberg and D Huszar, Int Rev Immunol. 1995; 13(1):65-93) or Phage Display Technologies (Clackson et al., Nature. 1991 Aug. 15; 352(6336):624-8). Antibodies of the invention may be obtained from recombinant antibody libraries consisting for example of the amino acid sequences of a multiplicity of antibodies compiled from a large number of healthy volunteers. Antibodies may also be produced by means of known recombinant DNA technologies. The nucleic acid sequence of an antibody can be obtained by routine sequencing or is available from publically accessible databases.

[0101] An "isolated" antibody or binder has been purified to remove other constituents of the cell. Contaminating constituents of a cell which may interfere with a diagnostic or therapeutic use are, for example, enzymes, hormones, or other peptidic or non-peptidic constituents of the cell. A preferred antibody or binder is one which has been purified to an extent of more than 95%, relative to the antibody or binder (determined for example by Lowry method, UV-Vis spectroscopy or by SDS capillary gel electrophoresis). Furthermore an antibody which has been purified to such an extent that it is possible to determine at least 15 amino acids of the amino terminus or of an internal amino acid sequence, or which has been purified to homogeneity, the homogeneity being determined by SDS-PAGE under reducing or non-reducing conditions (detection may be determined by means of Coomassie

Blau staining or preferably by silver coloration). However, an antibody is normally prepared by one or more purification steps.

[0102] The term “specific binding” or “binds specifically” refers to an antibody or binder which binds to a predetermined antigen/target molecule. Specific binding of an antibody or binder typically describes an antibody or binder having an affinity of at least 10^{-7} M (as Kd value; i.e. preferably those with smaller Kd values than 10^{-7} M), with the antibody or binder having an at least two times higher affinity for the predetermined antigen/target molecule than for a non-specific antigen/target molecule (e.g. bovine serum albumin, or casein) which is not the predetermined antigen/target molecule or a closely related antigen/target molecule.

[0103] Antibodies which are specific against a cancer cell antigen can be prepared by a person of ordinary skill in the art by means of methods with which he or she is familiar (such as recombinant expression, for example) or may be acquired commercially (as for example from Merck KGaA, Germany). Examples of known commercially available antibodies in cancer therapy are Erbitux® (cetuximab, Merck KGaA), Avastin® (bevacizumab, Roche) and Herceptin® (trastuzumab, Genentech). Trastuzumab is a recombinant humanized monoclonal antibody of the IgG1kappa type which in a cell-based assay (Kd=5 nM) binds the extracellular domains of the human epidermal growth receptor with high affinity. The antibody is produced recombinantly in CHO cells.

[0104] The compounds of the formula (I) represent a subgroup of the compounds of the formula (Ia).

[0105] A preferred subject of the invention are binder-drug conjugates of the general formula (Ia) in which

[0106] n is a number from 1 to 50,


[0107] AK is AK₁ or AK₂

[0108] where

[0109] AK₁ is a binder (preferably an anti-mesothelin antibody) which is bonded via a sulphur atom of the binder to the group G,

[0110] AK₂ is a binder (preferably an anti-mesothelin antibody) which is bonded via a nitrogen atom of the binder to the group G,

[0111] G when AK=AK₁, is a group of the formula



[0112] where


[0113] #¹ marks the linkage site with the sulphur atom of the binder,

[0114] #² marks the linkage site with the group L¹,

[0115] or

[0116] when AK=AK₂, is carbonyl,

[0117] L¹ is a bond, linear (C₁-C₁₀)-alkanediyl, a group of the formula



[0118] where


[0119] m is a number from 2 to 6,

[0120] #¹ marks the linkage site with the group G,

[0121] #² marks the linkage site with the group B,

[0122] L^{1A} is linear (C₂-C₁₀)-alkanediyl,

[0123] B¹ is a group of the formula




[0124] in which

[0125] #⁵ marks the linkage site with the group L^{1A},

[0126] #⁶ marks the linkage site with the group L^{1B},

[0127] L⁵ is a bond or (C₂-C₄)-alkanediyl,

[0128] L⁶ is a bond or a group of the formula



[0129] in which

[0130] #⁷ marks the linkage site with the carbonyl group,

[0131] #⁸ marks the linkage site with L^{1B},

[0132] R³³ is hydrogen, (C₁-C₄)-alkylcarbonyl, tert-butyloxycarbonyl or benzyloxycarbonyl,

[0133] R³⁴ is hydrogen or methyl,

[0134] R²⁹ is hydrogen or (C₁-C₄)-alkyl,

[0135] R³⁰ is hydrogen or (C₁-C₄)-alkyl,

[0136] or

[0137] R²⁹ and R³⁰ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

[0138] R³¹ is hydrogen or (C₁-C₄)-alkyl,

[0139] R³² is hydrogen or (C₁-C₄)-alkyl,

[0140] or

[0141] R³¹ and R³² together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

[0142] L^{1B} is linear (C₂-C₁₀)-alkanediyl,


[0143] and

[0144] where (C₁-C₁₀)-alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,

[0145] and

[0146] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,

[0147] B is a bond or a group of the formula



[0148] where


[0149] * marks the linkage site with L¹,

[0150] ** marks the linkage site with L²,

[0151] P is O or NH,

[0152] L³ is a bond or (C₂-C₄)-alkanediyl,

[0153] L⁴ is a bond or a group of the formula



[0154] in which

[0155] *** marks the linkage site with the carbonyl group,

[0156] **** marks the linkage site with L²,

[0157] R²⁵ is hydrogen or methyl,

[0158] R²⁸ is hydrogen, (C₁-C₄)-alkylcarbonyl, tert-butyloxycarbonyl or benzyloxycarbonyl,

[0159] Q¹ is a 4- to 7-membered heterocycle,

[0160] Q² is a 3- to 7-membered carbocycle or a 4- to 7-membered heterocycle,

[0161] R¹⁴ is hydrogen or (C₁-C₄)-alkyl,

[0162] R¹⁵ is hydrogen or (C₁-C₄)-alkyl,

[0163] or

[0164] R¹⁴ and R¹⁵ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

[0165] R¹⁶ is hydrogen or (C₁-C₄)-alkyl,

[0166] R¹⁷ is hydrogen or (C₁-C₄)-alkyl,

[0167] or

[0168] R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

[0169] R¹⁸ is hydrogen or (C₁-C₄)-alkyl,

[0170] R¹⁹ is hydrogen or the side group of a natural α-amino acid or of its homologues or isomers,

[0171] R²⁰ is hydrogen or (C₁-C₄)-alkyl,

[0172] or

[0173] R¹⁹ and R²⁰ together with the atoms to which they are bonded form a pyrrolidinyl ring,

[0174] R²¹ is hydrogen or (C₁-C₄)-alkyl,

[0175] R²² is hydrogen or (C₁-C₄)-alkyl,

[0176] or

[0177] R²¹ and R²² together with the atoms to which they are bonded form a 3- to 7-membered carbocycle,

[0178] R²³ is (C₁-C₄)-alkyl,

[0179] R²⁴ is hydrogen or (C₁-C₄)-alkyl,

[0180] R²⁷ is hydrogen or (C₁-C₄)-alkyl,


[0181] R³⁶ is hydrogen, (C₁-C₄)-alkylcarbonyl, tert-butyloxycarbonyl or benzyloxycarbonyl,

[0182] R³⁷ is hydrogen or methyl,

[0183] or

[0184] R³⁶ and R³⁷ together with the atoms to which they are bonded form a pyrrolidine ring,

[0185] L² is linear (C₂-C₁₀)-alkanediyl or is a group of the formula



[0186] where

[0187] p is a number from 2 to 6,

[0188] ##3 marks the linkage site with the group B,

[0189] ##4 marks the linkage site with the nitrogen atom,

[0190] where (C₂-C₁₀)-alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,

[0191] and


[0192] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,

[0233] where

[0234] AK_1 is a binder which is bonded via a sulphur atom of the binder to the group G,

[0235] AK_2 is a binder which is bonded via a nitrogen atom of the binder to the group G,

[0236] G when $AK=AK_1$, is a group of the formula



[0237] in which


[0238] #1 marks the linkage site with the sulphur atom of the binder,

[0239] #2 marks the linkage site with the group L^1 ,

[0240] or

[0241] when $AK=AK_2$, is carbonyl,

[0242] L^1 is a bond, linear (C_1-C_{10}) -alkanediyl, a group of the formula



[0243] where


[0244] m is a number from 2 to 6,

[0245] ##1 marks the linkage site with the group G,

[0246] ##2 marks the linkage site with the group B,

[0247] L^{1A} is linear (C_2-C_{10}) -alkanediyl,

[0248] B^1 is a group of the formula




[0249] in which

[0250] ##5 marks the linkage site with the group L^{1A} ,

[0251] ##6 marks the linkage site with the group L^{1B} ,

[0252] L^5 is a bond or (C_2-C_4) -alkanediyl,

[0253] L^6 is a bond or a group of the formula



[0254] in which

[0255] ##7 marks the linkage site with the carbonyl group,

[0256] ##8 marks the linkage site with L^{1B} ,

[0257] R^{33} is hydrogen (C_1-C_4) -alkylcarbonyl, tert-butylloxycarbonyl or benzyloxycarbonyl,

[0258] R^{34} is hydrogen or methyl,

[0259] R^{29} is hydrogen or (C_1-C_4) -alkyl,

[0260] R^{30} is hydrogen or (C_1-C_4) -alkyl,

[0261] or

[0262] R^{29} and R^{30} together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

[0263] R^{31} is hydrogen or (C_1-C_4) -alkyl,

[0264] R^{32} is hydrogen or (C_1-C_4) -alkyl,

[0265] or

[0266] R^{31} and R^{32} together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

[0267] L^{1B} is linear (C_2-C_{10}) -alkanediyl,


[0268] and

[0269] where (C_1-C_{10}) -alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,


[0270] and

[0271] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C_3-C_6) -cycloalkyl ring or a phenyl ring,


[0272] B is a bond or a group of the formula



-continued




- [0273] where
 [0274] * marks the linkage site with L¹,
 [0275] ** marks the linkage site with L²,
 [0276] P is O or NH,
 [0277] L³ is a bond or (C₂-C₄)-alkanediyl,
 [0278] L⁴ is a bond or a group of the formula




- [0279] in which
 [0280] *** marks the linkage site with the carbonyl group,
 [0281] **** marks the linkage site with L²,
 [0282] R²⁵ is hydrogen or methyl,
 [0283] R²⁸ is hydrogen, (C₁-C₄)-alkylcarbonyl, tert-butyloxycarbonyl or benzyloxycarbonyl,
 [0284] Q¹ is a 4- to 7-membered heterocycle,
 [0285] Q² is a 3- to 7-membered carbocycle or a 4- to 7-membered heterocycle,
 [0286] R¹⁴ is hydrogen or (C₁-C₄)-alkyl,
 [0287] R¹⁵ is hydrogen or (C₁-C₄)-alkyl,
 [0288] or
 [0289] R¹⁴ and R¹⁵ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 [0290] R¹⁶ is hydrogen or (C₁-C₄)-alkyl,
 [0291] R¹⁷ is hydrogen or (C₁-C₄)-alkyl,
 [0292] or
 [0293] R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 [0294] R¹⁸ is hydrogen or (C₁-C₄)-alkyl,
 [0295] R¹⁹ is hydrogen or the side group of a natural α-amino acid or of its homologues or isomers,
 [0296] R²⁰ is hydrogen or (C₁-C₄)-alkyl,
 [0297] or
 [0298] R¹⁹ and R²⁰ together with the atoms to which they are bonded form a pyrrolidinyl ring,
 [0299] R²¹ is hydrogen or (C₁-C₄)-alkyl,
 [0300] R²² is hydrogen or (C₁-C₄)-alkyl,
 [0301] or
 [0302] R²¹ and R²² together with the atoms to which they are bonded form a 3- to 7-membered carbocycle,
 [0303] R²³ is (C₁-C₄)-alkyl,
 [0304] R²⁴ is hydrogen or (C₁-C₄)-alkyl,
 [0305] R²⁷ is hydrogen or (C₁-C₄)-alkyl,


- [0306] R³⁶ is hydrogen, (C₁-C₄)-alkylcarbonyl, tert-butyloxycarbonyl or benzyloxycarbonyl,
 [0307] R³⁷ is hydrogen or methyl,
 [0308] or
 [0309] R³⁶ and R³⁷ together with the atoms to which they are bonded form a pyrrolidine ring,
 [0310] L² is linear (C₂-C₁₀)-alkanediyl or is a group of the formula



- [0311] where
 [0312] p is a number from 2 to 6,
 [0313] ##3 marks the linkage site with the group B,
 [0314] ##4 marks the linkage site with the nitrogen atom,
 [0315] where (C₂-C₁₀)-alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,
 [0316] and
 [0317] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,
 [0318] D is a group of the formula



- [0319] in which
 [0320] #3 marks the linkage site with the nitrogen atom,
 [0321] R¹ is hydrogen or methyl,
 [0322] R² is isopropyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,
 [0323] or
 [0324] R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[0325] in which

[0326] #⁴ marks the linkage site with the adjacent nitrogen atom,

[0327] #⁵ marks the linkage site with the carbonyl group,

[0328] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[0329] in which

[0330] #⁶ marks the linkage site with the carbonyl group,


[0331] R⁶ is hydrogen, hydroxy or benzyloxy,

[0332] R³ is hydrogen or methyl,

[0333] R⁴ is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

[0334] or

[0335] R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[0336] in which

[0337] #⁷ marks the linkage site with the adjacent nitrogen atom,

[0338] #⁸ marks the linkage site with the group T¹,

[0339] T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹,

[0340] in which

[0341] R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

[0342] R⁸ is hydrogen or methyl,

[0343] R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,


[0344] or

[0345] R⁸ and R⁹ together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

[0346] R¹⁰ is benzoyl,

[0347] R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

[0348] R⁵ is hydrogen, methyl or a group of the formula



[0349] in which

[0350] #⁹ marks the linkage site with —CHC(R²⁶)-T²,

[0351] R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

[0352] R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,

[0353] R²⁶ is hydrogen or hydroxy,

[0354] T² is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,

[0355] R³⁵ is methyl or hydroxy,

and also their salts, solvates and solvates of the salts.

[0356] Preferred subject of the invention are binder-drug conjugates of the general formula (Ia) in which

[0357] n is a number from 1 to 20,


[0358] AK is AK₁ or AK₂

[0359] where

[0360] AK₁ is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

[0361] AK₂ is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the NH side group of a lysine residue of the binder to the group G,

[0362] G when AK=AK₁, is a group of the formula



[0363] in which


[0364] #¹ marks the linkage site with the cysteine residue of the binder,

[0365] #² marks the linkage site with the group L¹,

[0366] or

[0367] when AK=AK₂, is carbonyl,


[0368] L¹ is a bond, linear (C₂-C₆)-alkanediyl, a group of the formula



[0369] where

[0370] m is a number from 2 to 6,

[0371] ##¹ marks the linkage site with the group G,



-

- Chemical structures (1) through (5) are defined as follows:
- (1)
- (2)
- (3)
- (4)


-continued-

10a, 10b, or 10c


-

- [0398] in which
[0399] *** marks the linkage site with the carbonyl group,
[0400] **** marks the linkage site with L²,
[0401] R²⁵ is hydrogen or methyl,
[0402] R²⁸ is hydrogen, methylcarbonyl or tert-butoxycarbonyl,
[0403] Q¹ is a 4- to 7-membered heterocycle,
[0404] R¹⁴ is hydrogen,
[0405] R¹⁵ is hydrogen,
[0406] R¹⁶ is hydrogen or methyl,
[0407] R¹⁷ is hydrogen or methyl,
[0408] or
[0409] R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a piperazinyl ring,
[0410] R¹⁸ is hydrogen,
[0411] R¹⁹ is hydrogen, methyl, propan-2-yl, 2-methylpropan-1-yl or 1-methylpropan-1-yl,
[0412] R²⁰ is hydrogen or methyl,
[0413] or
[0414] R¹⁹ and R²⁰ together with the atoms to which they are bonded form a pyrrolidinyl ring,
[0415] R²¹ is hydrogen or methyl,
[0416] R²² is hydrogen or methyl,
[0417] or
[0418] R²¹ and R²² together with the atoms to which they are bonded form a cyclopropyl ring.


- [0419] R^{23} is methyl,
 [0420] R^{24} is hydrogen or methyl,
 [0421] R^{27} is hydrogen,
 [0422] R^{36} is hydrogen, methylcarbonyl or tert-butoxycarbonyl,
 [0423] R^{37} is hydrogen or methyl,
 [0424] or
 [0425] R^{36} and R^{37} together with the atoms to which they are bonded form a pyrrolidine ring,
 [0426] L^2 is linear (C_2-C_6) -alkanediyl or is a group of the formula




- [0427] where
 [0428] p is a number from 2 to 6,
 [0429] ##3 marks the linkage site with the group B,
 [0430] ##4 marks the linkage site with the nitrogen atom, where (C_2-C_{10}) -alkanediyl may be substituted by 1 or 2 methyl substituents,
 [0431] D is a group of the formula




- [0432] where
 [0433] #3 marks the linkage site with the nitrogen atom,
 [0434] R^1 is hydrogen,
 [0435] R^2 is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,
 [0436] or
 [0437] R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




- [0438] in which
 [0439] #4 marks the linkage site with the adjacent nitrogen atom,
 [0440] #5 marks the linkage site with the carbonyl group,
 [0441] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula




- [0442] in which
 [0443] #6 marks the linkage site with the carbonyl group,
 [0444] R^6 is hydrogen, hydroxy or benzyloxy,
 [0445] R^3 is hydrogen,
 [0446] R^4 is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,
 [0447] or
 [0448] R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




- [0449] in which
 [0450] #7 marks the linkage site with the adjacent nitrogen atom,
 [0451] #8 marks the linkage site with the group T^1 ,
 [0452] T^1 is a group of the formula $-C(=O)-OR^7$, $-C(=O)-NR^8R^9$, $-C(=O)-NH-NH-R^{10}$ or $-CH_2-O-R^{11}$,
 [0453] in which
 [0454] R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,
 [0455] R^8 is hydrogen or methyl,
 [0456] R^9 is hydrogen, methyl, ethyl, n-propyl or benzyl,
 [0457] or
 [0458] R^8 and R^9 together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,
 [0459] R^{10} is benzoyl,
 [0460] R^{11} is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,
 [0461] R^5 is hydrogen, methyl or a group of the formula





- [0462] in which
 [0463] #⁹ marks the linkage site with —CHC(R²⁶)-T²,
 [0464] R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,
 [0465] R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,
 [0466] R²⁶ is hydrogen or hydroxy,
 [0467] T² is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,
 [0468] R³⁵ is methyl or hydroxy,
 and also their salts, solvates and solvates of the salts.
 [0469] Preferred subject matter of the invention are binder-drug conjugates of the general formula (Ia) in which
 [0470] n is a number from 1 to 20,
 [0471] AK is AK₁ or AK₂
 [0472] where
 [0473] AK₁ is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,
 [0474] AK₂ is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the NH side group of a lysine residue of the binder to the group G,
 [0475] G when AK=AK₁, is a group of the formula



- [0476] in which
 [0477] #¹ marks the linkage site with the cysteine residue of the binder,
 [0478] #² marks the linkage site with the group L¹,
 [0479] or
 [0480] when AK=AK₂, is carbonyl,
 [0481] L¹ is a bond, linear (C₂-C₆)-alkanediyl, a group of the formula




- [0482] where
 [0483] m is a number from 2 to 6,
 [0484] #¹ marks the linkage site with the group G,
 [0485] #² marks the linkage site with the group B,
 [0486] L^{1A} is linear (C₂-C₆)-alkanediyl,
 [0487] B¹ is a group of the formula




-continued

- [0488] in which
 [0489] #⁵ marks the linkage site with the group L^{1A},
 [0490] #⁶ marks the linkage site with the group L^{1B},
 [0491] L⁵ is a bond,
 [0492] L⁶ is a bond or a group of the formula




- [0493] in which
 [0494] #⁷ marks the linkage site with the carbonyl group,
 [0495] #⁸ marks the linkage site with L^{1B},
 [0496] R³³ is hydrogen, methylcarbonyl or tert-butyloxycarbonyl,
 [0497] R³⁴ is hydrogen or methyl,
 [0498] R²⁹ is hydrogen,
 [0499] R³⁰ is hydrogen,
 [0500] R³¹ is hydrogen or methyl,
 [0501] R³² is hydrogen or methyl,
 [0502] L^{1B} is linear (C₂-C₆)-alkanediyl,
 [0503] and
 [0504] where (C₂-C₆)-alkanediyl may be substituted by 1 or 2 methyl substituents,
 [0505] B is a bond or a group of the formula



- [0506] where
 [0507] * marks the linkage site with L¹,
 [0508] ** marks the linkage site with L²,
 [0509] L³ is a bond or ethane-1,2-diyl,

[0510] L^4 is a bond or a group of the formula



[0511] in which

[0512] *** marks the linkage site with the carbonyl group,

[0513] **** marks the linkage site with L^2 ,

[0514] R^{25} is hydrogen or methyl,

[0515] R^{28} is hydrogen, methylcarbonyl or tert-butyloxycarbonyl,

[0516] Q^1 is a 4- to 7-membered heterocycle,

[0517] R^{14} is hydrogen,

[0518] R^{15} is hydrogen,

[0519] R^{16} is hydrogen or methyl,

[0520] R^{17} is hydrogen or methyl,

[0521] or

[0522] R^{16} and R^{17} together with the atoms to which they are bonded form a piperazinyl ring,

[0523] R^{18} is hydrogen,

[0524] R^{19} is hydrogen, methyl, propan-2-yl, 2-methylpropan-1-yl or 1-methylpropan-1-yl,

[0525] R^{20} is hydrogen or methyl,

[0526] or

[0527] R^{19} and R^{20} together with the atoms to which they are bonded form a pyrrolidinyl ring,

[0528] R^{21} is hydrogen or methyl,

[0529] R^{22} is hydrogen or methyl,

[0530] or

[0531] R^{21} and R^{22} together with the atoms to which they are bonded form a cyclopropyl ring,

[0532] R^{23} is methyl,

[0533] R^{24} is hydrogen or methyl,

[0534] R^{27} is hydrogen,


[0535] R^{36} is hydrogen, methylcarbonyl or tert-butyloxycarbonyl,

[0536] R^{37} is hydrogen or methyl,

[0537] or

[0538] R^{36} and R^{37} together with the atoms to which they are bonded form a pyrrolidine ring,

[0539] L^2 is linear (C_2 - C_6)-alkanediyl or is a group of the formula



[0540] where


[0541] p is a number from 2 to 6,

[0542] ##3 marks the linkage site with the group B,

[0543] ##4 marks the linkage site with the nitrogen atom,

[0544] where (C_2 - C_{10})-alkanediyl may be substituted by 1 or 2 methyl substituents,

[0545] D is a group of the formula



[0546] where


[0547] #3 marks the linkage site with the nitrogen atom,

[0548] R^1 is hydrogen,

[0549] R^2 is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-yl-methyl,

[0550] or

[0551] R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[0552] in which

[0553] #4 marks the linkage site with the adjacent nitrogen atom,

[0554] #5 marks the linkage site with the carbonyl group,

[0555] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[0556] in which

[0557] #6 marks the linkage site with the carbonyl group,


[0558] R^6 is hydrogen, hydroxy or benzyloxy,

[0559] R^3 is hydrogen,

[0560] R^4 is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-yl-methyl,

[0561] or

[0562] R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[0563] in which

[0564] #7 marks the linkage site with the adjacent nitrogen atom,

[0565] #8 marks the linkage site with the group T^1 ,

[0566] T^1 is a group of the formula $-C(=O)-OR^7$, $-C(=O)-NR^8R^9$, $-C(=O)-NH-NH-R^{10}$ or $-CH_2-O-R^{11}$,

[0567] in which

[0568] R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

[0569] R^8 is hydrogen or methyl,

[0570] R^9 is hydrogen, methyl, ethyl, n-propyl or benzyl,


[0571] or

[0572] R^8 and R^9 together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

[0573] R^{10} is benzoyl,

[0574] R^{11} is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

[0575] R^5 is hydrogen, methyl or a group of the formula



[0576] in which

[0577] #9 marks the linkage site with $-CHC(R^{26})-T^2$,

[0578] R^{12} is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula $-S(O)_2OH$,

[0579] R^{13} is phenyl which may be substituted by methoxycarbonyl or carboxyl,

[0580] R^{26} is hydrogen or hydroxy,

[0581] T^2 is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,

[0582] R^{35} is methyl or hydroxy,

and also their salts, solvates and solvates of the salts.

[0583] Preferred subject matter of the invention are binder-drug conjugates of the general formula (Ia), in which

[0584] n is a number from 1 to 10,


[0585] AK is AK_1 or AK_2

[0586] where

[0587] AK_1 is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

[0588] AK_2 is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the NH side group of a lysine residue of the binder to the group G,

[0589] G when $AK=AK_1$, is a group of the formula



[0590] in which


[0591] #1 marks the linkage site with the cysteine residue of the binder,

[0592] #2 marks the linkage site with the group L^1 ,

[0593] or

[0594] when $AK=AK_2$, is carbonyl,

[0595] L^1 is a bond, linear (C_2-C_6) -alkanediyl, a group of the formula



[0596] where


[0597] m is a number 2 or 3,

[0598] ##1 marks the linkage site with the group G,

[0599] ##2 marks the linkage site with the group B,

[0600] where (C_2-C_6) -alkanediyl may be substituted by 1 or 2 methyl substituents,

[0601] B is a bond or a group of the formula




[0602] where

[0603] * marks the linkage site with L^1 ,

[0604] ** marks the linkage site with L^2 ,

[0605] L^3 is a bond or ethane-1,2-diyl,

[0606] L^4 is a bond or a group of the formula



[0607] in which

[0608] *** marks the linkage site with the carbonyl group,

[0609] *** marks the linkage site with L^2 ,

[0610] R^{25} is methyl,

[0611] R^{28} is hydrogen, methylcarbonyl or tert-butyloxycarbonyl,

[0612] Q^1 is piperidine-1,4-diyl,

[0613] R^{16} is hydrogen or methyl,

[0614] R^{17} is hydrogen or methyl,

[0615] or

[0616] R^{16} and R^{17} together with the atoms to which they are bonded form a piperazinyl ring,

[0617] R^{21} is hydrogen or methyl,

[0618] R^{22} is hydrogen or methyl,

[0619] or

[0620] R^{21} and R^{22} together with the atoms to which they are bonded form a cyclopropyl ring,


[0621] R^{23} is methyl,

[0622] R^{24} is hydrogen,

[0623] R^{36} is hydrogen, methylcarbonyl or tert-butyloxycarbonyl,

[0624] R^{37} is hydrogen or methyl,

[0625] L^2 is linear (C_2 - C_6)-alkanediyl or is a group of the formula




[0626] where

[0627] p is a number from 2 to 6,


[0628] ##3 marks the linkage site with the group B,

[0629] ##4 marks the linkage site with the nitrogen atom,

[0630] D is a group of the formula



-continued



[0631] where


[0632] #3 marks the linkage site with the nitrogen atom,

[0633] R^1 is hydrogen,

[0634] R^2 is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-yl-methyl,

[0635] or

[0636] R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[0637] in which

[0638] #4 marks the linkage site with the adjacent nitrogen atom,

[0639] #5 marks the linkage site with the carbonyl group,

[0640] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[0641] in which

[0642] #6 marks the linkage site with the carbonyl group,


[0643] R^6 is hydrogen, hydroxy or benzyloxy,

[0644] R^3 is hydrogen,

[0645] R^4 is benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

[0646] or

[0647] R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[0648] in which

[0649] #7 marks the linkage site with the adjacent nitrogen atom,

[0650] #8 marks the linkage site with the group T^1 ,

[0651] T^1 is a group of the formula $-C(=O)-OR^7$, $-C(=O)-NR^8R^9$ or $-CH_2-O-R^{11}$, in which


[0652] R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

[0653] R^8 is hydrogen or methyl,

[0654] R^9 is hydrogen, methyl, ethyl, n-propyl or benzyl,

[0655] R^{11} is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

[0656] R^5 is hydrogen, methyl or a group of the formula



[0657] in which

[0658] #9 marks the linkage site with $-CHCH_2$ phenyl,

[0659] R^{12} is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula $-S(O)_2OH$,

[0660] R^{13} is phenyl which may be substituted by methoxycarbonyl or carboxyl,

[0661] R^{35} is methyl or hydroxy, and also their salts, solvates and solvates of the salts.

[0662] Preferred subject matter of the invention are binder-drug conjugates of the general formula (Ia) in which

[0663] n is a number from 1 to 10,


[0664] AK is AK_1 or AK_2

[0665] where

[0666] AK_1 is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

[0667] AK_2 is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the NH side group of a lysine residue of the binder to the group G,

[0668] G when $AK=AK_1$, is a group of the formula



[0669] in which


[0670] #1 marks the linkage site with the cysteine residue of the binder,

[0671] #2 marks the linkage site with the group L^1 ,

[0672] or

[0673] when $AK=AK_2$, is carbonyl,

[0674] L^1 is a bond, linear (C_2-C_6) -alkanediyl, a group of the formula



[0675] where


[0676] m is a number 2 or 3,

[0677] ##1 marks the linkage site with the group G,

[0678] ##2 marks the linkage site with the group B,

[0679] where (C_2-C_6) -alkanediyl may be substituted by 1 or 2 methyl substituents,

[0680] B is a bond or a group of the formula




[0681] where

[0682] * marks the linkage site with L^1 ,


[0683] ** marks the linkage site with L^2 ,

[0684] L^3 is a bond or ethane-1,2-diyl,


[0685] L^4 is a bond or a group of the formula




- [0686] in which
 [0687] *** marks the linkage site with the carbonyl group,
 [0688] **** marks the linkage site with L^2 ,
 [0689] R^{25} is methyl,
 [0690] R^{28} is hydrogen, methylcarbonyl or tert-butoxycarbonyl,
 [0691] Q^1 is piperidine-1,4-diyl,
 [0692] R^{16} is hydrogen or methyl,
 [0693] R^{17} is hydrogen or methyl,
 [0694] or
 [0695] R^{16} and R^{17} together with the atoms to which they are bonded form a piperazinyl ring,
 [0696] R^{21} is hydrogen or methyl,
 [0697] R^{22} is hydrogen or methyl,
 [0698] or
 [0699] R^{21} and R^{22} together with the atoms to which they are bonded form a cyclopropyl ring,
 [0700] R^{23} is methyl,
 [0701] R^{24} is hydrogen,
 [0702] L^2 is linear (C_2-C_6)-alkanediyl or is a group of the formula




- [0703] where
 [0704] p is a number from 2 to 6,
 [0705] ##3 marks the linkage site with the group B,
 [0706] ##4 marks the linkage site with the nitrogen atom,
 [0707] D is a group of the formula




- [0708] where
 [0709] #3 marks the linkage site with the nitrogen atom,
 [0710] R^1 is hydrogen,
 [0711] R^2 is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-yl-methyl,
 [0712] or
 [0713] R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



- [0714] in which
 [0715] #4 marks the linkage site with the adjacent nitrogen atom,
 [0716] #5 marks the linkage site with the carbonyl group,
 [0717] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula




- [0718] in which
 [0719] #6 marks the linkage site with the carbonyl group,
 [0720] R^6 is hydrogen, hydroxy or benzyloxy,
 [0721] R^3 is hydrogen,
 [0722] R^4 is benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,
 [0723] or
 [0724] R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



- [0725] in which
 [0726] #7 marks the linkage site with the adjacent nitrogen atom,
 [0727] #8 marks the linkage site with the group T^1 ,
 [0728] T^1 is a group of the formula $-C(=O)-OR^7$, $-C(=O)-NR^8R^9$ or $-CH_2-O-R^{11}$, in which
 [0729] R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,
 [0730] R^8 is hydrogen or methyl,
 [0731] R^9 is hydrogen, methyl, ethyl, n-propyl or benzyl,
 [0732] R^{11} is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

[0733] R^5 is hydrogen, methyl or a group of the formula



[0734] in which

[0735] $\#^9$ marks the linkage site with $-\text{CHCH}_2\text{phenyl}$,

[0736] R^{12} is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula $-\text{S}(\text{O})_2\text{OH}$,

[0737] R^{13} is phenyl which may be substituted by methoxycarbonyl or carboxyl,

[0738] R^{35} is methyl or hydroxy,

[0739] and also their salts, solvates and solvates of the salts.

[0740] Preferred subject matter of the present invention are binder-drug conjugates of the general formula (Ia) as indicated above, in which

[0741] n is a number from 1 to 10,

[0742] AK is AK_2 ,

[0743] where


[0744] AK_2 is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the NH side group of a lysine residue of the binder to the group G,

[0745] G is carbonyl,

[0746] L^1 is a bond,

[0747] B is a bond,

[0748] L^2 is linear $(\text{C}_3\text{--}\text{C}_6)$ -alkanediyl or is a group of the formula




[0749] where

[0750] p is a number 2 or 3,

[0751] $\# \# 3$ marks the linkage site with the group B,

[0752] $\# \# 4$ marks the linkage site with the nitrogen atom,

[0753] D is a group of the formula



[0754] where


[0755] $\#^3$ marks the linkage site with the nitrogen atom,

[0756] R^1 is hydrogen,

[0757] R^2 is benzyl, 4-hydroxybenzyl or 1H-indol-3-yl-methyl,

[0758] or

[0759] R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[0760] in which

[0761] $\#^4$ marks the linkage site with the adjacent nitrogen atom,

[0762] $\#^5$ marks the linkage site with the carbonyl group,

[0763] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[0764] in which


[0765] $\#^6$ marks the linkage site with the carbonyl group,

[0766] R^3 is hydrogen,

[0767] R^4 is benzyl, 4-hydroxybenzyl or 1H-indol-3-yl-methyl,

[0768] or

[0769] R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[0770] in which

[0771] $\#^7$ marks the linkage site with the adjacent nitrogen atom,

[0772] $\#^8$ marks the linkage site with the group T^1 ,


[0773] T^1 is a group of the formula $-\text{C}(=\text{O})-\text{OR}^7$ or $-\text{C}(=\text{O})-\text{NR}^8\text{R}^9$

[0774] in which


[0775] R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

[0776] R^8 is hydrogen,


- [0777] R^9 is hydrogen or benzyl,
 [0778] R^{35} is methyl,
 and also their salts, solvates and solvates of the salts.
 [0779] Preferred subject matter of the invention are binder-drug conjugates of the general formula (Ia), in which
 [0780] n is a number from 1 to 10,
 [0781] AK is AK_2 ,
 [0782] where
 [0783] AK_2 is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the NH side group of a lysine residue of the binder to the group G,
 [0784] G is carbonyl,
 [0785] L^1 is a bond,
 [0786] B is a bond,
 [0787] L^2 is linear (C_3 - C_6)-alkanediyl or is a group of the formula



- [0788] where
 [0789] p is a number 2 or 3,
 [0790] ##3 marks the linkage site with the group B,
 [0791] ##4 marks the linkage site with the nitrogen atom,
 [0792] D is a group of the formula




- [0793] where
 [0794] #3 marks the linkage site with the nitrogen atom,
 [0795] R^1 is hydrogen,
 [0796] R^2 is benzyl or 1H-indol-3-ylmethyl,
 [0797] or
 [0798] R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




- [0799] in which
 [0800] #4 marks the linkage site with the adjacent nitrogen atom,
 [0801] #5 marks the linkage site with the carbonyl group,

- [0802] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



- [0803] in which
 [0804] #6 marks the linkage site with the carbonyl group,
 [0805] R^3 is hydrogen,
 [0806] R^4 is benzyl or 1H-indol-3-ylmethyl,
 [0807] or
 [0808] R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



- [0809] in which
 [0810] #7 marks the linkage site with the adjacent nitrogen atom,
 [0811] #8 marks the linkage site with the group T^1 ,
 [0812] T^1 is a group of the formula $-C(=O)-OR^7$ or $-C(=O)-NR^8R^9$
 [0813] in which
 [0814] R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,
 [0815] R^8 is hydrogen,
 [0816] R^9 is hydrogen or benzyl,
 [0817] R^{35} is methyl,


and also their salts, solvates and solvates of the salts.

- [0818] Preferred subject matter of the present invention are binder-drug conjugates of the general formula (Ia) as indicated above, in which

- [0819] n is a number from 1 to 10,
 [0820] AK is AK_1 ,

- [0821] where
 [0822] AK_1 is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

[0823] G is a group of the formula




[0824] where

[0825] #¹ marks the linkage site with the cysteine residue of the binder,

[0826] #² marks the linkage site with the group L¹,

[0827] L¹ is a bond, linear (C₃-C₅)-alkanediyl or a group of the formula



[0828] where


[0829] m is a number 2 or 3,

[0830] ##¹ marks the linkage site with the group G,

[0831] ##² marks the linkage site with the group B,

[0832] where (C₃-C₅)-alkanediyl may be substituted by 1 or 2 methyl substituents,

[0833] B is a bond or a group of the formula




[0834] where

[0835] * marks the linkage site with L¹,

[0836] ** marks the linkage site with L²,

[0837] L³ is a bond or ethane-1,2-diyl,

[0838] L⁴ is a bond or a group of the formula



[0839] in which

[0840] **** marks the linkage site with the carbonyl group,

[0841] **** marks the linkage site with L²,

[0842] R²⁵ is methyl,

[0843] R²⁸ is hydrogen, methylcarbonyl or tert-butyloxycarbonyl,


[0844] R¹⁶ is hydrogen or methyl,

[0845] R¹⁷ is hydrogen or methyl,

[0846] or

[0847] R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a piperazinyl ring,

[0848] L² is linear (C₃-C₅)-alkanediyl or is a group of the formula




[0849] where

[0850] p is a number 2 or 3,

[0851] ##³ marks the linkage site with the group B,

[0852] ##⁴ marks the linkage site with the nitrogen atom,

[0853] D is a group of the formula



[0854] where


[0855] #³ marks the linkage site with the nitrogen atom,

[0856] R¹ is hydrogen,

[0857] R² is benzyl, 4-hydroxybenzyl or 1H-indol-3-yl-methyl,

[0858] or

[0859] R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[0860] in which

[0861] #⁴ marks the linkage site with the adjacent nitrogen atom,

[0862] #⁵ marks the linkage site with the carbonyl group,

[0863] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula




[0864] in which

[0865] #⁶ marks the linkage site with the carbonyl group,


[0866] R³ is hydrogen,

[0867] R⁴ is benzyl, 4-hydroxybenzyl or 1H-indol-3-yl-methyl,


- [0868] or
 [0869] R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




- [0870] in which
 [0871] #7 marks the linkage site with the adjacent nitrogen atom,
 [0872] #8 marks the linkage site with the group T^1 ,
 [0873] T^1 is a group of the formula $-C(=O)-OR^7$ or $-C(=O)-NR^8R^9$, in which
 [0874] R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,
 [0875] R^8 is hydrogen,
 [0876] R^9 is hydrogen or benzyl,
 [0877] R^{35} is methyl,
 and also their salts, solvates and solvates of the salts.
 [0878] Preferred subject matter of the invention are binder-drug conjugates of the general formula (Ia), in which
 [0879] n is a number from 1 to 10,
 [0880] AK is AK_1 ,
 [0881] where
 [0882] AK_1 is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,
 [0883] G is a group of the formula




- [0884] where
 [0885] #1 marks the linkage site with the cysteine residue of the binder,
 [0886] #2 marks the linkage site with the group L^1 ,
 [0887] L^1 is a bond, linear (C_3-C_5) -alkanediyl or a group of the formula




- [0888] where
 [0889] m is a number 2 or 3,
 [0890] ##1 marks the linkage site with the group G,
 [0891] ##2 marks the linkage site with the group B,
 [0892] where (C_3-C_5) -alkanediyl may be substituted by 1 or 2 methyl substituents,
 [0893] B is a bond or a group of the formula




- [0894] where
 [0895] * marks the linkage site with L^1 ,
 [0896] ** marks the linkage site with L^2 ,
 [0897] L^3 is a bond or ethane-1,2-diyl,
 [0898] L^4 is a bond or a group of the formula



- [0899] in which
 [0900] *** marks the linkage site with the carbonyl group,
 [0901] **** marks the linkage site with L^2 ,
 [0902] R^{25} is methyl,
 [0903] R^{28} is hydrogen, methylcarbonyl or tert-butoxycarbonyl,
 [0904] R^{16} is hydrogen or methyl,
 [0905] R^{17} is hydrogen or methyl,
 [0906] or
 [0907] R^{16} and R^{17} together with the atoms to which they are bonded form a piperazinyl ring,
 [0908] L^2 is linear (C_3-C_5) -alkanediyl or is a group of the formula



- [0909] where
 [0910] p is a number 2 or 3,
 [0911] ##3 marks the linkage site with the group B,
 [0912] ##4 marks the linkage site with the nitrogen atom,
 [0913] D is a group of the formula



[0914] where


[0915] #³ marks the linkage site with the nitrogen atom,

[0916] R¹ is hydrogen,

[0917] R² is benzyl or 1H-indol-3-ylmethyl,


[0918] or

[0919] R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[0920] in which

[0921] #⁴ marks the linkage site with the adjacent nitrogen atom,



[0930] in which

[0931] #⁷ marks the linkage site with the adjacent nitrogen atom,

[0932] #⁸ marks the linkage site with the group T¹,

[0933] T¹ is a group of the formula —C(=O)—OR⁷ or —C(=O)—NR⁸R⁹,

[0934] in which

[0935] R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,


[0936] R⁸ is hydrogen,

[0937] R⁹ is hydrogen or benzyl,

[0938] R³⁵ is methyl,


and also their salts, solvates and solvates of the salts.

[0939] Additionally provided by the present invention are compounds of the formula (XXXa)



[0922] #⁵ marks the linkage site with the carbonyl group,

[0923] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[0924] in which

[0925] #⁶ marks the linkage site with the carbonyl group,

[0926] R³ is hydrogen,

[0927] R⁴ is benzyl or 1H-indol-3-ylmethyl,


[0928] or

[0929] R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula

in which

[0940] Cys is a cysteine residue which is bonded via the sulphur atom of the side chain to a carbon atom of the succinimide,

[0941] L¹ is a bond, linear (C₁-C₁₀)-alkanediyl, a group of the formula



[0942] where


[0943] m is a number from 2 to 6,

[0944] ##¹ marks the linkage site with the group G,


[0945] ##² marks the linkage site with the group B,

[0946] L^{1A} is linear (C₂-C₁₀)-alkanediyl,

[0947] B¹ is a group of the formula




- [0948] in which
 [0949] ##⁵ marks the linkage site with the group L^{1A},
 [0950] ##⁶ marks the linkage site with the group L^{1B},
 [0951] L⁵ is a bond or (C₂-C₄)-alkanediyl,
 [0952] L⁶ is a bond,
 [0953] R²⁹ is hydrogen or (C₁-C₄)-alkyl,
 [0954] R³⁰ is hydrogen or (C₁-C₄)-alkyl,
 [0955] or
 [0956] R²⁹ and R³⁰ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 [0957] R³¹ is hydrogen or (C₁-C₄)-alkyl,
 [0958] R³² is hydrogen or (C₁-C₄)-alkyl,
 [0959] or
 [0960] R³¹ and R³² together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 [0961] L^{1B} is linear (C₂-C₁₀)-alkanediyl,
 [0962] and
 [0963] where (C₁-C₁₀)-alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,
 [0964] and
 [0965] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,
 [0966] B is a bond or a group of the formula




- [0967] where
 [0968] * marks the linkage site with L¹,
 [0969] ** marks the linkage site with L²,
 [0970] P is O or NH,
 [0971] L³ is a bond or (C₂-C₄)-alkanediyl,
 [0972] L⁴ is a bond,
 [0973] Q¹ is a 4- to 7-membered heterocycle,
 [0974] Q² is a 3- to 7-membered carbocycle or a 4- to 7-membered heterocycle,
 [0975] R¹⁴ is hydrogen or (C₁-C₄)-alkyl,
 [0976] R¹⁵ is hydrogen or (C₁-C₄)-alkyl,
 [0977] or
 [0978] R¹⁴ and R¹⁵ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 [0979] R¹⁶ is hydrogen or (C₁-C₄)-alkyl,
 [0980] R¹⁷ is hydrogen or (C₁-C₄)-alkyl,
 [0981] or
 [0982] R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 [0983] R¹⁸ is hydrogen or (C₁-C₄)-alkyl,


- [0984] R¹⁹ is hydrogen or the side group of a natural α -amino acid or of its homologues or isomers,
 [0985] R²⁰ is hydrogen or (C₁-C₄)-alkyl,
 [0986] or
 [0987] R¹⁹ and R²⁰ together with the atoms to which they are bonded form a pyrrolidinyl ring,
 [0988] R²¹ is hydrogen or (C₁-C₄)-alkyl,
 [0989] R²² is hydrogen or (C₁-C₄)-alkyl,
 [0990] or
 [0991] R²¹ and R²² together with the atoms to which they are bonded form a 3- to 7-membered carbocycle,
 [0992] R²³ is (C₁-C₄)-alkyl,
 [0993] R²⁴ is hydrogen or (C₁-C₄)-alkyl,
 [0994] R²⁷ is hydrogen or (C₁-C₄)-alkyl,
 [0995] L² is linear (C₂-C₁₀)-alkanediyl or is a group of the formula




- [0996] where
 [0997] p is a number from 2 to 6,
 [0998] ##³ marks the linkage site with the group B,
 [0999] ##⁴ marks the linkage site with the nitrogen atom,
 [1000] where (C₂-C₁₀)-alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,
 [1001] and
 [1002] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,
 [1003] D is a group of the formula




- [1004] where
 [1005] #³ marks the linkage site with the nitrogen atom,
 [1006] R¹ is hydrogen or methyl,
 [1007] R² is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,
 [1008] or
 [1009] R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



- [1010] in which
 [1011] #⁴ marks the linkage site with the adjacent
 nitrogen atom,
 [1012] #⁵ marks the linkage site with the carbonyl
 group,
 [1013] the ring A with the N—O moiety present therein
 is a mono- or bicyclic, optionally substituted heterocycle
 of the formula




- [1014]** in which
[1015] #⁶ marks the linkage site with the carbonyl group,
[1016] R⁶ is hydrogen, hydroxy or benzyloxy,
[1017] R³ is hydrogen or methyl,
[1018] R⁴ is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,
[1019] or
[1020] R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




- [1021] in which
 [1022] #⁷ marks the linkage site with the adjacent nitrogen atom,
 [1023] #⁸ marks the linkage site with the group T¹,
 [1024] T¹ is a group of the formula —C(=O)—OR⁷,
 —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or
 —CH₂—O—R¹¹,
 [1025] in which
 [1026] R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,


- [1027] R⁸ is hydrogen or methyl,
[1028] R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,
[1029] or
[1030] R⁸ and R⁹ together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,
[1031] R¹⁰ is benzoyl,
[1032] R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,
[1033] R⁵ is hydrogen, methyl or a group of the formula



- [1034] in which
 [1035] #⁹ marks the linkage site with —CHC(R²⁶)-T²,
 [1036] R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,
 [1037] R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,
 [1038] R²⁶ is hydrogen or hydroxy,
 [1039] T² is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,
 [1040] R³⁵ is methyl or hydroxy,
 and also their salts, solvates and solvates of the salts.
 [1041] Preferred in the context of the present invention are additionally also compounds of the formula (XXXa) in which
 [1042] Cys is a cysteine residue which is bonded via the sulphur atom of the side chain via a carbon atom of the succinimide,
 [1043] L¹ is a bond, linear (C₂-C₆)-alkanediyl, a group of the formula




- [1044] where
[1045] m is a number 2 or 3,
[1046] ##1 marks the linkage site with the group G,
[1047] ##2 marks the linkage site with the group B,
[1048] L^{1,4} is linear (C₂-C₆)-alkanediyl,
[1049] B¹ is a group of the formula




- [1050] in which
 [1051] $\#^{\#}$ marks the linkage site with the group L^{1A} ,
 [1052] $\#^{\#}$ marks the linkage site with the group L^{1B} ,
 [1053] L^5 is a bond,
 [1054] L^6 is a bond.


- [1055] R^{29} is hydrogen,
 [1056] R^{30} is hydrogen,
 [1057] R^{31} is hydrogen or methyl,
 [1058] R^{32} is hydrogen or methyl,
 [1059] L^{1B} is linear (C_2-C_6) -alkanediyl,
 [1060] and
 [1061] where (C_2-C_6) -alkanediyl may be substituted by
 1 or 2 methyl substituents,
 [1062] B is a bond or a group of the formula



- [1063] where
 [1064] * marks the linkage site with L^1 ,
 [1065] ** marks the linkage site with L^2 ,
 [1066] L^3 is a bond or ethane-1,2-diyl,
 [1067] L^4 is a bond,
 [1068] R^{14} is hydrogen,
 [1069] R^{15} is hydrogen,
 [1070] R^{16} is hydrogen or methyl,
 [1071] R^{17} is hydrogen or methyl,
 [1072] or
 [1073] R^{16} and R^{17} together with the atoms to which they
 are bonded form piperazinyl ring,
 [1074] R^{23} is methyl,
 [1075] R^{24} is hydrogen or methyl,
 [1076] L^2 is linear (C_2-C_6) -alkanediyl or is a group of the
 formula




- [1077] where
 [1078] p is a number 2 or 3,
 [1079] ##3 marks the linkage site with the group B,
 [1080] ##4 marks the linkage site with the nitrogen atom,
 [1081] D is a group of the formula




- [1082] where
 [1083] #3 marks the linkage site with the nitrogen atom,
 [1084] R^1 is hydrogen,
 [1085] R^2 is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl,
 1-phenylethyl or 1H-indol-3-yl-methyl,


- [1086] or
 [1087] R^1 and R^2 together with the carbon atom to which
 they are bonded form a (1S,2R)-2-phenylcyclopropane-
 1,1-diyl group of the formula



- [1088] in which
 [1089] #4 marks the linkage site with the adjacent
 nitrogen atom,
 [1090] #5 marks the linkage site with the carbonyl
 group,
 [1091] the ring A with the N—O moiety present therein
 is a mono- or bicyclic, optionally substituted heterocycle
 of the formula



- [1092] in which
 [1093] #6 marks the linkage site with the carbonyl
 group,
 [1094] R^6 is hydrogen, hydroxy or benzyloxy,
 [1095] R^3 is hydrogen,
 [1096] R^4 is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl,
 1-phenylethyl or 1H-indol-3-yl-methyl,
 [1097] or
 [1098] R^3 and R^4 together with the carbon atom to which
 they are bonded form a (1S,2R)-2-phenylcyclopropane-
 1,1-diyl group of the formula



- [1099] in which
 [1100] #7 marks the linkage site with the adjacent
 nitrogen atom,
 [1101] #8 marks the linkage site with the group T^1 ,

[1102] T^1 is a group of the formula $-C(=O)-OR^7$, $-C(=O)-NR^8R^9$, $-C(=O)-NH-NH-R^{10}$ or $-CH_2-O-R^{11}$,

[1103] in which

[1104] R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

[1105] R^8 is hydrogen or methyl,

[1106] R^9 is hydrogen, methyl, ethyl, n-propyl or benzyl,


[1107] or

[1108] R^8 and R^9 together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

[1109] R^{10} is benzoyl,

[1110] R^{11} is benzyl which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

[1111] R^5 is hydrogen, methyl or a group of the formula



[1112] in which

[1113] $\#^9$ marks the linkage site with $-CHCH_2$ phenyl,

[1114] R^{12} is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula $-S(O)_2OH$,

[1115] R^{13} is phenyl which may be substituted by methoxycarbonyl or carboxyl,

[1116] R^{35} is methyl or hydroxy,


and also their salts, solvates and solvates of the salts.

[1117] Particularly preferred in the context of the present invention are additionally also compounds of the formula (XXXa) in which

[1118] Cys is a cysteine residue which is bonded via the sulphur atom of the side chain via a carbon atom of the succinimide,

[1119] L^1 is a bond or linear (C_2-C_6) -alkanediyl,

[1120] B is a bond or a group of the formula



[1121] where

[1122] * marks the linkage site with L^1 ,

[1123] ** marks the linkage site with L^2 ,


[1124] L^3 is a bond,

[1125] L^4 is a bond,

[1126] R^{16} is hydrogen or methyl,

[1127] R^{17} is hydrogen or methyl,

[1128] L^2 is linear (C_2-C_6) -alkanediyl or is a group of the formula




[1129] where

[1130] p is a number 2 or 3,

[1131] $\#^3$ marks the linkage site with the group B,

[1132] $\#^4$ marks the linkage site with the nitrogen atom,

[1133] D is a group of the formula



[1134] where


[1135] $\#^3$ marks the linkage site with the nitrogen atom,

[1136] R^1 is hydrogen,

[1137] R^2 is benzyl or 1H-indol-3-ylmethyl,

[1138] or

[1139] R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[1140] in which

[1141] $\#^4$ marks the linkage site with the adjacent nitrogen atom,

[1142] $\#^5$ marks the linkage site with the carbonyl group,

[1143] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[1144] in which


[1145] $\#^6$ marks the linkage site with the carbonyl group,

[1146] R^3 is hydrogen,

[1147] R^4 is benzyl or 1H-indol-3-ylmethyl,

[1148] or

[1149] R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[1150] in which

[1151] #7 marks the linkage site with the adjacent nitrogen atom,

[1152] #8 marks the linkage site with the group T^1 ,

[1153] T^1 is a group of the formula $-C(=O)-OR^7$ or $-C(=O)-NR^8R^9$,

[1154] in which

[1155] R^7 is hydrogen,

[1156] R^8 is hydrogen,

[1157] R^9 is hydrogen,

[1158] R^{35} is methyl,


and also their salts, solvates and solvates of the salts.

[1159] Preferred subject matter of the present invention are compounds of the formula (XXXa) as indicated above, in which

[1160] Cys is a cysteine residue which is bonded via the sulphur atom of the side chain via a carbon atom of the succinimide,

[1161] L^1 is a bond or linear (C_2-C_6) -alkanediyl,

[1162] B is a bond or a group of the formula



[1163] where

[1164] * marks the linkage site with L^1 ,

[1165] ** marks the linkage site with L^2 ,


[1166] L^3 is a bond,

[1167] L^4 is a bond,

[1168] R^{16} is hydrogen or methyl,

[1169] R^{17} is hydrogen or methyl,

[1170] L^2 is linear (C_2-C_6) -alkanediyl or is a group of the formula




[1171] where

[1172] p is a number 2 or 3,

[1173] ##3 marks the linkage site with the group B,

[1174] ##4 marks the linkage site with the nitrogen atom,

[1175] D is a group of the formula




[1176] where #3 marks the linkage site with the nitrogen atom, R^1 is hydrogen,

[1177] R^2 is benzyl, 4-hydroxybenzyl or 1H-indol-3-yl-methyl,

[1178] or

[1179] R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[1180] in which

[1181] #4 marks the linkage site with the adjacent nitrogen atom,

[1182] #5 marks the linkage site with the carbonyl group,

[1183] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[1184] in which


[1185] #6 marks the linkage site with the carbonyl group,

[1186] R^3 is hydrogen,


[1187] R^4 is benzyl, 4-hydroxybenzyl or 1H-indol-3-yl-methyl,

[1188] or


[1189] R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



- [1190] in which
 [1191] #⁷ marks the linkage site with the adjacent nitrogen atom,
 [1192] #⁸ marks the linkage site with the group T¹,
 [1193] T¹ is a group of the formula —C(=O)—OR⁷ or —C(=O)—NR⁸R⁹,
 [1194] in which
 [1195] R⁷ is hydrogen,
 [1196] R⁸ is hydrogen,
 [1197] R⁹ is hydrogen,
 [1198] R³⁵ is methyl,
 and also their salts, solvates and solvates of the salts.
 [1199] The present invention additionally provides compounds of the formula (XXXI)




- [1207] in which
 [1208] ##^5 marks the linkage site with the group L^{1A} ,
 [1209] ##^6 marks the linkage site with the group L^{1B} ,
 [1210] L^5 is a bond or $(\text{C}_2\text{-C}_4)$ -alkanediyl,
 [1211] L^6 is a bond,
 [1212] R^{29} is hydrogen or $(\text{C}_1\text{-C}_4)$ -alkyl,
 [1213] R^{30} is hydrogen or $(\text{C}_1\text{-C}_4)$ -alkyl,
 [1214] or
 [1215] R^{29} and R^{30} together with the atoms to which they
 are bonded form a 5- or 6-membered heterocycle,
 [1216] R^{31} is hydrogen or $(\text{C}_1\text{-C}_4)$ -alkyl,
 [1217] R^{32} is hydrogen or $(\text{C}_1\text{-C}_4)$ -alkyl,
 [1218] or
 [1219] R^{31} and R^{32} together with the atoms to which they
 are bonded form a 5- or 6-membered heterocycle,
 [1220] L^{1B} is linear $(\text{C}_2\text{-C}_{10})$ -alkanediyl,
 [1221] and




(XXXI)

- in which
- [1200]** L¹ is a bond, linear (C₁-C₁₀)-alkanediyl, a group of the formula




- [1201] where
 [1202] m is a number from 2 to 6,
 [1203] ##¹ marks the linkage site with the group G,
 [1204] ##² marks the linkage site with the group B,
 [1205] L^{1,4} is linear (C₂-C₁₀)-alkanediyl,
 [1206] B¹ is a group of the formula




- [1222] where (C₁-C₁₀)-alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,
- [1223] and
- [1224] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,
- [1225] B is a bond or a group of the formula


- [1225] B is a bond or a group of the formula




-continued



- [1226] where
 [1227] * marks the linkage site with L¹,
 [1228] ** marks the linkage site with L²,
 [1229] P is O or NH,
 [1230] Q¹ is a 4- to 7-membered heterocycle,
 [1231] Q² is a 3- to 7-membered carbocycle or a 4- to 7-membered heterocycle,
 [1232] R¹⁸ is hydrogen or (C₁-C₄)-alkyl,
 [1233] R¹⁹ is hydrogen or the side group of a natural α -amino acid or of its homologues or isomers,
 [1234] R²⁰ is hydrogen or (C₁-C₄)-alkyl,
 [1235] or
 [1236] R¹⁹ and R²⁰ together with the atoms to which they are bonded form a pyrrolidinyl ring,
 [1237] R²¹ is hydrogen or (C₁-C₄)-alkyl,
 [1238] R²² is hydrogen or (C₁-C₄)-alkyl,
 [1239] or
 [1240] R²¹ and R²² together with the atoms to which they are bonded form a 3- to 7-membered carbocycle,
 [1241] R²⁷ is hydrogen or (C₁-C₄)-alkyl,
 [1242] L² is linear (C₂-C₁₀)-alkanediyl or is a group of the formula



- [1243] where
 [1244] p is a number from 2 to 6,
 [1245] ##3 marks the linkage site with the group B,
 [1246] ##4 marks the linkage site with the nitrogen atom,
 [1247] where (C₂-C₁₀)-alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,
 [1248] and
 [1249] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,
 [1250] D is a group of the formula




- [1251] in which
 [1252] #3 marks the linkage site with the nitrogen atom,
 [1253] R¹ is hydrogen or methyl,

- [1254] R² is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

[1255] or

- [1256] R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[1257] in which

- [1258] #4 marks the linkage site with the adjacent nitrogen atom,

- [1259] #5 marks the linkage site with the carbonyl group,

- [1260] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[1261] in which

- [1262] #6 marks the linkage site with the carbonyl group,


- [1263] R⁶ is hydrogen, hydroxy or benzyloxy,

- [1264] R³ is hydrogen or methyl,

- [1265] R⁴ is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

[1266] or

- [1267] R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[1268] in which

[1269] #7 marks the linkage site with the adjacent nitrogen atom,

[1270] #8 marks the linkage site with the group T¹,

[1271] T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹,

[1272] in which

[1273] R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

[1274] R⁸ is hydrogen or methyl,

[1275] R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,


[1276] or

[1277] R⁸ and R⁹ together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

[1278] R¹⁰ is benzoyl,

[1279] R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

[1280] R⁵ is hydrogen, methyl or a group of the formula



[1281] in which

[1282] #9 marks the linkage site with —CHC(R²⁶)-T²,

[1283] R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

[1284] R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,

[1285] R²⁶ is hydrogen or hydroxy,


[1286] T² is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,

[1287] R³⁵ is methyl or hydroxy,

and also their salts, solvates and solvates of the salts.

[1288] Preferred in the context of the present invention are also compounds of the formula (XXXI) in which

[1289] L¹ is a bond, linear (C₂-C₆)-alkanediyl or a group of the formula



[1290] where


[1291] m is a number 2 or 3,

[1292] ##¹ marks the linkage site with the group G,

[1293] ##² marks the linkage site with the group B,

[1294] where (C₂-C₆)-alkanediyl may be substituted by 1 or 2 methyl substituents,

[1295] B is a bond or a group of the formula



[1296] where

[1297] * marks the linkage site with L¹,

[1298] ** marks the linkage site with L²,

[1299] R¹⁸ is hydrogen,

[1300] R¹⁹ is methyl, propan-2-yl, 2-methylpropan-1-yl or 1-methylpropan-1-yl,

[1301] R²⁰ is hydrogen or (C₁-C₄)-alkyl,

[1302] or

[1303] R¹⁹ and R²⁰ together with the atoms to which they are bonded form a pyrrolidiny ring,

[1304] R²¹ is hydrogen or methyl,


[1305] R²² is hydrogen or methyl,

[1306] or

[1307] R²¹ and R²² together with the atoms to which they are bonded form a cyclopropyl ring,

[1308] R²⁷ is hydrogen or methyl,

[1309] L² is linear (C₂-C₆)-alkanediyl or is a group of the formula



[1310] where

[1311] p is a number 2 or 3,

[1312] ##³ marks the linkage site with the group B,


[1313] ##⁴ marks the linkage site with the nitrogen atom,

[1314] where (C₂-C₁₀)-alkanediyl may be substituted by 1 or 2 methyl substituents,

[1315] and

[1316] where two carbon atoms of the alkanediyl chain in 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a phenyl ring,

[1317] D is a group of the formula



[1318] in which


[1319] #3 marks the linkage site with the nitrogen atom,

[1320] R¹ is hydrogen,

[1321] R² is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-yl-methyl,

[1322] or

[1323] R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[1324] in which

[1325] #4 marks the linkage site with the adjacent nitrogen atom,

[1326] #5 marks the linkage site with the carbonyl group,

[1327] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[1328] in which

[1329] #6 marks the linkage site with the carbonyl group,


[1330] R⁶ is hydrogen, hydroxy or benzyloxy,

[1331] R³ is hydrogen,

[1332] R⁴ is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-yl-methyl,

[1333] or

[1334] R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[1335] in which

[1336] #7 marks the linkage site with the adjacent nitrogen atom,

[1337] #8 marks the linkage site with the group T¹,

[1338] T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹,

[1339] in which

[1340] R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

[1341] R⁸ is hydrogen or methyl,

[1342] R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,


[1343] or

[1344] R⁸ and R⁹ together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

[1345] R¹⁰ is benzoyl,

[1346] R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

[1347] R⁵ is hydrogen, methyl or a group of the formula



[1348] in which

[1349] #9 marks the linkage site with —CHCH₂phenyl,

[1350] R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

[1351] R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,

[1352] R³⁵ is methyl or hydroxy,


and also their salts, solvates and solvates of the salts.

[1353] Particularly preferred in the context of the present invention are also compounds of the formula (XXXI) in which

[1354] L¹ is a bond,

[1355] B is a bond,

[1356] L² is linear (C₂–C₆)-alkanediyl or is a group of the formula




[1357] where

[1358] p is a number 2 or 3,

[1359] ##3 marks the linkage site with the group B,

[1360] ##4 marks the linkage site with the nitrogen atom,

[1361] D is the group of the formula



[1362] where


[1363] #3 marks the linkage site with the nitrogen atom,

[1364] R¹ is hydrogen,

[1365] R² is benzyl or 1H-indol-3-ylmethyl,

[1366] or

[1367] R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[1368] in which

[1369] #4 marks the linkage site with the adjacent nitrogen atom,

[1370] #5 marks the linkage site with the carbonyl group,

[1371] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[1372] in which

[1373] #6 marks the linkage site with the carbonyl group,


[1374] R⁶ is hydrogen, hydroxy or benzyloxy,

[1375] R³ is hydrogen,

[1376] R⁴ is benzyl or 1H-indol-3-ylmethyl,

[1377] or

[1378] R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[1379] in which

[1380] #7 marks the linkage site with the adjacent nitrogen atom,

[1381] #8 marks the linkage site with the group T¹,

[1382] T¹ is a group of the formula —C(=O)—OR⁷ or —C(=O)—NR⁸R⁹,

[1383] in which

[1384] R⁷ is hydrogen,

[1385] R⁸ is hydrogen,

[1386] R⁹ is hydrogen,

[1387] R³⁵ is methyl,


and also their salts, solvates and solvates of the salts.

[1388] Preferred subject matter of the present invention are compounds of the formula (XXXI) as indicated above, in which

[1389] L¹ is a bond,

[1390] B is a bond,

[1391] L² is linear (C₂-C₆)-alkanediyl or is a group of the formula




[1392] where

[1393] p is a number 2 or 3,

[1394] ##3 marks the linkage site with the group B,

[1395] ##4 marks the linkage site with the nitrogen atom,

[1396] D is a group of the formula



[1397] where


[1398] #3 marks the linkage site with the nitrogen atom,

[1399] R¹ is hydrogen,

[1400] R² is benzyl, 4-hydroxybenzyl or 1H-indol-3-ylmethyl,

[1401] or

[1402] R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[1403] in which

[1404] #⁴ marks the linkage site with the adjacent nitrogen atom,

[1405] #⁵ marks the linkage site with the carbonyl group,

[1406] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[1407] in which

[1408] #⁶ marks the linkage site with the carbonyl group,


[1409] R⁶ is hydrogen, hydroxy or benzyloxy,

[1410] R³ is hydrogen,

[1411] R⁴ is benzyl, 4-hydroxybenzyl or 1H-indol-3-yl-methyl,

[1412] or

[1413] R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[1414] in which

[1415] #⁷ marks the linkage site with the adjacent nitrogen atom,

[1416] #⁸ marks the linkage site with the group T¹,

[1417] T¹ is a group of the formula —C(=O)—OR⁷ or —C(=O)—NR⁸R⁹,

[1418] in which

[1419] R⁷ is hydrogen,

[1420] R⁸ is hydrogen,

[1421] R⁹ is hydrogen,

[1422] R³⁵ is methyl,

and also their salts, solvates and solvates of the salts.

[1423] Particularly preferred are the following compounds of the formulae (XXXa) and (XXXI) selected from the following group:

[1424] N-[6-(3-[(2R)-2-amino-2-carboxyethyl]sulphonyl)-2,5-dioxopyrrolidin-1-yl]hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(1S)-1-carboxy-2-(1H-indol-3-yl)ethyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide,


[1425] N-[6-(3-[(2R)-2-amino-2-carboxyethyl]sulphonyl)-2,5-dioxopyrrolidin-1-yl]hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide,

[1426] N-(6-{[(5S)-5-amino-5-carboxypentyl]amino}-6-oxohexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate,

[1427] N-(6-{[(5S)-5-amino-5-carboxypentyl]amino}-6-oxohexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(1S)-1-carboxy-2-(1H-indol-3-yl)ethyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide,

and also their salts, solvates and solvates of the salts.

[1428] The present invention additionally provides binder-drug conjugates of the general formula (I)



in which

[1429] n is a number from 1 to 50,

[1430] AK is a binder,


[1431] the group §-G-L¹-B-L²-§§ is a linker,

[1432] where

[1433] § marks the linkage site with the group AK and

[1434] §§ marks the linkage site with the nitrogen atom,

[1435] D is a group of the formula



[1436] where


[1437] #³ marks the linkage site with the nitrogen atom,

[1438] R¹ is hydrogen,

[1439] R² is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

[1440] or

[1441] R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[1442] in which

[1443] #⁴ marks the linkage site with the adjacent nitrogen atom,

[1444] #⁵ marks the linkage site with the carbonyl group,

[1445] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[1446] in which

[1447] #⁶ marks the linkage site with the carbonyl group,


[1448] R⁶ is hydrogen, hydroxy or benzyloxy,

[1449] R³ is hydrogen,

[1450] R⁴ 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

[1451] or

[1452] R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[1453] in which

[1454] #⁷ marks the linkage site with the adjacent nitrogen atom,

[1455] #⁸ marks the linkage site with the group T¹,

[1456] T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹,

[1457] in which

[1458] R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

[1459] R⁸ is hydrogen or methyl,

[1460] R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,


[1461] or

[1462] R⁸ and R⁹ together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

[1463] R¹⁰ is benzoyl,

[1464] R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

[1465] R⁵ is hydrogen, methyl or a group of the formula



[1466] in which

[1467] #⁹ marks the linkage site with —CHC(R²⁶)-T²,

[1468] R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

[1469] R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,

[1470] R²⁶ is hydrogen or hydroxy,

[1471] T² is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,

and also their salts, solvates and solvates of the salts.

[1472] Preferred subject matter of the invention are binder-drug conjugates of the general formula (I), in which

[1473] n is a number from 1 to 50,


[1474] AK is AK₁ or AK₂

[1475] where

[1476] AK_1 is a binder which is bonded via a sulphur atom of the binder to the group G,

[1477] AK_2 is a binder which is bonded via a nitrogen atom of the binder to the group G,

[1478] G when $AK=AK_1$, is a group of the formula



[1479] where


[1480] #¹ marks the linkage site with the sulphur atom of the binder,

[1481] #² marks the linkage site with the group L^1 ,

[1482] or

[1483] when $AK=AK_2$, is carbonyl,

[1484] L^1 is a bond, linear (C_1 - C_{10})-alkanediyl or is a group of the formula



[1485] where

[1486] m is a number from 2 to 6,

[1487] ##¹ marks the linkage site with the group G,


[1488] ##² marks the linkage site with the group B,

[1489] where (C_1 - C_{10})-alkanediyl may be substituted by 1 to 4 methyl substituents,


[1490] and

[1491] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C_3 - C_6)-cycloalkyl ring or a phenyl ring,

[1492] B is a bond or a group of the formula



-continued



[1493] where


[1494] * marks the linkage site with L^1 ,

[1495] ** marks the linkage site with L^2 ,

[1496] P is O or NH,

[1497] L^3 is a bond or (C_2 - C_4)-alkanediyl,

[1498] L^4 is a bond or a group of the formula



[1499] in which

[1500] *** marks the linkage site with the carbonyl group,

[1501] **** marks the linkage site with L^2 ,

[1502] R^{25} is hydrogen or methyl,

[1503] Q^1 is a 4- to 7-membered heterocycle,

[1504] Q^2 is a 3- to 7-membered carbocycle or a 4- to 7-membered heterocycle,

[1505] R^{14} is hydrogen or (C_1 - C_4)-alkyl,

[1506] R^{15} is hydrogen or (C_1 - C_4)-alkyl,

[1507] or

[1508] R^{14} and R^{15} together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

[1509] R^{16} is hydrogen or (C_1 - C_4)-alkyl,

[1510] R^{17} is hydrogen or (C_1 - C_4)-alkyl,

[1511] or

[1512] R^{16} and R^{17} together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

[1513] R^{18} is hydrogen or (C_1 - C_4)-alkyl,

[1514] R^{19} is hydrogen or the side group of a natural α -amino acid or of its homologues or isomers,

[1515] R^{20} is hydrogen or (C_1 - C_4)-alkyl,

[1516] or

[1517] R^{19} and R^{20} together with the atoms to which they are bonded form a pyrrolidinyl ring,

[1518] R^{21} is hydrogen or (C_1 - C_4)-alkyl,

[1519] R^{22} is hydrogen or (C_1 - C_4)-alkyl,

[1520] or


[1521] R^{21} and R^{22} together with the atoms to which they are bonded form a 3- to 7-membered carbocycle,

[1522] R^{23} is (C_1 - C_4)-alkyl,


[1523] R^{24} is hydrogen or (C_1 - C_4)-alkyl,

[1524] R^{27} is hydrogen or (C_1 - C_4)-alkyl,


[1525] L^2 is linear (C_2 - C_{10})-alkanediyl or is a group of the formula




- [1526] where
 [1527] p is a number from 2 to 6,
 [1528] ##³ marks the linkage site with the group B,
 [1529] ##⁴ marks the linkage site with the nitrogen atom,
 [1530] where (C₂-C₁₀)-alkanediyl may be substituted by 1 to 4 methyl substituents, and
 [1531] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,
 [1532] D has the definitions indicated above, and also their salts, solvates and solvates of the salts.
 [1533] Preferred subject matter of the invention are binder-drug conjugates of the general formula (I), in which
 [1534] n is a number from 1 to 50,
 [1535] AK is AK₁ or AK₂
 [1536] where
 [1537] AK₁ is an antibody or an antigen-binding antibody fragment and are bonded via a sulphur atom to the group G,
 [1538] AK₂ is an antibody or an antigen-binding antibody fragment and are bonded via a nitrogen atom to the group G,
 [1539] G, L¹, B, L² and D have the definitions indicated above, and also their salts, solvates and solvates of the salts.
 [1540] Preferred subject matter of the present invention are binder-drug conjugates of the general formula (I), in which
 [1541] n is a number from 1 to 20,
 [1542] AK is AK₁ or AK₂
 [1543] where
 [1544] AK₁ is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,
 [1545] AK₂ is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the NH side group of a lysine residue of the binder to the group G,
 [1546] G when AK=AK₁, is a group of the formula




- [1547] in which
 [1548] #¹ marks the linkage site with the cysteine residue of the binder,
 [1549] #² marks the linkage site with the group L¹,
 [1550] or
 [1551] when AK=AK₂, is carbonyl,
 [1552] L¹ is a bond, linear (C₂-C₆)-alkanediyl or is a group of the formula



- [1553] where
 [1554] m is a number from 2 to 6,
 [1555] ##¹ marks the linkage site with the group G,
 [1556] ##² marks the linkage site with the group B,
 [1557] where (C₂-C₆)-alkanediyl may be substituted by 1 or 2 methyl substituents,
 [1558] B is a bond or a group of the formula




- [1559] where
 [1560] * marks the linkage site with L¹,
 [1561] ** marks the linkage site with L²,
 [1562] L³ is a bond or ethane-1,2-diyl,
 [1563] L⁴ is a bond or a group of the formula




- [1564] in which
 [1565] *** marks the linkage site with the carbonyl group,
 [1566] **** marks the linkage site with L²,
 [1567] R²⁵ is methyl,
 [1568] Q¹ is a 4- to 6-membered carbocycle or piperidine-1,4-diyl,
 [1569] R¹⁴ is hydrogen,
 [1570] R¹⁵ is hydrogen,
 [1571] R¹⁶ is hydrogen or methyl,
 [1572] R¹⁷ is hydrogen or methyl,
 [1573] or
 [1574] R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a piperazinyl ring,
 [1575] R¹⁸ is hydrogen,
 [1576] R¹⁹ is hydrogen, methyl, propan-2-yl, 2-methylpropan-1-yl or 1-methylpropan-1-yl,
 [1577] R²⁰ is hydrogen or methyl,


- [1578] or
 [1579] R^{19} and R^{20} together with the atoms to which they are bonded form a pyrrolidinyl ring,
 [1580] R^{21} is hydrogen or methyl,
 [1581] R^{22} is hydrogen or methyl,
 [1582] or
 [1583] R^{21} and R^{22} together with the atoms to which they are bonded form a cyclopropyl ring,
 [1584] R^{23} is methyl,
 [1585] R^{24} is hydrogen or methyl,
 [1586] L^2 is linear (C_2-C_6)-alkanediyl or is a group of the formula



- [1587] where
 [1588] p is a number from 2 to 6,
 [1589] ##3 marks the linkage site with the group B,
 [1590] ##4 marks the linkage site with the nitrogen atom,
 [1591] where (C_2-C_6)-alkanediyl may be substituted by 1 or 2 methyl substituents,
 [1592] D is a group of the formula




- [1593] where
 [1594] #3 marks the linkage site with the nitrogen atom,
 [1595] R^1 is hydrogen,
 [1596] R^2 is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,
 [1597] or
 [1598] R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




- [1599] in which
 [1600] #4 marks the linkage site with the adjacent nitrogen atom,
 [1601] #5 marks the linkage site with the carbonyl group,

- [1602] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula




- [1603] in which
 [1604] #6 marks the linkage site with the carbonyl group,
 [1605] R^6 is hydrogen, hydroxy or benzyloxy,
 [1606] R^3 is hydrogen,
 [1607] R^4 is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,
 [1608] or
 [1609] R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



- [1610] in which
 [1611] #7 marks the linkage site with the adjacent nitrogen atom,
 [1612] #8 marks the linkage site with the group T^1 ,
 [1613] T^1 is a group of the formula $-C(=O)-OR^7$, $-C(=O)-NR^8R^9$, $-C(=O)-NH-NH-R^{10}$ or $-CH_2-O-R^{11}$,
 [1614] in which
 [1615] R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,
 [1616] R^8 is hydrogen or methyl,
 [1617] R^9 is hydrogen, methyl, ethyl, n-propyl or benzyl,
 [1618] or
 [1619] R^8 and R^9 together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,
 [1620] R^{10} is benzoyl,
 [1621] R^{11} is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

[1622] R^5 is hydrogen, methyl or a group of the formula



[1623] in which

[1624] #⁹ marks the linkage site with $-\text{CHC}(R^{26})-\text{T}^2$,

[1625] R^{12} is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula $-\text{S}(\text{O})_2\text{OH}$,

[1626] R^{13} is phenyl which may be substituted by methoxycarbonyl or carboxyl,

[1627] R^{26} is hydrogen or hydroxy,

[1628] T^2 is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,

and also their salts, solvates and solvates of the salts.

[1629] Particularly preferred subject matter of the present invention are binder-drug conjugates of the general formula (I), in which

[1630] n is a number from 1 to 10,


[1631] AK is AK_1 or AK_2

[1632] where

[1633] AK_1 is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

[1634] AK_2 is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the NH side group of a lysine residue of the binder to the group G,

[1635] G when $\text{AK}=\text{AK}_1$, is a group of the formula



[1636] in which


[1637] #¹ marks the linkage site with the cysteine residue of the binder,

[1638] #² marks the linkage site with the group L^1 ,

[1639] or

[1640] when $\text{AK}=\text{AK}_2$, is carbonyl,

[1641] L^1 is a bond, linear (C_2-C_6) -alkanediyl or is a group of the formula



[1642] where


[1643] m is a number 2 or 3,

[1644] ##¹ marks the linkage site with the group G,

[1645] ##² marks the linkage site with the group B,

[1646] where (C_2-C_6) -alkanediyl may be substituted by 1 or 2 methyl substituents,

[1647] B is a bond or a group of the formula




[1648] where

[1649] * marks the linkage site with L^1 ,

[1650] ** marks the linkage site with L^2 ,

[1651] L^3 is a bond or ethane-1,2-diyl,

[1652] L^4 is a bond or a group of the formula



[1653] where

[1654] *** marks the linkage site with the carbonyl group,

[1655] **** marks the linkage site with L^2 ,

[1656] R^{25} is methyl,

[1657] Q^1 is piperidine-1,4-diyl,

[1658] R^{16} is hydrogen or methyl,

[1659] R^{17} is hydrogen or methyl,

[1660] or

[1661] R^{16} and R^{17} together with the atoms to which they are bonded form a piperazinyl ring,


[1662] R^{21} is hydrogen or methyl,

[1663] R^{22} is hydrogen or methyl,


[1664] or

[1665] R^{21} and R^{22} together with the atoms to which they are bonded form a cyclopropyl ring,


- [1666] R^{23} is methyl,
 [1667] R^{24} is hydrogen,
 [1668] L^2 is linear (C_2-C_6)-alkanediyl,
 [1669] D is a group of the formula




- [1670] where
 [1671] #³ marks the linkage site with the nitrogen atom,
 [1672] R^1 is hydrogen,
 [1673] R^2 is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,
 [1674] or
 [1675] R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




- [1676] in which
 [1677] #⁴ marks the linkage site with the adjacent nitrogen atom,
 [1678] #⁵ marks the linkage site with the carbonyl group,
 [1679] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



- [1680] in which
 [1681] #⁶ marks the linkage site with the carbonyl group,
 [1682] R^6 is hydrogen, hydroxy or benzyloxy,
 [1683] R^3 is hydrogen,
 [1684] R^4 is benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,
 [1685] or
 [1686] R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




- [1687] in which
 [1688] #⁷ marks the linkage site with the adjacent nitrogen atom,
 [1689] #⁸ marks the linkage site with the group T,
 [1690] T^1 is a group of the formula $-C(=O)-OR^7$, $-C(=O)-NR^8R^9$ or $-CH_2-O-R^{11}$, in which
 [1691] R^7 is hydrogen, methyl, ethyl, n-propyl, benzyl or adamantylmethyl,
 [1692] R^8 is hydrogen or methyl,
 [1693] R^9 is hydrogen, methyl, ethyl, n-propyl or benzyl,
 [1694] R^{11} is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,
 [1695] R^5 is hydrogen or a group of the formula



- [1696] in which
 [1697] #⁹ marks the linkage site with $-CHC(R^{26})$ phenyl,
 [1698] R^{12} is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula $-S(O)_2OH$,
 [1699] R^{13} is phenyl which may be substituted by methoxycarbonyl or carboxyl,
 and also their salts, solvates and solvates of the salts.
 [1700] The present invention additionally provides compounds of the formula (XXX)


(XXX)



in which

[1701] Cys is a cysteine residue which is bonded via the sulphur atom of the side chain to a carbon atom of the succinimide.

[1702] L^1 is a bond, linear (C_1 - C_{10})-alkanediyl or is a group of the formula



[1703] in which

[1704] m is a number from 2 to 6,

[1705] ##¹ marks the linkage site with the group G,


[1706] ##² marks the linkage site with the group B,

[1707] where (C_1 - C_{10})-alkanediyl may be substituted by 1 to 4 methyl substituents,

[1708] and

[1709] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C_3 - C_6)-cycloalkyl ring or a phenyl ring,

[1710] B is a bond or a group of the formula



[1711] where


[1712] * marks the linkage site with L^1 ,

[1713] ** marks the linkage site with L^2 ,


[1714] P is O or NH,

[1715] L^3 is a bond or (C_2 - C_4)-alkanediyl,

[1716] L^4 is a bond or a group of the formula



-continued



[1717] in which

[1718] *** marks the linkage site with the carbonyl group,

[1719] **** marks the linkage site with L^2 ,

[1720] R^{25} is hydrogen or methyl

[1721] Q^1 is a 3- to 7-membered carbocycle or a 4- to 7-membered aza heterocycle,

[1722] Q^2 is a 3- to 7-membered carbocycle or a 4- to 7-membered aza heterocycle,

[1723] R^{14} is hydrogen or (C_1 - C_4)-alkyl,

[1724] R^{15} is hydrogen or (C_1 - C_4)-alkyl,

[1725] or

[1726] R^{14} and R^{15} together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

[1727] R^{16} is hydrogen or (C_1 - C_4)-alkyl,

[1728] R^{17} is hydrogen or (C_1 - C_4)-alkyl,

[1729] or

[1730] R^{16} and R^{17} together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

[1731] R^{18} is hydrogen or (C_1 - C_4)-alkyl,

[1732] R^{19} is hydrogen or the side group of a natural α -amino acid or of its homologues or isomers,

[1733] R^{20} is hydrogen or (C_1 - C_4)-alkyl,

[1734] or

[1735] R^{19} and R^{20} together with the atoms to which they are bonded form a pyrrolidiny ring,

[1736] R^{21} is hydrogen or (C_1 - C_4)-alkyl,

[1737] R^{22} is hydrogen or (C_1 - C_4)-alkyl,

[1738] or


[1739] R^{21} and R^{22} together with the atoms to which they are bonded form a 3- to 7-membered carbocycle,

[1740] R^{23} is (C_1 - C_4)-alkyl,

[1741] R^{24} is hydrogen or (C_1 - C_4)-alkyl,

[1742] R^{27} is hydrogen or (C_1 - C_4)-alkyl,

[1743] L^2 is linear (C_2 - C_{10})-alkanediyl or is a group of the formula



[1744] where

[1745] p is a number from 2 to 6,

[1746] ##³ marks the linkage site with the group B,


[1747] ##⁴ marks the linkage site with the nitrogen atom,

[1748] where (C_2 - C_{10})-alkanediyl may be substituted by 1 to 4 substituents,

[1749] and

[1750] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C_3 - C_6)-cycloalkyl ring or a phenyl ring,

[1751] D is a group of the formula



[1752] in which


[1753] #3 marks the linkage site with the nitrogen atom,

[1754] R¹ is hydrogen,

[1755] R² is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

[1756] or

[1757] R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[1758] in which

[1759] #4 marks the linkage site with the adjacent nitrogen atom,

[1760] #5 marks the linkage site with the carbonyl group,

[1761] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[1762] in which

[1763] #6 marks the linkage site with the carbonyl group,


[1764] R⁶ is hydrogen, hydroxy or benzyloxy,

[1765] R³ is hydrogen,

[1766] R⁴ is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

[1767] or

[1768] R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[1769] in which

[1770] #7 marks the linkage site with the adjacent nitrogen atom,

[1771] #8 marks the linkage site with the group T¹,

[1772] T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹, in which

[1773] R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

[1774] R⁸ is hydrogen or methyl,

[1775] R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,


[1776] or

[1777] R⁸ and R⁹ together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

[1778] R¹⁰ is benzoyl,

[1779] R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

[1780] R⁵ is hydrogen, methyl or a group of the formula



[1781] in which

[1782] #9 marks the linkage site with —CHC(R²⁶)-T²,

[1783] R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

[1784] R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,

[1785] R²⁶ is hydrogen or hydroxy,


[1786] T² is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,

and also their salts, solvates and solvates of the salts.

[1787] Particularly preferred in the context of the present invention in addition are also compounds of the formula (XXX), in which

[1788] Cys is a cysteine residue which is bonded via the sulphur atom of the side chain via a carbon atom of the succinimide,

[1789] L¹ is a bond, linear (C₂-C₆)-alkanediyl or is a group of the formula



[1790] in which


[1791] m is a number from 2 to 6,

[1792] ##¹ marks the linkage site with the group G,

[1793] ##² marks the linkage site with the group B,

[1794] where (C₂-C₆)-alkandiyl may be substituted by 1 or 2 methyl substituents,

[1795] B is a bond or a group of the formula



[1796] where

[1797] * marks the linkage site with L¹,

[1798] ** marks the linkage site with L²,

[1799] L³ is a bond or ethane-1,2-diyl,

[1800] L⁴ is a bond,

[1801] R¹⁴ is hydrogen,

[1802] R¹⁵ is hydrogen,

[1803] R¹⁶ is hydrogen or methyl,

[1804] R¹⁷ is hydrogen or methyl,


[1805] or

[1806] R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a piperazinyl ring,

[1807] R²³ is methyl,

[1808] R²⁴ is hydrogen or methyl,

[1809] L² is linear (C₂-C₆)-alkanediyl or is a group of the formula




[1810] where

[1811] p is a number 2 or 3,

[1812] ##³ marks the linkage site with the group B,

[1813] ##⁴ marks the linkage site with the nitrogen atom,

[1814] D is a group of the formula



[1815] in which


[1816] #³ marks the linkage site with the nitrogen atom,

[1817] R¹ is hydrogen,

[1818] R² is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

[1819] or

[1820] R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[1821] in which

[1822] #⁴ marks the linkage site with the adjacent nitrogen atom,

[1823] #⁵ marks the linkage site with the carbonyl group,

[1824] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[1825] in which

[1826] #⁶ marks the linkage site with the carbonyl group,


[1827] R⁶ is hydrogen, hydroxy or benzyloxy,

[1828] R³ is hydrogen,

[1829] R⁴ is benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

[1830] or

[1831] R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[1832] in which

[1833] #⁷ marks the linkage site with the adjacent nitrogen atom,

[1834] #⁸ marks the linkage site with the group T,

[1835] T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹,

[1836] in which

[1837] R⁷ is hydrogen, methyl, ethyl, n-propyl, benzyl or adamantylmethyl,


[1838] R⁸ is hydrogen or methyl,

[1839] R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,

[1840] R¹⁰ is benzoyl,

[1841] R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

[1842] R⁵ is hydrogen or a group of the formula



[1843] in which

[1844] #⁹ marks the linkage site with —CHC(R²⁶)phenyl,

[1845] R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

[1846] R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,

and also their salts, solvates and solvates of the salts.

[1847] Preferred in the context of the present invention are also compounds of the formula (Ia), in which n=1-20, more preferably n=1-10 and very preferably n=2-8.


[1848] Preferred in the context of the present invention are also compounds of the formula (Ia), in which

[1849] AK is AK₁

[1850] where

[1851] AK₁ is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

[1852] G is a group of the formula



[1853] where

[1854] #¹ marks the linkage site with the cysteine residue of the binder,

[1855] #² marks the linkage site with the group L¹,

and

[1856] n, L¹, B, L², D and R³⁵ have the definitions indicated above,

and also their salts, solvates and solvates of the salts.

[1857] Preferred in the context of the present invention are also compounds of the formula (Ia), in which

[1858] AK is AK₂

[1859] where

[1860] AK₂ is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the NH side group of a lysine residue of the binder to the group G,

[1861] G is carbonyl,

and

[1862] n, L¹, B, L², D and R³⁵ have the definitions indicated above,

and also their salts, solvates and solvates of the salts.


[1863] Preference in the context of the present invention is also given to compounds of the formula (Ia), in which

[1864] AK is AK₁

[1865] where

[1866] AK₁ is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and which is attached via the sulphur atom of a cysteine residue of the binder to the group G,

[1867] G is a group of the formula



[1868] where

[1869] #¹ marks the linkage site with the cysteine residue of the binder,

[1870] #² marks the linkage site with the group L¹,

and

[1871] n, L¹, B, L², D and R³⁵ have the definitions indicated above,

and also their salts, solvates and solvates of the salts.

[1872] Preference in the context of the present invention is also given to compounds of the formula (Ia), in which

[1873] AK is AK₂

[1874] where

[1875] AK₂ is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and which is bonded via the NH side group of a lysine residue of the binder to the group G,

[1876] G is carbonyl,
and

[1877] n, L¹, B, L², D and R³⁵ have the definitions indicated above,

and also their salts, solvates and solvates of the salts.

[1878] Preference in the context of the present invention is also given to compounds of the general formula (Ia), in which

[1879] AK is AK₂

[1880] where


[1881] AK₂ is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and which is bonded via the NH side group of a lysine residue of the binder to the group G,

[1882] G is carbonyl,

[1883] L¹ is a bond,

[1884] B is a bond,

[1885] L² is linear (C₃-C₆)-alkanediyl or is a group of the formula



[1886] where

[1887] p is a number 2 or 3,

[1888] ##³ marks the linkage site with the group B,

[1889] ##⁴ marks the linkage site with the nitrogen atom,

[1890] n, D and R³⁵ have the definitions indicated above,
and also their salts, solvates and solvates of the salts.


[1891] Preference in the context of the present invention is also given to compounds of the general formula (Ia), in which

[1892] AK is AK₁

[1893] where

[1894] AK₁ is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and which is attached via the sulphur atom of a cysteine residue of the binder to the group G,

[1895] G is a group of the formula




[1896] where

[1897] #¹ marks the linkage site with the cysteine residue of the binder,

[1898] #² marks the linkage site with the group L¹,

[1899] L¹ is a bond, linear (C₃-C₅)-alkanediyl or a group of the formula



[1900] where


[1901] m is a number 2 or 3,

[1902] ##¹ marks the linkage site with the group G,

[1903] ##² marks the linkage site with the group B,

[1904] where (C₃-C₅)-alkanediyl may be substituted by 1 or 2 methyl substituents,

[1905] B is a bond or a group of the formula




[1906] where

[1907] * marks the linkage site with L¹,

[1908] ** marks the linkage site with L²,

[1909] L³ is a bond or ethane-1,2-diyl,

[1910] L⁴ is a bond or a group of the formula



[1911] in which

[1912] *** marks the linkage site with the carbonyl group,

[1913] **** marks the linkage site with L²,

[1914] R²⁵ is methyl,

[1915] R²⁸ is hydrogen, methylcarbonyl or tert-butoxycarbonyl,


[1916] R¹⁶ is hydrogen or methyl,

[1917] R¹⁷ is hydrogen or methyl,


[1918] or

[1919] R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a piperaziny ring,


[1920] L² is linear (C₃-C₅)-alkanediyl or is a group of the formula




- [1921] where
 [1922] p is a number 2 or 3,
 [1923] ##^3 marks the linkage site with the group B,
 [1924] ##^4 marks the linkage site with the nitrogen atom,
 and
 [1925] n, D and R^{35} have the definitions indicated above,
 and also their salts, solvates and solvates of the salts.
 [1926] Preferred in the context of the present invention are
 also compounds of the formula (Ia), (XXXa) and (XXXI), in
 which
 [1927] L^1 is a bond,
 [1928] B is a bond,
 [1929] L^2 is linear ($\text{C}_3\text{-C}_6$)-alkanediyl or is a group of the
 formula




- [1930] where
 [1931] p is a number 2 or 3,
 [1932] ##^3 marks the linkage site with the group B,
 [1933] ##^4 marks the linkage site with the nitrogen atom,
 and
 [1934] n, AK, Cys, G, D and R^{35} have the definitions indicated above,
 and also their salts, solvates and solvates of the salts.
 [1935] Preferred in the context of the present invention are
 also compounds of the formula (Ia), in which
 [1936] L^1 is linear ($\text{C}_1\text{-C}_{10}$)-alkanediyl or a group of the
 formula




- [1937] where
 [1938] m is a number from 2 to 6,
 [1939] ##^1 marks the linkage site with the group G,
 [1940] ##^2 marks the linkage site with the group B,
 [1941] where ($\text{C}_1\text{-C}_{10}$)-alkanediyl may be substituted by
 1 to 4 substituents selected independently of one another
 from the group consisting of methyl, hydroxyl and benzyl,
 [1942] B is a bond or a group of the formula



- [1943] where
 [1944] * marks the linkage site with L^1 ,
 [1945] ** marks the linkage site with L^2 ,
 [1946] L^3 is a bond or ($\text{C}_2\text{-C}_4$)-alkanediyl,
 [1947] L^4 is a group of the formula




- [1948] in which
 [1949] *** marks the linkage site with the carbonyl
 group,
 [1950] **** marks the linkage site with L^2 ,
 [1951] R^{25} is hydrogen or methyl,
 [1952] R^{28} is hydrogen, ($\text{C}_1\text{-C}_4$)-alkylcarbonyl, tert-
 butyloxycarbonyl or benzyloxycarbonyl,
 [1953] Q^1 is a 4- to 7-membered heterocycle,
 [1954] R^{16} is hydrogen or ($\text{C}_1\text{-C}_4$)-alkyl,
 [1955] R^{17} is hydrogen or ($\text{C}_1\text{-C}_4$)-alkyl,
 [1956] or
 [1957] R^{16} and R^{17} together with the atoms to which they
 are bonded form a 5- or 6-membered heterocycle,
 [1958] R^{23} is ($\text{C}_1\text{-C}_4$)-alkyl,
 [1959] R^{24} is hydrogen or ($\text{C}_1\text{-C}_4$)-alkyl,
 [1960] R^{36} is hydrogen, ($\text{C}_1\text{-C}_4$)-alkylcarbonyl, tert-bu-
 tyloxycarbonyl or benzyloxycarbonyl,
 [1961] R^{37} is hydrogen or methyl,
 [1962] or
 [1963] R^{36} and R^{37} together with the atoms to which they
 are bonded form a pyrrolidine ring,
 [1964] L^2 is linear ($\text{C}_2\text{-C}_{10}$)-alkanediyl or is a group of the
 formula




- [1965] where
 [1966] p is a number from 2 to 6,
 [1967] ##^3 marks the linkage site with the group B,
 [1968] ##^4 marks the linkage site with the nitrogen atom,
 [1969] where ($\text{C}_2\text{-C}_{10}$)-alkanediyl may be substituted by
 1 to 4 substituents selected independently of one another
 from the group consisting of methyl, hydroxyl and benzyl,


- and
 [1970] n, AK, G, D and R^{35} have the definitions indicated
 above,
 and also their salts, solvates and solvates of the salts.
 [1971] Preferred in the context of the present invention are
 also compounds of the formula (Ia), in which
 [1972] L^1 is linear ($\text{C}_2\text{-C}_6$)-alkanediyl or a group of the
 formula




- [1973] where
 [1974] m is a number 2 or 3,
 [1975] ##¹ marks the linkage site with the group G,
 [1976] ##² marks the linkage site with the group B,
 [1977] B is a bond or a group of the formula



- [1978] where
 [1979] * marks the linkage site with L¹,
 [1980] ** marks the linkage site with L²,
 [1981] L³ is a bond or ethane-1,2-diyl,
 [1982] L⁴ is a group of the formula



- [1983] where
 [1984] *** marks the linkage site with the carbonyl group,
 [1985] **** marks the linkage site with L²,
 [1986] R²⁵ is hydrogen or methyl,
 [1987] R²⁸ is hydrogen, methylcarbonyl or tert-butyloxycarbonyl,
 [1988] R¹⁶ is hydrogen or methyl,
 [1989] R¹⁷ is hydrogen or methyl,
 [1990] or
 [1991] R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a piperaziny ring,
 [1992] R³⁶ is hydrogen, methylcarbonyl or tert-butyloxycarbonyl,
 [1993] R³⁷ is hydrogen or methyl,
 [1994] or
 [1995] R³⁶ and R³⁷ together with the atoms to which they are bonded form a pyrrolidine ring,
 [1996] L² is linear (C₂-C₆)-alkanediyl or is a group of the formula



- [1997] where
 [1998] p is a number 2 or 3,
 [1999] ##³ marks the linkage site with the group B,
 [2000] ##⁴ marks the linkage site with the nitrogen atom,


and

- [2001] n, AK, G, D and R³⁵ have the definitions indicated above,


and also their salts, solvates and solvates of the salts.

- [2002] Preferred in the context of the present invention are also compounds of the formula (1a) in which


- [2003] L¹ is a bond, linear (C₂-C₆)-alkanediyl or a group of the formula




- [2004] where
 [2005] m is a number 2 or 3,
 [2006] ##¹ marks the linkage site with the group G,
 [2007] ##² marks the linkage site with the group B,
 [2008] where (C₂-C₆)-alkanediyl may be substituted by 1 or 2 methyl substituents,
 [2009] B is a bond or a group of the formula




- [2010] where
 [2011] * marks the linkage site with L¹,
 [2012] ** marks the linkage site with L²,
 [2013] L³ is a bond or ethane-1,2-diyl,
 [2014] L⁴ is a bond or a group of the formula








- [2015] in which
 [2016] *** marks the linkage site with the carbonyl group,
 [2017] **** marks the linkage site with L²,
 [2018] R²⁵ is methyl,
 [2019] R²⁸ is hydrogen, methylcarbonyl or tert-butyloxy carbonyl,
 [2020] Q¹ is piperidine-1,4-diyl,
 [2021] R¹⁶ is hydrogen or methyl,
 [2022] R¹⁷ is hydrogen or methyl,
 [2023] or
 [2024] R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a piperaziny ring,
 [2025] R²¹ is hydrogen or methyl,
 [2026] R²² is hydrogen or methyl,
 [2027] or
 [2028] R²¹ and R²² together with the atoms to which they are bonded form a cyclopropyl ring,
 [2029] R²³ is methyl,
 [2030] R²⁴ is hydrogen,
 [2031] R³⁶ is hydrogen, methylcarbonyl or tert-butyloxy carbonyl,
 [2032] R³⁷ is hydrogen or methyl,
 [2033] L² is linear (C₂-C₆)-alkanediyl or is a group of the formula




- [2034] where
 [2035] p is a number from 2 to 6,
 [2036] ###³ marks the linkage site with the group B,
 [2037] ###⁴ marks the linkage site with the nitrogen atom, and also their salts, solvates and solvates of the salts.
 [2038] Preferred in the context of the present invention are also compounds of the formula (Ia) in which
 [2039] L¹ is linear (C₂-C₆)-alkanediyl,
 [2040] where (C₂-C₆)-alkanediyl may be substituted by 1 or 2 methyl substituents,
 [2041] B is a bond or a group of the formula




- [2042] where
 [2043] * marks the linkage site with L¹,
 [2044] ** marks the linkage site with L²,
 [2045] L³ is a bond,
 [2046] L⁴ is a group of the formula




- [2047] in which
 [2048] *** marks the linkage site with the carbonyl group,
 [2049] **** marks the linkage site with L²,
 [2050] R²⁵ is methyl,
 [2051] R²⁸ is hydrogen, methylcarbonyl or tert-butyloxy carbonyl,
 [2052] R¹⁶ is hydrogen or methyl,
 [2053] R¹⁷ is hydrogen or methyl,
 [2054] R²³ is methyl,
 [2055] R²⁴ is hydrogen,
 [2056] R³⁶ is hydrogen, methylcarbonyl or tert-butyloxy carbonyl,
 [2057] R³⁷ is hydrogen or methyl,
 [2058] L² is linear (C₂-C₆)-alkanediyl, and also their salts, solvates and solvates of the salts.
 [2059] Preferred in the context of the present invention are also compounds of the formula (Ia) and (XXXa), in which
 [2060] G is a group of the formula




- [2061] in which
 [2062] #¹ marks the linkage site with the cysteine residue of the binder,
 [2063] #² marks the linkage site with the group L¹,
 [2064] L¹ is linear (C₃-C₅)-alkanediyl or a group of the formula




- [2065] in which
 [2066] m is a number 2 or 3,
 [2067] ##¹ marks the linkage site with the group G,
 [2068] ##² marks the linkage site with the group B,
 [2069] where (C₃-C₅)-alkanediyl may be substituted by
 1 or 2 methyl substituents,
 [2070] B is a bond or a group of the formula




- [2071] where
 [2072] * marks the linkage site with L¹,
 [2073] ** marks the linkage site with L²,
 [2074] L³ is a bond or ethane-1,2-diyl,
 [2075] L⁴ is a bond,
 [2076] L² is linear (C₃-C₅)-alkanediyl or is a group of the
 formula




- [2077] where
 [2078] p is a number 2 or 3,
 [2079] ##³ marks the linkage site with the group B,
 [2080] ##⁴ marks the linkage site with the nitrogen atom,
 and
 [2081] n, AK₁, Cys, D, R¹⁶ and R¹⁷ have the definitions
 indicated above,
 and also their salts, solvates and solvates of the salts.
 [2082] Preferred in the context of the present invention are
 also compounds of the formula (Ia) and (XXXa), in which
 [2083] B is a bond or a group of the formula




- [2084] where
 [2085] * marks the linkage site with L¹,
 [2086] ** marks the linkage site with L²,
 [2087] L³ is a bond or ethane-1,2-diyl,
 [2088] L⁴ is a bond,
 [2089] n, AK, Cys, G, L¹, L², D, R¹⁶, R¹⁷ and R³⁵ have the
 definitions indicated above,
 and also their salts, solvates and solvates of the salts.
 [2090] Preferred in the context of the present invention are
 also compounds of the formula (Ia), (XXXa) and (XXXI), in
 which
 [2091] L¹ is a bond, linear (C₃-C₅)-alkanediyl or a group of the
 formula




- [2092] where
 [2093] m is a number 2 or 3,
 [2094] ##¹ marks the linkage site with the group G,
 [2095] ##² marks the linkage site with the group B,
 [2096] where (C₃-C₅)-alkanediyl may be substituted by
 1 or 2 methyl substituents,
 [2097] B is a bond or a group of the formula



- [2098] where
 [2099] * marks the linkage site with L¹,
 [2100] ** marks the linkage site with L²,
 [2101] L³ is a bond,
 [2102] L⁴ is a bond
 [2103] R¹⁶ is hydrogen,
 [2104] R¹⁷ is hydrogen,
 [2105] L² is linear (C₃-C₆)-alkanediyl or is a group of the
 formula




- [2106] where
 [2107] p is a number 2 or 3,
 [2108] ##³ marks the linkage site with the group B,
 [2109] ##⁴ marks the linkage site with the nitrogen atom,
 [2110] n, AK, Cys, G, D and R³⁵ have the definitions indi-
 cated above,
 and also their salts, solvates and solvates of the salts.
 [2111] Preferred in the context of the present invention are
 also compounds of the formula (Ia), (XXXa) and (XXXI), in
 which
 [2112] L¹ is a bond,
 [2113] B is a bond,
 [2114] L² is linear (C₃-C₆)-alkanediyl or is a group of the
 formula



- [2115] where
 [2116] p is a number 2 or 3,
 [2117] ##³ marks the linkage site with the group B,
 [2118] ##⁴ marks the linkage site with the nitrogen atom,
 [2119] n, AK, Cys, G, D and R³⁵ have the definitions indi-
 cated above,
 and also their salts, solvates and solvates of the salts.
 [2120] Preferred in the context of the present invention are
 also compounds of the formula (Ia), (XXXa) and (XXXI), in
 which

[2121] L^1 is linear (C_3-C_5) -alkanediyl or a group of the formula



[2122] where


[2123] m is a number 2 or 3,

[2124] $##^1$ marks the linkage site with the group G ,

[2125] $##^2$ marks the linkage site with the group B ,

[2126] where (C_3-C_5) -alkanediyl may be substituted by 1 or 2 methyl substituents,

[2127] B is a group of the formula



[2128] where

[2129] $*$ marks the linkage site with L^1 ,

[2130] $**$ marks the linkage site with L^2 ,


[2131] L^3 is a bond,

[2132] L^4 is a bond,

[2133] R^{16} is hydrogen,

[2134] R^{17} is hydrogen,

[2135] L^2 is linear (C_3-C_6) -alkanediyl or is a group of the formula



[2136] where

[2137] p is a number 2 or 3,

[2138] $##^3$ marks the linkage site with the group B ,

[2139] $##^4$ marks the linkage site with the nitrogen atom,

[2140] n , AK , Cys , G , D and R^{35} have the definitions indicated above,

and also their salts, solvates and solvates of the salts.

[2141] Preferred in the context of the present invention are also compounds of the formula (Ia), in which

[2142] n is a number from 2 to 8, preferably 2 to 5,


[2143] AK is AK_1 or AK_2

[2144] where

[2145] AK_1 is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G ,

[2146] AK_2 is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the NH side group of a lysine residue of the binder to the group G ,

[2147] G when $AK=AK_1$, is a group of the formula



[2148] in which


[2149] $#^1$ marks the linkage site with the cysteine residue of the binder,

[2150] $#^2$ marks the linkage site with the group L^1 ,

[2151] or

[2152] when $AK=AK_2$, is carbonyl,

[2153] L^1 is a bond, linear (C_3-C_5) -alkanediyl or a group of the formula



[2154] where


[2155] m is a number 2 or 3,

[2156] $##^1$ marks the linkage site with the group G ,

[2157] $##^2$ marks the linkage site with the group B ,

[2158] where (C_3-C_5) -alkanediyl may be substituted by 1 or 2 methyl substituents,

[2159] B is a bond or a group of the formula



[2160] where

[2161] $*$ marks the linkage site with L^1 ,

[2162] $**$ marks the linkage site with L^2 ,


[2163] L^3 is a bond,

[2164] L^4 is a bond,

[2165] R^{16} is hydrogen,

[2166] R^{17} is hydrogen,

[2167] L^2 is linear (C_3-C_6) -alkanediyl or is a group of the formula



[2168] where

[2169] p is a number 2 or 3,

[2170] $##^3$ marks the linkage site with the group B ,

[2171] $##^4$ marks the linkage site with the nitrogen atom, and

[2172] D and R^{35} have the definitions indicated above, and also their salts, solvates and solvates of the salts.

[2173] Preferred in the context of the present invention are also compounds of the formula (Ia), in which

[2174] n is a number from 2 to 8, preferably 2 to 5,


[2175] AK is AK_1 or AK_2 ,

[2176] where

[2177] AK_1 is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

[2178] AK_2 is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the NH side group of a lysine residue of the binder to the group G,

[2179] G when $AK=AK_1$, is a group of the formula



[2180] in which

[2181] #1 marks the linkage site with the cysteine residue of the binder,

[2182] #2 marks the linkage site with the group L^1 ,


[2183] or

[2184] when $AK=AK_2$, is carbonyl,

[2185] L^1 is a bond,

[2186] B is a bond,

[2187] L^2 is linear (C_3-C_6)-alkanediyl or is a group of the formula



[2188] where

[2189] p is a number 2 or 3,

[2190] ##3 marks the linkage site with the group B,

[2191] ##4 marks the linkage site with the nitrogen atom,

[2192] D and R^{35} have the definitions indicated above, and also their salts, solvates and solvates of the salts.

[2193] Preferred in the context of the present invention are also compounds of the formula (Ia), in which


[2194] n is a number from 2 to 8, preferably 2 to 5,

[2195] AK is AK_1 ,

[2196] where

[2197] AK_1 is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

[2198] G is a group of the formula




[2199] where

[2200] #1 marks the linkage site with the cysteine residue of the binder,

[2201] #2 marks the linkage site with the group L^1 ,

[2202] L^1 is linear (C_3-C_5)-alkanediyl or a group of the formula



[2203] where


[2204] m is a number 2 or 3,

[2205] ##1 marks the linkage site with the group G,

[2206] ##2 marks the linkage site with the group B,

[2207] where (C_3-C_5)-alkanediyl may be substituted by 1 or 2 methyl substituents,

[2208] B is a group of the formula



[2209] where

[2210] * marks the linkage site with L^1 ,

[2211] ** marks the linkage site with L^2 ,


[2212] L^3 is a bond,

[2213] L^4 is a bond,

[2214] R^{16} is hydrogen,

[2215] R^{17} is hydrogen,

[2216] L^2 is linear (C_3-C_6)-alkanediyl or is a group of the formula



[2217] where

[2218] p is a number 2 or 3,


[2219] ##3 marks the linkage site with the group B,

[2220] ##4 marks the linkage site with the nitrogen atom,

[2221] D and R^{35} have the definitions indicated above, and also their salts, solvates and solvates of the salts.

[2222] Preferred in the context of the present invention are also compounds of the formula (Ia), in which

[2223] L^1 is a bond, linear (C_3-C_5)-alkanediyl, a group of the formula



[2224] where


[2225] m is a number 2 or 3,

[2226] ##1 marks the linkage site with the group G,

[2227] ##2 marks the linkage site with the group B,

[2228] L^{1A} is linear (C_3-C_6)-alkanediyl,

[2229] B¹ is a group of the formula




[2230] in which

[2231] ##5 marks the linkage site with the group L^{1A},

[2232] ##6 marks the linkage site with the group L^{1B},

[2233] L⁵ is a bond or ethane-1,2-diyl,

[2234] L⁶ is a bond or a group of the formula



[2235] in which

[2236] ##7 marks the linkage site with the carbonyl group,

[2237] ##8 marks the linkage site with L^{1B},

[2238] R³³ is hydrogen, (C₁-C₄)-alkylcarbonyl or tert-butyloxycarbonyl,

[2239] R³⁴ is hydrogen or methyl,

[2240] R²⁹ is hydrogen or methyl,


[2241] R³⁰ is hydrogen or methyl,

[2242] R³¹ is hydrogen or methyl,


[2243] R³² is hydrogen or methyl,

[2244] L^{1B} is linear (C₃-C₆)-alkanediyl,

[2245] B is a bond or a group of the formula



-continued



[2246] where


[2247] * marks the linkage site with L¹,

[2248] ** marks the linkage site with L²,

[2249] P is O,

[2250] L³ is a bond or ethane-1,2-diyl,

[2251] L⁴ is a group of the formula



[2252] in which

[2253] *** marks the linkage site with the carbonyl group,

[2254] **** marks the linkage site with L²,

[2255] R²⁵ is hydrogen or methyl,

[2256] R²⁸ is hydrogen, (C₁-C₄)-alkylcarbonyl or tert-butyloxycarbonyl,

[2257] Q¹ is a 4- to 7-membered heterocycle,

[2258] Q² is a 3- to 7-membered carbocycle or a 4- to 7-membered heterocycle,

[2259] R¹⁶ is hydrogen or methyl,

[2260] R¹⁷ is hydrogen or methyl,

[2261] R²³ is (C₁-C₄)-alkyl,

[2262] R²⁴ is hydrogen or (C₁-C₄)-alkyl,


[2263] R³⁶ is hydrogen, (C₁-C₄)-alkylcarbonyl or tert-butyloxycarbonyl,


[2264] R³⁷ is hydrogen or methyl,

[2265] or

[2266] R³⁶ and R³⁷ together with the atoms to which they are bonded form a pyrrolidine ring,

[2267] L² is linear (C₂-C₆)-alkanediyl or is a group of the formula






[2309] in which

[2310] #4 marks the linkage site with the adjacent nitrogen atom,

[2311] #5 marks the linkage site with the carbonyl group,

[2312] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[2313] in which

[2314] #6 marks the linkage site with the carbonyl group,


[2315] R⁶ is hydrogen, hydroxy or benzyloxy,

[2316] R³ is hydrogen or methyl,

[2317] R⁴ is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

[2318] or

[2319] R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



[2320] in which

[2321] #7 marks the linkage site with the adjacent nitrogen atom,

[2322] #8 marks the linkage site with the group T¹,

[2323] T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹,

[2324] in which

[2325] R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

[2326] R⁸ is hydrogen or methyl,

[2327] R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,


[2328] or

[2329] R⁸ and R⁹ together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

[2330] R¹⁰ is benzoyl,

[2331] R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl

[2332] R⁵ is hydrogen, methyl or a group of the formula



[2333] in which

[2334] #9 marks the linkage site with —CHC(R²⁶)-T²,

[2335] R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

[2336] R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,

[2337] R²⁶ is hydrogen,

[2338] T² is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,


and

[2339] n, AK, Cys, G, L¹, B, L², D and R³⁵ have the definitions indicated above,


and also their salts, solvates and solvates of the salts.

[2340] Preferred in the context of the present invention are also compounds of the formula (Ia), (XXXa) and (XXXI), in which


[2341] D is a group of the formula




- [2342] where
 [2343] #³ marks the linkage site with the nitrogen atom,
 [2344] R¹ is hydrogen,
 [2345] R² is benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,
 [2346] the ring A with the N—O moiety present therein is a heterocycle of formula



- [2347] in which
 [2348] #⁶ marks the linkage site with the carbonyl group,
 [2349] R³ is hydrogen,
 [2350] R⁴ is benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,
 [2351] T¹ is a group of the formula —C(=O)—OR⁷ or —C(=O)—NR⁸R⁹,
 [2352] in which
 [2353] R⁷ is hydrogen,
 [2354] R⁸ is hydrogen,
 [2355] R⁹ is hydrogen,
 [2356] n, AK, Cys, G, L¹, B, L², D and R³⁵ have the definitions indicated above,
 and also their salts, solvates and solvates of the salts.
 [2357] Preferred in the context of the present invention are also compounds of the formula (Ia), (XXXa) and (XXXI), in which
 [2358] D is a group of the formula




- [2359] where
 [2360] #³ marks the linkage site with the nitrogen atom,
 [2361] R¹ is hydrogen,
 [2362] R² is benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,
 [2363] the ring A with the N—O moiety present therein is a heterocycle of the formula




- [2364] in which
 [2365] #⁶ marks the linkage site with the carbonyl group,


- [2366] R³ is hydrogen,
 [2367] R⁴ is benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,
 [2368] T¹ is a group of the formula —C(=O)—NR⁸R⁹,
 [2369] in which
 [2370] R⁸ is hydrogen,
 [2371] R⁹ is hydrogen,
 [2372] n, AK, Cys, G, L¹, B, L², D and R³⁵ have the definitions indicated above,
 and also their salts, solvates and solvates of the salts.
 [2373] Preferred in the context of the present invention are also compounds of the formula (Ia), (XXXa) and (XXXI), in which
 [2374] D is a group of the formula



- [2375] where
 [2376] #³ marks the linkage site with the nitrogen atom,
 [2377] R¹ is hydrogen,
 [2378] R² is 4-hydroxybenzyl or 1H-indol-3-ylmethyl,
 [2379] the ring A with the N—O moiety present therein is a heterocycle of the formula



- [2380] in which
 [2381] #⁶ marks the linkage site with the carbonyl group,
 [2382] R³ is hydrogen,
 [2383] R⁴ is 4-hydroxybenzyl or 1H-indol-3-ylmethyl,
 [2384] T¹ is a group of the formula —C(=O)—NR⁸R⁹,
 [2385] in which
 [2386] R⁸ is hydrogen,
 [2387] R⁹ is hydrogen,
 [2388] n, AK, Cys, G, L¹, B, L², D and R³⁵ have the definitions indicated above,
 and also their salts, solvates and solvates of the salts.
 [2389] Preferred in the context of the present invention are also compounds of the formula (Ia), (XXXa) and (XXXI), in which
 [2390] D is a group of the formula



- [2391] where
 [2392] #³ marks the linkage site with the nitrogen atom,
 [2393] R³ is hydrogen,
 [2394] R⁴ is 4-hydroxybenzyl or 1H-indol-3-ylmethyl,

[2395] T^1 is a group of the formula $-C(=O)-OR^7$ or $-C(=O)-NR^8R^9$,

[2396] in which

[2397] R^7 is hydrogen,

[2398] R^8 is hydrogen,


[2399] R^9 is hydrogen,

[2400] n, AK, Cys, G, L^1 , B, L^2 , D and R^{35} have the definitions indicated above,

and also their salts, solvates and solvates of the salts.

[2401] Preferred in the context of the present invention are also compounds of the formula (Ia), (XXXa) and (XXXI), in which

[2402] D is a group of the formula



[2403] where

[2404] #³ marks the linkage site with the nitrogen atom,

[2405] R^3 is hydrogen,

[2406] R^4 is 4-hydroxybenzyl or 1H-indol-3-ylmethyl,

[2407] T^1 is a group of the formula $-C(=O)-NR^8R^9$,

[2408] in which

[2409] R^8 is hydrogen,


[2410] R^9 is hydrogen,

[2411] n, AK, Cys, G, L^1 , B, L^2 , D and R^{35} have the definitions indicated above,

and also their salts, solvates and solvates of the salts.

[2412] Preferred in the context of the present invention are also compounds of the formula (Ia), (XXXa) and (XXXI), in which

[2413] D is a group of the formula



[2414] where


[2415] #³ marks the linkage site with the nitrogen atom,

[2416] R^1 is hydrogen or methyl,

[2417] R^2 is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

[2418] or

[2419] R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[2420] in which

[2421] #⁴ marks the linkage site with the adjacent nitrogen atom,

[2422] #⁵ marks the linkage site with the carbonyl group,

[2423] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[2424] in which

[2425] #⁶ marks the linkage site with the carbonyl group,

[2426] R^6 is hydrogen, hydroxy or benzyloxy,


and

[2427] n, AK, Cys, G, L^1 , B, L^2 and R^{35} have the definitions indicated above,

and also their salts, solvates and solvates of the salts.

[2428] Preferred in the context of the present invention are also compounds of the formula (Ia), (XXXa) and (XXXI), in which

[2429] D is a group of the formula



[2430] where


[2431] #³ marks the linkage site with the nitrogen atom,

[2432] R^1 is hydrogen,

[2433] R^2 is benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

[2434] or

[2435] R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[2436] in which

[2437] #4 marks the linkage site with the adjacent nitrogen atom,

[2438] #5 marks the linkage site with the carbonyl group,

[2439] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[2440] in which

[2441] #6 marks the linkage site with the carbonyl group,

and

[2442] n, AK, Cys, G, L^1 , B, L^2 and R^{35} have the definitions indicated above,

and also their salts, solvates and solvates of the salts.

[2443] Preferred in the context of the present invention are also compounds of the formula (Ia), (XXXa) and (XXXI), in which

[2444] R^{35} is hydroxyl,

and

[2445] n, AK, Cys, G, L^1 , B, L^2 , D and R^{35} have the definitions indicated above,

and also their salts, solvates and solvates of the salts.

[2446] Preferred in the context of the present invention are also compounds of the formula (Ia), (XXXa) and (XXXI), in which

[2447] R^{35} is methyl,

and

[2448] n, AK, Cys, G, L^1 , B, L^2 , D and R^{35} have the definitions indicated above,

and also their salts, solvates and solvates of the salts.


[2449] Particularly preferred in the context of the present invention are, furthermore, also compounds of the formula (XXXa), in which

[2450] Cys is an L-cysteine residue which is bonded via the sulphur atom of the side chain via a carbon atom of the succinimide,

and also their salts, solvates and solvates of the salts.

[2451] Preferred in the context of the present invention are also compounds of the formula (I) and (XXX), in which

[2452] D is a group of the formula



[2453] where


[2454] #3 marks the linkage site with the nitrogen atom,

[2455] R^1 is hydrogen,

[2456] R^2 is benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

[2457] or

[2458] R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[2459] in which

[2460] #4 marks the linkage site with the adjacent nitrogen atom,

[2461] #5 marks the linkage site with the carbonyl group,

[2462] the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



[2463] in which

[2464] #6 marks the linkage site with the carbonyl group,


[2465] R^6 is hydrogen, hydroxy or benzyloxy,

[2466] n, AK, Cys, G, L^1 , L^2 and B have the definitions indicated above,

and also their salts, solvates and solvates of the salts.

[2467] Particularly preferred in the context of the present invention are also compounds of the formula (I) and (XXX), in which

[2468] D is a group of the formula



[2469] where


[2470] #3 marks the linkage site with the nitrogen atom,

[2471] R¹ is hydrogen,

[2472] R² is benzyl or 1H-indol-3-ylmethyl,

[2473] or

[2474] R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




[2475] in which

[2476] #4 marks the linkage site with the adjacent nitrogen atom,

[2477] #5 marks the linkage site with the carbonyl group,

[2478] the ring A with the N—O moiety present therein is a heterocycle of the formula



[2479] in which

[2480] #6 marks the linkage site with the carbonyl group,

[2481] n, AK, Cys, G, L¹, L² and B have the definitions indicated above,


and also their salts, solvates and solvates of the salts.

[2482] Preferred in the context of the present invention are also compounds of the formula (I), in which

[2483] n=1-20, more preferably n=1-10 and very preferably n=2-8.

[2484] Preferred in the context of the present invention are also compounds of the formula (I) and (XXX), in which

[2485] B is a bond or a group of the formula



[2486] where

[2487] * marks the linkage site with L¹,

[2488] ** marks the linkage site with L²,


[2489] L³ is a bond or ethane-1,2-diyl,

[2490] L⁴ is a bond,

[2491] n, AK, Cys, G, L¹, L², D, R¹⁶ and R¹⁷ have the definitions indicated above, and also their salts, solvates and solvates of the salts.

[2492] Particularly preferred in the context of the present invention are also compounds of the formula (I) and (XXX), in which

[2493] B is a bond or a group of the formula



[2494] where

[2495] * marks the linkage site with L¹,

[2496] ** marks the linkage site with L²,

[2497] L³ and L⁴ is a bond,

[2498] n, AK, Cys, G, L¹, L², D, R¹⁶ and R¹⁷ have the definitions indicated above, and also their salts, solvates and solvates of the salts.


[2499] Preferred in the context of the present invention are also binder-drug conjugates of the general formula (I), in which

[2500] AK is AK₁,

[2501] where

[2502] AK₁ is a binder which is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

[2503] G is a group of the formula




[2504] where

[2505] #1 marks the linkage site with the cysteine residue of the binder,


[2506] #2 marks the linkage site with the group L¹,

[2507] L¹ is a bond, linear (C₁-C₁₀)-alkanediyl or is a group of the formula




- [2508] where
 [2509] m is a number from 2 to 6,
 [2510] ##¹ marks the linkage site with the group G,
 [2511] ##² marks the linkage site with the group B,
 [2512] where (C₁-C₁₀)-alkanediyl may be substituted by 1 to 4 methyl substituents,
 [2513] and
 [2514] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,


[2515] B is a bond or a group of the formula



- [2516] where
 [2517] * marks the linkage site with L¹,
 [2518] ** marks the linkage site with L²,
 [2519] L³ is a bond or (C₂-C₄)-alkanediyl,
 [2520] L⁴ is a bond or a group of the formula



- [2521] in which
 [2522] *** marks the linkage site with the carbonyl group,
 [2523] **** marks the linkage site with L²,
 [2524] R²⁵ is hydrogen or methyl,
 [2525] Q¹ is a 4- to 7-membered heterocycle,
 [2526] R¹⁴ is hydrogen or (C₁-C₄)-alkyl,
 [2527] R¹⁵ is hydrogen or (C₁-C₄)-alkyl,
 [2528] or
 [2529] R¹⁴ and R¹⁵ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 [2530] R¹⁶ is hydrogen or (C₁-C₄)-alkyl,
 [2531] R¹⁷ is hydrogen or (C₁-C₄)-alkyl,
 [2532] or
 [2533] R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 [2534] L² is linear (C₂-C₁₀)-alkanediyl or is a group of the formula




- [2535] where
 [2536] p is a number from 2 to 6,
 [2537] ##³ marks the linkage site with the group B,
 [2538] ##⁴ marks the linkage site with the nitrogen atom,
 [2539] where (C₂-C₁₀)-alkanediyl may be substituted by 1 to 4 methyl substituents,
 [2540] and
 [2541] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,


and also their salts, solvates and solvates of the salts.

[2542] Preferred in the context of the present invention are also binder-drug conjugates of the general formula (I), in which

- [2543] AK is AK₂,
 [2544] where
 [2545] AK₂ is a binder which is bonded via the NH side group of a lysine residue of the binder to the group G,
 [2546] G is carbonyl,
 [2547] L¹ is a bond, linear (C₁-C₁₀)-alkanediyl or is a group of the formula




- [2548] where
 [2549] m is a number from 2 to 6,
 [2550] ##¹ marks the linkage site with the group G,
 [2551] ##² marks the linkage site with the group B,
 [2552] where (C₁-C₁₀)-alkanediyl may be substituted by 1 to 4 methyl substituents,
 [2553] and
 [2554] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,
 [2555] B is a bond or a group of the formula




- [2556] where
 [2557] * marks the linkage site with L¹,
 [2558] ** marks the linkage site with L²,


- [2559] P is O or NH,
 [2560] Q² is a 3- to 7-membered carbocycle or a 4- to 7-membered heterocycle,
 [2561] R¹⁸ is hydrogen or (C₁-C₄)-alkyl,
 [2562] R¹⁹ is hydrogen or the side group of a natural α-amino acid or of its homologues or isomers,
 [2563] R²⁰ is hydrogen or (C₁-C₄)-alkyl,
 [2564] or
 [2565] R¹⁹ and R²⁰ together with the atoms to which they are bonded form a pyrrolidinyl ring,
 [2566] R²¹ is hydrogen or (C₁-C₄)-alkyl,
 [2567] R²² is hydrogen or (C₁-C₄)-alkyl,
 [2568] or
 [2569] R²¹ and R²² together with the atoms to which they are bonded form a 3- to 7-membered carbocycle,
 [2570] R²³ is (C₁-C₄)-alkyl,
 [2571] R²⁴ is hydrogen or (C₁-C₄)-alkyl,
 [2572] L² is linear (C₂-C₁₀)-alkanediyl or is a group of the formula




- [2573] where
 [2574] p is a number from 2 to 6,
 [2575] ##³ marks the linkage site with the group B,
 [2576] ##⁴ marks the linkage site with the nitrogen atom,
 [2577] where (C₂-C₁₀)-alkanediyl may be substituted by 1 to 4 methyl substituents,
 [2578] and
 [2579] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring, and also their salts, solvates and solvates of the salts.
 [2580] Particularly preferred in the context of the present invention are binder-drug conjugates of the formula (Ia), in which
 [2581] n is a number from 2 to 8,
 [2582] AK is AK₁ or AK₂,
 [2583] where
 [2584] AK₁ is a human or humanized antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,
 [2585] AK₂ is a human or humanized antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the NH side group of a lysine residue of the binder to the group G,
 [2586] G when AK=AK₁, is a group of the formula




- [2587] in which
 [2588] #¹ marks the linkage site with the cysteine residue of the binder,
 [2589] #² marks the linkage site with the group L¹,
 [2590] or
 [2591] when AK=AK₂, is carbonyl,
 [2592] L¹ is a bond, linear (C₂-C₆)-alkanediyl or is a group of the formula



- [2593] where
 [2594] m is a number 2 or 3,
 [2595] ##¹ marks the linkage site with the group G,
 [2596] ##² marks the linkage site with the group B,
 [2597] where (C₂-C₆)-alkanediyl may be substituted by 1 or 2 methyl substituents,
 [2598] B is a bond or a group of the formula




- [2599] where
 [2600] * marks the linkage site with L¹,
 [2601] ** marks the linkage site with L²,
 [2602] L³ is a bond or ethane-1,2-diyl,
 [2603] L⁴ is a bond or a group of the formula




- [2604] in which
 [2605] *** marks the linkage site with the carbonyl group,
 [2606] **** marks the linkage site with L²,
 [2607] R²⁵ is methyl,
 [2608] R²⁸ is hydrogen, methylcarbonyl or tert-butoxycarbonyl,


- [2609] Q^1 is piperidine-1,4-diyl,
 [2610] R^{16} is hydrogen or methyl,
 [2611] R^{17} is hydrogen or methyl,
 [2612] or
 [2613] R^{16} and R^{17} together with the atoms to which they are bonded form a piperazinyl ring,
 [2614] R^{21} is hydrogen or methyl,
 [2615] R^{22} is hydrogen or methyl,
 [2616] or
 [2617] R^{21} and R^{22} together with the atoms to which they are bonded form a cyclopropyl ring,
 [2618] R^{23} is methyl,
 [2619] R^{24} is hydrogen,
 [2620] L^2 is linear (C_2 - C_6)-alkanediyl or is a group of the formula



- [2621] where
 [2622] p is a number from 2 to 6,
 [2623] $##^3$ marks the linkage site with the group B,
 [2624] $##^4$ marks the linkage site with the nitrogen atom,
 [2625] D is a group of the formula



- [2626] where
 [2627] $##^3$ marks the linkage site with the nitrogen atom,
 [2628] R^1 is hydrogen,
 [2629] R^2 is 4-hydroxybenzyl or 1H-indol-3-ylmethyl,
 [2630] the ring A with the N—O moiety present therein is



- [2631] in which
 [2632] $##^6$ marks the linkage site with the carbonyl group,
 [2633] R^3 is hydrogen,
 [2634] R^4 is 4-hydroxybenzyl or 1H-indol-3-ylmethyl,
 [2635] T^1 is a group of the formula $-C(=O)-NR^8R^9$,
 [2636] R^8 is hydrogen or methyl,
 [2637] R^9 is hydrogen, methyl, or ethyl,
 [2638] R^{35} is methyl,
 and also their salts, solvates and solvates of the salts.

- [2639] Particularly preferred in the context of the present invention are binder-drug conjugates of the formula (Ia), in which

[2640] n is a number from 2 to 8, preferably from 2 to 5,


[2641] AK is AK_1 or AK_2 ,

[2642] where


[2643] AK_1 is a human or humanized antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

[2644] AK_2 is a human or humanized antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the NH side group of a lysine residue of the binder to the group G,


[2645] G when $AK=AK_1$ is a group of the formula




- [2646] where
 [2647] $##^1$ marks the linkage site with the cysteine residue of the binder,
 [2648] $##^2$ marks the linkage site with the group L^1 ,
 [2649] or
 [2650] when $AK=AK_2$, is carbonyl,
 [2651] L^1 is a bond,
 [2652] B is a bond
 [2653] R^{17} is hydrogen,
 [2654] L^2 is pentane-1,5-diyl, hexane-1,6-diyl or is a group of the formula




- [2655] where
 [2656] p is a number 2 or 3,
 [2657] $##^3$ marks the linkage site with the group B,
 [2658] $##^4$ marks the linkage site with the nitrogen atom,
 [2659] D is a group of the formula




- [2660] where
 [2661] $##^3$ marks the linkage site with the nitrogen atom,
 [2662] R^1 is hydrogen,
 [2663] R^2 is 4-hydroxybenzyl or 1H-indol-3-ylmethyl,
 [2664] the ring A with the N—O moiety present therein is



- [2665] in which
 [2666] #⁶ marks the linkage site with the carbonyl group,
 [2667] R³ is hydrogen,
 [2668] R⁴ is 4-hydroxybenzyl or 1H-indol-3-ylmethyl,
 [2669] T¹ is a group of the formula —C(=O)—NR⁸R⁹,
 [2670] R⁸ is hydrogen,
 [2671] R⁹ is hydrogen,
 [2672] R³⁵ is methyl,
 and also their salts, solvates and solvates of the salts.
 [2673] Preferred subject matter of the invention are binder-drug conjugates of the general formula (Ia), in which
 [2674] n is a number from 2 to 8, preferably from 2 to 5,
 [2675] AK is AK₁ or AK₂,
 [2676] where
 [2677] AK₁ is a human or humanized antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,
 [2678] AK₂ is a human or humanized antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the NH side group of a lysine residue of the binder to the group G,
 [2679] G when AK=AK₁, is a group of the formula




- [2680] where
 [2681] #¹ marks the linkage site with the cysteine residue of the binder,
 [2682] #² marks the linkage site with the group L¹,
 [2683] or
 [2684] when AK=AK₂, is carbonyl,
 [2685] L¹ is linear (C₂-C₆)-alkanediyl,
 [2686] B is a group of the formula




- [2687] where
 [2688] * marks the linkage site with L¹,
 [2689] ** marks the linkage site with L²,


- [2690] L³ is a bond,
 [2691] L⁴ is a group of the formula







- [2692] in which
 [2693] *** marks the linkage site with the carbonyl group,
 [2694] **** marks the linkage site with L²,
 [2695] R²⁵ is methyl,
 [2696] R²⁸ is hydrogen, methylcarbonyl or tert-butyloxy carbonyl,
 [2697] R¹⁶ is hydrogen,
 [2698] R¹⁷ is hydrogen,
 [2699] R³⁶ is hydrogen, methylcarbonyl or tert-butyloxy carbonyl,
 [2700] R³⁷ is methyl,
 [2701] L² is linear (C₂-C₄)-alkanediyl,
 [2702] D is a group of the formula




- [2703] where
 [2704] #³ marks the linkage site with the nitrogen atom,
 [2705] R¹ is hydrogen,
 [2706] R² is 4-hydroxybenzyl or 1H-indol-3-ylmethyl,
 [2707] the ring A with the N—O moiety present therein is



- [2708] in which
 [2709] #⁶ marks the linkage site with the carbonyl group,
 [2710] R³ is hydrogen,
 [2711] R⁴ is 4-hydroxybenzyl or 1H-indol-3-ylmethyl,
 [2712] T¹ is a group of the formula —C(=O)—NR⁸R⁹,
 [2713] R⁸ is hydrogen,
 [2714] R⁹ is hydrogen,
 [2715] R³⁵ is methyl or hydroxy,
 and also their salts, solvates and solvates of the salts.
 [2716] Particularly preferred in the context of the present invention are also drug-binder conjugates selected from the following compounds:




-continued




[2717] where in each case

[2718] n is a number from 2 to 8, preferably 2 to 5,[2719] AK_1 is a human or humanized antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,[2720] AK_2 is a human or humanized antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the NH side group of a lysine residue of the binder to the group G.

[2721] More particularly preferred in the context of the present invention are also binder-drug conjugates selected from the following compounds:



-continued



[2722] where in each case

[2723] *n* is a number from 2 to 8, preferably 2 to 5,

[2724] AK₁ is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the

sulphur atom of a cysteine residue of the binder to the group G,


[2725] AK₂ is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the NH side chain of a lysine residue of the binder to the group G.

[2726] The definitions of radicals that are indicated individually in the respective combinations and preferred combinations of radicals are also replaced arbitrarily by radical definitions of other combinations, independently of the respective combinations of radicals that are indicated.

[2727] Especially preferred are combinations of two or more of the abovementioned preference ranges.


[2728] Further provided by the invention is a process for preparing the compounds of the invention of the formula (Ia), characterized in that a solution of the binder in PBS buffer

[2729] [A] is admixed with a suitable reducing agent, such as, for example, dithiothreitol or tris(2-carboxyethyl)phosphine hydrochloride, and is subsequently reacted with a compound of the formula (IIa)




[2730] in which D, L¹, B, L² and R³⁵ each have the definitions indicated above,

[2731] to give a compound of the formula (Ia-A)




[2732] in which n, AK₂, D, L¹, B, L² and R³⁵ each have the definitions indicated above, or

[2733] [B] is reacted with a compound of formula (IIIa)



[2734] in which D, L¹, B, L² and R³⁵ each have the definitions indicated above,

[2735] to give a compound of the formula (Ia-B)




[2736] in which n , AK_2 , D , L^1 , B , L^2 and R^{35} each have the definitions indicated above.

[2737] Further provided by the invention is a process for preparing the compounds of the invention of the formula (I), characterized in that a solution of the binder in PBS buffer

[2738] [A] is admixed with a suitable reducing agent, such as, for example, dithiothreitol or tris(2-carboxyethyl)phosphine hydrochloride, and is subsequently reacted with a compound of the formula (II)


(II)



[2739] in which D , L^1 , B and L^2 each have the definitions indicated above,

[2740] to give a compound of the formula (I-A)

(I-A)




[2741] in which n , AK_1 , D , L^1 , B and L^2 each have the definitions indicated above,

or

[2742] [B] is reacted with a compound of the formula (III)


(III)



[2743] in which D , L^1 , B and L^2 each have the definitions indicated above,

[2744] to give a compound of the formula (I-B)

(I-B)



[2745] in which n, AK₂, D, L¹, B and L² each have the definitions indicated above.

Cysteine Coupling:

[2746] The partial reduction of the antibody and also the subsequent conjugation of the (partially) reduced antibody with a compound of the formula (II) or (IIa) takes place in accordance with the methods known to the skilled person, see e.g. Ducry et. al., Bioconj. Chem. 2010, 21, 5 and references herein, Klussman et. al., Bioconj. Chem. 2004, 15(4), 765-773. The mild reduction of the antibody is accomplished preferably by addition of 2-6 equivalents of TCEP to the antibody, which is present in a suitable buffer solution, preferably phosphate buffer, and by stirring for 30-180 minutes at temperatures between 15 and 40° C., preferably at RT. This is followed by the conjugation, by addition of a solution of a compound of the formula (II) or (IIa) in DMSO, acetonitrile


or DMF to the solution of the (partially) reduced antibody in PBS buffer, and subsequent reaction at a temperature of 0° C. to +40° C., more particularly of +10° C. to +30° C., for a period of 30 minutes to 6 hours, more particularly 1 to 2 hours.

Lysine Coupling:

[2747] First of all the compounds of the formula (III) or (IIa) or comparable activated carboxyl components are prepared by conventional methods of peptide chemistry. They are then taken up in inert solvents such as DMSO or DMF, for example, and added to the antibody, which is preferably present in phosphate buffer at a neutral pH. The solution is stirred for 1-16 h at a temperature between 15 and 40° C., preferably RT.


[2748] The preparation processes described above are elucidated by way of example using the schemes below (Scheme 1 and 2):


Scheme 1



[a]: 1. AK (antibody), TCEP, PBS buffer, RT; 2. Addition of the maleimide derivative in DMSO, RT.


Scheme 2





[a): AK (antibody), PBS buffer, RT admix with activated carboxyl derivative of the linker-drug components].

[2749] The compounds of the formula (II) in which L¹ and B are a bond can be prepared by subjecting a compound of the formula (IV)




in which D has the definition indicated above,
to reductive amination in an inert solvent with a compound of
the formula (V)

in which

[2750] L^{2A} has the above-defined definition of L^2 , but is shortened by one carbon atom in the alkyl chain length,


[2751] PG¹ is an amino-protective group such as, for example, (9H-fluoren-9-ylmethoxy)carbonyl, tert-butoxycarbonyl or benzyloxycarbonyl,

[2752] to give a compound of the formula (VI)




in which D , L^2 and PG^1 have the definition indicated above,

[2753] eliminating the protective group PG¹ from this compound by methods known to the skilled person, and reacting the deprotected compound in an inert solvent in the presence of a suitable base with methyl 2,5-dioxo-2,5-dihydro-1H-pyrrole-1-carboxylate to give a compound of the formula (II-A)




in which D and L^2 each have the definitions indicated above.

[2754] The compounds of the formula (II) in which B is a group of the formula (B¹)

(B¹)

in which *, **, R¹⁴ and R¹⁵ each have the conditions indicated above,

can be prepared by eliminating the protective group PG¹ from a compound of the formula (VI) by methods known to the skilled person, and reacting the deprotected compound in an inert solvent in the presence of a suitable base with a compound of the formula (VII)




(VII)

in which L¹ has the definition indicated above, to give a compound of the formula (II-B)


in which D, L¹ and L² each have the definitions indicated above.

[2755] The compounds of the formula (II) in which B is a group of the formula (B²)

(B²)

in which *, **, L³, R¹⁶ and R¹⁷ each have the conditions indicated above can be prepared by subjecting a compound of the formula (IV)


to reductive amination in an inert solvent with a compound of the formula (VIII)



(VIII)


in which

[2756] L²⁴ has the above-defined definition of L², but is shortened by one carbon atom in the alkyl chain length,




(II-B)


to give a compound of the formula (IX)



in which D and L² have the definitions indicated above,
and reacting this compound in an inert solvent in the presence
of a suitable coupling reagent and a suitable base with a
compound of the formula (X)





in which L¹ and L³ each have the definitions indicated above,
to give a compound of the formula (II-C)



in which D, L¹, L² and L³ each have the definitions indicated
above.

[2758] L^{4A} is a group of the formula

[2757] Compound of the formula (II), in which B is a group
of the formula (B³)



[2759] in which


[2760] *** marks the linkage site with the carbonyl
group,

[2761] **** marks the linkage site with L²,


[2762] R²⁵ is hydrogen or methyl,

in which *, **, L³, R¹⁶ and R¹⁷ each have the conditions
indicated above and

can be prepared by reacting a compound of the formula (IX) in an inert solvent in the presence of a suitable base and a suitable coupling reagent with a compound of the formula (XI-A) or (XI-B)




(XI-A)




-continued

(XI-B)

in which R²⁵ and PG¹ each have the definitions indicated above and PG² is a suitable carboxyl-protective group, more particularly benzyl, to give a compound (XII-A) or (XII-B)




(XII-A)



(XII-B)


in which D, PG¹, PG² and L² have the definitions indicated above, eliminating the protective group PG² from this compound subsequently, by methods known to the skilled person, and reacting the deprotected compound in an inert solvent in the presence of a suitable coupling reagent and a suitable base with a compound of the formula (X), and finally, eliminating the protective group PG¹ from this compound, by methods known to the skilled person, to give a compound of the formula (II-D-A) or (II-D-B)



(II-D-A)

-continued



(II-D-B)



in which D, L^1 , L^2 and L^3 have the definitions indicated above.

[2763] Compound of the formula (II), in which B is a group of the formula (B⁴)

can be prepared by reacting a compound of the formula (IX) in an inert solvent in the presence of a suitable base and a suitable coupling reagent with a compound of the formula (XXI)

(B⁴)


(XXI)

in which *, ** each have the conditions indicated above and
[2764] Q^{1A} is an N-linked 4- to 7-membered heterocycle,


in which PG^1 and Q^{1A} each have the definitions indicated above,

to give a compound of the formula (XXII)

(XXII)



in which PG^1 , Q^{1A} , D and L^2 have the definitions indicated above,
eliminating the protective group PG^1 from this compound, by methods known to the skilled person, and subsequently reacting the deprotected compound in an inert solvent in the presence of a suitable coupling reagent and a suitable base with a compound of the formula (XXIII)



(XXIII)


in which L^1 has the definition indicated above,
to give a compound of the formula (II-D)

[2766] The compounds of the formula (III), in which L^1 is a bond and B is a group of the formula (B^{5A})

(B^{5A})

in which *, ** and P each have the definitions indicated above and


[2767] Q^{2A} is a 3- to 7-membered carbocycle,
can be prepared by reacting a compound of the formula (IX) in an inert solvent in the presence of a suitable coupling reagent and a suitable base with a compound of the formula (XIII)




(II-D)

in which Q^{1A} , D, L^1 and L^2 have the definitions indicated above.

[2765] The compounds of the formula (III), in which L^1 and B are a bond can be prepared by reacting a compound of the formula (IX) in an inert solvent in the presence of a suitable coupling reagent and a suitable base with N-hydroxysuccinimide to give a compound of the formula (III-A)




(XIII)




(III-A)

in which D and L^2 each have the definitions indicated above.

in which P, Q^{2A} and PG^2 each have the definitions indicated above,
to give a compound of the formula (XIV)




in which D, P, Q^{2A} , L^2 and PG^2 each have the definitions indicated above,
eliminating the protective group PG^2 from this compound by methods known to the skilled person, and subsequently reacting the deprotected compound in an inert solvent in the presence of a suitable base with N-hydroxysuccinimide to give a compound of the formula (III-B)




in which D, P, Q^{2A} and L^2 each have the definitions indicated above.

[2768] The compounds of the formula (III), in which L^1 is a bond and B is a group of the formula (B⁶)




in which *, **, R^{18} , R^{19} and R^{20} each have the definitions indicated above,

can be prepared by reacting a compound of the formula (IX) in an inert solvent in the presence of a suitable coupling reagent and a suitable base with a compound of the formula (XV)




in which R^{18} , R^{19} , R^{20} and PG^2 each have the definitions indicated above,
to give a compound of the formula (XVI)



in which D, R¹⁸, R¹⁹, R²⁰, L² and PG² each have the definitions indicated above,



eliminating the protective group PG² from this compound by methods known to the skilled person, and subsequently reacting the deprotected compound in an inert solvent in the presence of a suitable coupling reagent and a suitable base with N-hydroxysuccinimide to give a compound of the formula (III-C)



in which D, R¹⁸, R¹⁹, R²⁰ and L² each have the definitions indicated above.


[2769] The compounds of the formula (III), in which L¹ is a bond and B is a group of the formula (B⁷)

can be prepared by eliminating the protective group PG¹ from a compound of the formula (VI) by methods known to the skilled person, and reacting the resultant deprotected compound in an inert solvent in the presence of a suitable base with a compound of the formula (XVII)




in which *, **, R²¹ and R²² each have the definitions indicated above,

in which R²¹ and R²² each have the definitions indicated above,
to give a compound of the formula (III-D)




in which D , R^{21} , R^{22} and L^2 each have the definitions indicated above.

[2770] The compounds of the formula (III), in which B is a group of the formula (B⁸)




in which $*$, $**$, R^{23} and R^{24} each have the definitions indicated above,

can be prepared by reacting a compound of the formula (IX) in an inert solvent in the presence of a suitable coupling reagent and a suitable base with a compound of the formula (XVIII)




in which R^{23} , R^{24} and PG^1 each have the definitions indicated above, to give a compound of the formula (XIX)



in which D , R^{23} , R^{24} , L^2 and PG^1 each have the definitions indicated above.


eliminating the protective group PG¹ from this compound by methods known to the skilled person, and subsequently reacting the deprotected compound in an inert solvent in the pres-

ence of a suitable coupling reagent and a suitable base with a compound of the formula (XX)



in which

$L^{1,4}$ is linear (C_1 - C_{10})-alkanediyl or is a group of the formula



[2771] where

[2772] m is a number from 2 to 6,

[2773] ##¹ marks the linkage site with the group G,


[2774] ##² marks the linkage site with the group B,

[2775] where (C₁-C₁₀)-alkanediyl may be substituted by 1 to 4 methyl substituents,

[2776] and

[2777] where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring.

to give a compound of the formula (III-E)



in which D, R²³, R²⁴, L¹⁴ and L² each have the definitions indicated above.

[2778] The compounds of the formula (III), in which B is a group of the formula (B^{5B})



in which * and ** each have the definitions indicated above and


[2779] Q^{2B} is an N-linked 4- to 7-membered heterocycle,

can be prepared by reacting a compound of the formula (IX) in an inert solvent in the presence of a suitable base and a suitable coupling reagent with a compound of the formula (XXIV)




in which PG¹ and Q^{2B} each have the definitions indicated above,

to give a compound of the formula (XXV)



in which PG¹, Q^{2B}, D and L² have the definitions indicated above,

eliminating the protective group PG¹ from this compound by methods known to the skilled person, and subsequently converting the deprotected compound in an inert solvent in the presence of a suitable base with a compound of the formula (XX) into a compound of the formula (III-F)



in which Q^{2B} , D, L^{1A} and L^2 have the definitions indicated above.

[2780] The reactions (IV)+(V)→(VI) and (IV)+(VIII)→(IX) take place in the solvents which are customary for a reductive amination and are inert under the reaction conditions, optionally in the presence of an acid and/or of a water-removing agent as catalyst. Such solvents include, for example, alcohols such as methanol, ethanol, n-propanol, isopropanol, n-butanol or tert-butanol, ethers such as tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane or bis(2-methoxyethyl)ether, or other solvents such as dichloromethane, 1,2-dichloroethane, N,N-dimethylformamide or else water. It is also possible to use mixtures of these solvents. As solvent it is preferred to use a 1,4-dioxane/water mixture, with addition of acetic acid or dilute hydrochloric acid as catalyst.

[2781] Reducing agents suitable for this reaction are, in particular, complex borohydrides, such as, for example, sodium borohydride, sodium cyanoborohydride, sodium triacetoxyborohydride, tetra-n-butylammonium borohydride or borane-pyridine complex. It is preferred to use sodium cyanoborohydride or borane-pyridine complex.

[2782] The reactions (IV)+(V)→(VI) and (IV)+(VIII)→(IX) take place in general in a temperature range from 0° C. to +120° C., preferably at +50° C. to +100° C. The reactions may be carried out under atmospheric, increased or reduced pressure (e.g. from 0.5 to 5 bar); it is usual to operate at atmospheric pressure.

[2783] The above-described coupling reactions (IX)+(X)→(II-C), (XII-A) or (XII-B)+(X)→(II-D-A) or (II-D-B), (IX)+(XIII)→(XIV), (IX)+(XV)→(XVI) and (XXII)+(XXIII)→(II-D) (amide formation from amine component and carboxylic acid component respectively) are carried out by standard methods of peptide chemistry [see e.g. M. Bodanszky, *Principles of Peptide Synthesis*, Springer-Verlag, Berlin, 1993; M. Bodanszky and A. Bodanszky, *The Practice of Peptide Synthesis*, Springer-Verlag, Berlin, 1984; H.-D. Jakubke and H. Jeschkeit, *Aminosäuren, Peptide, Proteine*, Verlag Chemie, Weinheim, 1982].

[2784] Examples of inert solvents for these coupling reactions are ethers such as diethyl ether, diisopropyl ether, tert-butyl methyl ether, tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane or bis(2-methoxyethyl)ether, hydrocarbons such as benzene, toluene, xylene, pentane, hexane, heptane, cyclohexane or petroleum fractions, halogenated hydrocarbons such as dichloromethane, trichloromethane, tetrachloromethane, 1,2-dichloroethane, trichloroethylene or chlorobenzene, or dipolar-aprotic solvents such as acetone, methyl ethyl ketone, acetonitrile, ethyl acetate, pyridine, dimethyl sulfoxide (DMSO), N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), N,N'-dimethylpropyleneurea (DMPU) or N-methylpyrrolidinone (NMP). It is also possible to use mixtures of such solvents. Preference is given to using N,N-dimethylformamide.

[2785] Examples of suitable activating/condensing agents for these couplings include carbodiimides such as N,N'-diethyl-, N,N'-dipropyl-, N,N'-diisopropyl-, N,N'-dicyclohexylcarbodiimide (DCC) or N-(3-dimethylaminoisopropyl)-N'-ethylcarbodiimide hydrochloride (EDC), phosgene derivatives such as N,N'-carbonyldiimidazole (CDI) or isobutyl chloroformate, 1,2-oxazolium compounds such as 2-ethyl-5-phenyl-1,2-oxazolium 3-sulphate or 2-tert-butyl-5-methylisoxazolium perchlorate, acylamino compounds such as 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline, phosphorus compounds such as propanephosphonic anhydride, diethyl cyanophosphonate, bis(2-oxo-3-oxazolidinyl)phosphoryl chloride, benzotriazol-1-yloxytris(dimethylamino) phosphonium hexafluorophosphate or benzotriazol-1-yloxytris(pyrrolidino)phosphonium hexafluorophosphate

(PyBOP), or uronium compounds such as O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU), O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU), 2-(2-oxo-1-(2H)-pyridyl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TPTU), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) or O-(1H-6-chlorobenzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TCTU), optionally in combination with further auxiliaries such as 1-hydroxybenzotriazole (HOBt) or N-hydroxysuccinimide (HOSu), and also, as bases, alkali metal carbonates, e.g. sodium or potassium carbonate, or tertiary amine bases such as triethylamine, N-methylmorpholine, N-methylpiperidine, N,N-diisopropylethylamine, pyridine or 4-N,N-dimethylaminopyridine.

[2786] In the context of the present invention, as activating/condensing agents for such coupling reactions, it is preferred to use N-(3-dimethylaminoisopropyl)-N'-ethylcarbodiimide hydrochloride (EDC) in combination with 1-hydroxybenzotriazole (HOBt) and N,N-diisopropylethylamine, or O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) likewise in conjunction with N,N-diisopropylethylamine.

[2787] The coupling reactions (IX)+(X)→(II-C), (XII-A) or (XII-B)+(X)→(II-D-A) or (II-D-B), (IX)+(XIII)→(XIV), (IX)+(XV)→(XVI) and (XXII)+(XXIII)→(II-D) are carried out in general in a temperature range from -20° C. to +60° C., preferably at 0° C. to +40° C. The reactions may take place under atmospheric, at increased or at reduced pressure (e.g. from 0.5 to 5 bar); it is usual to operate under atmospheric pressure.

[2788] The esterifications (IX)+(XVIII)→(XII) and (IX)+(XI-A) or (XI-B)→(XII-A) or (XII-B), (IX)+(XXIV)→(XXV) and also (IX)+(XXI)→(XXII) take place in analogy to the above-described amide coupling reactions. These reactions take place preferably in dichloromethane, using N-(3-dimethylaminoisopropyl)-N'-ethylcarbodiimide hydrochloride (EDC) and 4-dimethylaminopyridine at a temperature of +50° C. to 100° C. under atmospheric pressure.

[2789] The functional groups optionally present in the compounds—such as amino, hydroxyl and carboxyl groups in particular—may also be present in a temporarily protected form during the above-described process steps, if useful or necessary. In these cases, such protective groups are introduced and removed in accordance with customary methods known from peptide chemistry [see, for example, T. W. Greene and P. G. M. Wuts, *Protective Groups in organic Synthesis*, Wiley, New York, 1999; M. Bodanszky and A. Bodanszky, *The Practice of Peptide Synthesis*, Springer-Verlag, Berlin, 1984]. Where two or more protected groups are present, they can be liberated again optionally simultaneously in a one-pot reaction, or else liberated again in separate reaction steps.

[2790] As an amino-protective group PG¹ it is preferred to use tert-butoxycarbonyl (Boc), benzyloxycarbonyl (Z) or (9H-fluoren-9-ylmethoxy)carbonyl (Fmoc); for a hydroxyl or carboxyl function it is preferred to use tert-butyl or benzyl as protective group PG². The elimination of a tert-butyl or tert-butoxycarbonyl group is typically accomplished by treatment with a strong acid, such as hydrogen chloride, hydrogen bromide or trifluoroacetic acid, in an inert solvent such as diethyl ether, 1,4-dioxane, dichloromethane or acetic acid; this reaction may optionally also be carried out without addition of an inert solvent. In the case of benzyl or benzyloxycarbonyl as protective group, this group is removed preferably by hydrogenolysis in the presence of a suitable palladium catalyst, such as palladium on activated carbon, for example. The

(9H-fluoren-9-ylmethoxy)carbonyl group is generally eliminated using a secondary amine base such as diethylamine or piperidine.

[2791] The reaction (VI)→(II-A) takes place in a solvent which is inert under the reaction conditions, such as, for example, ethers such as tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane or bis(2-methoxyethyl)ether, alcohols such as methanol, ethanol, isopropanol, n-butanol or tert-butanol, or dipolar-aprotic solvents such as acetone, methyl ethyl ketone, acetonitrile, ethyl acetate, pyridine, dimethyl sulphoxide (DMSO), N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), N,N'-dimethylpropyleneurea (DMPU) or N-methylpyrrolidinone (NMP) or water. It is also possible to use mixtures of such solvents. Preference is given to using a mixture of 1,4-dioxane and water.

[2792] Suitable bases for the reaction (VI)→(II-A) are, for example, alkali metal carbonates such as potassium carbonate, sodium carbonate or lithium carbonate, alkali metal hydrogencarbonates such as sodium or potassium hydrogen carbonate or alkali metal alkoxides such as sodium methoxide, sodium ethoxide or potassium tert-butoxide. It is preferred to use sodium hydrogencarbonate.

[2793] The reaction (VI)→(II-A) takes place in a temperature range from 0° C. to +50° C., preferably at +10° C. to +30° C. The reaction may take place under atmospheric, under elevated or under reduced pressure (e.g. from 0.5 to 5 bar); it is usual to operate under atmospheric pressure.

[2794] The reaction (VI)+(VII)→(II-B) takes place in a solvent which is inert under the reaction conditions, such as, for example, ethers such as tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane or bis(2-methoxyethyl)ether, alcohols such as methanol, ethanol, isopropanol, n-butanol or tert-butanol, or dipolar-aprotic solvents such as acetone, methyl ethyl ketone, acetonitrile, ethyl acetate, pyridine, dimethyl sulphoxide (DMSO), N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), N,N'-dimethylpropyleneurea (DMPU) or N-methylpyrrolidinone (NMP) or water. It is also possible to use mixtures of such solvents. Preference is given to using DMF.

[2795] Suitable bases for the reaction (VI)+(VII)→(II-B) are, for example, tertiary amine bases such as triethylamine, N-methylmorpholine, N-methylpiperidine, N,N-diisopropylethylamine, pyridine or 4-N,N-dimethylaminopyridine. Preference is given to using N,N-diisopropylethylamine.

[2796] The reaction (VI)+(VII)→(II-B) takes place in a temperature range from 0° C. to +50° C., preferably at +10° C. to +30° C. The reaction may take place under atmospheric, under elevated or under reduced pressure (e.g. from 0.5 to 5 bar); it is usual to operate under atmospheric pressure.

[2797] The reactions (IX)→(III-A), (XIV)→(III-B) and (XVI)→(III-C) and also (VI)+(XVII)→(III-D), (XIX)+(XX)→(III-E) and (XXV)+(XX)→(III-F) take place in a solvent which is inert under the reaction conditions. Examples of suitable solvents are ethers such as diethyl ether, diisopropyl ether, tert-butyl methyl ether, tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane or bis(2-methoxyethyl)ether, hydrocarbons such as benzene, toluene, xylene, pentane, hexane, heptane, cyclohexane or petroleum fractions, halogenated hydrocarbons such as dichloromethane, trichloromethane, tetrachloromethane, 1,2-dichloroethane, trichloroethylene or chlorobenzene, or dipolar-aprotic solvents such as acetone, methyl ethyl ketone, acetonitrile, ethyl acetate, pyridine, dimethyl sulphoxide (DMSO), N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), N,N'-dimethylpropyleneurea (DMPU) or N-methylpyrrolidinone (NMP). It is also possible to use mixtures of such solvents. Preference is given to using N,N-dimethylformamide.


[2798] Suitable bases for these reactions are, for example, tertiary amines such as triethylamine, N-methylmorpholine, N-methylpiperidine, N,N-diisopropylethylamine, pyridine or 4-N,N-dimethylaminopyridine. Preference is given to using N,N-diisopropylethylamine, optionally with addition of 4-N,N-dimethylaminopyridine.

[2799] The reactions (IX)→(III-A), (XIV)→(III-B) and (XVI)→(III-C) and also (VI)+(XVII)→(III-D) and (XIX)+(XX)→(III-E) take place in a temperature range from 0° C. to +50° C., preferably at +10° C. to +30° C. The reaction may take place under atmospheric, under elevated or under reduced pressure (e.g. from 0.5 to 5 bar); it is usual to operate under atmospheric pressure.


[2800] The compounds of the formulae (II), (III), (I-A) and (I-B) are sub-quantities of the compounds of the formulae (IIa), (IIIa), (Ia-A) and (Ia-B), respectively, where R³⁵ is methyl. The preparation of the compounds (IIa) and (IIIa) takes place in analogy to the preparation of the compound of the formulae (II) and (III) as described above.

[2801] The above-described processes are illustrated by way of example by the following synthesis schemes (Scheme 3 to 13, 18):

Scheme 3




-continued




[a): 1. Water/dioxane, 1 N HCl, 100° C.; 2. H₂, Pd/C, methanol, RT; b): NaHCO₃, H₂O, dioxane, RT].

Scheme 4




-continued



[a): Diisopropylethylamine, DMF, RT].


Scheme 5




CN1CCN(C1)C(=O)CCCN(C)C(=O)C[C@H](C)C(=O)N[C@@H](C)C(=O)N(C)C[C@H](C)C(=O)OCC[C@H](C)C(=O)N1CC[C@H](C1)C[C@H](C)C(=O)N[C@@H]2C[C@H](C2)C(=O)N3CCOCC3

[a): Water/dioxane, 1 N HCl, 100° C.; b): HATU, diisopropylamine, DMF, RT].

Scheme 6




-continued




[a): 1. EDCI, DMAP, dichloromethane, RT; 2. H₂, methanol, RT; b): 1. EDCI, HOBT, diisopropylamine, DMF, RT; 2. dichloromethane, RT].

Scheme 7




-continued




[a): 1. EDCI, DMAP, dichloromethane, RT; 2. H₂, methanol, RT; b): HATU, diisopropylamine, DMF, RT].

Scheme 8




[a): EDCI, dichloromethane, RT].

Scheme 9




[a): HATU, diisopropylethylamine, DMF, RT; 2. H₂, methanol, RT; b): EDCl, DMAP, dichloromethane, RT].

Scheme 10




[a): 1. HATU, diisopropylethylamine, DMF, RT; 2. dichloromethane, RT; b): EDCI, DMAP, dichloromethane, RT].

Scheme 11






[a): Diisopropylethylamine, DMF, RT].

Scheme 12




-continued




[a): 1. EDCI, DMAP, dichloromethane, RT; 2. diisopropylamine, DMAP, dichloromethane, RT].

Scheme 13




-continued




[a): DMAP diisopropylamine, dichloromethane, RT].

Scheme 18




-continued



[a]: 1. Water/dioxane, 1N HCl, 100° C.; 2. H₂, Pd/C, methanol, RT; b): HATU, diisopropylethylamine, RT].


[2802] The compounds of the formula (IV) can be prepared from commercially available amino acid building blocks or those known from the literature (see, for example, Pettit et al., *Synthesis* 1996, 719; Shioiri et al., *Tetrahedron Lett.* 1991, 32, 931; Shioiri et al., *Tetrahedron* 1993, 49, 1913; Koga et al., *Tetrahedron Lett.* 1991, 32, 2395; Vidal et al., *Tetrahedron* 2004, 60, 9715; Poncet et al., *Tetrahedron* 1994, 50, 5345. Pettit et al., *J. org. Chem.* 1994, 59, 1796) in analogy to processes known from the literature, in accordance with customary methods of peptide chemistry, and as described in the present experimental section. The synthesis schemes below (Scheme 14 to 16) illustrate the preparation by way of example.

-continued




[a): Hydroxylamine hydrochloride, KOH, MeOH, 0° C. → RT; b): BrCH₂(CH₂)₂CH₂Br, K₂CO₃, acetone, reflux].


Scheme 14




Scheme 15



-continued

[a): 1. Diisopropylethylamine, BEP, dichloromethane, -10°C . $\rightarrow$ RT; 2. MeOH].

Scheme 16



[a): 1. Diisopropylethylamine, BEP, DMF, RT; 2. dichloromethane; b): 1. HATU, diisopropylethylamine, DMF, RT; 2. dichloromethane, RT; c): 1. HATU, diisopropylethylamine, DMF, RT; 2. piperidine, DMF, RT].


[2803] The compounds of the formulae (XI), (XIII), (XV), (XVII) and (XXI), including, where appropriate, chiral or diastereomeric forms thereof, are available commercially or are described as such in the literature, or they can be prepared by routes that are obvious to the skilled person, in analogy to methods published in the literature. Numerous comprehensive instructions and also literature information on the preparation of the starting materials are also given in the experimental section, in the section relating to the preparation of the starting compounds and intermediates.

[2804] The compounds of the formulae (V), (VII), (VIII), (X), (XVIII), (XX) and (XXIII), including, where appropriate,


ate, chiral or diastereomeric forms thereof, are known from the literature, or can be prepared by routes which are obvious to the skilled person, in analogy to methods published in the literature. Numerous comprehensive instructions and also literature information on the preparation of the starting materials are also given in the experimental section, in the section relating to the preparation of the starting compounds and intermediates.

[2805] Alternatively, individual steps of the preparation sequence may be carried out in a different order. This approach is illustrated by way of example in the synthesis schemes below (Scheme 17, 19 and 20).

Scheme 17




-continued




[a): Borane-pyridine complex, acetic acid, MeOH; b): 1. HOBt, EDCl, diisopropylethylamine, DMF, RT; 2. TFA, dichloromethane, RT; c): HATU, diisopropylethylamine, DMF, RT; d) 1. Pd/C, MeOH, RT; 2. NaHCO₃, dioxane, water].

Scheme 19




-continued




[a): 1. HATU, diisopropylethylamine, DMF, RT; 2. TFA, dichloromethane, RT; 3. ((H₃C)₃C(C=O))₂O, DMF, diisopropylethylamine; b) diisopropylethylamine, BEP, DMF, RT; c) 1. H₂, Pd/C (10%) methanol, RT; 2. HATU, diisopropylethylamine, DMF, RT; 3. TFA, dichloromethane, RT].


Scheme 20



-continued




O




[a): Borane-pyridine complex, acetic acid, MeOH; b): 1. HOBt, EDCI, diisopropylethylamine, DMF, RT; 2. TFA, dichloromethane, RT; c): 1. H_2 , Pd/C, MeOH, RT; 2. NaHCO_3 , dioxane, water; d) HATU, diisopropylethylamine, DMF, RT;].

[2806] Further ADCs may be prepared by way of example in accordance with the following methods:


Scheme 21



Scheme 22



-continued



[a]: 1. EDCI, DMAP, dichloromethane, RT; 2. H₂, methanol, RT; b): HATU, diisopropylethylamine, DMF, RT;

[2807] In one embodiment the binder binds to a target molecule which is present on a cancer cell. In one preferred embodiment the binder binds to a cancer target molecule.

[2808] In another preferred embodiment the target molecule is a selective cancer target molecule.

[2809] In one particularly preferred embodiment the target molecule is a protein.

[2810] In one embodiment the target molecule is an extracellular target molecule. In one preferred embodiment the extracellular target molecule is a protein.

[2811] Cancer target molecules are known to the skilled person. Examples thereof are listed below.

[2812] Examples of cancer target molecules are as follows: (1) EGF receptor (NCBI reference sequence NP_005219.2) Sequence (1210 amino acids):

```
>gi|29725609|ref|NP_005219.2|epidermal growth factor receptor
isoform a precursor [Homo sapiens]
MRPSGTAGAALLALLAALCPASRALEEKVCQGTSNKLTQLGTFEDHFLSLQRMFN
NCEVVLGNLEITYVQRNYDLSFLKTIQEVAGYVLIALNTVERIPLNLQIIRGNMYYE
NSYALAVLSNYDANKTGLKELPMRNLQEILHGAVRFSNNPALCNVESIQWRDIVSSD
FLSNMSMDFQNLHLSGCKDPCPNWSCWGAGEENCQKLTIIICAQQCSGRCRGKS
PSDCCHNQCAAGCTGPRESDCLVCRKFRDEATCKDTCPLMLYNPTTYQMDVNPGE
KYSFGATCVKCKPRNYVVDHGSVCVRACGADSYEMEEDGVRKCKKCEGPRCKVCN
GIGIGEFKDSLSINATNIHKFNCTSIGDLHLPLVAFRGDSFTHTPPLDPQELDILKTVK
EITGFLLIQAWPENRTDLHAFENLEIIRGRTKQHGGFSLAVVSLNITSGLRSLKEISDG
DVIISGNKNLCYANTINWKKLFGTSGQKTKIISNRGENSKATGQVCHALCSPGCGW
```

-continued

GPEPRDCVSCRNVSRGREGVDKCNLLEGEPRFVENSECIQCHPECLPQAMNITCTGR
GPDNICQCAHYIDGPHCVKTCPAGVMGENTLVWKYADAGHVCHLCHPNCTYGCT
GPGLGECPTNGPKIPSIATGMVGALLLVVALGIGLFMRRHIVRKRTLRLRLQERE
 LVEPLTPSGEAPNQALLRIKETEFKKIKVLGSGAFGTVYKGLWIPEGEKVKIPVAIKE
 LREATSPKANKEILDEAYVMASVDNPHVCRLGLCLTSTVQLITQLMPFGCLLDYVRE
 HKDNIGSQYLLNWCQVIAKGMNYLEDRLVHRDLAARNVLVKTQHVKITDFGLA
 KLLGAEKEYHAEGGKVPIKWMALESI LHRITYTHQSDVWSYGVTVWELMTFGSKPY
 DGIPASEISSILEKGERLPQPPICTIDVYMIMVKCWMIDASRPKFRELII EFSKMARDP
 QRYLVIQGDERMHLPSPTDSNFYRALMDEEDMDDVDADEYLIPQQGFFSSPSTRT
 PLLSSLSATSNNTVACIDRNLQSCPIKEDSFLQRYSSDPTGALTEDSIDDTFLPVPEY
 INQSVPKRPAGSVQNPVYHNQPLNPAPSRDPHYQDPHSTAVGNPEYLN TVQPTCVNS
 TFDSPAHWAKGSHQISLDNPDYQQDFPKEAKPNGIFKGSTAENAEYLRVAPQSSE
 FIGA

[2813] The extracellular domain is marked by underlining.

(2) Mesothelin (SwissProt reference Q¹³⁴²¹-3)

Sequence (622 amino acids):

>sp|Q13421-3|MSLN_HUMAN Isoform 2 of Mesothelin OS = *Homo sapiens*
 GN = MSLN
 MALPTARPLLGSCTPALGSLFLFLFSLGWVQPSRTLGETGQEAAPLDGVLANPPNISS
 LSPRQLLGFPACAEVSGLSTERVRELAVALAQKNVKLSTEQLRCLAHRLSEPPEDLDALPL
 DLLLFLNPDAFSGPQACTRFFSRITKANVDLLPRGAPERQRLPAALACWGVRSLLSEA
 DVRLGGLACDLGRFVAESA EVLLPRLVSCPGPLDQDQQAARAALQGGGPPYGGPSTW
 SVSTM DALRGLLPVLGQPIIRSI PQGIVAAWRQRSSRDPSWRQPRTILRPRFRREVEKT
 ACPSGKKAREIDESLI FYKKWELEACVDAALLATQMDRVNAIPFTYEQLDVLKHKLD ELY
 PQGYPESVIQHLGYLFLKMSPEDIRKWNVTSL ETLKALLEVNKGHEMSPQVATLIDRFVK
 GRGQLDKDTLDTLTA FYPGYLCSLSP EELSSVPPSSIWAVRPQDLDTCDPRQLDVLYPKA
 RLAFQNMNGSEYFVKIQSFLGGAPTE DLKALSQQNVSM DLATFMKLRTDAVLPLTVAEVQ
 KLLGPHVEGLKAERHRPVRDWILRQRQDDLTGLGLQGGINPYLVLDLSMQEALSGT
 PCLLGPGPVLTVLALLLASTLA

[2814] Mesothelin is encoded by amino acids 296-598.

Amino acids 37-286 code for “megakaryocyte-potentiating factor”. Mesothelin is anchored in the cell membrane by a GPI anchor and is localized extracellularly.

(3) Carboanhydrase IX (SwissProt reference Q¹⁶⁷⁹⁰)

Sequence (459 amino acids):

>sp|Q16790|CAH9_HUMAN Carbonic anhydrase 9 OS = *Homo sapiens*
 GN = CA9 PE = 1 SV = 2
 MAPLCSPWLP LLIPAPAGLTVQLLSLLLLVPVHPQRLPRMQEDSPLGGGSSGEDDPL
GEEDLPSEEDSPREEDPPGEEDLPGEEDLPGEEDLPEVKPKSEEGSLKLEDLPTVEAPG
DPQEPQNNHRDKEGDDQSHWRYGGDPWPFRVSPACAGRFQSPVDIRPQLAAFCPALRPL
ELLGFQLPPLPELRLRNNGHSVQLTLPPGLEMALGPGREYRALQLHLHWGAAGRPGSEHT

-continued

VEGHRFP AEIHVVHLSTAFARVDEALGRPGGLAVLA AFLEEGPEENSAYEQLLSRLEEIA
EEGSETQVPGLDISALLPSDFSRYFQYEGSLTTPPCAQGV IWT VFNQTVMLS AKQLHTLS
DTLWGP GDSRLQLNFRATQPLNGRVIEASFPAGVDSSPRAAEPVQLNSCLAAGDILALVF
 GLLFAVTSVAFLVQMRRQHRRGTKGGVSYRPAEVAETGA

[2815] The extracellular domain is marked by underlining.
 (4) C4.4a (NCBI reference sequence NP_055215.2; Synonym LYPD3)
 Sequence (346 amino acids):

>gi|93004088|ref|NP_055215.2| ly6/PLAUR domain-containing
 protein 3 precursor [*Homo sapiens*]
 MDPARKAGAAQAMIWTAGWLLLLLLLRGGAQALECYSCVQKADDGCGSPNKMKTVC
APGVDVCTEAVGAVETIHGQFSLAVRGCGSLPGKNDRGLDLHGLLAFIQ LQQCAQ
DRCNAKLNLT SRALDPAGNESAYPPNGVECYSCVGLSREACQGTSPPVVSCYNASD
HVYKGCDFGNVTLTAANVTVSLPVRGCVQDEFCTRDGVTGP GFTLSGSCCQGSRCN
SDLRNKTYFSPRIPPLVRLPPPEPTTVASTTSVTTSAPVRPTSTTKPMPAPTSQTPTQ
GVEHEASRDEEPRLTGGAGHQDRSNSGOYPAKGGPQQPHNKGCVAPTAGLAALLL
 AVAAGVLL

[2816] The matured, extracellular domain is marked by
 underlining (SEQ ID NO:1).
 (5) CD52 (NCBI reference sequence NP_001794.2)

>gi|68342030|ref|NP_001794.2|CAMPATH-1 antigen precursor
 [*Homo sapiens*]
 MKRFLFLLLTISLLVMVQIQ TGLSGQNDTSQTSSPSASSNISGGIFLFFVANAIHLFCFS

(6) Her2 (NCBI reference sequence NP_004439.2)

>gi|54792096|ref|NP_004439.2|receptor tyrosine-protein kinase
 erbB-2 isoform a [*Homo sapiens*]
 MELAALCRWGLLLALLPPGAASTQVCTGTMKLRLPASPETHLDMRLHLYQGCQV
 VQGNLELTYP LPTNASLSFLQDIQEVQGYVLI AHNQVRQVPLQRLRIVRG TQLFEDNY
 ALAVLDNGDPLNNTTPVTGASPGGLRELQLRSLTEILKGGVLIQRNPQLCYQDTILW
 KDIFHKNNQLALT LIDTNRSRACHPCSPMCKGSRGWGESSEDCQSLTRTV CAGGCAR
 CKGPLPTDCCHEQCAAGCTGPKHSDCLACLHFNHSGICELHCPALVTYNTDTFESMP
 NPEGRYTFGASCVTACPYNYLSTDVGSCTLV CPLHNQEVTAEDGTQRCEKCSKPCAR
 VCYGLGMEHLREVRVTSANIQEFAGCKKIFGSLAFLPESFDGDPASNTAPLQPEQLQ
 VFETLEEITGYLYISAWPDSL PDL SVFQNLQVIRGRILHNGAYSLTLQGLGISWGLRLS
 LRELGSGLALIHNNTHLCFVHTVPWDQLFRNPHQALLHTANRPEDECVGEGLACHQ
 LCARGHCWGPGPTQCVNCSQFLRGQECVEECRVLQGLPREYVNARHCLPCHPECQP
 QNGSVTCFGPEADQCVACAHYKDP PFCVARCP SGVKPDLSYMPIWKFPDEEGACQP
 CPINCTHSCVDLDDKGCPAEQRASPLTSII SAVVGILLVVVLGVVFGILIKRRQQKIRK
 YTMRRLLQETELVEPLTPSGAMPNQAQMRI LKETELRKVKVLGSGAFGTVYKGIWIP
 DGENVKIPVAIKVIRENTSPKANKEILDEAYVMAGVGSPYVSRLLGICLTSTVQLVTQ
 LMPYGCLLDHVREN RGR LGSQDLLNWC MQIAKGMSYLEDVRLVHRDLAARNVLV

-continued

KSPNHVKITDFGLARLLDIDETEHADGGKVPIKWMALLESILRRRFTHQSDVWSYGV
TVWELMTFGAKPYDGIPAREIPDLLEKGERLPQPPICTIDVYMIMVKCWMIDSECRPR
FRELVSFEFSRMARDPQRFVVIQNEDLGPASPLDSTFYRSLLEDDDMGDLVDAEYLV
PQQGFFCPDPAPGAGGMVHHRHRSSTRSGGDLTLGLEPSEEEAPRSPLAPSEGAGS
DVFDDGLGMGAAGLQSLPTHDPSPQLQRYSEDPTVPLPSETDGYVAPLTCSPQPEYV
NQPDVRPQPPSPREGPLPAARPAGATLERPKTLPKGKNGVVKDVFAFGGAVENPEYL
TPQGGAAPQPHPPAFSPAFDNLYYWDQDPPERGAPPSTFKGTPTAENPEYLGLDVPV

(7) CD20 (NCBI reference sequence NP_068769.2)

>gi|23110987|ref|NP_068769.2| B-lymphocyte antigen CD20 [*Homo sapiens*]
MTTPRNSVNGTFPAEPMKGPAMQSGPKPLFRMSSLVGPTQSFMRKSTLGAQVI
MNGLFHIALGGLLMIPAGIYAPICVTWVYPLWGGIMYIISGSLAATEKNRKLKLVKG
KMIMNSLSLFAAISGMILSMDILNIKISHFLKMESLNFIRAHTPYINIYNCEPANPSEKN
SPSTQYCYSIQSLFLGILSVMLIFAFQELVIAGIVENEWKRTCSRPKSNIVLLSAEEKK
EQTEIEIKEEVVGLTETSSQPKNEEDIEIPIQEEEEETETNFPEPPQDQESSPIENDSSP

(8) The lymphocyte-activating antigen CD30 (SwissProt ID P28908)

>gi|68348711|ref|NP_001234.2| tumor necrosis factor receptor
superfamily member 8 isoform 1 precursor [*Homo sapiens*]
MRVLLAALGLLFLGALRAFPQDRPFEDTCHGNPSHYDKAVRRCYRCMPGLFPTQ
QCPQRPDTCRKQCEPDYYLDEADRCTACVTCRDDLVEKTPCAWNSSRVCECRPGM
FCSTSAVNSCARCFHVSVCAGMIVKFPGTAQKNTVCEPASPGVSPACASPENCKEPS
SGTIPQAKPTPVSPATSSASTMPVRGGTRLAQEAASKLTRAPDSPSSVGRPSSDPGLSP
TQPCPEGSGDCRKQCEPDYYLDEAGRCTACVTCRDDLVEKTPCAWNSSRTCECRP
GMICATSATNSRARCVYPICAAETVTKPDMAEKDTTFEAPPLGTQPD CNTPENG
EAPASTSPTQSLLVDSQASKTLPIPTSAVALSSTGKPVLDAGPVLFVWVILVVLVVVGS
SAFLCHRRACRKRIRQKLHLCPVQTSQPKLELVDSRPRSSSTQLRSGASVTEPVAE
ERGLMSQPLMETCHSVGAAYLESPLQDASPAGEPSSPRDLPEPRVSTEHTNKNIEKI
YIMKADTVIVGTVKAELPEGRGLAGPAEPELEEELEADHTPHYPEQETEPPLGSCSDV
MLSVEEEGKEDPLPTAASGK

(9) The lymphocyte adhesion molecule CD22 (SwissProt ID P20273)

>gi|157168355|ref|NP_001762.2| B-cell receptor CD22 isoform 1
precursor [*Homo sapiens*]
MHLLGPWLLLLVLEYLAFSDSSKWFEPETLYAWEGACVWIPCTYRALDGDLESFI
LFHNPEYNKNTSKFDGTRLYESTKDGVPSQKRVQFLGDKNKNCTLSIHPVHLNDS
GQLGLRMESKTEKWMERIHLNVSERPPPHIQLPPEIQESQEVTLTCLLNFSCYGYPIQ
LQWLLEGVPMRQAAVTSLSLTIKSVFTRSELKFPQWSHHGKIVTCQLQDADGKFLS
NDTVQLNVKHTPKLEIKVTPSDAIVREGDSVTMTCEVSSSNPEYTTVSWLKDGTSLK
KQNTFTLNLREVTKDQSGKYCCQVSNVGPGRSEEVFLQVQYAPPEPSTVQILHSPAV

-continued

EGSQVEFLCMSLANPLPTNYTWYHNGKEMQGRTEEKVHIPKILPWHAGTYS CVAEN
 ILGTGQRGPGAELDVQYPPKKVTTVIQNMPPIREGDVTLS CYNSSNPSVTRYEWK
 PHGAWEEPSLGLVKIQNVGWDNTTIACAACNSWC SWASPVALNVQYAPRDVRVKI
 KPLSEIHSNGNSVSLQCD FSSSHPKEVQFFWEKNGRLLGKESQLNFD SISPEDAGSYSC
 WVNNISIGQTASKAWTLEVL YAPRRLRVSMSPGDQVM EGKSATLTCESDANPPVSHY
 TWFDWNNQSLPYHSQKLRL EPVKVQHSGAYWCQGTNSVGKGRSPLSTLT VVYSPET
 IGRRVAVGLGSLAILILAICGLKLQRRWKRTQSQQGLQENSSGQSFFVRNKKVRRAP
 LSEGPSHLGCYNPMMEDGISYTTLRFP EMNIPRTGDAESSEMQRPPDCDDTVTYSAL
 HKRQVG DYENVIPDFPEDEGIHYSELIQFGVGERPQAQENV DYVILKH

(10) The myeloid cell surface antigen CD33 (SwissProt ID P20138)

>gi|130979981|ref|NP_001763.3| myeloid cell surface antigen
 CD33 isoform 1 precursor [*Homo sapiens*]
 MPLLLLLPLLLWAGALAMPNFWLQVQESVTVQEGLCVLVPCTFFHPIPYDKNSPV
 HGYWFREGAII SRDSPVATNKLDQEVQEETQGRFRL LGDPSRNNCSLSIVDARRRDN
 GSYFFRMERGSTKYSYKSPQLSVHVTDLTHRPKILIPGTLEPGH SKNLTCSVSWACEQ
 GTPPIFSWLSAAPTSLGPRTHSSVLIITPRPDHGTNLTCQVKFAGAGVT TERTIQLN
 VTYVPQNPTTGIFPGDGS GKQETRAGVVHGAIGGAGVTALLALCLCLIFFIVKTHRRK
 AARTAVGRNDTHPTTGSASPKHQKSKLHGPTETSSCSGAAPT VEMDEELHYASLNF
 HGMNPSKDTSTEYSEVRTQ

(11) The transmembrane glycoprotein NMB (SwissProt ID Q¹⁴⁹⁵⁶)

>gi|52694752|ref|NP_001005340.1| transmembrane glycoprotein NMB
 isoform a precursor [*Homo sapiens*]
 MECLYYFLGFLLLAARLPLDAAKRFHDVLGNERPSAYMREHNQLNGWSSDENDWN
 EKLYPVWKRGD MRWKNWKGGRVQAVLTSDSPALVGSNITFAVNLI FPRCQKEDA
 NGNIVYEKNCRNEAGLSADPYVYNWTAWSESDGENG TGQSHNVFPDGKPPHH
 PGWRRWNFIYVFHTLGQYFQKLGRCSVRVSVNTANVT LGPQLMEVTVYRRHGRAY
 VPIAQVKDVVYVTDQIPVFVTMFQKNDRNSSDETFLKDLPI MFDVLIHDP SHFLNYST
 INYKWSFGDNTGLFVSTNHTVNH TYVLNGT FSLNLT VKAAAPGPCPPPPPPRPSKPT
 PSLATTLSKSYDSNTPGPAGDNPLELSRIPDENCQINRYGHFQATITIV EGILEVNI IQMT
 DVLMPVPWPESSLIDFVVT CQGSIPTEVCTIISDPTCEITQNTVCSPVDVDEMCLLTVR
 RTFNGSGTYCVNLT LGDDTSLALTSTLISVPDRDPASPLRMANSALISVGCLAIFVTVI
 SLLVYKHKHEYNPIENSPGNVVR SKGLSVFLNRAKAVFFPGNQEKDPLLKNQEFGKVS

(12) The adhesion molecule CD56 (SwissProt ID P13591)

>gi|94420689|ref|NP_000606.3| neural cell adhesion molecule 1
 isoform 1 [*Homo sapiens*]
 MLQTKDLIWT LFFLGTAVSLQVDIVPSQGEISVGESKFFLCQVAGDAKDKDISWFSPN
 GEKLTNPQQRISVVWNDSSSTLT IYANIDDA GIYKCVVTGEDGSESEATVNVKIFQ

-continued

KLMFKNAPTQPEFREGEDAIVIVCDVVSLLPPTIIWKHKGRDVI LKKDVRFIVLSNNYL
QIRGIKKTDEGTYRCEGRILARGEINFKDIQVIVNVPPTIQARQNIVNATANLGQSVTL
VCDAEGFPEPTMSWTKDGEQIEQEEDDEKYIFSDDSSQLTIKKVDKNDEAEYICIAEN
KAGEQDATIHLKVFAKPKITYVENQTAMELEEQVTLTCEASGDPIPSITWRTSTRNISS
EEKTLDGHMVVRSHARVSSLTKSIQYTDAGEYICTASNTIGQDSQSMYLEVQYAPK
LQGPVAVYTWEGNQVNITCEVFAYPSATISWFRDQQLPSSNYSNIKIYNTPSASYLEV
TPDSENDGNYNCTAVNRIGQESLEFILVQADTPSSPSIDQVEPYSSAQVQFDEPEAT
GGVPILKYKAEWRAGVEEVHWSKYDAKEASMEGIVTIVGLKPETTYAVRLAALNGKGL
GEISAASEFKTQPVQGEPSAPKLEGQMGEDGNSIKVNLIKQDDGGSPIRHYLVRYRAL
SSEWKPEIRLPSGSDHVMLKSLDWNAAEYEVVVAENQQGKSKAAHFVFR TSAQPTAIP
ANGSPTSGSLTGAIVGILIVIFVLLLVVDITCYFLNKCGLFMCIAVNL CGKAGPGAK
GKDMEEGKAASFSDSKESKEPIVEVRTEEERTPNHDGGKHTEPNETPLTEPEKGPVEAK
PECQETETKPAPAEVKTVPN DATQTKENESKA

(13) The surface molecule CD70 (SwissProt ID P32970)

>gi|4507605|ref|NP_001243.1| CD70 antigen [*Homo sapiens*]
MP EEGSGCSVRRRPYGCVLRAALVPLVAGLVI CLVVCIQRFQAQQQLPLESLGWD
VAELQLNHTGPQQDPRLYWQGGPALGRSFLHGP ELDKGQLRIHRDGIYMVHIQVTL
AICSSTTASRHHPTTLAVGICSPASRSISLLRLSFHQGTIASQRLTPLARGDTLCTNLT
GTLLPSRNTDETFFGVQWVRP

(14) The surface molecule CD74 (SwissProt ID P04233)

>gi|10835071|ref|NP_004346.1| HLA class II histocompatibility antigen
gamma chain isoform b [*Homo sapiens*]
MHRRRSRSCREDQKPVMDQQRDLISNNEQLPMLGRRPGAPESKCSR GALYTGFSILV
TLLLAGQATTAYFLYQQQGRLDKLTVTSTQNQLLENLRMKLPKPKPVSKMRMATPL
LMQALPMGALPQGP MQNATKYGNMTEDHVMHLLQNADPLKVYPPLKGSFPENLRH
LKNTMETIDWKVFESWMHHWLLFEMSRHSLEQKPTDAPPKESLELEDPS SGLGVTK
QDLGPVPM

(15) The B-lymphocyte antigen CD19 (SwissProt ID P15391)

>gi|296010921|ref|NP_001171569.1| B-lymphocyte antigen CD19 isoform 1
precursor [*Homo sapiens*]
MPPPRLLFFLLFLT PMEVRPEEPLVVKVEEGDNAVLQCLKGTSDGPTQQLTWSRESPL
KPFLKLSGLPLGLGIHMRPLAIWLFIFNVSQQMGGFYLCQPGPPSEKAWQPGWTVNV
EGSGELFRWNVSDLGGLGCKNRSSEGPSSPSGKLMSPKLYVWAKDRPEIWEGEPP
CLPPRDSL NQSLSQDLTMAPGSTLWLS CGVPPDSVSRGPLSWTHVHPKGPKSLLSLEL
KDDRPARDMWVMTGLLLPRATAQDAGKYXCHRGNTMSFHLEITARPVLWHWL
LRTGWKVS AVTTLAYLIFCLCSLVGILHLQRALVLRKRKRMTDPTRRFKVTPPPG
SGPQNQYGNVLSLPTPTSGLGRAQRWAAGLGGTAPSYGNPSSDVQADGALGSRSP
GVGPEEEEGEGYEEDSEEDSEFYENDSNLGQDQLSQDGS GYENPEDEPLGPEDEDSF

-continued

SNAESYENEDDELTQPVARTMDFLSPHGSADWPSREATSLAGSQSYEDMRGILYAAP

QLRSIRGQPGPNHEEDADSYENMDNPDGPDPAWGGGGRMGTTWSTR

(16) The surface protein mucin-1 (SwissProt ID P15941)

>gi|65301117|ref|NP_002447.4| mucin-1 isoform 1 precursor [*Homo sapiens*]
MTPGTQSPFFLLLLLTVLTVVTGSGHASSTPGGEKETSATQRSSVPSSTEKNALSTGVS

FFFLSPHISNLQFNSSLEDPSDYYQELQRDISEMFLQIYKQGGFLGLSNIKFRPGSVVV

QLTLAFREGTINVHDTVETQFNQYKTEAASRYNLTISDVSVDVPPFSAQSGAGVPG

WGIALLVLCVLVALAIVYLIALAVCCRRKNYGQLDIFPARDTYHPMSEYPTYHTH

GRYVPPSSTDRSPYEKVSAGNGGSSLSYTNPAVAATSANL

(17) The surface protein CD138 (SwissProt ID P18827)

>gi|29568086|ref|NP_002988.3| syndecan-1 precursor [*Homo sapiens*]
MRRALWLWLCALALSLQPALPQIVATNLPEDQDGGSDSDNFGSGAGALQDIT

LSQQTPSTWKDTQLLTAIPTSPPTGLEATAASTSTLPAGEGPKGEAVVLPVEVEPGLT

AREQEATPRPRETTQLPTHQASTTTATTAQEPAATSHPHRDMQPGHHETSTPAGPSQ

ADLHTPHTEDGGPSATERAEDGASSQLPAAEGSGEQDFTFETSGENTAVVAVEPDR

RNQSPVDQAGTASQGLLDRKEVLGGVIAGGLVGLIFAVCLVGFMLYRMKKKDEGS

YSLEEPKQANGGAYQKPTKQEEFYA

(18) The integrin alphaV (Genbank Accession No.:
NP_002201.1)

>gi|4504763|ref|NP_002201.1| integrin alpha-V isoform 1 precursor
[*Homo sapiens*]
MAFPFRRRLRLGPRGLPLLLSGLLLPLCRAFNLVDSPAESYSGPEGSYFGFAVDFVFP

SASSRMFLLVGAPKANTTQPGIVEGGQVLKCDWSSTRRCQPIEFDATGNRDYAKDDP

LEPKSHQWFGASVRSKQDKILACAPLYHWRTEMKQEREPVGTCLQDGTKTVEYAP

CRSQDIDADGGQFCQGGFSIDFTKADRVLLGGPGSFYWQQLISDQVAEIVSKYDPN

VYSIKYNNQLATRTAQAIFFDDSYLGYSVAVGDFNGDGIDDFVSGVPRAARTLGMVYI

YDGKNMSSLYNFTGEQMAAYFGFSAATDINGDDYADVFIGAPLFMDRGSDGKLQE

VGQVSVSLQRASGDFQTTKLNGFEVFARFGSAIAPLGDLQDGFNDIAIAAPYGGED

KKGIVYIFNGRSTGLNAVPSQILEGQWAARSMPPSPGYSMKGATDIDKNGYDPLIVG

AFGVDRAILYRARPVITVNAGLEVYPSILNQDNKTCSLPGTALKVSCFNVRFLKAD

GKGVLPKRLNFQVELLLDKLKQKGAIRRALFLYSRSPSHSKNMTISRGLMQCEELIA

YLRDESEFRDKLTPITIFMEYRLDYRTAADTTGLQPIILNQFTPANISRQAHILLDCGED

NVCKPKLEVSVDSDQKKIYIGDDNPLTLIVKAQNQEGEGAYEALIVSIPLQADFIGVV

RNNEALARLSAPKTENQTRQVVCDLGNPMKAGTQLLAGLRFVHQQSEMDTSVKF

DLQIQSSNLFDKVSPVVSFKVLDLAVLAAVEIRGVSSPDHIFLPINWEHKENPETEED

VGPVQHIYELRNNGPSSFSKAMLHLQWPYKYNNTLLYILHYDIDGPMNCTSDMEI

NPLRIKISSLQTTTEKNDTVAGQGERDHLITKRDALALSEGDIHTLGCQVAQCLKIVCQV

GRLDRGKSAILYVKSLLWTETFMNKENQNHSSYSLKSSASFNVIEFPYKNLPIDITNST

-continued

LVTNTVWGIQAPAMPVPVWVILAVLAGLLLLAVLVFVMYRMGFFKRVRPPEEQ
EREQLQPHENGEGNSET

(19) The teratocarcinoma-derived growth factor 1 protein
TDGF1 (Genbank Accession No.: NP_003203.1)

>gi|4507425|ref|NP_003203.1| teratocarcinoma-derived growth factor 1
isoform 1 precursor [*Homo sapiens*]
MDCRKMARFSYSYVIWIMAIKVFELGLVAGLGHQEFARPSRGYLAFRDDSIWPQEEP
AIRPRSSQVRPPMGIQHSKELNRTCCLNGGTCMLGSPCACPPSFYGRNCEHDVRKEN
CGSVPHDTWLPKKCSLCKCWHGQLRCFPQAFLPGCDGLVMDEHLVASRTPELPPSA
RTTTFMLVIGICLSIQSYY

(20) The prostate-specific membrane antigen PSMA (Swiss
Prot ID: Q04609)

>gi|4758398|ref|NP_004467.1| glutamate carboxypeptidase 2 isoform 1
[*Homo sapiens*]
MWNLHETDSAVATARRPRWL CAGALVLAGGFLLGLFGWFIKSSNEATNITPKH
NMKAFLDELKAENIKKFLYNFTQIPHLAGTEQNFQLAKQIQSQWKEGLDSVELAHY
DVLLSYPNKTHPNYISIIINEDGNEIFNTSLFEPPPPGYENVSDIVPPPSAFSPQGMPEGD
LVVYNYARTEDDFKLERDMKINCSGKIVIARYGKVFGRNKVKNAQLAGAKGVILYS
DPADYFAPGVKSYPDGWNLPGGGVQGRNINLNLGAGDPLTPGYPANFYAYRRGIAE
AVGLPSIPVHPIGYYDAQKLEKMGGSAPPDSSWRGSLKVPYNVGPFGFTGNFSTQKV
KMHISTNEVTRIYNVIGTLRGAVEPDRYVILGGHRDSWVFGIDPQSGAAVVHEIV
RSFGTLKKEGWRPRRTILFASWDAEEFGLLGSTEWAEENSRLQERG VAYINADSSIE
GNYTLRVDCTPLMYSLVHNLTKEKSPDEGFEGKSLYESWTKKSPSPFSGMPRISKL
GSGNDFEVFPQRLGIASGRARYTKNWETNKPSGYPLYHSVYETVELVEKFYDPMFK
YHLTVAQVRGGMVFELANSIVLPFDCRDYAVVLRYADKIYSISMKHPQEMKTYSV
SFDSLFSAVKNFTEIASKFSERLQDFDKSNPIVLRMMNDQLMFLERAFIDPLGLPDRPF
YRHVIYAPSSHNNKYAGESFPGIYDALFDIESKVDPSKAWGEVQRQIYVAAPTQAAA
ETLSEVA

(21) The tyrosine protein kinase EPHA2 (Swiss Prot ID:
P29317)

>gi|32967311|ref|NP_004422.2| ephrin type-A receptor 2 precursor
[*Homo sapiens*]
MELQAARACFALLWGCALAAAAAQGKEVVLLDFAAAGGELGWLTHPYGKGWDL
MQNIMNDMPIYMSVCNVMSGDQDNWLRNTNWVYRGEAERIFIELKFTVRDCNSFP
GGASSCKETFNLYAESDLDYGTNFQKRLFTKIDTIAPDEITVSSDFEARHVKLNVEE
RSVGPLTRKGFYLAQDIGACVALLSVRVYKKCPPELLQGLAHFPETIAGSDAPSLAT
VAGTCVDHAVVPPGGEEPRMHCAVDGEWLVPIGQCCLCQAGYEKVEDACQACSPGF
FKFEASESPCLECEHTLPSPEGATSCCEEGFFRAPQDPASMPCTRPPSAPHYLTAVG
MGAKVELRWTPQDSGGREDIVYSVTCEQCWPESGECGPCASVRYSEPPHGLTRTS
VTVSDLEPHMNYTFTVEARNGVSGLVTSRFRSTASVSINQTEPPKVRLEGRSTTSLSV

-continued

SWSIPPPQQSRVWKYEVTYRKKGDSNSYNVRRTEGFSVTLDDLAPDTTYLVQVQAL
 TQEGQGAGSKVHEFQTLSPGSGNLAVIGGVAVGVVLLVLAGVGFFIHRRRKNQR
 ARQSPEDVYFSKSEQLKPLKTYVDPHTYEDPNQAVLKFTTEIHPSCVTRQKVIGAGEF
 GEVYKGMKLTSSGKKEVPVAIKTLKAGYTEKQRVDFLGEAGIMGQFSHHNIIRLEGV
 ISKYKPMMIITEYMENGALDKFLREKDGESVLQLVGMLRGIAAGMKYLANMNYV
 HRDLAARNILVNSNLVCKVSDFGLSRVLEDDPEATYTTSGGKIPIRWTAPEAISYRKF
 TSASDVWSFGIVMWEVMTYGERPYWELSNHEVMKAINDFRLPTPMDCPSAIYQLM
 MQCWQQRARRPKFADIVSILDKLIRAPDSLKTLDADFDPVRSIRLPSTSGSEGVFPRTV
 SEWLESIKMQQYTEHFMAAGYTAIEKVVMQMTNDDIKRIGVRLPGHQKRIAYSLLGLK
 DQVNTVGIPI

(22) The surface protein SLC44A4 (Genbank Accession No: NP_001171515)

>gi|295849282|ref|NP_001171515.1| choline transporter-like protein 4
 isoform 2 [*Homo sapiens*]
 MGGKQRDEDEAYGKPVKYDPSFRGPIKNRSDTVICCVLPLLFIILGYIVVGIVAWL
 YGDPRQVLYPRNSTGAYCGMGENKDKPYLLYFNIFSCILSSNIISVAENGLQCPTQT
 VITSLQQELCPSFLLPSAPALGRCFPWNTVTPPALPGITNDTTIQQGISGLIDSLNARDIS
 VKIFEDFAQSWYIWLVALGVALVLSLLFILLRLVAGPLVLVLILGVLGVLAYGIYYC
 WEEYRVLRDKGASISQLGFTTNLSAYQSVQETWLAALIVLAVLEAILLLMLIFLRQRI
 RIAIALKEASKAVGQMMSTMFYPLVTFVLLLICIAYWAMTALYLATSGQPQYVLW
 ASNISSPGCEKVPINTSCNPTAHLVNSSCPGLMCFVFGYSSKGLIQRSVFNLQIYGVLG
 LFWTLNWLALGQCVLAGAFASFYWAFHKPDIPFPLISAFIRTLRYHTGSLAFGAL
 ILTLVQIARVILEYIDHKLGRGVQNPVARCIMCCFKCLWCLEKFIKFLNRNAYIMIAIY
 GKNFCVSAKNAPMLMRNIVRVVLDKVTDLLFPFGKLLVGGVGVLSFFFFSGRIP
 GLGKDFKSPHLNYYWLPIMTSILGAYVIASGFFSVFGMCDTLFLCFLEDLERNNGSL
 DRPPYMSKSLKILGKNEAPPDNKKRKK

(23) The surface protein BMPR1B (SwissProt: O00238)
 (24) The transport protein SLC7A5 (SwissProt: Q01650)
 (25) The epithelial prostate antigen STEAP1 (SwissProt: Q9UHE8)
 (26) The ovarian carcinoma antigen MUC16 (SwissProt: Q8WXI7)
 (27) The transport protein SLC34A2 (SwissProt: O95436)
 (28) The surface protein SEMA5b (SwissProt: Q9P283)
 (29) The surface protein LYPD1 (SwissProt: Q8N2G4)
 (30) The endothelin receptor type B EDNRB (SwissProt: P24530)
 (31) The ring finger protein RNF43 (SwissProt: Q68DV7)
 (32) The prostate carcinoma-associated protein STEAP2 (SwissProt: Q8NFT2)
 (33) The cation channel TRPM4 (SwissProt: Q8TD43)
 (34) The complement receptor CD21 (SwissProt: P20023)
 (35) The B-cell antigen receptor complex-associated protein CD79b (SwissProt: P40259)
 (36) The cell adhesion antigen CEACAM6 (SwissProt: P40199)

(37) The dipeptidase DPEP1 (SwissProt: P16444)
 (38) The interleukin receptor IL20Ralpha (SwissProt: Q9UHF4)
 (39) The proteoglycan BCAN (SwissProt: Q96GW7)
 (40) The ephrin receptor EPHB2 (SwissProt: P29323)
 (41) The prostate stem cell-associated protein PSCA (Genbank Accession No: NP_005663.2)
 (42) The surface protein LHFPL3 (SwissProt: Q86UP9)
 (43) The receptor protein TNFRSF13C (SwissProt: Q96RJ3)
 (44) The B-cell antigen receptor complex-associated protein CD79a (SwissProt: P11912)
 (45) The receptor protein CXCR5 (SwissProt: P32302)
 (46) The ion channel P2X5 (SwissProt: Q93086)
 (47) The lymphocyte antigen CD180 (SwissProt: Q99467)
 (48) The receptor protein FCRL1 (SwissProt: Q96LA6)
 (49) The receptor protein FCRL5 (SwissProt: Q96RD9)
 (50) The MHC class II molecule Ia antigen HLA-DOB (Genbank Accession No: NP_002111.1)
 (51) The T-cell protein VTCN1 (SwissProt: Q7Z7D3).

[2817] In one preferred subject of the invention the cancer target molecule is selected from the group consisting of the cancer target molecules (1)-(51).

[2818] In another particularly preferred subject of the invention the binder binds to an extracellular cancer target molecule which is selected from the group consisting of the cancer target molecules (1)-(51).

[2819] In another particularly preferred subject of the invention the binder binds specifically to an extracellular cancer target molecule which is selected from the group consisting of the cancer target molecules (1)-(51).

[2820] In one particularly preferred subject of the invention the cancer target molecule is selected from the group consisting of EGF receptor (NP_005219.2), mesothelin (Q¹³⁴²¹-3), C4.4a (NP_055215.2) and carboanhydrase IX (CA IX; NP_001207.2).

[2821] In another particularly preferred subject of the invention the binder binds to an extracellular cancer target molecule which is selected from the group consisting of EGF receptor (NP_005219.2), mesothelin (Q¹³⁴²¹-3), C4.4a (NP_055215.2) and carboanhydrase IX (CA IX; Q¹⁶⁷⁹⁰).

[2822] In another particularly preferred subject of the invention the binder binds specifically to an extracellular cancer target molecule which is selected from the group consisting of EGF receptor (NP_005219.2), mesothelin (Q¹³⁴²¹-3), C4.4a (NP_055215.2) and carboanhydrase IX (CA IX; Q¹⁶⁷⁹⁰).

[2823] In one preferred embodiment the binder, after binding to its extracellular target molecule on the target cell, is internalized by the target cell as a result of the binding. The effect of this is that the binder-drug conjugate, which may be an immunoconjugate or an ADC, is taken up by the target cell.

[2824] In one embodiment the binder is a binding protein. In one preferred embodiment the binder is an antibody, an antigen-binding antibody fragment, a multispecific antibody or an antibody mimetic.

[2825] Preferred antibody mimetics are affibodies, adnectins, anticalins, DARPin, avimers, or nanobodies. Preferred multispecific antibodies are bispecific and trispecific antibodies.

[2826] In one preferred embodiment the binder is an antibody or an antigen-binding antibody fragment, more preferably an isolated antibody or an isolated antigen-binding antibody fragment.

[2827] Preferred antigen-binding antibody fragments are Fab, Fab', F(ab')₂ and Fv fragments, diabodies, DABs, linear antibodies and scFv. Particularly preferred are Fab, diabodies and scFv.

[2828] In one particularly preferred embodiment the binder is an antibody. Particularly preferred are monoclonal antibodies or antigen-binding antibody fragments thereof. Further particularly preferred are human, humanized or chimeric antibodies or antigen-binding antibody fragments thereof.

[2829] Antibodies or antigen-binding antibody fragments which bind cancer target molecules may be prepared by a person of ordinary skill in the art using known processes, such as, for example, chemical synthesis or recombinant expression. Binders for cancer target molecules may be acquired commercially or may be prepared by a person of ordinary skill in the art using known processes, such as, for example, chemical synthesis or recombinant expression. Further processes for preparing antibodies or antigen-binding antibody fragments are described in WO 2007/070538 (see page 22 "Antibodies"). The skilled person knows how processes such

as phage display libraries (e.g. Morphosys HuCAL Gold) can be compiled and used for discovering antibodies or antigen-binding antibody fragments (see WO 2007/070538, page 24 ff and Example 1 on page 70, Example 2 on page 72). Further processes for preparing antibodies that use DNA libraries from B-cells are described for example on page 26 (WO 2007/070538). Processes for humanizing antibodies are described on page 30-32 of WO2007070538 and in detail in Queen, et al., Pros. Natl. Acad. Sci. USA 86:10029-10033, 1989 or in WO 90/0786. Furthermore, processes for the recombinant expression of proteins in general and of antibodies in particular are known to the skilled person (see, for example, in Berger and Kimmel (Guide to Molecular Cloning Techniques, Methods in Enzymology, Vol. 152, Academic Press, Inc.); Sambrook, et al., (Molecular Cloning: A Laboratory Manual, (Second Edition, Cold Spring Harbor Laboratory Press; Cold Spring Harbor, N.Y.; 1989) Vol. 1-3); Current Protocols in Molecular Biology, (F. M. Ausubel et al. [Eds.], Current Protocols, Green Publishing Associates, Inc./ John Wiley & Sons, Inc.); Harlow et al., (Monoclonal Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press (1988), Paul [Ed.]); Fundamental Immunology, (Lippincott Williams & Wilkins (1998)); and Harlow, et al., (Using Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press (1998)). The skilled person knows the corresponding vectors, promoters and signal peptides which are necessary for the expression of a protein/antibody. Commonplace processes are also described in WO 2007/070538 on pages 41-45. Processes for preparing an IgG1 antibody are described for example in WO 2007/070538 in Example 6 on page 74 ff. Processes which allow the determination of the internalization of an antibody after binding to its antigen are known to the skilled person and are described for example in WO 2007/070538 on page 80. The skilled person is able to use the processes described in WO 2007/070538 that have been used for preparing carboanhydrase IX (Mn) antibodies in analogy for the preparation of antibodies with different target molecule specificity.

EGFR Antibodies

[2830] Examples of antibodies which bind the cancer target molecules EGFR are cetuximab (INN number 7906), panitumumab (INN number 8499) and nimotuzumab (INN number 8545). Cetuximab (Drug Bank Accession Number DB00002) is a chimeric anti-EGFR1 antibody which is produced in SP2/0 mouse myeloma cells and is sold by ImClone Systems Inc/Merck KGaA/Bristol-Myers Squibb Co. Cetuximab is indicated for the treatment of metastasizing, EGFR expressing, colorectal carcinoma with wild type K-Ras gene. It has an affinity of 10⁻¹⁰M

Sequence:

Cetuximab Light Chain (Kappa):

[2831]

```
DILLTQSPVILSVSPGERVVFSCRASQSIGTNIHWYQQRNGSPRLLIK
ASESISGIPSRFSGSGSGTDFTLINSINVESEDIAYYCQNNNWPPTFGA
GTKLELKRTVAAPSVFIFPPSDEQLKSGTASVVCCLNNFYFPREKQVQWKV
```

-continued

DNALQSGNSQESVTEQDQSKDSTYLSSTLTLSKADYEKHKVYACEVTHQG
LSSPVTKSFNRGEC

Cetuximab Heavy Chain:

[2832]

QVQLKQSGPGLVQPSQSLITCTVSGFSLTNYGVHWRQSPGKGLEWLGV
IWSGNTDYNTPFTRLSINKDNSKQVFFKMNSLQSNDAIYYCARALT
YYDYEFAYWGQGLTVTSAASTKGPSVFPLAPSSKSTSGGTAALGCLVKD
YFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTY
ICNVNHKPSNTKVDKRVPEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPK
DTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS
TYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQV
YTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLT
DSGDSFFLYSLKLTVDKSRWQQGNVFCSCVMHEALHNYTQKSLSLSPGK

[2833] Panitumumab (INN number 8499) (Drug Bank Accession Number DB01269) is a recombinant monoclonal human IgG2 antibody which binds specifically to the human EGF receptor 1 and is sold by Abgenix/Amgen. Panitumumab originates from the immunization of transgenic mice (XenoMouse). These mice are capable of producing human immunoglobulin (light and heavy chains). A specific B-cell clone was selected which produces antibodies against EGFR, and this clone was immortalized with CHO cells (Chinese hamster ovary cells). These cells are now used for the production of a 100% human antibody. Panitumumab is indicated for the treatment of EGFR-expressing, metastasizing colorectal carcinoma, which is resistant to chemotherapeutic treatment with fluoropyrimidine, oxaliplatin and irinotecan. It has an affinity of 10^{-11} M.

Sequence:

Panitumumab Light Chain (Kappa):

[2834]

DIQMTQSPSSLSASVGRVTITCQASQDISNYLNWYQQKPKAPKLLIYD
ASNLETGVPSRFSGSGSGTDFTFTISLQPEDIATYFCQHFHDLPLAFGG
GTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCCLNFPYPREAKVQWKV
DNALQSGNSQESVTEQDQSKDSTYLSSTLTLSKADYEKHKVYACEVTHQG
LSSPVTKSFNRGEC

Panitumumab Heavy Chain:

[2835]

QVQLQESGPGLVKPSSETLSLTCTVSGGSVSSGDYYWTWIRQSPGKGLEWI
GHIYYSGNTNYPNPSLKSRLTISIDTSKTQFSLKLSSVTAADTAIYYCVRD
RVTGAFDIWGQGMVTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKD

-continued

YFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSNFGTQTY
TCNVDHKPSNTKVDKTVKCCVECPCCAPPVAGPSVFLFPPKPKDTLM
ISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRV
VSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLT
PSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSG
SFFLYSLKLTVDKSRWQQGNVFCSCVMHEALHNYTQKSLSLSPG

[2836] Nimotuzumab (INN number 8545) (EP 00586002, EP 00712863) is a humanized monoclonal IgG1 antibody which binds specifically to the human EGF receptor 1 and is sold by YM BioScienecs Inc. (Mississauga Canada). It is produced in non-secreting NSO cells (mammalian cell line). Nimotuzumab is approved for the treatment of head-and-neck tumours, highly malignant astrocytoma and glioblastoma multiforms (not in EU and US) and pancreatic carcinoma (orphan drug, EMA). It has an affinity of 10^{-8} M.

[2837] Further embodiments of EGFR antibodies are as follows:

[2838] Zalutumumab/2F8/HuMax-EGFr, from Genmab A/S (WO 02/100348, WO 2004/056847, INN number 8605)

[2839] Necitumumab/11F8, ImClone/IMC-11F8, from ImClone Systems Inc. [Eli Lilly & Co] (WO 2005/090407 (EP 01735348-A1, US 2007/0264253-A1, U.S. Pat. No. 7,598,350, WO 2005/090407-A1), INN number 9083)

[2840] Matuzumab/anti-EGFR MAb, Merck KGaA/anti-EGFR MAb, Takeda/EMD 72000/EMD-6200/EMD-72000 and EMD-55900/MAB 425/monoclonal antibody 425, from Merck KGaA/Takeda (WO 92/15683, INN number 8103 (Matuzumab))

[2841] RG-7160/GA-201/GA201/R-7160/R7160/RG7160/RO-4858696/RO-5083945/R04858696/R05083945, from Glycart Biotechnology AG (Roche Holding AG) (WO 2010/112413-A1, WO 2010/115554)

[2842] GT-MAB 5.2-GEX/CetuGEX, from Glycotope GmbH (WO 2008/028686-A2 (EP 01900750-A1, EP 01911766-A1, EP 02073842-A2, US 2010/0028947-A1)

[2843] ISU-101, from Isu Abxis Inc (ISU Chemical Co Ltd)/Scancell (WO 2008/004834-A1)

[2844] ABT-806/mAb-806/ch-806/anti-EGFR monoc. antibody 806, from Ludwig Institute for Cancer Research/Abbott/Life Science Pharmaceuticals (WO 02/092771, WO 2005/081854 and WO 2009/023265)

[2845] SYM-004 (consists of two chimeric IgG1 antibodies (992 and 1024)), from Symphogen A/S (WO 2010/022736-A2)

[2846] MR1-1/MR1-1 KDEL, from IVAX Corp (Teva Pharmaceutical Industries Ltd) (Duke University), (Patent: WO2001/062931-A2)

[2847] Antibody against the deletion mutant, EGFRvIII, from Amgen/Abgenix (WO 2005/010151, U.S. Pat. No. 7,628,986)

[2848] SC-100, from Scancell Ltd (WO 01/088138-A1)

[2849] MDX-447/EMD 82633/BAB-447/H 447/MAB, EGFR, Medarex/Merck KGaA, from Bristol-Myers Squibb (US)/Merck KGaA (DE)/Takeda (JP), (WO 91/05871, WO 92/15683)

[2850] Anti-EGFR-Mab, from Xencor (WO 2005/056606)

[2851] DXL-1218/anti-EGFR monoclonal antibody (cancer), InNexus, from InNexus Biotechnology Inc., Pharmaprojects PH048638

[2852] In one preferred embodiment the anti-EGFR antibodies are selected from the group consisting of cetuximab, panitumumab, nimotuzumab, zalutumumab, necitumumab, matuzumab, RG-716, GT-MAB 5.2-GEX, ISU-101, ABT-806, SYM-004, MR1-1, SC-100, MDX-447, and DXL-1218.

[2853] In one particularly preferred embodiment the anti-EGFR antibodies are selected from the group consisting of cetuximab, panitumumab, nimotuzumab, zalutumumab, necitumumab and matuzumab.

[2854] The skilled person knows of processes which can be used to prepare further antibodies, from the CDR regions of the abovementioned antibodies by means of sequence variations, these further antibodies having a similar or better affinity and/or specificity for the target molecule.

[2855] In a further embodiment, the anti-EGFR antibodies or antigen-binding antibody fragments are selected from the group consisting of

antibodies or antigen-binding antibody fragments comprising the three CDR regions of the light chain and the three CDR regions of the heavy chain of one of the following antibodies: cetuximab, panitumumab, nimotuzumab, zalutumumab, necitumumab, matuzumab, RG-716, GT-MAB 5.2-GEX, ISU-101, ABT-806, SYM-004, MR1-1, SC-100, MDX-447, and DXL-1218.

[2856] In another embodiment the anti-EGFR antibodies or antigen-binding antibody fragments are selected from the group consisting of

antibodies or antigen-binding antibody fragments comprising the three CDR regions of the light chain and the three CDR regions of the heavy chain of one of the following antibodies: cetuximab, panitumumab, nimotuzumab, zalutumumab, necitumumab, matuzumab.

Carboanhydrase IX Antibodies

[2857] Examples of antibodies which bind the cancer target molecule carbonanhydrase IX are described in WO 2007/070538-A2 (e.g. Claims 1-16).

[2858] In one preferred embodiment the anti-carboanhydrase IX antibodies or antigen-binding antibody fragments are selected from the group consisting of anti-carboanhydrase IX antibodies or antigen-binding antibody fragments 3ee9 (Claim 4 (a) in WO 2007/070538-A2), 3ef2 (Claim 4 (b) in WO 2007/070538-A2), 1e4 (Claim 4 (c) in WO 2007/070538-A2), 3a4 (Claim 4 (d) in WO 2007/070538-A2), 3ab4 (Claim 4 (e) in WO 2007/070538-A2), 3ah10 (Claim 4 (f) in WO 2007/070538-A2), 3bb2 (Claim 4 (g) in WO 2007/070538-A2), 1aa1 (Claim 4 (h) in WO 2007/070538-A2), 5a6 (Claim 4 (i) in WO 2007/070538-A2) and 5aa3 (Claim 4 (j) in WO 2007/070538-A2).

[2859] In one preferred embodiment the anti-carboanhydrase IX antibodies or antigen-binding antibody fragments are selected from the group consisting of:

anti-carboanhydrase IX antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody 3ee9 (from WO 2007/070538-A2),

anti-carboanhydrase IX antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the

three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody 3ef2 (from WO 2007/070538-A2),

anti-carboanhydrase IX antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody 1e4 (from WO 2007/070538-A2),

anti-carboanhydrase IX antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody 3a4 (from WO 2007/070538-A2),

anti-carboanhydrase IX antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody 3ab4 (from WO 2007/070538-A2),

anti-carboanhydrase IX antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody 3ah10 (from WO 2007/070538-A2),

anti-carboanhydrase IX antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody 3bb2 (from WO 2007/070538-A2),

anti-carboanhydrase IX antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody 1aa1 (from WO 2007/070538-A2),

anti-carboanhydrase IX antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody 5a6 (from WO 2007/070538-A2), and

anti-carboanhydrase IX antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody 5aa3 (from WO 2007/070538-A2).

[2860] The here-indicated sequences of the CDR regions are disclosed in FIGS. 2a-2c, page 128-130 in WO 2007/070538-A2.

[2861] In one preferred embodiment the anti-carboanhydrase IX antibodies or antigen-binding antibody fragments are selected from the group consisting of:

an antibody or antigen-binding fragment which comprises the amino acid sequence of the variable light and variable heavy chains of the antibody 3ee9, as indicated in WO 2007/070538-A2 in FIG. 4b on page 137,

an antibody or antigen-binding fragment which comprises the amino acid sequence of the variable light and variable heavy chains of the antibody 3ef2, as indicated in WO 2007/070538-A2 in FIG. 4c on page 138 and in FIG. 4b on page 137, an antibody or antigen-binding fragment which comprises the amino acid sequence of the variable light and variable heavy chains of the antibody 1e4, as indicated in WO 2007/070538-A2 in FIG. 4a on page 136,

an antibody or antigen-binding fragment which comprises the amino acid sequence of the variable light and variable heavy chains of the antibody 3a4, as indicated in WO 2007/070538-A2 in FIG. 4a on page 136,

an antibody or antigen-binding fragment which comprises the amino acid sequence of the variable light and variable heavy chains of the antibody 3ab4, as indicated in WO 2007/070538-A2 in FIG. 4a on page 136,

an antibody or antigen-binding fragment which comprises the amino acid sequence of the variable light and variable heavy chains of the antibody 3ah10, as indicated in WO 2007/070538-A2 in FIG. 4a on page 136,

an antibody or antigen-binding fragment which comprises the amino acid sequence of the variable light and variable heavy chains of the antibody 3bb2, as indicated in WO 2007/070538-A2 in FIG. 4b on page 137,

an antibody or antigen-binding fragment which comprises the amino acid sequence of the variable light and variable heavy chains of the antibody 1aa1, as indicated in WO 2007/070538-A2 in FIG. 4a on page 136,

an antibody or antigen-binding fragment which comprises the amino acid sequence of the variable light and variable heavy chains of the antibody 5a6, as indicated in WO 2007/070538-A2 in FIG. 4b on page 137, and

an antibody or antigen-binding fragment which comprises the amino acid sequence of the variable light and variable heavy chains of the antibody 5aa3, as indicated in WO 2007/070538-A2 in FIG. 4b on page 137.

[2862] In one particularly preferred embodiment the anti-carboanhydrase IX antibody is antibody 3ee9 from WO 2007/070538-A2.

[2863] In one particularly preferred embodiment the anti-carboanhydrase IX antibody or the antigen-binding antibody fragment comprises the amino acid sequences of the CDR regions of the variable heavy chain of the antibody 3ee9 (VH3-CDR1: GFTFSSYGMS; VH3-CDR2: GISSLGST-TYYADSVK; VH3-CDR3: TGSPGTFMHGDH, see FIG. 2a, page 128 in WO2007070538-A2) and the amino acid sequences of the CDR regions of the variable light chain of the antibody 3ee9 (VLk1-CDR1: RASQDINNYLS; VLk1-CDR2: YGASNLSQ; VLk1-CDR3: QYYGRPT, see FIG. 2b, page 129 in WO 2007/070538-A2).

[2864] In one particularly preferred embodiment the anti-carboanhydrase IX antibody or the antigen-binding antibody fragment comprises the amino acid sequences of a variable heavy chain of the antibody 3ee9

(VH3:ELVESGGGLVQPGGSLRLSCAASGFTF-SSYGMSWVRQAPGKGLEWVSGISSLG-STYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARTGSPGTFMHGDHW GQGTILTVSS, see FIG. 4b, page 137 in WO 2007070538-A2) and the amino acid sequences of the variable light chain of the antibody 3ee9 (VLk1:DIQMTQSPSSLSASVGDRTIT-CRAsQDINNYLSWYQQKPGKAPKLLIYGASNLQSGVPSRFSFGSGSGTDFTLTISLQPEDFAVYYCQYYGRPTTFGQGTKVEIKRT, see FIG. 4b, page 137 in WO 2007070538-A2).

[2865] In one preferred embodiment the anti-carboanhydrase IX antibody 3ee9 is an IgG antibody.

[2866] In one particularly preferred embodiment the anti-carboanhydrase IX antibody 3ee9 is an IgG1 antibody (3ee9-IgG1),

where the amino acid sequence of the heavy chain comprises the following sequence:

```
QVELVESGGGLVQPGGSLRLSCAASGFTFSSYGMSWVRQAPGKGLEWVSG
ISSLGSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARTG
SPGTFMHGDHWGQGTILTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLV
KDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQ
TYICNVNHKPSNTKVDKKVEPKSCDKHTHTCPPCPAPELLGGPSVFLFPPK
PKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY
NSTYRVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREP
QVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTP
VLDSGSGFFLYSKLTVDKSRWQQGNVFSCSVMEALHNYHTQKSLSLSP
GK
```

and the amino acid sequences of the light chain comprises the following sequence:

```
DIQMTQSPSSLSASVGDRTITCRASQDINNYLSWYQQKPGKAPKLLIYG
ASNLSQGVPSRFSFGSGSGTDFTLTISLQPEDFAVYYCQYYGRPTTFGQ
GTKVEIKRTVAAPSVEIFPPSDEQLKSGTASVVCCLNNFYPREAKVQWKV
DNALQSGNSQESVTEQDSKSTYSLSSTLTLSKADYEKHKVYACEVTHQG
LSSPVTKSFNRGEC
```

anti-carboanhydrase IX antibody 3ee9-IgG1:

[2867] A further aspect of the present invention is the provision of the anti-carboanhydrase IX antibody 3ee9-IgG1.

C4.4a Antibodies:

[2868] Examples of C4.4a antibodies and antigen-binding fragments are described below. The sequences of the antibodies are indicated in Table 1, with each line reproducing the respective CDR amino acid sequences of the variable light chain and of the variable heavy chain, respectively of the antibody listed in column 1. The amino acid sequences of the variable light chain and of the variable heavy chain, and the nucleic acid sequence of the antibody indicated in column 1 in each case, are also indicated.

[2869] In one embodiment the anti-C4.4a antibodies or antigen-binding antibody fragments bind to the S1 domain S1 (amino acid position 1-85 of SEQ ID NO: 1) of C4.4a.

[2870] In one embodiment the anti-C4.4a antibodies or antigen-binding antibody fragments are cross-reactive with human C4.4a (SEQ ID NO:1) and with murine C4.4a (SEQ ID NO:2).

[2871] In one embodiment the anti-C4.4a antibodies or antigen-binding antibody fragments thereof, after binding to a cell which expresses C4.4a, are internalized by the cell.

[2872] In another embodiment the anti-C4.4a antibodies or antigen-binding antibody fragments compete with the antibody M31-B01 and/or with the antibody M20-D02-S-A for binding to C4.4a. Antibodies M31-B01 and M20-D02-S-A compete for binding to C4.4a. The antibodies B01-1 to B01-12 were prepared from M31-B01 by means of affinity maturation and compete with M31-B01 for binding to C4.4a. The antibodies D02-1 to D02-13 were prepared from M20-D02-S-A by means of affinity maturation and compete with M20-D02-S-A for binding to C4.4a.

[2873] In a further embodiment the anti-C4.4a antibodies or antigen-binding antibody fragments comprise at least one, two or three of the CDR amino acid sequences given in Table 1 or Table 2.

[2874] In another embodiment the anti-C4.4a antibodies or antigen-binding antibody fragments comprise at least one, two or three CDR amino acid sequences of an antibody given in Table 1 or Table 2.

[2875] In a further embodiment the anti-C4.4a antibodies or antigen-binding antibody fragments comprise at least one, two or three CDR amino acid sequences of the variable light chain and at least one, two or three CDR amino acid sequences of the variable heavy chain of an antibody given in Table 1 or Table 2.

[2876] In another embodiment the anti-C4.4a antibodies or antigen-binding antibody fragments comprise which are at least 50%, 60%, 70%, 80%, 90% or 95% identical with the CDR amino acid sequences of the variable light chain and with the CDR amino acid sequences of the variable heavy chain, of an antibody given in Table 1 or Table 2.

[2877] In another embodiment the CDR sequences of the anti-C4.4a antibodies or antigen-binding antibody fragments comprise

CDR sequences of the heavy chain which conform to the CDR sequences SEQ ID NO: 297 (CDR H1), SEQ ID NO: 298 (CDR H2) and SEQ ID NO: 299 (CDR H3) and CDR sequences of the light chain which conform to the CDR sequences SEQ ID NO: 300 (CDR L1), SEQ ID NO: 22 (CDR L2) and SEQ ID NO: 301 (CDR L3), or

CDR sequences of the heavy chain which conform to the CDR sequences SEQ ID NO: 302 (CDR H1), SEQ ID NO: 303 (CDR H2) and SEQ ID NO: 304 (CDR H3) and CDR sequences of the light chain which conform to the CDR sequences SEQ ID NO: 305 (CDR L1), SEQ ID NO: 306 (CDR L2) and SEQ ID NO: 307 (CDR L3).

[2878] In another embodiment the anti-C4.4a antibodies or antigen-binding antibody fragments comprise which are at least 50%, 60%, 70%, 80%, 90% or 95% identical with the variable light chain and with the variable heavy chain, of an antibody given in Table 1 or Table 2.

[2879] In another embodiment the anti-C4.4a antibodies or antigen-binding antibody fragments comprise the three CDR amino acid sequences of the variable light chain and the three CDR amino acid sequences of the variable heavy chain of an antibody given in Table 1 or Table 2.

[2880] In another embodiment the anti-C4.4a antibodies or antigen-binding antibody fragments comprise a variable light chain and/or a variable heavy chain of an antibody given in Table 1 or Table 2.

[2881] In another embodiment the anti-C4.4a antibodies or antigen-binding antibody fragments comprise the variable light chain and the variable heavy chain of an antibody given in Table 1 or Table 2.

[2882] In one preferred embodiment the C4.4a antibodies and the antigen-binding antibody fragments are selected from the group consisting of

antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 75-77 and which comprises the CDR sequences of the variable light chain represented by the sequences SEQ ID NO: 78-80 (B01-10),

antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 5, 9

and 13 and which comprises the CDR sequences of the variable light chain represented by the sequences SEQ ID NO: 17, 21 and 25 (M31-B01),

antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 6, 10 and 14 and which comprises the CDR sequences of the variable light chain represented by the sequences SEQ ID NO: 18, 22 and 26 (M20-D02-S-A),

antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 7, 11 and 15 and which comprises the CDR sequences of the variable light chain represented by the sequences SEQ ID NO: 19, 23 and 27 (M60-G03),

antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 8, 12 and 16 and which comprise the CDR sequences of the variable light chain represented by the sequences SEQ ID NO: 20, 24 and 28 (36-H02),

antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 45-47 and which comprises the CDR sequences of the variable light chain represented by the sequences SEQ ID NO: 48-50 (B01-3),

antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 55-57 and which comprises the CDR sequences of the variable light chain represented by the sequences SEQ ID NO: 58-60 (B01-5),

antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 65-67 and which comprises the CDR sequences of variable light chain represented by the sequences SEQ ID NO: 68-70 (B01-7),

antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 85-87 and which comprises the CDR sequences of the variable light chain represented by the sequences SEQ ID NO: 88-90 (B01-12),

antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 95-97 and which comprises the CDR sequences of the variable light chain represented by the sequences SEQ ID NO: 98-100 (D02-4),

antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 105-107 and which comprises CDR sequences of the variable light chain represented by the sequences SEQ ID NO: 108-110 (D02-6),

antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 115-117 and which comprises the CDR sequences of the variable light chain represented by the sequences SEQ ID NO: 118-120 (D02-7),

antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 125-127 and which comprises the CDR sequences of the variable light chain represented by the sequences SEQ ID NO: 128-130 (D02-11),

and antibody which comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO: 135-137 which comprises the CDR sequences of the variable light chain represented by the sequences SEQ ID NO: 138-140 (D02-13).

[2883] In one preferred embodiment the C4.4a antibodies and the antigen-binding antibody fragments are selected from the group consisting of

antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 81 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 82 (B01-7),

antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 33 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 29 (M31-B01),

antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 34 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 30 (M20-D02 S-A),

antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 35 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 31 (M60-G03),

antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 36 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 32 (M36-H02),

antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 51 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 52 (B01-3),

antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 61 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 62 (B01-5),

antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 71 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 72 (B01-7)

antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 91 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 92 (B01-12),

antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 101 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 102 (D02-4),

antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 111 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 112 (D02-6),

antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 121 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 122 (D02-7),

antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 131 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 132 (D02-11),

and antibodies which comprise the amino acid sequence of the variable heavy chain represented by the sequence SEQ ID NO: 141 and which comprise the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO: 142 (D02-13).

[2884] In another embodiment the anti-C4.4a antibodies comprise the light chain and the heavy chain of an antibody given in Table 2.

[2885] In one preferred embodiment the anti-C4.4a antibodies comprise the light chain and the heavy chain of an antibody given in Table 2.

[2886] In one particularly preferred embodiment the C4.4a antibody is selected from the group consisting of antibody which comprises the amino acid sequence of the light chain represented by SEQ ID NO: 346 and which comprises the amino acid sequence of the heavy chain represented by SEQ ID NO: 347 (M31-B01),

antibody which comprises the amino acid sequence of the light chain represented by SEQ ID NO: 352 and which comprises the amino acid sequence of the heavy chain represented by SEQ ID NO: 353 (B01-3),

antibody which comprises the amino acid sequence of the light chain represented by SEQ ID NO: 364 and which comprises the amino acid sequence of the heavy chain represented by SEQ ID NO: 365 (B01-10), and

antibody which comprises the amino acid sequence of the light chain represented by SEQ ID NO: 382 and which comprises the amino acid sequence of the heavy chain represented by SEQ ID NO: 383 (D02-6).

TABLE 1

Antibody	Sequences of the C4.4a antibodies					
	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:
	HCDR1	HCDR2	HCDR3	LCDR1	LCDR2	LCDR3
M31-B01	5	9	13	17	21	25
M20-D02 S-A	6	10	14	18	22	26
M60-G03	7	11	15	19	23	27
M36-H02	8	12	16	20	24	28
B01-3	45	46	47	48	49	50
B01-5	55	56	57	58	59	60
B01-7	65	66	67	68	69	70
B01-10	75	76	77	78	79	80
B01-12	85	86	87	88	89	90
D02-4	95	96	97	98	99	100
D02-6	105	106	107	108	109	110
D02-7	115	116	117	118	119	120
D02-11	125	126	127	128	129	130
D02-13	135	136	137	138	139	140
B01-nn1	145	146	147	148	149	150
B01-nn2	153	154	155	156	157	158
B01-nn3	161	162	163	164	165	166
B01-nn4	169	170	171	172	173	174
B01-nn5	177	178	179	180	181	182
B01-2	185	186	187	188	189	190
B01-4	193	194	195	196	197	198
B01-6	201	202	203	204	205	206
B01-8	209	210	211	212	213	214
B01-9	217	218	219	220	221	222
B01-11	225	226	227	228	229	230
B01-12	233	234	235	236	237	238

TABLE 1-continued

Sequences of the C4.4a antibodies					
D02-ogl	241	242	243	244	245
D02-5	249	250	251	252	253
D02-8	257	258	259	260	261
D02-9	265	266	267	268	269
D02-10	273	274	275	276	277
D02-11	281	282	283	284	285
D02-12	289	290	291	292	293
D02-12	294				
Antibody	SEQ ID NO: VH Protein	SEQ ID NO: VL Protein	SEQ ID NO: VH Nucleotide	SEQ ID NO: VL Nucleotide	
M31-B01	33	29	41	37	
M20-	34	30	42	38	
D02 S-A					
M60-G03	35	31	43	39	
M36-H02	36	32	44	40	
B01-3	51	52	53	54	
B01-5	61	62	63	64	
B01-7	71	72	73	74	
B01-10	81	82	83	84	
B01-12	91	92	93	94	
D02-4	101	102	103	104	
D02-6	111	112	113	114	
D02-7	121	122	123	124	
D02-11	131	132	133	134	
D02-13	141	142	143	144	
B01-nn1	151	152	308	309	
B01-nn2	159	160	310	311	
B01-nn3	167	168	312	313	
B01-nn4	175	176	314	315	
B01-nn5	183	184	316	317	
B01-2	191	192	318	319	
B01-4	199	200	320	321	
B01-6	207	208	322	323	
B01-8	215	216	324	325	
B01-9	223	224	326	327	
B01-11	231	232	328	329	
B01-12	239	240	330	331	
D02-ogl	247	248	332	333	
D02-5	255	256	334	335	
D02-8	263	264	336	337	
D02-9	271	272	338	339	
D02-10	279	280	340	341	
D02-11	287	288	342	343	
D02-12	295	296	344	345	

TABLE 2

Sequences of the light and heavy chain of the C4.4a antibodies		
Antibody	Light chain SEQ ID NO:	Heavy chain SEQ ID NO:
M31-B01	346	347
B01-1	348	349
B01-2	350	351
B01-3	352	353
B01-4	354	355
B01-5	356	357
B01-6	358	359
B01-7	360	361
B01-8	362	363
B01-10	364	365
B01-11	366	367
B01-12	368	369
M20-D02 S-A	370	371
D02-1	372	373
D02-2	374	375
D02-3	376	377
D02-4	378	379
D02-5	380	381
D02-6	382	383

TABLE 2-continued

Sequences of the light and heavy chain of the C4.4a antibodies		
Antibody	Light chain SEQ ID NO:	Heavy chain SEQ ID NO:
D02-7	384	385
D02-8	386	387
D02-9	388	389
D02-10	390	391
D02-11	392	393
D02-12	394	395
D02-13	396	397

Anti-C4.4a Antibody IgG:

[2887] A further aspect of the present invention is the provision of an anti-C4.4a IgG1 antibody which comprises the amino acid sequence of the light chain and of the heavy chain of an antibody given in Table 2.

[2888] One example of an antibody which binds the cancer target molecule Her2 is trastuzumab (Genentech). Trastuzumab is a humanized antibody which is used for the treatment among other things of breast cancer. One example of an antibody which binds the cancer target molecule CD20 is rituximab (Genentech). Rituximab (CAS number: 174722-31-7) is a chimeric antibody which is used for the treatment of Non-Hodgkin's Lymphoma. One example of an antibody which binds the cancer target molecule CD52 is alemtuzumab (Genzyme). Alemtuzumab (CAS number: 216503-57-0) is a humanized antibody which is used for the treatment of chronic lymphatic leukaemia.

Mesothelin Antibody

[2889] Binders particularly preferred in accordance with the invention are anti-mesothelin antibodies, more particularly human or humanized anti-mesothelin antibodies. The antibodies preferably have an affinity of at least 10^{-7} M (as Kd value; in other words preferably those with smaller Kd values than 10^{-7} M), preferably of at least 10^{-8} M, more preferably in the range from 10^{-9} M to 10^{-11} M. The Kd values may be determined, for example, by means of surface plasmon resonance spectroscopy.

[2890] The antibody-drug conjugates of the invention likewise exhibit affinities in these ranges. The affinity is preferably not substantially affected by the conjugation of the drugs (in general, the affinity is reduced by less than one order of magnitude, in other words, for example, at most from 10^{-8} M to 10^{-7} M).

[2891] The antibodies used in accordance with the invention are also notable preferably for a high selectivity. A high selectivity exists when the antibody of the invention exhibits an affinity for the target protein which is better by a factor of at least 2, preferably by a factor of 5 or more preferably by a factor of 10, than for an independent other antigen, e.g. human serum albumin (the affinity may be determined, for example, by means of surface plasmon resonance spectroscopy).

[2892] Furthermore, the antibodies of the invention that are used are preferably cross-reactive. In order to be able to facilitate and better interpret preclinical studies, for example toxicological or activity studies (e.g. in xenograft mice), it is advantageous if the antibody used in accordance with the invention not only binds the human target protein but also

binds the species target protein in the species used for the studies. In one embodiment the antibody used in accordance with the invention, in addition to the human target protein, is cross-reactive to the target protein of at least one further species. For toxicological and activity studies it is preferred to use species of the families of rodents, dogs and non-human primates. Preferred rodent species are mouse and rat. Preferred non-human primates are rhesus monkeys, chimpanzees and long-tailed macaques.

[2893] In one embodiment the antibody used in accordance with the invention, in addition to the human target protein, is cross-reactive to the target protein of at least one further species selected from the group of species consisting of mouse, rat and long-tailed macaque (*Macaca fascicularis*). Especially preferred are antibodies used in accordance with the invention which in addition to the human target protein are at least cross-reactive to the mouse target protein. Preference is given to cross-reactive antibodies whose affinity for the target protein of the further non-human species differs by a factor of not more than 50, more particularly by a factor of not more than ten, from the affinity for the human target protein.

[2894] The antibodies of the invention that are used are additionally notable preferably for invariant binding to mesothelin. Invariant binding is notable, for example, by the fact that the antibody of the invention that is used binds to an epitope of mesothelin which cannot be masked by a further extracellular protein. Such a further extracellular protein is, for example, the protein Ovarian Cancer Antigen 125 (CA125). Antibodies used with preference are notable for the fact that their binding to mesothelin is not blocked by CA125.

[2895] Anti-mesothelin antibodies are described for example in WO 2009/068204. These antibodies can be used in accordance with the invention.

[2896] A further aspect of the present invention is the provision of a new anti-mesothelin antibody (MF-Ta) whose amino acid sequence comprises the CDR sequences of the variable heavy chain represented by the sequences SEQ ID NO:398 (HCDR1), SEQ ID NO:399 (HCDR2) and SEQ ID NO:400 (HCDR3) and the CDR sequences of the variable light chain represented by the sequences SEQ ID NO:401 (LCDR1), SEQ ID NO:402 (LCDR2) and SEQ ID NO:403 (LCDR3).

[2897] In one preferred embodiment the amino acid sequence of the anti-mesothelin antibody MF-Ta or antigen-binding antibody fragments comprises the sequence of the variable heavy chain represented by the sequences SEQ ID NO:404 and the sequence of the variable light chain represented by the sequence SEQ ID NO:405. In one preferred embodiment the amino acid sequence of the anti-mesothelin antibody MF-Ta or antigen-binding antibody fragments comprises the sequence of the variable heavy chain which is encoded by the nucleic acid sequence SEQ ID NO:406, and the sequence of the variable light chain which is encoded by the nucleic acid sequence SEQ ID NO:407.

[2898] In one particularly preferred embodiment the amino acid sequence of the anti-mesothelin antibody MF-Ta comprises the sequence of the heavy chain represented by the sequences SEQ ID NO:408 and the sequence of the light chain represented by the sequence SEQ ID NO:409.

[2899] In one particularly preferred embodiment the amino acid sequence of the anti-mesothelin antibody MF-Ta comprises the sequence of the heavy chain which is encoded with

a nucleic acid sequence SEQ ID NO:410, and the sequence of the light chain with is encoded with a nucleic acid sequence SEQ ID NO: 411.

[2900] Further examples of antibodies which bind the cancer target molecule mesothelin are known to the skilled person and are described for example in WO 2009/068204 and can be used for the binder-drug conjugates of the invention.

[2901] In one embodiment of the binder-drug conjugates, the binder is an anti-mesothelin antibody or antigen-binding antibody fragment, where the antibody binds to mesothelin and exhibits invariant binding.

[2902] In one embodiment of the binder-drug conjugates, an anti-mesothelin antibody or antigen-binding antibody fragment comprises the amino acid sequences of the three CDR regions of the light chain and the amino acid sequences of the three CDR regions of the heavy chain of an antibody described in WO 2009/068204-A1 (Table 7; page 61-63).

[2903] In one preferred embodiment the mesothelin antibodies or antigen-binding antibody fragments are selected from the group consisting of

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody MF-Ta, anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody MF-J (WO 2009/068204-A1; Table 7; page 61),

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody MOR06640 (WO 2009/068204-A1; Table 7; page 61),

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody MF-226 (WO 2009/068204-A1; Table 7; page 61) and

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody MOR06626 (WO 2009/068204-A1; Table 7; page 61).

[2904] In one particularly preferred embodiment the mesothelin antibodies or antigen-binding antibody fragments are selected from the group consisting of

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequence of the variable light chain and the sequence of the variable heavy chain of the antibody MF-Ta,

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequence of the variable light chain and the sequence of the variable heavy chain of the antibody MF-J (WO 2009/068204-A1; Table 7; page 61),

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequence of the variable light chain and the sequence of the variable heavy chain of the antibody MOR06640 (WO 2009/068204-A1; Table 7; page 61),

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequence of the variable light chain and the sequence of the variable heavy chain of the antibody MF-226 (WO 2009/068204-A1; Table 7; page 61),

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequence of the variable light chain and the sequence of the variable heavy chain of the antibody MOR06626 (WO 2009/068204-A1; Table 7; page 61).

Further Antibodies:

[2905] An example of an antibody which binds the cancer target molecule Her2 is trastuzumab (Genentech). Trastuzumab is a humanized antibody which is used for the treatment inter alia of breast cancer. One example of an antibody which binds the cancer target molecule CD20 is rituximab (Genentech). Rituximab (CAS number: 174722-31-7) is a chimeric antibody which is used for the treatment of non-Hodgkin's lymphoma. One example of an antibody which binds the cancer target molecule CD52 is alemtuzumab (Genzyme). Alemtuzumab (CAS number: 216503-57-0) is a humanized antibody which is used for the treatment of chronic lymphatic leukaemia.

[2906] Other examples of antibodies which bind to HER2, besides trastuzumab (INN 7637, CAS No: RN: 180288-69-1) and pertuzumab (Cas No: 380610-27-5), are antibodies as disclosed in WO 2009/123894-A2, WO 2008/140603-A2, or in WO 2011/044368-A2. An example of an anti-HER2 conjugate is trastuzumab-emtansine (INN No. 9295).

[2907] Examples of antibodies which bind the cancer target molecule CD30 and can be used for the treatment of cancer, e.g. Hodgkin's lymphoma, are brentuximab, iratumumab and antibodies as disclosed in WO 2008/092117, WO 2008/036688 or WO 2006/089232. An example of an anti-CD30 conjugate is brentuximab vedotone (INN No. 9144).

[2908] Examples of antibodies which bind the cancer target molecule CD22 and can be used for the treatment of cancer, e.g. lymphoma, are inotuzumab or epratuzumab. Examples of anti-CD22 conjugates are inotuzumab ozagamyacin (INN No. 8574), or anti-CD22-MMAE and anti-CD22-MC-MMAE (CAS RN: 139504-50-0 and 474645-27-7).

[2909] Examples of antibodies which bind the cancer target molecule CD33 and can be used for the treatment of cancer, e.g. leukaemia, are gemtuzumab or lintuzumab (INN 7580). An example of an anti-CD33 conjugate is gemtuzumab-ozagamyacin.

[2910] An example of an antibody which binds the cancer target molecule NMB and can be used for the treatment of cancer, e.g. melanoma or breast cancer, is glembatumumab (INN 9199). An example of an anti-NMB conjugate is glembatumumab vedotone (CAS RN: 474645-27-7).

[2911] An example of an antibody which binds the cancer target molecule CD56 and can be used for the treatment of cancer, e.g. multiple myeloma, small-cell carcinoma of the lung, MCC or ovarian carcinoma, is lorvotuzumab. An example of an anti-CD56 conjugate is lorvotuzumab mertansine (CAS RN: 139504-50-0).

[2912] Examples of antibodies which bind the cancer target molecule CD70 and can be used for the treatment of cancer, e.g. non-Hodgkin's lymphoma or kidney cell cancer, are disclosed in WO 2007/038637-A2 or WO 2008/070593-A2. An example of an anti-CD70 conjugate is SGN-75 (CD70 MMAF).

[2913] An example of an antibody which binds the cancer target molecule CD74 and can be used for the treatment of cancer, e.g. multiple myeloma, is milatuzumab. An example of an anti-CD74 conjugate is milatuzumab-doxorubicin (CAS RN: 23214-92-8).

[2914] An example of an antibody which binds the cancer target molecule CD 19 and can be used for the treatment of cancer, e.g. non-Hodgkin's lymphoma, is disclosed in WO 2008/031056-A2. Further antibodies and examples of an anti-CD19 conjugate (SAR3419) are disclosed in WO 2008/047242-A2.

[2915] Examples of antibodies which bind the cancer target molecule mucin-1 and can be used for the treatment of cancer, e.g. non-Hodgkin's lymphoma, are clivatuzumab or the antibodies disclosed in WO 2003/106495-A2, WO 2008/028686-A2. Examples of anti-mucin conjugates are disclosed in WO 2005/009369-A2.

[2916] Examples of antibodies which bind the cancer target molecule CD138 and conjugates thereof which can be used for the treatment of cancer, e.g. multiple myeloma, are disclosed in WO 2009/080829-A1, WO 2009/080830-A1.

[2917] Examples of antibodies which bind the cancer target molecule integrin alphaV and can be used for the treatment of cancer, e.g. melanoma, sarcoma or carcinoma, are intetumumab (Cas RN: 725735-28-4), abciximab (Cas-RN: 143653-53-6), etaracizumab (Cas-RN: 892553-42-3) or the antibodies disclosed in U.S. Pat. No. 7,465,449, EP 19859-A1, WO 2002/012501-A1 or WO 2006/062779-A2. Examples of anti-integrin alphaV conjugates are intetumumab-DM4 and other ADCs disclosed in WO 2007/024536-A2.

[2918] Examples of antibodies which bind the cancer target molecule TDGF1 and can be used for the treatment of cancer are the antibodies disclosed in WO 02/077033-A1, U.S. Pat. No. 7,318,924, WO 2003/083041-A2 and WO 2002/088170-A2. Examples of anti-TDGF1 conjugates are disclosed in WO 2002/088170-A2.

[2919] Examples of antibodies which bind the cancer target molecule PSMA and can be used for the treatment of cancer, e.g. prostate carcinoma, are the antibodies disclosed in WO 97/35616-A1, WO 99/47554-A1, and WO 01/009192-A1. Examples of anti-PSMA conjugates are disclosed in WO 2009/026274-A1.

[2920] Examples of antibodies which bind the cancer target molecule EPHA2, can be used for preparing a conjugate and can be used for the treatment of cancer are disclosed in WO 2004/091375-A2.

[2921] Examples of antibodies which bind the cancer target molecule SLC44A4, can be used for preparing a conjugate and can be used for the treatment of cancer, e.g. pancreatic or prostate carcinoma, are disclosed in WO 2009/033094-A2 and US 2009/0175796-A1.

[2922] An example of an antibody which binds the cancer target molecule HLA-DOB is the antibody lym-1 (Cas-RN: 301344-99-0), which can be used for the treatment of cancer, e.g. non-Hodgkin's lymphoma. Examples of anti-HLA-DOB conjugates are disclosed for example in WO 2005/081711-A2.

[2923] Examples of antibodies which bind the cancer target molecule VTCN1, can be used for preparing a conjugate and can be used for the treatment of cancer, e.g. ovarian carcinoma, pancreatic, lung or breast cancer, are disclosed in WO 2006/074418-A2.

[2924] The compounds of the invention possess valuable pharmacological properties and can be used for the prevention and treatment of diseases in humans and animals.

[2925] The binder-drug conjugates (ADCs) of the invention, of the formula (Ia), exhibit a high and specific cytotoxic activity with regard to tumour cells, as may be shown on the

basis of the assays set out in the present experimental section (C-1. to C-7e.). This high and specific cytotoxic activity on the part of the binder-drug conjugates (ADCs) of the invention, of the formula (Ia), is achieved through the appropriate combination of the new N,N-dialkylauristatin derivative and binder with linkers which exhibit not only an enzymatically, hydrolytically or reductively cleavable predetermined break point, for the release of the toxophores, but also no such predetermined break point. More particularly, through the use of stable linkers which have no enzymatically, hydrolytically or reductively cleavable predetermined break point for the release of the toxophores, and which, following uptake of the ADCs into the tumour cell and following complete intracellular, enzymatic breakdown of the antibody, still remain wholly or partly intact, the activity is confined very specifically to the tumour cell. Compatibility between ADCs and stable linkers presupposes, among other things, that the metabolites formed intracellularly can be formed with sufficient efficacy, are able to reach their target and are able there to develop their anti-proliferative activity on the target with sufficient potency, without being carried out of the tumour cell again beforehand by transporter proteins. The metabolites formed intracellularly after the compounds of the formula (Ia) of the invention have been taken up exhibit a reduced potential as a substrate with respect to transporter proteins, thereby suppressing their redistribution into the systemic circulation and hence the triggering of potential side effects by the toxophore itself. Furthermore, the innovative N-alkyl attachment maintains the basic character at the amino end of the monomethyl-auristatin peptide. Especially in the case of the binder-drug conjugates (ADCs) of the formula (Ia) of the invention that are linked via lysine side chains, the overall charge of the antibody remains constant, irrespective of the number of toxophore-linker loadings.

[2926] The compatibility of the ADCs with a stabile linker chemistry and with the target in question, in conjunction with metabolites which represent a substrate for transporter proteins to a relatively low degree, offers an enlarged therapeutic window.

[2927] More particularly, the binder-drug conjugates of the invention, of the formula (Ia), exhibit a high and specific cytotoxic activity with respect to tumour cells which express mesothelin. The activity with respect to tumour cells which do not express mesothelin is significantly weaker at the same time.

[2928] On the basis of this profile of properties, the compounds of the invention are therefore suitable to a particular degree for the treatment of hyperproliferative diseases in humans and in mammals generally. The compounds are able on the one hand to inhibit, block, reduce or lower cell proliferation and cell division, and on the other hand to increase apoptosis.

[2929] The hyperproliferative diseases for the treatment of which the compounds of the invention can be employed include in particular the group of cancer and tumour diseases. In the context of the present invention, these are understood as meaning, in particular, the following diseases, but without being limited to them: mammary carcinomas and mammary tumours (ductal and lobular forms, also in situ), tumours of the respiratory tract (parvicellular and non-parvicellular carcinoma, bronchial carcinoma), cerebral tumours (e.g. of the brain stem and of the hypothalamus, astrocytoma, medulloblastoma, ependymoma and neuro-ectodermal and pineal tumours), tumours of the digestive organs (oesophagus, stom-

ach, gall bladder, small intestine, large intestine, rectum), liver tumours (including hepatocellular carcinoma, cholangiocellular carcinoma and mixed hepatocellular and cholangiocellular carcinoma), tumours of the head and neck region (larynx, hypopharynx, nasopharynx, oropharynx, lips and oral cavity), skin tumours (squamous epithelial carcinoma, Kaposi sarcoma, malignant melanoma, Merkel cell skin cancer and non-melanomatous skin cancer), tumours of soft tissue (including soft tissue sarcomas, osteosarcomas, malignant fibrous histiocytomas, lymphosarcomas and rhabdomyosarcomas), tumours of the eyes (including intraocular melanoma and retinoblastoma), tumours of the endocrine and exocrine glands (e.g. thyroid and parathyroid glands, pancreas and salivary gland), tumours of the urinary tract (tumours of the bladder, penis, kidney, renal pelvis and ureter) and tumours of the reproductive organs (carcinomas of the endometrium, cervix, ovary, vagina, vulva and uterus in women and carcinomas of the prostate and testicles in men). These also include proliferative blood diseases in solid form and as circulating blood cells, such as lymphomas, leukaemias and myeloproliferative diseases, e.g. acute myeloid, acute lymphoblastic, chronic lymphocytic, chronic myelogenous and hair cell leukaemia, and also AIDS-correlated lymphomas, Hodgkin's lymphomas, non-Hodgkin's lymphomas, cutaneous T-cell lymphomas, Burkitt's lymphomas and lymphomas in the central nervous system.

Preferred Hyperproliferative Diseases for Anti-CA9 Binder-Drug Conjugates

[2930] Hyperproliferative diseases for the treatment of which the compounds of the invention can be preferably employed are CA9-overexpressing tumours, mammary carcinomas and mammary tumours (e.g. ductal and lobular forms, also in situ); tumours of the respiratory tract (e.g. parvicellular and non-parvicellular carcinoma, bronchial carcinoma), including preferably non-parvicellular carcinoma of the lung; cerebral tumours (e.g. of the brain stem and of the hypothalamus, astrocytoma, medulloblastoma, ependymoma and/or neuro-ectodermal and pineal tumours); tumours of the digestive organs (oesophagus, stomach, gall bladder, small intestine, large intestine, rectum), including more preferably stomach tumours and intestinal tumours; liver tumours (including hepatocellular carcinoma, cholangiocellular carcinoma and mixed hepatocellular and cholangiocellular carcinoma); tumours of the head and neck region (e.g. larynx, hypopharynx, nasopharynx, oropharynx, lips, oral cavity, tongue and oesophagus); tumours of the urinary tract (tumours of the bladder, penis, kidney, renal pelvis and ureter), including more preferably tumours of the kidneys and of the bladder; and/or tumours of the reproductive organs (carcinomas of the endometrium, cervix, ovary, vagina, vulva and uterus in women and/or carcinomas of the prostate and testicles in men), including more preferably carcinomas of the cervix and uterus.

Preferred Hyperproliferative Diseases for Anti-EGFR Binder-Drug Conjugates

[2931] Hyperproliferative diseases for the treatment of which the compounds of the invention can be preferably employed are EGFR-overexpressing tumours, respiratory tract tumours (e.g. parvicellular and non-pavicellular carcinomas, bronchial carcinoma), including preferably non-parvicellular carcinoma of the lung; tumours of the digestive

organs (e.g. oesophagus, stomach, gall bladder, small intestine, large intestine, rectum), including especially intestinal tumours; tumours of the endocrine and exocrine glands (e.g. thyroid and parathyroid glands, pancreas and salivary gland), including preferably pancreas; tumours of the head and neck region (e.g. larynx, hypopharynx, nasopharynx, oropharynx, lips, oral cavity, tongue and oesophagus); and/or gliomas.

Preferred Hyperproliferative Diseases for Anti-Mesothelin Binder-Drug Conjugates

[2932] Hyperproliferative diseases for the treatment of which the compounds of the invention can be preferably employed are mesothelin-overexpressing tumours, tumours of the reproductive organs (carcinomas of the endometrium, cervix, ovary, vagina, vulva and uterus in women and/or carcinomas of the prostate and testicles in men), including preferably ovarian carcinomas; tumours of the endocrine and exocrine glands (e.g. thyroid and parathyroid glands, pancreas and salivary gland), including preferably pancreas; respiratory tract tumours (e.g. parvicellular and non-parvicellular carcinoma, bronchial carcinoma), including preferably non-parvicellular carcinoma of the lung; and/or mesotheliomas.

Preferred Hyperproliferative Diseases for Anti-C4.4a Binder-Drug Conjugates

[2933] Hyperproliferative diseases for the treatment of which the compounds of the invention can be preferably employed are C4.4a-overexpressing tumours, squamous epithelial carcinomas (e.g. of the cervix, vulva, vagina, of the anal duct, endometrium, fallopian tube, penis, scrotum, of the oesophagus, breast, of the bladder, of the bile duct, endometrium, uterus and ovary); mammary carcinomas and mammary tumours (e.g. ductal and lobular forms, also in situ); tumours of the respiratory tract (e.g. parvicellular and non-parvicellular carcinoma, bronchial carcinoma), including preferably non-parvicellular carcinoma of the lung, squamous epithelial carcinoma and adenocarcinoma of the lung; tumours of the head and neck region (e.g. larynx, hypopharynx, nasopharynx, oropharynx, lips, oral cavity, tongue and oesophagus, squamous epithelial carcinomas of the head and neck region); tumours of the urinary tract (tumours of the bladder, penis, kidney, renal pelvis and ureter, squamous epithelial carcinomas of the bladder), including more preferably tumours of the kidneys and of the bladder; skin tumours (squamous epithelial carcinoma, Kaposi sarcoma, malignant melanoma, Merkel cell skin cancer and non-melanomatous skin cancer), including more preferably melanomas; tumours of the endocrine and exocrine glands (e.g. thyroid and parathyroid glands, pancreas and salivary gland), including preferably pancreas; tumours of the digestive organs (e.g. oesophagus, stomach, gall bladder, small intestine, large intestine, rectum), including especially colorectal carcinomas; and/or tumours of the reproductive organs (carcinomas of the endometrium, cervix, ovary, vagina, vulva and uterus in women and/or carcinomas of the prostate and testicles in men), including more preferably uterine carcinomas.

[2934] These well-described diseases in humans can also occur with a comparable aetiology in other mammals and can be treated there with the compounds of the present invention.

[2935] In the context of the present invention, the term “treatment” or “treat” encompasses an inhibiting, delaying, stemming, alleviating, attenuating, restricting, reducing, sup-

pressing, repressing or healing of an illness, a complaint, a disease, an injury or a disruption to health, the development, course or progress of such conditions and/or the symptoms of such conditions. The term “therapy” is understood here to be synonymous with the term “treatment”.

[2936] The terms “prevention”, “prophylaxis” or “preclusion” are used synonymously in the context of the present invention, and refer to the avoidance or reduction in the risk of acquiring, experiencing, suffering or having an illness, a complaint, a disease, an injury or a disruption to health, a development or progress of such conditions and/or the symptoms of such conditions.

[2937] The treatment or prevention of an illness, complaint, disease, injury or disruption to health may be partial or complete.

[2938] The present invention thus further provides the use of the compounds of the invention for the treatment and/or prevention of diseases, in particular the abovementioned diseases.

[2939] The present invention furthermore provides the use of the compounds of the invention for the preparation of a medicament for the treatment and/or prevention of diseases, in particular the abovementioned diseases.

[2940] The present invention furthermore provides the use of the compounds of the invention in a method for the treatment and/or prevention of diseases, in particular the abovementioned diseases.

[2941] The present invention furthermore provides a method for the treatment and/or prevention of diseases, in particular the abovementioned diseases, using an effective amount of at least one of the compounds of the invention.

[2942] The anti-mesothelin binder-drug conjugate of the invention is used preferably for treating cancer in a patient, where the cancer cells of the patient that are to be treated have mesothelin expression. Treatment is administered more preferably to patients whose mesothelin expression in cancer cells is higher than in healthy cells.

[2943] One method of identifying patients who respond advantageously to an anti-mesothelin binder-drug conjugate for the treatment of cancer involves determining the mesothelin expression in cancer cells of the patient. In one embodiment the mesothelin expression is determined by mesothelin gene expression analysis. The skilled person knows of methods for gene expression analysis such as, for example, RNA detection, quantitative or qualitative polymerase chain reaction or fluorescence in situ hybridization (FISH). In another preferred embodiment the mesothelin expression is determined by means of immunohistochemistry with an anti-mesothelin antibody. The immunohistochemistry is carried out preferably on formaldehyde-fixed tissue. The antibody for use in the immunohistochemistry is the same antibody which is also used in the conjugate. The antibody for use in the immunohistochemistry is a second antibody which—preferably specifically—recognizes the mesothelin target protein.

[2944] The compounds according to the invention can be employed by themselves or, if required, in combination with one or more other pharmacologically active substances, as long as this combination does not lead to undesirable and unacceptable side effects. The present invention furthermore therefore provides medicaments comprising at least one of the compounds of the invention and one or more further drugs, in particular for the treatment and/or prevention of the abovementioned diseases.

[2945] For example, the compounds of the present invention can be combined with known antihyperproliferative, cytostatic or cytotoxic substances for the treatment of cancer diseases. Suitable drugs in the combination which may be mentioned by way of example are as follows:

aldesleukin, alendronic acid, alfaferone, alitretinoin, allopurinol, aloprim, aloxi, altretamine, aminoglutethimide, amifostine, amrubicin, amsacrine, anastrozole, anzmec, aranesp, arglabin, arsenic trioxide, aromasin, 5-azacytidine, azathioprine, BCG or tice-BCG, bestatin, betamethasone acetate, betamethasone sodium phosphate, bexarotene, bleomycin sulphate, broxuridine, bortezomib, busulfan, calcitonin, campath, capecitabine, carboplatin, casodex, cefesone, celmoleukin, cerubidin, chlorambucil, cisplatin, cladribin, clodronic acid, cyclophosphamide, cytarabine, dacarbazine, dactinomycin, daunoxome, decadron, decadron phosphate, delestrogen, denileukin difitox, depomedrol, deslorelin, dexrazoxane, diethylstilbestrol, diflucan, docetaxel, doxifluridine, doxorubicin, dronabinol, DW-166HC, eligard, elitek, ellence, emend, epirubicin, epoetin-alfa, epogen, eptaplatin, ergamisol, estrace, estradiol, estramustine sodium phosphate, ethinylestradiol, ethylol, etidronic acid, etopophos, etoposide, fadrozole, farstone, filgrastim, finasteride, fligastim, floxuridine, fluconazole, fludarabine, 5-fluorodeoxyuridine monophosphate, 5-fluorouracil (5-FU), fluoxymesterone, flutamide, formestane, fosteabine, fotemustine, fulvestrant, gammagard, gemcitabine, gemtuzumab, gleevec, gliadel, goserelin, granisetron hydrochloride, histrelin, hycamtin, hydrocortone, erythro-hydroxynonyladenine, hydroxyurea, ibritumomab tiuxetan, idarubicin, ifosfamide, interferon-alpha, interferon-alpha-2, interferon-alpha-2 α , interferon-alpha-2 β , interferon-alpha-n1, interferon-alpha-n3, interferon-beta, interferon-gamma-1 α , interleukin-2, intron A, iressa, irinotecan, kytril, lentinan sulphate, letrozole, leucovorin, leuprolide, leuprolide acetate, levamisole, levofolic acid calcium salt, levotirodine, levoxyl, lomustine, lonidamine, marinol, mechlorethamine, mecobalamin, medroxyprogesterone acetate, megestrol acetate, melphalan, menest, 6-mercaptopurine, mesna, methotrexate, metvix, miltefosine, minocycline, mitomycin C, mitotane, mitoxantrone, modrenal, myocet, nedaplatin, neulasta, neumega, neupogen, nilutamide, nolvadex, NSC-631570, OCT-43, octreotide, ondansetron hydrochloride, orapred, oxaliplatin, paclitaxel, pediapred, pegaspargase, pegasys, pentostatin, picibanil, pilocarpine hydrochloride, pirarubicin, plicamycin, porfimer sodium, prednimustine, prednisolone, prednisone, premarin, procarbazine, procrit, raltitrexed, rebif, rhenium-186 etidronate, rituximab, roferon-A, romurtide, salagen, sandostatin, sargramostim, semustine, sizofuran, sobuzoxane, solumedrol, streptozocin, strontium-89 chloride, Synthroid, tamoxifen, tamsulosin, tasonermin, tastolactone, taxoter, teceleukin, temozolomide, teniposide, testosterone propionate, testred, thioguanine, thiotepa, thyrotropin, tiludronic acid, topotecan, toremifen, tositumomab, tastuzumab, teosulfan, tretinoin, trexall, trimethylmelamine, trimetrexate, triptorelin acetate, triptorelin pamoate, UFT, uridine, valrubicin, vesnarinone, vinblastine, vincristine, vindesine, vinorelbine, virulizin, zincard, zinostatin-stimalamer, zofran; ABI-007, acolbifen, actimmune, affinitak, aminopterin, arzoxifen, asoprisnil, atamestane, atrasentan, avastin, BAY 43-9006 (sorafenib), CCI-779, CDC-501, celebrex, cetuximab, crisnatol, cyproterone acetate, decitabine, DN-101, doxorubicin-MTC, dSLIM, dutasteride, edotecarin, eflornithine, exatecan, fenretinide, histamine dihydrochloride, histrelin hydrogel

implant, holmium-166 DOTMP, ibandronic acid, interferon-gamma, intron-PEG, ixabepilone, keyhole limpet hemocyanine, L-651582, lanreotide, lasofoxifen, libra, lonafarnib, miproxifen, minodronate, MS-209, liposomal MTP-PE, MX-6, nafarelin, nemorubicin, neovastat, nolatrexed, oblimersen, onko-TCS, osidem, paclitaxel polyglutamate, pamidronate disodium, PN-401, QS-21, quazepam, R-1549, raloxifen, ranpirnas, 13-cis-retic acid, satraplatin, seocalcitol, T-138067, tarceva, taxoprexin, thymosin-alpha-1, tiazofurin, tipifarnib, tirapazamine, TLK-286, toremifen, transMID-107R, valspodar, vaporeotide, vatalanib, verteporfin, vinflunin, Z-100, zoledronic acid and combinations of these.

[2946] In a preferred embodiment, the compounds of the present invention can be combined with antihyperproliferative agents, which can be, by way of example—without this list being conclusive as follows:

aminoglutethimide, L-asparaginase, azathioprine, 5-azacytidine, bleomycin, busulfan, carboplatin, carmustine, chlorambucil, cisplatin, colaspase, cyclophosphamide, cytarabine, dacarbazine, dactinomycin, daunorubicin, diethylstilbestrol, 2',2'-difluorodeoxycytidine, docetaxel, doxorubicin (adriamycin), epirubicin, epothilone and its derivatives, erythro-hydroxynonyladenine, ethinylestradiol, etoposide, fludarabine phosphate, 5-fluorodeoxyuridine, 5-fluorodeoxyuridine monophosphate, 5-fluorouracil, fluoxymesterone, flutamide, hexamethylmelamine, hydroxyurea, hydroxyprogesterone caproate, idarubicin, ifosfamide, interferon, irinotecan, leucovorin, lomustine, mechlorethamine, medroxyprogesterone acetate, megestrol acetate, melphalan, 6-mercaptopurine, mesna, methotrexate, mitomycin C, mitotane, mitoxantrone, paclitaxel, pentostatin, N-phosphonoacetyl L-aspartate (PALA), plicamycin, prednisolone, prednisone, procarbazine, raloxifen, semustine, streptozocin, tamoxifen, teniposide, testosterone propionate, thioguanine, thiotepa, topotecan, trimethylmelamine, uridine, vinblastine, vincristine, vindesine and vinorelbine.

[2947] The compounds of the invention can also be combined in a very promising manner with biological therapeutics such as antibodies (e.g. avastin, rituxan, erbitux, herceptin). The compounds of the invention can also achieve positive effects in combination with therapies directed against angiogenesis, such as, for example, with avastin, axitinib, recentin, regorafenib, sorafenib or sunitinib. Combinations with inhibitors of the proteasome and of mTOR and also with antihormones and steroidal metabolic enzyme inhibitors are likewise particularly suitable because of their favourable profile of side effects.

[2948] Generally, the following aims can be pursued with the combination of compounds of the present invention with other agents having a cytostatic or cytotoxic action:

[2949] an improved activity in slowing down the growth of a tumour, in reducing its size or even in its complete elimination compared with treatment with an individual drug;

[2950] the possibility of employing the chemotherapeutics used in a lower dosage than in monotherapy;

[2951] the possibility of a more tolerable therapy with few side effects compared with individual administration;

[2952] the possibility of treatment of a broader spectrum of tumour diseases;

[2953] the achievement of a higher rate of response to the therapy;

[2954] a longer survival time of the patient compared with present-day standard therapy.

[2955] The compounds according to the invention can moreover also be employed in combination with radiotherapy and/or surgical intervention.

[2956] The present invention furthermore provides medicaments which comprise at least one compound of the invention, conventionally together with one or more inert, non-toxic, pharmaceutically suitable excipients, and the use thereof for the abovementioned purposes.

[2957] The compounds of the invention can act systemically and/or locally. They can be administered in a suitable manner for this purpose, such as for example parenterally, possibly by means of inhalation, or as an implant or stent.

[2958] The compounds of the invention can be administered in suitable administration forms for these administration routes.

[2959] Parenteral administration can be effected with bypassing of an absorption step (e.g. intravenously, intraarterially, intracardially, intraspinally or intralumbally) or with inclusion of an absorption (e.g. intramuscularly, subcutaneously, intracutaneously, percutaneously or intraperitoneally). Administration forms which are suitable for parenteral administration include injection and infusion formulations in the form of solutions, suspensions, emulsions or lyophilizates. Parenteral administration is preferred, in particular intravenous administration.

[2960] In general, it has proved advantageous in the case of parenteral administration to administer amounts of from about 0.001 to 1 mg/kg, preferably about 0.01 to 0.5 mg/kg of body weight to achieve effective results.

[2961] Nevertheless it may be necessary to deviate from the amounts mentioned, and in particular depending on the body weight, administration route, individual behaviour towards the active compound, nature of the formulation and point of time or interval at which administration takes place. Thus in some cases it may be sufficient to manage with less than the abovementioned minimum amount, while in other cases the upper limit mentioned must be exceeded. In the case where relatively large amounts are administered, it may be advisable to distribute these into several individual doses over the day.

[2962] The following examples illustrate the invention. The invention is not limited to the examples.

[2963] The percentage figures in the following tests and examples are percentages by weight, unless stated otherwise; parts are parts by weight. Solvent ratios, dilution ratios and concentration data of liquid/liquid solutions in each case relate to the volume.

A. EXAMPLES

Abbreviations and Acronyms

- [2964] A431NS human tumour cell line
- [2965] A549 human tumour cell line
- [2966] ABCB1 ATP-binding cassette sub-family B member 1 (synonym for P-gp and MDR1)
- [2967] abs. absolute
- [2968] ADC antibody-drug-conjugate
- [2969] Ac acetyl
- [2970] aq. aqueous, aqueous solution
- [2971] ATP adenosine triphosphate
- [2972] BCRP breast cancer resistance protein, an efflux transporter
- [2973] Boc tert-butoxycarbonyl
- [2974] br. broad (in NMR)
- [2975] Ex. example
- [2976] ca. circa, approximately
- [2977] CAIX carboanhydrase IX
- [2978] CI chemical ionization (in MS)
- [2979] d doublet (in NMR)
- [2980] d day(s)
- [2981] TLC thin-layer chromatography
- [2982] DCI direct chemical ionization (in MS)
- [2983] dd doublet of a doublet (in NMR)
- [2984] DMAP 4-N,N-dimethylaminopyridine
- [2985] DME 1,2-dimethoxyethane
- [2986] DMEM Dulbecco's modified eagle medium (standardized nutrient medium for cell culture)
- [2987] DMF N,N-dimethylformamide
- [2988] DMSO dimethyl sulphoxide
- [2989] DPBS, D-PBS, Dulbecco's phosphate-buffered saline solution
- [2990] PBS PBS=DPBS=D-PBS, pH 7.4, from Sigma, No. D8537
- [2991] Composition:
- [2992] 0.2 g KCl
- [2993] 0.2 g KH_2PO_4 (anhydrous)
- [2994] 8.0 g NaCl
- [2995] 1.15 g Na_2HPO_4 (anhydrous)
- [2996] make up to 1 l with H_2O
- [2997] dt doublet of a triplet (in NMR)
- [2998] DTT DL-dithiothreitol
- [2999] EDC N'-(3-dimethylaminopropyl)-N-ethylcarbodiimide hydrochloride
- [3000] EGFR epidermal growth factor receptor
- [3001] EI electron impact ionization (in MS)
- [3002] ELISA enzyme-linked immunosorbent assay
- [3003] eq. equivalent(s)
- [3004] ESI electrospray ionization (in MS)
- [3005] ESI-MicroTofq ESI-MicroTofq (name of the mass spectrometer, with T_{of} =time of flight and q =quadrupole)
- [3006] FCS foetal calf serum
- [3007] Fmoc (9H-fluoren-9-ylmethoxy)carbonyl
- [3008] sat. saturated
- [3009] GTP guanosine 5'-triphosphate
- [3010] h hour(s)
- [3011] HATU O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate
- [3012] HCT-116 human tumour cell line
- [3013] HEPES 4-(2-hydroxyethyl)piperazine-1-ethanesulphonic acid
- [3014] HOAc acetic acid
- [3015] HOBt 1-hydroxy-1H-benzotriazole hydrate
- [3016] HOSu N-hydroxysuccinimide
- [3017] HPLC high-pressure, high-performance liquid chromatography
- [3018] HT29 human tumour cell line
- [3019] IC_{50} half-maximum inhibitory concentration
- [3020] i.m. intramuscular, administration into the muscle
- [3021] i.v. intravenous, administration into the vein
- [3022] conc. concentrated
- [3023] LC-MS liquid chromatography-coupled mass spectrometry
- [3024] LLC-PK1 cells Lewis lung carcinoma pork kidney cell line
- [3025] L-MDR human MDR1 transfected LLC-PK1 cells
- [3026] m multiplet (in NMR)
- [3027] MDR1 multidrug resistance protein 1

- [3028] min minute(s)
- [3029] MS mass spectrometry
- [3030] MTT 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide
- [3031] NCI-H292 human tumour cell line
- [3032] NCI-H520 human tumour cell line
- [3033] NMM N-methylmorpholine
- [3034] NMP N-methyl-2-pyrrolidinone
- [3035] NMR nuclear magnetic resonance spectrometry
- [3036] NMRI mouse strain, originating from Naval Medical Research Institute (NMRI)
- [3037] Nude mice experimental animals
- [3038] NSCLC non-small cell lung cancer (non-parvicel-lular bronchial carcinoma)
- [3039] PBS phosphate-buffered saline solution
- [3040] Pd/C palladium on activated carbon
- [3041] P-gp P-glycoprotein, a transporter protein
- [3042] PNGaseF enzyme for sugar elimination
- [3043] quant. quantitative (for yield)
- [3044] quart quartet (in NMR)
- [3045] quint quintet (in NMR)
- [3046] R_f retention index (for TLC)
- [3047] RT room temperature
- [3048] R_t retention time (for HPLC)
- [3049] s singlet (in NMR)
- [3050] s.c. subcutaneous, administration beneath the skin
- [3051] SCC-4 human tumour cell line
- [3052] SCC-9 human tumour cell line
- [3053] SCID mice experimental mice with a severe combined immunodeficiency
- [3054] t triplet (in NMR)
- [3055] tert tertiary
- [3056] TFA trifluoroacetic acid
- [3057] THF tetrahydrofuran
- [3058] UV ultraviolet spectrometry
- [3059] v/v volume to volume ratio (of a solution)
- [3060] Z benzyloxycarbonyl

HPLC and LC-MS Methods:

Method 1 (LC-MS):

[3061] Instrument: Waters Acquity SQD UPLC System; column: Waters Acquity UPLC HSS T3 1.8 μ 50 mm $\times$ 1 mm; eluent A: 1 l water+0.25 ml 99% strength formic acid, eluent B: 1 l acetonitrile+0.25 ml 99% strength formic acid; gradient: 0.0 min 90% A $\rightarrow$ 1.2 min 5% A $\rightarrow$ 2.0 min 5% A; flow rate: 0.40 ml/min; oven: 50° C.; UV detection: 210-400 nm.

Method 2 (LC-MS):

[3062] Instrument: Micromass QuattroPremier with Waters UPLC Acquity; column: Thermo Hypersil GOLD 1.9 μ 50 mm $\times$ 1 mm; eluent A: 1 l water+0.5 ml 50% strength formic acid, eluent B: 1 l acetonitrile+0.5 ml 50% strength formic acid; gradient: 0.0 min 90% A $\rightarrow$ 0.1 min 90% A $\rightarrow$ 1.5 min 10% A $\rightarrow$ 2.2 min 10% A; flow rate: 0.33 ml/min; oven: 50° C.; UV detection: 210 nm.

Method 3 (LC-MS):

[3063] Instrument: Micromass Quattro Micro MS with HPLC Agilent Series 1100; column: Thermo Hypersil GOLD 3 μ 20 mm $\times$ 4 mm; eluent A: 1 l water+0.5 ml 50% strength formic acid, eluent B: 1 l acetonitrile+0.5 ml 50% strength formic acid; gradient: 0.0 min 100% A $\rightarrow$ 3.0 min 10% A $\rightarrow$ 4.0

min 10% A $\rightarrow$ 4.01 min 100% A (flow rate 2.5 ml/min) $\rightarrow$ 5.00 min 100% A; oven: 50° C.; flow rate: 2 ml/min; UV detection: 210 nm.

Method 4 (LC-MS):

[3064] MS instrument: Micromass ZQ; HPLC instrument: HP 1100 Series; UV DAD; column: Phenomenex Gemini 3 μ 30 mm $\times$ 3.00 mm; eluent A: 1 l water+0.5 ml 50% strength formic acid, eluent B: 1 l acetonitrile+0.5 ml 50% strength formic acid; gradient: 0.0 min 90% A $\rightarrow$ 2.5 min 30% A $\rightarrow$ 3.0 min 5% A $\rightarrow$ 4.5 min 5% A; flow rate: 0.0 min 1 ml/min $\rightarrow$ 2.5 min/3.0 min/4.5 min 2 ml/min; oven: 50° C.; UV detection: 210 nm.

Method 5 (HPLC):

[3065] Instrument: HP 1090 Series II; column: Merck Chromolith SpeedROD RP-18e, 50 mm $\times$ 4.6 mm; preliminary column: Merck Chromolith Guard Cartridge Kit RP-18e, 5 mm $\times$ 4.6 mm; injection volume: 5 μ l; eluent A: 70% HClO₄ in water (4 ml/litre), eluent B: acetonitrile; gradient: 0.00 min 20% B $\rightarrow$ 0.50 min 20% B $\rightarrow$ 3.00 min 90% B $\rightarrow$ 3.50 min 90% B $\rightarrow$ 3.51 min 20% B $\rightarrow$ 4.00 min 20% B; flow rate: 5 ml/min; column temperature: 40° C.

Method 6 (HPLC):

[3066] Instrument: Waters 2695 with DAD 996; column: Merck Chromolith SpeedROD RP-18e, 50 mm $\times$ 4.6 mm; ord. No.: 1.51450.0001, preliminary column: Merck Chromolith Guard Cartridge Kit RP-18e, 5 mm $\times$ 4.6 mm; ord. No.: 1.51470.0001, eluent A: 70% HClO₄ in water (4 ml/litre), eluent B: acetonitrile; gradient: 0.00 min 5% B $\rightarrow$ 0.50 min 5% B $\rightarrow$ 3.00 min 95% B $\rightarrow$ 4.00 min 95% B; flow rate: 5 ml/min.

Method 7 (LC-MS):

[3067] MS instrument: Waters ZQ; HPLC instrument: Agilent 1100 Series; UV DAD; column: Thermo Hypersil GOLD 3 μ 20 mm $\times$ 4 mm; eluent A: 1 l water+0.5 ml 50% strength formic acid, eluent B: 1 l acetonitrile+0.5 ml 50% strength formic acid; gradient: 0.0 min 100% A $\rightarrow$ 3.0 min 10% A $\rightarrow$ 4.0 min 10% A $\rightarrow$ 4.1 min 100% A (flow rate 2.5 ml/min); oven: 55° C.; flow rate: 2 ml/min; UV detection: 210 nm.

Method 8 (LC-MS):

[3068] MS instrument: Waters ZQ; HPLC instrument: Agilent 1100 Series; UV DAD; column: Thermo Hypersil GOLD 3 μ 20 mm $\times$ 4 mm; eluent A: 1 l water+0.5 ml 50% strength formic acid, eluent B: 1 l acetonitrile+0.5 ml 50% strength formic acid; gradient: 0.0 min 100% A $\rightarrow$ 2.0 min 60% A $\rightarrow$ 2.3 min 40% A $\rightarrow$ 3.0 min 20% A $\rightarrow$ 4.0 min 10% A $\rightarrow$ 4.2 min 100% A (flow rate 2.5 ml/min); oven: 55° C.; flow rate: 2 ml/min; UV detection: 210 nm.

Method 9 (LC-MS):

[3069] Instrument: Waters Acquity SQD UPLC System; column: Waters Acquity UPLC HSS T3 1.8 μ 50 mm $\times$ 1 mm; eluent A: 1 l water+0.25 ml 99% strength formic acid, eluent B: 1 l acetonitrile+0.25 ml 99% strength formic acid; gradient: 0.0 min 95% A $\rightarrow$ 6.0 min 5% A $\rightarrow$ 7.5 min 5% A; oven: 50° C.; flow rate: 0.35 ml/min; UV detection: 210-400 nm.

Method 10 (HPLC):

[3070] Instrument: Agilent 1200 Series; column: Agilent Eclipse XDB-C18 5 μ 4.6 mm $\times$ 150 mm; preliminary column: Phenomenex KrudKatcher Disposable Pre-Column; injection volume: 5 μ l; eluent A: 1 l water+0.01% trifluoroacetic acid; eluent B: 1 l acetonitrile+0.01% trifluoroacetic acid; gradient: 0.00 min 10% B $\rightarrow$ 1.00 min 10% B $\rightarrow$ 1.50 min 90% B $\rightarrow$ 5.5 min 10% B; flow rate: 2 ml/min; column temperature: 30 $^{\circ}$ C.

[3071] For all reactants or reagents whose preparation is not explicitly described below, they were obtained commercially from generally available sources. For all other reactants or reagents whose preparation is likewise not described below, and which were not available commercially or were obtained from sources which are not generally available, a reference is given to the published literature in which their preparation is described.

Method 11 (LC-MS):

[3072] Instrument: Waters ACQUITY SQD UPLC System; column: Waters Acquity UPLC HSS T3 1.8 μ 130 $\times$ 2 mm; eluent A: 1 l water+0.25 ml 99% strength formic acid, eluent B: 1 l acetonitrile+0.25 ml 99% strength formic acid; gradient: 0.0 min 90% A $\rightarrow$ 1.2 min 5% A $\rightarrow$ 2.0 min 5% A oven: 50 $^{\circ}$ C.; flow rate: 0.60 ml/min; UV detection: 208-400 nm.

Method 12 (HPLC):

[3073] Instrument: Agilent 1200 Series with column oven and DAD; column: Merck Chromolith SpeedROD RP-18e, 50 mm $\times$ 4.6 mm; ord. No.: 1.51450.0001; preliminary column: Merck Chromolith Guard Cartridge Kit RP-18e, 5 mm $\times$ 4.6 mm; ord. No.: 1.51470.0001; eluent A: 70% HClO₄ in water (4 ml/litre), eluent B: acetonitrile; gradient: 0.00 min 5% B $\rightarrow$ 0.50 min 5% B $\rightarrow$ 3.00 min 95% B $\rightarrow$ 4.00 min 95% B; flow rate: 5 ml/min; column temperature: 30 $^{\circ}$ C.

Method 13 (LC-MS):


[3074] MS instrument: Waters (Micromass) Quattro Micro; Instrument HPLC: Agilent 1100 Series; column: YMC-Triart C18 3 μ 50 $\times$ 3 mm; eluent A: 1 l water+0.01 mol ammonium carbonate, eluent B: 1 l acetonitrile; gradient: 0.0 min 100% A $\rightarrow$ 2.75 min 5% A $\rightarrow$ 4.5 min 5% A; oven: 40 $^{\circ}$ C.; flow rate: 1.25 ml/min; UV detection: 210 nm.

Starting Compounds and Intermediates

Starting Compound 1

(2R,3R)-3-[(2S)-1-(tert-butoxycarbonyl)pyrrolidin-2-yl]-3-methoxy-2-methylpropanoic acid (Boc-dolaisoleucine)

[3075]




[3076] The title compound can be prepared in various ways according to literature methods; see, for example, Pettit et al., *Synthesis* 1996, 719; Shioiri et al., *Tetrahedron Lett.* 1991, 32, 931; Shioiri et al., *Tetrahedron* 1993, 49, 1913; Koga et al., *Tetrahedron Lett.* 1991, 32, 2395; Vidal et al., *Tetrahedron* 2004, 60, 9715; Poncet et al., *Tetrahedron* 1994, 50, 5345. It was prepared either as the free acid or as a 1:1 salt with dicyclohexylamine.

Starting Compound 2a

tert-butyl (3R,4S,5S)-3-methoxy-5-methyl-4-(methylamino)heptanoate hydrochloride (dolaisoleucine-OtBuxHCl)

[3077]




[3078] The title compound can be prepared in various ways according to literature methods; see, for example, Pettit et al., *J. org. Chem.* 1994, 59, 1796; Koga et al., *Tetrahedron Lett.* 1991, 32, 2395; Shioiri et al., *Tetrahedron Lett.* 1991, 32, 931; Shioiri et al., *Tetrahedron* 1993, 49, 1913.

Starting Compound 2b

tert-butyl (3R,4S,5S)-3-methoxy-5-methyl-4-(methylamino)heptanoate (dolaisoleucine-O^tBu)

[3079]




[3080] The compound was prepared in analogy to starting compound 2a, except that the hydrogenation was performed without addition of 1N hydrochloric acid.

Starting Compound 3

α -(tert-butoxycarbonyl)-N-hydroxy-L-phenylalanine
namide

[3081]



[3082] The title compound was prepared by the literature method (A. Ritter et al., *J. org. Chem.* 1994, 59, 4602).


[3083] Yield: 750 mg (75% of theory)

[3084] LC-MS (Method 3): R_f =1.67 min; MS (ESIpos): m/z =281 (M+H)⁺.

Starting Compound 4

1,2-oxazolidine hydrochloride

[3085]




[3086] The title compound can be prepared by literature methods (see, for example, H. King, *J. Chem. Soc.* 1942, 432); it is also commercially available.

Starting Compound 5

1,2-oxazinane hydrochloride

[3087]




[3088] The title compound can be prepared by literature methods (see, for example, H. King, *J. Chem. Soc.* 1942, 432).

Starting Compound 6

2-oxa-3-azabicyclo[2.2.2]oct-5-ene

[3089]




[3090] The title compound can be prepared in Boc-protected form by the literature method (see, for example, C. Johnson et al., *Tetrahedron Lett.* 1998, 39, 2059); the deprotection was effected in a customary manner by treatment with trifluoroacetic acid and subsequent neutralization.

[3091] Yield: 149 mg (89% of theory)

Starting Compound 7

tert-butyl
(1S,2R)-1-(hydroxycarbonyl)-2-phenylcyclopropyl
carbamate

[3092]



[3093] The title compound was prepared by a literature method (A. Ritter et al., *J. org. Chem.* 1994, 59, 4602) proceeding from commercially available (1S,2R)-1-[(tert-butoxycarbonyl)amino]-2-phenylcyclopropanecarboxylic acid (C. Cativiela et al., *Chirality* 1999, 11, 583).


[3094] Yield: 339 mg (59% of theory)

[3095] LC-MS (Method 1): R_f =0.82 min; MS (ESIpos): m/z =293 (M+H)⁺.

Intermediate 1

tert-butyl (3R,4S,5S)-4-[[N-[(benzyloxy)carbonyl]-valyl](methyl)amino]-3-methoxy-5-methylheptanoate

[3096]



[3097] 10.65 g (41.058 mmol) of tert-butyl (3R,4S,5S)-3-methoxy-5-methyl-4-(methylamino)heptanoate (starting compound 2b) were taken up in 250 ml of dichloromethane and the solution was cooled to -10°C . Then, while stirring, 10.317 g (41.058 mmol) of N-[(benzyloxy)carbonyl]-L-valine, 16.866 g (61.586 mmol) of 2-bromo-1-ethylpyridinium tetrafluoroborate (BEP) and 28.6 ml of N,N-diisopropylethylamine were added, and the mixture was subsequently stirred at RT for 20 h. The reaction mixture was then diluted with dichloromethane and shaken twice with saturated sodium chloride solution, dried over sodium sulphate, filtered and concentrated. The residue was purified by flash chromatography on silica gel with 4:1 petroleum ether/ethyl acetate as the eluent. The corresponding fractions were concentrated and the residue was dried under high vacuum overnight. 10.22 g (51% of theory) of the title compound were obtained as a yellowish oil.

[3098] HPLC (Method 5): $R_f=2.3$ min;

[3099] LC-MS (Method 2): $R_f=1.59$ min; MS (ESIpos): $m/z=493$ (M+H) $^{+}$.

[3101] 500 mg (1 mmol) of tert-butyl (3R,4S,5S)-4-[[N-[(benzyloxy)carbonyl]-L-valyl](methyl)amino]-3-methoxy-5-methylheptanoate (intermediate 1) were dissolved in 50 ml of methanol and, after addition of 100 mg of 10% palladium on activated carbon, hydrogenated under standard hydrogen pressure at RT for 1 h. The catalyst was then filtered off and the solvent was removed under reduced pressure. This gave 370 mg (quant.) of the title compound as a virtually colourless oil.


[3102] HPLC (Method 5): $R_f=1.59$ min;

[3103] LC-MS (Method 1): $R_f=0.74$ min; MS (ESIpos): $m/z=359$ (M+H) $^{+}$.

Intermediate 3

N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-tert-butoxy-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide


[3104]



Intermediate 2

tert-butyl (3R,4S,5S)-3-methoxy-5-methyl-4-[methyl(L-valyl)amino]heptanoate

[3100]



[3105] 4.64 g (13.13 mmol) of N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valine were dissolved in 20 ml of DMF and admixed successively with 4.28 g (11.94 mmol) of tert-butyl (3R,4S,5S)-3-methoxy-5-methyl-4-[methyl(L-valyl)amino]heptanoate (Intermediate 2), 2.75 g (14.33 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 2.2 g (14.33 mmol) of 1-hydroxy-1H-benzotriazole hydrate. The mixture was stirred at RT overnight. The reaction mixture was then poured into a mixture of semisaturated aqueous ammonium chloride solution and ethyl acetate. The organic phase was removed, washed successively with saturated sodium hydrogencarbonate solution and saturated sodium chloride solution, dried over magnesium sulphate, filtered and concentrated. The residue was used directly in the next stage, without further purification.

[3106] Yield: 9.1 g (quant., 60% purity)


[3107] HPLC (Method 5): $R_f=2.7$ min;

[3108] LC-MS (Method 2): $R_f=1.99$ min; MS (ESIpos): $m/z=694$ (M+H) $^{+}$.

Intermediate 4

N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide

[3109]



ride, 382 mg (2.8 mmol) of 1-hydroxy-1H-benzotriazole hydrate and 731 mg (5.7 mmol) of N,N-diisopropylethylamine. The mixture was stirred at RT overnight. The reaction mixture was then poured into a mixture of semisaturated aqueous ammonium chloride solution and ethyl acetate. The organic phase was removed, washed successively with saturated sodium hydrogencarbonate solution and saturated

[3110] 9.1 g of the crude product N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-tert-butoxy-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 3) were taken up in 56.6 ml of dichloromethane, 56.6 ml of trifluoroacetic acid were added, and the mixture was stirred at RT for 2 h. Subsequently, the reaction mixture was concentrated under reduced pressure and the remaining residue was purified by flash chromatography, using dichloromethane, 3:1 dichloromethane/ethyl acetate and 15:5:0.5 dichloromethane/ethyl acetate/methanol as eluent. After purification of the corresponding fractions and concentration, 5.8 g (86% of theory) of the title compound were obtained as a colourless foam.


[3111] HPLC (Method 5): R_f =2.2 min;

[3112] LC-MS (Method 1): R_f =1.3 min; MS (ESIpos): m/z =638 (M+H)⁺.

Intermediate 5

tert-butyl (2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl carbamate

[3113]



[3114] 500 mg (1.9 mmol) of N-(tert-butoxycarbonyl)-L-phenylalanine were dissolved in 10 ml of DMF and admixed successively with 466 mg (3.8 mmol) of 1,2-oxazinan hydrochloride (Starting Compound 5), 433 mg (2.3 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride,

sodium chloride solution, dried over magnesium sulphate, filtered and concentrated. 620 mg (98% of theory) of the title compound were obtained.


[3115] HPLC (Method 5): R_f =1.8 min;

[3116] LC-MS (Method 2): R_f =1.62 min; MS (ESIpos): m/z =235 (M-C₄H₈-CO₂+H)⁺.

Intermediate 6

(2S)-2-amino-1-(1,2-oxazinan-2-yl)-3-phenylpropan-1-one trifluoroacetate

[3117]



[3118] 620 mg (1.85 mmol) of tert-butyl (2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl carbamate (Intermediate 5) were taken up in 5 ml of dichloromethane, 10 ml of trifluoroacetic acid were added and the mixture was stirred at RT for 30 min. Subsequently, the reaction mixture was concentrated under reduced pressure and the remaining residue was lyophilized from water/acetonitrile. In this way, 779 mg (91% of theory) of the title compound were obtained as a colourless foam.


[3119] HPLC (Method 5): R_f =0.45 min;

[3120] LC-MS (Method 3): R_f =1.09 min; MS (ESIpos): m/z =235 (M+H)⁺.

Intermediate 7

(2R,3R)-3-methoxy-2-methyl-N-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]-3-[(2S)-pyrrolidin-2-yl]propanamide trifluoroacetate

[3121]



[3122] 360 mg (1.25 mmol) of (2R,3R)-3-[(2S)-1-(tert-butoxycarbonyl)pyrrolidin-2-yl]-3-methoxy-2-methylpropanoic acid (Starting Compound 1) were taken up in 10 ml of DMF and admixed successively with 579.2 mg (1.25 mmol) of (2S)-2-amino-1-(1,2-oxazinan-2-yl)-3-phenylpropan-1-one trifluoroacetate (Intermediate 6), 714.5 mg (1.88 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) and 655 μ l of N,N-diisopropylethylamine. The mixture was stirred at RT for 16 h. The reaction mixture was then concentrated, and the residue was taken up in ethyl acetate and extracted by shaking first with 5% aqueous citric acid solution, then with 5% aqueous sodium hydrogencarbonate solution and subsequently with saturated sodium chloride solution. The organic phase was concentrated and the residue was purified by flash chromatography on silica gel with 16:4 dichloromethane/methanol as the eluent. The corresponding fractions were combined and the solvent was removed under reduced pressure. After the residue had been dried under high vacuum, 503.5 mg (74% of theory) of the Boc-protected intermediate tert-butyl (2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidine-1-carboxylate were obtained.

[3123] HPLC (Method 12): R_f =2.0 min;

[3124] LC-MS (Method 1): R_f =1.12 min; MS (ESIpos): m/z =504 (M+H)⁺.

[3125] 503 mg (1 mmol) of this intermediate were taken up in 20 ml of dichloromethane, 10 ml of trifluoroacetic acid were added, and the mixture was stirred at RT for 30 min. Subsequently, the reaction mixture was concentrated under reduced pressure and redistilled with dichloromethane. The remaining residue was precipitated from ethyl acetate with n-pentane, and the solvent was decanted off. The residue thus obtained was dissolved in water and extracted by shaking with ethyl acetate, and the aqueous phase was subsequently lyophilized. In this way, 462 mg (89% of theory) of the title compound were obtained as a colourless foam.


[3126] HPLC (Method 12): R_f =1.53 min;

[3127] LC-MS (Method 11): R_f =0.57 min; MS (ESIpos): m/z =404 (M+H)⁺.

Intermediate 8

N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide

[3128]



[3129] 51 mg (0.08 mmol) of N-[(9H-fluoren-9-yl-methoxy)carbonyl]-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 4) were dissolved in 10 ml of DMF, and 0.5 ml of piperidine was added. After stirring at RT for 10 min, the reaction mixture was concentrated under reduced pressure and the residue was stirred with diethyl ether. The insoluble constituents were filtered off and washed repeatedly with diethyl ether. Then the filter residue was taken up in 5 ml of dioxane/water and the solution was adjusted to pH 11 with 1 N sodium hydroxide solution. Under ultrasound treatment, a total of 349 mg (1.6 mmol) of di-tert-butyl dicarbonate were added in several portions, in the course of which the pH of the solution was kept at 11. After the reaction had ended, the dioxane was evaporated off and the aqueous solution was adjusted to a pH of 2-3 with citric acid. The mixture was extracted twice with 50 ml each time of ethyl acetate. The organic phases were combined, dried over magnesium sulphate and concentrated under reduced pressure. The residue was taken up in diethyl ether and the oil of the title compound was precipitated with pentane. The solvent was removed by decantation. The residue was digested several times more with pentane and finally dried under high vacuum. 40 mg (97% of theory) of the title compound were thus obtained.


[3130] HPLC (Method 6): R_f =2.2 min;

[3131] LC-MS (Method 2): R_f =1.32 min; MS (ESIpos): m/z =516 (M+H)⁺.

Intermediate 9

tert-butyl (2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidine-1-carboxylate

[3132]



[3133] The title compound was prepared in analogy to the synthesis of Intermediates 5, 6 and 7 over three stages, by coupling of commercially available (1S,2R)-1-[(tert-butoxycarbonyl)amino]-2-phenylcyclopropanecarboxylic acid with 1,2-oxazinane hydrochloride (Starting Compound 5), subsequent deprotection with trifluoroacetic acid and coupling with Starting Compound 1. The end product was purified by preparative HPLC.


[3134] HPLC (Method 5): R_f =2.12 min;

[3135] LC-MS (Method 2): R_f =1.25 min; MS (ESIpos): m/z =516 (M+H)⁺.

Intermediate 10

N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3136]



[3137] 315 mg (0.494 mmol) of N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 4) were dissolved in 12 ml of DMF, and admixed with 104 mg (0.543 mmol) of 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride and 83 mg (0.543 mmol) of 1-hydroxy-1H-benzotriazole hydrate, and the mixture was stirred at RT for 90 min. Subsequently, 112 μ l of N,N-diisopropylethylamine and 149 mg (0.494 mmol) of (2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoic acid trifluoroacetate, which had been prepared beforehand from Starting Compound 1 by elimination of the Boc protecting group by means of trifluoroacetic acid, were added. The mixture was stirred at RT for 2 h and then concentrated under high vacuum. The remaining residue was purified twice by preparative HPLC. 140 mg (35% of theory) of the title compound were obtained in the form of a colourless foam.


[3138] HPLC (Method 5): R_f =2.40 min;

[3139] LC-MS (Method 1): R_f =1.38 min; MS (ESIpos): m/z =807 (M+H)⁺.

Intermediate 11

N-[(benzyloxy)carbonyl]-N-methyl-L-threonyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide

[3140]



[3141] First, N-[(benzyloxy)carbonyl]-N-methyl-L-threonine was released from 237 mg (0.887 mmol) of its dicyclohexylamine salt thereof by taking it up in ethyl acetate and extractive shaking with 5% aqueous sulphuric acid. The organic phase was dried over magnesium sulphate, filtered and concentrated. The residue was taken up in 16 ml of DMF and admixed successively with 365 mg (1 mmol) of tert-butyl

(3R,4S,5S)-3-methoxy-5-methyl-4-[methyl(L-valyl)amino]heptanoate (Intermediate 2), 185 mg (0.967 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 148 mg (0.967 mmol) of 1-hydroxy-1H-benzotriazole hydrate. The mixture was stirred at RT for 2 h. The reaction mixture was then poured into a mixture of semisaturated aqueous ammonium chloride solution and ethyl acetate. The organic phase was removed, washed successively with saturated sodium hydrogencarbonate solution and saturated sodium chloride solution, dried over magnesium sulphate, filtered and concentrated. The residue was purified by preparative HPLC. 283 mg (53% of theory) of the tert-butyl ester intermediate N-[(benzyloxy)carbonyl]-N-methyl-L-threonyl-N-[(3R,4S,5S)-1-tert-butoxy-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were thus obtained.

[3142] HPLC (Method 5): R_f =2.17 min.

[3143] 283 mg (0.466 mmol) of this intermediate were taken up in 5 ml of dichloromethane, 5 ml of anhydrous trifluoroacetic acid were added, and the mixture was stirred at RT for 2 h. Subsequently, the reaction mixture was concentrated under high vacuum and the remaining residue was purified by means of preparative HPLC. This gave 156 mg (61% of theory) of the title compound as a colourless foam.


[3144] HPLC (Method 5): R_f =1.50 min;

[3145] LC-MS (Method 2): R_f =1.09 min; MS (ESIpos): m/z =552 (M+H)⁺.

Intermediate 12

benzyl N-[(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoyl]-L-phenylalaninate trifluoroacetate

[3146]



[3147] In the first step, Starting Compound 1 was released from 600 mg (1.28 mmol) of the corresponding dicyclohexylammonium salt by dissolving the salt in 100 ml of ethyl acetate and extractive shaking, first with 50 ml of 0.5% sulphuric acid and then with saturated sodium chloride solution. Then the organic phase was dried over magnesium sulphate, filtered, concentrated and reacted immediately with benzyl L-phenylalaninate in analogy to the synthesis of Intermediate 7, and then deprotected.

[3148] Yield: 650 mg (94% over 2 stages)


[3149] HPLC (Method 6): R_f =1.76 min;

[3150] LC-MS (Method 2): R_f =1.68 min; MS (ESIpos): m/z =425 (M+H)⁺.

Intermediate 13

benzyl (βS)—N-[(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoyl]-β-methyl-L-phenylalaninate trifluoroacetate

[3151]



[3152] First, (2R,3R)-3-[(2S)-1-(tert-butoxycarbonyl)pyrrolidin-2-yl]-3-methoxy-2-methylpropanoic acid was released from 351 mg (0.75 mmol) of the dicyclohexylamine salt (Starting Compound 1) by taking it up in ethyl acetate and extractive shaking with aqueous 5% potassium hydrogensulphate solution. The organic phase was dried over magnesium sulphate, filtered and concentrated. The residue was taken up in 10 ml of DMF and admixed successively with 373 mg (0.75 mmol) of benzyl (βS)-β-methyl-L-phenylalaninate trifluoroacetate [prepared from commercially available (βS)—N-(tert-butoxycarbonyl)-β-methyl-L-phenylalanine by EDC/DMAP-mediated esterification with benzyl alcohol and subsequent detachment of the Boc protecting group with trifluoroacetic acid], 428 mg (1.125 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) and 392 μl of N,N-diisopropylethylamine. The mixture was stirred at RT for 20 h. The reaction mixture was then poured onto a mixture of semisaturated aqueous ammonium chloride solution and ethyl acetate. The organic phase was removed, washed successively with saturated sodium hydrogencarbonate solution and saturated sodium chloride solution, and subsequently concentrated. The residue was purified by means of preparative HPLC. This gave 230 mg (57% of theory) of the Boc-protected intermediate benzyl (βS)—N-[(2R,3R)-3-[(2S)-1-(tert-butoxycarbonyl)pyrrolidin-2-yl]-3-methoxy-2-methylpropanoyl]-β-methyl-L-phenylalaninate.

[3153] HPLC (Method 6): R_f =2.3 min;

[3154] LC-MS (Method 1): R_f =1.36 min; MS (ESIpos): m/z =539 (M+H)⁺.


[3155] 230 mg (0.42 mmol) of this intermediate were taken up in 5 ml of dichloromethane, 5 ml of trifluoroacetic acid were added, and the mixture was stirred at RT for 30 min. Subsequently, the reaction mixture was concentrated under reduced pressure. The remaining residue was the reaction mixture dried further under reduced pressure and then lyophilized from acetonitrile/water. In this way, 230 mg (quant.) of the title compound were obtained.

[3156] HPLC (Method 6): R_f =1.6 min.

Intermediate 14

N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate


[3157]

x CF₃COOH

[3158] 143 mg (0.223 mmol) of N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 4) were taken up in 15 ml of DMF and admixed successively with 141 mg (0.22 mmol) of (2R,3R)-3-methoxy-2-methyl-N-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]-3-[(2S)-pyrrolidin-2-yl]propanamide trifluoroacetate (Intermediate 7), 102 mg (0.27 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium

Intermediate 15

N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S,3S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylbutan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[3165]

x CF₃COOH

hexafluorophosphate (HATU) and 128 μ l (0.74 mmol) of N,N-diisopropylethylamine. The mixture was stirred at RT for 3 h. The reaction mixture was then poured into a mixture of semisaturated aqueous ammonium chloride solution and ethyl acetate. The organic phase was removed, washed successively with saturated sodium hydrogencarbonate solution and saturated sodium chloride solution, dried over magnesium sulphate, filtered and concentrated. This gave 275 mg (quant.) of the Fmoc-protected intermediate N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S,3S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide.

[3159] HPLC (Method 5): R_f =2.73 min;

[3160] LC-MS (Method 4): R_f =3.19 min; MS (ESIpos): m/z =1023 (M+H)⁺.

[3161] 46 mg (0.045 mmol) of this intermediate were dissolved in 4 ml of DMF. After 1 ml of piperidine had been added, the reaction mixture was stirred at RT for 1 h. Subsequently, the reaction mixture was concentrated under reduced pressure and the residue was purified by means of preparative HPLC (eluent: acetonitrile+0.01% TFA/water+0.01% TFA). 22 mg (54% of theory) of the title compound were obtained as a colourless foam.

[3162] HPLC (Method 5): R_f =1.68 min;

[3163] LC-MS (Method 2): R_f =1.03 min; MS (ESIpos): m/z =801 (M+H)⁺

[3164] ¹H NMR (600 MHz, DMSO-d₆): δ =8.8 (m, 2H), 8.7 (m, 1H), 8.42 and 8.15 (2d, 1H), 7.3-7.1 (m, 5H), 5.12 and 4.95 (2m, 1H), 4.70 and 4.62 (2m, 1H), 4.62 and 4.50 (2t, 1H), 4.1-3.9 (m, 3H), 3.85 (m, 1H), 3.75-3.6 (m, 2H), 3.23, 3.18, 3.17, 3.14, 3.02 and 2.96 (6s, 9H), 3.1-2.9 and 2.75 (2m, 2H), 2.46 (m, 3H), 2.4-2.1 (m, 2H), 2.05 (br. m, 2H), 1.85-1.55 (br. m, 6H), 1.5-1.2 (br. m, 3H), 1.1-0.8 (m, 18H), 0.75 (t, 3H) [further signals hidden under H₂O peak].

[3166] 126 mg (0.198 mmol) of N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 4) were taken up in 10 ml of DMF and admixed successively with 105 mg (0.198 mmol) of (2R,3R)-3-methoxy-2-methyl-N-[(2S,3S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylbutan-2-yl]-3-[(2S)-pyrrolidin-2-yl]propanamide trifluoroacetate (Intermediate 17), 41.6 mg (0.217 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 33 mg (0.217 mmol) of 1-hydroxy-1H-benzotriazole hydrate and 79 μ l (0.454 mmol) of N,N-diisopropylethylamine. The mixture was stirred at RT overnight. The reaction mixture was then poured into a mixture of semisaturated aqueous ammonium chloride solution and ethyl acetate. The organic phase was removed, washed successively with saturated sodium hydrogencarbonate solution and saturated sodium chloride solution, dried over magnesium sulphate, filtered and concentrated. This gave 220 mg (quant.) of the Fmoc-protected intermediate N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S,3S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylbutan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide.

[3167] HPLC (Method 5): R_f =2.77 min;

[3168] LC-MS (Method 1): R_f =1.5 min; MS (ESIpos): m/z =1037 (M+H)⁺.

[3169] 220 mg (0.212 mmol) of this intermediate were dissolved in 5 ml of DMF. After 1 ml of piperidine had been added, the reaction mixture was stirred at RT for 1 h. Subsequently, the reaction mixture was concentrated under reduced pressure and the residue was purified by means of preparative HPLC (eluent: acetonitrile+0.01% TFA/water+0.01% TFA). 91 mg (46% of theory) of the title compound were obtained as a colourless foam.

[3170] HPLC (Method 5): R_f =1.71 min;

[3171] LC-MS (Method 1): R_f =0.9 min; MS (ESIpos): m/z =815 (M+H)⁺


[3172] ¹H NMR (600 MHz, DMSO-d₆): δ =8.87 and 8.80 (2d, 2H), 8.75 (m, 1H), 8.40 and 7.98 (2d, 1H), 7.3-7.1 (m,

5H), 5.45 and 5.2 (2t, 1H), 4.78 and 4.62 (2m, 1H), 4.73 and 4.58 (2t, 1H), 4.2-4.0 (m, 3H), 3.7-3.6 (m, 1H), 3.35, 3.20, 3.18, 3.14, 3.12 and 3.00 (6s, 9H), 3.1 and 2.95 (2m, 2H), 2.46 (m, 3H), 2.4-2.0 (m, 4H), 1.9-1.6 (m, 4H), 1.6-1.2 (m, 5H), 1.1-0.75 (m, 21H), 0.80 (t, 3H) [further signals hidden under H₂O peak].

Intermediate 16

N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[3173]



[3174] 617 mg (1.2 mmol) of tert-butyl (2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidine-1-carboxylate (Intermediate 24) were taken up in 44 ml of dichloromethane, 4.4 ml of trifluoroacetic acid were added and the mixture was stirred at RT for 30 min. Subsequently, the reaction mixture was concentrated under reduced pressure and the remaining residue was lyophilized from dioxane/water. 702 mg (quant.) of the deprotected compound (2R,3R)-3-methoxy-2-methyl-N-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]-3-[(2S)-pyrrolidin-2-yl]propanamide trifluoroacetate were obtained as a crude product, which was used in the following stage without further purification.

[3175] 470 mg (0.74 mmol) of N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 4) were taken up in 57 ml of DMF and admixed successively with 390 mg (approx. 0.74 mmol) of the above-obtained (2R,3R)-3-methoxy-2-methyl-N-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]-3-[(2S)-pyrrolidin-2-yl]propanamide trifluoroacetate, 336 mg (0.88 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) and 423 μ l (2.4 mmol) of N,N-diisopropylethylamine. The mixture was stirred at RT for 2 h. The reaction mixture was then poured into a mixture of semisaturated aqueous ammonium chloride solution and ethyl acetate. The organic phase was removed, washed successively with saturated sodium hydrogencarbonate solution and saturated sodium chloride solution, dried over sodium sulphate, filtered and concentrated. The residue was purified by preparative HPLC. This gave 453

mg (59% of theory) of the Fmoc-protected intermediate N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide.

[3176] HPLC (Method 5): R_f =2.58 min;

[3177] LC-MS (Method 1): R_f =3.10 min; MS (ESIpos): m/z =1035 (M+H)⁺.

[3178] 453 mg (0.438 mmol) of this intermediate were dissolved in 24 ml of DMF. After 2.4 ml of piperidine had been added, the reaction mixture was stirred at RT for 30 min. Subsequently, the reaction mixture was concentrated under reduced pressure and the residue was purified by preparative HPLC (eluent: acetonitrile/0.1% TFA in water). 260 mg (64% of theory) of the title compound were obtained as a colourless foam.

[3179] HPLC (Method 5): R_f =1.64 min;


[3180] LC-MS (Method 1): R_f =0.86 min; MS (ESIpos): m/z =813 (M+H)⁺

[3181] ¹H NMR (400 MHz, DMSO-d₆): δ =8.8 (m, 2H), 8.65 (m, 2H), 7.3-7.1 (m, 5H), 4.8-4.05 (m, 2H), 4.0 and 3.82 (2m, 2H), 3.8-3.5 (m, 8H), 3.32, 3.29, 3.20, 3.19, 3.12 and 3.00 (6s, 9H), 2.65 (t, 1H), 2.5-2.45 (m, 3H), 2.4-1.3 (m, 15H), 1.15-0.85 (m, 18H), 0.8 and 0.75 (2d, 3H) [further signals hidden under H₂O peak].

Intermediate 17

N-benzyl-N-methyl-L-phenylalaninamide trifluoroacetate

[3182]



[3183] 1000 mg (3.77 mmol) of N-(tert-butoxycarbonyl)-L-phenylalanine were dissolved in 10 ml of DMF and admixed with 457 mg (3.77 mmol) of N-methylbenzylamine, 2150 mg (5.65 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'

N'-tetramethyluronium hexafluorophosphate and 657 μ l of N,N-diisopropylethylamine. The reaction mixture was stirred at RT for 30 min and then concentrated under reduced pressure. The residue was taken up in dichloromethane and extracted by shaking three times with water. The organic phase was dried over magnesium sulphate and concentrated. The residue was purified by flash chromatography on silica gel with 3:1 petroleum ether/ethyl acetate as the eluent. The product fractions were concentrated and the residue was dried under high vacuum. This gave 1110 mg (75% of theory) of the Boc-protected intermediate N-benzyl-N^α-(tert-butoxycarbonyl)-N-methyl-L-phenylalaninamide.

[3184] HPLC (Method 6): R_f =2.1 min;

[3185] LC-MS (Method 1): R_f =1.14 min; MS (ESIpos): m/z =369 (M+H)⁺.

[3186] 1108 mg (3.007 mmol) of this intermediate were taken up in 30 ml of dichloromethane, 10 ml of trifluoroacetic acid were added, and the mixture was stirred at RT for 30 min. Subsequently, the reaction mixture was concentrated under reduced pressure, the remaining residue was stirred with dichloromethane and the solvent was distilled off. The residue was stirred twice more with pentane, the solvent was decanted off again each time and the of the title compound was finally dried under high vacuum. 1075 mg (93% of theory) of the title compound were thus obtained as a resin.


[3187] HPLC (Method 6): R_f =1.6 min;

[3188] LC-MS (Method 1): R_f =0.6 min; MS (ESIpos): m/z =269 (M+H)⁺.

Intermediate 18

N-benzyl-N^α-{(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoyl}-N-methyl-L-phenylalaninamide trifluoroacetate

[3189]



[3190] First, (2R,3R)-3-[(2S)-1-(tert-butoxycarbonyl)pyrrolidin-2-yl]-3-methoxy-2-methylpropanoic acid (Starting Compound 1) was released from 141 mg (0.491 mmol) of its dicyclohexylamine salt by taking it up in ethyl acetate and extractive shaking with 5% aqueous sulphuric acid. The organic phase was dried over magnesium sulphate, filtered and concentrated. The residue was taken up in 10 ml of DMF and 187.6 mg (0.49 mmol) of N-benzyl-N-methyl-L-phenylalaninamide trifluoroacetate (Intermediate 9), 190.3 mg (1.47 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) and 256 μ l of N,N-diisopropylethylamine were added. The mixture was stirred at RT for 1 h. The reaction mixture was then concen-

trated, the residue was taken up in ethyl acetate and the solution was subsequently extracted by shaking successively with saturated ammonium chloride solution, saturated sodium hydrogencarbonate solution and water. The organic phase was dried over magnesium sulphate and concentrated. The residue was purified by flash chromatography on silica gel with 30:1 acetonitrile/water as the eluent. The product fractions were concentrated and the residue was dried under high vacuum. This gave 168 mg (64% of theory) of the Boc-protected intermediate tert-butyl (2S)-2-[(1R,2R)-3-[(2S)-1-[benzyl(methyl)amino]-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidine-1-carboxylate.

[3191] HPLC (Method 6): R_f =2.2 min;

[3192] LC-MS (Method 2): R_f =1.22 min; MS (ESIpos): m/z =538 (M+H)⁺.

[3193] 168 mg (0.312 mmol) of this intermediate were taken up in 15 ml of dichloromethane, 3 ml of trifluoroacetic acid were added, and the mixture was stirred at RT for 30 min. Subsequently, the reaction mixture was concentrated under reduced pressure. The remaining residue was stirred first with dichloromethane, then with diethyl ether, and the solvent was distilled off again each time. After drying under high vacuum, 170 mg (99% of theory) of the title compound were obtained as a resin.


[3194] HPLC (Method 6): R_f =1.7 min;

[3195] LC-MS (Method 1): R_f =0.73 min; MS (ESIpos): m/z =438 (M+H)⁺.

Intermediate 19

methyl N-{(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoyl}-L-phenylalaninate trifluoroacetate

[3196]



[3197] The title compound was prepared in analogy to the synthesis of Intermediate 18, proceeding from (2R,3R)-3-[(2S)-1-(tert-butoxycarbonyl)pyrrolidin-2-yl]-3-methoxy-2-methylpropanoic acid (Starting Compound 1), which was released from the dicyclohexylamine salt, and methyl L-phenylalaninate hydrochloride.


[3198] HPLC (Method 5): R_f =0.6 min;

[3199] LC-MS (Method 3): R_f =1.17 min; MS (ESIpos): m/z =349 (M+H)⁺.

Intermediate 20

benzyl N-[(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoyl]-L-tryptophanate trifluoroacetate

[3200]



[3201] The title compound was prepared in analogy to the synthesis of Intermediate 18, proceeding from (2R,3R)-3-[(2S)-1-(tert-butoxycarbonyl)pyrrolidin-2-yl]-3-methoxy-2-methylpropanoic acid (Starting Compound 1), which was released from the dicyclohexylamine salt, and benzyl L-tryptophanate.


[3202] HPLC (Method 6): R_f =2.0 min;

[3203] LC-MS (Method 1): R_f =0.8 min; MS (ESIpos): m/z =464 (M+H)⁺.

Intermediate 21

benzyl (1S,2R)-1-[(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoyl]amino)-2-phenylcyclopropanecarboxylate trifluoroacetate

[3204]



[3205] The title compound was prepared in analogy to the synthesis of Intermediate 18, proceeding from (2R,3R)-3-[(2S)-1-(tert-butoxycarbonyl)pyrrolidin-2-yl]-3-methoxy-2-methylpropanoic acid (Starting Compound 1), which was released from the dicyclohexylamine salt, and benzyl (1S,2R)-1-amino-2-phenylcyclopropanecarboxylate. Benzyl (1S,2R)-1-amino-2-phenylcyclopropanecarboxylate had been prepared beforehand by standard methods, by esterifying commercially available (1S,2R)-1-[(tert-butoxycarbonyl)amino]-2-phenylcyclopropanecarboxylic acid with benzyl alcohol and subsequent Boc detachment with trifluoroacetic acid.


[3206] HPLC (Method 5): R_f =1.5 min;

[3207] LC-MS (Method 2): R_f =0.93 min; MS (ESIpos): m/z =437 (M+H)⁺.

Intermediate 22

6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-N'-methyl-hexanehydrazide trifluoroacetate

[3208]



[3209] 100 mg (473 μ mol) of 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoic acid were dissolved in 71 μ l of DMF and then admixed with 139 mg (947 μ mol) of tert-butyl 1-methylhydrazinecarboxylate, 182 mg (947 μ mol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 145 mg (947 μ mol) of 1-hydroxy-1H-benzotriazole hydrate. The mixture was stirred at RT overnight and then concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization from dioxane/water, 129 mg (80% of theory) of the protected intermediate were obtained as a colourless foam.


[3210] Subsequently, the 129 mg (380 μ mol) were deblocked with 2 ml of trifluoroacetic acid in 8 ml of dichloromethane. After stirring at RT for 1 h, the reaction mixture was concentrated under reduced pressure. The residue was lyophilized from acetonitrile/water, which left 125 mg (83% of theory) of the title compound as a colourless foam.

[3211] LC-MS (Method 1): R_f =0.38 min; MS (ESIpos): m/z =240 (M+H)⁺

Intermediate 23

N-(2-aminoethyl)-4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-N-methylbutanamide trifluoroacetate

[3212]



[3213] First, 35 mg (164 μ mol) of tert-butyl 2-(methylamino)ethyl carbamate hydrochloride trifluoroacetate, 30 mg (164 μ mol) of 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)butanoic acid, mg (197 μ mol) of O-(7-azabenzotriazol-1-yl)-N, N, N', N'-tetramethyluronium hexafluorophosphate and 57 μ l of N,N-diisopropylethylamine were combined in 5 ml of DMF and stirred at RT overnight. Subsequently, the solvent was removed under reduced pressure and the remaining residue was purified by means of preparative HPLC. The corresponding fractions were concentrated and, by lyophilization from dioxane/water, 35 mg (63% of theory) of the protected intermediate were obtained.

[3214] HPLC (Method 12): R_f =1.6 min;

[3215] LC-MS (Method 1): R_f =0.71 min; MS (ESIpos): m/z =340 (M+H)⁺.


[3216] Subsequently, the entire amount of the protected intermediate was deblocked with 1 ml of trifluoroacetic acid in 5 ml of dichloromethane to obtain 28 mg (77% of theory) of the title compound.

[3217] LC-MS (Method 3): R_f =0.75 min; MS (ESIpos): m/z =240 (M+H)⁺.

Intermediate 24

4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-N-[2-(methylamino)ethyl]butanamide trifluoroacetate

[3218]



[3219] First, 35 mg (164 μ mol) of tert-butyl (2-aminoethyl) methyl carbamate hydrochloride trifluoroacetate, 30 mg (164 μ mol) of 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)butanoic acid, mg (197 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and 57 μ l of N,N-diisopropylethylamine were combined in 5 ml of DMF and stirred at RT for 30 min. Subsequently, the solvent was removed under reduced pressure and the remaining residue was purified by means of preparative HPLC. The corresponding fractions were concentrated and, by lyophilization from dioxane/water, 51 mg (91% of theory) of the protected intermediate were obtained.

[3220] HPLC (Method 12): R_f =1.6 min;

[3221] LC-MS (Method 1): R_f =0.77 min; MS (ESIpos): m/z =340 (M+H)⁺.


[3222] Subsequently, the entire amount was deprotected with 1 ml of trifluoroacetic acid in 5 ml of dichloromethane to obtain 45 mg (69% of theory) of the title compound.

[3223] LC-MS (Method 1): R_f =0.19 min; MS (ESIpos): m/z =240 (M+H)⁺.

Intermediate 25

benzyl (2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoate trifluoroacetate

[3224]



[3225] First, (2R,3R)-3-[(2S)-1-(tert-butoxycarbonyl)pyrrolidin-2-yl]-3-methoxy-2-methylpropanoic acid was released from 1.82 g (388 mmol) of its dicyclohexylamine salt by taking it up in ethyl acetate and extractive shaking with 100 ml of 0.5% sulphuric acid. The organic phase was dried over magnesium sulphate, filtered and concentrated. The residue was taken up in 10 ml of dioxane and 10 ml of water, 1517 mg (4.66 mmol) of caesium carbonate were added, and the mixture was treated in an ultrasound bath for 5 min and concentrated under reduced pressure and redistilled once with DMF. The residue was then taken up in 15 ml of dichloromethane, and 1990 mg (11.64 mmol) of benzyl bromide were added. The mixture was treated in an ultrasound bath for 15 min and then concentrated under reduced pressure. The residue was partitioned between ethyl acetate and water, and the organic phase was removed and extracted by shaking with saturated sodium chloride solution and then concentrated. The residue was then purified by preparative HPLC. This gave 1170 mg (80% of theory) of the Boc-protected intermediate.

[3226] Subsequently, these 1170 mg were deprotected immediately with 5 ml of trifluoroacetic acid in 15 ml of dichloromethane. After stirring at RT for 15 min, the reaction mixture was concentrated under reduced pressure. The residue was lyophilized from dioxane. After drying under high vacuum, there remained 1333 mg (84% of theory) of the title compound as a yellow oil.


[3227] HPLC (Method 6): R_f =1.5 min;

[3228] LC-MS (Method 1): R_f =0.59 min; MS (ESIpos): m/z =278 (M+H)⁺.

Intermediate 26

N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3229]



[3230] 1200 mg (2.33 mmol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 5) were combined with 910.8 mg (2.33 mmol) of benzyl (2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoate trifluoroacetate (Intermediate 14), 1327 mg (3.49 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and 2027 μ l of N,N-diisopropylethylamine in 50 ml of DMF, and the mixture was stirred at RT for 5 min. Thereafter, the solvent was removed under reduced pressure. The remaining residue was taken up in ethyl acetate and extracted by shaking successively with 5% aqueous citric acid solution and saturated sodium hydrogen-carbonate solution. The organic phase was removed and concentrated. The residue was purified by means of preparative HPLC. The product fractions were combined and concentrated, and the residue was dried under high vacuum. This gave 1000 mg (55% of theory) of the benzyl ester intermediate N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-(benzyloxy)-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide as a resin.

[3231] LC-MS (Method 1): R_t =1.56 min; MS (ESIpos): m/z =775 (M+H)⁺.

[3232] The entire amount of this intermediate obtained was taken up in 25 ml of a mixture of methanol and dichloromethane (20:1), and the benzyl ester group was removed by hydrogenation under standard hydrogen pressure with 10% palladium on activated carbon as a catalyst. After stirring at RT for 30 min, the catalyst was filtered off and the filtrate was concentrated under reduced pressure. This gave 803 mg (91% of theory) of the title compound as a white solid.


[3233] HPLC (Method 6): R_t =2.1 min;

[3234] LC-MS (Method 1): R_t =1.24 min; MS (ESIpos): m/z =685 (M+H)⁺.

Intermediate 27

(1S,2R)-1-amino-2-phenyl-N-propylcyclopropanecarboxamide trifluoroacetate

[3235]



[3236] The title compound was prepared by coupling commercially available (1S,2R)-1-[(tert-butoxycarbonyl)amino]-2-phenylcyclopropanecarboxylic acid with n-propylamine in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) and subsequent Boc detachment with trifluoroacetic acid (yield: 85% of theory over both stages).


[3237] HPLC (Method 6): R_t =1.2 min;

[3238] LC-MS (Method 1): R_t =0.52 min; MS (ESIpos): m/z =219 (M+H)⁺.

Intermediate 28

ethyl
(1S,2R)-1-amino-2-phenylcyclopropanecarboxylate
trifluoroacetate

[3239]




[3240] The title compound was prepared by standard methods, by esterifying commercially available (1S,2R)-1-[(tert-butoxycarbonyl)amino]-2-phenylcyclopropanecarboxylic acid with ethanol and subsequent Boc detachment with trifluoroacetic acid.

[3241] LC-MS (Method 1): R_t =0.50 min; MS (ESIpos): m/z =206 (M+H)⁺.

Intermediate 29

4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2,2-dimethylbutanoic acid

[3242]




[3243] To a solution of 1.39 g (8.95 mmol) of N-methoxycarbonylmaimide in 44 ml of saturated sodium hydrogen-carbonate solution were added, at 0° C., 1.5 g (8.95 mmol) of 4-amino-2,2-dimethylbutyric acid, and the mixture was stirred for 40 min. Subsequently, the cooling bath was removed and the reaction mixture was stirred for a further 1 h. While cooling with ice, the reaction mixture was then adjusted to pH 3 by adding sulphuric acid, and extracted with ethyl acetate. The combined organic phases were dried over magnesium sulphate and concentrated. 1.17 g (79% purity, 49% of theory) of the title compound were obtained.

[3244] LC-MS (Method 1): R_t =0.64 min; m/z =212 (M+H)⁺.

Intermediate 30

tert-butyl 2-[4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2,2-dimethylbutanoyl]hydrazinecarboxylate

[3245]




[3246] To a solution of 50 mg (237 μ mol) of 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2,2-dimethylbutanoic acid in 2 ml of THF were added, at 0° C., first 26 μ l (237 μ mol) of 4-methylmorpholine and then 31 μ l (237 μ mol) of isobutyl chloroformate. After removing the cooling bath and stirring at RT for a further 15 min, 31.3 mg (237 μ mol) of tert-butyloxy-carbonyl hydrazide were added. The reaction mixture was stirred overnight and then concentrated. The residue was purified by preparative HPLC. 50.8 mg (66% of theory) of the title compound were obtained.

[3247] LC-MS (Method 1): R_f =0.71 min; m/z =324 (M-H)⁻.

Intermediate 31

4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2,2-dimethylbutanehydrazide trifluoroacetate

[3248]




[3249] 50 mg (154 mmol) of tert-butyl 2-[4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2,2-dimethylbutanoyl]hydrazinecarboxylate were dissolved in 2 ml of dichloromethane, and 0.4 ml of trifluoroacetic acid was added. The reaction mixture was stirred at RT for 30 min and then concentrated. 55.2 mg (93% purity, 99% of theory) of the title compound were obtained.

[3250] LC-MS (Method 1): R_f =0.36 min; m/z =226 (M+H)⁺.

Intermediate 32

adamantan-1-ylmethyl
N-(tert-butoxycarbonyl)-L-phenylalaninate

[3251]




[3252] To a solution of 500 mg (1.89 mmol) of N-Boc-L-phenylalanine in 25 ml of dichloromethane were added, at RT, 1192 mg (6.2 mmol) of EDC, 578 μ l (4.1 mmol) of triethylamine, 345 mg (2.8 mmol) of DMAP and 345 mg (2.1 mmol) of 1-adamantylmethanol. The reaction mixture was stirred overnight, then diluted with 50 ml of dichloromethane, and was successively washed with 10% aqueous citric acid solution, water and saturated sodium chloride solution. The organic phase was dried over magnesium sulphate, then concentrated, and the residue was purified by preparative HPLC. 769 mg (90% of theory) of the title compound were obtained.

[3253] LC-MS (Method 2): R_f =1.84 min; m/z =414 (M+H)⁺.

Intermediate 33

adamantan-1-ylmethyl L-phenylalaninate
hydrochloride

[3254]




[3255] 769 mg (1.86 mmol) of adamantan-1-ylmethyl N-(tert-butoxycarbonyl)-L-phenylalaninate (Intermediate 13) were dissolved in 25 ml of a 4 N solution of hydrogen chloride in dioxane and stirred at RT for 1 h. Subsequently, the reaction mixture was concentrated and the residue was dried under reduced pressure. 619 mg (95% of theory) of the title compound were obtained.

[3256] LC-MS (Method 1): R_f =0.82 min; m/z =314 (M+H)⁺.

Intermediate 34

N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-(adamantan-1-ylmethoxy)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3257]




[3258] To a solution of 20 mg (29 μmol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide in 1 ml of DMF were added, at RT, 15.3 μl (88 μmol) of N,N-diisopropylethylamine, 6.7 mg (44 μmol) of HOBt and 6.7 mg (35 μmol) of EDC, and the mixture was stirred for 30 min. Subsequently, 10.1 mg (32 μmol) of adamantan-1-yl L-phenylalaninate hydrochloride were added. After stirring overnight, the reaction mixture was separated directly into its components via preparative HPLC. 27.5 mg (93% of theory) of the title compound were obtained.

Intermediate 35

N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-(adamantan-1-ylmethoxy)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[3259] LC-MS (Method 1): R_f =1.70 min; m/z =980 ($M+H$)⁺.

[3260]




[3261] 27.5 mg (28 μ mol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(adamantan-1-ylmethoxy)-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 1.8 ml of dichloromethane, and 361 μ l of TFA were added. The reaction mixture was stirred for 30 min and then concentrated. The residue was taken up in water and lyophilized. 22.7 mg (81% of theory) of the title compound were obtained.

[3262] LC-MS (Method 1): R_t =1.14 min; m/z =880 (M+H)⁺.

Intermediate 36

tert-butyl (2S)-1-(benzyloxy)-3-phenylpropan-2-yl carbamate

[3263]



[3264] Under an argon atmosphere, 500 mg (1.99 mmol) of N-Boc-L-phenylalaninol were dissolved in 5 ml of DMF and cooled to 0° C. Subsequently, 159 mg (3.98 mmol) of a 60% suspension of sodium hydride in paraffin oil were added. The reaction mixture was stirred until the evolution of gas had ended and then 260 μ l (2.19 mmol) of benzyl bromide were added. The cooling bath was removed and the reaction mixture was stirred at RT for 2 h. Thereafter, the reaction mixture was concentrated, the residue was taken up in ice-water and the mixture was extracted with dichloromethane. The organic phase was washed with saturated sodium chloride solution,


dried over magnesium sulphate and concentrated. The residue was purified by means of preparative HPLC. 226 mg (33% of theory) of the title compound were obtained.

[3265] LC-MS (Method 1): R_t =1.28 min; m/z =342 (M+H)⁺.

Intermediate 37

(2S)-1-(benzyloxy)-3-phenylpropan-2-amine hydrochloride

[3266]




[3267] 220 mg (644 μ mol) of tert-butyl (2S)-1-(benzyloxy)-3-phenylpropan-2-yl carbamate were dissolved in 11 ml of a 4 N solution of hydrogen chloride in dioxane and stirred at RT for 1 h. Then the reaction mixture was concentrated and the residue was dried under reduced pressure. 138 mg (77% of theory) of the title compound were obtained.

[3268] LC-MS (Method 1): R_t =0.65 min; m/z =242 (M+H)⁺.

Intermediate 38

N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3269]




[3270] To a solution of 20 mg (29 μmol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide in 1 ml of DMF were added, at RT, 15.3 μl (88 μmol) of N,N-diisopropylethylamine, 6.7 mg (44 μmol) of HOBt and 6.7 mg (35 μmol) of EDC, and the mixture was stirred for 30 min. Subsequently, 7.8 mg (32 μmol) of (2S)-1-(benzyloxy)-3-phenylpropan-2-amine hydrochloride were added. After stirring overnight, the reaction mixture was separated directly into its components via preparative HPLC. 26 mg (98% of theory) of the title compound were obtained.

[3271] LC-MS (Method 1): R_t =1.51 min; m/z =909 (M+H)⁺.

Intermediate 39

N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[3272]




[3273] 26 mg (29 μmol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 1.8 ml of dichloromethane, and 370 μl of TFA were added. The reaction mixture was stirred at RT for 30 min and then concentrated. The residue was taken up in water and lyophilized. 26.4 mg (quant.) of the title compound were obtained.

[3274] LC-MS (Method 1): R_t =0.97 min; m/z =809 (M+H)⁺.

Intermediate 40

N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S,2R)-1-hydroxy-1-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3275]



[3276] 50 mg (70 μ mol) of Intermediate 26 and 11 mg (70 μ mol) of (1S,2R)-2-amino-1-phenylpropan-1-ol in 10 ml of DMF were admixed with 42 mg (0.11 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and 25 μ l of N,N-diisopropylethylamine, and the reaction mixture was stirred at RT for 5 min. This was followed by concentration and purification of the residue by means of preparative HPLC. After combining the corresponding fractions, concentrating and drying under high vacuum, 49 mg (81%) of the protected intermediate were obtained. Subsequently, the Boc group was detached by known conditions with trifluoroacetic acid in dichloromethane. Concentration was followed by the purification of the title compound by preparative HPLC, and 26 mg (52%) of the title compound were obtained.


[3277] HPLC (Method 12): R_f =1.65 min;

[3278] LC-MS (Method 1): R_f =0.77 min; MS (ESIpos): m/z =718 (M+H)⁺.

Intermediate 41

3-{2-[2-(2-aminoethoxy)ethoxy]ethoxy}propanoic acid trifluoroacetate

[3279]



[3280] 150 mg (541 μ mol) of tert-butyl 3-{2-[2-(2-aminoethoxy)ethoxy]ethoxy}propanoate were dissolved in 3 ml of dichloromethane, 1.5 ml of trifluoroacetic acid were added, and the reaction mixture was stirred at RT for 1 h, then concentrated.


[3281] 181 mg (100% of theory) of the title compound were obtained.

[3282] MS (EI): m/z 222 (M+H)⁺

Intermediate 42

3-(2-{2-[2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethoxy]ethoxy}ethoxy)propanoic acid

[3283]



[3284] 186 mg (555 μ mol) of 3-{2-[2-(2-aminoethoxy)ethoxy]ethoxy}propanoic acid trifluoroacetate were dissolved in 2.6 ml of saturated sodium hydrogencarbonate solution and admixed at 0° C. with 86 mg (555 μ mol) of N-methoxycarbonylmaleimide. The reaction mixture was stirred at 0° C. for 40 min and at RT for 1 h, then cooled again to 0° C., adjusted to pH 3 with sulphuric acid and extracted 3 $\times$ with 25 ml of ethyl acetate. The combined organic phases were dried over magnesium sulphate and concentrated.


[3285] 126 mg (75% of theory) of the title compound were obtained.

[3286] LC-MS (Method 1): R_f =0.53 min; m/z =302 (M+H)⁺.

Intermediate 43

tert-butyl 15-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-4-oxo-7,10,13-trioxa-2,3-diazapentadecan-1-oate

[3287]



[3288] 125 mg (417 μ mol) of 3-(2-{2-[2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethoxy]ethoxy}ethoxy) propanoic acid were dissolved at 0° C. in 2.1 ml of THF and admixed with 46 μ l (417 μ mol) of 4-methylmorpholine and 54.5 μ l (417 μ mol) of isobutyl chloroformate. The ice bath was removed and the reaction mixture was stirred at RT for 30 min. Subsequently, at 0° C., 55 mg (417 μ mol) of tert-butylloxycarbonyl hydrazide were added. The reaction mixture was warmed to RT overnight, concentrated and purified by preparative HPLC.


[3289] 60 mg (33% of theory) of the title compound were obtained.

[3290] LC-MS (Method 1): R_f =0.66 min; m/z =416 (M+H)⁺.

Intermediate 44

3-(2-{2-[2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethoxy]ethoxy}ethoxy)propanehydrazide trifluoroacetate

[3291]



[3292] 60 mg (145 μ mol) of tert-butyl 15-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-4-oxo-7,10,13-trioxa-2,3-diazapentadecan-1-oate were dissolved in 1 ml of dichloromethane, and 0.2 ml of trifluoroacetic acid was added. The reaction mixture was stirred at RT for 30 min and then concentrated.


[3293] 62 mg (100% of theory) of the title compound were obtained.

[3294] LC-MS (Method 1): R_f =0.35 min; m/z =316 (M+H)⁺.

Intermediate 45

benzyl
(1S,2R)-1-amino-2-phenylcyclopropanecarboxylate
trifluoroacetate

[3295]




[3296] The title compound was prepared by standard methods, by esterifying commercially available (1S,2R)-1-[(tert-butoxycarbonyl)amino]-2-phenylcyclopropanecarboxylic acid with benzyl alcohol and subsequent Boc detachment with trifluoroacetic acid.

[3297] LC-MS (Method 1): R_f =0.72 min; MS (ESIpos): m/z =268 (M+H)⁺.

Intermediate 46

N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(1S)-1-carboxy-2-phenylethyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3298]



[3299] 383 mg (0.743 mmol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 8) were combined with 485 mg (0.743 mmol) of benzyl N-[(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]pro-

panoyl]-L-phenylalaninate trifluoroacetate (Intermediate 12), 424 mg (1.114 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and 388 μ l of N,N-diisopropylethylamine in 15 ml of DMF, and the mixture was stirred at RT for 10 min. Subsequently, the solvent was removed under reduced pressure. The remaining residue was taken up in ethyl acetate and extracted by shaking successively with 5% aqueous citric acid solution and saturated sodium hydrogencarbonate solution. The organic phase was removed and concentrated, and the residue was purified by means of preparative HPLC. The product fractions were combined and concentrated, and the residue was dried under high vacuum. 335 mg (48% of theory) of the benzyl ester intermediate were obtained as a foam.

[3300] LC-MS (Method 1): R_f =1.49 min; MS (ESIpos): m/z =922 (M+H)⁺.

[3301] 100 mg (0.11 mmol) of this intermediate were taken up in 15 ml of methanol and the benzyl ester group was removed by hydrogenation under standard hydrogen pressure with 10% palladium on activated carbon as a catalyst. After stirring at RT for 1 h, the catalyst was filtered off and the filtrate was concentrated under reduced pressure. After lyophilization from dioxane, 85 mg (94% of theory) of the title compound were obtained as a solid.


[3302] HPLC (Method 12): R_f =2.4 min;

[3303] LC-MS (Method 1): R_f =1.24 min; MS (ESIpos): m/z =832 (M+H)⁺.

Intermediate 47

N-benzyl-L-tryptophanamide trifluoroacetate

[3304]



[3305] 202 mg (0.5 mmol) of 2,5-dioxopyrrolidin-1-yl N-(tert-butoxycarbonyl)-L-tryptophanate and 45 mg (0.42 mmol) of benzylamine were dissolved in 10 ml of DMF, and 110 μ l (630 μ mol) of N,N-diisopropylethylamine were added. The reaction mixture was stirred at RT for 3 h. Subsequently,

the reaction mixture was concentrated under reduced pressure and the residue was purified by flash chromatography on silica gel (eluent: 20:0.5:0.05 dichloromethane/methanol/17% aq. ammonia). The corresponding fractions were combined and concentrated. The resulting residue was digested with diethyl ether and then dried under high vacuum. Subsequently, this residue was taken up in 10 ml of dichloromethane, and 3 ml of anhydrous trifluoroacetic acid were added. After stirring at RT for 45 minutes, the mixture was concentrated and the residue was purified by preparative HPLC. After drying under high vacuum, 117 mg (57% of theory over both stages) of the title compound were obtained.


[3306] HPLC (Method 12): $R_f=1.6$ min;

[3307] LC-MS (Method 1): $R_f=0.66$ min; MS (ESIpos): $m/z=294$ (M+H)⁺.

Intermediate 48

(1S,2R)-1-amino-2-phenylcyclopropanecarboxamide trifluoroacetate

[3308]



[3309] 50 mg (180 μ mol) of commercially available (1S,2R)-1-[(tert-butoxycarbonyl)amino]-2-phenylcyclopropanecarboxylic acid were dissolved in 5 ml of DMF, 94 μ l (541 μ mol) of N,N-diisopropylethylamine, 31 mg (270 μ mol) of N-hydroxysuccinimide and 41.5 mg (216 μ mol) of EDC were added, and then the mixture was stirred at RT overnight. The reaction mixture was then concentrated, the residue was taken up in dioxane, 71 mg (901 μ mol) of ammonium hydrogen carbonate were added, and the reaction mixture was then left to stand at RT for 3 days. The reaction mixture was then diluted with a 1:1 mixture of ethyl acetate and water. The organic phase was removed, dried over magnesium sulphate and concentrated. The resulting residue was subsequently taken up in 3 ml of dichloromethane, and 3 ml of anhydrous trifluoroacetic acid were added. Stirring at RT for 1 h was followed by concentration. The residue was stirred with pentane, filtered off with suction and lyophilized from dioxane. In this way, 32 mg (62% of theory over both stages) of the title compound were obtained.


[3310] HPLC (Method 6): $R_f=0.38$ min;

[3311] LC-MS (Method 1): $R_f=0.20$ min; MS (ESIpos): $m/z=177$ (M+H)⁺.

Intermediate 49

N^α-{(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoyl}-L-tryptophanamide trifluoroacetate

[3312]



[3313] The title compound was prepared in analogy to the synthesis of Intermediate 13 from Starting Compound 1 and L-tryptophanamide hydrochloride.


[3314] HPLC (Method 5): $R_f=1.4$ min;

[3315] LC-MS (Method 1): $R_f=0.92$ min; MS (ESIpos): $m/z=473$ (M+H)⁺.

Intermediate 50

4-nitrophenyl 2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethyl carbamate

[3316]



[3317] 813 mg (3.1 mmol) of triphenylphosphine were dissolved in 25 ml of THF and cooled to -70° C. under argon. After dropwise addition of 627 mg (3.1 mmol) of diisopropyl azodicarboxylate, the mixture was stirred for 5 min. Subsequently, 500 mg (3.1 mmol) of tert-butyl 2-aminoethyl carbamate dissolved in 5 ml of THF were added dropwise, and the reaction mixture was stirred at -70° C. for a further 5 min. Then 136.6 mg (1.55 mmol) of 2,2-dimethyl-1-propanol dissolved in 1 ml of THF and 301 mg (3.1 mmol) of maleimide were added, the reaction mixture was stirred at -70° C. for a further 10 min and then the mixture was warmed to RT. After stirring at RT for a further 16 h, the solvent was removed under reduced pressure and the residue was purified by means of preparative HPLC. This gave 463 mg (62%) of the protected intermediate.

[3318] After removing the Boc protecting group under standard conditions, 652 mg of 1-(2-aminoethyl)-1H-pyrrole-2,5-dione were obtained as the trifluoroacetate.

[3319] 112.9 mg (543 μ mol) of nitrophenyl chloroformate were dissolved in 30 ml of THF and, after addition of 100 mg (271.6 μ mol) of 1-(2-aminoethyl)-1H-pyrrole-2,5-dione trifluoroacetate, the mixture was stirred at RT for 30 min. The mixture was filtered and the filtrate was concentrated to dryness and then slurried with diethyl ether. After filtration with suction and drying, 60 mg (95% of theory) of the title compound were obtained.


[3320] HPLC (Method 5): R_t =0.65 min;

[3321] LC-MS (Method 1): R_t =0.74 min; MS (ESIpos): m/z =306 (M+H)⁺.

Intermediate 51

(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethanamine trifluoroacetate

[3322]



[3323] 200 mg (0.75 mmol) of N-(tert-butoxycarbonyl)-L-phenylalanine were initially charged at 0° C. in 5.5 ml of dichloromethane, and 128 mg (0.79 mmol) of 1,1'-carbonyldiimidazole were added. After 30 min, 103 mg (0.75 mmol) of benzoyl hydrazide were added. After a further 45 min at 0° C., 500 mg (1.5 mmol) of carbon tetrabromide and 395 mg (1.5 mmol) of triphenylphosphine were finally added. The reaction mixture was stirred first at 0° C. for 2 h and then at RT overnight. The mixture was subsequently concentrated on a rotary evaporator, and the residue was dried under high vacuum. The crude product thus obtained was purified by means of preparative HPLC. 217 mg (78% of theory) of the Boc-protected intermediate tert-butyl (1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl carbamate were obtained.

[3324] LC-MS (Method 12): R_t =1.15 min; MS (ESIpos): m/z =366 (M+H)⁺


[3325] 217 mg (0.59 mmol) of this intermediate were taken up in 3 ml of dichloromethane, 0.6 ml of trifluoroacetic acid were added, and the mixture was stirred at RT for 30 min. Subsequently, the reaction mixture was concentrated under reduced pressure. The remaining residue was the reaction mixture dried further under reduced pressure and then lyophilized from dioxane. In this way, 214 mg (90% of theory) of the title compound were obtained.

[3326] LC-MS (Method 11): R_t =0.62 min; MS (ESIpos): m/z =266 (M+H)⁺

Intermediate 52

(1R)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethanamine trifluoroacetate

[3327]



[3328] 200 mg (0.75 mmol) of N-(tert-butoxycarbonyl)-D-phenylalanine were initially charged at 0° C. in 5.5 ml of dichloromethane, and 128.3 mg (0.79 mmol) of 1,1'-carbonyldiimidazole were added. After 30 min, 103 mg (0.75 mmol) of benzoyl hydrazide were added. After a further 45 min at 0° C., 500 mg (1.5 mmol) of carbon tetrabromide and 395 mg (1.5 mmol) of triphenylphosphine were finally added. The reaction mixture was stirred first at 0° C. for 2 h and then at RT overnight. The mixture was subsequently concentrated on a rotary evaporator, and the residue was dried under high vacuum. The crude product thus obtained was purified by means of preparative HPLC. 219 mg (80% of theory) of the Boc-protected intermediate tert-butyl (1R)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl carbamate were obtained.

[3329] LC-MS (Method 2): R_t =1.36 min; MS (ESIpos): m/z =366 (M+H)⁺


[3330] 219 mg (0.6 mmol) of this intermediate were taken up in 3 ml of dichloromethane, 0.6 ml of trifluoroacetic acid were added, and the mixture was stirred at RT for 30 min. Subsequently, the reaction mixture was concentrated under reduced pressure. The remaining residue was the reaction mixture dried further under reduced pressure and then lyophilized from dioxane. In this way, 196 mg (86% of theory) of the title compound were obtained as a solid.

[3331] HPLC (Method 10): R_t =2.41 min

Intermediate 53

(2S)-1-(benzylsulphonyl)-3-phenylpropan-2-amine

[3332]



[3333] 200 mg (1.13 mmol) of (4S)-4-benzyl-1,3-oxazolidin-2-one were initially charged in 3 ml of tert-butanol, and 280 mg (2.26 mmol) of benzyl mercaptan were added. The mixture was subsequently heated under reflux for 2 days. Thereafter, the reaction mixture was concentrated on a rotary evaporator and the resulting (2S)-1-(benzylsulphonyl)-3-phenylpropan-2-amine intermediate was converted further directly, without workup.

[3334] HPLC (Method 10): R_f =2.63 min

[3335] LC-MS (Method 1): R_f =0.67 min; MS (ESIpos): m/z =258 (M+H)⁺

[3336] The crude intermediate obtained above was dissolved in a solution of 2 ml of 30% hydrogen peroxide and 5 ml of formic acid, and the mixture was stirred at RT for 12 h. Then the reaction mixture was added to saturated sodium sulphate solution and extracted three times with ethyl acetate. The organic phase was dried over magnesium sulphate and concentrated under reduced pressure. The crude product obtained was purified by means of preparative HPLC. 343 mg (61% of theory) of the title compound were thus obtained.


[3337] HPLC (Method 10): R_f =2.40 min;

[3338] LC-MS (Method 12): R_f =0.65 min; MS (ESIpos): m/z =290 (M+H)⁺

Intermediate 54

(2S,3E)-1,4-diphenylbut-3-en-2-amine


[3339]



Intermediate 55

N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[3345]



[3340] 552.7 mg (9.85 mmol) of potassium hydroxide were dissolved in methanol, adsorbed onto 1.1 g of neutral aluminium oxide and then dried under high vacuum. To a solution of 240 mg (0.82 mmol) of (2S)-1-(benzylsulphonyl)-3-phenylpropan-2-amine and 1.56 g of the potassium hydroxide on aluminium oxide thus prepared in 6.2 ml of n-butanol were added dropwise, at 5-10° C., 307 µl (3.3 mmol) of dibromodifluoromethane. The reaction mixture was stirred at RT for 2 h, then filtered through Celite, and the residue was washed thoroughly with dichloromethane. The filtrate was concentrated and the resulting residue was dried under reduced pressure. The crude product thus obtained was purified by means of preparative HPLC. 98 mg (35% of theory) of the title compound were obtained with an E/Z diastereomer ratio of 4:1.

[3341] HPLC (Method 10): R_f =2.46 min;

[3342] LC-MS (Method 12): R_f =0.75 min; MS (ESIpos): m/z =224 (M+H)⁺

[3346] 20 mg (29 µmol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 1 ml of DMF, 13.3 mg (35 µmol) of HATU and 15.3 µl (88 µmol) of N,N-diisopropylethylamine were added, and the mixture was stirred at RT for 30 min. Subsequently, 12.2 mg (32 µmol) of (1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethanamine trifluoroacetate were added. The reaction mixture was stirred at RT overnight and then separated by preparative HPLC. This gave 22 mg (81% of theory) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide.

[3347] LC-MS (Method 12): R_f =1.45 min; MS (ESIpos): m/z =933 (M+H)⁺


[3348] By subsequently detaching the BOC protecting group with trifluoroacetic acid, 22.4 mg (98% of theory) of the title compound were then obtained.

[3349] LC-MS (Method 11): $R_t=0.85$ min; MS (ESIpos): $m/z=833$ (M+H)⁺

Intermediate 56

N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1R)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[3350]



[3351] N-(tert-Butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1R)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared in analogy to the synthesis of Intermediate 55, by reaction of 20 mg (29 μ mol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide with 12.2 mg (32 μ mol) of (1R)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethanamine trifluoroacetate.

[3352] Yield: 17 mg (64% of theory)

[3353] HPLC (Method 10): $R_t=3.74$ min;

[3354] LC-MS (Method 1): $R_t=1.45$ min; MS (ESIpos): $m/z=933$ (M+H)⁺

[3355] By subsequently detaching the BOC protecting group with trifluoroacetic acid, 17.1 mg (99% of theory) of the title compound were then obtained.


[3356] HPLC (Method 10): $R_t=2.55$ min;

[3357] LC-MS (Method 11): $R_t=0.85$ min; MS (ESIpos): $m/z=833$ (M+H)⁺

Intermediate 57

N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzylsulphonyl)-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[3358]



[3359] N-(tert-Butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzylsulphonyl)-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared in analogy to the synthesis of Intermediate 55, by reaction of 20 mg (29 μ mol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide with 9.3 mg (20 μ mol) of (2S)-1-(benzylsulphonyl)-3-phenylpropan-2-amine.

[3360] Yield: 19.2 mg (68% of theory)

[3361] HPLC (Method 10): R_f =3.5 min;

[3362] LC-MS (Method 12): R_f =1.41 min; MS (ESIpos): m/z =957 (M+H)⁺

[3363] By subsequently detaching the BOC protecting group with trifluoroacetic acid, 19.3 mg (99% of theory) of the title compound were then obtained.


[3364] HPLC (Method 10): R_f =2.52 min;

[3365] LC-MS (Method 1): R_f =0.86 min; MS (ESIpos): m/z =857 (M+H)⁺

Intermediate 58

N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S,3E)-1,4-diphenylbut-3-en-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[3366]



[3367] N-(tert-Butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S,3E)-1,4-diphenylbut-3-en-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared in analogy to the synthesis of Intermediate 55, by reaction of 20 mg (29 μ mol) N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide with 7.1 mg (32 μ mol) of (2S,3E)-1,4-diphenylbut-3-en-2-amine.

[3368] Yield: 15.1 mg (58% of theory)

[3369] HPLC (Method 10): R_f =4.2 min;

[3370] LC-MS (Method 12): R_f =1.51 min; MS (ESIpos): m/z =891 (M+H)⁺

[3371] By subsequently detaching the BOC protecting group with trifluoroacetic acid, 15.7 mg (99% of theory) of the title compound were then obtained.


[3372] HPLC (Method 10): R_f =2.62 min;

[3373] LC-MS (Method 12): R_f =0.97 min; MS (ESIpos): m/z =791 (M+H)⁺

Intermediate 61

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3374]



[3375] 50 mg (0.054 mmol) of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate (Intermediate 16) were dissolved in 8 ml of dioxane/water, and 70 ml (0.108 mmol) of a 15% solution of 4-oxobutanoic acid in water were added. The reaction mixture was subsequently stirred at 100° C. for 1 h. After cooling to RT, 3.7 mg (0.059 mmol) of sodium cyanoborohydride were added and the mixture was adjusted to a pH of 3 by adding about 300 μ l of 0.1 N hydrochloric acid. The reaction mixture was then stirred at 100° C. for a further 2 h. After cooling, another 70 ml (0.108 mmol) of the 15% 4-oxobutanoic acid solution were added and the reaction mixture was again stirred at 100° C. for 1 h. Then a further 3.7 mg (0.059 mmol) of sodium cyanoborohydride were added and about 300 μ l of 0.1 N hydrochloric acid were subsequently used to adjust the pH back to 3. The reaction mixture was then stirred at 100° C. for another 2 h. In the event of conversion still being incomplete, this procedure was repeated for a third time. The reaction mixture was finally concentrated and the residue was

purified by means of preparative HPLC. In this way, 32 mg (65% of theory) of the title compound were obtained in the form of a colourless foam.

[3376] HPLC (Method 5): R_f =1.64 min;


[3377] LC-MS (Method 9): R_f =4.76 min; MS (ESIpos): m/z =899 (M+H)⁺

[3378] ¹H NMR (500 MHz, DMSO- d_6): δ =8.95 and 8.8 (2m, 1H), 8.88 and 8.65 (2s, 1H), 7.4-7.1 (m, 5H), 5.0, 4.78, 4.65 and 4.55 (4m, 2H), 4.1-3.7 (m, 5H), 3.32, 3.29, 3.20, 3.12, 3.1 and 3.0 (6s, 9H), 2.75 (m, 2H), 2.63 (t, 1H), 2.4-2.2 (m, 4H), 2.1-1.2 (m, 12H), 1.2-0.8 (m, 16H), 0.75 (m, 3H) [further signals hidden under H₂O and DMSO peaks].

Intermediate 62

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3379]



[3380] The title compound was prepared in analogy to the synthesis of Intermediate 61, by reaction of 50 mg of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate (Intermediate 14) with 4-oxobutanoic acid.

[3381] Yield: 34 mg (70% of theory)


[3382] HPLC (Method 5): R_f =1.64 min;

[3383] LC-MS (Method 9): R_f =4.77 min; MS (ESIpos): m/z =887 (M+H)⁺.

Intermediate 63

N-(4-carboxybenzyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3384]



[3385] The title compound was prepared in analogy to the synthesis of Intermediate 61, by reaction of 15 mg of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate (Intermediate 16) with 4-formylbenzoic acid.

[3386] Yield: 7.5 mg (48% of theory)


[3387] HPLC (Method 5): R_f =1.75 min;

[3388] LC-MS (Method 1): R_f =0.97 min; MS (ESIpos): m/z =947 (M+H)⁺.

Intermediate 64

N-(5-carboxypentyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3389]



[3390] 10 mg (0.011 mmol) of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate (Intermediate 16) were dissolved in 2 ml of dioxane/water, and 2.8 mg (0.022 mmol) of 6-oxohexanoic acid were added. The reaction mixture was subsequently stirred at 100° C. for 1 h. After cooling to RT, 0.75 mg (0.012 mmol) of sodium cyanoborohydride was added and the mixture was adjusted to a pH of 3 by adding 0.1 N hydrochloric acid. The reaction mixture was then stirred at 100° C. for a further hour. After cooling, another 2.8 mg (0.022 mmol) of 6-oxohexanoic acid were added and the reaction mixture was again stirred at 100° C. for 1 h. A further 0.75 mg (0.012 mmol) of sodium cyanoborohydride was added and 0.1 N hydrochloric acid was subsequently used to adjust the pH back to 3. The reaction mixture was then stirred at 100° C. for another 1 h. This procedure was then repeated for a third time.

The reaction mixture was finally concentrated and the crude product was purified by means of preparative HPLC. This gave 6.4 mg (64% of theory) of the title compound in the form of a colourless foam.


[3391] HPLC (Method 5): R_f =1.68 min;

[3392] LC-MS (Method 9): R_f =4.86 min; MS (ESIpos): m/z =927 (M+H)⁺.

Intermediate 65

N-(2-aminoethyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide bistrifluoroacetate

[3393]



[3394] The title compound was prepared by reaction of 68 mg of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate (Intermediate 14) with tert-butyl 2-oxoethyl carbamate and subsequent detachment of the Boc protecting group with trifluoroacetic acid.

[3395] Yield: 49 mg (62% of theory over two stages)

[3396] HPLC (Method 5): R_f =1.58 min;


[3397] LC-MS (Method 2): R_f =1.05 min; MS (ESIpos): m/z =844 (M+H)⁺

[3398] ¹H NMR (600 MHz, DMSO-d₆): δ=8.25 (m, 1H), 8.45 and 8.15 (2d, 1H), 7.65-7.55 (m, 3H), 7.23-7.1 (m, 5H), 5.12 and 4.95 (2m, 1H), 4.72 and 4.62 (2m, 1H), 4.6 and 4.52 (2t, 1H), 4.2-3.8 (m, 4H), 3.7 (d, 1H), 3.23, 3.20, 3.19, 3.18, 3.03 and 2.98 (6s, 9H), 3.0-2.7 (m, 6H), 2.4-1.2 (m, 15H), 1.05, 1.0, 0.88 and 0.82 (4d, 6H), 0.92 (m, 6H), 0.73 (m, 6H) [further signals hidden under H₂O peak].

Intermediate 66

N-(3-aminopropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3399]



[3400] The title compound was prepared in analogy to the synthesis of Intermediate 65, by reaction of 25 mg (0.027 mmol) of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate (Intermediate 16) with benzyl 3-oxopropyl carbamate and subsequent hydrolytic detachment of the Z protecting group (with 10% palladium on charcoal as a catalyst, in ethanol as a solvent).

[3401] Yield: 11 mg (41% of theory over two stages)


[3402] HPLC (Method 5): R_f =1.53 min;

[3403] LC-MS (Method 1): R_f =0.72 min; MS (ESIpos): m/z =870 (M+H)⁺.

Intermediate 67

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(adamantan-1-ylmethoxy)-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3404]



[3405] 26 mg (26 μ mol) of N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(adamantan-1-yl-methoxy)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate and 33.9 μ l of a 15% aqueous succinaldehydic acid solution (53 μ mol) were dissolved in 957 μ l of a 1:1-dioxane/water mixture and heated to 100° C. for 1 h. After brief cooling, 1.81 mg (29 μ mol) of sodium cyanoborohydride were added. The reaction mixture was adjusted to pH 3 by adding 0.1 N hydrochloric acid and the mixture was heated to 100° C. for a further 2 h. After again adding the same amounts of succinaldehydic acid solution, sodium cyanoborohydride and hydrochloric acid, the mixture was heated once again to 100° C. for 2 h. The reaction mixture was


then separated directly into its components by means of preparative HPLC. 18.5 mg (73% of theory) of the title compound were thus obtained.

[3406] LC-MS (Method 1): R_t =1.17 min; m/z =967 (M+H)⁺.

Intermediate 68

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3407]




[3408] mg (26 μ mol) of N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate and 33.7 μ l of a 15% aqueous succinaldehydic acid solution (52 μ mol) were dissolved in 953 μ l of a 1:1-dioxane/water mixture and heated to 100° C. for 1 h. After brief cooling, 1.80 mg (29 μ mol) of sodium cyanoborohydride were added. The reaction mixture was adjusted to pH 3 by adding 0.1 N hydrochloric acid and the mixture was heated to 100° C. for a further 2 h. After again adding the same amounts of succinaldehydic acid solution, sodium cyanoborohydride and hydrochloric acid, the mixture was heated once again to 100° C. for 2 h. The reaction mixture was then separated directly into its components by means of preparative HPLC. 15.2 mg (65% of theory) of the title compound were thus obtained.

[3409] LC-MS (Method 1): R_t =1.01 min; m/z =895 (M+H)⁺

Intermediate 69

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3410]




[3411] 53 mg (84 μ mol) of N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 4) and 45 mg (84 μ mol) of benzyl N-[(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoyl]-L-phenylalaninate trifluoroacetate (Intermediate 12) were taken up in 2 ml of DMF, 19 μ l of N,N-diisopropylethylamine, 14 mg (92 μ mol) of HOBt and 17.6 mg (92 μ mol) of EDC were

Intermediate 70

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(2S,3S)-1-(benzyloxy)-1-oxo-3-phenylbutan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3419]



added and then the mixture was stirred at RT overnight. Subsequently, the reaction mixture was concentrated and the residue was purified by means of preparative HPLC. This gave 59 mg (68% of theory) of the Fmoc-protected intermediate N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(2S,3S)-1-(benzyloxy)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide.

[3412] LC-MS (Method 1): R_t =1.55 min; m/z =1044 (M+H)⁺.

[3413] 57 mg (0.055 mmol) of this intermediate were treated with 1.2 ml of piperidine in 5 ml of DMF to detach the Fmoc protecting group. After concentration and purification by means of preparative HPLC, 39 mg (76% of theory) of the free amine intermediate N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(2S,3S)-1-(benzyloxy)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were obtained as the trifluoroacetate.

[3414] HPLC (Method 5): R_t =1.9 min;

[3415] LC-MS (Method 1): R_t =1.01 min; m/z =822 (M+H)⁺.

[3416] 37 mg (0.045 mmol) of this intermediate were dissolved in 5 ml of dioxane/water and, analogously to the preparation of the compound in Intermediate 66, reacted with a 15% aqueous solution of 4-oxobutanoic acid in the presence of sodium cyanoborohydride. 16 mg (39% of theory) of the title compound were obtained in the form of a colourless foam.

[3417] HPLC (Method 6): R_t =2.1 min;

[3418] LC-MS (Method 1): R_t =1.01 min; MS (ESIpos): m/z =908 (M+H)⁺.

[3420] First, in analogy to the synthesis described in Intermediate 14, proceeding from Intermediates 4 and 26, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(2S,3S)-1-(benzyloxy)-1-oxo-3-phenylbutan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared.

[3421] 30 mg (0.032 mmol) of this compound were dissolved in 6 ml of dioxane/water, and 41 μ l (0.063 mmol) of a 15% aqueous solution of 4-oxobutanoic acid were added. The reaction mixture was subsequently stirred at 100° C. for 1 h. After cooling to RT, 2.2 mg (0.035 mmol) of sodium cyanoborohydride were added and the mixture was adjusted to a pH of 3 by adding about 300 μ l of 0.1 N hydrochloric acid. The reaction mixture was then stirred at 100° C. for a further 2 h. After cooling, another 41 μ l (0.063 mmol) of the 15% 4-oxobutanoic acid solution were added and the reaction mixture was again stirred at 100° C. for 1 h. Then a further 2.2 mg (0.035 mmol) of sodium cyanoborohydride were added and about 300 μ l of 0.1 N hydrochloric acid were subsequently used to adjust the pH back to 3. The reaction mixture was then stirred at 100° C. for another 2 h. In the event of conversion still being incomplete, this procedure was repeated for a third time. The reaction mixture was finally concentrated and the crude product was purified by means of preparative HPLC. This gave 24 mg (82% of theory) of the title compound in the form of a colourless foam.


[3422] HPLC (Method 5): R_t =1.9 min;

[3423] LC-MS (Method 9): R_t =5.15 min; MS (ESIpos): m/z =922 (M+H)⁺.

Intermediate 71

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-3-[(2S)-1-methoxy-1-oxo-3-phenylpropan-2-yl]amino]-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3424]



[3425] First, in analogy to the synthesis described in Intermediate 14, proceeding from Intermediates 4 and 39, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-3-[(2S)-1-methoxy-1-oxo-3-phenylpropan-2-yl]amino]-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared. 7 mg (0.009 mmol) of this compound were then used, in analogy to the preparation of Intermediate 61, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride, to obtain 2 mg (22% of theory) of the title compound in the form of a colourless foam.


[3426] HPLC (Method 6): R_f =1.9 min;

[3427] LC-MS (Method 2): R_f =1.06 min; MS (ESIpos): m/z =832 (M+H)⁺.

Intermediate 72

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3428]



[3429] 212 mg (411 μ mol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 8) and 237 mg (411 mmol) of benzyl-N-[(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoyl]-L-tryptophan trifluoroacetate (Intermediate 20) were taken up in 30 ml of DMF, and 188 mg (493 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and 215 μ l N,N-diisopropylethylamine were added. The reaction mixture was stirred at RT for 20 h, then concentrated under reduced pressure, and the residue was purified by means of preparative HPLC. The product fractions were combined and concentrated, and the residue was dried under high vacuum. This gave 315 mg (80% of theory) of the Boc-protected intermediate N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino)-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide as a colourless foam.

[3430] LC-MS (Method 1): R_t =1.45 min; m/z =961 (M+H)⁺.

[3431] 50 mg (52 μ mol) of this intermediate were treated with 1 ml of trifluoroacetic acid in 9 ml of dichloromethane to detach the Boc protecting group. After concentration and purification by means of preparative HPLC, 29 mg (57% of theory) of the free amine intermediate N-methyl-L-valyl-N-

and about 300 μ l of 0.1 N hydrochloric acid were subsequently used to adjust the pH back to 3. The mixture was then stirred at 100° C. for another 2 h. Thereafter, the reaction mixture was poured onto a 1:1 mixture of semisaturated aqueous ammonium chloride solution and ethyl acetate. The organic phase was removed, washed with saturated sodium chloride solution, dried over sodium sulphate and concentrated. The residue was freeze-dried from water/acetonitrile. This gave 27 mg (94% of theory) of the title compound in the form of a colourless foam.


[3434] HPLC (Method 5): R_t =2.2 min;

[3435] LC-MS (Method 9): R_t =5.04 min; MS (ESIpos): m/z =947 (M+H)⁺.

Intermediate 73

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-[benzyl(methyl)amino]-1-oxo-3-phenylpropan-2-yl]amino)-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3436]



[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino)-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were obtained as the trifluoroacetate.

[3432] LC-MS (Method 1): R_t =0.99 min; m/z =861 (M+H)⁺.

[3433] 29 mg (0.03 mmol) of this intermediate were dissolved in 6 ml of dioxane/water, and 39 μ l (0.059 mmol) of a 15% aqueous solution of 4-oxobutanoic acid were added. The reaction mixture was subsequently stirred at 100° C. for 1 h. After cooling to RT, 2 mg (0.033 mmol) of sodium cyanoborohydride were added and the mixture was adjusted to a pH of 3 by adding about 300 μ l of 0.1 N hydrochloric acid. The reaction mixture was then stirred at 100° C. for a further 2 h. After cooling, another 39 μ l (0.059 mmol) of the 15% 4-oxobutanoic acid solution were added and the reaction mixture was again stirred at 100° C. for 1 h. Then a further 2 mg (0.033 mmol) of sodium cyanoborohydride were added

[3437] First, in analogy to the synthesis described in Intermediate 14, proceeding from Intermediates 4 and 38, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-[benzyl(methyl)amino]-1-oxo-3-phenylpropan-2-yl]amino)-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared. 25 mg (0.026 mmol) of this compound were then used, in analogy to the preparation of Intermediate 61, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride, to obtain 13 mg (54% of theory) of the title compound in the form of a colourless foam.


[3438] HPLC (Method 12): R_t =2.2 min;

[3439] LC-MS (Method 9): R_t =5.01 min; MS (ESIpos): m/z =921 (M+H)⁺.

Intermediate 74

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{(1S,2R)-1-[(benzyloxy)carbonyl]-2-phenylcyclopropyl}amino)-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3440]



[3441] 50 mg (73 μ mol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) and 28 mg (73 μ mol) of benzyl (1S,2R)-1-amino-2-phenylcyclopropanecarboxylate trifluoroacetate (Intermediate 45) were taken up in ml of DMF, and 42 mg (110 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and 38 μ l of N,N-diisopropylethylamine were added. The reaction mixture was stirred at RT for 5 h, then concentrated under reduced pressure, and the residue was purified by means of preparative HPLC. The product fractions were combined and concentrated. After lyophilization from dioxane/water, 35 mg (51% of theory) of the Boc-protected intermediate N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{(1S,2R)-1-[(benzyloxy)carbonyl]-2-phenylcyclopropyl}amino)-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were obtained as a colourless foam.

[3442] LC-MS (Method 1): R_t =1.52 min; m/z =934 (M+H)⁺.

[3443] 35 mg of this intermediate were treated with 1 ml of trifluoroacetic acid in 5 ml of dichloromethane to detach the Boc protecting group. After concentration and lyophilization from dioxane/water, 34 mg (97% of theory) of the free amine

intermediate N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{(1S,2R)-1-[(benzyloxy)carbonyl]-2-phenylcyclopropyl}amino)-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were obtained as the trifluoroacetate.

[3444] LC-MS (Method 1): R_t =0.91 min; m/z =834 (M+H)⁺.

[3445] 11 mg (0.011 mmol) of this intermediate were then used, in analogy to the preparation of Intermediate 66, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride, to obtain 2.5 mg (24% of theory) of the title compound in the form of a colourless foam.


[3446] HPLC (Method 12): R_t =2.2 min;

[3447] LC-MS (Method 9): R_t =5.1 min; MS (ESIpos): m/z =920 (M+H)⁺.

Intermediate 75

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-{(1S,2R)-2-phenyl-1-(propylcarbamoyl)cyclopropyl}amino}propyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3448]



[3449] First, in analogy to the synthesis described in Intermediate 74, by coupling of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) and (1S,2R)-1-amino-2-phenyl-N-propylcyclopropanecarboxamide trifluoroacetate (Intermediate 27) in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Boc protecting group by means of trifluoroacetic acid, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S,2R)-2-phenyl-1-(propylcarbamoyl)cyclopropyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate. 14 mg (0.016 mmol) of this compound were

then used, in analogy to the preparation of Intermediate 61, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride, to obtain 11.3 mg (83% of theory) of the title compound.


[3450] HPLC (Method 6): R_f =1.9 min;

[3451] LC-MS (Method 2): R_f =1.27 min; MS (ESIpos): m/z =871 (M+H)⁺.

Intermediate 76

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S,2R)-1-(ethoxycarbonyl)-2-phenylcyclopropyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3452]



[3453] First, by coupling of Intermediate 46 (N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide) with Intermediate 48 (ethyl (1S,2R)-1-amino-2-phenylcyclopropanecarboxylate trifluoroacetate) in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent Boc detachment, the starting compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S,2R)-1-(ethoxycarbonyl)-2-phenylcyclopropyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate was prepared. 70 mg (0.079 mmol) of this starting material were then used, by reaction with 4-oxobutanoic acid, in analogy to Intermediate 61, to obtain 46 mg (68% of theory) of the title compound.


[3454] HPLC (Method 6): R_f =1.9 min;

[3455] LC-MS (Method 2): R_f =1.28 min; MS (ESIpos): m/z =858 (M+H)⁺.

Intermediate 77

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-amino-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3456]



[3457] First, in analogy to the synthesis described in Intermediate 75, by coupling of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) and L-phenylalaninamide hydrochloride in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Boc protecting group by means of trifluoroacetic acid, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S)-1-amino-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate. 47 mg (0.049 mmol) of this compound were then used, in analogy to the preparation of Intermediate 61, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride, to obtain 39 mg (96% of theory) of the title compound.

[3458] HPLC (Method 6): $R_f=1.7$ min;


[3459] LC-MS (Method 9): $R_f=4.44$ min; MS (ESIpos): $m/z=817$ (M+H)⁻¹

[3460] ¹H NMR (500 MHz, DMSO-d₆): $\delta=8.95$ and 8.8 (2m, 1H), 8.25 and 8.0 (2d, 1H), 7.45 , 7.35 and 7.0 (3s, broad, 2H), $7.3-7.1$ (m, 5H), $4.8-4.4$ (2m, 3H), 3.95 (m, 1H), 3.82 (m, 1H), 3.72 (d, 1H), 3.22 , 3.18 , 3.15 , 3.05 and 3.00 (5s, 9H), $2.85-2.7$ (m, 4H), $2.45-1.6$ (m, 12H), $1.5-1.2$ (m, 3H), $1.1-0.7$ (m, 21H) [further signals hidden under solvent peaks].

Intermediate 78

N-(6-aminoheptyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3461]



[3462] This compound was prepared in analogy to Intermediate 66 over 2 stages, proceeding from 20 mg (16 μ mol) of the compound from Intermediate 14 and benzyl 6-oxohexyl carbamate, and the hydrogenation was performed in methanol as the solvent.

[3463] Yield: 7.6 mg (55% of theory over 2 stages)


[3464] HPLC (Method 6): $R_f=1.8$ min;

[3465] LC-MS (Method 1): $R_f=0.7$ min; MS (ESIpos): $m/z=901$ (M+H)⁺.

Intermediate 79

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S)-1-(benzylamino)-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3466]



[3467] 36 mg (43 μ mol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S)-1-carboxy-2-phenylethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 46) and 4.6 mg (43 μ mol) of benzylamine were taken up in 5 ml of DMF, 7.5 μ l (88 μ mol) of N,N-diisopropylethylamine, 10 mg (65 μ mol) of HOBt and 10 mg (52 μ mol) of EDC were added, and then the mixture was stirred at RT overnight. Subsequently, the reaction mixture was concentrated and the residue was purified by means of preparative HPLC. 29 mg (73% of theory) of the Boc-protected intermediate N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzylamino)-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were obtained.

[3468] LC-MS (Method 1): R_f =1.43 min; m/z =921 (M+H)⁺.

[3469] 29 mg of this intermediate were treated with 1 ml of trifluoroacetic acid in 6 ml of dichloromethane to detach the Boc protecting group. After concentration and lyophilization from dioxane/water, 30 mg (quant.) of the free amine intermediate N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,

2R)-3-[(2S)-1-(benzylamino)-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were obtained as the trifluoroacetate.

[3470] LC-MS (Method 1): R_f =0.95 min; m/z =821 (M+H)⁺.

[3471] 17 mg (0.018 mmol) of this intermediate were then used, in analogy to the preparation of Intermediate 61, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride, to obtain 13 mg (80% of theory) of the title compound in the form of a colourless foam.


[3472] HPLC (Method 5): R_f =1.7 min;

[3473] LC-MS (Method 9): R_f =4.97 min; MS (ESIpos): m/z =907 (M+H)⁺.

Intermediate 80

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzylamino)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3474]



[3475] First, in analogy to the synthesis described in Intermediate 74, by coupling of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) and N-benzyl-L-tryptophanamide trifluoroacetate (Intermediate 47) in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Boc protecting group by means of trifluoroacetic acid, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzylamino)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate. 10 mg (0.01 mmol) of this compound were then used, in analogy to the preparation of Intermediate 61, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride, to obtain 2.5 mg (26% of theory) of the title compound.


[3476] HPLC (Method 5): R_f =1.7 min;

[3477] LC-MS (Method 2): R_f =1.13 min; MS (ESIpos): m/z =946 (M+H)⁺.

Intermediate 81

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S,2R)-1-carbamoyl-2-phenylcyclopropyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3478]



[3479] First, in analogy to the synthesis described in Intermediate 74, by coupling of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) and (1S,2R)-1-amino-2-phenylcyclopropanecarboxamide trifluoroacetate (Intermediate 48) in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Boc protecting group by means of trifluoroacetic acid, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S,2R)-1-carbamoyl-2-phenylcyclopropyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate. 14 mg (0.0163 mmol) of this compound were then used, in analogy to the preparation of Intermediate 61, by reaction with 4-ox-

obutanoic acid in the presence of sodium cyanoborohydride, to obtain 8 mg (57% of theory) of the title compound.


[3480] HPLC (Method 5): R_f =1.6 min;

[3481] LC-MS (Method 9): R_f =4.64 min; MS (ESIpos): m/z =829 (M+H)⁺.

Intermediate 82

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3482]



[3483] First, in analogy to the synthesis described in Intermediate 69, by coupling of N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 4) and N^α-{(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoyl}-L-tryptophanamide trifluoroacetate (Intermediate 49) in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Fmoc protecting group by means of piperidine, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[[[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate. 78 mg (0.088 mmol) of this compound were then used, in anal-

ogy to the preparation of Intermediate 61, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride, to obtain 68 mg (90% of theory) of the title compound.


[3484] HPLC (Method 5): R_t=1.8 min;

[3485] LC-MS (Method 9): R_t=4.49 min; MS (ESIpos): m/z=856 (M+H)⁺.

Intermediate 83

N-(5-carboxypentyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[[[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3486]



[3487] This compound was prepared in analogy to the compound in Intermediate 82, proceeding from 20 mg (26 μmol) of N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[[[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride prepared.

[3488] Yield: 5 mg (25% of theory)


[3489] HPLC (Method 5): R_t=1.6 min;

[3490] LC-MS (Method 11): R_t=0.72 min; MS (ESIpos): m/z=884 (M+H)⁺.

Intermediate 84

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(2S)-1-(morpholin-4-yl)-1-oxo-3-phenylpropan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3491]



[3492] First, in analogy to the synthesis described in Intermediate 79, by coupling of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S)-1-carboxy-2-phenylethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 46) and morpholine in the presence of EDC and HOBT and subsequent detachment of the Boc protecting group by means of trifluoroacetic acid, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(morpholin-4-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate. 30 mg (0.033 mmol) of this compound were then used, in analogy to the preparation of Intermediate 61, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride, to obtain 22 mg (76% of theory) of the title compound.


[3493] HPLC (Method 5): $R_f=1.6$ min;

[3494] LC-MS (Method 9): $R_f=4.58$ min; MS (ESIpos): $m/z=887$ (M+H)⁺.

Intermediate 85

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S,3R)-1-(benzylamino)-3-hydroxy-1-oxobutan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3495]




[3496] First, in analogy to the synthesis described in Intermediate 79, by coupling of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S)-1-carboxy-2-phenylethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 46) and N-benzyl-L-threoninamide trifluoroacetate in the presence of HATU and subsequent detachment of the Boc protecting group by means of trifluoroacetic acid, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S,3R)-1-(benzylamino)-3-hydroxy-1-oxobutan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate. 21 mg (0.024 mmol) of this compound were then used, in analogy to the preparation of Intermediate 61, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride, to obtain 20 mg (97% of theory) of the title compound.

[3497] HPLC (Method 5): $R_f=1.54$ min;

[3498] LC-MS (Method 9): $R_f=4.49$ min; MS (ESIpos): $m/z=861$ (M+H)⁺.

Intermediate 86

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-tert-butoxy-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide
[3499]



[3500] First, in analogy to the synthesis described in Intermediate 74, by coupling of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) and tert-butyl-L-phenylalaninate hydrochloride in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Boc protecting group by means of trifluoroacetic acid to obtain the tert-butyl ester (stirring with trifluoroacetic acid in dichloromethane for 40 minutes), the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-tert-butoxy-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate. 22 mg (0.02 mmol) of this compound were then used, in analogy to the preparation

of Intermediate 61, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride, to obtain 16 mg (94% of theory) of the title compound.


[3501] HPLC (Method 5): R_f =2.0 min;

[3502] LC-MS (Method 9): R_f =5.05 min; MS (ESIpos): m/z =874 (M+H)⁺.

Intermediate 87

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-tert-butoxy-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3503]



[3504] This compound was prepared in analogy to the synthesis described in Intermediate 86, proceeding from 230 mg (336 μmol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) and tert-butyl-L-tryptophanate hydrochloride over 3 stages.

[3505] Yield: 95 mg (31% of theory over 3 stages)


[3506] HPLC (Method 5): $R_f=2.0$ min;

[3507] LC-MS (Method 9): $R_f=5.05$ min; MS (ESIpos): $m/z=913$ (M+H)⁺.

Intermediate 88

N-(6-aminoheptyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3508]



[3509] First, in analogy to the syntheses described in Intermediate 69, by coupling of N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 4) and N^α-{(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoyl}-L-tryptophanamide trifluoroacetate (Intermediate 49) in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Fmoc protecting group by means of piperidine, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate. 30 mg (0.03 mmol) of this compound were then used, in analogy to the preparation of the compound of Intermediate 61, by reaction with benzyl 6-oxohexyl carbamate, which had been

obtained beforehand by oxidation of benzyl 6-hydroxyhexyl carbamate, in the presence of sodium cyanoborohydride, to obtain 17 mg (45% of theory) of the Z-protected compound. Subsequently, hydrogenolysis in methanol over 10% palladium/activated carbon afforded the title compound.

[3510] Yield: 14 mg (95% of theory)


[3511] HPLC (Method 5): $R_f=1.5$ min;

[3512] LC-MS (Method 1): $R_f=0.73$ min; MS (ESIpos): $m/z=869$ (M+H)⁺.

Intermediate 89

N-(6-aminoheptyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-tert-butoxy-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3513]



[3514] First, in analogy to the synthesis described in Intermediate 86, by coupling of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) and tert-butyl-L-tryptophanate hydrochloride in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Boc protecting group by means of trifluoroacetic acid to obtain the tert-butyl ester (stirring with 1:10 trifluoroacetic acid/dichloromethane for 30 min), the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-tert-butoxy-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate. 71 mg (0.075 mmol) of this compound were then used, in analogy to the preparation of the compound of Intermediate 61, by reaction with benzyl 6-oxohexyl carbamate, which had been obtained

beforehand by oxidation of benzyl 6-hydroxyhexyl carbamate, in the presence of sodium cyanoborohydride, to obtain 35 mg (44% of theory) of the Z-protected compound. Subsequently, hydrogenolysis in methanol over 10% palladium/activated carbon afforded the title compound.

[3515] Yield: 30 mg (98% of theory)


[3516] HPLC (Method 5): $R_t=1.9$ min;

[3517] LC-MS (Method 1): $R_t=0.77$ min; MS (ESIpos): $m/z=926$ (M+H)⁺.

Intermediate 90

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[2-(1H-indol-3-yl)ethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3518]



[3519] First, in analogy to the synthesis described in Intermediate 74, by coupling of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) and 2-(1H-indol-3-yl)ethanamine in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Boc protecting group by means of trifluoroacetic acid, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[2-(1H-indol-3-yl)ethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate. 100 mg (0.119 mmol) of this compound were then used, in analogy to the preparation of Intermediate 61, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride, to obtain 50 mg (49% of theory) of the title compound. The title compound was purified here by flash chromatography on silica gel with dichloromethane/methanol/17% ammonia as the eluent, in the course of which the mixing ratio was switched from initially 15/2/0.2 to 15/4/0.5.


[3520] HPLC (Method 6): $R_t=1.8$ min;

[3521] LC-MS (Method 1): $R_t=0.87$ min; MS (ESIpos): $m/z=813$ (M+H)⁺.

Intermediate 91

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2-phenylethyl)amino]propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3522]



[3523] First, in analogy to the synthesis described in Intermediate 74, by coupling of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) and phenylethylamine in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Boc protecting group by means of trifluoroacetic acid, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2-phenylethyl)amino]propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate. 57 mg (0.071 mmol) of this compound were then used, in analogy to the preparation of Intermediate 61, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohy-

dride, to obtain 44 mg (80% of theory) of the title compound. The title compound can also be purified here by flash chromatography on silica gel with dichloromethane/methanol/17% ammonia as the eluent (15/2/0.2->15/4/0.5).


[3524] HPLC (Method 5): R_f =1.7 min;

[3525] LC-MS (Method 9): R_f =4.64 min; MS (ESIpos): m/z =774 (M+H)⁺.

Intermediate 92

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S,2R)-1-hydroxy-1-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3526]



[3527] 100 mg (0.139 mmol) of N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S,2R)-1-hydroxy-1-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 40) were used, in analogy to the preparation of Intermediate 61, by reaction with 4-oxobutanoic acid in the presence of sodium cyanoborohydride, to obtain 94 mg (84% of theory) of the title compound. The title compound was purified by preparative HPLC.


[3528] HPLC (Method 5): R_f =1.5 min;

[3529] LC-MS (Method 9): R_f =4.46 min; MS (ESIpos): m/z =804 (M+H)⁺.

Intermediate 93

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3530]



[3531] 22.4 mg (24 μ mol) of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate were dissolved in 1.4 ml of dioxane/water and, analogously to the preparation of Intermediate 61, reacted with 15% aqueous solution of 4-oxobutanoic acid in the presence of sodium cyanoborohydride. After lyophilization from dioxane, 8.2 mg (38% of theory) of the title compound were obtained in the form of a white solid.


[3532] HPLC (Method 10): R_f =2.54 min

[3533] LC-MS (Method 12): R_f =0.94 min; MS (ESIpos): m/z =919 (M+H)⁺

Intermediate 94

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1R)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3534]



[3535] 17.1 mg (18 μ mol) of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[[[(1R)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino]propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate were dissolved in 1.1 ml of dioxane/water and, analogously to the preparation of Intermediate 61, reacted with 15% aqueous solution of 4-oxobutanoic acid in the presence of sodium cyanoborohydride. After lyophilization from dioxane, 14.8 mg (89% of theory) of the title compound were obtained in the form of a white solid.


[3536] HPLC (Method 10): R_f =2.54 min;

[3537] LC-MS (Method 12): R_f =0.92 min; MS (ESIpos): m/z =919 (M+H)+

Intermediate 95

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzylsulphonyl)-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3538]




[3539] 19.3 mg (20 μ mol) N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzylsulphonyl)-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate were dissolved in 1.2 ml of dioxane/water and, analogously to the preparation of Intermediate 61, reacted with 15% aqueous solution of 4-oxobutanoic acid in the presence of sodium cyanoborohydride. After lyophilization from dioxane, 8.6 mg (45% of theory) of the title compound were obtained in the form of a solid.

[3540] LC-MS (Method 11): R_f =0.85 min; MS (ESIpos): m/z =943 (M+H)+

Intermediate 96

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S,3E)-1,4-diphenylbut-3-en-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3541]



[3542] 15.5 mg (10 μ mol) of N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S,3E)-1,4-diphenylbut-3-en-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate were dissolved in 1.0 ml of dioxane/water and, analogously to the preparation of Intermediate 61, reacted with 15% aqueous solution of 4-oxobutanoic acid in the presence of sodium cyanoborohydride. After lyophilization from dioxane, 10.3 mg (68% of theory) of the title compound were obtained in the form of a white solid.


[3543] HPLC (Method 10): R_t =2.59 min;

[3544] LC-MS (Method 11): R_t =0.94 min; MS (ESIpos): m/z =877 (M+H)+

Intermediate 97

N-(6-aminohexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3545]



[3546] The title compound was prepared in analogy to the synthesis of Intermediate 66, by reaction of 200 mg (0.108 mmol) of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate (Intermediate 16) with benzyl 6-oxohexyl carbamate and subsequent hydrogenolytic detachment of the Z protecting group (with 5% palladium on charcoal as a catalyst, in methanol as a solvent).

[3547] Yield: 69 mg (65% of theory over two stages)


[3548] HPLC (Method 5): R_t =1.7 min;

[3549] LC-MS (Method 1): R_t =0.76 min; MS (ESIpos): m/z =912 (M+H)+.

Intermediate 98

N-(5-carboxypentyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S)-1-(benzylamino)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3550]



[3551] This compound was prepared in analogy to the synthesis described in Intermediate 80. The purification was effected by preparative HPLC.

[3552] Yield: 40 mg (29% of theory over 3 stages)


[3553] HPLC (Method 5): $R_f=1.9$ min;

[3554] LC-MS (Method 1): $R_f=0.92$ min; MS (ESIpos): $m/z=974$ (M+H)⁺.

Intermediate 99

(2S)-2-amino-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)propan-1-one trifluoroacetate

[3555]



[3559] Using 166 mg (444.5 μ mol) of this intermediate, under standard conditions with 3 ml of trifluoroacetic acid in 20 ml of dichloromethane, the Boc protecting group was detached and, after HPLC purification, 155 mg (86% of theory) of the title compound were obtained.


[3560] HPLC (Method 12): $R_f=1.43$ min;

[3561] LC-MS (Method 11): $R_f=0.56$ min; MS (ESIpos): $m/z=274$ (M+H)⁺.

Intermediate 100

N-(6-{[(benzyloxy)carbonyl]amino}hexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3562]



[3556] 324 mg (0.81 mmol) of 2,5-dioxopyrrolidin-1-yl N-(tert-butoxycarbonyl)-L-tryptophanate were dissolved in 20 ml of DMF, and 200 mg (1.62 mmol) of 1,2-oxazinan-2-ylpropan-1-one trifluoroacetate (Starting Compound 5) and 850 μ l of N,N-diisopropylethylamine were added. The reaction mixture was stirred at 50° C. overnight and then concentrated under reduced pressure. The residue was taken up in dichloromethane and extracted with water. The organic phase was dried over magnesium sulphate and concentrated. The residue was purified by flash chromatography on silica gel with 4:1 dichloromethane/ethyl acetate as the eluent. The product fractions were concentrated and the residue was dried under high vacuum. This gave 147.5 mg (48% of theory) of the Boc-protected intermediate.

[3557] HPLC (Method 12): $R_f=1.9$ min;

[3558] LC-MS (Method 1): $R_f=1.03$ min; MS (ESIpos): $m/z=374$ (M+H)⁺.

[3563] 177 mg (260 μ mol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) and 100 mg (260 μ mol) of (2S)-2-amino-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)propan-1-one trifluoroacetate (Intermediate 99) were taken up in 15 ml of DMF, and 118 mg (310 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and 140 μ l of N,N-diisopropylethylamine were added. The reaction mixture was stirred at RT for 30 min, then concentrated under reduced pressure, and the residue was purified by means of preparative HPLC. The product fractions were combined and concentrated. After lyophilization from dioxane, 170 mg (68% of theory) of the Boc-protected intermediate were obtained.

[3564] LC-MS (Method 1): $R_f=1.36$ min; $m/z=940$ (M+H)⁺.

[3565] 170 mg of this intermediate were treated with 3 ml of trifluoroacetic acid in 30 ml of dichloromethane for 30 min

to detach the Boc protecting group. Then the reaction mixture was concentrated under reduced pressure and the residue was purified by means of preparative HPLC to obtain 155 mg (86% of theory) of the deprotected N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide intermediate.

[3566] HPLC (Method 12): $R_t=1.85$ min;

[3567] LC-MS (Method 1): $R_t=0.86$ min; MS (ESIpos): $m/z=840$ (M+H)⁺.

[3568] 50 mg (0.052 mmol) of this intermediate were then used, in analogy to the preparation of Intermediate 97, with benzyl 6-oxohexyl carbamate in the presence of sodium cyanoborohydride and subsequent hydrogenolytic detach-

ment of the Z protecting group (with 5% palladium on charcoal as a catalyst, in methanol as a solvent), prepared to prepare the title compound.

[3569] Yield: 21 mg (37% of theory)


[3570] HPLC (Method 12): $R_t=2.1$ min;

[3571] LC-MS (Method 1): $R_t=1.02$ min; MS (ESIpos): $m/z=1073$ (M+H)⁺.

Intermediate 101

N-(6-aminoheptyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3572]



[3573] 26.7 mg (24.87 μ mol) of Intermediate 100 were dissolved in 10 ml of methanol and hydrogenated over palladium/activated carbon (5%) under standard hydrogen pressure for 30 min. The catalyst was filtered off and the solvent was evaporated off under reduced pressure. After the residue had been dried under high vacuum, 22.5 mg (96% of theory) of the title compound were obtained.


[3574] HPLC (Method 5): $R_t=1.7$ min;

[3575] LC-MS (Method 1): $R_t=0.76$ min; MS (ESIpos): $m/z=939$ (M+H)⁺.

Intermediate 102

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(morpholin-4-yl)-1-oxo-3-phenylpropan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3576]



[3577] This compound was prepared in analogy to the synthesis described in Intermediate 157 from N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(2S)-1-(morpholin-4-yl)-1-oxo-3-phenylpropan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide and commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide.

[3578] Yield: 8 mg (71% of theory)


[3579] HPLC (Method 12): R_f =1.9 min;

[3580] LC-MS (Method 1): R_f =0.87 min; MS (ESIpos): m/z =1094 (M+H)⁺.

Intermediate 103

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S,3R)-1-(benzylamino)-3-hydroxy-1-oxobutan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3581]



[3582] This compound was prepared in analogy to the synthesis described in Intermediate 157 from N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S,3R)-1-(benzylamino)-3-hydroxy-1-oxobutan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide and commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide.

[3583] Yield: 3 mg (22% of theory)


[3584] HPLC (Method 5): R_f =1.6 min;

[3585] LC-MS (Method 1): R_f =0.78 min; MS (ESIpos): m/z =1069 (M+H)⁺.

Intermediate 104

N-{4-[(trans-4-{[(2,5-dioxopyrrolidin-1-yl)oxy]carbonyl}cyclohexyl)amino]-4-oxobutyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3586]



[3587] First, benzyl trans-4-aminocyclohexanecarboxylate trifluoroacetate was prepared from trans-4-aminocyclohexanecarboxylic acid by introducing the Boc protecting group, then introducing the benzyl ester protecting group and subsequently detaching the Boc protecting group by conventional peptide chemistry methods.

[3588] 15 mg (18 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were then dissolved in 5 ml of dimethylformamide and subsequently admixed with 13 mg (35 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, 9 μ l of N,N-diisopropylethylamine and with 15 mg (44 μ mol) of benzyl trans-4-aminocyclohexanecarboxylate trifluoroacetate. The mixture was stirred at RT for 1 h and then concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. The corresponding fractions were combined and the solvent was evaporated off under reduced pressure. After the residue had been dried under high vacuum, 14.7 mg (78% of theory) of the protected intermediate were obtained as a colourless foam.

[3589] HPLC (Method 6): R_f =2.0 min;

[3590] LC-MS (Method 1): R_f =0.95 min; MS (ESIpos): m/z =1072 (M+H)⁺.

[3591] From this protected intermediate, the benzyl ester was first removed by hydrogenolytic means and the free carboxyl component was obtained in quantitative yield. 14 mg (14 μ mol; 1 equiv.) of the deprotected compound were taken

up in 5 ml of DMF and admixed with 3.3 mg (29 μ mol; 2.1 equiv.) of N-hydroxysuccinimide in the presence of 4.1 mg (21 μ mol; 1.5 equiv.) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 7.5 μ l (44 μ mol; 3.1 equiv.) of N,N-diisopropylethylamine and 0.9 mg (7 μ mol; 0.5 equiv.) of 4-dimethylaminopyridine, and the mixture was stirred at RT overnight. Then another 10 equiv. of N-hydroxysuccinimide, 5 equiv. of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 5 equiv. of N,N-diisopropylethylamine and 0.5 equiv. of 4-dimethylaminopyridine were added and the reaction mixture was treated in an ultrasound bath for 5 h. Subsequently, the solvent was evaporated off, the residue was purified by means of preparative HPLC and the corresponding fractions were combined and concentrated. After lyophilization of the residue from dioxane, 9.7 mg (62% of theory) of the title compound were obtained as a colourless foam.


[3592] HPLC (Method 6): R_f =1.8 min;

[3593] LC-MS (Method 11): R_f =0.77 min; MS (ESIpos): m/z =1078 (M+H)⁺.

Intermediate 105

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S)-1-carboxy-2-phenylethyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3594]



[3595] This compound was prepared in analogy to the synthesis described in Intermediate 157, proceeding from 4-[(2S)-1-[(2S)-1-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-tert-butoxy-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl](methyl)amino}-3-methylbutan-2-yl]amino]-3-methyl-1-oxobutan-2-yl](methyl)amino}butanoic acid and commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide. The ester intermediate was obtained in 42% yield. In a second step, 6 mg (6 μ mol) of this intermediate were cleaved with trifluoroacetic acid the tert-butyl ester. After HPLC purification, 3.4 mg (48% of theory) of the title compound were obtained.


[3596] HPLC (Method 5): R_f =1.66 min;

[3597] LC-MS (Method 2): R_f =1.04 min; MS (ESIpos): m/z =1025 (M+H)⁺.

Intermediate 106

N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3598]



[3599] 14 mg (16 μ mol) of N-(6-aminoheptyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 88) were taken up in 750 μ l of dioxane and admixed with 1.5 ml of saturated sodium hydrogencarbonate solution and then with 3.2 mg (21 μ mol) of methyl 2,5-dioxo-2,5-dihydro-1H-pyrrole-1-carboxylate. The reaction mixture was stirred at RT for 1 h and then concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization, 5.5 mg (36% of theory) of the title compound were obtained.


[3600] HPLC (Method 5): R_t =1.7 min;

[3601] LC-MS (Method 1): R_t =0.84 min; MS (ESIpos): m/z =949 (M+H)⁺.

Intermediate 107

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-amino-3-(1H-indol-3-yl)ethyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3602]



[3603] 38 mg (47 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-L-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-{[2-(1H-indol-3-yl)ethyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 37 ml of DMF and then admixed with 71 mg (187 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, 33 μ l of N,N-diisopropylethylamine and with 37 mg (140 μ mol) of commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide. The mixture was stirred at RT for 1 h. This was followed by concentration under high vacuum and purification of the remaining residue by means of preparative HPLC. Thus, 12.2 mg (26% of theory) of the title compound were obtained as a colourless foam.


[3604] HPLC (Method 5): R_f =1.6 min;

[3605] LC-MS (Method 1): R_f =0.85 min; MS (ESIpos): m/z =1020 (M+H)⁺.

Intermediate 108

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2-phenylethyl)amino]propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3606]



[3607] The compound was prepared in analogy to Intermediate 107.

Yield: 2.5 mg (30% of theory)


[3608] HPLC (Method 12): R_f =1.9 min;

[3609] LC-MS (Method 1): R_f =0.9 min; MS (ESIpos): m/z =981 (M+H)⁺.

Intermediate 109

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S,2R)-1-hydroxy-1-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3610]



[3611] The compound was prepared in analogy to Intermediate 107 from the compound in Intermediate 92.

[3612] Yield: 35 mg (65% of theory)


[3613] HPLC (Method 5): $R_f=1.9$ min;

[3614] LC-MS (Method 11): $R_f=0.76$ min; MS (ESIpos): $m/z=1011$ (M+H)⁺.

Intermediate 110

N-{6-[(2,5-dioxopyrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3615]



[3616] This compound was prepared in analogy to Intermediate 147 from the compound in Intermediate 83.

[3617] Yield: 2.4 mg (24% of theory)


[3618] HPLC (Method 6): $R_f=1.8$ min;

[3619] LC-MS (Method 1): $R_f=0.87$ min; MS (ESIpos): $m/z=981$ (M+H)⁺.

Intermediate 111

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]-1-methylhydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3620]



[3621] This compound was prepared in analogy to Intermediate 140 from Intermediate 82 and Intermediate 22.

[3622] Yield: 6.5 mg (51% of theory)


[3623] HPLC (Method 6): R_t =1.8 min;

[3624] LC-MS (Method 1): R_t =4.71 min; MS (ESIpos): m/z =1077 (M+H)⁺.

Intermediate 112

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(1S,2R)-1-carbamoyl-2-phenylcyclopropyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3625]



[3626] This compound was prepared in analogy to Intermediate 157 from the compound in Intermediate 81.

[3627] Yield: 5.7 mg (57% of theory)


[3628] HPLC (Method 5): R_t =1.6 min;

[3629] LC-MS (Method 1): R_t =0.87 min; MS (ESIpos): m/z =1036 (M+H)⁺.

Intermediate 113

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(1S)-1-carboxy-2-(1H-indol-3-yl)ethyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3630]



[3631] 95 mg (104 μ mol) of 4-{[(2S)-1-{[(2S)-1-{[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-tert-butoxy-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl](methyl)amino}-3-methylbutan-2-yl]amino}-3-methyl-1-oxobutan-2-yl](methyl)amino}butanoic acid were dissolved in DMF and then admixed with 79.5 mg (209 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetram-

ethyluronium hexafluorophosphate, 73 μ l of N,N-diisopropylethylamine and with 68 mg (261 μ mol) of commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide. The mixture was stirred at RT for 2 h. This was followed by concentration under high vacuum and purification of the remaining residue by means of preparative HPLC. Thus, 104 mg (89% of theory) of the tert-butyl ester of the title compound were obtained as a colourless foam.

[3632] HPLC (Method 5): $R_t=2.0$ min;

[3633] LC-MS (Method 1): $R_t=0.93$ min; MS (ESIpos): $m/z=1121$ (M+H)⁺.

[3634] The intermediate was taken up in 33.4 ml of dichloromethane, 17 ml of trifluoroacetic acid were added, and the mixture was stirred at RT for 1 h. Subsequently, the reaction mixture was concentrated under reduced pressure and the residue was purified by preparative HPLC.

[3635] Thus, 61 mg (62% of theory) of the title compound were obtained as a colourless foam.


[3636] HPLC (Method 5): $R_t=1.7$ min;

[3637] LC-MS (Method 1): $R_t=0.86$ min; MS (ESIpos): $m/z=1064$ (M+H)⁺.

Intermediate 114

N-[6-({[2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethyl]carbamoyl}amino)hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3638]



[3639] 5 mg (5 μ mol) of N-(6-aminoheptyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were taken up in 885 μ l of DMF and admixed with 5.3 mg (8 μ mol) of 4-nitrophenyl 2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethyl carbamate and 2.8 μ l of N,N-diisopropylethylamine. The reaction mixture was stirred at RT for 2 h and then concentrated to dryness. The residue was purified by means of preparative HPLC.

[3640] Yield: 0.58 mg (11% of theory) of a colourless foam


[3641] HPLC (Method 5): $R_t=1.6$ min;

[3642] LC-MS (Method 1): $R_t=0.83$ min; MS (ESIpos): $m/z=1035$ (M+H)⁺.

Intermediate 115

N-{4-[(2,5-dioxopyrrolidin-1-yl)oxy]-4-oxobutyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-yl)carbonyl]-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3643]



[3644] This compound was prepared in analogy to the compound in Intermediate 147, proceeding from 8 mg (9 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide.

After concentration, the activated ester was purified by means of preparative HPLC and, after removal of the solvent under reduced pressure, reacted immediately with the antibody.

[3645] Yield: 3 mg (27% of theory) (hydrolysis-sensitive)


[3646] HPLC (Method 5): R_f =1.7 min;

[3647] LC-MS (Method 1): R_f =0.87 min; MS (ESIpos): m/z =996 (M+H)⁺.

Intermediate 116

N-{4-[(2,5-dioxopyrrolidin-1-yl)oxy]-4-oxobutyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3648]



[3649] This compound was prepared in analogy to the compound in Intermediate 147, proceeding from 5 mg (6 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide.

After concentration, the activated ester was purified by means of preparative HPLC and, after removal of the solvent under reduced pressure, reacted immediately with the antibody.

[3650] Yield: 3.2 mg (43% of theory) (hydrolysis-sensitive)


[3651] HPLC (Method 5): R_f =1.7 min;

[3652] LC-MS (Method 1): R_f =0.92 min; MS (ESIpos): m/z =984 (M+H)⁺.

Intermediate 117

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-tert-butoxy-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3653]



[3654] This compound was prepared in analogy to Intermediate 157 from the compound in Intermediate 86.

[3655] Yield: 7 mg (42% of theory)


[3656] HPLC (Method 5): R_f =1.6 min;

[3657] LC-MS (Method 1): R_f =0.94 min; MS (ESIpos): m/z =1081 (M+H)⁺.

Intermediate 118

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2R)-1-(benzyloxy)-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3658]




[3659] The target compound was prepared analogously to Intermediate 157 from 7 mg (7.8 μ mol) of the compound in Intermediate 68. Yield: 6.3 mg (53% of theory)

[3660] LC-MS (Method 1): R_f =1.00 min; MS (ESIpos): m/z =1102 (M+H)⁺.

Intermediate 119

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-{[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3661]



[3662] 7.4 mg (8.1 mmol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-{[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-


valinamide and 6.3 mg (24.2 mmol) of 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide hydrochloride were coupled and worked up in analogy to Intermediate 157. 1.6 mg (13% of theory) of the title compound were obtained as a solid.

[3663] LC-MS (Method 11): $R_f=0.89$ min; MS (ESIpos): $m/z=1126$ (M+H)⁺

Intermediate 120

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1R)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3664]




[3665] 12.8 mg (13.9 mmol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1R)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide and 10.9 mg (41.8 mmol) of 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide hydrochloride were coupled and worked up in analogy to Intermediate 157. 10.8 mg (59% of theory) of the title compound were obtained as a solid.

[3666] LC-MS (Method 11): $R_f=0.90$ min; MS (ESIpos): $m/z=1126$ (M+H)⁺

Intermediate 121

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(2S)-1-(benzylsulphonyl)-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3667]




[3668] 7.4 mg (7.9 mmol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-(benzylsulphonyl)-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide and 6.2 mg (23.5 mmol) of 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide hydrochloride were coupled and worked up in analogy to Intermediate 157. 6.9 mg (74% of theory) of the title compound were obtained as a solid.

[3669] LC-MS (Method 11): R_t =0.87 min; MS (ESIpos): m/z =1150 (M+H)+

Intermediate 122

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(2S,3E)-1,4-diphenylbut-3-en-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide




[3671] 8 mg (9.1 mmol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[(2S,3E)-1,4-diphenylbut-3-en-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide and 7.2 mg (27.4 mmol) of 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanhydrazide hydrochloride were coupled and worked up in analogy to Intermediate 157. 8.2 mg (82% of theory) of the title compound were obtained as a white solid.

[3672] LC-MS (Method 11): $R_t=0.95$ min; MS (ESIpos): $m/z=1083$ (M+H)⁺

Intermediate 123

N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexyl]-
N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,
2R)-3-{[(2S)-1-tert-butoxy-3-(1H-indol-3-yl)-1-
oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-
oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-
oxoheptan-4-yl]N-methyl-L-valinamide



[3674] 30 mg (30 μ mol) of Intermediate 89 were taken up in 2 ml of 1,4-dioxane and admixed with 4 ml of saturated sodium hydrogencarbonate solution and then with 7.5 mg (50 μ mol) of methyl 2,5-dioxo-2,5-dihydro-1H-pyrrole-1-carboxylate. The reaction mixture was stirred at RT for 1 h and then concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization, 24 mg (74% of theory) of the title compound were obtained.


[3675] HPLC (Method 5): R_f =2.2 min;

[3676] LC-MS (Method 1): R_f =1.01 min; MS (ESIpos): m/z =1006 (M+H)⁺.

Intermediate 124

N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(1S)-1-carboxy-2-(1H-indol-3-yl)ethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3677]



[3678] 22 mg (20 μ mol) of Intermediate 123 were reacted with 4 ml of trifluoroacetic acid in 8 ml of dichloromethane at RT for 1 h. Thereafter, the reaction mixture was concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization, 11 mg (54% of theory) of the title compound were obtained.


[3679] HPLC (Method 5): R_f =1.8 min;

[3680] LC-MS (Method 11): R_f =0.85 min; MS (ESIpos): m/z =950 (M+H)⁺.

Intermediate 125

N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3681]



[3682] 22.5 mg (20 μ mol) of Intermediate 101 were taken up in 2 ml of 1:1 dioxane/water and then admixed with 5.6 mg (40 μ mol) of methyl 2,5-dioxo-2,5-dihydro-1H-pyrrole-1-carboxylate and with 0.25 ml of saturated sodium hydrogen-carbonate solution. The reaction mixture was stirred at RT for 30 min. Then another 0.25 ml of the saturated sodium hydrogencarbonate solution was added and the reaction mixture was stirred at RT for a further 15 min and then concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization, 12.8 mg (50% of theory) of the title compound were obtained as a colourless foam.


[3683] HPLC (Method 5): R_t =1.9 min;

[3684] LC-MS (Method 1): R_t =0.95 min; MS (ESIpos): m/z =1019 (M+H)⁺.

Intermediate 126

N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3685]



[3686] 64 mg (70 μ mol) of N-(6-aminoheptyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 97) were taken up in 3 ml of 1:1 dioxane/water, then adjusted to pH 9 with 4 ml of saturated sodium hydrogencarbonate solution and subsequently admixed with 16.3 mg (110 μ mol) of methyl 2,5-dioxo-2,5-dihydro-1H-pyrrole-1-carboxylate. The reaction mixture was stirred at RT for 1 h and then concentrated under reduced pressure. Then another 8 mg (55 μ mol) of methyl 2,5-dioxo-2,5-dihydro-1H-pyrrole-1-carboxylate were added, and the reaction mixture was

adjusted again to pH 9 and stirred at RT for a further hour. This was followed by concentration and purification of the remaining residue by means of preparative HPLC. At first, 31 mg of an as yet uncyclized intermediate were obtained. 27 mg of this intermediate were taken up again in 2 ml of 1:1 dioxane/water and then admixed with 250 μ l of saturated sodium hydrogencarbonate solution. After stirring at RT for 2 hours, the reaction mixture was concentrated and the residue was purified by means of preparative HPLC. After lyophilization, 20 mg (29% of theory) of the title compound were obtained.


[3687] HPLC (Method 5): R_t =1.96 min;

[3688] LC-MS (Method 1): R_t =0.97 min; MS (ESIpos): m/z =992 (M+H)⁺.

Intermediate 127

N-{6-[(2,5-dioxypyrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzylamino)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3689]



[3690] 17 mg (18 μ mol) of N-(5-carboxypentyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzylamino)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 98) were dissolved in 2.8 ml of dichloromethane and admixed with 20 mg (174 μ mol) of 1-hydroxypyrrolidine-2,5-dione and then with 10 mg (52 μ mol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 0.21 mg (0.17 μ mol) of DMAP. After stirring at RT for 4 h, the reaction mixture was concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization, 8.2 mg (43% of theory) of the title compound were obtained.


[3691] HPLC (Method 5): R_f =2.0 min;

[3692] LC-MS (Method 1): R_f =0.98 min; MS (ESIpos): m/z =1071 ($M+H$)⁺.

Intermediate 128

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3693]



[3694] 5 mg (5.6 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 845 μ l of DMF and then admixed with 3.2 mg (17 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 2.6 mg (17 μ mol) of 1-hydroxy-1H-benzotriazole hydrate, 1.96 μ l of N,N-diisopropylethylamine and with 5.9 mg (22.5 μ mol) of commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide. The mixture was stirred at RT overnight and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. Thus, 2.2 mg (36% of theory) of the title compound were obtained as a colourless foam.


[3695] HPLC (Method 5): R_f =1.7 min;

[3696] LC-MS (Method 1): $R_t=0.88$ min; MS (ESIpos): $m/z=1094$ (M+H)⁺.

Intermediate 129

N-(6-[2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino]-6-oxohexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazin-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl)-N-methyl-L-valinamide

[3697]



[3698] 4 mg (4.3 μ mol) of N-(5-carboxypentyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarboxyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 646 μ l of DMF and then admixed with 2.5 mg (13 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 2.0 mg (13 μ mol) of 1-hydroxy-1H-benzotriazole hydrate, 2.25 μ l of N,N-diisopropylethylamine

and with 4.5 mg (17 μ mol) of commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide. The mixture was stirred at RT for 3 h and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. Thus, 1.9 mg (39% of theory) of the title compound were obtained as a colourless foam.


[3699] HPLC (Method 5): R_t =1.7 min;

[3700] LC-MS (Method 9): R_t =4.9 min; MS (ESIpos): m/z =1134 (M+H)⁺.

Intermediate 130

N-(4-[(2R)-1-({5-[(2,5-dioxopyrrolidin-1-yl)oxy]-5-oxopentanoyl}amino)propan-2-yl]oxy)-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3701]



[3702] 10.5 mg (11.7 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 3.7 ml of dichloromethane and then admixed with 6.7 mg (35 μ mol) of 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride, 0.7 mg (5.8 mmol) of 4-dimethylaminopyridine and with 8.2 mg (47 μ mol) of commercially available tert-butyl (2R)-2-hydroxypropyl carbamate. The mixture was stirred at RT overnight and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. Thus, 7.5 mg (61% of theory) of the Boc-protected intermediate were obtained as a colourless foam.

[3703] HPLC (Method 5): R_f =2.0 min;

[3704] LC-MS (Method 1): R_f =1.03 min; MS (ESIpos): m/z =1056 (M+H)⁺.

[3705] Subsequently, the Boc protecting group was detached with trifluoroacetic acid. 4.9 mg (0.005 mmol) of the deprotected crude product were then, without further purification, taken up in 1.8 ml of dichloromethane and admixed

with 3.7 mg (0.011 mmol) of 1,1'-[(1,5-dioxopentane-1,5-diyl)bis(oxy)]dipyrrolidine-2,5-dione, 2.4 μ l (0.014 mmol) of N,N-diisopropylethylamine and 0.6 mg (5 μ mol) of 4-dimethylaminopyridine. The mixture was stirred at RT for 2 h and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. Thus, 0.77 mg (15% of theory) of the title compound were obtained as a colourless foam.


[3706] HPLC (Method 5): R_f =1.8 min;

[3707] LC-MS (Method 1): R_f =0.93 min; MS (ESIpos): m/z =1167 (M+H)⁺.

Intermediate 131

N-{4-[(1-{5-[(2,5-dioxopyrrolidin-1-yl)oxy]-5-oxopentanoyl}piperidin-4-yl)oxy]-4-oxobutyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3708]



[3709] 10 mg (11 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-L-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 2 ml of dichloromethane and then admixed with 4.3 mg (22 μ mol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 0.88 mg (6 mmol) of 4-dimethylaminopyridine and with 5.2 mg (22 μ mol) of commercially available benzyl 4-hydroxypiperidine-1-carboxylate. The mixture was stirred at RT overnight and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. Thus, 5 mg (40% of theory) of the Z-protected intermediate were obtained as a colourless foam.

[3710] HPLC (Method 5): R_f =2.1 min;

[3711] LC-MS (Method 1): R_t =1.04 min; MS (ESIpos): m/z =1116 (M+H)⁺.

[3712] Subsequently, the Z protecting group was detached by hydrogenolytic means in ethanol over palladium/activated carbon. 4.6 mg (0.005 mmol) of the deprotected crude prod-

uct were then, without further purification, taken up in 1.8 ml of dichloromethane and admixed with 3.8 mg (0.012 mmol) of 1,1'-[(1,5-dioxopentane-1,5-diyl)bis(oxy)]dipyrrolidine-2,5-dione, 0.8 μ l (0.005 mmol) of N,N-diisopropylethylamine and 0.6 mg (5 μ mol) of 4-dimethylaminopyridine. The mixture was stirred at RT overnight and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. Thus, 0.96 mg (16% of theory) of the title compound were obtained as a colourless foam.


[3713] HPLC (Method 5): R_t =1.8 min:

[3714] LC-MS (Method 1): R_t =0.94 min; MS (ESIpos): m/z =1193 (M+H)⁺.

Intermediate 132

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazinyl}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl)-N-methyl-L-valinamide

[3715]



[3716] 15 mg (16.7 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 2500 μ l of DMF and then admixed with 9.6 mg (50 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 7.6 mg (50 μ mol) of 1-hydroxy-1H-benzotriazole hydrate, 5.8 μ l of N,N-diisopropylethylamine and with 17.4 mg (67 μ mol) of commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide. The mixture was stirred at RT overnight and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. Thus, 11.2 mg (52% of theory) of the title compound were obtained as a colourless foam.


[3717] HPLC (Method 5): R_f =1.7 min;

[3718] LC-MS (Method 2): R_t =1.09 min; MS (ESIpos): m/z =1106 (M+H)⁺.

Intermediate 133

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazinyl}-4-oxobutyl)-N-methyl-L-valyl-N-{(3R,4S,5S)-1-{(2S)-2-[1-(2R,3S)-1-(benzyloxy)-1-oxo-3-phenylbutan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl}pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl)-N-methyl-L-valinamide

[3719]



[3720] 5.8 mg (6.3 μmol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S,3S)-1-(benzyloxy)-1-oxo-3-phenylbutan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 943 μl of DMF and then admixed with 3.6 mg (19 μmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 2.9 mg (19 μmol) of 1-hydroxy-1H-benzotriazole hydrate, 2.2 μl of N,N-diisopropylethylamine and with 6.6 mg (25 μmol) of commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide. The mixture was stirred at RT overnight and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. Thus, 4.5 mg (64% of theory) of the title compound were obtained as a colourless foam.


[3721] HPLC (Method 5): R_t =2.0 min;

[3722] LC-MS (Method 1): R_t =1.03 min; MS (ESIpos): m/z =1129 (M+H)⁺.

Intermediate 134

N-[3-([2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethyl]carbonyl)amino]propyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3723]



[3724] First, 4-nitrophenyl 2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethyl carbamate was prepared under standard conditions, proceeding from commercially available 1-(2-aminoethyl)-1H-pyrrole-2,5-dione trifluoroacetate and 4-nitrophenyl chlorocarbonate.

[3725] 5 mg (6 μ mol) of N-(3-aminopropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 1000 μ l of DMF and then admixed with 2 μ l of N,N-diisopropylethylamine and with 2.2 mg (9 μ mol) of 4-nitrophenyl 2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethyl carbamate. The mixture was stirred at RT for 1 h and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. Thus, 1.6 mg (23% of theory) of the title compound were obtained as a colourless foam.


[3726] HPLC (Method 5): R_f =1.7 min;

[3727] LC-MS (Method 2): R_f =1.09 min; MS (ESIpos): m/z =1036 (M+H)⁺.

Intermediate 135

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3728]



[3729] 10 mg (11 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 4000 μ l of DMF and then admixed with 6.3 mg (33 μ mol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 4.5 mg (33 μ mol) of 1-hydroxy-1H-benzotriazole hydrate, 5.7 μ l of N,N-diisopropylethylamine and with 11.5 mg (44 μ mol) of commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide. The mixture was stirred at RT overnight and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. Thus, 2.6 mg (14% of theory) of the title compound were obtained as a colourless foam.


[3730] HPLC (Method 6): R_f =2.1 min;

[3731] LC-MS (Method 1): R_f =1.01 min; MS (ESIpos): m/z =1115 (M+H)⁺.

Intermediate 136

N-(4-{4-[4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)butanoyl]piperazin-1-yl}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3732]



[3733] First, 1-[4-oxo-4-(piperazin-1-yl)butyl]-1H-pyrrole-2,5-dione trifluoroacetate was prepared under standard conditions, proceeding from tert-butyl piperazine-1-carboxylate and 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)butanoic acid over 2 stages.

[3734] 5 mg (5.6 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 1000 μ l of DMF and then admixed with 2.1 mg (11 μ mol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 1.7 mg (11 μ mol) of 1-hydroxy-1H-benzotriazole hydrate, 2 μ l of N,N-diisopropylethylamine and with 3.5 mg (5.6 μ mol) of 1-[4-oxo-4-(piperazin-1-yl)butyl]-1H-pyrrole-2,5-dione trifluoroacetate. The mixture was stirred at RT overnight. Then 2.1 mg (5.6 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate were added and the reaction mixture was stirred at RT for a further 3 h. Subsequently, the solvent

was removed under reduced pressure and the remaining residue was purified by means of preparative HPLC. The corresponding fractions were concentrated and, by lyophilization from water, 0.6 mg (10% of theory) of the title compound was obtained as a colourless foam.


[3735] HPLC (Method 6): R_f =1.9 min;

[3736] LC-MS (Method 1): R_f =0.9 min; MS (ESIpos): m/z =1132 (M+H)⁺.

Intermediate 137

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]-1-methylhydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3737]



[3738] First, 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-N-methylhexanehydrazide trifluoroacetate was prepared under standard conditions, proceeding from commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoic acid and tert-butyl 1-methylhydrazinecarboxylate over 2 stages.

[3739] 6.9 mg (8 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 2540 μ l of DMF and then admixed with 3.6 mg (9 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, 3 μ l of N,N-diisopropylethylamine and with 4.1 mg (12 μ mol) of 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-N-methylhexanehydrazide trifluoroacetate. The mixture was stirred at RT overnight. Subsequently, the solvent was removed under reduced pressure and the remaining residue was purified by means of

preparative HPLC. Thus, 3.9 mg (45% of theory) of the title compound were obtained as a colourless foam.


[3740] HPLC (Method 5): R_f =1.8 min;

[3741] LC-MS (Method 1): R_f =0.93 min; MS (ESIpos): m/z =1108 (M+H)⁺.

Intermediate 138

N-{4-[(2-{[4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)butanoyl](methylamino)ethyl}(methylamino)-4-oxobutyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3742]



[3743] Proceeding from tert-butylmethyl 2-(methylamino)ethyl carbamate and 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)butanoic acid, over 2 stages, 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-N-methyl-N-[2-(methylamino)ethyl]butanamide trifluoroacetate was first prepared by.

[3744] 6.6 mg (7.3 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 2000 μ l of DMF and then admixed with 5.6 mg (14.7 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, 2.6 μ l of N,N-diisopropylethylamine and with 4.1 mg (9 μ mol) of 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-N-methyl-N-[2-(methylamino)ethyl]butanamide trifluoroacetate. After stirring at RT for 3 h, the same amounts of HATU and N,N-diisopropylethylamine were added once again, and the reaction mixture was then stirred at RT overnight. Subsequently, the solvent was removed under reduced pressure and the remaining residue was purified by means of preparative HPLC. Thus, 4 mg (44% of theory) of the title compound were obtained as a colourless foam.


[3745] HPLC (Method 6): R_f =2.0 min;

[3746] LC-MS (Method 1): R_f =0.91 min; MS (ESIpos): m/z =1134 (M+H)⁺.

Intermediate 139

(2R,3S)-3-amino-4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutan-2-yl [(3R,4S,7S,10S)-4-[(2S)-butan-2-yl]-7,10-diisopropyl-3-(2-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl}-2-oxoethyl)-5,11-dimethyl-6,9-dioxo-2-oxa-5,8,11-triazapentadecan-15-oate

[3747]



[3748] 13 mg (14.7 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 10 ml of dichloromethane and then admixed with 8.4 mg (44 μ mol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 5.4 mg (44 μ mol) of 4-dimethylaminopyridine and with 9 mg (29.3 μ mol) of commercially available benzyl N-(tert-butoxycarbonyl)-L-threoninate. The mixture was stirred at RT for 5 h. Subsequently, the reaction mixture was twice extracted by shaking with water and the organic phase was dried over sodium sulphate and concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization from dioxane/water, 14 mg (81% of theory) of the protected intermediate were obtained as a colourless foam.

[3749] HPLC (Method 12): R_f =2.3 min;

[3750] LC-MS (Method 1): R_f =1.13 min; MS (ESIpos): m/z =1178 (M+H)⁺.

[3751] Subsequently, the Z protecting group was detached by hydrogenolytic means in methanol over 10% palladium/activated carbon. 9.5 mg (0.0087 μ mol) of the deprotected crude product were then, without further purification, taken up in 5 ml of DMF, and admixed 5 mg (26.2 μ mol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 4 mg (26.2 μ mol) of 1-hydroxy-1H-benzotriazole hydrate, 54.6 μ l of N,N-diisopropylethylamine and with 9.1 mg (34.9

μ mol) of commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide. The mixture was stirred at RT for 1 h and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. After lyophilization from dioxane, 9.5 mg (84% of theory) of the Boc-protected intermediate were obtained.

[3752] HPLC (Method 12): R_f =2.1 min;

[3753] LC-MS (Method 1): R_f =0.97 min; MS (ESIpos): m/z =1295 (M+H)⁺.

[3754] Subsequently, 9.5 mg (7.3 μ mol) were deprotected with 0.5 ml of trifluoroacetic acid in 2 ml of dichloromethane of the Boc-protected intermediate and, after lyophilization from dioxane, 9 mg (82% of theory) of the title compound were obtained as a colourless foam.


[3755] HPLC (Method 12): R_f =2.1 min;

[3756] LC-MS (Method 1): R_f =0.84 min; MS (ESIpos): m/z =1195 (M+H)⁺.

Intermediate 140

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]-1-methylhydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3757]



[3758] 4.1 mg (12 μ mol) of 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-N-methylhexanehydrazide trifluoroacetate (Intermediate 22) were dissolved together with 6.9 mg (8 μ mol) of the compound from Intermediate 61 in 2.5 ml of DMF and then admixed with 3.5 mg (9 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and 3 μ l of N,N-diisopropylethylamine. The mixture was stirred at RT overnight and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. After lyophilization from dioxane, 2.6 mg (30% of theory) of the title compound were obtained.


[3759] HPLC (Method 5): R_f =1.8 min;

[3760] LC-MS (Method 1): R_f =0.90 and 0.91 min; MS (ESIpos): m/z =1120 (M+H)⁺.

Intermediate 141

N-[4-({1-[4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)butanoyl]piperidin-4-yl}oxy)-4-oxobutyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3761]



[3762] 44 mg (49 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 2 ml of dichloromethane and then admixed with 18.8 mg (98 μ mol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 3.8 mg (24 μ mol) of 4-dimethylaminopyridine and with 23 mg (98 μ mol) of commercially available benzyl 4-hydroxypiperidine-1-carboxylate. The mixture was stirred at RT overnight and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. Thus, 22 mg (40% of theory) of the Z-protected intermediate were obtained as a colourless foam.

[3763] HPLC (Method 5): R_f =2.1 min;

[3764] LC-MS (Method 1): R_f =1.04 min; MS (ESIpos): m/z =1116 (M+H)⁺.

[3765] Subsequently, the Z protecting group was detached by hydrogenolytic means in ethanol over palladium/activated carbon.

[3766] 19 mg (19 μ mol) of the deprotected crude product were then, without further purification, taken up in 4 ml of DMF and admixed with 7 mg (39 μ mol) of 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)butanoic acid, 11 mg (29 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and 5 μ l of N,N-diisopropylethylamine. The mixture was stirred at RT for 1 h and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. After lyophilization from dioxane, 7.5 mg (34% of theory) of the title compound were obtained.


[3767] HPLC (Method 5): R_f =1.8 min;

[3768] LC-MS (Method 1): R_f =0.94 min; MS (ESIpos): m/z =1147 (M+H)⁺.

Intermediate 142

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-(benzyloxy)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3769]



[3770] 9 mg (9.5 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-(benzyloxy)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 72) were dissolved in 1000 μ l of DMF and then admixed with 10 mg (38 μ mol) of commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanohydrazide, 7.2 mg (19 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and 8 μ l of N,N-diisopropylethylamine, and the reaction mixture was stirred at RT for 1 h. Subsequently, the solvent was removed under reduced pressure and the remaining residue was purified by means of preparative HPLC. The corresponding fractions were concentrated and, by lyophilization, 6.4 mg (58% of theory) of the title compound were obtained as a colourless foam.


[3771] HPLC (Method 5): R_f =1.9 min;

[3772] LC-MS (Method 1): R_f =0.99 min; MS (ESIpos): m/z =1154 (M+H)⁺.

Intermediate 143

N-(4-{2-[4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2,2-dimethylbutanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3773]



[3774] 6 mg (6.7 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 61) were reacted with 3 mg (8.7 μ mol) of 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2,2-dimethylbutanehydrazide trifluoroacetate in analogy to Intermediate 142 to give 2 mg (27% of theory) of the title compound.


[3775] HPLC (Method 12): R_f =2.1 min;

[3776] LC-MS (Method 3): R_f =1.92 min; MS (ESIpos): m/z =1106 (M+H)⁺.

Intermediate 144

N-(4-{2-[4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2,2-dimethylbutanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3777]



[3778] To a solution of 5 mg (5.6 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide in 1 ml of DMF were added 7.65 mg (22.5 μ mol) of 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2,2-dimethylbutanehydrazide trifluoroacetate, 3.2 mg (16.9 μ mol) of EDC, 1.96 μ l (11.3 μ mol) of diisopropylethylamine and 2.6 mg (16.9 μ mol) of HOBt. The reaction mixture was stirred at RT for 3 h. Subsequently, a further 0.95 mg (2.8 μ mol) of 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2,2-dimethylbutanehydrazide trifluoroacetate was added. After stirring overnight, the reaction mixture was concentrated and


purified by preparative HPLC. 3.5 mg (85% purity, 48% of theory) of the title compound were obtained.

[3779] LC-MS (Method 3): R_f =1.86 min; m/z =1094 (M+H)⁺.

Intermediate 145

N-[3-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)propyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3780]



[3781] 12 mg (14 μ mol) of N-(3-aminopropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarboxyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 66) were taken up in 750 μ l of dioxane and admixed with 1.5 ml of saturated sodium hydrogencarbonate solution and then with 3.2 mg (21 μ mol) of methyl 2,5-dioxo-2,5-dihydro-1H-pyrrole-1-carboxylate. The reaction mixture was stirred at RT for 1 h and then concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization, 4.2 mg (32% of theory) of the title compound were obtained.


[3782] HPLC (Method 5): R_f =1.7 min;

[3783] LC-MS (Method 1): R_f =0.94 min; MS (ESIpos): m/z =950 (M+H)⁺.

Intermediate 146

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-[benzyl(methyl)amino]-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3784]



[3785] 9 mg (9.8 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-[benzyl(methyl)amino]-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 73) were reacted in analogy to Intermediate 133 with 10 mg (39 μ mol) of 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide to give 1.8 mg (15% of theory) of the title compound.


[3786] HPLC (Method 12): R_f =2.2 min;

[3787] LC-MS (Method 9): R_f =5.11 min; MS (ESIpos): m/z =1128 (M+H)⁺.

Intermediate 147

N-{4-[(2,5-dioxopyrrolidin-1-yl)oxy]-4-oxobutyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S,3S)-1-(benzyloxy)-1-oxo-3-phenylbutan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3788]




[3789] 16 mg (17 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S,3S)-1-(benzyloxy)-1-oxo-3-phenylbutan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 70) were dissolved in 2 ml of dichloromethane and admixed with 2.6 mg (23 mmol) of 1-hydroxypyrrolidine-2,5-dione and then with 4 mg (21 μ mol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride. After stirring at RT for 2 h, the same amounts of 1-hydroxypyrrolidine-2,5-dione and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride were added once again. Then stirring at RT overnight, the reaction mixture was concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization, 10 mg (56% of theory) of the title compound were obtained.

[3790] HPLC (Method 5): R_t =2.0 min;

Intermediate 148

N-{4-[(2-{[4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)butanoyl](methyl)amino}ethyl)amino]-4-oxobutyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3791]



[3792] 6 mg (7 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 61) were combined with 2.8 mg (8 μ mol) of N-(2-aminoethyl)-4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-N-methylbutanamide trifluoroacetate, 10.1 mg (27 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and 5 μ l of N,N-diisopropylethylamine in 2 ml of DMF and stirred at RT overnight. Then

another 5 mg (23.5 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and 3 μ l of N,N-diisopropylethylamine were added. After stirring at RT for a further 5 h, the solvent was removed under reduced pressure and the remaining residue was purified by means of preparative HPLC. The corresponding fractions were concentrated and, by lyophilization from dioxane, 1.3 mg (15% of theory) of the title compound were obtained.


[3793] HPLC (Method 12): R_t =2.1 min;

[3794] LC-MS (Method 2): R_t =1.21 min; MS (ESIpos): m/z =1120 (M+H)⁺.

Intermediate 149

N-{4-[(2-{[4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)butanoyl]amino}ethyl)(methyl)amino]-4-oxobutyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3795]




[3796] 6 mg (7 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 61) were combined with 3.1 mg (9 μ mol) of 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-N-[2-(methylamino)ethyl]butanamide trifluoroacetate, 10.1 mg (27 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and 5 μ l of N,N-diisopropylethylamine in 2 ml of DMF, and the mixture was stirred at RT for 4 h. Then the solvent was removed under reduced pressure and the remaining residue was purified by means of preparative HPLC. The corresponding fractions were concentrated and, by lyophilization from dioxane, 1 mg (13.4% of theory) of the title compound were obtained.

[3797] HPLC (Method 12): R_t =2.1 min;[3798] LC-MS (Method 1): R_t =0.89 min; MS (ESIpos): m/z =1121 (M+H)⁺.

Intermediate 150

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-{[(1S,2R)-2-phenyl-1-(propylcarbonyl)cyclopropyl]amino}propyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3799]



[3800] 7.9 mg (9 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[[[1S,2R)-2-phenyl-1-(propylcarbamoyl)cyclopropyl]amino]propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 3 ml of DMF and then admixed with 10.4 mg (54 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 8.3 mg (54 μ mol) of 1-hydroxy-1H-benzotriazole hydrate, 9 μ l of N,N-diisopropylethylamine and with 9.5 mg (36 μ mol) of commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide. The mixture was stirred at RT overnight and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. Thus, 4.3 mg (22% of theory) of the title compound were obtained as a colourless foam.


[3801] HPLC (Method 6): R_f =1.9 min;

[3802] LC-MS (Method 9): R_f =4.93 min; MS (ESIpos): m/z =1078 (M+H)⁺.

Intermediate 151

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[1S,2R)-1-carbamoyl-2-phenylcyclopropyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3803]



[3804] The compound was prepared analogously to Intermediate 150, proceeding from the compound in Intermediate 81.


[3805] HPLC (Method 5): R_f =1.7 min;

[3806] LC-MS (Method 1): R_f =0.87 min; MS (ESIpos): m/z =1036 (M+H)⁺.

Intermediate 152

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[1S,2R)-1-(ethoxycarbonyl)-2-phenylcyclopropyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3807]



[3808] 10 mg (12 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S,2R)-1-(ethoxycarbonyl)-2-phenylcyclopropyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 3 ml of DMF and then admixed with 8.9 mg (23 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, 10 μ l of N,N-diisopropylethylamine and with 12 mg (47 μ mol) of commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide. The mixture was stirred at RT for 1 h. This was followed by concentration under high vacuum and purification of the remaining residue by means of preparative HPLC. Thus, 5.8 mg (37% of theory) of the title compound were obtained as a colourless foam.


[3809] HPLC (Method 6): R_t =2.0 min;

[3810] LC-MS (Method 9): R_t =4.99 min; MS (ESIpos): m/z =1066 (M+H)⁺.

Intermediate 153

N-[1-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-12,15-dioxo-3,6,9-trioxa-13,14-diazaoctadecan-18-yl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3811]



[3812] To a solution of 5 mg (5.6 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide in 1 ml of DMF were added 9.7 mg (22.5 μ mol) of 3-(2-{2-[2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethoxy]ethoxy}ethoxy)propanehydrazide trifluoroacetate, 3.2 mg (16.9 μ mol) of EDC, 1.96 μ l (11.3 μ mol) of N,N-diisopropylethylamine and 2.6 mg (16.9 μ mol) of

HOBt. The reaction mixture was stirred at RT for 3 h. Subsequently, a further 1.2 mg (2.8 μ mol) of 3-(2-{2-[2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethoxy]ethoxy}ethoxy)propanehydrazide trifluoroacetate were added. The reaction mixture was stirred at RT overnight and then purified by preparative HPLC.


[3813] 3.6 mg (51% of theory) of the title compound were obtained.

[3814] LC-MS (Method 1): R_t =0.90 min; m/z =1185 (M+H)⁺.

Intermediate 154

(2R,3S)-3-amino-4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutan-2-yl (3R,4S,7S,10S)-4-[(2S)-butan-2-yl]-7,10-diisopropyl-3-(2-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-2-oxoethyl)-5,11-dimethyl-6,9-dioxo-2-oxa-5,8,11-triazapentadecan-15-oate

[3815]



[3816] 15 mg (17 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-yl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 10 ml of dichloromethane and then admixed with 12.8 mg (67 μ mol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 10 mg (83 μ mol) of 4-dimethylaminopyridine and with 10.3 mg (33 μ mol) of commercially available benzyl N-(tert-butoxycarbonyl)-L-threoninate. The mixture was heated to reflux for 4 h. Then the same amounts of coupling reagent and 4-dimethylaminopyridine were added again and the reaction mixture was heated under reflux overnight. Subsequently, the reaction mixture was diluted with dichloromethane and extracted by shaking once with water, and the organic phase was removed and concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. Thus, 7.7 mg (37% of theory) of the protected intermediate were obtained as a colourless foam.

[3817] HPLC (Method 12): R_f =2.5 min;

[3818] LC-MS (Method 1): R_f =1.13 min; MS (ESIpos): m/z =1190 (M+H)⁺.

[3819] Subsequently, the benzyl ester protecting group was removed by hydrogenation under standard hydrogen pressure in methanol over 10% palladium/activated carbon, and the acid thus obtained, as described in Intermediate 151, was joined to 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanohydrazide. In a last step, the Boc protecting group was detached with trifluoroacetic acid. The remaining residue was purified by means of preparative HPLC. Thus, 0.22 mg (2.5% of theory over 3 stages) of the title compound was obtained as a colourless foam.


[3820] HPLC (Method 12): R_f =2.0 min;

[3821] LC-MS (Method 1): R_f =0.81 min; MS (ESIpos): m/z =1207 (M+H)⁺.

Intermediate 155

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3822]



[3823] This compound was prepared in analogy to the synthesis described in Intermediate 152, from N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S)-1-amino-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide and commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide.


[3824] HPLC (Method 5): R_f =1.6 min;

[3825] LC-MS (Method 1): R_f =0.82 min; MS (ESIpos): m/z =1024 (M+H)⁺.

Intermediate 156

N-(3-[[[1-[(2,5-dioxopyrrolidin-1-yl)oxy]carbonyl]cyclopropyl]carbonyl]amino]propyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3826]



[3827] This compound was prepared in analogy to the synthesis described in the last stage of Intermediate 131, from N-(3-aminopropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide and 1,1'-[cyclopropane-1,1-diylbis(carbonyloxy)]dipyrrolidine-2,5-dione, which had been obtained from the corresponding dicarboxylic acid beforehand.


[3828] HPLC (Method 12): R_f =2.0 min;

[3829] LC-MS (Method 1): R_f =0.92 min; MS (ESIpos): m/z =1080 (M+H)⁺.

Intermediate 157

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3830]



[3831] 15 mg (18 μ mol) of (N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 3.8 ml of DMF and then admixed with 27 mg (70 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, 12 μ l of N,N-diisopropylethylamine and with 14 mg (53 μ mol) of commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide. The reaction mixture was stirred at RT for 1 h. This was followed by concentration under high vacuum and purification of the remaining residue by means of preparative HPLC. Thus, 6.2 mg (33% of theory) of the title compound were obtained as a colourless foam.

[3832] HPLC (Method 5): R_f =1.6 min;


[3833] LC-MS (Method 1): R_f =0.85 min; MS (ESIpos): m/z =1063 (M+H)⁺.

[3834] ¹H-NMR (500 MHz, DMSO-d₆, characteristic signals): δ =10.8 (d, 1H), 9.8-9.7 (m, 2H), 9.6 and 9.4 (2m, 1H), 8.9, 8.88, 8.78 and 8.75 (4d, 1H), 8.08 and 7.85 (2d, 1H), 7.6-6.9 (m, 9H), 4.7-4.4 (m, 3H), 3.4 (t, 2H), 3.23, 3.2, 3.18, 3.0, and 2.99 (5s, 9H), 2.8 (m, 3H), 2.1 (t, 2H), 1.06 and 1.01 (2d, 3H), 0.95-0.8 (m, 15H), 0.8-0.75 (dd, 3H).

Intermediate 158

N-[4-[(2R)-1-[(2,5-dioxopyrrolidin-1-yl)oxy]-4-methyl-1-oxopentan-2-yl]amino]-4-oxobutyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzylamino)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3835]



[3836] 13 mg (14.7 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzylamino)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 4 ml of dimethylformamide and then admixed with 9.4 mg (25 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, 6 μ l of N,N-diisopropylethylamine and with 7 mg (31 μ mol) of commercially available tert-butyl D-leucinate hydrochloride. The mixture was stirred at RT for 5 h and then concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization from dioxane/water, 6.5 mg (49% of theory) of the protected intermediate were obtained as a colourless foam.

[3837] HPLC (Method 5): R_f =2.2 min;


[3838] LC-MS (Method 1): R_f =1.21 min; MS (ESIpos): m/z =1076 (M+H)⁺.

[3839] Trifluoroacetic acid in dichloromethane was first used to detach the Boc protecting group from this protected intermediate, giving 6.2 mg (99% of theory) of the deprotected compound. 5.2 mg (5 μ mol) of this intermediate were taken up in 1.5 ml of dichloromethane and reacted with 0.8 mg (7 μ mol) of N-hydroxysuccinimide in the presence of 1.2 mg (6 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 0.16 mg (1 mmol) of 4-dimethylaminopyridine. After stirring at RT for 2 h, the reaction mixture was concentrated and purified by means of preparative HPLC. 1.3 mg of the title compound were obtained, some of which was hydrolysed to the reactant.

Intermediate 159

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-(benzylamino)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3840]



[3841] This compound was prepared in analogy to the synthesis described in Intermediate 157, from N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-(benzylamino)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide and commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide.

[3842] Yield: 6 mg (53% of theory)


[3843] HPLC (Method 5): R_f =1.9 min;

[3844] LC-MS (Method 1): R_f =0.94 min; MS (ESIpos): m/z =1114 (M+H)⁺.

Intermediate 160

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-(benzylamino)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3845]



[3846] This compound was prepared in analogy to the synthesis described in Intermediate 157, from 20 mg (21 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-(benzylamino)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide and commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide.

[3847] Yield: 13 mg (52% of theory)


[3848] HPLC (Method 5): R_f =1.9 min;

[3849] LC-MS (Method 1): R_f =0.92 min; MS (ESIpos): m/z =1153 (M+H)⁺.

Intermediate 161

N-(6-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-6-oxohexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3850]



[3851] This compound was prepared in analogy to the synthesis described in Intermediate 157, from N-(5-carboxypentyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide and commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide.

[3852] Yield: 0.8 mg (16% of theory)


[3853] HPLC (Method 5): R_f =1.6 min;

[3854] LC-MS (Method 1): R_f =0.78 min; MS (ESIpos): m/z =1092 (M+H)⁺.

Intermediate 162

N-{6-[(2,5-dioxopyrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3855]



[3856] 18 mg (20 μ mol) of N-(5-carboxypentyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[[[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 64) were dissolved in 3.2 ml of dichloromethane and admixed with 22 mg (190 mmol) of 1-hydroxypyrrolidine-2,5-dione and then with 11 mg (60 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 0.24 mg (0.17 mmol) of DMAP. After stirring at RT for 2 h, another 22 mg (190 mmol) of 1-hydroxypyrrolidine-2,5-dione, 11 mg (60 μ mol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 0.24 mg (0.17 μ mol) of DMAP were added and the reaction mixture was stirred at RT for a further hour. This was followed by concentration under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization, 8.2 mg (41% of theory) of the title compound were obtained.


[3857] HPLC (Method 5): R_f =2.0 min;

[3858] LC-MS (Method 11): R_f =0.9 min; MS (ESIpos): m/z =1024 (M+H)⁺.

Intermediate 163

[(1S,2R)-1-amino-2-phenylcyclopropyl](1,4-dihydro-3H-2,3-benzoxazin-3-yl)methanone trifluoroacetate

[3859]



[3860] First, proceeding from 265 mg (0.82 mmol) of tert-butyl (1S,2R)-1-(hydroxycarbonyl)-2-phenylcyclopropyl carbamate (Starting Compound 7), by reaction with 1,2-bis(bromomethyl)benzene, analogously to a literature method (see H. King, *J. Chem. Soc.* 1942, 432), the Boc-protected tert-butyl (1S,2R)-1-(1,4-dihydro-3H-2,3-benzoxazin-3-ylcarbonyl)-2-phenylcyclopropyl carbamate intermediate was prepared.

[3861] Yield: 108 mg (34% of theory)

[3862] LC-MS (Method 2): R_f =1.3 min; MS (ESIpos): m/z =395 (M+H)⁺.

[3863] 108 mg (0.27 mmol) of this intermediate were taken up in 3.7 ml of dichloromethane, 1.8 ml of trifluoroacetic acid were added, and the mixture was stirred at RT for 15 min. This was followed by concentration under reduced pressure and


lyophilization of the remaining residue from dioxane. 112 mg of the title compound were obtained in quantitative yield as a colourless foam.

[3864] LC-MS (Method 1): R_f =0.7 min; MS (ESIpos): m/z =295 (M+H)⁺.

Intermediate 164

N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(1S,2R)-1-(1,4-dihydro-3H-2,3-benzoxazin-3-ylcarbonyl)-2-phenylcyclopropyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[3865]



[3866] 166 mg (0.196 mmol) of N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 10) were taken up in 40 ml of DMF and admixed successively with 80 mg (0.196 mmol) of [(1S,2R)-1-amino-2-phenylcyclopropyl](1,4-dihydro-3H-2,3-benzoxazin-3-yl)methanone trifluoroacetate (Intermediate 163), 112 mg (0.294 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) and 682 μ l (3.9 mmol) of N,N-diisopropylethylamine. The mixture was subsequently stirred at RT overnight. The reaction mixture was then concentrated under reduced pressure, the residue was taken up in ethyl acetate and the solution was washed with saturated aqueous sodium chloride solution. The organic phase was dried over magnesium sulphate, filtered and concentrated. The residue was finally purified by preparative HPLC. In this way, 19 mg (9% of theory) of the Fmoc-protected intermediate N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(1S,2R)-1-(1,4-dihydro-3H-2,3-benzoxazin-3-ylcarbonyl)-2-phenylcyclopropyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were obtained.

[3867] HPLC (Method 5): R_f =1.68 min;

[3868] LC-MS (Method 1): R_f =1.51 min; MS (ESIpos): m/z =1083 (M+H)⁺.

[3869] 19 mg (0.015 mmol) of this intermediate were dissolved in 4 ml of DMF. After 817 μ l of piperidine had been added, the reaction mixture was stirred at RT for 5 min. This was followed by concentration under reduced pressure, and the residue was first digested with diethyl ether and then

purified by means of preparative HPLC (eluent: acetonitrile+ 0.1% TFA/0.1% aq. TFA). The corresponding fractions were combined, the solvent was removed under reduced pressure and then the residue was lyophilized from dioxane/water. 12 mg (92% of theory) of the title compound were obtained as a colourless foam.


[3870] HPLC (Method 6): R_f =2.0 min;

[3871] LC-MS (Method 1): R_f =0.94 min; MS (ESIpos): m/z =861 (M+H)⁺.

Intermediate 165

N-(6-aminohexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(1S,2R)-1-(1,4-dihydro-3H-2,3-benzoxazin-3-ylcarbonyl)-2-phenylcyclopropyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3872]



[3873] 20 mg (0.021 mmol) of Intermediate 164 were used, in analogy to the preparation of Intermediate 97, with benzyl 6-oxohexyl carbamate in the presence of sodium cyanoborohydride and subsequent hydrogenolytic detachment of the Z protecting group (with 5% palladium on charcoal as a catalyst, in methanol as a solvent), to prepare the title compound.

[3874] Yield: 4.5 mg (23% of theory over 2 stages)


[3875] HPLC (Method 12): R_f =1.9 min;

[3876] LC-MS (Method 1): R_f =0.9 min; MS (ESIpos): m/z =960 (M+H)⁺.

Intermediate 166

N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(1S,2R)-1-(1,4-dihydro-3H-2,3-benzoxazin-3-ylcarbonyl)-2-phenylcyclopropyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3877]



[3878] 4.4 mg (4.5 μ mol) of Intermediate 165 were taken up in 1 ml of 1:1 dioxane/water and then admixed with 1 mg (6.8 μ mol) of methyl 2,5-dioxo-2,5-dihydro-1H-pyrrole-1-carboxylate and with 50 μ l of saturated aqueous sodium hydrogencarbonate solution. The reaction mixture was stirred at RT for 30 min. Then another 50 μ l of the saturated aqueous sodium hydrogencarbonate solution were added and the reaction mixture was stirred at RT for a further 15 min and then concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization, 1 mg (21% of theory) of the title compound were obtained as a colourless foam.


[3879] HPLC (Method 12): R_f =2.1 min;

[3880] LC-MS (Method 1): R_f =1.08 min; MS (ESIpos): m/z =1040 (M+H)⁺.

Intermediate 167

benzyl 3-{2-[2-(2-oxoethoxy)ethoxy]
ethoxy}propanoate

[3881]



[3882] The title compound was prepared from 6 g (21.55 mmol) of commercially available 3-{2-[2-(2-hydroxyethoxy)ethoxy]ethoxy}propanoic acid under standard conditions, first by esterification with benzyl chloride and caesium carbonate and subsequent oxidation with sulphur trioxide-pyridine complex.

[3883] Yield: 611 mg (10% of theory over 2 stages)

[3884] LC-MS (Method 2): R_f =1.69 min; MS (ESIpos): m/z =311 (M+H)⁺.

Intermediate 168

N-(2-{2-[2-(2-carboxyethoxy)ethoxy]ethoxy}ethyl)-
N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,
2R)-3-[[[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3885]

[3886] First, in analogy to the synthesis described in Intermediate 69, by coupling of N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 4) and N^α-{(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoyl}-L-tryptophanamide trifluoroacetate (Intermediate 49) in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Fmoc protecting group by means of piperidine, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate.

[3887] 25 mg (0.028 mmol) of this compound and 17.5 mg (0.06 mmol) of Intermediate 167 were combined in 2 ml of methanol and admixed with 12.6 mg (0.14 mmol) of borane-pyridine complex and 2.5 ml of acetic acid. The reaction mixture was stirred at RT overnight. Then the same amounts of borane-pyridine complex and acetic acid were added once again and the reaction mixture was stirred at RT for a further 24 h. This was followed by concentration under reduced pressure, and the residue was purified by means of preparative HPLC. After concentration of the corresponding fractions and lyophilization from 1:1 dioxane/water, 26.5 mg (88% of theory) of the Z-protected title compound were obtained.


[3888] HPLC (Method 12): R_f =2.04 min;

[3889] LC-MS (Method 1): R_f =0.97 min; MS (ESIpos): m/z =1064 (M+H)⁺.

[3890] 25 mg (0.024 mmol) of this intermediate were taken up in 10 ml of methanol and hydrogenated over 10% palladium on activated carbon under standard hydrogen pressure at RT for 45 min. The catalyst was then filtered off and the solvent was removed under reduced pressure. After lyophilization from dioxane, 19.7 mg (85% of theory) of the title compound were obtained.

[3891] HPLC (Method 12): R_f =1.8 min;


[3892] LC-MS (Method 1): R_f =0.83 min; MS (ESIpos): m/z =974 (M+H)⁺.



Intermediate 169

N-{2-[2-(2-{3-[(2,5-dioxopyrrolidin-1-yl)oxy]-3-oxopropoxy}ethoxy)ethoxy]ethyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3893]



[3894] 10 mg (10 μ mol) of Intermediate 168 were dissolved in 3 ml of DMF and admixed with 3.5 mg (30 mmol) of 1-hydroxypyrrolidine-2,5-dione and then with 2.4 mg (10 μ mol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 5 μ l of N,N-diisopropylethylamine. After stirring at RT for 20 h, 8 mg (0.02 mmol) of HATU were added and the reaction mixture was stirred once again at RT overnight and then concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization from dioxane, 8.6 mg (64% of theory) of the title compound were obtained.


[3895] HPLC (Method 12): R_f =1.9 min;

[3896] LC-MS (Method 11): R_f =0.81 min; MS (ESIpos): m/z =1071 (M+H)⁺.

Intermediate 170

N-(6-aminohexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(2S,3S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylbutan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3897]



[3898] This compound was prepared in analogy to Intermediate 101 over 2 stages, proceeding from 26 mg (0.028 mmol) of Intermediate 15.

[3899] Yield: 16.7 mg (63% of theory over 2 stages)


[3900] HPLC (Method 12): R_f =1.9 min;

[3901] LC-MS (Method 1): R_f =0.81 min; MS (ESIpos): m/z =914 (M+H)⁺.

Intermediate 171

N-(6-{[4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)butanoyl]amino}hexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(2S,3S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylbutan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3902]



[3903] 6.7 mg (7.3 μ mol) of the compound formed from Intermediate 170 and 3 mg (14.7 μ mol) of commercially available 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)butanoic acid were taken up in 2 ml of DMF and admixed with 5.6 mg (14.7 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) and 2 μ l of N,N-diisopropylethylamine. The mixture was stirred at RT for 30 min. The reaction mixture was concentrated and the residue was purified by means of preparative HPLC. The corresponding fractions were combined, the solvent was removed under reduced pressure and then the residue was lyophilized from dioxane. Thus, 4.5 mg (56% of theory) of the title compound were obtained.


[3904] HPLC (Method 12): R_f =2.0 min;

[3905] LC-MS (Method 1): R_f =1.12 min; MS (ESIpos): m/z =1079 (M+H)⁺.

Intermediate 172

benzyl 2-{2-[2-(2-oxoethoxy)ethoxy]ethoxy}ethyl carbamate

[3906]



[3907] The title compound was prepared from commercially available 2-{2-[2-(2-aminoethoxy)ethoxy]ethoxy}ethanol under standard conditions, by first introducing the Z protecting group and then oxidizing with sulphur trioxide-pyridine complex.


[3908] HPLC (Method 12): R_f =1.4 min;

[3909] LC-MS (Method 11): R_f =0.65 min; MS (ESIpos): m/z =326 (M+H)⁺.

Intermediate 173

benzyl 2-[2-(2-oxoethoxy)ethoxy]ethyl carbamate

[3910]



[3911] The title compound was prepared analogously to Intermediate 172 from commercially available 2-[2-(2-aminoethoxy)ethoxy]ethanol under standard conditions, by first introducing the Z protecting group and then oxidizing with sulphur trioxide-pyridine complex.


[3912] HPLC (Method 12): R_f =1.3 min;

[3913] LC-MS (Method 11): R_f =0.68 min; MS (ESIpos): m/z =282 (M+H)⁺.

Intermediate 174

N-(2-{2-[2-(2-aminoethoxy)ethoxy]ethoxy}ethyl)-
N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{
(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-
1-(1,2-oxazinan-2-ylcarbonyl)-2-phenyl cyclopropyl]
amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-
oxoheptan-4-yl]-N-methyl-L-valinamide

[3914]



[3915] 47 mg (0.05 mmol) of Intermediate 16 were reductively aminated in analogy to the preparation of Intermediate 167 with benzyl 2-{2-[2-(2-oxoethoxy)ethoxy]ethoxy}ethyl carbamate in the presence of borane-pyridine complex. Subsequently, the Z protecting group was removed by hydrogenolytic means with 5% palladium on charcoal as a catalyst and in methanol as a solvent, and 38 mg (66% of theory over 2 stages) of the title compound were prepared.


[3916] HPLC (Method 5): R_f =1.7 min;

[3917] LC-MS (Method 1): R_f =0.8 min; MS (ESIpos): m/z =988 (M+H)⁺.

Intermediate 175

N-[2-(2-{2-[2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethoxy]ethoxy}ethoxy)ethyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3918]



[3919] The preparation was effected in analogy zu Intermediate 166, proceeding from 34 mg (0.03 mmol) of Intermediate 174.

[3920] Yield: 8.3 mg (23% of theory)


[3921] HPLC (Method 5): R_f =1.9 min;

[3922] LC-MS (Method 1): R_f =0.97 min; MS (ESIpos): m/z =1068 (M+H)⁺.

Intermediate 176

N-(2-{2-[2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethoxy]ethoxy}ethyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3923]



[3924] The preparation was effected in analogy to Intermediates 174 and 175, commencing with the reductive amination of Intermediate 16 with Intermediate 173, subsequent deprotection and formation of the maleimide.


[3925] HPLC (Method 12): R_f =1.8 min;

[3926] LC-MS (Method 11): R_f =0.8 min; MS (ESIpos): m/z =981 (M+H)⁺.

Intermediate 177

N-[2-(2-{2-[2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethoxy]ethoxy}ethoxy)ethyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3927]



[3928] The preparation was effected in analogy to Intermediates 174 and 175, commencing with the reductive amination of Intermediate 16 with Intermediate 172, subsequent deprotection and formation of the maleimide.


[3929] HPLC (Method 12): R_f =1.9 min;

[3930] LC-MS (Method 1): R_f =0.86 min; MS (ESIpos): m/z =1025 (M+H)⁺.

Intermediate 178

N-{4-[(2,5-dioxopyrrolidin-1-yl)oxy]-4-oxobutyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3931]




[3932] The preparation was effected in analogy to Intermediates 162, proceeding from 6 mg of Intermediate 82.

[3933] LC-MS (Method 1): R_f =0.82 min; MS (ESIpos): m/z =953 (M+H)⁺.

Intermediate 179

4-[(1E,3S)-3-amino-4-phenylbut-1-en-1-yl]benzenesulphonic acid trifluoroacetate

[3934]



[3935] A mixture of 13.6 mg (0.06 mmol) of palladium(II) acetate, 469 mg (1.46 mmol) of potassium 4-iodobenzenesulphonate, 300 mg (1.21 mmol) of (S)-tert-butyl 1-phenylbut-3-en-2-yl carbamate, 16.5 mg (0.12 mmol) of phenylurea and 167.6 mg (1.21 mmol) of potassium carbonate in 7.5 ml of DMF was heated to 160°C. in a microwave for 15 min. The crude product was subsequently purified directly by preparative HPLC. This gave 312 mg of a mixture of 31% of the BOC-protected compound and 69% of the free amine.

[3936] This mixture was subsequently taken up in 30 ml of dichloromethane, admixed with 1 ml of trifluoroacetic acid and stirred at RT for 20 h. After concentrating under reduced pressure, the residue was stirred with diethyl ether, and the


precipitate formed was filtered off with suction and washed with diethyl ether. This gave 200 mg (62% of theory) of the title compound.

[3937] LC-MS (Method 11): R_f =0.44 min; MS (ESIpos): m/z =304 (M+H)⁺.

Intermediate 180

4-[(3R)-3-amino-4-phenylbutyl]benzenesulphonic acid

[3938]



[3939] 100 mg (0.25 mmol) of 4-[(1E,3S)-3-amino-4-phenylbut-1-en-1-yl]benzenesulphonic acid trifluoroacetate were suspended in 10 ml of acetic acid and a few drops of DMF and water, admixed with 70 mg (0.07 mmol) of palladium on charcoal (10%) and hydrogenated at hydrogen pressure 2.2 bar for 24 h. The solution was filtered and the filtrate purified by prep. HPLC.


[3940] 29 mg (76% purity, 21% of theory) of product were obtained.

[3941] LC-MS (Method 1): R_f =0.46 min; MS (ESIpos): m/z =306 (M+H)⁺.

Intermediate 181

N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2S,3E)-1-phenyl-4-(4-sulphophenyl)but-3-en-2-yl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3942]




[3943] To a solution of 90 mg (0.13 mmol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide in 4 ml of DMF were added 60 mg (0.16 mmol) of HATU and 69 μ l of (0.39 mmol) Hünig's base. The reaction mixture was stirred at RT for 30 min and then admixed with 60 mg (0.15 mmol) 60.3 mg (0.13 mmol) of 4-[(1E,3S)-3-amino-4-phenylbut-1-en-1-yl]benzenesulphonic acid trifluoroacetate. After stirring overnight, the reaction mixture was purified by prep. HPLC. This gave 127 mg of a 44:56 mixture of the title compound and of the already deprotected amine.

[3944] LC-MS (Method 1): R_f =1.21 min; MS (ESIpos): m/z =971 (M+H)⁺; R_f =0.84 min; MS (ESIpos): m/z =871 (M+H)⁺ for the deprotected compound.

Intermediate 182

N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2S,3E)-1-phenyl-4-(4-sulphophenyl)but-3-en-2-yl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[3945]



[3946] 90 mg of Intermediate 180 were dissolved in 4.6 ml of dichloromethane, and 0.92 ml of trifluoroacetic acid was added. The reaction mixture was stirred at RT for 30 min and then concentrated. The crude product obtained was purified by prep. HPLC.


[3947] 91 mg (98% of theory) of the target compound were obtained.

[3948] LC-MS (Method 1): R_f =0.85 min; MS (ESIpos): m/z =871 (M+H)⁺

Intermediate 183

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2S,3E)-1-phenyl-4-(4-sulphophenyl)but-3-en-2-yl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3949]



[3950] 16.7 μ l (0.03 mmol) of a 15% aqueous succinaldehyde solution were initially charged in 943 μ l of methanol and admixed with 17 mg (0.02 mmol) of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2S,3E)-1-phenyl-4-(4-sulphophenyl)but-3-en-2-yl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate (Intermediate 181) and 1.1 μ l (0.02 mmol) of acetic acid. The reaction mixture was stirred for 5 min at RT and then 2.9 μ l (0.02 mmol) of borane-pyridine complex were added. After 1 h, a further 2 equivalents each of succinaldehyde, acetic acid and borane-pyridine complex were added and the mixture was stirred at RT for 20 h. The reaction mixture was then purified by prep. HPLC.


[3951] This gave 20 mg (83% purity, 80% of theory) of the title compound.

[3952] LC-MS (Method 1): R_f =0.87 min; MS (ESIpos): m/z =957 (M+H)⁺

Intermediate 184

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2S,3E)-1-phenyl-4-(4-sulphophenyl)but-3-en-2-yl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3953]



[3954] 8 mg (7.5 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2S,3E)-1-phenyl-4-(4-sulphophenyl)but-3-en-2-yl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide, 2.8 mg (8.2 μ mol) of 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanhydrazide trifluoroacetate, 3.4 mg (9 μ mol) of HATU and 3.9 μ l of Hünig's base were stirred in 0.77 ml of DMF at RT for 20 h. Subsequently, the reaction mixture was purified by prep. HPLC.


[3955] 3 mg (31% of theory) of the title compound were obtained.

[3956] LC-MS (Method 1): R_t =0.90 min; MS (ESIpos): m/z =1164 (M+H)⁺

Intermediate 185

N-{4-[(2,5-dioxopyrrolidin-1-yl)oxy]-4-oxobutyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2S,3E)-1-phenyl-4-(4-sulphophenyl)but-3-en-2-yl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3957]




[3958] To a solution of 8 mg (7.5 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2S,3E)-1-phenyl-4-(4-sulphophenyl)but-3-en-2-yl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide in 2 ml of DMF were added 8.6 mg (74.8 μ mol) of N-hydroxysuccinimide, 8.5 mg (22.4 μ mol) of EDCI and 0.1 mg (0.75 μ mol) of DMAP. The reaction mixture was stirred at RT for 20 h. Subsequently, 1.3 μ l (7.5 μ mol) of Hünig's base were added and the mixture was stirred for 1 h. The reaction mixture was then purified by prep. HPLC. 2.6 mg (72% purity, 21% of theory) of the title compound were obtained.

[3959] LC-MS (Method 1): R_t =0.89 min; MS (ESIpos): m/z =1054 (M+H)⁺

Intermediate 186

N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2R)-1-phenyl-4-(4-sulphophenyl)butan-2-yl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3960]




[3961] To a solution of 43 mg (0.06 mmol) of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide in 1.9 ml of DMF were added 29 mg (0.07 mmol) of HATU and 33 μ l (0.19 mmol) of Hünig's base. The reaction mixture was stirred at RT for 30 min and then admixed with 29 mg (0.07 mmol) of 4-[(3R)-3-amino-4-phenylbutyl]benzenesulphonic acid trifluoroacetate. After stirring overnight, the reaction mixture was purified by prep. HPLC. This gave 58 mg of a 45:55 mixture of the title compound and of the already deprotected amine.

[3962] LC-MS (Method 1): R_f =1.09 min; MS (ESIpos): m/z =973 (M+H)⁺; R_f =0.87 min; MS (ESIpos): m/z =873 (M+H)⁺ for the deprotected compound.

Intermediate 187

N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2R)-1-phenyl-4-(4-sulphophenyl)butan-2-yl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[3963]



[3964] 58 mg of Intermediate 186 were dissolved in 4.1 ml of dichloromethane, 0.41 ml of trifluoroacetic acid was added and the mixture was stirred at RT for 30 min. After concentration under reduced pressure, the crude product was purified by prep. HPLC.


[3965] 50 mg (90% purity, 85% of theory) of the title compound were obtained.

[3966] LC-MS (Method 1): R_t =0.87 min; MS (ESIpos): m/z =873 (M+H)⁺

Intermediate 188

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[[[(2R)-1-phenyl-4-(4-sulphophenyl)butan-2-yl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3967]



[3968] 171 μ l (0.26 mmol) of a 15% aqueous succinaldehyde solution were initially charged in 2.5 ml of methanol and admixed with 50 mg (0.05 mmol) of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[[[(2R)-1-phenyl-4-(4-sulphophenyl)butan-2-yl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate and 11.6 μ l (0.2 mmol) of acetic acid. The reaction mixture was stirred for 5 min at RT and then 30 μ l (0.24 mmol) of borane-pyridine complex were added. After stirring for 24 hours, a further equivalent of borane-pyridine complex was added and the mixture was stirred for a further 2 h. The reaction mixture was then purified by prep. HPLC.


[3969] 40 mg (90% purity, 66% of theory) of the title compound were obtained.

[3970] LC-MS (Method 1): R_t =0.91 min; MS (ESIpos): m/z =959 (M+H)⁺

Intermediate 189

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2R)-1-phenyl-4-(4-sulphophenyl)butan-2-yl]amino}propyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3971]



[3972] 10 mg (9.3 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2R)-1-phenyl-4-(4-sulphophenyl)butan-2-yl]amino}propyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide, 3.5 mg (10.3 μ mol) of 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanohydrazide trifluoroacetate, 4.3 mg (11.2 μ mol) of HATU and 4.9 μ l (28 μ mol) of Hünig's base were stirred in 1 ml of DMF at RT for 20 h. Subsequently, the reaction mixture was purified by prep. HPLC.


[3973] 4.2 mg (92% purity, 33% of theory) of the title compound were obtained.

[3974] LC-MS (Method 1): R_t =0.91 min; MS (ESIpos): m/z =1166 (M+H)⁺

Intermediate 190

N-{4-[(2,5-dioxopyrrolidin-1-yl)oxy]-4-oxobutyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2R)-1-phenyl-4-(4-sulphophenyl)butan-2-yl]amino}propyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3975]



[3976] To a solution of 10 mg (9.3 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(2R)-1-phenyl-4-(4-sulphophenyl)butan-2-yl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide in 2.5 ml of DMF were added 10.7 mg (93 μ mol) of N-hydroxysuccinimide, 10.6 mg (28 μ mol) of EDCI and 0.12 mg (0.9 μ mol) of DMAP. The reaction mixture was stirred at RT for 20 h and then purified by prep. HPLC.


[3977] 3.8 mg (72% purity, 25% of theory) of the title compound were obtained.

[3983] LC-MS (Method 1): R_f =0.7 min; MS (ESIpos): m/z =443 (M+H)⁺.

Intermediate 192

N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3984]




[3978] LC-MS (Method 1): R_f =0.90 min; MS (ESIpos): m/z =1055 (M+H)⁺

Intermediate 191

(2R,3R)-N-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanamide trifluoroacetate

[3979]



[3980] The title compound was prepared in analogy to the synthesis of Intermediate 7 over two stages from Starting Compound 1 and (2S)-2-amino-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)propan-1-one trifluoroacetate (Intermediate 99).

[3981] Yield over 2 stages: 62 mg (67% of theory)

[3982] HPLC (Method 6): R_f =1.65 min;

[3985] 1015 mg (1.59 mmol) of N-[(9H-fluoren-9-yl-methoxy)carbonyl]-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 4) were taken up in 50 ml of DMF, admixed with 654 mg (2.39 mmol) of 2-bromo-1-ethylpyridinium tetrafluoroborate (BEP) and 2.8 ml of N,N-diisopropylethylamine, and stirred at RT for 10 min. Then 1083 mg (1.75 mmol) of (2R,3R)-N-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanamide trifluoroacetate (Intermediate 191) were added and then the mixture was treated in an ultrasound bath at RT for 30 min. The reaction mixture was then concentrated under reduced pressure and the residue was taken up in 300 ml of ethyl acetate. The organic phase was washed successively with 5% aqueous citric acid solution and 5% aqueous sodium hydrogencarbonate solution, dried over magnesium sulphate, filtered and concentrated. The crude product thus obtained (1684 mg), without further purification, was taken up in 20 ml of acetonitrile, 2 ml of piperidine were added and the reaction mixture was then stirred at RT for 10 min. Then the mixture was concentrated under reduced pressure and the residue was admixed with diethyl ether. The solvent was concentrated by evaporation again and the residue was purified by flash chromatography on silica gel (eluent: 15:1:0.1->15:2:0.2 dichloromethane/methanol/17% aqueous ammonia solution). The corresponding fractions were combined, the solvent was removed under reduced pressure and the residue was lyophilized from acetonitrile/water. Thus, 895 mg (67% over 2 stages) of the title compound were obtained.

[3986] HPLC (Method 12): R_f =1.8 min;


[3987] LC-MS (Method 1): R_f =0.84 min; MS (ESIpos): m/z =840 (M+H)⁺.

[3988] ¹H NMR (500 MHz, DMSO-d₆): δ =10.8 (d, 1H), 8.3 and 8.05 (2d, 1H), 8.0 (d, 1H), 7.5 (m, 1H), 7.3 (m, 1H), 7.15 and 7.08 (2s, 1H), 7.05-6.9 (m, 2H), 5.12 and 4.95 (2m, 1H), 4.65 (m, 1H), 4.55 (m, 1H), 4.1-3.8 (m, 4H), 3.75 (d, 1H), 3.23, 3.18, 3.17, 3.12, 2.95 and 2.88 (6s, 9H), 3.1-3.0 and 2.85 (2m, 2H), 2.65 (d, 1H), 2.4-2.2 (m, 3H), 2.15 (m, 3H), 1.95 (br. m, 2H), 1.85-0.8 (br. m, 11H), 1.08 and 1.04 (2d, 3H), 0.9-0.75 (m, 15H), 0.75-0.65 (dd, 3H) [further signals hidden under H₂O peak].

Intermediate 193

N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3989]



[3990] 50 mg (0.052 mmol) of N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

(Intermediate 192) and 204 μ l of a 15% aqueous solution of 4-oxobutanoic acid were combined in 2 ml of methanol and admixed with 23.4 mg (0.252 mmol) of borane-pyridine complex and 6 μ l of acetic acid. The reaction mixture was stirred at RT overnight. This was followed by concentration under reduced pressure, and the residue was purified by means of preparative HPLC. After concentration of the corresponding fractions, 38 mg (78% of theory) of the title compound were obtained.


[3991] HPLC (Method 5): R_t =1.7 min;

[3992] LC-MS (Method 9): R_t =4.7 min; MS (ESIpos): m/z =926 (M+H)⁺.

Intermediate 194

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3993]



[3994] This compound was prepared in analogy to the synthesis described in Intermediate 157 from 10 mg (11 μ mol) of N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide and commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide.

[3995] Yield: 4.4 mg (35% of theory)


[3996] HPLC (Method 5): R_f =1.8 min;

[3997] LC-MS (Method 1): R_f =0.90 min; MS (ESIpos): m/z =1133 (M+H)⁺.

Intermediate 195

N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S,3S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylbutan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[3998]



[3999] This compound was prepared in analogy to Intermediate 166, proceeding from 9 mg (0.010 mmol) of Intermediate 170.

[4000] Yield: 1.1 mg (10% of theory)


[4001] HPLC (Method 12): R_f =2.0 min;

[4002] LC-MS (Method 1): R_f =0.99 min; MS (ESIpos): m/z =994 (M+H)⁺.

Intermediate 196

(2S)-2-amino-1-(2-oxa-3-azabicyclo[2.2.2]oct-5-en-3-yl)-3-phenylpropan-1-one trifluoroacetate

[4003]



[4004] 41 mg (0.37 mmol) of 2,5-dioxopyrrolidin-1-yl N-(tert-butoxycarbonyl)-L-phenylalaninate were taken up in 10 ml of DMF and admixed with 149 mg (0.41 mmol) of 2-oxa-3-azabicyclo[2.2.2]oct-5-ene (Starting Compound 6) and 72 μ l (0.41 mmol) of N,N-diisopropylethylamine. The mixture was stirred at RT for 1 h. The solvent was removed under reduced pressure, and the residue was taken up in ethyl acetate and extracted by shaking with 5% aqueous citric acid solution and then with 5% aqueous sodium hydrogencarbonate solution. The organic phase was concentrated and the residue was purified by flash chromatography on silica gel with 10:1 toluene/ethanol as the eluent. The corresponding fractions were combined and the solvent was removed under reduced pressure. After the residue had been dried under high vacuum, 69 mg (47% of theory) of the Boc-protected intermediate tert-butyl (2S)-1-(2-oxa-3-azabicyclo[2.2.2]oct-5-en-3-yl)-1-oxo-3-phenylpropan-2-yl carbamate were thus obtained as a diastereomer mixture.

[4005] LC-MS (Method 1): R_f =1.1 min; MS (ESIpos): m/z =359 (M+H)⁺.

[4006] 64 mg (0.18 mmol) of this intermediate were taken up in 10 ml of dichloromethane, 1 ml of trifluoroacetic acid was added, and the mixture was stirred at RT for 30 min. This was followed by concentration under reduced pressure and

lyophilization of the remaining residue from water/dioxane. In this way, 66 mg (quant.) of the title compound were obtained as a foam.


[4007] HPLC (Method 6): R_f =1.45 min;

[4008] LC-MS (Method 3): R_f =1.12 min; MS (ESIpos): m/z =259 (M+H)⁺.

Intermediate 197

(2R,3R)-3-methoxy-2-methyl-N-[(2S)-1-(2-oxa-3-azabicyclo[2.2.2]oct-5-en-3-yl)-1-oxo-3-phenylpropan-2-yl]-3-[(2S)-pyrrolidin-2-yl]propanamide trifluoroacetate

[4009]



[4010] First, (2R,3R)-3-[(2S)-1-(tert-butoxycarbonyl)pyrrolidin-2-yl]-3-methoxy-2-methylpropanoic acid (Starting Compound 1) was released from 83 mg (0.18 mmol) of its dicyclohexylamine salt by taking it up in ethyl acetate and extractive shaking with 5% aqueous potassium hydrogensulphate solution. The organic phase was dried over magnesium sulphate, filtered and concentrated. The residue was taken up in 10 ml of DMF and admixed successively with mg (0.18 mmol) of (2S)-2-amino-1-(2-oxa-3-azabicyclo[2.2.2]oct-5-en-3-yl)-3-phenylpropan-1-one trifluoroacetate (Intermediate 196), 101 mg (0.266 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) and 93 μ l (0.53 mmol) of N,N-diisopropylethylamine. The mixture was stirred at RT for 30 min. The reaction mixture was then concentrated and the residue was purified by preparative HPLC. This gave 52 mg (56% of theory) of the Boc-protected intermediate tert-butyl (2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(2-oxa-3-azabicyclo[2.2.2]oct-5-en-3-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidine-1-carboxylate.

[4011] HPLC (Method 6): R_f =2.13 min;

[4012] LC-MS (Method 1): R_f =1.13 min; MS (ESIpos): m/z =528 (M+H)⁺.

[4013] 52 mg (0.1 mmol) of this intermediate were taken up in 10 ml of dichloromethane, 1 ml of trifluoroacetic acid was added, and the mixture was stirred at RT for 20 min. This was followed by concentration under reduced pressure and stirring of the remaining residue with 20 ml of diethyl ether. After 10 min, the mixture was filtered and the filter residue was dried under high vacuum. In this way, 39 mg (72% of theory) of the title compound were obtained.

[4014] HPLC (Method 6): R_f =1.62 min;

[4015] LC-MS (Method 1): R_f =0.68 min; MS (ESIpos): m/z =428 (M+H)⁺.

Intermediate 198

N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(2-oxa-3-azabicyclo[2.2.2]oct-5-en-3-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[4016]

[4017] 44.5 mg (0.071 mmol) of N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 4) were taken up in 10 ml of DMF and admixed successively with 38.6 mg (0.071 mmol) of (2R,3R)-3-methoxy-2-methyl-N-[(2S)-1-(2-oxa-3-azabicyclo[2.2.2]oct-5-en-3-yl)-1-oxo-3-phenylpropan-2-yl]-3-[(2S)-pyrrolidin-2-yl]propanamide trifluoroacetate (Intermediate 197), 32.5 mg (0.086 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) and 41 μ l (0.235 mmol) of N,N-diisopropylethylamine. The mixture was stirred at RT for 1 h. The reaction mixture was then concentrated under reduced pressure and the residue was taken up in ethyl acetate. The organic phase was washed successively with 5% aqueous citric acid solution and 5% aqueous sodium hydrogencarbonate solution, dried over magnesium sulphate, filtered and concentrated. This gave 73 mg (98% of theory) of the Fmoc-protected intermediate N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(2-oxa-3-azabicyclo[2.2.2]oct-5-en-3-yl)-1-oxo-3-phenylpropan-2-yl]amino]-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide.

[4018] HPLC (Method 6): R_f =2.78 min;


[4019] LC-MS (Method 3): R_f =2.96 min; MS (ESIpos): m/z =1047 (M+H)⁺.

[4020] 73 mg (0.071 mmol) of this intermediate were dissolved in 5 ml of DMF. After 0.5 ml of piperidine had been added, the reaction mixture was stirred at RT for 10 min. This was followed by concentration under reduced pressure, and the residue was digested repeatedly with diethyl ether. After the diethyl ether had been decanted off, the residue was purified by preparative HPLC (eluent: acetonitrile/0.1% aq. TFA). 16 mg (26% of theory) of the title compound were obtained as a foam.

[4021] HPLC (Method 6): R_f =1.94 min;

[4022] LC-MS (Method 3): R_f =1.71 min; MS (ESIpos): m/z =825 (M+H)⁺


[4023] ¹H NMR (400 MHz, DMSO-d₆): δ =8.9-8.6 (m, 3H), 8.4, 8.3, 8.1 and 8.0 (4d, 1H), 7.3-7.1 (m, 5H), 6.7-6.5 (m, 2H), 5.2-4.8 (m, 3H), 4.75-4.55 (m, 3H), 4.05-3.95 (m, 1H), 3.7-3.4 (m, 4H), 3.22, 3.17, 3.15, 3.05, 3.02 and 2.95 (6s, 9H), 3.0 and 2.7 (2 br. m, 2H), 2.46 (m, 3H), 2.4-1.2 (br. m, 13H), 1.1-0.85 (m, 18H), 0.75 (m, 3H) [further signals hidden under H₂O peak].



Intermediate 199

N-(4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(2-oxa-3-azabicyclo[2.2.2]oct-5-en-3-yl)-1-oxo-3-phenylpropan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl)-N-methyl-L-valinamide

[4024]



[4025] The title compound was prepared in analogy to Intermediates 193 and 194, proceeding from 23 mg (24 μ mol) of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S)-1-(2-oxa-3-azabicyclo[2.2.2]oct-5-en-3-yl)-1-oxo-3-phenylpropan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl)-N-methyl-L-valinamide trifluoroacetate (Intermediate 198).


[4026] HPLC (Method 12): R_f =1.9 min;

[4027] LC-MS (Method 2): R_f =2.1 min; MS (ESIpos): m/z =1118 (M+H)⁺.

Intermediate 200

N-[2-(2-{2-[2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethoxy]ethoxy}ethoxy)ethyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl)-N-methyl-L-valinamide

[4028]



[4029] The preparation was effected in analogy to Intermediates 174 and 175, commencing with the reductive alkylation of Intermediate 192 with Intermediate 172, subsequent deprotection and formation of the maleimide.


[4030] HPLC (Method 12): R_t =1.9 min;

[4031] LC-MS (Method 1): R_t =0.86 min; MS (ESIpos): m/z =1025 (M+H)⁺.

Intermediate 201

N-{6-[(bromoacetyl)amino]hexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4032]



[4033] 22 mg (0.023 mmol) of N-(6-aminoheptyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 101) were dissolved in 9.5 ml of THF and admixed at 0° C. with 4.2 µl of triethylamine. A solution of bromoacetyl chloride in THF was added dropwise and the reaction mixture was stirred at 0° C. for 30 min. The reaction mixture was concentrated and the residue was purified by preparative HPLC. Thus, 6.9 mg (26% of theory) of the title compound were obtained as a foam.


[4034] HPLC (Method 5): R_t =1.8 min;

[4035] LC-MS (Method 11): R_t =0.9 min; MS (ESIpos): m/z =1059 and 1061 (M+H)⁺.

Intermediate 202

N-{2-[2-(2-{3-[(2,5-dioxypyrrolidin-1-yl)oxy]-3-oxopropoxy}ethoxy)ethoxy]ethyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4036]



[4037] The preparation was at first effected in analogy to Intermediate 168, commencing with the reductive alkylation of Intermediate 192 with Intermediate 167 and subsequent hydrogenolytic cleavage of the benzyl ester of N-(2-{2-[2-(2-carboxyethoxy)ethoxy]ethoxy}ethyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide.

[4038] 13 mg (10 μ mol) of this intermediate were dissolved in 5 ml of DMF and admixed with 2.1 mg (20 mmol) of 1-hydroxypyrrolidine-2,5-dione, 6.5 μ l of N,N-diisopropylethylamine and 7.1 mg (0.02 mmol) of HATU. The reaction mixture was stirred at RT overnight and then concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization from acetonitrile/water, 9.2 mg (62% of theory) of the title compound were obtained.


[4039] HPLC (Method 12): R_f =2.0 min;

[4040] LC-MS (Method 2): R_f =2.1 min; MS (ESIpos): m/z =1141 (M+H)⁺.

Intermediate 203

tert-butyl 6-hydrazino-6-oxohexyl carbamate

[4041]




[4042] This compound was prepared by standard peptide chemistry methods, by coupling of 6-[(tert-butoxycarbonyl)amino]hexanoic acid with benzyl hydrazinecarboxylate in the presence of EDCI and HOBT, and subsequent hydrogenolytic cleavage of the benzyloxycarbonyl protecting group.

[4043] LC-MS (Method 11): R_f =0.59 min; MS (ESIpos): m/z =246 (M+H)⁺.

Intermediate 204

N-{4-[2-(6-aminohexanoyl)hydrazino]-4-oxobutyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[4044]



[4045] 146 mg (50 μ mol) of (N-(3-carboxypropyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were dissolved in 5 ml of DMF and then admixed with 30.6 mg (80 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, 19 μ l of N,N-diisopropylethylamine and with 22.4 mg (60 μ mol) of tert-butyl 6-hydrazino-6-oxohexyl carbamate. The reaction mixture was stirred at RT for 1.5 h. This was followed by concentration under high vacuum and purification of the remaining residue by means of preparative HPLC. Thus, 43 mg (68% of theory) of the protected intermediate were obtained, which were then taken up in 10 ml of dichloromethane and deprotected with 1 ml of trifluoroacetic acid. The reaction mixture was concentrated and the residue was stirred with dichloromethane, and the solvent was removed again under reduced pressure. Thus, 45 mg (68% of theory over 2 stages) of the title compound were obtained.


[4046] HPLC (Method 12): R_f =1.6 min;

[4047] LC-MS (Method 11): R_f =0.66 min; MS (ESIpos): m/z =983 (M+H)⁺.

Intermediate 205

N-(4-{2-[6-({[2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethyl]carbamoyl}amino)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4048]



[4049] This compound was prepared in analogy to Intermediate 114, proceeding from Intermediates 50 and 204.

[4050] Yield: 4 mg (78% of theory)


[4051] HPLC (Method 12): R_f =1.7 min;

[4052] LC-MS (Method 11): R_f =0.73 min; MS (ESIpos): m/z =1149 (M+H)⁺.

Intermediate 206

N-(6-{[3-({3-[(2,5-dioxopyrrolidin-1-yl)oxy]-3-oxopropyl}disulphanyl)propanoyl]amino}hexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4053]



[4054] 8 mg (10 μ mol) of Intermediate 101 were dissolved in 2 ml of DMF and admixed with 8.6 mg (20 μ mol) of 1,1'-{disulphanediylbis[(1-oxopropane-3,1-diyl)oxy]}dipyrrolidine-2,5-dione and 3.7 μ l of N,N-diisopropylethylamine. The reaction mixture was stirred at RT for 2 h and then the solvent was evaporated off under reduced pressure and the residue was purified by preparative HPLC. 7.2 mg (68% of theory) of the title compound were obtained.


[4055] HPLC (Method 5): R_f =1.9 min;

[4056] LC-MS (Method 11): R_f =0.94 min; MS (ESIpos): m/z =615 [$\frac{1}{2}$ (M+2H)⁺]

Intermediate 207

(1S,2R)-1-amino-2-phenylcyclopropanecarboxylic acid trifluoroacetate

[4057]



[4061] The title compound was prepared from 1 g (2.95 mmol) of commercially available 9H-fluoren-9-ylmethyl 6-hydroxyhexyl carbamate under standard conditions, by oxidation with sulphur trioxide-pyridine complex. 840 mg (85% of theory) of the title compound were obtained.


[4062] HPLC (Method 12): R_f =2.0 min;

[4063] LC-MS (Method 1): R_f =1.1 min; MS (ESIpos): m/z =338 (M+H)⁺.

Intermediate 209

N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S,2R)-1-carboxy-2-phenylcyclopropyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4064]




[4058] The title compound was obtained in quantitative yield by deprotecting 210 mg (0.76 mmol) of commercially available (1S,2R)-1-[(tert-butoxycarbonyl)amino]-2-phenylcyclopropanecarboxylic acid with trifluoroacetic acid.

[4059] LC-MS (Method 1): R_f =0.23 min; MS (ESIpos): m/z =178 (M+H)⁺.

Intermediate 208

9H-fluoren-9-ylmethyl 6-oxohexyl carbamate

[4060]



[4065] First, in analogy to the synthesis described in Intermediate 75, by coupling of N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) and (1S,2R)-1-amino-2-phenylcyclopropanecarboxylic acid trifluoroacetate (Intermediate 207) in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Boc protecting group by means of trifluoroacetic acid, the amine compound N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S,2R)-1-carboxy-2-phenylcyclopropyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide was prepared as the trifluoroacetate.

[4066] To 22 mg (0.026 mmol) of this compound in 10 ml of methanol were then added 17 mg (0.05 mmol) of 9H-fluoren-9-ylmethyl 6-oxohexyl carbamate (Intermediate 208) and 2.3 mg of acetic acid, and also 11.4 mg (0.12 mmol) of borane-pyridine complex. The reaction mixture was stirred at RT overnight. Then the same amounts of borane-pyridine complex and acetic acid, and also 8 mg of fluoren-9-ylmethyl 6-oxohexyl carbamate, were added once again and the reaction mixture was stirred at RT for a further 24 h. This was followed by concentration under reduced pressure, and the residue was purified by means of preparative HPLC. After concentration of the corresponding fractions, the product was used immediately in the next stage.

[4067] 33 mg of the still contaminated intermediate were taken up in 5 ml of DMF, and 1 ml of piperidine was added. After stirring at RT for 15 min, the reaction mixture was concentrated and the resulting residue was purified by preparative HPLC. Thus, 11 mg (55% of theory over 2 stages) of the aminocarboxylic acid intermediate were obtained.

[4068] HPLC (Method 12): $R_f=1.7$ min;

[4069] LC-MS (Method 11): $R_f=0.7$ min; MS (ESIpos): $m/z=843$ (M+H)⁺.

[4070] 6 mg (7.12 μ mol) of this intermediate were taken up in 1 ml of dioxane and then admixed with 6.6 mg (42.7 μ mol) of methyl 2,5-dioxo-2,5-dihydro-1H-pyrrole-1-carboxylate and with 5 μ l of saturated aqueous sodium hydrogencarbonate solution. The reaction mixture was stirred at RT for 1 h. Then another 3 portions each of 50 μ l of the saturated aqueous sodium hydrogencarbonate solution were added and the reaction mixture was stirred at RT for a further 30 min. Then the reaction mixture was acidified to pH 2 with trifluoroacetic acid and subsequently concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization from acetonitrile/water, 4 mg (60% of theory) of the title compound were obtained as a foam.


[4071] HPLC (Method 12): $R_f=1.9$ min;

[4072] LC-MS (Method 11): $R_f=0.88$ min; MS (ESIpos): $m/z=923$ (M+H)⁺.

Intermediate 210

N-{6-[(2,5-dioxopyrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4073]



[4074] First, 6-oxohexanoic acid was prepared by a literature method (J. org. Chem. 58, 1993, 2196-2200).

[4075] 80 mg (0.08 mmol) of N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 192) and 65.4 mg (0.5 mmol) of 6-oxohexanoic

acid were combined in 9 ml of methanol and admixed with 10 μ l of acetic acid and 37.4 mg (0.4 mmol) of borane-pyridine complex. The reaction mixture was stirred at RT overnight. This was followed by concentration under reduced pressure, and the residue was taken up in 1:1 acetonitrile/water and adjusted to pH 2 with trifluoroacetic acid. The reaction mixture was concentrated again and the residue was purified by means of preparative HPLC. After concentration of the corresponding fractions, 70 mg (86% of theory) of N-(5-carboxypentyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were obtained as the trifluoroacetate.

[4076] HPLC (Method 12): $R_f=1.9$ min;

[4077] LC-MS (Method 1): $R_f=0.87$ min; MS (ESIpos): $m/z=955$ (M+H)⁺.

[4078] ¹H NMR (500 MHz, DMSO-d₆, characteristic signals): $\delta=12.0$ (br. M, 1H), 10.8 (s, 1H), 9.4 (m, 1H), 8.9 and 8.8 (2d, 1H), 8.3 and 8.02 (2d, 1H), 7.5 (m, 1H), 7.3 (m, 1H), 7.15 and 7.1 (2s, 1H) 7.05-6.9 (m, 2H), 5.12 and 4.95 (2m, 1H), 4.7-4.5 (m, 2H), 4.1-3.8 (m, 4H), 3.75 (d, 1H), 3.25, 3.2, 3.18, 3.13, 2.98 and 2.88 (6s, 9H), 2.8 (m, 3H), 1.08 and 1.04 (2d, 3H), 0.95-0.8 (m, 15H), 0.8-0.65 (dd, 3H).

[4079] 22 mg (23 μ mol) of this intermediate were dissolved in 1.8 ml of dichloromethane and admixed with 13.2 mg (70 μ mol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 26.5 mg (230 μ mol) of 1-hydroxypyrrolidine-2,5-dione and 0.28 mg (2 μ mol) of dimethylaminopyridine, and the reaction mixture was stirred at RT for 2 h. Subsequently, the reaction mixture was concentrated under reduced

pressure and the remaining residue was purified by means of preparative HPLC. After lyophilization from acetonitrile/water, 21.3 mg (88% of theory) of the title compound were obtained.


[4080] HPLC (Method 12): $R_f=1.9$ min;

[4081] LC-MS (Method 1): $R_f=0.94$ min; MS (ESIpos): $m/z=1052$ (M+H)⁺.

Intermediate 211

N-{6-[(2,5-dioxopyrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(2S,3S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylbutan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4082]



[4083] 15 mg (20 μ mol) of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(2S,3S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylbutan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate (Intermediate 15) were reductively alkylated with 6-oxohexanoic acid, in analogy to Intermediate 210.

[4084] Yield: 9.2 mg (61% of theory)

[4085] HPLC (Method 12): R_f =1.9 min;

[4086] LC-MS (Method 1): R_f =0.87 min; MS (ESIpos): m/z =929 (M+H)⁺.

[4087] 9 mg (10 μ mol) of this intermediate were dissolved in 3 ml of DMF and admixed with 5.6 mg (48 μ mol) of 1-hydroxypyrrolidine-2,5-dione, 5 μ l of N,N-diisopropylethylamine and 5.5 mg (0.015 mmol) of HATU, and the reaction mixture was treated in an ultrasound bath for 6 h. In the course of this, 5.5 mg of HATU were added every hour. Subsequently, the reaction mixture was concentrated under reduced pressure, and the residue was taken up in acetonitrile/water and adjusted to pH 2 with trifluoroacetic acid. After

concentrating again under reduced pressure, the remaining residue was purified by means of preparative HPLC. After lyophilization from acetonitrile/water, 5.8 mg (57% of theory) of the title compound were obtained.


[4088] HPLC (Method 12): R_f =2.0 min;

[4089] LC-MS (Method 1): R_f =0.95 min; MS (ESIpos): m/z =1027 (M+H)⁺.

Intermediate 212

N-{2-[2-(2-{3-[(2,5-dioxopyrrolidin-1-yl)oxy]-3-oxopropoxy}ethoxy)ethoxy]ethyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(2S,3S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylbutan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4090]



[4091] The preparation was at first effected in analogy to Intermediate 168, commencing with the reductive alkylation of Intermediate 15 with Intermediate 167 and subsequent hydrogenolytic cleavage of the benzyl ester of N-(2-{2-[2-(2-carboxyethoxy)ethoxy]ethoxy}ethyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(2S,3S)-1-(1,2-oxazinan-2-yl)-1-oxo-3-phenylbutan-2-yl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide.

[4092] 8.4 mg (8 μ mol) of this intermediate were dissolved in 3 ml of DMF and admixed with 9.5 mg (80 μ mol) of 1-hydroxypyrrolidine-2,5-dione, 10 μ l of N,N-diisopropylethylamine and 9.4 mg (25 μ mol) of HATU, and the reaction mixture was stirred at RT overnight and then concentrated under reduced pressure. Subsequently, the reaction mixture was concentrated under reduced pressure, and the residue was taken up in acetonitrile/water and adjusted to pH 2 with trifluoroacetic acid. After concentrating again under reduced pressure, the remaining residue was purified by means of

preparative HPLC. After lyophilization from acetonitrile/water, 4 mg (32% of theory) of the title compound were obtained.


[4093] HPLC (Method 12): R_f =2.0 min;

[4094] LC-MS (Method 1): R_f =0.96 min; MS (ESIpos): m/z =1117 (M+H)⁺.

Intermediate 213

N-{6-[(trans-4-{[(2,5-dioxopyrrolidin-1-yl)oxy]carbonyl}cyclohexyl)amino]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4095]



[4096] This compound was prepared in analogy to Intermediate 104, proceeding from N-(5-carboxypentyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide, the synthesis of which was described under Intermediate 210. 9.3 mg of the title compound (37% of theory over 3 stages) were obtained.


[4097] HPLC (Method 12): R_f =1.9 min;

[4098] LC-MS (Method 1): R_f =0.9 min; MS (ESIpos): m/z =1177 (M+H)⁺.

Intermediate 214

N-{4-[(2,5-dioxopyrrolidin-1-yl)oxy]-4-oxobutyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[(1S,2R)-1-hydroxy-1-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4099]



[4100] This compound was prepared in analogy to Intermediate 210, by conversion of Intermediate 92 to the active ester.


[4101] HPLC (Method 5): $R_f=1.6$ min;

[4102] LC-MS (Method 11): $R_f=0.82$ min; MS (ESIpos): $m/z=901$ (M+H)⁺.

Intermediate 215

N-{6-[(2,5-dioxopyrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(1S,2R)-1-hydroxy-1-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4103]



[4104] First, Intermediate 40, in analogy to Intermediate 183, was used with borane-pyridine complex to prepare N-(5-carboxypentyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(1S,2R)-1-hydroxy-1-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide. From this compound, in analogy to Intermediate 210, the active ester was then generated. 34 mg (36% of theory over 2 stages) of the title compound were obtained.


[4105] HPLC (Method 5): $R_f=1.6$ min;

[4106] LC-MS (Method 1): $R_f=0.85$ min; MS (ESIpos): $m/z=930$ (M+H)⁺.

Intermediate 216

N-(4-{[(2,5-dioxopyrrolidin-1-yl)oxy]carbonyl}benzyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4107]



[4108] First, in analogy to the preparation of Intermediate 183, Intermediate 192 was reacted with 4-formylbenzoic acid with borane-pyridine complex to give N-(4-carboxybenzyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide. This compound was then used, in analogy to Intermediate 210, to generate 11 mg (68% of theory) of the title compound.


[4109] HPLC (Method 5): R_f =1.8 min;

[4110] LC-MS (Method 1): R_f =1.13 min; MS (ESIpos): m/z =1072 (M+H)⁺.

Intermediate 217

N-(5-carboxypentyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4111]



[4112] 53 mg (84 μ mol) of N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(2R,3S,4S)-1-carboxy-2-methoxy-4-methylhexan-3-yl]-N-methyl-L-valinamide (Intermediate 4) and 45 mg (84 μ mol) of benzyl N-[(2R,3R)-3-methoxy-2-methyl-3-[(2S)-pyrrolidin-2-yl]propanoyl]-L-phenylalaninate trifluoroacetate (Intermediate 12) were taken up in 2 ml of DMF, 19 μ l of N,N-diisopropylethylamine, 14 mg (92 μ mol) of HOBt and 17.6 mg (92 μ mol) of EDC were added and then the mixture was stirred at RT overnight. Subsequently, the reaction mixture was concentrated and the residue was purified by means of preparative HPLC. This gave 59 mg (68% of theory) of the Fmoc-protected intermediate N-[(9H-fluoren-9-ylmethoxy)carbonyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide.

[4113] LC-MS (Method 1): R_f =1.55 min; m/z =1044 (M+H)⁺.

[4114] 57 mg (0.055 mmol) of this intermediate were treated with 1.2 ml of piperidine in 5 ml of DMF to detach the Fmoc protecting group. After concentration and purification by means of preparative HPLC, 39 mg (76% of theory) of the free amine intermediate N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-(benzyloxy)-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide were obtained as the trifluoroacetate.

[4115] HPLC (Method 5): R_f =1.9 min;

[4116] LC-MS (Method 1): R_f =1.01 min; m/z =822 (M+H)⁺.

[4117] 60 mg (0.06 mmol) of this intermediate were reacted, in analogy to Intermediate 210, with 6-oxohexanoic acid in the presence of borane-pyridine complex. 45 mg (75% of theory) of the title compound were obtained as a foam.


[4118] HPLC (Method 5): R_f =1.9 min;

[4119] LC-MS (Method 1): R_f =0.97 min; MS (ESIpos): m/z =9936 (M+H)⁺.

Intermediate 218

N-{6-[(2,5-dioxopyrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-(benzyloxy)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4120]



[4121] This compound was prepared by conversion of 42 mg (0.05 mmol) of Intermediate 217 to the active ester.

[4122] Yield: 26 mg (54%)


[4123] HPLC (Method 5): R_t =2.1 min;

[4124] LC-MS (Method 1): R_t =1.01 min; MS (ESIpos): m/z =1034 (M+H)⁺.

Intermediate 219

N-{6-[(2,5-dioxopyrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(1S)-1-carboxy-2-phenylethyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4125]



[4126] 20 mg (0.02 mol) of the compound from Intermediate 218 were taken up in 2.4 ml of methanol and hydrogenated over 5% palladium on activated carbon under standard hydrogen pressure at RT for 30 min. The catalyst was then filtered off and the solvent was removed under reduced pressure. The residue was lyophilized from 1:1 acetonitrile/water. This gave 14 mg (92% of theory) of the title compound as a colourless foam.


[4127] HPLC (Method 5): R_t =1.7 min;

[4128] LC-MS (Method 1): R_t =0.86 min; MS (ESIpos): m/z =944 (M+H)⁺.

Intermediate 220

N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4129]



[4130] 0.5 g (1.01 mmol) of Intermediate 1 in 10 ml of dichloromethane were admixed with 1 ml of trifluoroacetic acid. After treatment in an ultrasound bath for 30 min, the mixture was concentrated and redistilled first with DCM and then with diethyl ether, and dried under high vacuum. The oily residue was used in the next stage, without further purification.

[4131] 500 mg of this intermediate were dissolved in 20 ml of DMF and admixed with 466 mg (3.8 mmol) of Intermediate 191, 382 mg (1.01 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) and 440 μ l (2.5 mmol) of N,N-diisopropylethylamine. The mixture was stirred at RT for 1 h and then concentrated. The residue was taken up in dichloromethane and extracted by shaking first twice with 5% aqueous citric acid solution and then with saturated aqueous sodium hydrogen-carbonate solution. The organic phase was concentrated and the residue was purified by flash chromatography on silica gel with 95:5 dichloromethane/methanol as the eluent. The corresponding fractions were combined and the solvent was removed under reduced pressure. After the residue had been dried under high vacuum, 562 mg (65% of theory over both stages) of the Z-protected intermediate were obtained.

[4132] 562 mg (0.57 mmol) of this intermediate were taken up in 50 ml of methanol and hydrogenated with 155 mg of 10% palladium on activated carbon under standard hydrogen pressure at RT for 20 min. The catalyst was then filtered off and the solvent was removed under reduced pressure. The residue was purified by preparative HPLC. The corresponding fractions were combined, the solvent was evaporated off

under reduced pressure and the residue was lyophilized from dioxane. This gave 361 mg (87% of theory) of the title compound as a foam.


[4133] HPLC (Method 5): double peak with R_f =1.75 and 1.86 min;

[4134] LC-MS (Method 1): double peak at R_f =0.84 min and 0.91 min with the same mass; MS (ESIpos): m/z =944 (M+H)⁺.

Intermediate 221

N-{(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropyl}-N-methyl-L-valine

[4135]



[4136] 100 mg (0.76 mmol) of commercially available N-methyl-L-valine and 285 mg (1.14 mmol) of commercially available tert-butyl (2S)-1-oxo-3-phenylpropan-2-yl carbamate were combined in 22 ml of methanol and admixed with 340 mg (3.66 mmol) of borane-pyridine complex and 70 μ l of acetic acid. The reaction mixture was stirred at RT overnight. This was followed by concentration under reduced pressure, and the residue was purified by flash chromatography on silica gel with dichloromethane/methanol/17% aqueous ammonia solution as the eluent. After concentration of the corresponding fractions and lyophilization from 1:1 dioxane/water, 259 mg (93% of theory) of the title compound were obtained.


[4137] HPLC (Method 12): R_f =1.6 min;

[4138] LC-MS (Method 11): R_f =0.76 min; MS (ESIpos): m/z =365 (M+H)⁺.

Intermediate 222

N-[(2S)-2-amino-3-phenylpropyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[4139]



[4140] 40 mg (0.11 mmol) of N-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropyl]-N-methyl-L-valine (Intermediate 221) were dissolved in 5 ml of DMF and admixed with 80 mg (0.11 mmol) of N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 220), 50 mg (0.13 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) and 57 μ l (2.5 mmol) of N,N-diisopropylethylamine. The mixture was stirred at RT for 1 h and then concentrated. The residue was taken up in ethyl acetate and washed first with 5% aqueous citric acid solution and then with water. The organic phase was concentrated and the residue was purified by means of preparative HPLC. The corresponding fractions were combined and the solvent was removed under reduced pressure. After lyophilization from dioxane, 60 mg (50% of theory) of the protected intermediate were obtained.

[4141] HPLC (Method 12): R_f =2.2 min;

[4142] LC-MS (Method 1): R_f =1.17 min; MS (ESIpos): m/z =1073 (M+H)⁺.

[4143] 60 mg (0.05 mmol) of this intermediate were taken up in 10 ml of dichloromethane, 2 ml of trifluoroacetic acid were added, and the reaction mixture was stirred at RT for 1.5

h. Subsequently, the reaction mixture was concentrated under reduced pressure and the remaining residue was purified by means of preparative HPLC. The corresponding fractions were combined, the solvent was removed under reduced pressure and the residue was lyophilized from dioxane/water. In this way, 25 mg (42% of theory) of the title compound were obtained as a foam.


[4144] HPLC (Method 12): R_f =1.9 min;

[4145] LC-MS (Method 1): R_f =0.95 min; MS (ESIpos): m/z =974 (M+H)⁺.

Intermediate 223

N-[(2S)-2-([2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethyl]carbamoyl)amino]-3-phenylpropyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4146]



[4147] The preparation was effected in analogy to Intermediate 134, proceeding from 5 mg (4.6 mmol) of Intermediate 222. 3.4 mg (65% of theory) of the title compound were obtained.


[4148] HPLC (Method 12): R_f =2.0 min;

[4149] LC-MS (Method 1): R_f =0.99 min; MS (ESIpos): m/z =1140 (M+H)⁺.

Intermediate 224

N-[(2S)-2-({[2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethyl]carbamoyl}amino)propyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4150]



[4151] The preparation was effected in analogy to the synthesis of Intermediate 223.


[4152] HPLC (Method 12): R_t =1.9 min;

[4153] LC-MS (Method 1): R_t =0.92 min; MS (ESIpos): m/z =1064 (M+H)⁺.

Intermediate 225

N-(2-aminoethyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[4154]



[4155] 100 mg (0.76 mmol) of commercially available N-methyl-L-valine and 182 mg (1.14 mmol) of commercially available tert-butyl 2-oxoethyl carbamate were combined in 20 ml of methanol and admixed with 340 mg (3.66 mmol) of borane-pyridine complex and 65 μ l of acetic acid. The reaction mixture was stirred at RT overnight. This was followed by concentration under reduced pressure, and the residue was purified by flash chromatography on silica gel with dichloromethane/methanol/17% aqueous ammonia solution (15/4/0.5) as the eluent. After concentration of the corresponding fractions and lyophilization from 1:1 dioxane/water, 190 mg

in 39% purity (35% of theory) of the intermediate were obtained, which were converted further without further purification.

[4156] 50 mg (0.07 mmol) of this intermediate were dissolved in 10 ml of DMF and admixed with 52 mg (0.07 mmol) of N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 220), 32 mg (0.09 mmol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate

phate (HATU) and 37 μ l (0.2 mmol) of N,N-diisopropylethylamine. The mixture was stirred at RT overnight and then concentrated. The residue was taken up in ethyl acetate and extracted by shaking first with 5% aqueous citric acid solution and then with water. The organic phase was concentrated and the residue was purified by means of preparative HPLC. The corresponding fractions were combined and the solvent was removed under reduced pressure. After lyophilization from dioxane, 53 mg (76% of theory) of the protected intermediate were obtained.

[4157] HPLC (Method 12): R_f =2.0 min;

[4158] LC-MS (Method 1): R_f =1.02 min; MS (ESIpos): m/z =984 (M+H)⁺.

[4159] 53 mg (0.05 mmol) of this intermediate were taken up in 10 ml of dichloromethane, 2 ml of trifluoroacetic acid were added, and the reaction mixture was stirred at RT for 30 min. Subsequently, the reaction mixture was concentrated under reduced pressure and the remaining residue was purified by means of preparative HPLC. The corresponding frac-

tions were combined, the solvent was removed under reduced pressure and the residue was lyophilized from dioxane/water. In this way, 21 mg (40% of theory) of the title compound were obtained in 65% purity.


[4160] HPLC (Method 12): R_f =1.7 min;

[4161] LC-MS (Method 1): R_f =0.87 min; MS (ESIpos): m/z =884 (M+H)⁺.

Intermediate 226

N-[2-({[2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethyl]carbamoyl}amino)ethyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4162]



[4163] The preparation was effected proceeding from Intermediate 225, in analogy to the synthesis of Intermediate 134. 11.6 mg (59% of theory) of the title compound were obtained.


[4164] HPLC (Method 12): R_f =1.9 min;

[4165] LC-MS (Method 1): R_f =0.90 min; MS (ESIpos): m/z =1050 (M+H)⁺.

Intermediate 227

N-{6-[(2,5-dioxopyrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-(benzyloxy)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4166]



[4167] This compound was prepared analogously to Intermediate 218, by conversion to the active ester.

[4168] Yield: 18 mg (51% of theory)


[4169] HPLC (Method 5): $R_f=2.1$ min;

[4170] LC-MS (Method 1): $R_f=0.98$ min; MS (ESIpos): $m/z=1073$ (M+H)⁺.

Intermediate 228

(2R,3S)-3-[(tert-butoxycarbonyl)amino]-4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutan-2-yl(3R,4S,7S,10S)-4-[(2S)-butan-2-yl]-7,10-diisopropyl-3-(2-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-2-oxoethyl)-5,11-dimethyl-6,9-dioxo-2-oxa-5,8,11-triazapentadecan-15-oate

[4171]



[4172] The title compound was prepared by coupling the Boc-protected intermediate obtained in the synthesis of Intermediate 154 with commercially available 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanehydrazide.


[4173] HPLC (Method 12): $R_f=2.1$ min;

[4174] LC-MS (Method 1): $R_f=0.97$ min; MS (ESIpos): $m/z=1308$ (M+H)⁺.

Intermediate 229

(2R,3S)-3-acetamido-4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutan-2-yl (3R,4S,7S,10S)-4-[(2S)-butan-2-yl]-7,10-diisopropyl-3-(2-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-2-oxoethyl)-5,11-dimethyl-6,9-dioxo-2-oxa-5,8,11-triazapentadecan-15-oate

[4175]



[4176] The title compound was prepared from 7.5 mg (2.5 μ mol) of Intermediate 154 by acetylation with 2.3 μ l of acetic anhydride in 1 ml of DMF in the presence of 0.4 μ l of N,N-diisopropylethylamine.

[4177] Yield: 1.4 mg (40% of theory)


[4178] HPLC (Method 12): R_t =1.9 min;

[4179] LC-MS (Method 1): R_t =0.86 min; MS (ESIpos): m/z =1250 (M+H)⁺.

Intermediate 230

(2R,3S)-3-[(tert-butoxycarbonyl)amino]-4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutan-2-yl (3R,4S,7S,10S)-4-[(2S)-butan-2-yl]-3-(2-{(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-2-oxoethyl)-7,10-diisopropyl-5,11-dimethyl-6,9-dioxo-2-oxa-5,8,11-triazapentadecan-15-oate

[4180]



[4181] This compound was prepared in analogy to Intermediate 228, proceeding from Intermediate 193. 16 mg (30% of theory over 3 stages) of the title compound were obtained.


[4182] HPLC (Method 12): R_t =2.0 min;

[4183] LC-MS (Method 1): R_t =1.02 min; MS (ESIpos): m/z =1335 (M+H)⁺.

Intermediate 231

(2R,3S)-3-acetamido-4-{2-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]hydrazino}-4-oxobutan-2-yl (3R,4S,7S,10S)-4-[(2S)-butan-2-yl]-3-(2-{(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-2-oxoethyl)-7,10-diisopropyl-5,11-dimethyl-6,9-dioxo-2-oxa-5,8,11-triazapentadecan-15-oate

[4184]



[4185] This compound was prepared from 8 mg (6 μ mol) of Intermediate 230, first by deprotection with trifluoroacetic acid and subsequent acetylation with acetic anhydride in DMF in the presence of N,N-diisopropylethylamine. 2 mg (37% of theory over 2 stages) of the title compound were obtained.


[4186] HPLC (Method 12): R_f =1.9 min;

[4187] LC-MS (Method 1): R_f =0.88 min; MS (ESIpos): m/z =1277 (M+H)⁺.

Intermediate 232

benzyl N-[(4-nitrophenoxy)carbonyl]-beta-alaninate

[4188]



[4189] 200 mg (0.57 mmol) of commercially available 4-methylbenzenesulphonic acid-benzyl beta-alaninate and 229 mg (1.14 mmol) of 4-nitrophenyl chlorocarbonate were taken up in 15 ml of tetrahydrofuran and the reaction mixture was then heated to reflux for 30 min. Subsequently, the reaction mixture was concentrated under reduced pressure and the residue was purified by means of preparative HPLC. After concentration of the corresponding fractions and drying of the residue under high vacuum, 86 mg (44% of theory) of the title compound were obtained.


[4190] HPLC (Method 12): R_f =1.8 min;

[4191] LC-MS (Method 1): R_f =1.07 min; MS (ESIpos): m/z =345 (M+H)⁺.

Intermediate 233

N-{2-[(3-[(2,5-dioxopyrrolidin-1-yl)oxy]-3-oxopropyl}carbamoyl)amino]ethyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4192]



[4193] 13 mg (10 μ mol) of Intermediate 225 and 6.7 mg (20 μ mol) of Intermediate 232 were dissolved in 3 ml of DMF, and then 7 μ l of N,N-diisopropylethylamine were added. The mixture was stirred at RT overnight and then concentrated under high vacuum. The remaining residue was purified by means of preparative HPLC. After concentration of the corresponding fractions and drying of the residue under high vacuum, 5.4 mg (38% of theory) of the protected intermediate were obtained.

[4194] HPLC (Method 5): R_f =2.1 min;

[4195] LC-MS (Method 1): R_f =0.6 in; MS (ESIpos): m/z =1089 (M+H)⁺.

[4196] 5.4 mg (5 μ mol) of this intermediate were dissolved in 5 ml of methanol and, after addition of 2 mg of 10% palladium on activated carbon, hydrogenated under standard hydrogen pressure at RT for 20 min. The catalyst was then filtered off and the solvent was removed under reduced pressure. After the residue had been dried under high vacuum, 5 mg (quant.) of the acid intermediate were obtained.

[4197] HPLC (Method 12): R_f =1.8 min;

[4198] LC-MS (Method 1): R_f =0.84 min; MS (ESIpos): m/z =999 (M+H)⁺.

[4199] 5 mg (10 μ mol) of this intermediate were dissolved in 1 ml of DMF and admixed with 5.8 mg (50 mmol) of 1-hydroxypyrrolidine-2,5-dione and then with 2.6 μ l of N,N-diisopropylethylamine and 3.8 mg (10 μ mol) of HATU. After stirring at RT for 20 h, the reaction mixture was concentrated under reduced pressure. The remaining residue was purified by means of preparative HPLC. After lyophilization from 1:1 dioxane/water, 1.1 mg (20% of theory) of the title compound were obtained.


[4200] HPLC (Method 12): R_f =1.9 min;

[4201] LC-MS (Method 1): R_f =0.87 min; MS (ESIpos): m/z =1096 (M+H)⁺.

Intermediate 234

N-(6-[[[(benzyloxy)carbonyl]amino]hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino]propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4202]



[4203] 25 mg (30 μ mol) of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino]propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 55) and 45 mg (180 μ mol) of benzyl 6-oxohexyl carbamate were taken up in 3 ml of methanol and acidified with acetic acid. At room temperature, 15 μ l (144 μ mol; 9.4M) of borane-pyridine complex were subsequently added. The mixture was subsequently stirred at RT for 24 h, and acetic acid and 15 μ l (144 μ mol; 9.4M) of borane-pyridine complex were added again after 8 h. The reaction mixture was subsequently adjusted to pH 2 with TFA and purified by means of preparative HPLC. The product fractions were combined and concentrated, and


the residue was dried under high vacuum. This gave 15 mg (46% of theory) of the title compound as a foam.

[4204] LC-MS (Method 1): R_t =1.03 min; m/z =1066 (M+H)⁺.

Intermediate 235

N-(6-aminoheptyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino]propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4205]




[4206] 15 mg (14 μ mol) of N-(6-[(benzyloxy)carbonyl]amino)hexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 234) were taken up in 3 ml of methanol, and 1.8 mg of palladium on charcoal (5%) were added. The reaction mixture was subsequently hydrogenated under standard hydrogen pressure at RT for 2 h. The catalyst was then filtered off and the solvent was removed under reduced pressure. The residue was lyophilized from 1:1 acetonitrile/water. 11 mg (86% of theory) of the title compound were obtained as a foam.

[4207] LC-MS (Method 1): R_t =0.81 min; m/z =932 (M+H)⁺.

Intermediate 236

N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4208]




[4209] 11 mg (12 μ mol) of N-(6-aminoheptyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 235) were taken up in 500 μ l of 1:1 dioxane/water and admixed with 253 μ l of 1M aqueous sodium hydrogencarbonate solution and then with 2.8 mg (18 μ mol) of methyl 2,5-dioxo-2,5-dihydro-1H-pyrrole-1-carboxylate. The reaction mixture was stirred at RT for 30 min and then acidified with trifluoroacetic acid. The reaction mixture was purified by means of preparative HPLC. After lyophilization, 0.8 mg (7% of theory) of the title compound was obtained.

[4210] LC-MS (Method 1): R_t =1.01 min; m/z =1012 (M+H)⁺.

Intermediate 237

N-(5-carboxypentyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4211]




[4212] 25 mg (30 μ mol) of N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 55) and 23 mg (180 μ mol) of 6-oxohexanoic acid were taken up in 3 ml of methanol and acidified with acetic acid. At room temperature, 15 μ l (144 μ mol; 9.4M) of borane-pyridine complex were subsequently added. The reaction mixture was subsequently stirred at RT for 20 h, and acetic acid and 15 μ l (144 μ mol; 9.4M) of borane-pyridine complex were added again after 8 h. The reaction mixture was subsequently adjusted to pH 2 with trifluoroacetic acid and purified by means of preparative HPLC. The product fractions were combined and concentrated, and the residue was lyophilized. 21 mg (74% of theory) of the title compound were thus obtained as a foam.

[4213] LC-MS (Method 1): R_t =0.91 min; m/z =947 (M+H)⁺.

Intermediate 238

N-{6-[(2,5-dioxopyrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(5-phenyl-1,3,4-oxadiazol-2-yl)ethyl]amino}propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4214]




[4215] 21 mg (22 μ mol) of Intermediate 237 were dissolved in 1 ml of DMF and admixed with 38 mg (333 μ mol) of 1-hydroxypyrrolidine-2,5-dione and then with 2.4 mg (10 μ mol) of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) and 19 μ l of N,N-diisopropylethylamine. After stirring at RT for 2 h, the reaction mixture was purified by means of preparative HPLC. After lyophilization from dioxane, 22 mg (96% of theory) of the title compound were obtained.

[4216] LC-MS (Method 1): R_t =0.95 min; m/z =1044 (M+H)⁺.

Intermediate 239

N-methyl-L-threonyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[4217]



[4218] First, N-[(benzyloxy)carbonyl]-N-methyl-L-threonine was released from 237 mg (0.887 mmol) of its dicyclohexylamine salt by taking it up in ethyl acetate and extractive shaking with 5% aqueous sulphuric acid. The organic phase was dried over magnesium sulphate, filtered and concentrated. 14.7 mg (0.055 mmol) of N-[(benzyloxy)carbonyl]-N-methyl-L-threonine were taken up in 3 ml of DMF and admixed successively with 40 mg (0.055 mmol) of Intermediate 220, 12.7 mg (0.066 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 10 mg (0.066 mmol) of 1-hydroxy-1H-benzotriazole hydrate. The mixture was subsequently stirred at RT for 2 h. The solvent was then removed under reduced pressure and the residue purified by preparative HPLC. 29 mg (54% of theory) of the Z-protected intermediate were thus obtained.

[4219] LC-MS (Method 1): $R_f=1.15$ min; MS (ESIpos): $m/z=976$ (M+H)⁺.

[4220] 29 mg (0.003 mmol) of this intermediate were dissolved in 5 ml of methanol and hydrogenated over 5 mg of 5%


palladium/charcoal at RT and standard pressure for 1 h. The catalyst was subsequently filtered off and the solvent was evaporated off. The remaining residue was purified by preparative HPLC. 17 mg (54% of theory) of the title compound were obtained.

[4221] LC-MS (Method 1): $R_f=0.77$ min; MS (ESIpos): $m/z=842$ (M+H)⁺.

Intermediate 240

N-{6-[(2,5-dioxopyrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-threonyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[[[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4222]



[4223] This compound was prepared in analogy to Intermediate 210 from 15.6 mg (0.016 mmol) of Intermediate 239. 10.8 mg (67% of theory over 2 stages) of the title compound were obtained.


[4224] HPLC (Method 5): $R_f=1.7$ min;

[4225] LC-MS (Method 1): $R_f=0.85$ min; MS (ESIpos): $m/z=1053$ (M+H)⁺.

Intermediate 241

N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[[[(2S)-3-(4-hydroxyphenyl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[4226]



[4227] First, in analogy to Intermediate 5, trifluoroacetic acid-(2S)-2-amino-3-(4-hydroxyphenyl)-1-(1,2-oxazinan-2-yl)propan-1-one (1:1) was prepared. This reagent was then used, in analogy to the synthesis described in Intermediate 75, by coupling with N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Boc protecting group by means of trifluoroacetic acid, to prepare the title compound.


[4228] HPLC (Method 12): R_f =1.7 min;

[4229] LC-MS (Method 1): R_f =0.85 min; MS (ESIpos): m/z =817 (M+H)⁺.

Intermediate 242

N-{6-[(2,5-dioxopyrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(4-hydroxyphenyl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4230]



[4231] 50 mg (0.05 mmol) of Intermediate 241 were reacted, in analogy to Intermediate 210, with 6-oxohexanoic acid in the presence of borane-pyridine complex. Subsequently, 22.5 mg (0.02 mmol) of the resulting acid were converted to the activated ester. 13.5 mg (36% of theory over 2 stages) of the title compound were obtained.


[4232] HPLC (Method 12): R_f =1.8 min;

[4233] LC-MS (Method 1): R_f =0.86 min; MS (ESIpos): m/z =1028 (M+H)⁺.

Intermediate 243

N-(6-aminoheptyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(4-hydroxyphenyl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4234]



[4235] The preparation was effected in analogy to Intermediate 78, by reductive alkylation of Intermediate 241 with benzyl 6-oxohexyl carbamate and borane-pyridine complex and subsequent hydrogenation in methanol as the solvent.

[4236] Yield: 17.5 mg (34% of theory over 2 stages)


[4237] HPLC (Method 12): R_f =1.7 min;

[4238] LC-MS (Method 1): R_f =0.63 min; MS (ESIpos): m/z =916 (M+H)⁺.

Intermediate 244

N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(4-hydroxyphenyl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4239]



[4240] The preparation was effected in analogy to Intermediate 166, proceeding from Intermediate 243.

[4241] Yield: 1.3 mg (12% of theory)


[4242] HPLC (Method 12): R_f =1.9 min;

[4243] LC-MS (Method 1): R_f =0.89 min; MS (ESIpos): m/z =996 (M+H)⁺.

Intermediate 245

2,5-dioxopyrrolidin-1-yl O-[(3R,4S,7S,10S)-4-[(2S)-butan-2-yl]-3-(2-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-2-oxoethyl)-7,10-diisopropyl-5,11-dimethyl-6,9,15-trioxo-2-oxa-5,8,11-triazapentadecan-15-yl]-N-(tert-butoxycarbonyl)-L-threonyl-beta-alaninate

[4244]



[4245] First, Intermediate 193, as described for Intermediate 154, was reacted with benzyl N-(tert-butoxycarbonyl)-L-threoninate and then the benzyl ester was removed by hydrogenolysis. 30 mg (0.027 mmol) of the N-[4-((1S,2R)-1-[(tert-butoxycarbonyl)amino]-1-carboxypropan-2-yl)oxy)-4-oxobutyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide thus obtained were then coupled with 4-methylbenzenesulphonic acid-benzyl beta-alaninate in the presence of HATU and the benzyl ester was removed again by hydrogenolysis (yield: 24 mg (71% of theory over 2 stages)). Finally, 10 mg (0.008 mmol) of the resulting acid were converted to the activated ester. After HPLC purification, 2.7 mg (23% of theory) of the title compound were obtained.


[4246] HPLC (Method 5): $R_t=1.9$ min;

[4247] LC-MS (Method 1): $R_t=1.01$ min; MS (ESIpos): $m/z=1295$ (M+H)⁺

Intermediate 246a

(2S)-2-amino-1-(4-hydroxy-1,2-oxazolidin-2-yl)-3-(1H-indol-3-yl)propan-1-one trifluoroacetate (Diastereomer 1)

[4248]



[4249] 1.6 g (3.982 mmol) of 2,5-dioxopyrrolidin-1-yl N-(tert-butoxycarbonyl)-L-tryptophanate were dissolved in 15 ml of DMF and admixed with 500 mg (3.982 mmol) of 1,2-oxazolidin-4-ol and 100 μ l of N,N-diisopropylethylamine. The reaction mixture was stirred at RT overnight. Then another 100 μ l of N,N-diisopropylethylamine were added, and the mixture was first treated in an ultrasound bath for 5 h, then stirred at RT overnight, and subsequently concentrated under reduced pressure. The remaining residue was taken up in ethyl acetate and extracted first twice with 5% aqueous citric acid solution, then with saturated aqueous sodium hydrogencarbonate solution and finally with water. The organic phase was concentrated and the residue was separated into the diastereomers by flash chromatography on

silica gel with 95:5 dichloromethane/methanol as the eluent. The corresponding fractions of both diastereomers were combined and the solvent was removed under reduced pressure. After the residues had been dried under high vacuum, 272 mg (18% of theory) of Diastereomer 1 ($R_f=0.18$ (95:5 dichloromethane/methanol) and 236 mg (16% of theory) of Diastereomer 2 ($R_f=0.13$ (95:5 dichloromethane/methanol), and also 333 mg (22% of theory) of a mixed fraction of the Boc-protected intermediates were obtained.

[4250] Under standard conditions, 5 ml of trifluoroacetic acid in 20 ml of dichloromethane were used to detach the Boc protecting group from 272 mg (725 μ mol) of Diastereomer 1 of this intermediate and, after lyophilization from dioxane/water, 290 mg (quant) of the title compound were obtained in 75% purity and used in the next stage without further purification.


[4251] HPLC (Method 12): $R_t=1.1$ min;

[4252] LC-MS (Method 13): $R_t=1.80$ min; MS (ESIpos): $m/z=276$ (M+H)⁺

Intermediate 246b

(2S)-2-amino-1-(4-hydroxy-1,2-oxazolidin-2-yl)-3-(1H-indol-3-yl)propan-1-one trifluoroacetate (Diastereomer 2)

[4253]




[4254] Under standard conditions, 5 ml of trifluoroacetic acid in 20 ml of dichloromethane were used to detach the Boc protecting group from 236 mg (630 μ mol) of Diastereomer 2 of the intermediate described in 246a and, after concentration, stirring with diethyl ether and drying of the residue under high vacuum, 214 mg (76%) of the title compound were obtained.

[4255] LC-MS (Method 13): $R_t=1.84$ min; MS (ESIpos): $m/z=276$ (M+H)⁺

Intermediate 247a

N-{6-[(2,5-dioxypyrrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[[[(2S)-1-(4-hydroxy-1,2-oxazolidin-2-yl)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Diastereomer 1)

[4256]



[4257] To synthesize this compound, the coupling of Intermediates 26 and 246a with subsequent detachment of the Boc protecting group was first performed as described for Intermediate 74. Subsequently, the alkylation with 6-oxohexanoic acid in the presence of borane-pyridine complex and subsequent conversion of the acid to the active ester were performed as described for Intermediate 210. The title compound was purified by preparative HPLC.


[4258] HPLC (Method 12): R_f =1.8 min;

[4259] LC-MS (Method 1): R_f =0.86 min; MS (ESIpos): m/z =1053 (M+H)⁺

Intermediate 247b

N-{6-[(2,5-dioxypyrrrolidin-1-yl)oxy]-6-oxohexyl}-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[[[(2S)-1-(4-hydroxy-1,2-oxazolidin-2-yl)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Diastereomer 2)

[4260]



[4261] To synthesize this compound, the coupling of Intermediates 26 and 246b with subsequent detachment of the Boc protecting group was first performed as described for Intermediate 74. Subsequently, the alkylation with 6-oxohexanoic acid in the presence of borane-pyridine complex and subsequent conversion of the acid to the active ester were performed as described for Intermediate 210. The title compound was purified by preparative HPLC.


[4262] HPLC (Method 12): $R_f=1.8$ min;

[4263] LC-MS (Method 1): $R_f=0.86$ min; MS (ESIpos): $m/z=1053$ (M+H)⁺

Intermediate 248

N-(5-carboxypentyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-tert-butoxy-3-(4-hydroxyphenyl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4264]



[4265] First, in analogy to the synthesis described in Intermediate 86, by coupling N-(tert-butoxycarbonyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-2-carboxy-1-methoxypropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide (Intermediate 26) and tert-butyl-L-tyrosinate in the presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and subsequent detachment of the Boc protecting group by means of trifluoroacetic acid to obtain the tert-butyl ester (stirring with trifluoroacetic acid in dichloromethane for 40 min), the amine compound tert-butyl N-[(2R,3R)-3-methoxy-3-{(2S)-1-[(3R,4S,5S)-3-methoxy-5-methyl-4-(methyl {(2S)-3-methyl-2-[(N-methyl-L-valyl)amino]butyl}amino)heptanoyl]pyrrolidin-2-yl}-2-methylpropanoyl]-tyrosinate was prepared as the trifluoroacetate. 38 mg (0.04 mmol) of this compound were then used, in analogy to the preparation of Intermediate 210, by reaction with 6-oxohexanoic acid in the presence of borane-pyridine complex, to obtain 31 mg (99% of theory) of the title compound.

[4266] HPLC (Method 12): $R_f=1.8$ min;

[4267] LC-MS (Method 1): $R_f=0.88$ min; MS (ESIpos): $m/z=918$ (M+H)⁺.

B: PREPARATION OF ANTIBODY-DRUG CONJUGATES (ADCs)

[4268] The intermediates described above were linked, for example, to the anti-mesothelin antibody MF-Ta, with the linkage having taken place in accordance with the methods

set out below, via cysteine or lysine side chains, alternatively, of the antibody protein. The anti-mesothelin antibody MF-Ta was prepared in analogy to the methods described in WO 2009/068204-A1. The antibody MF-Ta was expressed in eukaryotic CHO cells (stable cell line) and purified via protein A and gel filtration before it was subjected to the conjugation in D-PBS buffer.

B-1. General Procedure 1 (Coupling Via Cysteine):


[4269] Added to a solution of the corresponding antibody in PBS buffer in the concentration range between 1 mg/ml and 15 mg/ml were 3 equivalents of tris(2-carboxyethyl) phosphine hydrochloride (TCEP), in solution in PBS buffer, and the mixture was stirred at RT for 1 hour. Subsequently, depending on the desired loading, between 2 and 10 equivalents of the maleimide precursor compound or halide precursor compound for coupling (Intermediate 102, 103, 105-109, 111-114, 117-126, 128, 129, 132-146, 148-155, 157, 159-161, 166, 171, 175-177, 184, 189, 194-195, 199-201, 205, 209, 223-224, 226, 228-231, 236 and 244) were added as a solution in DMSO. The amount of DMSO here ought not to

exceed 10% of the overall volume. The batch was stirred at RT for 60-120 minutes and then applied to PD 10 columns (Sephadex® G-25, GE Healthcare) equilibrated with PBS, and eluted with PBS buffer. Optionally a concentration procedure was carried out additionally by means of ultracentrifugation. If necessary, for more effective removal of low molecular mass constituents, the concentration by ultrafiltration was repeated after re-dilution with PBS buffer.

[4270] Normally, unless otherwise indicated, 5 mg of the corresponding antibody in PBS buffer were used for the reduction and the subsequent coupling. Following purification via the PD10 column, this gave, in each case, solutions of the corresponding ADC in 3.5 ml of PBS buffer. The particular protein concentration indicated was then determined for these solutions. Furthermore, the loading of the antibody (drug/mAb ratio) was determined in accordance with the methods described in B-4.

[4271] This process was used to prepare the immunoconjugates represented in Examples 1, 2, 5-19, 21-28, 30, 31, 33-37, 39-46, 48, 51-57, 59, 60, 71, 73-76, 78-80, 82-84, 86-87, 96, 98-103, 108 and 112.

[4272] In the structural formulae illustrated, the definition of AK₁ is as follows



where

[4273] §1 denotes the link with the succinimide group,

[4274] MF-Ta (partially reduced) is the partially reduced MF-Ta antibody (heavy chain SEQ ID NO: 408 and light chain SEQ ID NO: 409)

and

[4275] S stands for the sulphur atom of a cysteine residue of the partially reduced antibody.


B-2. General Operating Procedure 2 (Coupling Via Lysine Side Chains):

[4276] Added to a solution of the corresponding antibody in PBS buffer in the concentration range between 1 mg/ml and 15 mg/ml were, depending on the desired loading, between 2 and 5 equivalents of the precursor compound for coupling (Intermediate 104, 110, 115, 116, 127, 130, 131, 147, 156, 158, 162, 169, 178, 185, 190, 202, 206, 210-216, 218, 219, 227, 233, 238, 240, 242, 245, 247a and 247b) as a solution in DMSO. After 30 minutes of stirring at RT, the same amount of precursor compound in DMSO was added again. Alternatively it was possible to add 4-10 equivalents of the precursor compound for coupling, in one go. The amount of DMSO here ought not to exceed 10% of the overall volume. After a further 30 minutes of stirring at RT, the batch was applied to PD 10 columns (Sephadex® G-25, GE Healthcare) equilibrated with PBS, and eluted with PBS buffer. Optionally a concentration procedure was carried out additionally by means of ultracentrifugation. If necessary, for more effective removal of low molecular mass constituents, the concentration by ultrafiltration was repeated after re-dilution with PBS buffer.

[4277] Normally, unless otherwise indicated, 5 mg of the corresponding antibody in PBS buffer were used for the coupling. Following purification via the PD10 column, this gave, in each case, solutions of the corresponding ADC in 3.5 ml of PBS buffer. The particular protein concentration indicated was then determined for these solutions and the loading of the antibody (drug/mAb ratio) was determined in accordance with the methods described in B-4.

[4278] This process was used to prepare the immunoconjugates represented in Examples 3, 4, 20, 29, 32, 38, 47, 49, 50, 58, 61, 72, 77, 81, 85, 88-95, 97, 104, 109-111 and 113-115.

[4279] In the structural formulae illustrated, the definition of AK_2 is as follows



where

[4280] §^2 denotes the link with the carbonyl group,

MF-Ta is the non-reduced MF-Ta antibody (heavy chain SEQ ID NO: 408 and light chain SEQ ID NO: 409)


and

NH stands for the side chain amino group of a lysine residue of the antibody.

B-3a. General Process for Preparing Cysteine Adducts:

[4281] 10 μmol of the above-described maleimide precursor compounds were taken up in 3 ml of DMF and admixed with 2.1 mg (20 μmol) of L-cysteine. The reaction mixture was stirred at RT for 2 hours, then concentrated under reduced pressure and subsequently purified by preparative HPLC.

[4282] In the structural formulae illustrated, the definition of Cys is as follows



where

§^3 denotes the link with the linker-toxophore unit.

B-3b. General Process for Preparing Lysine Adducts:

[4283] 10 μmol of the above-described active ester precursor compounds were taken up in 5 ml of DMF and admixed, in the presence of 30 μmol of N,N-diisopropylethylamine, with α -amino-protected L-lysine. The reaction mixture was stirred at RT for 2 hours, then concentrated under reduced pressure and subsequently purified by preparative HPLC. The protective group was then removed by known methods.

Further Purification and Characterization of the Conjugates of the Invention

[4284] After reaction had taken place, in certain cases the reaction mixture was concentrated, by ultrafiltration, for example, and then desalted and purified by means of chromatography, for example using a Sephadex® G-25. Elution took place with, for example, phosphate-buffered saline (PBS). The solution was subsequently subjected to sterile filtration and freezing. An alternative option is to lyophilize the conjugate.

[4285] B-4. Determination of the toxophore loading

[4286] The toxophore loading of the resultant solutions of the conjugates described in the working examples, in PBS buffer, was determined as follows:

[4287] The toxophore loading of lysine-linked ADCs was determined by mass-spectrometric determination of the molecular weights of the individual conjugate species. In this case, to start with, the antibody conjugates were deglycosylated by means of PNGaseF, and the sample was acidified and, following HPLC separation, was analysed by mass spectrometry using an ESI-MicroTofQ (Bruker Daltonik). All of the spectra were added via the signal in the TIC (Total Ion Chromatogram), and the molecular weight of the various conjugate species was calculated on the basis of MaxEnt Deconvolution. Following signal integration of the different species, the DAR (Drug/Antibody Ratio) was then calculated.

[4288] For protein identification, in addition to the molecular weight determination, a tryptic digestion was carried out after deglycosylation and/or denaturing, and this digestion, after denaturing, reduction and derivatization, confirmed the identity of the protein on the basis of the tryptic peptides detected.

[4289] The toxophore loading of cysteine-linked conjugates was determined via reversed-phase chromatography of the reduced and denatured ADC. The ADC solution (1 mg/mL, 50 μL) was admixed with guanidinium hydrochloride (GuHCl) (28.6 mg) and with a solution of DL-dithiothreitol (DTT) (500 mM, 3 μL). The mixture was incubated at 55° C. for an hour and analysed by HPLC.

[4290] The HPLC analysis was carried out on an Agilent 1260 HPLC System with detection at 220 nm. The column used was a Polymer Laboratories PLRP-S Polymeric Reversed Phase column (catalogue number PL1912-3802) (2.1×150 mm, 8 µm particle size, 1000 Å) with a flow rate of 1 mL/min, using the following gradient: 0 min, 25% B; 3 min, 25% B; 28 min, 50% B. Eluent A consisted of 0.05% trifluoroacetic acid (TFA) in water, eluent B of 0.05% trifluoroacetic acid in acetonitrile.

[4291] The peaks detected were assigned by retention time comparison with the light chain (L0) and the heavy chain (H0) of the unconjugated antibody. Peaks which were detected exclusively in the conjugated sample were assigned to the light chain, with a toxophore (L1), and to the heavy chains, with one, two and three toxophores (H1, H2, H3).

[4292] The average loading of the antibody with toxophores was calculated as follows: first of all, the light-chain loading was calculated from the peak areas—determined by integration—of the peaks L0 and L1 belonging to the light chains, as the sum of the toxophore number weighted integration results of L0 and L1, divided by the sum of the singularly weighted integration results of L0 and L1. In the same way, the heavy-chain loading was calculated from the peak areas—determined by integration—of the peaks H0, H1, H2 and H3, belonging to the heavy chains, as the sum of the toxophore number weighted integration results of H0, H1, H2 and H3, divided by the sum of the singularly weighted

integration results of H0, H1, H2 and H3. The DAR is given by the light-chain loading and the heavy-chain loading, as the twofold sum of light-chain loading and heavy-chain loading. The factor 2 takes account of the fact that an antibody consists of two light chains and two heavy chains. In certain individual cases it may be impossible exactly to determine the toxophore loading, owing to co-elutions of certain peaks.

B-5. Testing of the Antigen Binding of the ADC


[4293] The binding capacity of the binder to the target molecule was tested after coupling had taken place. The skilled worker knows of diverse methods for achieving this—for example, the affinity of the conjugate can be tested by means of ELISA technology or surface plasmon resonance analysis (BIAcore™ measurements). The conjugate concentration can be measured by the skilled person using common methods—for example, for antibody conjugates, by means of protein determination (see also Doronina et al.; *Nature Biotechnol.* 2003; 21:778-784 and Polson et al., *Blood* 2007; 110:616-623).

Working Examples

Immunoconjugates

Example 1

[4294]




[4295] Protein concentration: 0.96 mg/ml

[4296] Drug/mAb Ratio: 3.1

Example 2

[4297]




[4298] Protein concentration: 0.44 mg/ml

[4299] Drug/mAb Ratio: 4.6

Example 3

[4300]




[4301] Protein concentration: 1.09 mg/ml

[4302] Drug/mAb Ratio: 2.1

Example 4

[4303]




[4304] Protein concentration: 0.87 mg/ml

[4305] Drug/mAb Ratio: 3.8

Example 5

[4306]




[4307] Protein concentration: 0.45 mg/ml

[4308] Drug/mAb Ratio: 6.5

Example 6

[4309]




[4310] Protein concentration: 0.15 mg/ml

[4311] Drug/mAb Ratio: 3.1

Example 7

[4312]




[4313] Protein concentration: 0.94 mg/ml

[4314] Drug/mAb Ratio: 2.8

Example 8

[4315]




[4316] Protein concentration: 0.45 mg/ml

[4317] Drug/mAb Ratio: 0.9

Example 9

[4318]




[4319] Protein concentration: 0.51 mg/ml

[4320] Drug/mAb Ratio: 6.6

Example 10

[4321]




[4322] Protein concentration: 0.47 mg/ml

[4323] Drug/mAb Ratio: 4.2

Example 11


[4324]



[4326] Drug/mAb Ratio: 5.9

Example 12

[4327]




[4328] Protein concentration: 0.47 mg/ml

[4329] Drug/mAb Ratio: 3.3

Example 13

[4330]




[4331] Protein concentration: 0.53 mg/ml

[4332] Drug/mAb Ratio: 2.8

Example 14

[4333]




[4334] Protein concentration: 0.92 mg/ml

[4335] Drug/mAb Ratio: 3.5

Example 15

[4336]




[4337] Protein concentration: 0.09 mg/ml

[4338] Drug/mAb Ratio: nd

Example 16

[4339]





[4340] Protein concentration: 0.62 mg/ml

[4341] Drug/mAb Ratio: 1.8

Example 17

[4342]






[4352] Protein concentration: 1.1 mg/ml

[4353] Drug/mAb Ratio: 0.3

Example 21

[4354]




[4355] Protein concentration: 0.61 mg/ml

[4356] Drug/mAb Ratio: 0.9

Example 22

[4357]




[4358] Protein concentration: 0.57 mg/ml

[4359] Drug/mAb Ratio: 1.2

Example 23

[4360]




[4361] Coupling here was carried out using 100 mg of MF-Ta in PBS and following the Sephadex purification the reaction mixture was concentrated by ultracentrifugation.

[4362] Protein concentration: 11.2 mg/ml

[4363] Drug/mAb Ratio: 3.4

Example 24

[4364]




[4365] Protein concentration: 1.56

[4366] Drug/mAb Ratio: 2.8

Example 25

[4367]




[4368] Protein concentration: 0.60 mg/ml

[4369] Drug/mAb Ratio: 2.4

Example 26


[4370]



Drug/mAb-Ratio: 2.6

Example 27

[4372]




[4373] Protein concentration: 0.39 mg/ml

Drug/mAb-Ratio: 0.8

Example 28

[4374]




[4375] Coupling here was carried out using 70 mg of MF-Ta in PBS and following the Sephadex purification the reaction mixture was concentrated by ultracentrifugation.

[4376] Protein concentration: 13.2 mg/ml

[4377] Drug/mAb Ratio: 4.6

Example 29

[4378]




[4379] Protein concentration: 0.98 mg/ml

[4380] Drug/mAb Ratio: 1.1

Example 30

[4381]




[4382] Protein concentration: 0.55 mg/ml

[4383] Drug/mAb Ratio: not determinable

Example 31

[4384]




[4385] Coupling here was carried out using 40 mg of MF-Ta in PBS and following the Sephadex purification the reaction mixture was concentrated by ultracentrifugation.

[4386] Protein concentration: 10.6 mg/ml

[4387] Drug/mAb Ratio: 4.1

Example 32

[4388]




[4389] Protein concentration: 0.96 mg/ml

[4390] Drug/mAb Ratio: 0.4

Example 33

[4391]




[4392] Coupling here was carried out using 70 mg of MF-Ta in PBS and following the Sephadex purification the reaction mixture was concentrated by ultracentrifugation.

[4393] Protein concentration: 12.7 mg/ml

[4394] Drug/mAb Ratio: 3.6

Example 34

[4395]




[4396] Protein concentration: 1.1 mg/ml

[4397] Drug/mAb Ratio: 2.7

Example 35

[4398]




[4399] Protein concentration: 1.24 mg/ml

[4400] Drug/mAb Ratio: 2.6

Example 36

[4401]




[4402] Protein concentration: 0.99 mg/ml

[4403] Drug/mAb Ratio: 2.3

Example 37

[4404]




[4405] Protein concentration: 1.22 mg/ml

[4406] Drug/mAb Ratio: 3.3

Example 38

[4407]




[4408] Protein concentration: 1.34 mg/ml

[4409] Drug/mAb Ratio: 1.2

Example 39

[4410]




[4411] Protein concentration: 1.28 mg/ml

[4412] Drug/mAb Ratio: 3.2

Example 40

[4413]




[4414] Coupling here was carried out using 70 mg of MF-Ta in PBS and following the Sephadex purification the reaction mixture was concentrated by ultracentrifugation.

[4415] Protein concentration: 10.9 mg/ml

[4416] Drug/mAb Ratio: 5.1

Example 41

[4417]




[4418] Coupling here was carried out using 100 mg of MF-Ta in PBS and following the Sephadex purification the reaction mixture was concentrated by ultracentrifugation.

[4419] Protein concentration: 10.3 mg/ml

[4420] Drug/mAb Ratio: 4.3

Example 42

[4421]




[4422] Protein concentration: 1.08 mg/ml

[4423] Drug/mAb Ratio: 2.8

Example 43

[4424]




[4425] Protein concentration: 1.24 mg/ml

[4426] Drug/mAb Ratio: 2.8

Example 44

[4427]




[4428] Protein concentration: 1.28 mg/ml

[4429] Drug/mAb Ratio: 3.8

Example 45

[4430]




[4431] Protein concentration: 1.07 mg/ml

[4432] Drug/mAb Ratio: 3.0

Example 46

[4433]




[4434] Protein concentration: 1.35 mg/ml

[4435] Drug/mAb Ratio: 4.0

Example 47

[4436]




[4437] Coupling here was carried out using 100 mg of MF-Ta in PBS and following the Sephadex purification the reaction mixture was concentrated by ultracentrifugation.

[4438] Protein concentration: 12.2 mg/ml

[4439] Drug/mAb Ratio: 5.6

Example 48

[4440]




[4441] Protein concentration: 1.32 mg/ml

[4442] Drug/mAb Ratio: 3.2

Example 49

[4443]




[4444] Protein concentration: 1.01 mg/ml

[4445] Drug/mAb Ratio: 0.9

Example 50

[4446]




[4447] Protein concentration: 1.03 mg/ml

[4448] Drug/mAb Ratio: 0.3

Example 51

[4449]




[4450] Protein concentration: 0.62 mg/ml

[4451] Drug/mAb Ratio: 3.1

[4452] This ADC was concentrated by means of vivaspin centrifugation, re-diluted, and concentrated and re-diluted again.

Example 52

[4453]




[4454] Protein concentration: 1.26 mg/ml

[4455] Drug/mAb Ratio: not determinable


Example 53

[4456]




[4459] Drug/mAb Ratio: not determinable exactly

[4460]




[4463] Drug/mAb Ratio: 2.8

[4464]



Example 56

[4467]




[4468] Protein concentration: 0.92 mg/ml

[4469] Drug/mAb Ratio: 3.5

Example 57

[4470]




[4471] Coupling here was carried out using 5 mg of MF-Ta in PBS and following the Sephadex purification the batch was concentrated by ultracentrifugation and re-diluted.

[4472] Protein concentration: 0.77 mg/ml

[4473] Drug/mAb Ratio: >1.5 (not determinable exactly)

Example 58

[4474]




[4475] Protein concentration: 1.3 mg/ml

[4476] Drug/mAb Ratio: 2.0

Example 59

[4477]




[4478] Coupling here was carried out using 150 mg of MF-Ta in PBS and following the Sephadex purification the reaction mixture was concentrated by ultracentrifugation.

[4479] Protein concentration: 11.2 mg/ml

[4480] Drug/mAb Ratio: 3.7

Example 60

[4481]




[4482] Coupling here was carried out using 100 mg of MF-Ta in PBS and following the Sephadex purification the reaction mixture was concentrated by ultracentrifugation.

[4483] Protein concentration: 11.4 mg/ml

[4484] Drug/mAb Ratio: 3.9

Example 61

[4485]




[4486] Coupling here was carried out using 60 mg of MF-Ta in PBS and following the Sephadex purification the reaction mixture was concentrated by ultracentrifugation.

[4487] Protein concentration: 10.5 mg/ml

[4488] Drug/mAb Ratio: 4.4

Example 62

[4489] N-(4-{2-[6-(3-{[(2R)-2-Amino-2-carboxyethyl]sulphanyl}-2,5-dioxopyrrolidin-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide



[4490] 10 mg (10 μ mol) of Intermediate 157 were taken up in 5.2 ml of DMF and admixed with 2.28 mg (20 μ mol) of L-cysteine. The reaction mixture was stirred at RT for 2 hours, then concentrated under reduced pressure and subsequently purified by preparative HPLC. This gave 5.8 mg (48% of theory) of the title compound.


[4491] HPLC (Method 5): R_t =1.45 min;

[4492] LC-MS (Method 1): R_t =0.74 min; MS (ESIpos): m/z =1184 (M+H)⁺.

Example 63

N-(4-{2-[6-(3-{[(2R)-2-Amino-2-carboxyethyl]sulphanyl}-2,5-dioxopyrrolidin-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(1S)-1-carboxy-2-(1H-indol-3-yl)ethyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4493]



[4494] 10 mg (10 μ mol) of Intermediate 113 were taken up in 5.2 ml of DMF and admixed with 2.28 mg (20 μ mol) of L-cysteine. The reaction mixture was stirred at RT for 2 hours, then concentrated under reduced pressure and subsequently purified by preparative HPLC. This gave 6 mg (54% of theory) of the title compound.


[4495] HPLC (Method 5): R_t =1.5 min;

[4496] LC-MS (Method 1): R_t =0.77 min; MS (ESIpos): m/z =1185 (M+H)⁺.

Example 64

N-(4-{2-[6-(3-{[(2R)-2-Amino-2-carboxyethyl]sulphanyl}-2,5-dioxopyrrolidin-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4497]



[4498] 9 mg (8.3 μ mol) of Intermediate 132 were taken up in 4 ml of DMF and admixed with 3 mg (24.4 μ mol) of L-cysteine. The reaction mixture was stirred at RT overnight, then concentrated under reduced pressure and subsequently purified by preparative HPLC. This gave 6.8 mg (68% of theory) of the title compound.


[4499] HPLC (Method 12): R_t =1.8 min;

[4500] LC-MS (Method 1): R_t =0.78 min; MS (ESIpos): m/z =1227 (M+H)⁺.

Example 65

N-[6-(3-[[[(2R)-2-Amino-2-carboxyethyl]sulphonyl]-2,5-dioxopyrrolidin-1-yl]hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S)-1-amino-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4501]



[4502] 10 mg (10 μ mol) of Intermediate 106 were taken up in 5.8 ml of DMF and admixed with 2.5 mg (20 μ mol) of L-cysteine. The reaction mixture was stirred at RT for 2 hours, then concentrated under reduced pressure and subsequently purified by preparative HPLC. This gave 5.2 mg (46% of theory of the title compound.


[4503] HPLC (Method 5): R_f =1.5 min;

[4504] LC-MS (Method 11): R_f =0.71 min; MS (ESIpos): m/z =1070 (M+H)⁺.

Example 66

N-[6-(3-[[[(2R)-2-Amino-2-carboxyethyl]sulphonyl]-2,5-dioxopyrrolidin-1-yl]hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(1S)-1-carboxy-2-(1H-indol-3-yl)ethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4505]



[4506] 10 mg (10 μ mol) of Intermediate 124 were taken up in 4 ml of DMF and admixed with 2.5 mg (20 μ mol) of L-cysteine. The reaction mixture was stirred at RT for 2 hours, then concentrated under reduced pressure and subsequently purified by preparative HPLC. This gave 7.2 mg (64% of theory of the title compound.


[4507] HPLC (Method 5): R_f =1.6 min;

[4508] LC-MS (Method 1): R_f =0.8 min; MS (ESIpos): m/z =1071 (M+H)⁺.

Example 67

N-[6-(3-[[[(2R)-2-Amino-2-carboxyethyl]sulphonyl]-2,5-dioxopyrrolidin-1-yl]hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4509]



[4510] 10 mg (10 μ mol) of Intermediate 125 were taken up in 4 ml of DMF and admixed with 2.4 mg (20 μ mol) of L-cysteine. The reaction mixture was stirred at RT for 2 hours, then concentrated under reduced pressure and subsequently purified by preparative HPLC. This gave 7.7 mg (69% of theory of the title compound).


[4511] HPLC (Method 5): R_f =1.7 min;

[4512] LC-MS (Method 2): R_f =1.91 min; MS (ESIpos): m/z =1140 (M+H)⁺.

Example 68

N-(4-{2-[6-(3-[[[(2R)-2-Amino-2-carboxyethyl]sulphonyl]-2,5-dioxopyrrolidin-1-yl]hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S)-1-(benzylamino)-3-(1H-indol-3-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4513]



[4514] 10 mg (10 μ mol) of Intermediate 160 were taken up in 3 ml of DMF and admixed with 2.1 mg (20 μ mol) of L-cysteine. The reaction mixture was stirred at RT for 2 hours, then concentrated under reduced pressure and subsequently purified by preparative HPLC. This gave 8.1 mg (73% of theory of the title compound).


[4515] HPLC (Method 5): R_f =1.7 min;

[4516] LC-MS (Method 1): R_f =0.86 min; MS (ESIpos): m/z =1274 (M+H)⁺.

Example 69

N-(4-{2-[6-(3-{[(2R)-2-Amino-2-carboxyethyl]sulphanyl}-2,5-dioxopyrrolidin-1-yl)hexanoyl]hydrazino}-4-oxobutyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-1-(benzylamino)-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4517]



[4518] 3.5 mg (3 μ mol) of Intermediate 159 were taken up in 1 ml of DMF and admixed with 0.76 mg (6 μ mol) of L-cysteine. The reaction mixture was stirred at RT for 2 hours, then concentrated under reduced pressure and subsequently purified by preparative HPLC. This gave 2.6 mg (65% of theory of the title compound).


[4519] HPLC (Method 5): R_f =1.75 min;

[4520] LC-MS (Method 1): R_f =0.85 min; MS (ESIpos): m/z =1235 (M+H)⁺.

Example 70

N-(6-{2-[6-(3-{[(2R)-2-Amino-2-carboxyethyl]sulphanyl}-2,5-dioxopyrrolidin-1-yl)hexanoyl]hydrazino}-6-oxohexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-3-methoxy-1-{(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-{[(1S,2R)-1-(1,2-oxazinan-2-ylcarbonyl)-2-phenylcyclopropyl]amino}-3-oxopropyl]pyrrolidin-1-yl}-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4521]




[4522] 3.6 mg (3 μ mol) of Intermediate 129 were taken up in 1 ml of DMF and admixed with 0.77 mg (6 μ mol) of L-cysteine. The reaction mixture was stirred at RT for 2 hours, then concentrated under reduced pressure and subsequently purified by preparative HPLC. This gave 1.55 mg (39% of theory of the title compound).

[4523] HPLC (Method 5): R_f =1.6 min;

[4524] LC-MS (Method 1): R_f =0.87 min; MS (ESIpos): m/z =1255 (M+H)⁺.

Example 71

[4525]




[4526] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4527] Protein concentration: 0.9 mg/ml

[4528] Drug/mAb Ratio: 1

Example 72

[4529]




[4530] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4531] Protein concentration: 1.86 mg/ml

[4532] Drug/mAb Ratio: 2.9

Example 73

[4533]




[4534] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4535] Protein concentration: 1.05 mg/ml

[4536] Drug/mAb Ratio: 4.4

Example 74

[4537]




[4538] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4539] Protein concentration: 1.13 mg/ml

[4540] Drug/mAb Ratio: 2.8

Example 75

[4541]




[4542] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4543] Protein concentration: 1.41 mg/ml

[4544] Drug/mAb Ratio: 3.9

Example 76

[4545]




[4546] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4547] Protein concentration: 1.38 mg/ml

[4548] Drug/mAb Ratio: 4.3

Example 77

[4549]




[4550] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4551] Protein concentration: 1.32 mg/ml

[4552] Drug/mAb Ratio: 1

Example 78

[4553]




[4554] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4555] Protein concentration: 1.14 mg/ml

[4556] Drug/mAb Ratio: 5.3

Example 79

[4557]




[4558] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4559] Protein concentration: 1.25 mg/ml

[4560] Drug/mAb Ratio: 4.8

Example 80

[4561]




[4562] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4563] Protein concentration: 1.12 mg/ml

[4564] Drug/mAb Ratio: 1.7

Example 81

[4565]




[4566] Coupling here was carried out using 150 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation, re-diluted with PBS and concentrated again.

[4567] Protein concentration: 12.2 mg/ml

[4568] Drug/mAb Ratio: 4.1

Example 82

[4569]




[4570] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4571] Protein concentration: 0.86 mg/ml

[4572] Drug/mAb Ratio: 3.4

Example 83

[4573]




[4574] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4575] Protein concentration: 1.43 mg/ml

[4576] Drug/mAb Ratio: 3.7

Example 84

[4577]




[4578] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and rediluted.

[4579] Protein concentration: 0.8 mg/ml

[4580] Drug/mAb Ratio: 0.7

Example 85

[4581]




[4582] Coupling here was carried out using 50 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation, re-diluted with PBS and concentrated again.

[4583] Protein concentration: 9.5 mg/ml

[4584] Drug/mAb Ratio: 2.9


Example 86

[4585]



[4588] Drug/mAb Ratio: 3.2


[4589]



[4590] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4592] Drug/mAb Ratio: 4.6


[4593]



[4594] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4596] Drug/mAb Ratio: 1.6

[4597]




[4598] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4599] Protein concentration: 0.99 mg/ml

[4600] Drug/mAb Ratio: 5.5

Example 90

[4601]




[4602] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4603] Protein concentration: 1.02 mg/ml

[4604] Drug/mAb Ratio: 4.0

Example 91

[4605]




[4606] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4607] Protein concentration: 1.63 mg/ml

[4608] Drug/mAb Ratio: 3.8

Example 92

[4609]




[4610] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4611] Protein concentration: 1.27 mg/ml

[4612] Drug/mAb Ratio: 3.0

Example 93

[4613]




[4614] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4615] Protein concentration: 1.58 mg/ml

[4616] Drug/mAb-Ratio: 0.6

Example 94

[4617]




[4618] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4619] Protein concentration: 1.31 mg/ml

[4620] Drug/mAb Ratio: 6.6

Example 95

[4621]




[4622] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4623] Protein concentration: 1.75 mg/ml

[4624] Drug/mAb Ratio: 1.8

Example 96

[4625]




[4626] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4627] Protein concentration: 1.44 mg/ml

[4628] Drug/mAb Ratio: 2.5

Example 97

[4629]




[4630] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4631] Protein concentration: 1.96 mg/ml

[4632] Drug/mAb Ratio: 5.6

Example 98

[4633]




[4634] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4635] Protein concentration: 1.58 mg/ml

[4636] Drug/mAb Ratio: 4.2

Example 99

[4637]




[4638] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4639] Protein concentration: 1.48 mg/ml

[4640] Drug/mAb Ratio: 4.6

Example 100

[4641]




[4642] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4643] Protein concentration: 1.5 mg/ml

[4644] Drug/mAb Ratio: 3.1

Example 101

[4645]




[4646] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4647] Protein concentration: 1.3 mg/ml

[4648] Drug/mAb Ratio: 4.3

Example 102

[4649]




[4650] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4651] Protein concentration: 1.62 mg/ml


[4652] Drug/mAb Ratio: 2.2

Example 103

[4653]



-continued




[4654] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4655] Protein concentration: 1.37 mg/ml

[4656] Drug/mAb Ratio: 2.8

Example 104

[4657]




[4658] Coupling here was carried out using 5 mg of MF-Ta in PBS and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4659] Protein concentration: 1.43 mg/ml

[4660] Drug/mAb Ratio: 4.0

Example 105

[4661] N-(6-{[(5S)-5-Amino-5-carboxypentyl]amino}-6-oxohexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate



[4662] 15.5 mg (15 μ mol) of Intermediate 210 were taken up in 5 ml of DMF and admixed with 4.4 mg (18 μ mol) of N²-(tert-butoxycarbonyl)-L-lysine and also 7.7 μ L (44 μ mol) of N,N-diisopropylethylamine. The reaction mixture was stirred at RT overnight and then concentrated under reduced pressure. The residue was subsequently purified by preparative HPLC. This gave 14 mg (81% of theory) of the protected intermediate of the title compound, which was subsequently taken up in 1 ml of dichloromethane and deprotected with 1 ml of trifluoroacetic acid. The reaction mixture was concentrated and, following lyophilization of the residue from acetonitrile/water 1:1, 15 mg (97% of theory) of the title compound were obtained.


[4663] HPLC (Method 12): R_t =1.8 min;

[4664] LC-MS (Method 1): R_t =0.79 min; MS (ESIpos): m/z =1083 (M+H)⁺.

Example 106

N-(6-{[(5S)-5-Amino-5-carboxypentyl]amino}-6-oxohexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(1S)-1-carboxy-2-(1H-indol-3-yl)ethyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide

[4665]



[4666] 40 mg (40 μ mol) of Intermediate 227 were taken up in 5 ml of DMF and admixed with 11.5 mg (40 μ mol) of N²-(benzyloxycarbonyl)-L-lysine and also 13 μ L (80 μ mol) of N,N-diisopropylethylamine. The reaction mixture was stirred at RT overnight, then concentrated under reduced pressure and subsequently purified by preparative HPLC. This gave 32.5 mg (70% of theory) of the protected intermediate of the title compound.

[4667] This 32.5 mg of the intermediate were dissolved in 10 ml of methanol and, following addition of 2 mg of 10% palladium on activated carbon, were hydrogenated under standard hydrogen pressure at RT for 30 minutes. The catalyst was then removed by filtration and the solvent was removed under reduced pressure. Lyophilization of the residue from dioxane/water 1:1 gave 26 mg (99% of theory) of the title compound.


[4668] HPLC (Method 12): R_t =1.7 min;

[4669] LC-MS (Method 1): R_t =0.76 min; MS (ESIpos): m/z =1014 (M+H)⁺.

Example 107

N-[(18S)-18-Amino-18-carboxy-12-oxo-3,6,9-trioxo-13-azaoctadec-1-yl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[4670]



[4671] 3.5 mg (3 μ mol) of Intermediate 202 were taken up in 2 ml of DMF and admixed with 0.8 mg (3 μ mol) of N²-(tert-butoxycarbonyl)-L-lysine and also 1.6 μ L (10 μ mol) of N,N-diisopropylethylamine. The reaction mixture was stirred at RT overnight and then concentrated under reduced pressure. The residue was taken up in acetonitrile/water: 1:1, brought to a pH of 2 with trifluoroacetic acid and then purified by preparative HPLC. This gave 1 mg (25% of theory) of the protected intermediate of the title compound, which was subsequently taken up in 500 μ L of dichloromethane and deprotected with 500 μ L of trifluoroacetic acid. The reaction mix-


ture was concentrated and, following lyophilization of the residue from acetonitrile/water 1:1, 1 mg (89% of theory) of the title compound was obtained.

[4672] HPLC (Method 12): R_f =1.9 min;

[4673] LC-MS (Method 1): R_f =0.82 min; MS (ESIpos): m/z =1173 (M+H)⁺.

Example 108

[4674]




[4675] Coupling here was carried out using 5 mg of MF-Ta in PBS, and the batch, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4676] Protein concentration: 0.81 mg/ml

[4677] Drug/mAb Ratio: 2.5

Example 109

[4678]




[4679] Coupling here was carried out using 5 mg of MF-Ta and the batch, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4680] Protein concentration: 1.06 mg/ml

[4681] Drug/mAb Ratio: 1.8

Example 110

[4682]




[4683] Coupling here was carried out using 5 mg of MF-Ta and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted with PBS.

[4684] Protein concentration: 1.36 mg/ml

[4685] Drug/mAb Ratio: 7.2

Example 111

[4686]




[4687] Coupling here was carried out using 5 mg of MF-Ta and the reaction mixture, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted with PBS.

[4688] Protein concentration: 1.57 mg/ml

[4689] Drug/mAb Ratio: 2.9

Example 112

[4690]




[4691] Coupling here was carried out using 5 mg of MF-Ta in PBS, and the batch, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.

[4692] Protein concentration: 1.44 mg/ml

[4693] Drug/mAb Ratio: 2.5

Example 113

[4694]



[4695] Coupling here was carried out using 5 mg of MF-Ta in PBS, and the batch, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted.


[4696] Protein concentration: 1.74 mg/ml

[4697] Drug/mAb Ratio: 3.6

Example 114

Diastereomer 1

[4698]



[4699] Coupling here was carried out using Intermediate 247a and 5 mg of MF-Ta in PBS and the batch, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted with PBS.


[4700] Protein concentration: 1.57 mg/ml

[4701] Drug/mAb Ratio: 4.2

Example 115

Diastereomer 2

[4702]




[4703] Coupling here was carried out using Intermediate 247b and 5 mg of MF-Ta in PBS and the batch, following Sephadex purification, was concentrated by ultracentrifugation and re-diluted with PBS.

[4704] Protein concentration: 1.42 mg/ml

[4705] Drug/mAb Ratio: 4.0

Example 116

[4706] N-(6-[(5S)-5-Amino-5-carboxypentyl]amino)-6-oxohexyl)-N-methyl-L-threonyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate



[4707] 8.6 mg (8 μ mol) of Intermediate 240 were taken up in 5 ml of DMF and admixed with 4.0 mg (16 μ mol) of N²-(tert-butoxycarbonyl)-L-lysine and also 2 μ L (16 μ mol) of N,N-diisopropylethylamine. The reaction mixture was stirred at RT for 4 hours, then admixed again with the same amounts of N²-(tert-butoxycarbonyl)-L-lysine and N,N-diisopropylethylamine, and stirred at RT overnight. The batch was subsequently concentrated under reduced pressure. The residue was then purified by preparative HPLC. This gave 7 mg (72%

of theory) of the protected intermediate of the title compound, which was subsequently taken up in 1 ml of dichloromethane and deprotected with 0.5 ml of trifluoroacetic acid. The batch was concentrated and the residue was purified by preparative HPLC. Drying under a high vacuum gave 3.3 mg (47% of theory) of the title compound.


[4708] HPLC (Method 5): R_t=1.5 min;

[4709] LC-MS (Method 1): R_t=0.8 min; MS (ESIpos): m/z=1084 (M+H)⁺.

Example 117

N-6-{[(5S)-5-Amino-5-carboxypentyl]amino}-6-oxohexyl-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(2S)-3-(4-hydroxyphenyl)-1-(1,2-oxazin-2-yl)-1-oxopropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[4710]



[4711] 8 mg (8 μ mol) of Intermediate 242 were taken up in 3 ml of DMF and admixed with 2.9 mg (12 μ mol) of N²-(tert-butoxycarbonyl)-L-lysine and also 2.7 μ L (16 μ mol) of N,N-diisopropylethylamine. The reaction mixture was stirred at RT overnight, then admixed again with the same amounts of N²-(tert-butoxycarbonyl)-L-lysine and N,N-diisopropylethylamine, and stirred at RT for a further 4 hours. The batch was subsequently concentrated under reduced pressure. The residue was then purified by preparative HPLC. Lyophilization from acetonitrile/water gave 6.5 mg (72% of theory) of the protected intermediate of the title compound, which was subsequently taken up in 5 ml of dichloromethane and deprotected with 0.75 ml of trifluoroacetic acid. The batch was concentrated, and lyophilization of the residue from dioxane/water gave 5 mg (76% of theory) of the title compound.


[4712] HPLC (Method 12): R_t =1.7 min;

[4713] LC-MS (Method 1): $R_t=0.69$ min; MS (ESIpos): $m/z=1059$ (M+H)⁺.

Example 118

N-(6-{[(5S)-5-Amino-5-carboxypentyl]amino}-6-oxohexyl)-N-methyl-L-valyl-N-[(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-[[(1S)-1-carboxy-2-(4-hydroxyphenyl)ethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate

[4714]



[4715] 38 mg (41 μ mol) of Intermediate 248 were first converted into the N-hydroxysuccinimide ester. 72 mg of the crude product obtained were taken up in 5 ml of DMF and admixed with 24 mg (100 μ mol) of N²-(tert-butoxycarbonyl)-L-lysine and 23 μ L of N,N-diisopropylethylamine. The reaction mixture was stirred at RT overnight, and then admixed again with 16 mg of N²-(tert-butoxycarbonyl)-L-lysine and 12 μ L of N,N-diisopropylethylamine, and subsequently treated in an ultrasound bath for a further 2 hours. The batch was then concentrated under reduced pressure and the residue was purified by preparative HPLC. Lyophilization from acetonitrile/water gave 20 mg (50% of theory) of the protected intermediate of the title compound.

[4716] 15 mg (12 μ mol) of this intermediate were subsequently taken up in 3 ml of dichloromethane and admixed with 1 ml of trifluoroacetic acid. After 40 minutes of stirring at RT, a further 1.5 ml of trifluoroacetic acid were added and the batch was treated in an ultrasound bath for 1 hour. Thereafter the batch was concentrated, and lyophilization of the residue from dioxane/water gave 13 mg (90% of theory) of the title compound.

[4717] HPLC (Method 12): R_f=1.5 min;

[4718] LC-MS (Method 1): R_f=0.68 min; MS (ESIpos): m/z=990 (M+H)⁺.

C: EVALUATION OF BIOLOGICAL ACTIVITY

[4719] The biological effect of the compounds of the invention can be demonstrated by the assays described below

C-1. In Vitro Cell Proliferation Tests

[4720] The cytotoxic activity of the conjugates of the invention was tested in an in vitro cell proliferation test. For this purpose, a mammalian cell which expresses either the target molecule of the binder endogenously or recombinantly was incubated with the conjugate of the invention. Cell proliferation was determined after an incubation time of several hours to several days, on the basis of the cell count in comparison to a control to which no conjugate was added. As further controls it is possible to add the unconjugated toxophore alone, or to use cells which do not express the target molecule of the binder. The cell count was determined by methods known to the skilled person, as for example by counting or through the use of an assay kit which allows the cell count to be determined via an ATP measurement (e.g. ATPlite™, Perkin-Elmer). In this way the IC₅₀ of the conjugates of the invention was ascertained. By comparing the IC₅₀ of the conjugate in measurements on cells which carry the target molecule of the binder, and non-carrying cells, it was possible to determine the selectivity of the conjugate.

C-2. Determination of the Antiproliferative Activity of Anti-Mesothelin ADC in the Human Colonic Carcinoma Cell Line HT29

[4721] A defined number of cells from the human colonic carcinoma cell line HT29 wt (2500 c/well, wildtype) was seeded out into a 96-well MTP in complete medium (10% FCS-RPMI) and incubated overnight at 37° C./5% carbon dioxide. In parallel to this, transfected HT29 cells which stably express mesothelin were plated out in complete medium in a 96-well MTP and incubated overnight (2500c/well; 37° C./5% carbon dioxide).

[4722] After 18 hours, the seeding medium was replaced by fresh medium with 10% FCS. Treatment commenced with the

addition of the compounds of the invention. The transfected cells and the HT29 wt cells were treated identically here.

[4723] For the substances under investigation, dose-effect curves were determined in a concentration range of 10⁻⁵ M-10⁻¹⁴ M (1:10 dilution series).

[4724] Incubation times of 48 h-96 h were selected.

[4725] Proliferation was detected using the MTT assay (ATCC, Manassas, Va., USA, Catalogue No. 30-1010K). At the end of the selected incubation time, the HT29 wt cells were incubated with MTT for 4 h, followed by lysis of the cells overnight by addition of the detergent.

[4726] The dye formed was detected at 570 nm.

[4727] The proliferation of cells which were not treated with test substance but were otherwise identically treated was defined as the 100% figure. The data obtained from this test represents triplicate determinations, and at least two independent experiments were carried out.

[4728] Table 3 lists the IC₅₀ values¹⁾ of representative working examples from this assay:

¹⁾ The activity data reported relate to the working examples described in the present experimental section, with the drug/mAB ratios indicated. The values may possibly deviate for different drug/mAB ratios.

TABLE 3

Example	HT29 meso ⁺ IC ₅₀ [nM]	HT29 wt IC ₅₀ [nM]
1	0.17	5.3
2	0.4	162
3	5	22
4	1	13
5	0.06	8
5	0.16	5.4
6	0.019	0.27
7	20	>300
8	0.03	1
9	0.11	1.4
10	0.14	3.3
11	1.5	8.6
12	0.015	11
13	2	9
14	0.07	20
15	0.02	2
16	0.6	11
17	0.05	8
18	25	221
19	9	>50
20	0.2	0.29
21	0.35	3
22	0.8	3.6
23	1.1	6
24	0.5	5
25	4.4	14
26	0.02	1.4
27	24	>1000
28	0.3	31
29	6	14
30	0.2	2
31	0.45	>500
32	0.9	3
33	0.01	0.2
34	0.06	3.1
35	2.6	60
36	0.35	1.1
37	38	167
38	52	89
39	0.6	>500
40	0.013	4.9
41	25	>500
42	0.2	0.2
43	1.7	6.3
44	0.27	1.6
45	2	44

TABLE 3-continued

Example	HT29 meso ⁺ IC ₅₀ [nM]	HT29 wt IC ₅₀ [nM]
46	3	500
47	7.2	294
48	10	32
49	1.8	0.9
50	0.6	1.4
51	0.8	15
52	3	2
53	0.74	9.2
54	0.09	1.1
55	0.002	0.28
56	0.7	162
57	0.018	4
58	0.4	2.5
59	0.6	4.1
60	0.06	10
61	9.2	46
71	22	90
72	1.5	>1000
73	2	227
74	5.3	57
75	0.13	672
76	0.09	769
77	3.3	>1000
78	2.1	219
79	0.03	199
80	0.25	317
81	1.6	3000
82	0.04	10
83	0.3	>1000
84	70	>1000
85	6.8	>1000
86	43	>1000
87	0.27	554
88	0.3	54
89	2.7	74
90	7.4	135
91	13	>1000
92	5.6	18
93	425	>1000
94	3.1	33
95	<0.1	0.15
96	>1000	>1000
97	4.1	>1000
98	30	>1000
99	2.2	>1000
100	0.5	>500
101	4.6	>500
102	0.09	839
103	0.18	602
104	177	>500
108	1.3	16
109	0.06	10
110	3.1	>500
111	0.02	14
112	1.6	>500
113	0.04	>500
114	7	>500
115	15	>500

C-3. Determination of the Effect on Tubulin Polymerization

[4729] Cancer cells are denatured cells which frequently lead to the formation of tumours also as a result of increased cell division. Microtubuli form the spindle fibres of the spindle apparatus and are an essential constituent of the cell cycle. The regulated construction and breakdown of microtubuli allows the precise division of the chromosomes among the daughter cells, and constitutes a continuously dynamic process. Disruption to this dynamic process results in incorrect cell division and ultimately in cell death. The increased cell division of cancer cells, however, also makes them par-

ticularly sensitive towards spindle fibre poisons, which constitute a fixed constituent of chemotherapy. Spindle fibre poisons such as paclitaxel or epothilone lead to a sharply increased polymerization rate of the microtubuli, while vinca alkaloids or else monomethylauristatin E (MMAE) lead to a sharply reduced polymerization rate of the microtubuli. In both cases, the necessary dynamism of the cell cycle is critically disrupted. The compounds investigated in the context of the present invention result in a reduced polymerization rate of the microtubuli.

[4730] Tubulin polymerization was investigated using the "Fluorescence-based Microtubule Polymerisation Assay Kit" from Cytoskeleton (Denver, Colo., USA; order number: BK011). With this assay, GTP is added to unpolymerized tubulin, allowing polymerization to take place spontaneously. The assay is based on the binding of the fluorophore 4',6-diamidino-2-phenylindole (DAPI) to tubulin. Free and bound DAPI can be differentiated on the basis of different emission spectra. Since DAPI exhibits a significantly high affinity for polymerized tubulin in comparison to non-polymerized tubulin, the tubulin polymerization can be followed via the increase in the fluorescence of bound DAPI fluorophores.

[4731] For the implementation of this assay, the compounds of the invention, in solution in DMSO, were diluted from their initial concentration of 10 mM to 1 μ M in water. In addition to the buffer control, paclitaxel, with a polymerization-increasing effect, and vinblastin, with a polymerization-inhibiting effect, were run additionally as assay controls. Measurement was carried out using 96-well plates with a half base area. The kinetics of the tubulin polymerization were monitored in a Fluorimeter at 37° C. for 1 hour. The excitation wavelength was 355 nm, and emission was monitored at 460 nm. For the region of linear increase within the first 10 minutes, a calculation was made of the change in fluorescence per minute ($\Delta F/\text{min}$), which represents the polymerization rate of the microtubuli. The potency of the test substances was quantified on the basis of their respective reduction of the polymerization rate.

[4732] The reduction in the tubulin polymerization rate occurs in dependence on the concentration of the test substance added. Whereas the addition of 100 μ M MMAF causes complete cessation of polymerization, the addition of 1 μ M MMAF leads to a reduction in the polymerization rate of 40-45%, where a comparison of the activity of different substances is shown at best. The compound from Example 64 inhibited the polymerization rate to an extent of 41% at 1 μ M. The compounds from Examples 62-70 and 105-107 exhibited a level of tubulin polymerization inhibition comparable with MMAF.

[4733] The value for the inhibition of MMAF at a concentration of 1 μ M is set as being 100%.

[4734] Table 6 gives data for the inhibition of tubulin polymerization by representative working examples.

TABLE 6

Working example	Concentration of toxophore [μ M]	Tubulin polymerization in the presence of toxophore in [%]. Tubulin polymerization rate at 1 μ M MMAF set at 100%
MMAF	1	100
MMAF	10	34
MMAF	100	0

TABLE 6-continued

Working example	Concentration of toxophore [μ M]	Tubulin polymerization in the presence of toxophore in [%]. Tubulin polymerization rate at 1 μ M MMAF set at 100%
105	1	45
105	10	1
106	1	80
106	10	14
107	1	60
107	10	0
62	1	88
62	10	25
63	1	109
63	10	27
64	1	120
65	1	117
65	10	64
66	1	107
66	10	25
67	1	121
67	10	35
68	1	111
68	10	45
69	1	110
107	1	78
107	10	24
116	1	102
116	10	31
117	1	88
117	10	21
118	1	90
118	10	17

[4735] The MMAF toxophore and the working examples inhibit tubulin polymerization as a function of their concentration. At 100 μ M MMAF, the tubulin polymerization is inhibited completely. Working Example 105 inhibits the tubulin polymerization rate at 1 μ M to 45% of the value measured for 1 μ M MMAF.

C-4. In Vitro Tests for Determining Cell Permeability

[4736] The cell permeability of a substance can be investigated by means of in vitro testing in a flux assay using Caco-2 cells [M. D. Troutman and D. R. Thakker, *Pharm. Res.* 20 (8), 1210-1224 (2003)]. For this purpose, the cells were cultured for 15-16 days on 24-well filter plates. For the determination of permeation, the respective working example was applied in a HEPES buffer to the cells either apically (A) or basally (B) and incubated for 2 hours. After 0 hours and after 2 hours, samples were taken from the cis and trans compartments. The samples were separated by HPLC (Agilent 1200, Böblingen, Germany) using reverse phase columns. The HPLC system was coupled via a Turbo Ion Spray Interface to a Triple Quadropole mass spectrometer API 4000 (Applied Biosystems Applera, Darmstadt, Germany). The permeability was evaluated on the basis of a P_{app} value, which was calculated using the formula published by Schwab et al. [D. Schwab et al., *J. Med. Chem.* 46, 1716-1725 (2003)].

[4737] Of critical importance for toxophores which are released intracellularly is the permeability from B to A [P_{app} (B-A)]: the lower this permeability, the longer the residence time of the working example in the cell following intracellular release, and hence also the longer the time available for interaction with the biochemical target (in this case: tubulin).

[4738] Table 4 below sets out permeability data for representative working examples from this assay:

TABLE 4

Working example	P_{app} (B-A) [μ m/s]
62	2
63	1.6
64	2.5
65	5
66	1
68	7
105	2
106	1
116	1.8
117	1.5

[4739] The working examples exhibit a low permeability from B to A [P_{app} (B-A)] and therefore have a long residence time in the CaCo-2 cells. In comparison, monomethylauristatin E (MMAE) and monomethylauristatin F (MMAF) in this test exhibit a P_{app} (B-A) value of 73 nm/s, and therefore have a significantly shorter residence time in the Caco-2 cells.

C-5. In Vitro Tests for Determining the Substrate Properties for P-Glycoprotein (P-gp)

[4740] Many tumour cells express transporter proteins for drugs, and this frequently accompanies the development of resistance towards cytostatics. Substances which are not substrates of such transporter proteins, such as P-glycoprotein (P-gp) or BCRP, for example, could therefore exhibit an improved activity profile.

[4741] The substrate properties of a substance for P-gp (ABCB1) were determined by means of a flux assay using LLC-PK1 cells which overexpress P-gp (L-MDR1 cells) [A. H. Schinkel et al., *J. Clin. Invest.* 96, 1698-1705 (1995)]. For this purpose, the LLC-PK1 cells or L-MDR1 cells were cultured on 96-well filter plates for 3-4 days. For determination of the permeation, the respective test substance, alone or in the presence of an inhibitor (such as Ivermectin or Verapamil, for example), was applied in a HEPES buffer to the cells either apically (A) or basally (B) and incubated for 2 hours. After 0 hours and after 2 hours, samples were taken from the cis and trans compartments. The samples were separated by HPLC using reverse phase columns. The HPLC system was coupled via a Turbo Ion Spray Interface to a Triple Quadropole mass spectrometer API 3000 (Applied Biosystems Applera, Darmstadt, Germany). The permeability was evaluated on the basis of a P_{app} value which was calculated using the formula published by Schwab et al. [D. Schwab et al., *J. Med. Chem.* 46, 1716-1725 (2003)].

[4742] Of critical importance for toxophores which are released intracellularly is the permeability from B to A [P_{app} (B-A)]: the lower this permeability, the longer the residence time of the working example in the cell following intracellular release, and hence also the longer the time available for interaction with the biochemical target (in this case: tubulin).

[4743] Table 5 below lists permeability data for representative working examples from this assay, which was carried out in L-MDR1 cells:

TABLE 5

Working example	P_{app} (B-A) [μ m/s]
62	3
63	3.6

TABLE 5-continued

Working example	P_{app} (B-A) [nm/s]
64	2.1
65	3.6
66	4
68	2
105	6
106	4

[4744] The working examples exhibit a low permeability from B to A [P_{app} (B-A)] and therefore have a long residence time in the L-MDR1 cells.

C-6. Pharmacokinetics in the HT29 Tumour Model with Mesothelin-Transfected and Non-Transfected HT29 Cells

[4745] Following intravenous administration of 16 mg/kg of Example 5, the plasma concentrations and tumour concentrations of Example 5 and also of potential metabolites such as, for example, of Example 64 were measured. In the animals with mesothelin-transfected tumours, the area under the curve (AUC) for the compound from Example 64 in plasma was about 0.50 mg*h/L; in the tumour, the exposure of the compound from Example 64 was about 400 times higher (AUC=203 mg*h/L). In the animals with nontransfected tumour animals, the exposure of Example 64 in plasma was identical to the exposure in plasma in animals with transfected tumours. In contrast, the AUC in the tumour in the nontransfected animals was about 8 times lower than in the transfected animals. This shows a distinct targeting effect in the tumour in the presence of the antigen.

Analysis for Quantifying the Compound from Example 64

[4746] The measurement of the compound from Example 64 in plasma and tumour took place following precipitation of the proteins with methanol, by means of high-pressure liquid chromatography (HPLC) coupled to a tandem mass spectrometer (MS).

[4747] For the processing of 100 μ L of plasma, it was admixed with 400 μ L of methanol and 10 μ L of internal standard (ISTD, 50 ng/mL in methanol) and shaken for 10 seconds. After centrifuging for 5 minutes at 16 000 g, 250 μ L of supernatant were transferred to an autosampler vial, which was made up with 250 μ L of ammonium acetate buffer (AAC, 10 mM, pH 6.8) and shaken again.

[4748] For the processing of a tumour, it was admixed with 4 times the amount of methanol. In a TissueLyser II (Qiagen), the sample was comminuted at 30 impacts per minute for 6 minutes and then centrifuged off at 16 000 g for 5 minutes. 50 μ L of the supernatant were transferred to an autosampler vial and made up with 50 μ L of ammonium acetate buffer (10 mM, pH 6.8) and with 5 μ L of ISTD. After again being shaken, the tumour sample was ready for measurement.

[4749] The measurement of both matrix samples took place, lastly, on the HPLC-coupled, atmospheric pressure ionization/tandem mass spectrometer by means of a Turbo Ion Spray Interface (TISP) on an API4000 instrument from SCIEX. The m/z transitions measured were as follows:

Example 64 (quantifier) 614.652→570.9

Example 64 (qualifier 1) 614.652→555.0

Example 64 (qualifier 2) 614.652→500.4

Internal standard (ISTD) 726.665→694.5

[4750] HPLC/LC-MSMS (TISP) analysis ran on an HP1100 pump (Agilent) with a Gemini column (5 μ m C18 110 A, 50×3 mm, Phenomenex). Under the following gradi-

ent conditions: flow rate 0.4 mL/min; gradient: 0.0 min-1.0 min 10% acetonitrile/90% AAC, 1.0 min-3.0 min 10% acetonitrile/90% AAC→50% acetonitrile/50% AAC, 3.0 min-5.5 min 50% acetonitrile/50% AAC, 5.5 min-5.6 min 50% acetonitrile/50% AAC→10% acetonitrile/90% AAC, 5.6 min-6.0 min 10% acetonitrile/90% AAC.

[4751] For calibration, plasma samples were admixed with concentrations of 0.5-2000 μ g/L. The detection limit (LOQ) was 2 μ g/L. The linear range extended from 2 to 1000 μ g/L.

[4752] For the calibration of the tumour samples, the supernatant of untreated tumours was admixed with concentrations of 0.5-200 μ g/L. The detection limit was 5 μ g/L. The linear range extended from 5 to 200 μ g/L.

[4753] Quality controls for validity testing contained 5 and 50 μ g/L, with an additional 500 μ g/L in plasma. The concentrations found for these samples deviated by up to 20% from the intended value (data not attached).

C-7. Activity Test In Vivo

[4754] The activity of the conjugates of the invention was tested in vivo by means for example of xenograft models. The skilled person knows of methods in the prior art for testing the activity of the compounds of the invention (see, for example, WO 2005/081711; Polson et al., Cancer Res. 2009 Mar. 15; 69(6):2358-64). For this purpose, for example, rodents (e.g. mice) were implanted with a tumour cell line which expresses the target molecule of the binder. The implanted animals were subsequently administered either a conjugate of the invention or a control antibody, or isotonic salt solution. Administration took place singly or more often. Tumour size was determined after an incubation time of several days, in a comparison of conjugate-treated animals and the control group. The conjugate-treated animals showed a lower tumour size.

C-7a. Testing of Anti-Mesothelin ADCs in Experimental Tumours in the Mouse

[4755] Human tumour cells which express mesothelin are injected subcutaneously into the flank of immunosuppressed mice, such as nude mice or SCID mice. 1-10 million cells are detached from the cell culture, centrifuged and resuspended with 100 μ L of medium or 1:1 medium/Matrigel. The cell suspension is injected beneath the skin of the mouse.

[4756] Within a few days, a tumour grows. Treatment begins no earlier than at a tumour size of 20 mm². In order to investigate the effect on tumours that are already established, the treatment may also be commenced only at a tumour size of 50-100 mm².

[4757] Treatment with ADCs takes place via the intravenous route into the caudal vein of the mouse. The ADC is dissolved in PBS and is administered with a volume of 5 mL/kg.

[4758] The treatment scheme is governed by the pharmacokinetics of the antibody. As a standard, treatment takes place three times following every fourth day. Treatment, however, may also be continued further, or a second cycle with three days of treatment may follow at a later point in time.

[4759] As a standard basis, 8 animals are used per treatment group. This number may be higher if particularly strong fluctuations in tumour growth or after treatment are anticipated. As well as the groups which receive the active substances, one group, as a control group, is treated only with the buffer, in accordance with the same scheme.

[4760] In the course of the experiment, the area of the tumour is measured regularly using a sliding caliper in two dimensions (length/width).

[4761] At the end of the experiment, the tumours are removed and weighed. The ratio of the average tumour weights for the therapy group (T) to the control group (C) is expressed as T/C.

C-7b. Activity in the HT29 Colonic Carcinoma Tumour Model with Mesothelin-Transfected HT29 Cells

[4762] One million HT29 cells (stably transfected with mesothelin) are inoculated subcutaneously into the flank of NMRI nude mice. Treatment begins intravenously with an average tumour size of 30 mm² on day 6 (day 6, 10, 14). Following the treatment, the tumour growth is monitored up to day 26. For reasons of animal protection law, the animals of the control group must be properly sacrificed on day 26, owing to the large tumours.

[4763] The ADCs bring about a distinct inhibition of tumour growth. On day 26, the experiment is ended and the tumours of all the animals are weighed. From this, T/C is calculated, as the tumour weight of the treatment group in relation to the control group.

TABLE 7

ADC	Dose	T/C
Vehicle	—	1
Example 60	5 mg/kg	0.32
Example 81	5 mg/kg	0.33

C-7c. Activity in the Ovar3 Ovarian Carcinoma Tumour Model

[4764] Three million Ovar3 cells are inoculated subcutaneously into the flank of NMRI nude mice.

[4765] Treatment begins intravenously at a dose of 5 mg/kg with an average tumour size of 40 mm² on day 49 (day 38, 42, 46 and day 70, 74). Following the treatment, the tumour growth is monitored up to day 78. At the end of the experiment, the tumours are removed and weighed.

[4766] The ADCs bring about a marked inhibition of tumour growth. On day 78, the experiment is ended and the tumours of all of the animals are weighed. From this, T/C is calculated, as the tumour weight of the treatment group in relation to the control group.

TABLE 8

ADC	Dose	T/C
Vehicle	—	1
Example 60	5 mg/kg	0.3
Example 81	5 mg/kg	0.23
Example 60	10 mg/kg	0.32
Example 81	10 mg/kg	0.16

C-7D. Activity in the NCI-H322 Pulmonary Carcinoma Tumour Model, which Expresses Mesothelin Endogenously. Treatment Scheme q4dx3.

[4767] Five million NCI-H322 cells are inoculated subcutaneously into the flank of NMRI nude mice. Treatment begins intravenously with a tumour size of about 50 mm² on day 21 (day 21, 25, 29). Following the treatment, the tumour growth is monitored up to day 42. For reasons of animal protection law, the animals of the control group must be properly sacrificed on day 42, owing to the large tumours.

[4768] The ADCs bring about a distinct inhibition of tumour growth. On day 42, the experiment is ended and the

tumours of all the animals are weighed. From this, T/C is calculated, as the tumour weight of the treatment group in relation to the control group.

TABLE 9

ADC	Dose	T/C
Vehicle	—	1
Example 60	5 mg/kg	0.31
Example 60	15 mg/kg	0.09
Example 81	5 mg/kg	0.09

C-7e. Activity in the NCI-H322 Pulmonary Carcinoma Tumour Model which Expresses Mesothelin Endogenously. Treatment Scheme: Single Administration.

[4769] Five million NCI-H322 cells are inoculated subcutaneously into the flank of NMRI nude mice. Treatment with the ADCs or vehicle takes place intravenously once at a tumour size of about 45 mm² on day 22. Following the treatment, the tumour growth is monitored. For reasons of animal protection law, the animals of the control group must be properly sacrificed on day 50, owing to the large tumours.

[4770] The ADCs bring about a marked inhibition of tumour growth. The tumour areas (length×width) on day 50 are used in order to determine the T/C.

TABLE 10

ADC	Dose	T/C
Vehicle	—	1
Example 60	5 mg/kg 1x	0.73
Example 81	5 mg/kg 1x	0.33
Example 60	15 mg/kg 1x	0.32
Example 81	15 mg/kg 1x	0.23

D. WORKING EXAMPLES FOR PHARMACEUTICAL COMPOSITIONS

[4771] The compounds of the invention can be converted as follows into pharmaceutical preparations:

i.v. Solution:

[4772] The compound of the invention is dissolved at a concentration below the saturation solubility in a physiologically tolerated solvent (e.g. isotonic saline solution, D-PBS, or a formulation with glycine and sodium chloride in citrate buffer with addition of polysorbate 80). The solution is subjected to sterile filtration and dispensed into sterile and pyrogen-free injection containers.

i.v. Solution:

[4773] The compounds of the invention can be converted into the administration forms cited. This can be accomplished in a known way by “mixing with” or “dissolving in” inert, non-toxic, pharmaceutically suitable excipients (e.g. buffer substances, stabilizers, solubilizers, preservatives). The following, for example, may be present: amino acids (glycine, histidine, methionine, arginine, lysine, leucine, isoleucine, threonine, glutamic acid, phenylalanine and others), sugars and related compounds (glucose, saccharose, mannitol, trehalose, sucrose, mannose, lactose, sorbitol), glycerol, sodium salts, potassium, ammonium salts and calcium salts (e.g. sodium chloride, potassium chloride or disodiumhydrogenphosphate and many others), acetate/acetic acid buffer sys-

tems, phosphate buffer systems, citric acid and citrate buffer systems, trometamol (TRIS and TRIS salts), Polysorbates (e.g. Polysorbate 80 and Polysorbate 20), Poloxamers (e.g. Poloxamer 188 and Poloxamer 171), Macrogols (PEG derivatives, e.g. 3350), Triton X-100, EDTA salts, glutathione, albumins (e.g. human), urea, benzyl alcohol, phenol, chlorocresol, metacresol, benzalkonium chloride and many others.

Lyophilizate for Subsequent Conversion into an i.v., s.c. or i.m. Solution:

[4774] Alternatively the compounds of the invention may be converted into a stable lyophilizate (possibly with the aid of abovementioned excipients) and, before being administered, reconstituted with a suitable solvent (e.g. injection-grade water, isotonic saline solution) and administered.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 411

<210> SEQ ID NO 1

<211> LENGTH: 278

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 1

```

Leu Glu Cys Tyr Ser Cys Val Gln Lys Ala Asp Asp Gly Cys Ser Pro
1          5          10          15

Asn Lys Met Lys Thr Val Lys Cys Ala Pro Gly Val Asp Val Cys Thr
20          25          30

Glu Ala Val Gly Ala Val Glu Thr Ile His Gly Gln Phe Ser Leu Ala
35          40          45

Val Arg Gly Cys Gly Ser Gly Leu Pro Gly Lys Asn Asp Arg Gly Leu
50          55          60

Asp Leu His Gly Leu Leu Ala Phe Ile Gln Leu Gln Gln Cys Ala Gln
65          70          75          80

Asp Arg Cys Asn Ala Lys Leu Asn Leu Thr Ser Arg Ala Leu Asp Pro
85          90          95

Ala Gly Asn Glu Ser Ala Tyr Pro Pro Asn Gly Val Glu Cys Tyr Ser
100         105         110

Cys Val Gly Leu Ser Arg Glu Ala Cys Gln Gly Thr Ser Pro Pro Val
115         120         125

Val Ser Cys Tyr Asn Ala Ser Asp His Val Tyr Lys Gly Cys Phe Asp
130         135         140

Gly Asn Val Thr Leu Thr Ala Ala Asn Val Thr Val Ser Leu Pro Val
145         150         155         160

Arg Gly Cys Val Gln Asp Glu Phe Cys Thr Arg Asp Gly Val Thr Gly
165         170         175

Pro Gly Phe Thr Leu Ser Gly Ser Cys Cys Gln Gly Ser Arg Cys Asn
180         185         190

Ser Asp Leu Arg Asn Lys Thr Tyr Phe Ser Pro Arg Ile Pro Pro Leu
195         200         205

Val Arg Leu Pro Pro Pro Glu Pro Thr Thr Val Ala Ser Thr Thr Ser
210         215         220

Val Thr Thr Ser Thr Ser Ala Pro Val Arg Pro Thr Ser Thr Thr Lys
225         230         235         240

Pro Met Pro Ala Pro Thr Ser Gln Thr Pro Arg Gln Gly Val Glu His
245         250         255

Glu Ala Ser Arg Asp Glu Glu Pro Arg Leu Thr Gly Gly Ala Ala Gly
260         265         270

His Gln Asp Arg Ser Asn
275

```

-continued

```

<210> SEQ ID NO 2
<211> LENGTH: 293
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 2
Leu Glu Cys Tyr Ser Cys Val Gln Lys Ala Asp Asp Gly Cys Ser Pro
1          5          10          15
His Arg Met Lys Thr Val Lys Cys Gly Pro Gly Val Asp Val Cys Thr
20          25          30
Glu Ala Val Gly Ala Val Glu Thr Ile His Gly Gln Phe Ser Val Ala
35          40          45
Val Arg Gly Cys Gly Ser Gly Ile Pro Gly Lys Asn Asp Arg Gly Leu
50          55          60
Asp Leu His Gly Leu Leu Ala Phe Phe Gln Leu Gln Gln Cys Ser Glu
65          70          75          80
Asp Arg Cys Asn Ala Lys Leu Asn Leu Thr Leu Arg Gly Leu Asn Pro
85          90          95
Ala Gly Asn Glu Ser Ala Tyr Glu Pro Asn Gly Ala Glu Cys Tyr Ser
100         105         110
Cys Val Gly Leu Ser Arg Glu Lys Cys Gln Gly Ser Met Pro Pro Val
115         120         125
Val Asn Cys Tyr Asn Ala Ser Gly Arg Val Tyr Lys Gly Cys Phe Asp
130         135         140
Gly Asn Val Thr Leu Thr Ala Ala Asn Val Thr Val Ser Leu Pro Val
145         150         155         160
Arg Gly Cys Val Gln Asp Glu Thr Cys Thr Arg Asp Gly Val Thr Gly
165         170         175
Pro Gly Phe Thr Leu Ser Gly Ser Cys Cys Gln Gly Pro Arg Cys Asn
180         185         190
Ala Asp Leu Arg Asn Lys Thr Tyr Phe Ser Pro Arg Ile Pro Pro Leu
195         200         205
Val Leu Leu Pro Pro Pro Thr Thr Ala Ala Pro Ser Thr Arg Ala Gln
210         215         220
Asn Ser Ser Ser Thr Thr Ser Thr Ala Ala Pro Thr Thr Thr Thr Ser
225         230         235         240
Ile Ile Lys Pro Thr Thr Ala Gln Ala Ser His Thr Ser Pro His Glu
245         250         255
Met Asp Leu Glu Val Ile Gln Glu Glu Gly Ala Ser Leu Ser Gly Gly
260         265         270
Ala Ala Gly His Gly Gly Thr Ala Gly His Gly Gly Ala Ala Gly His
275         280         285
Gln Asp Arg Ser Asn
290

<210> SEQ ID NO 3
<211> LENGTH: 346
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 3
Met Asp Pro Ala Arg Lys Ala Gly Ala Gln Ala Met Ile Trp Thr Ala
1          5          10          15
Gly Trp Leu Leu Leu Leu Leu Arg Gly Gly Ala Gln Ala Leu Glu

```

-continued

					20						25						30								
Cys	Tyr	Ser	Cys	Val	Gln	Lys	Ala	Asp	Asp	Gly	Cys	Ser	Pro	Asn	Lys										
					35						40						45								
Met	Lys	Thr	Val	Lys	Cys	Ala	Pro	Gly	Val	Asp	Val	Cys	Thr	Glu	Ala										
					50						55						60								
Val	Gly	Ala	Val	Glu	Thr	Ile	His	Gly	Gln	Phe	Ser	Leu	Ala	Val	Arg										
					65						70						75								
Gly	Cys	Gly	Ser	Gly	Leu	Pro	Gly	Lys	Asn	Asp	Arg	Gly	Leu	Asp	Leu										
					85						90						95								
His	Gly	Leu	Leu	Ala	Phe	Ile	Gln	Leu	Gln	Gln	Cys	Ala	Gln	Asp	Arg										
					100						105						110								
Cys	Asn	Ala	Lys	Leu	Asn	Leu	Thr	Ser	Arg	Ala	Leu	Asp	Pro	Ala	Gly										
					115						120						125								
Asn	Glu	Ser	Ala	Tyr	Pro	Pro	Asn	Gly	Val	Glu	Cys	Tyr	Ser	Cys	Val										
					130						135						140								
Gly	Leu	Ser	Arg	Glu	Ala	Cys	Gln	Gly	Thr	Ser	Pro	Pro	Val	Val	Ser										
					145						150						155								
Cys	Tyr	Asn	Ala	Ser	Asp	His	Val	Tyr	Lys	Gly	Cys	Phe	Asp	Gly	Asn										
					165						170						175								
Val	Thr	Leu	Thr	Ala	Ala	Asn	Val	Thr	Val	Ser	Leu	Pro	Val	Arg	Gly										
					180						185						190								
Cys	Val	Gln	Asp	Glu	Phe	Cys	Thr	Arg	Asp	Gly	Val	Thr	Gly	Pro	Gly										
					195						200						205								
Phe	Thr	Leu	Ser	Gly	Ser	Cys	Cys	Gln	Gly	Ser	Arg	Cys	Asn	Ser	Asp										
					210						215						220								
Leu	Arg	Asn	Lys	Thr	Tyr	Phe	Ser	Pro	Arg	Ile	Pro	Pro	Leu	Val	Arg										
					225						230						235								
Leu	Pro	Pro	Pro	Glu	Pro	Thr	Thr	Val	Ala	Ser	Thr	Thr	Ser	Val	Thr										
					245						250						255								
Thr	Ser	Thr	Ser	Ala	Pro	Val	Arg	Pro	Thr	Ser	Thr	Thr	Lys	Pro	Met										
					260						265						270								
Pro	Ala	Pro	Thr	Ser	Gln	Thr	Pro	Arg	Gln	Gly	Val	Glu	His	Glu	Ala										
					275						280						285								
Ser	Arg	Asp	Glu	Glu	Pro	Arg	Leu	Thr	Gly	Gly	Ala	Ala	Gly	His	Gln										
					290						295						300								
Asp	Arg	Ser	Asn	Ser	Gly	Gln	Tyr	Pro	Ala	Lys	Gly	Gly	Pro	Gln	Gln										
					305						310						315								
Pro	His	Asn	Lys	Gly	Cys	Val	Ala	Pro	Thr	Ala	Gly	Leu	Ala	Ala	Leu										
					325						330						335								
Leu	Leu	Ala	Val	Ala	Ala	Gly	Val	Leu	Leu																
					340						345														

<210> SEQ ID NO 4

<211> LENGTH: 363

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 4

Met	Asp	Ala	Ala	Arg	Arg	Gly	Asp	Thr	Gln	Pro	Val	Met	Trp	Thr	Thr
1				5					10					15	
Gly	Trp	Leu	Leu	Leu	Leu	Pro	Leu	Leu	Leu	Cys	Glu	Gly	Ala	Gln	Ala
		20						25					30		
Leu	Glu	Cys	Tyr	Ser	Cys	Val	Gln	Lys	Ala	Asp	Asp	Gly	Cys	Ser	Pro

-continued

35					40					45					
His	Arg	Met	Lys	Thr	Val	Lys	Cys	Gly	Pro	Gly	Val	Asp	Val	Cys	Thr
50					55					60					
Glu	Ala	Val	Gly	Ala	Val	Glu	Thr	Ile	His	Gly	Gln	Phe	Ser	Val	Ala
65					70					75					80
Val	Arg	Gly	Cys	Gly	Ser	Gly	Ile	Pro	Gly	Lys	Asn	Asp	Arg	Gly	Leu
			85						90					95	
Asp	Leu	His	Gly	Leu	Leu	Ala	Phe	Phe	Gln	Leu	Gln	Gln	Cys	Ser	Glu
			100					105						110	
Asp	Arg	Cys	Asn	Ala	Lys	Leu	Asn	Leu	Thr	Leu	Arg	Gly	Leu	Asn	Pro
			115					120					125		
Ala	Gly	Asn	Glu	Ser	Ala	Tyr	Glu	Pro	Asn	Gly	Ala	Glu	Cys	Tyr	Ser
			130					135					140		
Cys	Val	Gly	Leu	Ser	Arg	Glu	Lys	Cys	Gln	Gly	Ser	Met	Pro	Pro	Val
145					150					155					160
Val	Asn	Cys	Tyr	Asn	Ala	Ser	Gly	Arg	Val	Tyr	Lys	Gly	Cys	Phe	Asp
				165					170					175	
Gly	Asn	Val	Thr	Leu	Thr	Ala	Ala	Asn	Val	Thr	Val	Ser	Leu	Pro	Val
			180					185						190	
Arg	Gly	Cys	Val	Gln	Asp	Glu	Thr	Cys	Thr	Arg	Asp	Gly	Val	Thr	Gly
			195					200					205		
Pro	Gly	Phe	Thr	Leu	Ser	Gly	Ser	Cys	Cys	Gln	Gly	Pro	Arg	Cys	Asn
			210					215					220		
Ala	Asp	Leu	Arg	Asn	Lys	Thr	Tyr	Phe	Ser	Pro	Arg	Ile	Pro	Pro	Leu
225					230					235					240
Val	Leu	Leu	Pro	Pro	Pro	Thr	Thr	Ala	Ala	Pro	Ser	Thr	Arg	Ala	Gln
				245					250					255	
Asn	Ser	Ser	Ser	Thr	Thr	Ser	Thr	Ala	Ala	Pro	Thr	Thr	Thr	Thr	Ser
				260					265					270	
Ile	Ile	Lys	Pro	Thr	Thr	Ala	Gln	Ala	Ser	His	Thr	Ser	Pro	His	Glu
			275					280					285		
Met	Asp	Leu	Glu	Val	Ile	Gln	Glu	Glu	Gly	Ala	Ser	Leu	Ser	Gly	Gly
			290					295					300		
Ala	Ala	Gly	His	Gly	Gly	Thr	Ala	Gly	His	Gly	Gly	Ala	Ala	Gly	His
305					310					315					320
Gln	Asp	Arg	Ser	Asn	Met	Glu	Lys	Tyr	Pro	Gly	Lys	Gly	Gly	Ala	Gln
				325					330					335	
Ile	Pro	Ala	Lys	Gly	Gly	Ser	Gly	Thr	Leu	Gly	Ser	Trp	Leu	Ser	Ala
			340					345						350	
Val	Leu	Leu	Thr	Val	Val	Ala	Gly	Ala	Met	Leu					
			355					360							

<210> SEQ ID NO 5

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 5

Phe Ser Asn Ala Trp Met Ser Trp Val
 1 5

<210> SEQ ID NO 6

-continued

<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 6

Phe Ser Asp Tyr Gln Met Thr Trp Ile
1 5

<210> SEQ ID NO 7
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 7

Phe Gly His Tyr Tyr Met Phe Trp Ile
1 5

<210> SEQ ID NO 8
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 8

Phe Ser Ser Asn Tyr Met Ser Trp
1 5

<210> SEQ ID NO 9
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 9

Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 10

Val Ser Gly Val Ser Trp Asn Gly Ala Arg Thr His Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 11
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

-continued

<400> SEQUENCE: 11

Val Ser Ala Ile Ser Gly Ser Gly Tyr Ser Thr His Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 12

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 12

Val Ser Ala Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 13

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 13

Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr
1 5 10

<210> SEQ ID NO 14

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 14

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ala Tyr Tyr Phe Asp Ser
1 5 10 15

<210> SEQ ID NO 15

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 15

Ala Arg Leu Pro Tyr Gly Ser Gln Ser Gly Val Asp Tyr
1 5 10

<210> SEQ ID NO 16

<211> LENGTH: 17

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 16

Ala Arg Glu Ser Gly Gly Ser Gly Pro Asn Tyr Tyr Tyr Gly Met Asp
1 5 10 15

Val

-continued

<210> SEQ ID NO 17
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 17

Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 18
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 18

Ser Gly Ser Ser Ser Asn Val Gly Ser Asn Pro Val Asn
1 5 10

<210> SEQ ID NO 19
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 19

Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 20
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 20

Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 21
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 21

Asp Asn Asn Lys Arg Pro Ser
1 5

<210> SEQ ID NO 22
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 22

Arg Asn Asn Gln Arg Pro Ser

-continued

1 5

<210> SEQ ID NO 23
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 23

Ser Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 24
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 24

Ser Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 25
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 25

Cys Ala Ala Trp Asp Asp Arg Leu Asn Gly Pro Val
1 5 10

<210> SEQ ID NO 26
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 26

Cys Ala Ala Trp Asp Asp Arg Leu Asn Gly Trp Val
1 5 10

<210> SEQ ID NO 27
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 27

Cys Gln Ser Tyr Asp Ser Ser His Val Leu
1 5 10

<210> SEQ ID NO 28
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 28

-continued

Cys Gln Ser Tyr Asp Arg Ser Leu Arg Gly Trp Val
1 5 10

<210> SEQ ID NO 29
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 29

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15
Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30
Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35 40 45
Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
50 55 60
Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65 70 75 80
Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg
85 90 95
Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100 105 110

Gln

<210> SEQ ID NO 30
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 30

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15
Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Val Gly Ser Asn
20 25 30
Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35 40 45
Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
50 55 60
Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg
65 70 75 80
Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg Leu
85 90 95
Asn Gly Trp Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln
100 105 110

<210> SEQ ID NO 31
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 31

-continued

```

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1          5          10          15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
          20          25          30

Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
          35          40          45

Leu Ile Tyr Ser Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe
          50          55          60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65          70          75          80

Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser
          85          90          95

His Val Leu Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln
          100          105          110

```

```

<210> SEQ ID NO 32
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 32

```

```

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1          5          10          15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
          20          25          30

Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
          35          40          45

Leu Ile Tyr Ser Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe
          50          55          60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65          70          75          80

Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Ser
          85          90          95

Leu Arg Gly Trp Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
          100          105          110

```

```

Gln

```

```

<210> SEQ ID NO 33
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 33

```

```

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1          5          10          15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
          20          25          30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
          35          40          45

Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
          50          55          60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr

```

-continued

65	70	75	80
Leu Gln Met Asn Ser	Leu Arg Ala Glu Asp Thr	Ala Val Tyr Tyr Cys	
	85	90	95
Ala Arg Glu Gly Leu Trp	Ala Phe Asp Tyr Trp	Gly Gln Gly Thr Leu	
	100	105	110
Val Thr Val Thr Ser			
	115		

<210> SEQ ID NO 34
 <211> LENGTH: 122
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 34

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly	
1 5 10 15	
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr	
20 25 30	
Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val	
35 40 45	
Ser Gly Val Ser Trp Asn Gly Ala Arg Thr His Tyr Ala Asp Ser Val	
50 55 60	
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr	
65 70 75 80	
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys	
85 90 95	
Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ala Tyr Tyr Phe Asp Ser Trp	
100 105 110	
Gly Gln Gly Thr Leu Val Thr Val Thr Ser	
115 120	

<210> SEQ ID NO 35
 <211> LENGTH: 120
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 35

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly	
1 5 10 15	
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Gly His Tyr	
20 25 30	
Tyr Met Phe Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val	
35 40 45	
Ser Ala Ile Ser Gly Ser Gly Tyr Ser Thr His Tyr Ala Asp Ser Val	
50 55 60	
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr	
65 70 75 80	
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys	
85 90 95	
Ala Arg Leu Pro Tyr Gly Ser Gln Ser Gly Val Asp Tyr Trp Gly Gln	
100 105 110	
Gly Thr Leu Val Thr Val Thr Ser	

115	120	
<p><210> SEQ ID NO 36 <211> LENGTH: 124 <212> TYPE: PRT <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: C4.4a binder</p>		
<p><400> SEQUENCE: 36</p>		
Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly		
1 5 10 15		
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Asn		
20 25 30		
Tyr Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val		
35 40 45		
Ser Ala Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser Val		
50 55 60		
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		
65 70 75 80		
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys		
85 90 95		
Ala Arg Glu Ser Gly Gly Ser Gly Pro Asn Tyr Tyr Tyr Gly Met Asp		
100 105 110		
Val Trp Gly Gln Gly Thr Leu Val Thr Val Thr Ser		
115 120		
<p><210> SEQ ID NO 37 <211> LENGTH: 339 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: C4.4a binder</p>		
<p><400> SEQUENCE: 37</p>		
cagtctgtgc tgactcagcc accctcagcg tctgggaccc ctgggcagag ggtcaccatc	60	
tcctgcactg ggagcagctc caacattggg gcggggtatg ttgtacattg gtatcagcag	120	
ctcccaggaa cggcccccaa actcctcatc tatgacaata ataagcgacc ctcagggggc	180	
cctgaccgat tctctggctc caagtctggc acctcagcct ccttgcccat cagtgggctc	240	
cgggtccgagg atgaggctga ttattactgt gcagcatggg atgacaggct gaatggtcctg	300	
gtgttcggcg gaggaaccaa gttaaccgtc ctaggtcag	339	
<p><210> SEQ ID NO 38 <211> LENGTH: 336 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: C4.4a binder</p>		
<p><400> SEQUENCE: 38</p>		
cagtctgtgc tgactcagcc accctcagcg tctgggaccc cgggcagag ggtcaccatc	60	
tcttgttctg gaagcagctc caacgtcggg agtaatcctg taaactggta tcagcagctc	120	
ccaggaaacgg cccccaaact cctcatctat aggaataatc agcggccctc aggggtccct	180	
gaccgattct ctggctccaa gtctggcacc tcagcctccc tggccatcag tgggtccgg	240	
tccgaggatg aggctgatta ttactgtgca gcatgggatg acaggctgaa tggttgggtg	300	

-continued

ttcggcggag gaaccaagct gacggtccta ggtcag 336

<210> SEQ ID NO 39
<211> LENGTH: 333
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 39

cagtctgtgc tgactcagcc accctcagcg tctgggaccc ccgggcagag ggtcaccatc 60
tcctgcactg ggagcagctc caacattggg gcgggttatg ttgtacattg gtatcagcag 120
ctcccaggaa cggcccccaa actcctcatc tatagtaata atcagcggcc ctccaggagtc 180
cctgaccgat tctctggctc caagtctggc acctcagcct ccctggccat cagtgggctc 240
cggtcaggag atgaggctga ttattactgc cagtcctatg acagcagcca tgttttattc 300
ggcggaggaa ccaagctgac ggtcctaggt cag 333

<210> SEQ ID NO 40
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 40

cagtctgtgc tgactcagcc accctcagcg tctgggaccc ccgggcagag ggtcaccatc 60
tcctgcactg ggagcagctc caacattggg gcgggttatg ttgtacattg gtatcagcag 120
ctcccaggaa cggcccccaa actcctcatc tatagtaata atcagcggcc ctccagggtc 180
cctgaccgat tctctggctc caagtctggc acctcagcct ccctggccat cagtgggctc 240
cggtcaggag atgaggctga ttattactgc cagtcctatg acagaagcct gcgtggttgg 300
gtgttcggcg gaggaaccaa gctgacggtc ctaggtcag 339

<210> SEQ ID NO 41
<211> LENGTH: 351
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 41

gaggtgcagc tgttgagtc tgggggaggc ttggtacagc ctggggggtc cctgagactc 60
tcctgtgcag cctctggatt caccttcagt aacgcctgga tgagctgggt ccgccaggct 120
ccaggaaggg ggctggagtg ggtttcatc attagtagta gtggtagtag catatactac 180
gcagactctg tgaagggccg attcaccatc tccagagaca attccaagaa cagcgtgtat 240
ctgcaaatga acagcctgag agccgaggac actgccgtgt attactgtgc gagagaaggg 300
ttatgggcct ttgactactg gggccagggt accctggtca ccgtgactag t 351

<210> SEQ ID NO 42
<211> LENGTH: 366
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

-continued

<400> SEQUENCE: 42

```

gaggtgcagc tgttgagtc tgggggaggc ttggtacagc ctgggggggc cctgagactc    60
tcctgtgcag cctctggatt caccttcagt gactatcaga tgacctggat ccgccagact    120
ccaggaagg ggctggagtg ggtatcgggt gttagttaga atggcgctag gacgcactat    180
gcagactctg tgaagggccg attcaccatc tccagagaca attccaagaa cacgctgtat    240
ctacaaatga acagcctgag agccgaggac actgccgtgt attactgtgc gaagggcgac    300
tacctggttt actccgata ctactttgac tcctggggcc aggggtaccct ggtcacctg    360
actagt                                           366

```

<210> SEQ ID NO 43

<211> LENGTH: 360

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 43

```

gaggtgcagc tgttgagtc tgggggaggc ttggtacagc ctgggggggc cctgagactc    60
tcctgtgcag cctctggatt caccttcggt cactactata tgttctggat ccgtcaggct    120
ccaggaagg ggctggagtg ggtctcagct attagtggta gtggttatag cacacactac    180
gcagactccg tgaagggccg gttcaccatc tccagagaca attccaagaa cacgctgtat    240
ctgcaaatga acagcctgag agccgaggac actgccgtgt attactgtgc gagactgcca    300
tatggttcgc agagtggcgt tgactactgg ggccagggtta ccctggtcac cgtgactagt    360

```

<210> SEQ ID NO 44

<211> LENGTH: 372

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 44

```

gaggtgcagc tgttgagtc tgggggaggc ttggtacagc ctgggggggc cctgagactc    60
tcctgtgcag cctctggatt caccttcagt agcaactaca tgagctgggt ccgccaggct    120
ccaggaagg ggctggagtg ggtctcagct attagtagta gtggtagtag cacatactac    180
gcagactccg tgaagggccg gttcaccatc tccagagaca attccaagaa cacgctgtat    240
ctgcaaatga acagcctgag agccgaggac actgccgtgt attactgtgc gagagaatct    300
ggtgggagcg gaccgaacta ctactacggt atggacgtct ggggccaagg taccctggtc    360
accgtgacta gt                                           372

```

<210> SEQ ID NO 45

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 45

```

Phe Ser Asn Ala Trp Met Ser Trp Val
1           5

```

<210> SEQ ID NO 46

-continued

<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 46

Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 47
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 47

Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr
1 5 10

<210> SEQ ID NO 48
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 48

Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 49
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 49

Asp Asn Asn Lys Arg Pro Ser
1 5

<210> SEQ ID NO 50
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 50

Cys Ala Ala Trp Asp Asp Arg Leu Asn Gly Pro Val
1 5 10

<210> SEQ ID NO 51
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 51

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

-continued

1	5	10	15
Ser Leu Arg	Leu Ser Cys Ala Ala	Ser Gly Phe Thr Phe Ser Asn Ala	
	20	25	30
Trp Met Ser	Trp Val Arg Gln Ala Pro	Gly Lys Gly Leu Glu Trp Val	
	35	40	45
Ser Tyr Ile	Ser Ser Ser Gly Ser Thr	Ile Tyr Tyr Ala Asp Ser Val	
	50	55	60
Lys Gly Arg	Phe Thr Ile Ser Arg Asp	Asn Ser Lys Asn Thr Leu Tyr	
	65	70	75
Leu Gln Met	Asn Ser Leu Arg Ala Glu	Asp Thr Ala Val Tyr Tyr Cys	
	85	90	95
Ala Arg Glu	Gly Leu Trp Ala Phe Asp	Tyr Trp Gly Gln Gly Thr Leu	
	100	105	110
Val Thr Val	Ser Ser		
	115		

<210> SEQ ID NO 52
 <211> LENGTH: 113
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 52

Glu Ser Val	Leu Thr Gln Pro Pro	Ser Val Ser Gly Ala Pro Gly Gln	
1	5	10	15
Arg Val Thr	Ile Ser Cys Thr Gly Ser Ser	Ser Asn Ile Gly Ala Gly	
	20	25	30
Tyr Val Val	His Trp Tyr Gln Gln Leu Pro	Gly Thr Ala Pro Lys Leu	
	35	40	45
Leu Ile Tyr	Asp Asn Asn Lys Arg Pro Ser	Gly Val Pro Asp Arg Phe	
	50	55	60
Ser Gly Ser	Lys Ser Gly Thr Ser Ala Ser	Leu Ala Ile Ser Gly Leu	
	65	70	75
Arg Ser Glu	Asp Glu Ala Asp Tyr Tyr Cys	Ala Ala Trp Asp Asp Arg	
	85	90	95
Leu Asn Gly	Pro Val Phe Gly Gly Gly Thr	Lys Leu Thr Val Leu Gly	
	100	105	110

Gln

<210> SEQ ID NO 53
 <211> LENGTH: 351
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 53

gaggtgcagc tgctggaaag cggcggagga ctggtgcagc ctggaggcag cctgagactg	60
tcttgccgcg ccagcggcct caccttcagc aacgcctgga tgagctgggt ccgacaggct	120
cctggcaagg gcctggaatg ggtgtcctac atcagcagca gcggcagcac catctactac	180
gccgacagcg tgaagggccg gttcaccatc agccgggaca acagcaagaa caccctgtac	240
ctgcagatga acagcctgcg ggccgaggac accgccgtgt actactgcgc cagagaaggc	300
ctgtgggcct tcgactactg gggccagggc accctggtca ccgtgtctag c	351

-continued

<210> SEQ ID NO 54
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 54

```
gagagcgtgc tgacccagcc tctagcgtg tccggcgctc ctggccagag agtgaccatc      60
agctgcaccg gcagcagcag caacatcgga gccggctacg tgggtgactg gtatcagcag      120
ctgcccggca ccgcccccaa gctgctgacg tacgacaaca acaagcggcc tagcggcgctg      180
cccgcacagat tcagcggcag caagagcggc accagcgcca gcctggccat cagcggcctg      240
agaagcgagg acgagggcca ctactactgc gccgcctggg acgacagact gaacggccct      300
gtgttcggcg gaggcaccaa gctgaccgtg ctgggacag                                339
```

<210> SEQ ID NO 55
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 55

```
Phe Ser Asn Ala Trp Met Ser Trp Val
1              5
```

<210> SEQ ID NO 56
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 56

```
Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser
1              5              10              15
```

```
Val Lys Gly Arg
                20
```

<210> SEQ ID NO 57
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 57

```
Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr
1              5              10
```

<210> SEQ ID NO 58
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 58

```
Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1              5              10
```

-continued

<210> SEQ ID NO 59
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 59

Asp Asn Asn Lys Arg Pro Ser
1 5

<210> SEQ ID NO 60
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 60

Cys Ala Ala Tyr Asp Asp Ser Leu Ser Gly Pro Val
1 5 10

<210> SEQ ID NO 61
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 61

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val Ser Ser
115

<210> SEQ ID NO 62
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 62

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30

-continued

Tyr	Val	Val	His	Trp	Tyr	Gln	Gln	Leu	Pro	Gly	Thr	Ala	Pro	Lys	Leu
		35					40					45			
Leu	Ile	Tyr	Asp	Asn	Asn	Lys	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	Phe
	50					55					60				
Ser	Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Ser	Gly	Leu
65					70					75				80	
Arg	Ser	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Ala	Ala	Tyr	Asp	Asp	Ser
			85						90					95	
Leu	Ser	Gly	Pro	Val	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly
			100					105					110		

Gln

<210> SEQ ID NO 63
 <211> LENGTH: 351
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 63

```

gagggtgcagc tgctggaatc cggcggaggc ctggtgcagc ctggcggatc tctgagactg      60
tctgcgcgcg ccagcgggett taccttctcc aacgcctgga tgcctctgggt ccgacaggcc      120
cctggcaagg gactggaatg ggtgtcctac atctcctcct ccggtccac catctactac      180
gccgactccg tgaaggcccg gttcaccatc tcccgggaca actccaagaa caccctgtac      240
ctgcagatga actcctgcg gcccgaggac accgccgtgt actactgcgc ccgagagggc      300
ctgtgggect tcgattattg gggccagggc accctggtea ccgtcagetc a                351
  
```

<210> SEQ ID NO 64
 <211> LENGTH: 339
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 64

```

cagtcctgtc tgaccagcc cctttctgtg tctggcgccc ctggccagag agtgaccatc      60
tcttgaccg gctctccag caacatcggc gctggctacg tggcgactg gtatcagcag      120
ctgcccggca ccgcccccaa gctgctgac tacgacaaca acaagcggcc ctccggcgtg      180
cccagacagat tctccggctc caagtccggc acctccgcct cctggccat ctccggcctg      240
agatctgagg acgaggccga ctactactgc gccgcctacg acgactccct gtcgggcctt      300
gtgttcggcg gaggcacaaa gttaaccgtg ctgggccag                                339
  
```

<210> SEQ ID NO 65
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 65

Phe Ser Asn Ala Trp Met Ser Trp Val
 1 5

<210> SEQ ID NO 66
 <211> LENGTH: 20

-continued

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 66

Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 67
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 67

Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr
1 5 10

<210> SEQ ID NO 68
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 68

Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 69
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 69

Asp Asn Asn Lys Arg Pro Ser
1 5

<210> SEQ ID NO 70
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 70

Cys Ala Ala Phe Asp Asp Ser Leu Asn Gly Pro Val
1 5 10

<210> SEQ ID NO 71
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 71

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

-continued

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
 20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys Trp Gly Gln Gly Thr Leu
 100 105 110

Val Thr Val Ser Ser
 115

<210> SEQ ID NO 72
 <211> LENGTH: 113
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 72

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
 1 5 10 15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
 20 25 30

Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
 35 40 45

Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
 50 55 60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
 65 70 75 80

Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Phe Asp Asp Arg
 85 90 95

Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 100 105 110

Gln

<210> SEQ ID NO 73
 <211> LENGTH: 351
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 73

gaggtgcagc tgctggaatc cggcggaggc ctggtgcagc ctggcggatc tctgagactg 60

tcctgcgccg cctccggcctt taccttctcc aacgcctgga tgcctcgggt ccgacaggct 120

cctggcaagg gcctggaatg ggtgtcctac atctcctcct ccggctccac catctactac 180

gccgactccg tgaagggccg gttcaccatc tcccgggaca actccaagaa caccctgtac 240

ctgcagatga actccctgcg ggccgaggac accgccgtgt actactgcgc ccgagagggc 300

ctgtgggcct tcgataagtg gggccagggc accctggtca ccgtcagctc a 351

-continued

<210> SEQ ID NO 74
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 74

```
cagtcctgtgc tgaccagcc tccttcgcc tctggcacc ctggccagag agtgaccatc      60
tcctgcaccg gctcctccag caacatcggc gctggctacg tgggtgactg gtatcagcag      120
ctgcccggca ccgccccaa gctgctgac tacgacaaca acaagcggcc ctcggcgctg      180
cccagacagat tctccggctc caagtccggc acctccgcct ccctggccat ctcggcgctg      240
agatctgagg acgaggccga ctactactgc gccgccttcg acgaccggct gaacgggcct      300
gtgttcggcg gaggcacaaa gttaaccgtg ctgggccag                               339
```

<210> SEQ ID NO 75
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 75

```
Phe Ser Ser Ala Trp Met Ser Trp Val
1                               5
```

<210> SEQ ID NO 76
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 76

```
Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser
1           5           10           15
```

```
Val Lys Gly Arg
                20
```

<210> SEQ ID NO 77
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 77

```
Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr
1           5           10
```

<210> SEQ ID NO 78
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 78

```
Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1           5           10
```

-continued

<210> SEQ ID NO 79
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 79

Asp Asn Asn Lys Arg Pro Ser
1 5

<210> SEQ ID NO 80
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 80

Cys Ala Ala Tyr Asp Asp Ser Leu Ser Gly Pro Val
1 5 10

<210> SEQ ID NO 81
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 81

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Ala
20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val Ser Ser
115

<210> SEQ ID NO 82
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 82

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30

Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu

-continued

35	40	45	
Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe			
50	55	60	
Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu			
65	70	75	80
Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Tyr Asp Asp Ser			
	85	90	95
Leu Ser Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly			
	100	105	110
Gln			
<210> SEQ ID NO 83			
<211> LENGTH: 351			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: C4.4a binder			
<400> SEQUENCE: 83			
gagggtgcagc tgctggaatc cggcggagggc ctggtgcagc ctggcggatc tctgagactg			60
tcttcgcgcc ccagcggcctt taccttctcc agcgcttggg tgcctctggg ccgacaggcc			120
cctggcaagg gactggaatg ggtgtcctac atctcctcct ccggctccac catctactac			180
gccgactccg tgaagggccg gttcaccatc tcccgggaca actccaagaa caccctgtac			240
ctgcagatga actccctgcg ggccgaggac accgccgtgt actactgcgc ccgagagggc			300
ctgtgggcct tcgattattg gggccagggc accctggcca ccgtcagctc a			351
<210> SEQ ID NO 84			
<211> LENGTH: 339			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: C4.4a binder			
<400> SEQUENCE: 84			
cagtcctgtc tgaccagcc cccttctgtg totggcgccc ctggccagag agtgaccatc			60
tcttgccacc gctcctccag caacatcggc gctggctacg tgggtgactg gtatcagcag			120
ctgcccgcca ccgccccaa gctgctgac tacgacaaca acaagcggcc ctccggcgtg			180
cccagacagat tctccggctc caagtccggc acctccgcct cctgggcat ctccggcctg			240
agatctgagg acgagggcca ctactactgc gccgcctacg acgactccct gtcgggcct			300
gtgttcggcg gaggcacaaa gttaaccgtg ctgggccag			339
<210> SEQ ID NO 85			
<211> LENGTH: 9			
<212> TYPE: PRT			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: C4.4a binder			
<400> SEQUENCE: 85			
Phe Ser Ser Ala Trp Met Ser Trp Val			
1	5		
<210> SEQ ID NO 86			
<211> LENGTH: 20			
<212> TYPE: PRT			

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 86

Val Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 87
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 87

Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys
1 5 10

<210> SEQ ID NO 88
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 88

Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 89
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 89

Asp Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 90
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 90

Cys Ala Ala Phe Asp Asp Arg Leu Ser Gly Pro Val
1 5 10

<210> SEQ ID NO 91
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 91

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

-continued

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Ala
 20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys Trp Gly Gln Gly Thr Leu
 100 105 110

Val Thr Val Ser Ser
 115

<210> SEQ ID NO 92
 <211> LENGTH: 113
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 92

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
 1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly
 20 25 30

Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
 35 40 45

Leu Ile Tyr Asp Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe
 50 55 60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
 65 70 75 80

Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Phe Asp Asp Arg
 85 90 95

Leu Ser Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 100 105 110

Gln

<210> SEQ ID NO 93
 <211> LENGTH: 351
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 93

gaggtgcagc tgctggaaag cggcggaggc ctggtgcagc ctggcggaag cctgagactg 60

agctgtgccg ccagcggcctt caccttcagc agcgccctgga tgagctgggt ccgacaggcc 120

cctggcaagg gcctggaatg ggtgtcctac atcagcagca gcggcagcag cacctactac 180

gccgacagcg tgaagggccg gtccaccatc agccgggaca acagcaagaa caccctgtac 240

ctgcagatga acagcctgcg ggccgaggac accgccgtgt actactgcgc cagagaaggc 300

ctgtgggcct tcgataagtg gggccagggc accctggtca ccgtcagctc a 351

-continued

<210> SEQ ID NO 94
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 94

```
cagagcgtgc tgaccagcc tcctagcgcc tctggcacc ctggccagag agtgaccatc      60
agctgcagcg gcagcagcag caacatcgga gccggctacg tgggtgactg gtatcagcag      120
ctgcccggca ccgcccccaa gctgctgacg tacgacaaca accagcggcc cagcggcgctg      180
cccgacagat ttccggcgag caagagcggc accagcgcca gcctggccat cagcggcctg      240
agaagcgagg acgagggcca ctactactgc gccgccttcg acgacagact gagcggccct      300
gtgttcggcg gaggcacaaa gttaaccgtg ctgggccag                               339
```

<210> SEQ ID NO 95
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 95

```
Phe Ser Asp Tyr Gln Met Thr Trp Ile
1                               5
```

<210> SEQ ID NO 96
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 96

```
Val Ser Gly Val Ser Trp Asn Gly Ala Arg Thr His Tyr Ala Asp Ser
1           5           10          15
Val Lys Gly Arg
                20
```

<210> SEQ ID NO 97
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 97

```
Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ala Tyr Tyr Phe Asp Ser
1           5           10          15
```

<210> SEQ ID NO 98
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 98

```
Ser Gly Ser Ser Ser Asn Val Gly Ser Asn Pro Val Asn
1           5           10
```

-continued

<210> SEQ ID NO 99
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 99

Arg Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 100
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 100

Cys Ala Ala Trp Asp Asp Arg Leu Asn Gly Trp Val
1 5 10

<210> SEQ ID NO 101
<211> LENGTH: 122
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 101

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30

Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ala Tyr Tyr Phe Asp Ser Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 102
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 102

Glu Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
20 25 30

Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35 40 45

-continued

Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg
65 70 75 80

Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg Leu
85 90 95

Asn Gly Trp Gly Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln
100 105 110

<210> SEQ ID NO 103

<211> LENGTH: 366

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 103

gaggtgcagc tgctggaaag cggcggagga ctggtgcagc ctggaggcag cctgagactg 60
tcttgccg ccagcggcctt caccttcagc gactaccaga tgacctggat ccgacagacc 120
cctggcaagg gcctggaatg ggtgtccggc atcagctgga acggaggcag caccactac 180
gccgacagcg tgaagggcgg gttcaccatc agccgggaca acagcaagaa caccctgtac 240
ctgcagatga acagcctgcg ggccgaggac accgccgtgt actactgcgc caagggcgac 300
tacctggtgt acagcgcta ctacttcgac agctggggcc agggcacccct ggtcaccgtg 360
tctagc 366

<210> SEQ ID NO 104

<211> LENGTH: 336

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 104

gagagcgtgc tgaccagcc tcctagcgcc tctggcacc ctggccagag agtgaccatc 60
agctgctctg gcagcagcag caacatcgga agcaaccccg tgaactggtg tcagcagctg 120
cccggcaccg cccccaagct gctgatctac cggaacaacc agcggcctag cggcgtgccc 180
gacagattca gcggcagcaa gagcggcacc agcggcagcc tggccatcag cggcctgaga 240
agcaggagcg agggcgacta ctactgcgcc gcctgggacg acagactgaa cggtcggggc 300
ttcgcgaggag gcaccaagct gaccgtgctg ggacag 336

<210> SEQ ID NO 105

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 105

Phe Ser Asp Tyr Gln Met Thr Trp Ile
1 5

<210> SEQ ID NO 106

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 106

Val Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 107
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 107

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ser Tyr Tyr Phe Lys Ser
1 5 10 15

<210> SEQ ID NO 108
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 108

Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn Pro Val Asn
1 5 10

<210> SEQ ID NO 109
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 109

Arg Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 110
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 110

Cys Ala Ala Trp Asp Asp Arg Leu Ser Gly Trp Ala
1 5 10

<210> SEQ ID NO 111
<211> LENGTH: 122
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 111

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr

-continued

20					25					30									
Gln	Met	Thr	Trp	Ile	Arg	Gln	Thr	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val				
35					40					45									
Ser	Gly	Ile	Ser	Trp	Asn	Gly	Gly	Ser	Thr	His	Tyr	Ala	Asp	Ser	Val				
50					55					60									
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr				
65					70					75					80				
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys				
85					90					95									
Ala	Lys	Gly	Asp	Tyr	Leu	Val	Tyr	Ser	Ser	Tyr	Tyr	Phe	Lys	Ser	Trp				
100					105					110									
Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser										
115					120														

<210> SEQ ID NO 112
 <211> LENGTH: 112
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 112

Gln	Ser	Val	Leu	Thr	Gln	Pro	Pro	Ser	Ala	Ser	Gly	Thr	Pro	Gly	Gln
1				5					10					15	
Arg	Val	Thr	Ile	Ser	Cys	Ser	Gly	Ser	Ser	Ser	Asn	Ile	Gly	Ser	Asn
		20					25					30			
Pro	Val	Asn	Trp	Tyr	Gln	Gln	Leu	Pro	Gly	Thr	Ala	Pro	Lys	Leu	Leu
		35				40					45				
Ile	Tyr	Arg	Asn	Asn	Gln	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	Phe	Ser
	50				55					60					
Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Ser	Gly	Leu	Arg
65				70					75					80	
Ser	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Ala	Ala	Trp	Asp	Asp	Arg	Leu
			85					90						95	
Ser	Gly	Trp	Ala	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly	Gln
	100						105					110			

<210> SEQ ID NO 113
 <211> LENGTH: 366
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 113

gaggtgcagc	tgctggaatc	cggcggaggc	ctggtgcagc	ctggcggatc	tctgagactg	60
tctgcgcgcg	cctccggctt	caccttctcc	gactaccaga	tgacctggat	cagacagacc	120
cccggaagg	gcctggaatg	ggtgtccggc	atctcctgga	acggcggctc	caccactac	180
gccgactctg	tgaagggcgc	gttcaccatc	tccggggaca	actccaagaa	caccctgtac	240
ctgcagatga	actcctgcgc	ggccgaggac	accgccgtgt	actactgcgc	caagggcgac	300
tacctgggtg	actcctccta	ctacttcaag	tcctggggcc	agggcaccct	ggtcaccgtc	360
agctca						366

<210> SEQ ID NO 114

-continued

<211> LENGTH: 336
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 114

```
cagtcctgtgc tgaccacgcc tccttcgcgc tctggcaccg ctggccagag agtgaccatc      60
tctgtctccg gctcctcctc caacatcggc tccaaccccg tgaactggta tcagcagctg      120
cccggcaccg cccccaagct gctgatctac cggaacaacc agcggccctc cggcgtgccc      180
gacagattct ccggtccaa gtccggcacc tccgcctccc tgccatctc cggcctgaga      240
tctgaggacg aggccgacta ctactgcgcc gctggggacg accgggtgtc tggtggggt      300
tttggcggcg gaacaaagtt aaccgtgctg ggccag                                336
```

<210> SEQ ID NO 115
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 115

```
Phe Ser Asp Tyr Gln Met Thr Trp Ile
1              5
```

<210> SEQ ID NO 116
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 116

```
Val Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser
1              5              10              15
```

```
Val Lys Gly Arg
                20
```

<210> SEQ ID NO 117
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 117

```
Ala Lys Gly Asp Tyr Leu Val Tyr Lys Ser Tyr Tyr Phe Lys Ser
1              5              10              15
```

<210> SEQ ID NO 118
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 118

```
Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn Pro Val Asn
1              5              10
```

<210> SEQ ID NO 119

-continued

<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 119

Arg Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 120
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 120

Cys Ala Ala Trp Asp Asp Ser Leu Ser Gly Trp Ala
1 5 10

<210> SEQ ID NO 121
<211> LENGTH: 122
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 121

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30

Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Gly Asp Tyr Leu Val Tyr Lys Ser Tyr Tyr Phe Lys Ser Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 122
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 122

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
20 25 30

Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35 40 45

-continued

Ile	Tyr	Arg	Asn	Asn	Gln	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	Phe	Ser
50						55					60				
Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Ser	Gly	Leu	Arg
65					70					75				80	
Ser	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Ala	Ala	Trp	Asp	Asp	Ser	Leu
			85						90					95	
Ser	Gly	Trp	Ala	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly	Gln
			100					105					110		

<210> SEQ ID NO 123
 <211> LENGTH: 366
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 123

gagggtgcagc tgctggaatc cggcggaggc ctggtgcagc ctggcggatc tctgagactg	60
tcttgccgccc cctccggctt caccttctcc gactaccaga tgacctggat cagacagacc	120
cccggcaagg gcctggaatg ggtgtccggc atctctctgga acggcggctc caccactac	180
gccgactctg tgaagggccg gttcaccatc tcccgggaca actccaagaa caccctgtac	240
ctgcagatga actccctgcg ggccgaggac accgccgtgt actactgcgc caagggcgac	300
tacctggtgt acaagtccta ctacttcaag tctggggccc agggcaccct ggtcaccgtc	360
agctca	366

<210> SEQ ID NO 124
 <211> LENGTH: 336
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 124

cagtccgtgc tgaccagcc tccttcgcc tctggcacc ctggccagag agtgaccatc	60
tctgtctccg gctctctcct caacatcgcc tccaaccccg tgaactggta tcagcagctg	120
cccggcaccg cccccaagct gctgatctac cggaacaacc agcggccctc cggcgtgccc	180
gacagattct ccggctccaa gtccggcacc tccgctccc tggccatctc cggcctgaga	240
tctgaggacg agggcgacta ctactgcgcc gcctgggacg actccctgctc tggctgggct	300
tttggcggcg gaacaaagtt aaccgtgctg ggccag	336

<210> SEQ ID NO 125
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 125

Phe	Ser	Asp	Tyr	Gln	Met	Thr	Trp	Ile
1				5				

<210> SEQ ID NO 126
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 126

Val Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser
1 5 10 15Val Lys Gly Arg
20

<210> SEQ ID NO 127

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 127

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ser Tyr Tyr Phe Lys Ser
1 5 10 15

<210> SEQ ID NO 128

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 128

Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn Pro Val Asn
1 5 10

<210> SEQ ID NO 129

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 129

Arg Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 130

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 130

Cys Ala Ala Trp Asp Asp Arg Leu Ser Gly Trp Gly
1 5 10

<210> SEQ ID NO 131

<211> LENGTH: 122

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 131

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30

-continued

Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ser Tyr Tyr Phe Lys Ser Trp
 100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> SEQ ID NO 132
 <211> LENGTH: 112
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 132

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
 1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
 20 25 30

Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45

Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg
 65 70 75 80

Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg Leu
 85 90 95

Ser Gly Trp Gly Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln
 100 105 110

<210> SEQ ID NO 133
 <211> LENGTH: 366
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 133

gagggtgcagc tgctggaatc cggcggaggc ctggtgcagc ctggcggatc tctgagactg 60

tcttcgcgcg cctccggctt caccttctcc gactaccaga tgacctggat cagacagacc 120

cccggcaagg gcctggaatg ggtgtccggc atctcctgga acggcggctc caccactac 180

gccgactctg tgaagggcgc gttcaccatc tcccgggaca actccaagaa caccctgtac 240

ctgcagatga actccctcgc gcccgaggac accgccgtgt actactgcgc caagggcgac 300

tacctggtgt actcctccta ctactcaag tcttggggcc agggcaccct ggtcaccgtc 360

agctca 366

<210> SEQ ID NO 134
 <211> LENGTH: 336

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 134

```
cagtccgtgc tgaccagcc tccttcgcc tctggcacc ctggccagag agtgaccatc      60
tcctgtctccg gctcctcctc caacatcggc tccaaccccg tgaactggta tcagcagctg    120
ccccgcaccg cccccaaagt gctgatctac cggaacaacc agcggccctc cggcgtgccc    180
gacagattct ccggctccaa gtccggcacc tccgcctccc tggccatctc cggcctgaga    240
tctgaggacg aggccgacta ctactgcgcc gcctgggacg accggctgtc tggtctgggga    300
tttggcggcg gaacaaagtt aaccgtgctg ggccag                                336
```

<210> SEQ ID NO 135
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 135

```
Phe Ser Ser Tyr Gln Met Thr Trp Ile
1                      5
```

<210> SEQ ID NO 136
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 136

```
Val Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser
1                      5                      10                     15
```

```
Val Lys Gly Arg
                20
```

<210> SEQ ID NO 137
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 137

```
Ala Lys Gly Asp Tyr Leu Val Tyr Lys Ser Tyr Tyr Phe Lys Ser
1                      5                      10                     15
```

<210> SEQ ID NO 138
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 138

```
Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn Pro Val Asn
1                      5                      10
```

<210> SEQ ID NO 139
<211> LENGTH: 7

-continued

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 139

Arg Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 140
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 140

Cys Ala Ala Trp Asp Asp Ser Leu Ser Gly Trp Ala
1 5 10

<210> SEQ ID NO 141
<211> LENGTH: 122
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 141

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Gln Met Thr Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Gly Asp Tyr Leu Val Tyr Lys Ser Tyr Tyr Phe Lys Ser Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 142
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 142

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
20 25 30

Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35 40 45

Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser

-continued

50	55	60
Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg		
65	70	75 80
Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Ser Leu		
	85	90 95
Ser Gly Trp Ala Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln		
	100	105 110

<210> SEQ ID NO 143
 <211> LENGTH: 366
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 143

```

gaggtgcagc tgctggaaag cggcggaggc ctggtgcagc ctggcggaag cctgagactg      60
agctgtgccg ccagcggett caccttcagc agctaccaga tgacctggat cagacaggcc      120
cctggcaagg gcctggaatg ggtgtccggc atcagctgga acggcggcag caccactac      180
gccgacagcg tgaagggccg gttcaccatc agccgggaca acagcaagaa caccctgtac      240
ctgcagatga acagcctgcg ggccgaggac accgccgtgt actactgcgc caagggcgac      300
tacctggtgt acaagagcta ctactcaag agctggggcc agggcacact ggtcaccgtc      360
agctca                                          366
  
```

<210> SEQ ID NO 144
 <211> LENGTH: 336
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 144

```

cagagcgtgc tgaccagacc tcctagcgcc tctggcacc ctggccagag agtgaccatc      60
agctgcagcg gcagcagcag caacatcggc agcaaccccg tgaactggta tcagcagctg      120
cccggcaccg cccccaagct gctgatctac cggaacaacc agcggcccag cggcgtgccc      180
gacagathtt ccggcagcaa gagcggcacc agcggccagc tggccatcag cggcctgaga      240
agcggaggacg aggcgcacta ctactgcgcc gcctgggacg atagcctgag cggctggggc      300
tttggcggcg gaacaaagtt aaccgtgctg ggccag                                          336
  
```

<210> SEQ ID NO 145
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 145

```

Phe Ser Asn Ala Trp Met Ser Trp Val
1          5
  
```

<210> SEQ ID NO 146
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

-continued

<400> SEQUENCE: 146

Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 147

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 147

Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys
1 5 10

<210> SEQ ID NO 148

<211> LENGTH: 14

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 148

Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 149

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 149

Asp Asn Asn Lys Arg Pro Ser
1 5

<210> SEQ ID NO 150

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 150

Cys Ala Ala Tyr Asp Asp Ser Leu Lys Gly Pro Val
1 5 10

<210> SEQ ID NO 151

<211> LENGTH: 117

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 151

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
20 25 30

-continued

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

 Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60

 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

 Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys Trp Gly Gln Gly Thr Leu
 100 105 110

 Val Thr Val Thr Ser
 115

<210> SEQ ID NO 152
 <211> LENGTH: 113
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 152

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
 1 5 10 15

 Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
 20 25 30

 Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
 35 40 45

 Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
 50 55 60

 Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
 65 70 75 80

 Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Tyr Asp Asp Ser
 85 90 95

 Leu Lys Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 100 105 110

Gln

<210> SEQ ID NO 153
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 153

Phe Ser Asn Ala Trp Met Ser Trp Val
 1 5

<210> SEQ ID NO 154
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 154

Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser
 1 5 10 15

-continued

Val Lys Gly Arg
20

<210> SEQ ID NO 155
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 155

Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys
1 5 10

<210> SEQ ID NO 156
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 156

Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 157
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 157

Asp Asn Asn Lys Arg Pro Ser
1 5

<210> SEQ ID NO 158
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 158

Cys Ala Ala Phe Asp Asp Ser Leu Asn Gly Pro Val
1 5 10

<210> SEQ ID NO 159
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 159

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
50 55 60

-continued

```

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65              70              75              80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
              85              90              95

Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys Trp Gly Gln Gly Thr Leu
              100              105              110

Val Thr Val Thr Ser
              115

```

```

<210> SEQ ID NO 160
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 160

```

```

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
1              5              10              15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
              20              25              30

Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
              35              40              45

Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
              50              55              60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65              70              75              80

Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Phe Asp Asp Ser
              85              90              95

Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
              100              105              110

```

```

Gln

```

```

<210> SEQ ID NO 161
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 161

```

```

Phe Ser Asn Ala Trp Met Ser Trp Val
1              5

```

```

<210> SEQ ID NO 162
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 162

```

```

Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser
1              5              10              15

```

```

Val Lys Gly Arg
              20

```

```

<210> SEQ ID NO 163
<211> LENGTH: 10

```

-continued

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 163

Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys
1 5 10

<210> SEQ ID NO 164
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 164

Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 165
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 165

Asp Asn Asn Lys Arg Pro Ser
1 5

<210> SEQ ID NO 166
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 166

Cys Ala Ala Tyr Asp Asp Ser Leu Ser Gly Pro Val
1 5 10

<210> SEQ ID NO 167
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 167

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

-continued

Ala	Arg	Glu	Gly	Leu	Trp	Ala	Phe	Asp	Lys	Trp	Gly	Gln	Gly	Thr	Leu
		100						105					110		

Val	Thr	Val	Thr	Ser
		115		

<210> SEQ ID NO 168
 <211> LENGTH: 113
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 168

Gln	Ser	Val	Leu	Thr	Gln	Pro	Pro	Ser	Val	Ser	Gly	Ala	Pro	Gly	Gln
1			5					10					15		
Arg	Val	Thr	Ile	Ser	Cys	Thr	Gly	Ser	Ser	Ser	Asn	Ile	Gly	Ala	Gly
		20					25					30			
Tyr	Val	Val	His	Trp	Tyr	Gln	Gln	Leu	Pro	Gly	Thr	Ala	Pro	Lys	Leu
		35				40					45				
Leu	Ile	Tyr	Asp	Asn	Asn	Lys	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	Phe
	50				55					60					
Ser	Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Ser	Gly	Leu
65				70					75				80		
Arg	Ser	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Ala	Ala	Phe	Asp	Asp	Ser
		85					90						95		
Leu	Asn	Gly	Pro	Val	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly
		100					105						110		

Gln

<210> SEQ ID NO 169
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 169

Phe	Ser	Ser	Ala	Trp	Met	Ser	Trp	Val
1			5					

<210> SEQ ID NO 170
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 170

Val	Ser	Tyr	Ile	Ser	Ser	Ser	Gly	Ser	Ser	Thr	Tyr	Tyr	Ala	Asp	Ser
1			5					10					15		

Val	Lys	Gly	Arg
		20	

<210> SEQ ID NO 171
 <211> LENGTH: 10
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 171

-continued

Ala Arg Glu Gly Leu Trp Ala Phe Asp Trp
1 5 10

<210> SEQ ID NO 172
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 172

Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 173
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 173

Asp Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 174
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 174

Cys Ala Ala Tyr Asp Asp Ser Leu Ser Gly Pro Val
1 5 10

<210> SEQ ID NO 175
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 175

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Ala
20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Glu Gly Leu Trp Ala Phe Asp Trp Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val Thr Ser
115

-continued

<210> SEQ ID NO 176
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 176
Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15
Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30
Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35 40 45
Leu Ile Tyr Asp Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe
50 55 60
Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65 70 75 80
Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Tyr Asp Asp Ser
85 90 95
Leu Ser Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100 105 110
Gln

<210> SEQ ID NO 177
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 177
Phe Ser Ser Ala Trp Met Ser Trp Val
1 5

<210> SEQ ID NO 178
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 178
Val Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser
1 5 10 15
Val Lys Gly Arg
20

<210> SEQ ID NO 179
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 179
Ala Arg Glu Gly Leu Trp Ala Phe Asp Asn
1 5 10

<210> SEQ ID NO 180

-continued

<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 180

Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 181
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 181

Asp Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 182
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 182

Cys Ala Ala Tyr Asp Asp Ser Leu Asn Gly Pro Val
1 5 10

<210> SEQ ID NO 183
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 183

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val Thr Ser
115

<210> SEQ ID NO 184
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 184

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15
Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30
Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35 40 45
Leu Ile Tyr Asp Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe
50 55 60
Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65 70 75 80
Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Tyr Asp Asp Ser
85 90 95
Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100 105 110

Gln

<210> SEQ ID NO 185

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 185

Phe Ser Ser Ala Trp Met Ser Trp Val
1 5

<210> SEQ ID NO 186

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 186

Val Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 187

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 187

Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr
1 5 10

<210> SEQ ID NO 188

<211> LENGTH: 14

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

-continued

<400> SEQUENCE: 188

Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
 1 5 10

<210> SEQ ID NO 189

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 189

Asp Asn Asn Gln Arg Pro Ser
 1 5

<210> SEQ ID NO 190

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 190

Cys Ala Ala Trp Asp Asp Arg Leu Asn Gly Pro Val
 1 5 10

<210> SEQ ID NO 191

<211> LENGTH: 117

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 191

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Ala
 20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr Trp Gly Gln Gly Thr Leu
 100 105 110

Val Thr Val Ser Ser
 115

<210> SEQ ID NO 192

<211> LENGTH: 113

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 192

Glu Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
 1 5 10 15

-continued

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30
Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35 40 45
Leu Ile Tyr Asp Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe
50 55 60
Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65 70 75 80
Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg
85 90 95
Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100 105 110

Gln

<210> SEQ ID NO 193
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 193

Phe Thr Phe Ser Asn Ala Trp Met Ser Trp Val
1 5 10

<210> SEQ ID NO 194
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 194

Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 195
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 195

Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr
1 5 10

<210> SEQ ID NO 196
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 196

Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

-continued

```

<210> SEQ ID NO 197
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 197

```

```

Asp Asn Asn Lys Arg Pro Ser
1             5

```

```

<210> SEQ ID NO 198
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 198

```

```

Cys Ala Ala Tyr Asp Asp Arg Leu Asn Gly Pro Val
1             5             10

```

```

<210> SEQ ID NO 199
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 199

```

```

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1             5             10             15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
                20             25             30
Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
        35             40             45
Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
        50             55             60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
        65             70             75             80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
                85             90             95
Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr Trp Gly Gln Gly Thr Leu
        100             105             110
Val Thr Val Ser Ser
        115

```

```

<210> SEQ ID NO 200
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 200

```

```

Glu Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
1             5             10             15
Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
        20             25             30
Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
        35             40             45

```

-continued

Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
50 55 60
Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65 70 75 80
Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Tyr Asp Asp Arg
85 90 95
Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100 105 110

Gln

<210> SEQ ID NO 201
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 201

Phe Thr Phe Ser Ser Ala Trp Met Ser Trp Val
1 5 10

<210> SEQ ID NO 202
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 202

Val Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 203
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 203

Ala Arg Glu Gly Leu Trp Ala Phe Asp Gly
1 5 10

<210> SEQ ID NO 204
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 204

Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 205
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

-continued

<400> SEQUENCE: 205

Asp Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 206

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 206

Cys Ala Ala Tyr Asp Asp Ser Leu Asn Arg Pro Val
1 5 10

<210> SEQ ID NO 207

<211> LENGTH: 117

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 207

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Ala
20 25 30Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser Val
50 55 60Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95Ala Arg Glu Gly Leu Trp Ala Phe Asp Gly Trp Gly Gln Gly Thr Leu
100 105 110Val Thr Val Ser Ser
115

<210> SEQ ID NO 208

<211> LENGTH: 113

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 208

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35 40 45Leu Ile Tyr Asp Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe
50 55 60Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65 70 75 80

-continued

Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Tyr Asp Asp Ser
85 90 95

Leu Asn Arg Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100 105 110

Gln

<210> SEQ ID NO 209
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 209

Phe Thr Phe Ser Asn Ala Trp Met Ser Trp Val
1 5 10

<210> SEQ ID NO 210
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 210

Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 211
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 211

Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr
1 5 10

<210> SEQ ID NO 212
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 212

Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 213
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 213

Asp Asn Asn Lys Arg Pro Ser
1 5

-continued

<210> SEQ ID NO 214
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 214

Cys Ala Ala Phe Asp Asp Ser Leu Asn Gly Pro Val
1 5 10

<210> SEQ ID NO 215
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 215

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
20 25 30
Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr Trp Gly Gln Gly Thr Leu
100 105 110
Val Thr Val Ser Ser
115

<210> SEQ ID NO 216
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 216

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
1 5 10 15
Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30
Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35 40 45
Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
50 55 60
Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65 70 75 80
Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Phe Asp Asp Ser
85 90 95
Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100 105 110

-continued

Gln

<210> SEQ ID NO 217
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 217

Phe Ser Ser Ala Trp Met Ser Trp Val
1 5

<210> SEQ ID NO 218
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 218

Val Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 219
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 219

Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys
1 5 10

<210> SEQ ID NO 220
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 220

Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 221
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 221

Asp Asn Asn Lys Arg Pro Ser
1 5

<210> SEQ ID NO 222
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 222

Cys Ala Ala Phe Asp Asp Ser Leu Asn Gly Pro Val
1 5 10

<210> SEQ ID NO 223

<211> LENGTH: 117

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 223

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Ala
20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val Thr Ser
115

<210> SEQ ID NO 224

<211> LENGTH: 113

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 224

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30

Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35 40 45

Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
50 55 60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65 70 75 80

Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Phe Asp Asp Ser
85 90 95

Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100 105 110

Gln

<210> SEQ ID NO 225

<211> LENGTH: 11

-continued

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 225

Phe Thr Phe Ser Ser Ala Trp Met Ser Trp Val
1 5 10

<210> SEQ ID NO 226
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 226

Val Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 227
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 227

Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys
1 5 10

<210> SEQ ID NO 228
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 228

Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 229
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 229

Asp Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 230
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 230

Cys Ala Ala Phe Asp Asp Ser Leu Asn Gly Pro Val
1 5 10

-continued

<210> SEQ ID NO 231
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 231

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Ala
20 25 30
Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys Trp Gly Gln Gly Thr Leu
100 105 110
Val Thr Val Ser Ser
115

<210> SEQ ID NO 232
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 232

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15
Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30
Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35 40 45
Leu Ile Tyr Asp Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe
50 55 60
Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65 70 75 80
Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Phe Asp Asp Ser
85 90 95
Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100 105 110
Gln

<210> SEQ ID NO 233
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 233

-continued

Phe Thr Phe Ser Ser Ala Trp Met Ser Trp Val
1 5 10

<210> SEQ ID NO 234
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 234

Val Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 235
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 235

Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys
1 5 10

<210> SEQ ID NO 236
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 236

Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 237
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 237

Asp Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 238
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 238

Cys Ala Ala Phe Asp Asp Arg Leu Ser Gly Pro Val
1 5 10

<210> SEQ ID NO 239
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 239

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Ala
20 25 30
Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
Ser Tyr Ile Ser Ser Ser Gly Ser Ser Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys Trp Gly Gln Gly Thr Leu
100 105 110
Val Thr Val Ser Ser
115

<210> SEQ ID NO 240
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 240

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15
Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30
Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35 40 45
Leu Ile Tyr Asp Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe
50 55 60
Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65 70 75 80
Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Phe Asp Asp Arg
85 90 95
Leu Ser Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100 105 110
Gln

<210> SEQ ID NO 241
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 241

Phe Ser Asp Tyr Gln Met Thr Trp Ile
1 5

<210> SEQ ID NO 242

-continued

<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 242

Val Ser Gly Val Ser Trp Asn Gly Ala Arg Thr His Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 243
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 243

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ser Tyr Tyr Phe Lys Ser
1 5 10 15

<210> SEQ ID NO 244
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 244

Ser Gly Ser Ser Ser Asn Val Gly Ser Asn Pro Val Asn
1 5 10

<210> SEQ ID NO 245
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 245

Arg Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 246
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 246

Cys Ala Ala Trp Asp Asp Arg Leu Asn Gly Trp Thr Gly
1 5 10

<210> SEQ ID NO 247
<211> LENGTH: 122
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 247

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

-continued

1	5	10	15
Ser Leu Arg	Leu Ser Cys Ala Ala Ser	Gly Phe Thr Phe Ser Asp Tyr	
	20	25	30
Gln Met Thr	Trp Ile Arg Gln Thr Pro	Gly Lys Gly Leu Glu Trp Val	
	35	40	45
Ser Gly Val	Ser Trp Asn Gly Ala Arg Thr His Tyr Ala Asp Ser Val		
	50	55	60
Lys Gly Arg	Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		
	65	70	75
Leu Gln Met	Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys		
	85	90	95
Ala Lys Gly	Asp Tyr Leu Val Tyr Ser Ser Tyr Tyr Phe Lys Ser Trp		
	100	105	110
Gly Gln Gly	Thr Leu Val Thr Val Thr Ser		
	115	120	

<210> SEQ ID NO 248
 <211> LENGTH: 112
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 248

Gln Ser Val	Leu Thr Gln Pro Pro Ser	Ala Ser Gly Thr Pro Gly Gln
1	5	10
Arg Val Thr	Ile Ser Cys Ser Gly Ser Ser Ser Asn Val Gly Ser Asn	
	20	25
Pro Val Asn	Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu	
	35	40
Ile Tyr Arg	Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser	
	50	55
Gly Ser Lys	Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg	
	65	70
Ser Glu Asp	Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg Leu	
	85	90
Asn Gly Trp	Gly Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln	
	100	105

<210> SEQ ID NO 249
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 249

Phe Thr Phe	Ser Ser Tyr Gln Met Thr Trp Ile
1	5
	10

<210> SEQ ID NO 250
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 250

-continued

Val	Ser	Gly	Ile	Ser	Trp	Asn	Gly	Gly	Ser	Thr	His	Tyr	Ala	Asp	Ser
1				5					10					15	

Val	Lys	Gly	Arg
			20

<210> SEQ ID NO 251
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 251

Ala	Lys	Gly	Asp	Tyr	Leu	Val	Tyr	Ser	Ala	Tyr	Tyr	Phe	Asp	Ser
1				5				10					15	

<210> SEQ ID NO 252
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 252

Ser	Gly	Ser	Ser	Ser	Asn	Ile	Gly	Ser	Asn	Pro	Val	Asn
1				5				10				

<210> SEQ ID NO 253
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 253

Arg	Asn	Asn	Gln	Arg	Pro	Ser
1				5		

<210> SEQ ID NO 254
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 254

Cys	Ala	Ala	Trp	Asp	Asp	Arg	Leu	Asn	Gly	Trp	Gly
1				5				10			

<210> SEQ ID NO 255
 <211> LENGTH: 122
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 255

Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly
1				5					10					15	

Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Ser	Tyr
			20					25					30		

Gln	Met	Thr	Trp	Ile	Arg	Gln	Thr	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
	35						40					45			

-continued

```

Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val
  50          55          60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
  65          70          75          80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
          85          90          95

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ala Tyr Tyr Phe Asp Ser Trp
          100          105          110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
  115          120

```

```

<210> SEQ ID NO 256
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 256

```

```

Glu Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
  1          5          10          15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
          20          25          30

Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
          35          40          45

Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
          50          55          60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg
  65          70          75          80

Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg Leu
          85          90          95

Asn Gly Trp Gly Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln
          100          105          110

```

```

<210> SEQ ID NO 257
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 257

```

```

Phe Thr Phe Ser Asp Tyr Gln Met Thr Trp Ile
  1          5          10

```

```

<210> SEQ ID NO 258
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 258

```

```

Val Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser
  1          5          10          15

Val Lys Gly Arg
          20

```

```

<210> SEQ ID NO 259

```

-continued

<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 259

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ser Tyr Tyr Phe Lys Tyr
1 5 10 15

<210> SEQ ID NO 260
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 260

Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn Pro Val Asn
1 5 10

<210> SEQ ID NO 261
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 261

Arg Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 262
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 262

Cys Ala Ala Trp Asp Asp Arg Leu Asn Gly Trp Ala
1 5 10

<210> SEQ ID NO 263
<211> LENGTH: 122
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 263

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30

Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

-continued

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ser Tyr Tyr Phe Lys Tyr Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 264
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 264

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
20 25 30

Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35 40 45

Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg
65 70 75 80

Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg Leu
85 90 95

Asn Gly Trp Ala Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln
100 105 110

<210> SEQ ID NO 265
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 265

Phe Thr Phe Ser Asp Tyr Gln Met Thr Trp Ile
1 5 10

<210> SEQ ID NO 266
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 266

Val Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 267
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 267

-continued

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ser Tyr Tyr Phe Lys Ser
1 5 10 15

<210> SEQ ID NO 268
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 268

Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn Pro Val Asn
1 5 10

<210> SEQ ID NO 269
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 269

Arg Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 270
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 270

Cys Ala Ala Trp Asp Asp Ser Leu Asn Gly Trp Gly
1 5 10

<210> SEQ ID NO 271
 <211> LENGTH: 122
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 271

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30

Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ser Tyr Tyr Phe Lys Ser Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

-continued

<210> SEQ ID NO 272
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 272

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15
Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
20 25 30
Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35 40 45
Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
50 55 60
Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg
65 70 75 80
Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Ser Leu
85 90 95
Asn Gly Trp Gly Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln
100 105 110

<210> SEQ ID NO 273
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 273

Phe Thr Phe Ser Asp Tyr Gln Met Thr Trp Ile
1 5 10

<210> SEQ ID NO 274
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 274

Val Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser
1 5 10 15
Val Lys Gly Arg
20

<210> SEQ ID NO 275
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 275

Ala Lys Gly Asp Tyr Leu Val Tyr Lys Ser Tyr Tyr Phe Lys Ser
1 5 10 15

<210> SEQ ID NO 276
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 276

Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn Pro Val Asn
1 5 10

<210> SEQ ID NO 277

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 277

Arg Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 278

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 278

Cys Ala Ala Trp Asp Asp Arg Leu Ser Gly Trp Gly
1 5 10

<210> SEQ ID NO 279

<211> LENGTH: 122

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 279

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30

Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Gly Asp Tyr Leu Val Tyr Lys Ser Tyr Tyr Phe Lys Ser Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 280

<211> LENGTH: 112

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 280

-continued

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15
Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
20 25 30
Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35 40 45
Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
50 55 60
Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg
65 70 75 80
Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg Leu
85 90 95
Ser Gly Trp Gly Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln
100 105 110

<210> SEQ ID NO 281
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 281

Phe Thr Phe Ser Asp Tyr Gln Met Thr Trp Ile
1 5 10

<210> SEQ ID NO 282
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 282

Val Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 283
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 283

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ser Tyr Tyr Phe Lys Ser
1 5 10 15

<210> SEQ ID NO 284
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 284

Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn Pro Val Asn
1 5 10

-continued

<210> SEQ ID NO 285
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 285

Arg Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 286
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 286

Cys Ala Ala Trp Asp Asp Arg Leu Ser Gly Trp Gly
1 5 10

<210> SEQ ID NO 287
<211> LENGTH: 122
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 287

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30

Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ser Tyr Tyr Phe Lys Ser Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 288
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 288

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
20 25 30

Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu

-continued

Ile	Tyr	Arg	Asn	Asn	Gln	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	Phe	Ser
50					55					60					
Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Ser	Gly	Leu	Arg
65				70					75					80	
Ser	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Ala	Ala	Trp	Asp	Asp	Arg	Leu
			85					90						95	
Ser	Gly	Trp	Gly	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly	Gln
	100						105					110			

<210> SEQ ID NO 289
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 289

Phe	Thr	Phe	Ser	Asp	Tyr	Gln	Met	Thr	Trp	Ile
1			5					10		

<210> SEQ ID NO 290
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 290

Val	Ser	Gly	Ile	Ser	Trp	Asn	Gly	Gly	Ser	Thr	His	Tyr	Ala	Asp	Ser
1			5					10						15	

Val	Lys	Gly	Arg
		20	

<210> SEQ ID NO 291
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 291

Ala	Lys	Gly	Asp	Tyr	Leu	Val	Tyr	Lys	Ser	Tyr	Tyr	Phe	Lys	Ser
1			5					10					15	

<210> SEQ ID NO 292
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 292

Ser	Gly	Ser	Ser	Ser	Asn	Ile	Gly	Ser	Asn	Pro	Val	Asn
1			5					10				

<210> SEQ ID NO 293
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

-continued

<400> SEQUENCE: 293

Arg Asn Asn Gln Arg Pro Ser
1 5

<210> SEQ ID NO 294

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 294

Cys Ala Ala Trp Asp Asp Ser Leu Asn Gly Trp Gly
1 5 10

<210> SEQ ID NO 295

<211> LENGTH: 122

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 295

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30

Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Gly Asp Tyr Leu Val Tyr Lys Ser Tyr Tyr Phe Lys Ser Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 296

<211> LENGTH: 112

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 296

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
20 25 30

Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35 40 45

Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg
65 70 75 80

-continued

Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Ser Leu
85 90 95

Asn Gly Trp Gly Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln
100 105 110

<210> SEQ ID NO 297
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: consensus sequence for M20 D02 S-A derived CDR
 H1
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (3)..(3)
 <223> OTHER INFORMATION: X = D or S
 <400> SEQUENCE: 297

Phe Ser Xaa Tyr Gln Met Thr Trp Ile
1 5

<210> SEQ ID NO 298
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: consensus sequence for M20 D02 S-A derived
 CDRH2
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (4)..(4)
 <223> OTHER INFORMATION: X = V or I
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (9)..(9)
 <223> OTHER INFORMATION: X = A or G
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (10)..(10)
 <223> OTHER INFORMATION: X = R or S
 <400> SEQUENCE: 298

Val Ser Gly Xaa Ser Trp Asn Gly Xaa Xaa Thr His Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 299
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: consensus for M20 D02 S-A derived CDRH3
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (9)..(9)
 <223> OTHER INFORMATION: X = S, K, or R
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (10)..(10)
 <223> OTHER INFORMATION: X = A, S or R
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (14)..(14)
 <223> OTHER INFORMATION: X = D, K, E or R
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (15)..(15)
 <223> OTHER INFORMATION: X = S or Y

-continued

<400> SEQUENCE: 299

Ala Lys Gly Asp Tyr Leu Val Tyr Xaa Xaa Tyr Tyr Phe Xaa Xaa
1 5 10 15

<210> SEQ ID NO 300

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: consensus sequence derived from M20 D02 S-A
CDRL1

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (7)..(7)

<223> OTHER INFORMATION: X = V or I

<400> SEQUENCE: 300

Ser Gly Ser Ser Ser Asn Xaa Gly Ser Asn Pro Val Asn
1 5 10

<210> SEQ ID NO 301

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: consensus sequence derived from M20 D02 S-A
derived CDR L3

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (3)..(3)

<223> OTHER INFORMATION: X = A, Q or R

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (5)..(5)

<223> OTHER INFORMATION: X = D or G

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (7)..(7)

<223> OTHER INFORMATION: X = R or S

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (9)..(9)

<223> OTHER INFORMATION: X = N, W or S

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (12)..(12)

<223> OTHER INFORMATION: X = V, A or G

<400> SEQUENCE: 301

Cys Ala Xaa Trp Xaa Asp Xaa Leu Xaa Gly Trp Xaa
1 5 10

<210> SEQ ID NO 302

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: consensus sequence derived from M31 B01 CDR H1

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (3)..(3)

<223> OTHER INFORMATION: X = N or S

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (4)..(4)

<223> OTHER INFORMATION: X = A or Y

<400> SEQUENCE: 302

Phe Ser Xaa Xaa Trp Met Ser Trp Val
1 5

-continued

<210> SEQ ID NO 303
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: consensus sequence derived from M31 B01 CDR H2
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: X = T or S
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: X = I or T

<400> SEQUENCE: 303

Val Ser Tyr Ile Ser Ser Ser Gly Ser Xaa Xaa Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly Arg
20

<210> SEQ ID NO 304
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: consensus sequence derived from M31 B01 CDR H3
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: X = Y, K, G, W or N

<400> SEQUENCE: 304

Ala Arg Glu Gly Leu Trp Ala Phe Asp Xaa
1 5 10

<210> SEQ ID NO 305
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: consensus sequence derived from M31 B01 CDR L1
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: X = T or S

<400> SEQUENCE: 305

Xaa Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Val Val His
1 5 10

<210> SEQ ID NO 306
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: consensus sequence derived from M31 B01 CDR L2
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: X = K or Q

<400> SEQUENCE: 306

Asp Asn Asn Xaa Arg Pro Ser
1 5

-continued

```

<210> SEQ ID NO 307
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: consensus sequence derived from M31 B01 CDR L3
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: X = W, E, F or Y
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: X = R, S or M
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: X = N, K or S
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: X = G or R
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: X = P or A

```

```

<400> SEQUENCE: 307

```

```

Cys Ala Ala Xaa Asp Asp Xaa Leu Xaa Xaa Xaa Val
1          5          10

```

```

<210> SEQ ID NO 308
<211> LENGTH: 351
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 308

```

```

gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg      60
agctgcgcgg cgagcggggt tacctttagc aacgcgtgga tgagctgggt gcgccaggcg      120
ccggggaaa ggcctggagtg ggtgagctat attagcagca gcggggagcac catttattat      180
gcggatagcg tgaaagggcg ctttaccatt agccgcgata acagcaaaaa caccctgtat      240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gcgcgagggg      300
ctgtgggcgt ttgataaatg ggggcagggg accctggtga ccgtgaccag c              351

```

```

<210> SEQ ID NO 309
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 309

```

```

cagagcgtgc tgaccagcc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt      60
agctgcaccg ggagcagcag caacattggg gcggggtatg tggtcattg gtatcagcag      120
ctgccgggga ccgcgccgaa actgctgatt tatgataaca acaaacgccc gagcgggggtg      180
ccgatcgct ttagcgggag caaaagcggg accagcgcga gcctggcgat tagcgggctg      240
cgcagcgagg atgaggcgga ttattattgc gcggcgatg atgatagcct gaaagggccg      300
gtgtttgggg gggggaccaa actgaccgtg ctggggcag              339

```

-continued

<210> SEQ ID NO 310
<211> LENGTH: 351
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 310

```
gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg      60
agctgcgcgg cgagcggggt taccttttagc aacgcgtgga tgagctgggt gcgccaggcg      120
ccggggaaaag ggctggagtg ggtgagctat attagcagca gcgggagcac cattttattat      180
gcggatagcg tgaaaggggc ctttaccatt agccgcgata acagcaaaaa caccctgtat      240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gcgcgagggg      300
ctgtgggcgt ttgataaatg ggggcagggg accctggtga ccgtgaccag c                351
```

<210> SEQ ID NO 311
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 311

```
cagagcgtgc tgaccagacc gccgagcgtg agcggggcgc cggggcagcg cgtgaccatt      60
agctgcaccg ggagcagcag caacattggg gcggggtatg tgggtcattg gtatcagcag      120
ctgccgggga ccgcgccgaa actgctgatt tatgataaca acaaacgccc gagcgggggtg      180
ccgcatcgct ttagcgggag caaaagcggg accagcgcga gcctggcgat tagcgggctg      240
cgcagcgagg atgaggcgga ttattattgc gcggcggttg atgatagcct gaacggggccg      300
gtgtttgggg gggggaccaa actgaccgtg ctggggcag                339
```

<210> SEQ ID NO 312
<211> LENGTH: 351
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 312

```
gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg      60
agctgcgcgg cgagcggggt taccttttagc aacgcgtgga tgagctgggt gcgccaggcg      120
ccggggaaaag ggctggagtg ggtgagctat attagcagca gcgggagcac cattttattat      180
gcggatagcg tgaaaggggc ctttaccatt agccgcgata acagcaaaaa caccctgtat      240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gcgcgagggg      300
ctgtgggcgt ttgataaatg ggggcagggg accctggtga ccgtgaccag c                351
```

<210> SEQ ID NO 313
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 313

```
cagagcgtgc tgaccagacc gccgagcgtg agcggggcgc cggggcagcg cgtgaccatt      60
```

-continued

agctgcaccg ggagcagcag caacattggg gcgggggatg tgggtgcattg gtatcagcag	120
ctgccgggga ccgcgccgaa actgctgatt tatgataaca acaaacgccc gagcgggggtg	180
ccggatcgct ttagcgggag caaaagcggg accagcgcgga gcctggcgat tagcggggctg	240
cgcagcgagg atgaggcgga ttattattgc gcggcggttg atgatagcct gaacggggccg	300
gtgtttgggg gggggaccaa actgaccgtg ctggggcag	339

<210> SEQ ID NO 314
 <211> LENGTH: 351
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 314

gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg	60
agctgcgcgg cgagcgggtt taccttttagc agcgcgtgga tgagctgggt gcgccaggcg	120
ccgggaaaag ggctggagtg ggtgagctat attagcagca gcgggagcag cacctattat	180
gcggatagcg tgaaaggcg ctttaccatt agccgcgata acagcaaaaa caccctgtat	240
ctgcagatga acagcctgcg cgcggaggat accgcgggtg attattgcgc gcgcgagggg	300
ctgtgggcgt ttgattggtg ggggcagggg accctggtga ccgtgaccag c	351

<210> SEQ ID NO 315
 <211> LENGTH: 339
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 315

cagagcgtgc tgaccagacc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt	60
agctgcagcg ggagcagcag caacattggg gcgggggatg tgggtgcattg gtatcagcag	120
ctgccgggga ccgcgccgaa actgctgatt tatgataaca accagcgcgc gagcgggggtg	180
ccggatcgct ttagcgggag caaaagcggg accagcgcgga gcctggcgat tagcggggctg	240
cgcagcgagg atgaggcgga ttattattgc gcggcgatg atgatagcct gagcggggccg	300
gtgtttgggg gggggaccaa actgaccgtg ctggggcag	339

<210> SEQ ID NO 316
 <211> LENGTH: 351
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 316

gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg	60
agctgcgcgg cgagcgggtt taccttttagc aacgcgtgga tgagctgggt gcgccaggcg	120
ccgggaaaag ggctggagtg ggtgagctat attagcagca gcgggagcac catttattat	180
gcggatagcg tgaaaggcg ctttaccatt agccgcgata acagcaaaaa caccctgtat	240
ctgcagatga acagcctgcg cgcggaggat accgcgggtg attattgcgc gcgcgagggg	300
ctgtgggcgt ttgattattg ggggcagggg accctggtga ccgtgaccag c	351

-continued

<210> SEQ ID NO 317
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 317

cagagcgtgc tgaccagccc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt 60
agctgcagcg ggagcagcag caacattggg gcggggtatg tggcgcattg gtatcagcag 120
ctgccgggga ccgcgccgaa actgctgatt tatgataaca accagcgcgc gagcgggggtg 180
ccggatcgct ttagcgggag caaaagcggg accagcgcga gcctggcgat tagcgggctg 240
cgagcagagg atgaggcgga ttattattgc gcggcgatg atgatgcct gaacggggccg 300
gtgtttgggg gggggaccaa actgaccgtg ctggggcag 339

<210> SEQ ID NO 318
<211> LENGTH: 351
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 318

gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg 60
agctgcgcgg cgagcgggtt tacctttagc agcgcgtgga tgagctgggt gcgccaggcg 120
ccggggaag ggctggagtg ggtgagctat attagcagca gcgggagcag cacctattat 180
gcgatagcgc tgaaggggcg ctttaccatt agcgcgcgata acagcaaaaa caccctgtat 240
ctgcagatga acagcctcgc cgcggaggat accgcggtgt attattgcgc gcgcgagggg 300
ctgtgggcgt ttgattattg ggggcagggg accctggtga ccgtgagcag c 351

<210> SEQ ID NO 319
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 319

gagagcgtgc tgaccagccc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt 60
agctgcagcg ggagcagcag caacattggg gcggggtatg tggcgcattg gtatcagcag 120
ctgccgggga ccgcgccgaa actgctgatt tatgataaca accagcgcgc gagcgggggtg 180
ccggatcgct ttagcgggag caaaagcggg accagcgcga gcctggcgat tagcgggctg 240
cgagcagagg atgaggcgga ttattattgc gcggcggtgg atgatgcct gaacggggccg 300
gtgtttgggg gggggaccaa actgaccgtg ctggggcag 339

<210> SEQ ID NO 320
<211> LENGTH: 351
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 320

gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg 60

-continued

```

agctgcgcgg cgagcggggtt tacctttagc aacgcgtgga tgagctgggt gcgccaggcg 120
ccggggaaaag ggctggagtg ggtgagctat attagcagca gcgggagcac catttattat 180
gcggatagcg tgaaggggcg ctttaccatt agccgcgata acagcaaaaa caccctgtat 240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gcgcgagggg 300
ctgtgggcgt ttgattattg ggggcagggg accctggtga ccgtgagcag c 351

```

```

<210> SEQ ID NO 321
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 321

```

```

gagagcgtgc tgaccagcc gccgagcgtg agcggggcgc cggggcagcg cgtgaccatt 60
agctgcaccg ggagcagcag caacattggg gcggggtatg tggtcattg gtatcagcag 120
ctgccgggga ccgcgccgaa actgctgatt tatgataaca acaaacgccc gagcggggtg 180
ccggatcgct ttagcgggag caaaagcggg accagcgcga gcctggcgat tagcgggctg 240
cgcagcgagg atgaggcgga ttattattgc gcggcgatg atgatcgct gaacggggccg 300
gtgtttgggg gggggaccaa actgaccgtg ctggggcag 339

```

```

<210> SEQ ID NO 322
<211> LENGTH: 351
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 322

```

```

gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg 60
agctgcgcgg cgagcggggtt tacctttagc agcgcgtgga tgagctgggt gcgccaggcg 120
ccggggaaaag ggctggagtg ggtgagctat attagcagca gcgggagcag caccattat 180
gcggatagcg tgaaggggcg ctttaccatt agccgcgata acagcaaaaa caccctgtat 240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gcgcgagggg 300
ctgtgggcgt ttgatgggtg ggggcagggg accctggtga ccgtgagcag c 351

```

```

<210> SEQ ID NO 323
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 323

```

```

cagagcgtgc tgaccagcc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt 60
agctgcagcg ggagcagcag caacattggg gcggggtatg tggtcattg gtatcagcag 120
ctgccgggga ccgcgccgaa actgctgatt tatgataaca accagcgcgc gagcggggtg 180
ccggatcgct ttagcgggag caaaagcggg accagcgcga gcctggcgat tagcgggctg 240
cgcagcgagg atgaggcgga ttattattgc gcggcgatg atgatagcct gaaccgcccc 300
gtgtttgggg gggggaccaa actgaccgtg ctggggcag 339

```

-continued

<210> SEQ ID NO 324

<211> LENGTH: 351

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 324

```
gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg    60
agctgcgcgg cgagcggggt taccttttagc aacgcgtgga tgagctgggt gcgccaggcg    120
ccggggaaaag ggctggagtg ggtgagctat attagcagca gcgggagcac catttattat    180
gcggatagcg tgaaaggggc ctttaccatt agccgcgata acagcaaaaa caccctgtat    240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gcgcgagggg    300
ctgtgggcgt ttgattattg ggggcagggg accctggtga ccgtgagcag c              351
```

<210> SEQ ID NO 325

<211> LENGTH: 339

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 325

```
cagagcgtgc tgaccagacc gccgagcgtg agcggggcgc cggggcagcg cgtgaccatt    60
agctgcaccg ggagcagcag caacattggg gcggggtatg tgggtcattg gtatcagcag    120
ctgccgggga ccgcgccgaa actgctgatt tatgataaca acaaacgccc gagcgggggtg    180
ccgcatcgct ttagcgggag caaaagcggg accagcgcga gcctggcgat tagcgggctg    240
cgcagcgagg atgaggcgga ttattattgc gcggcggttg atgatagcct gaacggggccg    300
gtgtttgggg gggggaccaa actgaccgtg ctggggcag              339
```

<210> SEQ ID NO 326

<211> LENGTH: 351

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 326

```
gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg    60
agctgcgcgg cgagcggggt taccttttagc agcgcgtgga tgagctgggt gcgccaggcg    120
ccggggaaaag ggctggagtg ggtgagctat attagcagca gcgggagcag cacctattat    180
gcggatagcg tgaaaggggc ctttaccatt agccgcgata acagcaaaaa caccctgtat    240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gcgcgagggg    300
ctgtgggcgt ttgataaatg ggggcagggg accctggtga ccgtgaccag c              351
```

<210> SEQ ID NO 327

<211> LENGTH: 339

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 327

```
cagagcgtgc tgaccagacc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt    60
```

-continued

agctgcaccg ggagcagcag caacattggg gcggggtatg tgggtgcattg gtatcagcag	120
ctgccgggga ccgcgccgaa actgctgatt tatgataaca acaaacgccc gagcggggtg	180
ccggatcgct ttagcgggag caaaagcggg accagcgcga gcctggcgat tagcgggctg	240
cgcagcgagg atgaggcgga ttattattgc gcggcgtttg atgatagcct gaacggggccg	300
gtgtttgggg gggggaccaa actgaccgtg ctggggcag	339

<210> SEQ ID NO 328
 <211> LENGTH: 351
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 328

gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg	60
agctgcgcgg cgagcgggtt taccttttagc agcgcgtgga tgagctgggt gcgccaggcg	120
ccgggaaaag ggctggagtg ggtgagctat attagcagca gcgggagcag cacctattat	180
gcggatagcg tgaaaggcg ctttaccatt agccgcgata acagcaaaaa caccctgtat	240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gcgcgagggg	300
ctgtgggcgt ttgataaatg ggggcagggg accctggtga ccgtgagcag c	351

<210> SEQ ID NO 329
 <211> LENGTH: 339
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a

<400> SEQUENCE: 329

cagagcgtgc tgaccagcc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt	60
agctgcagcg ggagcagcag caacattggg gcggggtatg tgggtgcattg gtatcagcag	120
ctgccgggga ccgcgccgaa actgctgatt tatgataaca accagcgcgc gagcggggtg	180
ccggatcgct ttagcgggag caaaagcggg accagcgcga gcctggcgat tagcgggctg	240
cgcagcgagg atgaggcgga ttattattgc gcggcgtttg atgatagcct gaacggggccg	300
gtgtttgggg gggggaccaa actgaccgtg ctggggcag	339

<210> SEQ ID NO 330
 <211> LENGTH: 351
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 330

gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg	60
agctgcgcgg cgagcgggtt taccttttagc agcgcgtgga tgagctgggt gcgccaggcg	120
ccgggaaaag ggctggagtg ggtgagctat attagcagca gcgggagcag cacctattat	180
gcggatagcg tgaaaggcg ctttaccatt agccgcgata acagcaaaaa caccctgtat	240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gcgcgagggg	300
ctgtgggcgt ttgataaatg ggggcagggg accctggtga ccgtgagcag c	351

-continued

<210> SEQ ID NO 331
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 331

cagagcgtgc tgaccagacc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt	60
agctgcagcg ggagcagcag caacattggg gcggggtatg tggcgcattg gtatcagcag	120
ctgccgggga ccgcgccgaa actgctgatt tatgataaca accagcgcgc gagcgggggtg	180
ccggatcgct ttagcgggag caaaagcggg accagcgcga gcctggcgat tagcgggctg	240
cgcagcgagg atgagcgga ttattattgc gcggcggttg atgatcgcc gagcggggccg	300
gtgtttgggg gggggaccaa actgaccgtg ctggggcag	339

<210> SEQ ID NO 332
<211> LENGTH: 366
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 332

gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg	60
agctgcgcgg cgagcgggtt tacctttagc gattatcaga tgacctggat tcgccagacc	120
ccggggaag ggctggagtg ggtgagcggg gtgagctgga acggggcgcg caccattat	180
gcgatagcg tgaaggcg ctttaccatt agcgcgata acagcaaaaa caccctgtat	240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gaaaggggat	300
tatctggtgt atagcagcta ttattttaaa agctgggggc aggggaccct ggtgaccgtg	360
accagc	366

<210> SEQ ID NO 333
<211> LENGTH: 336
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 333

cagagcgtgc tgaccagacc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt	60
agctgcagcg ggagcagcag caacgtggg agcaaccgg tgaactggta tcagcagctg	120
ccggggaccg cgcgaaact gctgatttat cgcaacaacc agcggccgag cggggtgccg	180
gatcgcttta gggggagcaa aagcgggacc agcgcgagcc tggcgattag cgggctgcgc	240
agcgaggatg aggcggatta ttattgcgcg gcgtgggatg atcgccgaa cgggtggggg	300
tttggggggg ggaccaaact gaccgtgctg gggcag	336

<210> SEQ ID NO 334
<211> LENGTH: 363
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 334

-continued

gagcagctgc tggagagcgg gggggggctg gtgcagccgg gggggagcct gcgcctgagc	60
tgcgcggcga gcgggtttac ctttagcagc taccagatga cctggattcg ccagaccccg	120
gggaaagggc tggagtgggt gagcgggatt agctggaacg gggggagcac ccattatgcg	180
gatagcgtga aagggcgctt taccattagc cgcgataaca gcaaaaacac cctgtatctg	240
cagatgaaca gcctgcgcgc ggaggatacc gcggtgtatt attgcgcgaa aggggattat	300
ctggtgtata gcgcgtatta ttttgatagc tgggggcagg ggaccctggt gaccgtgagc	360
agc	363

<210> SEQ ID NO 335
 <211> LENGTH: 333
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 335

caggtgctga cccagccgcc gagcgcgagc gggaccccg ggcagcgcgt gaccattagc	60
tgagcggga gcagcagcaa cattgggagc aaccgggtga actggtatca gcagctgccg	120
gggacccgcg cgaaactgct gatttatcgc aacaaccagc gcccgagcgg ggtgccggat	180
cgcttttagcg ggagcaaaag cgggaccagc gcgagcctgg cgattagcgg gctgcgcagc	240
gaggtgaggg cggattatta ttgcgcggcg tgggatgac gcctgaacgg gtggggggtt	300
ggggggggga ccaaactgac cgtgctgggg cag	333

<210> SEQ ID NO 336
 <211> LENGTH: 366
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 336

gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg	60
agctgcgcgg cgagcgggtt tacctttagc gattatcaga tgacctggat tcgccagacc	120
ccggggaaag ggctggagtg ggtgagcggg attagctgga acggggggag caccattat	180
gcgtagatcg tgaagggcg ctttaccatt agccgcgata acagcaaaaa caccctgtat	240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gaaaggggat	300
tatctggtgt atagcagcta ttattttaaa tattgggggc aggggaccct ggtgaccgtg	360
agcagc	366

<210> SEQ ID NO 337
 <211> LENGTH: 336
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 337

cagagcgtgc tgaccagcc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt	60
agctgcagcg ggagcagcag caacattggg agcaaccgg tgaactggta tcagcagctg	120
ccggggaccg cgccgaaact gctgatttat cgcaacaacc agcggccgag cggggtgccg	180
gatcgcttta gggggagcaa aagcgggacc agcgcgagcc tggcgattag cgggctgcgc	240

-continued

agcgaggatg aggcggatta ttattgcgcg gcgtgggatg atcgccctgaa cgggtgggcg 300
tttggggggg ggaccaaact gaccgtgctg gggcag 336

<210> SEQ ID NO 338
<211> LENGTH: 366
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 338

gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg 60
agctgcgcgg cgagcgggtt taccttttagc gattatcaga tgacctggat tcgccagacc 120
ccggggaaaag ggctggagtg ggtgagcggg attagctgga acgggggggag caccattat 180
gcggatagcg tgaaaggcg ctttaccatt agccgcgata acagcaaaaa caccctgtat 240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gaaaggggat 300
tatctggtgt atagcagcta ttattttaaa agctgggggc aggggaccct ggtgaccgtg 360
agcagc 366

<210> SEQ ID NO 339
<211> LENGTH: 336
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 339

cagagcgtgc tgaccagacc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt 60
agctgcagcg ggagcagcag caacattggg agcaaccggg tgaactggta tcagcagctg 120
ccggggaccg cgccgaaact gctgatttat cgcaacaacc agcggccgag cggggtgccg 180
gatcgcttta gcgggagcaa aagcgggacc agcgcgagcc tggcgattag cgggctgcgc 240
agcgaggatg aggcggatta ttattgcgcg gcgtgggatg atagccctgaa cgggtggggg 300
tttggggggg ggaccaaact gaccgtgctg gggcag 336

<210> SEQ ID NO 340
<211> LENGTH: 366
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 340

gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg 60
agctgcgcgg cgagcgggtt taccttttagc gattatcaga tgacctggat tcgccagacc 120
ccggggaaaag ggctggagtg ggtgagcggg attagctgga acgggggggag caccattat 180
gcggatagcg tgaaaggcg ctttaccatt agccgcgata acagcaaaaa caccctgtat 240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gaaaggggat 300
tatctggtgt ataaaagcta ttattttaaa agctgggggc aggggaccct ggtgaccgtg 360
agcagc 366

<210> SEQ ID NO 341

-continued

<211> LENGTH: 336
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 341

```
cagagcgtgc tgaccagacc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt    60
agctgcagcg ggagcagcag caacattggg agcaaccgga tgaactggta tcagcagctg    120
ccggggaccc cgccgaaact gctgatttat cgcaacaacc agcgcccgag cggggtgccg    180
gatcgcttta ggggagcaa aagcgggacc agcgcgagcc tggcgattag cgggctgcgc    240
agcgaggatg aggcggatta ttattgcgcg gcgtgggatg atcgctgag cgggtggggg    300
tttggggggg ggaccaaact gaccgtgctg gggcag                                336
```

<210> SEQ ID NO 342
<211> LENGTH: 366
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 342

```
gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg    60
agctgcgcgg cgagcgggtt tacctttagc gattatcaga tgacctggat tcgccagacc    120
ccggggaaaag ggctggagtg ggtgagcggg attagctgga acggggggag caccattat    180
gcggatagcg tgaaagggag ctttaccatt agccgcgata acagcaaaaa caccctgtat    240
ctgcagatga acagcctgcg cgccgaggat accgcggtgt attattgcgc gaaaggggat    300
tatctggtgt atagcagcta ttattttaa agctgggggc aggggaccct ggtgaccgtg    360
agcagc                                366
```

<210> SEQ ID NO 343
<211> LENGTH: 336
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 343

```
cagagcgtgc tgaccagacc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt    60
agctgcagcg ggagcagcag caacattggg agcaaccgga tgaactggta tcagcagctg    120
ccggggaccc cgccgaaact gctgatttat cgcaacaacc agcgcccgag cggggtgccg    180
gatcgcttta ggggagcaa aagcgggacc agcgcgagcc tggcgattag cgggctgcgc    240
agcgaggatg aggcggatta ttattgcgcg gcgtgggatg atcgctgag cgggtggggg    300
tttggggggg ggaccaaact gaccgtgctg gggcag                                336
```

<210> SEQ ID NO 344
<211> LENGTH: 366
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

<400> SEQUENCE: 344

```
gaggtgcagc tgctggagag cggggggggg ctggtgcagc cgggggggag cctgcgcctg    60
```

-continued

```

agctgcgcgg cgagcggggt taccttttagc gattatcaga tgacctggat tcgccagacc 120
ccgggggaaag ggctggagtg ggtgagcggg attagctgga acgggggggag caccattat 180
gcggatagcg tgaaggggcg ctttaccatt agccgcgata acagcaaaaa caccctgtat 240
ctgcagatga acagcctgcg cgcggaggat accgcggtgt attattgcgc gaaaggggat 300
tatctggtgt ataaaagcta ttattttaaa agctgggggc aggggaccct ggtgaccgtg 360
agcagc 366

```

```

<210> SEQ ID NO 345
<211> LENGTH: 336
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C4.4a binder

```

```

<400> SEQUENCE: 345

```

```

cagagcgtgc tgaccagacc gccgagcgcg agcgggaccc cggggcagcg cgtgaccatt 60
agctgcagcg ggagcagcag caacattggg agcaaccggg tgaactggtg tcagcagctg 120
ccggggaccg cgccgaaact gctgatttat cgcaacaacc agcgcccagag cggggtgccg 180
gatcgcttta gcgggagcaa aagcgggacc agcgcgagcc tggcgattag cgggctgcgc 240
agcagagatg aggcggatta ttattgcgcg gcgtgggatg atagcctgaa cgggtggggg 300
tttggggggg ggaccaaact gaccgtgctg gggcag 336

```

```

<210> SEQ ID NO 346
<211> LENGTH: 217
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

```

```

<400> SEQUENCE: 346

```

```

Asp Ile Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1           5           10          15
Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
20          25          30
Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35          40          45
Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
50          55          60
Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65          70          75          80
Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg
85          90          95
Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100         105         110
Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu
115         120         125
Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe
130         135         140
Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Gly Asp Ser Ser Pro Val
145         150         155         160
Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys
165         170         175
Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser

```

180							185							190						
His	Arg	Ser	Tyr	Ser	Cys	Gln	Val	Thr	His	Glu	Gly	Ser	Thr	Val	Glu					
195							200							205						
Lys	Thr	Val	Ala	Pro	Thr	Glu	Cys	Ser												
210							215													
<210> SEQ ID NO 347																				
<211> LENGTH: 447																				
<212> TYPE: PRT																				
<213> ORGANISM: Homo Sapiens																				
<400> SEQUENCE: 347																				
Gln	Val	Glu	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly					
1				5					10					15						
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asn	Ala					
			20					25					30							
Trp	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val					
		35					40					45								
Ser	Tyr	Ile	Ser	Ser	Ser	Gly	Ser	Thr	Ile	Tyr	Tyr	Ala	Asp	Ser	Val					
	50					55					60									
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr					
65					70				75					80						
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys					
				85					90					95						
Ala	Arg	Glu	Gly	Leu	Trp	Ala	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu					
			100					105					110							
Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu					
		115					120					125								
Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys					
	130					135				140										
Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser					
145					150					155				160						
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser					
				165					170					175						
Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser					
			180					185					190							
Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn					
		195					200					205								
Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His					
	210					215					220									
Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val					
225					230					235					240					
Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr					
				245					250					255						
Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu					
			260					265					270							
Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys					
		275					280					285								
Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser					
	290					295					300									
Val	Leu	Thr	Val																	

-continued

325					330					335					
Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro
			340					345					350		
Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu
			355				360					365			
Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn
		370					375					380			
Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser
		385					390					395			400
Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg
			405						410					415	
Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu
			420					425					430		
His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys	
			435				440					445			

<210> SEQ ID NO 348

<211> LENGTH: 217

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 348

Glu	Ser	Val	Leu	Thr	Gln	Pro	Pro	Ser	Ala	Ser	Gly	Thr	Pro	Gly	Gln
1				5					10					15	
Arg	Val	Thr	Ile	Ser	Cys	Thr	Gly	Ser	Ser	Ser	Asn	Ile	Gly	Ala	Gly
			20					25					30		
Tyr	Val	Val	His	Trp	Tyr	Gln	Gln	Leu	Pro	Gly	Thr	Ala	Pro	Lys	Leu
			35				40					45			
Leu	Ile	Tyr	Asp	Asn	Asn	Lys	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	Phe
		50				55					60				
Ser	Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Ser	Gly	Leu
		65			70					75				80	
Arg	Ser	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Ala	Ala	Trp	Asp	Asp	Arg
			85						90					95	
Leu	Asn	Gly	Pro	Val	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly
			100					105					110		
Gln	Pro	Lys	Ala	Ala	Pro	Ser	Val	Thr	Leu	Phe	Pro	Pro	Ser	Ser	Glu
		115					120					125			
Glu	Leu	Gln	Ala	Asn	Lys	Ala	Thr	Leu	Val	Cys	Leu	Ile	Ser	Asp	Phe
		130				135					140				
Tyr	Pro	Gly	Ala	Val	Thr	Val	Ala	Trp	Lys	Ala	Asp	Ser	Ser	Pro	Val
		145				150				155				160	
Lys	Ala	Gly	Val	Glu	Thr	Thr	Thr	Pro	Ser	Lys	Gln	Ser	Asn	Asn	Lys
			165						170					175	
Tyr	Ala	Ala	Ser	Ser	Tyr	Leu	Ser	Leu	Thr	Pro	Glu	Gln	Trp	Lys	Ser
			180					185					190		
His	Arg	Ser	Tyr	Ser	Cys	Gln	Val	Thr	His	Glu	Gly	Ser	Thr	Val	Glu
		195					200					205			
Lys	Thr	Val	Ala	Pro	Thr	Glu	Cys	Ser							
		210					215								

<210> SEQ ID NO 349

<211> LENGTH: 446

<212> TYPE: PRT

-continued

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 349

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
 20 25 30
 Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr Trp Gly Gln Gly Thr Leu
 100 105 110
 Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu
 115 120 125
 Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys
 130 135 140
 Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
 145 150 155 160
 Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
 165 170 175
 Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser
 180 185 190
 Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn
 195 200 205
 Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His
 210 215 220
 Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val
 225 230 235 240
 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
 245 250 255
 Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
 260 265 270
 Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
 275 280 285
 Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
 290 295 300
 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 305 310 315 320
 Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
 325 330 335
 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 340 345 350
 Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 355 360 365
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 370 375 380
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser

-continued

```

385              390              395              400
Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
              405              410              415

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
              420              425              430

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
              435              440              445

```

```

<210> SEQ ID NO 350
<211> LENGTH: 217
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

```

```

<400> SEQUENCE: 350

```

```

Glu Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1              5              10              15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly
              20              25              30

Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
              35              40              45

Leu Ile Tyr Asp Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe
              50              55              60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65              70              75              80

Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg
              85              90              95

Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
              100             105             110

Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu
              115             120             125

Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe
              130             135             140

Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val
145             150             155             160

Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys
              165             170             175

Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser
              180             185             190

His Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu
              195             200             205

Lys Thr Val Ala Pro Thr Glu Cys Ser
              210             215

```

```

<210> SEQ ID NO 351
<211> LENGTH: 446
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

```

```

<400> SEQUENCE: 351

```

```

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1              5              10              15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Ala
              20              25              30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
              35              40              45

```

Ser	Tyr	Ile	Ser	Ser	Ser	Gly	Ser	Ser	Thr	Tyr	Tyr	Ala	Asp	Ser	Val
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
Ala	Arg	Glu	Gly	Leu	Trp	Ala	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu
Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu
Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys
Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser
Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser
Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn
Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His
Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val
Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr
Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu
Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys
Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser
Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys
Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile
Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro
Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu
Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn
Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser
Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg
Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu
His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly		

-continued

```

<210> SEQ ID NO 352
<211> LENGTH: 217
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 352
Glu Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
1          5          10          15
Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
20          25          30
Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35          40          45
Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
50          55          60
Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65          70          75          80
Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg
85          90          95
Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100         105         110
Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu
115         120         125
Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe
130         135         140
Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val
145         150         155         160
Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys
165         170         175
Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser
180         185         190
His Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu
195         200         205
Lys Thr Val Ala Pro Thr Glu Cys Ser
210         215

```

```

<210> SEQ ID NO 353
<211> LENGTH: 446
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 353
Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1          5          10          15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
20          25          30
Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35          40          45
Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
50          55          60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65          70          75          80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85          90          95
Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr Trp Gly Gln Gly Thr Leu
100         105         110

```

```
<210> SEQ ID NO 354
<211> LENGTH: 217
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 354
```

Glu Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30

-continued

Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
 35 40 45
 Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
 50 55 60
 Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
 65 70 75 80
 Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Tyr Asp Asp Arg
 85 90 95
 Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 100 105 110
 Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu
 115 120 125
 Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe
 130 135 140
 Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val
 145 150 155 160
 Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys
 165 170 175
 Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser
 180 185 190
 His Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu
 195 200 205
 Lys Thr Val Ala Pro Thr Glu Cys Ser
 210 215

<210> SEQ ID NO 355

<211> LENGTH: 446

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 355

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
 20 25 30
 Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr Trp Gly Gln Gly Thr Leu
 100 105 110
 Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu
 115 120 125
 Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys
 130 135 140
 Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
 145 150 155 160
 Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
 165 170 175

-continued

Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser
 180 185 190
 Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn
 195 200 205
 Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His
 210 215 220
 Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val
 225 230 235 240
 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
 245 250 255
 Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
 260 265 270
 Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
 275 280 285
 Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
 290 295 300
 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 305 310 315 320
 Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
 325 330 335
 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 340 345 350
 Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 355 360 365
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 370 375 380
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 385 390 395 400
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
 405 410 415
 Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
 420 425 430
 His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440 445

<210> SEQ ID NO 356

<211> LENGTH: 217

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 356

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
 1 5 10 15
 Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
 20 25 30
 Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
 35 40 45
 Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
 50 55 60
 Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
 65 70 75 80
 Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Tyr Asp Asp Ser
 85 90 95

-continued

Leu Ser Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 100 105 110
 Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu
 115 120 125
 Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe
 130 135 140
 Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val
 145 150 155 160
 Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys
 165 170 175
 Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser
 180 185 190
 His Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu
 195 200 205
 Lys Thr Val Ala Pro Thr Glu Cys Ser
 210 215

<210> SEQ ID NO 357

<211> LENGTH: 446

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 357

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
 20 25 30
 Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Glu Gly Leu Trp Ala Phe Asp Tyr Trp Gly Gln Gly Thr Leu
 100 105 110
 Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu
 115 120 125
 Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys
 130 135 140
 Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
 145 150 155 160
 Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
 165 170 175
 Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser
 180 185 190
 Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn
 195 200 205
 Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His
 210 215 220
 Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val
 225 230 235 240

-continued

```

Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
      245                      250                255

Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
      260                      265                270

Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
      275                      280                285

Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
      290                      295                300

Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
      305                      310                315                320

Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
      325                      330                335

Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
      340                      345                350

Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
      355                      360                365

Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
      370                      375                380

Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
      385                      390                395                400

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
      405                      410                415

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
      420                      425                430

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
      435                      440                445

```

<210> SEQ ID NO 358

<211> LENGTH: 217

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 358

```

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1      5      10      15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ala Gly
      20      25      30

Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
      35      40      45

Leu Ile Tyr Asp Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe
      50      55      60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
      65      70      75      80

Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Tyr Asp Asp Ser
      85      90      95

Leu Asn Arg Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
      100     105     110

Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu
      115     120     125

Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe
      130     135     140

Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val
      145     150     155     160

```



```
<210> SEQ ID NO 359
<211> LENGTH: 446
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 359
```

Glu 1	Val	Gln	Leu	Leu 5	Glu	Ser	Gly	Gly 10	Leu	Val	Gln	Pro	Gly 15	Gly	
Ser	Leu	Arg	Leu 20	Ser	Cys	Ala	Ala	Ser 25	Gly	Phe	Thr	Phe	Ser 30	Ser	Ala
Trp	Met	Ser 35	Trp	Val	Arg	Gln	Ala 40	Pro	Gly	Lys	Gly	Leu 45	Glu	Trp	Val
Ser	Tyr 50	Ile	Ser	Ser	Ser	Gly 55	Ser	Ser	Thr	Tyr	Tyr 60	Ala	Asp	Ser	Val
Lys 65	Gly	Arg	Phe	Thr	Ile 70	Ser	Arg	Asp	Asn	Ser 75	Lys	Asn	Thr	Leu	Tyr 80
Leu	Gln	Met	Asn	Ser 85	Leu	Arg	Ala	Glu	Asp 90	Thr	Ala	Val	Tyr	Tyr 95	Cys
Ala	Arg	Glu	Gly 100	Leu	Trp	Ala	Phe	Asp 105	Gly	Trp	Gly	Gln	Gly 110	Thr	Leu
Val	Thr	Val 115	Ser	Ser	Ala	Ser	Thr 120	Lys	Gly	Pro	Ser	Val 125	Phe	Pro	Leu
Ala	Pro 130	Ser	Ser	Lys	Ser	Thr 135	Ser	Gly	Gly	Thr	Ala 140	Ala	Leu	Gly	Cys
Leu 145	Val	Lys	Asp	Tyr	Phe 150	Pro	Glu	Pro	Val	Thr 155	Val	Ser	Trp	Asn	Ser 160
Gly	Ala	Leu	Thr	Ser 165	Gly	Val	His	Thr	Phe 170	Pro	Ala	Val	Leu	Gln	Ser 175
Ser	Gly	Leu 180	Tyr	Ser	Leu	Ser	Ser	Val 185	Val	Thr	Val	Pro	Ser 190	Ser	Ser
Leu	Gly	Thr 195	Gln	Thr	Tyr	Ile	Cys 200	Asn	Val	Asn	His	Lys 205	Pro	Ser	Asn
Thr	Lys 210	Val	Asp	Lys	Lys	Val 215	Glu	Pro	Lys	Ser	Cys 220	Asp	Lys	Thr	His
Thr 225	Cys	Pro	Pro	Cys	Pro 230	Ala	Pro	Glu	Leu	Leu 235	Gly	Gly	Pro	Ser	Val 240
Phe	Leu	Phe	Pro 245	Pro	Lys	Pro	Lys	Asp	Thr 250	Leu	Met	Ile	Ser	Arg 255	Thr
Pro	Glu	Val	Thr 260	Cys	Val	Val	Val	Asp 265	Val	Ser	His	Glu	Asp	Pro	Glu
Val	Lys	Phe 275	Asn	Trp	Tyr	Val	Asp 280	Gly	Val	Glu	Val	His 285	Asn	Ala	Lys
Thr	Lys 290	Pro	Arg	Glu	Glu	Gln 295	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser

-continued

```

Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
305                      310                      315                      320

Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
                      325                      330                      335

Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
                      340                      345                      350

Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
                      355                      360                      365

Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
370                      375                      380

Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
385                      390                      395                      400

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
                      405                      410                      415

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
420                      425                      430

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
435                      440                      445

```

<210> SEQ ID NO 360

<211> LENGTH: 217

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 360

```

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1                      5                      10                      15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
20                      25                      30

Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35                      40                      45

Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
50                      55                      60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
65                      70                      75                      80

Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Phe Asp Asp Arg
85                      90                      95

Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100                      105                      110

Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu
115                      120                      125

Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe
130                      135                      140

Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val
145                      150                      155                      160

Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys
165                      170                      175

Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser
180                      185                      190

His Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu
195                      200                      205

Lys Thr Val Ala Pro Thr Glu Cys Ser
210                      215

```

-continued

<210> SEQ ID NO 361

<211> LENGTH: 446

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 361

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
20 25 30
Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Arg Glu Gly Leu Trp Ala Phe Asp Lys Trp Gly Gln Gly Thr Leu
100 105 110
Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu
115 120 125
Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys
130 135 140
Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
145 150 155 160
Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
165 170 175
Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser
180 185 190
Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn
195 200 205
Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His
210 215 220
Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val
225 230 235 240
Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
245 250 255
Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
260 265 270
Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
275 280 285
Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
290 295 300
Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
305 310 315 320
Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
325 330 335
Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
340 345 350
Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
355 360 365

-continued

Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 370 375 380

Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 385 390 395 400

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
 405 410 415

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
 420 425 430

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440 445

<210> SEQ ID NO 362
 <211> LENGTH: 217
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 362

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
 1 5 10 15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
 20 25 30

Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
 35 40 45

Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
 50 55 60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
 65 70 75 80

Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Phe Asp Asp Ser
 85 90 95

Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 100 105 110

Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu
 115 120 125

Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe
 130 135 140

Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val
 145 150 155 160

Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys
 165 170 175

Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser
 180 185 190

His Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu
 195 200 205

Lys Thr Val Ala Pro Thr Glu Cys Ser
 210 215

<210> SEQ ID NO 363
 <211> LENGTH: 446
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 363

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

-continued

Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asn	Ala
			20					25					30		
Trp	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
	35						40					45			
Ser	Tyr	Ile	Ser	Ser	Ser	Gly	Ser	Thr	Ile	Tyr	Tyr	Ala	Asp	Ser	Val
	50					55					60				
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65					70					75					80
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
				85					90					95	
Ala	Arg	Glu	Gly	Leu	Trp	Ala	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu
		100						105					110		
Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu
		115					120					125			
Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys
	130					135					140				
Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser
145					150					155					160
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser
				165					170					175	
Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser
		180						185					190		
Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn
		195					200					205			
Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His
	210					215					220				
Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val
225					230					235					240
Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr
				245					250					255	
Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu
		260						265					270		
Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys
		275					280					285			
Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser
	290					295					300				
Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys
305					310					315					320
Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile
				325					330					335	
Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro
		340						345					350		
Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu
		355					360					365			
Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn
	370					375					380				
Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser
385					390					395					400
Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg
				405					410				415		
Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu
			420					425					430		

-continued

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440 445

<210> SEQ ID NO 364
 <211> LENGTH: 217
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 364

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
 1 5 10 15
 Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
 20 25 30
 Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
 35 40 45
 Leu Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Val Pro Asp Arg Phe
 50 55 60
 Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu
 65 70 75 80
 Arg Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Tyr Asp Asp Ser
 85 90 95
 Leu Ser Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 100 105 110
 Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu
 115 120 125
 Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe
 130 135 140
 Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val
 145 150 155 160
 Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys
 165 170 175
 Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser
 180 185 190
 His Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu
 195 200 205
 Lys Thr Val Ala Pro Thr Glu Cys Ser
 210 215

<210> SEQ ID NO 365
 <211> LENGTH: 446
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 365

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Ala
 20 25 30
 Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

-continued

Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	
Ala	Arg	Glu	Gly	Leu	Trp	Ala	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu
		100						105				110			
Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu
	115						120					125			
Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys
	130					135					140				
Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser
145					150					155				160	
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser
			165					170						175	
Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser
		180						185					190		
Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn
	195						200					205			
Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His
	210					215					220				
Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val
225					230					235					240
Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr
			245					250						255	
Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu
		260						265					270		
Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys
	275						280					285			
Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser
	290					295					300				
Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys
305					310					315					320
Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile
			325					330						335	
Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro
		340						345					350		
Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu
		355					360					365			
Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn
	370					375					380				
Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser
385					390					395					400
Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg
			405						410				415		
Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu
			420					425					430		
His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly		
	435						440					445			

<210> SEQ ID NO 366

<211> LENGTH: 217

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 366

-continued

Gln	Ser	Val	Leu	Thr	Gln	Pro	Pro	Ser	Ala	Ser	Gly	Thr	Pro	Gly	Gln	1	5	10	15
Arg	Val	Thr	Ile	Ser	Cys	Ser	Gly	Ser	Ser	Ser	Asn	Ile	Gly	Ala	Gly	20	25	30	
Tyr	Val	Val	His	Trp	Tyr	Gln	Gln	Leu	Pro	Gly	Thr	Ala	Pro	Lys	Leu	35	40	45	
Leu	Ile	Tyr	Asp	Asn	Asn	Gln	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	Phe	50	55	60	
Ser	Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Ser	Gly	Leu	65	70	75	80
Arg	Ser	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Ala	Ala	Phe	Asp	Asp	Ser	85	90	95	
Leu	Asn	Gly	Pro	Val	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly	100	105	110	
Gln	Pro	Lys	Ala	Ala	Pro	Ser	Val	Thr	Leu	Phe	Pro	Pro	Ser	Ser	Glu	115	120	125	
Glu	Leu	Gln	Ala	Asn	Lys	Ala	Thr	Leu	Val	Cys	Leu	Ile	Ser	Asp	Phe	130	135	140	
Tyr	Pro	Gly	Ala	Val	Thr	Val	Ala	Trp	Lys	Ala	Asp	Ser	Ser	Pro	Val	145	150	155	160
Lys	Ala	Gly	Val	Glu	Thr	Thr	Thr	Pro	Ser	Lys	Gln	Ser	Asn	Asn	Lys	165	170	175	
Tyr	Ala	Ala	Ser	Ser	Tyr	Leu	Ser	Leu	Thr	Pro	Glu	Gln	Trp	Lys	Ser	180	185	190	
His	Arg	Ser	Tyr	Ser	Cys	Gln	Val	Thr	His	Glu	Gly	Ser	Thr	Val	Glu	195	200	205	
Lys	Thr	Val	Ala	Pro	Thr	Glu	Cys	Ser	210	215									

<210> SEQ ID NO 367

<211> LENGTH: 446

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 367

Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	1	5	10	15
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Ser	Ala	20	25	30	
Trp	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	35	40	45	
Ser	Tyr	Ile	Ser	Ser	Ser	Gly	Ser	Ser	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	50	55	60	
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr	65	70	75	80
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	85	90	95	
Ala	Arg	Glu	Gly	Leu	Trp	Ala	Phe	Asp	Lys	Trp	Gly	Gln	Gly	Thr	Leu	100	105	110	
Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	115	120	125	
Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	130	135	140	

-continued

Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser
145					150					155					160
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser
				165					170					175	
Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser
			180					185					190		
Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn
		195					200					205			
Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His
	210					215					220				
Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val
225					230					235					240
Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr
				245					250					255	
Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu
			260					265					270		
Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys
		275					280					285			
Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser
	290					295					300				
Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys
305					310					315					320
Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile
				325					330					335	
Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro
		340						345					350		
Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu
		355					360					365			
Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn
	370					375					380				
Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser
385					390					395					400
Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg
			405						410				415		
Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu
			420					425					430		
His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly		
	435						440					445			

<210> SEQ ID NO 368

<211> LENGTH: 217

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 368

Gln	Ser	Val	Leu	Thr	Gln	Pro	Pro	Ser	Ala	Ser	Gly	Thr	Pro	Gly	Gln
1				5					10					15	
Arg	Val	Thr	Ile	Ser	Cys	Ser	Gly	Ser	Ser	Ser	Asn	Ile	Gly	Ala	Gly
		20					25					30			
Tyr	Val	Val	His	Trp	Tyr	Gln	Gln	Leu	Pro	Gly	Thr	Ala	Pro	Lys	Leu
	35					40					45				
Leu	Ile	Tyr	Asp	Asn	Asn	Gln	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	Phe
	50				55					60					

-continued

Ser	Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Ser	Gly	Leu
65					70					75					80
Arg	Ser	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Ala	Ala	Phe	Asp	Asp	Arg
			85						90					95	
Leu	Ser	Gly	Pro	Val	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly
			100					105					110		
Gln	Pro	Lys	Ala	Ala	Pro	Ser	Val	Thr	Leu	Phe	Pro	Pro	Ser	Ser	Glu
		115					120					125			
Glu	Leu	Gln	Ala	Asn	Lys	Ala	Thr	Leu	Val	Cys	Leu	Ile	Ser	Asp	Phe
	130						135				140				
Tyr	Pro	Gly	Ala	Val	Thr	Val	Ala	Trp	Lys	Ala	Asp	Ser	Ser	Pro	Val
145					150					155					160
Lys	Ala	Gly	Val	Glu	Thr	Thr	Thr	Pro	Ser	Lys	Gln	Ser	Asn	Asn	Lys
				165					170					175	
Tyr	Ala	Ala	Ser	Ser	Tyr	Leu	Ser	Leu	Thr	Pro	Glu	Gln	Trp	Lys	Ser
			180					185					190		
His	Arg	Ser	Tyr	Ser	Cys	Gln	Val	Thr	His	Glu	Gly	Ser	Thr	Val	Glu
		195					200					205			
Lys	Thr	Val	Ala	Pro	Thr	Glu	Cys	Ser							
	210					215									

<210> SEQ ID NO 369

<211> LENGTH: 446

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 369

Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly
1				5					10					15	
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Ser	Ala
			20					25					30		
Trp	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
	35					40						45			
Ser	Tyr	Ile	Ser	Ser	Ser	Gly	Ser	Ser	Thr	Tyr	Tyr	Ala	Asp	Ser	Val
	50					55				60					
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65				70					75					80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	
Ala	Arg	Glu	Gly	Leu	Trp	Ala	Phe	Asp	Lys	Trp	Gly	Gln	Gly	Thr	Leu
		100						105					110		
Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu
		115					120					125			
Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys
	130					135					140				
Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser
145				150					155					160	
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser
			165					170						175	
Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser
		180						185					190		
Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn
	195					200						205			

Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His
210						215					220				
Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val
225					230					235					240
Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr
				245					250					255	
Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu
			260					265					270		
Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys
		275					280					285			
Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser
	290					295					300				
Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys
305					310					315					320
Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile
				325					330					335	
Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro
			340					345					350		
Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu
		355					360					365			
Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn
	370					375					380				
Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser
385					390					395					400
Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg
			405						410					415	
Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu
			420					425					430		
His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly		
	435						440					445			
<210> SEQ ID NO 370															
<211> LENGTH: 216															
<212> TYPE: PRT															
<213> ORGANISM: Homo Sapiens															
<400> SEQUENCE: 370															
Asp	Ile	Val	Leu	Thr	Gln	Pro	Ser	Ala	Ser	Gly	Thr	Pro	Gly	Gln	
1				5				10					15		
Arg	Val	Thr	Ile	Ser	Cys	Ser	Gly	Ser	Ser	Asn	Val	Gly	Ser	Asn	
			20				25					30			
Pro	Val	Asn	Trp	Tyr	Gln	Gln	Leu	Pro	Gly	Thr	Ala	Pro	Lys	Leu	Leu
		35					40					45			
Ile	Tyr	Arg	Asn												

-continued

```

Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr
 130                      135                      140

Pro Gly Ala Val Thr Val Ala Trp Lys Gly Asp Ser Ser Pro Val Lys
145                      150                      155                      160

Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr
                      165                      170                      175

Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His
                      180                      185                      190

Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys
                      195                      200                      205

Thr Val Ala Pro Thr Glu Cys Ser
 210                      215

```

<210> SEQ ID NO 371

<211> LENGTH: 452

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 371

```

Gln Val Glu Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1                      5                      10                      15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
                      20                      25                      30

Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
 35                      40                      45

Ser Gly Val Ser Trp Asn Gly Ala Arg Thr His Tyr Ala Asp Ser Val
 50                      55                      60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65                      70                      75                      80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
                      85                      90                      95

Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ala Tyr Tyr Phe Asp Ser Trp
                      100                      105                      110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro
                      115                      120                      125

Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr
                      130                      135                      140

Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr
145                      150                      155                      160

Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro
                      165                      170                      175

Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr
                      180                      185                      190

Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn
                      195                      200                      205

His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser
                      210                      215                      220

Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu
225                      230                      235                      240

Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu
                      245                      250                      255

Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser
                      260                      265                      270

```

His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	
		275					280					285				
Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	
		290				295					300					
Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	
305					310					315						320
Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	
				325					330					335		
Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	
			340					345					350			
Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	
		355					360					365				
Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	
		370				375					380					
Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	
385					390					395						400
Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	
			405						410					415		
Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	
			420					425					430			
Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	
		435					440					445				
Ser	Pro	Gly	Lys													
		450														

```
<210> SEQ ID NO 372
<211> LENGTH: 216
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens
```

<400> SEQUENCE: 372

Glu 1	Ser	Val	Leu	Thr 5	Gln	Pro	Pro	Ser	Ala 10	Ser	Gly	Thr	Pro	Gly 15	Gln
Arg	Val	Thr	Ile 20	Ser	Cys	Ser	Gly	Ser 25	Ser	Ser	Asn	Val	Gly 30	Ser	Asn
Pro	Val	Asn	Trp	Tyr	Gln	Gln	Leu 40	Pro	Gly	Thr	Ala	Pro 45	Lys	Leu	Leu
Ile 50	Tyr	Arg	Asn	Asn	Gln	Arg 55	Pro	Ser	Gly	Val	Pro 60	Asp	Arg	Phe	Ser
Gly 65	Ser	Lys	Ser	Gly	Thr 70	Ser	Ala	Ser	Leu	Ala 75	Ile	Ser	Gly	Leu	Arg 80
Ser	Glu	Asp	Glu	Ala 85	Asp	Tyr	Tyr	Cys	Ala 90	Ala	Trp	Asp	Asp	Arg 95	Leu
Asn	Gly	Trp	Val 100	Phe	Gly	Gly	Gly	Thr 105	Lys	Leu	Thr	Val	Leu 110	Gly	Gln
Pro	Lys	Ala 115	Ala	Pro	Ser	Val	Thr 120	Leu	Phe	Pro	Pro	Ser 125	Ser	Glu	Glu
Leu 130	Gln	Ala	Asn	Lys	Ala	Thr 135	Leu	Val	Cys	Leu	Ile 140	Ser	Asp	Phe	Tyr
Pro 145	Gly	Ala	Val	Thr 150	Val	Ala	Trp	Lys	Ala	Asp 155	Ser	Ser	Pro	Val	Lys 160
Ala	Gly	Val	Glu	Thr 165	Thr	Thr	Pro	Ser	Lys 170	Gln	Ser	Asn	Asn	Lys	Tyr 175

-continued

Ala	Ala	Ser	Ser	Tyr	Leu	Ser	Leu	Thr	Pro	Glu	Gln	Trp	Lys	Ser	His
			180					185					190		
Arg	Ser	Tyr	Ser	Cys	Gln	Val	Thr	His	Glu	Gly	Ser	Thr	Val	Glu	Lys
		195					200					205			
Thr	Val	Ala	Pro	Thr	Glu	Cys	Ser								
	210					215									

<210> SEQ ID NO 373
 <211> LENGTH: 451
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens
 <400> SEQUENCE: 373

Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly
1			5						10					15	
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asp	Tyr
			20					25					30		
Gln	Met	Thr	Trp	Ile	Arg	Gln	Thr	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
	35					40						45			
Ser	Gly	Val	Ser	Trp	Asn	Gly	Ala	Arg	Thr	His	Tyr	Ala	Asp	Ser	Val
	50				55					60					
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65				70				75						80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85					90					95		
Ala	Lys	Gly	Asp	Tyr	Leu	Val	Tyr	Ser	Ala	Tyr	Tyr	Phe	Asp	Ser	Trp
		100						105					110		
Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro
	115					120						125			
Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr
	130				135						140				
Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr
145				150				155						160	
Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro
			165					170					175		
Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr
	180					185							190		
Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn
	195					200						205			
His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser
	210				215						220				
Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu
225				230					235					240	
Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu
			245					250					255		
Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser
		260					265						270		
His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu
	275					280					285				
Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr
	290				295						300				
Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn
305				310					315					320	

-continued

Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro
 325 330 335
 Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln
 340 345 350
 Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val
 355 360 365
 Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val
 370 375 380
 Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro
 385 390 395 400
 Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr
 405 410 415
 Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val
 420 425 430
 Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
 435 440 445
 Ser Pro Gly
 450

<210> SEQ ID NO 374
 <211> LENGTH: 216
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 374

Glu Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
 1 5 10 15
 Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
 20 25 30
 Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45
 Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60
 Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg
 65 70 75 80
 Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg Leu
 85 90 95
 Asn Gly Trp Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln
 100 105 110
 Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu
 115 120 125
 Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr
 130 135 140
 Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val Lys
 145 150 155 160
 Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr
 165 170 175
 Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His
 180 185 190
 Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys
 195 200 205
 Thr Val Ala Pro Thr Glu Cys Ser
 210 215

-continued

```

<210> SEQ ID NO 375
<211> LENGTH: 451
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 375

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1          5          10
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20        25        30
Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
35        40        45
Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val
50        55        60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65        70        75        80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85        90        95
Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ala Tyr Tyr Phe Asp Ser Trp
100       105       110
Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro
115       120       125
Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr
130       135       140
Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr
145       150       155       160
Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro
165       170       175
Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr
180       185       190
Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn
195       200       205
His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser
210       215       220
Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu
225       230       235       240
Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu
245       250       255
Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser
260       265       270
His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu
275       280       285
Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr
290       295       300
Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn
305       310       315       320
Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro
325       330       335
Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln
340       345       350
Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val
355       360       365

```


-continued

Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val
 370 375 380

Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro
 385 390 395 400

Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr
 405 410 415

Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val
 420 425 430

Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
 435 440 445

Ser Pro Gly
 450

<210> SEQ ID NO 376
 <211> LENGTH: 216
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 376

Glu Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
 1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
 20 25 30

Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45

Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg
 65 70 75 80

Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg Leu
 85 90 95

Asn Gly Trp Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln
 100 105 110

Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu
 115 120 125

Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr
 130 135 140

Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val Lys
 145 150 155 160

Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr
 165 170 175

Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His
 180 185 190

Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys
 195 200 205

Thr Val Ala Pro Thr Glu Cys Ser
 210 215

<210> SEQ ID NO 377
 <211> LENGTH: 451
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 377

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

-continued

1	5	10	15
Ser Leu Arg	Leu Ser Cys Ala Ala	Ser Gly Phe Thr Phe Ser Ser Tyr	
	20	25	30
Gln Met Thr	Trp Ile Arg Gln Thr	Gly Lys Gly Leu Glu Trp Val	
	35	40	45
Ser Gly Ile	Ser Trp Asn Gly Gly	Ser Thr His Tyr Ala Asp Ser Val	
	50	55	60
Lys Gly Arg	Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		
	65	70	75
Leu Gln Met	Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys		
	85	90	95
Ala Lys Gly	Asp Tyr Leu Val Tyr Ser Ala Tyr Tyr Phe Asp Ser Trp		
	100	105	110
Gly Gln Gly	Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro		
	115	120	125
Ser Val Phe	Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr		
	130	135	140
Ala Ala Leu	Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr		
	145	150	155
Val Ser Trp	Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro		
	165	170	175
Ala Val Leu	Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr		
	180	185	190
Val Pro Ser	Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn		
	195	200	205
His Lys Pro	Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser		
	210	215	220
Cys Asp Lys	Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu		
	225	230	235
Gly Gly Pro	Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu		
	245	250	255
Met Ile Ser	Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser		
	260	265	270
His Glu Asp	Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu		
	275	280	285
Val His Asn	Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr		
	290	295	300
Tyr Arg Val	Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn		
	305	310	315
Gly Lys Glu	Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro		
	325	330	335
Ile Glu Lys	Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln		
	340	345	350
Val Tyr Thr	Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val		
	355	360	365
Ser Leu Thr	Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val		
	370	375	380
Glu Trp Glu	Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro		
	385	390	395
Pro Val Leu	Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr		
	405	410	415

-continued

Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val
 420 425 430

Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
 435 440 445

Ser Pro Gly
 450

<210> SEQ ID NO 378
 <211> LENGTH: 216
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 378

Glu Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
 1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
 20 25 30

Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45

Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg
 65 70 75 80

Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg Leu
 85 90 95

Asn Gly Trp Gly Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln
 100 105 110

Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu
 115 120 125

Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr
 130 135 140

Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val Lys
 145 150 155 160

Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr
 165 170 175

Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His
 180 185 190

Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys
 195 200 205

Thr Val Ala Pro Thr Glu Cys Ser
 210 215

<210> SEQ ID NO 379
 <211> LENGTH: 451
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 379

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
 20 25 30

Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val

50					55					60					
Lys 65	Gly	Arg	Phe	Thr	Ile 70	Ser	Arg	Asp	Asn 75	Ser	Lys	Asn	Thr	Leu	Tyr 80
Leu	Gln	Met	Asn 85	Ser	Leu	Arg	Ala	Glu	Asp 90	Thr	Ala	Val	Tyr	Tyr 95	Cys
Ala	Lys	Gly	Asp 100	Tyr	Leu	Val	Tyr	Ser	Ala 105	Tyr	Tyr	Phe	Asp 110	Ser	Trp
Gly	Gln	Gly 115	Thr	Leu	Val	Thr 120	Val	Ser	Ser	Ala	Ser	Thr 125	Lys	Gly	Pro
Ser 130	Val	Phe	Pro	Leu	Ala 135	Pro	Ser	Ser	Lys	Ser	Thr 140	Ser	Gly	Gly	Thr
Ala 145	Ala	Leu	Gly	Cys 150	Leu	Val	Lys	Asp	Tyr	Phe 155	Pro	Glu	Pro	Val	Thr 160
Val	Ser	Trp	Asn 165	Ser	Gly	Ala	Leu	Thr	Ser 170	Gly	Val	His	Thr	Phe 175	Pro
Ala	Val	Leu	Gln 180	Ser	Ser	Gly	Leu	Tyr 185	Ser	Leu	Ser	Ser	Val 190	Val	Thr
Val 195	Pro	Ser	Ser	Ser	Leu	Gly 200	Thr	Gln	Thr	Tyr	Ile 205	Cys	Asn	Val	Asn
His 210	Lys	Pro	Ser	Asn 215	Thr	Lys	Val	Asp	Lys	Lys	Val 220	Glu	Pro	Lys	Ser
Cys 225	Asp	Lys	Thr	His 230	Thr	Cys	Pro	Pro	Cys	Pro 235	Ala	Pro	Glu	Leu	Leu 240
Gly	Gly	Pro	Ser 245	Val	Phe	Leu	Phe	Pro	Pro 250	Lys	Pro	Lys	Asp 255	Thr	Leu
Met	Ile	Ser	Arg 260	Thr	Pro	Glu	Val	Thr 265	Cys	Val	Val	Val	Asp 270	Val	Ser
His 275	Glu	Asp	Pro	Glu	Val	Lys 280	Phe	Asn	Trp	Tyr	Val 285	Asp	Gly	Val	Glu
Val 290	His	Asn	Ala	Lys 295	Thr	Lys	Pro	Arg	Glu	Glu	Gln 300	Tyr	Asn	Ser	Thr
Tyr 305	Arg	Val	Val	Ser 310	Val	Leu	Thr	Val	Leu 315	His	Gln	Asp	Trp	Leu	Asn 320
Gly	Lys	Glu	Tyr 325	Lys	Cys	Lys	Val	Ser 330	Asn	Lys	Ala	Leu	Pro 335	Ala	Pro
Ile	Glu	Lys 340	Thr	Ile	Ser	Lys	Ala 345	Lys	Gly	Gln	Pro	Arg	Glu 350	Pro	Gln
Val 355	Tyr	Thr	Leu	Pro	Pro	Ser 360	Arg	Asp	Glu	Leu	Thr	Lys 365	Asn	Gln	Val
Ser 370	Leu	Thr	Cys	Leu	Val	Lys 375	Gly	Phe	Tyr	Pro	Ser 380	Asp	Ile	Ala	Val
Glu 385	Trp	Glu	Ser	Asn 390	Gly	Gln	Pro	Glu	Asn 395	Asn	Tyr	Lys	Thr	Thr	Pro 400
Pro	Val	Leu	Asp 405	Ser	Asp	Gly	Ser	Phe 410	Phe	Leu	Tyr	Ser	Lys 415	Leu	Thr
Val	Asp	Lys 420	Ser	Arg	Trp	Gln	Gln 425	Gly	Asn	Val	Phe	Ser 430	Cys	Ser	Val
Met 435	His	Glu	Ala	Leu	His	Asn 440	His	Tyr	Thr	Gln	Lys 445	Ser	Leu	Ser	Leu
Ser 450	Pro	Gly													

-continued

```

<210> SEQ ID NO 380
<211> LENGTH: 216
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 380
Glu Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1          5          10          15
Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
20        25        30
Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35        40        45
Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
50        55        60
Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg
65        70        75        80
Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg Leu
85        90        95
Asn Gly Trp Gly Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln
100       105       110
Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu
115       120       125
Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr
130       135       140
Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val Lys
145       150       155       160
Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr
165       170       175
Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His
180       185       190
Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys
195       200       205
Thr Val Ala Pro Thr Glu Cys Ser
210       215

<210> SEQ ID NO 381
<211> LENGTH: 451
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 381
Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1          5          10          15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20        25        30
Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
35        40        45
Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val
50        55        60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65        70        75        80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85        90        95
Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ala Tyr Tyr Phe Asp Ser Trp

```

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln

-continued

1	5	10	15
Arg Val Thr	Ile Ser Cys Ser Gly	Ser Ser Ser Asn Ile Gly	Ser Asn
	20	25	30
Pro Val Asn	Trp Tyr Gln Gln Leu	Pro Gly Thr Ala Pro	Lys Leu Leu
	35	40	45
Ile Tyr Arg	Asn Asn Gln Arg Pro	Ser Gly Val Pro Asp	Arg Phe Ser
	50	55	60
Gly Ser Lys	Ser Gly Thr Ser Ala	Ser Leu Ala Ile Ser	Gly Leu Arg
	65	70	75
Ser Glu Asp	Glu Ala Asp Tyr Tyr	Cys Ala Ala Trp Asp	Asp Arg Leu
	85	90	95
Ser Gly Trp	Ala Phe Gly Gly Gly	Thr Lys Leu Thr Val	Leu Gly Gln
	100	105	110
Pro Lys Ala	Ala Pro Ser Val Thr	Leu Phe Pro Pro Ser	Ser Ser Glu Glu
	115	120	125
Leu Gln Ala	Asn Lys Ala Thr Leu	Val Cys Leu Ile Ser	Asp Phe Tyr
	130	135	140
Pro Gly Ala	Val Thr Val Ala Trp	Lys Ala Asp Ser Ser	Pro Val Lys
	145	150	155
Ala Gly Val	Glu Thr Thr Thr Pro	Ser Lys Gln Ser Asn	Asn Lys Tyr
	165	170	175
Ala Ala Ser	Ser Ser Tyr Leu Ser	Leu Thr Pro Glu Gln	Trp Lys Ser His
	180	185	190
Arg Ser Tyr	Ser Cys Gln Val Thr	His Glu Gly Ser Thr	Val Glu Lys
	195	200	205
Thr Val Ala	Pro Thr Glu Cys Ser		
	210	215	

<210> SEQ ID NO 383

<211> LENGTH: 451

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 383

Glu Val Gln	Leu Leu Glu	Ser Gly Gly	Gly Leu Val	Gln Pro Gly	Gly
1	5	10	15		
Ser Leu Arg	Leu Ser Cys	Ala Ala Ser	Gly Phe Thr	Phe Ser Asp	Tyr
	20	25	30		
Gln Met Thr	Trp Ile Arg	Gln Thr Pro	Gly Lys Gly	Leu Glu Trp	Val
	35	40	45		
Ser Gly Ile	Ser Trp Asn	Gly Gly Ser	Thr His Tyr	Ala Asp Ser	Val
	50	55	60		
Lys Gly Arg	Phe Thr Ile	Ser Arg Asp	Asn Ser Lys	Asn Thr Leu	Tyr
	65	70	75	80	
Leu Gln Met	Asn Ser Leu	Arg Ala Glu	Asp Thr Ala	Val Tyr Tyr	Cys
	85	90	95		
Ala Lys Gly	Asp Tyr Leu	Val Tyr Ser	Ser Tyr Tyr	Phe Lys Ser	Trp
	100	105	110		
Gly Gln Gly	Thr Leu Val	Thr Val Ser	Ser Ala Ser	Thr Lys Gly	Pro
	115	120	125		
Ser Val Phe	Pro Leu Ala	Pro Ser Ser	Lys Ser Thr	Ser Gly Gly	Thr
	130	135	140		
Ala Ala Leu	Gly Cys Leu	Val Lys Asp	Tyr Phe Pro	Glu Pro Val	Thr

-continued

145		150		155		160
Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro	165		170		175	
Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr	180		185		190	
Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn	195		200		205	
His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser	210		215		220	
Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu	225		230		235	240
Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu	245		250		255	
Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser	260		265		270	
His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu	275		280		285	
Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr	290		295		300	
Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn	305		310		315	320
Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro	325		330		335	
Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln	340		345		350	
Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val	355		360		365	
Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val	370		375		380	
Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro	385		390		395	400
Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr	405		410		415	
Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val	420		425		430	
Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu	435		440		445	
Ser Pro Gly	450					

<210> SEQ ID NO 384

<211> LENGTH: 216

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 384

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln	1	5	10	15
Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn	20	25	30	
Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu	35	40	45	
Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser				

-continued

50	55	60
Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg		
65	70	75
Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Ser Leu		
	85	90
Ser Gly Trp Ala Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln		
	100	105
Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu		
	115	120
Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr		
	130	135
Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val Lys		
	145	150
Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr		
	165	170
Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His		
	180	185
Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys		
	195	200
Thr Val Ala Pro Thr Glu Cys Ser		
	210	215
<210> SEQ ID NO 385		
<211> LENGTH: 451		
<212> TYPE: PRT		
<213> ORGANISM: Homo Sapiens		
<400> SEQUENCE: 385		
Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly		
1	5	10
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr		
	20	25
Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val		
	35	40
Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val		
	50	55
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		
	65	70
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys		
	85	90
Ala Lys Gly Asp Tyr Leu Val Tyr Lys Ser Tyr Tyr Phe Lys Ser Trp		
	100	105
Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro		
	115	120
Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr		
	130	135
Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr		
	145	150
Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro		
	165	170
Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr		
	180	185
Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn		

-continued

195					200					205					
His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser
210						215					220				
Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu
225					230					235					240
Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu
				245					250					255	
Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser
			260					265					270		
His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu
	275						280					285			
Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr
	290					295					300				
Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn
305					310					315					320
Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro
				325					330					335	
Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln
			340					345					350		
Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val
	355						360					365			
Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val
	370					375					380				
Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro
385					390					395					400
Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr
				405				410						415	
Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val
			420					425					430		
Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu
	435					440						445			
Ser	Pro	Gly													
	450														

<210> SEQ ID NO 386

<211> LENGTH: 216

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 386

Gln	Ser	Val	Leu	Thr	Gln	Pro	Pro	Ser	Ala	Ser	Gly	Thr	Pro	Gly	Gln
1				5					10					15	
Arg	Val	Thr	Ile	Ser	Cys	Ser	Gly	Ser	Ser	Ser	Asn	Ile	Gly	Ser	Asn
			20					25					30		
Pro	Val	Asn	Trp	Tyr	Gln	Gln	Leu	Pro	Gly	Thr	Ala	Pro	Lys	Leu	Leu
			35				40					45			
Ile	Tyr	Arg	Asn	Asn	Gln	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	Phe	Ser
	50				55					60					
Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Ser	Gly	Leu	Arg
65					70					75				80	
Ser	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Ala	Ala	Trp	Asp	Asp	Arg	Leu
			85					90						95	
Asn	Gly	Trp	Ala	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly	Gln

-continued

100					105					110					
Pro	Lys	Ala	Ala	Pro	Ser	Val	Thr	Leu	Phe	Pro	Pro	Ser	Ser	Glu	Glu
115					120					125					
Leu	Gln	Ala	Asn	Lys	Ala	Thr	Leu	Val	Cys	Leu	Ile	Ser	Asp	Phe	Tyr
130					135					140					
Pro	Gly	Ala	Val	Thr	Val	Ala	Trp	Lys	Ala	Asp	Ser	Ser	Pro	Val	Lys
145					150					155					
Ala	Gly	Val	Glu	Thr	Thr	Thr	Pro	Ser	Lys	Gln	Ser	Asn	Asn	Lys	Tyr
165					170					175					
Ala	Ala	Ser	Ser	Tyr	Leu	Ser	Leu	Thr	Pro	Glu	Gln	Trp	Lys	Ser	His
180					185					190					
Arg	Ser	Tyr	Ser	Cys	Gln	Val	Thr	His	Glu	Gly	Ser	Thr	Val	Glu	Lys
195					200					205					
Thr	Val	Ala	Pro	Thr	Glu	Cys	Ser								
210					215										

<210> SEQ ID NO 387

<211> LENGTH: 451

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 387

Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly
1			5					10						15	
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asp	Tyr
		20						25					30		
Gln	Met	Thr	Trp	Ile	Arg	Gln	Thr	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
		35				40						45			
Ser	Gly	Ile	Ser	Trp	Asn	Gly	Gly	Ser	Thr	His	Tyr	Ala	Asp	Ser	Val
	50				55					60					
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65				70				75						80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
		85						90						95	
Ala	Lys	Gly	Asp	Tyr	Leu	Val	Tyr	Ser	Ser	Tyr	Tyr	Phe	Lys	Tyr	Trp
		100						105					110		
Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro
	115					120						125			
Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr
	130				135						140				
Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr
145				150					155					160	
Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro
		165						170						175	
Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr
		180					185						190		
Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn
	195					200						205			
His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser
	210				215						220				
Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu
225				230					235					240	
Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu

```
<210> SEQ ID NO 388
<211> LENGTH: 216
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 388
```

Gln 1	Ser	Val	Leu	Thr 5	Gln	Pro	Pro	Ser	Ala 10	Ser	Gly	Thr	Pro	Gly 15	Gln
Arg	Val	Thr	Ile 20	Ser	Cys	Ser	Gly	Ser 25	Ser	Ser	Asn	Ile	Gly 30	Ser	Asn
Pro	Val	Asn 35	Trp	Tyr	Gln	Gln	Leu 40	Pro	Gly	Thr	Ala	Pro 45	Lys	Leu	Leu
Ile	Tyr 50	Arg	Asn	Asn	Gln	Arg 55	Pro	Ser	Gly	Val	Pro 60	Asp	Arg	Phe	Ser
Gly 65	Ser	Lys	Ser	Gly	Thr 70	Ser	Ala	Ser	Leu	Ala 75	Ile	Ser	Gly	Leu	Arg 80
Ser	Glu	Asp	Glu 85	Ala	Asp	Tyr	Tyr	Cys	Ala 90	Ala	Trp	Asp	Asp 95	Ser	Leu
Asn	Gly	Trp	Gly 100	Phe	Gly	Gly	Gly	Thr 105	Lys	Leu	Thr	Val 110	Leu	Gly	Gln
Pro	Lys 115	Ala	Ala	Pro	Ser	Val	Thr 120	Leu	Phe	Pro	Pro	Ser 125	Ser	Glu	Glu
Leu 130	Gln	Ala	Asn	Lys	Ala	Thr 135	Leu	Val	Cys	Leu	Ile 140	Ser	Asp	Phe	Tyr
Pro	Gly	Ala	Val	Thr	Val	Ala	Trp	Lys	Ala	Asp	Ser	Ser	Pro	Val	Lys

-continued

145	150	155	160
Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr	165	170	175
Ala Ala Ser Ser Thr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His	180	185	190
Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys	195	200	205
Thr Val Ala Pro Thr Glu Cys Ser	210	215	

<210> SEQ ID NO 389
 <211> LENGTH: 451
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens
 <400> SEQUENCE: 389

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly	1	5	10	15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr	20	25	30	
Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val	35	40	45	
Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val	50	55	60	
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr	65	70	75	80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys	85	90	95	
Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ser Tyr Tyr Phe Lys Ser Trp	100	105	110	
Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro	115	120	125	
Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr	130	135	140	
Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr	145	150	155	160
Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro	165	170	175	
Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr	180	185	190	
Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn	195	200	205	
His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser	210	215	220	
Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu	225	230	235	240
Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu	245	250	255	
Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser	260	265	270	
His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu	275	280	285	
Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr				

-continued

290	295	300
Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn		
305	310	315
Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro		
	325	330
Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln		
	340	345
Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val		
	355	360
Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val		
	370	375
Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro		
	385	390
Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr		
	405	410
Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val		
	420	425
Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu		
	435	440
Ser Pro Gly		
450		
<210> SEQ ID NO 390		
<211> LENGTH: 216		
<212> TYPE: PRT		
<213> ORGANISM: Homo Sapiens		
<400> SEQUENCE: 390		
Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln		
1	5	10
Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn		
	20	25
Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu		
	35	40
Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser		
	50	55
Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg		
	65	70
Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Arg Leu		
	85	90
Ser Gly Trp Gly Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln		
	100	105
Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu		
	115	120
Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr		
	130	135
Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val Lys		
	145	150
Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr		
	165	170
Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His		
	180	185
Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys		
		190

-continued

195				200				205			
Thr	Val	Ala	Pro	Thr	Glu	Cys	Ser				
210				215							
<210> SEQ ID NO 391 <211> LENGTH: 451 <212> TYPE: PRT <213> ORGANISM: Homo Sapiens <400> SEQUENCE: 391											
Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val
1				5				10			
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr
	20							25			
Gln	Met	Thr	Trp	Ile	Arg	Gln	Thr	Pro	Gly	Lys	Gly
	35							40			
Ser	Gly	Ile	Ser	Trp	Asn	Gly	Gly	Ser	Thr	His	Tyr
	50							55			
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys
	65							70			
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala
								85			
Ala	Lys	Gly	Asp	Tyr	Leu	Val	Tyr	Lys	Ser	Tyr	Tyr
								100			
Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser
								115			
Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr
								130			
Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro
								145			
Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val
								165			
Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser
								180			
Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile
								195			
His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val
								210			
Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala
								225			
Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro
								245			
Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val
								260			
His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val
								275			
Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln
								290			
Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln
								305			
Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala
								325			
Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro

-continued

340					345					350						
Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	
355					360					365						
Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	
370					375					380						
Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	
385					390					395					400	
Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	
405					410					415						
Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	
420					425					430						
Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	
435					440					445						
Ser	Pro	Gly														
450																

<210> SEQ ID NO 392

<211> LENGTH: 216

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 392

Gln	Ser	Val	Leu	Thr	Gln	Pro	Pro	Ser	Ala	Ser	Gly	Thr	Pro	Gly	Gln
1				5					10					15	
Arg	Val	Thr	Ile	Ser	Cys	Ser	Gly	Ser	Ser	Ser	Asn	Ile	Gly	Ser	Asn
		20					25						30		
Pro	Val	Asn	Trp	Tyr	Gln	Gln	Leu	Pro	Gly	Thr	Ala	Pro	Lys	Leu	Leu
		35					40					45			
Ile	Tyr	Arg	Asn	Asn	Gln	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	Phe	Ser
	50				55					60					
Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Ser	Gly	Leu	Arg
65				70						75				80	
Ser	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Ala	Ala	Trp	Asp	Asp	Arg	Leu
			85					90						95	
Ser	Gly	Trp	Gly	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly	Gln
		100					105						110		
Pro	Lys	Ala	Ala	Pro	Ser	Val	Thr	Leu	Phe	Pro	Pro	Ser	Ser	Glu	Glu
		115				120						125			
Leu	Gln	Ala	Asn	Lys	Ala	Thr	Leu	Val	Cys	Leu	Ile	Ser	Asp	Phe	Tyr
	130				135						140				
Pro	Gly	Ala	Val	Thr	Val	Ala	Trp	Lys	Ala	Asp	Ser	Ser	Pro	Val	Lys
145					150					155				160	
Ala	Gly	Val	Glu	Thr	Thr	Thr	Pro	Ser	Lys	Gln	Ser	Asn	Asn	Lys	Tyr
			165						170					175	
Ala	Ala	Ser	Ser	Tyr	Leu	Ser	Leu	Thr	Pro	Glu	Gln	Trp	Lys	Ser	His
		180					185						190		
Arg	Ser	Tyr	Ser	Cys	Gln	Val	Thr	His	Glu	Gly	Ser	Thr	Val	Glu	Lys
		195				200						205			
Thr	Val	Ala	Pro	Thr	Glu	Cys	Ser								
	210					215									

<210> SEQ ID NO 393

<211> LENGTH: 451

<212> TYPE: PRT

-continued

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 393

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
 20 25 30
 Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Lys Gly Asp Tyr Leu Val Tyr Ser Ser Tyr Tyr Phe Lys Ser Trp
 100 105 110
 Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro
 115 120 125
 Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr
 130 135 140
 Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr
 145 150 155 160
 Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro
 165 170 175
 Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr
 180 185 190
 Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn
 195 200 205
 His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser
 210 215 220
 Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu
 225 230 235 240
 Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu
 245 250 255
 Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser
 260 265 270
 His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu
 275 280 285
 Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr
 290 295 300
 Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn
 305 310 315 320
 Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro
 325 330 335
 Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln
 340 345 350
 Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val
 355 360 365
 Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val
 370 375 380
 Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro

-continued

385	390	395	400
Pro Val Leu Asp	Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr		
	405	410	415
Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val			
	420	425	430
Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu			
	435	440	445
Ser Pro Gly			
450			

<210> SEQ ID NO 394

<211> LENGTH: 216

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 394

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln			
1	5	10	15
Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn			
	20	25	30
Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu			
	35	40	45
Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser			
	50	55	60
Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg			
	65	70	75
Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Ser Leu			
	85	90	95
Asn Gly Trp Gly Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln			
	100	105	110
Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu			
	115	120	125
Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr			
	130	135	140
Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val Lys			
	145	150	155
Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr			
	165	170	175
Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His			
	180	185	190
Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys			
	195	200	205
Thr Val Ala Pro Thr Glu Cys Ser			
210	215		

<210> SEQ ID NO 395

<211> LENGTH: 451

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 395

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly			
1	5	10	15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr			
	20	25	30

-continued

Gln Met Thr Trp Ile Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val	35	40	45	
Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val	50	55	60	
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr	65	70	75	80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys	85	90	95	
Ala Lys Gly Asp Tyr Leu Val Tyr Lys Ser Tyr Tyr Phe Lys Ser Trp	100	105	110	
Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro	115	120	125	
Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr	130	135	140	
Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr	145	150	155	160
Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro	165	170	175	
Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr	180	185	190	
Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn	195	200	205	
His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser	210	215	220	
Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu	225	230	235	240
Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu	245	250	255	
Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser	260	265	270	
His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu	275	280	285	
Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr	290	295	300	
Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn	305	310	315	320
Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro	325	330	335	
Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln	340	345	350	
Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val	355	360	365	
Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val	370	375	380	
Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro	385	390	395	400
Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr	405	410	415	
Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val	420	425	430	
Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu				

-continued

435	440	445
Ser Pro Gly		
450		
 <210> SEQ ID NO 396		
<211> LENGTH: 216		
<212> TYPE: PRT		
<213> ORGANISM: Homo Sapiens		
 <400> SEQUENCE: 396		
Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln		
1 5 10 15		
Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn		
20 25 30		
Pro Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu		
35 40 45		
Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser		
50 55 60		
Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg		
65 70 75 80		
Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Ser Leu		
85 90 95		
Ser Gly Trp Ala Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln		
100 105 110		
Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu		
115 120 125		
Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr		
130 135 140		
Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val Lys		
145 150 155 160		
Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr		
165 170 175		
Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His		
180 185 190		
Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys		
195 200 205		
Thr Val Ala Pro Thr Glu Cys Ser		
210 215		
 <210> SEQ ID NO 397		
<211> LENGTH: 451		
<212> TYPE: PRT		
<213> ORGANISM: Homo Sapiens		
 <400> SEQUENCE: 397		
Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly		
1 5 10 15		
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr		
20 25 30		
Gln Met Thr Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val		
35 40 45		
Ser Gly Ile Ser Trp Asn Gly Gly Ser Thr His Tyr Ala Asp Ser Val		
50 55 60		
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		
65 70 75 80		

-continued

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Lys Gly Asp Tyr Leu Val Tyr Lys Ser Tyr Tyr Phe Lys Ser Trp
 100 105 110
 Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro
 115 120 125
 Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr
 130 135 140
 Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr
 145 150 155 160
 Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro
 165 170 175
 Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr
 180 185 190
 Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn
 195 200 205
 His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser
 210 215 220
 Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu
 225 230 235 240
 Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu
 245 250 255
 Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser
 260 265 270
 His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu
 275 280 285
 Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr
 290 295 300
 Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn
 305 310 315 320
 Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro
 325 330 335
 Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln
 340 345 350
 Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val
 355 360 365
 Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val
 370 375 380
 Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro
 385 390 395 400
 Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr
 405 410 415
 Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val
 420 425 430
 Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
 435 440 445
 Ser Pro Gly
 450

<210> SEQ ID NO 398

<211> LENGTH: 10

<212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 398

Gly Tyr Ser Phe Thr Ser Tyr Trp Ile Gly
1 5 10

<210> SEQ ID NO 399

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 399

Trp Met Gly Ile Ile Asp Pro Gly Asp Ser Arg Thr Arg Tyr Ser Pro
1 5 10 15

Ser Phe Gln Gly
20

<210> SEQ ID NO 400

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 400

Gly Gln Leu Tyr Gly Gly Thr Tyr Met Asp Gly
1 5 10

<210> SEQ ID NO 401

<211> LENGTH: 14

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 401

Thr Gly Thr Ser Ser Asp Ile Gly Gly Tyr Asn Ser Val Ser
1 5 10

<210> SEQ ID NO 402

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 402

Leu Met Ile Tyr Gly Val Asn Asn Arg Pro Ser
1 5 10

<210> SEQ ID NO 403

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 403

Ser Ser Tyr Asp Ile Glu Ser Ala Thr Pro
1 5 10

<210> SEQ ID NO 404

<211> LENGTH: 120

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 404

Gln Val Glu Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu
1 5 10 15

Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Thr Ser Tyr

-continued

20					25					30						
Trp	Ile	Gly	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Met	
35					40					45						
Gly	Ile	Ile	Asp	Pro	Gly	Asp	Ser	Arg	Thr	Arg	Tyr	Ser	Pro	Ser	Phe	
50					55					60						
Gln	Gly	Gln	Val	Thr	Ile	Ser	Ala	Asp	Lys	Ser	Ile	Ser	Thr	Ala	Tyr	
65					70					75					80	
Leu	Gln	Trp	Ser	Ser	Leu	Lys	Ala	Ser	Asp	Thr	Ala	Met	Tyr	Tyr	Cys	
85					90					95						
Ala	Arg	Gly	Gln	Leu	Tyr	Gly	Gly	Thr	Tyr	Met	Asp	Gly	Trp	Gly	Gln	
100					105					110						
Gly	Thr	Leu	Val	Thr	Val	Ser	Ser									
115					120											

<210> SEQ ID NO 405

<211> LENGTH: 113

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 405

Asp	Ile	Ala	Leu	Thr	Gln	Pro	Ala	Ser	Val	Ser	Gly	Ser	Pro	Gly	Gln
1			5						10					15	
Ser	Ile	Thr	Ile	Ser	Cys	Thr	Gly	Thr	Ser	Ser	Asp	Ile	Gly	Gly	Tyr
		20						25					30		
Asn	Ser	Val	Ser	Trp	Tyr	Gln	Gln	His	Pro	Gly	Lys	Ala	Pro	Lys	Leu
		35					40					45			
Met	Ile	Tyr	Gly	Val	Asn	Asn	Arg	Pro	Ser	Gly	Val	Ser	Asn	Arg	Phe
	50					55					60				
Ser	Gly	Ser	Lys	Ser	Gly	Asn	Thr	Ala	Ser	Leu	Thr	Ile	Ser	Gly	Leu
	65				70					75				80	
Gln	Ala	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Ser	Ser	Tyr	Asp	Ile	Glu
			85						90					95	
Ser	Ala	Thr	Pro	Val	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly
			100					105						110	

Gln

<210> SEQ ID NO 406

<211> LENGTH: 360

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 406

caggtggaat	tggttcagag	cggcgcggaa	gtgaaaaaac	cgggcgaaag	cctgaaaatt	60
agctgcaaa	gttcgggata	ttcctttact	tcttattgga	ttggttggtg	gcgccaggcc	120
cctgggaagg	gtctcgatg	gatgggcatt	atcgatccgg	gtgatagccg	taccggttat	180
tctccgagct	ttcagggcc	ggtgaccatt	agcgcggata	aaagcattag	caccgcgtat	240
cttcaatgga	gcagcctgaa	agcgagcgat	acggccatgt	attattgcgc	gcgtggtcag	300
ctttatgggtg	gtacttatat	ggatgggttg	ggccaaggca	ccctgggtgac	ggtagctca	360

<210> SEQ ID NO 407

<211> LENGTH: 339

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 407

```

gatatcgcac tgaccagacc agcttcagtg agcggctcac caggtcagag cattaccatc      60
tcgtgtacgg gtactagcag cgatattggt ggttataatt ctgtgtcttg gtaccagcag      120
catcccggga aggcgcgcaa acttatgatt tatggtgtta ataatcgccc ctcaggcgtg      180
agcaaccggt ttagcggatc caaaagcggc aacaccgcga gcctgaccat tagcggcctg      240
caagcggaag acgaagcgga ttattattgc tcttcttatg atattgagtc tgctactcct      300
gtgtttggcg gcggcacgaa gttaaccgtc ctaggtcag                                339

```

<210> SEQ ID NO 408

<211> LENGTH: 450

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 408

```

Gln Val Glu Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu
1           5           10           15
Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Thr Ser Tyr
20          25          30
Trp Ile Gly Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Met
35          40          45
Gly Ile Ile Asp Pro Gly Asp Ser Arg Thr Arg Tyr Ser Pro Ser Phe
50          55          60
Gln Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Ser Thr Ala Tyr
65          70          75          80
Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys
85          90          95
Ala Arg Gly Gln Leu Tyr Gly Gly Thr Tyr Met Asp Gly Trp Gly Gln
100         105         110
Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
115         120         125
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
130         135         140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
145         150         155         160
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
165         170         175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
180         185         190
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
195         200         205
Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp
210         215         220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
225         230         235         240
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
245         250         255
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
260         265         270
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
275         280         285
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg

```


-continued

290					295					300					
Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys
305					310					315					320
Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu
				325					330					335	
Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr
			340					345					350		
Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu
	355						360					365			
Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp
	370					375					380				
Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val
385					390					395					400
Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp
				405				410						415	
Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His
			420					425					430		
Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro
	435						440					445			
Gly	Lys														
	450														

<210> SEQ ID NO 409

<211> LENGTH: 217

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 409

Asp	Ile	Ala	Leu	Thr	Gln	Pro	Ala	Ser	Val	Ser	Gly	Ser	Pro	Gly	Gln
1				5					10					15	
Ser	Ile	Thr	Ile	Ser	Cys	Thr	Gly	Thr	Ser	Ser	Asp	Ile	Gly	Gly	Tyr
		20						25					30		
Asn	Ser	Val	Ser	Trp	Tyr	Gln	Gln	His	Pro	Gly	Lys	Ala	Pro	Lys	Leu
		35					40					45			
Met	Ile	Tyr	Gly	Val	Asn	Asn	Arg	Pro	Ser	Gly	Val	Ser	Asn	Arg	Phe
	50					55					60				
Ser	Gly	Ser	Lys	Ser	Gly	Asn	Thr	Ala	Ser	Leu	Thr	Ile	Ser	Gly	Leu
65					70					75				80	
Gln	Ala	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Ser	Ser	Tyr	Asp	Ile	Glu
			85						90					95	
Ser	Ala	Thr	Pro	Val	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly
		100						105					110		
Gln	Pro	Lys	Ala	Ala	Pro	Ser	Val	Thr	Leu	Phe	Pro	Pro	Ser	Ser	Glu
		115					120					125			
Glu	Leu	Gln	Ala	Asn	Lys	Ala	Thr	Leu	Val	Cys	Leu	Ile	Ser	Asp	Phe
	130					135					140				
Tyr	Pro	Gly	Ala	Val	Thr	Val	Ala	Trp	Lys	Gly	Asp	Ser	Ser	Pro	Val
145						150				155					160
Lys	Ala	Gly	Val	Glu	Thr	Thr	Thr	Pro	Ser	Lys	Gln	Ser	Asn	Asn	Lys
			165						170					175	
Tyr	Ala	Ala	Ser	Ser	Tyr	Leu	Ser	Leu	Thr	Pro	Glu	Gln	Trp	Lys	Ser
		180						185					190		
His	Arg	Ser	Tyr	Ser	Cys	Gln	Val	Thr	His	Glu	Gly	Ser	Thr	Val	Glu


-continued

195	200	205	
Lys Thr Val Ala Pro Thr Glu Cys Ser			
210	215		
<210> SEQ ID NO 410			
<211> LENGTH: 1353			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 410			
cagggtggaat tgggttcagag cggcgcgga	gtgaaaaaac cggcggaag cctgaaaatt	60	
agctgcaaaag gttccggata ttcctttact	tcttattgga ttggttgggt gcgccaggcc	120	
cctgggaagg gtctcgagtg gatgggcatt	atcgatccgg gtgatagccg tacccggtat	180	
tctccgagct ttcagggccca ggtgaccatt	agcgcggaata aaagcattag caccgcgtat	240	
cttcaatgga gcagcctgaa agcgagcgat	acggccatgt attattgcgc gcgtggtcag	300	
ctttatgggt gtacttatat ggatgggttg	ggccaaggca ccctgggtgac ggtagetca	360	
gcctccacca agggctccatc ggtcttcccc	ctggcaccct cctccaagag cacctctggg	420	
ggcacagcgg ccctgggctg cctgggtcaag	gactacttcc ccgaaccggg gacgggtgctg	480	
tggaaactcag gcgcctgac cagcggcgtg	cacaccttcc cggtgtgect acagtccctca	540	
ggactctact ccctcagcag cgtgggtgacc	gtgccctcca gcagcttggg caccagacc	600	
tacatctgca acgtgaatca caagcccagc	aacaccaagg tggacaagaa agttgagccc	660	
aaatcttggt acaaaactca cacatgccca	ccgtgccag cactgaact cctgggggga	720	
ccgtcagtc tctcttcccc cccaaaacc	aaggacaccc tcatgatctc ccggaccct	780	
gaggtcacat gcgtgggtgt ggacgtgagc	cacgaagacc ctgaggtcaa gttcaactgg	840	
tacgtggacg gcgtggaggt gcataatgcc	aagacaaagc cgcgggagga gcagtacaac	900	
agcacgtacc ggggtggtcag cgtcctcacc	gtcctgcacc aggactgggt gaatggcaag	960	
gagtacaagt gcaaggtctc caacaaagcc	ctcccagccc ccctcgagaa aaccatctcc	1020	
aaagccaaag ggcagccccc agaaccacag	gtgtacaccc tgcccccatc ccgggatgag	1080	
ctgaccaaga accaggtcag cctgacctgc	ctgggtcaaag gcttctatcc cagcgacatc	1140	
gccgtggagt gggagagcaa tgggcagccg	gagaacaact acaagaccac gcctcccggtg	1200	
ctggactccg acggctcctt cttcctctac	agcaagctca ccgtggacaa gagcaggtgg	1260	
cagcagggga acgtcttctc atgctccgtg	atgcatgagg ctctgcacaa ccactacacg	1320	
cagaagagcc tctccctgtc tccgggtaaa	tga	1353	
<210> SEQ ID NO 411			
<211> LENGTH: 654			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 411			
gatatcgac tgaccagcc agcttcagtg	agcggctcac caggtcagag cattaccatc	60	
tcgtgtacgg gtactagcag cgatatgggt	ggttataatt ctgtgtcttg gtaccagcag	120	
catccccgga aggcgcgga acttatgatt	tatgggtgta ataactgtcc ctgagcgtg	180	
agcaaccgtt ttagcggatc caaaagcggc	aacaccgca gcctgacct tagcggcctg	240	
caagcggaag acgaagcgga ttattattgc	tcttcttatg atattgagtc tgctactcct	300	

-continued

gtgtttggcg ggggcacgaa gttaaccgtc ctaggtcagc ccaaggctgc cccctcggtc	360
actctgttcc cggcctcttc tgaggagctt caagccaaca aggccacact ggtgtgtctc	420
ataagtgact tctaccgggg agccgtgaca gtggcctgga agggagatag cagcccggtc	480
aaggcgggag tggagaccac cacaccctcc aaacaaagca acaacaagta cgcggccagc	540
agctatctga gcctgacgcc tgagcagtg gaaagtcacaga gaagctacag ctgccaggtc	600
acgcatagaag ggagaccgtt ggagaagaca gtggccccta cagaatgttc atag	654

1. Binder-drug conjugates of the general formula (Ia)



in which

n is a number from 1 to 50,

AK is a binder,


the group $\text{-G-L}^1\text{-B-L}^2\text{-}$ is a linker,

where

§ marks the linkage site with the group AK and

§§ marks the linkage site with the nitrogen atom,

D is a group of the formula




in which

#4 marks the linkage site with the adjacent nitrogen atom,

#5 marks the linkage site with the carbonyl group,

the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



where


#3 marks the linkage site with the nitrogen atom,

R¹ is hydrogen or methyl,


R² is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

or


R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which
 $\#^6$ marks the linkage site with the carbonyl group,
 R^6 is hydrogen, hydroxy or benzyloxy,
 R^3 is hydrogen or methyl,
 R^4 is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,
 or
 R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




in which
 $\#^7$ marks the linkage site with the adjacent nitrogen atom,
 $\#^8$ marks the linkage site with the group T^1 ,
 T^1 is a group of the formula $-C(=O)-OR^7$, $-C(=O)-NR^8R^9$, $-C(=O)-NH-NH-R^{10}$ or $-CH_2-O-R^{11}$,
 in which
 R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,
 R^8 is hydrogen or methyl,
 R^9 is hydrogen, methyl, ethyl, n-propyl or benzyl,
 or
 R^8 and R^9 together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,
 R^{10} is benzoyl,
 R^{11} is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,
 R^5 is hydrogen, methyl or a group of the formula




in which
 $\#^9$ marks the linkage site with $-CHC(R^{26})-T^2$,
 R^{12} is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula $-S(O)_2OH$,
 R^{13} is phenyl which may be substituted by methoxycarbonyl or carboxyl,


R^{26} is hydrogen or hydroxy,
 T^2 is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,
 R^{35} is methyl or hydroxy,
 and also their salts, solvates and solvates of the salts.
 2. Binder-drug conjugates of the general formula (1a) according to claim 1, in which
 n is a number from 1 to 50,
 AK is AK_1 or AK_2
 where
 AK_1 is a binder which is bonded via a sulphur atom of the binder to the group G ,
 AK_2 is a binder which is bonded via a nitrogen atom of the binder to the group G ,
 G when $AK=AK_1$, is a group of the formula



where
 $\#^1$ marks the linkage site with the sulphur atom of the binder,
 $\#^2$ marks the linkage site with the group L^1 ,
 or
 when $AK=AK_2$, is carbonyl,
 L^1 is a bond, linear (C_1-C_{10}) -alkanediyl, a group of the formula




where
 m is a number from 2 to 6,
 $\#^1$ marks the linkage site with the group G ,
 $\#^2$ marks the linkage site with the group B ,
 L^{1A} is linear (C_2-C_{10}) -alkanediyl,
 B^1 is a group of the formula



in which

##⁵ marks the linkage site with the group L^{1A},
 ##⁶ marks the linkage site with the group L^{1B},
 L⁵ is a bond or (C₂-C₄)-alkanediyl,
 L⁶ is a bond or a group of the formula



in which


##⁷ marks the linkage site with the carbonyl group,
 ##⁸ marks the linkage site with L^{1B},
 R³³ is hydrogen, (C₁-C₄)-alkylcarbonyl, tert-butyloxy carbonyl or benzyloxycarbonyl,
 R³⁴ is hydrogen or methyl,
 R²⁹ is hydrogen or (C₁-C₄)-alkyl,
 R³⁰ is hydrogen or (C₁-C₄)-alkyl,
 or
 R²⁹ and R³⁰ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 R³¹ is hydrogen or (C₁-C₄)-alkyl,
 R³² is hydrogen or (C₁-C₄)-alkyl,
 or
 R³¹ and R³² together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 L^{1B} is linear (C₂-C₁₀)-alkanediyl,
 and

where (C₁-C₁₀)-alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,


and

where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,

B is a bond or a group of the formula




-continued



where

* marks the linkage site with L¹,
 ** marks the linkage site with L²,
 P is O or NH,
 L³ is a bond or (C₂-C₄)-alkanediyl,
 L⁴ is a bond or a group of the formula



in which

*** marks the linkage site with the carbonyl group,
 **** marks the linkage site with L²,
 R²⁵ is hydrogen or methyl,
 R²⁸ is hydrogen, (C₁-C₄)-alkylcarbonyl, tert-butyloxy carbonyl or benzyloxycarbonyl,

Q¹ is a 4- to 7-membered heterocycle,

Q² is a 3- to 7-membered carbocycle or a 4- to 7-membered heterocycle,

R¹⁴ is hydrogen or (C₁-C₄)-alkyl,

R¹⁵ is hydrogen or (C₁-C₄)-alkyl,

or

R¹⁴ and R¹⁵ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

R¹⁶ is hydrogen or (C₁-C₄)-alkyl,

R¹⁷ is hydrogen or (C₁-C₄)-alkyl,

or

R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

R¹⁸ is hydrogen or (C₁-C₄)-alkyl,

R¹⁹ is hydrogen or the side group of a natural α -amino acid or of its homologues or isomers,

R²⁰ is hydrogen or (C₁-C₄)-alkyl,

or

R¹⁹ and R²⁰ together with the atoms to which they are bonded form a pyrrolidinyl ring,

R²¹ is hydrogen or (C₁-C₄)-alkyl,

R²² is hydrogen or (C₁-C₄)-alkyl,

or

R²¹ and R²² together with the atoms to which they are bonded form a 3- to 7-membered carbocycle,

R²³ is (C₁-C₄)-alkyl,

R²⁴ is hydrogen or (C₁-C₄)-alkyl,

R²⁷ is hydrogen or (C₁-C₄)-alkyl,


R³⁶ is hydrogen, (C₁-C₄)-alkylcarbonyl, tert-butyloxy-carbonyl or benzyloxycarbonyl,

R³⁷ is hydrogen or methyl,

or

R³⁶ and R³⁷ together with the atoms to which they are bonded form a pyrrolidine ring,

L² is linear (C₂-C₁₀)-alkanediyl or is a group of the formula



where

p is a number from 2 to 6,

##³ marks the linkage site with the group B,


##⁴ marks the linkage site with the nitrogen atom,

where (C₂-C₁₀)-alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,

and

where two carbon atoms of the alkanediyl chain in 1,2-, 1,3- or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,

D is a group of the formula



in which


##³ marks the linkage site with the nitrogen atom,

R¹ is hydrogen or methyl,

R² is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

or


R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

##⁴ marks the linkage site with the adjacent nitrogen atom,

##⁵ marks the linkage site with the carbonyl group, the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



in which

##⁶ marks the linkage site with the carbonyl group,


R⁶ is hydrogen, hydroxy or benzyloxy,

R³ is hydrogen or methyl,

R⁴ is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

or

R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

##⁷ marks the linkage site with the adjacent nitrogen atom,

##⁸ marks the linkage site with the group T¹,

T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹,

in which

R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

R⁸ is hydrogen or methyl,

R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,


or

R⁸ and R⁹ together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

R¹⁰ is benzoyl,

R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

R⁵ is hydrogen, methyl or a group of the formula



in which

#⁹ marks the linkage site with —CHC(R²⁶)-T²,

R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,

R²⁶ is hydrogen or hydroxy,

T² is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,

R³⁵ is methyl or hydroxy,

and also their salts, solvates and solvates of the salts.

3. Binder-drug conjugates of the general formula (Ia) according to claim 1 or 2, in which

n is a number from 1 to 20,


AK is AK₁ or AK₂

where

AK₁ is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

AK₂ is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the NH side group of a lysine residue of the binder to the group G,

G when AK=AK₁, is a group of the formula



in which


#¹ marks the linkage site with the cysteine residue of the binder,

#² marks the linkage site with the group L¹,

or

when AK=AK₂, is carbonyl,

L¹ is a bond, linear (C₂-C₆)-alkanediyl, a group of the formula



where


m is a number from 2 to 6,

##¹ marks the linkage site with the group G,

##² marks the linkage site with the group B,

L^{1A} is linear (C₂-C₆)-alkanediyl,

B¹ is a group of the formula




in which

##⁵ marks the linkage site with the group L^{1A},

##⁶ marks the linkage site with the group L^{1B},

L⁵ is a bond,

L⁶ is a bond or a group of the formula



in which

##⁷ marks the linkage site with the carbonyl group,

##⁸ marks the linkage site with L^{1B},

R³³ is hydrogen, methylcarbonyl or tert-butyloxy-carbonyl,

R³⁴ is hydrogen or methyl,

R²⁹ is hydrogen,

R³⁰ is hydrogen,

R³¹ is hydrogen or methyl,


R³² is hydrogen or methyl,


L^{1B} is linear (C₂-C₆)-alkanediyl,

and

where (C₂-C₆)-alkanediyl may be substituted by 1 or 2 methyl substituents,

B is a bond or a group of the formula






where

* marks the linkage site with L^1 ,

** marks the linkage site with L^2 ,

L^3 is a bond or ethane-1,2-diyl,

L^4 is a bond or a group of the formula



in which

*** marks the linkage site with the carbonyl group,

**** marks the linkage site with L^2 ,

R^{25} is hydrogen or methyl,

R^{28} is hydrogen, methylcarbonyl or tert-butyloxycarbonyl,

Q^1 is a 4- to 7-membered heterocycle,

R^{14} is hydrogen,

R^{15} is hydrogen,

R^{16} is hydrogen or methyl,

R^{17} is hydrogen or methyl,

or

R^{16} and R^{17} together with the atoms to which they are bonded form a piperazinyl ring,

R^{18} is hydrogen,


R^{19} is hydrogen, methyl, propan-2-yl, 2-methylpropan-1-yl or 1-methylpropan-1-yl,

R^{20} is hydrogen or methyl,

or

R^{19} and R^{20} together with the atoms to which they are bonded form a pyrrolidinyl ring,

R^{21} is hydrogen or methyl,
 R^{22} is hydrogen or methyl,
 or
 R^{21} and R^{22} together with the atoms to which they are bonded form a cyclopropyl ring,
 R^{23} is methyl,
 R^{24} is hydrogen or methyl,
 R^{27} is hydrogen,
 R^{36} is hydrogen, methylcarbonyl or tert-butyloxycarbonyl,
 R^{37} is hydrogen or methyl,
 or
 R^{36} and R^{37} together with the atoms to which they are bonded form a pyrrolidine ring,
 L^2 is linear (C_2-C_6) -alkanediyl or is a group of the formula



where


p is a number from 2 to 6,


##3 marks the linkage site with the group B,

##4 marks the linkage site with the nitrogen atom,

where (C_2-C_{10}) -alkanediyl may be substituted by 1 or 2 methyl substituents,

D is a group of the formula





in which

#6 marks the linkage site with the carbonyl group,


R⁶ is hydrogen, hydroxy or benzyloxy,

R³ is hydrogen,

R⁴ is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or

R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

#7 marks the linkage site with the adjacent nitrogen atom,

#8 marks the linkage site with the group T¹,

T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹,

in which

R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

R⁸ is hydrogen or methyl,

R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,


or

R⁸ and R⁹ together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

R¹⁰ is benzoyl,

R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

R⁵ is hydrogen, methyl or a group of the formula



in which

#9 marks the linkage site with —CHC(R²⁶)-T²,

R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,

R²⁶ is hydrogen or hydroxy,

T² is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,

R³⁵ is methyl or hydroxy,

and also their salts, solvates and solvates of the salts.

4. Binder-drug conjugates of the general formula (Ia) according to any of claims 1 to 3, in which

n is a number from 1 to 10,


AK is AK₁ or AK₂

where

AK₁ is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

AK₂ is an antibody which comprises the six CDR sequences of the antibody B01-3, B01-10, M31-B01 or D02-6, the variable light and variable heavy chain of the antibody B01-3, B01-10, M31-B01 or D02-6 or the light and heavy chain of the antibody B01-3, B01-10, M31-B01 or D02-6, and is bonded via the NH side group of a lysine residue of the binder to the group G,

G when AK=AK₁, is a group of the formula



in which


#1 marks the linkage site with the cysteine residue of the binder,

#2 marks the linkage site with the group L¹,

or

when AK=AK₂, is carbonyl,

L¹ is a bond, linear (C₂-C₆)-alkanediyl, a group of the formula



where


m is a number 2 or 3,

##1 marks the linkage site with the group G,


##2 marks the linkage site with the group B,

where (C₂-C₆)-alkanediyl may be substituted by 1 or 2 methyl substituents,

B is a bond or a group of the formula



* marks the linkage site with L¹,
 ** marks the linkage site with L²,
 L³ is a bond or ethane-1,2-diyl,
 L⁴ is a bond or a group of the formula



in which

*** marks the linkage site with the carbonyl group,
 **** marks the linkage site with L²,
 R²⁵ is methyl,
 R²⁸ is hydrogen, methylcarbonyl or tert-butyloxycarbonyl,

Q¹ is piperidine-1,4-diyl,

R¹⁶ is hydrogen or methyl,

R¹⁷ is hydrogen or methyl,

or

R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a piperazinyl ring,

R²¹ is hydrogen or methyl,

R²² is hydrogen or methyl,

or

R²¹ and R²² together with the atoms to which they are bonded form a cyclopropyl ring,


R²³ is methyl,

R²⁴ is hydrogen,

R³⁶ is hydrogen, methylcarbonyl or tert-butyloxycarbonyl,

R³⁷ is hydrogen or methyl,

L² is linear (C₂-C₆)-alkanediyl or is a group of the formula




where

p is a number from 2 to 6,

##3 marks the linkage site with the group B,

##4 marks the linkage site with the nitrogen atom,
 D is a group of the formula



where


##3 marks the linkage site with the nitrogen atom,

R¹ is hydrogen,

R² is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or

R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




in which

#4 marks the linkage site with the adjacent nitrogen atom,

#5 marks the linkage site with the carbonyl group,

the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



in which


#6 marks the linkage site with the carbonyl group,

R⁶ is hydrogen, hydroxy or benzyloxy,


R³ is hydrogen,

R⁴ is benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or
 R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which
 $\#^7$ marks the linkage site with the adjacent nitrogen atom,
 $\#^8$ marks the linkage site with the group T^1 ,
 T^1 is a group of the formula $-C(=O)-OR^7$,
 $-C(=O)-NR^8R^9$, or $-CH_2-O-R^{11}$,
 in which
 R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,
 R^8 is hydrogen or methyl,
 R^9 is hydrogen, methyl, ethyl, n-propyl or benzyl,
 R^{11} is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,
 R^5 is hydrogen, methyl or a group of the formula




in which
 $\#^9$ marks the linkage site with $-CHCH_2$ phenyl,
 R^{12} is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula $-S(O)_2OH$,
 R^{13} is phenyl which may be substituted by methoxycarbonyl or carboxyl,
 R^{35} is methyl or hydroxy,
 and also their salts, solvates and solvates of the salts.
5. Binder-drug conjugates of the general formula (Ia) according to any of claims 1 to 4, in which
 n is a number from 1 to 10,
 AK is AK_2
 where

AK_2 is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the NH side group of a lysine residue of the binder to the group G,

G is carbonyl,
 L^1 is a bond,
 B is a bond,

L^2 is linear (C_3-C_6)-alkanediyl or is a group of the formula




where

p is a number 2 or 3,

$\#^3$ marks the linkage site with the group B,

$\#^4$ marks the linkage site with the nitrogen atom,

D is a group of the formula



where


$\#^3$ marks the linkage site with the nitrogen atom,

R^1 is hydrogen,

R^2 is benzyl, 4-hydroxybenzyl or 1H-indol-3-ylmethyl,

or

R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




in which

$\#^4$ marks the linkage site with the adjacent nitrogen atom,

$\#^5$ marks the linkage site with the carbonyl group,

the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula




in which

$\#^6$ marks the linkage site with the carbonyl group,

R^3 is hydrogen,

R^4 is benzyl, 4-hydroxybenzyl or 1H-indol-3-ylmethyl,

or
 R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




in which
 $\#7$ marks the linkage site with the adjacent nitrogen atom,
 $\#8$ marks the linkage site with the group T^1 ,
 T^1 is a group of the formula $-C(=O)-OR^7$ or $-C(=O)-NR^8R^9$
 in which
 R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,
 R^8 is hydrogen,
 R^9 is hydrogen or benzyl,
 R^{35} is methyl,
 and also their salts, solvates and solvates of the salts.


6. Binder-drug conjugates of the general formula (Ia) according to any of claims 1 to 4, in which
 n is a number from 1 to 10,
 AK is AK_1 ,
 where

AK_1 is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta, and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

G is a group of the formula




where
 $\#1$ marks the linkage site with the cysteine residue of the binder,
 $\#2$ marks the linkage site with the group L^1 ,
 L^1 is a bond, linear (C_3-C_5) -alkanediyl or a group of the formula




where
 m is a number 2 or 3,
 $\#1$ marks the linkage site with the group G,
 $\#2$ marks the linkage site with the group B,

where (C_3-C_5) -alkanediyl may be substituted by 1 or 2 methyl substituents,
 B is a bond or a group of the formula



where

* marks the linkage site with L^1 ,
 ** marks the linkage site with L^2 ,
 L^3 is a bond or ethane-1,2-diyl,
 L^4 is a bond or a group of the formula



in which

*** marks the linkage site with the carbonyl group,
 **** marks the linkage site with L^2 ,
 R^{25} is methyl,
 R^{28} is hydrogen, methylcarbonyl or tert-butyloxycarbonyl,


R^{16} is hydrogen or methyl,

R^{17} is hydrogen or methyl,

or

R^{16} and R^{17} together with the atoms to which they are bonded form a piperazinyl ring,

L^2 is linear (C_3-C_5) -alkanediyl or is a group of the formula




where

p is a number 2 or 3,

##3 marks the linkage site with the group B,

##4 marks the linkage site with the nitrogen atom,

D is a group of the formula



where


##3 marks the linkage site with the nitrogen atom,

R^1 is hydrogen,

R^2 is benzyl, 4-hydroxybenzyl or 1H-indol-3-ylmethyl,

or

R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

#⁷ marks the linkage site with the adjacent nitrogen atom,

#⁸ marks the linkage site with the group T¹,

T¹ is a group of the formula —C(=O)—OR⁷ or —C(=O)—NR⁸R⁹,

in which

R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,


R⁸ is hydrogen,

R⁹ is hydrogen or benzyl,

R³⁵ is methyl,

and also their salts, solvates and solvates of the salts.


7. Compounds of the formula (XXXa)



in which

#⁴ marks the linkage site with the adjacent nitrogen atom,

#⁵ marks the linkage site with the carbonyl group, the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



in which


#⁶ marks the linkage site with the carbonyl group,

R³ is hydrogen,

R⁴ is benzyl, 4-hydroxybenzyl or 1H-indol-3-ylmethyl,

or


R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

Cys is a cysteine residue which is bonded via the sulphur atom of the side chain to a carbon atom of the succinimide,

L¹ is a bond, linear (C₁-C₁₀)-alkanediyl, a group of the formula



where


m is a number from 2 to 6,

##¹ marks the linkage site with the group G,

##² marks the linkage site with the group B,

L^{1A} is linear (C₂-C₁₀)-alkanediyl,

B¹ is a group of the formula




in which

##⁵ marks the linkage site with the group L^{1A},


##⁶ marks the linkage site with the group L^{1B},

L^5 is a bond or (C₂-C₄)-alkanediyl,
 L^6 is a bond,
 R^{29} is hydrogen or (C₁-C₄)-alkyl,
 R^{30} is hydrogen or (C₁-C₄)-alkyl,
 or
 R^{29} and R^{30} together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 R^{31} is hydrogen or (C₁-C₄)-alkyl,
 R^{32} is hydrogen or (C₁-C₄)-alkyl,
 or
 R^{31} and R^{32} together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 L^{1B} is linear (C₂-C₁₀)-alkanediyl,
 and
 where (C₁-C₁₀)-alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,
 and
 where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,
 B is a bond or a group of the formula




where
 * marks the linkage site with L^1 ,
 ** marks the linkage site with L^2 ,
 P is O or NH,
 L^3 is a bond or (C₂-C₄)-alkanediyl,
 L^4 is a bond,
 Q^1 is a 4- to 7-membered heterocycle,
 Q^2 is a 3- to 7-membered carbocycle or a 4- to 7-membered heterocycle,
 R^{14} is hydrogen or (C₁-C₄)-alkyl,
 R^{15} is hydrogen or (C₁-C₄)-alkyl,
 or
 R^{14} and R^{15} together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 R^{16} is hydrogen or (C₁-C₄)-alkyl,
 R^{17} is hydrogen or (C₁-C₄)-alkyl,
 or
 R^{16} and R^{17} together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

R^{18} is hydrogen or (C₁-C₄)-alkyl,
 R^{19} is hydrogen or the side group of a natural α -amino acid or of its homologues or isomers,
 R^{20} is hydrogen or (C₁-C₄)-alkyl,
 or
 R^{19} and R^{20} together with the atoms to which they are bonded form a pyrrolidinyl ring,
 R^{21} is hydrogen or (C₁-C₄)-alkyl,
 R^{22} is hydrogen or (C₁-C₄)-alkyl,
 or
 R^{21} and R^{22} together with the atoms to which they are bonded form a 3- to 7-membered carbocycle,
 R^{23} is (C₁-C₄)-alkyl,
 R^{24} is hydrogen or (C₁-C₄)-alkyl,
 R^{27} is hydrogen or (C₁-C₄)-alkyl,
 L^2 is linear (C₂-C₁₀)-alkanediyl or is a group of the formula




where
 p is a number from 2 to 6,
 ##³ marks the linkage site with the group B,
 ##⁴ marks the linkage site with the nitrogen atom,
 where (C₂-C₁₀)-alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,
 and
 where two carbon atoms of the alkanediyl chain in 1,2-, 1,3- or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,
 D is a group of the formula



where
 ##³ marks the linkage site with the nitrogen atom,
 R^1 is hydrogen or methyl,
 R^2 is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

or


R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

#4 marks the linkage site with the adjacent nitrogen atom,

#5 marks the linkage site with the carbonyl group, the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



in which

#6 marks the linkage site with the carbonyl group,


R⁶ is hydrogen, hydroxy or benzyloxy,

R³ is hydrogen or methyl,

R⁴ is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

or

R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

#7 marks the linkage site with the adjacent nitrogen atom,

#8 marks the linkage site with the group T¹,

T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹,

in which

R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

R⁸ is hydrogen or methyl,

R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,


or

R⁸ and R⁹ together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

R¹⁰ is benzoyl,

R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

R⁵ is hydrogen, methyl or a group of the formula



in which

#9 marks the linkage site with —CHC(R²⁶)-T²,

R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,

R²⁶ is hydrogen or hydroxy,

T² is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,


R³⁵ is methyl or hydroxy,

and also their salts, solvates and solvates of the salts.

8. Compounds of the formula (XXXa) according to claim 7, in which

Cys is a cysteine residue which is bonded via the sulphur atom of the side chain via a carbon atom of the succinimide,

L^1 is a bond, linear (C_2-C_6)-alkanediyl, a group of the formula



where


m is a number 2 or 3,

$##^1$ marks the linkage site with the group G,

$##^2$ marks the linkage site with the group B,

L^{1A} is linear (C_2-C_6)-alkanediyl,

B^1 is a group of the formula



in which

$##^5$ marks the linkage site with the group L^{1A} ,

$##^6$ marks the linkage site with the group L^{1B} ,

L^5 is a bond,

L^6 is a bond,

R^{29} is hydrogen,

R^{30} is hydrogen,

R^{31} is hydrogen or methyl,


R^{32} is hydrogen or methyl,

L^{1B} is linear (C_2-C_6)-alkanediyl,

and

where (C_2-C_6)-alkanediyl may be substituted by 1 or 2 methyl substituents,

B is a bond or a group of the formula



where

$*$ marks the linkage site with L^1 ,

$**$ marks the linkage site with L^2 ,

L^3 is a bond or ethane-1,2-diyl,

L^4 is a bond,

R^{14} is hydrogen,

R^{15} is hydrogen,

R^{16} is hydrogen or methyl,

R^{17} is hydrogen or methyl,


or

R^{16} and R^{17} together with the atoms to which they are bonded form piperazinyl ring,

R^{23} is methyl,

R^{24} is hydrogen or methyl,

L^2 is linear (C_2-C_6)-alkanediyl or is a group of the formula




where

p is a number 2 or 3,

$##^3$ marks the linkage site with the group B,

$##^4$ marks the linkage site with the nitrogen atom,

D is a group of the formula



where


$##^3$ marks the linkage site with the nitrogen atom,

R^1 is hydrogen,

R^2 is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or

R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




in which

$##^4$ marks the linkage site with the adjacent nitrogen atom,

$##^5$ marks the linkage site with the carbonyl group,

the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



in which

#⁶ marks the linkage site with the carbonyl group,


R⁶ is hydrogen, hydroxy or benzyloxy,

R³ is hydrogen,

R⁴ is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or

R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

#⁷ marks the linkage site with the adjacent nitrogen atom,

#⁸ marks the linkage site with the group T¹,

T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹,

in which

R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

R⁸ is hydrogen or methyl,

R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,


or

R⁸ and R⁹ together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

R¹⁰ is benzoyl,

R¹¹ is benzyl which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

R⁵ is hydrogen, methyl or a group of the formula



in which

#⁹ marks the linkage site to —CHCH₂phenyl,

R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,

R³⁵ is methyl or hydroxy,


and also their salts, solvates and solvates of the salts.

9. Compounds of the formula (XXXa) according to either of claim 7 or 8, in which

Cys is a cysteine residue which is bonded via the sulphur atom of the side chain via a carbon atom of the succinimide,

L¹ is a bond or linear (C₂-C₆)-alkanediyl,

B is a bond or a group of the formula



where

* marks the linkage site with L¹,

** marks the linkage site with L²,


L³ is a bond,

L⁴ is a bond,

R¹⁶ is hydrogen or methyl,

R¹⁷ is hydrogen or methyl,

L² is linear (C₂-C₆)-alkanediyl or is a group of the formula




where

p is a number 2 or 3,

##³ marks the linkage site with the group B,

##⁴ marks the linkage site with the nitrogen atom,

D is a group of the formula



where


#³ marks the linkage site with the nitrogen atom,

R¹ is hydrogen,

R² is benzyl or 1H-indol-3-ylmethyl,

or

R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




L^5 is a bond or (C₂-C₄)-alkanediyl,
 L^6 is a bond,
 R^{29} is hydrogen or (C₁-C₄)-alkyl,
 R^{30} is hydrogen or (C₁-C₄)-alkyl,
 or
 R^{29} and R^{30} together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 R^{31} is hydrogen or (C₁-C₄)-alkyl,
 R^{32} is hydrogen or (C₁-C₄)-alkyl,
 or
 R^{31} and R^{32} together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,
 L^{1B} is linear (C₂-C₁₀)-alkanediyl,
 and

where (C₁-C₁₀)-alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,

and

where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,

B is a bond or a group of the formula



where

* marks the linkage site with L^1 ,

** marks the linkage site with L^2 ,

P is O or NH,

Q^1 is a 4- to 7-membered heterocycle,

Q^2 is a 3- to 7-membered carbocycle or a 4- to 7-membered heterocycle,

R^{18} is hydrogen or (C₁-C₄)-alkyl,

R^{19} is hydrogen or the side group of a natural α -amino acid or of its homologues or isomers,

R^{20} is hydrogen or (C₁-C₄)-alkyl,

or

R^{19} and R^{20} together with the atoms to which they are bonded form a pyrrolidinyl ring,

R^{21} is hydrogen or (C₁-C₄)-alkyl,


R^{22} is hydrogen or (C₁-C₄)-alkyl,

or

R^{21} and R^{22} together with the atoms to which they are bonded form a 3- to 7-membered carbocycle,

R^{27} is hydrogen or (C₁-C₄)-alkyl,

L^2 is linear (C₂-C₁₀)-alkanediyl or is a group of the formula



where

p is a number from 2 to 6,

##³ marks the linkage site with the group B,


##⁴ marks the linkage site with the nitrogen atom,

where (C₂-C₁₀)-alkanediyl may be substituted by 1 to 4 substituents selected independently of one another from the group consisting of methyl, hydroxy and benzyl,

and

where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,

D is a group of the formula



in which


#³ marks the linkage site with the nitrogen atom,

R^1 is hydrogen or methyl,

R^2 is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

or

R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1, 1-diyl group of the formula




in which

#⁴ marks the linkage site with the adjacent nitrogen atom,

#⁵ marks the linkage site with the carbonyl group,

the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



in which

#6 marks the linkage site with the carbonyl group,


R⁶ is hydrogen, hydroxy or benzyloxy,

R³ is hydrogen or methyl,

R⁴ is isopropyl, isobutyl, sec-butyl, tert-butyl, phenyl, benzyl, 1-hydroxyethyl, 4-hydroxybenzyl, 4-hydroxy-3-nitrobenzyl, 4-hydroxy-3-aminobenzyl, 1-phenylethyl, diphenylmethyl, 1H-imidazol-4-ylmethyl or 1H-indol-3-ylmethyl,

or

R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1, 1-diyl group of the formula



in which

#7 marks the linkage site with the adjacent nitrogen atom,

#8 marks the linkage site with the group T¹,

T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹,

in which

R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

R⁸ is hydrogen or methyl,

R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,



or

R⁸ and R⁹ together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

R¹⁰ is benzoyl,

R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

R⁵ is hydrogen, methyl or a group of the formula



#9 marks the linkage site with —CHC(R²⁶)-T²,

R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,

R²⁶ is hydrogen or hydroxy,


T² is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,

R³⁵ is methyl or hydroxy,

and also their salts, solvates and solvates of the salts.

11. Compounds of the formula (XXXI) according to claim 10, in which

L¹ is a bond, linear (C₂-C₆)-alkanediyl or a group of the formula



where


m is a number 2 or 3,

##1 marks the linkage site with the group G,

##2 marks the linkage site with the group B,

where (C₂-C₆)-alkanediyl may be substituted by 1 or 2 methyl substituents,

B is a bond or a group of the formula



where

* marks the linkage site with L¹,

** marks the linkage site with L²,

R¹⁸ is hydrogen,

R¹⁹ is methyl, propan-2-yl, 2-methylpropan-1-yl or 1-methylpropan-1-yl,

R²⁰ is hydrogen or (C₁-C₄)-alkyl,

or

R¹⁹ and R²⁰ together with the atoms to which they are bonded form a pyrrolidinyl ring,

R²¹ is hydrogen or methyl,


R²² is hydrogen or methyl,

or

R²¹ and R²² together with the atoms to which they are bonded form a cyclopropyl ring,

R²⁷ is hydrogen or methyl,

L^2 is linear (C_2 - C_6)-alkanediyl or is a group of the formula



where

p is a number 2 or 3,

##3 marks the linkage site with the group B,


##4 marks the linkage site with the nitrogen atom,

where (C_2 - C_{10})-alkanediyl may be substituted by 1 or 2 methyl substituents,

and

where two carbon atoms of the alkanediyl chain in 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a phenyl ring,

D is a group of the formula



in which


#3 marks the linkage site with the nitrogen atom,

R^1 is hydrogen,

R^2 is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or

R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




in which

#4 marks the linkage site with the adjacent nitrogen atom,

#5 marks the linkage site with the carbonyl group,

the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



in which

#6 marks the linkage site with the carbonyl group,


R^6 is hydrogen, hydroxy or benzyloxy,

R^3 is hydrogen,

R^4 is 1-hydroxyethyl, benzyl, 4-hydroxybenzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or

R^3 and R^4 together with the carbon atoms to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

#7 marks the linkage site with the adjacent nitrogen atom,

#8 marks the linkage site with the group T^1 ,

T^1 is a group of the formula $-C(=O)-OR^7$, $-C(=O)-NR^8R^9$, $-C(=O)-NH-NH-R^{10}$ or $-CH_2-O-R^{11}$,

in which

R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

R^8 is hydrogen or methyl,

R^9 is hydrogen, methyl, ethyl, n-propyl or benzyl,


or

R^8 and R^9 together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,

R^{10} is benzoyl,

R^{11} is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

R^5 is hydrogen, methyl or a group of the formula




in which


#9 marks the linkage site with $-CHCH_2$ phenyl,

R^{12} is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula $-S(O)_2OH$,


R^{13} is phenyl which may be substituted by methoxy-carbonyl or carboxyl,
 R^{35} is methyl or hydroxy,
 and also their salts, solvates and solvates of the salts.
12. Compounds of the formula (XXXI) according to either of claim 10 or 11, in which
 L^1 is a bond,
 B is a bond,
 L^2 is linear (C_2 - C_6)-alkanediyl or is a group of the formula




where
 p is a number 2 or 3,
 $##^3$ marks the linkage site with the group B,
 $##^4$ marks the linkage site with the nitrogen atom,
 D is a group of the formula




where
 $##^3$ marks the linkage site with the nitrogen atom,
 R^1 is hydrogen,
 R^2 is benzyl or 1H-indol-3-ylmethyl,
 or
 R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which
 $##^4$ marks the linkage site with the adjacent nitrogen atom,
 $##^5$ marks the linkage site with the carbonyl group,
 the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



in which
 $##^6$ marks the linkage site with the carbonyl group,
 R^6 is hydrogen, hydroxy or benzyloxy,
 R^3 is hydrogen,
 R^4 is benzyl or 1H-indol-3-ylmethyl,
 or
 R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which
 $##^7$ marks the linkage site with the adjacent nitrogen atom,
 $##^8$ marks the linkage site with the group T^1 ,
 T^1 is a group of the formula $-C(=O)-OR^7$ or $-C(=O)-NR^8R^9$,
 in which
 R^7 is hydrogen,
 R^8 is hydrogen,
 R^9 is hydrogen,
 R^{35} is methyl,
 and also their salts, solvates and solvates of the salts.

13. Compounds of the formulae (XXXa) and (XXXI) selected from the following group:

N-[6-(3-[[[(2R)-2-amino-2-carboxyethyl]sulphanyl]-2,5-dioxopyrrolidin-1-yl]hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(1S)-1-carboxy-2-(1H-indol-3-yl)ethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide,


N-[6-(3-[[[(2R)-2-amino-2-carboxyethyl]sulphanyl]-2,5-dioxopyrrolidin-1-yl]hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide,

N-(6-[[[(5S)-5-amino-5-carboxypentyl]amino]-6-oxo-hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(2S)-3-(1H-indol-3-yl)-1-(1,2-oxazinan-2-yl)-1-oxopropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide trifluoroacetate,

N-(6-[[[(5S)-5-amino-5-carboxypentyl]amino]-6-oxo-hexyl]-N-methyl-L-valyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[[[(1S)-1-carboxy-2-(1H-indol-3-yl)ethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide,

and also their salts, solvates and solvates of the salts.

14. Binder-drug conjugates of the general formula (I)



in which

n is a number from 1 to 50,

AK is a binder,


the group $\text{\textcircled{G-L}^1\text{-B-L}^2\text{-}}$ is a linker,

where

$\text{\textcircled{}}$ marks the linkage site with the group AK and

$\text{\textcircled{N}}$ marks the linkage site with the nitrogen atom,

D is a group of the formula



where


$\text{\textcircled{N}}$ marks the linkage site with the nitrogen atom,

R^1 is hydrogen,

R^2 is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or


R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

$\text{\textcircled{N}}$ marks the linkage site with the adjacent nitrogen atom,

$\text{\textcircled{C=O}}$ marks the linkage site with the carbonyl group, the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



in which

$\text{\textcircled{C=O}}$ marks the linkage site with the carbonyl group,


R^6 is hydrogen, hydroxy or benzyloxy,

R^3 is hydrogen,

R^4 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or

R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

$\text{\textcircled{N}}$ marks the linkage site with the adjacent nitrogen atom,

$\text{\textcircled{C=O}}$ marks the linkage site with the group T^1 ,

T^1 is a group of the formula —C(=O)—OR^7 , $\text{—C(=O)—NR}^8\text{R}^9$, $\text{—C(=O)—NH—NH—R}^{10}$ or $\text{—CH}_2\text{—O—R}^{11}$,


in which

R^7 is hydrogen, methyl, ethyl, n-propyl, tert-butyl, benzyl or adamantylmethyl,

R^8 is hydrogen or methyl,

R^9 is hydrogen, methyl, ethyl, n-propyl or benzyl,

or
 R^8 and R^9 together with the nitrogen atom to which they are bonded form a 4- to 7-membered heterocycle,
 R^{10} is benzoyl,
 R^{11} is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,
 R^5 is hydrogen, methyl or a group of the formula




in which
 $\#^9$ marks the linkage site with $-\text{CHC}(R^{26})-\text{T}^2$,
 R^{12} is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula $-\text{S}(\text{O})_2\text{OH}$,
 R^{13} is phenyl which may be substituted by methoxycarbonyl or carboxyl,
 R^{26} is hydrogen or hydroxy,
 T^2 is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylmethyl,
 and also their salts, solvates and solvates of the salts.


15. Binder-drug conjugates according to claim **14** with the general formula (I),

in which
 n is a number from 1 to 50,
 AK is AK_1 or AK_2
 where

AK_1 is a binder which is bonded via a sulphur atom of the binder to the group G,
 AK_2 is a binder which is bonded via a nitrogen atom of the binder to the group G,
 G when $\text{AK}=\text{AK}_1$, is a group of the formula




where
 $\#^1$ marks the linkage site with the sulphur atom of the binder,
 $\#^2$ marks the linkage site with the group L^1 ,
 or
 when $\text{AK}=\text{AK}_2$, is carbonyl,
 L^1 is a bond, linear $(\text{C}_1-\text{C}_{10})$ -alkanediyl or is a group of the formula



where


m is a number from 2 to 6,
 $\#^1$ marks the linkage site with the group G,
 $\#^2$ marks the linkage site with the group B,
 where $(\text{C}_1-\text{C}_{10})$ -alkanediyl may be substituted by 1 to 4 methyl substituents,
 and
 where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C_3-C_6) -cycloalkyl ring or a phenyl ring,

B is a bond or a group of the formula



where

* marks the linkage site with L^1 ,
 ** marks the linkage site with L^2 ,
 P is O or NH,
 L^3 is a bond or (C_2-C_4) -alkanediyl,
 L^4 is a bond or a group of the formula



in which

*** marks the linkage site with the carbonyl group,
 **** marks the linkage site with L^2 ,
 R^{25} is hydrogen or methyl,
 Q^1 is a 4- to 7-membered heterocycle,
 Q^2 is a 3- to 7-membered carbocycle or a 4- to 7-membered heterocycle,
 R^{14} is hydrogen or (C_1-C_4) -alkyl,
 R^{15} is hydrogen or (C_1-C_4) -alkyl,

or

R¹⁴ and R¹⁵ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

R¹⁶ is hydrogen or (C₁-C₄)-alkyl,

R¹⁷ is hydrogen or (C₁-C₄)-alkyl,

or

R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

R¹⁸ is hydrogen or (C₁-C₄)-alkyl,

R¹⁹ is hydrogen or the side group of a natural α -amino acid or of its homologues or isomers,

R²⁰ is hydrogen or (C₁-C₄)-alkyl,

or

R¹⁹ and R²⁰ together with the atoms to which they are bonded form a pyrrolidinyl ring,

R²¹ is hydrogen or (C₁-C₄)-alkyl,

R²² is hydrogen or (C₁-C₄)-alkyl,

or


R²¹ and R²² together with the atoms to which they are bonded form a 3- to 7-membered carbocycle,

R²³ is (C₁-C₄)-alkyl,

R²⁴ is hydrogen or (C₁-C₄)-alkyl,

R²⁷ is hydrogen or (C₁-C₄)-alkyl,

L² is linear (C₂-C₁₀)-alkanediyl or is a group of the formula



where

p is a number from 2 to 6,

##³ marks the linkage site with the group B,

##⁴ marks the linkage site with the nitrogen atom,

where (C₂-C₁₀)-alkanediyl may be substituted by 1 to 4 methyl substituents, and

where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,

D has the definitions indicated in claim 14,

and also their salts, solvates and solvates of the salts.

16. Binder-drug conjugates according to claim 14 or 15 of the general formula (I),

in which

n is a number from 1 to 50,

AK is AK₁ or AK₂

where

AK₁ is an antibody or an antigen-binding antibody fragment and are bonded via a sulphur atom to the group G,

AK₂ is an antibody or an antigen-binding antibody fragment and are bonded via a nitrogen atom to the group G,

G, L¹, B, L² and D have the definitions indicated in the claim 14 or 15,

and also their salts, solvates and solvates of the salts.

17. Binder-drug conjugates according to any of claims 14 to 16 of the general formula (I), in which

n is a number from 1 to 20,


AK is AK₁ or AK₂

where

AK₁ is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

AK₂ is an antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the NH side group of a lysine residue of the binder to the group G,

G when AK=AK₁, is a group of the formula



in which


##¹ marks the linkage site with the cysteine residue of the binder,

##² marks the linkage site with the group L¹,

or

when AK=AK₂, is carbonyl,

L¹ is a bond, linear (C₂-C₆)-alkanediyl or is a group of the formula



where


m is a number from 2 to 6,

##¹ marks the linkage site with the group G,

##² marks the linkage site with the group B,

where (C₂-C₆)-alkanediyl may be substituted by 1 or 2 methyl substituents,

B is a bond or a group of the formula




where

* marks the linkage site with L¹,

** marks the linkage site with L²,

L³ is a bond or ethane-1,2-diyl,

L^4 is a bond or a group of the formula



in which

**** marks the linkage site with the carbonyl group,

**** marks the linkage site with L^2 ,

R^{25} is methyl,

Q^1 is a 4- to 6-membered carbocycle or piperidine-1,4-diyl,

R^{14} is hydrogen,

R^{15} is hydrogen,

R^{16} is hydrogen or methyl,

R^{17} is hydrogen or methyl,

or

R^{16} and R^{17} together with the atoms to which they are bonded form a piperazinyl ring,

R^{18} is hydrogen,

R^{19} is hydrogen, methyl, propan-2-yl, 2-methylpropan-1-yl or 1-methylpropan-1-yl,

R^{20} is hydrogen or methyl,

or

R^{19} and R^{20} together with the atoms to which they are bonded form a pyrrolidinyl ring,

R^{21} is hydrogen or methyl,

R^{22} is hydrogen or methyl,


or

R^{21} and R^{22} together with the atoms to which they are bonded form a cyclopropyl ring,

R^{23} is methyl,

R^{24} is hydrogen or methyl,

L^2 is linear (C_2-C_6) -alkanediyl or is a group of the formula



where


p is a number from 2 to 6,

##3 marks the linkage site with the group B,

##4 marks the linkage site with the nitrogen atom,

where (C_2-C_6) -alkanediyl may be substituted by 1 or 2 methyl substituents,

D is a group of the formula



where


#3 marks the linkage site with the nitrogen atom,

R^1 is hydrogen,

R^2 is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or


R^1 and R^2 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

#4 marks the linkage site with the adjacent nitrogen atom,

#5 marks the linkage site with the carbonyl group, the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



in which

#6 marks the linkage site with the carbonyl group,


R^6 is hydrogen, hydroxy or benzyloxy,

R^3 is hydrogen,

R^4 is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or

R^3 and R^4 together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

#7 marks the linkage site with the adjacent nitrogen atom,

#8 marks the linkage site with the group T^1 ,

T¹ is a group of the formula —C(=O)—OR⁷,
—C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or
—CH₂—O—R¹¹,

in which

R⁷ is hydrogen, methyl, ethyl, n-propyl, tert-butyl,
benzyl or adamantylmethyl,

R⁸ is hydrogen or methyl,

R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,


or

R⁸ and R⁹ together with the nitrogen atom to which
they are bonded form a 4- to 7-membered hetero-
cycle,

R¹⁰ is benzoyl,

R¹¹ is benzyl, which may be substituted in the phenyl
group by methoxycarbonyl or carboxyl,

R⁵ is hydrogen, methyl or a group of the formula



in which

#⁹ marks the linkage site with —CHC(R²⁶)-T²,

R¹² is phenyl which may be substituted by methoxy-
carbonyl, carboxyl or a group of the formula
—S(O)₂OH,

R¹³ is phenyl which may be substituted by methoxy-
carbonyl or carboxyl,

R²⁶ is hydrogen or hydroxy,

T² is phenyl, benzyl, 1H-indol-3-yl or 1H-indol-3-ylm-
ethyl,

and also their salts, solvates and solvates of the salts.

18. Binder-drug conjugates according to any of claims **14**
to **17** of the general formula (I)

in which

n is a number from 1 to 10,


AK is AK₁ or AK₂

where

AK₁ is an antibody which comprises the six CDR
sequences of the antibody MF-Ta, the variable light
and variable heavy chain of the antibody MF-Ta or the
light and heavy chain of the antibody MF-Ta, and is
bonded via the sulphur atom of a cysteine residue of
the binder to the group G,

AK₂ is an antibody which comprises the six CDR
sequences of the antibody MF-Ta, the variable light
and variable heavy chain of the antibody MF-Ta or the
light and heavy chain of the antibody MF-Ta, and is
bonded via the NH side group of a lysine residue of
the binder to the group G,

G when AK=AK₁, is a group of the formula



in which


#¹ marks the linkage site with the cysteine residue of the
binder,

#² marks the linkage site with the group L¹,

or

when AK=AK₂, is carbonyl,

L¹ is a bond, linear (C₂-C₆)-alkanediyl or is a group of the
formula



where


m is a number 2 or 3,

##¹ marks the linkage site with the group G,

##² marks the linkage site with the group B,

where (C₂-C₆)-alkanediyl may be substituted by 1 or 2
methyl substituents,

B is a bond or a group of the formula




where

* marks the linkage site with L¹,

** marks the linkage site with L²,

L³ is a bond or ethane-1,2-diyl,

L⁴ is a bond or a group of the formula



in which

*** marks the linkage site with the carbonyl group,

**** marks the linkage site with L²,

R²⁵ is methyl,

Q¹ is piperidine-1,4-diyl,

R¹⁶ is hydrogen or methyl,

R¹⁷ is hydrogen or methyl,

or

R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a piperazinyl ring,

R²¹ is hydrogen or methyl,

R²² is hydrogen or methyl,

or


R²¹ and R²² together with the atoms to which they are bonded form a cyclopropyl ring,

R²³ is methyl,

R²⁴ is hydrogen,

L² is linear (C₂-C₆)-alkanediyl,

D is a group of the formula



where


#3 marks the linkage site with the nitrogen atom,

R¹ is hydrogen,

R² is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or

R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula




in which

#4 marks the linkage site with the adjacent nitrogen atom,

#5 marks the linkage site with the carbonyl group,

the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



in which

#6 marks the linkage site with the carbonyl group,


R⁶ is hydrogen, hydroxy or benzyloxy,

R³ is hydrogen,

R⁴ is benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or

R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

#7 marks the linkage site with the adjacent nitrogen atom,

#8 marks the linkage site with the group T¹,

T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹ or —CH₂—O—R¹¹,

in which


R⁷ is hydrogen, methyl, ethyl, n-propyl, benzyl or adamantylmethyl,

R⁸ is hydrogen or methyl,

R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,

R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

R⁵ is hydrogen or a group of the formula



in which

#9 marks the linkage site with —CHC(R²⁶)phenyl,


R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,


and also their salts, solvates and solvates of the salts.

19. Process for preparing the compounds of the invention according to any of claims **14** to **18**, of the general formula (I), characterized in that a solution of the binder in a buffer


[A] is admixed with a suitable reducing agent, such as, for example, dithiothreitol or tris(2-carboxyethyl)phosphine hydrochloride, and is subsequently reacted with a compound of the formula (II)




in which D, L¹, B and L² each have the definitions indicated in claims 1 to 5, to give a compound of the formula (I-A)



in which n, AK₁, D, L¹, B and L² each have the definitions indicated in claims 1 to 5,
or
[B] is reacted with a compound of the formula (III)



in which D, L¹, B and L² each have the definitions indicated in claims 1 to 5, to give a compound of the formula (I-B)




in which n, AK₂, D, L¹, B and L² each have the definitions indicated in claims 1 to 5.

20. Compounds prepared by the process according to claim 19 or 54, where AK₁ and AK₂ is an antibody which comprises the six CDR sequences of the antibody MF-Ta, the variable light and variable heavy chain of the antibody MF-Ta or the light and heavy chain of the antibody MF-Ta,

and also their salts, solvates and solvates of the salts.


21. Compounds of the formula (XXX)



in which

Cys is a cysteine residue which is bonded via the sulphur atom of the side chain to a carbon atom of the succinimide,

L¹ is a bond, linear (C₁-C₁₀)-alkanediyl or is a group of the formula



where

m is a number from 2 to 6,

##1 marks the linkage site with the group G,



##2 marks the linkage site with the group B,

where (C₁-C₁₀)-alkandiyl may be substituted by 1 to 4 methyl substituents,

and

where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,

B is a bond or a group of the formula



where


* marks the linkage site with L¹,

** marks the linkage site with L²,

P is O or NH,

L³ is a bond or (C₂-C₄)-alkanediyl,

L⁴ is a bond or a group of the formula



in which

*** marks the linkage site with the carbonyl group,

**** marks the linkage site with L²,

R²⁵ is hydrogen or methyl,

Q¹ is a 3- to 7-membered carbocycle or a 4- to 7-membered aza heterocycle,

Q² is a 3- to 7-membered carbocycle or a 4- to 7-membered aza heterocycle,

R¹⁴ is hydrogen or (C₁-C₄)-alkyl,

R¹⁵ is hydrogen or (C₁-C₄)-alkyl,

or

R¹⁴ and R¹⁵ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

R¹⁶ is hydrogen or (C₁-C₄)-alkyl,

R¹⁷ is hydrogen or (C₁-C₄)-alkyl,

or

R¹⁶ and R¹⁷ together with the atoms to which they are bonded form a 5- or 6-membered heterocycle,

R¹⁸ is hydrogen or (C₁-C₄)-alkyl,

R¹⁹ is hydrogen or the side group of a natural α-amino acid or of its homologues or isomers,

R²⁰ is hydrogen or (C₁-C₄)-alkyl,

or

R¹⁹ and R²⁰ together with the atoms to which they are bonded form a pyrrolidinyl ring,

R²¹ is hydrogen or (C₁-C₄)-alkyl,

R²² is hydrogen or (C₁-C₄)-alkyl,

or


R²¹ and R²² together with the atoms to which they are bonded form a 3- to 7-membered carbocycle,

R²³ is (C₁-C₄)-alkyl,

R²⁴ is hydrogen or (C₁-C₄)-alkyl,

R²⁷ is hydrogen or (C₁-C₄)-alkyl,

L² is linear (C₂-C₁₀)-alkanediyl or is a group of the formula



where

p is a number from 2 to 6,

##³ marks the linkage site with the group B,


##⁴ marks the linkage site with the nitrogen atom,

where (C₂-C₁₀)-alkanediyl may be substituted by 1 to 4 methyl substituents,

and

where two carbon atoms of the alkanediyl chain in 1,2, 1,3 or 1,4 relation to one another, with inclusion of any carbon atoms situated between them, may be bridged to form a (C₃-C₆)-cycloalkyl ring or a phenyl ring,

D is a group of the formula



in which


##³ marks the linkage site with the nitrogen atom,

R¹ is hydrogen,

R² is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or


R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1, 1-diyl group of the formula



in which

##⁴ marks the linkage site with the adjacent nitrogen atom,

##⁵ marks the linkage site with the carbonyl group, the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



in which

##⁶ marks the linkage site with the carbonyl group,


R⁶ is hydrogen, hydroxy or benzyloxy,

R³ is hydrogen,

R⁴ is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or

R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1, 1-diyl group of the formula



in which

##⁷ marks the linkage site with the adjacent nitrogen atom,

##⁸ marks the linkage site with the group T¹,

T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹,

in which


#³ marks the linkage site with the nitrogen atom,

R¹ is hydrogen,

R² is 1-hydroxyethyl, benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or


R¹ and R² together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

#⁴ marks the linkage site with the adjacent nitrogen atom,

#⁵ marks the linkage site with the carbonyl group, the ring A with the N—O moiety present therein is a mono- or bicyclic, optionally substituted heterocycle of the formula



in which

#⁶ marks the linkage site with the carbonyl group,


R⁶ is hydrogen, hydroxy or benzyloxy,

R³ is hydrogen,

R⁴ is benzyl, 1-phenylethyl or 1H-indol-3-ylmethyl,

or

R³ and R⁴ together with the carbon atom to which they are bonded form a (1S,2R)-2-phenylcyclopropane-1,1-diyl group of the formula



in which

#⁷ marks the linkage site with the adjacent nitrogen atom,

#⁸ marks the linkage site with the group T¹,

T¹ is a group of the formula —C(=O)—OR⁷, —C(=O)—NR⁸R⁹, —C(=O)—NH—NH—R¹⁰ or —CH₂—O—R¹¹,

in which

R⁷ is hydrogen, methyl, ethyl, n-propyl, benzyl or adamantylmethyl,


R⁸ is hydrogen or methyl,

R⁹ is hydrogen, methyl, ethyl, n-propyl or benzyl,

R¹⁰ is benzoyl,

R¹¹ is benzyl, which may be substituted in the phenyl group by methoxycarbonyl or carboxyl,

R⁵ is hydrogen or a group of the formula



in which

#⁹ marks the linkage site with —CHC(R²⁶)phenyl,

R¹² is phenyl which may be substituted by methoxycarbonyl, carboxyl or a group of the formula —S(O)₂OH,

R¹³ is phenyl which may be substituted by methoxycarbonyl or carboxyl,

and also their salts, solvates and solvates of the salts.

23. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 where the binder binds to a cancer target molecule.

24. Binder-drug conjugate according to any of claim 1 to 6, 14 to 18 or 23, where the binder binds to an extracellular target molecule.

25. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 or 23 to 24, where the binder binds to an extracellular cancer target molecule.

26. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 or 23 to 25, where the target molecule, extracellular target molecule, cancer target molecule or extracellular cancer target molecule is a protein.

27. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 or 23 to 26, where the binder after binding to the extracellular target molecule is internalized by the cell expressing the target molecule.

28. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 or 23 to 27, where the binder is a binder protein.

29. Binder-drug conjugate according to any of claims 1, 2, 14, 15, 20 or 23 to 28, where the binder is an antibody or an antigen-binding antibody fragment thereof or an antibody mimetic.

30. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 or 23 to 29, 10 where the antibody is a monoclonal antibody.

31. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 and 23 to 30, where the antibody is a human, humanized or chimeric antibody.

32. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 or 23 to 31, where the antibody is an intact or a modified intact antibody.

33. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 or 23 to 32, where the antibody is an antibody of the IgG class.

34. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 or 23 to 33, where the binder binds to an extracellular cancer target molecule which is selected from the group consisting of EGF receptor (NP_005219.2), mesothelin (Q13421-3), C4.4a (NP_055215.2) and carboanhydrase IX (CA IX; NP_001207.2).

35. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 or 23 to 34, where the binder binds specifically to an extracellular cancer target molecule which is selected from the group consisting of EGF receptor (NP_005219.2), mesothelin (Q13421-3), C4.4a (NP_055215.2) and carboanhydrase IX (CA IX; NP_001207.2).

36. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 or 23 to 35, where the antibody binds to C4.4a.

37. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 or 23 to 36, where the antibody binds to mesothelin and exhibits invariant binding.

38. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 or 23 to 37, where the antibody binds specifically to mesothelin.

39. Binder-drug conjugate according to any of claims 1 to 6, 14 to 18 or 23 to 38, where the binder is an anti-mesothelin antibody or antigen-binding antibody fragment which comprises the amino acid sequences of the three CDR regions of the light chain and the amino acid sequences of the three CDR regions of the heavy chain of an antibody described in WO 2009068204-A1 (Table 7; pages 61-63).

40. Binder-drug conjugate according to any of claims 1 to 3, 14 to 17 or 23 to 39, where the antimesothelin antibody or antigen-binding antibody fragment is selected from the group consisting of

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody MF-Ta,

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody MF-J,

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody MOR06640,

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the

three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody MF-226, and

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequences of the three CDR regions of the light chain and the sequences of the three CDR regions of the heavy chain of the antibody MOR06626.

41. Binder-drug conjugate according to any of claims 1 to 3, 14 to 17 or 23 to 40, where the antimesothelin antibody or antigen-binding antibody fragment is selected from the group consisting of

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequence of the variable light chain and the sequence of the variable heavy chain of the antibody MF-Ta,

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequence of the variable light chain and the sequence of the variable heavy chain of the antibody MF-J,

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequence of the variable light chain and the sequence of the variable heavy chain of the antibody MOR06640,

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequence of the variable light chain and the sequence of the variable heavy chain of the antibody MF-226, and

anti-mesothelin antibodies or antigen-binding antibody fragments thereof which comprise the sequence of the variable light chain and the sequence of the variable heavy chain of the antibody MOR06626.

42. Antibodies or antigen-binding antibody fragments which bind to mesothelin, where the antibody sequence comprises the amino acid sequence of the variable heavy chain represented by the sequences SEQ ID NO:404 and the amino acid sequence of the variable light chain represented by the sequence SEQ ID NO:405.

43. Antibodies or antigen-binding antibody fragments which bind to mesothelin, where the antibody sequence comprises the amino acid sequence of the variable heavy chain which is encoded by the nucleic acid sequence SEQ ID NO:406, and the amino acid sequence of the variable light chain which is encoded by the nucleic acid sequence SEQ ID NO:407.

44. Antibodies which bind to mesothelin, where the antibody sequence comprises the amino acid sequence of the heavy chain represented by the sequences SEQ ID NO:408 and the amino acid sequence of the light chain represented by the sequence SEQ ID NO:409.

45. Antibodies or antigen-binding antibody fragments according to any of claims 42 to 44, which bind specifically to mesothelin.

46. Binder-drug conjugate according to any of claims 1 to 3, 14 to 17 or 23 to 38, where the anti-mesothelin antibody comprises the amino acid sequence of the heavy chain represented by the sequences SEQ ID NO:408 and the amino acid sequence of the light chain represented by the sequence SEQ ID NO:409.

47. Binder-drug conjugate or compounds according to any of claim 1 to 18, 21 to 41 or 46 for the treatment and/or prophylaxis of illnesses.

48. Binder-drug conjugate or a compound according to any of claim **1** to **18**, **21** to **41** or **46** for use in a method for the treatment and/or prophylaxis of hyperproliferative and/or angiogenic diseases.

49. Binder-drug conjugate or a compound according to any of claim **1** to **18**, **21** to **41** or **46** for producing a medicament for the treatment and/or prophylaxis of hyperproliferative and/or angiogenic diseases.

50. Medicament comprising a binder-drug conjugate or a compound according to any of claims **1** to **18**, **21** to **41** or **46** to **48**, in combination with an inert, non-toxic, pharmaceutically suitable excipient.


51. Medicament comprising a binder-drug conjugate or a compound according to any of claims **1** to **18**, **21** to **41** or **46** to **48**, in combination with one or more anti-hyperproliferative, cytostatic or cytotoxic substances.

52. Medicament according to claim **50** or **51** for the treatment and/or prophylaxis of hyperproliferative and/or angiogenic diseases.


53. Method for the treatment and/or prophylaxis of hyperproliferative and/or angiogenic diseases in humans and animals, using an effective amount of at least one binder-drug conjugate or a compound according to any of claims **1** to **18** or **21** to **41**, or of a medicament as defined in either of claims **50** and **51**.

54. Process for preparing the compounds of the invention according to any of claims **1** to **6**, of the general formula (Ia), characterized in that a solution of the binder in a buffer

[A] is admixed with a suitable reducing agent, such as, for example, dithiothreitol or tris(2-carboxyethyl)phosphine hydrochloride, and is subsequently reacted with a compound of the formula (IIa)




in which D, L¹, B, L² and R³⁵ each have the definitions indicated in claims **1** to **6**, to give a compound of the formula (Ia-A)




in which n, AK₁, D, L¹, B, L² and R³⁵ each have the definitions indicated in claims **1** to **6**.

or

[B] is reacted with a compound of the formula (IIIa)




in which D, L¹, B, L² and R³⁵ each have the definitions indicated in claims 1 to 6, to give a compound of the formula (Ia-B)




in which n, AK₂, D, L¹, B, L² and R³⁵ each have the definitions indicated in claims 1 to 6.

55. Binder-drug conjugates selected from the following compounds:



-continued



where in each case

n is a number from 2 to 8, preferably 2 to 5,

AK_1 is a human or humanized antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the sulphur atom of a cysteine residue of the binder to the group G,

AK_2 is a human or humanized antibody or an antigen-binding antibody fragment which binds to mesothelin and is bonded via the NH side group of a lysine residue of the binder to the group G.

* * * * *