



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2722085 C 2014/02/11

(11)(21) **2 722 085**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(22) Date de dépôt/Filing Date: 2008/07/09

(41) Mise à la disp. pub./Open to Public Insp.: 2009/01/18

(45) Date de délivrance/Issue Date: 2014/02/11

(62) Demande originale/Original Application: 2 637 084

(30) Priorité/Priority: 2007/07/18 (EP07290901.3)

(51) Cl.Int./Int.Cl. *A01N 43/80* (2006.01),
A01N 43/64 (2006.01), *A01P 1/00* (2006.01)

(72) Inventeurs/Inventors:
EL A'MMA, BEVERLY JEAN, US;
HEER, BEAT, CH;
LEVY, RICHARD, FR;
PAREEK, KIRAN, US

(73) Propriétaire/Owner:
ROHM AND HAAS COMPANY, US

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : COMPOSITION MICROBICIDE

(54) Title: AN N-METHYL-1,2 BENZISOTHIAZOLIN-3-ONE MICROBICIDAL COMPOSITION

(57) Abrégé/Abstract:

Synergistic microbicidal compositions containing N-methyl-1,2- benzisothiazolin- 3-one.



ABSTRACT OF THE DISCLOSURE

Synergistic microbicidal compositions containing N-methyl-1,2-benzisothiazolin-3-one.

An N-Methyl-1,2 Benzisothiazolin-3-one Microbicidal Composition

This invention relates to a synergistic combination of selected microbicides having greater activity than would be observed for the individual microbicides.

5 In some cases, commercial microbicides cannot provide effective control of microorganisms, even at high use concentrations, due to weak activity against certain types of microorganisms, e.g., those resistant to some microbicides, or due to aggressive environmental conditions. Combinations of different microbicides are sometimes used to provide overall control of microorganisms in
10 a particular end use environment. For example, U.S. Pat. App. Pub. No. 2007/0078118 discloses synergistic combinations of N-methyl-1,2-benzisothiazolin-3-one (MBIT) with other biocides. However, there is a need for additional combinations of microbicides having enhanced activity against various strains of microorganisms to provide effective control of the microorganisms.
15 Moreover, there is a need for combinations containing lower levels of individual microbicides for environmental and economic benefit. The problem addressed by this invention is to provide such additional combinations of microbicides.

STATEMENT OF THE INVENTION

20 The present invention is directed to a microbicidal composition comprising: (a) N-methyl-1,2-benzisothiazolin-3-one; and (b) at least one microbicide selected from the group consisting of hexahydro-1,3,5-tris(2-hydroxyethyl)-s-triazine; 4,4-dimethyloxazolidine; sodium hydroxymethylglycinate; myristamidopropyl PG-dimoonium chloride phosphate;
25 octanoic acid; 2-ethylhexyl glycerin; glyceryl monocaprylate; glyceryl monocaprate; 7 α -ethyldihydro-1H, 3H, 5H-oxazolo (3,4-C) oxazole; and 2-(thiocyanomethylthio)benzothiazole.

DETAILED DESCRIPTION OF THE INVENTION

As used herein, the following terms have the designated definitions, unless the context clearly indicates otherwise. "MBIT" is N-methyl-1,2-benzisothiazolin-3-one. The term "microbicide" refers to a compound capable of killing, inhibiting the growth of or controlling the growth of microorganisms at a locus; microbicides include bactericides, fungicides and algacides. The term "microorganism" includes, for example, fungi (such as yeast and mold), bacteria and algae. The term "locus" refers to an industrial system or product subject to contamination by microorganisms. The following abbreviations are used throughout the specification: ppm = parts per million by weight (weight/weight), mL = milliliter, ATCC = American Type Culture Collection, MBC = minimum biocidal concentration, and MIC = minimum inhibitory concentration. Unless otherwise specified, temperatures are in degrees centigrade (°C), and references to percentages (%) are by weight. Amounts of organic microbicides are given on an active ingredient basis in ppm (w/w).

The compositions of the present invention unexpectedly have been found to provide enhanced microbicidal efficacy at a combined active ingredient level lower than that of the individual microbicides. Additional microbicides beyond those listed in the claims may be present in the composition.

In one embodiment of the invention, the antimicrobial composition comprises N-methyl-1,2-benzisothiazolin-3-one and hexahydro-1,3,5-tris(2-hydroxyethyl)-s-triazine. Preferably, a weight ratio of hexahydro-1,3,5-tris(2-hydroxyethyl)-s-triazine to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.0014 to 1:150, more preferably from 1:0.012 to 1:150.

In one embodiment of the invention, the antimicrobial composition comprises N-methyl-1,2-benzisothiazolin-3-one and 4,4-dimethyloxazolidine. Preferably, a weight ratio of 4,4-dimethyloxazolidine to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.0011 to 1:3.5.

In one embodiment of the invention, the antimicrobial composition comprises N-methyl-1,2-benzisothiazolin-3-one and sodium hydroxymethylglycinate. Preferably, a weight ratio of sodium hydroxymethylglycinate to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.0005

to 1:2.3, more preferably from 1:0.14 to 1:2.3, more preferably from 1:0.29 to 1:2.3.

In one embodiment of the invention, the antimicrobial composition comprises N-methyl-1,2-benzisothiazolin-3-one and myristamidopropyl PG-dimoonium chloride phosphate. Preferably, a weight ratio of myristamidopropyl PG-dimoonium chloride phosphate to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.0011 to 1:150.

In one embodiment of the invention, the antimicrobial composition comprises N-methyl-1,2-benzisothiazolin-3-one and octanoic acid. Preferably, a weight ratio of octanoic acid to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.0046 to 1:0.68, more preferably from 1:0.029 to 1:0.68.

In one embodiment of the invention, the antimicrobial composition comprises N-methyl-1,2-benzisothiazolin-3-one and 2-ethylhexyl glycerin. Preferably, a weight ratio of 2-ethylhexyl glycerin to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.017 to 1:2.3.

In one embodiment of the invention, the antimicrobial composition comprises N-methyl-1,2-benzisothiazolin-3-one and glyceryl monocaprylate. Preferably, a weight ratio of glyceryl monocaprylate to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.0004 to 1:75, more preferably from 1:0.012 to 1:75.

In one embodiment of the invention, the antimicrobial composition comprises N-methyl-1,2-benzisothiazolin-3-one and glyceryl monocaprato. Preferably, a weight ratio of glyceryl monocaprato to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.0031 to 1:8.8, more preferably from 1:0.017 to 1:8.8.

In one embodiment of the invention, the antimicrobial composition comprises N-methyl-1,2-benzisothiazolin-3-one and 7 α -ethyldihydro-1H, 3H, 5H-oxazolo (3,4-C) oxazole. Preferably, a weight ratio of 7 α -ethyldihydro-1H, 3H, 5H-oxazolo (3,4-C) oxazole to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.0003 to 1:7.1.

In one embodiment of the invention, the antimicrobial composition comprises N-methyl-1,2-benzisothiazolin-3-one and 2-(thiocyanomethylthio)

benzothiazole. Preferably, a weight ratio of 2-(thiocyanomethylthio) benzothiazole to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.063 to 1:143.

The microbicides in the composition of this invention may be used "as is" or may first be formulated with a solvent or a solid carrier. Suitable solvents include, for example, water; glycols, such as ethylene glycol, propylene glycol, diethylene glycol, dipropylene glycol, polyethylene glycol, and polypropylene glycol; glycol ethers; alcohols, such as methanol, ethanol, propanol, phenethyl alcohol and phenoxypropanol; ketones, such as acetone and methyl ethyl ketone; esters, such as ethyl acetate, butyl acetate, triacetyl citrate, and glycerol triacetate; carbonates, such as propylene carbonate and dimethyl carbonate; and mixtures thereof. It is preferred that the solvent is selected from water, glycols, glycol ethers, esters and mixtures thereof. Suitable solid carriers include, for example, cyclodextrin, silicas, diatomaceous earth, waxes, cellulosic materials, alkali and alkaline earth (e.g., sodium, magnesium, potassium) metal salts (e.g., chloride, nitrate, bromide, sulfate) and charcoal.

When a microbicide component is formulated in a solvent, the formulation may optionally contain surfactants. When such formulations contain surfactants, they are generally in the form of emulsive concentrates, emulsions, microemulsive concentrates, or microemulsions. Emulsive concentrates form emulsions upon the addition of a sufficient amount of water. Microemulsive concentrates form microemulsions upon the addition of a sufficient amount of water. Such emulsive and microemulsive concentrates are generally well known in the art; it is preferred that such formulations are free of surfactants. U.S. Patent No. 5,444,078 may be consulted for further general and specific details on the preparation of various microemulsions and microemulsive concentrates.

A microbicide component also can be formulated in the form of a dispersion. The solvent component of the dispersion can be an organic solvent or water, preferably water. Such dispersions can contain adjuvants, for example, co-solvents, thickeners, anti-freeze agents, dispersants, fillers, pigments, surfactants, biodispersants, sulfosuccinates, terpenes, furanones, polycations, stabilizers, scale inhibitors and anti-corrosion additives.

When both microbicides are each first formulated with a solvent, the solvent used for the first microbicide may be the same as or different from the solvent used to formulate the other commercial microbicide, although water is preferred for most industrial biocide applications. It is preferred that the two solvents are miscible.

Those skilled in the art will recognize that the microbicide components of the present invention may be added to a locus sequentially, simultaneously, or may be combined before being added to the locus. It is preferred that the first microbicide and the second microbicide component be added to a locus simultaneously or sequentially. When the microbicides are added simultaneously or sequentially, each individual component may contain adjuvants, such as, for example, solvent, thickeners, anti-freeze agents, colorants, sequestrants (such as ethylenediamine-tetraacetic acid, ethylenediaminedisuccinic acid, iminodisuccinic acid and salts thereof), dispersants, surfactants, biodispersants, sulfosuccinates, terpenes, furanones, polycations, stabilizers, scale inhibitors and anti-corrosion additives.

The microbicidal compositions of the present invention can be used to inhibit the growth of microorganisms or higher forms of aquatic life (such as protozoans, invertebrates, bryozoans, dinoflagellates, crustaceans, mollusks, etc.) by introducing a microbicidally effective amount of the compositions onto, into, or at a locus subject to microbial attack. Suitable loci include, for example: industrial process water; electrocoat deposition systems; cooling towers; air washers; gas scrubbers; mineral slurries; wastewater treatment; ornamental fountains; reverse osmosis filtration; ultrafiltration; ballast water; evaporative condensers; heat exchangers; pulp and paper processing fluids and additives; starch; plastics; emulsions; dispersions; paints; latices; coatings, such as varnishes; construction products, such as mastics, caulks, and sealants; construction adhesives, such as ceramic adhesives, carpet backing adhesives, and laminating adhesives; industrial or consumer adhesives; photographic chemicals; printing fluids; household products, such as bathroom and kitchen cleaners and sanitary wipes; cosmetics; toiletries; shampoos; soaps; detergents; industrial cleaners; floor polishes; laundry rinse water; metalworking fluids; conveyor

lubricants; hydraulic fluids; leather and leather products; textiles; textile products; wood and wood products, such as plywood, chipboard, wallboard, flakeboard, laminated beams, oriented strandboard, hardboard, and particleboard; petroleum processing fluids; fuel; oilfield fluids, such as injection water, fracture fluids, and drilling muds; agriculture adjuvant preservation; surfactant preservation; medical devices; diagnostic reagent preservation; food preservation, such as plastic or paper food wrap; food, beverage, and industrial process pasteurizers; toilet bowls; recreational water; pools; and spas.

Preferably, the microbicidal compositions of the present invention are used to inhibit the growth of microorganisms at a locus selected from one or more of mineral slurries, pulp and paper processing fluids and additives, starch, emulsions, dispersions, paints, latices, coatings, construction adhesives, such as ceramic adhesives, carpet backing adhesives, photographic chemicals, printing fluids, household products such as bathroom and kitchen cleaners and sanitary wipes, cosmetics, toiletries, shampoos, soaps, detergents, industrial cleaners, floor polishes, laundry rinse water, metal working fluids, textile products, wood and wood products, agriculture adjuvant preservation, surfactant preservation, diagnostic reagent preservation, food preservation, and food, beverage, and industrial process pasteurizers.

The specific amount of the composition of this invention necessary to inhibit or control the growth of microorganisms and higher aquatic life forms in a locus depends upon the particular locus to be protected. Typically, the amount of the composition of the present invention to control the growth of microorganisms in a locus is sufficient if it provides from 0.1 to 1,000 ppm of the isothiazoline ingredient of the composition in the locus. It is preferred that the isothiazolone ingredients of the composition be present in the locus in an amount of at least 0.5 ppm, more preferably at least 4 ppm and most preferably at least 10 ppm. It is preferred that the isothiazolone ingredients of the composition be present in the locus in an amount of no more than 1000 ppm, more preferably no more than 500 ppm, and most preferably no more than 200 ppm.

EXAMPLES

Materials and Methods

The synergism of the combination of the present invention was demonstrated by testing a wide range of concentrations and ratios of the compounds.

One measure of synergism is the industrially accepted method described by Kull, F.C.; Eisman, P.C.; Sylwestrowicz, H.D. and Mayer, R.L., in Applied Microbiology 9:538-541 (1961), using the ratio determined by the formula:

$$Q_a/Q_A + Q_b/Q_B = \text{Synergy Index ("SI")}$$

wherein:

Q_A = concentration of compound A (first component) in ppm, acting alone, which produced an end point (MIC of Compound A).

Q_a = concentration of compound A in ppm, in the mixture, which produced an end point.

Q_B = concentration of compound B (second component) in ppm, acting alone, which produced an end point (MIC of Compound B).

Q_b = concentration of compound B in ppm, in the mixture, which produced an end point.

When the sum of Q_a/Q_A and Q_b/Q_B is greater than one, antagonism is indicated. When the sum is equal to one, additivity is indicated, and when less than one, synergism is demonstrated. The lower the SI, the greater the synergy shown by that particular mixture. The minimum inhibitory concentration (MIC) of a microbicide is the lowest concentration tested under a specific set of conditions that prevents the growth of added microorganisms.

Synergy tests were conducted using standard microtiter plate assays with media designed for optimal growth of the test microorganism. Minimal salt medium supplemented with 0.2% glucose and 0.1% yeast extract (M9GY medium) was used for testing bacteria; Potato Dextrose Broth (PDB medium) was used for testing yeast and mold. In this method, a wide range of combinations of microbicides was tested by conducting high resolution MIC

assays in the presence of various concentrations of MBIT. High resolution MICs were determined by adding varying amounts of microbicide to one column of a microtitre plate and doing subsequent ten-fold dilutions using an automated liquid handling system to obtain a series of endpoints ranging from 2 ppm to 10,000 ppm active ingredient.

The synergy of the combinations of the present invention was determined against several microorganisms, as described in the Tables below. The bacteria were used at a concentration of about 5×10^6 bacteria per mL and the yeast and mold at 5×10^5 fungi per mL. These microorganisms are representative of natural contaminants in many consumer and industrial applications. The plates were visually evaluated for microbial growth (turbidity) to determine the MIC after various incubation times at 25° C (yeast and mold) or 30° C (bacteria).

The test results for demonstration of synergy of the MBIT combinations of the present invention are shown below in Tables 1 through 10. In each test, Second Component (B) was MBIT and the First Component (A) was the other commercial microbicide. Each table shows the specific combinations of MBIT and the other component; results against the microorganisms tested with incubation times; the end-point activity in ppm measured by the MIC for MBIT alone (Q_B), for the other component alone (Q_A), for MBIT in the mixture (Q_b) and for the other component in the mixture (Q_a); the calculated SI value; and the range of synergistic ratios for each combination tested (other component/MBIT or A/B).

Table: 1

Ca: ppm AI of BIOBAN GK(hexahydro-1,3,5-tris(2-hydroxyethyl)-s-triazine)

Cb: ppm AI of MBIT(N-methyl-1,2-benzisothiazolin-3-one)

Ratio:Ca:Cb

5

Test Organisms	Contact Time	Ca	Cb	S.I.	Ca:Cb
<i>A. niger</i> ATCC # 16404	3 days	425	-	-	-
		-	37.5	-	-
		110	18.8	0.76	1:0.1709
		200	18.8	0.97	1:0.0940
		300	18.8	1.21	1:0.0627
		300	9.4	0.96	1:0.0313
		425	9.4	1.25	1:0.0221
		300	4.7	0.83	1:0.0157
		425	4.7	1.13	1:0.0111
		300	2.4	0.77	1:0.0080
		425	2.4	1.06	1:0.0056
		300	1.2	0.74	1:0.0040
		425	1.2	1.03	1:0.0028
	7 days	650	-	-	-
		-	150	-	-
		42.5	75	0.57	1:1.7647
		65	75	0.60	1:1.1538
		110	75	0.67	1:0.6818
		200	75	0.81	1:0.3750
		300	75	0.96	1:0.2500
		400	75	1.12	1:0.1875
		425	18.8	0.78	1:0.0442
		525	18.8	0.93	1:0.0358
		650	9.4	1.06	1:0.0145
		525	4.7	0.84	1:0.0090
<i>C. albicans</i> ATCC # 10231	48 hrs	650	4.7	1.03	1:0.0072
		525	2.4	0.82	1:0.0046
		650	2.4	1.02	1:0.0037
		650	1.2	1.01	1:0.0018
		875	-	-	-
		-	30	-	-
		87.5	15	0.60	1:0.1714
		110	15	0.63	1:0.1364
		200	15	0.73	1:0.0750
		300	15	0.84	1:0.0500
		425	15	0.99	1:0.0353
		110	7.5	0.38	1:0.0682
		200	7.5	0.48	1:0.0375
		300	7.5	0.59	1:0.0250
		425	7.5	0.74	1:0.0176
		525	7.5	0.85	1:0.0143
		650	7.5	0.99	1:0.0115
		425	3.75	0.61	1:0.0088
		525	3.75	0.73	1:0.0071
		650	3.75	0.87	1:0.0058

* Trade-mark

	72 hrs	875	3.75	1.13	1:0.0043
		650	1.86	0.80	1:0.0029
		875	1.86	1.06	1:0.0021
		650	0.94	0.77	1:0.0014
		875	0.94	1.03	1:0.0011
		875	-	-	-
		-	30	-	-
		87.5	15	0.60	1:0.1714
		110	15	0.63	1:0.1364
		200	15	0.73	1:0.0750
		300	15	0.84	1:0.0500
		425	15	0.99	1:0.0353
		425	7.5	0.74	1:0.0176
		525	7.5	0.85	1:0.0143
		650	7.5	0.99	1:0.0115
		875	1.86	1.06	1:0.0021
		875	0.94	1.03	1:0.0011
<i>Ps. aeruginosa</i> ATCC#9027	24hrs	55	-	-	-
		-	75	-	-
		32.5	37.5	1.09	1:1.1538
		43.75	18.75	1.05	1:0.4286
		43.75	9.4	0.92	1:0.2149
		43.75	4.6	0.86	1:0.1051
	48hrs	55	-	-	-
		-	125	-	-
		26.25	75	1.08	1:2.8571
		32.5	37.5	0.89	1:1.1538
		43.75	37.5	1.10	1:0.8571
		55	4.6	1.04	1:0.0836
<i>S.aureus</i> ATCC#6538	24hrs	55	-	-	-
		-	15	-	-
		15	7.5	0.77	1:0.5000
		32.5	3.8	0.84	1:0.1169
		43.75	3.8	1.05	1:0.0869
		32.5	1.8	0.71	1:0.0554
		43.75	1.8	0.92	1:0.0411
		55	1.8	1.12	1:0.0327
		55	0.94	1.06	1:0.0171
	48hrs	55	-	-	-
		-	30	-	-
		0.1	15	0.50	1:150.0000
		1.5	15	0.53	1:10.0000
		2.625	15	0.55	1:5.7143
		4.375	15	0.58	1:3.4286
		5.5	15	0.60	1:2.7273
		10	15	0.68	1:1.5000
		15	15	0.77	1:1.0000
		21.25	15	0.89	1:0.7059
		26.25	15	0.98	1:0.5714
		32.5	15	1.09	1:0.4615
		21.25	7.5	0.64	1:0.3529
		26.25	7.5	0.73	1:0.2857

	32.5	7.5	0.84	1:0.2308
	43.75	7.5	1.05	1:0.1714
	32.5	3.8	0.72	1:0.1169
	43.75	3.8	0.92	1:0.0869
	55	3.8	1.13	1:0.0691
	32.5	1.8	0.65	1:0.0554
	55	1.8	1.06	1:0.0327
	43.75	0.94	0.83	1:0.0215
	55	0.94	1.03	1:0.0171

Table: 2

*

Ca: ppm AI of BIOBAN 1135 (4,4-dimethyloxazolidine)

Cb: ppm AI of MBIT(N-methyl-1,2-benzisothiazolin-3-one)

5 Ratio:Ca:Cb

Test Organisms	Contact Time	Ca	Cb	S.I.	Ca:Cb
<i>A. niger</i> ATCC # 16404	3 days	525	-	-	-
		-	37.5	-	-
		200	18.8	0.88	1:0.0940
		300	18.8	1.07	1:0.0627
		300	9.4	0.82	1:0.0313
		425	9.4	1.06	1:0.0221
		425	4.7	0.93	1:0.0111
		525	4.7	1.13	1:0.0090
		425	2.4	0.87	1:0.0056
		525	2.4	1.06	1:0.0046
	7 days	425	1.2	0.84	1:0.0028
		525	1.2	1.03	1:0.0023
		875	-	-	-
		-	150	-	-
		525	75	1.1	1:0.1429
		650	37.5	0.99	1:0.0577
		875	9.4	1.06	1:0.0107
		650	4.7	0.77	1:0.0072
		875	4.7	1.03	1:0.0054
		650	2.4	0.76	1:0.0037
<i>C. albicans</i> ATCC # 10231	48 hrs	875	2.4	1.02	1:0.0027
		875	1.2	1.01	1:0.0014
		1100	-	-	-
		-	30	-	-
		425	15	0.89	1:0.0353
		525	15	0.98	1:0.0286
		650	15	1.09	1:0.0231
		650	7.5	0.84	1:0.0115
		875	7.5	1.05	1:0.0086
		875	3.75	0.92	1:0.0043
		1100	3.75	1.13	1:0.0034
		875	1.86	0.86	1:0.0021
		1100	1.86	1.06	1:0.0017
		875	0.94	0.83	1:0.0011

* Trade-mark

	72 hrs	1100	0.94	1.03	1:0.0009
		1100	-	-	-
		-	30	-	-
		425	15	0.89	1:0.0353
		525	15	0.98	1:0.0286
		650	15	1.09	1:0.0231
		875	7.5	1.05	1:0.0086
		875	3.75	0.92	1:0.0043
		1100	3.75	1.13	1:0.0034
		875	1.86	0.86	1:0.0021
		1100	1.86	1.06	1:0.0017
		875	0.94	0.83	1:0.0011
		1100	0.94	1.03	1:0.0009
<i>Ps. aeruginosa</i> ATCC#9027	24hrs	100	-	-	-
		-	75	-	-
		55	37.5	1.05	1:0.6818
		55	9.4	0.68	1:0.1709
		100	9.4	1.13	1:0.0940
		55	4.6	0.61	1:0.0836
		100	4.6	1.06	1:0.0460
	48hrs	100	-	-	-
		-	125	-	-
		21.25	75	0.81	1:3.5294
		26.25	75	0.86	1:2.8571
		32.5	75	0.93	1:2.3077
		43.75	75	1.04	1:1.7143
		55	37.5	0.85	1:0.6818
		100	9.4	1.08	1:0.0940
		100	4.6	1.04	1:0.0460
<i>S. aureus</i> ATCC#6538	24hrs	55	-	-	-
		-	30	-	-
		21.25	15	0.89	1:0.7059
		26.25	15	0.98	1:0.5714
		32.5	15	1.09	1:0.4615
		26.25	7.5	0.73	1:0.2857
		32.5	7.5	0.84	1:0.2308
		43.75	7.5	1.05	1:0.1714
		43.75	3.8	0.92	1:0.0869
		55	3.8	1.13	1:0.0691
		55	1.8	1.06	1:0.0327
		55	0.94	1.03	1:0.0171
	48hrs	55	-	-	-
		-	30	-	-
		21.25	15	0.89	1:0.7059
		26.25	15	0.98	1:0.5714
		32.5	15	1.09	1:0.4615
		26.25	7.5	0.73	1:0.2857
		32.5	7.5	0.84	1:0.2308
		43.75	7.5	1.05	1:0.1714
		43.75	3.8	0.92	1:0.0869
		55	3.8	1.13	1:0.0691
		55	1.8	1.06	1:0.0327

		55	0.94	1.03	1:0.0171
--	--	----	------	------	----------

Table: 3

*
Ca: ppm AI of SUTTOCIDE A (sodium hydroxymethylglycinate)

5 Cb: ppm AI of MBIT(N-methyl-1,2-benzisothiazolin-3-one)

Ratio:Ca:Cb

Test Organisms	Contact Time	Ca	Cb	S.I.	Ca:Cb
<i>A. niger</i> ATCC # 16404	3 days	875	-	-	-
		-	75	-	-
		425	18.8	0.74	1:0.0442
		525	18.8	0.85	1:0.0358
		650	18.8	0.99	1:0.0289
		425	9.4	0.61	1:0.0221
		525	9.4	0.73	1:0.0179
		650	9.4	0.87	1:0.0145
		875	9.4	1.13	1:0.0107
		650	4.7	0.81	1:0.0072
		875	4.7	1.06	1:0.0054
		525	2.4	0.63	1:0.0046
		650	2.4	0.77	1:0.0037
		875	2.4	1.03	1:0.0027
		650	1.2	0.76	1:0.0018
		875	1.2	1.02	1:0.0014
	7 days	1100	-	-	-
		-	150	-	-
		110	75	0.60	1:0.6818
		200	75	0.68	1:0.3750
		300	75	0.77	1:0.2500
		425	75	0.89	1:0.1765
		525	75	0.98	1:0.1429
		650	75	1.09	1:0.1154
		425	37.5	0.64	1:0.0882
		525	37.5	0.73	1:0.0714
		650	37.5	0.84	1:0.0577
		875	37.5	1.05	1:0.0429
		525	18.8	0.6	1:0.0358
		650	18.8	0.72	1:0.0289
		875	18.8	0.92	1:0.0215
		1100	18.8	1.13	1:0.0171
		525	9.4	0.54	1:0.0179
		650	9.4	0.65	1:0.0145
		875	9.4	0.86	1:0.0107
		1100	9.4	1.06	1:0.0085
		650	4.7	0.62	1:0.0072
		875	4.7	0.83	1:0.0054
		1100	4.7	1.03	1:0.0043
		650	2.4	0.61	1:0.0037
		875	2.4	0.81	1:0.0027

* Trade-mark

		1100	2.4	1.02	1:0.0022
		650	1.2	0.60	1:0.0018
		875	1.2	0.80	1:0.0014
		1100	1.2	1.01	1:0.0011
<i>C.albicans</i> ATCC # 10231	48 hrs	3000	-	-	-
		-	30	-	-
		525	15	0.68	1:0.0286
		650	15	0.72	1:0.0231
		875	15	0.79	1:0.0171
		1100	15	0.87	1:0.0136
		2000	15	1.17	1:0.0075
		525	7.5	0.43	1:0.0143
		650	7.5	0.47	1:0.0115
		875	7.5	0.54	1:0.0086
		1100	7.5	0.62	1:0.0068
		2000	7.5	0.92	1:0.0038
		3000	7.5	1.25	1:0.0025
		875	3.75	0.42	1:0.0043
		1100	3.75	0.49	1:0.0034
		2000	3.75	0.79	1:0.0019
		3000	3.75	1.13	1:0.0013
		2000	1.86	0.73	1:0.0009
		3000	1.86	1.06	1:0.0006
		2000	0.94	0.70	1:0.0005
	72 hrs	3000	0.94	1.03	1:0.0003
		3000	-	-	-
		-	30	-	-
		525	15	0.68	1:0.0286
		650	15	0.72	1:0.0231
		875	15	0.79	1:0.0171
		1100	15	0.87	1:0.0136
		2000	15	1.17	1:0.0075
		3000	15	1.5	1:0.0050
		525	7.5	0.43	1:0.0143
		650	7.5	0.47	1:0.0115
		875	7.5	0.54	1:0.0086
		1100	7.5	0.62	1:0.0068
		2000	7.5	0.92	1:0.0038
		3000	7.5	1.25	1:0.0025
		875	3.75	0.42	1:0.0043
		1100	3.75	0.49	1:0.0034
		2000	3.75	0.79	1:0.0019
		3000	3.75	1.13	1:0.0013
		2000	1.86	0.73	1:0.0009
		3000	1.86	1.06	1:0.0006
		2000	0.94	0.70	1:0.0005
		3000	0.94	1.03	1:0.0003
<i>Ps. aeruginosa</i> ATCC#9027	24hrs	87.5	-	-	-
		-	75	-	-
	48 hrs	65	9.4	0.87	1:0.1446
		87.5	9.4	1.13	1:0.1074
		110	-	-	-

		-	125	-	-
		87.5	9.4	0.87	1:0.1074
		110	9.4	1.08	1:0.0855
		110	4.6	1.04	1:0.0418
<i>S.aureus</i> ATCC#6538	24hrs	110	-	-	-
		-	15	-	-
		30	7.5	0.77	1:0.2500
		42.5	7.5	89	1:0.1765
		52.5	7.5	98	1:0.1429
		65	7.5	1.09	1:0.1154
		52.5	3.8	0.73	1:0.0724
		65	3.8	0.84	1:0.0585
		87.5	3.8	1.05	1:0.0434
		110	0.94	1.06	1:0.0085
	48hrs	110	-	-	-
		-	30	-	-
		6.5	15	0.56	1:2.3077
		8.75	15	0.58	1:1.7143
		11	15	0.6	1:1.3636
		20	15	0.68	1:0.7500
		30	15	0.77	1:0.5000
		42.5	15	0.89	1:0.3529
		52.5	15	0.98	1:0.2857
		65	15	1.09	1:0.2308
		42.5	7.5	0.64	1:0.1765
		52.5	7.5	0.73	1:0.1429
		65	7.5	0.84	1:0.1154
		87.5	7.5	1.05	1:0.0857
		65	3.8	0.72	1:0.0585
		110	1.8	1.06	1:0.0164
		110	0.94	1.03	1:0.0085

Table:4

Ca: ppm AI of Phospholipid PTM(myristamidopropyl PG-dimoonium chloride phosphate)

5 Cb: ppm AI of MBIT(N-methyl-1,2-benzisothiazolin-3-one)

Ratio:Ca:Cb

Test Organisms	Contact Time	Ca	Cb	S.I.	Ca:Cb
<i>A. niger</i> ATCC # 16404	3 days	1100	-	-	-
		-	37.5	-	-
		425	18.8	0.89	1:0.0442
		525	18.8	0.98	1:0.0358
		650	18.8	1.09	1:0.0289
		525	9.4	0.73	1:0.0179
		650	9.4	0.84	1:0.0145
		875	9.4	1.05	1:0.0107
		1100	2.4	1.06	1:0.0022
		1100	1.2	1.03	1:0.0011
	7 days	1100	-	-	-

		-	150	-	-
		2	75	0.50	1:37.5000
		20	75	0.52	1:3.7500
		200	75	0.68	1:0.3750
		525	75	0.98	1:0.1429
		650	75	1.09	1:0.1154
		20	37.5	0.27	1:1.8750
		52.5	37.5	0.3	1:0.7143
		87.5	37.5	0.33	1:0.4286
		200	37.5	0.43	1:0.1875
		425	37.5	0.64	1:0.0882
		650	37.5	0.84	1:0.0577
		875	37.5	1.05	1:0.0429
		650	18.8	0.72	1:0.0289
		875	18.8	0.92	1:0.0215
		1100	18.8	1.13	1:0.0171
		650	9.4	0.65	1:0.0145
		875	9.4	0.86	1:0.0107
		1100	9.4	1.06	1:0.0085
		1100	4.7	1.03	1:0.0043
		1100	2.4	1.02	1:0.0022
		1100	1.2	1.01	1:0.0011
<i>C.albicans</i> ATCC # 10231	48 hrs	1100	-	-	-
		-	30	-	-
		525	15	0.98	1:0.0286
		650	15	1.09	1:0.0231
		525	7.5	0.73	1:0.0143
		650	7.5	0.84	1:0.0115
		875	7.5	1.05	1:0.0086
		650	3.75	0.72	1:0.0058
		875	3.75	0.92	1:0.0043
		1100	3.75	1.13	1:0.0034
		875	1.86	0.86	1:0.0021
		1100	1.86	1.06	1:0.0017
		875	0.94	0.83	1:0.0011
		1100	0.94	1.03	1:0.0009
	72 hrs	1100	-	-	-
		-	60	-	-
		2	30	0.50	1:15.0000
		20	30	0.52	1:1.5000
		110	30	0.60	1:0.2727
		200	30	0.68	1:0.1500
		300	30	0.77	1:0.1000
		425	30	0.89	1:0.0706
		525	30	0.98	1:0.0571
		650	30	1.09	1:0.0462
		525	15	0.73	1:0.0286
		650	15	0.84	1:0.0231
		525	7.5	0.6	1:0.0143
		650	7.5	0.72	1:0.0115
		875	7.5	0.92	1:0.0086
		650	3.75	0.65	1:0.0058

		875	3.75	0.86	1:0.0043
		1100	3.75	1.06	1:0.0034
		875	1.86	0.83	1:0.0021
		1100	1.86	1.03	1:0.0017
		875	0.94	0.81	1:0.0011
		1100	0.94	1.02	1:0.0009
<i>Ps. aeruginosa</i> ATCC#9027	24hrs	425	-	-	-
		-	75	-	-
		425	4.6	1.06	1:0.0108
	48 hrs	525	-	-	-
		-	125	-	-
		525	4.6	1.04	1:0.0088
<i>S.aureus</i> ATCC#6538	24hrs	10	-	-	-
		-	15	-	-
		2.625	7.5	0.76	1:2.8571
		3.25	7.5	0.83	1:2.3077
		4.375	7.5	0.94	1:1.7143
		5.5	7.5	1.05	1:1.3636
		4.375	3.8	0.69	1:0.8686
		5.5	3.8	0.80	1:0.6909
		10	3.8	1.25	1:0.3800
		4.375	1.8	0.56	1:0.4114
		5.5	1.8	0.67	1:0.3273
		10	1.8	1.12	1:0.1800
		10	0.94	1.06	1:0.0940
	48hrs	10	-	-	-
		-	30	-	-
		0.1	15	0.51	1:150.0000
		0.5	15	0.55	1:30.0000
		1	15	0.6	1:15.0000
		1.5	15	0.65	1:10.0000
		2.125	15	0.71	1:7.0588
		2.625	15	0.76	1:5.7143
		3.25	15	0.83	1:4.6154
		4.375	15	0.94	1:3.4286
		5.5	15	1.05	1:2.7273
		4.375	7.5	0.69	1:1.7143
		5.5	7.5	0.80	1:1.3636
		10	7.5	1.25	1:0.7500
		4.375	3.8	0.56	1:0.8686
		5.5	3.8	0.68	1:0.6909
		10	3.8	1.13	1:0.3800
		5.5	1.8	0.61	1:0.3273
		10	1.8	1.06	1:0.1800
		10	0.94	1.03	1:0.0940

Table: 5

Ca: ppm AI of Caprylic acid (octanoic acid)

Cb: ppm AI of MBIT(N-methyl-1,2-benzisothiazolin-3-one)

Ratio:Ca:Cb

5

Test Organisms	Contact Time	Ca	Cb	S.I.	Ca:Cb
<i>A. niger</i> ATCC # 16404	3 days	650	-	-	-
		-	75	-	-
		425	18.8	0.9	1:0.0442
		525	18.8	1.06	1:0.0358
		525	9.4	0.93	1:0.0179
		650	9.4	1.13	1:0.0145
		525	4.7	0.87	1:0.0090
		525	2.4	0.84	1:0.0046
		650	2.4	1.03	1:0.0037
	7 days	650	1.2	1.02	1:0.0018
		875	-	-	-
		-	150	-	-
		110	75	0.63	1:0.6818
		200	75	0.73	1:0.3750
		300	75	0.84	1:0.2500
		425	75	0.99	1:0.1765
		650	37.5	0.99	1:0.0577
		425	18.8	0.61	1:0.0442
		650	18.8	0.87	1:0.0289
		875	18.8	1.13	1:0.0215
		875	9.4	1.06	1:0.0107
		875	4.7	1.03	1:0.0054
		875	2.4	1.02	1:0.0027
		875	1.2	1.01	1:0.0014
<i>C.albicans</i> ATCC # 10231	48 hrs	65	-	-	-
		-	30	-	-
		42.5	7.5	0.9	1:0.1765
		52.5	7.5	1.06	1:0.1429
		65	1.86	1.06	1:0.0286
	72 hrs	65	0.94	1.03	1:0.0145
		87.5	-	-	-
		-	30	-	-
		42.5	15	0.99	1:0.3529
		42.5	7.5	0.74	1:0.1765
		52.5	7.5	0.85	1:0.1429
		65	7.5	0.99	1:0.1154
		65	3.75	0.87	1:0.0577
		87.5	3.75	1.13	1:0.0429
		87.5	1.86	1.06	1:0.0213
		87.5	0.94	1.03	1:0.0107
<i>Ps. aeruginosa</i> ATCC#9027	48 hrs	>11000	-	-	-
		-	125	-	-
		8750	75	1.4	1:0.0086
		11000	37.5	1.3	1:0.0034

		11000	18.75	1.15	1:0.0017
		11000	9.4	1.08	1:0.0009
		11000	4.6	1.04	1:0.0004
<i>S.aureus</i> ATCC#6538	24hrs	425	-	-	-
		-	15	-	-
		425	0.94	1.06	1:0.0022
	48hrs	425	-	-	-
		-	30	-	-
		425	1.8	1.06	1:0.0042
		425	0.94	1.03	1:0.0022

Table: 6

Ca: ppm AI of SENSIVA SC 50(3[(2-ethylhexyl)oxy]1,2-propanediol) (2-ethylhexyl glycerin)

5 Cb: ppm AI of MBIT(N-methyl-1,2-benzisothiazolin-3-one)

Ratio:Ca:Cb

Test Organisms	Contact Time	Ca	Cb	S.I.	Ca:Cb
<i>A. niger</i> ATCC # 16404	3 days	875	-	-	-
		-	37.5	-	-
		650	9.4	0.99	1:0.0145
		650	4.7	0.87	1:0.0072
		875	4.7	1.13	1:0.0054
		875	2.4	1.06	1:0.0027
		875	1.2	1.03	1:0.0014
	7 days	1100	-	-	-
		-	150	-	-
		87.5	75	0.58	1:0.8571
		110	75	0.60	1:0.6818
		200	75	0.68	1:0.3750
		300	75	0.77	1:0.2500
		425	75	0.89	1:0.1765
		525	75	0.98	1:0.1429
		650	75	1.09	1:0.1154
		650	37.5	0.84	1:0.0577
		1100	9.4	1.06	1:0.0085
		1100	4.7	1.03	1:0.0043
		1100	2.4	1.02	1:0.0022
		1100	1.2	1.01	1:0.0011
<i>C.albicans</i> ATCC # 10231	48 hrs	2000	-	-	-
		-	30	-	-
		6.5	15	0.50	1:2.3077
		11	15	0.51	1:1.3636
		110	15	0.56	1:0.1364
		200	15	0.60	1:0.0750
		300	15	0.65	1:0.0500
		650	15	0.83	1:0.0231
		875	15	0.94	1:0.0171
		1100	15	1.05	1:0.0136
		2000	1.86	1.06	1:0.0009

* Trade-mark

	72 hrs	2000	0.94	1.03	1:0.0005
		2000	-	-	-
		-	30	-	-
		425	15	0.71	1:0.0353
		525	15	0.76	1:0.0286
		650	15	0.83	1:0.0231
		875	15	0.94	1:0.0171
		1100	15	1.05	1:0.0136
		2000	1.86	1.06	1:0.0009
		2000	0.94	1.03	1:0.0005
<i>Ps. aeruginosa</i> ATCC#9027	24hrs	11000	-	-	-
		-	75	-	-
		8750	37.5	1.3	1:0.0043
	48 hrs	11000	-	-	-
		-	125	-	-
		4250	75	0.99	1:0.0176
<i>S.aureus</i> ATCC#6538	24hrs	2000	-	-	-
		-	30	-	-
		8.75	15	0.5	1:1.7143
		20	15	0.51	1:0.7500
		42.5	15	0.52	1:0.3529
		65	15	0.53	1:0.2308
		110	15	0.56	1:0.1364
		200	15	0.6	1:0.0750
		425	15	0.71	1:0.0353
		525	15	0.76	1:0.0286
		650	15	0.83	1:0.0231
		875	15	0.94	1:0.0171
		1100	15	1.05	1:0.0136
		2000	1.8	1.06	1:0.0009
		2000	0.94	1.03	1:0.0005
	48hrs	2000	-	-	-
		-	30	-	-
		8.75	15	0.50	1:1.7143
		20	15	0.51	1:0.7500
		42.5	15	0.52	1:0.3529
		65	15	0.53	1:0.2308
		110	15	0.56	1:0.1364
		200	15	0.60	1:0.0750
		425	15	0.71	1:0.0353
		525	15	0.76	1:0.0286
		650	15	0.83	1:0.0231
		875	15	0.94	1:0.0171
		1100	15	1.05	1:0.0136
		2000	1.8	1.06	1:0.0009
		2000	0.94	1.03	1:0.0005

Table: 7

Ca: ppm AI of CAPMUL MCM C8 (glyceryl monocaprylate)

Cb: ppm AI of MBIT(N-methyl-1,2-benzisothiazolin-3-one)

Ratio:Ca:Cb

5

Test Organisms	Contact Time	Ca	Cb	S.I.	Ca:Cb
<i>A. niger</i> ATCC # 16404	3 days	4250	-	-	-
		-	75	-	-
		2000	37.5	0.97	1:0.0188
		3000	37.5	1.21	1:0.0125
		2000	18.8	0.72	1:0.0094
		3000	18.8	0.96	1:0.0063
		4250	18.8	1.25	1:0.0044
		2000	9.4	0.60	1:0.0047
		3000	9.4	0.83	1:0.0031
		4250	9.4	1.13	1:0.0022
		2000	4.7	0.53	1:0.0024
		3000	4.7	0.77	1:0.0016
		4250	4.7	1.06	1:0.0011
		2000	2.4	0.5	1:0.0012
		3000	2.4	0.74	1:0.0008
		4250	2.4	1.03	1:0.0006
	7 days	3000	1.2	0.72	1:0.0004
		4250	1.2	1.02	1:0.0003
		4250	-	-	-
		-	150	-	-
		1100	75	0.76	1:0.0682
		2000	75	0.97	1:0.0375
		3000	75	1.21	1:0.0250
		2000	37.5	0.72	1:0.0188
		3000	37.5	0.96	1:0.0125
		4250	37.5	1.25	1:0.0088
		2000	18.8	0.60	1:0.0094
		3000	18.8	0.83	1:0.0063
		4250	18.8	1.13	1:0.0044
		2000	9.4	0.53	1:0.0047
		3000	9.4	0.77	1:0.0031
		4250	9.4	1.06	1:0.0022
<i>C.albicans</i> ATCC # 10231	48 hrs	200	-	-	-
		-	30	-	-
		0.2	15	0.50	1:75.0000
		2	15	0.51	1:7.5000
		20	15	0.60	1:0.7500
		42.5	15	0.71	1:0.3529
		65	15	0.83	1:0.2308

* Trade-mark

	72 hrs	87.5	15	0.94	1:0.1714
		110	15	1.05	1:0.1364
		200	1.86	1.06	1:0.0093
		200	0.94	1.03	1:0.0047
		200	-	-	-
		-	30	-	-
		0.2	15	0.50	1:75.0000
		2	15	0.51	1:7.5000
		20	15	0.6	1:0.7500
		42.5	15	0.71	1:0.3529
		65	15	0.83	1:0.2308
		87.5	15	0.94	1:0.1714
		110	15	1.05	1:0.1364
		200	1.86	1.06	1:0.0093
		200	0.94	1.03	1:0.0047
<i>Ps. aeruginosa</i> ATCC#9027	48 hrs	>11000	-	-	-
		-	125	-	-
		2000	75	0.78	1:0.0375
		3000	75	0.87	1:0.0250
		4250	75	0.99	1:0.0176
<i>S. aureus</i> ATCC#6538	24hrs	650	-	-	-
		-	30	-	-
		0.2	15	0.50	1:75.0000
		2	15	0.50	1:7.5000
		6.5	15	0.51	1:2.3077
		11	15	0.52	1:1.3636
		20	15	0.53	1:0.7500
		30	15	0.55	1:0.5000
		42.5	15	0.57	1:0.3529
		52.5	15	0.58	1:0.2857
		65	15	0.60	1:0.2308
		87.5	15	0.63	1:0.1714
		110	15	0.67	1:0.1364
		200	15	0.81	1:0.0750
		300	15	0.96	1:0.0500
	48hrs	425	15	1.15	1:0.0353
		425	7.5	0.9	1:0.0176
		525	7.5	1.06	1:0.0143
		875	-	-	-
		-	30	-	-
		0.2	15	0.50	1:75.0000
		2	15	0.50	1:7.5000
		6.5	15	0.51	1:2.3077
		11	15	0.51	1:1.3636
		20	15	0.52	1:0.7500
		30	15	0.53	1:0.5000
		42.5	15	0.55	1:0.3529
		52.5	15	0.56	1:0.2857
		65	15	0.57	1:0.2308
		87.5	15	0.60	1:0.1714
		110	15	0.63	1:0.1364
		200	15	0.73	1:0.0750

	300	15	0.84	1:0.0500
	425	15	0.99	1:0.0353
	650	7.5	0.99	1:0.0115
	875	1.8	1.06	1:0.0021
	875	0.94	1.03	1:0.0011

Table:8

Ca: ppm AI of CapmulMCM C10 (Mono/diglyceride of capric acid (Glyceryl Monocaprate))

5 Cb: ppm AI of MBIT(N-methyl-1,2-benzisothiazolin-3-one)

Ratio:Ca:Cb

Test Organisms	Contact Time	Ca	Cb	S.I.	Ca:Cb
<i>A. niger</i> ATCC # 16404	3 days	2000	-	-	-
		-	75	-	-
		4.25	37.5	0.50	1:8.8235
		6.5	37.5	0.50	1:5.7692
		11	37.5	0.51	1:3.4091
		30	37.5	0.52	1:1.2500
		65	37.5	0.53	1:0.5769
		110	37.5	0.56	1:0.3409
		200	37.5	0.60	1:0.1875
		300	37.5	0.65	1:0.1250
		425	37.5	0.71	1:0.0882
		525	37.5	0.76	1:0.0714
		650	37.5	0.83	1:0.0577
		875	37.5	0.94	1:0.0429
		1100	37.5	1.05	1:0.0341
		1100	18.8	0.8	1:0.0171
		2000	4.7	1.06	1:0.0024
		2000	2.4	1.03	1:0.0012
	7 days	2000	1.2	1.02	1:0.0006
		2000	-	-	-
		-	150	-	-
		525	75	0.76	1:0.1429
		650	75	0.83	1:0.1154
		875	75	0.94	1:0.0857
		1100	75	1.05	1:0.0682
		1100	37.5	0.8	1:0.0341
		1100	18.8	0.68	1:0.0171
		2000	9.4	1.06	1:0.0047
<i>C. albicans</i> ATCC # 10231	48 hrs	2000	4.7	1.03	1:0.0024
		2000	2.4	1.02	1:0.0012
		2000	1.2	1.01	1:0.0006
		425	-	-	-
		-	30	-	-
		87.5	15	0.71	1:0.1714
		110	15	0.76	1:0.1364
		200	15	0.97	1:0.0750
		300	15	1.21	1:0.0500

* Trade-mark

		200	7.5	0.72	1:0.0375
		300	7.5	0.96	1:0.0250
		425	7.5	1.25	1:0.0176
		200	3.75	0.60	1:0.0188
		300	3.75	0.83	1:0.0125
		425	3.75	1.13	1:0.0088
		200	1.86	0.53	1:0.0093
		300	1.86	0.77	1:0.0062
		425	1.86	1.06	1:0.0044
		200	0.94	0.50	1:0.0047
		300	0.94	0.74	1:0.0031
		425	0.94	1.03	1:0.0022
	72 hrs	425	-	-	-
		-	30	-	-
		87.5	15	0.71	1:0.1714
		110	15	0.76	1:0.1364
		200	15	0.97	1:0.0750
		300	15	1.21	1:0.0500
		200	7.5	0.72	1:0.0375
		300	7.5	0.96	1:0.0250
		425	7.5	1.25	1:0.0176
		200	3.75	0.60	1:0.0188
		300	3.75	0.83	1:0.0125
		425	3.75	1.13	1:0.0088
		200	1.86	0.53	1:0.0093
		300	1.86	0.77	1:0.0062
		425	1.86	1.06	1:0.0044
		200	0.94	0.50	1:0.0047
		300	0.94	0.74	1:0.0031
		425	0.94	1.03	1:0.0022
<i>Ps. aeruginosa</i> ATCC#9027	24hrs	11000	-	-	-
		-	75	-	-
		-	37.5	-	-
		-	18.75	-	-
		-	9.4	-	-
		-	4.6	-	-
	48 hrs	11000	-	-	-
		-	125	-	-
		-	75	-	-
		-	37.5	-	-
		-	18.75	-	-
		-	9.4	-	-
		-	4.6	-	-
<i>S.aureus</i> ATCC#6538	24hrs	200	-	-	-
		-	30	-	-
		5.25	15	0.53	1:2.8571
		11	15	0.56	1:1.3636
		20	15	0.60	1:0.7500
		30	15	0.65	1:0.5000
		42.5	15	0.71	1:0.3529
		52.5	15	0.76	1:0.2857
		65	15	0.83	1:0.2308

.25

	48hrs	87.5	15	0.94	1:0.1714
		110	15	1.05	1:0.1364
		200	-	-	-
		-	30	-	-
		65	15	0.83	1:0.2308
		87.5	15	0.94	1:0.1714
		110	15	1.05	1:0.1364

Table:9

*

Ca: ppm AI of BIOBAN1246 (7 α -ethyldihydro-1H, 3H, 5H-oxazolo (3,4-C) oxazole)

5 Cb: ppm AI of MBIT(N-methyl-1,2-benzisothiazolin-3-one)

Ratio:Ca:Cb

Test Organisms	Contact Time	Ca	Cb	S.I.	Ca:Cb
<i>A. niger</i> ATCC # 16404	3 days	650	-	-	-
		-	37.5	-	-
		200	18.8	0.81	1:0.0940
		300	18.8	0.96	1:0.0627
		425	18.8	1.16	1:0.0442
		425	9.4	0.90	1:0.0221
		525	9.4	1.06	1:0.0179
		425	4.7	0.78	1:0.0111
		525	4.7	0.93	1:0.0090
		650	4.7	1.13	1:0.0072
		425	2.4	0.72	1:0.0056
		525	2.4	0.87	1:0.0046
		650	2.4	1.06	1:0.0037
		425	1.2	0.69	1:0.0028
		525	1.2	0.84	1:0.0023
		650	1.2	1.03	1:0.0018
	7 days	875	-	-	-
		-	150	-	-
		425	75	0.99	1:0.1765
		650	37.5	0.99	1:0.0577
		525	18.8	0.73	1:0.0358
		650	18.8	0.87	1:0.0289
		875	18.8	1.13	1:0.0215
		875	9.4	1.06	1:0.0107
		875	4.7	1.03	1:0.0054
		875	2.5	1.02	1:0.0029
		875	1.2	1.01	1:0.0014
<i>C. albicans</i> ATCC # 10231	48 hrs	3000	-	-	-
		-	30	-	-
		875	15	0.79	1:0.0171
		1100	15	0.87	1:0.0136
		2000	15	1.17	1:0.0075
		875	7.5	0.54	1:0.0086
		1100	7.5	0.62	1:0.0068
		2000	7.5	0.92	1:0.0038

* Trade-mark

	72 hrs	3000	7.5	1.25	1:0.0025
		2000	3.75	0.79	1:0.0019
		3000	3.75	1.13	1:0.0013
		2000	1.86	0.73	1:0.0009
		3000	1.86	1.06	1:0.0006
		3000	0.94	1.03	1:0.0003
		4250	-	-	-
		-	30	-	-
		875	15	0.71	1:0.0171
		1100	15	0.76	1:0.0136
		2000	15	0.97	1:0.0075
		3000	15	1.21	1:0.0050
		875	7.5	0.46	1:0.0086
		1100	7.5	0.51	1:0.0068
		2000	7.5	0.72	1:0.0038
		3000	7.5	0.96	1:0.0025
		4250	7.5	1.25	1:0.0018
		2000	3.75	0.6	1:0.0019
		3000	3.75	0.83	1:0.0013
		4250	3.75	1.13	1:0.0009
		2000	1.86	0.53	1:0.0009
		3000	0.94	0.74	1:0.0003
		4250	0.94	1.03	1:0.0002
<i>Ps. aeruginosa</i> ATCC#9027	24hrs	100	-	-	-
		-	75	-	-
		26.25	75	1.26	1:2.8571
		43.75	37.5	0.94	1:0.8571
		55	37.5	1.05	1:0.6818
		55	18.75	0.80	1:0.3409
		100	18.75	1.25	1:0.1875
		55	9.4	0.68	1:0.1709
		100	9.4	1.13	1:0.0940
		100	4.6	1.06	1:0.0460
	48 hrs	150	-	-	-
		-	125	-	-
		26.25	75	0.78	1:2.8571
		32.5	75	0.82	1:2.3077
		43.75	75	0.89	1:1.7143
		55	75	0.97	1:1.3636
		100	75	1.27	1:0.7500
		55	37.5	0.67	1:0.6818
		100	37.5	0.97	1:0.3750
		150	37.5	1.3	1:0.2500
		100	18.75	0.82	1:0.1875
		150	18.75	1.15	1:0.1250
		150	9.4	1.08	1:0.0627
		150	4.6	1.04	1:0.0307
<i>S. aureus</i> ATCC#6538	24hrs	55	-	-	-
		-	30	-	-
		2.125	15	0.54	1:7.0588
		5.5	15	0.60	1:2.7273
		10	15	0.68	1:1.5000

		26.25	15	0.98	1:0.5714
		32.5	15	1.09	1:0.4615
		21.25	7.5	0.64	1:0.3529
		26.25	7.5	0.73	1:0.2857
		32.5	7.5	0.84	1:0.2308
		43.75	7.5	1.05	1:0.1714
		32.5	3.8	0.72	1:0.1169
		43.75	3.8	0.92	1:0.0869
		55	3.8	1.13	1:0.0691
		55	1.8	1.06	1:0.0327
		55	0.94	1.03	1:0.0171
	48hrs	55	-	-	-
		-	30	-	-
		2.125	15	0.54	1:7.0588
		5.5	15	0.60	1:2.7273
		10	15	0.68	1:1.5000
		26.25	15	0.98	1:0.5714
		32.5	15	1.09	1:0.4615
		21.25	7.5	0.64	1:0.3529
		26.25	7.5	0.73	1:0.2857
		32.5	7.5	0.84	1:0.2308
		43.75	7.5	1.05	1:0.1714
		32.5	3.8	0.72	1:0.1169
		43.75	3.8	0.92	1:0.0869
		55	3.8	1.13	1:0.0691
		55	1.8	1.06	1:0.0327
		55	0.94	1.03	1:0.0171

Table:10

Ca: ppm AI of TCMTB (2-(thiocyanomethylthio) benzothiazole)

Cb: ppm AI of MBIT(N-methyl-1,2-benzisothiazolin-3-one)

5 Ratio:Ca:Cb

Test Organisms	Contact Time	Ca	Cb	S.I.	Ca:Cb
<i>A. niger</i> ATCC # 16404	3 days	3	-	-	-
		-	37.5	-	-
	7 days	3	2.4	1.06	1:0.8000
		3	1.2	1.03	1:0.4000
		8.75	-	-	-
		-	150	-	-
		0.525	75	0.56	1:142.8571
		0.65	75	0.57	1:115.3846
		0.875	75	0.60	1:85.7143
		1.1	75	0.63	1:68.1818
		2	75	0.73	1:37.5000
		3	75	0.84	1:25.0000
		4.25	75	0.99	1:17.6471
		8.75	9.4	1.06	1:1.0743
		6.5	2.4	0.76	1:0.3692
		8.75	2.4	1.02	1:0.2743

28 .

		8.75	1.2	1.01	1:0.1371
<i>C.albicans</i> ATCC # 10231	48 hrs	4.25	-	-	-
		-	30	-	-
	72 hrs	4.25	1.86	1.06	1:0.4376
		4.25	0.94	1.03	1:0.2212
		5.25	-	-	-
		-	60	-	-
		1.1	30	0.71	1:27.2727
		2	30	0.88	1:15.0000
		3	30	1.07	1:10.0000
		4.25	15	1.06	1:3.5294
		4.25	7.5	0.93	1:1.7647
		5.25	3.75	1.06	1:0.7143
		4.25	1.86	0.84	1:0.4376
		5.25	1.86	1.03	1:0.3543
		4.25	0.94	0.83	1:0.2212
		5.25	0.94	1.02	1:0.1790
<i>Ps. aeruginosa</i> ATCC#9027	48 hrs	212.5	-	-	-
		-	125	-	-
		21.25	75	0.70	1:3.5294
		26.25	75	0.72	1:2.8571
		43.75	75	0.81	1:1.7143
		55	75	0.86	1:1.3636
		100	75	1.07	1:0.7500
		150	37.5	1.01	1:0.2500
		150	18.75	0.86	1:0.1250
		212.5	18.75	1.15	1:0.0882
		150	9.4	0.78	1:0.0627
		212.5	9.4	1.08	1:0.0442
		212.5	4.6	1.04	1:0.0216
<i>S.aureus</i> ATCC#6538	24hrs	26.25	-	-	-
		-	15	-	-
		26.25	0.94	1.06	1:0.0358

CLAIMS

1. A microbicidal composition comprising:
 - (a) N-methyl-1,2-benzisothiazolin-3-one; and
 - (b) hexahydro-1,3,5-tris(2-hydroxyethyl)-s-triazine.
2. The microbicidal composition of claim 1 in which a weight ratio of hexahydro-1,3,5-tris(2-hydroxyethyl)-s-triazine to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.0014 to 1:150.
3. The microbicidal composition of claim 1 in which a weight ratio of hexahydro-1,3,5-tris(2-hydroxyethyl)-s-triazine to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.012 to 1:150.

4. The microbicidal composition of claim 1 in which a weight ratio of hexahydro-1,3,5-tris(2-hydroxyethyl)-s-triazine to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.0014 to 1:0.1714.
5. The microbicidal composition of claim 1 in which a weight ratio of hexahydro-1,3,5-tris(2-hydroxyethyl)-s-triazine to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.1169 to 1:0.5000
6. The microbicidal composition of claim 1 in which a weight ratio of hexahydro-1,3,5-tris(2-hydroxyethyl)-s-triazine to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.2500 to 1:1.7647
7. The microbicidal composition of claim 1 in which a weight ratio of hexahydro-1,3,5-tris(2-hydroxyethyl)-s-triazine to N-methyl-1,2-benzisothiazolin-3-one is from 1:0.5714 to 1:150.