



(72) HILLMAN, JENNIFER L., US

(72) CORLEY, NEIL C., US

(72) SHAH, PURVI, US

(71) INCYTE PHARMACEUTICALS, INC., US

(51) Int.Cl.⁷ C12N 5/12, C07K 14/705, C12Q 1/68, C12N 5/10, C07K 16/28,
A61K 38/17

(30) 1997/09/12 (08/928,612) US

(54) **RECEPTEUR SIGMA HUMAIN**

(54) **HUMAN SIGMA RECEPTOR**

(57) L'invention concerne un récepteur sigma humain (SIGR) et des polynucléotides identifiant et codant ce SIGR. L'invention concerne également des vecteurs d'expression, des cellules hôtes, des agonistes, des anticorps et des antagonistes. L'invention concerne en outre des procédés qui permettent de traiter les troubles associés à l'expression du SIGR.

(57) The invention provides a human sigma receptor (SIGR) and polynucleotides which identify and encode SIGR. The invention also provides expression vectors, host cells, agonists, antibodies and antagonists. The invention also provides methods for treating disorders associated with expression of SIGR.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 5/12, 5/10, C07K 14/705, 16/28, C12Q 1/68, A61K 38/17	A1	(11) International Publication Number: WO 99/13055 (43) International Publication Date: 18 March 1999 (18.03.99)
(21) International Application Number: PCT/US98/18941 (22) International Filing Date: 11 September 1998 (11.09.98) (30) Priority Data: 08/928,612 12 September 1997 (12.09.97) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/928,612 (CIP) Filed on 12 September 1997 (12.09.97) (71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HILLMAN, Jennifer, L. [US/US]; 230 Monroe Drive #12, Mountain View, CA 94040 (US). CORLEY, Neil, C. [US/US]; 1240 Dale Avenue #30, Mountain View, CA 94040 (US). SHAH, Purvi [IN/US]; 1608 Queen Charlotte #5, Sunnyvale, CA 94087 (US).		(74) Agents: BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: HUMAN SIGMA RECEPTOR		
(57) Abstract The invention provides a human sigma receptor (SIGR) and polynucleotides which identify and encode SIGR. The invention also provides expression vectors, host cells, agonists, antibodies and antagonists. The invention also provides methods for treating disorders associated with expression of SIGR.		

HUMAN SIGMA RECEPTOR

TECHNICAL FIELD

5 This invention relates to nucleic acid and amino acid sequences of a human sigma receptor and to the use of these sequences in the diagnosis, prevention, and treatment of vesicle trafficking, immunological, and neoplastic disorders.

BACKGROUND OF THE INVENTION

10 The activity of neurons in both the central and peripheral nervous system is affected by a large number of neuropeptides and neurohormones that act upon cells quite distant from their site of release. Neuropeptides perform many functions in the central nervous system (CNS) as neurotransmitters, neuromodulators, and neurotrophic factors. Neurohormones can modify the ability of nerve cells to respond to synaptic neurotransmitters.

15 Morphine and its phamacoanalogs bind selectively at many recognition sites throughout the body to produce pharmacologic effects. Brain loci involved in the transmission of pain and in the alteration of reactivity to painful stimuli appear to be the primary, but not the only sites, at which opioids act. In general, sites that display a high affinity for exogenous opioid ligands also contain high concentrations of several endogenous
20 peptides having opioid-like properties. These peptides have been named endorphins and are derived from larger precursor proteins. The peptide precursor molecules for the endorphins are present at brain sites that have been implicated in pain modulation and also in many other tissues. Although endorphins share many features in chemistry and pharmacology, they possess distinct differences with respect to their biochemical and neuronal pathways. For
25 example, they can be activated during stress such as that produced by pain or the anticipation of pain.

Activation of endorphins results in their interaction with cell surface receptors of the CNS. Both endogenous and exogenous opioids can produce opioid receptor mediated analgesia at sites outside the CNS. This analgesic action may occur because mu (μ), kappa
30 (κ), and delta (δ) receptors located on primary afferent neurons are activated. A fourth receptor, the sigma (σ) receptor, is more controversial but may be related to the dysphoric, hallucinogenic, and cardiac stimulant effects of opioids.

The σ binding sites have preferential affinity for the dextrorotary isomers of certain opiate benzomorphans, such as (+)SKF 10047, (+)cyclazocine, and (+)pentazocine and also for some narcoleptics such as haloperidol. The σ receptor has at least two subtypes, which may be discriminated by stereoselective isomers of these pharmacologically active drugs. SKF 10047
5 has nanomolar affinity for σ_1 site, and has micromolar affinity for the σ_2 site. Haloperidol has similar affinities for both subtypes. Endogenous σ ligands are not known, although progesterone has been suggested to be one of them. Possible σ -site-mediated drug effects include modulation of glutamate receptor function, neurotransmitter response, neuroprotection, behavior, and cognition (Quirion, R. et al. (1992) Trends Pharmacol. Sci.
10 13:85-86). Most studies have implied that σ binding sites (receptors) are plasmalemmal elements of the signal transduction cascade. Drugs reported to be selective σ ligands have been evaluated as antipsychotics (Hanner, M. et al. (1996) Proc. Natl. Acad. Sci. 93:8072-8077).

The cDNA encoding the σ_1 receptor has recently been isolated from guinea pig liver. Northern blot analysis showed that highest densities of σ_1 mRNA were present in brain, liver,
15 kidney, stomach, jejunum, placenta, ovary, and adrenal gland. The tissue distribution was in agreement with its proposed participation in sterol synthesis. Comparisons with coding sequences from other organisms indicated that the guinea pig σ_1 binding site of the receptor was similar to a fungal sterol isomerase (Hanner et al. (supra)).

Physiological data have suggested that the σ_1 receptor plays an important role in hippocampal function (Monnet, F.P. et al. (1994) Br. J. Pharmacol. 112:709-715). Sigma ligands potentiate the NMDA-induced excitation of CA3 pyramidal cell neurons in rat dorsal hippocampus. The effects upon hippocampal neurons are differentially affected by pertussis toxin (Monnet et al. (supra)). The existence of σ receptors in the CNS, immune, and
20 endocrine systems have suggested a likelihood that it may serve as a link between the three systems. However, lack of a highly selective σ ligand has hampered efforts to understand the possible physiological roles of σ receptors (Su, T.-P. (1991) Eur. J. Biochem. 200:633-642).

The discovery of a new human sigma receptor and the polynucleotides encoding it satisfies a need in the art by providing new compositions which are useful in the diagnosis,
30 prevention and treatment of vesicle trafficking, immunological, and neoplastic disorders.

SUMMARY OF THE INVENTION

The invention features a substantially purified polypeptide, human sigma receptor (SIGR), having the amino acid sequence shown in SEQ ID NO:1, or fragments thereof.

The invention further provides an isolated and substantially purified polynucleotide
5 sequence encoding the polypeptide comprising the amino acid sequence of SEQ ID NO:1 or fragments thereof and a composition comprising said polynucleotide sequence. The invention also provides a polynucleotide sequence which hybridizes under stringent conditions to the polynucleotide sequence encoding the amino acid sequence SEQ ID NO:1, or fragments of said polynucleotide sequence. The invention further provides a polynucleotide sequence
10 comprising the complement of the polynucleotide sequence encoding the amino acid sequence of SEQ ID NO:1, or fragments or variants of said polynucleotide sequence.

The invention also provides an isolated and purified sequence comprising SEQ ID NO:2 or variants thereof. In addition, the invention provides a polynucleotide sequence which hybridizes under stringent conditions to the polynucleotide sequence of SEQ ID NO:2.
15 The invention also provides a polynucleotide sequence comprising the complement of SEQ ID NO:2, or fragments or variants thereof.

The present invention further provides an expression vector containing at least a fragment of any of the claimed polynucleotide sequences. In yet another aspect, the expression vector containing the polynucleotide sequence is contained within a host cell.

20 The invention also provides a method for producing a polypeptide comprising the amino acid sequence of SEQ ID NO:1 or a fragment thereof, the method comprising the steps of: a) culturing the host cell containing an expression vector containing at least a fragment of the polynucleotide sequence encoding SIGR under conditions suitable for the expression of the polypeptide; and b) recovering the polypeptide from the host cell culture.

25 The invention also provides a pharmaceutical composition comprising a substantially purified SIGR having the amino acid sequence of SEQ ID NO:1 in conjunction with a suitable pharmaceutical carrier.

The invention also provides a purified antagonist of the polypeptide of SEQ ID NO:1. In one aspect the invention provides a purified antibody which binds to a polypeptide
30 comprising the amino acid sequence of SEQ ID NO:1.

Still further, the invention provides a purified agonist of the polypeptide of SEQ ID NO:1.

The invention also provides a method for treating or preventing a vesicle trafficking disorder comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising purified SIGR.

5 The invention also provides a method for treating or preventing an immunological disorder comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising purified SIGR.

The invention also provides a method for treating or preventing a neoplastic disorder comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising purified SIGR.

10 The invention also provides a method for detecting a polynucleotide which encodes SIGR in a biological sample comprising the steps of: a) hybridizing the complement of the polynucleotide sequence which encodes SEQ ID NO:1 to nucleic acid material of a biological sample, thereby forming a hybridization complex; and b) detecting the hybridization complex, wherein the presence of the complex correlates with the presence of a
15 polynucleotide encoding SIGR in the biological sample. In one aspect the nucleic acid material of the biological sample is amplified by the polymerase chain reaction prior to hybridization.

BRIEF DESCRIPTION OF THE FIGURES

20 Figures 1A, 1B, 1C, and 1D show the amino acid sequence (SEQ ID NO:1) and nucleic acid sequence (SEQ ID NO:2) of SIGR. The alignment was produced using MacDNASIS PRO™ software (Hitachi Software Engineering Co. Ltd. San Bruno, CA).

Figures 2A and 2B show the amino acid sequence alignments between SIGR (1542751; SEQ ID NO:1), and guinea pig σ_1 receptor (GI 1403300; SEQ ID NO:3), produced
25 using the multisequence alignment program of DNASTAR™ software (DNASTAR Inc, Madison WI).

Figures 3A and 3B show the hydrophobicity plots for SIGR, SEQ ID NO: 1 and guinea pig σ_1 receptor (SEQ ID NO:3), respectively; the positive X axis reflects amino acid position, and the negative Y axis, hydrophobicity (MacDNASIS PRO software).

30

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is

understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors, and reagents described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the
5 appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a", "an", and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to "a host cell" includes a plurality of such host cells, reference to the "antibody" is a reference to one or more antibodies and equivalents thereof known to
10 those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods, devices,
15 and materials are now described. All publications mentioned herein are incorporated herein by reference for the purpose of describing and disclosing the cell lines, vectors, and methodologies which are reported in the publications which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

20

DEFINITIONS

SIGR, as used herein, refers to the amino acid sequences of substantially purified SIGR obtained from any species, particularly mammalian, including bovine, ovine, porcine, murine, equine, and preferably human, from any source whether natural, synthetic,
25 semi-synthetic, or recombinant.

The term "agonist", as used herein, refers to a molecule which, when bound to SIGR, increases or prolongs the duration of the effect of SIGR. Agonists may include proteins, nucleic acids, carbohydrates, or any other molecules which bind to and modulate the effect of SIGR.

30

An "allele" or "allelic sequence", as used herein, is an alternative form of the gene encoding SIGR. Alleles may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or polypeptides whose structure or function may or may not

be altered. Any given natural or recombinant gene may have none, one, or many allelic forms. Common mutational changes which give rise to alleles are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

5 "Altered" nucleic acid sequences encoding SIGR as used herein include those with deletions, insertions, or substitutions of different nucleotides resulting in a polynucleotide that encodes the same or a functionally equivalent SIGR. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding SIGR, and improper or unexpected hybridization to
10 alleles, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding SIGR. The encoded protein may also be "altered" and contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent SIGR. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the
15 amphipathic nature of the residues as long as the biological or immunological activity of SIGR is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid; positively charged amino acids may include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values may include leucine, isoleucine, and valine, glycine and alanine, asparagine and glutamine, serine and
20 threonine, and phenylalanine and tyrosine.

"Amino acid sequence" as used herein refers to an oligopeptide, peptide, polypeptide, or protein sequence, and fragment thereof, and to naturally occurring or synthetic molecules. Fragments of SIGR are preferably about 5 to about 15 amino acids in length and retain the biological activity or the immunological activity of SIGR. Where "amino acid sequence" is
25 recited herein to refer to an amino acid sequence of a naturally occurring protein molecule, amino acid sequence, and like terms, are not meant to limit the amino acid sequence to the complete, native amino acid sequence associated with the recited protein molecule.

"Amplification" as used herein refers to the production of additional copies of a nucleic acid sequence and is generally carried out using polymerase chain reaction (PCR) technologies well known in the art (Dieffenbach, C.W. and G.S. Dveksler (1995) PCR
30 Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainview, NY).

The term "antagonist" as used herein, refers to a molecule which, when bound to

SIGR, decreases the amount or the duration of the effect of the biological or immunological activity of SIGR. Antagonists may include proteins, nucleic acids, carbohydrates, antibodies or any other molecules which decrease the effect of SIGR.

As used herein, the term "antibody" refers to intact molecules as well as fragments thereof, such as Fa, F(ab')₂, and Fv, which are capable of binding the epitopic determinant. Antibodies that bind SIGR polypeptides can be prepared using intact polypeptides or fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal can be derived from the translation of RNA or synthesized chemically and can be conjugated to a carrier protein, if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin and thyroglobulin, keyhole limpet hemocyanin. The coupled peptide is then used to immunize the animal (e.g., a mouse, a rat, or a rabbit).

The term "antigenic determinant", as used herein, refers to that fragment of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to a given region or three-dimensional structure on the protein; these regions or structures are referred to as antigenic determinants. An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense", as used herein, refers to any composition containing nucleotide sequences which are complementary to a specific DNA or RNA sequence. The term "antisense strand" is used in reference to a nucleic acid strand that is complementary to the "sense" strand. Antisense molecules include peptide nucleic acids and may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form duplexes and block either transcription or translation. The designation "negative" is sometimes used in reference to the antisense strand, and "positive" is sometimes used in reference to the sense strand.

The term "biologically active", as used herein, refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic SIGR, or any oligopeptide thereof, to induce a specific immune response in appropriate

animals or cells and to bind with specific antibodies.

The terms "complementary" or "complementarity", as used herein, refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base-pairing. For example, the sequence "A-G-T" binds to the complementary sequence "T-C-A".

5 Complementary between two single-stranded molecules may be "partial", in which only some of the nucleic acids bind, or it may be complete when total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, which depend upon
10 binding between nucleic acids strands and in the design and use of PNA molecules.

A "composition comprising a given polynucleotide sequence" as used herein refers broadly to any composition containing the given polynucleotide sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding SIGR (SEQ ID NO:1) or fragments thereof (e.g., SEQ ID
15 NO:2 and fragments thereof) may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., SDS) and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

20 "Consensus", as used herein, refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, has been extended using XL-PCR™ (Perkin Elmer, Norwalk, CT) in the 5' and/or the 3' direction and resequenced, or has been assembled from the overlapping sequences of more than one Incyte Clone using a computer program for fragment assembly (e.g., GELVIEW™ Fragment Assembly system, GCG, Madison, WI).
25 Some sequences have been both extended and assembled to produce the consensus sequence .

The term "correlates with expression of a polynucleotide", as used herein, indicates that the detection of the presence of ribonucleic acid that is similar to SEQ ID NO:2 by northern analysis is indicative of the presence of mRNA encoding SIGR in a sample and thereby correlates with expression of the transcript from the polynucleotide encoding the
30 protein.

A "deletion", as used herein, refers to a change in the amino acid or nucleotide sequence and results in the absence of one or more amino acid residues or nucleotides.

The term "derivative", as used herein, refers to the chemical modification of a nucleic acid encoding or complementary to SIGR or the encoded SIGR. Such modifications include, for example, replacement of hydrogen by an alkyl, acyl, or amino group. A nucleic acid derivative encodes a polypeptide which retains the biological or immunological function of the natural molecule. A derivative polypeptide is one which is modified by glycosylation, pegylation, or any similar process which retains the biological or immunological function of the polypeptide from which it was derived.

The term "homology", as used herein, refers to a degree of complementarity. There may be partial homology or complete homology (i.e., identity). A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to using the functional term "substantially homologous." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization and the like) under conditions of low stringency. A substantially homologous sequence or hybridization probe will compete for and inhibit the binding of a completely homologous sequence to the target sequence under conditions of low stringency. This is not to say that conditions of low stringency are such that non-specific binding is permitted; low stringency conditions require that the binding of two sequences to one another be a specific (i.e., selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% identity). In the absence of non-specific binding, the probe will not hybridize to the second non-complementary target sequence.

Human artificial chromosomes (HACs) are linear microchromosomes which may contain DNA sequences of 10K to 10M in size and contain all of the elements required for stable mitotic chromosome segregation and maintenance (Harrington, J.J. et al. (1997) Nat Genet. 15:345-355).

The term "humanized antibody", as used herein, refers to antibody molecules in which amino acids have been replaced in the non-antigen binding regions in order to more closely resemble a human antibody, while still retaining the original binding ability.

The term "hybridization", as used herein, refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing.

The term "hybridization complex", as used herein, refers to a complex formed

between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary G and C bases and between complementary A and T bases; these hydrogen bonds may be further stabilized by base stacking interactions. The two complementary nucleic acid sequences hydrogen bond in an antiparallel configuration. A hybridization
5 complex may be formed in solution (e.g., C_0t or R_0t analysis) or between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

An "insertion" or "addition", as used herein, refers to a change in an amino acid or
10 nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively, as compared to the naturally occurring molecule.

"Microarray" refers to an array of distinct polynucleotides or oligonucleotides synthesized on a substrate, such as paper, nylon or other type of membrane, filter, chip, glass slide, or any other suitable solid support.

15 The term "modulate", as used herein, refers to a change in the activity of SGR. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional or immunological properties of SGR.

"Nucleic acid sequence" as used herein refers to an oligonucleotide, nucleotide, or polynucleotide, and fragments thereof, and to DNA or RNA of genomic or synthetic origin
20 which may be single- or double-stranded, and represent the sense or antisense strand.

"Fragments" are those nucleic acid sequences which are greater than 60 nucleotides in length, and most preferably includes fragments that are at least 100 nucleotides or at least 1000 nucleotides, and at least 10,000 nucleotides in length.

The term "oligonucleotide" refers to a nucleic acid sequence of at least about 6
25 nucleotides to about 60 nucleotides, preferably about 15 to 30 nucleotides, and more preferably about 20 to 25 nucleotides, which can be used in PCR amplification or a hybridization assay, or a microarray. As used herein, oligonucleotide is substantially equivalent to the terms "amplimers", "primers", "oligomers", and "probes", as commonly defined in the art.

30 "Peptide nucleic acid", PNA as used herein, refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least five nucleotides in length linked to a peptide backbone of amino acid residues which ends in lysine. The terminal

lysine confers solubility to the composition. PNAs may be pegylated to extend their lifespan in the cell where they preferentially bind complementary single stranded DNA and RNA and stop transcript elongation (Nielsen, P.E. et al. (1993) Anticancer Drug Des. 8:53-63).

5 The term "portion", as used herein, with regard to a protein (as in "a portion of a given protein") refers to fragments of that protein. The fragments may range in size from five amino acid residues to the entire amino acid sequence minus one amino acid. Thus, a protein "comprising at least a portion of the amino acid sequence of SEQ ID NO:1" encompasses the full-length SIGR and fragments thereof.

10 The term "sample", as used herein, is used in its broadest sense. A biological sample suspected of containing nucleic acid encoding SIGR, or fragments thereof, or SIGR itself may comprise a bodily fluid, extract from a cell, chromosome, organelle, or membrane isolated from a cell, a cell, genomic DNA, RNA, or cDNA(in solution or bound to a solid support, a tissue, a tissue print, and the like.

15 The terms "specific binding" or "specifically binding", as used herein, refers to that interaction between a protein or peptide and an agonist, an antibody and an antagonist. The interaction is dependent upon the presence of a particular structure (i.e., the antigenic determinant or epitope) of the protein recognized by the binding molecule. For example, if an antibody is specific for epitope "A", the presence of a protein containing epitope A (or free, unlabeled A) in a reaction containing labeled "A" and the antibody will reduce the amount of
20 labeled A bound to the antibody.

The terms "stringent conditions" or "stringency", as used herein, refer to the conditions for hybridization as defined by the nucleic acid, salt, and temperature. These conditions are well known in the art and may be altered in order to identify or detect identical or related polynucleotide sequences. Numerous equivalent conditions comprising either low or high
25 stringency depend on factors such as the length and nature of the sequence (DNA, RNA, base composition), nature of the target (DNA, RNA, base composition), milieu (in solution or immobilized on a solid substrate), concentration of salts and other components (e.g., formamide, dextran sulfate and/or polyethylene glycol), and temperature of the reactions (within a range from about 5°C below the melting temperature of the probe to about 20°C to
30 25°C below the melting temperature). One or more factors be may be varied to generate conditions of either low or high stringency different from, but equivalent to, the above listed conditions.

The term “substantially purified”, as used herein, refers to nucleic or amino acid sequences that are removed from their natural environment, isolated or separated, and are at least 60% free, preferably 75% free, and most preferably 90% free from other components with which they are naturally associated.

5 A “substitution”, as used herein, refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

“Transformation”, as defined herein, describes a process by which exogenous DNA enters and changes a recipient cell. It may occur under natural or artificial conditions using various methods well known in the art. Transformation may rely on any known method for
10 the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. Such “transformed” cells include stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as
15 part of the host chromosome. They also include cells which transiently express the inserted DNA or RNA for limited periods of time.

A “variant” of SIGR, as used herein, refers to an amino acid sequence that is altered by one or more amino acids. The variant may have “conservative” changes, wherein a substituted amino acid has similar structural or chemical properties, e.g., replacement of
20 leucine with isoleucine. More rarely, a variant may have “nonconservative” changes, e.g., replacement of a glycine with a tryptophan. Analogous minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for
25 example, DNASTAR software.

THE INVENTION

The invention is based on the discovery of a new human sigma receptor (hereinafter referred to as “SIGR”), the polynucleotides encoding SIGR, and the use of these compositions
30 for the diagnosis, prevention, or treatment of vesicle trafficking, immunological, and neoplastic disorders.

Nucleic acids encoding the SIGR of the present invention were first identified in

Incyte Clone 1542751 from the prostate cDNA library (PROSTUT04) using a computer search for amino acid sequence alignments. A consensus sequence, SEQ ID NO:2, was derived from the following overlapping and/or extended nucleic acid sequences: Incyte Clones 1542751 (PROSTUT04), 1929428 (COLNTUT03), 1611269 (COLNTUT06).

5 In one embodiment, the invention encompasses a polypeptide comprising the amino acid sequence of SEQ ID NO:1, as shown in Figs. 1A, 1B, 1C, and 1D. SIGR is 223 amino acids in length and has four potential casein kinase II phosphorylation sites at residues S-99, T-141, T-160, and S-192, three protein kinase C phosphorylation sites at residues S-117, T-140, and T-202, and two potential hydrophobic transmembrane segments between residues A-10 12 and S-34, and F-83 and L-111. As shown in Figure 2, SIGR has chemical and structural homology with guinea pig σ_1 receptor (GI 1403300; SEQ ID NO:3). In particular, SIGR and guinea pig σ_1 receptor share 93% identity, four potential casein kinase II phosphorylation sites, three protein kinase C phosphorylation sites at residues, and two potential hydrophobic transmembrane segments. As illustrated by Figs. 3A and 3B, SIGR and guinea pig σ_1 15 receptor have rather similar hydrophobicity plots. SIGR and guinea pig σ_1 receptor have similar isoelectric points, pI = 5.57 and 5.38, respectively. Northern analysis shows the expression of this sequence in various libraries, at least 62% of which are immortalized or cancerous, at least 69% are secretory, and at least 25% of which involve immune response. Of particular note is the expression of SIGR in prostate, gastrointestinal, brain, liver, ovary, 20 uterus, lung, and breast tissues.

The invention also encompasses SIGR variants. A preferred SIGR variant is one having at least 80%, and more preferably at least 90%, amino acid sequence identity to the SIGR amino acid sequence (SEQ ID NO:1) and which retains at least one biological, immunological or other functional characteristic or activity of SIGR. A most preferred SIGR 25 variant is one having at least 95% amino acid sequence identity to SEQ ID NO:1.

The invention also encompasses polynucleotides which encode SIGR. Accordingly, any nucleic acid sequence which encodes the amino acid sequence of SIGR can be used to produce recombinant molecules which express SIGR. In a particular embodiment, the invention encompasses the polynucleotide comprising the nucleic acid sequence of SEQ ID 30 NO:2 as shown in Figs. 1A, 1B, 1C, and 1D.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of nucleotide sequences encoding SIGR, some bearing minimal

homology to the nucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied
5 to the nucleotide sequence of naturally occurring SIGR, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode SIGR and its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring SIGR under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide
10 sequences encoding SIGR or its derivatives possessing a substantially different codon usage. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding SIGR and its derivatives without altering the encoded amino
15 acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences, or fragments thereof, which encode SIGR and its derivatives, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell
20 systems using reagents that are well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding SIGR or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed nucleotide sequences, and in particular, those shown in SEQ ID NO:2, under various conditions of stringency as taught in Wahl, G.M. and S.L. Berger (1987;
25 Methods Enzymol. 152:399-407) and Kimmel, A.R. (1987; Methods Enzymol. 152:507-511).

Methods for DNA sequencing which are well known and generally available in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, Sequenase® (US
30 Biochemical Corp, Cleveland, OH), Taq polymerase (Perkin Elmer), thermostable T7 polymerase (Amersham, Chicago, IL), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE Amplification System marketed by

Gibco/BRL (Gaithersburg, MD). Preferably, the process is automated with machines such as the Hamilton Micro Lab 2200 (Hamilton, Reno, NV), Peltier Thermal Cycler (PTC200; MJ Research, Watertown, MA) and the ABI Catalyst and 373 and 377 DNA Sequencers (Perkin Elmer).

5 The nucleic acid sequences encoding SIGR may be extended utilizing a partial nucleotide sequence and employing various methods known in the art to detect upstream sequences such as promoters and regulatory elements. For example, one method which may be employed, "restriction-site" PCR, uses universal primers to retrieve unknown sequence adjacent to a known locus (Sarkar, G. (1993) PCR Methods Applic. 2:318-322). In particular,
10 genomic DNA is first amplified in the presence of primer to a linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

15 Inverse PCR may also be used to amplify or extend sequences using divergent primers based on a known region (Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186). The primers may be designed using commercially available software such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, MN), or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target
20 sequence at temperatures about 68°-72° C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

 Another method which may be used is capture PCR which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial
25 chromosome DNA (Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119). In this method, multiple restriction enzyme digestions and ligations may also be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

 Another method which may be used to retrieve unknown sequences is that of Parker,
30 J.D. et al. (1991; Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PromoterFinder™ libraries to walk genomic DNA (Clontech, Palo Alto, CA). This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Also, random-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity may be converted to electrical signal using appropriate software (e.g. GenotyperTM and Sequence NavigatorTM, Perkin Elmer) and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode SIGH may be used in recombinant DNA molecules to direct expression of SIGH, fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced, and these sequences may be used to clone and express SIGH.

As will be understood by those of skill in the art, it may be advantageous to produce SIGH-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter SIGH encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR

reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

5 In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding SIGR may be ligated to a heterologous sequence to encode a fusion protein. For example, to screen peptide libraries for inhibitors of SIGR activity, it may be useful to encode a chimeric SIGR protein that can be recognized by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site located between
10 the SIGR encoding sequence and the heterologous protein sequence, so that SIGR may be cleaved and purified away from the heterologous moiety.

 In another embodiment, sequences encoding SIGR may be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers, M.H. et al. (1980) Nucl. Acids Res. Symp. Ser. 215-223, Horn, T. et al. (1980) Nucl. Acids Res. Symp. Ser. 225-232).
15 Alternatively, the protein itself may be produced using chemical methods to synthesize the amino acid sequence of SIGR, or a fragment thereof. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge, J.Y. et al. (1995) Science 269:202-204) and automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer).

20 The newly synthesized peptide may be substantially purified by preparative high performance liquid chromatography (e.g., Creighton, T. (1983) Proteins, Structures and Molecular Principles, WH Freeman and Co., New York, NY). The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; Creighton, supra). Additionally, the amino acid sequence of SIGR, or
25 any part thereof, may be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

 In order to express a biologically active SIGR, the nucleotide sequences encoding SIGR or functional equivalents, may be inserted into appropriate expression vector, i.e., a
30 vector which contains the necessary elements for the transcription and translation of the inserted coding sequence.

 Methods which are well known to those skilled in the art may be used to construct

expression vectors containing sequences encoding SIGR and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described in Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview, NY, and Ausubel, F.M. et al. (1989) Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY.

A variety of expression vector/host systems may be utilized to contain and express sequences encoding SIGR. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

The "control elements" or "regulatory sequences" are those non-translated regions of the vector--enhancers, promoters, 5' and 3' untranslated regions--which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the Bluescript® phagemid (Stratagene, LaJolla, CA) or pSport1™ plasmid (Gibco BRL) and the like may be used. The baculovirus polyhedrin promoter may be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (e.g., heat shock, RUBISCO; and storage protein genes) or from plant viruses (e.g., viral promoters or leader sequences) may be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding SIGR, vectors based on SV40 or EBV may be used with an appropriate selectable marker.

In bacterial systems, a number of expression vectors may be selected depending upon the use intended for SIGR. For example, when large quantities of SIGR are needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are

readily purified may be used. Such vectors include, but are not limited to, the multifunctional E. coli cloning and expression vectors such as Bluescript® (Stratagene), in which the sequence encoding SIGR may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors (Promega, Madison, WI) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast, Saccharomyces cerevisiae, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al. (1987) Methods Enzymol. 153:516-544.

In cases where plant expression vectors are used, the expression of sequences encoding SIGR may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, for example, Hobbs, S. or Murry, L.E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, NY; pp. 191-196.

An insect system may also be used to express SIGR. For example, in one such system, Autographa californica nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in Spodoptera frugiperda cells or in Trichoplusia larvae. The sequences encoding SIGR may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of SIGR will

render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, S. frugiperda cells or Trichoplusia larvae in which SIGR may be expressed (Engelhard, E.K. et al. (1994) Proc. Nat. Acad. Sci. 91:3224-3227).

5 In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding SIGR may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing SIGR in infected
10 host cells (Logan, J. and Shenk, T. (1984) Proc. Natl. Acad. Sci. 81:3655-3659). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

 Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6 to 10M are
15 constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes.

 Specific initiation signals may also be used to achieve more efficient translation of sequences encoding SIGR. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding SIGR, its initiation codon, and upstream
20 sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational
25 elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf, D. et al. (1994) Results Probl. Cell Differ. 20:125-162).

 In addition, a host cell strain may be chosen for its ability to modulate the expression
30 of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which

cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC; Bethesda, MD) and may be
5 chosen to ensure the correct modification and processing of the foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express SGR may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector.
10 Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell
15 type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler, M. et al. (1977) *Cell* 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. et al. (1980) *Cell* 22:817-23) genes which can be employed in *tk⁻* or *aprt⁻* cells, respectively. Also,
20 antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, *dhfr* which confers resistance to methotrexate (Wigler, M. et al. (1980) *Proc. Natl. Acad. Sci.* 77:3567-70); *npt*, which confers resistance to the aminoglycosides neomycin and G-418 (Colbere-Garapin, F. et al (1981) *J. Mol. Biol.* 150:1-14) and *als* or *pat*, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, supra).
25 Additional selectable genes have been described, for example, *trpB*, which allows cells to utilize indole in place of tryptophan, or *hisD*, which allows cells to utilize histinol in place of histidine (Hartman, S.C. and R.C. Mulligan (1988) *Proc. Natl. Acad. Sci.* 85:8047-51). Recently, the use of visible markers has gained popularity with such markers as anthocyanins, β glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely
30 used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes, C.A. et al. (1995) *Methods Mol. Biol.* 55:121-131).

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be confirmed. For example, if the sequence encoding SGR is inserted within a marker gene sequence, transformed cells containing sequences encoding SGR can be identified by the absence of marker gene
5 function. Alternatively, a marker gene can be placed in tandem with a sequence encoding SGR under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

Alternatively, host cells which contain the nucleic acid sequence encoding SGR and express SGR may be identified by a variety of procedures known to those of skill in the art.
10 These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein.

The presence of polynucleotide sequences encoding SGR can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or
15 fragments of polynucleotides encoding SGR. Nucleic acid amplification based assays involve the use of oligonucleotides or oligomers based on the sequences encoding SGR to detect transformants containing DNA or RNA encoding SGR.

A variety of protocols for detecting and measuring the expression of SGR, using either polyclonal or monoclonal antibodies specific for the protein are known in the art.
20 Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on SGR is preferred, but a competitive binding assay may be employed. These and other assays are described, among other places, in Hampton, R. et al. (1990; Serological Methods, a Laboratory Manual,
25 APS Press, St Paul, MN) and Maddox, D.E. et al. (1983; J. Exp. Med. 158:1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding SGR include oligolabeling, nick translation, end-labeling or PCR amplification
30 using a labeled nucleotide. Alternatively, the sequences encoding SGR, or any fragments thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in

vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits (Pharmacia & Upjohn, (Kalamazoo, MI); Promega (Madison WI); and U.S. Biochemical Corp., Cleveland, OH). Suitable reporter molecules or labels, which may be used for ease of
5 detection, include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding SIGR may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or contained intracellularly
10 depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode SIGR may be designed to contain signal sequences which direct secretion of SIGR through a prokaryotic or eukaryotic cell membrane. Other constructions may be used to join sequences encoding SIGR to nucleotide sequence encoding a polypeptide domain which will facilitate purification
15 of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, WA). The inclusion of cleavable linker sequences such as
20 those specific for Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and SIGR may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing SIGR and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMAC (immobilized metal ion affinity
25 chromatography as described in Porath, J. et al. (1992, Prot. Exp. Purif. 3: 263-281) while the enterokinase cleavage site provides a means for purifying SIGR from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D.J. et al. (1993; DNA Cell Biol. 12:441-453).

In addition to recombinant production, fragments of SIGR may be produced by direct
30 peptide synthesis using solid-phase techniques Merrifield J. (1963) J. Am. Chem. Soc. 85:2149-2154). Protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems

431A Peptide Synthesizer (Perkin Elmer). Various fragments of SIGR may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

5 THERAPEUTICS

Chemical and structural homology exists between SIGR and σ_1 receptor from guinea pig (GI 1403300). In addition, SIGR is expressed in prostate, gastrointestinal, brain, liver, ovary, uterus, lung, and breast tissues. Therefore, SIGR appears to play a role in vesicle trafficking, immunological, and neoplastic disorders.

10 Therefore, in one embodiment, SIGR or a fragment or derivative thereof may be administered to a subject to treat or prevent a vesicle trafficking disorder. Such disorders include, but are not limited to, cystic fibrosis, glucose-galactose malabsorption syndrome, hypercholesterolemia, diabetes mellitus, diabetes insipidus, hyper- and hypoglycemia, Grave's disease, goiter, Cushing's disease, and Addison's disease; gastrointestinal disorders
15 including ulcerative colitis, gastric and duodenal ulcers; other conditions associated with abnormal vesicle trafficking including AIDS; allergies including hay fever, asthma, and urticaria (hives); autoimmune hemolytic anemia; proliferative glomerulonephritis; inflammatory bowel disease; multiple sclerosis; myasthenia gravis; rheumatoid and osteoarthritis; scleroderma; Chediak-Higashi and Sjogren's syndromes; systemic lupus
20 erythematosus; toxic shock syndrome; traumatic tissue damage; and viral, bacterial, fungal, helminth, and protozoal infections.

In another embodiment, a vector capable of expressing SIGR, or a fragment or a derivative thereof, may also be administered to a subject to treat or prevent a vesicle trafficking disorder including, but not limited to, those described above.

25 In still another embodiment, an agonist of SIGR may also be administered to a subject to treat or prevent a vesicle trafficking disorder including, but not limited to, those described above.

In one embodiment, SIGR or a fragment or derivative thereof may be administered to a subject to treat or prevent an immunological disorder. Such disorders include, but are not
30 limited to, AIDS, Addison's disease, adult respiratory distress syndrome, allergies, anemia, asthma, atherosclerosis, bronchitis, cholecystitis, Crohn's disease, ulcerative colitis, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythema nodosum, atrophic

gastritis, glomerulonephritis, gout, Graves' disease, hypereosinophilia, irritable bowel syndrome, lupus erythematosus, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, rheumatoid arthritis, scleroderma, Sjögren's syndrome, Werner syndrome, and autoimmune thyroiditis; 5 complications of cancer, hemodialysis, and extracorporeal circulation; viral, bacterial, fungal, parasitic, protozoal, and helminthic infections; and trauma.

In another embodiment, a vector capable of expressing SIGR, or a fragment or a derivative thereof, may also be administered to a subject to treat or prevent an immunological disorder including, but not limited to, those described above.

10 In still another embodiment, an agonist of SIGR may also be administered to a subject to treat or prevent an immunological disorder including, but not limited to, those described above.

In one embodiment, SIGR or a fragment or derivative thereof may be administered to a subject to treat or prevent a neoplastic disorder. Such disorders include, but are not limited 15 to, adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus.

20 In another embodiment, a vector capable of expressing SIGR, or a fragment or a derivative thereof, may also be administered to a subject to treat or prevent a neoplastic disorder including, but not limited to, those described above.

In still another embodiment, an agonist of SIGR may also be administered to a subject to treat or prevent a neoplastic disorder including, but not limited to, those described above.

25 In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act 30 synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of SIGR may be produced using methods which are generally known in the art. In particular, purified SIGR may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind SIGR.

Antibodies to SIGR may be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies, (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others, may be immunized by injection with SIGR or any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to SIGR have an amino acid sequence consisting of at least five amino acids and more preferably at least 10 amino acids. It is also preferable that they are identical to a portion of the amino acid sequence of the natural protein, and they may contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of SIGR amino acids may be fused with those of another protein such as keyhole limpet hemocyanin and antibody produced against the chimeric molecule.

Monoclonal antibodies to SIGR may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique (Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. 80:2026-2030; Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120).

In addition, techniques developed for the production of "chimeric antibodies", the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity can be used (Morrison, S.L. et al.

(1984) Proc. Natl. Acad. Sci. 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; Takeda, S. et al. (1985) Nature 314:452-454). Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce SIGR-specific single chain antibodies. Antibodies with related
5 specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries (Burton D.R. (1991) Proc. Natl. Acad. Sci. 88:11120-3).

Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding
10 reagents as disclosed in the literature (Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. 86: 3833-3837; Winter, G. et al. (1991) Nature 349:293-299).

Antibody fragments which contain specific binding sites for SIGR may also be generated. For example, such fragments include, but are not limited to, the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments
15 which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity (Huse, W.D. et al. (1989) Science 254:1275-1281).

Various immunoassays may be used for screening to identify antibodies having the
20 desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between SIGR and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering SIGR epitopes
25 is preferred, but a competitive binding assay may also be employed (Maddox, *supra*).

In another embodiment of the invention, the polynucleotides encoding SIGR, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding SIGR may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be
30 transformed with sequences complementary to polynucleotides encoding SIGR. Thus, complementary molecules or fragments may be used to modulate SIGR activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or

antisense oligonucleotides or larger fragments, can be designed from various locations along the coding or control regions of sequences encoding SIGR.

Expression vectors derived from retro viruses, adenovirus, herpes or vaccinia viruses, or from various bacterial plasmids may be used for delivery of nucleotide sequences to the
5 targeted organ, tissue or cell population. Methods which are well known to those skilled in the art can be used to construct vectors which will express nucleic acid sequence which is complementary to the polynucleotides of the gene encoding SIGR. These techniques are described both in Sambrook et al. (supra) and in Ausubel et al. (supra).

Genes encoding SIGR can be turned off by transforming a cell or tissue with
10 expression vectors which express high levels of a polynucleotide or fragment thereof which encodes SIGR. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector and even
15 longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5' or regulatory regions of the gene encoding SIGR (signal sequence, promoters, enhancers, and introns). Oligonucleotides derived from the transcription initiation site, e.g., between
20 positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature (Gee, J.E. et al. (1994) In: Huber,
25 B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing Co., Mt. Kisco, NY). The complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific
30 hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples which may be used include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic

cleavage of sequences encoding SGR.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of
5 between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

10 Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding SGR. Such DNA sequences may
15 be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA constitutively or inducibly can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at
20 the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which
25 are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections or polycationic amino
30 polymers (Goldman, C.K. et al. (1997) Nature Biotechnology 15:462-66; incorporated herein by reference) may be achieved using methods which are well known in the art.

Any of the therapeutic methods described above may be applied to any subject in need

of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

An additional embodiment of the invention relates to the administration of a pharmaceutical composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of SIGR, antibodies to SIGR, mimetics, agonists, antagonists, or inhibitors of SIGR. The compositions may be administered alone or in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium

carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

5 Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

10 Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable
15 liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium
20 carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents
25 which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a
30 manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of SIGR, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models, usually mice, rabbits, dogs, or pigs. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example SIGR or fragments thereof, antibodies of SIGR, agonists, antagonists or inhibitors of SIGR, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio between therapeutic and toxic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to

WO 99/13055

PCT/US98/18941

the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug
5 combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from 0.1 to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages
10 and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

15 **DIAGNOSTICS**

In another embodiment, antibodies which specifically bind SIGR may be used for the diagnosis of conditions or diseases characterized by expression of SIGR, or in assays to monitor patients being treated with SIGR, agonists, antagonists or inhibitors. The antibodies useful for diagnostic purposes may be prepared in the same manner as those described above
20 for therapeutics. Diagnostic assays for SIGR include methods which utilize the antibody and a label to detect SIGR in human body fluids or extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by joining them, either covalently or non-covalently, with a reporter molecule. A wide variety of reporter molecules which are known in the art may be used, several of which are described above.

A variety of protocols including ELISA, RIA, and FACS for measuring SIGR are known in the art and provide a basis for diagnosing altered or abnormal levels of SIGR expression. Normal or standard values for SIGR expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to SIGR under conditions suitable for complex formation. The amount of standard
30 complex formation may be quantified by various methods, but preferably by photometric, means. Quantities of SIGR expressed in subject, control and disease, samples from biopsied tissues are compared with the standard values. Deviation between standard and subject

values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding SIGR may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of SIGR may be correlated with disease. The diagnostic assay may be used to distinguish between absence, presence, and excess expression of SIGR, and to monitor regulation of SIGR levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding SIGR or closely related molecules, may be used to identify nucleic acid sequences which encode SIGR. The specificity of the probe, whether it is made from a highly specific region, e.g., 10 unique nucleotides in the 5' regulatory region, or a less specific region, e.g., especially in the 3' coding region, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low) will determine whether the probe identifies only naturally occurring sequences encoding SIGR, alleles, or related sequences.

Probes may also be used for the detection of related sequences, and should preferably contain at least 50% of the nucleotides from any of the SIGR encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and derived from the nucleotide sequence of SEQ ID NO:2 or from genomic sequence including promoter, enhancer elements, and introns of the naturally occurring SIGR.

Means for producing specific hybridization probes for DNAs encoding SIGR include the cloning of nucleic acid sequences encoding SIGR or SIGR derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, radionuclides such as ³²P or ³⁵S, or enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding SIGR may be used for the diagnosis of conditions or disorders which are associated with expression of SIGR. Examples of such conditions or disorders include a vesicle trafficking disorder such as cystic fibrosis, glucose-galactose

malabsorption syndrome, hypercholesterolemia, diabetes mellitus, diabetes insipidus, hyper- and hypoglycemia, Grave's disease, goiter, Cushing's disease, and Addison's disease; gastrointestinal disorders including ulcerative colitis, gastric and duodenal ulcers; other conditions associated with abnormal vesicle trafficking including AIDS; allergies including
5 hay fever, asthma, and urticaria (hives); autoimmune hemolytic anemia; proliferative glomerulonephritis; inflammatory bowel disease; multiple sclerosis; myasthenia gravis; rheumatoid and osteoarthritis; scleroderma; Chediak-Higashi and Sjogren's syndromes; systemic lupus erythematosus; toxic shock syndrome; traumatic tissue damage; and viral, bacterial, fungal, helminth, and protozoal infections; an immunological disorder such as adult
10 respiratory distress syndrome, anemia, atherosclerosis, bronchitis, cholecystitis, Crohn's disease, atopic dermatitis, dermatomyositis, emphysema, erythema nodosum, atrophic gastritis, glomerulonephritis, gout, hypereosinophilia, irritable bowel syndrome, myocardial or pericardial inflammation, osteoporosis, pancreatitis, polymyositis, Werner syndrome, and autoimmune thyroiditis; complications of cancer, hemodialysis, and extracorporeal
15 circulation; or a neoplastic disorder such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. The polynucleotide sequences
20 encoding SIGR may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; or in dipstick, pin, ELISA assays or microarrays utilizing fluids or tissues from patient biopsies to detect altered SIGR expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding SIGR may be useful in
25 assays that detect activation or induction of various cancers, particularly those mentioned above. The nucleotide sequences encoding SIGR may be labeled by standard methods, and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the
30 biopsied or extracted sample is significantly altered from that of a comparable control sample, the nucleotide sequences have hybridized with nucleotide sequences in the sample, and the presence of altered levels of nucleotide sequences encoding SIGR in the sample indicates the

WO 99/13055

PCT/US98/18941

presence of the associated disease. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or in monitoring the treatment of an individual patient.

In order to provide a basis for the diagnosis of disease associated with expression of SIGR, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, which encodes SIGR, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with those from an experiment where a known amount of a substantially purified polynucleotide is used. Standard values obtained from normal samples may be compared with values obtained from samples from patients who are symptomatic for disease. Deviation between standard and subject values is used to establish the presence of disease.

Once disease is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to evaluate whether the level of expression in the patient begins to approximate that which is observed in the normal patient. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding SIGR may involve the use of PCR. Such oligomers may be chemically synthesized, generated enzymatically, or produced *in vitro*. Oligomers will preferably consist of two nucleotide sequences, one with sense orientation (5'→3') and another with antisense (3'←5'), employed under optimized conditions for identification of a specific gene or condition. The same two oligomers, nested sets of oligomers, or even a degenerate pool of oligomers may be employed under less stringent conditions for detection and/or quantitation of closely related DNA or RNA sequences.

Methods which may also be used to quantitate the expression of SGR include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and standard curves onto which the experimental results are interpolated (Melby, P.C. et al. (1993) J. Immunol. Methods, 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 229-236). The speed of quantitation of multiple samples may be accelerated by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, an oligonucleotide derived from any of the polynucleotide sequences described herein may be used as a target in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously (to produce a transcript image), and to identify genetic variants, mutations and polymorphisms. This information will be useful in determining gene function, understanding the genetic basis of disease, diagnosing disease, and in developing and monitoring the activity of therapeutic agents (Heller, R. et al. (1997) Proc. Natl. Acad. Sci. 94:2150-55) .

In one embodiment, the microarray is prepared and used according to the methods described in PCT application WO95/11995 (Chee et al.), Lockhart, D. J. et al. (1996; Nat. Biotech. 14: 1675-1680) and Schena, M. et al. (1996; Proc. Natl. Acad. Sci. 93: 10614-10619), all of which are incorporated herein in their entirety by reference.

The microarray is preferably composed of a large number of unique, single-stranded nucleic acid sequences, usually either synthetic antisense oligonucleotides or fragments of cDNAs, fixed to a solid support. The oligonucleotides are preferably about 6-60 nucleotides in length, more preferably 15-30 nucleotides in length, and most preferably about 20-25 nucleotides in length. For a certain type of microarray, it may be preferable to use oligonucleotides which are only 7-10 nucleotides in length. The microarray may contain oligonucleotides which cover the known 5', or 3', sequence, sequential oligonucleotides which cover the full length sequence; or unique oligonucleotides selected from particular areas along the length of the sequence. Polynucleotides used in the microarray may be oligonucleotides that are specific to a gene or genes of interest in which at least a fragment of the sequence is known or that are specific to one or more unidentified cDNAs which are common to a particular cell type, developmental or disease state.

In order to produce oligonucleotides to a known sequence for a microarray, the gene of interest is examined using a computer algorithm which starts at the 5' or more preferably at

the 3' end of the nucleotide sequence. The algorithm identifies oligomers of defined length that are unique to the gene, have a GC content within a range suitable for hybridization, and lack predicted secondary structure that may interfere with hybridization. In certain situations it may be appropriate to use pairs of oligonucleotides on a microarray. The "pairs" will be
5 identical, except for one nucleotide which preferably is located in the center of the sequence. The second oligonucleotide in the pair (mismatched by one) serves as a control. The number of oligonucleotide pairs may range from two to one million. The oligomers are synthesized at designated areas on a substrate using a light-directed chemical process. The substrate may be paper, nylon or other type of membrane, filter, chip, glass slide or any other suitable solid
10 support.

In another aspect, an oligonucleotide may be synthesized on the surface of the substrate by using a chemical coupling procedure and an ink jet application apparatus, as described in PCT application WO95/251116 (Baldeschweiler et al.) which is incorporated herein in its entirety by reference. In another aspect, a "gridded" array analogous to a dot (or
15 slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such as those described above, may be produced by hand or by using available devices (slot blot or dot blot apparatus), materials (any suitable solid support), and machines (including robotic instruments), and may contain 8, 24, 96, 384, 1536 or 6144
20 oligonucleotides, or any other number between two and one million which lends itself to the efficient use of commercially available instrumentation.

In order to conduct sample analysis using a microarray, the RNA or DNA from a biological sample is made into hybridization probes. The mRNA is isolated, and cDNA is produced and used as a template to make antisense RNA (aRNA). The aRNA is amplified in
25 the presence of fluorescent nucleotides, and labeled probes are incubated with the microarray so that the probe sequences hybridize to complementary oligonucleotides of the microarray. Incubation conditions are adjusted so that hybridization occurs with precise complementary matches or with various degrees of less complementarity. After removal of nonhybridized probes, a scanner is used to determine the levels and patterns of fluorescence. The scanned
30 images are examined to determine degree of complementarity and the relative abundance of each oligonucleotide sequence on the microarray. The biological samples may be obtained from any bodily fluids (such as blood, urine, saliva, phlegm, gastric juices, etc.), cultured

cells, biopsies, or other tissue preparations. A detection system may be used to measure the absence, presence, and amount of hybridization for all of the distinct sequences simultaneously. This data may be used for large scale correlation studies on the sequences, mutations, variants, or polymorphisms among samples.

5 In another embodiment of the invention, the nucleic acid sequences which encode SIGH may also be used to generate hybridization probes which are useful for mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome or to artificial chromosome constructions, such as human artificial chromosomes (HACs), yeast artificial chromosomes (YACs),
10 bacterial artificial chromosomes (BACs), bacterial P1 constructions or single chromosome cDNA libraries as reviewed in Price, C.M. (1993) Blood Rev. 7:127-134, and Trask, B.J. (1991) Trends Genet. 7:149-154.

Fluorescent in situ hybridization (FISH as described in Verma et al. (1988) Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York, NY) may be
15 correlated with other physical chromosome mapping techniques and genetic map data. Examples of genetic map data can be found in various scientific journals or at Online Mendelian Inheritance in Man (OMIM). Correlation between the location of the gene encoding SIGH on a physical chromosomal map and a specific disease, or predisposition to a specific disease, may help delimit the region of DNA associated with that genetic disease.
20 The nucleotide sequences of the subject invention may be used to detect differences in gene sequences between normal, carrier, or affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques such as linkage analysis using established chromosomal markers may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian
25 species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms, or parts thereof, by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic
30 linkage to a particular genomic region, for example, AT to 11q22-23 (Gatti, R.A. et al. (1988) Nature 336:577-580), any sequences mapping to that area may represent associated or regulatory genes for further investigation. The nucleotide sequence of the subject invention

may also be used to detect differences in the chromosomal location due to translocation, inversion, etc. among normal, carrier, or affected individuals.

In another embodiment of the invention, SIGR, its catalytic or immunogenic fragments or oligopeptides thereof, can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes, between SIGR and the agent being tested, may be measured.

Another technique for drug screening which may be used provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO84/03564. In this method, as applied to SIGR large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with SIGR, or fragments thereof, and washed. Bound SIGR is then detected by methods well known in the art. Purified SIGR can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding SIGR specifically compete with a test compound for binding SIGR. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with SIGR.

In additional embodiments, the nucleotide sequences which encode SIGR may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

The examples below are provided to illustrate the subject invention and are not included for the purpose of limiting the invention.

EXAMPLES

I PROSTUT04 cDNA Library Construction

The PROSTUT04 cDNA library was constructed from prostate tumor tissue of a 57-year-old Caucasian male. The pathology report indicated an adenocarcinoma (Gleason grade 3+3) in both the left and right periphery of the prostate. The patient was taking insulin

for type I diabetes. The patient's family included a malignant neoplasm of the prostate in the patient's father and type I diabetes without complications in the mother.

The frozen tissue was immediately homogenized and lysed using a Brinkmann Homogenizer Polytron-PT 3000 (Brinkmann Instruments, Inc. Westbury, NY) in guanidinium isothiocyanate solution. 1.0 ml of 2 M sodium acetate was added to the lysate and the lysate was extracted once with phenol chloroform at pH 5.5 per Stratagene's RNA isolation protocol (Stratagene), and then with once with acid phenol at pH 4.7. The RNA was precipitated twice with an equal volume of isopropanol per Stratagene's protocol. The RNA pellet was resuspended in DEPC-treated water and DNase treated for 50 min at 37°C. The reaction was stopped with an equal volume of acid phenol. The RNA was precipitated using 0.3 M sodium acetate and 2.5 volume of ethanol, resuspended in DEPC-treated water. The RNA was isolated using the Qiagen Oligotex kit (QIAGEN Inc, Chatsworth, CA) and used to construct the cDNA library.

The RNA was handled according to the recommended protocols in the SuperScript Plasmid System (Cat. #18248-013; GIBCO-BRL). cDNA synthesis was initiated using a NotI-oligo d(T) primer. Double-stranded cDNA was blunted, ligated to SalI adaptors, digested with NotI, fractionated on a Sepharose CL4B column (Cat. #275105, Pharmacia), and those cDNAs exceeding 400 bp were ligated into the NotI and SalI sites of pSport I. The plasmid pSport I was subsequently transformed into DH5 α TM competent cells (Cat. #18258-012, GIBCO-BRL).

II Isolation and Sequencing of cDNA Clones

Plasmid DNA was released from the cells and purified using the REAL Prep 96 plasmid kit (Catalog #26173, QIAGEN, Inc.). The recommended protocol was employed except for the following changes: 1) the bacteria were cultured in 1 ml of sterile Terrific Broth (Catalog #22711, GIBCO-BRL) with carbenicillin at 25 mg/L and glycerol at 0.4%; 2) after inoculation, the cultures were incubated for 19 hours and at the end of incubation, the cells were lysed with 0.3 ml of lysis buffer; and 3) following isopropanol precipitation, the plasmid DNA pellet was resuspended in 0.1 ml of distilled water. After the last step in the protocol, samples were transferred to a 96-well block for storage at 4° C.

The cDNAs were sequenced by the method of Sanger et al. (1975, J. Mol. Biol. 94:441f), using a Hamilton Micro Lab 2200 (Hamilton, Reno, NV) in combination with

Peltier Thermal Cyclers (PTC200 from MJ Research, Watertown, MA) and Applied Biosystems 377 DNA Sequencing Systems; and the reading frame was determined.

III Homology Searching of cDNA Clones and Their Deduced Proteins

5 The nucleotide sequences and/or amino acid sequences of the Sequence Listing were used to query sequences in the GenBank, SwissProt, BLOCKS, and Pima II databases. These databases, which contain previously identified and annotated sequences, were searched for regions of homology using BLAST, which stands for Basic Local Alignment Search Tool (Altschul, S.F. (1993) J. Mol. Evol 36:290-300; Altschul, et al. (1990) J. Mol. Biol. 215:403-
10 410).

BLAST produced alignments of both nucleotide and amino acid sequences to determine sequence similarity. Because of the local nature of the alignments, BLAST was especially useful in determining exact matches or in identifying homologs which may be of prokaryotic (bacterial) or eukaryotic (animal, fungal, or plant) origin. Other algorithms such
15 as the one described in Smith, T. et al. (1992, Protein Engineering 5:35-51), incorporated herein by reference, could have been used when dealing with primary sequence patterns and secondary structure gap penalties. The sequences disclosed in this application have lengths of at least 49 nucleotides, and no more than 12% uncalled bases (where N is recorded rather than A, C, G, or T).

20 The BLAST approach searched for matches between a query sequence and a database sequence. BLAST evaluated the statistical significance of any matches found, and reported only those matches that satisfy the user-selected threshold of significance. In this application, threshold was set at 10^{-25} for nucleotides and 10^{-14} for peptides.

Incyte nucleotide sequences were searched against the GenBank databases for primate
25 (pri), rodent (rod), and other mammalian sequences (mam); and deduced amino acid sequences from the same clones were then searched against GenBank functional protein databases, mammalian (mamp), vertebrate (vrtp), and eukaryote (eukp) for homology.

IV Northern Analysis

30 Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound (Sambrook et al., supra).

Analogous computer techniques using BLAST (Altschul, S.F. (1993) J.Mol.Evol. 36:290-300; Altschul, S.F. et al. (1990) J.Mol.Evol. 215:403-410) are used to search for identical or related molecules in nucleotide databases such as GenBank or the LIFESEQ™ database (Incyte Pharmaceuticals). This analysis is much faster than multiple, membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or homologous.

The basis of the search is the product score which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1-2% error; and at 70, the match will be exact. Homologous molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analysis are reported as a list of libraries in which the transcript encoding SIGR occurs. Abundance and percent abundance are also reported. Abundance directly reflects the number of times a particular transcript is represented in a cDNA library, and percent abundance is abundance divided by the total number of sequences examined in the cDNA library.

20

V Extension of SIGR Encoding Polynucleotides

The nucleic acid sequence of the Incyte Clone 1542751 was used to design oligonucleotide primers for extending a partial nucleotide sequence to full length. One primer was synthesized to initiate extension in the antisense direction, and the other was synthesized to extend sequence in the sense direction. Primers were used to facilitate the extension of the known sequence "outward" generating amplicons containing new, unknown nucleotide sequence for the region of interest. The initial primers were designed from the cDNA using OLIGO 4.06 (National Biosciences), or another appropriate program, to be about 22 to about 30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures of about 68° to about 72° C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

30

Selected human cDNA libraries (Gibco/BRL) were used to extend the sequence If

more than one extension is necessary or desired, additional sets of primers are designed to further extend the known region.

High fidelity amplification was obtained by following the instructions for the XL-PCR kit (Perkin Elmer) and thoroughly mixing the enzyme and reaction mix. Beginning with 40 pmol of each primer and the recommended concentrations of all other components of the kit, PCR was performed using the Peltier Thermal Cycler (PTC200; M.J. Research, Watertown, MA) and the following parameters:

	Step 1	94° C for 1 min (initial denaturation)
	Step 2	65° C for 1 min
10	Step 3	68° C for 6 min
	Step 4	94° C for 15 sec
	Step 5	65° C for 1 min
	Step 6	68° C for 7 min
	Step 7	Repeat step 4-6 for 15 additional cycles
15	Step 8	94° C for 15 sec
	Step 9	65° C for 1 min
	Step 10	68° C for 7:15 min
	Step 11	Repeat step 8-10 for 12 cycles
	Step 12	72° C for 8 min
20	Step 13	4° C (and holding)

A 5-10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a low concentration (about 0.6-0.8%) agarose mini-gel to determine which reactions were successful in extending the sequence. Bands thought to contain the largest products were excised from the gel, purified using QIAQuick™ (QIAGEN Inc., Chatsworth, CA), and trimmed of overhangs using Klenow enzyme to facilitate religation and cloning.

After ethanol precipitation, the products were redissolved in 13 μ l of ligation buffer, 1 μ l T4-DNA ligase (15 units) and 1 μ l T4 polynucleotide kinase were added, and the mixture was incubated at room temperature for 2-3 hours or overnight at 16° C. Competent *E. coli* cells (in 40 μ l of appropriate media) were transformed with 3 μ l of ligation mixture and cultured in 80 μ l of SOC medium (Sambrook et al., supra). After incubation for one hour at 37° C, the *E. coli* mixture was plated on Luria Bertani (LB)-agar (Sambrook et al., supra) containing 2x Carb. The following day, several colonies were randomly picked from each plate and cultured in 150 μ l of liquid LB/2x Carb medium placed in an individual well of an appropriate, commercially-available, sterile 96-well microtiter plate. The following day, 5 μ l of each overnight culture was transferred into a non-sterile 96-well plate and after dilution 1:10 with water, 5 μ l of each sample was transferred into a PCR array.

For PCR amplification, 18 μ l of concentrated PCR reaction mix (3.3x) containing 4 units of rTth DNA polymerase, a vector primer, and one or both of the gene specific primers used for the extension reaction were added to each well. Amplification was performed using the following conditions:

5	Step 1	94° C for 60 sec
	Step 2	94° C for 20 sec
	Step 3	55° C for 30 sec
	Step 4	72° C for 90 sec
	Step 5	Repeat steps 2-4 for an additional 29 cycles
10	Step 6	72° C for 180 sec
	Step 7	4° C (and holding)

Aliquots of the PCR reactions were run on agarose gels together with molecular weight markers. The sizes of the PCR products were compared to the original partial cDNAs, and appropriate clones were selected, ligated into plasmid, and sequenced.

In like manner, the nucleotide sequence of SEQ ID NO:2 is used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for 5' extension, and an appropriate genomic library.

20 VI Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:2 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base-pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 (National Biosciences), labeled by combining 50 pmol of each oligomer and 250 μ Ci of [γ -³²P] adenosine triphosphate (Amersham) and T4 polynucleotide kinase (DuPont NEN®, Boston, MA). The labeled oligonucleotides are substantially purified with Sephadex G-25 superfine resin column (Pharmacia & Upjohn). A aliquot containing 10⁷ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases (Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II; DuPont NEN®).

The DNA from each digest is fractionated on a 0.7 percent agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham, NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1 x

saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMAT AR™ film (Kodak, Rochester, NY) is exposed to the blots in a Phosphoimager cassette (Molecular Dynamics, Sunnyvale, CA) for several hours, hybridization patterns are compared visually.

5 VII Microarrays

To produce oligonucleotides for a microarray, the nucleotide sequence described herein is examined using a computer algorithm which starts at the 3' end of the nucleotide sequence. The algorithm identifies oligomers of defined length that are unique to the gene, have a GC content within a range suitable for hybridization, and lack predicted secondary
10 structure that would interfere with hybridization. The algorithm identifies 20 sequence-specific oligonucleotides of 20 nucleotides in length (20-mers). A matched set of oligonucleotides is created in which one nucleotide in the center of each sequence is altered. This process is repeated for each gene in the microarray, and double sets of twenty 20 mers are synthesized and arranged on the surface of the silicon chip using a light-directed chemical
15 process (Chee, M. et al., PCT/WO95/11995, incorporated herein by reference).

In the alternative, a chemical coupling procedure and an ink jet device are used to synthesize oligomers on the surface of a substrate (Baldeschweiler, J.D. et al., PCT/WO95/25116, incorporated herein by reference). In another alternative, a "gridded" array analogous to a dot (or slot) blot is used to arrange and link cDNA fragments or
20 oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array may be produced by hand or using available materials and machines and contain grids of 8 dots, 24 dots, 96 dots, 384 dots, 1536 dots or 6144 dots. After hybridization, the microarray is washed to remove nonhybridized probes, and a scanner is used to determine the levels and patterns of fluorescence. The
25 scanned images are examined to determine degree of complementarity and the relative abundance of each oligonucleotide sequence on the micro-array.

VIII Complementary Polynucleotides

Sequence complementary to the SIGH-encoding sequence, or any part thereof, is
30 used to decrease or inhibit expression of naturally occurring SIGH. Although use of oligonucleotides comprising from about 15 to about 30 base-pairs is described, essentially the same procedure is used with smaller or larger sequence fragments. Appropriate

oligonucleotides are designed using Oligo 4.06 software and the coding sequence of Sigr, SEQ ID NO:1. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the Sigr-encoding transcript.

IX Expression of Sigr

Expression of Sigr is accomplished by subcloning the cDNAs into appropriate vectors and transforming the vectors into host cells. In this case, the cloning vector is also used to express Sigr in *E. coli*. Upstream of the cloning site, this vector contains a promoter for β -galactosidase, followed by sequence containing the amino-terminal Met, and the subsequent seven residues of β -galactosidase. Immediately following these eight residues is a bacteriophage promoter useful for transcription and a linker containing a number of unique restriction sites.

Induction of an isolated, transformed bacterial strain with IPTG using standard methods produces a fusion protein which consists of the first eight residues of β -galactosidase, about 5 to 15 residues of linker, and the full length protein. The signal residues direct the secretion of Sigr into the bacterial growth media which can be used directly in the following assay for activity.

X Demonstration of Sigr Activity

Human sigma receptor activity is analyzed using the (+)[³H]pentazocine binding assay described by Hanner et al. (1996, supra). 200 μ g Sigr in the absence or presence of 0.3 mg/ml phosphatidylcholine is incubated with 0.4 to 2.1 nM (+)[³H]pentazocine (32 Ci/mmol, NEN) in 0.25 or 0.5 ml 25 mM Tris-HCl (Tris buffer), pH 8.3 at 22°C, for between 4 and twelve hours at 4°C. Nonspecific binding defined by 1 μ M pentazocine is subtracted from the total binding to yield specific binding. Bound ligand is separated from free ligand by filtration over GF/C glass fiber filters (Whatman) followed by washing with Tris buffer. The glass filters are air dried, then counted in a liquid scintillation counter. Binding parameters are calculated by non-linear curve fitting to a rectangular hyperbola (K_d , B_{max}) or the general dose-response equation (IC_{50} , slope factor).

XI Production of SIGR Specific Antibodies

SIGR that is substantially purified using PAGE electrophoresis (Sambrook, supra), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols. The amino acid sequence deduced from SEQ ID NO:2 is analyzed using
5 DNASTAR software (DNASTAR Inc) to determine regions of high immunogenicity and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions, is described by Ausubel et al. (supra), and others.

Typically, the oligopeptides are 15 residues in length, synthesized using an Applied
10 Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry, and coupled to keyhole limpet hemocyanin (KLH, Sigma, St. Louis, MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS; Ausubel et al., supra). Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. The resulting antisera are tested for antipeptide activity, for example, by binding the peptide to plastic, blocking with 1%
15 BSA, reacting with rabbit antisera, washing, and reacting with radio iodinated, goat anti-rabbit IgG.

XII Purification of Naturally Occurring SIGR Using Specific Antibodies

Naturally occurring or recombinant SIGR is substantially purified by immunoaffinity
20 chromatography using antibodies specific for SIGR. An immunoaffinity column is constructed by covalently coupling SIGR antibody to an activated chromatographic resin, such as CNBr-activated Sepharose (Pharmacia & Upjohn). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing SIGR is passed over the immunoaffinity column, and the column is
25 washed under conditions that allow the preferential absorbance of SIGR (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/SIGR binding (eg, a buffer of pH 2-3 or a high concentration of a chaotrope, such as urea or thiocyanate ion), and SIGR is collected.

XIII Identification of Molecules Which Interact with SIGR

SIGR or biologically active fragments thereof are labeled with ¹²⁵I Bolton-Hunter reagent (Bolton et al. (1973) Biochem. J. 133: 529). Candidate molecules previously arrayed

WO 99/13055

PCT/US98/18941

in the wells of a multi-well plate are incubated with the labeled SIGR, washed and any wells with labeled SIGR complex are assayed. Data obtained using different concentrations of SIGR are used to calculate values for the number, affinity, and association of SIGR with the candidate molecules.

5 All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed
10 should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

WO 99/13055

PCT/US98/18941

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.
 HILLMAN, Jennifer >
 CORLEY, Neil C.
 SHAH, Purvi

<120> HUMAN SIGMA RECEPTOR

<130> PF-0383 PCT

<140> To Be Assigned

<141> Herewith

<150> 08/928,612

<151> 1997-09-12

<160>3

<170> PERL Program

<210> 1

<211> 223

<212> PRT

<213> Homo sapiens

<220> -

<223> 1542751

<400> 1

Met	Gln	Trp	Ala	Val	Gly	Arg	Arg	Trp	Ala	Trp	Ala	Ala	Leu	Leu	1
				5					10					15	
Leu	Ala	Val	Ala	Ala	Val	Leu	Thr	Gln	Val	Val	Trp	Leu	Trp	Leu	20
				20					25					30	
Gly	Thr	Gln	Ser	Phe	Val	Phe	Gln	Arg	Glu	Glu	Ile	Ala	Gln	Leu	35
				35					40					45	
Ala	Arg	Gln	Tyr	Ala	Gly	Leu	Asp	His	Glu	Leu	Ala	Phe	Ser	Arg	50
				50					55					60	
Leu	Ile	Val	Glu	Leu	Arg	Arg	Leu	His	Pro	Gly	His	Val	Leu	Pro	65
				65					70					75	
Asp	Glu	Glu	Leu	Gln	Trp	Val	Phe	Val	Asn	Ala	Gly	Gly	Trp	Met	80
				80					85					90	
Gly	Ala	Met	Cys	Leu	Leu	His	Ala	Ser	Leu	Ser	Glu	Tyr	Val	Leu	95
				95					100					105	
Leu	Phe	Gly	Thr	Ala	Leu	Gly	Ser	Arg	Gly	His	Ser	Gly	Arg	Tyr	110
				110					115					120	
Trp	Ala	Glu	Ile	Ser	Asp	Thr	Ile	Ile	Ser	Gly	Thr	Phe	His	Gln	125
				125					130					135	
Trp	Arg	Glu	Gly	Thr	Thr	Lys	Ser	Glu	Val	Phe	Tyr	Pro	Gly	Glu	140
				140					145					150	
Thr	Val	Val	His	Gly	Pro	Gly	Glu	Ala	Thr	Ala	Val	Glu	Trp	Gly	155
				155					160					165	
Pro	Asn	Thr	Trp	Met	Val	Glu	Tyr	Gly	Arg	Gly	Val	Ile	Pro	Ser	170
				170					175					180	
Thr	Leu	Ala	Phe	Ala	Leu	Ala	Asp	Thr	Val	Phe	Ser	Thr	Gln	Asp	185
				185					190					195	
Phe	Leu	Thr	Leu	Phe	Tyr	Thr	Leu	Arg	Ser	Tyr	Ala	Arg	Gly	Leu	200
				200					205					210	
Arg	Leu	Glu	Leu	Thr	Thr	Tyr	Leu	Phe	Gly	Gln	Asp	Pro			215
				215					220						

WO 99/13055

PCT/US98/18941

<210> 2
 <211> 1700
 <212> DNA
 <213> Homo sapiens

<220> -
 <223> 1542751

<400> 2
 cggctcgagg gcgggggagc gcttcgtggg cagccggcgg gctccgaggc cgtgagcgca 60
 aagcctcagg ccccggtcc ctctgagct gcgccgtgcc aggccgcccg ccgggatgca 120
 gtgggcccgtg ggccggcggt gggcgtgggc cgcgctgtct ctggctgtcg cagcgggtgct 180
 gacccaggtc gtctggctct ggctgggtac gcagagcttc gtcttccagc gcgaagagat 240
 agcgcagttg gcgcggcagt acgctgggct ggaccacgag ctggccttct ctcgtctgat 300
 cgtggagctg cggcggctgc acccaggcca cgtgctgccc gacgaggagc tgcagtgggt 360
 gttcgtgaat gcgggtggct ggatgggcgc catgtgcctt ctgcacgcct cgctgtccga 420
 gtatgtgctg ctcttcggca ccgccttggg ctcccgcggc cactcggggc gctactgggc 480
 tgagatctcg gataccatca tctctggcac ctccaccag tggagagagg gcaccaccaa 540
 aagtgaggtc ttctaccag gggagacggt agtacacggg cctggtgagg caacagctgt 600
 ggagtggggg ccaaacacat ggatggtgga gtacggccgg ggcgtcatcc catccaccct 660
 ggccttcgcg ctggccgaca ctgtcttcag caccaggac ttctcacc tcttctatac 720
 tcttcgctcc tatgctcggg gcctccggct tgagctcacc acctacctt ttggccagga 780
 cccttgacca gccaggcctg aaggaagacc tgcggatgga caggagcggg caggcccgca 840
 catatccact tgctggagcc catgtttaca gacagggaca tacaccatgc agatcctgag 900
 ttctgtctgt atgagcaggg atatccatgc ttatgtatcc aaacacagag acctatggga 960
 acaaatgaga cacatataga tactgagacc tgtgtgtaca gtaggacat gcactcacac 1020
 ccatctggag agggagcccc cggatatacca agggagccag ttgtgttcag acacacacat 1080
 cacagcttga ctactaact gaggccttcc catagctcca cagcttccca cctcctcccc 1140
 accaaaccgg ggttctagag ttaaggatgg gggagggtat tatactgcct cagtctgact 1200
 cctcaaccca gcagcaattt gaggggatga gggggaagag gagctgcctt ttggaggccc 1260
 ccttcacctg cagctatgat gcccttcccc ttctcccctg tcctcaccat atgccttacc 1320
 cccattctac tcccctgcta tgcaagtgcc cctgtggctt gtccccaacc ccctcagcaa 1380
 caaagctcag ctggggaacg agagtaattt gaagaatgct tgaagtcagc gtcttccatt 1440
 ccagaaagac cccattctt cctttggggg tatgatgtgg aagctggttt cagcccagga 1500
 cccaccactg aggagaggat ctagacaggt gggcctaatt ccaaggggcc cttcctggcc 1560
 tggagaaggc cttttacaca cacacaacac atacacacac acacacacac acacatatca 1620
 cagttttcac acagcccctg ctgcattctc tgtccatctg tctgtttcta ttaataaaga 1680
 tttgttgatc tgttcaaaaa 1700

<210> 3
 <211> 223
 <212> PRT
 <213> Homo sapiens

<220> -
 <223> g1403300

<400> 3
 Met Gln Trp Ala Val Gly Arg Arg Trp Leu Trp Val Ala Leu Phe
 1 5 10 15
 Leu Ala Ala Val Ala Val Leu Thr Gln Ile Val Trp Leu Trp Leu
 20 25 30
 Gly Thr Gln Asn Phe Val Phe Gln Arg Glu Glu Ile Ala Gln Leu
 35 40 45
 Ala Arg Gln Tyr Ala Gly Leu Asp His Glu Leu Ala Phe Ser Lys
 50 55 60
 Leu Ile Val Glu Leu Arg Arg Leu His Pro Val His Val Leu Pro
 65 70 75
 Asp Glu Glu Leu Gln Trp Val Phe Val Asn Ala Gly Gly Trp Met
 80 85 90
 Gly Ala Met Cys Leu Leu His Ala Ser Leu Ser Glu Tyr Val Leu
 95 100 105
 Leu Phe Gly Thr Ala Leu Gly Ser Pro Arg His Ser Gly Arg Tyr
 110 115 120

WO 99/13055

PCT/US98/18941

Trp	Ala	Glu	Ile	Ser	Asp	Thr	Ile	Ile	Ser	Gly	Thr	Phe	His	Gln
				125					130					135
Trp	Arg	Glu	Gly	Thr	Thr	Lys	Ser	Glu	Val	Phe	Tyr	Pro	Gly	Glu
				140					145					150
Thr	Val	Val	His	Gly	Pro	Gly	Glu	Ala	Thr	Ala	Val	Glu	Trp	Gly
				155					160					165
Pro	Asn	Thr	Trp	Met	Val	Glu	Tyr	Gly	Arg	Gly	Val	Ile	Pro	Ser
				170					175					180
Thr	Leu	Gly	Phe	Ala	Leu	Ala	Asp	Thr	Val	Phe	Ser	Thr	Gln	Asp
				185					190					195
Phe	Leu	Thr	Leu	Phe	Tyr	Thr	Leu	Arg	Val	Tyr	Ala	Arg	Ala	Leu
				200					205					210
Gln	Leu	Glu	Leu	Thr	Thr	Tyr	Leu	Phe	Gly	Gln	Asp	Pro		
				215					220					

WO 99/13055

PCT/US98/18941

What is claimed is:

1. A substantially purified human sigma receptor comprising the amino acid sequence of SEQ ID NO:1 or fragments thereof.
5
2. A substantially purified variant of human sigma receptor having at least 90% amino acid identity to SEQ ID NO:1 and which retains at least one functional characteristic of human sigma receptor.
- 10 3. An isolated and purified polynucleotide sequence encoding the human sigma receptor of claim 1 or fragments or variants of said polynucleotide sequence.
4. A composition comprising the polynucleotide sequence of claim 3.
- 15 5. A polynucleotide sequence which hybridizes to the polynucleotide sequence of claim 3.
6. A polynucleotide sequence which is complementary to the polynucleotide sequence of claim 3 or fragments or variants thereof.
20
7. An isolated and purified polynucleotide sequence comprising SEQ ID NO:2 or fragments or variants thereof.
8. A polynucleotide sequence which is complementary to the polynucleotide
25 sequence of claim 7.
9. An expression vector containing at least a fragment of the polynucleotide sequence of claim 3.
- 30 10. A host cell containing the vector of claim 9.
11. A method for producing a polypeptide comprising the amino acid sequence of

SEQ ID NO:1, or a fragment thereof, the method comprising the steps of:

- a) culturing the host cell of claim 10 under conditions suitable for the expression of the polypeptide; and
- b) recovering the polypeptide from the host cell culture.

5

12. A pharmaceutical composition comprising a substantially purified human sigma receptor having the amino acid sequence of SEQ ID NO:1 in conjunction with a suitable pharmaceutical carrier.

10

13. A purified antibody which specifically binds to the polypeptide of claim 1.

14. A purified agonist of the polypeptide of claim 1.

15. A purified antagonist of the polypeptide of claim 1.

15

16. A method for treating a vesicle trafficking disorder comprising administering to a subject in need of such treatment an effective amount of the pharmaceutical composition of claim 12.

20

17. A method for treating an immunological disorder comprising administering to a subject in need of such treatment an effective amount of the pharmaceutical composition of claim 12.

25

18. A method for treating a neoplastic disorder comprising administering to a subject in need of such treatment an effective amount of the pharmaceutical composition of claim 12.

19. A method for detecting a polynucleotide which encodes human sigma receptor in a biological sample comprising the steps of:

30

- a) hybridizing the polynucleotide of claim 6 to nucleic acid material of a biological sample, thereby forming a hybridization complex; and
- b) detecting said hybridization complex, wherein the presence of said

complex correlates with the presence of a polynucleotide encoding human sigma receptor in said biological sample.

20. The method of claim 19 wherein the nucleic acid material is amplified by the
5 polymerase chain reaction prior to hybridization.

9	18	27	36	45	54
C GGC TCG AGG GCG GGG GAG CGC TTC GTG GGC AGC CGG CGG GCT CCG AGG CCG					
63	72	81	90	99	108
TGA GCG CAA AGC CTC AGG CCC CGG CTC CCT CCT GAG GAG CTG CGC CGT GCC AGG CCG					
117	126	135	144	153	162
CCC GCC GGG ATG CAG TGG GCC GTG GGC CGG CGG TGG GCG TGG GCC CTG CTC					
	M Q W A V G R R A W A A L L				
171	180	189	198	207	216
CTG GCT GTC GCA GCG GTG CTG ACC CAG GTC GTC TGG CTC TGG CTG GGT ACG CAG					
L A V A A V L T Q V V W L W L G T Q					
225	234	243	252	261	270
AGC TTC GTC TTC CAG CGC GAA GAG ATA GCG CAG TTG GCG CGG CAG TAC GCT GGG					
S F V F Q R E E I A Q L A R Q Y A G					
279	288	297	306	315	324
CTG GAC CAC GAG CTG GCC TTC TCT CGT CGT ATC GTG GAG CTG GAG CTG CCG CTG CAC					
L D H E L A F S R L L I V E L R R L H					
333	342	351	360	369	378
CCA GGC CAC GTG CTG CCC GAC GAG GAG CTG CAG TGG GTG TTC GTG AAT GCG GGT					
P G H V L P D E E E L L Q W V F V N A G					

FIGURE 1A

387	GGC	TGG	ATG	GGC	GCC	ATG	TGC	CTT	CTG	CAC	GCC	TCG	CTG	TCC	GAG	TAT	GTG	CTG	432
	G	W	M	G	A	M	C	L	L	H	A	S	L	S	E	Y	V	L	
441	CTC	TTC	GGC	ACC	GCC	TTG	GGC	TCC	CGC	GGC	CAC	TCG	GGG	CGC	TAC	TGG	GCT	GAG	486
	L	F	G	T	A	L	G	S	R	G	H	S	G	R	Y	W	A	E	
495	ATC	TCG	GAT	ACC	ATC	ATC	TCT	GGC	ACC	TTC	CAC	CAG	TGG	AGA	GAG	GGC	ACC	ACC	540
	I	S	D	T	I	I	S	G	T	F	H	Q	W	R	E	G	T	T	
549	AAA	AGT	GAG	GTC	TTC	TAC	CCA	GGG	GAG	ACG	GTA	GTA	CAC	GGG	CCT	GGT	GAG	GCA	594
	K	S	E	V	F	Y	P	G	E	T	V	V	H	G	P	G	E	A	
603	ACA	GCT	GTG	GAG	TGG	GGG	CCA	AAC	ACA	TGG	ATG	GTG	GAG	TAC	GGC	CGG	GGC	GTC	648
	T	A	V	E	W	G	P	N	T	W	M	V	E	Y	G	R	G	V	
657	ATC	CCA	TCC	ACC	CTG	GCC	TTC	GCG	CTG	GCC	GAC	ACT	GTC	TTC	AGC	ACC	CAG	GAC	702
	I	P	S	T	L	A	F	A	L	A	D	T	V	F	S	T	Q	D	
711	TTC	CTC	ACC	CTC	TTC	TAT	ACT	CTT	CGC	TCC	TAT	GCT	CGG	GGC	CTC	CGG	CTT	GAG	756
	F	L	T	L	F	Y	T	L	R	S	Y	A	R	G	L	R	L	E	

FIGURE 1B

3/8

765	774	783	792	801	810
CTC ACC ACC TAC CTC TTT GGC CAG GAC CCT TGA CCA GCC AGG CCT GAA GGA AGA					
L T T Y L F G Q D P					
819	828	837	846	855	864
CCT GCG GAT GGA CAG GAG CGG GCA GGC CCG CAC ATA TCC ACT TGC TGG AGC CCA					
873	882	891	900	909	918
TGT TTA CAG ACA GGG ACA TAC ACC ATG CAG ATC CTG AGT TCC TGC TGT ATG AGC					
927	936	945	954	963	972
AGG GAT ATC CAT GCT TAT GTA TCC AAA CAC AGA GAC CCA TGG GAA CAA ATG AGA					
981	990	999	1008	1017	1026
CAC ATA TAG ATA CTG AGA CCT GTG TGT ACA GTA GGA CCA TGC ACT CAC ACC CAT					
1035	1044	1053	1062	1071	1080
CTG GAG AGG GAG CCC CCG GTA TAC CAA GGC AGC CAG TTG TGT TCA GAC ACA CAC					
1089	1098	1107	1116	1125	1134
ATC ACA GCT TGA CTC ACT AAC TGA GGC CTT TCC ATA GCT CCA CAG CTT CCC ACC					
1143	1152	1161	1170	1179	1188
TCC TCC CCA CCA AAC CGG GGT TCT AGA GTT AAG GAT GGG GGA GGG TAT TAT ACT					
1197	1206	1215	1224	1233	1242
GCC TCA GTC TGA CTC CTC AAC CCA GCA GCA ATT TGA GGG GAT GAG GGG GAA GAG					

FIGURE 1C

1251	1260	1269	1278	1287	1296
GAG CTG CCT TTT GGA GGC CCC CTT CAC CTG CAG CTA TGA TGC CCT TCC CCT TCT					
1305	1314	1323	1332	1341	1350
CCC CTG TCC TCA CCA TAT GCC TTA TCC CCA TTC TAC TCC CCT GCT ATG CAA GTG					
1359	1368	1377	1386	1395	1404
CCC CTG TGG CTT GTC CCC AAC CCC CTC AGC AAC AAA GCT CAG CTG GGG AAC GAG					
1413	1422	1431	1440	1449	1458
AGT AAT TTG AAG AAT GCT TGA AGT CAG CGT CTT CCA TTC CAG AAA GAC CCC CAT					
1467	1476	1485	1494	1503	1512
TCT TCC TTT GGG GGT ATG ATG TGG AAG CTG GTT TCA GCC CAG GAC CCA CCA CTG					
1521	1530	1539	1548	1557	1566
AGG AGA GGA TCT AGA CAG GTG GGC CTA ATT CCA AGG GGC CCT TCC TGG CCT GGA					
1575	1584	1593	1602	1611	1620
GAA GGC CTT TTA CAC ACA CAC AAC ACA TAC ACA CAC ACA CAC ACA TAT					
1629	1638	1647	1656	1665	1674
CAC AGT TTT CAC ACA GCC CCT GCT GCA TTC TCT GTC CAT CTG TCT GTT TCT ATT					
1683	1692	1701			
AAT AAA GAT TTG TTG ATC TGT TCA AAA A					

FIGURE 1D

1	M	Q	W	A	V	G	R	R	W	A	A	L	L	L	A	V	A	A	1542751
1	M	Q	W	A	V	G	R	R	W	L	W	V	A	L	F	L	A	V	GI 1403300
21	V	L	T	Q	V	V	W	L	W	L	G	T	Q	S	F	V	F	Q	1542751
21	V	L	T	Q	I	V	W	L	W	L	G	T	Q	N	F	V	F	Q	GI 1403300
41	E	I	A	Q	L	A	R	Q	Y	A	G	L	D	H	E	L	A	F	1542751
41	E	I	A	Q	L	A	R	Q	Y	A	G	L	D	H	E	L	A	F	GI 1403300
61	L	I	V	E	L	R	R	L	H	P	G	H	V	L	P	D	E	E	1542751
61	L	I	V	E	L	R	R	L	H	P	V	H	V	L	P	D	E	E	GI 1403300
81	W	V	F	V	N	A	G	G	W	M	G	A	M	C	L	L	H	A	1542751
81	W	V	F	V	N	A	G	G	W	M	G	A	M	C	L	L	H	A	GI 1403300
101	S	E	Y	V	L	L	F	G	T	A	L	G	S	R	G	H	S	G	1542751
101	S	E	Y	V	L	L	F	G	T	A	L	G	S	P	R	H	S	G	GI 1403300
121	W	A	E	I	S	D	T	I	I	S	G	T	F	H	Q	W	R	E	1542751
121	W	A	E	I	S	D	T	I	I	S	G	T	F	H	Q	W	R	E	GI 1403300

FIGURE 2A

141	T	K	S	E	V	F	Y	P	G	E	T	V	V	H	G	P	G	E	A	T	1542751
141	T	K	S	E	V	F	Y	P	G	E	T	V	V	H	G	P	G	E	A	T	GI 1403300
161	A	V	E	W	G	P	N	T	W	M	V	E	Y	G	R	G	V	I	P	S	1542751
161	A	V	E	W	G	P	N	T	W	M	V	E	Y	G	R	G	V	I	P	S	GI 1403300
181	T	L	A	F	A	L	A	D	T	V	F	S	T	Q	D	F	L	T	L	F	1542751
181	T	L	G	F	A	L	A	D	T	V	F	S	T	Q	D	F	L	T	L	F	GI 1403300
201	Y	T	L	R	S	Y	A	R	G	L	R	L	E	L	T	T	Y	L	F	G	1542751
201	Y	T	L	R	V	Y	A	R	A	L	Q	L	E	L	T	T	Y	L	F	G	GI 1403300
221	Q	D	P																		1542751
221	Q	D	P																		GI 1403300

FIGURE 2B

7/8




FIGURE 3A

8/8




FIGURE 3B