



(51) International Patent Classification:

C07D 313/08 (2006.01) *A61K 31/44* (2006.01)
C07D 405/04 (2006.01) *A61P 3/10* (2006.01)
A61K 31/335 (2006.01) *A61P 3/08* (2006.01)

(21) International Application Number:

PCT/CN2014/084034

(22) International Filing Date:

8 August 2014 (08.08.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(71) Applicant: **MERCK SHARP & DOHME CORP.**
 [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey
 07065 (US).

(72) Inventors; and

(71) Applicants (*for US only*): **PLUMMER, Christopher W.**
 [US/US]; Merck Sharp & Dohme Corp., 2000 Galloping
 Hill Road, Kenilworth, NJ 07090 (US). **DeMONG, Duane**
 [US/US]; Merck Sharp & Dohme Corp., 2000 Galloping
 Hill Road, Kenilworth, NJ 07090 (US). **FANG, Minghai**
 [CN/CN]; WuXi AppTec (Shanghai) Co., Ltd., 288 Fute
 Zhong Road, Waigaoqiao Free Trade Zone, Shanghai
 200131 (CN). **HU, Bin** [CN/CN]; WuXi AppTec (Shang-
 hai) Co., Ltd., 288 Fute Zhong Road, Waigaoqiao Free
 Trade Zone, Shanghai 200131 (CN).

(74) Agent: **WU, FENG & ZHANG CO.**; Room 305, Tower
 B, Beijing Aerospace CPMIEC Building, No. 30, Haidian
 South Road, Haidian District, Beijing 100080 (CN).


(81) Designated States (*unless otherwise indicated, for every
 kind of national protection available*): AE, AG, AL, AM,
 AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
 BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
 DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
 HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
 KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
 MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
 OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
 SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,
 TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
 ZW.

(84) Designated States (*unless otherwise indicated, for every
 kind of regional protection available*): ARIPO (BW, GH,
 GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
 UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
 TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
 EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
 MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
 TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
 KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: [7, 6]-FUSED BICYCLIC ANTIDIABETIC COMPOUNDS



(57) Abstract: Provided are compounds of the structural formula (I), the pharmaceutically acceptable salts thereof, and the use as agonists of G-protein coupled receptor 40 (GPR40) for the treatment of Type 2 diabetes mellitus, and of conditions that are often associated with this disease, including obesity and lipid disorders, such as mixed or diabetic dyslipidemia, hyperlipidemia, hypercholesterolemia, and hypertriglyceridemia.

[7,6]-FUSED BICYCLIC ANTIDIABETIC COMPOUNDS

BACKGROUND OF THE INVENTION

Diabetes mellitus is a disease derived from multiple causative factors and characterized by elevated levels of plasma glucose (hyperglycemia) in the fasting state or after administration of glucose during an oral glucose tolerance test. There are two generally recognized forms of diabetes. In Type 1 diabetes, or insulin-dependent diabetes mellitus (IDDM), patients produce little or no insulin, the hormone which regulates glucose utilization. In Type 2 diabetes, or noninsulin-dependent diabetes mellitus (NIDDM), insulin is still produced in the body, however patients have a resistance to the effects of insulin in stimulating glucose and lipid metabolism in the insulin-sensitive muscle, liver and adipose tissues. Type 2 diabetes patients often have normal levels of insulin, and may have hyperinsulinemia (elevated plasma insulin levels), as they compensate for the reduced effectiveness of insulin by secreting increased amounts of insulin. This lack of responsiveness to insulin results in insufficient insulin-mediated activation of uptake, oxidation and storage of glucose in muscle, and inadequate insulin-mediated repression of lipolysis in adipose tissue and of glucose production and secretion in the liver.

Persistent or uncontrolled hyperglycemia is associated with increased and premature morbidity and mortality. Abnormal glucose homeostasis is associated both directly and indirectly with obesity, hypertension, and alterations of the lipid, lipoprotein and apolipoprotein metabolism, as well as other metabolic and hemodynamic disease. Patients with Type 2 diabetes mellitus have a significantly increased risk of macrovascular and microvascular complications, including atherosclerosis, coronary heart disease, stroke, peripheral vascular disease, hypertension, nephropathy, neuropathy, and retinopathy. Therefore, therapeutic control of glucose homeostasis, lipid metabolism, obesity, and hypertension are critically important in the clinical management and treatment of diabetes mellitus.

Patients who have insulin resistance often have Metabolic Syndrome (as defined in the Third Report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III, or ATP III), National Institutes of Health, 2001, NIH Publication No. 01-3670). Patients with Metabolic Syndrome have an increased risk of developing the macrovascular and microvascular complications that occur with Type 2 diabetes, such as atherosclerosis and coronary heart disease.

There are several available treatments for Type 2 diabetes. Physical exercise and a reduction in dietary intake of calories are the recommended first-line treatment of Type 2 diabetes and of pre-diabetic conditions associated with insulin resistance, however compliance is generally poor. Pharmacologic treatments for diabetes have largely focused on three areas of pathophysiology: (1) hepatic glucose production (biguanides, such as phenformin and metformin), (2) insulin resistance

(PPAR agonists, such as rosiglitazone, troglitazone, englitazone, balaglitazone, and pioglitazone), (3) insulin secretion (sulfonylureas, such as tolbutamide, glipizide and glimepiride); (4) incretin hormone mimetics (GLP-1 derivatives and analogs, such as exenatide, liraglutide, dulaglutide, semaglutide, lixisenatide, albiglutide and taspoglutide); and (5) inhibitors of incretin hormone degradation (DPP-4 inhibitors, such as sitagliptin, alogliptin, vildagliptin, linagliptin, denagliptin and saxagliptin).

The two best known biguanides, phenformin and metformin, cause some correction of hyperglycemia, but can also induce lactic acidosis and nausea/diarrhea. PPAR gamma agonists, such as rosiglitazone and pioglitazone, are modestly effective in reducing plasma glucose and Hemoglobin A1C. However, the currently marketed glitazones do not greatly improve lipid metabolism and may negatively effect on the lipid profile. The administration of insulin secretagogues, such as the sulfonylureas (e.g. tolbutamide, glipizide, and glimepiride) can result in hypoglycemia; their administration must therefore be carefully controlled.

There has been a renewed focus on pancreatic islet-based insulin secretion that is controlled by glucose-dependent insulin secretion. This approach has the potential for stabilization and restoration of β -cell function. Several orphan G-protein coupled receptors (GPCR's) have been identified that are preferentially expressed in the β -cell and that are implicated in glucose stimulated insulin secretion (GSIS). GPR40 is a cell-surface GPCR that is highly expressed in human (and rodent) islets as well as in insulin-secreting cell lines. Several naturally-occurring medium to long-chain fatty acids (FA's) as well as synthetic compounds, including several members of the thiazolidinedione class of PPAR γ agonists, have recently been identified as ligands for GPR40 [Itoh, Y. et al., *Nature*, 422: 173 (2003); Briscoe, C.P. et al., *J. Biol. Chem.*, 278: 11303 (2003); Kotarsky, K. et al., *Biochem. Biophys. Res. Comm.*, 301: 406 (2003)]. Under hyperglycemic conditions, GPR40 agonists are capable of augmenting the release of insulin from islet cells. The specificity of this response is suggested by results showing that the inhibition of GPR40 activity by siRNA attenuates FA-induced amplification of GSIS. These findings indicate that, in addition to the intracellular generation of lipid-derivatives of FA's that are thought to promote insulin release, FA's (and other synthetic GPR40 agonists) may also act as extracellular ligands that bind to GPR40 in mediating FA-induced insulin secretion. There are several potential advantages of GPR40 as a target for the treatment of Type 2 diabetes. First, since GPR40-mediated insulin secretion is glucose dependent, there is little or no risk of hypoglycemia. Second, the limited tissue distribution of GPR40 (mainly in islets) suggests that there would be less chance for side effects associated with GPR40 activity in other tissues. Third, GPR40 agonists that are active in the islets may have the potential to restore or preserve islet function. This would be advantageous, because long term diabetes therapy often leads to the gradual diminution of islet activity; after extended periods of treatment, it is often necessary to treat Type 2 diabetic patients with daily insulin injections. By

restoring or preserving islet function, GPR40 agonists may delay or prevent the diminution and loss of islet function in a Type 2 diabetic patient.


Compounds that are agonists of G-protein-coupled receptor 40 (GPR40) may be useful to treat type 2 diabetes mellitus, obesity, hypertension, dyslipidemia, cancer, and metabolic syndrome, as well as cardiovascular diseases, such as myocardial infarction and stroke, by improving glucose and lipid metabolism and by reducing body weight. There is a need for potent GPR40 agonists that have pharmacokinetic and pharmacodynamic properties suitable for use as human pharmaceuticals.

G-protein-coupled receptor 40 (GPR40) agonists are disclosed in WO 2007/136572, WO 2007/136573, WO 2009/058237, WO 2006/083612, WO 2006/083781, WO 2010/085522, WO 2010/085525, WO 2010/085528, WO 2010/091176, WO 2004/041266, EP 2004/1630152, WO 2004/022551, WO 2005/051890, WO 2005/051373, EP 2004/1698624, WO 2005/086661, WO 2007/213364, WO 2005/063729, WO 2005/087710, WO 2006/127503, WO 2007/1013689, WO 2006/038738, WO 2007/033002, WO 2007/106469, WO 2007/123225, WO 2008/001931, WO 2008/030520, WO 2008/030618, WO 2008/054674, WO 2008/054675, WO 2008/066097, WO 2008/130514, WO 2009/048527, WO 2009/058237, WO 2009/111056, WO 2010/004347, WO 2010/045258, WO 2010/085522, WO 2010/085525, WO 2010/085528, WO 2010/091176, WO 2010/143733, WO 2012/0004187, WO 2012/011125, WO 2012/072691, WO2013/104257, WO 2013/122028, WO 2013/122029, WO 2013/128378, WO 2013/178575, WO 2014/073904, WO 2014/078608, WO 2014/078609, WO 2014/078610, WO 2014/078611, US 8030354, US 8450522, CN 103030646, CN 103012343, and GB 2498976.

GPR40 agonists are also disclosed in Negoro et al., ACS Medicinal Chemistry Letters (2010), 1(6), 290-294; Walsh et al., Bioorganic & Medicinal Chemistry Letters (2011), 21(11), 3390-3394; Zhou et al., Bioorganic & Medicinal Chemistry Letters (2010), 20(3), 1298-1301; Houze et al., Bioorganic & Medicinal Chemistry Letters (2012), 22(2), 1267-1270; Lu et al., Bioorganic & Medicinal Chemistry Letters (2013), 23(10), 2920-2924; Takano et al., Bioorganic & Medicinal Chemistry Letters (2014), 24(13), 2949-2953; Tan et al., Diabetes (2008), 57(8), 2211-2219; Brown et al., ACS Medicinal Chemistry Letters (2012), 3(9), 726-730; Lin et al., PloS One (2011), 6(11), e27270; Lou et al., PloS One (2012), 7(10), e46300; Lin et al., Molecular Pharmacology (2012), 82(5), 843-859; Yang, Lihu, Abstracts of Papers, 239th ACS Meeting, San Francisco, CA, USA March 21-25, 2010 MEDI-313; Houze et al., Abstracts of Papers, 243rd ACS National Meeting & Exposition, San Diego, CA, USA March 25-29, 2012, MEDI-265; Wang et al., ACS Medicinal Chemistry Letters (2013), 4(6), 551-555; and Du et al., ACS Medicinal Chemistry Letters (2014), 5(4), 384-389.

SUMMARY OF THE INVENTION

The present invention relates to novel substituted compounds of structural formula I:




and pharmaceutically acceptable salts thereof. The compounds of structural formula I, and
embodiments thereof, are agonists of G-protein-coupled receptor 40 (GPR40) and may be useful in
the treatment, prevention and suppression of diseases, disorders and conditions mediated by
agonism of the G-protein-coupled receptor 40, such as Type 2 diabetes mellitus, insulin resistance,
hyperglycemia, dyslipidemia, lipid disorders, obesity, hypertension, Metabolic Syndrome and
atherosclerosis.

The present invention also relates to pharmaceutical compositions comprising the
compounds of the present invention and a pharmaceutically acceptable carrier. The present
invention also relates to methods for the treatment, control or prevention of disorders, diseases, and
conditions that may be responsive to agonism of the G-protein-coupled receptor 40 in a subject in
need thereof by administering the compounds and pharmaceutical compositions of the present
invention. The present invention also relates to the use of compounds of the present invention for
manufacture of a medicament useful in treating diseases, disorders and conditions that may be
responsive to the agonism of the G-protein-coupled receptor 40. The present invention is also
concerned with treatment of these diseases, disorders and conditions by administering the
compounds of the present invention in combination with a therapeutically effective amount of
another agent that may be useful to treat the disease, disorder and condition. The invention is
further concerned with processes for preparing the compounds of this invention.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is concerned with novel compounds of structural Formula I:



5 or a pharmaceutically acceptable salt thereof; wherein

"a" is a single bond or a double bond, provided that if "a" is a double bond, then R⁵ and R⁶ are absent and Q is selected from the group consisting of: C-R⁴, -C-OC₁₋₆alkyl, and N, and further provided that if "a" is a single bond, then R⁵ and R⁶ are present and Q is selected from the group consisting of:

- 10 (1) oxygen,
- (2) sulfur,
- (3) -CR⁴R⁶,
- (4) S(O)₂,
- (5) C=O,
- 15 (6) -C(R⁹)OC₁₋₆alkyl, and
- (7) -NR^c;

T is selected from the group consisting of:

- (1) CH,
- (2) N, and
- 20 (3) N-oxide;

U is selected from the group consisting of:

- (1) CR¹,
- (2) N, and
- (3) N-oxide;

25 V is selected from the group consisting of:

- (1) CR²,
- (2) N, and
- (3) N-oxide;

W is selected from the group consisting of:

- 30 (1) CH,

- (2) N, and
- (3) N-oxide,

provided that no more than two of T, U, V and W are selected from N and N-oxide, further provided that if both T and W are N or N-oxide, then R^3 is absent, and further provided that both U and V are not N or N-oxide;

X is selected from the group consisting of:

- (1) oxygen,
- (2) sulfur,
- (3) $-CR^9R^9$,
- (4) $S(O)_2$,
- (5) $C=O$,
- (6) $-C(R^9)OC_{1-6}alkyl$, and
- (7) $-NR^c$,

provided that if Q is oxygen, sulfur or $-NR^c$, then X is not oxygen, sulfur or $-NR^c$, further provided that if Q is $C=O$, then X is not $C=O$ or $S(O)_2$, and further provided that if Q is $S(O)_2$, then X is not $S(O)_2$, $C=O$, oxygen or sulfur;

Y is selected from the group consisting of:

- (1) oxygen,
- (2) sulfur,
- (3) $-CR^9R^9$,
- (4) $S(O)_2$,
- (5) $C=O$,
- (6) $-C(R^9)OC_{1-6}alkyl$, and
- (7) $-NR^c$,

provided that if X is oxygen, sulfur or $-NR^c$, then Y is not oxygen, sulfur or $-NR^c$, further provided that if X is $C=O$, then Y is not $C=O$ or $S(O)_2$, and further provided that if X is $S(O)_2$, then Y is not $S(O)_2$, $C=O$, oxygen or sulfur;

Z is selected from the group consisting of:

- (1) oxygen,
- (2) sulfur,
- (3) $-CR^9R^9$,
- (4) $S(O)_2$,
- (5) $C=O$,
- (6) $-C(R^9)OC_{1-6}alkyl$, and
- (7) $-NR^c$;

A is selected from the group consisting of:

- (1) aryl,
- (2) heteroaryl,
- (3) C₃₋₆cycloalkyl, and
- (4) C₂₋₅cycloheteroalkyl,

5 wherein A is unsubstituted or substituted with one to five substituents selected from R^a;
B is selected from the group consisting of:

- (1) hydrogen,
- (2) aryl,
- (3) aryl-O-,
- 10 (4) aryl-C₁₋₁₀ alkyl-,
- (5) aryl-C₁₋₁₀ alkyl-O-,
- (6) C₃₋₆cycloalkyl,
- (7) C₃₋₆cycloalkyl-C₁₋₁₀alkyl-,
- (8) C₃₋₆cycloalkyl-C₁₋₁₀alkyl-O-,
- 15 (9) C₃₋₆cycloalkenyl,
- (10) C₃₋₆cycloalkenyl-C₁₋₁₀alkyl-,
- (11) C₃₋₆cycloalkenyl-C₁₋₁₀alkyl-O-,
- (12) C₂₋₅cycloheteroalkyl,
- (13) C₃₋₆cycloheteroalkyl-C₁₋₁₀alkyl-,
- 20 (14) C₃₋₆cycloheteroalkyl-C₁₋₁₀alkyl-O-,
- (15) heteroaryl,
- (16) heteroaryl-O-,
- (17) heteroaryl-C₁₋₁₀ alkyl-, and
- (18) heteroaryl-C₁₋₁₀ alkyl-O-,

25 wherein B is unsubstituted or substituted with one to five substituents selected from R^b;
R¹ and R² are each independently selected from:

- (1) a bond,
- (2) hydrogen,
- (3) halogen,
- 30 (4) -OR^k,
- (5) -CN,
- (6) -C₁₋₆alkyl,
- (7) -C₃₋₆cycloalkyl,
- (8) C₃₋₆cycloalkyl-C₁₋₃alkyl-,
- 35 (9) -C₂₋₆cycloheteroalkyl, and
- (10) C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-,

wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 , or R^1 and R^2 together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from

oxygen, sulfur and N-R_g, wherein each R^1 and R^2 is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 ;

R^3 is absent or selected from the group consisting of:

- (1) hydrogen,
- (2) halogen,
- (3) -OR^e,
- (4) -CN,
- (5) -C₁₋₆alkyl,
- (6) -C₃₋₆cycloalkyl, and
- (7) C₃₋₆cycloalkyl-C₁₋₃alkyl-,

wherein each alkyl and cycloalkyl is unsubstituted or substituted with one to three substituents selected from R^i ;

R^4 is selected from the group consisting of:

- (1) hydrogen,
- (2) halogen,
- (3) OR^e,
- (4) C₀₋₅alkylNR^cR^d,
- (5) C₁₋₆alkyl,
- (6) C₁₋₆alkyl-O-,
- (7) C₃₋₆cycloalkyl,
- (8) C₃₋₆cycloalkyl-O-,
- (9) C₃₋₆cycloalkyl-C₁₋₁₀alkyl-,
- (10) C₃₋₆cycloalkyl-C₁₋₁₀alkyl-O-,
- (11) C₂₋₅cycloheteroalkyl,
- (12) C₂₋₅cycloheteroalkyl-O-,
- (13) C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-,
- (14) C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-O-,
- (15) aryl,
- (16) aryl-O-,
- (17) aryl-C₁₋₁₀alkyl-,
- (18) heteroaryl,
- (19) heteroaryl-O-, and

(20) heteroaryl-C₁₋₁₀alkyl-,

wherein each alkyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or substituted with one to three substituents selected from R_j,

provided that when R⁴ is selected from the group consisting of:

- 5 (1) ORe,
- (2) C₀alkyl-NR^cR^d,
- (3) C₁₋₆alkyl-O-,
- (4) C₃₋₆cycloalkyl-O-,
- (5) C₃₋₆cycloalkyl-C₁₋₁₀alkyl-O-,
- 10 (6) C₂₋₅cycloheteroalkyl-O-,
- (7) C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-O-,
- (8) aryl-O-, and
- (9) heteroaryl-O-,

then X is selected from the group consisting of:

- 15 (1) -CR⁹R⁹,
- (2) C=O, and
- (3) -C(R⁹)OC₁₋₆alkyl;

R⁵ is absent or selected from the group consisting of:

- (1) hydrogen,
- 20 (2) -C₁₋₆alkyl, and
- (3) -C₃₋₆cycloalkyl,

wherein each alkyl and cycloalkyl is unsubstituted or substituted with one to three substituents selected from R_j;


R⁶ is absent or selected from the group consisting of:

- 25 (1) hydrogen,
- (2) -C₁₋₆alkyl, and
- (3) -C₃₋₆cycloalkyl,

wherein each alkyl and cycloalkyl is unsubstituted or substituted with one to three substituents selected from R_j;

30 R⁷ is selected from the group consisting of:

- (1) -CO₂R⁸,
- (2) -C₁₋₆alkyl-CO₂R⁸,
- (3) -C₁₋₆alkyl-CONHSO₂R^m,
- (4) -C₁₋₆alkyl-SO₂NHCOR^m,
- 35 (5) -C₁₋₆alkyl-tetrazolyl, and
- (6) a cycloheteroalkyl selected from the group consisting of:



R^8 is selected from the group consisting of:

- (1) hydrogen,
- (2) $-C_{1-6}$ alkyl,
- (3) $-C_{3-6}$ cycloalkyl, and
- (4) aryl- C_{1-6} alkyl,

wherein each alkyl, cycloalkyl and aryl is unsubstituted or substituted with one to three substituents selected from R^j ;

R^9 is selected from the group consisting of:

- (1) hydrogen,
- (2) $-C_{1-6}$ alkyl,
- (3) halogen,
- (4) $-(CH_2)_{0-5}OH$,
- (5) $-(CH_2)_{0-5}OC_{1-6}$ alkyl,
- (6) $-(CH_2)_{0-5}NR^cR^d$,
- (7) $-C_{3-6}$ cycloalkyl, and
- (8) aryl- C_{1-6} alkyl,

wherein CH_2 is unsubstituted or substituted with one or two substituents selected from R^j , and wherein alkyl, cycloalkyl and aryl is unsubstituted or substituted with one to three substituents selected from R^j ;

R^a is selected from the group consisting of:

- (1) $-C_{1-6}$ alkyl,
- (2) halogen,
- (3) $-C_{0-6}$ alkyl- OR^e ,
- (4) $-C_{0-6}$ alkyl- $NR^cS(O)_nR^e$,

- (5) -C₀₋₆alkyl-S(O)_nRe,
 (6) -C₀₋₆alkyl-S(O)_nNR^cR^d,
 (7) -C₀₋₆alkyl-NR^cR^d,
 (8) -C₀₋₆alkyl-C(O)Re,
 5 (9) -C₀₋₆alkyl-OC(O)Re,
 (10) -C₀₋₆alkyl-CO₂Re,
 (11) -C₀₋₆alkyl-CN,
 (12) -C₀₋₆alkyl-C(O)NR^cR^d,
 (13) -C₀₋₆alkyl-NR^cC(O)Re,
 10 (14) -C₀₋₆alkyl-NR^cC(O)ORE,
 (15) -C₀₋₆alkyl-NR^cC(O)NR^cR^d,
 (16) -CF₃,
 (17) -OCF₃,
 (18) -OCHF₂,
 15 (19) -C₀₋₆alkyl-aryl,
 (20) -C₀₋₆alkyl-heteroaryl,
 (21) -C₀₋₆alkyl -C₃₋₁₀cycloalkyl,
 (22) -C₀₋₆alkyl -C₃₋₁₀cycloalkenyl, and
 (23) -C₀₋₆alkyl -C₂₋₁₀cycloheteroalkyl,
- 20 wherein each alkyl, cycloalkyl, cycloalkenyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or substituted with one to five substituents independently selected from: -C₁₋₆alkyl, halogen, OH, -O-C₁₋₆alkyl, -S(O)₂-C₁₋₄alkyl, -CN, -OCHF₂, -OCF₃, -CF₃, and -C₀₋₆alkyl-NR^cR^d;
- R^b is selected from the group consisting of:
- (1) -C₁₋₁₀alkyl,
 25 (2) -C₂₋₁₀alkenyl,
 (3) -CF₃,
 (4) halogen,
 (5) -CN,
 (6) -OH,
 30 (7) -OC₁₋₁₀alkyl,
 (8) -OC₂₋₁₀alkenyl,
 (9) -O(CH₂)_pOC₁₋₁₀alkyl,
 (10) -O(CH₂)_pC₃₋₆cycloalkyl,
 (11) -O(CH₂)_pC₃₋₆ cycloalkyl-C₁₋₁₀alkyl,
 35 (12) -O(CH₂)_pC₂₋₅cycloheteroalkyl,
 (13) -O(CH₂)_pC₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl,

- (14) -O-aryl,
 (15) -O-heteroaryl,
 (16) -O-aryl-C₁₋₁₀alkyl,
 (17) -O-heteroaryl-C₁₋₁₀alkyl,
 5 (18) -O(CH₂)_pNR^cS(O)_mRe,
 (19) -O(CH₂)_pS(O)_mRe,
 (20) -O(CH₂)_pS(O)_mNR^cR^d,
 (21) -O(CH₂)_pNR^cR^d,
 (22) -C(O)Re,
 10 (23) -OC(O)Re,
 (24) -CO₂Re,
 (25) -C(O)NR^cR^d,
 (26) -NR^cC(O)Re,
 (27) -NR^cC(O)ORE,
 15 (28) -NR^cC(O)NR^cR^d,
 (29) -O(CH₂)_pO-C₃₋₆cycloalkyl,
 (30) -O(CH₂)_pO-C₂₋₅cycloheteroalkyl,
 (31) -OCF₃,
 (32) -OCHF₂,
 20 (33) -(CH₂)_pC₃₋₆cycloalkyl,
 (34) -(CH₂)_pC₂₋₅cycloheteroalkyl,
 (35) aryl,
 (36) heteroaryl,
 (37) aryl-C₁₋₁₀alkyl-, and
 25 (38) heteroaryl-C₁₋₁₀alkyl-,

wherein each CH, CH₂, alkyl, alkenyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or substituted with one to five substituents independently selected from: -C₁₋₆alkyl, halogen, OH, -O-C₁₋₆alkyl and -CF₃;

R^c and R^d are each independently selected from the group consisting of:

- 30 (1) hydrogen,
 (2) C₁₋₁₀alkyl,
 (3) C₂₋₁₀alkenyl,
 (4) C₃₋₆cycloalkyl,
 (5) C₃₋₆ cycloalkyl-C₁₋₁₀alkyl-,
 35 (6) C₂₋₅cycloheteroalkyl,
 (7) C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-,

- (8) aryl,
- (9) heteroaryl,
- (10) aryl-C₁₋₁₀alkyl-, and
- (11) heteroaryl-C₁₋₁₀alkyl-,

5 wherein each alkyl, alkenyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or substituted with one to three substituents independently selected from R^f, or R^c and R^d together with the atom(s) to which they are attached form a C₂₋₁₀ cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R^g, wherein each R^c and R^d is unsubstituted or substituted with one to three substituents independently
10 selected from R^f,

each R^e is independently selected from the group consisting of:

- (1) hydrogen,
- (2) -C₁₋₁₀alkyl,
- (3) -C₂₋₁₀ alkenyl,
- 15 (4) -C₃₋₆ cycloalkyl,
- (5) C₃₋₆ cycloalkyl-C₁₋₁₀alkyl-,
- (6) -C₂₋₅cycloheteroalkyl,
- (7) C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-,
- (8) aryl,
- 20 (9) aryl-C₁₋₁₀alkyl-,
- (10) heteroaryl, and
- (11) heteroaryl-C₁₋₁₀alkyl-,

wherein each alkyl, alkenyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or substituted with one to three substituents selected from R^h;

25 each R^f is selected from the group consisting of:

- (1) halogen,
- (2) C₁₋₁₀alkyl,
- (3) -OH,
- (4) -O-C₁₋₄alkyl,
- 30 (5) -S(O)_m-C₁₋₄alkyl,
- (6) -CN,
- (7) -CF₃,
- (8) -OCHF₂, and
- (9) -OCF₃,

35 wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from: -OH, halogen, C₁₋₆alkyl, cyano and S(O)₂C₁₋₆alkyl;

each R^g is independently selected from the group consisting of:

- (1) hydrogen,
- (2) $-C(O)R^e$, and
- (3) $-C_{1-10}alkyl$,

5 wherein alkyl is unsubstituted or substituted with one to five halogens;

each R^h is independently selected from the group consisting of:

- (1) halogen,
- (2) $C_{1-10}alkyl$,
- (3) $-OH$,
- 10 (4) $-O-C_{1-4}alkyl$,
- (5) $-S(O)_m-C_{1-4}alkyl$,
- (6) $-CN$,
- (7) $-CF_3$,
- (8) $-OCHF_2$, and
- 15 (9) $-OCF_3$,

wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from: $-OH$, halogen, $C_{1-6}alkyl$, cyano and $S(O)_2C_{1-6}alkyl$;

each R^i is independently selected from the group consisting of:

- (1) $-C_{1-6}alkyl$,
- 20 (2) $-OR^e$,
- (3) $-NR^cS(O)_mR^e$,
- (4) halogen,
- (5) $-S(O)_mR^e$,
- (6) $-S(O)_mNR^cR^d$,
- 25 (7) $-NR^cR^d$,
- (8) $-C(O)R^e$,
- (9) $-OC(O)R^e$,
- (10) $-CO_2R^e$,
- (11) $-CN$,
- 30 (12) $-C(O)NR^cR^d$,
- (13) $-NR^cC(O)R^e$,
- (14) $-NR^cC(O)OR^e$,
- (15) $-NR^cC(O)NR^cR^d$,
- (16) $-CF_3$,
- 35 (17) $-OCF_3$,
- (18) $-OCHF_2$,

(19) $-C_{3-6}$ cycloalkyl, and

(20) $-C_{2-5}$ cycloheteroalkyl;

each R^j is independently selected from the group consisting of:

(1) $-C_{1-6}$ alkyl,

(2) $-OR^e$,

(3) $-NR^cS(O)_mR^e$,

(4) halogen,

(5) $-S(O)_mR^e$,

(6) $-S(O)_mNR^cR^d$,

(7) $-NR^cR^d$,

(8) $-C(O)R^e$,

(9) $-OC(O)R^e$,

(10) $-CO_2R^e$,

(11) $-CN$,

(12) $-C(O)NR^cR^d$,

(13) $-NR^cC(O)R^e$,

(14) $-NR^cC(O)OR^e$,

(15) $-NR^cC(O)NR^cR^d$,

(16) $-CF_3$,

(17) $-OCF_3$,

(18) $-OCHF_2$,

(19) $-C_{3-6}$ cycloalkyl, and

(20) $-C_{2-5}$ cycloheteroalkyl;

each R^k is independently selected from the group consisting of:

(1) hydrogen,

(2) $-C_{1-6}$ alkyl,

(3) $-C_{1-6}$ alkyl- SO_2C_{1-6} alkyl,

(4) $-CF_3$, and

(5) $-CHF_2$,

wherein each alkyl is unsubstituted or substituted with one to three substituents independently selected from: $-OH$, $-OC_{1-6}$ alkyl, halogen, cyano, and $-S(O)_2C_{1-6}$ alkyl;

each R^L is independently selected from the group consisting of:

(1) $-CO_2C_{1-6}$ alkyl,

(2) $-C_{1-10}$ alkyl,

(3) $-C_{2-10}$ alkenyl,

(4) $-C_{2-10}$ alkynyl,

- (5) -C₃₋₆cycloalkyl,
- (6) -C₂₋₆cycloheteroalkyl,
- (7) aryl, and
- (8) heteroaryl,

5 wherein each alkyl, alkenyl, alkynyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or substituted with one to four substituents independently selected from C₁₋₆alkyl, halogen, and -OC₁₋₆alkyl;

each R^m is independently selected from the group consisting of:

- (1) -C₁₋₁₀alkyl,
- 10 (2) -C₂₋₁₀ alkenyl,
- (3) -C₃₋₆ cycloalkyl,
- (4) C₃₋₆ cycloalkyl-C₁₋₁₀alkyl-,
- (5) -C₂₋₅cycloheteroalkyl,
- (6) C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-,
- 15 (7) aryl,
- (8) heteroaryl,
- (9) aryl-C₁₋₁₀alkyl-, and
- (10) heteroaryl-C₁₋₁₀alkyl-;

each n is independently selected from: 0, 1 or 2;

20 each m is independently selected from: 0, 1 or 2;

each p is independently selected from: 0, 1, 2, 3, 4, 5 or 6; and

each r is independently selected from: 0, 1, 2 or 3.

The invention has numerous embodiments, which are summarized below. The invention includes the compounds as shown, and also includes individual diastereoisomers, enantiomers, and 25 epimers of the compounds, and mixtures of diastereoisomers and/or enantiomers thereof including racemic mixtures.

In one embodiment of the present invention, "a" is a single bond. In another embodiment of the present invention, "a" is a single bond; and R⁵ and R⁶ are present.

In another embodiment of the present invention, "a" is a double bond and R⁵ and R⁶ are 30 absent. In another embodiment, "a" is a double bond, R⁵ and R⁶ are absent and Q is selected from the group consisting of: C-R⁴, -C-OC₁₋₆alkyl, and N.

In another embodiment of the present invention, “a” is a single bond, and Q is selected from the group consisting of: oxygen, sulfur, $-CR^4R^6$, $S(O)_2$, $C=O$, $-C(R^9)OC_{1-6}alkyl$, and $-NR^c$. In another embodiment of the present invention, “a” is a single bond, R^5 and R^6 are present, and Q is selected from the group consisting of: oxygen, sulfur, $-CR^4R^6$, $S(O)_2$, $C=O$, $-C(R^9)OC_{1-6}alkyl$, and $-NR^c$.

In another embodiment of the present invention, T is selected from the group consisting of: CH, N and N-oxide. In a class of this embodiment, T is selected from the group consisting of: CH and N. In another class of this embodiment, T is CH. In another class of this embodiment, T is N or N-oxide. In another class of this embodiment, T is N. In another class of this embodiment, T is N-oxide.

In another embodiment of the present invention, U is selected from the group consisting of: CR^1 , N and N-oxide. In a class of this embodiment, U is selected from the group consisting of: CR^1 and N. In another class of this embodiment, U is CR^1 . In another class of this embodiment, U is N or N-oxide. In another class of this embodiment, U is N. In another class of this embodiment, U is N-oxide.

In another embodiment of the present invention, V is selected from the group consisting of: CR^2 , N and N-oxide. In a class of this embodiment, V is selected from the group consisting of: CR^2 and N. In another class of this embodiment, V is CR^2 . In another class of this embodiment, V is N or N-oxide. In another class of this embodiment, V is N. In another class of this embodiment, V is N-oxide.

In another embodiment of the present invention, W is selected from the group consisting of: CH, N and N-oxide, provided that no more than two of T, U, V and W are selected from N and N-oxide, further provided that if both T and W are N or N-oxide, then R^3 is absent, and further provided that both U and V are not N or N-oxide. In a class of this embodiment, W is selected from the group consisting of: CH and N, provided that no more than two of T, U, V and W are selected from N and N-oxide, further provided that if both T and W are N or N-oxide, then R^3 is absent, and further provided that both U and V are not N or N-oxide. In another class of this embodiment, W is CH, provided that no more than two of T, U, V and W are selected from N and N-oxide, further provided that if both T and W are N or N-oxide, then R^3 is absent, and further provided that both U and V are not N or N-oxide. In another class of this embodiment, W is N or N-oxide, provided that no more than two of T, U, V and W are selected from N and N-oxide, further provided that if both T and W are N or N-oxide, then R^3 is absent, and further provided that both U and V are not N or N-oxide. In another class of this embodiment, W is N, provided that no more than two of T, U, V and W are selected from N and N-oxide, further provided that if both T and W are N or N-oxide, then R^3 is absent, and further provided that both U and V are not N or N-oxide.

In another embodiment of the present invention, W is selected from the group consisting of: CH, N and N-oxide. In a class of this embodiment, W is selected from the group consisting of: CH and N. In another class of this embodiment, W is CH. In another class of this embodiment, W is N or N-oxide. In another class of this embodiment, W is N. In another class of this embodiment, W is N-oxide.

In another embodiment of the present invention, T is CH, U is CR¹, V is CR², and W is CH. In a class of this embodiment, T is CH, U is CR¹, V is CH, and W is CH. In another class of this embodiment, T is CH, U is CH, V is CR², and W is CH.

In another embodiment of the present invention, T is N or N-oxide, U is CR¹, V is CR², and W is CH. In a class of this embodiment, T is N, U is CR¹, V is CR², and W is CH. In another embodiment of the present invention, T is CH, U is N or N-oxide, and V is CR², and W is CH. In a class of this embodiment, T is CH, U is N, V is CR², and W is CH. In another embodiment of the present invention, T is CH, U is CR¹, V is N or N-oxide, and W is CH. In a class of this embodiment, T is CH, U is CR¹, and V is N or N-oxide, and W is CH. In another embodiment of the present invention, T is CH, U is CR¹, V is CR², and W is CH, N or N-oxide. In another embodiment of the present invention, T is CH, U is CR¹, V is CR², and W is N or N-oxide. In a class of this embodiment, T is CH, U is CR¹, V is CR², and W is N.

In another embodiment of the present invention, T is N or N-oxide, U is N or N-oxide, V is CR², and W is CH. In a class of this embodiment, T is N, U is N, V is CR², and W is CH. In another embodiment of the present invention, T is N or N-oxide, U is CR¹, V is N or N-oxide, and W is CH. In a class of this embodiment, T is N, U is CR¹, V is N, and W is CH. In another embodiment of the present invention, T is N or N-oxide, U is CR¹, V is CR², and W is N or N-oxide. In a class of this embodiment, T is N, U is CR¹, V is CR², and W is N.

In another embodiment of the present invention, T is N or N-oxide, U is CR¹, V is CR², and W is N or N-oxide; and R³ is absent. In a class of this embodiment, T is N, U is CR¹, V is CR², and W is N; and R³ is absent. In another embodiment of the present invention, T is CH, U is N or N-oxide, V is CR², and W is N or N-oxide. In a class of this embodiment, T is CH, U is N, V is CR², and W is N. In another embodiment of the present invention, T is CH, U is CR¹, V is N or N-oxide, and W is N or N-oxide. In a class of this embodiment, T is CH, U is CR¹, V is N, and W is N. In another embodiment of the present invention, T is CH; U is CR¹; V is CR²; and W is CH, N or N-oxide.

In another embodiment of the present invention, Q is selected from the group consisting of: oxygen, sulfur, -CR⁴R⁶, C=O, -C(R⁹)OC₁₋₆alkyl, and -NRC. In another embodiment of the present invention, Q is selected from the group consisting of: oxygen, sulfur, -CR⁴R⁶, -C(R⁹)OC₁₋₆alkyl, and -NRC. In another embodiment of the present invention, Q is selected from the group consisting of: oxygen, sulfur, -CR⁴R⁶, and -NRC. In another embodiment of the present invention, Q is

selected from the group consisting of: oxygen, $-\text{CR}^4\text{R}^6$, and $-\text{NR}^c$. In another embodiment of the present invention, Q is selected from the group consisting of: oxygen, and $-\text{CR}^4\text{R}^6$. In a class of this embodiment, Q is selected from the group consisting of: oxygen, and $-\text{CH}_2$.

In another embodiment of the present invention, Q is oxygen. In another embodiment of the present invention, Q is $-\text{CR}^4\text{R}^6$. In a class of this embodiment, Q is $-\text{CH}_2$. In another class of this embodiment, Q is $-\text{CH}$.

In another embodiment of the present invention, X is selected from the group consisting of: oxygen, $-\text{CR}^9\text{R}^9$, $\text{S}(\text{O})_2$, $\text{C}=\text{O}$, $-\text{C}(\text{R}^9)\text{OC}_{1-6}\text{alkyl}$, and $-\text{NR}^c$, provided that if Q is oxygen, sulfur or $-\text{NR}^c$, then X is not oxygen or $-\text{NR}^c$, further provided that if Q is $\text{C}=\text{O}$, then X is not $\text{C}=\text{O}$ or $\text{S}(\text{O})_2$, and further provided that if Q is $\text{S}(\text{O})_2$, then X is not $\text{S}(\text{O})_2$, $\text{C}=\text{O}$, or oxygen. In another embodiment of the present invention, X is selected from the group consisting of: oxygen, $-\text{CR}^9\text{R}^9$, $\text{S}(\text{O})_2$, $\text{C}=\text{O}$, and $-\text{NR}^c$, provided that if Q is oxygen, sulfur or $-\text{NR}^c$, then X is not oxygen or $-\text{NR}^c$, further provided that if Q is $\text{C}=\text{O}$, then X is not $\text{C}=\text{O}$ or $\text{S}(\text{O})_2$, and further provided that if Q is $\text{S}(\text{O})_2$, then X is not $\text{S}(\text{O})_2$, $\text{C}=\text{O}$, or oxygen.

In another embodiment of the present invention, X is selected from the group consisting of: oxygen, $-\text{CR}^9\text{R}^9$, $\text{C}=\text{O}$, and $-\text{NR}^c$, provided that if Q is oxygen, sulfur or $-\text{NR}^c$, then X is not oxygen or $-\text{NR}^c$, further provided that if Q is $\text{C}=\text{O}$, then X is not $\text{C}=\text{O}$, and further provided that if Q is $\text{S}(\text{O})_2$, then X is not $\text{C}=\text{O}$, or oxygen. In another embodiment of the present invention, X is selected from the group consisting of: oxygen, $-\text{CR}^9\text{R}^9$, and $-\text{NR}^c$, provided that if Q is oxygen, sulfur or $-\text{NR}^c$, then X is not oxygen or $-\text{NR}^c$, and further provided that if Q is $\text{S}(\text{O})_2$, then X is not oxygen. In another embodiment of the present invention, X is oxygen. In another embodiment of the present invention, X is $-\text{NR}^c$. In another embodiment of the present invention, X is $-\text{CR}^9\text{R}^9$. In a class of this embodiment, X is $-\text{CH}_2$.

In another embodiment of the present invention, X is selected from the group consisting of: $-\text{CR}^9$, $-\text{C-OC}_{1-6}\text{alkyl}$, and $-\text{N}$, provided that if Q is oxygen, sulfur or $-\text{NR}^c$, then X is not $-\text{N}$. In another embodiment of the present invention, X is selected from the group consisting of: $-\text{CR}^9$, and $-\text{N}$, provided that if Q is oxygen, sulfur or $-\text{NR}^c$, then X is not $-\text{N}$. In another embodiment of the present invention, X is $-\text{N}$. In another embodiment of the present invention, X is $-\text{CR}^9$. In a class of this embodiment, X is $-\text{CH}$.

In another embodiment of the present invention, Y is selected from the group consisting of: oxygen, $-\text{CR}^9\text{R}^9$, $\text{S}(\text{O})_2$, $\text{C}=\text{O}$, $-\text{C}(\text{R}^9)\text{OC}_{1-6}\text{alkyl}$, and $-\text{NR}^c$, that if X is oxygen, sulfur or $-\text{NR}^c$, then Y is not oxygen or $-\text{NR}^c$, further provided that if X is $\text{C}=\text{O}$, then Y is not $\text{C}=\text{O}$ or $\text{S}(\text{O})_2$, and further provided that if X is $\text{S}(\text{O})_2$, then Y is not $\text{S}(\text{O})_2$, $\text{C}=\text{O}$, or oxygen. In another embodiment of the present invention, Y is selected from the group consisting of: oxygen, $-\text{CR}^9\text{R}^9$, $\text{S}(\text{O})_2$, $\text{C}=\text{O}$, and $-\text{NR}^c$, provided that if X is oxygen, sulfur or $-\text{NR}^c$, then Y is not oxygen, or $-\text{NR}^c$, further provided that if X is $\text{C}=\text{O}$, then Y is not $\text{C}=\text{O}$ or $\text{S}(\text{O})_2$, and further provided that if X is $\text{S}(\text{O})_2$, then

Y is not $S(O)_2$, $C=O$, or oxygen. In another embodiment of the present invention, Y is selected from the group consisting of: oxygen, $-CR^9R^9$, $C=O$, and $-NRC$, provided that if X is oxygen, sulfur or $-NRC$, then Y is not oxygen, or $-NRC$, further provided that if X is $C=O$, then Y is not $C=O$, and further provided that if X is $S(O)_2$, then Y is not $C=O$, or oxygen.

5 In another embodiment of the present invention, Y is selected from the group consisting of: oxygen, $-CR^9R^9$, and $-NRC$, provided that if X is oxygen, sulfur or $-NRC$, then Y is not oxygen, or $-NRC$, and further provided that if X is $S(O)_2$, then Y is not oxygen. In another embodiment of the present invention, Y is oxygen. In another embodiment of the present invention, Y is $-NRC$. In another embodiment of the present invention, Y is $-CR^9R^9$. In a class of this embodiment, Y is $-CH_2$.

10 In another embodiment of the present invention, Y is selected from the group consisting of: $-CR^9$, $-C-OC_{1-6}alkyl$, and $-N$, provided that if X is oxygen, sulfur or $-NRC$, then Y is not $-N$. In another embodiment of the present invention, Y is selected from the group consisting of: $-CR^9$, and $-N$, provided that if X is oxygen, sulfur or $-NRC$, then Y is not $-N$. In another embodiment of the present invention, Y is $-N$. In another embodiment of the present invention, Y is $-CR^9R^9$. In a class of this embodiment, Y is $-CH_2$.

15 In another embodiment of the present invention, Z is selected from the group consisting of: oxygen, $-CR^9R^9$, $S(O)_2$, and $-NRC$. In another embodiment of the present invention, Z is selected from the group consisting of: oxygen, $-CR^9R^9$, and $-NRC$. In another embodiment of the present invention, Z is selected from the group consisting of: oxygen, and $-CR^9R^9$. In another embodiment of the present invention, Z is selected from the group consisting of: oxygen, and $-NRC$. In another embodiment of the present invention, Z is $-NRC$. In another embodiment of the present invention, Z is $S(O)_2$. In another embodiment of the present invention, Z is $-CR^9R^9$. In a class of this embodiment, Z is CH_2 . In another embodiment of the present invention, Z is oxygen.

20 In another embodiment of the present invention, Z is selected from the group consisting of: oxygen, $-CR^9R^9$, and $-NRC$. In another embodiment of the present invention, Z is selected from the group consisting of: oxygen, and $-CR^9R^9$. In another embodiment of the present invention, Z is selected from the group consisting of: oxygen, and $-NRC$. In another embodiment of the present invention, Z is $-NRC$. In another embodiment of the present invention, Z is $S(O)_2$. In another embodiment of the present invention, Z is $-CR^9R^9$. In a class of this embodiment, Z is CH_2 . In another embodiment of the present invention, Z is oxygen.

25 In another embodiment of the present invention, A is selected from the group consisting of: aryl, heteroaryl, and $C_{2-5}cycloheteroalkyl$, wherein each aryl, heteroaryl and cycloheteroalkyl is unsubstituted or substituted with one to five substituents selected from R^a . In a class of this embodiment, A is selected from the group consisting of: phenyl, pyridyl, and azetidine, wherein each phenyl, pyridyl and azetidine is unsubstituted or substituted with one to four substituents selected from R^a . In another embodiment of the present invention, A is selected from the group consisting of: aryl, and heteroaryl, wherein each aryl and heteroaryl is unsubstituted or substituted with one to five substituents selected from R^a . In a class of this embodiment, A is selected from the group consisting of: phenyl, and pyridyl, wherein phenyl, and pyridyl is unsubstituted or substituted with one to four substituents selected from R^a . In another embodiment of the present invention, A is selected from the group consisting of: aryl, wherein aryl is unsubstituted or substituted with one to five substituents selected from R^a . In a class of this embodiment, A is phenyl, wherein phenyl is

30

35

unsubstituted or substituted with one to five substituents selected from R^a. In another embodiment of the present invention, A is selected from the group consisting of: heteroaryl, wherein heteroaryl is unsubstituted or substituted with one to five substituents selected from R^a. In a class of this embodiment, A is pyridyl, wherein pyridyl is unsubstituted or substituted with one to four substituents selected from R^a. In another embodiment of the present invention, A is selected from the group consisting of: C₂-5cycloheteroalkyl, wherein each cycloheteroalkyl is unsubstituted or substituted with one to five substituents selected from R^a. In a class of this embodiment, A is azetidine, wherein azetidine is unsubstituted or substituted with one to four substituents selected from R^a.

In another embodiment of the present invention, B is selected from the group consisting of: aryl, aryl-O-, aryl-C₁₋₁₀ alkyl-, aryl-C₁₋₁₀ alkyl-O-, C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₁₀alkyl-, C₃₋₆cycloalkyl-C₁₋₁₀alkyl-O-, C₃₋₆cycloalkenyl, C₃₋₆cycloalkenyl-C₁₋₁₀alkyl-, C₃₋₆cycloalkenyl-C₁₋₁₀alkyl-O-, C₂₋₅cycloheteroalkyl, C₃₋₆cycloheteroalkyl-C₁₋₁₀alkyl-, C₃₋₆cycloheteroalkyl-C₁₋₁₀alkyl-O-, heteroaryl, heteroaryl-O-, heteroaryl-C₁₋₁₀ alkyl-, and heteroaryl-C₁₋₁₀ alkyl-O-, wherein B is unsubstituted or substituted with one to five substituents selected from R^b.

In another embodiment of the present invention, B is selected from the group consisting of: aryl, aryl-C₁₋₁₀ alkyl-, C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₁₀alkyl-, C₃₋₆cycloalkenyl, C₃₋₆cycloalkenyl-C₁₋₁₀alkyl-, C₂₋₅cycloheteroalkyl, C₃₋₆cycloheteroalkyl-C₁₋₁₀alkyl, heteroaryl, and heteroaryl-C₁₋₁₀ alkyl-, wherein B is unsubstituted or substituted with one to five substituents selected from R^b. In another embodiment of the present invention, B is selected from the group consisting of: aryl, aryl-C₁₋₁₀ alkyl-, C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₁₀alkyl-, C₂₋₅cycloheteroalkyl, C₃₋₆cycloheteroalkyl-C₁₋₁₀alkyl, heteroaryl, and heteroaryl-C₁₋₁₀ alkyl-, wherein B is unsubstituted or substituted with one to five substituents selected from R^b. In another embodiment of the present invention, B is selected from the group consisting of: aryl, aryl-C₁₋₁₀ alkyl-, C₃₋₆cycloalkyl, C₂₋₅cycloheteroalkyl, heteroaryl, and heteroaryl-C₁₋₁₀ alkyl-, wherein B is unsubstituted or substituted with one to five substituents selected from R^b.

In another embodiment of the present invention, B is selected from the group consisting of: aryl, aryl-C₁₋₁₀ alkyl-, and heteroaryl, wherein each alkyl, aryl and heteroaryl is unsubstituted or substituted with one to five substituents selected from R^b. In a class of this embodiment, B is selected from the group consisting of: phenyl, -CH₂-phenyl, and pyridyl, wherein each phenyl and pyridyl is unsubstituted or substituted with one to five substituents selected from R^b.

In another embodiment of the present invention, B is selected from the group consisting of: aryl, wherein each aryl is unsubstituted or substituted with one to five substituents selected from R^b. In a class of this embodiment, B is phenyl, wherein each phenyl is unsubstituted or substituted with one to five substituents selected from R^b. In another embodiment of the present

invention, B is selected from the group consisting of: aryl-C₁₋₁₀ alkyl-, wherein each alkyl and aryl is unsubstituted or substituted with one to five substituents selected from R^b. In a class of this embodiment, B is selected from the group consisting of: -CH₂-phenyl, wherein each phenyl is unsubstituted or substituted with one to five substituents selected from R^b. In another embodiment of the present invention, B is selected from the group consisting of: heteroaryl, wherein each heteroaryl is unsubstituted or substituted with one to five substituents selected from R^b. In a class of this embodiment, B is selected from the group consisting of: pyridyl, wherein each pyridyl is unsubstituted or substituted with one to five substituents selected from R^b.

In another embodiment of the present invention, R¹ and R² are each independently selected from: hydrogen, halogen, -OR^k, -CN, -C₁₋₆alkyl, -C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₃alkyl-, -C₂₋₆cycloheteroalkyl, and C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷, or R¹ and R² together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R^g, wherein each R¹ and R² is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷.

In another embodiment of the present invention, R¹ and R² are each independently selected from: hydrogen, halogen, -OR^k, -CN, -C₁₋₆alkyl, -C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₃alkyl-, -C₂₋₆cycloheteroalkyl, and C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷.

In another embodiment of the present invention, R¹ and R² are each independently selected from: hydrogen, halogen, -OR^k, -CN, and -C₁₋₆alkyl, wherein each alkyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷, or R¹ and R² together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R^g, wherein each R¹ and R² is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷.

In another embodiment of the present invention, R¹ and R² are each independently selected from: hydrogen, halogen, -OR^k, -CN, and -C₁₋₆alkyl, wherein each alkyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷.

In another embodiment of the present invention, R¹ and R² are each independently selected from: hydrogen and -C₁₋₆alkyl, wherein each alkyl is unsubstituted or substituted with one to three

substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 , or R^1 and R^2 together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R_g, wherein each R^1 and R^2 is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 .

In another embodiment of the present invention, R^1 and R^2 are each independently selected from: hydrogen, and -C₁₋₆alkyl, wherein each alkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 . In another embodiment of the present invention, R^1 and R^2 are each independently selected from: hydrogen, and ethyl, wherein ethyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 , or R^1 and R^2 together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R_g, wherein each R^1 and R^2 is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 .

In another embodiment of the present invention, R^1 and R^2 are each independently selected from: hydrogen, and ethyl, wherein ethyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 .

In another class of this embodiment, R^1 is selected from: a bond, hydrogen, halogen, -OR^k, -CN, -C₁₋₆alkyl, -C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₃alkyl-, -C₂₋₆cycloheteroalkyl, and C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 , or R^1 and R^2 together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R_g, wherein each R^1 and R^2 is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 .

In another embodiment of the present invention, R^1 is selected from: hydrogen, halogen, -OR^k, -CN, -C₁₋₆alkyl, -C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₃alkyl-, -C₂₋₆cycloheteroalkyl, and C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 , or R^1 and R^2 together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R_g, wherein each R^1 and R^2 is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 .

In another embodiment of the present invention, R^1 is $-\text{CH}(\text{cyclopropyl})-\text{CH}(\text{CH}_3)-\text{CO}_2\text{H}$. In another embodiment of the present invention, R^1 is $-\text{CH}(\text{cyclopropyl})-\text{CH}(\text{CH}_3)-\text{CO}_2\text{H}$, and R^2 is hydrogen. In another embodiment of the present invention, R^2 is $-\text{CH}(\text{cyclopropyl})-\text{CH}(\text{CH}_3)-\text{CO}_2\text{H}$. In another embodiment of the present invention, R^2 is $-\text{CH}(\text{cyclopropyl})-\text{CH}(\text{CH}_3)-\text{CO}_2\text{H}$, and R^1 is hydrogen.

In another embodiment of the present invention, R^1 is $-(\text{CH}_2)_2-\text{CO}_2\text{H}$. In another embodiment of the present invention, R^1 is $-(\text{CH}_2)_2-\text{CO}_2\text{H}$, and R^2 is hydrogen. In another embodiment of the present invention, R^2 is $-(\text{CH}_2)_2-\text{CO}_2\text{H}$. In another embodiment of the present invention, R^2 is $-(\text{CH}_2)_2-\text{CO}_2\text{H}$, and R^1 is hydrogen.

In another embodiment of the present invention, R^1 is selected from: hydrogen, halogen, $-\text{OR}^k$, $-\text{CN}$, $-\text{C}_{1-6}\text{alkyl}$, $-\text{C}_{3-6}\text{cycloalkyl}$, $\text{C}_{3-6}\text{cycloalkyl}-\text{C}_{1-3}\text{alkyl}$ -, $-\text{C}_{2-6}\text{cycloheteroalkyl}$, and $\text{C}_{2-6}\text{cycloheteroalkyl}-\text{C}_{1-3}\text{alkyl}$ -, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 . In another embodiment of the present invention, R^1 is selected from: hydrogen, halogen, $-\text{OR}^k$, and $-\text{CN}$.

In another embodiment of the present invention, R^1 is selected from: $-\text{C}_{1-6}\text{alkyl}$, $-\text{C}_{3-6}\text{cycloalkyl}$, $\text{C}_{3-6}\text{cycloalkyl}-\text{C}_{1-3}\text{alkyl}$ -, $-\text{C}_{2-6}\text{cycloheteroalkyl}$, and $\text{C}_{2-6}\text{cycloheteroalkyl}-\text{C}_{1-3}\text{alkyl}$ -, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein R^1 is substituted with R^7 , or R^1 and R^2 together with the atom(s) to which they are attached form a $\text{C}_{3-6}\text{cycloalkyl}$ ring or a $\text{C}_{2-5}\text{cycloheteroalkyl}$ ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and $\text{N}-\text{Rg}$, wherein each R^1 and R^2 is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 .

In another embodiment of the present invention, R^1 is selected from: $-\text{C}_{1-6}\text{alkyl}$, $-\text{C}_{3-6}\text{cycloalkyl}$, $\text{C}_{3-6}\text{cycloalkyl}-\text{C}_{1-3}\text{alkyl}$ -, $-\text{C}_{2-6}\text{cycloheteroalkyl}$, and $\text{C}_{2-6}\text{cycloheteroalkyl}-\text{C}_{1-3}\text{alkyl}$ -, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein R^1 is substituted with R^7 . In another embodiment of the present invention, R^1 is selected from: $-\text{C}_{1-6}\text{alkyl}$, $-\text{C}_{3-6}\text{cycloalkyl}$, $\text{C}_{3-6}\text{cycloalkyl}-\text{C}_{1-3}\text{alkyl}$ -, $-\text{C}_{2-6}\text{cycloheteroalkyl}$, and $\text{C}_{2-6}\text{cycloheteroalkyl}-\text{C}_{1-3}\text{alkyl}$ -, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein R^1 is substituted with R^7 ; and R^2 is hydrogen.

In another embodiment of the present invention, R^1 is selected from: hydrogen, halogen, $-\text{OR}^k$, $-\text{CN}$, and $-\text{C}_{1-6}\text{alkyl}$, wherein each alkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 , or R^1 and R^2 together with the atom(s) to which they are attached form a $\text{C}_{3-6}\text{cycloalkyl}$ ring or a $\text{C}_{2-5}\text{cycloheteroalkyl}$ ring containing 0-2 additional heteroatoms independently selected from oxygen,

sulfur and N-Rg, wherein each R¹ and R² is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷. In another embodiment of the present invention, R¹ is selected from: hydrogen, halogen, -OR^k, -CN, and -C₁₋₆alkyl, wherein each alkyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷.

In another embodiment of the present invention, R¹ is selected from: hydrogen, and -C₁₋₆alkyl, wherein each alkyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷, or R¹ and R² together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring

containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-Rg, wherein each R¹ and R² is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷. In another embodiment of the present invention, R¹ is selected from: hydrogen, and -C₁₋₆alkyl, wherein each alkyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷. In another embodiment of the present invention, R¹ is selected from: hydrogen and ethyl, wherein ethyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷. In another embodiment of the present invention, R¹ is hydrogen.

In another embodiment of the present invention, R¹ is selected from: -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein R¹ is substituted with R⁷, or R¹ and R² together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-Rg, wherein each R¹ and R² is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷. In another embodiment of the present invention, R¹ is ethyl, wherein ethyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein R¹ is substituted with R⁷, or R¹ and R² together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-Rg, wherein each R¹ and R² is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷.

In another embodiment of the present invention, R¹ is selected from: -C₁₋₆alkyl, wherein each alkyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein R¹ is substituted with R⁷. In another embodiment of the present invention, R¹ is ethyl, wherein ethyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein R¹ is substituted with R⁷.

In another embodiment of the present invention, R^1 is selected from: -C₁₋₆alkyl, wherein each alkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein R^1 is substituted with R^7 ; and R^2 is hydrogen. In another embodiment of the present invention, R^1 is ethyl, wherein ethyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein R^1 is substituted with R^7 ; and R^2 is hydrogen.

In another embodiment of the present invention, R^1 is hydrogen. In another embodiment, R^1 is independently selected from: a bond, halogen, -OR^k, -CN, -C₁₋₆alkyl, -C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₃alkyl-, -C₂₋₆cycloheteroalkyl, and C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein R^1 is substituted with R^7 ; and R^2 is hydrogen.

In another embodiment, R^2 is selected from: a bond, hydrogen, halogen, -OR^k, -CN, -C₁₋₆alkyl, -C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₃alkyl-, -C₂₋₆cycloheteroalkyl, and C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 , or R^1 and R^2 together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R_g, wherein each R^1 and R^2 is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 .

In another embodiment of the present invention, R^2 is selected from: hydrogen, halogen, -OR^k, -CN, -C₁₋₆alkyl, -C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₃alkyl-, -C₂₋₆cycloheteroalkyl, and C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 , or R^1 and R^2 together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R_g, wherein each R^1 and R^2 is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 .

In another embodiment of the present invention, R^1 and R^2 are each independently selected from: hydrogen, and -C₁₋₆alkyl, wherein one of R^1 and R^2 is -C₁₋₆alkyl substituted with R^7 , and wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from R^L ; or a pharmaceutically acceptable salt thereof. In another embodiment of the present invention, R^1 and R^2 are each independently selected from: hydrogen, and ethyl, wherein one of R^1 and R^2 is ethyl substituted with R^7 , and wherein ethyl is unsubstituted or substituted with one to three substituents independently selected from R^L ; or a pharmaceutically acceptable salt thereof.

In another embodiment of the present invention, R^2 is selected from: hydrogen, halogen, -OR^k, -CN, -C₁₋₆alkyl, -C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₃alkyl-, -C₂₋₆cycloheteroalkyl, and C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 . In another embodiment of the present invention, R^2 is selected from: hydrogen, halogen, -OR^k, and -CN.

In another embodiment of the present invention, R^2 is selected from: -C₁₋₆alkyl, -C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₃alkyl-, -C₂₋₆cycloheteroalkyl, and C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein R^2 is substituted with R^7 , or R^1 and R^2 together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R_g, wherein each R^1 and R^2 is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 .

In another embodiment of the present invention, R^2 is selected from: -C₁₋₆alkyl, -C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₃alkyl-, -C₂₋₆cycloheteroalkyl, and C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein R^2 is substituted with R^7 . In another embodiment of the present invention, R^2 is selected from: -C₁₋₆alkyl, -C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₃alkyl-, -C₂₋₆cycloheteroalkyl, and C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein R^2 is substituted with R^7 ; and R^1 is hydrogen.

In another embodiment of the present invention, R^2 is selected from: hydrogen, halogen, -OR^k, -CN, and -C₁₋₆alkyl, wherein each alkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 , or R^1 and R^2 together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R_g, wherein each R^1 and R^2 is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 . In another embodiment of the present invention, R^2 is selected from: hydrogen, halogen, -OR^k, -CN, and -C₁₋₆alkyl, wherein each alkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 .

In another embodiment of the present invention, R^2 is selected from: hydrogen, and -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^L , and wherein one of R^1 and R^2 is substituted with R^7 , or R^1 and R^2 together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2

additional heteroatoms independently selected from oxygen, sulfur and N-R^g, wherein each R¹ and R² is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷.

In another embodiment of the present invention, R² is selected from: hydrogen, and -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷. In another embodiment of the present invention, R² is selected from: hydrogen and ethyl, wherein ethyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷. In another embodiment of the present invention, R² is hydrogen. In another embodiment of the present invention, R² is ethyl, wherein ethyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein R² is substituted with R⁷, or R¹ and R² together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R^g, wherein each R¹ and R² is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷.

In another embodiment of the present invention, R² is selected from: ethyl, wherein ethyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein R² is substituted with R⁷, or R¹ and R² together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R^g, wherein each R¹ and R² is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷.

In another embodiment of the present invention, R² is -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein R² is substituted with R⁷. In another embodiment of the present invention, R² is ethyl, wherein ethyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein R² is substituted with R⁷.

In another embodiment of the present invention, R² is -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein R² is substituted with R⁷; and R¹ is hydrogen. In another embodiment of the present invention, R² is ethyl, wherein ethyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein R² is substituted with R⁷; and R¹ is hydrogen.

In another embodiment, R² is independently selected from: a bond, halogen, -OR^k, -CN, -C₁₋₆alkyl, -C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₃alkyl-, -C₂₋₆cycloheteroalkyl, and C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted

or substituted with one to three substituents selected from R^L , and wherein R^2 is substituted with R^7 ; and R^1 is hydrogen.

In one class of the embodiments of the present invention, at least one of R^1 and R^2 is selected from: $-C_{1-6}$ alkyl, $-C_{3-6}$ cycloalkyl, C_{3-6} cycloalkyl- C_{1-3} alkyl-, $-C_{2-6}$ cycloheteroalkyl, and C_{2-6} cycloheteroalkyl- C_{1-3} alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents independently selected from R^L , and wherein one alkyl, cycloalkyl or cycloheteroalkyl is substituted with R^7 . In another class of the embodiments of the present invention, one of R^1 and R^2 is selected from: $-C_{1-6}$ alkyl, $-C_{3-6}$ cycloalkyl, C_{3-6} cycloalkyl- C_{1-3} alkyl-, $-C_{2-6}$ cycloheteroalkyl, and C_{2-6} cycloheteroalkyl- C_{1-3} alkyl-, wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents independently selected from R^L , and wherein one alkyl, cycloalkyl or cycloheteroalkyl is substituted with R^7 . In another class of the embodiments of the present invention, at least one of R^1 and R^2 is $-C_{1-6}$ alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from R^L , and wherein one alkyl is substituted with R^7 . In another class of the embodiments of the present invention, one of R^1 and R^2 is $-C_{1-6}$ alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from R^L , and wherein alkyl is substituted with R^7 . In another class of the embodiments of the present invention, one of R^1 and R^2 is $-C_{1-6}$ alkyl, wherein alkyl is substituted with one or two substituents independently selected from R^L , and wherein alkyl is substituted with R^7 . In another class of the embodiments of the present invention, at least one of R^1 and R^2 is ethyl, wherein ethyl is unsubstituted or substituted with one to three substituents independently selected from R^L , and wherein one ethyl is substituted with R^7 . In another class of the embodiments of the present invention, one of R^1 and R^2 is ethyl, wherein ethyl is unsubstituted or substituted with one to three substituents independently selected from R^L , and wherein ethyl is substituted with R^7 . In another class of the embodiments of the present invention, one of R^1 and R^2 is ethyl, wherein ethyl is substituted with one or two substituents independently selected from R^L , and wherein ethyl is substituted with R^7 .

In another embodiment, R^3 is absent or when present is selected from the group consisting of: hydrogen, halogen, $-OR^e$, $-CN$, $-C_{1-6}$ alkyl, $-C_{3-6}$ cycloalkyl, and C_{3-6} cycloalkyl- C_{1-3} alkyl-, wherein each alkyl and cycloalkyl is unsubstituted or substituted with one to three substituents selected from R^i . In another embodiment of the present invention, R^3 is absent or selected from the group consisting of: hydrogen, halogen, and $-C_{1-6}$ alkyl, wherein each alkyl is unsubstituted or substituted with one to three substituents selected from R^i . In another embodiment of the present invention, R^3 is absent or selected from the group consisting of: hydrogen, and halogen, wherein each alkyl is unsubstituted or substituted with one to three substituents selected from R^i . In another embodiment of the present invention, R^3 is absent. In another embodiment of the present invention,

R³ is absent or hydrogen. In another embodiment of the present invention, R³ is hydrogen. In another embodiment of the present invention, R³ is absent or halogen. In another embodiment of the present invention, R³ is halogen.

In another embodiment of the present invention, R³ is absent or -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from Rⁱ. In another embodiment of the present invention, R³ is -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from Rⁱ.

In another embodiment of the present invention, R⁴ is selected from the group consisting of: hydrogen, halogen, OR^e, C₀₋₅alkylNR^cR^d, C₁₋₆alkyl, and C₁₋₆alkyl-O-, wherein alkyl is

unsubstituted or substituted with one to three substituents selected from R^j, provided that when R⁴ is selected from the group consisting of: OR^e, C₀alkyl-NR^cR^d, and C₁₋₆alkyl-O-, then X is selected from the group consisting of: -CR⁹R⁹, C=O, and -C(R⁹)OC₁₋₆alkyl. In another embodiment of the present invention, R⁴ is selected from the group consisting of: hydrogen, halogen, -C₀₋₅alkylNR^cR^d, and -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j, provided that when R⁴ is selected from the group consisting of: -C₀alkyl-NR^cR^d, then X is selected from the group consisting of: -CR⁹R⁹, C=O, and -C(R⁹)OC₁₋₆alkyl.

In another embodiment of the present invention, R⁴ is selected from the group consisting of: hydrogen, halogen, and C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j. In another embodiment of the present invention, R⁴ is selected from the group consisting of: hydrogen, and C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j. In another embodiment of the present invention, R⁴ is hydrogen. In another embodiment of the present invention, R⁴ is -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j.

In another embodiment of the present invention, R⁵ is absent or when present R⁵ is selected from the group consisting of: hydrogen, -C₁₋₆alkyl, and -C₃₋₆cycloalkyl, wherein each alkyl and cycloalkyl is unsubstituted or substituted with one to three substituents selected from R^j. In another embodiment of the present invention, R⁵ is absent or selected from the group consisting of: hydrogen, and -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j.

In another embodiment of the present invention, R⁵ is absent. In another embodiment of the present invention, R⁵ is absent or hydrogen. In another embodiment of the present invention, R⁵ is hydrogen. In another embodiment of the present invention, R⁵ is absent or -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j. In another embodiment of the present invention, R⁵ is selected from the group consisting of: -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j. In another embodiment of the present invention, R⁵ is -C₁₋₆alkyl.

In another embodiment of the present invention, R^6 is absent, or when present R^6 is selected from the group consisting of: hydrogen, $-C_{1-6}$ alkyl, and $-C_{3-6}$ cycloalkyl, wherein each alkyl and cycloalkyl is unsubstituted or substituted with one to three substituents selected from R^j . In another embodiment of the present invention, R^6 is absent or selected from the group consisting of: hydrogen, and $-C_{1-6}$ alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j . In another embodiment of the present invention, R^6 is absent. In another embodiment of the present invention, R^6 is absent or hydrogen. In another embodiment of the present invention, R^6 is hydrogen. In another embodiment of the present invention, R^6 is absent or $-C_{1-6}$ alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j . In another embodiment of the present invention, R^6 is $-C_{1-6}$ alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j . In another embodiment of the present invention, R^6 is $-C_{1-6}$ alkyl.

In another embodiment of the present invention, R^7 is $-CO_2R^8$. In a class of this embodiment, R^7 is $-CO_2H$.

In another embodiment of the present invention, R^8 is selected from the group consisting of: hydrogen, $-C_{1-6}$ alkyl, and $-C_{3-6}$ cycloalkyl, wherein each alkyl, and cycloalkyl is unsubstituted or substituted with one to three substituents selected from R^j . In another embodiment of the present invention, R^8 is selected from the group consisting of: hydrogen, and $-C_{1-6}$ alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j . In another embodiment of the present invention, R^8 is hydrogen. In another embodiment of the present invention, R^8 is $-C_{1-6}$ alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j . In another embodiment of the present invention, R^8 is $-C_{1-6}$ alkyl.

In another embodiment of the present invention, R^9 is selected from the group consisting of: hydrogen, $-C_{1-6}$ alkyl, halogen, $-(CH_2)_{0-5}OH$, $-(CH_2)_{0-5}OC_{1-6}alkyl$, and $-(CH_2)_{0-6}NR^cR^d$, wherein each CH_2 is unsubstituted or substituted with one or two substituents selected from R^j , and alkyl is unsubstituted or substituted with one to three substituents selected from R^j . In another embodiment of the present invention, R^9 is selected from the group consisting of: hydrogen, $-C_{1-6}$ alkyl, halogen, and $-(CH_2)_{0-6}NR^cR^d$, wherein each CH_2 is unsubstituted or substituted with one or two substituents selected from R^j , and alkyl is unsubstituted or substituted with one to three substituents selected from R^j . In another embodiment of the present invention, R^9 is selected from the group consisting of: hydrogen, $-C_{1-6}$ alkyl, and halogen, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j . In another embodiment of the present invention, R^9 is selected from the group consisting of: hydrogen, and halogen. In another embodiment of the present invention, R^9 is selected from the group consisting of: hydrogen, and F. In another embodiment of the present invention, R^9 is hydrogen. In another embodiment of the

present invention, R^9 is $-C_{1-6}$ alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^j . In another embodiment of the present invention, R^9 is halogen.

In another embodiment of the present invention, R^a is selected from the group consisting of: $-C_{1-6}$ alkyl, halogen, $-C_{0-6}$ alkyl-OR^e, $-C_{0-6}$ alkyl-NR^cS(O)_nRe, $-C_{0-6}$ alkyl-S(O)_nRe, $-C_{0-6}$ alkyl-S(O)_nNR^cR^d, $-C_{0-6}$ alkyl-NR^cR^d, $-C_{0-6}$ alkyl-C(O)Re, $-C_{0-6}$ alkyl-OC(O)Re, $-C_{0-6}$ alkyl-CO₂Re, $-C_{0-6}$ alkyl-CN, $-C_{0-6}$ alkyl-C(O)NR^cR^d, $-C_{0-6}$ alkyl-NR^cC(O)Re, $-C_{0-6}$ alkyl-NR^cC(O)OR^e, $-C_{0-6}$ alkyl-NR^cC(O)NR^cR^d, -CF₃, -OCF₃, -OCHF₂, aryl, heteroaryl, $-C_{3-10}$ cycloalkyl, $-C_{3-10}$ cycloalkenyl, and $-C_{2-10}$ cycloheteroalkyl, wherein each alkyl, cycloalkyl, cycloalkenyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or substituted with one to five substituents independently selected from: $-C_{1-6}$ alkyl, halogen, OH, $-O-C_{1-6}$ alkyl, $-S(O)_2-C_{1-4}$ alkyl, -CN, -OCHF₂, -OCF₃, -CF₃, and $-C_{0-6}$ alkyl-NR^cR^d;

In another embodiment of the present invention, R^a is selected from the group consisting of: $-C_{1-6}$ alkyl, halogen, $-C_{0-6}$ alkyl-OR^e, $-C_{0-6}$ alkyl-NR^cS(O)_nRe, $-C_{0-6}$ alkyl-S(O)_nRe, $-C_{0-6}$ alkyl-S(O)_nNR^cR^d, $-C_{0-6}$ alkyl-NR^cR^d, $-C_{0-6}$ alkyl-C(O)Re, $-C_{0-6}$ alkyl-OC(O)Re, $-C_{0-6}$ alkyl-CO₂Re, $-C_{0-6}$ alkyl-CN, $-C_{0-6}$ alkyl-C(O)NR^cR^d, $-C_{0-6}$ alkyl-NR^cC(O)Re, $-C_{0-6}$ alkyl-NR^cC(O)OR^e, $-C_{0-6}$ alkyl-NR^cC(O)NR^cR^d, -CF₃, -OCF₃, and -OCHF₂, wherein each alkyl is unsubstituted or substituted with one to five substituents independently selected from: $-C_{1-6}$ alkyl, halogen, OH, $-O-C_{1-6}$ alkyl, $-S(O)_2-C_{1-4}$ alkyl, -CN, -OCHF₂, -OCF₃, -CF₃, and $-C_{0-6}$ alkyl-NR^cR^d. In another embodiment of the present invention, R^a is selected from the group consisting of: $-C_{1-6}$ alkyl, halogen, -OR^e, $-C_{0-6}$ alkyl-NR^cR^d, $-C(O)Re$, $-OC(O)Re$, $-CO_2Re$, -CN, -CF₃, -OCF₃, and -OCHF₂, wherein each alkyl is unsubstituted or substituted with one to five substituents independently selected from: $-C_{1-6}$ alkyl, halogen, OH, $-O-C_{1-6}$ alkyl, $-S(O)_2-C_{1-4}$ alkyl, -CN, -OCHF₂, -OCF₃, -CF₃, and $-C_{0-6}$ alkyl-NR^cR^d.

In another embodiment of the present invention, R^a is selected from the group consisting of: $-C_{1-6}$ alkyl, halogen, -OR^e, $-C_{0-5}$ alkyl-NR^cR^d, -CN, -CF₃, -OCF₃, and -OCHF₂, wherein alkyl is unsubstituted or substituted with one to five substituents independently selected from: $-C_{1-6}$ alkyl, halogen, OH, $-O-C_{1-6}$ alkyl and -CF₃. In another embodiment of the present invention, R^a is selected from the group consisting of: $-C_{1-6}$ alkyl, and halogen, wherein alkyl is unsubstituted or substituted with one to five substituents independently selected from: $-C_{1-6}$ alkyl, halogen, OH, $-O-C_{1-6}$ alkyl and -CF₃. In another embodiment of the present invention, R^a is $-C_{1-6}$ alkyl, wherein alkyl is unsubstituted or substituted with one to five substituents independently selected from: $-C_{1-6}$ alkyl, halogen, OH, $-O-C_{1-6}$ alkyl and -CF₃. In another embodiment of the present invention, R^a is halogen. In a class of this embodiment, R^a is F.

In another embodiment of the present invention, R^b is selected from the group consisting of: $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, -CF₃, halogen, -CN, -OH, $-OC_{1-10}$ alkyl, $-OC_{2-10}$ alkenyl, $-O(CH_2)_pOC_{1-10}$ alkyl, $-O(CH_2)_pNR^cS(O)_mRe$, $-O(CH_2)_pS(O)_mRe$, $-O(CH_2)_pS(O)_mNR^cR^d$, -

O(CH₂)_pNR^cR^d, -C(O)R^e, -OC(O)R^e, -CO₂R^e, -C(O)NR^cR^d, -NR^cC(O)R^e, -NR^cC(O)OR^e, -NR^cC(O)NR^cR^d, -OCF₃, and -OCHF₂, wherein each CH, CH₂, alkyl, and alkenyl is unsubstituted or substituted with one to five substituents independently selected from: -C₁₋₆alkyl, halogen, OH, -O-C₁₋₆alkyl and -CF₃. In another embodiment of the present invention, R^b is selected from the group consisting of: -C₁₋₁₀alkyl, -CF₃, halogen, -CN, -OH, -OC₁₋₁₀alkyl, -O(CH₂)_pOC₁₋₁₀alkyl, -O(CH₂)_pNR^cR^d, -C(O)R^e, -OC(O)R^e, -CO₂R^e, -OCF₃, and -OCHF₂, wherein each CH, CH₂, alkyl, and alkenyl is unsubstituted or substituted with one to five substituents independently selected from: -C₁₋₆alkyl, halogen, OH, -O-C₁₋₆alkyl and -CF₃.

In another embodiment of the present invention, R^b is selected from the group consisting of: -C₁₋₁₀alkyl, -CF₃, halogen, -CN, -OH, -OC₁₋₁₀alkyl, -OCF₃, and -OCHF₂, wherein each CH, and alkyl is unsubstituted or substituted with one to five substituents independently selected from: -C₁₋₆alkyl, halogen, OH, -O-C₁₋₆alkyl and -CF₃. In another embodiment of the present invention, R^b is selected from the group consisting of: -C₁₋₁₀alkyl, -CF₃, halogen, and -OC₁₋₁₀alkyl, wherein each alkyl is unsubstituted or substituted with one to five substituents independently selected from: -C₁₋₆alkyl, halogen, OH, -O-C₁₋₆alkyl and -CF₃. In a class of this embodiment, R^b is selected from the group consisting of: -CH₃, -CF₃, F, and -OCH₃.

In another embodiment of the present invention, R^c and R^d are each independently selected from the group consisting of: hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₁₀alkyl-, C₂₋₅cycloheteroalkyl, C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-, aryl, heteroaryl, aryl-C₁₋₁₀alkyl-, and heteroaryl-C₁₋₁₀alkyl-, wherein each alkyl, alkenyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or substituted with one to three substituents independently selected from R^f. In another embodiment of the present invention, R^c and R^d are each independently selected from the group consisting of: hydrogen, C₁₋₁₀alkyl, and C₂₋₁₀alkenyl, wherein each alkyl, and alkenyl is unsubstituted or substituted with one to three substituents independently selected from R^f. In another embodiment of the present invention, R^c and R^d are each independently selected from the group consisting of: hydrogen, and C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from R^f. In another embodiment of the present invention, R^c and R^d are each hydrogen. In another embodiment of the present invention, R^c and R^d are each independently selected from the group consisting of: C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from R^f. In another embodiment of the present invention, R^c and R^d are each independently selected from the group consisting of: C₁₋₁₀alkyl.

In another embodiment of the present invention, R^c is selected from the group consisting of: hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₃₋₆cycloalkyl, C₃₋₆ cycloalkyl-C₁₋₁₀alkyl-, C₂₋₅cycloheteroalkyl, C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-, aryl, heteroaryl, aryl-C₁₋₁₀alkyl-, and heteroaryl-C₁₋₁₀alkyl-, wherein each alkyl, alkenyl, cycloalkyl, cycloheteroalkyl, aryl and

heteroaryl is unsubstituted or substituted with one to three substituents independently selected from R^f. In another embodiment of the present invention, R^c is selected from the group consisting of: hydrogen, C₁₋₁₀alkyl, and C₂₋₁₀alkenyl, wherein each alkyl, and alkenyl is unsubstituted or substituted with one to three substituents independently selected from R^f. In another embodiment of the present invention, R^c is selected from the group consisting of: hydrogen, and C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from R^f. In another embodiment of the present invention, R^c is hydrogen. In another embodiment of the present invention, R^c is -C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from R^f. In another embodiment of the present invention, R^c is -C₁₋₁₀alkyl.

In another embodiment of the present invention, R^d is selected from the group consisting of: hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₃₋₆cycloalkyl, C₃₋₆ cycloalkyl-C₁₋₁₀alkyl-, C₂₋₅cycloheteroalkyl, C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-, aryl, heteroaryl, aryl-C₁₋₁₀alkyl-, and heteroaryl-C₁₋₁₀alkyl-, wherein each alkyl, alkenyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or substituted with one to three substituents independently selected from R^f. In another embodiment of the present invention, R^d is selected from the group consisting of: hydrogen, C₁₋₁₀alkyl, and C₂₋₁₀alkenyl, wherein each alkyl and alkenyl is unsubstituted or substituted with one to three substituents independently selected from R^f. In another embodiment of the present invention, R^d is selected from the group consisting of: hydrogen, and C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from R^f. In another embodiment of the present invention, R^d is hydrogen. In another embodiment of the present invention, R^d is -C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from R^f. In another embodiment of the present invention, R^d is -C₁₋₁₀alkyl.

In another embodiment of the present invention, R^e is independently selected from the group consisting of: hydrogen, -C₁₋₁₀alkyl, and -C₂₋₁₀alkenyl, wherein each alkyl, and alkenyl is unsubstituted or substituted with one to three substituents selected from R^h. In another embodiment of the present invention, each R^e is independently selected from the group consisting of: hydrogen, and -C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^h. In another embodiment of the present invention, R^e is hydrogen. In another embodiment of the present invention, each R^e is -C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents selected from R^h. In another embodiment of the present invention, each R^e is -C₁₋₁₀alkyl.

In another embodiment of the present invention, each R^f is selected from the group consisting of: halogen, -C₁₋₁₀alkyl, -OH, -O-C₁₋₄alkyl, -CN, -CF₃, -OCHF₂, and -OCF₃, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from: -OH,

halogen, C₁₋₆alkyl, cyano and S(O)₂C₁₋₆alkyl. In another embodiment of the present invention, each R^f is selected from the group consisting of: halogen, -C₁₋₁₀alkyl, -CN, and -CF₃, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from: -OH, halogen, C₁₋₆alkyl, cyano and S(O)₂C₁₋₆alkyl.

5 In another embodiment of the present invention, each R^f is selected from the group consisting of: halogen, and C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from: -OH, halogen, C₁₋₆alkyl, cyano and S(O)₂C₁₋₆alkyl. In another embodiment of the present invention, each R^f is selected from the group consisting of: halogen. In another embodiment of the present invention, each R^f is -C₁₋₁₀alkyl, wherein alkyl is
10 unsubstituted or substituted with one to three substituents independently selected from: -OH, halogen, C₁₋₆alkyl, cyano and S(O)₂C₁₋₆alkyl. In another embodiment of the present invention, each R^f is -C₁₋₁₀alkyl.

In another embodiment of the present invention, each R^g is independently selected from the group consisting of: hydrogen, and -C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with
15 one to five halogens. In another embodiment of the present invention, R^g is hydrogen. In another embodiment of the present invention, each R^g is -C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with one to five halogens. In another embodiment of the present invention, each R^g is -C₁₋₁₀alkyl.

In another embodiment of the present invention, each R^h is independently selected from the
20 group consisting of: halogen, -C₁₋₁₀alkyl, -OH, -O-C₁₋₄alkyl, -CN, -CF₃, -OCHF₂, and -OCF₃, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from: -OH, halogen, C₁₋₆alkyl, cyano and S(O)₂C₁₋₆alkyl. In another embodiment of the present invention, each R^h is independently selected from the group consisting of: halogen, -C₁₋₁₀alkyl, -CN, and -CF₃, wherein alkyl is unsubstituted or substituted with one to three substituents
25 independently selected from: -OH, halogen, C₁₋₆alkyl, cyano and S(O)₂C₁₋₆alkyl.

In another embodiment of the present invention, each R^h is independently selected from the group consisting of: halogen, and C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from: -OH, halogen, C₁₋₆alkyl, cyano and S(O)₂C₁₋₆alkyl. In another embodiment of the present invention, each R^h is independently selected from the
30 group consisting of: halogen. In another embodiment of the present invention, each R^h is -C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from: -OH, halogen, C₁₋₆alkyl, cyano and S(O)₂C₁₋₆alkyl. In another embodiment of the present invention, each R^h is -C₁₋₁₀alkyl.

In another embodiment of the present invention, each Rⁱ is independently selected from the
35 group consisting of: -C₁₋₆alkyl, -ORE, -NR^cS(O)_mRe, halogen, -S(O)_mRe, -S(O)_mNR^cR^d, -NR^cR^d, -C(O)Re, -OC(O)Re, -CO₂Re, -CN, -C(O)NR^cR^d, -NR^cC(O)Re, -NR^cC(O)ORE, -

NR^cC(O)NR^cR^d, -CF₃, -OCF₃, and -OCHF₂. In another embodiment of the present invention, each Rⁱ is independently selected from the group consisting of: -C₁₋₆alkyl, -OR^e, halogen, -S(O)_mR^e, -NR^cR^d, -C(O)R^e, -OC(O)R^e, -CO₂R^e, -CN, -CF₃, -OCF₃, and -OCHF₂. In another embodiment of the present invention, each Rⁱ is independently selected from the group consisting of:
 5 -C₁₋₆alkyl, -OR^e, halogen, -NR^cR^d, -CN, -CF₃, -OCF₃, and -OCHF₂. In another embodiment of the present invention, each Rⁱ is independently selected from the group consisting of: -C₁₋₆alkyl, and halogen. In another embodiment of the present invention, each Rⁱ is independently selected from the group consisting of: halogen. In another embodiment of the present invention, each Rⁱ is independently selected from the group consisting of: -C₁₋₆alkyl.

10 In another embodiment of the present invention, each R^j is independently selected from the group consisting of: -C₁₋₆alkyl, -OR^e, -NR^cS(O)_mR^e, halogen, -S(O)_mR^e, -S(O)_mNR^cR^d, -NR^cR^d, -C(O)R^e, -OC(O)R^e, -CO₂R^e, -CN, -C(O)NR^cR^d, -NR^cC(O)R^e, -NR^cC(O)OR^e, -NR^cC(O)NR^cR^d, -CF₃, -OCF₃, and -OCHF₂. In another embodiment of the present invention, each R^j is independently selected from the group consisting of: -C₁₋₆alkyl, -OR^e, halogen, -
 15 S(O)_mR^e, -NR^cR^d, -C(O)R^e, -OC(O)R^e, -CO₂R^e, -CN, -CF₃, -OCF₃, and -OCHF₂. In another embodiment of the present invention, each R^j is independently selected from the group consisting of: -C₁₋₆alkyl, -OR^e, halogen, -NR^cR^d, -CN, -CF₃, -OCF₃, and -OCHF₂. In another embodiment of the present invention, each R^j is independently selected from the group consisting of: -C₁₋₆alkyl, and halogen. In another embodiment of the present invention, each R^j is halogen. In another
 20 embodiment of the present invention, each R^j is -C₁₋₆alkyl.

In another embodiment of the present invention, each R^k is independently selected from the group consisting of: hydrogen, -C₁₋₆alkyl, -CF₃, and -CHF₂, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from: -OH, -OC₁₋₆alkyl, halogen, cyano, and -S(O)₂C₁₋₆alkyl. In another embodiment of the present invention, each R^k is
 25 independently selected from the group consisting of: hydrogen, and -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from: -OH, -OC₁₋₆alkyl, halogen, cyano, and -S(O)₂C₁₋₆alkyl. In another embodiment of the present invention, R^k is hydrogen. In another embodiment of the present invention, each R^k is -C₁₋₆alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from: -OH, -
 30 OC₁₋₆alkyl, halogen, cyano, and -S(O)₂C₁₋₆alkyl. In another embodiment of the present invention, each R^k is -C₁₋₆alkyl.

In another embodiment of the present invention, each R^L is independently selected from the group consisting of: -CO₂C₁₋₆alkyl, -C₁₋₁₀alkyl, -C₂₋₁₀alkenyl, -C₂₋₁₀alkynyl, -C₃₋₆cycloalkyl, and -C₂₋₆cycloheteroalkyl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, and cycloheteroalkyl is
 35 unsubstituted or substituted with one to four substituents independently selected from -C₁₋₆alkyl,

halogen, and -OC₁₋₆alkyl. In another embodiment of the present invention, each R^L is independently selected from the group consisting of: -CO₂C₁₋₆alkyl, -C₁₋₁₀alkyl, -C₃₋₆cycloalkyl, and -C₂₋₆cycloheteroalkyl, wherein each alkyl, cycloalkyl, and cycloheteroalkyl is unsubstituted or substituted with one to four substituents independently selected from C₁₋₆alkyl, halogen, and -OC₁₋₆alkyl. In another embodiment of the present invention, each R^L is independently selected from the group consisting of: -C₁₋₁₀alkyl, -C₃₋₆cycloalkyl, and -C₂₋₆cycloheteroalkyl, wherein each alkyl, cycloalkyl, and cycloheteroalkyl is unsubstituted or substituted with one to four substituents independently selected from C₁₋₆alkyl, halogen, and -OC₁₋₆alkyl.

In another embodiment of the present invention, each R^L is independently selected from the group consisting of: -C₁₋₁₀alkyl, and -C₃₋₆cycloalkyl, wherein each alkyl, and cycloalkyl is unsubstituted or substituted with one to four substituents independently selected from -C₁₋₆alkyl, halogen, and -OC₁₋₆alkyl. In another embodiment of the present invention, each R^L is independently selected from the group consisting of: -CH₃, and -cyclopropyl. In another embodiment of the present invention, each R^L is -C₁₋₁₀alkyl, wherein alkyl is unsubstituted or substituted with one to four substituents independently selected from C₁₋₆alkyl, halogen, and -OC₁₋₆alkyl. In another embodiment of the present invention, each R^L is -CH₃. In another embodiment of the present invention, each R^L is independently selected from the group consisting of: -C₃₋₆cycloalkyl, wherein cycloalkyl is unsubstituted or substituted with one to four substituents independently selected from C₁₋₆alkyl, halogen, and -OC₁₋₆alkyl. In another embodiment of the present invention, each R^L is -cyclopropyl.

In another embodiment of the present invention, each R^m is independently selected from the group consisting of: -C₁₋₁₀alkyl, and -C₂₋₁₀alkenyl. In another embodiment of the present invention, each R^m is -C₁₋₁₀alkyl.

In another embodiment of the present invention, n is 0, 1 or 2. In a class of this embodiment, n is 0 or 1. In another class of this embodiment, n is 1 or 2. In another class of this embodiment, n is 0. In another class of this embodiment, n is 1. In another class of this embodiment, n is 2.


In another embodiment of the present invention, m is 0, 1 or 2. In a class of this embodiment, m is 0 or 1. In another class of this embodiment, m is 1 or 2. In another class of this embodiment, m is 0. In another class of this embodiment, m is 1. In another class of this embodiment, m is 2.

In another embodiment of the present invention, each p is independently selected from: 0, 1, 2, 3, 4, 5 or 6. In another embodiment of the present invention, p is 0, 1, 2, 3 or 4. In a class of this embodiment, p is 0, 1, 2 or 3. In a class of this embodiment, p is 0, 1 or 2. In another embodiment of the present invention, p is 1, 2, 3 or 4. In a class of this embodiment, p is 1, 2 or 3. In a class of this embodiment, p is 1 or 2. In another class of this embodiment, p is 0 or 1. In another class of this embodiment, p is 0 or 2. In another class of this embodiment, p is 0. In another class of this

embodiment, p is 1. In another class of this embodiment, p is 2. In another class of this embodiment, p is 3. In another class of this embodiment, p is 4. In another class of this embodiment, p is 5. In another class of this embodiment, p is 6.


In another embodiment of the present invention, each r is independently selected from: 0, 1, 2 or 3. In a class of this embodiment, r is 0, 1 or 2. In another class of this embodiment, r is 1, 2 or 3. In a class of this embodiment, r is 1 or 2. In another class of this embodiment, r is 0 or 1. In another class of this embodiment, r is 0 or 2. In another class of this embodiment, r is 0 or 1. In another class of this embodiment, r is 1 or 3. In another class of this embodiment, r is 2 or 3. In another class of this embodiment, r is 0 or 2. In another class of this embodiment, r is 0. In another class of this embodiment, r is 1. In another class of this embodiment, r is 2. In another class of this embodiment, r is 3.

In another embodiment of the present invention, the invention relates to compounds of structural formula Ia:




or a pharmaceutically acceptable salt thereof.

In another embodiment of the present invention, the invention relates to compounds of structural formula Ib:




or a pharmaceutically acceptable salt thereof.

In another embodiment of the present invention, the invention relates to compounds of structural formula Ic:




or a pharmaceutically acceptable salt thereof.

In another embodiment of the present invention, the invention relates to compounds of structural formula Id:




or a pharmaceutically acceptable salt thereof.

In another embodiment of the present invention, the invention relates to compounds of structural formula Ie:




or a pharmaceutically acceptable salt thereof.

In another embodiment of the present invention, the invention relates to compounds of structural formula If:




or a pharmaceutically acceptable salt thereof.

In another embodiment of the present invention, the invention relates to compounds of structural formula Ig:




or a pharmaceutically acceptable salt thereof.

In another embodiment of the present invention, the invention relates to compounds of structural formula Ih:




or a pharmaceutically acceptable salt thereof.

In another embodiment of the present invention, the invention relates to compounds of structural formula Ii:




or a pharmaceutically acceptable salt thereof.

In another embodiment of the present invention, the invention relates to compounds of structural formula Ij:




or a pharmaceutically acceptable salt thereof.

In another embodiment of the present invention, the invention relates to compounds of structural formula Ik:




or a pharmaceutically acceptable salt thereof.

In another embodiment of the present invention, the invention relates to compounds of structural formula II:



or a pharmaceutically acceptable salt thereof.


In another embodiment of the present invention, the invention relates to compounds of structural formula Im:



or a pharmaceutically acceptable salt thereof.

The compound of structural formula I includes the compounds of structural formulas Ia, Ib, Ic, Id, Ie, If, Ig, Ih, Ii, Ij, Ik, Il and Im, and pharmaceutically acceptable salts, hydrates and solvates thereof.

Another embodiment of the present invention relates to compounds of structural formula Ia:



wherein

"a" is a single bond;

Q is -CR⁴R⁶;

X is -CR⁹R⁹;

Y is $-\text{CR}^9\text{R}^9$;

Z is oxygen;

A is selected from the group consisting of:

- (1) aryl,
- 5 (2) heteroaryl, and
- (3) C_{2-5} cycloheteroalkyl,

wherein each aryl, heteroaryl and cycloheteroalkyl is unsubstituted or substituted with one to five substituents selected from R^a ;

B is selected from the group consisting of:

- 10 (1) aryl,
- (2) aryl- C_{1-10} alkyl-, and
- (3) heteroaryl,

wherein each alkyl, aryl and heteroaryl is unsubstituted or substituted with one to five substituents selected from R^b ;

15 R^1 and R^2 are each independently selected from:

- (1) hydrogen, and
- (2) $-\text{C}_{1-6}$ alkyl,

wherein one of R^1 and R^2 is $-\text{C}_{1-6}$ alkyl substituted with R^7 , and wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from R^L ;

20 R^3 is absent or hydrogen;

R^4 is hydrogen;

R^5 is absent or hydrogen;

R^6 is absent or hydrogen;

R^7 is $-\text{CO}_2\text{R}^8$;

25 R^8 is hydrogen;

R^9 is hydrogen;


each R^L is independently selected from the group consisting of: $-\text{C}_{1-10}$ alkyl, and $-\text{C}_{3-6}$ cycloalkyl,

wherein each alkyl, and cycloalkyl is unsubstituted or substituted with one to four substituents independently selected from C_{1-6} alkyl, halogen, and $-\text{OC}_{1-6}$ alkyl; and

30 each r is independently selected from: 0, 1, 2 or 3;

or a pharmaceutically acceptable salt thereof.

Another embodiment of the present invention relates to compounds of structural Formula Ia:



wherein

"a" is a single bond;

Q is $-\text{CR}^4\text{R}^6$;

5 X is $-\text{CR}^9\text{R}^9$;

Y is $-\text{CR}^9\text{R}^9$;

Z is oxygen;

A is selected from the group consisting of:

- (1) phenyl,
- 10 (2) pyridyl, and
- (3) azetidine,

wherein each phenyl, pyridyl and azetidine is unsubstituted or substituted with one to four substituents selected from R^a ;

B is selected from the group consisting of:

- 15 (1) phenyl,
- (2) $-\text{CH}_2$ -phenyl, and
- (3) pyridyl,

wherein each phenyl and pyridyl is unsubstituted or substituted with one to five substituents selected from R^b ;

20 R^1 and R^2 are each independently selected from:

- (1) hydrogen, and
- (2) ethyl,

wherein one of R^1 and R^2 is ethyl substituted with R^7 , and wherein ethyl is unsubstituted or substituted with one to three substituents independently selected from R^L ;

25 R^3 is absent or hydrogen;

R^4 is hydrogen;

R^5 is hydrogen;

R^6 is hydrogen;

R^7 is $-\text{CO}_2\text{R}^8$;

R⁸ is hydrogen;


R⁹ is hydrogen;

each R^L is independently selected from the group consisting of: -CH₃, and -cyclopropyl; and



each r is independently selected from: 0, 1, 2 or 3;

5 or a pharmaceutically acceptable salt thereof.

Illustrative, but non-limiting, examples of the compounds of the present invention that are useful as agonists of G-protein-coupled receptor 40 (GPR40) are the following compounds:



10



and ; and pharmaceutically acceptable salts thereof.

Although the specific stereochemistries described above are preferred, other stereoisomers, including diastereoisomers, enantiomers, epimers, and mixtures of these may also have utility in
15 treating GPR40 mediated diseases.

Synthetic methods for making the compounds are disclosed in the Examples shown below. Where synthetic details are not provided in the examples, the compounds are readily made by a person of ordinary skill in the art of medicinal chemistry or synthetic organic chemistry by applying the synthetic information provided herein. Where a stereochemical center is not defined, the
20 structure represents a mixture of stereoisomers at that center. For such compounds, the individual

stereoisomers, including enantiomers, diastereoisomers, and mixtures of these are also compounds of the invention.

Definitions:

5 "Ac" is acetyl, which is $\text{CH}_3\text{C}(=\text{O})-$.

"Alkyl" means saturated carbon chains which may be linear or branched or combinations thereof, unless the carbon chain is defined otherwise. Other groups having the prefix "alk", such as alkoxy and alkanoyl, also may be linear or branched, or combinations thereof, unless the carbon chain is defined otherwise. The term $-\text{C}_2\text{alkyl}$ is ethyl. Examples of alkyl groups include methyl, 10 ethyl, propyl, isopropyl, butyl, sec- and tert-butyl, pentyl, hexyl, heptyl, octyl, nonyl, and the like. In one embodiment of the present invention, alkyl is methyl. In another embodiment of the present invention, alkyl is ethyl.

"Alkenyl" means carbon chains which contain at least one carbon-carbon double bond, and which may be linear or branched, or combinations thereof, unless otherwise defined. Examples of 15 alkenyl include vinyl, allyl, isopropenyl, pentenyl, hexenyl, heptenyl, 1-propenyl, 2-butenyl, 2-methyl-2-butenyl, and the like.

"Alkynyl" means carbon chains which contain at least one carbon-carbon triple bond, and which may be linear or branched, or combinations thereof, unless otherwise defined. Examples of 20 alkynyl include ethynyl, propargyl, 3-methyl-1-pentyne, 2-heptyne and the like.

"Cycloalkyl" means a saturated monocyclic, bicyclic or bridged carbocyclic ring, having a specified number of carbon atoms. The term may also be used to describe a carbocyclic ring fused to an aryl group. Examples of cycloalkyl include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, and the like. In one embodiment of the present invention, cycloalkyl is cyclopropane.

"Cycloalkenyl" means a nonaromatic monocyclic or bicyclic carbocyclic ring containing at 25 least one double bond. Examples of cycloalkenyl include cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl and the like.

"Cycloheteroalkyl" means a saturated monocyclic, bicyclic or bridged carbocyclic ring or ring system containing at least one ring heteroatom selected from N, NH, S (including SO and SO_2) and O. The cycloheteroalkyl ring may be substituted on the ring carbons and/or the ring nitrogen(s). 30 Examples of cycloheteroalkyl include tetrahydrofuran, pyrrolidine, tetrahydrothiophene, azetidine, piperazine, piperidine, morpholine, oxetane and tetrahydropyran. In one embodiment of the present invention, cycloheteroalkyl is azetidine.

"Cycloheteroalkenyl" means a nonaromatic monocyclic, bicyclic or bridged carbocyclic ring or ring system containing at least one double bond and containing at least one heteroatom selected 35 from N, NH, S (including SO and SO_2) and O.

"Aryl" means a monocyclic, bicyclic or tricyclic carbocyclic aromatic ring or ring system containing 5-14 carbon atoms, wherein at least one of the rings is aromatic. Examples of aryl include phenyl and naphthyl. In one embodiment of the present invention, aryl is phenyl.

"Heteroaryl" means monocyclic, bicyclic or tricyclic ring or ring system containing 5-14 carbon atoms and containing at least one ring heteroatom selected from N, NH, S (including SO and SO₂) and O, wherein at least one of the heteroatom containing rings is aromatic. Examples of heteroaryl include pyrrolyl, isoxazolyl, isothiazolyl, pyrazolyl, pyridyl, oxazolyl, oxadiazolyl, thiadiazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, furanyl, triazinyl, thienyl, pyrimidyl, pyridazinyl, pyrazinyl, benzisoxazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, benzpyrazole (or indazole), benzothiophenyl (including S-oxide and dioxide), furo(2,3-b)pyridyl, quinolyl, indolyl, isoquinolyl, quinazolinyl, dibenzofuranyl, and the like. In one embodiment of the present invention, heteroaryl is pyridine.


"Halogen" includes fluorine, chlorine, bromine and iodine. In one embodiment of the present invention, halogen is bromine, chlorine or fluorine. In another embodiment of the present invention, halogen is chlorine or fluorine. In another embodiment of the present invention, halogen is bromine. In another embodiment of the present invention, halogen is chlorine. In another embodiment of the present invention, halogen is fluorine.

"Me" represents methyl.

"Oxo" is =O.

When any variable (e.g., R¹, R^a, etc.) occurs more than one time in any constituent or in formula I, its definition on each occurrence is independent of its definition at every other occurrence. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. A squiggly line across a bond in a substituent variable represents the point of attachment.

Under standard nomenclature used throughout this disclosure, the terminal portion of the designated side chain is described first, followed by the adjacent functionality toward the point of attachment. For example, a C₁₋₅ alkylcarbonylamino C₁₋₆ alkyl substituent is equivalent to:



For example, -NR^cC(O)R^e is equivalent to -N(R^c)C(O)R^e.

Unless expressly depicted or described otherwise, substituents depicted in a structural formula with a "floating" bond, such as but not limited to R³, is permitted on any available carbon atom in the ring to which the substituent is attached. In one embodiment of the present invention, R³ may be substituted on any CH in the ring to which R³ is attached.

In choosing compounds of the present invention, one of ordinary skill in the art will recognize that the various substituents, i.e. R^1 , R^2 , etc., are to be chosen in conformity with well-known principles of chemical structure connectivity and stability.

The term "substituted" shall be deemed to include multiple degrees of substitution by a named substituent. Where multiple substituent moieties are disclosed or claimed, the substituted compound can be independently substituted by one or more of the disclosed or claimed substituent moieties, singly or plurally. By independently substituted, it is meant that the (two or more) substituents can be the same or different.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, salts and/or dosage forms which are, using sound medical judgment, and following all applicable government regulations, safe and suitable for administration to a human being or an animal.

The term "% enantiomeric excess" (abbreviated "ee") shall mean the % major enantiomer less the % minor enantiomer. Thus, a 70% enantiomeric excess corresponds to formation of 85% of one enantiomer and 15% of the other. The term "enantiomeric excess" is synonymous with the term "optical purity."

Compounds of Formula I may contain one or more asymmetric centers and can thus occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. The present invention is meant to comprehend all such isomeric forms of the compounds of Formula I.

Some of the compounds described herein contain olefinic double bonds, and unless specified otherwise, are meant to include both E and Z geometric isomers.

Tautomers are defined as compounds that undergo rapid proton shifts from one atom of the compound to another atom of the compound. Some of the compounds described herein may exist as tautomers with different points of attachment of hydrogen. Such an example may be a ketone and its enol form known as keto-enol tautomers. The individual tautomers as well as mixture thereof are encompassed with compounds of Formula I.

In the compounds of general formula I, the atoms may exhibit their natural isotopic abundances, or one or more of the atoms may be artificially enriched in a particular isotope having the same atomic number, but an atomic mass or mass number different from the atomic mass or mass number predominately found in nature. The present invention is meant to include all suitable isotopic variations of the compounds of structural formula I. For example, different isotopic forms of hydrogen (H) include protium (^1H), deuterium (^2H), and tritium (^3H). Protium is the predominant hydrogen isotope found in nature. Enriching for deuterium may afford certain therapeutic advantages, such as increasing in vivo half-life or reducing dosage requirements, or may provide a compound useful as a standard for characterization of biological samples. Tritium is radioactive and

may therefore provide for a radiolabeled compound, useful as a tracer in metabolic or kinetic studies. Isotopically-enriched compounds within structural formula I, can be prepared without undue experimentation by conventional techniques well known to those skilled in the art or by processes analogous to those described in the Schemes and Examples herein using appropriate isotopically-enriched reagents and/or intermediates.

The independent syntheses of optical isomers and diastereoisomers or their chromatographic separations may be achieved as known in the art by appropriate modification of the methodology disclosed herein. Their absolute stereochemistry may be determined by the X-ray crystallography of crystalline products or crystalline intermediates which are derivatized, if necessary, with a reagent containing an asymmetric center of known absolute configuration.

If desired, racemic mixtures of the compounds may be separated so that the individual enantiomers are isolated. The separation can be carried out by methods well-known in the art, such as the coupling of a racemic mixture of compounds to an enantiomerically pure compound to form a diastereoisomeric mixture, followed by separation of the individual diastereoisomers by standard methods, such as fractional crystallization or chromatography. The coupling reaction is often the formation of salts using an enantiomerically pure acid or base. The diastereomeric derivatives may then be converted to the pure enantiomers by cleavage of the added chiral residue. The racemic mixture of the compounds can also be separated directly by chromatographic methods utilizing chiral stationary phases, which methods are well known in the art.

Alternatively, any enantiomer of a compound may be obtained by stereoselective synthesis using optically pure starting materials or reagents of known configuration by methods well known in the art.

Furthermore, some of the crystalline forms for compounds of the present invention may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds of the instant invention may form solvates with water or common organic solvents. Such solvates are encompassed within the scope of this invention.

It is generally preferable to administer compounds of the present invention as enantiomerically pure formulations. Racemic mixtures can be separated into their individual enantiomers by any of a number of conventional methods. These include chiral chromatography, derivatization with a chiral auxiliary followed by separation by chromatography or crystallization, and fractional crystallization of diastereomeric salts.

Salts:

It will be understood that, as used herein, references to the compounds of the present invention are meant to also include the pharmaceutically acceptable salts, and also salts that are not

pharmaceutically acceptable when they are used as precursors to the free compounds or their pharmaceutically acceptable salts or in other synthetic manipulations.

The compounds of the present invention may be administered in the form of a pharmaceutically acceptable salt. The term "pharmaceutically acceptable salt" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Salts of basic compounds encompassed within the term "pharmaceutically acceptable salt" refer to non-toxic salts of the compounds of this invention which are generally prepared by reacting the free base with a suitable organic or inorganic acid.

Representative salts of basic compounds of the present invention include, but are not limited to, the following: acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinolate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, N-methylglucamine ammonium salt, oleate, oxalate, pamoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, sulfate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodide and valerate. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof include, but are not limited to, salts derived from inorganic bases including aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic, mangamous, potassium, sodium, zinc, and the like. Particularly preferred are the ammonium, calcium, magnesium, potassium, and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, cyclic amines, and basic ion-exchange resins, such as arginine, betaine, caffeine, choline, N,N-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like.

Also, in the case of a carboxylic acid (-COOH) or alcohol group being present in the compounds of the present invention, pharmaceutically acceptable esters of carboxylic acid derivatives, such as methyl, ethyl, or pivaloyloxymethyl, or acyl derivatives of alcohols, such as *O*-acetyl, *O*-pivaloyl, *O*-benzoyl, and *O*-aminoacyl, can be employed. Included are those esters and acyl groups known in the art for modifying the solubility or hydrolysis characteristics for use as sustained-release or prodrug formulations.

Solvates, and in particular, the hydrates of the compounds of the present invention are included in the present invention as well.

Utilities

5 The compounds of the present invention are potent agonists of the GPR40 receptor. The compounds, and pharmaceutically acceptable salts thereof, may be efficacious in the treatment of diseases that are modulated by GPR40 ligands, which are generally agonists. Many of these diseases are summarized below.

10 One or more of these diseases may be treated by the administration of a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt thereof, to a patient in need of treatment. Also, the compounds of the present invention may be used for the manufacture of a medicament which may be useful for treating one or more of these diseases: (1) non-insulin dependent diabetes mellitus (Type 2 diabetes); (2) hyperglycemia; (3) insulin resistance; (4) Metabolic Syndrome; (5) obesity; (6) hypercholesterolemia; (7) hypertriglyceridemia (elevated levels of triglyceride-rich-lipoproteins); (8) mixed or diabetic dyslipidemia; (9) low HDL cholesterol; (10) high LDL cholesterol; (11) hyperapo-B lipoproteinemia; and (12) atherosclerosis.

15 Preferred uses of the compounds may be for the treatment of one or more of the following diseases by administering a therapeutically effective amount to a patient in need of treatment. The compounds may be used for manufacturing a medicament for the treatment of one or more of these diseases: (1) Type 2 diabetes, and specifically hyperglycemia associated with Type 2 diabetes; (2) Metabolic Syndrome; (3) obesity; and (4) hypercholesterolemia.

20 The compounds may be effective in lowering glucose and lipids in diabetic patients and in non-diabetic patients who have impaired glucose tolerance and/or are in a pre-diabetic condition. The compounds may ameliorate hyperinsulinemia, which often occurs in diabetic or pre-diabetic patients, by modulating the swings in the level of serum glucose that often occurs in these patients. The compounds may also be effective in treating or reducing insulin resistance. The compounds may be effective in treating or preventing gestational diabetes.

25 The compounds may also be effective in treating or preventing lipid disorders. The compounds may be effective in treating or preventing diabetes related disorders. The compounds may also be effective in treating or preventing obesity related disorders.

30 The compounds of this invention may also have utility in improving or restoring β -cell function, so that they may be useful in treating Type 1 diabetes or in delaying or preventing a patient with Type 2 diabetes from needing insulin therapy.

35 The invention also includes pharmaceutically acceptable salts of the compounds, and pharmaceutical compositions comprising the compounds and a pharmaceutically acceptable carrier.

The compounds may be useful in treating insulin resistance, Type 2 diabetes, hyperglycemia, and dyslipidemia that is associated with Type 2 diabetes and insulin resistance. The compounds may also be useful for the treatment of obesity

A compound of the present invention, or a pharmaceutically acceptable salt thereof, may be used in the manufacture of a medicament for the treatment of Type 2 diabetes in a human or other mammalian patient.

A method of treating Type 2 diabetes comprises the administration of a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition comprising the compound, to a human or other mammalian subject or patient in need of treatment. Other medical uses of the compounds of the present invention are described herein.

The term "diabetes," as used herein, includes both insulin-dependent diabetes mellitus (i.e., IDDM, also known as type 1 diabetes) and non-insulin-dependent diabetes mellitus (i.e., NIDDM, also known as Type 2 diabetes). Type 1 diabetes, or insulin-dependent diabetes, is the result of an absolute deficiency of insulin, the hormone which regulates glucose utilization. Type 2 diabetes, or insulin-independent diabetes (i.e., non-insulin-dependent diabetes mellitus), often occurs in the face of normal, or even elevated levels of insulin and appears to be the result of the inability of tissues to respond appropriately to insulin. Most of the Type 2 diabetics are also obese. The compositions of the present invention may be useful for treating both Type 1 and Type 2 diabetes. The term "diabetes associated with obesity" refers to diabetes caused by obesity or resulting from obesity.

Diabetes is characterized by a fasting plasma glucose level of greater than or equal to 126 mg/dl. A diabetic subject has a fasting plasma glucose level of greater than or equal to 126 mg/dl. A pre diabetic subject is someone suffering from prediabetes. Prediabetes is characterized by an impaired fasting plasma glucose (FPG) level of greater than or equal to 110 mg/dl and less than 126 mg/dl; or impaired glucose tolerance; or insulin resistance. A prediabetic subject is a subject with impaired fasting glucose (a fasting plasma glucose (FPG) level of greater than or equal to 110 mg/dl and less than 126 mg/dl); or impaired glucose tolerance (a 2 hour plasma glucose level of ≥ 140 mg/dl and < 200 mg/dl); or insulin resistance, resulting in an increased risk of developing diabetes.

Treatment of diabetes mellitus refers to the administration of a compound or combination of the present invention to treat a diabetic subject. One outcome of treatment may be decreasing the glucose level in a subject with elevated glucose levels. Another outcome of treatment may be decreasing insulin levels in a subject with elevated insulin levels. Another outcome of treatment may be decreasing plasma triglycerides in a subject with elevated plasma triglycerides. Another outcome of treatment is decreasing LDL cholesterol in a subject with high LDL cholesterol levels. Another outcome of treatment may be increasing HDL cholesterol in a subject with low HDL cholesterol levels. Another outcome of treatment is increasing insulin sensitivity. Another outcome

of treatment may be enhancing glucose tolerance in a subject with glucose intolerance. Yet another outcome of treatment may be decreasing insulin resistance in a subject with increased insulin resistance or elevated levels of insulin. Prevention of diabetes mellitus, in particular diabetes associated with obesity, refers to the administration of a compound or combination of the present invention to prevent the onset of diabetes in a subject in need thereof. A subject in need of preventing diabetes is a prediabetic subject that is overweight or obese.

The term "diabetes related disorders" should be understood to mean disorders that are associated with, caused by, or result from diabetes. Examples of diabetes related disorders include retinal damage, kidney disease, and nerve damage.

The term "atherosclerosis" as used herein encompasses vascular diseases and conditions that are recognized and understood by physicians practicing in the relevant fields of medicine. Atherosclerotic cardiovascular disease, coronary heart disease (also known as coronary artery disease or ischemic heart disease), cerebrovascular disease and peripheral vessel disease are all clinical manifestations of atherosclerosis and are therefore encompassed by the terms "atherosclerosis" and "atherosclerotic disease." The combination comprised of a therapeutically effective amount of a GPR40 agonist in combination with a therapeutically effective amount of an anti-hypertensive agent may be administered to prevent or reduce the risk of occurrence, or recurrence where the potential exists, of a coronary heart disease event, a cerebrovascular event, or intermittent claudication. Coronary heart disease events are intended to include CHD death, myocardial infarction (i.e., a heart attack), and coronary revascularization procedures. Cerebrovascular events are intended to include ischemic or hemorrhagic stroke (also known as cerebrovascular accidents) and transient ischemic attacks. Intermittent claudication is a clinical manifestation of peripheral vessel disease. The term "atherosclerotic disease event" as used herein is intended to encompass coronary heart disease events, cerebrovascular events, and intermittent claudication. It is intended that persons who have previously experienced one or more non-fatal atherosclerotic disease events are those for whom the potential for recurrence of such an event exists. The term "atherosclerosis related disorders" should be understood to mean disorders associated with, caused by, or resulting from atherosclerosis.

The term "hypertension" as used herein includes essential, or primary, hypertension wherein the cause is not known or where hypertension is due to greater than one cause, such as changes in both the heart and blood vessels; and secondary hypertension wherein the cause is known. Causes of secondary hypertension include, but are not limited to obesity; kidney disease; hormonal disorders; use of certain drugs, such as oral contraceptives, corticosteroids, cyclosporin, and the like. The term "hypertension" encompasses high blood pressure, in which both the systolic and diastolic pressure levels are elevated (≥ 140 mmHg/ ≥ 90 mmHg), and isolated systolic hypertension, in which only the systolic pressure is elevated to greater than or equal to 140 mm Hg, while the diastolic

pressure is less than 90 mm Hg. Normal blood pressure may be defined as less than 120 mmHg systolic and less than 80 mmHg diastolic. A hypertensive subject is a subject with hypertension. A pre-hypertensive subject is a subject with a blood pressure that is between 120 mmHg over 80 mmHg and 139 mmHg over 89 mmHg. One outcome of treatment is decreasing blood pressure in a subject with high blood pressure. Treatment of hypertension refers to the administration of the compounds and combinations of the present invention to treat hypertension in a hypertensive subject. Treatment of hypertension-related disorder refers to the administration of a compound or combination of the present invention to treat the hypertension-related disorder. Prevention of hypertension, or a hypertension related disorder, refers to the administration of the combinations of the present invention to a pre-hypertensive subject to prevent the onset of hypertension or a hypertension related disorder. The hypertension-related disorders herein are associated with, caused by, or result from hypertension. Examples of hypertension-related disorders include, but are not limited to: heart disease, heart failure, heart attack, kidney failure, and stroke.

Dyslipidemias and lipid disorders are disorders of lipid metabolism including various conditions characterized by abnormal concentrations of one or more lipids (i.e. cholesterol and triglycerides), and/or apolipoproteins (i.e., apolipoproteins A, B, C and E), and/or lipoproteins (i.e., the macromolecular complexes formed by the lipid and the apolipoprotein that allow lipids to circulate in blood, such as LDL, VLDL and IDL). Hyperlipidemia is associated with abnormally high levels of lipids, LDL and VLDL cholesterol, and/or triglycerides. Treatment of dyslipidemia refers to the administration of the combinations of the present invention to a dyslipidemic subject. Prevention of dyslipidemia refers to the administration of the combinations of the present invention to a pre-dyslipidemic subject. A pre-dyslipidemic subject is a subject with higher than normal lipid levels, that is not yet dyslipidemic.

The terms "dyslipidemia related disorders" and "lipid disorder related disorders" should be understood to mean disorders associated with, caused by, or resulting from dyslipidemia or lipid disorders. Examples of dyslipidemia related disorder and lipid disorder related disorders include, but are not limited to: hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, low high density lipoprotein (HDL) levels, high plasma low density lipoprotein (LDL) levels, atherosclerosis and its sequelae, coronary artery or carotid artery disease, heart attack, and stroke.

The term "obesity" as used herein is a condition in which there is an excess of body fat. The operational definition of obesity is based on the Body Mass Index (BMI), which is calculated as body weight per height in meters squared (kg/m^2). "Obesity" refers to a condition whereby an otherwise healthy subject has a Body Mass Index (BMI) greater than or equal to 30 kg/m^2 , or a condition whereby a subject with at least one co-morbidity has a BMI greater than or equal to 27 kg/m^2 . An "obese subject" is an otherwise healthy subject with a Body Mass Index (BMI) greater than or equal to 30 kg/m^2 or a subject with at least one co-morbidity with a BMI greater than or

equal to 27 kg/m^2 . An overweight subject is a subject at risk of obesity. A “subject at risk of obesity” is an otherwise healthy subject with a BMI of 25 kg/m^2 to less than 30 kg/m^2 or a subject with at least one co-morbidity with a BMI of 25 kg/m^2 to less than 27 kg/m^2 .

The increased risks associated with obesity occur at a lower Body Mass Index (BMI) in
5 Asians. In Asian countries, including Japan, “obesity” refers to a condition whereby a subject with at least one obesity-induced or obesity-related co-morbidity, that requires weight reduction or that would be improved by weight reduction, has a BMI greater than or equal to 25 kg/m^2 . In Asian countries, including Japan, an “obese subject” refers to a subject with at least one obesity-induced or obesity-related co-morbidity that requires weight reduction or that would be improved by weight
10 reduction, with a BMI greater than or equal to 25 kg/m^2 . In Asia-Pacific, a “subject at risk of obesity” is a subject with a BMI of greater than 23 kg/m^2 to less than 25 kg/m^2 .

As used herein, the term “obesity” is meant to encompass all of the above definitions of obesity.

Obesity-induced or obesity-related co-morbidities include, but are not limited to, diabetes
15 mellitus, non-insulin dependent diabetes mellitus - type 2, diabetes associated with obesity, impaired glucose tolerance, impaired fasting glucose, insulin resistance syndrome, dyslipidemia, hypertension, hypertension associated with obesity, hyperuricacidemia, gout, coronary artery disease, myocardial infarction, angina pectoris, sleep apnea syndrome, Pickwickian syndrome, fatty liver; cerebral infarction, cerebral thrombosis, transient ischemic attack, orthopedic disorders, arthritis deformans,
20 lumbodinia, emmeniopathy, and infertility. In particular, co-morbidities include: hypertension, hyperlipidemia, dyslipidemia, glucose intolerance, cardiovascular disease, sleep apnea, and other obesity-related conditions.

Treatment of obesity and obesity-related disorders refers to the administration of the compounds of the present invention to reduce or maintain the body weight of an obese subject. One
25 outcome of treatment may be reducing the body weight of an obese subject relative to that subject’s body weight immediately before the administration of the compounds of the present invention. Another outcome of treatment may be preventing body weight regain of body weight previously lost as a result of diet, exercise, or pharmacotherapy. Another outcome of treatment may be decreasing the occurrence of and/or the severity of obesity-related diseases. The treatment may suitably result
30 in a reduction in food or calorie intake by the subject, including a reduction in total food intake, or a reduction of intake of specific components of the diet such as carbohydrates or fats; and/or the inhibition of nutrient absorption; and/or the inhibition of the reduction of metabolic rate; and in weight reduction in patients in need thereof. The treatment may also result in an alteration of metabolic rate, such as an increase in metabolic rate, rather than or in addition to an inhibition of the
35 reduction of metabolic rate; and/or in minimization of the metabolic resistance that normally results from weight loss.

Prevention of obesity and obesity-related disorders refers to the administration of the compounds of the present invention to reduce or maintain the body weight of a subject at risk of obesity. One outcome of prevention may be reducing the body weight of a subject at risk of obesity relative to that subject's body weight immediately before the administration of the compounds of the present invention. Another outcome of prevention may be preventing body weight regain of body weight previously lost as a result of diet, exercise, or pharmacotherapy. Another outcome of prevention may be preventing obesity from occurring if the treatment is administered prior to the onset of obesity in a subject at risk of obesity. Another outcome of prevention may be decreasing the occurrence and/or severity of obesity-related disorders if the treatment is administered prior to the onset of obesity in a subject at risk of obesity. Moreover, if treatment is commenced in already obese subjects, such treatment may prevent the occurrence, progression or severity of obesity-related disorders, such as, but not limited to, arteriosclerosis, Type II diabetes, polycystic ovarian disease, cardiovascular diseases, osteoarthritis, dermatological disorders, hypertension, insulin resistance, hypercholesterolemia, hypertriglyceridemia, and cholelithiasis.

The obesity-related disorders herein are associated with, caused by, or result from obesity. Examples of obesity-related disorders include overeating and bulimia, hypertension, diabetes, elevated plasma insulin concentrations and insulin resistance, dyslipidemias, hyperlipidemia, endometrial, breast, prostate and colon cancer, osteoarthritis, obstructive sleep apnea, cholelithiasis, gallstones, heart disease, abnormal heart rhythms and arrhythmias, myocardial infarction, congestive heart failure, coronary heart disease, sudden death, stroke, polycystic ovarian disease, craniopharyngioma, the Prader-Willi Syndrome, Frohlich's syndrome, GH-deficient subjects, normal variant short stature, Turner's syndrome, and other pathological conditions showing reduced metabolic activity or a decrease in resting energy expenditure as a percentage of total fat-free mass, e.g, children with acute lymphoblastic leukemia. Further examples of obesity-related disorders are metabolic syndrome, also known as syndrome X, insulin resistance syndrome, sexual and reproductive dysfunction, such as infertility, hypogonadism in males and hirsutism in females, gastrointestinal motility disorders, such as obesity-related gastro-esophageal reflux, respiratory disorders, such as obesity-hypoventilation syndrome (Pickwickian syndrome), cardiovascular disorders, inflammation, such as systemic inflammation of the vasculature, arteriosclerosis, hypercholesterolemia, hyperuricaemia, lower back pain, gallbladder disease, gout, and kidney cancer. The compounds of the present invention are also useful for reducing the risk of secondary outcomes of obesity, such as reducing the risk of left ventricular hypertrophy.

The term "metabolic syndrome", also known as syndrome X, is defined in the Third Report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III, or ATP III), National Institutes of Health, 2001, NIH Publication No. 01-3670. E.S. Ford et al., JAMA, vol. 287 (3), Jan.

16, 2002, pp 356-359. Briefly, a person is defined as having metabolic syndrome if the person has three or more of the following disorders: abdominal obesity, hypertriglyceridemia, low HDL cholesterol, high blood pressure, and high fasting plasma glucose. The criteria for these are defined in ATP-III. Treatment of metabolic syndrome refers to the administration of the combinations of the present invention to a subject with metabolic syndrome. Prevention of metabolic syndrome refers to the administration of the combinations of the present invention to a subject with two of the disorders that define metabolic syndrome. A subject with two of the disorders that define metabolic syndrome is a subject that has developed two of the disorders that define metabolic syndrome, but has not yet developed three or more of the disorders that define metabolic syndrome.

The terms "administration of" and or "administering a" compound should be understood to mean providing a compound of the invention or a prodrug of a compound of the invention to a human or other mammal in need of treatment.

The term "patient" should be understood to mean a human or other mammal in need of treatment.

The administration of the compound of structural formula I in order to practice the present methods of therapy is carried out by administering a therapeutically effective amount of the compound of structural formula I to the mammal (human or other mammal) in need of such treatment or prophylaxis. The need for a prophylactic administration according to the methods of the present invention is determined via the use of well known risk factors. The therapeutically effective amount of an individual compound is determined, in the final analysis, by the physician or veterinarian in charge of the case, but depends on factors such as the exact disease to be treated, the severity of the disease and other diseases or conditions from which the patient suffers, the chosen route of administration other drugs and treatments which the patient may concomitantly require, and other factors in the physician's judgment.

The usefulness of the present compounds in these diseases or disorders may be demonstrated in animal disease models that have been reported in the literature.

Administration and Dose Ranges

Any suitable route of administration may be employed for providing a mammal, especially a human, with a therapeutically effective dose of a compound of the present invention. For example, oral, rectal, topical, parenteral, ocular, pulmonary, nasal, and the like may be employed. Dosage forms include tablets, troches, dispersions, suspensions, solutions, capsules, creams, ointments, aerosols, and the like. Preferably compounds of the present invention are administered orally.

In the treatment or prevention of conditions which require agonism of GPR40 receptor activity, an appropriate dosage level will generally be about 0.01 to 500 mg per kg patient body weight per day which may be administered in single or multiple doses. Preferably, the dosage level

will be about 0.1 to about 250 mg/kg per day; more preferably about 0.5 to about 100 mg/kg per day. A suitable dosage level may be about 0.01 to 250 mg/kg per day, about 0.05 to 100 mg/kg per day, or about 0.1 to 50 mg/kg per day. Within this range the dosage may be 0.05 to 0.5, 0.5 to 5 or 5 to 50 mg/kg per day. For oral administration, the compositions may preferably be provided in the form of tablets containing 1.0 to 1000 mg of the active ingredient, particularly 1.0, 5.0, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 100.0, 150.0, 200.0, 250.0, 300.0, 400.0, 500.0, 600.0, 750.0, 800.0, 900.0, and 1000.0 mg of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. The compounds may be administered on a regimen of 1 to 4 times per day, preferably once or twice per day.

When treating or preventing diabetes mellitus and/or hyperglycemia or hypertriglyceridemia or other diseases for which compounds of the present invention may be indicated, generally satisfactory results could be obtained when the compounds of the present invention are administered at a daily dosage of from about 0.1 mg to about 100 mg per kilogram of animal body weight, preferably given as a single daily dose or in divided doses two to six times a day, or in sustained release form. For most large mammals, the total daily dosage is from about 1.0 mg to about 1000 mg, preferably from about 1 mg to about 50 mg. In the case of a 70 kg adult human, the total daily dose will generally be from about 7 mg to about 350 mg. This dosage regimen may be adjusted to provide the optimal therapeutic response.

It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the patient undergoing therapy.

The compounds of this invention may be used in pharmaceutical compositions comprising (a) the compound(s) or a pharmaceutically acceptable salt thereof, and (b) a pharmaceutically acceptable carrier. The compounds of this invention may be used in pharmaceutical compositions that include one or more other active pharmaceutical ingredients. The compounds of this invention may also be used in pharmaceutical compositions in which the compound of the present invention or a pharmaceutically acceptable salt thereof is the only active ingredient.

The term "composition," as in pharmaceutical composition, is intended to encompass a product comprising the active ingredient(s), and the inert ingredient(s) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the present invention encompass any composition made by admixing a compound of the present invention and a pharmaceutically acceptable carrier.

Combination Therapy

The compounds of the present invention may be useful in methods for the prevention or treatment of the aforementioned diseases, disorders and conditions in combination with other therapeutic agents.

The compounds of the present invention may be useful in combination with one or more other drugs in the treatment, prevention, suppression or amelioration of diseases or conditions for which compounds of formula I or the other drugs may have utility, where the combination of the drugs together are safer, more effective or more therapeutically effective than either drug alone. Such other drug(s) may be administered, by a route and in an amount commonly used therefore, contemporaneously or sequentially with a compound of formula I. When a compound of formula I is used contemporaneously with one or more other drugs, a pharmaceutical composition in unit dosage form containing such other drugs and the compound of formula I is preferred. However, the combination therapy may also include therapies in which the compound of formula I and one or more other drugs are administered on different overlapping schedules. It is also contemplated that when used in combination with one or more other active ingredients, the compounds of the present invention and the other active ingredients may be used in lower doses than when each is used singly. Accordingly, the pharmaceutical compositions of the present invention include those that contain one or more other active ingredients, in addition to a compound of formula I.

Examples of other active ingredients that may be administered separately or in the same pharmaceutical composition in combination with a compound of the formulas described herein include, but are not limited to: (1) dipeptidyl peptidase-IV (DPP-4) inhibitors (e.g., sitagliptin, omarigliptin, trelagliptin, teneligliptin, bisegliptin, anagliptin, vildagliptin, saxagliptin, alogliptin, melogliptin, linagliptin, gosogliptin, evogliptin, and gemigliptin), (2) insulin sensitizers, including (i) PPAR γ agonists, such as the glitazones (e.g. pioglitazone, AMG 131, MBX2044, mitoglitazone, lobeglitazone, IDR-105, rosiglitazone, and balaglitazone), and other PPAR ligands, including (1) PPAR α/γ dual agonists (e.g., ZYH2, ZYH1, GFT505, chiglitazar, muraglitazar, aleglitazar, sodelglitazar, and naveglitazar); (2) PPAR α agonists such as fenofibric acid derivatives (e.g., gemfibrozil, clofibrate, ciprofibrate, fenofibrate, bezafibrate), (3) selective PPAR γ modulators (SPPAR γ M's), (e.g., such as those disclosed in WO 02/060388, WO 02/08188, WO 2004/019869, WO 2004/020409, WO 2004/020408, and WO 2004/066963); and (4) PPAR γ partial agonists; (ii) biguanides, such as metformin and its pharmaceutically acceptable salts, in particular, metformin hydrochloride, and extended-release formulations thereof, such as GlumetzaTM, FortametTM, and GlucophageXRTM; and (iii) protein tyrosine phosphatase-1B (PTP-1B) inhibitors (e.g., ISIS-113715 and TTP814); (3) insulin or insulin analogs (e.g., insulin detemir, insulin glulisine, insulin

degludec, insulin glargine, insulin lispro and inhalable formulations of each); (4) leptin and leptin derivatives and agonists; (5) amylin and amylin analogs (e.g., pramlintide); (6) sulfonylurea and non-sulfonylurea insulin secretagogues (e.g., tolbutamide, glyburide, glipizide, glimepiride, mitiglinide, meglitinides, nateglinide and repaglinide); (7) α -glucosidase inhibitors (e.g., acarbose, voglibose and miglitol); (8) glucagon receptor antagonists (e.g., MK-3577, MK-0893, LY-2409021 and KT6-971); (9) incretin mimetics, such as GLP-1, GLP-1 analogs, derivatives, and mimetics; and GLP-1 receptor agonists (e.g., dulaglutide, semaglutide, albiglutide, exenatide, liraglutide, lixisenatide, taspoglutide, CJC-1131, and BIM-51077, including intranasal, transdermal, and once-weekly formulations thereof); (10) LDL cholesterol lowering agents such as (i) HMG-CoA reductase inhibitors (e.g., simvastatin, lovastatin, pravastatin, crivastatin, fluvastatin, atorvastatin, pitavastatin and rosuvastatin), (ii) bile acid sequestering agents (e.g., colestilan, colestimide, colesevalam hydrochloride, colestipol, cholestyramine, and dialkylaminoalkyl derivatives of a cross-linked dextran), (iii) inhibitors of cholesterol absorption, (e.g., ezetimibe), and (iv) acyl CoA:cholesterol acyltransferase inhibitors, (e.g., avasimibe); (11) HDL-raising drugs, (e.g., niacin and nicotinic acid receptor agonists, and extended-release versions thereof; (12) antiobesity compounds; (13) agents intended for use in inflammatory conditions, such as aspirin, non-steroidal anti-inflammatory drugs or NSAIDs, glucocorticoids, and selective cyclooxygenase-2 or COX-2 inhibitors; (14) antihypertensive agents, such as ACE inhibitors (e.g., lisinopril, enalapril, ramipril, captopril, quinapril, and tandolapril), A-II receptor blockers (e.g., losartan, candesartan, irbesartan, olmesartan, medoxomil, valsartan, telmisartan, and eprosartan), renin inhibitors (e.g., aliskiren), beta blockers, and calcium channel blockers; (15) glucokinase activators (GKAs) (e.g., AZD6370); (16) inhibitors of 11β -hydroxysteroid dehydrogenase type 1, (e.g., such as those disclosed in U.S. Patent No. 6,730,690, and LY-2523199); (17) CETP inhibitors (e.g., anacetrapib, evacetrapib, torcetrapib, and AT-03); (18) inhibitors of fructose 1,6-bisphosphatase, (e.g., MB-07803, and such as those disclosed in U.S. Patent Nos. 6,054,587; 6,110,903; 6,284,748; 6,399,782; and 6,489,476); (19) inhibitors of acetyl CoA carboxylase-1 or 2 (ACC1 or ACC2); (20) AMP-activated Protein Kinase (AMPK) activators (e.g., MB-11055); (21) other agonists of the G-protein-coupled receptors: (i) GPR-109, (ii) GPR-119 (e.g., MBX2982 and PSN821), (iii) GPR-40 (e.g., fasiglifam, JTT-851, TAK-875, and P-11187, and (iv) GPR-120 (e.g., KDT-501); (22) SSTR3 antagonists (e.g., pasireotide, and such as those disclosed in WO 2009/011836); (23) neuromedin U receptor agonists (e.g., such as those disclosed in WO 2009/042053, including, but not limited to, neuromedin S (NMS)); (24) SCD inhibitors; (25) GPR-105 antagonists (e.g., such as those disclosed in WO 2009/000087); (26) SGLT inhibitors (e.g., LIK-066, ASP1941, SGLT-3, ertugliflozin, empagliflozin, dapagliflozin, canagliflozin, BI-10773, PF-04971729, remogliflozin, luseogliflozin, tofogliflozin,

ipragliflozin, and LX-4211); (27) inhibitors of (i) acyl coenzyme A:diacylglycerol acyltransferase 1, DGAT-1 (e.g., pradigastat, and P-7435) and acyl coenzyme A:diacylglycerol acyltransferase 2, DGAT-2; (28) inhibitors of fatty acid synthase; (29) inhibitors of acyl coenzyme A:monoacylglycerol acyltransferase 1 and 2 (MGAT-1 and MGAT-2); (30) agonists of the TGR5 receptor (also known as GPBAR1, BG37, GPCR19, GPR131, and M-BAR) (eg., sodium taurocholate); (31) ileal bile acid transporter inhibitors (eg., elobixibat); (32) PACAP, PACAP mimetics, and PACAP receptor 3 agonists; (33) PPAR agonists; (34) protein tyrosine phosphatase-1B (PTP-1B) inhibitors; (35) IL-1b antibodies and inhibitors, (e.g., gevokizumab, canakinumab, danazol, AC-201, and BLX-1002); and (36) bromocriptine mesylate and rapid-release formulations thereof.

Of particular interest are dipeptidyl peptidase-IV (DPP-4) inhibitors that may be useful in combination with compounds of the present invention. Such inhibitors include, without limitation, sitagliptin (disclosed in US Patent No. 6,699,871), omarigliptin, trelagliptin, teneligliptin, bisegliptin, anagliptin, LC15-0444, vildagliptin, saxagliptin, alogliptin, melogliptin, linagliptin, gosogliptin, evogliptin, gemigliptin, and pharmaceutically acceptable salts thereof, and fixed-dose combinations of these compounds with metformin hydrochloride, pioglitazone, rosiglitazone, simvastatin, atorvastatin, or a sulfonylurea.

Other GPR-40 agonists that may be useful in combination with compounds of the formulas described herein include, but are not limited to: (1) 5-[4-[[[(1R)-4-[6-(3-hydroxy-3-methylbutoxy)-2-methylpyridine-3-yl]-2,3-dihydro-1H-indene-1-yl]oxy]phenyl]isothiazole-3-ol 1-oxide; (2) 5-(4-((3-(2,6-dimethyl-4-(3-(methylsulfonyl)propoxy)phenyl)phenyl)-methoxy)-phenyl)isothiazole-3-ol 1-oxide; and (3) 5-(4-((3-(2-methyl-6-(3-hydroxypropoxy)-pyridine-3-yl)-2-methylphenyl)methoxy)-phenyl)isothiazole-3-ol 1-oxide; and (4) 5-[4-[[[3-[4-(3-aminopropoxy)-2,6-dimethylphenyl]phenyl]-methoxy]phenyl]isothiazole-3-ol 1-oxide, and pharmaceutically acceptable salts thereof.

Antiobesity compounds that may be combined with compounds of formula I include topiramate; zonisamide; naltrexone; phentermine; bupropion; the combination of bupropion and naltrexone; the combination of bupropion and zonisamide; the combination of topiramate and phentermine; fenfluramine; dexfenfluramine; sibutramine; lipase inhibitors, such as orlistat and cetilistat; melanocortin receptor agonists, in particular, melanocortin-4 receptor agonists; CCK-1 agonists; melanin-concentrating hormone (MCH) receptor antagonists; neuropeptide Y₁ or Y₅ antagonists (such as MK-0557); β_3 adrenergic receptor agonists; CB-1 receptor inverse agonists and antagonists; ghrelin antagonists; bombesin receptor agonists (such as bombesin receptor subtype-3 agonists); and 5-hydroxytryptamine-2c (5-HT_{2c}) agonists, such as lorcaserin. For a review of anti-obesity compounds that may be useful in combination with a compound of the present invention, see S. Chaki et al., "Recent advances in feeding suppressing agents: potential therapeutic strategy for

the treatment of obesity,” Expert Opin. Ther. Patents, 11: 1677-1692 (2001); D. Spanswick and K. Lee, “Emerging antiobesity drugs,” Expert Opin. Emerging Drugs, 8: 217-237 (2003); J.A. Fernandez-Lopez, et al., “Pharmacological Approaches for the Treatment of Obesity,” Drugs, 62: 915-944 (2002); and K.M. Gadde, et al., “Combination pharmaceutical therapies for obesity,” Exp. Opin. Pharmacother., 10: 921-925 (2009).

Glucagon receptor antagonists that may be useful in combination with the compounds of formula I include, but are not limited to: (1) *N*-[4-((1*S*)-1-{3-(3,5-dichlorophenyl)-5-[6-(trifluoromethoxy)-2-naphthyl]-1*H*-pyrazol-1-yl}ethyl)benzoyl]-β-alanine; (2) *N*-[4-((1*R*)-1-{3-(3,5-dichlorophenyl)-5-[6-(trifluoromethoxy)-2-naphthyl]-1*H*-pyrazol-1-yl}ethyl)benzoyl]-β-alanine; (3) *N*-(4-{1-[3-(2,5-dichlorophenyl)-5-(6-methoxy-2-naphthyl)-1*H*-pyrazol-1-yl]ethyl}benzoyl)-β-alanine; (4) *N*-(4-{(1*S*)-1-[3-(3,5-dichlorophenyl)-5-(6-methoxy-2-naphthyl)-1*H*-pyrazol-1-yl]ethyl}benzoyl)-β-alanine; (5) *N*-(4-{(1*S*)-1-[(*R*)-(4-chlorophenyl)(7-fluoro-5-methyl-1*H*-indol-3-yl)methyl]butyl}benzoyl)-β-alanine; and (6) *N*-(4-{(1*S*)-1-[(4-chlorophenyl)(6-chloro-8-methylquinolin-4-yl)methyl]butyl}benzoyl)-β-alanine; and pharmaceutically acceptable salts thereof.

Another embodiment of the present invention relates to a pharmaceutical composition comprising one or more of the following agents: (a) a compound of structural formula I; (b) one or more compounds selected from the group consisting of: (1) dipeptidyl peptidase-IV (DPP-4) inhibitors (e.g., sitagliptin, omarigliptin, trelagliptin, teneligliptin, bisegliptin, anagliptin, vildagliptin, saxagliptin, alogliptin, melogliptin, linagliptin, gosogliptin, evogliptin, and gemigliptin); (2) insulin sensitizers, including (i) PPARγ agonists, such as the glitazones (e.g. AMG 131, MBX2044, mitoglitazone, lobeglitazone, IDR-105, pioglitazone, rosiglitazone, and balaglitazone) and other PPAR ligands, including (1) PPARα/γ dual agonists, such as ZYH1, YYH2, chiglitazar, GFT505, muraglitazar, aleglitazar, sodelglitazar, and naveglitazar, (2) PPARα agonists, such as fenofibric acid derivatives (e.g., gemfibrozil, clofibrate, ciprofibrate, fenofibrate and bezafibrate), (3) selective PPARγ modulators (SPPARγM's), (4) PPARγ partial agonists; (ii) biguanides, such as metformin and its pharmaceutically acceptable salts, in particular, metformin hydrochloride, and extended-release formulations thereof, such as Glumetza®, Fortamet®, and GlucophageXR®; (iii) protein tyrosine phosphatase-1B (PTP-1B) inhibitors, such as ISI-113715, and TTP814; (3) sulfonylurea and non-sulfonylurea insulin secretagogues, (e.g., tolbutamide, glyburide, glipizide, glimepiride, mitiglinide, and meglitinides, such as nateglinide and repaglinide); (4) α-glucosidase inhibitors (e.g., acarbose, voglibose and miglitol); (5) glucagon receptor antagonists; (6) LDL cholesterol lowering agents such as (i) HMG-CoA reductase inhibitors (e.g., lovastatin, simvastatin, pravastatin, cerivastatin, fluvastatin, atorvastatin, pitavastatin, and rosuvastatin), (ii) bile acid sequestering agents (e.g., colestilan, cholestyramine, colestimide, colesevelam hydrochloride,

colestipol, and dialkylaminoalkyl derivatives of a cross-linked dextran), (iii) inhibitors of cholesterol absorption, (e.g., ezetimibe), and (iv) acyl CoA:cholesterol acyltransferase inhibitors (e.g., avasimibe); (7) HDL-raising drugs, such as niacin or a salt thereof and extended-release versions thereof; and nicotinic acid receptor agonists; (8) antiobesity compounds; (9) agents intended for use
5 in inflammatory conditions, such as aspirin, non-steroidal anti-inflammatory drugs (NSAIDs), glucocorticoids, and selective cyclooxygenase-2 (COX-2) inhibitors; (10) antihypertensive agents, such as ACE inhibitors (e.g., enalapril, lisinopril, ramipril, captopril, quinapril, and tandolapril), A-II receptor blockers (e.g., losartan, candesartan, irbesartan, olmesartan medoxomil, valsartan, telmisartan, and eprosartan), renin inhibitors (e.g., aliskiren), beta blockers (e.g., calcium channel
10 blockers); (11) glucokinase activators (GKAs) (e.g., AZD6370, GKM-001, TMG-123, HMS-5552, DS-7309, PF-04937319, TTP-399, ZYGK-1); (12) inhibitors of 11 β -hydroxysteroid dehydrogenase type 1 (e.g., such as those disclosed in U.S. Patent No. 6,730,690; WO 03/104207; and WO 04/058741); (13) inhibitors of cholesteryl ester transfer protein (CETP), (e.g., torcetrapib, evacetrapib, anacetrapib, and AT-03); (14) inhibitors of fructose 1,6-bisphosphatase (e.g., MB-
15 07803, and such as those disclosed in U.S. Patent Nos. 6,054,587; 6,110,903; 6,284,748; 6,399,782; and 6,489,476); (15) inhibitors of acetyl CoA carboxylase-1 or 2 (ACC1 or ACC2); (16) AMP-activated Protein Kinase (AMPK) activators (e.g., MB-11055); (17) agonists of the G-protein-coupled receptors: (i) GPR-109, (ii) GPR-119 (e.g., MBX2982, and PSN821), (iii) GPR-40 (e.g., fasiglifam, JTT-851, P-11187, 5-[4-[[[(1R)-4-[6-(3-hydroxy-3-methylbutoxy)-2-methylpyridine-3-yl]-2,3-dihydro-1H-indene-1-yl]oxy]phenyl]isothiazole-3-ol 1-oxide, 5-(4-((3-(2,6-dimethyl-4-(3-(methylsulfonyl)propoxy)phenyl)phenyl)-methoxy)phenyl)-isothiazole-3-ol 1-oxide, 5-(4-((3-(2-methyl-6-(3-hydroxypropoxy)pyridine-3-yl)-2-methyl-phenyl)methoxy)phenyl)-isothiazole-3-ol 1-oxide, and 5-[4-[[3-[4-(3-aminopropoxy)-2,6-dimethylphenyl]phenyl]-methoxy]phenyl]isothiazole-3-ol 1-oxide), and (iv) GPR-120 (e.g., KDT-501); (18) SSTR3 antagonists (e.g., pasireotide, and
25 such as those disclosed in WO 2009/011836); (19) neuromedin U receptor agonists (e.g., such as those disclosed in WO2009/042053, including, but not limited to, neuromedin S (NMS)); (20) inhibitors of stearoyl-coenzyme A delta-9 desaturase (SCD); (21) GPR-105 antagonists (e.g., such as those disclosed in WO 2009/000087); (22) inhibitors of glucose uptake, such as sodium-glucose transporter (SGLT) inhibitors and its various isoforms, such as SGLT-1; SGLT-2 (e.g., LIK-066, ertugliflozin, ASP1941, luseogliflozin, BI10773, tofogliflozin, LX4211, canagliflozin, dapagliflozin and remogliflozin; and SGLT-3); (23) inhibitors of (i) acyl coenzyme A:diacylglycerol
acyltransferase 1, DGAT-1 (e.g., pradigastat, and P-7435) and acyl coenzyme A:diacylglycerol
acyltransferase 2, DGAT-2; (24) inhibitors of fatty acid synthase; (25) inhibitors of acyl coenzyme
30 A:monoacylglycerol acyltransferase 1 and 2 (MGAT-1 and MGAT-2); (26) agonists of the TGR5

receptor (also known as GPBAR1, BG37, GPCR19, GPR131, and M-BAR) (eg., sodium taurocholate); (28) bromocriptine mesylate and rapid-release formulations thereof, and (29) IL-1b antibodies and inhibitors (e.g., gevokizumab, canakinumab, danazol, AC-201, and BLX-1002); and (c) a pharmaceutically acceptable carrier.

5 Specific compounds that may be useful in combination with a compound of the present invention include: simvastatin, mevastatin, ezetimibe, atorvastatin, rosuvastatin, sitagliptin, omarigliptin, metformin, sibutramine, orlistat, topiramate, naltrexone, bupropion, phentermine, losartan, losartan with hydrochlorothiazide, olmesartan, canagliflozin, dapagliflozin, ipragliflozin and ertugliflozin.

10 The above combinations include combinations of a compound of the present invention not only with one other active compound, but also with two or more other active compounds. Non-limiting examples include combinations of compounds with two or more active compounds selected from biguanides, sulfonylureas, HMG-CoA reductase inhibitors, PPAR γ agonists, DPP-4 inhibitors, anti-obesity compounds, and anti-hypertensive agents.

15 The present invention may also provide a method for the treatment or prevention of a G-protein coupled receptor 40 (GPR40) mediated disease, which method comprises administration to a patient in need of such treatment or at risk of developing a GPR40 mediated disease of an amount of a GPR40 agonist and an amount of one or more active ingredients, such that together they give effective relief.

20 In a further aspect of the present invention, there is provided a pharmaceutical composition comprising a GPR40 agonist and one or more active ingredients, together with at least one pharmaceutically acceptable carrier or excipient.

25 Thus, according to a further aspect of the present invention there is provided the use of a GPR40 agonist and one or more active ingredients for the manufacture of a medicament for the treatment or prevention of a GPR40 mediated disease. In a further or alternative aspect of the present invention, there is therefore provided a product comprising a GPR40 agonist and one or more active ingredients as a combined preparation for simultaneous, separate or sequential use in the treatment or prevention of a GPR40 mediated disease. Such a combined preparation may be, for example, in the form of a twin pack.

30 For the treatment or prevention of diabetes, obesity, hypertension, Metabolic Syndrome, dyslipidemia, cancer, atherosclerosis, and related disorders thereof, a compound of the present invention may be used in conjunction with another pharmaceutical agent effective to treat that disorder.

35 The present invention may also provide a method for the treatment or prevention of diabetes, obesity, hypertension, Metabolic Syndrome, dyslipidemia, cancer, atherosclerosis, and related disorders thereof, which method comprises administration to a patient in need of such treatment an

effective amount of a compound of the present invention and an amount of another pharmaceutical agent effective to treat that disorder, such that together they give effective relief.

The present invention may also provide a method for the treatment or prevention of diabetes, obesity, hypertension, Metabolic Syndrome, dyslipidemia, cancer, atherosclerosis, and related disorders thereof, which method comprises administration to a patient in need of such treatment an amount of a compound of the present invention and an amount of another pharmaceutical agent useful in treating that particular condition, such that together they give effective relief.

The term "therapeutically effective amount" or "a therapeutically effective dose" means the amount the compound of structural formula I that will elicit the biological or medical response of a tissue, system, animal, mammal or human that is being sought by the researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disorder being treated. The novel methods of treatment of this invention are for disorders known to those skilled in the art. The term "mammal" includes, but is not limited to, humans, and companion animals such as dogs and cats.

The weight ratio of the compound of the Formula I to the second active ingredient may be varied and will depend upon the effective dose of each ingredient. Generally, a therapeutically effective dose of each will be used. Thus, for example, when a compound of the Formula I is combined with a DPIV inhibitor the weight ratio of the compound of the Formula I to the DPIV inhibitor will generally range from about 1000:1 to about 1:1000, preferably about 200:1 to about 1:200. Combinations of a compound of the Formula I and other active ingredients will generally also be within the aforementioned range, but in each case, an effective dose of each active ingredient should be used.

Methods of Synthesis of the Compounds of the Present Invention:

The following reaction schemes and Examples illustrate methods which may be employed for the synthesis of the compounds of structural formula I described in this invention. Those skilled in the art will readily understand that known variations of protecting groups, as well as of the conditions and processes of the following preparative procedures, can be used to prepare these compounds. It is also understood that whenever a chemical reagent such as a boronic acid or a boronate is not commercially available, such a chemical reagent can be readily prepared following one of numerous methods described in the literature. All temperatures are degrees Celsius unless otherwise noted. Mass spectra (MS) were measured either by electrospray ion-mass spectroscopy (ESMS) or by atmospheric pressure chemical ionization mass spectroscopy (APCI). All temperatures are degrees Celsius unless otherwise noted.


List of Abbreviations

Ac is acetyl; AcCN is acetonitrile; AcO is acetoxy; Alk is alkyl; anh. is anhydrous; APCI is atmospheric pressure chemical ionization; aq or aq. is aqueous; Ar is aryl; atm is atmosphere; Boc is *tert*-butoxycarbonyl; Bn-O is phenyl-CH₂-O or benzyloxy; BrettPhos palladacycle precatalyst is Brettphos Pd G1 precatalyst (Aldrich); *n*-BuLi is *n*-butyl lithium; Bu₃P is tributylphosphine; *t*-buxhos palladacycle is chloro(2-di-*t*-butylphosphino-2',4',6'-tri-*i*-propyl-1,1'-biphenyl)[2-(2-aminoethyl)phenyl]palladium(II), min. 98% [t-BuXPhos Palladacycle]; *t*-BuOK is potassium *tert*-butoxide; °C is degrees celsius; Cataxium precatalyst or Cataxium Pd precat or precatalyst is cataCXium A Pd G3 (Aldrich); Cbz is benzyloxycarbonyl; conc or conc. is concentrated; CV is column volumes; DAST is (diethylamino)sulfur trifluoride; DIBAL-H ix diisobutylaluminum hydride; DIAD is diisopropyl azodicarboxylate; DCM is dichloromethane; DEA is diethyl amine; DIPEA is *N,N*-diisopropylethylamine; DIPA is diisopropyl amine; DMAP is 4-dimethylaminopyridine; DMF is *N,N*-dimethylformamide; DMS is dimethyl sulfide; DMSO is dimethylsulfoxide; dppf is 1,1'-Bis(diphenyl-phosphino)ferrocene; ESI is electrospray ionization; EA or EtOAc is ethyl acetate; Et is ethyl; Et₂O is diethyl ether; EtMgBr is ethyl magnesium bromide; EtOH is ethanol; g is gram(s); h or hr or hrs is hour(s); hex is hexanes; HPLC is high pressure liquid chromatography; HOAc or AcOH is acetic acid; kg is kilogram(s); IPA or *i*-PrOH is isopropanol; KOAc is potassium acetate; K₂CO₃ is potassium carbonate; K₂Cr₂O₇ is potassium dichromate; K₂Cr₂O₇ is potassium dichromate; K₂Cr₂O₇ is potassium dichromate; L is liter; LAH is lithium aluminum hydride; M is molar; LC-MS is liquid chromatography-mass spectroscopy; LDA is lithium diisopropyl amide; Me is methyl; MeO is methoxy; m-CPBA, MCPBA, or mCPBA is meta chloroperbenzoic acid; ml or mL is milliliter; min or mins is minute(s); mol is mole(s); mmol is mmole(s); mg is milligram(s); MeMgBr is methyl magnesium bromide; MeOH is methyl alcohol or methanol; MPLC is medium pressure liquid chromatography; MS is mass spectroscopy; MsCl or Ms-Cl is methane sulfonyl chloride; MeCN is acetonitrile; MeI is methyl iodide; MsCl is methane sulfonyl chloride; MTBE is methyl *tert*-butyl ether; MW is molecular weight; NaHMDS is sodium hexamethyl disilazide; NH₄OAc is ammonium acetate; NBS is *N*-bromo succinamide; NEt₃ is triethyl amine; NIS is *N*-iodo succinamide; NMO is 4-methyl morpholine *N*-oxide; NMP is 1-methyl-2-pyrrolidinone; NMR is nuclear magnetic resonance spectroscopy; o.n. or ON is overnight; PE is petroleum ether; PG is protecting group; Pd₂(dba)₃ is tris(dibenzylidene-acetone)-dipalladium(0); Pd(OAc)₂ is palladium acetate; Pd[P(*t*-Bu)₃]₂ is bis(tri-*tert*-butylphosphine)-palladium (0); Pd(dppf)Cl₂ is [1,1'-bis(diphenyl-phosphino)-ferrocene]dichloro-palladium (II); PdCl₂ (dppf)₂CH₂Cl₂ is [1,1'-Bis(diphenyl-phosphino)-ferrocene]dichloropalladium(II), complex with dichloromethane (Aldrich); Pd(PPh₃)₄ is tetrakis or tetrakis(triphenylphosphine)palladium (0); PPh₃ is triphenyl phosphine; Pd(*t*-Bu₂P)₂ .FerrCl₂ is bis-tri-*tert*-butylphosphino ferrocene dichloro palladium (II); precat is precatalyst; prep is preparative; prep. TLC or prep-TLC, or prep TCL is preparative thin layer chromatography; rt or r.t. or RT is room temperature; RuCl[(R,R)-TSDPEN](mesitylene) is [*N*-[(1*R*,2*R*)-2-(Amino-κ*N*)-1,2-diphenylethyl]-4-

methylbenzenesulfonamidato- κN]chloro-[(1,2,3,4,5,6- η)-1,3,5-trimethylbenzene]-ruthenium; Ru-Josiphos is generated using (Me-allyl) $_2$ Ru(COD) (Aldrich) and Josiphos SL-J502-2 (Aldrich); R_f is retention factor; sat or sat. is saturated; SEM is trimethylsilyl ethoxy methyl, SEMCl is trimethylsilyl ethoxy methyl chloride; SFC is supercritical fluid chromatography; S-Phos is 2-dicyclohexylphosphino-2',6'-dimethoxy-biphenyl; S-Phos(Pd) is chloro(2-dicyclohexylphosphino-2',6'-dimethoxy-1,1'-biphenyl)[2-(2-aminoethylphenyl)]-palladium(II) [CAS-No. 1028206-58-7]; S-Phos precatalyst is S-Phos Pd G2 precatalyst - Aldrich; S-Phos second generation precatalyst is Chloro(2-dicyclohexyl-phosphino-2',6'-dimethoxy-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium-(II), SPhos-Pd-G2 [CAS-No. 1375325-64-6]; TBAF is tetrabutylammonium fluoride; TBSCl is *tert*-butyl dimethylsilyl chloride; TEA is triethyl amine; Tf is trifluoromethane sulfonyl; THF is tetrahydrofuran; Ti(OiPr) $_4$ is titanium isopropoxide; TFA is trifluoroacetic acid; TLC is thin-layer chromatography; TosCl and TsCl is *p*-toluene sulfonyl chloride; pTSA, pTsOH and TsOH is *p*-toluenesulfonic acid, Ts $_2$ O is tosic anhydride or *p*-toluene sulfonic anhydride; and xphos is 2-dicyclohexylphosphino-2',4',6'-triisopropyl-biphenyl.

Several methods for preparing the compounds of this invention are illustrated in the following Schemes and Examples. Starting materials are either commercially available or made by known procedures in the literature or as illustrated. The present invention further provides processes for the preparation of compounds of structural formula I as defined above. In some cases the order of carrying out the foregoing reaction schemes may be varied to facilitate the reaction or to avoid unwanted reaction products. The following Schemes and Examples are provided to illustrate the invention and are not to be construed as limiting the invention in any manner. The scope of the invention is defined by the appended claims.

SCHEME 1




Step 1: MeI, K $_2$ CO $_3$, THF; **Step 2:** NaHMDS, Ts $_2$ O, MTBE; **Step 3:** Cataxium-Pd precat, K $_2$ PO $_3$, MeCN; **Step 4:** Ru-Josiphos, BF $_4$ H-Et $_2$ O, H $_2$ gas, MeOH; **Step 5:** Pd/Pt/C, H $_2$ gas, MeOH

Scheme 1 provides a route to Intermediate 1. The cyclopropyl β -ketoester was methylated and then trapped at the vinyl tosylate. Suzuki cross-coupling with *m*-benzyloxy phenylboronic acid delivered

the "Z" enoate. Asymmetric reduction of the double bond was followed by debenzylation under hydrogen pressure.

INTERMEDIATE 1

(2*R*,3*R*)-Methyl 3-cyclopropyl-3-(3-hydroxyphenyl)-2-methylpropanoate



Step 1: methyl 3-cyclopropyl-2-methyl-3-oxopropanoate

A slurry of NaH (0.563 g, 60% in mineral oil) in THF (20 mL) was cooled to 10 °C, and methyl 3-cyclopropyl-3-oxopropanoate (2 g) was added in portions at room temperature over 30 min. Then MeI (0.88 mL) was added over 5 min at room temperature and the reaction was stirred at room temperature overnight. The reaction was then reverse quenched into a half saturated sodium bicarbonate solution (100 mL) and EtOAc (100 mL). The aqueous layer was separated and extracted once with 50 mL EtOAc. The organic layer was washed with water and brine, and then concentrated to give the title compound, which was used in the next the step without further purification.

Step 2: (Z)-methyl 3-cyclopropyl-2-methyl-3-(tosyloxy)acrylate

To a solution of methyl 3-cyclopropyl-2-methyl-3-oxopropanoate (5 g) in MTBE (50 mL) was added a 1M solution of NaHMDS in hexane (41 mL), while keeping the internal temperature between 18-23 °C. The reaction was stirred at room temperature for 30 min. Then tosic anhydride (10g) was slurried in MTBE (200 mL). The reaction mixture was added to the tosic anhydride slurry, while keeping the internal temperature between 19 °C and 24 °C. After 30 min, additional tosic anhydride (3.3 g, 0.3 eq) was added, and the reaction was stirred for 1h. Then water (500 mL) and EtOAc (400 mL) were added to the reaction mixture. The aqueous layer was separated and extracted once with EtOAc (100 mL) The organic layer was separated, washed with water (200 mL) and brine (100 mL), and concentrated to give the crude product, which was re-crystallized from MTBE/heptanes (1:1, 50 mL) to give the title compound.

Step 3: (Z)-methyl 3-(3-(benzyloxy)phenyl)-3-cyclopropyl-2-methylacrylate

(Z)-methyl 3-cyclopropyl-2-methyl-3-(tosyloxy)acrylate (1.03 g) was dissolved in MeCN (10mL), then aqueous potassium phosphate (1 M, 10 mL) was added, followed by the addition of (3-(benzyloxy)phenyl)boronic acid (1 g). The resulting slurry was degassed with a nitrogen stream for 30 min, then cataCXium A Pd G3 (Aldrich # 761435, 100 mg) was added and the reaction was heated to 35 °C for 14 h. The resulting slurry was filtered through Celite™, and the Celite™ was washed with EtOAc (20 mL). Then EtOAc (40 mL) and water (50 mL) were added to the filtrate. The aqueous layer was separated, and extracted once with EtOAc (20 mL). The combined organic

layer was washed with water (50 mL) and brine (20 mL), and concentrated to give a crude oil, which was purified via ISCO silica column (40 g, diluted with 0-30% Hexanes/EtOAc) to give the title product.


Step 4: (2S,3R)-methyl 3-(3-(benzyloxy)phenyl)-3-cyclopropyl-2-methylpropanoate

5 Bis(2-methylallyl)(1,5-cyclooctadiene)ruthenium(II) (1.0 g) and (S)-1-[(R)-2-(Di-tert-butylphosphino)ferrocenyl]ethyldiphenylphosphine (Josiphos SL-J502-2, CAS# 223121-01-5, 1.86 g) were added to DCM (12 mL) and agitated for 20 min at rt. Then tetrafluoroboric acid-diethyl ether complex (1.0 g) was added slowly and stirred for 20 min at rt. The reaction mixture was diluted with DCM (100 mL) and added to a catalyst bomb with a MeOH rinse. Then (Z)-methyl 3-(3-(benzyloxy)phenyl)-3-cyclopropyl-2-methylacrylate (2.57 g, 7.97 mmol) was added to the catalyst bomb with MeOH (200 mL) and agitated to dissolve. The catalyst bomb was pressurized to 500 psi with hydrogen, then heated to 80 °C and shaken for 20 h. Then the reaction was cooled, filtered through Celite™ and washed with MeOH. The filtrate was concentrated to give the crude product.

Step 5: (2S,3R)-methyl 3-cyclopropyl-3-(3-hydroxyphenyl)-2-methylpropanoate


15 To a solution of (2S,3S)-methyl 3-(3-(benzyloxy)phenyl)-3-cyclopropyl-2-methylpropanoate (4.81 g, 13 mmol) in methanol (67 ml) in a 25 ml glass shaker vessel, was 8% Pd-2% Pt/C (1.5 g, 35 wt % loading, 50 w/w, Johnson Matthey lot# F27N23). Then the vessel was evacuated and purged with nitrogen three times, and charged with 50 psi of hydrogen gas. The reaction was heated to 25 °C for 6 h, then filtered through Celite™, which was washed with MeOH (50 mL). The filtrate is
20 concentrated to give the title compound.

INTERMEDIATE 2
(2S,3R)-methyl 3-cyclopropyl-3-(3-hydroxy-4-iodophenyl)-2-methylpropanoate




25 A solution of (2S,3R)-methyl 3-cyclopropyl-3-(3-hydroxyphenyl)-2-methylpropanoate (1 g, 4.27 mmol) in DCM (10 mL) was cooled to 0°C, then NIS (0.960 g, 4.27 mmol) was added slowly over several minutes. After 2h, the reaction was poured into saturated Na₂SO₃ (aq, 50 mL). The organic layer was separated, dried over Na₂SO₄ and concentrated. The resulting residue was purified by HPLC (ISCO 120 g, 0 to 50% EtOAc/Hex) to give the title compound. ¹H NMR (500 MHz, CDCl₃)
30 δ 7.60 (d, 1H), 6.85 (s, 1H), 6.50 (d, 1H), 5.30 (br, 1H), 3.75 (s, 3H), 2.80 (m, 1H), 1.90 (t, 1H), 1.05 (m, 1H), 0.97 (d, 3H), 0.60 (m, 1H), 0.37 (m, 1H), 0.27 (m, 1H), 0.01 (m, 1H). (m/z): 361.07 (M+H)⁺.

SCHEME 2



Scheme 2 provides a route to prepare the compounds of the present invention. Suzuki cross-coupling of boronic acid **1** and aldehyde **2**, followed by allyl grignard addition to aldehyde **3** delivered the homoallylic alcohol **4**. Heck reaction of aryl iodide **5** with homoallylic alcohol **4** gave diol **6**. Reduction of the alkene was achieved via palladium-mediated hydrogenation to give **7**. An intramolecular Mitsunobu reaction resulted in ring closure to deliver the tetrahydrobenzoxapine ring **8**. SFC separation of the benzylic diastereomers was followed by hydrolysis of the methyl ester to give the final compound **10**.

SCHEME 3




Step 1: Pd(PPh₃)₄, K₂CO₃, dioxane; **Step 2:** allylMgBr, THF; **Step 3:** PPh₃, DIAD, PhMe; **Step 4:** Hoveyda-Grubbs 2, DCM; **Step 5:** SPhos Pd precat, THF; **Step 6:** LiOH, THF/H₂O/MeOH; **Step 7:** H₂ (1 atm), Pd(OH)₂, EtOAc

Scheme 3 provides an alternative route to prepare the compounds of the present invention. Suzuki cross-coupling of boronic acid **1** and aldehyde **2**, followed by allyl grignard addition to aldehyde **3** delivered the homoallylic alcohol **4**. Mitsunobu coupling of the homoallylic alcohol **4** and 4-bromo-2-vinylphenol gave the aryl benzyl ether **5**. Aryl benzyl ether **5** was subjected to ring-closing metathesis with catalytic Hoveyda-Grubbs 2 reagent to prepare dihydrobenzoxapiene **6**. The propanoic acid moiety was installed using a two step sequence consisting of Negishi reaction with (3-ethoxy-3-oxopropyl)zinc (II) bromide followed by lithium hydroxide-mediated hydrolysis of the ethyl ester to give acid **8**. Finally, reduction of the dihydrobenzoxapiene double bond of **8** was achieved with hydrogenation in the presence of Pd(OH)₂ to provide compound **9**.

EXAMPLE 1

(2S,3R)-3-cyclopropyl-3-((R or S)-2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid



Step A: 1-(2-fluoro-4-(2-methoxypyridin-4-yl) phenyl) but-3-en-1-ol To a solution of 2-fluoro-4-(2-methoxypyridin-4-yl)benzaldehyde (1.00 g, 4.32 mmol) in THF (10 mL) was added allylmagnesium bromide (8.65 mL, 8.65 mmol, 1.0 M) dropwise at 0 °C under a N₂ atmosphere.

5 The reaction mixture was stirred at 0 °C for 30 min, and at 20 °C for 3 hours. Then the reaction mixture was quenched with saturated NH₄Cl solution (20 mL) and extracted with EtOAc (10 mL×3). The combined organic layers were washed with brine (30 mL), dried over anhydrous Na₂SO₄ and filtered. The filtrate was concentrated *in vacuo* to give a residue, which was purified by preparative TLC (PE: EtOAc = 5: 1, v/v) to afford the title compound. ¹H NMR (400 MHz, CDCl₃): δ = 8.23 (d, *J* = 5.2 Hz, 1H), 7.60 (d, *J* = 8.0 Hz, 1H), 7.44 (d, *J* = 7.6 Hz, 1H), 7.30 (d, *J* = 7.6 Hz, 1H), 7.09 (d, *J* = 4.4 Hz, 1H), 6.93 (s, 1H), 5.89-5.82 (m, 1H), 5.23-5.19 (m, 2H), 5.13-5.11 (m, 1H), 3.99 (s, 3H), 2.63-2.49 (m, 2H).

Step B: (2S, 3R)-methyl 3-cyclopropyl-3-(4-((Z)-4-(2-fluoro-4-(2-methoxypyridin-4-yl) phenyl)-4-hydroxybut-1-en-1-yl)-3-hydroxyphenyl)-2-methylpropanoate To a solution of 1-(2-fluoro-4-(2-


15 methoxypyridin-4-yl)phenyl)but-3-en-1-ol (804 mg, 2.94 mmol) and (2S, 3R)-methyl 3-cyclopropyl-3-(3-hydroxy-4-iodophenyl)-2-methylpropanoate (530 mg, 1.47 mmol) in toluene (10 mL) were added *N*-cyclohexyl-*N*-methylcyclohexanamine (575 mg, 2.94 mmol) and t-buxphos Pd G1 (CAS#1142811-12-8, 101 mg, 0.147 mmol) under a N₂ atmosphere. The reaction mixture was stirred at 100 °C for 2 hours, then cooled to room temperature and concentrated *in vacuo* to give a residue, which was purified by silica gel chromatography (eluted from PE: EtOAc = 100: 1 to 5: 1, v/v) to afford the title compound. ¹H NMR (400 MHz, CDCl₃): δ = 8.22 (d, *J* = 5.2 Hz, 1H), 7.65-7.61 (m, 1H), 7.45 (d, *J* = 7.2 Hz, 1H), 7.32-7.29 (m, 1H), 7.26-7.25 (m, 1H), 7.08 (d, *J* = 5.6 Hz, 1H), 6.94 (s, 1H), 6.73-6.69 (m, 2H), 6.61 (s, 1H), 6.24-6.16 (m, 1H), 5.21-5.18 (m, 1H), 3.99 (s, 3H), 3.73 (s, 3H), 2.80-2.66 (m, 3H), 2.32-2.26 (m, 1H), 1.88-1.83 (m, 1H), 1.08-1.07 (m, 1H), 0.84 (d, *J* = 4.0 Hz, 3H), 0.57-0.55 (m, 1H), 0.36-0.30 (m, 1H), 0.25-0.22 (m, 1H), 0.01--0.03 (m, 1H).

Step C: (2S, 3R)-methyl 3-cyclopropyl-3-(4-(4-(2-fluoro-4-(2-methoxypyridin-4-yl) phenyl) -4-hydroxybutyl)-3-hydroxyphenyl)-2-methylpropanoate To a solution of (2S, 3R)-methyl 3-cyclopropyl-3-(4-((Z)-4-(2-fluoro-4-(2-methoxypyridin-4-yl) phenyl)-4-hydroxybut-1-en-1-yl)-3-

hydroxyphenyl)-2-methylpropanoate (610 mg, 1.21 mmol) in MeOH (10 mL) was added Pd(OH)₂/C (85.0 mg, 20%). The reaction mixture was stirred for at 20°C for 3 hours under H₂ balloon pressure. Then the reaction mixture was filtered, and the filtrate was concentrated in vacuo to give the title compound, which was used in the next step. ¹H NMR (400 MHz, CDCl₃): δ = 8.20 (d, *J* = 6.0 Hz, 1H), 7.58 (t, *J* = 7.6 Hz, 1H), 7.41 (dd, *J*₁₃ = 8.0 Hz, *J*₁₂ = 2.0 Hz, 1H), 7.28 (dd, *J*₁₃ = 11.2 Hz, *J*₁₂ = 2.0 Hz, 1H), 7.07 (dd, *J*₁₃ = 7.2 Hz, *J*₁₂ = 1.6 Hz, 1H), 7.01 (d, *J* = 8.0 Hz, 1H), 6.92 (s, 1H), 6.65 (d, *J* = 8.0 Hz, 1H), 6.61 (s, 1H), 5.17 (t, *J* = 6.0 Hz, 1H), 3.99 (s, 3H), 3.72 (s, 3H), 2.80-2.65 (m, 3H), 1.88-1.74 (m, 6H), 1.05-0.97 (m, 1H), 0.93 (d, *J* = 6.8 Hz, 3H), 0.56-0.51 (m, 1H), 0.34-0.20 (m, 2H), 0.02--0.02 (m, 1H).

Step D: (2S, 3R)-methyl 3-cyclopropyl-3-(2-(2-fluoro-4-(2-methoxypyridin-4-yl) phenyl)-2, 3, 4, 5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate To a solution of (2S,3R)-methyl 3-cyclopropyl-3-(4-(4-(2-fluoro-4-(2-methoxypyridin-4-yl) phenyl)-4-hydroxybutyl)-3-hydroxyphenyl)-2-methylpropanoate (400 mg, 0.788 mmol) and triphenylphosphine (289 mg, 1.10 mmol) in DCM (3.0 mL) was added dropwise a solution of diisopropyl azodicarboxylate (0.215 mL, 1.10 mmol) in DCM (1.0 mL) at 0 °C under a N₂ atmosphere. The reaction mixture was stirred at 20 °C for 2 hours, then concentrated *in vacuo* to give a residue, which was purified by preparative TLC (PE: EtOAc = 5: 1, v/v) to afford the title compound. ¹H NMR (400 MHz, CDCl₃): δ = 8.25 (d, *J* = 5.2 Hz, 1H), 7.79 (t, *J* = 8.0 Hz, 1H), 7.41 (dd, *J*₁₃ = 8.4 Hz, *J*₁₂ = 2.0 Hz, 1H), 7.33-7.30 (m, 1H), 7.11-7.09 (m, 2H), 6.96 (s, 1H), 6.84-6.79 (m, 2H), 4.96-4.92 (m, 1H), 4.00 (s, 3H), 3.71 (s, 3H), 3.00 (t, *J* = 12.8 Hz, 1H), 2.83-2.76 (m, 2H), 2.24-2.21 (m, 1H), 2.15-2.09 (m, 2H), 1.87 (t, *J* = 10.0 Hz, 1H), 1.72-1.69 (m, 1H), 1.06-0.99 (m, 1H), 0.94 (dd, *J*₁₃ = 6.8 Hz, *J*₁₂ = 5.6 Hz, 3H), 0.57-0.50 (m, 1H), 0.36-0.29 (m, 1H), 0.25-0.19 (m, 1H), 0.04--0.04 (m, 1H).

Step E: (2S, 3R)-methyl 3-cyclopropyl-3-((R or S)-2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate; and
(2S, 3R)-methyl 3-cyclopropyl-3-((S or R)-2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate



(2S, 3R)-methyl 3-cyclopropyl-3-(2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate (200 mg, 0.409 mmol) was separated via SFC (using SFC conditions: "Column: Chiralpak AD-3 50*4.6mm I.D., 3 μ m Mobile phase: ethanol (0.05 % DEA) in CO₂ from 5 % to 40 % Flow rate: 4 mL/min Wavelength: 220 nm") to give (2S, 3R)-methyl 3-cyclopropyl-3-((R or S)-2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate, and (2S, 3R)-methyl 3-cyclopropyl-3-((S or R)-2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate.

Step F: (2S,3R)-3-cyclopropyl-3-((R or S)-2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid


To a solution of (2S, 3R)-methyl 3-cyclopropyl-3-((R or S)-2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate (100 mg, 0.204 mmol) in THF/MeOH/H₂O (1.0 mL/1.0 mL/1.0 mL) was added LiOH (48.9 mg, 2.04 mmol). The reaction mixture was stirred at 50 °C for 16 hours, and then cooled to room temperature. The reaction mixture was then acidified with HCl (1.0 N) to adjust the pH to 2, and was extracted with EtOAc (5.0 mL $\times$ 3). The combined organic layers were concentrated *in vacuo* to afford a residue, which was purified by preparative HPLC (on a GILSON 281 instrument fitted with a Phenomenex Gemini C₁₈ (250*21.2mm*5 μ m) using water and acetonitrile as the eluents. Mobile phase A: water (neutral), mobile phase B: acetonitrile.

Gradient: 33-63 % B, 0-10 min; 100 % B, 10.5-12.5 min; 5 % B, 13-15 min) to give (2S,3R)-3-cyclopropyl-3-((R or S)-2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid. ¹H NMR (400 MHz, CDCl₃): δ = 8.25 (d, J = 4.8 Hz, 1H), 7.79 (t, J = 8.0 Hz, 1H), 7.47 (d, J = 7.2 Hz, 1H), 7.30 (d, J = 11.2 Hz, 1H), 7.12-7.10 (m, 2H), 6.96 (s, 1H), 6.84-6.83 (m, 2H), 4.92 (d, J = 10.0 Hz, 1H), 4.00 (s, 3H), 3.00 (t, J =

13.2 Hz, 1H), 2.83-2.79 (m, 2H), 2.24-2.09 (m, 3H), 1.94 (t, $J = 10.0$ Hz, 1H), 1.75-1.66 (m, 1H), 1.10-1.08 (m, 1H), 0.98 (d, $J = 6.8$ Hz, 3H), 0.61-0.60 (m, 1H), 0.36-0.33 (m, 2H), 0.03-0.02 (m, 1H).

To a solution of (2S,3R)-3-cyclopropyl-3-((R or S)-2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid in MeCN (1.0 mL) and water (1.0 mL) was added aq. NaOH (1.0 eq, 0.5 M). The mixture was stirred at 20 °C for 1 hour. Then the reaction mixture was lyophilized to give the sodium salt of the title compound.


TABLE 1 Example 2 was prepared in a similar manner to Example 1 using the appropriate starting materials

Example	Structure	MW	Compound Name	LC/MS (ESI) observed $[M+1]^+$
2		475.55	(2S,3R)-3-cyclopropyl-3-((R or S)-2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid	476.2

*stereoisomer at this position

EXAMPLE 3

(2S, 3R)-3-((R or S)-2-(1-(2, 5-bis (trifluoromethyl) benzyl) azetidin-3-yl)-2, 3, 4, 5-tetrahydrobenzo[b]oxepin-8-yl)-3-cyclopropyl-2-methylpropanoic acid



Step A: *tert*-butyl 3-(1-hydroxybut-3-en-1-yl) azetidine-1-carboxylate

To a solution of *tert*-butyl 3-formylazetidine-1-carboxylate (2.00 g, 10.8 mmol) in THF (20 mL) was added dropwise allylmagnesium bromide (21.6 mL, 21.6 mmol) at 0 °C under a N₂ atmosphere. The reaction mixture was stirred for 30 min at 0 °C and 3 hours at 20 °C (room temperature). Then the reaction mixture was quenched with saturated NH₄Cl (20 mL) and extracted with EtOAc (10 mL×3). The combined organic layers were washed with brine (50 mL), dried over anhydrous Na₂SO₄ and filtered. The filtrate was concentrated *in vacuo* to give the title compound, which was used in the next step. ¹H NMR (400 MHz, CDCl₃): δ = 5.85-5.75 (m, 1H), 5.19-5.04 (m, 2H), 4.14-3.71 (m, 6H), 2.24-2.04 (m, 2H), 1.44 (s, 9H).

Step B: *tert*-butyl 3-(4-(4-((1R, 2S)-1-cyclopropyl-3-methoxy-2-methyl-3-oxopropyl)-2-hydroxyphenyl) butanoyl) azetidine-1-carboxylate

To a solution of *tert*-butyl 3-(1-hydroxybut-3-en-1-yl)azetidine-1-carboxylate (2.27 g, 9.99 mmol) in DMF (20 ml) were added lithium chloride (1.05 g, 24.9 mmol), lithium acetate (1.65 g, 24.9 mmol), (2S,3R)-methyl 3-cyclopropyl-3-(3-hydroxy-4-iodophenyl)-2-methyl propanoate (3.00 g, 8.33 mmol), tetrabutylammonium chloride (4.63 g, 16.6 mmol) and PdOAc₂ (0.374 g, 1.66 mmol) under a N₂ atmosphere. The reaction mixture was stirred for 2 hours at 100 °C, then diluted with water (20 mL). The mixture was extracted with EtOAc (30 mL×3). The combined organic layers were washed with water (100 mL) twice. Then the organic layer was concentrated *in vacuo* to give crude product, which was purified via silica gel column (PE: EtOAc=20: 1) to give the title compound. MS (ESI) m/z: 482.2 [M+Na⁺].

Step C: *tert*-butyl 3-(4-(4-((1R, 2S)-1-cyclopropyl-3-methoxy-2-methyl-3-oxopropyl)-2-hydroxyphenyl)-1-hydroxybutyl) azetidine-1-carboxylate

A mixture of compound *tert*-butyl 3-((E)-4-(4-((1R,2S)-1-cyclopropyl-3-methoxy-2-methyl-3-oxopropyl)-2-hydroxyphenyl)-1-hydroxybut-3-en-1-yl) azetidine-1-carboxylate (100 mg, 0.218 mmol) and PdOH₂/C (1.528 mg, 20%) in MeOH (10 ml) was stirred at 15 °C for 16 hours under H₂ atmosphere. The mixture was filtered, and the filtrate was dried over MgSO₄ and concentrated *in vacuo* to give the crude product. The crude product was purified by prep-TLC (PE: EtOAc = 3: 1) to give the title compound. MS (ESI) m/z: 462.3 [M+H⁺].

Step D: *tert*-butyl 3-(8-((1R, 2S)-1-cyclopropyl-3-methoxy-2-methyl-3-oxopropyl)-2, 3, 4, 5-tetrahydrobenzo[b]oxepin-2-yl) azetidine-1- carboxylate

To a solution *tert*-butyl 3-(4-(4-((1R, 2S)-1-cyclopropyl-3-methoxy-2-methyl-3-oxopropyl)-2- hydroxyphenyl)-1-hydroxybutyl) azetidine-1-carboxylate (453 mg, 0.981 mmol) and triphenylphosphine (515 mg, 1.96 mmol) in DCM (8.0 mL) was added a solution of (E)-diisopropyl diazene-1,2-dicarboxylate (397 mg, 1.96 mmol) in DCM

(2.0 mL) at 0 °C under N₂ atmosphere. The reaction mixture was stirred for 1 hour at 20 °C, then concentrated *in vacuo* to give a residue, which was purified by prep-TLC (PE: EtOAc = 5:1) to give the title compound.

Step E: (2S, 3R)-methyl 3-(2-(azetidin-3-yl)-2, 3, 4, 5-tetrahydrobenzo[b]oxepin-8-yl)-3-

cyclopropyl-2-methylpropanoate To a solution of *tert*-butyl 3-(8-(1-cyclopropyl-3-methoxy-2-methyl-3-oxopropyl)-2, 3, 4, 5-tetrahydrobenzo[b]oxepin-2-yl) azetidine-1-carboxylate (60.0 mg, 0.135 mmol) in DCM (10 ml) was added 2, 2, 2-trifluoroacetic acid (308 mg, 2.71 mmol). Then the mixture was stirred at 20 °C for 30 min, and concentrated *in vacuo* to give the title compound, which was used in next step. MS (ESI) *m/z*: 344.1 [M+H⁺].


Step F: (2S, 3R)-methyl 3-(2-(1-(2, 5-bis (trifluoromethyl) benzyl) azetidin-3-yl)-2, 3, 4, 5-
tetrahydrobenzo[b]oxepin-8-yl)-3-cyclopropyl-2-methylpropanoate A mixture of compound K₂CO₃ (66.0 mg, 0.477 mmol), 2-(bromomethyl)-1,4-bis (trifluoromethyl)benzene (110 mg, 0.358 mmol) and methyl 3-(2-(azetidin-3-yl)-2,3,4,5- tetrahydrobenzo[b]oxepin-8-yl)-3-cyclopropyl-2-methylpropanoate (41.0 mg, 0.119 mmol) in MeCN (10 ml) was stirred at 20 °C for 10 hours. Then water (10 mL) was added, and the reaction mixture was extracted with DCM (5 ml) twice. The combined organic layers were dried over Na₂SO₄, and concentrated *in vacuo* to give the crude product. The crude product was purified via prep-TLC (PE: EtOAc = 5: 1) to give the title compound. MS (ESI) *m/z*: 570.2 [M+H⁺].

Step G: (2S, 3R)-3-(2-(1-(2, 5- bis (trifluoromethyl) benzyl) azetidin-3-yl)-2, 3, 4, 5-

tetrahydrobenzo[b]oxepin-8-yl)-3-cyclopropyl-2-methylpropanoic acid A mixture of LiOH (17.24 mg, 0.720 mmol) and (2S,3R)-methyl 3-(2-(1-(2,5-bis(trifluoromethyl)benzyl)azetidin-3-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-3-cyclopropyl-2-methylpropanoate (41 mg, 0.072 mmol) in water (1.0 mL), THF (1.0 mL) and MeOH (1.0 mL) was stirred at 50 °C for 2 hours. The reaction mixture was then acidified with HCl (2N, 1.0 mL) and diluted with DCM (10 mL). Then the reaction mixture was washed with water (20 ml). The organic layer was separated, dried over anhydrous MgSO₄ and filtered. The filtrate was concentrated to give the crude product, which was purified by prep-HPLC (on a GILSON 281 instrument fitted with a Phenomenex Synergi C18 100*21.2mm*4um using water and acetonitrile as the eluents. Mobile phase A: water (containing 0.1% TFA, v/v), mobile phase B: acetonitrile. Gradient: 32-58% B, 0-10min; 100% B, 10.5-12.5min; 5% B, 13-15min) to give the title compound. ¹H NMR (400 MHz, CDCl₃): δ = 8.15 (s, 1H), 7.93-7.80 (m, 2H), 7.10 (d, *J* = 6.4 Hz, 2H), 6.87-6.78 (m, 2H), 4.74-4.58 (m, 4H), 4.12 (s, 2H),

3.65 (s, 1H), 3.22 (s, 1H), 2.87-2.76 (m, 3H), 2.00 (s, 2H), 1.81-1.78 (m, 1H), 1.61-1.51 (m, 2H), 1.01-1.00 (m, 4H), 0.63-0.62 (m, 1H), 0.38-0.37 (m, 2H), 0.05-0.04 (m, 1H).

Step H: (2S, 3R)-3-((R or S)-2-(1-(2,5-bis(trifluoromethyl)benzyl)azetidin-3-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-3-cyclopropyl-2-methylpropanoic acid and
 5 (2S, 3R)-3-((R or S)-2-(1-(2,5-bis(trifluoromethyl)benzyl)azetidin-3-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-3-cyclopropyl-2-methylpropanoic acid




(2S, 3R)-3-(2-(1-(2,5-bis(trifluoromethyl)benzyl)azetidin-3-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-3-cyclopropyl-2-methylpropanoic acid (25 mg, 0.045 mmol) was separated via SFC (SFC
 10 conditions: Instrument: Berger MultiGramTM SFC, Mettler Toledo Co, Ltd, column: AD (250*30mm 5um), mobile phase: 10 % ETOH+NH₃H₂O 50mL/min; wavelength: 220nm) to give 2 peaks: peak 1: (2S,3R)-3-((R or S)-2-(1-(2,5-bis(trifluoromethyl)benzyl)azetidin-3-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-3-cyclopropyl-2-methylpropanoic acid, and peak 2: (2S,3R)-3-((R or S)-2-(1-(2,5-bis(trifluoromethyl)benzyl)azetidin-3-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-3-cyclopropyl-2-methylpropanoic acid.
 15

To the first compound (2S,3R)-3-((R or S)-2-(1-(2,5-bis(trifluoromethyl)benzyl)azetidin-3-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-3-cyclopropyl-2-methylpropanoic acid (9 mg, 0.016 mmol) was added a solution of NaOH (aq. 33 mg, 0.5 M) in MeCN (0.5 mL). The mixture was stirred for 1 hour at room temperature. Then the reaction mixture was lyophilized to give the sodium salt of the
 20 title compound. ¹H NMR (400 MHz, CDCl₃): δ = 8.00 (s, 1H), 7.77 (d, *J* = 7.2 Hz, 1H), 7.62 (d, *J* = 6.8 Hz, 1H), 7.06 (d, *J* = 6.8 Hz, 1H), 6.84 (d, *J* = 6.8 Hz, 1H), 3.96 (s, 2H), 3.74-3.64 (m, 3H), 3.40-3.39 (m, 1H), 3.13-3.13 (m, 1H), 2.84-2.73 (m, 4H), 2.05-1.99 (m, 2H), 1.86-1.83 (m, 1H), 1.64-1.50 (m, 2H), 1.16-1.15 (m, 1H), 1.00 (d, *J* = 5.6 Hz, 3H), 0.61-0.60 (m, 1H), 0.38-0.37 (m, 2H), 0.06-0.05 (m, 1H).

To a solution of the second compound (2S,3R)-3-((R or S)-2-(1-(2,5-bis(trifluoro-methyl)benzyl)azetidin-3-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-3-cyclopropyl-2-methylpropanoic acid (10 mg, 0.017 mmol) in MeCN (0.5 mL) was added the solution of NaOH (aq. 37 mg, 0.5 M). The mixture was stirred for 1 hour at room temperature. Then the reaction mixture was lyophilized to give the sodium salt of the title compound. MS (ESI) *m/z*: 556.1 [M+H]⁺.

Table 2 Example 4 was prepared in a similar manner to Example 3 using the appropriate starting materials


Example	Structure	MW	Compound Name	LC/MS (ESI) observed [M+1] ⁺
4		555.55	(2S,3R)-3-((R or S)-2-(1-(2,5-bis(trifluoromethyl)-benzyl)azetidin-3-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-3-cyclopropyl-2-methylpropanoic acid	556.1

*stereoisomer at this position

5

EXAMPLE 5

(2S,3R)-3-cyclopropyl-3-((RS)-2-(4-(5-fluoro-2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid



10

Step A: 4-(5-fluoro-2-methoxypyridin-4-yl)benzaldehyde

To a solution of compounds (5-fluoro-2-methoxypyridin-4-yl)boronic acid (800 mg, 4.68 mmol) and 4-bromobenzaldehyde (1.03 g, 5.62 mmol) in THF (10 mL) and water (2.0 mL) were added K₂CO₃ (1.93 g, 14.0 mmol) and PdCl₂(dppf) (342 mg, 0.468 mmol) under a N₂ atmosphere. The reaction mixture was stirred at 100 °C for 2 hours, then water (10 mL) was added and the mixture was extracted with EtOAc (2 mL×3). The combined organic layers were washed with brine (10 mL), dried over anhydrous Na₂SO₄ and filtered. The filtrate was concentrated *in vacuo* to give a residue, which was purified by preparative TLC (PE/EtOAc = 5:1) to give the title compound. ¹H NMR

15

(400 MHz, CDCl₃): δ = 10.1 (s, 1H), 8.12 (s, 1H), 8.01 (d, J = 8.0 Hz, 2H), 7.77 (d, J = 7.2 Hz, 2H), 6.85 (d, J = 5.2 Hz, 1H), 3.97 (s, 3H). MS(ESI) m/z : 232.0 [M+H]⁺.

Step B: 1-(4-(5-fluoro-2-methoxypyridin-4-yl)phenyl)but-3-en-1-ol To a solution of 4-(5-fluoro-2-methoxypyridin-4-yl)benzaldehyde (650 mg, 2.81 mmol) in THF (10 mL) was added

5 allylmagnesium bromide (11.2 mL, 11.2 mmol) dropwise at 0 °C under a N₂ atmosphere. The reaction mixture was stirred for 30 mins at 0 °C, and 3 hours at room temperature. Then the reaction was quenched with saturated NH₄Cl (20 mL) and extracted with EtOAc (5.0 mL×3). The combined organic layers were washed with brine (30 mL), dried over anhydrous Na₂SO₄ and filtered. The filtrate was concentrated *in vacuo* to give a residue, which was purified by preparative TLC (PE: EtOAc = 5: 1, v/v) to afford the title compound. ¹H NMR (400 MHz, CDCl₃): δ = 8.07 (s, 1H), 7.60 (d, J = 7.6 Hz, 2H), 7.50 (d, J = 7.6 Hz, 1H), 6.85 (m, 1H), 5.88-5.80 (m, 1H), 6.93 (s, 1H), 5.24 (t, J = 10.4 Hz, 2H), 4.83 (d, J = 7.6 Hz, 1H), 3.97 (s, 3H), 2.61 (m, 2H). MS(ESI) m/z : 274.1 [M+H]⁺.

Step C: (2S,3R)-methyl-3-cyclopropyl-3-(4-((Z)-4-(4-(5-fluoro-2-methoxypyridin-4-yl)phenyl)-4-hydroxybut-1-en-1-yl)-3-hydroxyphenyl)-2-methylpropanoate To a solution of 1-(4-(5-fluoro-2-

15 methoxypyridin-4-yl)phenyl)but-3-en-1-ol (620 mg, 2.27 mmol) and (2S, 3R)-methyl 3-cyclopropyl-3-(3-hydroxy-4-iodophenyl)-2-methylpropanoate (545 mg, 1.51 mmol) in toluene (10 mL) were added *N*-cyclohexyl-*N*-methylcyclohexanamine (443 mg, 2.27 mmol) and t-buxhos-palladacycle (104 mg, 0.151 mmol) under a N₂ atmosphere. The reaction mixture was stirred for 2 hours at 100 °C, then cooled to room temperature and concentrated *in vacuo* to give a residue. The residue was purified via silica gel chromatography (eluted from PE: EtOAc = 100: 1 to 5: 1, v/v) to afford the title compound. MS(ESI) m/z : 506.2 [M+H]⁺.

Step D: (2S, 3R)-methyl 3-cyclopropyl-3-(4-(4-(2-fluoro-4-(2-methoxypyridin-4-yl) phenyl)-4-hydroxybutyl)-3-hydroxyphenyl)-2-methylpropanoate To a solution of (2S, 3R)-methyl (2S,3R)-

25 methyl 3-cyclopropyl-3-(4-((Z)-4-(4-(5-fluoro-2-methoxypyridin-4-yl)phenyl)-4-hydroxybut-1-en-1-yl)-3-hydroxyphenyl)-2-methylpropanoate (450 mg, 0.890 mmol) in MeOH (10 mL) was added Pd(OH)₂/C (13.0 mg, 0.0890 mmol, 20%). The reaction mixture was stirred for 3 hours at room temperature under a H₂ balloon. Then the reaction mixture was filtered, and the filtrate was concentrated *in vacuo* to give the title compound, which was used in the next step. ¹H NMR (400 MHz, CDCl₃): δ = 8.05 (s, 1H), 7.57 (d, J = 7.6 Hz, 2H), 7.46 (d, J = 8.4 Hz, 2H), 7.03 (d, J = 7.2 Hz, 1H), 6.81 (d, J = 5.2 Hz, 1H), 6.65-6.61 (m, 2H), 4.86 (d, J = 6.8 Hz, 1H), 3.94 (s, 3H), 3.74 (s, 3H), 2.80-2.72 (m, 3H), 1.86-1.72 (m, 4H), 1.03-0.93 (m, 4H), 0.54 (s, 1H), 0.32-0.21 (m, 2H), 0.01-0.004 (m, 1H). MS(ESI) m/z : 508.2 [M+H]⁺.

Step E: (2S,3R)-methyl 3-cyclopropyl-3-(2-(4-(5-fluoro-2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate

To a solution of (2S,3R)-methyl 3-cyclopropyl-3-(4-(4-(4-(5-fluoro-2-methoxypyridin-4-yl)phenyl)-4-hydroxybutyl)-3-hydroxyphenyl)-2-methylpropanoate (345 mg, 0.680 mmol) and triphenylphosphine (357 mg, 1.36 mmol) in DCM (3.0 mL) was added dropwise a solution of DIAD (275 mg, 1.36 mmol) in DCM (1.0 mL) at 0 °C under a N₂ atmosphere. The reaction mixture was stirred for 2 hours at room temperature, then concentrated *in vacuo* to give a residue, which was purified by preparative TLC (PE: EtOAc = 5: 1, v/v) to afford the title compound. ¹H NMR (400 MHz, CDCl₃): δ = 8.09 (s, 1H), 7.65-7.58(m, 4H), 7.12 (d, *J* = 7.2 Hz, 1H), 6.86-6.79 (m, 3H), 4.68 (t, *J* = 5.6 Hz, 1H), 3.97 (s, 3H), 3.72 (s, 3H), 3.00 (t, *J* = 13.2 Hz, 1H), 2.82-2.78 (m, 2H), 2.22-2.12 (m, 4H), 1.87 (t, *J* = 10.0 Hz, 1H), 1.72-1.62 (m, 1H), 1.04(d, *J* = 4.8 Hz, 1H), 0.94 (t, *J* = 5.6 Hz, 3H), 0.55 (d, *J* = 3.6 Hz, 1H), 0.34 (d, *J* = 4.0 Hz, 1H), 0.23 (d, *J* = 4.4 Hz, 1H), 0.12 (s, 1H). MS(ESI) *m/z*: 490.3[M+H]⁺.

Step F: (2S,3R)-methyl 3-cyclopropyl-3-((RS)-2-(4-(5-fluoro-2-methoxypyridin-4-yl)phenyl)-

2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate ((2S,3R)-methyl 3-cyclopropyl-3-((RS)-2-(4-(5-fluoro-2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate (60 mg, 0.123 mmol) was separated by SFC (SFC conditions: Column: Chiralpak AD-3 50*4.6mm I.D., 3 μm Mobile phase: ethanol (0.05 % DEA) in CO₂ from 5 % to 40 % Flow rate: 4 mL/min Wavelength: 220 nm) to give 2 peaks: peak 1: (2S,3R)-methyl 3-cyclopropyl-3-((RS)-2-(4-(5-fluoro-2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate; and peak 2: (2S,3R)-methyl 3-cyclopropyl-3-((RS)-2-(4-(5-fluoro-2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate.

Step G: sodium (2S,3R)-methyl 3-cyclopropyl-3-((RS)-2-(4-(5-fluoro-2-methoxypyridin-4-

yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate To a solution of (2S,3R)-methyl 3-cyclopropyl-3-((RS)-2-(4-(5-fluoro-2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate (30 mg, 0.0613 mmol) in THF/MeOH/H₂O (1.0 mL/1.0 mL/1.0 mL) was added LiOH (15.0 mg, 0.613 mmol). The reaction mixture was stirred at 50 °C for 16 hours, then cooled to room temperature, and acidified with HCl (1mol/L) to pH 2. The mixture was extracted with EtOAc (2 mL×3), and the combined organic layers were concentrated *in vacuo* to afford a residue. The residue was purified by preparative HPLC (on a GILSON 281 instrument fitted with a Phenomenex Gemini C₁₈ (250*21.2mm*5μm) using water and acetonitrile as the

eluent. Mobile phase A: water (neutral), mobile phase B: acetonitrile. Gradient: 33-63 % B, 0-10 min; 100 % B, 10.5-12.5 min; 5 % B, 13-15 min) to give (2S,3R)-3-cyclopropyl-3-(2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydro-benzo[b]oxepin-8-yl)-2-methylpropanoic acid.


To a solution of (2S,3R)-3-cyclopropyl-3-(2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid (12.6 mg) in MeCN (0.5 mL) was added a solution of NaOH (aq. 172 mg, 0.5 M). The mixture was stirred for 1 hour at room temperature, then the reaction mixture was lyophilized to give the sodium salt of (2S,3R)-3-cyclopropyl-3-(2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-

methylpropanoic acid. ^1H NMR (400 MHz, CDCl_3): δ = 8.13 (d, J = 1.2 Hz, 1H), 7.64-7.58 (m, 4H), 7.12 (d, J = 7.6 Hz, 1H), 6.88-6.83 (m, 3H), 4.69 (d, J = 10.4 Hz, 1H), 3.98 (s, 3H), 2.97 (t, J = 12.8 Hz, 1H), 2.82-2.78 (m, 2H), 2.21-2.11 (m, 3H), 1.94 (t, J = 10.0 Hz, 1H), 1.75-1.66 (m, 1H), 1.10-1.08 (m, 1H), 0.98 (d, J = 6.8 Hz, 3H), 0.62-0.61 (m, 1H), 0.37-0.32 (m, 2H), 0.03-0.02 (m, 1H).

MS(ESI) m/z : 476.2 $[\text{M}+\text{H}]^+$.

To a solution of (2S,3R)-3-cyclopropyl-3-(2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid (12.6 mg) in MeCN (0.5 mL) was added the solution of NaOH (aq. 172 mg, 0.5 M), the mixture was stirred for 1 hour at 20 °C (room temperature). Then the reaction mixture was lyophilized to give the sodium salt of (2S,3R)-3-cyclopropyl-3-(2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid.


Table 3 Example 6 was prepared in a similar manner to Example 5 using the appropriate starting materials.

Example	Structure	MW	Compound Name	LC/MS (ESI) observed $[\text{M}+1]^+$
6		475.55	(2S,3R)-3-cyclopropyl-3-((R or S)-2-(4-(5-fluoro-2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid	476.2

*stereoisomer at this position

EXAMPLE 7

5 (2S,3R)-3-cyclopropyl-3-((RS)-2-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid



Step A: 5'-fluoro-2'-methoxy-[3,4'-bipyridine]-6-carbaldehyde

To a solution of compounds (5-fluoro-2-methoxypyridin-4-yl)boronic acid (1.00 g, 5.85 mmol) and
 10 5-bromopicolinaldehyde (1.19 g, 6.44 mmol) in THF (10 mL) and water (2.0 mL) were added
 K_2CO_3 (2.42 g, 1.76 mmol) and $PdCl_2(dppf)$ (428 mg, 0.585 mmol) under a N_2 atmosphere. The
 reaction mixture was stirred at 100 °C for 2 hours, then water (10 mL) was added and the mixture
 was extracted with EtOAc (2 mL×3). The combined organic layers were washed with brine (10
 mL), dried over anhydrous Na_2SO_4 and filtered. The filtrate was concentrated *in vacuo* to give a
 15 residue, which was purified by preparative TLC (PE/EtOAc = 5:1, v/v) to give the title compound.
 ^1H NMR (400 MHz, $CDCl_3$): δ = 10.2 (s, 1H), 8.99 (s, 1H), 8.17 (m, 3H), 6.88 (d, J = 4.8Hz, 1H),
 3.98 (s, 3H). MS (ESI) m/z : 233.0 $[M+H]^+$.

Step B: 1-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)but-3-en-1-ol

To a solution of 5'-fluoro-2'-methoxy-[3,4'-bipyridine]-6-carbaldehyde (950 mg, 4.09 mmol) in THF
 20 (10 mL) was added allylmagnesium bromide (8.18 mL, 8.18 mmol) dropwise at 0 °C under a N_2
 atmosphere. The reaction mixture was stirred for 30 min at 0 °C and 3 hours at room temperature.
 Then the reaction mixture was quenched with saturated NH_4Cl (20 mL) and extracted with EtOAc
 (5.0 mL×3). The combined organic layers were washed with brine (30 mL), dried over anhydrous
 Na_2SO_4 and filtered. The filtrate was concentrated *in vacuo* to give a residue, which was purified by
 25 preparative TLC (PE: EtOAc = 5: 1, v/v) to give the title compound. ^1H NMR (400 MHz, $CDCl_3$):
 δ = 8.77 (s, 1H), 8.12 (s, 1H), 7.94 (d, J = 4.8Hz, 1H), 7.46 (d, J = 4.2Hz, 1H), 5.91-5.84 (m, 1H),
 5.19-5.15 (m, 2H), 4.91-4.88 (m, 1H), 3.97 (s, 3H), 2.73-2.69 (m, 1H), 2.58-2.52 (m, 1H). MS(ESI)
 m/z : 275.1 $[M+H]^+$.

Step C: (2S,3R)-methyl 3-cyclopropyl-3-(4-((Z)-4-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)-4-hydroxybut-1-en-1-yl)-3-hydroxyphenyl)-2-methylpropanoate

To a solution of 1-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)but-3-en-1-ol (320 mg, 1.17 mmol) and (2S, 3R)-methyl 3-cyclopropyl-3-(3-hydroxy-4-iodophenyl)-2-methyl-propanoate (280 mg, 0.778 mmol) in Toluene (10 mL) were added *N*-cyclohexyl-*N*-methylcyclohexanamine (229 mg, 1.17 mmol) and *t*-buxhos palladacycle (54.0 mg, 0.0778 mmol) under a N₂ atmosphere. The reaction mixture was stirred for 2 hours at 100 °C, then cooled to room temperature and concentrated *in vacuo* to give a residue. The residue was purified via silica gel chromatography (eluted from PE: EtOAc = 100: 1 to 5: 1, v/v) to give the title compound. ¹H NMR (400 MHz, CDCl₃): δ = 8.80 (s, 1H), 8.12 (s, 1H), 7.97 (d, *J* = 8.4 Hz, 1H), 7.51 (d, *J* = 8.0 Hz, 1H), 7.20 (d, *J* = 8.0 Hz, 1H), 6.85 (d, *J* = 5.2 Hz, 1H), 6.68-6.58 (m, 3H), 6.18-6.14 (m, 1H), 5.02 (s, 1H), 3.97 (s, 3H), 3.73 (s, 3H), 2.86-2.71 (m, 3H), 1.88-1.83 (m, 1H), 1.01 (s, 1H), 0.95 (d, *J* = 6.8 Hz, 3H), 0.57-0.55 (m, 1H), 0.33-0.21 (m, 2H), 0.01--0.03 (m, 1H).


Step D: (2S,3R)-methyl 3-cyclopropyl-3-(4-(4-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)-4-hydroxybutyl)-3-hydroxyphenyl)-2-methylpropanoate

To a solution of (2S,3R)-methyl 3-cyclopropyl-3-(4-((Z)-4-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)-4-hydroxybut-1-en-1-yl)-3-hydroxyphenyl)-2-methylpropanoate (290 mg, 0.572 mmol) in MeOH (10 mL) was added Pd(OH)₂/C (8.00 mg, 0.0572 mmol, 20%). The reaction mixture was stirred for 3 hours at room temperature under a H₂ balloon. Then the reaction mixture was filtered, and the filtrate was concentrated *in vacuo* to give the title compound, which was used in the next step.

Step E: (2S, 3R)-methyl 3-cyclopropyl-3-(2-(2-fluoro-4-(2-methoxypyridin-4-yl) phenyl)-2, 3, 4, 5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate

To a solution of (2S,3R)-methyl 3-cyclopropyl-3-(4-(4-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)-4-hydroxybutyl)-3-hydroxyphenyl)-2-methylpropanoate (225 mg, 0.442 mmol) and triphenylphosphine (232 mg, 0.884 mmol) in DCM (3.0 mL) was added dropwise a solution of DIAD (179 mg, 0.884 mmol) in DCM (1.0 mL) at 0 °C under a N₂ atmosphere. The reaction mixture was stirred for 2 hours at room temperature, then concentrated *in vacuo* to give a residue, which was purified by preparative TLC (PE: EtOAc = 5: 1, v/v) to give the title compound. ¹H NMR (400 MHz, CDCl₃): δ = 8.78 (s, 1H), 8.13 (s, 1H), 8.01 (s, 1H), 7.86 (d, *J* = 7.6 Hz, 1H), 7.13 (d, *J* = 6.8 Hz, 1H), 6.86 (d, *J* = 5.2 Hz, 1H), 6.82-6.81 (m, 3H), 4.77 (s, 1H), 3.97 (s, 3H), 3.71 (s, 3H), 2.84 (t, *J* = 27.8 Hz, 1H), 2.83-2.77 (m, 2H), 2.48 (s, 1H), 1.87 (t, *J* = 10.0 Hz, 1H), 1.74-1.70 (m, 1H), 1.03 (s, 1H), 0.94-0.92 (m, 3H), 0.54 (s, 1H), 0.33-0.21 (m, 2H), 0.02 (m, 1H). MS(ESI) *m/z*: 491.3 [M+H]⁺.

Step F: (2S,3R)-methyl 3-cyclopropyl-3-((RS)-2-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate



The (2S,3R)-methyl 3-cyclopropyl-3-((RS)-2-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate (190 mg, 0.386 mmol) was separated by SFC (SFC conditions: Column: Chiralpak AD-3 50*4.6mm I.D., 3 μ m Mobile phase: ethanol (0.05 % DEA) in CO₂ from 5 % to 40 % Flow rate: 4 mL/min Wavelength: 220 nm) to give (2S,3R)-methyl 3-cyclopropyl-3-((RS)-2-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate; and (2S,3R)-methyl 3-cyclopropyl-3-((RS)-2-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate.


Step G: (2S,3R)-3-cyclopropyl-3-(2-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid

To a solution of compound (2S,3R)-methyl 3-cyclopropyl-3-((RS)-2-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoate (90.0 mg, 0.184 mmol) in THF/MeOH/H₂O (1.0 mL/1.0 mL/1.0 mL) was added LiOH (44.0 mg, 1.84 mmol). The reaction mixture was stirred at 50 °C for 16 hours, then cooled to room temperature. The reaction mixture was acidified with HCl (1mol/L) to pH 2 and extracted with EtOAc (2 mL*3). The combined organic layers were concentrated *in vacuo* to afford a residue, which was purified by preparative HPLC (on a GILSON 281 instrument fitted with a Phenomenex Gemini C₁₈ (250*21.2 mm * 5 μ m) using Mobile phase A: water (neutral), mobile phase B: acetonitrile. Gradient: 33-63 % B, 0-10 min; 100 % B, 10.5-12.5 min; 5 % B, 13-15 min) to give the title compound. ¹H NMR (400 MHz, CDCl₃): δ = 8.13 (d, *J* = 1.2 Hz, 1H), 7.64-7.58 (m, 3H), 7.12 (d, *J* = 7.6 Hz, 1H), 6.88-6.83 (m, 3H), 4.69 (d, *J* = 10.8Hz, 1H), 3.98 (s, 3H), 2.97 (t, *J* = 12.8 Hz, 1H), 2.82-2.79 (m, 2H), 2.21-2.11 (m, 3H), 1.94 (t, *J* = 10.0 Hz, 1H), 1.72-1.69

(m, 1H), 1.10-1.08 (m, 1H), 0.99 (d, $J = 6.8$ Hz, 3H), 0.63-0.61 (m, 1H), 0.37-0.32 (m, 2H), 0.03--0.02 (m, 1H). MS(ESI) m/z : 477.2 $[M+H]^+$.

To a solution of (2S,3R)-3-cyclopropyl-3-(2-(2-fluoro-4-(2-methoxypyridin-4-yl)phenyl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid (68.9 mg) in MeCN (0.5 mL) was added a solution of NaOH (aq. 172 mg, 0.5 M). The mixture was stirred for 1 hour at room temperature. Then the reaction mixture was lyophilized to give the sodium salt of the title compound.


Table 4 Example 8 was prepared in a similar manner to Example 7 using the appropriate starting materials

Example	Structure	MW	Compound Name	LC/MS (ESI) observed $[M+1]^+$
8		476.55	(2S,3R)-3-cyclopropyl-3-((R or S)-2-(5'-fluoro-2'-methoxy-[3,4'-bipyridin]-6-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-8-yl)-2-methylpropanoic acid	477.2

*stereoisomer at this position

EXAMPLE 9

3-(2-(2',6'-dimethyl-[1,1'-biphenyl]-3-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-7-yl)propanoic acid



Step 1: 2',6'-dimethyl-[1,1'-biphenyl]-3-carbaldehyde

To a degassed solution of 3-bromobenzaldehyde (500 mg, 2.70 mmol), (2,6-dimethylphenyl)boronic acid (608 mg, 4.05 mmol), and $Pd(PPh_3)_4$ (156 mg, 0.14 mmol) in dioxane (5 mL) was added

K₂CO₃ (2N, 4.05 mL, 8.11 mmol). The reaction was heated to 100°C. After 16 h, the mixture was poured into NH₄Cl (sat, 25 mL) and extracted with EtOAc (2 x 25 mL). The combined organic layers were dried over MgSO₄ and concentrated. The resulting residue was purified by HPLC (ISCO 40 gram silica gel cartridge, 0 to 40% EtOAc/Hex) to give the title compound.

Step 2: 1-(2',6'-dimethyl-[1,1'-biphenyl]-3-yl)but-3-en-1-ol

To a solution of 2',6'-dimethyl-[1,1'-biphenyl]-3-carbaldehyde (850 mg, 4.04 mmol) in THF (10 mL) at 0°C was added allyl grignard (1M in THF, 4.85 mL). After 30 minutes, the reaction was quenched by the slow addition of 25 mL NH₄Cl (saturated, aqueous). The mixture was extracted with EtOAc (2 x 25mL). The combined organic layers were dried over MgSO₄ and concentrated. The resulting residue was purified by HPLC (40 gram silica gel ISCO cartridge, 0 to 40% EtOAc/hex) to give the title compound.

Step 3: 3'-(1-(4-bromo-2-vinylphenoxy)but-3-en-1-yl)-2,6-dimethyl-1,1'-biphenyl

To a solution of 1-(2',6'-dimethyl-[1,1'-biphenyl]-3-yl)but-3-en-1-ol (100 mg, 0.396 mmol), 4-bromo-2-vinylphenol (79 mg, 0.396 mmol), and triphenylphosphine (208 mg, 0.793 mmol) in toluene (4 mL) was slowly added DIAD (0.154 mL, 0.793 mmol) dropwise. After 16 h, the reaction was filtered and then concentrated. The resulting residue was purified by HPLC (40 gram silica gel ISCO cartridge, 0 to 40% EtOAc/hex) to give the title compound.

Step 4: 7-bromo-2-(2',6'-dimethyl-[1,1'-biphenyl]-3-yl)-2,3-dihydrobenzo[b]oxepine

To a solution of 3'-(1-(4-bromo-2-vinylphenoxy)but-3-en-1-yl)-2,6-dimethyl-1,1'-biphenyl (28 mg, 0.065 mmol) in DCM (6 mL) was added Hoveyda-Grubbs Catalyst 2nd Generation (CAS # 301224-40-8, 4 mg, 0.006 mmol) and the reaction was heated to reflux. After 1 hour, the reaction was cooled to room temperature, and the volatiles were removed *in vacuo*. The resulting residue was purified by HPLC (40 gram silica gel ISCO cartridge, 0 to 40% EtOAc/hex) to give the title compound.

Step 5: ethyl 3-(2-(2',6'-dimethyl-[1,1'-biphenyl]-3-yl)-2,3-dihydrobenzo[b]oxepin-7-yl)propanoate


To a degassed solution of 7-bromo-2-(2',6'-dimethyl-[1,1'-biphenyl]-3-yl)-2,3-dihydrobenzo[b]oxepine (25 mg, 0.062 mmol) and Sphos Pd G2 (CAS # 1028206-58-7, 4.7 mg, 0.006 mmol) in THF (3 mL) was added (3-ethoxy-3-oxopropyl)zinc(II) bromide (Rieke Metals, Inc., 0.5 M solution in THF, 0.370 mL, 0.185 mmol). The reaction was heated to 70 °C in a sealed vial with a teflon screw cap. After 16 h, the reaction was cooled to room temperature and then poured into NH₄Cl (saturated, 25 mL). The mixture was extracted with EtOAc (2 x 25 mL). The combined organic layers were dried over MgSO₄ and concentrated. The resulting residue was purified by HPLC (40 gram silica gel ISCO cartridge, 0 to 40% EtOAc/hex) to give the title compound.

Step 6: 3-(2-(2',6'-dimethyl-[1,1'-biphenyl]-3-yl)-2,3-dihydrobenzo[b]oxepin-7-yl)propanoic acid

To a slurry of ethyl 3-(2-(2',6'-dimethyl-[1,1'-biphenyl]-3-yl)-2,3-dihydrobenzo[b]oxepin-7-

yl)propanoate (17 mg, 0.04 mmol) and THF (1 mL) in LiOH (10 mg, 0.40 mmol) was added MeOH (1 ml) and water (1 ml). The resulting mixture was heated to 60 °C. After 1 hour, the reaction mixture was poured into 1N HCl (10 mL) and extracted with EtOAc (2 x 10 mL). The combined organic layers were dried over MgSO₄ and concentrated. The resulting residue was purified by HPLC (40 gram silica gel ISCO cartridge, 0 to 100% EtOAc/hex) to give the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.48 (m, 2H), 7.22 (s, 1H), 7.18 (m, 1H), 7.15 (m, 3H), 7.03 (s, 1H), 6.99 (m, 2H), 6.40 (d, 1H), 6.01 (dd, 1H), 5.00 (d, 1H), 3.05 (dd, 1H), 2.95 t, 2H), 2.81 (dd, 1H), 2.70 (t, 2H), 2.07 (s, 6H); (m/z): 381.2 (M-18+H)⁺.

Step 7: 3-(2-(2',6'-dimethyl-[1,1'-biphenyl]-3-yl)-2,3,4,5-tetrahydrobenzo[b]oxepin-7-yl)propanoic acid



To a solution of 3-(2-(2',6'-dimethyl-[1,1'-biphenyl]-3-yl)-2,3-dihydrobenzo[b]oxepin-7-yl)propanoic acid (10 mg, 0.025 mmol) in EtOAc (1 mL) was added Pd(OH)₂ (1 mg, 20 wt%, 0.001 mmol). The mixture was vigorously stirred under a balloon of hydrogen gas. After 2 hours, the reaction was filtered through CeliteTM and the filtrate concentrated to give the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.48 (br, 2H), 7.21 (s, 1H), 7.10-7.20 (m, 4H), 7.01 (s, 1H), 6.98 (s, 2H), 4.62 (d, 1H), 3.00 (t, 1H), 2.92 (t, 2H), 2.78 (dd, 1H), 2.70 (t, 2H), 2.24-2.10 (m, 2H), 2.04 (s, 6H), 1.62 (q, 2H); (m/z): 383.4 (M-18+H)⁺.

EXAMPLE OF A PHARMACEUTICAL COMPOSITION

As a specific embodiment of an oral pharmaceutical composition, a 100 mg potency tablet is composed of 100 mg of any one of Examples, 268 mg microcrystalline cellulose, 20 mg of croscarmellose sodium, and 4 mg of magnesium stearate. The active, microcrystalline cellulose, and croscarmellose are blended first. The mixture is then lubricated by magnesium stearate and pressed into tablets.

BIOLOGICAL ASSAYS

Generation of GPR40-Expressing Cells:

Human and mouse GPR40 stable cell-lines were generated in CHO cells stably expressing NFAT BLA (Beta-lactamase). A human GPR40 stable cell-line was generated in HEK cells stably expressing the aequorin expressing reporter. The expression plasmids were transfected using

lipofectamine (Life Technologies) following manufacturer's instructions. Stable cell-lines were generated following drug selection.

FLIPR Assays:

FLIPR (Fluorimetric Imaging Plate Reader, Molecular Devices) assays were performed to measure agonist-induced calcium mobilization of the stable clones. For the FLIPR assay, one day before assay, GPR40/CHO NFAT BLA cells were seeded into black-wall-clear-bottom 384-well plates (Costar) at 1.4×10^4 cells / 20 μ L medium / well. The cells were incubated with 20 μ L / well of the assay buffer (HBSS, 0.1 % BSA, 20 mM HEPES, 2.5 mM probenecid, pH 7.4) containing 8 μ M fluo-4,AM, 0.08 % pluronic acid at room temperature for 100 minutes. Fluorescence output was measured using FLIPR. Compounds were dissolved in DMSO and diluted to desired concentrations with assay buffer. 13.3 μ L/well of compound solution was added. The FLIPR Assay EC₅₀ values for specific compounds are listed in Table I.

Inositol Phosphate Turnover (IP1) Assay:

The assay is performed in 384-well format. HEK cells stably expressing human GPR40 are plated at 15,000 cells per well in growth medium (DMEM/10% fetal calf serum). Cell plates are then incubated 16 hours at 37 degrees in a 5% CO₂ incubator.

Measurement of Inositol Phosphate Turnover (IP1) is performed using the CisBio IP-One kit (Part number 62IPAPEB). After the 16 hour incubation, the cells are washed with HEPES buffer and 10 μ L of stimulation buffer (prepared as described in the kit) is added to each well. In a separate plate, compounds are diluted in DMSO (400-fold over the final concentration in the assay well) and 25 μ L is acoustically transferred to the appropriate well in the assay cell plate. The plates are then incubated for 60 minutes at 37 degrees. 10 μ L of detection buffer (also prepared as described in the IP-One kit) is added to each well and the plates are incubated for 60 minutes in the dark. The plates are then read in a Perkin Elmer EnVision or equivalent reader able to measure FRET. Fluorescent ratio of emission at 665 and 620 nm is then converted to IP1 concentration by back calculating from an IP1 standard curve prepared at the time of the assay. The Inositol Phosphate Turnover (IP1) Assay EC₅₀ values for specific compounds are listed in Table I.

The compounds of the present invention, including the compounds in Examples 1-9, have either EC₅₀ values less than 10 micromolar (μ M) in the FLIPR assay described above, or have EC₅₀ values less than 6500 nanomolar (nM) in the Inositol Phosphate Turnover (IP1) Assay described above.

Table I.

Example No.	Stereoisomers	Human FLIPR EC ₅₀ , nM or % activation	Human IP1 Assay, EC ₅₀ , nM
1	Isomer A	ND	3.6
2	Isomer B	ND	14
3	Isomer A	ND	13
4	Isomer B	ND	27
5	Isomer A	ND	3.1
6	Isomer B	ND	5.2
7	Isomer A	ND	3.0
8	Isomer B	ND	29
9	Mix of isomers	3300	ND

*Isomer A = first eluting peak from chiral column; Isomer B = second eluting peak from chiral column; "ND" is not determined

In Vivo Studies:


5 Male C57BL/6N mice (7-12 weeks of age) are housed 10 per cage and given access to normal diet rodent chow and water *ad libitum*. Mice are randomly assigned to treatment groups and fasted 4 to 6 h. Baseline blood glucose concentrations are determined by glucometer from tail nick blood. Animals are then treated orally with vehicle (0.25% methylcellulose) or test compound. Blood glucose concentration is measured at a set time point after treatment (t = 0 min) and mice are
 10 then intraperitoneally-challenged with dextrose (2 g/kg). One group of vehicle-treated mice is challenged with saline as a negative control. Blood glucose levels are determined from tail bleeds taken at 20, 40, 60 min after dextrose challenge. The blood glucose excursion profile from t = 0 to t = 60 min is used to integrate an area under the curve (AUC) for each treatment. Percent inhibition values for each treatment are generated from the AUC data normalized to the saline-challenged
 15 controls.

While the invention has been described and illustrated with reference to certain particular embodiments thereof, those skilled in the art will appreciate that various adaptations, changes, modifications, substitutions, deletions, or additions of procedures and protocols may be made without departing from the scope of the
 20 invention. For example, effective dosages other than the particular dosages as set forth herein above may be applicable as a consequence of variations in responsiveness of the mammal being treated for any of the indications with the compounds of the invention indicated above. The specific pharmacological responses observed may vary according to and depending upon the particular active compounds selected or whether

there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention.

CLAIMS

1. A compound of structural formula I:



5

or a pharmaceutically acceptable salt thereof; wherein

"a" is a single bond or a double bond, provided that if "a" is a double bond, then R^5 and R^6 are absent and Q is selected from the group consisting of: $C-R^4$, $-C-OC_{1-6}alkyl$, and N, and further

10 provided that if "a" is a single bond, then R^5 and R^6 are present, and Q is selected from the group consisting of:

- (1) oxygen,
- (2) sulfur,
- (3) $-CR^4R^6$,
- 15 (4) $S(O)_2$,
- (5) $C=O$,
- (6) $-C(R^9)OC_{1-6}alkyl$, and
- (7) $-NRC$;

20 T is selected from the group consisting of:

- (1) CH,
- (2) N, and
- (3) N-oxide;

U is selected from the group consisting of:

- 25 (1) CR^1 ,
- (2) N, and
- (3) N-oxide;

V is selected from the group consisting of:

- (1) CR^2 ,
- 30 (2) N, and

- (3) N-oxide;

W is selected from the group consisting of:

- (1) CH,
(2) N, and
5 (3) N-oxide,

provided that no more than two of T, U, V and W are selected from N and N-oxide, further provided that if both T and W are N or N-oxide, then R³ is absent, and further provided that both U and V are not N or N-oxide;

10 X is selected from the group consisting of:

- (1) oxygen,
(2) sulfur,
(3) -CR⁹R⁹,
(4) S(O)₂,
15 (5) C=O,
(6) -C(R⁹)OC₁₋₆alkyl, and
(7) -NR^c,

provided that if Q is oxygen, sulfur or -NR^c, then X is not oxygen, sulfur or -NR^c, further provided that if Q is C=O, then X is not C=O or S(O)₂, and further provided that if Q is S(O)₂, then X is not

20 S(O)₂, C=O, oxygen or sulfur;

Y is selected from the group consisting of:

- (1) oxygen,
(2) sulfur,
25 (3) -CR⁹R⁹,
(4) S(O)₂,
(5) C=O,
(6) -C(R⁹)OC₁₋₆alkyl, and
(7) -NR^c,

30 provided that if X is oxygen, sulfur or -NR^c, then Y is not oxygen, sulfur or -NR^c, further provided that if X is C=O, then Y is not C=O or S(O)₂, and further provided that if X is S(O)₂, then Y is not S(O)₂, C=O, oxygen or sulfur;

Z is selected from the group consisting of:

- 35 (1) oxygen,
(2) sulfur,

- (3) $-\text{CR}^9\text{R}^9$,
- (4) $\text{S}(\text{O})_2$,
- (5) $\text{C}=\text{O}$,
- (6) $-\text{C}(\text{R}^9)\text{OC}_{1-6}\text{alkyl}$, and
- 5 (7) $-\text{NRC}$;

A is selected from the group consisting of:

- (1) aryl,
- (2) heteroaryl,
- 10 (3) $\text{C}_{3-6}\text{cycloalkyl}$, and
- (4) $\text{C}_{2-5}\text{cycloheteroalkyl}$,

wherein A is unsubstituted or substituted with one to five substituents selected from R^a ;

B is selected from the group consisting of:

- 15 (1) hydrogen,
- (2) aryl,
- (3) aryl-O-,
- (4) aryl- C_{1-10} alkyl-,
- (5) aryl- C_{1-10} alkyl-O-,
- 20 (6) $\text{C}_{3-6}\text{cycloalkyl}$,
- (7) $\text{C}_{3-6}\text{cycloalkyl-}\text{C}_{1-10}\text{alkyl-}$,
- (8) $\text{C}_{3-6}\text{cycloalkyl-}\text{C}_{1-10}\text{alkyl-O-}$,
- (9) $\text{C}_{3-6}\text{cycloalkenyl}$,
- (10) $\text{C}_{3-6}\text{cycloalkenyl-}\text{C}_{1-10}\text{alkyl-}$,
- 25 (11) $\text{C}_{3-6}\text{cycloalkenyl-}\text{C}_{1-10}\text{alkyl-O-}$,
- (12) $\text{C}_{2-5}\text{cycloheteroalkyl}$,
- (13) $\text{C}_{3-6}\text{cycloheteroalkyl-}\text{C}_{1-10}\text{alkyl-}$,
- (14) $\text{C}_{3-6}\text{cycloheteroalkyl-}\text{C}_{1-10}\text{alkyl-O-}$,
- (15) heteroaryl,
- 30 (16) heteroaryl-O-,
- (17) heteroaryl- C_{1-10} alkyl-, and
- (18) heteroaryl- C_{1-10} alkyl-O-,

wherein B is unsubstituted or substituted with one to five substituents selected from R^b ;

35 R^1 and R^2 are each independently selected from:

- (1) a bond,

- (2) hydrogen,
- (3) halogen,
- (4) -OR^k,
- (5) -CN,
- 5 (6) -C₁₋₆alkyl,
- (7) -C₃₋₆cycloalkyl,
- (8) C₃₋₆cycloalkyl-C₁₋₃alkyl-,
- (9) -C₂₋₆cycloheteroalkyl, and
- (10) C₂₋₆cycloheteroalkyl-C₁₋₃alkyl-,

10 wherein each alkyl, cycloalkyl and cycloheteroalkyl is unsubstituted or substituted with one to three substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷, or R¹ and R² together with the atom(s) to which they are attached form a C₃₋₆cycloalkyl ring or a C₂₋₅cycloheteroalkyl ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R^g, wherein each R¹ and R² is unsubstituted or substituted with one to three
 15 substituents selected from R^L, and wherein one of R¹ and R² is substituted with R⁷;

R³ is absent or selected from the group consisting of:

- (1) hydrogen,
- (2) halogen,
- 20 (3) -OR^e,
- (4) -CN,
- (5) -C₁₋₆alkyl,
- (6) -C₃₋₆cycloalkyl, and
- (7) C₃₋₆cycloalkyl-C₁₋₃alkyl-,

25 wherein each alkyl and cycloalkyl is unsubstituted or substituted with one to three substituents selected from Rⁱ;

R⁴ is selected from the group consisting of:

- (1) hydrogen,
- 30 (2) halogen,
- (3) OR^e,
- (4) C₀₋₅alkylNR^cR^d,
- (5) C₁₋₆alkyl,
- (6) C₁₋₆alkyl-O-,
- 35 (7) C₃₋₆cycloalkyl,
- (8) C₃₋₆cycloalkyl-O-,

- (9) C₃₋₆cycloalkyl-C₁₋₁₀alkyl-,
(10) C₃₋₆cycloalkyl-C₁₋₁₀alkyl-O-,
(11) C₂₋₅cycloheteroalkyl,
(12) C₂₋₅cycloheteroalkyl-O-,
5 (13) C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-,
(14) C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-O-,
(15) aryl,
(16) aryl-O-,
(17) aryl-C₁₋₁₀alkyl-,
10 (18) heteroaryl,
(19) heteroaryl-O-, and
(20) heteroaryl-C₁₋₁₀alkyl-,

wherein each alkyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or substituted with one to three substituents selected from R_j,

15 provided that when R⁴ is selected from the group consisting of:

- (1) ORe,
(2) C₀alkyl-NR^cR^d,
(3) C₁₋₆alkyl-O-,
(4) C₃₋₆cycloalkyl-O-,
20 (5) C₃₋₆cycloalkyl-C₁₋₁₀alkyl-O-,
(6) C₂₋₅cycloheteroalkyl-O-,
(7) C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-O-,
(8) aryl-O-, and
(9) heteroaryl-O-,

25 then X is selected from the group consisting of:

- (1) -CR⁹R⁹,
(2) C=O, and
(3) -C(R⁹)OC₁₋₆alkyl;

30 R⁵ is absent or selected from the group consisting of:

- (1) hydrogen,
(2) -C₁₋₆alkyl, and
(3) -C₃₋₆cycloalkyl,

wherein each alkyl and cycloalkyl is unsubstituted or substituted with one to three substituents
35 selected from R_j;


R⁶ is absent or selected from the group consisting of:

- (1) hydrogen,
- (2) -C₁₋₆alkyl, and
- (3) -C₃₋₆cycloalkyl,

5 wherein each alkyl and cycloalkyl is unsubstituted or substituted with one to three substituents selected from R_J;

R⁷ is selected from the group consisting of:

- (1) -CO₂R⁸,
- 10 (2) -C₁₋₆alkyl-CO₂R⁸,
- (3) -C₁₋₆alkyl-CONHSO₂R^m,
- (4) -C₁₋₆alkyl-SO₂NHCOR^m,
- (5) -C₁₋₆alkyl-tetrazolyl, and
- (6) a cycloheteroalkyl selected from the group consisting of:



R⁸ is selected from the group consisting of:

- 20 (1) hydrogen,
- (2) -C₁₋₆alkyl,
- (3) -C₃₋₆cycloalkyl, and
- (4) aryl-C₁₋₆alkyl,

25 wherein each alkyl, cycloalkyl and aryl is unsubstituted or substituted with one to three substituents selected from R_J;

R⁹ is selected from the group consisting of:

- (1) hydrogen,
- (2) -C₁₋₆alkyl,
- (3) halogen,
- (4) -(CH₂)₀₋₅OH,
- (5) -(CH₂)₀₋₅OC₁₋₆alkyl,
- (6) -(CH₂)₀₋₅NR^cR^d,
- (7) -C₃₋₆cycloalkyl, and
- (8) aryl-C₁₋₆alkyl,

wherein CH₂ is unsubstituted or substituted with one or two substituents selected from R^j, and wherein alkyl, cycloalkyl and aryl is unsubstituted or substituted with one to three substituents selected from R^j;

R^a is selected from the group consisting of:

- (1) -C₁₋₆alkyl,
- (2) halogen,
- (3) -C₀₋₆alkyl-OR^e,
- (4) -C₀₋₆alkyl-NR^cS(O)_nR^e,
- (5) -C₀₋₆alkyl-S(O)_nR^e,
- (6) -C₀₋₆alkyl-S(O)_nNR^cR^d,
- (7) -C₀₋₆alkyl-NR^cR^d,
- (8) -C₀₋₆alkyl-C(O)R^e,
- (9) -C₀₋₆alkyl-OC(O)R^e,
- (10) -C₀₋₆alkyl-CO₂R^e,
- (11) -C₀₋₆alkyl-CN,
- (12) -C₀₋₆alkyl-C(O)NR^cR^d,
- (13) -C₀₋₆alkyl-NR^cC(O)R^e,
- (14) -C₀₋₆alkyl-NR^cC(O)OR^e,
- (15) -C₀₋₆alkyl-NR^cC(O)NR^cR^d,
- (16) -CF₃,
- (17) -OCF₃,
- (18) -OCHF₂,
- (19) -C₀₋₆alkyl-aryl,
- (20) -C₀₋₆alkyl-heteroaryl,
- (21) -C₀₋₆alkyl -C₃₋₁₀cycloalkyl,
- (22) -C₀₋₆alkyl -C₃₋₁₀cycloalkenyl, and

(23) -C₀₋₆alkyl -C₂₋₁₀cycloheteroalkyl,

wherein each alkyl, cycloalkyl, cycloalkenyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or substituted with one to five substituents independently selected from: -C₁₋₆alkyl, halogen, OH, -O-C₁₋₆alkyl, -S(O)₂-C₁₋₄alkyl, -CN, -OCHF₂, -OCF₃, -CF₃, and -C₀₋₆alkyl-NR^cR^d;

5

R^b is selected from the group consisting of:

- (1) -C₁₋₁₀alkyl,
- (2) -C₂₋₁₀alkenyl,
- (3) -CF₃,
- 10 (4) halogen,
- (5) -CN,
- (6) -OH,
- (7) -OC₁₋₁₀alkyl,
- (8) -OC₂₋₁₀alkenyl,
- 15 (9) -O(CH₂)_pOC₁₋₁₀alkyl,
- (10) -O(CH₂)_pC₃₋₆cycloalkyl,
- (11) -O(CH₂)_pC₃₋₆ cycloalkyl-C₁₋₁₀alkyl,
- (12) -O(CH₂)_pC₂₋₅cycloheteroalkyl,
- (13) -O(CH₂)_pC₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl,
- 20 (14) -O-aryl,
- (15) -O-heteroaryl,
- (16) -O-aryl-C₁₋₁₀alkyl,
- (17) -O-heteroaryl-C₁₋₁₀alkyl,
- (18) -O(CH₂)_pNR^cS(O)_mRe,
- 25 (19) -O(CH₂)_pS(O)_mRe,
- (20) -O(CH₂)_pS(O)_mNR^cR^d,
- (21) -O(CH₂)_pNR^cR^d,
- (22) -C(O)Re,
- (23) -OC(O)Re,
- 30 (24) -CO₂Re,
- (25) -C(O)NR^cR^d,
- (26) -NR^cC(O)Re,
- (27) -NR^cC(O)ORE,
- (28) -NR^cC(O)NR^cR^d,
- 35 (29) -O(CH₂)_pO-C₃₋₆cycloalkyl,
- (30) -O(CH₂)_pO-C₂₋₅cycloheteroalkyl,

- (31) -OCF₃,
(32) -OCHF₂,
(33) -(CH₂)_pC₃₋₆cycloalkyl,
(34) -(CH₂)_pC₂₋₅cycloheteroalkyl,
5 (35) aryl,
(36) heteroaryl,
(37) aryl-C₁₋₁₀alkyl-, and
(38) heteroaryl-C₁₋₁₀alkyl-,

wherein each CH, CH₂, alkyl, alkenyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is
10 unsubstituted or substituted with one to five substituents independently selected from: -C₁₋₆alkyl,
halogen, OH, -O-C₁₋₆alkyl and -CF₃;

R^c and R^d are each independently selected from the group consisting of:

- (1) hydrogen,
15 (2) C₁₋₁₀alkyl,
(3) C₂₋₁₀alkenyl,
(4) C₃₋₆cycloalkyl,
(5) C₃₋₆ cycloalkyl-C₁₋₁₀alkyl-,
(6) C₂₋₅cycloheteroalkyl,
20 (7) C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-,
(8) aryl,
(9) heteroaryl,
(10) aryl-C₁₋₁₀alkyl-, and
(11) heteroaryl-C₁₋₁₀alkyl-,

25 wherein each alkyl, alkenyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or
substituted with one to three substituents independently selected from R^f,
or R^c and R^d together with the atom(s) to which they are attached form a C₂₋₁₀ cycloheteroalkyl
ring containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R_g,
wherein each R^c and R^d is unsubstituted or substituted with one to three substituents independently
30 selected from R^f;

each R^e is independently selected from the group consisting of:

- (1) hydrogen,
35 (2) -C₁₋₁₀alkyl,
(3) -C₂₋₁₀ alkenyl,
(4) -C₃₋₆ cycloalkyl,

- (5) C₃₋₆ cycloalkyl-C₁₋₁₀alkyl-,
(6) -C₂₋₅cycloheteroalkyl,
(7) C₂₋₅cycloheteroalkyl-C₁₋₁₀alkyl-,
(8) aryl,
5 (9) aryl-C₁₋₁₀alkyl-,
(10) heteroaryl, and
(11) heteroaryl-C₁₋₁₀alkyl-,

wherein each alkyl, alkenyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or substituted with one to three substituents selected from R^h;

each R^f is selected from the group consisting of:

- (1) halogen,
(2) C₁₋₁₀alkyl,
(3) -OH,
15 (4) -O-C₁₋₄alkyl,
(5) -S(O)_m-C₁₋₄alkyl,
(6) -CN,
(7) -CF₃,
(8) -OCHF₂, and
20 (9) -OCF₃,

wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from: -OH, halogen, C₁₋₆alkyl, cyano and S(O)₂C₁₋₆alkyl;

each R^g is independently selected from the group consisting of:

- 25 (1) hydrogen,
(2) -C(O)R^e, and
(3) -C₁₋₁₀alkyl,

wherein alkyl is unsubstituted or substituted with one to five halogens;

each R^h is independently selected from the group consisting of:

- (1) halogen,
(2) C₁₋₁₀alkyl,
(3) -OH,
(4) -O-C₁₋₄alkyl,
35 (5) -S(O)_m-C₁₋₄alkyl,
(6) -CN,

- (7) $-\text{CF}_3$,
- (8) $-\text{OCHF}_2$, and
- (9) $-\text{OCF}_3$,

wherein alkyl is unsubstituted or substituted with one to three substituents independently selected
 5 from: $-\text{OH}$, halogen, C_{1-6} alkyl, cyano and $\text{S}(\text{O})_2\text{C}_{1-6}$ alkyl;

each R^{i} is independently selected from the group consisting of:

- (1) $-\text{C}_{1-6}$ alkyl,
- (2) $-\text{OR}^{\text{e}}$,
- 10 (3) $-\text{NR}^{\text{c}}\text{S}(\text{O})_{\text{m}}\text{R}^{\text{e}}$,
- (4) halogen,
- (5) $-\text{S}(\text{O})_{\text{m}}\text{R}^{\text{e}}$,
- (6) $-\text{S}(\text{O})_{\text{m}}\text{NR}^{\text{c}}\text{R}^{\text{d}}$,
- (7) $-\text{NR}^{\text{c}}\text{R}^{\text{d}}$,
- 15 (8) $-\text{C}(\text{O})\text{R}^{\text{e}}$,
- (9) $-\text{OC}(\text{O})\text{R}^{\text{e}}$,
- (10) $-\text{CO}_2\text{R}^{\text{e}}$,
- (11) $-\text{CN}$,
- (12) $-\text{C}(\text{O})\text{NR}^{\text{c}}\text{R}^{\text{d}}$,
- 20 (13) $-\text{NR}^{\text{c}}\text{C}(\text{O})\text{R}^{\text{e}}$,
- (14) $-\text{NR}^{\text{c}}\text{C}(\text{O})\text{OR}^{\text{e}}$,
- (15) $-\text{NR}^{\text{c}}\text{C}(\text{O})\text{NR}^{\text{c}}\text{R}^{\text{d}}$,
- (16) $-\text{CF}_3$,
- (17) $-\text{OCF}_3$,
- 25 (18) $-\text{OCHF}_2$,
- (19) $-\text{C}_{3-6}$ cycloalkyl, and
- (20) $-\text{C}_{2-5}$ cycloheteroalkyl;

each R^{j} is independently selected from the group consisting of:

- 30 (1) $-\text{C}_{1-6}$ alkyl,
- (2) $-\text{OR}^{\text{e}}$,
- (3) $-\text{NR}^{\text{c}}\text{S}(\text{O})_{\text{m}}\text{R}^{\text{e}}$,
- (4) halogen,
- (5) $-\text{S}(\text{O})_{\text{m}}\text{R}^{\text{e}}$,
- 35 (6) $-\text{S}(\text{O})_{\text{m}}\text{NR}^{\text{c}}\text{R}^{\text{d}}$,
- (7) $-\text{NR}^{\text{c}}\text{R}^{\text{d}}$,

- (8) $-\text{C}(\text{O})\text{Re}$,
 (9) $-\text{OC}(\text{O})\text{Re}$,
 (10) $-\text{CO}_2\text{Re}$,
 (11) $-\text{CN}$,
 5 (12) $-\text{C}(\text{O})\text{NR}^{\text{c}}\text{R}^{\text{d}}$,
 (13) $-\text{NR}^{\text{c}}\text{C}(\text{O})\text{Re}$,
 (14) $-\text{NR}^{\text{c}}\text{C}(\text{O})\text{OR}^{\text{e}}$,
 (15) $-\text{NR}^{\text{c}}\text{C}(\text{O})\text{NR}^{\text{c}}\text{R}^{\text{d}}$,
 (16) $-\text{CF}_3$,
 10 (17) $-\text{OCF}_3$,
 (18) $-\text{OCHF}_2$,
 (19) $-\text{C}_{3-6}\text{cycloalkyl}$, and
 (20) $-\text{C}_{2-5}\text{cycloheteroalkyl}$;

15 each R^{k} is independently selected from the group consisting of:

- (1) hydrogen,
 (2) $-\text{C}_{1-6}\text{alkyl}$,
 (3) $-\text{C}_{1-6}\text{alkyl}-\text{SO}_2\text{C}_{1-6}\text{alkyl}$,
 (4) $-\text{CF}_3$, and
 20 (5) $-\text{CHF}_2$,

wherein each alkyl is unsubstituted or substituted with one to three substituents independently selected from: $-\text{OH}$, $-\text{OC}_{1-6}\text{alkyl}$, halogen, cyano, and $-\text{S}(\text{O})_2\text{C}_{1-6}\text{alkyl}$;

each R^{L} is independently selected from the group consisting of:

- 25 (1) $-\text{CO}_2\text{C}_{1-6}\text{alkyl}$,
 (2) $-\text{C}_{1-10}\text{alkyl}$,
 (3) $-\text{C}_{2-10}\text{alkenyl}$,
 (4) $-\text{C}_{2-10}\text{alkynyl}$,
 (5) $-\text{C}_{3-6}\text{cycloalkyl}$,
 30 (6) $-\text{C}_{2-6}\text{cycloheteroalkyl}$,
 (7) aryl, and
 (8) heteroaryl,

wherein each alkyl, alkenyl, alkynyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl is unsubstituted or substituted with one to four substituents independently selected from $\text{C}_{1-6}\text{alkyl}$,
 35 halogen, and $-\text{OC}_{1-6}\text{alkyl}$;

each R^m is independently selected from the group consisting of:

- (1) $-C_{1-10}alkyl$,
- (2) $-C_{2-10}alkenyl$,
- (3) $-C_{3-6}cycloalkyl$,
- 5 (4) $C_{3-6}cycloalkyl-C_{1-10}alkyl-$,
- (5) $-C_{2-5}cycloheteroalkyl$,
- (6) $C_{2-5}cycloheteroalkyl-C_{1-10}alkyl-$,
- (7) $aryl$,
- (8) $heteroaryl$,
- 10 (9) $aryl-C_{1-10}alkyl-$, and
- (10) $heteroaryl-C_{1-10}alkyl-$;

each n is independently selected from: 0, 1 or 2;

each m is independently selected from: 0, 1 or 2;

15 each p is independently selected from: 0, 1, 2, 3, 4, 5 or 6; and

each r is independently selected from: 0, 1, 2 or 3.

2. The compound according to Claim 1 wherein "a" is a single bond; R^5 and R^6 are present, and Q is selected from the group consisting of:

- 20 (1) oxygen,
- (2) sulfur,
- (3) $-CR^4R^6$,
- (4) $S(O)_2$,
- (5) $C=O$,
- 25 (6) $-C(R^9)OC_{1-6}alkyl$, and
- (7) $-NRC$;

or a pharmaceutically acceptable salt thereof.

3. The compound according to Claim 1 wherein T is CH; U is CR^1 ; V is CR^2 ; and W is
30 CH; or a pharmaceutically acceptable salt thereof.

4. The compound according to Claim 1 wherein Q is $-CR^4R^6$; X is $-CR^9R^9$; and Y is $-CR^9R^9$; or a pharmaceutically acceptable salt thereof.

35 5. The compound according to Claim 1 wherein Z is oxygen; or a pharmaceutically acceptable salt thereof.

6. The compound according to Claim 1 wherein A is selected from the group consisting of:

- (1) aryl,
- (2) heteroaryl, and
- (3) C₂₋₅cycloheteroalkyl,

wherein each aryl, heteroaryl and cycloheteroalkyl is unsubstituted or substituted with one to five substituents selected from R^a; or a pharmaceutically acceptable salt thereof.

7. The compound according to Claim 1 wherein A is selected from the group consisting of:

- (1) phenyl,
- (2) pyridyl, and
- (3) azetidine,

wherein each phenyl, pyridyl and azetidine is unsubstituted or substituted with one to four substituents selected from R^a; or a pharmaceutically acceptable salt thereof.

8. The compound according to Claim 1 wherein B is selected from the group consisting of:

- (1) aryl,
- (2) aryl-C₁₋₁₀ alkyl-, and
- (3) heteroaryl,

wherein each alkyl, aryl and heteroaryl is unsubstituted or substituted with one to five substituents selected from R^b; or a pharmaceutically acceptable salt thereof.

9. The compound according to Claim 1 wherein B is selected from the group consisting of:

- (1) phenyl,
- (2) -CH₂-phenyl, and
- (3) pyridyl,

wherein each phenyl and pyridyl is unsubstituted or substituted with one to five substituents selected from R^b; or a pharmaceutically acceptable salt thereof.

10. The compound according to Claim 1 wherein R¹ and R² are each independently selected from:

- (1) hydrogen, and

(2) -C₁₋₆alkyl,
wherein one of R¹ and R² is -C₁₋₆alkyl substituted with R⁷, and wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from R^L; or a pharmaceutically acceptable salt thereof.

5

11. The compound according to Claim 1 wherein R¹ and R² are each independently selected from:

(1) hydrogen, and

(2) ethyl,


10 wherein one of R¹ and R² is ethyl substituted with R⁷, and wherein ethyl is unsubstituted or substituted with one to three substituents independently selected from R^L; or a pharmaceutically acceptable salt thereof.

12. The compound according to Claim 1 wherein
15 R³ is absent or hydrogen;
R⁴ is hydrogen;
R⁵ is absent or hydrogen;
R⁶ is absent or hydrogen; and
R⁹ is hydrogen;
20 or a pharmaceutically acceptable salt thereof.

13. The compound according to Claim 1 wherein R⁷ is -CO₂R⁸; or a pharmaceutically acceptable salt thereof.

25 14. The compound according to Claim 1 wherein R⁸ is hydrogen; or a pharmaceutically acceptable salt thereof.

15. The compound according to Claim 1 of structural Formula Ia:



wherein

"a" is a single bond;

Q is $-\text{CR}^4\text{R}^6$;

5 X is $-\text{CR}^9\text{R}^9$;

Y is $-\text{CR}^9\text{R}^9$;

Z is oxygen;

A is selected from the group consisting of:

- (1) aryl,
- 10 (2) heteroaryl, and
- (3) C₂₋₅cycloheteroalkyl,

wherein each aryl, heteroaryl and cycloheteroalkyl is unsubstituted or substituted with one to five substituents selected from R^a;

B is selected from the group consisting of:

- 15 (1) aryl,
- (2) aryl-C₁₋₁₀ alkyl-, and
- (3) heteroaryl,

wherein each alkyl, aryl and heteroaryl is unsubstituted or substituted with one to five substituents selected from R^b;

20 R¹ and R² are each independently selected from:

- (1) hydrogen, and
- (2) -C₁₋₆alkyl,

wherein one of R¹ and R² is -C₁₋₆alkyl substituted with R⁷, and wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from R^L;

25 R³ is absent or hydrogen;

R⁴ is hydrogen;

R⁵ is absent or hydrogen;

R⁶ is absent or hydrogen;

R⁷ is -CO₂R⁸;

R⁸ is hydrogen;

R⁹ is hydrogen;


each R^L is independently selected from the group consisting of: -C₁₋₁₀alkyl, and -C₃₋₆cycloalkyl, wherein each alkyl and cycloalkyl is unsubstituted or substituted with one to four substituents

5 independently selected from C₁₋₆alkyl, halogen, and -OC₁₋₆alkyl; and

each r is independently selected from: 0, 1, 2 or 3;

or a pharmaceutically acceptable salt thereof.

10 16. The compound according to Claim 1 of structural Formula Ia:



wherein

"a" is a single bond;

15 Q is -CR⁴R⁶;

X is -CR⁹R⁹;

Y is -CR⁹R⁹;

Z is oxygen;

A is selected from the group consisting of:

- 20 (1) phenyl,
(2) pyridyl, and
(3) azetidine,

wherein each phenyl, pyridyl and azetidine is unsubstituted or substituted with one to four substituents selected from R^a;

25 B is selected from the group consisting of:

- (1) phenyl,
(2) -CH₂-phenyl, and
(3) pyridyl,

wherein each phenyl and pyridyl is unsubstituted or substituted with one to five substituents selected from R^b;

R¹ and R² are each independently selected from:

(1) hydrogen, and

5 (2) ethyl,

wherein one of R¹ and R² is ethyl substituted with R⁷, and wherein ethyl is unsubstituted or substituted with one to three substituents independently selected from R^L;

R³ is absent or hydrogen;

R⁴ is hydrogen;

10 R⁵ is hydrogen;

R⁶ is hydrogen;

R⁷ is -CO₂R⁸;

R⁸ is hydrogen;


R⁹ is hydrogen;


15 each R^L is independently selected from the group consisting of: -CH₃, and -cyclopropyl; and

each r is independently selected from: 0, 1, 2 or 3;

or a pharmaceutically acceptable salt thereof.

20 17. The compound according to Claim 16 selected from:





or a pharmaceutically acceptable salt thereof.

5

18. A pharmaceutical composition comprising a compound of Claim 1, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

19. The use of a compound according to Claim 1 for the preparation of a medicament useful for the treatment of a disorder, condition, or disease that may be responsive to the agonism of the G-protein-coupled receptor 40 in a mammal in need thereof.

10

20. The use of a compound of Claim 1, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of Type 2 diabetes.

15

21. A method of treating or preventing a disorder, condition or disease that may be responsive to the agonism of the G-protein-coupled receptor 40 in a patient in need thereof comprising administration of a therapeutically effective amount of a compound according to Claim 1.

20

22. A method of treating type 2 diabetes mellitus in a patient in need of treatment comprising the administration to the patient of a therapeutically effective amount of a compound of Claim 1, or a pharmaceutically acceptable salt thereof.

25

23. A pharmaceutical composition comprising
(1) a compound of Claim 1, or a pharmaceutically acceptable salt thereof;
(2) one or more compounds selected from the group consisting of :

- (a) PPAR gamma agonists and partial agonists;
- (b) biguanides;
- (c) protein tyrosine phosphatase-1B (PTP-1B) inhibitors;
- (d) dipeptidyl peptidase IV (DP-IV) inhibitors;
- (e) insulin or an insulin mimetic;
- (f) sulfonylureas;
- (g) α -glucosidase inhibitors;

(h) agents which improve a patient's lipid profile, said agents being selected from the group consisting of (i) HMG-CoA reductase inhibitors, (ii) bile acid sequestrants, (iii) nicotinic alcohol, nicotinic acid or a salt thereof, (iv) PPAR α agonists, (v) cholesterol absorption inhibitors, (vi) acyl CoA:cholesterol acyltransferase (ACAT) inhibitors, (vii) CETP inhibitors, and (viii) phenolic anti-oxidants;

- (i) PPAR α/γ dual agonists,
- (j) PPAR δ agonists,
- (k) antiobesity compounds,
- (l) ileal bile acid transporter inhibitors;
- (m) anti-inflammatory agents;
- (n) glucagon receptor antagonists;
- (o) GLP-1;
- (p) GIP-1;
- (q) GLP-1 analogs;
- (r) HSD-1 inhibitors;
- (s) SGLT-2 inhibitors; and
- (t) SGLT-1/SGLT-2 inhibitors; and

(3) a pharmaceutically acceptable carrier.

24. A pharmaceutical composition comprising a compound according to Claim 1, or a pharmaceutically acceptable salt thereof, and a compound selected from simvastatin, ezetimibe and sitagliptin; and a pharmaceutically acceptable carrier.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CN2014/084034

A. CLASSIFICATION OF SUBJECT MATTER

C07D 313/08(2006.01)i; C07D 405/04(2006.01)i; A61K 31/335(2006.01)i; A61K 31/44(2006.01)i; A61P 3/10(2006.01)i; A61P 3/08(2006.01)i

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D; A61K; A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CNPAT, CNKI, EPODOC, WPI, REGISTRY(STN), CAPLUS(STN): benzoxepin, phenyl, pyridyl, antagonist, diabete?, G protein, couple? receptor, GPR40

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0388188 A1 (SUNTORY LTD) 19 September 1990 (1990-09-19) description, examples 1-4, reference examples 12, 17, claims 1, 5-6 and 9	1-7, 10-14, 18-24
X	EP 0250265 B1 (SUNTORY LTD) 05 September 1990 (1990-09-05) description, examples 1-33, claims 1, 4-5 and 13	1-7, 10-14, 18-24
A	US 6268354 B1 (TAKEDA CHEMICAL INDUSTRIES LTD) 31 July 2001 (2001-07-31) the whole document	1-24
X	SUCHAND B. et al. "An efficient intermolecular [Pd]-catalyzed C-C and intramolecular [Cu]-catalyzed C-O bonds formation: synthesis of functionalized flavans and benzoxepine" <i>Tetrahedron Letters</i> , Vol. 30, No. 53, 16 May 2012 (2012-05-16), ISSN: pages 3861-3864, page 3864, left column	1-7, 10-12, 18

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

08 April 2015

Date of mailing of the international search report

30 April 2015

Name and mailing address of the ISA/CN

STATE INTELLECTUAL PROPERTY OFFICE OF THE
P.R.CHINA(ISA/CN)
6,Xitucheng Rd., Jimen Bridge, Haidian District, Beijing
100088, China

Authorized officer

KANG,Lei

Facsimile No. (86-10)62019451

Telephone No. (86-10)61648356

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CN2014/084034

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **21-22**
because they relate to subject matter not required to be searched by this Authority, namely:
 - [1] The subject matter of claims 21-22 relates to methods for the treatment of human body by therapy as defined in PCT Rule 39.1(IV). This search has been carried out on the basis of the subject matter of the use in manufacture of medicaments for treating the alleged diseases.
2. ☒ Claims Nos.: **1-14**
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
 - [1] In claims 1-14, the definition of groups Q, T, U, V, W, X, Y, Z encompasses a very large number of possible embodiments, while the description discloses, and provides support for, only a relatively small proportion of those embodiments. Thus, claims 1-14 are not fully supported by description in the sense of PCT Article 6. The search has been carried out on the base of the definition of groups Q, T, U, V, W, X, Y, Z as occurred in claim 15.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.

PCT/CN2014/084034

Patent document cited in search report			Publication date (day/month/year)	Patent family member(s)			Publication date (day/month/year)
EP	0388188	A1	19 September 1990	AT	100097	T	15 January 1994
				US	5041444	A	20 August 1991
				DE	69005861	T2	28 April 1994
				JP	H02243687	A	27 September 1990
				ES	2062336	T3	16 December 1994
				EP	0388188	B1	12 January 1994
				DK	0388188	T3	14 February 1994
				DE	69005861	D1	24 February 1994
				JP	2851054	B2	27 January 1999
EP	0250265	B1	05 September 1990	HU	206108	B	28 August 1992
				DE	3764724	D1	11 October 1990
				US	4845094	A	04 July 1989
				HU	T48231	A	29 May 1989
				AU	597671	B2	07 June 1990
				JP	H0813807	B2	14 February 1996
				JP	S63107972	A	12 May 1988
				DK	314487	D0	19 June 1987
				AU	7424587	A	24 December 1987
				CA	1277320	C	04 December 1990
				DK	166676	B1	28 June 1993
				DK	314487	A	21 December 1987
				EP	0250265	A1	23 December 1987
US	6268354	B1	31 July 2001	US	6172061	B1	09 January 2001