



(51) International Patent Classification:

Not classified

(21) International Application Number:

PCT/US2017/048934

(22) International Filing Date:

28 August 2017 (28.08.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/401,687 29 September 2016 (29.09.2016) US
62/505,458 12 May 2017 (12.05.2017) US

(71) Applicants: **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; 1111 Franklin Street, 12th Floor, Oakland, CA 94607 (US). **FRED HUTCHINSON CANCER RESEARCH CENTER** [US/US]; 1100 Fairview Ave N, Seattle, WA 98109 (US).

(72) Inventors: **ABERGEL, Rebecca J.**; c/o The Regents of the University of California, 1111 Franklin Street, 12th Floor, Oakland, CA 94607 (US). **STRONG, Roland K.**; c/o Fred Hutchinson Cancer Research Center, 1100 Fairview Ave N, Seattle, WA 98109 (US). **DEBLONDE, Gauthier J.**; c/o

The Regents of the University of California, 1111 Franklin Street, 12th Floor, Oakland, CA 94607 (US).


(74) Agent: **MALLON, Joseph, J.**; Knobbe Martens Olson & Bear LLP, 2040 Main Street, Fourteenth Floor, Irvine, CA 92614 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: SEPARATION OF METAL IONS BY LIQUID-LIQUID EXTRACTION

FIG. 35



(57) Abstract: Provided herein are separation processes for metal ions present in aqueous solutions based on methods involving liquid-liquid extraction. The separation process involves a chelator that can selectively bind to at least one of the metals at a relatively low pH. This can be used, for example, for recovery and purification of actinides from lanthanides, separation of metal ions based on their valence, and separation of metal ions based on the pH of the extraction conditions.

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- *without international search report and to be republished upon receipt of that report (Rule 48.2(g))*
- *with information concerning incorporation by reference of missing parts and/or elements (Rule 20.6)*

SEPARATION OF METAL IONS BY LIQUID-LIQUID EXTRACTION

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application Nos. 62/505,458, filed on May 12, 2017 and 62/401,687, filed on Sept. 29, 2016, which is hereby incorporated by reference in its entirety.

STATEMENT REGARDING FEDERALLY SPONSORED R&D

[0002] The invention was made with government support under Contract No. DE-AC02-05CH 11231 between the U.S. Department of Energy and the University of California. The government has certain rights in the invention.

BACKGROUND

Field

[0003] The present disclosure is related to separation processes for metal ions.

Description of the Related Art

[0004] Owing to concerns related to global warming due to CO₂ emission by traditional energy sources, nuclear energy has emerged as an alternative energy source. However, disposing of nuclear waste is a major challenge.

[0005] A few processes, using liquid-liquid extraction, are under development with limited success in term of selectivity and efficacy: TALSPEAK process (USA), GANEX process (France), and DIAMEX-SANEX process (France). The purification of the element actinium (Ac³⁺) is also catching growing attention. In fact, actinium isotopes are emerging radioactive sources for cancer treatments such as the “targeted alpha therapies.” The methods of production of Ac³⁺ isotopes generate Ln³⁺ fission products in addition to the initial material that is usually made of thorium (Th⁴⁺). Current purification protocols for actinium involve numerous chromatographic steps that require different chemical media in order to obtain a purified fraction of Ac³⁺.

SUMMARY

[0006] In some embodiments, a method of processing spent nuclear fuel is provided. This comprises obtaining spent nuclear fuel dissolved in an acidic medium, contacting the dissolved spent nuclear fuel with an octadentate ligand and an organic phase to generate a mixture, and separating an aqueous phase enriched for Pu^{4+} from the mixture.

[0007] In some embodiments, a method for enriching a metal ion is provided. The method comprises contacting a first aqueous phase with an organic phase to generate a mixture, wherein the first aqueous phase comprises: a plurality of metal ions; and an octadentate ligand, and wherein the aqueous phase has an acidic pH of less than 1. The method further includes separating from the mixture a second aqueous phase enriched for a metal ion of the plurality of metal ions.

[0008] In some embodiments, a method of preparing a medical isotope is provided. The method comprises obtaining a metallic precursor dissolved under an acidic condition, wherein the metallic precursor comprises a medical isotope; contacting the dissolved metallic precursor with an octadentate ligand and an organic phase to generate a mixture; and separating the medical isotope from one or more metal ions in the dissolved metallic precursor based on an interaction between the octadentate ligand and the medical isotope.

[0009] In some embodiments, a method of separating metal ions for nuclear forensics is provided. The method comprises obtaining a sample derived from a nuclear material, wherein the sample comprises UO_2^{2+} , Ac^{3+} , Pu^{4+} , and Np^{4+} ; contacting the sample with an octadentate ligand to generate a first mixture, wherein the first mixture has an acidic pH of 1 or lower; separating UO_2^{2+} and Ac^{3+} from Pu^{4+} and Np^{4+} in the mixture based on an interaction or lack of interaction between the octadentate ligand and each of UO_2^{2+} , Ac^{3+} , Pu^{4+} , and Np^{4+} , to generate a second mixture comprising the Pu^{4+} and Np^{4+} ; and further separating each metal within both mixtures by classical techniques.

[0010] In some embodiments, a method of separating metal ions is provided. The method comprises contacting a liquid composition comprising a plurality of metal ions with an octadentate ligand to generate a mixture, under conditions sufficient to form a metal ion-ligand complex comprising a metal ion of the plurality of metal ions, and separating a first fraction of the mixture enriched for the metal ion-ligand complex from a second fraction

depleted for the metal ion-ligand complex, wherein the first fraction of the mixture has an acidic pH of less than 1. The plurality of metal ions are selected from the group consisting of: a p-, d- or f-block element of period 5 or greater, a group 3 element, or a group 4 element

[0011] In some embodiments, a method of separating trivalent metal ions from tetravalent metal ions is provided. The method comprises providing an aqueous solution comprising tetravalent metal ions and trivalent metal ions, wherein the solution has an acidic pH; adding a HOPO chelator to the aqueous solution; and performing an extraction against the aqueous solution, wherein the HOPO chelator functions as a holdback agent to keep the tetravalent metal ions in the aqueous solution, while allowing the trivalent metal ions to be removed via decanting at the end of the extraction.

[0012] In some embodiments, a method for separating metal ions is provided. The method comprises providing an aqueous solution comprising a first metal ion and a second metal ion, wherein the solution has an acidic pH; adding a HOPO chelator to the aqueous solution; and performing an extraction against the aqueous solution. The HOPO chelator functions as a holdback agent to selectively keep the first metal ion in the aqueous solution, while allowing the second metal ion to go into an organic phase during the extraction.

BRIEF DESCRIPTION OF THE DRAWINGS

[0013] FIG. 1 shows current state-of-the-art purification of 227/225Ac.

[0014] FIG. 2 shows extraction can be based on one or more principles.

[0015] FIG. 3 shows an embodiment of the principle of liquid-liquid extraction process.

[0016] FIG. 4 shows spectroscopic characterization of Sn^{4+} complexation in 2 M HCl.

[0017] FIG. 5 shows an embodiment of the PUREX process, which can be modified by one or more of the HOPO chelators as described herein.

[0018] FIG. 6 shows structures of compounds used in the TALSPEAK process, which can be modified by one or more of the HOPO chelators inspired by the siderophore Enterobactin, as described herein.

[0019] FIG. 7 shows an embodiment of the TALSPEAK process, which can be modified by one or more of the HOPO chelators as described herein.

[0020] FIG. 8 shows a Eu(III) speciation diagram in the presence of excess DTPA ($I=0.1$ M at 25°C).

[0021] FIG. 9 shows structures of the aqueous octadentate ligand chelator 3,4,3-LI(1,2-HOPO) (left) and of the binding moiety 1,2-HOPO (right). Particular protons are labile and lead to the formation of $[3,4,3\text{-LI}(1,2\text{-HOPO})]^{4-}$ upon deprotonation for binding to a metal ion. HOPO = hydroxypyridinone.

[0022] FIG. 10 shows binding of DTPA (left) to M^{n+} in acidic solutions.

[0023] FIG. 11 shows comparison of ratio Ac/Am in organic phase for 3,4,3-LI(1,2-HOPO) and DTPA.

[0024] FIG. 12 shows comparison of ratio Ac/Gd in organic phase for 3,4,3-LI(1,2-HOPO) and DTPA.

[0025] FIG. 13 shows comparison of ratio Gd/Am in organic phase for 3,4,3-LI(1,2-HOPO) and DTPA.

[0026] FIG. 14 shows a graph of extraction coefficient for DTPA for Gd, Ac and Am.

[0027] FIG. 15 shows a graph of extraction coefficient for 3,4,3-LI(1,2-HOPO) for Gd, Ac, and Am.

[0028] FIG. 16 shows percentage of extraction as a function of pH for trivalent Ac, Am and, Gd ions in the presence of DTPA as hold-back reagent.

[0029] FIG. 17 shows calculated separation factors (Ac/Gd and Ac/Am) for a 1-step extraction using 3,4,3-LI(1,2-HOPO) as a hold-back reagent.

[0030] FIG. 18 shows percentage of extraction of plutonium(IV) as a function of pH. For clarity, a logarithm scale is used on the Y-axis.

[0031] FIG. 19 shows extraction profile for Gd(III) and Am(III) along with the separation factor plotted at various pH with 3,4,3-LI(1,2-HOPO) as hold-back reagent.

[0032] FIG. 20 shows the stability constant values published for the complexes of 3,4,3-LI(1,2-HOPO).

[0033] FIG. 21 shows a speciation diagram for solutions containing 3,4,3-LI(1,2-HOPO) and one equivalent of Eu^{3+} or Th^{4+} ($I=0.1$ M at 25°C), showing use of 3,4,3-LI(1,2-HOPO) to suppress the extraction of M^{4+} and selective extraction of M^{3+} .

[0034] FIG. 22 shows relationship between ionic radius and stability constants for 3,4,3-LI(1,2-HOPO) for other actinide and lanthanide ions.

[0035] FIG. 23 shows percentage of extraction as a function of pH for trivalent actinium, americium and, gadolinium ions (Ac(III)-225, Am(III)-243, and Gd(III)-153, respectively) in the presence of 3,4,3-LI(1,2-HOPO) as hold-back reagent.

[0036] FIG. 24 shows an embodiment of a general process flowsheet proposed for the purification of actinium isotopes using 3,4,3-LI(1,2-HOPO) and HDEHP.

[0037] FIG. 25 shows an embodiment of another process proposed for the recovery of actinium using 3,4,3-LI(1,2-HOPO) and a non-selective extractant.

[0038] FIG. 26 shows percentage of extraction of plutonium(IV), americium(III), and gadolinium(III) by HDEHP at constant pH and using 3,4,3-LI(1,2-HOPO) as hold-back reagent.

[0039] FIG. 27 shows corresponding separation factors for FIG. 26.

[0040] FIG. 28 shows percentage of extraction of americium(III), and gadolinium(III) by HDEHP at constant pH and using 3,4,3-LI(1,2-HOPO) as hold-back reagent.

[0041] FIG. 29 shows the effects of acid and pH on Zr^{4+} -3,4,3-LI(1,2-HOPO) interactions.

[0042] FIG. 30 shows the effects of acid and pH on Ti^{4+} -3,4,3-LI(1,2-HOPO) interactions.

[0043] FIG. 31 shows the effects of acid and pH on Sn^{4+} -3,4,3-LI(1,2-HOPO) interactions.

[0044] FIG. 32 shows speciation of M^{3+} or M^{2+} versus M^{4+} ions in the presence of 3,4,3-LI(1,2-HOPO).

[0045] FIG. 33 shows an embodiment of a strategy for grouped separation of M^{4+} ions versus others ions at low pH.

[0046] FIG. 34 shows a flowsheet for acidic feed solutions for selective extraction step using 3,4,3-LI(1,2-HOPO) as hold-back reagent for thorium.

[0047] FIG. 35 shows a flowsheet for feed solutions at pH near 3 and using HDEHP as extractant for selective extraction step using 3,4,3-LI(1,2-HOPO) as hold-back reagent for thorium.

[0048] FIG. 36 shows an embodiment of a general process strategy for the recovery and purification of Ac-225 from a Th spallation target.

[0049] FIG. 37 shows an embodiment of the TALSPEAK process with a HOPO chelator.

FIG. 38A depicts an embodiment of a general peptoid synthesis.

FIG. 38B depicts an embodiment of a general peptoid synthesis.

FIG. 38C depicts some embodiments of a general peptoid synthesis.

FIG. 38D depicts some embodiment of a general peptoid synthesis.

FIG. 38E depicts some embodiments of a combination of products obtained from a general peptoid synthesis scheme.

FIG. 38F depicts some embodiments of a fluorescent tag.

FIG. 39A depicts some embodiments of a conjugation pathway.

FIG. 39B depicts some embodiments of a conjugation pathway.

FIG. 40A depicts reaction schemes for conjugation.

FIG. 40B depicts reaction schemes for a conjugation.

FIG. 40C depicts some embodiments of a fluorescent tag incorporation.

FIG. 41 depicts a representative reversed-phase HPLC trace

FIG. 42A depicts the mass spectra (HHHC Peptoid. MS1: free peptoid, MS2: peptoid-Na, MS3: peptoid-Fe and peptoid-Fe-K.

FIG. 42B depicts the mass spectra for CHHH Peptoid. MS1: free peptoid and peptoid-K, MS2: peptoid-Na, peptoid-Fe, and possible impurity.

FIG. 42C depicts the mass spectra for HCHH Peptoid. MS1: free peptoid and

peptoid-K, MS2: peptoid-Na, peptoid-Fe.

FIG. 42D depicts the mass spectra for HHCH Peptoid. MS1: free peptoid and possible impurity, MS2: peptoid-Na, MS3: Peptoid-Fe.

FIG. 42E depicts the mass spectra for CHHC Peptoid. MS1: free peptoid only and peptoid-K, MS2: peptoid-Na and peptoid-Fe.

FIG. 42F depicts the mass spectra for HHCC Peptoid. MS1: free peptoid and peptoid-Na, MS2: peptoid-Na, free peptoid, and peptoid-Fe.

FIG. 42G depicts the mass spectra for CCHH Peptoid. MS1: free peptoid and peptoid-Na, MS2: Peptoid-Fe and possible trimer.

FIG. 42H depicts the mass spectra for HCHC Peptoid. MS1: free peptoid, MS2: peptoid-Fe and peptoid-Na.

FIG. 42I depicts the mass spectra for HCCH Peptoid. MS1: free peptoid and small amount trimer, MS2: peptoid-Fe, MS3: peptoid-Na and possible fragments.

FIG. 42J depicts the mass spectra for CHCH Peptoid. MS1: free peptoid and peptoid-K, MS2: peptoid-Fe.

FIG. 42K depicts the mass spectra for HCCC Peptoid. MS1: free peptoid and trimer-Fe trace, MS2: free peptoid and peptoid-Fe.

FIG. 42L depicts the mass spectra for CHCC Peptoid. MS1: free peptoid and peptoid-Na, MS2: free peptoid and unidentified masses.

FIG. 42M depicts the mass spectra for CCHC Peptoid. MS1: free peptoid, MS2: free peptoid and peptoid-Na.

FIG. 42N depicts the mass spectra for CCCH Peptoid. MS1: free peptoid and peptoid-Na, MS2: free peptoid and peptoid-K/Na.

FIG. 42O depicts the mass spectra for CCCH Peptoid. MS1: free peptoid and peptoid-Na, MS2: free peptoid and peptoid-K/Na.

FIG. 42P is the mass spectra for HHHH Peptoid. MS1: free peptoid, MS2:

peptoid-Fe/Na, MS3: Peptoid-Fe-Na.

FIGs. 43A-43C depict the spectra that were obtained in negative mode. All peptoids are 1:1 Fe^{3+} complexes except for CCCC and HHHH. Top spectrum is calculated and bottom is obtained.

FIG. 44 depicts the TOF MSMS of select peptoids.

FIGs. 45A-45D depict NMR results of select peptoids.

DETAILED DESCRIPTION

[0050] Provided herein are separation processes for metal ions in aqueous solutions. In some embodiments, this can be based on liquid-liquid extraction. In some embodiments, this allows for recovery and/or purification of actinides from lanthanides, based on their valence, and the pH of the extraction conditions.

[0051] In some embodiments, the extraction includes dissolving the raw material in an acidic or basic matrix and providing a chelator or ligand (such as a HOPO chelator) that selectively extracts the metal(s) ion(s) of interest. This can be done at a relatively low pH in some embodiments. For example, in some embodiments, the first step of the extraction is to dissolve the raw material in an acidic or basic matrix and the second step is to find a soluble species that selectively extracts (or back-extracts) the metal(s) ion(s) of interest (FIG. 2), such as a HOPO chelator or ligand. The metal associated with this chelator can then be extracted (or retained) via the properties of the ligand, while the metal not associated with the chelator can be removed (or retained). Furthermore, as disclosed herein, chelators such as HOPO molecules, allow binding of various metal ions, as a function of pH, even at low pH. This allows for pH manipulation to drive selective binding of particular metal ion species over different metal ion species (e.g., M^{3+} vs M^{4+}). Thus, by using such a chelator, and adjusting the pH accordingly, one can selectively bind the metal ion desired, remove any remaining metal ion (via, e.g., a general “extracting agent” that will move to the organic phase), and then, once the organic phase is decanted, collect the metal desired metal ion and removing it from the chelator. The separation techniques can include liquid-liquid extraction (large scale), and/or chromatography or ion-exchange. In some embodiments, the various

metal separation techniques can employ chromatography and/or a liquid-liquid extraction. In some embodiments, further steps that can occur include, for example liquid-liquid extraction, chromatography, and/or analytical techniques.

Definitions

[0052] The terms "pure," "purified," "substantially purified," and "isolated" as used herein refer to the compound of the embodiment being free of other, dissimilar compounds with which the compound, if found in its natural state (e.g., natural ores), as raw material(s), in waste for recycling, or man-made materials, would be associated in its natural state or as raw material(s). In certain embodiments described as "pure," "purified," "substantially purified," or "isolated" herein, the compound can comprise at least 0.5% to 1%, 1% to 5%, 5% to 10%, 10% to 20%, 20% to 50%, 50% to 70%, 70% to 90%, 90% to 95%, 95% to 99%, and 99% to 100%. In some embodiments, the amount of the compound will be at least 50% or 75% of the mass, by weight, of a given sample. A "functional purity" is a measurement of the amount of a particular compound in a sample or product in relation to other compounds in a sample that can adversely impact the function of the compound. Thus, other components in a sample that do not interfere with the compound's activity (e.g., water), will not be used in determining the purity of a sample or product.

[0053] The terms "derivative," "variant," or other similar term refers to a compound that is an analog of the other compound.

[0054] The term "and/or" designates both the option of "and" as well as the option of "or" in that particular circumstance. However, unless otherwise specified in the specification, the use of the term "or" or "and" encompasses a description of both option as well. Thus, the use of the term "or" should not be taken as excluding the option of "and", unless additional context indicates that it should (this definition does not apply to the language in the claims). The use of the singular or plural forms of a term encompasses both options (singular or plural) as well as both options combined (singular and plural), unless indicated otherwise.

Methods

[0055] In some embodiments, a method of processing spent nuclear fuel is provided. This comprises obtaining spent nuclear fuel dissolved in an acidic medium,

contacting the dissolved spent nuclear fuel with an octadentate ligand and an organic phase to generate a mixture, and separating an aqueous phase enriched for Pu^{4+} from the mixture.

[0056] In some embodiments, a method for enriching a metal ion is provided. The method comprises contacting a first aqueous phase with an organic phase to generate a mixture, wherein the first aqueous phase comprises: a plurality of metal ions; and an octadentate ligand, and wherein the aqueous phase has an acidic pH of less than 1. The method further includes separating from the mixture a second aqueous phase enriched for a metal ion of the plurality of metal ions.

[0057] In some embodiments, a method of preparing a medical isotope is provided. The method comprises obtaining a metallic precursor dissolved under an acidic condition, wherein the metallic precursor comprises a medical isotope; contacting the dissolved metallic precursor with an octadentate ligand and an organic phase to generate a mixture; and separating the medical isotope from one or more metal ions in the dissolved metallic precursor based on an interaction between the octadentate ligand and the medical isotope.

[0058] In some embodiments, a method of separating metal ions for nuclear forensics is provided. The method comprises obtaining a sample derived from a nuclear material, wherein the sample comprises UO_2^{2+} , Ac^{3+} , Pu^{4+} , and Np^{4+} ; contacting the sample with an octadentate ligand to generate a first mixture, wherein the first mixture has an acidic pH of 1 or lower; separating UO_2^{2+} and Ac^{3+} from Pu^{4+} and Np^{4+} in the mixture based on an interaction or lack of interaction between the octadentate ligand and each of UO_2^{2+} , Ac^{3+} , Pu^{4+} , and Np^{4+} , to generate a second mixture comprising the Pu^{4+} and Np^{4+} ; and further separating each metal within both mixtures by classical techniques.

[0059] In some embodiments, a method of separating metal ions is provided. The method comprises contacting a liquid composition comprising a plurality of metal ions with an octadentate ligand to generate a mixture, under conditions sufficient to form a metal ion-ligand complex comprising a metal ion of the plurality of metal ions, and separating a first fraction of the mixture enriched for the metal ion-ligand complex from a second fraction depleted for the metal ion-ligand complex, wherein the first fraction of the mixture has an acidic pH of less than 1. The plurality of metal ions are selected from the group consisting of: a p-, d- or f-block element of period 5 or greater, a group 3 element, or a group 4 element.

[0060] In some embodiments, a method of separating trivalent metal ions from tetravalent metal ions is provided. The method comprises providing an aqueous solution comprising tetravalent metal ions and trivalent metal ions, wherein the solution has an acidic pH; adding a HOPO chelator to the aqueous solution; and performing an extraction against the aqueous solution, wherein the HOPO chelator functions as a holdback agent to keep the tetravalent metal ions in the aqueous solution, while allowing the trivalent metal ions to be removed via decanting at the end of the extraction.

[0061] In some embodiments, a method for separating metal ions is provided. The method comprises providing an aqueous solution comprising a first metal ion and a second metal ion, wherein the solution has an acidic pH; adding a HOPO chelator to the aqueous solution; and performing an extraction against the aqueous solution. The HOPO chelator functions as a holdback agent to selectively keep the first metal ion in the aqueous solution, while allowing the second metal ion to go into an organic phase during the extraction.

[0062] In some embodiments, the acidic denotes a pH of between about 1 to about 0.

[0063] In some embodiments, the octadentate ligand is a hydroxypyridonate ligand. In some embodiments, the octadentate ligand is built on a spermine scaffold. In some embodiments, the octadentate ligand is 3,4,3-LI(1,2-HOPO).

[0064] In some embodiments, the method further comprises dissolving a metal in an acidic solution to obtain the liquid composition.

[0065] In some embodiments, the first fraction is enriched for a tetravalent metal ion and the second fraction is enriched for a trivalent metal ion and/or a divalent metal ion.

[0066] In some embodiments, the contacting further comprises contacting the sample with an organic phase.

[0067] In some embodiments, a method of separating trivalent metal ions from tetravalent metal ions is provided. The method comprises providing an aqueous solution comprising tetravalent metal ions and trivalent metal ions, wherein the solution has an acidic pH; adding a HOPO chelator to the aqueous solution; and performing an extraction against the aqueous solution, wherein the HOPO chelator functions as a holdback agent to keep the

tetravalent metal ions in the aqueous solution, while allowing the trivalent metal ions to be removed via decanting at the end of the extraction.

[0068] In some embodiments, the method further comprises separating the HOPO chelator from the tetravalent metal to thereby collect the tetravalent metal.

[0069] In some embodiments, the method further comprises lowering a pH beneath 0 to separate tetravalent ions.

[0070] In some embodiments, separating comprises filtering, precipitating, liquid-liquid extraction or chromatography, or wherein the method further comprises filtering, precipitating, liquid-liquid extraction or chromatography.

[0071] In some embodiments, the first fraction is enriched for a first metal ion that has a charge that is different from a charge of a second metal ion enriched in the second fraction.

[0072] In some embodiments, 1) the separating comprises contacting the liquid composition with an organic phase or 2) further comprising contacting the liquid composition with an organic phase.

[0073] In some embodiments, the method does not comprise adjusting the pH of the liquid composition after dissolving the metal. In some embodiments, the method does not comprise raising the pH of the liquid composition after dissolving the metal.

[0074] In some embodiments, the organic phase comprises a non-selective extractant. In some embodiments, the organic phase does not comprise an extractant that selectively binds to the first or second metal ion. In some embodiments, the method does not comprise adjusting the pH of the dissolved spent nuclear fuel. In some embodiments, the extractant is selected from the group consisting of di-(2-ethylhexyl)phosphoric acid (HDEHP), a derivative of HDEHP, calixarenes, diglycoamide derivatives, carbamoylphosphine oxide derivatives, tributylphosphate (TBP), monoamide derivatives, and tertiary amines, and quaternary ammonium salts.

[0075] In some embodiments, the first metal ion is a tetravalent metal ion and the second metal ion is selected from the group consisting of a divalent metal ion and a trivalent metal ion. In some embodiments, the first metal ion is a trivalent metal ion and the second metal ion is a trivalent metal ion.

[0076] In some embodiments, a HOPO chelator binds to the first metal ion more effectively than it binds to the second metal ion at a given pH.

[0077] In some embodiments, the fuel, first aqueous phase, precursor, nuclear material, liquid composition, or solution comprises actinium. In some embodiments, the actinium is separated from a M^{4+} metal. Actinium can be included in and/or isolated as a result of, any of the methods provided herein.

[0078] In some embodiments, the chelators provided herein are distinct from classical aqueous chelators such as citrate, EDTA (Ethylenediaminetetraacetic acid), DTPA (Diethylenetriaminepentaacetic acid) as classical chelators were not able to complex metal ions under highly acidic conditions. Due to that, traditional separation of metal ions relied on the selectivity of the extractant (organic molecule, such as TBP, dissolved in an organic phase) or on the selectivity of a solid support (in the case of chromatography). This selectivity is not required in some of the embodiments provided herein.

[0079] In some embodiments, the use of 3,4,3-LI(1,2-HOPO) allows one to be able to selectively bind to the tetravalent metal ions under highly acidic conditions. Results show that 3,4,3-LI(1,2-HOPO) stays bound to Hf^{4+} and Zr^{4+} from pH 8 to strongly acidic solutions (at least 6 M HCl). The same compound stays from pH 6 to 10 M HCl (at least) in the case of Ti^{4+} and from pH 8 to 1 M HCl in the case of Sn^{4+} . Based on thermodynamic data (stability constant of the complex) one can expect Pu^{4+} , Th^{4+} and Ce^{4+} to behave like Zr^{4+} and Hf^{4+} .

[0080] In some embodiments, the chelating agents are capable of binding or chelating, or capable of forming stable complexes with actinides and/or lanthanides, such as the cations of Eu, Pu, Np, Th, Am, and/or Cf, such as of $^{152}Eu(III)$, $^{241}Am(III)$, $^{238}Pu(IV)$, $^{237}Np(IV)$, $^{237}Np(V)$, and $^{233}U(VI)$.

[0081] As noted above, selective pH dependent binding, at an acidic pH range, can be achieved using HOPO molecules. Furthermore, thermodynamic studies have shown that 3,4,3-LI(1,2-HOPO) releases the trivalent metal ions (M^{3+}) and divalent uranyl (UO_2^{2+}) as soon as the pH value is lower than about 1. Thus, in some embodiments M^{4+} metals can be separated from M^{3+} and/or M^{2+} via a HOPO ligand when subjected to lower pH values. In some embodiments, the ligand 3,4,3-LI(1,2-HOPO) could also be immobilized on a solid support (e.g., beads, nanoparticles, or other form of solid support).

[0082] In some embodiments, provided herein is a process involving a high acidity (directly after the metallic target dissolution) where 3,4,3-LI(1,2-HOPO) is used to selectively bind the tetravalent metal ions in the aqueous and acidic liquor. The separation of tetravalent ions (bound to the chelator) for the other ions (unbound) can then be achieved by using a non-selective extractant for the remaining metal ions and performing a liquid-liquid extraction step (for example). The tetravalent ions stay in the aqueous phase whereas the rest are extracted in the organic phase. The selectivity of such a process can be extremely high since 3,4,3-LI(1,2-HOPO) specifically binds the tetravalent metal ions under acidic condition (but not the trivalent metal ions). In some embodiments, the concentration of HCl can range from about 0 M to about 4 M. In some embodiments, the concentration of HCl can range from about 4 M to about 8 M. In some embodiments, the concentration of HCl can range from about 8 M to about 12 M. In some embodiments, the concentration of HCl is about 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 M, or a value within a range defined by any two of the aforementioned values. The selectivity of the process can be due to the presence of 3,4,3-LI(1,2-HOPO) in the aqueous phase (*See*, TABLE 0.2). A non-selective extractant can be used in the organic phase.

[0083] In some embodiments, selective complexation of tetravalent ions is achieved by 3,4,3-LI(1,2-HOPO) under very acidic conditions, e.g., less than pH 3, 2.9, 2.8, 2.7, 2.6, 2.5, 2.4, 2.3, 2.2, 2.1, 2.0, 1.9, 1.8, 1.7, 1.6, 1.5, 1.4, 1.3, 1.2, 1.1, 1.0, 0.9, 0.8, 0.7, 0.6, 0.5, 0.4, 0.3, 0.2, 0.1, 0, or lower.

[0084] In some embodiments, the strategy for metal ion separation provided herein aims at using innovative chelators (e.g., HOPO containing molecules), for metal ion separation processes. In some embodiments, processes that are more selective, more efficacious and more flexible than the current state-of-the-art are provided.

[0085] In some embodiments, one or more of the processes disclosed herein can be used to separate M^{2+} from M^{3+} from M^{4+} , where each M is a metal. In some embodiments, M^{2+} is first separated from a M^{3+}/M^{4+} mixture, and then the M^{3+} is separated from M^{4+} via a different technique, such as chromatography. In some embodiments, one or more of the processes disclosed herein can be employed to separate a first M^{4+} from a second M^{4+} or a first M^{3+} from a second M^{3+} . In some embodiments, the separation is of Actinium from other metals. The separation is achieved by different affinities of each metal ion for the

chelator. This may, or may not, also be driven in a pH dependent manner (that is, affinity changes as a function of pH for the various metal ions as well).

[0086] In some embodiments, the separation can be of lanthanides from actinides. In some embodiments, separation is of element actinium from both lanthanides and actinides.

[0087] The separation of trivalent lanthanide ions (examples: Gd^{3+} , Eu^{3+} , etc. herein referred to as Ln^{3+}) from the trivalent actinide ions (Examples: Ac^{3+} , Am^{3+} , Cm^{3+} , Bk^{3+} , Cf^{3+} , etc. Here after referred to as An^{3+}) can be challenging because these two classes of metal ions exhibit similar chemical properties. Nonetheless, there is value in efficient processes able to separate Ln^{3+} from An^{3+} . Indeed, Ln^{3+} and An^{3+} are both present in the nuclear spent fuels (civilian and military) and their separation could allow for two developments. First, the presence of An^{3+} impairs the long-term storage of the nuclear waste because it would generate too much heat in hypothetical underground repositories. The removal of An^{3+} from the main nuclear waste would therefore facilitate the storage and reduce the cost. Second, the minor actinides An^{3+} could be transmuted into less toxic elements in the next generation of nuclear reactors but this transmutation is only possible in the absence of lanthanide fission products Ln^{3+} . The separation Ln^{3+}/An^{3+} is therefore beneficial to obtain closed nuclear fuel cycles (with recycling of U and Pu, transmutation of An^{3+} , and storage of Ln^{3+}).

[0088] In some embodiments, separation of metal ions is beneficial during mining for natural resources. In some embodiments, separation of metal ions is involved during mining for recycling of valuable waste. In some embodiments, separation of metal ions is required during mining for purification of medical and/or research isotopes, e.g., ^{177}Lu , ^{223}Ra , ^{225}Ac , and ^{227}Th . In some embodiments, the metal ion separation can be for mining from natural stones, recycling of valuable wastes, purification of medical and/or research isotopes such as ^{177}Lu , ^{223}Ra , ^{225}Ac , and ^{227}Th . An example of purification of $^{227/225}Ac$ is shown in FIG. 1 (Radchenko et al 2015 Journal of Chromatography A, 55-63, which is hereby incorporated by reference in its entirety).

[0089] In some embodiments, the method for separation of metal ions include, without limitations, liquid-liquid extraction and chromatography or ion-exchange-based methods. In some embodiments, the methods are large scale. In some embodiments, the methods are small scale. In some embodiments, liquid-liquid extraction methods are large

scale. In some embodiments, chromatography or ion-exchange-based methods are small scale. In some embodiments, countercurrent liquid-liquid extraction can be employed.

[0090] Liquid-liquid extraction (FIG. 3) processes involve two immiscible phases (one aqueous, one organic) between which the elements of interest are selectively transferred. In some embodiments, one can use one or more chelators in the aqueous phase (also called hold-back reagent) and use, preferentially, a non-selective extractant to pull some or all of the other metals into the organic phase.

[0091] In some embodiments, the present disclosure provides liquid-liquid extraction with ligands based on the binding units 1,2-HOPO that can selectively bind or release targeted elements (Ac^{3+} , Am^{3+} , Gd^{3+} , etc.) in aqueous media. As disclosed herein, the affinity of the 1,2-HOPO ligands for a given metal ion also depends on the acidity of the aqueous media. Thus, modifying the acidity allows for selective binding or release of HOPO chelators to the metal of choice. Furthermore, the selectivity of the HOPO ligands is usually superior to that of the chelators currently used for this type of application (e.g., DTPA). Thus, in some embodiments, depending on the chemical conditions, a given element can be selectively extracted in the organic from a mixture of elements by using specific chemical conditions in the aqueous phase containing the HOPO ligand and a non-specific extractant in the organic phase.

[0092] In some embodiments, a new strategy for the complexation and purification of tetravalent metal ions from other ions (not exhibiting a $4+$ charge) is provided. Examples of elements that naturally form tetravalent ions are: plutonium (Pu^{4+}), thorium (Th^{4+}), titanium (Ti^{4+}), zirconium (Zr^{4+}), hafnium (Hf^{4+}), cerium (Ce^{4+}), tin (Sn^{4+}), platinum (Pt^{4+}), iridium (Ir^{4+}). Examples of elements that can be stabilized as tetravalent ions under specific conditions are: berkelium (Bk^{4+}), uranium (U^{4+}), and lead (Pb^{4+}). Thus, any of the elements can be used in the present methods. In some embodiments, actinium can be separated from any of these metals.

[0093] In some embodiments, the separation techniques can employ altering and/or controlling the pH of a solution so as to allow M^{3+} to be separated from M^{4+} . In some embodiments, the pH of a solution can be varied from ~ 8 to very strong acidic solutions ($\text{pH} \sim 0$) in order to separate various components within the solution. In some embodiments, a

metal species is formed when the pH of the solution is between 0 and 6 and other species are formed when the pH of the solution is between 6 and 8.

[0094] In some embodiments, a M^{3+} can be separated from a M^{4+} via a pH dependent process. In some embodiments, this can comprise dropping the pH of the solution to less than 2, which allows for trivalent metals to be separated from tetravalent metals. That is, in some embodiments, M^{4+} and M^{3+} can be separated by pH differences. While M^{4+} ions are still bound at $pH < 0$, M^{3+} ions are released at $pH < 2$, which is shown by the complexation of Sn^{4+} in 2 M HCl (FIG. 4). Other examples include Ce^{4+} and Th^{4+} complexes studied in 0.05 M H_2SO_4 , and Zr^{4+} and Pu^{4+} complexes studied in 0.05-0.5 M H_2SO_4 (*Inorg. Chem.* 2013 & *Inorg. Chem.* 2015 which are hereby incorporated by reference in their entireties). Thus, M^{4+} metals will remain associated with the chelator (e.g., 3,4,3-LI(1,2-HOPO)), while M^{3+} metals will still be free in solution. These M^{3+} metal ions can then be removed, for example, by addition of an organic metal binding compound that will be extracted into the organic phase of an extraction process. The HOPO ligand (with the M^{4+} metal ion) will stay in the aqueous phase). In some embodiments, the M^{3+} is actinium.

[0095] In some embodiments, any of these options involve the use of a highly acidic aqueous media as a feed solution. In some embodiments, where the HOPO ligand is adequately selective for the method, the aqueous solution need not be acidic.

[0096] Examples of commercial and non-selective extractant for liquid-liquid extraction under highly acidic conditions include, but are not limited to, trioctyl amine, quaternary ammonium salts (such as Aliquat® 336), TODGA, cyanex®, HDEHP, CMPO, etc.

[0097] In some embodiments, non-limiting characteristics of the extractant include an ability to transfer the targeted material from the aqueous phase to the organic phase, and solubility in an organic diluent and insoluble in aqueous media. In some embodiments, mixtures of two or more extractants can be used. In some embodiments, a pure form of the extractant can be expensive and, in most case, too viscous. Therefore, in some embodiments, the extractant is diluted in an organic diluent. In some embodiments, the diluent is usually an inert and inexpensive liquid (non-limiting examples include dodecane, toluene, etc.) and is only there to convey the diluted extractant.

[0098] In some embodiments, a solution comprising one or more of M^{2+} , M^{3+} , and M^{4+} can be separated into one or more part (e.g., M^{2+} only, M^{3+} only, M^{2+} and M^{3+} , M^{3+} and M^{4+} etc.), via the addition of a HOPO chelator, wherein the HOPO chelator will bind to the metal in a pH dependent manner.

[0099] In some embodiments, the metal ion is released from the chelator by lowering the pH beneath a level which allows for the metal binding. Thus, a pH of approximately -1 can be used to release the tetravalent metal from the HOPO molecule. Other non-limiting options include the pH be increased to about 10 in order to release the M^{4+} ions. In some embodiments, if the targeted metal is very valuable, the M^{4+} /HOPO solution can be evaporated and calcinated in order to burn the HOPO ligand and recover the metal. In some embodiments, the metal to be separated is Actinium.

Applications

[0100] In some embodiments, the process can be applied to the recycling of nuclear fuels (civilian and military) to the production of medical isotopes, and also analytical devices, production of radiation sources, etc. In some embodiments, typical examples of metal ions separation by liquid-liquid extraction would include the separation of plutonium (Pu^{4+}) from lanthanide fission products (Ln^{3+}), uranyl (UO_2^{2+}) and minor actinides (Am^{3+} , Cm^{3+}) in the frame of nuclear waste reprocessing.

[0101] This type of technology can be useful, for example, to reduce the environmental impact of nuclear energy the need for closed nuclear fuel cycles will become more critical. Thus far, the proposed plan for spent nuclear fuel in the United States has been long-term storage at a yet to be decided repository.¹ Storage of spent nuclear fuel becomes an issue because of longer-lived actinide (An) isotopes, like americium (Am) and curium (Cm), which are intense alpha-emitters and dominate the thermal heat load in spent fuels. *See*, Travis S. Grimes, Richard D. Tillotson, and Leigh R. Martin, "Trivalent Lanthanide/Actinide Separation Using Aqueous-Modified TALSPEAK Chemistry," *Solvent Extr. Ion Exch.* 32(4), 378-390 (2014); Kenneth L. Nash, "The Chemistry of TALSPEAK: A Review of the Science," *Solvent Extr. Ion Exch.* 33(1), 1-55 (2015).

[0102] Repositories must then be larger and better ventilated for longer periods of time, thus more expensive, to accommodate the minor actinide species from an open fuel

cycle. Fast-breeder reactors are the next generation of nuclear reactors and will have the potential to transmute Am and Cm into shorter-lived radionuclides, which will reduce the storage time, heat generated, and radiotoxicity of the spent fuels. However, lanthanide (Ln) fission products compete with An metals for neutrons and render the transmutation process less efficient. *See*, Kenneth L. Nash, “The Chemistry of TALSPEAK: A Review of the Science,” *Solvent Extr. Ion Exch.* 33(1), 1-55 (2015).

[0103] Minor actinides can be separated from lanthanides for the transmutation process but this separation can be challenging since Ln, Am, and Cm predominate as trivalent ions in solution and exhibit similar ionic-radii, thus almost identical chemistries. *See*, Travis S. Grimes, Richard D. Tillotson, and Leigh R. Martin, “Trivalent Lanthanide/Actinide Separation Using Aqueous-Modified TALSPEAK Chemistry,” *Solvent Extr. Ion Exch.* 32(4), 378-390 (2014). Some embodiments provided herein allow for the separation of some of the metals from one another.

[0104] Another typical example would be the purification of actinium (Ac^{3+}) thorium (Th^{4+}), and vice versa, in the frame of medical isotope production.

[0105] Another example would be separation of lutetium (Lu^{3+}) from hafnium (Hf^{4+}) or the separation of yttrium (Y^{3+}) from zirconium (Zr^{4+}) in the frame of medical isotope productions.

[0106] In some embodiments, the processes herein can be applied to separation of Sn/Zr or Sn/Hf in an acidic media. In some embodiments, the process can separate Sn/Ti at pH of about 7. In some embodiments, the process can separate Ti/Zr or Ti/Hf at pH 6-11. In some embodiments, the process can separate Sn/Zr or Sn/Hf in acidic media. In some embodiments, the process can further separate Sn/Ti at pH 7.

[0107] In some embodiments, the process provides for industrial processes that are simpler and more efficient than the current ones, for the separation and/or purification of metal ions.

[0108] The applications of the invention are numerous, going from the recycling of nuclear fuels (civilian and military) to the production of medical isotopes, and also analytical devices, production of radiation sources, etc. Typical examples of metal ions separation by liquid-liquid extraction can include the separation of lanthanide fission products (Ln^{3+}) from minor actinides (Am^{3+} , Cm^{3+}) in the frame of nuclear waste

reprocessing. Another typical example would be the purification of actinium (Ac^{3+}) from lanthanide (Ln^{3+} and Ce^{4+}) and actinides (Th^{4+} , Am^{3+} , etc) in the frame medical isotope production. In some embodiments, this can be scaled-up to an industrial process and be simpler and more efficient than the current ones, for the separation and/or purification of lanthanide and actinide elements.

[0109] In some embodiments, one or more of the processes disclosed herein can be used to separate Bk from Ce, such as Ce^{4+} from Bk^{4+} . In some embodiments, one or more of the processes disclosed herein can be used to separate Bk^{4+} from An^{3+} . In some embodiments, one or more of the processes disclosed herein can be used to separate Pu from other An, such as in nuclear fuel waste processing. In some embodiments, one or more of the processes disclosed herein can be used for nuclear forensics.

[0110] In some embodiments, one or more of the processes disclosed herein can be used to separate medical isotopes, such as Sc and/or Ac from Th, including Sc^{3+} and/or Ac^{3+} from Th^{4+} .

[0111] In some embodiments, one or more of the processes disclosed herein can be used to separate ions for nuclear forensics, including UO_2^{2+} from Pu^{4+} from Np^{4+} . Pu can be separated from Np via chromatography. In some embodiments, one or more of the processes disclosed herein can be used to separate UO_2^{2+} from Pu^{4+} and Np^{4+} .

[0112] In some embodiments, one or more of the processes disclosed herein can be used to separate Ce^{4+} from Tm^{3+} , such as in the processing of medical isotopes.

[0113] In some embodiments, one or more of the processes disclosed herein can be used to separate Zr^{4+} from Y^{3+} , such as in the processing of medical isotopes.

[0114] In some embodiments, a mixture of metal ions exhibiting different charges is obtained and these metal ions are to be separated in order to obtain the final product. For example, when reprocessing the nuclear spent fuels, the nuclear spent fuel rods are dissolved in highly acidic solutions (usually concentrated HNO_3). These acidic solutions contain Pu^{4+} but also trivalent lanthanide fission products (Ln^{3+}), trivalent actinides (Am^{3+} , Cm^{3+}), divalent uranyl ions (UO_2^{2+}) and monovalent neptunyl ($[\text{NpO}_2]^+$). Pu^{4+} eventually has to be separated from the other ions in order to reuse the plutonium material for other applications. The separation of Pu^{4+} from the other ions is currently performed at an industrial scale by using the liquid-liquid extraction process called "PUREX."

[0115] In some embodiments, the separation of metal ions involves a variation of the PUREX process (FIG. 5). The PUREX process involves recovery of U and Pu from nuclear spent fuel. The basic principle of the PUREX process involves the use of TBP which extracts Pu^{4+} and UO_2^{2+} but not M^{3+} . It is estimated that approximately 5040 t of nuclear fuel reprocessed annually in the UK, France, Japan, and Russia.

[0116] The PUREX process is based on the simultaneous extraction of Pu^{4+} and UO_2^{2+} in an organic phase comprising the molecule TBP (tributylphosphophate) while letting the Ln^{3+} , Am^{3+} , Cm^{3+} and $[\text{NpO}_2]^+$ ions in the original solution comprising nitric acid. The separation of Pu^{4+} from UO_2^{2+} requires further liquid-liquid extraction or back-extraction steps. The selectivity of such a process is based on differences in the affinity of TBP (present in the organic phase) for the different metal ions (initially in the organic phase). No reagent in the aqueous phase need be added to improve the selectivity of the process. In some embodiments, the PUREX process is modified in that one employs pH dependent HOPO chelating agent instead of TBP, under acidic conditions, to allow the HOPO ligand to selectively chelate to Pu^{4+} under acidic conditions, while not chelating the trivalent metals. The Pu^{4+} will remain in the aqueous phase, while the other metals will instead be moved to the organic phase. Thus, in this modified PUREX approach, the Pu^{4+} will remain in the acidic aqueous phase.

[0117] In some embodiments, trivalent actinium ions (Ac^{3+}) can be separated from thorium (Th^{4+}) in the frame of medical isotope production. The goal can be either to recover Ac^{3+} isotopes or Th^{4+} isotopes. Medical isotope productions sometimes also require to separate Zr^{4+} from yttrium (Y^{3+}). These processes usually involve dissolving a metallic target (containing the isotopes of interests, the initial metallic target material, and the impurities generated during the isotope production) in an acidic solution, such as HCl or HNO_3 , before separating the different metal ions by chromatographic or ion-exchange techniques.


[0118] TALSPEAK was developed in the 60's at Oak Ridge National Lab. *See*, Kenneth L. Nash, "The Chemistry of TALSPEAK: A Review of the Science," Solvent Extr. Ion Exch. 33(1), 1-55 (2015). TALSPEAK is a liquid-liquid extraction process that, in principle, allows the complete extraction of lanthanides from actinides by using a nonselective extractant and a hold-back reagent.

[0119] In some embodiments, TALSPEAK is used for the separation of An^{3+} or Ln^{3+} after PUREX process. The TALSPEAK process can also be varied given the disclosure herein. In some embodiments, the goal is to recover Am, Cm, Bk, Cf isotopes to ease the radioactive waste storage. The TALSPEAK process is still under development and has not been applied at an industrial level because of a few drawbacks: i) extraction efficiency decreases with increasing pH, ii) slow extraction kinetics, and iii) narrow set of conditions for optimum efficiency. *See*, Kenneth L. Nash, "The Chemistry of TALSPEAK: A Review of the Science," Solvent Extr. Ion Exch. 33(1), 1-55 (2015). Much of the effort in improving the TALSPEAK process has revolved around tackling the aforementioned issues and simplifying the implementation of this process into the nuclear cycle scheme. FIG. 7 shows an embodiment of the TALSPEAK process (adapted from Johnson and Maloney 2013 US Patent 8475747 B1, which is hereby incorporated by reference in its entirety).

[0120] The metal ion separation process can be described according to Equation (1),³ where $M = An^{3+}$ or Ln^{3+} , H_5DTPA is the pentaprotic hold-back reagent, and HDEHP is the non-selective extractant. The selectivity in the process arises from the hold-back reagent, which can be for example, Diethylenetriamine-N, N, N', N'', N'''-pentaacetic acid (DTPA). Actinides exhibit a higher degree of covalency with respect to their 4-f analogues⁴, thus will form stronger complexes with DTPA. Therefore, for trivalent actinides and lanthanides the equilibrium in the Equation (1) will lie to the left and right, respectively.



[0121] The structures of DTPA and HDEHP are as follows:



with Diethylenetriamine-N, N, N', N'', N''-pentaacetic acid (DTPA) shown to the left and di(2-ethylhexyl)phosphoric acid (HDEHP) to the right.

[0122] In some embodiments, 3,4,3-LI(1,2-HOPO) can be used as a replacement for DTPA in the TALSPEAK process. Non-limiting example of the TALSPEAK process with a HOPO chelator is shown in FIG. 37. In some embodiments, the process can be a TALSPEAK process, but be at a pH of 0-3 (rather than 3-4), e.g., a pH of 0.2.9, 0-2.8, 0-2.7, 0-2.6, 0-2.5, 0.5-2, etc.

[0123] The Trivalent Actinide-Lanthanide Separation by Phosphorous reagent Extraction from Aqueous Komplexes (TALSPEAK) process addresses the issue by selectively removing the long-lived minor actinides (Am, Cm) from fission products using liquid-liquid extractions, thus decreasing the storage size and time, and radiotoxicity of the raffinate. In some embodiments, this TALSPEAK process can be modified by using a HOPO chelator shown to form stronger complexes than the current hold-back reagent: diethylenetriamine-N,N,N',N'',N''-pentaacetic acid (DTPA). Evidence was observed for the selective extraction Ac(III) over Gd(III) and Am(III) at pH >2.5. Furthermore, the separation of Gd(III) over Am(III) can be achieved when using HOPO chelators under conditions similar to that of the TALKSPEAK process.

[0124] In the TALSPEAK scheme, lactate serves as a buffering agent and to speed up transfer kinetics, bis-2-ethylhexyl phosphoric acid (HDEHP) is a nonselective extractant in the organic phase, DTPA is the current TALSPEAK hold-back reagent in the aqueous phase, and enterobactin is a naturally found iron chelator (FIG 6).

[0125] Preliminary studies were done by modeling the species in solution as a function of pH for 40 mM hold-back reagent and 500 μ M Eu(III) (FIG. 8 and FIG. 9). Europium was used in the speciation diagram since the stability constants with 3,4,3-LI(1,2-HOPO) have been well characterized. *See*, Rebecca J. Abergel, Anthony D'Aleo, Clara Ng Pak Leung, David K. Shuh, and Kenneth N. Raymond, "Using the Antenna Effect as a Spectroscopic Tool: Photophysics and Solution Thermodynamics of the Model Luminescent Hydroxypyridonate Complex [EuIII(3,4,3-LI(1,2-HOPO))]-," *Inorg. Chem.* 48, 10868-10870 (2009). The binding of Eu(III) by DTPA and 3,4,3-LI(1,2-HOPO) by using available thermodynamic data at 0.1 M ionic strength at 25°C were compared. Typical TALSPEAK conditions are > 0.1 M ionic strength, due to needing strict pH control which is provided by

sodium lactate (0.5-2.0 M). *See*, Kenneth L. Nash, “The Chemistry of TALSPEAK: A Review of the Science,” *Solvent Extr. Ion Exch.* 33(1), 1-55 (2015). Therefore, extensive investigations of Ln-lactate stability constants have been obtained at high ionic strength, which are lacking for HOPO chelators. Nonetheless, this can be used to get an overview of the region where metal complexation is expected to occur. The calculations were made in the absence of lactate since lactate ($\log\beta_{130} = 5.88$ for Eu(III), $I = 2.0$ M at 25°C) (*See*, Kenneth L. Nash, “The Chemistry of TALSPEAK: A Review of the Science,” *Solvent Extr. Ion Exch.* 33(1), 1-55 (2015)) is a much weaker chelator than 3,4,3-LI(1,2-HOPO) ($\log\beta_{111} = 24.8$ for Eu(III), $I = 0.1$ M at 25°C)⁶ and DTPA ($\log\beta_{111} = 22.56$ for Eu(III), $I = 2.0$ M at 25°C) (*See*, Kenneth L. Nash, “The Chemistry of TALSPEAK: A Review of the Science,” *Solvent Extr. Ion Exch.* 33(1), 1-55 (2015)). In FIG. 26, 3,4,3-LI(1,2-HOPO) binds Eu(III) at pH as low as 0, whereas with DTPA (FIG. 8) free metal was present below pH 1. Thus, FIG. 8 shows that DTPA shows no binding at $\text{pH} < 1$, i.e., Free Eu(III) is observed at $\text{pH} < 1$. Stability constants obtained from NIST database. *See*, Rebecca J. Abergel, Anthony D’Aleo, Clara Ng Pak Leung, David K. Shuh, and Kenneth N. Raymond, “Using the Antenna Effect as a Spectroscopic Tool: Photophysics and Solution Thermodynamics of the Model Luminescent Hydroxypyridonate Complex $[\text{Eu}^{\text{III}}(3,4,3\text{-LI}(1,2\text{-HOPO}))]$,” *Inorg. Chem.* 48, 10868-10870 (2009). In contrast, FIG 26 shows that 3,4,3-LI(1,2-HOPO) shows evidence for binding Eu(III) at low pH.

[0126] The model supports the use of 3,4,3-LI(1,2-HOPO) as metal complexing agent even at low pH, which is advantageous to the TALSPEAK process since the raffinate begins in high acid media. Moreover, working at lower pH would be beneficial from an industrial point of view since the use of a buffering agent may be avoided and the pH control would more robust. Additionally, the protonated 3,4,3-LI(1,2-HOPO):Eu complex is the dominant species over a relatively large pH range (0-4), forming a neutral species and potentially allowing for more favorable phase-transfer kinetics, which in the classic process lactate has been implicated in. *See*, Kenneth L. Nash, “The Chemistry of TALSPEAK: A Review of the Science,” *Solvent Extr. Ion Exch.* 33(1), 1-55 (2015)

[0127] FIG 21 shows a pH extraction profile of Gd(III)-153, Ac(III)-225, and Am(III)-243 with $[\text{Gd(III)-153}] = 0.21$ nM DTPA as hold-back reagent. At $[\text{Ac(III)-225}] = 0.4$

nM, $[Gd(III)] = 500 \mu M$, $[Am(III)-243] = 40.4 \text{ nM}$, selective extraction of Gd(III) at pH 3-4, just as expected for this system under TALSPEAK conditions is observed. In some embodiments, the same applies to different isotopes, including, for example, 153, 225, and 243.

[0128] FIG 28 shows a pH extraction profile of Gd(III)-153, Ac(III)-225, and Am(III)-243 with 3,4,3-LI(1,2-HOPO) as hold-back reagent. With $[Gd(III)-153] = 0.21 \text{ nM}$, $[Ac(III)-225] = 0.4 \text{ nM}$, $[Gd(III)_{Total}] = 500 \mu M$, $[Am(III)-243] = 40.4 \text{ nM}$, selective extraction of Ac(III) over Gd(III) and Am(III)-243 is seen above pH 2.5. FIG 24 shows evidence for the selective extraction of Gd(III) over Am(III) with 3,4,3-LI(1,2-HOPO) as hold-back reagent. In some embodiments, the same applies to different isotopes, including, for example, 153, 225, and 243.

[0129] In some embodiments, DTPA chelator is a relatively weak complexing agent for Ac(III). Separation of Ac(III) from Gd(III) and Am(III) can be achieved by using a HOPO chelator at pH>2.5. Also observed was evidence for the selective extraction of Gd(III) over Am(III) between pH 1-2.5.

[0130] In some embodiments, one can separate Nd(III) and Am(III), which is often the limiting pair in the Ln(III)/An(III) extractions.

[0131] As described above, metal extraction processes rely on the selectivity of the ligands and organic extractants. In some embodiments, the initial feeds in a metal extraction process are highly acidic in order to allow for dissolution of the feeds. Ligands used for the aqueous phases in the metal extraction processes are essentially carboxylic acids. However, aqueous ligands are usually not able to bind M^{n+} in acidic solutions (e.g., FIG. 10 for DTPA). An exception is HOPO ligands (FIG. 9). HOPO ligands are suitable for nuclear operations, are highly soluble, and need not have F or Cl atoms. In addition, 3,4,3-LI(1,2-HOPO) has an extreme affinity and selectivity for 4+ metal ions (TABLE 1).

TABLE 1 – COMPARISON OF HOPO WITH DTPA

	Log β_{110} (Th ⁴⁺)	Log β_{110} (Eu ³⁺)	$\Delta \log \beta$	Last pKa
3,4,3-LI(1,2-HOPO) ⁴⁻	40.1	20.2	19.9	6.64
DTPA ⁵⁻	28.7	22.4	6.3	10.4

[0132] In some embodiments, the TALSPEAK process can be modified by using 3,4,3-LI(1,2-HOPO), a promising f-element chelator shown to form stronger complexes than the current hold-back reagent: diethylenetriamine-N,N,N',N'',N''-pentaacetic acid (DTPA).

[0133] Given the lack of data on the lightest actinide element, actinium (Ac), extraction experiments under TALSPEAK conditions were also run for this actinide (Ac(III)-225 and Ac(III)-227). Evidence for the quantitative and selective extraction of Ac(III) over Gd(III) and Am(III) at pH >3 when using 3,4,3-LI(1,2-HOPO) was obtained. Thus, separation between metals of the same charge is also possible via this technique, as the various metals display varied affinity for the 3,4,3-LI(1,2-HOPO) ligand.

[0134] In some embodiments, one or more of the processes disclosed herein can be employed to separate a first M³⁺ from a second M³⁺. FIG. 11, FIG. 12 and FIG. 13 display the selectivity of 3,4,3-LI(1,2-HOPO) compared to DTPA. They give the calculated ratio between metals in the organic phase after extraction (what one recovers if one wants to purify something) if one had started with mixtures of Ac/Gd, Ac/Am or Gd/Am, respectively. FIG. 14 and FIG. 15 outline the extraction coefficient values comparing DTPA and 3,4,3-LI(1,2-HOPO), respectively.

[0135] In some embodiments, it is possible to obtain selective extraction of Ac(III) over Gd(III) and Am(III) at pH >3. Furthermore, it was possible to obtain higher extraction of Gd(III) over Am(III) at a lower pH range when using 3,4,3-LI(HOPO) as the hold-back reagent compared to DTPA and upon further optimization of the extraction condition expect to extract trivalent lanthanides over trivalent actinides


[0136] Furthermore, in some embodiments, higher extraction of Gd(III) over Am(III) at a lower pH range when using 3,4,3-LI(1,2-HOPO) as the hold-back reagent compared to DTPA. The separation factors Gd/Am were also higher in the case of 3,4,3-LI(1,2-HOPO). In some embodiments, trivalent lanthanides are extracted over trivalent actinides at lower pH which is a significant optimization of the TALSPEAK process.

[0137] Traditionally, actinium cannot be purified from a mixture of Ac^{3+} , Ln^{3+} and An^{3+} when using DTPA as a hold-back reagent because the metals would be partially extracted at the same time. Chelators of the DTPA family (i.e., carboxylic acids) that are commonly used in separation processes, such as EDTA, NTA and, DOTA, are likely to exhibit a similar non-selectivity toward actinium. In contrast, in some embodiments, a HOPO-based method combines two features: (i) quantitative extraction for actinium, and (ii) a very high decontamination capability. In some embodiments, HOPO ligands provide very high separation factors between actinium and the lanthanides or actinides, after only 1 extraction step.

[0138] 3,4,3-LI(1,2-HOPO) was investigated as a replacement for DTPA in the TALSPEAK process along with the behavior of Ac(III) with both hold-back reagents. Ac(III) was observed to behave similarly to Am(III) when using DTPA as the aqueous complexing agent and serves to show the dexterity of DTPA in the TALSPEAK process but lacks a pH range where selective extraction of Ac(III) over Am(III) is observed. Complete and selective extraction of Ac(III) in the liquid-liquid extractions with 3,4,3-LI(1,2-HOPO) was observed while retaining both Am(III) and Gd(III) in the aqueous phase. A selective, effective, and simplified separation process of Ac(III) from An(III) and Ln(III) can be envisioned from 3,4,3-LI(1,2-HOPO)'s behavior. Additionally, evidence for the selective extraction of Gd(III) over Am(III) at low pH is seen, a practical advantage to DTPA. Further probing of 3,4,3-LI(1,2-HOPO) with other f-elements is warranted to fully characterize it as a hold-back reagent. In some embodiments, the conditions used in the liquid-liquid extractions were from the classical TALSPEAK process but serve as a good starting point for the replacement of DTPA. Upon optimization of extraction conditions, 3,4,3-LI(1,2-HOPO) is expected to selectively separate An(III) (Am , Cm) from Ln fission products.



HOPO ligands for metal ion separation processes

[0139] A structure of an embodiment of a HOPO molecule is as follows:




Formula I


[0140] Suitable 1,2-HOPO chelating agent include, but are not limited to, molecules defined by the structure:




wherein R is a hydroxy group or R_1 where R_1 and R_2 are selected from the group consisting of H, $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$ and $-\text{CH}_2-\phi$, and X is either hydrogen, an alkali metal ion, or a quaternary ammonium ion.

[0141] Suitable 1,2-HOPO chelating agent include, but are not limited to, molecules incorporating a plurality of HOPO-type structures, including:






In some embodiments, the $-(CH_2)_m-$ chains can contain ether bonds $-(CH_2-O-CH_2)-$ or ramifications. $-(CH_2-CH(R')-CH_2)-$, or hydroxyl groups $-(CH_2-CH(OH)-CH_2)-$ etc. In some embodiments, one can substitute one or both of the oxygens on the HOPO ring with a sulfur.




In a particular embodiment of the invention, m is three. In a particular embodiment of the invention, m is three and n is four. In a particular embodiment of the invention, l and n are three, and m is four. The 1,2-HOPO and 3,2-HOPO chelating agents suitable for use are also taught in U.S. Pat. Nos. 4,698,431 ("Hydroxypyridonate Chelating Agents"), 5,634,901 ("3-Hydroxy-2(1H)-pyridonate Chelating Agents"), and 5,892,029 ("3-Hydroxy-2(1H)-

pyridonate Chelating Agents"), all of which are hereby incorporated by reference. Examples of appropriate chelators useful for the present methods described herein are shown in the Figures, Structure I, exemplary embodiments 1-51, and can include any of the HOPO formula provided herein, or other disclosed chelating formula, including those in figures 38B-38E, 39, and 44-45D and the claims as filed. Any of the chelators provided herein can be applied to the various methods of extraction provided herein relating to the various pH manipulations.

[0142] Suitable 1,2-HOPO and 3,2-HOPO chelating agents include, but are not limited to, a chelating agent comprised of a plurality of chelating functional units joined by one or more linking members, said chelating functional units independently selected from the group consisting of




in which at least one of said plurality of chelating functional units on said chelating agent is



wherein R₁ and R₂ are independently selected from the group consisting of hydrogen, C₁₋₄ aliphatic hydrocarbon groups, and C₁₋₄ aliphatic hydrocarbon groups substituted by a single halide, hydroxy, carboxy, acrylamido group or an aryl group, and R' is a member selected

from the group consisting of a bond to a linking member, a hydrogen atom, C₁₋₈ aliphatic hydrocarbon groups, aryl groups, and C₁₋₈ aliphatic hydrocarbon groups substituted by amino, carboxy, or hydroxy groups.

[0143] Suitable 3,2-HOPO chelating agents include, but are not limited to, a chelating agent having the structure:



wherein R₁ is a member selected from the group consisting of hydrogen, C₁₋₄ aliphatic hydrocarbon groups, and C₁₋₄ aliphatic hydrocarbon groups substituted by a single halide, hydroxy, carboxy, or aryl group; Z is a member selected from the group consisting of O, NH, N--alkyl, and N--aryl; a is 2-4; and b is 2-4.

[0144] The methods for synthesizing the 1,2-HOPO and 3,2-HOPO chelating agents are taught in U.S. Pat. Nos. 4,698,431; 5,634,901; and 5,892,029, all of which are hereby incorporated by reference.

[0145] In some embodiments, any siderophore based chelator that binds at a low pH can be employed in any of the processes provided herein.

[0146] In some embodiments, “3,4,3-LI(1,2-HOPO)” (FIG. 9; left) can be used as the chelator. This water-soluble chelator has previously been investigated for the complexation, *in vitro* or *in vivo*, of various metal ions, including: uranyl, lanthanide(III) ions, americium(III), curium(III), zirconium(IV), cerium(IV), thorium(IV), plutonium(IV) and, berkelium(IV). The solution thermodynamic properties of this molecule are therefore well documented. *See*, M. Sturzbecher-Hoehne, G.J.-P. Deblonde, R.J. Abergel, Solution thermodynamic evaluation of hydroxypyridinonate chelators 3,4,3-LI(1,2-HOPO) and 5-LIO(Me-3,2-HOPO) for UO₂(VI) and Th(IV) decorporation, *Radiochim. Acta.* 101 (2013)

359–366. doi:10.1524/ract.2013.2047; R.J. Abergel, A. D'Aléo, C. Ng Pak Leung, D.K. Shuh, K.N. Raymond, Using the Antenna Effect as a Spectroscopic Tool: Photophysics and Solution Thermodynamics of the Model Luminescent Hydroxypyridonate Complex [Eu(III)(3,4,3-LI(1,2-HOPO))]-, *Inorg. Chem.* 48 (2009) 10868–10870. doi:10.1021/ic9013703; M. Sturzbecher-Hoehne, C. Ng Pak Leung, A. D'Aléo, B. Kullgren, A.-L. Prigent, D.K. Shuh, K.N. Raymond, R.J. Abergel, 3,4,3-LI(1,2-HOPO): In vitro formation of highly stable lanthanide complexes translates into efficacious in vivo europium decorporation, *Dalton Trans.* 40 (2011) 8340. doi:10.1039/c1dt10840a; M. Sturzbecher-Hoehne, P. Yang, A. D'Aléo, R.J. Abergel, Intramolecular sensitization of americium luminescence in solution: shining light on short-lived forbidden 5f transitions, *Dalton Trans.* (2016). doi:10.1039/C6DT00328A; M. Sturzbecher-Hoehne, B. Kullgren, E.E. Jarvis, D.D. An, R.J. Abergel, Highly Luminescent and Stable Hydroxypyridinonate Complexes: A Step Towards New Curium Decontamination Strategies, *Chem. - Eur. J.* 20 (2014) 9962–9968. doi:10.1002/chem.201402103; M. Sturzbecher-Hoehne, T.A. Choi, R.J. Abergel, Hydroxypyridinonate Complex Stability of Group (IV) Metals and Tetravalent f-Block Elements: The Key to the Next Generation of Chelating Agents for Radiopharmaceuticals, *Inorg. Chem.* 54 (2015) 3462–3468. doi:10.1021/acs.inorgchem.5b00033; G.J.-P. Deblonde, M. Sturzbecher-Hoehne, R.J. Abergel, Solution Thermodynamic Stability of Complexes Formed with the Octadentate Hydroxypyridinonate Ligand 3,4,3-LI(1,2-HOPO): A Critical Feature for Efficient Chelation of Lanthanide(IV) and Actinide(IV) Ions, *Inorg. Chem.* 52 (2013) 8805–8811. doi:10.1021/ic4010246; G.J.-P. Deblonde, M. Sturzbecher-Hoehne, P.B. Rupert, D.D. An, M.-C. Illy, C.Y. Ralston, J. Brabec, W.A. de Jong, R.K. Strong, R.J. Abergel, Chelation and stabilization of berkelium in oxidation state +IV, *Nat. Chem.* (2017). doi:10.1038/nchem.2759.

[0147] The ligand 3,4,3-LI(1,2-HOPO) exhibits an unprecedented affinity for a broad variety of metal ions and it is particularly efficient for tetravalent ions such as Th⁴⁺ and Pu⁴⁺. FIG. 20 shows the stability constant values published for the complexes of 3,4,3-LI(1,2-HOPO). Stability constant values (log β_{ML}) for various complexes of 3,4,3-LI(1,2-HOPO). FIG. 20 shows that the ligand forms extremely stable complexes with trivalent and tetravalent ions but it also keeps a high selectivity toward tetravalent ions. $\beta_{ML} = [ML]_{eq} / ([M^{x+}]_{eq}[L^{n-}]_{eq})$. Data collected at 25 °C.

[0148] This particular chelator combines several features, including but not limited to (i) formation of extremely stable complexes for both trivalent and tetravalent ions, (ii) high selectivity toward tetravalent metal ions, and (iii) very fast complexation kinetics (metal ions bound within a few seconds). In some embodiments, the ligand has one or more of these properties.

[0149] Another feature of the 3,4,3-LI(1,2-HOPO) molecule is that it is able to bind the tetravalent ions (e.g., Th^{4+} , Pu^{4+}) even at high acidity. In some embodiments, results 3,4,3-LI(1,2-HOPO) stays bound to M^{4+} ions in solutions containing, at least, up to 6 mol/L H_3O^+ . This feature is unprecedented and unmatched by any of the common chelators used in the nuclear chemistry field such as ethylenediamine-tetraacetic acid (EDTA), diethylenetriamine-pentaacetic acid (DTPA), or tetraaza-cyclododecane-1,4,7,10-tetraacetic acid (DOTA).

Separation aspects

[0150] FIG. 21 shows the speciation for solutions containing the ligand and one equivalent of Eu^{3+} or Th^{4+} showing what species are found in aqueous solutions of 3,4,3-LI(1,2-HOPO). FIG. 21 shows a speciation diagram for 3,4,3-LI(1,2-HOPO) solutions of Eu^{3+} (left) and Th^{4+} (right). Conditions: $[\text{Metal}] = [\text{ligand}] = 1 \text{ mM}$. Hydroxide stability constants taken from the NIST database. See, A.E. Martell, R.M. Smith, R.J. Motekaitis, NIST Standard Reference Database 46, (n.d.). In the near-neutral pH region, the trivalent metal ions form a negatively charged complex $[\text{M(III)L}]^-$ whereas the tetravalent metal ions form a neutral complex $[\text{M(IV)L}]$. However, in the acidic pH range, below pH ~ 2 , the trivalent lanthanide ions are released by 3,4,3-LI(1,2-HOPO). In contrast, the tetravalent ions remain complexed to the ligand even at pH lower than 0. This clear difference in terms of metal speciation (*i.e.* free M^{3+} versus bound M^{4+}) can be leveraged for separating trivalent ions from tetravalent ions as, for example, when purifying actinium from $\text{Ac}^{3+}/\text{Th}^{4+}$ mixtures.

[0151] Thus far, no stability constant has been published for the actinium (Ac^{3+}) complex of 3,4,3-LI(1,2-HOPO). Based on the ionic radius of Ac^{3+} (1.22 Å; See, D. Lundberg, I. Persson, The size of actinoid(III) ions - structural analysis vs. common misinterpretations, *Coord. Chem. Rev.* 318 (2016) 131–134. doi:10.1016/j.ccr.2016.04.003) and stability constants determined for other actinide and lanthanide ions, the stability of the

Ac^{3+} complex is expected to be very low, with a $\log \beta_{[\text{AcL}]}$ value of about 12 (FIG. 22). Stability of constant of 3,4,3-LI(1,2-HOPO) complexes with trivalent ions as a function of the ionic radius of the metal is shown in FIG. 22. This graph shows that the stability of the actinium complex is expected to be low. Ionic radius values taken from Lundberg et al. for nona-coordinated metal ions. The Ac^{3+} data point is an estimate. *See*, Lundberg, I. Persson, The size of actinoid(III) ions - structural analysis vs. common misinterpretations, *Coord. Chem. Rev.* 318 (2016) 131–134. doi:10.1016/j.ccr.2016.04.003; D. Lundberg, I. Persson, L. Eriksson, P. D'Angelo, S. De Panfilis, Structural Study of the N,N'-Dimethylpropyleneurea Solvated Lanthanoid(III) Ions in Solution and Solid State with an Analysis of the Ionic Radii of Lanthanoid(III) Ions, *Inorg. Chem.* 49 (2010) 4420–4432. doi:10.1021/ic100034q. Hence, this weak stability compared to the other complexes can be used for separation purposes.

[0152] Given the solution thermodynamics of the 3,4,3-LI(1,2-HOPO) complexes, some embodiments provided involve highly selective processes based on the difference in affinity that 3,4,3-LI(1,2-HOPO) exhibits for the metal ions. The affinity of the ligand for the metals ions follows the order: $\text{M}^{4+} \gg \text{Cm}^{3+} > \text{Am}^{3+} > \text{Lu}^{3+} - \text{La}^{3+} \gg \text{Ac}^{3+}$. One can then expect to sequester selectively the impurities present in a given media (Th^{4+} , Ln^{3+} , etc.) while leaving the actinium ions unbound. In some embodiments, any of the metal ions can be separated from one another via their differences in affinity to the HOPO molecule, based on variables such as the amount of time of the extraction, amount of chelator added, pH value, amount of extractant in the organic phase, ratio between the organic phase and aqueous phase.

[0153] In the case of a liquid-liquid extraction process, the unbound Ac^{3+} ions can then be extracted in an organic phase while letting the impurities remain in the aqueous phase. Since the selectivity of the process comes from the aqueous chelator 3,4,3-LI(1,2-HOPO), any extractant able to transfer Ac^{3+} into an organic phase can be used and there is no need to develop a selective organic extractant for actinium. The separation method can then rely on commercially available compounds that are usually used in hydrometallurgy processes.

[0154] In some embodiments, the extract for Ac^{3+} ions into an organic phase can be one or more of: di-(2-ethylhexyl)phosphoric acid (HDEHP), calixarenes (*See*, C. Wai, D. Fisher, others, Carboxylate-derived calixarenes with high selectivity for actinium-225,

Chem. Commun. (1998) 377–378), diglycoamide derivatives (*See*, V. Radchenko, J.W. Engle, J.J. Wilson, J.R. Maassen, F.M. Nortier, W.A. Taylor, E.R. Birnbaum, L.A. Hudston, K.D. John, M.E. Fassbender, Application of ion exchange and extraction chromatography to the separation of actinium from proton-irradiated thorium metal for analytical purposes, J. Chromatogr. A. 1380 (2015) 55–63. doi:10.1016/j.chroma.2014.12.045), carbamoylphosphine oxide derivatives (*See*, V. Ostapenko, A. Vasiliev, E. Lapshina, S. Ermolaev, R. Aliev, Y. Totskiy, B. Zhuikov, S. Kalmykov, Extraction chromatographic behavior of actinium and REE on DGA, Ln and TRU resins in nitric acid solutions, J. Radioanal. Nucl. Chem. 306 (2015) 707–711. doi:10.1007/s10967-015-4331-y), and quaternary ammonium salts (*See*, L.R. Morss, N.M. Edelstein, J. Fuger, The Chemistry of the Actinide and Transactinide Elements, 4th ed., Springer, 2010).

[0155] In some embodiments, HOPO can be used for Zr^{4+} extraction. In some embodiments, Zr^{4+} -3,4,3-Ll(1,2-HOPO) complexes are stable from 6 M HCl (FIG. 29). In some embodiments, Zr^{4+} -3,4,3-Ll(1,2-HOPO) complexes are stable to pH 11 (FIG. 29).

[0156] In some embodiments, HOPO can be used for Hf^{4+} extraction. In some embodiments, Hf^{4+} -3,4,3-Ll(1,2-HOPO) complexes are stable from 6 M HCl. In some embodiments, Hf^{4+} -3,4,3-Ll(1,2-HOPO) complexes are stable to pH 11.

[0157] In some embodiments, Ti^{4+} -3,4,3-Ll(1,2-HOPO) complexes are stable, at least, from ~6 M HCl (FIG. 30). In some embodiments, Ti^{4+} -3,4,3-Ll(1,2-HOPO) complexes are stable to ~pH 6 (FIG. 30). In some embodiments, separation of Ti/Zr or Ti/Hf is possible at pH 6-11 using HOPO.

[0158] In some embodiments, Sn^{4+} -3,4,3-Ll(1,2-HOPO) complexes are stable from ~1 M HCl (FIG. 31). In some embodiments, Sn^{4+} -3,4,3-Ll(1,2-HOPO) complexes are stable to ~pH 8 (FIG. 31). In some embodiments, separation of Sn/Zr or Sn/Hf is possible in acidic media using HOPO. In some embodiments, separation of Sn/Ti is possible at pH ~7 using HOPO.

[0159] In some embodiments, speciation of M^{3+} or M^{2+} versus M^{4+} ions is readily different in the presence of 3,4,3-Ll(1,2-HOPO) (FIG. 32). In some embodiments, Ln^{3+} , An^{3+} and UO_2^{2+} are released below pH ~1 (FIG. 32).

[0160] In some embodiments, a possible application of 3,4,3-Ll(1,2-HOPO) is for grouped separation of M^{4+} ions versus others ions at low pH. In some embodiments, a

separation strategy is shown in FIG. 33. Non-limiting examples of applications of the separation strategy of FIG. 33 include separation of $^{227}\text{Th}^{4+}$ from $^{227}\text{Ac}^{3+}$ and $^{223}\text{Ra}^{2+}$, Pu^{4+} from Ln^{3+} , An^{3+} , TcO_4^- , UO_2^{2+} , NpO_2^+ , Sn^{4+} from Pb^{2+} , Hf^{4+} from $^{177}\text{Lu}^{3+}$, Ti^{4+} from Sc^{3+} , and Bk^{4+} from Cf^{3+} .

[0161] In some embodiments, 3,4,3-LI(1,2-HOPO) is a chelator that binds M^{4+} in really acidic conditions. The speciation of $\text{M}^{4+}/\text{M}^{3+}$ is readily different under acidic conditions. Therefore, in some embodiments, 3,4,3-LI(1,2-HOPO) can be used to suppress the extraction of M^{4+} .

[0162] In some embodiments, selective extraction of M^{3+} in the presence of 3,4,3-LI(1,2-HOPO) is possible (FIG. 26), for example, with classical REE extractants (e.g., HDEHP, CMPO, etc.).

[0163] In some embodiments, based on the results provided in the examples below, processes can be established for the recovery and/or purification of actinium from a mixture of actinium, thorium and lanthanides. The starting material is considered to be an acidic solution since spallation Th target are likely to be dissolved in a mineral acid aqueous solution.

[0164] In some embodiments, a first strategy would involve extracting solely actinium in an organic phase using HDEHP while letting the 3,4,3-LI(1,2-HOPO) complexes of lanthanide(III) and thorium(IV) in the aqueous phase (FIG. 24). A pH adjustment to ~ 3 would be necessary due to the use of HDEHP. The actinium ions would then be back-extracted by common procedures (contact with HCl or HNO_3) and the solvent would be recycled into the extraction step. The aqueous phase containing Th^{4+} and Ln^{3+} complexes could then be neutralized in order to precipitate the Th^{4+} and Ln^{3+} hydroxides and to recycle 3,4,3-LI(1,2-HOPO). The solid material could then be let for decay of the short-lived isotopes (^{227}Th and lanthanide fission products) and then reprocessed to make a new ^{232}Th spallation target.

[0165] In some embodiments, another strategy (FIG. 25) would involve of working directly on the acidic feed solution without pH adjustment. Under highly acidic conditions, (e.g., 3-7.5 M HNO_3), the lanthanide and actinium complexes of 3,4,3-LI(1,2-HOPO) will not be formed whereas the thorium(IV) complex will be formed. Due to this, Ac^{3+} ions could be extracted selectively from Th^{4+} but the Ln^{3+} ions are also likely to be

transferred into the organic phase (depending on the acidity and extractant used). In that case, the process would serve as an actinium recovery method as well as an $\text{Ac}^{3+}/\text{Th}^{4+}$ separation step. Further processing of the organic phase can be done in order to do the final $\text{Ac}^{3+}/\text{Ln}^{3+}$ separation, such as the one described in FIG. 31 or by the mean of a chromatography column (See, J.T. Harvey, Production of Actinium-225 via High Energy Proton Induced Spallation of Thorium-232. Final Technical Report DE-SC0003602, NorthStar Medical Radioisotopes, LLC, [https://world wide web .osti.gov/scitech/servlets/purl/1032445/](https://worldwideweb.osti.gov/scitech/servlets/purl/1032445/)).

Additional embodiments

[0166] TABLE 3 gives the formation constants of 3,4,3-LI(1,2-HOPO) complexes with various metal ions. As provided herein, the high affinity of 3,4,3-LI(1,2-HOPO) toward Ln^{3+} , An^{3+} and An^{4+} ions so that the chelator acts as a hold-back reagent for these ions while letting actinium being transferred into an organic phase.

TABLE 3
Stability constant ($\beta_{\text{ML}} = [\text{ML}]/([\text{M}][\text{L}])$) of different complexes
of the ligand 3,4,3-LI(1,2-HOPO).

Ion	Ionic radius (Å. CN = 8)	log β_{ML}
UO_2^{2+}	/	18.0 $\pm$ 0.4
Pb^{2+}	1.29	18.0 $\pm$ 0.2
Cd^{2+}	1.10	16.1 $\pm$ 0.3
La^{3+}	1.16	16.4 $\pm$ 0.4
Gd^{3+}	1.05	20.5 $\pm$ 0.1
Lu^{3+}	0.98	21.2 $\pm$ 0.1
Am^{3+}	1.09	20.4 $\pm$ 0.2
Th^{4+}	1.05	40.1 $\pm$ 0.5
Ac^{3+}	1.12	10-14
Ra^{2+}	1.48	Unknown

[0167] The chemical interactions between radium ions (Ra^{2+}) and the chelator 3,4,3-LI(1,2-HOPO) or its derivatives can be extrapolated from data related to interaction between 3,4,3-LI(1,2-HOPO) with other metal ions (i.e. Cd^{2+} , Pb^{2+} , UO_2^{2+} , and Sr^{2+}).

[0168] The behavior of Ra^{2+} ions is driven by two phenomena: (1) the ability of 3,4,3-LI(1,2-HOPO) to bind Ra^{2+} ions in the aqueous phase; (2) the ability of the chosen extractant to transfer non-complexed Ra^{2+} ions into the organic phase.

[0169] The ligand 3,4,3-LI(1,2-HOPO) usually exhibits a lower affinity for divalent cations compared to trivalent and tetravalent ions (TABLE 3). This ligand is also a hard donor and has a lower affinity for soft metal ions, as exemplified by the lower stability of the Cd^{2+} complex when compared to the Pb^{2+} complex (TABLE 3). With their large ionic radius, Ra^{2+} ions are expected to be weakly bound, or even not bound at all, by 3,4,3-LI(1,2-HOPO). Thus, the ligand is not expected to hold-back Ra^{2+} ions throughout the pH scale and especially not in the acidic region where competition with protons can occur. Hence, the main parameter to take into account for predicting the behavior of radium is the ability of the extractant to transfer Ra^{2+} ions into the organic phase.

[0170] Ra^{2+} ions do not readily form coordination compounds with organic chelators. Conversely, the purification of radium is usually done by extracting the other ions present in the media into the organic phase while letting radium in the aqueous phase. With the exception of crown ethers that form fairly strong complexes with radium and also strontium, classical organic extractants, like HDEHP, are not able to extract Ra^{2+} ions. Considering the process flowsheets proposed in the prior note, radium will most likely not be extracted and will be separated from actinium. FIG. 34 and FIG. 35 highlight the most likely behavior of radium ions during the extraction step of the proposed processes. As radium will end up in the aqueous raffinate, additional steps could eventually be developed for the selective recovery of radium. For example, selective precipitation of Ra^{2+} using barium sulfate is a common technique for the selective recovery of radium.

[0171] Another option, would involve storing the aqueous raffinate (containing thorium, radium and eventually the lanthanide fission products (FIG. 34 and FIG. 35) until the in-growth of Ac-225 from Ra-225 is deemed adequate and simply perform another extraction step in order to harvest a second batch of Ac-225. The cycle can eventually be repeated until the radium parent isotope has completely decayed.

[0172] In some embodiments, one lets thorium in the aqueous phase instead of extracting it. Likewise, actinium can be extracted in the proposed process, in some embodiments.

[0173] When thorium represents the main metal ion present in the spallation target, the volume of the feed aqueous solution (coming from the dissolution step) will be dictated by the solubility of the complex formed by 3,4,3-LI(1,2-HOPO) and Th^{4+} .

[0174] The solubility of the complex is also expected to vary with the pH value.

[0175] The aqueous solubility of the ligand itself was determined experimentally as 1 mol/L at room temperature. It has also been observed experimentally that a few percent's of dimethylformamide (DMF) or dimethylsulfoxide (DMSO) can increase the solubility of the ligand and its metal complexes.

[0176] In some embodiments, the dissolution step itself can be performed on a lower volume and the addition of the ligand and the downstream steps can be performed independently. A separate dissolution step could allow the use of a higher concentration of acid and consequently fasten the dissolution of the metallic target.

[0177] In some embodiments, since the process can extract microgram-quantities of actinium, the subsequent extraction steps can use a low ratio O/A (organic flow/aqueous flow). This has the advantage of minimizing the volume of actinium-containing solutions (loaded organic phase) as it concentrates the actinium flow. The downstream process would therefore require volumes in the range of hundreds of mL.

[0178] Some embodiments for the recovery and purification of actinium starting from a thorium spallation target is given in FIG. 36. In some embodiments, the actinium extraction part would use a low ratio O/A as mentioned above. In some embodiments, the number of stages to obtain a quantitative extraction of actinium can be determined by the McCabe & Thiele method. In some embodiments, the number of stages is low (1 to 3 stages). Indeed, experimental results already obtained show very high extraction efficiencies for actinium due to the low affinity of 3,4,3-LI(1,2-HOPO) for Ac^{3+} ions. In some embodiments, only micrograms of actinium need to be extracted, which is favorable for not saturating the organic phase and, as a consequence, will allow using a low number of extraction stages and a low O/A ratio. In some embodiments, larger amounts of Ac^{3+} ions can be extracted

[0179] Regarding the back extraction of actinium, since only micrograms quantities will need to be stripped from the loaded organic phase, a low number of back-extraction stages is expected (1 to 3). The dimensioning of the actinium back-extraction can also be evaluated by the McCabe & Thiele method in a near future. A high ratio O/A could potentially be used so that the final Ac-225 purified solution is as concentrated as possible. Preliminary results show the potential for using a biologically compatible buffer for the stripping of actinium from HDEHP-based organic solvents.

Extraction aspects

[0180] In some embodiments, the aqueous phase can be made up of water. In some embodiments, the aqueous phase also includes a HOPO-type chelator, a buffering agent to control the pH, dissolved inorganic salts or mineral acids (ex: NaNO₃, HNO₃, HCl, NaCl, H₂SO₄ etc.), and/or solubility improver (a few percent's of DMF or DMSO).

[0181] In some embodiments, the organic phase can be any organic solution or solvent, as long as it does not ineffectively dissolve or merge into the aqueous phase. In some embodiments, the organic phase can also include one or several extractant(s), an inert diluent (Ex: dodecane, kerosene, TPH, Elixore®, toluene, etc.), and/or a phase modifier in order to help the phases decant. In some embodiments, the volume ratio (or flows) of the organic phase and aqueous phase (usually called "O/A") can be different from 1. In some embodiments, the organic phase is selected from the group consisting of octanol, dodecanol, dichloromethane, chloroform, toluene, dimethylformamide, dimethylsulfoxide, octane, dodecane, kerosene.

[0182] In some embodiments, the aqueous volume is at least 1 mL, e.g., 1-100 or 1-1000 mL. In some embodiments, the aqueous volume is at least 1 L, e.g., 1-10L, 1-100, 1-1000, 1-10000, 10-10000, 10-100000, or 100-1000000L, or larger volumes. In some embodiments, the organic volume is at least 1 mL, e.g., 1-100 or 1-1000 mL. In some embodiments, the organic volume is at least 1 L, e.g., 1-10L, 1-100, 1-1000, 1-10000, 10-10000, 10-100000, or 100-1000000L, or larger volumes. In some embodiments, the organic phase is larger than the aqueous phase, e.g., 1.1, 1.5, 2, 5, 10, 20, 100, 1000 fold larger. In some embodiments, the volumes of the two phases are about the same. In some

embodiments, the aqueous phase is larger than the organic phase, e.g., 1.1, 1.5, 2, 5, 10, 20, 100, 1000 fold larger.

[0183] In some embodiments, the two phases are mixed at room temperature. In some embodiments, they are mixed at a temperature of freezing to just below boiling. In some embodiments, the mixture occurs at 1, 10, 20, 30, 40, 50, 60, 70, 80, 90, or 99 degrees Celsius.

[0184] In some embodiments, the extraction and/or mixing occurs for as long as effective or necessary and can be repeated as many times as desired in order to obtain the purification and/or enrichment desired. In some embodiments, this is for 1-10 minutes, 1-60 minutes, 1 hour or more, e.g., 1, 2, 3, 4, 5, 6, 10, 20 or 24 hours, or 1 day or more. In some embodiments, the extraction mixing can occur 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more times. In some embodiments, the extraction process (add, agitate, decant) can occur 1 or more times, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more times.


[0185] As will be appreciated by one of skill in the art, often the present disclosure refers to the metal being chelated by the HOPO molecule as the desired metal for enrichment, isolation, collection. However, each of these embodiments can equally be applied to the other metal that is separated from the HOPO chelator (into the organic phase) for various embodiments. Thus, each process disclosed herein regarding a chelator separation, is envisioned not only as an option for collecting the metal associated with the chelator (in the aqueous phase) but also as an option for collecting the metal that is separated therefrom (in the organic phase). This applies for all embodiments provided herein.

[0186] Although this disclosure is in the context of certain embodiments and examples, those skilled in the art will understand that the present disclosure extends beyond the specifically disclosed embodiments to other alternative embodiments and/or uses of the embodiments and obvious modifications and equivalents thereof. In addition, while several variations of the embodiments have been shown and described in detail, other modifications, which are within the scope of this disclosure, will be readily apparent to those of skill in the art based upon this disclosure. It is also contemplated that various combinations or sub-combinations of the specific features and aspects of the embodiments may be made and still fall within the scope of the disclosure. It should be understood that various features and aspects of the disclosed embodiments can be combined with, or substituted for, one another

in order to form varying modes or embodiments of the disclosure. Thus, it is intended that the scope of the present disclosure herein disclosed should not be limited by the particular disclosed embodiments described above. For example, chemical conditions (e.g., acidity, concentrations, etc.) could be carried to match those of interest for various applications. In some embodiments, the ligand 3,4,3-LI(1,2-HOPO) could also be immobilized on a solid support if the chromatographic technique is preferred over the liquid-liquid extraction methods. Thus, in some embodiments, any extraction process described herein can instead be performed instead via a solid support and a chromatographic technique. Additionally, one or more embodiments herein can be combined with any one or more of the other embodiments herein.


In particular embodiments, chelators can include a number of metal-coordinating atoms that bond with a metal. The metal-coordinating atoms can bond with metals having cations with a +1 charge. The metal-coordinating atoms can also bond with metals having cations with a +2 charge. Additionally, the metal-coordinating atoms can bond with metals having cations with a +3 charge. Further, the metal-coordinating atoms can bond with metals having cations with a +4 charge. The chelators described herein can, in some cases, include siderophores.

In particular embodiments, the metal-coordinating atoms of the chelators described herein can be included in one or more functional groups of the chelators. In some examples, the metal-coordinating atoms of the chelators can be included in one or more catecholate (CAM) groups. A CAM group can include at least a phenyl ring substituted by hydroxyl groups on adjacent carbon atoms. According to some illustrative embodiments, a CAM group can include:




In some embodiments, the metal-coordinating atoms of the chelators can be included in one

or more hydroxamate (HA) groups. According to some embodiments, a HA group can include:



where R_a can include H or an alkyl group including no greater than 5 carbon atoms. For example, R_a can include a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an iso-butyl group, a tert-butyl group, a pentyl group, a tert-pentyl group, a neopentyl group, or an iso-pentyl group.

In some embodiments, the metal-coordinating atoms of the chelators can be included in one or more hydroxypyridinone (HOPO) groups. A HOPO group can include a pyridinone ring substituted by a hydroxyl group on the N atom. In some cases, a HOPO group can include a 1,2-HOPO group. According to some illustrative embodiments, a 1,2-HOPO group can include:



The metal-coordinating atoms of the chelators can be included in combinations of two or more of one or more CAM groups, one or more HA groups, or one or more HOPO groups. In illustrative examples, the metal-coordinating atoms of the chelators can be included in one or more CAM groups and one or more HA groups. In other illustrative examples, the metal-coordinating atoms of the chelators can be included in one or more CAM groups and one or more HOPO groups. In additional illustrative examples, the metal-coordinating atoms of the chelators can be included in one or more HA groups and one or more HOPO groups. In

further illustrative examples, the metal-coordinating atoms of the chelators can be included in one or more HA groups, one or more CAM groups, and one or more HOPO groups.

The chelators can include a number of functional groups having metal-coordinating atoms with the functional groups being bonded to a linear scaffold or a branched scaffold. The functional groups and/or substituents described herein may be substituted or unsubstituted. Substituted functional groups and/or substituents can be substituted by one or more hydroxyl groups, one or more alkyl groups having no greater than 10 carbon atoms, one or more amine groups, one or more thiol groups, one or more ester groups, or combinations thereof.


The scaffold can include one or more amine groups. An amine group can include a nitrogen atom bonded to three substituents. In particular embodiments, an amine group can include a nitrogen atom bonded at least one carbon atom of substituent. In various embodiments, an amine group can include a nitrogen atom bonded to at least a first carbon atom of a first substituent and a second carbon atom of a second substituent. In further embodiments, an amine group can include a nitrogen atom bonded to a first carbon atom of a first substituent, a second carbon atom of a second substituent and a third carbon atom of a third substituent. In certain embodiments, an amine group can include a nitrogen atom bonded to one or more hydrogen atoms.

In some embodiments, the scaffold can include one or more amide groups. An amide group can include a nitrogen atom bonded to a carbonyl group and two additional substituents. In various examples, an amide group can include a nitrogen atom bonded to a carbonyl group and a carbon atom of a first additional substituent. In other examples, an amide group can include a nitrogen atom bonded to a carbonyl group and a first carbon atom of a first additional substituent and a second carbon atom of a second additional substituent. In certain embodiments, an amine group can include a nitrogen atom bonded to one or more hydrogen atoms.

In particular embodiments, the scaffold can include one or more amine groups and one or more amide groups. The scaffold can include one or more carbon-based chains bonded between amine groups, a carbon-based chain bonded between amide groups, or one or more carbon-based chains bonded between a combination of one or more amine groups and one or

more amide groups. The carbon-based chains can include at least one carbon atom, at least 2 carbon atoms, at least 3 carbon atoms, at least 4 carbon atoms, or at least 5 carbon atoms. In addition, the carbon-based chains can include no greater than 10 carbon atoms, no greater than 9 carbon atoms, no greater than 8 carbon atoms, no greater than 7 carbon atoms, or no greater than 6 carbon atoms. In various embodiments, the carbon-based chains can include from 1 carbon atom to 10 carbon atoms, from 2 carbon atoms to 7 carbon atoms, or from 3 carbon atoms to 6 carbon atoms. In illustrative embodiments, the carbon-based chains can include alkane chains having carbon-carbon single bonds. In some cases, the carbon-based chains can include alkene chains having at least one carbon-carbon double bond. The carbon-based chains can be substituted or unsubstituted.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure I:



In some examples, A1, A2, A3, and A4 can, individually, include a CAM group, a HOPO group, or a HA group. Additionally, B1, B2, B3, and B4 can, individually, include an amide group or an amine group. Further, at least one of C1, C2, C3, C4, C5, or C6 can, individually, include NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. Also, in various examples, at least another one of C1, C2, C3, C4, C5, or C6 can be optional. In particular examples, at least one of L1, L2, L3, L4, L5, L6, L7, L8, L9, L10, L11, L12, or L13 can, individually, include H, an alkyl group having no greater than 10 carbon atoms, an alkylamino group having no greater than 10 carbon atoms and no greater than 2 nitrogen atoms; an alkylamido group having no greater than 10 carbon atoms and no greater than 2 nitrogen atoms; an alkyl ether group having no greater than 10 carbon atoms, a hydroxy ester group, or an alkyl ester group having no greater than 10 carbon atoms. In certain examples, at least one of L1, L5, L6, L7, L8, L9, L10, L11, L12, or L13 can be optional.

In illustrative examples, at least another one of L2, L3, or L4, can, individually, include an amine group or an amide group. In additional illustrative examples, L1, C1, L7, C2, L9, C3, L11, C4, and L13, C5 can be absent, L5 can include an alkyl group having no greater than 5 carbon atoms, and C6 can include NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. In further illustrative examples, L2, L3, L4, L6, L8, L10, and L12 can, individually, include an alkyl group having no greater than 5 carbon atoms. Also, A1 can include a CAM group or a 1, 2-HOPO group; A2 can include a HA group, A3 can include a HA group, and A4 can include a CAM group, a 1,2-HOPO group, or a HA group. In other illustrative examples, at least one of L2, L3, or L4 includes an alkylamino or alkylamido group.


In various illustrative examples, B1, B2, and B3 can, individually, include an amide group and B4 can include an amino group, L2 and L3 can include an amino group, and L4 can include an alky group having no greater than 5 carbon atoms. Additionally, C1, C2, C3, C4, C5, L1, A1, A2, A3, L1, L6, L7, L8, L9, L10, L11, L12, and L13 can be absent; A4 can include a CAM group, a 1,2-HOPO group, or a HA group; and L5 can include an alkyl group having no greater than 5 carbon atoms.

In certain illustrative examples, B1, B2, and B3 can include an amide group and B4 can

include an amide group; L2 and L3 can, individually, include an amino group; and L4 includes an alkyl group having no greater than 5 carbon atoms. Further, C1, C2, C3, C4, C5, A1, A2, A3, L1, L6, L7, L8, L9, L10, L11, and L13 can be absent, L12 can include an amino group, L5 can include an ether group having no greater than 10 carbon atoms, and A4 can include a CAM group, a 1,2-HOPO group, or a HA group.


In particular illustrative examples, C1, C2, C5, C6, L1, L2, L3, L4, L5, L7, L13, B2, and B4 can be absent; B1 and B3 can, individually, include an amide group; L6, L8, L10, and L12 can, individually, include an amino group, A1, A2, A3, and A4 can, individually, include a CAM group, a 1,2-HOPO group, or a HA group; and L9 and L11 can, individually, include an alkyl group having no greater than 5 carbon atoms.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure II:




R₁, R₂, R₃, R₄, and R₅ can, individually, include H, an alkyl group having from 1 to 10 carbon atoms, a CAM group, a HA group, or a 1,2-HOPO group. R₆ can include H, an alkyl group having from 1 to 10 carbon atoms, or an alkyl group having from 1 to 10 carbon atoms and substituted by at least one of NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. m can be from 1 to 6; n can be from 1 to 6; and o can be from 1 to 6. In particular embodiments, at least one of R₁, R₂, R₃, R₄, or R₅ can, individually, include a CAM group, a HA group, or a 1,2-HOPO group. In various embodiments, Structure II can include a linear, spermine-based backbone.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure III:




At least one of R_1 , R_3 , R_4 , or R_5 can, individually, include a CAM group, a HA group, or a 1,2-HOPO group. Optionally, another one of R_1 , R_3 , R_4 , or R_5 can, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R_2 can include H, OH, or an alkyl group including from 1 to 5 carbon atoms. p can be from 0 to 4. R_7 can include NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. In illustrative embodiments, R_1 can include a CAM group or a 1,2-HOPO group, R_3 and R_4 can, individually, include a HA group, and R_5 can include a CAM group, a 1,2-HOPO group, or a HA group.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure IV:




R_7 can include NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. R_2 , R_8 , and R_9 can, individually, include H, OH, or an alkyl group including from 1 to 5 carbon atoms. p can be from 0 to 4.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure V:




R_7 can include NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. R_2 , R_8 , and R_9 can, individually, include H, OH, or an alkyl group including from 1 to 5 carbon atoms. p can be from 0 to 4.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure VI:




R_7 can include NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. R_2 , R_8 , R_9 , and R_{10} can, individually, include H, OH, or an alkyl group including from 1 to 5 carbon atoms. p can be from 0 to 4.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure VII:




R_7 can include NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. R_2 , R_8 , and R_9 can, individually, include H, OH, or an alkyl group including from 1 to 5 carbon atoms. p can be from 0 to 4.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure VIII:




R_7 can include NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. R_2 , R_8 , and R_9 can, individually, include H, OH, or an alkyl group including from 1 to 5 carbon atoms. p can be from 0 to 4.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure IX:




R_7 can include NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. R_2 , R_8 , R_9 , and R_{10} can, individually, include H, OH, or an alkyl group including from 1 to 5 carbon atoms. p can be from 0 to 4.


In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure X:



In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XI:




In particular embodiments, compositions can have a branched backbone rather than the linear, spermine-based backbone of Structures III-XI. In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XII:



At least one of R₁₁, R₁₂, R₁₃, or R₁₅, can, individually, include a CAM group, a HA group, or a 1,2-HOPO group. Optionally, at least another one of R₁₁, R₁₂, R₁₃, or R₁₅ can, individually,


include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R_{17} can include NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. r can be from 0 to 6. R_2 , R_{14} , and R_{16} can, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms. In illustrative embodiments, R_{11} can include a CAM group or a 1,2-HOPO group, R_{12} and R_{15} can, individually, include a HA group, and R_{13} can include a CAM group, a 1,2-HOPO group, or a HA group.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XIII:




R_2 , R_{14} , R_{16} , R_{18} , and R_{19} can, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R_{17} can include NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. r can be from 0 to 4.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XIV:



R₂, R₁₄, R₁₆, R₁₈, and R₁₉ can, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R₁₇ can include NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. r can be from 0 to 4.


In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XV:



R₂, R₁₄, R₁₆, R₁₈, R₁₉, and R₂₀ can, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R₁₇ can include NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. r can be from 0 to 4.


In particular embodiments, compositions that function as chelators for radionuclides can

have the following structure, referred to herein as Structure XVI:



R₂, R₁₄, R₁₆, R₁₈, R₁₉, and R₂₀ can, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R₁₇ can include NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. r can be from 0 to 4.


In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XVII:



R₂, R₁₄, R₁₆, R₁₈, and R₁₉ can, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R₁₇ can include NH₂, C(=O)OH, maleimide, dibromo-maleimide,


isothiocyanate, alkyne, or azide. r can be from 0 to 4.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XVIII:



R_2 , R_{14} , R_{16} , R_{18} , and R_{19} can include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R_{17} can, individually, include NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. r can be from 0 to 4.


In particular embodiments, compositions can have a backbone that includes a number of amide groups and a number of amine groups. In some embodiments, the backbone of compositions that function as chelators for radionuclides can be based on Desferrioxamine B. In particular embodiments, compositions that function as chelators can have the following structure, referred to herein as Structure XIX:



R_{21} and R_{22} can include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R_{23} can include H, OH, an alkyl group having from 1 to 10 carbon atoms, or $(CH_2)_eR_a$, where R_a is NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide and e is


from 1 to 10. R_{24} can include a substituent that includes a CAM group, a 1,2-HOPO group, or a HA group. Optionally, R_{24} can include NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. a, b, and c can include from 1 to 10 and d can include from 1 to 4.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XX:



R_{25} , R_{26} , and R_{27} can, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R_{28} can include H, an alkyl group having from 1 to 5 carbon atoms, NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. s can be from 0 to 4.


In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XXI:



R_{25} , R_{26} , R_{27} , and R_{30} can, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R_{28} and R_{29} can, individually, include H, an alkyl group having from 1 to 5


carbon atoms, NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. s can be from 0 to 4. t can be from 0 to 4.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XXII:



R_{25} , R_{26} , R_{27} , and R_{30} can, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R_{28} and R_{29} can, individually, include H, an alkyl group having from 1 to 5 carbon atoms, NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. s can be from 0 to 4. t can be from 0 to 4.


In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XXIII:



R_{25} , R_{26} , R_{27} , R_{30} , and R_{31} can, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R_{28} and R_{29} can, individually, include H, an alkyl group having from 1 to


5 carbon atoms, NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. s can be from 0 to 4. t can be from 0 to 4.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XXIV:




R_{25} , R_{26} , and R_{27} can, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R_{28} can include H, an alkyl group having from 1 to 5 carbon atoms, NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. s can be from 0 to 4.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XXV:




R_{25} , R_{26} , R_{27} , and R_{32} can, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms. R_{28} can include H, an alkyl group having from 1 to 5 carbon atoms, NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. s can be from 0 to 4.

In particular embodiments, compositions can have an amide-based backbone. In particular embodiments, compositions that function as chelators can have the following structure, referred to herein as Structure XXVI:




A, B, C, and D can, individually, include one or more amide groups, one or more amine groups, or an alkyl group having from 1 to 10 carbon atoms. R_{33} , R_{34} , R_{35} , and R_{36} can, individually, include a CAM group, a 1,2-HOPO group, or a HA group. g , h , i , and j can, individually, be from 1 to 10.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XXVII:




R_{37} and R_{42} can, individually, include H, an alkyl group having from 1 to 5 carbon atoms, NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. R_{38} , R_{39} , R_{40} , and R_{41} can, individually, include H, OH, or an alkyl group having from 1 to 5 carbon atoms. u and v can, individually, be from 0 to 5.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XXVIII:




R_{37} and R_{42} can, individually, include H, an alkyl group having from 1 to 5 carbon atoms, NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. R_{38} , R_{39} , R_{40} , and R_{41} can, individually, include H, OH, or an alkyl group having from 1 to 5 carbon atoms. u and v can, individually, be from 0 to 5.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure XXIX:



R_{37} and R_{42} can, individually, include H, an alkyl group having from 1 to 5 carbon atoms, NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide. R_{38} , R_{39} , R_{40} , and R_{41} can, individually, include H, OH, or an alkyl group having from 1 to 5 carbon atoms. u and v can, individually, be from 0 to 5.

In particular embodiments, compositions that function as chelators for radionuclides can have the following structure, referred to herein as Structure L:



wherein: at least one of R_1 , R_2 , R_3 , R_4 , R_5 , individually, comprise a CAM group, a HA group, or a 1,2-HOPO group; at least another one of R_1 , R_2 , R_3 , R_4 , and R_5 , individually, comprise H or an alkyl group having from 1 to 10 carbon atoms; R_6 comprises (i) H, (ii) an alkyl group having from 1 to 10 carbon atoms, or (iii) an alkyl group having from 1 to 100 carbon atoms and no greater than 2 nitrogen atoms, and substituted by at least one of NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, amide, fluorescent moiety, or azide; m can be from 1 to 6; n can be from 1 to 6; o can be from 1 to 6; p can be from 1 to 6; q can be from 0 to 6; r can be from 1 to 6; s can be from 1 to 6; t can be from 1 to 6.

In some embodiments, the chelator is any one or more of the chelators in any of the figures. In some embodiments, the chelator has a structure as depicted in FIG. 38E. In some embodiments, the chelator is made by any one or more of the flow charts provided in the present figures. In some embodiments, the chelator is a peptoid chelator.

Examples of radioisotopes useful for isolation, collection, enrichment, and/or use etc. include ^{225}Ac , ^{226}Ac , ^{228}Ac , ^{105}Ag , ^{106}mAg , ^{110}mAg , ^{111}Ag , ^{112}Ag , ^{113}Ag , ^{239}Am , ^{240}Am , ^{242}Am , ^{244}Am , ^{37}Ar , ^{71}As , ^{72}As , ^{73}As , ^{74}As , ^{76}As , ^{77}As , ^{209}At , ^{210}At , ^{191}Au , ^{192}Au , ^{193}Au , ^{194}Au , ^{195}Au , ^{196}Au , ^{196m}Au , ^{198}Au , ^{198m}Au , ^{199}Au , ^{200m}Au , ^{128}Ba , ^{131}Ba , ^{133m}Ba , ^{135m}Ba , ^{140}Ba , ^7Be , ^{203}Bi , ^{204}Bi , ^{205}Bi , ^{206}Bi , ^{210}Bi , ^{212}Bi , ^{243}Bk , ^{244}Bk , ^{245}Bk , ^{246}Bk , ^{248m}Bk , ^{250}Bk , ^{76}Br , ^{77}Br , ^{80m}Br , ^{82}Br , ^{11}C , ^{14}C , ^{45}Ca , ^{47}Ca , ^{107}Cd , ^{115}Cd , ^{115m}Cd , ^{117m}Cd , ^{132}Ce , ^{133m}Ce , ^{134}Ce , ^{135}Ce , ^{137}Ce , ^{137m}Ce , ^{139}Ce , ^{141}Ce , ^{143}Ce , ^{144}Ce , ^{246}Cf , ^{247}Cf , ^{253}Cf , ^{254}Cf , ^{240}Cm , ^{241}Cm , ^{242}Cm , ^{252}Cm , ^{55}Co , ^{56}Co , ^{57}Co , ^{58}Co , ^{58m}Co , ^{60}Co , ^{48}Cr , ^{51}Cr , ^{127}Cs , ^{129}Cs , ^{131}Cs , ^{132}Cs , ^{136}Cs , ^{137}Cs , ^{61}Cu , ^{62}Cu , ^{64}Cu , ^{67}Cu , ^{153}Dy , ^{155}Dy , ^{157}Dy , ^{159}Dy , ^{165}Dy , ^{166}Dy , ^{160}Er , ^{161}Er , ^{165}Er , ^{169}Er , ^{171}Er , ^{172}Er , ^{250}Es , ^{251}Es , ^{253}Es , ^{254}Es , ^{254m}Es , ^{255}Es , ^{256m}Es , ^{145}Eu , ^{146}Eu , ^{147}Eu ,


¹⁴⁸Eu, ¹⁴⁹Eu, ¹⁵⁰mEu, ¹⁵²mEu, ¹⁵⁶Eu, ¹⁵⁷Eu, ⁵²Fe, ⁵⁹Fe, ²⁵¹Fm, ²⁵²Fm, ²⁵³Fm, ²⁵⁴Fm, ²⁵⁵Fm, ²⁵⁷Fm, ⁶⁶Ga, ⁶⁷Ga, ⁶⁸Ga, ⁷²Ga, ⁷³Ga, ¹⁴⁶Gd, ¹⁴⁷Gd, ¹⁴⁹Gd, ¹⁵¹Gd, ¹⁵³Gd, ¹⁵⁹Gd, ⁶⁸Ge, ⁶⁹Ge, ⁷¹Ge, ⁷⁷Ge, ¹⁷⁰Hf, ¹⁷¹Hf, ¹⁷³Hf, ¹⁷⁵Hf, ¹⁷⁹m²Hf, ¹⁸⁰mHf, ¹⁸¹Hf, ¹⁸⁴Hf, ¹⁹²Hg, ¹⁹³Hg, ¹⁹³mHg, ¹⁹⁵Hg, ¹⁹⁵mHg, ¹⁹⁷Hg, ¹⁹⁷mHg, ²⁰³Hg, ¹⁶⁰mHo, ¹⁶⁶Ho, ¹⁶⁷Ho, ¹²³I, ¹²⁴I, ¹²⁶I, ¹³⁰I, ¹³²I, ¹³³I, ¹³⁵I, ¹⁰⁹In, ¹¹⁰In, ¹¹¹In, ¹¹⁴mIn, ¹¹⁵mIn, ¹⁸⁴Ir, ¹⁸⁵Ir, ¹⁸⁶Ir, ¹⁸⁷Ir, ¹⁸⁸Ir, ¹⁸⁹Ir, ¹⁹⁰Ir, ¹⁹⁰m²Ir, ¹⁹²Ir, ¹⁹³mIr, ¹⁹⁴Ir, ¹⁹⁴m²Ir, ¹⁹⁵mIr, ⁴²K, ⁴³K, ⁷⁶Kr, ⁷⁹Kr, ⁸¹mKr, ⁸⁵mKr, ¹³²La, ¹³³La, ¹³⁵La, ¹⁴⁰La, ¹⁴¹La, ²⁶²Lr, ¹⁶⁹Lu, ¹⁷⁰Lu, ¹⁷¹Lu, ¹⁷²Lu, ¹⁷⁴mLu, ¹⁷⁶mLu, ¹⁷⁷Lu, ¹⁷⁷mLu, ¹⁷⁹Lu, ²⁵⁷Md, ²⁵⁸Md, ²⁶⁰Md, ²⁸Mg, ⁵²Mn, ⁹⁰Mo, ⁹³mMo, ⁹⁹Mo, ¹³N, ²⁴Na, ⁹⁰Nb, ⁹¹mNb, ⁹²mNb, ⁹⁵Nb, ⁹⁵mNb, ⁹⁶Nb, ¹³⁸Nd, ¹³⁹mNd, ¹⁴⁰Nd, ¹⁴⁷Nd, ⁵⁶Ni, ⁵⁷Ni, ⁶⁶Ni, ²³⁴Np, ²³⁶mNp, ²³⁸Np, ²³⁹Np, ¹⁵O, ¹⁸²Os, ¹⁸³Os, ¹⁸³mOs, ¹⁸⁵Os, ¹⁸⁹mOs, ¹⁹¹Os, ¹⁹¹mOs, ¹⁹³Os, ³²P, ³³P, ²²⁸Pa, ²²⁹Pa, ²³⁰Pa, ²³²Pa, ²³³Pa, ²³⁴Pa, ²⁰⁰Pb, ²⁰¹Pb, ²⁰²mPb, ²⁰³Pb, ²⁰⁹Pb, ²¹²Pb, ¹⁰⁰Pd, ¹⁰¹Pd, ¹⁰³Pd, ¹⁰⁹Pd, ¹¹¹mPd, ¹¹²Pd, ¹⁴³Pm, ¹⁴⁸Pm, ¹⁴⁸mPm, ¹⁴⁹Pm, ¹⁵¹Pm, ²⁰⁴Po, ²⁰⁶Po, ²⁰⁷Po, ²¹⁰Po, ¹³⁹Pr, ¹⁴²Pr, ¹⁴³Pr, ¹⁴⁵Pr, ¹⁸⁸Pt, ¹⁸⁹Pt, ¹⁹¹Pt, ¹⁹³mPt, ¹⁹⁵mPt, ¹⁹⁷Pt, ²⁰⁰Pt, ²⁰²Pt, ²³⁴Pu, ²³⁷Pu, ²⁴³Pu, ²⁴⁵Pu, ²⁴⁶Pu, ²⁴⁷Pu, ²²³Ra, ²²⁴Ra, ²²⁵Ra, ⁸¹Rb, ⁸²Rb, ⁸²mRb, ⁸³Rb, ⁸⁴Rb, ⁸⁶Rb, ¹⁸¹Re, ¹⁸²Re, ¹⁸²mRe, ¹⁸³Re, ¹⁸⁴Re, ¹⁸⁴mRe, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁸⁹Re, ¹⁹⁰mRe, ⁹⁹Rh, ⁹⁹mRh, ¹⁰⁰Rh, ¹⁰¹mRh, ¹⁰²Rh, ¹⁰³mRh, ¹⁰⁵Rh, ²¹¹Rn, ²²²Rn, ⁹⁷Ru, ¹⁰³Ru, ¹⁰⁵Ru, ³⁵S, ¹¹⁸mSb, ¹¹⁹Sb, ¹²⁰Sb, ¹²⁰mSb, ¹²²Sb, ¹²⁴Sb, ¹²⁶Sb, ¹²⁷Sb, ¹²⁸Sb, ¹²⁹Sb, ⁴³Sc, ⁴⁴Sc, ⁴⁴mSc, ⁴⁶Sc, ⁴⁷Sc, ⁴⁸Sc, ⁷²Se, ⁷³Se, ⁷⁵Se, ¹⁵³Sm, ¹⁵⁶Sm, ¹¹⁰Sn, ¹¹³Sn, ¹¹⁷mSn, ¹¹⁹mSn, ¹²¹Sn, ¹²³Sn, ¹²⁵Sn, ⁸²Sr, ⁸³Sr, ⁸⁵Sr, ⁸⁹Sr, ⁹¹Sr, ¹⁷³Ta, ¹⁷⁵Ta, ¹⁷⁶Ta, ¹⁷⁷Ta, ¹⁸⁰Ta, ¹⁸²Ta, ¹⁸³Ta, ¹⁸⁴Ta, ¹⁴⁹Tb, ¹⁵⁰Tb, ¹⁵¹Tb, ¹⁵²Tb, ¹⁵³Tb, ¹⁵⁴Tb, ¹⁵⁴mTb, ¹⁵⁴m²Tb, ¹⁵⁵Tb, ¹⁵⁶Tb, ¹⁵⁶mTb, ¹⁵⁶m²Tb, ¹⁶⁰Tb, ¹⁶¹Tb, ⁹⁴Tc, ⁹⁵Tc, ⁹⁵mTc, ⁹⁶Tc, ⁹⁷mTc, ⁹⁹mTc, ¹¹⁸Te, ¹¹⁹Te, ¹¹⁹mTe, ¹²¹Te, ¹²¹mTe, ¹²³mTe, ¹²⁵mTe, ¹²⁷Te, ¹²⁷mTe, ¹²⁹mTe, ¹³¹mTe, ¹³²Te, ²²⁷Th, ²³¹Th, ²³⁴Th, ⁴⁵Ti, ¹⁹⁸Tl, ¹⁹⁹Tl, ²⁰⁰Tl, ²⁰¹Tl, ²⁰²Tl, ²⁰⁴Tl, ¹⁶⁵Tm, ¹⁶⁶Tm, ¹⁶⁷Tm, ¹⁶⁸Tm, ¹⁷⁰Tm, ¹⁷²Tm, ¹⁷³Tm, ²³⁰U, ²³¹U, ²³⁷U, ²⁴⁰U, ⁴⁸V, ¹⁷⁸W, ¹⁸¹W, ¹⁸⁵W, ¹⁸⁷W, ¹⁸⁸W, ¹²²Xe, ¹²⁵Xe, ¹²⁷Xe, ¹²⁹mXe, ¹³¹mXe, ¹³³Xe, ¹³³mXe, ¹³⁵Xe, ⁸⁵mY, ⁸⁶Y, ⁸⁷Y, ⁸⁷mY, ⁸⁸Y, ⁹⁰Y, ⁹⁰mY, ⁹¹Y, ⁹²Y, ⁹³Y, ¹⁶⁶Yb, ¹⁶⁹Yb, ¹⁷⁵Yb, ⁶²Zn, ⁶⁵Zn, ⁶⁹mZn, ⁷¹mZn, ⁷²Zn, ⁸⁶Zr, ⁸⁸Zr, ⁸⁹Zr, ⁹⁵Zr, and ⁹⁷Zr. In some embodiments, the following can be used/and/or collected and/or isolated and/or enriched: ¹⁴⁷Sm, ²²⁷Ac, ²³²Th, ²³²U, ²³³U, ²³⁴U, ²³⁵U, ²³⁶U, ²³⁷U, ²³⁸U, ²³⁷Np, ²³⁸Pu, ²³⁹Pu, ²⁴⁰Pu, ²⁴²Pu, ²⁴⁴Pu, ²⁴¹Am, ²⁴³Am, ²⁴²Cm, ²⁴³Cm, ²⁴⁴Cm, ²⁴⁵Cm, ²⁴⁶Cm, ²⁴⁷Cm, ²⁴⁸Cm, ²⁴⁷Bk, ²⁴⁹Bk, ²⁴⁹Cf, ²⁵²Cf.

The term analog (also structural analog or chemical analog) is used to refer to a compound that is structurally similar to another compound but differs with respect to a certain component, such as an atom, a functional group, or a substructure. The term derivative refers

to a compound that is obtained from a similar compound or a precursor compound by a chemical reaction. As used herein, analogs and derivatives retain the therapeutic effectiveness of the parent compound (i.e., there is no statistically significant difference in therapeutic activity according to an imaging assay or assessment of clinical improvement) or have improved therapeutic effectiveness as defined elsewhere herein.

Exemplary Embodiments.

1. A composition having a structure including:



wherein:

- (i) A1, A2, A3, and A4, individually, include a CAM group, a 1,2-HOPO group, or a HA group;
- (ii) B1, B2, B3, and B4, individually, include an amide group or an amine group;
- (iii) at least one of C1, C2, C3, C4, C5, or C6, individually, include NH_2 , $\text{C}(=\text{O})\text{OH}$,

maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;


(iv) at least another one of C1, C2, C3, C4, C5, or C6 is optional;

(v) at least one of L1, L2, L3, L4, L5, L6, L7, L8, L9, L10, L11, L12, or L13, individually, include H, an alkyl group having no greater than 10 carbon atoms, an alkylamino group having no greater than 10 carbon atoms and no greater than 2 nitrogen atoms; an alkylamido group having no greater than 10 carbon atoms and no greater than 2 nitrogen atoms; an alkyl ether group having no greater than 10 carbon atoms, a hydroxy ester group, or an alkyl ester group having no greater than 10 carbon atoms; and

(vi) at least one of L1, L5, L6, L7, L8, L9, L10, L11, L12, or L13 is optional.

2. A composition of embodiment 1, wherein at least another one of L2, L3, or L4, individually, include an amine group or an amide group.
3. A composition of embodiment 1 or embodiment 2, wherein L1, C1, L7, C2, L9, C3, L11, C4, and L13, C5 are absent, L5 includes an unsubstituted alkyl group having no greater than 5 carbon atoms, and C6 includes NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide.
4. A composition of embodiment 3, wherein L2, L3, L4, L6, L8, L10, and L12, individually, include an unsubstituted alkyl group having no greater than 5 carbon atoms.
5. A composition of embodiment 4, wherein A1 includes a CAM group or a HOPO group; A2 includes a HA group, A3 includes a HA group, and A4 includes a CAM group, a HOPO group, or a HA group.
6. A composition of any one of embodiments 1-5, wherein at least one of L2, L3, or L4, individually, include an alkylamino or alkylamido group.
7. A composition of embodiment 1, wherein B1, B2, and B3, individually, include an amide group and B4 includes an amino group, L2 and L3 include an amino group, and L4 includes an alkyl group having no greater than 5 carbon atoms
8. A composition of embodiment 7, wherein:
C1, C2, C3, C4, C5, L1, A1, A2, A3, L1, L6, L7, L8, L9, L10, L11, L12, and L13 are absent, A4 includes a CAM group, a HOPO group, or a HA group; and
L5 includes an alkyl group having no greater than 5 carbon atoms.
9. A composition of embodiment 1, wherein B1, B2, and B3, individually, include an

- amide group and B4 includes an amide group, L2 and L3, individually, include an amino group, and L4 includes an alkyl group having no greater than 5 carbon atoms.
10. A composition of embodiment 9, wherein C1, C2, C3, C4, C5, A1, A2, A3, L1, L6, L7, L8, L9, L10, L11, and L13 are absent, L12 includes an amino group, L5 includes an ether group having no greater than 10 carbon atoms, and A4 includes a CAM group, a HOPO group, or a HA group.
11. A composition of embodiment 1, wherein C1, C2, C5, C6, L1, L2, L3, L4, L5, L7, L13, B2, and B4 are absent, B1 and B3, individually, include an amide group, L6, L8, L10, and L12, individually, include an amino group, A1, A2, A3, and A4, individually, include a CAM group, a HOPO group, or a HA group, L9 and L11, individually, include an alkyl group having no greater than 5 carbon atoms.
12. A composition, including a structure:



wherein:

at least one of R₁, R₂, R₃, R₄, and R₅, individually, include a CAM group, a HA group, or a 1,2-HOPO group;

at least another one of R₁, R₂, R₃, R₄, and R₅, individually, include H or an alkyl group having from 1 to 10 carbon atoms;


R₆ includes (i) H, (ii) an alkyl group having from 1 to 10 carbon atoms, or (iii) an alkyl group having from 1 to 10 carbon atoms and substituted by at least one of NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;

m can be from 1 to 6;

n can be from 1 to 6;

o can be from 1 to 6.

13. A composition of embodiment 12, including a structure:



wherein:

at least one of R_1 , R_3 , R_4 , or R_5 ; R_1 , R_2 , R_3 , R_4 , and R_5 , individually, include a CAM group, a HA group, or a 1,2-HOPO group;

optionally, another one of R_1 , R_3 , R_4 , or R_5 ; R_1 , R_2 , R_3 , R_4 , and R_5 , individually, include H or an alkyl group having from 1 to 10 carbon atoms;

R_2 includes H or an alkyl group including from 1 to 5 carbon atoms;

R_7 includes NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide; and

p is from 0 to 4.


14. A composition of embodiment 13, wherein:

R_1 includes a CAM group or a 1,2-HOPO group;

R_3 and R_4 , individually, include a HA group; and

R_5 includes a CAM group, a 1,2-HOPO group, or a HA group.

15. A composition of embodiment 12, including a structure:




wherein:

R_7 includes NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;

R_2 , R_8 , and R_9 , individually, include H, OH, or an alkyl group including from 1 to 5 carbon atoms; and

p is from 0 to 4.

16. A composition of embodiment 12, including a structure:




wherein:

R_7 includes NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;

R_2 , R_8 , and R_9 , individually, include H, OH, or an alkyl group including from 1 to 5 carbon atoms; and

p is from 0 to 4.

17. A composition of embodiment 12, including a structure:




wherein:

R_7 includes NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;

R_2 , R_8 , R_9 , and R_{10} , individually, include H, OH, or an alkyl group including from 1 to 5 carbon atoms; and

p is from 0 to 4.

18. A composition of embodiment 12, including a structure:




wherein:

R_7 includes NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;

R_2 , R_8 , and R_9 , individually, include H, OH, or an alkyl group including from 1 to 5 carbon atoms; and

p is from 0 to 4.

19. A composition of embodiment 12, including a structure:




wherein:

R_7 includes NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;

R_2 , R_8 , and R_9 , individually, include H, OH, or an alkyl group including from 1 to 5 carbon atoms; and

p is from 0 to 4.

20. A composition of embodiment 12, including a structure:



wherein:


R_7 includes NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;

R_2 , R_8 , R_9 , and R_{10} , individually, include H, OH, or an alkyl group including from 1 to 5


carbon atoms; and

p is from 0 to 4.


21. A composition of embodiment 12, including a structure:



22. A composition of embodiment 12, including a structure:



23. A composition, including a structure:



wherein:

at least one of R_{11} , R_{12} , R_{13} , or R_{15} , individually, include a CAM group, a HA group, or a 1,2-HOPO group;

optionally, at least another one of R_{11} , R_{12} , R_{13} , or R_{15} , individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{17} includes NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;

R_2 , R_{14} , and R_{16} , individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms; and

r can be from 0 to 6.


24. A composition of embodiment 23, wherein:

R_{11} includes a CAM group or a 1,2-HOPO group;

R_{12} and R_{15} , individually, include a HA group; and

R_{13} includes a CAM group, a 1,2-HOPO group, or a HA group.

25. A composition of embodiment 23, including a structure:




wherein:

R_2 , R_{14} , R_{16} , R_{18} , and R_{19} , individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{17} includes NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide; and

r can be from 0 to 4.

26. A composition of embodiment 23, including a structure:




wherein

R_2 , R_{14} , R_{16} , R_{18} , and R_{19} , individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{17} includes NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide; and

r is from 0 to 4.

27. A composition of embodiment 23, including a structure:




wherein:

R₂, R₁₄, R₁₆, R₁₈, R₁₉, and R₂₀, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R₁₇ includes NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide; and

r can be from 0 to 4.

28. A composition of embodiment 23, including a structure:




wherein:

R_2 , R_{14} , R_{16} , R_{18} , R_{19} , and R_{20} , individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{17} includes NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide; and

r can be from 0 to 4.

29. A composition of embodiment 23, including a structure:




wherein:

R_2 , R_{14} , R_{16} , R_{18} , and R_{19} , individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{17} includes NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide; and

r is from 0 to 4.

30. A composition of embodiment 23, including a structure:




wherein:

R₂, R₁₄, R₁₆, R₁₈, and R₁₉, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R₁₇ includes NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;

r is from 0 to 4.

31. A composition, including a structure:



wherein:

R₂₁ and R₂₂, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R₂₃ includes H, OH, an alkyl group having from 1 to 10 carbon atoms, or (CH₂)_eR_a, where R_a is NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;


R_{25} , R_{26} , R_{27} , and R_{30} , individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{28} and R_{29} , individually, include H, an alkyl group having from 1 to 5 carbon atoms, NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;

s is from 0 to 4; and

t is from 0 to 4.

35. A composition of embodiment 31, including a structure:



wherein:


R_{25} , R_{26} , R_{27} , and R_{30} , individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{28} and R_{29} , individually, include H, an alkyl group having from 1 to 5 carbon atoms, NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, or azide;

s is from 0 to 4; and

t is from 0 to 4.

36. A composition of embodiment 31, including a structure:



wherein:


R₂₅, R₂₆, R₂₇, R₃₀, and R₃₁, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R₂₈ and R₂₉, individually, include H, an alkyl group having from 1 to 5 carbon atoms, NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, or azide;

s is from 0 to 4; and

t is from 0 to 4.

37. A composition of embodiment 31, including a structure:




wherein

R₂₅, R₂₆, and R₂₇, individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R₂₈ includes H, an alkyl group having from 1 to 5 carbon atoms, NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide; and

s is from 0 to 4.

38. A composition of embodiment 31, including a structure:




wherein:

R_{25} , R_{26} , R_{27} , and R_{32} , individually, include H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{28} includes H, an alkyl group having from 1 to 5 carbon atoms, NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide; and

s is from 0 to 4.

39. A composition, including a structure:




wherein:

A, B, C, and D, individually, include one or more amide groups, one or more amine groups, or an alkyl group having from 1 to 10 carbon atoms;

R_{33} , R_{34} , R_{35} , and R_{36} , individually, include a CAM group, a 1,2-HOPO group, or a HA group; and

g , h , i , and j , individually, are from 1 to 10.

40. A composition of embodiment 39, including a structure:




wherein:

R₃₇ and R₄₂, individually, include H, an alkyl group having from 1 to 5 carbon atoms, NH₂, C(=O)OH, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;

R₃₈, R₃₉, R₄₀, and R₄₁, individually, include H, OH, or an alkyl group having from 1 to 5 carbon atoms; and

u and v, individually, are from 0 to 5.

41. A composition of embodiment 39, including a structure:




wherein:

R_{37} and R_{42} , individually, include H, an alkyl group having from 1 to 5 carbon atoms, NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;

R_{38} , R_{39} , R_{40} , and R_{41} , individually, include H, OH, or an alkyl group having from 1 to 5 carbon atoms; and

u and v , individually, are from 0 to 5.

42. A composition of embodiment 39, including a structure:



wherein:


R_{37} and R_{42} , individually, include H, an alkyl group having from 1 to 5 carbon atoms, NH_2 , $C(=O)OH$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, or azide;

R_{38} , R_{39} , R_{40} , and R_{41} , individually, include H, OH, or an alkyl group having from 1 to 5 carbon atoms; and

u and v , individually, are from 0 to 5.

43. A method of synthesizing a siderophore using dichlorodiphenylmethane.

44. A method of embodiment 43, wherein a siderophore or siderophore-like ligand is synthesized according to the following pathway:



45. A composition of any one of embodiment s1-42, wherein the metal is a radionuclide.


46. A composition of embodiment 53, wherein the radionuclide includes ^{225}Ac , ^{226}Ac , ^{228}Ac , ^{105}Ag , ^{106}mAg , ^{110}mAg , ^{111}Ag , ^{112}Ag , ^{113}Ag , ^{239}Am , ^{240}Am , ^{242}Am , ^{244}Am , ^{37}Ar , ^{71}As , ^{72}As , ^{73}As , ^{74}As , ^{76}As , ^{77}As , ^{209}At , ^{210}At , ^{191}Au , ^{192}Au , ^{193}Au , ^{194}Au , ^{195}Au , ^{196}Au , ^{196}mAu , ^{198}Au , ^{198}mAu , ^{199}Au , ^{200}mAu , ^{128}Ba , ^{131}Ba , ^{133}mBa , ^{135}mBa , ^{140}Ba , ^7Be , ^{203}Bi , ^{204}Bi , ^{205}Bi , ^{206}Bi , ^{210}Bi , ^{212}Bi , ^{243}Bk , ^{244}Bk , ^{245}Bk , ^{246}Bk , ^{248}mBk , ^{250}Bk , ^{76}Br , ^{77}Br , ^{80}mBr , ^{82}Br , ^{11}C , ^{14}C , ^{45}Ca , ^{47}Ca , ^{107}Cd , ^{115}Cd , ^{115}mCd , ^{117}mCd , ^{132}Ce , ^{133}mCe , ^{134}Ce , ^{135}Ce , ^{137}Ce , ^{137}mCe , ^{139}Ce , ^{141}Ce , ^{143}Ce , ^{144}Ce , ^{246}Cf , ^{247}Cf , ^{253}Cf , ^{254}Cf , ^{240}Cm , ^{241}Cm , ^{242}Cm , ^{252}Cm , ^{55}Co , ^{56}Co , ^{57}Co , ^{58}Co , ^{58}mCo , ^{60}Co , ^{48}Cr , ^{51}Cr , ^{127}Cs , ^{129}Cs , ^{131}Cs , ^{132}Cs , ^{136}Cs , ^{137}Cs , ^{61}Cu , ^{62}Cu , ^{64}Cu , ^{67}Cu , ^{153}Dy , ^{155}Dy , ^{157}Dy , ^{159}Dy , ^{165}Dy , ^{166}Dy , ^{160}Er , ^{161}Er , ^{165}Er , ^{169}Er , ^{171}Er , ^{172}Er , ^{250}Es , ^{251}Es , ^{253}Es , ^{254}Es , ^{254}mEs , ^{255}Es , ^{256}mEs , ^{145}Eu , ^{146}Eu , ^{147}Eu , ^{148}Eu , ^{149}Eu , ^{150}mEu , ^{152}mEu , ^{156}Eu , ^{157}Eu , ^{52}Fe , ^{59}Fe , ^{251}Fm , ^{252}Fm , ^{253}Fm , ^{254}Fm , ^{255}Fm , ^{257}Fm , ^{66}Ga , ^{67}Ga , ^{68}Ga , ^{72}Ga , ^{73}Ga , ^{146}Gd , ^{147}Gd , ^{149}Gd , ^{151}Gd , ^{153}Gd , ^{159}Gd , ^{68}Ge , ^{69}Ge , ^{71}Ge , ^{77}Ge , ^{170}Hf , ^{171}Hf , ^{173}Hf ,

¹⁷⁵Hf, ¹⁷⁹m²Hf, ¹⁸⁰mHf, ¹⁸¹Hf, ¹⁸⁴Hf, ¹⁹²Hg, ¹⁹³Hg, ¹⁹³mHg, ¹⁹⁵Hg, ¹⁹⁵mHg, ¹⁹⁷Hg, ¹⁹⁷mHg, ²⁰³Hg, ¹⁶⁰mHo, ¹⁶⁶Ho, ¹⁶⁷Ho, ¹²³I, ¹²⁴I, ¹²⁶I, ¹³⁰I, ¹³²I, ¹³³I, ¹³⁵I, ¹⁰⁹In, ¹¹⁰In, ¹¹¹In, ¹¹⁴mIn, ¹¹⁵mIn, ¹⁸⁴Ir, ¹⁸⁵Ir, ¹⁸⁶Ir, ¹⁸⁷Ir, ¹⁸⁸Ir, ¹⁸⁹Ir, ¹⁹⁰Ir, ¹⁹⁰m²Ir, ¹⁹²Ir, ¹⁹³mIr, ¹⁹⁴Ir, ¹⁹⁴m²Ir, ¹⁹⁵mIr, ⁴²K, ⁴³K, ⁷⁶Kr, ⁷⁹Kr, ⁸¹mKr, ⁸⁵mKr, ¹³²La, ¹³³La, ¹³⁵La, ¹⁴⁰La, ¹⁴¹La, ²⁶²Lr, ¹⁶⁹Lu, ¹⁷⁰Lu, ¹⁷¹Lu, ¹⁷²Lu, ¹⁷⁴mLu, ¹⁷⁶mLu, ¹⁷⁷Lu, ¹⁷⁷mLu, ¹⁷⁹Lu, ²⁵⁷Md, ²⁵⁸Md, ²⁶⁰Md, ²⁸Mg, ⁵²Mn, ⁹⁰Mo, ⁹³mMo, ⁹⁹Mo, ¹³N, ²⁴Na, ⁹⁰Nb, ⁹¹mNb, ⁹²mNb, ⁹⁵Nb, ⁹⁵mNb, ⁹⁶Nb, ¹³⁸Nd, ¹³⁹mNd, ¹⁴⁰Nd, ¹⁴⁷Nd, ⁵⁶Ni, ⁵⁷Ni, ⁶⁶Ni, ²³⁴Np, ²³⁶mNp, ²³⁸Np, ²³⁹Np, ¹⁵O, ¹⁸²Os, ¹⁸³Os, ¹⁸³mOs, ¹⁸⁵Os, ¹⁸⁹mOs, ¹⁹¹Os, ¹⁹¹mOs, ¹⁹³Os, ³²P, ³³P, ²²⁸Pa, ²²⁹Pa, ²³⁰Pa, ²³²Pa, ²³³Pa, ²³⁴Pa, ²⁰⁰Pb, ²⁰¹Pb, ²⁰²mPb, ²⁰³Pb, ²⁰⁹Pb, ²¹²Pb, ¹⁰⁰Pd, ¹⁰¹Pd, ¹⁰³Pd, ¹⁰⁹Pd, ¹¹¹mPd, ¹¹²Pd, ¹⁴³Pm, ¹⁴⁸Pm, ¹⁴⁸mPm, ¹⁴⁹Pm, ¹⁵¹Pm, ²⁰⁴Po, ²⁰⁶Po, ²⁰⁷Po, ²¹⁰Po, ¹³⁹Pr, ¹⁴²Pr, ¹⁴³Pr, ¹⁴⁵Pr, ¹⁸⁸Pt, ¹⁸⁹Pt, ¹⁹¹Pt, ¹⁹³mPt, ¹⁹⁵mPt, ¹⁹⁷Pt, ²⁰⁰Pt, ²⁰²Pt, ²³⁴Pu, ²³⁷Pu, ²⁴³Pu, ²⁴⁵Pu, ²⁴⁶Pu, ²⁴⁷Pu, ²²³Ra, ²²⁴Ra, ²²⁵Ra, ⁸¹Rb, ⁸²Rb, ⁸²mRb, ⁸³Rb, ⁸⁴Rb, ⁸⁶Rb, ¹⁸¹Re, ¹⁸²Re, ¹⁸²mRe, ¹⁸³Re, ¹⁸⁴Re, ¹⁸⁴mRe, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁸⁹Re, ¹⁹⁰mRe, ⁹⁹Rh, ⁹⁹mRh, ¹⁰⁰Rh, ¹⁰¹mRh, ¹⁰²Rh, ¹⁰³mRh, ¹⁰⁵Rh, ²¹¹Rn, ²²²Rn, ⁹⁷Ru, ¹⁰³Ru, ¹⁰⁵Ru, ³⁵S, ¹¹⁸mSb, ¹¹⁹Sb, ¹²⁰Sb, ¹²⁰mSb, ¹²²Sb, ¹²⁴Sb, ¹²⁶Sb, ¹²⁷Sb, ¹²⁸Sb, ¹²⁹Sb, ⁴³Sc, ⁴⁴Sc, ⁴⁴mSc, ⁴⁶Sc, ⁴⁷Sc, ⁴⁸Sc, ⁷²Se, ⁷³Se, ⁷⁵Se, ¹⁵³Sm, ¹⁵⁶Sm, ¹¹⁰Sn, ¹¹³Sn, ¹¹⁷mSn, ¹¹⁹mSn, ¹²¹Sn, ¹²³Sn, ¹²⁵Sn, ⁸²Sr, ⁸³Sr, ⁸⁵Sr, ⁸⁹Sr, ⁹¹Sr, ¹⁷³Ta, ¹⁷⁵Ta, ¹⁷⁶Ta, ¹⁷⁷Ta, ¹⁸⁰Ta, ¹⁸²Ta, ¹⁸³Ta, ¹⁸⁴Ta, ¹⁴⁹Tb, ¹⁵⁰Tb, ¹⁵¹Tb, ¹⁵²Tb, ¹⁵³Tb, ¹⁵⁴Tb, ¹⁵⁴mTb, ¹⁵⁴m²Tb, ¹⁵⁵Tb, ¹⁵⁶Tb, ¹⁵⁶mTb, ¹⁵⁶m²Tb, ¹⁶⁰Tb, ¹⁶¹Tb, ⁹⁴Tc, ⁹⁵Tc, ⁹⁵mTc, ⁹⁶Tc, ⁹⁷mTc, ⁹⁹mTc, ¹¹⁸Te, ¹¹⁹Te, ¹¹⁹mTe, ¹²¹Te, ¹²¹mTe, ¹²³mTe, ¹²⁵mTe, ¹²⁷Te, ¹²⁷mTe, ¹²⁹mTe, ¹³¹mTe, ¹³²Te, ²²⁷Th, ²³¹Th, ²³⁴Th, ⁴⁵Ti, ¹⁹⁸Tl, ¹⁹⁹Tl, ²⁰⁰Tl, ²⁰¹Tl, ²⁰²Tl, ²⁰⁴Tl, ¹⁶⁵Tm, ¹⁶⁶Tm, ¹⁶⁷Tm, ¹⁶⁸Tm, ¹⁷⁰Tm, ¹⁷²Tm, ¹⁷³Tm, ²³⁰U, ²³¹U, ²³⁷U, ²⁴⁰U, ⁴⁸V, ¹⁷⁸W, ¹⁸¹W, ¹⁸⁵W, ¹⁸⁷W, ¹⁸⁸W, ¹²²Xe, ¹²⁵Xe, ¹²⁷Xe, ¹²⁹mXe, ¹³¹mXe, ¹³³Xe, ¹³³mXe, ¹³⁵Xe, ⁸⁵Y, ⁸⁶Y, ⁸⁷Y, ⁸⁷mY, ⁸⁸Y, ⁹⁰Y, ⁹⁰mY, ⁹¹Y, ⁹²Y, ⁹³Y, ¹⁶⁶Yb, ¹⁶⁹Yb, ¹⁷⁵Yb, ⁶²Zn, ⁶⁵Zn, ⁶⁹mZn, ⁷¹mZn, ⁷²Zn, ⁸⁶Zr, ⁸⁸Zr, ⁸⁹Zr, ⁹⁵Zr, and/or ⁹⁷Zr.

47. A composition of embodiment 46, wherein the radionuclide includes ⁹⁰Y, ⁶⁷Cu, ²¹³Bi, ²¹²Bi, ¹⁸⁶Re, ⁶⁷Cu ⁹⁰Y, ²¹³Bi, ¹⁷⁷Lu, ¹⁸⁶Re, and/or ⁶⁷Ga.
48. A composition of embodiment 46, wherein the radionuclide includes ⁸⁹Zr, ²²⁵Ac, and/or ²²⁷Th.
49. A composition of any one of embodiments 1-42, wherein the metal includes a daughter isotope of a radionuclide.

50. A composition of embodiment 49, wherein the daughter isotope of the radionuclide includes ^{89}Y , ^{18}O , ^{221}Fr , ^{213}Bi , and/or ^{209}Pb .

51. A composition, comprising a structure:



wherein:

at least one of R_1 , R_2 , R_3 , R_4 , R_5 , individually, comprise a CAM group, a HA group, or a 1,2-HOPO group;

at least another one of R_1 , R_2 , R_3 , R_4 , and R_5 , individually, comprise H or an alkyl group having from 1 to 10 carbon atoms;

R_6 comprises (i) H, (ii) an alkyl group having from 1 to 10 carbon atoms, or (iii) an alkyl group having from 1 to 100 carbon atoms and no greater than 2 nitrogen atoms, and substituted by at least one of NH_2 , $\text{C}(=\text{O})\text{OH}$, maleimide, dibromo-maleimide, isothiocyanate, alkyne, amide or fluorescent moiety or azide;

m can be from 1 to 6;

n can be from 1 to 6;

o can be from 1 to 6;

p can be from 1 to 6;

q can be from 0 to 6;

r can be from 1 to 6;

s can be from 1 to 6; and

t can be from 1 to 6.

In some embodiments, the chelator 3,4,3-LI(1,2-HOPO) is an octadentate, tetraprotic compound including 4 bidentate 1,2-HOPO metal binding units attached onto a spermine ("3,4,3-LI") scaffold, which was recently modified to enable monoclonal antibody attachment and form a bioconjugate chelator that displayed great properties for positron emission tomography (PET) when bound to ^{89}Zr (Deri, et al., *Bioconjugate Chemistry*, 2015, 26 (12): 2579-2591; Deri, et al., *J. Med. Chem.*, 2014, 57 (11): 4849-4860). Methyl 2,3-dihydroxybenzoate (2). A stirred suspension of 1 (8.06 g, 52.3 mmol) in 100 mL of MeOH was treated with 2.00 mL of concentrated sulfuric acid. The suspension warmed and clarified 2 minutes after the addition. The reaction was equipped with a reflux condenser and was heated to 65°C overnight. The next morning the conversion was verified by LC-MS and the volatiles were removed under reduced pressure. The crude was partitioned between H₂O (100 mL) and ethyl acetate (100 mL) and the aqueous layer was extracted with ethyl acetate (3x50 mL). The organic extracts were combined, dried over MgSO₄, and concentrated under reduced pressure. The crude was passed through a plug of silica using 10% ethyl acetate in hexanes as eluent. The eluent was concentrated under reduced pressure and dried under high vacuum for 2 hours to yield 2 (7.66 g, 45.6 mmol, 88%) as a white solid, the spectral properties of which matched previous reports (Weitl, et al., *J. Am. Chem. Soc.*, 1980, 102 (7): 2289-2293).

Methyl 2,2-diphenylbenzo[d][1,3]dioxole-4-carboxylate (3). Precursor 2 (5.00 g, 29.7 mmol) was mixed with dichlorodiphenylmethane (8.56 mL, 44.6 mol) under an argon atmosphere; the resulting suspension was stirred and heated to 160°C for 1 hour. The mixture was allowed to cool to room temperature and was diluted with 100 mL of ethyl acetate. The solution was washed with sat. NaHCO₃ (30 mL), brine (30 mL), dried over MgSO₄, and then concentrated under reduced pressure. The ensuing greyish oil was dissolved in 30 mL of hot MeOH (65°C) and was slowly cooled to 5°C, which resulted in the formation of white crystals. The crystals were a mixture of 3 and benzophenone that could not be easily separated; the crude product was used as is for the subsequent step.

2,2-diphenylbenzo[d][1,3]dioxole-4-carboxylic acid (4). The mixture from the previous step was dissolved in 100 mL of THF and was treated with 100 mL of 0.9 M LiOH. The emulsion was rapidly stirred and heated to reflux for 5 hours. Conversion was verified by LC-MS and


the reaction was cooled to room temperature. The solution was neutralized with 10% v/v aqueous acetic acid and was extracted with ethyl acetate (3x50 mL). The organic extracts were combined, dried over MgSO_4 , and concentrated under reduced pressure. The crude was chromatographed using 25% ethyl acetate in hexanes as eluent. Volatiles were then removed under reduced pressure followed by high vacuum to yield 4 (7.6 g, 24.06 mmol, 81% over 2 steps) as a white solid, the spectral properties of which matched previous reports (Weitl, et al., *J. Am. Chem. Soc.*, 1980, 102 (7): 2289-2293).

3,4,3-LI(2,2-diphenylbenzo[d][1,3]-2,3-catecholamide) (5). Precursor 4 (746 mg, 2.33 mmol) was suspended in 10 mL of dry toluene under an argon atmosphere and was treated with oxalyl chloride (220 μL , 2.55 mmol). Catalytic *N,N*-dimethylformamide was added and the suspension was heated to 40°C. The solution was stirred until the evolution of gas ceased and was concentrated on the manifold vacuum at the same temperature. The resulting brown oil was dissolved in 10 mL of dry THF. In a separate container a solution of spermine (118 mg, 0.583 mmol), triethylamine (356 μL , 2.56 mmol), and THF (5 mL) was prepared. The solutions were combined and heated to 50°C overnight in a sealed flask. The following day the reaction was filtered and concentrated under reduced pressure. The resulting crude oil was chromatographed using 3% MeOH in CH_2Cl_2 as eluent. The volatiles were then removed under reduced pressure and dried under vacuum, yielding 5 as a white foam (641 mg, 0.457 mmol, 78 % yield). ^1H NMR (300 MHz, CDCl_3) δ 7.88 (1H, t, $J = 5.7$ Hz), 7.66-7.76 (6H, br t), 7.60 (1H, br s), 7.57 (1H, br s), 7.43-7.53 (10H, br s), 7.33-7.40 (4H, br s), 7.19-7.31 (20H, br s), 7.01 (2H, d, $J = 7.6$ Hz), 6.91 (4H, dd, $J = 12.1$ Hz, 8.0 Hz), 6.80 (2H, br s), 6.72 (2H, br s), 3.85 (4H, br s), 3.43 (2H br s), 3.21 (2H, br s), 3.06 (1H, br s), 2.96 (1H, br s), 2.80 (2H, br s), 1.81 (4H, br s), 1.54 (1H, br s), 1.43 (1H, br s), 1.19 (1H, br s), 0.89 (2H, br s). ^{13}C NMR (75 MHz, CDCl_3) δ 167.5, 163.7, 147.3, 147.1, 145.0, 142.8, 139.7, 139.4, 138.9, 129.7, 129.2, 128.4, 128.3, 126.4, 126.3, 126.1, 126.0, 122.3, 122.2, 121.7, 120.4, 118.4, 118.1, 116.0, 111.8, 111.4, 111.3, 109.4, 47.9, 41.8, 36.5, 27.9, 25.5.

3,4,3-LI(CAM) (6). The protected chelator 5 (411 mg, 0.293 mmol) was dissolved in a mixture of 5 mL acetic acid, 0.5 mL H_2O , and 0.1 mL concentrated HCl. The solution was stirred in a sealed container at 60°C overnight. The next day the conversion was confirmed by LC-MS and the volatiles were removed under vacuum. A portion of the crude was

purified using reverse-phase prep-HPLC using at 10→50% MeOH in H₂O + 0.1 % trifluoroacetic acid as eluent. The solvent was removed on a Genevac centrifugal evaporator followed by lyophilization of residual H₂O. 6 was obtained as a pure white powder (90% yield). ¹H NMR (600 MHz, DMSO-d₆) δ 12.82 (1H, br s), 12.69 (1H, br s), 9.52 (2H, br s), 9.11 (2H, br s), 8.78 (1H, br s), 8.60 (3H, br s), 7.26 (1H, br s), 7.12 (1H, br s), 6.90 (2H, br s), 6.77 (1H, br s), 6.66 (4H, br s), 6.56 (2H, br s), 6.44 (1H, br s), 2.88-3.52 (12H, overlapping aliphatic signals), 1.16-1.83 (8H, overlapping aliphatic signals); ¹³C NMR (125 MHz, MeOD-d₄) δ 172.9, 171.5, 150.4, 147.3, 146.6, 125.6, 125.4, 121.0, 119.6, 119.1, 118.8, 118.6, 116.9, 116.6, 47.7, 44.9, 43.2, 37.8, 37.5, 29.3, 28.2, 26.5, 25.5. MS-ESI (m/z) [M + H] Calcd for C₃₈H₄₃N₄O₁₂, 747.2878; found 747.2922 and [M – H] Calcd. for C₃₈H₄₁N₄O₁₂, 745.2721; found 745.2774.

Another pathway for synthesizing a chelator that includes a carboxyl group for binding with another compound, such as a protein or a dye, can include:



Metal, Chelator Solutions. The trivalent lanthanide Ln(III) working stock solutions were prepared in standardized 0.1 M HCl. A Zr(IV) stock solution was prepared by dissolving ZrCl₄ in 3.0 M H₂SO₄, to prevent hydrolysis. The metal salt ZrCl₄ was handled and stored in a glovebox kept under inert atmosphere. The Zr(IV) stock solution was standardized against EDTA, with xylene orange as the indicator (Welcher, F. J. *The analytical uses of ethylenediamine tetraacetic acid*; 1958). A Th(IV) stock solution was prepared in 0.1 M


H₂SO₄. Stock solutions (4 mM) of Ent, and 3,4,3-LI(CAM) were prepared by direct dissolution of a weighed portion of chelator in DMSO and aliquots were removed prior to each set of experiments.

Solution Thermodynamics. All titrant solutions were degassed by boiling for 1 h while being purged under Ar. Carbonate-free 0.1 M KOH was prepared from concentrate (J.T Baker Dilut-It) and was standardized by titrating against 0.1 M potassium hydrogen phthalate (99.95%, Sigma Aldrich). Solutions of 0.1 M HCl were similarly prepared and were standardized by titrating against TRIS (99.9%, J.T. Baker). The glass electrode (Metrohm - Micro Combi - response to [H⁺]) used for the pH measurements was calibrated at 25.0°C and at an ionic strength of 0.1 M (KCl) before each potentiometric or spectrophotometric titration. The calibration data were analyzed using the program GLEE (Gans & O'Sullivan, *Talanta*, 2000, 51 (1): 33-37) to refine for the E° and slope. All thermodynamic measurements were conducted at 25.0°C, in 0.1 M KCl supporting electrolyte under positive Ar gas pressure. The automated titration system was controlled by an 867 pH Module (Metrohm). Two-milliliter Dosino 800 burets (Metrohm) dosed the titrant (0.1 M KOH or 0.1 M HCl) into the thermostated titration vessel (5–90 mL). UV–visible spectra were acquired with an Ocean Optics USB4000-UV–vis spectrometer equipped with a TP-300 dip probe (Ocean Optics; path length of 10 mm), fiber optics and a DH-2000 light source (deuterium and halogen lamps). The fully automated titration system and the UV-vis spectrophotometer were coordinated by LBNL titration system, a computer program developed in house.

Incremental Spectrophotometric Titrations. This method was used to determine the protonation constants of 3,4,3-LI(CAM) as well as the stability constants of its complexes formed with Eu(III), Zr(IV) and ²³²Th(IV). The experimental titration setup is similar to previously described systems (Sturzbecher-Hoehne, et al., *Radiochimica Acta.*, 2013, 101 (6): 359-366). For the 3,4,3-LI(CAM) protonation (and Eu(III)-3,4,3-LI(CAM) complexes), titrations were performed with an initial concentration of 50 μM of 3,4,3-LI(CAM) (and 50 μM of Eu(III)) resulting in absorbance values included between 0 and 1.0 throughout the titration. Typically, 9 mL of a sample containing 3,4,3-LI(CAM) (and Eu(III)) and the supporting electrolyte (KCl/HCl) were incrementally perturbed by addition of 0.025 mL of carbonate-free 0.1 M KOH followed by a time delay of 80 s. Buffering of the solution was

ensured by the addition of 10 mM of HEPES, 10 mM of CHES and 10 mM of MES. Between 130 and 250 data points were collected per titration, each data point including a pH measurement and a UV-Vis spectrum (250-450 nm) over the pH range 1.5 to 12.0. All spectra were corrected for dilution before data fitting. The entire procedure (electrode calibrate, titration and data treatment) was performed independently five times for the protonation constants and four times for the Eu(III)-3,4,3-LI(CAM) complexes. For the Zr(IV) and Th(IV) complexes, titrations were performed similarly but in the presence of DTPA to avoid the formation of metal hydroxides at low pH, before the uptake by 3,4,3-LI(CAM). For each metal, three titrations were performed independently in the presence of 1.1 to 40 equivalents of DTPA. Examples of titrations are displayed in the Supporting Information.

Data Treatment. Thermodynamic constants and spectral deconvolution were refined using the nonlinear least-squares fitting program HypSpec (Gans, et al., *Talanta*, 1996, 43 (10): 1739-1753). All equilibrium constants were defined as cumulative formation constants, β_{mlh} according to Equation (1), where the metal and chelator are designated as M and L, respectively. All metal and chelator concentrations were held at estimated values determined from the volume of standardized stock solutions. All species formed with 3,4,3-LI(CAM) were considered to have significant absorbance to be observed in the UV-vis spectra and were therefore included in the refinement process. The refinements of the overall formation constants β included in each case with previously determined chelator protonation constants and the metal hydrolysis products, whose equilibrium constants were fixed to the literature values (Smith, et al., NIST standard reference database 46. NIST Critically selected stability constants of metal complexes database ver 2004, 2) The speciation diagrams were calculated using the modeling program Hyss (Alderighi, et al., *Coordination Chemistry Reviews*, 1999, 184 (1): 311-318). Errors on $\log \beta_{mlh}$ and pK_a values presented in this Example correspond to the standard deviation observed over the n replicates ($n = 3$ to 5) of the entire procedure (electrode calibrate, titration and data treatment).



3,4,3-LI(CAM) was synthesized from readily available building blocks. The new preparation moves away from using harsh reaction conditions by using the protected diphenylmethylene

acetal derivative (5), which greatly simplifies purification of the final product.

Affinity of 3,4,3-LI(CAM) Toward 3+ and 4+ Metals. A comprehensive solution thermodynamic analysis was performed to characterize the affinity of 3,4,3-LI(CAM) for trivalent and tetravalent metals and the effect of substituting 1,2-HOPO for CAM binding units on the octadentate spermine scaffold. The protonation constants of 3,4,3-LI(CAM) were determined by spectrophotometric titrations, and eight protonation equilibria were assigned to sequential removal of two protons from each of the four CAM units. Previous studies of Ent and other CAM-containing synthetic analogs established that the protonation constants (pK_{a1} - pK_{a4}) of the meta-hydroxyl oxygen atoms are well separated from the ortho-hydroxyl oxygen atoms (pK_{a5} - pK_{a8}) (Loomis & Raymond, *Inorganic Chemistry*, 1991, 30 (5): 906-911). The last four pK_a values are most relevant to metal binding as moieties corresponding to these values have to be deprotonated at physiological pH in order to bind the metal ions. The overall acidity of 3,4,3-LI(CAM) can be defined as $\sum pK_{a5-8} = 45.4$ versus 3,4,3-LI(1,2-HOPO)'s 21.2 (Abergel, et al., *Inorganic chemistry* 2009, 48 (23): 10868-10870) with lower values representing higher acidity. 3,4,3-LI(CAM) is therefore less prone to bind hard Lewis acids at low pH than its 1,2-HOPO analog, due to competition between metal uptake and protonation of the CAM moieties.

Incremental spectrophotometric titrations were then carried out to determine the formation of Eu^{III} , Zr^{IV} or Th^{IV} complexes with 3,4,3-LI(CAM). Because of the very short half-life of ^{225}Ac and the scarce availability of the longer-lived ^{227}Ac , Eu^{III} was used here as a non-radioactive Ln surrogate for Ac^{III} . Based on previous solution thermodynamic studies of Ln^{III} complexes of 3,4,3-LI(1,2-HOPO) and other common polyaminocarboxylate chelators,¹⁵ it is reasonable to expect similar stability constants for Eu^{III} and Ac^{III} complexes of 3,4,3-LI(CAM). The CAM octadentate chelator showed a very high affinity for both 3+ and 4+ ions. The stability constants of $[Eu-3,4,3-LI(CAM)]^{5-}$, $[Th-3,4,3-LI(CAM)]^{4-}$ and $[Zr-3,4,3-LI(CAM)]^{4-}$ are several orders of magnitude higher than those of their 1,2-HOPO counterparts, with log β_{110} values of 29.7, 47.7 and 57.3, respectively. Consequently, 3,4,3-LI(CAM) is one of the strongest chelators ever reported for the chelation of both trivalent and tetravalent *f*-elements. For comparison, a cyclic octadentate terephthalamide derivative was recently designed to bind Th^{4+} *in vivo* and showed an unprecedented affinity for Th^{4+}

with a $\log \beta_{110}$ (ThL4-) value of 53.7 (Pham, et al., *J. Am. Chem. Soc.*, 2014, 136 (25): 9106-9115). To inspect the pH dependency of metal complex formation, speciation diagrams were calculated for 3,4,3-LI(CAM) in the presence of 1 equivalent of Eu(III), Zr(IV) or Th(IV). Both Zr(IV) and Th(IV) complexes start forming at around pH 3, with the mono and fully deprotonated species, $[\text{MIVLH}]^{3-}$ and $[\text{MIVL}]^{4-}$, being predominant at physiological pH (7.4). This behavior departs from that of 3,4,3-LI(1,2-HOPO), with which 4+ metal complexes are formed even under very acidic conditions ($\text{pH} < 0$) (Deblonde, et al., *Inorganic chemistry*, 2013, 52 (15): 8805-8811; Sturzbecher-Hoehne, et al., *Inorganic chemistry*, 2015, 54 (7): 3462-3468). For Eu(III), complexation by 3,4,3-LI(CAM) starts at pH 5 and the mono-protonated complex, $[\text{Eu}^{\text{III}}\text{LH}]^{4-}$, is the only species present at pH 7.4. Similar to what is observed with 4+ metals, the pH at which Eu(III)-3,4,3-LI(CAM) complexes start forming is higher than in the case of Eu(III)-3,4,3-LI(1,2-HOPO) species that already appear at pH 1 under those same conditions (Abergel, et al., *Inorganic chemistry*, 2009, 48 (23): 10868-10870). Additional embodiments of various chelators are shown in FIG. 38A, which depicts an embodiment of a general peptoid synthesis. Submonomer units are generally depicted as "R" group, and can be metal chelating units substituted with an available primary amine. These submonomer units can be used to prepare the peptoids on Rink Amide resin employing standard coupling chemistry. An unlimited number of submonomer units can be added, with each submonomer unit providing 1, 2, 3, or more metal chelating atoms such as oxygen, nitrogen, or sulfur donors. Peptoid structures formed with one, two, three, four, or five submonomer CAM- or HOPO-based units will result in bidentate, tetradentate, hexadentate, Octadentate, or pentadentate ligands. Unless noted otherwise, all synthetic aspects may be carried out in fritted polypropylene syringes, which allowed for recovery of submonomer for re-use. Synthesis can include: 1. Adding 100-150 mg of Rink amide resin to a fritted syringe. Swelling the resin by adding 2 mL of DMF and rock for 30 minutes. Ejecting the solution to isolate the swelled resin. 2.

Adding 1 mL of 20 % 4-methylpiperidine in DMF (v/v) to deprotect the Fmoc group. Agitating for 2 minutes, draining, and repeating for 12 minutes. 3. Rinsing the resin with DMF (2 mL, 5 times for 1 minute). 4. Bromoacetylation. Premixing 0.8 M bromoacetic acid in DMF with 0.8 M N,N-diisopropylcarbodiimide (DIC), 2 mL total solution with 0.4 M of each reagent. Drawing the solution into the syringe, agitating for 5 minutes, and rinsing (2

mL DMF 5x1 minute). 5. Displacement. Drawing in 1.5 mL of submonomer solution (0.2 M in DMF), agitating for 1 hour at 45 °C, then rinsing (DMF 5x1 minute). 6. Repeating bromoacetylation and displacement until synthesis is finished. Washing with DCM (2 mL 3x1 minute) after last DMF wash and drying resin by pulling the plunger out and applying a gentle vacuum onto the syringe needle.

FIG. 38B depicts an embodiment of a general peptoid synthesis, where only one same submonomer unit is used throughout four coupling steps. The submonomer unit comprises a 1,2-HOPO chelating functionality linked to a primary amine through an ethylene bridge to allow for coupling to the peptoid scaffold. The resulting peptoid ligand is octadentate, with four HOPO chelating units available for metal binding. In some embodiments, different units can be added for different functionality, such as to allow cross-linking or other functions.

FIG. 38C depicts an embodiment of a general peptoid synthesis where two different submonomer units are used throughout four coupling steps. The submonomer units comprise either a 1,2-HOPO or CAM chelating functionality linked to a primary amine through an ethylene bridge to allow for coupling to the peptoid scaffold. The combinatorial nature of peptoid synthesis provides a unique opportunity to control binding moiety sequence. The modularity of the scaffold allows for the preparation of up to 16 possible tetrapeptoid, octadentate ligands, with zero, one, two, three, or four HOPO or CAM chelating units available for metal binding. “H” and “C” are defined in FIG. 38D. The option of this approach allows for the generation of a wide variety of resulting chelators.

FIG. 38D depicts an embodiment of a general peptoid synthesis, where two different submonomer units are each used throughout two coupling steps. The submonomer units can comprise a 1,2-HOPO or CAM chelating functionality linked to a primary amine through an ethylene bridge to allow for coupling to the peptoid scaffold. The particular embodiment depicts the “CHHC” peptoid ligand, which incorporates directionally from the rink amide resin a CAM, two HOPO, and a CAM chelating units available for metal binding.

FIG. 38E depicts an embodiment of the 16 products obtained from a general peptoid synthesis, where two different submonomer units are used throughout four coupling steps. The submonomer units comprise a 1,2-HOPO or CAM chelating functionality linked to a

primary amine through an ethylene bridge to allow for coupling to the peptoid scaffold. The modularity of the scaffold allows for the preparation of up to 16 possible tetrapeptoid, octadentate ligands, with zero, one, two, three, or four HOPO or CAM chelating units available for metal binding.

FIG. 38F depicts an embodiment of an organic fluorophore, FITC, that can be used as a fluorescent tag. The tag, similarly to the majority of well-characterized commercially-available fluorophores, can be incorporated onto the peptoid structures through standard amine coupling conditions, using its available primary-amine-reactive isothiocyanate functionality.

FIG. 39A depicts an embodiment of a conjugation pathway for attachment to biologically-relevant targeting molecules, solid resins, or nanoparticles, that contain thiol reactive sites (such as those in cysteine residues). A maleimide or dibromo-maleimide functionality can be attached to the peptoid ligands through standard amide coupling conditions as a subsequent step following the addition of the last chelating submonomer. This embodiment specifically depicts the functionalization of the available peptoid secondary amine by a carboxylic group linked to maleimide or di-bromomaleimide groups through an alkyl chain. In this case, the alkyl chain comprises 5 carbons, but its length can vary from 2 to 10 carbon atoms.

FIG. 39B depicts an embodiment of conjugation pathways that use click chemistry methods for attachment to biologically-relevant targeting molecules, solid resins, or nanoparticles. A functionality comprising of isothiocyanate, azide, or alkyne, can be attached to the peptoid ligands through standard amide coupling conditions as a subsequent step following the addition of the last chelating submonomer. This embodiment specifically depicts the addition of an available peptoid primary amine linked to the scaffold through an alkyl chain and its subsequent reaction to form the isothiocyanate, azide, or alkyne moiety. In this case, the alkyl chain comprises 5 carbons, but its length can vary from 2 to 10 carbon atoms.

FIG. 40A depicts reaction schemes for the conjugation of maleimide- or dibromomaleimide-substituted peptoids to the disulfide bridges available on antibodies. Reduction of the disulfide bridges results in the formation of thiol functionalities on the targeting antibodies, which then readily react with the substituted peptoid.

FIG. 40B depicts the reaction schemes for the conjugation of isothiocyanate- or azide-substituted peptoids to antibody substitutions sites through standard click chemistry methods.

FIG. 40C depicts some embodiments of a fluorescent tag incorporation. A fluorophore can be attached to the peptoid ligands through standard amide coupling conditions as a subsequent step following the addition of the last chelating submonomer. This embodiment specifically depicts the functionalization of the peptoid through additional bromoacetylation and displacement steps with the FITC fluorophore that is substituted with a primary amine through an alkyl chain. In this case, the alkyl chain comprises 2 carbons, but its length can vary from 2 to 10 carbon atoms. Any fluorophore can be used in some embodiments.

EXAMPLES

[0187] The following examples are non-limiting and other exemplary variants as contemplated by one of ordinary skill in the art are acceptable.

Example 1

[0188] HDEHP was 95 % pure and received from Merck and dissolved in kerosene to form an organic stock solution of 0.5 M HDEHP. Sodium nitrate (99 %) was purchased from Alfa Aesar and dissolved in Milli-Q water to form a stock of 5.168 M. Gd(III) chloride hexahydrate was purchased from Aldrich at a purity of 99.999 % and dissolved in 0.1 M nitric acid to form a 5.0 mM stock solution. Nitric acid was purchased from BDH at 70 % concentration. Racemic sodium lactate was purchased from Sigma Aldrich at a purity of 99 %. Stock solutions of DTPA (98 %, Tokyo Chemical Industry) were made by dissolving the solid acid in water and adjusting the pH to 5.4 by 37 % NaOH. 3,4,3-LI(1,2-HOPO) was prepared by Synthetech, Inc using previously reported procedures^{11, 12}. The HOPO monomer, 98 % pure and purchased from Tokyo Chemical Industry, was dissolved in Milli-Q water. One mCi of $^{225}\text{Ac}(\text{NO}_3)_3$ dissolved in 50 μL of 50 mM HCl was received from Oak Ridge National Lab at a purity of 99 %. 326.85 μCi of Gd-153 was received from Oak Ridge National Laboratory and dissolved in 1.0 M HCl. CmCl_3 used for the experiments was 95.78 % ^{248}Cm , 4.12 % ^{246}Cm , 0.06 % ^{245}Cm , 0.02% $^{244}\text{Cm}/^{247}\text{Cm}$ by isotopic distribution.

[0189] All liquid-liquid extractions were conducted using a 1:1 organic to aqueous phase ratio for a total volume of 1 mL. Samples were pH adjusted by HNO_3 and prepared for a final concentration of 1.0 M sodium lactate, 40 mM of hold-back reagent (160 mM in the case of the HOPO monomer), and 2.0 M Na^+ . The concentration of the trivalent f-element was dependent on the availability and radioactivity of the isotopes. The organic phase consisted of 0.5 M HDEHP in kerosene. All samples were incubated in a thermostatic shaker held at 28°C and rocked at 280 rpm for 90 minutes. To ensure complete phase separation, samples were left overnight at room temperature to equilibrate. Radioactive

samples were analyzed by liquid scintillation counting (LSC). The analytical protocol consisted of diluting 10 μ L of radioactive sample into 10 mL of Ultima Gold™. Ultima Gold™ is a commercially available analytical reagent commonly used for LSC. The resulting analytical samples were shaken for 10 min prior to counting on a Perkin-Elmer Packard Tri-Carb instrument (model B4430).

[0190] Data was analyzed by the distribution ratio (D_m), separation factor ($SF_{m1/m2}$) and extraction percentage (% Extraction) as defined by Equations (6), (7) and (8), respectively. Equation (8) uses concentrations of the metal in each phase since the aqueous: organic ratio was 1. Both the organic and aqueous phases were analyzed after liquid-liquid extraction and phase separation. These calculation methods rely on the assumptions that no metal (M) is lost to sorption or precipitation and that no volume variation occurred during the mass transfer between the aqueous and organic phases.

$$D_m = \frac{[M]_{f,org}}{[M]_{f,aq}} = \frac{[M]_{i,aq} - [M]_{f,aq}}{[M]_{f,aq}} \quad (6)$$

$$SF_{m1/m2} = \frac{D_{m1}}{D_{m2}} \quad (7)$$

$$\% \text{ Extraction} = \frac{[M]_{f,org}}{[M]_{i,aq}} = \frac{[M]_{i,aq} - [M]_{f,aq}}{[M]_{i,aq}} = 1 - \frac{[M]_{f,aq}}{[M]_{i,aq}} \quad (8)$$

[0191] The results are shown in FIGs. 11 and 12.

Example 2

[0192] Series of liquid-liquid extraction experiments were first performed with the radionuclide actinium-225. Actinium is the lightest element of the actinide series and its chemistry was expected to be similar to that of americium and curium because the three elements exhibit the same oxidation state in solution: Ac(III), Am(III), and Cm(III) but with ionic radii of 1.26 Å, 1.115 Å, and 1.11 Å,⁷ respectively. The main difference among Ac, Am and Cm is that the only isotopes of actinium available for bulk chemistry (Ac-225 and Ac-227) are short-lived isotopes ($t_{1/2}$ = 9.95 days and 21.8 years, respectively) whereas long-

lived isotopes of americium and curium are available for research purposes ($t_{1/2} = 7,388$ years for Am-243 and 3.5×10^5 years for Cm-248). The practical consequence is that samples of Ac-225 are far more radioactive than the corresponding Am-243 and Cm-248 samples. So far, no data have been published on the behavior of actinium in the TALSPEAK process.

[0193] The behavior of Ac(III) was probed during liquid-liquid extraction conditions similar to that of TALSPEAK and compare it to Am(III) and Gd(III). Gd(III) was used as a surrogate for Eu(III) and was expected to behave similarly based on the similarity in ionic radii of 1.08 Å and 1.09 Å for Gd(III) and Eu(III), respectively.⁷ Graphs summarizing the extraction of Ac, Am, and Gd are shown in FIG. 8 and FIG. 26 for DTPA and 3,4,3-LI(1,2-HOPO), respectively.

[0194] FIG. 8 shows the extraction profile of Ac(III)-225, Am(III)-243, and Gd(III) (with [Gd(III)-153] = 0.21 nM) with DTPA as the hold-back reagent. [Ac(III)-225] = 0.4 nM, [Gd(III)] = 500 μM, [Am(III)-243] = 40.4 nM. Selective extraction of Gd(III) at pH 3-4 was observed, just as expected for this system under TALSPEAK conditions. FIG. 26 shows extraction profile of Ac(III)-225, Am(III)-243, and Gd(III) (with [Gd(III)-153] = 0.21 nM) with 3,4,3-LI(1,2-HOPO) as the hold-back reagent. [Ac(III)-225] = 0.4 nM, [Gd(III)Total] = 500 μM, [Am(III)-243] = 40.4 nM. In this case, evidence for the selective extraction of Ac(III) over Gd(III) and Am(III)-243 was seen.

[0195] The separation factors of Ac(III)/M(III) (M = Am, Gd) are compared for DTPA and 3,4,3-LI(1,2-HOPO) in TABLE 2. A comparison of the behavior between Am and Gd for 3,4,3-LI(1,2-HOPO) is provided in FIG. 19, which shows the extraction profile for Gd(III) and Am(III) along with the separation factor are plotted at various pH. Selective extraction for Gd(III) was observed at pH 1-2.5.

Example 3

[0196] FIG. 18 shows that, in the presence of 3,4,3-LI(1,2-HOPO), the extraction capacity of HDEHP is totally suppressed and that no Pu^{4+} is transferred into the organic phase. Literally no activity was detected in the organic phase after extraction meaning that the $[\text{Pu(IV)3,4,3-LI(1,2-HOPO)}]$ is too stable to be challenged by HDEHP. FIG. 18 shows percentage of extraction of plutonium(IV) as a function of pH, i.e., extraction yields obtained for tetravalent plutonium ions from nitric media in the presence of 3,4,3-LI(1,2-HOPO). The

extractant used in this example was HDEHP since this commercially available molecule is very common in the nuclear chemistry field and in hydrometallurgy processes. *See*, K.L. Nash, The Chemistry of TALSPEAK: A Review of the Science, Solvent Extr. Ion Exch. 33 (2015) 1–55. doi:10.1080/07366299.2014.985912. Similar experiments were also performed with DTPA instead of 3,4,3-LI(1,2-HOPO) since DTPA is often taken as a reference chelator for actinide ions. Note that the plutonium(IV) case was studied here because it is a good chemical surrogate for thorium(IV) and because of the availability of Pu-242 in our laboratory. Pu-242 ($t_{1/2} = 3.7 \times 10^5$ y) was also more convenient to analyze by liquid scintillation counting than the natural and long-lived isotope of thorium (Th-232; $t_{1/2} = 1.4 \times 10^{10}$ y).

[0197] A value of 0.03% was plotted on FIG. 18, which corresponds to our instrumental detection limit under the tested conditions, but this extraction yield has to be taken as an upper limit for Pu⁴⁺.

[0198] In contrast, when DTPA is present in the aqueous phase, Pu⁴⁺ ions are extracted with an increase in the extraction yield from 0.30 to 95.2% when the pH decreases from 4.2 to 0.4 due to the release of Pu⁴⁺ by DTPA (FIG. 18). These results clearly show that DTPA cannot be used to hold-back Pu⁴⁺ in the aqueous phase.

[0199] In the case of 3,4,3-LI(1,2-HOPO), the ligand affords a total sequestration of Pu⁴⁺ in the aqueous phase. Based on the solution thermodynamics data (FIG. 20), a similar behavior is expected in the case of the other tetravalent ions such as Zr⁴⁺, Ce⁴⁺, and Th⁴⁺.

[0200] In FIG. 23, aqueous phase: [ligand] = 40 mM, buffer sodium lactate/sodium nitrate. Organic phase: [HDEHP] = 0.5 M, kerosene diluent. T = 25 °C. O/A = 1, one contact. Initial metal concentrations: [Ac-225] = 0.4 nM, [Am-243] = 40.4 nM, or [Gd] = 500 μM (spiked with Gd-153 for LSC measurements). For actinium, data acquired in the absence or presence of 500 μM of Eu(III) did not show any significant influence on the extraction yields.


[0201] FIG. 23 shows percentage of extraction as a function of pH for trivalent actinium, americium and, gadolinium ions in the presence of 3,4,3-LI(1,2-HOPO), i.e., extraction yields obtained for the trivalent ions Ac³⁺, Gd³⁺ and, Am³⁺ using 3,4,3-LI(1,2-HOPO) and chemical conditions similar to those described above for Pu⁴⁺. The equation for release from the ligand is given by Equation (4) main phase transfer equation for trivalent

ions and HDEHP is given by Equation (5). *See*, K.L. Nash, The Chemistry of TALSPEAK: A Review of the Science, Solvent Extr. Ion Exch. 33 (2015) 1–55. doi:10.1080/07366299.2014.985912. At pH below 1, low extraction yields are obtained due the protonation of HDEHP which competes with the extraction of the trivalent ions. *See*, K.L. Nash, The Chemistry of TALSPEAK: A Review of the Science, Solvent Extr. Ion Exch. 33 (2015) 1–55. doi:10.1080/07366299.2014.985912; K.V. Lohithakshan, P. Patil, S.K. Aggarwal, Solvent extraction studies of plutonium(IV) and americium(III) in room temperature ionic liquid (RTIL) by di-2-ethyl hexyl phosphoric acid (HDEHP) as extractant, J. Radioanal. Nucl. Chem. 301 (2014) 153–157. doi:10.1007/s10967-014-3119-9. This behavior is specific to the extractant HDEHP and is not a limitation to the present invention. Other extractants that work at low pH values, such as the tertiaryalkyl amines or alkylphosphates, could be used in lieu of HDEHP.

[0202] Equation (4) (release from the ligand):



[0203] Equation (5) (extraction):



[0204] When 3,4,3-LI(1,2-HOPO) is present and for a pH higher than 1, Am³⁺ and Gd³⁺ ions are retained in the aqueous phase. This behavior was expected from the solution thermodynamic data in FIG. 20. A slight extraction of Gd³⁺ ions is observed in the pH range 1-2 due to the release of Gd³⁺ ions by 3,4,3-LI(1,2-HOPO). No extraction is observed for Am³⁺ which is in line with the higher stability of the Am³⁺ complex (*See*, M. Sturzbecher-Hoehne, P. Yang, A. D'Aléo, R.J. Abergel, Intramolecular sensitization of americium luminescence in solution: shining light on short-lived forbidden 5f transitions, Dalton Trans. (2016). doi:10.1039/C6DT00328A) when compared to the Gd³⁺ complex (*See*, M. Sturzbecher-Hoehne, C. Ng Pak Leung, A. D'Aléo, B. Kullgren, A.-L. Prigent, D.K. Shuh, K.N. Raymond, R.J. Abergel, 3,4,3-LI(1,2-HOPO): In vitro formation of highly stable lanthanide complexes translates into efficacious in vivo europium decorporation, Dalton Trans. 40 (2011) 8340. doi:10.1039/c1dt10840a). In the case of actinium, a quantitative extraction is observed at pH higher than 1.5 suggesting that the 3,4,3-LI(1,2-HOPO) complex

of Ac^{3+} is not stable enough to compete with the extraction by HDEHP (Equation (5)). This distinct behavior for Ac^{3+} represents a powerful leverage for separation methods.

[0205] Combining the results in FIG. 18 and FIG. 23, it is clear that actinium can be selectively extracted from a mixture of trivalent lanthanide, trivalent actinide and, tetravalent ions when using 3,4,3-LI(1,2-HOPO) as a hold-back reagent. A one-step process would afford a direct and very selective extraction of actinium.

Example 4

[0206] The specificity of the 3,4,3-LI(1,2-HOPO) chelator for actinium was demonstrated. Similar experiments were performed for the DTPA ligand and the trivalent metal ions. FIG. 16 shows that DTPA is not strong enough to scavenge Ac^{3+} , Gd^{3+} or Am^{3+} ions in the aqueous phase. In FIG. 16 aqueous phase: [ligand] = 40 mM, buffer sodium lactate/sodium nitrate. Organic phase: [HDEHP] = 0.5 M, kerosene diluent. T = 25 °C. O/A = 1, one contact. Initial metal concentrations: [Ac-225] = 0.4 nM, [Am-243] = 40.4 nM, or [Gd] = 500 μM (spiked with Gd-153 for LSC measurements). For actinium, data was acquired in the absence or presence of 500 μM of Eu(III) did not show any significant influence on the extraction yields. Therefore, actinium cannot be purified from a mixture of Ac^{3+} , Ln^{3+} and An^{3+} when using DTPA as a hold-back reagent because the metals would be partially extracted at the same time. The extraction yields for Ac^{3+} are also not quantitative (under the conditions tested) and would lead to a partial recovery of Ac isotopes. Chelators of the DTPA family (*i.e.* carboxylic acids) that are commonly used in separation processes, such as EDTA, NTA and, DOTA, are likely to exhibit a similar non-selectivity toward actinium. For comparison purposes, the separation factors that would be obtained with the method proposed here are plotted in FIG. 17.

[0207] In FIG. 17, aqueous phase: [ligand] = 40 mM, buffer sodium lactate/sodium nitrate, organic phase: [HDEHP] = 0.5 M, kerosene diluent. T = 25 °C. O/A = 1, one contact. Initial metal concentrations: [Ac-225] = 0.4 nM, [Am-243] = 40.4 nM, or [Gd] = 500 μM (spiked with Gd-153 for LSC measurements). Separation factor = ratio M_1/M_2 in the final organic phase / ratio M_1/M_2 in the initial aqueous phase.

[0208] In FIG. 18, aqueous phase: [ligand] = 40 mM, buffer sodium lactate/sodium nitrate, organic phase: [HDEHP] = 0.5 M, kerosene diluent. T = 25 °C. O/A =

1, one contact. Initial metal concentrations: [Pu-242] = 5 μ M. Phases analyzed after one contact. Contact time: 30 min. Limit of detection: 0.03 % of the initial Pu-242 activity. Data for separation of tetravalent metal ions from trivalent metal ions using 3,4,3-LI(1,2-HOPO) as a hold-back reagent for tetravalent metal ions are shown in TABLE 0.1.

TABLE 0.1
Percentage of extraction of plutonium(IV) by HDEHP as a function of pH
and using either diethylenetriaminepentaacetic acid (DTPA) or
3,4,3-LI(1,2-HOPO) as hold-back reagent

Hold-back reagent	pH	Activity in organic phase (cpm)	Activity in aqueous phase (cpm)	%Extraction
DTPA	0.40	4181	209	95.24%
	1.27	110	2585	4.08%
	1.84	35	2951	1.17%
	2.55	19	2995	0.63%
	2.75	23	3121	0.73%
	3.45	22	3379	0.65%
	4.29	10	3491	0.29%
3,4,3-LI(1,2-HOPO)	0.25	0	2791	0.00%
	0.85	0	3325	0.00%
	1.78	0	3310	0.00%
	2.71	0	3219	0.00%
	2.87	0	3339	0.00%
	3.62	0	3185	0.00%
	4.52	0	2896	0.00%


[0209] Very high separation factors could be obtained between actinium and the lanthanides or actinides, after only 1 extraction step. The separation factors Ac^{3+}/Pu^{4+} were not calculated since the extraction yield of Pu^{4+} is null (FIG. 18) in the presence of 3,4,3-LI(1,2-HOPO), meaning a virtually infinite separation factor.

[0210] Thus, in some embodiments, the proposed method combines two essential features: (i) quantitative extraction for actinium, and (ii) a very high decontamination capability.

[0211] For DTPA extractions, the expected behavior of the system with selective extraction of Gd(III) at pH 3-4 under TALSPEAK conditions was observed. Below pH ~1.5 (FIG. 21), retention of Ac(III) and Am(III) in the aqueous phase was observed and corresponding to Equation (2), where the equilibrium lies to the left. Protonation of the extractant in the organic phase prevents metal complexation and does not necessarily indicate DTPA binding. In the speciation diagram in FIG. 8, free metal or nitrate complexes are expected to be present below pH 1. As pH increases in the extraction samples, the selectivity of DTPA begins to emerge with preferential binding to the Ac(III) and Am(III), while Gd(III) is extracted by HDEHP. Additionally, a high pH dependence of the process in the 2.5-3.5 pH range was observed and this can be attributed to the shift from the protonated metal complex to the proton independent complex, as modeled in FIG. 8.

[0212] Extraction of Gd(III) was above 90 % until after pH 4, while for Ac(III) (49 %) and Am(III) (78 %) maximum metal extraction is around pH 1.5 and decreases to less than 10 % at pH 4. Above pH 3.5, a decrease in extraction of Gd(III) is observed, hence the decrease in selectivity for the current TALSPEAK model.

[0213] Metals with larger ionic radii was expected to form weaker complexes relative to smaller metals. Therefore, higher retention of Am(III) (1.115 Å) relative to Ac(III) (1.26 Å) in the aqueous phase. However, the reverse was observed.



[0214] Based on the speciation diagram for 3,4,3-LI(1,2-HOPO) in FIG. 26, metal complexation was expected to persist even at low pH. However, the complete extraction of Ac(III) above pH 1.5 (FIG. 28) was unexpected and indicated relatively weak binding of Ac(III) by the HOPO chelator compared to DTPA.

[0215] Without being bound any particular theory, upon examining the structures of DTPA and 3,4,3-LI(1,2-HOPO), an argument can be made for the relative cavity limitations when binding metals. Each of the chelating moieties in 3,4,3-LI(1,2-HOPO) is bidentate and the ligand scaffold is not directly involved in metal binding. On the other hand, DTPA chelates through the three nitrogen atoms in the diethylenetriamine backbone and the

five carboxylic acid groups, potentially giving it more flexibility in the size of metal it can chelate. 3,4,3-LI(1,2-HOPO) may be limited by the length of the scaffold and chelate a metal as large as Ac(III) less efficiently than a more flexible ligand like DTPA.

[0216] In FIG. 28, a striking contrast in the behavior of Ac(III) and Gd(III)/Am(III) was observed, where complete retention of Gd(III) and Am(III) is seen above pH 2.5. TABLE 2 helps accentuate the difference in the selective extraction of Ac(III) when using 3,4,3-LI(1,2-HOPO) and DTPA.

TABLE 2
Comparison of the separation factors for Ac(III)/M(III)
(M = Am, Gd) for 3,4,3-LI(1,2-HOPO) and DTPA at various pH.

$SF_{Ac/Gd}$		
pH*	HOPO	DTPA
0.8	$1.23 \cdot 10^0$	$< 10^{-3}$
1.5	$6.55 \cdot 10^2$	$< 10^{-3}$
3.5	$4.61 \cdot 10^5$	$< 10^{-3}$
4.0	$1.33 \cdot 10^6$	$< 10^{-3}$
$SF_{Ac/Am}$		
pH*	HOPO	DTPA
0.9	$3.99 \cdot 10^{-2}$	$5.83 \cdot 10^{-2}$
1.5	$9.06 \cdot 10^3$	$2.70 \cdot 10^{-1}$
3.5	$8.13 \cdot 10^4$	$6.86 \cdot 10^{-2}$
4.0	$1.28 \cdot 10^5$	$1.11 \cdot 10^{-1}$


[0217] The data in TABLE 2 show that Ac extraction is more selective when using 3,4,3-LI(1,2-HOPO) and the selectivity increases with increasing pH. pH values are ± 0.1 .

[0218] In some embodiments, DTPA conferred no selectivity between Ac and Gd(III)/Am(III), whereas with 3,4,3-LI(1,2-HOPO) selective extraction of Ac(III) was observed.

[0219] As previously mentioned, 3,4,3-LI(1,2-HOPO) was expected to form complexes at lower pH and a DTPA-like extraction profile shifted to lower pH regions was expected. Low extraction of Gd(III) and Am(III) (FIG. 19) was observed even at pH below

1, and slight increase at pH slightly above 1, where the latter corresponds to the deprotonation of HDEHP and extraction of the metal as in Equation (2). FIG. 19 shows selective extraction for Gd(III) is observed at pH 1-2.5.

[0220] As in the DTPA system, higher affinity to metals at higher pH was observed and reached almost complete extraction above pH 2.5. The equilibria of the process are described by Equation (3), where the protonated complex is the primary species in solution. Gd(III) selective extraction over Am(III) is seen at pH 1-2.5 and albeit a small difference under the current conditions it serves as evidence for selective extraction of Gd(III).



Example 5

[0221] FIG. 26 and FIG. 27 show data related to separation of tetravalent metal ions from trivalent metal ions using a HOPO chelator as a hold-back reagent for tetravalent metal ions.

[0222] FIG. 26 shows percentage of extraction of plutonium(IV), americium(III), and gadolinium(III) by HDEHP at constant pH and using 3,4,3-LI(1,2-HOPO) as hold-back reagent. In FIG. 26, aqueous phase: [ligand] = 1 mM, buffer 0.1 M HNO₃/1.9 M NaNO₃. Organic phase: HDEHP diluted in kerosene. [HDEHP] = 0.05, 0.10, 0.25, 0.50, 1.0, or 2.0 M. T = 25 °C. Volume organic phase/Volume aqueous phase = 1. Initial metal concentrations: [Pu-242] = 10 µM, [Am-243] = 0.25 µM, or [Gd-153] = 0.16 nM. Phases analyzed after one contact. Contact time: 30 min.

[0223] FIG. 27 shows corresponding separation factors: SF M₁/M₂ = Ratio [M₁]/[M₂] in the final organic phase / Ratio [M₁]/[M₂] in the initial aqueous solution. The separation factor values given here must be seen as lower limit since the extraction of Pu(IV) is near zero. Data for FIG. 26 and FIG. 27 are shown in TABLE 0.2

[0224] FIG. 28 shows data related to separation of trivalent actinides from trivalent lanthanides using 3,4,3-LI(1,2-HOPO) in the aqueous phase and a non-selective

extractant in the organic phase. FIG. 28 shows percentage of extraction of americium(III), and gadolinium(III) by HDEHP at constant pH and using 3,4,3-LI(1,2-HOPO) as hold-back reagent. In FIG. 28, aqueous phase: [ligand] = 1 mM, buffer 0.1 M HNO₃/1.9 M NaNO₃. Organic phase: HDEHP diluted in kerosene. [HDEHP] = 0.05, 0.10, 0.25, 0.50, 1.0, or 2.0 M. T = 25 °C. Volume organic phase/Volume aqueous phase = 1. Initial metal concentrations: [Am-243] = 0.25 μM, or [Gd-153] = 0.16 nM. Phases analyzed after one contact. Contact time: 30 min. Corresponding separation factors (SF) are given on the right Y-axis. $SF_{Gd/Am} = \text{Ratio [Gd]/[Am] in the final organic phase} / \text{Ratio [Gd]/[Am] in the initial aqueous solution}$. Data for FIG. 28 are shown in TABLE 0.2

TABLE 0.2

	Concentrat ion HDEHP	Activity in organic phase (cpm)	Activity in aqueous phase (cpm)	%Extracti on	SF Gd ³⁺ /A m ³⁺	
²⁴³ Americium(III)	0.05	5	2326	0.21%	24.1	
	0.10	30	2237	1.32%	23.3	
	0.25	246	2195	10.08%	25.3	
	0.50	735	1693	30.27%	27.0	
	1.00	1296	1297	49.98%	45.6	
	2.00	1277	952	57.29%	99.8	SF Gd ³⁺ /Pu 4+
¹⁵³ Gadolinium(III)	0.05	2505	48361	4.92%		331
	0.10	12170	38983	23.79%		1948
	0.25	40716	14376	73.91%		5976
	0.50	51171	4362	92.15%		39182
	1.00	55033	1207	97.85%		28460
	2.00	56084	419	99.26%		64723
²⁴³ Plutonium(IV)	0.05	1	6382	0.02%		14
	0.10	1	6241	0.02%		84
	0.25	3	6330	0.05%		236
	0.50	2	6680	0.03%		1450
	1.00	10	6242	0.16%		624
	2.00	13	6286	0.21%		649
						SF Am ³⁺ /P u ⁴⁺

*SF = separation factor; cpm = count per minute

Example 6

[0225] Metal ions can be separated in an extraction. One first provides an aqueous solution that comprises a first metal ion (M^{4+}) and a second metal ion (M^{3+}). The solution is at a pH of between 0 and 1. One then adds a HOPO chelator (3,4,3-LI(1,2-HOPO)) to the aqueous solution. One then performs an extraction against the aqueous solution using an organic solution, wherein the HOPO chelator functions as a holdback agent to selectively keep the first metal ion in the aqueous solution. This allows the second metal

ion to go into an organic phase during the extraction. One then decants the organic solution, keeping the first metal ion in the aqueous phase. One can then optionally release the first metal ion from the chelator by further lowering the pH. Optionally, a non-specific extractant can be used to assist in moving the M^{3+} metal ions to the organic phase.

Example 7

[0226] Metal ions can be separated in an extraction. One first provides an aqueous solution that comprises a first metal ion (M^{3+}) and a second metal ion (M^{3+}). The solution is at a pH of between 0 and 1. One then adds a HOPO chelator (3,4,3-LI(1,2-HOPO)) to the aqueous solution. The first metal ion has a stronger affinity for the HOPO chelator than the second metal ion does. One then performs an extraction against the aqueous solution using an organic solution, wherein the HOPO chelator functions as a holdback agent to selectively keep the first metal ion in the aqueous solution. This allows for more of the second metal ion to go into an organic phase during the extraction. One then decants the organic solution, keeping the first metal ion in the aqueous phase. One can then optionally release the first metal ion from the chelator by further lowering the pH. Optionally, a non-specific extractant can be used to assist in moving the M^{3+} metal ions to the organic phase.

Example 8

[0227] A method of processing spent nuclear fuel is provided. It involves obtaining spent nuclear fuel that comprises Pu^{4+} and dissolving the metal in an acidic medium at a pH of less than 2. One then contacts the dissolved spent nuclear fuel with a spermine-based octadentate ligand and an organic phase to generate a mixture (and agitating the mixture such that the organic and aqueous phases mix). One then separates an aqueous phase enriched for Pu^{4+} from the mixture by decanting. The organic phase will contain various M^{3+} and/or M^{2+} metal ions. Optionally, a non-specific extractant can be used to assist in moving the M^{3+}/M^{2+} metal ions to the organic phase.

Example 8

[0228] A method for enriching a metal ion is provided. One contacts a first aqueous phase with an organic phase to generate a mixture (e.g., during an extraction). The first aqueous phase comprises a plurality of metal ions (both M^{3+} and M^{4+}) and a spermine-based octadentate ligand (e.g., a HOPO chelator). The aqueous phase has an acidic pH of less than 1. One then separates from the mixture a second aqueous phase. The second aqueous phase is only different from the first in that it is, relatively enriched (or has a greater purity) for a M^{4+} metal ion of the plurality of metal ions. The organic phase will be enriched for the M^{3+} metal ions. Optionally, a non-specific extractant can be used to assist in moving the M^{3+} metal ions to the organic phase.

Example 9

[0229] A method of preparing a medical isotope is provided. The method involves obtaining a metallic precursor dissolved under an acidic condition. The metallic precursor comprises a medical isotope (M^{4+} or M^{3+}), while the precursor is a mixture of various metal ions (M^{4+} and M^{3+} or different M^{3+}). One then contacts the dissolved metallic precursor with a spermine-based octadentate ligand (e.g., 3,4,3-LI(1,2-HOPO)) and an organic phase to generate a mixture. One then separates the medical isotope from the one or more metal ions in the dissolved metallic precursor based on an interaction between the octadentate ligand and the medical isotope. The metal that has a higher affinity for the HOPO ligand will remain in the aqueous phase while the metal with a lower affinity (that is not chelated) can be moved to the organic phase via a non-specific extractant. The desired medical isotope will then be in one or the other of the aqueous phase or the organic phase.

Example 10

[0230] A method of separating metal ions for nuclear forensics is provided. The method includes obtaining a sample derived from a nuclear material, wherein the sample comprises UO_2^{2+} , Pu^{4+} , and Np^{4+} . One then contacts the sample with a spermine-based octadentate ligand (e.g., 3,4,3-LI(1,2-HOPO)) to generate a first mixture. The first mixture has an acidic pH of 1 or lower. One then separates UO_2^{2+} from Pu^{4+} and Np^{4+} in the mixture based on an interaction between the 3,4,3-LI(1,2-HOPO) and Pu^{4+} , and Np^{4+} . A non-specific extractant can be used to assist in moving the UO_2^{2+} to the organic phase. This results in

altering the first mixture to form a second mixture comprising the Pu⁴⁺ and Np⁴⁺ (but it is still the aqueous phase). One can then chromatographically separate Pu⁴⁺ from Np⁴⁺ in the second mixture.

Example 11

[0231] A method of separating metal ions is provided. The method involves contacting a liquid composition comprising a plurality of metal ions with a spermine-based octadentate ligand (e.g., 3,4,3-LI(1,2-HOPO)) to generate a mixture, under conditions sufficient to form a metal ion-ligand complex comprising a metal ion of the plurality of metal ions. One then separates a first fraction of the mixture enriched for the metal ion-ligand complex from a second fraction depleted for the metal ion-ligand complex. The first fraction of the mixture has an acidic pH of less than 1 and will include the 3,4,3-LI(1,2-HOPO) and any metal chelated thereto. A non-specific extractant can be used to assist in moving any unchelated metal ions to the organic phase. The plurality of metal ions are selected from the group consisting of: a p-, d- or f-block element of period 5 or greater, a group 3 element, or a group 4 element.


Example 12

Experimental Procedures


General Considerations. Chemicals were obtained from commercial suppliers and were used as received unless stated otherwise. ¹H NMR spectra were recorded on Bruker instruments; ¹³C NMR spectra were recorded on Bruker instruments with tetramethylsilane as an internal reference. SilicaFlash G60 (particle size 60-200 μm) was used for flash column chromatography. LC-MS was performed on an Agilent LC/MS system consisting of an Agilent 1200 binary LC pump, a temperature-controlled autosampler, a PDA UV detector, and a 6530 Accurate Mass Q-TOF mass spectrometer (Wilmington, DE, USA). The mass spectrometer was equipped with a JetStream® ESI probe operating at atmospheric pressure. The ESI source parameter settings were: mass range m/z 100–1200, gas temperature 350 °C, gas flow 10 L/min, nebulizer 50 psi, sheath gas temperature 400 °C, sheath gas flow 12 L/min, capillary voltage (Vcap) 3500 V, nozzle voltage 500 V, fragmentor 200 V, skimmer 65 V, octopole RF (OCT 1 RF Vpp) 750 V. Reverse phase preparatory HPLC was performed on

a Varian ProStar system with a Vydac C18 column. HRMS and MS-MS were obtained on a Waters Xevo G2 Qtof mass spectrometer, leucine encephalin lockspray with mass correction was used for HRMS.

Example 13--Synthetic Procedures--Submonomer




Methyl 2,3-dihydroxybenzoate, 4. A stirred suspension of **3** (8.06 g, 52.3 mmol) in 100ml of methanol was treated with 2.00 ml of concentrated sulfuric acid. The suspension warmed and clarified 2 minutes after the addition. The reaction was equipped with a reflux condenser and was heated to 65 °C overnight. The next morning the conversion was verified by LC-MS and the volatiles were removed under reduced pressure. The crude was partitioned between water (100 ml) and ethyl acetate (100 ml) and the aqueous layer was extracted with ethyl acetate (3x50 ml). The organic extracts were combined, dried over MgSO₄, and concentrated under reduced pressure. The crude was passed through a plug of silica using 10% ethyl acetate in hexanes as eluent. The eluent was concentrated under reduced pressure and dried under high vacuum for 2 hours to yield **4** (7.66g, 45.6 mmol, 88%) as a white solid, the spectral properties of which matched previous reports.¹



Methyl 2,2-diphenylbenzo[d][1,3]dioxole-4-carboxylate, 5. **4** (5.00g, 29.7 mmol) was mixed with dichlorodiphenylmethane (8.56 ml, 44.6 mmol) under an argon atmosphere, the


resulting suspension was stirred and heated to 160°C for 1 hour. The mixture was allowed to cool to room temperature and was diluted with 100 ml of ethyl acetate. The solution was washed with sat. NaHCO₃ (30 mL) then brine (30 mL), dried over MgSO₄, then concentrated under reduced pressure. The ensuing greyish oil was dissolved in 30 mL of hot methanol (65 °C) and was slowly cooled to 5 °C, which resulted in the formation of white crystals. The crystals were a mixture of **5** and benzophenone dimethyl acetal that could not be easily separated; product was used as is for the subsequent step.

2,2-diphenylbenzo[d][1,3]dioxole-4-carboxylic acid, 6. The mixture from the previous step was dissolved in 100 mL of THF and was treated with 100 mL of 0.9 M LiOH. The emulsion was rapidly stirred and heated to reflux for 5 hours. Conversion was verified by LC-MS and the reaction was cooled to room temperature. The solution was neutralized with 10 % v/v aqueous acetic acid and was extracted with ethyl acetate (3x50 mL). The organic extracts were combined, dried over MgSO₄, and concentrated under reduced pressure. The crude was chromatographed using 25% ethyl acetate in hexanes as eluent. Volatiles were then removed under reduced pressure followed by high vacuum to yield **6** (7.6 g, 24 mmol, 81% over 2 steps) as a white solid, the spectral properties of which matched previous reports.¹



“CAM Submonomer, 2.” **1** (3.84 g, 12.1 mmol) was suspended in 30 mL of toluene under an argon atmosphere. Oxalyl chloride (1.14 mL, 13.3 mmol) was then added followed by a catalytic amount of N,N-dimethylformamide. The suspension was heated to 40 °C and was stirred until it became clear and the evolution of gas has ceased (~1 hour). The volatiles were then removed under reduced pressure and the resulting white solid was dissolved in dry dichloromethane. A separate 1 L roundbottom flask outfitted with an addition funnel was charged with ethylenediamine (8 mL, 120 mmol) and 50 mL dry dichloromethane; the resulting solution was cooled to 0 °C using an ice bath. The aforementioned solution of acyl

chloride was transferred into the addition funnel and was diluted with dichloromethane to a total volume of 700 mL. The acyl chloride solution was then added into the vigorously stirred ethylenediamine over 1.5 hours at 0 °C. Following the addition, the reaction solution was transferred into a separatory funnel and was washed with 0.5 M NaOH in 50% saturated aq. NaCl (50 mL x2). The organic phase was dried over MgSO₄ and was concentrated on a rotary evaporator yielding the crude. The crude was purified using silica column chromatography (5→10% MeOH in DCM with 1% Et₃N, R_f = 0.35 in 10% MeOH in DCM). The desired fractions were combined, concentrated under reduced pressure, and dried under vacuum yielding the CAM submonomer as a sticky yellow oil (3.49 g, 9.68 mmol, 80% yield). ¹H NMR (300 MHz, CDCl₃) δ 7.64 (1H, br s, NH), 7.56-7.61 (5H, m, ArH), 7.37-7.42 (6H, m, ArH), 7.01 (1H, dd, *J* = 7.7, 1.4 Hz, ArH), 6.94 (1H, t, *J* = 7.9 Hz, ArH), 3.56 (2H, q, *J* = 6.0 Hz, NHCH₂), 2.97 (2H, t, *J* = 6.0 Hz, NH₂CH₂), 2.75 (2H, s, NH₂). ¹³C NMR (75 MHz, CDCl₃) δ 163.7, 147.2, 144.7, 139.4, 129.5, 128.4, 126.2, 122.4, 122.0, 118.0, 116.0, 111.6, 42.3, 41.5. HRMS-ESI (*m/z*) [M+H]⁺ Calcd. For C₂₂H₂₀N₂O₃+H, 361.1563; found, 361.1581.



“HOPO Submonomer, 3.” **3** was synthesized in an identical fashion to **2** above, using 1-benzyloxy-6-carboxy-2 (1H) pyridinone (*J. Labelled Cpd. Radiopharm.* **2001**, *44*, 13-19, CAS 210366-15-7, US Patent US6846915) as the starting material. The crude product was purified using silica column chromatography (10% NH₄OH (10%) in MeOH) in DCM, R_f = 0.08). The desired fractions were combined, concentrated under reduced pressure, and dried under vacuum yielding the HOPO submonomer **3** as a sticky yellow oil in ~80% yield). ¹H NMR (300 MHz, CDCl₃) δ 7.40-7.48 (3H, m, ArH), 7.35-7.38 (3H, m, ArH and NH), 7.24-7.29 (1H, m, ArH), 6.66 (1H, dd, *J* = 9.0, 1.5 Hz, CHCH₂CH), 6.45 (1H, dd, *J* = 6.9, 1.8 Hz,

CHCHCH), 5.29 (2H, s, CH₂Ph), 3.36 (2H, q, $J = 6.0$ Hz, NHCH₂), 2.80 (2H, t, $J = 6.0$ Hz, NH₂CH₂). ¹³C NMR (75 MHz, CDCl₃) δ 160.3, 158.5, 142.5, 138.0, 133.2, 130.1, 129.4, 128.6, 124.0, 106.4, 79.3, 77.4, 77.0, 76.6, 42.7, 40.7. Calcd. For C₁₅H₁₇N₃O₃+H, 288.1328; found, 288.1361.

Example 14--Synthesis of Peptoids

Unless noted otherwise, all steps were carried out in fritted polypropylene syringes, which allowed for recovery of submonomer for re-use. Automatic peptoid synthesis was not an option for this work due to difficulty of sub-monomer preparation.

1. Add 100-150 mg of Rink amide resin to a fritted syringe. Swell the resin by adding 2 mL of DMF and rock for 30 minutes. Eject the solution to isolate the swelled resin.
2. Add 1 mL of 20 % 4-methylpiperidine in DMF (v/v) to deprotect the Fmoc group. Agitate for 2 minutes, drain, and repeat for 12 minutes.
3. Rinse the resin with DMF (2 mL, 5 times for 1 minute)
4. Bromoacetylation. Premix 0.8 M bromoacetic acid in DMF with 0.8 M N,N-diisopropylcarbodiimide (DIC), 2 mL total solution with 0.4 M of each reagent. Draw the solution into the syringe, agitate for 5 minutes, and rinse (2 mL DMF 5x1 minute).
5. Displacement. Draw in 1.5 mL of submonomer solution (0.2 M in DMF), agitate for 1 hour at 45 °C, then rinse (2 mL DMF 5x1 minute).
6. Repeat bromoacetylation and displacement until synthesis is finished. Wash with DCM (2 mL 3x1 minute) after last DMF wash and dry resin by pulling the plunger out and applying a gentle vacuum onto the syringe needle.

Example 15--Deprotection, cleavage, and purification

Dry resin (100-150 mg) was placed into a scintillation vial and was swollen in 9 mL of DCM by shaking for 30 minutes. 1 mL of 1.0 M BBr₃ in hexanes was added via a syringe, the vial was capped and shaken for 60 minutes ensuring that all of the resin was thoroughly submerged; this removes benzyl protecting groups from HOPO units. The solvent was

carefully removed with a glass pipette and the resin was washed with DCM (2 mL) methanol (2x2 mL) followed by DCM (2x2 mL). The peptoid was then cleaved from resin by treatment with cleavage cocktail for 60 minutes (the treatment also deprotects CAM units). The cleavage cocktail (95 % trifluoroacetic acid, 2.5 % water, 2.5% triethylsilane) was filtered from resin and a small aliquot was removed and diluted with methanol for LC-MS analysis (1→30% MeCN in H₂O over 20 minutes, both with 0.1 % formic acid); the resin was washed of TFA traces and discarded. Most LC-MS analyses showed a relatively clean desired compound; iron complexes were sometimes seen, which we believe came from stainless steel components of the instrument. Volatiles were then removed from the filtrate using a vacuum pump. The resulting residue was dissolved in 90/10 acetic acid/water (0.5-1 mL) and the resulting clear solution was stirred at 42 °C and treated with water in 0.5 mL increments. The solution turned turbid upon addition of water and slowly clarified with continued stirring (5-15 min between additions). A total of ~2.5 mL of water was added, at which point the solution remained turbid even with prolonged stirring.

The turbid solution was taken up into a syringe and injected onto reverse-phase prep-HPLC through a 0.45 µm filter in no more than 2.0 mL batches (~ 2 injections per peptoid).

Example 16--Reversed-Phase Preparative HPLC Method

Time (min)	% Acetonitrile (with 0.1 % TFA)	% Water (with 0.1 % TFA)	Flow (ml/min)
0	5	95	10
5	5	95	10
25	20	80	15
70	40	20	15
75	90	10	15
80	5	95	15
85	5	95	15

Most peptoids had peak maxima between 20 and 40 minutes, methods were typically terminated once the target material was collected. The column was flushed with 50/50 solvent composition for 5 minutes and equilibrated to initial condition for at least 20 minutes before every injection; insufficient equilibration leads to low column loading and very low yields. Peptoids with higher CAM compositions tended to be less polar and thus came out later than HOPO-heavy analogs.

A representative reversed-phase preparative HPLC trace is presented in FIG. 41.

Example 17--Peptoid LC-MS of purified peptoids:

LC traces of peptoids A@320. Ion counts along the entire peak including shoulders were used to generate mass spectra. Spectra were obtained in negative mode on Agilent 6530 mass spectrometer.

MS labels from left to right and are shown in FIG. 42A (HHHC Peptoid. MS1: free peptoid, MS2: peptoid-Na, MS3: peptoid-Fe and peptoid-Fe-K. FIG. 42B is the mass spectra for CHHH Peptoid. MS1: free peptoid and peptoid-K, MS2: peptoid-Na, peptoid-Fe, and possible impurity. FIG. 42C is the mass spectra for HCHH Peptoid. MS1: free peptoid and peptoid-K, MS2: peptoid-Na, peptoid-Fe. FIG. 42D is the mass spectra for HHCH Peptoid. MS1: free peptoid and possible impurity, MS2: peptoid-Na, MS3: Peptoid-Fe. FIG. 42E is the mass spectra for CHHC Peptoid. MS1: free peptoid only and peptoid-K, MS2: peptoid-Na and peptoid-Fe. FIG. 42F is the mass spectra for HHCC Peptoid. MS1: free peptoid and peptoid-Na, MS2: peptoid-Na, free peptoid, and peptoid-Fe. FIG. 42G is the mass spectra for CCHH Peptoid. MS1: free peptoid and peptoid-Na, MS2: Peptoid-Fe and possible trimer. FIG. 42H is the mass spectra for HCHC Peptoid. MS1: free peptoid, MS2: peptoid-Fe and peptoid-Na. FIG. 42I is the mass spectra for HCCH Peptoid. MS1: free peptoid and small amount trimer, MS2: peptoid-Fe, MS3: peptoid-Na and possible fragments. FIG. 42J is the mass spectra for CHCH Peptoid. MS1: free peptoid and peptoid-K, MS2: peptoid-Fe. FIG. 42K is the mass spectra for HCCC Peptoid. MS1: free peptoid and trimer-Fe trace, MS2: free peptoid and peptoid-Fe. FIG. 42L is the mass spectra for CHCC Peptoid. MS1: free peptoid and peptoid-Na, MS2: free peptoid and unidentified masses. FIG. 42M is the mass spectra

for CCHC Peptoid. MS1: free peptoid, MS2: free peptoid and peptoid-Na. FIG. 42N is the mass spectra for CCCH Peptoid. MS1: free peptoid and peptoid-Na, MS2: free peptoid and peptoid-K/Na. FIG. 42O is the mass spectra for CCCH Peptoid. MS1: free peptoid and peptoid-Na, MS2: free peptoid and peptoid-K/Na. FIG. 42P is the mass spectra for HHHH Peptoid. MS1: free peptoid, MS2: peptoid-Fe/Na, MS3: Peptoid-Fe-Na.

Example 18--High Resolution Mass Spectrometry Data

FIGs. 43A-43C depict the spectra that were obtained in negative mode. All peptoids are 1:1 Fe^{3+} complexes except for CCCC and HHHH. Top spectrum is calculated and bottom is obtained.

Example 19--TOF MSMS of Select Peptoids

FIG. 44 depicts the positive mode, fragmentation of molecular ion, masses of interest. Differences in peptoids are illustrated based on both composition and sequence. H and C units differ by a mass of 1 amu. FIGs. 45A-45D depict the NMR results.

[0232] The references cited in this disclosure are incorporated by reference herein in their entireties.

References

[0233] “Designing a Process for Selecting a Site for a Deep-Mined, Geologic Repository for High Level Radioactive Waste and Spent Nuclear Fuel,” United States Nuclear Waste Technical Review Board, 1-228 (2015).

[0234] Travis S. Grimes, Richard D. Tillotson, and Leigh R. Martin, “Trivalent Lanthanide/Actinide Separation Using Aqueous-Modified TALSPEAK Chemistry,” Solvent Extr. Ion Exch. 32(4), 378-390 (2014).

[0235] Kenneth L. Nash, “The Chemistry of TALSPEAK: A Review of the Science,” Solvent Extr. Ion Exch. 33(1), 1-55 (2015).

[0236] Gregg J. Lumetta, Amanda J. Casella, Brian M. Rapko, Tatiana G., Levitskaia, et al., “An Advanced TALSPEAK Concept Using 2-Ethylhexylphosphonic Acid Mono-2-Ethylhexyl Ester as the Extractant,” *Solvent Extr. Ion Exch.* 33(3), 211-223 (2015).

[0237] Manuel Sturzbecher-Hoehne, Clara Ng Pak Leung, Anthony D'Aleo, et al. “ 3,4,3-LI(1,2-HOPO): *In vitro* formation of highly stable lanthanide complexes translates into efficacious *in vivo* europium decorporation,” *Dalton Trans.* 40, 8340-8346 (2011).

[0238] Ilya Captain, Gauthier J.-P. Deblonde, Peter B. Rupert, et al. “Engineered Recognition of Tetravalent Zirconium and Thorium by Chelator—Protein Systems: Toward Flexible Radiotherapy and Imaging Platforms,” *Inorg. Chem.* 55(22), 11930-11936 (2016).

[0239] R.D Shannon, “Revised Effective Ionic Radii and Systematic Studies of Interatomic Distances in Halides and Chalcogenides,” *Acta Cryst.* A32, 751-767 (1976).

[0240] Manuel Sturzbecher-Hoehne, Birgitta Kullgren, Erin E. Jarvis, et al. “Highly Luminescent and Stable Hydroxypyridinonate Complexes: A Step Towards New Curium Decontamination Strategies,” *Chem.—Eur. J.* 20 (32), 9962-9968 (2014).

[0241] Manuel Sturzbecher-Hoehne, P. Yang, A. D'Aleo, Rebecca Abergel, “Intramolecular sensitization of americium luminescence in solution: shining light on short-lived forbidden 5f transitions,” *Dalton Trans.* 45, 9912-9919 (2016).

[0242] David L. White, Patricia W. Durbin, Nylan Jeung, and Kenneth N. Raymond, “Specific Sequestering Agents for the Actinides. 16. Synthesis and Initial Biological Testing of Polydentate Oxohydroxypyridinecarboxylate Ligands,” *J. Med. Chem.* 31 (1), 11-18 (1988).

[0243] Jide Xy, Birgitta Kullgren, Patricia W. Durbin, and Kenneth N. Raymond, “Specific Sequestering Agents for the Actinides. 28. Synthesis and Initial Evaluation of Multidentate 4-Carbamoyl-3-hydroxy-1-methyl-2(1H)-pyridinone Ligands for *in Vivo* Plutonium (IV) Chelation,” *J. Med. Chem.* 38 (14), 2606-2614 (1988).

[0244] Martell, A. E.; Smith, R. M.; Motekaitis, R. J. NIST Standard Reference Database; National Institute of Standards and Technology: Gaithersburg, MD.

[0245] Rebecca J. Abergel, Anthony D'Aleo, Clara Ng Pak Leung, David K. Shuh, and Kenneth N. Raymond, “Using the Antenna Effect as a Spectroscopic Tool:

Photophysics and Solution Thermodynamics of the Model Luminescent Hydroxypyridonate Complex [Eu(III)(3,4,3-LI(1,2-HOPO))],” *Inorg. Chem.* 48, 10868-10870 (2009).

[0246] J.T. Harvey, Production of Actinium-225 via High Energy Proton Induced Spallation of Thorium-232. Final Technical Report DE-SC0003602, NorthStar Medical Radioisotopes, LLC, [https:// world wide web .osti.gov/scitech/servlets/purl/1032445/](https://worldwideweb.osti.gov/scitech/servlets/purl/1032445/).

[0247] M. Sturzbecher-Hoehne, G.J.-P. Deblonde, R.J. Abergel, Solution thermodynamic evaluation of hydroxypyridinonate chelators 3,4,3-LI(1,2-HOPO) and 5-LIO(Me-3,2-HOPO) for UO₂(VI) and Th(IV) decorporation, *Radiochim. Acta.* 101 (2013) 359–366. doi:10.1524/ract.2013.2047.

[0248] R.J. Abergel, A. D’Aléo, C. Ng Pak Leung, D.K. Shuh, K.N. Raymond, Using the Antenna Effect as a Spectroscopic Tool: Photophysics and Solution Thermodynamics of the Model Luminescent Hydroxypyridonate Complex [Eu(III)(3,4,3-LI(1,2-HOPO))], *Inorg. Chem.* 48 (2009) 10868–10870. doi:10.1021/ic9013703.

[0249] M. Sturzbecher-Hoehne, C. Ng Pak Leung, A. D’Aléo, B. Kullgren, A.-L. Prigent, D.K. Shuh, K.N. Raymond, R.J. Abergel, 3,4,3-LI(1,2-HOPO): In vitro formation of highly stable lanthanide complexes translates into efficacious in vivo europium decorporation, *Dalton Trans.* 40 (2011) 8340. doi:10.1039/c1dt10840a.

[0250] M. Sturzbecher-Hoehne, P. Yang, A. D’Aléo, R.J. Abergel, Intramolecular sensitization of americium luminescence in solution: shining light on short-lived forbidden 5f transitions, *Dalton Trans.* (2016). doi:10.1039/C6DT00328A.

[0251] M. Sturzbecher-Hoehne, B. Kullgren, E.E. Jarvis, D.D. An, R.J. Abergel, Highly Luminescent and Stable Hydroxypyridinonate Complexes: A Step Towards New Curium Decontamination Strategies, *Chem. - Eur. J.* 20 (2014) 9962–9968. doi:10.1002/chem.201402103.

[0252] M. Sturzbecher-Hoehne, T.A. Choi, R.J. Abergel, Hydroxypyridinonate Complex Stability of Group (IV) Metals and Tetravalent f-Block Elements: The Key to the Next Generation of Chelating Agents for Radiopharmaceuticals, *Inorg. Chem.* 54 (2015) 3462–3468. doi:10.1021/acs.inorgchem.5b00033.

[0253] G.J.-P. Deblonde, M. Sturzbecher-Hoehne, R.J. Abergel, Solution Thermodynamic Stability of Complexes Formed with the Octadentate Hydroxypyridinonate

Ligand 3,4,3-LI(1,2-HOPO): A Critical Feature for Efficient Chelation of Lanthanide(IV) and Actinide(IV) Ions, *Inorg. Chem.* 52 (2013) 8805–8811. doi:10.1021/ic4010246.

[0254] G.J.-P. Deblonde, M. Sturzbecher-Hoehne, P.B. Rupert, D.D. An, M.-C. Illy, C.Y. Ralston, J. Brabec, W.A. de Jong, R.K. Strong, R.J. Abergel, Chelation and stabilization of berkelium in oxidation state +IV, *Nat. Chem.* (2017). doi:10.1038/nchem.2759.

[0255] A.E. Martell, R.M. Smith, R.J. Motekaitis, NIST Standard Reference Database 46, (n.d.).

[0256] D. Lundberg, I. Persson, The size of actinoid(III) ions - structural analysis vs. common misinterpretations, *Coord. Chem. Rev.* 318 (2016) 131–134. doi:10.1016/j.ccr.2016.04.003.

[0257] D. Lundberg, I. Persson, L. Eriksson, P. D'Angelo, S. De Panfilis, Structural Study of the N,N'-Dimethylpropyleneurea Solvated Lanthanoid(III) Ions in Solution and Solid State with an Analysis of the Ionic Radii of Lanthanoid(III) Ions, *Inorg. Chem.* 49 (2010) 4420–4432. doi:10.1021/ic100034q.

[0258] C. Wai, D. Fisher, others, Carboxylate-derived calixarenes with high selectivity for actinium-225, *Chem. Commun.* (1998) 377–378.

[0259] V. Radchenko, J.W. Engle, J.J. Wilson, J.R. Maassen, F.M. Nortier, W.A. Taylor, E.R. Birnbaum, L.A. Hudston, K.D. John, M.E. Fassbender, Application of ion exchange and extraction chromatography to the separation of actinium from proton-irradiated thorium metal for analytical purposes, *J. Chromatogr. A.* 1380 (2015) 55–63. doi:10.1016/j.chroma.2014.12.045.

[0260] V. Ostapenko, A. Vasiliev, E. Lapshina, S. Ermolaev, R. Aliev, Y. Totskiy, B. Zhuikov, S. Kalmykov, Extraction chromatographic behavior of actinium and REE on DGA, Ln and TRU resins in nitric acid solutions, *J. Radioanal. Nucl. Chem.* 306 (2015) 707–711. doi:10.1007/s10967-015-4331-y.

[0261] L.R. Morss, N.M. Edelstein, J. Fuger, *The Chemistry of the Actinide and Transactinide Elements*, 4th ed., Springer, 2010.

[0262] K.L. Nash, *The Chemistry of TALSPEAK: A Review of the Science*, *Solvent Extr. Ion Exch.* 33 (2015) 1–55. doi:10.1080/07366299.2014.985912.

[0263] K.V. Lohithakshan, P. Patil, S.K. Aggarwal, Solvent extraction studies of plutonium(IV) and americium(III) in room temperature ionic liquid (RTIL) by di-2-ethyl hexyl phosphoric acid (HDEHP) as extractant, J. Radioanal. Nucl. Chem. 301 (2014) 153–157. doi:10.1007/s10967-014-3119-9.

[0264] J.T. Harvey, Production of Actinium-225 via High Energy Proton Induced Spallation of Thorium-232. Final Technical Report DE-SC0003602, NorthStar Medical Radioisotopes, LLC, n.d. [https:// world wide web.osti.gov/scitech/servlets/purl/1032445/](https://world.wide.web.osti.gov/scitech/servlets/purl/1032445/).

What is claimed is:

1. A method of processing spent nuclear fuel, comprising:
obtaining spent nuclear fuel dissolved in an acidic medium;
contacting the dissolved spent nuclear fuel with an octadentate ligand and an organic phase to generate a mixture; and
separating an aqueous phase enriched for Pu^{4+} from the mixture.
2. A method for enriching a metal ion comprising:
contacting a first aqueous phase with an organic phase to generate a mixture, wherein the first aqueous phase comprises:
a plurality of metal ions; and
an octadentate ligand, and wherein the aqueous phase has an acidic pH of less than 1;
and
separating from the mixture a second aqueous phase enriched for a metal ion of the plurality of metal ions.
3. A method of preparing a medical isotope, the method comprising:
obtaining a metallic precursor dissolved under an acidic condition, wherein the metallic precursor comprises a medical isotope;
contacting the dissolved metallic precursor with an octadentate ligand and an organic phase to generate a mixture; and
separating the medical isotope from one or more metal ions in the dissolved metallic precursor based on an interaction between the octadentate ligand and the medical isotope.
4. A method of separating metal ions for nuclear forensics, comprising:
obtaining a sample derived from a nuclear material, wherein the sample comprises UO_2^{2+} , Ac^{3+} , Pu^{4+} , and Np^{4+} ; contacting the sample with an octadentate ligand to generate a first mixture, wherein the first mixture has an acidic pH of 1 or lower; separating UO_2^{2+} and Ac^{3+} from Pu^{4+} and Np^{4+} in the mixture based on an interaction or lack of interaction between the octadentate ligand and each of UO_2^{2+} , Ac^{3+} , Pu^{4+} , and Np^{4+} , to generate a second mixture comprising the Pu^{4+} and Np^{4+} ; and further separating each metal within both mixtures by classical techniques.
5. A method of separating metal ions, comprising:
contacting a liquid composition comprising a plurality of metal ions with an octadentate ligand to generate a mixture, under conditions sufficient to form a metal ion-ligand complex comprising a metal ion of the plurality of metal ions; and
separating a first fraction of the mixture enriched for the metal ion-ligand complex from a second fraction depleted for the metal ion-ligand complex, wherein the first fraction of the mixture has an acidic pH of less than 1,

wherein the plurality of metal ions are selected from the group consisting of: a p-, d- or f-block element of period 5 or greater, a group 3 element, or a group 4 element.

6. The method of any one of the preceding claims, wherein acidic denotes a pH of between about 1 to about 0.

7. The method of any one of the preceding claims, wherein the octadentate ligand is a hydroxypyridonate ligand.

8. The method of claim 7, wherein the octadentate ligand is built on a spermine scaffold.

9. The method of claim 7, wherein the octadentate ligand is 3,4,3-LI(1,2-HOPO).

10. The method of any one of the preceding claims, further comprising dissolving a metal in an acidic solution to obtain the liquid composition.

11. The method of claim 5, wherein the first fraction is enriched for a tetravalent metal ion and the second fraction is enriched for a trivalent metal ion and/or a divalent metal ion.

12. The method of claim 4 or 5, wherein the contacting further comprises contacting the sample with an organic phase.

13. A method of separating trivalent metal ions from tetravalent metal ions, the method comprising:
providing an aqueous solution comprising tetravalent metal ions and trivalent metal ions, wherein the solution has an acidic pH;

adding a HOPO chelator to the aqueous solution;

performing an extraction against the aqueous solution, wherein the HOPO chelator functions as a holdback agent to keep the tetravalent metal ions in the aqueous solution, while allowing the trivalent metal ions to be removed via decanting at the end of the extraction.

14. The method of claim 13, further comprising separating the HOPO chelator from the tetravalent metal to thereby collect the tetravalent metal.

15. The method of any one of the preceding claims, further comprising lowering a pH beneath 0 to separate tetravalent ions.

16. The method of any one of the preceding claims, further comprising an extractant.

17. The method of claim 16, wherein the extractant is selected from the group consisting of di-(2-ethylhexyl)phosphoric acid (HDEHP), a derivative of HDEHP, calixarenes, diglycoamide derivatives, carbamoylphosphine oxide derivatives, tributylphosphate (TBP), monoamide derivatives, and tertiary amines, and quaternary ammonium salts.

18. The method of any one of the preceding claims, wherein separating comprises filtering, precipitating, liquid-liquid extraction or chromatography, or wherein the method further comprises filtering, precipitating, liquid-liquid extraction or chromatography.

19. The method of claim 5, wherein the first fraction is enriched for a first metal ion that has a charge that is different from a charge of a second metal ion enriched in the second fraction.


20. The method of any one of the preceding claims, wherein

- 1) the separating comprises contacting the liquid composition with an organic phase or
 - 2) further comprising contacting the liquid composition with an organic phase.
21. The method of any one of the preceding claims, wherein the method does not comprise adjusting the pH of the liquid composition after dissolving the metal.
 22. The method of any one of the preceding claims, wherein the method does not comprise raising the pH of the liquid composition after dissolving the metal.
 23. The method of any one of the preceding claims, wherein the organic phase comprises a non-selective extractant.
 24. The method of any one of the preceding claims, wherein the organic phase does not comprise an extractant that selectively binds to the first or second metal ion.
 25. The method of any one of the preceding claims, wherein the method does not comprise adjusting the pH of the dissolved spent nuclear fuel.
 26. A method for separating metal ions, the method comprising:
 - providing an aqueous solution comprising a first metal ion and a second metal ion, wherein the solution has an acidic pH;
 - adding a HOPO chelator to the aqueous solution;
 - performing an extraction against the aqueous solution, wherein the HOPO chelator functions as a holdback agent to selectively keep the first metal ion in the aqueous solution, while allowing the second metal ion to go into an organic phase during the extraction.
 27. The method of claim 26, wherein the first metal ion is a tetravalent metal ion and the second metal ion is selected from the group consisting of a divalent metal ion and a trivalent metal ion.
 28. The method of claim 26, wherein the first metal ion is a trivalent metal ion and the second metal ion is a trivalent metal ion.
 29. The method of claim 28, wherein the HOPO chelator binds to the first metal ion more effectively than it binds to the second metal ion at a given pH.
 30. The method of any one of the preceding claims, wherein the fuel, first aqueous phase, precursor, nuclear material, liquid composition, or solution comprises actinium.
 31. The method of claim 30, wherein the actinium is separated from a M^{4+} metal.
 - ~~32. A composition having a structure comprising:----- }~~

~~Deleted by RO/US~~

^ RO/US

^ 32. A composition having a structure comprising::




wherein:

- (i) A1, A2, A3, and A4, individually, comprise a CAM group, a 1,2-HOPO group, or a HA group;
- (ii) B1, B2, B3, and B4, individually, comprise an amide group or an amine group;
- (iii) at least one of C1, C2, C3, C4, C5, or C6, individually, comprise SH, C(=O)OH, a NH₂;
- (iv) at least another one of C1, C2, C3, C4, C5, or C6 is optional;
- (v) at least one of L1, L2, L3, L4, L5, L6, L7, L8, L9, L10, L11, L12, or L13, individually, comprise H, an alkyl group having no greater than 10 carbon atoms, an alkylamino group having no greater than 10 carbon atoms and no greater than 2 nitrogen atoms; an alkyl ether group having no greater than 10 carbon atoms, a hydroxy ester group, or an alkyl ester group having no greater than 10 carbon atoms; and
- (vi) at least one of L1, L5, L6, L7, L8, L9, L10, L11, L12, or L13 is optional.

33. A composition of claim 32, wherein at least another one of L2, L3, or L4, individually, comprise an amine group or an amide group.

34. A composition of claim 32, wherein L1, C1, L7, C2, L9, C3, L11, C4, and L13, C5 are absent, L5 comprises an unsubstituted alkyl group having no greater than 5 carbon atoms, and C6 comprises SH, C(=O)OH, or NH₂.
35. A composition of claim 34, wherein L2, L3, L4, L6, L8, L10, and L12, individually, comprise an unsubstituted alkyl group having no greater than 5 carbon atoms.
36. A composition of claim 35, wherein A1 comprises a CAM group or a HOPO group; A2 comprises a HA group, A3 comprises a HA group, and A4 comprises a CAM group, a HOPO group, or a HA group.
37. A composition of claims 32, wherein at least one of L2, L3, or L4, individually, comprise an alkylamino group.
38. A composition of claim 32, wherein B1, B2, and B3, individually, comprise an amide group and B4 comprises an amino group, L2 and L3 comprise an amino group, and L4 comprises an alkyl group having no greater than 5 carbon atoms
39. A composition of claim 38, wherein:
C1, C2, C3, C4, C5, L1, A1, A2, A3, L1, L6, L7, L8, L9, L10, L11, L12, and L13 are absent,
A4 comprises a CAM group, a HOPO group, or a HA group; and
L5 comprises an alkyl group having no greater than 5 carbon atoms.
40. A composition of claim 32, wherein B1, B2, and B3, individually, comprise an amide group and B4 comprises an amide group, L2 and L3, individually, comprise an amino group, and L4 comprises an alkyl group having no greater than 5 carbon atoms.
41. A composition of claim 40, wherein C1, C2, C3, C4, C5, A1, A2, A3, L1, L6, L7, L8, L9, L10, L11, and L13 are absent, L12 comprises an amino group, L5 comprises an ether group having no greater than 10 carbon atoms, and A4 comprises a CAM group, a HOPO group, or a HA group.
42. A composition of claim 32, wherein C1, C2, C5, C6, L1, L2, L3, L4, L5, L7, L13, B2, and B4 are absent, B1 and B3, individually, comprise an amide group, L6, L8, L10, and L12, individually, comprise an amino group, A1, A2, A3, and A4, individually, comprise a CAM group, a HOPO group, or a HA group, L9 and L11, individually, comprise an alkyl group having no greater than 5 carbon atoms.
43. A composition, comprising a structure:



wherein:

at least one of R₁, R₂, R₃, R₄, and R₅, individually, comprise a CAM group, a HA group, or a 1,2-HOPO group;

at least another one of R_1 , R_2 , R_3 , R_4 , and R_5 , individually, comprise H or an alkyl group having from 1 to 10 carbon atoms;


R_6 comprises (i) H, (ii) an alkyl group having from 1 to 10 carbon atoms, or (iii) an alkyl group having from 1 to 10 carbon atoms and substituted by at least one of SH, NH_2 , or $C(=O)OH$;

m can be from 1 to 6;

n can be from 1 to 6;

o can be from 1 to 6.

44. A composition of claim 43, comprising a structure:



wherein:

at least one of R_1 , R_3 , R_4 , or R_5 ; R_1 , R_2 , R_3 , R_4 , and R_5 , individually, comprise a CAM group, a HA group, or a 1,2-HOPO group;

optionally, another one of R_1 , R_3 , R_4 , or R_5 ; R_1 , R_2 , R_3 , R_4 , and R_5 , individually, comprise H or an alkyl group having from 1 to 10 carbon atoms;

R_2 comprises H or an alkyl group including from 1 to 5 carbon atoms;

R_7 comprises SH, $C(=O)OH$, or NH_2 ; and

p is from 0 to 4.


45. A composition of claim 44, wherein:

R_1 comprises a CAM group or a 1,2-HOPO group;

R_3 and R_4 , individually, comprise a HA group; and

R_5 comprises a CAM group, a 1,2-HOPO group, or a HA group.

46. A composition of claim 43, comprising a structure:




wherein:

R₇ comprises SH, NH₂, or C(=O)OH;

R₂, R₈, and R₉, individually, comprise H, OH, or an alkyl group including from 1 to 5 carbon atoms; and p is from 0 to 4.

47. A composition of claim 43, comprising a structure:




wherein:

R₇ comprises SH, NH₂, or C(=O)OH;

R₂, R₈, and R₉, individually, comprise H, OH, or an alkyl group including from 1 to 5 carbon atoms; and p is from 0 to 4.

48. A composition of claim 43, comprising a structure:




wherein:

R₇ comprises SH, NH₂, or C(=O)OH;

R₂, R₈, R₉, and R₁₀, individually, comprise H, OH, or an alkyl group including from 1 to 5 carbon atoms; and

p is from 0 to 4.

49. A composition of claim 43, comprising a structure:




wherein:

R₇ comprises SH, NH₂, or C(=O)OH;

R₂, R₈, and R₉, individually, comprise H, OH, or an alkyl group including from 1 to 5 carbon atoms; and p is from 0 to 4.

50. A composition of claim 43, comprising a structure:




wherein:

R₇ comprises SH, NH₂, or C(=O)OH;

R₂, R₈, and R₉, individually, comprise H, OH, or an alkyl group including from 1 to 5 carbon atoms; and
p is from 0 to 4.

51. A composition of claim 43, comprising a structure:




wherein:


R₇ comprises SH, NH₂, or C(=O)OH;

R₂, R₈, R₉, and R₁₀, individually, comprise H, OH, or an alkyl group including from 1 to 5 carbon atoms;
and
p is from 0 to 4.


52. A composition of claim 43, comprising a structure:



53. A composition of claim 43, comprising a structure:



54. A composition, comprising a structure:



wherein:

at least one of R_{11} , R_{12} , R_{13} , or R_{15} , individually, comprise a CAM group, a HA group, or a 1,2-HOPO group;

optionally, at least another one of R_{11} , R_{12} , R_{13} , or R_{15} , individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{17} comprises SH, NH_2 , or $C(=O)OH$;

R_2 , R_{14} , and R_{16} , individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms; and r can be from 0 to 6.


55. A composition of claim 54, wherein:

R_{11} comprises a CAM group or a 1,2-HOPO group;

R_{12} and R_{15} , individually, comprise a HA group; and

R_{13} comprises a CAM group, a 1,2-HOPO group, or a HA group.

56. A composition of claim 54, comprising a structure:




wherein:

R_2 , R_{14} , R_{16} , R_{18} , and R_{19} , individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{17} comprises SH, NH_2 , or $C(=O)OH$; and

r can be from 0 to 4.

57. A composition of claim 54, comprising a structure:




wherein

R₂, R₁₄, R₁₆, R₁₈, and R₁₉, individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R₁₇ comprises SH, NH₂, or C(=O)OH; and

r is from 0 to 4.

58. A composition of claim 54, comprising a structure:




wherein:

R_2 , R_{14} , R_{16} , R_{18} , R_{19} , and R_{20} , individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{17} comprises SH, NH_2 , or $C(=O)OH$; and

r can be from 0 to 4.

59. A composition of claim 54, comprising a structure:




wherein:

R_2 , R_{14} , R_{16} , R_{18} , R_{19} , and R_{20} , individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{17} comprises SH, NH_2 , or $C(=O)OH$; and

r can be from 0 to 4.

60. A composition of claim 54, comprising a structure:




wherein:

R₂, R₁₄, R₁₆, R₁₈, and R₁₉, individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R₁₇ comprises SH, NH₂, or C(=O)OH; and

r is from 0 to 4.

61. A composition of claim 54, comprising a structure:




wherein:

R₂, R₁₄, R₁₆, R₁₈, and R₁₉, individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R₁₇ comprises SH, NH₂, or C(=O)OH;

r is from 0 to 4.

62. A composition, comprising a structure:



wherein:

R₂₁ and R₂₂, individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R₂₃ comprises H, OH, an alkyl group having from 1 to 10 carbon atoms, or (CH₂)_eR_a, where R_a is SH, C(=O)OH, or NH₂;

R₂₄ comprises a substituent having a CAM group, a 1,2-HOPO group, or a HA group;


a, b, and c, individually, are from 1 to 10;

d is from 1 to 4; and

e is from 1 to 10.

63. A composition of claim 62, wherein R₂₄ comprises a substituent having SH, C(=O)OH, or NH₂.

64. A composition of claim 62, comprising a structure:




wherein:

R₂₅, R₂₆, and R₂₇, individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R₂₈ comprises H, an alkyl group having from 1 to 5 carbon atoms, SH, NH₂, or C(=O)OH; and

s is from 0 to 4.

65. A composition of claim 62, comprising a structure:




wherein:

R_{25} , R_{26} , R_{27} , and R_{30} , individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;
 R_{28} and R_{29} , individually, comprise H, an alkyl group having from 1 to 5 carbon atoms, SH, NH_2 , or $\text{C}(=\text{O})\text{OH}$;

s is from 0 to 4; and

t is from 0 to 4.

66. A composition of claim 62, comprising a structure:




wherein:

R_{25} , R_{26} , R_{27} , and R_{30} , individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;
 R_{28} and R_{29} , individually, comprise H, an alkyl group having from 1 to 5 carbon atoms, SH, NH_2 , or $\text{C}(=\text{O})\text{OH}$;

s is from 0 to 4; and

t is from 0 to 4.

67. A composition of claim 62, comprising a structure:



wherein:


R_{25} , R_{26} , R_{27} , R_{30} , and R_{31} , individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{28} and R_{29} , individually, comprise H, an alkyl group having from 1 to 5 carbon atoms, SH, NH_2 , or $\text{C}(=\text{O})\text{OH}$;

s is from 0 to 4; and

t is from 0 to 4.

68. A composition of claim 62, comprising a structure:




wherein

R_{25} , R_{26} , and R_{27} , individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;

R_{28} comprises H, an alkyl group having from 1 to 5 carbon atoms, SH, NH_2 , or $\text{C}(=\text{O})\text{OH}$; and

s is from 0 to 4.


69. A composition of claim 62, comprising a structure:



wherein:

R_{25} , R_{26} , R_{27} , and R_{32} , individually, comprise H, OH, or an alkyl group having from 1 to 10 carbon atoms;
 R_{28} comprises H, an alkyl group having from 1 to 5 carbon atoms, SH, NH_2 , or $C(=O)OH$; and
 s is from 0 to 4.

70. A composition, comprising a structure:




wherein:

A, B, C, and D, individually, comprise one or more amide groups, one or more amine groups, or an alkyl group having from 1 to 10 carbon atoms;

R_{33} , R_{34} , R_{35} , and R_{36} , individually, comprise a CAM group, a 1,2-HOPO group, or a HA group; and

g , h , i , and j , individually, are from 1 to 10.

71. A composition of claim 70, comprising a structure:




wherein:

R_{37} and R_{42} , individually, comprise H, an alkyl group having from 1 to 5 carbon atoms, SH, $C(=O)OH$, or NH_2 ;

R_{38} , R_{39} , R_{40} , and R_{41} , individually, comprise H, OH, or an alkyl group having from 1 to 5 carbon atoms;
 and

u and v, individually, are from 0 to 5.

72. A composition of claim 70, comprising a structure:




wherein:

R₃₇ and R₄₂, individually, comprise H, an alkyl group having from 1 to 5 carbon atoms, SH, C(=O)OH, or NH₂;

R₃₈, R₃₉, R₄₀, and R₄₁, individually, comprise H, OH, or an alkyl group having from 1 to 5 carbon atoms; and

u and v, individually, are from 0 to 5.

73. A composition of claim 70, comprising a structure:



wherein:

R_{37} and R_{42} , individually, comprise H, an alkyl group having from 1 to 5 carbon atoms, SH, C(=O)OH, or NH_2 ;

R_{38} , R_{39} , R_{40} , and R_{41} , individually, comprise H, OH, or an alkyl group having from 1 to 5 carbon atoms; and

u and v , individually, are from 0 to 5.

74. A composition comprising a siderocalin bound to any of the structures of claims 32-73.

FIG. 1




FIG. 2




FIG. 4





FIG. 5



1: World nuclear association
2: International Atomic Energy Agency, DOC(2012)15.

FIG. 6







FIG. 7



Adapted from Johnson and Maloney, 2013, US patent 8475747 B1

FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27




FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32




FIG. 33




FIG. 34




FIG. 35




FIG. 36





FIG. 37






FIG. 38A




FIG. 38B




FIG. 38C

FIG. 38D





FIG 38E






FIG 38F.




FIG. 39A





FIG. 39B

FIG. 40A

Disulfide / Thiol Conjugation







FIG. 40B

FIG. 40C






FIG. 41




FIG. 42A




FIG. 42B




FIG. 42C




FIG. 42D




FIG. 42E




FIG. 42F




FIG. 42G




FIG. 42H




FIG. 42I




FIG. 42J





FIG. 42K

FIG. 42L






FIG. 42M




FIG. 42N






FIG. 420


FIG. 42P






Left to right and top to bottom: HHHH, CHHH, HCHH, HHCH, CHHC, HHCC

FIG. 43A



Left to right and top to bottom: CCHH, HCHC, HCCH, CHCH, HCCC, CHCC

FIG. 43B



Left to right and top to bottom: CCHC, CCCH, CCCC (uncomplexed), HHHH:Zr:Na

FIG. 43C

FIG. 44




FIG. 45A




FIG. 45B




FIG. 45C




FIG 45D.

