



- (51) **International Patent Classification:**
A61K 48/00 (2006.01) *C12N 15/861* (2006.01)
- (21) **International Application Number:**
PCT/US2017/016303
- (22) **International Filing Date:**
2 February 2017 (02.02.2017)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
62/290,187 2 February 2016 (02.02.2016) US
62/290,209 2 February 2016 (02.02.2016) US
- (71) **Applicants:** MEIRAGTX UK II LIMITED [GB/GB]; 92 Britannia Walk, London N1 7NQ (GB). MEIRAGTX, LLC [US/US]; 450 East 29th Street, 15th Floor, New York, NY 10016 (US).
- (72) **Inventors:** VOLLES, Michael, J.; 69 Bolton Street, Apt. 102, Cambridge, MA 02140 (US). DANOS, Olivier, F.; 49 Hancock Street, Boston, MA 02114 (US). BOYNE, Alex, K.; 198 Van Vorst Street, Apt. 415, Jersey City, NJ 07302 (US). ZENNOU, Veronique; 22 Mercer Street, Apt. 2, Jersey City, NJ 07302 (US). GUO, Xuecui; 1 Bungtown Road, Cold Spring Harbor, NY 11724 (US).
- (74) **Agents:** SOMERVILLE, Deborah, A. et al.; Fox Rothschild, LLP, Princeton Pike Corporate Center, 997 Lenox Drive, Building 3, Lawrenceville, NJ 08648-2311 (US).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))



WO 2017/136608 A2

(54) **Title:** REGULATION OF GENE EXPRESSION VIA APTAMER-MEDIATED CONTROL OF SELF-CLEAVING RIBOZYMES

(57) **Abstract:** The invention provides polynucleotide constructs for the regulation of gene expression by aptamer-based modulation of self-cleaving ribozymes and methods of using the constructs to regulate gene expression in response to the presence or absence of a ligand that binds the aptamer. The invention further provides methods for making and using riboswitches that decrease target gene expression in response to an aptamer ligand as well as riboswitches that increase target gene expression in response to an aptamer ligand.

REGULATION OF GENE EXPRESSION VIA APTAMER-MEDIATED CONTROL OF SELF-CLEAVING RIBOZYMES

FIELD OF THE INVENTION

[0001] The invention provides polynucleotide constructs for the regulation of gene expression by aptamer-based modulation of self-cleaving ribozymes and methods of using the constructs to regulate gene expression in response to the presence or absence of a ligand that binds the aptamer.

BACKGROUND OF THE INVENTION

[0002] Small endonucleolytic ribozymes comprise RNA motifs of about 50-150 nucleotides (“nt”) in length that have intrinsic RNA cleavage activity. There are several classes of small endonucleolytic ribozymes including hammerhead, and hairpin. Recently, new classes of self-cleaving ribozymes have been identified including twister, twister sister, pistol and hatchet.

[0003] The present invention utilizes ribozymes linked to aptamers to create polynucleotide cassettes for the regulation of target gene expression in response to the presence or absence of a ligand that binds the aptamer.

SUMMARY OF THE INVENTION

[0004] In one aspect, the present invention provides a polynucleotide cassette for the regulation of the expression of a target gene comprising a riboswitch that comprises a twister ribozyme linked by a stem to an aptamer, wherein the stem linking the twister ribozyme to the aptamer attaches to the ribozyme at the location of the P3 stem of the twister ribozyme and wherein the target gene is linked to the P1 stem of the twister ribozyme. In this context, the riboswitch is an “off riboswitch” because aptamer/ligand binding increases the ribonuclease function of the twister ribozyme leading to cleavage of the target gene RNA that contains the polynucleotide cassette and thereby reduces target gene expression. In one embodiment, the aptamer binds a small molecule ligand.

[0005] In one embodiment, the twister ribozyme is from *Nasonia vitripennis*. In one embodiment, the twister ribozyme comprises SEQ ID NO.:38. In one embodiment, the twister ribozyme is from an environmental sample. In one embodiment, the twister ribozyme comprises SEQ ID NO.:39.

[0006] In one embodiment, the P1 stem comprises sequence that lengthens the P1 stem of the twister ribozyme by 1 to 3 base pairs. In one embodiment, the P1 stem of the twister ribozyme is 3 to 9 base pairs. In one embodiment, the P1 stem of the twister ribozyme is 4 to 7 base pairs. In one embodiment, the P1 stem of the twister ribozyme is 6 base pairs. In one embodiment, the P1 stem of the twister ribozyme is 7 base pairs. In one embodiment, the P1 stem of the twister ribozyme is 8 base pairs.

[0007] In one embodiment, the stem linking the ribozyme to the aptamer comprises sequence from the ribozyme P3 stem and/or sequence from the aptamer P1 stem. In one, embodiment, the stem linking the ribozyme to the aptamer does not comprises sequence from the ribozyme P3 stem and/or does not comprises sequence from the aptamer P1 stem. In one embodiment, the stem linking the ribozyme to the aptamer is about 3 to about 7 base pairs long. In one embodiment, the stem linking the ribozyme to the aptamer is 3 to 7 base pairs long. In one embodiment, the stem linking the ribozyme to the aptamer is 4 base pairs long. In one embodiment, the stem linking the ribozyme to the aptamer is 3 base pairs long.

[0008] In another aspect, the present invention provides a method of modulating the expression of a target gene comprising:

[0009] (a) inserting a polynucleotide cassette comprising an off riboswitch described herein into a target gene,

[0010] (b) introducing the target gene comprising the polynucleotide cassette into a cell, and

[0011] (c) exposing the cell to a small molecule ligand that specifically binds the aptamer in an amount effective to decrease expression of the target gene.

[0012] In one embodiment, the polynucleotide cassette is inserted into the 5' untranslated region of the target gene, the 3' untranslated region of the target gene, and/or downstream of the translation start codon of the target gene.

[0013] In one embodiment, when the polynucleotide cassette is inserted downstream (i.e., 3') of the translation start codon of the target gene, the riboswitch is adjacent to the start codon. In other embodiments, the riboswitch is about 1 to about 100, about 1 to about 50, or about 1 to about 20 nucleotides from the start codon of the target gene. In one embodiment, the polynucleotide cassette comprises a 2A peptide sequence between the 3' end of the riboswitch and the target gene coding sequence. In one embodiment, the polynucleotide cassette comprises a linker sequence adjacent to the 2A peptide sequence.

[0014] In one embodiment, two or more of the polynucleotide cassettes are inserted into the target gene. In one embodiment, the two or more polynucleotide cassettes comprise different aptamers that specifically bind to different small molecule ligands. In one embodiment, the two or more polynucleotide cassettes comprise the same aptamer.

[0015] In one embodiment, the target gene comprising the polynucleotide cassette is incorporated in a vector for the expression of the target gene. In one embodiment, the vector is a viral vector. In one embodiment, the viral vector is selected from the group consisting of adenoviral vector, adeno-associated virus vector, and lentiviral vector.

[0016] In another aspect, the present invention provides a polynucleotide cassette for the regulation of the expression of a target gene comprising a riboswitch that comprises a twister ribozyme linked to an aptamer, wherein the aptamer is linked to the 3' or 5' end of the twister ribozyme P1 stem, wherein when the aptamer is linked to the 3' end of the twister ribozyme P1 stem, a portion of the 3' arm of the twister ribozyme P1 stem is alternatively the 5' arm of the aptamer P1 stem, and wherein when the aptamer is linked to the 5' end of the twister ribozyme P1 stem, a portion of the 5' arm of the twister ribozyme P1 stem is alternatively the 3' arm of the aptamer P1 stem (*see, e.g.*, Figs. 6a-6b). In this context, the riboswitch is an "on riboswitch" because aptamer/ligand binding inhibits the ribonuclease function of the twister ribozyme decreasing cleavage of the target gene RNA that contains the polynucleotide cassette and thereby increasing target gene expression. In one embodiment, the aptamer binds a small molecule ligand.

[0017] In one embodiment, the twister ribozyme is from *Nasonia vitripennis*. In one embodiment, the twister ribozyme comprises SEQ ID NO.:38. In one embodiment, the twister ribozyme is from an environmental sample. In one embodiment, the twister ribozyme comprises SEQ ID NO.:39.

[0018] In one embodiment, the portion of the arm of the twister ribozyme P1 stem that is alternatively an arm of the aptamer P1 stem is 4 to 8 nucleotides, 5 to 7 nucleotides, 6 nucleotides, or 7 nucleotides. In one embodiment, the P1 stem of the twister ribozyme is 4 to 9 base pairs, 5 to 8 base pairs, 6 base pairs, or 7 base pairs. In one embodiment, the P1 stem of the aptamer is 6 to 11 base pairs, 7 to 10 base pairs, 8 base pairs, or 9 base pairs.

[0019] In another aspect, the present invention provides a method of modulating the expression of a target gene comprising:

[0020] (a) inserting the polynucleotide cassette comprising an on riboswitch described herein into a target gene,

[0021] (b) introducing the target gene comprising the polynucleotide cassette into a cell, and

[0022] (c) exposing the cell to a small molecule ligand that specifically binds the aptamer in an amount effective to increase expression of the target gene.

[0023] In one embodiment, the polynucleotide cassette is inserted into the 5' untranslated region of the target gene, the 3' untranslated region of the target gene, and/or downstream of the translation start codon.

[0024] In one embodiment, when the polynucleotide cassette is inserted downstream (i.e., 3') of the translation start codon of the target gene, the on riboswitch is adjacent to the start codon. In other embodiments, the riboswitch is about 1 to about 100, about 1 to about 50, or about 1 to about 20 nucleotides from the start codon of the target gene. In one embodiment, the polynucleotide cassette comprises a 2A peptide sequence between the 3' end of the riboswitch and the target gene coding sequence. In one embodiment, the polynucleotide cassette comprises a linker sequence adjacent to the 2A peptide sequence.

[0025] In one embodiment, two or more of the polynucleotide cassettes are inserted into the target gene. In one embodiment, the two or more polynucleotide cassettes comprise different aptamers that specifically bind to different small molecule ligands. In one embodiment, the two or more polynucleotide cassettes comprise the same aptamer.

[0026] In one embodiment, the target gene comprising the polynucleotide cassette is incorporated in a vector for the expression of the target gene. In one embodiment, the vector is a viral vector. In one embodiment, the viral vector is selected from the group consisting of adenoviral vector, adeno-associated virus vector, and lentiviral vector.

[0027] In another aspect, the present invention provides vector comprising a target gene that contains a polynucleotide cassette of the present invention. In one embodiment, the vector is a viral vector. In one embodiment, the viral vector is selected from the group consisting of adenoviral vector, adeno-associated virus vector, and lentiviral vector.

BRIEF DESCRIPTION OF THE FIGURES

[0028] Figs. 1a-1b. Twister ribozymes self-cleave and suppress gene expression in mammalian cells.

[0029] Fig. 1a. Twister ribozyme sequence from *N. vitripennis* (wasp-twister) or from environmental sample (es-twister) was cloned into the 3' untranslated region ("UTR") in pEGFP-C1 vector. HEK 293 cells were transfected with the indicated constructs and fluorescence intensity of GFP measured. Wasp-twister caused approximately 97% or 36-fold reduction in GFP expression when compared to pEGFP-C1 control. The es-twister when cloned into the 3' UTR of GFP significantly reduced target gene expression in HEK 293 cells, but not to the same extent as the wasp twister.

[0030] Fig. 1b. Mutant twister sequence was generated by mutating the A6 to G, as indicated by the wasp-twister secondary structure embedded in the graph. HEK 293 cells were transfected with the indicated constructs and fluorescence intensity of GFP measured. The wasp-twister again reduced expression of GFP when cloned into the 3' UTR. The wasp-twister Mutant, however, did not cause a reduction in GFP expression.

[0031] Figs. 2a-2b. Modification of P1 stem sequence improves twister ribozyme cleavage activity.

[0032] Figs. 2a. The sequence of 3 base pair extension (3HP) of the wasp twister P1 stem.

[0033] Fig. 2b. Deletion of SacII site from the initial construct GFP_wasp-twister and addition of 3 bp stem to twister P1 stem improved cleavage by twister ribozyme, leading to further reduction in the GFP expression.

[0034] Fig. 2c. Shown is the result of luciferase assay. HEK 293 cells were transfected with the indicated constructs containing firefly luciferase gene and twister with or without 3HP in 3' UTR.

[0035] Fig. 3a-3e. Generation of guanine-responsive twister ribozyme.

[0036] Fig. 3a. The schematics of the linking aptamer sequence to twister ribozyme and the rationale for aptamer ligand responsive twister ribozyme. Aptamer sequence was shown in thicker line. In the absence of aptamer ligand, the ligand free aptamer disrupts twister ribozyme cleavage, leading to intact mRNA and subsequent gene expression. In the presence of aptamer ligand, ligand bound aptamer facilitates the P3 stem formation, thus restoring the twister ribozyme cleavage activity, leading to inhibition of gene expression.

[0037] Fig. 3b. The results of firefly luciferase assay. Twenty constructs were made by linking the P1 stem of guanine aptamer to the P3 stem of the twister ribozyme, as shown


in Fig. 3a, lower panel. The additional 3HP was counted in as part of the twister P1 stem. The stem that connects the Loop 2 and aptamer was truncated sequentially, generating different length of stems as indicated in the round brackets. HEK 293 cells were transfected with the indicated constructs, treated with or without guanine. The fold reduction for construct Luci-waspGua16 is indicated in the graph.

[0038] Fig. 3c. Validation of luciferase constructs containing waspGua16 or a mutant waspGua16. Wild type waspGua16 reduced luciferase activity upon treatment with 500 μ M guanine, while the mutant waspGua16 did not.

[0039] Fig. 3d. shown is the results of luciferase assay. HEK 293 cells were transfected with the indicated constructs, and treated with or without 500 μ M guanine or 1mM guanosine.

[0040] [0040] Fig. 3e. HEK 293 cells were transfected with the indicated constructs, and treated with 500 μ M guanosine. DMSO was used as solvent control. Supernatant was collected 18 hours after guanosine treatment, the Epo was measured using ELISA kit. The results were expressed as mean \pm SD (n=2). Shown is the representative of two independent experiments.

[0041] Figs. 4a-4b. Generation of theophylline responsive twister riboswitches.

[0042] Fig. 4a. Schematic of the linking theophylline aptamer to the loop 2 sequence in twister ribozyme. The sequences and the length of the stem connecting theophylline aptamer and twister loop 2 were shown. The aptamer and ligand was indicated as .

[0043] Fig. 4b. Shown are the results of the luciferase assay. HEK 293 cells were transfected with the indicated constructs, treated with and without 2 mM theophylline 4 hours after transfection.

[0044] Fig. 5a-5b. The portability of twister riboswitch in mammalian cells.

[0045] Fig. 5a. The schematics of constructs with twisterGua16 riboswitch and a linker and 2A peptide sequence (linker2A) inserted downstream of ATG and upstream of the remaining luciferase coding sequence (top panel), or twisterGua16 and linker2A sequence inserted downstream the coding sequence for the first 10 amino acids of GFP (lower panel).

[0046] Fig. 5b. Result of a luciferase assay are shown. HEK 293 cells were transfected with the indicated constructs. The transfected cells were treated with or without guanine 4 hours after transfection. Results is expressed as mean \pm SD (n=3).

[0047] Figs. 6a-6b. Generation of aptamer ligand responsive ON twister riboswitch.

[0048] Fig. 6a. Shows the schematics of the ON twister guanine riboswitch (where the aptamer is linked to the 3' end of twister P1). The configuration of the wasp twister and aptamer sequence is shown as an example, in which the 3' end of the P1 stem (including the 3 bp stem) was linked to the aptamer sequence. In this configuration, the stem sequence was shared between the twister P1 stem and the aptamer P1 stem. The shared stem sequence is indicated as in thicker line. Presumably, in the absence of the aptamer ligand (top panel), aptamer sequence linked to the twister P1 stem does not disrupt twister structure and its ribozyme cleavage activity, therefore resulting in cleavage of mRNA and subsequent inhibition of gene expression. In the presence of aptamer ligand (lower panel), aptamer forms P1 stem and makes the shared stem sequence (thicker line) unavailable for twister P1 stem formation, therefore disrupting twister structure and its ribozyme activity, resulting in intact mRNA and subsequent gene expression.

[0049] Fig. 6b. HEK 293 cells were transfected with the indicated luciferase constructs. The transfected cells were treated with or without 500 μ M guanosine. DMSO was used as solvent control. Results are expressed as mean \pm SD (n=3). The induction folds were indicated in the graph for each construct.

[0050] Figure 7a-7b. Hammerhead ribozyme sequence (7a) and Hairpin ribozyme sequence (7b) placed downstream of translation start codon. The top panel shows the nucleotide sequence, and the bottom panel shows the translated peptide sequences using different reading frames. The top line in each case is the parent ribozyme, not translated-through; dots in the nucleotide sequence are used to indicate "same as the top line." The 2A peptide sequence from *Thoseaasigna* virus (T2A) was indicated as 2A. Mutations were made to eliminate potential stop codons in the ribozyme sequences.

[0051] Figure 7c. Results of Beta-galactosidase activity generated from HEK 293 cells transfected with the indicated constructs. The left half of the graph shows the results from constructs with hammerhead ribozyme sequence, and the right half indicates the results from constructs with Hairpin ribozyme sequence. The fold reduction of the LacZ activity was expressed as the ratio of the value generated from ribozyme inactive mutant construct to the value generated from the ribozyme active construct. The highest fold was indicated by arrow for each ribozyme.

DETAILED DESCRIPTION OF THE INVENTION

[0052] The invention provides polynucleotide constructs for the regulation of gene expression by aptamer-based modulation of small endonucleolytic ribozymes and methods of using the constructs to regulate gene expression in response to the presence or absence of a ligand that binds the aptamer. The polynucleotide construct contains at least one riboswitch that contains a ribozyme and an aptamer. The way the ribozyme is attached to the aptamer provides for riboswitches that activate ribozyme self-cleavage in the presence of aptamer ligand (“off” riboswitches) or riboswitches that inhibit ribozyme self-cleavage in the presence of aptamer (“on” riboswitches).

[0053] The gene regulation polynucleotide cassette refers to a recombinant DNA construct that, when incorporated into the DNA of a target gene, provides the ability to regulate expression of the target gene by aptamer/ligand mediated regulation of ribozyme self-cleavage. As used herein, a polynucleotide cassette or construct is a nucleic acid (e.g., DNA or RNA) comprising elements derived from different sources (e.g., different organisms, different genes from the same organism, and the like). The polynucleotide cassette comprises a riboswitch. The riboswitch in the context of the present invention contains a sensor region (e.g., an aptamer) and a ribozyme that together are responsible for sensing the presence of a ligand that binds the sensor region and altering the conformation of ribozyme binding site. In one embodiment, the target gene’s expression is increased when the aptamer ligand is present and decreased when the ligand is absent. In another embodiment, the target gene’s expression is decreased when the aptamer ligand is present and increased when the ligand is absent.

[0054] Ribozymes

[0055] The ribozyme used in the present invention can be any small endonucleolytic ribozyme that will self-cleave in the target cell type including, for example, hammerhead, hairpin, the hepatitis delta virus, the Varkud satellite, twister, twister sister, pistol and hatchet. *See, e.g.,* Roth et al., Nat Chem Biol. 10(1):56-60; and Weinberg et al., Nat Chem Biol. 2015 Aug;11(8):606-10, both incorporated herein by reference. U.S. 2015/0056174 provides modified hammerhead ribozymes with enhanced endonucleolytic activity (*see, e.g.,* Figs. 5A-F, incorporated herein by reference). Preferred ribozymes are hammerhead, hairpin, and twister. In one embodiment, the ribozyme is the twister ribozyme is from *Nasonia vitripenni*

(SEQ. ID NO.:38). In one embodiment, the ribozyme is the twister ribozyme identified from an environmental sample (SEQ. ID NO.:39).

[0056] Riboswitch

[0057] The term “riboswitch” as used herein refers to a regulatory segment of a RNA polynucleotide. A riboswitch in the context of the present invention contains a sensor region (e.g., an aptamer) a self-cleaving ribonuclease that together are responsible for sensing the presence of a ligand (e.g., a small molecule) and modulating the conformation of a self-cleaving ribonuclease and thereby its activity. In one embodiment, the riboswitch is recombinant, utilizing polynucleotides from two or more sources. The term “synthetic” as used herein in the context of a riboswitch refers to a riboswitch that is not naturally occurring. In one embodiment, the sensor (e.g., aptamer) and ribonuclease regions are joined by a polynucleotide linker. In one embodiment, the polynucleotide linker forms a RNA stem (i.e., a region of the RNA polynucleotide that is double-stranded).

[0058] Off riboswitches

[0059] In one aspect, the present invention provides a polynucleotide cassette for the regulation of the expression of a target gene comprising a riboswitch that comprises a twister ribozyme linked by a stem to an aptamer, wherein the stem linking the twister ribozyme to the aptamer attaches to the ribozyme at the location of the P3 stem of the twister ribozyme, and wherein the target gene is linked to the P1 stem of the twister ribozyme (*see, e.g.,* Fig. 3a). In this context, the riboswitch is an “off riboswitch” because aptamer/ligand binding increases the ribonuclease function of the twister ribozyme leading to cleavage of the target gene RNA that contains the polynucleotide cassette thereby reducing target gene expression.

[0060] In this configuration, where both arms of an aptamer stem are linked to both arms of the ribozyme P3 stem, this stem may contain additional sequence capable of forming a stem when the aptamer binds its ligand. This additional stem sequence may include sequence from the aptamer P1 stem and/or additional sequence added to facilitate stem formation when the aptamer is bound by its ligand. The P3 stem may comprise one, two, three or more changes from the wild-type P3 sequence for the ribozyme. In some embodiments, the stem linking the aptamer to the ribozyme does not contain sequence from either the P3 stem of twister or from the aptamer stem. In addition, the stem linking the ribozyme to the aptamer (including the P3 stem) may comprise one or more mismatches where a nucleotide is not complementary to its counterpart on the other stem arm.

[0061] The stem linking the aptamer to the ribozyme should be of a sufficient length (and GC content) to form a stem upon ligand binding the aptamer thereby promoting the ribozyme endonuclease activity, while also, when the ligand is not present in sufficient quantities, taking a conformation that inhibits ribozyme endonuclease activity. The length and sequence of the stem can be modified using known techniques in order to identify stems that allow acceptable background expression of the target gene when ligand is present and acceptable expression levels of the target gene when the ligand is not present. If the stem is, for example, too long it may provide an active ribozyme conformation in the presence or absence of ligand. If the stem is too short, it may not provide an active ribozyme even in the presence of ligand. In certain embodiments, the stem linking the ribozyme to the aptamer is about 3 to about 7 base pairs. In one embodiment, the stem linking the ribozyme to the aptamer is 3 to 7 base pairs. In one embodiment, the stem linking the ribozyme to the aptamer is 4 base pairs long. In one embodiment, the stem linking the ribozyme to the aptamer is 3 base pairs long.

[0062] It should be understood that the stem linking the aptamer and ribozyme is not always in a double-stranded stem configuration. As shown in Fig. 3a, when the aptamer is not bound to ligand, the structure of the stem is disrupted inhibiting ribozyme endonucleolytic activity. When the aptamer binds its ligand the stem is stabilized promoting ribozyme endonucleolytic activity.

[0063] In this off riboswitch configuration, the P1 stem of the ribozyme is linked to the nucleotide sequence of the target gene (*see, e.g.*, Fig. 3a). This ribozyme P1 stem may comprise all or some of the P1 sequence of the naturally-occurring ribozyme. This P1 stem may contain additional stem sequence from another source. The P1 stem may comprise one, two, three or more changes from the wild-type P1 sequence for the ribozyme. In some embodiments, the P1 stem of the ribozyme does not contain any sequence from the wild-type P1 sequence for the ribozyme. The stem is still referred to as the P1 stem of the ribozyme by virtue of its location. The length and sequence of the P1 stem can be modified using known techniques in order to identify stems that allow, e.g., acceptable background expression of the target gene. In one embodiment, the P1 stem of the twister ribozyme is 3 to 9 base pairs. In one embodiment, the P1 stem of the twister ribozyme is 4 to 7 base pairs. In one embodiment, the P1 stem of the twister ribozyme is 7 base pairs. In one embodiment, the P1 stem of the twister ribozyme is 8 base pairs.

[0064] The polynucleotide cassette comprising the off riboswitch may be placed at one or more locations in the target gene, including, but not limited to the 5' and 3' UTR and downstream of the start codon.

[0065] On riboswitches

[0066] In another aspect, the present invention provides a polynucleotide cassette for the regulation of the expression of a target gene comprising a riboswitch that comprises a twister ribozyme linked to an aptamer, wherein the aptamer is linked to the 3' or 5' end of the twister ribozyme P1 stem, wherein when the aptamer is linked to the 3' end of the twister ribozyme P1 stem, a portion of the 3' arm of the twister ribozyme P1 stem is alternatively the part of the 5' arm of the aptamer P1 stem, and wherein when the aptamer is linked to the 5' end of the twister ribozyme P1 stem, a portion of the 5' arm of the twister ribozyme P1 stem is alternatively part of the 3' arm of the aptamer P1 stem (*see, e.g.*, Figs. 6a-6b). In this context, the riboswitch is an “on riboswitch” because aptamer/ligand binding inhibits the ribonuclease function of the twister ribozyme decreasing cleavage of the target gene RNA that contains the polynucleotide cassette and thereby increasing target gene expression in the presence of ligand.

[0067] As with the aptamer and ribozyme stems discussed with respect to off riboswitch, the ribozyme P1 stem and aptamer P1 stem may or may not comprise wild-type stem sequence. As such, the stem names are based on their location on the ribozyme or aptamer and does not imply that the stems comprise any particular sequence. The P1 ribozyme stem and the P1 aptamer stem may each independently comprise stem sequence in addition to the shared stem sequence. This additional stem sequence may include sequence from the respective ribozyme P1 or aptamer P1 stem and/or additional sequence added to facilitate stem formation. In addition, ribozyme P1 and/or aptamer P1 stem may comprise one or more mismatches where a nucleotide is not complementary to its counterpart on the other stem arm.

[0068] The ribozyme P1 and aptamer P1 stem should be of a sufficient length (and GC content) such that the aptamer forms the P1 stem in the presence of aptamer ligand and the ribozyme forms the P1 stem when ligand is not present. The length and sequence of the stems can be modified using known techniques in order to identify stems that allow acceptable background expression of the target gene when ligand is not present and acceptable expression levels of the target gene when the ligand is present. In one

embodiment, P1 stem of the twister ribozyme is 4 to 9 base pairs, 5 to 8 base pairs, 6 base pairs, or 7 base pairs. In one embodiment, the P1 stem of the aptamer is 6 to 11 base pairs, 7 to 10 base pairs, 8 base pairs, or 9 base pairs.

[0069] In this on riboswitch configuration, the aptamer is adjacent to the ribozyme (either 5' or 3') and the aptamer and ribozyme both comprise an alternatively shared stem arm. When the aptamer is bound to its ligand, it stabilizes the aptamer P1 stem, preventing its use as part of the ribozyme structure and inhibiting ribozyme endonuclease activity. When ligand is not present, the shared stem is free to form part of the ribozyme P1 stem, thereby allowing ribozyme endonuclease activity and cleavage of the target RNA leading to reduced target gene expression. In embodiments, the alternatively shared portion of the ribozyme and aptamer stems is 4 to 8 nucleotides, 5 to 7 nucleotides, 6 nucleotides, or 7 nucleotides.

[0070] The polynucleotide cassette comprising the on riboswitch may be placed at one or more locations in the target gene, including, but not limited to the 5' and 3' UTR and downstream of the start codon.

[0071] Aptamer/Ligand

[0072] In one embodiment, the sensor region comprises an aptamer. The term "aptamer" as used herein refers to an RNA polynucleotide that specifically binds to a ligand. The term "ligand" refers to a molecule that is specifically bound by an aptamer. In one embodiment, the ligand is a low molecular weight (less than about 1,000 Daltons) molecule including, for example, lipids, monosaccharides, second messengers, co-factors, metal ions, other natural products and metabolites, nucleic acids, as well as most therapeutic drugs. In one embodiment, the ligand is a polynucleotide with two or more nucleotide bases.

[0073] In one embodiment, the ligand is selected from the group consisting of 8-azaguanine, adenosine 5'-monophosphate monohydrate, amphotericin B, avermectin B1, azathioprine, chlormadinone acetate, mercaptopurine, moricizine hydrochloride, N6-methyladenosine, nadide, progesterone, promazine hydrochloride, pyrvinium pamoate, sulfaguanidine, testosterone propionate, thioguanosine, tyloxapol and vorinostat.

[0074] Aptamer ligands can also be cell endogenous components that increase significantly under specific physiological/pathological conditions, such as oncogenic transformation – these may include second messenger molecules such as GTP or GDP, calcium; fatty acids, or fatty acids that are incorrectly metabolized such as 13-HODE in

breast cancer (Flaherty, JT et al., Plos One, Vol. 8, e63076, 2013, incorporated herein by reference); amino acids or amino acid metabolites; metabolites in the glycolysis pathway that usually have higher levels in cancer cells or in normal cells in metabolic diseases; and cancer-associated molecules such as Ras or mutant Ras protein, mutant EGFR in lung cancer, indoleamine-2,3-dioxygenase (IDO) in many types of cancers. Endogenous ligands include progesterone metabolites in breast cancer as disclosed by JP Wiebe (Endocrine-Related Cancer (2006) 13:717–738, incorporated herein by reference). Endogenous ligands also include metabolites with increased levels resulting from mutations in key metabolic enzymes in kidney cancer such as lactate, glutathione, kynurenine as disclosed by Minton, DR and Nanus, DM (Nature Reviews, Urology, Vol. 12, 2005, incorporated herein by reference).

[0075] Aptamers have binding regions that are capable of forming complexes with an intended target molecule (i.e., the ligand). The specificity of the binding can be defined in terms of the comparative dissociation constants (K_d) of the aptamer for its ligand as compared to the dissociation constant of the aptamer for unrelated molecules. Thus, the ligand is a molecule that binds to the aptamer with greater affinity than to unrelated material. Typically, the K_d for the aptamer with respect to its ligand will be at least about 10-fold less than the K_d for the aptamer with unrelated molecules. In other embodiments, the K_d will be at least about 20-fold less, at least about 50-fold less, at least about 100-fold less, and at least about 200-fold less. An aptamer will typically be between about 15 and about 200 nucleotides in length. More commonly, an aptamer will be between about 30 and about 100 nucleotides in length.

[0076] The aptamers that can be incorporated as part of the riboswitch can be a naturally occurring aptamer, or modifications thereof, or aptamers that are designed de novo and/or screened through systemic evolution of ligands by exponential enrichment (SELEX) or other screening methods. Examples of aptamers that bind small molecule ligands include, but are not limited to theophylline, dopamine, sulforhodamine B, cellobiose, kanamycin A, lividomycin, tobramycin, neomycin B, viomycin, chloramphenicol, streptomycin, cytokines, cell surface molecules, and metabolites. For a review of aptamers that recognize small molecules, *see, e.g.*, Famulok, Science 9:324-9 (1999) and McKeague, M. & DeRosa, M.C. J. Nuc. Aci. 2012 (both of which are incorporated herein by reference). In another embodiment, the aptamer is a complementary polynucleotide.

[0077] Methods for identifying aptamer/ligand

[0078] In one embodiment, the aptamer is designed to bind a particular small molecule ligand. Methods for designing and selecting aptamers that bind particular ligands are disclosed in 62/370,599, incorporated herein by reference. Other methods for screening aptamers include, for example, SELEX. Methods for designing aptamers that selectively bind a small molecule using SELEX are disclosed in, e.g., U.S. Patent Nos. 5,475,096, 5,270,163, and Abdullah Ozer, et al. Nuc. Aci. 2014, which are incorporated herein by reference. Modifications of the SELEX process are described in U.S. Patent Nos. 5,580,737 and 5,567,588, which are incorporated herein by reference.

[0079] Selection techniques for identifying aptamers generally involve preparing a large pool of DNA or RNA molecules of the desired length that contain a region that is randomized or mutagenized. For example, an oligonucleotide pool for aptamer selection might contain a region of 20-100 randomized nucleotides flanked by regions of defined sequence that are about 15-25 nucleotides long and useful for the binding of PCR primers. The oligonucleotide pool is amplified using standard PCR techniques, or other means that allow amplification of selected nucleic acid sequences. The DNA pool may be transcribed *in vitro* to produce a pool of RNA transcripts when an RNA aptamer is desired. The pool of RNA or DNA oligonucleotides is then subjected to a selection based on their ability to bind specifically to the desired ligand. Selection techniques include, for example, affinity chromatography, although any protocol which will allow selection of nucleic acids based on their ability to bind specifically to another molecule may be used. Selection techniques for identifying aptamers that bind small molecules and function within a cell may involve cell based screening methods. In the case of affinity chromatography, the oligonucleotides are contacted with the target ligand that has been immobilized on a substrate in a column or on magnetic beads. The oligonucleotide is preferably selected for ligand binding in the presence of salt concentrations, temperatures, and other conditions which mimic normal physiological conditions. Oligonucleotides in the pool that bind to the ligand are retained on the column or bead, and nonbinding sequences are washed away. The oligonucleotides that bind the ligand are then amplified (after reverse transcription if RNA transcripts were utilized) by PCR (usually after elution). The selection process is repeated on the selected sequences for a total of about three to ten iterative rounds of the selection procedure. The resulting oligonucleotides are then amplified, cloned, and sequenced using standard procedures to identify the sequences of the oligonucleotides that are capable of binding the target ligand.

Once an aptamer sequence has been identified, the aptamer may be further optimized by performing additional rounds of selection starting from a pool of oligonucleotides comprising a mutagenized aptamer sequence.

[0080] In vivo aptamer screening may be used following one or more rounds of in vitro selection (e.g., SELEX). For example, Konig, J. et al. (RNA. 2007, 13(4):614–622, incorporated herein by reference) describe combining SELEX and a yeast three-hybrid system for in vivo selection of aptamer.

[0081] Target genes

[0082] The gene regulation cassette of the present invention is a platform that can be used to regulate the expression of any target gene that can be expressed in a target cell, tissue or organism. The term “target gene” refers to a polynucleotide that is introduced into a cell and is capable of being transcribed into RNA and translated and/or expressed under appropriate conditions. Alternatively, the target gene is endogenous to the target cell and the gene regulation cassette of the present invention is positioned into the target gene (for example into the 5' or 3' UTR of an endogenous target gene). An example of a target gene is a polynucleotide encoding a therapeutic polypeptide. In another embodiment, the target gene encodes an RNA such as a miRNA, rRNA, small or long noncoding RNAs, short hairpin RNA (shRNA) and any other regulatory RNAs. In one embodiment, the target gene is exogenous to the cell in which the recombinant DNA construct is to be transcribed. In another embodiment, the target gene is endogenous to the cell in which the recombinant DNA construct is to be transcribed.

[0083] The target gene according to the present invention may be a gene encoding a protein, or a sequence encoding a non-protein coding RNA. The target gene may be, for example, a gene encoding a structural protein, an enzyme, a cell signaling protein, a mitochondrial protein, a zinc finger protein, a hormone, a transport protein, a growth factor, a cytokine, an intracellular protein, an extracellular protein, a transmembrane protein, a cytoplasmic protein, a nuclear protein, a receptor molecule, an RNA binding protein, a DNA binding protein, a transcription factor, translational machinery, a channel protein, a motor protein, a cell adhesion molecule, a mitochondrial protein, a metabolic enzyme, a kinase, a phosphatase, exchange factors, a chaperone protein, and modulators of any of these. In embodiments, the target gene encodes erythropoietin (Epo), human growth hormone (hGH), transcription activator-like effector nucleases (TALEN), human insulin, CRISPR associated

protein 9 (cas9), or an immunoglobulin (or portion thereof), including, e.g., a therapeutic antibody.

[0084] Expression Constructs

[0085] The present invention contemplates the use of a recombinant vector for introduction into target cells a polynucleotide encoding a target gene and containing the gene regulation cassette described herein. In many embodiments, the recombinant DNA construct of this invention includes additional DNA elements including DNA segments that provide for the replication of the DNA in a host cell and expression of the target gene in that cell at appropriate levels. The ordinarily skilled artisan appreciates that expression control sequences (promoters, enhancers, and the like) are selected based on their ability to promote expression of the target gene in the target cell. “Vector” means a recombinant plasmid, yeast artificial chromosome (YAC), mini chromosome, DNA mini-circle or virus (including virus derived sequences) that comprises a polynucleotide to be delivered into a host cell, either in vitro or in vivo. In one embodiment, the recombinant vector is a viral vector or a combination of multiple viral vectors.

[0086] Viral vectors for the aptamer-mediated expression of a target gene in a target cell, tissue, or organism are known in the art and include adenoviral (AV) vectors, adeno-associated virus (AAV) vectors, retroviral and lentiviral vectors, and Herpes simplex type 1 (HSV1) vectors.

[0087] Adenoviral vectors include, for example, those based on human adenovirus type 2 and human adenovirus type 5 that have been made replication defective through deletions in the E1 and E3 regions. The transcriptional cassette can be inserted into the E1 region, yielding a recombinant E1/E3-deleted AV vector. Adenoviral vectors also include helper-dependent high-capacity adenoviral vectors (also known as high-capacity, “gutless” or “guttled” vectors), which do not contain viral coding sequences. These vectors, contain the cis-acting elements needed for viral DNA replication and packaging, mainly the inverted terminal repeat sequences (ITR) and the packaging signal (Ψ). These helper-dependent AV vector genomes have the potential to carry from a few hundred base pairs up to approximately 36 kb of foreign DNA.

[0088] Recombinant adeno-associated virus “rAAV” vectors include any vector derived from any adeno-associated virus serotype, including, without limitation, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-7 and AAV-8, AAV-9, AAV-10, and the like.

rAAV vectors can have one or more of the AAV wild-type genes deleted in whole or in part, preferably the Rep and/or Cap genes, but retain functional flanking ITR sequences. Functional ITR sequences are retained for the rescue, replication, packaging and potential chromosomal integration of the AAV genome. The ITRs need not be the wild-type nucleotide sequences, and may be altered (e.g., by the insertion, deletion or substitution of nucleotides) so long as the sequences provide for functional rescue, replication and packaging.

[0089] Alternatively, other systems such as lentiviral vectors can be used in embodiments of the invention. Lentiviral-based systems can transduce non-dividing as well as dividing cells making them useful for applications targeting, for examples, the non-dividing cells of the CNS. Lentiviral vectors are derived from the human immunodeficiency virus and, like that virus, integrate into the host genome providing the potential for long-term gene expression.

[0090] Polynucleotides, including plasmids, YACs, minichromosomes and minicircles, carrying the target gene containing the gene regulation cassette can also be introduced into a cell or organism by nonviral vector systems using, for example, cationic lipids, polymers, or both as carriers. Conjugated poly-L-lysine (PLL) polymer and polyethylenimine (PEI) polymer systems can also be used to deliver the vector to cells. Other methods for delivering the vector to cells includes hydrodynamic injection and electroporation and use of ultrasound, both for cell culture and for organisms. For a review of viral and non-viral delivery systems for gene delivery see Nayerossadat, N. et al. (Adv Biomed Res. 2012; 1:27) incorporated herein by reference.

[0091] Methods of Modulating Expression of a Target Gene

[0092] In one aspect, this invention provides a method of modulating expression of a target gene (e.g., a therapeutic gene), by (a) inserting the gene regulation polynucleotide cassette of the present invention into a target gene; (b) introducing the target gene comprising the gene regulation cassette into a cell; and (c) exposing the cell to a ligand that binds the aptamer in an effective amount to modulate target gene expression. In one embodiment, the polynucleotide cassette comprises an off riboswitch and exposure to the aptamer ligand reduces target gene expression. In one embodiment, the polynucleotide cassette comprises an on riboswitch and exposure to the aptamer ligand increases target gene expression. In one embodiment, the ligand is a small molecule. In aspects, expression of the target gene in

target cells confers a desired property to a cell into which it was introduced, or otherwise leads to a desired therapeutic outcome.

[0093] In one embodiment, one or more gene regulation cassettes are inserted into the 3' UTR and/or the 5' UTR of the target gene. In one embodiment, a single gene regulation cassette is inserted into the 3' UTR of a target gene. In other embodiments 2, 3, 4, or more gene regulation cassettes are inserted in the target gene. In one embodiment, two gene regulation cassettes are inserted into the target gene. When multiple gene regulation cassettes are inserted into a target gene, they each can contain the same aptamer such that a single ligand can be used to modulate ribonuclease cleavage of the multiple cassettes and thereby modulate target gene expression. In other embodiments, multiple gene regulation cassettes are inserted into a target gene, each can contain a different aptamer so that exposure to multiple different small molecule ligands modulates target gene expression. In other embodiments, multiple gene regulation cassettes are inserted into a target gene, each containing different ribonuclease substrate sequences. This may be useful in reducing recombination and improving ease of incorporation into viral vectors.

[0094] In one embodiment, a polynucleotide cassette of the present invention is effective at modulating target gene expression when placed downstream (i.e., 3') of the target gene start codon. In one embodiment, when the polynucleotide cassette is inserted downstream of the translation start codon of the target gene, the on riboswitch is adjacent to the start codon. In other embodiments, the riboswitch is about 1 to about 100, about 1 to about 50, or about 1 to about 20 nucleotides from the start codon of the target gene. In one embodiment, the polynucleotide cassette comprises a 2A peptide sequence between the 3' end of the riboswitch and the target gene coding sequence. In one embodiment, the polynucleotide cassette comprises a 2A peptide sequence and a linker ("2A linker") between the 3' end of the riboswitch and the target gene coding sequence. The 2A sequence used in the Examples was identified from *Thoseaasigna* virus, therefore it is sometimes called "T2A." In place of T2A, other sequences that separate leader poly peptides from the remaining gene product may be used. Other 2A peptides include those identified in Picornaviruses: foot-and-mouth disease virus (F2A, the first one discovered), equine rhinitis A virus (E2A), and porcine teschovirus (P2A). Other 2A peptides include the 2A peptide from type C rotaviruses.

[0095] The polynucleotide cassette of the present invention can be used in combination with other mechanisms for the regulation of expression of the target gene. In

one embodiment, a polynucleotide cassette of the present invention is used in combination with a gene regulation cassette that modulates target gene expression by aptamer-mediated regulation of alternative splicing as described in WO 2016/126747, incorporated herein by reference.

[0096] Methods of Treatment and Pharmaceutical Compositions

[0097] One aspect of the invention provides a method of regulating the level of a therapeutic protein delivered by gene therapy. In this embodiment, the “target gene” may encode the therapeutic protein. The “target gene” may encode a protein that is endogenous or exogenous to the cell.

[0098] The therapeutic gene sequence containing the regulatory cassette with aptamer-driven riboswitch is delivered to target cells in vitro or ex vivo, e.g., by a vector. The cell specificity of the “target gene” may be controlled by promoter or other elements within the vector. Delivery of the vector construct containing the target gene and the polynucleotide cassette, and the transfection of the target tissues resulting in stable transfection of the regulated target gene, is the first step in producing the therapeutic protein.

[0099] However, due to the presence of the regulatory cassette within the target gene sequence, the target gene is not expressed at significant levels. In some embodiments, the target gene is in the “off state” in the absence of the specific ligand that binds to the aptamer contained within in the regulatory cassette riboswitch. Only when the aptamer specific ligand is administered (or otherwise present in sufficient quantities) is the target gene expression activated. In other embodiments, the target gene is in the “off state” in the presence of the specific ligand that binds to the aptamer contained within in the regulatory cassette riboswitch. When administration stops (or the ligand is otherwise present at sufficiently low levels), target gene expression will occur.

[0100] The delivery of the vector construct containing the target gene and the delivery of the activating ligand generally are separated in time. The delivery of the activating ligand will control when the target gene is expressed, as well as the level of protein expression. The ligand may be delivered by a number of routes including, but not limited to, oral, intramuscular (IM), intravenous (IV), intraocular, or topically.

[0101] The timing of delivery of the ligand will depend on the requirement for activation of the target gene. For example, if the therapeutic protein encoded by the target gene is required constantly, in the case of the “on” riboswitches, an oral small molecule

ligand may be delivered daily, or multiple times a day, to ensure continual activation of the target gene, and thus continual expression of the therapeutic protein. If the target gene has a long acting effect, the inducing ligand may be dosed less frequently.

[0102] This invention allows the expression of the therapeutic transgene to be controlled temporally, in a manner determined by the temporal dosing of the ligand specific to the aptamer within the riboswitch of the regulatory polynucleotide cassette. The increased expression of the therapeutic transgene only on ligand administration (in the case of the “on” riboswitches), increases the safety of a gene therapy treatment by allowing the target gene to be off in the absence of the ligand.

[0103] Different aptamers can be used to allow different ligands to activate target genes or repress target genes. In certain embodiments of the invention, each therapeutic gene containing a regulatory cassette will have a specific aptamer within the cassette that will be activated by a specific small molecule. This means that each therapeutic gene can be activated only by the ligand specific to the aptamer housed within it. In these embodiments, each ligand will only activate one therapeutic gene. This allows for the possibility that several different “target genes” may be delivered to one individual and each will be activated on delivery of the specific ligand for the aptamer contained within the regulatory cassette housed in each target gene.

[0104] This invention allows any therapeutic protein whose gene can be delivered to the body (such as erythropoietin (EPO) or a therapeutic antibody) to be produced by the body when the activating ligand is delivered. This method of therapeutic protein delivery may replace the manufacture of such therapeutic proteins outside of the body which are then injected or infused, e.g., antibodies used in cancer or to block inflammatory or autoimmune disease. The body containing the regulated target gene becomes the biologics manufacturing factory, which is switched on when the gene-specific ligand is administered.

[0105] Dosing levels and timing of dosing of a therapeutic protein may be important to therapeutic effect. For example, in the delivery of AVASTIN (anti-VEGF antibody) for cancer. The present invention increases the ease of dosing in response to monitoring for therapeutic protein levels and effects.

[0106] In one embodiment, the target gene may encode a nuclease that can target and edit a particular DNA sequence. Such nucleases include Cas9, zinc finger containing nucleases, or TALENs. In the case of these nucleases, the nuclease protein may be required

for only a short period of time that is sufficient to edit the target endogenous genes. However, if an unregulated nuclease gene is delivered to the body, this protein may be present for the rest of the life of the cell. In the case of nucleases, there is an increasing risk of off-target editing the longer the nuclease is present. Regulation of expression of such proteins has a significant safety advantage. In this case, vector containing the nuclease target gene containing a regulatory cassette could be delivered to the appropriate cells in the body. The target gene is in the “off” state in the absence of the cassette-specific ligand, so no nuclease is produced. Only when the activating ligand is administered, is the nuclease produced. When sufficient time has elapsed allowing sufficient editing to occur, the ligand will be withdrawn and not administered again. Thus, the nuclease gene is thereafter in the “off” state and no further nuclease is produced and editing stops. This approach may be used to correct genetic conditions, including a number of inherited retinopathies such as LCA10 caused by mutations in CEP290 and Stargardt’s Disease caused by mutations in ABCA4.

[0107] Administration of a regulated target gene encoding a therapeutic protein which is activated only on specific ligand administration may be used to regulate therapeutic genes to treat many different types of diseases, e.g., cancer with therapeutic antibodies, immune disorders with immune modulatory proteins or antibodies, metabolic diseases, rare diseases such as PNH with anti-C5 antibodies or antibody fragments as the regulated gene, or ocular angiogenesis with therapeutic antibodies, and dry AMD with immune modulatory proteins.

[0108] A wide variety of specific target genes, allowing for the treatment of a wide variety of specific diseases and conditions, are suitable for use in the present invention. For example, insulin or an insulin analog (preferably human insulin or an analog of human insulin) may be used as the target gene to treat type I diabetes, type II diabetes, or metabolic syndrome; human growth hormone may be used as the target gene to treat children with growth disorders or growth hormone-deficient adults; erythropoietin (preferably human erythropoietin) may be used as the target gene to treat anemia due to chronic kidney disease, anemia due to myelodysplasia, or anemia due to cancer chemotherapy.

[0109] The present invention may be especially suitable for treating diseases caused by single gene defects such as cystic fibrosis, hemophilia, muscular dystrophy, thalassemia, or sickle cell anemia. Thus, human β -, γ -, δ -, or ζ -globin may be used as the target gene to treat β -thalassemia or sickle cell anemia; human Factor VIII or Factor IX may be used as the target gene to treat hemophilia A or hemophilia B.

[0110] The ligands used in the present invention are generally combined with one or more pharmaceutically acceptable carriers to form pharmaceutical compositions suitable for administration to a patient. Pharmaceutically acceptable carriers include solvents, binders, diluents, disintegrants, lubricants, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, generally used in the pharmaceutical arts. Pharmaceutical compositions may be in the form of tablets, pills, capsules, troches, and the like, and are formulated to be compatible with their intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, intranasal, subcutaneous, oral, inhalation, transdermal (topical), transmucosal, and rectal.

[0111] The pharmaceutical compositions comprising ligands are administered to a patient in a dosing schedule such that an amount of ligand sufficient to desirably regulate the target gene is delivered to the patient. When the ligand is a small molecule and the dosage form is a tablet, capsule, or the like, preferably the pharmaceutical composition comprises from 0.1 mg to 10 g of ligand; from 0.5 mg to 5 g of ligand; from 1 mg to 1 g of ligand; from 2 mg to 750 mg of ligand; from 5 mg to 500 mg of ligand; or from 10 mg to 250 mg of ligand.

[0112] The pharmaceutical compositions may be dosed once per day or multiple times per day (e.g., 2, 3, 4, 5, or more times per day). Alternatively, pharmaceutical compositions may be dosed less often than once per day, e.g., once every 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, or 14 days, or once a month or once every few months. In some embodiments of the invention, the pharmaceutical compositions may be administered to a patient only a small number of times, e.g., once, twice, three times, etc.

[0113] The present invention provides a method of treating a patient in need of increased expression of a therapeutic protein encoded by a target gene, the method comprising administering to the patient a pharmaceutical composition comprising a ligand for an aptamer, where the patient previously had been administered a recombinant DNA comprising the target gene, where the target gene contains a gene regulation cassette of the present invention that provides the ability to regulate expression of the target gene by the ligand of the aptamer by alternative splicing of pre-mRNA of the target gene, thereby increasing expression of the therapeutic protein.

[0114] Articles of Manufacture and Kits

[0115] Also provided are kits or articles of manufacture for use in the methods described herein. In aspects, the kits comprise the compositions described herein (e.g., for compositions for delivery of a vector comprising the target gene containing the gene regulation cassette) in suitable packaging. Suitable packaging for compositions (such as ocular compositions for injection) described herein are known in the art, and include, for example, vials (such as sealed vials), vessels, ampules, bottles, jars, flexible packaging (e.g., sealed Mylar or plastic bags), and the like. These articles of manufacture may further be sterilized and/or sealed.

[0116] The present invention also provides kits comprising compositions described herein and may further comprise instruction(s) on methods of using the composition, such as uses described herein. The kits described herein may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for performing the administration, including e.g., any methods described herein. For example, in some embodiments, the kit comprises rAAV for expression of the target gene comprising the gene regulation cassette of the present invention, a pharmaceutically acceptable carrier suitable for injection, and one or more of: a buffer, a diluent, a filter, a needle, a syringe, and a package insert with instructions for performing the injections. In some embodiments, the kit is suitable for intraocular injection, intramuscular injection, intravenous injection and the like.

[0117] “Homology” and “homologous” as used herein refer to the percent of identity between two polynucleotide sequences or between two polypeptide sequences. The correspondence between one sequence to another can be determined by techniques known in the art. For example, homology can be determined by a direct comparison of two polypeptide molecules by aligning the sequence information and using readily available computer programs. Two polynucleotide or two polypeptide sequences are “substantially homologous” to each other when, after optimally aligned with appropriate insertions or deletions, at least about 80%, at least about 85%, at least about 90%, and at least about 95% of the nucleotides or amino acids, respectively, match over a defined length of the molecules, as determined using the methods above.

[0118] “Percent sequence identity” with respect to a reference polypeptide or nucleic acid sequence is defined as the percentage of amino acid residues or nucleotides in a candidate sequence that are identical with the amino acid residues or nucleotides in the reference polypeptide or nucleic acid sequence, after aligning the sequences and introducing

gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent amino acid or nucleic acid sequence identity can be achieved in ways known to the ordinarily-skilled artisan, for example, using publicly available computer software programs including BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software.

[0119] The term “polynucleotide” or “nucleic acid” as used herein refers to a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. Thus, this term includes, but is not limited to, single-, double- or multi- stranded DNA or RNA, genomic DNA, cDNA, DNA-RNA hybrids, or a polymer comprising purine and pyrimidine bases, or other natural, chemically or biochemically modified, non-natural, or derivatized nucleotide bases.

[0120] “Heterologous” or “exogenous” means derived from a genotypically distinct entity from that of the rest of the entity to which it is compared or into which it is introduced or incorporated. For example, a polynucleotide introduced by genetic engineering techniques into a different cell type is a heterologous polynucleotide (and, when expressed, can encode a heterologous polypeptide). Similarly, a cellular sequence (e.g., a gene or portion thereof) that is incorporated into a viral vector is a heterologous nucleotide sequence with respect to the vector.

[0121] The below table contains a listing of the DNA sequences of the constructs described herein as well as other sequences as described. The ribozyme sequence is in underlined, italicized lower case letters; the additional 3 base pair stem (3HP) is in double underlined lower case letters; the aptamer sequences are in wave underlined lower case letters; the stem sequences shared between twister stem and aptamer stem are grey shaded lower case letters; and the linker2A sequence is in thick underlined lower case letters; the coding sequences are in upper case letters.

SEQ ID NO.:	Description	Sequence
1	wasp-twister in pEGFP-C1	<i>GTGACCGCCGCGGGATCACTCTCGGCATGGACGAGC</i> <i>TGTACAAGTAA</i> <u><i>accgcggcttgtaatgcggccgigttaataatttacgctg</i></u> <u><i>gtctcaagcccataaacgcagagagcaaggcgccgccataccacattt</i></u> <u><i>gtaga</i></u> <i>ggttttacttgcttaaaaaacctcccacacctcccctgaacctgaaacataaaatgaat</i> <i>gc</i>
2	es-twister in pEGFPC1	<i>GTGACCGCCGCGGGATCACTCTCGGCATGGACGAG</i> <i>CTGTACAAGTAA</i> <u><i>accgcggcaataaagcggttacaagcccgcaaaaat</i></u> <u><i>agcagagtaatgtcgcgatagcgcggcattaatgcagctttattggcgccgccata</i></u>

		ccacatttgtagaggtttacttgcttataaaaacctcccacacctccccctgaacctgaa acataaaatgaatgc
3	wasp-twister-mut in pEGFP-C1	GTGACCGCCGCGGGATCACTCTCGGCATGGACGAG CTGTACAAGTAAaccgcggttgatgcgccgtgtaataatttacac <u>gtcggctcaagcccgataaacgcagagagcaag</u> GCGGCCGCATAC CACATTTGTAGAGGTTTTACTTGCTTTAAAAAACCTC CCACACCTCCCCCTGAACCTGAAACATAAAATGAAT GC
4	wasp_no Sac 0HP in pEGFP-C1	GTGACCGCCGCGGGATCACTCTCGGCATGGACGAG CTGTACAAGTAAacttgtaatgcgccgtgtaataatttacacgtcggctc <u>caagcccgataaacgcagagagcaag</u> gcggccgccataccacatttgtagaggtt tacttgcttataaaaacctcccacacctccccctgaacctgaaacataaaatgaatgc
5	wasp_no Sac 3HP in pEGFP-C1	GTGACCGCCGCGGGATCACTCTCGGCATGGACGAG CTGTACAAGTAAaccgcttgtaatgcgccgtgtaataatttacacgtc <u>gtcgaagcccgataaacgcagagagcaag</u> gcggccgccataccacatttgtaga ggttttacttgcttataaaaacctcccacacctccccctgaacctgaaacataaaatgaat gc
6	wasp_No Sac 0HP in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAActtgtaatgcgccgtgtaataatttacacgt <u>cggtctcaagcccgataaacgcagagagcaag</u> gcggccgccataccacatttgta gaggtttacttgcttataaaaacctcccacacctccccctgaacctgaaacataaaatg aatg
7	wasp_No Sac 3HP in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAaccgcttgtaatgcgccgtgtaataatttacac <u>gtcggctcaagcccgataaacgcagagagcaag</u> gcggccgccataccacattt gtagaggtttacttgcttataaaaacctcccacacctccccctgaacctgaaacataaa atgaatgc
8	waspGua1 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAaccgcttgtaatgcgccgtgtaataactcatat <u>aatcgcgtggatatggcacgcaagtttctaccgggcaccgtaaatgtccgactatgggt</u> <u>ttacacgtcggctcaagcccgataaacgcagagagcaag</u> gcggccgccata ccacatttgtagaggtttacttgcttataaaaacctcccacacctccccctgaacctgaa acataaaatgaatgc
9	waspGua2 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAaccgcttgtaatgcgccgtgtaaaactcatata <u>atcgcgtggatatggcacgcaagtttctaccgggcaccgtaaatgtccgactatgggt</u> <u>ttacacgtcggctcaagcccgataaacgcagagagcaag</u> gcggccgccatacca catttgtagaggtttacttgcttataaaaacctcccacacctccccctgaacctgaaaca taaaatgaatgc
10	waspGua3 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAaccgcttgtaatgcgccgtgtaactcatataatc <u>gcgtggatatggcacgcaagtttctaccgggcaccgtaaatgtccgactatgggttac</u> <u>acgtcggctcaagcccgataaacgcagagagcaag</u> gcggccgccataccacatt ttagaggtttacttgcttataaaaacctcccacacctccccctgaacctgaaacataaa atgaatgc
11	waspGua4 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAaccgcttgtaatgcgccgtgatcatataatcgc <u>gtggatatggcacgcaagtttctaccgggcaccgtaaatgtccgactatggtacacgtc</u> <u>ggctcaagcccgataaacgcagagagcaag</u> gcggccgccataccacatttgtag aggtttacttgcttataaaaacctcccacacctccccctgaacctgaaacataaaatga

		atgc
12	waspGua5 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggccgtg</u> tcataataatcgctg tggatatggcacgcaagtttctaccgggcaccgtaaatgtccgactat <u>gcacacgctc</u> <u>gtctcaagcccgataaacgcagagagcaaggcggcc</u> gccataccacattttaga ggttttacttgcTTAAAAAacctcccacacctccccctgaacctgaaacataaaatgaat gc
13	waspGua6 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggccgtg</u> catataatcgctg ggatatggcacgcaagtttctaccgggcaccgtaaatgtccgactat <u>gcacacgctc</u> <u>ctcaagcccgataaacgcagagagcaaggcggcc</u> gccataccacattttagaggg tttacttgcTTAAAAAacctcccacacctccccctgaacctgaaacataaaatgaatgc
14	waspGua7 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggccgtg</u> catataatcgctg ggatatggcacgcaagtttctaccgggcaccgtaaatgtccgactat <u>gcacacgctc</u> <u>caagcccgataaacgcagagagcaaggcggcc</u> gccataccacattttagaggttt tacttgcTTAAAAAacctcccacacctccccctgaacctgaaacataaaatgaatgc
15	waspGua8 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggccgtg</u> atataatcgctg gatatggcacgcaagtttctaccgggcaccgtaaatgtccgactat <u>cacgctcgtctca</u> <u>agcccgataaacgcagagagcaaggcggcc</u> gccataccacattttagaggtttta cttgcTTAAAAAacctcccacacctccccctgaacctgaaacataaaatgaatgc
16	waspGua9 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggccgtg</u> atataatcgctgg atatggcacgcaagtttctaccgggcaccgtaaatgtccgactat <u>acgctcgtctcaag</u> <u>cccgataaacgcagagagcaaggcggcc</u> gccataccacattttagaggttttactt gctTTAAAAAacctcccacacctccccctgaacctgaaacataaaatgaatgc
17	waspGua12 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggccgata</u> atcgctggata tggcacgcaagtttctaccgggcaccgtaaatgtccgact <u>gctcgtctcaagcccga</u> <u>taaacgcagagagcaaggcggcc</u> gccataccacattttagaggttttacttgcTTA AAAAAacctcccacacctccccctgaacctgaaacataaaatgaatgc
18	waspGua13 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggccata</u> atcgctgggat ggcacgcaagtttctaccgggcaccgtaaatgtccgact <u>gctcgtctcaagcccgata</u> <u>aacgcagagagcaaggcggcc</u> gccataccacattttagaggttttacttgcTTAAA Aacctcccacacctccccctgaacctgaaacataaaatgaatgc
19	waspGua14 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggccata</u> atcgctgggat gcacgcaagtttctaccgggcaccgtaaatgtccgact <u>gctcgtctcaagcccgataa</u> <u>acgcagagagcaaggcggcc</u> gccataccacattttagaggttttacttgcTTAAAA Aacctcccacacctccccctgaacctgaaacataaaatgaatgc
20	waspGua15 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggccata</u> atcgctgggat atggcacgcaagtttctaccgggcaccgtaaatgtccgactat <u>gctcgtctcaagccc</u> <u>gataaacgcagagagcaaggcggcc</u> gccataccacattttagaggttttacttgcTT AAAAAacctcccacacctccccctgaacctgaaacataaaatgaatgc
21	waspGua16 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggccata</u> atcgctgggat

		<u>ggcacgcaagtttctaccgggcaccgtaaagtccgactatcgggtcaagcccgat</u> <u>aaacgcagagagcaaggcggccgccataccacattttagaggtttacttgctttaa</u> <u>aaaacctcccacacctccccctgaacctgaaacataaaatgaatgc</u>
22	waspGua17 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggc</u> tataatcgcgtggatg gcacgcaagtttctaccgggcaccgtaaagtccgactatcgggtcaagcccgataa <u>acgcagagagcaaggcggccgccataccacattttagaggtttacttgcttataaa</u> <u>aacctcccacacctccccctgaacctgaaacataaaatgaatgc</u>
23	waspGua18 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggc</u> gtataatcgcgtgg atatggcacgcaagtttctaccgggcaccgtaaagtccgactacacgctggctcaa <u>gcccgataaacgcagagagcaaggcggccgccataccacattttagaggtttact</u> <u>tgcttataaaaacctcccacacctccccctgaacctgaaacataaaatgaatgc</u>
24	waspGua19 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggc</u> gtataatcgcgtggat atggcacgcaagtttctaccgggcaccgtaaagtccgactacacgctggctcaa <u>cgataaacgcagagagcaaggcggccgccataccacattttagaggtttactgc</u> <u>ttataaaaacctcccacacctccccctgaacctgaaacataaaatgaatgc</u>
25	waspGua20 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggc</u> gtataatcgcgtggat atggcacgcaagtttctaccgggcaccgtaaagtccgactacacgctggctcaa <u>gataaacgcagagagcaaggcggccgccataccacattttagaggtttactgctt</u> <u>tataaaaacctcccacacctccccctgaacctgaaacataaaatgaatgc</u>
26	Epo-waspGua16	GCTGTACACGGGAGAGGTCTGCAGGAGAGGGGACA GGTGAggatccac <u>gccttgatgcggc</u> atataatcgcgtggatggcacgcaa gtttctaccgggcaccgtaaagtccgactatcgggtcaagcccgataaacgcaga <u>gagcaaggcgtc</u> gaggcccagatctaattcaccaccagtgaggtgcctatc agaaagtgggtgggtggctaatgcctggcccacaagtataactaagctcgttt cttgctgccaattctattaaagggttcctt
27	Epo-mutwaspGua16	GCTGTACACGGGAGAGGTCTGCAGGAGAGGGGACA GGTGAggatccac <u>gccttgatgcggc</u> atataatcgcgtggatggcacgcaa gtttctaccgggcaccgtaaagtccgactatcgggtcaagcccgataaacgcaga <u>gagcaaggcgtc</u> gaggcccagatctaattcaccaccagtgaggtgcctatc agaaagtgggtgggtggctaatgcctggcccacaagtataactaagctcgttt cttgctgccaattctattaaagggttcctt
28	twister_theo_1 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggc</u> atgataaccagccgaaa ggcccttggcagc <u>atcgggtcaagcccgataaacgcagagagcaaggcggcc</u> ccataccacattttagaggtttacttgcttataaaaacctcccacacctccccctgaac ctgaaacataaaatgaatgc
29	twister_theo_2 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggc</u> gataaccagccgaa aggcccttggcagc <u>gtcgggtcaagcccgataaacgcagagagcaaggcggcc</u> cgccataccacattttagaggtttacttgcttataaaaacctcccacacctccccctga acctgaaacataaaatgaatgc
30	twister_theo_3 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAA <u>acgccttgtaatgcggc</u> gataaccagccgaaa ggcccttggcagc <u>gtcgggtcaagcccgataaacgcagagagcaaggcggcc</u> gccataccacattttagaggtttacttgcttataaaaacctcccacacctccccctgaa cctgaaacataaaatgaatgc

31	twister_theo_4 in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAAC <u>gccttgtaatgcggcct</u> gataaccagccgaaa <u>ggcccttggcagcagtcggtctcaagccc</u> <u>gataaacgcagagagcaaggcggcc</u> gceataccacattttagaggttttacttcttataaaaacctcccacacctccccctgaa cctgaaacataaaatgaatgc
32	ATG_wtWaspGua16 in pLuc	ATGAc <u>gccttgtaatgcggc</u> catataatcgcgtggatgacgcaagtttaccg ggcaccgtaaatgtccgactat <u>tcggtctcaagccc</u> <u>gataaacgcagagagcaagg</u> <u>cgaagcggatcttcaggtatggc</u> <u>gagggcagaggaagtctt</u> <u>taacatgcggtga</u> <u>cgtgaggagaatccc</u> <u>gcccc</u> GAAGACGCCAAAAACATAAAG AAAGGCCCGGCGCCA
33	ATG_mutWaspGua16 in pLuc	ATGAc <u>gccttgatgcggc</u> catataatcgcgtggatgacgcaagtttaccg ggcaccgtaaatgtccgactat <u>tcggtctcaagccc</u> <u>gataaacgcagagagcaagg</u> <u>cgaagcggatcttcaggtatggc</u> <u>gagggcagaggaagtctt</u> <u>taacatgcggtga</u> <u>cgtgaggagaatccc</u> <u>gcccc</u> GAAGACGCCAAAAACATAAAG AAAGGCCCGGCGCCA
34	10aa_wtWaspGua16 in pLuc	ATGGTGAGCAAGGGCGAGGAGCTGTTACC <u>gccttgta</u> <u>atgcggc</u> catataatcgcgtggatggcagcaagtttaccgggaccgtaaatgtc <u>cgactatcggctcaagccc</u> <u>gataaacgcagagagcaaggc</u> <u>gaa</u> <u>gcggatctc</u> <u>aggtatggc</u> <u>gagggcagaggaagtctt</u> <u>taacatgcggtgac</u> <u>gtggaggagaatc</u> <u>ccggcccc</u> GAAGACGCCAAAAACATAAAGAAAGGCCCG GCGCCA
35	10aa_mutWaspGua16 in pLuc	ATGGTGAGCAAGGGCGAGGAGCTGTTACC <u>gccttgat</u> <u>atgcggc</u> catataatcgcgtggatggcagcaagtttaccgggaccgtaaatgtc <u>cgactatcggctcaagccc</u> <u>gataaacgcagagagcaaggc</u> <u>gaa</u> <u>gcggatctc</u> <u>aggtatggc</u> <u>gagggcagaggaagtctt</u> <u>taacatgcggtgac</u> <u>gtggaggagaatc</u> <u>ccggcccc</u> GAAGACGCCAAAAACATAAAGAAAGGCCCG GCGCCA
36	GT_8bp in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAATA <u>aaaggc</u> gtataatcgcgtggatggcagcaa gtttaccgggaccgtaaatgtccgactac <u>gccttgtaatgcggcgtgtaataat</u> <u>ttacagtcggtctcaagccc</u> <u>gataaacgcagagagcaaggcggcc</u> <u>gccatacca</u> cattttagaggttttacttcttataaaaacctcccacacctccccctgaacctgaaaca taaatgaatgc
37	TG_9b in pLuc	TCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGA AAGATCGCCGTGTAAC <u>gccttgtaatgcggc</u> ggtgtaataatttaca <u>cgtcggtctcaagccc</u> <u>gataaacgcagagagcaaggc</u> gtataatcgcgtggatg gcacgcaagtttaccgggaccgtaaatgtccgactac <u>gccttg</u> ataaccatttga gaggtttacttcttataaaaacctcccacacctccccctgaacctgaaacataaatg aatgc
38	Twister ribozyme is from <i>Nasonia vitripennis</i> (wasp twister)	cttgaatgcgccgtgtaataatttacacgtcggctcaagcccgataaacgcagag agcaag
39	Twister ribozyme is from an environmental sample (es twister)	ggcaataaagcggttacaagcccgcaaaaatagcagagtaatgtcgcgatagcgcg gcattaatgcagctttattg

[0122] It is to be understood and expected that variations in the principles of invention herein disclosed can be made by one skilled in the art and it is intended that such modifications are to be included within the scope of the present invention. The following Examples further illustrate the invention, but should not be construed to limit the scope of the invention in any way. All references cited herein are hereby incorporated by reference in their entirety.

EXAMPLE 1

[0123] Twister ribozymes self-cleave and suppress gene expression in mammalian cells

[0124] Experimental procedures:

[0125] Plasmid constructs: oligonucleotides containing wild type or mutant twister ribozyme sequences, either from *Nasonia Vitripennis* (wasp) or from environmental samples, flanked by BsrGI and MfeI sites, were synthesized (IDT). The synthesized oligonucleotides (“oligos”) were digested with BsrGI and MfeI and cloned into pEGFP-C1 (Clontech) digested with the compatible restriction enzymes. The sequences of the constructed vectors were verified by DNA sequencing (Genewiz).

[0126] Transfection and measurement of GFP fluorescence: 3.5×10^4 HEK 293 cells were plated in a 96-well flat bottom plate the day before transfection. Plasmid DNA (500 ng) was added to a tube or a 96-well U-bottom plate. Separately, TransIT-293 reagent (Mirus; 1.4 μ l) was added to 50 μ l Opti-mem I media (Life Technologies), and allowed to sit for 5 minutes at room temperature (“RT”). Then, 50 μ l of this diluted transfection reagent was added to the DNA, mixed, and incubated at RT for 20 min. Finally, 7 μ l of this solution was added to a well of cells in a 96-well plate. A LacZ construct that does not express GFP was used as background negative control. GFP fluorescence intensity was measured by Tecan plate reader, using Excitation wavelength at 484 nm, Emission wavelength at 510 nm and Excitation bandwidth at 5 nm. The GFP fluorescence intensity was expressed as the value generated by GFP constructs subtracted by the value generated by LacZ construct.

[0127] Results:

[0128] Twister ribozymes exist in many species of bacteria and eukarya. However, their self-cleaving activity in mammalian cells has not been determined. To investigate whether twister ribozyme could cleave mRNA and suppress gene expression in mammalian

cells, the twister ribozyme sequence from *N. Vitripennis* (wasp) and twister ribozyme sequence from environmental samples was each inserted into the 3' UTRs of the EGFP gene in the pEGFP-C1 vector. As shown in Figure 1a, insertion of wasp-twister ribozyme in the 3' UTR resulted in 97% and 36-fold inhibition of GFP expression when comparing with the pEGFP-C1 control construct. The twister sequence from environmental sample (es-twister) did not lead to as robust suppression of GFP expression as wasp-twister, generating about 50% suppression of GFP expression when comparing with the control GFP vector. To further determine that the suppression of GFP expression was due to the twister ribozyme cleavage, a mutant construct was generated where A6 was replaced with G, making twister ribozyme cleavage-incapable. This A6G mutation completely abolished the suppression in GFP expression, while the wild type twister ribozyme caused 44-fold reduction, as shown in Figure 1b. These results indicate that in mammalian cells twister ribozymes self-cleave when present in the target gene RNA and thereby suppress target gene expression.

EXAMPLE 2

[0129] Modification of twister ribozyme P1 stem sequence improves twister ribozyme cleavage activity

[0130] Experimental procedures:

[0131] Plasmid constructs: The following primers were used to generate constructs GFP_wasp_No Sac 0HP or GFP_wasp_No Sac 3HP: forward primers NoSac_Fwd (5' GCTGTACAAGTAAACTTGTAATGCGGCCGTGTAAATAATTTAC) and NoSac3HP_Fwd (5' AGCTGTACAAGTAAACGCCTTGTAATGCGGCCGTGTAAATAATTTAC) and reverse primer MfeI Rev (5' ACAACAACAATTGCATTCATTTTATG). GFP_wasp-twister vector was used as template and the PCR-amplified DNA fragments were digested with BsrGI and MfeI and cloned into pEGFP-C1 digested with BsrGI and MfeI enzymes. The EGFP gene in pEGFP-C1 was replaced with the firefly luciferase gene to generate the pFLuc vector using Gibson cloning strategy and cloning kit (NEB). The same strategy was used to generate constructs Luc_wasp_No Sac 0HP and Luc_wasp_No Sac 3 HP by replacing the EGFP gene in constructs GFP_wasp_no Sac 0HP and GFP_wasp_no Sac 3HP with the firefly luciferase gene.

[0132] Measurement of GFP fluorescence intensity was performed as described in Example 1.

[0133] Firefly luciferase assay of cultured cells: 24 hours after media change, plates were removed from the incubator, equilibrated to RT for several minutes on a lab bench, and then aspirated. Glo-lysis buffer (Promega, 100 μ L, RT) was added, and the plates allowed to remain at RT for at least 5 minutes. The well contents were mixed by 50 μ L trituration, and 20 μ L of each sample was mixed with 20 μ L of bright-glo reagent (Promega) that had been diluted to 10% in glo-lysis buffer. 96 wells were spaced on an opaque white 384-well plate. Following a 5-minute incubation at RT, luminescence was measured using a Tecan machine with 500 mSec read time. The luciferase activity was expressed as mean relative light unit (RLU) \pm S.D.

[0134] Results:

[0135] To further enhance the cleavage activity of the twister ribozyme in mammalian cells, and thereby increase the dynamic range of suppression of target gene expression by the ribozyme, the sequences adjacent to the twister P1 stem were modified in the GFP-twister construct. First, the SacII sequence located at the 5' end of the twister sequence was deleted, generating GFP_wasp_no Sac 0HP. To determine whether a longer P1 stem could stabilize the twister ribozyme and thus enhance its cleavage activity, the sequence of SacII site was replaced with CGC to create a 3 base pair stem (3HP) at the base of the P1 stem of the twister ribozyme (Fig. 2a), generating construct GFP_wasp_no Sac 3HP. Interestingly, as shown in Fig. 2b, deletion of SacII sequence increased the GFP suppression, from 37-fold by the construct with the SacII sequence to 68-fold by the construct GFP_wasp_no Sac 0HP without the SacII site, suggesting the SacII sequence adjacent to the 5' of the P1 sequence impaired the ribozyme activity. Addition of a 3 bp stem to the base of the P1 stem further enhanced the ribozyme activity, generating 104-fold suppression of GFP expression. The activity of this modified twister construct was also tested using firefly luciferase gene as a reporter. As shown in Fig. 2c, deletion of SacII sequence resulted in 101-fold reduction of luciferase activity when comparing to pFLuc control construct. Addition of a 3bp stem at the P1 stem further enhanced the reduction of luciferase activity to 254 fold compared to the control construct.

EXAMPLE 3

[0136] Use of xpt-guanine aptamer to regulate target gene expression via modulating twister ribozyme

[0137] Experimental procedures:

[0138] Oligos containing the twister ribozyme sequence linked with an xpt-guanine aptamer were synthesized (IDT), and cloned into the pLuc vector using the Gibson cloning strategy and kit (NEB). Construct sequences were verified by DNA sequencing (Genewiz).

[0139] Transfection and aptamer ligand treatment: HEK 293 cells were transfected as described in Example 1. Four hours after transfection, the media was aspirated, and new media with or without 500 μ M guanine or 1 mM guanosine (Sigma) was added. Luciferase assay was performed 20 to 24 hours after guanine or guanosine treatment as described in Example 2. The reduction fold was expressed as the quotient of luciferase activity obtained in the absence of aptamer ligand divided by the value obtained in the presence of the aptamer ligand.

[0140] ELISA assay for measuring mouse Epo in culture media: Supernatant was collected 18 hours after drug treatment, and an ELISA was performed using ELISA kit (R&D) according to manufacturer's instructions. The data was analyzed using 4-parameter curve fitting.

[0141] Results:

[0142] In order to regulate the cleavage activity of twister ribozyme, and thereby regulate target gene expression, aptamer sequence was linked to the P3 stem of the twister ribozyme. When aptamer ligand is not present, the structure of the ribozyme is disrupted, preventing cleavage, and allowing target gene expression (Fig. 3a, top panel). Addition of the aptamer ligand leads to a conformational change allowing twister ribozyme cleavage and reducing target gene expression (Fig. 3a, bottom panel). The P1 stem of the xpt-guanine aptamer was grafted to the P3 stem of twister ribozyme to generate guanine-responsive twister ribozyme. In this configuration (Fig. 3a), in the absence of aptamer ligand, insertion of aptamer sequence between the two arms of the twister P3 stem disrupts the P3 stem and the twister ribozyme structure, disrupting cleavage activity and allowing the target gene to be expressed. In the presence of aptamer ligand, aptamer/ligand binding brings together the arms of the stem, facilitating the P3 stem formation and twister cleavage activity, thereby suppressing target gene expression.

[0143] In addition, we speculated that if the length of the stem connecting the aptamer and twister is too long, the stem formation could be independent of the existence of aptamer sequence and occur in the absence of aptamer ligand. However, if the stem is too short, even in the presence of aptamer ligand, a stable stem structure may not be achieved. Therefore, to

determine an optimal length of the stem (to achieve optimal responsiveness to aptamer ligand), the length of the stem connecting aptamer and ribozyme was optimized. We made serial truncations to the stem, generating a total of 20 constructs with the length of the stem ranging from 17 to 1 bp. As shown in the Fig. 3b, of all the 20 constructs constructed, Luci-waspGua16 which has 3 bp of stem connecting the guanine aptamer directly to twister loop 2, showed the most profound reduction in luciferase gene expression (15 fold). Constructs with stem length shorter than 3 bp or longer than 7 bp showed no significant reduction upon guanine treatment.

[0144] The Luci-waspGua16 construct was then validated in a separate experiment, in which a control construct containing the cleavage-incapable mutant was included. As shown in Fig. 3c, upon guanine treatment, Luci-waspGua16 generated 92% (12-fold) reduction in luciferase activity when compared to the luciferase activity when no guanine was added. In contrast, neither construct pLuc nor Luci-mutwaspGua16 showed any reduction in luciferase activity upon guanine treatment. These results indicate that upon aptamer ligand treatment, the twister ribozyme activity was restored, leading to the reduced production of target gene expression. When guanosine was used at 1 mM as the aptamer ligand, construct Luci-waspG16 yielded 25-fold reduction in luciferase expression (Fig. 3d).

[0145] When the waspGua16 was inserted into the 3' UTR of a mouse erythropoietin (Epo) gene, upon guanine treatment, the production of Epo was reduced approximately 30% compared to Epo produced from the solvent-treated control cells. Our results demonstrate that we have generated guanine/guanosine responsive twister ribozyme, a synthetic twister riboswitch which suppresses target gene expression in response to aptamer ligand treatment.

EXAMPLE 4

[0146] Use of theophylline aptamer to regulate target gene expression via modulating twister ribozyme

[0147] Experimental procedure:

[0148] Oligos containing the twister ribozyme sequence linked with theophylline aptamer were synthesized (IDT), and cloned into pLuc vector using Gibson cloning strategy and kit (NEB). Construct sequences were verified by DNA sequencing (Genewiz).

[0149] Transfection and firefly luciferase assay: HEK 293 cells were transfected as described in Example 1. Four hours after transfection, the media was aspirated, and new

media with or without 2 mM theophylline (Sigma) was added. Luciferase assay was performed 20 to 24 hours after theophylline treatment as described in Example 2. The reduction fold was expressed as the quotient of luciferase activity obtained in the absence of aptamer ligand divided by the value obtained in the presence of the aptamer ligand.

[0150] Results:

[0151] We tested the use of an additional aptamer in modulating cleavage by twister ribozyme, thereby regulating target gene expression. Based on the 20 Luci-waspGua constructs (1-20) as described in Example 3, either 3 or 4 bp additional stem sequence was used in connecting the theophylline aptamer to loop 2 of twister ribozyme. In addition, the sequence of the stems was varied as shown in Fig. 4a. As shown in Fig. 4b, theophylline treatment resulted in reduction in the luciferase activity in all the constructs when compared to samples without theophylline treatment. Construct *twister_theo_3* had a 62% reduction in gene expression upon theophylline treatment. These data demonstrate that we have generated a theophylline responsive twister ribozyme—a synthetic twister riboswitch.

[0152] Grafting theophylline aptamer and stem sequences to loop 2 sequence in twister caused different effect on the disruption of twister ribozyme activity, as demonstrated in Fig. 4b in the absence of theophylline treatment. Construct *twister_theo_3* showed the most profound disruption of ribozyme activity, and therefore the most robust luciferase activity, in the absence of aptamer ligand. The difference between *twister_theo_2* and *twister_theo_3* is the length of the stem connecting aptamer and twister, which lead to a difference in disrupting the ribozyme activity. When comparing construct *twister_theo_1*, 3 and 4 that all have same length of stem (3 bp), the difference resides in the composition of the stem. These results indicate that the sequence composition of the stem also has impact on the ribozyme activity. Therefore, aptamer ligand-responsive ribozyme can be achieved through optimizing the length and the sequence composition of the effector stem region that connects aptamer and twister ribozyme.

EXAMPLE 5

[0153] Self-cleaving ribozymes suppress target gene expression when placed downstream of the translation start codon.

[0154] Experimental procedures:

[0155] Plasmid constructs: Synthetic dsDNA (IDT) and standard restriction cloning techniques were used to introduce test sequences into the previously created vector

pHDM.2b-LacZ (layout: CMV promoter – intron – LacZ – polyA). All constructs were verified by DNA sequencing.

[0156] Transfection: HEK293 cells (ATCC CRL-1573; grown in DMEM, 10% FBS, 37°C, 10% CO₂) were plated in 96-well flat bottom plates (100 µL) the day before transfection, such that confluence was about 70% at the time of transfection. Plasmid DNA (500 ng) was added to a tube or a 96-well U-bottom plate. Separately, TransIT-293 reagent (Mirus; 1.4 µL) was added to 50 µL Optimem I media (Life Technologies), and allowed to sit for 5 minutes at RT. Then, 50 µL of this diluted transfection reagent was added to the DNA, mixed, and incubated at RT for 20 min. Finally, 7 µL of this solution was added to a well of cells in a 96-well plate.

[0157] Beta-galactosidase (LacZ) assay: 24 hours after transfection, plates were removed from the incubator, and equilibrated to RT for several minutes on a lab bench, then aspirated. Glo-lysis buffer (Promega, 100 µL, RT) was added, and the plates allowed to remain at room temp at least 5 minutes. Then, the well contents were mixed by 50 µL trituration. A sample of 2 µL was taken from each well and diluted into 200 µL of glo-lysis buffer (101-fold dilution). The dilutions were mixed by trituration, and then 20 µL of each dilution was mixed with 20 µL of beta-glo reagent (Promega) in a solid-white 384-well plate. Thirty minutes later, luminescence was measured for relative light unit (RLU). At most, 96 wells of the 384 were used, in a pattern such that every assayed well was surrounded by either empty wells or the edge of the plate, to reduce cross-talk. All of the measurements fall in the linear range of the assay (determined by comparing various undiluted and diluted samples; data not shown).

[0158] Results: Constructs were made in which the only usable in-frame ATG for a LacZ reporter was upstream of either a hammerhead or hairpin ribozyme (disclosed in U.S. 2015/0056174, *see e.g.*, Fig. 1E, incorporated herein by reference) and for each we made three variants by putting the ribozyme sequence in each possible frame. In all constructs, the ribozyme was followed by linker-2A-peptide-LacZ. Versions with mutant (inactive) ribozymes were also made as controls. Mutations were introduced into each ribozyme to remove potential stop codons. The nucleotide and protein sequences are provided in Figs. 7a and 7b. (the top line in each case is the parent ribozyme, not translated through; dots in the nucleotide sequence are used to indicate “same as the top line.”)

[0159] These constructs and their inactivated ribozyme versions were transfected into HEK293 cells, and after 24 hours measured LacZ expression, as shown in Fig. 7c. The results from the hammerhead (HH) ribozymes are shown on the left in Fig 7c. Placing the HH ribozyme in the 5' UTR gave 162-fold reduction in expression compared to the inactive mutant ribozyme. However, translation-through (frame 2, arrow) gave a much improved 513-fold reduction in expression. Frame 1 does not function, which was expected due to in frame stop codons in the ribozyme sequence that could not be removed. The hairpin (HP) ribozymes are shown on the right in Fig. 7c. Placing the HP ribozyme in the 5' UTR gave 133-fold, but translation in frame 1 resulted in a 502-fold (arrow) reduction in expression. Other frames worked as well. We have made novel ribozyme constructs that show superior gene-control ability compared to published work. Gene expression constructs are described herein where the ribozyme is located not in the UTR, but rather located after the start codon. Control of target gene expression can be significantly enhanced by placing the ribozyme 3' of the translation start as compared to ribozyme placement in the 5' UTR. When the ribozyme is placed downstream of the translation start, self-cleavage removes the start codon from the rest of the mRNA of the target gene, resulting in decreased residual background expression.

EXAMPLE 6

[0160] Guanine-responsive twister riboswitch functions downstream translation start codon.

[0161] Experimental procedures:

[0162] Oligos containing either a start codon or coding sequence for the first 10 amino acids of GFP and waspGua16 followed by a linker2A sequence were synthesized (IDT) and cloned into the pLuc vector using the Gibson cloning strategy and cloning kit (NEB). Transfection and luciferase assay of cultured cells were performed as described in Example 3.

[0163] Results:

[0164] To test the portability of aptamer ligand-responsive twister ribozyme, we placed the waspGua16 sequence downstream of either (i) the translation start codon of the luciferase gene or (ii) the coding sequence for the first 10 aa of GFP, as illustrated in Fig. 5a. Fusing the twister riboswitch sequence with a downstream target gene would create an N-terminal tag when the target gene product is expressed, which may potentially affect the protein function. To remove this N-terminal tag, a linker2A sequence was inserted between

twister riboswitch and downstream target gene sequence. As shown in Fig. 5b, insertion of the twister riboswitch polynucleotide cassette did not impair luciferase gene expression in the absence of aptamer ligand, guanine, indicating the disrupted ribozyme activity. However, in the presence of guanine, luciferase gene expression was reduced 68% of the untreated control for both ATG_waspGua16 and 10aa_waspGua16 constructs, while the control constructs that contain the mutant twister sequence showed no reduction in luciferase expression in response to guanine treatment. This result demonstrates that the reduction in target gene expression with the polynucleotide cassette containing the functional twister riboswitches was due to restoration of twister ribozyme activity by aptamer/ligand binding. Our results indicate that twister riboswitch functions downstream start codon in the coding sequence, demonstrating the portability of the polynucleotide cassette containing the synthetic twister riboswitch.

EXAMPLE 7

[0165] Use of guanine aptamer to generate guanine responsive ON twister riboswitch

[0166] Experimental procedures:

[0167] Plasmid constructs: Oligos containing aptamer sequence linker to the P1 stem of twister ribozyme were synthesized (IDT) and cloned into pLuc vector using Gibson cloning strategy and cloning kit (NEB).

[0168] Transfection was performed as described in Example 1, and firefly luciferase assay was performed as described in Example 3.

[0169] Results:

[0170] The twister riboswitches described in Examples 3 and 4 suppress target gene expression in response to the aptamer ligand treatment, and thus are OFF twister riboswitches. Here, a different strategy was employed for linking the aptamer sequence to the twister ribozyme generating ON twister riboswitches that induce target gene expression upon aptamer ligand treatment. As shown in Fig. 6a, the 3' end of the twister P1 stem (including the additional 3 bp stem) is adjacent to the aptamer sequence. In this configuration, stem sequence is shared between the twister P1 stem and the aptamer P1 stem such that the shared sequence can either form the twister P1 stem or the aptamer P1 stem. In the absence of the aptamer ligand (top panel), the aptamer sequence adjacent to the twister P1 stem does not disrupt twister structure and its ribozyme cleavage activity remains intact, leading to cleavage of the mRNA and inhibition of target gene expression. In the presence of aptamer ligand

(lower panel), the ligand-bound aptamer forms the aptamer P1 stem with the shared sequence from the twister P1 stem thereby disrupting twister P1 stem formation, and disrupting its ribozyme activity and increasing target gene expression. As described in Examples 3 and 4, the length of the aptamer P1 stem is important in its regulating function in the context of a riboswitch. If the stem length is too long, the aptamer P1 formation becomes aptamer ligand independent, leading to unregulatable target gene expression. In addition, we speculated that the length of the shared sequence would be important as well. If the length of the shared stem sequence is too short, the twister ribozyme may not be affected by aptamer P1 stem formation.

[0171] ON twister riboswitches were created having either 6 or 7 nt of the twister P1 stem sequence shared with aptamer P1 stem. With the 6 or 7 nt shared stem sequences, the aptamer forms an 8 or 9 bp P1 stem. An xpt-guanine aptamer was linked to either the 5' or 3' arm of the twister P1 stem sequence, generating GT8 or TG8 and TG9 constructs. Linking the guanine aptamer to the 5' arm of the twister P1 stem (construct GT8) provided 1.4 fold induction of luciferase activity upon guanosine treatment, as shown in Fig. 6b, left panel. However, when aptamer was linked to the 3' end of the twister P1 stem as illustrated in Fig. 6a, the basal level of luciferase activity was reduced to 0.7% of the control vector that does not contain twister/aptamer sequences in 3' UTR. Upon guanosine treatment, luciferase activity was induced generating 8.6-fold and 9.2-fold induction from construct TG8 and TG9, respectively, as shown in Fig. 6b (middle and right panels). The low basal level expression of luciferase from construct TG8 and TG9 indicates that linking aptamer to the 3' end of P1 stem did not disrupt ribozyme activity, and upon aptamer/ligand binding, the twister ribozyme activity was impaired, thus leading to increased target gene expression. These results demonstrate that we have generated novel twister ON riboswitches that induce target gene expression in response to aptamer ligand treatment.

We claim:

1. A polynucleotide cassette for the regulation of the expression of a target gene comprising a riboswitch that comprises a twister ribozyme linked by a stem to an aptamer, wherein the stem linking the twister ribozyme to the aptamer attaches to the ribozyme at the location of the P3 stem of the twister ribozyme and wherein the target gene is linked to the P1 stem of the twister ribozyme.
2. The polynucleotide cassette of claim 1, wherein the aptamer binds a small molecule ligand.
3. The polynucleotide cassette of claim 1, wherein the twister ribozyme is from *Nasonia vitripennis*.
4. The polynucleotide cassette of claim 1, wherein the twister ribozyme comprises SEQ ID NO.:38.
5. The polynucleotide cassette of claim 1, wherein the twister ribozyme is from an environmental sample.
6. The polynucleotide cassette of claim 1, wherein the twister ribozyme comprises SEQ ID NO.:39.
7. The polynucleotide cassette of claims 1-6, comprising sequence that lengthens the P1 stem of the twister ribozyme by 1 to 3 base pairs.
8. The polynucleotide cassette of claims 1-7, wherein the P1 stem of the twister ribozyme is 4 to 7 base pairs.
9. The polynucleotide cassette of claims 1 to 8, wherein the stem linking the ribozyme to the aptamer comprises sequence from the ribozyme P3 stem and/or sequence from the aptamer P1 stem.
10. The polynucleotide cassette of claims 1 to 8, wherein the stem linking the ribozyme to the aptamer is 3 to 7 base pairs long.
11. A method of modulating the expression of a target gene comprising
 - a. inserting the polynucleotide cassette of claims 1-10 into a target gene,
 - b. introducing the target gene comprising the polynucleotide cassette into a cell,and

- c. exposing the cell to a small molecule ligand that specifically binds the aptamer in an amount effective to decrease expression of the target gene.
12. The method of claim 11, wherein the polynucleotide cassette is inserted into the 5' untranslated region of the target gene.
13. The method of claim 11, wherein the polynucleotide cassette is inserted into the 3' untranslated region of the target gene.
14. The method of claim 11, wherein the polynucleotide cassette is inserted downstream of the translation start codon.
15. The method of claim 14, wherein the polynucleotide cassette comprises a 2A peptide sequence between the 3' end of the riboswitch and the target gene coding sequence.
16. The method of claims 11-15, wherein two or more of the polynucleotide cassettes are inserted into the target gene.
17. The method of claim 16, wherein the two or more polynucleotide cassettes comprise different aptamers that specifically bind to different small molecule ligands.
18. The method of claim 16, wherein the two or more polynucleotide cassettes comprise the same aptamer.
19. The method of claim 11, wherein the target gene comprising the polynucleotide cassette is incorporated in a vector for the expression of the target gene.
20. The method of claim 19, wherein the vector is a viral vector.
21. The method of claim 20, wherein the viral vector is selected from the group consisting of adenoviral vector, adeno-associated virus vector, and lentiviral vector.
22. A vector comprising a target gene that contains a polynucleotide cassette of claims 1-10.
23. The vector of claim 22, wherein the vector is a viral vector.
24. The vector of claim 23, wherein the viral vector is selected from the group consisting of adenoviral vector, adeno-associated virus vector, and lentiviral vector.
25. A polynucleotide cassette for the regulation of the expression of a target gene comprising a riboswitch that comprises a twister ribozyme linked to an aptamer,

wherein the aptamer is linked to the 3' or 5' end of the twister ribozyme P1 stem, wherein when the aptamer is linked to the 3' end of the twister ribozyme P1 stem, a portion of the 3' arm of the twister ribozyme P1 stem is alternatively part of the 5' arm of the aptamer P1 stem, and wherein when the aptamer is linked to the 5' end of the twister ribozyme P1 stem, a portion of the 5' arm of the twister ribozyme P1 stem is alternatively part of the 3' arm of the aptamer P1 stem.

26. The polynucleotide cassette of claim 25, wherein the aptamer binds a small molecule ligand.
27. The polynucleotide cassette of claims 25-26, wherein the twister ribozyme is from *Nasonia vitripennis*.
28. The polynucleotide cassette of claims 25-26, wherein the twister ribozyme comprises SEQ ID NO.:38.
29. The polynucleotide cassette of claims 25-26, wherein the twister ribozyme is from an environmental sample.
30. The polynucleotide cassette of claims 25-26, wherein the twister ribozyme is SEQ ID NO.:39.
31. The polynucleotide cassette of claims 25-30, wherein the portion of the arm of the twister ribozyme P1 stem that is alternatively an arm of the aptamer P1 stem is 5 to 8 nucleotides.
32. The polynucleotide cassette of claims 25-31, wherein the P1 stem of the twister ribozyme is 4 to 9 base pairs.
33. The polynucleotide cassette of claims 25-31, wherein the P1 stem of the aptamer is 5 to 8 base pairs.
34. A method of modulating the expression of a target gene comprising
 - a. inserting the polynucleotide cassette of claims 25-33 into a target gene,
 - b. introducing the target gene comprising the polynucleotide cassette into a cell, and
 - c. exposing the cell to a small molecule ligand that specifically binds the aptamer in an amount effective to increase expression of the target gene.

35. The method of claim 34, wherein the polynucleotide cassette is inserted into the 5' untranslated region of the target gene.
36. The method of claim 34, wherein the polynucleotide cassette is inserted into the 3' untranslated region of the target gene.
37. The method of claim 34, wherein the polynucleotide cassette is inserted downstream of the translation start codon.
38. The method of claim 37, wherein the polynucleotide cassette comprises a 2A peptide sequence between the 3' end of the riboswitch and the target gene coding sequence.
39. The method of claims 34 to 38, wherein two or more of the polynucleotide cassettes are inserted into the target gene.
40. The method of claim 39, wherein the two or more polynucleotide cassettes comprise different aptamers that specifically bind to different small molecule ligands.
41. The method of claim 39, wherein the two or more polynucleotide cassettes comprise the same aptamer.
42. The method of claims 34-41, wherein the target gene comprising the polynucleotide cassette is incorporated in a vector for the expression of the target gene.
43. The method of claim 42, wherein the vector is a viral vector.
44. The method of claim 43, wherein the viral vector is selected from the group consisting of adenoviral vector, adeno-associated virus vector, and lentiviral vector.
45. A vector comprising a target gene that contains a polynucleotide cassette of claims 25-33.
46. The vector of claim 45, wherein the vector is a viral vector.
47. The vector of claim 46, wherein the viral vector is selected from the group consisting of adenoviral vector, adeno-associated virus vector, and lentiviral vector.




Fig. 1a.




Fig. 2a.




Fig. 2b.




Fig. 2c.




Fig. 3a.




Fig. 3b.




Fig. 3c.




Fig. 3d.




Fig. 3e.




Fig. 4a.




Fig. 4b.

ATG_waspGua16



10aa_waspGua16




Fig. 5a.




Fig. 5b.




Fig. 6a.




Fig. 6b.




Fig. 7a.




Fig. 7b.




Fig. 7c.